



HU000030426T2

(19) **HU**(11) Lajstromszám: **E 030 426**(13) **T2****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 08 797046**(22) A bejelentés napja: **2008. 08. 01.**(51) Int. Cl.: **C12N 9/24** (2006.01)**C11D 33/86** (2006.01)**C12P 7/10** (2006.01)**C12P 19/14** (2006.01)(96) Az európai bejelentés bejelentési száma:
EP 20080797046(97) Az európai bejelentés közzétételi adatai:
EP 2183363 A2 **2009. 02. 05.**

(86) A nemzetközi (PCT) bejelentési szám:

PCT/US 08/071982(97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2183363 B1 **2016. 06. 08.**

(87) A nemzetközi közzétételi szám:

WO 09018537

(30) Elsőbbségi adatok:

833133**2007. 08. 02.****US**

(73) Jogosult(ak):

Danisco US Inc., Palo Alto, CA 94304 (US)

(72) Feltaláló(k):

GUSAKOV, Alexander, Vasilievich, Moscow 115054 (RU)**PUNT, Peter, J., NL-3994 XT Houten (NL)****VERDOES, Jan, Cornelis, deceased (NL)****SINITSYN, Arkady, Panteleimonovich, Moscow 119590 (RU)****VLASENKO, Elena, Davis, CA 95616 (US)****HINZ, Sandra Wihelmina, Agnes, NL-6708 SN****Wageningen (NL)****GOSINK, Mark, Wellington, FL 33414 (US)****JIANG, Zhijie, West Palm Beach, FL 33401 (US)**

(74) Képviselő:

**Danubia Szabadalmi és Jogi Iroda Kft.,
Budapest**

(54)

Gombaxilanáz

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmat az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

(19)



(11)

EP 2 183 363 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
08.06.2016 Bulletin 2016/23

(51) Int Cl.:
C12N 9/24 (2006.01) **C12P 7/10** (2006.01)
C12P 19/14 (2006.01) **C11D 3/386** (2006.01)

(21) Application number: **08797046.3**

(86) International application number:
PCT/US2008/071982

(22) Date of filing: **01.08.2008**

(87) International publication number:
WO 2009/018537 (05.02.2009 Gazette 2009/06)

(54) **FUNGAL XYLANASE**

XYLANASE AUS PILZ

XYLANASE FONGIQUES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT
RO SE SI SK TR**

(30) Priority: **02.08.2007 US 833133**

(43) Date of publication of application:
12.05.2010 Bulletin 2010/19

(73) Proprietor: **Danisco US Inc.**
Palo Alto, CA 94304 (US)

(72) Inventors:

- **GUSAKOV, Alexander, Vasilievich**
Moscow 115054 (RU)
- **PUNT, Peter, J.**
NL-3994 XT Houten (NL)
- **VERDOES, Jan, Cornelis**
deceased (NL)
- **SINITSYN, Arkady, Panteleimonovich**
Moscow 119590 (RU)
- **VLASENKO, Elena**
Davis, CA 95616 (US)
- **HINZ, Sandra Wihelmina, Agnes**
NL-6708 SN Wageningen (NL)
- **GOSINK, Mark**
Wellington, FL 33414 (US)
- **JIANG, Zhijie**
West Palm Beach, FL 33401 (US)

(74) Representative: **Nederlandsch Octrooibureau**
P.O. Box 29720
2502 LS The Hague (NL)

(56) References cited:

WO-A1-98/15633	WO-A2-00/20555
WO-A2-01/79507	WO-A2-2008/008070
US-A1- 2004 002 136	US-A1- 2006 053 514
US-A1- 2006 134 747	US-A1- 2006 218 671
US-A1- 2007 077 630	

- **GUSAKOV ALEXANDER V ET AL: "Purification, cloning and characterisation of two forms of thermostable and highly active cellobiohydrolase I (Cel7A) produced by the industrial strain of Chrysosporium lucknowense" ENZYME AND MICROBIAL TECHNOLOGY, vol. 36, no. 1, 6 January 2005 (2005-01-06), pages 57-69, XP002608583 ISSN: 0141-0229**
- **GUSAKOV ALEXANDER V ET AL: "Design of highly efficient cellulase mixtures for enzymatic hydrolysis of cellulose" BIOTECHNOLOGY AND BIOENGINEERING, WILEY & SONS, HOBOKEN, NJ, US LNKD- DOI:10.1002/BIT.21329, vol. 97, no. 5, 1 August 2007 (2007-08-01) , pages 1028-1038, XP002565359 ISSN: 0006-3592 [retrieved on 2007-01-12]**
- **DATABASE UniProt [Online] 1 May 1997 (1997-05-01), likura et al.: "Cloning of a Gene Encoding a Putative Xylanase with a Cellulose-Binding Domain from Humicola grisea" XP002608584 retrieved from EBI Database accession no. P79046**
- **DATABASE UniProt [Online] 21 March 1996 (1996-03-21), Birren et al.: "Annotation of the Chaetomium globosum CBS 148.51 genome" XP002608585 retrieved from EBI Database accession no. Q2HIC4**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 183 363 B1

- DATABASE Geneseq [Online] 19 February 1997 (1997-02-19), Ikura et al.: "Cloning of a Gene Encoding a Putative Xylanase with a Cellulose-Binding Domain from *Humicola grisea*" XP002608586 retrieved from EBI Database accession no. AB001030
- DATABASE Geneseq [Online] 18 October 2007 (2007-10-18), "Acremonium thermophilum XYN60/Xyn10A gene." XP002608587 retrieved from EBI accession no. GSN:AGD76041 Database accession no. AGD76041
- MARKOV A V ET AL: "Properties of hemicellulases of the enzyme complex from *Trichoderma longibrachiatum*" APPLIED BIOCHEMISTRY AND MICROBIOLOGY, KLUWER ACADEMIC PUBLISHERS-PLENUM PUBLISHERS, NE LNKD- DOI:10.1134/S000368380606007X, vol. 42, no. 6, 1 December 2006 (2006-12-01), pages 573-583, XP019432666 ISSN: 1608-3024
- USTINOV B B ET AL: "Comparison of properties and mode of action of six secreted xylanases from *Chrysosporium lucknowense*" ENZYME AND MICROBIAL TECHNOLOGY, STONEHAM, MA, US LNKD- DOI:10.1016/J.ENZMICTEC.2008.01.017, vol. 43, no. 1, 7 July 2008 (2008-07-07), pages 56-65, XP022683403 ISSN: 0141-0229 [retrieved on 2008-02-07]
- HINZ SANDRA W A ET AL: "Hemicellulase production in *Chrysosporium lucknowense* C1" JOURNAL OF CEREAL SCIENCE, ACADEMIC PRESS LTD, XX LNKD- DOI:10.1016/J.JCS.2009.07.005, vol. 50, no. 3, 1 November 2009 (2009-11-01), pages 318-323, XP002593773 ISSN: 0733-5210 [retrieved on 2009-08-20]

Description**FIELD OF THE INVENTION**

[0001] This invention relates to novel enzymes and novel methods for producing the same. More specifically this invention relates to enzymes produced by fungi. The invention also relates to a method to convert lignocellulosic biomass to fermentable sugars with enzymes that degrade the lignocellulosic material and novel combinations of enzymes, including those that provide a synergistic release of sugars from plant biomass. The invention also relates to methods to use the novel enzymes and compositions of such enzymes in a variety of other processes, including washing of clothing, detergent processes, deinking and biobleaching of paper and pulp, and treatment of waste streams.

BACKGROUND OF THE INVENTION

[0002] Large amounts of carbohydrates in plant biomass provide a plentiful source of potential energy in the form of sugars (both five carbon and six carbon sugars) that can be utilized for numerous industrial and agricultural processes. However, the enormous energy potential of these carbohydrates is currently under-utilized because the sugars are locked in complex polymers, and hence are not readily accessible for fermentation. These complex polymers are often referred to collectively as lignocellulose. Sugars generated from degradation of plant biomass potentially represent plentiful, economically competitive feedstocks for fermentation into chemicals, plastics, and fuels, including ethanol as a substitute for petroleum.

[0003] For example, distillers' dried grains (DDG) are lignocellulosic byproducts of the corn dry milling process. Milled whole corn kernels are treated with amylases to liquefy the starch within the kernels and hydrolyze it to glucose. The glucose so produced is then fermented in a second step to ethanol. The residual solids after the ethanol fermentation and distillation are centrifuged and dried, and the resulting product is DDG, which is used as an animal feed stock. Although DDG composition can vary, a typical composition for DDG is: 32% hemicellulose, 22% cellulose, 30% protein, 10% lipids, 4% residual starch, and 4% inorganics. In theory, the cellulose and hemicellulose fractions, comprising about 54% of the weight of the DDG, can be efficiently hydrolyzed to fermentable sugars by enzymes; however, it has been found that the carbohydrates comprising lignocellulosic materials in DDG are more difficult to digest. To date, the efficiency of hydrolysis of these (hemi) cellulosic polymers by enzymes is much lower than the hydrolytic efficiency of starch, due to the more complex and recalcitrant nature of these substrates. Accordingly, the cost of producing the requisite enzymes is higher than the cost of producing amylases for starch hydrolysis.

[0004] Major polysaccharides comprising lignocellulosic materials include cellulose and hemicelluloses. The enzymatic hydrolysis of these polysaccharides to soluble sugars (and finally to monomers such as glucose, xylose and other hexoses and pentoses) is catalyzed by several enzymes acting in concert. For example, endo-1,4- β -glucanases (EGs) and exo-cellobiohydrolases (CBHs) catalyze the hydrolysis of insoluble cellulose to cellobiosaccharides (with cellobiose the main product), while β -glucosidases (BGLs) convert the oligosaccharides to glucose. Similarly, xylanases, together with other enzymes such as α -L-arabinofuranosidases, feruloyl and acetyl xylan esterases and β -xylosidases, catalyze the hydrolysis of hemicelluloses.

[0005] Regardless of the type of cellulosic feedstock, the cost and hydrolytic efficiency of enzymes are major factors that restrict the widespread use of biomass bioconversion processes. The hydrolytic efficiency of a multi-enzyme complex in the process of lignocellulosic saccharification depends both on properties of the individual enzymes and the ratio of each enzyme within the complex.

[0006] Enzymes useful for the hydrolysis of complex polysaccharides are also highly useful in a variety of industrial textile applications, as well as industrial paper and pulp applications, and in the treatment of waste streams. For example, as an alternative to the use of pumice in the stone washing process, methods for treating cellulose-containing fabrics for clothing with hydrolytic enzymes, such as cellulases, are known to improve the softness or feel of such fabrics. Cellulases are also used in detergent compositions, either for the purpose of enhancing the cleaning ability of the composition or as a softening agent. Cellulases are also used in combination with polymeric agents in processes for providing a localized variation in the color density of fibers. Such enzymes can also be used for the saccharification of lignocellulosic biomass in waste streams, such as municipal solid waste, for biobleaching of wood pulp, and for deinking of recycled print paper. As with the hydrolysis of these polysaccharides in lignocellulosic materials for use as feedstocks described above, the cost and hydrolytic efficiency of the enzymes are major factors that control the use of enzymes in these processes.

[0007] Filamentous fungi are a source of cellulases and hemicellulases, as well as other enzymes useful in the enzymatic hydrolysis of major polysaccharides. In particular, strains of *Trichoderma* sp., such as *T. viride*, *T. reesei* and *T. longibrachiatum*, and *Penicillium* sp., and enzymes derived from these strains, have previously been used to hydrolyze crystalline cellulose. However, the costs associated with producing enzymes from these fungi, as well as the presence of additional, undesirable enzymes, remains a drawback. It is therefore desirable to produce inexpensive enzymes and

enzyme mixtures that efficiently degrade cellulose and hemicellulose for use in a variety of agricultural and industrial applications.

[0008] GUSAKOV ALEXANDER V ET AL: ENZYME AND MICROBIAL TECHNOLOGY, vol 36, No. 1, 6 January 2005, pages 57-69, discloses the purification and cloning of two thermostable cellobiohydrolases from a strain of the fungus *Chrysosporium lucknowense*, which fungus is reported to be an industrial producer of cellulases and hemicellulases.

[0009] GUSAKOV ALEXANDER V ET AL: BIOTECHNOLOGY AND BIOENGINEERING, vol. 97, No. 5, 1 August 2007, pages 1028-1038, discloses various enzymes from *C. lucknowense*, including cellobiohydrolase, endoglucanases and xylanase II.

[0010] WO 98/15633 A1 discloses enzymes and genes of *F. luckowense* and mentions the presence of xylanases in said fungus.

[0011] WO 01/79507 A2 discloses the DNA sequence of the *C. lucknowense* xylanase XylII gene and the amino acid sequence of the encoded XylII protein.

SUMMARY OF THE INVENTION

[0012] The present invention provides isolated nucleic acid molecules comprising a nucleic acid sequence selected from the group consisting of:

- a) a nucleic acid sequence encoding a protein consisting of an amino acid sequence selected from the group consisting of: SEQ ID NO:32;
- b) a nucleic acid sequence encoding an amino acid sequence with xylanase activity that is at least 90% identical to a full length amino acid sequence of (a);
- c) a nucleic acid sequence selected from the group consisting of: SEQ ID NO:31, and SEQ ID NO:33;
- d) a nucleic acid sequence that is fully complementary to the nucleic acid sequences of (a), (b) or (c)

[0013] Such an isolated nucleic acid molecule is defined in independent claim 1 appended hereto. Embodiments of the invention are defined in appended claims which are dependent on claim 1 and to which reference should now be made.

[0014] In some embodiments, the nucleic acid sequence of (b) encodes an amino acid sequence that is at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of (a) and has a biological activity of the protein comprising the amino acid sequence.

[0015] The invention also provides a protein as defined in claim 2 appended hereto.

[0016] The invention also provides a composition for the degradation of a lignocellulosic material comprising at least the isolated protein of the present invention.

[0017] The invention also provides a recombinant nucleic acid molecule comprising an isolated nucleic acid molecule of the present invention, operatively linked to at least one expression control sequence.

[0018] In some embodiments, the recombinant nucleic acid molecule comprises an expression vector. In some embodiments, the recombinant nucleic acid molecule comprises a targeting vector.

[0019] The invention also provides an isolated host cell transfected with a nucleic acid molecule of the present invention.

[0020] In some embodiments, the host cell is a fungus. In some embodiments, the host cell is a filamentous fungus. In some embodiments, the filamentous fungus is from a genus selected from the group consisting of: *Chrysosporium*, *Thielavia*, *Neurospora*, *Aureobasidium*, *Filibasidium*, *Piromyces*, *Corynascus*, *Cryptococcus*, *Acremonium*, *Tolyocladium*, *Scytalidium*, *Schizophyllum*, *Sporotrichum*, *Penicillium*, *Gibberella*, *Myceliophthora*, *Mucor*, *Aspergillus*, *Fusarium*, *Humicola*, and *Trichoderma*, and anamorphs and teleomorphs thereof. In some embodiments, the host cell is a bacterium.

[0021] The invention also provides a method for producing a protein of the present invention, comprising culturing a cell that has been transfected with a nucleic acid molecule comprising a nucleic acid sequence encoding the protein, and expressing the protein with the transfected cell.

[0022] In some embodiments, the method further comprises recovering the protein from the cell or from a culture comprising the cell.

[0023] The invention also provides a genetically modified organism comprising components suitable for degrading a lignocellulosic material to fermentable sugars, wherein the organism has been genetically modified to express at least the protein of the present invention.

[0024] In some embodiments, the genetically modified organism is a microorganism. In some embodiments, the microorganism is a filamentous fungus.

[0025] In some embodiments, the filamentous fungus is from a genus selected from the group consisting of: *Chrysosporium*, *Thielavia*, *Neurospora*, *Aureobasidium*, *Filibasidium*, *Piromyces*, *Corynascus*, *Cryptococcus*, *Acremonium*, *Tolyocladium*, *Scytalidium*, *Schizophyllum*, *Sporotrichum*, *Penicillium*, *Talaromyces*, *Gibberella*, *Myceliophthora*, *Mucor*, *Aspergillus*, *Fusarium*, *Humicola*, and *Trichoderma*.

[0026] In some embodiments, the filamentous fungus is selected from the group consisting of: *Trichoderma reesei*,

Chrysosporium lucknowense, *Aspergillus japonicus*, *Penicillium canescens*, *Penicillium solitum*, *Penicillium funiculosum*, and *Talaromyces flavus*.

[0027] In some embodiments, the organism has been genetically modified to express at least one additional enzyme.

[0028] In some embodiments, the additional enzyme is an accessory enzyme selected from the group consisting of: cellulase, xylanase, ligninase, glucuronidase, arabinofuranosidase, arabinogalactanase, ferulic acid esterase, lipase, pectinase, glucomannase, amylase, laminarinase, xyloglucanase, galactanase, glucoamylase, pectate lyase, chitinase, exo- β -D-glucosaminidase, cellobiose dehydrogenase, and acetylxyylan esterase.

[0029] In some embodiments, the genetically modified organism is a plant.

[0030] The invention also provides a multi-enzyme composition comprising at least the protein of the present invention and at least one additional protein for degrading a lignocellulosic material or a fragment thereof that has biological activity.

[0031] In some embodiments, the at least one additional protein for degrading a lignocellulosic material to fermentable sugars is selected from the group consisting of: SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:17, SEQ ID NO:20, SEQ ID NO:26 and SEQ ID NO:47, or a fragment thereof that has biological activity.

[0032] In some embodiments, the multi-enzyme composition comprises at least one cellobiohydrolase, at least one xylanase, at least one endoglucanase, at least one β -glucosidase, at least one β -xylosidase, and at least one accessory enzyme.

[0033] In some embodiments, between about 50% and about 70% of the enzymes in the composition are cellobiohydrolases.

[0034] In some embodiments, between about 10% and about 30% of the enzymes in the composition are xylanases.

[0035] In some embodiments, between about 5% and about 15% of the enzymes in the composition are endoglucanases.

[0036] In some embodiments, between about 1% and about 5% of the enzymes in the composition are β -glucosidases.

[0037] In some embodiments, between about 1% and about 3% of the enzymes in the composition are β -xylosidases.

[0038] In some embodiments, the composition comprises about 60% cellobiohydrolases, about 20% xylanases, about 10% endoglucanases, about 3% β -glucosidases, about 2% β -xylosidases, and about 5% accessory enzymes.

[0039] In some embodiments, the xylanases are selected from the group consisting of: endoxylanases, exoxylanases, and β -xylosidases.

[0040] In some embodiments, the accessory enzymes include an enzyme selected from the group consisting of: ligninase, glucuronidase, arabinofuranosidase, arabinogalactanase, ferulic acid esterase, lipase, pectinase, glucomannase, amylase, laminarinase, xyloglucanase, galactanase, glucoamylase, pectate lyase, chitinase, exo- β -D-glucosaminidase, cellobiose dehydrogenase, and acetylxyylan esterase.

[0041] In some embodiments, the composition comprises at least one protein comprising the amino acid sequence of SEQ ID NO:11, wherein the protein has cellobiohydrolase activity, at least one protein comprising an amino acid sequence selected from the group consisting of: SEQ ID NO:94 and SEQ ID NO:23, wherein the protein has endoglucanase activity, at least one protein comprising an amino acid sequence selected from the group consisting of: SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38 and SEQ ID NO:41, wherein the protein has xylanase activity, at least one protein comprising an amino acid sequence of SEQ ID NO:44, wherein the protein has β -glucosidase activity, and at least one protein comprising an amino acid sequence selected from the group consisting of: SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:62, SEQ ID NO:65, SEQ ID NO:68, SEQ ID NO:71, SEQ ID NO:74, SEQ ID NO:77, SEQ ID NO:80, SEQ ID NO:83, SEQ ID NO:86, SEQ ID NO:89, and SEQ ID NO:92, wherein the protein has hemicellulase activity.

[0042] In some embodiments, the composition comprises at least one first protein comprising an amino acid sequence that is at least 95% identical to an amino acid sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 and SEQ ID NO:11 and has cellobiohydrolase activity, or a fragment thereof that has cellobiohydrolase activity; at least one second protein comprising an amino acid sequence that is at least 95% identical to an amino acid sequence selected from the group consisting of: SEQ ID NO:14, SEQ ID NO:94, SEQ ID NO:17, SEQ ID NO:20 and SEQ ID NO:23 and has endoglucanase activity, or a fragment thereof that has endoglucanase activity; and at least one third protein comprising an amino acid sequence that is at least 95% identical to an amino acid sequence selected from the group consisting of: SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38 and SEQ ID NO:41 and has xylanase activity, or a fragment thereof that has xylanase activity.

[0043] In some embodiments, the composition further comprises a fourth protein comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:44 and has β -glucosidase activity, or a fragment thereof that has β -glucosidase activity.

[0044] In some embodiments, the composition further comprises at least one fifth protein comprising an amino acid sequence that is at least 95% identical to an amino acid sequence selected from the group consisting of: SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:62, SEQ ID NO:65, SEQ ID NO:68, SEQ ID NO:71, SEQ ID NO:74, SEQ ID NO:77, SEQ ID NO:80, SEQ ID NO:83, SEQ ID NO:86, SEQ ID NO:89, and SEQ ID NO:92, and has hemicellulase or chitinase activity, or a fragment thereof that has hemicellulase or chitinase activity.

[0045] The invention also provides a method for degrading a lignocellulosic material to fermentable sugars, comprising

contacting the lignocellulosic material with at least the isolated protein of the present invention.

[0046] In some embodiments, the isolated protein is part of a multi-enzyme composition.

[0047] The invention also provides a method for degrading a lignocellulosic material to fermentable sugars, comprising contacting the lignocellulosic material with at least one multi-enzyme composition of the present invention.

[0048] The invention also provides a method for producing an organic substance, comprising:

- a) saccharifying a lignocellulosic material with a multi-enzyme composition of the present invention;
- b) fermenting the saccharified lignocellulosic material obtained with one or more fermentating microorganisms; and
- c) recovering the organic substance from the fermentation.

[0049] In some embodiments, the steps of saccharifying and fermenting are performed simultaneously.

[0050] In some embodiments, the organic substance is an alcohol, organic acid, ketone, amino acid, or gas.

[0051] In some embodiments, the organic substance is an alcohol. In some embodiments, the alcohol is ethanol.

[0052] In some embodiments, the lignocellulosic material is selected from the group consisting of her-
baceous material, agricultural residue, forestry residue, municipal solid waste, waste paper, and pulp and paper mill
residue.

[0053] In some embodiments, the lignocellulosic material is distiller's dried grains or distiller's dried grains with solubles.

[0054] In some embodiments, the distiller's dried grains or distiller's dried grains with solubles is derived from corn.

[0055] The invention also provides a method for degrading a lignocellulosic material consisting of distiller's dried grains
or distiller's dried grains with solubles to sugars, the method comprising contacting the distiller's dried grains or distiller's
dried grains with solubles with a multi-enzyme composition, whereby at least 10% of the fermentable sugars are liberated,
wherein the multi-enzyme composition is the multi-enzyme composition of the present invention.

[0056] In some embodiments, at least 15% of the sugars are liberated. In some embodiments, at least 20% of the
sugars are liberated. In some embodiments, at least 23% of the sugars are liberated.

[0057] In some embodiments, the distiller's dried grains or distiller's dried grains with solubles is derived from corn.

[0058] In some embodiments, the method further comprises a pretreatment process for pretreating the lignocellulosic
material.

[0059] In some embodiments, the pretreatment process is selected from the group consisting of physical treatment,
metal ion, ultraviolet light, ozone, organosolv treatment, steam explosion treatment, lime impregnation with steam ex-
plosion treatment, hydrogen peroxide treatment, hydrogen peroxide/ozone (peroxone) treatment, acid treatment, dilute
acid treatment, and base treatment.

[0060] In some embodiments, the pretreatment process is selected from the group consisting of organosolv, steam
explosion, heat treatment and AFEX.

[0061] In some embodiments, the heat treatment comprises heating the lignocellulosic material to 121°C for 15 minutes.

[0062] In some embodiments, the method further comprises detoxifying the lignocellulosic material.

[0063] In some embodiments, the method further comprises recovering the fermentable sugar.

[0064] In some embodiments, the sugar is selected from the group consisting of glucose, xylose, arabinose, galactose,
mannose, rhamnose, sucrose and fructose.

[0065] In some embodiments, the method further comprises recovering the contacted lignocellulosic material after the
fermentable sugars are degraded.

[0066] The invention also provides a method for stonewashing a fabric, comprising contacting the fabric with at least
the isolated protein of the present invention..

[0067] The invention also provides a method for stonewashing a fabric, comprising contacting the fabric with at least
one multi-enzyme composition of the present invention.

[0068] In some embodiments, the fabric is denim.

[0069] The invention also provides a method for enhancing the softness or feel of a fabric or depilling a fabric,
comprising contacting the fabric with at least the isolated protein of the present inventionThe invention also provides a
method for enhancing the softness or feel of a fabric or depilling a fabric, comprising contacting the fabric with at least
one multi-enzyme composition of the present invention.

[0070] The invention also provides a method for restoring color to or brightening a fabric, comprising contacting the
fabric with at least the isolated protein of the present invention.

[0071] The invention also provides a method for restoring color to or brightening a fabric, comprising contacting the
fabric with at least one multi-enzyme composition of the present invention.

[0072] The invention also provides a method of biopolishing, defibrillating, bleaching, dyeing or desizing a fabric,
comprising contacting the fabric with at least the isolated protein of the present invention.

[0073] The invention also provides a method of biopolishing, defibrillating, bleaching, dyeing or desizing a fabric,
comprising contacting the fabric with at least one multi-enzyme composition of the present invention.

[0074] The invention also provides a method of deinking or biobleaching paper or pulp, comprising contacting the

paper or pulp with at least the isolated protein of the present invention.

[0075] The invention also provides a method of deinking or biobleaching paper or pulp, comprising contacting the paper or pulp with at least one multi-enzyme composition of the present invention.

[0076] The invention also provides a detergent composition, comprising at least the isolated protein of the present invention and at least one surfactant.

[0077] The invention also provides a detergent composition, comprising at least one multi-enzyme composition of the present invention and at least one surfactant.

BRIEF DESCRIPTION OF THE FIGURES

[0078]

Figure 1 shows the specific activity of cellobiohydrolases and endoglucanases toward the indicated substrates. Specific activities (Units/mg) were determined at pH 5.0 and 50°C.

Figure 2 illustrates the enzymatic activity (percentage of maximum) of cellobiohydrolases at varying pH levels. The activities were determined using an Avicel substrate at 40°C.

Figure 3 illustrates the enzymatic activity (percentage of maximum) of cellobiohydrolases at varying temperature levels. The activities were determined using an Avicel substrate at pH 5.0.

Figure 4 illustrates the enzymatic activity (percentage of maximum) of endoglucanases at varying pH levels. The activities were determined using a CMC substrate at 50°C.

Figure 5 illustrates the enzymatic activity (percentage of maximum) of endoglucanases at varying temperature levels. The activities were determined using a CMC substrate at pH 5.0.

Figure 6 shows the specific activity of the xylanase of the present invention and xylanases of the disclosure toward the indicated substrates. Specific activities (Units/mg) were determined at pH 5.0 and 50°C.

Figure 7 illustrates the enzymatic activity (percentage of maximum) of the xylanase of the present invention and xylanases of the disclosure at varying pH levels. The activities were determined using a soluble xylan substrate at 40°C.

Figure 8 illustrates the enzymatic activity (percentage of maximum) of the xylanase of the present invention and xylanases of the disclosure at varying temperature levels. The activities were determined using a soluble xylan substrate at pH 5.0.

Figure 9 illustrates the enzymatic activity (percentage of maximum) of enzymes at varying pH levels.

Figure 10 illustrates the enzymatic activity (percentage of maximum) of enzymes at varying temperature levels.

Figure 11 shows a standard calibration plot for the Somogyi-Nelson assay used for the measurement of reducing sugars in enzymatic hydrolyzates.

Figure 12 shows a standard calibration plot with glucose standards from 0.025 g/l to 0.5 g/l for a glucosooxidase-peroxidase assay used to determine glucose levels.

Figure 13 shows the relative activity over time of xylanase at 20-, 10- and 5-fold dilutions in a thermoshock assay.

Figure 14 shows the pelletizing stability of the xylanase of the present invention and xylanases of disclosure. The percentage of activity after 20 seconds at 80°C relative to an unheated sample is shown.

Figure 15 shows the relative viscosity of water/rye mixes treated with the indicated enzyme(s). To compare the efficiency of the preparations, the xylanase activities of the samples (units per flask) are also indicated.

Figure 16 shows a plot of the residual viscosity (%) of water/rye mixes) after enzymatic treatment.

Figure 17 shows a graph of the residual viscosity (%) of rye meal water extract after a 1-hour incubation with 1 unit of the indicated xylanase (*in vitro* feed test).

Figure 18 shows the biobleaching ability (A_{235}) of the xylanase of the present invention and xylanases of the disclosure on soft wood pulp at pH 7.5 and 50°C for 2 hours. For each xylanase, the assay was conducted using 0.01, 0.003, 0.001, 0.0003 mg/ml of enzyme in the reaction mixture, as indicated.

Figure 19 shows the biobleaching ability (A_{235}) of the xylanase of the present invention and xylanases of the disclosure on hard wood pulp at pH 7.5 and 50°C for 2 hours. For each xylanase, the assay was conducted using 0.01, 0.003, 0.001, 0.0003 mg/ml of enzyme in the reaction mixture, as indicated.

Figure 20 illustrates the bioscouring ability of cellobiohydrolases and endoglucanases of the present invention. The reciprocal of the time ($1/\text{time}$; sec^{-1}) of water drop adsorption on a textile after treatment at pH 5.0 with 0.6 mg/l of enzyme followed by heating is shown. The water drop adsorption time for the untreated control sample is 2400 sec ($1/\text{time} = 0.0004$).

Figure 21 illustrates a comparison of the pectin content in an unsoured sample, an alkaline scoured sample, and bioscoured samples.

Figure 22 shows a diagram of cells for the enzymatic and mechanical stress treatment of fabric. In Figure 22A, the following numbers represent the indicated structure: 1: glass vessel, 2: plastic tube, 3: denim swatch, 4: rubber ring,

5: buckshot balls, and 6: teflon disk. In Figure 22B, the following numbers represent the indicated structure: 1: lid with a thread inside, 2: rubber ring, 3: cell body (stainless steel), 4: cylindrical cartridge, 5: stainless steel cylinders, 6: circular denim swatch.

Figure 23 illustrates the biopolishing ability of cellobiohydrolases and endoglucanases of the present invention. The dye release (D 400 nm) after treatment with 0.2g/l enzyme at pH 5.0, 50°C is shown.

Figure 24 shows an example histogram generated from the color intensity analysis of a denim swatch.

Figure 25 illustrates the denim-washing (abrasion) ability of cellobiohydrolases and endoglucanases of the present invention. The relative units per mg of enzyme determined at 50°C, pH 5.0 is shown.

Figure 26 shows a graph of the saccharification ability of cellobiohydrolases and endoglucanases. The amount of glucose (g/l) produced after the hydrolysis of 5 g/l Avicel at pH 5.0, 40°C for 120 hours in the presence of 0.1 mg/ml enzyme and 0.2 U/ml cellobiase is shown.

DETAILED DESCRIPTION OF THE INVENTION

[0079] The present disclosure relates generally to proteins that play a role in the degradation of cellulose and hemicellulose and nucleic acids encoding the same. In particular, the present invention relates to a xylanase isolated from a filamentous fungal strain denoted herein as C1 (Accession No. VKM F-3500-D), nucleic acids encoding the enzyme, and methods of producing and using the enzyme. The invention also provides compositions that include at least the xylanase of the invention described herein for uses including, but not limited to, the hydrolysis of lignocellulose. The invention and disclosure stems, in part, from the discovery of a variety of novel cellulases and hemicellulases produced by the C1 fungus that exhibit high activity toward cellulose and other components of biomass.

[0080] The present invention also provides methods and compositions for the conversion of plant biomass to fermentable sugars that can be converted to useful products. The methods include methods for degrading lignocellulosic material using enzyme mixtures to liberate sugars. The compositions of the invention include enzyme combinations that break down lignocellulose. As used herein the terms "biomass" or "lignocellulosic material" includes materials containing cellulose and/or hemicellulose. Generally, these materials also contain xylan, lignin, protein, and carbohydrates, such as starch and sugar. Lignocellulose is generally found, for example, in the stems, leaves, hulls, husks, and cobs of plants or leaves, branches, and wood of trees. The process of converting a complex carbohydrate (such as starch or cellulose) into fermentable sugars is also referred to herein as "saccharification." Fermentable sugars, as used herein, refers to simple sugars, such as glucose, xylose, arabinose, galactose, mannose, rhamnose, sucrose and fructose.

[0081] Biomass can include virgin biomass and/or non-virgin biomass such as agricultural biomass, commercial organics, construction and demolition debris, municipal solid waste, waste paper and yard waste. Common forms of biomass include trees, shrubs and grasses, wheat, wheat straw, sugar cane bagasse, corn, corn husks, corn kernel including fiber from kernels, products and by-products from milling of grains such as corn, wheat and barley (including wet milling and dry milling) as well as municipal solid waste, waste paper and yard waste. The biomass can also be, but is not limited to, herbaceous material, agricultural residues, forestry residues, municipal solid wastes, waste paper, and pulp and paper mill residues. "Agricultural biomass" includes branches, bushes, canes, corn and corn husks, energy crops, forests, fruits, flowers, grains, grasses, herbaceous crops, leaves, bark, needles, logs, roots, saplings, short rotation woody crops, shrubs, switch grasses, trees, vegetables, fruit peels, vines, sugar beet pulp, wheat midlings, oat hulls, and hard and soft woods (not including woods with deleterious materials). In addition, agricultural biomass includes organic waste materials generated from agricultural processes including farming and forestry activities, specifically including forestry wood waste. Agricultural biomass may be any of the aforestated singularly or in any combination or mixture thereof.

[0082] Biomass high in starch, sugar, or protein such as corn, grains, fruits and vegetables are usually consumed as food. Conversely, biomass high in cellulose, hemicellulose and lignin are not readily digestible and are primarily utilized for wood and paper products, animal feed, fuel, or are typically disposed. Generally, the substrate is of high lignocellulose content, including distillers' dried grains corn stover, corn cobs, rice straw, hay, sugarcane bagasse, and other agricultural biomass, switchgrass, forestry wastes, poplar wood chips, pine wood chips, sawdust, yard waste, and the like, including any combination of substrate.

[0083] In one embodiment, the lignocellulosic material is distillers' dried grains (DDG). DDG (also known as dried distiller's grain, or distiller's spent grain) is spent, dried grains recovered after alcohol fermentation. The lignocellulosic material can also be distiller's dried grain with soluble material recycled back (DDGS). While reference will be made herein to DDG for convenience and simplicity, it should be understood that both DDG and DDGS are contemplated as desired lignocellulosic materials. These are largely considered to be waste products and can be obtained after the fermentation of the starch derived from any of a number of grains, including corn, wheat, barley, oats, rice and rye. In one embodiment the DDG is derived from corn.

[0084] It should be noted that the distiller's grains do not necessarily have to be dried. Although the grains normally, currently dried, water and enzymes are added to the DDG substrate in the present invention. If the saccharification were

done on site, the drying step could be eliminated and enzymes could be added to the distiller's grains without drying.

[0085] Due in part to the many components that comprise biomass and lignocellulosic materials, enzymes or a mixture of enzymes capable of degrading xylan, lignin, protein, and carbohydrates are needed to achieve saccharification. The present invention and disclosure includes enzymes or compositions thereof with, for example, cellobiohydrolase, endoglucanase, xylanase, β -glucosidase, hemicellulase and chitinase activities.

[0086] The enzymes of the present invention and disclosure may also be used for stone washing cellulosic fabrics such as cotton (e.g., denim), linen, hemp, ramie, cupro, lyocell, newcell, rayon and the like. See, for example, U.S. Patent No. 6,015,707. The enzymes and compositions of the present invention and disclosure are suitable for industrial textile applications in addition to the stone washing process. For example, cellulases are used in detergent compositions, either for the purpose of enhancing the cleaning ability of the composition or as a softening agent. When so used, the cellulase will degrade a portion of the cellulosic material, e.g., cotton fabric, in the wash, which facilitates the cleaning and/or softening of the fabric. The endoglucanase components of fungal cellulases have also been used for the purposes of enhancing the cleaning ability of detergent compositions, for use as a softening agent, and for use in improving the feel of cotton fabrics, and the like. Enzymes and compositions of the present invention and disclosure may also be used in the treatment of paper pulp (e.g., for improving the drainage or for de-inking of recycled paper) or for the treatment of wastewater streams (e.g., to hydrolyze waste material containing cellulose, hemicellulose and pectins to soluble lower molecular weight polymers).

[0087] In one aspect, the present invention includes a protein isolated from, or derived from the knowledge of enzymes from, a fungus such as *C. luceknowense* or a mutant or other derivative thereof, and more particularly, from the fungal strain denoted herein as C1 (Accession No. VKM F-3500-D). Preferably, the protein of the invention possess enzymatic activity. As described in U.S. Patent No. 6,015,707 or U.S. Patent No. 6,573,086, a strain called C1 (Accession No. VKM F-3500-D), was isolated from samples of forest alkaline soil from Sola Lake, Far East of the Russian Federation. This strain was deposited at the All-Russian Collection of Microorganisms of Russian Academy of Sciences (VKM), Bakhurhina St. 8, Moscow, Russia, 113184, under the terms of the Budapest Treaty on the International Regulation of the Deposit of Microorganisms for the Purposes of Patent Procedure on August 29, 1996, as *Chrysosporium luceknowense* Garg 27K, VKM-F 3500 D. Various mutant strains of *C. luceknowense* C1 have been produced and these strains have also been deposited at the All-Russian Collection of Microorganisms of Russian Academy of Sciences (VKM), Bakhurhina St. 8, Moscow, Russia, 113184, under the terms of the Budapest Treaty on the International Regulation of the Deposit of Microorganisms for the Purposes of Patent Procedure on September 2, 1998. For example, Strain C1 was mutagenised by subjecting it to ultraviolet light to generate strain UV13-6 (Accession No. VKM F-3632 D). This strain was subsequently further mutated with N-methyl-N'-nitro-N-nitrosoguanidine to generate strain NG7C-19 (Accession No. VKM F-3633 D). This latter strain in turn was subjected to mutation by ultraviolet light, resulting in strain UV18-25 (VKM F-3631 D). Strain C1 was classified as a *Chrysosporium luceknowense* based on morphological and growth characteristics of the microorganism, as discussed in detail in U.S. Patent No. 6,015,707 and U.S. Patent No. 6,573,086.

[0088] In certain embodiments of the present invention, a protein of the invention consists of the amino acid sequence SEQ ID NO: 32.

[0089] In general, the proteins disclosed herein possess carbohydrase enzymatic activity, or the ability to degrade carbohydrate-containing materials. More specifically, the proteins may possess cellulase activity such as endoglucanase activity (e.g., 1,4- β -D-glucan-4-glucanohydrolases), exoglucanase activity (e.g., 1,4- β -D-glucan cellobiohydrolases), and β -glucosidase activity. The proteins may possess hemicellulase activity such as endoxylanase activity, exoxylanase activity, or β -xylosidase activity. The proteins may possess laminarinase, xyloglucanase, galactanase, glucoamylase, pectate lyase, chitinase, exo- β -D-glucosaminidase, cellobiose dehydrogenase, acetyl xylan esterase, ligninase, amylase, glucuronidase, ferulic acid esterase, arabinofuranosidase, pectin methyl esterase, arabinase, lipase, glucosidase or glucomannanase activities. Physical properties, biochemical characteristics and substrate specificities of proteins of the present invention are illustrated below.

[0090] As used herein, "carbohydrase" refers to any protein that catalyzes the hydrolysis of carbohydrates, "Glycoside hydrolase" or "glycosidase" refers to a protein that catalyzes the hydrolysis of the glycosidic bonds between carbohydrates or between a carbohydrate and a non-carbohydrate residue. Endoglucanases, cellobiohydrolases, β -glucosidases, α -glucosidases, xylanases, β -xylosidases, galactanases, α -galactosidases, β -galactosidases, α -amylases, glucoamylases, endo-arabinases, arabinofuranosidases, mannanases, β -mannosidases, pectinases, acetyl xylan esterases, acetyl mannan esterases, ferulic acid esterases, coumaric acid esterases, pectin methyl esterases, and chitinases are examples of glycosidases.

[0091] "Cellulase" refers to a protein that catalyzes the hydrolysis of 1,4- β -D-glycosidic linkages in cellulose (such as bacterial cellulose, cotton, filter paper, phosphoric acid swollen cellulose, Avicel); cellulose derivatives (such as carboxymethylcellulose and hydroxyethylcellulose); plant lignocellulosic materials, beta-D-glucans or xyloglucans. Cellulose is a linear beta-(1-4) glucan consisting of anhydrocellobiose units. Endoglucanases, cellobiohydrolases, and β -glucosidases are examples of cellulases. "Endoglucanase" refers to a protein that catalyzes the hydrolysis of cellulose to oligosaccharide chains at random locations by means of an endoglucanase activity. "Cellobiohydrolase" refers to a

protein that catalyzes the hydrolysis of cellulose to cellobiose via an exoglucanase activity, sequentially releasing molecules of cellobiose from the reducing or non-reducing ends of cellulose or cellooligosaccharides. " β -glucosidase" refers to an enzyme that catalyzes the conversion of cellobiose and oligosaccharides to glucose.

[0092] "Hemicellulase" refers to a protein that catalyzes the hydrolysis of hemicellulose, such as that found in lignocellulosic materials. Hemicellulose is a complex polymer, and its composition often varies widely from organism to organism, and from one tissue type to another. Hemicelluloses include a variety of compounds, such as xylans, arabinoxylans, xyloglucans, mannans, glucomannans, and galactomannans. Hemicellulose can also contain glucan, which is a general term for beta-linked glucose residues. In general, a main component of hemicellulose is beta-1,4-linked xylose, a five carbon sugar. However, this xylose is often branched as beta-1,3 linkages or beta-1,2 linkages, and can be substituted with linkages to arabinose, galactose, mannose, glucuronic acid, or by esterification to acetic acid. The composition, nature of substitution, and degree of branching of hemicellulose is very different in dicotyledonous plants (dicots, *i.e.*, plant whose seeds have two cotyledons or seed leaves such as lima beans, peanuts, almonds, peas, kidney beans) as compared to monocotyledonous plants (monocots; *i.e.*, plants having a single cotyledon or seed leaf such as corn, wheat, rice, grasses, barley). In dicots, hemicellulose is comprised mainly of xyloglucans that are 1,4-beta-linked glucose chains with 1,6-alpha-linked xylosyl side chains. In monocots, including most grain crops, the principal components of hemicellulose are heteroxylans. These are primarily comprised of 1,4-beta-linked xylose backbone polymers with 1,2- or 1,3-beta linkages to arabinose, galactose and mannose as well as xylose modified by ester-linked acetic acids. Also present are branched beta glucans comprised of 1,3- and 1,4-beta-linked glucosyl chains. In monocots, cellulose, heteroxylans and beta glucans are present in roughly equal amounts, each comprising about 15-25% of the dry matter of cell walls. Hemicellulolytic enzymes, *i.e.* hemicellulases, include both endo-acting and exo-acting enzymes, such as xylanases, β -xylosidases, galactanases, α -galactosidases, β -galactosidases, endo-arabinases, arabinofuranosidases, mannanases, β -mannosidases. Hemicellulases also include the accessory enzymes, such as acetylerases, ferulic acid esterases, and coumaric acid esterases. Among these, xylanases and acetyl xylan esterases cleave the xylan and acetyl side chains of xylan and the remaining xylo-oligomers are unsubstituted and can thus be hydrolysed with β -xylosidase only. In addition, several less known side activities have been found in enzyme preparations which hydrolyze hemicellulose. Accordingly, xylanases, acetylerases and β -xylosidases are examples of hemicellulases.

[0093] "Xylanase" specifically refers to an enzyme that hydrolyzes the β -1,4 bond in the xylan backbone, producing short xylooligosaccharides.

[0094] β -Mannanases hydrolyze mannan-based hemicelluloses (mannan, glucomannan, galactomannan) and produce short β -1,4-mannooligosaccharides.

[0095] "Galactanase" or "arabinogalactan endo-1,4- β -galactosidase" refers to a protein that catalyzes the hydrolysis of endo-1,4- β -D-galactosidic linkages in arabinogalactans.

[0096] "Glucoamylase" refers to a protein that catalyzes the hydrolysis of terminal 1,4-linked α -D-glucose residues successively from non-reducing ends of the chains with the release of β -D-glucose.

[0097] "Chitinase" or "exo- β -D-chitinase" refers to a protein that catalyzes the hydrolysis of *N*-acetyl- β -D-glucosaminide 1,4- β -linkages in chitin and chitodextrins.

[0098] " α -L-arabinofuranosidase" or "arabinofuranosidase" refers to a protein that hydrolyzes arabinofuranosyl-containing hemicelluloses. Some of these enzymes remove arabinofuranoside residues from O-2 and/or O-3 single substituted xylose residues, as well as from O-2 and O-3 doubly substituted xylose residues

[0099] "Endo-arabinase" refers to a protein that catalyzes the hydrolysis of 1,5- α -arabinofuranosidic linkages in 1,5-arabinans.

[0100] " β -xylosidase" refers to a protein that hydrolyzes short 1,4- β -D-xylooligomers into xylose.

[0101] "Acetyl xylan esterase" refers to a protein that catalyzes the removal of the acetyl groups from xylose residues.

"Acetyl mannan esterase" refers to a protein that catalyzes the removal of the acetyl groups from mannose residues.

"Feruloyl esterase" or "ferulic acid esterase" refers to a protein that hydrolyzes the ester bond between the arabinose substituent group and ferulic acid. "Coumaric acid esterase" refers to a protein that hydrolyzes the ester bond between the arabinose substituent group and coumaric acid. Acetyl xylan esterases, ferulic acid esterases and pectin methyl esterases are examples of carbohydrate esterases.

[0102] "Pectate lyase" refers to a protein that catalyzes the cleavage of 1,4- α -D-galacturonan in oligosaccharide substrates.

[0103] Glycosidases (glycoside hydrolases; GH), a large family of enzymes that includes cellulases and hemicellulases, catalyze the hydrolysis of glycosidic linkages, predominantly in carbohydrates. Glycosidases such as the proteins of the present disclosure may be assigned to families on the basis of sequence similarities, and there are now over 100 different such families defined (see the CAZy (Carbohydrate Active EnZymes database) website, maintained by the Architecture of Fonction de Macromolécules Biologiques of the Centre National de la Recherche Scientifique, which describes the families of structurally-related catalytic and carbohydrate-binding modules (or functional domains) of enzymes that degrade, modify, or create glycosidic bonds; Coutinho, P.M. & Henrissat, B. (1999) Carbohydrate-active enzymes: an integrated database approach. In "Recent Advances in Carbohydrate Bioengineering", H.J. Gilbert, G. Davies, B. Hen-

rissat and B. Svensson eds., The Royal Society of Chemistry, Cambridge, pp. 3-12). Because there is a direct relationship between the amino acid sequence of a protein and its folding similarities, such a classification reflects the structural features of these enzymes and their substrate specificity. Such a classification system can help to reveal the evolutionary relationships between these enzymes and provide a convenient tool to determine information such as an enzyme's activity and function. Thus, enzymes assigned to a particular family based on sequence homology with other members of the family are expected to have similar enzymatic activities and related substrate specificities. CAZy family classifications also exist for glycosyltransferases (GT), polysaccharide lyases (PL), and carbohydrate esterases (CE). Likewise, sequence homology may be used to identify particular domains within proteins, such as cellulose binding modules (CBMs; also known as cellulose binding domains (CBDs)). The CAZy homologies of proteins of the present invention are disclosed below.

[0104] Proteins of the present disclosure may also include homologues and fragments of the proteins disclosed herein. The protein fragments include, but are not limited to, fragments comprising a catalytic domain (CD) and/or a cellulose-binding domain (also known as a cellulose binding module (CBM)); both are referred to herein as CBD. The identity and location of domains within proteins of the present disclosure are disclosed in detail below. The present disclosure encompasses all combinations of the disclosed domains. For example, a protein fragment may comprise a CD of a protein but not a CBD of the protein or a CBD of a protein but not a CD. Similarly, domains from different proteins may be combined. Protein fragments comprising a CD, CBD or combinations thereof for each protein disclosed herein can be readily produced using standard techniques known in the art. In some embodiments, a protein fragment comprises a domain of a protein that has at least one biological activity of the full-length protein. Homologues of proteins of the invention and disclosure that have at least one biological activity of the full-length protein are described in detail below. As used herein, the phrase "biological activity" of a protein refers to any function(s) exhibited or performed by the protein that is ascribed to the naturally occurring form of the protein as measured or observed *in vitro* or *in vivo*. In certain embodiments, a protein fragment comprises a domain of a protein that has the catalytic activity of the full-length enzyme. Specific biological activities of the proteins of the disclosure and invention, and structures within the proteins that are responsible for the activities, are described below.

Cellobiohydrolases

[0105] Certain proteins disclosed herein possess cellobiohydrolase enzymatic activity. For example, the polypeptides denoted herein as CBH Ia (SEQ ID NO:2), CBH Ib (SEQ ID NO:5), CBH IIa (SEQ ID NO:8), and CBH IIb (SEQ ID NO:11) possess cellobiohydrolase activity. These enzymes participate in the hydrolytic conversion of insoluble cellulose to cellooligosaccharides, with cellobiose being a primary product of the enzymatic catalysis.

[0106] The enzyme denoted CBH Ia is encoded by the genomic nucleic acid sequence represented herein as SEQ ID NO:1 and the cDNA sequence represented herein as SEQ ID NO:3. The CBH Ia nucleic acid sequence encodes a 526 amino acid sequence, represented herein as SEQ ID NO:2. The signal peptide for CBH Ia is located from positions 1 to about position 17 of SEQ ID NO:2, with the mature protein spanning from about position 18 to position 526 of SEQ ID NO:2. Within CBH Ia are two domains: a catalytic domain (CD) and a cellulose-binding domain (CBD). The amino acid sequence containing the CD of CBH Ia spans from a starting point of about position 19 of SEQ ID NO:2 to an ending point of about position 453 of SEQ ID NO:2. The amino acid sequence containing the CBD of CBH Ia spans from a starting point of about position 490 of SEQ ID NO:2 to an ending point of about position 519 of SEQ ID NO:2. Based on homology, CBH Ia can be assigned to CAZy Families GH 7 and CBM 1. As evidenced below, CBH Ia exhibits cellobiohydrolase activity.

[0107] The enzyme denoted CBH Ib is encoded by the nucleic acid sequence represented herein as SEQ ID NO:4 and the cDNA sequence represented herein as SEQ ID NO:6. The CBH Ib nucleic acid sequence encodes a 450 amino acid sequence, represented herein as SEQ ID NO:5. The signal peptide for CBH Ia is located from positions 1 to about position 20 of SEQ ID NO:5, with the mature protein spanning from about position 21 to position 450 of SEQ ID NO:5. Within CBH Ia is a catalytic domain (CD). The amino acid sequence containing the CD of CBH Ib spans from a starting point of about position 22 of SEQ ID NO:5 to an ending point of about position 450 of SEQ ID NO:5. Based on homology, CBH Ib can be assigned to CAZy Family GH 7. As evidenced below, CBH Ib exhibits cellobiohydrolase activity.

[0108] The enzyme denoted CBH IIa is encoded by the nucleic acid sequence represented herein as SEQ ID NO:7 and the cDNA sequence represented herein as SEQ ID NO:9. The CBH IIa nucleic acid sequences encodes a 395 amino acid sequence, represented herein as SEQ ID NO:8. The signal peptide for CBH IIa is located from positions 1 to about position 17 of SEQ ID NO:8, with the mature protein spanning from about position 18 to position 395 of SEQ ID NO:8. Within CBH IIa is a catalytic domain (CD). The amino acid sequence containing the CD of CBH IIa spans from a starting point of about position 40 of SEQ ID NO:8 to an ending point of about position 395 of SEQ ID NO:8. Based on homology, CBH IIa can be assigned to CAZy Family GH 6. As evidenced below, CBH IIa exhibits cellobiohydrolase activity.

[0109] The enzyme denoted CBH IIb is encoded by the nucleic acid sequence represented herein as SEQ ID NO:10

and the cDNA sequence represented herein as SEQ ID NO:12. The CBH IIb nucleic acid sequence encodes a 482 amino acid sequence, represented herein as SEQ ID NO: 11. The signal peptide for CBH IIb is located from positions 1 to about position 17 of SEQ ID NO: 11, with the mature protein spanning from about position 18 to position 482 of SEQ ID NO: 11. Within CBH IIb are two domains: a catalytic domain (CD) and a cellulose-binding domain (CBD). The amino acid sequence containing the CBD of CBH IIb spans from a starting point of about position 27 of SEQ ID NO:11 to an ending point of about position 62 of SEQ ID NO: 11. The amino acid sequence containing the CD of CBH IIb spans from a starting point of about position 125 of SEQ ID NO:11 to an ending point of about position 482 of SEQ ID NO: 11. Based on homology, CBH IIb can be assigned to CAZy Families GH 6 and CBM 1. As evidenced below, CBH IIb exhibits cellobiohydrolase activity.

[0110] Physical properties, biochemical characteristics and substrate specificities of cellobiohydrolases disclosed herein are illustrated in Tables 1 and 2 below. Physical and biochemical characteristics include molecular weight, isoelectric point, pH and temperature optima of enzymatic activity, and the stability of the enzymes after heating to 50°C for five hours at pH 5.0 or 7.0. Substrate specificities demonstrating enzyme activity for the indicated substrates were determined using standard assays known in the art, such as those disclosed in the Examples below. Figure 1 illustrates the specific activity of cellobiohydrolases of the present disclosure toward substrates selected from Table 2.

Table 1. Physical and Biochemical Properties of C1 Cellobiohydrolases

Enzyme	SEQ ID NO	MW (kDa)	pI	pH (50%)	T °C (50%)	Stability pH 5.0	Stability pH 7.0
CBH Ia	2	65	4.5	5.0 (3.5-6.5)	60 (45->80)	99%	96%
CBH Ia (CD)	2 (CD Only)	52	4.5	5.0 (3.5-6.5)	60 (45->80)	97%	95%
CBH Ib	5	60	3.8	4.7 (3.3-6.3)	65 (50->80)	80%	80%
CBH IIa	8	43	4.2	5.0 (3.5-6.5)	60 (50-72)	98%	70%
CBH IIb	11	70	5.6	5.0 (<3.0-6.5)	60 (50-75)	100%	80%

MW = Molecular Weight, kiloDaltons (kDa)
pI = isoelectric point
pH (50%) = pH optimum of enzyme activity (range of at least 50% activity)
T °C (50%) = Temperature optimum of enzyme activity (range of at least 50% activity)
Stability pH 5.0 = residual enzyme activity after 5 hour incubation at 50 °C, pH 5.0
Stability pH 7.0 = residual enzyme activity after 5 hour incubation at 50 °C, pH 7.0

Table 2. Substrate Specificities of C1 Cellobiohydrolases (Activity Towards Specific Substrates (units/mg))

Substrate	CBH Ia	CBH Ia (CD)	CBH Ib	CBH IIa	CBH IIb
CMC (S-N Assay)	0.2	0.2	0.3	1.1	0.1
CMC (Viscometric)	0.01	0.02	0.02	0.6	0.01
β-glucan	<0.1	<0.1	<0.1	2.0	0.2
Avicel	0.21	0.10	0.12	0.08	0.22
Birchwood Xylan	0	0	0	1.4	0.03
Xyloglucan	0	0	0	0	0
Laminarin	0	0	0	0	0
Arabinan	0	0	0	0.1	0
Galactan	0	0	0	0	0
Galactomannan	0	0	0	0	0
p-NPh-β-D-cellobioside	0.021	0.025	0.02	0	0
p-NPh-β-D-lactoside	0.12	0.11	0.09	0	0

[0111] pH and temperature dependencies of cellobiohydrolases disclosed herein are illustrated in Tables 3 and 4 below. pH and temperature dependencies were determined using the Avicel cellulase assay described in Example 5

below. pH dependencies were assayed by varying the pH at 40°C for one hour, while temperature dependencies were assayed by varying the temperature at pH 5.0 for one hour. The temperature/pH that exhibited the highest enzyme activity was set at 100% in the tables below, with the remaining values expressed a percentage of this standard. The pH and temperature dependencies of the cellobiohydrolases are also illustrated graphically in Figures 2 and 3, respectively.

Table 3. pH Dependency of C1 Cellobiohydrolases (% Activity)

pH	CBH Ia	CBH Ia (CD)	CBH Ib	CBH IIa	CBH IIb
3	23	35	22	21	72
3.55	50	50	48	50.0	87
3.9	58	61	85	68	89
4.3	75	80	95	85	91
4.7	95	95	100	99	95
5.08	100	100	92	100	100
5.6	98	98	75	98	98
6	77	78	58	78	94
6.6	48	49	46	51.6	50
7.1	17.4	20	20	42	30
7.5	2	8		23	10

Table 4. Temperature Dependency of C1 Cellobiohydrolases (% Activity)

Temp °C	CBH Ia	CBH Ia (CD)	CBH Ib	CBH IIa	CBH IIb
30	11.0				10
35	20	20		20.0	18
40	39	32		28	30
45	50	48		43	41
50	62	55	25	55	55.6
55	82	67	54	78	73
60	95	82	92	100	89
65	100	100	100	90	100
70	85	92	29	64	76
75	40	75	6	40	36.5
80	15	20	2	20	23

Endoglucanases

[0112] Some proteins disclosed herein possess endo-1,4- β -glucanase enzymatic activity. For example, the polypeptides EG I (SEQ ID NO: 14), EG II (SEQ ID NO:94), EG III (SEQ ID NO: 17), EG V (SEQ ID NO:20), and EG VI (SEQ ID NO:23) possess endo-1,4- β -glucanase activity. As with the cellobiohydrolases described above, these endoglucanases also participate in the hydrolytic conversion of insoluble cellulose to cellobiosaccharides, with cellobiose being a primary product of the enzymatic catalysis.

[0113] The enzyme denoted EG I is encoded by the nucleic acid sequence represented herein as SEQ ID NO:13 and the cDNA sequence represented herein as SEQ ID NO:15. The EG I nucleic acid sequence encodes a 456 amino acid sequence, represented herein as SEQ ID NO: 14. The signal peptide for EG I is located from positions 1 to about position 20 of SEQ ID NO: 14, with the mature protein spanning from about position 21 to position 456 of SEQ ID NO:14. Within

EG I is a catalytic domain (CD). The amino acid sequence containing the CD of EG I spans from a starting point of about position 22 of SEQ ID NO: 14 to an ending point of about position 420 of SEQ ID NO: 14. Based on homology, EG I can be assigned to CAZy Family GH 7. As evidenced below, EG I exhibits endoglucanase activity.

[0114] The enzyme denoted EG II is a 389 amino acid sequence, represented herein as SEQ ID NO:94. The signal peptide for EG II is located from positions 1 to about position 16 of SEQ ID NO:94, with the mature protein spanning from about position 17 to position 389 of SEQ ID NO:94. Within EG II are two domains: a catalytic domain (CD) and a cellulose-binding domain (CBD). The amino acid sequence containing the CBD of EG II spans from a starting point of about position 17 of SEQ ID NO:94 to an ending point of about position 52 of SEQ ID NO:94. The amino acid sequence containing the CD of EG II spans from a starting point of about position 84 of SEQ ID NO:94 to an ending point of about position 388 of SEQ ID NO:94. As evidenced below, EG II exhibits endoglucanase activity.

[0115] The enzyme denoted EG III is encoded by the nucleic acid sequence represented herein as SEQ ID NO:16 and the cDNA sequence represented herein as SEQ ID NO:18. The EG III nucleic acid sequence encodes a 247 amino acid sequence, represented herein as SEQ ID NO:17. The signal peptide for EG III is located from positions 1 to about position 15 of SEQ ID NO: 17, with the mature protein spanning from about position 16 to position 247 of SEQ ID NO:17. Within EG III is a catalytic domain (CD). The amino acid sequence containing the CD of EG III spans from a starting point of about position 91 of SEQ ID NO: 17 to an ending point of about position 247 of SEQ ID NO: 17. Based on homology, EG III can be assigned to CAZy Family GH 12. As evidenced below, EG III exhibits endoglucanase activity.

[0116] The enzyme denoted EG V is encoded by the nucleic acid sequence represented herein as SEQ ID NO:19 and the cDNA sequence represented herein as SEQ ID NO:21. The EG V nucleic acid sequence encodes a 225 amino acid sequence, represented herein as SEQ ID NO:20. The signal peptide for EG V is located from positions 1 to about position 18 of SEQ ID NO:20, with the mature protein spanning from about position 19 to position 225 of SEQ ID NO:20. Within EG V is a catalytic domain (CD). The amino acid sequence containing the CD of EG V spans from a starting point of about position 23 of SEQ ID NO:20 to an ending point of about position 224 of SEQ ID NO:20. Based on homology, EG V can be assigned to CAZy Family GH 45. As evidenced below, EG V exhibits endoglucanase activity.

[0117] The enzyme denoted EG VI is encoded by the nucleic acid sequence represented herein as SEQ ID NO:22 and the cDNA sequence represented herein as SEQ ID NO:24. The EG VI nucleic acid sequence encodes a 381 amino acid sequence, represented herein as SEQ ID NO:23. The signal peptide for EG VI is located from positions 1 to about position 18 of SEQ ID NO:23, with the mature protein spanning from about position 19 to position 381 of SEQ ID NO:23. Within EG VI is a catalytic domain (CD). The amino acid sequence containing the CD of EG VI spans from a starting point of about position 33 of SEQ ID NO:23 to an ending point of about position 378 of SEQ ID NO:23. Based on homology, EG VI can be assigned to CAZy Family GH 6. As evidenced below, EG VI exhibits endoglucanase activity.

[0118] Physical properties, biochemical characteristics and substrate specificities of endoglucanases of the present disclosure are illustrated in Tables 5 and 6 below. Physical and biochemical characteristics include molecular weight, isoelectric point, pH and temperature optima of enzymatic activity, and the stability of the enzymes after heating to 50°C for five hours at pH 5.0 or 7.0. Substrate specificities demonstrating enzyme activity for the indicated substrates were determined using standard assays known in the art, such as those disclosed in the Examples below. Figure 1 illustrates the specific activity of endoglucanases of the present invention toward substrates selected from Table 6.

Table 5, Physical and Biochemical Properties of C1 Endoglucanases

Enzyme	SEQ ID NO	MW (kDa)	pI	pH (50%)	T °C (50%)	Stability pH 5.0	Stability pH 7.0
EG I	14	60	3.7	4.8 (3.5-7.5)	60 (43-76)	80%	80%
EG II	97	51	4.8	4.7 (3.5-7.5)	70 (53-77)	93%	93%
EG II (CD)	97 (CD Only)	44	6.0	5.2 (3.5-7.5)	70 (53-77)	90%	90%
EG III	17	28	5.7	5.3 (3.8-7.0)	60 (30-75)	92%	74%
EG V	20	25	4.0	5.5 (3.5-8.5)	65 (50-75)	95%	92%
EG VI	23	47	5.7	6.0 (4.5->8.5)	70 (40-82)	60%	75%
MW = Molecular Weight, kiloDaltons (kDa) pI = isoelectric point pH (50%) = pH optimum of enzyme activity (range of at least 50% activity) T °C (50%) = Temperature optimum of enzyme activity (range of at least 50% activity) Stability pH 5.0 = residual enzyme activity after 5 hour incubation at 50 °C, pH 5.0 Stability pH 7.0 = residual enzyme activity after 5 hour incubation at 50 °C, pH 7.0							

Table 6. Substrate Specificities of C1 Endoglucanases (Activity Towards Specific Substrates (units/mg))

Substrate	EG I	EG II	EG II (CD)	EG III	EG V	EG VI
CMC (S-N Assay)	12	52	59	11	17	14
CMC (Viscometric)	5	46	39	19	20	6
β -glucan	18	75	74	125	5.3	18
Avicel	0.08	0.19	0.06	0.02	0.06	0.07
Birchwood Xylan	0	0.18	0.07	0.2	0	0.08
Xyloglucan	0	0	0	15.6	1.9	0
Laminarin	0	0.1	0.1	0	0	0
Arabinan	0	0	0.3	0	0	0
Galactan	0	0	0.6	0	0	0.1
Galactomannan	0	1.1	1	0	0.1	0
p-NPh- β -D-cellobioside	0.12	0	0	0	0	0
p-NPh- β -D-lactoside	0.27	0	0	0	0	0

[0119] pH and temperature dependencies of endoglucanases are illustrated in Tables 7 and 8 below. pH and temperature dependencies were determined using the CMC cellulase assay described in Example 3 below. pH dependencies were assayed by varying the pH at 50°C for five minutes, while temperature dependencies were assayed by varying the temperature at pH 5.0 for five minutes. The temperature/pH that exhibited the highest enzyme activity was set at 100% in the tables below, with the remaining values expressed a percentage of this standard. The pH and temperature dependencies of the endoglucanases are also illustrated graphically in Figures 4 and 5, respectively.

Table 7. pH Dependency of C1 Endoglucanases (% Activity)

pH	EG I	EG II	EG II (CD)	EG III	EG V	EG VI
2.7	9	23	16	0	16.5	8
3.34	30	39	29	10	48	17
4.04	88	87	77	60	81	37
4.777	100	100	95	98	94.5	75
5.36	98	96	100	100	100	97
6.06	87	86	96	89	98	100
6.73	74	75	81	70	88	96
7.38	55	58	64	38	76.5	91
8	33	43	38	15	67	75
8.44	23	32	25	8	52	68

Table 8. Temperature Dependency of C1 Endoglucanases (% Activity)

Temp °C	EG I	EG II	EG II (CD)	EG III	EG V	EG VI
30	22.5	19	10.5	55.2	22	24.7
35	30.5	20	12	69.1	26	37.0
40	42.3	24	20.7	80.6	37	52
45	61.1	27	25	86.3	45	68.5

(continued)

Temp °C	EG I	EG II	EG II (CD)	EG III	EG V	EG VI
50	83.3	35.4	32	93.2	51	79.8
55	95.5	55	56	97.5	70	89.0
60	100	72	71	100	83.3	96.1
65	93.5	90	92	95.9	100	99
70	77.6	100	100	87.9	88	100
75	57.3	62	70	57.5	60	82.2
80	32.4	31	35	13.4	33	57.5

Xylanuses

[0120] Certain proteins of the present disclosure and the protein of the present invention possess xylanase enzymatic activity. For example, the polypeptides Xyl 1 (SEQ ID NO:26), Xyl 2 (SEQ ID NO:29), Xyl 3 (SEQ ID NO:32), Xyl 4 (SEQ ID NO:35), Xyl 5 (SEQ ID NO:38), and Xyl 6 (SEQ ID NO:41) possess xylanase activity. These xylanases participate in the hydrolytic conversion of hemicellulose into xylose.

[0121] The enzyme denoted Xyl 1 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:25 and the cDNA sequence represented herein as SEQ ID NO:27. The Xyl 1 nucleic acid sequence encodes a 384 amino acid sequence, represented herein as SEQ ID NO:26. The signal peptide for Xyl 1 is located from positions 1 to about position 17 of SEQ ID NO:26, with the mature protein spanning from about position 18 to position 384 of SEQ ID NO:26. Within Xyl 1 are two domains: a catalytic domain (CD) and a cellulose-binding domain (CBD). The amino acid sequence containing the CBD of Xyl 1 spans from a starting point of about position 18 of SEQ ID NO:26 to an ending point of about position 53 of SEQ ID NO:26. The amino acid sequence containing the CD of Xyl 1 spans from a starting point of about position 83 of SEQ ID NO:26 to an ending point of about position 384 of SEQ ID NO:26. Based on homology, Xyl 1 can be assigned to CAZy Families GH 10 and CBM 1. As evidenced below, Xyl 1 exhibits xylanase activity.

[0122] The enzyme denoted Xyl 2 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:28 and the cDNA sequence represented herein as SEQ ID NO:30. The Xyl 2 nucleic acid sequence encodes a 221 amino acid sequence, represented herein as SEQ ID NO:29. The signal peptide for Xyl 2 is located from positions 1 to about position 18 of SEQ ID NO:29, with the mature protein spanning from about position 19 to position 221 of SEQ ID NO:29. Within Xyl 2 is a catalytic domain (CD). The amino acid sequence containing the CD of Xyl 2 spans from a starting point of about position 25 of SEQ ID NO:29 to an ending point of about position 219 of SEQ ID NO:29. Based on homology, Xyl 2 can be assigned to CAZy Family GH 11. As evidenced below, Xyl 2 exhibits xylanase activity.

[0123] The enzyme denoted Xyl 3 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:31 and the cDNA sequence represented herein as SEQ ID NO:33. The Xyl 3 nucleic acid sequence encodes a 413 amino acid sequence, represented herein as SEQ ID NO:32. The signal peptide for Xyl 3 is located from positions 1 to about position 17 of SEQ ID NO:32, with the mature protein spanning from about position 18 to position 413 of SEQ ID NO:32. Within Xyl 3 are two domains: a catalytic domain (CD) and a cellulose-binding domain (CBD). The amino acid sequence containing the CD of Xyl 3 spans from a starting point of about position 18 of SEQ ID NO:32 to an ending point of about position 336 of SEQ ID NO:32. The amino acid sequence containing the CBD of Xyl 3 spans from a starting point of about position 377 of SEQ ID NO:32 to an ending point of about position 413 of SEQ ID NO:32. Based on homology, Xyl 3 can be assigned to CAZy Family GH 10. As evidenced below, Xyl 3 exhibits xylanase activity. The enzyme denoted Xyl 4 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:34 and the cDNA sequence represented herein as SEQ ID NO:36. The Xyl 4 nucleic acid sequence encodes a 375 amino acid sequence, represented herein as SEQ ID NO:35. The signal peptide for Xyl 4 is located from positions 1 to about position 19 of SEQ ID NO:35, with the mature protein spanning from about position 20 to position 375 of SEQ ID NO:35. Within Xyl 4 is a catalytic domain (CD). The amino acid sequence containing the CD of Xyl 4 spans from a starting point of about position 31 of SEQ ID NO:35 to an ending point of about position 354 of SEQ ID NO:35. Based on homology, Xyl 4 can be assigned to CAZy Family GH 10. As evidenced below, Xyl 4 exhibits xylanase activity.

[0124] The enzyme denoted Xyl 5 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:37 and the cDNA sequence represented herein as SEQ ID NO:39. The Xyl 5 nucleic acid sequence encodes a 226 amino acid sequence, represented herein as SEQ ID NO:38. The signal peptide for Xyl 5 is located from positions 1 to about position 21 of SEQ ID NO:38, with the mature protein spanning from about position 22 to position 226 of SEQ ID NO:38. Within Xyl 5 is a catalytic domain (CD). The amino acid sequence containing the CD of Xyl 5 spans from a starting point of

about position 42 of SEQ ID NO:38 to an ending point of about position 223 of SEQ ID NO:38. Based on homology, Xyl 5 can be assigned to CAZy Families GH 11 and CE 4. As evidenced below, Xyl 5 exhibits xylanase activity.

[0125] The enzyme denoted Xyl 6 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:40 and the cDNA sequence represented herein as SEQ ID NO:42. The Xyl 6 nucleic acid sequence encodes a 228 amino acid sequence, represented herein as SEQ ID NO:41. The signal peptide for Xyl 6 is located from positions 1 to about position 20 of SEQ ID NO:41, with the mature protein spanning from about position 21 to position 228 of SEQ ID NO:41. Within Xyl 6 is a catalytic domain (CD). The amino acid sequence containing the CD of Xyl 6 spans from a starting point of about position 37 of SEQ ID NO:41 to an ending point of about position 226 of SEQ ID NO:41. Based on homology, Xyl 6 can be assigned to CAZy Family GH 11. As evidenced below, Xyl 6 exhibits xylanase activity.

[0126] Physical properties, biochemical characteristics and substrate specificities of xylanases of the present disclosure and the xylanase of the present invention are illustrated in Tables 9, 10, 11 and 12 below. Physical and biochemical characteristics include molecular weight, isoelectric point, pH and temperature optima of enzymatic activity, and the stability of the enzymes after heating to 50°C for five hours at pH 5.0 or 7.0, or after heating to 80°C for twenty seconds. Substrate specificities demonstrating enzyme activity for the indicated substrates (specific and non-specific) were determined using standard assays known in the art, such as those disclosed in the Examples below. The degree of xylan exhaustive hydrolysis of specific substrates is also indicated. Figure 6 illustrates the specific activity of xylanases of the present disclosure and the xylanase of the present invention toward substrates selected from Table 10.

Table 9. Physical and Biochemical Properties of C1 Xylanases

Enzyme	SEQ ID NO	MW (kDa)	pI	pH	T °C	Stability pH 5.0	Stability pH 7.0	Stability 80 °C
Xyl 1	26	42	7.8	5.5-7.0	65-70	100% (81%)	100% (85%)	12%
Xyl 1 (CD)	26 (CD Only)	31	8.9	5.5-7.0	65-70	100% (76%)	100% (88%)	19%
Xyl 2	29	24	7.9	5.7-7.5	70	84% (35%)	71% (25%)	23%
Xyl 3	32	57	4.4	5.5-6.5	80-85	100% (90%)	100% (92%)	75%
Xyl 3 (CD)	32(CD Only)	46	4.3	5.5-6.5	80-85	100% (95%)	100% (97%)	98%
Xyl 4	35	38	4.8	5.0	80	100% (82%)	100% (70%)	82%
Xyl 5	38	22	6.7	4.5	65	45% (35%)	Unstable at pH ≥ 7.0	68%
Xyl 6	41	23	8.4	6.0	65-70	90% (12%)	95% (20%)	60%

MW = Molecular Weight, kiloDaltons (kDa)
 pI = isoelectric point
 pH = pH optimum of enzyme activity
 T °C = Temperature optimum of enzyme activity
 Stability pH 5.0 = residual enzyme activity after 1 hour incubation at 50 °C (60 °C), pH 5.0
 Stability pH 7.0 = residual enzyme activity after 1 hour incubation at 50 °C (60 °C), pH 7.0
 Stability 80 °C = residual enzyme activity after 20 seconds at 80 °C

Table 10. Substrate Specificities of C1 Xylanases (Activity Towards Specific Substrates (units/mg))

Substrate	Xyl 1	Xyl 1 (CD)	Xyl 2	Xyl 3	Xyl 3 (CD)	Xyl 4	Xyl 5	Xyl 6
Birchwood Xylan	65	83	395	39	85	32	300	169
Beechwood Xylan	66	71	329	47	88	36	300	188
Wheat Arabinoxylan	83	102	494	52	98	40	143	268
Wheat Arabinoxylan (Insoluble)	83	82	198	52	78	24	57	107
Arabinoxylan from Oat Spelts	96	100	485	63	107	42	320	329
Viscometric Activity, Wheat Arabinoxylan	1311	1382	2158	807	1210	234	2700	2452

Table 11. Degree of Xylan Exhaustive Hydrolysis of Specific Substrates by C1 Xylanases (%)

Substrate	Xyl 1	Xyl 1 (CD)	Xyl 2	Xyl 3	Xyl 3 (CD)	Xyl 4	Xyl 5	Xyl 6
Birchwood Xylan	26	24	19	24	23	25	22	21
Beechwood Xylan	24	22	15	22	20	23	18	17
Wheat Arabinoxylan	26	28	15	30	29	24	18	17
Wheat Arabinoxylan (Insoluble)	12	13	7	14	11	12	9	8
Arabinoxylan from Oat Spelts	28	27	19	29	25	26	20	19

Table 12. Substrate Specificities of C1 Xylanases (Activity Towards Non-Specific Substrates (units/mg))

Substrate	Xyl 1	Xyl 1 (CD)	Xyl 2	Xyl 3	Xyl 3 (CD)	Xyl 4	Xyl 5	Xyl 6
CMC	0.65	1	0	0.12	0.2	0	0	0
Avicel	0	0	0	0	0	0	0	0
p-NPh- β -D-xylopyranoside	0	0	0	0.3	0.6	0.3	0	0
p-NPh- β -D-cellobioside	0.36	0.42	0	0.84	1.6	4.0	0	0

[0127] pH and temperature dependencies of xylanases of the present disclosure and the xylanase of the present invention are illustrated in Tables 13 and 14 below. pH and temperature dependencies were determined using the xylanase assay (on soluble xylan) described in Example 7 below. pH dependencies were assayed by varying the pH at 40°C for 10 minutes, while temperature dependencies were assayed by varying the temperature at pH 5.0 for 10 minutes. The temperature/pH that exhibited the highest enzyme activity was set at 100% in the table below, with the remaining values expressed a percentage of this standard. The pH and temperature dependencies of the xylanases are also illustrated graphically in Figures 7 and 8, respectively.

Table 13. pH Dependency of C1 Xylanases (% Activity)

pH	Xyl 1	Xyl 1 (CD)	Xyl 3	Xyl 3 (CD)	pH	Xyl 4	pH	Oxy 2	pH	Xyl 6	pH	Xyl 5
2.5	0.00	0.46	1.83	0.42	3.0	0.37	3.1	0	2.35	0.00	2.44	0.00
3.53	22.00	25.48	39.99	25.32	3.5	36.33	3.8	5.89	3.53	14.25	3	13.65
4.67	70.23	73.56	74.68	71.26	4.0	78.51	4.2	26.66	4.42	45.85	3.5	36.97
4.92	85.63	83.69	84.23	86.62	4.5	94.89	4.8	70.06	4.92	75.55	4	80.39
5.57	100	97.00	97.22	95.00	5.0	100	5.3	90.00	5.57	94.76	4.5	100
6.03	99.59	100	98.33	97.53	5.5	89.22	5.8	97.26	6.03	100	5.1	70.32
6.64	97.33	95.34	100	100	6.0	73.65	6.4	100	6.64	96.92	5.5	45.14
7.1	95.46	90.08	90.43	91.99	6.5	52.16	6.9	99.05	7.1	78.61	5.81	29.11
7.61	90.76	85.16	79.49	79.90	7.0	40.83	7.9	89.62	7.61	60.77	6.75	0.00
8.71	72.12	62.86	61.10	63.44	8.0	19.64	9.5	4.40	8.71	21.99	7.2	0.00

Table 14. Temperature Dependency of C1 Xylanases (% Activity)

Temp °C	Xyl 1	Xyl 1 (CD)	Xyl 2	Xyl 3	Xyl 3 (CD)	Xyl 4	Xyl 5	Xyl 6
30	37.5	37.5	8.3	27.3	26.5	18.6	44.8	35.4
40	46.8	46.8	22.7	34.1	31.7	27.4	56.3	57.0
45	63.1	63.1		40.0	30.0	36.4		62.6

(continued)

Temp °C	Xyl 1	Xyl 1 (CD)	Xyl 2	Xyl 3	Xyl 3 (CD)	Xyl 4	Xyl 5	Xyl 6
50	73.1	73.1	47.2	46.4	36.5	40.3	75.9	76.1
55	82.9	82.9		59.5	63.8	50.5		85.0
60	88.1	88.1	79.0	63.3	67.5	70.3	98.3	87.5
65	98.0	98.0	92.5	76.1	80.4	81.8	100.0	97.2
70	100.0	100.0	100.0	81.5	87.5	86.4	92.2	97.3
75	95.0	95.0	95.2	92.0	88.8	92.6	70.1	71.6
80	63.7	63.7	56.4	100.0	98.3	100.0	30.2	46.2
85	41.0	41.0		100.0	100.0	85.3		31.1
90	21.1	21.1		71.9	69.8	65.6		20.8

Additional Glycosidases

[0128] Certain proteins of the present disclosure possess glycosidase enzymatic activity, including β -glucosidase, galactanase, glucoamylase, endo-arabinase, and β -xylosidase activities. The polypeptides denoted herein as Bgl3A (SEQ ID NO:44), Gal53A (SEQ ID NO:47), Gla15A (SEQ ID NO:50), Abn1 (SEQ ID NO:89) and bx12 (SEQ ID NO:92) possess glycosidase activity, as indicated below.

[0129] The enzyme Bgl3A is encoded by the nucleic acid sequence represented herein as SEQ ID NO:43 and the cDNA sequence represented herein as SEQ ID NO:45. The Bgl3A nucleic acid sequence encodes a 871 amino acid sequence, represented herein as SEQ ID NO:44. The signal peptide for Bgl3A is located from positions 1 to about position 19 of SEQ ID NO:44, with the mature protein spanning from about position 20 to position 871 of SEQ ID NO:44. Within Bgl3A is a catalytic domain (CD). The amino acid sequence containing the CD of Bgl3A spans from a starting point of about position 87 of SEQ ID NO:44 to an ending point of about position 643 of SEQ ID NO:44. Based on homology, Bgl3A can be assigned to CAZy Family GH 3.

[0130] The enzyme Gal53A is encoded by the nucleic acid sequence represented herein as SEQ ID NO:46 and the cDNA sequence represented herein as SEQ ID NO:48. The Gal53A nucleic acid sequence encodes a 350 amino acid sequence, represented herein as SEQ ID NO:47. The signal peptide for Gal53A is located from positions 1 to about position 18 of SEQ ID NO:47, with the mature protein spanning from about position 19 to position 350 of SEQ ID NO:47. Within Gal53A is a catalytic domain (CD). The amino acid sequence containing the CD of Gal53A spans from a starting point of about position 19 of SEQ ID NO:47 to an ending point of about position 334 of SEQ ID NO:47. Based on homology, Gal53A can be assigned to CAZy Family GH 53. As evidenced below, Gal53A exhibits galactanase (arabinogalactan endo-1,4- β -galactosidase) activity.

[0131] The enzyme Gla15A is encoded by the nucleic acid sequence represented herein as SEQ ID NO:49 and the cDNA sequence represented herein as SEQ ID NO:51. The Gla15A nucleic acid sequence encodes a 628 amino acid sequence, represented herein as SEQ ID NO:50. The signal peptide for Gla15A is located from positions 1 to about position 20 of SEQ ID NO:50, with the mature protein spanning from about position 21 to position 628 of SEQ ID NO:50. Within Gla15A are two domains: a catalytic domain (CD) and a starch-binding domain (CBM 20). The amino acid sequence containing the CD of Gla15A spans from a starting point of about position 28 of SEQ ID NO:50 to an ending point of about position 461 of SEQ ID NO:50. The amino acid sequence containing the starch-binding domain of Gla15A spans from a starting point of about position 518 of SEQ ID NO:50 to an ending point of about position 628 of SEQ ID NO:50. Based on homology, Gla15A can be assigned to CAZy Families GH 15 and CBM 20. As evidenced below, Gla15A exhibits glucoamylase activity.

[0132] The enzyme Abn1 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:88 and the cDNA sequence represented herein as SEQ ID NO:90. The Abn1 nucleic acid sequence encodes a 321 amino acid sequence, represented herein as SEQ ID NO:89. The signal peptide for Abn1 is located from positions 1 to about position 20 of SEQ ID NO:89, with the mature protein spanning from about position 21 to position 321 of SEQ ID NO:89. Within Abn1 is a catalytic domain (CD). The amino acid sequence containing the CD of Abn1 spans from a starting point of about position 27 of SEQ ID NO:89 to an ending point of about position 321 of SEQ ID NO:89. Based on homology, Abn1 can be assigned to CAZy Family GH 43. As illustrated below in Example 23, Abn1 exhibits endo-arabinase activity as measured by the ability to hydrolyze AZCL-arabinan (debranched). Abn1 also possesses significant homology (about 61% from amino acids 13 to 321 of Abn1) with an endo-arabinase from *Aspergillus niger* (Genbank Accession No.

AAA32682; see also Flippin et al., Appl. Microbiol. Biotechnol. 40:318 (1993)). Based on this degree of homology, Abn1 is expected to exhibit similar endo-arabinase enzymatic activity.

[0133] The enzyme Bx12 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:91 and the cDNA sequence represented herein as SEQ ID NO:93. The Bx12 nucleic acid sequence encodes a 733 amino acid sequence, represented herein as SEQ ID NO:92. The signal peptide for Bx12 is located from positions 1 to about position 17 of SEQ ID NO:92, with the mature protein spanning from about position 18 to position 733 of SEQ ID NO:92. Within Bx12 is a catalytic domain (CD). The amino acid sequence containing the CD of Bx12 spans from a starting point of about position 26 of SEQ ID NO:92 to an ending point of about position 602 of SEQ ID NO:92. Based on homology, Bx12 can be assigned to CAZy Family GH 3. As illustrated below in Example 24, Bx12 exhibits β -xylosidase activity. Bx12 also possesses significant homology (about 70% from amino acids 28 to 733 of Bx12) with a β -glucosidase from *Hypocrea jecorina* (Genbank Accession No. 1713235A; see also Bamett et al., Biotechnol. 9:562 (1991)). Based on this degree of homology, Bx12 is also expected to exhibit similar β -glucosidase enzymatic activity.

Pectate Lyases

[0134] Certain proteins of the present disclosure possess pectate lyase enzymatic activity. The polypeptide denoted herein as Pell (SEQ ID NO:53) possesses pectate lyase activity.

[0135] The enzyme Pell is encoded by the nucleic acid sequence represented herein as SEQ ID NO:52 and the cDNA sequence represented herein as SEQ ID NO:54. The Pell nucleic acid sequence encodes a 375 amino acid sequence, represented herein as SEQ ID NO:53. The signal peptide for Pell is located from positions 1 to about position 20 of SEQ ID NO:53, with the mature protein spanning from about position 21 to position 375 of SEQ ID NO:53. Within Pell is a catalytic domain (CD). The amino acid sequence containing the CD of Pell spans from a starting point of about position 21 of SEQ ID NO:53 to an ending point of about position 375 of SEQ ID NO:53. As evidenced below, Pell exhibits pectate lyase activity.

Chitinases

[0136] Certain proteins of the present disclosure possess chitinase enzymatic activity. The polypeptides denoted herein as Chi18A (SEQ ID NO:56) and Gls2A (SEQ ID NO:59) possess chitinase activity.

[0137] The enzyme Chi18A is encoded by the nucleic acid sequence represented herein as SEQ ID NO:55 and the cDNA sequence represented herein as SEQ ID NO:57. The Chi18A nucleic acid sequence encodes a 426 amino acid sequence, represented herein as SEQ ID NO:56. The signal peptide for Chi18A is located from positions 1 to about position 23 of SEQ ID NO:56, with the mature protein spanning from about position 24 to position 426 of SEQ ID NO:56. Within Chi18A is a catalytic domain (CD). The amino acid sequence containing the CD of Chi18A spans from a starting point of about position 42 of SEQ ID NO:56 to an ending point of about position 386 of SEQ ID NO:56. Based on homology, Chi18A can be assigned to CAZy Family GH 18. As evidenced below, Chi18A exhibits chitinase activity.

[0138] The enzyme Gls2A is encoded by the nucleic acid sequence represented herein as SEQ ID NO:58 and the cDNA sequence represented herein as SEQ ID NO:60. The Gls2A nucleic acid sequence encodes a 907 amino acid sequence, represented herein as SEQ ID NO:59. The signal peptide for Gls2A is located from positions 1 to about position 20 of SEQ ID NO:59, with the mature protein spanning from about position 21 to position 907 of SEQ ID NO:59. Within Gls2A is a catalytic domain (CD). The amino acid sequence containing the CD of Gls2A spans from a starting point of about position 210 of SEQ ID NO:59 to an ending point of about position 764 of SEQ ID NO:59. Based on homology, Gls2A can be assigned to CAZy Family GH 2. As evidenced below, Gls2A exhibits chitinase (exo- β -D-chitinase) activity.

Carbohydrate Esterases

[0139] Certain proteins of the present disclosure possess carbohydrate esterase enzymatic activity, such as acetyl xylan esterase, ferulic acid esterase and pectin methyl esterase activity. The polypeptides denoted herein as Axe3 (SEQ ID NO:62), FaeA1 (SEQ ID NO:65), FaeA2 (SEQ ID NO:68), FaeB2 (SEQ ID NO:71), Axe1 (SEQ ID NO:80), Axe2 (SEQ ID NO:83) and Pme1 (SEQ ID NO:86) possess carbohydrate esterase activity.

[0140] The enzyme Axe3 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:61 and the cDNA sequence represented herein as SEQ ID NO:63. The Axe3 nucleic acid sequence encodes a 313 amino acid sequence, represented herein as SEQ ID NO:62. The signal peptide for Axe3 is located from positions 1 to about position 21 of SEQ ID NO:62, with the mature protein spanning from about position 22 to position 313 of SEQ ID NO:62. Within Axe3 is a catalytic domain (CD). The amino acid sequence containing the CD of Axe3 spans from a starting point of about position 22 of SEQ ID NO:62 to an ending point of about position 255 of SEQ ID NO:62. Based on homology, Axe3 can be assigned to CAZy Family CE 1. As illustrated below in Example 19, Axe3 exhibits acetyl esterase enzymatic

activity. Axe3 also possesses significant homology (about 69% from amino acids 41 to 312 of Axe3) with an acetyl xylan esterase from *Penicillium purpurogenum* (Genbank Accession No. AAM93261.1; see also Gordillo et al., Mycol. Res. 110:1129 (2006)). Based on this degree of homology, Axe3 is expected to exhibit similar acetyl xylan esterase enzymatic activity.

[0141] The enzyme FaeA1 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:64 and the cDNA sequence represented herein as SEQ ID NO:66. The FaeA1 nucleic acid sequence encodes a 279 amino acid sequence, represented herein as SEQ ID NO:65. The signal peptide for FaeA1 is located from positions 1 to about position 20 of SEQ ID NO:65, with the mature protein spanning from about position 21 to position 279 of SEQ ID NO:65. Within FaeA1 is a catalytic domain (CD). The amino acid sequence containing the CD of FaeA1 spans from a starting point of about position 33 of SEQ ID NO:65 to an ending point of about position 216 of SEQ ID NO:65. Based on homology, FaeA1 can be assigned to CAZy Family CE 1. As illustrated below in Example 20, FaeA1 exhibits ferulic acid esterase enzymatic activity. FaeA1 also possesses significant homology (about 51% from amino acids 22 to 264 of FaeA1) with a ferulic acid esterase from *Aspergillus clavatus* (Genbank Accession No. CAC85738). Based on this degree of homology, FaeA1 is expected to exhibit similar ferulic acid esterase enzymatic activity.

[0142] The enzyme FaeA2 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:67 and the cDNA sequence represented herein as SEQ ID NO:69. The FaeA2 nucleic acid sequence encodes a 302 amino acid sequence, represented herein as SEQ ID NO:68. The signal peptide for FaeA2 is located from positions 1 to about position 26 of SEQ ID NO:68, with the mature protein spanning from about position 27 to position 302 of SEQ ID NO:68. As illustrated below in Example 20, FaeA2 exhibits ferulic acid esterase enzymatic activity. FaeA2 also possesses significant homology (about 45% from amino acids 32 to 301 of FaeA2) with a ferulic acid esterase from *Aspergillus clavatus* (Genbank Accession No. CAC85738). Based on this degree of homology, FaeA2 is expected to exhibit similar ferulic acid esterase enzymatic activity.

[0143] The enzyme FaeB2 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:70 and the cDNA sequence represented herein as SEQ ID NO:72. The FaeB2 nucleic acid sequence encodes a 319 amino acid sequence, represented herein as SEQ ID NO:71. The signal peptide for FaeB2 is located from positions 1 to about position 22 of SEQ ID NO:71, with the mature protein spanning from about position 23 to position 319 of SEQ ID NO:71. Within FaeB2 is a catalytic domain (CD). The amino acid sequence containing the CD of FaeB2 spans from a starting point of about position 63 of SEQ ID NO:71 to an ending point of about position 278 of SEQ ID NO:71. As illustrated below in Example 20, FaeB2 exhibits ferulic acid esterase enzymatic activity. FaeB2 also possesses significant homology (about 68% from amino acids 46 to 319 of FaeB2) with a type B ferulic acid esterase from *Neurospora crassa* (Genbank Accession No. CAC05587). Based on this degree of homology, FaeB2 is expected to exhibit similar ferulic acid esterase enzymatic activity.

[0144] The enzyme Axe1 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:79 and the cDNA sequence represented herein as SEQ ID NO:81. The Axe1 nucleic acid sequence encodes a 303 amino acid sequence, represented herein as SEQ ID NO:80. The signal peptide for Axe1 is located from positions 1 to about position 21 of SEQ ID NO: 80, with the mature protein spanning from about position 22 to position 303 of SEQ ID NO:80. Within Axe1 are two domains: a catalytic domain (CD) and a cellulose-binding domain (CBD). The amino acid sequence containing the CD of Axe1 spans from a starting point of about position 37 of SEQ ID NO:80 to an ending point of about position 246 of SEQ ID NO:80. The amino acid sequence containing the CBD of Axe1 spans from a starting point of about position 271 of SEQ ID NO:80 to an ending point of about position 303 of SEQ ID NO:80. Based on homology, Axe1 can be assigned to CAZy Families CE 5 and CBM 1. As illustrated below in Example 19, Axe1 exhibits acetyl esterase enzymatic activity. Axe1 also possesses significant homology (about 57% from amino acids 3 to 303 of Axe1) with a acetyl xylan esterase from *Hypocrea jecorina* (Genbank Accession No. CAA93247; see also Margolles-Clark et al., Eur. J. Biochem. 237:553 (1996)). Based on this degree of homology, Axe1 is expected to exhibit similar acetyl xylan esterase enzymatic activity.

[0145] The enzyme Axe2 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:82 and the cDNA sequence represented herein as SEQ ID NO:84. The Axe2 nucleic acid sequence encodes a 228 amino acid sequence, represented herein as SEQ ID NO:83. The signal peptide for Axe2 is located from positions 1 to about position 17 of SEQ ID NO: 83, with the mature protein spanning from about position 18 to position 228 of SEQ ID NO:83. Within Axe2 is a catalytic domain (CD). The amino acid sequence containing the CD of Axe2 spans from a starting point of about position 26 of SEQ ID NO:83 to an ending point of about position 242 of SEQ ID NO:83. Based on homology, Axe2 can be assigned to CAZy Family CE 5. As illustrated below in Example 19, Axe2 exhibits acetyl esterase enzymatic activity. Axe2 also possesses significant homology (about 58% from amino acids 7 to 225 of Axe2) with a acetyl xylan esterase from *Penicillium purpurogenum* (Genbank Accession No. 059893; see also Egana et al., Biotechnol. Appl. Biochem. 24(1):33 (1996)). Based on this degree of homology, Axe2 is expected to exhibit similar acetyl xylan esterase enzymatic activity.

[0146] The enzyme Pme1 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:85 and the cDNA sequence represented herein as SEQ ID NO:87. The Pme1 nucleic acid sequence encodes a 331 amino acid

sequence, represented herein as SEQ ID NO:86. The signal peptide for Pmel is located from positions 1 to about position 27 of SEQ ID NO: 86, with the mature protein spanning from about position 28 to position 331 of SEQ ID NO:86. Within Pmel is a catalytic domain (CD). The amino acid sequence containing the CD of Pmel spans from a starting point of about position 43 of SEQ ID NO:86 to an ending point of about position 327 of SEQ ID NO:86. Based on homology, Pme1 can be assigned to CAZy Family CE 8. As illustrated below in Example 22, Pmel exhibits pectin methyl esterase enzymatic activity on substrates such as methylated pectin (e.g., the ability to hydrolyze citrus pectin with a degree of methylation of 65 leading to the formation of unsubstituted galacturonic acid). Pectin methyl esterases catalyze the hydrolysis of methylester groups of pectins such as those found in the cell walls of plants. Pmel also possesses significant homology (about 55% from amino acids 7 to 331 of Pmel) with a pectin methyl esterase from *Emericella nidulans* (Genbank Accession No. ABF50865.1; see also Bauer et al., Proc. Natl. Acad. Sci. U.S.A. 103:11417 (2006)). Based on this degree of homology, Pmel is expected to exhibit similar pectin methyl esterase enzymatic activity.

Arabinofuranosidases

[0147] Certain proteins of the present disclosure possess arabinofuranosidase enzymatic activity. The polypeptides denoted herein as Abf1 (SEQ ID NO:74) and Abf2 (SEQ ID NO:77) possess arabinofuranosidase activity.

[0148] The enzyme Abf1 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:73 and the cDNA sequence represented herein as SEQ ID NO:75. The Abf1 nucleic acid sequence encodes a 370 amino acid sequence, represented herein as SEQ ID NO:74. The signal peptide for Abf1 is located from positions 1 to about position 22 of SEQ ID NO:74, with the mature protein spanning from about position 23 to position 370 of SEQ ID NO:74. Within Abf1 are two domains: a catalytic domain (CD) and a cellulose-binding domain (CBD). The amino acid sequence containing the CD of Abf1 spans from a starting point of about position 24 of SEQ ID NO:74 to an ending point of about position 294 of SEQ ID NO:74. The amino acid sequence containing the CBD of Abf1 spans from a starting point of about position 339 of SEQ ID NO:74 to an ending point of about position 370 of SEQ ID NO:74. Based on homology, Abf1 can be assigned to CAZy Families CBM 1, CBM 13 and GH 62. As illustrated below in Example 21, Abf1 exhibits α -arabinofuranosidase enzymatic activity. Abf1 also possesses significant homology (about 70% from amino acids 19 to 322 of Abf1) with a arabinofuranosidase from *Streptomyces thermoviolaceus* (Genbank Accession No. JC7820; see also Tsujibo et al., Biosci. Biotechnol. Biochem. 66:434 (2002)). Based on this degree of homology, Abf1 is expected to exhibit similar arabinofuranosidase enzymatic activity.

[0149] The enzyme Abf2 is encoded by the nucleic acid sequence represented herein as SEQ ID NO:76 and the cDNA sequence represented herein as SEQ ID NO:78. The Abf2 nucleic acid sequence encodes a 321 amino acid sequence, represented herein as SEQ ID NO:77. The signal peptide for Abf2 is located from positions 1 to about position 19 of SEQ ID NO:77, with the mature protein spanning from about position 20 to position 321 of SEQ ID NO:77. Within Abf2 is a catalytic domain (CD). The amino acid sequence containing the CD of Abf2 spans from a starting point of about position 22 of SEQ ID NO:77 to an ending point of about position 291 of SEQ ID NO:77. Based on homology, Abf2 can be assigned to CAZy Family GH 62. As illustrated below in Example 21, Abf2 exhibits α -arabinofuranosidase enzymatic activity. Abf2 also possesses significant homology (about 75% from amino acids 21 to 315 of Abf2) with a arabinofuranosidase from *Streptomyces thermoviolaceus* (Genbank Accession No. JC7820; see also Tsujibo et al., Biosci. Biotechnol. Biochem. 66:434 (2002)). Based on this degree of homology, Abf2 is expected to exhibit similar arabinofuranosidase enzymatic activity.

[0150] Physical properties, biochemical characteristics and substrate specificities of the glycosidases, pectate lyases, chitinases, carbohydrate esterases, and arabinofuranosidases of the present disclosure are illustrated in Tables 15, 16, 17 and 18 below. Physical and biochemical characteristics include molecular weight, isoelectric point, pH and temperature optima of enzymatic activity, and the stability of the enzymes after heating to 50°C for five hours at pH 5.0. Substrate specificities demonstrating enzyme activity for the indicated substrates (specific and non-specific) were determined using standard assays known in the art, such as those disclosed in the Examples below.

Table 15. Physical and Biochemical Properties of C1 Enzymes

Enzyme	SEQ ID NO	MW (kDa)	pI	pH (50%)	T °C (50%)	Stability pH 5.0
Bgl3A	44	106	4.8	4.0 (2.5-7.0)	60 (45-70)	64%
Gal53A	47	30	8.2	4.5 (2.7-6.0)	55-60 (30-70)	100%
Gla15A	50	68	4.3	5.5 (3.5-7.0)	70	50%
Pe1	53	33	5.3	8.5 (7.8-8.8)	60-65 (30-70)	100% **
Chi18A	56	43	4.0	7.5 (5.7-8.7)	70 (60-75)	76%

EP 2 183 363 B1

(continued)

Enzyme	SEQ ID NO	MW (kDa)	pI	pH (50%)	T °C (50%)	Stability pH 5.0
Gls2A	59	80	6.6	5.9 (5.0-7.0)	70 (55-75)	100%
Axe3	62	35	6.5-6.8			
FaeA1	65	25-40	3.5-4.0			
FaeA2	68	29.1*	4.87*			
FaeB2	71	29.3*	4.28*			
Abf1	74	35	7.3 and 8.5-9.0			
Abf2	77	33	4.5-5.0			
Axe1	80	28*	7.2*			
Axe2	83	27	6			
Pme1	86	32.6*	8.79*			
Abn1	89	32.8*	5.57*			
bx12	95	75.9*	4.8*			
MW = Molecular Weight, kiloDaltons (kDa) (* = Predicted) pI = isoelectric point (* = Predicted) pH (50%) = pH optimum of enzyme activity (range of at least 50% activity) T °C (50%) = Temperature optimum of enzyme activity (range of at least 50% activity) Stability pH 5.0 = residual enzyme activity after 3 hour incubation at 50 °C, pH 5.0 ** pH 8.5, 1mM CaCl ₂						

Table 16. Substrate Specificities of C1 Enzymes (Activity Towards Substrates (units/mg))

Substrate	Bgl3A	Gal53A	Pel1
CMC (S-N Assay)	1.5	0	0
β-glucan from Barley	18	0	0
Birchwood GlucuronoXylan	0	0	0
Wheat ArabinoXylan	0	0	0
Xyloglucan from Tamarind	0	0	0
Laminarin (β-1,3; β-1,6)	109	0	0
Polygalacturonic acid (RS)	0	0	70
Citrus Pectin (Viscosimetry, A235)	0	0	525*
Citrus Pectin (Etherification 89%, A235)	0	0	82*
Citrus Pectin (Etherification 65%, A235)	0	0	126*
Citrus Pectin (Etherification 26%, A235)	0	0	133*
Galactan	0	620	0
Arabinan Debranched	0	97	0
Arabinan Branched	0	90	0
Arabinogalactan	0	0	0
Galactomannan	0	0	0
p-NPh-β-D-cellobioside	0.03	0	0
p-NPh-β-D-lactoside	0	0	0

EP 2 183 363 B1

(continued)

Substrate	Bgl3A	Gal53A	Pel1
p-NPh-β-L-arabinofuranoside	0	0	0
p-NPh-β-L-arabinopyranoside	0	0	0
o-NPh-β-D-xylopyranoside	0	0	0
p-NPh-β-D-glucopyranoside	39	0	0
p-NPh-β-L-galactopyranoside	0	0	0
Cellobiose (β-1,4)	52	0	0
Laminaribiose (β-1,3)	86	0	0
α-Sophorose (β-1,2)	50	0	0
Gentibiose (β-1,6)	107	0	0
Maltose	0	0	0
* C1 pectate lyase is a Ca ⁺² -dependent enzyme, so activities were measured in 1 mM of CaCl ₂ .			

Table 17. Substrate Specificities of C1 Gla15A (Activity Towards Substrates (units/mg))

Substrate	Glucoamylase Activity
Starch soluble (potato) NS*	150
Amylose soluble (potato), NS	87
Amylopectin soluble (potato), NS	104
Pullulan (Aureobasidium pullulans)	11
Starch insoluble (potato) NS	35
Amylose insoluble (potato), NS	59
Amylopectin insoluble (potato), NS	43
Maltose**	18
Isomaltose**	1.2
Maltotriose**	48
Maltohexaose**	61
pNph-α-D-glucopiranoside	0

Table 18. Substrate Specificities of C1 Enzymes (Activity Towards Substrates (units/mg))

Substrate	Chi18A	Gls2A
Chitosan soluble, 140kDa	35	28
Chitin Colloid	8	0
CMC	0	0
β-glucan	0	0
Birchwood GlucuronoXylan	0	0
p-NPh-β-D-N-acetylglucosamine	0.1	0
Chitosan, 140kDa viscometry	210	0

EP 2 183 363 B1

[0151] pH and temperature dependencies of certain enzymes of the present disclosure are illustrated in Tables 19 and 20 below. pH and temperature dependencies were determined using the standard assays with the substrate and conditions listed below:

Enzyme	pH	Temperature
Bgl3A	p-NPh-β-D-glucopyranoside	Cellobiose
Gla15A	Potato Starch, 50°C, 10 minutes	Potato Starch, pH 5.0, 10 minutes
Chi18A	Colloidal chitin, 50°C, 10 minutes	Chitosan (140 kD), pH 5.0, 10 minutes
Gls2A	Chitosan (140 kD), 50°C, 10 minutes	Chitosan (140 kD), pH 5.0, 10 minutes
Ga153A ¹	Arabinan branched, 50°C, 10 minutes	Arabinan branched, pH 5.0, 10 minutes
Ga153A ²	Galactan, 50°C, 10 minutes	Not Applicable
Pel1	A235, PGA + 1 mM CaCl ₂	A235, PGA + 1 mM CaCl ₂

[0152] The temperature/pH that exhibited the highest enzyme activity was set at 100% in the tables below, with the remaining values expressed a percentage of this standard. The pH and temperature dependencies of certain enzymes are also illustrated graphically in Figures 9 and 10, respectively.

Table 19. pH Dependency of C1 Enzymes (% Activity)

pH	Bgl3A	Gla15 A	Chi18A	Gls2 a	pH	Gal 53A ¹	pH	Gal 53A ²	pH	Pel1
2.0	2									
2.6	50	37			2.45	39.5	2.9	0.3		
3.0	70	43	2	0	2.73	55.6	3.51	26.5		
3.5 3	90	51	2	3	3.5	70.4	4.22	93.9	7.25	2.8
4.0	100	78	2	10.3	4.05	89.3	4.62	97.4	7.75	41.9
4.5 7	98	90	8	40	4.5	100.0	4.90	97.7	8.05	70.5
5.0	93	97	20	50	4.95	75.6	5.45	100	8.18	88.8
5.5	83	100	30	64	5.5	70.6	6.02	95.8	8.42	100
6.0	72	96	57	100	6.02	48.2	6.51	91.6	8.88	49.6
6.5	58	83	62	65	6.47	34.5	7.03	71.9	8.99	33
7.0	47	50	83	37.8	7.03	19.8	7.50	51.7	9.5	5
7.5	30	25	100		7.5	17.8	8.00	42.2		
8	17	1	98		9.04	17.1	9.11	1.7		
9			25							

Table 20. Temperature Dependency of C1 Enzymes (% Activity)

Temp °C	Bgl3A	Gla15 A	Chi18A	Gls2 a	Gal 53A ¹	Temp °C	Pel1
25				10		21	6.9
30	10		20	15	46.6	30	15.2
35	18		21	18			
40	36	20	25	22	80.3	39	29.6
45	48	19	36	32		47	51.5
50	65	20	40	43	92.1		
55	95	44	42	53		56	82.5
60	100	62	47	60	100	61	100

(continued)

Temp °C	Bgl3A	Gla15 A	Chi18A	Gls2 a	Gal 53A ¹	Temp °C	Pel1
65	80	88	60	80	97.1	67	94.8
70	48	100	100	100	59.8	72	33.9
75	15	5	53	53			
80			12	15			
85							
90							

[0153] More specifically, an isolated protein, such as an enzyme according to the present invention, is a protein (including a polypeptide or peptide) that has been removed from its natural milieu (*i.e.*, that has been subject to human manipulation) and can include purified proteins, partially purified proteins, recombinantly produced proteins, synthetically produced proteins, proteins complexed with lipids, soluble proteins, and isolated proteins associated with other proteins, for example. As such, "isolated" does not reflect the extent to which the protein has been purified. Preferably, an isolated protein of the present invention is produced recombinantly. In addition, and by way of example, a "*C. lucknowens* protein" or "*C. lucknowens* enzyme" refers to a protein (generally including a homologue of a naturally occurring protein) from *Chrysosporium lucknowense* or to a protein that has been otherwise produced from the knowledge of the structure (*e.g.*, sequence) and perhaps the function of a naturally occurring protein from *Chrysosporium lucknowense*. In other words, a *C. lucknowens* protein includes any protein that has substantially similar structure and function of a naturally occurring *C. lucknowens* protein or that is a biologically active (*i.e.*, has biological activity) homologue of a naturally occurring protein from *C. lucknowens* as described in detail herein. As such, a *C. lucknowens* protein can include purified, partially purified, recombinant, mutated/modified and synthetic proteins. Herein, the terms "modification" and "mutation" can be used interchangeably, particularly with regard to the modifications/mutations to the amino acid sequence of a *C. lucknowens* protein (or nucleic acid sequences) described herein. An isolated protein according to the present invention can be isolated from its natural source, produced recombinantly or produced synthetically.

[0154] Herein, the terms "modification" and "mutation" can be used interchangeably, particularly with regard to the modifications/mutations to the primary amino acid sequences of a protein or peptide (or nucleic acid sequences) described herein. The term "modification" can also be used to describe post-translational modifications to a protein or peptide including, but not limited to, methylation, farnesylation, carboxymethylation, geranyl geranylation, glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitation, and/or amidation. Modifications can also include, for example, complexing a protein or peptide with another compound. Such modifications can be considered to be mutations, for example, if the modification is different than the post-translational modification that occurs in the natural, wild-type protein or peptide.

[0155] As used herein, the term "homologue" is used to refer to a protein or peptide which differs from a naturally occurring protein or peptide (*i.e.*, the "prototype" or "wild-type" protein) by minor modifications to the naturally occurring protein or peptide, but which maintains the basic protein and side chain structure of the naturally occurring form. Such changes include, but are not limited to: changes in one or a few amino acid side chains; changes one or a few amino acids, including deletions (*e.g.*, a truncated version of the protein or peptide), insertions and/or substitutions; changes in stereochemistry of one or a few atoms; and/or minor derivatizations, including but not limited to: methylation, glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitation, amidation and/or addition of glycosylphosphatidyl inositol. A homologue can have either enhanced, decreased, or substantially similar properties as compared to the naturally occurring protein or peptide. A homologue can include an agonist of a protein or an antagonist of a protein.

[0156] Homologues can be the result of natural allelic variation or natural mutation. A naturally occurring allelic variant of a nucleic acid encoding a protein is a gene that occurs at essentially the same locus (or loci) in the genome as the gene which encodes such protein, but which, due to natural variations caused by, for example, mutation or recombination, has a similar but not identical sequence. Allelic variants typically encode proteins having similar activity to that of the protein encoded by the gene to which they are being compared. One class of allelic variants can encode the same protein but have different nucleic acid sequences due to the degeneracy of the genetic code. Allelic variants can also comprise alterations in the 5' or 3' untranslated regions of the gene (*e.g.*, in regulatory control regions). Allelic variants are well known to those skilled in the art.

[0157] Homologues can be produced using techniques known in the art for the production of proteins including, but not limited to, direct modifications to the isolated, naturally occurring protein, direct protein synthesis, or modifications to the nucleic acid sequence encoding the protein using, for example, classic or recombinant DNA techniques to effect random or targeted mutagenesis.

[0158] Modifications in protein homologues, as compared to the wild-type protein, either agonize, antagonize, or do not substantially change, the basic biological activity of the homologue as compared to the naturally occurring protein. Modifications of a protein, such as in a homologue, may result in proteins having the same biological activity as the naturally occurring protein, or in proteins having decreased or increased biological activity as compared to the naturally occurring protein. Modifications which result in a decrease in protein expression or a decrease in the activity of the protein, can be referred to as inactivation (complete or partial), down-regulation, or decreased action of a protein. Similarly, modifications which result in an increase in protein expression or an increase in the activity of the protein, can be referred to as amplification, overproduction, activation, enhancement, up-regulation or increased action of a protein.

[0159] According to the present disclosure, an isolated protein, including a biologically active homologue or fragment thereof, has at least one characteristic of biological activity of a wild-type, or naturally occurring, protein. As discussed above, in general, the biological activity or biological action of a protein refers to any function(s) exhibited or performed by the protein that is ascribed to the naturally occurring form of the protein as measured or observed *in vivo* (i.e., in the natural physiological environment of the protein) or *in vitro* (i.e., under laboratory conditions). The biological activity of a protein of the present disclosure can include an enzyme activity (catalytic activity and/or substrate binding activity), such as cellulase activity, hemicellulase activity, β -glucanase activity, β -glucosidase activity, α -galactosidase activity, β -galactosidase activity, xylanase activity or any other activity disclosed herein. Specific biological activities of the proteins disclosed herein are described in detail above and in the Examples. Methods of detecting and measuring the biological activity of a protein of the invention and disclosure include, but are not limited to, the assays described in the Examples section below. Such assays include, but are not limited to, measurement of enzyme activity (e.g., catalytic activity), measurement of substrate binding, and the like. It is noted that an isolated protein of the present disclosure (including homologues) is not required to have a biological activity such as catalytic activity. A protein can be a truncated, mutated or inactive protein, or lack at least one activity of the wild-type enzyme, for example. Inactive proteins may be useful in some screening assays, for example, or for other purposes such as antibody production.

[0160] Methods to measure protein expression levels of a protein according to the invention and disclosure include, but are not limited to: western blotting, immunocytochemistry, flow cytometry or other immunologic-based assays; assays based on a property of the protein including but not limited to, ligand binding or interaction with other protein partners. Binding assays are also well known in the art. For example, a BIAcore machine can be used to determine the binding constant of a complex between two proteins. The dissociation constant for the complex can be determined by monitoring changes in the refractive index with respect to time as buffer is passed over the chip (O'Shannessy et al. Anal. Biochem. 212:457-468 (1993); Schuster et al., Nature 365:343-347 (1993)). Other suitable assays for measuring the binding of one protein to another include, for example, immunoassays such as enzyme linked immunoabsorbent assays (ELISA) and radioimmunoassays (RIA), or determination of binding by monitoring the change in the spectroscopic or optical properties of the proteins through fluorescence, UV absorption, circular dichroism, or nuclear magnetic resonance (NMR).

[0161] The enzyme and protein of the present invention may be a desirable target for modification and use in the processes described herein. The protein has been described in terms of function and amino acid sequence (and nucleic acid sequence encoding the same) of the representative wild-type protein. In one embodiment of the invention, homologues of the protein (which can include related proteins from other organisms or modified forms of the given protein) are encompassed for use in the invention. Homologues of a protein encompassed by the present invention can comprise, consist essentially of, or consist of, in one embodiment, an amino acid sequence that is at least 90% identical, and more preferably at least about 95% identical, and more preferably at least about 96% identical, and more preferably at least about 97% identical, and more preferably at least about 98% identical, and more preferably at least about 99% identical, to an amino acid sequence disclosed herein that represents the amino acid sequence of an enzyme or protein according to the invention (including a biologically active domain of a full-length protein). Preferably, the amino acid sequence of the homologue has a biological activity of the wild-type or reference protein or of a biologically active domain thereof (e.g., a catalytic domain).

[0162] In one embodiment, a protein of the present invention comprises, consists essentially of, or consists of an amino acid sequence that is less than 100% identical to an amino acid sequence of SEQ ID NO:32 (i.e., a homologue). In another aspect of the invention, a homologue according to the present invention has an amino acid sequence that is less than about 99% identical to any of such amino acid sequences, and in another embodiment, is less than about 98% identical to any of such amino acid sequences, and in another embodiment, is less than about 97% identical to any of such amino acid sequences, and in another embodiment, is less than about 96% identical to any of such amino acid sequences, and in another embodiment, is less than about 95% identical to any of such amino acid sequences, and in another embodiment, is less than about 94% identical to any of such amino acid sequences, and in another embodiment, is less than about 93% identical to any of such amino acid sequences, and in another embodiment, is less than about 92% identical to any of such amino acid sequences, and in another embodiment, is less than about 91% identical to any of such amino acid sequences.

[0163] As used herein, unless otherwise specified, reference to a percent (%) identity refers to an evaluation of homology which is performed using: (1) a BLAST 2.0 Basic BLAST homology search using blastp for amino acid searches

and blastn for nucleic acid searches with standard default parameters, wherein the query sequence is filtered for low complexity regions by default (described in Altschul, S.F., Madden, T.L., Schääffer, A.A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D.J. (1997) "Gapped BLAST and PSI-BLAST: a new generation of protein database search program." *Nucleic Acids Res.* 25:3389-3402); (2) a BLAST 2 alignment (using the parameters described below); (3) PSI-BLAST with the standard default parameters (Position-Specific Iterated BLAST; and/or (4) CAZy homology determined using standard default parameters from the Carbohydrate Active EnZymes database (Coutinho, P.M. & Henrissat, B. (1999) *Carbohydrate-active enzymes: an integrated database approach*. In "Recent Advances in Carbohydrate Bioengineering", H.J. Gilbert, G. Davies, B. Henrissat and B. Svensson eds., The Royal Society of Chemistry, Cambridge, pp. 3-12).

[0164] It is noted that due to some differences in the standard parameters between BLAST 2.0 Basic BLAST and BLAST 2, two specific sequences might be recognized as having significant homology using the BLAST 2 program, whereas a search performed in BLAST 2.0 Basic BLAST using one of the sequences as the query sequence may not identify the second sequence in the top matches. In addition, PSI-BLAST provides an automated, easy-to-use version of a "profile" search, which is a sensitive way to look for sequence homologues. The program first performs a gapped BLAST database search. The PSI-BLAST program uses the information from any significant alignments returned to construct a position-specific score matrix, which replaces the query sequence for the next round of database searching. Therefore, it is to be understood that percent identity can be determined by using any one of these programs.

[0165] Two specific sequences can be aligned to one another using BLAST 2 sequence as described in Tatusova and Madden, (1999), "Blast 2 sequences - a new tool for comparing protein and nucleotide sequences", FEMS Microbiol Lett. 174:247-250. BLAST 2 sequence alignment is performed in blastp or blastn using the BLAST 2.0 algorithm to perform a Gapped BLAST search (BLAST 2.0) between the two sequences allowing for the introduction of gaps (deletions and insertions) in the resulting alignment. For purposes of clarity herein, a BLAST 2 sequence alignment is performed using the standard default parameters as follows.

For blastn, using 0 BLOSUM62 matrix:

Reward for match = 1
Penalty for mismatch = -2
Open gap (5) and extension gap (2) penalties
gap x_dropoff (50) expect (10) word size (11) filter (on)

For blastp, using 0 BLOSUM62 matrix:

Open gap (11) and extension gap (1) penalties
gap x dropoff (50) expect (10) word size (3) filter (on).

[0166] A protein of the present disclosure can also include proteins having an amino acid sequence comprising at least 10 contiguous amino acid residues of any of the sequences described herein (*i.e.*, 10 contiguous amino acid residues having 100% identity with 10 contiguous amino acids of SEQ ID NO:2). In other embodiments, a homologue of a protein amino acid sequence includes amino acid sequences comprising at least 20, or at least 30, or at least 40, or at least 50, or at least 75, or at least 100, or at least 125, or at least 150, or at least 175, or at least 150, or at least 200, or at least 250, or at least 300, or at least 350 contiguous amino acid residues of any of the amino acid sequence represented disclosed herein. Even small fragments of proteins without biological activity are useful in the present disclosure, for example, in the preparation of antibodies against the full-length protein or in a screening assay (*e.g.*, a binding assay). Fragments can also be used to construct fusion proteins, for example, where the fusion protein comprises functional domains from two or more different proteins (*e.g.*, a CBD from one protein linked to a CD from another protein). In one embodiment, a homologue has a measurable or detectable biological activity associated with the wild-type protein (*e.g.*, enzymatic activity).

[0167] As used herein, the term "contiguous" or "consecutive", with regard to nucleic acid or amino acid sequences described herein, means to be connected in an unbroken sequence. For example, for a first sequence to comprise 30 contiguous (or consecutive) amino acids of a second sequence, means that the first sequence includes an unbroken sequence of 30 amino acid residues that is 100% identical to an unbroken sequence of 30 amino acid residues in the second sequence. Similarly, for a first sequence to have "100% identity" with a second sequence means that the first sequence exactly matches the second sequence with no gaps between nucleotides or amino acids.

[0168] In another embodiment, a protein of the present disclosure, including a homologue, includes a protein having an amino acid sequence that is sufficiently similar to a natural amino acid sequence that a nucleic acid sequence encoding the homologue is capable of hybridizing under moderate, high or very high stringency conditions (described below) to (*i.e.*, with) a nucleic acid molecule encoding the natural protein (*i.e.*, to the complement of the nucleic acid strand encoding the natural amino acid sequence). Preferably, a homologue of a protein of the present disclosure is encoded by a nucleic acid molecule comprising a nucleic acid sequence that hybridizes under low, moderate, or high stringency conditions

to the complement of a nucleic acid sequence that encodes a protein comprising, consisting essentially of, or consisting of, an amino acid sequence represented by any of: SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, SEQ ID NO:14, SEQ ID NO:17, SEQ ID NO:20, SEQ ID NO:23, SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:62, SEQ ID NO:65, SEQ ID NO:68, SEQ ID NO:71, SEQ ID NO:74, SEQ ID NO:77, SEQ ID NO:80, SEQ ID NO:83, SEQ ID NO:86, SEQ ID NO:89, SEQ ID NO:92, or SEQ ID NO:94. Such hybridization conditions are described in detail below.

[0169] A nucleic acid sequence complement of nucleic acid sequence encoding a protein of the present invention refers to the nucleic acid sequence of the nucleic acid strand that is complementary to the strand which encodes the protein. It will be appreciated that a double stranded DNA which encodes a given amino acid sequence comprises a single strand DNA and its complementary strand having a sequence that is a complement to the single strand DNA. As such, nucleic acid molecules of the present invention can be either double-stranded or single-stranded. Methods to deduce a complementary sequence are known to those skilled in the art. It should be noted that since nucleic acid sequencing technologies are not entirely error-free, the sequences presented herein, at best, represent apparent sequences of the proteins of the present invention.

[0170] As used herein, reference to hybridization conditions refers to standard hybridization conditions under which nucleic acid molecules are used to identify similar nucleic acid molecules. Such standard conditions are disclosed, for example, in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Labs Press, 1989. Sambrook et al., *ibid.* (see specifically, pages 9.31-9.62). In addition, formulae to calculate the appropriate hybridization and wash conditions to achieve hybridization permitting varying degrees of mismatch of nucleotides are disclosed, for example, in Meinkoth et al., 1984, *Anal. Biochem.* 138, 267-284; Meinkoth et al., *ibid.*

[0171] More particularly, moderate stringency hybridization and washing conditions, as referred to herein, refer to conditions which permit isolation of nucleic acid molecules having at least about 70% nucleic acid sequence identity with the nucleic acid molecule being used to probe in the hybridization reaction (i.e., conditions permitting about 30% or less mismatch of nucleotides). High stringency hybridization and washing conditions, as referred to herein, refer to conditions which permit isolation of nucleic acid molecules having at least about 80% nucleic acid sequence identity with the nucleic acid molecule being used to probe in the hybridization reaction (i.e., conditions permitting about 20% or less mismatch of nucleotides). Very high stringency hybridization and washing conditions, as referred to herein, refer to conditions which permit isolation of nucleic acid molecules having at least about 90% nucleic acid sequence identity with the nucleic acid molecule being used to probe in the hybridization reaction (i.e., conditions permitting about 10% or less mismatch of nucleotides). As discussed above, one of skill in the art can use the formulae in Meinkoth et al., *ibid.* to calculate the appropriate hybridization and wash conditions to achieve these particular levels of nucleotide mismatch. Such conditions will vary, depending on whether DNA:RNA or DNA:DNA hybrids are being formed. Calculated melting temperatures for DNA:DNA hybrids are 10°C less than for DNA:RNA hybrids. In particular embodiments, stringent hybridization conditions for DNA:DNA hybrids include hybridization at an ionic strength of 6X SSC (0.9 M Na⁺) at a temperature of between about 20°C and about 35°C (lower stringency), more preferably, between about 28°C and about 40°C (more stringent), and even more preferably, between about 35°C and about 45°C (even more stringent), with appropriate wash conditions. In particular embodiments, stringent hybridization conditions for DNA:RNA hybrids include hybridization at an ionic strength of 6X SSC (0.9 M Na⁺) at a temperature of between about 30°C and about 45°C, more preferably, between about 38°C and about 50°C, and even more preferably, between about 45°C and about 55°C, with similarly stringent wash conditions. These values are based on calculations of a melting temperature for molecules larger than about 100 nucleotides, 0% formamide and a G + C content of about 40%. Alternatively, T_m can be calculated empirically as set forth in Sambrook et al., *supra*, pages 9.31 to 9.62. In general, the wash conditions should be as stringent as possible, and should be appropriate for the chosen hybridization conditions. For example, hybridization conditions can include a combination of salt and temperature conditions that are approximately 20-25°C below the calculated T_m of a particular hybrid, and wash conditions typically include a combination of salt and temperature conditions that are approximately 12-20°C below the calculated T_m of the particular hybrid. One example of hybridization conditions suitable for use with DNA:DNA hybrids includes a 2-24 hour hybridization in 6X SSC (50% formamide) at about 42°C, followed by washing steps that include one or more washes at room temperature in about 2X SSC, followed by additional washes at higher temperatures and lower ionic strength (e.g., at least one wash as about 37°C in about 0.1X-0.5X SSC, followed by at least one wash at about 68°C in about 0.1X-0.5X SSC).

[0172] The minimum size of a protein and/or homologue of the present disclosure is a size sufficient to have biological activity or, when the protein is not required to have such activity, sufficient to be useful for another purpose associated with a protein of the present disclosure, such as for the production of antibodies that bind to a naturally occurring protein. In one embodiment, the protein of the present disclosure is at least 20 amino acids in length, or at least about 25 amino acids in length, or at least about 30 amino acids in length, or at least about 40 amino acids in length, or at least about 50 amino acids in length, or at least about 60 amino acids in length, or at least about 70 amino acids in length, or at least about 80 amino acids in length, or at least about 90 amino acids in length, or at least about 100 amino acids in

length, or at least about 125 amino acids in length, or at least about 150 amino acids in length, or at least about 175 amino acids in length, or at least about 200 amino acids in length, or at least about 250 amino acids in length, and so on up to a full length of each protein, and including any size in between in increments of one whole integer (one amino acid). There is no limit, other than a practical limit, on the maximum size of such a protein in that the protein can include

a portion of a protein or a full-length protein, plus additional sequence (e.g., a fusion protein sequence), if desired.

[0173] The present invention also includes a fusion protein that includes a protein of the present invention (including a homologue) attached to one or more fusion segments, which are typically heterologous in sequence to the protein sequence (*i.e.*, different than protein sequence). Suitable fusion segments for use with the present invention include, but are not limited to, segments that can: enhance a protein's stability; provide other desirable biological activity; and/or assist with the purification of the protein (e.g., by affinity chromatography). A suitable fusion segment can be a domain of any size that has the desired function (e.g., imparts increased stability, solubility, action or biological activity; and/or simplifies purification of a protein). Fusion segments can be joined to amino and/or carboxyl termini of the domain of a protein of the present invention and can be susceptible to cleavage in order to enable straight-forward recovery of the protein. Fusion proteins are preferably produced by culturing a recombinant cell transfected with a fusion nucleic acid molecule that encodes a protein including the fusion segment attached to either the carboxyl and/or amino terminal end of a domain of a protein of the present invention. Accordingly, proteins of the present invention also include expression products of gene fusions (for example, used to overexpress soluble, active forms of the recombinant protein), of mutagenized genes (such as genes having codon modifications to enhance gene transcription and translation), and of truncated genes (such as genes having membrane binding domains removed to generate soluble forms of a membrane protein, or genes having signal sequences removed which are poorly tolerated in a particular recombinant host).

[0174] In one embodiment of the present invention, any of the amino acid sequences described herein can be produced with from at least one, and up to about 20, additional heterologous amino acids flanking each of the C- and/or N-terminal ends of the specified amino acid sequence. The resulting protein or polypeptide can be referred to as "consisting essentially of" the specified amino acid sequence. According to the present invention, the heterologous amino acids are a sequence of amino acids that are not naturally found (*i.e.*, not found in nature, *in vivo*) flanking the specified amino acid sequence, or that are not related to the function of the specified amino acid sequence, or that would not be encoded by the nucleotides that flank the naturally occurring nucleic acid sequence encoding the specified amino acid sequence as it occurs in the gene, if such nucleotides in the naturally occurring sequence were translated using standard codon usage for the organism from which the given amino acid sequence is derived.

[0175] The present invention also provides enzyme combinations that break down lignocellulose material. Such enzyme combinations or mixtures can include a multi-enzyme composition that contains at least the protein of the present invention in combination with one or more additional proteins of the present disclosure or one or more enzymes or other proteins from other microorganisms, plants, or similar organisms. Synergistic enzyme combinations and related methods are contemplated. The disclosure includes methods to identify the optimum ratios and compositions of enzymes with which to degrade each lignocellulosic material. These methods entail tests to identify the optimum enzyme composition and ratios for efficient conversion of any lignocellulosic substrate to its constituent sugars. The Examples below include assays that may be used to identify optimum ratios and compositions of enzymes with which to degrade lignocellulosic materials.

[0176] Any combination of the proteins disclosed herein is suitable for use in the multi-enzyme compositions of the present disclosure. Due to the complex nature of most biomass sources, which can contain xylan, lignin, protein, and carbohydrates, among other components, preferred enzyme combinations may contain enzymes with a range of substrate specificities that work together to degrade biomass into fermentable sugars in the most efficient manner. One example of a multi-enzyme complex for lignocellulose saccharification is a mixture of cellobiohydrolase(s), xylanase(s), endoglucanase(s), β -glucosidase(s), β -xylosidase(s), and accessory enzymes. However, it is to be understood that the enzymes described specifically herein can be combined with any one or more of the enzymes described herein or with any other available and suitable enzymes, to produce a multi-enzyme composition. The disclosure is not restricted or limited to the specific exemplary combinations listed below.

[0177] In one embodiment, the cellobiohydrolase(s) comprise between about 30% and about 90% or between about 50% and about 70% of the enzymes in the composition, and more preferably, between about 55% and 65%, and more preferably, about 60% of the enzymes in the composition (including any percentage between 50% and 70% in 0.5% increments (e.g., 50%, 50.5%, 51%, etc.).

[0178] In one embodiment, the xylanase(s) comprise between about 10% and about 30% of the enzymes in the composition, and more preferably, between about 15% and about 25%, and more preferably, about 20% of the enzymes in the composition (including any percentage between 10% and 30% in 0.5% increments).

[0179] In one embodiment, the endoglucanase(s) comprise between about 5% and about 15% of the enzymes in the composition, and more preferably, between about 7% and about 13%, and more preferably, about 10% of the enzymes in the composition (including any percentage between 5% and 15% in 0.5% increments).

[0180] In one embodiment, the β -glucosidase(s) comprise between about 1% and about 5% of the enzymes in the

composition, and preferably between about 2% and 4%, and more preferably, about 3% of the enzymes in the composition (including any percentage between 1% and 5% in 0.5% increments).

[0181] In one embodiment, the β -xylosidase(s) comprise between about 1% and about 3% of the enzymes in the composition, and preferably, between about 1.5% and about 2.5%, and more preferably, about 2% of the enzymes in the composition (including any percentage between 1% and 3% in 0.5% increments).

[0182] In one embodiment, the accessory enzymes comprise between about 2% and about 8% of the enzymes in the composition, and preferably, between about 3% and about 7%, and more preferably, about 5% of the enzymes in the composition (including any percentage between 2% and 8% in 0.5% increments).

[0183] One particularly preferred example of a multi-enzyme complex for lignocellulose saccharification is a mixture of about 60% cellobiohydrolase(s), about 20% xylanase(s), about 10% endoglucanase(s), about 3% β -glucosidase(s), about 2% β -xylosidase(s) and about 5% accessory enzyme(s).

[0184] The multi-enzyme composition may comprise at least one cellobiohydrolase. In some embodiments, the cellobiohydrolase may have an amino acid sequence selected from SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11 or a homologue, fusion protein, or fragment thereof that has cellobiohydrolase activity. Compositions comprising at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 or more cellobiohydrolases are contemplated by the disclosure.

[0185] The multi-enzyme composition may comprise at least one endoglucanase. In certain embodiments, the endoglucanase may have an amino acid sequence selected from SEQ ID NO: 14, SEQ ID NO:94, SEQ ID NO:17, SEQ ID NO:20 or SEQ ID NO:23 or a homologue, fusion protein, or fragment thereof that has endoglucanase activity. Compositions comprising at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 or more endoglucanases are contemplated by the disclosure.

[0186] The multi-enzyme composition may comprise at least one xylanase. In some embodiments, the xylanase may have an amino acid sequence selected from SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38 or SEQ ID NO:41 or a homologue, fusion protein, or fragment thereof that has xylanase activity. Compositions comprising at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 or more xylanases are contemplated by the disclosure.

[0187] The multi-enzyme composition may further comprise a β -glucosidase. In one embodiment, the β -glucosidase may have an amino acid sequence of SEQ ID NO:44 or a homologue, fusion protein, or fragment thereof that has β -glucosidase activity. Compositions comprising at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 or more β -glucosidases are contemplated by the disclosure.

[0188] The multi-enzyme composition may further comprise a hemicellulase. In certain embodiments, the hemicellulase may have an amino acid sequence selected from SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:62, SEQ ID NO:65, SEQ ID NO:68, SEQ ID NO:71, SEQ ID NO:74, SEQ ID NO:77, SEQ ID NO:80, SEQ ID NO:83, SEQ ID NO:86, SEQ ID NO:89, SEQ ID NO:92, or a homologue, fusion protein, or fragment thereof that has a hemicellulase activity. Compositions comprising at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 or more hemicellulases are contemplated by the disclosure.

[0189] One or more components of a multi-enzyme composition (other than the protein of the present invention) can be obtained from or derived from a microbial, plant, or other source or combination thereof, and will contain enzymes capable of degrading lignocellulosic material. Examples of enzymes included in the multi-enzyme compositions of the invention include cellulases, hemicellulases (such as xylanases, including endoxylanases, exoxylanase, and β -xylosidase), ligninases, amylases, glucuronidases, proteases, esterases (including ferulic acid esterase), lipases, glucosidases (such as β -glucosidase), glucomannanases, and xylogluconases.

[0190] While the multi-enzyme composition may contain many types of enzymes, mixtures comprising enzymes that increase or enhance sugar release from biomass are preferred, including hemicellulases. In one embodiment, the hemicellulase is selected from a xylanase, an arabinofuranosidase, an acetyl xylan esterase, a glucuronidase, an endogalactanase, a mannanase, an endo-arabinase, an exo-arabinase, an exo-galactanase, a ferulic acid esterase, a galactomannanase, a xylogluconase, or mixtures of any of these. In particular, the enzymes can include glucoamylase, β -xylosidase and/or β -glucosidase. The enzymes of the multi-enzyme composition can be provided by a variety of sources. In one embodiment, the enzymes can be produced by a growing microorganisms or plants which produce the enzymes naturally or by virtue of being genetically modified to express the enzyme or enzymes. In another embodiment, at least one enzyme of the multi-enzyme composition is a commercially available enzyme.

[0191] In some embodiments, the multi-enzyme compositions comprise an accessory enzyme. An accessory enzyme is any additional enzyme capable of hydrolyzing lignocellulose or enhancing or promoting the hydrolysis of lignocellulose, wherein the accessory enzyme is typically provided in addition to a core enzyme or core set of enzymes. An accessory enzyme can have the same or similar function or a different function as an enzyme or enzymes in the core set of enzymes. These enzymes have been described elsewhere herein, and can generally include cellulases, xylanases, ligninases, amylases, lipases, or glucuronidases, for example. Accessory enzymes can include enzymes that when contacted with biomass in a reaction, allow for an increase in the activity of enzymes (e.g., hemicellulases) in the multi-enzyme composition. An accessory enzyme or enzyme mix may be composed of enzymes from (1) commercial suppliers; (2) cloned genes expressing enzymes; (3) complex broth (such as that resulting from growth of a microbial strain in media, wherein the strains secrete proteins and enzymes into the media); (4) cell lysates of strains grown as in (3); and, (5)

plant material expressing enzymes capable of degrading lignocellulose. In some embodiments, the accessory enzyme is a glucoamylase, a pectinase, or a ligninase.

[0192] As used herein, a ligninase is an enzyme that can hydrolyze or break down the structure of lignin polymers, including lignin peroxidases, manganese peroxidases, laccases, and other enzymes described in the art known to depolymerize or otherwise break lignin polymers. Also included are enzymes capable of hydrolyzing bonds formed between hemicellulosic sugars (notably arabinose) and lignin.

[0193] In one embodiment, the multi-enzyme composition comprises the enzyme CBH Ia (SEQ ID NO:2), a biologically active (e.g., has enzyme activity) homologue thereof, or a fragment comprising, consisting essentially of, or consisting of the CD thereof. In another embodiment, the multi-enzyme composition comprises the enzyme CBH IIb (SEQ ID NO:11), a biologically active homologue thereof, or a fragment comprising, consisting essentially of, or consisting of the CD thereof. In another embodiment, the multi-enzyme composition comprises the enzyme EG II (SEQ ID NO:94), a biologically active homologue thereof, or a fragment comprising, consisting essentially of, or consisting of the CD thereof. In a further embodiment, the multi-enzyme composition comprises the enzyme Xyl 2 (SEQ ID NO:29), a biologically active homologue thereof, or a fragment comprising, consisting essentially of, or consisting of the CD thereof. In yet another embodiment, the multi-enzyme composition comprises the enzyme β -glucosidase (SEQ ID NO:44), a biologically active homologue thereof, or a fragment comprising, consisting essentially of, or consisting of the CD thereof. In certain embodiments, the multi-enzyme composition comprises any combination of the enzymes listed above. In other embodiments, the multi-enzyme composition comprises at least one of the enzymes listed above in combination with a β -xylosidase enzyme.

[0194] In one embodiment, the multi-enzyme composition comprises CBH Ia (SEQ ID NO:2), CBH IIb (SEQ ID NO:11), EG II (SEQ ID NO:94), the CD of Xyl 2 (SEQ ID NO:29) and the CD of β -glucosidase (SEQ ID NO:44). This preferred enzyme combination may further comprise a β -xylosidase enzyme. One of skill in the art will appreciate, however, that any enzyme in the combinations described above may be replaced with an enzyme exhibiting a similar substrate specificity. Further, the optimal enzyme combination for a particular use (e.g., for degradation of a biomass derived from a specific source) can be determined by one of skill in the art using routine experimentation and assays known in the art, such as those disclosed herein.

[0195] In one embodiment, the multi-enzyme composition comprises the enzyme CBH Ib (SEQ ID NO:5), a biologically active homologue thereof, or a fragment comprising, consisting essentially of, or consisting of the CD thereof. In an additional embodiment, the multi-enzyme composition comprises the enzyme EG V (SEQ ID NO:20), a biologically active homologue thereof, or a fragment comprising, consisting essentially of, or consisting of the CD thereof. In still another embodiment, the multi-enzyme composition comprises the enzyme Xyl 1 (SEQ ID NO:26), a biologically active homologue thereof, or a fragment comprising, consisting essentially of, or consisting of the CD thereof. In further embodiments, the multi-enzyme composition comprises the enzyme arabinogalactanase (SEQ ID NO:47), a biologically active homologue thereof, or a fragment comprising, consisting essentially of, or consisting of the CD thereof. In certain embodiments, the multi-enzyme composition comprises any combination of the enzymes listed above. In other embodiments, the multi-enzyme composition comprises at least one of the enzymes listed above in combination with at least one accessory enzyme such as, for example, α -L-arabinofuranosidase.

[0196] One of skill in the art will appreciate, however, that any enzyme in the combinations described above may be replaced with an enzyme exhibiting a similar substrate specificity. Further, the optimal enzyme combination for a particular use (e.g., for degradation of a biomass derived from a specific source) can be determined by one of skill in the art using routine experimentation and assays known in the art, such as those disclosed herein.

[0197] The multi-enzyme compositions, in some embodiments, comprise a biomass comprising microorganisms or a crude fermentation product of microorganisms. A crude fermentation product refers to the fermentation broth which has been separated from the microorganism biomass (by filtration, for example). In general, the microorganisms are grown in fermentors, optionally centrifuged or filtered to remove biomass, and optionally concentrated, formulated, and dried to produce an enzyme(s) or a multi-enzyme composition that is a crude fermentation product. In other embodiments, enzyme(s) or multi-enzyme compositions produced by the microorganism (including a genetically modified microorganism as described below) are subjected to one or more purification steps, such as ammonium sulfate precipitation, chromatography, and/or ultrafiltration, which result in a partially purified or purified enzyme(s). If the microorganism has been genetically modified to express the enzyme(s), the enzyme(s) will include recombinant enzymes. If the genetically modified microorganism also naturally expresses the enzyme(s) or other enzymes useful for lignocellulosic saccharification, the enzyme(s) may include both naturally occurring and recombinant enzymes.

[0198] Another embodiment of the present invention relates to a composition comprising at least about 500 ng, and preferably at least about 1 μ g, and more preferably at least about 5 μ g, and more preferably at least about 10 μ g, and more preferably at least about 25 μ g, and more preferably at least about 50 μ g, and more preferably at least about 75 μ g, and more preferably at least about 100 μ g, and more preferably at least about 250 μ g, and more preferably at least about 500 μ g, and more preferably at least about 750 μ g, and more preferably at least about 1 mg, and more preferably at least about 5 mg, of an isolated protein of the invention. Such a composition of the present invention may include any

carrier with which the protein is associated by virtue of the protein preparation method, a protein purification method, or a preparation of the protein for use in any method according to the present invention. For example, such a carrier can include any suitable buffer, extract, or medium that is suitable for combining with the protein of the present invention so that the protein can be used in any method described herein according to the present invention.

[0199] In one embodiment of the invention, one or more enzymes of the invention is bound to a solid support, *i.e.*, an immobilized enzyme. As used herein, an immobilized enzyme includes immobilized isolated enzymes, immobilized microbial cells which contain the enzyme of the invention, other stabilized intact cells that produce the enzyme of the invention, and stabilized cell/membrane homogenates. Stabilized intact cells and stabilized cell/membrane homogenates include cells and homogenates from naturally occurring microorganisms expressing the enzyme of the invention and preferably, from genetically modified microorganisms as disclosed elsewhere herein. Thus, although methods for immobilizing enzymes are discussed below, it will be appreciated that such methods are equally applicable to immobilizing microbial cells and in such an embodiment, the cells can be lysed, if desired.

[0200] A variety of methods for immobilizing an enzyme are disclosed in Industrial Enzymology 2nd Ed., Godfrey, T. and West, S. Eds., Stockton Press, New York, N.Y., 1996, pp. 267-272; Immobilized Enzymes, Chibata, I. Ed., Halsted Press, New York, N.Y., 1978; Enzymes and Immobilized Cells in Biotechnology, Laskin, A. Ed., Benjamin/Cummings Publishing Co., Inc., Menlo Park, California, 1985; and Applied Biochemistry and Bioengineering, Vol. 4, Chibata, I. and Wingard, Jr., L. Eds, Academic Press, New York, N.Y., 1983.

[0201] Briefly, a solid support refers to any solid organic, biopolymer or inorganic supports that can form a bond with an enzyme without significantly effecting the activity of the enzyme. Exemplary organic solid supports include polymers such as polystyrene, nylon, phenol-formaldehyde resins, acrylic copolymers (*e.g.*, polyacrylamide), stabilized intact whole cells, and stabilized crude whole cell/membrane homogenates. Exemplary biopolymer supports include cellulose, polydextrans (*e.g.*, Sephadex®), agarose, collagen and chitin. Exemplary inorganic supports include glass beads (porous and nonporous), stainless steel, metal oxides (*e.g.*, porous ceramics such as ZrO₂, TiO₂, Al₂O₃, and NiO) and sand. In one embodiment, the solid support is selected from the group consisting of stabilized intact cells and/or crude cell homogenates (*e.g.*, produced from the microbial host cells expressing recombinant enzymes, alone or in combination with natural enzymes). Preparation of such supports requires a minimum of handling and cost. Additionally, such supports provide excellent stability of the enzyme.

[0202] Stabilized intact cells and/or cell/membrane homogenates can be produced, for example, by using bifunctional crosslinkers (*e.g.*, glutaraldehyde) to stabilize cells and cell homogenates. In both the intact cells and the cell membranes, the cell wall and membranes act as immobilizing supports. In such a system, integral membrane proteins are in the "best" lipid membrane environment. Whether starting with intact cells or homogenates, in this system the cells are either no longer "alive" or "metabolizing", or alternatively, are "resting" (*i.e.*, the cells maintain metabolic potential and active enzyme, but under the culture conditions are not growing); in either case, the immobilized cells or membranes serve as biocatalysts.

[0203] An enzyme of the invention can be bound to a solid support by a variety of methods including adsorption, cross-linking (including covalent bonding), and entrapment. Adsorption can be through van der Waal's forces, hydrogen bonding, ionic bonding, or hydrophobic binding. Exemplary solid supports for adsorption immobilization include polymeric adsorbents and ion-exchange resins. Solid supports in a bead form are particularly well-suited. The particle size of an adsorption solid support can be selected such that the immobilized enzyme is retained in the reactor by a mesh filter while the substrate is allowed to flow through the reactor at a desired rate. With porous particulate supports it is possible to control the adsorption process to allow enzymes or cells to be embedded within the cavity of the particle, thus providing protection without an unacceptable loss of activity.

[0204] Cross-linking of an enzyme to a solid support involves forming a chemical bond between a solid support and the enzyme. It will be appreciated that although cross-linking generally involves linking the enzyme to a solid support using an intermediary compound, it is also possible to achieve a covalent bonding between the enzyme and the solid support directly without the use of an intermediary compound. Cross-linking commonly uses a bifunctional or multifunctional reagent to activate and attach a carboxyl group, amino group, sulfur group, hydroxy group or other functional group of the enzyme to the solid support. The term "activate" refers to a chemical transformation of a functional group which allows a formation of a bond at the functional group. Exemplary amino group activating reagents include water-soluble carbodiimides, glutaraldehyde, cyanogen bromide, N-hydroxysuccinimide esters, triazines, cyanuric chloride, and carbonyl diimidazole. Exemplary carboxyl group activating reagents include water-soluble carbodiimides, and N-ethyl-5-phenylisoxazolium-3-sulfonate. Exemplary tyrosyl group activating reagents include diazonium compounds. And exemplary sulfhydryl group activating reagents include dithiobis-5,5'-(2-nitrobenzoic acid), and glutathione-2-pyridyl disulfide. Systems for covalently linking an enzyme directly to a solid support include Eupergit®, a polymethacrylate bead support available from Rohm Pharma (Darmstadt, Germany), kieselguhl (Macrosorbs), available from Sterling Organics, kaolinite available from English China Clay as "Biofix" supports, silica gels which can be activated by silanization, available from W.R. Grace, and high-density alumina, available from UOP (Des Plains, IL).

[0205] Entrapment can also be used to immobilize an enzyme. Entrapment of an enzyme involves formation of, *inter*

alia, gels (using organic or biological polymers), vesicles (including microencapsulation), semipermeable membranes or other matrices. Exemplary materials used for entrapment of an enzyme include collagen, gelatin, agar, cellulose triacetate, alginate, polyacrylamide, polystyrene, polyurethane, epoxy resins, carrageenan, and egg albumin. Some of the polymers, in particular cellulose triacetate, can be used to entrap the enzyme as they are spun into a fiber. Other materials such as polyacrylamide gels can be polymerized in solution to entrap the enzyme. Still other materials such as polyglycol oligomers that are functionalized with polymerizable vinyl end groups can entrap enzymes by forming a cross-linked polymer with UV light illumination in the presence of a photosensitizer.

[0206] Further embodiments of the present invention include nucleic acid molecules that encode a protein of the present invention. A nucleic acid molecule of the present disclosure includes a nucleic acid molecule comprising, consisting essentially of, or consisting of, a nucleic acid sequence encoding any of the isolated proteins disclosed herein, including a fragment or a homologue of such proteins, described above. Nucleic acid molecules can include a nucleic acid sequence that encodes a fragment of a protein that does not have biological activity, and can also include portions of a gene or polynucleotide encoding the protein that are not part of the coding region for the protein (e.g., introns or regulatory regions of a gene encoding the protein). Nucleic acid molecules can include a nucleic acid sequence that is useful as a probe or primer (oligonucleotide sequences).

[0207] In one embodiment, a nucleic acid molecule of the present invention includes a nucleic acid molecule comprising, consisting essentially of, or consisting of, a nucleic acid sequence represented by SEQ ID NO:31, or SEQ ID NO:33. In one embodiment, a nucleic acid molecule of the present invention includes a nucleic acid molecule comprising, consisting essentially of, or consisting of, a nucleic acid sequence encoding an amino acid sequence represented by SEQ ID NO:32 or homologues thereof. Preferably, the nucleic acid sequence encodes a protein (including homologues thereof) useful in the invention, or encompasses complementary nucleic acid sequences.

[0208] In accordance with the present invention, an isolated nucleic acid molecule is a nucleic acid molecule (polynucleotide) that has been removed from its natural milieu (*i.e.*, that has been subject to human manipulation) and can include DNA, RNA, or derivatives of either DNA or RNA, including cDNA. As such, "isolated" does not reflect the extent to which the nucleic acid molecule has been purified. Although the phrase "nucleic acid molecule" primarily refers to the physical nucleic acid molecule, and the phrase "nucleic acid sequence" primarily refers to the sequence of nucleotides on the nucleic acid molecule, the two phrases can be used interchangeably, especially with respect to a nucleic acid molecule, or a nucleic acid sequence, being capable of encoding a protein. An isolated nucleic acid molecule of the present invention can be isolated from its natural source or produced using recombinant DNA technology (e.g., polymerase chain reaction (PCR) amplification, cloning) or chemical synthesis. Isolated nucleic acid molecules can include, for example, genes, natural allelic variants of genes, coding regions or portions thereof, and coding and/or regulatory regions modified by nucleotide insertions, deletions, substitutions, and/or inversions in a manner such that the modifications do not substantially interfere with the nucleic acid molecule's ability to encode a protein of the present invention or to form stable hybrids under stringent conditions with natural gene isolates. An isolated nucleic acid molecule can include degeneracies. As used herein, nucleotide degeneracy refers to the phenomenon that one amino acid can be encoded by different nucleotide codons. Thus, the nucleic acid sequence of a nucleic acid molecule that encodes a protein of the present invention can vary due to degeneracies. It is noted that a nucleic acid molecule of the present disclosure is not required to encode a protein having protein activity. A nucleic acid molecule can encode a truncated, mutated or inactive protein, for example. In addition, nucleic acid molecules of the disclosure are useful as probes and primers for the identification, isolation and/or purification of other nucleic acid molecules. If the nucleic acid molecule is an oligonucleotide, such as a probe or primer, the oligonucleotide preferably ranges from about 5 to about 50 or about 500 nucleotides, more preferably from about 10 to about 40 nucleotides, and most preferably from about 15 to about 40 nucleotides in length.

[0209] As used herein, reference to a gene includes all nucleic acid sequences related to a natural (*i.e.* wild-type) gene, such as regulatory regions that control production of the protein encoded by that gene (such as, but not limited to, transcription, translation or post-translation control regions) as well as the coding region itself. In another embodiment, a gene can be a naturally occurring allelic variant that includes a similar but not identical sequence to the nucleic acid sequence encoding a given protein. Allelic variants have been previously described above. The phrases "nucleic acid molecule" and "gene" can be used interchangeably when the nucleic acid molecule comprises a gene as described above.

[0210] Preferably, an isolated nucleic acid molecule of the present invention is produced using recombinant DNA technology (e.g., polymerase chain reaction (PCR) amplification, cloning, etc.) or chemical synthesis. Isolated nucleic acid molecules include any nucleic acid molecules and homologues thereof that are part of a gene described herein and/or that encode a protein described herein, including, but not limited to, natural allelic variants and modified nucleic acid molecules (homologues) in which nucleotides have been inserted, deleted, substituted, and/or inverted in such a manner that such modifications provide the desired effect on protein biological activity or on the activity of the nucleic acid molecule. Allelic variants and protein homologues (e.g., proteins encoded by nucleic acid homologues) have been discussed in detail above.

[0211] A nucleic acid molecule homologue (*i.e.*, encoding a homologue of a protein of the present invention) can be

produced using a number of methods known to those skilled in the art (see, for example, Sambrook *et al.*). For example, nucleic acid molecules can be modified using a variety of techniques including, but not limited to, by classic mutagenesis and recombinant DNA techniques (e.g., site-directed mutagenesis, chemical treatment, restriction enzyme cleavage, ligation of nucleic acid fragments and/or PCR amplification), or synthesis of oligonucleotide mixtures and ligation of mixture groups to "build" a mixture of nucleic acid molecules and combinations thereof. Another method for modifying a recombinant nucleic acid molecule encoding a protein is gene shuffling (i.e., molecular breeding) (See, for example, U.S. Patent No. 5,605,793 to Stemmer; Minshull and Stemmer; 1999, *Curr. Opin. Chem. Biol.* 3:284-290; Stemmer, 1994, P.N.A.S. USA 91:10747-10751). This technique can be used to efficiently introduce multiple simultaneous changes in the protein. Nucleic acid molecule homologues can be selected by hybridization with a gene or polynucleotide, or by screening for the function of a protein encoded by a nucleic acid molecule (*i.e.*, biological activity).

[0212] The minimum size of a nucleic acid molecule of the present disclosure is a size sufficient to encode a protein (including a fragment or homologue of a full-length protein) having biological activity, sufficient to encode a protein comprising at least one epitope which binds to an antibody, or sufficient to form a probe or oligonucleotide primer that is capable of forming a stable hybrid with the complementary sequence of a nucleic acid molecule encoding a natural protein (e.g., under moderate, high, or high stringency conditions). As such, the size of the nucleic acid molecule encoding such a protein can be dependent on nucleic acid composition and percent homology or identity between the nucleic acid molecule and complementary sequence as well as upon hybridization conditions per se (e.g., temperature, salt concentration, and formamide concentration). The minimal size of a nucleic acid molecule that is used as an oligonucleotide primer or as a probe is typically at least about 12 to about 15 nucleotides in length if the nucleic acid molecules are GC-rich and at least about 15 to about 18 bases in length if they are AT-rich. There is no limit, other than a practical limit, on the maximal size of a nucleic acid molecule of the present invention, in that the nucleic acid molecule can include a portion of a protein encoding sequence, a nucleic acid sequence encoding a full-length protein (including a gene), including any length fragment between about 20 nucleotides and the number of nucleotides that make up the full length cDNA encoding a protein, in whole integers (e.g., 20, 21, 22, 23, 24, 25..... nucleotides), or multiple genes, or portions thereof.

[0213] The phrase "consisting essentially of", when used with reference to a nucleic acid sequence herein, refers to a nucleic acid sequence encoding a specified amino acid sequence that can be flanked by from at least one, and up to as many as about 60, additional heterologous nucleotides at each of the 5' and/or the 3' end of the nucleic acid sequence encoding the specified amino acid sequence. The heterologous nucleotides are not naturally found (*i.e.*, not found in nature, *in vivo*) flanking the nucleic acid sequence encoding the specified amino acid sequence as it occurs in the natural gene or do not encode a protein that imparts any additional function to the protein or changes the function of the protein having the specified amino acid sequence.

[0214] In one embodiment, the polynucleotide probes or primers of the disclosure are conjugated to detectable markers. Detectable labels suitable for use in the present disclosure include any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means. Useful labels in the present disclosure include biotin for staining with labeled streptavidin conjugate, magnetic beads (e.g., Dynabeads™), fluorescent dyes (e.g., fluorescein, texas red, rhodamine, green fluorescent protein, and the like), radiolabels (e.g., ³H, ¹²⁵I, ³⁵S, ¹⁴C, or ³²P), enzymes (e.g., horse radish peroxidase, alkaline phosphatase and others commonly used in an ELISA), and colorimetric labels such as colloidal gold or colored glass or plastic (e.g., polystyrene, polypropylene, latex, etc.) beads. Preferably, the polynucleotide probes are immobilized on a substrate such as: artificial membranes, organic supports, biopolymer supports and inorganic supports.

[0215] One embodiment of the present invention relates to a recombinant nucleic acid molecule which comprises the isolated nucleic acid molecule described above which is operatively linked to at least one expression control sequence. More particularly, according to the present invention, a recombinant nucleic acid molecule typically comprises a recombinant vector and any one or more of the isolated nucleic acid molecules as described herein. According to the present invention, a recombinant vector is an engineered (*i.e.*, artificially produced) nucleic acid molecule that is used as a tool for manipulating a nucleic acid sequence of choice and/or for introducing such a nucleic acid sequence into a host cell. The recombinant vector is therefore suitable for use in cloning, sequencing, and/or otherwise manipulating the nucleic acid sequence of choice, such as by expressing and/or delivering the nucleic acid sequence of choice into a host cell to form a recombinant cell. Such a vector typically contains heterologous nucleic acid sequences, that is, nucleic acid sequences that are not naturally found adjacent to nucleic acid sequence to be cloned or delivered, although the vector can also contain regulatory nucleic acid sequences (e.g., promoters, untranslated regions) which are naturally found adjacent to nucleic acid sequences of the present invention or which are useful for expression of the nucleic acid molecules of the present invention (discussed in detail below). The vector can be either RNA or DNA, either prokaryotic or eukaryotic, and typically is a plasmid. The vector can be maintained as an extrachromosomal element (e.g., a plasmid) or it can be integrated into the chromosome of a recombinant host cell, although it is preferred if the vector remain separate from the genome for most applications of the invention. The entire vector can remain in place within a host cell, or under certain conditions, the plasmid DNA can be deleted, leaving behind the nucleic acid molecule of the present

invention. An integrated nucleic acid molecule can be under chromosomal promoter control, under native or plasmid promoter control, or under a combination of several promoter controls. Single or multiple copies of the nucleic acid molecule can be integrated into the chromosome. A recombinant vector of the present invention can contain at least one selectable marker.

[0216] In one embodiment, a recombinant vector used in a recombinant nucleic acid molecule of the present invention is an expression vector. As used herein, the phrase "expression vector" is used to refer to a vector that is suitable for production of an encoded product (e.g., a protein of interest, such as an enzyme of the present invention). In this embodiment, a nucleic acid sequence encoding the product to be produced (e.g., the protein or homologue thereof) is inserted into the recombinant vector to produce a recombinant nucleic acid molecule. The nucleic acid sequence encoding the protein to be produced is inserted into the vector in a manner that operatively links the nucleic acid sequence to regulatory sequences in the vector which enable the transcription and translation of the nucleic acid sequence within the recombinant host cell.

[0217] Typically, a recombinant nucleic acid molecule includes at least one nucleic acid molecule of the present invention operatively linked to one or more expression control sequences (e.g., transcription control sequences or translation control sequences). As used herein, the phrase "recombinant molecule" or "recombinant nucleic acid molecule" primarily refers to a nucleic acid molecule or nucleic acid sequence operatively linked to a transcription control sequence, but can be used interchangeably with the phrase "nucleic acid molecule", when such nucleic acid molecule is a recombinant molecule as discussed herein. According to the present invention, the phrase "operatively linked" refers to linking a nucleic acid molecule to an expression control sequence in a manner such that the molecule is able to be expressed when transfected (i.e., transformed, transduced, transfected, conjugated or conducted) into a host cell. Transcription control sequences are sequences which control the initiation, elongation, or termination of transcription. Particularly important transcription control sequences are those which control transcription initiation, such as promoter, enhancer, operator and repressor sequences. Suitable transcription control sequences include any transcription control sequence that can function in a host cell or organism into which the recombinant nucleic acid molecule is to be introduced.

[0218] Recombinant nucleic acid molecules of the present invention can also contain additional regulatory sequences, such as translation regulatory sequences, origins of replication, and other regulatory sequences that are compatible with the recombinant cell. In one embodiment, a recombinant molecule of the present invention, including those which are integrated into the host cell chromosome, also contains secretory signals (i.e., signal segment nucleic acid sequences) to enable an expressed protein to be secreted from the cell that produces the protein. Suitable signal segments include a signal segment that is naturally associated with the protein to be expressed or any heterologous signal segment capable of directing the secretion of the protein according to the present invention. In another embodiment, a recombinant molecule of the present invention comprises a leader sequence to enable an expressed protein to be delivered to and inserted into the membrane of a host cell. Suitable leader sequences include a leader sequence that is naturally associated with the protein, or any heterologous leader sequence capable of directing the delivery and insertion of the protein to the membrane of a cell.

[0219] According to the present invention, the term "transfection" is generally used to refer to any method by which an exogenous nucleic acid molecule (i.e., a recombinant nucleic acid molecule) can be inserted into a cell. The term "transformation" can be used interchangeably with the term "transfection" when such term is used to refer to the introduction of nucleic acid molecules into microbial cells or plants and describes an inherited change due to the acquisition of exogenous nucleic acids by the microorganism that is essentially synonymous with the term "transfection." Transfection techniques include, but are not limited to, transformation, particle bombardment, electroporation, microinjection, lipofection, adsorption, infection and protoplast fusion.

[0220] One or more recombinant molecules of the present invention can be used to produce an encoded product (e.g., a protein) of the present invention. In one embodiment, an encoded product is produced by expressing a nucleic acid molecule as described herein under conditions effective to produce the protein. A preferred method to produce an encoded protein is by transfecting a host cell with one or more recombinant molecules to form a recombinant cell. Suitable host cells to transfect include, but are not limited to, any bacterial, fungal (e.g., filamentous fungi or yeast), plant, insect, or animal cell that can be transfected. Host cells can be either untransfected cells or cells that are already transfected with at least one other recombinant nucleic acid molecule.

[0221] Suitable cells (e.g., a host cell or production organism) include any microorganism (e.g., a bacterium, a protist, an alga, a fungus, or other microbe), and is preferably a bacterium, a yeast or a filamentous fungus. Suitable bacterial genera include, but are not limited to, *Escherichia*, *Bacillus*, *Lactobacillus*, *Pseudomonas* and *Streptomyces*. Suitable bacterial species include, but are not limited to, *Escherichia coli*, *Bacillus subtilis*, *Bacillus licheniformis*, *Lactobacillus brevis*, *Pseudomonas aeruginosa* and *Streptomyces lividans*. Suitable genera of yeast include, but are not limited to, *Saccharomyces*, *Schizosaccharomyces*, *Candida*, *Hansenula*, *Pichia*, *Kluyveromyces*, and *Phaffia*. Suitable yeast species include, but are not limited to, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Candida albicans*, *Hansenula polymorpha*, *Pichia pastoris*, *P. canadensis*, *Kluyveromyces marxianus* and *Phaffia rhodozyma*.

[0222] Suitable fungal genera include, but are not limited to, *Chrysosporium*, *Thielavia*, *Neurospora*, *Aureobasidium*,

Filibasidium, *Piromyces*, *Corynascus*, *Cryptococcus*, *Acremonium*, *Tolypocladium*, *Scytalidium*, *Schizophyllum*, *Sporotrichum*, *Penicillium*, *Gibberella*, *Myceliophthora*, *Mucor*, *Aspergillus*, *Fusarium*, *Humicola*, and *Trichoderma*, and anamorphs and teleomorphs thereof. Suitable fungal species include, but are not limited to, *Aspergillus niger*, *Aspergillus nidulans*, *Aspergillus japonicus*, *Absidia coerulea*, *Rhizopus oryzae*, *Chrysosporium lucknowense*, *Neurospora crassa*, *Neurospora intermedia*, *Trichoderma reesei*, *Penicillium canescens*, *Penicillium solitum*, *Penicillium funiculosum*, and *Talaromyces flavus*. In one embodiment, the host cell is a fungal cell of the species *Chrysosporium lucknowense*. In one embodiment, the host cell is a fungal cell of Strain C1 (VKM F-3500-D) or a mutant strain derived therefrom (e.g., UV13-6 (Accession No. VKM F-3632 D); NG7C-19 (Accession No. VKM F-3633 D); or UV18-25 (VKM F-3631D)). Host cells can be either untransfected cells or cells that are already transfected with at least one other recombinant nucleic acid molecule. Additional embodiments of the present invention include any of the genetically modified cells described herein.

[0223] In one embodiment, one or more protein(s) expressed by an isolated nucleic acid molecule of the present invention are produced by culturing a cell that expresses the protein (i.e., a recombinant cell or recombinant host cell) under conditions effective to produce the protein. In some instances, the protein may be recovered, and in others, the cell may be harvested in whole, either of which can be used in a composition.

[0224] Microorganisms used in the present invention (including recombinant host cells or genetically modified microorganisms) are cultured in an appropriate fermentation medium. An appropriate, or effective, fermentation medium refers to any medium in which a cell of the present invention, including a genetically modified microorganism (described below), when cultured, is capable of expressing enzymes useful in the present invention and/or of catalyzing the production of sugars from lignocellulosic biomass. Such a medium is typically an aqueous medium comprising assimilable carbon, nitrogen and phosphate sources. Such a medium can also include appropriate salts, minerals, metals and other nutrients. Microorganisms and other cells of the present invention can be cultured in conventional fermentation bioreactors. The microorganisms can be cultured by any fermentation process which includes, but is not limited to, batch, fed-batch, cell recycle, and continuous fermentation. The fermentation of microorganisms such as fungi may be carried out in any appropriate reactor, using methods known to those skilled in the art. For example, the fermentation may be carried out for a period of 1 to 14 days, or more preferably between about 3 and 10 days. The temperature of the medium is typically maintained between about 25 and 50°C, and more preferably between 28 and 40°C. The pH of the fermentation medium is regulated to a pH suitable for growth and protein production of the particular organism. The fermentor can be aerated in order to supply the oxygen necessary for fermentation and to avoid the excessive accumulation of carbon dioxide produced by fermentation. In addition, the aeration helps to control the temperature and the moisture of the culture medium. In general the fungal strains are grown in fermentors, optionally centrifuged or filtered to remove biomass, and optionally concentrated, formulated, and dried to produce an enzyme(s) or a multi-enzyme composition that is a crude fermentation product. Particularly suitable conditions for culturing filamentous fungi are described, for example, in U.S. Patent No. 6,015,707 and U.S. Patent No. 6,573,086, *supra*.

[0225] Depending on the vector and host system used for production, resultant proteins of the present invention may either remain within the recombinant cell; be secreted into the culture medium; be secreted into a space between two cellular membranes; or be retained on the outer surface of a cell membrane. The phrase "recovering the protein" refers to collecting the whole culture medium containing the protein and need not imply additional steps of separation or purification. Proteins produced according to the present invention can be purified using a variety of standard protein purification techniques, such as, but not limited to, affinity chromatography, ion exchange chromatography, filtration, electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and differential solubilization.

[0226] Proteins of the present invention are preferably retrieved, obtained, and/or used in "substantially pure" form. As used herein, "substantially pure" refers to a purity that allows for the effective use of the protein in any method according to the present invention. For a protein to be useful in any of the methods described herein or in any method utilizing enzymes of the types described herein according to the present invention, it is substantially free of contaminants, other proteins and/or chemicals that might interfere or that would interfere with its use in a method disclosed by the present invention (e.g., that might interfere with enzyme activity), or that at least would be undesirable for inclusion with a protein of the present invention (including homologues) when it is used in a method disclosed by the present invention (described in detail below). Preferably, a "substantially pure" protein, as referenced herein, is a protein that can be produced by any method (i.e., by direct purification from a natural source, recombinantly, or synthetically), and that has been purified from other protein components such that the protein comprises at least about 80% weight/weight of the total protein in a given composition (e.g., the protein of interest is about 80% of the protein in a solution/composition/buffer), and more preferably, at least about 85%, and more preferably at least about 90%, and more preferably at least about 91%, and more preferably at least about 92%, and more preferably at least about 93%, and more preferably at least about 94%, and more preferably at least about 95%, and more preferably at least about 96%, and more preferably at least about 97%, and more preferably at least about 98%, and more preferably at least about 99%, weight/weight of the total protein in a given composition.

[0227] It will be appreciated by one skilled in the art that use of recombinant DNA technologies can improve control of expression of transfected nucleic acid molecules by manipulating, for example, the number of copies of the nucleic acid molecules within the host cell, the efficiency with which those nucleic acid molecules are transcribed, the efficiency with which the resultant transcripts are translated, and the efficiency of post-translational modifications. Additionally, the promoter sequence might be genetically engineered to improve the level of expression as compared to the native promoter. Recombinant techniques useful for controlling the expression of nucleic acid molecules include, but are not limited to, integration of the nucleic acid molecules into one or more host cell chromosomes, addition of vector stability sequences to plasmids, substitutions or modifications of transcription control signals (e.g., promoters, operators, enhancers), substitutions or modifications of translational control signals (e.g., ribosome binding sites), modification of nucleic acid molecules to correspond to the codon usage of the host cell, and deletion of sequences that destabilize transcripts.

[0228] Another aspect of the present invention relates to a genetically modified microorganism that has been transfected with one or more nucleic acid molecules of the present invention. As used herein, a genetically modified microorganism can include a genetically modified bacterium, yeast, filamentous fungus, or other microbe. Such a genetically modified microorganism has a genome which is modified (i.e., mutated or changed) from its normal (i.e., wild-type or naturally occurring) form such that the desired result is achieved (i.e., increased or modified activity and/or production of at least one an enzyme or a multi-enzyme composition for the conversion of lignocellulosic material to fermentable sugars). Genetic modification of a microorganism can be accomplished using classical strain development and/or molecular genetic techniques. Such techniques known in the art and are generally disclosed for microorganisms, for example, in Sambrook et al., 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Labs Press or *Molecular Cloning: A Laboratory Manual*, third edition (Sambrook and Russel, 2001), (jointly referred to herein as "Sambrook"). The references of Sambrook, *ibid.* A genetically modified microorganism can include a microorganism in which nucleic acid molecules have been inserted, deleted or modified (i.e., mutated; e.g., by insertion, deletion, substitution, and/or inversion of nucleotides), in such a manner that such modifications provide the desired effect within the microorganism.

[0229] In one embodiment, a genetically modified microorganism can endogenously contain and express an enzyme or a multi-enzyme composition for the conversion of lignocellulosic material to fermentable sugars, and the genetic modification can be a genetic modification of one or more of such endogenous enzymes, whereby the modification has some effect on the ability of the microorganism to convert lignocellulosic material to fermentable sugars (e.g., increased expression of the protein by introduction of promoters or other expression control sequences, or modification of the coding region by homologous recombination to increase the activity of the encoded protein).

[0230] In another embodiment, a genetically modified microorganism can endogenously contain and express an enzyme for the conversion of lignocellulosic material to fermentable sugars, and the genetic modification can be an introduction of at least one exogenous nucleic acid sequence (e.g., a recombinant nucleic acid molecule), wherein the exogenous nucleic acid sequence encodes at least one additional enzyme useful for the conversion of lignocellulosic material to fermentable sugars and/or a protein that improves the efficiency of the enzyme for the conversion of lignocellulosic material to fermentable sugars. In this aspect of the invention, the microorganism can also have at least one modification to a gene or genes comprising its endogenous enzyme(s) for the conversion of lignocellulosic material to fermentable sugars.

[0231] In yet another embodiment, the genetically modified microorganism does not necessarily endogenously (naturally) contain an enzyme for the conversion of lignocellulosic material to fermentable sugars, but is genetically modified to introduce at least one recombinant nucleic acid molecule encoding at least one enzyme or a multiplicity of enzymes for the conversion of lignocellulosic material to fermentable sugars. Such a microorganism can be used in a method of the invention, or as a production microorganism for crude fermentation products, partially purified recombinant enzymes, and/or purified recombinant enzymes, any of which can then be used in a method of the present invention.

[0232] Once the proteins (enzymes) are expressed in a host cell, a cell extract that contains the activity to test can be generated. For example, a lysate from the host cell is produced, and the supernatant containing the activity is harvested and/or the activity can be isolated from the lysate. In the case of cells that secrete enzymes into the culture medium, the culture medium containing them can be harvested, and/or the activity can be purified from the culture medium. The extracts/activities prepared in this way can be tested using assays known in the art. Accordingly, methods to identify multi-enzyme compositions capable of degrading lignocellulosic biomass are provided.

[0233] Artificial substrates, or complex mixtures of polymeric carbohydrates and lignin, or actual lignocellulose can be used in such tests. One assay that may be used to measure the release of sugars and oligosaccharides from these complex substrates is the dinitrosalicylic acid assay (DNS). In this assay, the lignocellulosic material such as DDG is incubated with enzymes(s) for various times and reducing sugars are measured.

[0234] The disclosure also contemplates genetically modified plants transformed with one or more nucleic acid molecules of the invention. The plants may be used for production of the enzymes, and/or as the lignocellulosic material used as a substrate in the methods of the invention. Methods to generate recombinant plants are known in the art. For instance, numerous methods for plant transformation have been developed, including biological and physical transfor-

mation protocols. See, for example, Miki et al., "Procedures for Introducing Foreign DNA into Plants" in Methods in Plant Molecular Biology and Biotechnology, Glick, B.R. and Thompson, J.E. Eds. (CRC Press, Inc., Boca Raton, 1993) pp. 67-88. In addition, vectors and *in vitro* culture methods for plant cell or tissue transformation and regeneration of plants are available. See, for example, Gruber et al., "Vectors for Plant Transformation" in Methods in Plant Molecular Biology and Biotechnology, Glick, B.R. and Thompson, J.E. Eds. (CRC Press, Inc., Boca Raton, 1993) pp. 89-119.

[0235] The most widely utilized method for introducing an expression vector into plants is based on the natural transformation system of *Agrobacterium*. See, for example, Horsch et al., Science 227:1229 (1985). *A. tumefaciens* and *A. rhizogenes* are plant pathogenic soil bacteria which genetically transform plant cells. The Ti and Ri plasmids of *A. tumefaciens* and *A. rhizogenes*, respectively, carry genes responsible for genetic transformation of the plant. See, for example, Kado, C.I., Crit. Rev. Plant. Sci. 10:1 (1991). Descriptions of *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer are provided by numerous references, including Gruber et al., supra, Miki et al., supra, Moloney et al., Plant Cell Reports 8:238 (1989), and U.S. Patents Nos. 4,940,838 and 5,464,763.

[0236] Another generally applicable method of plant transformation is microprojectile-mediated transformation wherein DNA is carried on the surface of microprojectiles. The expression vector is introduced into plant tissues with a biolistic device that accelerates the microprojectiles to speeds sufficient to penetrate plant cell walls and membranes. Sanford et al., Part. Sci. Technol. 5:27 (1987), Sanford, J.C., Trends Biotech. 6:299 (1988), Sanford, J.C., Physiol. Plant 79:206 (1990), Klein et al., Biotechnology 10:268 (1992).

[0237] Another method for physical delivery of DNA to plants is sonication of target cells. Zhang et al., Bio/Technology 9:996 (1991). Alternatively, liposome or spheroplast fusion have been used to introduce expression vectors into plants. Deshayes et al., EMBO J., 4:2731 (1985), Christou et al., Proc Natl. Acad. Sci. USA 84:3962 (1987). Direct uptake of DNA into protoplasts using CaCl_2 precipitation, polyvinyl alcohol or poly-L-ornithine have also been reported. Hain et al., Mol. Gen. Genet. 199:161 (1985) and Draper et al., Plant Cell Physiol. 23:451 (1982). Electroporation of protoplasts and whole cells and tissues have also been described. Donn et al., In Abstracts of VIIth International Congress on Plant Cell and Tissue Culture IAPTC, A2-38, p. 53 (1990); D'Halluin et al., Plant Cell 4:1495-1505 (1992) and Spencer et al., Plant Mol. Biol. 24:51-61 (1994).

[0238] Another embodiment of the present invention relates to an isolated antibody or antigen binding fragment thereof capable of selectively binding to a protein of the present invention. The binding agent selectively binds to an amino acid sequence selected from SEQ ID NO:32.

[0239] According to the present invention, the phrase "selectively binds to" refers to the ability of an antibody, antigen binding fragment of the present invention to preferentially bind to specified proteins. More specifically, the phrase "selectively binds" refers to the specific binding of one protein to another (e.g., an antibody, fragment thereof, or binding partner to an antigen), wherein the level of binding, as measured by any standard assay (e.g., an immunoassay), is statistically significantly higher than the background control for the assay. For example, when performing an immunoassay, controls typically include a reaction well/tube that contain antibody or antigen binding fragment alone (i.e., in the absence of antigen), wherein an amount of reactivity (e.g., non-specific binding to the well) by the antibody or antigen binding fragment thereof in the absence of the antigen is considered to be background. Binding can be measured using a variety of methods standard in the art including enzyme immunoassays (e.g., ELISA), immunoblot assays, etc.).

[0240] Antibodies are characterized in that they comprise immunoglobulin domains and as such, they are members of the immunoglobulin superfamily of proteins. An antibody of the invention includes polyclonal and monoclonal antibodies, divalent and monovalent antibodies, bi- or multi-specific antibodies, serum containing such antibodies, antibodies that have been purified to varying degrees, and any functional equivalents of whole antibodies. Isolated antibodies of the present invention can include serum containing such antibodies, or antibodies that have been purified to varying degrees. Whole antibodies of the present invention can be polyclonal or monoclonal. Alternatively, functional equivalents of whole antibodies, such as antigen binding fragments in which one or more antibody domains are truncated or absent (e.g., Fv, Fab, Fab', or F(ab)₂ fragments), as well as genetically-engineered antibodies or antigen binding fragments thereof, including single chain antibodies or antibodies that can bind to more than one epitope (e.g., bi-specific antibodies), or antibodies that can bind to one or more different antigens (e.g., bi- or multi-specific antibodies), may also be employed in the invention. Methods for the generation and production of antibodies are well known in the art.

[0241] Monoclonal antibodies may be produced according to the methodology of Kohler and Milstein (Nature 256:495-497, 1975). Non-antibody polypeptides, sometimes referred to as binding partners, are designed to bind specifically to a protein of the invention. Examples of the design of such polypeptides, which possess a prescribed ligand specificity are given in Beste et al. (Proc. Natl. Acad. Sci. 96:1898-1903, 1999). In one embodiment, a binding agent of the invention is immobilized on a substrate such as: artificial membranes, organic supports, biopolymer supports and inorganic supports such as for use in a screening assay.

[0242] Proteins of the present invention, at least one protein of the present invention, compositions comprising such protein(s) of the present invention, and multi-enzyme compositions (examples of which are described above) may be used in any method where it is desirable to hydrolyze glycosidic linkages in lignocellulosic material, or any other method wherein enzymes of the same or similar function are useful.

[0243] In one embodiment, the present invention includes the use of at least the protein of the present invention, compositions comprising at least the protein of the present invention, or multi-enzyme compositions in methods for hydrolyzing lignocellulose and the generation of fermentable sugars therefrom. In one embodiment, the method comprises contacting the lignocellulosic material with an effective amount of one or more proteins of the present invention, composition comprising at least the protein of the present invention, or a multi-enzyme composition, whereby at least one fermentable sugar is produced (liberated). The lignocellulosic material may be partially or completely degraded to fermentable sugars. Economical levels of degradation at commercially viable costs are contemplated.

[0244] Typically, the amount of enzyme or enzyme composition contacted with the lignocellulose will depend upon the amount of glucan present in the lignocellulose. In some embodiments, the amount of enzyme or enzyme composition contacted with the lignocellulose may be from about 0.1 to about 200 mg enzyme or enzyme composition per gram of glucan; in other embodiments, from about 3 to about 20 mg enzyme or enzyme composition per gram of glucan. The disclosure encompasses the use of any suitable or sufficient amount of enzyme or enzyme composition between about 0.1 mg and about 200 mg enzyme per gram glucan, in increments of 0.05 mg (*i.e.*, 0.1 mg, 0.15 mg, 0.2 mg... 199.9 mg, 199.95 mg, 200 mg).

[0245] In a further embodiment, the invention provides a method for degrading DDG, preferably, but not limited to, DDG derived from corn, to sugars. The method comprises contacting the DDG with a protein of the present invention, a composition comprising at least the protein of the present invention, or a multi-enzyme composition of the present invention. In certain embodiments, at least 10% of fermentable sugars are liberated. In other embodiment, the at least 15% of the sugars are liberated, or at least 20% of the sugars are liberated, or at least 23% of the sugars are liberated, or at least 24% of the sugars are liberated, or at least 25% of the sugars are liberated, or at least 26% of the sugars are liberated, or at least 27% of the sugars are liberated, or at least 28% of the sugars are liberated.

[0246] In another embodiment, the invention provides a method for producing fermentable sugars comprising cultivating a genetically modified microorganism of the present invention in a nutrient medium comprising a lignocellulosic material, whereby fermentable sugars are produced.

[0247] Also provided are methods that comprise further contacting the lignocellulosic material with at least one accessory enzyme. Accessory enzymes have been described elsewhere herein. The accessory enzyme or enzymes may be added at the same time, prior to, or following the addition of a protein of the present invention, a composition comprising at least the protein of the present invention, or a multi-enzyme composition of the present invention, or can be expressed (endogenously or overexpressed) in a genetically modified microorganism used in a method of the invention. When added simultaneously, the protein of the present invention, a composition comprising at least the protein of the present invention, or a multi-enzyme composition of the present invention will be compatible with the accessory enzymes selected. When the enzymes are added following the treatment with the protein of the present invention, a composition comprising at least the protein of the present invention, or a multi-enzyme composition of the present invention, the conditions (such as temperature and pH) may be altered to those optimal for the accessory enzyme before, during, or after addition of the accessory enzyme. Multiple rounds enzyme addition are also encompassed. The accessory enzyme may also be present in the lignocellulosic material itself as a result of genetically modifying the plant. The nutrient medium used in a fermentation can also comprise one or more accessory enzymes.

[0248] In some embodiments, the method comprises a pretreatment process. In general, a pretreatment process will result in components of the lignocellulose being more accessible for downstream applications or so that it is more digestible by enzymes following treatment in the absence of hydrolysis. The pretreatment can be a chemical, physical or biological pretreatment. The lignocellulose may have been previously treated to release some or all of the sugars, as in the case of DDG. Physical treatments, such as grinding, boiling, freezing, milling, vacuum infiltration, and the like may also be used with the methods of the invention. In one embodiment, the heat treatment comprises heating the lignocellulosic material to 121°C for 15 minutes. A physical treatment such as milling can allow a higher concentration of lignocellulose to be used in the methods of the invention. A higher concentration refers to about 20%, up to about 25%, up to about 30%, up to about 35%, up to about 40%, up to about 45%, or up to about 50% lignocellulose. The lignocellulose may also be contacted with a metal ion, ultraviolet light, ozone, and the like. Additional pretreatment processes are known to those skilled in the art, and can include, for example, organosolv treatment, steam explosion treatment, lime impregnation with steam explosion treatment, hydrogen peroxide treatment, hydrogen peroxide/ozone (peroxone) treatment, acid treatment, dilute acid treatment, and base treatment, including ammonia fiber explosion (AFEX) technology. Details on pretreatment technologies and processes can be found in Wyman et al., *Bioresource Tech.* 96:1959 (2005); Wyman et al., *Bioresource Tech.* 96:2026(2005); Hsu, "Pretreatment of biomass" In *Handbook on Bioethanol: Production and Utilization*, Wyman, Taylor and Francis Eds., p. 179-212 (1996); and Mosier et al., *Bioresource Tech.* 96:673 (2005).

[0249] In an additional embodiment, the method comprises detoxifying the lignocellulosic material. Dextoxification may be desirable in the event that inhibitors are present in the lignocellulosic material. Such inhibitors can be generated by a pretreatment process, deriving from sugar degradation or are direct released from the lignocellulose polymer. Detoxifying can include the reduction of their formation by adjusting sugar extraction conditions; the use of inhibitor-tolerant or inhibitor-degrading strains of microorganisms. Detoxifying can also be accomplished by the addition of ion

exchange resins, active charcoal, enzymatic detoxification using, e.g., laccase, and the like. In some embodiments, the proteins, compositions or products of the present invention further comprises detoxifying agents.

[0250] In some embodiments, the methods may be performed one or more times in whole or in part. That is, one may perform one or more pretreatments, followed by one or more reactions with a protein of the present invention, composition or product of the present invention and/or accessory enzyme. The enzymes may be added in a single dose, or may be added in a series of small doses. Further, the entire process may be repeated one or more times as necessary. Therefore, one or more additional treatments with heat and enzymes are contemplated.

[0251] The methods described above result in the production of fermentable sugars. During, or subsequent to the methods described, the fermentable sugars may be recovered. In the case of a cultivation of microorganisms, the sugars can be recovered through a continuous, batch or fed-batch method. The sugars recovered can be concentrated or purified. Recovery may occur by any method known in the art, including, but not limited to, washing, gravity flow, pressure, chromatography, extraction, crystallization (e.g., evaporative crystallization), membrane separation, reverse osmosis, distillation, and filtration. The sugars can be subjected further processing; e.g., they can also be sterilized, for example, by filtration.

[0252] In a related embodiment, the disclosure provides means for for improving quality of lignocellulosic material, including DDG for for animal nutrition. In one embodiment, the treated lignocellulosic material (e.g., a lignocellulosic material which has been saccharified) is recovered (e.g., has the soluble sugars removed). The recovered material can be used as an animal feed additive. It is believed that the recovered material will have beneficial properties for animal nutrition, possibly due to a higher protein content. In some embodiments, the amount of enzyme or enzyme composition contacted with the lignocellulosic material may be from about 0.0001 % to about 1.0 % of the weight of the lignocellulosic material; in other embodiments, from about 0.005 % to about 0.1 % of the weight of the lignocellulosic material. The disclosure includes the use of any amount of enzyme or enzyme composition between about 0.0001 % and about 1.0 %, in increments of 0.0001 (*i.e.*, 0.0001, 0.0002, 0.0003...etc.).

[0253] In an additional embodiment, the invention provides a method for producing an organic substance, comprising saccharifying a lignocellulosic material with an effective amount of a protein of the present invention or a composition comprising at least the protein of the present invention, fermenting the saccharified lignocellulosic material obtained with one or more fermentating microorganisms, and recovering the organic substance from the fermentation. Sugars released from biomass can be converted to useful fermentation products including but not limited to amino acids, vitamins, pharmaceuticals, animal feed supplements, specialty chemicals, chemical feedstocks, plastics, solvents, fuels, or other organic polymers, lactic acid, and ethanol, including fuel ethanol. Specific products that may be produced by the methods of the invention include, but not limited to, biofuels (including ethanol); lactic acid; plastics; specialty chemicals; organic acids, including citric acid, succinic acid and maleic acid; solvents; animal feed supplements; pharmaceuticals; vitamins; amino acids, such as lysine, methionine, tryptophan, threonine, and aspartic acid; industrial enzymes, such as proteases, cellulases, amylases, glucanases, lactases, lipases, lyases, oxidoreductases, and transferases; and chemical feedstocks. The methods of the invention are also useful to generate feedstocks for fermentation by fermenting microorganisms. In one embodiment, the method further comprises the addition of at least one fermenting organism. As used herein, "fermenting organism" refers to an organism capable of fermentation, such as bacteria and fungi, including yeast. Such feedstocks have additional nutritive value above the nutritive value provided by the liberated sugars.

[0254] The protein of the present invention and compositions comprising at least the protein of the present invention are also useful in a variety of other applications involving the hydrolysis of glycosidic linkages in lignocellulosic material, such as stone washing, color brightening, depilling and fabric softening, as well as other applications well known in the art. The proteins of the present invention and compositions comprising at least the protein of the present invention are also readily amenable to use as additives in detergent and other media used for such applications. These and other methods of use will readily suggest themselves to those of skill in the art based on the invention described herein.

[0255] In one embodiment of this invention, the protein and compositions of the present invention can be used in stone washing procedures for fabrics or other textiles. In some embodiments, the protein and compositions can be used in stone washing procedures for denim jeans. By way of example, the method for stone washing the fabric comprises contacting the fabric with a protein or composition of the present invention. In an additional embodiment, the protein or composition of the present invention is included in a detergent composition, as described below. A preferred pH range of stone wash applications is between about 5.5 to 7.5, most preferably at about pH 6 to about 7. One of skill in the art will know how to regulate the amount or concentration of the protein or composition produced by this invention based on such factors as the activity of the enzyme and the wash conditions, including but not limited to temperature and pH. Examples of these uses can be found in U.S. Patent Application Publication No. 2003/0157595.

[0256] In yet another embodiment of this disclosure, the cellulase compositions of this invention can be used to reduce or eliminate the harshness associated with a fabric or textile by contacting the fabric or textile with a protein or composition of the present invention. In some embodiments, the fabric or textile may be made from cellulose or cotton. By way of example, a preferred range for reducing or eliminating the harshness associated with a fabric or textile is between about pH 8 to about 12, or between about pH 10 to about 11.

[0257] The protein or compositions of the subject invention can be used in detergent compositions. In one embodiment, the detergent composition may comprise at least one protein or composition of the present invention and one or more surfactants. The detergent compositions may also include any additional detergent ingredient known in the art. Detergent ingredients contemplated for use with the detergent compositions of the subject invention include, but are not limited to, detergents, buffers, surfactants, bleaching agents, softeners, solvents, solid forming agents, abrasives, alkalis, inorganic electrolytes, cellulase activators, antioxidants, builders, silicates, preservatives, and stabilizers. The detergent compositions of this invention preferably employ a surface active agent, i.e., surfactant, including anionic, non-ionic, and ampholytic surfactants well known for their use in detergent compositions. In addition to the protein or composition of the present invention and the surface active agent, the detergent compositions of this invention can additionally contain one or more of the following components: the enzymes amylases, cellulases, proteinase, lipases, oxido-reductases, peroxidases and other enzymes; cationic surfactants and long-chain fatty acids; builders; antiredeposition agents; bleaching agents; bluing agents and fluorescent dyes; caking inhibitors; masking agents for factors inhibiting the cellulase activity; cellulase activators; antioxidants; and solubilizers. In addition, perfumes, preservatives, dyes, and the like can be used, if desired, with the detergent compositions of this invention. Examples of detergent compositions employing cellulases are exemplified in U.S. Pat. Nos. 4,435,307; 4,443,355; 4,661,289; 4,479,881; 5,120,463.

[0258] When a detergent base used in the present invention is in the form of a powder, it may be one which is prepared by any known preparation method including a spray-drying method and/or a granulation method. The granulation method are the most preferred because of the non-dusting nature of granules compared to spray dry products. The detergent base obtained by the spray-drying method is hollow granules which are obtained by spraying an aqueous slurry of heat-resistant ingredients, such as surface active agents and builders, into a hot space. The granules have a size of from about 50 to about 2000 micrometers. After the spray-drying, perfumes, enzymes, bleaching agents, and/or inorganic alkaline builders may be added. With a highly dense, granular detergent base obtained by such as the spray-drying-granulation method, various ingredients may also be added after the preparation of the base. When the detergent base is a liquid, it may be either a homogenous solution or an inhomogeneous solution.

[0259] Other textile applications in which proteins and compositions of the present disclosure may be used include, but are not limited to, garment dyeing applications such as enzymatic mercerizing of viscose, bio-polishing applications, enzymatic surface polishing; biowash (washing or washing down treatment of textile materials), enzymatic microfibrillation, enzymatic "cottonization" of linen, ramie and hemp; and treatment of Lyocel or Newcell (i.e., "TENCEL" from Courtauld's), Cupro and other cellulosic fibers or garments, dye removal from dyed cellulosic substrates such as dyed cotton (Leisola & Linko--(1976) Analytical Biochemistry, v. 70, p. 592. Determination Of The Solubilizing Activity Of A Cellulase Complex With Dyed Substrates; Blum & Stahl--Enzymic Degradation Of Cellulose Fibers; Reports of the Shizuoka Prefectural Hamamatsu Textile Industrial Research Institute No. 24 (1985)), as a bleaching agent to make new indigo dyed denim look old (Fujikawa--Japanese Patent Application Kokai No. 50-132269), to enhance the bleaching action of bleaching agents (Suzuki--Great Britain Patent No. 2 094 826), and in a process for compositions for enzymatic desizing and bleaching of textiles (Windbichtler et al., U.S. Pat. No. 2,974,001. Another example of enzymatic desizing using cellulases is provided in Bhatawadekar (May 1983) Journal of the Textile Association, pages 83-86.

[0260] The amount of enzyme or enzyme composition contacted with a textile may vary with the particular application. Typically, for biofinishing and denim washing applications, from about 0.02 wt. % to about 5 wt. % of an enzyme or enzyme composition may be contacted with the textile. In some embodiments, from about 0.5 wt. % to about 2 wt. % of an enzyme or enzyme composition may be contacted with the textile. For bioscouring, from about 0.1 to about 10, or from about 0.1 to about 1.0 grams of an enzyme or enzyme composition per kilogram of textile may be used, including any amount between about 0.1 grams and about 10 grams, in increments of 0.1 grams.

[0261] In other embodiments, the protein or compositions of the present invention can be used in the saccharification of lignocellulose biomass from agriculture, forest products, municipal solid waste, and other sources, for biobleaching of wood pulp, and for de-inking of recycled print paper all by methods known to one skilled in the art.

[0262] The amount of enzyme or enzyme composition used for pulp and paper modification (e.g., biobleaching of wood pulp, de-inking of paper, or biorefining of pulp for paper making) typically varies depending upon the stock that is used, the pH and temperature of the system, and the retention time. In certain embodiments, the amount of enzyme or enzyme composition contacted with the paper or pulp may be from about 0.01 to about 50 U; from about 0.1 to about 15 U; or from about 0.1 to about 5 U of enzyme or enzyme composition per dry gram of fiber, including any amount between about 0.01 and about 50 U, in 0.01 U increments. In other embodiments, the amount of enzyme or enzyme composition contacted with the paper or pulp may be from about 1 to about 2000 grams or from about 100 to about 500 grams enzyme or enzyme composition per dry ton of pulp, including any amount between about 1 and about 2000 grams, in 1 gram increments.

[0263] Proteins or compositions of the present disclosure can added to wastewater to reduce the amount of solids such as sludge or to increase total biochemical oxygen demand (BOD) and chemical oxygen demand (COD) removal. For example, proteins or compositions of the present invention can be used to transform particulate COD to soluble COD in wastewater produced from grain/fruit/cellulose industrial processes or to increase the BOD/COD ratio by in-

creasing waste biodegradability (soluble lower molecular weight polymers in cellulosic/hemicellulosic wastes are typically more readily biodegradable than non-soluble material). In biological wastewater treatment systems, proteins or compositions of the present invention can also be used to increase waste digestion by aerobic and/or anaerobic bacteria.

[0264] Chitinases of the present disclosure (e.g., SEQ ID NOS: 56 and 59) can hydrolyze the P-1,4-glycosidic linkage present in chitin and thus may be used to degrade chitin-containing materials. Examples of chitin-containing materials include fungal cell walls, insect exoskeletons, the eggs of parasitic worms, and crustacean shells.

[0265] Chitinases may be used to inhibit or reduce fungal growth, including the treatment of fungal infections such as those caused by nail fungi. For example, chitinases of the present disclosure may be applied to any fungus or area susceptible to fungal growth. Chitinases may also be used to coat or treat seeds and flower bulbs to prevent the growth of fungi. Further, chitinases may be added to fungal cultures to lower culture viscosity by increasing cell wall degradation. Chitinases may also be used as lysing enzymes for the generation of protoplasts from fungi (see, e.g., Yano et al., *Biosci Biotechnol Biochem.* 70:1754 (2006)).

[0266] Chitinases or compositions containing chitinases may be used as a biological control agent such as an insecticide (see, e.g., Kramer et al., *Insect Biochem Mol Biol.* 27:887 (1997)). Chitinases of the present disclosure have been shown to be effective for controlling white-fly larvae in laboratory tests. Thus, chitinases may be applied to crops, plants and the like to control insect infestations.

[0267] Chitin has also been suggested to play a role in inducing allergic inflammation and asthma (see Reese et al., *Nature* 447:92 (2007)). Accordingly, chitinases of the present disclosure may be administered to a subject to reduce allergic inflammatory responses induced by chitin or to reduce the symptoms of asthma.

[0268] Exemplary methods according to the invention and disclosure are presented below. Examples of the methods described above may also be found in the following references: *Trichoderma & Gliocladium*. Volume 2. Enzymes, biological control and commercial applications. Editors: Gary E. Harman, Christian P. Kubicek. Taylor & Francis Ltd 1998, 393 (in particular, chapters 14, 15 and 16) Helmut Uhlig. *Industrial enzymes and their applications*. Translated and updated by Elfriede M. Linsmaier-Bednar. John Wiley & Sons, Inc 1998, p. 454 (in particular, chapters 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.9, 5.10, 5.11, and 5.13) For saccharification applications: Hahn-Hagerdal, B., Galbe, M., Gorwa-Grauslund, M.F. Lidén, Zacchi, G. *Bio-ethanol - the fuel of tomorrow from the residues of today*. *Trends in Biotechnology*. 2006, 24 (12), 549-556; Mielenz, J.R. *Ethanol production from biomass: technology and commercialization status*. *Current Opinion in Microbiology*. 2001, 4, 324-329; Himmel, M.E., Ruth, M.F., Wyman, C.E. *Cellulase for commodity products from cellulosic biomass*. *Current Opinion in Biotechnology*. 1999, 10, 358-364; Sheehan, J., Himmel, M. *Enzymes, energy, and the environment: a strategic perspective on the U.S. Department of Energy's Research and Development Activities for Bioethanol*. *Biotechnology Progress*. 1999, 15, 817-827. For textile processing applications: Galante, Y.M., Formantici, C. *Enzyme applications in detergency and in manufacturing industries*. *Current Organic Chemistry*. 2003, 7, 1399-1422. For pulp and paper applications: Bajpai, P., Bajpai, P.K. *Deinking with enzymes: a review*. *TAPPI Journal*. 1998, 81(12), 111-117; Viikari, L., Pere, J., Suurnäkki, A., Oksanen, T., Buchert, J. *Use of cellulases in pulp and paper applications*. In: "Carbohydrates from *Trichoderma reesei* and other microorganisms. Structure, Biochemistry, Genetics and Applications." Editors: Mark Claessens, Wim Nerinckx, and Kathleen Piens. The Royal Society of Chemistry 1998, 245-254. For food and beverage applications: Roller, S., Dea, I.C.M. *Biotechnology in the production and modification of biopolymers for foods*. *Critical Reviews in Biotechnology*. 1992, 12(3), 261-277.

[0269] The following examples are provided for the purpose of illustration and are not intended to limit the scope of the present invention.

EXAMPLES

Example 1

[0270] The following example illustrates the Somogyi-Nelson assay used to measure beta-glucanase enzymatic activity.

[0271] This assay measures the release of reducing sugars (as glucose equivalents) by the action of β -glucanase on a soluble β -glucan. One β -glucanase unit of activity is the amount of enzyme that liberates 1 micromole of reducing sugars, expressed as glucose equivalents, in one minute at 50° C and pH 5.0.

Reagents

[0272] Acetate buffer (0.05 M, pH 5.0) is prepared as follows. 4.1 g of anhydrous sodium acetate or 6.8 g of sodium acetate $\cdot$ 3H₂O is dissolved in distilled water so that the final volume of the solution is 1000 mL (Solution A). In a separate flask, 3.0 g (2.86 mL) of glacial acetic acid is mixed with distilled water to make a total volume of 1000 mL (Solution B). The final 0.05 M acetate buffer, pH 5.0, is prepared by mixing Solution A with Solution B until the pH of the resulting solution is equal to 5.0.

EP 2 183 363 B1

[0273] β -glucan from barley (medium viscosity, Megazyme, P-BGBM) is used as the assay substrate. The substrate is milled in a mortar before dissolving. 1% w/v stock solution is prepared as follows. 500 mg of β -glucan is mixed with 50 mL of distilled water and agitated on a magnetic stirrer for 1 hour. After that, the suspension is placed into a Bandelin SONOREX TK52 ultrasonic bath for 10 minutes to destroy undissolved particles and then into a boiling water bath until the solution becomes clear. Finally, the solution is stirred on a magnetic stirrer for 30 minutes. The solution is typically stable for 2 days.

[0274] The Somogyi reagent is prepared as follows. 24 g of anhydrous sodium carbonate and 12 g of sodium-potassium tartrate tetrahydrate are dissolved in 250 mL of distilled water. A solution of copper sulfate pentahydrate (4 g in 40 mL of distilled water) is added on stirring, and then 16 g of sodium bicarbonate is dissolved to obtain Solution A. In a separate flask, 180 g of sodium sulfate is dissolved in 500 mL of hot ($\sim 80^\circ\text{C}$) distilled water and boiled for 5 minutes (Solution B). Solution A is mixed with Solution B and the final volume is adjusted to 1000 mL. The reagent is stable for 2-3 months on storing in a dark glass vessel.

[0275] The Nelson reagent is prepared as follows. 50 g of ammonium molybdate is dissolved in 900 mL of hot ($\sim 60^\circ\text{C}$) distilled water. The solution is cooled to $5\text{--}10^\circ\text{C}$ and 42 g of concentrated sulfuric acid containing 6 g of sodium arsenate are carefully added on stirring. The volume of the mixture is adjusted to 1000 mL. The flask is incubated at 40°C for 48 hours and then the solution is filtered if necessary. The reagent is stable for 2-3 months.

[0276] Using the above reagents, the assay is performed as detailed below.

Reagent Blank

[0277] 0.5 mL of 0.05 M sodium acetate buffer is mixed in a graduated glass test tube (1.5 cm x 15 cm) with 0.5 mL of Somogyi reagent and incubated in a boiling water bath for 40 minutes. Then, the test tube is cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water.

Enzyme Sample

[0278] 0.25 mL of 1% w/v β -glucan stock solution is mixed with 0.15 mL of 0.05 M acetate buffer, pH 5.0, in graduated glass test tubes (1.5 cm x 15 cm) and preheated at 50°C for 5 minutes. Enzyme sample is suitably diluted by the 0.05 M sodium acetate buffer (the dilution of the enzyme sample is chosen so that A_{610} should be between 0.05 and 0.4 units of optical density). 0.1 mL of suitably diluted enzyme sample is added to 0.4 mL of the preheated substrate solution, mixed and incubated at 50°C for 10 minutes. After exactly 10 minutes of incubation, 0.5 mL of Somogyi reagent is added, and test tubes are placed into a boiling water bath for 40 minutes. Then, the test tubes are cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the reagent blank as A_s (enzyme sample).

Enzyme Blank

[0279] 0.4 mL of 0.05 M sodium acetate buffer is mixed in a graduated glass test tube (1.5 cm x 15 cm) with 0.1 mL of enzyme sample having the same dilution as above. 0.5 mL of Somogyi reagent is added and the test tube is incubated in a boiling water bath for 40 minutes. Then, the test tube is cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the reagent blank as A_{EB} (enzyme blank).

Substrate Blank

[0280] 0.25 mL of 1% w/v β -glucan stock solution is mixed with 0.25 mL of 0.05 M acetate buffer, pH 5.0, in a graduated glass test tube (1.5 cm x 15 cm), 0.5 mL of Somogyi reagent is added, and the test tube is placed into a boiling water bath for 40 minutes. Then, the test tube is cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the reagent blank as A_{SB} (substrate blank). The value of A_{SB} should not exceed 0.05 units of optical density.

[0281] All enzyme samples and enzyme and substrate blanks should be boiled together.

Standard Calibration Plot

[0282] Stock glucose solution (1 mg/mL) is prepared by dissolving 100 mg of D-glucose in 100 mL of 0.05 M acetate buffer, pH 5.0. Then, different dilutions of stock glucose solution are prepared as follows:

5

10

15

Stock solution (mL)	0.05 M acetate buffer (mL)	Glucose concentration (mg/mL)
0.05	4.95	0.01
0.10	4.90	0.02
0.15	4.85	0.03
0.20	4.80	0.04
0.25	4.75	0.05
0.30	4.70	0.06
0.35	4.65	0.07
0.40	4.60	0.08
0.45	4.55	0.09
0.50	4.50	0.10

[0283] Samples (0.5 mL) of glucose solution with different dilutions are placed to graduated glass test tubes (1.5 cm x 15 cm). 0.5 mL of Somogyi reagent is added and test tubes are placed into a boiling water bath for 40 minutes. Then, the test tubes are cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the reagent blank. The following typical A_{610} data obtained with different glucose dilutions are given below:

25

Glucose, mg/mL	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	0.10
A_{610}	0.11	0.20	0.28	0.41	0.48	0.60	0.72	0.79	0.93	1.06

Absorbance (A_{610}) on the y-axis versus glucose concentration (G, mg/mL) on the x-axis is plotted to generate a standard calibration plot such as that shown in the Figure 11. From Figure 11, the inverse of the slope ($1/a = 0.098$) is calculated.

30

Calculation of Activity

[0284] Activity is calculated as follows: activity (IU/mL) = $1/a * \Delta A_{610} * DF * 2.78$, where $1/a$ - inverse of the slope calculated from the standard calibration plot, $\Delta A_{610} = A_s$ (enzyme sample) - A_{EB} (enzyme blank) - A_{SB} (substrate blank), DF is the enzyme dilution factor, and 2.78 is the coefficient that accounts for additional 5-fold enzyme dilution in the reaction mixture (0.1 mL enzyme + 0.4 mL substrate), reaction time (10 minutes) and molecular weight of glucose (0.18016 mg/micromole), i.e. $5/(10*0.18016) = 2.78$

[0285] An example calculation is supplied below.

40

B4 UF concentrate	DF	ΔA_{610}
	20,000	0.545
	40,000	0.316
	80,000	0.154

45

$$\beta\text{-glucanase activity (IU/mL)} = 0.098 * 0.316 * 40000 * 2.78 = 3444$$

50

Example 2

[0286] The following example illustrates the assays used to measure beta-glucosidase, α -galactosidase and β -galactosidase enzymatic activities.

[0287] This assay measures the release of *p*-nitrophenol by the action of β -glucosidase on *p*-nitrophenyl β -D-glucopyranoside (PNPG). Modifications to the assay for measuring α -galactosidase and β -galactosidase enzymatic activities are discussed below. One β -glucosidase unit of activity is the amount of enzyme that liberates 1 micromole of *p*-nitrophenol in one minute at 40 °C and pH 5.0.

55

EP 2 183 363 B1

Reagents

[0288] Acetate buffer (0.1 M, pH 5.0) is prepared as follows. 8.2 g of anhydrous sodium acetate or 13.6 g of sodium acetate * 3H₂O is dissolved in distilled water so that the final volume of the solution to be 1000 mL (Solution A). In a separate flask, 6.0 g (5.72 mL) of glacial acetic acid is mixed with distilled water to make the total volume of 1000 mL (Solution B). The final 0.1 M acetate buffer, pH 5.0, is prepared by mixing Solution A with Solution B until the pH of the resulting solution is equal to 5.0.

[0289] PNPG from Sigma (USA) is used as the assay substrate. 6 mg of PNPG is dissolved in 2 mL of distilled water using magnetic stirrer to obtain 10 mM stock solution. The solution is stable for 2 days on storage at 4 °C.

[0290] The stop reagent (1 M sodium carbonate solution) is prepared as follows. 10.6 g of anhydrous sodium carbonate is dissolved in 80 mL of distilled water, and the solution volume is adjusted to 100 mL. This reagent is used to terminate the enzymatic reaction.

[0291] Using the above reagents, the assay is performed as detailed below.

Enzyme Sample

[0292] 0.09 mL of 10 mM PNPG stock solution is mixed with 0.81 mL of 0.1 M acetate buffer, pH 5.0, and preheated at 40° C for 5 minutes. The enzyme sample is suitably diluted with the 0.1 M sodium acetate buffer (the dilution of the enzyme sample is chosen so that A₄₀₀ should be between 0.05 and 0.35 units of optical density). 0.1 mL of suitably diluted enzyme sample is added to 0.9 mL of the preheated substrate solution, mixed and incubated at 40 °C for 10 minutes. After exactly 10 minutes of incubation, 0.5 mL of 1 M sodium carbonate solution is added and then the absorbance at 400 nm (A₄₀₀) is measured as A_s (enzyme sample).

Substrate Blank

[0293] 0.09 mL of 10 mM PNPG stock solution is mixed with 0.91 mL of 0.1 M acetate buffer, pH 5.0. Then, 0.5 mL of 1 M sodium carbonate solution is added and the absorbance at 400 nm (A₄₀₀) is measured as A_{SB} (substrate blank).

Calculation of Activity

[0294] Activity is calculated as follows: activity (IU/mL) = ΔA₄₀₀ * DF * 0.082, where ΔA₄₀₀ = A_s (enzyme sample) - A_{SB} (substrate blank), DF is the enzyme dilution factor, and 0.082 is the coefficient obtained as follows:

$$0.082 = \frac{10^6}{18300 * 10 * (0.1/0.0015)}$$

where 10⁶ is factor used to convert moles/L in the extinction coefficient to micromoles/L, 18300 M⁻¹ cm⁻¹ is the extinction coefficient of *p*-nitrophenol released, 10 minutes is the reaction time, and 0.1 mL is the volume of enzyme solution added to 0.0015 L of total assay volume.

[0295] An example calculation is supplied below.

B4 UF concentrate	DF	ΔA ₄₀₀
	5,000	0.495
	10,000	0.268
	20,000	0.144

For

$$\text{DF}=10,000: \beta\text{-glucosidase activity (IU/mL)} = 0.268 * 10000 * 0.082 = 220$$

For

$$\text{DF}=20,000: \beta\text{-glucosidase activity (IU/mL)} = 0.144 * 20000 * 0.082 = 236$$

Finally, β -glucosidase activity = 228 IU/mL, as a mean value between two measurements.

α - and β -galactosidase assays

[0296] The assay procedures for α - and β -galactosidases are the same as the β -glucosidase assay described above, except that *p*-nitrophenyl α -D-galactopyranoside and *p*-nitrophenyl β -D-galactopyranoside are used as substrates, respectively, instead of *p*-nitrophenyl β -D-glucopyranoside.

Example 3

[0297] The following example illustrates the Somogyi-Nelson assay used to measure CMCase enzymatic activity.

[0298] This assay measures the release of reducing sugars (as glucose equivalents) by the action of cellulase on a soluble cellulosic substrate (CMC). One CMCase unit of activity is the amount of enzyme that liberates 1 micromole of reducing sugars, expressed as glucose equivalents, in one minute at 50° C, pH 5.0, 0.05 M sodium acetate buffer.

Reagents

[0299] Acetate buffer (0.05 M, pH 5.0) is prepared as follows. 4.1 g of anhydrous sodium acetate or 6.8 g of sodium acetate $\cdot$ 3H₂O is dissolved in distilled water so that the final volume of the solution to be 1000 mL (Solution A). In a separate flask, 3.0 g (2.86 mL) of glacial acetic acid is mixed with distilled water to make the total volume of 1000 mL (Solution B). The final 0.05 M acetate buffer, pH 5.0, is prepared by mixing Solution A with Solution B until the pH of the resulting solution is equal to 5.0.

[0300] The sodium salt of carboxymethyl cellulose (CMC, medium viscosity, Sigma, C 4888) is used as the substrate. A 1% w/v stock solution is prepared in 0.05 M acetate buffer, pH 5.0, by adding gradually 1.0 g of CMC to 99 mL of the buffer agitated on a magnetic stirrer. The mixture is stirred until the CMC is completely dissolved. The solution is stable for 2 days at 4° C.

[0301] The Somogyi reagent is prepared as follows. 24 g of anhydrous sodium carbonate and 12 g of sodium-potassium tartrate tetrahydrate are dissolved in 250 mL of distilled water. The solution of copper sulfate pentahydrate (4 g in 40 mL of distilled water) is added on stirring, and then 16 g of sodium bicarbonate is dissolved to obtain Solution A. In a separate flask, 18.0 g of sodium sulfate is dissolved in 500 mL of hot (~80° C) distilled water and boiled for 5 min (Solution B). Solution A is mixed with Solution B and the final volume is adjusted to 1000 mL. The reagent is stable for 2-3 months on storing in a dark glass vessel at room temperature.

[0302] The Nelson reagent is prepared as follows. 50 g of ammonium molybdate is dissolved in 900 mL of hot (~60° C) distilled water. The solution is cooled to 5-10° C and 42 g of concentrated sulfuric acid containing 6 g of sodium arsenate are carefully added on stirring. The volume of the mixture is adjusted to 1000 mL. The flask is incubated at 40° C for 48 hours and then the solution is filtered if necessary. The reagent is stable for 2-3 months at room temperature.

[0303] Using the above reagents, the assay is performed as detailed below.

Reagent Blank

[0304] 0.5 mL of 0.05 M sodium acetate buffer is mixed in a graduated glass test tube (1.5 cm x 15 cm) with 0.5 mL of Somogyi reagent and incubated in a boiling water bath for 40 minutes. Then, the test tube is cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water.

Enzyme Sample

[0305] 0.25 mL of 1% w/v CMC stock solution is preheated in graduated glass test tubes (1.5 cm x 15 cm) at 50° C for 5 minutes. Enzyme sample is suitably diluted by the 0.05 M sodium acetate buffer (the dilution of the enzyme sample is chosen so that A₆₁₀ should be between 0.05 and 0.35 units of optical density) and preheated at 50° C for 5 minutes. 0.25 mL of suitably diluted enzyme sample is added to 0.25 mL of preheated CMC stock solution, mixed and incubated at 50° C for 5 minutes. After exactly 5 minutes of incubation, 0.5 mL of Somogyi reagent is added, and test tubes are placed into a boiling water bath for 40 minutes. Then, the test tubes are cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, 1 mL of acetone is added to dissolve CMC precipitated during the boiling procedure and the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A₆₁₀) is measured against the reagent blank as A_s (enzyme sample).

EP 2 183 363 B1

Enzyme Blank

[0306] 0.25 mL of 0.05 M sodium acetate buffer is mixed in a graduated glass test tube (1.5 cm x 15 cm) with 0.25 mL of enzyme sample having the same dilution as above. 0.5 mL of Somogyi reagent is added and the test tube is incubated in a boiling water bath for 40 minutes. Then, the test tube is cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the reagent blank as A_B (enzyme blank).

[0307] All enzyme samples and enzyme blanks should be boiled together.

Standard Calibration Plot

[0308] A stock glucose solution (1 mg/mL) is prepared by dissolving 100 mg of D-glucose in 100 mL of 0.05 M acetate buffer, pH 5.0. Then, different dilutions of stock glucose solution are prepared as follows:

Stock solution (mL)	0.05 M acetate buffer (mL)	Glucose concentration (mg/mL)
0.05	4.95	0.01
0.10	4.90	0.02
0.15	4.85	0.03
0.20	4.80	0.04
0.25	4.75	0.05
0.30	4.70	0.06
0.35	4.65	0.07
0.40	4.60	0.08
0.45	4.55	0.09
0.50	4.50	0.10

[0309] Samples (0.5 mL) of glucose solution with different dilutions are placed to graduated glass test tubes (1.5 cm x 15 cm). 0.5 mL of Somogyi reagent is added and test tubes are placed into a boiling water bath for 40 minutes. Then, the test tubes are cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the reagent blank. The following typical A_{610} data obtained with different glucose dilutions are given below:

Glucose, mg/mL	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	0.10
A_{610}	0.11	0.20	0.28	0.41	0.48	0.60	0.72	0.79	0.93	1.06

Absorbance (A_{610}) on the y-axis versus glucose concentration (G, mg/mL) on the x-axis is plotted to generate a standard calibration plot such as that shown in the Figure 11. From Figure 11, the inverse of the slope ($1/a = 0.098$) is calculated.

Calculation of Activity

[0310] Activity is calculated as follows: $CMCase\ (IU/mL) = 1/a * \Delta A_{610} * DF * 2.22$, where $1/a$ - inverse of the slope calculated from the standard calibration plot, $\Delta A_{610} = A_s$ (enzyme sample) - A_B (enzyme blank), DF - enzyme dilution factor, and 2.22 is the coefficient that accounts for additional 2-fold enzyme dilution in the reaction mixture (0.25 mL enzyme + 0.25 mL substrate), reaction time (5 minutes) and molecular weight of glucose (0.18016 mg/micromole), i.e. $2/(5*0.18016) = 2.22$.

[0311] An example calculation is supplied below.

ACE sample	DF	ΔA_{610}
	40,000	0.41
	80,000	0.25

$$\text{CMCase activity (IU/mL)} = 0.098 \times 0.25 \times 80000 \times 2.22 = 4351$$

Example 4

[0312] The following example illustrates the Lowry protein assay.

[0313] The Lowry procedure is one of the most known and widely used protein assays, being first described in 1951 (Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) J. Biol. Chem. 193: 265-275). Under alkaline conditions, copper complexes with protein. When Folin's phenol reagent (phospho-molybdic-phosphotungstic reagent) is added, it binds to the protein. Bound reagent is slowly reduced and changes color from yellow to blue.

Stock Solutions and Reagents

[0314] Lowry A stock solution is 2% Na_2CO_3 in 0.1 M NaOH. Lowry B stock solution is 1% $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ in distilled water. Lowry C stock solution is 2% sodium potassium tartrate ($\text{NaKC}_4\text{H}_4\text{O}_6 \cdot 4\text{H}_2\text{O}$) in distilled water. Lowry stock reagent is prepared fresh daily by mixing 49 ml Lowry A, 0.5 ml Lowry B and 0.5 ml Lowry C. Commercial Folin's Reagent: Phenol reagent - 2N (Folin - Ciocalteu reagent) is diluted 1:1 in distilled water before use and stored in a refrigerator.

[0315] Bovine serum albumin (BSA) is used as the assay standard. When BSA is weighed out before use, the protein preparation used to prepare the standard should be salt-free to avoid inaccurate results. BSA is dissolved at a concentration of 1 mg/ml in a buffer similar to the solution for analysis. A series of standard dilutions is prepared where the protein (BSA) concentration is varied in the range of 0.05-0.5 mg/ml.

Assay Procedure

[0316] Using the above reagents, the assay is performed as follows. 100 μl of sample is added to each tube followed by 1.0 ml of Lowry stock reagent and the mixture is incubated 10 minutes at room temperature. 100 μl of Folin's reagent is added to each tube and the tubes are further incubated for 40 minutes at room temperature. An absorbance at 750 nm is read on a spectrophotometer against a blank (control) prepared in the same way as the sample except the buffer is added to the tube instead of the analyzed protein. Protein concentration is determined by reading the absorbance at 750 nm for each sample and plotting the result on a BSA calibration graph prepared from the standard dilutions discussed above.

[0317] The Lowry procedure should be carried out with caution since it is subjected to interference by a wide variety of chemicals. Among the chemicals reported to interfere with the Lowry procedure are barbital, CAPS, cesium chloride, citrate, cysteine, diethanolamine, dithiothreitol, EDTA, EGTA, HEPES, mercaptoethanol, Nonidet P-40, phenol, polyvinyl pyrrolidone, sodium deoxycholate, sodium salicylate, thimerosal, Tricine, TRIS and Triton X-100.

Example 5

[0318] The following example illustrates the Somogyi-Nelson assay used to measure avicelase activity.

[0319] The assay measures the velocity of release of reducing sugars by the action of cellulases on insoluble Avicel cellulose (micro crystalline cellulose) when reducing sugars are assayed by Somogyi-Nelson method as glucose equivalents. One unit of avicelase activity is equal to the amount of enzyme that liberates 1 micromole of reducing sugars, expressed as glucose equivalents (when reducing sugars assayed by Somogyi-Nelson method) in one minute at 40° C, pH 5.0 (0.05 M sodium acetate buffer).

Reagents

[0320] Acetate buffer (0.05 M, pH 5.0) is prepared as follows. 4.1 g of anhydrous sodium acetate or 6.8 g of sodium acetate $\cdot 3\text{H}_2\text{O}$ is dissolved in distilled water so that the final volume of the solution is 1000 mL (Solution A). In a separate flask, 3.0 g (2.86 mL) of glacial acetic acid is mixed with distilled water to make the total volume of 1000 mL (Solution B). The final 0.05 M acetate buffer, pH 5.0, is prepared by mixing Solution A with Solution B until the pH of the resulting solution is equal to 5.0.

[0321] Avicel PH 105 (Serva, FRG) (or any other available source of Avicel) is used as the assay substrate. 1% w/v of stock suspension of Avicel is prepared in 0.05 M acetate buffer, pH 5.0, by adding 50 mg of dry Avicel gradually to a vessel with 5 mL of the buffer agitated on a magnetic stirrer. After that the obtained Avicel suspension is mixed on a magnetic stirrer during 1 hour at room temperature (for swelling of Avicel). 1% w/v stock suspension of Avicel can be stored in covered vessel for 5-7 days at 4°C.

EP 2 183 363 B1

[0322] The Somogyi reagent is prepared as follows. 24 g of anhydrous sodium carbonate and 12 g of sodium-potassium tartrate $\cdot 4\text{H}_2\text{O}$ are dissolved in 250 mL of distilled water. The solution of copper sulfate $\cdot 5\text{H}_2\text{O}$ (4 g in 40 mL of distilled water) is added on stirring, and then 16 g of sodium bicarbonate is dissolved (Solution A). In a separate flask, 18.0 g of sodium sulfate is dissolved in 500 mL of hot ($\sim 80^\circ\text{C}$) distilled water and boiled for 5 minutes (Solution B). Solution A is mixed with Solution B and the final volume is adjusted to 1000 mL by distilled water. The reagent is stable for 2-3 months on storing in a dark glass vessel at room temperature.

[0323] The Nelson reagent is prepared as follows. 50 g of ammonium molybdate is dissolved in 900 mL of hot ($\sim 60^\circ\text{C}$) distilled water. The solution is cooled to $5\text{--}10^\circ\text{C}$ and 42 mL of concentrated sulfuric acid containing 6 g of sodium arsenate are carefully added with stirring. The volume of the mixture is adjusted to 1000 mL with distilled water. The flask with this solution is incubated at 40°C for 48 hours and then the solution is filtered if necessary. The reagent is stable for 2-3 months at room temperature.

[0324] Using the above reagents, the assay is performed as detailed below.

Buffer Blank

[0325] 0.2 mL of 0.05 M sodium acetate buffer (pH 5.0) is mixed in a graduated glass test tube with 0.2 mL of Somogyi reagent, covered, and incubated in a boiling water bath for 40 minutes. Then, the test tube is cooled in a cold-water bath and 0.2 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 2 mL with distilled water. The obtained solution is used as a buffer blank.

Determination of Avicelase activity

[0326] Avicelase activity is determined by carrying out of the reaction of Avicel hydrolysis at 40°C and pH 5.0 for 60 minutes while mixing. A 1% w/v stock suspension of Avicel is mixed on magnetic stirrer for 10 minutes, then 0.25 mL of this 1% w/v Avicel stock solution is placed in a graduated glass test tube. 0.15 mL of 0.05 M acetate buffer (pH 5.0) is added to the tube and the mixture is heated to 40°C for 10 minutes with stirring. 0.1 mL of enzyme sample, suitably diluted and preheated to 40°C for 10 minutes, is placed into the tube. The reaction mixture is then incubated for 1 hour at 40°C with stirring. After 1 hour, the reaction mixture is centrifuged 2 min at 12000 rpm, 0.2 mL of supernatant is withdrawn and placed in a graduated glass test tube. 0.2 mL of Somogyi reagent is added and the mixture incubated, covered, in a boiling water bath for 40 minutes. Then, the tube is cooled in a cold water bath and 0.2 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 2 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the buffer blank as A_{ES} (enzyme sample). The enzyme sample must be diluted with the 0.05 M sodium acetate buffer so that A_{610} is between 0.5 and 1.5 units of optical density at 610 nm.

Enzyme Blank

[0327] 0.16 mL of 0.05 M sodium acetate buffer (pH 5.0) is mixed in a graduated glass test tube with 0.04 mL of enzyme sample having the same dilution rate as above. 0.2 mL of Somogyi reagent is added to the test tube, and the tube is incubated, covered, in a boiling water bath for 40 minutes. Then, the test tube is cooled in a cold water bath, 0.2 mL of Nelson reagent is added and the tube is mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 2 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the buffer blank as A_{E} (enzyme blank).

Substrate Blank

[0328] A 1% w/v stock suspension of Avicel is mixed using a magnetic stirrer for 10 minutes, then 0.25 mL of the 1% w/v Avicel stock solution is placed into a graduated glass test tube. 0.25 mL of 0.05 M acetate buffer (pH 5.0) is added to the tube and the mixture is heated to 40°C for 10 minutes with stirring. 0.1 mL of enzyme sample, suitably diluted and preheated to 40°C for 10 minutes, is placed into the tube. The reaction mixture is then incubated for 1 hour at 40°C with stirring. After 1 hour, the reaction mixture is centrifuged 2 min at 12000 rpm, 0.2 mL of supernatant is withdrawn and placed in a graduated glass test tube. 0.2 mL of Somogyi reagent is added and the mixture incubated, covered, in a boiling water bath for 40 minutes. Then, the tube is cooled in a cold water bath and 0.2 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 2 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the buffer blank as A_{S} (substrate blank).

Standard Calibration Plot

[0329] Stock glucose solution (0.5 mg/mL) is prepared by dissolving 10 mg of D-glucose in 20 mL of 0.05 M acetate buffer, pH 5.0. Then, different dilutions of stock glucose solution are prepared as follows:

5

10

15

Stock solution (mL)	0.05 M acetate buffer (mL)	Glucose concentration (mg/mL)
0.020	0.180	0.02
0.040	0.160	0.04
0.060	0.140	0.06
0.080	0.120	0.08
0.100	0.100	0.10
0.120	0.080	0.12
0.140	0.060	0.14
0.160	0.040	0.16
0.180	0.020	0.18
0.200	0	0.20

20 Samples (0.2 mL) of glucose solution with different dilutions are placed to graduated glass test tubes. 0.2 mL of Somogyi reagent is added and the test tubes are placed into a boiling water bath for 40 minutes. Then, the test tubes are cooled in a cold water bath and 0.2 mL of Nelson reagent is added to each tube, and the tubes mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume in each tube is adjusted to 2 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the buffer blank. The following typical A_{610} data obtained with different glucose dilutions are given below:

25

Glucose, mg/mL	0.02	0.04	0.06	0.08	0.10	0.12	0.14	0.16	0.18	0.20
A_{610}	0.02	0.12	0.38	0.50	0.70	0.92	1.10	1.38	1.54	1.67

30

Absorbance (A_{610}) on the y-axis versus glucose concentration (G, mg/mL) on the x-axis is plotted to generate a standard calibration plot such as that shown in the Figure 11.

[0330] The dependence of A_{610} from G is described by the equation $y = ax + b$. The concentration of reducing sugars (C) in an assayed sample can be calculated from a calibration plot using the equation $C = (\Delta A_{610} - b)/a$, where $b = -0.2239$ and $1/a = 0.204$. In other words, $C = (\Delta A_{610} + 0.2239) * 0.104$ (in mg/mL).

35

Calculation of Activity

[0331] Activity is calculated as follows: Avicelase (U/mL) = $(\Delta A_{610} - b) * (1/a) * 0.463 * DR$, where $\Delta A_{610} = A_{ES}$ (enzyme sample) - A_E (enzyme blank) - A_S (substrate blank), $1/a$ is 0.204, b is -0.2239, the coefficients from the glucose calibration plot (see above), DR is the enzyme dilution rate (dilution rate of enzyme sample before addition into the reaction mixture), and 0.463 is the coefficient obtained as $5/(0.180 * 60)$, where 5 is the additional dilution of enzyme sample in the reaction mixture (0.1 mL of enzyme + 0.4 mL of reaction mixture), 0.180 mg/mkmol is the transition coefficient for glucose from mg to mkmol (molecular weight of glucose is 180) and 60 is the reaction time (60 minutes). The enzyme sample must be diluted so that ΔA_{610} should be between 0.5 and 1.5 units of optical density.

45

[0332] An example calculation is supplied below. The sample the dilution rate of 500, A_{ES} is 1.436, A_E is 0.102, A_S is 0.070, so $\Delta A_{610} = 1.436 - 0.102 - 0.070 = 1.264$. The Avicelase activity is: $(1.264 + 0.2239) * 0.104 * 0.463 * 500 = 35.8$ U/mL.

Example 6



50

[0333] The following example illustrates glucose determination by a glucosooxidase-peroxidase assay.

[0334] As a result of β -D-Glucose oxidation by glucose oxidase (GOD), hydrogen peroxide H_2O_2 and glucono-lactone are formed. Then, H_2O_2 is used by peroxidase (POD) as an oxidizing agent for 4-aminoantipyrine, oxidizing in the presence of phenolic substances. As a result, dyed products accumulate in the reaction mixture and can be detected at 490 nm spectrophotometrically. The amount of dyed products corresponds to glucose concentration as shown in the reaction scheme below:

55

EP 2 183 363 B1



[0335] The assay is carried out using a Fotoglucose kit (Impact Ltd., Russia) with the general conditions: 40°C, pH 7.0, duration - 15min, A_{490} , light path 1cm.

Reagents

[0336] Reagent 1 (R1) is prepared as follows. Two "Buffer-Substrate" tablets (0.1mmol/L potassium or ammonium salts of phosphoric acid, 50 mmol/l 4-aminoantipyrine, 0.75 mmol/L 8-oxyquinolyn) are dissolved in 200 mL of distilled water, filtered through a cellulose filter, and the pH adjusted to 7.0 if necessary. The solution can be stored in the dark at 4° C and should be made fresh weekly.

[0337] Reagent 2 (R2) is prepared as follows. One "Enzyme" tablet (2500 units of glucose oxidase and 500 units of peroxidase) is dissolved in 5 mL of distilled water. The solution can be stored in the dark at 4° C and should be made fresh weekly.

[0338] A glucose solution of 10 mmol/L glucose in 0.15% benzoic acid is used as the calibration standard. The solution is diluted with distilled water to obtain glucose solutions of 0 to 5 g/L.

[0339] Using the above reagents, the assay is performed as detailed below.

[0340] R1 and R2 are mixed at a ratio of 40:1 (this solution may be stored in the dark at 4° C and should be made fresh weekly) and 1 mL of the mixture is preheated to 40° C for 5-10 seconds. 0.1 mL of the sample (or glucose standard) is then added and the mixture is incubated at 40° C for 15 minutes. The mixtures are then placed at room temperature and the A_{490} determined on spectrophotometer. Prolongation of the incubation time to 20 minutes has little or no effect on the results, while decreasing it to less than 15 minutes can result in a lower optical density. Thus, 15-20 min of incubation at 40°C is preferred for glucose determination.

Calculation of Glucose Concentration

[0341] As shown in Table 21, calibration with standards containing varying concentrations of glucose reveals an optical density A_{490} in the range from 0.06 to 1.15 corresponds to glucose concentrations from 0.025 g/L to 0.5 g/L. These data are plotted in Figure 12, which demonstrates a linear proportion and high reproducibility at these conditions.

[0342] Using the calibration results, the following formula is obtained for glucose determination in samples from an A_{490} reading:

$$\text{GLUCOSE, g/L} = (A_{490\text{sample}} - A_{490\text{blank}}) - 0.0058) / 2.3029$$

Table 21. Calibration of the reagents using standard glucose solutions (two experiments)

Glucose, g/L	A_{490}	A_{490}
0	0.05	0.05
0.0025	0.05	0.05
0.005	0.06	0.06
0.01	0.07	0.07
0.025	0.11	0.11
0.05	0.17	0.17
0.1	0.29	0.29
0.25	0.65	0.64
0.5	1.20	1.20

EP 2 183 363 B1

(continued)

Glucose, g/L	A ₄₉₀	A ₄₉₀
1	1.71	1.72
2.5	1.93	1.94
5	2.06	2.06

Example 7

[0343] The following example illustrates the Somogyi-Nelson assay used to measure xylanase activity.

[0344] This assay measures the release of reducing sugars (as glucose equivalents) by the action of xylanase on a soluble xylan. One xylanase unit of activity is the amount of enzyme that liberates 1 micromole of reducing sugars, expressed as glucose equivalents, in one minute at 50° C and pH 5.0.

Reagents

[0345] Acetate buffer (0.05 M, pH 5.0) is prepared as follows. 4.1 g of anhydrous sodium acetate or 6.8 g of sodium acetate * 3H₂O is dissolved in distilled water so that the final volume of the solution is 1000 mL (Solution A). In a separate flask, 3.0 g (2.86 mL) of glacial acetic acid is mixed with distilled water to make a total volume of 1000 mL (Solution B). The final 0.05 M acetate buffer, pH 5.0, is prepared by mixing Solution A with Solution B until the pH of the resulting solution is equal to 5.0.

[0346] Xylan from birchwood (Sigma, X 0502) is used as the substrate. A 1% w/v stock solution is prepared by dissolving 500 mg of xylan in 50 mL of distilled water with agitation on a magnetic stirrer for 1 hour. The solution is stable for 2 days at 4° C.

[0347] The Somogyi reagent is prepared as follows. 24 g of anhydrous sodium carbonate and 12 g of sodium-potassium tartrate tetrahydrate are dissolved in 250 mL of distilled water. A solution of copper sulfate pentahydrate (4 g in 40 mL of distilled water) is added with stirring, and then 16 g of sodium bicarbonate is dissolved to obtain Solution A. In a separate flask, 180 g of sodium sulfate is dissolved in 500 mL of hot (~80° C) distilled water and boiled for 5 min (Solution B). Solution A is mixed with Solution B and the final volume is adjusted to 1000 mL. The reagent is stable for 2-3 months on storing in a dark glass vessel.

[0348] The Nelson reagent is prepared as follows. 50 g of ammonium molybdate is dissolved in 900 mL of hot (~60° C) distilled water. The solution is cooled to 5-10° C and 42 g of concentrated sulfuric acid containing 6 g of sodium arsenate are carefully added with stirring. The volume of the mixture is adjusted to 1000 mL. The flask is incubated at 40° C for 48 hours and then the solution is filtered if necessary. The reagent is stable for 2-3 months.

[0349] Using the above reagents, the assay is performed as detailed below.

Reagent Blank

[0350] 0.5 mL of 0.05 M sodium acetate buffer is mixed in a graduated glass test tube (1.5 cm x 15 cm) with 0.5 mL of Somogyi reagent and incubated in a boiling water bath for 40 minutes. Then, the test tube is cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water.

Enzyme Sample

[0351] 0.25 mL of 1% w/v xylan stock solution is mixed with 0.15 mL of 0.05 M acetate buffer, pH 5.0, in graduated glass test tubes (1.5 cm x 15 cm) and preheated at 50° C for 5 minutes. Enzyme sample is suitably diluted by the 0.05 M sodium acetate buffer (the dilution of the enzyme sample is chosen so that A₆₁₀ should be between 0.95 and 1.05 units of optical density). 0.1 mL of suitably diluted enzyme sample is added to 0.4 mL of the preheated substrate solution, mixed and incubated at 50° C for 10 minutes. After exactly 10 minutes of incubation, 0.5 mL of Somogyi reagent is added, and test tubes are placed into a boiling water bath for 40 minutes. Then, the test tubes are cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A₆₁₀) is measured against the reagent blank as As (enzyme sample).

EP 2 183 363 B1

Enzyme Blank

[0352] 0.4 mL of 0.05 M sodium acetate buffer is mixed in a graduated glass test tube (1.5 cm x 15 cm) with 0.1 mL of enzyme sample having the same dilution as above. 0.5 mL of Somogyi reagent is added and the test tube is incubated in a boiling water bath for 40 minutes. Then, the test tube is cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the reagent blank as A_{EB} (enzyme blank).

Substrate Blank

[0353] 0.25 mL of 1% w/v xylan stock solution is mixed with 0.25 mL of 0.05 M acetate buffer, pH 5.0, in a graduated glass test tube (1.5 cm x 15 cm), 0.5 mL of Somogyi reagent is added, and the test tube is placed into a boiling water bath for 40 minutes. Then, the test tube is cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the reagent blank as A_{SB} (substrate blank). The value of A_{SB} should not exceed 0.05 units of optical density.

[0354] All enzyme samples and enzyme and substrate blanks should be boiled together.

Standard Calibration Plot

[0355] Stock glucose solution (1 mg/mL) is prepared by dissolving 100 mg of D-glucose in 100 mL of 0.05 M acetate buffer, pH 5.0. Then, different dilutions of stock glucose solution are prepared as follows:

Stock solution (mL)	0.05 M acetate buffer (mL)	Glucose concentration (mg/mL)
0.05	4.95	0.01
0.10	4.90	0.02
0.15	4.85	0.03
0.20	4.80	0.04
0.25	4.75	0.05
0.30	4.70	0.06
0.35	4.65	0.07
0.40	4.60	0.08
0.45	4.55	0.09
0.50	4.50	0.10

Samples (0.5 mL) of glucose solution with different dilutions are placed in graduated glass test tubes (1.5 cm x 15 cm). 0.5 mL of Somogyi reagent is added and test tubes are placed into a boiling water bath for 40 minutes. Then, the test tubes are cooled in an ice or cold water bath and 0.5 mL of Nelson reagent is added and mixed by shaking manually. After 10 minutes incubation at room temperature, the solution volume is adjusted to 5 mL with distilled water. Then, the absorbance at 610 nm (A_{610}) is measured against the reagent blank. The following typical A_{610} data obtained with different glucose dilutions are given below:

Glucose, mg/mL	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	0.10
A_{610}	0.11	0.20	0.28	0.41	0.48	0.60	0.72	0.79	0.93	1.06

Absorbance (A_{610}) on the y-axis versus glucose concentration (G, mg/mL) on the x-axis is plotted to demonstrate a standard calibration plot such as that shown in Figure 11. From Figure 11, the inverse of the slope ($1/a = 0.098$) is calculated.

Calculation of Activity

[0356] Activity is calculated as follows: activity (IU/mL) = $1/a * \Delta A_{610} * DF * 2.78$, where $1/a$ is the inverse of the slope calculated from the standard calibration plot, ΔA_{610} is A_s (enzyme sample) - A_{EB} (enzyme blank) - A_{SB} (substrate blank), DF is the enzyme dilution factor, and 2.78 is the coefficient that accounts for additional 5-fold enzyme dilution in the reaction mixture (0.1 mL enzyme + 0.4 mL substrate), reaction time (10 minutes) and molecular weight of glucose

(0.18016 mg/micromole), i.e. $5/(10 \times 0.18016) = 2.78$

[0357] An example calculation is supplied below.

B4 UF concentrate	DF	ΔA_{610}
	5,000	1.61
	10,000	1.04
	20,000	0.64

$$\text{Xylanase activity (IU/mL)} = 0.098 * 1.04 * 10000 * 2.78 = 2833$$

Example 8

[0358] The following example illustrates the assay used to measure the stability of enzymes of the present disclosure and the protein of the present invention to thermoshock.

[0359] The following assay determines the stability of enzymes after short-term treatment at elevated temperature (e.g. temperature of pelletization of animal feed). A solution of enzyme is incubated for approximately two minutes in a glass tube in a thermostat at 80°C. The gradual drop of enzyme activity is recorded over time and the resultant curve is used to determine how long the enzyme is stable (results usually range from 10 - 60 seconds).

[0360] The use of glass tubes that fit tightly in a regular thermostat or eppendorf-type tubes help ensure proper thermal conditions and fast heating of the tube content. The enzyme dilution used is selected so that after additional dilution in reaction mixture the optical density is as high as possible (depends on particular method used for activity determination). For example, when determining xylanase activity by the Somogyi-Nelson assay disclosed in Example 7, the enzyme selected should give an optical density A_{610} 1.0-1.1 after being diluted 5 times in reaction mixture.

[0361] For faster equilibration at 80°C, 0.4-0.5 ml water may be placed in the well of thermostat, prior to fitting in a glass tube. The temperature is set at 80.3 - 80.4°C to compensate for cooling. 360 μ l of desired buffer is placed in the tube and heated for 2-3 minutes until it reaches 80°C. It is advisable to check the temperature in the tube (using a thermometer) to be sure the temperature has been reached. 40 μ l of enzyme solution is then added and the tube is mixed rapidly. After 20, 40, 60, 90 and 120 seconds, aliquots are removed from the tube and immediately transferred in the reaction mixture for determination of activity (assay components are prepared beforehand). To determine zero-point activity (without heating), the same solution (360 μ l buffer+40 μ l enzyme) is prepared in the separate tube at room temperature. Determined activities at every time point are then plotted as relative activity (in %) vs. time, with zero time point (without heating) representing 100% activity. Figure 13 demonstrates these data for stability of xylanase at 20-, 10- and 5-fold dilutions.

[0362] In this protocol, a 10-fold dilution of enzyme is used (9:1 preheated buffer solution to cold enzyme solution). In principle one can use higher dilution rates (e.g., 1:20), but typically not lower dilution factors (e.g., 1:5) because mixing of larger volumes of cold enzyme solution can produce inaccurate results. The plot below illustrates this principle. The stability of xylanase was tested using 20-, 10- and 5-fold dilutions, and the results shown in Figure 13 demonstrate that dilutions of 10- and 20-fold (and likely higher) gave consistent results. In contrast, use of a 5-fold dilution generates overestimated values.

[0363] The percentages of enzymatic activity of xylanases of the present invention after 20 seconds of incubation at 80°C (also known as the pelletizing stabilities) are shown below in Table 22 and in the graph depicted in Figure 14.

Table 22. Pelletizing Stability of C1 Xylanases

Enzyme	% Activity
Xyl 1	19
Xyl 1 (CD)	12
Xyl 2	98
Xyl 3	75
Xyl 3 (CD)	82
Xyl 4	23
Xyl 5	60

(continued)

Enzyme	% Activity
Xyl 6	68

Example 9

[0364] The following example illustrates the *in vitro*, semi-quantitative assay used to measure the feed activity of enzymes of the present disclosure and the protein of the present invention.

[0365] The assay was developed to simulate the enzymatic treatment of feed. The assay detects the dynamics of liquid and sludge fraction formation from a uniform slurry of milled rye bran by the action of the enzyme samples in narrow test tubes.

[0366] A mixture of enzyme solution and uniform rye bran suspension in acetate buffer (pH 5.0) is placed into a long glass tube (e.g., 2 ml calibrated glass pipettes) and stored at room temperature (or at 40°C). The action of enzymes on non-starch rye bran generates polysaccharides (e.g. arabinoxylan, β -glucan and others) causing the original uniform rye bran slurry to separate into a liquid phase (containing the polysaccharides) on the top and an insoluble sludge on the bottom of. The volume of liquid phase formed on the top of tubes over the reaction is measured every 10-15 min during 1.5-3 hours (the total assay time depends on enzymatic activity). The data obtained are plotted as "liquid formed" (ml) versus time (minutes) and the slope of the line calculated. The higher the slope of the line on the obtained graph, the higher the *in vitro* feed activity of the tested enzyme sample.

Substrate

[0367] Rye bran is used as a substrate in this example. However, wheat and barley bran, rye, wheat, barley, and soy meal may also be used as assay substrates. Partially de-starched rye bran samples are carefully milled and passed through the fine sieve. The fine fraction is taken as the substrate. Substrate is accurately weighed and placed into a dry 20-ml screw cap vial. The sample of substrate is suspended in acetate buffer to get 30% w/v suspension. The substrate suspension is mixed vigorously with a magnetic stirrer for 30-45 minutes at room temperature. The resulting suspension obtained can be stored at 4°C in a sealed vial for about 2 days.

Assay procedure

[0368] The reaction mixture contains the following components:

- 1500 μ l of substrate suspension (300 mg/ml)
- 750 μ l of enzyme solution

[0369] The blank solution contains the same amount of substrate, but the enzyme solution is substituted with the same volume of acetate buffer.

[0370] The reaction mixture is subjected to vigorous agitation and placed in appropriate calibrated glass tubes (e.g., 2 ml glass pipettes). Glass tubes or pipettes should be filled with the reaction mixture immediately and placed vertically. The top end of the tube is left open while the bottom end is sealed with a piece of Parafilm or by dipping in melted paraffin.

[0371] The concentration of enzyme samples can be measured either in mg/ml of protein or in xylanase units/ml in the reaction mixture (or any other enzyme activities, e.g., β -glucanase, α -galactosidase, protease, etc.). Protein content or xylanase (or other activities) of the enzyme samples are measured separately prior to the experiments. In the experiments, enzyme concentration was equalized to 1 mg/ml of protein in the reaction mixture, or in xylanase activity (2 to 3 U/ml).

Materials and instruments**[0372]**

- Thermostat for 2 ml Eppendorf type test tubes, set at 40°C
- Set of calibrated glass tubes (e.g., 2-ml glass pipettes)
- UV-Vis spectrophotometer
- Reagents to perform Nelson-Somogyi assay
- Analytical electronic balances

EP 2 183 363 B1

- Adjustable volume automated pipettes 40-200 µl and 200-1000 µl
- Rye bran
- 0.05 M acetate buffer, pH 5.0

Data analysis

[0373] Data obtained is presented on a graph as liquid formed (ml) versus time (minutes). The higher the slope of the line on the obtained graph, the higher the *in vitro* feed activity of the tested enzyme sample.

Example 10

[0374] The following example illustrates the assay used to measure the ability of enzymes to reduce the viscosity of feed materials.

[0375] Enzymatic treatment of barley, rye or wheat can increase the nutritive value by hydrolysis of viscous polysaccharides (beta-glucan, arabino xylan) within cereals. Table 23 below illustrates typical viscous polysaccharide profiles of sample feedstuffs compared with the viscosity of water/meal mixtures.

Table 23. Viscous Polysachharide Profiles of Feedstuffs

	Arabino Xylans g/kg of dry matter	Beta-Glucans, g/kg of dry matter	Relative Viscosity of water/meal mixes, U
RYE	14.4	7.6	9.71
BARLEY	3.3	24.3	4.21
WHEAT	3.2	5.6	2.25

[0376] As shown in the table, the high viscosity of a water/rye meal mixture is predominantly due to the presence of arabinoxylans. Similarly, the viscosity of a water/barley mixture is predominantly due to the presence of beta-glucans, while the viscosity of water/wheat mixture close to water. However, cereals such as barley, rye and wheat often contain specific inhibitors of the hydrolytic enzymes. Thus, preferred enzymatic preparations that effectively decrease viscosity should be resistant to cereal inhibitors and be enzymatically active at low dosage.

Materials and Instruments

[0377]

- 100 ml glass flasks with caps
- Ostwald capillary viscometer with thermostat at 40° C
- Shaker for meal incubation at 40° C
- Whole rye, barley, wheat grains or hulls or their mixtures
- Rotary mill and 0.5 mm sieve to prepare whole meal
- Adjustable volume automated pipettes 40-200 and 200-1000 µl
- Distilled water or 0.1M sodium acetate buffer, pH 5.0

Sample Preparation

[0378] To prepare the crude barley, wheat or rye meals, whole grains are milled on a rotary mill and passed through a 0.5 mm sieve. 6 g of crude barley or rye meal or 12 g of wheat meal is added to a 100 ml glass flask. 30 ml of distilled water or buffer (blank flask) or 30 ml of the enzyme water solution (experimental flasks) is then added and the flasks are covered. The concentration of 6 g of meal per 30 ml of water is 200 g per liter, while 12 g per 30 ml is 400 g/l. The flasks are shaken well by hand then placed on a shaker at 40° C, 250 rpm.

[0379] After a 1-hour incubation, the mixtures are centrifuged for 5 min at 5000 rpm to separate the insoluble precipitate of the crude meal. The supernatant is then filtered through a cotton pad. 5 ml of the filtered supernatant is placed into an Ostwald viscometer at 40° C and incubated for at least 3 minutes before the viscosity of the sample is determined. The measurement of the efflux time of the samples from the viscometer bulb should be done later than 80 minutes after starting the incubation.

[0380] A dosage of 0.15 kg/t of the meal approximates the dosage used in real practice. For liquid preparations, a

dosage of 0.15 kg/t equals 90 µl of the preparation in a 100-fold dilution per flask (30ml of water/meal mix). For dry preparations, a dosage of 0.15 kg/t equals 90 µl of the 10 g/l solution per flask (30 ml of water/meal mix).

[0381] Figure 15 shows the relative viscosity of water/rye mixes treated with the indicated enzyme(s). To compare the efficiency of the preparations, the xylanase activities of the samples (units per flask) are also indicated.

[0382] The efficiency of the preparations may also be compared by plotting the residual viscosity (%) of the water/meal mix after enzymatic treatment, as shown in Figure 16.

[0383] Residual viscosity is calculated as follows:

$$\text{ResV} = (\text{Rvi} - \text{Rvwater}) / (\text{Rvnative} - \text{Rvwater}) \times 100\%$$

The final formula is:

$$\text{ResV} = (\text{Rvi} - 1) / (\text{Rvnative} - 1) \times 100\%$$

Where ResV is residual viscosity, Rvi is the relative viscosity of the sample with enzyme, Rvnative is the relative viscosity of the native water/meal mix without enzyme, and Rvwater is 1.

[0384] Table 24 shows of the residual viscosity (%) of rye meal water extract after a 1-hour incubation with 1 unit of the xylanase of the present invention and the xylanases of the present disclosure. These data are presented graphically in Figure 17.

Table 24. Residual Feed Viscosities of C1 Xylanases

Enzyme	% Residual Viscosity
Xyl 1	59
Xyl 1 (CD)	59
Xyl 2	27
Xyl 3	25
Xyl 3 (CD)	76
Xyl 4	77
Xyl 5	94
Xyl 6	91

[0385] Table 25 shows of the residual viscosity (%) of barley meal after a 1-hour incubation with 1 unit of the endoglucanases of the present disclosure.

Table 25. Residual Feed Viscosities of C1 Endoglucanases

Enzyme	% Residual Viscosity
EG I	18
EG II	13
EG II (CD)	14
EG III	10
EG V	18
EG VI	20

Example 11

[0386] The following example illustrates an assay used to measure the biobleaching ability of enzymes of the present disclosure and the enzyme of the present invention.

[0387] Ability of different crude enzyme preparations or purified enzymes to (bio)bleach cellulosic pulp may be evaluated by the assay below. The assay monitors the release of chromogenic phenolic substances from raw cellulosic pulp by the action of enzymes. The phenolic substances are detected by the absorbance at 237 nm (A_{237}) or 235 nm (A_{235}).

Mini biobleaching assay

[0388] Raw (wet) eucalyptus cellulosic pulp with natural pH 7.7 (National Paper Fabric, Uruguay) is placed into 5 ml reaction tubes, and the tubes are placed into a water bath shaker, (50°C, 200 rpm). The reaction volume consists 2 ml of enzyme sample appropriately diluted in appropriate buffer solution and 2 ml of cellulosic pulp suspension (0.04 g of cellulosic pulp by dry weight). The concentration of cellulosic pulp is 10 mg/ml. The biobleaching process is started by the addition of 2 ml of enzyme solution to the pulp and the assay is conducted for 2 hours. After that reaction, the mixture is centrifuged and A_{237} of the supernatant is monitored in a spectrophotometer. Since some phenolic compounds are washed out from the raw cellulosic pulp even in the absence of enzymes, the blank experiments with buffer solution only (with no enzymes) are carried out. Some input to A_{237} value could be made by chromogenic substances, existing in the raw enzyme sample, so another blank experiments were carried out to estimate this input. All results take into account both of those blanks as ΔA_{237} . The higher the ΔA_{237} value, the better was biobleaching performance.

[0389] The assay may also be performed using soft or hard wood pulp at pH 7.5 and 50°C for 2 hours, with the absorbance determined at 235 nm. Table 26 and Figure 18 show the biobleaching ability (A_{235}) of the xylanase of the present invention and xylanases of the present disclosure on soft wood pulp under these conditions, while Table 27 and Figure 19 show the same biobleaching ability on hard wood pulp. For each xylanase, the assay was conducted using 0.01, 0.003, 0.001, 0.0003 mg/ml of enzyme in the reaction mixture, as indicated. A positive control xylanase preparation (Luminase, Verenum Corp., San Diego, CA USA) was also used for comparison.

Table 26. Biobleaching Ability of C1 Xylanases on Soft Wood Pulp (A_{235})

Enzyme (mg/ml)	Xyl 1 (CD)	Xyl 1	Xyl 3	Xyl 3 (CD)	Xyl 4	Xyl 2	Xyl 6	Luminase
0.01	3	1.37	2.28	2.2	1.73	3.66	2.62	3
0.003	2.65	1.015	2.7	1.43	1.1	2.8	2.03	2.445
0.001	2.385	1.07	0.85	1.05	0.455	2.22	1.7	2.14
0.0003	1.45	0.615	0.575	0.31	0.2	1.615	1.4	1

Table 27. Biobleaching Ability of C1 Xylanases on Hard Wood Pulp (A_{235})

Enzyme (mg/ml)	Xyl 1 (CD)	Xyl 1	Xyl 3	Xyl 3 (CD)	Xyl 4	Xyl 2	Xyl 6	Luminase
0.01	8.5	2.355	4.2	3.715	1.9	5	6.5	7.585
0.003	6.25	1.82	2.24	2.27	0.82	4.45	4.5	5.6
0.001	5.4	1.37	1.05	1.5	0.5	2.355	3	4.8
0.0003	3.5	0.4	0.61	0.675	0.3	1.5	1.26	2.785

Example 12

[0390] The following example illustrates the Saccharification of pretreated lignocellulosic feedstock using crude enzyme preparations.

[0391] The assay involves a protocol for enzymatic saccharification of pretreated lignocellulosic biomass with a known moisture content and composition in order to determine the extent of cellulose and hemicellulose digestibility under concrete conditions of saccharification of the feedstock with crude cellulase and hemicellulase enzyme preparations.

Equipment

[0392]

- INNOVA 40 Thermo Shaker (New Jersey, USA), where agitation of a slurry in plastic vessels is carried out using 250 rpm vibrations (each vessel supplied by one 0.5 cm metal ball to improve mixing of slurry)

- Micro centrifuge
- pH meter
- Vortex mixer
- Analytical balance, sensitive to 0.1 mg
- UV/Vis spectrophotometer
- HPLC with refractive index detector and Silica column with bonded amino phase (4 x 250 mm) and guard column
- Pipettes with variable sample volume (5-5000 μ L)

Reagents and Materials

[0393]

- Lignocellulosic feedstock with known moisture content and composition (cellulose, hemicellulose, lignin, ash, extractives)
- Sodium acetate buffer (0.1 M, pH 5.0) containing 1 mM of sodium azide to prevent microflora growth
- Enzyme cellulase and hemicellulase preparations from different fungal strains, either in soluble or in dry form
- Reagents for protein determination (either Lowry or Pierce BCA kit)
- Water, distilled and/or deionized
- Ampiox, sodium-salt.
- "Fotoglucose" kit (reagents for glucose determination using glucose oxidase - peroxidase assay) from Impact Ltd., Russia
- Somogyi and Nelson reagents for reducing sugars' assay
- Acetonitrile, HPLC grade
- NaOH, acetic acid for pH adjustments

Assay Procedure

[0394] The procedure described below assumes that the hydrolysis is carried out at 5% solids. If the process should be carried out at different substrate concentration, then recalculations should be done for the initial biomass sample weight (on a dry weight basis) using the known moisture content in the sample.

[0395] A biomass sample containing 1g of solids (on a dry weight basis) is weighed and placed into a 50 ml plastic vessel equipped with a hermetic lid and one metal ball (0.5 cm in diameter) to improve mixing (the moisture content is determined in preliminary experiments by drying the biomass samples at 105°C until a constant weight is reached). The weight (volume) of water in the sample is noted as W_s .

[0396] The volume(s) of enzyme(s) necessary to add to the reaction mixture is calculated taking into account the protein concentration (enzyme activity) in a stock solution and the final protein concentration (enzyme activity) in the reaction mixture. Using these data, the volume of buffer (W_a) that is necessary for adjusting the total volume of the reaction system to 20 ml is calculated, taking into account water contained in the biomass sample (W_s).

[0397] The calculated amount (W_a) of 0.1 M, pH 5.0 Na-acetate buffer containing 200mg/l (4 mM) of sodium azide is added to the biomass slurry, and the slurry is thoroughly mix in the 50 ml vessel using a Vortex. The pH is measure and adjusted pH to 5.0 using concentrated acetic acid or NaOH, if necessary. The lid is closed and the vessels incubated in an INNOVA 40 Thermo Shaker for 15 minutes to preheat the sample to 50°C.

[0398] The calculated amount(s) of enzyme stock solution(s) is added to the reaction mixture to start the enzymatic reaction. 20 μ l of Ampiox solution (100g/l) is then added to the final reaction mixture, which is allowed to proceed for 72 hours. (50°C and 250 rpm vibrations).

[0399] Aliquots of the reaction slurry (50-100 μ L) are typically taken at 6, 24, 48, 72 or 96 hour time points, using a pipette with a cut tip. The aliquots are centrifuged at 15,000 rpm in a micro centrifuge for 3 minutes, and the supernatants are analyzed for glucose by the glucose oxidase - peroxidase assay (see Example 6) and for reducing sugars by the Nelson-Somogyi assay (see, for example, Example 1). Samples (supernatants) are typically analyzed immediately. In some cases, they are subjected to boiling for 5 minutes, then frozen and analyzed later. The final time point (72 hours) is also analyzed for sugars by HPLC.

[0400] A substrate blank (control) is prepared in the same way as described above, except that distilled water is added to the vessel instead of the enzyme(s) stock solution(s). The supernatant resulting from the substrate blank is analyzed for glucose by the glucose oxidase - peroxidase assay, reducing sugars by the Nelson-Somogyi assay and also for sugars by HPLC.

[0401] Reactions are typically carried out in duplicate. The results of the analysis for sugars are determined as average values together with standard deviations. Typically, the protein content in the reaction system (mg protein/g solids) is used as a measure of enzyme loading. In the experiments with crude enzyme preparations, the enzyme loading is 5,

2.5 and 1 mg protein/g solids. In enzymatic reactions may be carried out in the presence of β -glucosidase preparations (40 Units of p-NPh- β -glucosidase activity per 1 g solids) in order to convert all soluble oligosaccharides to glucose.

Calculations

[0402] To calculate the percent digestibility of cellulose (glucan), the glucose concentration in the final (72 or 96 hour) supernatant of the reaction mixture is determined (data from the glucose oxidase - peroxidase assay and/or HPLC assay), subtract the glucose concentration from the substrate blank (control). The initial cellulose (glucan) content in the reaction system is also calculated. For example, if the cellulose content in the pretreated feedstock is 35% of the total solids (dry matter) and the hydrolysis is carried out at 100 g solids/L, then the initial cellulose concentration would be 35 mg/mL.

[0403] The percent digestion of cellulose is calculated as follows:

$$\text{Digestion (\%)} = \frac{\text{Glucose concentration (mg/mL)} \times 100\%}{\text{Initial cellulose concentration (mg/mL)} \times 1.11}$$

The factor 1.11 is a result of hydration of anhydroglucose residues in cellulose (162 Da per one residue) that yields a glucose molecule (180 Da).

[0404] To calculate the percent digestibility of hemicellulose (xylan), the total concentration of pentoses (xylose + arabinose) in the final (72 or 96 hour) supernatant of the reaction mixture is determined (data from HPLC assay), subtracting the pentose (xylose + arabinose) concentration from the substrate blank (control). The initial hemicellulose (xylan) content in the reaction system is also calculated. For example, if hemicellulose content in the pretreated feedstock is 30% of the total solids (dry matter) and the hydrolysis is carried out at 100 g solids/l, then the initial hemicellulose concentration would be 30 mg/ml.

[0405] The percent digestion of hemicellulose is calculated as follows:

$$\text{Digestion (\%)} = \frac{\text{Total pentose concentration (mg/mL)} \times 100\%}{\text{Initial hemicellulose concentration (mg/mL)} \times 1.136}$$

The factor 1.136 is a result of hydration of anhydropentose (xylose or arabinose) residue in hemicellulose (132 Da per one residue) that yields a pentose molecule (150 Da).

References

[0406]

1. Berezin IV, Rabinovich ML, Sinitsyn AP (1977) Study of applicability of quantitative kinetic spectrophotometric method for glucose determination. *Biokhimiya (Moscow)* 42:1631-1636.
2. Ghose TK (1987) Measurement of cellulase activities. *Pure Appl. Chem.* 59:257-268.
3. Gusakov AV, Salanovich TN, Antonov AI, Ustinov BB, Okunev ON, Burlingame R, Emalfarb M, Baez M, Sinitsyn AP (2007) Design of highly efficient cellulase mixtures for enzymatic hydrolysis of cellulose. *Biotechnol. Bioeng.* On-line: <http://www3.interscience.wiley.com/cgi-bin/abstract/114064699/>, DOI: 10.1002/bit.21329
4. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193:265-275.
5. Sinitsyn AP, Chemoglazov VM, Gusakov AV (1990) Methods of investigation and properties of cellulolytic enzymes (in Russian), *Biotechnology Series*, v.25. Moscow: VINITI Press. 220 p.
6. Somogyi M (1952) Notes on sugar determination. *J. Biol. Chem.* 195:19-23.

Example 13

[0407] The following example illustrates the saccharification of pretreated lignocellulosic feedstock with individual purified enzymes and enzyme mixtures.

[0408] The assay involves a protocol for the enzymatic saccharification of pretreated lignocellulosic biomass with known moisture content and composition in order to determine the extent of cellulose and hemicellulose digestibility

under concrete conditions of saccharification of the feedstock with individual purified enzymes (monocomponents) and their compositions (mixes of monocomponents).

Equipment

[0409]

- TS-100 Thermo Shaker (Riga, Latvia), where agitation of a slurry in plastic tubes is carried out using 1200 rpm vibrations
- Micro centrifuge
- pH meter
- Vortex mixer
- Analytical balance, sensitive to 0.1 mg
- UV/Vis spectrophotometer
- HPLC with refractive index detector and Silica column with bonded amino phase (4 x 250 mm) and guard column
- Pipettes with variable sample volume (5-1000 μ L)

Reagents and Materials

[0410]

- Lignocellulosic feedstock with known moisture content and composition (cellulose, hemicellulose, lignin, ash, extractives)
- Sodium acetate buffer (0.1 M, pH 5.0) containing 1 mM of sodium azide to prevent microflora growth
- Enzyme monocomponents (cellulases, hemicellulases, glycosidases) isolated from different fungal strains, either in solution or freeze-dried
- Reagents for protein determination (either Lowry or Pierce BCA kit)
- Water, distilled and/or deionized
- "Fotoglucose" kit (reagents for glucose determination using glucose oxidase - peroxidase assay) from Impact Ltd., Russia
- Somogyi and Nelson reagents for reducing sugars' assay
- Acetonitrile, HPLC grade
- NaOH, acetic acid for pH adjustments.

Assay Procedure

[0411] The procedure described below assumes that the hydrolysis is carried out at 10% solids. If the process should be carried out at different substrate concentration, then recalculations should be done for the initial biomass sample weight (on a dry weight basis) using the known moisture content in the sample.

[0412] A biomass sample containing 200 mg of solids (on a dry weight basis) is weighed and placed into a 2 ml plastic tube equipped with a hermetic lid (the moisture content is determined in preliminary experiments by drying the biomass samples at 105°C until a constant weight is reached). The weight (volume) of water in the sample is noted as W_s .

[0413] 1 ml of Na-acetate buffer (0.1 M, pH 5.0) containing 1 mM of sodium azide is added, and the slurry is thoroughly mixed using a vortex. The pH is measured and adjusted to 5.0 using concentrated acetic acid or NaOH, if necessary.

[0414] The volume(s) of enzyme(s) that is necessary to add to the 2 ml tube is calculated, taking into account the protein concentration (enzyme activity) in a stock solution and the final protein concentration (enzyme activity) in the reaction mixture. Using these data, the volume of water (buffer), that is necessary for adjusting the total volume of the reaction system to 2 mL (W_a) is calculated, taking into account water contained in the biomass sample (W_s) and the previously added 1 ml of buffer.

[0415] The calculated amount of the acetate buffer (W_a) is added to the biomass slurry, the lid closed and the tube incubated in a TS-100 Thermo Shaker for 5 minutes to preheat the sample to 50°C. The calculated amount(s) of enzyme stock solution(s) is added to the tube to start the enzymatic reaction. The reaction is allowed to proceed at 50°C and 1200 rpm vibrations in a TS-100 Thermo Shaker for 96 hours.

[0416] Aliquots of the reaction slurry (50-100 μ L) are typically taken at 3 or 6, 24, 48, 72, and 96 hours, using a pipette with a cut tip. The aliquots are centrifuged at 15,000 rpm in a micro centrifuge for 3 minutes, and the supernatants are analyzed for glucose by the glucose oxidase - peroxidase assay (see Example 6) and for reducing sugars by the Nelson-Somogyi assay (see, for example, Example 1). Samples (supernatants) are typically analyzed immediately. In some cases, they are boiled for 5 minutes, frozen and analyzed later. The final time point (96 hours) is also analyzed for sugars

by HPLC.

[0417] A substrate blank (control) is prepared in the same way as above, except that distilled water is added to the tube instead of the enzyme(s) stock solution(s). The supernatant resulting from the substrate blank is analyzed for glucose by the glucose oxidase - peroxidase assay, reducing sugars by the Nelson-Somogyi assay and also for sugars by HPLC. Typically, there is no need to prepare enzyme blanks since purified enzymes (monocomponents) usually don't contain sugars.

[0418] All reactions are carried out in duplicate, and the results of the analysis for sugars are given as average values together with standard deviations. Typically, protein content in the reaction system (mg protein/g solids) is used as a measure of enzyme loading. In the case of purified enzymes where the amino acid sequences are known, the protein concentration is calculated from the UV absorption at 280 nm using enzyme extinction coefficients predicted by the ProtParam tool (<http://www.expasy.ch/tools/protparam.html>). Otherwise, the protein concentration is determined using the Lowry (Pierce BCA) assay.

[0419] In the experiments with individual purified enzymes (monocomponents), the enzyme loading is 2 mg protein/g solids. In the case of monocomponent cellulases (endoglucanases and cellobiohydrolases), the enzymatic reactions are carried out in the presence of purified β -glucosidase (0.5 U/ml cellobiase activity) in order to convert all soluble oligosaccharides to glucose.

[0420] In the experiments with mixes of monocomponents, the enzyme loading is typically 2 mg of total protein per 1 g of solids, when preliminary screening of mixes is carried out. With selected mixes of monocomponents, the enzyme (protein) loading may be varied (up to 5-10 mg protein/g solids).

Calculations

[0421] To calculate the percent digestibility of cellulose (glucan), the glucose concentration in the final (96 hour) supernatant of the reaction mixture is calculated (data from the glucose oxidase - peroxidase assay and/or HPLC assay), subtract the glucose concentration from the substrate blank (control). The initial cellulose (glucan) content in the reaction system is also calculated. For example, if the cellulose content in the pretreated feedstock was 35% of the total solids (dry matter) and the hydrolysis was carried out at 100 g solids/l, then the initial cellulose concentration would be 35 mg/ml.

[0422] The percent digestion of cellulose is calculated as follows:

$$\text{Digestion (\%)} = \frac{\text{Glucose concentration (mg/mL)} \times 100\%}{\text{Initial cellulose concentration (mg/mL)} \times 1.11}$$

The factor 1.11 is a result of hydration of anhydroglucose residue in cellulose (162 Da per one residue) that yields a glucose molecule (180 Da).

[0423] To calculate the percent digestibility of hemicellulose (xylan), the total concentration of pentoses (xylose + arabinose) in the final (96 hour) supernatant of the reaction mixture is determined (data from HPLC assay), subtract the pentose (xylose + arabinose) concentration from the substrate blank (control). The initial hemicellulose (xylan) content in the reaction system is also calculated. For example, if the hemicellulose content in the pretreated feedstock is 30% of the total solids (dry matter) and the hydrolysis is carried out at 100 g solids/l, then the initial hemicellulose concentration would be 30 mg/ml.

[0424] The percent digestion of hemicellulose is calculated as follows:

$$\text{Digestion (\%)} = \frac{\text{Total pentose concentration (mg/mL)} \times 100\%}{\text{Initial hemicellulose concentration (mg/mL)} \times 1.136}$$

The factor 1.136 is a result of hydration of anhydropentose (xylose or arabinose) residue in hemicellulose (132 Da per one residue) that yields in pentose molecule (150 Da).

References

1. Berezin IV, Rabinovich ML, Sinitsyn AP (1977) Study of applicability of quantitative kinetic spectrophotometric method for glucose determination. Biokhimiya (Moscow) 42:1631-1636.

2. Ghose TK (1987) Measurement of cellulase activities. Pure Appl. Chem. 59:257-268.
3. Gusakov AV, Salanovich TN, Antonov AI, Ustinov BB, Okunev ON, Burlingame R, Emalfarb M, Baez M, Sinitsyn AP (2007) Design of highly efficient cellulase mixtures for enzymatic hydrolysis of cellulose. Biotechnol. Bioeng. On-line: <http://www3.interscience.wiley.com/cgi-bin/abstract/114064699/>, DOI: 10.1002/bit.21329
4. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the Folin phenol reagent. J. Biol. Chem. 193:265-275.
5. Sinitsyn AP, Chemoglazov VM, Gusakov AV (1990) Methods of investigation and properties of cellulolytic enzymes (in Russian), Biotechnology Series, v.25. Moscow: VINITI Press. 220 p.
6. Somogyi M (1952) Notes on sugar determination. J. Biol. Chem. 195:19-23.

Example 14

[0426] The following example illustrates an assay used to measure the thermal stability of enzymes of the present disclosure and the enzyme of the invention.

[0427] This assay allows the determination of the long-term thermal stability of an enzyme at 40, 50 and 70°C and at different pH values (the preferred values may vary with the area of use of the enzyme and its pH-profile, which is typically from pH 5.0 or 7.0). A solution of enzyme is incubated from 3-24 hours in a thermostat. The gradual drop of enzyme activity is recorded. The resulting data are used to determine enzyme half-life or enzyme activity after 3 hours of incubation under specified conditions (temperature and pH).

[0428] An enzyme dilution that gives a high enough optical density to be determined when the enzyme is diluted (e.g., in the range 0.8-1.2 using the Somogyi-Nelson assay described above) should be selected. 500-800 µl of this dilution (prepared using preheated buffer and 10- to 20-fold dilution factors (e.g., 450 µl preheated buffer + 50 µl of enzyme) is placed into a 1.5-2.0 ml eppendorf tube (capped) in a thermostat set at 40, 50 or 60°C. Aliquots are removed at 15, 30, 45, 60, 90, 120, 180 minutes and activity determined simultaneously using any of the enzyme activity assays disclosed in this application or known in the art. A zero-point activity is determined by preparing the same enzyme dilution at room temperature. Relative activity (in %) is plotted versus time, with the activity of the enzyme at the zero-time point (without heating) set to 100%.

Example 15

[0429] The following example illustrates methods of bioscouring textiles using enzymes of the present disclosure and enzyme of the invention.

[0430] The methods below are primarily intended for testing small amount of enzymes (mini-testing in the range of 0.5-5 mg of enzyme sample).

Bioscouring assays

[0431] Scouring removes hydrophobic waxes and pectic substance from the surface of raw cotton fabric. The following methods, which allow the evaluation of the scouring efficiency of pure enzymes and crude enzyme preparations, are listed below:

- 1) Scouring and evaluating procedure for strips of 2.5 x 16 cm textile strips,
- 2) Micromethod of scouring and evaluating procedure for circular pieces of textile, diameter 1.4 cm,
- 3) Method for determination (staining) of remaining pectic substances in scoured textile (qualitative),
- 4) Method for determination of remaining waxes in scoured textile.

Method A. Scouring and evaluation procedure for the strips of 2.5 x 16 cm textile strips

[0432] Sample preparation: For all testing, raw weave 100% cotton fabric is used (density of 142g/m²). Cotton fabric is desized by α-amylase overnight with surfactant added at room temperature, then rinsed and dried at room temperature. For 1-2 m² fabric, about 5 liters of solution containing 5 g/l textile α-amylase preparation and 1g/l nonionic surfactant for scouring is needed. The fabric is rinsed thoroughly with tap water and dried at room temperature overnight. The fabric should not be ironed. The fabric is then cut into strips (2.5 x 16 cm).

[0433] Scouring procedure: The strips are treated in 100 mL plastic vessels on an Elpan water bath shaker at 50°C at 250 rpm for one hour. Two 16 x 2.5 cm strips of fabric (1.1-1.2 g total) are loaded in one vessel. The liquor ratio is 20:1, using 0.1M acetate buffer, pH 5.0.

[0434] In detail, the strips, usually 20 strips (in 10 vessels), are marked with a textile marker, then dipped into the water to completely wet each strip. An enzyme solution (20 ml) is poured in the vessels. Then, two wet strips are rapidly

placed into each vessel using tweezers. The vessels are capped and placed into a preheated water bath shaker (set at 50°C, 250 rpm, amplitude 4) for one hour.

[0435] After the treatment, the strips are rinsed with cold tap water for 2-4 minutes, usually in one bath, and then dried at room temperature overnight. More intense rinses have been tested (hot water, sequestrates e.g. EDTA, surfactants) but are not recommended since they provide less reproducible results.

[0436] Heating: Dry strips are heated for 1 hour at 105°C, allowed to remain overnight at room temperature, and then assayed for wicking height.

[0437] Analysis of wettability: Wettability (wicking height) is measured according to the Russian textile standard. One end of the vertical strip of dry fabric is immersed in a water bath. Wicking height (cm) is measured after 30 minutes. The higher the value of wicking height - the better the scouring result. A value of 8 cm or higher (for heated samples) indicates good scouring. All experiments are conducted in duplicate or triplicate.

[0438] Enzyme dosages for commercial preparations are usually within the range 5-60 g preparation per kg textile. Preferably, the preparations should be normalized by protein. The dose of 0.6 mg protein per 1 ml solution (12 g protein per kg textile) proves to be suitable when comparing *T.reesei* preparations and pure enzymes using this method and micromethod B below.

[0439] When comparing preparations in one dose, 4 repetitions for each preparation (two vessels with two strips in each) are carried out. It is often more convenient to run a series of 10-16 vessels, with 1 blank (buffer treatment without enzyme) and 1-2 vessels with some standard preparation in each series.

[0440] Preferably, an alkaline scoured strip should also be added in the heating and evaluation procedure. For alkaline scouring, the strips are treated with a solution of 40 g/l NaOH in a boiling water bath for 2 hours at a liquor ratio of 5-20. Adding surfactant is typically not needed for these samples, but low doses of 1g/l of EM-3 surfactant or similar products may be beneficial.

[0441] The strips can be analyzed without heating after drying overnight. However, heating of the strips provides more reproducible result and leads to a greater difference between well and poorly scoured strips. Drying and conditioning of the strips overnight or longer after heating is an important step and should not be shortened. Results may also be improved by using a special conditioning chamber with constant humidity, rather than leaving the samples at room conditions.

Method B. Micromethod of scouring and evaluation procedure for the circular pieces of textile, diameter 1.4 cm

[0442] Treatment procedure: In this method, circular pieces (diameter 1.4 cm), of the same desized fabric from method A are used. The fabric should be completely wetted in distilled water before the analysis, with excess water removed using filter paper.

[0443] A circular piece of textile is placed in a small cylindrical vial ("penicillin vial", 2.2 cm diameter and 5 cm height) containing 2.0 ml of enzyme solution in 0.1 M Na-acetate buffer (pH 5.0). The vials are placed into a preheated water bath (50°C) on a shaker (250 rpm) and incubated for 1 hour. The fabric swatch is removed with tweezers, washed 2-3 minutes with cold tap water, and then dried at room temperature conditions overnight. For each enzyme sample and control sample (treatment with no enzyme in buffer), the assay procedure is typically carried out in 2 repetitions (2 pieces).

[0444] Heating: Overnight-dried textile swatches are heated for 1 hour at 105°C, then room conditioned overnight prior to being assayed for wettability. Heating generally provides better results, but may be omitted.

[0445] Analysis of wettability: Wettability is measured using a drop test. A drop of distilled water, 5 ul, is placed on the surface of fabric and time it takes the drop to adsorb into the material is measured. For each circular piece, 5 measurements are taken, placing each new drop in a dry area within the fabric swatch. The average value for each sample is then calculated.

[0446] Enzyme preparations and pure enzymes are tested at 0.6 mg protein per ml solution, but the assays are typically reproducible at doses between 0.6-2.4 mg protein per ml solution. This method usually gives greater deviations than method A, but 10-fold less enzyme is needed. Thus, method B is usually recommended where enzyme amounts are limiting.

[0447] Testing at one dose with heat treatment reveals preparations of 8-10 cm (e.g. classical ACE, BioACE, IndiAGE 44L, Ecoston L 350, Celloviridin) provide comparative results.

[0448] Table 28 and Figure 20 show the bioscouring ability of cellobiohydrolases and endoglucanases of the present disclosure. The reciprocal of the time ($1/\text{time}$; sec^{-1}) of water drop adsorption on a textile after treatment at pH 5.0 with 0.6 mg/l of enzyme followed by heating is shown. The water drop adsorption time for the untreated control sample is 2400 sec ($1/\text{time} = 0.0004$).

Table 28. Bioscourins Abilities of C1 Cellobiohydrolases and Endoglucanases

Enzyme	1/Time (sec ⁻¹)
Blank	0.0004
EG I	0.0006
EG II	0.0006
EG II (CD)	0.0007
EG III	0.0008
EG V	0.0007
EG VI	0.0007
CBH Ia	0.0006
CBH Ia (CD)	0.0006
CBH Ib	0.0006
CBH IIa	0.0006
CBH IIb	0.0006

Qualitative method for determination (staining) of remaining pectic substances in a scoured fabric

[0449] Ruthenium Red Dye (Sigma, R-2751) is used for specifically staining pectic substances in fabric, by forming a complex with charged carboxyl groups of pectins.

[0450] Small swatches (2.5 x 6 or 2.5 x 2.5 cm) of scoured fabric are marked using a textile marker. The scoured samples remaining after the wettability analyses described above can be used.

[0451] The following controls are included:

- 1) desized unscoured fabric,
- 2) control treated with buffer (without an enzyme),
- 3) alkaline scoured fabric.

[0452] All the samples are dyed in one bath and weighed together. The volume of the solution is calculated using the following formula:

$$V(\text{ml}) = m \text{ samples (g)} \times 30.$$

[0453] The desired volume of 1g/l solution of the dye is prepared. The dye solution is stirred and warmed in a boiling water bath for 5 minutes until the dye dissolves completely. The dye solution is warmed in a glass vessel on a water bath shaker set at 50°C, 200 rpm. The samples are then placed in the solution and some surfactant is added if the samples are not wetted. The samples are dyed for 30 minutes at 50°C with shaking at about 200 rpm for even dyeing.

[0454] Then, the samples are rinsed under tap water followed 3-4 rinses (about 5 minutes each) with hot water (about 70°C in a glass vessel). After this, the samples are dried overnight at room temperature. The samples should not be iron dried as it may destroy the dye. The samples are generally compared visually, although quantitative colorimetric evaluation can be used..

[0455] Example of stained strips are presented in Figure 21. Figure 21 illustrates a comparison of the pectin content in an unscoured sample, an alkaline scoured sample, a sample bioscoured by BioPrep (Novo Nordisk, Denmark) at 20 g preparation per 1 kg fabric, and samples bioscoured by ACE at 40 g/kg for 1 or 3 hours.

[0456] The unscoured sample and control (treated without enzyme) are stained at high and equal intensity - that means no pectins are removed during buffer treatment. Alkaline scoured sample is very faint stained - 100 % pectin removed (by definition). All three enzyme scoured samples are also faint, but little bit more intensive than alkaline scoured. This means that cellulase scouring and scouring by BioPrep removes the majority of surface (stainable) pectins from the primary wall, similar to alkaline scouring.

[0457] Other positively charged dyes can be used for staining pectins due to the negatively charged polymer in cotton. We have tested methylene blue, but it resulted in less distinction between scoured and unscoured fabric.

Method for determination of remaining waxes in scoured textiles

[0458] The presence of remaining waxes in scoured textiles is determined using an extraction procedure. 3-12 g of scoured fabric is weighed and then extracted with boiling benzene in a Soxhlet apparatus for 4 hours. The extract is evaporated in a rotary evaporator. Then, a flask containing the evaporated, extracted waxes is brought to a constant weight at 105°C (0.75-1 hour of incubation time is usually necessary). The weight of extracted waxes is then determined and the wax content is calculated.

[0459] We compared bioscouring (ACE preparation) and alkaline scouring. Desized cotton strips were treated, dried at room conditions overnight and divided into two parts: one part was evaluated for wicking height, another part were heated to 105°C for 1 hour, equilibrated at room conditions overnight and then evaluated for wicking height.

[0460] Wax content was determined by the method described above, and the results are presented in Table 29 below.

[0461] Bioscouring by ACE and by the alkaline procedure gave comparable results based on wicking height changes, but bioscouring removed less wax. Alkaline scouring, however, did not remove all of the wax.

Table 29. Bioscourins by ACE versus alkaline scouring.

Type of treatment	Wicking height, cm Room drying	Wicking height, cm heating 105°C, 1h	Waxes, %
Untreated cotton	6.0	0	0.62 ± 0.03
Alkaline scouring	13.0	8.4	0.37 ± 0.03
Bioscouring (ACE)	12.4	8.6	0.57 ± 0.03

Example 16

[0462] The following example illustrates a method of biopolishing (biofinishing) textiles using enzymes of the present disclosure and the enzyme of the invention.

[0463] The method allows for testing small amount of enzymes (mini-testing in the range of 0.5-5 mg of enzyme sample). The method for testing of pure enzymes and crude enzyme preparations is carried out in specially designed cells (essentially a micro-washing machine) also used for the mini-denim washing assay (see Figure 22B).

[0464] Sample preparation: Circular pieces of textile (diameter of 28 mm) are excised from TIC-460 cotton interlock dyed Red 80 fabric. The samples are marked with a textile marker on the side that will be treated with an enzyme and mechanical stress. Each sample is then weighed to an accuracy 1 mg. One piece generally weighs 160 to 180 mg.

[0465] Treatment procedure: The sample is fixed in the cell and three metal cylinders are placed into it. When all cells are assembled, 3 ml of enzyme solution are added to each cell. The enzyme solutions contain 0.1M Na-acetate buffer, pH 5.0, to control the pH. The cells are placed into a preheated Elpan water bath shaker (50°C, 300 rpm, amplitude 4) and the samples are treated for 1 hour.

[0466] The efficiency of biopolishing is assayed by measuring the optical density of the reaction solution. Immediately after the treatment, the solution should be thoroughly mixed, using a pipette, to resuspend any possible sediment of removed fuzz. A 1-1.5 ml aliquot of reaction solution is then taken from each cell. The optical density of the solution is then measured on a spectrophotometer at 400 nm in a 1 ml cuvette (1 cm optical path). The solution is resuspended and dosed into the cuvette and the optical density determined rapidly (within 10 seconds), as the suspension sediments gradually and the optical density changes due to sedimentation. If the value of A_{400} exceeds 1 unit, the sample is diluted 2-4 times carefully and the optical density determined once again. If a sample was diluted, the resultant value should be multiplied by the dilution factor.

[0467] Rinse and drying: The circular pieces of textile should be rinsed immediately after taking the aliquots of reaction solution, since the enzymatic reaction will proceed otherwise. The sample is removed from the cell, and washed briefly in distilled water. All the samples are then rinsed together thoroughly in distilled water for 3-5 min, stirring gently and changing water periodically. After rinsing, the samples are blotted using filter paper and dried at room temperature for at least 24 hours.

[0468] Measurement of the efficiency of biopolishing by determination of the weight loss: Each sample is weighed to a 1 mg accuracy. The resulting values are compared with that of the same sample before the treatment and weight loss is calculated as below:

$$\text{Weight loss (mg)} = m \text{ before treatment (mg)} - m \text{ after treatment (mg)}$$

$$\% \text{ Weight loss} = \text{Weight loss/m before treatment (mg)} \times 100\%$$

[0469] Qualitative evaluation of the surface appearance: The image of the edge of the folded fabric is taken using suitable device at a magnification about x 60. Images of the samples treated by different enzymes and blank runs are compared visually.

[0470] In detail, the circular piece is folded (for all samples in a same manner- e.g., so that the surface threads are perpendicular to the edge line). All samples should be folded so that the treated surface is up. The samples can be stapled to a piece of paper for more convenient operation. Images are then taken using a QX3 Intel Play microscope at x 60 magnification and recorded in JPEG format.

[0471] The samples should be run in series up to 8 samples in each. In each series, there should be a control treatment (without an enzyme) and a treatment with a standard preparation. For each preparation under investigation, at least 2 repetitions should be made.

[0472] Weight determination is influenced by the drying conditions. Drying of the samples before the treatment and after in reproducible conditions is recommended (e.g., heating in an oven and drying in a dessicator). Drying at room conditions can result in slight deviations between series run on different days.

[0473] The effect of enzymatic treatment along with mechanical stress in our test is similar to the widely used Launderometer and other laboratory testing machines, but it has different geometry. Due to this geometry, the system is amenable to further miniaturization and require less than 1 mg quantities of an enzyme.

[0474] Evaluation of the biopolished fabric can be problematic. The most important (and direct) parameters are pilling resistance and fuzziness of a treated fabric. Weight loss, color clarification, and improvement of softness are less direct but valuable parameters. Greater pilling resistance correlates with greater weight losses and gives less fuzz, clarification of color and softer feel. Fuzziness is evaluated qualitatively in our method on 3 levels: good, moderate and poor. Optical density should correlate directly both to weight loss (weight loss is due mainly to the removed fuzz making turbid suspension in a solution) and fuzziness of fabric.

[0475] Table 30 and Figure 23 illustrate the biopolishing ability of cellobiohydrolases and endoglucanases of the present disclosure. The dye release (D 400 nm) after treatment with 0.2 g/l enzyme at pH 5.0 and 50°C is shown.

Table 30. Biopolishing Abilities of C1 Cellobiohydrolases and Endoglucanases

Enzyme	% Dye Release
EG I	0.17
EG II	0.2
EG II (CD)	0.1
EG III	0.16
EG V	0.2
EG VI	0.1
CBH Ia	0.1
CBH Ia (CD)	0.2
CBH Ib	0.3
CBH IIa	0.2
CBH IIb	0.1

Example 17

[0476] The following example illustrates methods of biostoning (abrasion) textiles using enzymes of the present disclosure and the enzyme of the invention.

[0477] The method allow for testing small amount of enzymes (mini-testing in the range of 0.5-5 mg of enzyme sample). The method is available in two scales, requiring approximately 10 and 1 mg of protein, respectively (Method A and Method B). Both scales utilize the same principle and gives comparable results. Method B requires specially designed cells but proved easier when assaying pure enzymes.

Method A

[0478] 250 ml vessels (internal diameter 56 mm, height 110 mm) with Teflon disks at the bottom (thickness 4 mm) are used for the treatment of denim fabric with cellulase (see Figure 22A). A swatch of desized denim fabric (10 x 10 cm) is pulled over the butt of a plastic tube (diameter 50 mm, height 100 mm) and fixed with a rubber ring. The tube is inserted into the 250 ml vessel so that the swatch attaches tightly and uniformly to the Teflon disk at the bottom of the glass. Ten metal buck-shot balls (diameter 7 mm) are placed inside the tube to provide mechanical stress to the surface of fabric, and 30 ml of enzyme solution in the appropriate buffer is added. The vessel is placed onto a ELPAN water bath shaker type 357 (Poland) at 50°C. The enzymatic treatment of the denim is carried out for 1 hour at the shaker speed of 300 rpm and amplitude of vibrations adjusted to 4 units. The swatch is then removed and washed for 1 minute with running water. An excess of water on the surface of the swatch is removed by squeezing it between two sheets of blotting paper, and then the swatch is dried overnight in the open air at room temperature. In a separate experiment, a control assay is carried out as described above, but in the absence of enzyme. For the control, 30 ml of the same buffer used for enzyme dilution is added to the vessel.

Method B

[0479] This assay procedure uses specially designed cylindrical cells (external diameter 35 mm, internal diameter 30 mm, height 48 mm) made of stainless steel and equipped with tightened lids (see Figure 22B). This method also uses 10-fold less enzyme than Method A. A circular swatch of denim fabric (diameter of 30 mm) is placed at the bottom of a cell, and a stainless steel cylindrical cartridge is inserted to clamp the swatch. Three metal cylinders (7 x 7 mm), made of stainless steel, are placed into the cell to provide mechanical stress, and 3 ml of enzyme solution in the appropriate buffer is added. A lid, equipped with a rubber ring inside, is quickly tightened to seal the cell hermetically, and the cell is placed in a water bath on a shaker. Then, the procedure is carried under the same conditions as described above for Method A. In the control assay, afacric swatch is treated under the same conditions but in the absence of enzyme. To this end, 3 ml of the same buffer used for enzyme dilution is added to the cell.

[0480] Both assays (Method A or Method B) are carried out at least in two repetitions for each particular enzyme (usually three repetitions). Typically, when different cellulase preparations are compared, the conditions in the reaction system are equalized by the CMCase activity (1-3 U/ml). Acid cellulase samples are assayed at pH 5.0 using 0.1 M acetate buffer. Neutral cellulases are assayed at pH 6.0 using 0.1 M phosphate buffer.

[0481] Analysis of color intensity on denim swatches is carried out using a Mustek MFS-12000SP (USA) scanner. A circular area on the swatch, subjected to the combined enzyme and mechanical action as above, is scanned at 300 dpi resolution. This area is usually more light than the denim surface near the swatch edges. Each swatch is scanned twice: first, when the threads on the surface of the fabric are parallel to the movement of the light source on scanning, and second, when the threads are directed perpendicular to the movement of the light source on scanning.

[0482] The JPG images obtained as a result of scanning are analyzed using Adobe Photoshop software (versions 3.0 or later can be used). For each particular scan, a histogram is opened. The blue channel is selected and the percentile at level 32 (P32) is read. This parameter shows the percentage of image pixels having the intensity of color higher than at the level 32 of histogram (a total of 256 shades (levels) of blue color as the abscissa axis). The higher is the P32, the darker is the denim swatch. On the contrary, the parameter (100 - P32, %) shows the percentage of pixels having the intensity of color lower than at the level 32 of the histogram. Figure 24 shows an example histogram generated from the color intensity analysis of a denim swatch.

[0483] For each particular enzyme tested, an average value of the P32 and standard deviation are calculated using the data of all scans for this enzyme. For example, if the assay is performed in two repetitions, four P32 values are taken into account, since for each repetition two scans are carried out.

[0484] The difference between the average value of P32 obtained in the control (without cellulase) and the average value of P32 for the enzyme is taken as a criterion of the abrasive activity of the enzyme.

[0485] Table 31 and Figure 25 illustrate the denim-washing (abrasion) ability of cellobiohydrolases and endoglucanases of the present disclosure using Assay B above. The relative units per mg of enzyme determined at 50°C, pH 5.0 is shown.

Table 31. Abrasion Abilities of C1 Cellobiohydrolases and Endoglucanases

Enzyme	Relative Units per mg
EG I	14
EG II	62
EG II (CD)	54

(continued)

Enzyme	Relative Units per mg
EG III	25
EG V	65
EG VI	12
CBH Ia	3
CBH Ia (CD)	11
CBH Ib	10
CBH IIa	22
CBH IIb	15

[0486] Additional examples and discussion concerning methods A and B are presented in A.Gusakov, A.Sinitsyn, S.Grishutin, D.Tikhomirov, D.Shook, D.Sheer, M.Emlfarb Microassays to control the results of cellulase treatment of denim fabrics; Textile Chemist and Colorist and American Dyestuff Reporter, (2000), V.32, N. 5, P.42.

Example 18

[0487] The following example illustrates methods of backstaining textiles using enzymes of the present disclosure and the enzyme of the present invention.

[0488] The methods allow for testing small amount of enzymes (mini-testing in the range of 0.5-5 mg of enzyme sample).

[0489] Standard Indigo backstaining Method A requires larger enzyme quantities (typically 2-5 mg of protein for one assay performed with 3.5 x 3.5 cm swatches in 2-3 repetitions). When purification of enzyme components is performed on a laboratory scale and it is necessary to assay the backstaining index (BSI) for the purified enzyme, much lower quantities of an enzyme are typically available. In such case, a modification of this method (Method B) should be used, where smaller swatches of standard garment (1.4 x 1.4 cm) are used, and the total volume of Indigo suspension is 2 ml. This minimized microassay procedure requires ten times lower amounts of protein (0.2-0.5 mg).

Method A

[0490] Indigo reagent preparation: 250 mg of indigo is added to 50 ml of distilled water in a flask and stirred vigorously for 15-20 minutes on a magnetic stirrer. Before taking aliquots for staining experiments, the suspension is placed into a Bandelin SONOREX TK52 ultrasonic bath for 5 minutes to disrupt large indigo aggregates and to make the suspension more uniform. 2 ml aliquots are used for staining experiments (see below) under magnetic stirring in order to provide a uniform indigo suspension in the reaction mixture.

[0491] Assay procedure: A piece of white fabric (3.5 x 3.5 cm) is added to 18 ml of 0.1 M Na-acetate buffer (pH 5.0) in a 250 ml glass vessel and incubated at room temperature for 10 minutes (control). A solution of enzyme preparation in the same amount of the buffer is used in the enzyme assay procedure. 2 ml of indigo suspension (5 mg/ml) is then added, and the vessel is placed into a water bath (50°C) on a shaker (300 rpm) and agitated for 30 min. The piece of fabric is removed with tweezers and washed twice for 5 minutes with 50 ml of distilled water (on a shaker at 300 rpm) and then dried at room temperature overnight. For each enzyme sample and control, the assay procedure is carried out at least in 3 repetitions.

[0492] Samples of stained fabric are scanned at 300 dpi resolution on a "Paragon" ("Muztek", USA) scanner. Two images of the front side and two images of the back side (one direct and one rotated 90°) are obtained for each sample. Images are processed using an Adobe Photoshop software. A histogram of colour intensities is calculated for each sample using a blue channel, and a percentile at level 120 is used as a criterion of staining. An average value and standard deviation are calculated for each enzyme preparation and control.

[0493] *T.reesei* (ACE) is usually used as a control (internal standard) for analyzing backstaining properties of a particular enzyme sample, since ACE is characterized by a high backstaining index. If possible, a IndiAge Super GX sample should be used as a second internal standard since it has a low backstaining index.

Method B

[0494] The Indigo stock suspension (5 g/l) is prepared and the image processing is carried out as described above.

[0495] Assay procedure: A piece of white fabric (1.4 x 1.4 cm) is placed in a small cylindrical vial ("penicillin vial", 2.2 cm diameter and 5 cm height) containing 1.8 ml of enzyme solution in 0.1 M Na-acetate buffer (pH 5.0) and incubated at room temperature for 10 min. 0.2 ml of Indigo suspension is then added, and the vial is placed into a water bath (50°C) on a shaker (300 rpm) and agitated for 30 minutes. The fabric swatch is removed with tweezers and washed twice for 5 minutes with 5 ml of distilled water at 50°C (on a shaker at 300 rpm), then dried at room temperature overnight. For each enzyme sample and control (internal standard), the assay procedure is typically carried out in 3 repetitions. In most experiments, 1.5 CMCase units per ml of reaction system are used. In cases where specific CMCase activity is very low (or absent), 0.05-0.1 mg of protein per ml of reaction system is used.

[0496] It is convenient to place three vials (3 repetitions) into a 250 ml vessel containing 15-20 ml of water at the bottom and then to place the vessel into a water bath on a shaker.

[0497] Additional examples and discussion concerning methods A and B are presented in A.Gusakov, A.Sinitsyn, S.Grishutin, D.Tikhomirov, D.Shook, D.Sheer, M.Emalfarb Microassays to control the results of cellulase treatment of denim fabrics; Textile Chemist and Colorist and American Dyestuff Reporter, (2000), V.32, N. 5, P.42.

Example 19

[0498] The following example illustrates assays used to measure acetyl esterase enzymatic activity.

Activity towards *p*-nitrophenyl acetate substrate

[0499] This assay measures the release of *p*-nitrophenol by the action of an acetyl esterase on *p*-nitrophenyl acetate (PNPac). One acetyl esterase unit of activity is the amount of enzyme that liberates 1 micromole of *p*-nitrophenol in one minute at 37°C and pH 7.2.

[0500] Phosphate buffer (0.01 M, pH 7.2) is prepared as follows: 0.124 g of NaH₂PO₄ * H₂O and 0.178 g Na₂HPO₄ are dissolved in distilled water so that the final volume of the solution 500 ml and the pH of the resulting solution is 7.2.

[0501] PNPac (Fluka Chemie, Switzerland, cat. # 46021) is used as the assay substrate. 3.6 mg of PNPac is dissolved in 10 ml of 0.01 M phosphate buffer using a magnetic stirrer to obtain a 2 mM stock solution. The solution is stable for 2 days with storage at 4 °C.

[0502] The stop reagent (0.25 M Tris-HCl, pH 8.5) is prepared as follows: 30.29 g of Tris is dissolved in 900 ml of distilled water (Solution A). The final 0.25 M Tris-HCl pH 8.5 is prepared by mixing solution A with 37% HCl until the pH of the resulting solution is equal to 8.5. The solution volume is then adjusted to 1000 ml. This reagent is used to terminate the enzymatic reaction. Using the above reagents, the assay is performed as detailed below.

[0503] For the enzyme sample, 0.10 ml of 2 mM PNPac stock solution is mixed with 0.01 ml of an enzyme sample and incubated at 37°C for 5 minutes (Axe2 and Axe3) or 10 minutes (Axe1). After exactly 5 or 10 minutes of incubation, 0.1 ml of 0.25 M Tris-HCl solution is added and then the absorbance at 405 nm (A₄₀₅) is measured in microtiter plates as A_S.

[0504] For the substrate blank, 0.10 mL of 2 mM PNPac stock solution is mixed with 0.01 ml of 0.01 M phosphate buffer, pH 7.2. 0.1 mL of 0.25 M Tris-HCl solution is then added and the absorbance at 405 nm (A₄₀₅) is measured in microtiter plates as A_{SB}.

[0505] Activity is calculated as follows:

$$\text{Activity (IU/ml)} = \frac{\Delta A_{405} * \text{DF} * 21 * 1.33}{13.700 * 5}$$

where $\Delta A_{405} = A_S - A_{SB}$, DF is the enzyme dilution factor, 21 is the dilution of 10 µl enzyme solution in 210 µl reaction volume, 1.33 is the conversion factor of microtiter plates to cuvettes, 13.700 is the extinction coefficient (13700 M⁻¹ cm⁻¹ of *p*-nitrophenol released corrected for mol/L to µmol/mL), and 5 minutes is the reaction time.

[0506] Using the assay above, acetyl esterase activity of Axe1 was found to be 0.23 IU/mL ($\Delta A_{405} = 1.11$, DF=1), Axe2 was found to be 2.80 U/ml ($\Delta A_{405} = 0.1714$, DF=40), and Axe3 was found to be 39.97 U/ml ($\Delta A_{405} = 0.4901$, DF=200).

Activity towards arabinoxylan oligosaccharide substrates

[0507] This assay measures the release of acetate by the action of a xylan acetyl esterase on the arabinoxylan oligosaccharides from *Eucalyptus* wood.

[0508] 0.01 M sodium hydroxide is prepared as follows: 0.4 g of sodium hydroxide is dissolved in distilled water so that the final volume of the solution to be 1000 ml

[0509] Arabinoxylan oligosaccharides from *Eucalyptus* wood were prepared by dissolving 5 mg of arabinoxylan oli-

gosaccharides in 1 ml distilled water using a magnetic stirrer as described in Kabel et al. Carbohydr. Polym. 50:191 (2002).

[0510] For the enzyme sample, 1.0 mL of arabinoxylan oligosaccharides stock solution is mixed with 0.005 mg of the enzyme sample and incubated at 35 °C. The pH was maintained at 5.0 by the addition of 0.01 M NaOH. The added volume of 0.01 M NaOH is proportional to the breakdown of ester linkage and release of acetate.

[0511] For the substrate blank, 1.0 ml of arabinoxylan oligosaccharides stock solution is mixed with 0.005 mg of distilled water and incubated at 35°C. The pH was maintained at 5.0 by addition of 0.01 M NaOH. The added volume of 0.01 M NaOH is proportional to the breakdown of ester linkage and release of acetate.

[0512] Using the assay above, both Axe2 and Axe3 were found to exhibit xylan acetyl esterase activity, as measured by the release of acetate from eucalyptus wood arabinoxylan oligosaccharides. For Axe2, 0.7 ml NaOH was used after 200 minutes of incubation. For Axe3, 0.35 ml was used after 200 minutes.

Example 20

[0513] The following example illustrates assays used to measure ferulic acid esterase enzymatic activity.

Activity towards *p*-nitrophenyl butyrate substrate

[0514] This assay measures the release of *p*-nitrophenol by the action of a ferulic acid esterase on *p*-nitrophenyl butyrate (PNPBU). One acetyl esterase unit of activity is the amount of enzyme that liberates 1 micromole of *p*-nitrophenol in one minute at 37°C and pH 7.2.

[0515] Phosphate buffer (0.01 M, pH 7.2) is prepared as follows: 0.124 g of NaH₂PO₄ * H₂O and 0.178 g Na₂HPO₄ are dissolved in distilled water so that the final volume of the solution is 500 ml and the pH of the resulting solution is equal to 7.2.

[0516] PNPBU (Sigma, USA, cat. # N9876-5G) is used as the assay substrate. 10 µl of PNPBU is mixed with 25 ml of 0.01 M phosphate buffer using a magnetic stirrer to obtain a 2 mM stock solution. The solution is stable for 2 days with storage at 4°C.

[0517] The stop reagent (0.25 M Tris-HCl, pH 8.5) is prepared as follows: 30.29 g of Tris is dissolved in 900 ml of distilled water (Solution A). The final 0.25 M Tris-HCl pH 8.5 is prepared by mixing solution A with 37% HCl until the pH of the resulting solution is equal to 8.5. The solution volume is adjusted to 1000 ml. This reagent is used to terminate the enzymatic reaction. Using the above reagents, the assay is performed as detailed below.

[0518] For the enzyme sample, 0.10 ml of 2 mM PNPBU stock solution is mixed with 0.01 ml of the enzyme sample and incubated at 37°C for 10 minutes. After exactly 10 minutes of incubation, 0.1 ml of 0.25 M Tris-HCl solution is added and then the absorbance at 405 nm (A₄₀₅) is measured in microtiter plates as A_S.

[0519] For the substrate blank, 0.10 ml of 2 mM PNPBU stock solution is mixed with 0.01 ml of 0.01 M phosphate buffer, pH 7.2. 0.1 ml of 0.25 M Tris-HCl solution is then added and the absorbance at 405 nm (A₄₀₅) is measured microtiter plates as A_{SB}.

[0520] Activity is calculated as follows:

$$\text{Activity (IU/ml)} = \frac{\Delta A_{405} * \text{DF} * 21 * 1.33}{13.700 * 10}$$

where $\Delta A_{405} = A_S - A_{SB}$, DF is the enzyme dilution factor, 21 is the dilution of 10 µl enzyme solution in 210 µl reaction volume, 1.33 is the conversion factor of microtiter plates to cuvettes, 13.700 is the extinction coefficient (13700 M⁻¹ cm⁻¹ of *p*-nitrophenol released corrected for mol/l to µmol/ml), and 10 minutes is the reaction time.

[0521] Using the assay above, the ferulic acid esterase activity of FaeA1 was found to be 18.5 IU/ml ($\Delta A_{405} = 0.906$, DF=10), FaeA2 was found to be 0.40 U/ml ($\Delta A_{405} = 1.976$, DF=1) and FaeB2 was found to be 0.02 U/ml ($\Delta A_{405} = 0.111$, DF=1).

Activity towards *p*-nitrophenyl butyrate substrate

[0522] The following assay is used to measure the enzymatic activity of a ferulic acid esterase towards wheat bran (WB) oligosaccharides by measuring the release of ferulic acid.

[0523] Wheat bran oligosaccharides are prepared by degradation of wheat bran (obtained from Nedalco, The Netherlands) by endo-xylanase III from *A. niger* (enzyme collection Laboratory of Food Chemistiy, Wageningen University, The Netherlands). 50 mg of WB is dissolved in 10 ml of 0.05 M acetate buffer pH 5.0 using a magnetic stirrer. 1.0 ml of WB stock solution is mixed with 0.0075 mg of the enzyme and incubated at 35°C for 24 hours. The reaction is stopped by heating the samples for 10 minutes at 100°C. The residual material is removed by centrifugation (15 minutes at 14000

rpm), and the supernatant is used as the substrate in the assay detailed below.

[0524] For the enzyme sample, 1.0 ml of wheat bran oligosaccharides stock solution is mixed with 0.005 mg of the enzyme sample and incubated at 35°C for 24 hours. The reaction is stopped by heating the samples for 10 minutes at 100°C. The release of ferulic acid is analyzed by measuring the absorbance at 335 nm.

[0525] For the substrate blank, 1.0 ml of wheat bran oligosaccharides stock solution is mixed with 0.005 mg of 0.05 M acetate buffer, pH 5.0, and incubated at 35°C for 24 hours. The reaction is stopped by heating the samples for 10 minutes at 100°C. The release of ferulic acid is analyzed by measuring the absorbance at 335 nm.

[0526] Using the assay above, FaeA1 was found to release ferulic acid from wheat bran arabinoxylan oligosaccharides, releasing 75% of the ferulic acid present in wheat bran. FaeA1 also exhibited activity towards sugar beet pulp oligosaccharides (20% hydrolysis). FaeB2 was found to release 80% of the ferulic acid present in wheat bran. FaeB2 also exhibited activity towards sugar beet pulp oligosaccharides (16% hydrolysis).

Example 21

[0527] The following example illustrates an assay used to measure α -arabinofuranosidase enzymatic activity.

[0528] This assay measures the release of arabinose by the action of the α -arabinofuranosidase on wheat arabinoxylan oligosaccharides (WAX).

[0529] Acetate buffer (0.05 M, pH 5.0) is prepared as follows: 4.1 g of anhydrous sodium acetate or 6.8 g of sodium acetate * 3H₂O is dissolved in distilled water so that the final volume of the solution to be 1000 ml (Solution A). In a separate flask, 3.0 g (2.86 mL) of glacial acetic acid is mixed with distilled water to make the total volume 1000 ml (Solution B). The final 0.05 M acetate buffer, pH 5.0, is prepared by mixing Solution A with Solution B until the pH of the resulting solution is equal to 5.0.

[0530] Wheat arabinoxylan oligosaccharides are prepared by degradation of Wheat arabinoxylan (Megazyme, Bray Ireland, Cat. #P-WAXYI) by endo-xylanase I from *A. niger* (enzyme collection Laboratory of Food Chemistry, Wageningen University, The Netherlands). 50 mg of WAX is dissolved in 10 ml 0.05 M acetate buffer, pH 5.0, using a magnetic stirrer. 1.0 ml of WAX stock solution is mixed with 0.0075 mg of the enzyme and incubated at 35°C for 24 hours. The reaction is stopped by heating the samples for 10 minutes at 100°C. The residual material is removed by centrifugation (15 minutes at 14000 rpm), and the supernatant is used as the substrate in the assay detailed below.

[0531] For the enzyme sample, 1.0 ml of wheat arabinoxylan oligosaccharides stock solution is mixed with 0.0075 mg of the enzyme sample and incubated at 35°C for 24 hours. The reaction is stopped by heating the samples for 10 minutes at 100°C. The release of arabinose and formation of new arabinoxylan oligosaccharides were analyzed by High Performance Anion Exchange Chromatography.

[0532] For the substrate blank, 1.0 ml of wheat arabinoxylan oligosaccharides stock solution is mixed with 0.0075 mg of the 0.05 M acetate buffer, pH 5.0, and incubated at 35°C for 24 hours. The reaction is stopped by heating the samples for 10 minutes at 100°C. The release of arabinose and formation of new arabinoxylan oligosaccharides were analyzed by High Performance Anion Exchange Chromatography.

[0533] High Performance Anion Exchange Chromatography analysis is performed using a Dionex HPLC system equipped with a Dionex CarboPac PA-1 (2 mm ID x 250 mm) column in combination with a CarboPac PA guard column (1 mm ID x 25 mm) and a Dionex EDetI PAD-detector (Dionex Co., Sunnyvale, CA, USA). A flow rate of 0.3 ml/min was used with the following gradient of sodium acetate in 0.1 M NaOH: 0-5 min, 0-100 mM; 5-45 min, 100-400 mM. Each elution was followed by a washing step of 5 minutes 1,000 mM sodium acetate in 0.1 M NaOH and an equilibration step of 15 minutes 0.1 M NaOH.

[0534] Using the above assay, both Abf1 and Abf2 were found to release arabinose from wheat arabinoxylan oligosaccharides. Both enzymes removed arabinose residues, which were monosubstituted to the xylose backbone. Arabinose residues linked to O2 and O3 position of the xylose residue were released by both Abf1 and Abf2.

Example 22

[0535] The following example illustrates an assay used to measure pectin methyl esterase enzymatic activity.

[0536] The following assay is used to measure the enzymatic activity of a pectin methyl esterase towards highly methylated citrus pectin (DM 65) in combination with methyl red. The assay measures the release of methanol by the action of pectin methyl esterases on methylated citrus pectin. The release of methanol will acidify a solution (non-esterified galacturonic acid), which makes the methyl red indicator change color at a pH of less than 4.4.

[0537] Pectin-methyl red solution is prepared as follows: 0.5 g of methylated citrus pectin (DM 65, Sigma, USA, Cat. # P9436-5G), 2 mg methyl red (Sigma, USA) and 0.435 g NaCl is dissolved in distilled water so that the final volume of the solution is 50 ml. The final pH is adjusted to 5.0 by adding 0.1 M NaOH.

[0538] For the enzyme sample, 0.1 ml of pectin-methyl red stock solution is mixed with 0.01 ml of the enzyme sample and incubated at 37°C overnight. The absorbance at 520 nm (A_{520}) is measured as A_s .

[0539] For the substrate blank, 0.1 ml of pectin/methyl red stock solution is mixed with 0.01 ml of distilled water and incubated at 37°C overnight. The absorbance at 520 nm (A_{520}) is measured as A_{SB} .

[0540] Using the assay above, the ΔA_{520} of Pme1 was found to be 0.401, demonstrating that Pme1 is active towards highly methylated pectin.

Example 23

[0541] The following example illustrates an assay used to measure endo-arabinase enzymatic activity.

[0542] The assay is used to measure the enzymatic activity of an endo-arabinase towards AZCL-arabinan (debranched). This substrate is insoluble in buffered solutions, but rapidly hydrates to form gel particles that are hydrolysed by specific *endo*-hydrolases, releasing soluble dye-labeled fragments.

[0543] Acetate buffer (0.05 M, pH 5.0) is prepared as follows: 4.1 g of anhydrous sodium acetate or 6.8 g of sodium acetate * 3H₂O is dissolved in distilled water so that the final volume of the solution is 1000 ml (Solution A). In a separate flask, 3.0 g (2.86 ml) of glacial acetic acid is mixed with distilled water to yield a total volume of 1000 ml (Solution B). The final 0.05 M acetate buffer, pH 5.0, is prepared by mixing Solution A with Solution B until the pH of the resulting solution is equal to 5.0.

[0544] AZCL-arabinan (debranched) from Megazyme (Bray, Ireland, Cat. # I-AZDAR) is used as the assay substrate. 50 mg of AZCL-arabinan is dissolved in 10 ml of 0.05 M acetate buffer, pH 5.0, using a magnetic stirrer. 96% Ethanol is used to terminate the enzymatic reaction.

[0545] For the enzyme sample, 0.2 ml of 5 mg/ml AZCL-arabinan stock solution is preheated to 40°C for 10 minutes. This preheated stock solution is mixed with 200 µl of the enzyme sample (preheated at 40°C for 10 minutes) and incubated at 40°C for 10 minutes. After exactly 10 minutes of incubation, 1.0 ml of 96% ethanol is added and the absorbance at 590 nm (A_{590}) is then measured as A_S .

[0546] For the substrate blank, 1.2 ml of 5 mg/ml AZCL-arabinan stock solution is preheated at 40°C for 10 minutes. This preheated stock solution is mixed with 200 µl of 0.05 M acetate buffer pH 5.0 (preheated at 40°C for 10 minutes) and incubated at 40°C for 10 minutes. After exactly 10 minutes of incubation, 1.0 ml of 96% ethanol is added and the absorbance at 590 nm (A_{590}) is then measured as A_{SB} .

[0547] Endo-arabinase activity is determined using a standard curve produced from an endo-arabinase with known activity towards AZCL-arabinan.. Activity (IU/ml) = $\Delta A_{595} \cdot DF$,

where $\Delta A_{595} = A_S - A_{SB}$ and DF is the enzyme dilution factor.

[0548] Using the above assay, the ΔA_{595} was found to be 1.53 with a DF of 1, indicating that Abnl exhibits endo-arabinase activity towards AZCL-arabinan.

Example 24

[0549] The following example illustrates an assay used to measure β -xylosidase enzymatic activity.

[0550] This assay measures the release of *p*-nitrophenol by the action of a β -xylosidase on *p*-nitrophenyl β -D-xylopyranoside (PNPX). One β -xylosidase unit of activity is the amount of enzyme that liberates 1 micromole of *p*-nitrophenol in one minute at 37°C and pH 5.0.

[0551] Acetate buffer (0.1 M, pH 5.0) is prepared as follows: 8.2 g of anhydrous sodium acetate or 13.6 g of sodium acetate * 3H₂O is dissolved in distilled water so that the final volume of the solution is 1000 ml (Solution A). In a separate flask, 6.0 g (5.72 ml) of glacial acetic acid is mixed with distilled water to make the total volume of 1000 ml (Solution B). The final 0.1 M acetate buffer, pH 5.0, is prepared by mixing Solution A with Solution B until the pH of the resulting solution is equal to 5.0.

[0552] PNPX (Extrasynthese, France, Cat. # 4244) is used as the assay substrate. 16.5 mg of PNPX is dissolved in 5 ml of distilled water and 5 ml 0.1 M acetate buffer using a magnetic stirrer to obtain 2 mM stock solution. The solution is stable for 2 days with storage at 4°C.

[0553] The stop reagent (0.25 M sodium carbonate solution) is prepared as follows: 26.5 g of anhydrous sodium carbonate is dissolved in 800 ml of distilled water, and the solution volume is adjusted to 1000 ml. This reagent is used to terminate the enzymatic reaction.

[0554] For the enzyme sample, 0.10 ml of 2 mM PNPX stock solution is mixed with 0.01 ml of the enzyme sample and incubated at 37°C for 30 minutes. After exactly 30 minutes of incubation, 0.1 ml of 0.25 M sodium carbonate solution is added and the absorbance at 405 nm (A_{405}) is then measured in microtiter plates as A_S .

[0555] For the substrate blank, 0.10 ml of 2 mM PNPX stock solution is mixed with 0.01 ml of 0.05 M acetate buffer, pH 5.0. 0.1 mL of 0.25 M sodium carbonate solution is then added and the absorbance at 405 nm (A_{405}) is measured microtiter plates as A_{SB} .

[0556] Activity is calculated as follows:

$$\text{Activity (IU/ml)} = \frac{\Delta A_{405} * \text{DF} * 21 * 1.33}{13.700 * 30}$$

where $\Delta A_{405} = A_S - A_{SB}$, DF is the enzyme dilution factor, 21 is the dilution of 10 μl enzyme solution in 210 μl reaction volume, 1.33 is the conversion factor of microtiter plates to cuvettes, 13.700 is the extinction coefficient $13700 \text{ M}^{-1} \text{ cm}^{-1}$ of *p*-nitrophenol released corrected for mol/l to $\mu\text{mol/ml}$, and 30 minutes is the reaction time.

[0557] Using the above assay, Bx12 was found to exhibit a β -xylosidase activity of 0.09 IU/ml ($\Delta A_{405} = 1.36$, DF=1).

Example 25

[0558] The following example illustrates an assay used to measure the saccharification ability of enzymes of the present disclosure and the enzyme of the present invention.

[0559] The hydrolysis of 5 g/l Avicel catalyzed by purified cellulases was performed for 120 hours at 40°C and pH 5.0. Avicel substrate (10 mg) was mixed with 1.0 ml of 0.1 M Na-acetate buffer, pH 5.0, containing 1 mM of sodium azide to prevent microbial growth, in a 2 ml vial. The vial was incubated in a thermostatic water bath equipped with a magnetic stirrer. The enzyme solution (1 ml), diluted as needed to reach a particular concentration, was added and the mixture was stirred. Equal protein concentrations (0.1 mg/ml) were used for all enzyme preparations. To convert celooligosaccharides, which are soluble products of cellulose hydrolysis catalyzed by endoglucanases and cellobiohydrolases, into a final hydrolysis product (glucose) and thus simplify the analysis of hydrolyzate, highly purified beta-glucosidase from *A. japonicas* (0.2 U/ml; see below) was added to the reaction mixture. Aliquots of the suspension (0.1 ml) were collected from the reaction mixture and centrifuged after fixed time intervals. Glucose concentration in the supernatant was determined by the glucose oxidase-peroxidase method (see Example 6).

[0560] Beta-glucosidase was purified from *A. japonicas* culture filtrates using anion-exchange chromatography on SOURCE Q media (pH 5.5, gradient of NaCl; GE Healthcare, USA) followed by hydrophobic chromatography on Phenyl SUPEROSE media (GE Healthcare, USA) in a gradient of ammonium sulfate.

[0561] Table 32 and Figure 26 show a graph of the saccharification ability of cellobiohydrolases and endoglucanases of the present disclosure. The amount of glucose (g/l) produced after the hydrolysis of 5 g/l Avicel at pH 5.0, 40°C for 120 hours in the presence of 0.1 mg/ml enzyme and 0.2 U/ml beta-glucosidase is shown.

Table 32. Saccharification Abilities of C1 Cellobiohydrolases and Endoglucanases

Enzyme	Glucose Yield (g/l)
EG I	1.2
EG II	0.97
EG II (CD)	0.4
EG III	0.42
EG V	0.51
EG VI	0.56
CBH Ia	2.54
CBH Ia (CD)	1.31
CBH Ib	1.25
CBH IIa	1.16
CBH IIb	3.2

SEQUENCE LISTING

[0562]

<110> Dyadic International, Inc.
Nedwin, Glenn

EP 2 183 363 B1

<120> Novel Fungal Enzymes

<130> 5671-5-PCT

5 <150> 11 833,133
<151> 2007-08-02

<160> 94

10 <170> PatentIn version 3.5

<210> 1

<211> 1836

<212> DNA

15 <213> Chrysosporium lucknowense

<220>

<221> CDS

<222> (98) .. (506)

20

<220>

<221> CDS

<222> (574)..(1742)

25 <400> 1

atataaattg ggtgattccc agctcttgat gggcgtgtct tctgcctggc agccctcgtc 60

30 ttcagatcaa gcaactgtgt gctgacctc ttccgcc atg tac gcc aag ttc gcg 115
Met Tyr Ala Lys Phe Ala
1 5

acc ctc gcc gcc ctt gtg gct ggc gcc gct gct cag aac gcc tgc act 163
Thr Leu Ala Ala Leu Val Ala Gly Ala Ala Ala Gln Asn Ala Cys Thr
10 15 20

ctg acc gct gag aac cac ccc tcg ctg acg tgg tcc aag tgc acg tct 211
Leu Thr Ala Glu Asn His Pro Ser Leu Thr Trp Ser Lys Cys Thr Ser
25 30 35

40 ggc ggc agc tgc acc agc gtc cag ggt tcc atc acc atc gac gcc aac 259
Gly Gly Ser Cys Thr Ser Val Gln Gly Ser Ile Thr Ile Asp Ala Asn
40 45 50

45 tgg cgg tgg act cac cgg acc gat agc gcc acc aac tgc tac gag ggc 307
Trp Arg Trp Thr His Arg Thr Asp Ser Ala Thr Asn Cys Tyr Glu Gly
55 60 65 70

50 aac aag tgg gat act tcg tac tgc agc gat ggt cct tct tgc gcc tcc 355
Asn Lys Trp Asp Thr Ser Tyr Cys Ser Asp Gly Pro Ser Cys Ala Ser
75 80 85

aag tgc tgc atc gac ggc gct gac tac tcg agc acc tat ggc atc acc 403
Lys Cys Cys Ile Asp Gly Ala Asp Tyr Ser Ser Thr Tyr Gly Ile Thr
90 95 100

55

EP 2 183 363 B1

	acg agc ggt aac tcc ctg aac ctc aag ttc gtc acc aag ggc cag tac	451
	Thr Ser Gly Asn Ser Leu Asn Leu Lys Phe Val Thr Lys Gly Gln Tyr	
	105 110 115	
5	tcg acc aac atc ggc tcg cgt acc tac ctg atg gag agc gac acc aag	499
	Ser Thr Asn Ile Gly Ser Arg Thr Tyr Leu Met Glu Ser Asp Thr Lys	
	120 125 130	
10	tac cag a gtaagttcct ctgcacccg gccgccggga gatgatggcg cccagcccgc	556
	Tyr Gln	
	135	
15	tgacgcgaat gacacag tg ttc cag ctc ctc ggc aac gag ttc acc ttc	605
	Met Phe Gln Leu Leu Gly Asn Glu Phe Thr Phe	
	140 145	
20	gat gtc gac gtc tcc aac ctc ggc tgc ggc ctc aat ggc gcc ctc tac	653
	Asp Val Asp Val Ser Asn Leu Gly Cys Gly Leu Asn Gly Ala Leu Tyr	
	150 155 160	
25	ttc gtg tcc atg gat gcc gat ggt ggc atg tcc aag tac tcg ggc aac	701
	Phe Val Ser Met Asp Ala Asp Gly Gly Met Ser Lys Tyr Ser Gly Asn	
	165 170 175	
30	aag gca ggt gcc aag tac ggt acc ggc tac tgt gat tct cag tgc ccc	749
	Lys Ala Gly Ala Lys Tyr Gly Thr Gly Tyr Cys Asp Ser Gln Cys Pro	
	180 185 190 195	
35	cgc gac ctc aag ttc atc aac ggc gag gcc aac gta gag aac tgg cag	797
	Arg Asp Leu Lys Phe Ile Asn Gly Glu Ala Asn Val Glu Asn Trp Gln	
	200 205 210	
40	agc tcg acc aac gat gcc aac gcc ggc acg ggc aag tac ggc agc tgc	845
	Ser Ser Thr Asn Asp Ala Asn Ala Gly Thr Gly Lys Tyr Gly Ser Cys	
	215 220 225	
45	tgc tcc gag atg gac gtc tgg gag gcc aac aac atg gcc gcc gcc ttc	893
	Cys Ser Glu Met Asp Val Trp Glu Ala Asn Asn Met Ala Ala Ala Phe	
	230 235 240	
50	act ccc cac cct tgc acc gtg atc ggc cag tcg cgc tgc gag ggc gac	941
	Thr Pro His Pro Cys Thr Val Ile Gly Gln Ser Arg Cys Glu Gly Asp	
	245 250 255	
55	tcg tgc ggc ggt acc tac agc acc gac cgc tat gcc ggc atc tgc gac	989
	Ser Cys Gly Gly Thr Tyr Ser Thr Asp Arg Tyr Ala Gly Ile Cys Asp	
	260 265 270 275	
60	ccc gac gga tgc gac ttc aac tcg tac cgc cag ggc aac aag acc ttc	1037
	Pro Asp Gly Cys Asp Phe Asn Ser Tyr Arg Gln Gly Asn Lys Thr Phe	
	280 285 290	
65	tac ggc aag ggc atg acg gtc gac acg acc aag aag atc acg gtc gtc	1085
	Tyr Gly Lys Gly Met Thr Val Asp Thr Thr Lys Lys Ile Thr Val Val	
	295 300 305	
70	acc cag ttc ctc aag aac tcg gcc ggc gag ctc tcc gag atc aag cgg	1133
	Thr Gln Phe Leu Lys Asn Ser Ala Gly Glu Leu Ser Glu Ile Lys Arg	
	310 315 320	
75	ttc tac gtc cag aac ggc aag gtc atc ccc aac tcc gag tcc acc atc	1181

EP 2 183 363 B1

	Phe	Tyr	Val	Gln	Asn	Gly	Lys	Val	Ile	Pro	Asn	Ser	Glu	Ser	Thr	Ile	
	325						330					335					
5	ccg	ggc	gtc	gag	ggc	aac	tcc	atc	acc	cag	gac	tgg	tgc	gac	cgc	cag	1229
	Pro	Gly	Val	Glu	Gly	Asn	Ser	Ile	Thr	Gln	Asp	Trp	Cys	Asp	Arg	Gln	
	340					345					350					355	
10	aag	gcc	gcc	ttc	ggc	gac	gtg	acc	gac	ttc	cag	gac	aag	ggc	ggc	atg	1277
	Lys	Ala	Ala	Phe	Gly	Asp	Val	Thr	Asp	Phe	Gln	Asp	Lys	Gly	Gly	Met	
					360					365						370	
15	gtc	cag	atg	ggc	aag	gcc	ctc	gcg	ggg	ccc	atg	gtc	ctc	gtc	atg	tcc	1325
	Val	Gln	Met	Gly	Lys	Ala	Leu	Ala	Gly	Pro	Met	Val	Leu	Val	Met	Ser	
				375					380					385			
	atc	tgg	gac	gac	cac	gcc	gtc	aac	atg	ctc	tgg	ctc	gac	tcc	acc	tgg	1373
	Ile	Trp	Asp	Asp	His	Ala	Val	Asn	Met	Leu	Trp	Leu	Asp	Ser	Thr	Trp	
			390					395					400				
20	ccc	atc	gac	ggc	gcc	ggc	aag	ccg	ggc	gcc	gag	cgc	ggt	gcc	tgc	ccc	1421
	Pro	Ile	Asp	Gly	Ala	Gly	Lys	Pro	Gly	Ala	Glu	Arg	Gly	Ala	Cys	Pro	
		405					410					415					
25	acc	acc	tcg	ggc	gtc	ccc	gct	gag	gtc	gag	gcc	gag	gcc	ccc	aac	tcc	1469
	Thr	Thr	Ser	Gly	Val	Pro	Ala	Glu	Val	Glu	Ala	Glu	Ala	Pro	Asn	Ser	
	420					425					430					435	
30	aac	gtc	atc	ttc	tcc	aac	atc	cgc	ttc	ggc	ccc	atc	ggc	tcc	acc	gtc	1517
	Asn	Val	Ile	Phe	Ser	Asn	Ile	Arg	Phe	Gly	Pro	Ile	Gly	Ser	Thr	Val	
					440					445					450		
	tcc	ggc	ctg	ccc	gac	ggc	ggc	agc	ggc	aac	ccc	aac	ccg	ccc	gtc	agc	1565
	Ser	Gly	Leu	Pro	Asp	Gly	Gly	Ser	Gly	Asn	Pro	Asn	Pro	Pro	Val	Ser	
				455					460					465			
35	tcg	tcc	acc	ccg	gtc	ccc	tcc	tcg	tcc	acc	aca	tcc	tcc	ggt	tcc	tcc	1613
	Ser	Ser	Thr	Pro	Val	Pro	Ser	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ser	Ser	
			470					475						480			
40	ggc	ccg	act	ggc	ggc	acg	ggt	gtc	gct	aag	cac	tat	gag	caa	tgc	gga	1661
	Gly	Pro	Thr	Gly	Gly	Thr	Gly	Val	Ala	Lys	His	Tyr	Glu	Gln	Cys	Gly	
		485					490					495					
45	gga	atc	ggg	ttc	act	ggc	cct	acc	cag	tgc	gag	agc	ccc	tac	act	tgc	1709
	Gly	Ile	Gly	Phe	Thr	Gly	Pro	Thr	Gln	Cys	Glu	Ser	Pro	Tyr	Thr	Cys	
	500					505					510					515	
50	acc	aaa	ctg	aat	gac	tgg	tac	tcg	cag	tgc	ctg	taa	acga	acc	tctctga	agg	1762
	Thr	Lys	Leu	Asn	Asp	Trp	Tyr	Ser	Gln	Cys	Leu						
					520					525							
55	aggttctgag	acacgcgcga	ttcttctgta	tatagtttta	tttttca	ctc	tggagtgc	tt									1822
	cgctccacca	gtac															1836
	<210> 2																
	<211> 526																
	<212> PRT																
	<213> Chrysosporium lucknowense																

EP 2 183 363 B1

<400> 2

5	Met	Tyr	Ala	Lys	Phe	Ala	Thr	Leu	Ala	Ala	Leu	Val	Ala	Gly	Ala	Ala	1	5	10	15
10	Ala	Gln	Asn	Ala	Cys	Thr	Leu	Thr	Ala	Glu	Asn	His	Pro	Ser	Leu	Thr	20	25	30	
15	Trp	Ser	Lys	Cys	Thr	Ser	Gly	Gly	Ser	Cys	Thr	Ser	Val	Gln	Gly	Ser	35	40	45	
20	Ile	Thr	Ile	Asp	Ala	Asn	Trp	Arg	Trp	Thr	His	Arg	Thr	Asp	Ser	Ala	50	55	60	
25	Thr	Asn	Cys	Tyr	Glu	Gly	Asn	Lys	Trp	Asp	Thr	Ser	Tyr	Cys	Ser	Asp	65	70	75	80
30	Gly	Pro	Ser	Cys	Ala	Ser	Lys	Cys	Cys	Ile	Asp	Gly	Ala	Asp	Tyr	Ser	85	90	95	
35	Ser	Thr	Tyr	Gly	Ile	Thr	Thr	Ser	Gly	Asn	Ser	Leu	Asn	Leu	Lys	Phe	100	105	110	
40	Val	Thr	Lys	Gly	Gln	Tyr	Ser	Thr	Asn	Ile	Gly	Ser	Arg	Thr	Tyr	Leu	115	120	125	
45	Met	Glu	Ser	Asp	Thr	Lys	Tyr	Gln	Met	Phe	Gln	Leu	Leu	Gly	Asn	Glu	130	135	140	
50	Phe	Thr	Phe	Asp	Val	Asp	Val	Ser	Asn	Leu	Gly	Cys	Gly	Leu	Asn	Gly	145	150	155	160
55	Ala	Leu	Tyr	Phe	Val	Ser	Met	Asp	Ala	Asp	Gly	Gly	Met	Ser	Lys	Tyr	165	170	175	
60	Ser	Gly	Asn	Lys	Ala	Gly	Ala	Lys	Tyr	Gly	Thr	Gly	Tyr	Cys	Asp	Ser	180	185	190	
65	Gln	Cys	Pro	Arg	Asp	Leu	Lys	Phe	Ile	Asn	Gly	Glu	Ala	Asn	Val	Glu	195	200	205	
70	Asn	Trp	Gln	Ser	Ser	Thr	Asn	Asp	Ala	Asn	Ala	Gly	Thr	Gly	Lys	Tyr	210	215	220	
75	Gly	Ser	Cys	Cys	Ser	Glu	Met	Asp	Val	Trp	Glu	Ala	Asn	Asn	Met	Ala	225	230	235	240

EP 2 183 363 B1

	Ala	Ala	Phe	Thr	Pro	His	Pro	Cys	Thr	Val	Ile	Gly	Gln	Ser	Arg	Cys	
					245					250					255		
5	Glu	Gly	Asp	Ser	Cys	Gly	Gly	Thr	Tyr	Ser	Thr	Asp	Arg	Tyr	Ala	Gly	
				260					265					270			
10	Ile	Cys	Asp	Pro	Asp	Gly	Cys	Asp	Phe	Asn	Ser	Tyr	Arg	Gln	Gly	Asn	
			275					280					285				
15	Lys	Thr	Phe	Tyr	Gly	Lys	Gly	Met	Thr	Val	Asp	Thr	Thr	Lys	Lys	Ile	
		290					295					300					
20	Thr	Val	Val	Thr	Gln	Phe	Leu	Lys	Asn	Ser	Ala	Gly	Glu	Leu	Ser	Glu	
	305					310					315					320	
25	Ile	Lys	Arg	Phe	Tyr	Val	Gln	Asn	Gly	Lys	Val	Ile	Pro	Asn	Ser	Glu	
					325					330					335		
30	Ser	Thr	Ile	Pro	Gly	Val	Glu	Gly	Asn	Ser	Ile	Thr	Gln	Asp	Trp	Cys	
				340					345					350			
35	Asp	Arg	Gln	Lys	Ala	Ala	Phe	Gly	Asp	Val	Thr	Asp	Phe	Gln	Asp	Lys	
			355					360					365				
40	Gly	Gly	Met	Val	Gln	Met	Gly	Lys	Ala	Leu	Ala	Gly	Pro	Met	Val	Leu	
		370					375					380					
45	Val	Met	Ser	Ile	Trp	Asp	Asp	His	Ala	Val	Asn	Met	Leu	Trp	Leu	Asp	
	385					390					395					400	
50	Ser	Thr	Trp	Pro	Ile	Asp	Gly	Ala	Gly	Lys	Pro	Gly	Ala	Glu	Arg	Gly	
					405					410					415		
55	Ala	Cys	Pro	Thr	Thr	Ser	Gly	Val	Pro	Ala	Glu	Val	Glu	Ala	Glu	Ala	
				420					425					430			
60	Pro	Asn	Ser	Asn	Val	Ile	Phe	Ser	Asn	Ile	Arg	Phe	Gly	Pro	Ile	Gly	
			435					440					445				
65	Ser	Thr	Val	Ser	Gly	Leu	Pro	Asp	Gly	Gly	Ser	Gly	Asn	Pro	Asn	Pro	
		450					455					460					
70	Pro	Val	Ser	Ser	Ser	Thr	Pro	Val	Pro	Ser	Ser	Ser	Thr	Thr	Ser	Ser	
	465					470					475					480	

EP 2 183 363 B1

Gly Ser Ser Gly Pro Thr Gly Gly Thr Gly Val Ala Lys His Tyr Glu
485 490 495

5 Gln Cys Gly Gly Ile Gly Phe Thr Gly Pro Thr Gln Cys Glu Ser Pro
500 505 510

10 Tyr Thr Cys Thr Lys Leu Asn Asp Trp Tyr Ser Gln Cys Leu
515 520 525

<210> 3

<211> 1578

<212> DNA

15 <213> Chrysosporium lucknowense

<400> 3

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	atgtacgcca agttcgcgac cctcgccgcc cttgtggtg gcgccgctgc tcagaacgcc	60
	tgcactctga ccgctgagaa ccacccctcg ctgacgtggt ccaagtgcac gtctggcggc	120
5	agctgcacca gcgtccaggg ttccatcacc atcgacgcca actggcggtg gactcaccgg	180
	accgatagcg ccaccaactg ctacgagggc aacaagtggg atacttcgta ctgcagcgat	240
10	ggtccttctt gcgcctccaa gtgctgcatc gacggcgctg actactcgag cacctatggc	300
	atcaccacga gcggtaaact cctgaacctc aagttcgtca ccaagggcca gtactcgacc	360
	aacatcgggt cgcgtaacta cctgatggag agcgacacca agtaccagat gttccagctc	420
15	ctcggcaacg agttcacctt cgatgtcgac gtctccaacc tcggctgcgg cctcaatggc	480
	gccctctact tcgtgtccat ggatgccgat ggtggcatgt ccaagtactc gggcaacaag	540
	gcaggtgcca agtacggtag cggctactgt gattctcagt gccccgcga cctcaagttc	600
20	atcaacggcg aggccaacgt agagaactgg cagagctcga ccaacgatgc caacgccggc	660
	acgggcaagt acggcagctg ctgctccgag atggacgtct gggaggccaa caacatggcc	720
25	gccgccttca ctccccaccc ttgcaccgtg atcggccagt cgcgctgcga gggcgactcg	780
	tgcggcggtg cctacagcac cgaccgctat gccggcatct gcgaccccg cggatgcgac	840
	ttcaactcgt accgccaggg caacaagacc ttctacggca agggcatgac ggtcgacacg	900
30	accaagaaga tcacggtcgt caccagttc ctcaagaact cggccggcga gctctccgag	960
	atcaagcggg tctacgtcca gaacggcaag gtcacccca actccgagtc caccatcccg	1020
35	ggcgtcgagg gcaactccat caccaggac tgggtgcgacc gccagaaggc cgccttcggc	1080
	gacgtgaccg acttccagga caaggcggc atggtccaga tgggcaaggc cctcgcgggg	1140
	cccatggtcc tcgtcatgtc catctgggac gaccacgccg tcaacatgct ctggctcgac	1200
40	tccacctggc ccatcgacgg cgccggcaag ccgggcgccg agcgcggtgc ctgccccacc	1260
	acctcggggc tccccgctga ggtcgaggcc gaggcccca actccaacgt catcttctcc	1320
45	aacatccgct tcggcccat cggtccacc gtctccggcc tgcccgacgg cggcagcggc	1380
	aaccccaacc cgccgctcag ctcggtccacc ccggtccct cctcggtccac cacatcctcc	1440
	ggttcctccg gcccgactgg cggcacgggt gtcgctaagc actatgagca atgcggagga	1500
50	atcgggttca ctggccctac ccagtgcgag agcccctaca cttgcaccaa actgaatgac	1560
	tggtactcgc agtgccctg	1578
55	<210> 4 <211> 2029 <212> DNA <213> Chrysosporium lucknowense	

EP 2 183 363 B1

<220>
 <221> CDS
 <222> (108)..(674)

5

<220>
 <221> CDS
 <222> (762)..(898)

10

<220>
 <221> CDS
 <222> (974)..(1186)

15

<220>
 <221> CDS
 <222> (1284)..(1530)

20

<220>
 <221> CDS
 <222> (1638)..(1768)

25

<400> 4

atataaaaaag gcaaattgagg cggcgccttg gacaggtcca ttctcccacc gctcaaccag 60
 cctccaattc ctcagaagtc tgttgctctc tcgcagtcgc agtcaag atg aag cag 116
 Met Lys Gln
 1

30

tac ctc cag tac ctc gcg gcg acc ctg ccc ctg gtg ggc ctg gcc acg 164
 Tyr Leu Gln Tyr Leu Ala Ala Thr Leu Pro Leu Val Gly Leu Ala Thr
 5 10 15

35

gcc cag cag gcg ggt aac ctg cag acc gag act cac ccc aag ctc act 212
 Ala Gln Gln Ala Gly Asn Leu Gln Thr Glu Thr His Pro Lys Leu Thr
 20 25 30 35

40

tgg tcg aag tgc acg gcc ccg gga tcc tgc caa cag gtc aac ggc gag 260
 Trp Ser Lys Cys Thr Ala Pro Gly Ser Cys Gln Gln Val Asn Gly Glu
 40 45 50

45

50

55

EP 2 183 363 B1

	gtc gtc atc gac tcc aac tgg cgc tgg gtg cac gac gag aac gcg cag	308
	Val Val Ile Asp Ser Asn Trp Arg Trp Val His Asp Glu Asn Ala Gln	
	55 60 65	
5	aac tgc tac gac ggc aac cag tgg acc aac gct tgc agc tct gcc acc	356
	Asn Cys Tyr Asp Gly Asn Gln Trp Thr Asn Ala Cys Ser Ser Ala Thr	
	70 75 80	
10	gac tgc gcc gag aat tgc gcg ctc gag ggt gcc gac tac cag ggc acc	404
	Asp Cys Ala Glu Asn Cys Ala Leu Glu Gly Ala Asp Tyr Gln Gly Thr	
	85 90 95	
15	tat ggc gcc tcg acc agc ggc aat gcc ctg acg ctc acc ttc gtc act	452
	Tyr Gly Ala Ser Thr Ser Gly Asn Ala Leu Thr Leu Thr Phe Val Thr	
	100 105 110 115	
20	aag cac gag tac ggc acc aac att ggt tgc cgc ctc tac ctc atg aac	500
	Lys His Glu Tyr Gly Thr Asn Ile Gly Ser Arg Leu Tyr Leu Met Asn	
	120 125 130	
25	ggc gcg aac aag tac cag atg ttc acc ctc aag ggc aac gag ctg gcc	548
	Gly Ala Asn Lys Tyr Gln Met Phe Thr Leu Lys Gly Asn Glu Leu Ala	
	135 140 145	
30	ttc gac gtc gac ctc tcg gcc gtc gag tgc ggc ctc aac agc gcc ctc	596
	Phe Asp Val Asp Leu Ser Ala Val Glu Cys Gly Leu Asn Ser Ala Leu	
	150 155 160	
35	tac ttc gtg gcc atg gag gag gat ggc ggt gtg tgc agc tac ccg acc	644
	Tyr Phe Val Ala Met Glu Glu Asp Gly Gly Val Ser Ser Tyr Pro Thr	
	165 170 175	
40	aac acg gcc ggt gct aag ttc ggc act ggg gtaagttcaa cgacccgaga	694
	Asn Thr Ala Gly Ala Lys Phe Gly Thr Gly	
	180 185	
45	cgggtgccct tattatctgc tgcgaaaacg gacgggtcccc ttttgctaac taccctctc	754
	caaacag tac tgc gac gcc caa tgc gca cgc gac ctc aag ttc gtc ggc	803
	Tyr Cys Asp Ala Gln Cys Ala Arg Asp Leu Lys Phe Val Gly	
	190 195 200	
50	ggc aag ggc aac atc gag ggc tgg aag ccg tcc acc aac gat gcc aat	851
	Gly Lys Gly Asn Ile Glu Gly Trp Lys Pro Ser Thr Asn Asp Ala Asn	
	205 210 215	
55	gcc ggt gtc ggt cct tat ggc ggg tgc tgc gct gag atc gac gtc tg	898
	Ala Gly Val Gly Pro Tyr Gly Gly Cys Cys Ala Glu Ile Asp Val Trp	
	220 225 230	
60	gtaagttttg ttgcctgggc agcaatggta tattagctcg agtggttccc gtcggtgctg	958
	accctctctt accag g gag tgc aac aag tat gct ttc gct ttc acc ccg	1007
	Glu Ser Asn Lys Tyr Ala Phe Ala Phe Thr Pro	
	240 245	
65	cac ggt tgc gag aac cct aaa tac cac gtc tgc gag acc acc aac tgc	1055
	His Gly Cys Glu Asn Pro Lys Tyr His Val Cys Glu Thr Thr Asn Cys	
	250 255 260	
70	ggt ggc acc tac tcc gag gac cgc ttc gct ggt gac tgc gat gcc aac	1103

EP 2 183 363 B1

	Gly	Gly	Thr	Tyr	Ser	Glu	Asp	Arg	Phe	Ala	Gly	Asp	Cys	Asp	Ala	Asn	
			265					270					275				
5	ggc	tgc	gac	tac	aac	ccc	tac	cgc	atg	ggc	aac	cag	gac	ttc	tac	ggt	1151
	Gly	Cys	Asp	Tyr	Asn	Pro	Tyr	Arg	Met	Gly	Asn	Gln	Asp	Phe	Tyr	Gly	
		280					285					290					
	ccc	ggc	ttg	acg	gtc	gat	acc	agc	aag	aag	ttc	ac	gtgagtacac				1196
10	Pro	Gly	Leu	Thr	Val	Asp	Thr	Ser	Lys	Lys	Phe	Thr					
		295				300					305						
	cgtgcttgaa	gccccctccc	cccccccccc	caaaaaaaaa	aagaaaaaag	aagtcaaatg											1256
	attgatgcta	accaaataca	ataacag	c	gtc	gtc	agc	cag	ttc	gag	gag	aac					1308
15								Val	Val	Ser	Gln	Phe	Glu	Glu	Asn		
											310						
	aag	ctc	acc	cag	ttc	ttc	gtc	cag	gac	ggc	aag	aag	att	gag	atc	ccc	1356
	Lys	Leu	Thr	Gln	Phe	Phe	Val	Gln	Asp	Gly	Lys	Lys	Ile	Glu	Ile	Pro	
20		315				320					325					330	
	ggc	ccc	aag	gtc	gag	ggc	atc	gat	gcg	gac	agc	gcc	gct	atc	acc	cct	1404
	Gly	Pro	Lys	Val	Glu	Gly	Ile	Asp	Ala	Asp	Ser	Ala	Ala	Ile	Thr	Pro	
					335					340				345			
	gag	ctg	tgc	agt	gcc	ctg	ttc	aag	gcc	ttc	gat	gac	cgt	gac	cgc	ttc	1452
25	Glu	Leu	Cys	Ser	Ala	Leu	Phe	Lys	Ala	Phe	Asp	Asp	Arg	Asp	Arg	Phe	
				350					355					360			
	tcg	gag	gtt	ggc	ggc	ttc	gat	gcc	atc	aac	acg	gcc	ctc	agc	act	ccc	1500
30	Ser	Glu	Val	Gly	Gly	Phe	Asp	Ala	Ile	Asn	Thr	Ala	Leu	Ser	Thr	Pro	
			365				370						375				
	atg	gtc	ctc	gtc	atg	tcc	atc	tgg	gat	gat	gtacgttacc	taaccccccc					1550
	Met	Val	Leu	Val	Met	Ser	Ile	Trp	Asp	Asp							
		380				385											
35	ccccctttttt	tttcccgcctt	ctctccccga	aactgccact	acttatatac	gtccccgcgc											1610
	catgatgctt	acctttttctc	cttcacag	cac	tac	gcc	aat	atg	ctc	tgg	ctc	gac					1664
								His	Tyr	Ala	Asn	Met	Leu	Trp	Leu	Asp	
40								390				395					
	tcg	agc	tac	ccc	cct	gag	aag	gct	ggc	cag	cct	ggc	ggt	gac	cgt	ggc	1712
	Ser	Ser	Tyr	Pro	Pro	Glu	Lys	Ala	Gly	Gln	Pro	Gly	Gly	Asp	Arg	Gly	
			400				405					410					
	ccg	tgt	cct	cag	gac	tct	ggc	gtc	ccg	gcc	gac	gtt	gag	gct	cag	tac	1760
45	Pro	Cys	Pro	Gln	Asp	Ser	Gly	Val	Pro	Ala	Asp	Val	Glu	Ala	Gln	Tyr	
		415				420						425					
	cct	aat	gc	gtgagtcgaa	accgtaaaat	gtcgggcaaa	aaaaagatcg										1808
50	Pro	Asn	Ala														
		430															
	ctcaagctaa	cgaataata	tgattag	c	aag	gtc	atc	tgg	tcc	aac	atc	cgc					1860
								Lys	Val	Ile	Trp	Ser	Asn	Ile	Arg		
									435					440			
55	ttc	ggc	ccc	atc	ggc	tgc	act	gtc	aac	gtc	taaactgcaa	cctgaccggg					1910
	Phe	Gly	Pro	Ile	Gly	Ser	Thr	Val	Asn	Val							

EP 2 183 363 B1

445

450

ccctttctct ccacccccac ccctctcaag ttctctctgg tggagccctc gtgtccttct 1970

5 tttcctaggt tcgcgaacct ttgagcttgt gtatcgtagg gtcattgtgt acatacaca 2029

<210> 5

<211> 450

<212> PRT

10 <213> Chrysosporium lucknowense

<400> 5

15 Met Lys Gln Tyr Leu Gln Tyr Leu Ala Ala Thr Leu Pro Leu Val Gly
1 5 10 15

20 Leu Ala Thr Ala Gln Gln Ala Gly Asn Leu Gln Thr Glu Thr His Pro
20 25 30

25 Lys Leu Thr Trp Ser Lys Cys Thr Ala Pro Gly Ser Cys Gln Gln Val
35 40 45

Asn Gly Glu Val Val Ile Asp Ser Asn Trp Arg Trp Val His Asp Glu
50 55 60

30 Asn Ala Gln Asn Cys Tyr Asp Gly Asn Gln Trp Thr Asn Ala Cys Ser
65 70 75 80

35 Ser Ala Thr Asp Cys Ala Glu Asn Cys Ala Leu Glu Gly Ala Asp Tyr
85 90 95

40 Gln Gly Thr Tyr Gly Ala Ser Thr Ser Gly Asn Ala Leu Thr Leu Thr
100 105 110

Phe Val Thr Lys His Glu Tyr Gly Thr Asn Ile Gly Ser Arg Leu Tyr
115 120 125

45 Leu Met Asn Gly Ala Asn Lys Tyr Gln Met Phe Thr Leu Lys Gly Asn
130 135 140

50 Glu Leu Ala Phe Asp Val Asp Leu Ser Ala Val Glu Cys Gly Leu Asn
145 150 155 160

Ser Ala Leu Tyr Phe Val Ala Met Glu Glu Asp Gly Gly Val Ser Ser
165 170 175

55 Tyr Pro Thr Asn Thr Ala Gly Ala Lys Phe Gly Thr Gly Tyr Cys Asp
180 185 190

EP 2 183 363 B1

	Ala	Gln	Cys	Ala	Arg	Asp	Leu	Lys	Phe	Val	Gly	Gly	Lys	Gly	Asn	Ile
			195					200					205			
5	Glu	Gly	Trp	Lys	Pro	Ser	Thr	Asn	Asp	Ala	Asn	Ala	Gly	Val	Gly	Pro
		210					215					220				
10	Tyr	Gly	Gly	Cys	Cys	Ala	Glu	Ile	Asp	Val	Trp	Glu	Ser	Asn	Lys	Tyr
	225					230					235					240
15	Ala	Phe	Ala	Phe	Thr	Pro	His	Gly	Cys	Glu	Asn	Pro	Lys	Tyr	His	Val
				245						250					255	
20	Cys	Glu	Thr	Thr	Asn	Cys	Gly	Gly	Thr	Tyr	Ser	Glu	Asp	Arg	Phe	Ala
				260					265					270		
25	Gly	Asp	Cys	Asp	Ala	Asn	Gly	Cys	Asp	Tyr	Asn	Pro	Tyr	Arg	Met	Gly
			275					280					285			
30	Asn	Gln	Asp	Phe	Tyr	Gly	Pro	Gly	Leu	Thr	Val	Asp	Thr	Ser	Lys	Lys
		290					295					300				
35	Phe	Thr	Val	Val	Ser	Gln	Phe	Glu	Glu	Asn	Lys	Leu	Thr	Gln	Phe	Phe
	305					310					315					320
40	Val	Gln	Asp	Gly	Lys	Lys	Ile	Glu	Ile	Pro	Gly	Pro	Lys	Val	Glu	Gly
				325						330					335	
45	Ile	Asp	Ala	Asp	Ser	Ala	Ala	Ile	Thr	Pro	Glu	Leu	Cys	Ser	Ala	Leu
			340						345					350		
50	Phe	Lys	Ala	Phe	Asp	Asp	Arg	Asp	Arg	Phe	Ser	Glu	Val	Gly	Gly	Phe
			355					360					365			
55	Asp	Ala	Ile	Asn	Thr	Ala	Leu	Ser	Thr	Pro	Met	Val	Leu	Val	Met	Ser
		370					375					380				
60	Ile	Trp	Asp	Asp	His	Tyr	Ala	Asn	Met	Leu	Trp	Leu	Asp	Ser	Ser	Tyr
	385					390					395					400
65	Pro	Pro	Glu	Lys	Ala	Gly	Gln	Pro	Gly	Gly	Asp	Arg	Gly	Pro	Cys	Pro
					405					410					415	
70	Gln	Asp	Ser	Gly	Val	Pro	Ala	Asp	Val	Glu	Ala	Gln	Tyr	Pro	Asn	Ala
				420					425					430		

EP 2 183 363 B1

Lys Val Ile Trp Ser Asn Ile Arg Phe Gly Pro Ile Gly Ser Thr Val
435 440 445

Asn Val

5

450

<210> 6

<211> 1350

10

<212> DNA

<213> Chrysosporium lucknowense

<400> 6

15

atgaagcagt acctccagta cctcgcggcg accctgcccc tgggtgggcct ggccacggcc 60

cagcaggcgg gtaacctgca gaccgagact caccccaagc tcacttggtc gaagtgcacg 120

20

gccccgggat cctgccaaca ggtcaacggc gaggtcgtca tcgactccaa ctggcgctgg 180

gtgcacgacg agaacgcgca gaactgctac gacggcaacc agtggacca cgcttgcagc 240

tctgccaccg actgcgccga gaattgcgcg ctcgaggggtg ccgactacca gggcacctat 300

25

ggcgcctcga ccagcggcaa tgccctgacg ctcaccttcg tcactaagca cgagtacggc 360

accaacattg gttcgcgcct ctacctcatg aacggcgcga acaagtacca gatgttcacc 420

ctcaagggca acgagctggc cttcgacgtc gacctctcgg ccgtcgagtg cggcctcaac 480

30

agcgccctct acttcgtggc catggaggag gatggcgggtg tgtcgagcta cccgaccaac 540

acggccgggtg ctaagttcgg cactgggtac tgcgacgccc aatgcgcacg cgacctcaag 600

35

ttcgtcggcg gcaagggcaa catcgagggc tggaagccgt ccaccaacga tgccaatgcc 660

gggtgtcggtc cttatggcgg gtgctgcgct gagatcgacg tctgggagtc gaacaagtat 720

gctttcgctt tcaccccgca cggttgcgag aaccctaaat accacgtctg cgagaccacc 780

40

aactgcgggtg gcacctactc cgaggaccgc ttcgctgggtg actgcgatgc caacggctgc 840

gactacaacc cctaccgcat gggcaaccag gacttctacg gtcccggctt gacggtcgat 900

45

accagcaaga agttcacctg cgtcagccag ttcgaggaga acaagctcac ccagttcttc 960

gtccaggacg gcaagaagat tgagatcccc ggccccaagg tcgagggcat cgatgcggac 1020

agcgccgcta tcaccctga gctgtgcagt gccctgttca aggccttcga tgaccgtgac 1080

50

cgcttctcgg aggttggcgg cttcgatgcc atcaacacgg ccctcagcac tcccatggtc 1140

ctcgtcatgt ccatctggga tgatcactac gccaatatgc tctggctcga ctcgagctac 1200

ccccctgaga aggctggcca gcctggcggc gaccgtggcc cgtgtcctca ggactctggc 1260

55

gtcccggccg acgttgaggc tcagtaccct aatgccaagg tcatctggtc caacatccgc 1320

ttcggcccca tcggctcgac tgtcaacgtc 1350

<210> 7
 <211> 1592
 <212> DNA
 <213> Chrysosporium lucknowense

5

<220>
 <221> CDS
 <222> (78) .. (519)

10

<220>
 <221> CDS
 <222> (630)..(954)

15

<220>
 <221> CDS
 <222> (1048)..(1465)

<400> 7

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	tgcttgaggaa gaaaggatct ctcgaccatg caccacagcc tagctctaac ccagcttgtc	60
5	gtgtgttggtt gcccagc atg aag ttc gtg cag tcc gcc acc ctg gcg ttc Met Lys Phe Val Gln Ser Ala Thr Leu Ala Phe 1 5 10	110
10	gcc gcc acg gcc ctc gct gcg ccc tcg cgc acg act ccc cag aag ccc Ala Ala Thr Ala Leu Ala Ala Pro Ser Arg Thr Thr Pro Gln Lys Pro 15 20 25	158
	cgc cag gcc tcg gcg ggc tgc gcg tcg gcc gtg acg ctc gat gcc agc Arg Gln Ala Ser Ala Gly Cys Ala Ser Ala Val Thr Leu Asp Ala Ser 30 35 40	206
15	acc aac gtg ttc cag cag tac acg ctg cac ccc aac aac ttc tac cgt Thr Asn Val Phe Gln Gln Tyr Thr Leu His Pro Asn Asn Phe Tyr Arg 45 50 55	254
20	gcc gag gtc gag gct gcc gcc gag gcc atc tcc gac tcg gcg ctg gcc Ala Glu Val Glu Ala Ala Ala Glu Ala Ile Ser Asp Ser Ala Leu Ala 60 65 70 75	302
25	gag aag gcc cgc aag gtc gcc gac gtc ggt acc ttc ctg tgg ctc gac Glu Lys Ala Arg Lys Val Ala Asp Val Gly Thr Phe Leu Trp Leu Asp 80 85 90	350
	acc atc gag aac att ggc cgg ctg gag ccc gcg ctc gag gac gtg ccc Thr Ile Glu Asn Ile Gly Arg Leu Glu Pro Ala Leu Glu Asp Val Pro 95 100 105	398
30	tgc gag aac atc gtg ggt ctc gtc atc tac gac ctc ccg ggc cgt gac Cys Glu Asn Ile Val Gly Leu Val Ile Tyr Asp Leu Pro Gly Arg Asp 110 115 120	446
35	tgc gcg gcc aag gcc tcc aac ggc gag ctc aag gtc ggc gag ctc gac Cys Ala Ala Lys Ala Ser Asn Gly Glu Leu Lys Val Gly Glu Leu Asp 125 130 135	494
40	agg tac aag acc gag tac atc gac a gtgagttaac cctttgtggc Arg Tyr Lys Thr Glu Tyr Ile Asp 140 145	539
45		
50		
55		

EP 2 183 363 B1

	cccttctttt cccccgagag agcgtctggt tgagtggggt tgtgagagag aaaatggggc	599
	gagcttaaag actgacgtgt tggctcgcag ag atc gcc gag atc ctc aag gcc	652
5	Lys Ile Ala Glu Ile Leu Lys Ala 150 155	
	cac tcc aac acg gcc ttc gcc ctc gtc atc gag ccc gac tcg ctc ccc	700
	His Ser Asn Thr Ala Phe Ala Leu Val Ile Glu Pro Asp Ser Leu Pro 160 165 170	
10	aac ctg gtc acc aat agc gac ctg cag acg tgc cag cag agc gct tcc	748
	Asn Leu Val Thr Asn Ser Asp Leu Gln Thr Cys Gln Gln Ser Ala Ser 175 180 185	
15	ggc tac cgc gag ggt gtc gcc tat gcc ctc aag cag ctc aac ctc ccc	796
	Gly Tyr Arg Glu Gly Val Ala Tyr Ala Leu Lys Gln Leu Asn Leu Pro 190 195 200	
20	aac gtg gtc atg tac atc gat gcc ggc cac ggt ggc tgg ctc ggc tgg	844
	Asn Val Val Met Tyr Ile Asp Ala Gly His Gly Gly Trp Leu Gly Trp 205 210 215	
	gac gcc aac ctc aag ccc ggc gcc cag gag ctc gcc agc gtc tac aag	892
	Asp Ala Asn Leu Lys Pro Gly Ala Gln Glu Leu Ala Ser Val Tyr Lys 220 225 230 235	
25	tct gct ggt tcg ccc tcg caa gtc cgc ggt atc tcc acc aac gtg gct	940
	Ser Ala Gly Ser Pro Ser Gln Val Arg Gly Ile Ser Thr Asn Val Ala 240 245 250	
30	ggt tgg aac gcc tg gtaagacact ctatgtcccc ctcgtcggtc aatggcgagc	994
	Gly Trp Asn Ala Trp 255	
35	ggaatggcgt gaaatgcatg gtgctgacct ttgatctttt cccctccta tag g gac	1051
	Asp	
	cag gag ccc ggt gag ttc tcg gac gcc tcg gat gcc cag tac aac aag	1099
	Gln Glu Pro Gly Glu Phe Ser Asp Ala Ser Asp Ala Gln Tyr Asn Lys 260 265 270	
40	tgc cag aac gag aag atc tac atc aac acc ttt ggc gct gag ctc aag	1147
	Cys Gln Asn Glu Lys Ile Tyr Ile Asn Thr Phe Gly Ala Glu Leu Lys 275 280 285	
45	tct gcc ggc atg ccc aac cac gcc atc atc gac act ggc cgc aac ggt	1195
	Ser Ala Gly Met Pro Asn His Ala Ile Ile Asp Thr Gly Arg Asn Gly 290 295 300 305	
50	gtc acc ggt ctc cgc gac gag tgg ggt gac tgg tgc aac gtc aac ggc	1243
	Val Thr Gly Leu Arg Asp Glu Trp Gly Asp Trp Cys Asn Val Asn Gly 310 315 320	
	gcc ggc ttc ggt gtg cgc cgc act gcc aac act ggc gac gag ctc gcc	1291
	Ala Gly Phe Gly Val Arg Pro Thr Ala Asn Thr Gly Asp Glu Leu Ala 325 330 335	
55	gac gcc ttc gtg tgg gtc aag ccc ggt ggc gag tcc gac ggc acc agc	1339
	Asp Ala Phe Val Trp Val Lys Pro Gly Gly Glu Ser Asp Gly Thr Ser 340 345 350	

EP 2 183 363 B1

gac tcg tcg gcg gcg cgc tac gac agc ttc tgc ggc aag ccc gac gcc 1387
 Asp Ser Ser Ala Ala Arg Tyr Asp Ser Phe Cys Gly Lys Pro Asp Ala
 355 360 365

5 ttc aag ccc agc ccc gag gcc ggt acc tgg aac cag gcc tac ttc gag 1435
 Phe Lys Pro Ser Pro Glu Ala Gly Thr Trp Asn Gln Ala Tyr Phe Glu
 370 375 380 385

10 atg ctc ctc aag aac gcc aac ccg tcc ttc taagctcctc gacggcttct 1485
 Met Leu Leu Lys Asn Ala Asn Pro Ser Phe
 390 395

tgctgtcagt cgctctgacg gtgggtgtgct ggtgggtgccc ctgctcctgc tgctgctgct 1545

15 ccgcgggggag gggaggcaac gaaaatgaag tcctgcttca aaacaaa 1592

<210> 8
 <211> 395
 <212> PRT
 20 <213> Chrysosporium lucknowense

<400> 8

25 Met Lys Phe Val Gln Ser Ala Thr Leu Ala Phe Ala Ala Thr Ala Leu
 1 5 10 15

Ala Ala Pro Ser Arg Thr Thr Pro Gln Lys Pro Arg Gln Ala Ser Ala
 20 25 30

30 Gly Cys Ala Ser Ala Val Thr Leu Asp Ala Ser Thr Asn Val Phe Gln
 35 40 45

35 Gln Tyr Thr Leu His Pro Asn Asn Phe Tyr Arg Ala Glu Val Glu Ala
 50 55 60

40 Ala Ala Glu Ala Ile Ser Asp Ser Ala Leu Ala Glu Lys Ala Arg Lys
 65 70 75 80

45 Val Ala Asp Val Gly Thr Phe Leu Trp Leu Asp Thr Ile Glu Asn Ile
 85 90 95

Gly Arg Leu Glu Pro Ala Leu Glu Asp Val Pro Cys Glu Asn Ile Val
 100 105 110

50 Gly Leu Val Ile Tyr Asp Leu Pro Gly Arg Asp Cys Ala Ala Lys Ala
 115 120 125

55 Ser Asn Gly Glu Leu Lys Val Gly Glu Leu Asp Arg Tyr Lys Thr Glu
 130 135 140

EP 2 183 363 B1

	Tyr	Ile	Asp	Lys	Ile	Ala	Glu	Ile	Leu	Lys	Ala	His	Ser	Asn	Thr	Ala	
	145					150					155					160	
5	Phe	Ala	Leu	Val	Ile	Glu	Pro	Asp	Ser	Leu	Pro	Asn	Leu	Val	Thr	Asn	
					165					170					175		
10	Ser	Asp	Leu	Gln	Thr	Cys	Gln	Gln	Ser	Ala	Ser	Gly	Tyr	Arg	Glu	Gly	
				180					185					190			
15	Val	Ala	Tyr	Ala	Leu	Lys	Gln	Leu	Asn	Leu	Pro	Asn	Val	Val	Met	Tyr	
			195					200					205				
20	Ile	Asp	Ala	Gly	His	Gly	Gly	Trp	Leu	Gly	Trp	Asp	Ala	Asn	Leu	Lys	
		210					215					220					
25	Pro	Gly	Ala	Gln	Glu	Leu	Ala	Ser	Val	Tyr	Lys	Ser	Ala	Gly	Ser	Pro	
	225					230					235					240	
30	Ser	Gln	Val	Arg	Gly	Ile	Ser	Thr	Asn	Val	Ala	Gly	Trp	Asn	Ala	Trp	
					245					250					255		
35	Asp	Gln	Glu	Pro	Gly	Glu	Phe	Ser	Asp	Ala	Ser	Asp	Ala	Gln	Tyr	Asn	
				260					265					270			
40	Lys	Cys	Gln	Asn	Glu	Lys	Ile	Tyr	Ile	Asn	Thr	Phe	Gly	Ala	Glu	Leu	
			275					280					285				
45	Lys	Ser	Ala	Gly	Met	Pro	Asn	His	Ala	Ile	Ile	Asp	Thr	Gly	Arg	Asn	
		290					295					300					
50	Gly	Val	Thr	Gly	Leu	Arg	Asp	Glu	Trp	Gly	Asp	Trp	Cys	Asn	Val	Asn	
	305					310					315					320	
55	Gly	Ala	Gly	Phe	Gly	Val	Arg	Pro	Thr	Ala	Asn	Thr	Gly	Asp	Glu	Leu	
					325					330					335		
60	Ala	Asp	Ala	Phe	Val	Trp	Val	Lys	Pro	Gly	Gly	Glu	Ser	Asp	Gly	Thr	
				340					345					350			
65	Ser	Asp	Ser	Ser	Ala	Ala	Arg	Tyr	Asp	Ser	Phe	Cys	Gly	Lys	Pro	Asp	
			355					360					365				
70	Ala	Phe	Lys	Pro	Ser	Pro	Glu	Ala	Gly	Thr	Trp	Asn	Gln	Ala	Tyr	Phe	
		370					375					380					
75	Glu	Met	Leu	Leu	Lys	Asn	Ala	Asn	Pro	Ser	Phe						

EP 2 183 363 B1

	385	390	395	
5	<210> 9 <211> 1185 <212> DNA <213> Chrysosporium lucknowense			
10	<400> 9			
	atgaagttcg tgcagtcgcg caccctggcg ttcgccgccg cgcccttcgc tgcgccctcg			60
	cgcacgactc cccagaagcc ccgccaggcc tcggcgggct gcgcgtcggc cgtgacgctc			120
15	gatgccagca ccaacgtgtt ccagcagtac acgtcgcacc ccaacaactt ctaccgtgcc			180
	gaggtcgagg ctgccgccga ggccatctcc gactcggcgc tggccgagaa ggcccgaag			240
	gtcgccgacg tcggtacctt cctgtggctc gacaccatcg agaacattgg ccggttgag			300
20	cccgcgctcg aggacgtgcc ctgcgagAAC atcgtgggtc tcgtcatcta cgacctcccg			360
	ggccgtgact gcgcggccaa ggcctccaac ggcgagctca aggtcggcga gctcgacagg			420
25	tacaagaccg agtacatcga caagatcgcc gagatcctca aggccactc caacacggcc			480
	ttcgccctcg tcatcgagcc cgactcgtc cccaacctgg tcaccaatag cgacctgcag			540
	acgtgccagc agagcgcttc cggctaccgc gaggggtgtc cctatgccct caagcagctc			600
30	aacctcccca acgtgggtcat gtacatcgat gccggccacg gtgggtgggt cggctgggac			660
	gccaacctca agcccggcgc ccaggagctc gccagcgtct acaagtctgc tggttcgccc			720
35	tcgcaagtcc gcggtatctc caccaacgtg gctggttga acgcctggga ccaggagccc			780
	ggtgagttct cggacgcctc ggatgccag tacaacaagt gccagaacga gaagatctac			840
	atcaacacct ttggcgctga gctcaagtct gccggcatgc ccaaccacgc catcatcgac			900
40	actggccgca acggtgtcac cggctctccg gacgagtggg gtgactggtg caacgtcaac			960
	ggcgccggct tcggtgtgcg cccgactgcc aacactggcg acgagctcgc cgacgccttc			1020
	gtgtgggtca agcccgggtg cgagtccgac ggcaccagcg actcgtcggc ggcgcgctac			1080
45	gacagcttct gcggcaagcc cgacgccttc aagcccagcc ccgaggccgg tacctggaac			1140
	caggcctact tcgagatgct cctcaagaac gccaacccgt ccttc			1185
50	<210> 10 <211> 2115 <212> DNA <213> Chrysosporium lucknowense			
55	<220> <221> CDS <222> (90)..(181)			

<220>
<221> CDS
<222> (278)..(786)

5 <220>
 <221> CDS
 <222> (871)..(1455)

10 <220>
 <221> CDS
 <222> (1629)..(1888)

<400> 10

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	tgtgtggttt ctctccctcg tgccaaacca ctccacctc ccgccccgag atagttgctt	60
5	gtttcgctcc gtgagaggga cacacacca atg gcc aag aag ctt ttc atc acc Met Ala Lys Lys Leu Phe Ile Thr 1 5	113
10	gcc gcg ctt gcg gct gcc gtg ttg gcg gcc ccc gtc att gag gag cgc Ala Ala Leu Ala Ala Ala Val Leu Ala Ala Pro Val Ile Glu Glu Arg 10 15 20	161
	cag aac tgc ggc gct gtg tg gtaagaaagc ccggtccgag tctcccatga Gln Asn Cys Gly Ala Val Trp 25 30	211
15	ttttctcgtc gagtaatggc ataagggcca ccccttcgac tgaccgtgag aatcgatcaa	271
	atccag g act caa tgc ggc ggt aac ggg tgg caa ggt ccc aca tgc tgc Thr Gln Cys Gly Gly Asn Gly Trp Gln Gly Pro Thr Cys Cys 35 40 45	320
20	gcc tcg ggc tcg acc tgc gtt gcg cag aac gag tgg tac tct cag tgc Ala Ser Gly Ser Thr Cys Val Ala Gln Asn Glu Trp Tyr Ser Gln Cys 50 55 60	368
25	ctg ccc aac agc cag gtg acg agt tcc acc act ccg tcg tcg act tcc Leu Pro Asn Ser Gln Val Thr Ser Ser Thr Thr Pro Ser Ser Thr Ser 65 70 75	416
30	acc tcg cag cgc agc acc agc acc tcc agc agc acc acc agg agc ggc Thr Ser Gln Arg Ser Thr Ser Thr Ser Ser Ser Thr Thr Arg Ser Gly 80 85 90	464
35	agc tcc tcc tcc tcc tcc acc acg ccc ccg ccc gtc tcc agc ccc gtg Ser Ser Ser Ser Ser Thr Thr Pro Pro Pro Val Ser Ser Pro Val 95 100 105	512
	acc agc att ccc ggc ggt gcg acc tcc acg gcg agc tac tct ggc aac Thr Ser Ile Pro Gly Gly Ala Thr Ser Thr Ala Ser Tyr Ser Gly Asn 110 115 120 125	560
40	ccc ttc tcg ggc gtc cgg ctc ttc gcc aac gac tac tac agg tcc gag Pro Phe Ser Gly Val Arg Leu Phe Ala Asn Asp Tyr Tyr Arg Ser Glu 130 135 140	608
45	gtc cac aat ctc gcc att cct agc atg act ggt act ctg gcg gcc aag Val His Asn Leu Ala Ile Pro Ser Met Thr Gly Thr Leu Ala Ala Lys 145 150 155	656
50	gct tcc gcc gtc gcc gaa gtc cct agc ttc cag tgg ctc gac cgg aac	704
55		

EP 2 183 363 B1

	Ala	Ser	Ala	Val	Ala	Glu	Val	Pro	Ser	Phe	Gln	Trp	Leu	Asp	Arg	Asn	
			160					165					170				
5	gtc	acc	atc	gac	acc	ctg	atg	gtc	cag	act	ctg	tcc	cag	gtc	cgg	gct	752
	Val	Thr	Ile	Asp	Thr	Leu	Met	Val	Gln	Thr	Leu	Ser	Gln	Val	Arg	Ala	
		175					180					185					
10	ctc	aat	aag	gcc	ggc	gcc	aat	cct	ccc	tat	gct	g	gtgagttaca				796
	Leu	Asn	Lys	Ala	Gly	Ala	Asn	Pro	Pro	Tyr	Ala						
	190					195					200						
	tggcgacttg	ccttctcgtc	ccctaccttt	cttgacggga	tcggttacct	gacctggagg											856
15	caaaacaaca	acag	cc	caa	ctc	gtc	gtc	tac	gac	ctc	ccc	gac	cgt	gac			905
			Ala	Gln	Leu	Val	Val	Tyr	Asp	Leu	Pro	Asp	Arg	Asp			
							205					210					
20	tgt	gcc	gcc	gct	gcg	tcc	aac	ggc	gag	ttt	tcg	att	gca	aac	ggc	ggc	953
	Cys	Ala	Ala	Ala	Ala	Ser	Asn	Gly	Glu	Phe	Ser	Ile	Ala	Asn	Gly	Gly	
		215					220					225					
25	gcc	gcc	aac	tac	agg	agc	tac	atc	gac	gct	atc	cgc	aag	cac	atc	att	1001
	Ala	Ala	Asn	Tyr	Arg	Ser	Tyr	Ile	Asp	Ala	Ile	Arg	Lys	His	Ile	Ile	
		230					235					240					
30	gag	tac	tcg	gac	atc	cgg	atc	atc	ctg	gtt	atc	gag	ccc	gac	tcg	atg	1049
	Glu	Tyr	Ser	Asp	Ile	Arg	Ile	Ile	Leu	Val	Ile	Glu	Pro	Asp	Ser	Met	
	245					250					255					260	
35	gcc	aac	atg	gtg	acc	aac	atg	aac	gtg	gcc	aag	tgc	agc	aac	gcc	gcg	1097
	Ala	Asn	Met	Val	Thr	Asn	Met	Asn	Val	Ala	Lys	Cys	Ser	Asn	Ala	Ala	
				265					270						275		
40	tcg	acg	tac	cac	gag	ttg	acc	gtg	tac	gcg	ctc	aag	cag	ctg	aac	ctg	1145
	Ser	Thr	Tyr	His	Glu	Leu	Thr	Val	Tyr	Ala	Leu	Lys	Gln	Leu	Asn	Leu	
			280						285					290			
45	ccc	aac	gtc	gcc	atg	tat	ctc	gac	gcc	ggc	cac	gcc	ggc	tgg	ctc	ggc	1193
	Pro	Asn	Val	Ala	Met	Tyr	Leu	Asp	Ala	Gly	His	Ala	Gly	Trp	Leu	Gly	
		295					300						305				
50	tgg	ccc	gcc	aac	atc	cag	ccc	gcc	gcc	gag	ctg	ttt	gcc	ggc	atc	tac	1241
	Trp	Pro	Ala	Asn	Ile	Gln	Pro	Ala	Ala	Glu	Leu	Phe	Ala	Gly	Ile	Tyr	
		310				315						320					
55	aat	gat	gcc	ggc	aag	ccg	gct	gcc	gtc	cgc	ggc	ctg	gcc	act	aac	gtc	1289
	Asn	Asp	Ala	Gly	Lys	Pro	Ala	Ala	Val	Arg	Gly	Leu	Ala	Thr	Asn	Val	
	325					330					335					340	
60	gcc	aac	tac	aac	gcc	tgg	agc	atc	gct	tcg	gcc	ccg	tcg	tac	acg	tcg	1337
	Ala	Asn	Tyr	Asn	Ala	Trp	Ser	Ile	Ala	Ser	Ala	Pro	Ser	Tyr	Thr	Ser	
				345						350					355		
65	cct	aac	cct	aac	tac	gac	gag	aag	cac	tac	atc	gag	gcc	ttc	agc	ccg	1385
	Pro	Asn	Pro	Asn	Tyr	Asp	Glu	Lys	His	Tyr	Ile	Glu	Ala	Phe	Ser	Pro	
				360					365					370			
70	ctc	ttg	aac	tcg	gcc	ggc	ttc	ccc	gca	cgc	ttc	att	gtc	gac	act	ggc	1433
	Leu	Leu	Asn	Ser	Ala	Gly	Phe	Pro	Ala	Arg	Phe	Ile	Val	Asp	Thr	Gly	
		375						380					385				

EP 2 183 363 B1

	cgc aac ggc aaa caa cct acc g gtatgttttt ttttcttttg tctctgtccc Arg Asn Gly Lys Gln Pro Thr 390 395	1485
5	cccccttttct ccccccttcag ttggcgtcca caaggtctct tagtcctgct tcatctgtga ccaacctccc cccccccggc accgcccaca accgtttgac tctatactct tgggaatggg	1545 1605
10	cgccgaaaact gaccgttcca cag gc caa caa cag tgg ggt gac tgg tgc aat Gly Gln Gln Gln Trp Gly Asp Trp Cys Asn 400 405	1657
15	gtc aag ggc acc ggc ttt ggc gtg cgc ccg acg gcc aac acg ggc cac Val Lys Gly Thr Gly Phe Gly Val Arg Pro Thr Ala Asn Thr Gly His 410 415 420	1705
20	gag ctg gtc gat gcc ttt gtc tgg gtc aag ccc ggc ggc gag tcc gac Glu Leu Val Asp Ala Phe Val Trp Val Lys Pro Gly Gly Glu Ser Asp 425 430 435	1753
25	ggc aca agc gac acc agc gcc gcc cgc tac gac tac cac tgc ggc ctg Gly Thr Ser Asp Thr Ser Ala Ala Arg Tyr Asp Tyr His Cys Gly Leu 440 445 450	1801
30	tcc gat gcc ctg cag cct gcc ccc gag gct gga cag tgg ttc cag gcc Ser Asp Ala Leu Gln Pro Ala Pro Glu Ala Gly Gln Trp Phe Gln Ala 455 460 465	1849
35	tac ttc gag cag ctg ctc acc aac gcc aac ccg ccc ttc taaacctcgt Tyr Phe Glu Gln Leu Leu Thr Asn Ala Asn Pro Pro Phe 470 475 480	1898
40	cataaagaga gagagatggc gggcatgggc ctgattgggt tcattgacca tgcggctctt ctgggggttac atattttacc tacctaccta taaataaggc ggcctatcgg gctctcgctt cgtttattag gtacttggtc ttgtacatac tttgtttata catacagcag ttagcatcca ctattcgttt cgacaaagcg gaactttcca gaaaaaa	1958 2018 2078 2115
45	<210> 11 <211> 482 <212> PRT <213> Chrysosporium lucknowense	
50	<400> 11	
55		

EP 2 183 363 B1

Met Ala Lys Lys Leu Phe Ile Thr Ala Ala Leu Ala Ala Ala Val Leu
1 5 10 15

5 Ala Ala Pro Val Ile Glu Glu Arg Gln Asn Cys Gly Ala Val Trp Thr
20 25 30

10 Gln Cys Gly Gly Asn Gly Trp Gln Gly Pro Thr Cys Cys Ala Ser Gly
35 40 45

Ser Thr Cys Val Ala Gln Asn Glu Trp Tyr Ser Gln Cys Leu Pro Asn
15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	50					55					60					
5	Ser 65	Gln	Val	Thr	Ser	Ser 70	Thr	Thr	Pro	Ser	Ser 75	Thr	Ser	Thr	Ser	Gln 80
10	Arg	Ser	Thr	Ser	Thr 85	Ser	Ser	Ser	Thr	Thr 90	Arg	Ser	Gly	Ser	Ser 95	Ser
15	Ser	Ser	Ser	Thr 100	Thr	Pro	Pro	Pro	Val 105	Ser	Ser	Pro	Val	Thr 110	Ser	Ile
20	Pro	Gly	Gly 115	Ala	Thr	Ser	Thr	Ala 120	Ser	Tyr	Ser	Gly	Asn 125	Pro	Phe	Ser
25	Gly	Val 130	Arg	Leu	Phe	Ala	Asn 135	Asp	Tyr	Tyr	Arg	Ser 140	Glu	Val	His	Asn
30	Leu 145	Ala	Ile	Pro	Ser	Met 150	Thr	Gly	Thr	Leu	Ala 155	Ala	Lys	Ala	Ser	Ala 160
35	Val	Ala	Glu	Val	Pro 165	Ser	Phe	Gln	Trp	Leu 170	Asp	Arg	Asn	Val	Thr 175	Ile
40	Asp	Thr	Leu	Met 180	Val	Gln	Thr	Leu	Ser 185	Gln	Val	Arg	Ala	Leu 190	Asn	Lys
45	Ala	Gly	Ala 195	Asn	Pro	Pro	Tyr	Ala 200	Ala	Gln	Leu	Val	Val 205	Tyr	Asp	Leu
50	Pro	Asp 210	Arg	Asp	Cys	Ala	Ala 215	Ala	Ala	Ser	Asn	Gly 220	Glu	Phe	Ser	Ile
55	Ala	Asn	Gly	Gly	Ala	Ala 230	Asn	Tyr	Arg	Ser	Tyr 235	Ile	Asp	Ala	Ile	Arg 240
60	Lys	His	Ile	Ile	Glu 245	Tyr	Ser	Asp	Ile	Arg 250	Ile	Ile	Leu	Val	Ile 255	Glu
65	Pro	Asp	Ser	Met 260	Ala	Asn	Met	Val	Thr 265	Asn	Met	Asn	Val	Ala 270	Lys	Cys
70	Ser	Asn	Ala 275	Ala	Ser	Thr	Tyr	His 280	Glu	Leu	Thr	Val	Tyr 285	Ala	Leu	Lys
75	Gln	Leu	Asn	Leu	Pro	Asn 295	Val	Ala	Met	Tyr	Leu	Asp 300	Ala	Gly	His	Ala

EP 2 183 363 B1

	Gly Trp Leu Gly Trp Pro Ala Asn Ile Gln Pro Ala Ala Glu Leu Phe	
	305 310 315 320	
5	Ala Gly Ile Tyr Asn Asp Ala Gly Lys Pro Ala Ala Val Arg Gly Leu	
	325 330 335	
10	Ala Thr Asn Val Ala Asn Tyr Asn Ala Trp Ser Ile Ala Ser Ala Pro	
	340 345 350	
15	Ser Tyr Thr Ser Pro Asn Pro Asn Tyr Asp Glu Lys His Tyr Ile Glu	
	355 360 365	
20	Ala Phe Ser Pro Leu Leu Asn Ser Ala Gly Phe Pro Ala Arg Phe Ile	
	370 375 380	
25	Val Asp Thr Gly Arg Asn Gly Lys Gln Pro Thr Gly Gln Gln Gln Trp	
	385 390 395 400	
30	Gly Asp Trp Cys Asn Val Lys Gly Thr Gly Phe Gly Val Arg Pro Thr	
	405 410 415	
35	Ala Asn Thr Gly His Glu Leu Val Asp Ala Phe Val Trp Val Lys Pro	
	420 425 430	
40	Gly Gly Glu Ser Asp Gly Thr Ser Asp Thr Ser Ala Ala Arg Tyr Asp	
	435 440 445	
45	Tyr His Cys Gly Leu Ser Asp Ala Leu Gln Pro Ala Pro Glu Ala Gly	
	450 455 460	
50	Gln Trp Phe Gln Ala Tyr Phe Glu Gln Leu Leu Thr Asn Ala Asn Pro	
	465 470 475 480	

Pro Phe

<210> 12
 <211> 1446
 <212> DNA
 <213> Chrysosporium lucknowense

atggccaaga agcttttcat caccgccgcg cttgcggctg ccgtgttggc ggcccccgtc	60
attgaggagc gccagaactg cggcgctgtg tggactcaat gcggcggtaa cgggtggcaa	120
gggtccacat gctgcgcctc gggctcgacc tgcgttgcgc agaacgagtg gtactctcag	180

EP 2 183 363 B1

	tgcttgccca acagccaggt gacgagttcc accactccgt cgtcgacttc cacctcgag	240
	cgcagcacca gcacctccag cagcaccacc aggagcggca gtcctctctc ctctccacc	300
5	acgccccgc cgtctccag ccccgtagc agcattcccg gcggtgcgac ctccacggcg	360
	agctactctg gcaacccctt ctcgggcgtc cggctcttcg ccaacgacta ctacaggtcc	420
10	gaggtccaca atctcgccat tcctagcatg actggtactc tggcggccaa ggcttcgcgc	480
	gtcgccgaag tccctagctt ccagtggctc gaccggaacg tcaccatcga caccctgatg	540
	gtccagactc tgtcccaggt ccgggctctc aataaggccg gtgccaatcc tccctatgct	600
15	gcccactcg tegtctacga cctccccgac cgtgactgtg ccgccgctgc gtccaacggc	660
	gagttttcga ttgcaaacgg cggcgccgcc aactacagga gctacatcga cgctatccgc	720
	aagcacatca ttgagtactc ggacatccgg atcatcctgg ttatcgagcc cgactcgatg	780
20	gccaacatgg tgaccaacat gaacgtggcc aagtgcagca acgccgcgtc gacgtaccac	840
	gagttgaccg tgtacgcgct caagcagctg aacctgcccc acgtcgccat gtatctcgac	900
25	gccggccacg ccggttggt cggctggccc gccaacatcc agcccgccgc cgagctgttt	960
	gccggcatct acaatgatgc cggcaagccg gctgccgtcc gcggcctggc cactaacgtc	1020
	gccaactaca acgcctggag catcgcttcg gccccgtcgt acacgtcgcc taaccctaac	1080
30	tacgacgaga agcactacat cgaggccttc agcccgtct tgaactcggc cggttcccc	1140
	gcacgcttca ttgtcgacac tggccgcaac ggcaaacaac ctaccggcca acaacagtgg	1200
35	ggtgactggt gcaatgtcaa gggcaccggc tttggcgtgc gcccgacggc caacacgggc	1260
	cacgagctgg tcgatgcctt tgtctgggtc aagcccggcg gcgagtcgga cggcacaagc	1320
	gacaccagcg ccgcccgtc cgactaccac tgcggcctgt ccgatgccct gcagcctgcc	1380
40	cccgaggctg gacagtgggt ccaggcctac ttcgagcagc tgctcaccaa cgccaacccg	1440
	cccttc	1446
45	<210> 13 <211> 2507 <212> DNA <213> Chrysosporium lucknowense	
50	<220> <221> CDS <222> (967)..(1572)	
55	<220> <221> CDS <222> (1666)..(2427)	
	<400> 13	

EP 2 183 363 B1

	ctatataaag gtacttacag ataaaatccc cagggttgaca gcccttttoga gcataagaat	60
	ccaggctcta tagtggccgg ctogaagccc agatttgagc cctattgaga ataactatat	120
5	ggcactggat gaaggattgg cttgaacatc gctacgatgt ttaatcgtg agtcttacag	180
	ggcttcgact ggtgtttctg gaggcgtggg aggagagccc ccccgagttt cttcagtgcc	240
	tggcacactc aatgcctoga cggctagcaa aaaccagggg ggagagactg ggtattgata	300
10	cttctgtatt atagaaattt gtattgggtt ggttgtagct agacttcttg gctacttttc	360
	tgatttgctg tagttgtggg gcatgtatcg cactaggatt aataatgctg gaagaggtgt	420
	agagatgttg cgtcgcgatg ccaccgagat ccggcctggc gtcgtgcctt gcagctcgca	480
15	cgatgcgcgc gacgatgcgg gacacccctc cccctcccc cctccccct acattaggag	540
	gggatccggg gccgctgccc gttttccgtc ttggtattcc cggctgacac aacggttctc	600
20	cagagtgata accgagggcac accagtagca gtccacgatg gcacggcagg ggatacattc	660
	cgcccggaaac aacccaagat ctgggaatcc cagtgcgcgc cccgggggtcc tccgggggtcc	720
	cccgatctcg tctgaaccga cgaactggaa ggaggcgagc cttggaacga tgggggatat	780
25	gatatatatt aagatgcaac attctctccc tccctcctct ctctctccct cccccctct	840
	ctctcctct tccctcctc cccgcggttc tctctccatg agcctcaatt cttgctccga	900
	gcgcggtgta tttccccac gaggaattga caaaagaaaa agaaaaagac aagactctcg	960
30	agaacg atg ggt cgc ggc gct gct ttc cta ggc ctc gcc tcg ctc ctc	1008
	Met Gly Arg Gly Ala Ala Phe Leu Gly Leu Ala Ser Leu Leu	
	1 5 10	
35	gtg ggc gcg gcc aag gcc cag acg ccc ggc gag ggc gag gag gtg cac	1056
	Val Gly Ala Ala Lys Ala Gln Thr Pro Gly Glu Gly Glu Glu Val His	
	15 20 25 30	
	ccg cag atc acg acg tac cgc tgc acc aag gcg gac ggg tgc gag gag	1104
	Pro Gln Ile Thr Thr Tyr Arg Cys Thr Lys Ala Asp Gly Cys Glu Glu	
40	35 40 45	
	aag acc aac tac atc gtg ctg gac gcc cta tcg cac ccg gtc cac cag	1152
	Lys Thr Asn Tyr Ile Val Leu Asp Ala Leu Ser His Pro Val His Gln	
	50 55 60	
45	gtc gac aac ccg tac aac tgc ggc gac tgg ggc cag aag ccc aac gag	1200
	Val Asp Asn Pro Tyr Asn Cys Gly Asp Trp Gly Gln Lys Pro Asn Glu	
	65 70 75	
50	acg gcc tgc ccg gac ctc gag tcg tgc gcc agg aac tgc atc atg gac	1248
	Thr Ala Cys Pro Asp Leu Glu Ser Cys Ala Arg Asn Cys Ile Met Asp	
	80 85 90	
	ccg gtc tcg gac tac ggc cgg cac ggg gtc tcg acc gac ggc acc tcg	1296
	Pro Val Ser Asp Tyr Gly Arg His Gly Val Ser Thr Asp Gly Thr Ser	
55	95 100 105 110	
	ctg cgc ctc aag cag cta gtc ggc ggc aac gtc gtc agc ccg cgc gtc	1344

EP 2 183 363 B1

	Leu	Arg	Leu	Lys	Gln	Leu	Val	Gly	Gly	Asn	Val	Val	Ser	Pro	Arg	Val	
					115					120					125		
5	tac	ctg	ctc	gac	gag	acc	aag	gag	cgc	tac	gag	atg	ctc	aag	ctg	acc	1392
	Tyr	Leu	Leu	Asp	Glu	Thr	Lys	Glu	Arg	Tyr	Glu	Met	Leu	Lys	Leu	Thr	
				130					135					140			
10	ggc	aac	gag	ttc	acc	ttt	gac	gtc	gac	gcc	acc	aag	ctg	ccc	tgc	ggc	1440
	Gly	Asn	Glu	Phe	Thr	Phe	Asp	Val	Asp	Ala	Thr	Lys	Leu	Pro	Cys	Gly	
			145					150					155				
15	atg	aac	agc	gcc	ctc	tac	ctc	tcc	gag	atg	gac	gcc	acc	ggc	gcc	cgg	1488
	Met	Asn	Ser	Ala	Leu	Tyr	Leu	Ser	Glu	Met	Asp	Ala	Thr	Gly	Ala	Arg	
		160					165					170					
20	agc	gag	ctc	aac	ccg	ggc	ggc	gcc	acc	ttt	ggc	acc	ggc	tac	tgc	gac	1536
	Ser	Glu	Leu	Asn	Pro	Gly	Gly	Ala	Thr	Phe	Gly	Thr	Gly	Tyr	Cys	Asp	
	175					180					185					190	
25	gcc	cag	tgc	tac	gtc	acc	ccc	ttc	atc	aac	ggc	ctc	gtacgtattc				1582
	Ala	Gln	Cys	Tyr	Val	Thr	Pro	Phe	Ile	Asn	Gly	Leu					
					195					200							
30	tcctattaat	cttggttctt	ttttcctttt	ctttttttga	tataatagaa	ttaagaagaa											1642
35	ctcgggtgggc	tgacatgaca	cag	ggc	aac	atc	gag	ggc	aag	ggc	gcg	tgc	tgc				1695
				Gly	Asn	Ile	Glu	Gly	Lys	Gly	Ala	Cys	Cys				
					205						210						
40	aac	gag	atg	gat	atc	tgg	gag	gcc	aac	gcg	cgg	gcg	cag	cac	atc	gcg	1743
	Asn	Glu	Met	Asp	Ile	Trp	Glu	Ala	Asn	Ala	Arg	Ala	Gln	His	Ile	Ala	
		215						220					225				
45	ccg	cac	ccg	tgc	agc	aag	gcg	ggg	ccg	tac	ctg	tgc	gag	ggc	gcc	gag	1791
	Pro	His	Pro	Cys	Ser	Lys	Ala	Gly	Pro	Tyr	Leu	Cys	Glu	Gly	Ala	Glu	
		230					235					240					
50	tgc	gag	ttc	gac	ggc	gtg	tgc	gac	aag	aac	ggc	tgc	gcc	tgg	aac	ccg	1839
	Cys	Glu	Phe	Asp	Gly	Val	Cys	Asp	Lys	Asn	Gly	Cys	Ala	Trp	Asn	Pro	
	245					250					255					260	
55	tac	cgg	gtc	aac	gtg	acg	gac	tac	tac	ggc	gag	ggc	gcc	gag	ttc	agg	1887
	Tyr	Arg	Val	Asn	Val	Thr	Asp	Tyr	Tyr	Gly	Glu	Gly	Ala	Glu	Phe	Arg	
					265					270					275		
60	gtg	gac	acg	acc	ccg	ccc	ttc	tcg	gtc	gtc	acg	cag	ttc	cgc	gcc	ggc	1935
	Val	Asp	Thr	Thr	Arg	Pro	Phe	Ser	Val	Val	Thr	Gln	Phe	Arg	Ala	Gly	
				280					285					290			
65	ggc	gac	gcg	ggg	ggc	ggc	aag	ctc	gag	agc	atc	tac	cgg	ctc	ttc	gtc	1983
	Gly	Asp	Ala	Gly	Gly	Gly	Lys	Leu	Glu	Ser	Ile	Tyr	Arg	Leu	Phe	Val	
			295					300					305				
70	cag	gac	ggc	agg	gtg	atc	gag	tcg	tac	gtc	gtc	gac	aag	ccc	ggc	ctg	2031
	Gln	Asp	Gly	Arg	Val	Ile	Glu	Ser	Tyr	Val	Val	Asp	Lys	Pro	Gly	Leu	
		310					315					320					
75	ccc	ccg	acg	gac	cgc	atg	acg	gac	gag	ttc	tgc	gcc	gcc	acc	ggc	gcc	2079
	Pro	Pro	Thr	Asp	Arg	Met	Thr	Asp	Glu	Phe	Cys	Ala	Ala	Thr	Gly	Ala	
	325					330					335					340	

EP 2 183 363 B1

	gcc cgc ttc acg gag ctc ggc gcc atg gag gcc atg ggc gac gcc ctg	2127
	Ala Arg Phe Thr Glu Leu Gly Ala Met Glu Ala Met Gly Asp Ala Leu	
	345 350 355	
5	acg cgc ggc atg gtc ctc gcc ctc agc atc tgg tgg agc gag ggc gac	2175
	Thr Arg Gly Met Val Leu Ala Leu Ser Ile Trp Trp Ser Glu Gly Asp	
	360 365 370	
10	aac atg aac tgg ctc gac tcg ggc gag gcc ggc ccc tgc gac ccg gac	2223
	Asn Met Asn Trp Leu Asp Ser Gly Glu Ala Gly Pro Cys Asp Pro Asp	
	375 380 385	
15	gag ggc aac ccg tcc aac atc atc cgc gtc cag ccc gac ccg gag gtc	2271
	Glu Gly Asn Pro Ser Asn Ile Ile Arg Val Gln Pro Asp Pro Glu Val	
	390 395 400	
20	gtc ttc agc aac ctg cgc tgg ggc gag atc ggc tca acc tac gag tcc	2319
	Val Phe Ser Asn Leu Arg Trp Gly Glu Ile Gly Ser Thr Tyr Glu Ser	
	405 410 415 420	
25	gcc gtc gac ggg ccc gtc ggc aag ggc aag ggc aag ggc aag ggc aag	2367
	Ala Val Asp Gly Pro Val Gly Lys Gly Lys Gly Lys Gly Lys Gly Lys	
	425 430 435	
30	gct ccc gcc ggc gac ggc aac ggg aag gag aag agc aat ggc aag cgc	2415
	Ala Pro Ala Gly Asp Gly Asn Gly Lys Glu Lys Ser Asn Gly Lys Arg	
	440 445 450	
35	ttc agg agg ttc tgagcaacct tgatattatt tttttctttc tttccttcac	2467
	Phe Arg Arg Phe	
	455	
40	cagttaatta gttgcctttg attagaaaga gagagagaaa	2507
45	<210> 14	
	<211> 456	
	<212> PRT	
	<213> Chrysosporium lucknowense	
50	<400> 14	
55	Met Gly Arg Gly Ala Ala Phe Leu Gly Leu Ala Ser Leu Leu Val Gly	
	1 5 10 15	
60	Ala Ala Lys Ala Gln Thr Pro Gly Glu Gly Glu Glu Val His Pro Gln	
	20 25 30	
65	Ile Thr Thr Tyr Arg Cys Thr Lys Ala Asp Gly Cys Glu Glu Lys Thr	
	35 40 45	
70	Asn Tyr Ile Val Leu Asp Ala Leu Ser His Pro Val His Gln Val Asp	
	50 55 60	
75	Asn Pro Tyr Asn Cys Gly Asp Trp Gly Gln Lys Pro Asn Glu Thr Ala	
	65 70 75 80	

EP 2 183 363 B1

	Cys	Pro	Asp	Leu	Glu	Ser	Cys	Ala	Arg	Asn	Cys	Ile	Met	Asp	Pro	Val	
					85					90					95		
5	Ser	Asp	Tyr	Gly	Arg	His	Gly	Val	Ser	Thr	Asp	Gly	Thr	Ser	Leu	Arg	
				100					105					110			
10	Leu	Lys	Gln	Leu	Val	Gly	Gly	Asn	Val	Val	Ser	Pro	Arg	Val	Tyr	Leu	
			115					120					125				
15	Leu	Asp	Glu	Thr	Lys	Glu	Arg	Tyr	Glu	Met	Leu	Lys	Leu	Thr	Gly	Asn	
		130					135					140					
20	Glu	Phe	Thr	Phe	Asp	Val	Asp	Ala	Thr	Lys	Leu	Pro	Cys	Gly	Met	Asn	
	145					150					155					160	
25	Ser	Ala	Leu	Tyr	Leu	Ser	Glu	Met	Asp	Ala	Thr	Gly	Ala	Arg	Ser	Glu	
					165					170					175		
30	Leu	Asn	Pro	Gly	Gly	Ala	Thr	Phe	Gly	Thr	Gly	Tyr	Cys	Asp	Ala	Gln	
				180					185					190			
35	Cys	Tyr	Val	Thr	Pro	Phe	Ile	Asn	Gly	Leu	Gly	Asn	Ile	Glu	Gly	Lys	
			195					200					205				
40	Gly	Ala	Cys	Cys	Asn	Glu	Met	Asp	Ile	Trp	Glu	Ala	Asn	Ala	Arg	Ala	
		210					215					220					
45	Gln	His	Ile	Ala	Pro	His	Pro	Cys	Ser	Lys	Ala	Gly	Pro	Tyr	Leu	Cys	
	225					230				235					240		
50	Glu	Gly	Ala	Glu	Cys	Glu	Phe	Asp	Gly	Val	Cys	Asp	Lys	Asn	Gly	Cys	
					245					250					255		
55	Ala	Trp	Asn	Pro	Tyr	Arg	Val	Asn	Val	Thr	Asp	Tyr	Tyr	Gly	Glu	Gly	
				260					265					270			
60	Ala	Glu	Phe	Arg	Val	Asp	Thr	Thr	Arg	Pro	Phe	Ser	Val	Val	Thr	Gln	
			275					280					285				
65	Phe	Arg	Ala	Gly	Gly	Asp	Ala	Gly	Gly	Gly	Lys	Leu	Glu	Ser	Ile	Tyr	
		290					295					300					
70	Arg	Leu	Phe	Val	Gln	Asp	Gly	Arg	Val	Ile	Glu	Ser	Tyr	Val	Val	Asp	
	305					310					315					320	

EP 2 183 363 B1

	Lys	Pro	Gly	Leu	Pro	Pro	Thr	Asp	Arg	Met	Thr	Asp	Glu	Phe	Cys	Ala	
				325						330					335		
5	Ala	Thr	Gly	Ala	Ala	Arg	Phe	Thr	Glu	Leu	Gly	Ala	Met	Glu	Ala	Met	
				340					345					350			
10	Gly	Asp	Ala	Leu	Thr	Arg	Gly	Met	Val	Leu	Ala	Leu	Ser	Ile	Trp	Trp	
			355					360					365				
15	Ser	Glu	Gly	Asp	Asn	Met	Asn	Trp	Leu	Asp	Ser	Gly	Glu	Ala	Gly	Pro	
		370					375					380					
20	Cys	Asp	Pro	Asp	Glu	Gly	Asn	Pro	Ser	Asn	Ile	Ile	Arg	Val	Gln	Pro	
	385					390					395					400	
25	Asp	Pro	Glu	Val	Val	Phe	Ser	Asn	Leu	Arg	Trp	Gly	Glu	Ile	Gly	Ser	
					405					410					415		
30	Thr	Tyr	Glu	Ser	Ala	Val	Asp	Gly	Pro	Val	Gly	Lys	Gly	Lys	Gly	Lys	
				420					425					430			
35	Gly	Lys	Gly	Lys	Ala	Pro	Ala	Gly	Asp	Gly	Asn	Gly	Lys	Glu	Lys	Ser	
		435						440					445				
40	Asn	Gly	Lys	Arg	Phe	Arg	Arg	Phe									
		450					455										
45	<210> 15																
	<211> 1368																
	<212> DNA																
	<213> Chrysosporium lucknowense																
50	<400> 15																
55																	

EP 2 183 363 B1

atggggtcgcg gcgctgcttt cctaggcctc gcctcgctcc tcgtgggcgc ggccaaggcc 60
 cagacgcccc gcgagggcga ggaggtgcac ccgcagatca cgacgtaccg ctgcaccaag 120
 5 gcggacgggt gcgaggagaa gaccaactac atcgtgctgg acgccctatc gcacccggtc 180
 caccaggtcg acaacccgta caactgcggc gactggggcc agaagcccaa cgagacggcc 240
 10 tgccccggacc tcgagtcgtg cgccaggaac tgcacatgg acccggtctc ggactacggc 300
 cggcacgggg tctcgaccga cggcacctcg ctgcgcctca agcagctagt cggcggcaac 360
 gtcgtcagcc cgcgcgtcta cctgctcgac gagaccaagg agcgtacga gatgctcaag 420
 15 ctgaccggca acgagttcac ctttgacgtc gacgccacca agctgccctg cggcatgaac 480
 agcgccctct acctctccga gatggacgcc accggcgccc ggagcgagct caacccgggc 540
 ggcgccacct ttggcaccgg ctactgcgac gccagtgct acgtcacccc cttcatcaac 600
 20 ggcctcggca acatcgagg caagggcgcg tgctgcaacg agatggatat ctgggaggcc 660
 aacgcgcggg cgcagcacat cgcgccgcac ccgtgcagca aggcggggcc gtacctgtgc 720
 25 gagggcgccg agtgcgagtt cgacggcgtg tgcgacaaga acggctgcgc ctggaacccg 780
 taccgggtca acgtgacgga ctactacggc gagggcgccg agttcagggt ggacacgacc 840
 30 cggcccttct cggtcgtcac gcagttccgc gccggcgggc acgcgggggg cggcaagctc 900
 gagagcatct accggctctt cgtccaggac ggaggggtga tcgagtcgta cgtcgtcgac 960
 aagcccggcc tgcccccgac ggaccgcatg acggacgagt tctgcgccgc caccggcgcc 1020
 35 gcccgcttca cggagctcgg cgccatggag gccatggggc acgccctgac gcgcggcatg 1080
 gtccctcgccc tcagcatctg gtggagcgag ggcgacaaca tgaactggct cgactcgggc 1140
 gaggccggcc cctgcgaccc ggacgagggc aaccgtcca acatcatccg cgtccagccc 1200
 40 gacccggagg tcgtcttcag caacctgcgc tggggcgaga tcggctcaac ctacgagtc 1260
 gccgtcgacg ggcccgtcgg caagggcaag ggcaagggca agggcaaggc tcccgcgggc 1320
 45 gacggcaacg ggaaggagaa gagcaatggc aagcgcttca ggaggttc 1368

<210> 16

<211> 1305

<212> DNA

50 <213> Chrysosporium lucknowense

<220>

<221> CDS

<222> (103)..(545)

55

<220>

<221> CDS

<222> (683)..(903)

EP 2 183 363 B1

<220>

<221> CDS

<222> (986)..(1062)

5 <400> 16

cgacaaagac ccgtcagtga ttaataataa ttagtagcag tttctttctt tcaagactca 60

10 agaataactcc tttccgccat cgtggcagcg tttagattca tc atg cag ccg ttt 114
Met Gln Pro Phe
1

15 ctg ctc ttg ttc ctc tcg tcg gtc acg gcg gcg agc ccc ctg acg gcg 162
Leu Leu Leu Phe Leu Ser Ser Val Thr Ala Ala Ser Pro Leu Thr Ala
5 10 15 20

ctc gac aag cgg cag cag gcg acg ttg tgc gag cag tac ggc tac tgg 210
Leu Asp Lys Arg Gln Gln Ala Thr Leu Cys Glu Gln Tyr Gly Tyr Trp
25 30 35

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	tcg ggc aac ggt tac gag gtc aac aac aac aac tgg ggc aag gat tcg Ser Gly Asn Gly Tyr Glu Val Asn Asn Asn Asn Trp Gly Lys Asp Ser 40 45 50	258
5	gcc tcg ggc ggc cat cag tgc acc tac gtc gac agc agc agc tcc agc Ala Ser Gly Gly His Gln Cys Thr Tyr Val Asp Ser Ser Ser Ser 55 60 65	306
10	ggc gtc gcc tgg cac acg acc tgg cag tgg gaa gga ggc cag aac cag Gly Val Ala Trp His Thr Thr Trp Gln Trp Glu Gly Gly Gln Asn Gln 70 75 80	354
15	gtc aag agc ttc gcc aac tgc ggc ctg cag gtg ccc aag ggc agg acc Val Lys Ser Phe Ala Asn Cys Gly Leu Gln Val Pro Lys Gly Arg Thr 85 90 95 100	402
	atc tcg tcc atc agc aac ctg cag acc tcc atc tcg tgg tcc tac agc Ile Ser Ser Ile Ser Asn Leu Gln Thr Ser Ile Ser Trp Ser Tyr Ser 105 110 115	450
20	aac acc aac atc cgc gcc aac gtg gcc tac gac ctc ttc acc gcg gca Asn Thr Asn Ile Arg Ala Asn Val Ala Tyr Asp Leu Phe Thr Ala Ala 120 125 130	498
25	gac ccg aac cac gcg acc agc agc ggc gac tac gag ctc atg atc tg Asp Pro Asn His Ala Thr Ser Ser Gly Asp Tyr Glu Leu Met Ile Trp 135 140 145	545
30	gtcagttttt tttttctttt ttcttttctt ctcttttctt ttcttttctt ttctctgtt ttattttctt atccattgct tcgcccctctt tccttaaccc tgctgactct ctcttcttgt caatgatact gtaatag g ctg gcg aga ttc ggc gac gtc tac ccc atc ggc Leu Ala Arg Phe Gly Asp Val Tyr Pro Ile Gly 150 155	605 665 716
35	tcg tcc cag ggc cac gtc aac gtg gcc ggc cag gac tgg gag ctg tgg Ser Ser Gln Gly His Val Asn Val Ala Gly Gln Asp Trp Glu Leu Trp 160 165 170 175	764
40	acg ggc ttc aac ggc aac atg cgg gtc tac agc ttc gta gcg ccc agc Thr Gly Phe Asn Gly Asn Met Arg Val Tyr Ser Phe Val Ala Pro Ser 180 185 190	812
45	ccc cgc aac agc ttc agc gcc aac gtc aag gac ttc ttc aac tat ctc Pro Arg Asn Ser Phe Ser Ala Asn Val Lys Asp Phe Phe Asn Tyr Leu 195 200 205	860
50	cag tcc aac cag ggc ttc ccg gcc agc agc caa tac ctt ctc a Gln Ser Asn Gln Gly Phe Pro Ala Ser Ser Gln Tyr Leu Leu 210 215 220	903
	gtaaggagac gagatctcga acagcatacc atatatgcgt gcggtacaag tgcactaacc cccttttttt tcccgttcgc ag tc ttc cag gcg ggc acc gag ccc ttc acc Ile Phe Gln Ala Gly Thr Glu Pro Phe Thr 225 230	963 1014
55	ggc ggc gag acc acc ctt acc gtc aac aac tac tct gca agg gtt gct Gly Gly Glu Thr Thr Leu Thr Val Asn Asn Tyr Ser Ala Arg Val Ala 235 240 245	1062

EP 2 183 363 B1

taaacaggaa ggccgaggat ggcccccaag gccgttgccg gttcacgagc tctcttcttt 1122
 tcaagtgctg tacatacata attagcgtac caagtcatag ctgtttgtca gcttcaaact 1182
 5 aagtgctcgc ccacaaaaga ggggggaggg gaaaataaca aattgccgaa cgcagtgata 1242
 agcttctggg agcgttgaaa gcagtctaca gtaggtggct gtacgaagga aaagagtgcc 1302
 tta 1305
 10

<210> 17
 <211> 247
 <212> PRT
 <213> Chrysosporium lucknowense

15

<400> 17

20

Met Gln Pro Phe Leu Leu Leu Phe Leu Ser Ser Val Thr Ala Ala Ser
 1 5 10 15

Pro Leu Thr Ala Leu Asp Lys Arg Gln Gln Ala Thr Leu Cys Glu Gln
 20 25 30

25

Tyr Gly Tyr Trp Ser Gly Asn Gly Tyr Glu Val Asn Asn Asn Trp
 35 40 45

30

Gly Lys Asp Ser Ala Ser Gly Gly His Gln Cys Thr Tyr Val Asp Ser
 50 55 60

35

Ser Ser Ser Ser Gly Val Ala Trp His Thr Thr Trp Gln Trp Glu Gly
 65 70 75 80

Gly Gln Asn Gln Val Lys Ser Phe Ala Asn Cys Gly Leu Gln Val Pro
 85 90 95

40

Lys Gly Arg Thr Ile Ser Ser Ile Ser Asn Leu Gln Thr Ser Ile Ser
 100 105 110

45

Trp Ser Tyr Ser Asn Thr Asn Ile Arg Ala Asn Val Ala Tyr Asp Leu
 115 120 125

50

Phe Thr Ala Ala Asp Pro Asn His Ala Thr Ser Ser Gly Asp Tyr Glu
 130 135 140

55

Leu Met Ile Trp Leu Ala Arg Phe Gly Asp Val Tyr Pro Ile Gly Ser
 145 150 155 160

Ser Gln Gly His Val Asn Val Ala Gly Gln Asp Trp Glu Leu Trp Thr
 165 170 175

EP 2 183 363 B1

	Gly	Phe	Asn	Gly	Asn	Met	Arg	Val	Tyr	Ser	Phe	Val	Ala	Pro	Ser	Pro
				180					185					190		
5	Arg	Asn	Ser	Phe	Ser	Ala	Asn	Val	Lys	Asp	Phe	Phe	Asn	Tyr	Leu	Gln
			195					200					205			
10	Ser	Asn	Gln	Gly	Phe	Pro	Ala	Ser	Ser	Gln	Tyr	Leu	Leu	Ile	Phe	Gln
		210					215					220				
15	Ala	Gly	Thr	Glu	Pro	Phe	Thr	Gly	Gly	Glu	Thr	Thr	Leu	Thr	Val	Asn
	225					230					235					240
	Asn	Tyr	Ser	Ala	Arg	Val	Ala									
					245											

20 <210> 18
 <211> 741
 <212> DNA
 <213> Chrysosporium lucknowense

25 <400> 18

	atgcagccgt	ttctgctctt	gttcctctcg	tcggtcacgg	cggcgagccc	cctgacggcg	60
30	ctcgacaagc	ggcagcaggc	gacgttgtgc	gagcagtacg	gctactggtc	gggcaacggc	120
	tacgaggtca	acaacaacaa	ctggggcaag	gattcggcct	cgggcggcca	tcagtgcacc	180
	tacgtcgaca	gcagcagctc	cagcggcgtc	gcctggcaca	cgacctggca	gtgggaagga	240
35	ggccagaacc	aggtcaagag	cttcgccaac	tgcggcctgc	aggtgcccaa	gggcaggacc	300
	atctcgtcca	tcagcaacct	gcagacctcc	atctcgtggt	cctacagcaa	caccaacatc	360
	cgcgccaacg	tggcctacga	cctcttcacc	gcggcagacc	cgaaccacgc	gaccagcagc	420
40	ggcgactacg	agctcatgat	ctggctggcg	agattcggcg	acgtctaccc	catcggtctg	480
	tcccagggcc	acgtcaacgt	ggccggccag	gactgggagc	tgtggacggg	cttcaacggc	540
45	aacatgcggg	tctacagctt	cgtagcgccc	agccccgca	acagcttcag	cgccaacgtc	600
	aaggacttct	tcaactatct	ccagtccaac	cagggcttcc	cggccagcag	ccaatacctt	660
	ctcatcttcc	aggcgggcac	cgagcccttc	accggcgggc	agaccaccct	taccgtcaac	720
50	aactactctg	caagggttgc	t				741

55 <210> 19
 <211> 1448
 <212> DNA
 <213> Chrysosporium lucknowense

<220>

<221> CDS
<222> (166)..(597)

5 <220>
<221> CDS
<222> (708)..(738)

10 <220>
<221> CDS
<222> (846)..(972)

15 <220>
<221> CDS
<222> (1074)..(1158)

<400> 19

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	atatataagt gagagtgttt tttgactgcc ccgggttctg gtagttgaag ggaagttcga	60
	tgtctctctgc tgtcgtcgtc ctcgtcgtc tcgtcggcat cctccatccg tccgcctttg	120
5	ataacccgct ccccgactca gtcaagacga cgcatacttg gcacc atg cat ctc tcc Met His Leu Ser 1	177
10	gcc acc acc ggg ttc ctc gcc ctc ccg gcc ctg gcc ctg gcc cag ctc Ala Thr Thr Gly Phe Leu Ala Leu Pro Ala Leu Ala Leu Ala Gln Leu 5 10 15 20	225
15	tcg ggc agc ggc cag acg acc cgg tac tgg gac tgc tgc aag ccg agc Ser Gly Ser Gly Gln Thr Thr Arg Tyr Trp Asp Cys Cys Lys Pro Ser 25 30 35	273
20	tgc gcc tgg ccc ggc aag ggc ccc tcg tct ccg gtg cag gcc tgc gac Cys Ala Trp Pro Gly Lys Gly Pro Ser Ser Pro Val Gln Ala Cys Asp 40 45 50	321
25	aag aac gac aac ccg ctc aac gac ggc ggc tcc acc ccg tcc ggc tgc Lys Asn Asp Asn Pro Leu Asn Asp Gly Gly Ser Thr Arg Ser Gly Cys 55 60 65	369
30	gac gcg ggc ggc agc gcc tac atg tgc tcc tcc cag agc ccc tgg gcc Asp Ala Gly Gly Ser Ala Tyr Met Cys Ser Ser Gln Ser Pro Trp Ala 70 75 80	417
35	gtc agc gac gag ctg tcg tac ggc tgg gcg gcc gtc aag ctc gcc ggc Val Ser Asp Glu Leu Ser Tyr Gly Trp Ala Ala Val Lys Leu Ala Gly 85 90 95 100	465
40	agc tcc gag tcg cag tgg tgc tgc gcc tgc tac gag ctg acc ttc acc Ser Ser Glu Ser Gln Trp Cys Cys Ala Cys Tyr Glu Leu Thr Phe Thr 105 110 115	513
45	agc ggg ccg gtc gcg ggc aag aag atg att gtg cag gcg acc aac acc Ser Gly Pro Val Ala Gly Lys Lys Met Ile Val Gln Ala Thr Asn Thr 120 125 130	561
50	ggc ggc gac ctg ggc gac aac cac ttt gac ctg gcc gtgagttgcc Gly Gly Asp Leu Gly Asp Asn His Phe Asp Leu Ala 135 140	607

EP 2 183 363 B1

	tcccccttctc cccggaccgc tcagattaga tgagattaga ctttgctcgt aaatcggtcc	667
5	aagattccccg ttgactgacc aacaaacatc atacgggcag atc ccc ggt ggc ggt Ile Pro Gly Gly Gly 145	722
10	gtc ggt att ttc aac g gtaagctggt gcccccgac ccctcccccg acccctcccc Val Gly Ile Phe Asn 150	778
	cttttcctcc agcgagccga gttgggatcg ccgagatcga gaactcacac aacttctctc	838
15	tcgacag cc tgc acc gac cag tac ggc gct ccc ccg aac ggc tgg ggc Ala Cys Thr Asp Gln Tyr Gly Ala Pro Pro Asn Gly Trp Gly 160 165	886
20	gac cgc tac ggc ggc atc cat tcc aag gaa gag tgc gaa tcc ttc ccg Asp Arg Tyr Gly Gly Ile His Ser Lys Glu Glu Cys Glu Ser Phe Pro 170 175 180	934
	gag gcc ctc aag ccc ggc tgc aac tgg cgc ttc gac tg gtacgttgct Glu Ala Leu Lys Pro Gly Cys Asn Trp Arg Phe Asp Trp 185 190 195	982
25	ttgacatacc ggaacccaat tcctccaacc ccccccttt tctcccccaa ctccgggggt	1042
	agtcggaatg tcgcgactga ccctatttca g g ttc caa aac gcc gac aac ccg Phe Gln Asn Ala Asp Asn Pro 200	1095
30	tcg gtc acc ttc cag gag gtg gcc tgc ccg tcg gag ctc acg tcc aag Ser Val Thr Phe Gln Glu Val Ala Cys Pro Ser Glu Leu Thr Ser Lys 205 210 215 220	1143
35	agc ggc tgc tcc cgt taagagggaa gagagggggc tggaaggacc gaaagattca Ser Gly Cys Ser Arg 225	1198
	acctctgctc ctgctgggga agctcggggc cgagtgtgaa actggtgtaa atattgtggc	1258
40	acacacaagc tactacagtc cgtctcgccg tccggctaac tagccttgct gcggatctgt	1318
	ccatcttcgg tccgaactgt ccgttgctgt tttggctcgg tgccatcatct tctcccaacc	1378
45	tagtcaagaa tgaatcgtga gagaggctga gagagataag atcgacttca gaaatccagg	1438
	gttgaaagca	1448
50	<210> 20 <211> 225 <212> PRT <213> Chrysosporium lucknowense	
55	<400> 20	

EP 2 183 363 B1

	Met	His	Leu	Ser	Ala	Thr	Thr	Gly	Phe	Leu	Ala	Leu	Pro	Ala	Leu	Ala	
	1				5					10					15		
5	Leu	Ala	Gln	Leu	Ser	Gly	Ser	Gly	Gln	Thr	Thr	Arg	Tyr	Trp	Asp	Cys	
					20				25					30			
10	Cys	Lys	Pro	Ser	Cys	Ala	Trp	Pro	Gly	Lys	Gly	Pro	Ser	Ser	Pro	Val	
			35					40					45				
15	Gln	Ala	Cys	Asp	Lys	Asn	Asp	Asn	Pro	Leu	Asn	Asp	Gly	Gly	Ser	Thr	
		50					55					60					
20	Arg	Ser	Gly	Cys	Asp	Ala	Gly	Gly	Ser	Ala	Tyr	Met	Cys	Ser	Ser	Gln	
	65					70					75					80	
25	Ser	Pro	Trp	Ala	Val	Ser	Asp	Glu	Leu	Ser	Tyr	Gly	Trp	Ala	Ala	Val	
					85					90					95		
30	Lys	Leu	Ala	Gly	Ser	Ser	Glu	Ser	Gln	Trp	Cys	Cys	Ala	Cys	Tyr	Glu	
				100					105					110			
35	Leu	Thr	Phe	Thr	Ser	Gly	Pro	Val	Ala	Gly	Lys	Lys	Met	Ile	Val	Gln	
			115					120					125				
40	Ala	Thr	Asn	Thr	Gly	Gly	Asp	Leu	Gly	Asp	Asn	His	Phe	Asp	Leu	Ala	
		130					135					140					
45	Ile	Pro	Gly	Gly	Gly	Val	Gly	Ile	Phe	Asn	Ala	Cys	Thr	Asp	Gln	Tyr	
	145					150					155				160		
50	Gly	Ala	Pro	Pro	Asn	Gly	Trp	Gly	Asp	Arg	Tyr	Gly	Gly	Ile	His	Ser	
					165					170					175		
55	Lys	Glu	Glu	Cys	Glu	Ser	Phe	Pro	Glu	Ala	Leu	Lys	Pro	Gly	Cys	Asn	
				180					185					190			
60	Trp	Arg	Phe	Asp	Trp	Phe	Gln	Asn	Ala	Asp	Asn	Pro	Ser	Val	Thr	Phe	
			195					200					205				
65	Gln	Glu	Val	Ala	Cys	Pro	Ser	Glu	Leu	Thr	Ser	Lys	Ser	Gly	Cys	Ser	
		210					215					220					
70	Arg																
	225																

EP 2 183 363 B1

<210> 21
 <211> 675
 <212> DNA
 <213> Chrysosporium lucknowense

5
 <400> 21
 atgcatctct cgcaccac cgggttcctc gccctcccg cctggccct ggcccagctc 60
 10 tggggcagcg gccagacgac ccggtactgg gactgctgca agccgagctg cgcctggccc 120
 ggcaagggcc cctcgtctcc ggtgcaggcc tgcgacaaga acgacaaccc gctcaacgac 180
 15 ggcggtcca cccggtccgg ctgacgacg ggcggcagcg cctacatgtg ctctccag 240
 agccctggg ccgtcagcga cgagctgtcg tacggctggg cggccgtcaa gctcgccggc 300
 agctccgagt cgcagtggcg ctgcgcctgc tacgagctga ccttcaccag cgggccggctc 360
 20 gcgggcaaga agatgattgt gcaggcgacc aacaccggcg gcgacctggg cgacaaccac 420
 ttgacctgg ccatcccccg tggcggtgtc ggtattttca acgctgcac cgaccagtac 480
 ggcgtcccc cgaacggctg ggcgaccgc tacggcgga tccattcaa ggaagagtgc 540
 25 gaatccttcc cggaggccct caagcccggc tgcaactggc gcttcgactg gttccaaaac 600
 gccgacaacc cgtcggtcac cttccaggag gtggcctgcc cgtcgagct caggtccaag 660
 30 agcggtgct cccgt 675

<210> 22
 <211> 1658
 <212> DNA
 <213> Chrysosporium lucknowense

35
 <220>
 <221> CDS
 <222> (211)..(1139)
 40
 <220>
 <221> CDS
 <222> (1217)..(1430)
 45
 <400> 22

50
 55

EP 2 183 363 B1

	aggaacgggg aggaggaggg cttggttagg gtcgcgttcg tttggagatt gctgagctct	60
	gagccttcgg tccttggatc cctgcggtcc ccggtctcct ctctctctct ctctctctct	120
5	ctctctctct cttcttccca cgctcgttcg acagacgcct ccccttcttc gctctccttt	180
	ccctcgcacg tagcacacta atagtgcacc atg cgc gtc tct agt ttg gtc gcg	234
	Met Arg Val Ser Ser Leu Val Ala	
10	1 5	
	gcc ctt gct acc ggt ggt ctt gtc gcc gcc acg cct aag ccc aag ggg	282
	Ala Leu Ala Thr Gly Gly Leu Val Ala Ala Thr Pro Lys Pro Lys Gly	
	10 15 20	
15	tcg tcg ccc cct ggg gcc gtg gac gcg aac cct ttc aag ggc aag acg	330
	Ser Ser Pro Pro Gly Ala Val Asp Ala Asn Pro Phe Lys Gly Lys Thr	
	25 30 35 40	
	cag ttc gtc aac ccg gca tgg gcg gcc aag ctg gaa cag acc aaa aag	378
20	Gln Phe Val Asn Pro Ala Trp Ala Ala Lys Leu Glu Gln Thr Lys Lys	
	45 50 55	
25		
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	gcg ttc ctg gcc agg aac gac acc gtc aat gcc gcc aag acg gag aag Ala Phe Leu Ala Arg Asn Asp Thr Val Asn Ala Ala Lys Thr Glu Lys 60 65 70	426
5	gtc cag cag acc agc tcg ttc gtc tgg gtc tcg agg atc gcc gag ctc Val Gln Gln Thr Ser Ser Phe Val Trp Val Ser Arg Ile Ala Glu Leu 75 80 85	474
10	tcc aac atc gac gac gcc atc gcg gct gcc cgc aag gcg cag aag aag Ser Asn Ile Asp Asp Ala Ile Ala Ala Arg Lys Ala Gln Lys Lys 90 95 100	522
15	acg ggc agg agg cag atc gtc ggc ctg gtg ctc tac aac ctt ccg gac Thr Gly Arg Arg Gln Ile Val Gly Leu Val Leu Tyr Asn Leu Pro Asp 105 110 115 120	570
	cgc gac tgc agc gcg ggc gag agc gcg ggc gag ctc agc agc gac aag Arg Asp Cys Ser Ala Gly Glu Ser Ala Gly Glu Leu Ser Ser Asp Lys 125 130 135	618
20	aac ggg ctc gag atc tac aag act gag ttc gtc aag ccc ttc gcc gac Asn Gly Leu Glu Ile Tyr Lys Thr Glu Phe Val Lys Pro Phe Ala Asp 140 145 150	666
25	aag gtg gcg gcc gca aag gac ctc gac ttc gcc atc gtc ctg gag ccc Lys Val Ala Ala Ala Lys Asp Leu Asp Phe Ala Ile Val Leu Glu Pro 155 160 165	714
30	gac tcg ctg gcc aac ctg gtc acc aac ctg ggc atc gag ttc tgc gcc Asp Ser Leu Ala Asn Leu Val Thr Asn Leu Gly Ile Glu Phe Cys Ala 170 175 180	762
	aac gcc gcc ccc gtc tac cgc gag ggc atc gcc tat gcc atc tcc agc Asn Ala Ala Pro Val Tyr Arg Glu Gly Ile Ala Tyr Ala Ile Ser Ser 185 190 195 200	810
35	ctt cag cag cca aac gtg cac ttg tac atc gat gct gcc cac ggc ggc Leu Gln Gln Pro Asn Val His Leu Tyr Ile Asp Ala Ala His Gly Gly 205 210 215	858
40	tgg ctc ggc tgg gac gac aac ctg ccg ctg gcc gcc aag gag ttt gcc Trp Leu Gly Trp Asp Asp Asn Leu Pro Leu Ala Ala Lys Glu Phe Ala 220 225 230	906
45	gag gtg gtc aag ctt gcc ggc gag ggc aag aag atc cgc ggc ttc gtc Glu Val Val Lys Leu Ala Gly Glu Gly Lys Lys Ile Arg Gly Phe Val 235 240 245	954
	acc aac gtg tcc aac tac aac ccc ttc cac gcc gtc gtg cgc gag aac Thr Asn Val Ser Asn Tyr Asn Pro Phe His Ala Val Val Arg Glu Asn 250 255 260	1002
50	ttt acc gag tgg agc aac tcg tgg gac gag tct cac tac gcc tcc tcg Phe Thr Glu Trp Ser Asn Ser Trp Asp Glu Ser His Tyr Ala Ser Ser 265 270 275 280	1050
55	ctc aca ccg ttc ctc gag aaa gag ggg ctg ccg gca cgc ttc atc gtc Leu Thr Pro Phe Leu Glu Lys Glu Gly Leu Pro Ala Arg Phe Ile Val 285 290 295	1098

EP 2 183 363 B1

	gac cag ggt cgc gtt gcc ctc ccg gga gcc cgc aag gag tg	1139
	Asp Gln Gly Arg Val Ala Leu Pro Gly Ala Arg Lys Glu Trp	
	300 305	
5	gtgagtttcg accagattga ccctcgaccc atgcgaccga gattgctgac gattgaattg	1199
	cgtgtcccgt cccccag g ggt gaa tgg tgc aac gtg gca ccc gcc gga ttt	1250
	Gly Glu Trp Cys Asn Val Ala Pro Ala Gly Phe	
	315 320	
10	ggc ccc gcg ccc acg acc agg gtc aac aac acc gtc gtc gat gct ctc	1298
	Gly Pro Ala Pro Thr Thr Arg Val Asn Asn Thr Val Val Asp Ala Leu	
	325 330 335	
15	gtc tgg gtc aag cct ggc ggc gag agc gac ggc gag tgt ggc ttg gct	1346
	Val Trp Val Lys Pro Gly Gly Glu Ser Asp Gly Glu Cys Gly Leu Ala	
	340 345 350	
20	ggc gcc ccc aag gcc ggc cag tgg ttc gac gag tac gcc cag atg ctg	1394
	Gly Ala Pro Lys Ala Gly Gln Trp Phe Asp Glu Tyr Ala Gln Met Leu	
	355 360 365	
25	gtc gag aat gcc cac ccg tct gtc gtc cac aag tgg tagataaatt	1440
	Val Glu Asn Ala His Pro Ser Val Val His Lys Trp	
	370 375 380	
30	ttggaggtccg agaagggtcc cagatagact tttgttttaa aacaaaatgc aaggtgtcga	1500
	cagatactgg cttaacatta accaagcacc atgaacatga cttgtcaaca tattgataca	1560
	ttccgctgct ttcccatacg tgctctcagg tctcagggat caaatggata ggtcggtaat	1620
	gcaaaacgat ccattggata tccagaagag agaaaaaa	1658
35	<210> 23	
	<211> 381	
	<212> PRT	
	<213> Chrysosporium lucknowense	
40	<400> 23	
	Met Arg Val Ser Ser Leu Val Ala Ala Leu Ala Thr Gly Gly Leu Val	
	1 5 10 15	
45	Ala Ala Thr Pro Lys Pro Lys Gly Ser Ser Pro Pro Gly Ala Val Asp	
	20 25 30	
50	Ala Asn Pro Phe Lys Gly Lys Thr Gln Phe Val Asn Pro Ala Trp Ala	
	35 40 45	
55	Ala Lys Leu Glu Gln Thr Lys Lys Ala Phe Leu Ala Arg Asn Asp Thr	
	50 55 60	
	Val Asn Ala Ala Lys Thr Glu Lys Val Gln Gln Thr Ser Ser Phe Val	
	65 70 75 80	

EP 2 183 363 B1

	Trp	Val	Ser	Arg	Ile	Ala	Glu	Leu	Ser	Asn	Ile	Asp	Asp	Ala	Ile	Ala
					85					90					95	
5	Ala	Ala	Arg	Lys	Ala	Gln	Lys	Lys	Thr	Gly	Arg	Arg	Gln	Ile	Val	Gly
				100					105					110		
10	Leu	Val	Leu	Tyr	Asn	Leu	Pro	Asp	Arg	Asp	Cys	Ser	Ala	Gly	Glu	Ser
			115					120					125			
15	Ala	Gly	Glu	Leu	Ser	Ser	Asp	Lys	Asn	Gly	Leu	Glu	Ile	Tyr	Lys	Thr
		130					135					140				
20	Glu	Phe	Val	Lys	Pro	Phe	Ala	Asp	Lys	Val	Ala	Ala	Ala	Lys	Asp	Leu
	145					150					155					160
25	Asn	Leu	Gly	Ile	Glu	Phe	Cys	Ala	Asn	Ala	Ala	Pro	Val	Tyr	Arg	Glu
			180					185						190		
30	Gly	Ile	Ala	Tyr	Ala	Ile	Ser	Ser	Leu	Gln	Gln	Pro	Asn	Val	His	Leu
		195						200					205			
35	Tyr	Ile	Asp	Ala	Ala	His	Gly	Gly	Trp	Leu	Gly	Trp	Asp	Asp	Asn	Leu
	210						215					220				
40	Pro	Leu	Ala	Ala	Lys	Glu	Phe	Ala	Glu	Val	Val	Lys	Leu	Ala	Gly	Glu
	225					230					235					240
45	Gly	Lys	Lys	Ile	Arg	Gly	Phe	Val	Thr	Asn	Val	Ser	Asn	Tyr	Asn	Pro
				245						250					255	
50	Phe	His	Ala	Val	Val	Arg	Glu	Asn	Phe	Thr	Glu	Trp	Ser	Asn	Ser	Trp
			260					265						270		
55	Asp	Glu	Ser	His	Tyr	Ala	Ser	Ser	Leu	Thr	Pro	Phe	Leu	Glu	Lys	Glu
		275						280					285			
60	Gly	Leu	Pro	Ala	Arg	Phe	Ile	Val	Asp	Gln	Gly	Arg	Val	Ala	Leu	Pro
	290						295					300				
65	Gly	Ala	Arg	Lys	Glu	Trp	Gly	Glu	Trp	Cys	Asn	Val	Ala	Pro	Ala	Gly
	305					310					315					320

EP 2 183 363 B1

Phe Gly Pro Ala Pro Thr Thr Arg Val Asn Asn Thr Val Val Asp Ala
325 330 335

5 Leu Val Trp Val Lys Pro Gly Gly Glu Ser Asp Gly Glu Cys Gly Leu
340 345 350

10 Ala Gly Ala Pro Lys Ala Gly Gln Trp Phe Asp Glu Tyr Ala Gln Met
355 360 365

15 Leu Val Glu Asn Ala His Pro Ser Val Val His Lys Trp
370 375 380

<210> 24

<211> 1143

<212> DNA

<213> Chrysosporium lucknowense

<400> 24

EP 2 183 363 B1

	atgcgcgtct ctagtttggg cgcgccctt gctaccggtg gtcttgtcgc cgccacgcct	60
	aagcccaagg ggtcgtcgcc ccctggggcc gtggacgcga accctttcaa gggcaagacg	120
5	cagttcgtca acccggcatt ggcgccaag ctggaacaga ccaaaaaggc gttcctggcc	180
	aggaacgaca ccgtcaatgc cgccaagacg gagaagggtcc agcagaccag ctcgttcgtc	240
10	tgggtctcga ggatcgccga gctctccaac atcgacgacg ccatcgcggc tgcccgaag	300
	gcgcagaaga agacgggcag gaggcagatc gtcggcctgg tgctctacaa ccttcggac	360
	cgcgactgca gcgcgggcga gagcgcgggc gagctcagca gcgacaagaa cgggctcgag	420
15	atctacaaga ctgagttcgt caagcccttc gccgacaagg tggcgggccgc aaaggacctc	480
	gacttcgcca tcgtcctgga gcccgactcg ctggccaacc tggtcacca cctgggcac	540
	gagttctgcg ccaacgcccgc cccgtctac cgcgagggca tcgcctatgc catctccagc	600
20	cttcagcagc caaacgtgca cttgtacatc gatgctgccc acggcggtg gctcggtg	660
	gacgacaacc tgccgctggc cgccaaggag tttgccgagg tggtaagct tgccggcgag	720
25	ggcaagaaga tccgcggctt cgtcaccaac gtgtccaact acaaccctt ccacgcccgc	780
	gtgcgcgaga actttaccga gtggagcaac tcgtgggacg agtctcacta cgctcctcg	840
	ctcacaccgt tcctcgagaa agaggggctg ccggcacgct tcctcgtcga ccagggtcgc	900
30	gttgccctcc cgggagcccg caaggagtgg ggtgaatggt gcaacgtggc acccgccgga	960
	tttgggccccg cgcccacgac cagggtaaac aacaccgtcg tcgatgctct cgtctgggtc	1020
	aagcctggcg gcgagagcga cggcgagtgt ggcttggctg gcgcccccaa ggccggccag	1080
35	tggttcgacg agtacgcca gatgctggtc gagaatgccc acccgctctgt cgtccacaag	1140
	tgg	1143
40	<210> 25 <211> 1683 <212> DNA <213> Chrysosporium lucknowense	
45	<220> <221> CDS <222> (76) .. (148)	
50	<220> <221> CDS <222> (223)..(286)	
55	<220> <221> CDS <222> (438)..(700)	
	<220> <221> CDS	

EP 2 183 363 B1

<222> (780)..(1531)

<400> 25

5	atctgccggt tgcccacgtc gccgtcgact gcttgtccgc ttcctacctg cagcctcttt	60
	cagagaccat caaac atg cgt act ctt acg ttc gtg ctg gca gcc gcc ccg	111
	Met Arg Thr Leu Thr Phe Val Leu Ala Ala Ala Pro	
	1 5 10	
10	gtg gct gtg ctt gcc caa tct cct ctg tgg ggc cag t gtatgtaatt	158
	Val Ala Val Leu Ala Gln Ser Pro Leu Trp Gly Gln	
	15 20	
15	gccttactcg gaaaatagtc accactagag ggacttaagc tcactacttc ctgtttcaca	218
	atag gc ggc ggt caa ggc tgg aca ggt ccc acg acc tgc gtt tct ggc	266
	Cys Gly Gly Gln Gly Trp Thr Gly Pro Thr Thr Cys Val Ser Gly	
	30 35	
20	gca gta tgc caa ttc gtc aa gtcagtaact gcttttattt cttttctctc	316
	Ala Val Cys Gln Phe Val Asn	
	40 45	
25	tgggattacg atttcgtttt gcacttagct tggttctgca tttcattggt gtattgttct	376
	ctttttgtgt gtgagagggt ttattaccac ctaaaggcca tttgctaaca aatctcccca	436
	g t gac tgg tac tcc caa tgc gtg ccc gga tgc agc aac cct cct acg	483
	Asp Trp Tyr Ser Gln Cys Val Pro Gly Ser Ser Asn Pro Pro Thr	
30	50 55 60	
	ggc acc acc agc agc acc act gga agc acc ccg gct cct act ggc ggc	531
	Gly Thr Thr Ser Ser Thr Thr Gly Ser Thr Pro Ala Pro Thr Gly Gly	
	65 70 75	
35	ggc ggc agc gga acc ggc ctc cac gac aaa ttc aag gcc aag ggc aag	579
	Gly Gly Ser Gly Thr Gly Leu His Asp Lys Phe Lys Ala Lys Gly Lys	
	80 85 90	
40		
45		
50		
55		

EP 2 183 363 B1

	ctc tac ttc gga acc gag atc gat cac tac cat ctc aac aac aat gcc	627
	Leu Tyr Phe Gly Thr Glu Ile Asp His Tyr His Leu Asn Asn Asn Ala	
	95 100 105	
5	ttg acc aac att gtc aag aaa gac ttt ggt caa gtc act cac gag aac	675
	Leu Thr Asn Ile Val Lys Lys Asp Phe Gly Gln Val Thr His Glu Asn	
	110 115 120 125	
10	agc ttg aag tgg gat gct act gag c gtgagtgacc tctcctcctt	720
	Ser Leu Lys Trp Asp Ala Thr Glu	
	130	
	ctcccgaaca ataatagata attacgagcc ggttcgaggc tgacattgcg cgattctag	779
15	cg agc cgc aat caa ttc aac ttt gcc aac gcc gac gcg gtt gtc aac	826
	Pro Ser Arg Asn Gln Phe Asn Phe Ala Asn Ala Asp Ala Val Val Asn	
	135 140 145	
20	ttt gcc cag gcc aac ggc aag ctc atc cgc ggc cac acc ctc ctc tgg	874
	Phe Ala Gln Ala Asn Gly Lys Leu Ile Arg Gly His Thr Leu Leu Trp	
	150 155 160 165	
	cac tct cag ctg ccg cag tgg gtg cag aac atc aac gac cgc aac acc	922
	His Ser Gln Leu Pro Gln Trp Val Gln Asn Ile Asn Asp Arg Asn Thr	
	170 175 180	
25	ttg acc cag gtc atc gag aac cac gtc acc acc ctt gtc act cgc tac	970
	Leu Thr Gln Val Ile Glu Asn His Val Thr Thr Leu Val Thr Arg Tyr	
	185 190 195	
30	aag ggc aag atc ctc cac tgg gac gtc gtt aac gag atc ttt gcc gag	1018
	Lys Gly Lys Ile Leu His Trp Asp Val Val Asn Glu Ile Phe Ala Glu	
	200 205 210	
35	gac ggc tcg ctc cgc gac agc gtc ttc agc cgc gtc ctc ggc gag gac	1066
	Asp Gly Ser Leu Arg Asp Ser Val Phe Ser Arg Val Leu Gly Glu Asp	
	215 220 225	
	ttt gtc ggc atc gcc ttc cgc gcc gcc cgc gcc gcc gat ccc aac gcc	1114
	Phe Val Gly Ile Ala Phe Arg Ala Ala Arg Ala Ala Asp Pro Asn Ala	
	230 235 240 245	
40	aag ctc tac atc aac gac tac aac ctc gac att gcc aac tac gcc aag	1162
	Lys Leu Tyr Ile Asn Asp Tyr Asn Leu Asp Ile Ala Asn Tyr Ala Lys	
	250 255 260	
45	gtg acc cgg ggc atg gtc gag aag gtc aac aag tgg atc gcc cag ggc	1210
	Val Thr Arg Gly Met Val Glu Lys Val Asn Lys Trp Ile Ala Gln Gly	
	265 270 275	
50	atc ccg atc gac ggc atc ggc acc cag tgc cac ctg gcc ggg ccc ggc	1258
	Ile Pro Ile Asp Gly Ile Gly Thr Gln Cys His Leu Ala Gly Pro Gly	
	280 285 290	
	ggg tgg aac acg gcc gcc ggc gtc ccc gac gcc ctc aag gcc ctc gcc	1306
	Gly Trp Asn Thr Ala Ala Gly Val Pro Asp Ala Leu Lys Ala Leu Ala	
	295 300 305	
55	gcg gcc aac gtc aag gag atc gcc atc acc gag ctc gac atc gcc ggc	1354
	Ala Ala Asn Val Lys Glu Ile Ala Ile Thr Glu Leu Asp Ile Ala Gly	
	310 315 320 325	

EP 2 183 363 B1

gcc tcc gcc aac gac tac ctc acc gtc atg aac gcc tgc ctc cag gtc 1402
Ala Ser Ala Asn Asp Tyr Leu Thr Val Met Asn Ala Cys Leu Gln Val
330 335 340

5 tcc aag tgc gtc ggc atc acc gtc tgg ggc gtc tct gac aag gac agc 1450
Ser Lys Cys Val Gly Ile Thr Val Trp Gly Val Ser Asp Lys Asp Ser
345 350 355

10 tgg agg tcg agc agc aac ccg ctc ctc ttc gac agc aac tac cag cca 1498
Trp Arg Ser Ser Ser Asn Pro Leu Leu Phe Asp Ser Asn Tyr Gln Pro
360 365 370

15 aag gcg gca tac aat gct ctg att aat gcc ttg taagaggagg tatattattt 1551
Lys Ala Ala Tyr Asn Ala Leu Ile Asn Ala Leu
375 380

ttagaggcaa tgaagctagg aggaaagagg ggaagtgagg taattagcta ggacaggcaa 1611
atctagcagc aattataagt caacactata taaaatattc ctataatggc ttgtgcttcg 1671
20 gtgtgcaaaa aa 1683

<210> 26
<211> 384
25 <212> PRT
<213> Chrysosporium lucknowense
<400> 26

30 Met Arg Thr Leu Thr Phe Val Leu Ala Ala Ala Pro Val Ala Val Leu
1 5 10 15

Ala Gln Ser Pro Leu Trp Gly Gln Cys Gly Gly Gln Gly Trp Thr Gly
35 20 25 30

Pro Thr Thr Cys Val Ser Gly Ala Val Cys Gln Phe Val Asn Asp Trp
40 35 40 45

Tyr Ser Gln Cys Val Pro Gly Ser Ser Asn Pro Pro Thr Gly Thr Thr
50 50 55 60

45 Ser Ser Thr Thr Gly Ser Thr Pro Ala Pro Thr Gly Gly Gly Gly Ser
65 70 75 80

50 Gly Thr Gly Leu His Asp Lys Phe Lys Ala Lys Gly Lys Leu Tyr Phe
85 90 95

Gly Thr Glu Ile Asp His Tyr His Leu Asn Asn Asn Ala Leu Thr Asn
55 100 105 110

Ile Val Lys Lys Asp Phe Gly Gln Val Thr His Glu Asn Ser Leu Lys
115 120 125

EP 2 183 363 B1

	Trp	Asp	Ala	Thr	Glu	Pro	Ser	Arg	Asn	Gln	Phe	Asn	Phe	Ala	Asn	Ala	
	130						135					140					
5	Asp	Ala	Val	Val	Asn	Phe	Ala	Gln	Ala	Asn	Gly	Lys	Leu	Ile	Arg	Gly	
	145					150					155					160	
	His	Thr	Leu	Leu	Trp	His	Ser	Gln	Leu	Pro	Gln	Trp	Val	Gln	Asn	Ile	
10					165					170					175		
	Asn	Asp	Arg	Asn	Thr	Leu	Thr	Gln	Val	Ile	Glu	Asn	His	Val	Thr	Thr	
15				180					185					190			
	Leu	Val	Thr	Arg	Tyr	Lys	Gly	Lys	Ile	Leu	His	Trp	Asp	Val	Val	Asn	
			195					200					205				
20	Glu	Ile	Phe	Ala	Glu	Asp	Gly	Ser	Leu	Arg	Asp	Ser	Val	Phe	Ser	Arg	
	210						215					220					
	Val	Leu	Gly	Glu	Asp	Phe	Val	Gly	Ile	Ala	Phe	Arg	Ala	Ala	Arg	Ala	
25	225					230					235					240	
	Ala	Asp	Pro	Asn	Ala	Lys	Leu	Tyr	Ile	Asn	Asp	Tyr	Asn	Leu	Asp	Ile	
30				245						250					255		
	Ala	Asn	Tyr	Ala	Lys	Val	Thr	Arg	Gly	Met	Val	Glu	Lys	Val	Asn	Lys	
				260					265					270			
35	Trp	Ile	Ala	Gln	Gly	Ile	Pro	Ile	Asp	Gly	Ile	Gly	Thr	Gln	Cys	His	
			275					280					285				
40	Leu	Ala	Gly	Pro	Gly	Gly	Trp	Asn	Thr	Ala	Ala	Gly	Val	Pro	Asp	Ala	
	290						295					300					
	Leu	Lys	Ala	Leu	Ala	Ala	Ala	Asn	Val	Lys	Glu	Ile	Ala	Ile	Thr	Glu	
45	305					310					315					320	
	Leu	Asp	Ile	Ala	Gly	Ala	Ser	Ala	Asn	Asp	Tyr	Leu	Thr	Val	Met	Asn	
					325					330					335		
50	Ala	Cys	Leu	Gln	Val	Ser	Lys	Cys	Val	Gly	Ile	Thr	Val	Trp	Gly	Val	
				340					345					350			
55	Ser	Asp	Lys	Asp	Ser	Trp	Arg	Ser	Ser	Ser	Asn	Pro	Leu	Leu	Phe	Asp	
			355					360					365				

EP 2 183 363 B1

Ser Asn Tyr Gln Pro Lys Ala Ala Tyr Asn Ala Leu Ile Asn Ala Leu
370 375 380

<210> 27

<211> 1152

<212> DNA

<213> Chrysosporium lucknowense

<400> 27

```

atgcgtactc ttacgttcgt gctggcagcc gccccggtgg ctgtgcttgc ccaatctcct      60
ctgtgggggcc agtgcggcgg tcaaggctgg acagggtccca cgacctgcgt ttctggcgca      120
gtatgccaat tCGTcaatga ctggtactcc caatgcgtgc ccggatcgag caaccctcct      180
acggggcacca ccagcagcac cactggaagc accccggctc ctactggcgg cggcggcagc      240
ggaaccgggcc tccacgacaa attcaaggcc aagggaagc tctacttcgg aaccgagatc      300
gatcactacc atctcaacaa caatgccttg accaacattg tcaagaaaga ctttgggtcaa      360
gtcactcacg agaacagctt gaagtgggat gctactgagc cgagccgcaa tcaattcaac      420
tttgccaacg ccgacgcggt tgtcaacttt gcccaggcca acggcaagct catccgcggc      480
cacaccctcc tctggcactc tcagctgccg cagtgggtgc agaacaatcaa cgaccgcaac      540
accttgaccc aggtcatcga gaaccacgtc accacccttg tcactcgcta caagggaag      600
atcctccact gggacgtcgt taacgagatc ttgtccgagg acggctcgct ccgcgacagc      660
gtcttcagcc gcgtcctcgg cgaggacttt gtcggcatcg ccttcgcgcg cggccgcgcc      720
gccgatccca acgccaagct ctacatcaac gactacaacc tcgacattgc caactacgcc      780
aagggtgaccc ggggcatggt cgagaaggct aacaagtga tgcgccaggg catcccgatc      840
gacggcatcg gcacccagtg ccacctggcc gggcccggcg ggtggaacac ggccgcccggc      900
gtccccgacg ccctcaaggc cctcgccgcg gccaacgtca aggagatcgc catcaccgag      960
ctcgacatcg ccggcgctc cgccaacgac tacctcaccg tcatgaacgc ctgcctccag     1020
gtctccaagt gcgtcggcat caccgtctgg ggcgtctctg acaaggacag ctggaggctc     1080
agcagcaacc cgctcctctt cgacagcaac taccagccaa aggcggcata caatgctctg     1140
attaatgcct tg                                     1152

```

<210> 28

<211> 1194

<212> DNA

<213> Chrysosporium lucknowense

<220>

<221> CDS

<222> (76) .. (367)

<220>
<221> CDS
<222> (480)..(850)

5 <400> 28

10

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	atgaatcaga agacgacact gggcaactaa acagcttgca gcagagtttt gtgccttgca	60
	taggccctcg atatc atg gtc tcg ttc act ctc ctc ctc acg gtc atc gcc	111
5	Met Val Ser Phe Thr Leu Leu Leu Thr Val Ile Ala	
	1 5 10	
	gct gcg gtg acg acg gcc agc cct ctc gag gtg gtc aag cgc ggc atc	159
	Ala Ala Val Thr Thr Ala Ser Pro Leu Glu Val Val Lys Arg Gly Ile	
10	15 20 25	
	cag ccg ggc acg ggc acc cac gag ggg tac ttc tac tcg ttc tgg acc	207
	Gln Pro Gly Thr Gly Thr His Glu Gly Tyr Phe Tyr Ser Phe Trp Thr	
	30 35 40	
15	gac ggc cgt ggc tcg gtc gac ttc aac ccc ggg ccc cgc ggc tcg tac	255
	Asp Gly Arg Gly Ser Val Asp Phe Asn Pro Gly Pro Arg Gly Ser Tyr	
	45 50 55 60	
	agc gtc acc tgg aac aac gtc aac aac tgg gtt ggc ggc aag ggc tgg	303
20	Ser Val Thr Trp Asn Asn Val Asn Asn Trp Val Gly Gly Lys Gly Trp	
	65 70 75	
	aac ccg ggc ccg ccg cgc aag att gcg tac aac ggc acc tgg aac aac	351
	Asn Pro Gly Pro Pro Arg Lys Ile Ala Tyr Asn Gly Thr Trp Asn Asn	
25	80 85 90	
	tac aac gtg aac agc t gtgcgttgtc ctctcttttc tcccttttcgc ttgttttct	407
	Tyr Asn Val Asn Ser	
	95	
30	tgatgattgg gatccatttt aaaagagaag gaaaaaaaaa acaaaggaaa atagaagata	467
	actaacgccca ag ct ctg gca gac ctc gcc ctg tac ggc tgg act cgc aac	517
	Ser Leu Ala Asp Leu Ala Leu Tyr Gly Trp Thr Arg Asn	
35	100 105 110	
	ccg ctg gtc gag tat tac atc gtg gag gca tac ggc acg tac aac ccc	565
	Pro Leu Val Glu Tyr Tyr Ile Val Glu Ala Tyr Gly Thr Tyr Asn Pro	
	115 120 125	
40	tcg tcg ggc acg gcg cgg ctg ggc acc atc gag gac gac ggc ggc gtg	613
	Ser Ser Gly Thr Ala Arg Leu Gly Thr Ile Glu Asp Asp Gly Gly Val	
	130 135 140	
	tac gac atc tac aag acg acg cgg tac aac cag ccg tcc atc gag ggg	661
45	Tyr Asp Ile Tyr Lys Thr Thr Arg Tyr Asn Gln Pro Ser Ile Glu Gly	
	145 150 155	
	acc tcc acc ttc gac cag tac tgg tcc gtc cgc cgc cag aag cgc gtc	709
	Thr Ser Thr Phe Asp Gln Tyr Trp Ser Val Arg Arg Gln Lys Arg Val	
50	160 165 170	
	ggc ggc act atc gac acg ggc aag cac ttt gac gag tgg aag cgc cag	757
	Gly Gly Thr Ile Asp Thr Gly Lys His Phe Asp Glu Trp Lys Arg Gln	
	175 180 185 190	
55		

EP 2 183 363 B1

ggc aac ctc cag ctc ggc acc tgg aac tac atg atc atg gcc acc gag 805
 Gly Asn Leu Gln Leu Gly Thr Trp Asn Tyr Met Ile Met Ala Thr Glu
 195 200 205

5 ggc tac cag agc tct ggt tcg gcc act atc gag gtc cgg gag gcc 850
 Gly Tyr Gln Ser Ser Gly Ser Ala Thr Ile Glu Val Arg Glu Ala
 210 215 220

10 taaagaagcc aggcgccttt cttttgtttt gcaggagggg gtagaggggg ggggggaggg 910
 aaaacgaaaa gtagcagggt ggttttatgc cggcagccgt gggccattcg agtgcaacct 970
 gtatctctct ctctcccaag tctccgggct ccttctcaga gaacttcaat atgtctgggg 1030

15 acaaaccacc ttgtgaaata caacggtaat tatctaagtt tgagtgccct atcgtatgct 1090
 tctgaaaatt tcctgctcct tgatacaagt cggtttgagc cgagccaatg agactgtgtc 1150
 gattgataga ggccctgaag gatcaagcgc gatgcaacaa ttaa 1194

20 <210> 29
 <211> 221
 <212> PRT
 <213> *Chrysosporium lucknowense*

25 <400> 29

30 Met Val Ser Phe Thr Leu Leu Leu Thr Val Ile Ala Ala Ala Val Thr
 1 5 10 15

Thr Ala Ser Pro Leu Glu Val Val Lys Arg Gly Ile Gln Pro Gly Thr
 20 25 30

35 Gly Thr His Glu Gly Tyr Phe Tyr Ser Phe Trp Thr Asp Gly Arg Gly
 35 40 45

40 Ser Val Asp Phe Asn Pro Gly Pro Arg Gly Ser Tyr Ser Val Thr Trp
 50 55 60

45 Asn Asn Val Asn Asn Trp Val Gly Gly Lys Gly Trp Asn Pro Gly Pro
 65 70 75 80

Pro Arg Lys Ile Ala Tyr Asn Gly Thr Trp Asn Asn Tyr Asn Val Asn
 85 90 95

50 Ser Ser Leu Ala Asp Leu Ala Leu Tyr Gly Trp Thr Arg Asn Pro Leu
 100 105 110

55 Val Glu Tyr Tyr Ile Val Glu Ala Tyr Gly Thr Tyr Asn Pro Ser Ser
 115 120 125

EP 2 183 363 B1

	Gly Thr Ala Arg Leu Gly Thr Ile Glu Asp Asp Gly Gly Val Tyr Asp	
	130 135 140	
5	Ile Tyr Lys Thr Thr Arg Tyr Asn Gln Pro Ser Ile Glu Gly Thr Ser	
	145 150 155 160	
10	Thr Phe Asp Gln Tyr Trp Ser Val Arg Arg Gln Lys Arg Val Gly Gly	
	165 170 175	
15	Thr Ile Asp Thr Gly Lys His Phe Asp Glu Trp Lys Arg Gln Gly Asn	
	180 185 190	
20	Leu Gln Leu Gly Thr Trp Asn Tyr Met Ile Met Ala Thr Glu Gly Tyr	
	195 200 205	
25	Gln Ser Ser Gly Ser Ala Thr Ile Glu Val Arg Glu Ala	
	210 215 220	
30	<210> 30	
	<211> 663	
	<212> DNA	
	<213> Chrysosporium lucknowense	
35	<400> 30	
	atgggtctcgt tcaactctcct cctcacggtc atcgccgctg cggtgacgac ggccagccct	60
	ctcgaggtgg tcaagcgcg g catccagccg ggcacgggca cccacgaggg gtacttctac	120
40	tcgttctgga ccgacggccg tggctcggtc gacttcaacc ccgggccccg cggctcgtac	180
	agcgtcacct ggaacaacgt caacaactgg gttggcggca agggctggaa cccgggccccg	240
45	ccgcgcaaga ttgcgtacaa cggcacctgg aacaactaca acgtgaacag ctctctggca	300
	gacctcgccc tgtacggctg gactcgcaac ccgctggctg agtattacat cgtggaggga	360
50	tacggcacgt acaaccctc gtcgggcacg gcgcggctgg gcaccatcga ggacgacggc	420
	ggcgtgtacg acatctacaa gacgacgcg tacaaccagc cgtccatcga ggggacctcc	480
	accttcgacc agtactggtc cgtccgccc cagaagcgcg tcggcgggcac tatcgacacg	540
	ggcaagcact ttgacgagtg gaagcgccag ggcaacctcc agctcggcac ctggaactac	600
	atgatcatgg ccaccgaggg ctaccagagc tctggttcgg ccactatcga ggtccgggag	660
	gcc	663
55	<210> 31	
	<211> 1598	
	<212> DNA	
	<213> Chrysosporium lucknowense	

<220>
<221> CDS
<222> (69) .. (398)

5 <220>
 <221> CDS
 <222> (467)..(1375)

10 <400> 31

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	tggtctctcca acaccctgc ccctcagcct cccgctcgtc agtgttgtat cctcttctgg	60
5	ccggtatc atg cac tcc aaa gct ttc ttg gca gcg ctt ctt gcg cct gcc Met His Ser Lys Ala Phe Leu Ala Ala Leu Leu Ala Pro Ala 1 5 10	110
10	gtc tca ggg caa ctg aac gac ctc gcc gtc agg gct gga ctc aag tac Val Ser Gly Gln Leu Asn Asp Leu Ala Val Arg Ala Gly Leu Lys Tyr 15 20 25 30	158
15	ttt ggt act gct ctt agc gag agc gtc atc aac agt gat act cgg tat Phe Gly Thr Ala Leu Ser Glu Ser Val Ile Asn Ser Asp Thr Arg Tyr 35 40 45	206
20	gct gcc atc ctc agc gac aag agc atg ttc ggc cag ctc gtc ccc gag Ala Ala Ile Leu Ser Asp Lys Ser Met Phe Gly Gln Leu Val Pro Glu 50 55 60	254
25	aat ggc atg aag tgg gat gct act gag ccg tcc cgt ggc cag ttc aac Asn Gly Met Lys Trp Asp Ala Thr Glu Pro Ser Arg Gly Gln Phe Asn 65 70 75	302
30	tac gcc tcg ggc gac atc acg gcc aac acg gcc aag aag aat ggc cag Tyr Ala Ser Gly Asp Ile Thr Ala Asn Thr Ala Lys Lys Asn Gly Gln 80 85 90	350
35	ggc atg cgt tgc cac acc atg gtc tgg tac agc cag ctc ccg agc tgg Gly Met Arg Cys His Thr Met Val Trp Tyr Ser Gln Leu Pro Ser Trp 95 100 105 110	398
40	gtgagtgcag ccccccctcat cacatccacc agtctgcagt catactcacg agtataataa	458
45	caccgcag gtc tcc tcg ggc tcg tgg acc agg gac tcg ctc acc tcg gtc Val Ser Ser Gly Ser Trp Thr Arg Asp Ser Leu Thr Ser Val 115 120	508
50	atc gag acg cac atg aac aac gtc atg ggc cac tac aag ggc caa tgc Ile Glu Thr His Met Asn Asn Val Met Gly His Tyr Lys Gly Gln Cys 125 130 135 140	556
55	tac gcc tgg gat gtc atc aac gag gcc atc aat gac gac ggc aac tcc Tyr Ala Trp Asp Val Ile Asn Glu Ala Ile Asn Asp Asp Gly Asn Ser 145 150 155	604
60	tgg cgc gac aac gtc ttt ctc cgg acc ttt ggg acc gac tac ttc gcc Trp Arg Asp Asn Val Phe Leu Arg Thr Phe Gly Thr Asp Tyr Phe Ala 160 165 170	652
65	ctg tcc ttc aac cta gcc aag aag gcc gat ccc gat acc aag ctg tac Leu Ser Phe Asn Leu Ala Lys Lys Ala Asp Pro Asp Thr Lys Leu Tyr 175 180 185	700

EP 2 183 363 B1

	tac aac gac tac aac ctc gag tac aac cag gcc aag acg gac cgc gct Tyr Asn Asp Tyr Asn Leu Glu Tyr Asn Gln Ala Lys Thr Asp Arg Ala 190 195 200	748
5	gtt gag ctc gtc aag atg gtc cag gcc gcc gcc gcg ccc atc gac ggt Val Glu Leu Val Lys Met Val Gln Ala Ala Gly Ala Pro Ile Asp Gly 205 210 215 220	796
10	gtc gcc ttc cag gcc cac ctc att gtc gcc tcg acc ccg acg cgc tcg Val Gly Phe Gln Gly His Leu Ile Val Gly Ser Thr Pro Thr Arg Ser 225 230 235	844
15	cag ctg gcc acc gcc ctc cag cgc ttc acc gcg ctc gcc ctc gag gtc Gln Leu Ala Thr Ala Leu Gln Arg Phe Thr Ala Leu Gly Leu Glu Val 240 245 250	892
	gcc tac acc gag ctc gac atc cgc cac tcg agc ctg ccg gcc tct tcg Ala Tyr Thr Glu Leu Asp Ile Arg His Ser Ser Leu Pro Ala Ser Ser 255 260 265	940
20	tcg gcg ctc gcg acc cag gcc aac gac ttc gcc aac gtg gtc gcc tct Ser Ala Leu Ala Thr Gln Gly Asn Asp Phe Ala Asn Val Val Gly Ser 270 275 280	988
25	tgc ctc gac acc gcc gcc tgc gtc gcc gtc acc gtc tgg gcc ttc acc Cys Leu Asp Thr Ala Gly Cys Val Gly Val Thr Val Trp Gly Phe Thr 285 290 295 300	1036
30	gat gcg cac tcg tgg atc ccg aac acg ttc ccc gcc cag gcc gac gcc Asp Ala His Ser Trp Ile Pro Asn Thr Phe Pro Gly Gln Gly Asp Ala 305 310 315	1084
	ctg atc tac gac agc aac tac aac aag aag ccc gcg tgg acc tcg atc Leu Ile Tyr Asp Ser Asn Tyr Asn Lys Lys Pro Ala Trp Thr Ser Ile 320 325 330	1132
35	tcg tcc gtc ctg gcc gcc aag gcc acc gcc gcc ccg ccc gcc tcg tcc Ser Ser Val Leu Ala Ala Lys Ala Thr Gly Ala Pro Pro Ala Ser Ser 335 340 345	1180
40	tcc acc acc ctc gtc acc atc acc acc cct ccg ccg gca tcc acc acc Ser Thr Thr Leu Val Thr Ile Thr Thr Pro Pro Pro Ala Ser Thr Thr 350 355 360	1228
	gcc tcc tcc tcc tcc agt gcc acg ccc acg agc gtc ccg acg cag acg Ala Ser Ser Ser Ser Ser Ala Thr Pro Thr Ser Val Pro Thr Gln Thr 365 370 375 380	1276
45	agg tgg gga cag tgc gcc gcc atc gga tgg acg ggg ccg acc cag tgc Arg Trp Gly Gln Cys Gly Gly Ile Gly Trp Thr Gly Pro Thr Gln Cys 385 390 395	1324
50	gag agc cca tgg acc tgc cag aag ctg aac gac tgg tac tgg cag tgc Glu Ser Pro Trp Thr Cys Gln Lys Leu Asn Asp Trp Tyr Trp Gln Cys 400 405 410	1372
55	ctg taagggacga ggtcccgtg agggcgagtc ttctggattg ttgaagttgg Leu	1425
	cagcataatt tggtagggtt cctagacaga gaccatcggg ggccgatcca ggaaacactg	1485

EP 2 183 363 B1

gctgcatccg accgggattt tctgacttct tctgcatgta ttgaagggtg tgttgagaac 1545

caactgtaaa taagggacca gcagatcaaa gctggagggtt caagcagaat aaa 1598

5

<210> 32

<211> 413

<212> PRT

<213> Chrysosporium lucknowense

10

<400> 32

15

Met His Ser Lys Ala Phe Leu Ala Ala Leu Leu Ala Pro Ala Val Ser
1 5 10 15

20

Gly Gln Leu Asn Asp Leu Ala Val Arg Ala Gly Leu Lys Tyr Phe Gly
20 25 30

Thr Ala Leu Ser Glu Ser Val Ile Asn Ser Asp Thr Arg Tyr Ala Ala
35 40 45

25

Ile Leu Ser Asp Lys Ser Met Phe Gly Gln Leu Val Pro Glu Asn Gly
50 55 60

30

Met Lys Trp Asp Ala Thr Glu Pro Ser Arg Gly Gln Phe Asn Tyr Ala
65 70 75 80

35

Ser Gly Asp Ile Thr Ala Asn Thr Ala Lys Lys Asn Gly Gln Gly Met
85 90 95

Arg Cys His Thr Met Val Trp Tyr Ser Gln Leu Pro Ser Trp Val Ser
100 105 110

40

Ser Gly Ser Trp Thr Arg Asp Ser Leu Thr Ser Val Ile Glu Thr His
115 120 125

45

Met Asn Asn Val Met Gly His Tyr Lys Gly Gln Cys Tyr Ala Trp Asp
130 135 140

50

Val Ile Asn Glu Ala Ile Asn Asp Asp Gly Asn Ser Trp Arg Asp Asn
145 150 155 160

Val Phe Leu Arg Thr Phe Gly Thr Asp Tyr Phe Ala Leu Ser Phe Asn
165 170 175

55

Leu Ala Lys Lys Ala Asp Pro Asp Thr Lys Leu Tyr Tyr Asn Asp Tyr
180 185 190

EP 2 183 363 B1

	Asn	Leu	Glu	Tyr	Asn	Gln	Ala	Lys	Thr	Asp	Arg	Ala	Val	Glu	Leu	Val	
			195					200					205				
5	Lys	Met	Val	Gln	Ala	Ala	Gly	Ala	Pro	Ile	Asp	Gly	Val	Gly	Phe	Gln	
		210					215					220					
10	Gly	His	Leu	Ile	Val	Gly	Ser	Thr	Pro	Thr	Arg	Ser	Gln	Leu	Ala	Thr	
	225					230					235					240	
15	Ala	Leu	Gln	Arg	Phe	Thr	Ala	Leu	Gly	Leu	Glu	Val	Ala	Tyr	Thr	Glu	
				245						250					255		
20	Leu	Asp	Ile	Arg	His	Ser	Ser	Leu	Pro	Ala	Ser	Ser	Ser	Ala	Leu	Ala	
				260					265					270			
25	Thr	Gln	Gly	Asn	Asp	Phe	Ala	Asn	Val	Val	Gly	Ser	Cys	Leu	Asp	Thr	
			275					280					285				
30	Ala	Gly	Cys	Val	Gly	Val	Thr	Val	Trp	Gly	Phe	Thr	Asp	Ala	His	Ser	
	290						295					300					
35	Trp	Ile	Pro	Asn	Thr	Phe	Pro	Gly	Gln	Gly	Asp	Ala	Leu	Ile	Tyr	Asp	
	305					310					315					320	
40	Ser	Asn	Tyr	Asn	Lys	Lys	Pro	Ala	Trp	Thr	Ser	Ile	Ser	Ser	Val	Leu	
				325						330					335		
45	Ala	Ala	Lys	Ala	Thr	Gly	Ala	Pro	Pro	Ala	Ser	Ser	Ser	Thr	Thr	Leu	
				340					345					350			
50	Val	Thr	Ile	Thr	Thr	Pro	Pro	Pro	Ala	Ser	Thr	Thr	Ala	Ser	Ser	Ser	
			355					360					365				
55	Ser	Ser	Ala	Thr	Pro	Thr	Ser	Val	Pro	Thr	Gln	Thr	Arg	Trp	Gly	Gln	
	370						375					380					
60	Cys	Gly	Gly	Ile	Gly	Trp	Thr	Gly	Pro	Thr	Gln	Cys	Glu	Ser	Pro	Trp	
	385					390					395					400	
65	Thr	Cys	Gln	Lys	Leu	Asn	Asp	Trp	Tyr	Trp	Gln	Cys	Leu				
				405						410							

<210> 33

<211> 1239

<212> DNA

<213> Chrysosporium lucknowense

EP 2 183 363 B1

<400> 33

	atgcactcca aagctttctt ggcagcgctt cttgcgcctg ccgtctcagg gcaactgaac	60
5	gacctcgccg tcagggctgg actcaagtac tttggtactg ctcttagcga gagcgatcatc	120
	aacagtgata ctcggtatgc tgccatcctc agcgacaaga gcatgttcgg ccagctcgtc	180
10	cccgagaatg gcatgaagtg ggatgctact gagccgtccc gtggccagtt caactacgcc	240
	tcgggcgaca tcacggccaa cacggccaag aagaatggcc agggcatgcg ttgccacacc	300
	atggtctggt acagccagct cccgagctgg gtctcctcgg gctcgtggac cagggactcg	360
15	ctcacctcgg tcatcgagac gcacatgaac aacgtcatgg gccactacaa gggccaatgc	420
	tacgcctggg atgtcatcaa cgaggccatc aatgacgacg gcaactcctg gcgcgacaac	480
20	gtctttctcc ggacctttgg gaccgactac ttcgccctgt ccttcaacct agccaagaag	540
	gccgatcccg ataccaagct gtactacaac gactacaacc tcgagtacaa ccaggccaag	600
	acggaccgcg ctgttgagct cgtcaagatg gtccaggccg ccggcgcgcc catcgacggt	660
25	gtcggcttcc agggccacct cattgtcggc tcgaccccg aacgctcga gctggccacc	720
	gccctccagc gcttcaccgc gtcgggctc gaggtcgcct acaccgagct cgacatccgc	780
	cactcgagcc tgccggcctc ttcgtcggcg ctgcgcgacc agggcaacga cttcgccaac	840
30	gtggtcgggt cttgcctcga caccgcccgc tgcgtcggcg tcaccgtctg gggcttcacc	900
	gatgcgcact cgtggatccc gaacacgttc cccggccagg gcgacgccct gatctacgac	960
35	agcaactaca acaagaagcc cgcgtggacc tcgatctcgt ccgtcctggc cgccaaggcc	1020
	accggcgccc cgcccgctc gtcctccacc accctcgtca ccatcaccac ccctccgccg	1080
	gcatccacca ccgcctcctc ctccctccagt gccacgccc aagagcgtccc gacgcagacg	1140
40	aggtggggac agtgcggcgg catcggatgg acggggccga cccagtgcga gagcccatgg	1200
	acctgccaga agctgaacga ctggtactgg cagtgcctg	1239

<210> 34

45 <211> 1664

<212> DNA

<213> Chrysosporium lucknowense

<220>

50 <221> CDS

<222> (136)..(393)

<220>

55 <221> CDS

<222> (496)..(1131)

<220>

<221> CDS

<222> (1256) .. (1486)

<400> 34

5

10

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	gattaaatag ggtcgctctg ctcgccgctc tcttgggttt gctctcgtca agcatcctcc	60
	cccacctcgc gccgcaagta cggtaacaggc ctgcaaccga ggatcgagcc cgtcaaagaa	120
5	cgagtcaaaa tcaag atg cat ctc tcc tcg tct ctc ctc ctc ctc gcc gcc Met His Leu Ser Ser Ser Leu Leu Leu Leu Ala Ala 1 5 10	171
10	ttg ccc ctg ggc atc gcc ggc aag ggc aag ggc cac ggc cac ggc ccc Leu Pro Leu Gly Ile Ala Gly Lys Gly Lys Gly His Gly His Gly Pro 15 20 25	219
15	cat acc ggg ctc cac acc ctc gcc aag cag gcc ggc ctc aag tac ttc His Thr Gly Leu His Thr Leu Ala Lys Gln Ala Gly Leu Lys Tyr Phe 30 35 40	267
20	ggc tct gcc acc gac tct ccc ggc cag cgt gag cgc gcc ggc tac gag Gly Ser Ala Thr Asp Ser Pro Gly Gln Arg Glu Arg Ala Gly Tyr Glu 45 50 55 60	315
25	gac aag tac gcc cag tac gac cag atc atg tgg aag tcg ggc gag ttc Asp Lys Tyr Ala Gln Tyr Asp Gln Ile Met Trp Lys Ser Gly Glu Phe 65 70 75	363
30	ggc ctg acg acc ccg acc aac ggc caa aag gttcgacatt agacctcctg Gly Leu Thr Thr Pro Thr Asn Gly Gln Lys 80 85	413
35	cctgcgctcc cttcttaacc ctacacctgc tttcgccccg ggaaggccgc ctcgttgcta	473
40	accagttctc ggctcattct ag tgg ctg ttt act gag ccc gag cgt ggc gtg Trp Leu Phe Thr Glu Pro Glu Arg Gly Val 90 95	525
45	ttc aac ttc acc gag ggt gac atc gtg acg aac ctg gcc cgg aag cac Phe Asn Phe Thr Glu Gly Asp Ile Val Thr Asn Leu Ala Arg Lys His 100 105 110	573
50	ggt ttc atg cag cgg tgc cac gcg ctc gtc tgg cac agc cag ctc gcc Gly Phe Met Gln Arg Cys His Ala Leu Val Trp His Ser Gln Leu Ala 115 120 125	621
55	cct tgg gtc gag tcg acc gag tgg acg ccc gag gag ctg cgc cag gtc Pro Trp Val Glu Ser Thr Glu Trp Thr Pro Glu Glu Leu Arg Gln Val 130 135 140	669
60	att gtc aac cac atc acc cac gtg gcc ggc tac tac aag ggc aag tgc Ile Val Asn His Ile Thr His Val Ala Gly Tyr Tyr Lys Gly Lys Cys 145 150 155 160	717
65	tat gcc tgg gac gtc gtc aac gag gcc ctg aac gag gac ggc acc tac Tyr Ala Trp Asp Val Val Asn Glu Ala Leu Asn Glu Asp Gly Thr Tyr 165 170 175	765
70	cgc gag tcc gtc ttc tac aag gtg ctc ggc gag gac tac atc aag ctg Arg Glu Ser Val Phe Tyr Lys Val Leu Gly Glu Asp Tyr Ile Lys Leu 180 185 190	813
75	gcc ttc gag acg gcc gcc aag gtc gac ccc cac gcc aag ctc tac tac	861

EP 2 183 363 B1

	Ala	Phe	Glu	Thr	Ala	Ala	Lys	Val	Asp	Pro	His	Ala	Lys	Leu	Tyr	Tyr	
			195					200					205				
5	aac	gac	tac	aac	ctc	gag	tcc	ccc	agc	gcc	aag	acc	gag	ggc	gcc	aag	909
	Asn	Asp	Tyr	Asn	Leu	Glu	Ser	Pro	Ser	Ala	Lys	Thr	Glu	Gly	Ala	Lys	
		210					215					220					
	cgc	atc	gtc	aag	atg	ctc	aag	gac	gcc	ggc	atc	cgc	atc	gac	ggc	gtc	957
10	Arg	Ile	Val	Lys	Met	Leu	Lys	Asp	Ala	Gly	Ile	Arg	Ile	Asp	Gly	Val	
	225					230					235					240	
	ggc	ctg	cag	gcc	cac	ctc	gtc	gcc	gag	agc	cac	ccg	acc	ctc	gac	gag	1005
	Gly	Leu	Gln	Ala	His	Leu	Val	Ala	Glu	Ser	His	Pro	Thr	Leu	Asp	Glu	
15					245					250					255		
	cac	atc	gat	gcc	atc	aag	ggc	ttc	acc	gag	ctc	ggc	gtc	gag	gtc	gcc	1053
	His	Ile	Asp	Ala	Ile	Lys	Gly	Phe	Thr	Glu	Leu	Gly	Val	Glu	Val	Ala	
				260					265					270			
20	ctg	acc	gag	ctc	gac	atc	cgc	ctc	tcc	atc	ccg	gcc	aac	gcc	acc	aac	1101
	Leu	Thr	Glu	Leu	Asp	Ile	Arg	Leu	Ser	Ile	Pro	Ala	Asn	Ala	Thr	Asn	
			275					280					285				
	ctc	gcc	cag	cag	agg	gag	gcg	tac	aag	aac	gtgagttttt	tttattttat					1151
25	Leu	Ala	Gln	Gln	Arg	Glu	Ala	Tyr	Lys	Asn							
		290					295										
	ttttatcttc	cctcttggtc	ttccatccaa	ccccccccc	ttttgacggt	gcccagtcag											1211
	attgctcttg	atcgatgcta	acctttgttc	cttctctttg	atag	gtc	gtc	ggc	gct								1267
30						Val	Val	Gly	Ala								
							300										
	tgc	gtc	cag	gtt	cgc	ggc	tgc	att	ggc	gtg	gag	atc	tgg	gac	ttc	tat	1315
35	Cys	Val	Gln	Val	Arg	Gly	Cys	Ile	Gly	Val	Glu	Ile	Trp	Asp	Phe	Tyr	
			305					310					315				
	gac	ccc	ttc	agc	tgg	gtc	cct	gcc	acc	ttc	ccc	ggc	cag	ggc	gcc	ccc	1363
	Asp	Pro	Phe	Ser	Trp	Val	Pro	Ala	Thr	Phe	Pro	Gly	Gln	Gly	Ala	Pro	
40			320				325					330					
	ctg	ctc	tgg	ttc	gag	gac	ttt	tcc	aag	cac	ccc	gcc	tac	gac	ggc	gtc	1411
	Leu	Leu	Trp	Phe	Glu	Asp	Phe	Ser	Lys	His	Pro	Ala	Tyr	Asp	Gly	Val	
	335					340					345					350	
45	gtc	gag	gcc	ctg	acc	aac	agg	acc	acg	ggc	ggg	tgc	aag	ggc	aag	ggc	1459
	Val	Glu	Ala	Leu	Thr	Asn	Arg	Thr	Thr	Gly	Gly	Cys	Lys	Gly	Lys	Gly	
					355					360					365		
	aag	ggc	aag	ggc	aag	gtt	tgg	aag	gcc	taataagtgg	tgttggtgcgg						1506
50	Lys	Gly	Lys	Gly	Lys	Val	Trp	Lys	Ala								
				370					375								
	gtcgtgtaga	gatttaggga	gaggattggg	tccttcggtt	cttgtatcta	attattattc											1566
	cgtactatta	attgcatttc	gatctaaagt	tagtggtgag	cgggtcgccg	cagtcatgcg											1626
55																	
	acatgcatgg	cacgctaagc	tcggccgagc	tctcggga													1664

<210> 35

EP 2 183 363 B1

<211> 375

<212> PRT

<213> Chrysosporium lucknowense

5 <400> 35

	Met	His	Leu	Ser	Ser	Ser	Leu	Leu	Leu	Leu	Ala	Ala	Leu	Pro	Leu	Gly	
	1				5					10					15		
10	Ile	Ala	Gly	Lys	Gly	Lys	Gly	His	Gly	His	Gly	Pro	His	Thr	Gly	Leu	
				20					25					30			
15	His	Thr	Leu	Ala	Lys	Gln	Ala	Gly	Leu	Lys	Tyr	Phe	Gly	Ser	Ala	Thr	
			35					40					45				
20	Asp	Ser	Pro	Gly	Gln	Arg	Glu	Arg	Ala	Gly	Tyr	Glu	Asp	Lys	Tyr	Ala	
		50					55					60					
25	Gln	Tyr	Asp	Gln	Ile	Met	Trp	Lys	Ser	Gly	Glu	Phe	Gly	Leu	Thr	Thr	
	65					70					75					80	
30	Pro	Thr	Asn	Gly	Gln	Lys	Trp	Leu	Phe	Thr	Glu	Pro	Glu	Arg	Gly	Val	
					85					90					95		
35	Phe	Asn	Phe	Thr	Glu	Gly	Asp	Ile	Val	Thr	Asn	Leu	Ala	Arg	Lys	His	
				100					105					110			
40	Gly	Phe	Met	Gln	Arg	Cys	His	Ala	Leu	Val	Trp	His	Ser	Gln	Leu	Ala	
			115					120					125				
45	Pro	Trp	Val	Glu	Ser	Thr	Glu	Trp	Thr	Pro	Glu	Glu	Leu	Arg	Gln	Val	
		130					135					140					
50	Ile	Val	Asn	His	Ile	Thr	His	Val	Ala	Gly	Tyr	Tyr	Lys	Gly	Lys	Cys	
	145					150					155					160	
55	Tyr	Ala	Trp	Asp	Val	Val	Asn	Glu	Ala	Leu	Asn	Glu	Asp	Gly	Thr	Tyr	
					165					170					175		
60	Arg	Glu	Ser	Val	Phe	Tyr	Lys	Val	Leu	Gly	Glu	Asp	Tyr	Ile	Lys	Leu	
				180					185					190			
65	Ala	Phe	Glu	Thr	Ala	Ala	Lys	Val	Asp	Pro	His	Ala	Lys	Leu	Tyr	Tyr	
			195					200					205				
70	Asn	Asp	Tyr	Asn	Leu	Glu	Ser	Pro	Ser	Ala	Lys	Thr	Glu	Gly	Ala	Lys	
		210					215					220					

EP 2 183 363 B1

	Arg	Ile	Val	Lys	Met	Leu	Lys	Asp	Ala	Gly	Ile	Arg	Ile	Asp	Gly	Val	
	225					230					235					240	
5	Gly	Leu	Gln	Ala	His	Leu	Val	Ala	Glu	Ser	His	Pro	Thr	Leu	Asp	Glu	
					245					250					255		
10	His	Ile	Asp	Ala	Ile	Lys	Gly	Phe	Thr	Glu	Leu	Gly	Val	Glu	Val	Ala	
				260					265					270			
15	Leu	Thr	Glu	Leu	Asp	Ile	Arg	Leu	Ser	Ile	Pro	Ala	Asn	Ala	Thr	Asn	
			275				280						285				
20	Leu	Ala	Gln	Gln	Arg	Glu	Ala	Tyr	Lys	Asn	Val	Val	Gly	Ala	Cys	Val	
	290						295					300					
25	Gln	Val	Arg	Gly	Cys	Ile	Gly	Val	Glu	Ile	Trp	Asp	Phe	Tyr	Asp	Pro	
	305					310					315					320	
30	Phe	Ser	Trp	Val	Pro	Ala	Thr	Phe	Pro	Gly	Gln	Gly	Ala	Pro	Leu	Leu	
				325						330					335		
35	Trp	Phe	Glu	Asp	Phe	Ser	Lys	His	Pro	Ala	Tyr	Asp	Gly	Val	Val	Glu	
			340					345					350				
40	Ala	Leu	Thr	Asn	Arg	Thr	Thr	Gly	Gly	Cys	Lys	Gly	Lys	Gly	Lys	Gly	
		355						360					365				
45	Lys	Gly	Lys	Val	Trp	Lys	Ala										
	370					375											

<210> 36
 <211> 1125
 <212> DNA
 <213> Chrysosporium lucknowense

45	atgcatctct	cctcgtctct	cctcctctc	gccgccttgc	ccctgggcat	cgccggcaag	60
	ggcaagggcc	acggccacgg	cccccatacc	gggctccaca	ccctcgccaa	gcaggccggc	120
50	ctcaagtact	tcggctctgc	caccgactct	cccggccagc	gtgagcgcgc	cggctacgag	180
	gacaagtacg	cccagtacga	ccagatcatg	tggaagtcgg	gcgagttcgg	cctgacgacc	240
55	ccgaccaacg	gccaaaagtg	gctgtttact	gagcccagac	gtggcgtgtt	caacttcacc	300
	gagggtgaca	tcgtgacgaa	cctggcccgg	aagcacgggt	tcatgcagcg	gtgccacgcg	360
	ctcgtctggc	acagccagct	cgccccttgg	gtcgagtcga	ccgagtggac	gcccgaggag	420

EP 2 183 363 B1

	ctgcgccagg tcattgtcaa ccacatcacc cacgtggccg gctactacaa gggcaagtgc	480
	tatgcctggg acgtcgtcaa cgaggccctg aacgaggacg gcacctaccg cgagtcgctc	540
5	ttctacaagg tgctcggcga ggactacatc aagctggcct tcgagacggc cgccaaggtc	600
	gacccccacg ccaagctcta ctacaacgac tacaacctcg agtccccag cgccaagacc	660
10	gagggcgcca agcgcacgtg caagatgctc aaggacgccg gcatccgcat cgacggcgctc	720
	ggcctgcagg cccacctcgt cgccgagagc caccgaccc tcgacgagca catcgatgcc	780
	atcaagggct tcaccgagct cggcgctcgag gtcgccctga ccgagctcga catccgcctc	840
15	tccatcccgg ccaacgccac caacctcgcc cagcagaggg aggcgtacaa gaacgtcgctc	900
	ggcgcttgcg tccagggttcg cggctgcatt ggcgtggaga tctgggactt ctatgacccc	960
20	ttcagctggg tccctgccac cttccccggc cagggcgccc cctgctctg gttcgaggac	1020
	ttttccaagc accccgccta cgacggcgctc gtcgaggccc tgaccaacag gaccacgggc	1080
	gggtgcaagg gcaagggcaa gggcaagggc aaggtttgga aggcc	1125
25	<210> 37 <211> 1192 <212> DNA <213> Chrysosporium lucknowense	
30	<220> <221> CDS <222> (121)..(386)	
35	<220> <221> CDS <222> (531)..(942)	
	<400> 37	
40		
45		
50		
55		

EP 2 183 363 B1

	acataaatag agccactggc tcccctcaag ggattggcgt tcgaagcatc gtcagaaccg	60
	caatatacat attagtgtaa tatcttgcac ctcatatgat ccacaatcat catcagcatc	120
5	atg gtt acc ctc act cgc ctg gcg gtc gcc gcg gcg gcc atg atc tcc Met Val Thr Leu Thr Arg Leu Ala Val Ala Ala Ala Ala Met Ile Ser 1 5 10 15	168
10	agc act ggc ctg gct gcc ccg acg ccc gaa gct ggc ccc gac ctt ccc Ser Thr Gly Leu Ala Ala Pro Thr Pro Glu Ala Gly Pro Asp Leu Pro 20 25 30	216
15	gac ttt gag ctc ggg gtc aac aac ctc gcc cgc cgc gcg ctg gac tac Asp Phe Glu Leu Gly Val Asn Asn Leu Ala Arg Arg Ala Leu Asp Tyr 35 40 45	264
20	aac cag aac tac agg acc agc ggc aac gtc aac tac tcg ccc acc gac Asn Gln Asn Tyr Arg Thr Ser Gly Asn Val Asn Tyr Ser Pro Thr Asp 50 55 60	312
	aac ggc tac tcg gtc agc ttc tcc aac gcg gga gat ttt gtc gtc ggg	360
25		
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	Asn Gly Tyr Ser Val Ser Phe Ser Asn Ala Gly Asp Phe Val Val Gly	
	65 70 75 80	
5	aag ggc tgg agg acg gga gcc acc ag gtaagacagc agcgtacaat Lys Gly Trp Arg Thr Gly Ala Thr Arg	406
	85	
	cgtcagggtt attggttata tggttatcca gacttgagcc gtggatttct tttatttcat	466
10	ttttcctttc ctccactctt aaacacgctt cgggcagcgt tgacaaacct tctaatatct	526
	tcag a aac atc acc ttc tcg gga tcg aca cag cat acc tcg ggc acc	573
	Asn Ile Thr Phe Ser Gly Ser Thr Gln His Thr Ser Gly Thr	
	90 95 100	
15	gtg ctg gtc tcc gtc tac ggc tgg acc cgg aac ccg ctg atc gag tac	621
	Val Leu Val Ser Val Tyr Gly Trp Thr Arg Asn Pro Leu Ile Glu Tyr	
	105 110 115	
20	tac gtg cag gag tac acg tcc aac ggg gcc ggc tcc gct cag ggc gag	669
	Tyr Val Gln Glu Tyr Thr Ser Asn Gly Ala Gly Ser Ala Gln Gly Glu	
	120 125 130 135	
	aag ctg ggc acg gtc gag agc gac ggg ggc acg tac gag atc tgg cgg	717
25	Lys Leu Gly Thr Val Glu Ser Asp Gly Gly Thr Tyr Glu Ile Trp Arg	
	140 145 150	
	cac cag cag gtc aac cag ccg tcg atc gag ggc acc tcg acc ttc tgg	765
	His Gln Gln Val Asn Gln Pro Ser Ile Glu Gly Thr Ser Thr Phe Trp	
	155 160 165	
30	cag tac atc tcg aac cgc gtg tcc ggc cag cgg ccc aac ggc ggc acc	813
	Gln Tyr Ile Ser Asn Arg Val Ser Gly Gln Arg Pro Asn Gly Gly Thr	
	170 175 180	
35	gtc acc ctc gcc aac cac ttc gcc gcc tgg cag aag ctc ggc ctg aac	861
	Val Thr Leu Ala Asn His Phe Ala Ala Trp Gln Lys Leu Gly Leu Asn	
	185 190 195	
	ctg ggc cag cac gac tac cag gtc ctg gcc acc gag ggc tgg ggc aac	909
40	Leu Gly Gln His Asp Tyr Gln Val Leu Ala Thr Glu Gly Trp Gly Asn	
	200 205 210 215	
	gcc ggc ggc agc tcc cag tac acc gtc agc ggc tgaccgttgg gtggtggggg	962
45	Ala Gly Gly Ser Ser Gln Tyr Thr Val Ser Gly	
	220 225	
	gcaaagagct ggtcttagaa ccatccaacg atccttacca tgaaagagct tgtgacttag	1022
	tcgctgttca tgaaaacatg tctgttttac acagacaaga tttaccaatt gcaatgaagc	1082
50	atacgtcaac tcgaacgtgt ttgtgtttct cgtgcctctc gtatgaatat gctatgctga	1142
	tggagtatca ctcttgatgt caaacatgct atattaacga cataactata	1192
55	<210> 38 <211> 226 <212> PRT <213> Chrysosporium lucknowense	

EP 2 183 363 B1

<400> 38

5	Met	Val	Thr	Leu	Thr	Arg	Leu	Ala	Val	Ala	Ala	Ala	Ala	Met	Ile	Ser	1	5	10	15
10	Ser	Thr	Gly	Leu	Ala	Ala	Pro	Thr	Pro	Glu	Ala	Gly	Pro	Asp	Leu	Pro	20	25	30	
15	Asp	Phe	Glu	Leu	Gly	Val	Asn	Asn	Leu	Ala	Arg	Arg	Ala	Leu	Asp	Tyr	35	40	45	
20	Asn	Gln	Asn	Tyr	Arg	Thr	Ser	Gly	Asn	Val	Asn	Tyr	Ser	Pro	Thr	Asp	50	55	60	
25	Asn	Gly	Tyr	Ser	Val	Ser	Phe	Ser	Asn	Ala	Gly	Asp	Phe	Val	Val	Gly	65	70	75	80
30	Lys	Gly	Trp	Arg	Thr	Gly	Ala	Thr	Arg	Asn	Ile	Thr	Phe	Ser	Gly	Ser	85	90	95	
35	Thr	Gln	His	Thr	Ser	Gly	Thr	Val	Leu	Val	Ser	Val	Tyr	Gly	Trp	Thr	100	105	110	
40	Arg	Asn	Pro	Leu	Ile	Glu	Tyr	Tyr	Val	Gln	Glu	Tyr	Thr	Ser	Asn	Gly	115	120	125	
45	Ala	Gly	Ser	Ala	Gln	Gly	Glu	Lys	Leu	Gly	Thr	Val	Glu	Ser	Asp	Gly	130	135	140	
50	Gly	Thr	Tyr	Glu	Ile	Trp	Arg	His	Gln	Gln	Val	Asn	Gln	Pro	Ser	Ile	145	150	155	160
55	Glu	Gly	Thr	Ser	Thr	Phe	Trp	Gln	Tyr	Ile	Ser	Asn	Arg	Val	Ser	Gly	165	170	175	
60	Gln	Arg	Pro	Asn	Gly	Gly	Thr	Val	Thr	Leu	Ala	Asn	His	Phe	Ala	Ala	180	185	190	
65	Trp	Gln	Lys	Leu	Gly	Leu	Asn	Leu	Gly	Gln	His	Asp	Tyr	Gln	Val	Leu	195	200	205	
70	Ala	Thr	Glu	Gly	Trp	Gly	Asn	Ala	Gly	Gly	Ser	Ser	Gln	Tyr	Thr	Val	210	215	220	
75	Ser	Gly															225			

EP 2 183 363 B1

<210> 39
 <211> 678
 <212> DNA
 <213> Chrysosporium lucknowense

5
 <400> 39

 atggttacct tcaactcgct gccggtcgcc gccggcgcca tgatctccag cactggcctg 60
 10 gctgccccga cgcccgaaac tggccccgac cttcccgact ttgagctcgg ggtcaacaac 120
 ctgccccgcc gcgcgctgga ctacaaccag aactacagga ccagcggcaa cgtcaactac 180
 15 tcgcccaccg acaacggcta ctcggtcagc ttctccaacg cgggagattt tgcgtcggg 240
 aagggctgga ggacgggagc caccagaaac atcaccttct cgggatcgac acagcatacc 300
 tcgggcaccg tgctcgtctc cgtctacggc tggaccgga acccgctgat cgagtactac 360
 20 gtgcaggagt acacgtccaa cggggccggc tccgctcagg gcgagaagct gggcacggtc 420
 gagagcgacg ggggcacgta cgagatctgg cggcaccagc aggtcaacca gccgtcgatc 480
 gagggcacct cgaccttctg gcagtacatc tcgaaccgcg tgtccggcca gcggcccaac 540
 25 gggggcaccg tcacctctgc caaccacttc gccgcctggc agaagctcgg cctgaacctg 600
 ggccagcacg actaccaggt cctggccacc gagggctggg gcaacgccgg cggcagctcc 660
 30 cagtacaccg tcagcggc 678

<210> 40
 <211> 1194
 <212> DNA
 <213> Chrysosporium lucknowense

35
 <220>
 <221> CDS
 <222> (205)..(485)

40
 <220>
 <221> CDS
 <222> (584)..(986)

45
 <400> 40

50

55

EP 2 183 363 B1

	gtataaatgc atcaaggagc gactgcccc ccataataa ccacccggtt gcttgagtct	60
	ctcgcaactcg cggccccttc ttctctgctt cgcacgcac tcgctgtctc gctgtctcgc	120
5	tgtctcactg tctcgctgtc tcaactgtctc gctgtctcac tgtctcgctg tctcactgtc	180
	tcaactcgtcc atcagagcaa aacc atg gtc tcg ctc aag tcc ctc ctc ctc	231
	Met Val Ser Leu Lys Ser Leu Leu Leu	
	1 5	
10		
	gcc gcg gcg gcg acg ttg acg gcg gtg acg gcg cgc ccg ttc gac ttt	279
	Ala Ala Ala Ala Thr Leu Thr Ala Val Thr Ala Arg Pro Phe Asp Phe	
	10 15 20 25	
15		
20		
25		
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	gac gac ggc aac tcg acc gag gcg ctg gcc aag cgc cag gtc acg ccc Asp Asp Gly Asn Ser Thr Glu Ala Leu Ala Lys Arg Gln Val Thr Pro 30 35 40	327
5	aac gcg cag ggc tac cac tcg ggc tac ttc tac tcg tgg tgg tcc gac Asn Ala Gln Gly Tyr His Ser Gly Tyr Phe Tyr Ser Trp Trp Ser Asp 45 50 55	375
10	ggc ggc ggc cag gcc acc ttc acc ctg ctc gag ggc agc cac tac cag Gly Gly Gly Gln Ala Thr Phe Thr Leu Leu Glu Gly Ser His Tyr Gln 60 65 70	423
15	gtc aac tgg agg aac acg ggc aac ttt gtc ggt ggc aag ggc tgg aac Val Asn Trp Arg Asn Thr Gly Asn Phe Val Gly Gly Lys Gly Trp Asn 75 80 85	471
20	ccg ggt acc ggc cg gtaagttgtg tttcttttct tgaactattg actactaatt Pro Gly Thr Gly Arg 90	525
25	atcgtatctt tccagtcaac gggggtgaaa ggttggttgg ctgactgccg gatcgag g acc atc aac tac ggc ggc tcg ttc aac ccg agc ggc aac ggc tac ctg Thr Ile Asn Tyr Gly Gly Ser Phe Asn Pro Ser Gly Asn Gly Tyr Leu 95 100 105 110	584
30	gcc gtc tac ggc tgg acg cac aac ccg ctg atc gag tac tac gtg gtc Ala Val Tyr Gly Trp Thr His Asn Pro Leu Ile Glu Tyr Tyr Val Val 115 120 125	632
35	gag tcg tac ggg acc tac aac ccg ggc agc cag gcc cag tac aag ggc Glu Ser Tyr Gly Thr Tyr Asn Pro Gly Ser Gln Ala Gln Tyr Lys Gly 130 135 140	680
40	agc ttc cag agc gac ggc ggc acc tac aac atc tac gtc tcg acc cgc Ser Phe Gln Ser Asp Gly Gly Thr Tyr Asn Ile Tyr Val Ser Thr Arg 145 150 155	728
45	tac aac gcg ccc tcg atc gag ggc acc cgc acc ttc cag cag tac tgg Tyr Asn Ala Pro Ser Ile Glu Gly Thr Arg Thr Phe Gln Gln Tyr Trp 160 165 170	776
50	tcc atc cgc acc tcc aag cgc gtc ggc ggc tcc gtc acc atg cag aac Ser Ile Arg Thr Ser Lys Arg Val Gly Gly Ser Val Thr Met Gln Asn 175 180 185 190	824
55	cac ttc aac gcc tgg gcc cag cac ggc atg ccc ctc ggc tcc cac gac His Phe Asn Ala Trp Ala Gln His Gly Met Pro Leu Gly Ser His Asp 195 200 205	872
60	tac cag atc gtc gcc acc gag ggc tac cag agc agc ggc tcc tcc gac Tyr Gln Ile Val Ala Thr Glu Gly Tyr Gln Ser Ser Gly Ser Ser Asp 210 215 220	920
65	atc tac gtc cag act cac taggtacctt aggtattcaa gcgatccagt Ile Tyr Val Gln Thr His 225	968
70	cggtcagaca gactottcca ggcacgccgc tgtccatcag tgggtagtat tatcccatga tgggcgccgg cggtgacggg aaaagggggc attcgtctgc agtaccttat ttactatta 1076 1136	1016

EP 2 183 363 B1

atactgccta tctgccagca taacaatagg tagcacgcgt gcacttacca ggaattca 1194

<210> 41

5 <211> 228

<212> PRT

<213> Chrysosporium lucknowense

<400> 41

10

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	Met	Val	Ser	Leu	Lys	Ser	Leu	Leu	Leu	Ala	Ala	Ala	Ala	Thr	Leu	Thr
	1				5					10					15	
5	Ala	Val	Thr	Ala	Arg	Pro	Phe	Asp	Phe	Asp	Asp	Gly	Asn	Ser	Thr	Glu
				20				25						30		
10	Ala	Leu	Ala	Lys	Arg	Gln	Val	Thr	Pro	Asn	Ala	Gln	Gly	Tyr	His	Ser
			35					40					45			
15	Gly	Tyr	Phe	Tyr	Ser	Trp	Trp	Ser	Asp	Gly	Gly	Gly	Gln	Ala	Thr	Phe
		50					55					60				
20	Thr	Leu	Leu	Glu	Gly	Ser	His	Tyr	Gln	Val	Asn	Trp	Arg	Asn	Thr	Gly
	65					70					75					80
25	Asn	Phe	Val	Gly	Gly	Lys	Gly	Trp	Asn	Pro	Gly	Thr	Gly	Arg	Thr	Ile
					85					90					95	
30	Asn	Tyr	Gly	Gly	Ser	Phe	Asn	Pro	Ser	Gly	Asn	Gly	Tyr	Leu	Ala	Val
			100						105					110		
35	Tyr	Gly	Trp	Thr	His	Asn	Pro	Leu	Ile	Glu	Tyr	Tyr	Val	Val	Glu	Ser
			115					120					125			
40	Tyr	Gly	Thr	Tyr	Asn	Pro	Gly	Ser	Gln	Ala	Gln	Tyr	Lys	Gly	Ser	Phe
		130					135					140				
45	Gln	Ser	Asp	Gly	Gly	Thr	Tyr	Asn	Ile	Tyr	Val	Ser	Thr	Arg	Tyr	Asn
	145					150					155					160
50	Ala	Pro	Ser	Ile	Glu	Gly	Thr	Arg	Thr	Phe	Gln	Gln	Tyr	Trp	Ser	Ile
				165						170					175	
55	Arg	Thr	Ser	Lys	Arg	Val	Gly	Gly	Ser	Val	Thr	Met	Gln	Asn	His	Phe
				180					185					190		
60	Asn	Ala	Trp	Ala	Gln	His	Gly	Met	Pro	Leu	Gly	Ser	His	Asp	Tyr	Gln
			195					200					205			
65	Ile	Val	Ala	Thr	Glu	Gly	Tyr	Gln	Ser	Ser	Gly	Ser	Ser	Asp	Ile	Tyr
		210					215					220				
70	Val	Gln	Thr	His												
	225															

<210> 42

EP 2 183 363 B1

<211> 684
 <212> DNA
 <213> Chrysosporium lucknowense

5 <400> 42

atggtctcgc tcaagtccct cctcctcgcc gcggcggcga cgttgacggc ggtgacggcg 60

10 cgcccgttcg actttgacga cggcaactcg accgaggcgc tggccaagcg ccaggtcacg 120

cccaacgcgc agggctacca ctcgggctac ttctactcgt ggtggtccga cggcggcggc 180

caggccacct tcacctgct cgagggcagc cactaccagg tcaactggag gaacacgggc 240

15 aactttgtcg gtggcaaggg ctggaacccg ggtaccggcc ggaccatcaa ctacggcggc 300

tcgttcaacc cgagcggcaa cggctacctg gccgtctacg gctggacgca caaccgctg 360

20 atcgagtact acgtggtcga gtcgtacggg acctacaacc cgggcagcca ggcccagtac 420

aagggcagct tccagagcga cggcggcacc tacaacatct acgtctcgac ccgctacaac 480

gcgccctcga tcgagggcac ccgcaccttc cagcagtact ggtccatccg cacctccaag 540

25 cgcgtcggcg gctccgtcac catgcagaac cacttcaacg cctggggcca gcacggcatg 600

ccccctgggt cccacgacta ccagatcgtc gccaccgagg gctaccagag cagcggctcc 660

tccgacatct acgtccagac tcac 684

30 <210> 43
 <211> 2740
 <212> DNA
 <213> Chrysosporium lucknowense

35 <220>
 <221> CDS
 <222> (1)..(230)

40 <220>
 <221> CDS
 <222> (294)..(682)

45 <220>
 <221> CDS
 <222> (744)..(2737)

<400> 43

50 atg caa ctt cca gcc gca gcc caa tgg ctg ctc acg ctt ccc gcg aaa 48

55

EP 2 183 363 B1

	Met	Gln	Leu	Pro	Ala	Ala	Ala	Gln	Trp	Leu	Thr	Leu	Pro	Ala	Lys			
	1				5					10				15				
5	gcc	tca	ctt	gct	gac	aat	cat	cgt	cag	gtt	cac	cag	aag	ccc	ctc	gcg	96	
	Ala	Ser	Leu	Ala	Asp	Asn	His	Arg	Gln	Val	His	Gln	Lys	Pro	Leu	Ala		
				20				25					30					
10	aga	tct	gaa	cct	ttt	tac	ccg	tcg	cca	tgg	atg	aat	ccc	aac	gcc	gac	144	
	Arg	Ser	Glu	Pro	Phe	Tyr	Pro	Ser	Pro	Trp	Met	Asn	Pro	Asn	Ala	Asp		
			35				40					45						
	ggc	tgg	gcg	gag	gcc	tat	gcc	cag	gcc	aag	tcc	ttt	gtc	tcc	caa	atg	192	
	Gly	Trp	Ala	Glu	Ala	Tyr	Ala	Gln	Ala	Lys	Ser	Phe	Val	Ser	Gln	Met		
		50				55					60							
15	act	ctg	cta	gag	aag	gtc	aac	ttg	acc	acg	gga	gtc	gg	gtaagttttg			240	
	Thr	Leu	Leu	Glu	Lys	Val	Asn	Leu	Thr	Thr	Gly	Val	Gly					
	65					70				75								
20	tcattttgtc caggtaacat gcaaatgggt ctgctaacaa taacttaccg tag c tgg																297	
	Trp																	
	ggg	gct	gag	cag	tgc	gtc	ggc	caa	gtg	ggc	gcg	atc	cct	cgc	ctt	gga	345	
	Gly	Ala	Glu	Gln	Cys	Val	Gly	Gln	Val	Gly	Ala	Ile	Pro	Arg	Leu	Gly		
		80				85					90							
25	ctt	cgc	agt	ctg	tgc	atg	cat	gac	tcc	cct	ctc	ggc	atc	cga	gga	gcc	393	
	Leu	Arg	Ser	Leu	Cys	Met	His	Asp	Ser	Pro	Leu	Gly	Ile	Arg	Gly	Ala		
	95					100				105						110		
30	gac	tac	aac	tca	gcg	ttc	ccc	tct	ggc	cag	acc	gtt	gct	gct	acc	tgg	441	
	Asp	Tyr	Asn	Ser	Ala	Phe	Pro	Ser	Gly	Gln	Thr	Val	Ala	Ala	Thr	Trp		
				115					120						125			
	gat	cgc	ggt	ctg	atg	tac	cgt	cgc	ggc	tac	gca	atg	ggc	cag	gag	gcc	489	
35	Asp	Arg	Gly	Leu	Met	Tyr	Arg	Arg	Gly	Tyr	Ala	Met	Gly	Gln	Glu	Ala		
				130				135					140					
	aaa	ggc	aag	ggc	atc	aat	gtc	ctt	ctc	gga	cca	gtc	gcc	ggc	ccc	ctt	537	
	Lys	Gly	Lys	Gly	Ile	Asn	Val	Leu	Leu	Gly	Pro	Val	Ala	Gly	Pro	Leu		
			145				150					155						
	ggc	cgc	atg	ccc	gag	ggc	ggt	cgt	aac	tgg	gaa	ggc	ttc	gct	ccg	gat	585	
	Gly	Arg	Met	Pro	Glu	Gly	Gly	Arg	Asn	Trp	Glu	Gly	Phe	Ala	Pro	Asp		
		160				165					170							
45	ccc	gtc	ctt	acc	ggc	atc	ggc	atg	tcc	gag	acg	atc	aag	ggc	att	cag	633	
	Pro	Val	Leu	Thr	Gly	Ile	Gly	Met	Ser	Glu	Thr	Ile	Lys	Gly	Ile	Gln		
		175				180				185						190		
	gat	gct	ggc	gtc	atc	gct	tgt	gcg	aag	cac	ttt	att	gga	aac	gag	cag	g	682
50	Asp	Ala	Gly	Val	Ile	Ala	Cys	Ala	Lys	His	Phe	Ile	Gly	Asn	Glu	Gln		
				195				200					205					
	gtgagtagtc aaagacgggc cgtctcggac ccgcggcttc aagctgctga ctctgctgca																742	
55	g	ag	cac	ttc	aga	cag	gtg	cca	gaa	gcc	cag	gga	tac	ggt	tac	aac	atc	790
	Glu	His	Phe	Arg	Gln	Val	Pro	Glu	Ala	Gln	Gly	Tyr	Gly	Tyr	Asn	Ile		
				210				215					220					

EP 2 183 363 B1

	agc gaa acc ctc tcc tcc aac att gac gac aag acc atg cac gag ctc	838
	Ser Glu Thr Leu Ser Ser Asn Ile Asp Asp Lys Thr Met His Glu Leu	
	225 230 235	
5	tac ctt tgg ccg ttt gcc gat gcc gtc cgg gcc ggc gtc ggc tct gtc	886
	Tyr Leu Trp Pro Phe Ala Asp Ala Val Arg Ala Gly Val Gly Ser Val	
	240 245 250	
10	atg tgc tcg tac cag cag gtc aac aac tcg tac gcc tgc cag aac tcg	934
	Met Cys Ser Tyr Gln Gln Val Asn Asn Ser Tyr Ala Cys Gln Asn Ser	
	255 260 265 270	
15	aag ctg ctg aac gac ctc ctc aag aac gag ctt ggg ttt cag ggc ttc	982
	Lys Leu Leu Asn Asp Leu Leu Lys Asn Glu Leu Gly Phe Gln Gly Phe	
	275 280 285	
20	gtc atg agc gac tgg cag gca cag cac act ggc gca gca agc gcc gtg	1030
	Val Met Ser Asp Trp Gln Ala Gln His Thr Gly Ala Ala Ser Ala Val	
	290 295 300	
25	gct ggt ctc gat atg tcc atg ccg ggc gac acc cag ttc aac act ggc	1078
	Ala Gly Leu Asp Met Ser Met Pro Gly Asp Thr Gln Phe Asn Thr Gly	
	305 310 315	
30	gtc agt ttc tgg ggc gcc aat ctc acc ctc gcc gtc ctc aac ggc aca	1126
	Val Ser Phe Trp Gly Ala Asn Leu Thr Leu Ala Val Leu Asn Gly Thr	
	320 325 330	
35	gtc cct gcc tac cgt ctc gac gac atg gcc atg cgc atc atg gcc gcc	1174
	Val Pro Ala Tyr Arg Leu Asp Asp Met Ala Met Arg Ile Met Ala Ala	
	335 340 345 350	
40	ctc ttc aag gtc acc aag acc acc cac ctg gaa ccc atc aac ttc tcc	1222
	Leu Phe Lys Val Thr Lys Thr Thr His Leu Glu Pro Ile Asn Phe Ser	
	355 360 365	
45	ttc tgg acc gac gac act tat ggc ccg atc cac tgg gcc gcc aag cat	1270
	Phe Trp Thr Asp Asp Thr Tyr Gly Pro Ile His Trp Ala Ala Lys His	
	370 375 380	
50	ggc tac cag aag att aat tcc cac gtt gac gtc cgc gcc gac cac ggc	1318
	Gly Tyr Gln Lys Ile Asn Ser His Val Asp Val Arg Ala Asp His Gly	
	385 390 395	
55	aac ctc atc cgg gag att gcc gcc aag ggt acg gtg ctg ctg aag aat	1366
	Asn Leu Ile Arg Glu Ile Ala Ala Lys Gly Thr Val Leu Leu Lys Asn	
	400 405 410	
60	acc ggc tct cta ccc ctg aac aag cca aag ttc gtg gcc gtc atc ggc	1414
	Thr Gly Ser Leu Pro Leu Asn Lys Pro Lys Phe Val Ala Val Ile Gly	
	415 420 425 430	
65	gag gat gct ggg tcg agc ccc aac ggg ccc aac ggc tgc agc gac cgc	1462
	Glu Asp Ala Gly Ser Ser Pro Asn Gly Pro Asn Gly Cys Ser Asp Arg	
	435 440 445	
70	ggc tgt aac gaa ggc acg ctc gcc atg ggc tgg gga tcc ggc aca gcc	1510
	Gly Cys Asn Glu Gly Thr Leu Ala Met Gly Trp Gly Ser Gly Thr Ala	
	450 455 460	
75	aac tat ccg tac ctc gtt tcc ccc gac gcc gcg ctc cag gcc cgg gcc	1558

EP 2 183 363 B1

	Asn	Tyr	Pro	Tyr	Leu	Val	Ser	Pro	Asp	Ala	Ala	Leu	Gln	Ala	Arg	Ala	
			465					470					475				
5	atc	cag	gac	ggc	acg	agg	tac	gag	agc	gtc	ctg	tcc	aac	tac	gcc	gag	1606
	Ile	Gln	Asp	Gly	Thr	Arg	Tyr	Glu	Ser	Val	Leu	Ser	Asn	Tyr	Ala	Glu	
		480					485					490					
10	gaa	aag	aca	aag	gct	ctg	gtc	tcg	cag	gcc	aat	gca	acc	gcc	atc	gtc	1654
	Glu	Lys	Thr	Lys	Ala	Leu	Val	Ser	Gln	Ala	Asn	Ala	Thr	Ala	Ile	Val	
	495					500					505					510	
	ttc	gtc	aat	gcc	gac	tca	ggc	gag	ggc	tac	atc	aac	gtg	gac	ggc	aac	1702
	Phe	Val	Asn	Ala	Asp	Ser	Gly	Glu	Gly	Tyr	Ile	Asn	Val	Asp	Gly	Asn	
					515					520					525		
15	gag	ggc	gac	cgt	aag	aac	ctg	act	ctc	tgg	aac	aac	ggc	gat	act	ctg	1750
	Glu	Gly	Asp	Arg	Lys	Asn	Leu	Thr	Leu	Trp	Asn	Asn	Gly	Asp	Thr	Leu	
				530					535					540			
20	gtc	aag	aac	gtc	tcg	agc	tgg	tgc	agc	aac	acc	atc	gtc	gtc	atc	cac	1798
	Val	Lys	Asn	Val	Ser	Ser	Trp	Cys	Ser	Asn	Thr	Ile	Val	Val	Ile	His	
			545				550						555				
25	tcg	gtc	ggc	ccg	gtc	ctc	ctg	acc	gat	tgg	tac	gac	aac	ccc	aac	atc	1846
	Ser	Val	Gly	Pro	Val	Leu	Leu	Thr	Asp	Trp	Tyr	Asp	Asn	Pro	Asn	Ile	
		560				565						570					
	acg	gcc	att	ctc	tgg	gct	ggc	ctt	ccg	ggc	cag	gag	tcg	ggc	aac	tcc	1894
	Thr	Ala	Ile	Leu	Trp	Ala	Gly	Leu	Pro	Gly	Gln	Glu	Ser	Gly	Asn	Ser	
	575					580					585					590	
30	atc	acc	gac	gtg	ctt	tac	ggc	aag	gtc	aac	ccc	gcc	gcc	cgc	tcg	ccc	1942
	Ile	Thr	Asp	Val	Leu	Tyr	Gly	Lys	Val	Asn	Pro	Ala	Ala	Arg	Ser	Pro	
					595					600					605		
35	ttc	act	tgg	ggc	aag	acc	cgc	gaa	agc	tat	ggc	gcg	gac	gtc	ctg	tac	1990
	Phe	Thr	Trp	Gly	Lys	Thr	Arg	Glu	Ser	Tyr	Gly	Ala	Asp	Val	Leu	Tyr	
			610						615					620			
40	aag	ccg	aat	aat	ggc	aat	ggc	gag	ccc	caa	cag	gac	ttc	acc	gag	ggc	2038
	Lys	Pro	Asn	Asn	Gly	Asn	Gly	Ala	Pro	Gln	Gln	Asp	Phe	Thr	Glu	Gly	
			625				630						635				
	gtc	ttc	atc	gac	tac	cgc	tac	ttc	gac	aag	gtt	gac	gat	gac	tcg	gtc	2086
	Val	Phe	Ile	Asp	Tyr	Arg	Tyr	Phe	Asp	Lys	Val	Asp	Asp	Asp	Ser	Val	
		640					645					650					
45	atc	tac	gag	ttc	ggc	cac	ggc	ctg	agc	tac	acc	acc	ttc	gag	tac	agc	2134
	Ile	Tyr	Glu	Phe	Gly	His	Gly	Leu	Ser	Tyr	Thr	Thr	Phe	Glu	Tyr	Ser	
	655					660					665					670	
50	aac	atc	cgc	gtc	gtc	aag	tcc	aac	gtc	agc	gag	tac	cgg	ccc	acg	acg	2182
	Asn	Ile	Arg	Val	Val	Lys	Ser	Asn	Val	Ser	Glu	Tyr	Arg	Pro	Thr	Thr	
					675					680					685		
	ggc	acc	acg	gcc	cag	gcc	ccg	acg	ttt	ggc	aac	ttc	tcc	acc	gac	ctc	2230
	Gly	Thr	Thr	Ala	Gln	Ala	Pro	Thr	Phe	Gly	Asn	Phe	Ser	Thr	Asp	Leu	
				690					695					700			
55	gag	gac	tat	ctc	ttc	ccc	aag	gac	gag	ttc	ccc	tac	atc	tac	cag	tac	2278
	Glu	Asp	Tyr	Leu	Phe	Pro	Lys	Asp	Glu	Phe	Pro	Tyr	Ile	Tyr	Gln	Tyr	

EP 2 183 363 B1

	705	710	715	
5	atc tac ccg tac ctc aac acg acc gac ccc cgg agg gcc tcg gcc gat Ile Tyr Pro Tyr Leu Asn Thr Thr Asp Pro Arg Arg Ala Ser Ala Asp 720 725 730			2326
10	ccc cac tac ggc cag acc gcc gag gag ttc ctc ccg ccc cac gcc acc Pro His Tyr Gly Gln Thr Ala Glu Glu Phe Leu Pro Pro His Ala Thr 735 740 745 750			2374
15	gat gac gac ccc cag ccg ctc ctc cgg tcc tcg ggc gga aac tcc ccc Asp Asp Asp Pro Gln Pro Leu Leu Arg Ser Ser Gly Gly Asn Ser Pro 755 760 765			2422
20	ggc ggc aac cgc cag ctg tac gac att gtc tac aca atc acg gcc gac Gly Gly Asn Arg Gln Leu Tyr Asp Ile Val Tyr Thr Ile Thr Ala Asp 770 775 780			2470
25	atc acg aat acg ggc tcc gtt gta ggc gag gag gta ccg cag ctc tac Ile Thr Asn Thr Gly Ser Val Val Gly Glu Glu Val Pro Gln Leu Tyr 785 790 795			2518
30	gtc tcg ctg ggc ggt ccc gag gat ccc aag gtg cag ctg cgc gac ttt Val Ser Leu Gly Gly Pro Glu Asp Pro Lys Val Gln Leu Arg Asp Phe 800 805 810			2566
35	gac agg atg cgg atc gaa ccc gcc gag acg agg cag ttc acc gcc cgc Asp Arg Met Arg Ile Glu Pro Gly Glu Thr Arg Gln Phe Thr Gly Arg 815 820 825 830			2614
40	ctg acg cgc aga gat ctg agc aac tgg gac gtc acg gtg cag gac tgg Leu Thr Arg Arg Asp Leu Ser Asn Trp Asp Val Thr Val Gln Asp Trp 835 840 845			2662
45	gtc atc agc agg tat ccc aag acg gca tat gtt ggg agg agc agc cgg Val Ile Ser Arg Tyr Pro Lys Thr Ala Tyr Val Gly Arg Ser Ser Arg 850 855 860			2710
50	aag ttg gat ctc aag att gag ctt cct tga Lys Leu Asp Leu Lys Ile Glu Leu Pro 865 870			2740
55	<210> 44 <211> 871 <212> PRT <213> Chrysosporium lucknowense <400> 44			
50	Met Gln Leu Pro Ala Ala Ala Gln Trp Leu Leu Thr Leu Pro Ala Lys 1 5 10 15			
55	Ala Ser Leu Ala Asp Asn His Arg Gln Val His Gln Lys Pro Leu Ala 20 25 30			
	Arg Ser Glu Pro Phe Tyr Pro Ser Pro Trp Met Asn Pro Asn Ala Asp 35 40 45			

EP 2 183 363 B1

	Gly	Trp	Ala	Glu	Ala	Tyr	Ala	Gln	Ala	Lys	Ser	Phe	Val	Ser	Gln	Met
	50						55					60				
5	Thr	Leu	Leu	Glu	Lys	Val	Asn	Leu	Thr	Thr	Gly	Val	Gly	Trp	Gly	Ala
	65					70					75					80
10	Glu	Gln	Cys	Val	Gly	Gln	Val	Gly	Ala	Ile	Pro	Arg	Leu	Gly	Leu	Arg
					85					90					95	
15	Ser	Leu	Cys	Met	His	Asp	Ser	Pro	Leu	Gly	Ile	Arg	Gly	Ala	Asp	Tyr
				100					105					110		
20	Asn	Ser	Ala	Phe	Pro	Ser	Gly	Gln	Thr	Val	Ala	Ala	Thr	Trp	Asp	Arg
			115					120					125			
25	Gly	Leu	Met	Tyr	Arg	Arg	Gly	Tyr	Ala	Met	Gly	Gln	Glu	Ala	Lys	Gly
			130				135					140				
30	Lys	Gly	Ile	Asn	Val	Leu	Leu	Gly	Pro	Val	Ala	Gly	Pro	Leu	Gly	Arg
	145					150					155					160
35	Met	Pro	Glu	Gly	Gly	Arg	Asn	Trp	Glu	Gly	Phe	Ala	Pro	Asp	Pro	Val
					165					170					175	
40	Leu	Thr	Gly	Ile	Gly	Met	Ser	Glu	Thr	Ile	Lys	Gly	Ile	Gln	Asp	Ala
				180					185					190		
45	Gly	Val	Ile	Ala	Cys	Ala	Lys	His	Phe	Ile	Gly	Asn	Glu	Gln	Glu	His
			195					200					205			
50	Phe	Arg	Gln	Val	Pro	Glu	Ala	Gln	Gly	Tyr	Gly	Tyr	Asn	Ile	Ser	Glu
		210					215					220				
55	Thr	Leu	Ser	Ser	Asn	Ile	Asp	Asp	Lys	Thr	Met	His	Glu	Leu	Tyr	Leu
	225					230					235					240
60	Trp	Pro	Phe	Ala	Asp	Ala	Val	Arg	Ala	Gly	Val	Gly	Ser	Val	Met	Cys
				245						250					255	
65	Ser	Tyr	Gln	Gln	Val	Asn	Asn	Ser	Tyr	Ala	Cys	Gln	Asn	Ser	Lys	Leu
				260					265					270		
70	Leu	Asn	Asp	Leu	Leu	Lys	Asn	Glu	Leu	Gly	Phe	Gln	Gly	Phe	Val	Met
			275					280					285			

EP 2 183 363 B1

	Ser	Asp	Trp	Gln	Ala	Gln	His	Thr	Gly	Ala	Ala	Ser	Ala	Val	Ala	Gly	
	290						295					300					
5	Leu	Asp	Met	Ser	Met	Pro	Gly	Asp	Thr	Gln	Phe	Asn	Thr	Gly	Val	Ser	
	305					310					315					320	
	Phe	Trp	Gly	Ala	Asn	Leu	Thr	Leu	Ala	Val	Leu	Asn	Gly	Thr	Val	Pro	
10					325					330					335		
	Ala	Tyr	Arg	Leu	Asp	Asp	Met	Ala	Met	Arg	Ile	Met	Ala	Ala	Leu	Phe	
				340					345					350			
15	Lys	Val	Thr	Lys	Thr	Thr	His	Leu	Glu	Pro	Ile	Asn	Phe	Ser	Phe	Trp	
			355					360					365				
	Thr	Asp	Asp	Thr	Tyr	Gly	Pro	Ile	His	Trp	Ala	Ala	Lys	His	Gly	Tyr	
20		370				375					380						
	Gln	Lys	Ile	Asn	Ser	His	Val	Asp	Val	Arg	Ala	Asp	His	Gly	Asn	Leu	
25	385				390					395						400	
	Ile	Arg	Glu	Ile	Ala	Ala	Lys	Gly	Thr	Val	Leu	Leu	Lys	Asn	Thr	Gly	
				405					410						415		
30	Ser	Leu	Pro	Leu	Asn	Lys	Pro	Lys	Phe	Val	Ala	Val	Ile	Gly	Glu	Asp	
				420					425					430			
	Ala	Gly	Ser	Ser	Pro	Asn	Gly	Pro	Asn	Gly	Cys	Ser	Asp	Arg	Gly	Cys	
35			435					440					445				
	Asn	Glu	Gly	Thr	Leu	Ala	Met	Gly	Trp	Gly	Ser	Gly	Thr	Ala	Asn	Tyr	
40		450				455						460					
	Pro	Tyr	Leu	Val	Ser	Pro	Asp	Ala	Ala	Leu	Gln	Ala	Arg	Ala	Ile	Gln	
	465				470						475					480	
45	Asp	Gly	Thr	Arg	Tyr	Glu	Ser	Val	Leu	Ser	Asn	Tyr	Ala	Glu	Glu	Lys	
				485					490					495			
	Thr	Lys	Ala	Leu	Val	Ser	Gln	Ala	Asn	Ala	Thr	Ala	Ile	Val	Phe	Val	
50				500					505					510			
	Asn	Ala	Asp	Ser	Gly	Glu	Gly	Tyr	Ile	Asn	Val	Asp	Gly	Asn	Glu	Gly	
		515				520						525					
55	Asp	Arg	Lys	Asn	Leu	Thr	Leu	Trp	Asn	Asn	Gly	Asp	Thr	Leu	Val	Lys	

EP 2 183 363 B1

	530					535						540					
5	Asn	Val	Ser	Ser	Trp	Cys	Ser	Asn	Thr	Ile	Val	Val	Ile	His	Ser	Val	
	545					550					555					560	
	Gly	Pro	Val	Leu	Leu	Thr	Asp	Trp	Tyr	Asp	Asn	Pro	Asn	Ile	Thr	Ala	
10					565					570					575		
	Ile	Leu	Trp	Ala	Gly	Leu	Pro	Gly	Gln	Glu	Ser	Gly	Asn	Ser	Ile	Thr	
				580					585					590			
15																	
	Asp	Val	Leu	Tyr	Gly	Lys	Val	Asn	Pro	Ala	Ala	Arg	Ser	Pro	Phe	Thr	
			595					600					605				
	Trp	Gly	Lys	Thr	Arg	Glu	Ser	Tyr	Gly	Ala	Asp	Val	Leu	Tyr	Lys	Pro	
20		610					615					620					
	Asn	Asn	Gly	Asn	Gly	Ala	Pro	Gln	Gln	Asp	Phe	Thr	Glu	Gly	Val	Phe	
	625					630					635					640	
25																	
	Ile	Asp	Tyr	Arg	Tyr	Phe	Asp	Lys	Val	Asp	Asp	Asp	Ser	Val	Ile	Tyr	
					645					650					655		
30																	
	Glu	Phe	Gly	His	Gly	Leu	Ser	Tyr	Thr	Thr	Phe	Glu	Tyr	Ser	Asn	Ile	
				660					665					670			
	Arg	Val	Val	Lys	Ser	Asn	Val	Ser	Glu	Tyr	Arg	Pro	Thr	Thr	Gly	Thr	
35			675					680					685				
	Thr	Ala	Gln	Ala	Pro	Thr	Phe	Gly	Asn	Phe	Ser	Thr	Asp	Leu	Glu	Asp	
		690					695					700					
40																	
	Tyr	Leu	Phe	Pro	Lys	Asp	Glu	Phe	Pro	Tyr	Ile	Tyr	Gln	Tyr	Ile	Tyr	
	705					710					715					720	
45																	
	Pro	Tyr	Leu	Asn	Thr	Thr	Asp	Pro	Arg	Arg	Ala	Ser	Ala	Asp	Pro	His	
				725						730					735		
	Tyr	Gly	Gln	Thr	Ala	Glu	Glu	Phe	Leu	Pro	Pro	His	Ala	Thr	Asp	Asp	
50				740					745					750			
	Asp	Pro	Gln	Pro	Leu	Leu	Arg	Ser	Ser	Gly	Gly	Asn	Ser	Pro	Gly	Gly	
			755					760					765				
55																	
	Asn	Arg	Gln	Leu	Tyr	Asp	Ile	Val	Tyr	Thr	Ile	Thr	Ala	Asp	Ile	Thr	
		770					775					780					

EP 2 183 363 B1

	Asn Thr Gly Ser Val Val Gly Glu Glu Val Pro Gln Leu Tyr Val Ser	
	785	790 795 800
5	Leu Gly Gly Pro Glu Asp Pro Lys Val Gln Leu Arg Asp Phe Asp Arg	
		805 810 815
10	Met Arg Ile Glu Pro Gly Glu Thr Arg Gln Phe Thr Gly Arg Leu Thr	
		820 825 830
15	Arg Arg Asp Leu Ser Asn Trp Asp Val Thr Val Gln Asp Trp Val Ile	
		835 840 845
20	Ser Arg Tyr Pro Lys Thr Ala Tyr Val Gly Arg Ser Ser Arg Lys Leu	
		850 855 860
25	Asp Leu Lys Ile Glu Leu Pro	
		865 870
30	<210> 45	
	<211> 2613	
	<212> DNA	
	<213> Chrysosporium lucknowense	
35	<400> 45	
40		
45		
50		
55		

EP 2 183 363 B1

	atgcaacttc cagccgcagc ccaatggctg ctcacgcttc ccgcgaaagc ctcaacttgct	60
	gacaatcatc gtcagggttca ccagaagccc ctcgcgagat ctgaaccttt ttaccgcgctg	120
5	ccatggatga atcccaacgc cgacggctgg gcggaggcct atgccaggc caagtccttt	180
	gtctcccaa tgactctgct agagaaggtc aacttgacca cgggagtcgg ctggggggct	240
10	gagcagtgcg tcggccaagt gggcgcgatc cctcgccttg gacttcgcag tctgtgcatg	300
	catgactccc ctctcggcat ccgaggagcc gactacaact cagcgttccc ctctggccag	360
	accgttgctg ctacctggga tcgcggtctg atgtaccgtc gcggctacgc aatgggccag	420
15	gaggccaaag gcaagggcat caatgtcctt ctcggaccag tcgccggccc ccttgggcgc	480
	atgcccagag gcggtcgtaa ctgggaaggc ttcgctccgg atcccgtcct taccggcatc	540
20	ggcatgtccg agacgatcaa gggcattcag gatgctggcg tcatcgcttg tgcgaagcac	600
	tttattggaa acgagcagga gcacttcaga cagggtccag aagcccaggg atacggttac	660
	aacatcagcg aaacctctc ctccaacatt gacgacaaga ccatgcacga gctctacctt	720
25	tggccgtttg ccgatgccgt ccggggccggc gtcggctctg tcatgtgctc gtaccagcag	780
	gtcaacaact cgtacgcctg ccagaactcg aagctgctga acgacctcct caagaacgag	840
30	cttgggtttc agggcttcgt catgagcgac tggcaggcac agcacactgg cgcagcaagc	900
35		
40		
45		
50		
55		

EP 2 183 363 B1

gccgtggctg gtctcgatat gtccatgccg ggcgacaccc agttcaacac tggcgtcagt 960

ttctggggcg ccaatctcac cctcgccgtc ctcaacggca cagtccttgc ctaccgtctc 1020

5 gacgacatgg ccatgcgcat catggccgcc ctcttcaagg tcaccaagac caccacactg 1080

gaacccatca acttctcctt ctggaccgac gacacttatg gcccgatcca ctgggcccgc 1140

aagcatggct accagaagat taattccac gttagcgtcc gcgccgacca cggcaacctc 1200

10 atccgggaga ttgccgcaa gggtaggtg ctgctgaaga ataccggctc tctaccctg 1260

aacaagccaa agttcgtggc cgtcatcggc gaggatgctg ggtcgagccc caacgggccc 1320

15 aacggctgca gcgaccgcg ctgtaacgaa ggcacgtcg ccattgggctg gggatccggc 1380

acagccaact atccgtacct cgtttcccc gacgccgcgc tccaggcccg ggccatccag 1440

gacggcacga ggtacgagag cgtcctgtcc aactacgccg aggaaaagac aaaggctctg 1500

20 gtctcgagcagg ccaatgcaac cgccatcgtc ttctgtcaatg ccgactcagg cgagggttac 1560

atcaacgtgg acggtaacga gggcgaccgt aagaacctga ctctctggaa caacggtgat 1620

actctggtca agaactctc gagctggtgc agcaacacca tcgtcgtcat cactcggtc 1680

25 ggcccgggtcc tctgaccga ttggtacgac aacccaaca tcacggccat tctctgggt 1740

ggctctcccg gccaggagtc gggcaactcc atcaccgacg tgctttacgg caaggtaaac 1800

30 cccgccgcc gctcgccctt cacttggggc aagaccgcg aaagctatgg cgcgacgtc 1860

ctgtacaagc cgaataatgg caatggtgcg cccaacagg acttcaccga gggcgtcttc 1920

atcgactacc gctacttca caaggttgac gatgactcgg tcatctacga gttcgccac 1980

35 ggcctgagct acaccacctt cgagtacagc aacatccgcg tcgtcaagtc caacgtcagc 2040

gagtaccggc ccacgacggg caccacggcc caggccccga cgtttggcaa cttctccacc 2100

40 gacctcgagg actatctctt cccaaggac gagttccctt acatctacca gtacatctac 2160

ccgtacctca acacgaccga ccccgagg gcctcggccg atccccacta cggccagacc 2220

gccgaggagt tctcccgcc ccacgccacc gatgacgacc ccagccgct cctccggtcc 2280

45 tcgggcgga actcccccg cggaaccgc cagctgtacg acattgtcta cacaatcacg 2340

gccgacatca cgaatacggg ctccgttgta ggcgaggagg taccgcagct ctacgtctcg 2400

ctgggcggtc ccgaggatcc caaggtgcag ctgcgcgact ttgacaggat gcggatcgaa 2460

50 cccggcgaga cgaggcagtt caccggccgc ctgacgcga gagatctgag caactgggac 2520

gtcacggtgc aggactgggt catcagcagg tatcccaaga cggcatatgt tgggaggagc 2580

55 agccggaagt tggatctcaa gattgagctt cct 2613

<210> 46

<211> 1226

<212> DNA
 <213> Chrysosporium lucknowense

5 <220>
 <221> CDS
 <222> (89) .. (690)

10 <220>
 <221> CDS
 <222> (776)..(1223)

<400> 46

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	agacagcttg agagacctcc taatcctaga tctcgaggct caacctgagg agagacggtc	60
	tgtaattacc cgctgcacga caccaaag atg atg ctc aca cgc ttc gtg gct	112
5	Met Met Leu Thr Arg Phe Val Ala	
	1 5	
	ggc ctg ctc ggc atc tcc gcc gcg gat gcc gcc ctc acc tac aga ggc	160
	Gly Leu Leu Gly Ile Ser Ala Ala Asp Ala Ala Leu Thr Tyr Arg Gly	
10	10 15 20	
	gtg gat tgg tcc tca gtg gtg gtt gag gaa cgg gcc ggc gtc tcg tac	208
	Val Asp Trp Ser Ser Val Val Val Glu Glu Arg Ala Gly Val Ser Tyr	
	25 30 35 40	
15	aag aac acc aac ggg aat gcc caa ccg ctt gag aac atc ctg gct gcc	256
	Lys Asn Thr Asn Gly Asn Ala Gln Pro Leu Glu Asn Ile Leu Ala Ala	
	45 50 55	
	aat ggc gtc aac acg gtg cgg cag cga gtc tgg gtt aac ccc gcg gac	304
20	Asn Gly Val Asn Thr Val Arg Gln Arg Val Trp Val Asn Pro Ala Asp	
	60 65 70	
	ggc aac tac aac ctc gac tac aac atc gcg atc gcg aag agg gcg aag	352
	Gly Asn Tyr Asn Leu Asp Tyr Asn Ile Ala Ile Ala Lys Arg Ala Lys	
25	75 80 85	
	gct gcc ggg ctt ggc gtg tac atc gac ttc cac tac agc gac acc tgg	400
	Ala Ala Gly Leu Gly Val Tyr Ile Asp Phe His Tyr Ser Asp Thr Trp	
	90 95 100	
30	gcc gat cct gct cat cag acc atg ccc gct ggg tgg ccg agc gac att	448
	Ala Asp Pro Ala His Gln Thr Met Pro Ala Gly Trp Pro Ser Asp Ile	
	105 110 115 120	
	gac aac ctc tcc tgg aag ctc tac aac tac act ctg gac gcc gcc aac	496
35	Asp Asn Leu Ser Trp Lys Leu Tyr Asn Tyr Thr Leu Asp Ala Ala Asn	
	125 130 135	
	aag ctc cag aac gcg ggt atc cag ccc acc atc gtg tcc atc ggt aac	544
40	Lys Leu Gln Asn Ala Gly Ile Gln Pro Thr Ile Val Ser Ile Gly Asn	
	140 145 150	
	gag atc cgg gcc ggt ctg cta tgg ccc aca ggg aga acc gag aac tgg	592
	Glu Ile Arg Ala Gly Leu Leu Trp Pro Thr Gly Arg Thr Glu Asn Trp	
	155 160 165	
45	gcc aac att gcc cgg ttg ttg cac tcc gct gct tgg ggt atc aag gac	640

50

55

EP 2 183 363 B1

	Ala Asn Ile Ala Arg Leu Leu His Ser Ala Ala Trp Gly Ile Lys Asp	
	170 175 180	
5	tcg tcg ctc agc ccg aag cca aag atc atg atc cac ctc gac aac gga Ser Ser Leu Ser Pro Lys Pro Lys Ile Met Ile His Leu Asp Asn Gly	688
	185 190 195 200	
10	tg gtgagttata tcacccctttc ctgtgccgat aacctgttcc tcagcctggg Trp	740
	cgatgtggag agctgatggc gggggatcca aacag g gac tgg ggt acc cag aat Asp Trp Gly Thr Gln Asn	794
15		205
	tgg tgg tac acg aat gtc ttg aag cag ggt aca ctt gag ctg tcc gac Trp Trp Tyr Thr Asn Val Leu Lys Gln Gly Thr Leu Glu Leu Ser Asp	842
	210 215 220	
20	ttt gac atg atg ggc gtc tcg ttc tac ccc ttt tac tcg tcg tcg gca Phe Asp Met Met Gly Val Ser Phe Tyr Pro Phe Tyr Ser Ser Ser Ala	890
	225 230 235	
25	acc ttg agc gcc ctg aaa tcg agc ttg gac aac atg gcc aaa acc tgg Thr Leu Ser Ala Leu Lys Ser Ser Leu Asp Asn Met Ala Lys Thr Trp	938
	240 245 250 255	
30	aac aag gag att gcc gtg gtc gag acc aat tgg cca atc tct tgt ccc Asn Lys Glu Ile Ala Val Val Glu Thr Asn Trp Pro Ile Ser Cys Pro	986
	260 265 270	
	aac cca agg tac agt ttc ccc tcg gac gtc aag aac atc ccc ttc tcg Asn Pro Arg Tyr Ser Phe Pro Ser Asp Val Lys Asn Ile Pro Phe Ser	1034
	275 280 285	
35	ccg gaa gga cag acg acc ttc atc acc aac gtg gcc aac atc gtg tcc Pro Glu Gly Gln Thr Thr Phe Ile Thr Asn Val Ala Asn Ile Val Ser	1082
	290 295 300	
40	tcg gta agc cgc ggc gta ggc ctg ttt tat tgg gaa ccc gct tgg att Ser Val Ser Arg Gly Val Gly Leu Phe Tyr Trp Glu Pro Ala Trp Ile	1130
	305 310 315	
45	cac aat gca aac ctg ggc tcg tcg tgc gcc gac aac acc atg ttt tcg His Asn Ala Asn Leu Gly Ser Ser Cys Ala Asp Asn Thr Met Phe Ser	1178
	320 325 330 335	
50	caa tcc ggg cag gct ttg tcc agc ttg tcc gtt ttc cag aga atc tga Gln Ser Gly Gln Ala Leu Ser Ser Leu Ser Val Phe Gln Arg Ile	1226
	340 345 350	
55	<210> 47 <211> 350 <212> PRT <213> Chrysosporium lucknowense <400> 47	

EP 2 183 363 B1

Met Met Leu Thr Arg Phe Val Ala Gly Leu Leu Gly Ile Ser Ala Ala
1 5 10 15

5

10

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	Asp	Ala	Ala	Leu	Thr	Tyr	Arg	Gly	Val	Asp	Trp	Ser	Ser	Val	Val	Val	
				20					25					30			
5	Glu	Glu	Arg	Ala	Gly	Val	Ser	Tyr	Lys	Asn	Thr	Asn	Gly	Asn	Ala	Gln	
			35					40					45				
10	Pro	Leu	Glu	Asn	Ile	Leu	Ala	Ala	Asn	Gly	Val	Asn	Thr	Val	Arg	Gln	
		50					55					60					
15	Arg	Val	Trp	Val	Asn	Pro	Ala	Asp	Gly	Asn	Tyr	Asn	Leu	Asp	Tyr	Asn	
	65				70					75						80	
20	Ile	Ala	Ile	Ala	Lys	Arg	Ala	Lys	Ala	Ala	Gly	Leu	Gly	Val	Tyr	Ile	
					85				90						95		
25	Asp	Phe	His	Tyr	Ser	Asp	Thr	Trp	Ala	Asp	Pro	Ala	His	Gln	Thr	Met	
				100					105					110			
30	Pro	Ala	Gly	Trp	Pro	Ser	Asp	Ile	Asp	Asn	Leu	Ser	Trp	Lys	Leu	Tyr	
			115					120					125				
35	Asn	Tyr	Thr	Leu	Asp	Ala	Ala	Asn	Lys	Leu	Gln	Asn	Ala	Gly	Ile	Gln	
		130					135					140					
40	Pro	Thr	Ile	Val	Ser	Ile	Gly	Asn	Glu	Ile	Arg	Ala	Gly	Leu	Leu	Trp	
	145					150					155					160	
45	Pro	Thr	Gly	Arg	Thr	Glu	Asn	Trp	Ala	Asn	Ile	Ala	Arg	Leu	Leu	His	
					165					170					175		
50	Ser	Ala	Ala	Trp	Gly	Ile	Lys	Asp	Ser	Ser	Leu	Ser	Pro	Lys	Pro	Lys	
				180					185					190			
55	Ile	Met	Ile	His	Leu	Asp	Asn	Gly	Trp	Asp	Trp	Gly	Thr	Gln	Asn	Trp	
			195					200					205				
60	Trp	Tyr	Thr	Asn	Val	Leu	Lys	Gln	Gly	Thr	Leu	Glu	Leu	Ser	Asp	Phe	
		210					215					220					
65	Asp	Met	Met	Gly	Val	Ser	Phe	Tyr	Pro	Phe	Tyr	Ser	Ser	Ser	Ala	Thr	
	225					230					235					240	
70	Leu	Ser	Ala	Leu	Lys	Ser	Ser	Leu	Asp	Asn	Met	Ala	Lys	Thr	Trp	Asn	
				245						250					255		

EP 2 183 363 B1

	Lys	Glu	Ile	Ala	Val	Val	Glu	Thr	Asn	Trp	Pro	Ile	Ser	Cys	Pro	Asn	
				260					265					270			
5	Pro	Arg	Tyr	Ser	Phe	Pro	Ser	Asp	Val	Lys	Asn	Ile	Pro	Phe	Ser	Pro	
			275					280					285				
10	Glu	Gly	Gln	Thr	Thr	Phe	Ile	Thr	Asn	Val	Ala	Asn	Ile	Val	Ser	Ser	
		290					295					300					
15	Val	Ser	Arg	Gly	Val	Gly	Leu	Phe	Tyr	Trp	Glu	Pro	Ala	Trp	Ile	His	
	305					310					315					320	
20	Asn	Ala	Asn	Leu	Gly	Ser	Ser	Cys	Ala	Asp	Asn	Thr	Met	Phe	Ser	Gln	
					325					330					335		
25	Ser	Gly	Gln	Ala	Leu	Ser	Ser	Leu	Ser	Val	Phe	Gln	Arg	Ile			
				340					345					350			
30	<210> 48																
	<211> 1050																
	<212> DNA																
	<213> Chrysosporium lucknowense																
35	<400> 48																
40																	
45																	
50																	
55																	

EP 2 183 363 B1

	atgatgctca cacgcttcgt ggctggcctg ctcggcattct ccgccgcgga tgccgccttc	60
	acctacagag gcgtggattg gtcctcagtg gtggttgagg aacggggccgg cgtctcgtac	120
5	aagaacacca acgggaatgc ccaaccgctt gagaacatcc tggctgccaa tggcgtcaac	180
	acggtgcggc agcgagtctg ggttaacccc gcggacggca actacaacct cgactacaac	240
10	atcgcgatcg cgaagagggc gaaggctgcc gggcttggcg tgtacatcga cttccactac	300
	agcgacacct gggccgatcc tgctcatcag accatgcccc ctgggtggcc gagcgacatt	360
	gacaacctct cctggaagct ctacaactac actctggacg ccgccaacaa gctccagaac	420
15	gcgggtatcc agcccaccat cgtgtccatc ggtaacgaga tccggggccgg tctgctatgg	480
	cccacagggga gaaccgagaa ctgggccaac attgcccgtt tgttgactc cgctgcttgg	540
20	ggtatcaagg actcgtcgtc cagcccgaag ccaaagatca tgatccacct cgacaacgga	600
	tgggactggg gtaccacagaa ttggtgttac acgaatgtct tgaagcaggg tacacttgag	660
	ctgtccgact ttgacatgat gggcgtctcg ttctaccctt tttactcgtc gtcggcaacc	720
25	ttgagcgccc tgaaatcgag cttggacaac atggccaaaa cctggaacaa ggagattgcc	780
	gtggtcgaga ccaattggcc aatctcttgt cccaacccaa ggtacagttt cccctcggac	840
	gtcaagaaca tccccttctc gccggaagga cagacgacct tcatcaccaa cgtggccaac	900
30	atcgtgtcct cggtaaagccg cggcgtaggc ctgttttatt gggaaccgc ttggattcac	960
	aatgcaaacc tgggctcgtc gtgcgccgac aacaccatgt tttcgcaatc cgggcaggct	1020
35	ttgtccagct tgtccgtttt ccagagaatc	1050

<210> 49

<211> 2344

<212> DNA

40 <213> Chrysosporium lucknowense

<220>

<221> CDS

<222> (54) .. (276)

45

<220>

<221> CDS

<222> (348)..(717)

50

<220>

<221> CDS

<222> (814)..(887)

55

<220>

<221> CDS

<222> (947)..(1078)

<220>

EP 2 183 363 B1

<221> CDS

<222> (1144)..(2228)

<400> 49

5	agctccagga ccacctccca tccgtaaacg gatctggcct cgtcacgccc gcc atg	56
	Met	
	1	
10	cac gct ctc tcg tcg ctc gct gtc ctc ggc gcc tgg gcc gtc cag acg	104
	His Ala Leu Ser Ser Leu Ala Val Leu Gly Ala Trp Ala Val Gln Thr	
	5 10 15	
15	gtc ttg ggc cgt ccg gcc acc ctc tca aag cgg gcc acc gac tcc ttc	152
	Val Leu Gly Arg Pro Ala Thr Leu Ser Lys Arg Ala Thr Asp Ser Phe	
	20 25 30	
20	atc gag acc gag acg cct atc gca tgg gaa aag ctg cgg tgc aac atc	200
	Ile Glu Thr Glu Thr Pro Ile Ala Trp Glu Lys Leu Arg Cys Asn Ile	
	35 40 45	
25	ggc gct aac ggc tgt gcg gct tcc gga gcc gct gcc ggt gtg gtc att	248
	Gly Ala Asn Gly Cys Ala Ala Ser Gly Ala Ala Ala Gly Val Val Ile	
	50 55 60 65	
30	gcc agc ccg tcc aag tcg gat cca gac t gtgcgttgac agcggcccct	296
	Ala Ser Pro Ser Lys Ser Asp Pro Asp	
	70	
35	accccttatt atcgcaactt cggtcgggtgt tgagatgctg agacgcgaca g ac ttc	352
	Tyr Phe	
40		
45		
50		
55		

EP 2 183 363 B1

	tac acc tgg act cga gat gcc ggc ctg gtc ctg acg ggt atc gtg gac	400
	Tyr Thr Trp Thr Arg Asp Ala Gly Leu Val Leu Thr Gly Ile Val Asp	
	80 85 90	
5	gcc ctg tcc caa aac tac tcg gcg gcc ctg cag acc aac att cag gac	448
	Ala Leu Ser Gln Asn Tyr Ser Ala Ala Leu Gln Thr Asn Ile Gln Asp	
	95 100 105	
10	tac atc atc gcc cag gcc aag ctc cag ggt gtt tcg aac ccc tcc ggt	496
	Tyr Ile Ile Ala Gln Ala Lys Leu Gln Gly Val Ser Asn Pro Ser Gly	
	110 115 120	
15	agc ctc tcg gac ggc acc ggt ctt ggc gag ccc aag ttc aat gtc gat	544
	Ser Leu Ser Asp Gly Thr Gly Leu Gly Glu Pro Lys Phe Asn Val Asp	
	125 130 135 140	
	ctc acc cag ttc acg ggc gac tgg ggc cgg ccg cag cgc gac ggt ccg	592
	Leu Thr Gln Phe Thr Gly Asp Trp Gly Arg Pro Gln Arg Asp Gly Pro	
	145 150 155	
20	cct ctc cgg gcc atc gcc ctc atc cgc tac gcc aag tgg ctg gct tcc	640
	Pro Leu Arg Ala Ile Ala Leu Ile Arg Tyr Ala Lys Trp Leu Ala Ser	
	160 165 170	
25	aac ggt tac aag gac acg gcc aac agc gtc gtc tgg ccc gtc atc aag	688
	Asn Gly Tyr Lys Asp Thr Ala Asn Ser Val Val Trp Pro Val Ile Lys	
	175 180 185	
30	aac gac ctg gcc tat gcc gct cag tat tg gtgagtgcc cttctacttc	737
	Asn Asp Leu Ala Tyr Ala Ala Gln Tyr Trp	
	190 195	
	tcgcccttgg gggagtttct tcccgtcctg acgccggagg cacgctaaca cacgccacac	797
35	acacacgcac atacag g aac gag act ggt ttc gac ctg tgg gag gag gtt	847
	Asn Glu Thr Gly Phe Asp Leu Trp Glu Glu Val	
	200 205	
	ccc ggc agc tcg ttc ttc acc att gcc agc acg cac cga g gtgggtgggc	897
	Pro Gly Ser Ser Phe Phe Thr Ile Ala Ser Thr His Arg	
	210 215 220	
40	cccgttgtac atatatgggt tgatcaacga attactgacg cagcgacag cc ttg gta	954
	Ala Leu Val	
	225	
45	gag gga gct gcc ctc gct gcc cag ctc ggc acc gaa tgc agc gcc tgc	1002
	Glu Gly Ala Ala Leu Ala Ala Gln Leu Gly Thr Glu Cys Ser Ala Cys	
	230 235 240	
50	atc acc gtc gcg ccc caa gtc ctc tgc ttc cag cag agc ttc tgg aac	1050
	Ile Thr Val Ala Pro Gln Val Leu Cys Phe Gln Gln Ser Phe Trp Asn	
	245 250 255	
	ccg tcg ggc ggt tac gtt gtc tcg aac a gtaagtctct tctcgtggtg	1098
	Pro Ser Gly Gly Tyr Val Val Ser Asn	
	260 265	
55	gtgccgctcc cagcctcaat ttccacacaa ctaaacaacg gcaag tc aac ggc ggc	1154
	Ile Asn Gly Gly	

EP 2 183 363 B1

270

5	aac aac cgg tcc ggc aag gat ctc aac tcg gtc ctg gcc tcc atc cac Asn Asn Arg Ser Gly Lys Asp Leu Asn Ser Val Leu Ala Ser Ile His 275 280 285	1202
10	acc ttt gac ccg gcg gtc ggc tgc gac tcg gtc acc ttc cag ccc tgc Thr Phe Asp Pro Ala Val Gly Cys Asp Ser Val Thr Phe Gln Pro Cys 290 295 300	1250
15	agc gac aag gcg ctc tcc aac cac aag gcc tac gtc gac tcc ttc cgc Ser Asp Lys Ala Leu Ser Asn His Lys Ala Tyr Val Asp Ser Phe Arg 305 310 315	1298
20	agc gtc tac gcc atc aac tcg ggc att gcc cag ggc aag gcc gtc gcc Ser Val Tyr Ala Ile Asn Ser Gly Ile Ala Gln Gly Lys Ala Val Ala 320 325 330	1346
25	gtg ggc cgc tac tcg gag gac gtc tac tac aac ggc aac ccg tgg tac Val Gly Arg Tyr Ser Glu Asp Val Tyr Tyr Asn Gly Asn Pro Trp Tyr 335 340 345 350	1394
30	ctg gcc aac ttc gcg gcc gcc gag cag ctc tac gac gcc gtc ttc gtc Leu Ala Asn Phe Ala Ala Ala Glu Gln Leu Tyr Asp Ala Val Phe Val 355 360 365	1442
35	tgg aag aag cag cag tcc atc gag gtc acc cag ctg tcc ctc ccc ttc Trp Lys Lys Gln Gln Ser Ile Glu Val Thr Gln Leu Ser Leu Pro Phe 370 375 380	1490
40	ttc aag gac ctg ctc ccc ggc atc tcc acc ggc acc tac acc ccg tcg Phe Lys Asp Leu Leu Pro Gly Ile Ser Thr Gly Thr Tyr Thr Pro Ser 385 390 395	1538
45	tcg tcg acg tac cag cag atc ctc gac gcc gtc tcg gcc tac gcc gac Ser Ser Thr Tyr Gln Gln Ile Leu Asp Ala Val Ser Ala Tyr Ala Asp 400 405 410	1586
50	ggc ttc atc gac gtc gcg gcc aag tac acc ccc tcg gac ggc tcc ctg Gly Phe Ile Asp Val Ala Ala Lys Tyr Thr Pro Ser Asp Gly Ser Leu 415 420 425 430	1634
55	gcc gag cag tac acg cgc gac tcg ggc cag ccg atc tcg gcc aag gac Ala Glu Gln Tyr Thr Arg Asp Ser Gly Gln Pro Ile Ser Ala Lys Asp 435 440 445	1682
60	ctg acc tgg tcc tac gcc gcc ttc ctc tcg gcc gcc gac cgc cgc gcg Leu Thr Trp Ser Tyr Ala Ala Phe Leu Ser Ala Ala Asp Arg Arg Ala 450 455 460	1730
65	ggc atc gtc ccg gcc ggc tgg tcc gcc gag cac ggc aag acg ctg ccc Gly Ile Val Pro Ala Gly Trp Ser Ala Glu His Gly Lys Thr Leu Pro 465 470 475	1778
70	ggc tcg tgc tcg gcc gtc cag gtc gcc ggc acc tac acc cag gcc acc Gly Ser Cys Ser Ala Val Gln Val Ala Gly Thr Tyr Thr Gln Ala Thr 480 485 490	1826
75	gcc acc tcc ttc ccg ccc ggc cag acg ccc aac ccg acc agc gac acc Ala Thr Ser Phe Pro Pro Gly Gln Thr Pro Asn Pro Thr Ser Asp Thr 495 500 505 510	1874

EP 2 183 363 B1

	ccg gcc ccg ttc ccc acg gcc tgc gcc gac tcc acc cag gtc ttc gtc	1922
	Pro Ala Pro Phe Pro Thr Ala Cys Ala Asp Ser Thr Gln Val Phe Val	
	515 520 525	
5	acc ttc cgc gcc gag gtg acc acc cag tgg ggc cag tcg gtc aag gtc	1970
	Thr Phe Arg Ala Glu Val Thr Thr Gln Trp Gly Gln Ser Val Lys Val	
	530 535 540	
10	gtc ggc agc tcg tcc gag ctc ggc aac tgg gac gtc tcc aag gcc ccg	2018
	Val Gly Ser Ser Ser Glu Leu Gly Asn Trp Asp Val Ser Lys Ala Pro	
	545 550 555	
15	cgc ctg tcc gcg tcg gcc tac acg gcg tcg gac ccg ctc tgg gcc atc	2066
	Arg Leu Ser Ala Ser Ala Tyr Thr Ala Ser Asp Pro Leu Trp Ala Ile	
	560 565 570	
20	acg gtg ccc atg aag gcc ggc cag tcg gtg cag tac aag ttc gtc aag	2114
	Thr Val Pro Met Lys Ala Gly Gln Ser Val Gln Tyr Lys Phe Val Lys	
	575 580 585 590	
25	gtc aac ggg gac ggg tcg atc cag tgg gag tcg gac ccg aac cgc cag	2162
	Val Asn Gly Asp Gly Ser Ile Gln Trp Glu Ser Asp Pro Asn Arg Gln	
	595 600 605	
30	ttc acg gtc agc tct tcc tcc acg gcc agc ggc tgt gcc tgg cag acc	2210
	Phe Thr Val Ser Ser Ser Ser Thr Ala Ser Gly Cys Ala Trp Gln Thr	
	610 615 620	
35	atc gag gcg acc tgg cgg tagatatggg agtactggag caaaagcctc	2258
	Ile Glu Ala Thr Trp Arg	
	625	
40	attcggagcc ttcggtggct cttgggtggg ggggggggtg gaaaagaggg gaggtaatag	2318
	gatgatcatg agcgtttggg aaaaaa	2344
	<210> 50	
	<211> 628	
	<212> PRT	
	<213> Chrysosporium lucknowense	
45	Met His Ala Leu SerSer Leu Ala Val Leu Gly Ala Trp Ala Val Gln	
	1 5 10 15	
50	Thr Val Leu Gly Arg Pro Ala Thr Leu Ser Lys Arg Ala Thr Asp Ser	
	20 25 30	
55	Phe Ile Glu Thr Glu Thr Pro Ile Ala Trp Glu Lys Leu Arg Cys Asn	
	35 40 45	
60	Ile Gly Ala Asn Gly Cys Ala Ala Ser Gly Ala Ala Ala Gly Val Val	
	50 55 60	

EP 2 183 363 B1

	Ile	Ala	Ser	Pro	Ser	Lys	Ser	Asp	Pro	Asp	Tyr	Phe	Tyr	Thr	Trp	Thr	65	70	75	80
5	Arg	Asp	Ala	Gly	Leu	Val	Leu	Thr	Gly	Ile	Val	Asp	Ala	Leu	Ser	Gln		85	90	95
10	Asn	Tyr	Ser	Ala	Ala	Leu	Gln	Thr	Asn	Ile	Gln	Asp	Tyr	Ile	Ile	Ala	100	105	110	
15	Gln	Ala	Lys	Leu	Gln	Gly	Val	Ser	Asn	Pro	Ser	Gly	Ser	Leu	Ser	Asp	115	120	125	
20	Gly	Thr	Gly	Leu	Gly	Glu	Pro	Lys	Phe	Asn	Val	Asp	Leu	Thr	Gln	Phe	130	135	140	
25	Thr	Gly	Asp	Trp	Gly	Arg	Pro	Gln	Arg	Asp	Gly	Pro	Pro	Leu	Arg	Ala	145	150	155	160
	Ile	Ala	Leu	Ile	Arg	Tyr	Ala	Lys	Trp	Leu	Ala	Ser	Asn	Gly	Tyr	Lys		165	170	175
30	Asp	Thr	Ala	Asn	Ser	Val	Val	Trp	Pro	Val	Ile	Lys	Asn	Asp	Leu	Ala	180	185	190	
35	Tyr	Ala	Ala	Gln	Tyr	Trp	Asn	Glu	Thr	Gly	Phe	Asp	Leu	Trp	Glu	Glu	195	200	205	
40	Val	Pro	Gly	Ser	Ser	Phe	Phe	Thr	Ile	Ala	Ser	Thr	His	Arg	Ala	Leu	210	215	220	
	Val	Glu	Gly	Ala	Ala	Leu	Ala	Ala	Gln	Leu	Gly	Thr	Glu	Cys	Ser	Ala	225	230	235	240
45	Cys	Ile	Thr	Val	Ala	Pro	Gln	Val	Leu	Cys	Phe	Gln	Gln	Ser	Phe	Trp	245	250	255	
50	Asn	Pro	Ser	Gly	Gly	Tyr	Val	Val	Ser	Asn	Ile	Asn	Gly	Gly	Asn	Asn	260	265	270	
	Arg	Ser	Gly	Lys	Asp	Leu	Asn	Ser	Val	Leu	Ala	Ser	Ile	His	Thr	Phe	275	280	285	
55	Asp	Pro	Ala	Val	Gly	Cys	Asp	Ser	Val	Thr	Phe	Gln	Pro	Cys	Ser	Asp	290	295	300	
	Lys	Ala	Leu	Ser	Asn	His	Lys	Ala	Tyr	Val	Asp	Ser	Phe	Arg	Ser	Val				

EP 2 183 363 B1

	305					310					315					320
5	Tyr	Ala	Ile	Asn	Ser	Gly	Ile	Ala	Gln	Gly	Lys	Ala	Val	Ala	Val	Gly
				325						330					335	
10	Arg	Tyr	Ser	Glu	Asp	Val	Tyr	Tyr	Asn	Gly	Asn	Pro	Trp	Tyr	Leu	Ala
				340					345					350		
15	Asn	Phe	Ala	Ala	Ala	Glu	Gln	Leu	Tyr	Asp	Ala	Val	Phe	Val	Trp	Lys
			355					360					365			
20	Lys	Gln	Gln	Ser	Ile	Glu	Val	Thr	Gln	Leu	Ser	Leu	Pro	Phe	Phe	Lys
		370					375					380				
25	Asp	Leu	Leu	Pro	Gly	Ile	Ser	Thr	Gly	Thr	Tyr	Thr	Pro	Ser	Ser	Ser
	385					390					395					400
30	Thr	Tyr	Gln	Gln	Ile	Leu	Asp	Ala	Val	Ser	Ala	Tyr	Ala	Asp	Gly	Phe
					405					410					415	
35	Ile	Asp	Val	Ala	Ala	Lys	Tyr	Thr	Pro	Ser	Asp	Gly	Ser	Leu	Ala	Glu
				420					425					430		
40	Gln	Tyr	Thr	Arg	Asp	Ser	Gly	Gln	Pro	Ile	Ser	Ala	Lys	Asp	Leu	Thr
			435					440					445			
45	Trp	Ser	Tyr	Ala	Ala	Phe	Leu	Ser	Ala	Ala	Asp	Arg	Arg	Ala	Gly	Ile
	450						455					460				
50	Val	Pro	Ala	Gly	Trp	Ser	Ala	Glu	His	Gly	Lys	Thr	Leu	Pro	Gly	Ser
	465					470					475					480
55	Cys	Ser	Ala	Val	Gln	Val	Ala	Gly	Thr	Tyr	Thr	Gln	Ala	Thr	Ala	Thr
					485					490					495	
60	Ser	Phe	Pro	Pro	Gly	Gln	Thr	Pro	Asn	Pro	Thr	Ser	Asp	Thr	Pro	Ala
				500					505					510		
65	Pro	Phe	Pro	Thr	Ala	Cys	Ala	Asp	Ser	Thr	Gln	Val	Phe	Val	Thr	Phe
			515					520					525			
70	Arg	Ala	Glu	Val	Thr	Thr	Gln	Trp	Gly	Gln	Ser	Val	Lys	Val	Val	Gly
		530					535					540				
75	Ser	Ser	Ser	Glu	Leu	Gly	Asn	Trp	Asp	Val	Ser	Lys	Ala	Pro	Arg	Leu
	545					550					555					560

EP 2 183 363 B1

	Ser	Ala	Ser	Ala	Tyr	Thr	Ala	Ser	Asp	Pro	Leu	Trp	Ala	Ile	Thr	Val	
					565					570					575		
5	Pro	Met	Lys	Ala	Gly	Gln	Ser	Val	Gln	Tyr	Lys	Phe	Val	Lys	Val	Asn	
				580					585					590			
10	Gly	Asp	Gly	Ser	Ile	Gln	Trp	Glu	Ser	Asp	Pro	Asn	Arg	Gln	Phe	Thr	
			595					600					605				
15	Val	Ser	Ser	Ser	Ser	Thr	Ala	Ser	Gly	Cys	Ala	Trp	Gln	Thr	Ile	Glu	
		610					615					620					
	Ala	Thr	Trp	Arg													
		625															
20	<210> 51																
	<211> 1884																
	<212> DNA																
	<213> Chrysosporium lucknowense																
25	<400> 51																
	atgcacgctc	tctcgtcgct	cgctgtcctc	ggcgctggg	ccgtccagac	ggtcttgggc											60
30	cgctccggcca	ccctctcaaa	gcggggcacc	gactccttca	tcgagaccga	gacgcctatc											120
	gcatgggaaa	agctgcggtg	caacatcggc	gctaacggct	gtgcggcttc	cggagccgct											180
	gccggtgtgg	tcattgccag	cccgtccaag	tcggatccag	actacttcta	cacctggact											240
35	cgagatgccg	gcctggtcct	gacgggtatc	gtggacgccc	tgtcccaaaa	ctactcggcg											300
	gccctgcaga	ccaacattca	ggactacatc	atcgcccagg	ccaagctcca	gggtgtttcg											360
	aaccctccg	gtagcctctc	ggacggcacc	ggtcttggcg	agcccaagtt	caatgtcgat											420
40	ctcaccagct	tcacgggcga	ctggggccgg	ccgcagcgcg	acggtcggcc	tctccggggc											480
	atcgccctca	tccgctacgc	caagtggctg	gcttccaacg	gttacaagga	cacggccaac											540
45	agcgtcgtct	ggcccgtcat	caagaacgac	ctggcctatg	ccgctcagta	ttggaacgag											600
	actggtttcg	acctgtggga	ggaggttccc	ggcagctcgt	tcttcaccat	tgccagcacg											660
	caccgagcct	tggtagaggg	agctgccctc	gctgcccagc	tcggcaccga	atgcagcgcc											720
50	tgcatacccg	tcgcgcccc	agtcctctgc	ttccagcaga	gcttctggaa	cccgtcgggc											780
	ggttacgttg	tctcgaacat	caacggcggc	aacaaccggt	ccggcaagga	tctcaactcg											840
	gtcctggcct	ccatccacac	ctttgacccg	gcggtcggct	gcgactcggg	caccttccag											900
55	ccctgcagcg	acaaggcgct	ctccaaccac	aaggcctacg	tcgactcctt	ccgcagcgtc											960
	tacgccatca	actcgggcat	tgcccagggc	aaggccgtcg	ccgtggggccg	ctactcggag											1020

EP 2 183 363 B1

	gacgtctact acaacggcaa cccgtggtac ctggccaact tcgcggccgc cgagcagctc	1080
	tacgacgccg tcttcgtctg gaagaagcag cagtccatcg aggtcaccca gctgtccctc	1140
5	cccttcttca aggacctgct ccccggcacg tccaccggca cctacacccc gtcgtcgctg	1200
	acgtaccagc agatcctcga cgccgtctcg gcctacgccg acggcttcat cgacgtcgcg	1260
10	gccaagtaca cccctcggga cggctccctg gccgagcagt acacgcgcga ctcgggccag	1320
	ccgatctcgg ccaaggacct gacctggtcc tacgccgctt tcctctcggc cgccgaccgc	1380
	cgcgcgggca tcgtcccggc cggctggtcc gccgagcacg gcaagacgct gcccggtcgc	1440
15	tgctcggccg tccaggctcg cgccacctac acccaggcca ccgccacctc cttcccgccc	1500
	ggccagacgc ccaacccgac cagcgacacc ccggccccgt tccccacggc ctgcgccgac	1560
20	tccacccagg tcttcgtcac cttccgcgcc gaggtgacca ccagtgagg ccagtcggtc	1620
	aaggctcgct gcagctcgtc cgagctcggc aactgggacg tctccaaggc cccgcgcctg	1680
	tccgcgtcgg cctacacggc gtcggacccg ctctggggca tcacggtgcc catgaaggcc	1740
25	ggccagtcgg tgcagtacaa gttcgtcaag gtcaacgggg acgggtcgat ccagtgagg	1800
	tcggacccga accgccagtt cacggtcagc tcttcctcca cggccagcgg ctgtgcctgg	1860
	cagaccatcg aggcgacctg gcgg	1884

30

<210> 52
 <211> 1637
 <212> DNA
 <213> Chrysosporium lucknowense

35

<220>
 <221> CDS
 <222> (148)..(864)

40

<220>
 <221> CDS
 <222> (1194)..(1601)

45

<400> 52

50

55

EP 2 183 363 B1

	gtataagatt gacgacgac cgggtagaca aaccgacttg cagatctttg caacccatat	60
	ttgtactagc ctccagggtc atcctgccca ggtctcctgg cctagagttg tatacatttg	120
5	ctttgttaac ggagcctttt cctcgtc atg aga tcc tcc gcc atc tgg tcc acc	174
	Met Arg Ser Ser Ala Ile Trp Ser Thr	
	1 5	
10	gcc ttc gcg gcg ttg gca cac act gcc gcc gcc gtc act ggt gct gcc	222
	Ala Phe Ala Ala Leu Ala His Thr Ala Ala Ala Val Thr Gly Ala Ala	
	10 15 20 25	
15	gag ggc ttc gcc aaa ggg gtc acg gga ggt ggc aat gcg act ccc gtc	270
	Glu Gly Phe Ala Lys Gly Val Thr Gly Gly Gly Asn Ala Thr Pro Val	
20		
25		
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

EP 2 183 363 B1

	gaatagaata atgcctaaat aggtacagat gctaacggaa ttgcttccac ccaatccag	1193
5	atg atc acc ttc aag aac aac ctc atc acc cac atg tcg ggc cgc agc Met Ile Thr Phe Lys Asn Asn Leu Ile Thr His Met Ser Gly Arg Ser 240 245 250 255	1241
10	ccc aag gtg gcg ggc aac acg ctg ctg cac gcg gtc aac aac tac tgg Pro Lys Val Ala Gly Asn Thr Leu Leu His Ala Val Asn Asn Tyr Trp 260 265 270	1289
	tac gac agc aag ggc cac aac ttc gag ctg gac gcg ggg gcc atg gtg Tyr Asp Ser Lys Gly His Asn Phe Glu Leu Asp Ala Gly Ala Met Val 275 280 285	1337
15	ctc gtc gag ggc agc gtg ttc cag aac gtc gag gcg ccg ctc cag tcc Leu Val Glu Gly Ser Val Phe Gln Asn Val Glu Ala Pro Leu Gln Ser 290 295 300	1385
20	ggc ggc gcc ggc cgg ctc ttc gcc agc tcc ggc acc gcc tcg gcc tgc Gly Gly Ala Gly Arg Leu Phe Ala Ser Ser Gly Thr Ala Ser Ala Cys 305 310 315	1433
25	gcc ccg tac ctc ggc cgc gtg tgc cag gcc aac ggc ttc ggc agc tcc Ala Pro Tyr Leu Gly Arg Val Cys Gln Ala Asn Gly Phe Gly Ser Ser 320 325 330 335	1481
	ggc tcc ctc tcc ggc tcc gac acc tcc ttc ctg cag tac ttc cag aac Gly Ser Leu Ser Gly Ser Asp Thr Ser Phe Leu Gln Tyr Phe Gln Asn 340 345 350	1529
30	aag aac atc gcc ggc gcg aca gac tac aac acg gcc aag aac gtc ctc Lys Asn Ile Ala Gly Ala Thr Asp Tyr Asn Thr Ala Lys Asn Val Leu 355 360 365	1577
35	gac acc gcc gga ttt ggc aca atc tgatcgatcg tgttgcgttt ccttgga Asp Thr Ala Gly Phe Gly Thr Ile 370 375	1631
	ggtgga	1637
40	<210> 53 <211> 375 <212> PRT <213> Chrysosporium lucknowense	
45	<400> 53	
	Met Arg Ser Ser Ala Ile Trp Ser Thr Ala Phe Ala Ala Leu Ala His 1 5 10 15	
50	Thr Ala Ala Ala Val Thr Gly Ala Ala Glu Gly Phe Ala Lys Gly Val 20 25 30	
55	Thr Gly Gly Gly Asn Ala Thr Pro Val Tyr Pro Lys Thr Thr Ala Glu 35 40 45	

EP 2 183 363 B1

	Leu	Val	Ser	Tyr	Leu	Gly	Asp	Ser	Glu	Pro	Arg	Val	Ile	Met	Leu	Asp
	50						55					60				
5	Arg	Thr	Phe	Asp	Phe	Thr	Gly	Thr	Glu	Gly	Thr	Ala	Thr	Glu	Thr	Gly
	65					70					75					80
10	Cys	Ala	Pro	Trp	Gly	Thr	Gly	Ser	Gly	Cys	Gln	Leu	Ala	Leu	Asn	Lys
					85					90					95	
15	Asn	Asp	Trp	Cys	Thr	Arg	Glu	Gln	Pro	Asn	Ala	Pro	Lys	Val	Ser	Ile
				100					105					110		
20	Thr	Tyr	Asp	Lys	Ala	Gly	Ile	Leu	Gly	Ile	Thr	Val	Lys	Ser	Asn	Lys
			115					120					125			
25	Ser	Leu	Ile	Gly	Val	Gly	Asn	Lys	Gly	Val	Ile	Lys	Gly	Lys	Gly	Ile
	130						135					140				
30	Arg	Met	Val	Ser	Gly	Thr	Ser	Asn	Val	Ile	Ile	Gln	Asn	Ile	His	Ile
	145					150					155					160
35	Thr	Asn	Leu	Asn	Pro	Gln	Tyr	Val	Trp	Gly	Gly	Asp	Ala	Ile	Thr	Leu
				165						170					175	
40	Asp	Asn	Thr	Glu	Met	Ile	Trp	Ile	Asp	His	Val	Thr	Thr	Ser	Leu	Ile
				180					185					190		
45	Gly	Arg	Gln	His	Val	Val	Leu	Gly	Asn	Asn	Pro	Ser	Gly	Arg	Val	Thr
			195					200					205			
50	Ile	Ser	Asn	Ser	Lys	Phe	Asp	Gly	Gln	Thr	Ser	Trp	Ser	Ala	Thr	Cys
	210						215					220				
55	Asn	Thr	Tyr	His	Tyr	Trp	Gly	Leu	Tyr	Phe	Thr	Gly	Ser	Asn	Asp	Met
	225					230					235					240
60	Ile	Thr	Phe	Lys	Asn	Asn	Leu	Ile	Thr	His	Met	Ser	Gly	Arg	Ser	Pro
				245						250					255	
65	Lys	Val	Ala	Gly	Asn	Thr	Leu	Leu	His	Ala	Val	Asn	Asn	Tyr	Trp	Tyr
				260					265					270		
70	Asp	Ser	Lys	Gly	His	Asn	Phe	Glu	Leu	Asp	Ala	Gly	Ala	Met	Val	Leu
			275					280					285			
75	Val	Glu	Gly	Ser	Val	Phe	Gln	Asn	Val	Glu	Ala	Pro	Leu	Gln	Ser	Gly

EP 2 183 363 B1

	290	295	300
5	Gly Ala Gly Arg Leu Phe Ala Ser Ser Gly Thr Ala Ser Ala Cys Ala 305 310 315 320		
10	Pro Tyr Leu Gly Arg Val Cys Gln Ala Asn Gly Phe Gly Ser Ser Gly 325 330 335		
15	Ser Leu Ser Gly Ser Asp Thr Ser Phe Leu Gln Tyr Phe Gln Asn Lys 340 345 350		
20	Asn Ile Ala Gly Ala Thr Asp Tyr Asn Thr Ala Lys Asn Val Leu Asp 355 360 365		
	Thr Ala Gly Phe Gly Thr Ile 370 375		

<210> 54

<211> 1125

25 <212> DNA

<213> Chrysosporium lucknowense

<400> 54

30

35

40

45

50

55

EP 2 183 363 B1

	atgagatcct cgcacatctg gtccaccgcc ttcgcggcgt tggcacacac tgccgcgcgc	60
	gtcactgggtg ctgccgaggg cttcgccaaa ggggtcacgg gaggtggcaa tgcgactccc	120
5	gtctatccca aaaccacggc cgagctggtc tcgtacctgg gtgactcgga gcctcgtgtc	180
	atcatgctgg accggacatt cgacttcacc ggaaccgagg gcaccgcgac cgagacgggc	240
10	tgcgccccct ggggcaccgg ctcgggctgc cagctggccc tgaacaagaa cgactggtgc	300
	acccgggagc agccgaacgc gcccaaagtg agcatcacgt acgacaaggc cggcattctc	360
	ggcatcaccg tcaagtccaa caagtccctc atcggcgtcg gcaacaaggg cgtcatcaag	420
15	ggcaaaggca tccgcatggt cagcggcacc agcaacgtca tcatccagaa catccacatc	480
	accaacctca acccgagta cgtctggggc ggcgacgcca tcacgctcga caataccgag	540
20	atgatctgga tcgaccacgt caccacctcc ctcatcggcc gccagcacgt cgtcctgggt	600
	aacaacccga gcggccgcgt caccatctcc aacagcaagt tcgacggcca gacctcctgg	660
	tccgccacct gcaacaccta ccaactactg ggctgtact ttacagggtc caacgacatg	720
25	atcaccttca agaacaacct catcacccac atgtcgggcc gcagcccaa ggtggcgggc	780
	aacacgctgc tgcacgcggt caacaactac tggtagcaca gcaagggcca caacttcgag	840
	ctggacgcgg gggccatggt gctcgtcgag ggcagcgtgt tccagaacgt cgaggcgccg	900
30	ctccagtcgg gcggcgccgg ccggtcttcc gccagctccg gcaccgcctc ggctgcgcc	960
	ccgtacctcg gccgcgtgtg ccaggccaac ggcttcggca gctccggctc cctctccggc	1020
35	tccgacacct ccttcctgca gtacttccag aacaagaaca tcgccggcgc gacagactac	1080
	aacacggcca agaacgtcct cgacaccgcc ggatttggca caatc	1125
40	<210> 55 <211> 1904 <212> DNA <213> Chrysosporium lucknowense	
45	<220> <221> CDS <222> (179)..(330)	
50	<220> <221> CDS <222> (443)..(591)	
55	<220> <221> CDS <222> (732)..(1708)	
	<400> 55	

EP 2 183 363 B1

	gtataagagc actgtgatcg ttcacaaagc cggcgctactc ctctagagca tctatcatca	60
	acatcaccag aaagggtcaag accaggtggt tgccatatcc agtcgcaaaa gagccaaaga	120
5	gcgaaggagc acgaaagcac agcccaatca ttccctgctt tgctacttct tctccacc	178
	atg gtg act ccc tct ctg aag aag gca gct ctg gct gcc ctg tcc ctc	226
	Met Val Thr Pro Ser Leu Lys Lys Ala Ala Leu Ala Ala Leu Ser Leu	
	1 5 10 15	
10	ttc ccg ctc ctc tcc cta gct tcg ccg gtg cca gca aca gcc gaa gct	274
	Phe Pro Leu Leu Ser Leu Ala Ser Pro Val Pro Ala Thr Ala Glu Ala	
	20 25 30	
15	agc gta cag acc cgg cag tca tcg gga tac aag aac atc gtg tac ttt	322
	Ser Val Gln Thr Arg Gln Ser Ser Gly Tyr Lys Asn Ile Val Tyr Phe	
	35 40 45	
	acc aac tg gatatgtacgc ccggccgtca acccctccgc cctccgctcc	370
20	Thr Asn Trp	
	50	
	ctctctccac cgtttccggg ccagacgaac cactcaagac ttactttaat tactcactgg	430
	ccaatcgaac ag g ggt att tac ggc cgc aac tac cag cca gac cag ctg	479
25	Gly Ile Tyr Gly Arg Asn Tyr Gln Pro Asp Gln Leu	
	55 60	
	ccc gcg tct cag ctc acc cac gtg ctc tat tcc ttt gcc aac atc agg	527
	Pro Ala Ser Gln Leu Thr His Val Leu Tyr Ser Phe Ala Asn Ile Arg	
30	65 70 75	
	tcc aac ggc gag gta ttc ctc tcc gac acg tat gcg gac cta gag aag	575
	Ser Asn Gly Glu Val Phe Leu Ser Asp Thr Tyr Ala Asp Leu Glu Lys	
	80 85 90 95	
35		
40		
45		
50		
55		

EP 2 183 363 B1

	cac tat ccc aac gac t gtaagattaa ttatatccct cttccatcca cccatgtact	631
	His Tyr Pro Asn Asp	
	100	
5	cccaacccaa caaccagtca tccattttat ccacccccctt tcttttgtaat cttctctctt	691
	cccccttct gttccactga cagcacggct cccccggcag ca tgg aat gac gtc	745
	Ser Trp Asn Asp Val	
	105	
10	gga aac aac gtg tac ggc tgc gtc aag cag ctc ttc ctg ctc aag aag	793
	Gly Asn Asn Val Tyr Gly Cys Val Lys Gln Leu Phe Leu Leu Lys Lys	
	110 115 120	
15	gcg aac cgc cag ctc aag acg ctg ctc tcc atc ggc ggc tgg acc tac	841
	Ala Asn Arg Gln Leu Lys Thr Leu Leu Ser Ile Gly Gly Trp Thr Tyr	
	125 130 135	
	tcg gcc acc ttc ccc gcg gcg gcc agc aca gcc gag agc cga gcc ctc	889
	Ser Ala Thr Phe Pro Ala Ala Ala Ser Thr Ala Glu Ser Arg Ala Leu	
20	140 145 150	
	ttc gcg tcg tcg gcg gtc cgg ctc ctg gcc gac ctg ggc ttc gac ggc	937
	Phe Ala Ser Ser Ala Val Arg Leu Leu Ala Asp Leu Gly Phe Asp Gly	
	155 160 165	
25	ctc gac atc gac tgg gag tac ccg gcc aac gag cag gag gcg gcc aac	985
	Leu Asp Ile Asp Trp Glu Tyr Pro Ala Asn Glu Gln Glu Ala Ala Asn	
	170 175 180 185	
30	ttc gtc ctc ctg ctc aag gcg gtc cgc tcg gcc ctg gac gac tac gcg	1033
	Phe Val Leu Leu Leu Lys Ala Val Arg Ser Ala Leu Asp Asp Tyr Ala	
	190 195 200	
	gcc cag cac gcg ccc ggc tac cac ttc ctc ctc acc atc gcc tcg ccg	1081
	Ala Gln His Ala Pro Gly Tyr His Phe Leu Leu Thr Ile Ala Ser Pro	
35	205 210 215	
	gcc ggc ccg agc aac tac ggg cac ctg ccg ctg cgc gac atc gcc ggc	1129
	Ala Gly Pro Ser Asn Tyr Gly His Leu Pro Leu Arg Asp Ile Ala Gly	
	220 225 230	
40	gtg atc gac ttc ttc aac ttc atg ggc tac gac tac gcc ggc tcg tgg	1177
	Val Ile Asp Phe Phe Asn Phe Met Gly Tyr Asp Tyr Ala Gly Ser Trp	
	235 240 245	
45	agc acc gcg gcg gga cac cag gcc aac ctg tac ccg acc gcc gac gcg	1225
	Ser Thr Ala Ala Gly His Gln Ala Asn Leu Tyr Pro Thr Ala Asp Ala	
	250 255 260 265	
	ggc agg acg ccc ttc tcg acc gac aag gcc ctg tcc gac tac gtc gcc	1273
	Gly Arg Thr Pro Phe Ser Thr Asp Lys Ala Leu Ser Asp Tyr Val Ala	
50	270 275 280	
	gcc ggc gtc gac ccg gcc aag atc gtg ctc ggc atg ccc atc tac ggc	1321
	Ala Gly Val Asp Pro Ala Lys Ile Val Leu Gly Met Pro Ile Tyr Gly	
	285 290 295	
55	cgc tcc ttc gag gcc acc gac ggc ctg ggc aag ccc ttc acc ggc gtc	1369
	Arg Ser Phe Glu Ala Thr Asp Gly Leu Gly Lys Pro Phe Thr Gly Val	

EP 2 183 363 B1

	300	305	310	
5	ggc cag ggc agc tgg gag agc ggc gtg tgg gac tac aag gtg ctg ccc Gly Gln Gly Ser Trp Glu Ser Gly Val Trp Asp Tyr Lys Val Leu Pro 315 320 325	1417		
10	cgg gcc ggc gcc acc gtc cag tac gac gag gag gcc ggc gcg acc tac Arg Ala Gly Ala Thr Val Gln Tyr Asp Glu Glu Ala Gly Ala Thr Tyr 330 335 340 345	1465		
15	agc tac gac ccg gcc acg cgc gag ctc atc agc ttc gac acg gtc gac Ser Tyr Asp Pro Ala Thr Arg Glu Leu Ile Ser Phe Asp Thr Val Asp 350 355 360	1513		
20	atg gtc aag aag aag gtc gac tac gtc aag cag aag ggc ttc gcc ggc Met Val Lys Lys Lys Val Asp Tyr Val Lys Gln Lys Gly Phe Ala Gly 365 370 375	1561		
25	agc atg ttc tgg gag gcc tcc gcc gac cgc acc ggc gac cag agc ctc Ser Met Phe Trp Glu Ala Ser Ala Asp Arg Thr Gly Asp Gln Ser Leu 380 385 390	1609		
30	atc ggc gcc agc ttc ggc gcc ctc ggc ggc atc gac cag tcc cag aac Ile Gly Ala Ser Phe Gly Ala Leu Gly Gly Ile Asp Gln Ser Gln Asn 395 400 405	1657		
35	cag ctg tcg tac ccg gac agc aag tat gat aac ctc agg gcc ggc ttc Gln Leu Ser Tyr Pro Asp Ser Lys Tyr Asp Asn Leu Arg Ala Gly Phe 410 415 420 425	1705		
40	ccg tgaggcccta tgacctcccg agtcctgggc gagagagggg agctggagag Pro	1758		
45	agagagagag agagagagag agagggagcc tgaagccgta ctgattcggc aaaccatttt gaagcctgga catgtttacc ttcgttttct cttcttggat ggtttgcatt attattagcc gcttttcttt tttctctcca ttcggt	1818 1878 1904		
50	<210> 56 <211> 426 <212> PRT <213> Chrysosporium lucknowense			
55	<400> 56			

EP 2 183 363 B1

Met Val Thr Pro Ser Leu Lys Lys Ala Ala Leu Ala Ala Leu Ser Leu
1 5 10 15

5 Phe Pro Leu Leu Ser Leu Ala Ser Pro Val Pro Ala Thr Ala Glu Ala
20 25 30

10 Ser Val Gln Thr Arg Gln Ser Ser Gly Tyr Lys Asn Ile Val Tyr Phe
35 40 45

15 Thr Asn Trp Gly Ile Tyr Gly Arg Asn Tyr Gln Pro Asp Gln Leu Pro

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	50					55					60					
5	Ala 65	Ser	Gln	Leu	Thr	His 70	Val	Leu	Tyr	Ser	Phe 75	Ala	Asn	Ile	Arg	Ser 80
10	Asn	Gly	Glu	Val	Phe 85	Leu	Ser	Asp	Thr	Tyr 90	Ala	Asp	Leu	Glu	Lys 95	His
15	Tyr	Pro	Asn	Asp 100	Ser	Trp	Asn	Asp	Val 105	Gly	Asn	Asn	Val	Tyr 110	Gly	Cys
20	Val	Lys	Gln 115	Leu	Phe	Leu	Leu	Lys 120	Lys	Ala	Asn	Arg	Gln 125	Leu	Lys	Thr
25	Leu 130	Leu	Ser	Ile	Gly	Gly	Trp 135	Thr	Tyr	Ser	Ala	Thr 140	Phe	Pro	Ala	Ala
30	Ala 145	Ser	Thr	Ala	Glu	Ser 150	Arg	Ala	Leu	Phe	Ala 155	Ser	Ser	Ala	Val	Arg 160
35	Leu	Leu	Ala	Asp	Leu 165	Gly	Phe	Asp	Gly	Leu 170	Asp	Ile	Asp	Trp	Glu 175	Tyr
40	Pro	Ala	Asn	Glu 180	Gln	Glu	Ala	Ala	Asn 185	Phe	Val	Leu	Leu	Leu	Lys 190	Ala
45	Val	Arg	Ser 195	Ala	Leu	Asp	Asp	Tyr 200	Ala	Ala	Gln	His 205	Ala	Pro	Gly	Tyr
50	His	Phe 210	Leu	Leu	Thr	Ile	Ala 215	Ser	Pro	Ala	Gly	Pro 220	Ser	Asn	Tyr	Gly
55	His	Leu	Pro	Leu	Arg	Asp 230	Ile	Ala	Gly	Val	Ile 235	Asp	Phe	Phe	Asn	Phe 240
60	Met	Gly	Tyr	Asp	Tyr 245	Ala	Gly	Ser	Trp	Ser 250	Thr	Ala	Ala	Gly	His 255	Gln
65	Ala	Asn	Leu	Tyr 260	Pro	Thr	Ala	Asp	Ala 265	Gly	Arg	Thr	Pro	Phe 270	Ser	Thr
70	Asp	Lys	Ala 275	Leu	Ser	Asp	Tyr	Val 280	Ala	Ala	Gly	Val	Asp 285	Pro	Ala	Lys
75	Ile	Val	Leu	Gly	Met	Pro	Ile 295	Tyr	Gly	Arg	Ser	Phe 300	Glu	Ala	Thr	Asp

EP 2 183 363 B1

	Gly	Leu	Gly	Lys	Pro	Phe	Thr	Gly	Val	Gly	Gln	Gly	Ser	Trp	Glu	Ser	
	305					310					315					320	
5	Gly	Val	Trp	Asp	Tyr	Lys	Val	Leu	Pro	Arg	Ala	Gly	Ala	Thr	Val	Gln	
					325					330					335		
10	Tyr	Asp	Glu	Glu	Ala	Gly	Ala	Thr	Tyr	Ser	Tyr	Asp	Pro	Ala	Thr	Arg	
				340					345					350			
15	Glu	Leu	Ile	Ser	Phe	Asp	Thr	Val	Asp	Met	Val	Lys	Lys	Lys	Val	Asp	
			355					360					365				
20	Tyr	Val	Lys	Gln	Lys	Gly	Phe	Ala	Gly	Ser	Met	Phe	Trp	Glu	Ala	Ser	
		370					375						380				
25	Ala	Asp	Arg	Thr	Gly	Asp	Gln	Ser	Leu	Ile	Gly	Ala	Ser	Phe	Gly	Ala	
	385					390					395					400	
30	Leu	Gly	Gly	Ile	Asp	Gln	Ser	Gln	Asn	Gln	Leu	Ser	Tyr	Pro	Asp	Ser	
					405					410					415		
35	Lys	Tyr	Asp	Asn	Leu	Arg	Ala	Gly	Phe	Pro							
				420					425								
40	<210> 57																
	<211> 1278																
	<212> DNA																
	<213> Chrysosporium lucknowense																
45	<400> 57																
50	atggtgactc	cctctctgaa	gaaggcagct	ctggctgccc	tgtccctctt	cccgtcctc											60
55	tccctagctt	cgccggtgcc	agcaacagcc	gaagctagcg	tacagaccg	gcagtcacg											120
	ggatacaaga	acatcgtgta	ctttaccaac	tggggtattt	acggccgcaa	ctaccagcca											180
	gaccagctgc	ccgctctca	gtcaccac	gtgctctatt	cctttgcaa	catcaggtcc											240
	aacggcgagg	tattcctctc	cgacacgtat	gcggacctag	agaagcacta	tcccaacgac											300
	tcatggaatg	acgtcggaaa	caacgtgtac	ggctgcgtca	agcagctctt	cctgctcaag											360
	aaggcgaacc	gccagctcaa	gacgtgctc	tccatcggcg	gctggaccta	ctcggccacc											420
	ttccccgcgg	cggccagcac	agccgagagc	cgagccctct	tcgctcgtc	ggcggtcgg											480
	ctcctggccg	acctgggctt	cgacggcctc	gacatcgact	gggagtacc	ggccaacgag											540
	caggaggcgg	ccaacttcgt	cctcctgctc	aaggcggctc	gctcggccct	ggacgactac											600
	gcggcccagc	acgcgcccgg	ctaccacttc	ctcctcacca	tcgcctcgcc	ggccggcccg											660

EP 2 183 363 B1

	agcaactacg ggcacctgcc gctgcgcgac atcgccggcg tgatecgactt cttcaacttc	720
	atgggctacg actacgccgg ctctgtggagc accgcggcgg gacaccaggc caacctgtac	780
5	ccgaccgccg acgcgggcag gacgcccttc tcgaccgaca aggccctgtc cgactacgtc	840
	gccgcccggcg tcgaccggc caagatcgtg ctcggcgatgc ccatctacgg ccgctccttc	900
10	gaggccaccg acggcctggg caagcccttc accggcgctcg gccagggcag ctgggagagc	960
	ggcgtgtggg actacaaggt gctgccccgg gccggcgcca ccgtccagta cgacgaggag	1020
	gccggcgcgga cctacagcta cgaccgggcc acgcgcgagc tcatcagctt cgacacggtc	1080
15	gacatggtca agaagaaggt cgactacgtc aagcagaagg gcttcgccgg cagcatgttc	1140
	tgggaggcct ccgccgaccg caccggcgac cagagcctca tcggcgccag cttcggcgcc	1200
	ctcggcggca tcgaccagtc ccagaaccag ctgtcgtacc cggacagcaa gtatgataac	1260
20	ctcagggccg gcttcccg	1278

	<210> 58
	<211> 3200
25	<212> DNA
	<213> Chrysosporium lucknowense
	<220>
	<221> CDS
30	<222> (118)..(2838)
	<400> 58

35

40

45

50

55

EP 2 183 363 B1

	aggcccttcc cttgtgccta ttaccttacc cgaattgtta attatgtgtc taccaggcac	60
	ccaggagggc attgtactca ctcactctcc gagagaacat cgaatccgga aaacaga	117
5	atg cgg gcg aaa agg gcg gcc att gcc ctt ttc cta gct tgg ccg gcc Met Arg Ala Lys Arg Ala Ala Ile Ala Leu Phe Leu Ala Trp Pro Ala 1 5 10 15	165
10	ata aca cgg gcc gcg tcc ctc gtc tct ttt ccg gga aac acg gcc gcg Ile Thr Arg Ala Ala Ser Leu Val Ser Phe Pro Gly Asn Thr Ala Ala 20 25 30	213
15	att cca tcg tgg gaa ctg cag tct tct gtg gag gcc ggc act gac ctc Ile Pro Ser Trp Glu Leu Gln Ser Ser Val Glu Ala Gly Thr Asp Leu 35 40 45	261
20	gaa ggt ctt tca caa gtt ggg gtc gat aca aag tca tgg cac cac atc Glu Gly Leu Ser Gln Val Gly Val Asp Thr Lys Ser Trp His His Ile 50 55 60	309
	aaa aca tgg aga tgt acg ctc atg ggc tgc ctg atc gag gcc ggc gtc Lys Thr Trp Arg Cys Thr Leu Met Gly Cys Leu Ile Glu Ala Gly Val 65 70 75 80	357
25	tac aat gaa gat caa ctc ttc tac tcg gag aac ctg cgc aaa gtc aac Tyr Asn Glu Asp Gln Leu Phe Tyr Ser Glu Asn Leu Arg Lys Val Asn	405
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	85								90					95					
5	gag	acg	caa	ttc	ctg	gtg	cca	tgg	att	tat	cgc	aac	gag	ttc	tcc	ctg	453		
	Glu	Thr	Gln	Phe	Leu	Val	Pro	Trp	Ile	Tyr	Arg	Asn	Glu	Phe	Ser	Leu			
				100					105					110					
	cat	ccc	ggc	ccg	ggc	agg	cac	ttc	ttc	ctg	cag	aca	cac	ggc	atc	tct	501		
	His	Pro	Gly	Pro	Gly	Arg	His	Phe	Phe	Leu	Gln	Thr	His	Gly	Ile	Ser			
			115					120					125						
10	tcg	cga	gcc	gac	atc	ttt	ctg	aac	ggc	aaa	cga	gtg	gcc	aat	tct	tcg	549		
	Ser	Arg	Ala	Asp	Ile	Phe	Leu	Asn	Gly	Lys	Arg	Val	Ala	Asn	Ser	Ser			
			130				135					140							
15	gag	cag	gcg	ggt	tcg	tac	gtt	ggc	cgg	att	tac	gac	atc	acc	aaa	cag	597		
	Glu	Gln	Ala	Gly	Ser	Tyr	Val	Gly	Arg	Ile	Tyr	Asp	Ile	Thr	Lys	Gln			
						150					155					160			
	gtt	cac	aga	gaa	aat	gcg	ctg	gca	att	cag	gtg	tac	ccg	acc	aac	tac	645		
	Val	His	Arg	Glu	Asn	Ala	Leu	Ala	Ile	Gln	Val	Tyr	Pro	Thr	Asn	Tyr			
20					165					170					175				
	tac	tac	gac	ctg	gcg	ctc	gga	tgg	gtg	gat	tgg	aac	cca	tgg	ccg	gcc	693		
	Tyr	Tyr	Asp	Leu	Ala	Leu	Gly	Trp	Val	Asp	Trp	Asn	Pro	Trp	Pro	Ala			
				180					185				190						
25	gac	aac	ggc	acc	ggc	gtc	tgg	cgg	gac	gtc	gag	atc	aag	cag	acc	ggg	741		
	Asp	Asn	Gly	Thr	Gly	Val	Trp	Arg	Asp	Val	Glu	Ile	Lys	Gln	Thr	Gly			
			195					200					205						
	gcg	gtc	atg	ctg	gag	ccg	ttg	cgg	gtt	gtg	acg	cac	ttg	ggg	ccg	acg	789		
30	Ala	Val	Met	Leu	Glu	Pro	Leu	Arg	Val	Val	Thr	His	Leu	Gly	Pro	Thr			
			210				215					220							
	ctc	gga	gat	gag	cca	gca	aac	gtg	acc	ctc	aag	gcc	caa	gca	cat	aac	837		
	Leu	Gly	Asp	Glu	Pro	Ala	Asn	Val	Thr	Leu	Lys	Ala	Gln	Ala	His	Asn			
35						230					235					240			
	ctt	gaa	aac	tac	acc	gtc	acc	atc	acg	gcc	acc	ggc	cac	atc	gcc	ctg	885		
	Leu	Glu	Asn	Tyr	Thr	Val	Thr	Ile	Thr	Ala	Thr	Gly	His	Ile	Ala	Leu			
					245					250					255				
40	gac	ccc	cac	tct	gat	aac	caa	caa	cac	cat	cac	cac	cac	cac	cct	att	933		
	Asp	Pro	His	Ser	Asp	Asn	Gln	Gln	His	His	His	His	His	His	Pro	Ile			
				260					265					270					
	acc	tgg	cac	cga	acg	tac	acg	ctc	cca	ccc	ctt	tcc	acc	acc	gag	ctc	981		
45	Thr	Trp	His	Arg	Thr	Tyr	Thr	Leu	Pro	Pro	Leu	Ser	Thr	Thr	Glu	Leu			
			275					280					285						
	gcc	ctc	tcc	gcg	acc	att	gcc	gac	ccc	cag	atc	tgg	tgg	ccc	cgc	caa	1029		
	Ala	Leu	Ser	Ala	Thr	Ile	Ala	Asp	Pro	Gln	Ile	Trp	Trp	Pro	Arg	Gln			
50			290				295					300							
	tgg	ggt	ggt	caa	ccg	ctc	tac	acc	gcc	gcc	ttg	acc	gtc	acc	acg	cgc	1077		
	Trp	Gly	Gly	Gln	Pro	Leu	Tyr	Thr	Ala	Ala	Leu	Thr	Val	Thr	Thr	Arg			
			305			310					315					320			
55	ggc	acc	acc	acg	ccg	tcc	gac	cgg	tcc	acg	gcc	acc	ttc	ggc	ctg	cgc	1125		
	Gly	Thr	Thr	Thr	Pro	Ser	Asp	Arg	Ser	Thr	Ala	Thr	Phe	Gly	Leu	Arg			
					325					330					335				

EP 2 183 363 B1

	acc gtc gag gcg agg ctg aac ccg acc tac aac gac acc acc ttc cac	1173
	Thr Val Glu Ala Arg Leu Asn Pro Thr Tyr Asn Asp Thr Thr Phe His	
	340 345 350	
5	gtc aac tcg cgc ccc ttc cag gtc ctg ggc gcc ggc tac tcc ccc aac	1221
	Val Asn Ser Arg Pro Phe Gln Val Leu Gly Ala Gly Tyr Ser Pro Asn	
	355 360 365	
10	gtc ttc ctc cgc gcg ccg cca cgg acc gcc gcg gcc gac tac acg gcc	1269
	Val Phe Leu Arg Ala Pro Pro Arg Thr Ala Ala Ala Asp Tyr Thr Ala	
	370 375 380	
15	acg ctc cag tac gta ctc gac ctg ggc ctg aac acg atc cgg ctc gag	1317
	Thr Leu Gln Tyr Val Leu Asp Leu Gly Leu Asn Thr Ile Arg Leu Glu	
	385 390 395 400	
	ggc aag gac gag cac tcg gaa ctg tac gcg gcg gcg gac cgg ctg ggc	1365
	Gly Lys Asp Glu His Ser Glu Leu Tyr Ala Ala Ala Asp Arg Leu Gly	
	405 410 415	
20	gtc atg atc ctt gcc ggc tgg gag tgc tgc gac aag tgg gag gcg tgg	1413
	Val Met Ile Leu Ala Gly Trp Glu Cys Cys Asp Lys Trp Glu Ala Trp	
	420 425 430	
25	agc tac aac ggc gac ctg gcc atc gac ccg gtg ccg gtc tgg gac gac	1461
	Ser Tyr Asn Gly Asp Leu Ala Ile Asp Pro Val Pro Val Trp Asp Asp	
	435 440 445	
30	gcc gac tac gcc gtc gcg cgt gcc gcc atg gcc cac gag gcc cgc atg	1509
	Ala Asp Tyr Ala Val Ala Arg Ala Ala Met Ala His Glu Ala Arg Met	
	450 455 460	
35	ctc cag ccc cat ccc agc ctg ctg ggc ttc ctc atc ggc agc gac ttc	1557
	Leu Gln Pro His Pro Ser Leu Leu Gly Phe Leu Ile Gly Ser Asp Phe	
	465 470 475 480	
40	tgg ccc gac gac cgc gcc acc gcc gcc tac ctc gat gcc ctg gaa gcc	1605
	Trp Pro Asp Asp Arg Ala Thr Ala Ala Tyr Leu Asp Ala Leu Glu Ala	
	485 490 495	
45	gcc gac tgg gcc gcc ccc gtg ctg gcc tcg gcc tcg aag cgc ggg tac	1653
	Ala Asp Trp Ala Ala Pro Val Leu Ala Ser Ala Ser Lys Arg Gly Tyr	
	500 505 510	
50	ccc gag cgg acc ggt ccg gga ggg ctc aag atg gac ggg ccg tac gac	1701
	Pro Glu Arg Thr Gly Pro Gly Gly Leu Lys Met Asp Gly Pro Tyr Asp	
	515 520 525	
55	tgg gtg cca ccg ggg tac tgg tgg gac acg gag ccg acc ggg gag agg	1749
	Trp Val Pro Pro Gly Tyr Trp Trp Asp Thr Glu Pro Thr Gly Glu Arg	
	530 535 540	
50	ctc gga gcc gcg ttc ggt ttt ggg tcc gag ctg ggc gcg ggc gtg ggc	1797
	Leu Gly Ala Ala Phe Gly Phe Gly Ser Glu Leu Gly Ala Gly Val Gly	
	545 550 555 560	
55	acg ccg gag ctg ggg tcg ctg agg cgg ttc ctg agc gag ggg gac ctg	1845
	Thr Pro Glu Leu Gly Ser Leu Arg Arg Phe Leu Ser Glu Gly Asp Leu	
	565 570 575	

EP 2 183 363 B1

[illegible]

EP 2 183 363 B1

	Val	Thr	Val	His	Asn	Val	Trp	His	Gly	Ser	Gly	Val	Asp	Met	Thr	Leu	
				820					825					830			
5	gag	aac	aaa	tct	aag	gtg	cca	gcg	ttc	ttt	gtc	agc	ttg	acg	ctg	gtg	2661
	Glu	Asn	Lys	Ser	Lys	Val	Pro	Ala	Phe	Phe	Val	Ser	Leu	Thr	Leu	Val	
			835					840					845				
10	gat	gta	cac	ggg	aac	gag	gtc	ctg	ccg	gtg	acg	tgg	gag	gac	aac	tat	2709
	Asp	Val	His	Gly	Asn	Glu	Val	Leu	Pro	Val	Thr	Trp	Glu	Asp	Asn	Tyr	
		850					855					860					
15	gtc	acc	cta	tgg	ccg	ggt	gag	aaa	ttg	aca	ctt	cgg	gga	agg	tca	atg	2757
	Val	Thr	Leu	Trp	Pro	Gly	Glu	Lys	Leu	Thr	Leu	Arg	Gly	Arg	Ser	Met	
	865					870					875				880		
20	ggg	gta	gat	gca	aag	ccg	tgg	gag	gtc	att	gtg	aga	gga	aag	aac	gtt	2805
	Gly	Val	Asp	Ala	Lys	Pro	Trp	Glu	Val	Ile	Val	Arg	Gly	Lys	Asn	Val	
					885				890						895		
	gaa	gca	aag	aga	gtc	cgt	cac	ccg	tgg	cat	gta	tgagagaaga	ccaacattac				2858
	Glu	Ala	Lys	Arg	Val	Arg	His	Arg	Trp	His	Val						
				900					905								
25	atggatagcc	ttaatggtat	agatcctacc	gaatgaaatc	actcattagc	gaggaagtac											2918
	tttgtgagtt	gggtgagcat	attccctaaa	ttcattctac	tggggggttgc	attagaggca											2978
	cctcacgtgg	aacaactggt	aggattcaat	ttcgcagcaa	gaaccgtagt	cccactcggt											3038
	ttcaaatatc	catgtgagac	atgtgatctg	tctgtataac	ggggcggcta	cttgcccact											3098
30	tcaccaaca	cggaagacct	agggttggtg	ggccaccata	ttgcctttgc	atgactat											3158
	cacaccccaa	cgtgaaacga	tgctctaaaa	ccaacgaaaa	aa												3200
35	<210>	59															
	<211>	907															
	<212>	PRT															
	<213>	Chrysosporium lucknowense															
40	<400>	59															
	Met	Arg	Ala	Lys	Arg	Ala	Ala	Ile	Ala	Leu	Phe	Leu	Ala	Trp	Pro	Ala	
	1			5						10					15		
45	Ile	Thr	Arg	Ala	Ala	Ser	Leu	Val	Ser	Phe	Pro	Gly	Asn	Thr	Ala	Ala	
				20					25					30			
50	Ile	Pro	Ser	Trp	Glu	Leu	Gln	Ser	Ser	Val	Glu	Ala	Gly	Thr	Asp	Leu	
			35					40					45				
	Glu	Gly	Leu	Ser	Gln	Val	Gly	Val	Asp	Thr	Lys	Ser	Trp	His	His	Ile	
		50					55					60					
55	Lys	Thr	Trp	Arg	Cys	Thr	Leu	Met	Gly	Cys	Leu	Ile	Glu	Ala	Gly	Val	
	65					70					75					80	

EP 2 183 363 B1

	Tyr	Asn	Glu	Asp	Gln	Leu	Phe	Tyr	Ser	Glu	Asn	Leu	Arg	Lys	Val	Asn	
					85					90					95		
5	Glu	Thr	Gln	Phe	Leu	Val	Pro	Trp	Ile	Tyr	Arg	Asn	Glu	Phe	Ser	Leu	
				100					105					110			
10	His	Pro	Gly	Pro	Gly	Arg	His	Phe	Phe	Leu	Gln	Thr	His	Gly	Ile	Ser	
			115					120					125				
15	Ser	Arg	Ala	Asp	Ile	Phe	Leu	Asn	Gly	Lys	Arg	Val	Ala	Asn	Ser	Ser	
		130					135					140					
20	Glu	Gln	Ala	Gly	Ser	Tyr	Val	Gly	Arg	Ile	Tyr	Asp	Ile	Thr	Lys	Gln	
	145					150					155					160	
25	Val	His	Arg	Glu	Asn	Ala	Leu	Ala	Ile	Gln	Val	Tyr	Pro	Thr	Asn	Tyr	
					165					170					175		
30	Tyr	Tyr	Asp	Leu	Ala	Leu	Gly	Trp	Val	Asp	Trp	Asn	Pro	Trp	Pro	Ala	
				180					185					190			
35	Asp	Asn	Gly	Thr	Gly	Val	Trp	Arg	Asp	Val	Glu	Ile	Lys	Gln	Thr	Gly	
			195					200					205				
40	Ala	Val	Met	Leu	Glu	Pro	Leu	Arg	Val	Val	Thr	His	Leu	Gly	Pro	Thr	
		210					215					220					
45	Leu	Gly	Asp	Glu	Pro	Ala	Asn	Val	Thr	Leu	Lys	Ala	Gln	Ala	His	Asn	
	225					230					235					240	
50	Leu	Glu	Asn	Tyr	Thr	Val	Thr	Ile	Thr	Ala	Thr	Gly	His	Ile	Ala	Leu	
					245					250					255		
55	Asp	Pro	His	Ser	Asp	Asn	Gln	Gln	His	His	His	His	His	His	Pro	Ile	
				260					265					270			
60	Thr	Trp	His	Arg	Thr	Tyr	Thr	Leu	Pro	Pro	Leu	Ser	Thr	Thr	Glu	Leu	
			275					280					285				
65	Ala	Leu	Ser	Ala	Thr	Ile	Ala	Asp	Pro	Gln	Ile	Trp	Trp	Pro	Arg	Gln	
		290					295					300					
70	Trp	Gly	Gly	Gln	Pro	Leu	Tyr	Thr	Ala	Ala	Leu	Thr	Val	Thr	Thr	Arg	
	305					310					315					320	

EP 2 183 363 B1

	Gly	Thr	Thr	Thr	Pro	Ser	Asp	Arg	Ser	Thr	Ala	Thr	Phe	Gly	Leu	Arg	
					325					330					335		
5	Thr	Val	Glu	Ala	Arg	Leu	Asn	Pro	Thr	Tyr	Asn	Asp	Thr	Thr	Phe	His	
				340					345					350			
10	Val	Asn	Ser	Arg	Pro	Phe	Gln	Val	Leu	Gly	Ala	Gly	Tyr	Ser	Pro	Asn	
			355					360					365				
15	Val	Phe	Leu	Arg	Ala	Pro	Pro	Arg	Thr	Ala	Ala	Ala	Asp	Tyr	Thr	Ala	
		370					375					380					
20	Thr	Leu	Gln	Tyr	Val	Leu	Asp	Leu	Gly	Leu	Asn	Thr	Ile	Arg	Leu	Glu	
	385					390					395					400	
25	Gly	Lys	Asp	Glu	His	Ser	Glu	Leu	Tyr	Ala	Ala	Ala	Asp	Arg	Leu	Gly	
					405					410					415		
30	Val	Met	Ile	Leu	Ala	Gly	Trp	Glu	Cys	Cys	Asp	Lys	Trp	Glu	Ala	Trp	
				420					425					430			
35	Ser	Tyr	Asn	Gly	Asp	Leu	Ala	Ile	Asp	Pro	Val	Pro	Val	Trp	Asp	Asp	
			435					440					445				
40	Ala	Asp	Tyr	Ala	Val	Ala	Arg	Ala	Ala	Met	Ala	His	Glu	Ala	Arg	Met	
		450					455					460					
45	Leu	Gln	Pro	His	Pro	Ser	Leu	Leu	Gly	Phe	Leu	Ile	Gly	Ser	Asp	Phe	
	465					470					475					480	
50	Trp	Pro	Asp	Asp	Arg	Ala	Thr	Ala	Ala	Tyr	Leu	Asp	Ala	Leu	Glu	Ala	
					485					490					495		
55	Ala	Asp	Trp	Ala	Ala	Pro	Val	Leu	Ala	Ser	Ala	Ser	Lys	Arg	Gly	Tyr	
				500					505					510			
60	Pro	Glu	Arg	Thr	Gly	Pro	Gly	Gly	Leu	Lys	Met	Asp	Gly	Pro	Tyr	Asp	
			515					520					525				
65	Trp	Val	Pro	Pro	Gly	Tyr	Trp	Trp	Asp	Thr	Glu	Pro	Thr	Gly	Glu	Arg	
		530					535					540					
70	Leu	Gly	Ala	Ala	Phe	Gly	Phe	Gly	Ser	Glu	Leu	Gly	Ala	Gly	Val	Gly	
	545					550					555					560	

EP 2 183 363 B1

	Thr	Pro	Glu	Leu	Gly	Ser	Leu	Arg	Arg	Phe	Leu	Ser	Glu	Gly	Asp	Leu	
					565					570					575		
5	Glu	Ala	Leu	Trp	Lys	Arg	Pro	Asn	Val	Ser	Leu	Phe	His	Met	Ser	Arg	
				580					585					590			
	Glu	Thr	Ser	Gln	Phe	Thr	Thr	Arg	Val	Ile	Tyr	Asn	Ala	Gly	Leu	Trp	
10			595					600					605				
	Asn	Arg	Trp	Gly	Ala	Pro	Ala	Ser	Leu	Glu	Asp	Tyr	Leu	Met	Lys	Ala	
		610					615					620					
15	Gln	Leu	Met	Asp	Tyr	Glu	Ala	Thr	Arg	Ala	Gln	Phe	Asp	Ala	Tyr	Thr	
	625					630					635					640	
	Ala	Met	Trp	Asn	Ala	Glu	Arg	Pro	Ala	Thr	Gly	Leu	Ile	Tyr	Trp	Met	
20				645						650					655		
	Leu	Asn	Asn	Ala	Trp	Pro	Ser	Leu	His	Trp	Asn	Leu	Trp	Asp	Tyr	Tyr	
25				660					665					670			
	Met	Arg	Pro	Ala	Gly	Ser	Tyr	Phe	Gly	Ala	Lys	Ala	Gly	Ser	Arg	Val	
			675					680					685				
30	Glu	Asn	Val	Val	Phe	Asp	Tyr	Val	Arg	Arg	Ala	Val	Trp	Leu	Val	Asn	
		690					695					700					
	Arg	Ser	Leu	Asp	Lys	Ser	Gly	Thr	Arg	Arg	Val	Gln	Val	Asp	Val	Met	
35	705					710					715					720	
	Asp	Lys	His	Gly	Lys	Val	Leu	Tyr	Lys	Asp	Thr	Thr	Val	Thr	Thr	Thr	
40				725						730					735		
	Val	Pro	Asn	Thr	Ser	Lys	Asp	Ile	Leu	Ser	Leu	Ala	Gly	Pro	Leu	Gly	
				740					745					750			
45	Asn	Ile	Thr	Asp	Val	Val	Phe	Leu	Arg	Leu	Val	Leu	Tyr	Asp	Ala	Gln	
			755					760					765				
	Arg	Gln	Gly	His	Ala	Leu	Ser	Arg	Asn	Val	Tyr	Trp	Val	Ala	Lys	Glu	
50			770				775					780					
	Pro	Asp	Val	Leu	Asp	Trp	Asp	Glu	Ser	Asp	Trp	Tyr	Phe	Thr	Pro	Val	
	785					790					795					800	
55	Thr	Ser	Tyr	Ser	Asp	Tyr	Thr	Ala	Leu	Asn	Asn	Met	Ala	Gln	Ala	Asp	

EP 2 183 363 B1

805

810

815

5 Val Thr Val His Asn Val Trp His Gly Ser Gly Val Asp Met Thr Leu
820 825 830

10 Glu Asn Lys Ser Lys Val Pro Ala Phe Phe Val Ser Leu Thr Leu Val
835 840 845

15 Asp Val His Gly Asn Glu Val Leu Pro Val Thr Trp Glu Asp Asn Tyr
850 855 860

Val Thr Leu Trp Pro Gly Glu Lys Leu Thr Leu Arg Gly Arg Ser Met
865 870 875 880

20 Gly Val Asp Ala Lys Pro Trp Glu Val Ile Val Arg Gly Lys Asn Val
885 890 895

25 Glu Ala Lys Arg Val Arg His Arg Trp His Val
900 905

<210> 60

<211> 2721

<212> DNA

30 <213> Chrysosporium lucknowense

<400> 60

35

40

45

50

55

EP 2 183 363 B1

	atgcggggcga aaagggcggc cattgccctt ttcctagctt ggccggccat aacacggggc	60
	gcgtccctcg tctcttttcc gggaaacacg gccgcgattc catcgtggga actgcagtct	120
5	tctgtggagg ccggcactga cctcgaaggt ctttcacaag ttgggggtcga tacaaagtca	180
	tggcaccaca tcaaaacatg gagatgtacg ctcatgggct gcctgatcga ggccggcgctc	240
10	tacaatgaag atcaactctt ctactcggag aacctgcgca aagtcaacga gacgcaattc	300
	ctgggtgccat ggatttatcg caacgagttc tccctgcatc ccggcccggg caggcacttc	360
	ttcctgcaga cacacggcat ctcttcgcga gccgacatct ttctgaacgg caaacgagtg	420
15	gccaattctt cggagcaggc gggttcgtac gttggccgga ttacgacat caccaaacag	480
	gttcacagag aaaatgcgct ggcaattcag gtgtaccgga ccaactacta ctacgacctg	540
20	gcgctcggat ggggtgattg gaacccatgg ccggccgaca acggcaccgg cgtctggcgg	600
	gacgtcgaga tcaagcagac cggggcggtc atgctggagc cgttgcggggt tgtgacgcac	660
	ttggggccga cgctcggaga tgagccagca aacgtgacct tcaaggccca agcacataac	720
25	cttgaaaact acaccgtcac catcacggcc accggccaca tcgccctgga cccccactct	780
	gataaccaac aacaccatca ccaccaccac cctattacct ggcaccgaac gtacacgctc	840
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	ccaccccttt ccaccaccga gctcgccctc tccgcgacca ttgcgcaccc ccagatctgg	900
	tggccccgcc aatgggggtgg tcaaccgctc tacaccgccg ccttgaccgt caccacgcgc	960
5	ggcaccacca cgccgtccga cgggtccacg gccaccttcg gcctgcgcac cgtcgaggcg	1020
	aggctgaacc cgacctacaa cgacaccacc ttccacgtca actcgcgccc cttccaggtc	1080
	ctggggcgccg gctactcccc caacgtcttc ctccgcgcgc cgccacggac cgccgcggcc	1140
10	gactacacgg ccacgctcca gtacgtactc gacctgggcc tgaacacgat ccggctcgag	1200
	ggcaaggacg agcactcgga actgtacgcg gcggcggacc ggctgggctg catgatcctt	1260
	gccggctggg agtgctgcga caagtgggag gcgtggagct acaacggcga cctggccatc	1320
15	gacccgggtgc cggctctggga cgacgccgac tacgccgtcg cgcgtgccgc catggcccac	1380
	gaggccccga tgctccagcc ccattccagc ctgctgggct tcctcatcgg cagcgacttc	1440
	tggccccgacg accgcgccac cgccgcctac ctcgatgcc tggaaagccgc cgactgggcc	1500
20	gcccccgctgc tggcctcggc ctccaagcgc gggtagcccg agcggaccgg tccgggaggg	1560
	ctcaagatgg acgggcccga cgactgggtg ccaccggggt actggtggga cacggagccg	1620
25	accgggggaga ggctcggagc cgcgttcggg tttgggtccg agctgggcgc gggcgtgggc	1680
	acgccggagc tggggtcgct gaggcggttc ctgagcgagg gggacctgga ggcctgtg	1740
	aagcggccga acgtgagcct attccacatg tcgagggaga cctcccaatt tacgacgagg	1800
30	gtgatctaca acgccgggct ctggaaccgg tggggcgccc ccgcgagcct ggaagactac	1860
	ctgatgaagg cgcagctgat ggactacgag gccacgcggg ctgagttcga cgcctacacg	1920
	gccatgtgga acgccgagag gccggccacg ggctgatat actggatgct caacaacgcc	1980
35	tggccgagcc tgcaactggaa cctgtgggac tactacatgc gtcccgccgg tagttacttt	2040
	ggggcaaagg ccggcagcag ggtggaaaat gtcgtgtttg actacgtcag gagggcggtc	2100
40	tggctggtca accggtccct ggacaagagc gggacgagga gagtgcagggt ggacgtcatg	2160
	gataagcatg gtaaagtcct atacaaggac acgaccgtca ccacgactgt tccaaacacg	2220
	agcaaggaca ttctttcact agcgggacct ctgggcaaca tcacgcagct ggtgtttttg	2280
45	cggctggttc tctacgatgc acaacggcaa gggcatgcgc tgagtcgaaa cgtgtactgg	2340
	gttgccaagg agccagatgt gctggactgg gacgagtcgg attggtactt cacgccggtg	2400
	accagctact cggactacac tgcaactaac aacatggctc aggcagacgt cacggttcac	2460
50	aatgtctggc acggctcagg ggttgacatg acgctggaga acaaactaa ggtgccagcg	2520
	ttctttgtca gcttgacgct ggtggatgta cacgggaacg aggtcctgcc ggtgacgtgg	2580
55	gaggacaact atgtcaccct atggccgggt gagaaattga cacttcgggg aaggtcaatg	2640
	ggggtagatg caaagccgtg ggaggtcatt gtgagaggaa agaacgttga agcaaagaga	2700

gtccgtcacc ggtggcatgt a

2721

<210> 61

<211> 1275

<212> DNA

<213> Chrysosporium lucknowense

<220>

<221> CDS

<222> (180)..(668)

<220>

<221> CDS

<222> (776)..(1225)

<400> 61

20	gtataattct acgtcatccc ctccgttccc ttcccagctc ccctctccta ctgctgagtg	60

	atgttaataa tagtacgagt cgtggaattt gtggatactg atccaggata aagtctcttc	120

	cattcggccc gacctcctaa aactcccgac aaagctaaac cgccccgtgt gacgacaag	179

25	atg aag ctc ctg ggc aaa ctc tcg gcg gca ctc gcc ctc gcg ggc agc	227
	Met Lys Leu Leu Gly Lys Leu Ser Ala Ala Leu Ala Leu Ala Gly Ser	
	1 5 10 15	

30	agg ctg gct gcc gcg cac ccg gtc ttc gac gag ctg atg cgg ccg acg	275
	Arg Leu Ala Ala Ala His Pro Val Phe Asp Glu Leu Met Arg Pro Thr	
	20 25 30	

35	gcg ccg ctg gtg cgc ccg cgg gcg gcc ctg cag cag gtg acc aac ttt	323
	Ala Pro Leu Val Arg Pro Arg Ala Ala Leu Gln Gln Val Thr Asn Phe	
	35 40 45	

	ggc agc aac ccg tcc aac acg aag atg ttc atc tac gtg ccc gac aag	371
	Gly Ser Asn Pro Ser Asn Thr Lys Met Phe Ile Tyr Val Pro Asp Lys	
	50 55 60	

40	ctg gcc ccc aac ccg ccc atc ata gtg gcc atc cac tac tgc acc ggc	419
	Leu Ala Pro Asn Pro Pro Ile Ile Val Ala Ala Ile His Tyr Cys Thr Gly	
	65 70 75 80	

45	acc gcc cag gcc tac tac tcg ggc tcc cct tac gcc cgc ctc gcc gac	467
	Thr Ala Gln Ala Tyr Tyr Ser Gly Ser Pro Tyr Ala Arg Leu Ala Asp	
	85 90 95	

50	cag aag ggc ttc atc gtc atc tac ccg gag tcc ccc tac agc ggc acc	515
	Gln Lys Gly Phe Ile Val Ile Tyr Pro Glu Ser Pro Tyr Ser Gly Thr	
	100 105 110	

	tgt tgg gac gtc tcg tcg cgc gcc gcc ctg acc cac aac ggc ggc ggc	563
	Cys Trp Asp Val Ser Ser Arg Ala Ala Leu Thr His Asn Gly Gly Gly	
	115 120 125	

55	gac agc aac tcg atc gcc aac atg gtc acc tac acc ctc gaa aag tac	611
	Asp Ser Asn Ser Ile Ala Asn Met Val Thr Tyr Thr Leu Glu Lys Tyr	
	130 135 140	

EP 2 183 363 B1

	aat ggc gac gcc agc aag gtc ttt gtc acc ggc tcc tcg tcc ggc gcc	659
	Asn Gly Asp Ala Ser Lys Val Phe Val Thr Gly Ser Ser Ser Gly Ala	
	145 150 155 160	
5	atg atg acg gtgcgttcta attacccctc ccactccttt tttttttttt	708
	Met Met Thr	
10	tttgttttcc ttccttttct ttttcccagg agcgtcttgg ttgattgact gacttgttca	768
	aatccag aac gtg atg gcc gcc gcg tac ccg gaa ctg ttc gcg gca gga	817
	Asn Val Met Ala Ala Ala Tyr Pro Glu Leu Phe Ala Ala Gly	
	165 170 175	
15	atc gcc tac tcg ggc gtg ccc gcc ggc tgc ttc tac agc cag tcc gga	865
	Ile Ala Tyr Ser Gly Val Pro Ala Gly Cys Phe Tyr Ser Gln Ser Gly	
	180 185 190	
20	ggc acc aac gcg agg aac agc tcg tgc gcc aac ggg cag atc aac tcg	913
	Gly Thr Asn Ala Arg Asn Ser Ser Cys Ala Asn Gly Gln Ile Asn Ser	
	195 200 205	
25	acg ccc cag gtg tgg gcc aag atg gtc ttc gac atg tac ccg gaa tac	961
	Thr Pro Gln Val Trp Ala Lys Met Val Phe Asp Met Tyr Pro Glu Tyr	
	210 215 220 225	
30	gac ggc ccg cgc ccc aag atg cag atc tac cac ggc tcg gcc gac ggc	1009
	Asp Gly Pro Arg Pro Lys Met Gln Ile Tyr His Gly Ser Ala Asp Gly	
	230 235 240	
35	acg ctc aga ccc agc aac tac aac gag acc atc aag cag tgg tgc ggc	1057
	Thr Leu Arg Pro Ser Asn Tyr Asn Glu Thr Ile Lys Gln Trp Cys Gly	
	245 250 255	
40	gtc ttc ggc ttc gac tac acc cgc ccc gac acc acc cag gcc aac tcc	1105
	Val Phe Gly Phe Asp Tyr Thr Arg Pro Asp Thr Thr Gln Ala Asn Ser	
	260 265 270	
45	ccg cag gcc ggc tac acc acc tac acc tgg ggc gag cag cag ctc gtc	1153
	Pro Gln Ala Gly Tyr Thr Thr Tyr Thr Trp Gly Glu Gln Gln Leu Val	
	275 280 285	
50	ggc atc tac gcc cag ggc gtc gga cac acg gtc ccc atc cgc ggc agc	1201
	Gly Ile Tyr Ala Gln Gly Val Gly His Thr Val Pro Ile Arg Gly Ser	
	290 295 300 305	
55	gac gac atg gcc ttc ttt ggc ctg tgatgatgag aggttcggtg ggctgggaac	1255
	Asp Asp Met Ala Phe Phe Gly Leu	
	310	
60	cagagcggaa gtacggttta	1275
	<210> 62	
	<211> 313	
	<212> PRT	
	<213> Chrysosporium lucknowense	
	<400> 62	

EP 2 183 363 B1

Met Lys Leu Leu Gly Lys Leu Ser Ala Ala Leu Ala Leu Ala Gly Ser

5

10

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	1				5					10					15	
5	Arg	Leu	Ala	Ala	Ala	His	Pro	Val	Phe	Asp	Glu	Leu	Met	Arg	Pro	Thr
				20					25					30		
10	Ala	Pro	Leu	Val	Arg	Pro	Arg	Ala	Ala	Leu	Gln	Gln	Val	Thr	Asn	Phe
			35					40					45			
15	Gly	Ser	Asn	Pro	Ser	Asn	Thr	Lys	Met	Phe	Ile	Tyr	Val	Pro	Asp	Lys
		50					55					60				
20	Leu	Ala	Pro	Asn	Pro	Pro	Ile	Ile	Val	Ala	Ile	His	Tyr	Cys	Thr	Gly
	65					70					75					80
25	Thr	Ala	Gln	Ala	Tyr	Tyr	Ser	Gly	Ser	Pro	Tyr	Ala	Arg	Leu	Ala	Asp
					85					90					95	
30	Gln	Lys	Gly	Phe	Ile	Val	Ile	Tyr	Pro	Glu	Ser	Pro	Tyr	Ser	Gly	Thr
				100					105					110		
35	Cys	Trp	Asp	Val	Ser	Ser	Arg	Ala	Ala	Leu	Thr	His	Asn	Gly	Gly	Gly
			115					120					125			
40	Asp	Ser	Asn	Ser	Ile	Ala	Asn	Met	Val	Thr	Tyr	Thr	Leu	Glu	Lys	Tyr
		130					135					140				
45	Asn	Gly	Asp	Ala	Ser	Lys	Val	Phe	Val	Thr	Gly	Ser	Ser	Ser	Gly	Ala
	145					150					155					160
50	Met	Met	Thr	Asn	Val	Met	Ala	Ala	Ala	Tyr	Pro	Glu	Leu	Phe	Ala	Ala
				165						170					175	
55	Gly	Ile	Ala	Tyr	Ser	Gly	Val	Pro	Ala	Gly	Cys	Phe	Tyr	Ser	Gln	Ser
				180					185					190		
60	Gly	Gly	Thr	Asn	Ala	Arg	Asn	Ser	Ser	Cys	Ala	Asn	Gly	Gln	Ile	Asn
			195					200					205			
65	Ser	Thr	Pro	Gln	Val	Trp	Ala	Lys	Met	Val	Phe	Asp	Met	Tyr	Pro	Glu
		210					215					220				
70	Tyr	Asp	Gly	Pro	Arg	Pro	Lys	Met	Gln	Ile	Tyr	His	Gly	Ser	Ala	Asp
	225					230					235					240
75	Gly	Thr	Leu	Arg	Pro	Ser	Asn	Tyr	Asn	Glu	Thr	Ile	Lys	Gln	Trp	Cys
				245						250					255	

EP 2 183 363 B1

	Gly	Val	Phe	Gly	Phe	Asp	Tyr	Thr	Arg	Pro	Asp	Thr	Thr	Gln	Ala	Asn	
				260					265					270			
5	Ser	Pro	Gln	Ala	Gly	Tyr	Thr	Thr	Tyr	Thr	Trp	Gly	Glu	Gln	Gln	Leu	
			275					280					285				
10	Val	Gly	Ile	Tyr	Ala	Gln	Gly	Val	Gly	His	Thr	Val	Pro	Ile	Arg	Gly	
		290					295					300					
15	Ser	Asp	Asp	Met	Ala	Phe	Phe	Gly	Leu								
	305					310											
	<210> 63																
	<211> 939																
	<212> DNA																
	<213> Chrysosporium lucknowense																
20	<400> 63																
	atgaagctcc	tgggcaaact	ctcggcggca	ctcgccctcg	cgggcagcag	gctggctgcc											60
25	gcgcaccccg	tcttcgacga	gctgatgcgg	ccgacggcgc	cgctggtgcg	cccgcgggcg											120
	gccctgcagc	aggtgaccaa	ctttggcagc	aaccctcca	acacgaagat	gttcatctac											180
	gtgcccgcga	agctggcccc	caaccggccc	atcatagtgg	ccatccacta	ctgcaccggc											240
30	accgcccagg	cctactactc	gggctcccct	tacgcccgcc	tcgcccacca	gaagggcttc											300
	atcgatcatc	accgaggatc	cccctacagc	ggcacctgtt	gggacgtctc	gtcgcgcgcc											360
35	gccctgaccc	acaacggcgg	cggcgacagc	aactcgatcg	ccaacatggg	cacctacacc											420
	ctcgaaaagt	acaatggcga	cgccagcaag	gtctttgtca	ccggctcctc	gtccggcgcc											480
	atgatgacga	acgtgatggc	cgccgcgtac	ccggaactgt	tcgcggcagg	aatcgccctac											540
40	tcggggcgtg	ccgccggctg	cttctacagc	cagtccggag	gcaccaacgc	gaggaacagc											600
	tcgtgcgcca	acgggcagat	caactcgacg	cccaggtgt	gggccaagat	ggctcttcgac											660
45	atgtaccccg	aatacgacgg	cccgcgcccc	aagatgcaga	tctaccacgg	ctcggccgac											720
	ggcacgctca	gaccagcaa	ctacaacgag	accatcaagc	agtgggtgcg	cgtcttcggc											780
	ttcgactaca	cccgccccga	caccaccag	gccaaactccc	cgcaggcccg	ctacaccacc											840
50	tacacctggg	gcgagcagca	gctcgtcggc	atctacgccc	agggcgctcg	acacacggtc											900
	cccatccgcg	gcagcgacga	catggccttc	tttggcctg													939
	<210> 64																
55	<211> 1182																
	<212> DNA																
	<213> Chrysosporium lucknowense																

EP 2 183 363 B1

<220>
<221> CDS
<222> (162)..(621)

5 <220>
<221> CDS
<222> (793)..(1169)

10 <400> 64

atataaataa gccgggaggg catcaggaca tgacctctac ctctctgtaa tcagacgctg	60
gtggcttcct cccacatatc ggggttaata tttattagct cgtcattgtc tctgcgcgcc	120
ttgccgcaaa acatcacccg tatattgcgt atcagcaaac a atg atc tcg gtt cct	176
	Met Ile Ser Val Pro
	1 5
gct ctc gct ctg gcc ctt ctg gcc gcc gtc cag gtc gtc gag tct gcc	224
Ala Leu Ala Leu Ala Leu Leu Ala Ala Val Gln Val Val Glu Ser Ala	
	10 15 20
tcg gct ggc tgt ggc aag gcg ccc cct tcc tcg ggc acc aag tcg atg	272
Ser Ala Gly Cys Gly Lys Ala Pro Pro Ser Ser Gly Thr Lys Ser Met	
	25 30 35
acg gtc aac ggc aag cag gcg cag tac att ctc cag ctg ccc aac aac	320
Thr Val Asn Gly Lys Gln Arg Gln Tyr Ile Leu Gln Leu Pro Asn Asn	
	40 45 50
tac gac gcc aac aag gcc cac agg gtg gtg atc ggg tac cac tgg cgc	368
Tyr Asp Ala Asn Lys Ala His Arg Val Val Ile Gly Tyr His Trp Arg	
	55 60 65
gac gga tcc atg aac gac gtg gcc aac ggc ggc ttc tac gat ctg cgg	416
Asp Gly Ser Met Asn Asp Val Ala Asn Gly Gly Phe Tyr Asp Leu Arg	
	70 75 80 85
tcc cgg gcg ggc gac agc acc atc ttc gtt gcc ccc aac ggc ctc aat	464
Ser Arg Ala Gly Asp Ser Thr Ile Phe Val Ala Pro Asn Gly Leu Asn	
	90 95 100
gcc gga tgg gcc aac gtg ggc ggc gag gac atc acc ttt acg gac cag	512
Ala Gly Trp Ala Asn Val Gly Gly Glu Asp Ile Thr Phe Thr Asp Gln	
	105 110 115
atc gta gac atg ctc aag aac gac ctc tgc gtg gac gag acc cag ttc	560
Ile Val Asp Met Leu Lys Asn Asp Leu Cys Val Asp Glu Thr Gln Phe	
	120 125 130
ttt gct acg ggc tgg agc tat ggc ggt gcc atg agc cat agc gtg gct	608
Phe Ala Thr Gly Trp Ser Tyr Gly Gly Ala Met Ser His Ser Val Ala	
	135 140 145
tgt tct cgg cca g gtaaatattga ttttgtcatc ctgtcttttt tcttcttctt	661
Cys Ser Arg Pro	
	150
cttctcttttc ttccctctgg cattcttctc ttctcttctc ttctcttctc ctcttttatt	721

EP 2 183 363 B1

	tccccgtcat ttccccccac aaagaaacac ctggacaagc caatgacatc cattaatata	781
5	tataaatgca g ac gtc ttc aag gcc gtc gcg gtc atc gcc ggg gcc cag Asp Val Phe Lys Ala Val Ala Val Ile Ala Gly Ala Gln 155 160 165	830
10	ctg tcc ggc tgc gcc ggc ggc acg acg ccc gtg gcg tac cta ggc atc Leu Ser Gly Cys Ala Gly Gly Thr Thr Pro Val Ala Tyr Leu Gly Ile 170 175 180	878
	cac gga gcc gcc gac aac gtc ctg ccc atc gac ctc ggc cgc cag ctg His Gly Ala Ala Asp Asn Val Leu Pro Ile Asp Leu Gly Arg Gln Leu 185 190 195	926
15	cgc gac aag tgg ctg cag acc aac ggc tgc aac tac cag ggc gcc cag Arg Asp Lys Trp Leu Gln Thr Asn Gly Cys Asn Tyr Gln Gly Ala Gln 200 205 210	974
20	gac ccc gcg ccg ggc cag cag gcc cac atc aag acc acc tac agc tgc Asp Pro Ala Pro Gly Gln Gln Ala His Ile Lys Thr Thr Tyr Ser Cys 215 220 225 230	1022
25	tcc cgc gcg ccc gtc acc tgg atc ggc cac ggg ggc ggc cac gtc ccc Ser Arg Ala Pro Val Thr Trp Ile Gly His Gly Gly Gly His Val Pro 235 240 245	1070
	gac ccc acg ggc aac aac ggc gtc aag ttt gcg ccc cag gag acc tgg Asp Pro Thr Gly Asn Asn Gly Val Lys Phe Ala Pro Gln Glu Thr Trp 250 255 260	1118
30	gac ttc ttt gat gcc gcc gtc gga gcg gcc ggc gcg cag agc ccg atg Asp Phe Phe Asp Ala Ala Val Gly Ala Ala Gly Ala Gln Ser Pro Met 265 270 275	1166
35	aca taagcccgaa gga Thr	1182
40	<210> 65 <211> 279 <212> PRT <213> Chrysosporium lucknowense <400> 65	
45	Met Ile Ser Val Pro Ala Leu Ala Leu Ala Leu Leu Ala Ala Val Gln 1 5 10 15	
50	Val Val Glu Ser Ala Ser Ala Gly Cys Gly Lys Ala Pro Pro Ser Ser 20 25 30	
55	Gly Thr Lys Ser Met Thr Val Asn Gly Lys Gln Arg Gln Tyr Ile Leu 35 40 45	
	Gln Leu Pro Asn Asn Tyr Asp Ala Asn Lys Ala His Arg Val Val Ile 50 55 60	

EP 2 183 363 B1

	Gly	Tyr	His	Trp	Arg	Asp	Gly	Ser	Met	Asn	Asp	Val	Ala	Asn	Gly	Gly	
	65					70					75					80	
5	Phe	Tyr	Asp	Leu	Arg	Ser	Arg	Ala	Gly	Asp	Ser	Thr	Ile	Phe	Val	Ala	
					85					90					95		
10	Pro	Asn	Gly	Leu	Asn	Ala	Gly	Trp	Ala	Asn	Val	Gly	Gly	Glu	Asp	Ile	
				100					105					110			
15	Thr	Phe	Thr	Asp	Gln	Ile	Val	Asp	Met	Leu	Lys	Asn	Asp	Leu	Cys	Val	
			115					120					125				
20	Asp	Glu	Thr	Gln	Phe	Phe	Ala	Thr	Gly	Trp	Ser	Tyr	Gly	Gly	Ala	Met	
		130					135					140					
25	Ser	His	Ser	Val	Ala	Cys	Ser	Arg	Pro	Asp	Val	Phe	Lys	Ala	Val	Ala	
	145					150					155					160	
30	Val	Ile	Ala	Gly	Ala	Gln	Leu	Ser	Gly	Cys	Ala	Gly	Gly	Thr	Thr	Pro	
				165						170					175		
35	Val	Ala	Tyr	Leu	Gly	Ile	His	Gly	Ala	Ala	Asp	Asn	Val	Leu	Pro	Ile	
				180					185					190			
40	Asp	Leu	Gly	Arg	Gln	Leu	Arg	Asp	Lys	Trp	Leu	Gln	Thr	Asn	Gly	Cys	
			195					200					205				
45	Asn	Tyr	Gln	Gly	Ala	Gln	Asp	Pro	Ala	Pro	Gly	Gln	Gln	Ala	His	Ile	
		210					215					220					
50	Lys	Thr	Thr	Tyr	Ser	Cys	Ser	Arg	Ala	Pro	Val	Thr	Trp	Ile	Gly	His	
	225					230					235					240	
55	Gly	Gly	Gly	His	Val	Pro	Asp	Pro	Thr	Gly	Asn	Asn	Gly	Val	Lys	Phe	
				245						250					255		
60	Ala	Pro	Gln	Glu	Thr	Trp	Asp	Phe	Phe	Asp	Ala	Ala	Val	Gly	Ala	Ala	
				260					265					270			
65	Gly	Ala	Gln	Ser	Pro	Met	Thr										
				275													
70	<210>	66															
75	<211>	837															
80	<212>	DNA															
85	<213>	Chrysosporium lucknowense															

EP 2 183 363 B1

<400> 66

5	atgatctcgg ttctgtctct cgctctggcc cttctggccg ccgtccaggt cgtcagagtct	60
	gcctcggctg gctgtggcaa ggcgccccct tctcgggca ccaagtcgat gacggtcaac	120
	ggcaagcagc gccagtacat tctccagctg cccaacaact acgacgcca caaggccac	180
10	aggggtggtga tcgggtacca ctggcgcgac ggatccatga acgacgtggc caacggcggc	240
	ttctacgata tgcgggtcccg ggcgggcgac agcaccatct tcgttgcccc caacggcctc	300
	aatgccggat gggccaacgt ggcggcgag gacatcacct ttacggacca gatcgtagac	360
15	atgctcaaga acgacctctg cgtggacgag acccagttct ttgctacggg ctggagctat	420
	ggcggtgcca tgagccatag cgtggcttgt tctcggccag acgtcttcaa ggccgtcgcg	480
20	gtcatcgccg gggcccagct gtccggctgc gccggcgga cgacgcccggt ggcgtaccta	540
	ggcatccacg gagccgccga caacgtcctg cccatcgacc tcggccgcca gctgcgcgac	600
	aagtggctgc agaccaacgg ctgcaactac cagggcgccc aggaccccgcc gccggggccag	660
25	caggcccaca tcaagaccac ctacagctgc tcccgcgcgc ccgtcacctg gatcgggccac	720
	gggggcggcc acgtccccga cccacgggc aacaacggcg tcaagtttgc gcccaggag	780
30	acctgggact tctttgatgc cgccgtcgga gcggccggcg cgagagccc gatgaca	837

<210> 67

<211> 1537

<212> DNA

<213> Chrysosporium lucknowense

35

<220>

<221> CDS

<222> (394)..(1299)

40

<400> 67

45

50

55

EP 2 183 363 B1

	ccttccgagt cccaatgccg agtaattaac ccaaacaaca accccagtca ctttagccca	60
	ggttttgcgg cattcgtag agtaccgtta cagccaagcg ttgaaatatg ctagagaaac	120
5	cggaggatgt tccccaggtt ggttctcagc ggagagagca attgcaaccc tggaagaaaa	180
	cgtccgtggc acttcggcaa ctattcttat catatagcac cgtcaatcct tcaccagcat	240
10	gccaaaggac tgtataatat ctttgacata aacttgggcc accattcttg gaacacgagg	300
	atgggctgaa ccctttgttt ccctcttgcc caccatttaa ttacgttcta caaccaatct	360
	ttggcaacat agcaaggacc aagagaagca aag atg tcc cgg tac ctt ctc ggt	414
15	Met Ser Arg Tyr Leu Leu Gly	
	1 5	
	cga cta ata gtg gcc aca gtg gcc atc acg gtg gcc tct ctc cgc ggc	462
	Arg Leu Ile Val Ala Thr Val Ala Ile Thr Val Ala Ser Leu Arg Gly	
20	10 15 20	
25		
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	gtg acg gcc gcc acg ccg tcg ccg gga tgc ggg aag acg ccg acg ctg	510
	Val Thr Ala Ala Thr Pro Ser Pro Gly Cys Gly Lys Thr Pro Thr Leu	
	25 30 35	
5	atc acg gac ggg tcc gcc acg act ccc ctg acg ctg acc tcc aac ggc	558
	Ile Thr Asp Gly Ser Ala Thr Thr Pro Leu Thr Leu Thr Ser Asn Gly	
	40 45 50 55	
10	aag acg cgg cgg ttc tac gtc aag ctt ccc gac gac tac gac aac agc	606
	Lys Thr Arg Arg Phe Tyr Val Lys Leu Pro Asp Asp Tyr Asp Asn Ser	
	60 65 70	
15	cac ccg tac cgg ctc atc ttc gcg ctg cac gcc ctc ggc ggg acc gcc	654
	His Pro Tyr Arg Leu Ile Phe Ala Leu His Ala Leu Gly Gly Thr Ala	
	75 80 85	
20	cag cag gtc acc acc ggg acg ggc ggc tac ctg ccg tgg tac ggg atc	702
	Gln Gln Val Thr Thr Gly Thr Gly Gly Tyr Leu Pro Trp Tyr Gly Ile	
	90 95 100	
25	ccc gac ctc gcg gcc aac gac acg gtg ggc gcc gtc tac gtg gcg ccc	750
	Pro Asp Leu Ala Ala Asn Asp Thr Val Gly Ala Val Tyr Val Ala Pro	
	105 110 115	
30	gac ggg ctc aac aac ggc tgg gcc aac cag ggg ggc gag gac gtg gcc	798
	Asp Gly Leu Asn Asn Gly Trp Ala Asn Gln Gly Gly Glu Asp Val Ala	
	120 125 130 135	
35	ttc ctg gag gcc gtg atg gag acg gtg gag cag gac gtg tgc gtc gac	846
	Phe Leu Glu Ala Val Met Glu Thr Val Glu Gln Asp Val Cys Val Asp	
	140 145 150	
40	cgg gac ctg cgc ttc tcg acc ggc ttc agc tac ggc gcc gcc atg tcg	894
	Arg Asp Leu Arg Phe Ser Thr Gly Phe Ser Tyr Gly Ala Ala Met Ser	
	155 160 165	
45	tac acg ctc gcg tgc gcg ctc ggc cgc cgc atc cgg gcc gtc gcc gtg	942
	Tyr Thr Leu Ala Cys Ala Leu Gly Arg Arg Ile Arg Ala Val Ala Val	
	170 175 180	
50	ctc tcg ggc agc ccc gtc atc agc ggc ggc tgc gcc ggc gcc ggc tcc	990
	Leu Ser Gly Ser Pro Val Ile Ser Gly Gly Cys Ala Gly Ala Gly Ser	
	185 190 195	
55	ggc gcc tcc gag ccc gtc gcc tac tac ggc cag cac ggc atg tcc gac	1038
	Gly Ala Ser Glu Pro Val Ala Tyr Tyr Gly Gln His Gly Met Ser Asp	
	200 205 210 215	
60	ccg gtc ctc ccc gtc gcc ggc ggc cgc gag atg cgc gac cac ttc gtc	1086
	Pro Val Leu Pro Val Ala Gly Gly Arg Glu Met Arg Asp His Phe Val	
	220 225 230	
65	cgc acc aac ggc tgc gac gcc ggc cgg ggc ccc ccc cgg gag ccc gcc	1134
	Arg Thr Asn Gly Cys Asp Ala Gly Arg Gly Pro Pro Arg Glu Pro Ala	
	235 240 245	
70	cgc ggc agc ggc acc cac gtc aag acg gtc tac gac ggc tgc gac ccg	1182
	Arg Gly Ser Gly Thr His Val Lys Thr Val Tyr Asp Gly Cys Asp Pro	
	250 255 260	

EP 2 183 363 B1

	gac tac ccc gtc gtg tgg aac gcg ttc gac ggc gac cac acc ccg cag	1230
	Asp Tyr Pro Val Val Trp Asn Ala Phe Asp Gly Asp His Thr Pro Gln	
	265 270 275	
5	ccc gtc gac cgg ggc gcc acc acc act ttc tcg gcc gtc gag acc tgg	1278
	Pro Val Asp Arg Gly Ala Thr Thr Thr Phe Ser Ala Val Glu Thr Trp	
	280 285 290 295	
10	gag ttc ttc tcg cag ttc aag taattaagcc cgagaggaat cccctgagca	1329
	Glu Phe Phe Ser Gln Phe Lys	
	300	
	ttccctttttg gcacaatatg tgaacccgctc tcgcaacttc ggttcggttt ccgatgatgt	1389
15	taggtattttt acgttccggt catgaccgct gccctgatat tcttgatcgt gtttcaatct	1449
	ggacaagtaa agaacattcg acctttgtgc ctgactgact tcggggccga tgtgccgcgc	1509
	ctgatcacca ggttgtaagt aattaatt	1537
20		
	<210> 68	
	<211> 302	
	<212> PRT	
	<213> Chrysosporium lucknowense	
25		
	<400> 68	
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	Met	Ser	Arg	Tyr	Leu	Leu	Gly	Arg	Leu	Ile	Val	Ala	Thr	Val	Ala	Ile
	1				5					10					15	
5	Thr	Val	Ala	Ser	Leu	Arg	Gly	Val	Thr	Ala	Ala	Thr	Pro	Ser	Pro	Gly
				20					25					30		
10	Cys	Gly	Lys	Thr	Pro	Thr	Leu	Ile	Thr	Asp	Gly	Ser	Ala	Thr	Thr	Pro
			35					40					45			
15	Leu	Thr	Leu	Thr	Ser	Asn	Gly	Lys	Thr	Arg	Arg	Phe	Tyr	Val	Lys	Leu
	50						55					60				
20	Pro	Asp	Asp	Tyr	Asp	Asn	Ser	His	Pro	Tyr	Arg	Leu	Ile	Phe	Ala	Leu
	65					70					75					80
25	His	Ala	Leu	Gly	Gly	Thr	Ala	Gln	Gln	Val	Thr	Thr	Gly	Thr	Gly	Gly
				85						90					95	
30	Tyr	Leu	Pro	Trp	Tyr	Gly	Ile	Pro	Asp	Leu	Ala	Ala	Asn	Asp	Thr	Val
				100					105					110		
35	Gly	Ala	Val	Tyr	Val	Ala	Pro	Asp	Gly	Leu	Asn	Asn	Gly	Trp	Ala	Asn
			115					120					125			
40	Gln	Gly	Gly	Glu	Asp	Val	Ala	Phe	Leu	Glu	Ala	Val	Met	Glu	Thr	Val
45																
50																
55																

EP 2 183 363 B1

[illegible]

EP 2 183 363 B1

	atgtcccggg accttctcgg tcgactaata gtggccacag tggccatcac ggtggcctct	60
	ctccgcggcg tgacggccgc cacgccgtcg ccgggatgcg ggaagacgcc gacgctgac	120
5	acggacgggt ccgccacgac tcccctgacg ctgacctcca acggcaagac gcggcggttc	180
	tacgtcaagc ttcccgacga ctacgacaac agccaccgt accggctcat cttcgcgctg	240
10	cacgccctcg gcgggaccgc ccagcaggtc accaccggga cgggcggcta cctgccgtgg	300
	tacgggatcc ccgacctgc ggccaacgac acggtgggcg ccgtctacgt ggcgcccgc	360
	gggctcaaca acggctgggc caaccagggg ggcgaggacg tggccttcct ggaggccgtg	420
15	atggagacgg tggagcagga cgtgtgcgtc gaccgggacc tgcgcttctc gaccggcttc	480
	agctacggcg ccgccatgtc gtacacgctc gcgtgcgcgc tcggccgccc catccgggcc	540
20	gtcgccgtgc tctcgggcag ccccgatc agcggcggt gcgccggcg cggctccggc	600
	gcctccgagc ccgtcgccta ctacggccag cacggcatgt ccgaccgggt cctccccgtc	660
	gccggcggcc gcgagatgcg cgaccacttc gtccgcacca acggctgcga cgccggccgg	720
25	ggcccccccc gggagcccg ccgcggcagc ggcaccacg tcaagacggt ctacgacggc	780
	tgcgaccgg actaccccg cgtgtggaac gcgttcgacg gcgaccacac ccgcagccc	840
30	gtcgaccgg gcgccaccac cactttctcg gccgtcgaga cctgggagtt cttctcgag	900
	ttcaag	906
	<210> 70	
	<211> 1919	
35	<212> DNA	
	<213> Chrysosporium lucknowense	
	<220>	
	<221> CDS	
40	<222> (279)..(340)	
	<220>	
	<221> CDS	
45	<222> (541)..(718)	
	<220>	
	<221> CDS	
	<222> (802)..(915)	
50	<220>	
	<221> CDS	
	<222> (1028)..(1630)	
55	<400> 70	

EP 2 183 363 B1

	agcactgttg aagggctgtt tggaagtttc ttattgtgag agggggctcg agtcacggca	60
	accaagttcg gcgcctggtc atctctggag ctgaccacgc ctttggccgt cgtcgattcc	120
5	gcgtctcaga gccgagttgg cgggccccgt gcagggttaa ttgtctgcag cgcccggcta	180
	cctggaactc cgagcgatta ccagtttcgg ctcggagcac gaccttcgcc aaccccagat	240
10	cggccacttc aagggaccga aaccgttgcc gtggcccg atg ccg tgc aag tcg agc	296
	Met Pro Cys Lys Ser Ser	
	1 5	
	ctc ctc ctc ctc ctc ctc tcc tcc ttc ttc tcc tcc gag gcc aa	340
15	Leu Leu Leu Leu Leu Leu Ser Ser Phe Phe Ser Ser Glu Ala Asn	
	10 15 20	
	gtccgtgaat acaccaacga cgccgttacg ttgctgatac ttaagtaagg aagggcgcg	400
20		
25		
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	cgctcgctgc	tgctgatgtg	aagacgagga	gcagacttga	acattggtaa	agcagagcac	460	
	tgtttgattg	cgctcgtgca	ctcgccacca	tcatccaatt	aacataccta	gcccataatt	520	
5	atcggacact	ggcgttcgag	c	acc gcc cac	cgc cca cca	tgc gtc tct gcg	571	
				Thr Ala His	Arg Pro Pro	Cys Val Ser Ala		
					25	30		
	atc gct cac	caa gat ggc	cgc cat ggc	cgg cct gcg	cgg ggc gcg	tca	619	
10	Ile Ala His	Gln Asp Gly	Arg His Gly	Arg Pro Ala	Arg Gly Ala	Ser		
		35		40		45		
	ctg cag cta	gtg acc aac	ttt ggc gac	aat ccg acc	ggg ctg cag	aag	667	
	Leu Gln Leu	Val Thr Asn	Phe Gly Asp	Asn Pro Thr	Gly Leu Gln	Lys		
		50		55		60		
15	tat gtg tac	gtt ccg gac	aaa gtt gcg	gtc tcg ccc	gcc atc atc	gtt	715	
	Tyr Val Tyr	Val Pro Asp	Lys Val Ala	Val Ser Pro	Ala Ile Ile	Val		
		65		70		75		
20	gct gtaagtcaac	agagcttgat	taacattatt	atgtgtttga	gttggattga		768	
	Ala							
	80							
	gtgggtgatga	ccgctaactt	tgttgtcgat	cag	ctg cat ccc	tgt gga gga	tcg	822
25					Leu His Pro	Cys Gly Gly	Ser	
							85	
	gct cag gga	tgg tac tcg	cag aca cgg	ctc ccg tcg	tac gcg gac	cag	870	
	Ala Gln Gly	Trp Tyr Ser	Gln Thr Arg	Leu Pro Ser	Tyr Ala Asp	Gln		
		90		95		100		
30	ctt ggg ttc	atc ctc atc	tac gcc ggc	acg acc aag	atg agc aac		915	
	Leu Gly Phe	Ile Leu Ile	Tyr Ala Gly	Thr Thr Lys	Met Ser Asn			
		105		110		115		
35	gtacgcatag	tttttttttt	tttttgcttt	tttttgcttg	ctgtattaca	tagagagtta	975	
	taccettact	aaccataatt	attttcttcc	tcgatatttt	ctttgccaac	ag tgc tgg	1033	
						Cys Trp		
						120		
40	gac gtc cag	aac ccg gcc	agc ctg aca	cac ggc ggg	ggc gac gcc	ggg	1081	
	Asp Val Gln	Asn Pro Ala	Ser Leu Thr	His Gly Gly	Gly Asp Ala	Gly		
			125		130	135		
	ggc atc gtc	agc atg gtc	aag tac gcg	ctg aag cag	tac aac ggc	gac	1129	
45	Gly Ile Val	Ser Met Val	Lys Tyr Ala	Leu Lys Gln	Tyr Asn Gly	Asp		
		140		145		150		
	gcg tcg agg	gtg tac gtc	atg ggg ggc	tcg tcg ggg	gcc atg atg	acc	1177	
	Ala Ser Arg	Val Tyr Val	Met Gly Gly	Ser Ser Gly	Ala Met Met	Thr		
50		155		160		165		
	aac gtg ctg	gcg ggc tcg	tac ccg gac	gtg ttc gag	gcc ggc gcg	gcc	1225	
	Asn Val Leu	Ala Gly Ser	Tyr Pro Asp	Val Phe	Glu Ala Gly	Ala Ala		
		170		175		180		
55	ttc tcc ggg	gtg gcg cac	gcg tgc ttc	ctc ggg gcc	gac tcg gcg	acg	1273	
	Phe Ser Gly	Val Ala His	Ala Cys Phe	Leu Gly Ala	Asp Ser Ala	Thr		

EP 2 183 363 B1

	185		190		195		200	
5	ccc ttc tcg cct aac cag acg tgc gcg cag gcc cgg atc cag cgg tcg							1321
	Pro Phe Ser Pro Asn Gln Thr Cys Ala Gln Gly Arg Ile Gln Arg Ser							
			205		210		215	
10	gcg cgc gag tgg ggg gac ctg gtg cgc aac tcg ttc ccg gcc tac gac							1369
	Ala Arg Glu Trp Gly Asp Leu Val Arg Asn Ser Phe Pro Ala Tyr Asp							
			220		225		230	
15	ggc cgc cgc ccg cgc atg cag atc ttc cac gcc aac gcc gac ttc ctc							1417
	Gly Arg Arg Pro Arg Met Gln Ile Phe His Gly Asn Ala Asp Phe Leu							
			235		240		245	
20	gtc cat ccc gag tgc gcc cac cag gcc ctc gcc cag tgg gcc gac gtc							1465
	Val His Pro Glu Cys Ala His Gln Ala Leu Ala Gln Trp Ala Asp Val							
			250		255		260	
25	ctc ggc ctc cag ctg acc cag acc aac aag gcc gtg ccc tcg gcc gag							1513
	Leu Gly Leu Gln Leu Thr Gln Thr Asn Lys Gly Val Pro Ser Ala Glu							
			265		270		275	280
30	tac acc cag gag gtc tac ggc gac ggc acc cag ctg cag gcc ttc ttt							1561
	Tyr Thr Gln Glu Val Tyr Gly Asp Gly Thr Gln Leu Gln Gly Phe Phe							
			285		290		295	
35	ggc gac ggc gtc ggc cac atc gcc ccg gtc aac gag ccc gtc atg ctc							1609
	Gly Asp Gly Val Gly His Ile Ala Pro Val Asn Glu Pro Val Met Leu							
			300		305		310	
40	cgc ttc ttt ggc ctg atg aac tagctcgtac cggatgtact gtaccgtaaa							1660
	Arg Phe Phe Gly Leu Met Asn							
			315					
45	tgatccatca tttaatagcg aatagtacat agttacatac cagtatgtaa ctgacatgca							1720
	tgtcaagatc ccgacattgt gctccgagtc aaccttggat gggaggtgat gacagcctgg							1780
	aggacgaaag ttgatttgac cgtagcatgc tgtctgtgct cttttgaggt ctgaacgcct							1840
50	cattattacc gcgaacggtg cattgcatta tgggggggtcg ataaacttcc aaggggtata							1900
	gttatacgcg cgtagaaat							1919
55	<210> 71							
	<211> 319							
	<212> PRT							
	<213> Chrysosporium lucknowense							
	<400> 71							

EP 2 183 363 B1

Met Pro Cys Lys Ser Ser Leu Leu Leu Leu Leu Leu Ser Ser Phe Phe
1 5 10 15

5 Ser Ser Glu Ala Asn Thr Ala His Arg Pro Pro Cys Val Ser Ala Ile
20 25 30

10 Ala His Gln Asp Gly Arg His Gly Arg Pro Ala Arg Gly Ala Ser Leu

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	35					40					45				
5	Gln	Leu	Val	Thr	Asn	Phe	Gly	Asp	Asn	Pro	Thr	Gly	Leu	Gln	Tyr
	50						55					60			
	Val	Tyr	Val	Pro	Asp	Lys	Val	Ala	Val	Ser	Pro	Ala	Ile	Ile	Val
10	65					70					75				80
	Leu	His	Pro	Cys	Gly	Gly	Ser	Ala	Gln	Gly	Trp	Tyr	Ser	Gln	Thr
					85					90					95
15	Leu	Pro	Ser	Tyr	Ala	Asp	Gln	Leu	Gly	Phe	Ile	Leu	Ile	Tyr	Ala
				100					105					110	Gly
	Thr	Thr	Lys	Met	Ser	Asn	Cys	Trp	Asp	Val	Gln	Asn	Pro	Ala	Ser
20			115					120					125		Leu
	Thr	His	Gly	Gly	Gly	Asp	Ala	Gly	Gly	Ile	Val	Ser	Met	Val	Lys
25			130				135					140			Tyr
	Ala	Leu	Lys	Gln	Tyr	Asn	Gly	Asp	Ala	Ser	Arg	Val	Tyr	Val	Met
	145					150					155				Gly
30	Gly	Ser	Ser	Gly	Ala	Met	Met	Thr	Asn	Val	Leu	Ala	Gly	Ser	Tyr
					165					170					175
	Asp	Val	Phe	Glu	Ala	Gly	Ala	Ala	Phe	Ser	Gly	Val	Ala	His	Ala
35				180					185					190	Cys
	Phe	Leu	Gly	Ala	Asp	Ser	Ala	Thr	Pro	Phe	Ser	Pro	Asn	Gln	Thr
			195					200					205		Cys
40	Ala	Gln	Gly	Arg	Ile	Gln	Arg	Ser	Ala	Arg	Glu	Trp	Gly	Asp	Leu
		210					215					220			Val
45	Arg	Asn	Ser	Phe	Pro	Ala	Tyr	Asp	Gly	Arg	Arg	Pro	Arg	Met	Gln
	225					230					235				240
	Phe	His	Gly	Asn	Ala	Asp	Phe	Leu	Val	His	Pro	Glu	Cys	Ala	His
50				245						250					255
	Ala	Leu	Ala	Gln	Trp	Ala	Asp	Val	Leu	Gly	Leu	Gln	Leu	Thr	Gln
				260					265					270	Thr
55	Asn	Lys	Gly	Val	Pro	Ser	Ala	Glu	Tyr	Thr	Gln	Glu	Val	Tyr	Gly
			275					280					285		Asp

EP 2 183 363 B1

Gly Thr Gln Leu Gln Gly Phe Phe Gly Asp Gly Val Gly His Ile Ala
290 295 300

5 Pro Val Asn Glu Pro Val Met Leu Arg Phe Phe Gly Leu Met Asn
305 310 315

<210> 72

<211> 957

10 <212> DNA

<213> Chrysosporium lucknowense

<400> 72

15 atgccgtgca agtcgagcct cctcctcctc ctctctctct ccttcttctc ctccgaggcc 60
aacaccgccc accgcccacc atgcgtctct gcgatcgctc accaagatgg ccgccatggc 120
cggcctgcgc ggggcgcgtc actgcagcta gtgaccaact ttggcgacaa tccgaccggt 180
20 ctgcagaagt atgtgtacgt tccggacaaa gttgcggtct cgcccgccat catcggtgct 240
ctgcatacct gtggaggatc ggctcaggga tggactctgc agacacggct cccgtcgtac 300
25 gcggaccagc ttgggttcat cctcatctac gccggcacga ccaagatgag caactgctgg 360
gacgtccaga acccggccag cctgacacac ggcgggggcg acgccggggg catcgtcagc 420
atgggtcaagt acgcgctgaa gcagtacaac ggcgacgcgt cgagggtgta cgtcatgggg 480
30 ggctcgctcg gggccatgat gaccaacgtg ctggcgggct cgtaccggga cgtgttcgag 540
gccggcgcg g ccttctccgg ggtggcgcac gcgtgcttcc tgggggccga ctggcgacg 600
cccttctcgc ctaaccagac gtgcgcgcag ggccggatcc agcggtcggc gcgcgagtgg 660
35 ggggacctgg tgcgcaactc gttcccggcc tacgacggcc gccgcccgcg catgcagatc 720
ttccacggca acgccgactt cctcgtccat cccgagtgcg cccaccaggc cctcgcccag 780
40 tgggcccgacg tcctcggcct ccagctgacc cagaccaaca agggcggtgcc ctgggccgag 840
tacaccaggg aggtctacgg cgacggcacc cagctgcagg gcttctttgg cgacggcgctc 900
ggccacatcg ccccgggtcaa cgagcccgtc atgctccgct tctttggcct gatgaac 957

45 <210> 73

<211> 1560

<212> DNA

<213> Chrysosporium lucknowense

50

<220>

<221> CDS

<222> (221)..(1330)

55

<400> 73

gtgagttaca catagttata taaaggataa ccccatcggg ggaaccagtc ctcgctgcat 60

EP 2 183 363 B1

	atctgcttcc ggcacggatc ttcagttatt tcgcctcgcg cttggccggc tgctccccct	120
	gttcccaccg ccgaagtacg aagtaattaa tagtcattgc gcggacccaa ctgattgtgc	180
5	ctgctccgtg tgaagcagag aaaaggagga caccgccatc atg cgt ctc acc aag Met Arg Leu Thr Lys 1 5	235
10	acg gac ctc agc gtc gcg ctg ctc gcg tcg gcc cta ccc gca acc cag Thr Asp Leu Ser Val Ala Leu Leu Ala Ser Ala Leu Pro Ala Thr Gln 10 15 20	283
15	gcg gcg tgc agc ctc ccc tcg tcc tac cgg tgg acg tcg acg ggc gcg Ala Ala Cys Ser Leu Pro Ser Ser Tyr Arg Trp Thr Ser Thr Gly Ala 25 30 35	331
	ttg gct cag ccc aaa tcg ggg tgg gct aac ctc aag gac ttc acc cac Leu Ala Gln Pro Lys Ser Gly Trp Ala Asn Leu Lys Asp Phe Thr His 40 45 50	379
20	gtg cct tac aac ggc aag cac ctt gtg tac ggg agc tac tac tct aac Val Pro Tyr Asn Gly Lys His Leu Val Tyr Gly Ser Tyr Tyr Ser Asn 55 60 65	427
25	gcc tac ggc tca atg aac ttc ggc ctc ttc tcg gac tgg agc gag atg Ala Tyr Gly Ser Met Asn Phe Gly Leu Phe Ser Asp Trp Ser Glu Met 70 75 80 85	475
30	ggc tcg gcc agc cag aac cag atg agc cag gcc gcg gtc gcg ccc acg Gly Ser Ala Ser Gln Asn Gln Met Ser Gln Ala Ala Val Ala Pro Thr 90 95 100	523
	ctc ttc tac ttc tcc cca aag aac atc tgg gtc ctg gcc tac cag tgg Leu Phe Tyr Phe Ser Pro Lys Asn Ile Trp Val Leu Ala Tyr Gln Trp 105 110 115	571
35	ggt gcc acc cct ttc tcc tac cgc acc tcg acc gat ccg acc aac ccc Gly Ala Thr Pro Phe Ser Tyr Arg Thr Ser Thr Asp Pro Thr Asn Pro 120 125 130	619
40	aac ggc tgg tcc gct ccc cag ccg ctc ttc tcg ggc tcc atc tcc gac Asn Gly Trp Ser Ala Pro Gln Pro Leu Phe Ser Gly Ser Ile Ser Asp 135 140 145	667
45	tcg gac acg ggc ccc atc gac cag acc ctg atc ggc gac gac aag aac Ser Asp Thr Gly Pro Ile Asp Gln Thr Leu Ile Gly Asp Asp Lys Asn 150 155 160 165	715
	atg tac ctg ttc ttc gcc ggc gac aac ggc aag atc tac cgc gcc agc Met Tyr Leu Phe Phe Ala Gly Asp Asn Gly Lys Ile Tyr Arg Ala Ser 170 175 180	763
50	atg ccg ctc ggc aac ttc ccg ggc aac ttt ggc tcg gcc gcc acc gtc Met Pro Leu Gly Asn Phe Pro Gly Asn Phe Gly Ser Ala Ala Thr Val 185 190 195	811
55	gtc ctg agc ggc gcc cgc tac gac ctg ttc gag gcc gtg cag gta tac Val Leu Ser Gly Ala Arg Tyr Asp Leu Phe Glu Ala Val Gln Val Tyr 200 205 210	859

EP 2 183 363 B1

	acg gtc gtc ggc cag agc tcg ccg acc tat ctg atg atc gtc gag tcc	907
	Thr Val Val Gly Gln Ser Ser Pro Thr Tyr Leu Met Ile Val Glu Ser	
	215 220 225	
5	atc ggc gcg cgc ggg cgc tac ttc cgc tcc tac acg gcc agc cgg ctc	955
	Ile Gly Ala Arg Gly Arg Tyr Phe Arg Ser Tyr Thr Ala Ser Arg Leu	
	230 235 240 245	
10	gac ggc gcc tgg acg gcc cag gcg gcc agc gag agc cag ccg ttc gcc	1003
	Asp Gly Ala Trp Thr Ala Gln Ala Ala Ser Glu Ser Gln Pro Phe Ala	
	250 255 260	
15	ggc gcg gcc aac agc ggc gcc tcc tgg acc agc gac atc agc cac ggc	1051
	Gly Ala Ala Asn Ser Gly Ala Ser Trp Thr Ser Asp Ile Ser His Gly	
	265 270 275	
20	gac ctc atc cgc atc gac gcc aac cac acc atg ccc gtc gac ccc tgc	1099
	Asp Leu Ile Arg Ile Asp Ala Asn His Thr Met Pro Val Asp Pro Cys	
	280 285 290	
25	cgc ctc cag ctg ctc tac cag ggc cgc tcc ggc gac agc gcc gac tac	1147
	Arg Leu Gln Leu Leu Tyr Gln Gly Arg Ser Gly Asp Ser Ala Asp Tyr	
	295 300 305	
30	aac tcc ctc ccc tac cgc ccg ggc ctg ctc acc ctg cag ggc gct tct	1195
	Asn Ser Leu Pro Tyr Arg Pro Gly Leu Leu Thr Leu Gln Gly Ala Ser	
	310 315 320 325	
35	tcc ggc ggc ggc gac aac ggc ggc ggc ggc ggc ggc gta acc gtc ccc	1243
	Ser Gly Gly Gly Asp Asn Gly Gly Gly Gly Gly Gly Val Thr Val Pro	
	330 335 340	
40	cgg tac ggc cag tgc ggt ggc cag ggc tac acg ggg ccg aca acg tgc	1291
	Arg Tyr Gly Gln Cys Gly Gly Gln Gly Tyr Thr Gly Pro Thr Thr Cys	
	345 350 355	
45	gag agc cca tac acg tgc aag tac cag aat gat tgt aag tgatgtctta	1340
	Glu Ser Pro Tyr Thr Cys Lys Tyr Gln Asn Asp Cys Lys	
	360 365 370	
50	tttaactctg gtttttttctt ttcttccctc tcttgtgaat cctgtgggta gtagccagtc	1400
	tgaatcaatt tactaattaa caattcttga aagggtactc ccaatgcctc tgacaagcga	1460
	cggaataatt aataagtaag taaaagtggg tgagagcggc cttgaagctg cgggttcctt	1520
55	tttttactgc tgagttcaga tgtatgcata cgtaggtata	1560
	<210> 74	
	<211> 370	
	<212> PRT	
	<213> Chrysosporium lucknowense	
	<400> 74	

EP 2 183 363 B1

Met Arg Leu Thr Lys Thr Asp Leu Ser Val Ala Leu Leu Ala Ser Ala
1 5 10 15

5 Leu Pro Ala Thr Gln Ala Ala Cys Ser Leu Pro Ser Ser Tyr Arg Trp

10

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	20							25					30				
5	Thr	Ser	Thr	Gly	Ala	Leu	Ala	Gln	Pro	Lys	Ser	Gly	Trp	Ala	Asn	Leu	
			35					40					45				
10	Lys	Asp	Phe	Thr	His	Val	Pro	Tyr	Asn	Gly	Lys	His	Leu	Val	Tyr	Gly	
		50					55					60					
15	Ser	Tyr	Tyr	Ser	Asn	Ala	Tyr	Gly	Ser	Met	Asn	Phe	Gly	Leu	Phe	Ser	
	65					70					75					80	
20	Asp	Trp	Ser	Glu	Met	Gly	Ser	Ala	Ser	Gln	Asn	Gln	Met	Ser	Gln	Ala	
					85					90					95		
25	Ala	Val	Ala	Pro	Thr	Leu	Phe	Tyr	Phe	Ser	Pro	Lys	Asn	Ile	Trp	Val	
				100					105					110			
30	Leu	Ala	Tyr	Gln	Trp	Gly	Ala	Thr	Pro	Phe	Ser	Tyr	Arg	Thr	Ser	Thr	
			115					120					125				
35	Asp	Pro	Thr	Asn	Pro	Asn	Gly	Trp	Ser	Ala	Pro	Gln	Pro	Leu	Phe	Ser	
		130					135					140					
40	Gly	Ser	Ile	Ser	Asp	Ser	Asp	Thr	Gly	Pro	Ile	Asp	Gln	Thr	Leu	Ile	
	145					150					155					160	
45	Gly	Asp	Asp	Lys	Asn	Met	Tyr	Leu	Phe	Phe	Ala	Gly	Asp	Asn	Gly	Lys	
					165					170					175		
50	Ile	Tyr	Arg	Ala	Ser	Met	Pro	Leu	Gly	Asn	Phe	Pro	Gly	Asn	Phe	Gly	
				180					185					190			
55	Ser	Ala	Ala	Thr	Val	Val	Leu	Ser	Gly	Ala	Arg	Tyr	Asp	Leu	Phe	Glu	
			195				200						205				
60	Ala	Val	Gln	Val	Tyr	Thr	Val	Val	Gly	Gln	Ser	Ser	Pro	Thr	Tyr	Leu	
		210					215					220					
65	Met	Ile	Val	Glu	Ser	Ile	Gly	Ala	Arg	Gly	Arg	Tyr	Phe	Arg	Ser	Tyr	
	225					230					235					240	
70	Thr	Ala	Ser	Arg	Leu	Asp	Gly	Ala	Trp	Thr	Ala	Gln	Ala	Ala	Ser	Glu	
					245					250					255		
75	Ser	Gln	Pro	Phe	Ala	Gly	Ala	Ala	Asn	Ser	Gly	Ala	Ser	Trp	Thr	Ser	
				260					265					270			

EP 2 183 363 B1

	Asp Ile Ser His Gly Asp Leu Ile Arg Ile Asp Ala Asn His Thr Met	
	275	280 285
5	Pro Val Asp Pro Cys Arg Leu Gln Leu Leu Tyr Gln Gly Arg Ser Gly	
	290	295 300
10	Asp Ser Ala Asp Tyr Asn Ser Leu Pro Tyr Arg Pro Gly Leu Leu Thr	
	305	310 315 320
15	Leu Gln Gly Ala Ser Ser Gly Gly Gly Asp Asn Gly Gly Gly Gly Gly	
	325	330 335
20	Gly Val Thr Val Pro Arg Tyr Gly Gln Cys Gly Gly Gln Gly Tyr Thr	
	340	345 350
25	Gly Pro Thr Thr Cys Glu Ser Pro Tyr Thr Cys Lys Tyr Gln Asn Asp	
	355	360 365
30	Cys Lys	
	370	
	<210> 75	
	<211> 1110	
	<212> DNA	
	<213> Chrysosporium lucknowense	
	<400> 75	
35	atgcgtctca ccaagacgga cctcagcgtc gcgctgctcg cgtcggccct acccgcaacc	60
	caggcggcgt gcagcctccc ctctgcctac cgggtggacgt cgacggggcg gttggctcag	120
	cccaaatcgg ggtgggctaa cctcaaggac ttcaccacg tgccttaciaa cggcaagcac	180
40	cttgtgtacg ggagctacta ctctaacgcc tacggctcaa tgaacttcgg cctcttctcg	240
	gactggagcg agatgggctc ggccagccag aaccagatga gccaggccgc ggtcgcgccc	300
45	acgtctttct acttctcccc aaagaacatc tgggtcctgg cctaccagtg ggggtgccacc	360
	cctttctcct accgcacctc gaccgatccg accaacccca acggctggtc cgctccccag	420
	ccgctcttct cgggctccat ctccgactcg gacacgggcc ccatcgacca gaccctgac	480
50	ggcgacgaca agaacatgta cctgttcttc gccggcgaca acggcaagat ctaccgcgcc	540
	agcatgccgc tcggcaactt cccgggcaac tttggctcgg ccgccaccgt cgtcctgagc	600
	ggcgcccgt acgacctgtt cgaggccgtg caggtataca cggtcgtcgg ccagagctcg	660
55	ccgacctatc tgatgatcgt cgagtccatc ggcgcgcgcg ggcgctactt ccgctcctac	720
	acggccagcc ggctcgacgg cgcttgacg gccaggcg ccagcgagag ccagccgttc	780

EP 2 183 363 B1

	gccggcgcg	ccaacagcg	cgctcctgg	accagcgaca	tcagccacgg	cgacctcatc	840
	cgcatcgacg	ccaaccacac	catgcccgtc	gacccctgcc	gcctccagct	gctctaccag	900
5	ggccgctccg	gcgacagcgc	cgactacaac	tccctcccct	accgcccggg	cctgctcacc	960
	ctgcagggcg	cttcttccgg	cggcggcgac	aacggcgggc	gcggcggcgg	cgtaaccgtc	1020
10	ccccggtacg	gccagtgcgg	tggccagggc	tacacggggc	cgacaacgtg	cgagagccca	1080
	tacacgtgca	agtaccagaa	tgattgtaag				1110
	<210> 76						
	<211> 1310						
15	<212> DNA						
	<213> Chrysosporium lucknowense						
	<220>						
	<221> CDS						
20	<222> (114)..(1076)						
	<400> 76						
25	gtatatatca	gcgcgtagac	agccccgctc	tctccccggc	gttcatgtac	tcgtcgacaa	60
	cgctctctg	atctctctat	atcgccacgt	ggaacaaaca	cccccatccc	acc atg	116
						Met	
						1	
30	agg cca aca	cgc ccc ggt	atc gtg ctc	ctc gcc acc	gcg acc tcg	gtt	164
	Arg Pro Thr	Arg Pro Gly	Ile Val Leu	Leu Ala Thr	Ala Thr Ser	Val	
		5		10	15		
	gcc gga cag	tca tgc gag	ctc ccc tcc	tcg tac cgc	tgg aca tca	acc	212
35	Ala Gly Gln	Ser Cys Glu	Leu Pro Ser	Ser Tyr Arg	Trp Thr Ser	Thr	
		20	25	30			
	ggc gcg ctg	gca cag ccg	aaa agc ccg	tgg gtg tcg	ctc aag gac	ttc	260
40	Gly Ala Leu	Ala Gln Pro	Lys Ser Pro	Trp Val Ser	Leu Lys Asp	Phe	
		35	40	45			
	acc gtc gtc	ccg tac gac	ggc cag cac	ctg gtc tac	gca acg aca	aac	308
	Thr Val Val	Pro Tyr Asp	Gly Gln His	Leu Val Tyr	Ala Thr Thr	Asn	
	50		55	60		65	
45	gac ggg acc	aac tgg ggc	tcc atg ggc	ttc agc ctc	ttc agc gac	tgg	356
	Asp Gly Thr	Asn Trp Gly	Ser Met Gly	Phe Ser Leu	Phe Ser Asp	Trp	
		70	75	80			
	gat gcc atg	ggc tcg gcg	acg cag acg	ggc atg agc	tcc agc gcg	gtg	404
50	Asp Ala Met	Gly Ser Ala	Thr Gln Thr	Gly Met Ser	Ser Ser Ala	Val	
		85	90	95			
	gcg ccc acc	ctg ttc tac	ttc gag ccc	aag aac gtt	tgg atc ctc	gcc	452
55	Ala Pro Thr	Leu Phe Tyr	Phe Glu Pro	Lys Asn Val	Trp Ile Leu	Ala	
		100	105	110			
	cac cag tgg	ggc ccg acg	gcc ttc tcg	tac cgg acg	tcc aac gat	tcc	500
	His Gln Trp	Gly Pro Thr	Ala Phe Ser	Tyr Arg Thr	Ser Asn Asp	Ser	

EP 2 183 363 B1

	115	120	125	
5	acc aac ccc aac agc tgg tcc cag ccc cag ccc ctc ttc acc ggc acc Thr Asn Pro Asn Ser Trp Ser Gln Pro Gln Pro Leu Phe Thr Gly Thr 130 135 140 145	548		
10	atc tcg ggc tcg agc acc ggc ccc atc gac cag acc ctc atc ggc gac Ile Ser Gly Ser Ser Thr Gly Pro Ile Asp Gln Thr Leu Ile Gly Asp 150 155 160	596		
15	gac cag aat atg tac ctc ttc ttc tgc ggc gac aac ggc aag atc tac Asp Gln Asn Met Tyr Leu Phe Phe Cys Gly Asp Asn Gly Lys Ile Tyr 165 170 175	644		
20	cgc gcc agc atg ccc atc ggc aac ttc ccc gga aac ttt ggc agc gag Arg Ala Ser Met Pro Ile Gly Asn Phe Pro Gly Asn Phe Gly Ser Glu 180 185 190	692		
25	tcg acc gtc gtc atg agc gac agc acc aac aac ctc ttc gag gcc gtc Ser Thr Val Val Met Ser Asp Ser Thr Asn Asn Leu Phe Glu Ala Val 195 200 205	740		
30	cag gtg tac aag gtg aac ggg agg cag cag tac ctc atg ctc gtc gag Gln Val Tyr Lys Val Asn Gly Arg Gln Gln Tyr Leu Met Leu Val Glu 210 215 220 225	788		
35	gcc atc ggc gcc cag ggc cgc tac ttc cgc tcc ttc acg gcc acg agg Ala Ile Gly Ala Gln Gly Arg Tyr Phe Arg Ser Phe Thr Ala Thr Arg 230 235 240	836		
40	ctc gac ggc gag tgg acg ccc aac gcg gtc agc gag gct agt ccc ttc Leu Asp Gly Glu Trp Thr Pro Asn Ala Val Ser Glu Ala Ser Pro Phe 245 250 255	884		
45	gcc ggc aag gcc aac tcg ggc gcc agc tgg acc aac gac atc agc cac Ala Gly Lys Ala Asn Ser Gly Ala Ser Trp Thr Asn Asp Ile Ser His 260 265 270	932		
50	ggc gag ctc gtc cgc ctc agc gcc gac cag acc ttc ccc atc gac ccc Gly Glu Leu Val Arg Leu Ser Ala Asp Gln Thr Phe Pro Ile Asp Pro 275 280 285	980		
55	tgc aac ctg cag ctg ctc tac cag ggc cgc gac ccc agc tcg gga ggc Cys Asn Leu Gln Leu Tyr Gln Gly Arg Asp Pro Ser Ser Gly Gly 290 295 300 305	1028		
60	gat tac aac cgc ctg ccc tac cgc ccc ggc gtc ctt acc ctg cag cgc Asp Tyr Asn Arg Leu Pro Tyr Arg Pro Gly Val Leu Thr Leu Gln Arg 310 315 320	1076		
65	tgatctgata ggctctcttt tcgtactttc atccccaggt gcgtcaaattg gggtagctgg	1136		
70	tcgatggggc gggtcagaga aaccctgggc acgcggggaa gttcaaattcg agatgctttg	1196		
75	ttctatacct cacttttttg cggttagcag ccctagtctg tcgaggggatg tcagagcaat	1256		
80	ttctctaaca atttcgactc actaagggtg caacgttcat ctgcaatcgc tgca	1310		
	<210> 77			
	<211> 321			

<210> 77

<211> 321

EP 2 183 363 B1

<212> PRT

<213> Chrysosporium lucknowense

<400> 77

```

5      Met Arg Pro Thr Arg Pro Gly Ile Val Leu Leu Ala Thr Ala Thr Ser
      1          5          10          15

10     Val Ala Gly Gln Ser Cys Glu Leu Pro Ser Ser Tyr Arg Trp Thr Ser
      20          25          30

15     Thr Gly Ala Leu Ala Gln Pro Lys Ser Pro Trp Val Ser Leu Lys Asp
      35          40          45

20     Phe Thr Val Val Pro Tyr Asp Gly Gln His Leu Val Tyr Ala Thr Thr
      50          55          60

25     Asn Asp Gly Thr Asn Trp Gly Ser Met Gly Phe Ser Leu Phe Ser Asp
      65          70          75          80

30     Trp Asp Ala Met Gly Ser Ala Thr Gln Thr Gly Met Ser Ser Ser Ala
      85          90          95

35     Val Ala Pro Thr Leu Phe Tyr Phe Glu Pro Lys Asn Val Trp Ile Leu
      100         105         110

40     Ala His Gln Trp Gly Pro Thr Ala Phe Ser Tyr Arg Thr Ser Asn Asp
      115         120         125

45     Ser Thr Asn Pro Asn Ser Trp Ser Gln Pro Gln Pro Leu Phe Thr Gly
      130         135         140

50     Thr Ile Ser Gly Ser Ser Thr Gly Pro Ile Asp Gln Thr Leu Ile Gly
      145         150         155         160

55     Asp Asp Gln Asn Met Tyr Leu Phe Phe Cys Gly Asp Asn Gly Lys Ile
      165         170         175

60     Tyr Arg Ala Ser Met Pro Ile Gly Asn Phe Pro Gly Asn Phe Gly Ser
      180         185         190

65     Glu Ser Thr Val Val Met Ser Asp Ser Thr Asn Asn Leu Phe Glu Ala
      195         200         205

70     Val Gln Val Tyr Lys Val Asn Gly Arg Gln Gln Tyr Leu Met Leu Val
      210         215         220

```

EP 2 183 363 B1

	Glu	Ala	Ile	Gly	Ala	Gln	Gly	Arg	Tyr	Phe	Arg	Ser	Phe	Thr	Ala	Thr	
	225					230					235					240	
5	Arg	Leu	Asp	Gly	Glu	Trp	Thr	Pro	Asn	Ala	Val	Ser	Glu	Ala	Ser	Pro	
					245					250					255		
10	Phe	Ala	Gly	Lys	Ala	Asn	Ser	Gly	Ala	Ser	Trp	Thr	Asn	Asp	Ile	Ser	
				260					265					270			
15	His	Gly	Glu	Leu	Val	Arg	Leu	Ser	Ala	Asp	Gln	Thr	Phe	Pro	Ile	Asp	
			275					280					285				
20	Pro	Cys	Asn	Leu	Gln	Leu	Leu	Tyr	Gln	Gly	Arg	Asp	Pro	Ser	Ser	Gly	
		290					295					300					
25	Gly	Asp	Tyr	Asn	Arg	Leu	Pro	Tyr	Arg	Pro	Gly	Val	Leu	Thr	Leu	Gln	
	305					310					315					320	
30	Arg																
	<210>	78															
	<211>	963															
	<212>	DNA															
	<213>	Chrysosporium lucknowense															
	<400>	78															
35																	
40																	
45																	
50																	
55																	

EP 2 183 363 B1

	atgaggccaa cacgccccg tatcgtgctc ctcgccaccg cgacctcggg tgccggacag	60
	tcatgcgagc tccccctctc gtaccgctgg acatcaaccg gcgcgctggc acagccgaaa	120
5	agcccgtggg tgtcgtctaa ggacttcacc gtcgtcccg acgacggcca gcacctgggc	180
	tacgcaacga caaacgacgg gaccaactgg ggctccatgg gcttcagcct cttcagcgac	240
10	tgggatgcca tgggctcggc gacgcagacg ggcatgagct ccagcgcggg ggcccccacc	300
	ctgtttctact tcgagcccaa gaacgtttgg atcctcgccc accagtgggg cccgacggcc	360
	ttctcgtacc ggacgtccaa cgattccacc aacccaaca gctgggtcca gccccagccc	420
15	ctcttcaccg gcaccatctc gggctcgagc accggcccca tcgaccagac cctcatcggc	480
	gacgaccaga atatgtacct cttcttctgc ggcgacaacg gcaagatcta ccgcgccagc	540
20	atgcccacgc gcaacttccc cggaaacttt ggcagcgagt cgaccgtcgt catgagcgac	600
	agcaccaaca acctcttcga ggccgtccag gtgtacaagg tgaacgggag gcagcagtac	660
	ctcatgctcg tcgaggccat cggcgcccag ggccgctact tccgctcctt cacggccacg	720
25	aggctcgacg gcgagtggac gcccaacgcg gtcagcgagg ctagtccctt cgccggcaag	780
	gccaactcgg gcgccagctg gaccaacgac atcagccacg gcgagctcgt ccgcctcagc	840
30	gccgaccaga ccttccccat cgaccctgc aacctgcagc tgctctacca gggccgcgac	900
	cccagctcgg gaggcgatta caaccgctg ccctaccgcc ccggcgctct taccctgcag	960
	cgc	963
35	<210> 79 <211> 1422 <212> DNA <213> Chrysosporium lucknowense	
40	<220> <221> CDS <222> (68)..(221)	
45	<220> <221> CDS <222> (288)..(411)	
50	<220> <221> CDS <222> (465)..(570)	
55	<220> <221> CDS <222> (648)..(827)	
	<220> <221> CDS <222> (890)..(1234)	

EP 2 183 363 B1

<400> 79

	aaaccttcta tcaccacctc acttcaactt cactctgccg tcgttttaaa ccggcaactc	60
5	gcccaga atg tgg cct act ccc aag gtt gct tac ctc ttg ctg gca gct	109
	Met Trp Pro Thr Pro Lys Val Ala Tyr Leu Leu Leu Ala Ala	
	1 5 10	
10	cac tcg ctg ctc gtc agc ggg act ccc gtc gac gcc gag gcc gtg gac	157
	His Ser Leu Leu Val Ser Gly Thr Pro Val Asp Ala Glu Ala Val Asp	
	15 20 25 30	
15	acg gcc atc ctc gcc aag cgc caa tgc ccg cag atc cat atc ttc gga	205
	Thr Ala Ile Leu Ala Lys Arg Gln Cys Pro Gln Ile His Ile Phe Gly	
	35 40 45	
	gct cga gaa acg aca g gtttgtcgta ttcttgccag gttactaatg tgcacttgag	261
	Ala Arg Glu Thr Thr	
	50	
20	gtcgtctagc tgacaatacg ggacag ca cca ccc gga tat gga act gcc ggt	313
	Ala Pro Pro Gly Tyr Gly Thr Ala Gly	
	55 60	
25	acc gtc gtc aac ctg att ctc aat gcg ttc cct ggg gcc acg gcc gaa	361
	Thr Val Val Asn Leu Ile Leu Asn Ala Phe Pro Gly Ala Thr Ala Glu	
	65 70 75	
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	gca atc aac tat ccc gcc tgc ggt gga caa gca tct tgt gga gga gtc Ala Ile Asn Tyr Pro Ala Cys Gly Gly Gln Ala Ser Cys Gly Gly Val 80 85 90	409
5	ca gaacgtgatc ccttctatctt ggcgccatgat atcgctcgact aaccatccta cag g Gln	465
10	tac ggt gac tcg gca agg cag ggc acc gac gcg gtg gct tcg gcg gtg Tyr Gly Asp Ser Ala Arg Gln Gly Thr Asp Ala Val Ala Ser Ala Val 95 100 105	513
15	aac agc ttc aac cag cgg tgt ccc aac act cag atc gtg ctt gtt ggc Asn Ser Phe Asn Gln Arg Cys Pro Asn Thr Gln Ile Val Leu Val Gly 110 115 120 125	561
	tac tcc cag gtgcgtgtcg atgagaatga caagacacat ctgggaaagg Tyr Ser Gln	610
20	acaaagatga gaaggaacgg ggttctaaca aacgcag ggc ggt caa atc atc gac Gly Gly Gln Ile Ile Asp 130	665
25	aat gct gtt tgt ggt ggc cct gac acg ggc tct gga atc acg acc acg Asn Ala Val Cys Gly Gly Pro Asp Thr Gly Ser Gly Ile Thr Thr Thr 135 140 145 150	713
30	act cct ccc atc tcg gct gcg gcc ctg aac cag atc aag gcc gtc atc Thr Pro Pro Ile Ser Ala Ala Ala Leu Asn Gln Ile Lys Ala Val Ile 155 160 165	761
	gaa atg ggc tct cca cgg ttc gtc gcc ggc ttg tcc tac gac gtc ggt Glu Met Gly Ser Pro Arg Phe Val Ala Gly Leu Ser Tyr Asp Val Gly 170 175 180	809
35	acc tgc acg gca caa ggt gtaagtggct tcgtcctctc tcgaatgagg Thr Cys Thr Ala Gln Gly 185	857
40	tattctctag tttctaacac acgcattcct ag ttt gct gcg cgc ccg cgt ggc Phe Ala Ala Arg Pro Arg Gly 190 195	910
	tac gtc tgc ggc tcc aac tcg gcg tcc aag atc cag agc tac tgc gat Tyr Val Cys Gly Ser Asn Ser Ala Ser Lys Ile Gln Ser Tyr Cys Asp 200 205 210	958
45	tcg acc gac ccc tac tgc tgc acc ggc aac gac gcc aac agc cac cag Ser Thr Asp Pro Tyr Cys Cys Thr Gly Asn Asp Ala Asn Ser His Gln 215 220 225	1006
50	cag tat ggc aac aag tac gga cag cag gcc ctg gct ttt gtc aag gcc Gln Tyr Gly Asn Lys Tyr Gly Gln Gln Ala Leu Ala Phe Val Lys Ala 230 235 240	1054
55	agg ctc agc ggc agc ggc gga act ccg acg tcg tcg gcc ggc ggc tcc Arg Leu Ser Gly Ser Gly Gly Thr Pro Thr Ser Ser Ala Gly Gly Ser 245 250 255	1102
	gtt ccg acc ggc ggc aac ggc ggc act tgc agc ccg ctg tac gga cag	1150

EP 2 183 363 B1

	Val	Pro	Thr	Gly	Gly	Asn	Gly	Gly	Thr	Cys	Ser	Pro	Leu	Tyr	Gly	Gln	
	260					265				270					275		
5	tgc	ggt	ggc	cag	gga	tgg	acg	ggt	ccc	acg	tgc	tgc	tct	cag	gga	acc	1198
	Cys	Gly	Gly	Gln	Gly	Trp	Thr	Gly	Pro	Thr	Cys	Cys	Ser	Gln	Gly	Thr	
					280					285					290		
	tgc	agg	gca	tcc	aac	cag	tgg	tac	tcc	cag	tgc	ttg	taa	atcc	agg		1244
10	Cys	Arg	Ala	Ser	Asn	Gln	Trp	Tyr	Ser	Gln	Cys	Leu					
				295					300								
	gcg	tgactcc	gtg	agga	aaca	tgg	gtt	ccaa	ggact	cgacc	tctt	cttgta	aata	tcttga			1304
	ttg	cggc	cagt	gtga	atac	ag	ccc	ggcacat	agc	ggaattc	cttt	ctgccc	ctt	ctgacta			1364
15	acac	cgagag	cag	cata	aagt	agata	tatgc	ttag	gaac	ag	acc	gagccag	ata	aaaaaa			1422
	<210>	80															
	<211>	303															
20	<212>	PRT															
	<213>	Chrysosporium lucknowense															
	<400>	80															
25	Met	Trp	Pro	Thr	Pro	Lys	Val	Ala	Tyr	Leu	Leu	Leu	Ala	Ala	His	Ser	
	1				5					10					15		
30	Leu	Leu	Val	Ser	Gly	Thr	Pro	Val	Asp	Ala	Glu	Ala	Val	Asp	Thr	Ala	
				20					25					30			
35	Ile	Leu	Ala	Lys	Arg	Gln	Cys	Pro	Gln	Ile	His	Ile	Phe	Gly	Ala	Arg	
			35					40					45				
	Glu	Thr	Thr	Ala	Pro	Pro	Gly	Tyr	Gly	Thr	Ala	Gly	Thr	Val	Val	Asn	
	50						55					60					
40																	
	Leu	Ile	Leu	Asn	Ala	Phe	Pro	Gly	Ala	Thr	Ala	Glu	Ala	Ile	Asn	Tyr	
	65					70					75				80		
45	Pro	Ala	Cys	Gly	Gly	Gln	Ala	Ser	Cys	Gly	Gly	Val	Gln	Tyr	Gly	Asp	
				85						90					95		
	Ser	Ala	Arg	Gln	Gly	Thr	Asp	Ala	Val	Ala	Ser	Ala	Val	Asn	Ser	Phe	
50				100					105					110			
	Asn	Gln	Arg	Cys	Pro	Asn	Thr	Gln	Ile	Val	Leu	Val	Gly	Tyr	Ser	Gln	
				115				120					125				
55																	
	Gly	Gly	Gln	Ile	Ile	Asp	Asn	Ala	Val	Cys	Gly	Gly	Pro	Asp	Thr	Gly	
	130						135					140					

EP 2 183 363 B1

	Ser	Gly	Ile	Thr	Thr	Thr	Thr	Pro	Pro	Ile	Ser	Ala	Ala	Ala	Leu	Asn
	145						150				155					160
5	Gln	Ile	Lys	Ala	Val	Ile	Glu	Met	Gly	Ser	Pro	Arg	Phe	Val	Ala	Gly
					165					170					175	
10	Leu	Ser	Tyr	Asp	Val	Gly	Thr	Cys	Thr	Ala	Gln	Gly	Phe	Ala	Ala	Arg
				180					185					190		
15	Pro	Arg	Gly	Tyr	Val	Cys	Gly	Ser	Asn	Ser	Ala	Ser	Lys	Ile	Gln	Ser
			195					200					205			
20	Tyr	Cys	Asp	Ser	Thr	Asp	Pro	Tyr	Cys	Cys	Thr	Gly	Asn	Asp	Ala	Asn
	210						215					220				
25	Ser	His	Gln	Gln	Tyr	Gly	Asn	Lys	Tyr	Gly	Gln	Gln	Ala	Leu	Ala	Phe
	225					230					235					240
30	Val	Lys	Ala	Arg	Leu	Ser	Gly	Ser	Gly	Gly	Thr	Pro	Thr	Ser	Ser	Ala
					245					250					255	
35	Gly	Gly	Ser	Val	Pro	Thr	Gly	Gly	Asn	Gly	Gly	Thr	Cys	Ser	Pro	Leu
				260					265					270		
40	Tyr	Gly	Gln	Cys	Gly	Gly	Gln	Gly	Trp	Thr	Gly	Pro	Thr	Cys	Cys	Ser
			275					280					285			
45	Gln	Gly	Thr	Cys	Arg	Ala	Ser	Asn	Gln	Trp	Tyr	Ser	Gln	Cys	Leu	
	290						295					300				

<210> 81

<211> 909

<212> DNA

<213> Chrysosporium lucknowense

<400> 81

EP 2 183 363 B1

	atgtggccta ctcccaaggt tgcttacctc ttgctggcag ctcaactcgct gctcgtcagc	60
	gggactcccg tcgacgccga ggccgtggac acggccatcc tcgccaagcg ccaatgcccc	120
5	cagatccata tcttcggagc tcgagaaacg acagcaccac ccggatatgg aactgccggt	180
	accgtcgtca acctgattct caatgcgttc cctggggcca cggccgaagc aatcaactat	240
10	cccgcttgcg gtggacaagc atcttgtgga ggagtccagt acggtgactc ggcaaggcag	300
	ggcaccgacg cgggtggcttc ggcggtgaac agcttcaacc agcgggtgtcc caacactcag	360
	atcgtgcttg ttggctactc ccagggcggt caaatcatcg acaatgctgt ttgtggtggc	420
15	cctgacacgg gctctggaat cacgaccacg actcctccca tctcggctgc ggccctgaac	480
	cagatcaagg ccgtcatcga aatgggctct ccacggttcg tcgccggctt gtcctacgac	540
20	gtcgggtacct gcacggcaca aggttttgtc gcgcgcccgc gtgggtacgt ctgcggctcc	600
	aactcggcgt ccaagatcca gagctactgc gattcgaccg acccctactg ctgcaccggc	660
	aacgacgcca acagccacca gcagtatggc aacaagtacg gacagcaggc cctggctttt	720
25	gtcaaggcca ggctcagcgg cagcggcgga actccgacgt cgtcggccgg cggctccgtt	780
	ccgaccggcg gcaacggcgg cacttgacgc ccgctgtacg gacagtgcgg tggccaggga	840
	tggacgggtc ccacgtgctg ctctcaggga acctgcaggg catccaacca gtggtactcc	900
30	cagtgccttg	909

	<210> 82
	<211> 1440
35	<212> DNA
	<213> Chrysosporium lucknowense

	<220>
	<221> CDS
40	<222> (282)..(402)

	<220>
	<221> CDS
45	<222> (470)..(594)

	<220>
	<221> CDS
50	<222> (672)..(929)

	<220>
	<221> CDS
55	<222> (987)..(1166)

	<400> 82

EP 2 183 363 B1

	cccaaagaga ttaaagcgac cttccggcgt gccattaggc agccttcgc cgacccttc	60
	tccgtcttcc ctatttcgaa caccgcgcg tccgggctcc ggggtgtaac atagggctc	120
5	ggacgagggg catcagacgg tgcgagtg tgctctggt tgtaatgtag ttatataata	180
	taagaatcca cccgactctt cgggtgtactt gtagccgatg tttgatcagg gcacatctca	240
10	agtccatcta tactcttcaa tcagaataac atatacacgc c atg aag gtc acc gcc	296
	Met Lys Val Thr Ala	
	1 5	
	ggt gcc gtt ccc gtc ctg gct ggc att gcc agc gcc aac ccc gtc gac	344
15	Val Ala Val Pro Val Leu Ala Gly Ile Ala Ser Ala Asn Pro Val Asp	
	10 15 20	
	gtc gag gcc cgc cag tcg tgc ccc gag gtc cac gta ttc ggc gcc cgg	392
	Val Glu Ala Arg Gln Ser Cys Pro Glu Val His Val Phe Gly Ala Arg	
20	25 30 35	
25		
30		
35		
40		
45		
50		
55		

EP 2 183 363 B1

	gaa act aca g gtaattaagc ttaggggttg ggttgtctca ggggtaatta	442
	Glu Thr Thr	
	40	
5	cggctaccta actaacgtca acatcag ca ccc cct ggc tat ggc aca tcc caa	495
	Ala Pro Pro Gly Tyr Gly Thr Ser Gln	
	45	
	ggc ctc gtc aac atg gtc gtg cag gcg tac ccc gga gca aca agc gag	543
10	Gly Leu Val Asn Met Val Val Gln Ala Tyr Pro Gly Ala Thr Ser Glu	
	50 55 60 65	
	gcc ata aac tac ccc gca tgc ggc ggt cag gcg tct tgc ggt gga atc	591
	Ala Ile Asn Tyr Pro Ala Cys Gly Gly Gln Ala Ser Cys Gly Gly Ile	
	70 75 80	
15	gat gcaagtgacc ccgaaaatgc cgctgactgt aaatcaggtg catccgttgt	644
	Asp	
20	tcctaacgag taactcgggg ctttttag tac aat acc tct gcg aac cag ggc act	698
	Tyr Asn Thr Ser Ala Asn Gln Gly Thr	
	85 90	
	cag gcg gtt gtc tcc gct gtg acg agc ttt aac cag cgc tgc cct gac	746
25	Gln Ala Val Val Ser Ala Val Thr Ser Phe Asn Gln Arg Cys Pro Asp	
	95 100 105	
	aca aag atc gtc ttg atc ggc tat tcg cag ggt ggc cag atc atg gac	794
	Thr Lys Ile Val Leu Ile Gly Tyr Ser Gln Gly Gly Gln Ile Met Asp	
	110 115 120	
30	aac gcc tac tgc ggc ggc gcc ggt gcc acc ctc agc ggc agc gcc ctc	842
	Asn Ala Tyr Cys Gly Gly Ala Gly Ala Thr Leu Ser Gly Ser Ala Leu	
	125 130 135	
35	aac gcc gtt aaa gcc acg gtc tgg ttc gga aat ccg cat tat ctg agc	890
	Asn Ala Val Lys Ala Thr Val Trp Phe Gly Asn Pro His Tyr Leu Ser	
	140 145 150 155	
	caa tta agt tat cga gtt ggg aca tgc cag gcc gga ggg gtaggttcct	939
40	Gln Leu Ser Tyr Arg Val Gly Thr Cys Gln Ala Gly Gly	
	160 165	
	ctattgtcga tgtatttttcg tgagacacga catgctgacc gagatag ttc gcg gcc	995
	Phe Ala Ala	
	170	
45	cgc cct ccc gga ttc caa tgt tcg cct ggt aac cca gat aac atc aag	1043
	Arg Pro Pro Gly Phe Gln Cys Ser Pro Gly Asn Pro Asp Asn Ile Lys	
	175 180 185	
	tcg tac tgc gat gcc gag gat ccg tac tgc tgc aat gga aac gat gca	1091
50	Ser Tyr Cys Asp Ala Glu Asp Pro Tyr Cys Cys Asn Gly Asn Asp Ala	
	190 195 200	
	aac cac cac caa cag tac gtc acc atc tac ggc cag cag gcg ctc gcc	1139
	Asn His His Gln Gln Tyr Val Thr Ile Tyr Gly Gln Gln Ala Leu Ala	
	205 210 215	
55	ttc atc aag tcg aag ctc gac gct gct taagcgtagg gagatacgct	1186

EP 2 183 363 B1

Phe Ile Lys Ser Lys Leu Asp Ala Ala
220 225

5 gcggttggaag gggggcgatg gagcacatta ccaagtaaag acagtgccac gtccgtttgt 1246
atgtacacag gtagccaaaa aaacagccat gtcgggatcc atatgtttgc ccctctctct 1306
tggagtgacc cgactctgtt gaggttggtg gtatcgcggt acgagaaagt ggacgaccaa 1366
10 ggcaccaatg agctgctgct ctttgcaatg gacatttctg ggcgatattt gccgtgaact 1426
atcacactcg ccat 1440

<210> 83

15 <211> 228

<212> PRT

<213> Chrysosporium lucknowense

<400> 83

20 Met Lys Val Thr Ala Val Ala Val Pro Val Leu Ala Gly Ile Ala Ser
1 5 10 15
25 Ala Asn Pro Val Asp Val Glu Ala Arg Gln Ser Cys Pro Glu Val His
20 25 30
30 Val Phe Gly Ala Arg Glu Thr Thr Ala Pro Pro Gly Tyr Gly Thr Ser
35 40 45
35 Gln Gly Leu Val Asn Met Val Val Gln Ala Tyr Pro Gly Ala Thr Ser
50 55 60
40 Glu Ala Ile Asn Tyr Pro Ala Cys Gly Gly Gln Ala Ser Cys Gly Gly
65 70 75 80
45 Ile Asp Tyr Asn Thr Ser Ala Asn Gln Gly Thr Gln Ala Val Val Ser
85 90 95
50 Ala Val Thr Ser Phe Asn Gln Arg Cys Pro Asp Thr Lys Ile Val Leu
100 105 110
Ile Gly Tyr Ser Gln Gly Gly Gln Ile Met Asp Asn Ala Tyr Cys Gly
115 120 125
55 Gly Ala Gly Ala Thr Leu Ser Gly Ser Ala Leu Asn Ala Val Lys Ala
130 135 140
Thr Val Trp Phe Gly Asn Pro His Tyr Leu Ser Gln Leu Ser Tyr Arg
145 150 155 160

EP 2 183 363 B1

	Val Gly Thr Cys Gln Ala Gly Gly Phe Ala Ala Arg Pro Pro Gly Phe	
	165 170 175	
5	Gln Cys Ser Pro Gly Asn Pro Asp Asn Ile Lys Ser Tyr Cys Asp Ala	
	180 185 190	
10	Glu Asp Pro Tyr Cys Cys Asn Gly Asn Asp Ala Asn His His Gln Gln	
	195 200 205	
15	Tyr Val Thr Ile Tyr Gly Gln Gln Ala Leu Ala Phe Ile Lys Ser Lys	
	210 215 220	
	Leu Asp Ala Ala	
	225	
20	<210> 84	
	<211> 684	
	<212> DNA	
	<213> Chrysosporium lucknowense	
25	<400> 84	
	atgaaggtca ccgccgttgc cgttccccgtc ctggctggca ttgccagcgc caaccccgtc	60
30	gacgtcgagg ccgccagtc gtgccccgag gtccacgtat tcggcgcccc ggaaactaca	120
	gcaccccctg gctatggcac atcccaaggc ctcgtcaaca tggtcgtgca ggcgtacccc	180
	ggagcaacaa gcgaggccat aaactacccc gcatgcggcg gtcaggcgtc ttgcggtgga	240
35	atcgattaca atacctctgc gaaccagggc actcaggcgg ttgtctccgc tgtgacgagc	300
	tttaaccagc gctgccctga cacaaagatc gtcttgatcg gctattcgca gggtgggcag	360
40	atcatggaca acgcctactg cggcggcgcc ggtgccaccc tcagcggcag cgccctcaac	420
	gccgttaaag ccacgggtctg gttcggaat ccgcattatc tgagccaatt aagttatcga	480
	gttgggacat gccaggccgg agggttcgcg gcccgccctc ccggattcca atgttcgcct	540
45	ggtaacccag ataacatcaa gtcgtactgc gatgccgagg atccgtactg ctgcaatgga	600
	aacgatgcaa accaccacca acagtacgtc accatctacg gccagcaggc gctcgccttc	660
	atcaagtcga agctcgacgc tgct	684
50	<210> 85	
	<211> 1401	
	<212> DNA	
	<213> Chrysosporium lucknowense	
55	<220>	
	<221> CDS	
	<222> (85)..(371)	

<220>
 <221> CDS
 <222> (432)..(849)

5 <220>
 <221> CDS
 <222> (927)..(1075)

10 <220>
 <221> CDS
 <222> (1143)..(1198)

15 <220>
 <221> CDS
 <222> (1263)..(1345)

<400> 85

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	agcttctcag catggacatg ttgaatgcct agcagacgga aatcccatca gtaatatact	60
5	tagtcttctg gaaacttgct cgtc atg agg cta tcc ctt tgc agg ttg gca Met Arg Leu Ser Leu Cys Arg Leu Ala 1 5	111
10	tgc gcg gtg cta tcc ctg acc ttt gtg ggc gat gct ttg cct ggt gct Ser Ala Val Leu Ser Leu Thr Phe Val Gly Asp Ala Leu Pro Gly Ala 10 15 20 25	159
	tct gcg cag gcc gcg agc cgg acc act ccg ccg agc gga tgt ctg tca Ser Ala Gln Ala Ala Ser Arg Thr Thr Pro Ser Gly Cys Leu Ser 30 35 40	207
15	gta gga agc ggc agg cag tac agc acc atc acg gaa gcg ata act gcc Val Gly Ser Gly Arg Gln Tyr Ser Thr Ile Thr Glu Ala Ile Thr Ala 45 50 55	255
20	ttg ggt tct ggc act tca gct gcc tgc att ttc atc tac ccg gga acc Leu Gly Ser Gly Thr Ser Ala Ala Cys Ile Phe Ile Tyr Pro Gly Thr 60 65 70	303
25	tac aac gtt gct gat ggt gtt tcg atc aag tac aag ggc ccg ctg aca Tyr Asn Val Ala Asp Gly Val Ser Ile Lys Tyr Lys Gly Pro Leu Thr 75 80 85	351
30	ctg tat ggg tca acc tcg ga gtgagtactt acataataac tgccaaatgc Leu Tyr Gly Ser Thr Ser Asp 90 95	401
	cgtattcacc actattaatc accatggcag t acc agc aag cag agc gcc aac Thr Ser Lys Gln Ser Ala Asn 100	453
35	cag gtg acc ttt acc agg aac aag ggt tcg gcc gat gct ggg agc ctg Gln Val Thr Phe Thr Arg Asn Lys Gly Ser Ala Asp Ala Gly Ser Leu 105 110 115	501
40	gac gcg tct gcc acg ttc aac atc gtg agc agc aac ttc ccg gcg tac Asp Ala Ser Ala Thr Phe Asn Ile Val Ser Ser Asn Phe Arg Ala Tyr 120 125 130 135	549
45	aac atc aac ttt cgt aac aca tac ggc acg cag ggc cag gcc gtg gcc Asn Ile Asn Phe Arg Asn Thr Tyr Gly Thr Gln Gly Gln Ala Val Ala	597
50		
55		

EP 2 183 363 B1

	140	145	150	
5	gtg gcc gcc aac ggc gac aag cag gcc tac tat gcg tgc ggt ttc tac Val Ala Ala Asn Gly Asp Lys Gln Ala Tyr Tyr Ala Cys Gly Phe Tyr 155 160 165			645
10	ggc tac cag gac acg ctg tac gcc aag agc ggt cgc cag tac tac tcc Gly Tyr Gln Asp Thr Leu Tyr Ala Lys Ser Gly Arg Gln Tyr Tyr Ser 170 175 180			693
15	aac tgc tac atc gag ggc gcc gtc gac ttc atc ttt ggc gcc gcg gct Asn Cys Tyr Ile Glu Gly Ala Val Asp Phe Ile Phe Gly Ala Ala Ala 185 190 195			741
20	gcc tgg ttt ggc gag tgt acc gtt gcc tcc aac ggc gcc ggg tac atc Ala Trp Phe Gly Glu Cys Thr Val Ala Ser Asn Gly Gly Gly Tyr Ile 200 205 210 215			789
25	acc gct aac tct cgc tcc act acg gct gat act acc tgg tat gct ttt Thr Ala Asn Ser Arg Ser Thr Thr Ala Asp Thr Thr Trp Tyr Ala Phe 220 225 230			837
30	gat cac agt act gtaagttcac cgtccaaagc aagtccaacg ctccgtgccc Asp His Ser Thr 235			889
35	acaactatta acgattgtcc cgcttcgcta ttatcag atc cgg gcg gct gcc gga Ile Arg Ala Ala Ala Gly 240			944
40	att agc ttg gcc ggc aaa gtg ttc ctc ggc cga ccg tgg cgc gta ctt Ile Ser Leu Ala Gly Lys Val Phe Leu Gly Arg Pro Trp Arg Val Leu 245 250 255			992
45	gcc cgc gtt atc ttc caa aat tcc gag ttg acc gac gtc gtg cac cct Ala Arg Val Ile Phe Gln Asn Ser Glu Leu Thr Asp Val Val His Pro 260 265 270			1040
50	gag ggc tgg acc aga atg gct gca gga gct act cc gtaagtgatt Glu Gly Trp Thr Arg Met Ala Ala Gly Ala Thr Pro 275 280			1085
55	ctctatccgc gacttctggg gaccagatg ctgacaaata tggcgaaact cttgcag c gag ttc cgg gag ttc caa aac act ggt gca gga tct aac aca tcc cag Glu Phe Arg Glu Phe Gln Asn Thr Gly Ala Gly Ser Asn Thr Ser Gln 290 295 300			1143
60	aga aag t gtaagtgaat gtatcttcta ctctagaggg cactgaattg ctaacgctct Arg Lys			1191
65	tgggtgtgtat gcag gg ctc act ttc cca acg act gct ggt gtc act aag Trp Leu Thr Phe Pro Thr Thr Ala Gly Val Thr Lys 305 310 315			1248
70	acc cag ctc tgg ggc agt gac tgg aag act tgg ata gat act gca tgg Thr Gln Leu Trp Gly Ser Asp Trp Lys Thr Trp Ile Asp Thr Ala Trp 320 325 330			1297
75	tgagaggacg tgtccaaggt tggcaatcag cacctcaaata caacggacat aattaa			1345
80				1401

EP 2 183 363 B1

<210> 86
 <211> 331
 <212> PRT
 <213> Chrysosporium lucknowense

5

<400> 86

10

Met Arg Leu Ser Leu Cys Arg Leu Ala Ser Ala Val Leu Ser Leu Thr
 1 5 10 15

15

Phe Val Gly Asp Ala Leu Pro Gly Ala Ser Ala Gln Ala Ala Ser Arg
 20 25 30

20

Thr Thr Pro Pro Ser Gly Cys Leu Ser Val Gly Ser Gly Arg Gln Tyr
 35 40 45

25

Ser Thr Ile Thr Glu Ala Ile Thr Ala Leu Gly Ser Gly Thr Ser Ala
 50 55 60

Ala Cys Ile Phe Ile Tyr Pro Gly Thr Tyr Asn Val Ala Asp Gly Val
 65 70 75 80

30

Ser Ile Lys Tyr Lys Gly Pro Leu Thr Leu Tyr Gly Ser Thr Ser Asp
 85 90 95

Thr Ser Lys Gln Ser Ala Asn Gln Val Thr Phe Thr Arg Asn Lys Gly
 100 105 110

35

Ser Ala Asp Ala Gly Ser Leu Asp Ala Ser Ala Thr Phe Asn Ile Val
 115 120 125

40

Ser Ser Asn Phe Arg Ala Tyr Asn Ile Asn Phe Arg Asn Thr Tyr Gly
 130 135 140

45

Thr Gln Gly Gln Ala Val Ala Val Ala Ala Asn Gly Asp Lys Gln Ala
 145 150 155 160

Tyr Tyr Ala Cys Gly Phe Tyr Gly Tyr Gln Asp Thr Leu Tyr Ala Lys
 165 170 175

50

Ser Gly Arg Gln Tyr Tyr Ser Asn Cys Tyr Ile Glu Gly Ala Val Asp
 180 185 190

55

Phe Ile Phe Gly Ala Ala Ala Ala Trp Phe Gly Glu Cys Thr Val Ala
 195 200 205

EP 2 183 363 B1

	Ser	Asn	Gly	Gly	Gly	Tyr	Ile	Thr	Ala	Asn	Ser	Arg	Ser	Thr	Thr	Ala	
	210						215					220					
5	Asp	Thr	Thr	Trp	Tyr	Ala	Phe	Asp	His	Ser	Thr	Ile	Arg	Ala	Ala	Ala	
	225					230					235					240	
10	Gly	Ile	Ser	Leu	Ala	Gly	Lys	Val	Phe	Leu	Gly	Arg	Pro	Trp	Arg	Val	
					245					250					255		
15	Leu	Ala	Arg	Val	Ile	Phe	Gln	Asn	Ser	Glu	Leu	Thr	Asp	Val	Val	His	
				260					265					270			
20	Pro	Glu	Gly	Trp	Thr	Arg	Met	Ala	Ala	Gly	Ala	Thr	Pro	Glu	Phe	Arg	
			275					280					285				
25	Glu	Phe	Gln	Asn	Thr	Gly	Ala	Gly	Ser	Asn	Thr	Ser	Gln	Arg	Lys	Trp	
		290					295					300					
30	Leu	Thr	Phe	Pro	Thr	Thr	Ala	Gly	Val	Thr	Lys	Thr	Gln	Leu	Trp	Gly	
	305					310					315					320	
35	Ser	Asp	Trp	Lys	Thr	Trp	Ile	Asp	Thr	Ala	Trp						
					325					330							
40																	
	<210> 87																
	<211> 993																
	<212> DNA																
	<213> Chrysosporium lucknowense																
45																	
	<400> 87																
50																	
55																	

EP 2 183 363 B1

	atgaggctat ccctttgcag gttggcatcg gcggtgctat ccctgacctt tgtgggcat	60
	gctttgcctg gtgcttctgc gcaggccgcg agccggacca ctccgccgag cggatgtctg	120
5	tcagtaggaa gcggcaggca gtacagcacc atcacggaag cgataactgc cttgggttct	180
	ggcacttcag ctgcctgcat tttcatctac ccgggaacct acaacgttgc tgatgggtgtt	240
10	tcgatcaagt acaagggccc gctgacactg tatgggtcaa cctcggatac cagcaagcag	300
	agcgccaacc aggtgacctt taccaggaac aagggttcgg ccgatgctgg gagcctggac	360
	gcgtctgcca cgttcaacat cgtgagcagc aacttccggg cgtacaacat caactttcgt	420
15	aacacatacg gcacgcaggg ccaggccgtg gccgtggccg ccaacggcga caagcaggcc	480
	tactatgcgt gcggtttcta cggctaccag gacacgctgt acgccaagag cggtcgccag	540
20	tactactcca actgctacat cgagggcgcc gtcgacttca tctttggcgc cgcggctgcc	600
	tggtttggcg agtgtaccgt tgcctccaac ggcgggcggg acatcacgcg taactctcgc	660
	tccactacgg ctgatactac ctggtatgct tttgatcaca gtactatccg ggcggtgcc	720
25	ggaattagct tggccggcaa agtgttcctc ggccgacctg ggcgcgact tgcccgcgtt	780
	atcttccaaa attccgagtt gaccgacgtc gtgcaccctg agggctggac cagaatggct	840
30	gcaggagcta ctcccagatt ccgggagttc caaaacactg gtgcaggatc taacacatcc	900
	cagagaaaagt ggctcacttt cccaacgact gctggtgtca ctaagacca gctctggggc	960
	agtgactgga agacttggat agatactgca tgg	993
35	<210> 88 <211> 1504 <212> DNA <213> Chrysosporium lucknowense	
40	<220> <221> CDS <222> (296)..(565)	
45	<220> <221> CDS <222> (650)..(1240)	
50	<220> <221> CDS <222> (1351)..(1452)	
	<400> 88	

55

EP 2 183 363 B1

	acaccctcac gacaagtcgc aacctcgatt tcccgtgtgc ccggagcttg gcaacgcgtc	60
	acgacccata gttagtagta tcgtcaaggg ccgtctcgct aaaagtagcg tggtttcaga	120
5	aaccgtttat ttagcgacct taaataatta ttattgatta tctcccgaca acctccgggc	180
	ctcgtcggcg ggtttctgtc taactgctgc tgcaaacttc aggcggctgt ccgtcgtggt	240
10	cggtcattct caccttgcca gctttggcta attaccttac cttgctctac caacg atg Met 1	298
	aag aga tcc ggc ctt ctg ggg acg gtg gct ctt ctg ttc ccg gcc ctc	346
15	Lys Arg Ser Gly Leu Leu Gly Thr Val Ala Leu Leu Phe Pro Ala Leu 5 10 15	
	ggt tct gcc tat gcg ctc ccg ggc gaa tgt act gga gtc tgc aac aac	394
20	Val Ser Ala Tyr Ala Leu Pro Gly Glu Cys Thr Gly Val Cys Asn Asn 20 25 30	
	tcc cac gac ccg tca atc atc cgc cgc gca gat ggg acg tac ttc cgc	442
25	Ser His Asp Pro Ser Ile Ile Arg Arg Ala Asp Gly Thr Tyr Phe Arg 35 40 45	
	ttc tcg act ggt ggc aag atc gcc att cac acg gcg ccg agc ctg acc	490
30	Phe Ser Thr Gly Gly Lys Ile Ala Ile His Thr Ala Pro Ser Leu Thr 50 55 60 65	
	ggc ccg tgg acc tat agg ggc gcc gcc ttg ccg gcc ggg tcc tcc atc	538
35	Gly Pro Trp Thr Tyr Arg Gly Ala Ala Leu Pro Ala Gly Ser Ser Ile	
40		
45		
50		
55		

EP 2 183 363 B1

	70	75	80	
5	aac ttg gac ggg aac cag gac ctc tgg	gtaagaagaa caaggcgcttc	585	
	Asn Leu Asp Gly Asn Gln Asp Leu Trp			
	85	90		
	acggccccgca gcctggatgt aattttattta tgggtgatga tgatgttgac gacgagctcg		645	
10	ccag gcc ccc gat gtt gca cag gtc gga gac cag tac tac ctg tac tac		694	
	Ala Pro Asp Val Ala Gln Val Gly Asp Gln Tyr Tyr Leu Tyr Tyr			
	95	100	105	
15	tcc gtc agc acc ttc ggc gtc cag aac tcg gcc atc ggc ata gcc cgc		742	
	Ser Val Ser Thr Phe Gly Val Gln Asn Ser Ala Ile Gly Ile Ala Arg			
	110	115	120	
	tca tcg agc ctg gac gcc ggc acc tgg acc gac ctc ggg tct acc ggt		790	
	Ser Ser Ser Leu Asp Ala Gly Thr Trp Thr Asp Leu Gly Ser Thr Gly			
	125	130	135	
20	gtc acc tcc agc acg ggc tcg gcg tac aac gcc atc gac ggc aac ctg		838	
	Val Thr Ser Ser Thr Gly Ser Ala Tyr Asn Ala Ile Asp Gly Asn Leu			
	140	145	150	
25	atc aac gag ccg ggg acc agc aac tac ttc ctc acg ttt ggc agc ttc		886	
	Ile Asn Glu Pro Gly Thr Ser Asn Tyr Phe Leu Thr Phe Gly Ser Phe			
	155	160	165	
30	tgg aac ggc atc cat cgc gtc cgc atg acg ccc acc aag acg aac ggc		934	
	Trp Asn Gly Ile His Arg Val Arg Met Thr Pro Thr Lys Thr Asn Gly			
	170	175	180	185
	aac gtg tac cag gtc gcc ttt gat ccc aac gac ccg gcc atg gaa ggc		982	
	Asn Val Tyr Gln Val Ala Phe Asp Pro Asn Asp Pro Ala Met Glu Gly			
	190	195	200	
35	ccg acc gtc ttc aag tac ggc gac tac tac tac ctt ttc ttc tcc aag		1030	
	Pro Thr Val Phe Lys Tyr Gly Asp Tyr Tyr Tyr Leu Phe Phe Ser Lys			
	205	210	215	
40	ggc acg tgc tgc ggc tac gac cag aac agg ccg ccc gcc gga cag gag		1078	
	Gly Thr Cys Cys Gly Tyr Asp Gln Asn Arg Pro Pro Ala Gly Gln Glu			
	220	225	230	
	tac agg atc atg gtg tgc agg tcg acc tcg ccc acc ggc ccc ttc gag		1126	
	Tyr Arg Ile Met Val Cys Arg Ser Thr Ser Pro Thr Gly Pro Phe Glu			
	235	240	245	
45	gac agg gac ggc aag tcg tgc cgc tcg ggc ggt gga act ctg gtg ctc		1174	
	Asp Arg Asp Gly Lys Ser Cys Arg Ser Gly Gly Gly Thr Leu Val Leu			
	250	255	260	265
50	ccg tct cac gac tgg gtg tac ggc ccc ggc ggc caa ggt gtc tac cag		1222	
	Pro Ser His Asp Trp Val Tyr Gly Pro Gly Gly Gln Gly Val Tyr Gln			
	270	275	280	
55	gac ccc gaa cac gga ccg gttagttccc cgtcctttct cccgtacggg		1270	
	Asp Pro Glu His Gly Pro			
	285			
	cctagcattc agcatcgggg atcgagcatc aagcatcacc ctgcttgaac ccatctcttg		1330	

EP 2 183 363 B1

	ctaacgctct tccttcccag gtc ttg tac tac cat tat gtt gat act cgc att	1383
	Val Leu Tyr Tyr His Tyr Val Asp Thr Arg Ile	
	290 295	
5	ggc tat gcg gac ggc cag aag aag ttt ggt tgg aat cgc atc gac ttt	1431
	Gly Tyr Ala Asp Gly Gln Lys Lys Phe Gly Trp Asn Arg Ile Asp Phe	
	300 305 310	
10	tca agc ggt tgg ccg gtt gtc tgagaaaccc gagacgcttg acttgtttat	1482
	Ser Ser Gly Trp Pro Val Val	
	315 320	
	ttacaggtat taccggtatt aa	1504
15	<210> 89	
	<211> 321	
	<212> PRT	
	<213> Chrysosporium lucknowense	
20	<400> 89	
	Met Lys Arg Ser Gly Leu Leu Gly Thr Val Ala Leu Leu Phe Pro Ala	
	1 5 10 15	
25	Leu Val Ser Ala Tyr Ala Leu Pro Gly Glu Cys Thr Gly Val Cys Asn	
	20 25 30	
30	Asn Ser His Asp Pro Ser Ile Ile Arg Arg Ala Asp Gly Thr Tyr Phe	
	35 40 45	
35	Arg Phe Ser Thr Gly Gly Lys Ile Ala Ile His Thr Ala Pro Ser Leu	
	50 55 60	
40	Thr Gly Pro Trp Thr Tyr Arg Gly Ala Ala Leu Pro Ala Gly Ser Ser	
	65 70 75 80	
45	Ile Asn Leu Asp Gly Asn Gln Asp Leu Trp Ala Pro Asp Val Ala Gln	
	85 90 95	
50	Val Gly Asp Gln Tyr Tyr Leu Tyr Tyr Ser Val Ser Thr Phe Gly Val	
	100 105 110	
55	Gln Asn Ser Ala Ile Gly Ile Ala Arg Ser Ser Ser Leu Asp Ala Gly	
	115 120 125	
	Thr Trp Thr Asp Leu Gly Ser Thr Gly Val Thr Ser Ser Thr Gly Ser	
	130 135 140	
	Ala Tyr Asn Ala Ile Asp Gly Asn Leu Ile Asn Glu Pro Gly Thr Ser	
	145 150 155 160	

EP 2 183 363 B1

	Asn	Tyr	Phe	Leu	Thr	Phe	Gly	Ser	Phe	Trp	Asn	Gly	Ile	His	Arg	Val	
					165					170					175		
5	Arg	Met	Thr	Pro	Thr	Lys	Thr	Asn	Gly	Asn	Val	Tyr	Gln	Val	Ala	Phe	
				180					185				190				
10	Asp	Pro	Asn	Asp	Pro	Ala	Met	Glu	Gly	Pro	Thr	Val	Phe	Lys	Tyr	Gly	
			195					200					205				
15	Asp	Tyr	Tyr	Tyr	Leu	Phe	Phe	Ser	Lys	Gly	Thr	Cys	Cys	Gly	Tyr	Asp	
	210						215					220					
20	Gln	Asn	Arg	Pro	Pro	Ala	Gly	Gln	Glu	Tyr	Arg	Ile	Met	Val	Cys	Arg	
	225					230					235					240	
25	Ser	Thr	Ser	Pro	Thr	Gly	Pro	Phe	Glu	Asp	Arg	Asp	Gly	Lys	Ser	Cys	
					245					250					255		
30	Arg	Ser	Gly	Gly	Gly	Thr	Leu	Val	Leu	Pro	Ser	His	Asp	Trp	Val	Tyr	
			260						265					270			
35	Gly	Pro	Gly	Gly	Gln	Gly	Val	Tyr	Gln	Asp	Pro	Glu	His	Gly	Pro	Val	
			275					280					285				
40	Leu	Tyr	Tyr	His	Tyr	Val	Asp	Thr	Arg	Ile	Gly	Tyr	Ala	Asp	Gly	Gln	
	290						295					300					
45	Lys	Lys	Phe	Gly	Trp	Asn	Arg	Ile	Asp	Phe	Ser	Ser	Gly	Trp	Pro	Val	
	305					310					315					320	
50	Val																

<210> 90

<211> 963

<212> DNA

<213> Chrysosporium lucknowense

<400> 90

50	atgaagagat	ccggccttct	ggggacggtg	gctcttctgt	tcccggccct	cgtttctgcc	60
	tatgcgctcc	cgggcgaatg	tactggagtc	tgcaacaact	cccacgaccc	gtcaatcatc	120
	cgccgcgcag	atgggacgta	cttccgcttc	tcgactgggtg	gcaagatcgc	cattcacacg	180
55	gcgcccagacc	tgaccggccc	gtggacctat	aggggcgccg	ccttgccggc	cggttctctcc	240
	atcaacttgg	acgggaacca	ggacctctgg	gccccgatg	ttgcacaggt	cggagaccag	300

EP 2 183 363 B1

	tactacctgt actactccgt cagcaccttc ggcggtccaga actcggccat cggcatagcc	360
	cgctcatcga gcctggacgc cggcacctgg accgacctcg ggtctaccgg tgtcacctcc	420
5	agcacgggct cggcgtacaa cgccatcgac ggcaacctga tcaacgagcc ggggaccagc	480
	aactacttcc tcacgtttgg cagcttctgg aacggcatcc atcgcgtccg catgacgccc	540
10	accaagacga acggcaacgt gtaccaggtc gcctttgatc ccaacgaccc ggccatggaa	600
	ggccccgaccg tcttcaagta cggcgactac tactaccttt tcttctccaa gggcacgtgc	660
	tgcggctacg accagaacag gccgccccgcc ggacaggagt acaggatcat ggtgtgcagg	720
15	tcgacctcgc ccaccggccc cttcgaggac agggacggca agtcgtgccg ctcgggcggt	780
	ggaactctgg tgctcccgtc tcacgactgg gtgtacggcc ccggcgccca aggtgtctac	840
	caggacccccg aacacggacc ggtcttgtac taccattatg ttgatactcg cattggctat	900
20	gcggacggcc agaagaagtt tggttggaat cgcacgact tttcaagcgg ttggccggtt	960
	gtc	963
25	<210> 91 <211> 2451 <212> DNA <213> Chrysosporium lucknowense	
30	<220> <221> CDS <222> (57)..(1851)	
35	<220> <221> CDS <222> (1954)..(2357)	
	<400> 91	
40	aggaccgtca aacgtaaggg cagctctagc tcaccagata cgagacgtcg cccatc atg	59
		Met
		1
45	acc ctt caa gcc ttt gcg ctg ctg gcg gcg gcg gcc ctt gta cgg ggc	107
	Thr Leu Gln Ala Phe Ala Leu Leu Ala Ala Ala Ala Leu Val Arg Gly	
	5 10 15	
	gag acc ccg acc aag gtc cct cgc gac gct ccg aga gga gct gct gct	155
50	Glu Thr Pro Thr Lys Val Pro Arg Asp Ala Pro Arg Gly Ala Ala Ala	
	20 25 30	
	tgg gaa gct gcc cac tcc tca gca gcc gct gcc ttg gga aag ctg tcc	203
	Trp Glu Ala Ala His Ser Ser Ala Ala Ala Ala Leu Gly Lys Leu Ser	
	35 40 45	
55	cag cag gac aag atc aac atc gtg acg ggc gtc ggc tgg aac aag ggg	251
	Gln Gln Asp Lys Ile Asn Ile Val Thr Gly Val Gly Trp Asn Lys Gly	
	50 55 60 65	

EP 2 183 363 B1

	ccc tgc gtg ggc aat act ccc gct atc agc tcc atc aac tac ccg cag	299
	Pro Cys Val Gly Asn Thr Pro Ala Ile Ser Ser Ile Asn Tyr Pro Gln	
	70 75 80	
5	ctc tgc ctc caa gac gga ccc ctg ggc gtc cgc ttc ggt tcc agc atc	347
	Leu Cys Leu Gln Asp Gly Pro Leu Gly Val Arg Phe Gly Ser Ser Ile	
	85 90 95	
10	acc gct ttc act cca ggc att cag gcc gcg tcg acg tgg gat gtg gac	395
	Thr Ala Phe Thr Pro Gly Ile Gln Ala Ala Ser Thr Trp Asp Val Asp	
	100 105 110	
15	ctg atc cga caa cga ggc gag tac atg ggc gcc gag ttc aag ggg tgt	443
	Leu Ile Arg Gln Arg Gly Glu Tyr Met Gly Ala Glu Phe Lys Gly Cys	
	115 120 125	
	ggc atc cac gtc cag cta ggc ccc gtg gcc ggg ccc ttg ggt aag gtt	491
	Gly Ile His Val Gln Leu Gly Pro Val Ala Gly Pro Leu Gly Lys Val	
	130 135 140 145	
20	ccg caa ggc ggc cgc aac tgg gag ggc ttt ggc gtg gat ccc tac ctc	539
	Pro Gln Gly Gly Arg Asn Trp Glu Gly Phe Gly Val Asp Pro Tyr Leu	
	150 155 160	
25	acc ggc att gcc atg gcc gag acc atc gag ggg ata cag tca gcg ggc	587
	Thr Gly Ile Ala Met Ala Glu Thr Ile Glu Gly Ile Gln Ser Ala Gly	
	165 170 175	
30	gtg caa gcc acc gcc aag cat tac atc ctc aac gag cag gag ctc aac	635
	Val Gln Ala Thr Ala Lys His Tyr Ile Leu Asn Glu Gln Glu Leu Asn	
	180 185 190	
	cgc gag acc atg agc agc aat gtc gac gac cgc acc ttg cac gag ctg	683
	Arg Glu Thr Met Ser Ser Asn Val Asp Asp Arg Thr Leu His Glu Leu	
	195 200 205	
35	tac ctc tgg cct ttt gca gac gct gtt cac tct aac gtg gcc agc gtc	731
	Tyr Leu Trp Pro Phe Ala Asp Ala Val His Ser Asn Val Ala Ser Val	
	210 215 220 225	
40	atg tgc agt tac aac aag atc aac ggc acg tgg gcg tgc gag aac gac	779
	Met Cys Ser Tyr Asn Lys Ile Asn Gly Thr Trp Ala Cys Glu Asn Asp	
	230 235 240	
	cgg gtg ctg aac gtc att ctg aag cag gag ctt ggg ttt ccg ggc tat	827
	Arg Val Leu Asn Val Ile Leu Lys Gln Glu Leu Gly Phe Pro Gly Tyr	
	245 250 255	
45	gtg atg agc gac tgg aac gcg cag cac tcg acc gac gac gcg gcc aac	875
	Val Met Ser Asp Trp Asn Ala Gln His Ser Thr Asp Asp Ala Ala Asn	
	260 265 270	
50	cac ggc atg gac atg acg atg ccc ggc agc gac ttc aac gga ggg acg	923
	His Gly Met Asp Met Thr Met Pro Gly Ser Asp Phe Asn Gly Gly Thr	
	275 280 285	
55	atc ctc tgg ggc ccg cag ctc gac agc gcc gtc aac agc ggc cgg gtg	971
	Ile Leu Trp Gly Pro Gln Leu Asp Ser Ala Val Asn Ser Gly Arg Val	
	290 295 300 305	
	ccc aag tcc cgc ctc gac gac atg gtc gag cgc atc ctc gcg gcg tgg	1019

EP 2 183 363 B1

	Pro	Lys	Ser	Arg	Leu	Asp	Asp	Met	Val	Glu	Arg	Ile	Leu	Ala	Ala	Trp	
					310					315					320		
5	tac	ctt	ctc	ggt	cag	gac	agc	aat	tac	ccg	gct	atc	aac	atc	ggc	gcc	1067
	Tyr	Leu	Leu	Gly	Gln	Asp	Ser	Asn	Tyr	Pro	Ala	Ile	Asn	Ile	Gly	Ala	
				325					330					335			
10	aac	gta	cag	ggc	aac	cac	aag	gag	aac	gtg	cgg	gcg	gtc	gcg	cgc	gat	1115
	Asn	Val	Gln	Gly	Asn	His	Lys	Glu	Asn	Val	Arg	Ala	Val	Ala	Arg	Asp	
			340					345					350				
	ggc	atc	gtg	ctc	ctc	aag	aac	gac	gac	ggc	atc	ctc	ccg	ctc	aag	aag	1163
	Gly	Ile	Val	Leu	Leu	Lys	Asn	Asp	Asp	Gly	Ile	Leu	Pro	Leu	Lys	Lys	
		355					360					365					
15	ccc	gcc	aag	ctc	gcc	ctc	atc	ggt	tcg	gcc	gcc	gtc	gtc	aac	ccg	cag	1211
	Pro	Ala	Lys	Leu	Ala	Leu	Ile	Gly	Ser	Ala	Ala	Val	Val	Asn	Pro	Gln	
	370					375					380					385	
20	gga	ctg	aac	tcc	tgc	cag	gat	cag	ggc	tgc	aac	aaa	ggc	gcg	ctg	ggc	1259
	Gly	Leu	Asn	Ser	Cys	Gln	Asp	Gln	Gly	Cys	Asn	Lys	Gly	Ala	Leu	Gly	
				390					395						400		
25	atg	gga	tgg	ggg	tcc	ggg	gcg	gtc	aac	tac	ccg	tac	ttt	gtc	gcg	ccg	1307
	Met	Gly	Trp	Gly	Ser	Gly	Ala	Val	Asn	Tyr	Pro	Tyr	Phe	Val	Ala	Pro	
				405					410					415			
	tac	gac	gcg	ctc	aag	gcc	cgc	gcc	caa	gag	gac	ggg	acg	acg	gtc	agc	1355
	Tyr	Asp	Ala	Leu	Lys	Ala	Arg	Ala	Gln	Glu	Asp	Gly	Thr	Thr	Val	Ser	
			420					425					430				
30	ctg	cac	aac	tcg	gac	agc	acg	agc	ggg	gtg	gcg	aac	gtg	gcc	tcg	gac	1403
	Leu	His	Asn	Ser	Asp	Ser	Thr	Ser	Gly	Val	Ala	Asn	Val	Ala	Ser	Asp	
		435					440					445					
35	gcg	gac	gcg	gcc	atc	gtg	gtc	atc	acg	gcc	gac	tcg	ggc	gag	ggg	tac	1451
	Ala	Asp	Ala	Ala	Ile	Val	Val	Ile	Thr	Ala	Asp	Ser	Gly	Glu	Gly	Tyr	
	450					455					460					465	
40	atc	aca	gtc	gaa	ggc	gcc	gcc	ggg	gac	cgg	ctg	aac	ctc	gac	ccg	tgg	1499
	Ile	Thr	Val	Glu	Gly	Ala	Ala	Gly	Asp	Arg	Leu	Asn	Leu	Asp	Pro	Trp	
				470					475					480			
	cac	aac	ggc	aac	gag	ctg	gta	aag	gcg	gtg	gcg	gcg	gcc	aac	aag	aac	1547
	His	Asn	Gly	Asn	Glu	Leu	Val	Lys	Ala	Val	Ala	Ala	Ala	Asn	Lys	Asn	
				485					490				495				
45	acc	atc	gtc	gtc	gtg	cac	agc	gtc	ggg	ccc	atc	atc	ctc	gag	acc	atc	1595
	Thr	Ile	Val	Val	Val	His	Ser	Val	Gly	Pro	Ile	Ile	Leu	Glu	Thr	Ile	
			500					505					510				
50	ctg	gca	acc	gaa	ggg	gtc	aag	gcg	att	gtg	tgg	gcc	ggc	ctg	cca	agc	1643
	Leu	Ala	Thr	Glu	Gly	Val	Lys	Ala	Ile	Val	Trp	Ala	Gly	Leu	Pro	Ser	
		515					520					525					
55	cag	gag	aat	ggt	aac	gcg	ctg	gtg	gac	atc	ctg	tac	ggg	ctg	gcc	tcg	1691
	Gln	Glu	Asn	Gly	Asn	Ala	Leu	Val	Asp	Ile	Leu	Tyr	Gly	Leu	Ala	Ser	
	530					535					540					545	
	ccg	tcg	ggc	aag	ctg	gtc	tac	acc	atc	gcc	aag	cga	gaa	cag	gac	tac	1739
	Pro	Ser	Gly	Lys	Leu	Val	Tyr	Thr	Ile	Ala	Lys	Arg	Glu	Gln	Asp	Tyr	

EP 2 183 363 B1

	550	555	560	
5	ggc act gcg gtc gtt cgc gga gac gac aca ttt ccg gag ggc ctg ttt Gly Thr Ala Val Val Arg Gly Asp Asp Thr Phe Pro Glu Gly Leu Phe 565 570 575	1787		
10	gtc gac tac cgg cac ttt gac aag gag aac atc gag ccg cgg tac gag Val Asp Tyr Arg His Phe Asp Lys Glu Asn Ile Glu Pro Arg Tyr Glu 580 585 590	1835		
15	ttt ggg ttt ggg ctg t gtaagttgtt ctctgtcata aagagtgtcc ccgtccactc Phe Gly Phe Gly Leu 595	1891		
20	cgccatcttt tttttttttt cagttttacc tcgtcctaac agcaagtga acaaactggc ag ca tac acc aac ttc acg tac gcc gac ctc gaa ctc acc tcg acc Ser Tyr Thr Asn Phe Thr Tyr Ala Asp Leu Glu Leu Thr Ser Thr 600 605 610	1951	1997	
25	gcc acg gcc ggc cca gcg acg ggc gag acc atc ccc ggc ggc gcg gcc Ala Thr Ala Gly Pro Ala Thr Gly Glu Thr Ile Pro Gly Gly Ala Ala 615 620 625	2045		
30	gac ctc tgg gag gag gtg gcc acg gtc acg gcg acc atc acc aac agc Asp Leu Trp Glu Glu Val Ala Thr Val Thr Ala Thr Ile Thr Asn Ser 630 635 640 645	2093		
35	ggc ggc gtg gac ggc gcc gag gtg gcc cag ctg tac ctg acg ttg ccg Gly Gly Val Asp Gly Ala Glu Val Ala Gln Leu Tyr Leu Thr Leu Pro 650 655 660	2141		
40	tcg tcg gcg ccg gcg acc ccg ccc aag cag ctc cgc ggc ttc gcc aag Ser Ser Ala Pro Ala Thr Pro Pro Lys Gln Leu Arg Gly Phe Ala Lys 665 670 675	2189		
45	ctc aag ctg gcg gcc ggc gcc agc ggg acc gcg acg ttt agt ctg cgc Leu Lys Leu Ala Ala Gly Ala Ser Gly Thr Ala Thr Phe Ser Leu Arg 680 685 690	2237		
50	agg cgc gac ctc agc tac tgg gac acc ggc cgc ggg cag tgg gtg gtg Arg Arg Asp Leu Ser Tyr Trp Asp Thr Gly Arg Gly Gln Trp Val Val 695 700 705	2285		
55	ccg gag ggc gag ttt ggc gtc tcg gtc ggg gcg agc tcg cgc gat atc Pro Glu Gly Glu Phe Gly Val Ser Val Gly Ala Ser Ser Arg Asp Ile 710 715 720 725	2333		
60	cgg ctg acg ggg agc ttc cga gta tgaaaggaaa tatcctgggg cggaatagttt Arg Leu Thr Gly Ser Phe Arg Val 730	2387		
65	ctgcttcttc gatggagtga actatatata ggtgcgcccg gacgccagag atatggacaa taac	2447	2451	
70	<210> 92 <211> 733 <212> PRT <213> Chrysosporium lucknowense			

EP 2 183 363 B1

<400> 92

5	Met	Thr	Leu	Gln	Ala	Phe	Ala	Leu	Leu	Ala	Ala	Ala	Ala	Leu	Val	Arg	1	5	10	15
10	Gly	Glu	Thr	Pro	Thr	Lys	Val	Pro	Arg	Asp	Ala	Pro	Arg	Gly	Ala	Ala	20	25	30	
15	Ala	Trp	Glu	Ala	Ala	His	Ser	Ser	Ala	Ala	Ala	Ala	Leu	Gly	Lys	Leu	35	40	45	
20	Ser	Gln	Gln	Asp	Lys	Ile	Asn	Ile	Val	Thr	Gly	Val	Gly	Trp	Asn	Lys	50	55	60	
25	Gly	Pro	Cys	Val	Gly	Asn	Thr	Pro	Ala	Ile	Ser	Ser	Ile	Asn	Tyr	Pro	65	70	75	80
30	Gln	Leu	Cys	Leu	Gln	Asp	Gly	Pro	Leu	Gly	Val	Arg	Phe	Gly	Ser	Ser	85	90	95	
35	Ile	Thr	Ala	Phe	Thr	Pro	Gly	Ile	Gln	Ala	Ala	Ser	Thr	Trp	Asp	Val	100	105	110	
40	Asp	Leu	Ile	Arg	Gln	Arg	Gly	Glu	Tyr	Met	Gly	Ala	Glu	Phe	Lys	Gly	115	120	125	
45	Cys	Gly	Ile	His	Val	Gln	Leu	Gly	Pro	Val	Ala	Gly	Pro	Leu	Gly	Lys	130	135	140	
50	Val	Pro	Gln	Gly	Gly	Arg	Asn	Trp	Glu	Gly	Phe	Gly	Val	Asp	Pro	Tyr	145	150	155	160
55	Leu	Thr	Gly	Ile	Ala	Met	Ala	Glu	Thr	Ile	Glu	Gly	Ile	Gln	Ser	Ala	165	170	175	
60	Gly	Val	Gln	Ala	Thr	Ala	Lys	His	Tyr	Ile	Leu	Asn	Glu	Gln	Glu	Leu	180	185	190	
65	Asn	Arg	Glu	Thr	Met	Ser	Ser	Asn	Val	Asp	Asp	Arg	Thr	Leu	His	Glu	195	200	205	
70	Leu	Tyr	Leu	Trp	Pro	Phe	Ala	Asp	Ala	Val	His	Ser	Asn	Val	Ala	Ser	210	215	220	
75	Val	Met	Cys	Ser	Tyr	Asn	Lys	Ile	Asn	Gly	Thr	Trp	Ala	Cys	Glu	Asn	225	230	235	240

EP 2 183 363 B1

	Asp	Arg	Val	Leu	Asn	Val	Ile	Leu	Lys	Gln	Glu	Leu	Gly	Phe	Pro	Gly	
					245					250					255		
5	Tyr	Val	Met	Ser	Asp	Trp	Asn	Ala	Gln	His	Ser	Thr	Asp	Asp	Ala	Ala	
				260					265					270			
10	Asn	His	Gly	Met	Asp	Met	Thr	Met	Pro	Gly	Ser	Asp	Phe	Asn	Gly	Gly	
			275					280					285				
15	Thr	Ile	Leu	Trp	Gly	Pro	Gln	Leu	Asp	Ser	Ala	Val	Asn	Ser	Gly	Arg	
		290					295					300					
20	Val	Pro	Lys	Ser	Arg	Leu	Asp	Asp	Met	Val	Glu	Arg	Ile	Leu	Ala	Ala	
	305					310					315					320	
25	Trp	Tyr	Leu	Leu	Gly	Gln	Asp	Ser	Asn	Tyr	Pro	Ala	Ile	Asn	Ile	Gly	
					325					330					335		
30	Ala	Asn	Val	Gln	Gly	Asn	His	Lys	Glu	Asn	Val	Arg	Ala	Val	Ala	Arg	
				340					345					350			
35	Asp	Gly	Ile	Val	Leu	Leu	Lys	Asn	Asp	Asp	Gly	Ile	Leu	Pro	Leu	Lys	
			355					360					365				
40	Lys	Pro	Ala	Lys	Leu	Ala	Leu	Ile	Gly	Ser	Ala	Ala	Val	Val	Asn	Pro	
		370					375					380					
45	Gln	Gly	Leu	Asn	Ser	Cys	Gln	Asp	Gln	Gly	Cys	Asn	Lys	Gly	Ala	Leu	
		385				390					395					400	
50	Gly	Met	Gly	Trp	Gly	Ser	Gly	Ala	Val	Asn	Tyr	Pro	Tyr	Phe	Val	Ala	
					405					410					415		
55	Pro	Tyr	Asp	Ala	Leu	Lys	Ala	Arg	Ala	Gln	Glu	Asp	Gly	Thr	Thr	Val	
				420					425					430			
60	Ser	Leu	His	Asn	Ser	Asp	Ser	Thr	Ser	Gly	Val	Ala	Asn	Val	Ala	Ser	
			435					440					445				
65	Asp	Ala	Asp	Ala	Ala	Ile	Val	Val	Ile	Thr	Ala	Asp	Ser	Gly	Glu	Gly	
		450					455					460					
70	Tyr	Ile	Thr	Val	Glu	Gly	Ala	Ala	Gly	Asp	Arg	Leu	Asn	Leu	Asp	Pro	
	465					470					475					480	

EP 2 183 363 B1

	Trp	His	Asn	Gly	Asn	Glu	Leu	Val	Lys	Ala	Val	Ala	Ala	Ala	Asn	Lys
					485					490					495	
5	Asn	Thr	Ile	Val	Val	Val	His	Ser	Val	Gly	Pro	Ile	Ile	Leu	Glu	Thr
				500					505					510		
10	Ile	Leu	Ala	Thr	Glu	Gly	Val	Lys	Ala	Ile	Val	Trp	Ala	Gly	Leu	Pro
			515					520					525			
15	Ser	Gln	Glu	Asn	Gly	Asn	Ala	Leu	Val	Asp	Ile	Leu	Tyr	Gly	Leu	Ala
		530					535					540				
20	Ser	Pro	Ser	Gly	Lys	Leu	Val	Tyr	Thr	Ile	Ala	Lys	Arg	Glu	Gln	Asp
	545					550					555					560
25	Tyr	Gly	Thr	Ala	Val	Val	Arg	Gly	Asp	Asp	Thr	Phe	Pro	Glu	Gly	Leu
					565					570					575	
30	Phe	Val	Asp	Tyr	Arg	His	Phe	Asp	Lys	Glu	Asn	Ile	Glu	Pro	Arg	Tyr
				580					585					590		
35	Glu	Phe	Gly	Phe	Gly	Leu	Ser	Tyr	Thr	Asn	Phe	Thr	Tyr	Ala	Asp	Leu
			595					600					605			
40	Glu	Leu	Thr	Ser	Thr	Ala	Thr	Ala	Gly	Pro	Ala	Thr	Gly	Glu	Thr	Ile
		610					615					620				
45	Pro	Gly	Gly	Ala	Ala	Asp	Leu	Trp	Glu	Glu	Val	Ala	Thr	Val	Thr	Ala
	625					630					635					640
50	Thr	Ile	Thr	Asn	Ser	Gly	Gly	Val	Asp	Gly	Ala	Glu	Val	Ala	Gln	Leu
				645						650					655	
55	Tyr	Leu	Thr	Leu	Pro	Ser	Ser	Ala	Pro	Ala	Thr	Pro	Pro	Lys	Gln	Leu
				660					665					670		
60	Arg	Gly	Phe	Ala	Lys	Leu	Lys	Leu	Ala	Ala	Gly	Ala	Ser	Gly	Thr	Ala
			675					680					685			
65	Thr	Phe	Ser	Leu	Arg	Arg	Arg	Asp	Leu	Ser	Tyr	Trp	Asp	Thr	Gly	Arg
		690					695					700				
70	Gly	Gln	Trp	Val	Val	Pro	Glu	Gly	Glu	Phe	Gly	Val	Ser	Val	Gly	Ala
	705					710					715					720

EP 2 183 363 B1

Ser Ser Arg Asp Ile Arg Leu Thr Gly Ser Phe Arg Val
725 730

<210> 93

5 <211> 2199

<212> DNA

<213> Chrysosporium lucknowense

<400> 93

10

15

20

25

30

35

40

45

50

55

EP 2 183 363 B1

	atgacccttc aagcctttgc gctgctggcg gcggcgggcc ttgtacgggg cgagacccccg	60
	accaaggtcc ctgcgcacgc tccgagagga gctgctgctt gggaagctgc ccactcctca	120
5	gcagccgctg ccttgggaaa gctgtcccag caggacaaga tcaacatcgt gacgggctgc	180
	ggctggaaca aggggccctg cgtgggcaat actcccgcta tcagctccat caactacccg	240
10	cagctctgcc tccaagacgg acccctgggc gtccgcttcg gttccagcat caccgctttc	300
	actccaggca ttcaggccgc gtcgacgtgg gatgtggacc tgatccgaca acgaggcgag	360
	tacatgggcy ccgagttcaa ggggtgtggc atccacgtcc agctaggccc cgtggccggg	420
15	cccttgggta aggttccgca aggcggccgc aactgggagg gctttggcgt ggatccctac	480
	ctcaccggca ttgccatggc cgagaccatc gaggggatac agtcagcggg cgtgcaagcc	540
	accgccaagc attacatcct caacgagcag gagctcaacc gcgagaccat gagcagcaat	600
20	gtcgacgacc gcaccttgca cgagctgtac ctctggcctt ttgcagacgc tgttactct	660
	aacgtggcca gcgtcatgtg cagttacaac aagatcaacg gcacgtgggc gtgcgagaac	720
25	gaccgggtgc tgaacgtcat tctgaagcag gagcttgggt ttccgggcta tgtgatgagc	780
	gactggaacg cgcagcactc gaccgacgac gcggccaacc acggcatgga catgacgatg	840
	cccggcagcg acttcaacgg agggacgac ctctggggcc cgcagctcga cagcgccgtc	900
30	aacagcggcc ggggtgccc aa gtccgcctc gacgacatgg tcgagcgcat cctcgcggcg	960
	tggtaccttc tcggtcagga cagcaattac ccggctatca acatcggcgc caacgtacag	1020
35	ggcaaccaca aggagaacgt gcgggcggtc gcgcgcgatg gcatcgtgct cctcaagaac	1080
	gacgacggca tcctcccgt caagaagccc gccaaagctcg cctcatcgg ttcgcccgcc	1140
	gtcgtcaacc cgcagggact gaactcctgc caggatcagg gctgcaacaa aggcgcgctg	1200
40	ggcatgggat ggggggtccg ggcggtcaac taccgtact ttgtcgcgcc gtacgacgcg	1260
	ctcaaggccc gcgcccaga ggacgggacg acggtcagcc tgcacaactc ggacagcacg	1320
	agcgggggtg cgaacgtggc ctcgacgcg gacgcggcca tcgtgggtcat cacggccgac	1380
45	tcgggcgagg gttacatcac agtcgaaggc gccgccgggg accgggtgaa cctcgacccg	1440
	tggcacaacg gcaacgagct ggtaaaggcg gtggcgggcg ccaacaagaa caccatcgtc	1500
50	gtcgtgcaca gcgtcgggcc catcatcctc gagaccatcc tggcaaccga aggggtcaag	1560

55

EP 2 183 363 B1

gcgatttgtgt gggccggcct gccaaagccag gagaatggta acgcgctggt ggacatcctg 1620
 tacggtcttgg cctcgcctgc gggcaagctg gtctacacca tcgccaagcg agaacaggac 1680
 5 tacggcactg cggtcgttcg cggagacgac acatttccgg agggcctggt tgtcgactac 1740
 cggcactttg acaaggagaa catcgagccg cggtagcagt ttgggttttg gctgtcatac 1800
 10 accaacttca cgtacgccga cctcgaactc acctcgaccg ccacggcccg cccagcgacg 1860
 ggcgagacca tccccggcgg cgcggccgac ctctgggagg aggtggccac ggtcacggcg 1920
 accatcacca acagcggcgg cgtggacggc gccgaggtgg cccagctgta cctgacgttg 1980
 15 ccgtcgtcgg cgccggcgac cccgccaag cagctcccg gcttcgcaa gctcaagctg 2040
 gcggccggcg ccagcgggac cgcgacgtt agtctgcgca ggcgcgacct cagctactgg 2100
 gacaccggcc gcgggcagt ggtggtgccg gagggcgagt ttggcgcttc ggtcggggcg 2160
 20 agctcgcgcg atatccggct gacggggagc ttccgagta 2199

<210> 94

<211> 389

25 <212> PRT

<213> Chrysosporium lucknowense

<400> 94

30 Met Lys Ser Ser Ile Leu Ala Ser Val Phe Ala Thr Gly Ala Val Ala
 1 5 10 15
 Gln Ser Gly Pro Trp Gln Gln Cys Gly Gly Ile Gly Trp Gln Gly Ser
 35 20 25 30
 Thr Asp Cys Val Ser Gly Tyr His Cys Val Tyr Gln Asn Asp Trp Tyr
 40 35 40 45
 Ser Gln Cys Val Pro Gly Ala Ala Ser Thr Thr Leu Gln Thr Ser Thr
 50 50 55 60
 45 Thr Ser Arg Pro Thr Ala Thr Ser Thr Ala Pro Pro Ser Ser Thr Thr
 65 70 75 80
 Ser Pro Ser Lys Gly Lys Leu Lys Trp Leu Gly Ser Asn Glu Ser Gly
 50 85 90 95
 Ala Glu Phe Gly Glu Gly Asn Tyr Pro Gly Leu Trp Gly Lys His Phe
 55 100 105 110
 Ile Phe Pro Ser Thr Ser Ala Ile Gln Thr Leu Ile Asn Asp Gly Tyr
 115 120 125

EP 2 183 363 B1

	Asn	Ile	Phe	Arg	Ile	Asp	Phe	Ser	Met	Glu	Arg	Leu	Val	Pro	Asn	Gln	
	130						135					140					
5	Leu	Thr	Ser	Ser	Phe	Asp	Glu	Gly	Tyr	Leu	Arg	Asn	Leu	Thr	Glu	Val	
	145					150					155					160	
10	Val	Asn	Phe	Val	Thr	Asn	Ala	Gly	Lys	Tyr	Ala	Val	Leu	Asp	Pro	His	
					165					170					175		
15	Asn	Tyr	Gly	Arg	Tyr	Tyr	Gly	Asn	Val	Ile	Thr	Asp	Thr	Asn	Ala	Phe	
				180					185						190		
20	Arg	Thr	Phe	Trp	Thr	Asn	Leu	Ala	Lys	Gln	Phe	Ala	Ser	Asn	Ser	Leu	
			195					200					205				
25	Val	Ile	Phe	Asp	Thr	Asn	Asn	Glu	Tyr	Asn	Thr	Met	Asp	Gln	Thr	Leu	
	210						215					220					
30	Val	Leu	Asn	Leu	Asn	Gln	Ala	Ala	Ile	Asp	Gly	Ile	Arg	Ala	Ala	Gly	
	225					230					235					240	
35	Ala	Thr	Ser	Gln	Tyr	Ile	Phe	Val	Glu	Gly	Asn	Ala	Trp	Ser	Gly	Ala	
				245						250					255		
40	Trp	Ser	Trp	Asn	Thr	Thr	Asn	Thr	Asn	Met	Ala	Ala	Leu	Thr	Asp	Pro	
				260					265					270			
45	Gln	Asn	Lys	Ile	Val	Tyr	Glu	Met	His	Gln	Tyr	Leu	Asp	Ser	Asp	Ser	
			275					280					285				
50	Ser	Gly	Thr	His	Ala	Glu	Cys	Val	Ser	Ser	Asn	Ile	Gly	Ala	Gln	Arg	
	290						295					300					
55	Val	Val	Gly	Ala	Thr	Gln	Trp	Leu	Arg	Ala	Asn	Gly	Lys	Leu	Gly	Val	
	305					310					315					320	
60	Leu	Gly	Glu	Phe	Ala	Gly	Gly	Ala	Asn	Ala	Val	Cys	Gln	Gln	Ala	Val	
				325						330					335		
65	Thr	Gly	Leu	Leu	Asp	His	Leu	Gln	Asp	Asn	Ser	Glu	Val	Trp	Leu	Gly	
				340					345					350			
70	Ala	Leu	Trp	Trp	Ala	Ala	Gly	Pro	Trp	Trp	Gly	Asp	Tyr	Met	Tyr	Ser	
			355					360					365				

Phe Glu Pro Pro Ser Gly Thr Gly Tyr Val Asn Tyr Asn Ser Ile Leu
370 375 380

5 Lys Lys Tyr Leu Pro
385

Claims

- 10 1. An isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:
 - 15 a) a nucleic acid sequence encoding a protein consisting of the amino acid sequence SEQ ID NO:32;
 - b) a nucleic acid sequence encoding an amino acid sequence with xylanase activity that is at least 90% identical to a full length amino acid sequence of (a);
 - c) a nucleic acid sequence selected from the group consisting of: SEQ ID NO:31, and SEQ ID NO:33;
 - d) a nucleic acid sequence that is fully complementary to the nucleic acid sequences of (a), (b) or (c).
- 20 2. An isolated protein comprising a) an amino acid sequence encoded by the nucleic acid molecule of claim 1; b) a fusion protein comprising the isolated protein of (a) fused to a protein comprising an amino acid sequence that is heterologous to the isolated protein of (a); or c) an antibody or antigen binding fragment thereof that selectively binds to the protein of (a).
- 25 3. A composition for the degradation of a lignocellulosic material comprising at least one isolated protein of Claim 2 part (a).
4. A recombinant nucleic acid molecule comprising the isolated nucleic acid molecule of claim 1, operatively linked to at least one expression control sequence.
- 30 5. A recombinant host cell transfected with the nucleic acid molecule of claim 1.
6. A method for producing the protein of Claim 2 part (a), comprising culturing a cell that has been transfected with a nucleic acid molecule comprising a nucleic acid sequence encoding the protein, and expressing the protein with the transfected cell, optionally recovering the protein from the cell or from a culture comprising the cell.
- 35 7. A genetically modified microorganism comprising components suitable for degrading a lignocellulosic material to fermentable sugars, wherein the organism has been genetically modified to express at least one protein of Claim 2 part (a).
- 40 8. The genetically modified microorganism of Claim 7, wherein the organism has been genetically modified to express at least one additional enzyme; wherein the additional enzyme is an accessory enzyme selected from the group consisting of: cellulase, xylanase, ligninase, glucuronidase, arabinofuranosidase, arabinogalactanase, ferulic acid esterase, lipase, pectinase, glucomannase, amylase, laminarinase, xyloglucanase, galactanase, glucoamylase, pectate lyase, chitinase, exo- β -D-glucosaminidase, cellobiose dehydrogenase, and acetylxylan esterase.
- 45 9. The composition of claim 3, further comprising at least one additional enzyme for degrading a lignocellulosic material or a fragment of said enzyme that retains biological activity; a) wherein the multi-enzyme composition comprises at least one cellobiohydrolase, at least one xylanase, at least one endoglucanase, at least one β -glucosidase, at least one β -xylosidase, and at least one accessory enzyme; b) wherein between about 50% and about 70% of the enzymes in the composition are cellobiohydrolases; c) wherein between about 10% and about 30% of the enzymes in the composition are xylanases; d) wherein between about 5% and about 15% of the enzymes in the composition are endoglucanases; e) wherein between about 1% and about 5% of the enzymes in the composition are β -glucosidases; f) wherein between about 1% and about 3% of the enzymes in the composition are β -xylosidases; or g) wherein the composition comprises about 60% cellobiohydrolases, about 20% xylanases, about 10% endoglucanases, about 3% β -glucosidases, about 2% β -xylosidases, and about 5% accessory enzymes.
- 50 10. The multi-enzyme composition of Claim 9, wherein the composition comprises a) at least one first protein comprising an amino acid sequence that is at least 95% identical to an amino acid sequence selected from the group consisting

of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 and SEQ ID NO:11 and has cellobiohydrolase activity, or a fragment thereof that has cellobiohydrolase activity; b) at least one second protein comprising an amino acid sequence that is at least 95% identical to an amino acid sequence selected from the group consisting of: SEQ ID NO:14, SEQ ID NO:94, SEQ ID NO:17, SEQ ID NO:20 and SEQ ID NO:23 and has endoglucanase activity, or a fragment thereof that has endoglucanase activity; c) at least one third protein comprising an amino acid sequence that is at least 95% identical to an amino acid sequence selected from the group consisting of: SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38 and SEQ ID NO:41 and has xylanase activity, or a fragment thereof that has xylanase activity; d) at least one fourth protein comprising an amino acid sequence that is at least 95% identical to SEQ ID NO:44 and has β -glucosidase activity, or a fragment thereof that has β -glucosidase activity; and e) at least one fifth protein comprising an amino acid sequence that is at least 95% identical to an amino acid sequence selected from the group consisting of: SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:62, SEQ ID NO:65, SEQ ID NO:68, SEQ ID NO:71, SEQ ID NO:74, SEQ ID NO:77, SEQ ID NO:80, SEQ ID NO:83, SEQ ID NO:86, SEQ ID NO:89, and SEQ ID NO:92, and has hemicellulase or chitinase activity, or a fragment thereof that has hemicellulase or chitinase activity.

11. The multi-enzyme composition of any one of Claims 9 to 10, further comprising one or more accessory enzymes; wherein the accessory enzymes includes at least one enzyme selected from the group consisting of: cellulase, xylanase, ligninase, glucuronidase, arabinofuranosidase, arabinogalactanase, furilic acid esterase, lipase, pectinase, glucomannase, amylase, laminarinase, xyloglucanase, galactanase, glucoamylase, pectate lyase, chitinase, exo- β -D-glucosaminidase, cellobiose dehydrogenase, and acetylxylose esterase.

12. A method for degrading a lignocellulosic material to fermentable sugars, comprising contacting the lignocellulosic material with at least one isolated protein of Claim 2 part (a) or at least one multi-enzyme composition of any one of Claims 9 to 11.

13. A method for producing an organic substance, comprising:

- a) saccharifying a lignocellulosic material with a multi-enzyme composition of any one of Claims 9 to 11;
- b) fermenting the saccharified lignocellulosic material obtained with one or more fermenting microorganisms; and
- c) recovering the organic substance from the fermentation.

14. A method for degrading a lignocellulosic material consisting of distiller's dried grains or distiller's dried grains with solubles to sugars, the method comprising contacting the distiller's dried grains or distiller's dried grains with solubles with a multi-enzyme composition, whereby at least 10% of the fermentable sugars are liberated, and wherein the multi-enzyme composition is the multi-enzyme composition of any one of Claims 9 to 11.

15. The method of any one of Claims 12 to 14, further comprising recovering the contacted lignocellulosic material after the fermentable sugars are degraded.

16. A method of deinking or biobleaching paper or pulp, comprising contacting the paper or pulp with at least one isolated protein of claim 2 part (a) or at least one multi-enzyme composition of any one of claims 9 to 11.

17. A method for treating a fabric, wherein the treating is selected from the group consisting of stonewashing a fabric; enhancing the softness or feel of a fabric or depilling a fabric; restoring color to or brightening a fabric; biopolishing, defibrillating, bleaching, dyeing or desizing a fabric; wherein the method comprises contacting the fabric with at least one isolated protein of Claim 2 part (a) or at least one multi-enzyme composition of any one of Claims 9 to 11.

18. A detergent composition, comprising at least one isolated protein of Claim 2 part (a) or at least one multi-enzyme composition of any one of Claims 9 to 11, optionally comprising a surfactant.

Patentansprüche

1. Isoliertes Nukleinsäuremolekül, welches eine der folgenden Nukleinsäuresequenzen aufweist:

- a) eine Nukleinsäuresequenz, welche ein Protein kodiert, welches die Aminosäuresequenz SEQ ID NO:32 aufweist;

- b) eine Nukleinsäuresequenz, welche eine Aminosäuresequenz mit Xylanaseaktivität kodiert, welche mindestens zu 90% identisch mit einer Aminosäuresequenz voller Länge aus (a) ist;
 c) eine Nukleinsäuresequenz, welche aus Folgenden ausgewählt wird: SEQ ID NO:31 und SEQ ID NO:33;
 d) eine Nukleinsäuresequenz, welche vollständig komplementär zu der Nukleinsäuresequenz aus (a), (b) oder (c) ist.

2. Isoliertes Protein, welches Folgendes aufweist:

- a) eine Aminosäuresequenz, welche durch das Nukleinsäuremolekül nach Anspruch 1 kodiert ist;
 b) ein Fusionsprotein, welches das isolierte Protein aus (a) aufweist, welches mit einem Protein vereinigt ist, welches eine Aminosäuresequenz aufweist, welche heterolog zu dem isolierten Protein aus (a) ist; oder
 c) ein Antikörper oder ein antigenbindender Teil davon, welcher das Protein aus (a) selektiv bindet.

3. Zusammensetzung zum Zersetzen eines Lignocellulose-Materials, welche mindestens ein isoliertes Protein nach Anspruch 2 Abschnitt (a) aufweist.

4. Rekombinantes Nukleinsäuremolekül, welches das isolierte Nukleinsäuremolekül nach Anspruch 1 aufweist, welches operativ mit mindestens einer Expressionssteuersequenz verbunden ist.

5. Rekombinante Wirtszelle, welche mit dem Nukleinsäuremolekül aus Anspruch 1 transfiziert ist.

6. Verfahren zum Herstellen des Proteins nach Anspruch 2 Abschnitt (a), welches folgende Schritte beinhaltet: Kultivieren einer Zelle, welche mit einem Nukleinsäuremolekül transfiziert worden ist, welches eine Nukleinsäuresequenz aufweist, welche das Protein kodiert, und Exprimieren des Proteins mit der transfizierten Zelle und optional folgenden Schritt aufweist: Zurückgewinnen des Proteins aus den Zellen oder aus einer Kultur, welche die Zelle aufweist.

7. Genetisch modifizierter Mikroorganismus, welcher Bestandteile aufweist, welche geeignet sind zum Zersetzen eines Lignocellulose-Materials in fermentierbare Zucker, wobei der Organismus genetisch modifiziert worden ist, um mindestens ein Protein nach Anspruch 2 Abschnitt (a) zu exprimieren.

8. Genetisch modifizierter Mikroorganismus nach Anspruch 7, wobei der Organismus genetisch modifiziert worden ist, um mindestens ein zusätzliches Enzym zu exprimieren; wobei das zusätzliche Enzym ein akzessorisches Enzym ist, welches aus Folgenden ausgewählt wird: Cellulase, Xylanase, Ligninase, Glucuronidase, Arabinofuranosidase, Arabinogalactanase, Ferulasäure-Esterase, Lipase, Pektinase, Glucomannase, Amylase, Laminarinase, Xyloglucanase, Galactanase, Glucoamylase, Pektatlyase, Chitinase, exo- β -D-Glucosaminidase, Cellobiose Dehydrogenase und Acetylxylan-Esterase.

9. Zusammensetzung nach Anspruch 3, welche ferner mindestens ein zusätzliches Enzym aufweist zum Zersetzen eines Lignocellulose-Materials oder einen Teil des Enzyms, welcher biologische Aktivität beinhaltet;

- a) wobei die Multi-Enzym-Zusammensetzung mindestens eine Cellobiohydrolase, mindestens eine Xylanase, mindestens eine Endoglucanase, mindestens eine β -Glucosidase, mindestens eine β -Xylosidase, und mindestens ein akzessorisches Enzym aufweist;
 b) wobei zwischen etwa 50% und etwa 70% der Enzyme in der Zusammensetzung Cellobiohydralasen sind;
 c) wobei zwischen etwa 10% und etwa 30% der Enzyme in der Zusammensetzung Xylanasen sind;
 d) wobei zwischen etwa 5% und etwa 15% der Enzyme in der Zusammensetzung Endoglucanasen sind;
 e) wobei zwischen etwa 1% und etwa 5% der Enzyme in der Zusammensetzung β -Glucosidasen sind;
 f) wobei zwischen etwa 1% und etwa 3% der Enzyme in der Zusammensetzung β -Xylosidasen sind;
 g) wobei die Zusammensetzung etwa 60% Cellobiohydrolasen, etwa 20% Xylanasen, etwa 10% Endoglucanasen, etwa 3% β -Glucosidasen, etwa 2% β -Xylosidasen und etwa 5% akzessorische Enzyme aufweist.

10. Multi-Enzym-Zusammensetzung nach Anspruch 9, wobei die Zusammensetzung Folgendes aufweist:

- a) mindestens ein erstes Protein, welches eine Aminosäuresequenz aufweist, welche mindestens zu 95% identisch mit einer Aminosäuresequenz ist, welche aus Folgenden ausgewählt wird: SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 und SEQ ID NO:11 und Cellobiohydrolasenaktivität aufweist, oder einen Teil davon aufweist, welcher Cellobiohydrolasenaktivität aufweist;

b) mindestens ein zweites Protein, welches eine Aminosäuresequenz aufweist, welche mindestens zu 95% identisch mit einer Aminosäuresequenz ist, welche aus Folgenden ausgewählt wird: SEQ ID NO:14, SEQ ID NO:94, SEQ ID NO:17, SEQ ID NO:20 und SEQ ID NO:23 und Endoglucanaseaktivität aufweist, oder einen Teil davon aufweist, welcher Endoglucanaseaktivität aufweist;

c) mindestens ein drittes Protein, welches eine Aminosäuresequenz aufweist, welche mindestens zu 95% identisch mit einer Aminosäuresequenz ist, welche aus Folgenden ausgewählt wird: SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38 und SEQ ID NO:41 und Xylanaseaktivität aufweist, oder einen Teil davon aufweist, welcher Xylanaseaktivität aufweist;

d) mindestens ein viertes Protein, welches eine Aminosäuresequenz aufweist, welche mindestens zu 95% identisch mit SEQ ID NO:44 ist und β -Glucosidaseaktivität aufweist, oder einen Teil davon aufweist, welcher β -Glucosidaseaktivität aufweist; und

e) mindestens ein fünftes Protein, welches eine Aminosäuresequenz aufweist, welche mindestens zu 95% identisch mit einer Aminosäuresequenz ist, welche aus Folgenden ausgewählt wird: SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:62, SEQ ID NO:65, SEQ ID NO:68, SEQ ID NO:71, SEQ ID NO:74, SEQ ID NO:77, SEQ ID NO:80, SEQ ID NO:83, SEQ ID NO:86, NO:89 und SEQ ID NO:92 und Hemicellulase- oder Chitinaseaktivität aufweist, oder einen Teil davon aufweist, welcher Hemicellulase- oder Chitinaseaktivität aufweist.

11. Multi-Enzym-Zusammensetzung nach einem der Ansprüche 9 bis 10, welche ferner ein oder mehrere akzessorische Enzyme aufweist; wobei die akzessorischen Enzyme mindestens ein Enzym aufweisen, welches aus Folgenden ausgewählt wird: Cellulase, Xylanase, Ligninase, Glucuronidase, Arabinofuranosidase, Arabinogalactanase, Furiclsäure-Esterase, Lipase, Pektinase, Glucomannase, Amylase, Laminarinase, Xyloglucanase, Galactanase, Glucoamylase, Pektatlyase, Chitinase, exo- β -D-Glucosaminidase, Cellobiose Deyhydrogenase und Acetylxytan-Esterase.

12. Verfahren zum Zersetzen eines Lignocellulose-Materials in fermentierbare Zucker, welches ein Verbinden des Lignocellulose-Materials mit mindestens einem isolierten Protein nach Anspruch 2 Abschnitt (a) oder mit mindestens einer Multi-Enzym-Zusammensetzung nach einem der Ansprüche von 9 bis 11 aufweist.

13. Verfahren zum Erzeugen einer organischen Substanz, welches folgende Schritte aufweist:

a) Verzuckern eines Lignocellulose-Materials mit einer Multi-Enzym-Zusammensetzung nach einem der Ansprüche 9 bis 11;

b) Fermentieren des verzuckerten Lignocellulose-Materials, welches mit Hilfe eines oder mehrerer Fermentierungsmikroorganismen erreicht wird; und

c) Zurückgewinnen der organischen Substanz von der Fermentation.

14. Verfahren zum Zersetzen eines Lignocellulose-Materials, welches Trockenschlempe (DDG) oder Trockenschlempe mit löslichem Anteil (DDGS) aufweist, in Zucker, wobei das Verfahren ein Verbinden der Trockenschlempe oder der Trockenschlempe mit löslichem Anteil mit einer Multi-Enzym-Zusammensetzung aufweist, wobei mindestens 10% der fermentierbaren Zucker gelöst sind, und wobei die Multi-Enzym-Zusammensetzung die Multi-Enzym-Zusammensetzung nach einem der Ansprüche 9 bis 11 ist.

15. Verfahren nach einem der Ansprüche 12 bis 14, welches ferner das Zurückgewinnen des verbundenen Lignocellulose-Materials aufweist, nachdem die fermentierbaren Zucker zersetzt sind.

16. Verfahren zum Deinking oder zum biologischen Bleichen von Papier oder Zellstoff, welches ein Verbinden des Papiers oder des Zellstoffes mit mindestens einem Protein nach Anspruch 2 Abschnitt (a) oder mindestens einer Multi-Enzym-Zusammensetzung nach einem der Ansprüche 9 bis 11 aufweist.

17. Verfahren zum Behandeln eines Gewebes, wobei das Behandeln aus den Folgenden ausgewählt wird: Stonewash-Behandeln eines Gewebes, Verbessern der Weichheit oder der Haptik eines Gewebes oder Entfusseln eines Gewebes; Wiederherstellen der Farbe oder Aufhellen eines Gewebes; Biopolieren, Defibrillieren, Bleichen, Färben oder Entschlichten eines Gewebes, wobei das Verfahren das Verbinden des Gewebes mit mindestens einem isolierten Protein nach Anspruch 2 Abschnitt (a) oder mit mindestens einer Multi-Enzym-Zusammensetzung nach einem der Ansprüche 9 bis 11 aufweist.

18. Reinigungsmittel-Zusammensetzung, welche mindestens ein isoliertes Protein nach Anspruch 2 Abschnitt (a) oder

mindestens eine Multi-Enzym-Zusammensetzung nach einem der Ansprüche 9 bis 11 und optional ein Tensid aufweist.

5 Revendications

1. Une molécule d'acide nucléique isolée comprenant une séquence d'acide nucléique choisie dans le groupe constitué par:

- 10 a) une séquence d'acide nucléique codant pour une protéine consistant en la séquence d'acide aminé SEQ ID NO: 32;
 b) une séquence d'acide nucléique codant pour une séquence d'acide aminé présentant une activité de xylanase qui est au moins identique à 90% à la longueur totale de la séquence d'acides aminés de (a);
 15 c) une séquence d'acide nucléique choisie dans le groupe consistant en SEQ ID NO: 31 et SEQ ID NO: 33;
 d) une séquence d'acide nucléique qui est totalement complémentaire aux séquences d'acides nucléiques de (a), (b) et (c).

2. Une protéine isolée comprenant

- 20 a) une séquence d'acide aminé codée par la molécule d'acide nucléique selon la revendication 1;
 b) une protéine de fusion comprenant la protéine isolée de (a) fusionnée à une protéine comprenant une séquence d'acide aminé qui est hétérologue à la protéine isolée de (a) ou
 c) un anticorps ou un fragment de liaison d'antigène en dérivant qui est lié sélectivement à la protéine (a).

25 3. Une composition destinée à la dégradation d'une matière lignocellulosique comprenant au moins une protéine isolée selon la partie (a) de la revendication 2.

4. Une molécule d'acide nucléique recombinante comprenant la molécule d'acide nucléique isolée selon la revendication 1, lié de manière fonctionnelle à au moins une séquence de contrôle d'expression.

30 5. Une cellule hôte recombinante transfectée avec la molécule d'acide nucléique selon la revendication 1.

6. Un procédé de production de la protéine selon la partie (a) de la revendication 2, comprenant la culture d'une cellule qui a été transfectée avec une molécule d'acide nucléique comprenant une séquence d'acide nucléique codant pour la protéine, et l'expression de la protéine par la cellule transfectée et éventuellement la récupération de la protéine à partir de la cellule ou d'une culture comprenant la cellule.

7. Un microorganisme modifié génétiquement comprenant des composants appropriés pour dégrader un matériau lignocellulosique en sucres fermentescibles, dans lequel l'organisme a été génétiquement modifié pour exprimer au moins une protéine de la partie (a) de la revendication 2.

8. Le microorganisme modifié génétiquement selon la revendication 7, dans lequel l'organisme a été génétiquement modifié pour exprimer au moins une enzyme supplémentaire, dans lequel l'enzyme supplémentaire est une enzyme accessoire choisie dans le groupe constitué de cellulase, xylanase, ligninase, glucuronidase, arabinofuranosidase, arabinogalactanase, estérase d'acide férulique, lipase, pectinase, glucomannase, amylase, laminarinase, xyloglucanase, galactanase, glucoamylase, pectate lyase, chitinase, exo- β -D-glucosaminidase, cellobiose déshydrogénase et acétylxylane estérase.

9. La composition selon la revendication 3, comprenant en outre au moins une enzyme supplémentaire pour la dégradation d'une matière lignocellulosique ou un fragment de ladite enzyme qui présente une activité biologique,

- a) dans laquelle la composition multi-enzymatique comprend au moins une cellobiohydrolase, au moins une xylanase, au moins une endoglucanase, au moins une β -glucosidase, au moins une β -xylosidase, et au moins une enzyme accessoire;
 55 b) dans laquelle entre environ 50% et environ 70% des enzymes dans la composition sont des cellobiohydrolases;
 c) dans laquelle entre environ 10% et environ 30% des enzymes dans la composition sont des xylanases;
 d) dans laquelle entre environ 5% et environ 15% des enzymes dans la composition sont des endoglucanases;

- e) dans laquelle entre environ 1% et environ 5% des enzymes dans la composition sont des β -glucosidase;
 f) dans laquelle entre environ 1% et environ 3% des enzymes dans la composition sont des β -xylosidases; ou
 g) dans laquelle la composition comprend environ 60% de cellobiohydrolases, environ 20% de xylanases, environ 10% d'endoglucanases, environ 3% de β -glucosidases, environ 2% de β -xylosidases, et environ 5% d'enzymes accessoires

10. La composition multi-enzymatique selon la revendication 9, dans lequel la composition comprend:

- a) au moins une première protéine comprenant une séquence d'acides aminés qui est au moins identique à 95% à une séquence d'acides aminés choisie dans le groupe constitué par: SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 et SEQ ID NO:11 et présente une activité cellobiohydrolase, ou un fragment en dérivant présentant une activité cellobiohydrolase;
 b) au moins une seconde protéine comprenant une séquence d'acides aminés qui est au moins identique à 95% à une séquence d'acides aminés choisie dans le groupe constitué par: SEQ ID NO:14, SEQ ID NO:94, SEQ ID NO:17, SEQ ID NO:20 et SEQ ID NO:23 et présente une activité d'endoglucanase, ou un fragment de celui-ci qui a une activité d'endoglucanase; et
 c) au moins une troisième protéine comprenant une séquence d'acides aminés qui est au moins identique à 95% à une séquence d'acides aminés choisie dans le groupe constitué par: SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38 et SEQ ID NO:41 et présente une activité de xylanase, ou un fragment en dérivant présentant une activité de xylanase;
 d) au moins une quatrième protéine comprenant une séquence d'acides aminés qui est au moins identique à 95% à SEQ ID NO:44 et présente une activité de β -glucosidase, ou un fragment en dérivant présentant une activité de β -glucosidase;
 e) au moins une cinquième protéine comprenant une séquence d'acides aminés qui est au moins identique à 95% à une séquence d'acides aminés choisie dans le groupe constitué par: SEQ ID NO: 47, SEQ ID NO: 50, SEQ ID NO: 53, SEQ ID NO: 56, SEQ ID NO: 59, SEQ ID NO: 62, SEQ ID NO: 65, SEQ ID NO: 68, SEQ ID NO: 71, SEQ ID NO: 74, SEQ ID NO: 77, SEQ ID NO: 80, SEQ ID NO: 83, SEQ ID NO: 86, SEQ ID NO: 89 et SEQ ID NO: 92, et présente une activité de hémicellulase ou de chitinase, ou un fragment en dérivant présentant une activité d'hémicellulase ou de chitinase.

11. La composition multi-enzymatique selon une quelconque des revendications 9 à 10, comprenant une ou plusieurs enzymes accessoires, dans laquelle les enzymes accessoires incluent au moins une enzyme choisie dans le groupe constitué de cellulase, xylanase, ligninase, glucuronidase, arabinofuranosidase, arabinogalactanase, estérase d'acide férulique, lipase, pectinase, glucomannase, amylase, laminarinase, xyloglucanase, galactanase, glucoamylase, pectate lyase, chitinase, exo- β -D-glucosaminidase, cellobiose déshydrogénase et acétylxylane estérase.

12. Un procédé de dégradation d'une matière lignocellulosique en sucres fermentescibles, qui comprend la mise en contact de la matière lignocellulosique avec au moins une protéine isolée selon la partie (a) de la revendication 2 ou au moins une composition multi-enzymatique selon une quelconque des revendications 9 à 11.

13. Procédé de production d'une substance organique, comprenant:

- a) la saccharification d'une matière lignocellulosique avec une composition multi-enzymatique selon l'une quelconque des revendications 9 à 11;
 b) la fermentation de la matière lignocellulosique saccharifiée obtenue avec un ou plusieurs microorganismes en fermentation; et
 c) la récupération de la substance organique à partir de la fermentation.

14. Un procédé pour dégrader un matériau lignocellulosique comprenant des grains séchés de distillerie ou des grains séchés de distillerie additionnés de composés solubles dans les sucres, le procédé comprenant la mise en contact des grains séchés de distillerie ou des grains séchés de distillerie additionnés de composés solubles avec une composition d'enzyme multiples, de sorte qu'au moins 10% des sucres fermentescibles sont libérés, et dans lequel la composition multi-enzymatique est la composition multi-enzymatique selon l'une quelconque des revendications 9 à 11.

15. Un procédé selon l'une quelconque des revendications 12 à 14, comprenant en outre la récupération de la matière lignocellulosique contactée après que les sucres fermentescibles sont dégradés.

16. Un procédé de désencrage ou bio-blanchiment de papier ou de pâte à papier, comprenant la mise en contact du papier ou de la pâte avec au moins une protéine isolée selon la partie (a) de la revendication 2 ou au moins une composition multi-enzymatique selon l'une quelconque des revendications 9 à 11.

5 **17.** Un procédé de traitement d'un tissu, dans lequel le traitement est choisi dans le groupe consistant en délavage d'un tissu, améliorer la souplesse ou le toucher d'un tissu ou déboulochage d'un tissu, restaurer la couleur ou azurer un tissu, bio-polissage, défibrillation, blanchiment, teinture ou désencollage d'un tissu, dans lequel le procédé comprend la mise en contact du tissu avec au moins une protéine isolée selon la partie (a) de la revendication 2 ou au moins une composition multi-enzyme selon l'une quelconque des revendications 9 à 11.

10 **18.** Une composition détergente, comprenant au moins une protéine isolée selon la partie (a) de la revendication 2 ou au moins une composition multi-enzyme selon l'une quelconque des revendications 9 à 11, comprenant de façon optionnelle un surfactant.

15

20

25

30

35

40

45

50

55




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9








Figure 10

**Figure 11**

**Figure 12**

**Figure 13**

**Figure 14**




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21





Figure 22

**Figure 23**




Figure 24




Figure 25




Figure 26

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9815633 A1 [0010]
- WO 0179507 A2 [0011]
- US 6015707 A [0086] [0087] [0224]
- US 6573086 B [0087] [0224]
- US 5605793 A, Stemmer [0211]
- US 4940838 A [0235]
- US 5464763 A [0235]
- US 20030157595 A [0255]
- US 4435307 A [0257]
- US 4443355 A [0257]
- US 4661289 A [0257]
- US 4479881 A [0257]
- US 5120463 A [0257]
- JP 50132269 A [0259]
- GB 2094826 A [0259]
- US 2974001 A, Windbichtler [0259]
- WO 11833133 A [0562]

Non-patent literature cited in the description

- GUSAKOV ALEXANDER V et al. *ENZYME AND MICROBIAL TECHNOLOGY*, 06 January 2005, vol. 36 (1), 57-69 [0008]
- GUSAKOV ALEXANDER V et al. *BIOTECHNOLOGY AND BIOENGINEERING*, 01 August 2007, vol. 97 (5), 1028-1038 [0009]
- Carbohydrate-active enzymes: an integrated database approach. COUTINHO, P.M. ; HENRISSAT, B. Recent Advances in Carbohydrate Bioengineering. The Royal Society of Chemistry, 1999, 3-12 [0103] [0163]
- FLIPPIN et al. *Appl. Microbiol. Biotechnol.*, 1993, vol. 40, 318 [0132]
- BAMETT et al. *Biotechnol.*, 1991, vol. 9, 562 [0133]
- GORDILLO et al. *Mycol. Res.*, 2006, vol. 110, 1129 [0140]
- MARGOLLES-CLARK et al. *Eur. J. Biochem.*, 1996, vol. 237, 553 [0144]
- EGANA et al. *Biotechnol. Appl. Biochem.*, 1996, vol. 24 (1), 33 [0145]
- BAUER et al. *Proc. Natl. Acad. Sci. U.S.A.*, 2006, vol. 103, 11417 [0146]
- TSUJIBO et al. *Biosci. Biotechnol. Biochem.*, 2002, vol. 66, 434 [0148] [0149]
- O'SHANNESY et al. *Anal. Biochem.*, 1993, vol. 212, 457-468 [0160]
- SCHUSTER et al. *Nature*, 1993, vol. 365, 343-347 [0160]
- ALTSCHUL, S.F. ; MADDEN, T.L. ; SCHÄFFER, A.A. ; ZHANG, J. ; ZHANG, Z. ; MILLER, W. ; LIPMAN, D.J. Gapped BLAST and PSI-BLAST: a new generation of protein database search program. *Nucleic Acids Res.*, 1997, vol. 25, 3389-3402 [0163]
- TATUSOVA ; MADDEN. Blast 2 sequences - a new tool for comparing protein and nucleotide sequences. *FEMS Microbiol Lett.*, 1999, vol. 174, 247-250 [0165]
- SAMBROOK et al. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Labs Press, 1989 [0170] [0228]
- SAMBROOK. *MOLECULAR CLONING: A LABORATORY MANUAL* [0170]
- MEINKOTH et al. *Anal. Biochem.*, 1984, vol. 138, 267-284 [0170]
- MEINKOTH. *ANAL. BIOCHEM.* [0170]
- *Industrial Enzymology*. Stockton Press, 1996, 267-272 [0200]
- *Immobilized Enzymes*. Halsted Press, 1978 [0200]
- *Enzymes and Immobilized Cells in Biotechnology*. Benjamin/Cummings Publishing Co., Inc, 1985 [0200]
- *Applied Biochemistry and Bioengineering*. Academic Press, 1983, vol. 4 [0200]
- MINSHULL ; STEMMER. *Curr. Opin. Chem. Biol.*, 1999, vol. 3, 284-290 [0211]
- STEMMER. *P.N.A.S. USA*, 1994, vol. 91, 10747-10751 [0211]
- SAMBROOK ; RUSSEL. *Molecular Cloning: A Laboratory Manual*. 2001 [0228]
- *Procedures for Introducing Foreign DNA into Plants*. MIKI et al. *Methods in Plant Molecular Biology and Biotechnology*. CRC Press, Inc, 1993, 67-88 [0234]
- *Vectors for Plant Transformation*. GRUBER et al. *Methods in Plant Molecular Biology and Biotechnology*. CRC Press, Inc, 1993, 89-119 [0234]
- HORSCH et al. *Science*, 1985, vol. 227, 1229 [0235]
- KADO, C.I. *Crit. Rev. Plant. Sci.*, 1991, vol. 10, 1 [0235]
- MOLONEY et al. *Plant Cell Reports*, 1989, vol. 8, 238 [0235]
- SANFORD et al. *Part. Sci. Technol.*, 1987, vol. 5, 27 [0236]

- **SANFORD, J.C.** *Trends Biotech.*, 1988, vol. 6, 299 [0236]
- **SANFORD, J.C.** *Physiol. Plant*, 1990, vol. 79, 206 [0236]
- **KLEIN et al.** *Biotechnology*, 1992, vol. 10, 268 [0236]
- **ZHANG et al.** *Bio/Technology*, 1991, vol. 9, 996 [0237]
- **DESHAYES et al.** *EMBO J.*, 1985, vol. 4, 2731 [0237]
- **CHRISTOU et al.** *Proc Natl. Acad. Sci. USA*, 1987, vol. 84, 3962 [0237]
- **HAIN et al.** *Mol. Gen. Genet.*, 1985, vol. 199, 161 [0237]
- **DRAPER et al.** *Plant Cell Physiol.*, 1982, vol. 23, 451 [0237]
- **DONN et al.** *Abstracts of VIIth International Congress on Plant Cell and Tissue Culture IAPTC*, 1990, vol. A2-38, 53 [0237]
- **D'HALLUIN et al.** *Plant Cell*, 1992, vol. 4, 1495-1505 [0237]
- **SPENCER et al.** *Plant Mol. Biol.*, 1994, vol. 24, 51-61 [0237]
- **KOHLER ; MILSTEIN.** *Nature*, 1975, vol. 256, 495-497 [0241]
- **BESTE et al.** *Proc. Natl. Acad. Sci.*, 1999, vol. 96, 1898-1903 [0241]
- **WYMAN et al.** *Bioresource Tech.*, 2005, vol. 96, 1959 [0248]
- **WYMAN et al.** *Bioresource Tech.*, 2005, vol. 96, 2026 [0248]
- Pretreatment of biomass. **HSU.** *Handbook on Bioethanol: Production and Utilization*. 1996, 179-212 [0248]
- **MOSIER et al.** *Bioresource Tech.*, 2005, vol. 96, 673 [0248]
- **LEISOLA ; LINKO.** *Analytical Biochemistry*, 1976, vol. 70, 592 [0259]
- Determination Of The Solubilizing Activity Of A Cellulase Complex With Dyed Substrates. Blum & Stahl--Enzymic Degradation Of Cellulose Fibers. Reports of the Shizuoka Prefectural Hamamatsu Textile Industrial Research Institute, 1985 [0259]
- **BHATAWADEKAR.** *Journal of the Textile Association*, May 1983, 83-86 [0259]
- **YANO et al.** *Biosci Biotechnol Biochem.*, 2006, vol. 70, 1754 [0265]
- **KRAMER et al.** *Inspect Biochem Mol Biol.*, 1997, vol. 27, 887 [0266]
- **REESE et al.** *Nature*, 2007, vol. 447, 92 [0267]
- *Trichoderma & Gliocladium. Enzymes, biological control and commercial applications.* Taylor & Francis Ltd, 1998, vol. 2, 393 [0268]
- **HELMUT UHLIG ; ELFRIEDE M. LINSMAIER-BEDNAR.** *Industrial enzymes and their applications.* John Wiley & Sons, Inc, 1998, 454 [0268]
- **HAHN-HAGERDAL, B. ; GALBE, M. ; GORWA-GRAUSLUND ; M.F. LIDÉN ; ZACCHI, G.** Bio-ethanol - the fuel of tomorrow from the residues of today. *Trends in Biotechnology.*, 2006, vol. 24 (12), 549-556 [0268]
- **MIELENZ, J.R.** Ethanol production from biomass: technology and commercialization status. *Current Opinion in Microbiology*, 2001, vol. 4, 324-329 [0268]
- **HIMMEL, M.E. ; RUTH, M.F. ; WYMAN, C.E.** Cellulase for commodity products from cellulosic biomass. *Current Opinion in Biotechnology*, 1999, vol. 10, 358-364 [0268]
- **SHEEHAN, J. ; HIMMEL, M.** Enzymes, energy, and the environment: a strategic perspective on the U.S. Department of Energy's Research and Development Activities for Bioethanol. *Biotechnology Progress.*, 1999, vol. 15, 817-827 [0268]
- **GALANTE, Y.M. ; FORMANTICI, C.** Enzyme applications in detergency and in manufacturing industries. *Current Organic Chemistiy.*, 2003, vol. 7, 1399-1422 [0268]
- **BAJPAI, P. ; BAJPAI, P.K.** Deinking with enzymes: a review. *TAPPI Journal.*, 1998, vol. 81 (12), 111-117 [0268]
- Use of cellulases in pulp and paper applications. In: "Carbohydrates from *Trichoderma reesei* and other microorganisms. **VIKARI, L. ; PERE, J. ; SUURNÄKKI, A. ; OKSANEN, T. ; BUCHERT, J.** Structure, Biochemistry, Genetics and Applications. The Royal Society of Chemistry, 1998, 245-254 [0268]
- **ROLLER, S. ; DEA, I.C.M.** Biotechnology in the production and modification of biopolymers for foods. *Critical Reviews in Biotechnology*, 1992, vol. 12 (3), 261-277 [0268]
- **LOWRY OH ; ROSEBROUGH NJ ; FARR AL ; RANDALL RJ.** *J. Biol. Chem.*, 1951, vol. 193, 265-275 [0313]
- **BEREZIN IV ; RABINOVICH ML ; SINITSYN AP.** Study of applicability of quantitative kinetic spectrophotometric method for glucose determination. *Biokhimiya (Moscow)*, 1977, vol. 42, 1631-1636 [0406]
- **GHOSE TK.** Measurement of cellulase activities. *Pure Appl. Chem*, 1987, vol. 59, 257-268 [0406]
- **GUSAKOV AV ; SALANOVICH TN ; ANTONOV AI ; USTINOV BB ; OKUNEV ON ; BURLINGAME R ; EMALFARB M ; BAEZ M ; SINITSYN AP.** Design of highly efficient cellulase mixtures for enzymatic hydrolysis of cellulose. *Biotechnol. Bioeng*, 2007, <http://www3.interscience.wiley.com/cgi-bin/abstract/114064699> [0406] [0425]
- **LOWRY OH ; ROSEBROUGH NJ ; FARR AL ; RANDALL RJ.** Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*, 1951, vol. 193, 265-275 [0406] [0425]

- Methods of investigation and properties of cellulolytic enzymes. **SINITSYN AP ; CHEMOGLAZOV VM ; GUSAKOV AV.** Biotechnology Series. VINITI Press, 1990, vol. 25, 220 [0406] [0425]
- **SOMOGYI M.** Notes on sugar determination. *J. Biol. Chem.*, 1952, vol. 195, 19-23 [0406] [0425]
- **BEREZIN IV ; RABINOVICH ML ; SINITSYN AP.** Study of applicability of quantitative kinetic spectrophotometric method for glucose determination. *Biokhimiya (Moscow)*, 1977, vol. 42, 1631-1636 [0425]
- **A.GUSAKOV ; A.SINITSYN ; S.GRISHUTIN ; D.TIKHOMIROV ; D.SHOOK ; D.SHEER ; M.EMALFARB.** Microassays to control the results of cellulase treatment of denim fabrics. *Textile Chemist and Colorist and American Dyestuff Reporter*, 2000, vol. 32 (5), 42 [0486] [0497]
- **KABEL et al.** *Carbohydr. Polym.*, 2002, vol. 50, 191 [0509]

GOMBAXILANÁZ

Szabadalmi igénypontok

1. Izolált nukleinsav-molekula, amely a következők alkotta csoportból választott nukleinsav-szekvenciát tartalmaz:

- a) SEQ ID NO. 32 szerinti aminosav-szekvencia alkotta fehérjét kódoló nukleinsav-szekvencia;
- b) az (a) szerinti, teljes hosszúságú aminosav-szekvenciával legalább 90%-ban azonos, xilanázaktivitású aminosav-szekvenciát kódoló nukleinsav-szekvencia,
- c) a SEQ ID NO. 31 és SEQ ID NO. 33 szerinti szekvenciák alkotta csoportból választott nukleinsav-szekvencia;
- d) nukleinsav-szekvencia, amely teljesen komplementer az (a), (b) vagy (c) szerinti nukleinsav-szekvenciával.

2. Izolált fehérje, amely a) az 1. igénypont szerinti nukleinsav-molekula által kódolt aminosav-szekvenciát; b) az (a) szerinti izolált fehérjét az (a) szerinti izolált fehérjéhez képest heterológ aminosav-szekvenciát tartalmazó fehérjéhez fuzionáltan tartalmazó fúziós fehérjét; vagy c) antitestet vagy annak antigénkötő fragmentumát, amely az (a) szerinti fehérjéhez szelektíven kötődik.

3. Készítmény lignocellulóz-anyag lebontására, amely legalább egy, a 2. igénypont (a) pontja szerinti izolált fehérjét tartalmaz.

4. Rekombináns nukleinsav-molekula, amely az 1. igénypont szerinti, izolált nukleinsav-molekulát tartalmazza legalább egyetlen, expressziót kontrolláló szekvenciához működőképesen kapcsolt formában.

5. Rekombináns gazdasejt, amely az 1. igénypont szerinti nukleinsav-molekulával transzfektált.

6. Eljárás a 2. igénypont (a) pontja szerinti fehérje előállítására, amely tartalmazza a fehérjét kódoló nukleinsav-szekvenciát tartalmazó nukleinsav-molekulával transzfektált sejt tenyésztését és a fehérje expresszálasát a transzfektált sejttel, adott esetben a fehérje kinyerését a sejtből vagy a sejtet tartalmazó tenyészetből.

7. Genetikailag módosított mikroorganizmus, amely lignocellulóz anyag fermentálható cukrokká történő lebontására alkalmas összetevőket tartalmaz, ahol az organizmus 2. igénypont (a) pontja szerinti, legalább egy fehérje expresszálasára van genetikailag módosítva.

8. A 7. igénypont szerinti, genetikailag módosított mikroorganizmus, ahol az organizmus genetikailag módosított legalább egy további enzim expresszálasára; ahol a további enzim a következők alkotta csoportból kiválasztott kiegészítő enzim: celluláz, xilanáz, lignináz, glükuronidáz, arabinofuranosidáz, arabinogalaktanáz, ferulasav-észteráz, lipáz, pektináz, glükomannáz, amiláz, laminarináz, xiloglukanáz, galaktanáz, glükóamiláz, pektát-liaáz, kitináz, exo- β -D-glükózaminidáz, cellobioz-dehidrogenáz és acetilxilán-észteráz.

9. A 3. igénypont szerinti készítmény, amely tartalmaz továbbá legalább egy további enzimet lignocellulóz-anyag lebontására vagy tartalmazza az enzim olyan fragmentumát, amely megtartja biológiai aktivitását; a) ahol ez a multienzim-készítmény tartalmaz legalább egy cellobiohidrolázt, legalább egy xilanázt, legalább egy endoglukanázt, legalább egy β -glükozidázt, legalább egy β -xilozidázt és legalább egy kiegészítő enzimet; b) ahol a készítményben az enzimek mintegy 50-70%-a cellobiohidroláz; c) ahol a készítményben az

enzimek mintegy 10-30%-a xilanáz; d) ahol a készítményben az enzimek mintegy 5-15%-a endoglukanáz; e) ahol a készítményben az enzimek mintegy 1-5%-a β -glükozidáz; f) ahol a készítményben az enzimek mintegy 1-3%-a β -xilozidáz; vagy g) ahol a készítmény tartalmaz mintegy 60% cellobiohidrolázi, mintegy 20% xilanázi, mintegy 10% endoglukanázi, mintegy 3% β -glükozidázi, mintegy 2% β -xilozidázi és mintegy 5% kiegészítő enzimet.

10. A 9. igénypont szerinti multienzim-készítmény, ahol a készítmény tartalmaz:

- a) legalább egy első fehérjét, amely aminosav-szekvenciát tartalmaz, amely legalább 95%-ban azonos egy, a következők alkotta csoportból választott aminosav-szekvenciával: SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8 és SEQ ID NO:11, és cellobiohidroláz aktivitása van; vagy fragmentumát, amelynek cellobiohidroláz aktivitása van;
- b) legalább egy második fehérjét, amely aminosav-szekvenciát tartalmaz, amely legalább 95%-ban azonos egy, a következők alkotta csoportból választott aminosav-szekvenciával: SEQ ID NO:14, SEQ ID NO:94, SEQ ID NO:17, SEQ ID NO:20 és SEQ ID NO:23, és endoglukanáz aktivitása van, vagy fragmentumát, amelynek endoglukanáz aktivitása van;
- c) legalább egy harmadik fehérjét, amely aminosav-szekvenciát tartalmaz, amely legalább 95%-ban azonos egy, a következők alkotta csoportból választott aminosav-szekvenciával: SEQ ID NO:26, SEQ ID NO:29, SEQ ID NO:32, SEQ ID NO:35, SEQ ID NO:38 és SEQ ID NO:41, és xilanáz aktivitása van, vagy fragmentumát, amelynek xilanáz aktivitása van;
- d) legalább egy negyedik fehérjét, amely aminosav-szekvenciát tartalmaz, amely legalább 95%-ban azonos a következő aminosav-szekvenciával: SEQ ID NO:44 és β -glükozidáz aktivitása van, vagy fragmentumát, amelynek β -glükozidáz aktivitása van; és
- e) legalább egy ötödik fehérjét, amely aminosav-szekvenciát tartalmaz, amely legalább 95%-ban azonos egy, a következők alkotta csoportból választott aminosav-szekvenciával: SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:62, SEQ ID NO:65, SEQ ID NO:68, SEQ ID NO:71, SEQ ID NO:74, SEQ ID NO:77, SEQ ID NO:80, SEQ ID NO:83, SEQ ID NO:86, SEQ ID NO:89 és SEQ ID NO:92, és hemicelluláz vagy kitináz aktivitása van, vagy fragmentumát, amelynek hemicelluláz vagy kitináz aktivitása van.

11. A 9-10. igénypontok bármelyike szerinti multienzim-készítmény, amely tartalmaz továbbá egy vagy több kiegészítő enzimet, ahol a kiegészítő enzim(ek) magában(magukban) foglal(nak) legalább egy enzimet, amely a következők alkotta csoportból választott: celluláz, xilanáz, lignináz, glükuronidáz, arabinofuranozidáz, arabinogalaktanáz, ferulasav-észteráz, lipáz, pektináz, glükomannáz, amiláz, laminarináz, xiloglukanáz, galaktanáz, glükooamiláz, pektát-láz, kitináz, exo- β -D-glükózaminidáz, cellobioz-dehidrogenáz és acetilxilán-észteráz.

12. Eljárás lignocellulóz-anyag lebontására fermentálható cukrokká, amely tartalmazza a lignocellulóz-anyag érintkeztetését legalább egy, a 2. igénypont (a) pontja szerinti izolált fehérjével vagy legalább egy, a 9-11. igénypontok bármelyike szerinti multienzim-készítménnyel.

13. Eljárás szerves anyag előállítására, amely eljárás tartalmazza a következőket:

- a) lignocellulóz-anyag cukrosítása a 9-11. igénypontok bármelyike szerinti multienzim-készítménnyel;
- b) a cukrosított lignocellulóz-anyag fermentálása egy vagy több fermentáló mikroorganizmussal elvégezve; és

c) a szerves anyag kinyerése a fermentálásból.

14. Eljárás szeszipari száraz alapanyagból vagy oldható anyagokat is tartalmazó szeszipari száraz alapanyagból álló lignocellulóz-anyag lebontására cukrokká, amely tartalmazza a szeszipari száraz alapanyag vagy az oldható anyagokat is tartalmazó szeszipari száraz alapanyag érintkeztetését multienzim-készítménnyel, miáltal a fermentálható cukrok legalább 10%-a felszabadul, és ahol a multienzim-készítmény a 9-11. igénypontok bármelyike szerinti multienzim-készítmény.

15. A 12-14. igénypontok bármelyike szerinti eljárás, amely tartalmazza továbbá az érintkeztetett lignocellulóz anyag visszanyerését, a fermentálható cukrok lebontása után.

16. Eljárás papír vagy pép szintelenítésére vagy biológiai fehérítésére, amely tartalmazza a papír vagy pép érintkeztetését a 2. igénypont (a) pontja szerinti, legalább egy izolált fehérjével vagy a 9-11. igénypontok bármelyike szerinti legalább egy multienzim-készítménnyel.

17. Eljárás szövet kezelésére, ahol a kezelés a következők alkotta csoportból választott: szövet kömosása; szövet lágyságának vagy tapintásának fokozása vagy szövet depilálása; szövet színének helyreállítása vagy világosítása; szövet biológiai fényesítése, defibrillálása, fehérítése, festése vagy irtelenítése; ahol az eljárás tartalmazza a szövet érintkeztetését a 2. igénypont (a) pontja szerinti, legalább egy izolált fehérjével vagy a 9-11. igénypontok bármelyike szerinti legalább egy multienzim-készítménnyel.

18. Detergenskészítmény, amely legalább egy, a 2. igénypont (a) pontja szerinti, izolált fehérjét vagy legalább egy, a 9-11. igénypontok bármelyike szerinti multienzim-készítményt tartalmaz, és adott esetben felületaktív anyagot tartalmaz.