

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199527072 B2
(10) Patent No. 703359

(54) Title
Ribozymes with optimized hybridizing arms, stems, and loops, tRNA embedded
ribozymes and compositions thereof

(51)⁶ International Patent Classification(s)
C07H 021/02 C12N 015/64

(21) Application No: 199527072 (22) Application Date: 1995 .06 .21

(87) WIPO No: W096/00232

(30) Priority Data

(31) Number (32) Date (33) Country
08/265484 1994 .06 .24 US

(43) Publication Date : 1996 .01 .19
(43) Publication Journal Date : 1996 .02 .29
(44) Accepted Journal Date : 1999 .03 .25

(71) Applicant(s)
Gene Shears Pty. Ltd.

(72) Inventor(s)
Paul Keese; Marianne Stapper; Rhonda Perriman

(74) Agent/Attorney
DAVIES COLLISON CAVE,1 Little Collins Street,MELBOURNE VIC 3000

(56) Related Art
AU 18249/92
AU 79976/91
WO 95/06764

OPI DATE 19/01/96 APPLN. ID 27072/95
AOJP DATE 29/02/96 PCT NUMBER PCT/AU95/00359



AU9527072

IN

(51) International Patent Classification ⁶ : C07H 21/02, C12N 15/64		A1	(11) International Publication Number: WO 96/00232
			(43) International Publication Date: 4 January 1996 (04.01.96)
(21) International Application Number: PCT/AU95/00359		(81) Designated States: AU, CA, JP, RU, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 21 June 1995 (21.06.95)		Published With international search report.	
(30) Priority Data: 08/265,484 24 June 1994 (24.06.94) US			
(60) Parent Application or Grant (63) Related by Continuation US 08/265,484 (CIP) Filed on 24 June 1994 (24.06.94)			
(71) Applicant (for all designated States except US): GENE SHEARS PTY. LTD. [AU/AU]; Level 3, 154-156 Military Road, Neutral Bay, NSW 2089 (AU). Cnr Barry Drive & Clunies Ross Street, Canberra ACT 2601 (AU)			
(72) Inventors; and (75) Inventors/Applicants (for US only): KEESE, Paul [AU/AU]; 10 Morehead Street, Curtin, ACT 2605 (AU). STAPPER, Marianne [NL/AU]; 29 Kauper Street, Scullin, ACT 2614 (AU). PERRIMAN, Rhonda [AU/AU]; 6 Belah Street, O'Connor, ACT 2601 (AU).			
(74) Agents: SLATTERY, John, Michael et al.; Davies Collison Cave, 1 Little Collins Street, Melbourne, VIC 3000 (AU).			
(54) Title: RIBOZYMES WITH OPTIMIZED HYBRIDIZING ARMS, STEMS, AND LOOPS, tRNA EMBEDDED RIBOZYMES AND COMPOSITIONS THEREOF			
(57) Abstract This invention is directed to improved catalytic compounds, hammerhead ribozymes, capable of hybridizing with a target RNA to be cleaved. These improved compounds have optimized stems (X)m * (X)m', loops (X)b and hybridizing arms. The invention is also directed to compositions for enhanced RNA cleavage which comprise a first synthetic non-naturally occurring oligonucleotide compound which comprises nucleotides whose sequence defines a conserved catalytic region and nucleotides whose sequence is capable of hybridizing with a predetermined target sequence and a second synthetic non-naturally occurring oligonucleotide which does not contain the predetermined target sequence and is complementary to at least a portion of the first oligonucleotide compound. The invention is also directed to synthetic non-naturally occurring oligonucleotide compounds embedded in a tRNA. The ribozymes and compositions of the present invention may be used <i>in vitro</i> or <i>in vivo</i>. They may be used as diagnostic or therapeutic agents.			

- 1 -

**RIBOZYMES WITH OPTIMIZED HYBRIDIZING ARMS, STEMS,
AND LOOPS, tRNA EMBEDDED RIBOZYMES AND COMPOSITIONS
THEREOF.**

Throughout this application various references are cited in bracket by author and publication year. The full citations are listed alphabetically and may be found immediately preceding the claims. These publications are hereby incorporated by reference into the present application.

Background of the Invention

The discovery that RNA molecules can function as enzymes revolutionized the understanding of chemistry in biological systems. It has now been demonstrated that RNA molecules can catalyze chemical transformations on themselves as well as on other RNA molecules [Castanotto, 1992]. The term ribozyme has been given to these RNA molecules. It has now become apparent that ribozymes play an important role in the biochemical function of many organisms.

Several types of ribozymes have been identified in living organisms. The first ribozyme to show catalytic turnover was RNA of ribonucleases P. Ribonucleases P (RNase P) cleaves precursor tRNAs (pre-tRNAs) at their 5' ends to give the mature 5'-termini of tRNAs. In *Escherichia coli* and *Bacillus subtilis*, the RNase P holoenzyme is composed of one basic protein subunit of approximate M_r 14,000 (119 amino acids) and one single stranded RNA molecule of 377 and 401 nucleotides, respectively [Baer, 1990; Altman 1987; Waugh, 1989; Pace, 1990; Nichols, 1988]. The second ribozyme to show turnover was the L-19 intervening sequence (IVS) from *tetrahymena*. The 413 nucleotide intervening sequence (IVS) in the nuclear rRNA precursor from *Tetrahymena thermophila* can be excised and the two exons ligated in



the complete absence of any protein [Kruger, 1982; Cech, 1981]. Unique to the *Tetrahymena thermophila* self-splicing reaction is the requirement of a guanosine or 5' guanosine nucleotide cofactor. The hammerhead self-cleavage reactions constitutes a third class of ribozymes. A number of plant pathogenic RNAs [Symons, 1989; Symons, 1990; Bruening, 1989; Bruening 1990], one animal viral RNA [Taylor, 1990] and a transcript from satellite II of DNA of the newt [Epstein, 1987; Epstein 1989] and from a *Neurospora* DNA plasmid [Saville, 1990] undergo a site specific self-cleavage reaction *in vitro* to produce cleavage fragments with a 2',3'-cyclic phosphate and a 5'-hydroxyl group. This self-cleavage reaction is non-hydrolytic, unlike RNases P RNA cleavage of pre-tRNAs, where the internucleotide bond undergoes a phosphoryl transfer reaction in the presence of Mg^{++} or other divalent cations. Metal cations may be essential to RNA catalysis [Pyle, 1993]. Other reactions documented to date show that ribozymes can catalyze the cleavage of DNA [Robertson, 1990; Herschlag 1990], the replication of RNA strands [Green, 1992], the opening of 2'-3'-cyclic phosphate rings [Pan, 1992], as well as react with phosphate monoesters [Zaug, 1986] and carbon centers [Noller, 1992; Piccirilli, 1992]. Finally, ribozymes with new kinds of catalytic reactivity are being created through techniques of *in vitro* selection and evolution [Joyce, 1992].

The ability to specifically target and cleave a designated RNA sequence has led to much interest in the potential application of hammerhead ribozymes as gene therapy agents or drugs. One component to the success of treating a disease or targeting a specific RNA or DNA strand (substrate) to be cleaved is the optimization of the ribozyme/substrate complex. Optimization may be performed *in vitro* whereby a ribozyme is modified until it achieves maximum chemical activity on a target RNA or DNA molecule. While much success has been achieved *in vitro* in targeting and cleaving a number of designated RNA sequences (Saxena and Ackerman, 1990; Lamb and Hayes, 1991; Evans, et al., 1992; Mazzolini, et al., 1992; Homann, et al., 1993), the translation of this success into ribozyme action in the whole cell has been limited. This is particularly the case for plant systems, with only one example of ribozyme

induced cleavage presently reported (Steinecke, et al., 1992). Thus, it is not entirely clear how the success in vitro would correlate to the success in vivo.

5 Previous reports have demonstrated that high levels of ribozyme expression are required to achieve reduced accumulation of target sequence in vivo [Cameron and Jennings, 1989; Cotten and Birnsteil, 1989; Sioud and Drilca, 1991; L'Huillier, et al., 1992; Perriman et al., 1993]. Additionally, a recent article suggests a necessity for the target and ribozyme to be sequestered in the same cellular compartment [Sullenger and Cech, 1993]. The results of several reports suggest that while the ribozyme
10 molecule is clearly capable of inducing specific cleavage of a designated target RNA within a biological system, the rate limiting step, under in vivo conditions, is the formation of the active substrate/ribozyme hybrid. Such approaches towards optimizing the formation of the active substrate/ribozyme hybrid include attempts at "stabilizing" the ribozyme transcript by embedding the ribozyme within a stable RNA
15 (Cameron and Jennings, 1989; Sioud, et al., 1992; Cotten and Birnsteil, 1989). By providing the ribozyme transcript with a longer half-life, the chances of forming the desired hybrid with the target RNA are optimized.

20 In the approach of Cotten and Birnsteil (1989) a tRNA motif was used to deliver a hammerhead ribozyme to *Xenopus* oocytes and reduce the accumulation of a cytoplasmic target RNA. The embedded ribozyme was more effective than the analogous non-embedded ribozyme.

Summary of the Invention

This invention is directed to improved catalytic compounds, hammerhead ribozymes, capable of hybridizing with a target RNA to be cleaved. These improved compounds have optimized stems (X)_m * (X)_m', loops (X)_b and hybridizing arms. The invention
5 is also directed to compositions for enhanced RNA cleavage which comprise a first synthetic non-naturally occurring oligonucleotide compound which comprises nucleotides whose sequence defines a conserved catalytic region and nucleotides whose sequence is capable of hybridizing with a predetermined target sequence and a second
10 synthetic non-naturally occurring oligonucleotide which does not contain the predetermined target sequence and is complementary to at least a portion of the first oligonucleotide compound. The invention is also directed to synthetic non-naturally occurring oligonucleotide compounds embedded in a tRNA.

The ribozymes and compositions of the present invention may be used in vitro or in
15 vivo. They may be used as diagnostic or therapeutic agents.

Brief Description of the Figures

Figure 1. Shows the cleavage efficiency of helix II ((X)_m and (X)_m of formulas 1 and 2) stem-length variants by plotting % cleavage as a function time in minutes. The data plotted is tabulated in Table 1 and each entry in Table 1 corresponds to each graph in the plot by number.

Figure 2. Shows the cleavage efficiency of helix II ((X)_m and (X)_m of formulas 1 and 2) conserved G-C base pair by plotting % cleavage as a function time in minutes. The data plotted is tabulated in Table 2 and each entry in Table 2 corresponds to each graph in the plot by number.

Figures 3A-3C. Shows the cleavage efficiency of helix II ((X)_m and (X)_m of formulas 1 and 2) loop variants by plotting % cleavage as a function time in minutes. The data plotted is tabulated in Table 3 and each entry in Table 3 corresponds to each graph in the plot by number. Figure 3A shows the control and loop variants 1-6, Figure 3B shows the control and loop variants 7-11, and Figure 3C shows the control and loop variants 14, 15 and 21-22.

Structure of the ribozyme/substrate complex represented as a wishbone, as determined by X-ray crystallography (Pley et al. 1994 A) and numbered according to Hertel et al. (1992). Nucleotides in bold correspond to the highly conserved catalytic core of naturally occurring hammerhead self-cleaving domains (Bruening 1989). Dashed lines depict non-canonical base pairs. (Fig. 4B) Time course of the parent ribozyme cleavage reaction, with error bars that correspond to standard deviation from ten experiments.

Figure 5. Relative rate constants, k^{rel} of ribozymes with mutations in stem II. Rz 1 is the parent sequence, Rz 2-18 have sequence variations in the loop only. Free energy calculations according to Frier et al. (1986; kcal/mol at 37°C in 1 M NaCl) were only for Watson Crick base paired sequences. Dashed lines depict non-canonical base-pairs.

Figure 6. Shows the cleavage efficiency of helix I ((X)_a of formulas 1 and 2; or -N-N-(X)_a of formula 3) sequence variants by plotting % cleavage as a function time in minutes. The data plotted is tabulated in Table 6 and each entry in Table 6 corresponds to each graph in the plot by number.

Figure 7. Shows the cleavage efficiency of helix III ((X)_a



of formulas 1 and 2) sequence variants by plotting % cleavage as a function time in minutes. The data plotted is tabulated in Table 7 and each entry in Table 7 corresponds to each graph in the plot by number.

5 **Figure 8.** Shows the effect of oligonucleotides, complementary to the ribozyme, on cleavage efficiency by plotting cleavage as % of control verses a 10 minute interval. The data plotted is tabulated in Table 8 and each entry in Table 8 corresponds to each graph in the plot by number.

10 **Figures 9A-9D.** Construction of plasmids for in vitro and in vivo analysis of non-embedded and tRNA-embedded ribozyme and antisense; and CAT and CM2, mutant CAT targets; (9a). 266bp tyrosine tRNA sequence subcloned as SspI AccI fragment into pGEM3zf- (pGEMtRNA). Hatched box represents site of insertion of 10 bp Bam HI.SmaI.EcoRI polylinker at position 112 (pGEMtRNApoly). This is at the 5' end of the anticodon loop. Clover-leaf motif shows site of insertion of polylinker as well as 13 base loop out which is the endogenous intron. (9b). Sequence of hammerhead ribozyme Rz12 which is designed to anneal and cleave at position 464 on the CAT gene. (9c). 815 bp CAT sequence. Arrowed site is position 464, a GUC triplet targeted by Rz12. (9d). pACMV vector used for expression of ribozyme/antisense and target sequences in vivo. The RNA polymerase II coat protein promoter is indicated 5' of the polylinker region with termination and polyadenylation signals indicated 3'.
20 All ribozyme/antisense and target sequences are subcloned within this region. The 670 bp duplication contains viral origin of replication. The pUC19-derived bacterial plasmid inserted at the ClaI site contains bacterial replication origin ColEI and chloramphenicol resistance gene.

25 **Figures 10A-10B.** In vitro cleavage by linear and tRNA embedded ribozymes. (10a). Autoradiograph of in vitro cleavage reaction at 30°C for 1 hour. The CAT substrate but not the ribozymes were radioactively labelled in these reactions. The lanes labelled 1 to 3 represent: 1, CAT RNA incubated alone; 2, CAT RNA + Rz12; 3, CAT RNA + tRNARz12. The RNA species are as follows: Sub, CAT substrate RNA; 5'P, cleavage product 5' of target site; 3'P, cleavage product 3' of cleavage site. (10b) Bar graph of in vitro cleavage reactions showing average % cleavage of
30

CAT transcript in three independent cleavage reactions. Y axis shows % cleavage and error bars represent 2 standard deviations.

Figures 11A-11B. Relative expression of the wildtype (CAT) (11A) and mutant (CATMUT2) (11B) genes in the presence of pACMVtRNA (control), pACMVAs24 or pACMVtRNAAs24 (antisense), pACMVRz12 or pACMVtRNARz12 (ribozyme) in *Nicotiana tabacum* protoplasts. The figures show the results from seven separate protoplast isolations. Each bar represents the mean relative value of CAT or mutant CAT expression, upon addition of the antisense or ribozyme in the tRNA-embedded form. Control CAT or CATMUT2 activity has been normalized as 100%. Each protoplast isolation and subsequent electroporation involved at least triplicate samples of each of the construct combinations: i.e.; CAT (or CATMUT2) + control, CAT (or CATMUT2) + antisense (tRNA embedded or non-embedded) and CAT (or CATMUT2) + ribozyme (tRNA embedded or non-embedded).

Figures 12A-12B. Autoradiograph of thin layer chromatography plate showing relative CAT (12A) or CATMUT2 (12B) activity in the presence of pACMVtRNA or pACMVtRNARz12. For CAT + tRNA or CM2 + tRNA, three independent electroporations are shown (labelled 2-4). For CAT + tRNARz12 or CM2 + tRNARz12, four independent electroporations are shown (labelled 5-8). Control lanes (1) are mock inoculated protoplasts.


Figures 13A-13B. Analysis of CAT mRNA level in vivo. Southern blot of RT-PCR products from total RNA isolated from electroporated *Nicotiana tabacum* protoplasts. The two blots (Figures 11a and 11b) show accumulated products of full length CAT mRNA and CAT mRNA 3' of the GUC cleavage site. The lanes for the CAT and CM2 full length CAT mRNA, 4 + tRNARz12 -3' CAT mRNA, 5, + tRNAAs24 - full length CAT mRNA, 6 + tRNAAs24 -3' CAT mRNA, 7 + Rz12 - full length CAT mRNA, 8 - Rz12 -3' CAT mRNA, 9 + As24 - full length CAT mRNA, 10 + As24 -3' CAT mRNA. Blots were probed with ssDNA representing full length CAT sequence. Also, see Table 10: Ratio of accumulation of 3' product to full length CAT product. Table 10 shows the ratio of these products for each of the construct combinations. These ratios are the accumulated data from two independent experiments.




Figures 14A-14B. Lane 1: 20ng pACMVCAT or pACMVCM2 MboI restricted. Lane 2: 20ng pACMVCAT or pACMVCM2 DpnI restricted. Lane 3: 1ug transfected DNA (time zero), pACMVCAT/CM2 input only, DpnI restricted. Lane 4: 1ug transfected DNA (time zero), pACMVCAT/CM2 input only, MboI restricted. Lane 5: 1ug transfected DNA (time: 3 days), pACMVCAT/CM2 input only, DpnI restricted. Lane 6: 1ug transfected DNA (time: 3 days), pACMVCAT/CM2 input only, MboI restricted. Lane 7: 1ug transfected DNA (time zero), pACMVCAT/CM2 + pACMVtRNARz12, DpnI restricted. Lane 8: 1ug transfected DNA (time zero), pACMVCAT/CM2 + pACMVtRNARz12, MboI restricted. Lane 9: 1ug transfected DNA (time: 3 days), pACMVCAT/CM2 + pACMVtRNARz12, DpnI restricted. Lane 10: 1ug transfected DNA (time zero), pACMVCAT/CM2 + pACMVtRNARz12, MboI restricted. (14a) CAT: Southern blot of DNA from 3 day old or time zero cells with either DpnI or mboI. Any replicated DNA will be cut with mboI. Band 2 represents CAT/CM2 insert and is maintained in mboI digests signifying replicated DNA. Band 2 is present in equal amounts in Lanes 6 and 10 representing pACMVCAT/CM2 input only and pACMVCAT/CM2 + pACMVtRNARz12 respectively. Band 1 is on input DNA only and is lost in mboI digests. (14b) CATMUT2: Southern blot of DNA from 3 day old or time zero cells with either DpnI or mboI. Any replicated DNA will be cut with mboI. Band 2 represents CAT/CM2 insert and is maintained in mboI digests signifying replicated DNA. Band 2 is present in equal amounts in Lanes 6 and 10 representing pACMVCAT/CM2 input only and pACMVCAT/CM2 + pACMVtRNARz12 respectively. Band 1 is on input DNA only and is lost in mboI digests. This blot doesn't show that lane 1 DNA is completely resistant to mboI digestion.

Detailed Description of the Invention


In several embodiments there is provided compounds that are capable of cleaving a RNA target molecule. These compounds are represented below as formulas 1, 2, 3 and 4. Generally these compounds are known as ribozymes. More specifically, compounds of formula 1, 2, and 3 are known as ribozymes; and compounds of formula 4 are known as tRNA embedded ribozymes.



Formula 1



Formula 2



Formula 3



Formula 4

In the compounds of formulas 1, 2, and 3: The symbol X represents a nucleotide which may be the same or different and may be substituted or modified in its sugar, base or phosphate. The solid lines drawn in formulas 1, 2 and 3 represent a chemical linkage providing covalent bonds between the nucleotides located on either side thereof. The integer a defines the number of nucleotides in $(X)_a$ such that a may be 0 or 1 and if 0, the A located 5' of $(X)_a$ is bonded to the X located 3' of $(X)_a$.

In the above compounds of formulas 1 and 2: The symbols $(X)_n$ and $(X)_{n'}$ represent an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved. The oligonucleotides, $(X)_n$ and $(X)_{n'}$, do not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, and such RNA target sequences are not present within formula 1 or formula 2. The symbols n and n' represent integers which define the number of nucleotides in the oligonucleotides $(X)_n$ and $(X)_{n'}$ respectively, and the sum of n + n' is greater than 14.

In the compounds of formulas 1 and 3: The oligonucleotide $(X)_b$ may be present or absent, and b represents an integer which is greater than or equal to four if $(X)_b$ is present. In the compounds of formulas 2 and 3: The symbol * represents base pairing between the nucleotides located on either side thereof. In the above compounds of formula 1: The oligonucleotides, $(X)_m$ and $(X)_{m'}$, represent an oligonucleotide having the sequences (a) 3'-CUG-5' and 5'-GAC-3', (b) 3'-GCG-5' and 5'-AGC-3', (c) 3'-CGG-5' and 5'-GCC-3' or (d) 3'-CGC-5' and 5'-GCG-3', respectively and each * represents base pairing between the nucleotides located on either side thereof.

In the above compounds of formula 2: The integers m and m' range from two to eight inclusive. The oligonucleotide $(X)_b$ has one of the following sequences: 3'-AGGUU-5', 3'-AGGG-5', 3'-AAGGC-5' or 3'-GGUG-5'.

In the above compounds of formula 3: The symbol $(X)_n$ represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target

sequence to be cleaved. The oligonucleotides, $(X)_n$ and $(X)_{n'}$, do not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, and such RNA target sequences are not present within formula 3. The segment -N-N- are nucleotides that have a predetermined sequence to the RNA target sequence. The symbols n and n' represent integers which define the number of nucleotides in the oligonucleotides $(X)_n$ and $(X)_{n'}$ respectively, and the sum of $n + n'$ is greater than 12. The integers m and m' are greater than or equal to 4.

In the above compounds of formula 4: The symbol X represents a nucleotide which may be the same or different. The oligonucleotides, $(X)_t$ and $(X)_{t'}$, have a sequence from a portion of a tRNA molecule. The integers, t and t', define the number of nucleotides in the oligonucleotide with the proviso that the sum of $t + t'$ is greater than 50. The symbol Y represents a synthetic non-naturally occurring oligonucleotide compound which compound comprises nucleotides whose sequence defines a conserved catalytic region and nucleotides whose sequence is capable of hybridizing with a predetermined target sequence to be cleaved.

In another embodiment, the oligonucleotides, $(X)_t$ and $(X)_{t'}$ of formula 4, are derived from a tyrosine tRNA molecule of *Nicotiana rustica*.

In several embodiments, each of m and m' of formulas 2 or 3 are four.

In several embodiments, the nucleotide $(X)_b$ of formula 1 or 3 is present. In several embodiments, if $(X)_b$ of formula 1 or 3 is present, then the integer b of $(X)_b$ is equal to four. In several embodiments, the oligonucleotide $(X)_b$ of formula 1 or 3 has the sequence 5'-G-U-G-A-3'. In several embodiments, the oligonucleotide 3'-($X)_n$ - of formula 1, 2 or 3 is 3'-($X)_{n-1}$ -A-. In several embodiments, the oligonucleotide 3'-($X)_n$ - of formula 1, 2 or 3 is 3'-($X)_{n-2}$ -C-A-. In several embodiments, the nucleotide $(X)_a$ of formula 1, 2 or 3 is absent. The oligonucleotides $(X)_n$, $(X)_{n'}$, N-N-($X)_n$ (of formula 1, 2, 3, or 4) represents the arms or flanking sequences of a ribozyme which hybridize to respective portions of a target RNA sequence. The arms may hybridize

along the full length of the target RNA or part thereof. The sequence 3'-...AAG-(X)_m-(X)_n-(X)_m-(X)_n-AGXAGUC-...-5' of formulas 1-3 represents the catalytic region. The catalytic region may also form part of the hybridizing region. The catalytic region may contain one or more additional nucleotides which do not adversely effect catalytic activity. Such additions could be readily tested for ribozyme activity without undue experimentation.

In another embodiment, there is provided a composition which comprises a compound of any of formulas 1, 2, or 3 in association with an acceptable carrier. In several embodiments, there is provided a oligonucleotide transfer vector containing a nucleotide sequence which on transcription gives rise to a compound of formula 1, 2, 3, or 4.

In several embodiments, there is provided a transfer vector, as mentioned hereinabove, that may be a bacterial plasmid, a bacteriophage DNA, a cosmid, or an eukaryotic viral DNA. In one embodiment, there is provided a transfer vector that may be a bacterial plasmid, a bacteriophage DNA, a cosmid, or plant viral DNA. In one embodiment, there is provided a transfer vector, as mentioned hereinabove, that contains the promoter sequences for RNA polymerase II or RNA polymerase III. In several embodiments, there is provided a host cell transformed by the transfer vector as mentioned hereinabove. In several embodiments, there is provided a host cell, as mentioned hereinabove, which may be a prokaryotic host cell or an eukaryotic host cell. In several embodiments, the prokaryotic host cell, as mentioned hereinabove, is an E. coli host cell. In several embodiments, the eukaryotic host cell, as mentioned hereinabove, is a monkey COS host cell, a Chinese hamster ovary host cell, a mammalian host cell or a plant host cell. In one embodiment, the eukaryotic host cell, as mentioned hereinabove, is a monkey COS host cell, a Chinese hamster ovary host cell, a mammalian host cell, a plant host cell or a plant protoplast host cell.


In one embodiment, there is provided a packaged oligonucleotide transfer vector, as mentioned hereinabove, that contains a nucleotide sequence which on transcription

gives rise to the compound of formula 4. In one embodiment, there is provided a packaged oligonucleotide transfer vector, as mentioned hereinabove, which is a plant DNA virus, a geminivirus or an infective phage particle. In one embodiment, there is provided a packaged oligonucleotide transfer vector, as mentioned hereinabove, which contains the promoter sequences for RNA polymerase II or RNA polymerase III.

In accordance with another aspect of this invention, there is provided a composition for enhanced RNA cleavage comprising a first synthetic non-naturally occurring oligonucleotide compound which compound comprises nucleotides whose sequence defines a conserved catalytic region and nucleotides whose sequence is capable of hybridizing with a predetermined target sequence to be cleaved and a second synthetic non-naturally occurring oligonucleotide which does not contain the predetermined target sequence and is complementary to at least a portion of the compound. In another embodiment, the composition, as discussed hereinabove, is in association with an acceptable carrier. Additionally, the second synthetic non-naturally occurring oligonucleotide, as discussed hereinabove, may include, but are not limited to, one of the sequences listed in Table 8. Furthermore, the second synthetic non-naturally occurring oligonucleotide may be replaced with a solvent which may also enhance ribozymes (of the present invention) cleavage rates and may include, but are not limited to, one of the solvents listed in Table 8.

An aspect of this invention provides a composition as discussed hereinabove wherein the oligonucleotide is an oligoribonucleotide or an oligodeoxyribonucleotide which may be substituted or modified in its sugars, bases or phosphodiester linkages. Accordingly, oligonucleotide derivatives or modifications may be made at the level of the sugar, base, monophosphate groupings or phosphodiester linkages. It is preferred that the oligonucleotide be an oligoribonucleotide, however, other substitutions or modifications in the nucleotide are possible providing that endonuclease activity is not lost. Such derivatives or modifications are described hereinbelow.

In several embodiments, the synthetic non-naturally occurring oligonucleotide compound of (a) the composition and (b) of formula 4, discussed hereinabove, has the structure:



15

20

25


30

wherein each X represents a nucleotide which is the same or different and may be modified or substituted in its sugar, phosphate or base; wherein each of A, C, U, and G represents a ribonucleotide which may be unmodified or modified or substituted in its sugar, phosphate or base; wherein 3'—AAG....AGUCX — 5' defines a conserved catalytic region; wherein each of (X)_nA and (X)_n represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved, and does not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, such RNA target sequence not being present within the compound; wherein each * represents base pairing between the nucleotides located on either side thereof; wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof; wherein a represents an integer which defines a number of nucleotides with the proviso that a may be 0 or 1 and if 0, the A located 5' of (X)_a is bonded to the X located 3' of (X)_a; wherein each of m and m' represents an integer which is greater than or equal to 1; and wherein (X)_b represents an oligonucleotide and b represents an integer which is greater than or equal to four.

35


In several embodiments, the synthetic non-naturally occurring oligonucleotide compound of (a) the composition and (b) of formula 4, discussed hereinabove, has the structure:

-15-



wherein each X is the same or different and represents a ribonucleotide or a deoxyribonucleotide which may be modified or substituted in its sugar, phosphate or base; wherein each of A, C, U, and G represents a ribonucleotide which may be unmodified or modified or substituted in its sugar, phosphate or base; wherein 3'—AAG....AGUCX—5' defines a conserved catalytic region; wherein each of $(X)_nA$ and $(X)_{n'}$ represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved, and does not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, such RNA target sequence not being present within the compound; wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof; wherein m represents an integer from 2 to 20; and wherein none of the nucleotides $(X)_m$ are Watson-Crick base paired to any other nucleotide within the compound.

In several embodiments, the synthetic non-naturally occurring oligonucleotide compound of (a) the composition and (b) of formula 4, discussed hereinabove, has the structure:



wherein each X is the same or different and represents a ribonucleotide or a deoxyribonucleotide which may be modified or substituted in its sugar, phosphate or base; wherein each of A, C, U, and G represents a ribonucleotide which may be unmodified or modified or substituted in its sugar, phosphate or base; wherein 3'(X)_{F4}....(X)_{P1}— 5' defines a conserved catalytic region; wherein each of (X)_{F4} and (X)_{F3} represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved; wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof; wherein F3 represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that F3 is greater than or equal to 3; wherein F4 represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that F4 is from 3 to 5; wherein each of (X)_{P1} and (X)_{P4} represents an oligonucleotide having a predetermined sequence such that (X)_{P4} base-pairs with 3-6 bases of (X)_{P1}; wherein P1 represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that P1 is from 3 to 6 and the sum of P1 and F4 equals 9; wherein each of (X)_{P2} and (X)_{P3} represents an oligonucleotide having a predetermined sequence such that (X)_{P2} base-pairs with at least 3 bases of (X)_{P3}; wherein each * represents base pairing between the nucleotides located on either side thereof; wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof; wherein each of the dashed lines independently represents either a chemical linkage providing covalent bonds between the nucleotides located on either side thereof or the absence of any such chemical linkage; and wherein (X)_{L2} represents an oligonucleotide which may be present or absent with the proviso that L2 represents an integer which is greater than or equal to 3 if (X)_{L2} is present.

In formulas 1, 2, 3, 1a, and 1b, the integers n and n' that define the number of nucleotides in $(X)_n$ and $(X)_{n'}$, respectively, may range from five to two hundred, and preferably may range from six to twenty. In formulas 1, 2, 3, and 1a, the integers m and m' that define the number of nucleotides in $(X)_m$ and $(X)_{m'}$, respectively, may range from four to twenty, and preferably may range from four to five. In formulas 1, 2, 3, and 1a, the integer b that defines the number of nucleotides in $(X)_b$ may range from one to eight, but preferably may range from one to four. In formula 1b, the integer m that define the number of nucleotides in $(X)_m$ may range from two to thirty, and preferably may range from four to ten. In formula 1c, the integer $P1$ that defines the number of nucleotides in $(X)_{P1}$ may range from three to ten, but preferably may range from three to six. In formula 1c, the integers $P2$ and $P3$ that define the number of nucleotides in $(X)_{P2}$ and $(X)_{P3}$, respectively, may range from three to twenty, and preferably may range from three to ten. In formula 1c, the integer $P4$ that defines the number of nucleotides in $(X)_{P4}$ may range from three to ten, but preferably may range from three to six. In formula 1c, the integer $L2$ that defines the number of nucleotides in $(X)_{L2}$ may range, if $(X)_{L2}$ present, from three to twenty, but preferably may range from three to ten. In formula 1c, the integer $F3$ that defines the number of nucleotides in $(X)_{F3}$ may range from three to twenty, but preferably may range from three to ten. In formula 1c, the integer $F4$ that defines the number of nucleotides in $(X)_{F4}$ may range from three to twenty, but preferably may range from three to five. Additionally, each of $(X)_m$ and $(X)_{m'}$ of formulas 1, 2, 3, or 1a may be, but is not limited to, an oligonucleotide having one of the sequences listed in Table 1, Table 2, and Table 4. Additionally, the oligonucleotide $(X)_b$ of formulas 1, 2, 3, or 1a may include, but are not limited to, one of the sequences listed in Table 3. Additionally, each of $(X)_n$ and $(X)_{n'}$ of formulas 1, 2, 3, 1a, or 1b may include, but are not limited to, one of the sequences listed in Tables 6 and Table 7 wherein $(X)_n$ and $(X)_{n'}$ in the tables are helix III and helix I respectively.

The sequence $(X)_m-(X)_b-(X)_{m'}$ as represented in formulas 1, 2, 3 or 1a, may be short in length and may be absent of base pairing between $(X)_m$ and $(X)_{m'}$. For convenience, these types of ribozymes are called minizymes. The sequences $(X)_m-(X)_b-(X)_{m'}$ of

formulas 1-4 may include, but are not limited to, one of the sequences listed as minizyme variants in Table 5.

5 Endonuclease activity of the ribozyme (of formula 1, 2, 3, or 4) is readily, simply and routinely tested by incubating the ribozyme with its substrate and thereafter assessing whether cleavage of a the substrate takes place. For example, cleavage of a target mRNA takes place after the trinucleotide sequence XUX' where X and X' represent any ribonucleotide, and which may be the same or different and U represents a ribonucleotide having the base uridine. Preferred cleavage sites, XUX', include
10 GUC, GUU, GUA and UUC. By way of example, suitable reaction conditions may comprise a temperature from about 4 degree(s) C. to about 60 degree(s) C. (preferably about 20 degree(s) to 55 degree(s) C.), pH from about 7.0 to about 9.0 and salt (such as Mg^{2+}) from about 1 to about 100 mM (preferably 1 to 20 mM). Ribozymes containing a small number of nucleotides in each of the groups $(X)_n$ and
15 $(X')_n$ of formulas 1-3 (such as four nucleotides) would generally be incubated at lower temperatures, such as about 20 degree(s) C. to about 25 degree(s) C. to aid duplexing of complementary nucleotide sequences in the ribozyme sequences $(X)_n$ and $(X')_n$ and the substrate. The ribozyme would generally be in an equimolar ratio to the substrate or in excess thereof. However, as the ribozyme may act as an enzyme, cleaving
20 substrate without consumption, the ratio of ribozyme to substrate is not of importance.

A target RNA containing a suitable cleavage site such as GUC site may be incubated with a ribozyme (of formula 1, 2, 3, or 4) or a composition (as described hereinabove) wherein one of the components is a ribozyme (which may be a
25 compound of formula 1, 2, 3, 4, 1a, 1b, or 1c) which, for example, may contain one or more modifications within the catalytic region of the ribozyme. The nucleotide sequences $(X)_n$ and $(X')_n$ of formulas 1-2 are selected so as to be complementary (that is, capable of forming base pairs) to nucleotide sequences flanking the cleavage site in the target RNA. The nucleotide sequences $(X)_n$ of formula 3 are selected so as to
30 be complementary (that is, capable of forming base pairs) to nucleotide sequences flanking the cleavage site in the target RNA, whereas the nucleotide sequences -N-N-

(X)_n of formula 3 are selected so as to be complementary (that is, capable of forming base pairs) to part of nucleotide sequences flanking the cleavage site in the target RNA. On incubation of the ribozyme or ribozyme composition and its substrate, an enzyme/substrate complex is formed as a result of base pairing between complementary nucleotides in the ribozyme and the substrate. Nucleotide sequences (X)_n and (X)_n of the formulas 1-3 and nucleotide sequences flanking the cleavage site in the substrate may form a double stranded duplex as a result of base pairing, which base pairing is well known in the art [See for example: Sambrook, 1989]. The formation of a double stranded duplex between complementary nucleotides may be referred to as hybridization [Sambrook, 1989]. Hybridization or duplex formation between the ribozyme and its substrate can be readily assessed, for example, by labelling one or both components, such as with a radiolabel, and then subjecting the reaction mixture to polyacrylamide gel electrophoresis under non-denaturing conditions [Sambrook, 1989]. If the target is cleaved on incubation with the ribozyme it is active and is within the scope of this invention. Accordingly, a ribozyme containing a nucleotide derivative may be simply tested for endonuclease activity in a routine manner.

As will be readily appreciated by workers in the field to which this invention relates, the cleavage of a target RNA may be readily assessed by various methods well known in the art [See for example: Sambrook, 1989]. Cleavage may, for example, be assessed by running the reaction products (where the substrate is radioactively labelled) on acrylamide, agarose, or other gel systems, and then subjecting the gel to autoradiography or other analytical technique to detect cleavage fragments [Sambrook, 1989].

The nucleotides of the sequences (X)_n and (X)_n of formulas 1-4 may be of any length and sequence sufficient to enable hybridization formation with complementary nucleotides in the target RNA, as described herein. The nucleotides may be in the form of deoxyribonucleotides, ribonucleotides, deoxyribonucleotide ribonucleotide hybrids, or derivatives thereof as herein described. These flanking sequences ((X)_n

and $(X)_n$ of formulas 1-4) may be chosen to optimize stability of the ribozyme from degradation. For example, deoxyribonucleotides are resistant to the action of ribonucleases. Modified bases, sugars or phosphate linkages of nucleotides, such as phosphoramidate, or phosphorothioate linkages in the phosphate backbone of the nucleotide sequences, may also provide resistance to nuclease attack. Binding affinity may also be optimized in particular circumstances, by providing nucleotides solely in the form of ribonucleotides, deoxyribonucleotides, or combinations thereof. In some circumstances it may be necessary to optimize the composition of the sequences $(X)_n$ and $(X)_m$ of formulas 1-4, to maximize target RNA cleavage. The cleavage activity of ribozymes having flanking nucleotide sequences which hybridize to target sequences and which are comprised wholly of deoxyribonucleotides may, in some circumstances, have reduced activity. In such circumstances optimization may involve providing a mixture of deoxyribonucleotides and ribonucleotides in the nucleotide sequences $(X)_n$ and $(X)_m$ of formulas 1-4. For example, nucleotides in the ribozyme which are proximal to the cleavage site in a target RNA may be in the form of ribonucleotides.

The respective 3' and 5' termini of the sequences $(X)_n$ and $(X)_m$ (of formula 1, 2, 3, 4, 1a, 1b, or 1c) or alternatively the 3' and 5' end termini of the ribozyme, may be modified to stabilize the ribozyme from degradation. For example, blocking groups may be added to prevent terminal nuclease attack, in particular 3'-5' progressive exonuclease activity. By way of example, blocking groups may be selected from optionally substituted alkyl, optionally substituted phenyl, optionally substituted alkanoyl. Optional substituents may be selected from C_1 - C_5 alkyl; halogen such as F, Cl or Br; hydroxy; amino; C_1 - C_5 alkoxy and the like. Alternatively, nucleotide analogues such as phosphothioates, methylphosphonates or phosphoramidates or nucleoside derivatives (such as alpha -anomers of the ribose moiety) which are resistant to nuclease attack may be employed as terminal blocking groups.

Alternatively, groups which alter the susceptibility of the ribozyme molecule to other nucleases may be inserted into the 3' and/or 5' end of the ribozyme (of formula 1, 2, 3, 4, 1a, 1b, or 1c). For example, 9-amino-acridine attached to the ribozyme may

act as a terminal blocking group to generate resistance to nuclease attack on the ribozyme molecules and/or as an intercalating agent to aid endonucleolytic activity. It will be readily appreciated that a variety of other chemical groups, e.g. spermine or spermidine could be used in a related manner.

5

It is also possible to stabilize the ribozyme (of formula 1, 2, 3, 4, 1a, 1b, or 1c) from degradation by embedding it in a RNA molecule, for example, as described hereinabove in formula 4. Such RNA molecules embedded with a ribozyme molecule may include, but are not limited to, tRNA, mRNA, rRNA or other RNA molecules.

10

Further, it is possible to insert the ribozyme in a DNA molecule as well.

15

Ribozymes of this invention (of formula 1, 2, 3, 4, 1a, 1b, or 1c) may be covalently or non-covalently associated with affinity agents such as proteins, steroids, hormones, lipids, nucleic acid sequences, intercalating molecules (such as acridine derivatives, for example 9-amino acridine) or the like to modify binding affinity for a substrate nucleotide sequence or increase affinity for target cells, or localization in cellular compartments or the like. For example, the ribozymes of the present invention may be associated with RNA binding peptides or proteins which may assist in bringing the ribozyme into juxtaposition with a target nucleic acid such that hybridization and cleavage of the target sequence may take place. Nucleotide sequences may be added to the respective 3' and 5' termini of the sequences (X)_n and (X)_n' of formulas 1-4 or alternatively the 3' and 5' end termini of the ribozyme to increase affinity for substrates. Such additional nucleotide sequences may form triple helices with target sequences [Strobel, 1991] which may enable interaction with intramolecularly folded substrate. Alternatively, modified bases (non-natural or modified bases as described in Principles of Nucleic Acid Structure [Sangar, 1984]) within the additional nucleotide sequences may be used that will associate with either single stranded or duplex DNA generating base pair, triplet, or quadruplet, interactions with nucleotides in the substrate. Suitable bases would include inosine, 5'-methylcytosine, 5'-bromouracil and other such bases as are well known in the art, as described, for example, in Principles of Nucleic Acid Structure [Sangar, 1984].

25

30

The ribozymes of this invention (of formula 1, 2, 3, 4, 1a, 1b, or 1c) may be produced by nucleotide synthetic techniques which are well known in the art, and described for example by Carruthers et al., Foehler et al. and Sprat et al. [Carruthers, 1987; Foehler, 1986; Sprat, 1984]. Generally, such synthetic procedures involve the sequential coupling of activated and protected nucleotide bases to give a protected nucleotide chain, whereafter protecting groups may be removed by suitable treatment. Alternatively, the ribozymes in accordance with this invention may be produced by transcription of nucleotide sequences encoding said ribozymes in host-cells or in cell free systems utilizing enzymes such as T3, SP6 or T7 RNA-polymerase. Further means for producing the ribozymes of this invention are further discussed hereinbelow [Sambrook, 1989].

Nucleotides represented in formulas 1, 2, 3, 4, 1a, 1b, and 1c comprise a sugar, base, and a monophosphate group or a phosphodiester linkages. Accordingly, nucleotide derivatives or modifications may be made at the level of the sugar, base, monophosphate groupings or phosphodiester linkages. It is preferred that the nucleotides in formulas 1-4 be ribonucleotides, however, other substitutions or modifications in the nucleotide are possible providing that endonuclease activity is not lost. More specifically, it is preferred that the catalytic region be comprised of ribonucleotides. In one aspect of this invention, the sugar of the nucleotide may be a ribose or a deoxyribose such that the nucleotide is either a ribonucleotide or a deoxyribonucleotide, respectively. Furthermore, the sugar moiety of the nucleotide may be modified according to well known methods in the art [See for example: Sangar, 1984; Sober, 1970]. This invention embraces various modifications to the sugar moiety of nucleotides as long as such modifications do not abolish cleavage activity of the ribozyme. Examples of modified sugars include replacement of secondary hydroxyl groups with halogen, amino or azido groups; 2'-methylation; conformational variants such as the O2'-hydroxyl being cis-oriented to the glycosyl C_{1'}-N link to provide arabinonucleosides, and conformational isomers at carbon C_{1'} to give alpha -nucleosides, and the like.

Accordingly, the base of the nucleotide may be adenine, guanine, cytosine, methyl cytosine, uracil, thymine xanthine, hypoxanthine, inosine, or other methylated bases. Nucleotide bases, deoxynucleotide bases, and ribonucleotide bases are well known in the art and are described, for example in Principles of Nucleic Acid Structure [Sangar, 1984]. Furthermore, nucleotide, ribonucleotide, and deoxyribonucleotide derivatives, substitutions and/or modifications are well known in the art [See for example: Sangar, 1984; Sober, 1970], and may be made with the proviso that endonuclease activity of the ribozyme is not lost. As mentioned previously, endoribonuclease activity may be readily and routinely assessed.

In addition, a large number of modified bases are found in nature, and a wide range of modified bases have been synthetically produced [See for example: Sangar, 1984; Sober, 1970]. For example, amino groups and ring nitrogens may be alkylated, such as alkylation of ring nitrogen atoms or carbon atoms such as N₁ and N₇ of guanine and C₅ of cytosine; substitution of keto by thioketo groups; saturation of carbon double bond carbon double bonds, and introduction of a C-glycosyl link in pseudouridine. Examples of thioketo derivatives are 6-mercaptapurine and 6-mercaptoguanine. Bases may be substituted with various groups, such as halogen, hydroxy, amine, alkyl, azido, nitro, phenyl and the like. The phosphate moiety of nucleosides or the phosphodiester linkages of oligonucleotides are also subject to derivatization or modifications, which are well known in the art. For example, replacement of oxygen with nitrogen, sulphur or carbon derivatives to respectively give phosphoramidates, phosphorothioates and phosphonates. Substitutions of oxygen with nitrogen, sulphur or carbon derivatives may be made in bridging or non bridging positions. It has been well established from work involving antisense oligonucleotides [Uhlmann, 1990] that phosphodiester and phosphorothioate derivatives may efficiently enter cells (particularly when of short length), possibly due to association with a cellular receptor. Methylphosphonates are probably readily taken up by cells by virtue of their electrical neutrality.

A further aspect of the invention provides alternative linkages such as an amide, a sulfonamide, a hydroxylamine, a thioformal, a formacetal, a 3'-thioformacetal, a sulfide, or an ethylene glycol function to replace the conventional phosphodiester linkage. These modifications may increase resistance towards cellular nucleases and/or improved pharmacokinetics.

Any combination of the above listed nucleotide modifications, substitutions, or derivatizations, made at the level of the sugar, base, or monophosphate groupings or phosphophodiester linkages may be made providing that endonuclease activity of the ribozymes (of formula 1, 2, 3, 4, 1a, 1b, or 1c) included in this invention are not lost.

As hereinabove discussed in formulas 1, 2, 3, 4, 1a, 1b, or 1c, base pairing between nucleotides are depicted through the use of the symbol "**". Base pairing may be Watson-Crick base pairing or Non-Watson-Crick base pairing. In some instances, an admixture of Watson-Crick and Non-Watson-Crick base pairing yields ribozymes with increased catalytic activity.

Ribozymes of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be incorporated and expressed in cells as a part of a DNA or RNA transfer vector, or a combination thereof, for the maintenance, replication and transcription of the ribozyme sequences of this invention. Nucleotide sequences encoding the ribozymes of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be integrated into the genome of a eukaryotic or prokaryotic host cell for subsequent expression (for example as described [Sambrook, 1989]. Genomic integration may be facilitated by transfer vectors which integrate into the host genome. Such vectors may include nucleotide sequences, for example of viral or regulatory origin, which facilitate genomic integration. Methods for the insertion of nucleotide sequences into a host genome are described for example in Sambrook et al. and Hogan et al. [Sambrook, 1989; Hogan, 1986; 1989].

Genomically integrated nucleic acid sequences encoding the ribozymes of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) generally comprise a promoter operably linked to the nucleotide sequence encoding the ribozyme of this invention, and capable of expressing said ribozyme in a eukaryotic (such as animal or plant cells) or prokaryotic (such as bacteria) host cells. Additionally, the ribozymes of the present invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be prepared by methods known per se in the art for the synthesis of RNA molecules. (For example, according to recommended protocols of Promega, Madison, Wis., USA). In particular, the ribozymes of the invention may be prepared from a corresponding DNA sequence (DNA which on transcription yields a ribozyme, and which may be synthesized according to methods known per se in the art for the synthesis of DNA) operably linked to an RNA polymerase promoter such as a promoter for T7 RNA polymerase or SP6 RNA polymerase. A DNA sequence corresponding to a ribozyme of the present invention may be ligated into a DNA transfer vector, such as plasmid or bacteriophage DNA. Where the transfer vector contains an RNA polymerase promoter operably linked to DNA corresponding to a ribozyme, the ribozyme may be conveniently produced upon incubation with an RNA polymerase. Ribozymes may, therefore, be produced in vitro by incubation of RNA polymerase with an RNA polymerase promoter operably linked to DNA corresponding to a ribozyme, in the presence of ribonucleotides. In vivo, prokaryotic or eukaryotic cells (including mammalian and plant cells) may be transfected with an appropriate transfer vector containing genetic material corresponding to a ribozyme in accordance with the present invention, operably linked to an RNA polymerase polymer such that the ribozyme is transcribed in the host cell. Transfer vectors may be bacterial plasmids or viral RNA or DNA. Nucleotide sequences corresponding to ribozymes are generally placed under the control of strong promoters such as, for example, the lac, SV40 late, SV40 early, metallothionein, or lambda promoters. Ribozymes may be directly transcribed in-vivo from a transfer vector, or alternatively, may be transcribed as part of a larger RNA molecule. For example, DNA corresponding to ribozyme sequences may be ligated into the 3' end of a carrier gene, for example, after a translation stop signal. Larger RNA molecules may help to stabilize the ribozyme

molecules against nuclease digestion within cells. On translation the carrier gene may give rise to a protein, whose presence can be directly assayed, for example, by enzymatic reaction. The carrier gene may, for example, encode an enzyme.

- 5 Ribozymes of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be involved in gene therapy techniques, where, for example, cells from a human suffering from a disease, such as HIV are removed from a patient, treated with the ribozyme to inactivate the infectious agent, and then returned to the patient to repopulate a target site with resistant cells. In the case of HIV, nucleotide sequences encoding ribozymes
10 of this invention capable of inactivating the HIV virus may be integrated into the genome of lymphocytes or be present in the cells a transfer vector capable of expressing ribozymes of this invention. Such cells would be resistant to HIV infection and the progeny thereof would also confer such resistance.
- 15 Ribozymes of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be incorporated and expressed in cells as a part of a DNA or RNA transfer vector, or a combination thereof, for the maintenance, replication and transcription of the ribozyme sequences of this invention. In-vivo, that is, within the cell or cells of an organism, a transfer vector such as a bacterial plasmid or viral RNA or DNA, encoding one or more
20 ribozymes (formulas 1, 2, 3, 4, 1a, 1b, or 1c), may be transfected into cells [See for example: Llewellyn, 1987; Hanahan, 1983]. Once inside the cell, the transfer vector may replicate and be transcribed by cellular polymerases to produce ribozyme RNAs which may have ribozyme sequences of this invention; the ribozyme RNAs produced may then inactivate a desired target RNA. Alternatively, a transfer vector containing
25 one or more ribozyme sequences may be transfected into cells or introduced into cells by way of micromanipulation techniques such as microinjection, such that the transfer vector or a part thereof becomes integrated into the genome of the host cell. Transcription of the integrated genetic material gives rise to ribozymes, which act to inactivate a desired target RNA.

Transfer vectors expressing ribozymes of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be capable of replication in a host cell for stable expression of ribozyme sequences. Alternatively, transfer vectors encoding ribozyme sequences of this invention may be incapable of replication in host cells, and thus may result in transient expression of ribozyme sequences. Methods for the production of DNA and RNA transfer vectors, such as plasmids and viral constructs are well known in the art and are described for example by Sambrook et al. [Sambrook, 1989]. Transfer vectors would generally comprise the nucleotide sequence encoding the ribozyme of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c), operably linked to a promoter and other regulatory sequences required for expression and optionally replication in prokaryotic and/or eukaryotic cells. Suitable promoters and regulatory sequences for transfer vector maintenance and expression in plant, animal, bacterial, and other cell types are well known in the art and are described for example in Hogan [Hogan, 1986; 1989].

The ribozymes of the present invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) have extensive therapeutic and biological applications. For example, disease causing viruses in man and animals may be inactivated by administering to a subject infected with a virus, a ribozyme in accordance with the present invention (such as given in formulas 1, 2, 3, 4, 1a, 1b, or 1c) adapted to hybridize to and cleave RNA transcripts of the virus. Such ribozymes may be delivered by parenteral or other means of administration. Alternatively, a subject infected with a disease causing virus may be administered a non-virulent virus such as vaccinia or adenovirus which has been genetically engineered to contain DNA corresponding to a ribozyme operably linked to an RNA promoter, such that the ribozyme is transcribed in the cells of the host animal, transfected with the engineered virus, to effect cleavage and/or inactivation of the target RNA transcript of the disease causing virus.

The ribozymes of the present invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) have particular application to viral diseases caused for example, by the herpes simplex virus (HSV) or the AIDS virus (HIV). Further examples of human and animal disease which may be treated with the ribozymes of this invention include psoriasis, cervical

preneoplasia, papilloma disease, bacterial and prokaryotic infection, viral infection and neoplastic conditions associated with the production of aberrant RNAs such as occurs in chronic myeloid leukemia. Diseases or infections which may be treated in plants with ribozymes of this invention include fungal infection, bacterial infections (such as Crown-Gall disease) and disease associated with plant viral infection.

The period of treatment would depend on the particular disease being treated and could be readily determined by a physician. Generally treatment would continue until the disease being treated was ameliorated. The ribozymes of the present invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) also have particular application to the inactivation of RNA transcripts in bacteria and other prokaryotic cells, plants and animals. In bacteria, RNA transcripts of, for example, bacteriophage, (which cause bacterial cell death) may be inactivated by transfecting a cell with a DNA transfer vector which is capable of producing a ribozyme in accordance with the present invention which inactivates the phage DNA. Alternatively, the ribozyme itself may be added to and taken up by the bacterial cell to effect cleavage of the phage RNA. Similarly, eukaryotic and prokaryotic cells in culture may, for example, be protected from infection or disease associated with mycoplasma infection, phage infection, fungal infection and the like.

RNA transcripts in plants may be inactivated using ribozymes (such as those of formulas 1, 2, 3, 4, 1a, 1b, or 1c) encoded by a transfer vector such as the Ti plasmid of *Agrobacterium tumefaciens*. When such vectors are transfected into a plant cell, the ribozymes are produced under the action of RNA polymerase and may effect cleavage of a specific target RNA sequence. Accordingly, plant viruses whose RNA sequence are known, or the RNA transcripts of plant genes, may be inactivated using ribozymes. Endogenous gene transcripts in plants, animals or other cell types may be inactivated using the ribozymes of the present invention (such as those of formulas 1, 2, 3, 4, 1a, 1b, or 1c). Accordingly, undesirable phenotypes or characteristics may be modulated. It may, for example, be possible using the ribozymes of the present invention to remove stones from fruit or treat hereditary diseases in humans which are

caused by the production of a deleterious protein, or over production of a particular protein. Furthermore, for the in-vivo applications of the ribozymes of this invention in humans, animals, plants, and eukaryotic and prokaryotic cells, such as in phenotypic modification and the treatment of disease, it is necessary to introduce the
5 ribozyme into cells whereafter, cleavage of target RNAs takes place. In vivo applications are highly suitable to the ribozyme of formula 4 as discussed in more detail hereinbelow.

Methods for the introduction of RNA and DNA sequences into cells, and the
10 expression of the same in prokaryotic and eukaryotic cells are well known in the art for example as discussed by Cotten and Friedman [Cotten, 1990; Friedman, 1989]. The same widely known methods may be utilized in the present invention.

The ribozymes of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be
15 incorporated into cells by direct cellular uptake, where the ribozymes of this invention would cross the cell membrane or cell wall from the extracellular environment. Agents may be employed to enhance cellular uptake, such as liposomes or lipophilic vehicle, cell permeability agents, such as dimethylsulfoxide, and the like. Ribozymes of the present invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be combined with
20 pharmaceutically and veterinarily acceptable carriers and excipients which are well known in the art, and include carriers such as water, saline, dextrose and various sugar solutions, fatty acids, liposomes, oils, skin penetrating agents, gel forming agents and the like, as described for example in Remington's Pharmaceutical Sciences, 17th Edition, Mack Publishing Co., Easton, Pa., Edited by Ostol et al., which is
25 incorporated herein by reference.

Agriculturally acceptable carriers and excipients are well known in the art and include water; surfactants; detergents; particularly biodegradable detergents; talc; inorganic and/or organic nutrient solutions; mineral earths and clays; calcium carbonate;
30 gypsum; calcium sulfate; fertilizers such as ammonium sulfate, ammonium phosphate and urea; and natural products of vegetable origin such as, for example, grain, meals

and flours, bark meals; and the like. The ribozymes of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be provided in a composition with one or more anti-viral, anti-fungal, anti-bacterial, anti-parasitic, anti-protazoan or anthelmintic agents, herbicides, pesticides or the like, for example as described in the Merck Index (1989) 11th Edition, Merck & Co. Inc.

By way of example only, therapeutic compositions of this invention may be directed against Herpes Simplex virus types 1 and 2, psoriasis, cervical preneoplasia, papilloma disease, and bacterial and prokaryotic infection. Such treatments may, for example, involve topical application of ribozyme to the site of disease. For example, in the treatment of Herpes virus lesions, ribozymes may be formulated into a cream containing a concentration of 1 nM to 1 mM ribozyme. The cream may then be applied to the site of infection over a 1 to 14 day period in order to cause amelioration of symptoms of the infection. Prior to the final development of topical formulations for the treatment of Herpes virus infection, effectiveness and toxicity of the ribozymes and formulations involving them may, for example, be tested on an animal model, such as scarified mouse ear, to which virus particles, such as 2×10^6 plaque forming units are added. A titre of infectious virus particles in the ear after treatment can then be determined to investigate effectiveness of treatment, amount of nuclease required and like considerations. Similar investigations in animal models prior to human trials may also be conducted, for example, in respect of the treatment of psoriasis, papilloma disease, cervical preneoplasia, and in diseases such as HIV infection, bacterial or prokaryotic infection, viral infection and various neoplastic conditions, which neoplastic conditions involve a deleterious RNA species.

Compositions for topical application are generally in the form of creams, where the ribozymes of this invention may be mixed with viscous components. The ribozymes of this invention (formulas 1, 2, 3, 4, 1a, 1b, or 1c) may be incorporated into liposomes or other barrier type preparations to shield the ribozymes from nuclease attack or other degradative agents (such as ribozymes and adverse environmental conditions such as UV light). Compositions may be provided as unit dosages, such

as capsules (for example gelatin capsules), tablets, suppositories and the like. Injectable compositions may be in the form of sterile solutions of ribozyme in saline, dextrose or other media. Compositions for oral administration may be in the form of suspensions, solutions, syrups, capsules, tablets and the like. Ribozymes may also be provided in the form of sustained release articles, impregnated bandages, patches and the like. Pharmaceutical compositions which may be used in this invention are described, for example, in Remington's Pharmaceutical Sciences.

The present invention is further directed to a DNA cassette for a plant, said cassette comprising a genetic sequence and a promoter capable of directing expression of said genetic sequences wherein said genetic sequence on expression provides anti-sense or ribozyme RNA to (-) RNA or a portion thereof associated with a viroid. The DNA cassette may further be part of a DNA transfer vector suitable for transferring the DNA cassette into a plant cell and insertion into a plant genome. In a most preferred embodiment of the present invention, the DNA cassette is carried by broad host range plasmid pGA470 and which is capable of transformation into plant cells using Agrobacterium. The present invention, however, extends to other means of transfer such as genetic bullets (e.g. DNA-coated tungsten particles, high-velocity micro projectile bombardment) and electroporation amongst others (Maliga, 1993; Bryant, 1992; or Shimamoto, 1989).

The transgenic plant resistant to a virus characterized in that it contains in its genome a sequence which gives rise, on transcription, to the nucleic acid molecule mentioned above. This transgenic plant, including fruits, and seeds thereof, may be from alfalfa, apple, bean, canola (oilseed rape), cantaloupe, corn, cotton, courgette, cucumber, melon, papaya, pepper, potato, rice, soybean, squash, strawberry, sunflower, sweet pepper, tobacco, tomato, or walnut. Also included are the plant cells transformed by the above-mentioned transfer vector, as well as a prokaryotic or eukaryotic cell, plant or animal, comprising a nucleotide sequence which is, or on transcription gives rise to, nucleic acid molecule.

The present invention will now be illustrated by way of non-limiting Examples only, with reference to the following non-limiting Examples, and Figures.

Throughout EXAMPLES 1-3, various Tables (1-8) are referenced by number. These
5 tables are hereby incorporated by number into the present application. Tables 1-8 can be found immediately preceding the references.

In EXAMPLES 1-3 below: The ribozyme variants were examined for cleavage
10 efficiency relative to the control sequence from satellite of tobacco ringspot virus (satTobRSV). The ribozymes were generated as in vitro T7 RNA polymerase transcripts and purified by denaturing polyacrylamide gel electrophoresis. The
15 standard reaction conditions were 50mM Tris-HCl pH8, 10mM MgCl₂, with a ribozyme to substrate ratio of about 10:1. Reactions were initiated by adding 1-4ng of substrate to ribozyme premixed with buffer and incubated at 37°C. Reactions were
20 stopped by adding formamide loading buffer with EDTA and the mixtures then fractionated by denaturing polyacrylamide gel electrophoresis. The products were quantified by phosphorimaging. The target RNA was a RNA 21 residue synthetic
substrate (5'-UUGCCCCUUCGCCAGCUC-3' (SEQ. ID. NO: XX)) and is based on part of the lacZ gene.

20 EXAMPLE 1

A. Helix II variants (Tables 1-5):

Most ribozyme variants of the satTobRSV helix II sequence reduced rates of cleavage. The exceptions fall into classes: loop sequence variants that probably increase the rate
25 and stability of stem formation, and two stem-length variants that give less stable stem structures.

i) Cleavage efficiency of helix II stem-length variants:

The middle two base-pairs of helix II were replaced with 0-6-A-U base-pairs and the
30 variants tested for cleavage efficiency (Table 1). The two outer G-C base-pairs were

retained in all cases. Internal A-U base-pairs were chosen as providing the smallest increments in stem stability.

The most notable feature is the lack of any simple correlation between stem-length and cleavage rates (2bp-3bp > control > 4bp=7bp=8bp > 5bp=6bp; Figure 1). Several

5 competing factors appear to influence cleavage efficiency:-

ii) stem-length variants that give increased cleavage efficiency (Table 1):

The two base-pair stem variant G-C, C-G (Table 1, #1) had a faster initial rate of cleavage than the control but the final proportion of cleavage products was reduced.

10 One three base-pair variant G-C, A-U, C-G (Table 1, #2) cleaved more rapidly and to the same extent as the satTobRSV-based ribozyme. However, the sequence of the stem appears to be important, as the three base-pair stem G-C, C-G, G-C (Table 1, #3) reduced cleavage efficiency.

15 iii) mutations of conserved G-C base pair in helix II (Table 2):

The G-C base-pair adjacent to the catalytic core is strictly conserved in naturally occurring hammerhead ribozyme sequences. Most mutations, however, could cleave the substrate with moderate efficiency, including one variant (A-G; Table 2, #1) that had similar cleavage rates to the control. Intriguingly, this variant gave negligible

20 cleavage in the system reported by Ruffner et al. [Ruffner, 1990].

A. Helix II variants (Tables 1-5):

i) Cleavage efficiency of helix II stem-length variants:

The middle two base-pairs of helix II were replaced with 0-6-A-U base-pairs and the variants tested for cleavage efficiency (Table 1). The two outer G-C base-pairs were

25 retained in all cases. Internal A-U base-pairs were chosen as providing the smallest increments in stem stability.

The most notable feature is the lack of any simple correlation between stem-length and cleavage rates (2bp-3bp > control > 4bp=7bp=8bp > 5bp=6bp; fig. 1). Several

30 competing factors appear to influence cleavage efficiency:-

- 1) stem stability, the 7 and 8 base-pair stems cleave faster than the 5 and 6 base-pair stems;
- 2) stem flexibility, the fastest ribozymes are those with 2 or 3 base-pairs (however the 2bp stem variant yielded a lower overall percentage of cleavage products); and
- 5 3) sequence context, ribozymes with the same number of base-pairs or similar thermodynamic stability cleave at different rates.

Presumably the 4 base-pair stem is the optimal solution in nature. Clearly, however, faster cleaving variants can be devised in any one particular case or conditions. It
10 would be worthwhile to test the robustness of the faster 3 base-pair variant (Table 1, #2) against other targets.

- ii) Cleavage efficiency of the conserved G-C base-pair in helix II. (Table 2):
The 5' G-C 3' base-pair in helix II, that is adjacent to the catalytic core, is conserved
15 in nature. Ruffner et al. [Ruffner, 1990] extensively mutated this ribonucleotide pair and found several examples with moderate but lower rates of cleavage. The results herein results correlate well with those findings (Table 2), including elimination of cleavage activity with a 5' C-G 3' base-pair (Table 2, #10). One notable exception was a cytosine to adenine substitution. Indeed, the 5' G-A 3' nucleotide pair gave
20 comparable cleavage rates relative to the control (Table 2, #1); whereas, Ruffner al. [Ruffner, 1990] reported almost no activity with this nucleotide pair.

As the 5'-G-C-3' base-pair eliminates cleavage activity while all other nucleotide pairs retain low to moderate cleavage efficiency, base-pairing of these two nucleotides of
25 helix II may not occur in the active ribozyme conformation. This notion is consistent with minizyme activity.

- iii) Cleavage efficiency of helix II loop variants. (Table 3):
Helix II loops that differed in both sequence and size were tested for cleavage
30 efficiency. The structure of RNA tetraloops have been studied extensively in vitro, showing differences in conformation and stem stability. Based primarily on ribosomal

RNA sequences, the tetraloops most commonly favored in nature are 5'-UUCG-3' and 5'-GNRA-3': both of which enhance stem stability. The loop of helix II from satTobRSV (5'-GUGA-3') conforms to the latter class. Not unexpectedly, one tetraloop variant with improved cleavage efficiency, 5'-GGGA-3' (Table 3, #1), also fits this pattern.

Interestingly, two penta-loops also increased cleavage rates (5'-CGGAA-3', Table 3, #14; and 5'-UUGGA-3', Table 3, #15). Unfortunately, the effect of penta-loops on stem stability has not been formally examined either structurally or thermodynamically.

All tri-loop variants reduced cleavage efficiency, probably by preventing formation of the adjacent base pair, leading to a penta-loop bounded by an A-U base pair and resulting in a stem of reduced stability.

In conclusion, although the loop sequence is redundant for cleavage activity, mutants can vary considerably in cleavage efficiency (Table 3): a few may actually increase cleavage rates, but many are likely to reduce cleavage rates.

iv) Cleavage efficiency of helix II stem mismatches:

The introduction of mismatches into the stem of helix II lowered cleavage efficiency (Table 4). There appears to be a good correlation between stem stability and cleavage efficiency (Table 4).

v) Cleavage efficiency of minizyme variants:

Of seven minizyme variants tested, only the UUUU (Table 5, #1) variant cleaved with moderate efficiency. In previous experiments with ribozyme and substrate arranged in cis, one minizyme 5'-GCGC-3', gave a higher percentage of cleave products after one hour than the UUUU minizyme. In trans, cleavage with the GCGC minizyme occurred only in the first of several experiments. This particular ribozyme will need to be resequenced.

In view of the large variation in cleavage efficiencies between minizymes, and previous results with the GCGC variant, it is possible that there are minizymes capable of faster cleavage rates than the UUUU (Table 5, #1) minizyme. Interestingly, the 5' and 3' nucleotides of the UUUU (Table 5, #1) and GCGC minizymes also correspond in sequence and position to highly favored nucleotide pairs in the full-sized helix II (see A.ii and Table 2; and [Ruffner, 1990]).

EXAMPLE 1A

Additional Ribozyme assays of stem II variants

10

Stem II consists of highly conserved tandem ((anti)-A(anti) base-pairs proximal to the catalytic core and a helical extension with canonical Watson-Crick base pairing (Haseloff et al. 1988). In many hammerhead self-cleaving domains, stem II is linked by a three to five nucleotide loop (Hertel et al. 1992). The nucleotide sequences of the stem extension and loop are highly variable. To assess the importance of the stem II variable region for cleavage activity, mutations were created by nucleotide substitutions, additions or deletions in either the stem or the loop (Figure 4).

15

Mutant ribozymes with shorter stems of either five or six base pairs (Rz 2-5; Figure 4) had rate constants up to 4.4 times greater than the seven base pair parent sequence. Reduced stem stability of the shorter stem ribozymes is likely to contribute to the increases rate constants, but further destabilization can greatly reduce cleavage rates. For example, increased loop size of five or six base paired ribozyme stem II variants (Rz 6-12; Figure 4) sometimes gave very low relative rate constants (e.g. Rz 7-10). Similarly, Tuschel and Eckstein (1993) reported a modest 1.5 fold rate increase for one five base pair stem II ribozyme variant, but different loop sequences in this variant gave ribozymes with slower cleavage rates. In addition, low stem II stability in the form of a three base pair minizyme (McCall 1992) resulted in a 20 fold reduction in rate constant (Figure 2: Rz 13), similar to reports by Tushel and Eckstein (1993) and Long and Uhlenbeck (1994).

25

30

Besides stability, the sequence and size of stem II appears to have additional effects that influence cleavage efficiency. For example, the length of stem II was increased incrementally with A-U base pairs (Figure 4: Rz 14-18) but there was no linear relationship between stem length and rate constant. The ribozyme with ten base pairs (Rz 17) had faster cleavage rates than the eight (Rz 15), nine (Rz 16) and eleven (Rz 18) base paired variants but slower than the seven base paired forms (Rz 1, 14; Figure 4).

Variations in the size and sequence of the loop were also examined by randomly substituting three, four or five nucleotides for the parental tetraloop sequence 5'-GUGA-3' (Figure 4; Rz 19-38). Although the majority of ribozyme loop variants had reduced rate constants, two ribozymes with tetraloops (Rz 22, 23) and two with pentaloops (Rz 33, 34) had increased cleavage rates.

Naturally occurring stem-loop structures within ribosomal RNAs strongly favor tetraloops with the consensus sequence 5'-GNRA-3' (N, any nucleotide; R, purine; 25). The tetraloop sequence of the parent ribozyme 5'-GUGA-3' fits this pattern, as does one of the faster cleaving variants, Rz 22, 5'-GGGA-3'. However, a second faster cleaving tetraloop variant, Rz 23, 5'-GUGG-3', differs slightly from the 5'-GNRA-3' pattern. None (All) of the slower cleaving variants (Rz 24-32) do not conform to the 5'-GNRA-3' pattern. Similarly, Tuschel and Eckstein (1993) reported faster cleavage rates with a loop variant 5'-GAAA-3' than two other non-5'-GNRA-3' variants. It has been suggested that 5'-GNRA-3' sequences are biologically favored due to increased stability of the adjoining stem (Uhlenbeck 1990) or to specific three-dimensional conformation (SantaLucia et al. 1992). Another possibility is a greater capability of 5'-GNRA-3' sequences to form tertiary interactions (Biou et al. 1994; Pley et al. 1994 B). Interestingly, one tetraloop sequence, 5'-UUCG-3', that is also overrepresented in ribosomal RNAs (Woese et al. 1990) and adopts a similar conformation to the 5'-GNRA-3' tetraloop (Cheong, C. et al. 1990; Heus et al. 1991), give rise to a ribozyme variant (Rz 32) with the lowest rate constant of all tetraloop mutants examined (Figure 4).

Systematic structural and functional studies of RNA triloops and pentaloops have not been reported. However, in our random sampling of six pentaloop variants (out of 1024 possible combinations), 33% gave increased rate constants. All three triloop variants that were tested, gave reduced rate constants.

5

EXAMPLE 2

B. Helix I and III variants (Tables 6 and Table 7 respectively):

10 i) The helix I (Table 6) variants demonstrate that maintenance of a stable base-pair immediately adjacent to the cleaved ribonucleotide residue is critical for good cleavage rates.

15 ii) One helix I variant (Table 6, #1), in which continuous base-pairing extends for only three base-pairs, gave markedly increased cleavage rates.

15 iii) Increased stem stability of helix III generally correlated with increased cleavage rates, so long as the base-pairs proximal to the catalytic core were maintained.

B Helix I and III variants. (Tables 6 and Table 7):

20 Table 6 depicts cleavage efficiency of helix I variants. The two most significant findings are the crucial role of the base pair adjacent to the cleavage site and the possibility of significantly improving cleavage rates by reducing stem stability.

25 The substitution of the C residue with either A or G in the base pair adjacent to the cleavage site eliminated cleavage activity (Table 6, Table 7, and Table 8). Mutating the C residue to a U, allowing a G-U bond to form, greatly reduced cleavage efficiency (Table 6, #2). Even the strictly conserved G-C pair in helix II can be more readily modified (Table 2).

30 Stem mismatches elsewhere in helix I, however, sometimes gave increased cleavage rates. The most notable example includes one mutant in which only the first three base

pairs adjacent to the cleavage site are retained before mismatching occurs (Table 6, #1). It is uncertain if the marked improvement in cleavage efficiency is due to structural properties of this modified helix I mutant or reduced stem stability. One simple test would be to use a shorter substrate as the target.

5

Unlike helix I, all helix III stem mismatches gave reduced cleavage rates. Cleavage efficiency seems most closely correlated with increased stem stability; with particular emphasis on maintenance of base pairing adjacent to the cleavage site. One minor anomaly appears to be the sequence 5'-AAUUGUGGGA-3' (Table 7, #3) which appears to have poor base pairing possibilities and yet retains moderate cleavage efficiency.

10

In conclusion, helix III appears to have greater sequence specificity than helix I. It is uncertain, however, if this applies to other substrate sequences. At present, most short-armed ribozymes target an equal number of base pairs either side of the target. An uneven distribution of base pairs may prove optimal.

15

EXAMPLE 3

C. Enhancement of cleavage rates by complementary oligonucleotides. (Table 8):

20

i) DNA oligonucleotides of 8-10 residues complementary to either the catalytic core or the hybridizing arms of the ribozyme, increased cleavage rates.

ii) This enhancement of cleavage efficiency could be mimicked by complementary RNA oligonucleotides of six residues, but not by free nucleotides, spermine or polyethylene glycol, or denaturants such as formamide or urea.

25

C. Enhancement of cleavage rates by complementary oligonucleotides. (Table 8):

A series of different sized oligodeoxynucleotides were devised to complement different regions of the ribozyme. The aim was to test which regions of the ribozyme are most sensitive to interference by competing oligonucleotides. The oligodeoxynucleotides

30

were pre-incubated with ribozyme in 5X excess, at 100°C/1' and snap cooled on ice. Specific binding of the oligonucleotides to the ribozyme was tested by RNaseH digestion. Surprisingly, these substrate analogs, rather than acting as typical enzyme inhibitors, actually increased cleavage efficiency (Table 8, #1-11). However, there
5 appears to be an optimal stability range for the oligonucleotide-ribozyme interaction: high enough to allow binding, but low enough to allow displacement by the substrate. Improved cleavage efficiencies could also be achieved with complementary oligoribonucleotides (Table 8, #12-16). One possible explanation is that the oligonucleotides prevent formation or stabilization of inactive ribozyme conformers.

10 Enhancement of cleavage efficiency could not be achieved by the addition of free deoxynucleotide residues, spermine or polyethylene glycol, or by denaturants such as urea or formamide (Table 8, #17-28).

15 It may be useful to determine if the complementary oligonucleotide sequences could enhance activity when added in cis as part of the ribozyme transcript.

EXAMPLE 4

20 In vivo studies on the ribozymes of formula 4:

The present invention is directed to a plant tyrosine tRNA (Stange and Beier, 1988) to deliver hammerhead ribozyme and antisense RNAs against a reporter gene, chloramphenicol acetyl transferase (CAT), to plant protoplasts using a vector derived
25 from African cassava mosaic geminivirus (ACMV - Stanley, et al.). Geminiviruses are plant DNA viruses that rely on host components for replication, are localized in the nucleus and can replicate to high levels. For these reasons geminiviruses provide the basis for excellent vector systems for assaying plant gene expression.

30 Compared to work of Cotten and Birnsteil (1989), the present invention is different in several respects. First, in the present invention, a tyrosine tRNA from Nicotiana

(tobacco) was used instead of methionine tRNA-Xenopus. Second, the present invention uses a CAT mRNA target which is synthesized in vivo (nuclear/cytoplasmic). Cotten and Birnsteil used a U7snRNA which was synthesized in vitro and then microinjected (cytoplasmic). The target site on the RNA target molecule was different as well; i.e. - a GUC (present invention) verses a CUC (Cotten and Birnsteil). The present invention used an antisense control as well as a mutant target in order to test the activity of the tRNA embedded ribozyme. Cotten and Birnsteil did not use such controls. Finally, most of the active tRNARz12 transcripts of the present invention are RNA polymerase III derived.

The present invention is directed to the design of improved hammerhead ribozymes (a) in vitro and (b) in vivo.

a) in vitro

In the present invention, new hammerhead ribozymes were designed from experiments performed on the satellite RNAs of tobacco ringspot virus (sTobRSV). Some of the ribozymes developed show higher rates of cleavage *in vitro* than the native sTobRSV ribozyme. The present invention is also directed to compositions partly consisting of these new hammerhead ribozymes.

b) in vivo

The present invention is also directed to tRNA embedded hammerhead ribozymes. A tRNA embedded hammerhead ribozyme is created by embedding a hammerhead ribozyme into a tRNA molecule. The tRNA embedded hammerhead ribozyme shows decreased effectiveness in vitro but significant enhancement in its activity *in vivo*, relative to the non-embedded ribozyme. These results were based on in vivo cleavage of a target CAT mRNA molecule. Control studies further support the observed reductions in CAT activity as due to tRNA embedded hammerhead ribozyme function in vivo. The controls included the use of antisense molecules.

Materials and Methods:

Plasmid constructions:

All oligodeoxyribonucleotides were synthesized on an Applied Biosystems model 392 DNA synthesizer (Applied Biosystems). Molecular cloning and related techniques were carried out essentially as described by Sambrook, et al. 1989).

5

(i) **tyrosine tRNA.** A clone of the tyrosine tRNA was obtained from the laboratory of Hildberg Beier [Stange and Beier, 1986]; Figure 9a) and subcloned as an *AccI/SspI* fragment into like digested pGEM3zf-(Promega) to make pGEMtRNA. A short oligodeoxyribonucleotide, containing three restriction enzyme recognition sites [BamHI, *SmaI*, *EcoRI*], was inserted within the anti codon loop of the tRNA by site directed mutagenesis [Kunkel, et al., 1987] to make pGEMtRNApoly. This is the site of subsequent insertions of both ribozyme and antisense sequences.

10

(ii) **Ribozyme/Antisense (Figure 9b).** Overlapping oligodeoxyribonucleotides encoding ribozyme Rz12 contained the hammerhead motif plus 12 bases upstream and downstream of this motif. The double stranded oligodeoxyribonucleotide was cloned into *EcoRI PstI* digested pGEM4z to make pGEMRz12. This ribozyme is designed to cleave CAT mRNA at position 464 as shown in Figure 9c. Antisense control, representing the equivalent hybridization to the CAT mRNA as the ribozyme sequence, is as described in Cameron and Jennings (1989). The ribozyme was subcloned into *SmaI* digested pGEMtRNApoly as an end-filled *EcoRI PstI* fragment to make the clone pGEMtRNARz12. The antisense sequence was subcloned as a BamHI *SnaBI* fragment into BamHI *SmaI* digested pGEMtRNA poly to make the clone pGEMtRNAAs24.

20

25

(iii) **CAT substrate.** The CAT sequence was as described in Haseloff and Gerlach, 1988 and Perriman, et al., 1993. Sequence length and ribozyme target site are diagrammed in Figure 9c.

30

(iv) **Mutant CAT substrate.** The GUC target site at position 464 on the CAT gene was mutated to a GUG by site directed mutagenesis [Kunkel, et al., 1987] using a

specifically designed oligodeoxyribonucleotide. This mutation maintains codon usage and does not affect in vivo CAT activity. The plasmid bearing mutated CAT sequence is called pGEMCM-2.

- 5 (v) **Geminivirus vector construction.** The infectious clone of the A component of the geminivirus African cassava mosaic virus (ACMV) was a gift from Stanley, et al., (1988). Modification to this vector included the insertion of a pUC19-derived plasmid carrying, bacterial origin of replication and chloramphenicol resistance gene [Chay and Feldstein, unpublished; Figure 9d). In this construction a pUC19-derived poly
10 linker ant T7 and SP6 RNA polymerase recognition sites were inserted in place of the coat protein open reading frame. Into this region were subcloned the wildtype (pACMVCAT) and mutant (pACMVCM2) targets as well as non-embedded (pACMVRz12, pACMV As24) and tRNA-embedded (pACMVtRNARz12, PACMVtRNAAs24) ribozyme and antisense sequences. Upstream of all sequences
15 is the endogenous coat protein promoter, while downstream are coat protein termination and polyadenylation signals.

In vitro cleavage assay:

- 20 RNAs for the in vitro assays were prepared by in vitro transcription as follows: 1-2mg linearized template DNA, 40mM Tris-HCl (pH 7.5) 6mM MgCl₂, 2mM spermidine, 10mM NaCl, 10mM DTT, 8 units RNasin (Promega), 1mM rATP, rGTP, rUTP and 0.25mM rCTP (Pharmacia) in a final volume of 50ml. CAT and CM2 target RNA were transcribed in the presence of 60pmoles of [α -³²P] CTP (Amersham; 400Ci/mmol) using 25 units of T7 RNA polymerase (Bethesda Research Labs). The
25 ribozyme sequences, pGEMRz12 and pGEMtRNARz12, were transcribed in the presence of 1mM CTP and 25 units of RQ-1 DNaseI (Promega) to remove the DNA template. Transcript yields and the size and fidelity of RNAs were confirmed by electrophoresis through 6% polyacrylamide gel in 7M urea (denaturing PAGE).
- 30 CAT or CM2 target RNAs and ribozyme (Rz12 or tRNARz12) were mixed on ice prior to the addition of buffer (10mM MgCl₂, 50mM Tris-HCl, pH7.4). The

reactions were immediately transferred to 30°C for one hour. Reactions were terminated by the addition of EDTA to a final concentration of 50mM followed by ethanol/sodium acetate precipitation. Cleavage rates were quantitated using an AMBIS Image Acquisition analyzer (AMBIS, Inc.).

5

Analysis of in vivo expression:

(i) **Transfection.** A cell culture of *Nicotiana tabacum* (cv. Xanthi) was used for protoplast isolation essentially as described by Howard, et al. (1987). Protoplasts were suspended at a concentration of 1.2×10^6 /ml, with 700ml aliquot used per transfection. Transfection was obtained by electroporation using a Hoefer PG200 progenitor II apparatus (Hoefer Scientific Instruments, San Francisco, California). Each electroporation involved the co-electroporation of the target construct (either pACMVCAT or pACMVCM2) with (a) control pACMVtRNA or either (b) the antisense constructs (pACMVAs24 or pACMVtRNAAs24), or (c) the ribozyme constructs (pACMVRz12 or pACMVtRNARz12), so that each event involved the same amount of input DNA. The amount and ratio of target to control, antisense or ribozyme construct was 1:3 (5mg pACMVCAT or pACMVCM2: 15mg pACMVtRNA, pACMVAs24, pACMVtRNAAs24, pACMVRz12 or pACMVtRNARz12). This results in approximately a six-fold molar excess of tRNA-embedded antisense/ribozyme inserts over the target molecule. Each construct pairing was replicated a minimum of three times for each protoplast isolation. The conditions for electroporation were 490mF capacitance, 330 volts, single 8msec pulse with the electrodes in the electroporation chamber separated by 0.4mm. Following electroporation, cells were transferred to culture dishes and incubated in 3ml growth media (5% coconut water, 95% MS organics [Gibco-BRL], 265mM mannitol, 0.1mg/ml kinetin, 0.2mg/ml 2-4-D, 1.5mM KH_2PO_4 , 0.5mg/ml each of nicotinic acid, pyroxidine and thiamine) at 26°C for three days. At this time cells were harvested and assayed for CAT activity, ACMV DNA replication and mRNA levels.

(ii) **CAT assays.** Following incubation, 0.5 ml of cell suspension were harvested for CAT assays. The remaining 2.5ml were used to isolate nucleic acids. Cells were

30

collected at 80 x g for 10 mins and resuspended in 200 ml 0.25M Tris-HCl (pH7.4) before being transferred to 1.5ml eppendorf tubes. The cells were sonicated and extracts were centrifuged at 14,000 x g for 10 mins to remove cell debris. Protein concentrations were determined using Bradford assay kit (Biorad). Equivalent
5 amounts of protein for each extract were incubated at 37°C for 60 mins in the presence of 5ml of 10mM acetyl-Coenzyme A (Sigma) and 18 picomoles of C¹⁴ chloramphenicol (Amersham). Reactions were stopped by the addition of 700ml ethyl acetate. The ethyl acetate phase was dried and suspended in 10ml of ethyl acetate for separation on silica gel thin layer chromatography in 5% methanol and 95%
10 chloroform. Radioactive products were detected by autoradiograph and then quantitated using an AMBIS Image Acquisition analyzer to determine the proportion of C¹⁴ chloramphenicol that had been acetylated. The average rates of acetylation were plotted using Lotus Freelance graphics (Lotus, Cambridge, MA).

15 (iii) DNA isolation. Cells were collected as for the CAT assays and resuspended in 200ml TE. An equal volume of 2 x SDS extraction buffer (0.1mM Tris, pH7.4, 2mM EDTA, 2% SDS) was added and the solution mixed. Proteinase K (Boehringer Mannheim) was added to 100mg/ml and the solution incubated at 37°C for 30 minutes. Following this, samples were extracted with equal volumes of
20 phenol/chloroform and then nucleic acids precipitated in ethanol/sodium acetate. Nucleic acid samples were treated with 10mg RNase A (Sigma) to remove contaminating RNA prior to restriction enzyme digestion. DNAs were suspended at a concentration of 0.5mb/ml. Routinely 1-2mg of total DNA was digested with either DpnI or MboI and analyzed by southern blot analysis.

25 (iv) RNA isolation. Total RNA was isolated from cells using Trizol solution (Gibco BRL) as per manufacturers specifications. RNA was suspended at a final concentration of 1mg/ml and was analyzed by electrophoresis through non-denaturing 1% agarose.

30

(v) **Reverse transcriptase-PCR.** (RT-PCR) analysis of in vivo CAT mRNA. First strand cDNA synthesis was carried out with 5mg of total RNA using 200 units of Superscript reverse transcriptase (Gibco-BRL). cDNA synthesis was carried out at 45°C to minimize any interference by RNA secondary structure. The first strand primer was oligo dT with an 18 base unique sequence ("TAG" - Lincoln & Karrer, unpublished) at the 5' end of the primer. This TAG sequence was used to prime for subsequent PCT amplifications. Dual PCR reactions were carried out on reaction samples. Primer 1 which primes at positions 198 to 216 on the CAT sequence was designed to amplify full length CAT sequence. Primer 2 anneals at positions 470 to 488, just 3' of the ribozyme cleavage site and acts as a positive control for the presence of CAT RNA, cleaved or uncleaved. Conditions for the PCR reaction were verified to be within linear range of amplification and were as follows: 52°C 22 mins, 72°C 40 mins, 30 cycles of 94°C 40secs; 52°C 2 mins; 72°C 3 mins followed by 72°C 15 mins. Amplification was carried out using either 1 + TAG or 2 + TAG primer pairs. 1/6 of the PCR reactions were loaded on agarose gels and southern blotted to determine the extent of amplification of the CAT sequence by the two primer pairs. Southern blotting was carried out as described in Sambrood, et al. (1989). Filters were analyzed by autoradiograph and total counts determined using an AMBIS image acquisition analyzer.

Results:

Efficiency of in vitro cleavage by the linear and tRNA embedded ribozymes:

In vitro cleavage assays were carried out to test the efficiency of the linear and tRNA embedded ribozyme constructs in cleaving the CAT transcript. As shown in Figure 10a, cleavage of the CAT RNA was dependent upon the addition of either the linear or tRNA-embedded in ribozyme sequences. Both constructs cleaved the CAT RNA efficiently although as shown in Figure 10b, the non-embedded ribozyme (linear) cleaved more efficiently than the tRNA-embedded form. In the present example, a reduction in substrate was used as a measurement of cleavage in these in vitro assays. This quantity is the percentage of substrate calculated to have been cleaved to form

the two product bands visible in Figure 10a after one hour incubation at 30°C, based on radioactivity associated with bands for transcript and the two cleavage products.

In three independent experiments CAT RNA was cleaved to the average extent of 75% when incubated with the linear ribozyme. For the tRNA-embedded ribozyme, the average extent of CAT RNA cleavage was only 38%. These results, including standard deviations, are shown in Figure 10b. Incubation of the CAT transcript with the linear and tRNA-embedded antisense transcripts did not yield these cleavage products. Additionally, incubation of the mutant target transcript, CM2, with the two ribozyme transcripts also showed no accumulated cleavage products (data not shown).

Significantly reduced levels of in vitro CAT activity in the presence of the tRNARz12 ribozyme:

Having shown that both ribozymes are capable of cleaving the substrate RNA in vitro, the relative effects of these on CAT gene expression from Xanthi tobacco suspension cells was examined. In seven separate experiments using different protoplast preparations, the addition of the tRNA-embedded ribozyme gene construct resulted in greater suppression of CAT activity than was observed after the addition of the constructions expressing the antisense sequence in the non-embedded or tRNA-embedded form or a construction bearing the non-embedded ribozyme (Figures 10a and 10b). These results are in contrast to those obtained for the in vitro assays, in which the non-embedded ribozyme was more efficient at cleaving the CAT transcript. The mutated CAT target, CM2, did not show the enhanced CAT gene suppression observed for the wildtype CAT target. A CAT assay showing the effect on CAT activity compared with the effect on CM2 activity of the tRNARz12 transcript is shown in Figures 12a and 12b. Results obtained for each experiment were normalized by assignment of a value of 100% to the expression of pACMVCAT or pACMVCM2 in the presence of pACMVtRNA and in the absence of any of the antisense or ribozyme constructs (Figures 11a and 11b).

As shown in Figures 11a and 11b, the average reduction of the CAT gene expression, when the antisense gene construct is delivered as a non-embedded sequence is to about 60% of the control value. By comparison, the non-embedded and tRNA-embedded ribozymes were able to reduce CAT activity to 40% and 15% of controls, respectively. When the target sequence was CM2, which is the CAT target, all four constructs reduce CAT activity to the 60% level. Table 9 shows T-test analysis of pair-wise comparisons of CAT activity in the presence of either of the ribozyme or antisense constructs and the CAT and CM2 target. These T tests include each data point which make up the means shown in Figures 12a and 12b. For each construct pairing this represents 24 separate electroporation events. At > 99% confidence level,

1. The tRNARz12 construct caused significantly lower levels of CAT activity than either of the antisense or the non-embedded ribozyme sequences when targeting the CAT construct,
2. CAT activity for the CAT construct in the presence of the tRNARz12 construct was significantly lower than that for the CM2 mutant target construct, with the same ribozyme constructs produced significantly different levels of CAT activity.
3. For the CM2 target, none of the two antisense or two ribozyme constructs produced significantly different levels of CAT activity.

Table 9: T-test analysis of co-transfected CAT (CATMUT2) + antisense or ribozyme constructs. T-tests were carried out comparing each of the points which make up the means shown in Figures 11a and 11b. Partwise comparisons of each of two construct combinations were carried out. A T-value greater than 99% indicates the values are significantly different in one another.

	Co-transfections	T value > 99% level of Probability
	CAT + Rz12	YES
	v CAT + As24	
5	CAT + tRNAAs24	YES
	v CAT + tRNARz12	
	CAT + tRNARz12	YES
	v CAT + Rz12	
10	CAT + tRNAAs24	NO
	v CAT + As24	
	CAT + tRNARz12	YES
	v CM2 + tRNARz12	
	CAT + tRNAAs24	NO
	v CM2 + tRNAAs24	
15	CAT + As24	NO
	v CM2 + As24	
	CAT + Rz12	NO
	v CM2 + As24	
	CM2 + tRNARz12	NO
20	v CM2 + tRNAAs24	
	CM2 + Rz12	NO
	v CM2 + As24	

Analysis of replication of pACMV target, antisense and ribozyme constructs To ensure that the observed reduction in CAT activity was not due to inhibition of replication of the pACMVCAT or pACMVCM2 construct, in the presence of the tRNARz12 ribozyme, DNA was analysed at one and three days post transfection in the presence and absence of the tRNARz12 ribozyme. Methylation sensitive isoschizomers, DpnI and MboI were used to differentiate replicated from input pACMV sequences. As, shown in Figures 14a and 14b, both the CAT and CM2 constructs replicate to equivalent levels in the presence or absence of the tRNARz12 construct. Also, all four pACMV expressing ribozyme and antisense constructs replicate. Reduced levels of accumulated CAT mRNA correlate with reduced CAT activity: The effect the ribozymes had on the accumulation of CAT mRNA within the cell was examined using Figures 13a and 13b which shows a southern blot of two PCR amplified products derived from total RNA isolated from protoplasts electroporated with CAT target and either of the two antisense or two ribozyme constructs. The blots, which were probed with radiolabelled CAT sequence, show amplification products for both the CAT and CM2 target sequences. The two amplified products represent either full length (primers 1 + TAG) or truncated (primers 2 + TAG) CAT mRNA. The truncated is derived from amplification of CAT message downstream of the GUC target site and therefore provides an internal control to observe ribozyme mediated cleavage. As shown in Table 10, it was observed that amplification of these two products approximated a 1:1 ratio in the

presence of all antisense/ribozyme constructs for both the CAT and CM2 targets except in the case of CAT + tRNARz12. For the tRNARz12 ribozyme, the ratio of the downstream product to the full length product was 3.3. That is, the product amplifying downstream of the cleavage site was amplified three times more than the full length product. Amplification has been determined to be in the linear range by carrying out a time course of cycles from 10 through 40 with aliquots removed at 10 cycle intervals (data not shown). With confidence, it may be said that the amplification obtained is representative of the actual amounts of CAT mRNA present within each of the isolated RNA samples.

Table 10: Ratio of accumulation of 3' product to full length CAT product. (An explanation of these products is given hereinabove as well as in Figures 13a and 13b which illustrate a southern blot of these products.) The Table below shows the ratio of these products for each of the construct combinations. These ratios are the accumulated data from two independent experiments.

+	CAT	tRNA	tRNA	Rz12	As24
	tRNA	Rz12	As24		
	1.05	3.3	0.85	1.24	0.77

+	CM2	tRNA	tRNA	Rz12	As24
	tRNA	Rz12	As24		
	0.75	1.28	0.78	0.83	0.73

Discussion:

The present invention shows that transfection of tobacco protoplasts with self-replicating vectors expressing both a target sequence, CAT, and a tRNA-embedded ribozyme that is capable of cleaving the CAT message, results in significant reduction in CAT activity. Unlike previous *in vivo* ribozyme mediated destruction over substrate sequence to the cell [Cameron and Jennings, 1989; Cotten and Birnsteil, 1989; Sioud, et al., 1992; Steinecke, et al., 1992; Perriman, et al., 1993]. The results obtained herein from the internal control, in which the GUC target site on the CAT mRNA has been mutated to a GUG, suggest that the reduction observed for the wildtype CAT sequence is indeed a ribozyme mediated one. That is, the mutant target, CM2, showed an inhibition in the presence of the tRNARz12 sequence comparable to what was observed for antisense constructions, suggesting that the

increased reduction observed for the wildtype target is due to the ribozyme cleavage of the CAT mRNA.

5 This conclusion is also supported from the analysis on the accumulation of CAT mRNA within the electroporated protoplasts. Reverse-transcriptase PCR on RNA obtained from transfected tobacco cells showed that the ratio of the downstream PCR product to the full length product was markedly elevated, demonstrating intracellular cleavage. The ratio of 3.3 reported herein is similar to that obtained by Cantor, et al. 1993] and Dropulic, et al. (1992) in assays using a ribozyme against HIV. While 10 it was possible to observe reduction in full length CAT mRNA, it was not possible to detect cleavage products using an RNase protection assay. Reports on the stability of CAT message suggest that in stably transformed cells, the half-life is around 70 minutes. In transiently transformed cells, the half-life may be shorter [Gallie, et al., 1991]. In considering that the cleavage products would also have elevated 15 susceptibility to exonuclease activity, the chances are low of protecting, and therefore observing intracellular accumulation of cleavage products. Instability of the cleavage products also suggests that the observed ratio for the RT-PCR amplification in the presence of the tRNARz12 ribozyme may be greater than that pertaining, if all cleavage products were detected. Much of the 3' cleavage product, which is 20 represented by the truncated PCR product (primers 2 + TAG), once produced intracellularly, is probably rapidly degraded.

Unlike the results of Cotten and Birnsteil (1989), a reduced level of in vitro cleavage by the tRNA-embedded ribozyme over that obtained for the non-embedded ribozyme 25 was observed. Despite this reduced level in vitro, the tRNA-embedded ribozyme presented herein significantly reduced CAT activity in vivo. The tRNA constructions were designed so that two putatively active promoter sequences are present. The first being the endogenous RNA polymerase II based coat protein promoter and the second is the RNA polymerase III promoter internal to the tRNA sequence. Although Cotten and Birnsteil observed reduced transcription of their tRNARz construct over that of 30 the wildtype tRNA sequence, it was noted that no such reductions occurred. The

predominant transcript in the chimeric system presented herein was the RNA polymerase III transcript (manuscript in preparation). This is similar to results obtained by Kinsey and Sandemeyer (1991) when delivering tyrosine tRNA to yeast cells containing an upstream RNA polymerase II promoter. Additionally, it was found that processing rates of the tRNARz12 and tRNAAs24 transcripts were comparable to wildtype tRNA in S100 wheat germ extracts (manuscript in preparation). From these results one can conclude that most of the active tRNARz12 transcripts are RNA polymerase III derived.

Demonstration that hammerhead ribozymes can specifically target and cleave specific mRNA sequences in vivo in both plant and animal systems has been limited. While some success has been obtained in delivering single non-embedded ribozymes in animal systems [Sarver, et al., 1990; Droupilic, et al., 1992; L'Huillier, et al., 1992; Cantor, et al., 1993; Zhao, et al., 1993], it is likely that the ribozyme requires additional stabilizing sequence 5' and/or 3' to enable sufficient time to anneal to and cleave the designated target. While Sullenger and Cech (1993) show an enhanced effect of ribozymes when the target and ribozyme are sequestered within the same molecule, this approach is certainly not applicable to many therapeutic situations. This is particularly the case if the target sequence is plant or animal genome derived or of viral origin.

One could believe that tRNA molecules are excellent candidates for enhancing the ribozymes effectiveness in vivo. The results of Cotten and Birnstein (1989) in *Xenopus* oocytes combined with the results presented herein on plant cells suggest that a tRNA delivery system has a number of advantages over traditional approaches to gene stability. Firstly, the tRNA is a small and ubiquitous sequence. Secondly, RNA polymerase III may transcribe to high levels within the cell, thus providing us with a means of producing large amounts of the desired transcript. Finally measurements of tRNA stability within plant systems estimate the half-life to be in the order of 45 hours [Karnail and Wasternack, 1992] making them ideal for delivery of ribozyme sequences.

- 52A -

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group

5 of integers or steps.

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
22

Table 1. Cleavage efficiency of helix II stem-length variants.
 In Formula 1: (X)_n is 5'-AAAGGGGAA-3' (SEQ ID NO: 12); (X)_m is 5'-GGGAGCUGGC-3' (SEQ ID NO: 2); (X)_a is absent; (X)_b is 5'-GUGA-3'; 5'-...AXG...-3' is 5'-...AUG...-3'; and (X)_m and (X)_n are given below. See Figure 1 for Plot.

	% cleavage products									
	0	2	5	10	15	30	60	120 min		
control (X) _m 5'-GUCC-3' (X) _n 5'-GGAC-3'	3	15	41	60	72	88	92	94		
stem length variants										
1) 2 base pairs (X) _m 5'-GC-3' (X) _n 5'-GC-3'	4	24	60	68	70	73	76			
2) 3 base pairs (X) _m 5'-GAC-3' (X) _n 5'-GUC-3'	2	36	68	83	88	91	91	93		
3) 3 base pairs (X) _m 5'-GCC-3' (X) _n 5'-GGC-3'	3	26	44	68	69	74	79			
4) 4 base pairs (X) _m 5'-GAUC-3' (X) _n 5'-GAUC-3'	1	6	15	31	47	68	86			

Table 1. (Cont.)

	% cleavage products									
	0	2	5	10	15	30	60	120 min		
5) 5 base pairs (X) _{m'} 5'-GAUAC-3', (X) _m 5'-GUAUC-3'	0	3	8	15	23	44	68			
6) 6 base pairs (X) _{m'} 5'-GAUAC-3', (X) _m 5'-GAUAC-3'	0	3	8	20	32	48	75	89		
7) 7 base pairs (X) _{m'} 5'-GAUAC-3', (X) _m 5'-GUAUAC-3'	1	7	15	28	41	67	85	94		
8) 8 base pairs (X) _{m'} 5'-GAUUAUC-3', (X) _m 5'-GAUUAUC-3'	1	6	13	27	38	60	83	91		
9) 3 base pairs (X) _{m'} 5'-GCG-3', (X) _m 5'-CGC-3'	1	29	53	78	82	89	91			

Table 2. Cleavage efficiency of helix II conserved G-C base pair.
 In Formula 1: (X)_n is 5'-AAAGGGGAA-3' (SEQ ID NO: 12); (X)_n is 5'-GGGAGCUGGC-3' (SEQ ID NO: 2); (X)_a is absent; (X)_b is 5'-GUGA-3'; 5'-...AXG...-3' is 5'-...AUG...-3'; and (X)_m and (X)_n are given below. See Figure 2 for Plot.

	% cleavage products										
	0	2	5	10	15	30	60	120	min		
control											
(X) _m 5'-GUCC-3'	3	15	41	60	72	88	92	94			
(X) _m 5'-GGAC-3'											
variants of conserved G-C base pair											
1) A-G	3	21	39	58	72	85	90	92			
(X) _m 5'-GUCC-3'											
(X) _m 5'-GGAA-3'											
2) U-A	1	7	13	24	35	55	78	88			
(X) _m 5'-AUCC-3'											
(X) _m 5'-GGAU-3'											
3) C-A	0	4	8	16	22	41	58	65			
(X) _m 5'-AUCC-3'											
(X) _m 5'-GGAC-3'											
4) C-C	0	6	8	-	20	33	52	66			
(X) _m 5'-CUCC-3'											
(X) _m 5'-GGAC-3'											
5) A-U	0	3	7	12	16	30	51	75			
(X) _m 5'-UUCC-3'											
(X) _m 5'-GGAA-3'											

Table 2. (Cont.)

	% cleavage products										
	0	2	5	10	15	30	60	120 min			
6) U-C	0	1	3	5	7	13	23	43			
(X) _m 5'-CUCC-3'											
(X) _m 5'-GGAU-3'											
7) G-A	0	1	2	4	6	12	23	40			
(X) _m 5'-AUCC-3'											
(X) _m 5'-GGAG-3'											
8) G-U	0	0	2	3	4	9	15	30			
(X) _m 5'-UUCC-3'											
(X) _m 5'-GGAG-3'											
9) U-U	1	5	10	21	25	48	66	80			
(X) _m 5'-UUCC-3'											
(X) _m 5'-GGAU-3'											
10) G-C											
(X) _m 5'-CUCC-3'											
(X) _m 5'-GGAG-3'											

0

Table 3. Cleavage efficiency of helix II loop variants.
(In Formula 1: (X)_n is 5'-AAAGGGGAA-3' (SEQ ID NO: 12); (X)_n is 5'-GGGAGCUGGC-3' (SEQ ID NO: 2); (X)_n is absent; (X)_m is 5'-GUCC-3'; (X)_m is 5'-GGAC-3'; 5'-...AXG...-3' is 5'-...AUG...-3'; and (X)_b is given below. See Figures 3A-3C for Plots.

	% cleavage products									
	0	2	5	10	15	30	60	120 min		
control	3	15	41	60	72	88	92	94		
(X) _b 5'-GUGA-3'										
loop variants										
1) (X) _b 5'-GGGA-3'	4	36	68	80	85	90	91	93		
2) (X) _b 5'-GUGG-3'	2	27	53	71	79	87	90	92		
3) (X) _b 5'-GUGU-3'	2	23	41	57	69	83	90	91		
4) (X) _b 5'-UGGC-3'	2	19	38	61	67	81	83	86		
5) (X) _b 5'-GGCA-3'	0	21	38	53	64	76	82	85		
6) (X) _b 5'-GUGC-3'	2	19	33	53	61	76	81	88		
7) (X) _b 5'-GUUA-3'	0	14	26	47	61	74	81	83		
8) (X) _b 5'-GGGC-3'	0	12	25	39	48	65	67	74		
9) (X) _b 5'-UCAG-3'	0	9	19	36	53	80	90	92		
10) (X) _b 5'-GACU-3'	0	8	17	26	44	65	84	92		

Table 3. (Cont.)

		% cleavage products											
		0	2	5	10	15	30	60	120 min				
11)	(X) _b 5'-UUCG-3'	0	6	15	25	39	62	83	93				
12)	(X) _b 5'-CCGG-3'	0	6	15	25	39	62	83	93				
13)	(X) _b 5'-UUUU-3'							84					
14)	(X) _b 5'-CGGA-3'							84					
15)	(X) _b 5'-UUGGA-3'	5	28	54	79	85	92	94	96				
16)	(X) _b 5'-GUGUU-3'	5	29	49	67	75	88	90	93				
17)	(X) _b 5'-UUCGA-3'							85					
18)	(X) _b 5'-GGGGU-3'							84					
19)	(X) _b 5'-GCUUA-3'							81					
20)	(X) _b 5'-GAC-3'							66					
21)	(X) _b 5'-GAG-3'	3	17	33	50	64	76	82	85				
22)	(X) _b 5'-GUG-3'	3	16	31	51	59	75	84	88				
23)	(X) _b 5'-ACA-3'	0	14	29	45	54	68	75	83				
24)	(X) _b 5'-UGA-3'							80					
25)	(X) _b 5'-GCU-3'							77					
								68					

Table 4. Cleavage efficiency of helix II stem mismatches.
 In Formula 1: (X)_n is 5'-AAAGGGGAA-3' (SEQ ID NO: 12); (X)_n is 5'-GGAGGCTGGC-3' (SEQ ID NO: 2); (X)_a is absent; (X)_b is 5'-GUGA-3'; 5'-...AXG...-3' is 5'-...AUG...-3'; and (X)_m and (X)_n are given below.

	% cleavage products									
	0	2	5	10	15	30	60	120 min		
control	3	15	41	60	72	88	92	94		
(X) _m 5'-GUCC-3'										
(X) _m 5'-GGAC-3'										
stem mismatches										
1) (X) _m 5'-GUCG-3'	0	9	28	49	53	65	65	71		
(X) _m 5'-GGAC-3'										
2) (X) _m 5'-GUCC-3'	0	1	2	3	6	8	11	12		
(X) _m 5'-CAAC-3'										
3) (X) _m 5'-GUCC-3'									51	
(X) _m 5'-CGAC-3'										

Table 4. (Cont.)

		% cleavage products									
		0	2	5	10	15	30	60	120 min		
4)	(X) _m 5'-GUCU-3'								50		
	(X) _m 5'-GUAC-3'										
5)	(X) _m 5'-GUCC-3'								37		
	(X) _m 5'-AGAC-3'										
6)	(X) _m 5'-GUCC-3'								28		
	(X) _m 5'-UAAC-3'										
7)	(X) _m 5'-GUCC-3'								5		
	(X) _m 5'-UUAC-3'										
8)	(X) _m 5'-GUCC-3'								4		
	(X) _m 5'-ACAC-3'										

Table 5. Cleavage efficiency of minizyme variants.
 In Formula 1: (X)_n is 5'-AAAGGGGAA-3' (SEQ ID NO: 12); (X)_m is 5'-GGAGCUGGC-3' (SEQ ID NO: 2); (X)_a is absent; 5'-...AXG...-3' is 5'-...AUG...-3'; and 5'-(X)_m-(X)_b-(X)_m-3' is given below.

	% cleavage products										
	0	2	5	10	15	30	60	120	min		
control	0	2	5	10	15	30	60	120	min		
(X) _m 5'-GUCC-3'	3	15	41	60	72	88	92	94			
(X) _m 5'-GGAC-3'											
(X) _b 5'-GUGA-3'											
minizyme variants											
5'-(X) _m -(X) _b -(X) _m -3'											
1) 5'-UUUU-3'	0	0	0	2	6	12	26	48			
2) 5'-UUCG-3'							6				
3) 5'-GUGA-3'							5				
4) 5'-GGGC-3'							1				
5) 5'-UCAG-3'							0				
6) 5'-AAAC-3'							0				
7) 5'-ACGC-3'							0				

Table 6. Cleavage efficiency of helix I sequence variants.
In Formulas 1, 2, or 3: (X)_n is 5'-AAAGGGGAA-3' (SEQ ID NO: 12); (X)_n is absent; (X)_n is 5'-GUGA-3'; (X)_m is 5'-GUCC-3'; (X)_m 5'-GGAC-3'; 5'-...AXG...-3' is 5'-...AUG...-3'; and 3'-(X)_n-5' or 3'-N-N-(X)_n-5' is given below. See Figure 6 for Plot.

	% cleavage products							
	0	2	5	10	15	30	60	120 min
control	3	15	41	60	72	88	92	94
substrate helix I:								
5'-GCCAGUCCC-3', (SEQ ID NO: 1)								
ribozyme helix I: 3'-(X) _n -5' or 3'-N-N-(X) _n -5'								
5'-GGGAGCUGGC-3', (SEQ ID NO: 2)								
ribozyme helix I sequence variants								
1) 5'-GGUGAGCGGC-3' (SEQ ID NO: 3)	8	72	83	88	90	92	94	95
2) 5'-GGGAGCUGGU-3' (SEQ ID NO: 4)	0	0	3	3	5	6	9	17
3) 5'-GGAGCUCUAU-3' (SEQ ID NO: 5)	0	0	0	0	1	1	2	4
4) 5'-GGUCUGUGGC-3' (SEQ ID NO: 6)	-	25	51	66	77	83	88	
5) 5'-GGGGAGCGGC-3' (SEQ ID NO: 7)	-	22	37	64	73	84	90	92
6) 5'-GGACGGUGCC-3' (SEQ ID NO: 8)	-	2	4	8	11	20	40	61
7) 5'-GGGAGCUGGA-3' (SEQ ID NO: 9)								0
8) 5'-GGGAGCUGGG-3' (SEQ ID NO: 10)								0

Table 7. Cleavage efficiency of helix III sequence variants.
 In Formula 1: (X)_n is 5'-GGAGCUGGC-3' (SEQ ID NO: 2); (X)_n is absent; (X)_b is 5'-GUGA-3'; (X)_m is 5'-GUCC-3'; (X)_m 5'-GGAC-3'; 5'-...AXG...-3' is 5'-...AUG...-3', and (X)_n is given below. See Figure 7 for Plot.

	% cleavage products										
	0	2	5	10	15	30	60	120 min			
control	3	15	41	60	72	88	92	94			
substrate helix III											
5'-UCCCCUUU-3' (SEQ ID NO: 11)											
ribozyme helix III: (X) _n											
5'-AAAGGGGAA-3' (SEQ ID NO: 12)											
ribozyme helix III sequence variants: (X) _n											
1) 5'-AAAGUGGAAUG-3' (SEQ ID NO: 13)	1	15	33	53	63	84	86	90			
2) 5'-AAGGGGGAA-3' (SEQ ID NO: 14)	1	14	28	50	62	81	87	89			
3) 5'-AAUUGUGGA-3' (SEQ ID NO: 15)	0	5	11	21	25	42	60	70			
4) 5'-AAUGGGUGAA-3' (SEQ ID NO: 16)	0	4	7	11	19	36	62	81			
5) 5'-AAUAGGGGAA-3' (SEQ ID NO: 17)	0	2	5	8	12	20	41	64			
6) 5'-ACAGGGGGAA-3' (SEQ ID NO: 18)	0	2	4	6	9	17	28	49			
7) 5'-AAGCGGAGUG-3' (SEQ ID NO: 19)	0	0	1	2	2	3	6	9			
8) 5'-AUAGGGGGAA-3' (SEQ ID NO: 20)	0	0	1	1	2	3	7	12			
9) 5'-AAGGGGAAU-3' (SEQ ID NO: 21)	0	0	0	0	1	2	3	5			
10) 5'-AAGAUGGGG-3' (SEQ ID NO: 22)	0	0	0	0	0	1	1	1			
11) 5'-CAAGGGGAA-3' (SEQ ID NO: 23)								0			

Table 8. Effect of oligonucleotides, complementary to the ribozyme, on cleavage efficiency. (In Formula 1; (X)_n is 5'-AAGGGGAA-3' (SEQ ID NO: 12), (X)_n is 5'-GGAGCUGGC-3' (SEQ ID NO: 2), (X)₆ is absent, (X)₆ is 5'-GUGA-3', (X)_n is 5'-GUCC-3', (X)_m is 5'-GGAC-3', and 5'-...AXG...-3' is 5'-...AUG...-3'). See Figure 8 for Plot. cleavage as % of control (37°C/10 min)

Complementary Oligonucleotide

1) 5'-dCCCCCTTT-3'	158
2) 5'-dTTTTCGTC-3'	190
3) 5'-dCTCATCAG-3'	177
4) 5'-dGCCAGCTC-3'	203
5) 5'-dTTCCTT-3' (SEQ ID NO: 24)	152
6) 5'-dCCCTTTTCG-3' (SEQ ID NO: 25)	197
7) 5'-dATCAGGCCAG-3' (SEQ ID NO: 26)	194
8) 5'-dGCCAGCTCCC-3' (SEQ ID NO: 27)	161
9) 5'-dNNNNNNNT-3' (SEQ ID NO: 31)	147
(N = dA, dC, dG or dT)	
10) 5'-dAGCCAGCTCCC-3' (SEQ ID NO: 28)	122
11) 5'-dTTCCTT-3' (SEQ ID NO: 29)	71
12) 5'-CUUUUCG-3'	157
13) 5'-GCCAGC-3'	170
14) 5'-GGCCAGCU-3'	130
15) 5'-AGGCCAGCUC-3' (SEQ. ID NO. 30)	130

Table 8. (Cont.)

	cleavage as % of control (37°C/10 min)
16) 5'-NNNNNNG-3'	109
-(N = A, C, G or U)	
17) 0.1mM dATP, dCTP, dGTP, dTTP	92
18) 5mM spermine	32
19) 10mM spermine	30
20) 50mM spermine	20
21) 5% polyethylene glycol 8000	108
22) 10% polyethylene glycol 8000	80
23) 15% polyethylene glycol 8000	60
24) 0.5M urea	85
25) 1M urea	80
26) 3M urea	39
27) 10% formamide	71
28) 20% formamide	51

References

- Altman, S. (1987) Adv. Enzymol. 62:1.
- Baer, M.F., et al. (1990) Methods Enzymol. 181:569.
- Been, M.D., and Cech, T.R. (1988) Science 239:1412
- 5 Biou, V. et al. (1994) Science 263:1404-1410.
- Boehm, S., (1987) FEBS Letters. 220: 283-287.
- 10 Bruening, G., (1989) Methods in Enzymology. 180:546-558.
- Bruening, G. (1990) Seminars in Virol. 1:127.
- Bryant, J. (1992) Tibtech 10:342-343.
- 15 Buzayán et al., (1986) Proc. Natl. Acad. Sci. USA 83:8859-8862.
- Cameron et al., (1989) Proc. Natl. Acad. Sci. USA, 86:9139-9143.
- 20 Cantor, G.H., et al. (1993) Proc. Natl. Acad. Sci. U.S.A. 90:10932-10936.
- Carruthers et al. (1987) Methods in Enzymology. 154: 287-313).
- Castanotto, D., et al. (1992) Critical Rev. in Eukaryotic gene Exp. 2:331-337.
- 25 Cech, T. R., (1987) Science. 236: 1532-1539.
- Cech, T.R., et al. (1981) Cell 27:487.
- 30 Cheong, C. et al. (1990) Nature 346:680-682.
- Chuat, J. et al., (1989) Biochemical and Biophysical Research Communications, 162: 1025-1029.
- 35 Cotten, M., (1990) Tibtech 8: 174-178.
- Cotten, M. et al., (1989) The EMBO Journal, 8: 3861-3866.
- Dahm, S. C., et al., (1990) Biochimie 72:819-823.
- 40 Dropulic, B., et al. (1992) J. Virol. 66:1432-1441.
- Eckner, R. et al., (1991) The EMBO Journal, 10: 3513-3522.
- 45 Epstein, L.M., and Gall, J.G., (1987) Cell 48:535.

- Epstein, L.M., and Gall, J.G., (1989) Cold Spring Harbour Symp. Quant. Biol. 52:261.
- 5 Evans, G.J., et al. (1992) Biochem. Soc. Trans. 20:344S.
- Foehler et al. (1986) Nucleic Acids Research 14: 5399-407.
- Forster et al., (1987) Cell, 50:9-16.
- 10 Forster, A. C. et al., (1988) Nature, 334:265-267.
- Forster, A.C. and Symons, R. H., (1987) Cell, 49: 211-220.
- Freier, S.M. et al. (1986) Proc. Nat. Acad. Sci. USA 83:9373-9377.
- 15 Friedman, T., (1989) Science 244: 1275-1280.
- Gallie, D.R., et al. (1991) Mol. Gen. Genet. 228:258-264.
- 20 Goodchild, J., et al., (1991) Archives to Biochem & Biophys. 284:386-391.
- Green, R., and Szostak, J.W. (1992) Science 258:1910.
- 25 Goodchild, J. et al., (1990) Poster No. 12 at Conf in San Diego.
- Haseloff, J. and W. L. Gerlach, (1988) Nature, 334:585-591.
- Haseloff et al., (1989), Gene, 82:43-52.
- 30 Herschlag, D., and Cech, T.R. (1990) Nature 344:405.
- Hertel, K.J. et al. (1992) Nucleic Acids Res. 20:3252.
- 35 Heus, H.A. et al. (1991) Science 253:191-194.
- Hogan, B. et al., (1986) Manipulating the Mouse Embryo, A Laboratory Manual, Cold Spring Harbor.
- 40 Hogan, B. et al., (1989) Science, 244: 1275.
- Homann, M., et al. (1993) Nucleic Acid Res. 28:2809-2814.
- Howard, E.A., et al. (1987) Planta 170:535-540.
- 45 Huillier, A. et al., Ribozyme Mediated Suppression of Lactalbumin Expressed in C1271 Cells by T7/Vaccinia Virus, (Abstract from conference proceedings).

- Hutchins, C. J. et al., (1986) Nucleic Acids Res. 14:3627-3635.
- Jefferies, A. C., et al., (1989) Nucleic Acids Res. 17:1371-1377.
- 5 Joyce, G.F. (1992) Sci. Am. 267:90.
- Karnail, U. and Wasternack, C. (1992) J. Biochem. 24:493-497.
- Kikuchi, Y. et al., (1991) Nucleic Acids Res. 19:6751-6755.
- 10 Kinsey, P.T. and Sandmeyer, S.S. (1991) Nucleic Acid Res. 19:1317-1324.
- Koizumi et al., (1988) FEBS Letters, 228:228-230.
- 15 Koizumi et al., (1989) Nucleic Acids Res. 17:7059-7071.
- Kruger, K., et al. (1982) Cell 31:147.
- Kunkel, T.A., et al. (1987) Methods in Enzymology 154:367-382.
- 20 Lamb, J. W. & Hay, R. T., (1990) J. Gen. Virol., 71:2257-2264.
- L'Huillier, P.J.L., et al. (1992) The EMBO J. 11:4411-4418.
- 25 Llewellyn et al., (1987), J. Mol. Biol., 195: 115-123.
- Long, D.M. et al. (1994) Proc. Natl. Acad. Sci. U.S.A. 91:6977-6981.
- Maliga, P. (1993) Tibtech 11:101-106.
- 30 Mazzolini, L., et al. (1992) Plant Molecular Biology 20:715-731.
- McCall, M.J. et al. (1992) Proc. Natl. Acad. Sci. U.S.A. 89:5710-5714.
- 35 McClain et al., (1987) Science 238:527-530.
- Miller, W. A. et al., (1991) Virology, 183:711-720.
- Nichols, M., et al. (1988) Proc. Natl. Acad. Sci. U.S.A. 85:1379.
- 40 Noller, H.F., et al. (1992) Science 256:1416.
- Pace, N.R., and Smith, D., (1990) J. Biol. Chem. 265:3587.
- 45 Pan, T., and Uhlenbeck, O.C. (1992) Nature 358:560.
- Perriman et al., (1992), Gene, 113:157.

- Perriman, R., et al. (1993) Antisense Res. & Dev. 3:253-263.
- Perreault et al., (1990), Nature, 344:565-567.
- 5 Perreault, J. P., et al., (1991) Biochem. 30:4020-4025.
- Piccirilli, J.A., et al. (1992) Science 256:1420.
- Pley, H.W., et al. (1994 A) Nature 372:68-74.
- 10 Pley, H.W., et al. (1994 B) Nature 372:111-113.
- Prody et al., (1986), Science 231: 1577-1580.
- 15 Pyle, A.M. (1993) Science 261:709-714.
- Robertson, D.L., and Joyce, G.F. (1990) Nature 344:467.
- Ruffner, D. E. et al., (1990) Biochemistry, 29: 10695-10702.
- 20 Ruffner, D. E. et al., (1989) Gene, 82:31-41.
- Sambrook, J. et al., (1989), Molecular Cloning, A Laboratory Manual, 2nd Edition, Cold Spring Harbor Press.
- 25 Sampson, et al., (1987) Cold Spring Harbor Sym Quant. Biol. 52:267-275.
- Sangar, W. (1984), Principles of Nucleic Acid Structure, Springer-Verlag, N. Y.
- 30 Sarver, N. et al., (1990) Science, 247:1222-1224.
- Saville, B.J. and Collins, R.A. (1990) Cell 61:685.
- Saxena, S. et al., (1990) J. Biol. Chem., 265:17106-17109.
- 35 Scanlon, K. et al., (1991) Proc. Natl. Acad. Sci. USA, 88:10591-10595.
- Sheldon, C. C. & Symons, R. H., (1989) Nucleic Acids Res. 17:5679-5686.
- 40 Sheldon, C. C. & Symons, R. H., (1989) Nucleic Acids Res. 17:5665-5678.
- Shimamoto, K., Terada, R., Izawa, T., and Fujimoto, H. (1989) Nature 338:274-276.
- 45 Sioud, M. and Drlica, K. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:7303-7307.

- Sioud, M., et al. (1992) J. Mol. Biol. 223:831-835.
- Strobel, S.A. et al., (1991) Nature 350: 172-174 and references therein.
- 5 Sober, H. (1970), CRC Handbook of Biochemistry, Second edition.
- Sprout et al. (Oligonucleotide Synthesis--A Practical Approach, IRL Press, Oxford (1984) M. J. Gait--Editor, pp. 83-115).
- 10 Sullenger, B.A. and Cech, T.R. (1993) Science 262:1566-1569.
- Stange, N. and Beier, D. (1986) Nucleic Acid Res. 14:8961.
- Steinecke, P., et al. (1992) The EMBO J. 11:1525-1530.
- 15 Symons, R. H., (1989) TIBS, 14:445-450.
- Symons, R.H., (1990) Seminars in Virol. 1:117.
- 20 Tabler, M. & Tsagris, M., (1991) Gene, 108:175-183.
- Taylor, J., (1990) Seminars in Virol. 1:135.
- Tuschl, T. et al. (1993) Proc. Nat. Acad. Sci. USA 90:6991-6994.
- 25 Uhlenbeck et al., (1987) Nature, 328:596-600.
- Uhlmann et al., (1990), Chem. Revs., 90:544-584.
- 30 Waugh, D.S., et al. (1989) Science 244:1569.
- Woese, C.R. et al. (1990) Proc. Nat. Acad. Sci. USA 90:8467-8471.
- Yang, et al., (1990) Biochemistry 29:11156-11160.
- 35 Zaug, A.J. et al, (1984), Science, 224:574-578
- Zaug, A. J. & Cech, T.R., (1986a), Science, 231:473-474.
- 40 Zaug, A. J. et al., (1986b) Nature, 324:429-433.
- Zaug, A.J., and Cech, T.R. (1986) Biochemistry 25:4478.
- Zhao, J.J. and Pick, L. (1993) Nature 365:448-451.

- 71 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Keese, Paul
Stapper, Marianne
Perriman, Rhonda
- (ii) TITLE OF INVENTION: Ribozymes With Optimized Hybridizing Arms,
Stems And Loops, tRNA Embedded Ribozymes
and Compositions Thereof
- (iii) NUMBER OF SEQUENCES: 31
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Cooper & Dunham
 - (B) STREET: 30 Rockefeller Plaza
 - (C) CITY: New York
 - (D) STATE: New York
 - (E) COUNTRY: U.S.A.
 - (F) ZIP: 10112
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: 3.5 INCH, 1.44Mb
 - (B) COMPUTER: IBM PC
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.24
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: June 24, 1994
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: White, John P.
 - (B) REGISTRATION NUMBER: 28,678
 - (C) REFERENCE/DOCKET NUMBER: 45284
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 212 977 9550
 - (B) TELEFAX: 212 977 9809

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCCAGCUC

10

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

SUBSTITUTE SHEET (RULE 26)

- 72 -

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GGGAGCUGGC

10

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGUGAGCGGC

10

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGGAGCUGGU

10

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GGAGCUCUAU

10

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GGUCUGUGGC

10

SUBSTITUTE SHEET (RULE 26)

- 73 -

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GGGGAUGCGC

10

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GGACGGUGCC

10

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GGGAGCUGGA

10

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGGAGCUGGG

10

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:

SUBSTITUTE SHEET (RULE 26)

- 74 -

- (A) LENGTH: 10 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

UUCCCCUUU

10

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AAAGGGGGAA

10

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

AAAGUGGAUG

10

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AAGGGGGGAA

10

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

SUBSTITUTE SHEET (RULE 26)

- 75 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AAUUGUGGGA

10

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

AAUGGGUGAA

10

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

AAUAGGGGAA

10

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

ACAGGGGGAA

10

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

SUBSTITUTE SHEET (RULE 20)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

AAGCGGAGUG

10

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

AUAGGGGGAA

10

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

AAGGGGGUAU

10

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

AAGAUGGGGG

10

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

SUBSTITUTE SHEET (RULE 26)

CAAGGGGGAA

10

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

TTCCCCCTTT

10

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CCCTTTTTCG

10

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

ATCAGGCCAG

10

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GCCAGCTCCC

10

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other Nucleic Acid


SUBSTITUTE SHEET (RULE 26)

- 78 -

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:
AGGCCAGCTC CC 12
- (2) INFORMATION FOR SEQ ID NO:29:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 13 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other Nucleic Acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:
TTCCCCCTTT TTG 13
- (2) INFORMATION FOR SEQ ID NO:30:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other Nucleic Acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:
AGGCCAGCUC 10
- (2) INFORMATION FOR SEQ ID NO:31:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Other Nucleic Acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:
NNNNNNNNNT 10

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A compound having the formula:



wherein each X represents a nucleotide which may be the same or different and may be substituted or modified in its sugar, base or phosphate;

wherein each of A, C, U, and G represents a ribonucleotide which may be substituted or modified in its sugar, base or phosphate;

wherein each of $(X)_n$ and $(X)_{n'}$ represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved, and does not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, such RNA target sequence not being present within the compound;

wherein each of n and n' represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that the sum of $n + n'$ is greater than 14;

wherein each of $(X)_m$ and $(X)_{m'}$ represents an oligonucleotide having the sequences selected from the group consisting of (a) 3'-CUG-5' and 5'-GAC-3', (b) 3'-CGG-5' and 5'-GCC-3' and (c) 3'-CGC-5' and 5'-GCG-3', respectively, and * represents base pairing between each nucleotide of $(X)_m$ and $(X)_{m'}$ located on



either side thereof;

wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof;


wherein a represents an integer which defines a number of nucleotides with the proviso that a may be 0 or 1 and if 0, the A located 5' of $(X)_a$ is bonded to the X located 3' of $(X)_a$; and

wherein $(X)_b$ represents an oligonucleotide with the proviso that b represents an integer which is greater than or equal to 4 if $(X)_b$ is present.

2. The compound of claim 1, wherein the oligonucleotide $3'-(X)_n$ is $3'-(X)_{n-1}-A-$.
3. The compound of claim 1, wherein the oligonucleotide $3'-(X)_n$ is $3'-(X)_{n-2}-C-A-$.
4. The compound of claim 1, wherein $(X)_a$ is absent.
5. The compound of claim 1, wherein the integer b of $(X)_b$ is equal to 4.
6. The compound of claim 1, wherein the oligonucleotide $(X)_b$ has the sequence $5'-G-U-G-A-3'$.
7. The compound of claim 1, wherein $(X)_b$ represents an oligonucleotide having one of the following sequences: $3'-AAGGC-5'$, $3'-AGGUU-5'$, $3'-AGGG-5'$, or $3'-GGUG-5'$.
8. A composition which comprises the compound of claim 1 and an acceptable carrier.



9. A compound having the formula:



wherein each X represents a nucleotide which may be the same or different and may be substituted or modified in its sugar, base or phosphate;

wherein each of A, C, U, and G represents a ribonucleotide which may be substituted or modified in its sugar, base or phosphate;

wherein each of $(X)_n$ and $(X)_{n'}$ represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved, and does not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, such RNA target sequence not being present within the compound;

wherein each of n and n' represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that the sum of $n + n'$ is greater than 14;

wherein * represents base pairing between each nucleotide of $(X)_m$ and $(X)_{m'}$ located on either side thereof;

wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof;




wherein a represents an integer which defines a number of nucleotides with the proviso that a may be 0 or 1 and if 0, the A located 5' of $(X)_a$ is bonded to the X located 3' of $(X)_a$;

wherein each m and m' represents an integer which is from two to eight inclusive; and

wherein $(X)_b$ represents an oligonucleotide having a sequence selected from the group consisting of 3'-AAGGC-5', 3'-AGGUU-5', 3'-AGGG-5', and 3'-GGUG-5'.

10. The compound of claim 9, wherein the oligonucleotide 3'-(X)_n- is 3'-(X)_{n-1}-A-.
11. The compound of claim 9, wherein the oligonucleotide 3'-(X)_n- is 3'-(X)_{n-2}-C-A-.
12. The compound of claim 9, wherein $(X)_a$ is absent.
13. The compound of claim 9, wherein each of m and m' are 4.
14. A composition which comprises the compound of claim 9 and an acceptable carrier.
15. A compound having the formula:



wherein each X and each N represents a nucleotide which may be the same or different and may be substituted or modified



in its sugar, base or phosphate;

wherein each of A, C, U, and G represents a ribonucleotide which may be substituted or modified in its sugar, base or phosphate;

wherein each of $(X)_n$ and $(X)_{n'}$ represents an oligonucleotide having a predetermined sequence which hybridizes with an RNA target sequence to be cleaved, and each of $(X)_n$ and $N-N-(N)_d-(X)_{n'}$ does not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, such RNA target sequence not being present within the compound;

wherein each N represents a nucleotide which hybridizes with the RNA target sequence;


wherein d is either zero or an integer between 1 and 5, and if d is zero, then $(N)_d$ is absent;

wherein one or more nucleotides at the 3' end of $(X)_{n'}$ do not hybridize to the RNA target sequence and at least one nucleotide directly 5' of such nucleotides hybridizes to the RNA target sequence;

wherein each of n and n' represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that n' is at least 2 and the sum of n + n' is greater than 12;

wherein * represents base pairing between each nucleotide of $(X)_m$ and $(X)_{m'}$ located on either side thereof;

wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on



either side thereof;

wherein a represents an integer which defines a number of nucleotides with the proviso that a may be 0 or 1 and if 0, the A located 5' of $(X)_a$ is bonded to the X located 3' of $(X)_a$;

wherein each m and m' represents an integer which is greater than or equal to 4; and

wherein $(X)_b$ represents an oligonucleotide which may be present or absent with the proviso that b represents an integer which is greater than or equal to 4 if $(X)_b$ is present.

16
17
18
19
20

16. The compound of claim 15, wherein d is 0.
17. The compound of claim 15, wherein d is 2 to 5.
18. The compound of claim 15, wherein the oligonucleotide 3'-(X)_n- is 3'-(X)_{n-1}-A-.
19. The compound of claim 15, wherein the oligonucleotide 3'-(X)_n- is 3'-(X)_{n-2}-C-A-.
20. The compound of claim 15, wherein $(X)_a$ is absent.
21. The compound of claim 15, wherein $(X)_b$ is present.
22. The compound of claim 15, wherein the integer b of $(X)_b$ is equal to 4.
23. The compound of claim 15, wherein each of m and m' are 4.
24. The compound of claim 15, wherein the oligonucleotide $(X)_b$ has




the sequence 5'-G-U-G-A-3'.

25. A composition which comprises a compound of claim 15 in association with a carrier.
26. An oligonucleotide transfer vector containing a nucleotide sequence which on transcription gives rise to any of the compounds recited in claim 15.
27. The transfer vector of claim 26, wherein the transfer vector is selected from a group consisting of a bacterial plasmid, a bacteriophage DNA, a cosmid, and an eukaryotic viral DNA.
28. A host cell transformed by the transfer vector of claim 26.
29. The host cell of claim 28, wherein the host cell is a prokaryotic host cell or an eukaryotic host cell.
30. The prokaryotic host cell of claim 29, wherein the prokaryotic host cell is an E. coli host cell.
31. The eukaryotic host cell of claim 29, wherein the eukaryotic host cell is a monkey COS host cell, a Chinese hamster ovary host cell, a mammalian host cell or a plant host cell.
32. A composition which comprises the compound of claim 15 and an acceptable carrier.
33. A composition comprising a first synthetic non-naturally occurring oligonucleotide compound which compound comprises nucleotides whose sequence defines a conserved catalytic region and nucleotides whose sequence is capable of hybridizing with a predetermined target sequence and a second synthetic non-naturally occurring oligonucleotide compound which does not contain the predetermined target sequence and is complementary



to at least a portion of the first compound.

34. The composition of claim 33, wherein the first synthetic non-naturally occurring oligonucleotide compound has the structure:



wherein each X represents a nucleotide which is the same or different and may be modified or substituted in its sugar, phosphate or base;

wherein each of A, C, U, and G represents a ribonucleotide which may be modified or substituted in its sugar, phosphate or base;

wherein 3'-AAG...AGUCX-5' defines a conserved catalytic region;

wherein each of $(X)_nA$ and $(X)_n$ represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved, and does not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, such RNA target sequence not being present within the compound;

wherein each * represents base pairing between the nucleotides located on either side thereof;




wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof;

wherein a represents an integer which defines a number of nucleotides with the proviso that a may be 0 or 1 and if 0, the A located 5' of $(X)_a$ is bonded to the X located 3' of $(X)_a$;

wherein each of m and m' represents an integer which is greater than or equal to 1; and

wherein $(X)_b$ represents an oligonucleotide and b represents an integer which is greater than or equal to 4.

35. The composition of claim 33, wherein the first synthetic non-naturally occurring oligonucleotide compound of has the structure:



wherein each X is the same or different and represents a ribonucleotide or a deoxyribonucleotide which may be modified or substituted in its sugar, phosphate or base;

wherein each of A, C, U, and G represents a ribonucleotide which may be modified or substituted in its sugar, phosphate or base;

wherein 3'—AAG...AGUCX—5' defines a conserved catalytic region;




wherein each of $(X)_nA$ and $(X)_n$ represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved, and does not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, such RNA target sequence not being present within the compound;

wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof;

wherein m represents an integer from 2 to 20; and

wherein none of the nucleotides $(X)_m$ are Watson-Crick base paired to any other nucleotide within the compound.

36. The composition of claim 33, wherein the first synthetic non-naturally occurring oligonucleotide compound has the structure:



wherein each X is the same or different and represents a ribonucleotide or a deoxyribonucleotide which may be modified or substituted in its sugar, phosphate or base;

wherein each of A, C, U, and G represents a ribonucleotide which may be modified or substituted in its sugar, phosphate or base;

wherein 3' $(X)_{P4} \dots (X)_{P1}$ --- 5' defines a conserved catalytic region;



wherein each of $(X)_{P_4}$ and $(X)_{F_3}$ represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved;

wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof;

wherein F_3 represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that F_3 is greater than or equal to 3;

wherein F_4 represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that F_4 is from 3 to 5;

wherein each of $(X)_{P_1}$ and $(X)_{P_4}$ represents an oligonucleotide having a predetermined sequence such that $(X)_{P_4}$ base-pairs with 3-6 bases of $(X)_{P_1}$;

wherein P_1 represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that P_1 is from 3 to 6 and the sum of P_1 and F_4 equals 9;

wherein each of $(X)_{P_2}$ and $(X)_{P_3}$ represents an oligonucleotide having a predetermined sequence such that $(X)_{P_2}$ base-pairs with at least 3 bases of $(X)_{P_3}$;

wherein each * represents base pairing between the nucleotides located on either side thereof;


wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof; and



wherein $(X)_{L2}$ represents an oligonucleotide which may be present or absent with the proviso that $L2$ represents an integer which is greater than or equal to 3 if $(X)_{L2}$ is present.

37. The composition of any one of claims 33-36, wherein the composition is in association with an acceptable carrier.

38. A compound having the formula:




wherein each $(X)_t$ and $(X)_{t'}$ represents an oligonucleotide having a sequence which corresponds to a portion of a plant tRNA molecule;

wherein each of t and t' represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that the sum of $t + t'$ is greater than 50; and

wherein Y is a synthetic non-naturally occurring oligonucleotide moiety which comprises nucleotides whose sequence defines a conserved catalytic region and nucleotides whose sequence is capable of hybridizing with a predetermined target sequence to be cleaved.



39. The compound of claim 38, wherein the synthetic non-naturally occurring oligonucleotide moiety has the structure:



wherein each X represents a nucleotide which is the same or different and may be modified or substituted in its sugar, phosphate or base;

wherein each of A, C, U, and G represents a ribonucleotide which may be modified or substituted in its sugar, phosphate or base;

wherein 3'—AAG....AGUCX—5' defines a conserved catalytic region;

wherein each of (X)nA and (X)n' represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved, and does not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, such RNA target sequence not being present within the compound;

wherein each * represents base pairing between the nucleotides located on either side thereof;

wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side




thereof;

wherein a represents an integer which defines a number of nucleotides with the proviso that a may be 0 or 1 and if 0, the A located 5' of $(X)_a$ is bonded to the X located 3' of $(X)_a$;

wherein each of m and m' represents an integer which is greater than or equal to 1; and

wherein $(X)_b$ represents an oligonucleotide and b represents an integer which is greater than or equal to 4.

40. The compound of claim 38, wherein the synthetic non-naturally occurring oligonucleotide moiety has the structure:



wherein each X is the same or different and represents a ribonucleotide or a deoxyribonucleotide which may be modified or substituted in its sugar, phosphate or base;

wherein each of A, C, U, and G represents a ribonucleotide which may be unmodified or modified or substituted in its sugar, phosphate or base;

wherein 3'—AAG...AGUCX—5' defines a conserved catalytic region;

wherein each of $(X)_nA$ and $(X)_{n'}$ represents an oligonucleotide




having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved, and does not naturally occur covalently bound to the sequences 3'-A-A-G-5' and 5'-C-U-G-A-3', respectively, such RNA target sequence not being present within the compound;

wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof;

wherein m represents an integer from 2 to 20; and

wherein none of the nucleotides (X)_m are Watson-Crick base paired to any other nucleotide within the compound.

41. The compound of claim 38, wherein the synthetic non-naturally occurring oligonucleotide moiety has the structure:



wherein each X is the same or different and represents a ribonucleotide or a deoxyribonucleotide which may be modified or substituted in its sugar, phosphate or base;

wherein each of A, C, U, and G represents a ribonucleotide which may be unmodified or modified or substituted in its sugar, phosphate or base;

wherein 3' (X)_{P4}....(X)_{P1}--- 5' defines a conserved catalytic region;



wherein each of $(X)_{F4}$ and $(X)_{F3}$ represents an oligonucleotide having a predetermined sequence which is capable of hybridizing with an RNA target sequence to be cleaved;

wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof;

wherein F3 represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that F3 is greater than or equal to 3;

wherein F4 represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that F4 is from 3 to 5;

wherein each of $(X)_{P1}$ and $(X)_{P4}$ represents an oligonucleotide having a predetermined sequence such that $(X)_{P4}$ base-pairs with 3-6 bases of $(X)_{P1}$;

wherein P1 represents an integer which defines the number of nucleotides in the oligonucleotide with the proviso that P1 is from 3 to 6 and the sum of P1 and F4 equals 9;

wherein each of $(X)_{P2}$ and $(X)_{P3}$ represents an oligonucleotide having a predetermined sequence such that $(X)_{P2}$ base-pairs with at least 3 bases of $(X)_{P3}$;

wherein each * represents base pairing between the nucleotides located on either side thereof;

wherein each solid line represents a chemical linkage providing covalent bonds between the nucleotides located on either side thereof; and

1
2
3
4
5
6
7
8
9
10



wherein $(X)_{L2}$ represents an oligonucleotide which may be present or absent with the proviso that $L2$ represents an integer which is greater than or equal to 3 if $(X)_{L2}$ is present.

42. The compound of any one of claims 38-41, wherein each of $(X)_t$ and $(X)_c$ correspond to a portion of a tyrosine tRNA molecule of Nicotiana rustica.
43. A oligonucleotide transfer vector containing a nucleotide sequence which on transcription gives rise to the compound of any one of claims 38-41.
44. The transfer vector of claim 43, wherein the transfer vector is a bacterial plasmid, a bacteriophage DNA, a cosmid, or plant viral DNA.
45. The transfer vector of claim 44, wherein the transfer vector contains the promoter sequences for RNA polymerase II or RNA polymerase III.
46. A host cell transformed by the transfer vector of any one of claims 43-44.
47. The host cell of claim 46, wherein the host cell is a prokaryotic host cell or an eukaryotic host cell.
48. The prokaryotic host cell of claim 47, wherein the prokaryotic host cell is an E. coli host cell.
49. The eukaryotic host cell of claim 47, wherein the eukaryotic host cell is a monkey COS host cell, a Chinese hamster ovary host cell, a mammalian host cell, a plant host cell or a plant protoplast host cell.
50. A packaged oligonucleotide transfer vector containing a



nucleotide sequence which on transcription gives rise to the compound of any one of claims 38-41.

51. The packaged oligonucleotide transfer vector of claim 50, wherein the packaged oligonucleotide transfer vector is selected from the group consisting of a plant DNA virus, a geminivirus and an infective phage particle.
52. The packaged oligonucleotide transfer vector of claim 51, wherein the packaged oligonucleotide transfer vector contains the promoter sequences for RNA polymerase II or RNA polymerase III.

Dated this 9th day of December, 1998

GENE SHEARS PTY.LTD.

By its Patent Attorneys
Davies Collison Cave



FIGURE 1






FIGURE 2




3/23

FIGURE 3A




4/23

FIGURE 3B



5/23

FIGURE 3C



6/23

FIGURE 4A





FIGURE 4B




SUBSTITUTE SHEET (RULE 26)

7/23

FIGURE 5


stem II



SUBSTITUTE SHEET (RULE 26)


8/23

FIGURE 6




9/23

FIGURE 7




10/23

FIGURE 8



11/23

FIGURE 9A



12/23

FIGURE 9B






FIGURE 9C




14/23



15/23


FIGURE 10A

CAT	CAT + Rz12	CAT + tRNARz12
-----	---------------	-------------------




16/23

FIGURE 10B




17/23

FIGURE 11A




18/23

FIGURE 11B




19/23

FIGURE 12A




20/23

FIGURE 12B




21/23

FIGURE 13A



22/23

FIGURE 13B



23/23

FIGURE 14A

CAT




FIGURE 14B

CATMUT2

