




- (51) **International Patent Classification:**
A61K 47/68 (2017.01) *A61P 35/00* (2006.01)
A61K 47/65 (2017.01)
- (21) **International Application Number:**
PCT/EP2020/057697
- (22) **International Filing Date:**
19 March 2020 (19.03.2020)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
19163810.5 19 March 2019 (19.03.2019) EP
- (71) **Applicant:** PAUL SCHERRER INSTITUT [CH/CH];
OIPA/10, 5232 Villigen PSI (CH).
- (72) **Inventors:** SCHIBLI, Roger; Schartenrainstrasse 42, 5400
Baden (CH). BEHE, Roger; Rauracherweg 3, 4460 Gel-
terkinden (CH). SPYCHER, Philipp; Rotachstrasse 44,
8003 Zürich (CH). FREI, Julia; Seidenstrasse 6, 5200
Brugg AG (CH). WEHRMÜLLER, Jöri; Dialogweg 6,
8050 Zürich (CH).
- (74) **Agent:** VOSSIUS & PARTNER (NO 31); Siebertstraße 3,
81675 München (DE).
- (81) **Designated States** (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

(54) **Title:** TRANSGLUTAMINASE CONJUGATION METHOD WITH A GLYCINE BASED LINKER

Fig. 1



(57) **Abstract:** The present invention relates to a method for generating an antibody-payload conjugate by means of a microbial transglutaminase (MTG). The method comprises a step of conjugating a linker comprising or having the peptide structure (shown in N->C direction) Gly-(Aax)_m-B-(Aax)_n via the N-terminal primary amine of the N-terminal glycine (Gly) residue to a glutamine (Gln) residue comprised in the heavy or light chain of an antibody.

Published:

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*

TRANSGLUTAMINASE CONJUGATION METHOD WITH A GLYCINE BASED LINKER

FIELD OF THE INVENTION

The present invention relates to methods for generating an antibody-payload conjugate by means of a microbial transglutaminase. The invention further provides linkers, linker-payload constructs and/or antibody-payload constructs.

BACKGROUND OF THE INVENTION

Attaching highly potent payloads to antibodies finds increasing interest for the targeted treatment of cancer or inflammatory diseases. The constructs this produces are called antibody-payload conjugates, or antibody-drug conjugates (ADC).

Currently, seven ADCs have gained FDA-approval (Adcetris, Kadcyła, Besponsa, Mylotarg, Polivy, Padcev, Enhertu), all of which have their payload chemically attached to the antibody in a non-site specific manner. Hence, the resulting product is highly heterogeneous, both with respect to the stoichiometric relationship between antibody and payload (payload-antibody ratio, or drug-to-antibody ratio, DAR), as well concerning the conjugation sites on the antibody. Each of the resulting species, although in the same drug product, may have distinct properties that could potentially lead to a wide range of different *in vivo* pharmacokinetic properties and activities.

In a previous *in vivo* study (Lhospice et al., 2015), it was shown that a site-specific drug attachment led to a significant higher tumor uptake (~2x) and a decreased uptake in non-targeted tissues compared to the FDA-approved ADC, also the maximal tolerated dose was at least 3x higher. These data suggest that stoichiometrically well-defined ADCs display

improved pharmacokinetics and better therapeutic indexes compared to chemically modified ADCs.

As a site-specific technology, enzymatic conjugation has gained great interest since these conjugation reactions are typically fast and can be performed under physiological conditions. Among the available enzymes, microbial transglutaminase (MTG) from the species *Streptomyces mobaraensis* has gained increasing interest as an attractive alternative to conventional chemical protein conjugation of functional moieties including antibodies. The MTG catalyzes under physiological conditions a transamidation reaction between a 'reactive' glutamine of a protein or peptide and a 'reactive' lysine residue of a protein or peptide, whereas the latter can also be a simple, low molecular weight primary amine such as a 5-aminopentyl group (Jeger et al., 2010, Strop et al., 2014).

The bond formed is an isopeptide bond which is an amide bond that does not form part of the peptide-bond backbone of the respective polypeptide or protein. It is formed between the γ -carboxamide of the glutamyl residue of the acyl glutamine-containing amino acid donor sequence and a primary (1°) amine of the amino donor-comprising substrate according to the invention.

From the inventor's experience as well as from others, it seems that only few glutamines are typically targeted by MTG, thus making the MTG an attractive tool for site-specific and stoichiometric protein modifications.

Previously, glutamine 295 (Q295) was identified as the only reactive glutamine on the heavy chain of different IgG types to be specifically targeted by MTG with low-molecular weight primary amine substrates (Jeger et al. 2010).

Quantitative conjugation to Q295, however, was only possible upon removal of the glycan moiety at the asparagine residue 297 (N297) with PNGase F, while glycosylated antibodies could not be conjugated efficiently (conjugation efficiency <20%). This finding is also supported by the studies of Mindt et al. (2008) and Jeger et al. (2010) and Dickgiesser et al. 2020

In order to obviate deglycosylation it is also possible to insert a point mutation at the residue N297 which results in the ablation of the glycosylation called aglycosylation.

However, both approaches come with significant disadvantages. An enzymatic deglycosylation step is undesired under GMP aspects, because it has to be made sure that both the deglycosylation enzyme (e.g., PNGase F) as well as the cleaved glycan are removed from the medium, to ensure a high purity product.

The substitution of N297 against another amino acid has unwanted effects, too, because it may affect the overall stability of the C_H2 domain, and the efficacy of the entire conjugate as a consequence. Further, the glycan that is present at N297 has important immunomodulatory effects, as it triggers antibody dependent cellular cytotoxicity (ADCC) and the like. These immunomodulatory effects would get lost upon deglycosylation or substitution of N297 against another amino acid.

Furthermore, the genetic engineering of an antibody for payload attachment may have disadvantages in that the sequence insertion may increase immunogenicity and decrease the overall stability of the antibody.

It is hence one object of the present invention to provide a transglutaminase based antibody conjugation approach which does not require prior deglycosylation of the antibody, in particular of N297.

It is another object of the present invention to provide a transglutaminase based antibody conjugation approach which does not require the substitution or modification of N297 in the C_H2 domain.

It is one further object of the present invention to provide an antibody conjugation technology that allows the manufacture of highly homogenous conjugation products, both as regards stoichiometry as well as site-specificity of the conjugation.

These and further objects are met with methods and means according to the independent claims of the present invention. The dependent claims are related to specific embodiments.

SUMMARY OF THE INVENTION

The present invention relates to methods and linker structures for generating an antibody-linker conjugate and/or an antibody-payload conjugate by means of a microbial transglutaminase (MTG). The invention and general advantages of its features will be discussed in detail below.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows an illustration of one aspect of the present invention. MTG = microbial transglutaminase. The star symbol illustrates the payload or linking moiety B. Gp is a Gly residue, which is N-terminally in a peptide, and which is the substrate for MTG. Note that this process allows to maintain the glycosylation at N297. Note that in case B/star is a linking moiety, the actual payload still has to be conjugated to this moiety.

As discussed elsewhere herein, B/star can be or comprise a linking moiety, like e.g. a bio-orthogonal group (e.g., an azide/ N_3 -group) that is suitable for strain-promoted alkyne-azide cycloaddition (SPAAC) click-chemistry reaction to a DBCO-containing payload (e.g. a toxin or a fluorescent dye or a metal chelator, like DOTA or NODA-GA). This click-chemistry-based “two-step chemoenzymatic”-approach to attach the functional moiety to the antibody has the major advantage that it can be clicked at low molecular excess compared to the antibody, typically e.g. at 5eq per conjugation site or lower (Dennler et al. 2014). This allows for a cost-effective generation of ADCs. In addition, virtually any probe can be clicked with this approach ranging from fluorescent dyes to metal chelators (cf. Spycher et al. 2017, Dennler et al. 2015).

B/star can also be the actual payload, e.g., a toxin. Such embodiment allows the rapid manufacture of the resulting compound in one step, facilitating purification and production.

Fig. 2 shows an example of a linker peptide comprising an oligopeptide according to the invention. The sequence is GlyAlaArgLys(N_3) (GARK₁ with K₁ = Lys(N_3)). Lys(N_3) is a Lys residue in which the primary amine has been replaced by an azide group ($-N-N\equiv N$, or $-N_3$). According to the nomenclature of the present invention, either Lys(N_3) or N_3 alone can be regarded as the linking moiety B (in this example, N_3 is suitable for click-chemistry).

The peptide efficiently conjugates to native IgG1 antibodies (~77.3 % as estimated from LC-MS analysis under non-optimized conditions) at position Q295.

It is important to understand that in some linker peptides shown herein, the moiety at the C-terminus is simply designated as N₃. However, this should be understood as an abbreviation of Lys(N₃). For example, GAR(N₃) corresponds to the peptide GlyAlaArgLys(N₃) or GARK(N₃). That is, 6-azido-L-lysine may be abbreviated as Lys(N₃) in three letter code or as K(N₃) or (N₃) in single letter code. It is thus to be understood that K(N₃) when part of a peptide always relates to the single amino acid residue Lys(N₃) but not to the dipeptide Lys-Lys(N₃). A dipeptide Lys-Lys(N₃), on the other hand, would be designated KK(N₃) in single letter code.

It is furthermore important to understand that in different linker peptides shown herein, the C-terminus or primary amines on side chains may or may not be protected, even if shown otherwise. Protection can be accomplished by, e.g., amidation of the former, and/or acetylation of the latter. In the context of the present invention, both the protected and unprotected linker peptides are encompassed.

For example, GARK(N₃) does indeed encompass two variants, with the C-terminus protected or unprotected. The following figure shows a C-terminal Lys(N₃) residue wherein the C-terminus is protected by amidation:




Fig. 3 shows results of the screening of a small given peptide library to native IgG1 antibody. Different peptides were screened that contained MTG-reactive N-terminal amino acid residues or derivatives (beta-alanine). As can be seen, single or double N-terminal glycine works most efficient. LC-MS was used for analysis.

Figs. 4 and 5 show an embodiment wherein the linker comprises a Cys residue with a free sulfhydryl group, suitable to conjugate a maleimide-comprising toxin linker construct thereto.

Fig. 4 shows the binding reaction, and Fig. 5 some potential linker constructs.

Fig. 6 shows a two-step conjugation process (Fig. 6A) with the peptide being conjugated to the Gln of the antibody (e.g. Q295 of IgG or molecularly engineered) and a one-step conjugation process (Fig. 6B) according to the present invention. The following table 1 clarifies the two terms as used herein:

Table 1: One- and two step conjugation

Linker peptide	Process type	Steps
Gly(Aax) _m -Payload	One-step conjugation	step 1: conjugation of linker comprising the payload to Gln residue in antibody
Gly(Aax) _n -Cys-Linking moiety	Two-step conjugation	step 1: conjugation of linker comprising the Cys-Linking moiety to Gln residue in antibody step 2: conjugation of payload to Cys-Linking moiety

In the two-step process, the linker peptide is Gly-(Aax)_n-Cys-linking moiety. The Gly residue is conjugated to a Gln residue in the antibody via microbial transglutaminase, and the linking moiety – in this case a Cys residue with a free sulfhydryl group - is then conjugated to the payload, in this case a MMAE toxin carrying a MC/VC/PABDC linker structure, via the maleimide.

In the one-step process, the linker peptide Gly-(Aax)_m is already conjugated to the payload. The Gly residue is conjugated to a Gln residue in the antibody, and the payload consists of an MMAE toxin carrying a VC/PABC structure. The valine residue of the VC structure is conjugated to the last amino acid of the linker peptide by means of a peptide bond.

Fig.7 shows two examples of linkers comprising a linker suitable for dual-payload attachment.

Fig. 7A shows a peptide that has a first linking moiety which is an azide (N₃), while a second linking moiety is a tetrazine (both bio-orthogonal). The structure of the oligopeptide is GlyAlaArgLys(N₃)Lys(Tetrazine) (GARK₁K₂, with K₁=Lys(N₃), K₂ = Lys(Tetrazine)).

Figure 7B shows a peptide carrying an azide (N₃) and a free sulfhydryl-group from the Cys-moiety. The structure of the oligopeptide is GlyAlaArgLys(N₃)Cys (GARK₁C, with K₁=Lys(N₃)).

Each of the linking moieties are bio-orthogonally compatible groups that can be clicked simultaneously.

These linkers thus allow to conjugate two different payloads to the Q295 of the C_H2 domain of an antibody. Using a second payload allows for the development of a completely new class of antibody-payload conjugates that go beyond current therapeutic approaches with respect to efficacy and potency. Also new application fields are envisioned, for example, dual-type imaging for imaging and therapy or intra-/postoperative surgery (cf. Azhdarinia A. et al., *Molec Imaging and Biology*, 2012). For example, dual-labeled antibodies encompassing a molecular imaging agent for preoperative positron emission tomography (PET) and a near-infrared fluorescent (NIRF)-dye for guided delineation of surgical margins could greatly enhance the diagnosis, staging, and resection of cancer (cf. Houghton JL. et al., *PNAS* 2015). PET and NIRF optical imaging offer complementary clinical applications, enabling the non-invasive whole-body imaging to localize disease and identification of tumor margins during surgery, respectively. However, the generation of such dual-labeled probes up to date has been difficult due to a lack of suitable site-specific methods; attaching two different probes by chemical means results in an almost impossible analysis and reproducibility due to the random conjugation of the probes. Furthermore, in a study of Levengood M. et al., (*Angewandte Chemie*, 2016) a dual-drug labeled antibody, having attached two different auristatin toxins (having differing physiochemical properties and exerting complementary anti-cancer activities) imparted activity in cell line and xenograft models that were refractory to ADCs comprised of the individual auristatin components. This suggests that dual-labeled ADCs enable to address cancer heterogeneity and resistance more effectively than the single, conventional ADCs alone. Since one resistance mechanism towards ADCs include the active pumping-out of the cytotoxic moiety from the cancer cell, another dual-drug application may include the additional and simultaneous delivery of a drug that specifically blocks the efflux

mechanism of the cytotoxic drug. Such a dual-labeled ADC could thus help to overcome cancer resistance to the ADC more effectively than conventional ADCs.

Similar structures in which alkynes or tetrazine/trans-cyclooctenes are being used as linker are equally suitable and covered by the scope and gist of the present invention.

It is important to understand that in some linker peptides shown herein, the moiety at the C-terminus is simply designated as N₃. However, this should be understood as an abbreviation of Lys(N₃). For example, GAR(N₃) or GARK(N₃) does actually mean GARK₁, with K₁ = Lys(N₃), or GlyAlaArgLys(N₃).

It is furthermore important to understand that in different linker peptides shown herein, the C-terminus may or may not be protected, even if shown otherwise. Protection may be accomplished by amidation of the C-terminus. Since conjugation of the linker to an antibody is achieved via the primary amine of the N-terminal glycine residue of the linker, the N-terminus of the linker is preferably unprotected. In the context of the present invention, both the protected and unprotected linker peptides are encompassed. For example, GARK(N₃) does indeed encompass two variants, with a) both termini unprotected as discussed above, or b) only the C-terminus protected as discussed above.

The question whether or not the C-terminus is amidated is a practical question, depending on the conjugation conditions (buffer, medium, reactivity of the other reaction components, etc).

Fig. 8A and B show two possible linker structures with two Azide linker moieties, respectively. Fig. 8A shows GlyGlyAlaArgLys(N₃)Lys(N₃) (GGARK₁K₂, with K₁ and K₂=Lys(N₃)). Fig. 8B shows GlyGlyAlaArgLys(N₃)ArgLys(N₃) (GGARK₁RK₂; with K₁ and K₂=Lys(N₃)). In such way, an antibody payload ratio of 4 can be obtained. The presence of the charged Arg residues helps to keep hydrophobic payloads in solution.

It is important to understand that in some linker peptides shown herein, the moiety at the C-terminus is simply designated as N₃. However, this should be understood as an abbreviation of Lys(N₃). For example, GAR(N₃) or GARK(N₃) does actually mean GARK₁, with K₁ = Lys(N₃), or GlyAlaArgLys(N₃).

Fig. 9 shows further linkers that are suitable for MTG-mediated conjugation to native antibodies. These linker structures contain a linking moiety (azide, N₃) suitable for click-chemistry based attachment of the functional payload in a second step, or a Cys-residue which provides a thiol group suitable for attachment to a maleimide. Since these structures are based on peptides, that chemistry is well-understood and which is assembled from building blocks of single amino acids, new linkers can rapidly and easily be synthesized and evaluated. The following table 2 gives an overview:

Table 2

Structure	Sequence, residue for transglutaminase reaction in bold print		Linking moiety B
1	GlyAlaArgLys (N ₃)	GARK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)
2	GlyAlaArgXaa (N ₃)	GARX, with X = Xaa(N ₃), Xaa is 4-Azido-L-homoalanine	Xaa(N ₃)
3	GlyAlaArg [PEG] ₃ (N ₃)	GAR[PEG] ₃ N ₃ , with [PEG] ₃ = triethylenglycol	[PEG] ₃ N ₃
4	GlyAlaArgCys	GARC	Cysteine
5	GlyGlyAlaArgLys (PEG) _n ArgLys(N ₃)	GGAR[PEG] _n RK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)

Fig. 10 shows that the light chain of IgG1 antibodies is not modified by the conjugation. Shown is the deconvoluted LC-MS spectra of a IgG1 light chain.

Fig. 11 A shows deconvoluted LC-MS spectra of Trastuzumab native IgG1 heavy chain selectively modified with the N₃-functional linker GGARK(N₃). From the spectra it can be seen that the heavy chain got selectively and quantitatively (>95%) modified with only one peptide-linker since the observed mass difference corresponds to the expected peptide mass shift (Mw unmodified heavy chain = 50595 Da, expected Mw = 51091 Da, measured Mw = 51092 Da)

Fig. 11 B shows deconvoluted LC-MS spectra of Trastuzumab native IgG1 heavy chain selectively clicked with DBCO-PEG4-Ahx-DM1 to the N₃-functional linker GGARK(N₃) pre-installed on the heavy chain. From the spectra, it can be seen that the heavy chain got selectively and quantitatively (>95%) clicked.

Fig. 11 C shows the deconvoluted LC-MS of another IgG1 heavy chain modified with GGARK(N₃) under non-optimized conjugation conditions. Conjugation ratio: 83%

Fig. 12 A shows the deconvoluted LC-MS of Trastuzumab heavy chain modified with GARK(N₃). >95% conjugation efficiency was achieved.

Fig. 12 B shows the deconvoluted LC-MS of Trastuzumab heavy chain modified with GARK(N₃), clicked with DBCO-PEG4-Ahx-DM1 >95% clicking efficiency was achieved, resulting in an ADC with DAR 2.

Fig. 13 shows an overview of the Ig C_{H2} domain with the different numbering schemes. For the purposes of the present invention, the EU numbering is being used.

Fig. 14 shows a transglutaminase reaction to conjugate a linker having an N-terminal Gly residue with a free primary amine to the free primary amine of the Q295 residue of an antibody.

Fig. 15. Click chemistry reaction scheme (strain-promoted alkyne-azide cycloaddition (SPAAC)) to conjugate the linker GlyAlaArgLys(N₃) (GARK₁ with K₁ = Lys(N₃)) to dibenzocyclooctyne labelled with a payload.

Fig. 16 shows different peptide linkers that can be used in the context of the present invention, comprising a non-natural amino acid each.

Fig. 17 shows different linker toxin constructs that can be conjugated to an antibody according to the method described herein. In all cases, the Gly residues carry the primary amine for transglutaminase conjugation

Fig. 17A This Figure shows the non-cleavable GGARR-Ahx-May peptide-toxin conjugate with two arginine-groups serving to increase the solubility of the hydrophobic payload Maytansine (May). The primary amine of the N-terminal glycine residue serves for the conjugation to the antibody via MTG. The Ahx-spacer serves to decouple the positively-charged arginine from the May, helping the latter to more efficiently bind its target since the linker is not cleavable.

Fig. 17B This Figure shows the non-cleavable GGARR-PEG4-May peptide-toxin conjugate with two arginine-groups and a PEG4-spacer, all three moieties serving to increase the solubility of the hydrophobic payload May. The primary amine of the N-terminal glycine residue serves for the conjugation to the antibody via MTG. The PEG4 furthermore helps to decouple the positively-charged arginine from the May, helping the latter to more efficiently bind its target since the linker is not cleavable.

Fig. 17C This Figure shows the cleavable GGARR-PEG4-VC-MMAE peptide-toxin conjugate with two arginine-groups, a PEG4-spacer, a PABC-group and a val-cit sequence (VC). The primary amine of the N-terminal glycine residue serves for the conjugation to the antibody via MTG, the arginine-groups and the PEG4-spacer to increase the solubility and the PABC-group and the val-cit sequence help to release the toxin.

Fig. 17D This Figure shows the cleavable GGARR-MMAE peptide-toxin conjugate with two arginine-groups and a PABC-group with no PEG-spacer and val-cit sequence. Since the GGARR-group is intrinsically degradable by peptidases, no val-cit sequence might be necessary for toxin release through the self-immolative PABC-moiety, and as the two arginine-groups are very hydrophilic no PEG-spacer may be needed, keeping thus the whole peptide-toxin conjugate as small as possible to minimize undesired interactions with other molecules while in blood circulation.

Fig. 18 shows results of a cellular toxicity assay as performed according to example 3. The Her-GARK(N₃) (P684) and Her-GGARK(N₃) (P579) N-terminal Glycine ADCs which have been generated with the method according to the invention and comprise a May-moiety click-attached to each linker have similar potency against SK-BR3 cells as Kadcyla. Hence, the advantages provided by the novel linker technology (ease of manufacture, site specificity, stable stoichiometry, no need to deglycosylate that antibody) do not come at any disadvantage regarding the cellular toxicity.

Fig. 19: Structure of β Ala-Gly-Ala-Arg-Lys(N₃). β Ala designates β -alanine, which is structurally similar to glycine. However the said linker has inferior conjugation efficiency compared to GGARK(N₃) (see example 2), which has an N-terminal glycine.


DETAILED DESCRIPTION OF THE INVENTION

Before the invention is described in detail, it is to be understood that this invention is not limited to the particular components or process steps of the methods described as such devices and methods may vary. It is also to be understood that the terminology used herein is for purposes of describing particular embodiments only, and is not intended to be limiting. It must be noted that, as used in the specification and the appended claims, the singular forms "a", "an", and "the" include singular and/or plural referents unless the context clearly dictates otherwise. It is moreover to be understood that, in case parameter ranges are given which are delimited by numeric values, the ranges are deemed to include these limitation values.

It is further to be understood that embodiments disclosed herein are not meant to be understood as individual embodiments which would not relate to one another. Features discussed with one embodiment are meant to be disclosed also in connection with other embodiments shown herein. If, in one case, a specific feature is not disclosed with one embodiment, but with another, the skilled person would understand that does not necessarily mean that said feature is not meant to be disclosed with said other embodiment. The skilled person would understand that it is the gist of this application to disclose said feature also for the other embodiment, but that just for purposes of clarity and to keep the specification in a manageable volume this has not been done.

Furthermore, the content of the documents referred to herein is incorporated by reference. This refers, particularly, for documents that disclose standard or routine methods. In that case, the incorporation by reference has mainly the purpose to provide sufficient enabling disclosure, and avoid lengthy repetitions.

According to a first aspect, a method for generating an antibody-payload conjugate or an antibody-linker conjugate by means of a microbial transglutaminase (MTG) is provided, which method comprises the step of conjugating a linker comprising the peptide structure (shown in N->C direction)




via the N-terminal primary amine of the N-terminal glycine (Gly) residue to a glutamine (Gln) residue comprised in the heavy or light chain of an antibody,

wherein

- m is an integer between ≥ 0 and ≤ 12 ,
- n is an integer between ≥ 0 and ≤ 12 ,
- $m + n \geq 0$,
- Aax is an amino acid or an amino acid derivative, and
- B is a payload or a linking moiety.

As used herein, the term “primary amine” relates to an amine substituted with two hydrogen atoms, of the general formula R-NH₂.


In certain embodiments, the peptide linker may comprise two or more linking moieties and/or payloads. That is, the linker may have the peptide structure (shown in N -> C direction)



wherein

- m , n and o are integers between ≥ 0 and ≤ 12 ,
- $m + n + o \geq 0$,
- Aax is an amino acid or an amino acid derivative, and
- B₁ and B₂ are payloads and/or linking moieties, wherein B₁ and B₂ may be identical or different from each other.

In other embodiments, the peptide linker may comprise three linking moieties and/or payloads. That is, the linker may have the peptide structure (shown in N -> C direction)




wherein

- m , n , o and p are integers between ≥ 0 and ≤ 12 ,
- $m + n + o + p \geq 0$,
- Aax is an amino acid or an amino acid derivative, and

- B₁, B₂ and B₃ are payloads and/or linking moieties, wherein B₁, B₂ and B₃ may be identical or different from each other.

It is to be understood, that the invention also encompasses linkers comprising more than three linking moieties and/or payloads, such as 4, 5 or 6 linking moieties and/or payloads. In this case, the peptide structure of the linkers follows the same pattern as described above for the linkers comprising 2 or 3 linking moieties and/or payloads.

In certain embodiments, a method for generating an antibody-payload conjugate by means of a microbial transglutaminase (MTG) is provided, which method comprises the step of conjugating a linker having the peptide structure (shown in N->C direction)




via the N-terminal primary amine of the N-terminal glycine (Gly) residue to a glutamine (Gln) residue comprised in the heavy or light chain of an antibody,

wherein

- m is an integer between ≥ 0 and ≤ 12
- n is an integer between ≥ 0 and ≤ 12
- $m + n \geq 0$,
- Aax can be any naturally or non-naturally occurring L- or D-amino acid, or amino acid derivative or mimetic, and
- B is a payload or a linking moiety.

In certain embodiments, the invention relates to a method for generating an antibody-payload conjugate or an antibody-linker conjugate by means of a microbial transglutaminase (MTG), which method comprises the step of conjugating a linker having the peptide structure (shown in N->C direction)



via the N-terminal primary amine of the N-terminal glycine (Gly) residue to a glutamine (Gln) residue comprised in the heavy or light chain of an antibody. In this case, it is to be understood that the moiety B may comprise more than one payload and/or linking moiety. For example, B may stand for $(B'-(Aax)_o-B'')$, wherein B' and B'' are payloads and/or linking moieties and wherein o is an integer between ≥ 0 and ≤ 12 . Alternatively, B may stand for $(B'-(Aax)_o-B''-(Aax)_p-B''')$, wherein B', B'' and B''' are payloads and/or linking moieties and wherein o and p are integers between ≥ 0 and ≤ 12 .

Thus, in a particular embodiment, the invention relates to a method according to the invention, wherein the linker comprises two or more payloads and/or linking moieties. In another embodiment, the invention relates to a method according to the invention, wherein the two or more payloads and/or linking moieties B differ from one another.

That is, the linker according to the invention may comprise a single payload or linking moiety. In certain embodiments, the linker comprises two linking moieties, wherein the two linking moieties are identical. In other embodiments, the linker comprises two linking moieties, wherein the two linking moieties are different. In yet another embodiment, the linker comprises two identical or different payloads. The invention further encompasses linkers comprising one or more payload and one or more linking moiety.

It is further to be understood that not all payloads or linking moieties can function as an intrachain payload or linking moiety, for example, because they do not have the functional groups to form peptide or amide bonds with the C-terminal carboxyl group of a first Aax moiety and the N-terminal amine group of a second Aax moiety. In this case, it is preferred that such payload or linking moieties are located at the C-terminal end of the linker, where they preferably are attached to the carboxyl group of the C-terminal Aax moiety of the linker. In cases where the payload or linking moiety are at an intrachain position of the linker, it is preferred that the payload or linking moiety is an amino acid, an amino acid derivative or attached to a molecule having the general structure $—NH—CHR—CO—$.

In preferred embodiments, m and/or n is $\geq 1, \geq 2, \geq 3, \geq 4, \geq 5, \geq 6, \geq 7, \geq 8, \geq 9, \geq 10$, or ≥ 11 . In other preferred embodiments, m and/or n is $\leq 12, \leq 11, \leq 10, \leq 9, \leq 8, \leq 7, \leq 6, \leq 5, \leq 4, \leq 3, \leq 2$, or ≤ 1 . In further preferred embodiments, m + n is $\geq 1, \geq 2, \geq 3, \geq 4, \geq 5, \geq 6, \geq 7, \geq 8, \geq 9, \geq 10$, or ≥ 11 .

In still further preferred embodiments $m + n$ is $\leq 12, \leq 11, \leq 10, \leq 9, \leq 8, \leq 7, \leq 6, \leq 5, \leq 4, \leq 3, \leq 2,$ or ≤ 1 .

Members of both ranges can be combined with another to disclose a preferred length range with lower and upper limit.

Accordingly, in a particular embodiment, the invention relates to a method according to the invention, wherein $m+n$, and optionally $m + n + o$ and $m + n + o + p$, is $\leq 12, \leq 11, \leq 10, \leq 9, \leq 8, \leq 7, \leq 6, \leq 5$ or ≤ 4 .

It is important to understand that in different linker peptides shown herein, the C-terminus may or may not be protected, even if shown otherwise. Protection can be accomplished by amidation of the former. In the context of the present invention, both the protected and unprotected linker peptides are encompassed.

The inventors have shown that this process is suitable to very cost effectively and quickly produce site-specific antibody-payload conjugates (24 – 36 hrs, or optionally 48 hrs), and hence allows the production of large libraries of such molecules, and subsequent screening thereof in high throughput screening systems.

In contrast thereto, a Cys engineering process in which an antibody payload conjugate is produced where the payload is conjugated to the antibody via a genetically (molecularly) engineered Cys residue needs at least about 3 – 4 weeks.

In general, the method allows conjugation of a large number of payloads to an antibody. For each payload, a suitable peptide linker structure can be identified from a large linker pool to deliver optimal clinical and non-clinical characteristics. This is not possible in other methods where the linker structure is fixed. In addition, the method according to the invention allows to generate antibody-payload conjugates comprising two or more different payloads, wherein each payload is conjugated to the antibody in a site-specific manner. Thus, the method according to the invention may be used to generate antibodies with novel and/or superior therapeutic or diagnostic capacities.

The linker may comprise any amino acid, including, without limitation, α -, β -, γ -, δ - and ϵ -amino acids. In the case of α -amino acids, the linker may comprise any naturally occurring L- or D-amino acid. A naturally occurring L- or D-amino acid encompasses any L- or D-amino acid that can be found in nature. That is, the term “naturally occurring L- or D-amino” acid encompasses all canonical or proteogenic amino acids that are used as building blocks in naturally-occurring proteins. In addition, the term “naturally occurring L- or D-amino” acid encompasses all non-canonical L- or D-amino acids that can be found in nature, for example as metabolic intermediates or degradation products or as building blocks for other non-proteogenic macromolecules.

Further, the linker may comprise non-naturally occurring L- or D-amino acids. A non-naturally occurring L- or D-amino acid encompasses any molecule having the general structure $\text{H}_2\text{N}-\text{CHR}-\text{COOH}$, which has not previously been found in nature.

The skilled person is aware of resources and databases to consult when determining whether an L- or D-amino acid is naturally or non-naturally occurring. However, in cases of doubt, it is to be understood that the term “naturally or non-naturally occurring L- or D-amino acid” encompasses the L- and D-isomer of any molecule having the general structure $\text{H}_2\text{N}-\text{CHR}-\text{COOH}$, irrespective of the origin of said molecule.

In certain embodiments, the linker of the invention may also comprise naturally or non-naturally occurring, non-chiral amino acids having the general structure $\text{H}_2\text{N}-\text{CR}_1\text{R}_2-\text{COOH}$.

Furthermore, the linker of the invention may comprise amino acid derivatives. An amino acid derivative is a compound that has been derived from a naturally or non-naturally occurring amino acid by one or more chemical reactions, such as chemical reactions of the α -amino group, the α -carboxylic acid group and/or the amino acid side chain. That is, the term amino acid derivative encompasses any molecule having the structure $-\text{NH}-\text{CHR}-\text{CO}-$, which has been derived from a naturally or non-naturally occurring L- or D-amino acid. Since it is envisioned that the amino acid derivative of the invention is part of a peptide-based linker, it is preferred that the amino acid derivative has been obtained by one or more chemical reactions of the amino acid side chain of a naturally or non-naturally occurring L- or D-amino acid, or, in cases where the amino acid derivative is located at the C-terminal end of the

peptide, the alpha-carboxylic acid group of a naturally or non-naturally occurring L- or D-amino acid. It is to be noted that naturally and non-naturally occurring amino acids can be amino acid derivatives and vice versa.

Examples of non-canonical amino acids, non-naturally occurring amino acids and amino acid derivatives that may be comprised in the linker of the invention include, but are not limited to, α -aminobutyric acid, α -aminoisobutyric acid, ornithine, hydroxyproline, agmatine, (S)-2-amino-4-((2-amino)pyrimidinyl)butanoic acid, α -aminoisobutyric acid, *p*-benzoyl-L-phenylalanine, t-butylglycine, citrulline, cyclohexylalanine, desamino tyrosine, L-(4-guanidino)phenylalanine, homoarginine, homocysteine, homoserine, homolysine, n-formyl tryptophan, norleucine, norvaline, phenylglycine, (S)-4-piperidyl-(N-amidino)glycine, parabenzoyl-L-phenylalanine, sarcosine and 2-thienyl alanine.

Besides alpha-amino acids as described above, the linker of the invention may also comprise one or more β -, γ -, δ - or ϵ -amino acids. Thus, in certain embodiments, the linker may be a peptidomimetic. The peptidomimetic may not exclusively contain classical peptide bonds that are formed between two α -amino acids but may additionally or instead comprise one or more amide bonds that are formed between an alpha amino acid and a β -, γ -, δ - or ϵ -amino acid, or between two β -, γ -, δ - or ϵ -amino acids. An example of a linker that is a peptidomimetic and comprises an amide bond between an α -amino acid and a β -amino acid is shown in Figure 16 (Gly- β -Ala-Arg-Lys(N₃)). Accordingly, in any instance of the present invention where the linker is described as a peptide, it is to be understood that the linker may also be a peptidomimetic and thus not exclusively consist of α -amino acids, but may instead comprise one or more β -, γ -, δ - or ϵ -amino acids or molecules that are not classified as an amino acid. Examples of β -, γ -, δ - or ϵ -amino acids that may be comprised in the linker of the present invention include, but are not limited to, β -alanine, γ -aminobutyric acid, 4-amino-3-hydroxy-5-phenylpentanoic acid, 4-amino-3-hydroxy-6-methylheptanoic acid, 6-aminohexanoic acid and statine.

The term "D-amino acid" is understood to comprise the D-counterparts of both naturally occurring amino acids as well as of non-naturally occurring amino acids.

Since the peptide linkers of the present invention are peptide-based, they are likely to be hydrolyzed by a host cell peptidase once the antibody-payload conjugate has been internalized

into a target cell. Accordingly, in certain embodiments, the linker does not necessarily need to comprise a cathepsin cleavage site. Thus, in one embodiment, the linker comprising or having the peptide structure is not cleavable by cathepsin. This includes, in particular, cathepsin B. In one further embodiment, the linker comprising or having the peptide structure does not comprise a valine-alanine motif or a valine-citrulline motif. However, it is to be understood that the invention also encompasses linkers that comprise a cathepsin cleavage site, such as valine-alanine or valine-citrulline. For example, linkers comprising non-canonical or D-amino acids may not be cleaved efficiently by host cell peptidases. In this case, a cathepsin cleavage site in the linker may improve the release of the payload after internalization into the host cell. The linker may further comprise other motifs or self-immolative groups that allow efficient release of the payload inside a target cell if required.

One typical dipeptide structure used in ADC linkers, yet devoid of a Lys residue, is the valine-citrulline motif, as e.g. provided in Brentuximab Vedotin, and discussed in Dubowchik and Firestone 2002. This linker can be cleaved by cathepsin B to release the toxin at the site of disease. The same applies to the valine-alanine motif, which is for example provided in SGN-CD33A.

In one further embodiment, the linker does not comprise polyethylene glycol or a polyethylene glycol derivative.

Polyethylene glycol (PEG) is a polyether compound with many applications from industrial manufacturing to medicine. PEG is also known as polyethylene oxide (PEO) or polyoxyethylene (POE), depending on its molecular weight. The structure of PEG is commonly expressed as $H-(O-CH_2-CH_2)_n-OH$. However, it is to be understood that the linkers of the invention may comprise PEG or a PEG-derivative.

It is hence important to understand that, because B can either be a payload or a linking moiety, the method according to the invention has two major embodiments, as shown in the following table 3:

Table 3

Linker peptide	Process type	Steps
Gly-(Aax) _m -Payload-(Aax) _n	One-step conjugation	step 1: conjugation of linker comprising

Gly-(Aax) _m -linking moiety-(Aax) _n	Two-step conjugation	the payload to Gln residue in antibody step 1: conjugation of linker comprising the linking moiety to Gln residue in antibody step 2: conjugation of payload to Linking moiety
---	----------------------	--

That is, in certain embodiments, the payload is coupled to the linker by chemical synthesis. Accordingly, the linker may have the structure Gly-(Aax)_m-Payload or Gly-(Aax)_m-Payload-(Aax)_n. For example, the payload may be coupled to the C-terminus of a peptide by chemical synthesis. Thus, in certain embodiments the linker may have the structure Gly-Ala-Arg-Payload, Gly-Ala-Arg-Arg-Payload, Gly-Gly-Ala-Arg-Payload, Gly-Gly-Ala-Arg-Arg-Payload or Gly-Gly-Gly-Payload.

According to one further embodiment of the invention, the antibody is at least one selected from the group consisting of

- IgG, IgE, IgM, IgD, IgA and IgY
- IgG1, IgG2, IgG3, IgG4, IgA1 and IgA, and/or
- a fragment or recombinant variant thereof retaining target binding properties and comprising the C_H2 domain

The antibody is preferably a monoclonal antibody.

The antibody can be of human origin, but likewise from mouse, rat, goat, donkey, hamster, or rabbit. In case the conjugate is for therapy, a murine or rabbit antibody can optionally be chimerized or humanized.

Fragment or recombinant variants of antibodies comprising the C_H2 domain are, for example,

- antibody formats comprising mere heavy chain domains (shark antibodies/IgNAR (V_H-C_H1-C_H2-C_H3-C_H4-C_H5)₂ or camelid antibodies/hcIgG (V_H-C_H2-C_H3)₂)
- scFv-Fc (V_H-V_L-C_H2-C_H3)₂
- Fc fusion peptides, comprising an Fc domain and one or more receptor domains.

The antibody can also be bispecific (e.g., DVD-IgG, crossMab, appended IgG – HC fusion) or biparatopic. See Brinkmann and Kontermann (2017) for an overview.

Accordingly, in a particular embodiment, the invention relates to the method according to the invention, wherein the antibody is an IgG, IgE, IgM, IgD, IgA or IgY antibody, or a fragment or recombinant variant thereof, wherein the fragment or recombinant variant thereof retains target binding properties and comprises a C_H2 domain.

In a preferred embodiment, the antibody is an IgG antibody. That is, the antibody may be an IgG antibody that is glycosylated, preferably at residue N297. Alternatively, the antibody may be a deglycosylated antibody, preferably wherein the glycan at residue N297 has been cleaved off with the enzyme PNGase F. Further, the antibody may be an aglycosylated antibody, preferably wherein residue N297 has been replaced with a non-asparagine residue. Methods for deglycosylating antibodies and for generating aglycosylated antibodies are known in the art.

As discussed herein, IgG antibodies that are glycosylated at residue N297 have several advantages over non-glycosylated antibodies. In addition, it has been demonstrated that the linkers of the invention can be conjugated to antibodies that are glycosylated at residue N297 with unexpectedly high efficiency. Accordingly, in an even more preferred embodiment, the antibody is an IgG antibody that is glycosylated at residue N297 (EU numbering) of the C_H2 domain.

In a particular embodiment, the invention relates to the method according to the invention, wherein (a) the linker including the payload or linking moiety B is conjugated to a Gln residue which has been introduced into the heavy or light chain of the antibody by molecular engineering or (b) the linker including the payload or linking moiety B is conjugated to a Gln residue in the Fc domain of the antibody.

According to one further embodiment of the invention, the payload or linking moiety is conjugated to a Gln residue which was introduced into the heavy or light chain of the antibody by molecular engineering.

The term “molecular engineering,” as used herein, refers to the use of molecular biology methods to manipulate nucleic acid sequences. Within the present invention, molecular engineering may be used to introduce Gln residues into the heavy or light chain of an antibody. In general, two different strategies to introduce Gln residues into the heavy or light chain of an antibody are envisioned within the present invention. First, single residues of the heavy or light chain of an antibody may be substituted with a Gln residue. Second, Gln-containing peptide tags consisting of two or more amino acid residues may be integrated into the heavy or light chain of an antibody. For that, the peptide tag may either be integrated into an internal position of the heavy or light chain, that is, between two existing amino acid residues of the heavy or light chain or by replacing them, or the peptide tag may be fused (appended) to the N- or C-terminal end of the heavy or light chain of the antibody.

In the first case, any amino residue of the heavy or light chain of an antibody may be substituted with a Gln residue, provided that the resulting antibody can be conjugated with the linkers of the invention by a microbial transglutaminase. In certain embodiments, the antibody is an antibody wherein amino acid residue N297 (EU numbering) of the C_H2 domain of an IgG antibody is substituted, in particular wherein the substitution is an N297Q substitution. Antibodies comprising an N297Q mutation may be conjugated to more than one linker per heavy chain of the antibody. For example, antibodies comprising an N297Q mutation may be conjugated to four linkers, wherein a one linker is conjugated to residue Q295 of the first heavy chain of the antibody, one linker is conjugated to residue N297Q of the first heavy chain of the antibody, one linker is conjugated to residue Q295 of the second heavy chain of the antibody and one linker is conjugated to residue N297Q of the second heavy chain of the antibody. The skilled person is aware that replacement of residue N297 of an IgG antibody with a Gln residue results in an aglycosylated antibody.

In a particular embodiment, the invention relates to the method according to the invention, wherein the Gln residue that has been introduced into the heavy or light chain of the antibody by molecular engineering is comprised in a peptide that has been (a) integrated into the heavy or light chain of the antibody or (b) fused to the N- or C-terminal end of the heavy or light chain of the antibody.

Thus, instead of substituting single amino acid residues of an antibody, peptide tags comprising a Gln residue that is accessible for a transglutaminase may be introduced into the

heavy or light chain of the antibody. Such peptide tags may be fused to the N- or C-terminus of the heavy or light chain of the antibody. Preferably, peptide tags comprising a transglutaminase-accessible Gln residue are fused to the C-terminus of the heavy chain of the antibody. Even more preferably, the peptide tags comprising a transglutaminase-accessible Gln residue are fused to the C-terminus of the heavy chain of an IgG antibody. Several peptide tags that may be fused to the C-terminus of the heavy chain of an antibody and serve as substrate for a microbial transglutaminase are described in WO 2012/059882, WO 2016/144608, WO 2016/100735, WO 2016/096785 and by Steffen et al. (JBC, 2017) and Malesevic et al. (Chembiochem, 2015).

Exemplary peptide linkers that may be introduced into the heavy or light chain of an antibody, in particular fused to the C-terminus of the heavy chain of the antibody, are LLQGG, LLQG, LSLSQG, GGLLQGG, GLLQG, LLQ, GSPLAQSHGG, GLLQGGG, GLLQGG, GLLQ, LLQLLQGA, LLQGA, LLQYQGA, LLQGSG, LLQYQG, LLQLLQG, SLLQG, LLQLQ, LLQLLQ, LLQGR, EEQYASTY, EEQYQSTY, EEQYNSTY, EEQYQS, EEQYQST, EQYQSTY, QYQS, QYQSTY, YRYRQ, DYALQ, FGLQRPY, EQKLISEEDL, LQR and YQR.

The skilled person is aware of methods to substitute amino acid residues of antibodies or to introduce peptide tags into antibodies, for example by methods of molecular cloning as described in Sambrook, Joseph. (2001). Molecular cloning : a laboratory manual. Cold Spring Harbor, N.Y. :Cold Spring Harbor Laboratory Press.

According to one further embodiment of the invention, the payload or linking moiety is conjugated to a Gln in the Fc domain of the antibody.

That is, the linkers of the invention may be conjugated to any Gln residue in the Fc domain of the antibody that can serve as a substrate for a microbial transglutaminase.

Typically, the term Fc domain as used herein refers to the last two constant region immunoglobulin domains of IgA, IgD and IgG (C_{H2} and C_{H3}) and the last three constant region domains of IgE, IgY and IgM (C_{H2}, C_{H3} and C_{H4}). That is, the linker comprising the payload or linking moiety B may be conjugated to the C_{H2}, C_{H3} and, where applicable, C_{H4} domains of the antibody.

According to one further embodiment of the invention, the payload or linking moiety is conjugated to the Gln residue Q295 (EU numbering) of the C_H2 domain of the antibody. In a particular embodiment, the invention relates to the method according to the invention, wherein the Gln residue in the Fc domain of the antibody is Gln residue Q295 (EU numbering) of the C_H2 domain of an IgG.

It is important to understand that Q295 is an extremely conserved amino acid residue in IgG type antibodies. It is conserved in human IgG1, 2, 3, 4, as well as in rabbit and rat antibodies amongst others. Hence, being able to use Q295 is a considerable advantage for making therapeutic antibody-payload conjugates, or diagnostic conjugates where the antibody is often of non-human origin. The method according to the invention does hence provide an extremely versatile and broadly applicable tool. Even though residue Q295 is extremely conserved among IgG type antibodies, some IgG type antibodies do not possess this residue, such as mouse IgG2a or IgG2b. Thus, it is to be understood that the antibody used in the method of the present invention is preferably an IgG type antibody comprising residue Q295 (EU numbering) of the C_H2 domain.

Further, it has been shown that engineered conjugates using Q295 for payload attachment demonstrate good pharmacokinetics and efficacy (Lhospice et al. 2015), and are capable of carrying even unstable toxins prone for degradation (Dorywalska et al. 2015). It is thus expected that similar effects will be seen with this site-specific method since the same residue is modified, but of glycosylated antibodies. Glycosylation may further contribute to overall ADC stability, removal of the glycan moieties as with the mentioned approaches has been shown to result in less-stable antibodies (Zheng et al. 2011).

According to one further embodiment of the invention, the antibody to which the payload or linking moiety is conjugated is glycosylated.

Typical IgG shaped antibodies are N-glycosylated in position N297 (Asp-X-Ser/Thr-motif) of the C_H2 domain.

Accordingly, in a particular embodiment, the invention relates to the method according to the invention, wherein the Gln residue in the Fc domain of the antibody is Gln residue Q295 (EU

numbering) of the C_H2 domain of an IgG antibody that is glycosylated at residue N297 (EU numbering) of the C_H2 domain.

In the literature discussing the conjugation of linkers to a C_H2 Gln residue by means of a transglutaminase, the focus has been on small, low-molecular weight substrates. However, in the prior art literature, to accomplish such conjugation, a deglycosylation step in position N297, or the use of an aglycosylated antibody, is always described as necessary (WO 2015/015448; WO 2017/025179; WO 2013/092998).

Quite surprisingly, and against all expectations, however, site-specific conjugation to Q295 of glycosylated antibodies is indeed efficiently possible by using the above discussed oligopeptide structure.

Though Q295 is very close to N297, which is, in its native state, glycosylated, the method according to the invention, using the specified linker, still allows the conjugation of the linker or payload thereto.

However, as shown, the method according to the invention does not require an upfront enzymatic deglycosylation of Q295, nor the use of an aglycosylated antibody, nor a substitution of N297 against another amino acid, nor the introduction of a T299A mutation to prevent glycosylation.

These two points provide significant advantages under manufacturing aspects. An enzymatic deglycosylation step is undesired under GMP aspects, because it has to be made sure that the both the deglycosylation enzyme (e.g., PNGase F) as well as the cleaved glycan have to be removed from the medium.

Furthermore, no genetic engineering of the antibody for payload attachment is necessary, so that sequence insertions which may increase immunogenicity and decrease the overall stability of the antibody can be avoided.

The substitution of N297 against another amino acid has unwanted effects, too, because it may affect the overall stability of the entire Fc domain (Subedi et al, 2015), and the efficacy of the entire conjugate as a consequence that can lead to increased antibody aggregation and a

decreased solubility (Zheng et al. 2011) that particularly gets important for hydrophobic payloads such as PBDs. Further, the glycan that is present at N297 has important immunomodulatory effects, as it triggers antibody dependent cellular cytotoxicity (ADCC) and the like. These immunomodulatory effects would get lost upon deglycosylation or any of the other approaches discussed above to obtain an aglycosylated antibody. Further, any sequence modification of an established antibody can also lead to regulatory problems, which is problematic because often times an accepted and clinically validated antibody is used as a starting point for ADC conjugation.

Hence, the method according to the invention allows to easily and without disadvantages make stoichiometrically well-defined ADCs with site specific payload binding.

In view of the above, it is stated that the method of the present invention is preferably used for the conjugation of an IgG antibody at residue Q295 (EU numbering) of the C_{H2} domain of the antibody, wherein the antibody is glycosylated at residue N297 (EU numbering) of the C_{H2} domain. However, it is expressly stated that the method of the invention also encompasses the conjugation of deglycosylated or aglycosylated antibodies at residue Q295 or any other suitable Gln residue of the antibody, wherein the Gln residue may be an endogenous Gln residue or a Gln residue that has been introduced by molecular engineering.

The invention also encompasses the conjugation of antibodies of other isotypes than IgG antibodies, such as IgA, IgE, IgM, IgD or IgY antibodies. Conjugation of these antibodies may take place at an endogenous Gln residue, for example an endogenous Gln residue in the Fc domain of the antibody, or at a Gln residue that has been introduced into the antibody by molecular engineering.

In general, the skilled person is aware of methods to determine at which position of an antibody a linker is conjugated. For example, the conjugation site may be determined by proteolytic digestion of the antibody-payload conjugate and LC-MS/MS analysis of the resulting fragments. For example, samples may be deglycosylated with GlycINATOR (Genovis) according to the instruction manual and subsequently digested with trypsin gold (mass spectrometry grade, Promega), respectively. Therefore, 1 µg of protein may be incubated with 50 ng trypsin at 37 °C overnight. LC-MS/MS analysis may be performed using a nanoAcquity HPLC system coupled to a Synapt-G2 mass spectrometer (Waters). For

that, 100 ng peptide solution may be loaded onto an Acquity UPLC Symmetry C18 trap column (Waters, part no. 186006527) and trapped with 5 $\mu\text{L}/\text{min}$ flow rate at 1 % buffer A (Water, 0.1 % formic acid) and 99 % buffer B (acetonitrile, 0.1 % formic acid) for 3 min. Peptides may then be eluted with a linear gradient from 3 % to 65 % Buffer B within 25 min. Data may be acquired in resolution mode with positive polarity and in a mass range from 50 to 2000 m/z . Other instrument settings may be as follows: capillary voltage 3,2 kV, sampling cone 40 V, extraction cone 4.0 V, source temperature 130 $^{\circ}\text{C}$, cone gas 35 L/h, nano flow gas 0.1 bar, and purge gas 150 L/h. The mass spectrometer may be calibrated with [Glu1]-Fibrinopeptide.

Further, the skilled person is aware of methods to determine the drug-to-antibody (DAR) ratio or payload-to-antibody ratio of an antibody-payload construct. For example, the DAR may be determined by hydrophobic interaction chromatography (HIC) or LC-MS.

For hydrophobic interaction chromatography (HIC), samples may be adjusted to 0.5 M ammonium sulfate and assessed *via* a MAB PAK HIC Butyl column (5 μm , 4.6 x 100 mm, Thermo Scientific) using a full gradient from A (1.5 M ammonium sulfate, 25 mM Tris HCl, pH 7.5) to B (20 % isopropanol, 25 mM Tris HCl, pH 7.5) over 20 min at 1 mL/min and 30 $^{\circ}\text{C}$. Typically, 40 μg sample may be used and signals may be recorded at 280 nm. Relative HIC retention times (HIC-RRT) may be calculated by dividing the absolute retention time of the ADC DAR 2 species by the retention time of the respective unconjugated mAb.

For LC-MS DAR determination, ADCs may be diluted with NH_4HCO_3 to a final concentration of 0.025 mg/mL. Subsequently, 40 μL of this solution may be reduced with 1 μL TCEP (500 mM) for 5 min at room temperature and then alkylated by adding 10 μL chloroacetamide (200 mM), followed by overnight incubation at 37 $^{\circ}\text{C}$ in the dark. For reversed phase chromatography, a Dionex U3000 system in combination with the software Chromeleon may be used. The system may be equipped with a RP-1000 column (1000 \AA , 5 μm , 1.0 x 100 mm, Sepax) heated to 70 $^{\circ}\text{C}$, and an UV-detector set to a wavelength of 214 nm. Solvent A may consist of water with 0.1 % formic acid and solvent B may comprise 85 % acetonitrile with 0.1 % formic acid. The reduced and alkylated sample may be loaded onto the column and separated by a gradient from 30 – 55 % solvent B over the course of 14 min. The liquid chromatography system may be coupled to a Synapt-G2 mass spectrometer for identification of the DAR species. The capillary voltage of the mass spectrometer may be set

to 3 kV, the sampling cone to 30 V and the extraction cone may add up to a value of 5 V. The source temperature may be set to 150 °C, the desolvation temperature to 500 °C, the cone gas to 20 l/h, the desolvation gas to 600 l/h, and the acquisition may be made in positive mode in a mass range from 600-5000 Da with 1 s scan time. The instrument may be calibrated with sodium iodide. Deconvolution of the spectra may be performed with the MaxEnt1 algorithm of MassLynx until convergence. After assignment of the DAR species to the chromatographic peaks, the DAR may be calculated based on the integrated peak areas of the reversed phase chromatogram.

According to one further embodiment of the invention, the net charge of the linker is neutral or positive.

The net charge of a peptide is usually calculated at neutral pH (7.0). In the simplest approach, the net charge is determined by adding the number of positively charged amino acid residues (Arg and Lys and optionally His) and the number of negatively charged ones (Asp and Glu), and calculate the difference of the two groups. In cases where the linker comprises non-canonical amino acids, the skilled person is aware of methods to determine the charge of the non-canonical amino acid at neutral pH.

According to one further embodiment of the invention, the linker does not comprise negatively charged amino acid residues.

Preferably, the oligopeptide does not comprise the negatively charged amino acid residues Glu and Asp or negatively charged non-canonical amino acids.

According to one further embodiment of the invention, the linker comprises positively charged amino acid residues.

According to one embodiment of the invention, the linker comprises at least two amino acid residues selected from the group consisting of

- Lysine or a Lysine derivative or a Lysine mimetic,
- Arginine, and/or
- Histidine.

In certain embodiments, the linker comprises at least one amino acid residue selected from the group consisting of

- Lysine or a Lysine derivative or a Lysine mimetic,
- Arginine, and
- Histidine.

In certain embodiments, the linker comprises at least one amino acid residue selected from the group consisting of

- Lysine,
- Arginine, and
- Histidine.

In certain embodiments, the linker comprises at least one amino acid residue selected from the group consisting of

- Arginine, and
- Histidine.

In certain embodiments, the linker comprises at least one arginine residue.

Table 8 shows that linkers with negative, neutral and positive net charge can be conjugated to a glycosylated antibody with the method of the invention. In particular, linkers comprising a positively charged arginine residue can be conjugated to the glycosylated antibody with high efficiency.

That is, in certain embodiments, the linker according to the invention has a neutral or positive net charge. In certain embodiments, the linker according to the invention has a neutral or positive net charge and comprises at least one arginine and/or histidine residue. In certain embodiments, the linker according to the invention has a neutral or positive net charge and comprises at least one arginine residue. In certain embodiments, the linker according to the

invention does not comprise a lysine residue. In certain embodiments, the linker according to the invention has a neutral or positive net charge and does not comprise a lysine residue.

Table 8 further shows that linkers with the amino acid sequence Gly-[Gly/Ala]-Arg-B can be efficiently conjugated to a glycosylated antibody. Accordingly, in certain embodiments, the linker according to the invention has the sequence Gly-[Gly/Ala]-Arg-B or Gly-[Gly/Ala]-Arg-B-(Aax)_n.

In certain embodiments, the linker comprising one or more linking moiety B is selected from a group consisting of: GDC, GRCD, GRDC, GGDC, GGCD, GGEC, GGK(N₃)D, GGRCD, GGGDC, GC, GRC, GGRC, GRAC, GARC, GGHK(N₃), GGK(N₃)RC, GARK(N₃) and GGARK(N₃). In a preferred embodiment, the linker comprising one or more linking moiety B is selected from a group consisting of: GGK(N₃)D, GGRCD, GC, GRC, GGRC, GARC, GGK(N₃)RC, GARK(N₃) and GGARK(N₃). In a more preferred embodiment, the linker comprising one or more linking moiety B is selected from a group consisting of: GGRCD, GC, GGRC, GARC, GGK(N₃)RC, GARK(N₃) and GGARK(N₃). In a most preferred embodiment, the linker comprising one or more linking moiety B is selected from a group consisting of: GC, GGRC, GARC and GGARK(N₃). In certain embodiments, the linker comprising one or more linking moiety B is GGGK(N₃).

According to one further embodiment of the invention, the antibody comprises the Asn residue N297 (EU numbering) in the C_H2 domain of the antibody.

According to one further embodiment of the invention, the N297 residue is glycosylated.

According to one further embodiment of the invention, the linker including the payload or linking moiety B is conjugated to the amide side chain of the Gln residue. That is, the amide side chain of the Gln residue of the antibody is conjugated to the N-terminal amino group of the linker via an isopeptide bond.

According to one further embodiment of the invention, the microbial transglutaminase is derived from a *Streptomyces* species, in particular from *Streptomyces mobaraensis*, preferentially with a sequence identity of 80% to the native enzyme. Accordingly, the MTG may be a native enzyme or may be an engineered variant of a native enzyme. As shown in

Fig.8, high conjugation efficiencies have been obtained with a native MTG variant that has not been optimized for the conjugation of glycosylated antibodies.

One such microbial transglutaminase is commercially available from Zedira (Germany). It is recombinantly produced in *E. coli*. *Streptomyces mobaraensis* transglutaminase has an amino acid sequence as disclosed in SEQ ID NO 48. *S. mobaraensis* MTG variants with other amino acid sequences have been reported and are also encompassed by this invention (SEQ ID NO:28 and 49).

In another embodiment, a microbial transglutaminase *Streptomyces ladakanum* (formerly known as *Streptoverticillium ladakanum*) is being used. *Streptomyces ladakanum* transglutaminase (US Pat No US 6,660,510 B2) has an amino acid sequence as disclosed in SEQ ID NO 27.

Both the above transglutaminases can be sequence modified. In several embodiments, transglutaminases can be used which have 80 % or more sequence identity with SEQ ID NOs 27, 28, 48 and 49.

Another suitable microbial transglutaminase is commercially from Ajinomoto, called ACTIVA TG. In comparison to the transglutaminase from Zedira, ACTIVA TG lacks 4 N terminal amino acids, but has similar activity.

Further microbial transglutaminases which can be used in the context of the present invention are disclosed in Kieliszek and Misiewicz 2014, WO 2015/191883 A1, WO 2008/102007 A1 and US 2010/0143970, the content of which is fully incorporated herein by reference.

In certain embodiments, a mutant variant of a microbial transglutaminase is used for the conjugation of a linker to an antibody. That is, the microbial transglutaminase that is used in the method of the present invention may be a variant of *S. mobaraensis* transglutaminase as set forth in SEQ ID NOs: 27 or 29. In certain embodiments, the recombinant *S. morabaensis* transglutaminase as set forth in SEQ ID NO:29 comprises the mutation G250D. In certain embodiments, the recombinant *S. morabaensis* transglutaminase as set forth in SEQ ID NO:29 comprises the mutations G250D and E300D. In certain embodiments, the recombinant *S. morabaensis* transglutaminase as set forth in SEQ ID NO:29 comprises the mutations D4E

and G250D. In certain embodiments, the recombinant *S. morabaensis* transglutaminase as set forth in SEQ ID NO:29 comprises the mutations E120A and G250D. In certain embodiments, the recombinant *S. morabaensis* transglutaminase as set forth in SEQ ID NO:29 comprises the mutations A212D and G250D. In certain embodiments, the recombinant *S. morabaensis* transglutaminase as set forth in SEQ ID NO:29 comprises the mutations G250D and K327T.

Microbial transglutaminase may be added to the conjugation reaction at any concentration that allows efficient conjugation of an antibody with a linker. In certain embodiments, microbial transglutaminase may be added to the conjugation reaction at a concentration of less than 100 U/mL, 90 U/mL, 80 U/ml, 70 U/mL, 60 U/mL, 50 U/mL, 40 U/mL, 30 U/mL, 20 U/mL, 10 U/mL or 7 U/mL.

The method according to the invention comprises the use of a microbial transglutaminase. However, it is to be noted that an equivalent reaction may be carried out by an enzyme comprising transglutaminase activity that is of a non-microbial origin. Accordingly, also the antibody-payload conjugates according to the invention may be generated with an enzyme comprising transglutaminase activity that is of a non-microbial origin.

To obtain efficient conjugation, it is preferred that the linker is added to the antibody in molar excess. That is, in certain embodiments, the antibody is mixed with at least 5, 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 molar equivalents excess of peptide linker versus the antibody.

The method according to the invention is preferably carried out at a pH ranging from 6 to 8.5. Examples 1 and 2 show that the conjugation efficiency is highest at an pH of 7.6. Thus, in a preferred embodiment, the invention relates to a method according to the invention, wherein the conjugation of the linker to the antibody is achieved at a pH ranging from 6 to 8.5, more preferably at a pH ranging from 7 to 8. In a most preferred embodiment, the invention relates to a method according to the invention, wherein the conjugation of the linker to the antibody is achieved at pH 7.6.

The method of the invention may be carried out in any buffer that is suitable for the conjugation of a linker or linker-payload construct to an antibody with the method of the invention. Buffers that are suitable for the method of the invention include, without limitation, Tris, MOPS, HEPES, PBS or BisTris. Further, the buffer may comprise any salt concentration

that is suitable for carrying out the method of the invention. For example, the buffer used in the method of the invention may have a salt concentration ≤ 150 mM, ≤ 140 mM, ≤ 130 mM, ≤ 120 mM, ≤ 110 mM, ≤ 100 mM, ≤ 90 mM, ≤ 80 mM, ≤ 70 mM, ≤ 60 mM, ≤ 50 mM, ≤ 40 mM, ≤ 30 mM, ≤ 20 mM, ≤ 10 mM or 1 mM. In certain embodiments, the buffer may be salt free.

It is to be noted that the optimal reaction conditions (e.g. pH, buffer, salt concentration) may vary between payloads and to some degree depend on the physicochemical properties of the linkers and/or payloads. However, no undue experimentation is required by the skilled person to identify reaction conditions that are suitable for carrying out the method of the invention.

According to one further embodiment of the invention, the linking moiety B is at least one selected from the group consisting of

- bioorthogonal marker group, or
- other non-bio-orthogonal entities for crosslinking

In certain embodiments of the invention, the linking moiety B comprises

- a bioorthogonal marker group, or
- a non-bio-orthogonal entity for crosslinking.

The term “bioorthogonal marker group” has been established by Sletten and Bertozzi (2011) to designate reactive groups that can lead to chemical reactions to occur inside of living systems without interfering with native biochemical processes. A “non-bio-orthogonal entity for crosslinking” may be any molecule that comprises or consists of a first functional group, wherein the first functional group can be chemically or enzymatically crosslinked to a payload comprising a compatible second functional group. Even in cases where the crosslinking reaction is a non-bio-orthogonal reaction, it is preferred that the reaction does not introduce additional modifications to the antibody other than the crosslinking of the payload to the linker. In view of the above, the linking moiety B may either consist of the “bioorthogonal marker group” or the “non-bio-orthogonal entity” or may comprise the “bioorthogonal marker group” or the “non-bio-orthogonal entity”. For example, in case of the



linking moiety Lys(N₃), both the entire Lys(N₃) and the azide group alone may be seen as a bioorthogonal marker group within the present invention.


According to one further embodiment of the invention, the bioorthogonal marker group or the non-bio-orthogonal entity is at least one selected from the group consisting of:

- -N-N≡N, or -N₃
- Lys(N₃)
- Tetrazine
- Alkyne
- DBCO
- BCN
- Norborene
- Transcyclooctene
- -RCOH (aldehyde),
- Acyltrifluoroborates,
- -SH, and
- Cysteine

These groups can for example engage in any of the binding reactions shown in table 4:

Table 4

binding partner 1	binding partner 2	reaction type
-N-N≡N	cyclooctyne derivatives (e.g. DIFO, BCN, DIBAC, DIBO, ADIBO/DBCO)	SPAAC
-N-N≡N	Alkyne	CuAAC
-N-N≡N	Triarylphosphines	Staudinger ligation
tetrazine	Cyclopropene Norborene Cyclooctyne (BCN)	tetrazine ligation
-SH, e.g., of a Cys residue	Maleimide	Thiol-Maleimide conjugation
Amine	N-hydroxysuccinimid	
-O-carbamoylhydroxylamines 	Acyltrifluoroborates 	KAT-ligation (potassium acyl-trifluoroborate)

$R_x-S-S-R_y$	R_z-SH + reducing agent (e.g. TCEP, DTT)	Direct disulfide bioconjugation
-CHO (aldehyde)	HIPS-probe 	Hydrazino-iso-Pictet-Spengler (HIPS)
-CHO (aldehyde)	<i>N</i> -pyrrolyl alanine derivative	pyrrolyl alanine Pictet-Spengler (PAPS)
-CHO (aldehyde)	$R_1-N-N-R_2$ $HO-N-R_1$ $H_2N-CHR_1-CH_2-SH$	Hydrazone-ligation Oxime-ligation Thiazolidine-Ligation
maleimide	-SH, e.g., of a Cys residue	Thiol-Maleimide conjugation

In the above table 4, the said linking moieties can either be or comprise what is called therein “binding partner 1” or “binding partner 2”.

According to one further embodiment of the invention, the linking moiety B is a Cys residue with a free sulfhydryl group.

The free sulfhydryl group of such Cys residue (or derivative) can be used to conjugate a maleimide-comprising linker toxin construct thereto. See Fig. 5 for some more details of the conjugation reaction, and some potential linker constructs.

Toxins comprising a maleimide linker have frequently been used, and also approved by medical authorities, like Adcetris. Thus drugs comprising a MMAE toxin are conjugated to a linker comprising (i) a *p*-aminobenzyl spacer, (ii) a dipeptide and (iii) a maleimidocaproyl linker, which enables the conjugation of the construct to the free sulfhydryl group of a Cys residue in the antibody.

Providing a Cys-residue in the linker according to the present invention does therefore have the advantage to be able to use off-the-shelf-toxin-maleimide constructs to create antibody-payload conjugates, or, more generally, to be able to fully exploit the advantages of Cys-maleimide binding chemistry. At the same time, off-the-shelf antibodies can be used, which do not have to be deglycosylated.

In specific embodiments, the Cys residue is C-terminal, or intrachain in the peptide linker.

In another embodiment, the linking moiety B comprises an azide group. The skilled person is aware of molecules comprising an azide group which may be incorporated into a linker according to the invention, such as 6-azido-lysine (Lys(N₃)) or 4-azido-homoalanine (Xaa(N₃)). Linking moieties comprising an azide group may be used as substrates in various bio-orthogonal reactions, such as strain-promoted azide-alkyne cycloaddition (SPAAC), copper-catalyzed azide-alkyne cycloaddition (CuAAC) or Staudinger ligation. For example, in certain embodiments, payloads comprising a cyclooctene derivative, such as DBCO, may be coupled to a linker comprising an azide group by SPAAC (see FIG.15).

In yet another embodiment, the linking moiety B comprises a tetrazine. The skilled person is aware of tetrazine-comprising molecules which may be incorporated into a linker according to the invention, preferably amino acid derivatives comprising a tetrazine group (see for example Fig.7A). Linking moieties comprising a tetrazine may be used as substrates in a bio-orthogonal tetrazine ligation. For example, in certain embodiments, payloads comprising a cyclopropene, a norbornene or a cyclooctyne group, such as bicyclo[6.1.0]nonyne (BCN), may be coupled to a linker comprising a tetrazine group.

The invention further encompasses linkers comprising two different bio-orthogonal marker groups and/or non-bio-orthogonal entities. For example, a linker according to the invention may comprise an azide-comprising linking moiety, such as Lys(N₃) or Xaa(N₃), and a sulfhydryl-comprising linking moiety, such as cysteine. In certain embodiments, the linker according to the invention may comprise an azide-comprising linking moiety, such as Lys(N₃) or Xaa(N₃), and a tetrazine-comprising linking moiety, such as a tetrazine-modified amino acid. In certain embodiments, the linker according to the invention may comprise a sulfhydryl-comprising linking moiety, such as cysteine, and a tetrazine-comprising linking moiety, such as a tetrazine-modified amino acid. Linkers comprising two different bio-orthogonal marker groups and/or non-bio-orthogonal entities have the advantage that they can accept two distinct payloads and thus result in antibody-payload conjugates comprising more than one payload.

According to one further embodiment of the invention, it is provided that, in case B is a linking moiety, a further step of linking the actual payload to the linking moiety is carried out.

A number of chemical ligation strategies have been developed that fulfill the requirements of bio-orthogonality, including the 1,3-dipolar cycloaddition between azides and cyclooctynes (also termed copper-free click chemistry, Baskin et al (2007)), between nitrones and cyclooctynes (Ning et al (2010)), oxime/hydrazone formation from aldehydes and ketones (Yarema, et al (1998)), the tetrazine ligation (Blackman et al (2008)), the isonitrile-based click reaction (Stöckmann et al (2011)), and most recently, the quadricyclane ligation (Sletten & Bertozzi (JACS, 2011)), Copper(I)-catalyzed azide-alkyne cycloaddition (CuAAC, Kolb & Sharpless (2003)), Strain-promoted azide-alkyne cycloaddition (SPAAC, Agard et al (2004)), or Strain-promoted alkyne-nitrone cycloaddition (SPANAC, MacKenzie et al (2014)).

All these documents are incorporated by reference herein to provide sufficient enabling disclosure, and avoid lengthy repetitions.

It is to be understood that the payload is preferably coupled to the bio-orthogonal marker group or the non-bio-orthogonal entity of the linker according to the invention after said linker has been conjugated to a Gln residue of an antibody by means of a microbial transglutaminase. However, the invention also encompasses antibody-payload conjugates wherein a payload has been coupled to a linker comprising a linking moiety in a first step and wherein the resulting linker-payload construct is conjugated to the antibody by a microbial transglutaminase in a second step.

In a particular embodiment, the invention relates to the method according to the invention, wherein the payload is linked to the linking moiety B of the antibody-linker conjugate via a click-reaction, for example any one of the click reactions mentioned above. In a preferred embodiment, the click reaction is SPAAC.

According to one further embodiment of the invention, the payload B is at least one selected from the group consisting of:

- toxin
- cytokine
- growth factor

- radionuclide
- hormone
- anti-viral agent
- anti-bacterial agent
- fluorescent dye
- immunoregulatory/immunostimulatory agent
- half-life increasing moiety
- solubility increasing moiety
- a polymer-toxin conjugate
- a nucleic acid
- a biotin or streptavidin moiety
- a vitamin
- a target binding moiety, and
- anti-inflammatory agent.

Half-life increasing moieties are, for example, PEG-moieties (polyethylenglycol moieties; PEGylation), other polymer moieties, PAS moieties (oligopeptides comprising Proline, Alanine and Serine; PASylation), or Serum albumin binders. Solubility increasing moieties are, for example PEG-moieties (PEGylation) or PAS moieties (PASylation).

Polymer-toxin conjugates are polymers that are capable of carrying many payload molecules. Such conjugates are sometimes also called fleximers, as e.g. marketed by Mersana therapeutics.

One example of a nucleic acid payload is MCT-485, which is a very small non-coding double stranded RNA which has oncolytic and immune activating properties, developed by MultiCell Technologies, Inc.

Anti-inflammatory agents are for example anti-inflammatory cytokines, which, when conjugated to a target specific antibody, can ameliorate inflammations caused, e.g., by autoimmune diseases.

The term "fluorescent dye" as used herein refers to a dye that absorbs light at a first wavelength and emits at second wavelength that is longer than the first wavelength. In certain embodiment, the fluorescent dye is a near-infrared fluorescent dye, which emits light at a wavelength between 650 and 900 nm. In this region, tissue autofluorescence is lower, and less fluorescence extinction enhances deep tissue penetration with minimal background interference. Accordingly, near-infrared fluorescent imaging may be used to make tissues that are bound by the antibody-payload conjugate of the invention visible during surgery. "Near-infrared fluorescent dyes" are known in the art and commercially available. In certain embodiments, the near-infrared fluorescent dye may be IRDye 800CW, Cy7, Cy7.5, NIR CF750/770/790, DyLight 800 or Alexa Fluor 750.

The term "radionuclide", as used herein, relates to medically useful radionuclides, including, for example, positively charged ions of radiometals such as Y, In, Tb, Ac, Cu, Lu, Tc, Re, Co, Fe and the like, such as ^{90}Y , ^{111}In , ^{67}Cu , ^{77}Lu , ^{99}Tc , ^{161}Tb , ^{225}Ac and the like. The radionuclide may be comprised in a chelating agent. Further, the radionuclide may be a therapeutic radionuclide or a radionuclide that can be used as contrast agent in imaging techniques as discussed below. Radionuclides or molecules comprising radionuclides are known in the art and commercially available.

The term "toxin" as used herein relates to any compound that is poisonous to a cell or organism. A toxin, thus can be, e.g. small molecules, nucleic acids, peptides, or proteins. Specific examples are neurotoxins, necrotoxins, hemotoxins and cytotoxins. According to one further embodiment of the invention, the toxin is at least one selected from the group consisting of

- Pyrrolbenzodiazepines (PBD)
- Auristatins (e.g., MMAE, MMAF)
- Maytansinoids (Maytansine, DM1, DM4, DM21)
- Duocarmycins
- Tubulysins
- Eneidyenes (e.g. Calicheamicin)
- PNUs, doxorubicins
- Pyrrole-based kinesin spindle protein (KSP) inhibitors

- Calicheamicins
- Amanitins (e.g. α -Amanitin), and/or
- Camptothecins (e.g. exatecans, deruxtecans)

In certain embodiments, the payload is an auristatin. As used herein, the term "auristatin" refers to a family of anti-mitotic agents. Auristatin derivatives are also included within the definition of the term "auristatin". Examples of auristatin include, but are not limited to, synthetic analogues of auristatin E (AE), monomethyl auristatin E (MMAE), monomethyl auristatin F (MMAF) and dolastatin.

In certain embodiments, the payload is a maytansinoid. In the context of the present invention, the term "maytansinoid" refers to a class of highly cytotoxic drugs originally isolated from the African shrub *Maytenus ovatus* and further maytansinol (Maytansinol) and C-3 ester of natural maytansinol (US Pat. No. 4,151,042); C-3 ester analog of synthetic maytansinol (Kupchan et al., J. Med. Chem. 21: 31-37, 1978; Higashide et al., Nature 270: 721-722, 1977; Kawai et al., Chem. Farm. Bull. 32: 3441-3451; and US Pat. No. 5,416,064); C-3 esters of simple carboxylic acids (US Pat. 4,248,870; 4,265,814; 4,308,268; 4,308,269; 4,309,428; 4,317,821; 4,322,348; and 4,331,598); and C-3 esters with derivatives of N-methyl-L-alanine (U.S. Pat. Nos. 4,137,230; 4,260,608; and Kawai et al., Chem. Pharm Bull. 12: 3441, 1984). Exemplary maytansinoids that may be used in the method of the invention or that may be comprised in the antibody-payload conjugate of the invention are DM1, DM3, DM4 and/or DM21.

In certain embodiments, the toxic payload molecule is duocarmycin. Suitable duocarmycins may be e.g. duocarmycin A, duocarmycin B1, duocarmycin B2, duocarmycin C1, duocarmycin C2, duocarmycin D, duocarmycin SA, duocarmycin MA, and CC-1065. The term "duocarmycin" should be understood as referring also to synthetic analogs of duocarmycins, such as adozelesin, bizelesin, carzelesin, KW-2189 and CBI-TMI.

The toxin, in the sense of the present invention may also be an inhibitor of a drug efflux transporter. Antibody-payload conjugates comprising a toxin and an inhibitor of a drug efflux transporter may have the advantage that, when internalized into a cell, the inhibitor of the drug efflux transporter prevents efflux of the toxin out of the cell. Within the present invention, the drug efflux transporter may be P-glycoprotein. Some common pharmacological

inhibitors of P-glycoprotein include: amiodarone, clarithromycin, ciclosporin, colchicine, diltiazem, erythromycin, felodipine, ketoconazole, lansoprazole, omeprazole and other proton-pump inhibitors, nifedipine, paroxetine, reserpine, saquinavir, sertraline, quinidine, tamoxifen, verapamil, and duloxetine. Elacridar and CP 100356 are other common P-gp inhibitors. Zosuquidar and tariquidar were also developed with this in mind. Lastly, valspodar and reversan are other examples of such agents.

The vitamin can be selected from the group consisting of folates, including folic acid, folacin, and vitamin B9.

The target binding moiety can be a protein or small molecule being capable of specifically binding to a protein or non-protein target. In one embodiment, such target binding moiety is a scFv shaped antibody, a Fab fragment, a F(ab)₂ fragment, a nanobody, affibody, a diabody, a VHH shaped antibody, or an antibody mimetic, including a DARPIN.

It is to be understood that the payload can be coupled to a linking moiety of a linker by any suitable reaction, such as a click reaction, or may be attached to the linker by chemical synthesis.

According to one further embodiment of the invention, the linker has two or more linking moieties B.

In such embodiment, an antibody-payload conjugate can be created with, for example, an antibody to payload ratio of 4, with two payloads conjugated to each Q295 residue.

According to one further embodiment of the invention, the two or more linking moieties B differ from one another.

In such embodiment, a first linking moiety could for example be or comprise an azide (N₃), while a second linking moiety could be or comprise a tetrazine. Such oligopeptide linker thus allows to conjugate two different payloads to two Gln residues of the antibody, i.e., the Q295 residues of the two C_H2 domains of the antibody.

In such way, an antibody payload ratio of 2+2 can be obtained. Using a second payload allows for the development of a completely new class of antibody payload conjugates that go beyond current therapeutic approaches with respect to efficacy and potency.


Such embodiment allows, *inter alia*, to target two different structures in a cell, like, e.g., the DNA and microtubule. Because some cancers can be resistant to one drug, like e.g., a microtubule toxin, the DNA-toxin can still kill the cancer cells.

According to another embodiment, two drugs could be used that are only fully potent when they are released at the same time and in the same tissue. This may lead to reduced off-target toxicity in case the antibody is partially degraded in healthy tissues or one drug is prematurely lost.

Furthermore, dual-labeled probes can be used for non-invasive imaging and therapy or intra/post-operative imaging/surgery. In such embodiment, a tumor patient can be selected by means of the non-invasive imaging. Then, the tumor can be removed surgically using the other imaging agent (e.g., a fluorescent dye), which helps the surgeon or robot to identify all cancerous tissue.

According to another aspect of the invention, an antibody-payload conjugate is provided which has been generated with a method according to any one of the aforementioned steps.

According to another aspect of the invention, a linker is provided comprising the peptide structure (shown in N->C direction)



wherein Gly comprises an N-terminal primary amine, and wherein

- m is an integer between ≥ 0 and ≤ 12
- n is an integer between ≥ 0 and ≤ 12
- $m + n \geq 0$,
- Aax is an amino acid or an amino acid derivative, and
- B is a payload or a linking moiety,

and wherein the linker can be conjugated to an antibody by a microbial transglutaminase via the N-terminal primary amine of the N-terminal Gly of the linker.


Said linker is suitable to be conjugated, via the N-terminal primary amine of the N-terminal glycine (Gly) residue, to a glutamine (Gln) residue comprised in the heavy or light chain of an antibody, by means of a transglutaminase enzyme.

Generally, the advantages and embodiments discussed above in accordance with the method of the present invention do also apply to this aspect. i.e., the linker as composition of matter. Hence, those embodiments shall be deemed disclosed also with the linker as composition of matter.

It is important to understand that in different linker peptides shown herein, the C-terminus may or may not be protected, even if shown otherwise. Protection can be accomplished by amidation. In the context of the present invention, linker peptides that are protected and unprotected at the C-terminus are encompassed.

In a particular embodiment, the invention relates to the linker according to the invention, wherein the linker comprises two or more payloads and/or linking moieties B.

In certain embodiments, the linker may comprise two or more linking moieties and/or payloads. That is the linker may have the peptide structure (shown in N -> C direction)




wherein

- m, n and o are integers between ≥ 0 and ≤ 12 ,
- $m + n + o \geq 0$,
- Aax is an amino acid or an amino acid derivative, and
- B₁ and B₂ are payloads and/or linking moieties, wherein B₁ and B₂ may be identical or different from each other

and wherein the linker can be conjugated to an antibody by a microbial transglutaminase via the N-terminal primary amine of the N-terminal Gly of the linker.

In other embodiments, the linker may comprise three linking moieties and/or payloads. That is the linker may have the peptide structure (shown in N → C direction)




wherein

- m, n, o and p are integers between ≥ 0 and ≤ 12 ,
- $m + n + o + p \geq 0$,
- Aax is an amino acid or an amino acid derivative, and
- B₁, B₂ and B₃ are payloads and/or linking moieties, wherein B₁, B₂ and B₃ may be identical or different from each other

and wherein the linker can be conjugated to an antibody by a microbial transglutaminase via the N-terminal primary amine of the N-terminal Gly of the linker.

It is to be understood, that the invention also encompasses linkers comprising more than three linking moieties and/or payloads, such as 4, 5 or 6 linking moieties and/or payloads. In this case, the peptide structure of the linkers follows the same pattern as described above for the linkers comprising 2 or 3 linking moieties and/or payloads.

In certain embodiments, the invention relates to a linker having the peptide structure (shown in N→C direction)



wherein

- m is an integer between ≥ 0 and ≤ 12
- n is an integer between ≥ 0 and ≤ 12
- $m + n \geq 0$,
- Aax is an amino acid or an amino acid derivative, and

- B is a payload or a linking moiety,

and wherein the linker can be conjugated to an antibody by a microbial transglutaminase via the N-terminal primary amine of the N-terminal Gly of the linker.

In this case, it is to be understood that the moiety B may comprise more than one payload and/or linking moiety. For example, B may stand for $(B'-(Aax)_o-B'')$, wherein B' and B'' are payloads and/or linking moieties and wherein o is an integer between ≥ 0 and ≤ 12 . Alternatively, B may stand for $(B'-(Aax)_o-B''-(Aax)_p-B''')$, wherein B', B'' and B''' are payloads and/or linking moieties and wherein o and p are integers between ≥ 0 and ≤ 12 .

In preferred embodiments, m and/or n is $\geq 1, \geq 2, \geq 3, \geq 4, \geq 5, \geq 6, \geq 7, \geq 8, \geq 9, \geq 10$, or ≥ 11 . In other preferred embodiments, m and/or n is $\leq 12, \leq 11, \leq 10, \leq 9, \leq 8, \leq 7, \leq 6, \leq 5, \leq 4, \leq 3, \leq 2$, or ≤ 1 . In further preferred embodiments, m + n is $\geq 1, \geq 2, \geq 3, \geq 4, \geq 5, \geq 6, \geq 7, \geq 8, \geq 9, \geq 10$, or ≥ 11 . In still further preferred embodiments m + n is $\leq 12, \leq 11, \leq 10, \leq 9, \leq 8, \leq 7, \leq 6, \leq 5, \leq 4, \leq 3, \leq 2$, or ≤ 1 .

Members of both ranges can be combined with another to disclose a preferred length range with lower and upper limit.

Accordingly, in a particular embodiment, the invention relates to a linker according to the invention, wherein m+n is $\leq 12, \leq 11, \leq 10, \leq 9, \leq 8, \leq 7, \leq 6, \leq 5$ or ≤ 4 .

In further embodiments, the linker is not cleavable by cathepsin B, and/or the linker does not comprise a valine-alanine motif or a valine-citrulline motif, and/or the linker does not comprise Polyethylenglycol or a Polyethylenglycol derivative.

According to one embodiment, the linking moiety B is at least one selected from the group consisting of

- bioorthogonal marker group
- other non-bio-orthogonal entities for crosslinking.

In certain embodiments, at least one linking moiety B of the linker comprises or consists of

- a bioorthogonal marker group; or
- a non-bio-orthogonal entity for crosslinking.

According to one embodiment, the bioorthogonal marker group or the non-bio-orthogonal entity is at least one selected from the group consisting of

- $-N-N\equiv N$, or $-N_3$
- Lys(N_3)
- tetrazine
- alkyne
- DBCO
- BCN
- norborene
- transcyclooctene
- $-RCOH$ (aldehyde),
- acyltrifluoroborates,
- $-SH$, and
- cysteine.

In further embodiments, the net charge of the linker is neutral or positive, and/or the linker does not comprise negatively charged amino acid residues, and/or the linker comprises positively charged amino acid residues, and/or the linker comprises at least two amino acid residues selected from the group consisting of

- Lysine,
- Arginine, and/or
- Histidine.

In certain embodiments, the linker comprises at least one amino acid residue selected from the group consisting of

- Lysine,

- Arginine, and
- Histidine.

In certain embodiments, the linker comprises at least one amino acid residue selected from the group consisting of

- Arginine, and
- Histidine.

That is, in certain embodiments, the linker according to the invention has a neutral or positive net charge. In certain embodiments, the linker according to the invention has a neutral or positive net charge and comprises at least one arginine and/or histidine residue. In certain embodiments, the linker according to the invention does not comprise a lysine residue. In certain embodiments, the linker has a neutral or positive net charge and does not comprises a lysine residue.

According to one embodiment, the primary amine group is suitable to serve as the substrate of a microbial transglutaminase (MTG).

According to one further embodiment, the linker is suitable for generating an antibody-payload conjugate by means of a microbial transglutaminase (MTG).

According to one further embodiment, the linker is selected from

- a) the list as shown in table 5 and/or
- b) any one of SEQ ID NO 1 – 25

In a particular embodiment, the invention relates to a linker according to the invention, wherein the linker is selected from the list as shown in table 5.

According to yet another aspect of the invention, a linker-payload construct is provided, comprising at least

- a) a linker according to the above description, and

b) one or more payloads,

wherein, in said construct, the linker and/or the payload have optionally been chemically modified during binding to allow covalent or non-covalent binding, to form said construct.

In certain embodiments, a linker-payload construct is provided, comprising at least

a) a linker according to the above description, and

b) one or more payloads,

wherein the one or more payloads are covalently or non-covalently bound to the linker.

In a particular embodiment, the invention relates to the linker-payload construct according to the invention, wherein in said construct, the one or more payloads have been covalently bound to the linking moiety B of the linker with a click reaction. That is, the one or more payloads may be attached to a linking moiety B by any of the click reactions discussed above, such as, without limitation, SPAAC, tetrazine ligation or thiol-maleimide conjugation.

Besides a click reaction between the linking moiety in the linker and the payload, the payload may be covalently bound to the linker by any enzymatic or non-enzymatic reaction known in the art. For that, the payload may be bound to the C-terminus of the linker or to an amino acid side chain of the linker.

In certain embodiments, the payload is coupled to a linker by chemical synthesis. The skilled person is aware of methods to couple a payload to a peptide linker by chemical synthesis. For example, an amine-comprising payload, or a thiol-comprising payload (for e.g. maytansine analogs), or an hydroxyl-containing payload (for e.g. SN-38 analogs) may be attached to the C-terminus of a peptide linker by chemical synthesis to obtain, for example, the linkers shown in Fig.17. However, the skilled person is aware of further reactions and reactive groups that may be utilized for coupling a payload to the C-terminus or the side chain of an amino acid or amino acid derivative by chemical synthesis. Typical reactions that may be used to couple a payload to a peptide linker by chemical synthesis include, without limitation: peptide coupling, activated ester coupling (NHS ester, PFP ester), click reaction (CuAAC, SPAAC), michael addition (thiol maleimide conjugation). The coupling of payloads to peptides has

been extensively described in the prior art, for example by Costoplus et al. (ACS Med Chem, 2019), Sonzini et al. (Bioconj Chem, 2019), Boderio et al. (Belstein, 2018), Nunes et al. (RSC Adv, 2017), Doronina et al. (Bioconj Chem, 2006), Nakada et al. (Bioorg Med Chem, 2016) and Dickgiesser et al. (Bioconj Chem, 2020).

In a particular embodiment, the invention relates to the linker-payload construct according to the invention, wherein in said construct, the linker and/or the payload have optionally been chemically modified during binding to allow covalent or non-covalent binding, to form said construct.

In case two or more payloads are being used, the latter can be identical or different from one another.

In one embodiment, the payload is at least one selected from the group consisting of

- a toxin
- a cytokine
- a growth factor
- a radionuclide
- a hormone
- an anti-viral agent
- an anti-bacterial agent
- a fluorescent dye
- an immunoregulatory/immunostimulatory agent
- a half-life increasing moiety
- a solubility increasing moiety
- a polymer-toxin conjugate
- a nucleic acid
- a biotin or streptavidin moiety
- a vitamin
- a target binding moiety, and
- an anti-inflammatory agent.
- a protein degrader (PROTACs)

In another embodiment, the toxin is at least one selected from the group consisting of

- pyrrolobenzodiazepines (PBD)
- auristatins (e.g., MMAE, MMAF)
- maytansinoids (Maytansine, DM1, DM4, DM21)
- duocarmycins
- tubulysins
- enediyenes (e.g. Calicheamicin)
- PNUs, doxorubicins
- pyrrole-based kinesin spindle protein (KSP) inhibitors
- calicheamicins
- amanitins (e.g. α -amanitin), and/or
- camptothecins (e.g. exatecans, deruxtecans)

According to another aspect of the invention, an antibody-payload conjugate is provided comprising

- a) one or more linker-payload constructs according to the above description, and
- b) an antibody comprising at least one Gln residue in the heavy or light chain,

wherein, in said conjugate, the linker-payload constructs and/or the antibody have optionally been chemically modified during conjugation to allow covalent or non-covalent conjugation, to form said conjugate.

In a particular embodiment, the invention relates to an antibody payload conjugate comprising

- a) one or more linker-payload constructs according to the above description, and
- b) an antibody comprising at least one Gln residue in the heavy or light chain,

wherein, the linker-payload construct is conjugated to the amide side chain of a Gln residue in the heavy or light chain of the antibody via an N-terminal primary amine of the N-terminal glycine residue comprised in the linker-payload construct

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the conjugation has been achieved with a microbial transglutaminase (MTG).

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the conjugation has been achieved before or after formation of the linker-payload construct. That is, the invention encompasses antibody-payload conjugates wherein the linkers have been conjugated to the antibody in a first step before the one or more payloads are coupled to the linking moieties of the linkers in a second step. However, the invention also encompasses antibody-payload conjugates, wherein the one or more payloads are coupled to the linking moieties of the linkers in a first step, and wherein the resulting linker-payload constructs are then conjugated to the antibody in a second step. Further, the one or more payloads may be attached to a peptide linker by means of chemical synthesis and the resulting linker-payload construct may then be conjugated to the antibody in a one-step reaction.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody is an IgG, IgE, IgM, IgD, IgA or IgY antibody, or a fragment or recombinant variant thereof, wherein the fragment or recombinant variant thereof retains target binding properties and comprises a C_H2 domain.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody is an IgG antibody.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody is a glycosylated antibody, a deglycosylated antibody or an aglycosylated antibody.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the glycosylated antibody is an IgG antibody that is glycosylated at residue N297(EU numbering) of the C_H2 domain or wherein the glycosylated antibody is an antibody of a different isotype that is glycosylated at a residue that is homologous to residue N297 (EU numbering) of the C_H2 domain of an IgG antibody.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein (a) the linker-payload construct is conjugated to a Gln residue which has been introduced into the heavy or light chain of the antibody by molecular engineering or (b) the linker-payload construct is conjugated to a Gln residue in the Fc domain of the antibody.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the Gln residue in the Fc domain of the antibody is Gln residue Q295 (EU numbering) of the C_H2 domain of an IgG antibody or a homologous Gln residue of an antibody of a different isotype.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the Gln residue in the Fc domain of the antibody is Gln residue Q295 (EU numbering) of the C_H2 domain of an IgG antibody that is glycosylated at residue N297(EU numbering) of the C_H2 domain.

The antibody of the method or the antibody-payload conjugate of the invention may be any antibody, preferably any IgG type antibody. For example, the antibody may be, without limitation Brentuximab, Trastuzumab, Gemtuzumab, Inotuzumab, Avelumab, Cetuximab, Rituximab, Daratumumab, Pertuzumab, Vedolizumab, Ocrelizumab, Tocilizumab, Ustekinumab, Golimumab, Obinutuzumab, Polatuzumab or Enfortumab.

That is, in certain embodiments, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Brentuximab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Trastuzumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Gemtuzumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Inotuzumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Avelumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Cetuximab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the

antibody is Rituximab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Daratumumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Pertuzumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Vedolizumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Ocrelizumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Tocilizumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Ustekinumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Golimumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Obinutuzumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Polatuzumab. In a further embodiment, the invention relates to an antibody-payload conjugate according to the invention, wherein the antibody is Enfortumab.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the Gln residue that has been introduced into the heavy or light chain of the antibody by molecular engineering is N297Q (EU numbering) of the C_H2 domain of an aglycosylated IgG antibody.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the Gln residue that has been introduced into the heavy or light chain of the antibody by molecular engineering is comprised in a peptide that has been (a) integrated into the heavy or light chain of the antibody or (b) fused to the N- or C-terminal end of the heavy or light chain of the antibody.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the peptide comprising the Gln residue has been fused to the C-terminal end of the heavy chain of the antibody.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the peptide comprising the Gln residue is selected from a group consisting of: LLQGG, LLQG, LSLSQG, GGGLLQGG, GLLQG, LLQ, GSPLAQSHGG, GLLQGGG, GLLQGG, GLLQ, LLQLLQGA, LLQGA, LLQYQGA, LLQGSG, LLQYQG, LLQLLQG, SLLQG, LLQLQ, LLQLLQ, LLQGR, EEQYASTY, EEQYQSTY, EEQYNSTY, EEQYQS, EEQYQST, EQYQSTY, QYQS, QYQSTY, YRYRQ, DYALQ, FGLQRPY, EQKLISEEDL, LQR and YQR.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody-payload conjugate comprises at least one toxin.

That is, the antibody-payload conjugate of the invention comprises an antibody that is conjugated to at least one linker, wherein the one linker comprises at least one toxin. In certain embodiments, the antibody-payload conjugate comprises two linkers, wherein each heavy chain of the antibody is conjugated to one linker, respectively. In certain embodiments, the antibody-payload conjugate comprises four linkers, wherein each heavy chain of the antibody is conjugated to two linkers, respectively. In such cases, each linker may contain one or more payloads, such as toxins.

In certain embodiments, the antibody-payload conjugate according to the invention comprises two linkers, wherein each linker comprises one payload, for example a toxin. In other embodiments, the antibody-payload conjugate according to the invention comprises two linkers, wherein each linker comprises two payloads, for example one toxin and one other payload or two identical or different toxins. In embodiments where the antibody-payload conjugate comprises two linkers, it is preferred that the linkers are conjugated to residue Q295 of the two heavy chains of an IgG antibody. Even more preferably, the antibody is an IgG antibody that is glycosylated at residue N297.

In certain embodiments, the antibody-payload conjugate according to the invention comprises four linkers, wherein each linker comprises one payload, for example a toxin. In other embodiments, the antibody-payload conjugate according to the invention comprises four linkers, wherein each linker comprises two payloads, for example one toxin and one other payload or two identical or different toxins. In embodiments where the antibody-payload

conjugate comprises four linkers, it is preferred that the linkers are conjugated to residues Q295 and N297Q of the two heavy chains of an IgG antibody.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody-payload conjugate comprises two different toxins.

In certain embodiments, the antibody-payload conjugate according to the invention comprises two different toxins. That is, in certain embodiments, the antibody-payload conjugate may comprise two linkers, wherein each linker comprises two different toxins. Antibody-payload conjugates comprising two different toxins have the advantage that they may have increased cytotoxic activity. Such increased cytotoxic activity may be achieved by combining two toxins that target two different cellular mechanisms. For example, the antibody-payload conjugates according to the invention may comprise a first toxin that inhibits cell division and a second toxin is a toxin that interferes with replication and/or transcription of DNA.

Accordingly, in a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein a first toxin is a toxin that inhibits cell division and a second toxin is a toxin that interferes with replication and/or transcription of DNA.

A toxin that inhibits cell division, such as an anti-mitotic agent or a spindle poison, is an agent that has the potential to inhibit or prevent mitotic division of a cell. A spindle poison is a poison that disrupts cell division by affecting the protein threads that connect the centromere regions of chromosomes, known as spindles. Spindle poisons effectively cease the production of new cells by interrupting the mitosis phase of cell division at the spindle assembly checkpoint (SAC). The mitotic spindle is composed of microtubules (polymerized tubulin) that aid, along with regulatory proteins; each other in the activity of appropriately segregating replicated chromosomes. Certain compounds affecting the mitotic spindle have proven highly effective against solid tumors and hematological malignancies.

Two specific families of antimitotic agents — vinca alkaloids and taxanes — interrupt the cell's division by the agitation of microtubule dynamics. The vinca alkaloids work by causing the inhibition of the polymerization of tubulin into microtubules, resulting in the G2/M arrest within the cell cycle and eventually cell death. In contrast, the taxanes arrest the mitotic cell cycle by stabilizing microtubules against depolymerization. Even though numerous other

spindle proteins exist that could be the target of novel chemotherapeutics, tubulin-binding agents are the only types in clinical use. Agents that affect the motor protein kinesin are beginning to enter clinical trials. Another type, paclitaxel, acts by attaching to tubulin within existing microtubules. Preferred toxins that inhibit cell division within the present invention are auristatins, such as MMAE and MMAF, and maytansinoids, such as DM1, DM3, DM4 and/or DM21.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein at least one of the toxins is an auristatin or a maytansinoid.

Several agents that prevent the correct replication and/or transcription of DNA molecules and have been shown to be suitable in cancer treatment are known to the person skilled in the art. For example, antimetabolites such as nucleotide or nucleoside analogs which are misincorporated into newly formed DNA and/or RNA molecules are known in the art and have been summarized by Tsesmetzis et al, *Cancers (Basel)*, 2018, 10(7): 240. Other toxins that are known to interfere with the replication and/or transcription of DNA are duromycins.

Accordingly, in certain embodiments, the antibody-payload conjugate according to the invention comprises two different toxins, wherein the first toxin is a duromycin and wherein the second payload is an auristatin or a maytansinoid.

In certain embodiments, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody-payload conjugate comprises two different auristatins.

One main advantage of antibody-payload conjugates comprising two different toxins is that the antibody-payload conjugates may still act against target cells that have escaped the mechanism of action of one of the toxins and/or that the antibody-payload conjugate may have a higher efficacy against heterogenous tumors.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody-payload conjugate comprises a toxin and an inhibitor of a drug efflux transporter.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody-payload conjugate comprises a toxin and a solubility increasing moiety.

That is, the antibody-payload conjugate may comprise two payloads, wherein the first payload is a toxin and the second payload is a solubility increasing moiety. Structure 5 in Fig.9 shows a peptide linker comprising a solubility increasing moiety coupled to a lysine side chain. Accordingly, an antibody-payload conjugate comprising a toxin and a solubility increasing moiety may be obtained by clicking a toxin to the azide group of the linker shown in Structure 5 in Fig.9. Alternatively, an antibody-linker conjugate may be obtained by clicking a toxin to an azide-comprising linking moiety of a linker and by clicking a maleimide-comprising solubility increasing moiety to a cysteine side chain of the same linker. Alternatively, the toxin and/or the solubility increasing moiety may be attached to the linker by chemical synthesis.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody-payload conjugate comprises a toxin and an immunostimulatory agent.

As used herein and depending on context, the term “immunostimulatory agent” includes compounds that increase a subject's immune response to an antigen. Examples of immunostimulatory agents include immune stimulants and immune cell activating compounds. Antibody-payload conjugates of the present invention may contain immunostimulatory agents that help program the immune cells to recognize ligands and enhance antigen presentation. Immune cell activating compounds include Toll-like receptor (TLR) agonists. Such agonists include pathogen associated molecular patterns (PAMPs), e.g., an infection-mimicking composition such as a bacterially-derived immunomodulator (a.k.a., danger signal) and damage associated molecular pattern (DAMPs), e.g. a composition mimicking a stressed or damaged cell. TLR agonists include nucleic acid or lipid compositions (e.g., monophosphoryl lipid A (MPLA)). In one example, the TLR agonist comprises a TLR9 agonist such as a cytosine-guanosine oligonucleotide (CpG-ODN), a poly(ethylenimine) (PEI)-condensed oligonucleotide (ODN) such as PEI-CpG-ODN, or double stranded deoxyribonucleic acid (DNA). In another example, the TLR agonist comprises a TLR3 agonist such as polyinosine-polycytidylic acid (poly (I:C)), PEI-poly (I:C),

polyadenylic-polyuridylic acid (poly (A:U)), PEI-poly (A:U), or double stranded ribonucleic acid (RNA). Other exemplary vaccine immunostimulatory compounds include lipopolysaccharide (LPS), chemokines/cytokines, fungal beta-glucans (such as lentinan), imiquimod, CRX-527, and OM-174.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody-payload conjugate comprises two different immunostimulatory agents.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the at least one immunostimulatory agent is a TLR agonist.

The term "TLR agonist", as used herein, refers to a molecule which is capable of causing a signaling response through a TLR signaling pathway, either as a direct ligand or indirectly through generation of endogenous or exogenous. Agonistic ligands of TLR receptors are (i) natural ligands of the actual TLR receptor, or functionally equivalent variants thereof which conserve the capacity to bind to the TLR receptor and induce co-stimulation signals thereon, or (ii) an agonist antibody against the TLR receptor, or a functionally equivalent variant thereof capable of specifically binding to the TLR receptor and, more particularly, to the extracellular domain of said receptor, and inducing some of the immune signals controlled by this receptor and associated proteins. The binding specificity can be for the human TLR receptor or for a TLR receptor homologous to the human one of a different species.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the antibody-payload conjugate comprises a radionuclide and a fluorescent dye.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, wherein the radionuclide is a radionuclide that is suitable for use in tomography, in particular single-photon emission computed tomography (SPECT) or positron emission tomography (PET), and wherein the fluorescent dye is a near-infrared fluorescent dye.

The term "radionuclide" as used herein has the same meaning as radioactive nuclide, radioisotope or radioactive isotope.

The radionuclide is preferably detectable by nuclear medicine molecular imaging technique(s), such as, Positron Emission Tomography (PET), Single Photon Emission Computed Tomography (SPECT), an hybrid of SPECT and/or PET or their combinations. Single Photon Emission Computed Tomography (SPECT) herein includes planar scintigraphy (PS).

An hybrid of SPECT and/or PET is for example SPECT/CT, PET/CT, PET/IRM or SPECT/IRM.

SPECT and PET acquire information on the concentration (or uptake) of radionuclides introduced into a subject's body. PET generates images by detecting pairs of gamma rays emitted indirectly by a positron-emitting radionuclide. A PET analysis results in a series of thin slice images of the body over the region of interest (e.g., brain, breast, liver,...). These thin slice images can be assembled into a three dimensional representation of the examined area. SPECT is similar to PET, but the radioactive substances used in SPECT have longer decay times than those used in PET and emit single instead of double gamma rays. Although SPECT images exhibit less sensitivity and are less detailed than PET images, the SPECT technique is much less expensive than PET and offers the advantage of not requiring the proximity of a particle accelerator. Actual clinical PET presents higher sensitivity and better spatial resolution than SPECT, and presents the advantage of an accurate attenuation correction due to the high energy of photons; so PET provides more accurate quantitative data than SPECT. Planar scintigraphy (PS) is similar to SPECT in that it uses the same radionuclides. However, PS only generates 2D-information.

SPECT produces computer-generated images of local radiotracer uptake, while CT produces 3-D anatomic images of X ray density of the human body. Combined SPECT/CT imaging provides sequentially functional information from SPECT and the anatomic information from CT, obtained during a single examination. CT data are also used for rapid and optimal attenuation correction of the single photon emission data. By precisely localizing areas of abnormal and/or physiological tracer uptake, SPECT/CT improves sensitivity and specificity, but can also aid in achieving accurate dosimetric estimates as well as in guiding interventional

procedures or in better defining the target volume for external beam radiation therapy. Gamma camera imaging with single photon emitting radiotracers represents the majority of procedures.

The radionuclide may be selected in the group consisting of technetium-99m (^{99m}Tc), gallium-67 (^{67}Ga), gallium-68 (^{68}Ga), yttrium-90 (^{90}Y), indium-111 (^{111}In), rhenium-186 (^{186}Re), fluorine-18 (^{18}F), copper-64 (^{64}Cu), terbium-149 (^{149}Tb) or thallium-201 (^{201}Tl). The radionuclide may be comprised in a molecule or bound to a chelating agent.

According to another aspect of the invention, a pharmaceutical composition is provided, the composition comprising the linker according to the above description, the linker-payload construct according to the above description, and/or the antibody-payload conjugate according to the above description.

According to another aspect of the invention, a pharmaceutical product is provided, the product comprising the antibody-payload conjugate according to the above description, or the pharmaceutical composition according to the above description, and at least one further pharmaceutically acceptable carrier.

A pharmaceutically acceptable carrier refers to an ingredient in a pharmaceutical formulation, other than an active ingredient, which is nontoxic to a subject. A pharmaceutically acceptable carrier includes, but is not limited to, a buffer, excipient, stabilizer, or preservative.

Pharmaceutical formulations of the antibody-payload conjugates described herein are prepared by mixing such conjugates having the desired degree of purity with one or more optional pharmaceutically acceptable carriers (Flemington's Pharmaceutical Sciences 16th edition, Oslo, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Pharmaceutically acceptable carriers are generally nontoxic to recipients at the dosages and concentrations employed, and include, but are not limited to: buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride; benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides;

proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn protein complexes); and/or non-ionic surfactants such as polyethylene glycol (PEG). Exemplary pharmaceutically acceptable carriers herein further include interstitial drug dispersion agents such as soluble neutral-active hyaluronidase glycoproteins (sHASEGP), for example, human soluble PH-20 hyaluronidase glycoproteins, such as rHuPH20 (HYLENEX®, Baxter International, Inc.). Certain exemplary sHASEGPs and methods of use, including rHuPH20, are described in US Patent Publication Nos. 2005/0260186 and 2006/0104968. In one aspect, a sHASEGP is combined with one or more additional glycosaminoglycanases such as chondroitinases.

In one embodiment, the invention relates to the antibody-payload conjugate according to the invention, the pharmaceutical composition according to the invention or the pharmaceutical product according to the invention for use in therapy and/or diagnostics.

That is, the antibody-payload conjugates of the invention may be used in the treatment of a subject or in diagnosing a disease or condition in a subject. An individual or subject is a mammal. Mammals include, but are not limited to, domesticated animals (e.g., cows, sheep, cats, dogs, and horses), primates (e.g., humans and non human primates such as macaques), rabbits, and rodents (e.g., mice and rats). In certain embodiments, the individual or subject is a human.

According to another aspect of the invention, the pharmaceutical composition according to the above description or the product according to the above description is provided (for the manufacture of a medicament) for the treatment of a patient

- suffering from,
- being at risk of developing, and/or
- being diagnosed for

a neoplastic disease, neurological disease, an autoimmune disease, an inflammatory disease or an infectious disease, or the prevention or for the prevention of such condition.

Preferably, the invention relates to the antibody-payload conjugate according to the invention, the pharmaceutical composition according to the invention or the pharmaceutical product according to the invention for use in treatment of a patient suffering from a neoplastic disease.

The term “neoplastic disease” as used herein refers to a condition characterized by uncontrolled, abnormal growth of cells. Neoplastic diseases include cancer. Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include breast cancer, prostate cancer, colon cancer, squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, ovarian cancer, cervical cancer, gastrointestinal cancer, pancreatic cancer, glioblastoma, liver cancer, bladder cancer, hepatoma, colorectal cancer, uterine cervical cancer, endometrial carcinoma, salivary gland carcinoma, kidney cancer, vulval cancer, thyroid cancer, hepatic carcinoma, skin cancer, melanoma, brain cancer, ovarian cancer, neuroblastoma, myeloma, various types of head and neck cancer, acute lymphoblastic leukemia, acute myeloid leukemia, Ewing sarcoma and peripheral neuroepithelioma. Preferred cancers include liver cancer, lymphoma, acute lymphoblastic leukemia, acute myeloid leukemia, Ewing sarcoma and peripheral neuroepithelioma.

That is, the antibody-payload conjugates of the invention are preferably used for the treatment of cancer. As such, in certain embodiments, the antibody-payload conjugates comprise an antibody that specifically binds to an antigen that is present on a tumor cell. In certain embodiments, the antigen is an antigen on the surface of a tumor cell. In certain embodiments, the antigen on the surface of the tumor cell is internalized into the cell together with the antibody-payload conjugate upon binding of the antibody-payload conjugate to the antigen.

If the antibody-payload conjugate is used in the treatment of cancer, it is preferred that the antibody-payload conjugate comprises at least one payload that has the potential to kill or inhibit the proliferation of the tumor cell to which the antibody-drug conjugate binds to. In certain embodiments, the at least one payload exhibits its cytotoxic activity after the antibody-payload conjugate has been internalized into the tumor cell. In certain embodiments, the at least one payload is a toxin.

According to another aspect of the invention, a method of treating or preventing a neoplastic disease is provided, said method comprising administering to a patient in need thereof the antibody-payload conjugate according to the above description, the pharmaceutical composition according to the above description, or the product according to the above description.

The inflammatory disease can be an autoimmune disease. The infectious disease can be a bacterial infection or a viral infection.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, the pharmaceutical composition according to the invention or the pharmaceutical product according to the invention for use in pre, intra- and/or postoperative imaging.

That is, the antibody-payload conjugate according to the invention may be used in imaging. For that, the antibody-payload conjugate may be visualized while binding to a specific target molecule, cell or tissue. Different techniques are known in the art to visualize particular payloads. For example, if the payload is a radionuclide, the molecules, cells, or tissues to which the antibody-payload conjugate binds may be visualized by PET or SPECT. If the payload is a fluorescent dye, the molecules, cells, or tissues to which the antibody-payload conjugate binds may be visualized by fluorescence imaging. In certain embodiments, the antibody-payload conjugate according to the invention comprises two different payloads, for example a radionuclide and a fluorescent dye. In this case, the molecule, cell or tissue to which the antibody-payload conjugate binds may be visualized using two different and/or complementary imaging techniques, for example PET/SPECT and fluorescence imaging.

The antibody-payload conjugate may be used for pre- intra- and/or post-operative imaging.

Pre-operative imaging encompasses all imaging techniques that may be performed before a surgery to make specific target molecules, cells or tissues visible when diagnosing a certain disease or condition and, optionally, to provide guidance for a surgery. Preoperative imaging may comprise a step of making a tumor visible by PET or SPECT before a surgery is performed by using an antibody-payload conjugate that comprises an antibody that

specifically binds to an antigen on the tumor and is conjugated to a payload that comprises a radionuclide.

Intra-operative imaging encompasses all imaging techniques that may be performed during a surgery to make specific target molecules, cells or tissues visible and thus provide guidance for the surgery. In certain embodiments, an antibody-payload conjugate comprising a near-infrared fluorescent dye may be used to visualize a tumor during surgery by near-infrared fluorescent imaging. Intraoperative imaging allows the surgeon to identify specific tissues, for example tumor tissue, during surgery and thus may allow complete removal of tumor tissue.

Post-operative imaging encompasses all imaging techniques that may be performed after a surgery to make specific target molecules, cells or tissues visible and to evaluate the result of the surgery. Post-operative imaging may be performed similarly as pre-operative surgery.

In certain embodiments, the invention relates to antibody-payload conjugates comprising two or more different payloads. For example, the antibody-payload conjugate may comprise a radionuclide and a near-infrared fluorescent dye. Such an antibody-payload conjugate may be used for imaging by PET/SPECT and near-infrared fluorescent imaging. The advantage of such an antibody is that it may be used to visualize the target tissue, for example a tumor before and after a surgery by PET or SPECT. At the same time, the tumor may be visualized during the surgery by near-fluorescent infrared imaging.

In a particular embodiment, the invention relates to the antibody-payload conjugate according to the invention, the pharmaceutical composition according to the invention or the pharmaceutical product according to the invention for use in intraoperative imaging-guided cancer surgery.

As mentioned above, the antibody-payload conjugate of the invention may be used to visualize a target molecule, cell or tissue and to guide a surgeon or robot during a surgery. That is, the antibody-payload conjugate may be used to visualize tumor tissue during a surgery, for example by near-infrared imaging and to allow complete removal of the tumor tissue.

Said conjugate or product is administered to the human or animal subject in an amount or

dosage that efficiently treats the disease. Alternatively, a corresponding method of treatment is provided.

An antibody-payload conjugate of the invention can be administered by any suitable means, including parenteral, intrapulmonary, and intranasal, and, if desired for local treatment, intralesional, intrauterine or intravesical administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. Dosing can be by any suitable route, e.g. by injections, such as intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic. Various dosing schedules including but not limited to single or multiple administrations over various time-points, bolus administration, and pulse infusion are contemplated herein.

Antibody-payload conjugates of the invention would be formulated, dosed, and administered in a fashion consistent of the invention would be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The antibody-payload conjugate need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of antibody-payload conjugate present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as described herein, or about from 1 to 99% of the dosages described herein, or in any dosage and by any route that is empirically/clinically determined to be appropriate.

For the prevention or treatment of disease, the appropriate dosage of an antibody-payload conjugate of the invention (when used alone or in combination with one or more other additional therapeutic agents) will depend on the type of disease to be treated, the type of antibody-payload conjugate, the severity and course of the disease, whether the antibody-payload conjugate is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody-payload conjugate, and the discretion of the attending physician. The antibody-payload conjugate is suitably administered to the patient at one time or over a series of treatments.

The following table 5 shows different linkers that can be used in the context of the present invention, and their SEQ ID Numbers. For the avoidance of doubt, if there is a discrepancy with the electronic WIPO ST 25 sequence listing, the sequences of this table are to be deemed the correct ones.

It is important to understand that in some linker peptides shown herein, the moiety at the C-terminus is simply designated as N₃. However, this should be understood as an abbreviation of Lys(N₃). For example, GARK(N₃) or GlyAlaArgLys(N₃) does actually mean GARK₁ (with K₁ = Lys(N₃)).

It is furthermore important to understand that in different linker peptides shown herein, the C-terminus may or may not be protected, even if shown otherwise.

Protection can be accomplished by amidation of the former. In the context of the present invention, both the protected and unprotected linker peptides are encompassed.

For example GARK(N₃) does indeed encompass two variants, with the C-terminus protected or unprotected. On the other hand, GARK(N₃)-COOH for example explicitly specifies a peptide which is not protected, i.e., has an unprotected C terminus.

The following table 5 shows some linkers that are encompassed, and suitable to be used, in the context of the present invention:

Table 5

Three letter code	One letter code	Linking moiety B	peptide length	number of positive amino acids (Lys/Arg/His)
GlyAlaArgLys(N ₃)	GARK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyAlaArgLys(N ₃)Lys(Tetrazine)	GARK ₁ K ₂ with K ₁ = Lys(N ₃) and K ₂ = Lys(tetrazine)	Lys(N ₃) and Lys(tetrazine)	5	1
GlyAlaArgLys(N ₃)Cys	GARK ₁ C with K ₁ = Lys(N ₃)	Lys(N ₃), Cys-	5	1

		SH		
GlyGlyAlaArgLys(N ₃) Lys(N ₃)	GGARK ₁ K ₂ with K _{1,2} = Lys(N ₃)	Lys(N ₃)	6	1
GlyGlyAlaArgLys(N ₃)ArgLys(N ₃)	GGARK ₁ RK ₂ with K _{1,2} = Lys(N ₃)	Lys(N ₃)	7	2
GlyAlaArgXaa(N ₃)	GARX with X = Xaa(N ₃), Xaa is 4-Azido-L-homoalanine	Xaa(N ₃)	4	1
GlyAlaArg[PEG]3(N ₃)	GAR [PEG]3N ₃ , with [PEG]3 = triethylenglycol	N ₃	3	1
GlyAlaArgCys	GARC	Cys-SH	4	1
GlyGlyAlaArgLys(PEG) _n ArgLys(N ₃)	GGARK(PEG) _n RK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	7	2
GlyβAlaArgLys(N ₃)	GβARK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyAlahomoArgLys(N ₃)	GAhRK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyβAlahomoArgLys(N ₃)	GβAhRK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyGlyAlaArgArg-B	GGARR-B	N/A	5	2
GlyArgAlaLys(N ₃)	GRAK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyArgAlaCys	GRAC	Cys-SH	4	1
GlyGlyArgLys(N ₃)	GGRK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyArgLys(N ₃)	GRK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	3	1
GlyGlyArgLys(N ₃)Arg	GGRK ₁ R with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyGlyLys(N ₃)ArgCys	GGK ₁ RC with K ₁ = Lys(N ₃)	Lys(N ₃), Cys-SH	5	1
GlyGlyArgArgLys(N ₃)	GGRRK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	5	2
GlyGlyArgLys(N ₃)Arg	GGRK ₁ R with K ₁ = Lys(N ₃)	Lys(N ₃)	5	2
GlyAlaHisLys(N ₃)	GAHK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyGlyHisLys(N ₃)	GGHK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyCys	GC	Cys-SH	2	0
GlyGlyArgCys	GGRC	Cys-SH	4	1
GlyArgCys	GRC	Cys-SH	3	1
GlyGlyArgLys(N ₃)	GGRK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	4	1
GlyGlyAlaLysLys(N ₃)	GGAKK ₁ with K ₁ = Lys(N ₃)	Lys(N ₃)	5	1

EXAMPLES

While the invention has been illustrated and described in detail in the drawings and foregoing description, such illustration and description are to be considered illustrative or exemplary and not restrictive; the invention is not limited to the disclosed embodiments. Other variations to the disclosed embodiments can be understood and effected by those skilled in the art in

practicing the claimed invention, from a study of the drawings, the disclosure, and the appended claims. In the claims, the word “comprising” does not exclude other elements or steps, and the indefinite article “a” or “an” does not exclude a plurality. The mere fact that certain measures are recited in mutually different dependent claims does not indicate that a combination of these measures cannot be used to advantage. Any reference signs in the claims should not be construed as limiting the scope.

All amino acid sequences disclosed herein are shown from N-terminus to C-terminus; all nucleic acid sequences disclosed herein are shown 5'→3'.

Example 1: Conjugation efficiency

Peptides were used as obtained and dissolved at a suitable stock concentration (e.g. 25mM) following the manufacturers instruction, aliquots were prepared and stored at -20°C. Two antibodies of IgG-subclass (antibody 1: anti Her2 IgG1, antibody 2: anti CD38 IgG1) were modified as follows: 1 mg/mL of non-deglycosylated antibody (~6.67μM) was mixed with 80 molar equivalents of peptide linker (i.e. ~533μM), 6 U/mL MTG and buffer. The reaction mixture was incubated for 20 h at 37°C and then subjected for LC-MS analysis under reducing conditions.

The following table 6 shows the conjugation efficiency of a linker according to the present invention (marked with a (*)) vs other linkers:

Table 6

Linker (one letter code)	pH6	pH7.6	pH8.5
GARK(N3)*	0	77.3	34.2
AARK(N3)	0	10.7	8.1
SARK(N3)	0	5.2	3.6
AGRK(N3)	3.6	22.3	14.7

It is clearly visible that the peptide comprising a N-terminal Gly residue, with no further primary amine except the N-terminal primary amine, has by far the best conjugation

efficiency with the Q295 residue in the glycosylated antibody, even though the other peptides likewise comprise N-terminal primary amines, yet not comprised in a Gly residue

Example 2: Conjugation efficiency

Peptides were used as obtained and dissolved at a suitable stock concentration (e.g. 25mM) following the manufacturers instruction, aliquots were prepared and stored at -20°C. Two antibodies of IgG-subclass (antibody 1: anti Her2 IgG1, antibody 2: anti CD38 IgG1) were modified as follows: 1 mg/mL of non-deglycosylated antibody (~6.67μM) was mixed with 80 molar equivalents of peptide linker (i.e. ~533μM), 6 U/mL MTG and buffer. The reaction mixture was incubated for 20 h at 37°C and then subjected for LC-MS analysis under reducing conditions.

The following table 7 shows the conjugation efficiency of a linker according to the present invention (marked with a (*)) vs another linker βAGARK(N3) is shown in Fig 19 (note that βA designates β-Alanine).

Table 7

Linker (One letter code)	Buffer 1 (pH 6)	Buffer 2 (pH 7.6)	Buffer 3 (pH 8.5)	Buffer 4 (pH 7.6)	Buffer 5 (pH 7.6)
βAGARK(N3)	10	57	0	67	63
GGARK(N3)*	50	84	0	86	85


It is clearly visible that the peptide comprising a N-terminal Gly residue, with no further primary amine except the N-terminal primary amine, has by far the better conjugation efficiency with the Q295 residue in the glycosylated antibody, compared to the structurally similar linker which has an N-terminal β-Ala residue.

Example 3: Cell toxicity assay

Cell lines and culture: MDA-MB-231, and SK-BR-3 were obtained from the American Type Culture Collection (ATCC) and cultured in RPMI-1640 following standard cell-culture protocols.

SK-BR-3 is a breast cancer cell line isolated by the Memorial Sloan-Kettering Cancer Center in 1970 that is used in therapeutic research, especially in context of HER2 targeting. MDA-

MB-231 cells are derived from human breast adenocarcinoma of the "basal" type, and are triple negative (ER, PR and HER2 negative). Adcetris (Brentuximab Vedotin) is a commercially available antibody drug conjugate that targets CD30 and is hence expected to not be active against cells which do not express CD30, e.g., MDA-MB-231, and SK-BR-3. Kadcylla (Trastuzumab emtansine) is a commercially available antibody drug conjugate that targets Her2 and is hence expected to be active against cells which express Her2 (e.g., SK-BR-3), and not active against cells which do not express Her2 (e.g., MDA-MB-231). p684 and p579 are antibody drug conjugates produced with the linker technology as specified herein, with linkers having an N-terminal glycine (GARK(N₃) (P684) and GGARK(N₃) (P579)). To generate the antibody-payload conjugate, a May (Maytansine) molecule coupled to a DBCO group (see below) has been clicked to the azide groups of the linkers. Both conjugates use a non-deglycosylated antibody, and target Her2, having a Drug to Antibody Ratio of 2, hence bearing two May (Maytansine) molecules. Herceptin is a non-deglycosylated, unconjugated antibody, targeting Her2.



Cell toxicity assay: Cells were seeded into 96 well plates (white walled, clear flat bottom plates) at densities of 10,000 cells per well and incubated overnight at 37°C and 5% CO₂. Monoclonal antibodies (mAbs) and antibody-drug conjugates (ADCs) were serially diluted 1:4 in media at a starting concentration of 10 µg/mL (66.7 nM). Media was removed from cells, and mAb/ADC dilutions were added. Cells treated with media only served as the reference for 100% viability. Cells were incubated with antibodies for three days at 37°C and 5% CO₂.

Cell viability was assessed by Cell Titer-Glo[®] (Promega) following manufacturer's instructions and as briefly outlined here. Plates were equilibrated to room temperature for 30 minutes. Cell Titer-Glo[®] reagent was made by addition of Cell Titer-Glo buffer to substrate.

50 μ L per well of Cell Titer-Glo® reagent was added and incubated at room temperature with shaking for two minutes followed by an additional 30 minutes incubation at room temperature. Luminescence was detected on a Perkin Elmer 2030 Multilabel Reader Victor™ X3 plate reader using an integration time of 1 second.

The data were processed as follows: luminescence values of wells treated with media only were averaged and served as the reference for 100% viability. Percent viability of mAb/ADC treated wells was calculated using the following equation:

$$\% \text{ viability} = \left(\frac{\text{Luminescence of treated well}}{\text{Average luminescence of media treated wells}} \right) * 100\%$$

Normalized percent viability was plotted versus the logarithm of mAb/ADC concentration and the data were fit using GraphPad Prism 7.00.

As can be seen in Fig. 18, P684 and P579 have the same potency against SK-BR3 cells as Kadcylla. Hence, the advantages provided by the novel linker technology (ease of manufacture, site specificity, stable stoichiometry, no need to deglycosylate that antibody) do not come at any disadvantage regarding the cellular toxicity. This is even more important as P684 and P579 have a DAR of 2, while Kadcylla has an average DAR of 3.53 ± 0.05 , hence is capable to deliver more toxin to the target cells. The following table show the potencies (IC50):

Her-P684-May	1.4 nM
Her-P579-May	0.50 nM
Kadcylla	0.33 nM

Example 4: Conjugation efficiency

Peptides were used as obtained and dissolved at a suitable stock concentration (e.g. 25mM) following the manufacturers instruction, aliquots were prepared and stored at -20°C. The anti-Her2 IgG1 antibody (Trastuzumab) was modified as follows: 1 mg/mL of non-deglycosylated antibody ($\sim 6.67 \mu\text{M}$) was mixed with 80 molar equivalents of peptide linker (i.e. $\sim 533 \mu\text{M}$), 6 U/mL MTG and buffer. The reaction mixture was incubated for 20 h at 37°C and then subjected for LC-MS analysis under reducing conditions.

The following table 8 shows the conjugation efficiencies (CE in %) of various linkers falling within the scope of the present invention.

Sequence	Charge	C-Terminal	Formula	MW	CE in %
GDC	negative	amidation	C ₉ H ₁₆ N ₄ O ₅ S ₁	292,31	9,7
GRCD	neutral	amidation	C ₁₅ H ₂₈ N ₈ O ₆ S ₁	448,49	11,2
GRDC	neutral	amidation	C ₁₅ H ₂₈ N ₈ O ₆ S ₁	448,49	4,4
GGDC	negative	amidation	C ₁₁ H ₁₉ N ₅ O ₆ S ₁	349,36	25,1
GGCD	negative	amidation	C ₁₁ H ₁₉ N ₅ O ₆ S ₁	349,36	49,1
GGEC	negative	amidation	C ₁₂ H ₂₁ N ₅ O ₆ S ₁	363,39	23,7
GGK(N ₃)D	negative	amidation	C ₁₄ H ₂₄ N ₈ O ₆	400,4	68,4
GGRC	neutral	amidation	C ₁₇ H ₃₁ N ₉ O ₇ S ₁	505,54	78,8
GGGDC	negative	amidation	C ₁₃ H ₂₂ N ₆ O ₇ S ₁	406,41	35,6
GC	neutral	amidation	C ₅ H ₁₀ N ₂ O ₃ S ₁	178,21	92
GRC	positive	amidation	C ₁₁ H ₂₃ N ₇ O ₃ S ₁	333,41	51,7
GGRC	positive	amidation	C ₁₃ H ₂₆ N ₈ O ₄ S ₁	390,46	91,4
GRAC	positive	amidation	C ₁₄ H ₂₈ N ₈ O ₄ S ₁	404,49	47,7
GARC	positive	amidation	C ₁₄ H ₂₈ N ₈ O ₄ S ₁	404,49	88,5
GGHK(N ₃)	positive	amidation	C ₁₆ H ₂₆ N ₁₀ O ₄	422,45	16,1
GGK(N ₃)RC	positive	amidation	C ₁₉ H ₃₆ N ₁₂ O ₅ S ₁	544,63	79,5
GARK(N ₃)	positive	amidation	C ₁₉ H ₃₇ N ₉ O ₄	455,56	77,3
AARK(N ₃)	positive	amidation	C ₂₀ H ₃₉ N ₉ O ₄	469,59	10,7
SARK(N ₃)	positive	amidation	C ₂₀ H ₃₉ N ₉ O ₅	485,59	5,2
AGRK(N ₃)	positive	amidation	C ₁₉ H ₃₇ N ₉ O ₄	455,56	22,3
βAGARK(N ₃)	positive	amidation	C ₂₀ H ₃₈ N ₁₂ O ₅	526,6	57
GGARK(N ₃)	positive	amidation	C ₁₉ H ₃₆ N ₁₂ O ₅	512,58	84

REFERENCES

Dorywalska et al (2015), Site-Dependent Degradation of a Non-Cleavable Auristatin-Based Linker-Payload in Rodent Plasma and Its Effect on ADC Efficacy. PLoS ONE 10(7): e0132282

Sletten & Bertozzi, A Bioorthogonal Quadricyclane Ligation. J Am Chem Soc 2011, 133 (44), 17570-17573.

Agard et al, J Am Chem Soc. 2004 Nov 24;126(46):15046-7.

Stöckmann et al (2011). "Exploring isonitrile-based click chemistry for ligation with biomolecules". Organic & Biomolecular Chemistry. 9 (21): 7303.

Blackman et al (2008). "The Tetrazine Ligation: Fast Bioconjugation based on Inverse-electron-demand Diels-Alder Reactivity". Journal of the American Chemical Society. 130 (41): 13518–9.

Yarema, et al (1998). "Metabolic Delivery of Ketone Groups to Sialic Acid Residues. Application To Cell Surface Glycoform Engineering". Journal of Biological Chemistry. 273 (47): 31168–79.

Ning et al (2010). "Protein Modification by Strain-Promoted Alkyne-Nitrone Cycloaddition". Angewandte Chemie International Edition. 49 (17): 3065.

Baskin et al (2007). "Copper-free click chemistry for dynamic in vivo imaging". Proceedings of the National Academy of Sciences. 104 (43): 16793–7.

MacKenzie, DA; Sherratt, AR; Chigrinova, M; Cheung, LL; Pezacki, JP (Aug 2014). "Strain-promoted cycloadditions involving nitrones and alkynes—rapid tunable reactions for bioorthogonal labeling". Curr Opin Chem Biol. 21: 81–8.

Agard, N. J.; Baskin, J. M.; Prescher, J. A.; Lo, A.; Bertozzi, C. R. (2006). "A Comparative Study of Bioorthogonal Reactions with Azides". ACS Chem. Biol. 1: 644–648

Kolb, H.C.; Sharpless, B.K. (2003). "The growing impact of click chemistry on drug discovery". *Drug Discov Today*. **8** (24): 1128–1137.

Lhospice et al., Site-Specific Conjugation of Monomethyl Auristatin E to Anti-Cd30 Antibodies Improves Their Pharmacokinetics and Therapeutic Index in Rodent Models, *Mol Pharm* 12 (6), 1863-1871. 2015

Jeger et al, Site-specific and stoichiometric modification of antibodies by bacterial transglutaminase. *Angew Chem Int Ed Engl*. 2010 Dec 17;49(51):9995-7

Strop, et al., Versatility of Microbial Transglutaminase. *Bioconjugate Chemistry* 2014, 25 (5), 855-862.

Spycher et al., Dual Site-Specifically Modified Antibodies With Solid-Phase Immobilized Microbial Transglutaminase. *Chembiochem*. 2017 Aug 03; 18(19): 1923–1927

Dennler et al., Transglutaminase-based chemo-enzymatic conjugation approach yields homogeneous antibody-drug conjugates. *Bioconjug Chem*. 2014 Mar 19;25(3):569-78

Dennler et al. Microbial transglutaminase and c-myc-tag: a strong couple for the functionalization of antibody-like protein scaffolds from discovery platforms. *Chembiochem*. 2015 Mar 23;16(5):861-7

Mindt, et al., Modification of different IgG1 antibodies via glutamine and lysine using bacterial and human tissue transglutaminase. *Bioconjugate chemistry* 2008, 19 (1), 271-8.

Dubowchik et al., Cathepsin B-labile dipeptide linkers for lysosomal release of doxorubicin from internalizing immunoconjugates: model studies of enzymatic drug release and antigen-specific in vitro anticancer activity. *Bioconjug Chem*. 2002 Jul-Aug; 13(4):855-69.

Zheng, et al., The impact of glycosylation on monoclonal antibody conformation and stability. *Mabs-Austin* 2011, 3 (6), 568-576.

Subedi, et al., The Structural Role of Antibody N-Glycosylation in Receptor Interactions. *Structure* 2015, 23 (9), 1573-1583.

Kieliszek and Misiewicz, *Folia Microbiol (Praha)*. 2014; 59(3): 241–250

Brinkmann and Kontermann, The making of bispecific antibodies. *MAbs*. 2017 Feb-Mar; 9(2): 182–212.

Azhdarinia A. et al., Dual-Labeling Strategies for Nuclear and Fluorescence Molecular Imaging: A Review and Analysis. *Mol Imaging Biol*. 2012 Jun; 14(3): 261–276.

Houghton JL. et al., Site-specifically labeled CA19.9-targeted immunoconjugates for the PET, NIRF, and multimodal PET/NIRF imaging of pancreatic cancer. *Proc Natl Acad Sci U S A*. 2015 Dec 29;112(52):15850-5

Levengood M. et al., Orthogonal Cysteine Protection Enables Homogeneous Multi-Drug Antibody–Drug Conjugates. *Angewandte Chemie*, Volume56, Issue3, January 16, 2017

Costoplus JA. et al., Peptide-Cleavable Self-immolative Maytansinoid Antibody-Drug Conjugates Designed To Provide Improved Bystander Killing. *ACS Med Chem Lett*. 2019 Sep 27;10(10):1393-1399.

Sonzini S. et al., Improved Physical Stability of an Antibody-Drug Conjugate Using Host-Guest Chemistry. *Bioconjug Chem*. 2020 Jan 15;31(1):123-129.

Bodero L. et al., Synthesis and biological evaluation of RGD and isoDGR peptidomimetic- α -amanitin conjugates for tumor-targeting. *Beilstein J. Org. Chem*. 2018, 14, 407–415.

Nunes JPM. et al., Use of a next generation maleimide in combination with THIOMAB™ antibody technology delivers a highly stable, potent and near homogeneous THIOMAB™ antibody-drug conjugate (TDC). *RSC Adv.*, 2017,7, 24828-24832.

Doronina SO. et al., Enhanced activity of monomethylauristatin F through monoclonal antibody delivery: effects of linker technology on efficacy and toxicity. *Bioconjug Chem.* 2006 Jan-Feb;17(1):114-24.

Nakada T. et al., Novel antibody drug conjugates containing exatecan derivative-based cytotoxic payloads. *Bioorg Med Chem Lett.* 2016 Mar 15;26(6):1542-1545.

Dickgiesser S. et al., Site-Specific Conjugation of Native Antibodies Using Engineered Microbial Transglutaminases. *Bioconjug Chem.* 2020 Mar 12. doi: 10.1021/acs.bioconjchem.0c00061.

Steffen W. et al., Discovery of a microbial transglutaminase enabling highly site-specific labeling of proteins. *The Journal of Biological Chemistry.* July 27, 2017 doi: 10.1074/jbc.M117.797811.

Malesevic M. et al., A fluorescence-based array screen for transglutaminase substrates. *Chembiochem.* 2015 May 26;16(8):1169-74.


DISCLAIMER

It is important to understand that in some linker peptides shown herein, the moiety at the C-terminus is simply designated as N₃. However, this should be understood as an abbreviation of Lys(N₃). For example, GAR(N₃) does actually mean GARK₁, with K₁ = Lys(N₃), or GlyAlaArgLys(N₃).

It is furthermore important to understand that in different linker peptides shown herein, the C-terminus may or may not be protected, even if shown otherwise. Protection can be accomplished by amidation. In the context of the present invention, both the protected and unprotected linker peptides are encompassed. For example GARK(N₃) does indeed encompass two variants, with the C-terminus protected or unprotected.

CLAIMS

1. A method for generating an antibody-linker conjugate by means of a microbial transglutaminase (MTG), which method comprises the step of conjugating a linker comprising the peptide structure (shown in N → C direction)



via the N-terminal primary amine of the N-terminal glycine (Gly) residue to a glutamine (Gln) residue comprised in the heavy or light chain of an antibody,

wherein

- m is an integer between ≥ 0 and ≤ 12
- n is an integer between ≥ 0 and ≤ 12
- $m + n \geq 0$,
- Aax is an amino acid or an amino acid derivative, and
- B is a linking moiety.

2. The method according to claim 1, wherein the linker comprises two or more linking moieties B.

3. The method according to claim 2, wherein the two or more linking moieties B differ from one another.

4. The method according to any one of claims 1 to 3, wherein at least one of the one or more linking moieties B comprises

- a bioorthogonal marker group, or
- a non-bio-orthogonal entity for crosslinking.

5. The method according to claim 4, wherein the bioorthogonal marker group or the non-bio-orthogonal entity is at least one selected from a group consisting of:


- $-N-N\equiv N$, or $-N_3$
- Lys(N_3)
- tetrazine
- alkyne
- DBCO
- BCN
- norborene
- transcyclooctene
- $-RCOH$ (aldehyde),
- acyltrifluoroborates,
- $-SH$, and
- cysteine.

6. A method for generating an antibody-payload conjugate, the method comprising the steps of

- a) generating an antibody-linker conjugate according to any one of claims 1 to 5, and
- b) linking a payload to the one or more linking moieties B of the antibody-linker conjugate.

7. The method according to claim 6, wherein the payload is linked to the linking moiety B of the antibody-linker conjugate via a click-reaction.

8. A method for generating an antibody-payload conjugate by means of a microbial transglutaminase (MTG), which method comprises the step of conjugating a linker comprising the peptide structure (shown in N -> C direction)



via the N-terminal primary amine of the N-terminal glycine (Gly) residue to a glutamine (Gln) residue comprised in the heavy or light chain of an antibody,

wherein

- m is an integer between ≥ 0 and ≤ 12
- n is an integer between ≥ 0 and ≤ 12
- $m + n \geq 0$,
- Aax is an amino acid or an amino acid derivative, and
- B is a payload.

9. The method according to claim 8, wherein the linker comprises two or more payloads B.

10. The method according to claim 9, wherein the two or more payloads B differ from one another.

11. The method according to any one of claims 6 to 10, wherein the one or more payloads is selected from a group consisting of:

- a toxin
- a cytokine
- a growth factor
- a radionuclide
- a hormone
- an anti-viral agent
- an anti-bacterial agent
- a fluorescent dye
- an immunoregulatory/immunostimulatory agent
- a half-life increasing moiety
- a solubility increasing moiety
- a polymer-toxin conjugate

- a nucleic acid
- a biotin or streptavidin moiety
- a vitamin
- a target binding moiety, and
- an anti-inflammatory agent.

12. The method according to claim 11, wherein the toxin is at least one selected from the group consisting of

- pyrrolbenzodiazepines (PBD)
- auristatins (e.g., MMAE, MMAF)
- maytansinoids (maytansine, DM1, DM4, DM21)
- duocarmycins
- tubulysins
- enediyenes (e.g. calicheamicin)
- PNUs, doxorubicins
- pyrrole-based kinesin spindle protein (KSP) inhibitors
- calicheamicins
- amanitins (e.g. α -amanitin), and
- camptothecins (e.g. exatecans, deruxtecans).

13. The method according to any one of claims 1 to 12, wherein the linker is not cleavable by cathepsin.

14. The method according to any one of claims 1 to 13, wherein the linker does not comprise a valine-alanine motif or a valine-citrulline motif.

15. The method according to any one of claims 1 to 14, wherein the antibody is an IgG, IgE, IgM, IgD, IgA or IgY antibody, or a fragment or recombinant variant thereof, wherein the fragment or recombinant variant thereof retains target binding properties and comprises a C_H2 domain.

16. The method according to claim 15, wherein the antibody is an IgG antibody.

17. The method according to claim 15 or 16, wherein the antibody is a glycosylated antibody, a deglycosylated antibody or an aglycosylated antibody.

18. The method according to claim 17, wherein the glycosylated antibody is an IgG antibody that is glycosylated at residue N297 (EU numbering) of the C_H2 domain.

19. The method according to any one of claims 1 to 18, wherein (a) the linker including the payload or linking moiety B is conjugated to a Gln residue which has been introduced into the heavy or light chain of the antibody by molecular engineering or (b) the linker including the payload or linking moiety B is conjugated to a Gln residue in the Fc domain of the antibody.

20. The method according to claim 19, wherein the Gln residue in the Fc domain of the antibody is Gln residue Q295 (EU numbering) of the C_H2 domain of an IgG antibody.

21. The method according to claim 19, wherein the Gln residue that has been introduced into the heavy or light chain of the antibody by molecular engineering is N297Q (EU numbering) of the C_H2 domain of an aglycosylated IgG antibody.

22. The method according to claim 19 wherein the Gln residue that has been introduced into the heavy or light chain of the antibody by molecular engineering is comprised in a peptide that has been (a) integrated into the heavy or light chain of the antibody or (b) fused to the N- or C-terminal end of the heavy or light chain of the antibody.

23. The method according to claim 22, wherein the peptide comprising the Gln residue has been fused to the C-terminal end of the heavy chain of the antibody.

24. The method according to claim 22 or 23, wherein the peptide comprising the Gln residue is selected from a group consisting of:

- LLQGG,
- LLQG,
- LSLSQG,

- GGLLQGG,
- GLLQG,
- LLQ,
- GSPLAQSHGG,
- GLLQGGG,
- GLLQGG,
- GLLQ,
- LLQLLQGA,
- LLQGA,
- LLQYQGA,
- LLQGSG,
- LLQYQG,
- LLQLLQG,
- SLLQG,
- LLQLQ,
- LLQLLQ,
- LLQGR,
- EEQYASTY,
- EEQYQSTY,
- EEQYNSTY,
- EEQYQS,
- EEQYQST,
- EQYQSTY,
- QYQS,
- QYQSTY,
- YRYRQ,
- DYALQ,
- FGLQRPY,
- EQKLISEEDL,
- LQR, and
- YQR.

25. The method according to any one of claims 1 to 24, wherein $m + n \leq 12, 11, 10, 9, 8, 7, 6, 5$ or 4.

26. The method according to any one of claims 1 to 25, wherein the net charge of the linker is neutral or positive.

27. The method according to any one of claims 1 to 26, wherein the linker does not comprise negatively charged amino acid residues.

28. The method according to any one of claims 1 to 27, wherein the linker comprises at least one positively charged amino acid residue.

29. The method according to any one claims 1 to 28, wherein the linker comprises at least one amino acid residue selected from a group consisting of


- lysine,
- arginine, and
- histidine

30. The method according to any one of claims 1 to 29, wherein the linker comprising the at least one payload or linking moiety B is conjugated to the amide side chain of the Gln residue.

31. The method according to any one of claims 1 to 30, wherein the microbial transglutaminase is derived from a *Streptomyces* species, in particular *Streptomyces mobaraensis*.

32. An antibody-payload conjugate which has been generated with a method according to any one of claims 6 to 31.

33. A linker comprising the peptide structure (shown in N -> C direction)



wherein Gly comprises an N-terminal primary amine, and wherein

- m is an integer between ≥ 0 and ≤ 12
- n is an integer between ≥ 0 and ≤ 12
- $m + n \geq 0$,
- Aax is an amino acid or an amino acid derivative, and
- B is a payload or a linking moiety,

wherein the linker can be conjugated to an antibody by a microbial transglutaminase via the N-terminal primary amine of the N-terminal Gly of the linker.

34. The linker according to claim 33, wherein the linker comprises two or more payloads and/or linking moieties B.

35. The linker according to claim 33 or 34, wherein at least one of the one or more linking moieties B comprises

- a bioorthogonal marker group, or
- a non-bio-orthogonal entity for crosslinking.

36. The linker according to claim 35, wherein the bioorthogonal marker group or the non-bio-orthogonal entity is at least one selected from a group consisting of:

- $-N-N\equiv N$, or $-N_3$
- Lys(N_3)
- tetrazine
- alkyne
- DBCO
- BCN
- norborene
- transcyclooctene
- $-RCOH$ (aldehyde),
- acyltrifluoroborates,

- -SH, and
- cysteine.

37. The linker according to claim 33 or 34, wherein the one or more payloads is selected from a group consisting of:

- a toxin
- a cytokine
- a growth factor
- a radionuclide
- a hormone
- an anti-viral agent
- an anti-bacterial agent
- a fluorescent dye
- an immunoregulatory/immunostimulatory agent
- a half-life increasing moiety
- a solubility increasing moiety
- a polymer-toxin conjugate
- a nucleic acid
- a biotin or streptavidin moiety
- a vitamin
- a target binding moiety, and
- an anti-inflammatory agent.

38. The linker according to claim 37, wherein the toxin is at least one selected from the group consisting of

- pyrrolbenzodiazepines (PBD)
- auristatins (e.g., MMAE, MMAF)
- maytansinoids (maytansine, DM1, DM4, DM21)
- duocarmycins
- tubulysins
- enediyenes (e.g. calicheamicin)

- PNUs, doxorubicins
- pyrrole-based kinesin spindle protein (KSP) inhibitors
- calicheamicins
- amanitins (e.g. α -amanitin), and
- camptothecins (e.g. exatecans, deruxtecans).

39. The linker according to any one of claims 33 to 38, wherein the linker is not cleavable by cathepsin.

40. The linker according to any one of claims 33 to 39, wherein the linker does not comprise a valine-alanine motif or a valine-citrulline motif.

41. The linker according to any one of claims 33 to 40, wherein $m + n \leq 12, 11, 10, 9, 8, 7, 6, 5$ or 4.

42. The linker according to any one of claims 33 to 41, wherein the net charge of the linker is neutral or positive.

43. The linker according to any one of claims 33 to 42, wherein the linker does not comprise negatively charged amino acid residues.

44. The linker according to any one of claims 33 to 43, wherein the linker comprises at least one positively charged amino acid residue.

45. The linker according to any one of claims 33 to 44, wherein the linker comprises at least one amino acid residue selected from a group consisting of

- lysine,
- arginine, and
- histidine.

46. The linker according to any one of claims 33 to 45, wherein the linker is selected from the list as shown in table 5.

47. A linker-payload construct comprising at least

- a) a linker according to any one of claims 33 to 46, and
- b) one or more payloads,

wherein the one or more payloads are covalently or non-covalently bound to the linker.

48. The linker-payload construct according to claim 47, wherein in said construct, the one or more payloads have been covalently bound to the linking moiety B of the linker with a click reaction.

49. The linker-payload construct according to claim 47, wherein the linker-payload construct has been obtained by chemical synthesis.

50. The linker-payload construct according to any one of claims 47 to 49, wherein in said construct, the linker and/or the one or more payloads have been chemically modified during binding to allow covalent or non-covalent binding to form said construct.

51. An antibody-payload conjugate comprising

- a) one or more linker-payload construct according to any one of claims 47 to 50, and
- b) an antibody comprising at least one Gln residue in the heavy or light chain,

wherein the linker-payload construct is conjugated to the amide side chain of a Gln residue in the heavy or light chain of the antibody via an N-terminal primary amine of the N-terminal glycine residue comprised in the linker-payload construct.

52. The antibody-payload conjugate according to claim 51, wherein the conjugation has been achieved with a microbial transglutaminase (MTG).

53. The antibody-payload conjugate according to claim 51 or 52, wherein the conjugation has been achieved before or after formation of the linker-payload construct.

54. The antibody-payload conjugate according to any one of claims 51 to 53, wherein in said conjugate, the linker-payload constructs and/or the antibody have optionally been chemically modified during conjugation to allow covalent conjugation, to form said conjugate.

55. The antibody-payload conjugate according to any one of claims 51 to 54, wherein the antibody is an IgG, IgE, IgM, IgD, IgA or IgY antibody, or a fragment or recombinant variant thereof, wherein the fragment or recombinant variant thereof retains target binding properties and comprises a C_H2 domain.

56. The antibody-payload conjugate according to claim 55, wherein the antibody is an IgG antibody.

57. The antibody-payload conjugate according to claim 55 or 56, wherein the antibody is a glycosylated antibody, a deglycosylated antibody or an aglycosylated antibody.

58. The antibody-payload conjugate according to claim 57, wherein the glycosylated antibody is an IgG antibody that is glycosylated at residue N297 (EU numbering) of the C_H2 domain.

59. The antibody-payload conjugate according to any one of claims 51 to 58, wherein (a) the linker-payload construct is conjugated to a Gln residue which has been introduced into the heavy or light chain of the antibody by molecular engineering or (b) the linker-payload construct is conjugated to a Gln residue in the Fc domain of the antibody.

60. The antibody-payload conjugate according to claim 59, wherein the Gln residue in the Fc domain of the antibody is Gln residue Q295 (EU numbering) of the C_H2 domain of an IgG antibody.

61. The antibody-payload conjugate according to claim 59, wherein the Gln residue that has been introduced into the heavy or light chain of the antibody by molecular engineering is N297Q (EU numbering) of the C_{H2} domain of an aglycosylated antibody.

62. The antibody-payload conjugate according to claim 59, wherein the Gln residue that has been introduced into the heavy or light chain of the antibody by molecular engineering is comprised in a peptide that has been (a) integrated into the heavy or light chain of the antibody or (b) fused to the N- or C-terminal end of the heavy or light chain of the antibody.

63. The antibody-payload conjugate according to claim 62, wherein the peptide comprising the Gln residue has been fused to the C-terminal end of the heavy chain of the antibody.

64. The antibody-payload conjugate according to claim 62 or 63, wherein the peptide comprising the Gln residue is selected from a group consisting of:

- LLQGG,
- LLQG,
- LSLSQG,
- GGGLLQGG,
- GLLQG,
- LLQ,
- GSPLAQSHGG,
- GLLQGGG,
- GLLQGG,
- GLLQ,
- LLQLLQGA,
- LLQGA,
- LLQYQGA,
- LLQGSG,
- LLQYQG,
- LLQLLQG,

- SLLQG,
- LLQLQ,
- LLQLLQ,
- LLQGR,
- EEQYASTY,
- EEQYQSTY,
- EEQYNSTY,
- EEQYQS,
- EEQYQST,
- EQYQSTY,
- QYQS,
- QYQSTY,
- YRYRQ,
- DYALQ,
- FGLQRPY,
- EQKLISEEDL,
- LQR, and
- YQR.

65. The antibody-payload conjugate according to any one of claims 51 to 64, wherein the antibody-payload conjugate comprises at least one toxin.

66. The antibody-payload conjugate according to claim 65, wherein the antibody-payload conjugate comprises a toxin and an inhibitor of a drug efflux transporter.

67. The antibody-payload conjugate according to claim 65, wherein the antibody-payload conjugate comprises a toxin and a solubility increasing moiety.

68. The antibody-payload conjugate according to claim 65, wherein the antibody-payload conjugate comprises a toxin and an immunostimulatory agent.

69. The antibody-payload conjugate according to claim 65, wherein the antibody-payload conjugate comprises two different toxins.

70. The antibody-payload conjugate according to claim 69, wherein a first toxin is a toxin that inhibits cell division and a second toxin is a toxin that interferes with replication and/or transcription of DNA.

71. The antibody-payload conjugate according to any one of claims 65 to 70, wherein at least one of the toxins is an auristatin or a maytansinoid.

72. The antibody-payload conjugate according to any one of claims 51 to 64, wherein the antibody-payload conjugate comprises two immunostimulatory agents.

73. The antibody-payload conjugate according to claim 68 to 72, wherein the at least one immunostimulatory agent is a TLR agonist.

74. The antibody-payload conjugate according to any one of claims 51 to 64, wherein the antibody-payload conjugate comprises a radionuclide and a fluorescent dye.

75. The antibody-payload conjugate according to claim 74, wherein the radionuclide is a radionuclide that is suitable for use in tomography, in particular single-photon emission computed tomography (SPECT) or positron emission tomography (PET), and wherein the fluorescent dye is a near-infrared fluorescent dye.

76. A pharmaceutical composition comprising the linker according to any one of claims 33 – 46, the linker-payload construct according to any one of claims 47 to 50, and/or the antibody-payload conjugate according to any one of claims 51 to 75.

77. A pharmaceutical product comprising the antibody-payload conjugate according to any one of claims 51 to 75 or the pharmaceutical composition according to claim 76 and at least one further pharmaceutically acceptable ingredient.

78. The antibody-payload conjugate according to any one of claims 51 to 75, the pharmaceutical composition according to claim 76 or the pharmaceutical product according to claim 77 for use in therapy and/or diagnostics.

79. The antibody-payload conjugate according to any one of claims 51 to 75, the pharmaceutical composition according to claim 76 or the pharmaceutical product according to claim 77 for use in treatment of a patient

- suffering from,
- being at risk of developing, and/or
- being diagnosed for

a neoplastic disease, neurological disease, an autoimmune disease, an inflammatory disease or an infectious disease.

80. The antibody-payload conjugate according to any one of claims 51 to 75, the pharmaceutical composition according to claim 76 or the pharmaceutical product according to claim 77 for use in treatment of a patient suffering from a neoplastic disease.

81. Use of the antibody-payload conjugate according to any one of claims 51 to 75, the pharmaceutical composition according to claim 76 or the pharmaceutical product according to claim 77 for the manufacture of a medicament for the treatment of a patient

- suffering from,
- being at risk of developing, and/or
- being diagnosed for

a neoplastic disease, neurological disease, an autoimmune disease, an inflammatory disease or an infectious disease.

82. A method of treating or preventing a neoplastic disease, said method comprising administering to a patient in need thereof the antibody-payload conjugate according to any one of claims 51 to 75, the pharmaceutical composition according to claim 76 or the pharmaceutical product according to claim 77.

83. The antibody-payload conjugate according to any one of claims 51 to 75, the pharmaceutical composition according to claim 76 or the pharmaceutical product according to claim 77 for use in pre-, intra- or post-operative imaging.

84. The antibody-payload conjugate according to any one of claims 51 to 75, the pharmaceutical composition according to claim 76 or the pharmaceutical product according to claim 77 for use in intraoperative imaging-guided cancer surgery.

Fig. 1




Fig. 2




Fig. 3




Fig.4







Fig.5




maleimidocaproyl (mc)



maleimidomethyl cyclohexane-1-carboxylate (mcc)



mc-like linker used in SYD985



self-stabilizing maleimide

Fig. 6






Fig. 7

A



B





Fig. 8

A



B




Fig. 9



Structure 1**Structure 2**

Fig. 9 ctd'



Structure 3**Structure 4**

Fig. 9 ctd'


Structure 5

Fig. 10




Fig. 11





Fig. 11 ctd'




Fig. 12






Fig. 13

IMGT unique numbering for C-DOMAIN	CH2 IGHG1 amino acid translation J00228 (1)	IMGT exon numbering 1-110	Eu numbering [1] (1) (110 aa) [2]	Kabat numbering [2]	Strands, turns and loops for C-DOMAIN (5)
1,8	-	-	-	-	
1,7	-	-	-	-	
1,6	(A)	1	231	244	
1,5	P	2	232	245	
1,4	E	3	233	246	
1,3	L	4	234	247	
1,2	L	5	235	248	
1,1	G	6	236	249	
1	G	7	237	250	
2	P	8	238	251	
3	S	9	239	252	
4	V	10	240	253	
5	F	11	241	254	
6	L	12	242	255	
7	F	13	243	256	
8	P	14	244	257	A-STRAND
9	P	15	245	258	
10	K	16	246	259	
11	P	17	247	260	
12	K	18	248	261	
13	D	19	249	262	
14	T	20	250	263	
15	L	21	251	264	
15,1	M	22	252	265	
15,2	I	23	253	266	AB-TURN
15,3	-	-	-	-	
16	S	24	254	267	
17	R	25	255	268	
18	T	26	256	269	
19	P	27	257	270	
20	E	28	258	271	
21	V	29	259	272	B-STRAND
22	T	30	260	273	
23	C	31	261	274	
24	V	32	262	275	
25	V	33	263	276	
26	V	34	264	277	
27	D	35	265	278	
28	V	36	266	279	
29	S	37	267	280	
30	H	38	268	281	
31	E	39	269	282	
34	D	40	270	283	BC-LOOP
35	P	41	271	284	
36	E	42	272	285	
37	V	43	273	286	
38	K	44	274	287	
39	F	45	275	288	
40	N	46	276	289	
41	W	47	277	290	
42	Y	48	278	291	C-STRAND
43	V	49	279	292	
44	D	50	280	295	
45	G	51	281	296	
45,1	V	52	282	299	
45,2	E	53	283	300	
45,3	V	54	284	301	
45,4	H	55	285	302	CD-STRAND
45,5	-	-	-	-	
45,6	-	-	-	-	
45,7	-	-	-	-	

IMGT unique numbering for C-DOMAIN	CH2 IGHG1 amino acid translation J00228 (1)	IMGT exon numbering 1-110	Eu numbering [1] (1) (110 aa) [2]	Kabat numbering [2]	Strands, turns and loops for C-DOMAIN (5)
77	N	56	286	303	
78	A	57	287	304	
79	K	58	288	305	
80	T	59	289	306	
81	K	60	290	307	D-STRAND
82	P	61	291	308	
83	R	62	292	309	
84	E	63	293	310	
84,1	E	64	294	311	
84,2	Q	65	295	312	
84,3	Y	66	296	313	
84,4	N	67	297	314	
84,5	-	-	-	-	
84,6	-	-	-	-	
84,7	-	-	-	-	DE-TURN
85,7	-	-	-	-	
85,6	-	-	-	-	
85,5	-	-	-	-	
85,4	S	68	298	317	
85,3	T	69	299	318	
85,2	Y	70	300	319	
85,1	R	71	301	320	
85	V	72	302	321	
86	V	73	303	322	
87	S	74	304	323	
88	V	75	305	324	
89	L	76	306	325	
90	T	77	307	326	
91	V	78	308	327	E-STRAND
92	L	79	309	328	
93	H	80	310	329	
94	Q	81	311	330	
95	D	82	312	331	
96	W	83	313	332	
96,1	-	-	-	-	
96,2	-	-	-	-	EF-TURN
97	L	84	314	333	
98	N	85	315	334	
99	G	86	316	335	
100	K	87	317	336	
101	E	88	318	337	F-STRAND
102	Y	89	319	338	
103	K	90	320	339	
104	C	91	321	340	
105	K	92	322	341	
106	V	93	323	342	
107	S	94	324	343	
108	N	95	325	344	
109	K	96	326	345	
110	A	97	327	346	
111	-	-	-	-	FG-LOOP
112	-	-	-	-	
113	L	98	328	347	
114	P	99	329	348	
115	A	100	330	349	
116	P	101	331	350	
117	I	102	332	351	
118	E	103	333	352	
119	K	104	334	353	
120	T	105	335	354	
121	I	106	336	355	
122	S	107	337	357	G-STRAND
123	K	108	338	358	
124	A	109	339	359	
125	K	110	340	360	

Fig. 14



Antibody with Q295 having free primary amine (NH_2)

Oligopeptide having an N-terminal Gly residue

Formation of crosslinking isopeptidyl bond

Fig. 15




Fig. 16





	GlyAlaArgLys(N3)	Lys(N3) comprises an Azide for click chemistry instead of a primary amine.
	GlyβAlaArgLys(N3)	βAla does not make a peptide bond, hence this is an example for a peptidomimetic
	GlyAla-homoArg-Lys(N3)	homoArg is a non naturally occurring amino acid
	GlyβAla-homoArg-Lys(N3)	Combination of the upper two

Fig. 17




Fig. 18





Fig. 19



SEQUENCE LISTING

<110> Paul Scherrer Institut

<120> Transglutaminase conjugation method with a glycine based linker

<130> AD1457 PCT BS

<160> 83

<170> BiSSAP 1.3.6

<210> 1

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 4..4

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 1

Gly Ala Arg Xaa

1

<210> 2

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 4..4

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<220>

<221> MOD_RES

<222> 5..5

<223> Xaa is Lys(tetrazine)

<400> 2

Gly Ala Arg Xaa Xaa

1

5

<210> 3

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 4..4

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 3

Gly Ala Arg Xaa Cys
1 5

<210> 4
<211> 6
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 5..5
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<220>
<221> MOD_RES
<222> 6..6
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 4
Gly Gly Ala Arg Xaa Xaa
1 5

<210> 5
<211> 7
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 5..5
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<220>
<221> MOD_RES
<222> 7..7
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 5
Gly Gly Ala Arg Xaa Arg Xaa
1 5

<210> 6
<211> 4
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 4..4
<223> Xaa is 4-azido-L-homoalanine

<400> 6
Gly Ala Arg Xaa
1

<210> 7
<211> 4

<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<400> 7
Gly Ala Arg Cys
1

<210> 8
<211> 4
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 2..2
<223> Xaa = β -Alanine

<400> 8
Gly Xaa Arg Lys
1

<210> 9
<211> 4
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 3..3
<223> Xaa = homoarginine

<220>
<221> MOD_RES
<222> 4..4
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 9
Gly Ala Xaa Xaa
1

<210> 10
<211> 4
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 2..2
<223> Xaa = β -Alanine

<220>
<221> MOD_RES
<222> 3..3

<223> Xaa = Homoarginine

<220>

<221> MOD_RES

<222> 4..4

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 10

Gly Xaa Xaa Xaa

1

<210> 11

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 11

Gly Gly Ala Arg Arg

1 5

<210> 12

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 4..4

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 12

Gly Arg Ala Xaa

1

<210> 13

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 13

Gly Arg Ala Cys

1

<210> 14

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 4..4

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 14

Gly Gly Arg Xaa

1

<210> 15

<211> 3

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 3..3

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 15

Gly Arg Xaa

1

<210> 16

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 4..4

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 16

Gly Gly Arg Xaa Arg

1

5

<210> 17

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 3..3

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 17

Gly Gly Xaa Arg Cys

1

5

<210> 18

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 5..5
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 18
Gly Gly Arg Arg Xaa
1 5

<210> 19
<211> 5
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 4..4
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 19
Gly Gly Arg Xaa Arg
1 5

<210> 20
<211> 4
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 4..4
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 20
Gly Ala His Xaa
1

<210> 21
<211> 4
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 4..4
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 21
Gly Gly His Xaa
1

<210> 22
<211> 2
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<400> 22

Gly Cys

1

<210> 23

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 23

Gly Gly Arg Cys

1

<210> 24

<211> 3

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 24

Gly Arg Cys

1

<210> 25

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 4..4

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 25

Gly Gly Arg Xaa

1

<210> 26

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 5..5

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 26

Gly Gly Ala Lys Xaa

1

5

<210> 27

<211> 395

<212> PRT

<213> Streptoverticillium ladakanum

<400> 27

Met His Arg Arg Ile His Ala Val Gly Gln Ala Arg Pro Pro Pro Thr
1 5 10 15
Met Ala Arg Gly Lys Glu Thr Lys Ser Tyr Ala Glu Thr Tyr Arg Leu
20 25 30
Thr Ala Asp Asp Val Ala Asn Ile Asn Ala Leu Asn Glu Ser Ala Pro
35 40 45
Ala Ala Ser Ser Ala Gly Pro Ser Phe Arg Ala Pro Asp Ser Asp Asp
50 55 60
Arg Val Thr Pro Pro Ala Glu Pro Leu Asp Arg Met Pro Asp Pro Tyr
65 70 75 80
Arg Pro Ser Tyr Gly Arg Ala Glu Thr Val Val Asn Asn Tyr Ile Arg
85 90 95
Lys Trp Gln Gln Val Tyr Ser His Arg Asp Gly Arg Lys Gln Gln Met
100 105 110
Thr Glu Glu Gln Arg Glu Trp Leu Ser Tyr Gly Cys Val Gly Val Thr
115 120 125
Trp Val Asn Ser Gly Gln Tyr Pro Thr Asn Arg Leu Ala Phe Ala Ser
130 135 140
Phe Asp Glu Asp Arg Phe Lys Asn Glu Leu Lys Asn Gly Arg Pro Arg
145 150 155 160
Ser Gly Glu Thr Arg Ala Glu Phe Glu Gly Arg Val Ala Lys Glu Ser
165 170 175
Phe Asp Glu Glu Lys Gly Phe Gln Arg Ala Arg Glu Val Ala Ser Val
180 185 190
Met Asn Arg Ala Leu Glu Asn Ala His Asp Glu Ser Ala Tyr Leu Asp
195 200 205
Asn Leu Lys Lys Glu Leu Ala Asn Gly Asn Asp Ala Leu Arg Asn Glu
210 215 220
Asp Ala Arg Ser Pro Phe Tyr Ser Ala Leu Arg Asn Thr Pro Ser Phe
225 230 235 240
Lys Glu Arg Asn Gly Gly Asn His Asp Pro Ser Arg Met Lys Ala Val
245 250 255
Ile Tyr Ser Lys His Phe Trp Ser Gly Gln Asp Arg Ser Ser Ser Ala
260 265 270
Asp Lys Arg Lys Tyr Gly Asp Pro Asp Ala Phe Arg Ser Ala Pro Gly
275 280 285
Thr Gly Leu Val Asp Met Ser Arg Asp Arg Asn Ile Pro Arg Ser Pro
290 295 300
Thr Ser Pro Gly Glu Gly Phe Val Asn Phe Asp Tyr Gly Trp Phe Gly
305 310 315 320
Ala Gln Thr Glu Ala Asp Ala Asp Lys Thr Val Trp Thr His Gly Asn
325 330 335
His Tyr His Ala Pro Asn Gly Ser Leu Gly Cys His Ala Cys Leu Thr
340 345 350
Arg Ala Ser Ser Ala Thr Gly Ser Glu Gly Tyr Ser Asp Phe Asp Arg
355 360 365
Gly Glu Pro Tyr Val Val Ser Pro Ser Pro Ser Pro Arg Met Leu Glu
370 375 380
His Arg Pro Arg Gln Gly Lys Ala Gly Leu Ala
385 390 395

<210> 28

<211> 335

<212> PRT

<213> Streptomyces mobaraensis

<400> 28

Phe Arg Ala Pro Asp Ser Asp Glu Arg Val Thr Pro Pro Ala Glu Pro
1 5 10 15
Leu Asp Arg Met Pro Asp Pro Tyr Arg Pro Ser Tyr Gly Arg Ala Glu
20 25 30
Thr Ile Val Asn Asn Tyr Ile Arg Lys Trp Gln Gln Val Tyr Ser His
35 40 45

Arg Asp Gly Arg Lys Gln Gln Met Thr Glu Glu Gln Arg Glu Trp Leu
 50 55 60
 Ser Tyr Gly Cys Val Gly Val Thr Trp Val Asn Ser Gly Gln Tyr Pro
 65 70 75 80
 Thr Asn Arg Leu Ala Phe Ala Phe Phe Asp Glu Asp Lys Tyr Lys Asn
 85 90 95
 Glu Leu Lys Asn Gly Arg Pro Arg Ser Gly Glu Thr Arg Ala Glu Phe
 100 105 110
 Glu Gly Arg Val Ala Lys Asp Ser Phe Asp Glu Ala Lys Gly Phe Gln
 115 120 125
 Arg Ala Arg Asp Val Ala Ser Val Met Asn Lys Ala Leu Glu Asn Ala
 130 135 140
 His Asp Glu Gly Ala Tyr Leu Asp Asn Leu Lys Lys Glu Leu Ala Asn
 145 150 155 160
 Gly Asn Asp Ala Leu Arg Asn Glu Asp Ala Arg Ser Pro Phe Tyr Ser
 165 170 175
 Ala Leu Arg Asn Thr Pro Ser Phe Lys Asp Arg Asn Gly Gly Asn His
 180 185 190
 Asp Pro Ser Lys Met Lys Ala Val Ile Tyr Ser Lys His Phe Trp Ser
 195 200 205
 Gly Gln Asp Arg Ser Gly Ser Ser Asp Lys Arg Lys Tyr Gly Asp Pro
 210 215 220
 Glu Ala Phe Arg Pro Asp Arg Gly Thr Gly Leu Val Asp Met Ser Arg
 225 230 235 240
 Asp Arg Asn Ile Pro Arg Ser Pro Thr Ser Pro Gly Glu Ser Phe Val
 245 250 255
 Asn Phe Asp Tyr Gly Trp Phe Gly Ala Gln Thr Glu Ala Asp Ala Asp
 260 265 270
 Lys Thr Val Trp Thr His Gly Asn His Tyr His Ala Pro Asn Gly Ser
 275 280 285
 Leu Gly Ala Met His Val Tyr Glu Ser Lys Phe Arg Asn Trp Ser Asp
 290 295 300
 Gly Tyr Ser Asp Phe Asp Arg Gly Ala Tyr Val Val Thr Phe Val Pro
 305 310 315 320
 Lys Ser Trp Asn Thr Ala Pro Asp Lys Val Thr Gln Gly Trp Pro
 325 330 335

<210> 29
 <211> 4
 <212> PRT
 <213> artificial sequence

<220>
 <223> linker sequence

<220>
 <221> MOD_RES
 <222> 4..4
 <223> Xaa is [PEG]3N3

<400> 29
 Gly Ala Arg Xaa
 1

<210> 30
 <211> 7
 <212> PRT
 <213> artificial sequence

<220>
 <223> linker sequence

<220>
 <221> MOD_RES
 <222> 5..5
 <223> Xaa is Lys(PEG)n

<220>
<221> MOD_RES
<222> 7..7
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 30
Gly Gly Ala Arg Xaa Arg Xaa
1 5

<210> 31
<211> 5
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 5..5
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 31
Gly Gly Ala Arg Xaa
1 5

<210> 32
<211> 5
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<220>
<221> MOD_RES
<222> 1..1
<223> Xaa is beta-alanine

<220>
<221> MOD_RES
<222> 5..5
<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 32
Xaa Gly Ala Arg Xaa
1 5

<210> 33
<211> 3
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<400> 33
Gly Ala Arg
1

<210> 34
<211> 4
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<400> 34
Gly Ala Arg Arg
1

<210> 35
<211> 4
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<400> 35
Gly Gly Ala Arg
1

<210> 36
<211> 5
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<400> 36
Gly Gly Ala Arg Arg
1 5

<210> 37
<211> 3
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<400> 37
Gly Gly Gly
1

<210> 38
<211> 3
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<400> 38
Gly Asp Cys
1

<210> 39
<211> 4
<212> PRT
<213> artificial sequence

<220>
<223> linker sequence

<400> 39

Gly Arg Cys Asp

1

<210> 40

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 40

Gly Arg Asp Cys

1

<210> 41

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 41

Gly Gly Asp Cys

1

<210> 42

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 42

Gly Gly Cys Asp

1

<210> 43

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 43

Gly Gly Glu Cys

1

<210> 44

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 3..3

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 44

Gly Gly Xaa Asp

1

<210> 45

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 45

Gly Gly Arg Cys Asp

1 5

<210> 46

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<400> 46

Gly Gly Gly Asp Cys

1 5

<210> 47

<211> 4

<212> PRT

<213> artificial sequence

<220>

<223> linker sequence

<220>

<221> MOD_RES

<222> 4..4

<223> Xaa is 6-azido-L-lysine (Lys(N3))

<400> 47

Gly Gly Gly Xaa

1

<210> 48

<211> 335

<212> PRT

<213> Artificial Sequence

<220>

<223> Streptomyces mobaraensis MTG Zedira

<400> 48

Phe Arg Ala Pro Asp Ser Asp Asp Arg Val Thr Pro Pro Ala Glu Pro
1 5 10 15

Leu Asp Arg Met Pro Asp Pro Tyr Arg Pro Ser Tyr Gly Arg Ala Glu
20 25 30

Thr Val Val Asn Asn Tyr Ile Arg Lys Trp Gln Gln Val Tyr Ser His
35 40 45

Arg Asp Gly Arg Lys Gln Gln Met Thr Glu Glu Gln Arg Glu Trp Leu
50 55 60

Ser Tyr Gly Cys Val Gly Val Thr Trp Val Asn Ser Gly Gln Tyr Pro
65 70 75 80

Thr Asn Arg Leu Ala Phe Ala Ser Phe Asp Glu Asp Arg Phe Lys Asn

85 90 95
 Glu Leu Lys Asn Gly Arg Pro Arg Ser Gly Glu Thr Arg Ala Glu Phe
 100 105 110
 Glu Gly Arg Val Ala Lys Glu Ser Phe Asp Glu Glu Lys Gly Phe Gln
 115 120 125
 Arg Ala Arg Glu Val Ala Ser Val Met Asn Arg Ala Leu Glu Asn Ala
 130 135 140
 His Asp Glu Ser Ala Tyr Leu Asp Asn Leu Lys Lys Glu Leu Ala Asn
 145 150 155 160
 Gly Asn Asp Ala Leu Arg Asn Glu Asp Ala Arg Ser Pro Phe Tyr Ser
 165 170 175
 Ala Leu Arg Asn Thr Pro Ser Phe Lys Glu Arg Asn Gly Gly Asn His
 180 185 190
 Asp Pro Ser Arg Met Lys Ala Val Ile Tyr Ser Lys His Phe Trp Ser
 195 200 205
 Gly Gln Asp Arg Ser Ser Ser Ala Asp Lys Arg Lys Tyr Gly Asp Pro
 210 215 220
 Asp Ala Phe Arg Pro Ala Pro Gly Thr Gly Leu Val Asp Met Ser Arg
 225 230 235 240
 Asp Arg Asn Ile Pro Arg Ser Pro Thr Ser Pro Gly Glu Gly Phe Val
 245 250 255
 Asn Phe Asp Tyr Gly Trp Phe Gly Ala Gln Thr Glu Ala Asp Ala Asp
 260 265 270
 Lys Thr Val Trp Thr His Gly Asn His Tyr His Ala Pro Asn Gly Ser
 275 280 285
 Leu Gly Ala Met His Val Tyr Glu Ser Lys Phe Arg Asn Trp Ser Glu
 290 295 300
 Gly Tyr Ser Asp Phe Asp Arg Gly Ala Tyr Val Ile Thr Phe Ile Pro
 305 310 315 320
 Lys Ser Trp Asn Thr Ala Pro Asp Lys Val Lys Gln Gly Trp Pro
 325 330 335

<210> 49

<211> 407

<212> PRT

<213> Artificial Sequence

<220>

<223> Streptomyces mobaraensis MTG P81453

<400> 49

Met Arg Ile Arg Arg Arg Ala Leu Val Phe Ala Thr Met Ser Ala Val
 1 5 10 15
 Leu Cys Thr Ala Gly Phe Met Pro Ser Ala Gly Glu Ala Ala Ala Asp
 20 25 30
 Asn Gly Ala Gly Glu Glu Thr Lys Ser Tyr Ala Glu Thr Tyr Arg Leu
 35 40 45
 Thr Ala Asp Asp Val Ala Asn Ile Asn Ala Leu Asn Glu Ser Ala Pro
 50 55 60
 Ala Ala Ser Ser Ala Gly Pro Ser Phe Arg Ala Pro Asp Ser Asp Asp
 65 70 75 80
 Arg Val Thr Pro Pro Ala Glu Pro Leu Asp Arg Met Pro Asp Pro Tyr
 85 90 95
 Arg Pro Ser Tyr Gly Arg Ala Glu Thr Val Val Asn Asn Tyr Ile Arg
 100 105 110
 Lys Trp Gln Gln Val Tyr Ser His Arg Asp Gly Arg Lys Gln Gln Met
 115 120 125
 Thr Glu Glu Gln Arg Glu Trp Leu Ser Tyr Gly Cys Val Gly Val Thr
 130 135 140
 Trp Val Asn Ser Gly Gln Tyr Pro Thr Asn Arg Leu Ala Phe Ala Ser
 145 150 155 160
 Phe Asp Glu Asp Arg Phe Lys Asn Glu Leu Lys Asn Gly Arg Pro Arg
 165 170 175
 Ser Gly Glu Thr Arg Ala Glu Phe Glu Gly Arg Val Ala Lys Glu Ser
 180 185 190
 Phe Asp Glu Glu Lys Gly Phe Gln Arg Ala Arg Glu Val Ala Ser Val
 195 200 205
 Met Asn Arg Ala Leu Glu Asn Ala His Asp Glu Ser Ala Tyr Leu Asp

210						215						220					
Asn	Leu	Lys	Lys	Glu	Leu	Ala	Asn	Gly	Asn	Asp	Ala	Leu	Arg	Asn	Glu		
225					230					235					240		
Asp	Ala	Arg	Ser	Pro	Phe	Tyr	Ser	Ala	Leu	Arg	Asn	Thr	Pro	Ser	Phe		
				245						250					255		
Lys	Glu	Arg	Asn	Gly	Gly	Asn	His	Asp	Pro	Ser	Arg	Met	Lys	Ala	Val		
			260					265					270				
Ile	Tyr	Ser	Lys	His	Phe	Trp	Ser	Gly	Gln	Asp	Arg	Ser	Ser	Ser	Ala		
	275						280					285					
Asp	Lys	Arg	Lys	Tyr	Gly	Asp	Pro	Asp	Ala	Phe	Arg	Pro	Ala	Pro	Gly		
290						295					300						
Thr	Gly	Leu	Val	Asp	Met	Ser	Arg	Asp	Arg	Asn	Ile	Pro	Arg	Ser	Pro		
305					310						315				320		
Thr	Ser	Pro	Gly	Glu	Gly	Phe	Val	Asn	Phe	Asp	Tyr	Gly	Trp	Phe	Gly		
				325						330					335		
Ala	Gln	Thr	Glu	Ala	Asp	Ala	Asp	Lys	Thr	Val	Trp	Thr	His	Gly	Asn		
			340					345						350			
His	Tyr	His	Ala	Pro	Asn	Gly	Ser	Leu	Gly	Ala	Met	His	Val	Tyr	Glu		
	355						360					365					
Ser	Lys	Phe	Arg	Asn	Trp	Ser	Glu	Gly	Tyr	Ser	Asp	Phe	Asp	Arg	Gly		
	370				375						380						
Ala	Tyr	Val	Ile	Thr	Phe	Ile	Pro	Lys	Ser	Trp	Asn	Thr	Ala	Pro	Asp		
385					390						395				400		
Lys	Val	Lys	Gln	Gly	Trp	Pro											
				405													

<210> 50
 <211> 5
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Q-Tag 1

<400> 50
 Leu Leu Gln Gly Gly
 1 5

<210> 51
 <211> 4
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Q-Tag 2

<400> 51
 Leu Leu Gln Gly
 1

<210> 52
 <211> 6
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Q-Tag 3

<400> 52
 Leu Ser Leu Ser Gln Gly
 1 5

<210> 53
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Q-Tag 4

<400> 53

Gly Gly Gly Leu Leu Gln Gly Gly
1 5

<210> 54

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 5

<400> 54

Gly Leu Leu Gln Gly
1 5

<210> 55

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 6

<400> 55

Leu Leu Gln
1

<210> 56

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 7

<400> 56

Gly Ser Pro Leu Ala Gln Ser His Gly Gly
1 5 10

<210> 57

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 8

<400> 57

Gly Leu Leu Gln Gly Gly Gly
1 5

<210> 58

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 9

<400> 58
Gly Leu Leu Gln Gly Gly
1 5

<210> 59
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 10

<400> 59
Gly Leu Leu Gln
1

<210> 60
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 11

<400> 60
Leu Leu Gln Leu Leu Gln Gly Ala
1 5

<210> 61
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 12

<400> 61
Leu Leu Gln Gly Ala
1 5

<210> 62
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 13

<400> 62
Leu Leu Gln Tyr Gln Gly Ala
1 5

<210> 63
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 14

<400> 63
Leu Leu Gln Gly Ser Gly
1 5

<210> 64
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 15

<400> 64
Leu Leu Gln Tyr Gln Gly
1 5

<210> 65
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 16

<400> 65
Leu Leu Gln Leu Leu Gln Gly
1 5

<210> 66
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 17

<400> 66
Ser Leu Leu Gln Gly
1 5

<210> 67
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 18

<400> 67
Leu Leu Gln Leu Gln
1 5

<210> 68
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 19

<400> 68
Leu Leu Gln Leu Leu Gln
1 5

<210> 69
<211> 5
<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 20

<400> 69

Leu Leu Gln Gly Arg

1 5

<210> 70

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 21

<400> 70

Glu Glu Gln Tyr Ala Ser Thr Tyr

1 5

<210> 71

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 22

<400> 71

Glu Glu Gln Tyr Gln Ser Thr Tyr

1 5

<210> 72

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 23

<400> 72

Glu Glu Gln Tyr Asn Ser Thr Tyr

1 5

<210> 73

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 24

<400> 73

Glu Glu Gln Tyr Gln Ser

1 5

<210> 74

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 25

<400> 74

Glu Glu Gln Tyr Gln Ser Thr
1 5

<210> 75

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 26

<400> 75

Glu Gln Tyr Gln Ser Thr Tyr
1 5

<210> 76

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 27

<400> 76

Gln Tyr Gln Ser
1

<210> 77

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 28

<400> 77

Gln Tyr Gln Ser Thr Tyr
1 5

<210> 78

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 29

<400> 78

Tyr Arg Tyr Arg Gln
1 5

<210> 79

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Q-Tag 30

<400> 79

Asp Tyr Ala Leu Gln

1 5

<210> 80
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 31

<400> 80
Phe Gly Leu Gln Arg Pro Tyr
1 5

<210> 81
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 32

<400> 81
Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu
1 5 10

<210> 82
<211> 3
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 33

<400> 82
Leu Gln Arg
1

<210> 83
<211> 3
<212> PRT
<213> Artificial Sequence

<220>
<223> Q-Tag 34

<400> 83
Tyr Gln Arg
1