



Office de la Propriété

Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2306053 C 2003/01/21

(11)(21) 2 306 053

(12) BREVET CANADIEN  
CANADIAN PATENT

(13) C

(86) Date de dépôt PCT/PCT Filing Date: 1998/11/16  
(87) Date publication PCT/PCT Publication Date: 1999/05/27  
(45) Date de délivrance/Issue Date: 2003/01/21  
(85) Entrée phase nationale/National Entry: 2000/05/17  
(86) N° demande PCT/PCT Application No.: US 1998/024486  
(87) N° publication PCT/PCT Publication No.: 1999/025855  
(30) Priorités/Priorities: 1997/11/18 (60/065,613) US;  
1997/11/18 (60/065,627) US; 1998/09/08 (60/099,461) US

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> C12N 15/83, A01H 4/00, C12N 5/10,  
C12N 15/84, C12N 15/63

(72) Inventeurs/Inventors:  
BASZCZYNSKI, CHRISTOPHER L., US;  
LYZNIK, LESZEK ALEXANDER, US;  
GORDON-KAMM, WILLIAM J., US;  
GUAN, XUENI, US

(73) Propriétaire/Owner:  
PIONEER HI-BRED INTERNATIONAL, INC., US

(74) Agent: SMART & BIGGAR

(54) Titre : TRANSFERT DE GENOMES VIRAUX PROVENANT DE L'ADN-T AU MOYEN DE SYSTEMES DE  
RECOMBINAISON SPECIFIQUES DE SITES

(54) Title: MOBILIZATION OF VIRAL GENOMES FROM T-DNA USING SITE-SPECIFIC RECOMBINATION SYSTEMS

(57) Abrégé/Abstract:

The invention relates to methods and compositions for site-specific recombinase-mediated mobilization of viral replicons and associated DNAs of interest from T-DNA. The methods of the invention comprise Agrobacterium-mediated transfer recombinase. The nucleic acids of the invention may additionally contain expression cassettes encoding the cognate site-specific recombinase for the target sites flanking the viral genome. The compositions of the invention further comprise Agrobacterium containing the nucleic acids of the invention. The compositions and methods of the invention have use in increasing the efficiency of agroinfection, providing high copy numbers of a DNA of interest for transient expression or for integration into a plant chromosome, and in simplifying the construction and stable maintenance of vectors for agroinfection and transformation.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/83, 15/84, 15/63, 5/10, A01H 4/00</b>		A1	(11) International Publication Number: <b>WO 99/25855</b>
(21) International Application Number: <b>PCT/US98/24486</b>		(43) International Publication Date: <b>27 May 1999 (27.05.99)</b>	
(22) International Filing Date: <b>16 November 1998 (16.11.98)</b>			(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(30) Priority Data: 60/065,627 18 November 1997 (18.11.97) US 60/065,613 18 November 1997 (18.11.97) US 60/099,461 8 September 1998 (08.09.98) US			(71) Applicant (for all designated States except US): <b>PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50319 (US).</b>
(72) Inventors; and (75) Inventors/Applicants (for US only): <b>BASZCZYNSKI, Christopher, L. [US/US]; 7305 Benton Drive, Urbandale, IA 50322 (US). LYZNIK, Leszek, Alexander [PL/US]; 5960 Greendale Place #105, Johnston, IA 50131 (US). GORDON-KAMM, William, J. [US/US]; 3916 67th Street, Urbandale, IA 50322 (US). GUAN, Xueni [CN/US]; 3668 Ruette de Ville, San Diego, CA 82139 (US).</b>			(74) Agents: <b>SPRUILL, W., Murray et al.; Alston &amp; Bird LLP, Bell Seltzer Intellectual Property Law Group, P.O. Drawer 34009, Charlotte, NC 28234 (US).</b>

(54) Title: MOBILIZATION OF VIRAL GENOMES FROM T-DNA USING SITE-SPECIFIC RECOMBINATION SYSTEMS

## (57) Abstract

The invention relates to methods and compositions for site-specific recombinase-mediated mobilization of viral replicons and associated DNAs of interest from T-DNA. The methods of the invention comprise *Agrobacterium*-mediated transfer recombinase. The nucleic acids of the invention may additionally contain expression cassettes encoding the cognate site-specific recombinase for the target sites flanking the viral genome. The compositions of the invention further comprise *Agrobacterium* containing the nucleic acids of the invention. The compositions and methods of the invention have use in increasing the efficiency of agroinfection, providing high copy numbers of a DNA of interest for transient expression or for integration into a plant chromosome, and in simplifying the construction and stable maintenance of vectors for agroinfection and transformation.

## MOBILIZATION OF VIRAL GENOMES FROM T-DNA USING SITE-SPECIFIC RECOMBINATION SYSTEMS

### FIELD OF THE INVENTION

The present invention relates to plant molecular biology. More specifically, the invention relates to methods and compositions for increasing the efficiency of excision of viral replicon from T-DNA that is transferred to a plant by agroinfection.

### BACKGROUND OF THE INVENTION

*Agrobacterium* harboring a Ti or a Ri plasmid can efficiently transfer a portion of these plasmids, the T-DNA, into plant cells at the site of a wound. Transfer of the T-DNA into the plant cell is induced by signal compounds present at the site of a plant wound and requires T-DNA border sequences at both ends of the T-DNA and trans-acting virulence gene products (*vir*) encoded by the Ti or Ri plasmid. The transferred T-DNA is then targeted to the nucleus and integrated into the plant genome. A tumor, crown gall, forms at the site of inoculation in most dicotyledonous plants.

Tumor formation results from the expression of T-DNA oncogenes encoding the growth factors auxin and cytokinin which promote plant cell proliferation. In order for these oncogenes to be expressed, the T-DNA must first integrate into the genome of the plant. Tumor formation is limited to dicotyledonous plants because the T-DNA, while transferred into the cells of monocots, is either not normally integrated into the genome of monocotyledonous plants or is integrated and silenced.

By inserting a viral genome into the T-DNA, *Agrobacterium* can be used to mediate the viral infection of plants. Following transfer of the T-DNA to the plant cell, excision of the viral genome from the T-DNA (mobilization) is required for successful viral infection. This *Agrobacterium*-mediated method for introducing a virus into a plant host is known as agroinfection (for a review, see

Grimsley, "Agroinfection" pp. 325-342, in *Methods in Molecular Biology, vol 44: Agrobacterium Protocols*, ed. Gartland and Davey, Humana Press, Inc., Totowa, NJ; and Grimsley (1990) *Physiol. Plant.* 79:147-153). Upon entry into the plant cell nucleus, a unit

5 length circular viral genome that is capable of initiating systemic infection is mobilized from the T-DNA. Integration of the T-DNA into the plant genome is not required for this event. Two non-exclusive mechanisms, intramolecular homologous recombination and replicative release, have been proposed for this release of circular viral genomes or replication intermediates from T-DNA. The 10 replicative release of viral genomes by the rolling-circle replication mechanism has been demonstrated for the mobilization of geminivirus genomes from T-DNA (Stenger *et al.* (1991) *Proc. Natl. Acad. Sci.* 88:8029-8033). Release of geminivirus via homologous recombination between tandemly repeated genomes has also been demonstrated (Lazarowitz *et al.* (1989) *EMBO J.* 8:1023-1032).

15 Release by either of the above mechanisms requires the presence of tandem directly repeated copies of the viral replicon in the T-DNA. A circular viral replicon may be excised from the T-DNA by intramolecular homologous recombination between the repeated genomes. For replicative release, two origin of replication sequences must be present to initiate and complete the replication 20 process. Both mechanisms of mobilization are complex biochemical processes that may be attenuated by a number of factors which in turn affect the efficiency of viral excision. Furthermore, tandem dimers of viral DNA are often difficult to construct and are unstable in recombination proficient host cells.

Agroinfection has been reported in a number of publications as a 25 successful method for inducing systemic viral infections in plant cells, including monocotyledonous plants such as maize (Heath *et al.* (1997) *Mol. Plant-Microbe Interact.* 10:221-227, Grimsley *et al.* (1989) *Mol. Gen. Genet.* 217:309-316). In many instances, 30 particularly when naked viral nucleic acid is non-infectious, agroinfection is the only way of transforming a plant with cloned viral DNA. Even where naked viral nucleic acid is infectious, agroinfection is frequently used because it is relatively

efficient and does not require the production of large amounts of plasmid or viral DNA. Agroinfection has been used to study viral replication and recombination, in the investigation of viral gene functions, for the production of autonomously replicating viral vectors, for transient expression of genes inserted into T-DNA, for 5 integration of DNA into a plant genome, for production of virus resistant plants, for the study of transposable elements and for the determination of tissue-specific susceptibility to T-DNA transfer.

The development of plant virus gene vectors for expression of foreign genes in plants provides a means to provide high levels of gene expression 10 within a short time. The benefits of virus-based transient RNA and DNA replicons include rapid and convenient engineering coupled with flexibility for expeditious application in various plant species. In this manner, autonomously replicating viruses offer numerous advantages for use as vehicles for transient expression of foreign genes, including their characteristic high levels of multiplication and 15 concomitant levels of transient gene expression. Accordingly, it would be beneficial to provide methods that facilitate the construction of vectors for agroinfection, provide flexibility in designing viral vectors for genetic transformation of plant cells and increase the efficiency of mobilization of viral replicons from T-DNA and the copy number of a DNA sequence of interest 20 associated with the viral replicon.

The present invention accomplishes these objectives by providing methods and compositions for the mobilization of viral replicon from T-DNA via site-specific recombination systems. Site-specific recombination-mediated excision of DNA fragments from chromosomal or extrachromosomal DNA 25 molecules has been described for a number of site-specific recombination systems and plant species. See Russell *et al.* (1992) *Mol. Gen. Genet.* 234:49-59; Lyznik *et al.* (1996) *Nucleic Acids Res.* 24:3784-3789; and Dale *et al.* (1991) *Proc. Natl. Acad. Sci.* 88:10558-10562.

However, mobilization of viral vectors from T-DNA via site-specific 30 recombination has not previously been applied to agro-mediated transformation.

62451-850 (S)

## SUMMARY OF THE INVENTION

The invention relates to methods and compositions for site-specific recombinase-mediated mobilization of viral replicons and associated DNAs of interest from T-DNA. The 5 methods of the invention comprise *Agrobacterium*-mediated transfer of T-DNA to a plant cell, wherein the T-DNA contains a viral replicon flanked by directly repeated target sites for a site-specific recombinase and optionally a DNA of interest linked to the viral replicon. The DNA of interest may also 10 contain a non-identical target site for the recombinase. an expression cassette for the site-specific recombinase is present on the T-DNA or the plant genome, or is transiently introduced into the plant cell. Expression of the site-specific recombinase in the plant cell results in excision of 15 the viral replicon and the associated DNA of interest. The viral replicon and DNA of interest are then replicated to high copy number in the host plant cell.

The compositions of the invention comprise nucleic acids, such as T-DNAs containing a viral DNA flanked by 20 directly repeated target sites for a site-specific recombinase. The nucleic acids of the invention may additionally contain expression cassettes encoding the cognate site-specific recombinase for the target sites flanking the viral genome. The compositions of the invention further comprise 25 *Agrobacterium* containing the nucleic acids of the invention.

The compositions and methods of the invention have use in providing high copy numbers of a DNA of interest for transient expression or for integration into a plant chromosome, in simplifying the construction and stable 30 maintenance of vectors for agro-mediated transformation of plants and in increasing the efficiency of agroinfection.

62451-850 (S)

4a

One aspect of the invention provides a method for mobilizing in a plant cell a viral replicon from a T-DNA, comprising: (a) infecting a plant cell with an *Agrobacterium* carrying an *Agrobacterium* replicon having a T-DNA which 5 contains a viral replicon flanked by directly repeated target sites for a site-specific recombinase; and (b) providing in the cell a nucleotide sequence encoding a site-specific recombinase or an active fragment or variant thereof, the nucleotide sequence being operably linked to a promoter for expressing the 10 recombinase in a plant cell; wherein expression of said recombinase mobilizes the viral replicon.

The invention also provides a T-DNA of an *Agrobacterium* replicon, wherein the T-DNA contains a viral replicon flanked by directly repeated target sites for a site- 15 specific recombinase.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 schematically illustrates a model of mobilization of a viral replicon from T-DNA by *loxP*/*Cre* site-specific recombination.

20 Figure 2 schematically illustrates *Cre*-mediated site-specific excision of a viral DNA and DNA of interest from a T-DNA.

Figure 3 schematically illustrates the integration of DNA into a chromosome by double site-specific recombination.

Figure 4 shows the 5' to 3' sequences of the *loxP* and FRT sites.

Figure 5 schematically illustrates a PCR protocol for construction of 5 a viral DNA flanked by *loxP* target sites and *Xba*I restriction sites.

#### DETAILED DESCRIPTION OF THE INVENTION

The invention is directed to methods and compositions for the mobilization of viral DNA from T-DNA. In the methods of the invention, a viral 10 replicon flanked by directly repeated target sites for a site-specific recombinase is inserted into a T-DNA carried by an *Agrobacterium* replicon. The T-DNA is transferred into the cell of a plant by agro-mediated transfer. Expression of the site-specific recombinase in the infected plant cell will result in excision of a circular viral replicon from the transferred T-DNA (Figure 1). Replication of this 15 viral replicon will result in a high number of copies of the replicon. Systemic infection of the plant may take place if the replicon encodes infectious viral particles. Accordingly, the methods of the invention have use in the production of viral DNA and/or in promoting systemic viral infection of a plant.

Thus, the invention is directed to a method for mobilizing a viral 20 replicon from a T-DNA, comprising:

- a) providing an *Agrobacterium* replicon having a T-DNA, wherein said T-DNA contains a viral replicon flanked by directly repeated target sites for a site-specific recombinase; and
- b) infecting a cell of a plant with an *Agrobacterium* 25 carrying said *Agrobacterium* replicon under conditions that allow transfer of said T-DNA and expression of said recombinase in said cell; wherein said cell, said T-DNA, or said viral replicon contains a nucleotide sequence encoding said recombinase or an active fragment or variant thereof, and said nucleotide sequence is operably linked to a promoter that 30 drives expression in said cell.

If a DNA of interest has been inserted into the viral replicon, or

between the viral replicon and a target site for a site-specific recombinase, the DNA of interest will also be replicated to high copy. The high copy number of the DNA of interest increases the efficiency of integration of this DNA into the genome, or the level of transient expression of a gene encoded by the DNA of interest. Thus, in another aspect, the invention provides a method for providing to a plant cell a plurality of copies of a DNA sequence of interest, comprising:

- 5 a) providing an *Agrobacterium* replicon having a T-DNA, wherein said T-DNA contains in a 5' to 3' or 3' to 5' orientation, a first target site for a site-specific recombinase, a viral replicon, said DNA sequence of interest, and a second target site for said recombinase in direct repeat with said first target site, wherein said first and second target sites are identical; and
- 10 b) infecting a cell of a plant with an *Agrobacterium* carrying said *Agrobacterium* replicon under conditions that allow transfer of said T-DNA and expression of said recombinase in said cell; wherein said cell, said T-DNA, or said viral replicon contains a nucleotide sequence encoding said recombinase or an active fragment or variant thereof, and said nucleotide sequence is operably linked to a promoter that drives expression in said cell.

The methods of the invention are useful for providing a high copy number of a DNA of interest for site-specific integration into a plant chromosome.

- 20 In this case, the plant chromosome will contain one or more target sites for a site-specific recombinase. Depending on the substrates, the recombination event could consist of a single or a double cross-over event. In the simplest case, the plant chromosome and the excised viral replicon each contain a single target site. Site-specific recombination between these two target sites results in the insertion of the
- 25 viral replicon and any associated DNA of interest into the plant chromosome.

The inclusion of a non-identical target site between the viral replicon and the DNA of interest allows integration of the DNA of interest into a plant genome having a corresponding target site, without concomitant insertion of the viral replicon. In this method, the distal ends of the viral replicon and the DNA of interest are flanked by identical target sites. Recombination between the

- 30 identical target sites results in excision of a circular viral replicon containing the

DNA of interest flanked by non-identical target sites for the recombinase (Figure 2). For targeted insertion of the DNA of interest, the same two non-identical target sites are present in the target organisms genome, thereby establishing a target site for insertion of the DNA of interest. A double crossover event resulting from site-specific recombination between the identical target sites on the host genome and the DNA of interest results in the insertion of the DNA of interest into the chromosome of the target organism, free of the viral replicon (Figure 3). Such methods are discussed in detail in Publication WO99/25854.

10

Thus, in a further aspect, the invention provides a method for providing to a plant cell a plurality of copies of a DNA sequence of interest flanked by nonidentical target sites for a site-specific recombinase, comprising:

- a) providing an *Agrobacterium* replicon having a T-DNA, wherein said T-DNA contains in a 5' to 3' or 3' to 5' orientation, a first target site for said recombinase, a viral replicon, a second target site for said recombinase, said DNA sequence of interest, and a third target site for said recombinase, wherein said first and third target sites are directly repeated and identical with respect to each other, and said second target site is nonidentical to said first and third target sites; and
- b) infecting a cell of a plant with an *Agrobacterium* carrying said *Agrobacterium* replicon under conditions that allow transfer of said T-DNA and expression of said recombinase in said cell;  
wherein said cell, said T-DNA, or said viral replicon  
25 contains a nucleotide sequence encoding said or an active fragment or variant thereof, and said nucleotide sequence is operably linked to a promoter that drives expression in said cell.

The compositions of the invention comprise T-DNAs containing a viral replicon flanked by directly repeated target sites for a site-specific recombinase. In another aspect, the compositions of the invention comprise a T-DNA containing in a 5' to 3' or 3' to 5' orientation, a first target site for said

recombinase, a viral replicon, a second target site for said recombinase, said DNA sequence of interest, and a third target site for said recombinase, wherein said first and third target sites are directly repeated and identical with respect to each other, and said second target site is nonidentical to said first and third target sites. The T-

5 DNAs of the invention may further comprise nucleotide sequence encoding a site-specific recombinase, wherein the nucleotide sequence is operably linked to a promoter that drives expression in a plant cell. The compositions of the invention further comprise replicons containing these T-DNAs and Agrobacterium containing these replicons.

10 By "mobilization of a viral replicon" is meant excision of a viral replicon from a T-DNA sequence after it has been transferred to a plant cell. In the methods of the invention, mobilization of viral DNA is accomplished by conservative site-specific recombination between directly repeated target sites flanking the viral DNA. In one embodiment, the resulting product is a circularized 15 viral DNA containing one copy of the target site (see Figure 1).

By "viral replicon" is meant double-stranded DNA from a virus having a double stranded DNA genome or replication intermediate. The excised viral DNA is capable of acting as a replicon or replication intermediate, either independently, or with factors supplied *in trans*. The viral DNA may or may not 20 encode infectious viral particles and furthermore may contain insertions, deletions, substitutions, rearrangements or other modifications. The viral DNA may contain heterologous DNA. In this case, heterologous DNA refers to any non-viral DNA or DNA from a different virus. For example, the heterologous DNA may comprise an expression cassette for a protein or RNA of interest.

25 Viral replicons suitable for use in the methods and compositions of the invention include those of viruses having a circular DNA genome or replication intermediate, such as: Abutilon mosaic virus (AbMV), African cassava mosaic virus (ACMV), banana streak virus (BSV), bean dwarf mosaic (BDMV), bean golden mosaic virus (BGMV), beet curly top virus (BCTV), beet western yellows 30 virus (BWYV) and other luteoviruses, cassava latent virus (CLV), carnation etched virus (CERV), cauliflower mosaic virus (CaMV), chloris striate mosaic (CSMV),

commelina yellow mottle virus (CoYMV), cucumber mosaic virus (CMV), dahlia mosaic virus (DaMV), digitaria streak virus (DSV), figwort mosaic virus (FMV), hop stunt viroid (HSV), maize streak virus (MSV), mirabilis mosaic virus (MMV), miscanthus streak virus (MiSV), potato stunt tuber virus (PSTV),

5      panicum streak virus (PSV), potato yellow mosaic virus (PYMV), rice tungro bacilliform virus (RTBV), soybean chlorotic mottle virus (SoyCMV), squash leaf curl virus (SqLCV), strawberry vein banding virus (SVBV), sugarcane streak virus (SSV), thistle mottle virus (ThMV), tobacco mosaic virus (TMV), tomato golden mosaic virus (TGMV) , tomato mottle virus (TMoV), tobacco ringspot virus

10     (TobRV), tobacco yellow dwarf virus (TobYDV), tomato leaf curl virus (TLCV), tomato yellow leaf curl virus (TYLCV), tomato yellow leaf curl virus-Thailand (TYLCV-t) and wheat dwarf virus (WDV) and derivatives thereof. Preferably the viral replicon is from MSV, WDV, TGMV or TMV.

By "T-DNA" is meant the T-DNA of an *Agrobacterium tumefaciens*

15     Ti plasmid or from an *Agrobacterium rhizogenes* Ri plasmid, or a derivative thereof. The T-DNA may comprise an entire T-DNA, but need only comprise the minimal sequences required *in cis* for transfer (i.e., the right and the left T-DNA border sequences). The T-DNAs of the invention have inserted into them, anywhere between the right and left border sequences, a viral DNA flanked by

20     target sites for a site-specific recombinase. The T-DNA may contain deletions, substitutions and/or additional insertions of DNA other than the viral genome and target sites. Preferably the T-DNA contains an expression cassette for the cognate recombinase of the target sites flanking the viral DNA. The sequences encoding factors required *in trans* for transfer of the T-DNA into a plant cell, such as *vir* genes, may be inserted into the T-DNA, or may be present on the same replicon as the T-DNA, or *in trans* on a compatible replicon in the *Agrobacterium* host. Preferably the *trans* acting factors required for T-DNA transfer are present on the same replicon as the T-DNA.

By "Agrobacterium replicon" is meant any replicon (e.g., plasmid or other vector) that is capable of being stably maintained in an *Agrobacterium* host. Such replicons include the *Agrobacterium* chromosome, *Agrobacterium*

plasmids, cosmids, phagemids, etc., derivatives thereof, and any other vector capable of stable replication in *Agrobacterium*. For example, a binary vector suitable for both *Agrobacterium*-mediated transfer and for facile recombinant manipulations and replication in other organisms is useful in the methods and compositions of the invention.

5 Preferably the replicon is a Ti or Ri plasmid or a derivative thereof.

In the compositions and methods of the invention, the viral replicon is flanked by directly repeated target sites for a site-specific recombinase. By "flanked by" is meant that the target sites may be directly contiguous with the viral DNA or there may be one or more intervening sequences present between one or both ends of the viral DNA 10 and the target sites. Intervening sequences of particular interest would include linkers, adapters, selectable markers and/or other sites which aid in vector construction or analysis and expression cassette for a gene of interest. Target sites for site-specific recombinases are known to those skilled in the art and are discussed in Publication WO99/25854.

15 Examples of target sites include, but are not limited to FRT, FRT1, FRT5 (SEQ ID NO:5), FRT6 (SEQ ID NO:6), FRT7 (SEQ ID NO:7), other FRT mutants, loxP, loxP mutants, and the like. The loxP and FRT sites are shown in Figure 4.

20 By "directly repeated" is meant that the target sites that flank the viral DNA are arranged in the same orientation, so that recombination between these sites results in excision, rather than inversion, of the viral DNA.

By "site-specific recombinase" is meant any enzyme capable of being functionally expressed in plants, that catalyzes conservative site-specific recombination between its corresponding target sites. For reviews of site-specific recombinases, see Sauer (1994) *Current Opinion in Biotechnology* 5:521-527; Sadowski (1993) *FASEB* 7:760-767.

25 Methods of using site-specific recombination systems to excise DNA fragments from chromosomal or extrachromosomal plant DNA are known to those skilled in the art. The bacteriophage *Pl*-*loxP*-*Cre* and the *Saccharomyces* 2 $\mu$  plasmid FRT/FLP site-specific recombinations systems have been extensively studied. For example, Russell *et al.* (1992, *Mol. Gen. 30 Genet.* 234:49-59) describe

the excision of selectable markers from tobacco and *Arabidopsis* genomes using the *loxP*-Cre site-specific recombination system.

It is furthermore recognized that the recombinase, which is used in the invention, will depend upon the target sites of the transformed organism and the targeting cassette. That is, if *FRT* sites are utilized, the FLP recombinase will be needed. In the same manner, where *lox* sites are utilized, the Cre recombinase is required. If the non-identical target sites comprise both a *FRT* and a *lox* site, both the FLP and Cre recombinase or a chimeric recombinase having both Cre and FLP functions, as described in Publication WO99/25840 entitled "A NOVEL METHOD FOR 10 THE INTEGRATION OF FOREIGN DNA INTO EUKARYOTIC GENOMES," will be required in the plant cell.

Methods of the invention comprise the introduction of T-DNA to a plant cell, wherein the T-DNA contains a viral replicon flanked by directly repeated target sites for a site-specific recombinase and optionally a DNA of interest linked to the viral replicon. The DNA of interest may also contain a non-identical target site for the recombinase. An expression cassette for the site-specific recombinase is present on the T-DNA or the plant genome, or is transiently introduced into the plant cell. Expression of the site-specific recombinase in the plant cell results in excision of the viral replicon and the associated DNA of interest. The viral replicon and DNA of interest are then replicated to high copy number in the host plant cell.

The compositions of the invention comprise nucleic acids, such as T-DNAs containing a viral DNA flanked by directly repeated target sites for a site-specific recombinase. The site-specific recombinase may be a naturally occurring recombinase or an active fragment derivative thereof. Site-specific recombinases useful in the methods and compositions of the invention, include recombinases from the integrase and resolvase families, derivatives thereof, and any other naturally occurring or recombinantly produced enzyme or derivative thereof, that catalyzes conservative site-specific recombination between specified DNA sites. The integrase family of recombinases has over thirty members and includes FLP, Cre, Int and R. The resolvase family includes  $\gamma\delta$  resolvase. Recombinant enzymes that catalyze site-specific conservative recombination include moFLP (SEQ ID NOS:1 and 2), described in U.S. patent number 30 5,929,301.

62451-850 (S)

-12-

issued on July 27, 1999, and moCre (SEQ ID NOS:3 and 4). Preferably, the recombinase is one that does 5 not require cofactors or a supercoiled substrate. Such recombinases include Cre, FLP, moflp, and moCre.

10 moflp is derived from the *Saccharomyces* 2 $\mu$  plasmid FLP recombinase, but is encoded by a nucleic acid sequence utilizing maize-preferred codons. While the moflp nucleic acid sequence includes preferred codons for expression of amino acids in maize, it is understood that a useful sequence may contain codons occurring in maize with less than the highest reported maize codon frequencies.

15 The site-specific recombinases and sequences encoding them that are used in the methods and compositions of the invention may be variants of naturally occurring recombinases and the genes encoding them. The term "conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or conservatively modified variants of the amino acid sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and 20 GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations" and represent one species of conservatively modified variation. One of ordinary skill will recognize that each codon in a nucleic acid (except AUG, which is 25 ordinarily the only codon for methionine) can be modified to yield a functionally identical molecule.

As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid

or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Thus, any number of amino acid residues selected from the group of integers consisting of from 1 to 15 can be so altered.

5 Thus, for example, 1, 2, 3, 4, 5, 7, or 10 alterations can be made. Conservatively modified variants typically provide similar biological activity as the unmodified polypeptide sequence from which they are derived. For example, substrate specificity, enzyme activity, or ligand/receptor binding is generally at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% of the native protein for its native substrate.

10 Conservative substitution tables providing functionally similar amino acids are well known in the art.

The following six groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Serine (S), Threonine (T);
- 15 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

20 See Creighton (1984) *Proteins*, W.H. Freeman and Company.

When the nucleic acid is prepared or altered synthetically, advantage can be taken of known codon preferences of the intended host where the nucleic acid is to be expressed. For example, although nucleic acid sequences of the present invention may be expressed in both monocotyledonous and

25 dicotyledonous plant species, sequences can be modified to account for the specific codon preferences and GC content preferences of monocotyledons or dicotyledons as these preferences have been shown to differ (Murray *et al.* (1989) *Nucl. Acids Res.* 17:477-498; and Campbell *et al.* (1990) *Plant Physiol.* 92:1).

Thus, the maize preferred codon for a particular amino acid may be derived from

30 known gene sequences from maize. Maize codon usage for 28 genes from maize plants are listed in Table 4 of Murray *et al.*, *supra*.

The FLP recombinase gene from yeast (*Saccharomyces cerevisiae*) is commercially available in plasmid pOG44 from Stratagene Cloning Systems (11011 North Torrey Pines Road, La Jolla, CA 92037). Similarly, the sequences of many other site specific recombinases and their cognate target sites are publicly 5 or commercially available.

Genes encoding site-specific recombinases, viral DNA, T-DNA and target sites can be made using (a) standard recombinant methods, (b) synthetic techniques, or combinations thereof. Use of cloning vectors, expression vectors, adapters, and linkers is well known in the art and can be found in such references 10 as Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd ed. (Cold Spring Harbor, New York, 1989). A variety of strategies are available for ligating fragments of DNA, the choice of which depends on the nature of the termini of the DNA fragments and which choices can be readily made by those of skill in the art. For a description of various nucleic acids see, for example, Stratagene Cloning 15 Systems, Catalogs 1995, 1996, 1997 (La Jolla, CA); and, Amersham Life Sciences, Inc, Catalog =97 (Arlington Heights, IL). Genes encoding FLP can be obtained, for example, by synthesizing the genes with mutually priming long oligonucleotides. See, for example, Ausubel *et al.* (eds.), *Current Protocols In Molecular Biology*, pages 8.2.8 to 8.2.13, Wiley Interscience (1990). Also, see 20 Wosniak *et al.* (1987) *Gene* 60:115. Moreover, current techniques using the polymerase chain reaction provide the ability to synthesize genes as large as 1.8 kilobases in length (Adang *et al.* (1993) *Plant Mol. Biol.* 21:1131; Bombat *et al.* (1993) *PCR Methods and Applications* 2:266).

Rather than use full length recombinases, functional fragments of 25 site-specific recombinases may be used in the methods and compositions of the invention. Functional fragments of site-specific recombinases can be identified using a variety of techniques such as restriction analysis, Southern analysis, primer extension analysis, and DNA sequence analysis. Primer extension analysis or S1 nuclease protection analysis, for example, can be used to localize the putative start 30 site of transcription of the cloned gene. Ausubel at pages 4.8.1 to 4.8.5; Walmsley *et al.*, *Quantitative and Qualitative Analysis of Exogenous Gene Expression by the*

S1 Nuclease Protection Assay, in Methods In Molecular Biology, Vol. 7: Gene Transfer and Expression Protocols, Murray (ed.), pages 271-281 (Humana Press, Inc. 1991). For examples, functional fragments of the FLP protein may be identified by their ability, upon introduction to cells containing appropriate FRT substrates, to catalyze site-specific recombination (for example, excision of an FRT-flanked sequence which upon removal will activate an assayable marker gene).

The general approach of such functional analysis involves subcloning DNA fragments of a genomic clone, cDNA clone or synthesized gene sequence into an expression vector, introducing the expression vector into a heterologous host, and screening to detect the product of recombination (*i.e.* using restriction analysis to verify the product of recombination at the nucleic acid level, or relying on an assay system for recombination as described above). Methods for generating fragments of a CDNA or genomic clone are well known. Variants of an isolated DNA encoding a site-specific recombinase can be produced by deleting, adding and/or substituting nucleotides. Such variants can be obtained, for example, by oligonucleotide-directed mutagenesis, linker-scanning mutagenesis, mutagenesis using the polymerase chain reaction, and the like. See, for example, Ausubel, pages 8.0.3 - 8.5.9. Also, see generally, McPherson (ed.), Directed Mutagenesis: A Practical approach, (IRL Press, 1991).

The nucleic acid sequence encoding the site-specific recombinase is operably linked to a promoter that drives expression in a plant. As used herein "operably linked" includes reference to a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. Generally, operably linked means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame.

As used herein "promoter" includes reference to a region of DNA upstream from the start of transcription and involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is

a promoter capable of initiating transcription in plant cells. Exemplary plant promoters include, but are not limited to, those that are obtained from plants, plant viruses, and bacteria genes that are expressed in plant cells such as those of *Agrobacterium* or *Rhizobium*. Both heterologous and non-heterologous (i.e., 5 endogenous) promoters can be employed to direct expression of a sequence encoding a site-specific recombinase. The promoter may be constitutive, inducible or tissue specific.

Many different constitutive promoters can be utilized in the instant invention. Exemplary constitutive promoters include the promoters from plant 10 viruses such as the 35S promoter from CaMV (Odell *et al.* (1985) *Nature* 313:810-812) and the promoters from such gene as rice actin (McElroy *et al.* (1990) *Plant Cell* 2:163-171); ubiquitin (Christensen *et al.* (1989) *Plant Mol. Biol.* 12:619-632 and Christensen *et al.* (1992) *Plant Mol. Biol.* 18:675-689); pEMU (Last *et al.* (1991) *Theor. Appl. Genet.* 81:581-588); MAS (Velten *et al.* (1984) *EMBO J.* 15 3:2723-2730); maize H3 histone (Lepetit *et al.* (1992) *Mol. Gen. Genet.* 231: 276-285 and Atanassova *et al.* (1992) *Plant Journal* 2(3):291-300); the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, the ubiquitin 1 promoter, the Smas promoter, the cinnamyl alcohol dehydrogenase promoter (U.S. Patent No. 5,683,439), the Nos promoter, the Pemu promoter, the rubisco 20 promoter, the GRP1-8 promoter, and other transcription initiation regions from various plant genes known to those of skill. The ALS promoter, a XbaI/NcoI fragment 5-prime to the *Brassica napus* ALS3 structural gene (or a nucleotide sequence that has substantial sequence similarity to said XbaI/NcoI fragment), represents a particularly useful constitutive promoter for dicots. (See 25 Pioneer Hi-Bred International US Patent number 5,659,029 (WO96/30530)).

A variety of inducible promoters can be used in the instant invention. See Ward *et al.* (1993) *Plant Mol. Biol.* 22:361-366. Exemplary inducible promoters include that from the ACE1 system which responds to copper 30 (Mett *et al.* (1993) *PNAS* 90:4567-4571); In2 gene from maize which responds to benzenesulfonamide herbicide safeners (Hershey *et al.* (1991) *Mol. Gen. Genetics* 227:229-237 and Gatz *et al.* (1994) *Mol. Gen. Genetics* 243:32-38); the Adh1

promoter which is inducible by hypoxia or cold stress, the Hsp70 promoter which is inducible by heat stress, and the PPDK promoter which is inducible by light; or Tet repressor from Tn10 (Gatz *et al.* (1991) *Mol. Gen. Genet.* 227:229-237. A particularly preferred inducible promoter is a promoter that responds to an 5 inducing agent to which plants do not normally respond. An exemplary inducible promoter is the inducible promoter from a steroid hormone gene the transcriptional activity of which is induced by a glucocorticosteroid hormone (Schena *et al.* (1991) *Proc. Natl. Acad. Sci. U.S.A.* 88:10421).

Examples of promoters under developmental control include 10 promoters that initiate transcription only, or preferentially, in certain tissues, such as leaves, roots, fruit, seeds, or flowers. The operation of a promoter may also vary depending on its location in the genome. Thus, an inducible promoter may become fully or partially constitutive in certain locations.

The site-specific recombinase must be expressed in the plant cell in 15 order for mobilization of the viral DNA from the T-DNA. Accordingly, the recombinase protein must be expressed so that it is present in the cell at an effective concentration sometime between when the T-DNA is transferred to the plant cell and before non-integrated T-DNA is lost from the plant cells.

Accordingly, the expression cassette encoding the site-specific 20 recombinase may be supplied on the T-DNA *in cis* to the viral DNA; *in trans* on a plant chromosome or extrachromosomal replicon; or may be transferred to the plant near to the time of agro-mediated transformation. The site-specific recombinase gene may be constitutively or transiently expressed, or may be inducible. If cofactors are required for a particular site-specific recombination 25 system, they may be encoded on the T-DNA of the invention or in the plant host chromosomal or extrachromosomal DNA.

By "target site for a site-specific recombinase" is meant a DNA sequence that is recognized by a particular site-specific recombinase. A variety of target sites are known to those skilled in the art and may be used in the methods 30 and compositions of the invention. The site may have the sequence of the cognate site for a given recombinase, or may be modified, so long as it is capable of acting

62451-850 (S)

-18-

as a target site. The site may be contain the minimal sequences necessary for recombination, or it may contain additional sequences that enhance recombination. Examples of target sites for use in the invention are known in the art and include FRT and *loxP* sites (See, for example, Schlake and Bode (1994) *Biochemistry* 5: 33:12746-12751; Huang *et al.* (1991) *Nucleic Acids Research* 19:443-448; Paul D. Sadowski (1995) *In Progress in Nucleic Acid Research and Molecular Biology* vol. 51, pp. 53-91; Michael M. Cox (1989) *In Mobile DNA*, Berg and Howe (eds) American Society of Microbiology, Washington D.C., pp. 116-670; Dixon *et al.* (1995) 18:449-458; Umlauf and Cox (1988) *The EMBO Journal* 7:1845-1852; 10 Buchholz *et al.* (1996) *Nucleic Acids Research* 24:3118-3119; Kilby *et al.* (1993) *Trends Genet.* 9:413-421; Rossant and Geagy (1995) *Nat. Med.* 1: 592-594; *Lox* Albert *et al.* (1995) *The Plant J.* 7:649-659; Bayley *et al.* (1992) *Plant Mol. Biol.* 18:353-361; Odell *et al.* (1990) *Mol. Gen. Genet.* 223:369-378; and Dale and Ow (1991) *Proc. Natl. Acad. Sci. USA* 88:10558-105620; Qui *et al.* (1994) *Proc. Natl. 15 Acad. Sci. USA* 91:1706-1710; Stuurman *et al.* (1996) *Plant Mol. Biol.* 32:901-913; Hartley *et al.* (1980) *Nature* 286: 860-864; Sauer (1994) *Current Opinion in Biotechnology* 5:521-527; and Dale *et al.* (1990) *Gene* 91:79-85).

Each *loxP* and FRT site contains two 13 base pair inverted repeats 20 which flank an 8 base pair spacer. The FRT site contains an additional non-essential 13 base pair repeat. The sequences of the *loxP* and FRT sites are shown in Figure 4. A minimal FRT site comprising two 13 base pair repeats, separated by an 8 base spacer, is:

5'-GAAGTTCCCTATTC[TCTAGAAA]GTATAGGAACCTC3'

25 wherein the nucleotides within the brackets indicate the spacer region. The nucleotides in the spacer region can be replaced with a combination of nucleotides, so long as the two 13-base repeats are separated by eight nucleotides. FLP is a conservative, site-specific recombinase, capable of catalyzing inversion of a nucleic acid sequence positioned between two inversely oriented FRTs; 30 recombination between two molecules each containing a FRT site; and excision between FRT sites. The core region is not symmetrical, and its asymmetry dictates

the directionality of the reaction. Recombination between inverted FRT sites causes inversion of a DNA sequence between them, whereas recombination between directly oriented sites leads to excision of the DNA between them.

T-DNAs containing viral DNA flanked by recombinations sites, 5 expression cassettes for site-specific recombinases and vectors carrying these sequences may be constructed using standard molecular biology techniques. See, for example, Sambrook *et al.* (eds.) *Molecular Cloning: A Laboratory Manual*, Second Edition, (Cold Spring Harbor Laboratory Press, cold Spring Harbor, NY 1989).

10 Typical vectors useful in the methods and compositions of the invention are well known in the art and include vectors derived from the tumor-inducing (Ti) plasmid of *Agrobacterium tumefaciens* described by Rogers *et al.* (1987) *Meth. in Enzymol.*, 153:253-277. These vectors are plant integrating vectors in that on transformation, the vectors integrate a portion of vector DNA 15 into the genome of the host plant. Exemplary *A. tumefaciens* vectors useful herein are plasmids pKYLX6 and pKYLX7 of Schardl *et al.*, *Gene*, 61:1-11 (1987) and Berger *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 86:8402-8406 (1989). Another useful vector herein is plasmid pBI101.2 that is available from Clontech Laboratories, Inc. (Palo Alto, CA).

20 Techniques for transforming a wide variety of higher plant species are well known and described in the technical, scientific, and patent literature. See, for example, Weising *et al.*, *Ann. Rev. Genet.* 22: 421-477 (1988). These methods are useful for transforming a plant cell with a site-specific recombinase expression cassette. This step will be necessary if the cassette is not included in the 25 transferred T-DNA. The expression cassette encoding the site-specific recombinase may be present in the plant genome prior to Agroinfection or may be transformed into the plant around the time of transfer of the T-DNA to the plant cell so that it will be transiently expressed.

For example, the DNA construct may be introduced directly into the genomic 30 DNA of the plant cell using techniques such as electroporation, PEG poration, particle bombardment, silicon fiber delivery, microinjection of plant cell

protoplasts or embryogenic callus, or *Agrobacterium*-mediated transformation (Hiei *et al.* (1994) *Plant J.* 6:271-282).

The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski *et al.*, *Embo J.* 3: 2717-2722 (1984).

5 Electroporation techniques are described in Fromm *et al.*, *Proc. Natl. Acad. Sci.* 82: 5824 (1985). Ballistic transformation techniques are described in Klein *et al.*, *Nature* 327: 70-73 (1987).

DNA can also be introduced into plants by direct DNA transfer into pollen as described by Zhou *et al.*, *Methods in Enzymology*, 101:433 (1983); D. 10 Hess, *Intern Rev. Cytol.*, 107:367 (1987); Luo *et al.*, *Plane Mol. Biol. Reporter*, 6:165 (1988). Expression of polypeptide coding genes can be obtained by injection of the DNA into reproductive organs of a plant as described by Pena *et al.*, *Nature*, 325.:274 (1987). DNA can also be injected directly into the cells of immature embryos and the rehydration of desiccated embryos as described by Neuhaus *et al.*, 15 *Theor. Appl. Genet.*, 75:30 (1987); and Benbrook *et al.*, in *Proceedings Bio Expo* 1986, Butterworth, Stoneham, Mass., pp. 27-54 (1986). A variety of plant viruses that can be employed as vectors are known in the art and include cauliflower mosaic virus (CaMV), geminivirus, brome mosaic virus, and tobacco mosaic virus.

Plants cells stably transformed with a site-specific recombinase 20 expression cassette can be regenerated, e.g., from single cells, callus tissue or leaf discs according to standard plant tissue culture techniques. It is well known in the art that various cells, tissues, and organs from almost any plant can be successfully cultured to regenerate an entire plant. Plant regeneration from cultured protoplasts is described in Evans *et al.*, *Protoplasts Isolation and Culture, Handbook of Plant* 25 *Cell Culture*, Macmillan Publishing Company, New York, pp. 124-176 (1983); and Binding, *Regeneration of Plants, Plant Protoplasts*, CRC Press, Boca Raton, pp. 21-73 (1985).

The regeneration of plants containing the recombinase gene introduced by *Agrobacterium* from leaf explants can be achieved as described by 30 Horsch *et al.*, *Science*, 227:1229-1231 (1985). In this procedure, transformants are grown in the presence of a selection agent and in a medium that induces the

regeneration of shoots in the plant species being transformed as described by Fraley *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 80:4803 (1983). This procedure typically produces shoots within two to four weeks and these transformant shoots are then transferred to an appropriate root-inducing medium containing the 5 selective agent and an antibiotic to prevent bacterial growth. Transgenic plants of the present invention may be fertile or sterile.

Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee *et al.*, *Ann. Rev. of Plant Phys.* 38: 467-486 (1987). The regeneration of 10 plants from either single plant protoplasts or various explants is well known in the art. See, for example, *Methods for Plant Molecular Biology*, A. Weissbach and H. Weissbach, eds., Academic Press, Inc., San Diego, Calif. (1988). This regeneration and growth process includes the steps of selection of transformant cells and shoots, rooting the transformant shoots and growth of the plantlets in soil. For maize cell 15 culture and regeneration see generally, *The Maize Handbook*, Freeling and Walbot, Eds., Springer, New York (1994); *Corn and Corn Improvement*, 3<sup>rd</sup> edition, Sprague and Dudley Eds., American Society of Agronomy, Madison, Wisconsin (1988).

One of skill will recognize that after the site-specific recombinase 20 expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

The methods and compositions of the invention are useful to 25 mobilize viral DNA from T-DNAs transferred by Agroinfection to any plant host. As used herein, the term "plant" includes reference to whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds and plant cells and progeny of same. Plant cell, as used herein includes, without limitation, seeds suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, 30 gametophytes, sporophytes, pollen, and microspores. The class of plants which can be used in the methods of the invention is generally as broad as the class of

62451-850 (S)

-22-

higher plants amenable to *Agrobacterium* transformation techniques, including both monocotyledonous and dicotyledonous plants. A particularly preferred monocot is maize. Other monocots of particular interest include wheat, rice, barley, sorghum and rye. Dicots of particular interest include soybean, *Brassica*,  
 5 sunflower, alfalfa, and safflower.

The T-DNA containing the viral DNA flanked by sites for a site-specific recombinase is transferred to a plant cell by agroinfection. *Agrobacterium tumefaciens*-mediated transformation techniques are well described in the scientific literature. See, for example Horsch *et al.*, *Science* 233: 496-498 (1984),  
 10 Fraley *et al.*, *Proc. Natl. Acad. Sci.* 80: 4803 (1983) and Kado, (1991), *Crit. Rev. Plant Sci.* 10:1. Descriptions of the *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer are provide in Gruber *et al.*, *supra*; Miki, *et al.*, *supra*; and Moloney *et al.* (1989), *Plant Cell Reports* 8:238. Although *Agrobacterium* is useful primarily in dicots, certain monocots can be  
 15 transformed by *Agrobacterium*. For instance, *Agrobacterium* transformation of maize is described in U.S. Patent No. 5,550,318. Other methods of agroinfection include *Agrobacterium rhizogenes*-mediated transformation (see, e.g., Lichtenstein and Fuller In: *Genetic Engineering*, vol. 6, PWJ Rigby, Ed., London, Academic Press, 1987; and Lichtenstein, C. P., and Draper, J., In: *DNA Cloning*, Vol. II, D.  
 20 M. Glover, Ed., Oxford, IRI Press, 1985), Application PCT/US87/02512 (WO 88/02405 published Apr. 7, 1988) describes the use of *A. rhizogenes* strain A4 and its Ri plasmid along with *A. tumefaciens* vectors pARC8 or pARC16.

Optimized methods and vectors for *Agrobacterium*-mediated transformation of plants in the family Graminae, such as rice and maize have been  
 25 described by Heath *et al.* (1997) *Mol. Plant-Microbe Interact.* 10:221-227; Hiei *et al.* (1994) *Plant J.* 6:271-282 and Ishida *et al.* (1996) *Nat. Biotech.* 14:745-750. The efficiency of maize transformation is affected by a variety of factors including the types and stages of tissue infected, the concentration of *Agrobacterium*, the tissue culture media, the Ti vectors and the maize genotype. Super binary vectors  
 30 carrying the vir genes of *Agrobacterium* strains A281 and A348 are useful for high efficiency

transformation of monocots. However, even without the use of high efficiency vectors, it has been demonstrated that T-DNA is transferred to maize at an efficiency that results in systemic infection by viruses introduced by agroinfection, although tumors are not formed (Grimsley *et al.* (1989) *Mol. Gen. Genet.* 217:309-5 316, the contents of which are incorporated herein by reference). This is because integration of the T-DNA containing the viral genome is not required for viral multiplication, since the excised viral genome acts as an independent replicon.

Another useful basic transformation protocol involves a combination of wounding by particle bombardment, followed by use of 10 *Agrobacterium* for DNA delivery, as described by Bidney *et al.* (1992) *Plant Mol. Biol.* 18:301-313. In general, the intact meristem transformation method involves imbibing seed for 24 hours in the dark, removing the cotyledons and root radical, followed by culturing of the meristem explants. Twenty-four hours later, the primary leaves are removed to expose the apical meristem. The explants are 15 placed apical dome side up and bombarded, *e.g.*, twice with particles, followed by co-cultivation with *Agrobacterium*. To start the co-cultivation for intact meristems, *Agrobacterium* is placed on the meristem. After about a 3-day co-cultivation period the meristems are transferred to culture medium with selection agents.

20 The split meristem method involves imbibing seed, breaking of the cotyledons to produce a clean fracture at the plane of the embryonic axis, excising the root tip and then bisecting the explants longitudinally between the primordial leaves. The two halves are placed cut surface up on the medium then bombarded twice with particles, followed by co-cultivation with *Agrobacterium*. For split 25 meristems, after bombardment, the meristems are placed in an *Agrobacterium* suspension for 30 minutes. They are then removed from the suspension onto solid culture medium for three day co-cultivation. After this period, the meristems are transferred to fresh medium with cefotaxime (plus selection agents).

30 The excision of viral DNA from the T-DNA may be determined by a variety of means including but not limited to observation of viral lesions or other signs of viral infection on the plant, by antibody-based detection of viral proteins,

by amplification of PCR products specific to a circular genome, or by Northern analysis for viral RNA. In addition, assays for FLP recombinase activity are known and generally measure the overall activity of the enzyme on DNA substrates containing FRT sites. In this manner, a frequency of excision of the 5 sequence can be determined. Excision of DNA from a linear molecule or intermolecular recombination frequency induced by the enzyme may be assayed, as described, for example, in Babineau *et al.* (1985) *JBC* 260:12313; Meyer-Leon *et al.* (1987) *NA Res* 15:6469; and Gronostajski *et al.* (1985) *JBC* 260:12328.

10 The following examples are offered by way of illustration not by way of limitation.

## EXPERIMENTAL

### Example 1

15 Construction of T-DNA Vectors Containing Viral DNA Flanked By Site-Specific Recombination Sites

DNA fragments containing viral DNA flanked by FRT or *loxP* site-specific recombination sites are constructed either by synthesizing, annealing and 20 ligating complementary oligonucleotides or by creating primers for PCR amplification of a DNA product containing the FRT or *loxP* site and restriction sites useful for cloning into T-DNA near the 5' end of the PCR product.

For example, long PCR primers may be designed wherein the 3' end of the primer hybridizes to the 5' end of the viral genome of interest and the 5' end 25 of the primer further contains a *loxP* site and useful cloning sites (see Figure 5). The resulting PCR product is digested with the appropriate restriction enzyme and inserted into the T-DNA of an *Agrobacterium* binary vector as described by Bevan, M. (1984) *Nucl. Acids Res.* 12:8711-8721.

An expression cassette for Cre is also inserted into the T-DNA. The 30 resulting constructs are transformed into *E. coli*, screened, and then transferred to *Agrobacterium*.

Example 2

Transfer of T-DNA to Dicotyledonous Plant Cells

5                   *Nicotiana tabacum* BY-2 cells are propagated in Murashige and Skoog medium (Gibco BRL) containing 3% sucrose, 1  $\mu$ g/ml thiamine, 0.2  $\mu$ g/ml 2,4-D, and 370  $\mu$ g/ml  $\text{KH}_2\text{PO}_4$ . *Zea mays* Black Mexican Sweet (BMS) cells are propagated in Murashige and Skoog medium containing 2% sucrose, 2  $\mu$ g/ml 2,4-D, 0.2 mg/ml myoinositol, 0.13 mg/ml L-asparagine, 0.13  $\mu$ g/ml nicotinic acid, 10 and 0.25  $\mu$ g/ml each of thiamine, pyridoxine, and pantothenic acid. The cultures are shaken at 140 rpm at 25°C in continuous light.

To infect plant cells, virulence (*vir*) gene activity is induced by treatment with acetosyringone in *Agrobacterium* carrying the vector constructed in Example 1. *Agrobacterium* cells are grown to a density of  $2 \times 10^9$  cells per ml (A = 100, using a Klett-Summerson spectrophotometer, red filter) in AB-sucrose medium. The cells are centrifuged at 10,000g, suspended at a concentration of  $1 \times 10^9$  cells per ml (A = 50) in induction medium (AB salts, 0.5% glucose, 2mM sodium phosphate, 50 Mm Mes, Ph 5.6, 50°C acetosyringone), and incubated with gentle shaking at 25°C for 14 to 18 hr. After washing the bacterial cells in plant 15 culture medium, plant cells are inoculated with induced *Agrobacterium* (-20 bacterial cells per plant cell, except where noted otherwise) and cocultivated at 25°C with shaking at 140 rpm for various periods of time. Most of the bacteria is washed off by centrifugation of the cocultivation mixture at 300 rpm (model 20 GLC-2 clinical centrifuge; Beckman Sorvall, Newtown, CT) for 2 min. The plant cell pellet is suspended and washed once more in plant culture medium and then 25 resuspended in culture containing either 100  $\mu$ g/ml timentin or 200  $\mu$ g/mL cefotaxime. Mobilization of viral genomes from the T-DNA is monitored by quantitating infectious viral particles.

Example 3

Transfer of T-DNA to Monocotyledonous Plants

5 Prior to inoculation, strains of *Agrobacterium* carrying the vector constructed in Example 1 are streaked out on YEB (Grimsley *et al.*(1986) *Proc. Natl. Acad. Sci. USA*, 83:3282-3286) plates solidified with 1.5% agar and supplemented with 100 $\mu$ g/ml rifampicin and 25 $\mu$ g/ml kanamycin and allowed to grow for 48h. A single colony is used to inoculate 10 ml of liquid YEB medium  
10 in a 100ml Erlenmeyer flask supplemented with antibiotics as previously. Growth is continued with shaking at 200 r.p.m. for 24h, then 500 $\mu$ l of this culture is used to inoculate a similar flask and growth continued for a further 20h. This procedure yields a final density of viable *Agrobacterium* cells in the region of 10<sup>9</sup>/ml (estimated by plating). The cells are then harvested by centrifugation and  
15 resuspended in an equal volume of 10mM MgSO<sub>4</sub> without antibiotics; such a suspension is subsequently referred to as undiluted or 10<sup>0</sup> dilution; for experiments involving a dilution series 10mM MgSO<sub>4</sub> was also used as the diluent.

Maize seeds for 10-day old plants are sown in pots in a phytotron  
20 in a 12 hour light/dark cycle at 25°C in a light intensity of about 10000 lux (Sylvania 215W fluorescent lamps type F96T12/CW/VHO) then moved to the BL3 containment laboratory immediately prior to inoculation; subsequent growth conditions have been described (Grimsley *et al.*(1987) *Nature* 325:177-179). Three-day old seedlings are prepared by (i) sterilization by stirring for 20min in  
25 0.7% calcium hypochlorite solution, (ii) washing three times (stirring for 20min each time) in sterile distilled water (iii) preparing 9cm diameter presterilized Petri dishes with 3 sheets of sterile 8.5cm diameter Macherey-Nagel (Germany) filter paper in the bottom and *ca.* 10ml of sterile water per dish, (iv) putting *ca.* 20 seeds into each geranium dish, and (v) incubating in the dark at 28EC for 3 days, or until  
30 the distance between the scutellar node and the apical tip of the coleoptiles is 1-2 cm.

For inoculation of plants, a 50 $\mu$ l or a 100 $\mu$ l Hamilton syringe fitted with a 0.4mm diameter disposable needle is loaded with the bacterial suspension

-27-

avoiding trapped air bubbles. Between inoculations with different bacterial strains the needle is discarded and the syringe flushed out 3 times with 100% ethanol and 3 times with sterile distilled water. 10-day old plants are inoculated by (i) abrasion of an upper leaf, applying 20 $\mu$ l of suspension, and rubbing in with carborundum powder until the leaf appears wet all over, (ii) injection of 10 $\mu$ l of bacterial suspension into the central part of the plant either just above the first leaf blade, or 1cm below the first leaf blade, or at the base of the plant, in the meristematic region where adventitious roots later begin to appear. Three-day old seedlings are injected with 10 $\mu$ l of bacterial suspension in different ways by (i) pushing the needle down through the apical tip of the coleoptile to the coleoptilar node, (ii) injecting 2mm below the apical tip of the coleoptile, (iii) 2 mm above the coleoptilar node, (iv) at the coleoptilar node, (iv) 2mm below the coleoptilar node, (v) at the scutellar node, and by pushing the needle up through the primary root to a region close to the scutellar node. Ten  $\mu$ l is used as a standard inoculum of bacterial suspension, but only 1-2 $\mu$ l routinely remains in the inoculation site, the rest is forced out, usually coming out from the point of entry of the inoculating needle. Following inoculation seedlings are planted immediately in damp soil, incubated as before (Grimsley *et al.*(1987) *Nature* 325:177-179), and observed daily for the appearance of symptoms of viral infection, characterized by the appearance of yellow spots and/or stripes at the base of new leaves.

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains.

25

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the 30 scope of the appended claims.

## SEQUENCE LISTING

<110> Baszczynski, Christopher L.

Lyznik, Leszek

Gordon-Kamm, William J.

Guan, Xueni

<120> Mobilization of Viral Genomes From T-DNA

Using Site-Specific Recombination Systems

<130> 5718-61-1-PC

10 <140> PCT/US98/24486

<141> 1998-11-16

<160> 7

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1272

<212> DNA

<213> Artificial Sequence

<220>

<223> moFLP:

20 FLP recombinase with maize preferred codons

<221> CDS

<222> (1) ... (1272)

<400> 1

atg ccc cag ttc gac atc ctc tgc aag acc ccc ccc aag gtg ctc gtg  
 Met Pro Gln Phe Asp Ile Leu Cys Lys Thr Pro Pro Lys Val Leu Val  
 1 5 10 15

48

agg cag ttc gtg gag agg ttc gag agg ccc tcc ggc gag aag atc gcc  
 Arg Gln Phe Val Glu Arg Phe Glu Arg Pro Ser Gly Glu Lys Ile Ala  
 20 25 30

96

ctc tgc gcc gcc gag ctc acc tac ctc tgc tgg atg atc acc cac aac  
 Leu Cys Ala Ala Glu Leu Thr Tyr Leu Cys Trp Met Ile Thr His Asn  
 35 40 45

144

ggc acc gcc att aag agg gcc acc ttc atg tca tac aac acc atc atc  
 Gly Thr Ala Ile Lys Arg Ala Thr Phe Met Ser Tyr Asn Thr Ile Ile  
 50 55 60

192

	tcc aac tcc ctc tcc ttc gac atc gtg aac aag tcc ctc cag ttc aaa		240
	Ser Asn Ser Leu Ser Phe Asp Ile Val Asn Lys Ser Leu Gln Phe Lys		
	65 70 75 80		
	tac aag acc cag aag gcc acc atc ctc gag gcc tcc ctc aag aag ctc		288
	Tyr Lys Thr Gln Lys Ala Thr Ile Leu Glu Ala Ser Leu Lys Leu		
	85 90 95		
10	atc ccc gcc tgg gag ttc acc atc atc ccc tac tac ggc cag aag cac		336
	Ile Pro Ala Trp Glu Phe Thr Ile Ile Pro Tyr Tyr Gly Gln Lys His		
	100 105 110		
	cag tcc gac atc acc gac atc gtg tca tcc ctc cag ctt cag ttc gag		384
	Gln Ser Asp Ile Thr Asp Ile Val Ser Ser Leu Gln Leu Gln Phe Glu		
	115 120 125		
	tcc tcc gag gag gct gac aag ggc aac tcc cac tcc aag aag atg ctg		432
	Ser Ser Glu Glu Ala Asp Lys Gly Asn Ser His Ser Lys Lys Met Leu		
20	130 135 140		
	aag gcc ctc ctc tcc gag ggc gag tcc atc tgg gag atc acc gag aag		480
	Lys Ala Leu Leu Ser Glu Gly Glu Ser Ile Trp Glu Ile Thr Glu Lys		
	145 150 155 160		
	atc ctc aac tcc ttc gag tac acc tcc agg ttc act aag acc aag acc		528
	Ile Leu Asn Ser Phe Glu Tyr Thr Ser Arg Phe Thr Lys Thr Lys Thr		
	165 170 175		
30	ctc tac cag ttc ctc ttc ctc gcc acc ttc atc aac tgc ggc agg ttc		576
	Leu Tyr Gln Phe Leu Phe Leu Ala Thr Phe Ile Asn Cys Gly Arg Phe		
	180 185 190		
	tca gac atc aag aac gtg gac ccc aag tcc ttc aag ctc gtg cag aac		624
	Ser Asp Ile Lys Asn Val Asp Pro Lys Ser Phe Lys Leu Val Gln Asn		
	195 200 205		
	aag tac ctc ggc gtg atc atc cag tgc ctc gtg acc gag acc aag acc		672
	Lys Tyr Leu Gly Val Ile Ile Gln Cys Leu Val Thr Glu Thr Lys Thr		
40	210 215 220		
	tcc gtg tcc agg cac atc tac ttc ttc tcc gct cgc ggc agg atc gac		720
	Ser Val Ser Arg His Ile Tyr Phe Ser Ala Arg Gly Arg Ile Asp		
	225 230 235 240		
	ccc ctc gtg tac ctc gac gag ttc ctc agg aac tca gag ccc gtg ctc		768
	Pro Leu Val Tyr Leu Asp Glu Phe Leu Arg Asn Ser Glu Pro Val Leu		
	245 250 255		
50	aag agg gtg aac agg acc ggc aac tcc tcc aac aag cag gag tac		816
	Lys Arg Val Asn Arg Thr Gly Asn Ser Ser Asn Lys Gln Glu Tyr		
	260 265 270		
	cag ctc ctc aag gac aac ctc gtg agg tcc tac aac aag gcc ctc aag		864
	Gln Leu Leu Lys Asp Asn Leu Val Arg Ser Tyr Asn Lys Ala Leu Lys		
	275 280 285		
	aag aac gcc ccc tac tcc atc ttc gcc atc aag aac ggc ccc aag tcc		912
	Lys Asn Ala Pro Tyr Ser Ile Phe Ala Ile Lys Asn Gly Pro Lys Ser		
60	290 295 300		

305	cac atc ggt agg cac ctc atg acc tcc ttc ctc tca atg aag ggc ctc His Ile Gly Arg His Leu Met Thr Ser Phe Leu Ser Met Lys Gly Leu 310	315	320	960
325	acc gag ctc acc aac gtg gtg ggc aac tgg tcc gac aag agg gcc tcc Thr Glu Leu Thr Asn Val Val Gly Asn Trp Ser Asp Lys Arg Ala Ser 330	335		1008
10	1056	gcc gtg gcc agg acc acc tac acc cac cag atc acc gcc atc ccc gac Ala Val Ala Arg Thr Thr Tyr Thr His Gln Ile Thr Ala Ile Pro Asp 340	345	350
355	cac tac ttc gcc ctc gtg tca agg tac tac gcc tac gac ccc atc tcc His Tyr Phe Ala Leu Val Ser Arg Tyr Tyr Ala Tyr Asp Pro Ile Ser 360	365		1104
20	1152	aag gag atg atc gcc ctc aag gac gag act aac ccc atc gag gag tgg Lys Glu Met Ile Ala Leu Lys Asp Glu Thr Asn Pro Ile Glu Glu Trp 370	375	380
385	cag cac atc gag cag ctc aag ggc tcc gcc gag ggc tcc atc agg tac Gln His Ile Glu Gln Leu Lys Gly Ser Ala Glu Gly Ser Ile Arg Tyr 390	395	400	1200
405	ccc gcc tgg aac ggc atc atc tcc cag gag gtg ctc gac tac ctc tcc Pro Ala Trp Asn Gly Ile Ile Ser Gln Glu Val Leu Asp Tyr Leu Ser 410	415		1248
30	1272	tcc tac atc aac agg agg atc tga Ser Tyr Ile Asn Arg Arg Ile 420		

&lt;210&gt; 2

&lt;211&gt; 423

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;400&gt; 2

40	40	Met Pro Gln Phe Asp Ile Leu Cys Lys Thr Pro Pro Lys Val Leu Val 1 5 10 15
	20	Arg Gln Phe Val Glu Arg Phe Glu Arg Pro Ser Gly Glu Lys Ile Ala 25 30
	35	Leu Cys Ala Ala Glu Leu Thr Tyr Leu Cys Trp Met Ile Thr His Asn 40 45
50	50	Gly Thr Ala Ile Lys Arg Ala Thr Phe Met Ser Tyr Asn Thr Ile Ile 55 60
	65	Ser Asn Ser Leu Ser Phe Asp Ile Val Asn Lys Ser Leu Gln Phe Lys 70 75 80

85	90	95
Ile Pro Ala Trp Glu Phe Thr Ile Ile Pro Tyr Tyr Gly Gln Lys His		
100	105	110
Gln Ser Asp Ile Thr Asp Ile Val Ser Ser Leu Gln Leu Gln Phe Glu		
115	120	125
Ser Ser Glu Glu Ala Asp Lys Gly Asn Ser His Ser Lys Lys Met Leu		
130	135	140
Lys Ala Leu Leu Ser Glu Gly Glu Ser Ile Trp Glu Ile Thr Glu Lys		
145	150	160
Ile Leu Asn Ser Phe Glu Tyr Thr Ser Arg Phe Thr Lys Thr Lys Thr		
165	170	175
Leu Tyr Gln Phe Leu Phe Leu Ala Thr Phe Ile Asn Cys Gly Arg Phe		
180	185	190
Ser Asp Ile Lys Asn Val Asp Pro Lys Ser Phe Lys Leu Val Gln Asn		
195	200	205
Lys Tyr Leu Gly Val Ile Ile Gln Cys Leu Val Thr Glu Thr Lys Thr		
210	215	220
Ser Val Ser Arg His Ile Tyr Phe Phe Ser Ala Arg Gly Arg Ile Asp		
225	230	240
Pro Leu Val Tyr Leu Asp Glu Phe Leu Arg Asn Ser Glu Pro Val Leu		
245	250	255
Lys Arg Val Asn Arg Thr Gly Asn Ser Ser Asn Lys Gln Glu Tyr		
260	265	270
Gln Leu Leu Lys Asp Asn Leu Val Arg Ser Tyr Asn Lys Ala Leu Lys		
275	280	285
Lys Asn Ala Pro Tyr Ser Ile Phe Ala Ile Lys Asn Gly Pro Lys Ser		
290	295	300
His Ile Gly Arg His Leu Met Thr Ser Phe Leu Ser Met Lys Gly Leu		
305	310	320
Thr Glu Leu Thr Asn Val Val Gly Asn Trp Ser Asp Lys Arg Ala Ser		
325	330	335
Ala Val Ala Arg Thr Thr Tyr Thr His Gln Ile Thr Ala Ile Pro Asp		
340	345	350
His Tyr Phe Ala Leu Val Ser Arg Tyr Tyr Ala Tyr Asp Pro Ile Ser		
355	360	365
Lys Glu Met Ile Ala Leu Lys Asp Glu Thr Asn Pro Ile Glu Glu Trp		
370	375	380
Gln His Ile Glu Gln Leu Lys Gly Ser Ala Glu Gly Ser Ile Arg Tyr		
385	390	400

Pro Ala Trp Asn Gly Ile Ile Ser Gln Glu Val Leu Asp Tyr Leu Ser  
 405 410 415

Ser Tyr Ile Asn Arg Arg Ile  
 420

<210> 3

10 <211> 1032

<212> DNA

<213> Artificial Sequence

<220>

<223> moCre:

Cre recombinase with maize preferred codons

<221> CDS

<222> (1)...(1032)

<400> 3

20 atg tcc aac ctg ctc acg gtt cac cag aac ctt ccg gct ctt cca gtg 48  
 Met Ser Asn Leu Leu Thr Val His Gln Asn Leu Pro Ala Leu Pro Val  
 1 5 10 15

gac gcg acg tcc gat gaa gtc agg aag aac ctc atg gac atg ttc cgc 96  
 Asp Ala Thr Ser Asp Glu Val Arg Lys Asn Leu Met Asp Met Phe Arg  
 20 25 30

gac agg caa gcg ttc agc gag cac acc tgg aag atg ctg ctc tcc gtc 144  
 Asp Arg Gln Ala Phe Ser Glu His Thr Trp Lys Met Leu Leu Ser Val  
 35 40 45

30 tgc cgc tcc tgg gct gca tgg tgc aag ctg aac aac agg aag tgg ttc 192  
 Cys Arg Ser Trp Ala Ala Trp Cys Lys Leu Asn Asn Arg Lys Trp Phe  
 50 55 60

ccc gct gag ccc gag gac gtg agg gat tac ctt ctg tac ctg caa gct 240  
 Pro Ala Glu Pro Glu Asp Val Arg Asp Tyr Leu Leu Tyr Leu Gln Ala  
 65 70 75 80

40 cgc ggg ctg gca gtg aag acc atc cag caa cac ctt gga caa ctg aac 288  
 Arg Gly Leu Ala Val Lys Thr Ile Gln Gln His Leu Gly Gln Leu Asn  
 85 90 95

atg ctt cac agg cgc tcc ggc ctc ccg cgc ccc agc gac tcg aac gcc 336  
 Met Leu His Arg Arg Ser Gly Leu Pro Arg Pro Ser Asp Ser Asn Ala  
 100 105 110

gtg agc ctc gtc atg cgc cgc atc agg aag gaa aac gtc gat gcc ggc 384  
 Val Ser Leu Val Met Arg Arg Ile Arg Lys Glu Asn Val Asp Ala Gly  
 115 120 125

50

gaa	agg	gca	aag	cag	gcc	ctc	gcg	ttc	gag	agg	acc	gat	ttc	gac	cag	432	
Glu	Arg	Ala	Lys	Gln	Ala	Leu	Ala	Phe	Glu	Arg	Thr	Asp	Phe	Asp	Gln		
130					135				140								
gtc	cgc	agc	ctg	atg	gag	aac	agc	gac	agg	tgc	cag	gac	att	agg	aac	480	
Val	Arg	Ser	Leu	Met	Glu	Asn	Ser	Asp	Arg	Cys	Gln	Asp	Ile	Arg	Asn		
145				150				155			160						
10	ctg	gcg	ttc	ctc	gga	att	gca	tac	aac	acg	ctc	ctc	agg	atc	gcg	528	
Leu	Ala	Phe	Leu	Gly	Ile	Ala	Tyr	Asn	Thr	Leu	Leu	Arg	Ile	Ala	Glu		
					165			170			175						
att	gcc	cgc	att	cgc	gtg	aag	gac	att	agc	cgc	acc	gac	ggc	ggc	agg	576	
Ile	Ala	Arg	Ile	Arg	Val	Lys	Asp	Ile	Ser	Arg	Thr	Asp	Gly	Gly	Arg		
					180			185			190						
20	atg	ctt	atc	cac	att	ggc	agg	acc	aag	acg	ctc	gtt	tcc	acc	gca	ggc	624
Met	Leu	Ile	His	Ile	Gly	Arg	Thr	Lys	Thr	Leu	Val	Ser	Thr	Ala	Gly		
					195			200			205						
gtc	gaa	aag	gcc	ctc	agc	ctc	gga	gtg	acc	aag	ctc	gtc	gaa	cgc	tgg	672	
Val	Glu	Lys	Ala	Leu	Ser	Leu	Gly	Val	Thr	Lys	Leu	Val	Glu	Arg	Trp		
					210			215			220						
atc	tcc	gtg	tcc	ggc	gtc	gcg	gac	gac	cca	aac	aac	tac	ctc	ttc	tgc	720	
Ile	Ser	Val	Ser	Gly	Val	Ala	Asp	Asp	Pro	Asn	Asn	Tyr	Leu	Phe	Cys		
					225			230			235			240			
30	cgc	gtc	cgc	aag	aac	ggg	gtg	gct	gcc	cct	agc	gcc	acc	agc	caa	ctc	768
Arg	Val	Arg	Lys	Asn	Gly	Val	Ala	Ala	Pro	Ser	Ala	Thr	Ser	Gln	Leu		
					245			250			255						
agc	acg	agg	gcc	ttg	gaa	ggt	att	ttc	gag	gcc	acc	cac	cgc	ctg	atc	816	
Ser	Thr	Arg	Ala	Leu	Glu	Gly	Ile	Phe	Glu	Ala	Thr	His	Arg	Leu	Ile		
					260			265			270						
40	tac	ggc	gcf	aag	gat	gac	agc	ggt	caa	cgc	tac	ctc	gca	tgg	tcc	ggg	864
Tyr	Gly	Ala	Lys	Asp	Asp	Ser	Gly	Gln	Arg	Tyr	Leu	Ala	Trp	Ser	Gly		
					275			280			285						
cac	tcc	gcc	cgc	gtt	gga	gct	gct	agg	gac	atg	gcc	cgc	gcc	ggt	gtt	912	
His	Ser	Ala	Arg	Val	Gly	Ala	Ala	Arg	Asp	Met	Ala	Arg	Ala	Gly	Val		
					290			295			300						
tcc	atc	ccc	gaa	atc	atg	cag	gcg	ggt	gga	tgg	acg	aac	gtg	aac	att	960	
Ser	Ile	Pro	Glu	Ile	Met	Gln	Ala	Gly	Gly	Trp	Thr	Asn	Val	Asn	Ile		
					305			310			315			320			
50	gtc	atg	aac	tac	att	cgc	aac	ctt	gac	agc	gag	acg	ggc	gca	atg	gtt	1008
Val	Met	Asn	Tyr	Ile	Arg	Asn	Leu	Asp	Ser	Glu	Thr	Gly	Ala	Met	Val		
					325			330			335						
cgc	ctc	ctg	gaa	gat	ggt	gac	tga									1032	
Arg	Leu	Leu	Glu	Asp	Gly	Asp											
					340												

&lt;210&gt; 4

&lt;211&gt; 343

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;400&gt; 4

Met	Ser	Asn	Leu	Leu	Thr	Val	His	Gln	Asn	Leu	Pro	Ala	Leu	Pro	Val
1															15

Asp	Ala	Thr	Ser	Asp	Glu	Val	Arg	Lys	Asn	Leu	Met	Asp	Met	Phe	Arg
10															30

Asp	Arg	Gln	Ala	Phe	Ser	Glu	His	Thr	Trp	Lys	Met	Leu	Leu	Ser	Val
35															45

Cys	Arg	Ser	Trp	Ala	Ala	Trp	Cys	Lys	Leu	Asn	Asn	Arg	Lys	Trp	Phe
50															60

Pro	Ala	Glu	Pro	Glu	Asp	Val	Arg	Asp	Tyr	Leu	Leu	Tyr	Leu	Gln	Ala
65															80

20	Arg	Gly	Leu	Ala	Val	Lys	Thr	Ile	Gln	Gln	His	Leu	Gly	Gln	Leu	Asn
															95	

Met	Leu	His	Arg	Arg	Ser	Gly	Leu	Pro	Arg	Pro	Ser	Asp	Ser	Asn	Ala
100															110

Val	Ser	Leu	Val	Met	Arg	Arg	Ile	Arg	Lys	Glu	Asn	Val	Asp	Ala	Gly
115															125

30	Glu	Arg	Ala	Lys	Gln	Ala	Leu	Ala	Phe	Glu	Arg	Thr	Asp	Phe	Asp	Gln
															140	

Val	Arg	Ser	Leu	Met	Glu	Asn	Ser	Asp	Arg	Cys	Gln	Asp	Ile	Arg	Asn
145															160

Leu	Ala	Phe	Leu	Gly	Ile	Ala	Tyr	Asn	Thr	Leu	Leu	Arg	Ile	Ala	Glu
165															175

40	Ile	Ala	Arg	Ile	Arg	Val	Lys	Asp	Ile	Ser	Arg	Thr	Asp	Gly	Gly	Arg
															190	

Met	Leu	Ile	His	Ile	Gly	Arg	Thr	Lys	Thr	Leu	Val	Ser	Thr	Ala	Gly
195															205

Val	Glu	Lys	Ala	Leu	Ser	Leu	Gly	Val	Thr	Lys	Leu	Val	Glu	Arg	Trp
210															220

50	Ile	Ser	Val	Ser	Gly	Val	Ala	Asp	Asp	Pro	Asn	Asn	Tyr	Leu	Phe	Cys
															240	

Arg	Val	Arg	Lys	Asn	Gly	Val	Ala	Ala	Pro	Ser	Ala	Thr	Ser	Gln	Leu
245															255

Ser	Thr	Arg	Ala	Leu	Glu	Gly	Ile	Phe	Glu	Ala	Thr	His	Arg	Leu	Ile
260															270

Tyr Gly Ala Lys Asp Asp Ser Gly Gln Arg Tyr Leu Ala Trp Ser Gly  
 275 280 285

His Ser Ala Arg Val Gly Ala Ala Arg Asp Met Ala Arg Ala Gly Val  
 290 295 300

10 Ser Ile Pro Glu Ile Met Gln Ala Gly Gly Trp Thr Asn Val Asn Ile  
 305 310 315 320

Val Met Asn Tyr Ile Arg Asn Leu Asp Ser Glu Thr Gly Ala Met Val  
 325 330 335

Arg Leu Leu Glu Asp Gly Asp  
 340

<210> 5

20 <211> 69

<212> DNA

<213> Artificial Sequence

<220>

<223> FRT5

<400> 5

ccatggctag cgaagttcct attccgaagt tcctattctt caaaaggtat aggaacttca 60

gtactcgag 69

30 <210> 6

<211> 72

<212> DNA

<213> Artificial Sequence

<220>

<223> FRT6

<400> 6

ccatggctag cgaagttcct attccgaagt tcctattctt caaaaaggtat aggaacttca 60

gacgtcctcg ag 72

<210> 7

<211> 72

<212> DNA

<213> Artificial Sequence

<220>

<223> FRT7

<400> 7

ccatggctag cgaagttcct attccgaagt tccttattctt caataagttat aggaacttca 60

10 ctagttctcg ag 72

62451-850 (S)

37

CLAIMS:

1. A method for mobilizing in a plant cell a viral replicon from a T-DNA, comprising:

(a) infecting a plant cell with an *Agrobacterium* 5 carrying an *Agrobacterium* replicon having a T-DNA which contains a viral replicon flanked by directly repeated target sites for a site-specific recombinase; and

(b) providing in the cell a nucleotide sequence 10 encoding a site-specific recombinase or an active fragment or variant thereof, the nucleotide sequence being operably linked to a promoter for expressing the recombinase in a plant cell;

wherein expression of said recombinase mobilizes the viral replicon.

2. The method according to claim 1 wherein the 15 nucleotide sequence is provided in the cell, the T-DNA or the viral replicon.

3. The method according to claim 1 or 2, wherein said method provides to said plant cell a plurality of copies of a 20 DNA sequence of interest and wherein said T-DNA contains in a 5' to 3' or 3' to 5' orientation, a first target site for a site-specific recombinase, said viral replicon, said DNA sequence of interest, and a second target site for said recombinase in direct repeat with said first target site, wherein said first and second target sites are identical.

25 4. The method according to claim 1 or 2, wherein said method provides to a plant cell a plurality of copies of a DNA sequence of interest flanked by nonidentical target sites for a site-specific recombinase, and wherein said T-DNA contains in a 30 5' to 3' or 3' to 5' orientation, a first target site for said recombinase, said viral replicon, a second target site for said

62451-850 (S)

38

recombinase, said DNA sequence of interest, and a third target site for said recombinase, wherein said first and third target sites are directly repeated and identical with respect to each other, and said second target site is nonidentical to said 5 first and third target sites.

5. The method according to claim 3, wherein the genome of said plant cell comprises at least one target site for a site-specific recombinase which corresponds to the identical target sites in said T-DNA, and wherein said DNA of interest is 10 inserted into the genome of said plant cell.

6. The method according to claim 4, wherein the genome of said plant cell comprises at least two non-identical target sites for a site-specific recombinase which correspond to the non-identical second and third target sites flanking the DNA 15 sequence of interest, and wherein said DNA of interest is inserted into the genome of said plant cell.

7. The method according to any one of claims 1-6, wherein said site-specific recombinase is a member of the integrase family.

20 8. The method according to claim 7, wherein said site-specific recombinase is FLP, Cre, Int, SSVI, integrase R or an active fragment or variant thereof.

9. The method according to claim 8, wherein said target sites are selected from the group consisting of FRT and mutant 25 FRT sites.

10. The method according to claim 9, wherein said mutant FRT site is FRT5, FRT6 or FRT7.

11. The method according to claim 9, wherein said recombinase is encoded by moFLP.

62451-850 (S)

39

12. The method according to claim 8, wherein said target sites are loxP and a mutant loxP.

13. The method according to claim 12, wherein said recombinase is encoded by moCre.

5 14. The method according to any one of claims 4 to 6, wherein said *Agrobacterium* replicon is a Ti plasmid or a derivative thereof.

15. The method according to any one of claims 4 to 6, wherein said *Agrobacterium* replicon is a Ri plasmid or a derivative thereof.

16. The method according to any one of claims 4 to 6, wherein said T-DNA contains said nucleotide sequence encoding said recombinase or an active fragment or variant thereof.

17. The method according to any one of claims 4 to 6, 15 wherein said viral replicon contains said nucleotide sequence encoding said recombinase or an active fragment or variant thereof.

18. The method according to any one of claims 4 to 6, wherein said plant cell is from a monocot.

20 19. The method according to claim 18, wherein said monocot is a member of the family *Gramineae*.

20. The method according to claim 18, wherein said monocot is maize, wheat, rice, barley, sorghum or rye.

21. The method according to any one of claims 4 to 6, 25 wherein said viral replicon is a maize streak virus (MSV), wheat dwarf virus (WDV), tomato golden mosaic virus (TGMV) or tobacco mosaic virus (TMV) replicon.

62451-850 (S)

40

22. The method according to any one of claims 4 to 6, wherein said plant cell is from a dicot.

23. The method according to claim 22, wherein said dicot is soybean, canola, *Brassica*, sunflower, alfalfa or safflower.

5 24. A T-DNA of an *Agrobacterium* replicon, wherein the T-DNA contains a viral replicon flanked by directly repeated target sites for a site-specific recombinase.

10 25. The T-DNA according to claim 24, wherein said T-DNA contains in a 5' to 3' or 3' to 5' orientation, a first target site for a site-specific recombinase, said viral replicon, a DNA sequence of interest, and a second target site for said recombinase in direct repeat with said first target site, wherein said first and second target sites are identical.

15 26. The T-DNA according to claim 24, wherein said T-DNA contains in a 5' to 3' or 3' to 5' orientation, a first target site for said recombinase, said viral replicon, a second target site for said recombinase, a DNA sequence of interest, and a third target site for said recombinase, wherein said first and third target sites are directly repeated and identical with respect to each other, and said second target site is nonidentical to said first and third target sites.

27. The T-DNA according to claim 26, wherein said viral replicon contains a nucleotide sequence encoding said recombinase or an active fragment or variant thereof, and said nucleotide sequence is operably linked to a promoter that drives expression in a plant cell.

28. The T-DNA according to claim 27, wherein said recombinase is encoded by FLP or moFLP.

29. The T-DNA according to claim 27, wherein said recombinase is encoded by Cre or moCre.

62451-850 (S)

41

30. The T-DNA according to claim 26, wherein said site-specific recombinase is a member of the integrase family.

31. The T-DNA according to claim 30, wherein said site-specific recombinase is FLP, Cre, Int, SSVI, integrase R or an 5 active fragment or variant thereof.

32. The T-DNA according to claim 31, wherein said target sites are selected from the group consisting of FRT and mutant FRT sites.

33. The T-DNA according to claim 32, wherein said mutant 10 FRT site is FRT5, FRT6 or FRT7.

34. The T-DNA according to claim 31, wherein said target sites are loxP and a mutant loxP.

35. The T-DNA according to claim 26, further comprising a nucleotide sequence encoding said site-specific recombinase, 15 wherein said nucleotide sequence is operably linked to a promoter that drives expression in a plant cell.


36. An *Agrobacterium* replicon containing the T-DNA according to any one of claims 24 to 35.

37. An *Agrobacterium* containing the replicon according to 20 claim 36.

SMART & BIGGAR  
OTTAWA, CANADA


PATENT AGENTS

1/5




**FIG. 1.**

2/5




3/5

FIG. 3.

WO 99/25855

PCT/US98/24486

4/5

FIG. 4.

5/5




FIG. 5.