



US 20060024725A1

(19) **United States**

(12) **Patent Application Publication**
Hussa et al.

(10) **Pub. No.: US 2006/0024725 A1**

(43) **Pub. Date: Feb. 2, 2006**

(54) **ONCOFETAL FIBRONECTIN AS A MARKER FOR PREGNANCY-RELATED INDICATIONS**

sional application No. 60/592,824, filed on Jul. 30, 2004.

(76) Inventors: **Robert Hussa**, Sunnyvale, CA (US);
Mark Fischer-Colbrie, Cupertino, CA (US);
Jerome P. LaPointe, Oakland, CA (US);
Durlin Hickok, Seattle, WA (US)

Publication Classification

(51) **Int. Cl.**
C12Q *1/68* (2006.01)
G01N *33/53* (2006.01)
G01N *33/567* (2006.01)
(52) **U.S. Cl.** **435/6; 435/7.2**

Correspondence Address:
FISH & RICHARDSON, PC
P.O. BOX 1022
MINNEAPOLIS, MN 55440-1022 (US)

(57) **ABSTRACT**

Methods and products for the detection of oncofetal fibronectin indicating molecules in samples are provided. Methods for imaging of oncofetal fibronectin are provided. In some methods provided herein, the sample is treated with a reagent and/or contacted with a non-specific binder. Provided are methods for testing subjects to ascertain health and disease status and to assess the risk of developing a disease or condition. Methods for detecting the presence of oncofetal fibronectin indicating molecules by a variety of methods such as immunoassays and mass spectrometry also are provided. Methods and products for detection of oncofetal fibronectin for selection of concepti are provided.

(21) Appl. No.: **11/193,857**

(22) Filed: **Jul. 29, 2005**

Related U.S. Application Data

(60) Provisional application No. 60/592,823, filed on Jul. 30, 2004. Provisional application No. 60/592,803, filed on Jul. 30, 2004. Provisional application No. 60/592,825, filed on Jul. 30, 2004. Provisional application No. 60/592,804, filed on Jul. 30, 2004. Provi-





FIG. 1A



Fibronectin repeats			SH- Free sulfhydryl group
□ Type I	○ Type II	○ Type III	☆ N-linked glycosylation sites
▨ Exposed binding site	▩ Cryptic binding site	☆ O-linked glycosylation site	

FIG. 1B




FIG. 1C

ONCOFETAL FIBRONECTIN AS A MARKER FOR PREGNANCY-RELATED INDICATIONS

RELATED APPLICATIONS

[0001] Benefit of priority under 35 U.S.C. §119(e) is claimed to U.S. Provisional Application Ser. No. 60/592,823, filed Jul. 30, 2004, to Robert Husa, Mark Fischer-Colbrie, Jerome Lapointe, Simon Shorter and Andrew Senyei entitled "Methods for Detecting Oncofetal Fibronectin;" to U.S. Provisional Application Ser. No. 60/592,803, to Robert Husa, Mark Fischer-Colbrie, Jerome Lapointe, and Durlin Hickok, filed Jul. 30, 2004, entitled "Oncofetal Fibronectin as a Marker for Pregnancy-Related Indications;" to U.S. Provisional Application Ser. No. 60/592,825, filed Jul. 30, 2004, to Mark Fischer-Colbrie, Jerome Lapointe, and Durlin Hickok, entitled "Samples for Detection of Oncofetal Fibronectin and Uses Thereof;" to U.S. Provisional Application Ser. No. 60/592,804, filed Jul. 30, 2004, to Robert Husa, Mark Fischer-Colbrie, Jerome Lapointe, and Simon Shorter, entitled "Oncofetal Fibronectin as a Marker for Health and Disease;" and to U.S. Provisional Application Ser. No. 60/592,824, filed Jul. 30, 2004, to Robert Husa and Simon Shorter, entitled "Detection of Oncofetal Fibronectin for Selection of Concepti."

[0002] This application also is related to International PCT application No. _____ (attorney docket no. 17101-029WO1/830PC); and to each of U.S. application Ser. No. _____ (attorney docket nos. 17101-027001/828, 17101-028001/829, 17101-029001/830 and 17101-080001/831), each filed the same day herewith.

[0003] The subject matter of each of the above noted provisional applications, applications and international application is incorporated by reference in its entirety by reference thereto.

FIELD OF THE INVENTION

[0004] Methods and products for the detection of oncofetal fibronectin protein and nucleic acid molecule encoding oncofetal fibronectin in samples are provided. In particular, methods and products for the detection of oncofetal fibronectin proteins or nucleic acids, and/or autoantibodies therefor, in body tissue and fluid samples, such as lavage samples, cervicovaginal and urine samples are provided. The methods permit screening or indicating of risk for or for diseases including cancers, inflammatory diseases and pregnancy-related conditions.

BACKGROUND

[0005] Fibronectins constitute a family of proteins expressed from a single gene. Various isoforms of fibronectin are present in plasma and adult tissue, including connective tissue, skin, colon, liver, spleen and kidney (Matsuura and Hakomori, *Proc. Natl. Acad. Sci. USA* 82:6517-6521 (1985)). Fetal tissues and some tumor cells and other cells contain or express fibronectin isoforms collectively called "fetal" or "oncofetal" fibronectins. For example, oncofetal fibronectin (onFFN) is present in placenta, amniotic fluid, fetal tissue and cell lines from hepatomas and sarcomas (Matsuura and Hakomori, *Proc. Natl. Acad. Sci. USA*, 82:6517-6521 (1985)). Oncofetal fibronectin has been used as a marker for pre-term delivery by a pregnant woman and also as a marker for some cancers.

[0006] Early detection of cancer and other diseases and prediction of risk factors associate pregnancy-related conditions is important to developing effective treatment strategies, and oncofetal fibronectin is associated with some of these conditions, a need exists for improved tests to detect oncofetal fibronectin, for improved sampling methods and oncofetal fibronectin detection methods in order to exploit its use as a marker. Therefore, among the objects herein, it is an object herein to provide methods and products for detection of oncofetal fibronectin proteins and nucleic acids encoding the proteins, and to provide sampling methods and to provide diagnostic tests and products therefor.

SUMMARY

[0007] Provided are methods for detection of an oncofetal fibronectin indicating molecule and methods for obtaining samples for use in methods of detection. Also provided are products for use in methods of detection of oncofetal fibronectin or of molecules indicative thereof in samples.

[0008] Provided are methods that include detection of oncofetal fibronectin for any of a variety of indications and uses related to pregnancy or delivery, including, but not limited to, risk of preterm, impending and/or imminent delivery, prediction of delivery date, prediction of maintenance of pregnancy, use in methods of preventing preterm delivery and use in inducing delivery.

[0009] Among the methods provided herein, are methods for assessing whether a subject has an increased likelihood of imminent or preterm delivery, by detecting an oncofetal fibronectin indicating molecule in a sample from a pregnant subject, wherein presence of the oncofetal fibronectin indicating molecule in the sample indicates that the subject has an increased likelihood of imminent or preterm delivery. In such methods, presence of an amount of oncofetal fibronectin indicating molecule at or above a threshold level can indicate that the subject has an increased likelihood of imminent or preterm delivery. The threshold level for such methods can be 1 ng/ml or about 1 ng/ml, 2 ng/ml or about 2 ng/ml, 3 ng/ml or about 3 ng/ml, 4 ng/ml or about 4 ng/ml, 5 ng/ml or about 5 ng/ml, for a buffer-treated sample, or 1 ng/ml or about 1 ng/ml, 3 ng/ml or about 3 ng/ml, 5 ng/ml or about 5 ng/ml, 7 ng/ml or about 7 ng/ml, 10 ng/ml or about 10 ng/ml, for an untreated sample.

[0010] In other embodiments, provided herein are methods for determining whether to administer oxytocin for induction of delivery, by determining the amount of an oncofetal fibronectin indicating molecule in a sample of a pregnant subject, and if the amount of oncofetal fibronectin indicating molecule is equal to or above threshold level, identifying the subject as one for whom oxytocin induction is likely to result in vaginal delivery. Such methods can be used to identify a subject as one for whom oxytocin induction of delivery is favorable, and to administer to the subject a dose of oxytocin effective to induce delivery. According to such methods when the sample is positive for an oncofetal fibronectin indicating molecule, the subject is likely to vaginally deliver within 24 or 48 hours and/or is likely to vaginally deliver after a single induction procedure. In such methods, the induction procedure can be vaginal ripening, administration of a pre-induction agent, or administration of an induction agent. In some instances, subject can be likely to deliver after a single administration of a pre-induction

agent. Also provided are methods for evaluating the effectiveness of induction, by determining the amount of an oncofetal fibronectin indicating molecule in a sample from a pregnant subject who has undergone an induction procedure, and if the amount of oncofetal fibronectin indicating molecule is equal to or above threshold level, identifying the subject as one who is likely to vaginally deliver. In such methods, the subject can be likely to vaginally deliver within 48 hours or 24 hours, after a single induction procedure, or after a single administration of a pre-induction agent. Induction procedures can be selected from vaginal ripening, administration of a pre-induction agent and administration of an induction agent.

[0011] Also provided herein are methods for identifying a subject for induction of delivery, by determining the amount of an oncofetal fibronectin indicating molecule in a sample from a pregnant subject, determining a second indicator of induction outcome for the subject, and if the amount of oncofetal fibronectin indicating molecule is above threshold level and the second indicator indicates favorable induction outcome, identifying the subject as one for whom induction is likely to be successful. Such methods can be used to identify a subject as one for whom induction of delivery is likely to be successful, and to administer to the subject an induction procedure. Likelihood of successful induction can be indicated by any of the following: increased likelihood of vaginal delivery upon induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between initiating induction and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between administering a parturifacient and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between administering oxytocin and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, increased likelihood of delivering within 24 hours of induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, increased likelihood of delivering within 48 hours of induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome and decreased likelihood of more than one administration of pre-induction agent to the subject relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicators of induction outcome, and combinations thereof. The second indicator of induction outcome can be any of a measurement or observation of the pregnant subject, a measurement or observation of the fetus(es), and medical history of the pregnant subject. Such indicators include, but are not limited to, cervical length, Bishop score, effacement, parity, cervical dilation, gestational age, body mass index, station, consistency, transvaginal ultrasound, and digital examination, or a combination thereof.

[0012] Also provided are methods for determining the likelihood of successful induction of a subject, by determining the amount of an oncofetal fibronectin indicating molecule in a sample of a pregnant subject, determining the highest of multi-tiered threshold levels equal to or less than the amount of oncofetal fibronectin indicating molecule in

the sample, where each higher threshold indicates an increased likelihood of successful induction relative to each lower threshold, and identifying the likelihood of successful induction for the subject according to likelihood indicated by the highest threshold. Also provided are methods for identifying a subject for whom induction of delivery is unfavorable, by determining the amount of an oncofetal fibronectin indicating molecule in a sample of a pregnant subject, and if the amount of oncofetal fibronectin indicating molecule is below threshold level, identifying the subject as one for whom oxytocin induction is unlikely to result in vaginal delivery.

[0013] In the methods provided, successful induction can be indicated by each increasing amount in the multi-tiered thresholds indicates, relative to lower thresholds: increased likelihood of vaginal delivery upon induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between initiating induction and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between administering a parturifacient and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between administering oxytocin and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, increased likelihood of delivering within 24 hours of induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, increased likelihood of delivering within 48 hours of induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome and decreased likelihood of more than one administration of pre-induction agent to the subject relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicators of induction outcome, or combinations thereof.

[0014] In such methods, the sample can be any one of a swab of the point of a possible cervicovaginal lesion, urine, blood, plasma, serum, a body tissue, lavage and cervical vaginal fluid, sampled from among the cervical canal, cervical os, ectocervix, transition zone on the cervix between squamous and columnar cells, posterior fornix, a portion of the vagina below the posterior fornix, lower third of the vagina, labia, cervical interstitial fluid and combinations thereof. In a particular embodiment, the sample is any one of a swab of the cervical canal, a swab of the cervical os, a swab of the ectocervix, a swab of the transition zone on the cervix between squamous and columnar cells, a swab of the vagina, a swab of the posterior fornix, a swab of the portion of the vagina below the posterior fornix, a swab of the lower third of the vagina, a swab of the labia, and combinations thereof. A sample can be collected with a polyester swab, a cotton swab or a rayon swab. When the sample is a cotton swab, the method can be conducted on the swab. When the sample is tested by vertical flow, the sample essentially does not contain blood, or contain 5% or about 5% or less, 1% or about 1% or less, 0.5% or about 0.5% or less, 0.1% or about 0.1% or less blood.

[0015] Further in such methods testing for the presence of oncofetal fibronectin indicating molecule can include testing for the presence of an oncofetal fibronectin protein, or a fragment thereof, testing for the presence of a nucleic acid molecule encoding oncofetal fibronectin, a nucleic acid molecule complementary to a nucleic acid molecule encoding oncofetal fibronectin, or a fragment thereof, or testing for the presence of an autoantibody for oncofetal fibronectin protein or an autoantibody for a nucleic acid molecule encoding oncofetal fibronectin and fragments thereof. When the indicating molecule is a nucleic acid molecule, the methods can further include treating the sample under nucleic acid synthesis conditions such as reverse transcriptase polymerase chain reaction.

[0016] Such methods can further include contacting the sample with a fibronectin or oncofetal fibronectin binding partner, and detecting complexes of the binding partner and oncofetal fibronectin, whereby detection of a complex is indicative of the amount of oncofetal fibronectin indicating molecule in the sample. The method can further include contacting the sample with a first fibronectin or oncofetal fibronectin binding partner, contacting the sample with a second fibronectin or oncofetal fibronectin binding partner, wherein the second fibronectin or oncofetal fibronectin binding partner is conjugated to a detectable or bindable moiety, or the second fibronectin or oncofetal fibronectin binding partner is immobilized to a solid support, and detecting complexes of the first binding partner, oncofetal fibronectin indicating molecule, and the second binding partner, whereby detection of a complex is indicative of the amount of oncofetal fibronectin indicating molecule in the sample. The methods further can include contacting a sample from the subject with a non-specific binding compound, contacting the sample with a second fibronectin or oncofetal fibronectin binding partner, such as a conjugate or immobilized binding partner. In one aspect, the first binding partner is conjugated to a moiety such as, for example, colloidal metal, photodetectable latex microsphere, chromophore, fluorescent moiety, quantum dot, and detectable enzyme. The method can further include contacting the sample with a detectable compound that specifically binds the first binding partner, wherein the detectable compound is an antibody conjugate or a nucleic acid conjugate. The first binding partner or second binding partner can be an anti-fibronectin antibody or an antigen-binding fragment thereof.

[0017] The methods provided herein can be performed in any of numerous manners. For example, complexes can be detected by determining if any first binding partner is in spatial proximity to the second binding partner, whereby detection of any first and second binding partners in spatial proximity indicates presence of an oncofetal fibronectin indicating molecule in a sample, where spatial proximity can be determined by a non-radioactive energy transfer reaction, such as fluorescence energy transfer (FET), fluorescence resonance energy transfer (FRET), and homogeneous time-resolved fluorescence (HTRF). In other methods, oncofetal fibronectin indicating molecule or fragment thereof or binding partner can be detected by mass spectrometry or gel electrophoresis. In some methods, the amount of oncofetal fibronectin indicating molecule detected can be compared to one or more thresholds, wherein the sample is classified according to the highest threshold that is less than or equal to the detected amount of oncofetal fibronectin indicating molecule. In some methods, the sample is contacted with a

non-specific binding compound or with a non-specific binding surface of a solid support. In the methods described herein, the complex can be detected by measuring the oncofetal fibronectin indicating molecule that bound to the fibronectin or oncofetal fibronectin binding partner, or to a fragment of the oncofetal fibronectin indicating molecule that bound to the fibronectin or oncofetal fibronectin binding partner. In the methods described herein, the complex can be detected by detecting the weight of a compound bound to the fibronectin or oncofetal fibronectin binding partner, wherein a detected weight that corresponds to oncofetal fibronectin indicating molecule indicates the presence of oncofetal fibronectin in the sample. In the methods described herein, the complex can be measured by detecting the fibronectin or oncofetal fibronectin binding partner. The fibronectin or oncofetal fibronectin binding partner can be measured by detecting fluorescence, reflectance, absorption, bioluminescence, enzyme-linked detectable signal, or radioactive decay. In particular methods provided herein, at least one fibronectin or oncofetal fibronectin binding partner is immobilized to a test strip.

[0018] The fibronectin or oncofetal fibronectin binding partner can bind to the EDA-specific portion of an oncofetal fibronectin indicating molecule and the EDA-specific portion of an oncofetal fibronectin indicating molecule is any of an EDA portion of an oncofetal fibronectin protein, an EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, a portion of an autoantibody that binds to EDA of oncofetal fibronectin protein, and a portion of an autoantibody that binds to the EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin.

[0019] The fibronectin or oncofetal fibronectin binding partner can bind to the EDB-specific portion of an oncofetal fibronectin indicating molecule and the EDB-specific portion of an oncofetal fibronectin indicating molecule is any of an EDB portion of an oncofetal fibronectin protein, an EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, a portion of an autoantibody that binds to EDB of oncofetal fibronectin protein, and a portion of an autoantibody that binds to the EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin.

[0020] The fibronectin or oncofetal fibronectin binding partner can bind to the IIICS-specific portion of an oncofetal fibronectin indicating molecule and the IIICS-specific portion of an oncofetal fibronectin indicating molecule is any of a IIICS portion of an oncofetal fibronectin protein, a IIICS-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, a portion of an autoantibody that binds to IIICS of oncofetal fibronectin protein, and a portion of an autoantibody that binds to the IIICS-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin. The IIICS portion can be any one of V64, V89, V95 and V120. When the oncofetal fibronectin indicating molecule is a IIICS portion of oncofetal fibronectin protein, the fibronectin or oncofetal fibronectin binding partner can recognize a post-translational modification of oncofetal fibronectin protein. In one aspect, the post-translational modification can be O-glycosylation of threonine 33 of IIICS.

[0021] In one aspect the oncofetal fibronectin indicating molecule is identified as lacking EDA, EDB or IIICS. If the oncofetal fibronectin indicating molecule is identified as lacking, the portion of IIICS can be amino acids 1-25 of IIICS, amino acids 90-120 of IIICS or both.

[0022] Also provided are methods that include contacting the sample with a test strip containing a mobilizable oncofetal fibronectin binding partner conjugated to a detectable moiety and a fibronectin or oncofetal fibronectin binding partner immobilized to the test strip, and detecting complexes of the first binding partner, oncofetal fibronectin indicating molecule, and the second binding partner, whereby detection of a complex is indicative of the amount of oncofetal fibronectin indicating molecule in the sample. The oncofetal fibronectin binding partner specifically binds a oncofetal fibronectin indicating molecule in preference to a non-oncofetal fibronectin indicating molecule.

[0023] Also provided are methods that include contacting the sample with a test strip containing a mobilizable fibronectin or oncofetal fibronectin binding partner conjugated to a detectable moiety and an oncofetal fibronectin binding partner immobilized to the test strip, and detecting complexes of the first binding partner, oncofetal fibronectin indicating molecule, and the second binding partner, whereby detection of a complex is indicative of the amount of oncofetal fibronectin indicating molecule in the sample. The oncofetal fibronectin binding partner specifically binds a oncofetal fibronectin indicating molecule in preference to a non-oncofetal fibronectin indicating molecule.

[0024] Also provided herein are methods for increasing the accuracy of delivery date prediction, by measuring an oncofetal fibronectin indicating molecule in a sample from a pregnant subject, wherein an amount of oncofetal fibronectin indicating molecule in the sample at or above a threshold level indicates an increased likelihood that the subject will deliver within a particular time period, relative to a pregnant subject having a sample with an amount of oncofetal fibronectin indicating molecule amount below the threshold level. Also provided are methods for increasing the accuracy of pregnancy maintenance prediction, by measuring an oncofetal fibronectin indicating molecule in a sample from a pregnant subject, wherein an amount of oncofetal fibronectin indicating molecule in the sample below a threshold level indicates an increased likelihood that the subject will maintain her pregnancy for a particular time period, relative to a pregnant subject having a sample with an amount of oncofetal fibronectin indicating molecule amount at or above the threshold level. The particular time period can be any one of 3 weeks or less, 2 weeks or less, 10 days or less, 1 week or less, 6 days or less, 5 days or less, 4 days or less, 3 days or less, 2 days or less, and 1 day or less. The subject can be at least 50% more likely to deliver within the particular time period, relative to a pregnant subject having a sample with an amount of oncofetal fibronectin indicating molecule amount below the threshold level.

[0025] Also provided are combinations and kits, including combinations and kits for performing the methods provided herein. In one embodiment, a combination is provided, containing a fibronectin or oncofetal fibronectin binding partner, a parturifacient, and optionally instructions for use of the combination. Combinations also can include a non-specific binding compound. The kits provided herein can include the combinations provided herein and also a system for classifying the sample according to one or more threshold levels, and/or instructions for use.

[0026] Provided herein is the use of any of the products provided herein for use in the preparation of a medicament

for any of the methods provided herein, including, but not limited to, diagnosis, imaging, and/or treatment of a health disorder characterized by the presence of, or elevated levels of oncofetal fibronectin.

[0027] Provided are methods for detection of an oncofetal fibronectin indicating molecule and methods for obtaining samples for use in methods of detection. Also provided are products for use in methods of detection of oncofetal fibronectin or of molecules indicative thereof in samples.

[0028] Provided are methods for detecting the presence of an oncofetal fibronectin indicating molecule in a variety of samples. In some methods provided herein, the sample is treated with a reagent and/or contacted with a non-specific binder.

[0029] Provided herein are methods for detecting the presence of an oncofetal fibronectin indicating molecule in a sample, by contacting a sample with a substance that reduces non-specific binding of background material to a fibronectin or oncofetal fibronectin binding partner, contacting the sample with a fibronectin or oncofetal fibronectin binding partner, and detecting any complex formed between an oncofetal fibronectin indicating molecule and the fibronectin or oncofetal fibronectin binding partner, whereby detection of complex is indicative of the presence of oncofetal fibronectin indicating molecule in the sample. Also provide are methods for detecting the presence of an oncofetal fibronectin protein or fragment thereof in a sample, by contacting a sample with a substance that reduces non-specific binding of background material to a fibronectin or oncofetal fibronectin binding partner, wherein the fibronectin or oncofetal fibronectin binding partner is a fibronectin or oncofetal fibronectin binding protein, contacting the sample with a fibronectin or oncofetal fibronectin binding protein, and detecting any complex formed between an oncofetal fibronectin indicating molecule and the fibronectin or oncofetal fibronectin binding protein, wherein the oncofetal fibronectin indicating molecule is an oncofetal fibronectin protein or fragment thereof, whereby detection of complex is indicative of the presence of oncofetal fibronectin protein or fragment thereof in the sample. Also provide are methods for detecting the presence of an oncofetal fibronectin indicating molecule in a sample, by contacting a sample with solution that reduces the ionic strength of a sample, whereby specific binding of an oncofetal fibronectin indicating molecule to a fibronectin or oncofetal fibronectin binding partner is increased, contacting the solution-contacted sample with a fibronectin or oncofetal fibronectin binding partner, and detecting any complex formed between the oncofetal fibronectin indicating molecule and the fibronectin or oncofetal fibronectin binding partner, whereby detection of complex is indicative of the presence of oncofetal fibronectin indicating molecule in the sample.

[0030] In such methods, the substance can be a solid support, and can contain a non-specific binder to which oncofetal fibronectin indicating molecule in the sample does not specifically bind. In some methods, the non-specific binding compound can be a non-specific binding protein or a non-specific binding nucleic acid molecule. In some methods, the ionic strength of the sample after contacting with the substance is at least 150 μ or about 150 μ , or is less than or equal to 500 μ or about 500 μ . For example, the ionic strength can range from 50 μ to 350 μ , or from about 50 μ to about

350 μ , or from 150 μ to 250 μ , or ranging from about 150 μ to about 250 μ . In such methods, the amount of background material in the sample can be decreased relative to the amount of oncofetal fibronectin indicating molecule in the sample, or the amount of oncofetal fibronectin indicating molecule in the sample can be increased relative to the amount of background material in the sample. In some methods, the sample is a liquid sample, and the solute concentration of oncofetal fibronectin indicating molecule in the sample is unchanged. In some such methods, background material binds no more than 10% of the fibronectin or oncofetal fibronectin binding partner. These methods can further include contacting the sample with a second fibronectin or oncofetal fibronectin binding partner, wherein the second fibronectin or oncofetal fibronectin binding partner is conjugated to a detectable or bindable moiety, or the second fibronectin or oncofetal fibronectin binding partner is immobilized on a solid support; and the detecting step includes detecting complexes of the first binding partner, the second binding partner, and the oncofetal fibronectin indicating molecule.

[0031] Samples used in the methods provided herein can be any of urine, lymph, blood, plasma, serum, saliva, seminal fluid, lavages, cervical fluid, cervicovaginal fluid, vaginal fluid, breast fluid, breast milk, synovial fluid, semen, seminal fluid, stool, sputum, cerebral spinal fluid, tears, mucus, interstitial fluid, follicular fluid, amniotic fluid, aqueous humor, vitreous humor, peritoneal fluid, ascites, sweat, lymphatic fluid, lung sputum, lavage, or fractions or components thereof. The sample can be any of urine, lavage, breast milk, cervicovaginal swab, saliva, serum, plasma, blood, and interstitial fluid. In a particular embodiment, the sample is a urine sample. Exemplary urine samples are a neat (as obtained, i.e., unmodified or untreated) urine sample, and a frozen urine sample. In some methods, at least 30 minutes or about 30 minutes, or at least 12 hours or about 12 hours, prior to the contacting or detecting steps, the urine sample is collected from a subject.

[0032] In the methods provided herein, the non-specific binding compound is any of albumin, casein, fetal calf serum, gelatin, or an antibody that does not specifically bind an oncofetal fibronectin indicating molecule; and can be, for example, bovine serum albumin (BSA). In some of the methods provided herein, the second binding partner is immobilized to the solid support of a test strip. The second binding partner can be immobilized to a first region of the test strip, and a non-specific binding compound can be immobilized to a second region of the test strip, wherein the first region is downstream of the sample fluid flow pathway relative to the second region.

[0033] In some embodiments, the methods include normalizing the amount of oncofetal fibronectin indicating molecule in the sample according to the concentration of one or more normalization analytes in the sample; and exemplary normalization analyte is creatinine. The methods provided herein can further include contacting a sample with a non-specific binding compound, and separating the sample from the non-specific binding compound; and additionally further include, after separating the sample from the non-specific binding compound, contacting the sample with a solid support whereby protein and/or nucleic acid components of the sample are immobilized on the solid support, and contacting the solid support with a fibronectin or

oncofetal fibronectin binding partner. In some cases, background material is removed from the sample. The non-specific binding compound can be immobilized on a solid support.

[0034] In some methods, a sample positive for oncofetal fibronectin can identify the subject from whom the sample was collected as having cancer (e.g., malignant neoplastic or metastatic) cells; and in some instances, the cancer cells can originate from bladder, kidney, prostate, cervix or ovary. In a particular aspect, the cancerous cells originate from bladder.

[0035] In other methods, a sample positive for oncofetal fibronectin identifies the subject from whom the sample was collected as having an increased risk of imminent or preterm delivery. For example, a sample positive for oncofetal fibronectin identifies the subject from whom the sample was collected as one for whom induction is likely to be successful. In the methods provided, successful induction can be indicated by each increasing amount relative to lower thresholds: increased likelihood of vaginal delivery upon induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between initiating induction and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between administering a parturifacient and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between administering oxytocin and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, increased likelihood of delivering within 24 hours of induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, increased likelihood of delivering within 48 hours of induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome and decreased likelihood of more than one administration of pre-induction agent to the subject relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicators of induction outcome, or combinations thereof.

[0036] Also provided herein are methods for detecting the presence of an oncofetal fibronectin indicating molecule in a urine sample, by determining the amount of oncofetal fibronectin indicating molecule present in a buffer-treated urine sample, whereby 60 ng/ml or more, or about 60 ng/ml or more oncofetal fibronectin indicating molecule present in the buffer-treated sample identifies the sample as positive for oncofetal fibronectin. Also provided herein are methods for detecting the presence of an oncofetal fibronectin indicating molecule in a urine sample, by thawing a frozen urine sample, and determining the amount of oncofetal fibronectin indicating molecule present in the thawed urine sample. In some such methods, the determining step can include contacting the sample with a first fibronectin or oncofetal fibronectin binding partner, contacting the sample with a second fibronectin or oncofetal fibronectin binding partner, wherein: the second fibronectin or oncofetal fibronectin binding partner is conjugated to a detectable or bindable

moiety, or the second fibronectin or oncofetal fibronectin binding partner is immobilized to a solid support, and detecting complexes of the first binding partner, oncofetal fibronectin indicating molecule and the second binding partner.

[0037] In these embodiments, the methods also can include contacting the sample with a non-specific binding compound. Such methods can be used to identify the subject from whom the sample was collected as having cancerous (e.g., neoplastic, malignant or metastatic) cells; for example the cancerous cells can originate from bladder, kidney, prostate, cervix or ovary. In a particular aspect, the cancerous cells originate from bladder.

[0038] In another embodiment, provided herein are methods for detecting the presence of an oncofetal fibronectin indicating molecule in an interstitial fluid sample, by detecting any oncofetal fibronectin indicating molecule in an interstitial fluid sample.

[0039] In yet another embodiment, provided herein are methods for detecting the presence of an oncofetal fibronectin indicating molecule in a lavage sample, by detecting any oncofetal fibronectin indicating molecule in a lavage sample.

[0040] In such embodiments, the methods can further include contacting the sample with a fibronectin or oncofetal fibronectin binding partner, and detecting any complex formed between an oncofetal fibronectin indicating molecule and the fibronectin or oncofetal fibronectin binding partner, whereby detection of complex is indicative of the presence of oncofetal fibronectin indicating molecule in the sample. A lavage sample can be a ductal lavage sample. In such methods, a sample positive for oncofetal fibronectin can identify the subject from whom the sample was collected as having cancerous (e.g., neoplastic, malignant or metastatic) cells, and in some cases the cells can be from breast.

[0041] In the methods provided herein, complexes can be detected by determining if any first binding partner is in spatial proximity to the second binding partner, whereby detection of any first and second binding partners in spatial proximity indicates presence of an oncofetal fibronectin indicating molecule in a sample. In such methods, spatial proximity can be detected as a result of a non-radioactive energy transfer reaction, where the non-radioactive energy transfer reaction can be any one of fluorescence energy transfer (FET), fluorescence resonance energy transfer (FRET), homogenous time-resolved fluorescence (HTRF).

[0042] Also in such methods, the oncofetal fibronectin indicating molecule can be any of an oncofetal fibronectin protein, a nucleic acid molecule encoding oncofetal fibronectin, a nucleic acid molecule complementary to a nucleic acid molecule encoding oncofetal fibronectin, an autoantibody for oncofetal fibronectin protein, an autoantibody for a nucleic acid molecule encoding oncofetal fibronectin, and fragments thereof. In the methods provided herein, the binding partner can be an anti-fibronectin antibody, or a nucleic acid molecule, or a fragment thereof. In any of the methods provided herein, the oncofetal fibronectin indicating molecule can be any of an oncofetal fibronectin protein or a fragment thereof, a nucleic acid molecule encoding oncofetal fibronectin or a fragment thereof, a

nucleic acid molecule complementary to a nucleic acid molecule encoding oncofetal fibronectin or a fragment thereof, an autoantibody for oncofetal fibronectin protein or a fragment thereof, or an autoantibody for a nucleic acid molecule encoding oncofetal fibronectin or a fragment thereof.

[0043] Also in the methods provided herein, presence of an oncofetal fibronectin indicating molecule below a threshold amount can classify the sample as oncofetal fibronectin negative; and presence of an oncofetal fibronectin indicating molecule equal to or above a threshold amount can classify the sample as oncofetal fibronectin positive. A threshold amount of oncofetal fibronectin protein can be 50 ng/mL or about 50 ng/mL. In some such methods, the amount of oncofetal fibronectin detected is compared to two or more thresholds, wherein the sample can be classified according to the highest threshold that is less than or equal to the detected amount of oncofetal fibronectin indicating molecule. For example, a first threshold is 50 ng/mL and a second threshold is 150 ng/mL. In methods where the subject from whom the sample is collected is pregnant, the two or more threshold amounts can be a function of pregnancy gestational duration. In the methods provided herein, the complex can be detected by measuring the oncofetal fibronectin indicating molecule that bound to the fibronectin or oncofetal fibronectin binding partner, or a fragment of the oncofetal fibronectin indicating molecule that bound to the fibronectin or oncofetal fibronectin binding partner.

[0044] The methods provided herein can be used to detect oncofetal fibronectin by mass spectrometry or gel electrophoresis, by measuring the fibronectin or oncofetal fibronectin binding partner bound to the oncofetal fibronectin indicating molecule, and/or by detecting fluorescence, reflectance, absorption, bioluminescence, enzyme-linked detectable signal, or radioactive decay. In one aspect, at least one fibronectin or oncofetal fibronectin binding partner is immobilized to a test strip. The fibronectin or oncofetal fibronectin binding partner can bind to the EDA-specific portion of an oncofetal fibronectin indicating molecule and the EDA-specific portion of an oncofetal fibronectin indicating molecule is any of an EDA portion of an oncofetal fibronectin protein, an EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, a portion of an autoantibody that binds to EDA of oncofetal fibronectin protein, and a portion of an autoantibody that binds to the EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin.

[0045] The fibronectin or oncofetal fibronectin binding partner can bind to the EDB-specific portion of an oncofetal fibronectin indicating molecule and the EDB-specific portion of an oncofetal fibronectin indicating molecule is any of an EDB portion of an oncofetal fibronectin protein, an EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, a portion of an autoantibody that binds to EDB of oncofetal fibronectin protein, and a portion of an autoantibody that binds to the EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin.

[0046] The fibronectin or oncofetal fibronectin binding partner can bind to the IIICS-specific portion of an oncofetal fibronectin indicating molecule and the IIICS-specific portion of an oncofetal fibronectin indicating molecule is any of a IIICS portion of an oncofetal fibronectin protein, a IIICS-

encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, a portion of an autoantibody that binds to IIICS of oncofetal fibronectin protein, and a portion of an autoantibody that binds to the IIICS-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin. The IIICS portion can be any one of V64, V89, V95 and V120. When the oncofetal fibronectin indicating molecule is a IIICS portion of oncofetal fibronectin protein, the fibronectin or oncofetal fibronectin binding partner can recognize a post-translational modification of oncofetal fibronectin protein. In one aspect, the post-translational modification can be O-glycosylation of threonine 33 of IIICS.

[0047] In one aspect the oncofetal fibronectin indicating molecule is identified as lacking EDA, EDB or IIICS. If the oncofetal fibronectin indicating molecule is identified as lacking, the portion of IIICS can be amino acids 1-25 of IIICS, amino acids 90-120 of IIICS or both.

[0048] In another embodiment, provided herein are test strips, containing a non-specific binding region, and an analyte binding region containing a first fibronectin or oncofetal fibronectin binding partner immobilized thereon, wherein the analyte binding region is downstream of the sample fluid flow pathway relative to the non-specific binding region. In yet another embodiment, provided herein are test strips for detecting the presence of an oncofetal fibronectin indicating molecule in a sample, containing a non-specific binding region, and an analyte binding region containing a first fibronectin or oncofetal fibronectin binding partner immobilized thereon, wherein the analyte binding region is downstream of the sample fluid flow pathway relative to the non-specific binding region.

[0049] The test strips provided herein can also contain a conjugate pad, which serves as a sample application component, an absorbent pad, which serves to draw liquid continuously through the device, wherein the materials of the membrane system form a single fluid flow pathway, and a porous or bibulous member in fluid communication with the absorbent pad and conjugate pad, which porous or bibulous member accommodates a liquid sample, wherein the porous or bibulous member contains the analyte binding region. The test strips provided herein also can contain a mobilization region containing a second fibronectin or oncofetal fibronectin binding partner, wherein the second fibronectin or oncofetal fibronectin binding partner is mobilized upon contact with the sample, and the mobilization region is upstream of the analyte binding region. The test strips also can contain a control region containing a biomolecule that specifically binds the second fibronectin or oncofetal fibronectin binding partner, wherein the control region is downstream of the analyte binding region.

[0050] In some of the test strips provided herein the first fibronectin or oncofetal fibronectin binding partner binds oncofetal fibronectin in preference to fibronectin. In some of the test strips provided herein, the non-specific binding region contains a non-specific binding protein immobilized thereon, wherein the non-specific binding protein can be, for example, BSA, methylated BSA, W632 or mouse IgG.

[0051] Also provided herein are combinations containing a fibronectin or oncofetal fibronectin binding partner and a non-specific binding compound. The combinations provided herein can be used for performing the methods provided herein. In such combinations, the non-specific binder can be

a non-specific binding compound. Also in such combinations, the non-specific binder can be a non-specific binding surface of a solid support. The combinations can further contain a fibronectin or oncofetal fibronectin binding partner and a solid support containing a non-specific binding surface. The combinations can contain a test strip with a non-specific binding compound immobilized thereto. The combinations can contain a sample collection device.

[0052] Also provided herein are kits containing the combinations provided herein, and also containing instructions for use, and/or a system for classifying the subject with respect to multiple thresholds.

[0053] Also provided herein are methods for indicating oncofetal fibronectin in a subject, by collecting a lower vaginal sample from a subject, and testing for the presence of an oncofetal fibronectin indicating molecule in the sample, wherein the presence of oncofetal fibronectin indicating molecule in the sample indicates the presence of oncofetal fibronectin in the subject. In such methods, the lower vaginal sample can include a sample collected from any portion of the vagina below the posterior fornix. In some methods, the lower vaginal sample can be collected from the lower third of the vagina, can be collected with a swab. In some methods, the sample can include a labial sample. In some methods, the sample is collected by an individual unskilled in the medical profession, including, in some instances, the subject.

[0054] Also provided herein are methods for indicating oncofetal fibronectin in a subject, by collecting a labial sample from a subject, and testing for the presence of an oncofetal fibronectin indicating molecule in the sample, wherein the presence of oncofetal fibronectin indicating molecule in the sample indicates the presence of oncofetal fibronectin in the subject. In such methods, the sample can further include a lower vaginal sample. In some methods, the labial sample can be collected from the lower third of the vagina, can be collected with a swab. In some methods, the sample is collected by an individual unskilled in the medical profession, including, in some instances, the subject.

[0055] Also provided herein are methods for indicating oncofetal fibronectin in a subject, by passively collecting a cervicovaginal sample from a subject, and testing for the presence of an oncofetal fibronectin indicating molecule in the sample, wherein the presence of oncofetal fibronectin indicating molecule in the sample indicates the presence of oncofetal fibronectin in the subject. In such methods, the step of passively collecting a cervicovaginal sample can further comprise inserting a sample collection device into the vagina. In such methods, the sample collection device can be maintained in the vagina for at least 5 minutes or about 5 minutes, at least 15 minutes or about 15 minutes, at least 1 hour or about 1 hour, or at least 2 hours or about 2. In such methods, the sample collection device can be inserted into the lower third of the vagina. In such methods, the sample collection device can be inserted into the vagina in the same manner as a tampon is inserted. The sample collection device can be absorptive, and/or can be a tampon-like device or a sanitary napkin-like device. In such methods, the step of passively collecting a cervicovaginal sample can further include placing a sample collection device outside of and below the vagina. In some methods, the sample collection device can contact the labia. In some

methods, the sample collection device can be placed between the labia or vaginal orifice and the subject's undergarment. In some methods, the sample collection device can be maintained in place for 5 minutes or more, 10 minutes or more, 15 minutes or more, 1 hour or more, or 2 hours or more.

[0056] In such methods, the presence of oncofetal fibronectin indicating molecule above a threshold level can indicate that the sample is positive for oncofetal fibronectin. A threshold level can be 1 ng/ml, 2 ng/ml, 3 ng/ml, 5 ng/ml, 7 ng/ml, 10 ng/ml, 15 ng/ml, or 20 ng/ml for a buffer-treated sample. A threshold level can be 1 ng/ml, 3 ng/ml, 5 ng/ml, 10 ng/ml, 25 ng/ml, 35 ng/ml, or 50 ng/ml for an untreated sample.

[0057] In the methods provided herein, presence of oncofetal fibronectin in the sample can indicate a risk of imminent or preterm delivery, the likelihood of successful induction, the presence of cancerous cells in a subject, the risk of a subject developing cancerous cells, the aggressiveness of cancerous cells in a subject, or the effectiveness of treating cancerous cells in a subject.

[0058] Also provided herein are combinations containing a sample collection device and a fibronectin or oncofetal fibronectin binding partner. In such combinations, the sample collection device includes, but is not limited to, a urine collection device, a dipstick, a swab and a passive cervicovaginal fluid collection device. In one combination, the passive cervicovaginal sample collection device is insertable into the vagina. In another combination the passive cervicovaginal sample collection device is placed between the labia or vaginal orifice and the undergarment of a subject. The combination also can include a swab long enough to insert into the vagina, but not long enough to contact the cervix, which can be a swab 10 cm or shorter. In another combination, the fibronectin or oncofetal fibronectin binding partner can be immobilized onto the sample collection device.

[0059] Combinations provided herein also can include a second fibronectin or oncofetal fibronectin binding partner. In such combinations, the second fibronectin or oncofetal fibronectin binding partner can be conjugated to a detectable label.

[0060] The combinations provided herein can be configured to indicate a positive result when the amount of oncofetal fibronectin indicating molecule in the sample is above a threshold level. The combinations provided herein also can contain a test strip reader configured to indicate a positive result when the amount of oncofetal fibronectin indicating molecule in the sample is above a threshold level. In some such combinations, the threshold level is about or is 1 ng/ml, is about or is 2 ng/ml, is about or is 3 ng/ml, is about or is 5 ng/ml, is about or is 7 ng/ml, is about or is 10 ng/ml, is about or is 15 ng/ml, or is about or is 20 ng/ml for a buffer-treated sample. In some such combinations, the threshold level is about or is 1 ng/ml, is about or is 3 ng/ml, is about or is 5 ng/ml, is about or is 10 ng/ml, is about or is 15 ng/ml, is about or is 25 ng/ml, is about or is 35 ng/ml, or is about or is 50 ng/ml for an untreated sample. Also provided herein are kits containing the combinations provided herein, and optionally one or more of instructions for collecting and/or measuring the oncofetal fibronectin indicating molecule, and reagents therefor.

[0061] Also provided herein are methods for indicating oncofetal fibronectin in a subject, comprising collecting the sample and testing the sample for the presence of an oncofetal fibronectin indicating molecule with the combinations provided herein. In some such methods, the sample is collected from the portion of the vagina below the posterior fornix. In some such methods, the sample is collected from the lower third of the vagina. In some such methods, the sample is collected from the labia. In the methods provided herein, the sample can be collected with a passive sample collection device. In some methods provided herein, the sample is urine. In such methods, the sample can be collected by an individual not skilled in medical practice. In such methods, the sample can be collected by the subject. Provided herein are any of the methods where the step of testing the sample for the presence of oncofetal fibronectin indicating molecule is performed by an individual not skilled in medical practice or by the subject.

[0062] Also provided are methods for determining the presence and/or amount of an oncofetal fibronectin indicating molecule in a sample, by treating a urine sample under conditions for fragmentation of an oncofetal fibronectin indicating molecule, and detecting any fragments of oncofetal fibronectin in the sample, whereby detected oncofetal fibronectin fragments indicate the presence and/or amount of oncofetal fibronectin indicating molecule in a sample.

[0063] Also provided are methods for determining the presence and/or amount of an oncofetal fibronectin indicating molecule in a sample, by contacting a body surface or cavity with a lavage fluid, and detecting any oncofetal fibronectin indicating molecule in the lavage fluid.

[0064] Provided herein is the use of any of the products provided herein for use in the preparation of a medicament for any of the methods provided herein, including, but not limited to, diagnosis, imaging, and/or treatment of a health disorder characterized by the presence of, or elevated levels of oncofetal fibronectin.

[0065] Provided are methods for detection of an oncofetal fibronectin indicating molecule and methods for obtaining samples for use in methods of detection. Also provided are products for use in methods of detection of oncofetal fibronectin or of molecules indicative thereof in samples. Provided are methods for testing subjects to ascertain health and disease status and to assess the risk of developing a disease or condition.

[0066] Provided herein are methods for identifying the presence of cervical cancer in a subject by testing for an oncofetal fibronectin indicating molecule in a sample from a subject, wherein an oncofetal fibronectin positive sample identifies the presence of cervical cancer in the subject. Also provided herein are methods of detecting the presence of cancerous (e.g., neoplastic, malignant or metastatic) cervical cells in a subject, by testing for an oncofetal fibronectin indicating molecule in a sample from a subject, wherein an oncofetal fibronectin positive sample indicates the presence of cancerous (e.g., malignant neoplastic or metastatic) cervical cells in the subject.

[0067] In the methods provided herein, the sample can be, for example, a swab of the point of a possible cervicovaginal lesion, a swab of the cervical canal, a swab of the cervical

os, a swab of the ectocervix, a swab of the transition zone on the cervix between squamous and columnar cells, a swab of the vagina, a swab of the posterior fornix, a swab of the portion of the vagina below the posterior fornix, a swab of the lower third of the vagina, a swab of the labia, cervical interstitial fluid, urine, blood, plasma, serum and combinations thereof. In one aspect, the sample is a cervicovaginal sample, and the sample is one or more of the following: a swab of cervical os, a swab of cervical lesion, a swab of ectocervix, a swab of transition zone between squamous and columnar cells of cervix, or a combination thereof. In another aspect, the sample is a swab of the portion of the vagina below the posterior fornix. In some methods, the sample is collected with a polyester swab, cotton swab or rayon swab. When the sample is a cotton swab, the method can be conducted on the swab.

[0068] In some methods provided herein, presence of oncofetal fibronectin indicating molecule in the sample identifies the sample as oncofetal fibronectin positive. In some methods, absence of oncofetal fibronectin indicating molecule in the sample identifies the sample as oncofetal fibronectin negative. In other of the methods provided herein, an amount of oncofetal fibronectin indicating molecule in the sample at or above a threshold identifies the sample as oncofetal fibronectin positive. In some methods, an amount of oncofetal fibronectin indicating molecule in the sample below a threshold identifies the sample as oncofetal fibronectin negative. In some methods, the threshold amount is 40 ng/ml or about 40 ng/ml, 10 ng/ml or about 10 ng/ml, or 5 ng/ml or about 5 ng/ml.

[0069] When the sample is assayed using vertical flow, the sample essentially does not contain blood. In such methods, the sample can contain 1% or about 1% or less blood, 0.5% or about 0.5% or less blood, or 0.1% or about 0.1% or less blood.

[0070] The methods provided herein also include methods in which the step of testing for the oncofetal fibronectin indicating molecule further includes testing for an oncofetal fibronectin protein, or a fragment thereof, testing for a nucleic acid molecule encoding oncofetal fibronectin, a nucleic acid molecule complementary to a nucleic acid molecule encoding oncofetal fibronectin, or a fragment thereof, or testing for an autoantibody for oncofetal fibronectin protein, an autoantibody for a nucleic acid molecule encoding oncofetal fibronectin, and fragments thereof. In some methods, the step of testing further includes contacting the sample with a fibronectin or oncofetal fibronectin binding partner, and detecting complexes of the binding partner and oncofetal fibronectin indicating molecule. In some methods, the step of testing further includes contacting the sample with a second fibronectin or oncofetal fibronectin binding partner, wherein the second fibronectin or oncofetal fibronectin binding partner is conjugated to a detectable or bindable moiety, or the second fibronectin or oncofetal fibronectin binding partner is immobilized to a solid support. The methods provided herein also can include, prior to detecting complexes, contacting the sample with a non-specific binding compound.

[0071] In the methods provided herein, a binding partner can be conjugated to a moiety such as, for example, colloidal metal, photodetectable latex bead, chromophore, fluorescent moiety, quantum dot and detectable enzyme. The methods

provided herein can further include, after contacting the first binding partner and sample, contacting the sample with a detectable compound that specifically binds the first binder. In such methods, the detectable compound is an antibody conjugate or a nucleic acid conjugate. In some methods provided herein, a binding partner can be an anti-fibronectin antibody, or a fragment thereof.

[0072] In the methods provided herein, complexes can be detected by determining if any first binding partner is in spatial proximity to the second binding partner, whereby detection of any first and second binding partners in spatial proximity indicates presence of an oncofetal fibronectin indicating molecule in a sample. In some such methods, spatial proximity is detected as a result of a non-radioactive energy transfer reaction. In such methods, the non-radioactive energy transfer reaction can be, for example, fluorescence energy transfer (FET), fluorescence resonance energy transfer (FRET), or homogenous time-resolved fluorescence (HTRF).

[0073] In some methods provided herein, the sample is contacted with a non-specific binding compound. In some methods, the sample is contacted with a non-specific binding surface of a solid support.

[0074] The methods provided herein include methods in which the amount of oncofetal fibronectin indicating molecule detected is compared to two or more thresholds, wherein the sample is classified according to the highest threshold that is less than or equal to the detected amount of oncofetal fibronectin indicating molecule.

[0075] In some methods provided herein, the complex is detected by measuring the oncofetal fibronectin indicating molecule that bound to the fibronectin or oncofetal fibronectin binding partner, or a fragment of the oncofetal fibronectin indicating molecule that bound to the fibronectin or oncofetal fibronectin binding partner. In some methods, complex is detected by detecting the molecular weight of compounds bound to the fibronectin or oncofetal fibronectin binding partner; wherein a molecular weight that corresponds to an oncofetal fibronectin indicating molecule indicates the presence of the oncofetal fibronectin indicating molecule in the sample. In some methods, the oncofetal fibronectin indicating molecule is detected by mass spectrometry or gel electrophoresis. In some methods, the complex is detected by detecting the fibronectin or oncofetal fibronectin binding partner bound to the oncofetal fibronectin indicating molecule. In some such methods, the fibronectin or oncofetal fibronectin binding partner is detected by detecting fluorescence, reflectance, absorption, bioluminescence, enzyme-linked detectable signal, or radioactive decay. In some methods, at least one fibronectin or oncofetal fibronectin binding partner is immobilized to a test strip.

[0076] The fibronectin or oncofetal fibronectin binding partner can bind to the EDA-specific portion of an oncofetal fibronectin indicating molecule and the EDA-specific portion of an oncofetal fibronectin indicating molecule is any of an EDA portion of an oncofetal fibronectin protein, an EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, a portion of an autoantibody that binds to EDA of oncofetal fibronectin protein, and a portion of an autoantibody that binds to the EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin.

[0077] The fibronectin or oncofetal fibronectin binding partner can bind to the EDB-specific portion of an oncofetal

fibronectin indicating molecule and the EDB-specific portion of an oncofetal fibronectin indicating molecule is any of an EDB portion of an oncofetal fibronectin protein, an EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, a portion of an autoantibody that binds to EDB of oncofetal fibronectin protein, and a portion of an autoantibody that binds to the EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin.

[0078] The fibronectin or oncofetal fibronectin binding partner can bind to the IIICS-specific portion of an oncofetal fibronectin indicating molecule and the IIICS-specific portion of an oncofetal fibronectin indicating molecule is any of a IIICS portion of an oncofetal fibronectin protein, a IIICS-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, a portion of an autoantibody that binds to IIICS of oncofetal fibronectin protein, and a portion of an autoantibody that binds to the IIICS-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin. The IIICS portion can be any one of V64, V89, V95 and V120. When the oncofetal fibronectin indicating molecule is a IIICS portion of oncofetal fibronectin protein, the fibronectin or oncofetal fibronectin binding partner can recognize a post-translational modification of oncofetal fibronectin protein. In one aspect, the post-translational modification can be O-glycosylation of threonine 33 of IIICS.

[0079] In one aspect the oncofetal fibronectin indicating molecule is identified as lacking EDA, EDB or IIICS. If the oncofetal fibronectin indicating molecule is identified as lacking, the portion of IIICS can be amino acids 1-25 of IIICS, amino acids 90-120 of IIICS or both.

[0080] Also provided herein are methods of detecting the location in a subject of cervical cancer or cancerous cervical cells, by identifying the presence of cancer in a subject, or the presence of cancerous cells in a subject, according to the methods provided herein, administering to a subject a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety, whereby the conjugate localizes to tissue or cells in the subject containing an oncofetal fibronectin indicating molecule, and detecting the localization of the conjugate within the subject, thereby detecting the oncofetal fibronectin indicating molecule in tissue or cells of the subject, wherein detection is indicative of cancer or a disease state characterized by the presence of oncofetal fibronectin. In one embodiment, the tissues or cells are cervical tissues or cells. Such methods can further include detecting the location in a subject of cervical cancer or cancerous cervical cells, and administering to a subject a treatment fibronectin or oncofetal fibronectin binding partner, whereby the treatment binding partner localizes to regions in the subject containing an oncofetal fibronectin indicating molecule, whereby the localized treatment binding partner causes cell death or inhibits cell growth of cervical cancer or cancerous cervical cells.

[0081] Also provided herein are methods of treating a subject having cervical cancer or cancerous cervical cells, by identifying the presence of cervical cancer in a subject, or the presence of cancerous cervical cells in a subject, according to the methods provided herein, and administering to a subject a treatment fibronectin or oncofetal fibronectin binding partner, whereby the treatment binding partner localizes to regions in the subject containing an oncofetal fibronectin indicating molecule, whereby the localized binding partner

causes cell death or inhibits cell growth of cervical cancer or cancerous cervical cells. In such methods, the treatment binding partner can be conjugated to a therapeutic moiety.

[0082] In some methods, the imaging binding partner conjugate or treatment binding partner can be, for example, a nucleic acid molecule, a binding partner that binds the IIICS region of an oncofetal fibronectin protein, and a binding partner that binds the EDA region of an oncofetal fibronectin protein. In some methods, the imaging conjugate or treatment binding partner is administered topically.

[0083] Also provided herein are test strips for performing the methods provided herein. Also provided herein are test strips containing an analyte binding region containing a first fibronectin or oncofetal fibronectin binding partner immobilized to a solid support. The test strips provided herein can further contain a non-specific binding region, wherein the analyte binding region is downstream of the sample fluid flow pathway relative to the non-specific binding region. The test strips provided herein can further contain a conjugate pad, which serves as a sample application component, an absorbent pad, which serves to draw liquid continuously through the device, wherein the materials of the membrane system form a single fluid flow pathway, and a porous or bibulous member in fluid communication with the absorbent pad and conjugate pad, which porous or bibulous member accommodates a liquid sample and serves as the solid support upon which biomolecule interactions occur, wherein the porous or bibulous member contains the analyte binding region.

[0084] Some test strips provided herein can further contain an immobilization region containing a second fibronectin or oncofetal fibronectin binding partner, wherein the second fibronectin or oncofetal fibronectin binding partner is mobilized upon contact with the sample, and the immobilization region is upstream of the analyte binding region. Some test strips can further contain a control region containing a biomolecule that specifically binds the second fibronectin or oncofetal fibronectin binding partner, wherein the control region is downstream of the analyte binding region. In some test strips, the first fibronectin or oncofetal fibronectin binding partner binds oncofetal fibronectin in preference to fibronectin. In some test strips, the non-specific binding region contains a non-specific binding protein immobilized to the solid support, wherein the non-specific binding protein can be, for example, BSA, methylated BSA, W632 or mouse IgG.

[0085] Also provided herein are combinations for performing the methods provided herein. Also provided herein are combinations containing a fibronectin or oncofetal fibronectin binding partner and a non-specific binding compound. Some combinations further contain a test strip containing the non-specific binding partner. Also provided herein are combinations containing an fibronectin or oncofetal fibronectin binding partner and a solid support containing a non-specific binding surface. Such combinations can also contain a test strip containing the solid support. Also provided herein are combinations containing a fibronectin or oncofetal fibronectin binding partner and a sample collection device. The combinations provided herein also can contain a sample collection device.

[0086] Also provided herein are kits that contain the combinations provided herein, and optionally further com-

prising instructions for use. Also provided herein are kits that contain the combinations provided herein, and also a system for classifying the subject with respect to multiple thresholds.

[0087] Also provided herein are methods of detecting an oncofetal fibronectin indicating molecule in a subject, by administering to a subject a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety, whereby the conjugate localizes to regions in the subject containing an oncofetal fibronectin indicating molecule, and detecting the localization of the conjugate within the subject, thereby detecting the oncofetal fibronectin indicating molecule in the subject, wherein the oncofetal fibronectin binding partner is, for example, a nucleic acid molecule, a binding partner that binds the IIIICS region of an oncofetal fibronectin protein and a binding partner that binds the EDA region of an oncofetal fibronectin protein. Also provided are methods of imaging tumorous tissue in a subject, by detecting an oncofetal fibronectin indicating molecule in a subject, thereby imaging tumorous tissue in the subject. Also provided herein are methods of imaging cancerous cells in a subject, by detecting an oncofetal fibronectin indicating molecule in a subject, thereby imaging tumorous or cancerous cells in the subject.

[0088] Also provided herein are methods of detecting an oncofetal fibronectin indicating molecule in cervical tissue or cells of a subject, by administering to a subject a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety, whereby the conjugate localizes to cervical tissue or cells in the subject containing an oncofetal fibronectin indicating molecule, and detecting the localization of the conjugate within the subject, thereby detecting the oncofetal fibronectin indicating molecule in cervical tissue or cells of the subject. Also provided herein are methods of indicating the presence of cancerous (e.g., hyperplastic or malignant neoplastic) cervical cells in a subject, by detecting an oncofetal fibronectin indicating molecule in cervical tissue or cells of a subject, thereby indicating the presence of cancerous cervical cells in the subject. Also provided are methods of imaging tumorous cervical tissue or cancerous cervical cells in a subject comprising detecting an oncofetal fibronectin indicating molecule in cervical tissue or cells of a subject, thereby imaging tumorous cervical tissue or cancerous cervical cells in the subject.

[0089] In the methods provided herein, the fibronectin or oncofetal fibronectin binding partner can specifically bind the EDA, EDB or IIIICS region of an oncofetal fibronectin protein or oncofetal fibronectin-encoding nucleic acid molecule as described above. In the methods provided herein, the fibronectin or oncofetal fibronectin binding partner can be an antibody, or fragment thereof, or a nucleic acid molecule.

[0090] In the methods provided herein the tumorous tissue or cells can include, for example, lung, breast, ovary, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, prostate, thyroid, pituitary, eye, brain, oral, skin, head and neck cancer, lymphoma, leukemia, squamous cell carcinoma, adenocarcinoma, small cell carcinoma, melanoma, glioma, sarcoma and neuroblastoma.

[0091] In the methods provided herein, the conjugate can be administered intravenously, topically, or orally. In some

methods, the imaging method can be, for example, magnetic resonance imaging, ultrasonic imaging, fluorescence imaging, scintigraphy, computed tomography, computerized axial tomography, positron emission tomography, single photon emission computed tomography, ultrasound tomography and x-ray tomography. In some methods, the imaging moiety can be, for example, fluorescent moieties, radionuclides, magnetically detectable isotopes or compounds, sonographic imaging agents, chromophores, latex microspheres, or quantum dots.

[0092] Also provided herein are methods of detecting an oncofetal fibronectin indicating molecule in a subject, by topically administering to a subject a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety, whereby the conjugate localizes to surfaces on the subject containing an oncofetal fibronectin indicating molecule, and detecting the localization of the conjugate on the subject, thereby detecting the oncofetal fibronectin indicating molecule on the subject. Also provided herein are methods of indicating the presence of cancerous cells in a subject, by detecting an oncofetal fibronectin indicating molecule in a subject, wherein the detected oncofetal fibronectin indicating molecule indicates the presence of cancerous cells on the subject. In some methods, cancerous cells on the surface of the cervix are indicated.

[0093] The methods of treating a subject provided herein can also include indicating the location in a subject of cancer cells, and administering to a subject a treatment fibronectin or oncofetal fibronectin binding partner, whereby the treatment binding partner localizes to regions in the subject containing oncofetal fibronectin, whereby the localized treatment binding partner causes cell death or inhibits cell growth of cancer cells. In some such methods, the cancer is cervical cancer and the cancerous cells are cervical malignant, neoplastic or hyperplastic cells.

[0094] Also provided herein are methods of treating a health problem associated with oncofetal fibronectin in a subject, by administering to a subject a fibronectin or oncofetal fibronectin binding partner, whereby the binding partner localizes to regions in the subject containing an oncofetal fibronectin indicating molecule, whereby the localized binding partner causes cell death or inhibits cell growth, whereby the cell death or cell growth inhibition caused by the binding partner treats the health problem associated with oncofetal fibronectin, wherein the fibronectin or oncofetal fibronectin binding partner is not conjugated to a therapeutic moiety. Also provided herein are methods of treating a health problem associated with oncofetal fibronectin in a subject, by administering to a subject a fibronectin or oncofetal fibronectin binding partner, whereby the binding partner localizes to regions in the subject containing an oncofetal fibronectin indicating molecule, whereby the localized binding partner causes cell death or inhibits cell growth, whereby the cell death or cell growth inhibition caused by the binding partner treats the health problem associated with oncofetal fibronectin, wherein the fibronectin or oncofetal fibronectin binding partner is, for example, a fibronectin or oncofetal fibronectin binding partner nucleic acid molecule, a binding partner that binds the IIIICS region of a fibronectin or oncofetal fibronectin protein and a binding partner that binds the EDA region of a fibronectin or

oncofetal fibronectin protein. In some methods, tumorous tissue or malignant, hyperplastic or neoplastic cells of a subject no longer proliferate.

[0095] Also provided herein are methods of treating a cervical cancer subject, by administering to a subject a fibronectin or oncofetal fibronectin binding partner, whereby the binding partner localizes to regions in the subject containing an oncofetal fibronectin indicating molecule, whereby the localized binding partner causes cervical cell death or inhibits cell growth, whereby the cell death or cell growth inhibition caused by the binding partner stops the proliferation of cancerous cervical cells in the subject. Also provided herein are methods of treating tumorous cervical tissue in a subject, by stopping the proliferation of cancerous cervical cells in the subject, thereby treating tumorous cervical tissue in the subject.

[0096] In some methods, the fibronectin or oncofetal fibronectin binding partner specifically binds the EDA region, the EDB region, or the IIICS region of an oncofetal fibronectin protein or oncofetal fibronectin-encoding nucleic acid molecule as described above. In some methods, the fibronectin or oncofetal fibronectin binding partner is an antibody, or fragment thereof, or a nucleic acid molecule.

[0097] In some methods provided herein, the tumorous tissue or cells are from, for example, lung, breast, ovary, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, prostate, thyroid, pituitary, eye, brain, oral, skin, head and neck cancer, lymphoma, leukemia, squamous cell carcinoma, adenocarcinoma, small cell carcinoma, melanoma, glioma, sarcoma and neuroblastoma.

[0098] In some methods, the binding partner is administered intravenously, topically or orally. In some methods, the binding partner is conjugated to a therapeutic moiety. In some such methods, the therapeutic moiety is, for example, a biological toxin, a cytokine, a photosensitizing agent, a toxin, an anticancer antibiotic, a chemotherapeutic compound, a radionuclide, a binding partner and a bioluminescent compound.

[0099] Also provided herein are methods of indicating the presence of cancerous cells in a subject, by topically administering to a subject a fibronectin or oncofetal fibronectin binding partner, whereby the binding partner localizes to surfaces on the subject containing an oncofetal fibronectin indicating molecule, and detecting the localization of the conjugate on the subject, thereby indicating presence of cancerous cells in the subject. Also provided herein are methods of treating tumorous tissue in a subject, by topically administering to a subject a fibronectin or oncofetal fibronectin binding partner, whereby the binding partner localizes to surfaces on the subject containing an oncofetal fibronectin indicating molecule, whereby the localized binding partner causes cell death or inhibits cell growth, whereby the cell death or cell growth inhibition caused by the binding partner inhibits tumor proliferation in the subject. In some such methods, cancerous cells on the surface of the cervix are treated.

[0100] Also provided herein are methods of testing for an oncofetal fibronectin indicating molecule in a sample, thereby detecting oncofetal fibronectin indicating molecule, if present, in the sample. As provided herein, such methods

can be used for a variety of applications, examples of which are included in the following paragraph.

[0101] Some oncofetal fibronectin indicating molecule testing methods can be used in methods of indicating cancerous cells in a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample identifies the presence of cancerous cells in the subject. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of determining the risk of a subject developing cancerous, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample identifies a risk of the subject developing neoplastic, malignant or metastatic cells. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of predicting the development of cancerous cells in a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample identifies the likelihood of a subject developing cancerous cells. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of assessing the aggressiveness of cancerous cells in a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample identifies the cancerous cells as aggressive. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of predicting the outcome of a treatment of a cancerous disease in a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample indicates that a treatment of a cancerous disease is predicted to be successful. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of predicting the outcome of a treatment of a cancerous disease in a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample indicates that a treatment of a cancerous disease is predicted to be unsuccessful. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of monitoring a treatment of a cancerous disease in a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample indicates that a treatment of a cancerous disease is effective. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of monitoring a treatment of a cancerous disease in a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample indicates that a treatment of a cancerous disease is ineffective.

[0102] Some oncofetal fibronectin indicating molecule testing methods can be used in methods for determining the risk of cells becoming cancerous, by detecting the presence of the oncofetal fibronectin indicating molecule in a sample from cells, wherein the presence of oncofetal fibronectin indicating molecule in the sample identifies a risk of the cells

becoming cancerous. In some methods, the cells have an abnormal morphology. In some methods, the cells are dysplastic cells.

[0103] Also provided herein are methods for inhibiting the development of cancer in a subject, by treating a subject for cancer, and administering to the treated subject a fibronectin or oncofetal fibronectin binding partner, whereby development of cancer is inhibited. Also provided herein are methods for inhibiting the recurrence of cancer in a subject, by treating a subject for cancer, and administering to the treated subject a fibronectin or oncofetal fibronectin binding partner, whereby recurrence of cancer is inhibited.

[0104] In some methods, the cancer cells are, for example, lung, breast, ovary, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, prostate, thyroid, pituitary, eye, brain, oral, skin, head and neck cancer, lymphoma, leukemia, squamous cell carcinoma, adenocarcinoma, small cell carcinoma, melanoma, glioma, sarcoma and neuroblastoma cells.

[0105] Also provided herein are methods of testing for an oncofetal fibronectin indicating molecule in a sample, thereby detecting oncofetal fibronectin indicating molecule, if present, in the sample. As provided herein, such methods can be used for a variety of applications, examples of which are included in the following paragraph.

[0106] Some oncofetal fibronectin indicating molecule testing methods can be used in methods of determining the overall health state of a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample indicates that the subject is not free of disease. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of determining that a subject is disease free, by testing a sample for the presence or absence of an oncofetal fibronectin indicating molecule, wherein absence of oncofetal fibronectin indicating molecule indicates that the subject is free of disease. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of screening a subject for a health problem associated with oncofetal fibronectin, comprising testing a sample from a subject for the presence or absence of an oncofetal fibronectin indicating molecule, wherein presence of oncofetal fibronectin indicating molecule indicates that the subject has a health problem associated with oncofetal fibronectin. In some methods, the disease is, for example, cancer, pregnancy-related disorder, arthritis, diabetic retinopathy and Dupuytren's contracture. In some methods, prior to testing for the presence or absence of oncofetal fibronectin indicating molecule, the subject has not been diagnosed with a disease. Some methods further include performing one or more additional tests to identify the disease. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of indicating arthritis in a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample identifies the presence of arthritis in the subject. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of indicating diabetic retinopathy in a subject, by testing for the presence of an oncofetal fibronectin indicating

molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample identifies the presence of diabetic retinopathy in the subject. Some oncofetal fibronectin indicating molecule testing methods can be used in methods of indicating Dupuytren's contracture in a subject, by testing for the presence of an oncofetal fibronectin indicating molecule in a sample from a subject, whereby presence of oncofetal fibronectin indicating molecule in the sample identifies the presence of Dupuytren's contracture in the subject.

[0107] In some methods the sample is, for example, urine, lymph, lymphatic fluid, blood, plasma, serum, saliva, cervical fluid, cervicovaginal fluid, vaginal fluid, breast fluid, synovial fluid, semen, seminal fluid, stool, sputum, cerebral spinal fluid, tears, mucus, interstitial fluid, follicular fluid, amniotic fluid, aqueous humor, vitreous humor, lavage, tissue, peritoneal fluid, ascites and sweat. In some methods the oncofetal fibronectin indicating molecule is detected by a method such as, for example, mass spectrometry, sandwich assay, Western blot, dot blot, FRET, fluorescence polarization, fluorimetry, flow cytometry, RT-PCR, Southern blot, Northern blot, fluorescence in situ and in vivo imaging. In some methods, the oncofetal fibronectin indicating molecule is, for example, an oncofetal fibronectin protein, an oncofetal fibronectin-encoding nucleic acid and an autoantibody that preferentially binds oncofetal fibronectin.

[0108] Provided herein is the use of any of the products provided herein for use in the preparation of a medicament for any of the methods provided herein, including, but not limited to, diagnosis, imaging, and/or treatment of a health disorder characterized by the presence of, or elevated levels of oncofetal fibronectin.

[0109] Provided are methods for detection of an oncofetal fibronectin indicating molecule and methods for obtaining samples for use in methods of detection. Also provided are products for use in methods of detection of oncofetal fibronectin or of molecules indicative thereof in samples.

[0110] Provided herein are methods for detecting an oncofetal fibronectin indicating molecule in a sample, by treating a sample under conditions that separate one or more first sample components from one or more second sample components, wherein an oncofetal fibronectin indicating molecule, if present, is among the one or more first sample components, and detecting the oncofetal fibronectin indicating molecule or fragment thereof by its molecular weight.

[0111] In such methods, the treating step can further comprise contacting the sample with a fibronectin or oncofetal fibronectin binding partner immobilized on a solid support to form a complex thereof and then treating the solid support to release oncofetal fibronectin indicating molecule or a fragment thereof from the complex, and the detecting step can further comprise detecting the released oncofetal fibronectin indicating molecule or fragment thereof. In some methods, the detecting step can further comprise calculating the molecular weight of the detected oncofetal fibronectin indicating molecule or fragment thereof. In the methods provided herein, the detecting step can further comprise comparing detected first sample components to one or more references, wherein a reference that matches a detected first sample component corresponds to a fibronectin indicating molecule or fragment thereof. The methods provided herein also can include, prior to treating the solid support to release

oncofetal fibronectin indicating molecule from the binding partner, treating the solid support under conditions that separate the solid support from sample components not specifically bound to the binding partner. In some of the methods provided herein, prior to detecting and subsequent to contacting the sample with a fibronectin or oncofetal fibronectin binding partner, the method further comprises a step of contacting the sample with a fragmentation reagent. In such methods, the fragmentation reagent can be a protease or a nuclease. In some methods, the fragmentation reagent is immobilized onto a second solid support and the step of contacting the sample with a fragmentation reagent further comprises contacting the sample with the second solid support. In some methods, prior to detecting and subsequent to contacting the sample with a fragmentation reagent, the method further comprises a step of treating the sample under conditions whereby the oncofetal fibronectin indicating molecule or fragment thereof is released from the second solid support. In some methods, release of the oncofetal fibronectin indicating molecule or fragment thereof from the second solid support is accomplished by matrix-assisted laser desorption or electrospray desorption.

[0112] Also provided herein are methods where the mass corresponding to the mass of a fragment of an oncofetal fibronectin indicating molecule can be, for example, 14 kDa, 35 kDa, 55 kDa, 65 kDa, 85 kDa, 110 kDa, 120 kDa, 160 kDa, 200 kDa or 235 kDa. In some methods, the oncofetal fibronectin binding partner binds IIICS, the sample is contacted with trypsin, and the mass corresponding to the mass of a fragment of an oncofetal fibronectin indicating molecule can be, for example, 55 kDa, 65 kDa, 120 kDa, 160 kDa, 200 kDa or 235 kDa. In some other methods, the oncofetal fibronectin binding partner binds IIICS, the sample is contacted with cathepsin D, and the mass corresponding to the mass of a fragment of an oncofetal fibronectin indicating molecule can be, for example, 85 kDa or 110 kDa. In some other methods, the oncofetal fibronectin binding partner binds EDB, the sample is contacted with thermolysin, and the mass corresponding to the mass of a fragment of an oncofetal fibronectin indicating molecule can be, for example, 35 kDa, 85 kDa or 120 kDa. In some other methods, the oncofetal fibronectin binding partner binds IIICS, wherein the sample is contacted with *Aerhobacter* protease I and wherein the mass corresponding to the mass of a fragment of an oncofetal fibronectin indicating molecule is 14 kDa.

[0113] Some of the methods provided herein further include separating DNA from RNA in a sample. Such methods further include contacting the sample with a primer complementary to an oncofetal fibronectin-encoding nucleotide sequence, which primer contacts RNA in the sample, and treating the sample with one or more nucleic acid synthesis steps. In some methods, the primer is complementary to mRNA encoding oncofetal fibronectin. In some methods, a first nucleic acid synthesis step includes nucleic acid synthesis by reverse transcriptase. In some such methods, the binding partner specifically binds to a nucleotide sequence encoding oncofetal fibronectin, a nucleotide sequence complementary to a nucleotide sequence encoding oncofetal fibronectin, or a fragment thereof.

[0114] The methods provided herein also can be directed to methods wherein the binding partner specifically binds to a nucleotide sequence encoding the EDA region of oncofetal

fibronectin, a nucleotide sequence complementary to a nucleotide sequence encoding the EDA region of oncofetal fibronectin, a nucleotide sequence encoding the EDB region of oncofetal fibronectin, a nucleotide sequence complementary to a nucleotide sequence encoding the EDB region of oncofetal fibronectin, a nucleotide sequence encoding the IIICS region of oncofetal fibronectin, a nucleotide sequence complementary to a nucleotide sequence encoding the IIICS region of oncofetal fibronectin, or a fragment thereof.

[0115] In another embodiment, provided herein are methods of detecting an oncofetal fibronectin indicating molecule in a sample, by contacting the sample with a first fibronectin or oncofetal fibronectin binding partner immobilized to a solid support, contacting the solid support with a second fibronectin or oncofetal fibronectin binding partner, contacting the solid support with a third fibronectin or oncofetal fibronectin binding partner, and detecting any complex formed between an oncofetal fibronectin indicating molecule in the sample and the first fibronectin or oncofetal fibronectin binding partner, and either the second fibronectin or oncofetal fibronectin binding partner or the third fibronectin or oncofetal fibronectin binding partner, or both, whereby detection of complex is indicative of the presence of the oncofetal fibronectin indicating molecule in the sample, and wherein at least one of the first fibronectin or oncofetal fibronectin binding partner, the second fibronectin or oncofetal fibronectin binding partner, and the third fibronectin or oncofetal fibronectin binding partner is an oncofetal fibronectin binding partner. In some methods, the first oncofetal fibronectin binding partner binds a region of an oncofetal fibronectin indicating molecule can be, for example, EDA, EDB, IIICS, or a combination thereof. In some methods, the first oncofetal fibronectin binding partner binds a region of an oncofetal fibronectin indicating molecule other than EDA, EDB or IIICS. In some methods, the second oncofetal fibronectin binding partner binds a region of an oncofetal fibronectin indicating molecule such as, for example, EDA, EDB, IIICS or a combination thereof. In some methods, third oncofetal fibronectin binding partner binds a region of an oncofetal fibronectin indicating molecule such as, for example, EDA, EDB, IIICS or a combination thereof. In some methods, the second and third oncofetal fibronectin binding partners bind to a region of an oncofetal fibronectin indicating molecule such as, for example, EDA, EDB, IIICS or a combination thereof, wherein the second and third oncofetal fibronectin binding partners do not bind to the same oncofetal fibronectin indicating molecule region. Some methods further include contacting the sample with a fourth fibronectin or oncofetal fibronectin binding partner. In some such methods, the fourth fetal or oncofetal fibronectin binding partner binds a region of an oncofetal fibronectin indicating molecule such as, for example, EDA, EDB, IIICS or a combination thereof. In some methods, the second, third and fourth fetal or oncofetal fibronectin binding partners bind to a region of an oncofetal fibronectin indicating molecule such as, for example, EDA, EDB, IIICS or a combination thereof, wherein the second, third and fourth fetal or oncofetal fibronectin binding partners do not bind to the same oncofetal fibronectin indicating molecule region.

[0116] The IIICS portion can be any one of V64, V89, V95 and V120. When the oncofetal fibronectin indicating molecule is a IIICS portion of oncofetal fibronectin protein, the fibronectin or oncofetal fibronectin binding partner can

recognize a post-translational modification of oncofetal fibronectin protein. In one aspect, the post-translational modification can be O-glycosylation of threonine 33 of IIICS.

[0117] In one aspect the oncofetal fibronectin indicating molecule is identified as lacking EDA, EDB or IIICS. If the oncofetal fibronectin indicating molecule is identified as lacking, the portion of IIICS can be amino acids 1-25 of IIICS, amino acids 90-120 of IIICS or both.

[0118] In some methods, the complex is detected by measuring products of a nucleic acid synthesis reaction. In some methods, the first binding partner is immobilized to a test strip.

[0119] In another embodiment, provided herein are methods of detecting an oncofetal fibronectin indicating molecule in a sample, by contacting the sample with a first fibronectin or oncofetal fibronectin binding partner, contacting the sample with a second oncofetal fibronectin binding partner, and detecting by flow cytometry any complex formed between an oncofetal fibronectin indicating molecule in the sample, the first fetal or oncofetal fibronectin binding partner, and the second oncofetal fibronectin binding partner, whereby detection of complex is indicative of the presence of the oncofetal fibronectin indicating molecule in the sample. In some methods, complexes are detected by detecting spatial proximity between the first binding partner and the second binding partner. In some methods, complexes are detected by detecting two spatially proximal signals, wherein the first detected signal arises from the first binding partner and the second detected signal arises from the second binding partner. In some methods, the first binding partner and the second binding partner are bound to the same or different oncofetal fibronectin indicating molecules on the surface of a cell.

[0120] In the methods provided herein, the sample can be, for example a tissue or cell sample, or a liquid sample. A tissue or cell sample can be any of the following: lung, breast, ovary, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, prostate, thyroid, pituitary, eye, brain, oral, skin, head and neck cancer, lymphoma, leukemia, squamous cell carcinoma, adenocarcinoma, small cell carcinoma, melanoma, glioma, sarcoma neuroblastoma, semen, stool and a fraction or component thereof. A liquid sample can be a body fluid or tissue such as, for example, urine, blood, plasma, serum, saliva, lavage, cervical fluid, cervicovaginal fluid, vaginal fluid, breast fluid, breast milk, synovial fluid, seminal fluid, sputum, cerebral spinal fluid, tears, interstitial fluid, follicular fluid, amniotic fluid, aqueous humor, vitreous humor, peritoneal fluid, ascites, sweat, lymphatic fluid, and a fraction or component thereof. In some methods, the sample is treated with one or more reagents prior to the detecting step. Some methods further include removing sample components not bound to the binding partner prior to the step of removing oncofetal fibronectin indicating molecule or fragment thereof from the complex.

[0121] In the methods provided herein, the binding partner can contain an energy absorbing moiety. In some methods, the step of removing is accomplished by ultraviolet Matrix-Assisted Laser Desorption/Ionization, infrared Matrix-Assisted Laser Desorption/Ionization, or electrospray ioniza-

tion. In some methods, the oncofetal fibronectin indicating molecule or fragment thereof is detected by a mass spectrometry method such as, but not limited to, time-of-flight, Fourier transform, or magnetic sector/magnetic deflection. Some methods provided herein can further include the step of enhancing a signal from the oncofetal fibronectin indicating molecule bound to the binding partner. The methods provided herein can further include quantitating the amount of oncofetal fibronectin indicating molecule in the sample.

[0122] In some of the methods provided herein, presence of the oncofetal fibronectin indicating molecule in a sample determines a risk or identifies a health problem associated with oncofetal fibronectin, or indicates an increased risk of imminent or preterm delivery.

[0123] In some embodiments, methods are provided for indicating oncofetal fibronectin in a subject by detecting the presence of autoantibodies specific for oncofetal fibronectin in a sample from a subject, wherein the presence of anti-oncofetal fibronectin autoantibodies in the sample indicates the presence oncofetal fibronectin in the subject. Some methods can include indicating an oncofetal fibronectin associated health problem in a subject by detecting the presence of autoantibodies specific for oncofetal fibronectin in a sample from a subject according to the methods provided herein, wherein the presence of anti-oncofetal fibronectin autoantibodies in the sample indicates the presence of an oncofetal fibronectin associated health problem in the subject. Some methods can include indicating oncofetal fibronectin in a subject by detecting the presence of autoantibodies specific for oncofetal fibronectin in a sample from a subject according to the methods provided herein, wherein the presence of anti-oncofetal fibronectin autoantibodies in the sample indicates the presence oncofetal fibronectin in the subject. Such methods can further include contacting the sample with an anti-oncofetal fibronectin autoantibody binding partner, and detecting complexes formed between the binding partner and an anti-oncofetal fibronectin autoantibody.

[0124] Also provided herein are methods for indicating an oncofetal fibronectin associated health problem in a subject by indicating the presence of autoantibodies specific for oncofetal fibronectin in a subject according to the methods provided herein, wherein the presence of anti-oncofetal fibronectin autoantibodies in the subject indicates the presence of an oncofetal fibronectin associated health problem in the subject. In some such methods, the binding partner is oncofetal fibronectin protein or a fragment thereof, or an antibody that specifically binds human antibodies, or a fragment thereof. Some methods further include a second binding partner, wherein the second binding partner is oncofetal fibronectin protein or a fragment thereof, or an antibody that specifically binds human antibodies, or a fragment thereof.

[0125] The methods provided herein can include methods in which the sample is contacted with a non-specific binding compound under conditions in which substantially no oncofetal fibronectin indicating molecule in the sample binds to the non-specific binding compound. In some methods, the sample is contacted with a non-specific binding surface of a solid support under conditions in which substantially no oncofetal fibronectin indicating molecule in the sample binds to the non-specific binding surface. In some

methods, the non-specific binding surface contains a non-specific binding compound immobilized thereto. In some methods, 10% or less of the oncofetal fibronectin indicating molecule in the sample binds to the non-specific binding compound or surface. Some methods can further include contacting a sample with a solution that reduces non-specific binding of background material to a fibronectin or oncofetal fibronectin binding partner. In some methods, the ionic strength of the sample when contacted by the binding partner ranges from 50μ to 350μ or from about 50μ to about 350μ . Some methods can further include contacting a sample with solution that increases specific binding of oncofetal fibronectin indicating molecule in the sample to a fibronectin or oncofetal fibronectin binding partner. Some methods can further include a step of increasing the relative amount of oncofetal fibronectin indicating molecule in a sample. Some methods can further include decreasing the amount of background material in a sample that non-specifically binds to oncofetal fibronectin binding partner.

[0126] In some methods provided herein, the sample is, for example, urine, lavage, breast milk, cervicovaginal swab, saliva, serum, plasma, blood and interstitial fluid. In some methods, prior to the detecting step, one or more reagents is added to a urine sample. In some methods, the non-specific binding compound can be, for example, albumin, casein, fetal calf serum, gelatin and an antibody that does not specifically bind to an oncofetal fibronectin indicating molecule, for example, a non-specific binding compound can be methylated bovine serum albumin.

[0127] In some of the methods provided herein, a sample positive for oncofetal fibronectin identifies the subject from whom the sample was collected as having cancerous (e.g., neoplastic, malignant or metastatic) cells. In some such methods, the cancerous (e.g., neoplastic, malignant or metastatic) cells originate from bladder, kidney, prostate, cervix or ovary. In a particular method, the cancerous cells originate from bladder.

[0128] In other methods provided herein, a sample positive for oncofetal fibronectin identifies the subject from whom the sample was collected as having an increased risk of imminent or preterm delivery. In some such methods, a sample positive for oncofetal fibronectin identifies the subject from whom the sample was collected as one for whom induction is likely to be successful. Such methods can be used to identify a subject as one for whom induction of delivery is likely to be successful, and to administer to the subject an induction procedure. Likelihood of successful induction can be indicated by any of the following: increased likelihood of vaginal delivery upon induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between initiating induction and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between administering a parturifacient and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, likely decreased time interval between administering oxytocin and delivery relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, increased likelihood of delivering within 24 hours of

induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome, increased likelihood of delivering within 48 hours of induction relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicator of induction outcome and decreased likelihood of more than one administration of pre-induction agent to the subject relative to subjects negative for oncofetal fibronectin and/or having a negative result for the second indicators of induction outcome, and combinations thereof. The second indicator of induction outcome can be any of a measurement or observation of the pregnant subject, a measurement or observation of the fetus(es), and medical history of the pregnant subject. Such indicators include, but are not limited to, cervical length, Bishop score, effacement, parity, cervical dilation, gestational age, body mass index, station, consistency, transvaginal ultrasound, and digital examination, or a combination thereof.

[0129] In some methods, the first binding partner is conjugated to a moiety such as, but not limited to, colloidal metal, photodetectable latex microsphere, chromophore, fluorescent moiety, quantum dot and detectable enzyme. Some methods further include contacting the sample with a binding partner that specifically binds the first fibronectin or oncofetal fibronectin binding partner. In some such methods, the binding partner that specifically binds the first fibronectin or oncofetal fibronectin binding partner is an antibody or a nucleic acid molecule.

[0130] In some methods provided herein, the first fibronectin or oncofetal fibronectin binding partner is an anti-fibronectin antibody, or a fragment thereof. In some methods, the second fibronectin or oncofetal fibronectin binding partner is an anti-fibronectin antibody, or a fragment thereof.

[0131] In some of the methods provided herein, complexes are detected by determining if any first binding partner is in spatial proximity to the second binding partner, whereby detection of any first and second binding partners in spatial proximity indicates presence of an oncofetal fibronectin molecule in a sample. In some such methods, spatial proximity is detected as a result of a non-radioactive energy transfer reaction. In some such methods, the non-radioactive energy transfer reaction is, for example, fluorescence energy transfer (FET), fluorescence resonance energy transfer (FRET), or homogeneous time-resolved fluorescence (HTRF).

[0132] In the methods provided herein, the oncofetal fibronectin indicating molecule can be, for example, an oncofetal fibronectin protein, a nucleic acid molecule encoding oncofetal fibronectin, a nucleic acid molecule complementary to a nucleic acid molecule encoding oncofetal fibronectin, an autoantibody for oncofetal fibronectin protein, an autoantibody for a nucleic acid molecule encoding oncofetal fibronectin and a fragment thereof.

[0133] In some methods, presence of an oncofetal fibronectin indicating molecule below a threshold level classifies the sample as oncofetal fibronectin negative. In some methods, presence of an oncofetal fibronectin indicating molecule equal to or above a threshold level classifies the sample as oncofetal fibronectin positive. In some such methods, the oncofetal fibronectin indicating molecule is an oncofetal fibronectin protein and the threshold level of oncofetal fibronectin protein is 50 ng/mL.

[0134] In some methods provided herein, the amount of oncofetal fibronectin indicating molecule detected is compared to one or more thresholds, wherein the sample is classified according to the highest threshold that is less than or equal to the detected amount of oncofetal fibronectin indicating molecule. In some such methods, a first threshold is 50 ng/mL and a second threshold is 150 ng/mL. In some methods, the subject from whom the sample is collected is pregnant and the one or more threshold levels correspond to increasing likelihood of imminent pregnancy termination.

[0135] In some methods, the complex between the oncofetal fibronectin indicating molecule and the fibronectin or oncofetal fibronectin binding partner is measured by detecting the oncofetal fibronectin indicating molecule or a fragment thereof. In some methods, the oncofetal fibronectin indicating molecule is detected by mass spectrometry or gel electrophoresis. In some methods, the complex between the oncofetal fibronectin indicating molecule and the fibronectin or oncofetal fibronectin binding partner is measured by detecting the fibronectin or oncofetal fibronectin binding partner. In such methods, the fibronectin or oncofetal fibronectin binding partner is detected by detecting fluorescence, reflectance, absorption, bioluminescence, enzyme-linked detectable signal, or radioactive decay. In some methods, the fibronectin or oncofetal fibronectin binding partner is immobilized to a test strip.

[0136] The methods provided herein can include methods in which the fibronectin or oncofetal fibronectin binding partner binds to the EDA-specific portion, EDB-specific portion, or IIICS-specific portion of an oncofetal fibronectin indicating molecule as described above.

[0137] Also provided herein are test strips. Exemplary test strips include an analyte binding region containing a first fibronectin or oncofetal fibronectin binding partner immobilized on a solid support. Such test strip can contain a non-specific binding region, wherein the analyte binding region is downstream of the sample fluid flow pathway relative to the non-specific binding region. Some test strips can further contain a conjugate pad, which serves as a sample application component, an absorbent pad, which serves to draw liquid continuously through the device, wherein the materials of the membrane system form a single fluid flow pathway, and a porous or bibulous member in fluid communication with the absorbent pad and conjugate pad, which porous or bibulous member accommodates a liquid sample and serves as the solid support upon which molecule interactions occur, wherein the porous or bibulous member contains the analyte binding region. Some test strips can further contain a mobilization region containing a second fibronectin or oncofetal fibronectin binding partner, wherein the second fibronectin or oncofetal fibronectin binding partner is mobilized upon contact with the sample and the mobilization region is upstream of the analyte binding region. Some test strips can further contain a control region containing a molecule that specifically binds the second fibronectin or oncofetal fibronectin binding partner, wherein the control region is downstream of the analyte binding region. In some test strips, the first fibronectin or oncofetal fibronectin binding partner binds oncofetal fibronectin in preference to fibronectin. In some test strips, the non-specific binding region contains a non-specific binding pro-

tein immobilized to the solid support, wherein the non-specific binding protein is, for example, BSA, methylated BSA, W632 or mouse IgG.

[0138] Also provided herein are combinations containing a first binding partner and a sample collection device. Combinations provided herein can contain a first binding partner and a solid support. Combinations provided herein can contain a first binding partner and a second binding partner. Combinations provided herein can contain a first binding partner and a non-specific binding compound. Combinations provided herein can contain a first binding partner and a parturifacient. The combinations provided herein also can contain a non-specific binding compound. Also provided herein are kits containing the combinations provided herein and further containing instructions for use and/or a system for classifying the sample according to one or more threshold levels.

[0139] In another embodiment, provided herein are probes for detecting an oncofetal fibronectin indicating molecule, containing a mass spectrometry substrate, and a fibronectin or oncofetal fibronectin binding partner immobilized on the substrate. In some probes, the substrate contains a substance such as, but not limited to, glass, metal, ceramic, Teflon coated magnetic material, organic polymer, biopolymer and inorganic polymer.

[0140] Also provided herein are methods of characterizing oncofetal fibronectin protein in a sample, by contacting a sample with a first oncofetal fibronectin binding partner that binds oncofetal fibronectin protein in preference to non-oncofetal fibronectin protein, contacting a sample with a second oncofetal fibronectin binding partner that binds oncofetal fibronectin protein in preference to non-oncofetal fibronectin protein, and detecting complexes between oncofetal fibronectin protein and either the first oncofetal fibronectin binding partner, the second oncofetal fibronectin binding partner, or both binding partners, whereby the presence or absence of two domains of oncofetal fibronectin protein in the sample is determined. In such methods, the presence or absence of two domains of individual oncofetal fibronectin proteins in the sample is determined.

[0141] Also provided herein are methods of characterizing oncofetal fibronectin protein in a sample, by contacting a sample with a first oncofetal fibronectin binding partner that binds oncofetal fibronectin protein in preference to non-oncofetal fibronectin protein, contacting a sample with a second oncofetal fibronectin binding partner that binds oncofetal fibronectin protein in preference to non-oncofetal fibronectin protein, contacting a sample with a third oncofetal fibronectin binding partner that binds oncofetal fibronectin protein in preference to non-oncofetal fibronectin protein, and detecting complexes between oncofetal fibronectin and any combination of the first oncofetal fibronectin binding partner, the second oncofetal fibronectin binding partner and the third oncofetal fibronectin binding partner, whereby the presence or absence of three domains of oncofetal fibronectin protein in the sample is determined. In such methods, the presence or absence of three domains of individual oncofetal fibronectin proteins in the sample can be determined. In some methods, the first oncofetal fibronectin binding partner preferentially binds oncofetal fibronectin protein containing an EDA domain, an EDB domain, a IIICS domain, or a combination thereof. In some

such methods, the second oncofetal fibronectin binding partner preferentially binds oncofetal fibronectin protein containing an EDA domain, an EDB domain, a HIICS domain, or a combination thereof and wherein the first and the second binding partners do not bind to the same region of oncofetal fibronectin protein. In some such methods, the third oncofetal fibronectin binding partner preferentially binds oncofetal fibronectin protein containing an EDA domain, an EDB domain, a HIICS domain, or a combination thereof and wherein the first, the second and the third binding partners do not bind to the same region of oncofetal fibronectin protein. In some methods, complexes are detected by measuring oncofetal fibronectin protein, or a fragment thereof. In some such methods, oncofetal fibronectin is dissociated from one or more of the fibronectin or oncofetal fibronectin binding partners and a mass of oncofetal fibronectin protein, or a fragment thereof, is measured. In some methods, complexes are measured by detecting one or more of the fibronectin or oncofetal fibronectin binding partners. In some such methods, the first, second and/or third oncofetal fibronectin binding partners are measured by detecting fluorescence, reflectance, absorption, bioluminescence, enzyme-linked detectable signal or radioactive decay. In some methods, at least one oncofetal fibronectin binding partner is immobilized to a solid support.

[0142] Also provided herein are methods for classifying the level of oncofetal fibronectin in a sample, by measuring the amount of an oncofetal fibronectin indicating molecule in a sample, comparing the amount of oncofetal fibronectin indicating molecule in a sample to two or more threshold levels, and classifying the amount of oncofetal fibronectin indicating molecule in a sample according to the highest threshold level that is less than or equal to amount of oncofetal fibronectin indicating molecule in the sample. In some such methods, each threshold level is correlated with a higher risk of preterm, impending and/or imminent delivery, increased ability to predict delivery date, decreased likelihood of maintaining pregnancy, increased benefit from using methods of preventing preterm delivery, increased likelihood of success in inducing delivery, increased likelihood of delivery within a predetermined time, presence of cancerous (e.g., malignant neoplastic) disease, an increased risk of developing cancerous disease or recurrence of cancerous disease, a faster progression of the cancerous disease, or a more aggressive cancerous disease, or a decreased efficacy of cancerous disease therapy for the subject that provided the sample. In some methods, one threshold level is about or is 50 ng/mL, or is about or is 150 ng/mL.

[0143] Also provided herein are methods for classifying a sample as oncofetal fibronectin positive or negative, by measuring the amount of an oncofetal fibronectin indicating molecule in a sample, comparing the measured sample oncofetal fibronectin indicating molecule amount to a threshold level of 150 ng/ml, and classifying the sample as oncofetal fibronectin positive if the amount of oncofetal fibronectin indicating molecule is equal to or greater than the threshold level, otherwise classifying the sample as oncofetal fibronectin negative. Also provided herein are methods for detecting the presence of an oncofetal fibronectin indicating molecule in a sample, by contacting a sample with a first fibronectin or oncofetal fibronectin binding partner and a second fibronectin or oncofetal fibronectin binding partner, and measuring formation of a complex of oncofetal fibronectin, the first binding partner and the second binding

partner by detecting fluorescence from non-radioactive energy transfer. In some methods, the non-radioactive energy transfer occurs by fluorescence energy transfer (FET), fluorescence resonance energy transfer (FRET), or homogeneous time-resolved fluorescence (HTRF). In some methods, a binding partner is conjugated to a fluorescent dye or quantum dot.

[0144] Also provided herein are methods for detecting the presence of an oncofetal fibronectin indicating molecule in a sample, by contacting a sample with a fibronectin or oncofetal fibronectin binding partner, measuring formation of a complex of an oncofetal fibronectin indicating molecule and the binding partner by detecting fluorescence polarization indicative of complex formation. In some methods, the fluorescence polarization measurement indicates the quantity of oncofetal fibronectin indicating molecule in the sample. In some methods, the fluorescence polarization measurement indicates the mass of the complex. In some methods, the binding partner is conjugated to a fluorescent dye or quantum dot.

[0145] Some methods provided herein can further include, prior to contacting the sample, contacting the fibronectin or oncofetal fibronectin binding partner with an oncofetal fibronectin indicating molecule or analog thereof conjugated to a fluorescent dye or quantum dot, wherein signal dissipation or change indicates complex formation of a sample oncofetal fibronectin indicating molecule and the binding partner.

[0146] Also provided herein are conjugates that contain a fibronectin or oncofetal fibronectin binding partner linked to a leukocyte binding partner. In some such conjugates, the leukocyte binding partner is a natural killer cell binding partner. In some conjugates the leukocyte binding partner is an antibody. In some conjugates, the fibronectin or oncofetal fibronectin binding partner is FDC-6.

[0147] Also provided herein are methods for detecting an oncofetal fibronectin indicating molecule in a sample, by detecting the molecular weight of an oncofetal fibronectin indicating molecule or fragment thereof, by mass spectrometry to thereby detect the presence of an oncofetal fibronectin indicating molecule. The oncofetal fibronectin indicating molecule can be, for example, a oncofetal fibronectin protein or fragment thereof or an oncofetal fibronectin nucleic acid or fragment thereof.

[0148] Provided herein is the use of any of the products provided herein for use in the preparation of a medicament for any of the methods provided herein, including, but not limited to, diagnosis, imaging, and/or treatment of a health disorder characterized by the presence of, or elevated levels of oncofetal fibronectin.

[0149] Provided are methods for detection of an oncofetal fibronectin indicating molecule and methods for obtaining samples for use in methods of detection. Also provided are products for use in methods of detection of oncofetal fibronectin and/or of molecules indicative thereof in samples.

[0150] Provided are methods for assessing the suitability of concepti for implantation and selecting concepti for implantation. Concepti that produce oncofetal fibronectin can be used in implantation methods with greater success than concepti that do not produce oncofetal fibronectin.

[0151] Provided herein are methods for assessing a conceptus for implantation. These methods include testing a conceptus sample to detect an oncofetal fibronectin indicating molecule, wherein a conceptus for which a sample is oncofetal fibronectin positive is suitable for implantation. Also provided herein are methods for selecting concepti for implantation, by testing one or more conceptus samples to detect oncofetal fibronectin indicating molecule, and selecting a conceptus or concepti for implantation that yielded an oncofetal fibronectin positive sample.

[0152] In some such methods, a conceptus for which a sample is oncofetal fibronectin negative is not suitable for implantation. In some methods, presence of any oncofetal fibronectin indicating molecule in the sample identifies the sample as oncofetal fibronectin positive. In other methods, an amount of oncofetal fibronectin indicating molecule in the sample equal to or greater than a predetermined threshold level identifies the sample as oncofetal fibronectin positive. The threshold level can be a level predetermined to indicate that the conceptus will implant. Provided herein, the oncofetal fibronectin indicating molecule comprises a fibronectin III connecting segment (IIICS), an EDA or EDB segment or an autoantibody that specifically binds to IIICS, EDA or EDB.

[0153] In some embodiments of the methods provided herein, samples from two or more concepti are assayed, and a first conceptus for which a sample contains a higher amount of oncofetal fibronectin indicating molecule is identified as more suitable for implantation than a second conceptus for which a sample contains a lower amount of oncofetal fibronectin indicating molecule. In other embodiments of the methods provided herein, subsequent to testing the first conceptus sample, testing a second conceptus sample from the same conceptus to detect an oncofetal fibronectin indicating molecule, wherein a conceptus for which the second sample contains more oncofetal fibronectin indicating molecule than the first sample is suitable for implantation. In such embodiments, two or more samples from each of two or more concepti are assayed, and a first conceptus for which a sample contains a higher rate of increasing amount of oncofetal fibronectin indicating molecule is identified as more suitable for implantation than a second conceptus for which a sample contains a lower rate of increasing amount or a decreasing amount of oncofetal fibronectin indicating molecule. Suitability for implantation of a conceptus can increase with increasing amounts of oncofetal fibronectin indicating molecule or increasing rate of increase of oncofetal fibronectin indicating molecule can indicate suitability of a conceptus for implantation.

[0154] In the methods provided herein, the conceptus sample can be selected from among conceptus extract, sample from outside of the conceptus, and an extract of a cell from the conceptus. In some methods the sample can be treated with a reagent and/or fractionated prior to the step of testing for an oncofetal fibronectin indicating molecule in a conceptus sample. In one exemplary sample, from outside the conceptus, the sample is an extract of culture medium in which a single conceptus has been cultured.

[0155] The methods provided herein can further include determining an additional maternal or conceptus marker, wherein an oncofetal fibronectin positive conceptus having a favorable additional markers is identified as a conceptus favorable for implantation. In such methods the additional marker can be detected in a conceptus sample, can be determined by visual inspection of the conceptus, or can be

detected in a maternal sample. Exemplary additional markers can be selected from among genetic composition of the conceptus, gene expression of the conceptus, and morphology of the conceptus. In methods in which the additional marker is morphology of the conceptus, the morphology of the conceptus can be graded according to factors, such as cell number, degree of fragmentation, cell regularity, symmetry, pronuclear morphology, follicle size, follicular fluid volume, multi-nucleation, presence of vacuoles, granularity and combinations thereof.

[0156] Provided herein is the use of any of the products provided herein for use in the preparation of a medicament for any of the methods provided herein, including, but not limited to, diagnosis, imaging, and/or treatment of a health disorder characterized by the presence of, or elevated levels of oncofetal fibronectin.

BRIEF DESCRIPTION OF THE DRAWINGS

[0157] FIG. 1 depicts a schematic of fibronectin, including domain organization, protein interaction sites, proteolysis sites and ligand interaction sites. This figure is reproduced from the publication by Pankov et al., *J. Cell Science* 2002 115:3861-3863.

[0158] FIG. 2 depicts an overhead view of an exemplary test strip for oncofetal fibronectin indicating molecule detection.

[0159] FIG. 3 depicts a side view of an exemplary test strip for oncofetal fibronectin indicating molecule detection.

[0160] FIG. 4 depicts an exemplary test strip holder for oncofetal fibronectin indicating molecule detection.

DETAILED DESCRIPTION

[0161]

Outline

- A. Definitions
- B. Detection of Oncofetal Fibronectin
- C. Structure and Properties of Fibronectin
 - 1. Structural Characteristics of Fibronectin
 - 2. Binding Properties and Proteolysis of Fibronectin
 - 3. Oncofetal Fibronectin
 - a. Structural Features of Oncofetal Fibronectin
 - b. Molecules that Bind to Oncofetal Fibronectin
 - c. Proteolysis of Oncofetal Fibronectin
- D. Use of Oncofetal Fibronectin as a Biological Marker
 - 1. Pregnancy Indications
 - a. Likelihood of Pre-term delivery
 - b. Preventing Pre-term delivery
 - c. Predictor of Delivery Date
 - d. Use with Inducing Delivery
 - i. Induction Methods and Compounds
 - ii. Post-Induction Measurements
 - e. Conceptus Indications
 - i. Detection of Oncofetal Fibronectin Production by a Conceptus
 - ii. Assisted Reproduction Technology Related Uses
 - iii. Post-Measurement Steps
 - a. Increasing Oncofetal Fibronectin Production
 - b. Identify Conceptus Based on Oncofetal Fibronectin Production
 - i. Selection of a Conceptus
 - ii. Criteria for Selection
 - c. Other Markers Used in Conjunction with Oncofetal Fibronectin
 - i. Conceptus Markers
 - ii. Maternal Markers
 - iv. Stem Cell Related Uses

-continued

Outline
2. Indicator of Cancer
a. Bladder Cancer
b. Breast Cancer
c. Cervical Cancer
d. Ovarian Cancer
e. Prostate Cancer
f. Lung Cancer
g. Colorectal Cancer
h. Additional Cancers
3. Health State Assessment
4. Other Health Problems
a. Arthritis
b. Diabetic Retinopathy
c. Dupuytren's Contracture
E. Collection of Samples
1. Swab and Cervicovaginal Samples
2. Lavage Samples
a. Sample Collection
b. Lavage Fluid
c. Applying a Label with Lavage
d. Ductal Lavage
i. Sample Collection
ii. Applying a Label to a Duct
iii. Lavage Fluid
3. Collection of Urine Samples
a. Sample Handling
b. Sample Condition Modification
i. Ionic Strength
ii. Ionic Strength Testing
iii. Normalization
c. Sample Treatment
i. Non-specific Binding
ii. Filtration
4. Interstitial Fluid
F. Methods of Detecting Oncofetal Fibronectin
1. Compounds and Compositions in Detecting Oncofetal Fibronectin
a. Molecules that Indicate the Presence of Oncofetal Fibronectin
b. Fibronectin Portions Indicative of Oncofetal Fibronectin
c. Autoantibody to Oncofetal Fibronectin or to a Nucleic Acid Molecule Encoding Oncofetal Fibronectin
d. Binding Partner
i. Antibodies
a. Antibodies for Oncofetal Fibronectin
b. Conjugation of the Antibody to a Label
ii. Nucleic Acid Molecules
iii. Binding Partners to Autoantibodies
iv. Additional Binding Partners
v. Binding Partners That Bind a Region of Oncofetal Fibronectin
e. Non-Specific Binder
2. Assays for Detecting Oncofetal Fibronectin Complexed with a Binding Partner
a. Solution Detection
i. Signal Indicative of Complex Formation
ii. Molecular Weight Corresponding to Complex
b. Immobilized Sample
i. Dot Blot Analysis
ii. Western Blot Analysis
iii. Southern and Northern Blot Analyses
iv. In Situ Analysis
a. Tissue or Organ Samples
b. Detection in a Subject
c. Treatment in a Subject
c. Immobilized Binding Partner
i. Sandwich Assay
ii. Test Device
a. Test strip
b. Test Strip Housing
c. Analysis with a Test Device
iii. Quantitation
iv. Affinity-Based Isolation of Oncofetal Fibronectin
d. Detection of Regions of Oncofetal Fibronectin

-continued

Outline
3. Detection of Oncofetal Fibronectin by Mass Spectrometry
a. Sample Manipulation
i. Contact with Binding Partner
ii. Contact with Fragmentation Compound
a. Trypsin Proteolysis
b. Cathepsin D Proteolysis
c. Thermolysin Proteolysis
d. <i>Achromobacter</i> Protease I Proteolysis
iii. Solid Support
iv. Conditioning
v. Combinations of Sample Manipulation Steps
b. Substrate for Mass Spectrometry
c. Mass Spectrometric Analysis
i. Formation of Ions in the Gas Phase
ii. Detection
iii. Use of Mass Spectrometry for Detecting Oncofetal Fibronectin in a Sample
a. Direct Measurement
b. With Signal Enhancement
iv. Detection of Regions of Oncofetal Fibronectin
v. Quantitation of Oncofetal Fibronectin
4. Detecting Nucleic Acid Molecules
a. Detection Methods
b. Detection of RNA
i. Reverse Transcription
ii. cDNA Amplification
iii. Additional Components
iv. Nucleic Acid Synthesis Steps
v. Detection
a. DNA Detection Methods
b. Quantitation
c. Detection of Regions of Oncofetal Fibronectin
5. Detection of Autoantibodies to Oncofetal Fibronectin
6. Measurement of Other Analytes
a. Normalization
b. Combination with Other Markers
i. Indicators of membrane rupture
a. Insulin-Like Growth Factor Binding Protein
b. Hypochlorous Acid
ii. Estriol
iii. Other Tumor Indicators
G. Analysis of Detection Measurements
1. Quantitation
2. Thresholds
3. Identification of Tissue Source
H. Combinations, Probes, Conjugates and Kits
I. Examples

A. Definitions

[0162] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which the invention(s) belong. All patents, patent applications, published applications and publications, Genbank sequences, websites and other published materials referred to throughout the entire disclosure herein, unless noted otherwise, are incorporated by reference in their entirety. In the event that there are a plurality of definitions for terms herein, those in this section prevail. Where reference is made to a URL or other such identifier or address, it is understood that such identifiers can change and particular information on the internet can come and go, but equivalent information is known and can be readily accessed, such as by searching the internet and/or appropriate databases. Reference thereto evidences the availability and public dissemination of such information.

[0163] As used herein, fibronectins refer to a family of high molecular weight glycoproteins encoded by a single

gene. They occur in a soluble forms in plasma and other body fluids and in cellular forms in the extracellular matrices. The family of fibronectins includes at least twenty isoforms. These arise from alternative splicing in regions of the primary transcript of a single gene (see, e.g., SEQ ID NO: 37). and from post-translational modifications, generally in the EDA, EDB and IIIICS regions. The amino acid sequence of an exemplary fibronectin protein containing EDA, EDB and IIIICS (V64 splice variant) regions is set forth in SEQ ID NO: 38. SEQ ID NO: 15 sets forth the amino acid sequence of a fibronectin containing the complete EDA, EDB and IIIICS (V120 splice variant) domains.

[0164] Cellular and plasma fibronectins often occur as heterodimers containing similar polypeptides. Alternative splicing occurs in at least two regions of the pre-mRNA, causing variability in internal primary sequences. This, results in differences between the fibronectin subunits (Kornblihtt et al. *EMBO J.* 4(7): 1755-1759 (1985)). Plasma fibronectins generally lack extradomain A (EDA) and extradomain B (EDB) regions of fibronectin. Cellular fibronectins can have EDA and/or EDB included or excluded, depending on the cellular and extracellular contents. Another variable region, the fibronectin III connecting segment (IIIICS) region occurs in certain fibronectins and in variant forms by virtue of alternative splicing. The encoding nucleic acid contains multiple splice sites leading to variants in humans that include those designated V0, V120, V95, V89 and V64 (see, e.g., Khan et al. *Investigative Ophthalmology & Visual Science* 45(1): 287-295 (2004)).

[0165] Fibronectins designated oncofetal fibronectins, constitute a subset of cellular fibronectins. The oncofetal fibronectins include one or more of an EDA, EDB and/or IIIICS region or are determined to be oncofetal fibronectin by virtue of expression or altered expression, such by alternative splicing of the encoding gene to produce a truncated fibronectin (see, e.g. Schor et al. (2003) *Cancer Research* 63:8827-883; see, e.g., SEQ ID Nos. 26 and 27). Oncofetal fibronectins can be expressed at higher levels in tumor cells than normal cells and tissues and also in fetal cells and tissues (Kaczmarek et al. *Int. J. Cancer* 58: 11-16 (1994); Castellani et al., *Int. J. Cancer* 59: 612-618 (1994); and Taviani et al. *Int. J. Cancer* 56: 820-825 (1994)) or can be produced by virtue of alternative splicing in tumor cells or other cells and tissues is subjects with particular diseases. Fibronectins also can be associated with tumors and acquire metastatic potential by virtue of mutations. For example, tumor cells containing a fibronectin containing a single point mutation were found to have reduced fibronectin matrix formation and an increase in metastatic potential relative to tumor cells expressing wild-type fibronectin (Wang et al., *J. Exp. Med.*, 195: 1397-1406 (2002)).

[0166] As used herein, oncofetal fibronectin refers to this heterogeneous subset of fibronectin proteins that share characteristics domains and/or expression patterns. Oncofetal fibronectins generally are cellular fibronectins. In tumor cells and tissues the extracellular domain portion can be shed (see, e.g., Mardon et al., *J. Cell Sci.* 104: 783-792 (1993)) or splicing of the encoding nucleic acid can be altered. Oncofetal fibronectins include fibronectins that are expressed in or shed by tumors, by tissues or cells involved in other disease states and also in fetal cells tissues (also referred to as fetal fibronectins). Hence oncofetal fibronectins, while typically cellular fibronectins, can be detected in

connection with disease states and pregnancy-related conditions by virtue of expression of particular splice variants, increased expression and/or shedding from such tissues and cells by virtue of overexpression or proteolytic cleavage or other mechanism.

[0167] For purposes herein, oncofetal fibronectin proteins contain extra-domain A (EDA), extra-domain B (EDB), or fibronectin III connecting segment (IIIICS), or any combination thereof or are produced as a result of alternative splicing or post-translational events in tumor cells and tissues to produce for example truncated fibronectins. The group of oncofetal fibronectins includes fibronectins that result from alternative splicing of these three regions (EDA, EDB and IIIICS) in fibronectin and also from post-translational modifications and other variations in splicing of the encoding nucleic acid. The splicing and expression of oncofetal fibronectin is differently regulated in cells and tissues and at different developmental stages. In fetal tissues and in some abnormal cells and tissues, expression of fibronectin is altered to produce what is designated an oncofetal fibronectin or the expression of an oncofetal fibronectin is increased relative to the corresponding normal adult cells and tissues. In some normal adult cells, tissues and sample types, oncofetal fibronectin is not present in amounts detectable by antibody assay. Accordingly, for purposes of detection of oncofetal fibronectin, abnormal levels of oncofetal fibronectin can be determined by comparing the detected amount to a control or a to predetermined amount. The amino acid sequence of human fibronectins including EDA, EDB and IIIICS regions (SEQ ID NOS: 4, 6 and 8, respectively) and encoding nucleic acid molecules (SEQ ID NOS: 3, 5 and 7, respectively) are known in the art and are available in public databases. An exemplary sequence of a human EDA region also is set forth as amino acid residues 1432-1621 of SEQ ID NO: 2. An exemplary sequence of a human EDB region also is set forth as amino acid residues 963-1109 of SEQ ID NO: 2. An exemplary sequence of a human IIIICS region also is set forth as amino acid residues 1830-1949 of SEQ ID NO: 2.

[0168] The amino acid sequences of exemplary oncofetal fibronectins and the sequences of encoding nucleic acid molecules are set forth in SEQ ID NO. 1, 14, 16, 18, 20, 22, 24 and 26 and the encoded amino acid molecules are set forth in SEQ ID NOS: 2, 15, 17, 19, 21, 23, 25 and 27. Examples of oncofetal fibronectin variants that generally are absent in normal tissues and cells as detected by antibody-based assays include oncofetal fibronectins generated by O-glycosylation in the IIIICS splicing region and oncofetal ED-B containing fibronectin such as, for example, the amino acid sequence and the sequence of encoding nucleic acid molecule set forth in SEQ ID NOS: 25 and 24, respectively (see, e.g., Kaczmarek et al. *Int. J. Cancer* 58: 11-16 (1994); Castellani et al., *Int. J. Cancer* 59: 612-618 (1994); Midulla et al., *Cancer Res.* 60:164-169 (2000)).

[0169] The nucleic acid sequence for the entire human fn1 gene is set forth in SEQ ID NO: 37. The protein and encoding nucleic acid molecules for a variety of species including, for example, rat, mouse, chicken, cow and *Xenopus laevis* also are known and readily available in public databases. Oncofetal fibronectins include fibronectins that binds specifically to the FDC-6 monoclonal antibody (see, e.g., Matsuura and S. Hakomori, *Proc. Natl. Acad. Sci. USA*, 82:6517-6521 (1985)), which is produced by the hybridoma

(deposited at the American Type Culture Collection under accession number ATCC HB 9018; see also in U.S. Pat. No. 4,894,326, issued Jan. 16, 1990, to Matsuura et al.)

[0170] The fibronectin III connecting segment (IIICS) contains three separate splice regions that can be expressed in particular combinations, resulting in a variety of different sizes and sequences of a IIICS region present in oncofetal fibronectin. The three segments encode an N-terminal segment of 25 amino acids, a middle segment encoding 64 amino acids and a C-terminal segment encoding 31 amino acids, resulting in IIICS regions that, when present can contain 64 amino acids, 89 amino acids, 95 amino acids, or 120 amino acids. Exemplary sequences are set forth in SEQ ID Nos. 35, 33, 31 and 29, respectively, which are encoded by nucleic acid sequences set forth in SEQ ID Nos. 34, 32, 30 and 28, respectively.

[0171] An oncofetal fibronectin can be identified by specific binding of one or more anti-oncofetal fibronectin antibodies. Such antibodies bind with less affinity or do not bind to non-oncofetal fibronectin. A variety of anti-oncofetal fibronectin antibodies are known in the art, including, for example, IST-9 (Carnemolla et al., *FEBS Lett.* 215:269-273 (1987); available at Accurate Chemical & Sci. Corp., Westbury, N.Y.), DH1 (Vartio et al., *J. Cell Sci.* 88:419-430 (1987)), BC-1 (Carnemolla et al., *J. Cell Biol.* 108:1139-1148 (1989)), L19 (U.S. Pat. App. No. 20030176663), ME4C (Giovannoni et al., *Nucleic Acids Res.* 29:e27 (2001); the nucleic acid encoding sequence and the amino acid sequence for ME4C scFv recombinant antibody are provided as SEQ ID Nos.9 and 10, respectively; the sequences also are available at GenBank accession no. AJ297960), A134 (Islami et al., *Eur. J. Obstet. Gynecol. Reprod. Biol.*, 97:40-45 (2001)) FDC-6 (U.S. Pat. No. 4,894,326; ATCC HB 9018), 5C10 (Mandel et al., *APMIS* 100:817-826 (1992)) and X18A4, X20C4 and X8E3 (U.S. Pat. No. 5,523,229; ATCC Nos. HB-11587, HB-11589 and HB-11588, respectively). Antibodies that bind specifically and preferentially to oncofetal fibronectin also can be prepared. Methods for preparing such anti-oncofetal fibronectin antibodies are known in the art as exemplified in U.S. Pat. No. 4,894,326 and WO 02/46455.

[0172] Oncofetal fibronectin can be captured with a fibronectin binding partner that binds to most or all fibronectins, such as an integrin, heparin or an anti-fibronectin antibody, or oncofetal fibronectin can be captured with an oncofetal fibronectin binding partner, such as an anti-oncofetal fibronectin antibody.

[0173] As used herein, use of oncofetal fibronectin as a marker refers to detection of oncofetal fibronectin. This refers to detection of any oncofetal fibronectin indicating molecule, including an oncofetal fibronectin protein, a nucleic acid molecule encoding oncofetal fibronectin or a complement thereto, or an autoantibody that specifically binds oncofetal fibronectin protein or a nucleic acid molecule encoding oncofetal fibronectin, and fragments thereof indicative of oncofetal fibronectin.

[0174] As used herein, a fetal-restricted antigen refers to an antigen present in pregnant women uniquely, or in substantially elevated amounts compared to non-pregnant women. The fetal restricted antigen can be present in maternal serum, plasma, urine, saliva, sweat, tears and other bodily fluids. Oncofetal fibronectin can contain a fetal

restricted antigen and can be found in placenta, amniotic fluid and fetal connective tissue.

[0175] As used herein, a binding partner is a compound that specifically binds to a particular molecule or class of molecules. Binding partners can include proteins, nucleic acid molecules, carbohydrates, lipids, ligands, drugs, ions and any other compound that can specifically bind to a particular molecule. A fibronectin binding partner specifically binds to any fibronectin indicating molecule, including a fibronectin protein, an oncofetal fibronectin protein, an autoantibody to a fibronectin protein or fibronectin-encoding nucleic acid or a complement thereto, an autoantibody to an oncofetal fibronectin protein or oncofetal fibronectin-encoding nucleic acid, a nucleic acid encoding fibronectin or a complement thereto, a nucleic acid encoding oncofetal fibronectin or a complement thereto, and fragments of any of these.

[0176] As used herein, an oncofetal fibronectin binding partner is a molecule that specifically binds to an oncofetal fibronectin protein, an autoantibody to an oncofetal fibronectin protein or oncofetal fibronectin-encoding nucleic acid, a nucleic acid encoding oncofetal fibronectin or a complement thereto, and fragments of any of these. In particular, an oncofetal fibronectin binding partner binds to portions of an oncofetal fibronectin indicating molecule that are unique to oncofetal fibronectin, such as EDA, EDB or IIICS and also can bind to portions of oncofetal fibronectin that are present in non-oncofetal fibronectin protein and nucleic acid molecules, such as an FNIII₅ region, where binding of such a region occurs by virtue of the presence of EDA, EDB or IIICS in the molecule (see, e.g., Carnemolla et al. *J. Biol. Chem.* 267:24689-24692 (1992)).

[0177] As used herein, selective binding of a binding partner to the binding of a binding partner to a particular molecule with at least about 2-fold and typically at least about 5-fold, 10-fold, 50-fold, 100-fold, or more, greater affinity (K_a or K_{eq}) than for another molecule, or at least 2-fold and typically at least 5-fold, 10-fold, 50-fold, 100-fold, or more, greater affinity (K_a or K_{eq}) than for another molecule. Typical conditions for detecting and determining binding affinity constants or for determining the selectivity of binding include physiological conditions, such as PBS (137 mM NaCl, 2.7 mM KCl, 10 mM phosphate buffer pH 7.4). Binding partners that specifically bind, bind with a binding affinity K_a of typically at least about 10^7 l/mol, 10^8 l/mol or more. Generally, it refers to binding partners that selectively and specifically bind.

[0178] As used herein, a compound that binds preferentially to an oncofetal fibronectin indicating molecule, is a compound that binds to an oncofetal fibronectin indicating molecule in preference to binding to a non-oncofetal fibronectin molecule, where the preference can be manifested as at least about 2-fold higher affinity and typically at least about 5-fold, 10-fold, 50-fold, 100-fold, or more, higher affinity, or at least 2-fold higher affinity and typically at least 5-fold, 10-fold, 50-fold, 100-fold, or more, higher affinity. Preferential binding is selective and also, typically is specific, and demonstrates less than about 25% or 10% and typically less than about 5%, or less than 25% or 10% and typically less than 5%, cross-reactive nonspecific binding. For example, an antibody such as FDC-6 preferentially binds to oncofetal fibronectin protein over non-oncofetal

fibronectin protein because FDC-6 can bind to a IIICS-containing oncofetal fibronectin protein with much higher affinity than FDC-6 binds to fibronectin protein not containing the IIICS region. Typical conditions for performing such binding or for determining preferential binding include physiological conditions, such as PBS (137 mM NaCl, 2.7 mM KCl, 10 mM phosphate buffer pH 7.4).

[0179] As used herein, non-oncofetal fibronectin refers to a fibronectin protein or nucleic acid molecule that does not contain or encode an EDA, EDB and IIICS domains as detected by methods herein.

[0180] As used herein, an oncofetal fibronectin indicating molecule refers to any molecule associated with the expression or presence of oncofetal fibronectin. For example, an oncofetal fibronectin indicating molecule includes an oncofetal fibronectin protein or a fragment thereof, a nucleic acid encoding oncofetal fibronectin such as RNA or cDNA or a complement thereto, or an autoantibody to an oncofetal fibronectin protein or oncofetal fibronectin encoding nucleic acid molecule or an antibody fragment thereof and a fragment or fragments thereof.

[0181] As used herein, cancer refers to the growth of abnormal cells in the body in an uncontrolled manner; unlike benign tumors, these tend to invade surrounding tissues, and spread to distant sites of the body via the blood stream and lymphatic system. Cancer also refers to any of various malignant neoplasms characterized by the proliferation of anaplastic cells that tend to invade surrounding tissue and metastasize to new body sites. A cancer can be a solid tumor or a blood born cancer. As used herein, a tumor refers to an abnormal growth of tissue resulting from uncontrolled, progressive multiplication of cells and serving no physiological function or a neoplasm. A cancer cell, as used herein, refers to malignant neoplastic, anaplastic, metastatic, hyperplastic, dysplastic, neoplastic, malignant tumor (solid or blood-borne) cells that display abnormal growth in the body in an uncontrolled manner. Cancer can be of lung, prostate, bladder, cervical, kidney or ovarian tissue.

[0182] As used herein, neoplasm refers to new and abnormal growth of tissue, which can be cancerous, such as a malignant tumor.

[0183] As used herein, neoplastic disease, means a disease brought about by the existence of a neoplasm in the body.

[0184] As used herein, metastasis refers to the migration of cancerous cells to other parts of the body. As used herein, hyperplasia refers to an abnormal increase in the number of cells in an organ or a tissue with consequent enlargement. As used herein, neoplasm and dysplasia refer to abnormal growth of tissues, organs or cells. As used herein, malignant means an cancerous or tending to metastasize. As used herein, anaplastic means cells that have become less differentiated.

[0185] As used herein, leukemia refers to a cancer of blood cells. Any of various acute or chronic neoplastic diseases of the bone marrow in which unrestrained proliferation of white blood cells occurs, usually accompanied by anemia, impaired blood clotting, and enlargement of the lymph nodes, liver and spleen. Leukemia occurs when bone marrow cells multiply abnormally caused by mutations in the DNA of stem cells. Bone marrow stem cells, as used herein, refer to undifferentiated stem cells that differentiate into red

blood cells and white blood cells. Leukemia is characterized by an excessive production of abnormal white blood cells, overcrowding the bone marrow and spilling into peripheral blood. Various types of leukemias spread to lymph nodes, spleen, liver, the central nervous system and other organs and tissues.

[0186] As used herein, lymphoma generally refers to a malignant tumor that arises in the lymph nodes or other lymphoid tissue.

[0187] As used herein, detection of oncofetal fibronectin refers to detection of an oncofetal fibronectin indicating molecule, where a fragment can be formed using the methods described herein or known in the art, such as, but not limited to, proteolysis or PCR. Further in regard to this phrase, one skilled in the art recognizes that, even if not explicitly provided for herein, methods for detecting oncofetal fibronectin proteins or fragments also can be used for detecting other oncofetal fibronectin indicating molecules such as oncofetal fibronectin-encoding nucleic acid molecules or complements thereto, or fragments thereof, or autoantibodies to oncofetal fibronectin proteins or nucleic acids or antibody fragments thereof. Selection of any particular method for detecting an oncofetal fibronectin indicating molecule can be a matter of design choice, where one skilled in the art will know which method or methods (e.g., PCR, mass spectrometry, sandwich assay) to select according to the nature (e.g., protein, nucleic acid) of the oncofetal fibronectin indicating molecule detected.

[0188] As used herein, a subject includes any animal for whom diagnosis, screening, monitoring or treatment is contemplated. Animals include mammals such as primates and domesticated animals. An exemplary primate is human. A patient refers to a subject such as a mammal, primate, human or livestock subject afflicted with a disease condition or for which a disease condition is to be determined or risk of a disease condition is to be determined.

[0189] As used herein, sample generally refers to anything that contains an analyte for which an analyte assay is to be performed. For example, a sample can be a specimen from a subject, where presence or absence of an oncofetal fibronectin indicating molecule in the specimen is to be determined using, for example, the oncofetal fibronectin indicating molecule detection methods provided herein. A sample can be used in neat form (e.g., unmodified) or can be modified by adding one or more reagents such as a buffer and/or by one or more fractionation or separation steps. The sample can be a biological sample, such as a biological or body fluid sample or a biological tissue sample. Examples of biological or body fluid samples include urine, lymph, blood, plasma, serum, saliva, cervical fluid, cervicovaginal fluid, vaginal fluid, breast fluid, breast milk, synovial fluid, semen, seminal fluid, stool, sputum, cerebral spinal fluid, tears, mucus, interstitial fluid, follicular fluid, amniotic fluid, aqueous humor, vitreous humor, peritoneal fluid, ascites, sweat, lymphatic fluid, lung sputum and lavage or samples derived therefrom (e.g., reagent-modified and/or fractionated samples). Urine samples can be neat or frozen. Fluid samples can be analyzed as it is being provided (e.g., a urine stream dipstick), can be collected in a container, or can be collected with a swab. Exemplary swab samples include cervicovaginal swab samples, including, but not limited to swab of the point of a possible cervicovaginal lesion, the

cervical canal, the cervical os, the ectocervix, the transition zone on the cervix between squamous and columnar cells (i.e., the squamocolumnar junction), the vagina, the posterior fornix, the portion of the vagina below the posterior fornix such as the lower third of the vagina, the labia, or combinations thereof. Biological tissue samples are samples containing an aggregate of cells, usually of a particular kind, together with intercellular substances that form one of the structural materials of a human, animal, plant, bacterial, fungal or viral structure, including connective, epithelium, muscle and nerve tissues. Examples of biological tissue samples also include organs (e.g., breasts), tumors, lymph nodes, arteries and individual cell(s). For example, the sample can be a tissue sample suspected of being cancerous. Reference herein to any of the above fluid types or any tissue or cell sample also includes reagent-modified and fractionated samples. Thus, reference to a cervicovaginal sample also includes a buffer-treated cervicovaginal sample, and reference to a tissue sample includes the supernatant from a homogenate of that tissue.

[0190] As used herein, a normalizing analyte refers to an analyte used to normalizing the amount of oncofetal fibronectin indicating molecule in the sample according to the concentration of one or more normalization analytes in the sample. A normalizing agent is, for example, creatinine. For a vaginal swab sample, for example, the sample can be a swab of any portion of the vagina, including the posterior fornix or the portion of the vagina below the posterior fornix, such as, for example, the lower third of the vagina.

[0191] As used herein, below the posterior fornix refers to the portion of the vagina that includes the vaginal vestibule and regions of the vagina superior to the vaginal vestibule but inferior to the posterior fornix, which can include the vaginal vault, the lower third of the vagina, and the vaginal sphincter. Thus, a vaginal swab below the posterior fornix refers to a swab of the vaginal vestibule and regions of the vagina superior to the vaginal vestibule, but inferior to the posterior fornix, which can include the vaginal vault, the lower third of the vagina, and the vaginal sphincter. In the case of a labial swab, the swab can be from the labia minora or labia majora and typically includes a swab of the labia minora.

[0192] As used herein, cervicovaginal fluid can contain cervical fluid, vaginal fluid, or combinations thereof.

[0193] As used herein, an in vivo method refers to a method performed within the living body of a subject.

[0194] As used herein, a conjugate or a binding partner conjugated to a moiety refers to a complex that includes a binding partner bound to a moiety, where the binding between the binding partner and the moiety can arise from one or more covalent bonds or non-covalent interactions such as hydrogen bonds, or electrostatic interactions. A conjugate also can include a linker that connects the binding partner to the moiety. Included within the scope of conjugates are binding partners immobilized on a support such as a solid support. Exemplary conjugates include binding partners conjugated to a detectable moiety such as a detectable label, or a bindable moiety such as a bindable compound.

[0195] As used herein, a detectable moiety or an imaging moiety refer to moieties used to image an oncofetal fibronectin indicating molecule in any of the methods pro-

vided herein. Imaging moieties include, for example, fluorescent moieties, radionuclides, magnetically detectable isotopes or compounds, sonographic imaging agents, chromophores, latex microspheres, or quantum dots.

[0196] As used herein, a binding partner immobilized on a support such as a solid support refers to a binding partner bound to a support by covalent or non-covalent interactions. Binding to a support can be accomplished by a linker connected to the binding partner and the support, or the binding partner can be bound directly to the support.

[0197] As used herein, a detectable label or detectable moiety refers to an atom, molecule or composition, wherein the presence of the atom, molecule or composition can be directly or indirectly measured. Such a label can be detected, for example, by visual inspection, by fluorescence spectroscopy, by reflectance measurement, by flow cytometry, or by mass spectrometry. Direct detection of a detectable label refers to measurement of a physical phenomenon, such as energy or particle emission or absorption, of the moiety itself. Indirect detection refers to measurement of a physical phenomenon, such as energy or particle emission or absorption, of an atom, molecule or composition that binds directly or indirectly to the detectable moiety. In an example of indirect detection, a detectable label can be biotin, which can be detected by binding to avidin and avidin can be detected by, for example, binding avidin with a second biotin molecule linked to fluorescein. Thus, included within the scope of a detectable label or detectable moiety is a bindable label or bindable moiety, which refers to an atom, molecule or composition, wherein the presence of the atom, molecule or composition can be detected as a result of the label or moiety binding to another atom, molecule or composition.

[0198] A detectable label can be conjugated to a fibronectin or oncofetal fibronectin binding partner, or can specifically bind to a fibronectin or oncofetal fibronectin binding partner. For example, a detectable label such as colloidal gold or a dye in a latex microsphere can be conjugated to an anti-oncofetal fibronectin antibody. In another example, a detectable label such as a goat anti-mouse IgG antibody conjugated to horseradish peroxidase can specifically bind to a mouse IgG antibody fibronectin or oncofetal fibronectin binding partner.

[0199] As used herein, bind, bound and binding refer to the binding between atoms or molecules with a K_d in the range of 10^{-2} to 10^{-15} mole/L, generally, 10^{-6} to 10^{-15} , 10^{-7} to 10^{-15} and typically 10^{-8} to 10^{-15} (and/or a K_a of 10^5 - 10^{12} , 10^7 - 10^{12} , 10^8 - 10^{12} L/mole).

[0200] As used herein, complex refers generally to an association between two or more species regardless of the nature of the interaction between the species (i.e., ionic, covalent, or electrostatic).

[0201] As used herein, mass spectrometry encompasses any mass spectrometric format known to those of skill in the art. In particular, the phrases mass spectrometry include time-of-flight, Fourier transform, inductively coupled plasma, ion trap, magnetic sector/magnetic deflection instruments in single or triple quadrupole mode (MS/MS) and also can include combinations thereof. SELDI and MALDI mass spectrometry can be particularly attractive when a time-of-flight (TOF) configuration is used as a mass analyzer. The MALDI-TOF mass spectrometry was introduced by Hillen-

kamp et al., "Matrix Assisted UV-Laser Desorption/Ionization: A New Approach to Mass Spectrometry of Large Biomolecules," *Biological Mass Spectrometry*, (Burlingame and McCloskey, editors), Elsevier Science Publishers, Amsterdam, pp. 49-60 (1990). SELDI-TOF methods are summarized in Merchant et al., *Electrophoresis* 21:1164-1177 (2000).

[0202] As used herein, desorb, desorbed and desorbing refer to the departure of a species from a surface and/or entry of the species into the gaseous phase. In particular, analytes can be desorbed from substrates using any of a variety of techniques, such as, for example, ultraviolet (UV) and infrared (IR) Matrix-Assisted Laser Desorption/Ionization (MALDI; see, e.g., published International PCT Application No. WO 99/57318 and U.S. Pat. No. 5,118,937), field desorption (FD) or fast atom bombardment (FAB). For the MALDI desorption/ionization process, numerous matrix/laser combinations can be used. Additional desorption methods include surface-enhanced neat desorption (SEND; see, e.g., U.S. Pat. No. 5,894,063) and surface-enhanced photo-labile attachment and release (SEPAR; see, e.g., U.S. Pat. No. 6,124,137).

[0203] As used herein, ionization in the context of mass spectrometry refers to methods of creating charged particles in the gaseous phase. Ionization methods include desorption methods provided above such as SELDI, MALDI, FD and FAB. Ionization methods also include non-desorption methods such as electrospray (ES), electron impact (EI) and chemical ionization (CI). For ES, the samples, dissolved in water or in a volatile buffer, are injected either continuously or discontinuously into an atmospheric pressure ionization interface (API). Such ions can be mass analyzed by a quadrupole. The generation of multiple ion peaks which can be obtained using ES mass spectrometry can increase the accuracy of the mass determination.

[0204] As used herein, matrix material refers to any one of several small, photon absorbing compounds that can be mixed in solution with an analyte (e.g., an oncofetal fibronectin indicating molecule) in such a manner so that, upon drying on the mass spectrometry substrate, the matrix-embedded analyte molecules are successfully desorbed and ionized from the solid phase (e.g., crystals) into the gaseous or vapor phase and accelerated as intact molecular ions. For MALDI, sample can be mixed with a prepared solution of the chemical matrix (e.g., at a matrix-to-sample molar ratio of about 10,000:1, or 10,000:1) and placed on the mass spectrometry substrate and dried. Alternatively, a sample can be placed on a mass spectrometry substrate containing matrix and then dried. The large fold excess of matrix, present at concentrations near saturation, facilitates crystal formation and entrapment of analyte.

[0205] As used herein, energy absorbing molecule refers to any one of several small, photon absorbing compounds that, when presented on the surface of a mass spectrometry substrate, facilitate the neat desorption of molecules from the solid phase (i.e., the surface of the probe) into the gaseous or vapor phase for subsequent detection.

[0206] As used herein, substrate when used in the context of mass spectrometry, refers to an insoluble support that can serve as a surface from which a sample is desorbed and ionized in the process of mass measurement of sample components.

[0207] As used herein, a combination refers to any association between two or among more items.

[0208] As used herein, detect, detected and detecting refer generally to any manner of discovering or determining the presence of a signal, such as fluorescence or absorption, or a substance such as an oncofetal fibronectin indicating molecule or a binding partner.

[0209] As used herein, lavage refers generally to a method for obtaining a sample from a biological region or surface by contacting the region or surface with a fluid.

[0210] As used herein, ductal lavage refers generally to a method for obtaining a sample from a biological passageway through which excretions or secretions can pass (e.g., a sample from a milk duct of a breast).

[0211] As used herein, fine needle aspiration refers to a technique whereby a lumen-containing needle is used to obtain a sample. The needle is typically passed through the skin into the tissue to be sampled (e.g., a suspected tumor). A negative pressure in the needle can be formed to draw a small amount of tissue fluid, typically together with loose cells, into the needle. The needle then is removed from the tissue. Fine needle aspiration is described, for example, in U.S. Pat. No. 5,964,735 (Oct. 12, 1999) and U.S. Pat. No. 5,645,537 (Jul. 8, 1997).

[0212] As used herein, a fragmentation compound refers to a compound that can be used to fragment a molecule such as a protein or a nucleic acid molecule. For example a fragmentation compound can be used to fragment an oncofetal fibronectin protein or nucleic acid encoding oncofetal fibronectin or complement thereto. A fragmentation compound can be a protein, peptide, oligonucleotide, or other compound that can fragment molecules, particularly biomolecules, including macromolecules. For example, a fragmentation compound can be a protease or other compound that can be used to fragment an oncofetal fibronectin protein. Exemplary compounds for fragmenting oncofetal fibronectin protein include cathepsin D, trypsin, thermolysin, 2-nitro-5-thiocyanobenzoic acid (for S-cyanylation), *Achromobacter* protease 1, *S. aureus* V8 protease and hydroxylamine. In another example, a fragmentation compound can be nuclease, ribozyme, DNase, or other compound that can be used to fragment an oncofetal fibronectin nucleic acid molecule or complement thereto. Exemplary compounds for fragmenting nucleic acid molecules include restriction endonucleases, exonucleases, hammerhead ribozymes and RNases.

[0213] As used herein, fragment refers to a derivative of a species that is less than the full species. For example, a fragment of an oncofetal fibronectin protein is typically a polypeptide containing fewer than the total amount of amino acids present in a translated fibronectin protein. In another example, a fragment of an oncofetal fibronectin-encoding nucleic acid molecule or complement thereto is typically an oligonucleotide containing fewer nucleic acids than the total amount of nucleic acids present in a transcribed oncofetal fibronectin-encoding nucleic acid molecule.

[0214] As used herein, an immunoassay is defined as any method using a specific or preferential binding of an antigen with a second material (i.e., a binding partner, usually an antibody, antibody fragment or another substance having an antigen binding site) that specifically or preferentially binds

to an epitope of the antigen. The immunoassay methods provided herein include any known to those of skill in the art, including, but not limited to, sandwich, competition, agglutination, or precipitation assays.

[0215] As used herein, antibody refers to an immunoglobulin, whether natural or partially or wholly recombinantly or synthetically produced, including any derivative thereof that retains the specific binding ability of the antibody. Hence, antibody includes any protein having an immunoglobulin binding domain or a binding domain that is homologous or substantially homologous to an immunoglobulin binding domain. For purposes herein, antibody includes antibody fragments, such as Fab fragments, which are composed of a light chain and the variable region of a heavy chain. Antibodies include members of any immunoglobulin class, including IgG, IgM, IgA, IgD and IgE.

[0216] As used herein, a monoclonal antibody refers to an antibody secreted by a transfected or cloned cell such as a hybridoma clone. Each such hybridoma clone is derived from a single B cell and, therefore, all of the antibody molecules are identical. Monoclonal antibodies can be prepared using standard methods known to those with skill in the art (see, e.g., Kohler et al., *Nature* 256:495-497 (1975), Kohler et al., *Eur. J. Immunol.* 6:511-519 (1976) and WO 02/46455). For example, an animal is immunized by standard methods to produce antibody-secreting somatic cells. These cells then are removed from the immunized animal for fusion to myeloma cells.

[0217] Somatic cells with the potential to produce antibodies, particularly B cells, can be used for fusion with a myeloma cell line. These somatic cells can be derived from the lymph nodes, spleens and peripheral blood of primed animals. Specialized myeloma cell lines have been developed from lymphocytic tumors for use in hybridoma-producing fusion procedures (Kohler and Milstein, *Eur. J. Immunol.* 6:511-519 (1976); Shulman et al., *Nature*, 276:269-282 (1978); Volk et al., *J. Virol.*, 42:220-227 (1982)). These cell lines have three useful properties. The first is they facilitate the selection of fused hybridomas from unfused and similarly indefinitely self-propagating myeloma cells by having enzyme deficiencies that render them incapable of growing in selective medium that support the growth of hybridomas. The second is they have the ability to produce antibodies and are incapable of producing endogenous light or heavy immunoglobulin chains. A third property is they efficiently fuse with other cells. Other methods for producing hybridomas and monoclonal antibodies are well known to those of skill in the art.

[0218] As used herein, an antibody fragment refers to any derivative of an antibody that is less than a full length antibody, retaining at least a portion of the full-length antibody's specific binding ability. Examples of antibody fragments include, but are not limited to, Fab, Fab', F(ab)₂, single-chain Fvs (scFv), small immune proteins, Fv, dsFv diabody and Fd fragments. The fragment can include multiple chains linked together, such as by disulfide bridges. An antibody fragment generally contains at least about 50 amino acids and typically at least about 200 amino acids, or at least 50 amino acids and typically at least 200 amino acids.

[0219] As used herein, a Fv antibody fragment is composed of one variable heavy domain (V_H) and one variable light (V_L) domain linked by non-covalent interactions.

[0220] As used herein, a dsFv refers to a Fv with an engineered intermolecular disulfide bond, which stabilizes the V_H-V_L pair.

[0221] As used herein, scFvs refer to antibody fragments that contain a variable light chain domain (V_L) and variable heavy chain domain (V_H) covalently connected by a polypeptide linker in any order. The linker is of a length such that the two variable domains are bridged without substantial interference. Exemplary linkers are (Gly-Ser)_n residues with some Glu or Lys residues dispersed throughout to increase solubility.

[0222] As used herein, diabodies are dimeric scFv; diabodies typically have shorter peptide linkers than scFvs and they preferentially dimerize.

[0223] As used herein, small immune proteins (SIP) are scFv fragments connected to a dimerization domain of an antibody, such as an IgG CH₃ domain. For example and SIP can be formed by connecting scFvs through a short linker to the CH3 domain of the human immunoglobulin 1γ H-chain, or a similar domain such as the CH4 domain of human IgE (see, e.g., Li et al., *Protein Engineering* 10:731-736 (1997) and Borsi et al., *Int. J. Cancer* 102:75-85 (2002)).

[0224] As used herein, a Fab fragment is an antigen-binding antibody fragment containing one variable heavy domain (V_H), one variable light (V_L) domain, one constant heavy domain 1 (C_{H1}) and one constant light (C_L) domain. A Fab fragment can be produced by digestion of an immunoglobulin with papain; a Fab fragment also can be recombinantly produced.

[0225] As used herein, hsFv refers to antibody fragments in which the constant domains normally present in a Fab fragment have been substituted with a heterodimeric coiled-coil domain (see, e.g., Arndt et al. *J. Mol. Biol.* 7:312:221-228 (2001)).

[0226] As used herein, an F(ab)₂ fragment is an antibody fragment containing two variable heavy domains (V_H), two variable light (V_L) domains, two constant heavy domains 1 (C_{H1}) and two constant light (C_L) domains. An F(ab)₂ can be produced by digestion of an immunoglobulin with pepsin at pH 4.0-4.5; an F(ab)₂ fragment also can be recombinantly produced.

[0227] As used herein, humanized antibodies refer to antibodies that are modified to include "human" sequences of amino acids so that administration to a human does not provoke an immune response. Methods for preparation of such antibodies are known. For example, a hybridoma that expresses a monoclonal antibody is altered by recombinant DNA techniques to express an antibody in which the amino acid composition of the non-variable regions is based on human antibodies. Computer programs that identify such regions are known in the art.

[0228] As used herein, autoantibody refers to an antibody produced by a subject that binds to an endogenous antigen of the subject. For example, an autoantibody can be produced in response to presence of a tumor, cancer, or cancerous condition with the subject. Autoantibodies, although produced by the subject in response to an endogenous antigen, can be detected or measured by reaction of the autoantibody with a binding partner, such as a test antigen produced or obtained from a variety of sources including by

recombinant techniques. An anti-fibronectin autoantibody refers to an antibody that specifically binds fibronectin. An anti-oncofetal fibronectin autoantibody refers to an antibody that specifically and preferentially binds to oncofetal fibronectin protein or nucleic acid molecule; that is, the autoantibody specifically binds an oncofetal fibronectin molecule in preference to a non-oncofetal fibronectin molecule.

[0229] As used herein in regard to nucleic acid molecules, amplify, amplified and amplifying refer to methods for increasing the number of copies of a specific nucleic acid molecule, such as a DNA fragment. In particular, amplify, amplified and amplifying include processes wherein a nucleic acid molecule is increased in copy number using techniques such as, for example, cloning, transcription, the polymerase chain reaction (PCR), the ligase chain reaction (LCR) and strand displacement.

[0230] As used herein, an amplified nucleic acid molecule corresponding to an oncofetal fibronectin encoding nucleic acid molecule refers to an amplified nucleic acid molecule formed using any amplification method and using an oncofetal fibronectin encoding nucleic acid molecule as the template nucleic acid molecule. Such an amplified nucleic acid molecule can contain all or a portion of the nucleic acid molecule of an oncofetal fibronectin encoding nucleic acid molecule or all or a portion of the nucleotide molecule complementary to an oncofetal fibronectin encoding nucleic acid molecule. For example, an amplified nucleic acid molecule can contain all or a portion of the nucleic acid molecule encoding the EDA, EDB or IIICS regions of fibronectin. In another example, a complement to an oncofetal fibronectin encoding nucleic acid molecule can contain all or a portion of the nucleic acid molecule complementary to the nucleic acid molecule encoding the EDA, EDB or IIICS regions of fibronectin.

[0231] As used herein, the terms convert, converted and converting refer to processes wherein species are converted using, for example, chemical, physical and/or biological reactions.

[0232] As used herein, risk refers to a predictive process in which the probability of a particular outcome is assessed.

[0233] As used herein, the phrases impending delivery and imminent delivery refer to delivery within a predetermined time frame, such as within about 7, 14, 21, or 28 days, or within 7, 14, 21, or 28 days.

[0234] As used herein, the phrase pre-term delivery refers to delivery that occurs from about 20 weeks gestation to about 37 weeks gestation, or from 20 weeks gestation to 37 weeks gestation. The number of weeks gestation (i.e., gestational age) can be determined using any of a number of conventional methods. For example, the gestational age can be calculated from the first day of the last menstruation.

[0235] As used herein, a support refers to any solid or semisolid or insoluble support to which a molecule of interest, typically a biological molecule or organic molecule or biospecific ligand, is linked or contacted. Typically, a support contains immobilized thereto one or more fibronectin or oncofetal fibronectin binding partners. Support materials include any material that can be used as affinity matrices or supports for chemical and biological molecule syntheses and analyses, such as, but are not limited to:

organic or inorganic polymers, biopolymers, natural and synthetic polymers, including, but not limited to, agarose, cellulose, nitrocellulose, cellulose acetate, other cellulose derivatives, dextran, dextran-derivatives and dextran copolymers, other polysaccharides, gelatin, polyvinyl pyrrolidone, rayon, nylon, polyethylene, polypropylene, polybutylene, polycarbonate, polyesters, polyamides, vinyl polymers, polyvinylalcohols, polyvinylidenedifluoride (PVDF), polystyrene and polystyrene copolymers, polystyrene cross-linked with divinylbenzene or the like, acrylic resins, acrylates and acrylic acids, acrylamides, polyacrylamides, polyacrylamide blends, co-polymers of vinyl and acrylamide, methacrylates, methacrylate derivatives and copolymers, other polymers and co-polymers with various functional groups, rubber, latex, butyl rubber and other synthetic rubbers, silicon, glass (e.g. controlled-pore glass (CPG)), silica gels, ceramics, paper, natural sponges, insoluble protein, surfactants, red blood cells, metals (including metal ions; e.g., steel, gold, silver, aluminum and copper), metalloids, magnetic materials (including Teflon® coated magnetic materials and magnetic beads), Wang resin, Merrifield resin, Sephadex®, Sepharose®, nylon, dextran, chitin, sand, pumice, dendrimers, buckyballs, or other commercially available medium. Exemplary supports include, but are not limited to flat supports such as glass fiber filters, silicon surfaces, glass surfaces, magnetic beads, metal surfaces (steel, gold, silver, aluminum and copper) and plastic materials.

[0236] The support can take any of a variety of forms. For example, the substrate can be formed as plates; whiskers; single crystals; ceramics; self-assembling monolayers; beads or microbeads (e.g., silica gel, controlled pore glass, magnetic, Sephadex/Sepharose, cellulose); flat surfaces or chips (e.g., glass fiber filters, glass surfaces, glass slides, metal surfaces (steel, gold, silver, aluminum, copper and silicon)); capillaries; membranes or microtiter plates (e.g., nylon, polyethylene, polypropylene, polyamide, polyvinylidenedifluoride, or nitrocellulose membranes or microtiter plates); pins or combs; wafers (e.g., silicon wafers); and combinations thereof (e.g., beads placed into pits in flat surfaces).

[0237] When particulate, typically the supports have at least one dimension in the 5-100 μm range or smaller. Such supports, referred collectively herein as beads, are often, but not necessarily, spherical. Such reference, however, does not constrain the geometry of the support, which can be any shape, including random shapes, needles, fibers and elongated. Roughly spherical beads, particularly microspheres that can be used in the liquid phase, can be employed. Beads can include additional components, such as magnetic or para-magnetic particles (e.g., Dynabeads® (Dynal, Oslo, Norway)) for separation using magnets, as long as the additional components do not interfere with the methods and analyses herein.

[0238] As used herein, support particles refer to support materials that are in the form of discrete particles. The particles can have any shape and dimensions, but typically have at least one dimension that is 100 μm or less, 50 μm or less, 10 μm or less, 5 μm or less, 4 μm or less, 3 μm or less, 2 μm or less, 1 μm or less, 900 μm or less, 800 μm or less, 700 μm or less, 600 μm or less, 500 μm or less, 400 μm or less, 300 μm or less, 200 μm or less, 100 μm or less, 50 μm or less, 40 μm or less, 30 μm or less, 20 μm or less,

10 μm or less, 5 μm or less, 4 μm or less, 3 μm or less, 2 μm or less, 1 μm or less, 900 nm or less, 800 nm or less, 700 nm or less, 600 nm or less, 500 nm or less, 400 nm or less, 300 nm or less, 200 nm or less, 100 nm or less, 50 nm or less, 40 nm or less, 30 nm or less, 20 nm or less and 10 nm or less. The particles typically have a size that is 100 nm^3 or less, 50 nm^3 or less, 10 nm^3 or less and 5 nm^3 or less, 4 nm^3 or less, 3 nm^3 or less, 2 nm^3 or less and 1 nm^3 or less, 900 μm^3 or less, 800 μm^3 or less, 700 μm^3 or less, 600 μm^3 or less, 500 μm^3 or less, 400 μm^3 or less, 300 μm^3 or less, 200 μm^3 or less, 100 μm^3 or less, 50 μm^3 or less, 40 μm^3 or less, 30 μm^3 or less, 20 μm^3 or less, 10 μm^3 or less, 5 μm^3 or less, 4 μm^3 or less, 3 μm^3 or less, 2 μm^3 or less, 1 μm^3 or less, 900 nm^3 or less, 800 nm^3 or less, 700 nm^3 or less, 600 nm^3 or less, 500 nm^3 or less, 400 nm^3 or less, 300 nm^3 or less, 200 nm^3 or less, 100 nm^3 or less, 50 nm^3 or less, 40 nm^3 or less, 30 nm^3 or less, 20 nm^3 or less, 10 nm^3 or less, 5 nm^3 and can be on the order of cubic nanometers; typically the particles have a diameter of about 1.5 microns and less than about 15 microns, such as about 4-6 microns, or 1.5 microns and less than 15 microns, such as 4-6 microns. Such particles are collectively called beads.

[0239] As used herein in the context of a test strip, upstream describes a relationship between at least two regions, where a first region that is upstream of a second region is a first region that is contacted by the sample prior to sample contact with the second region. Similarly, downstream describes the relationship between two or more regions where a first region that is downstream of a second region is a first region that is contacted by the sample subsequent to sample contact with the second region.

[0240] As used herein, an epitope present in fibronectin refers to any region present in fibronectin that binds to an antibody or fragment thereof. For example, an epitope present in fibronectin can include an epitope that is present in a non-oncofetal fibronectin, for example, an epitope in the FNIII₇ repeat. A large number of antibodies that bind to epitopes present on fibronectins are known in the art, including antibodies HFN 36.3 and HFN 7.1 (Schoen R C, et al. "Monoclonal antibody against human fibronectin which inhibits cell attachment." *Hybridoma* 1: 99-108, 1982; ATCC Nos. CRL-1605 and CRL-1606, respectively), P3NP/PFn (ATCC No. HB-91), 3E3 (Borsi et al., *FEBS Lett.* 192:71-74 (1985)) and IST-4 (Sekiguchi et al., *J. Biol. Chem.* 260:5105-5114 (1985)); Accurate Chemical and Scientific Corp., Westbury, N.Y.).

[0241] As used herein, an epitope present in oncofetal fibronectin refers to any region present in oncofetal fibronectin that binds to an anti-oncofetal fibronectin antibody or fragment thereof. For example, an epitope present in oncofetal fibronectin can include a splice region specific to an oncofetal fibronectin indicating molecule, for example, an epitope in EDA, EDB or IIICS and also can include other regions in an oncofetal fibronectin indicating molecule to which an anti-oncofetal fibronectin antibody binds by virtue of the presence of one or more of EDA, EDB or IIICS, for example the FNIII₆ repeat when EDB is present. A large number of antibodies that bind to oncofetal fibronectin-specific are known in the art, including IST-9 (Carnemolla et al., *FEBS Lett.* 215:269-273 (1987)); available at Accurate Chemical & Sci. Corp., Westbury, N.Y.), DH1 (Vartio et al., *J. Cell Sci.* 88:419-430 (1987)), BC-1 (Carnemolla et al., *J. Cell Biol.* 108:1139-1148 (1989)), L19 (U.S. Pat. App. No.

20030176663), ME4C (SEQ ID NO: 9) (Giovannoni et al., *Nucleic Acids Res.* 29:e27 (2001)); the ME4C scFv recombinant antibody sequence is provided as SEQ ID No:10 (see, also GenBank accession no. AJ297960), H10 (U.S. Pat. App. No. 20030176663), A134 (Islami et al., *Eur. J. Obstet. Gynecol. Reprod. Biol.*, 97:40-45 (2001)) FDC-6 (U.S. Pat. No. 4,894,326; ATCC HB 9018), 5C10 (Mandel et al., *APMIS* 100:817-826 (1992)) and X18A4, X20C4 and X8E3 (U.S. Pat. No. 5,523,229; ATCC Nos. HB-11587, HB-11589 and HB-11588, respectively).

[0242] As used herein, non-radioactive energy transfer reactions, such as FET (fluorescence energy transfer) assays, FRET (fluorescence resonance energy transfer) assays, fluorescence polarization assays and HTRF (homogeneous time-resolved fluorescence), are homogeneous luminescence assays based on energy transfer between a donor luminescent label and an acceptor label (see, e.g., Cardullo et al. *Proc. Natl. Acad. Sci. U.S.A.* 85:8790-8794 (1988); Pearce et al. *Proc. Natl. Acad. Sci. U.S.A.* 83:8092-8096 (1986); U.S. Pat. No. 4,777,128; U.S. Pat. No. 5,162,508; U.S. Pat. No. 4,927,923; U.S. Pat. No. 5,279,943; and International PCT Application No. WO 92/01225).

[0243] As used herein, Fluorescence Resonance Energy Transfer (FRET) refers to non-radioactive energy transfer between chemical and/or protein fluorophores. Fluorescent resonance energy transfer (FRET) is an art-recognized process in which one fluorophore (the acceptor) can be promoted to an excited electronic state through quantum mechanical coupling with and receipt of energy from an electronically excited second fluorophore (the donor). This transfer of energy results in a decrease in visible fluorescence emission by the donor and an increase in fluorescent energy emission by the acceptor.

[0244] For FRET to occur efficiently, the absorption and emission spectra between the donor and acceptor have to overlap. Dye pairs are characterized by their spectral overlap properties. Emission spectrum of donor must overlap acceptor absorption spectrum. Extent of overlap determines the efficiency of energy transfer. Extent of overlap also determines the optimal distance for which the assay is sensitive. Where the overlap of spectra is large, the transfer is efficient, so it is sensitive to longer distances. The selection of donor/acceptor depends upon the distances considered.

[0245] Significant energy transfer can only occur when the donor and acceptor are sufficiently closely positioned since the efficiency of energy transfer is highly dependent upon the distance between donor and acceptor fluorophores. The fluorophores can be chemical fluorophores and/or protein fluorophores. For example, FRET energy transfer between two fluorescent proteins as a physiological reporter has been reported (Miyawaki et al. *Nature* 388:882-887 (1997)), in which two different fluorescent proteins were fused to the carboxyl and amino termini of calmodulin. Changes in calcium ion concentration caused a sufficient conformational change in calmodulin to alter the level of energy transfer between the fluorescent protein moieties.

[0246] As used herein, fluorescence polarization or fluorescence polarization anisotropy (see, e.g., Jameson et al. *Methods Enzymol.* 246:283-300 (1995)) refers to procedures in which fluorescently labeled molecules are illuminated in solution with plane-polarized light. When fluorescently labeled molecules in solution are so-illuminated, the emitted

fluorescence is in the same plane provided the molecules remain stationary. Since all molecules in solution tumble as a result of collisional motion, depolarization phenomenon is proportional to the rotational relaxation time of the molecule, which is defined by the expression $3\eta V/RT$. At constant viscosity (η) and temperature (T) of the solution, polarization is directly proportional to the molecular volume (V) (R is the universal gas constant). Hence, changes in molecular volume or molecular weight due to binding interactions can be detected as a change in polarization. For example, the binding of a fluorescently labeled ligand to its receptor results in significant changes in measured fluorescence polarization values for the ligand. Measurements can be made in a "mix and measure" mode without physical separation of the bound and free ligands. The polarization measurements are relatively insensitive to fluctuations in fluorescence intensity when working in solutions with moderate optical intensity.

[0247] As used herein, luminescence refers to the detectable EM radiation, generally, UV, IR or visible EM radiation that is produced when the excited product of an exergic chemical process reverts to its ground state with the emission of light. Chemiluminescence is luminescence that results from a chemical reaction. Bioluminescence is chemiluminescence that results from a chemical reaction using biological molecules (or synthetic versions or analogs thereof) as substrates and/or enzymes.

[0248] As used herein, chemiluminescence refers to a chemical reaction in which energy is specifically channeled to a molecule causing it to become electronically excited and subsequently to release a photon thereby emitting visible light. Temperature does not contribute to this channeled energy. Thus, chemiluminescence involves the direct conversion of chemical energy to light energy.

[0249] As used herein, bioluminescence, which is a type of chemiluminescence, refers to the emission of light by biological molecules, particularly proteins. The essential condition for bioluminescence is molecular oxygen, either bound or free in the presence of an oxygenase, a luciferase, which acts on a substrate, a luciferin. Bioluminescence is generated by an enzyme or other protein (luciferase) that is an oxygenase that acts on a substrate luciferin (a bioluminescence substrate) in the presence of molecular oxygen and transforms the substrate to an excited state, which, upon return to a lower energy level releases the energy in the form of light.

[0250] As used herein, the substrates and enzymes for producing bioluminescence are generically referred to as luciferin and luciferase, respectively. When reference is made to a particular species thereof, for clarity, each generic term is used with the name of the organism from which it derives, for example, bacterial luciferin or firefly luciferase.

[0251] As used herein, luciferase refers to oxygenases that catalyze a light emitting reaction. For instance, bacterial luciferases catalyze the oxidation of flavin mononucleotide (FMN) and aliphatic aldehydes, which reaction produces light. Another class of luciferases, found among marine arthropods, catalyzes the oxidation of *Cypridina* (*Vargula*) luciferin and another class of luciferases catalyzes the oxidation of Coleoptera luciferin.

[0252] Thus, luciferase refers to an enzyme or photoprotein that catalyzes a bioluminescent reaction (a reaction that

produces bioluminescence). The luciferases, such as firefly and *Gaussia* and *Renilla* luciferases, are enzymes which act catalytically and are unchanged during the bioluminescence generating reaction. The luciferase photoproteins, such as the aequorin photoprotein to which luciferin is non-covalently bound, are changed, such as by release of the luciferin, during bioluminescence generating reaction. The luciferase is a protein that occurs naturally in an organism or a variant or mutant thereof, such as a variant produced by mutagenesis that has one or more properties, such as thermal stability, that differ from the naturally-occurring protein. Luciferases and modified mutant or variant forms thereof are well known. For purposes herein, reference to luciferase refers to either the photoproteins or luciferases.

[0253] Thus, reference, for example, to *Renilla* luciferase refers to an enzyme isolated from member of the genus *Renilla* or an equivalent molecule obtained from any other source, such as from another related copepod, or that has been prepared synthetically. It is intended to encompass *Renilla* luciferases with conservative amino acid substitutions that do not substantially alter activity. Conservative substitutions of amino acids are known to those of skill in this art and can be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson et al. *Molecular Biology of the Gene*, 4th Edition, 1987, The Benjamin/Cummings Pub. co., p. 224).

[0254] As used herein, bioluminescence substrate refers to the compound that is oxidized in the presence of a luciferase and any necessary activators and generates light. These substrates are referred to as luciferins herein, are substrates that undergo oxidation in a bioluminescence reaction. These bioluminescence substrates include any luciferin or analog thereof or any synthetic compound with which a luciferase interacts to generate light. Typical substrates include those that are oxidized in the presence of a luciferase or protein in a light-generating reaction. Bioluminescence substrates, thus, include those compounds that those of skill in the art recognize as luciferins. Luciferins, for example, include firefly luciferin, *Cypridina* (also known as *Vargula*) luciferin (coelenterazine), bacterial luciferin, as well as synthetic analogs of these substrates or other compounds that are oxidized in the presence of a luciferase in a reaction the produces bioluminescence.

[0255] As used herein, capable of conversion into a bioluminescence substrate refers to being susceptible to chemical reaction, such as oxidation or reduction, that yields a bioluminescence substrate. For example, the luminescence producing reaction of bioluminescent bacteria involves the reduction of a flavin mononucleotide group (FMN) to reduced flavin mononucleotide (FMNH₂) by a flavin reductase enzyme. The reduced flavin mononucleotide (substrate) then reacts with oxygen (an activator) and bacterial luciferase to form an intermediate peroxy flavin that undergoes further reaction, in the presence of a long-chain aldehyde, to generate light. With respect to this reaction, the reduced flavin and the long chain aldehyde are substrates.

[0256] As used herein, a bioluminescence generating system refers to the set of reagents required to conduct a bioluminescent reaction. Thus, the specific luciferase,

luciferin and other substrates, solvents and other reagents that can be required to complete a bioluminescent reaction form a bioluminescence system. Thus a bioluminescence generating system refers to any set of reagents that, under appropriate reaction conditions, yield bioluminescence. Appropriate reaction conditions refers to the conditions necessary for a bioluminescence reaction to occur, such as pH, salt concentrations and temperature. In general, bioluminescence systems include a bioluminescence substrate, luciferin, a luciferase, which includes enzymes luciferases and photoproteins and one or more activators. A specific bioluminescence system can be identified by reference to the specific organism from which the luciferase derives; for example, the *Renilla* bioluminescence system includes a *Renilla* luciferase, such as a luciferase isolated from the *Renilla* or produced using recombinant methods or modifications of these luciferases. This system also includes the particular activators necessary to complete the bioluminescence reaction, such as oxygen and a substrate with which the luciferase reacts in the presence of the oxygen to produce light.

[0257] As used herein, a fluorescent protein (FP) refers to a protein that possesses the ability to fluoresce (i.e., to absorb energy at one wavelength and emit it at another wavelength). For example, a green fluorescent protein (GFP) refers to a polypeptide that has a peak in the emission spectrum at 510 nm or about 510 nm. A variety of FPs that emit at various wavelengths are known in the art.

[0258] As used herein, *Aequora* GFP refers to GFPs from the genus *Aequora* and to mutants or variants thereof. Such variants and GFPs from other species are well known and are available and known to those of skill in the art.

[0259] As used herein, quantitation of an oncofetal fibronectin indicating molecule refers to the calculation of the concentration, mass, or molar quantity of an oncofetal fibronectin indicating molecule in a sample.

[0260] As used herein, a threshold level of an oncofetal fibronectin indicating molecule refers to a level of an oncofetal fibronectin indicating molecule that is compared to a measured amount of an oncofetal fibronectin indicating molecule, where a measured amount above or equal to the threshold level is categorized differently than a measured amount below a threshold level. For example, a measured amount above or equal to a threshold level can be categorized as oncofetal fibronectin positive and a measured amount below the threshold level can be categorized as oncofetal fibronectin negative. In the case of multiple thresholds, the measured amount can be categorized according to the highest level that is less than or equal to the measured amount. One or more thresholds as used herein, refers to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more thresholds. A level of an oncofetal fibronectin indicating molecule, such as a threshold level, can be a reference amount represented as a raw concentration (i.e., not normalized), normalized concentration, mass quantity, molar quantity, or other quantitative amount. For example, the threshold level can be the level of an oncofetal fibronectin indicating molecule that is present in a sample from a population of normal individuals or from the subject at different time points. For example, a subject that is negative for oncofetal fibronectin, as used herein, refers to a subject that does not exhibit oncofetal fibronectin indicating molecule levels significantly above

normal oncofetal fibronectin indicating molecule levels. As is understood by one skilled in the art, the threshold level can vary depending on the tissue or fluid sampled, depending on the sample type, depending on the detection method, depending on the age, gender or biological state (e.g., pregnant or not pregnant) of a subject. In some instances, the threshold level for an oncofetal fibronectin indicating molecule is zero (i.e., when any oncofetal fibronectin indicating molecule is present, the sample is positive for oncofetal fibronectin).

[0261] As used herein, a primer refers to an oligonucleotide to which can be enzymatically added one or more additional nucleotides. Typically a primer contains a free 3' hydroxy moiety.

[0262] As used herein, an amplifiable signaling nucleic acid refers to a nucleic acid that can be amplified using known amplification methods such as polymerase chain reaction (PCR) and the presence of which indicates complex formation between an oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner.

[0263] As used herein, a health problem associated with oncofetal fibronectin refers to an adverse health condition, such as a disease or pregnancy trouble, that is more common in subjects having the presence of an oncofetal fibronectin indicating molecule or an amount at or over a threshold relative to subjects having absence of an oncofetal fibronectin indicating molecule or an amount below a threshold, or that is less common in subjects having an absence of an oncofetal fibronectin indicating molecule or an amount below a threshold relative to subjects having the presence of an oncofetal fibronectin indicating molecule or an amount at or over a threshold. An oncofetal fibronectin-associated health problem is characterized by the presence of, or elevated levels of, an oncofetal fibronectin indicating molecule in a body tissue or fluid sample. The presence of, or elevated levels of, an oncofetal fibronectin indicating molecule does not necessarily indicate that the health problem is caused by an oncofetal fibronectin indicating molecule, but that elevated levels of an oncofetal fibronectin indicating molecule are observed in tissue and/or fluid samples. For example, an oncofetal fibronectin indicating molecule can serve as an indicator of cancer, can serve as an indicator of pre-term or imminent delivery and also can serve as an indicator of a condition such as, but not limited to, arthritis, diabetic retinopathy, renal disease and Dupuytren's contracture. Detection of an oncofetal fibronectin indicating molecule in a body tissue or fluid sample at or above one or more thresholds or at a level above a baseline for a particular individual can be an indicator of a variety of health problems or risk therefor. Similarly, its absence or presence below one or more thresholds can be indicative of the absence of any of these variety of diseases and disorders (i.e., health problems).

[0264] As used herein, progesterational therapy refers to one or more therapeutic methods that favors, or is conducive to, gestation, or inhibits premature labor, or increases the viability of an infant after birth, particularly the viability of a pre-term infant. Progesterational therapy can include methods such as bedrest for the pregnant subject and also can include administration of one or more agents that reduce or inhibit uterine contractions, that prolongs the pregnancy, or that

increases the viability of an infant delivered pre-term. For example, progestational therapy can include administration of a tocolytic agent.

[0265] As used herein, a tocolytic agent refers to a compound or composition that, upon administration to a subject, reduces or inhibits uterine contractions, or otherwise inhibits premature labor.

[0266] As used herein, conceptus refers to any cells, cell masses, and tissues resulting from fertilization of an ova by a sperm from the moment of fertilization through birth. These include, but not limited to, zygotes, embryos, blastocysts, and fetuses.

[0267] As used herein, concepti is the plural of conceptus.

[0268] As used herein, a conceptus sample refers to a sample that contains compounds produced by a conceptus. Conceptus samples include conceptus extracts, samples from outside of the conceptus, such as culture medium, cell and tissue extracts, and cells, where one or more cells is removed from a conceptus, leaving the remainder of the conceptus competent for subsequent culture, implantation and/or development. A conceptus sample can be analyzed neat, or can be reagent-treated and/or fractionated prior to detection of an oncofetal fibronectin indicating molecule.

[0269] As used herein, an additional maternal or conceptus marker refers to a marker that is predetermined to be marker for successful implantation. The additional marker can be detected in a conceptus sample, is determined by visual inspection of the conceptus or is detected in a maternal sample. Any such marker can be employed. Exemplary markers include, but are not limited to, genetic composition of the conceptus, gene expression of the conceptus and morphology of the conceptus. One additional marker can also be, for example morphology of the conceptus, and the morphology of the conceptus is graded according to factors such as cell number, degree of fragmentation, cell regularity, symmetry, pronuclear morphology, follicle size, follicular fluid volume, multi-nucleation, presence of vacuoles, granularity, and combinations thereof.

[0270] As used herein, gametes refer to ova and sperm.

[0271] As used herein, fertilization refers to the fusion of a sperm cell with an ova.

[0272] As used herein, implantation with reference to the uterus, uterine wall, or endometrial layer, refers to the penetration and/or attachment of a conceptus (or concepti) and/or cells of a conceptus (or concepti) into or onto such cells and/or tissues.

[0273] As used herein, trophoblast refers to the outer layer of epithelial cells surrounding the inner cell mass of a blastocyst. The trophoblast also is referred to as the outer cell mass. Trophoblast cells can develop into extra-embryonic cells and tissues including placenta, amnion and umbilical cord.

[0274] As used herein a non-specific binder, or a substance that reduces non-specific binding, is a substance that binds to at least a portion of background material in a sample without binding more than a small amount, typically less than or less than about 1%, 2%, 5% or 10% depending upon the assay or application, of oncofetal fibronectin indicating molecule in the sample. A non-specific binder can include,

for example, a non-specific binding compound, or a solid support containing a non-specific binding surface.

[0275] As used herein, a non-specific binding compound can bind to at least a portion of background material in a sample without binding more than a small amount (e.g., less than 10%) of oncofetal fibronectin indicating molecule in the sample. A non-specific binding compound can be in any of a variety of forms, including, but not limited to, soluble in solution, mobile in a solvent, present in an emulsion, present in a gel, present on a solid support (including, e.g., immobilized on a solid support). Exemplary, non-limiting, non-specific binding compounds that can be used include non-specific binding proteins, including albumins such as bovine serum albumin (BSA), human, rabbit, goat, sheep and horse serum albumins; and other proteins such as ovalbumin, fibrinogen, thrombin, transferrin, glycoproteins, casein, antibodies not specific for an oncofetal fibronectin indicating molecule and other proteins. Non-specific binding proteins also can include water-soluble polyamino acids such as, for example, polymers of one or more amino acids such as lysine, glutamic acid, alanine, histidine, methionine and proline. Non-specific binding compounds also can be protein-containing compositions including serum such as fetal calf serum, gelatin and dried milk.

[0276] Non-specific binders can include non-specific binding surfaces, which are solid structures that can contain one or more components, where the non-specific binding surface binds to at least a portion of background material in a sample while not binding more than a small amount (e.g., less than 10%) of oncofetal fibronectin indicating molecule in the sample. Exemplary solid supports for non-specific binding surfaces include paper and cellulose derivatives, such as cellulose esters and ethers, natural and synthetic polymeric materials, such as latex, vinyl polymers, polypropylene, polyethylene and partially hydrolyzed derivatives, polycondensates, copolymers and inorganic materials. For example, a non-specific binding surface is a porous or bibulous member capable of transporting a liquid sample along a test strip. Non-specific binding surfaces can have immobilized thereon one or more non-specific binding compounds such as, but not limited to, albumin (including bovine serum albumin, or BSA), antibodies not specific for an oncofetal fibronectin indicating molecule and others provided herein or known in the art.

[0277] As used herein, therapeutic agents are agents that ameliorate the symptoms of a disease or disorder or ameliorate the disease or disorder. Therapeutic agents include, but are not limited to, moieties that inhibit cell growth or promote cell death, that can be activated to inhibit cell growth or promote cell death, or that activate another agent to inhibit cell growth or promote cell death. Optionally, the therapeutic agent can exhibit or manifest additional properties, such as, properties that permit its use as an imaging agent, as described elsewhere herein. Therapeutic agents include, but are not limited to, for example, cytokines and growth factors, photosensitizing agents, toxins, anticancer antibiotics, a chemotherapeutic compound, a radionuclide,

an angiogenesis inhibitor, a signaling modulator, a bioluminescent compound or a combination thereof.

[0278] Cytokines and growth factors include, but are not limited to, interleukins, such as, for example, interleukin-1, interleukin-2, interleukin-6 and interleukin-12, tumor necrosis factors, such as tumor necrosis factor alpha (TNF- α), interferons such as interferon gamma (IFN- γ), granulocyte macrophage colony stimulating factors (GM-CSF), angiogenins, and tissue factors.

[0279] Exemplary photosensitizing agents include, but are not limited to, for example, indocyanine green, toluidine blue, aminolevulinic acid, texaphyrins, benzoporphyrins, phenothiazines, phthalocyanines, porphyrins such as sodium porfimer, chlorins such as tetra(m-hydroxyphenyl)chlorin or tin(IV) chlorin e6, purpurins such as tin ethyl etiopurpurin, purpurinimides, bacteriochlorins, pheophorbides, pyropheophorbides or cationic dyes.

[0280] Anti-cancer agents, include, but are not limited to, for example, porfiromycin, doxorubicin, dactinomycin, pliamycin, mitomycin, bleomycin, actinomycin, or daunorubicin and chemotherapeutic compounds.

[0281] Radionuclides, which depending up the radionuclide, amount and application can be used for diagnosis and/or for treatment. They include, but are not limited to, for example, a compound or molecule containing ³²Phosphate, ⁶⁰Cobalt, ⁹⁰Yttrium, ⁹⁹Technicium, ¹⁰³ Palladium, ¹⁰⁶Ruthenium, ¹¹¹Indium, ¹¹⁷Lutetium, ¹²⁵Iodine, ¹³¹Iodine, ¹³⁷Cesium, ¹⁵³Samarium, ¹⁸⁶Rhenium, ¹⁸⁸Rhenium, ¹⁹²Iridium, ¹⁹⁸Gold, ²¹¹Astatine, ²¹²Bismuth or ²¹³Bismuth. Toxins include, but are not limited to, chemotherapeutic compounds such as, but not limited to, 5-fluorouridine, calicheamicin and maytansine. Signaling modulators include, but are not limited to, for example, inhibitors of macrophage inhibitory factor, toll-like receptor agonists and stat 3 inhibitors.

[0282] Chemotherapeutic compounds include, but are not limited to, alkylating agents such as thiotepa and cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramidate and trimethylolomelamine nitrogen mustards such as chlorambucil, chlormaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carubicin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitab-

ine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; polysaccharide-K; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; cytosine arabinoside; cyclophosphamide; thiotepa; taxoids, e.g. paclitaxel and doxetaxel; chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone and toremifene (Fareston); and antiandrogens such as flutamide, nilutamide, bicalutamide, leuprolide and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Such chemotherapeutic compounds that can be used herein include compounds whose toxicities preclude use of the compound in general systemic chemotherapeutic methods.

[0283] As used herein, a pre-induction agent or procedure refers to an agent or process, such as cervical and vaginal ripening, prostaglandin, a foley catheter, or dinoprostone agents such as Prepidil and Cervidil used to prepare a pregnant subject for induction of labor.

[0284] As used herein, an induction agent refers to an agent in is administered to causes labor to begin.

[0285] As used herein, an induction procedure refers any procedure that is used to induce labor. The procedures include, but are not limited to, balloon catheterization such as foley balloon catheterization or Atad balloon catheterization, amniotic membrane stripping, extra-amniotic saline infusion, amniotomy and/or nipple stimulation, and administration of an induction agent.

[0286] As used herein, an induction agent, refers to an agent that induces labor, and includes, for example, oxytocin. Oxytocin plays a role in the initiation of labor, stimulates the contraction of smooth muscle of the uterus during labor and facilitates ejection of milk from the breast during nursing.

[0287] As used herein, a parturifacient refers to any of a variety of compounds or compositions known in the art for pre-inducement, cervical ripening, or inducement. Exem-

plary parturifacients include, but are not limited to, prostaglandins such as PGE1 (misoprostol) and PGE2 (dinoprostone), oxytocic hormones such as oxytocin and steroids such as RU486 (mifepristone).

[0288] As used herein, successful induction is an induction that, for example, results in vaginal delivery, or a shorter time to delivery, or fewer administrations of induction or pre-induction agents compared to in the absence of induction. The likelihood of a successful induction refers to a subject who has been induced and the likelihood that induction will be successful. In this context, a positive test for oncofetal fibronectin is correlated with successful induction, which can be manifested by exhibiting a mean time interval between a first dose of pre-induction agent and delivery that is shorter than the mean time interval between first dose of parturifacient and delivery. A positive test can be measured relative to a threshold amount or compared to similar subjects. A negative test for oncofetal fibronectin is correlated with an observation of a mean time interval between first dose of pre-induction agent and delivery that is longer than the mean time interval between first dose of parturifacient and delivery for subjects testing positive for oncofetal fibronectin or a sample having an amount of oncofetal fibronectin indicating molecule above a threshold.

[0289] As used herein, a second indicator of induction outcome refers to measurements or observations of the pregnant subject, a measurement or observation of the fetus(es), and/or a medical history of the pregnant subject. Second indicators include, but are not limited to, for example, cervical length, Bishop score, effacement, parity, cervical dilation, gestational age, body mass index, station, consistency, transvaginal ultrasound, and/or digital examination.

[0290] As used herein, the stage and grading of bladder cancer is in accordance with the Union Internationale Centre le Cancer (UICC) staging from 1997. Tis: carcinoma in situ (CIS), Ta: papillary, Stage 1 (T1) occurs when papillary invade lamina propria, stage 2 (T2a) occurs when the papillary invade superficial muscle, stage 3 (T2b) occurs when the papillary invade deep muscle, stage 3 occurs when there is microscopic invasion of perivesical tissue (T3a) or gross invasions of perivesical tissue (T3b), and stage 4 occurs when there is invasion pelvic organs (prostate, uterus, vagina; T4a) or the pelvic wall or abdominal wall (T4b). The N stage (status of lymph nodes) and M stage (metastatic sites) also are described. Transitional cell carcinoma can be divided into grade 1 (well-differentiated), grade 2 (moderately differentiated) and grade 3 (poorly differentiated).

[0291] For clarity of disclosure and not by way of limitation, the detailed description is divided into the subsections that follow.

B. Detection of Oncofetal Fibronectin

[0292] Provided herein are methods of detecting oncofetal fibronectin indicating molecules. Oncofetal fibronectin indicating molecules can serve as a marker for health state such as general health, cancer, pregnancy and delivery. Detection of an oncofetal fibronectin indicating molecule by the methods provided herein can improve the accuracy, speed and/or convenience of oncofetal fibronectin indicating molecule detection. The detection methods herein also can provide further information, such as the amount or level of an

oncofetal fibronectin indicating molecule present in the sample, the domains present in the detected oncofetal fibronectin indicating molecule and post-translational modifications of a detected oncofetal fibronectin protein.

[0293] In some embodiments, detection methods can be performed by contacting an oncofetal fibronectin indicating molecule with one or more binding partners and detecting complex formation between the oncofetal fibronectin indicating molecule and one or more binding partners. The detection methods can be performed in any of a variety of ways, for example either the sample components or a fibronectin or oncofetal fibronectin binding partner can be immobilized on a solid support; or neither can be immobilized. In another example, detection can be performed in vivo, for example, in an in vivo diagnostic method. In vivo methods also can be used in treatment of a health problem associated with oncofetal fibronectin.

[0294] Presence of oncofetal fibronectin can be determined by detecting an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, nucleic acid molecules encoding oncofetal fibronectin or complements thereto, or autoantibodies specific for oncofetal fibronectin protein or nucleic acid molecules encoding oncofetal fibronectin, or a fragment thereof. Any of a variety of protein, nucleic acid molecule and antibody detection methods can be used to detect an oncofetal fibronectin indicating molecule. Exemplary detection methods include RT-PCR for detecting mRNA encoding oncofetal fibronectin or fragments thereof and mass spectrometry for detecting oncofetal fibronectin proteins or fragments thereof. In addition, methods such as immunoassays are provided herein for the detection of autoantibodies to oncofetal fibronectin.

[0295] In some embodiments, an oncofetal fibronectin indicating molecule can be detected such that the presence or absence of portions of the oncofetal fibronectin indicating molecule can be identified. As described herein, an oncofetal fibronectin indicating molecule can contain one or more of the domains EDA, EDB and IIICS. The methods provided herein can be used to determine the presence or absence of EDA, EDB and/or IIICS in an oncofetal fibronectin indicating molecule. Such an identification can be used, for example, to identify the tissue source of the oncofetal fibronectin indicating molecule.

[0296] Detection of an oncofetal fibronectin indicating molecule can be used to determine whether or not an oncofetal fibronectin indicating molecule is present in a sample, or can be used to measure the amount of an oncofetal fibronectin indicating molecule present in a sample. When the amount is measured, the measured amount can be compared to one or more threshold levels. When a single threshold level is used, an measured amount of an oncofetal fibronectin indicating molecule above the threshold can indicate, for example, overall health state, imminent or pre-term delivery, delivery date, or a health problem such as a cancerous condition or arthritis. When two or more threshold levels are used, the amount of oncofetal fibronectin indicating molecule measured can be used to classify the subject that provided the sample according to the highest threshold value less than or equal to the measured oncofetal fibronectin indicating molecule amount, where such classification can indicate, for example, different overall health state, different expected outcomes of preg-

nancy and delivery, different accuracies of delivery date prediction, or different severities of a health problem such as a cancerous condition or arthritis.

C. Structure and Properties of Fibronectins

[0297] Methods are provided herein that include detection of oncofetal fibronectin proteins, oncofetal fibronectin-encoding nucleic acids or complements thereto, autoantibodies to oncofetal fibronectin and fragments thereof. Hence, knowledge of the structure and properties and identity of oncofetal fibronectin proteins or nucleic acids encoding the proteins can aid in practice of the methods herein. For example, knowledge of proteolytic fragments of the molecule can aid in mass spectrometric detection as can knowledge of glycosylation patterns. Knowledge of the sequences of nucleic acid encoding oncofetal fibronectin molecules and/or domains thereof can aid in methods requiring specific amplification. Knowledge of oncofetal fibronectin domains and molecules that specifically and preferentially bind to oncofetal fibronectin domains can aid in design of apparatuses for detecting oncofetal fibronectin, in methods such as reflectance methods for detecting oncofetal fibronectin and in methods for characterizing oncofetal fibronectin.

[0298] Fibronectin (FN) is one of a largest multi-domain proteins (Pankov et al., *Journal of Cell Science*, 115:3861-3863, (2002)). Fibronectin (FN) mediates a variety of cellular interactions with the extracellular matrix (ECM) and is important for cell adhesion, migration, growth and differentiation (Mosher, D. F., "Fibronectin," San Diego: Academic Press, Inc., (1989); Carsons, S. E., "Fibronectin in Health and Disease," Florida: CRC Press, Inc., (1989); Hynes, R. O., "Fibronectins," New York: Springer-Verlag, (1990); Yamada and Clark, "The Molecular and Cellular Biology of Wound Repair," (ed. R. A. F. Clark) pp 51-93, New York: Plenum Press, (1996). FN is expressed by multiple cell types and is important in vertebrate development, as demonstrated by the early embryonic lethality of mice with targeted inactivation of the FN gene (George et al., "Defects in mesoderm, neural tube and vascular development in mouse embryos lacking fibronectin," *Development*, 119:1079-1091, (1993).

[0299] 1. Structural Characteristics of Fibronectin

[0300] A variety of features of the fibronectin structure are known; a summary of such features is available at (Pankov et al., *Journal of Cell Science*, 115:3861-3863, (2002)) and summarized herein. Fibronectin usually exists as a dimer containing two nearly identical approximately 250 kDa subunits linked covalently near their C-termini by a pair of disulfide bonds. Each monomer includes three types of repeating units (termed FN repeats): type I, type II and type III. Fibronectin contains 12 type I repeats, two type II repeats and 15-17 type III repeats, which together account for approximately 90% of the fibronectin sequence. Type I repeats are about 40 amino-acid residues in length and contain two disulfide bonds; type II repeats contain a stretch of approximately 60 amino acids and two intra-chain disulfide bonds; and type III repeats are about 90 residues long without any disulfide bonds.

[0301] Fibronectin is encoded by a single gene, the product of which can exist in multiple forms resulting from alternative splicing of a single pre-mRNA that can generate as many as 20 variants in human fibronectin (see, e.g.,

French-Constant, C., "Alternative splicing of fibronectin—many different proteins but few different functions," *Exp. Cell Res.*, 221:261-271, (1995); Kosmehl et al., "Molecular variants of fibronectin and laminin: structure, physiological occurrence and histopathological aspects," *Virchows Arch*, 429:311-322, (1996)). Splicing occurs within the central set of type III repeats, FN III7 to FN III15. Exon usage or skipping leads to inclusion or exclusion of two type III repeats—EDB (also termed EIIIB or EDII and located between FN repeats III7 and III8) and EDA (also called EIIIA or EDI and located between FN repeats III11 and III12), or both. This splicing of fibronectin ED domains is found in many vertebrates, including *Xenopus*, chickens, rats and humans.

[0302] A third region of alternative splicing is localized to a portion of fibronectin not homologous to FNIII repeats, called the V (variable in length) or IIIICS (type III connecting segment) region. The structural variations in this region can include splice variants and are species dependent. In most species, this region can be either partially or completely included or excluded; for example, in human fibronectin, there can be five different V region variants. In chicken, the whole 120 amino acid residues of the V region can be included or a 44 amino acid segment from the 5' end can be excluded (creating V76), but the whole V region of chicken fibronectin is never missing. Splicing in rat leads to exclusion of a 25 amino acid fragment, generating V95 that can be detected, as can be the V0 and V120 forms. Splicing of the V region in human can include combinations of three different regions, the first containing 25 amino acids, the second containing 64 amino acids and the third containing 31 amino acids. Differential splicing in humans leads to at least five variants where segments from the 5' (25aa) and 3' (31aa) ends can be omitted independently (creating V95 and V89 correspondingly) or together (V64), or can both be present (V120), or all three regions can be absent (V0), producing five different V splice variants.

[0303] FNs are glycoproteins that contain 4-9% carbohydrate, depending on the cell source. N-linked and O-linked glycosylation sites are located mainly in type III repeats and the collagen-binding domain.

[0304] Some forms of fibronectin are abundant and soluble in plasma (300 $\mu\text{g}/\text{ml}$) and other body fluids and also part of the insoluble extracellular matrix.

[0305] 2. Binding Properties and Proteolysis of Fibronectin

[0306] FN can be a ligand for numerous members of the integrin receptor family (see, e.g., Plow et al., "Ligand binding to integrins," *J. Biol. Chem.*, 275:21785-21788, (2000)). Integrins are structurally and functionally related cell-surface heterodimeric receptors that link the ECM with the intracellular cytoskeleton. A number of different integrins bind to FN, including the FN receptor $\alpha_5\beta_1$. Several integrin-recognition sequences are known. For example, integrin $\alpha_5\beta_1$ is recognized by the RGD sequence located in FN repeat III10. The recognition of this simple tripeptide sequence can be influenced by flanking residues, the tripeptide's three-dimensional presentation and individual features of the integrin-binding pockets. A second site, in FN repeat III9 (the 'synergy site' PHSRN), promotes specific $\alpha_5\beta_1$ integrin binding to FN via interactions with the α_5 subunit. The FN receptor $\alpha_5\beta_1$ also can interact with an

N-terminal fragment containing repeats I1-9 and III,2, which also promotes $\alpha_5\beta_1$ -integrin-mediated cell adhesion. Interaction with this N-terminal region triggers integrin-mediated intracellular signals that are distinct from those generated in response to ligation with the RGD sequence.

[0307] A second set of fibronectin sequences, which are bound by the $\alpha_4\beta_1$ integrin, also are known. Two such cell-recognition sequences (LDV and REDV) are present in the alternatively spliced V region. Both are recognized by $\alpha_4\beta_1$ and $\alpha_4\beta_7$. Additional sites recognized by the $\alpha_4\beta_1$ integrin—IDAPS and KLDAPT—also are present in repeats III14 and III5, respectively (KLDAPT also binds to the $\alpha_4\beta_7$ integrin). The EDGIHEL sequence of EDA can bind to $\alpha_4\beta_1$ as well as $\alpha_9\beta_1$ (Liao et al., "The EIIIA segment of fibronectin is a ligand for integrins $\alpha_9\beta_1$ and $\alpha_4\beta_1$ providing a novel mechanism for regulating cell adhesion by alternative splicing," *J. Biol. Chem.*, 277:14467-14474, (2002)).

[0308] Fibronectins can be cleaved at known locations when subjected to limited proteolytic digestion (reviewed by Mosher, D. F., "Fibronectin," San Diego: Academic Press, Inc., (1989); Hynes, R. O., "Fibronectin," New York: Springer-Verlag, (1990)). Even a protease capable of cleaving proteins at many sites (such as pronase) initially cleaves FN at highly specific locations. A simplified scheme of major proteolytic cleavage sites is shown in FIG. 1. The binding activities of FN can be preserved after such proteolysis and identified within particular fragments.

[0309] Fibronectin has a variety of functional activities besides binding to cell surfaces through integrins. It also can bind to biologically important molecules such as heparin, collagen/gelatin and fibrin. These interactions are mediated by several distinct structural and functional domains, which have been defined by proteolytic fragmentation or recombinant DNA analyses (see FIG. 1 and Mosher, D. F., "Fibronectin," San Diego: Academic Press, Inc., (1989); Hynes, R. O., "Fibronectins," New York: Springer-Verlag, (1990); and Yamada and Clark, "The Molecular and Cellular Biology of Wound Repair," (ed. R. A. F. Clark), pp. 51-93, New York: Plenum Press, (1996)).

[0310] Fibronectin contains two major heparin-binding domains that interact with heparin sulfate proteoglycans. A strong heparin-binding site is located in the C-terminal part (Heparin II) and a weaker binding domain is situated at the N-terminal end of the protein (Heparin I). The high-affinity heparin II domain also can bind to a widely distributed glycosaminoglycan, chondroitin sulfate; the weaker heparin-binding domain contains a *Staphylococcus aureus*-binding site that mediates FN interactions with bacteria. A glycosaminoglycan-binding site is located within the V region of fibronectin (Mostafavi-Pour et al., 2001) (marked as Heparin at the V domain). In at least some cell types, the heparin-binding domains of fibronectin mediate cell adhesion.

[0311] The collagen-binding domain includes repeats I6-9 and III,2 and these repeats bind more effectively to denatured collagen (gelatin) than to native collagen. This fibronectin domain also can interact with native collagen *in vivo*.

[0312] Fibronectin also contains two major fibrin-binding sites (Fibrin I and Fibrin II). The Fibrin I binding site is in the N-terminal domain and is formed by type I repeats 4 and

5. The interaction of fibronectin with fibrin is involved in cell adhesion or cell migration into fibrin clots. In both instances, cross-linking between fibronectin and fibrin is mediated by factor XIII transglutaminase (the cross-linking site on the fibronectin molecule of FIG. 1 is marked by factor XIIIa and an arrow).

[0313] Fibronectin also can self-associate into aggregates and fibrils, at multiple binding sites that have been identified along the molecule (FIG. 1). Some of these self-interaction sites are exposed and available for binding, while others are cryptic and become accessible only after conformational changes, for example, mechanical stretching of the fibronectin molecule.

[0314] 3. Oncofetal Fibronectins

[0315] Oncofetal fibronectins constitute a heterogeneous group of fibronectin proteins that share certain characteristics. As noted oncofetal fibronectin proteins contain extra-domain A (EDA), extra-domain B (EDB), or fibronectin III connecting segment (IIICS), or any combination thereof. They also are expressed in or shed from certain cells or tissues, and their pattern of expression or shedding or level of expression can vary in tissues. The group of oncofetal fibronectins results from alternative splicing of these three regions (EDA, EDB and IIICS) in fibronectin and also from post-translational modifications. The splicing and expression of oncofetal fibronectin is differently regulated in cells and tissues and at different developmental stages. In fetal tissues and in some abnormal cells and tissues, expression of oncofetal fibronectin is increased relative to the corresponding normal adult cells and tissues. In some normal adult cells, tissues and sample types, oncofetal fibronectin is not present in amounts detectable by antibody assay. Accordingly, for purposes of detection of oncofetal fibronectin, abnormal levels of oncofetal fibronectin can be determined by comparing the detected amount to a control or a to predetermined amount. The amino acid sequence of human fibronectin including EDA, EDB and IIICS and the fibronectin encoding nucleic acid molecule are known in the art and are available in public databases. Exemplary sequences of human oncofetal fibronectins, and EDA, EDB and IIICS regions are set forth in SEQ ID Nos. 4, 6 and 8. Exemplary sequences of human oncofetal fibronectin IIICS regions V120, V95, V89 and V64 are set forth in SEQ ID Nos. 29, 31, 33 and 35, respectively. The protein and encoding nucleic acid molecules from a variety of additional species including, for example, rat, mouse, chicken, cow and *Xenopus laevis* also are known and readily available in public databases. An example of oncofetal fibronectin is a protein that binds specifically to the FDC-6 monoclonal antibody (see, Matsuura and S. Hakomori, *Proc. Natl. Acad. Sci. USA*, 82:6517-6521 (1985). Production of the hybridoma (deposited at the American Type Culture Collection as accession number ATCC HB 9018) which produces FDC-6 antibody is described in detail in U.S. Pat. No. 4,894,326, issued Jan. 16, 1990, to Matsuura et al. Another example of oncofetal fibronectin is a protein that binds preferentially with the BC-1 monoclonal antibody described by Carnemolla et al., *J. Cell. Biol.*, 108:1139-1148 (1989). Another example of oncofetal fibronectin is a protein that binds preferentially with the IST-9 monoclonal antibody described by Carnemolla et al., *FEBS Lett.*, 215:269-273 (1987).

[0316] a. Structural Features of Oncofetal Fibronectin

[0317] Oncofetal fibronectin contains extra-domain A (EDA), extra-domain B (EDB), or fibronectin III connecting segment (IIICS), or any combination thereof. The amino acid sequence of human fibronectin including EDA, EDB and IIICS and the fibronectin encoding nucleic acid molecule are known in the art and are available in public databases and the nucleotide and amino acid sequences also are provided herein as SEQ ID NOS: 1 and 2, 14 and 15, 16 and 17, 18 and 19, 20 and 21, 22 and 23, and 24 and 25, respectively. Nucleic acid molecules encoding a variety of oncofetal fibronectins from other species including, but not limited to, rat, mouse, chicken, cow and *Xenopus laevis* also are known and readily available in public databases.

[0318] Human fibronectin is encoded by the nucleotide and amino acid sequences of SEQ ID NOS: 1 and 2, respectively. EDA is encoded by nucleotides 4405 to 4674 of SEQ ID NO: 1 (SEQ ID NO: 3) and amino acids 1432 to 1621 of SEQ ID NO: 2 (SEQ ID NO: 4). EDB is encoded by nucleotides 3037 to 3309 of SEQ ID NO: 1 (SEQ ID NO: 5) and amino acid 963 to 1109 of SEQ ID NO: 2 (SEQ ID NO: 6). Full length IIICS is encoded by nucleotides 5488 to 5847 of SEQ ID NO: 1 (SEQ ID NO: 7) and amino acid 1830 to 1949 of SEQ ID NO: 2 (SEQ ID NO: 8).

[0319] IIICS can contain various combinations of splice regions resulting in five different splice variants (see Table 1). Amino acid positions 1-25 of IIICS make up splice region A (A). Amino acid positions 26-89 of IIICS make up splice region B (B). Amino acid positions 90-120 of IIICS make up splice region C (C). IIICS, which also is termed the variable or V domain, can be any of at least five different splice variants, including V0 which contains 0 amino acids of IIICS, V64 which contains amino acids 26-89 of IIICS (D; SEQ ID NO: 35), V89 which contains amino acids 1-89 of IIICS (E; SEQ ID NO: 33), V95 which contains amino acids 26-120 of IIICS (F; SEQ ID NO: 31) or V120 which contains amino acids 1-120 of IIICS (G; SEQ ID NO: 29), (see, e.g., Pankov et al., *J. Cell Science* 115:3861-3863 (2002)).

[0320] Portions of IIICS also can be represented as CS1 which contains amino acids 1-25 of IIICS (H), CS2 which contains amino acids 23-47 of IIICS (I), CS3 which contains amino acids 45-68 (J), CS4 which contains amino acids 66-92 of IIICS (K), CS5 which contains amino acids 90-109 of IIICS (L) and CS6 which contains amino acids 107-120 of IIICS (M) (see Table 1). IIICS can be glycosylated at one or more sites. One site for glycosylation is threonine 33 of IIICS, which is O-glycosylated (see Table 1, N). The enzyme that glycosylates threonine 33 is N-acetylgalactosaminyltransferase-T2. The genomic, mRNA, and predicted amino acid sequences of N-acetylgalactosaminyltransferase-T2 are provided in SEQ ID Nos: 11, 12 and 13, respectively, and also are available at GenBank Accession Numbers Y10345 (genomic sequence), X92689 (mRNA sequence), or CAA63371 (amino acid sequence predicted from mRNA sequence); see Wandall et al., *J. Biol. Chem.* 272: 23503-23514 (1997)).

TABLE 1

Splice regions and variable domains of IIICS			
Reference letter	Region of IIICS	Nucleotides (See Seq ID NO: 7)	Amino Acids (See Seq ID NO: 8)
A	Splice region A	1-75	1-25
B	Splice region B	76-267	26-89
C	Splice region C	268-360	90-120
D	V64	76-360	26-89
E	V89	1-267	1-89
F	V95	76-360	26-120
G	V120	1-360	1-120
H	CS1	1-75	1-25
I	CS2	67-141	23-47
J	CS3	133-204	45-68
K	CS4	196-276	66-92
L	CS5	268-327	90-109
M	CS6	319-360	107-120
N	O-glycosylation	97-99	33

[0321] b. Molecules that Bind to Oncofetal Fibronectin

[0322] Oncofetal fibronectin can be specifically bound by one or more anti-oncofetal fibronectin antibodies. A variety of anti-oncofetal fibronectin antibodies are known in the art, including IST-9 (Carnemolla et al., *FEBS Lett.* 215:269-273 (1987)); available at Accurate Chemical & Sci. Corp., Westbury, N.Y.), DH1 (Vartio et al., *J. Cell Sci.* 88:419-430 (1987)), BC-1 (Carnemolla et al., *J. Cell Biol.* 108:1139-1148 (1989)), L19 (U.S. Pat. App. No. 20030176663), ME4C (Giovannoni et al., *Nucleic Acids Res.* 29:e27 (2001)); the ME4C scFv recombinant antibody sequence is available at GenBank accession no. AJ297960), H10 (U.S. Pat. App. No. 20030176663), FDC-6 (U.S. Pat. No. 4,894,326), 5C10 (Mandel et al., *APMIS*, 100:817-826 (1992)) and X18A4, X20C4 and X8E3 (U.S. Pat. No. 5,523,229, ATCC Nos. HB-11587, HB-11589 and HB-11588, respectively).

[0323] IST-9 and DH1 can bind to oncofetal fibronectin when EDA is present. IST-9 and DH1 can bind to at least amino acids Ile-43 and His-44 of EDA. IST-9 and DH1 can bind to the region in EDA containing amino acids 31-44.

[0324] BC-1, L19, ME4C, H10, C6 and A134 can bind to oncofetal fibronectin when EDB is present. BC-1 can bind to oncofetal fibronectin when fibronectin repeat III-7 (FNIII-7) and EDB are present. C6 can bind to oncofetal fibronectin when EDB and fibronectin repeat III-8 (FNIII-8) are present. L19 can bind to EDB.

[0325] FDC-6 can bind to IIICS. FDC-6 can bind to the hexapeptide VTHPGY of IIICS (IIICS amino acids 32-37; SEQ ID NO: 39) when the hexapeptide is O-glycosylated at Thr-33. Typically, the glycosylation of Thr-33 contains an α -N-acetylgalactosamine bonded to the oxygen atom of the threonine side chain. The Thr-33 glycosyl moiety can be NeuAc α 2->3Gal β 1->3GalNac, or 3Gal β 1->3GalNac. 5C10 can bind to a IIICS sequence that overlaps with the FDC-6 hexapeptide. X18A4 can bind to a IIICS sequence different than the hexapeptide bound by FDC-6.

[0326] EDB can contain one or more N-linked glycosylation sites. IIICS can contain one or more O-linked glycosylation sites and from 1 to 6 or about 6 N-linked glycosylation sites.

[0327] EDA can bind to $\alpha_4\beta_1$ integrin and $\alpha_5\beta_1$ integrin. The amino acid sequence EDGIHEL of EDA (EDA amino

acids 40-46) can bind to $\alpha_4\beta_1$ integrin and $\alpha_9\beta_1$ integrin. IIICS can bind to $\alpha_4\beta_1$ integrin, $\alpha_4\beta_7$ integrin and heparin. The V95 splice variant of IIICS can bind to heparin. CS1 and CS5 of IIICS can bind to $\alpha_4\beta_1$ integrin and $\alpha_4\beta_7$ integrin. The IIICS amino acid sequence LDV (IIICS amino acids 20-22) can bind to $\alpha_4\beta_1$ integrin and $\alpha_4\beta_7$ integrin. The IIICS amino acid sequence REDV (IIICS amino acids 100-103) can bind to $\alpha_4\beta_1$ integrin and $\alpha_4\beta_7$ integrin.

[0328] c. Proteolysis of Oncofetal Fibronectin

[0329] Oncofetal fibronectin can be detected according to a variety of properties of oncofetal fibronectin. One method for identifying oncofetal fibronectin is by characteristic proteolysis patterns, including protein fragments with characteristic masses and/or binding properties. Masses of fragments can be measured by any of a variety of methods known in the art or provided elsewhere herein; an exemplary mass measurement method is mass spectrometry.

[0330] One exemplary proteolysis pattern can be generated using trypsin. Trypsin digest of oncofetal fibronectin can yield trypsin fragments that are 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa and/or 55 kDa. Typically, each of these six trypsin fragments binds to the antibody FDC-6. In one example, trypsin fragments from oncofetal fibronectin can be 200 kDa, 120 kDa or 55 kDa, where each smaller fragment represents a product of further trypsin cleavage of a larger fragment. In another example, trypsin fragments from an oncofetal fibronectin can be 235 kDa, 160 kDa or 65 kDa, where each smaller fragment represents a product of further trypsin cleavage of a larger fragment.

[0331] Another exemplary proteolysis pattern can be generated using cathepsin D. Cathepsin D digest of oncofetal fibronectin can yield fragments of 110 kDa and/or 85 kDa. Typically these two cathepsin D fragments can bind to the antibody FDC-6.

[0332] Another exemplary proteolysis pattern can be generated using thermolysin. Thermolysin digest of oncofetal fibronectin can yield fragments of 120 kDa, 85 kDa and/or 35 kDa. Typically the 0.120 kDa and 85 kDa can bind to the antibody BC-1 and the 85 kDa fragment represents a product of further thermolysin cleavage of the 120 kDa fragment.

[0333] Another exemplary proteolysis pattern can be generated using *Achromobacter* protease I. *Achromobacter* protease I digest of oncofetal fibronectin can yield a 14 kDa fragment, where this fragment typically can bind to the antibody FDC-6.

D. Use of Oncofetal Fibronectin as a Biological Marker

[0334] Detection of an oncofetal fibronectin indicating molecule serves as a biological marker for a variety of current or future health conditions such as general health state, cancer, pregnancy and delivery. Any of the uses of oncofetal fibronectin as a biological marker provided herein can be performed using any of sample types provided herein or known in the art and in conjunction with any of the oncofetal fibronectin indicating molecule detection methods provided herein or known in the art. For example, any of a variety of samples can be measured for the presence of an oncofetal fibronectin indicating molecule, including, but not limited to tissue samples, organ samples, urine, lymph, blood, plasma, serum, saliva, cervical fluid, cervicovaginal fluid, vaginal fluid, breast fluid, breast milk, synovial fluid,

semen, seminal fluid, stool, sputum, cerebral spinal fluid, tears, mucus, interstitial fluid, follicular fluid, amniotic fluid, aqueous humor, vitreous humor, peritoneal fluid, ascites, sweat, lymphatic fluid, lung sputum and lavage. Further, any of a variety of methods provided herein or otherwise known in the art for detecting an oncofetal fibronectin indicating molecule in a sample can be used, including, but not limited to, dot blot analysis, western blot analysis, northern blot analysis, southern blot analysis, RT-PCR methods, mass spectrometric methods, sandwich assays such as test strip-based sandwich assays, ELISA methods, fluorescence polarization methods, FRET methods and flow cytometry methods. Selection of any particular any particular method for detecting an oncofetal fibronectin indicating molecule is a matter of design choice, where one skilled in the art can select an appropriate assay or detection (e.g., PCR, mass spectrometry, sandwich assay) based upon the nature (e.g., protein, nucleic acid) of the oncofetal fibronectin indicating molecule to be detected. Similarly, selection of a particular sample type can be a matter of choice to one skilled in the art, and can be based on any of a variety of criteria, for example, based on the relevance of the sample type to the diagnostic purpose, on the ease of sample collection or handling or on the detection method to be used.

[0335] A sample or subject can be categorized according to the presence and/or amount of an oncofetal fibronectin indicating molecule measured. Categorization of an oncofetal fibronectin indicating molecule measurement can vary according to a variety of factors known to one skilled in the art, including the tissue or fluid sampled, the sample type, the detection method, the age, gender or biological state (e.g., pregnant or not pregnant) of a subject.

[0336] In some cases a measurement is considered positive for oncofetal fibronectin when any oncofetal fibronectin indicating molecule is detected in a sample. In other cases, a measurement is considered positive for oncofetal fibronectin when the presence of an oncofetal fibronectin indicating molecule in a sample is equal to or above one or more threshold levels. In one example, a threshold level of oncofetal fibronectin protein in a buffer-treated cervicovaginal sample assayed using a test strip can be 50 ng/mL. In another example, a threshold level for oncofetal fibronectin protein in a buffer-treated cervicovaginal sample assayed using a test strip can be 150 ng/mL.

[0337] In embodiments that compare the amount of an oncofetal fibronectin indicating molecule in a sample to a threshold level, the threshold level can be the amount of oncofetal fibronectin indicating molecule present in an unmodified sample, or the threshold level can be the amount of oncofetal fibronectin indicating molecule present in a modified sample (e.g., the concentration of an oncofetal fibronectin indicating molecule of a cervicovaginal swab sample after mixture with a buffer solution). Reference herein to the level of an oncofetal fibronectin indicating molecule in a sample or the threshold level of an oncofetal fibronectin indicating molecule typically refers to the level of an oncofetal fibronectin indicating molecule in a modified sample. For example, some oncofetal fibronectin indicating molecule measurements, such as measurement of an oncofetal fibronectin indicating molecule in a cervicovaginal swab sample, are known in the art according to the sample-modified form; thus, oncofetal fibronectin indicating mol-

ecule levels and threshold levels for a cervicovaginal swab sample typically refer to the sample modified level.

[0338] In some embodiments, the measured amount of an oncofetal fibronectin indicating molecule can be compared to one or more thresholds. Typically, an oncofetal fibronectin indicating molecule concentration in the sample equal to or above a threshold level indicates that the sample is oncofetal fibronectin positive. In one embodiment, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 50 ng/ml or more (or 500 ng/ml untreated swab sample or more), or about 50 ng/ml or more (or about 500 ng/ml untreated swab sample or more) indicates that the sample is oncofetal fibronectin positive. Typically, an oncofetal fibronectin indicating molecule concentration in the sample below a threshold level indicates that the sample is oncofetal fibronectin negative. In one embodiment, an oncofetal fibronectin indicating molecule concentration in a buffer-treated cervicovaginal swab sample of less than 50 ng/ml (or less than 500 ng/ml untreated swab sample), or about 50 ng/ml (or less than about 500 ng/ml untreated swab sample) indicates that the sample is oncofetal fibronectin negative.

[0339] Different sample types can have different threshold levels. Provided herein, different sample types also can have related threshold levels. For example, the amount of an oncofetal fibronectin indicating molecule in a cervicovaginal swab sample collected from the portion of the vagina below the posterior fornix, such as the lower third of the vagina, can be one-third or about one-third the amount of an oncofetal fibronectin indicating molecule in a cervicovaginal swab of the posterior fornix collected from the same subject. Accordingly, in methods provided herein in which the level of an oncofetal fibronectin indicating molecule in a sample is compared to a threshold level, the threshold level for a swab of the lower portion of the vagina, such as the lower third of the vagina, can be one-third or about one-third of the threshold level for a swab of the posterior fornix. For example, when the threshold level for a buffer-treated swab of the posterior fornix is 60 ng/ml (or 600 ng/ml for an untreated sample), or about 60 ng/ml (or about 600 ng/ml for an untreated sample), the threshold level of a buffer-treated swab of the lower portion of the vagina such as the lower third of the vagina can be 20 ng/ml (or 200 ng/ml for an untreated sample), or about 20 ng/ml (or about 200 ng/ml for an untreated sample). Similarly, when the threshold level for a buffer-treated swab of the posterior fornix is 300 ng/ml (or 3000 ng/ml for an untreated sample), 200 ng/ml (or 2000 ng/ml for an untreated sample), 150 ng/ml (or 1500 ng/ml for an untreated sample), 100 ng/ml (or 1000 ng/ml for an untreated sample), 50 ng/ml (or 500 ng/ml for an untreated sample), 30 ng/ml (or 300 ng/ml for an untreated sample), 15 ng/ml (or 150 ng/ml for an untreated sample) or 10 ng/ml (or 100 ng/ml for an untreated sample), the threshold level of a buffer-treated swab of the lower portion of the vagina such as the lower third of the vagina can respectively be 100 ng/ml (or 1000 ng/ml for an untreated sample), 60-70 ng/ml (or 600-700 ng/ml for an untreated sample), 50 ng/ml (or 500 ng/ml for an untreated sample), 30-40 ng/ml (or 300-400 ng/ml for an untreated sample), 15-20 ng/ml (or 150-200 ng/ml for an untreated sample), 10 ng/ml (or 100 ng/ml for an untreated sample), 5 ng/ml (or 50 ng/ml for an untreated sample) or 3-4 ng/ml (or 30-40 ng/ml for an untreated sample). Similarly, when the threshold level for a

buffer-treated swab of the posterior fornix is about 300 ng/ml (or about 3000 ng/ml for an untreated sample), about 200 ng/ml (or about 2000 ng/ml for an untreated sample), about 150 ng/ml (or about 1500 ng/ml for an untreated sample), about 100 ng/ml (or about 1000 ng/ml for an untreated sample), about 50 ng/ml (or about 500 ng/ml for an untreated sample), about 30 ng/ml (or about 300 ng/ml for an untreated sample), about 15 ng/ml (or about 150 ng/ml for an untreated sample) or about 10 ng/ml (or about 100 ng/ml for an untreated sample), the threshold level of a buffer-treated swab of the lower portion of the vagina such as the lower third of the vagina can respectively be about 100 ng/ml (or about 1000 ng/ml for an untreated sample), about 60-70 ng/ml (or about 600-700 ng/ml for an untreated sample), about 50 ng/ml (or about 500 ng/ml for an untreated sample), about 30-40 ng/ml (or about 300-400 ng/ml for an untreated sample), about 15-20 ng/ml (or about 150-200 ng/ml for an untreated sample), about 10 ng/ml (or about 100 ng/ml for an untreated sample), about 5 ng/ml (or about 50 ng/ml for an untreated sample) or about 3-4 ng/ml (or about 30-40 ng/ml for an untreated sample).

[0340] In another example, the amount of an oncofetal fibronectin indicating molecule in a urine sample can be one-tenth or about one-tenth the amount of an oncofetal fibronectin indicating molecule in a cervicovaginal swab of the posterior fornix collected from the same subject. Accordingly, in methods provided herein in which the level of an oncofetal fibronectin indicating molecule in a sample is compared to a threshold level, the threshold level for a urine sample can be one-tenth or about one-tenth of the threshold level for a swab of the posterior fornix. For example, when the threshold level for a buffer-treated swab of the posterior fornix is 60 ng/ml (or 600 ng/ml for an untreated sample) or about 60 ng/ml (or about 600 ng/ml for an untreated sample), the threshold level of a buffer-treated swab of the lower portion of the vagina such as the lower third of the vagina can be 6 ng/ml (or 60 ng/ml for an untreated sample) or about 6 ng/ml (or about 60 ng/ml for an untreated sample). Similarly, when the threshold level for a buffer-treated swab of the posterior fornix is 300 ng/ml (or 3000 ng/ml for an untreated sample), 200 ng/ml (or 2000 ng/ml for an untreated sample), 150 ng/ml (or 1500 ng/ml for an untreated sample), 100 ng/ml (or 1000 ng/ml for an untreated sample), 50 ng/ml (or 500 ng/ml for an untreated sample), 30 ng/ml (or 300 ng/ml for an untreated sample), 15 ng/ml (or 150 ng/ml for an untreated sample) or 10 ng/ml (or 100 ng/ml for an untreated sample), the threshold level of a urine sample can respectively be 30 ng/ml (or 300 ng/ml for an untreated sample), 20 ng/ml (or 200 ng/ml for an untreated sample), 15 ng/ml (or 150 ng/ml for an untreated sample), 10 ng/ml (or 100 ng/ml for an untreated sample), 5 ng/ml (or 50 ng/ml for an untreated sample), 3 ng/ml (or 30 ng/ml for an untreated sample), 1.5 ng/ml (or 15 ng/ml for an untreated sample) or 1 ng/ml (or 10 ng/ml for an untreated sample). Similarly, when the threshold level for a buffer-treated swab of the posterior fornix is about 300 ng/ml (or about 3000 ng/ml for an untreated sample), about 200 ng/ml (or about 2000 ng/ml for an untreated sample), about 150 ng/ml (or about 1500 ng/ml for an untreated sample), about 100 ng/ml (or about 1000 ng/ml for an untreated sample), about 50 ng/ml (or about 500 ng/ml for an untreated sample), about 30 ng/ml (or about 300 ng/ml for an untreated sample), about 15 ng/ml (or about 150 ng/ml for an untreated sample) or about 10 ng/ml (or about

100 ng/ml for an untreated sample), the threshold level of a urine sample can respectively be about 30 ng/ml (or about 300 ng/ml for an untreated sample), about 20 ng/ml (or about 200 ng/ml for an untreated sample), about 15 ng/ml (or about 150 ng/ml for an untreated sample), about 10 ng/ml (or about 100 ng/ml for an untreated sample), about 5 ng/ml (or about 50 ng/ml for an untreated sample), about 3 ng/ml (or about 30 ng/ml for an untreated sample), about 1.5 ng/ml (or about 15 ng/ml for an untreated sample) or about 1 ng/ml (or about 10 ng/ml for an untreated sample).

[0341] In another embodiment, a threshold oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample is 150 ng/ml (or 1500 ng/ml untreated swab sample) or about 150 ng/ml (or about 1500 ng/ml untreated swab sample), where a measured amount in a subject's sample at or above the 150 ng/ml threshold indicates that the sample is oncofetal fibronectin positive and a measured amount in a subject's sample below the 150 ng/ml threshold indicates that the sample is oncofetal fibronectin negative.

[0342] Exemplary threshold values for buffer-treated samples that can indicate different likelihoods of imminent or pre-term delivery include 50 ng/ml, 150 ng/ml, 200 ng/ml, 300 ng/ml, 500 ng/ml, 750 ng/ml and 1000 ng/ml, or about 50 ng/ml, about 150 ng/ml, about 200 ng/ml, about 300 ng/ml, about 500 ng/ml, about 750 ng/ml and about 1000 ng/ml. Exemplary threshold values for untreated samples that can indicate different likelihoods of imminent or pre-term delivery include 500 ng/ml, 1500 ng/ml, 2000 ng/ml, 3000 ng/ml, 5000 ng/ml, 7500 ng/ml and 10000 ng/ml, or about 500 ng/ml, about 1500 ng/ml, about 2000 ng/ml, about 3000 ng/ml, about 5000 ng/ml, about 7500 ng/ml and about 10000 ng/ml.

[0343] In other cases, multi-tiered thresholds can be applied to the oncofetal fibronectin indicating molecule measurement, where multi-tiered thresholds include two or more threshold levels, where each larger threshold level indicates a separate health state categorization; for example each larger threshold level can indicate a more severe health problem, an increased likelihood of imminent delivery, increased certainty of delivery date, or increased aggressiveness of a tumor. An exemplary multi-tiered threshold is a two-tiered threshold for oncofetal fibronectin protein, where the lower threshold is 50 ng/mL and the higher threshold is 150 ng/mL for buffer-treated samples. Another exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 500 ng/mL and the higher threshold level is 1500 ng/mL for untreated samples. Another exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 50 ng/mL and the higher threshold level is 200 ng/mL for buffer-treated samples. An exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 500 ng/mL and the higher threshold level is 2000 ng/mL for untreated samples. Another exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 50 ng/mL and the higher threshold level is 300 ng/mL for buffer-treated samples. An exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 500 ng/mL and the higher threshold level is 3000 ng/mL for untreated samples.

[0344] In some embodiments, the threshold level can vary over time, for example, as a function of the gestation period

of pregnancy, the progression of disease, or the age of the subject. The varying threshold level can be expressed as a threshold curve where the threshold level of an oncofetal fibronectin indicating molecule varies as a function of time (e.g., week of pregnancy term). In some cases, a threshold level can decrease with increasing time, such as, for example, in weeks 12 to 20 of a pregnancy. In other cases, a threshold level can increase with increasing time, such as for example, over the progression of a cancerous condition. Thus, in one example, a measured amount of an oncofetal fibronectin indicating molecule can be classified as greater than a threshold level or less than a threshold level, depending on the point along a defined time period that the sample was collected. Similarly two or more threshold levels can vary over time, resulting in two or more threshold curves that each separate different health categories. The two or more threshold levels can increase with increasing time or can decrease with increasing time. Thus, in one example, a measured amount of an oncofetal fibronectin indicating molecule can be differently categorized depending on the point along a defined time period that the sample was collected.

[0345] Thus, provided herein are methods for categorizing the health state of a subject, by measuring the amount of an oncofetal fibronectin indicating molecule in a sample and comparing the amount to two or more threshold levels or curves, where a measured amount below the lowest threshold indicates a more favorable health state and a measured amount higher than respectively higher threshold levels indicates increasingly less favorable health state, or increasingly unfavorable health problems.

[0346] In some embodiments, one or more threshold levels or one or more threshold curves applied to a measured amount in a subject's sample can be determined according to any of a variety of subject-specific factors. In one example, a subject-specific factor can be the measured amount of one or more samples from a subject. In some instances, a single sample measurement can be used to define one or more subject-specific threshold levels or one or more subject-specific threshold curves. A single sample measurement can be used, for example, to modify one or more pre-defined threshold levels or one or more threshold curves. For example, a measured sample amount can be compared to the mean or median normal amount and the ratio of the sample amount:normal amount can be applied to one or more pre-defined threshold levels or one or more threshold curves to either increase or decrease the levels or curves (e.g., a sample amount that is twice the normal amount can be applied to double one or more standard threshold levels or curves).

[0347] In some instances, the rate of change of the amount of an oncofetal fibronectin indicating molecule in a particular sample type (e.g., cervicovaginal swab) from a subject can be used to identify a sample as oncofetal fibronectin positive or negative, or to categorize the sample into two or more populations. The rate of change of the amount of an oncofetal fibronectin indicating molecule in a type of sample can indicate a stable, increasing or decreasing amount of oncofetal fibronectin indicating molecule in a sample. In some cases, when the rate of change is equal to or greater than one or more threshold rates, the rate of change can be categorized according to the highest threshold rate less than or equal to the rate of change measured in the samples.

Exemplary rates of change include an increase of 10% or more per week, an increase of 20% or more per week, an increase of 30% or more per week, an increase of 40% or more per week, an increase of 50% or more per week, an increase of 60% or more per week, an increase of 70% or more per week, an increase of 80% or more per week, an increase of 90% or more per week, or an increase of 100% or more per week. In other cases, the measured rate of change can be compared to one or more threshold curves or one or more threshold rates of change and a measured rate of change can be categorized according to the highest slope of a threshold curve or the highest rate equal to or less than the measured rate of change over the same time period.

[0348] Additional factors also can be applied to increase or decrease one or more pre-defined threshold levels, one or more threshold curves, or one or more threshold rates of change. Such additional factors can include other health state markers, for example, overall health markers, cancer markers, pregnancy or delivery markers, or genetic markers, as are exemplified herein or otherwise known in the art.

[0349] Also provided herein, methods for indicating health state of a subject can include measurement of an oncofetal fibronectin indicating molecule in a sample and also can include consideration of one or more other health markers. Health markers can include any of a variety of known markers, including markers related to overall health, pregnancy or delivery markers, or cancer or tumor markers. Any of a variety of markers related to overall health are known in the art or are provided elsewhere herein, exemplary markers include, but are not limited to, blood pressure, pulse, body weight, health history, family history or sample tests. A variety of detectable tumor markers are known in the art or are provided elsewhere herein, exemplary markers include, but are not limited to, AE1/AE3, BCA-225, Cathepsin D, E-Cadherin, Epidermal Growth Factor Receptor (EGFR), Estrogen receptor (ER), Gross Cystic Disease Fluid Protein 15 (GCDFFP-15), HOX-B3, Ki-67, p65, Progesterone Receptor (PR), Retinoblastoma (Rb) and Transglutaminase K (TGK), p21, DCC, NF-1, NF-2, BRCA-3, p16, FHIT, WT-1, MEN-I, MEN-IIa, MEN-IIb, VHL, FCC, MCC, raf, erb, src, fms, jun, trk, ret, gsp, hst, bcr/ab1, p53, c-erbB2, c-myc, MUC1, BRCA1, BRCA2, Her-2/neu, bcl-2, bax, PSA, CYFRA 21-1, PTH-RP, CA125, CEA gene family members, pro-gastrin, gastrin G17, gastrin G34, CA 19-9, CA 15-3, CA 27-29, CA 72-4, APC, SCC, HPV subtypes, TK, alphaFP, p62, Kallikrein, ras, vasopressin, gastrin releasing peptide, annexin I, annexin II, Hu and KOC. A variety of markers associated with pregnancy or delivery are known in the art or are provided elsewhere herein, exemplary markers include, but are not limited to, multiple fetus gestations, incompetent cervix, uterine anomalies, polyhydramnios, previous pre-term rupture of membranes or pre-term labor, pre-eclampsia, first trimester vaginal bleeding, little or no antenatal care, cervical length, Bishop score, effacement, parity (i.e., previous vaginal delivery by the subject), cervical dilation, gestational age, body mass index (BMI), station, consistency, transvaginal ultrasound, digital examination, maternal obesity, fetus size, maternal age, previous post-date delivery, gender of fetus, particular genetic disorders, fetal anomalies, abnormal placental formation, maternal infectious disease, endocrine disorder, cardiovascular renal hypertension, autoimmune and other immunologic disease, malnutrition and symptoms such as abdominal pain, low backache, passage of cervical mucus

and contractions. Methods of indicating health state of a subject that include measurement of an oncofetal fibronectin indicating molecule in a sample and also consideration of one or more other health markers are known in the art, including decision support systems. In one example, a decision support system, such as a neural network can analyze patient data or information, typically patient history or clinical data, to guide further testing or treatment of a subject. (see, U.S. Pat. Nos. 6,678,669 and 6,267,722).

[0350] 1. Pregnancy Indications

[0351] The present methods and probes can be used to determine whether a pregnant woman is at risk of pre-term, impending and/or imminent delivery, to predict delivery date, to predict maintenance of pregnancy, for use in methods of preventing pre-term delivery, or for use in inducing delivery.

[0352] Oncofetal fibronectin (onfFN) can contain a fetal restricted antigen and can be found in placenta, amniotic fluid and fetal connective tissue. The presence of an oncofetal fibronectin indicating molecule, for example, in cervicovaginal fluid samples in subjects after week 12 of pregnancy, is associated with a risk of impending delivery, including spontaneous abortions (12-20 weeks), pre-term delivery (20-37 weeks), term (37-42 weeks) and post-date delivery (after 42 weeks), in pregnant women. In addition, the presence of an oncofetal fibronectin indicating molecule, for example, in a cervicovaginal sample, provides a method for determining increased risk of labor and fetal membrane rupture after week 20 of pregnancy. Indication of rupture of the amniotic membrane is important in distinguishing true and false labor and when the rupture is small and the volume of amniotic liquid escaping is small, the rupture is often undetermined. The methods and systems herein provide a manner to reliably assess the risk of pregnancy and delivery-related conditions.

[0353] Any of a variety of samples can be used for pregnancy-related indications, where exemplary samples include blood, plasma, serum, interstitial fluid, urine, cervicovaginal lavage, cervicovaginal swab, swab of the lower portion of the vagina, swab of the lower third of the vagina, swab of the labia, passive cervicovaginal fluid collection, or other collection of cervical fluid and/or vaginal fluid. For example, the sample can be a cervicovaginal swab. For pregnancy-related indications, any of a variety of methods provided herein or otherwise known in the art for detecting an oncofetal fibronectin indicating molecule in a sample can be used, including, but not limited to, dot blot analysis, western blot analysis, northern blot analysis, southern blot analysis, RT-PCR methods, mass spectrometric methods, sandwich assays such as test strip-based sandwich assays and ELISA methods. For example, a test strip containing mobilizable mouse anti-oncofetal fibronectin antibody conjugated to a blue latex particle and polyclonal anti-human fibronectin antibody immobilized to the test strip, can be used to detect oncofetal fibronectin protein in conjunction with pregnancy-related indications.

[0354] Low or undetectable amounts of an oncofetal fibronectin indicating molecule indicate decreased risk of pre-term, impending and/or imminent delivery, decrease the ability to predict delivery date, predict increased likelihood of maintaining pregnancy, indicate decreased benefit from using methods of preventing pre-term delivery, or indicate

decreased likelihood of success in inducing delivery. The methods provided herein can be sensitive and specific and have a high negative predictive value. For example, a large percentage of subjects who do not deliver early, who have a less certain predicted delivery date, who maintain pregnancy, who do not require methods of preventing pre-term delivery, or with less successful induction outcomes, have low oncofetal fibronectin indicating molecule values. As such, the test is an effective screening procedure for pregnant women.

[0355] In one embodiment, an elevated amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman indicates that the woman has an increased risk of pre-term, impending and/or imminent delivery, increased accuracy in predicted delivery date, decreased likelihood of maintaining pregnancy, increased benefit from using methods of preventing pre-term delivery, or increased likelihood of success in inducing delivery, relative to women with a lower level. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a full 39-week term who will soon deliver is elevated over the level for pregnant women at or near a full 39-week term who will not soon deliver. In another example an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman beyond full term who will soon deliver is elevated over the level for pregnant women beyond a full term who will not soon deliver. In another example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman with an increased risk of pre-term delivery is elevated over the level for pregnant women at the same stage of gestation with a decreased risk of pre-term delivery. In another example, an amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman whose delivery date can be more accurately predicted is elevated over the level for pregnant women whose delivery date is less accurately predicted. In another example, an amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman with an increased likelihood of maintaining her pregnancy is lower than the level for pregnant women with a decreased likelihood of maintaining their pregnancy. In another example, an amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman with a decreased benefit from methods of preventing pre-term delivery is lower than the level for pregnant women with an increased benefit from methods of preventing pre-term delivery. In another example, the amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman with an increased likelihood of successful induction is elevated over the level for pregnant women with a decreased likelihood of successful induction.

[0356] In some embodiments, the measured amount of an oncofetal fibronectin indicating molecule can be compared to one or more thresholds. Typically, an oncofetal fibronectin indicating molecule concentration in the sample equal to or above a threshold level indicates that the sample is oncofetal fibronectin positive. In one embodiment, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 50 ng/ml or more (or 500 ng/ml untreated swab sample or more) or about 50 ng/ml or more (or about 500 ng/ml untreated swab sample or more) indicates a

pregnant woman has an increased risk of pre-term, impending and/or imminent delivery, has an increased accuracy of delivery date, has an increased likelihood of maintaining her pregnancy, has a decreased benefit from methods of preventing pre-term delivery, or has an increased likelihood of success in inducing delivery.

[0357] In another embodiment, a lower amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman indicates that the woman has a decreased risk of pre-term, impending and/or imminent delivery, decreased accuracy in predicted delivery date, increased likelihood of maintaining pregnancy, decreased benefit from using methods of preventing pre-term delivery, or decreased likelihood of success in inducing delivery, relative to women with an elevated level. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a full 39-week term who will not soon deliver is lower than the level for pregnant women at or near a full 39-week term who will soon deliver. In another example an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman beyond full term who will not soon deliver is lower than the level for pregnant women beyond a full term who will soon deliver. In another example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman with a decreased risk of pre-term delivery is lower than the level for pregnant women at the same stage of gestation with an increased risk of pre-term delivery. In another example, an amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman whose delivery date can be less accurately predicted is lower than the level for pregnant women whose delivery date is more accurately predicted. In another example, an amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman with a decreased likelihood of maintaining her pregnancy is elevated over the level for pregnant women with an increased likelihood of maintaining their pregnancy. In another example, an amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman with an increased benefit from methods of preventing pre-term delivery is elevated over the level for pregnant women with a decreased benefit from methods of preventing pre-term delivery. In another example, the amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman with a decreased likelihood of successful induction is lower than the level for pregnant women with an increased likelihood of successful induction.

[0358] Typically, an oncofetal fibronectin indicating molecule concentration in the sample below a threshold level indicates that the sample is oncofetal fibronectin negative. In one embodiment, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of less than 50 ng/ml (or less than 500 ng/ml untreated swab sample) or about 50 ng/ml (or less than about 500 ng/ml untreated swab sample) indicates a pregnant woman has a decreased risk of pre-term, impending and/or imminent delivery, has a decreased accuracy of delivery date, has a decreased likelihood of maintaining her pregnancy, has an increased benefit from methods of preventing pre-term delivery, or has a decreased likelihood of success in inducing delivery.

[0359] In another embodiment, multi-tiered thresholds can be applied to the oncofetal fibronectin indicating molecule measurement, where multi-tiered thresholds can include two or more thresholds, where each larger threshold indicates a further increased risk of pre-term, impending and/or imminent delivery, increased accuracy in predicted delivery date, decreased likelihood of maintaining pregnancy, increased benefit from using methods of preventing pre-term delivery, or increased likelihood of success in inducing delivery, relative to each lower threshold. An exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 50 ng/mL and the higher threshold level is 150 ng/mL for buffer-treated samples. An exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 500 ng/mL and the higher threshold level is 1500 ng/mL for untreated samples. Another exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 50 ng/mL and the higher threshold level is 200 ng/mL for buffer-treated samples. An exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 500 ng/mL and the higher threshold level is 2000 ng/mL for untreated samples. Another exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 50 ng/mL and the higher threshold level is 300 ng/mL for buffer-treated samples. An exemplary multi-tiered threshold contains two threshold levels where the lower threshold level is 500 ng/mL and the higher threshold level is 3000 ng/mL for untreated samples.

[0360] In accordance with the methods that include multi-tiered thresholds, methods are provided herein for classifying a sample, by measuring the amount of an oncofetal fibronectin indicating molecule in a sample and comparing the sample to two or more thresholds, where classification in each larger threshold indicates a further increased risk of pre-term, impending and/or imminent delivery, increased accuracy in predicted delivery date, decreased likelihood of maintaining pregnancy, increased benefit from using methods of preventing pre-term delivery, or increased likelihood of success in inducing delivery, relative to each lower threshold.

[0361] Also provided herein, methods for pregnancy and delivery-related indications can include measurement of an oncofetal fibronectin indicating molecule in a sample and also can include consideration of one or more other pregnancy or delivery markers. A variety of markers associated with pregnancy or delivery are known in the art or are provided elsewhere herein, exemplary markers include, but are not limited to, multiple fetus gestations, incompetent cervix, uterine anomalies, polyhydramnios, previous pre-term rupture of membranes or pre-term labor, pre-eclampsia, first trimester vaginal bleeding, little or no antenatal care, cervical length, Bishop score, effacement, parity (i.e., previous vaginal delivery by the subject), cervical dilation, gestational age, body mass index (BMI), station, consistency, transvaginal ultrasound, digital examination, maternal obesity, fetus size, maternal age, previous post-date delivery, gender of fetus, particular genetic disorders, fetal anomalies, abnormal placental formation, maternal infectious disease, endocrine disorder, cardiovascular renal hypertension, autoimmune and other immunologic disease, malnutrition and symptoms such as abdominal pain, low backache, passage of cervical mucus and contractions. Thus, provided herein are methods of determining risk of pre-term, impending

and/or imminent delivery, accuracy in predicted delivery date, likelihood of maintaining pregnancy, benefit from using methods of preventing pre-term delivery, or likelihood of success in inducing delivery, where the methods include detecting an oncofetal fibronectin indicating molecule in a sample and determining one or more additional pregnancy or delivery-related markers, where presence or higher amount of an oncofetal fibronectin indicating molecule and one or more pregnancy or delivery-related markers can indicate, relative to absence or lower amount of an oncofetal fibronectin indicating molecule and/or a positive result for one or more additional pregnancy or delivery-related markers, increased risk of pre-term, impending and/or imminent delivery, increased accuracy in predicted delivery date, decreased likelihood of maintaining pregnancy, increased benefit from using methods of preventing pre-term delivery, or increased likelihood of success in inducing delivery. Also provided herein are methods of determining risk of pre-term, impending and/or imminent delivery, accuracy in predicted delivery date, likelihood of maintaining pregnancy, benefit from using methods of preventing pre-term delivery, or likelihood of success in inducing delivery, where the methods include detecting an oncofetal fibronectin indicating molecule in a sample and determining one or more additional pregnancy or delivery-related markers, where absence or lower amount of an oncofetal fibronectin indicating molecule and one or more pregnancy or delivery-related markers can indicate, relative to presence or higher amount of an oncofetal fibronectin indicating molecule and/or a positive result for one or more additional pregnancy or delivery-related markers, decreased risk of pre-term, impending and/or imminent delivery, decreased accuracy in predicted delivery date, increased likelihood of maintaining pregnancy, decreased benefit from using methods of preventing pre-term delivery, or decreased likelihood of success in inducing delivery.

[0362] Methods of indicating health state of a subject that include measurement of an oncofetal fibronectin indicating molecule in a sample and also consideration of one or more other pregnancy or delivery-related markers are known in the art, including decision support systems. In one example, a decision support system, such as a neural network can analyze patient data or information, typically patient history or clinical data, to guide further testing or treatment of a subject. (see, U.S. Pat. Nos. 6,678,669 and 6,267,722).

[0363] The methods can be performed for all pregnant women following 12 weeks or about 12 weeks gestation until delivery. The present methods can be used for any pregnant woman after about 12 weeks, after about 13 weeks, after about 14 weeks, after about 15 weeks, after about 16 weeks, after about 17 weeks, after about 18 weeks, after about 19 weeks, after about 20 weeks, after about 21 weeks, after about 22 weeks, after about 23 weeks, after about 24 weeks, after about 25 weeks, after about 26 weeks, after about 27 weeks, after about 28 weeks, after about 29 weeks, after about 30 weeks, after about 31 weeks, after about 32 weeks, after about 33 weeks, after about 34 weeks, after about 35 weeks, after about 36 weeks, after about 37 weeks, after about 38 weeks, after about 39 weeks, after about 40 weeks, after about 41 weeks, after about 42 weeks, after about 43 weeks, after about 44 weeks, or after about 45 weeks.

[0364] For example, the present methods can be used for any pregnant woman after about day 80, after about day 81, after about day 82, after about day 83, after about day 84, after about day 85, after about day 86, after about day 87, after about day 88, after about day 89, after about day 90, after about day 91, after about day 92, after about day 93, after about day 94, after about day 95, after about day 96, after about day 97, after about day 98, after about day 99, after about day 100, after about day 101, after about day 102, after about day 103, after about day 104, after about day 105, after about day 106, after about day 107, after about day 108, after about day 109, after about day 110, after about day 111, after about day 112, after about day 113, after about day 114, after about day 115, after about day 116, after about day 117, after about day 118, after about day 119, after about day 120, after about day 121, after about day 122, after about day 123, after about day 124, after about day 125, after about day 126, after about day 127, after about day 128, after about day 129, after about day 130, after about day 131, after about day 132, after about day 133, after about day 134, after about day 135, after about day 136, after about day 137, after about day 138, after about day 139, after about day 140, after about day 141, after about day 142, after about day 143, after about day 144, after about day 145, after about day 146, after about day 147, after about day 148, after about day 149, after about day 150, after about day 151, after about day 152, after about day 153, after about day 154, after about day 155, after about day 156, after about day 157, after about day 158, after about day 159, after about day 160, after about day 161, after about day 162, after about day 163, after about day 164, after about day 165, after about day 166, after about day 167, after about day 168, after about day 169, after about day 170, after about day 171, after about day 172, after about day 173, after about day 174, after about day 175, after about day 176, after about day 177, after about day 178, after about day 179, after about day 180, after about day 181, after about day 182, after about day 183, after about day 184, after about day 185, after about day 186, after about day 187, after about day 188, after about day 189, after about day 190, after about day 191, after about day 192, after about day 193, after about day 194, after about day 195, after about day 196, after about day 197, after about day 198, after about day 199, after about day 200, after about day 201, after about day 202, after about day 203, after about day 204, after about day 205, after about day 206, after about day 207, after about day 208, after about day 209, after about day 210, after about day 211, after about day 212, after about day 213, after about day 214, after about day 215, after about day 216, after about day 217, after about day 218, after about day 219, after about day 220, after about day 221, after about day 222, after about day 223, after about day 224, after about day 225, after about day 226, after about day 227, after about day 228, after about day 229, after about day 230, after about day 231, after about day 232, after about day 233, after about day 234, after about day 235, after about day 236, after about day 237, after about day 238, after about day 239, after about day 240, after about day 241, after about day 242, after about day 243, after about day 244, after about day 245, after about day 246, after about day 247, after about day 248, after about day 249, after about day 250, after about day 251, after about day 252, after about day 253, after about day 254, after about day 255, after about day 256, after about day 257, after about day 258, after about day 259, after about day 260, after about day 261, after about day 262, after about day 263, after about day 264, after about day 265, after about day 266, after about day 267,

after about day 268, after about day 269, after about day 270, after about day 271, after about day 272, after about day 273, after about day 274, after about day 275, after about day 276, after about day 277, after about day 278, after about day 279, after about day 280, after about day 182, after about day 282, after about day 283, after about day 284, after about day 285, after about day 286, after about day 287, after about day 288, after about day 289, after about day 290, after about day 291, after about day 292, after about day 293, after about day 294, after about day 295, after about day 296, after about day 297, after about day 298, after about day 299, after about day 300, after about day 301, after about day 302, after about day 303, after about day 304, after about day 305, after about day 306, after about day 307, after about day 308, after about day 309, after about day 310, after about day 311, after about day 312, after about day 313, after about day 314, or after about day 315 of pregnancy.

[0365] a. Likelihood of Pre-Term Delivery

[0366] The methods provided herein can be used to determine the likelihood of pre-term delivery for a pregnant subject. For example, a large percentage of subjects who do not deliver pre-term have low oncofetal fibronectin indicating molecule values. Thus, the methods provided herein can be sensitive and specific and have a high negative predictive value. The methods provided herein can indicate an increased or decreased likelihood of imminent or pre-term delivery, according to the amount of an oncofetal fibronectin indicating molecule measured in a sample. For example, a subject with a decreased likelihood of imminent or pre-term delivery can have measured amounts of an oncofetal fibronectin indicating molecule lower than levels in women with an increased likelihood of imminent or pre-term delivery and a subject with an increased likelihood of imminent or pre-term delivery can have measured amounts of an oncofetal fibronectin indicating molecule higher than levels in women with a decreased likelihood of imminent or pre-term delivery. A measured amount of an oncofetal fibronectin indicating molecule also can be compared to one or more thresholds, where the likelihood of imminent or pre-term delivery increases for each increasing threshold level. A measured amount of an oncofetal fibronectin indicating molecule also can be considered along with one or more other pregnancy or delivery-related markers in determining the likelihood of imminent or pre-term delivery.

[0367] Elevated levels of an oncofetal fibronectin indicating molecule can indicate increased risk of pre-term delivery. In one embodiment, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman who will soon deliver is elevated over the level for pregnant women who will not soon deliver. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at increased risk of pre-term delivery is elevated over the level for pregnant women at the same stage of gestation with a decreased risk of pre-term delivery.

[0368] Typically, an oncofetal fibronectin indicating molecule concentration in the sample above or equal to a threshold indicates an increased likelihood of imminent or pre-term delivery relative to an oncofetal fibronectin indicating molecule concentration in the sample below the threshold. For example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentra-

tion in a buffer-treated cervicovaginal swab sample of 50 ng/ml or more (or 500 ng/ml untreated swab sample or more) or about 50 ng/ml or more (or about 500 ng/ml untreated swab sample or more) indicates an increased likelihood of imminent or pre-term delivery relative to an oncofetal fibronectin indicating molecule concentration in a sample below 50 ng/ml or about 50 ng/ml.

[0369] In another embodiment, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman who will not soon deliver is below the level for pregnant women who will soon deliver. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman with a decreased risk of pre-term delivery is below the level for pregnant women at the same stage of gestation with an increased risk of pre-term delivery.

[0370] Typically, an oncofetal fibronectin indicating molecule concentration in the sample below a threshold value indicates a decreased likelihood of imminent or pre-term delivery relative to an oncofetal fibronectin indicating molecule concentration in the sample at or above the threshold. For example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample less than 50 ng/ml (or 500 ng/ml untreated swab sample) or about 50 ng/ml (or about 500 ng/ml untreated swab sample) indicates a decreased likelihood of imminent or pre-term delivery relative to an oncofetal fibronectin indicating molecule concentration in the sample at or above 50 ng/ml or about 50 ng/ml.

[0371] In another embodiment, a threshold oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample is 150 ng/ml (or 1500 ng/ml untreated swab sample) or about 150 ng/ml (or about 1500 ng/ml untreated swab sample), where a measured amount in a subject's sample at or above the 150 ng/ml threshold indicates a higher risk of imminent or pre-term delivery and a measured amount in a subject's sample below the 150 ng/ml threshold indicates a lower risk of imminent or pre-term delivery. For example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 150 ng/ml or more (or 1500 ng/ml untreated swab sample or more) or about 150 ng/ml or more (or about 1500 ng/ml untreated swab sample or more) can indicate a 5%, 10%, 15%, 20% or greater likelihood of imminent or pre-term delivery. In one example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 150 ng/ml or more (or 1500 ng/ml untreated swab sample or more) or about 150 ng/ml or more (or about 1500 ng/ml untreated swab sample or more) can indicate a 5% or greater likelihood of delivery within six weeks or about six weeks, a 5% or greater likelihood of delivery within four weeks or about four weeks, a 5% or greater likelihood of delivery within two weeks or about two weeks, or a 5% or greater likelihood of delivery within a week or about a week. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein concentration in a buffer-treated cervicovaginal swab sample of 150 ng/ml or more (or 1500 ng/ml untreated swab sample or more) or about 150 ng/ml or more (or about 1500 ng/ml untreated swab sample

or more) can indicate a 10% or greater likelihood of delivery within six weeks or about six weeks, a 10% or greater likelihood of delivery within four weeks or about four weeks, or a 10% or greater likelihood of delivery within two weeks or about two weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 150 ng/ml or more (or 1500 ng/ml untreated swab sample or more) or about 150 ng/ml or more (or about 1500 ng/ml untreated swab sample or more) can indicate a 20% or greater likelihood of delivery within eight weeks or about eight weeks, a 20% or greater likelihood of delivery within six weeks or about six weeks, or a 20% or greater likelihood of delivery within four weeks or about four weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 150 ng/ml or more (or 1500 ng/ml untreated swab sample or more) or about 150 ng/ml or more (or about 1500 ng/ml untreated swab sample or more) can indicate a 25% or greater likelihood of delivery within ten weeks or about ten weeks, a 25% or greater likelihood of delivery within eight weeks or about eight weeks, or a 25% or greater likelihood of delivery within six weeks or about six weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 150 ng/ml or more (or 1500 ng/ml untreated swab sample or more) or about 150 ng/ml or more (or about 1500 ng/ml untreated swab sample or more) can indicate a 30% or greater likelihood of delivery within ten weeks or about ten weeks, or a 30% or greater likelihood of delivery within eight weeks or about eight weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 150 ng/ml or more (or 1500 ng/ml untreated swab sample or more) or about 150 ng/ml or more (or about 1500 ng/ml untreated swab sample or more) can indicate a 35% or greater likelihood of delivery within ten weeks or about ten weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 150 ng/ml or more (or 1500 ng/ml untreated swab sample or more) or about 150 ng/ml or more (or about 1500 ng/ml untreated swab sample or more) can indicate a 40% or greater likelihood of delivery within twelve weeks or about twelve weeks.

[0372] In another embodiment, a threshold oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample is 200 ng/ml (or 2000 ng/ml untreated swab sample) or about 200 ng/ml (or about 2000 ng/ml untreated swab sample), where a measured amount in a subject's sample at or above the 200 ng/ml threshold indicates a higher risk of imminent or pre-term delivery and a measured amount in a subject's sample below the 200 ng/ml threshold indicates a lower risk of imminent or pre-term delivery. For example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in buffer-treated cervicovaginal swab sample of 200 ng/ml (or 2000 ng/ml untreated swab sample) or about 200 ng/ml (or about 2000 ng/ml untreated swab sample) can indicate a 5%, 10%, 15%, 20% or greater likelihood of imminent or pre-term delivery. In one example, an oncofetal fibronectin indicating

molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml (or 2000 ng/ml untreated swab sample) or about 200 ng/ml (or about 2000 ng/ml untreated swab sample) can indicate a 5% or greater likelihood of delivery within six weeks or about six weeks, a 5% or greater likelihood of delivery within four weeks or about four weeks, a 5% or greater likelihood of delivery within two weeks or about two weeks, or a 5% or greater likelihood of delivery within a week or about a week. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml (or 2000 ng/ml untreated swab sample) or about 200 ng/ml (or about 2000 ng/ml untreated swab sample) can indicate a 10% or greater likelihood of delivery within six weeks or about six weeks, a 10% or greater likelihood of delivery within four weeks or about four weeks, or a 10% or greater likelihood of delivery within two weeks or about two weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml (or 2000 ng/ml untreated swab sample) or about 200 ng/ml (or about 2000 ng/ml untreated swab sample) can indicate a 20% or greater likelihood of delivery within eight weeks or about eight weeks, a 20% or greater likelihood of delivery within six weeks or about six weeks, or a 20% or greater likelihood of delivery within four weeks or about four weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml (or 2000 ng/ml untreated swab sample) or about 200 ng/ml (or about 2000 ng/ml untreated swab sample) can indicate a 25% or greater likelihood of delivery within ten weeks or about ten weeks, a 25% or greater likelihood of delivery within eight weeks or about eight weeks, or a 25% or greater likelihood of delivery within about six weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml (or 2000 ng/ml untreated swab sample) or about 200 ng/ml (or about 2000 ng/ml untreated swab sample) can indicate a 30% or greater likelihood of delivery within ten weeks or about ten weeks, or a 30% or greater likelihood of delivery within eight weeks or about eight weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml (or 2000 ng/ml untreated swab sample) or about 200 ng/ml (or about 2000 ng/ml untreated swab sample) can indicate a 35% or greater likelihood of delivery within ten weeks or about ten weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml (or 2000 ng/ml untreated swab sample) or about 200 ng/ml (or about 2000 ng/ml untreated swab sample) can indicate a 40% or greater likelihood of delivery within twelve weeks or about twelve weeks.

[0373] Exemplary subjects for pre-term delivery screening are those subjects with a gestational age of 80 days or about 80 days gestation or later, or 12 weeks or about 12 weeks gestation or later, until delivery, or at least until the risk of premature delivery (i.e., until week 37 or about week 37) ceases. Typically, the subjects have intact membranes.

[0374] In some instances, a subject can be tested on multiple occasions. For example, if the oncofetal fibronectin indicating molecule amount is above a threshold value, the subject can be subsequently tested again for the presence of oncofetal fibronectin indicating molecule in her cervicovaginal secretions. For subjects that are oncofetal fibronectin positive, testing can be performed more frequently than performed for subjects testing negative for oncofetal fibronectin. In addition, for oncofetal fibronectin positive subjects, measures to determine or enhance fetal lung maturity, or to prolong the pregnancy, can be undertaken. If the oncofetal fibronectin indicating molecule assay is negative, the subject can be monitored and repeated evaluations of the subject's oncofetal fibronectin indicating molecule levels can be performed on subsequent visits. In general, subjects can be examined every two weeks from 12 to 36 or about 12 to 36 weeks and weekly from week 36 or about week 36. If the oncofetal fibronectin indicating molecule test is negative, the test can be repeated on each subsequent antenatal visit until either the test is positive or the subject has delivered her baby.

[0375] b. Preventing Pre-Term Delivery

[0376] The methods of detecting the presence of an oncofetal fibronectin indicating molecule also can be used to prevent pre-term delivery of a pregnant woman. As with other pregnancy or delivery-related methods, a sample from a pregnant woman that is oncofetal fibronectin positive can indicate that the woman has an increased likelihood of imminent or pre-term delivery relative to pregnant women with samples that are oncofetal fibronectin negative. For such women with an increased likelihood of imminent or pre-term delivery, methods can be performed that can favor or extend pregnancy, or increase the viability of an infant delivered pre-term. For example, by monitoring the amount of an oncofetal fibronectin indicating molecule in a subject, when an elevated level of an oncofetal fibronectin indicating molecule indicative of increased likelihood of pre-term or imminent delivery is measured, progestational therapy such as a tocolytic agent can be administered to the subject. A measured amount of an oncofetal fibronectin indicating molecule also can be compared to one or more thresholds, where the benefit from using methods of preventing pre-term delivery increases for each increasing threshold level. A measured amount of an oncofetal fibronectin indicating molecule also can be considered along with one or more other pregnancy or delivery-related markers in determining the likelihood of imminent or pre-term delivery.

[0377] Thus, provided herein is a method of screening and, if appropriate, treating a pregnant subject, by obtaining a sample from the subject, measuring the amount of an oncofetal fibronectin indicating molecule in the sample and assessing whether the level of an oncofetal fibronectin indicating molecule is equal to or above a threshold level that is indicative of an increased risk of pre-term or imminent delivery and if the amount of an oncofetal fibronectin indicating molecule is equal to or above the threshold level, administering progestational therapy, such as a therapeutically effective amount of a progestational agent to the subject. Methods provided herein for screening a pregnant subject at risk of pre-term delivery and treating the subject with progestational therapy such as a tocolytic agent also can be used in conjunction with the methods provided in

copending application U.S. patent application Ser. No. 10/774,144, which is incorporated by reference herein in its entirety.

[0378] Progestational therapy such as administration of a tocolytic agent, favors, or is conducive to, gestation, or inhibits premature labor by, for example, inhibiting uterine contractions, or increases the viability of an infant born pre-term. Progestational therapy that can be administered in accordance with the methods provided herein can include any of a variety of techniques for prolonging gestation, inhibiting premature labor, or increasing the viability of an infant born pre-term. Progestational therapy can include methods such as bedrest for the pregnant subject and also can include administration of one or more agents that reduce or inhibits uterine contractions, that prolongs the pregnancy, or that increases the viability of an infant delivered pre-term. For example, progestational therapy can include administration of a tocolytic agent.

[0379] Tocolytic agents include any of a group of agents known to inhibit pre-term labor. Any such agent can be employed. Exemplary tocolytic agents for use in accord with the methods herein include, but are not limited to, any of the following: magnesium salts such as magnesium sulfate; prostaglandin synthesis inhibitors, including non-steroidal anti-inflammatory compounds such as indomethacin, sulindac, naproxen, aspirin and fenoprofen; β -adrenergic agonists such as ritodrine, terbutaline, albuterol, fenoterol, hexoprenaline, isoxuprine, metaproterenol, nylidrin, orciprenaline and salbutamol, or other epinephrin or norepinephrine analogs or derivatives; calcium channel blockers such as nifedipine and nicardipine; oxytocin antagonists such as atosiban; nitric oxide donors such as glyceryl trinitrate; hormones secreted by the corpus luteum, placenta and adrenal cortex and derivatives thereof, including dydrogesterone, ethynodiol diacetate, hydroxyprogesterone caproate, medroxyprogesterone acetate, norethindrone, norethindrone acetate, norethynodrel, norgestrel, megestrol acetate, gestodene, desogestrel, cingestol, lynestrenol, quingestanol acetate, levonorgestrel, 3-ketodesogestrel, norgestimate, osaterone, cyproterone acetate, trimegestone, dienogest, drospirenone, nomegestrol, (17-deacetyl)norgestimnate, 19-norprogesterone, melengestrol, ethisterone, medroxyprogesterone acetate, 17- α -hydroxyprogesterone, dime-thisterone, ethinylestrenol, demegestone, promegestone, chlormadinone, pregn-4-ene-3,20-dione (progesterone), 19-nor-pregn-4-ene-3,20-dione, 17-hydroxy-19-nor-17 α -pregn-5(10)-ene-20-yn-3-one, dl-11 α -ethyl-17-ethinyl-17- α -hydroxygon-4-ene-3-one, 17-ethinyl-17-hydroxy-5(10)-estren-3-one, 17 α -ethinyl-19-norestosterone, 6-chloro-17-hydroxypregna-4,6-diene-3,20-dione, 17 α -hydroxy-6 α -methyl-17(-1-propynyl)-androst-4-ene-3-one, 9 α ,10 α -pregna-4,6-diene-3,20-dione, 17-hydroxy-17 α -pregn-4-en-20-yne-3-one, 19-nor-17 α -preg-4-en-20-yen-3,17-diol, 17-hydroxy-pregn-4-ene-3,20-dione, 1-7-hydroxy-6 α -methylpregn-4-ene-3,20-dione and derivatives and mixtures thereof (see, e.g., U.S. Pat. No. 5,211,952). Tocolytic agents also include omega-3 fatty acids, whether naturally or synthetically produced and derivatives thereof. Exemplary omega-3 fatty acids include, for example, docosahexaenoic acid (DHA).

[0380] Tocolytic agents can be administered by any of a variety of methods known in the art. For example, the tocolytic agent can be administered orally, parenterally by

injection (e.g., by bolus injection or continuous infusion), transdermally, intranasally, or by inhalation. The therapeutically effective amount of tocolytic agent will vary according to, for example, the particular agent and/or pharmaceutical composition being used, the mode of administration and the course of treatment. Optimal dosages for a given set of conditions can be ascertained using conventional dosage-determination tests. Further, administration of the tocolytic agent can be repeated at appropriate intervals (e.g., daily, weekly, etc.). In one embodiment, the dose is determined by measuring the concentration of tocolytic agent in the circulating blood and adjusting the mode of administration and/or course of treatment accordingly.

[0381] c. Predictor of Delivery Date

[0382] In another embodiment, the methods of detecting the presence of an oncofetal fibronectin indicating molecule can be used to predict the delivery date of a pregnant woman. As with the methods provided for prediction of pre-term or imminent delivery provided herein, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman who will soon deliver is elevated over the level for pregnant women who will not soon deliver. The methods provided herein can be used to indicate a subject with an increased likelihood of imminent delivery, as well as an increased likelihood of delivery within a particular time frame. The methods provided herein also can be used to indicate the likelihood in which a subject will soon deliver or will deliver within a particular time frame. The methods provided herein also can include comparing a measured amount of an oncofetal fibronectin indicating molecule to one or more thresholds, where the likelihood of imminent delivery increases for each increasing threshold level, or where the likelihood of delivery within a particular time frame increases with each increasing threshold level. A measured amount of an oncofetal fibronectin indicating molecule also can be considered along with one or more other pregnancy or delivery-related markers in determining the likelihood of imminent delivery or the likelihood of delivery within a particular time frame.

[0383] Additionally, provided herein are methods in which an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman who will maintain her pregnancy is lower than the level for pregnant women who will soon deliver. The methods provided herein can be used to indicate a subject with an increased likelihood of maintaining her pregnancy until full term, as well as an increased likelihood of maintaining pregnancy for a particular time frame. The methods provided herein also can be used to indicate the likelihood in which a subject will maintain her pregnancy until full term or will maintain her pregnancy for a particular time frame. The methods provided herein also can include comparing a measured amount of an oncofetal fibronectin indicating molecule to one or more thresholds, where the likelihood of maintaining pregnancy increases for each decreasing threshold level, or where the likelihood of maintaining pregnancy for a particular time frame increases with each decreasing threshold level. A measured amount of an oncofetal fibronectin indicating molecule also can be considered along with one or more other pregnancy or delivery-related markers in determining the likelihood of maintaining a pregnancy to full term or the likelihood of maintaining a pregnancy for a particular time frame.

[0384] In one embodiment, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman who will soon deliver is elevated over the level for pregnant women who will not soon deliver. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 24-week gestation who will soon deliver is elevated over the level for pregnant women at or near a 24-week gestation who will not soon deliver. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 35-week gestation who will soon deliver is elevated over the level for pregnant women at or near a 35-week gestation who will not soon deliver. In another example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 37-week gestation who will soon deliver is elevated over the level for pregnant women at or near a 37-week gestation who will not soon deliver. In another example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 39-week gestation who will soon deliver is elevated over the level for pregnant women at or near a 39-week gestation who will not soon deliver. In another example an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman beyond full term who will soon deliver is elevated over the level for pregnant women beyond a full term who will not soon deliver. In such examples, a threshold level can be defined, where an oncofetal fibronectin indicating molecule amount in a sample that is equal to or above the threshold level indicates an increased likelihood that the subject will soon deliver and an oncofetal fibronectin indicating molecule amount in a sample that is below the threshold level indicates an increased likelihood that the subject will not soon deliver.

[0385] In one embodiment, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman who will maintain her pregnancy is lower than the level for pregnant women who will not maintain her pregnancy. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 24-week gestation who will maintain her pregnancy is lower than the level for pregnant women at or near a 24-week gestation who will not maintain her pregnancy. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 35-week gestation who will maintain her pregnancy is lower than the level for pregnant women at or near a 35-week gestation who will not maintain her pregnancy. In another example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 37-week gestation who will maintain her pregnancy is lower than the level for pregnant women at or near a 37-week gestation who maintain her pregnancy. In another example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 39-week gestation who will maintain her pregnancy is lower than the level for pregnant women at or near a 39-week gestation who will not maintain her pregnancy. In another example an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman beyond full term who will maintain her pregnancy is lower than the level for pregnant women beyond a full term who will not

maintain her pregnancy. In such examples, a threshold level can be defined, where an oncofetal fibronectin indicating molecule amount in a sample that is below the threshold level indicates an increased likelihood that the subject will maintain her pregnancy and an oncofetal fibronectin indicating molecule amount in a sample that is equal to or above the threshold level indicates an increased likelihood that the subject will not maintain her pregnancy.

[0386] Typically, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman who will deliver within a particular time period is elevated over the level for pregnant women who will not deliver within that time period. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy who will deliver within a particular time period is elevated over the level for pregnant women at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy who will not soon deliver. In such examples, a threshold level can be defined, where an oncofetal fibronectin indicating molecule amount in a sample that is equal to or above the threshold level indicates an increased likelihood that the subject will deliver within a particular time period and an oncofetal fibronectin indicating molecule amount in a sample that is below the threshold level indicates an increased likelihood that the subject will not deliver within that time period. Exemplary time periods for which a likelihood of delivery can be indicated include 5 months or less, 4 months or less, 14 weeks or less, 3 months or less, 12 weeks or less, 11 weeks or less, 10 weeks or less, 9 weeks or less, 2 months or less, 8 weeks or less, 7 weeks or less, 6 weeks or less, 5 weeks or less, 1 month or less, 4 weeks or less, 3 weeks or less, 2 weeks or less, 10 days or less, 1 week or less, 6 days or less, 5 days or less, 4 days or less, 3 days or less, 2 days or less, or 1 day or less.

[0387] In related methods, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman who will maintain pregnancy for a particular time period is lower than the level for pregnant women who will not maintain pregnancy over that time period. For example, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy who will maintain pregnancy for a particular time period is lower than the level for pregnant women at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy who will not maintain pregnancy over this time period. In such examples, a threshold level can be defined, where an oncofetal fibronectin indicating molecule amount in a sample that is below the threshold level indicates an increased likelihood that the subject will maintain pregnancy for a particular time period and an oncofetal fibronectin indicating molecule amount in a sample that is equal to or greater than the threshold level indicates an increased likelihood that the subject will not maintain pregnancy over that time period. Exemplary time periods for which a likelihood of delivery can be indicated include at least 1 day, at least 2 days, at least 3 days, at least 4 days, at least 5 days, at least 6 days, at least 1 week, at least 10 days, at least 2 weeks, at least 3 weeks, at least 4 weeks, at least 1 month, at least 5 weeks, at least 6 weeks, at least 7 weeks, at least 8 weeks, at least 2 months, at least 9 weeks,

at least 10 weeks, at least 11 weeks, at least 12 weeks, at least 3 months, at least 14 weeks, at least 4 months, or at least 5 months.

[0388] In some instances, the degree of likelihood of delivering can be indicated according to the amount of an oncofetal fibronectin indicating molecule present in the sample of a subject. For example, a sample from a pregnant woman with an elevated amount of an oncofetal fibronectin indicating molecule indicates a higher likelihood of imminent delivery relative to pregnant women with lower levels of the oncofetal fibronectin indicating molecule. For example, an elevated amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy indicates a higher likelihood of imminent delivery relative to pregnant women at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy with lower levels of the oncofetal fibronectin indicating molecule. In another example, a sample from a pregnant woman with an elevated amount of an oncofetal fibronectin indicating molecule indicates a higher likelihood of delivery within a particular time period relative to pregnant women with lower levels of the oncofetal fibronectin indicating molecule. For example, an elevated amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy indicates a higher likelihood of delivery within a particular time period relative to pregnant women at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy with lower levels of the oncofetal fibronectin indicating molecule. Exemplary higher likelihoods of delivery can be at least 5%, 10%, 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% (e.g., 2-fold), 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 12-fold, 15-fold, 20-fold, 25-fold, 30-fold, 35-fold, 40-fold, 45-fold, or 50-fold higher likelihood of delivery. Exemplary higher likelihoods of delivery for particular time periods include at least 2-fold, 5-fold, 10-fold, 15-fold, 20-fold higher, 30-fold, or 40-fold higher likelihood of delivery within two weeks or about two weeks; at least 50%, 75%, 2-fold, 5-fold, 10-fold, or 15-fold higher likelihood of delivery within six weeks or about six weeks; at least 10%, 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% or 3-fold higher likelihood of delivery within three months or about three months.

[0389] Similarly, the degree of likelihood of maintaining pregnancy can be indicated according to the amount of an oncofetal fibronectin indicating molecule present in the sample of a subject. For example, a sample from a pregnant woman with a lower amount of an oncofetal fibronectin indicating molecule indicates a higher likelihood of maintaining pregnancy relative to pregnant women with elevated levels of the oncofetal fibronectin indicating molecule. For example, a lower amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy indicates a higher likelihood of maintaining pregnancy relative to pregnant women at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy with elevated levels of the oncofetal fibronectin indicating molecule. In another example, a sample from a pregnant woman with a lower amount of an oncofetal fibronectin indicating molecule indicates a higher likelihood of maintaining pregnancy for a particular time period rela-

tive to pregnant women with elevated levels of the oncofetal fibronectin indicating molecule. For example, a lower amount of an oncofetal fibronectin indicating molecule in the sample from a pregnant woman at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy indicates a higher likelihood of maintaining pregnancy for a particular time period relative to pregnant women at or near a 24 week, 35 week, 37 week, 39 week, or beyond full-term pregnancy with elevated levels of the oncofetal fibronectin indicating molecule. Exemplary higher likelihoods of maintaining pregnancy can be at least 1%, 2%, 3%, 4%, 5%, 7%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100%, 110%, 120%, 130%, 150%, 175%, or 200% higher likelihood of maintaining pregnancy. Exemplary likelihoods of maintaining pregnancy for particular time periods include at least 1%, 2%, 3%, 5%, 7%, 10%, 15%, 20%, 25%, 30%, 40%, or 50% higher likelihood of maintaining pregnancy for at least two weeks or about two weeks; at least 2%, 5%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, or 75% higher likelihood of maintaining pregnancy for at least six weeks or about six weeks; at least 10%, 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% higher likelihood of delivery maintaining pregnancy for at least three months or about three months.

[0390] In such examples, a threshold level can be defined, where an oncofetal fibronectin indicating molecule amount in a sample that is equal to or above the threshold level indicates an increased likelihood that the subject will soon deliver and an oncofetal fibronectin indicating molecule amount in a sample that is below the threshold level indicates an increased likelihood that the subject will not soon deliver.

[0391] For use in conjunction with delivery date and pregnancy maintenance prediction methods herein, a subject is considered positive for oncofetal fibronectin when the presence of an oncofetal fibronectin indicating molecule in a sample is equal to or above one or more threshold levels. As one skilled in the art will recognize, a threshold level can vary according to the type of sample measured and the selected stringency of the test. In one example, a threshold level for a cervicovaginal sample assayed using a test strip can be 50 ng/mL. In another example, a threshold level for a cervicovaginal sample assayed using a test strip can be 150 ng/mL.

[0392] In another embodiment, multi-tiered thresholds can be applied to the oncofetal fibronectin indicating molecule measurement, where multi-tiered thresholds include two or more threshold levels, where each larger threshold level indicates a further increased likelihood of imminent delivery or an increased likelihood of delivery within a particular time period, or each lower threshold level indicates a further increased likelihood of maintaining pregnancy or an increased likelihood of maintaining pregnancy for a particular time period. An exemplary multi-tiered threshold is a two-tiered threshold where the lower threshold is 50 ng/mL and the higher threshold is 150 ng/mL for buffer-treated samples. An exemplary multi-tiered threshold is a two-tiered threshold where the lower threshold is 500 ng/mL and the higher threshold is 1500 ng/mL for untreated samples.

[0393] As provided herein, particular threshold levels can be identified with particular likelihoods of delivery or like-

tin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml or more (or 2000 ng/ml untreated swab sample or more) or about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can indicate a 20% likelihood of delivery within 4 weeks or about 4 weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml or more (or 2000 ng/ml untreated swab sample or more) or about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can indicate a 25% likelihood of delivery within 6 weeks or about 6 weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml or more (or 2000 ng/ml untreated swab sample or more) or about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can indicate a 30% likelihood of delivery within 8 weeks or about 8 weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml or more (or 2000 ng/ml untreated swab sample or more) or about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can indicate a 40% likelihood of delivery within 8 weeks or about 8 weeks.

[0396] In one example, a woman can be tested for the presence of oncofetal fibronectin indicating molecule at 24 weeks or about 24 weeks into her pregnancy and a woman having an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of 200 ng/ml or more (or 2000 ng/ml untreated swab sample or more) or about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can have a higher risk of imminent or pre-term delivery. For example, a woman at 24 weeks or about 24 weeks in her pregnancy and having an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can have a 5% or about a 5% likelihood of delivery within a week or about a week. For example, a woman at 24 weeks or about 24 weeks in her pregnancy and having an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can have a 10% or about a 10% likelihood of delivery within two weeks or about two weeks. For example, a woman at 24 weeks or about 24 weeks in her pregnancy and having an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can have a 20% or about a 20% likelihood of delivery within four weeks or about four weeks. For example, a woman at 24 weeks or about 24 weeks in her pregnancy and having an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can have a 25% or about a 25% likelihood of delivery within six weeks or about six weeks. For example, a woman at 24 weeks or about 24 weeks in her

pregnancy and having an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can have a 30% or about a 30% likelihood of delivery within eight weeks or about eight weeks. For example, a woman at 24 weeks or about 24 weeks in her pregnancy and having an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of about 200 ng/ml or more (or about 2000 ng/ml untreated swab sample or more) can have a 40% or about a 40% likelihood of delivery within twelve weeks or about twelve weeks.

[0397] In other examples, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in buffer-treated cervicovaginal swab sample less than 150 ng/ml (or 1500 ng/ml untreated swab sample) or about 150 ng/ml (or about 1500 ng/ml untreated swab sample) can indicate the likelihood of maintaining pregnancy for a defined amount of time. For example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in buffer-treated cervicovaginal swab sample of less than 150 ng/ml (or 1500 ng/ml untreated swab sample) or about 150 ng/ml (or about 1500 ng/ml untreated swab sample) can indicate a 99% likelihood of maintaining pregnancy for at least a week or about a week. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of less than 150 ng/ml (or 1500 ng/ml untreated swab sample) or about 150 ng/ml (or about 1500 ng/ml untreated swab sample) can indicate a 98% likelihood of maintaining pregnancy for at least two weeks or about two weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of less than 150 ng/ml (or 1500 ng/ml untreated swab sample) or about 150 ng/ml (or about 1500 ng/ml untreated swab sample) can indicate a 97% likelihood of maintaining pregnancy for at least 4 weeks or about 4 weeks. In another example, an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, concentration in a buffer-treated cervicovaginal swab sample of less than 150 ng/ml (or 1500 ng/ml untreated swab sample) or about 150 ng/ml (or about 1500 ng/ml untreated swab sample) can indicate a 95% likelihood of maintaining pregnancy for at least 10 weeks or about 10 weeks.

[0398] d. Use with Inducing Delivery

[0399] The methods of detecting the presence of an oncofetal fibronectin indicating molecule also can be used in conjunction with inducing delivery in a pregnant woman. As provided herein, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman with increased risk of delivering is elevated over the level for pregnant women with less risk of delivering. Similarly, as provided herein, subjects testing positive for oncofetal fibronectin have an increased risk of delivering after induction, or have a shorter time period between induction and delivery, or require fewer administrations of a parturifacient, or have a decreased likelihood of cesarean delivery, relative to subjects testing negative for oncofetal fibronectin.

[0400] In some embodiments, an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman with increased risk of delivering is increased compared to a threshold, or when the presence of an oncofetal fibronectin indicating molecule in a sample is equal to or above one or more threshold levels, render the sample positive for oncofetal fibronectin indicating molecule. In another embodiment, a sample is considered negative for oncofetal fibronectin when an amount of an oncofetal fibronectin indicating molecule detected in the sample from a pregnant woman is decreased compared to a threshold. As one skilled in the art will recognize, a threshold level can vary according to the type of sample measured and the selected stringency of the test.

[0401] Accordingly, by detecting an oncofetal fibronectin indicating molecule in a subject prior to performing an induction method, the likely effect of inducing delivery can be predicted and the likelihood that induction will lead to a prompt delivery can be estimated. For example, by detecting an oncofetal fibronectin indicating molecule in a subject prior to performing an induction method, it is possible to estimate the likely amount of time between administration of a parturifacient or induction procedure and delivery, or to estimate the likely amount of time between oxytocin administration and delivery. Further, by detecting an oncofetal fibronectin indicating molecule in a subject prior to performing an induction method, it is possible to estimate the likelihood of the subject delivering within 24 hours or about 24 hours after induction of delivery, or to estimate the likelihood of the subject delivering within 48 hours or about 48 hours after induction of delivery. Further, by detecting an oncofetal fibronectin indicating molecule in a subject prior to performing an induction method, it is possible to estimate the likelihood that more than a single administration of a parturifacient or induction procedure will be required to induce delivery. Further, by detecting an oncofetal fibronectin indicating molecule in a subject prior to performing an induction method, it is possible to estimate the likelihood that vaginal delivery will be performed and childbirth will not require cesarean section, or, alternatively, to estimate the likelihood that vaginal delivery will not be performed and childbirth will instead be accomplished by cesarean section.

[0402] The methods provided herein include a method for determining whether or not to perform a labor induction method, by detecting an oncofetal fibronectin indicating molecule in a sample from a subject and, if the oncofetal fibronectin indicating molecule is present in the sample at a level equal to or above a threshold level, inducing delivery in a subject. Also provided herein is a method of identifying a subject as a good candidate for induction of delivery, the method including detecting an oncofetal fibronectin indicating molecule in a sample from a subject and, if the oncofetal fibronectin indicating molecule is present in the sample at a level equal to or above a threshold level, identifying the subject as a good candidate for induction of delivery. Also provided herein is a method of identifying a subject for whom induction of delivery is likely to be successful, the method including detecting an oncofetal fibronectin indicating molecule in a sample from a subject and, if the oncofetal fibronectin indicating molecule is present in the sample at a level equal to or above a threshold level, identifying the subject as one for whom induction of delivery is likely to be successful. Also provided herein is a method of identifying a subject for whom the time period after initiation of

induction or administration of a parturifacient such as a pre-induction agent or oxytocin is likely to be relatively shorter, the method including detecting an oncofetal fibronectin indicating molecule in a sample from a subject and, if the oncofetal fibronectin indicating molecule is present in the sample at a level equal to or above a threshold level, identifying the subject as one for whom the time period after initiation of induction or administration of a parturifacient such as a pre-induction agent or oxytocin is likely to be shorter relative to subjects with oncofetal fibronectin indicating molecule levels below the threshold level. Also provided herein is a method of identifying a subject for whom the time period after initiation of induction is likely to be within 24 hours, the method including detecting an oncofetal fibronectin indicating molecule in a sample from a subject and, if the oncofetal fibronectin indicating molecule is present in the sample at a level equal to or above a threshold level, identifying the subject as one for whom the time period after initiation of induction is likely to be within 24 hours. Also provided herein is a method of identifying a subject for whom the time period after initiation of induction is likely to be within 48 hours, the method including detecting an oncofetal fibronectin indicating molecule in a sample from a subject and, if the oncofetal fibronectin indicating molecule is present in the sample at a level equal to or above a threshold level, identifying the subject as one for whom the time period after initiation of induction is likely to be within 48 hours. Also provided herein is a method of identifying a subject for whom induction of delivery is likely to lead to vaginal delivery, the method including detecting an oncofetal fibronectin indicating molecule in a sample from a subject and, if the oncofetal fibronectin indicating molecule is present in the sample at a level equal to or above a threshold level, identifying the subject as one for whom induction of delivery is likely to lead to vaginal delivery. Also provided herein is a method of identifying a subject for whom the number of induction procedures or administrations of a parturifacient is likely to be relatively fewer, the method including detecting an oncofetal fibronectin indicating molecule in a sample from a subject and, if the oncofetal fibronectin indicating molecule is present in the sample at a level equal to or above a threshold level, identifying the subject as one for whom the number of induction procedures or administrations of a parturifacient is likely to be relatively fewer relative to subjects with oncofetal fibronectin indicating molecule levels below the threshold level.

[0403] As provided herein, subjects testing positive for oncofetal fibronectin are more likely to deliver vaginally after induction relative to subjects testing negative for oncofetal fibronectin or to a sample having an amount of oncofetal fibronectin indicating molecule below a threshold. For example, subjects testing positive for oncofetal fibronectin can be at least about 7%, at least about 8%, at least about 9%, at least about 10%, at least about 11%, at least about 12%, at least about 13%, at least about 14%, at least about 15%, at least about 16%, at least about 18%, at least about 20%, at least about 25%, or at least about 30%, more likely to deliver vaginally after induction relative to subjects testing negative for oncofetal fibronectin. Typically, subjects testing positive for oncofetal fibronectin about 3-30%, about 5-25%, about 7-22%, about 8-20%, about 9-18%, about 10-16%, about 11-15%, or about 12-14%, or

about 13%, more likely to deliver vaginally after induction relative to subjects testing negative for oncofetal fibronectin.

[0404] Analogously, subjects testing negative for oncofetal fibronectin are more likely to require cesarean section after induction relative to subjects testing positive for oncofetal fibronectin or to a sample having an amount of oncofetal fibronectin indicating molecule above a threshold. For example, subjects testing negative for oncofetal fibronectin can be at least about 25%, at least about 26%, at least about 27%, at least about 28%, at least about 29%, at least about 30%, at least about 32%, at least about 34%, at least about 36%, at least about 38%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, or at least about 60%, more likely to require cesarean section after induction relative to subjects testing positive for oncofetal fibronectin. Typically, subjects testing negative for oncofetal fibronectin are about 20-60%, about 21-55%, about 22-50%, about 23-45%, about 24-40%, about 25-38%, about 26-36%, about 27-34%, about 28-32%, or about 29-30%, more likely to require cesarean section after induction relative to subjects testing positive for oncofetal fibronectin.

[0405] Also provided herein, subjects testing positive for oncofetal fibronectin are more likely to deliver within 24 hours of induction relative to subjects testing negative for oncofetal fibronectin or to a sample having an amount of oncofetal fibronectin indicating molecule below a threshold. For example, subjects testing positive for oncofetal fibronectin can be at least about 30%, at least about 31%, at least about 32%, at least about 33%, at least about 34%, at least about 35%, at least about 36%, at least about 37%, at least about 38%, at least about 40%, at least about 42%, at least about 44%, at least about 46%, or at least about 50%, more likely to deliver within 24 hours of induction relative to subjects testing negative for oncofetal fibronectin. Typically, subjects testing positive for oncofetal fibronectin are about 20-60%, about 22-55%, about 24-50%, about 26-48%, about 28-46%, about 30-44%, about 32-43%, about 34-42%, about 35-41%, about 36-40%, about 37-39%, or about 38%, more likely to deliver within 24 hours of induction relative to subjects testing negative for oncofetal fibronectin.

[0406] Analogously, subjects testing negative for oncofetal fibronectin are more likely to deliver after more than 24 hours after induction relative to subjects testing positive for oncofetal fibronectin or to a sample having an amount of oncofetal fibronectin indicating molecule above a threshold. For example, subjects testing negative for oncofetal fibronectin can be at least about 40%, at least about 41%, at least about 42%, at least about 43%, at least about 44%, at least about 45%, at least about 46%, at least about 48%, at least about 50%, at least about 52%, at least about 55%, at least about 60%, at least about 65%, at least about 75%, or at least about 90%, more likely to deliver after more than 24 hours after induction relative to subjects testing positive for oncofetal fibronectin. Typically, subjects testing negative for oncofetal fibronectin are about 30-90%, about 32-75%, about 34-65%, about 36-60%, about 38-55%, about 39-53%, about 40-51%, about 41-50%, about 42-49%, about 43-48%, about 44-47% or about 45-46%, more likely to deliver after more than 24 hours after induction relative to subjects testing positive for oncofetal fibronectin.

[0407] Also provided herein, subjects testing positive for oncofetal fibronectin are more likely to deliver within 48

hours of induction relative to subjects testing negative for oncofetal fibronectin or to a sample having an amount of oncofetal fibronectin indicating molecule below a threshold. For example, subjects testing positive for oncofetal fibronectin can be at least about 9%, at least about 10%, at least about 11%, at least about 12%, at least about 13%, at least about 14%, at least about 15%, at least about 16%, at least about 17%, at least about 18%, at least about 20%, at least about 22%, at least about 25%, at least about 30%, or at least about 35%, more likely to deliver within 48 hours of induction relative to subjects testing negative for oncofetal fibronectin. Typically, subjects testing positive for oncofetal fibronectin are about 5-35%, about 7-30%, about 8-25%, about 9-22%, about 10-20%, about 11-19%, about 12-18%, about 13-17%, about 14-16%, or about 15%, more likely to deliver within 48 hours of induction relative to subjects testing negative for oncofetal fibronectin.

[0408] Analogously, subjects testing negative for oncofetal fibronectin are more likely to deliver after more than 48 hours after induction relative to subjects testing positive for oncofetal fibronectin or to a sample having an amount of oncofetal fibronectin indicating molecule above a threshold. For example, subjects testing negative for oncofetal fibronectin can be at least about 25%, at least about 26%, at least about 27%, at least about 28%, at least about 29%, at least about 30%, at least about 32%, at least about 34%, at least about 36%, at least about 38%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, or at least about 60%, more likely to deliver after more than 48 hours after induction relative to subjects testing positive for oncofetal fibronectin. Typically, subjects testing negative for oncofetal fibronectin are about 20-60%, about 21-55%, about 22-50%, about 23-45%, about 24-40%, about 25-38%, about 26-36%, about 27-34%, about 28-32% or about 29-30%, more likely to deliver after more than 48 hours after induction relative to subjects testing positive for oncofetal fibronectin.

[0409] Also provided herein, subjects testing positive for oncofetal fibronectin have a mean time interval between first dose of pre-induction agent and delivery that is shorter than the mean time interval between first dose of parturifacient and delivery for subjects testing negative for oncofetal fibronectin or a sample having an amount of oncofetal fibronectin indicating molecule below a threshold. For example, subjects testing positive for oncofetal fibronectin can have a mean time interval between first dose of pre-induction agent and delivery that is at least about 28%, at least about 29%, at least about 30%, at least about 31%, at least about 32%, at least about 33%, at least about 34%, at least about 35%, at least about 36%, at least about 38%, at least about 40%, at least about 42%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, or at least about 75%, or at least about 6 hours, at least about 6.5 hours, at least about 7 hours, at least about 7.5 hours, at least about 8 hours, at least about 8.5 hours, at least about 9 hours, at least about 9.5 hours, at least about 10 hours, at least about 11 hours, at least about 13 hours, or at least about 15 hours, shorter than the mean time interval between first dose of parturifacient and delivery for subjects testing negative for oncofetal fibronectin. Typically, subjects testing positive for oncofetal fibronectin have a mean time interval between first dose of pre-induction agent and delivery that is 25-75% or about 25-75%, 26-60% or about 26-60%, 27-50% or about 27-50%, 28-45% or about 28-45%,

29-40% or about 29-40%, 30-38% or about 30-38%, 31-36% or about 31-36%, 32-34% or about 32-34%, or 33% or about 33%, or 6-20 hours or about 6-20 hours, 6.5-15 hours or about 6.5-15 hours, 7-14 hours or about 7-14 hours, 7.5-13 hours or about 7.5-13 hours, 8-12 hours or about 8-12 hours, 8.5-11 hours or about 8.5-11 hours, 9-10 hours or about 9-10 hours, or 9.5 or about 9.5 hours, shorter than the mean time interval between first dose of parturifacient and delivery for subjects testing negative for oncofetal fibronectin.

[0410] Analogously, subjects testing negative for oncofetal fibronectin have a mean time interval between first dose of pre-induction agent and delivery that is longer than the mean time interval between first dose of parturifacient and delivery for subjects testing positive for oncofetal fibronectin or a sample having an amount of oncofetal fibronectin indicating molecule above a threshold. For example, subjects testing negative for oncofetal fibronectin can have a mean time interval between first dose of pre-induction agent and delivery that is at least 45%, at least 46%, at least 47%, at least 48%, at least 49%, at least 50%, at least 52%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, or at least 90%, or at least about 45%, at least about 46%, at least about 47%, at least about 48%, at least about 49%, at least about 50%, at least about 52%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, or at least about 90%, or at least 6 hours, at least 6.5 hours, at least 7 hours, at least 7.5 hours, at least 8 hours, at least 8.5 hours, at least 9 hours, at least 9.5 hours, at least 10 hours, at least 11 hours, at least 13 hours, or at least 15 hours, or at least about 6 hours, at least about 6.5 hours, at least about 7 hours, at least about 7.5 hours, at least about 8 hours, at least about 8.5 hours, at least about 9 hours, at least about 9.5 hours, at least about 10 hours, at least about 11 hours, at least about 13 hours, or at least about 15 hours, longer than the mean time interval between first dose of parturifacient and delivery for subjects testing positive for oncofetal fibronectin. Typically, subjects testing negative for oncofetal fibronectin have a mean time interval between first dose of pre-induction agent and delivery that is 30-90% or about 30-90%, 35-80% or about 35-80%, 40-75% or about 40-75%, 42-70% or about 42-70%, 44-65% or about 44-65%, 45-60% or about 45-60%, 46-58% or about 46-58%, 47-56% or about 47-56%, 48-54% or about 48-54%, 49-52% or about 49-52%, or 50% or about 50%, or 6-20 hours or about 6-20 hours, 6.5-15 hours or about 6.5-15 hours, 7-14 hours or about 7-14 hours, 7.5-13 hours or about 7.5-13 hours, 8-12 hours or about 8-12 hours, 8.5-11 hours or about 8.5-11 hours, 9-10 hours or about 9-10 hours, or 9.5 or about 9.5 hours, longer than the mean time interval between first dose of parturifacient and delivery for subjects testing positive for oncofetal fibronectin.

[0411] Also provided herein, subjects testing positive for oncofetal fibronectin have a mean time interval between oxytocin administration and delivery that is shorter than the mean time interval between oxytocin administration and delivery for subjects testing negative for oncofetal fibronectin or a sample having an amount of oncofetal fibronectin indicating molecule below a threshold. For example, subjects testing positive for oncofetal fibronectin can have a mean time interval between oxytocin administration and delivery that is at least 20%, at least 21%, at least 22%, at least 23%, at least 24%, at least 25%, at least 26%, at least

28%, at least 30%, at least 32%, at least 34%, at least 36%, at least 38%, at least 40%, at least 45%, at least 50%, or at least 60%, or at least about 20%, at least about 21%, at least about 22%, at least about 23%, at least about 24%, at least about 25%, at least about 26%, at least about 28%, at least about 30%, at least about 32%, at least about 34%, at least about 36%, at least about 38%, at least about 40%, at least about 45%, at least about 50%, or at least about 60%, or at least 3 hours, at least 3.2 hours, at least 3.4 hours, at least 3.6 hours, at least 3.8 hours, at least 4 hours, at least 4.2 hours, at least 4.4 hours, at least 4.6 hours, at least 4.8 hours, at least 5 hours, at least 5.5 hours, at least 6 hours, at least 6.5 hours, at least 7 hours, or at least 8 hours, or at least about 3 hours, at least about 3.2 hours, at least about 3.4 hours, at least about 3.6 hours, at least about 3.8 hours, at least about 4 hours, at least about 4.2 hours, at least about 4.4 hours, at least about 4.6 hours, at least about 4.8 hours, at least about 5 hours, at least about 5.5 hours, at least about 6 hours, at least about 6.5 hours, at least about 7 hours, or at least about 8 hours, shorter than the mean time interval between oxytocin administration and delivery for subjects testing negative for oncofetal fibronectin. Typically, subjects testing positive for oncofetal fibronectin have a mean time interval between oxytocin administration and delivery that is 15-60% or about 15-60%, 18-50% or about 18-50%, 20-45% or about 20-45%, 21-40% or about 21-40%, 22-35% or about 22-35%, 23-32% or about 23-32%, 24-30% or about 24-30%, 25-28% or about 25-28%, or 2-10 hours or about 2-10 hours, 3-8 hours or about 3-8 hours, 3.5-7 hours or about 3.5-7 hours, 3.8-6 hours or about 3.8-6 hours, 4-5 hours or about 4-5 hours, 4.1-4.8 hours or about 4.1-4.8 hours, 4.2-4.6 hours or about 4.2-4.6 hours, or 4.4 hours or about 4.4 hours, shorter than the mean time interval between oxytocin administration and delivery for subjects testing negative for oncofetal fibronectin.

[0412] Analogously, subjects testing negative for oncofetal fibronectin have a mean time interval between oxytocin administration and delivery that is longer than the mean time interval between oxytocin administration and delivery for subjects testing positive for oncofetal fibronectin or a sample having an amount of oncofetal fibronectin indicating molecule above a threshold. For example, subjects testing negative for oncofetal fibronectin can have a mean time interval between oxytocin administration and delivery that is at least 30%, at least 31%, at least 32%, at least 33%, at least 34%, at least 35%, at least 36%, at least 37%, at least 38%, at least 40%, at least 42%, at least 45%, at least 50%, at least 55%, at least 60%, or at least 75%, or at least about 30%, at least about 31%, at least about 32%, at least about 33%, at least about 34%, at least about 35%, at least about 36%, at least about 37%, at least about 38%, at least about 40%, at least about 42%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, or at least about 75%, or at least 3 hours, at least 3.2 hours, at least 3.4 hours, at least 3.6 hours, at least 3.8 hours, at least 4 hours, at least 4.2 hours, at least 4.4 hours, at least 4.6 hours, at least 4.8 hours, at least 5 hours, at least 5.5 hours, at least 6 hours, at least 6.5 hours, at least 7 hours, or at least 8 hours, or at least about 3 hours, at least about 3.2 hours, at least about 3.4 hours, at least about 3.6 hours, at least about 3.8 hours, at least about 4 hours, at least about 4.2 hours, at least about 4.4 hours, at least about 4.6 hours, at least about 4.8 hours, at least about 5 hours, at least about 5.5 hours, at least about 6

hours, at least about 6.5 hours, at least about 7 hours, or at least about 8 hours, longer than the mean time interval between oxytocin administration and delivery for subjects testing positive for oncofetal fibronectin. Typically, subjects testing negative for oncofetal fibronectin have a mean time interval between oxytocin administration and delivery that is 20-75% or about 20-75%, 25-60% or about 25-60%, 28-55% or about 28-55%, 30-50% or about 30-50%, 31-45% or about 31-45%, 32-40% or about 32-40%, 33-38% or about 33-38%, 34-36% or about 34-36%, 35% or about 35%, or 2-10 hours or about 2-10 hours, 3-8 hours or about 3-8 hours, 3.5-7 hours or about 3.5-7 hours, 3.8-6 hours or about 3.8-6 hours, 4-5 hours or about 4-5 hours, 4.1-4.8 hours or about 4.1-4.8 hours, 4.2-4.6 hours or about 4.2-4.6 hours, or 4.4 hours or about 4.4 hours, longer than the mean time interval between oxytocin administration and delivery for subjects testing positive for oncofetal fibronectin.

[0413] Also provided herein, subjects testing positive for oncofetal fibronectin are likely to receive fewer pre-induction agent administrations relative to subjects testing negative for oncofetal fibronectin or to a sample having an amount of oncofetal fibronectin indicating molecule below a threshold. For example, subjects testing positive for oncofetal fibronectin can be predicted to receive at least 50%, at least 51%, at least 52%, at least 53%, at least 54%, at least 55%, at least 56%, at least 57%, at least 58%, at least 59%, at least 60%, at least 62%, at least 64%, at least 66%, at least 70%, at least 75%, at least 80%, or at least 90%, or at least about 50%, at least about 51%, at least about 52%, at least about 53%, at least about 54%, at least about 55%, at least about 56%, at least about 57%, at least about 58%, at least about 59%, at least about 60%, at least about 62%, at least about 64%, at least about 66%, at least about 70%, at least about 75%, at least about 80%, or at least about 90%, or at least 0.4, at least 0.45, at least 0.5, at least 0.55, at least 0.6, at least 0.65, at least 0.7, at least 0.8, at least 0.9, at least 1.0, or at least 1.1, or at least about 0.4, at least about 0.45, at least about 0.5, at least about 0.55, at least about 0.6, at least about 0.65, at least about 0.7, at least about 0.8, at least about 0.9, at least about 1.0, or at least about 1.1, fewer pre-induction agent administrations relative to subjects testing negative for oncofetal fibronectin. Typically, subjects testing positive for oncofetal fibronectin are predicted to receive 40-90% or about 40-90%, 45-80% or about 45-80%, 48-70% or about 48-70%, 50-68% or about 50-68%, 52-66% or about 52-66%, 54-64% or about 54-64%, 55-62% or about 55-62%, 56-60% or about 56-60%, or 57-58% or about 57-58%, or 0.3-1.1 or about 0.3-1.1, 0.35-1.0 or about 0.35-1.0, 0.4-0.85 or about 0.4-0.85, 0.45-0.75 or about 0.45-0.75, 0.5-0.7 or about 0.5-0.7, 0.52-0.68 or about 0.52-0.68, 0.54-0.66 or about 0.54-0.66, 0.58-0.62 or about 0.58-0.62, or 0.6 or about 0.6, fewer pre-induction agent administrations relative to subjects testing negative for oncofetal fibronectin.

[0414] Analogously, subjects testing negative for oncofetal fibronectin are likely to receive more pre-induction agent administrations relative to subjects testing positive for oncofetal fibronectin or to a sample having an amount of oncofetal fibronectin indicating molecule above a threshold. For example, subjects testing negative for oncofetal fibronectin can be likely to receive at least 65%, at least 66%, at least 67%, at least 68%, at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 75%, at least

77%, at least 80%, at least 85%, at least 90%, at least 100%, at least 110%, at least 120%, or at least 130%, or at least about 65%, at least about 66%, at least about 67%, at least about 68%, at least about 69%, at least about 70%, at least about 71%, at least about 72%, at least about 73%, at least about 75%, at least about 77%, at least about 80%, at least about 85%, at least about 90%, at least about 100%, at least about 110%, at least about 120%, or at least about 130%, or at least 0.4, at least 0.45, at least 0.5, at least 0.55, at least 0.6, at least 0.65, at least 0.7, at least 0.8, at least 0.9, at least 1.0, or at least 1.1, or at least about 0.4, at least about 0.45, at least about 0.5, at least about 0.55, at least about 0.6, at least about 0.65, at least about 0.7, at least about 0.8, at least about 0.9, at least about 1.0, or at least about 1.1, more pre-induction agent administrations relative to subjects testing positive for oncofetal fibronectin. Typically, subjects testing negative for oncofetal fibronectin are likely to receive 50-150% or about 50-150%, 55-125% or about 55-125%, 60-110% or about 60-110%, 62-100% or about 62-100%, 64-90% or about 64-90%, 66-85% or about 66-85%, 68-80% or about 68-80%, 70-75% or about 70-75%, 71-73% or about 71-73%, or 72% or about 72%, 0.3-1.1 or about 0.3-1.1, 0.35-1.0 or about 0.35-1.0, 0.4-0.9 or about 0.4-0.9, 0.45-0.75 or about 0.45-0.75, 0.5-0.7 or about 0.5-0.7, 0.52-0.68 or about 0.52-0.68, 0.54-0.66 or about 0.54-0.66, 0.58-0.62 or about 0.58-0.62, or 0.6 or about 0.6, more pre-induction agent administrations relative to subjects testing positive for oncofetal fibronectin.

[0415] In one embodiment, for purposes of use in conjunction with induction methods, a subject is considered positive for oncofetal fibronectin when the presence of an oncofetal fibronectin indicating molecule in a sample is equal to or above one or more threshold levels. As one skilled in the art will recognize, a threshold level can vary according to the type of sample measured and the selected stringency of the test. In one example, a threshold level for a buffer-treated cervicovaginal sample assayed using a test strip can be 50 ng/mL. In another example, a threshold level for a buffer-treated cervicovaginal sample assayed using a test strip can be 150 ng/mL.

[0416] In another embodiment, multi-tiered thresholds can be applied to the oncofetal fibronectin indicating molecule measurement, where multi-tiered thresholds include two or more threshold levels, where each larger threshold level indicates a further increased likelihood of success of induction, increased likelihood of vaginal delivery upon induction, likely decreased time interval between initiating induction and delivery, likely decreased time interval between administering a pre-induction agent and delivery, likely decreased time interval between administering oxytocin and delivery, increased likelihood of delivering within 24 hours of induction, increased likelihood of delivering within 48 hours of induction and decreased likelihood of more than one administration of pre-induction agent to the subject. An exemplary multi-tiered threshold is a two-tiered threshold where the lower threshold is 50 ng/mL and the higher threshold is 150 ng/mL for buffer-treated samples. An exemplary multi-tiered threshold is a two-tiered threshold where the lower threshold is 500 ng/mL and the higher threshold is 1500 ng/mL for untreated samples.

[0417] In accordance with the methods that include multi-tiered thresholds, methods are provided herein for classifying the likely outcome of induction, where the methods

include measuring the amount of an oncofetal fibronectin indicating molecule in a sample from a subject and determining which, if any, multi-tiered threshold levels the sample is equal to or greater than and classifying the likely outcome of induction, where each increasing amount in the multi-tiered thresholds classifies the outcome as: increased likelihood of success of induction, increased likelihood of vaginal delivery upon induction, likely decreased time interval between initiating induction and delivery, likely decreased time interval between administering a parturifacient and delivery, likely decreased time interval between administering oxytocin and delivery, increased likelihood of delivering within 24 hours of induction, increased likelihood of delivering within 48 hours of induction and decreased likelihood of more than one administration of pre-induction agent to the subject, or combinations thereof.

[0418] In another embodiment, methods for detecting an oncofetal fibronectin indicating molecule in conjunction with induction of labor can be further coupled with one or more additional indicators or markers for induction outcome. Thus, provided herein are methods of determining the success of induction, where the methods include detecting an oncofetal fibronectin indicating molecule in a sample and determining one or more additional indicators or markers of induction outcome, where presence (or level above threshold) of an oncofetal fibronectin indicating molecule and one or more indicators or markers of induction outcome can indicate, relative to absence (or level below threshold) of the oncofetal fibronectin indicating molecule and/or a positive result for one or more additional indicators or markers of induction outcome, an increased likelihood of success of induction, increased likelihood of vaginal delivery upon induction, likely decreased time interval between initiating induction and delivery, likely decreased time interval between administering a parturifacient and delivery, likely decreased time interval between administering oxytocin and delivery, increased likelihood of delivering within 24 hours of induction, increased likelihood of delivering within 48 hours of induction and decreased likelihood of more than one administration of pre-induction agent to the subject, or combinations thereof.

[0419] A variety of indicators or markers of induction outcome are provided herein or are otherwise known in the art, and typically include measurements or observations of the pregnant subject or the fetus(es), or medical history of the pregnant subject. Exemplary indicators include, but are not limited to, cervical length, Bishop score, effacement, parity (i.e., previous vaginal delivery by the subject), cervical dilation, gestational age, body mass index (BMI), station, consistency, transvaginal ultrasound, and digital examination. In one example, an indicator is parity. In another example, the indicator is BMI.

[0420] i. Induction Methods and Compounds

[0421] Induction can be performed by any of a variety of methods known in the art, including administering a parturifacient and performing an induction procedure. A variety of induction procedures are known in the art, including, but not limited to, balloon catheterization such as foley balloon catheterization or Atad balloon catheterization, amniotic membrane stripping, extra-amniotic saline infusion, amniotomy, or nipple stimulation. In addition, a parturifacient can be administered, where the parturifacient can be

any of a variety of compounds or compositions known in the art for pre-inducement, cervical ripening, or inducement. Exemplary parturifacients include, but are not limited to, prostaglandins such as PGE1 (misoprostol) and PGE2 (dinoprostone), oxytocic hormones such as oxytocin and steroids such as RU486 (mifepristone).

[0422] The methods provided herein for predicting outcome of induction of a subject or identifying a subject with higher likelihood of successful induction can apply to any of the above induction methods and compounds and combinations thereof. In one example, presence (or level above threshold) of an oncofetal fibronectin indicating molecule in a sample from a subject can indicate a decreased time interval after induction with oxytocin, relative to subjects negative (or level below threshold) for the oncofetal fibronectin indicating molecule.

[0423] ii. Post-Induction Measurements

[0424] In another embodiment, the success of induction can be monitored by measuring the amount of an oncofetal fibronectin indicating molecule. A pregnant mother can be induced into labor by various methods including administering a parturifacient such as an oxytocic hormone or a prostaglandin. After induction, the amount of an oncofetal fibronectin indicating molecule can be measured and an amount of an oncofetal fibronectin indicating molecule that is above a threshold level can indicate that the induction was effective and that delivery by the pregnant mother is imminent. Thus, also provided herein is a method for monitoring the effectiveness of induction of a pregnant woman by monitoring the post-induction amount of an oncofetal fibronectin indicating molecule in a pregnant woman. For example, a post-induction measurement of an oncofetal fibronectin indicating molecule that is above a threshold level can indicate that the induction of the pregnant woman is effective and the pregnant woman will soon deliver. Typically imminent delivery or likelihood that a woman will soon deliver, when used in the context of post-induction, refers to an indication that a pregnant woman will deliver within 48 hours or about 48 hours after induction, or within 48 hours or about 48 hours after a positive oncofetal fibronectin measurement. A post-induction measurement of an oncofetal fibronectin indicating molecule that is below a threshold level can indicate that the induction of the pregnant woman is ineffective and the pregnant woman will not soon deliver. In response to a negative post-induction oncofetal fibronectin measurement, a decision can be made whether to administer an additional induction procedure in order to accomplish vaginal delivery by the pregnant woman, or to instead choose to pursue childbirth by cesarean section. Thus, provided herein is a method for determining treatment method after induction of a subject, the method including measuring the amount of an oncofetal fibronectin indicating molecule in a sample from an induced subject and if the amount of the oncofetal fibronectin indicating molecule is at or above a threshold level, determining that additional induction is not required for vaginal delivery.

[0425] Also provided herein is a method for determining treatment method after induction of a subject, the method including measuring the amount of an oncofetal fibronectin indicating molecule in a sample from an induced subject and if the amount of the oncofetal fibronectin indicating molecule is below a threshold level, determining to administer

one or more additional administrations of induction, or determining to proceed with childbirth by cesarean section.

[0426] Post-induction oncofetal fibronectin indicating molecule measurements can be performed on any of the samples and by any of the methods used in pre-induction methods provided herein. Additionally, post-induction oncofetal fibronectin indicating molecule measurements can be used in conjunction with multi-tiered thresholds as provided herein in regard to pre-induction measurements and also can be used in conjunction with one or more additional indicators of induction outcome, as provided herein in regard to pre-induction measurements.

[0427] e. Conceptus Indications

[0428] Provided herein is a method for selecting a conceptus for implantation. In the method, a conceptus sample, such as from a well or other receptacle containing the conceptus or from the conceptus itself is tested for an oncofetal fibronectin indicating molecule. A conceptus with a sample having an amount greater than a pre-determined level, or an amount greater than the amount for other concepti, or a rate of increase of such amount that is greater than a pre-determined level or compared to other concepti in a group can be identified as favorable for implantation and/or selected for implantation.

[0429] The methods, combinations, compositions and kits provided herein can be used for a variety of conceptus indications, where the methods include determining the presence or amount of an oncofetal fibronectin indicating molecule, where the conceptus is identified and/or selected according to the amount of oncofetal fibronectin indicating molecule produced by the conceptus. For example, a conceptus identified as producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule at or above a threshold level can be identified as a conceptus with increased competence for or increased likelihood of successfully implanting, relative to a conceptus not producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule below a threshold level. In another example, a conceptus identified as producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule at or above a threshold level can be identified as a conceptus with increased competence for or increased likelihood of successful subsequent fetal development once transferred, relative to a conceptus not producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule below a threshold level. In another example, a conceptus identified as producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule at or above a threshold level can be selected for use in implantation. In another example, a conceptus identified as producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule at or above a threshold level can be selected for use in a cell culture. In another example, a conceptus identified as producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule at or above a threshold level can be selected for use as a stem cell. In another example, a conceptus identified as not producing oncofetal fibronectin indicating molecule or pro-

ducing an amount of oncofetal fibronectin indicating molecule below a threshold level can be identified as a conceptus with decreased competence for or decreased likelihood of successfully implanting, relative to a conceptus producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule at or above a threshold level. In another example, a conceptus identified as not producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule below a threshold level can be identified as a conceptus with decreased competence for or decreased likelihood of successful subsequent fetal development once transferred, relative to a conceptus producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule at or above a threshold level. In another example, a conceptus identified as not producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule below a threshold level can be selected to not use for implantation. In another example, a conceptus identified as not producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule below a threshold level can be selected for use in cell culture. In another example, a conceptus identified as not producing oncofetal fibronectin indicating molecule or producing an amount of oncofetal fibronectin indicating molecule below a threshold level can be selected for use as a stem cell.

[0430] Following penetration of the ovum by a sperm, fertilization is completed and the male and female pronuclei fuse, forming a zygote. The zygote undergoes rapid cell divisions and begins to develop. Approximately 4-5 days post-fertilization, the cells have developed into a ball of cells known as a morula. Further development at approximately days 5-6, the cells develop into a blastocyst. As the blastocyst develops further, distinct cell layers are formed. The inner cell mass gives rise to the embryo proper (also known as an embryoblast). The outer cell mass, also known as the trophoblast, develops as a layer of epithelium cells surrounding the embryo.

[0431] A conceptus attaches to the uterine wall, generally 6 to 7 days post-fertilization. The trophoblast, a layer of epithelial cells at the outer surface of the blastocyst, participates in the attachment of the blastocyst to the endometrial layer of cells in the uterus. Cells at the embryonic pole of the trophoblast layer differentiate to produce syncytiotrophoblasts, which begin the process of uterine implantation.

[0432] The trophoblast layer is the primary source for subsequent development of extraembryonic membranes including placental tissue, amnion and umbilical cord. The trophoblast cells of the conceptus synthesize and secrete oncofetal fibronectin protein. Oncofetal fibronectin is produced in early stage concepti including early stage embryos, and plays a role in the attachment of the conceptus to the uterine wall and/or vascularization of the attached conceptus. Oncofetal fibronectin is present at placental attachments sites at the placental-uterine junction.

[0433] The methods provided herein can be used to measure the presence and or amount of an oncofetal fibronectin indicating molecule produced by a conceptus, at one or more time points, and thereby to identify a conceptus with increased likelihood of successful uterine implantation.

Accordingly, as provided herein, a conceptus that produces an oncofetal fibronectin indicating molecule has an increased likelihood of successful uterine implantation relative to a conceptus that does not produce the oncofetal fibronectin indicating molecule. Also, as provided herein, a conceptus that produces higher amounts of an oncofetal fibronectin indicating molecule relative to a conceptus that produces lower amounts of the oncofetal fibronectin indicating molecule, has an increased likelihood of successful uterine implantation. Also, as provided herein, a conceptus that produces an amount of an oncofetal fibronectin indicating molecule at or above a threshold level has an increased likelihood of successful uterine implantation relative to a conceptus that produces an amount of the oncofetal fibronectin indicating molecule below a threshold level. Also, as provided herein, a conceptus that produces a more rapidly increasing amount of an oncofetal fibronectin indicating molecule relative to a conceptus that produces a less rapidly increasing or a decreasing amount of the oncofetal fibronectin indicating molecule, has an increased likelihood of successful uterine implantation.

[0434] Similarly, as provided herein, a conceptus that does not produce an oncofetal fibronectin indicating molecule has a decreased likelihood of successful uterine implantation relative to a conceptus that produces the oncofetal fibronectin indicating molecule. Also, as provided herein, a conceptus that produces lower amounts of an oncofetal fibronectin indicating molecule has a decreased likelihood of successful uterine implantation relative to a conceptus that produces higher amounts of the oncofetal fibronectin indicating molecule. Also, as provided herein, a conceptus that produces an amount of an oncofetal fibronectin indicating molecule below a threshold level has a decreased likelihood of successful uterine implantation relative to a conceptus that produces an amount of the oncofetal fibronectin indicating molecule at or above a threshold level. Also, as provided herein, a conceptus that produces a less rapidly increasing or decreasing amount of an oncofetal fibronectin indicating molecule has a decreased likelihood of successful uterine implantation relative to a conceptus that produces a more rapidly increasing amount of the oncofetal fibronectin indicating molecule. The methods provided herein can be used to measure the presence and or amount of an oncofetal fibronectin indicating molecule produced by a conceptus, at one or more time points, and thereby to identify a conceptus with decreased likelihood of successful uterine implantation.

[0435] In some cases a conceptus can be categorized as oncofetal fibronectin positive when any oncofetal fibronectin indicating molecule is detected in a conceptus sample. In other cases, a conceptus can be categorized according to the amount of oncofetal fibronectin indicating molecule present in a sample. In other cases, a conceptus can be categorized as oncofetal fibronectin positive when the presence of an oncofetal fibronectin indicating molecule in a conceptus sample is equal to or above one or more threshold levels. In one example, a threshold level of an oncofetal fibronectin indicating molecule such as oncofetal fibronectin protein in conceptus culture medium assayed using a test strip can be 50 ng/mL. In another example, a threshold level of an oncofetal fibronectin indicating molecule such as oncofetal fibronectin protein in conceptus culture medium assayed using a test strip can be 150 ng/mL. Exemplary threshold values for categorizing a conceptus include 50 ng/ml, 150 ng/ml, 200 ng/ml, 300 ng/ml, 500 ng/ml, 750 ng/ml and

1000 ng/ml, or about 50 ng/ml, about 150 ng/ml, about 200 ng/ml, about 300 ng/ml, about 500 ng/ml, about 750 ng/ml and about 1000 ng/ml.

[0436] i. Detection of Oncofetal Fibronectin Production by a Conceptus

[0437] Any of a variety of methods provided herein or otherwise known in the art for detecting an oncofetal fibronectin indicating molecule in a sample can be used for detecting an oncofetal fibronectin indicating molecule produced by a conceptus, including, but not limited to, dot blot analysis, western blot analysis, northern blot analysis, southern blot analysis, RT-PCR methods, mass spectrometric methods, sandwich assays such as test strip-based sandwich assays and ELISA methods. For example, concepti (e.g., zygotes and embryos) can be placed in culture medium individually.

[0438] An oncofetal fibronectin indicating molecule can be detected in any of a variety of conceptus samples. Conceptus samples include conceptus extracts, samples from outside of the conceptus, such as culture medium, cell and tissue extracts, and cells, where one or more cells is removed from a conceptus, leaving the remainder of the conceptus competent for subsequent culture, implantation and/or development. The samples can be analyzed neat, or can be reagent-treated and/or fractionated prior to detection of an oncofetal fibronectin indicating molecule. In one example, an oncofetal fibronectin indicating molecule can be detected in vitro. In vitro detection includes the detection of an oncofetal fibronectin indicating molecule in isolated cells and tissues and cultured cells and tissues, including an oncofetal fibronectin indicating molecule secreted by such cells and tissues. Cells and tissues for detection include, but are not limited to, any and all stages of a conceptus, including zygotes, morulas, blastocysts, embryoblasts, embryo, and placenta, cells and cells layers of a conceptus, including but not limited to, cytotrophoblasts, trophoblasts, syncytiotrophoblasts, hypoblast and epiblast.

[0439] In some embodiments, an oncofetal fibronectin indicating molecule produced by a conceptus can be measured by measuring oncofetal fibronectin indicating molecule outside of the conceptus. For example, an oncofetal fibronectin indicating molecule can be present in the culture medium that contains the conceptus. An oncofetal fibronectin indicating molecule in medium that contains the conceptus can include an oncofetal fibronectin indicating molecule secreted by the conceptus. In one example, the presence and/or amount of an oncofetal fibronectin indicating molecule produced and/or secreted by a conceptus can be measured by detecting an oncofetal fibronectin indicating molecule in the culture medium that contains the conceptus. Detection of an oncofetal fibronectin indicating molecule in culture medium can include the comparison to controls including culture medium without any added embryos or zygotes, as will be apparent to one skilled in the art.

[0440] In one embodiment, the medium for such oncofetal fibronectin indicating molecule detection methods contains a single conceptus. In such methods, measurement of an oncofetal fibronectin indicating molecule in the medium can indicate the production of the oncofetal fibronectin indicating molecule by the conceptus.

[0441] In another embodiment, the medium for such oncofetal fibronectin indicating molecule detection methods

contains two or more concepti. As provided herein, presence of an oncofetal fibronectin indicating molecule in culture medium can stimulate production of oncofetal fibronectin in a conceptus. Accordingly, presence of a conceptus in medium containing two or more concepti can indicate that the concepti are stimulated to produce oncofetal fibronectin and, therefore, can be identified as producing oncofetal fibronectin.

[0442] In other embodiments, an oncofetal fibronectin indicating molecule produced by a conceptus can be measured by measuring an oncofetal fibronectin indicating molecule present in a cell of a conceptus. For example, the oncofetal fibronectin indicating molecule present in a cell extract can be measured. A cell extract can be collected by removing a portion of cellular fluid such as cytoplasm using, for example, a syringe, or by cell harvesting. Cell harvesting can be performed, for example, by removing one or more cells from a conceptus, while leaving the conceptus sufficiently intact and competent for subsequent development and implantation. Oncofetal fibronectin indicating molecule detection in the removed cell can be performed immediately after removing the cell (e.g., by harvesting the cell), or after culturing the removed cell, where oncofetal fibronectin indicating molecule detection can be performed at one or more time points.

[0443] In other embodiments, conceptus extracts can be used for detecting an oncofetal fibronectin indicating molecule. A conceptus can contain one or more cavities, including, but not limited to, the blastocoel. An oncofetal fibronectin indicating molecule can be present in these cavities. A sample of the liquid in a conceptus cavity can be collected using known methods, for example, by extracting with a microsyringe.

[0444] Detection includes, but is not limited to, measurement of the presence, amount, rate of production and/or rate of secretion of an oncofetal fibronectin indicating molecule. An amount or rate can be compared to a threshold amount or can be compared to one or more amounts or rates from other concepti. For example, amounts or rates of oncofetal fibronectin indicating molecule production can be compared for two or more concepti, and the concepti can be classified from highest to lowest in amount oncofetal fibronectin indicating molecule produced or rate of increase of oncofetal fibronectin indicating molecule production.

[0445] A sample for which an oncofetal fibronectin indicating molecule is to be detected can be collected at any time post-isolation of the gametes and includes detection immediately after fertilization, and times following fertilization up to and including the time of implantation of a conceptus. For example, oncofetal fibronectin indicating molecule detection can be performed immediately following fertilization, and at 1, 2, 3, 4, 5, 6, 7, 8, 16, 24, 30, 36, 40, 48, 60, 72, 84, 96, 108, 120, 132, 144, and 168 hours and at any intermediate times post-fertilization. Detection of an oncofetal fibronectin indicating molecule also can be performed at times prior to transfer to a female including, but not limited to, the day of transfer, and 1 day, 2 days, 3 days, 4 days and 5 days pre-transfer. Oncofetal fibronectin indicating molecule detection can be performed at one or more of any such time points.

[0446] ii. Assisted Reproduction Technology Related Uses

[0447] The methods, compositions and kits provided herein for detecting an oncofetal fibronectin indicating mol-

ecule can be used in Assisted Reproduction Technologies (ART), including as predictive tests and for indicating success of ART. ART includes but is not limited to in vitro fertilization (IVF), zygote intra-fallopian transfer (ZIFT), gamete intra-fallopian transfer (GIFT), blastocyst transfer, in vitro maturation of oocytes coupled with IVF, and frozen embryo transfer (FET); these methods are known in the art. In these procedures gametes (ova and sperm) are collected, manipulated in vitro and then one or more gametes or cells derived from gametes are implanted into the female. One example of ART, in vitro fertilization (IVF), includes the collection of ova and sperm, fertilization of the ova in vitro, culturing the fertilized cells and allowing the fertilized cells to divide and develop in vitro, and then transferring one or more cultured fertilized cells (typically embryos) into the uterus. Another example of ART, zygote intra-fallopian transfer (ZIFT) includes collection of ova and sperm, fertilization of the ova in vitro, and transfer of a zygote (or more than one zygote) into the fallopian tubes. Following transfer of the conceptus (e.g., zygote or embryo) to the female, the conceptus may or may not subsequently attach to the uterine wall, and a conceptus that attaches to the uterine wall can further develop. The ability of the conceptus to successfully implant in the uterine wall plays a key role in the success of these procedures, because if the conceptus does not successfully implant, the development will arrest and the pregnancy will not go to term.

[0448] Accordingly, provided herein are methods of performing ART by transferring to a female a conceptus identified using the methods provided herein as having an increased competence for or increased likelihood of successful implantation in the uterus, and/or improved likelihood of successful subsequent fetal development once transferred.

[0449] Development of the conceptus in in vitro fertilization and other ART procedures parallels in vivo fertilization in development, with the exception that some of the stages occur in vitro prior to implantation and the remainder are completed post-implantation. In ART procedures, the conceptus is transferred to the uterus generally at approximately 2 to 6 days post-fertilization, usually at approximately 3 to 5 days post-fertilization. The conceptus also can be transferred into the fallopian tubes, generally 1, 2 and/or 3 days post-fertilization. Transfer of conceptus includes transfer of a 2 cell, 2-4 cell, 4 cell, 4-8 cell and greater than 8 cell stage conceptus and can include transfer at the blastocyst stage. Transfer includes transfer of one or more concepti, usually 1-5, 2-5, 3-5 and 2-3 conceptus are transferred. In some cases, such as single embryo transfer, only one conceptus is transferred to the female. ART also includes frozen embryo transfer where the embryos, following fertilization and optionally a period of cell culture, are frozen for later use, e.g., implantation into a female at a later time. Such embryos can be stored indefinitely under frozen conditions, such as storage in liquid nitrogen.

[0450] iii. Post-Measurement Steps

[0451] As provided herein, production of an oncofetal fibronectin indicating molecule can be measured. Oncofetal fibronectin indicating molecule measurements can be used to characterize a conceptus. For example, a conceptus for which an oncofetal fibronectin indicating molecule is present, is at or above a threshold level or has a higher

increasing rate of production can be characterized as having an increased competence for or increased likelihood of successful implantation in the uterus, and/or improved likelihood of successful subsequent fetal development once transferred, relative to a conceptus for which the oncofetal fibronectin indicating molecule is absent, is below a threshold level or has a decreasing rate of production.

[0452] a. Increasing Oncofetal Fibronectin Production

[0453] Also provided herein are methods for increasing production of oncofetal fibronectin in a conceptus. As provided herein, a conceptus that produces oncofetal fibronectin at a sufficiently low level or rate, is not typically selected for implantation. Methods provided herein can be used to select a conceptus producing an oncofetal fibronectin indicating molecule below a threshold level or rate for implantation, and treating the conceptus with one or more methods for increasing production of an oncofetal fibronectin indicating molecule in a conceptus. In one example, a conceptus can be monitored on one or more occasions after such treatment for production of an oncofetal fibronectin indicating molecule, and upon measurement of an amount of rate of production above a threshold level or rate, the conceptus can be as suitable for implantation.

[0454] Oncofetal fibronectin synthesis can be increased in amount and/or rate of synthesis by external stimuli. Synthesis of oncofetal fibronectin can increase over time in culture. For example, some isolated human trophoblasts secrete barely detectable oncofetal fibronectin in culture shortly after isolation. The synthesis of oncofetal fibronectin increases over time of culturing in vitro. A variety of constituents of the culture medium can stimulate oncofetal fibronectin production, including but not limited to serum, such as human and other mammalian serum, and transforming growth factors, including TGF- α and TGF- β . Oncofetal fibronectin synthesis also can be stimulated by cell attachment including, but not limited to, incubation of cells with extracellular matrix, synthetic matrices, or plastic.

[0455] b. Identify Conceptus Based on Oncofetal Fibronectin Production

[0456] In the methods provided herein, detection of an oncofetal fibronectin indicating molecule can be coupled with classification of a conceptus according to the oncofetal fibronectin indicating molecule production by the conceptus. Presence, amount or rate of oncofetal fibronectin indicating molecule production can indicate the competence of a conceptus for implantation, or the likelihood of a conceptus successfully implanting or the likelihood of successful subsequent fetal development once transferred. Accordingly, the methods provided herein can be used to classify a conceptus as having an increased competence for, or increased likelihood of, successful implantation in the uterus, and/or improved likelihood of successful subsequent fetal development once transferred, where a conceptus with the presence of, an increased amount of, or a higher rate of increase of oncofetal fibronectin indicating molecule has an increased competence for, or increased likelihood of, successful implantation in the uterus, and/or improved likelihood of successful subsequent fetal development once transferred, relative to a conceptus with an absence of, a decreased amount of, or a lower rate of increase or a decrease of oncofetal fibronectin indicating molecule. Similarly, methods provided herein can be used to identify a

conceptus as suitable for implantation, where a conceptus in which an oncofetal fibronectin indicating molecule is present or is present at or above a threshold level or is increasing in production can be identified as having a sufficiently high rate competence for, or likelihood of, successful implantation in the uterus, and/or likelihood of successful subsequent fetal development once transferred, and therefore as suitable for implantation.

[0457] i. Selection of a Conceptus

[0458] Detection of an oncofetal fibronectin indicating molecule can be used to predict and to improve conceptus implantation and viability in ART procedures including, but not limited to, in vitro fertilization procedures, zygote intrafallopian transfer and frozen embryo transfer. As provided herein, the presence, amount, rate of production and/or rate of secretion of an oncofetal fibronectin indicating molecule can be used as an indicator for the ability of a conceptus to implant in the uterus and/or for such implantation to be successfully developed and maintained. A conceptus having an increased competence for or increased likelihood of successful implantation in the uterus, and/or improved likelihood of successful subsequent fetal development once transferred, can be selected for transfer to a female (e.g., transfer to the uterus or fallopian tubes).

[0459] In one embodiment, detection of an oncofetal fibronectin indicating molecule is used to select one or more concepti (e.g., embryos and zygotes) for implantation. For example, an ovum can be collected and fertilized in vitro, to form a conceptus. Following fertilization, an oncofetal fibronectin indicating molecule can be detected in one or more samples corresponding to one or more concepti and the presence or amount of oncofetal fibronectin indicating molecule in the samples can be used to select one or more concepti having an increased competence for or increased likelihood of successful implantation in the uterus, and/or improved likelihood of successful subsequent fetal development once transferred.

[0460] In yet another embodiment, detection of an oncofetal fibronectin indicating molecule can be used to select a conceptus for frozen storage. In ART procedures, often more concepti are produced in vitro than will be transferred to a female. Concepti that are not transferred can be stored indefinitely in frozen storage such as liquid nitrogen tanks. Concepti can be selected for frozen storage, for example, based on detection of an amount of an oncofetal fibronectin indicating molecule, a threshold level of an oncofetal fibronectin indicating molecule, rate of production and/or secretion of an oncofetal fibronectin indicating molecule, and/or detection coupled with selection in comparison to production of an oncofetal fibronectin indicating molecule by other concepti.

[0461] ii. Criteria for Selection

[0462] In one aspect of the embodiment, a conceptus with the highest oncofetal fibronectin indicating molecule level and/or concepti (more than one conceptus) with the highest levels of oncofetal fibronectin indicating molecule relative to one or more other concepti are selected for transfer. In another aspect, one or more concepti with the lowest oncofetal fibronectin indicating molecule levels are discarded and/or are not selected for transfer. In yet another aspect, concepti with the lowest oncofetal fibronectin indicating molecule levels are stimulated for oncofetal fibronectin production before transfer.

[0463] In another embodiment, a threshold level of detectable oncofetal fibronectin indicating molecule is used to select concepti (e.g., embryos and zygotes) for transfer. In one aspect of the embodiment the presence of an amount equal to or above a threshold level of an oncofetal fibronectin indicating molecule indicates a conceptus to select for transfer. In another aspect of the embodiment, the presence of an amount less than a threshold level is used to discard a conceptus and/or identify conceptus not preferred for transfer. In yet another embodiment, the presence of an amount less than a threshold level is used to select a conceptus for stimulation of oncofetal fibronectin production prior to transfer. In such methods, a positive result can be the presence of any amount of an oncofetal fibronectin indicating molecule equal to or above a threshold level, and a negative result can be any amount of an oncofetal fibronectin indicating molecule below a threshold level.

[0464] In another embodiment, the rate of increase of oncofetal fibronectin indicating molecule production is detected and used to select a conceptus for transfer. For example, an oncofetal fibronectin indicating molecule can be detected at two or more time points post-fertilization and a rate of oncofetal fibronectin indicating molecule synthesis and/or oncofetal fibronectin indicating molecule secretion can be determined, based on the change of oncofetal fibronectin indicating molecule levels over time. In one aspect of the embodiment, one or more concepti are chosen that exhibit the highest rate of oncofetal fibronectin indicating molecule production relative to other concepti. In another aspect, one or more concepti are chosen that exhibit a rate equal to or above a chosen threshold rate of oncofetal fibronectin indicating molecule production. In another aspect of the embodiment, concepti that exhibit a lower rate of oncofetal fibronectin indicating molecule relative to other concepti or as compared to a threshold rate are discarded and/or are selected as concepti not for transfer.

[0465] c. Other Markers Used in Conjunction with Oncofetal Fibronectin

[0466] In any of the embodiments herein, oncofetal fibronectin indicating molecule detection can be coupled with other testing procedures which monitor and/or select criteria in the conceptus and/or female recipient.

[0467] i. Conceptus Markers

[0468] For example, oncofetal fibronectin indicating molecule detection can be coupled with pre-implantation characterization and/or diagnosis of the conceptus. Such characterization can be based on genetic characteristics, gene expression, and/or morphology of the conceptus. Characterization and/or diagnosis can be performed at any time after fertilization until implantation, and is typically performed between 1 and 7 days after fertilization, for example from 2-3 days after fertilization.

[0469] In one example, genetic characterization and/or diagnosis can be conducted. In such procedures, one or more cells, generally a single cell, is removed from the conceptus and tested for the absence and/or presence of genetic markers. In one embodiment, a conceptus can be selected for transfer based on detection of an oncofetal fibronectin indicating molecule and detection of at least one genetic marker. In another example of coupled detection, oncofetal fibronectin (onfFN) detection is coupled with detection of

additional implantation competence markers such as additional extracellular matrix proteins. In another example of coupled detection, onfFN detection is coupled with detection of at least one quality marker including, but not limited to, expression of oxidative stress genes (e.g., MnSOD, CuZnSOD, SOX), apoptosis genes (e.g., Bax), maternal recognition of pregnancy genes (e.g., INF-tau), genes related to communication through gap junctions (e.g., Cx31 and Cx43), and/or differentiation and implantation genes (e.g., LIF and LR-beta). A variety of markers are known in the art, as exemplified in Reese et al., *J. Biol. Chem.*, 276:44137-44145 (2001); and Yoshioka et al., *Biochem. Biophys. Res. Commun.* 272:531-538 (2000).

[0470] In another embodiment, oncofetal fibronectin indicating molecule detection can be combined with morphological characterization and/or diagnosis of a conceptus. Morphological features of a conceptus can indicate the likelihood of successful implantation and development to pregnancy of a conceptus. A variety of morphological features characteristic of concepti with an increased likelihood of successful implantation and development to pregnancy are known in the art, and include, but are not limited to, cell number, degree of fragmentation, cell regularity, symmetry, pronuclear morphology, follicle size, follicular fluid volume, multi-nucleation, presence of vacuoles, and granularity.

[0471] ii. Maternal Markers

[0472] In any of the embodiments herein, oncofetal fibronectin indicating molecule detection can be coupled with other testing procedures which monitor and/or select criteria in the female recipient. For example, a female recipient can be tested for markers correlative with receptivity to successful transfer and subsequent implantation of a conceptus. Such markers can include detection of mucin glycoproteins (e.g., MUC1) and heparin sulfate-binding proteins. Additional markers for use include phenotypes of follicles that can contribute to the competence and viability of the oocyte to participate in successful fertilization and conceptus development, including, but not limited to, expression of 11 β -hydroxysteroid dehydrogenase by granulosa cells, adhesion and proliferation of cumulus cells, steroidogenic activity of cumulus and perifollicular vascularity and vascular epithelial growth factors bound to granulosa and cumulus cells.

[0473] iv. Cell Culture and Stem Cell Uses

[0474] Methods and probes for detection of an oncofetal fibronectin indicating molecule can be used to select cells and tissues for non-implantation uses, including for use in cell culture and as stem cells. In one example, a cell or conceptus not selected for implantation can be selected for use in further in vitro culture as a source of embryonic stem cells. Such cells are useful in the research and treatment of diseases and conditions, including, but not limited to immunological and neurodegenerative diseases such as Parkinson's disease, lupus, diabetes, stroke, rheumatoid arthritis, heart trauma, and in cell replacement therapies and tissue regeneration.

[0475] In one embodiment, detection of an oncofetal fibronectin indicating molecule is used to select a conceptus not competent or less competent for transfer to a female. A conceptus further can be selected for use in cell culture or as stem cells, for example, based on their viability in culture.

Detection of an oncofetal fibronectin indicating molecule can include any of the detection methods, assays and selections herein, including selections based on detection of an amount, a threshold level, a rate of synthesis and/or a rate of secretion of an oncofetal fibronectin indicating molecule.

[0476] 2. Indicator of Cancer

[0477] As provided herein, the presence or an amount of an oncofetal fibronectin indicating molecule in body tissue or fluid sample can be associated with tumor, cancer, metastasis or neoplasia. An oncofetal fibronectin indicating molecule can serve as a marker for the presence of, the risk of developing, the progression or regression of, the recurrence of, aggressiveness of, or treatment of, tumor, cancer, metastasis or neoplasia in a subject. The methods provided herein can be used to determine the presence or absence, or the amount of an oncofetal fibronectin indicating molecule in a sample, and also can be used to monitor levels of an oncofetal fibronectin indicating molecule in a subject over time. Presence or an elevated amount above a threshold of an oncofetal fibronectin indicating molecule can indicate the presence of, an increased risk of developing, the progression of, the recurrence of, increased aggressiveness of, or ineffectiveness of treatment of, tumor, cancer, metastasis or neoplasia in a subject. Absence or a lower amount below a threshold of an oncofetal fibronectin indicating molecule can indicate the absence of, a decreased risk of developing, the regression of, the lack of recurrence of, decreased aggressiveness of, or effectiveness of treatment of, tumor, cancer, metastasis or neoplasia in a subject.

[0478] An oncofetal fibronectin indicating molecule can be used as a biological marker for a variety of cancer (e.g., neoplastic) diseases, including, but not limited to, carcinomas of the lung, breast, ovary, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, prostate, thyroid, pituitary, eye, brain, oral, skin, head and neck cancer, lymphoma, leukemia, squamous cell carcinomas, adenocarcinomas, small cell carcinomas, melanomas, gliomas, sarcomas and neuroblastomas. An oncofetal fibronectin indicating molecule indicating various cancerous (e.g., neoplastic) conditions can be detected in any of a variety of body tissue and fluid samples, including those provided herein or are otherwise known in the art; and any method for collecting a sample can be used that is appropriate for the tissue or organ under examination, as will be understood by one of skill in the art. Any oncofetal fibronectin indicating molecule detection method provided herein or otherwise known in the art can be used to detect the presence or amount of an oncofetal fibronectin indicating molecule in a sample. Provided herein below as non-exclusive examples of cancer-related oncofetal fibronectin indicating molecule detection methods are descriptions of detection of an oncofetal fibronectin indicating molecule in blood, plasma, serum, urine, lavage, sputum, tissue, aspirate, stool, or swab samples that can be used to indicate the presence, risk of, development of, likelihood of recurrence of, progression or regression of, aggressiveness of, or efficacy of treatment of, bladder, breast, cervical, ovarian, prostate, lung, or colorectal cancers.

[0479] The methods provided herein can include collecting a sample, measuring for the presence of an oncofetal fibronectin indicating molecule in a sample and/or measur-

ing the amount of an oncofetal fibronectin indicating molecule in the sample, optionally characterizing the oncofetal fibronectin indicating molecule in the sample. Presence of an oncofetal fibronectin indicating molecule, or an amount of oncofetal fibronectin indicating molecule equal to or above one or more thresholds can characterize the sample as oncofetal fibronectin positive, or as falling within a particular group, according to the highest threshold equal to or less than the measured oncofetal fibronectin indicating molecule amount. Absence of an oncofetal fibronectin indicating molecule, or an amount of oncofetal fibronectin indicating molecule below one or more thresholds can characterize the sample as oncofetal fibronectin negative, or as falling within a particular group, according to the lowest threshold greater than the measured oncofetal fibronectin indicating molecule amount.

[0480] A sample collected can be taken from any source, as provided herein or otherwise known in the art. Exemplary sources for samples include, but are not limited to, a tissue sample, swab of a region suspected of being cancerous, a lavage sample, a blood sample, a plasma sample, a serum sample, an interstitial fluid sample, a lymph sample, a lymphatic fluid sample, and a urine sample. The methods provided herein for collecting samples permit increased sensitivity, increased ease of use, increased sample quality or combinations thereof, relative to previously used methods. Samples can be stored and/or treated using the methods known in the art or provided herein. Presence of an oncofetal fibronectin indicating molecule in the sample can be determined using the methods provided herein or otherwise known in the art. Exemplary methods include dot blot analysis, western blot analysis, northern blot analysis, southern blot analysis, RT-PCR methods, mass spectrometric methods, sandwich assays such as test strip-based sandwich assays and ELISA methods. The oncofetal fibronectin indicating molecule measurement methods provided herein permit increased sensitivity in oncofetal fibronectin indicating molecule detection and provide additional advantages as well. Accordingly, the methods disclosed herein can be used for early indication of the presence of cancer (e.g., hyperplastic, neoplastic, malignant or metastatic) cells, by, for example being able to detect lower levels of an oncofetal fibronectin indicating molecule in a sample, or by facilitating more frequent testing for cancer.

[0481] Detection of an oncofetal fibronectin indicating molecule in a sample can indicate the presence of a solid tumor, a leukemia (i.e., a blood-born cancer), cancer, metastasis, hyperplasia or neoplasia in a subject. Detection of an oncofetal fibronectin indicating molecule in a sample can indicate the stage or severity of a solid tumor, a leukemia, cancer, metastasis, hyperplasia or neoplasia in a subject. As provided herein, an oncofetal fibronectin indicating molecule is present in a variety of cancers (solid and blood-born), tumors, metastases and neoplasias. Thus, an oncofetal fibronectin positive sample can be used to indicate the presence and/or stage of a tumor, cancer, metastasis or neoplasia in a subject, or can be used to indicate the presence of cancer (e.g., neoplastic, malignant or metastatic) cells in a subject. Detection of the presence of an oncofetal fibronectin indicating molecule can indicate the presence and/or stage of a tumor, cancer, metastasis or neoplasia in a subject, or can indicate the presence of cancer (e.g., neoplastic, malignant or metastatic) cells in a subject. In another embodiment, detection of an oncofetal fibronectin indicating

molecule at or above one or more threshold levels can indicate the presence and/or stage of a tumor, cancer, metastasis or neoplasia in a subject, or the presence of cancer (e.g., neoplastic, malignant or metastatic) cells in a subject.

[0482] Absence or level below a threshold of an oncofetal fibronectin indicating molecule in the sample can indicate the absence of tumor, cancer, metastasis or neoplasia in a subject. As provided herein, since an oncofetal fibronectin indicating molecule is present in body tissues and fluids in numerous cancer (neoplastic) diseases, absence of an oncofetal fibronectin indicating molecule can indicate that the subject is free of a cancer (neoplastic) disease. For example, an oncofetal fibronectin negative sample can be used to indicate absence of a tumor, cancer, metastasis or neoplasia in a subject, or can be used to indicate the absence of cancer (e.g., neoplastic, malignant or metastatic) cells in a subject. The absence or level below a threshold of an oncofetal fibronectin indicating molecule can indicate the absence of tumor, cancer, metastasis or neoplasia in a subject, or can indicate the absence of cancer (e.g., hyperplastic, neoplastic, malignant or metastatic) cells in a subject. Also, an oncofetal fibronectin indicating molecule below one or more threshold levels can indicate the absence of tumor, cancer, metastasis or neoplasia in a subject, or the absence of cancer (e.g., neoplastic, malignant or metastatic) cells in a subject.

[0483] In another embodiment, the methods provided herein can be used to indicate an increased risk of a subject developing a tumor, cancer, metastasis or malignant neoplasia. For example, the methods provided herein can be used to indicate an increased risk of an organ, tissue or cell becoming cancerous (neoplastic, malignant or metastatic), relative to a normal population or relative to the individual. An organ, tissue or cell that contains or is adjacent a region containing an oncofetal fibronectin indicating molecule or an amount of an oncofetal fibronectin indicating molecule above normal levels can indicate an increased risk of developing cancer (neoplastic, malignant or metastatic growth) relative to an organ, tissue or cell that does not contain or is not adjacent to a region containing an oncofetal fibronectin indicating molecule or levels of an oncofetal fibronectin indicating molecule above normal. Accordingly, using the methods provided herein, an oncofetal fibronectin positive sample can be used to indicate an increased risk of a subject developing tumor, cancer, metastasis or malignant neoplasia, or to indicate an increased risk of cells, including normal, abnormal, dysplastic or hyperplastic cells, developing into cancer (neoplastic, malignant or metastatic) cells, relative to a subject with an oncofetal fibronectin negative sample or relative to a sample having an amount of oncofetal fibronectin below a threshold. The presence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a threshold can indicate an increased risk of an organ, tissue or cell in a subject becoming cancerous (neoplastic, malignant or metastatic), or an increased risk of a normal, abnormal, dysplastic or hyperplastic cell becoming cancerous (neoplastic, malignant or metastatic), relative to a subject with a sample in which an oncofetal fibronectin indicating molecule is absent or below a threshold level. In one aspect, the presence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a

threshold can indicate an increased risk of an organ, tissue or cell in a subject becoming cancerous (neoplastic, malignant or metastatic), or an increased risk of a normal, abnormal, dysplastic or hyperplastic cell becoming cancerous (neoplastic, malignant or metastatic), relative to a baseline sample of the subject.

[0484] In another embodiment, the methods provided herein can be used to indicate a decreased risk of a subject developing a tumor, cancer, metastasis or neoplasia. For example, the methods provided herein can be used to indicate a decreased risk of an organ, tissue or cell becoming cancerous (e.g., neoplastic, malignant or metastatic). An organ, tissue or cell that lacks or is adjacent a region lacking an oncofetal fibronectin indicating molecule or has an amount of an oncofetal fibronectin indicating molecule at or below normal levels can indicate a decreased risk of future development of cancerous (e.g., neoplastic, malignant or metastatic) growth relative to an organ, tissue or cell that does contain or is adjacent to a region containing an oncofetal fibronectin indicating molecule or amounts of an oncofetal fibronectin indicating molecule above normal. Accordingly, using the methods provided herein, an oncofetal fibronectin negative sample can be used to indicate a decreased risk of a subject developing tumor, cancer, metastasis or neoplasia, or to indicate a decreased risk of cells, including normal, abnormal, dysplastic or hyperplastic cells, developing into cancerous (e.g., neoplastic, malignant or metastatic) cells, relative to a subject with an oncofetal fibronectin positive sample or relative to a sample having an amount of oncofetal fibronectin equal to or above a threshold. The presence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample below a threshold can indicate a decreased risk of an organ, tissue or cell in a subject becoming of cancerous (e.g., neoplastic, malignant or metastatic), or a decreased risk of a normal, abnormal, dysplastic or hyperplastic cell becoming of cancerous (e.g., neoplastic, malignant or metastatic), relative to a subject with a sample in which an oncofetal fibronectin indicating molecule is present or at or above a threshold level, or relative to a baseline sample from the subject.

[0485] Further provided herein is a method for indicating the development of cancer by measuring oncofetal fibronectin indicating molecule in a sample, where the presence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a threshold can indicate development of an organ, tissue or cell into a cancerous organ, tissue or cell. Standard methods for defining cancerous organs, tissues or cells require histological examination of the morphology of the organ, tissue or cells. As provided herein, measurement of an oncofetal fibronectin indicating molecule can function to indicate that organs, tissues or cells that may not be morphologically defined as cancerous, are developing or have developed into cancerous or pre-cancerous organs, tissues or cells, notwithstanding any morphological classification.

[0486] Further provided herein is a method for indicating the development of cancer cells (including normal cells, abnormal cells, dysplastic cells, hyperplastic cells, pre-cancerous neoplastic cells, malignant cells or metastatic cells) into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth. Normal

cells can develop into cancerous cells, and the process of this change can take place by normal cells becoming more primitive, undifferentiated, anaplastic, and/or unregulated in growth. The development of normal cells into cancerous cells can include a variety of transitions such as, for example, normal cells developing into abnormal cells, abnormal cells developing into dysplastic cells, dysplastic cells developing into hyperplastic cells, hyperplastic cells developing into neoplastic cells, neoplastic cells developing into malignant cells, and malignant cells developing into metastatic cells. As provided herein, an oncofetal fibronectin positive result indicates that cells are developing or have an increased likelihood of developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth. Accordingly, methods are provided herein for indicating that cells are developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth, by measuring the oncofetal fibronectin indicating molecule in a sample, where presence, or an amount at or above a threshold indicates that the cells are developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth. Also provided herein are methods for indicating that cells have an increased likelihood of developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth, relative to a control sample in which an oncofetal fibronectin indicating molecule is absent or is below a threshold. In one example, a sample of abnormal, dysplastic or hyperplastic cells, or a swab of a region containing abnormal, dysplastic or hyperplastic cells, can be examined for the presence and/or amount of an oncofetal fibronectin indicating molecule, where an oncofetal fibronectin positive sample can indicate that the abnormal, dysplastic or hyperplastic cells are developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth.

[0487] Similarly, absence of an oncofetal fibronectin indicating molecule, or oncofetal fibronectin indicating molecule amounts below a threshold indicate that cells are not developing or have a decreased likelihood of developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth. Accordingly, methods are provided herein for indicating that cells are not developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth, by measuring an oncofetal fibronectin indicating molecule in a sample, where absence, or an amount below a threshold indicates that the cells are not developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth. Also provided herein are methods for indicating that cells have a decreased likelihood of developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth, by measuring the oncofetal fibronectin indicating molecule in a sample, where absence, or an amount below a threshold indicates that the cells have a decreased likelihood of developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth, relative to a sample in which an oncofetal fibronectin indicating

molecule is present or is at or above a threshold. In one example, a sample of abnormal, dysplastic or hyperplastic cells, or a swab of a region containing abnormal, dysplastic or hyperplastic cells, can be examined for the presence and/or amount of an oncofetal fibronectin indicating molecule, where an oncofetal fibronectin negative sample can indicate that the abnormal, dysplastic or hyperplastic cells are not developing into cells that are increasingly primitive, undifferentiated, anaplastic, and/or unregulated in growth.

[0488] In another embodiment, the methods provided herein can be used to indicate progression of a tumor, cancer, metastasis or neoplasia in a subject. For example, the methods provided herein can be used to indicate progression of cancer of an organ, tissue or cell. Presence of an oncofetal fibronectin indicating molecule or an amount at or above a threshold can indicate progression of a cancer or metastasis where a cancer or metastasis in a subject continues to be, or is increasingly, malignant or metastatic. Accordingly, using the methods provided herein, an oncofetal fibronectin positive sample can be used to indicate progression of a tumor, cancer, metastasis or neoplasia in a subject, or to indicate progression of cancer cells, in a subject. The presence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a threshold can indicate progression of a tumor, cancer, metastasis or neoplasia in a subject, or to indicate progression of cancer cells, in a subject.

[0489] In another embodiment, the methods provided herein can be used to indicate regression of a tumor, cancer, metastasis or neoplasia in a subject. For example, the methods provided herein can be used to indicate regression of cancer of an organ, tissue or cell. Absence of an oncofetal fibronectin indicating molecule or an amount below a threshold can indicate regression of a cancer or metastasis, where a cancer or metastasis in a subject ceases to be, or is decreasingly, malignant or metastatic. Accordingly, using the methods provided herein, an oncofetal fibronectin negative sample can be used to indicate regression of a cancer or metastasis in a subject, or to indicate regression of cancer cells, in a subject. The absence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample below a threshold can indicate regression of a tumor, cancer, metastasis or neoplasia in a subject, or to indicate regression of cancer, in a subject.

[0490] In another embodiment, the methods provided herein can be used to distinguish between aggressive and non-aggressive cancer (e.g., a solid tumor or a leukemia). Cancer (solid or leukemia) with rapid growth properties can have different compositions relative to cancer with slow growth properties. For example, aggressive or rapid growth or high grade cancer such as astrocytomas can contain an oncofetal fibronectin indicating molecule or can contain an amount of an oncofetal fibronectin indicating molecule at or above a threshold, while non-aggressive or slow growth or low grade cancer can contain no oncofetal fibronectin indicating molecule, or amounts below a threshold. Accordingly, using the methods provided herein, measurement of an oncofetal fibronectin indicating molecule in a sample can be used to distinguish between an aggressive or rapid growth or high grade cancer and a non-aggressive or slow growth or low grade cancer. The methods can include measuring an oncofetal fibronectin indicating molecule in a sample, where

the presence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a threshold can indicate an aggressive or rapid growth or high grade cancer; similarly, absence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample below a threshold can indicate a non-aggressive or slow growth or low grade cancer. The methods provided herein also can be used to distinguish between a aggressive or rapid growth cells, including normal, abnormal, dysplastic, hyperplastic, neoplastic, malignant or metastatic cells and a non-aggressive or slow growth or low grade cell. The methods include measuring an oncofetal fibronectin indicating molecule in a sample, where the presence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a threshold can indicate aggressive or rapid growth normal, abnormal, dysplastic, hyperplastic, neoplastic, malignant or metastatic cancer cells; similarly, absence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample below a threshold can indicate non-aggressive or slow growth normal, abnormal, dysplastic, hyperplastic, neoplastic, malignant or metastatic cancer cells.

[0491] Further provided herein is a method for indicating the recurrence or likelihood of recurrence of cancer. Cancer can spontaneously subside or can subside as a result of therapy. A cancer also can recur in a subject. As provided herein, an oncofetal fibronectin positive sample can indicate that cancer has recurred in a subject. An oncofetal fibronectin positive sample also can indicate that cancer has an increased likelihood of recurring in a subject. Also provided herein, an oncofetal fibronectin positive sample can indicate that cancer cells are again present in a subject, or have an increased likelihood of being present again in a subject. The methods include measuring an oncofetal fibronectin indicating molecule in a sample, where the presence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a threshold indicates recurrence of cancer) in a subject or recurrence of cancer cells in the subject. The methods also include measuring an oncofetal fibronectin indicating molecule in a sample, where the presence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a threshold indicates an increased likelihood of recurrence of cancer in a subject or an increased likelihood of recurrence of cancer cells in the subject, relative to the likelihood of recurrence in a subject with a sample in which an oncofetal fibronectin indicating molecule is absent or is below a threshold, or relative to the likelihood of recurrence in a control sample in which an oncofetal fibronectin indicating molecule is present or is absent or is below a threshold.

[0492] Similarly, an oncofetal fibronectin negative sample can indicate that cancer (tumorous, metastatic or neoplastic disease) has not recurred in a subject. An oncofetal fibronectin negative sample also can indicate that the cancer has a decreased likelihood of recurring in a subject. Also provided herein, an oncofetal fibronectin negative sample can indicate that cancer cells are still not present in a subject, or have a decreased likelihood of being present again in a subject. The methods include measuring an oncofetal fibronectin indicat-

ing molecule in a sample, where the absence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample below a threshold indicates a lack of recurrence of cancer in a subject or a lack of recurrence of cancer cells in the subject. The methods also include measuring an oncofetal fibronectin indicating molecule in a sample, where the absence of an oncofetal fibronectin indicating molecule in a sample or an amount of an oncofetal fibronectin indicating molecule in a sample below a threshold indicates a decreased likelihood of recurrence of cancer, in a subject or a decreased likelihood of recurrence of cancer cells in the subject, relative to the likelihood of recurrence in a subject with a sample in which an oncofetal fibronectin indicating molecule is present or is at or above a threshold, or relative to the likelihood of recurrence in a control sample in which an oncofetal fibronectin indicating molecule is present or is at or above a threshold.

[0493] In another embodiment, the methods provided herein can be used to determine the success or the likelihood of success of treating a cancer (e.g., a solid tumor, leukemia, metastasis, or neoplastic disease). While some cancers can be successfully treated with therapy such as administration of one or more compounds such as chemotherapeutic compounds, other cancers are less responsive to such compounds or are responsive to different therapies. A cancer in which an oncofetal fibronectin indicating molecule is present or is at or above a threshold level can have a different sensitivity to treatment relative to a cancer in which an oncofetal fibronectin indicating molecule is absent or is below a threshold level. Accordingly, using the methods provided herein, measurement of an oncofetal fibronectin indicating molecule in a sample can be used to predict the success or to indicate the likelihood of success of treating a cancer. The likelihood of success can be a function of the particular therapy. For example a therapy can target neovascularization, or a therapy can target nucleotide synthesis; one skilled in the art can recognize how the presence or amount of an oncofetal fibronectin indicating molecule in sample can affect the likelihood of success of cancer therapy. Presence of an oncofetal fibronectin indicating molecule or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a threshold level can indicate that a particular therapeutic treatment has an increased likelihood of being successful, relative to a sample in which an oncofetal fibronectin indicating molecule is absent or is below a threshold level. In another example, presence of an oncofetal fibronectin indicating molecule or an amount of an oncofetal fibronectin indicating molecule in a sample at or above a threshold level can indicate that a particular therapeutic treatment has a decreased likelihood of being successful, relative to a sample in which an oncofetal fibronectin indicating molecule is absent or is below a threshold level. In another example, absence of an oncofetal fibronectin indicating molecule or an amount of an oncofetal fibronectin indicating molecule in a sample below a threshold level can indicate that a particular therapeutic treatment has a decreased likelihood of being successful, relative to a sample in which an oncofetal fibronectin indicating molecule is present or is at or above a threshold level. In another example, absence of an oncofetal fibronectin indicating molecule or an amount of an oncofetal fibronectin indicating molecule in a sample below a threshold level can indicate that a particular therapeutic treatment has an increased

likelihood of being successful, relative to a sample in which an oncofetal fibronectin indicating molecule is present or is at or above a threshold level.

[0494] In another embodiment, measurement of an oncofetal fibronectin indicating molecule in a sample can be used to monitor the success of therapeutic treatment of a cancer (e.g., a solid tumor, leukemia, metastasis, or malignant neoplasia). For example, after therapeutic treatment, presence of an oncofetal fibronectin indicating molecule or a level of an oncofetal fibronectin indicating molecule in a sample at or above a threshold level can indicate that a therapeutic treatment being conducted is ineffective. In another example, presence of an oncofetal fibronectin indicating molecule or a level of an oncofetal fibronectin indicating molecule in a sample at or above a threshold level can indicate that a therapeutic treatment being conducted is effective. In another example, absence of an oncofetal fibronectin indicating molecule or a level of an oncofetal fibronectin indicating molecule in a sample below a threshold level can indicate that a therapeutic treatment being conducted is effective. In another example, absence of an oncofetal fibronectin indicating molecule or a level of an oncofetal fibronectin indicating molecule in a sample below a threshold level can indicate that a therapeutic treatment being conducted is ineffective.

[0495] In another embodiment, measurement of an oncofetal fibronectin indicating molecule in a sample can be used to select a method of treating a cancer (e.g., a solid tumor, leukemia, metastasis or malignant neoplasia). For example, presence of an oncofetal fibronectin indicating molecule or a level of an oncofetal fibronectin indicating molecule in a sample at or above a threshold level can be used to select a method of treatment that is more effective for cancer associated with oncofetal fibronectin. In another example, absence of an oncofetal fibronectin indicating molecule or a level of an oncofetal fibronectin indicating molecule in a sample below a threshold level can be used to select a method of treatment that is more effective for cancer associated with oncofetal fibronectin.

[0496] In some embodiments, detection of an oncofetal fibronectin indicating molecule can be performed in conjunction with detection of one or more additional cancer (i.e., tumor-associated) markers. A variety of detectable cancer markers are known in the art or are provided elsewhere herein, exemplary markers include, but are not limited to, AE1/AE3, BCA-225, Cathespin D, E-Cadherin, Epidermal Growth Factor Receptor (EGFR), Estrogen receptor (ER), Gross Cystic Disease Fluid Protein 15 (GCDFP-15), HOX-B3, Ki-67, p65, Progesterone Receptor (PR), Retinoblastoma (Rb) and Transglutaminase K (TGK), p21, DCC, NF-1, NF-2, BRCA-3, p16, FHIT, WT-1, MEN-1, MEN-IIa, MEN-IIb, VHL, FCC, MCC, raf, erb, src, fms, jun, trk, ret, gsp, hst, bcr/ab1, p53, c-erbB2, c-myc, MUC1, BRCA1, BRCA2, Her-2/neu, bcl-2, bax, PSA, CYFRA 21-1, PTH-RP, CA125, CEA gene family members, progastrin, gastrin G17, gastrin G34, CA 19-9, CA 15-3, CA 27-29, CA 72-4, APC, SCC, HPV subtypes, TK, alphaFP, p62, Kallikrein, ras, vasopressin, gastrin-releasing peptide, annexin I, annexin II, Hu and KOC. Additional cancer markers, occurrence of cancer markers in particular cancers, and occurrence of cancer markers with other cancer markers are known in the art, as exemplified in Rhodes et al., *Proc. Natl. Acad. Sci. USA* 2004 101: 9309-9314. For example

presence of an oncofetal fibronectin indicating molecule and Her-2/neu can indicate that a subject has breast cancer, and presence of an oncofetal fibronectin indicating molecule and PSA can indicate that a subject has prostate cancer. Among the methods provided herein are methods in which the presence and/or level of an oncofetal fibronectin indicating molecule in a sample is measured and the presence and/or level of one or more additional cancer markers is determined. Such methods can serve to further characterize the cancer or identify the cell or tissue source of cancer (dysplastic, neoplastic, malignant or metastatic) cells.

[0497] In some instances detection of an oncofetal fibronectin indicating molecule in a sample can have a strong positive predictive value of a present or future cancerous (neoplastic) condition, but a less strong negative predictive value; in such instances, a method for identifying or predicting a present or future cancerous (neoplastic) condition by detecting an oncofetal fibronectin indicating molecule, such as the methods provided herein, can be coupled with use of a second marker with a strong negative predictive value. Agreement of the oncofetal fibronectin indicating molecule measurement and measurement of the second marker can indicate with increased certainty relative to either marker when used alone, the present or future cancer (neoplastic) condition of the subject. For example, presence of oncofetal fibronectin protein in a sample can have a 90% or about a 90% positive predictive value for bladder cancer and absence of bladder tumor antigen in a sample can have a 90% or about a 90% negative predictive value for bladder cancer; when both markers are used in conjunction with each other, results that are either both tests are either positive or negative will be correct for 95% or about 95% of subjects tested.

[0498] In other instances detection of an oncofetal fibronectin indicating molecule in a sample can have a strong negative predictive value of a present or future cancerous (neoplastic) condition, but a less strong positive predictive value; in such instances, a method for identifying or predicting a present or future cancerous (neoplastic) condition by detecting an oncofetal fibronectin indicating molecule, such as the methods provided herein, can be coupled with use of a second marker with a strong positive predictive value. Agreement of the oncofetal fibronectin indicating molecule measurement and measurement of the second marker can indicate with increased certainty relative to either marker when used alone, the present or future cancerous (neoplastic) condition of the subject.

[0499] In some cases a measurement is considered positive for oncofetal fibronectin when the presence of an oncofetal fibronectin indicating molecule in a sample is equal to or above one or more threshold levels. In some embodiments, the threshold level can vary, for example, as a function of the progression of disease, a subject-specific classification, or the age of the subject. A threshold level that varies as a function of a second factor can be expressed as a threshold curve. In some instances, the rate of change of the amount of an oncofetal fibronectin indicating molecule in a particular sample type from a subject can be used to identify a sample as oncofetal fibronectin positive or negative, or to categorize the sample into two or more populations. The rate of change of the amount of an oncofetal fibronectin indicating molecule in a type of sample can

indicate a stable, increasing or decreasing amount of an oncofetal fibronectin indicating molecule in a sample.

[0500] In some embodiments, the cancerous (malignant neoplastic, tumorous or metastatic) disease indicating methods provided herein can further include monitoring the presence of an oncofetal fibronectin indicating molecule over time. For example, the same type of sample can be collected from a subject every day, every week, every month or every year, and the oncofetal fibronectin indicating molecule measurements can be compared. In such cases, it is possible to identify an increasing amount of an oncofetal fibronectin indicating molecule in a subject, a decreasing amount of an oncofetal fibronectin indicating molecule in a subject, or a constant level of an oncofetal fibronectin indicating molecule in a subject. As provided herein, an increasing rate of an oncofetal fibronectin indicating molecule, or a increasing rate equal to or greater than a threshold rate, can be considered an oncofetal fibronectin positive measurement for the cancerous disease indicating methods provided herein. Similarly, a decreasing rate of an oncofetal fibronectin indicating molecule, or a decreasing rate equal to or greater than a threshold rate, can be considered an oncofetal fibronectin negative measurement for the cancerous disease indicating methods provided herein. The size of the rate of increase or decrease also can indicate an increasing or decreasing severity or likelihood of the cancerous indication. For example, a greater increasing rate can indicate a more severe neoplastic disease, a higher risk of developing cancer or recurrence of cancer, a faster progression of the cancer, or a more aggressive cancer, or a decreased efficacy of cancer therapy, relative to a smaller increasing rate or a decreasing rate. In another example, a smaller increasing rate can indicate a less severe cancer, a lower risk of developing cancer or recurrence of cancer, a slower progression of the cancer, or a less aggressive cancer, or an increased efficacy of cancer therapy relative to a greater increasing rate. In another example, a decreasing rate can indicate a less severe cancer, a lower risk of developing cancer or recurrence of cancer, a slower progression of the cancer, or a less aggressive cancer, or an increased efficacy of cancer therapy, relative to an increasing rate.

[0501] In some instances, multi-tiered thresholds can be applied to the oncofetal fibronectin indicating molecule measurement, where multi-tiered thresholds include two or more threshold levels, where each larger threshold level indicates a separate cancer categorization; for example each larger threshold level can indicate an increased likelihood of having a cancer, an increased risk of a cancer, an increased degree of development of cells into cancer, an increased likelihood of recurrence of a cancer, an increased aggressiveness of a cancer, or an increased likelihood of successful or unsuccessful cancer therapy. In another example, each smaller threshold level indicates a separate cancer categorization; for example each smaller threshold level can indicate a decreased likelihood of having a cancer, a decreased risk of a cancer, a decreased degree of development of cells into cancer cells, a decreased likelihood of recurrence of a cancer, a decreased aggressiveness of a cancer, or a decreased likelihood of successful or unsuccessful cancer therapy. An exemplary multi-tiered threshold is a two-tiered threshold for oncofetal fibronectin protein, where the lower threshold is 50 ng/mL and the higher threshold is 150 ng/mL for buffer-treated samples.

[0502] In another embodiment, any detected oncofetal fibronectin indicating molecule can be characterized according to the oncofetal fibronectin domains and/or post-translational modifications present in the oncofetal fibronectin indicating molecule. For example, an oncofetal fibronectin indicating molecule can be characterized as containing one or more of EDA, EDB and IIICS. In another example, the fibronectin or oncofetal fibronectin indicating molecule can be characterized as the EDA portion of an oncofetal fibronectin protein, the EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, the portion of an autoantibody that binds to EDA of oncofetal fibronectin protein, and the portion of an autoantibody that binds to the EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin. In another example, the fibronectin or oncofetal fibronectin indicating molecule can be characterized as the EDB portion of an oncofetal fibronectin protein, the EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, the portion of an autoantibody that binds to EDB of oncofetal fibronectin protein, and the portion of an autoantibody that binds to the EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin. In another example, the fibronectin or oncofetal fibronectin indicating molecule can be characterized as the IIICS portion of an oncofetal fibronectin protein, the IIICS-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, the portion of an autoantibody that binds to IIICS of oncofetal fibronectin protein, and the portion of an autoantibody that binds to the IIICS-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin. In another example, an oncofetal fibronectin indicating molecule can be characterized as containing the IIICS splice variant V64, V89, V95 or V120. In another example, an oncofetal fibronectin protein can be characterized as containing one or more post-translational modifications such as O-glycosylation of threonine 33 of IIICS. In another example, an oncofetal fibronectin protein can be characterized as lacking EDA, EDB or IIICS. In another example, IIICS is identified as lacking amino acids 1-25 of IIICS, or 90-120 of IIICS, or both. Characterization of an oncofetal fibronectin indicating molecule present in a sample can be used, for example, to identify the cell or tissue source of the oncofetal fibronectin indicating molecule.

[0503] a. Bladder Cancer

[0504] The methods provided herein include use of an oncofetal fibronectin indicating molecule as a marker for bladder cancer. The commonest site of occurrence of bladder cancer is on the trigone and lateral walls of the bladder. The growth can be papillary, sessile, solid or ulcerative. Most bladder cancers are transitional cell carcinoma (TCC). There can be a considerable amount of metaplasia, especially among muscle invasive tumors. Five percent of bladder cancers are pure squamous cell carcinoma, which are usually associated with irritative factors such as chronic indwelling catheters, stones or schistosomiasis infestation. Pure adenocarcinoma of the bladder is rare and metastasis from another primary should be excluded. Carcinoma in situ (CIS) is a flat epithelia lesion which displays a lack of cellular polarity and has anaplastic features. The lesion can be local or diffuse. When bladder cancer presents as a diffuse lesion, it usually follows an aggressive course.

[0505] An oncofetal fibronectin indicating molecule can be present in subjects that have bladder cancer. Oncofetal

fibronectin indicating molecule measurements can be used as a marker for bladder cancer in any of the cancer indications provided herein. For example, the oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has cancerous (malignant neoplastic or metastatic) bladder cells. The methods include determining the presence or amount of an oncofetal fibronectin indicating molecule in a sample, and characterizing the sample according to the presence or absence of an oncofetal fibronectin indicating molecule in the sample or according to the amount of an oncofetal fibronectin indicating molecule in the sample, where presence or an amount of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate that a subject has cancerous bladder cells, and absence or an amount of an oncofetal fibronectin indicating molecule below a threshold level can indicate that a subject does not have cancerous bladder cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the risk or lack of risk of bladder cells such as normal, abnormal, dysplastic or hyperplastic bladder cells, becoming cancerous. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the development or lack of development of bladder cells such as normal, abnormal, dysplastic or hyperplastic bladder cells, into less developed or anaplastic bladder cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for indicating the progression of a bladder tumor, cancer, metastasis or neoplasia. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for distinguishing between aggressive and non-aggressive bladder tumors. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has or is likely to have a recurrence of cancerous bladder cells. The oncofetal fibronectin indicating molecule detection methods provided herein can indicate the likely or actual efficacy or lack of efficacy of bladder cancer treatment.

[0506] A sample collected for bladder cancer determination can be taken from any source, as provided herein. Exemplary sources for samples include a bladder tissue, a urine sample, a lymphatic sample, a blood sample, a serum sample, a plasma sample and an interstitial fluid sample. For example, the presence of an oncofetal fibronectin indicating molecule can be detected in the urine of subjects with bladder cancer. In one embodiment, detection of any oncofetal fibronectin indicating molecule in a sample can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous bladder cells. In another embodiment, detection of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous bladder cells. An exemplary threshold value for oncofetal fibronectin protein in buffer-treated urine sample (diluted from 250 μ l to 1000 μ l or about 250 μ l to about 1000 μ l) as an indicator of bladder cancer is 5 ng/ml, 8 ng/ml, 10 ng/ml, 15 ng/ml or 20 ng/ml, or about 5 ng/ml, about 8 ng/ml, about 10 ng/ml, about 15 ng/ml or about 20 ng/ml. All forms of bladder cancer can be indicated using the methods provided herein. Exemplary forms of bladder cancer include transitional cell carcinoma, squamous cell carcinoma and adenocarcinoma.

[0507] In one embodiment, characterization of an oncofetal fibronectin indicating molecule in a sample can indicate whether or not the oncofetal fibronectin indicating molecule in the sample was produced by cancerous bladder cells. In some cases, one or more forms of oncofetal fibronectin indicating molecule (varying, e.g., at EDA, EDB, IIICS and/or in post-translational modifications) present in a sample such as urine or bladder tissue are known to be produced by cancerous bladder cells. For example, oncofetal fibronectin protein in urine samples from subjects with bladder cancer contain O-glycosylated threonine 33 in the IIICS region of fibronectin in the urine (Wunderlich et al., *Oncol. Rep.* 8:669-672 (2001)), and fibronectin protein in urine samples from subjects with bladder cancer can contain abnormal glycosylation (Guo et al., *J. Cancer Res. Clin. Oncol.* 127:512-519(2001)). In other cases, one or more forms of oncofetal fibronectin indicating molecule, although present in a sample such as urine or bladder tissue, are not produced by cancerous bladder cells, but instead are produced by a different tissue or organ source. The methods herein can be used to characterize the oncofetal fibronectin indicating molecule in a sample, and such characterization can indicate whether or not the oncofetal fibronectin indicating molecule observed in the sample is an oncofetal fibronectin indicating molecule form observed in bladder cancer. When a sample contains an oncofetal fibronectin indicating molecule observed in bladder cancer, the presence of the oncofetal fibronectin indicating molecule is consistent with a subject having bladder cancer. When a sample contains an oncofetal fibronectin indicating molecule not observed in bladder cancer, the presence of the oncofetal fibronectin indicating molecule is not consistent with a subject having bladder cancer.

[0508] b. Breast Cancer

[0509] The methods provided herein include use of an oncofetal fibronectin indicating molecule as a marker for breast cancer. An oncofetal fibronectin indicating molecule can be present in subjects that have breast cancer. Oncofetal fibronectin indicating molecule measurements can be used as a marker for breast cancer in any of the cancer (tumor, metastasis or malignant neoplastic) indications provided herein. For example, the oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has cancerous breast cells. The methods include determining the presence or amount of an oncofetal fibronectin indicating molecule in a sample, and characterizing the sample according to the presence or absence of an oncofetal fibronectin indicating molecule in the sample or according to the amount of an oncofetal fibronectin indicating molecule in the sample, where presence or an amount of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate that a subject has cancerous breast cells, and absence or an amount of an oncofetal fibronectin indicating molecule below a threshold level can indicate that a subject does not have cancerous breast cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the risk or lack of risk of breast cells such as normal, abnormal, dysplastic or hyperplastic breast cells, becoming cancerous. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the development or lack of development of breast cells such as normal, abnormal, dysplastic or hyperplastic breast cells, into cancerous cells. The

oncofetal fibronectin indicating molecule detection methods provided herein can be used for distinguishing between aggressive and non-aggressive breast tumors. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for indicating the progression of a breast cancer (e.g., tumor or metastasis). The oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has or is likely to have a recurrence of cancerous breast cells. The oncofetal fibronectin indicating molecule detection methods provided herein can indicate the likely or actual efficacy or lack of efficacy of breast cancer treatment.

[0510] A sample collected for breast cancer determination can be taken from any source, as provided herein. Exemplary sources for samples include a breast tissue sample, a fine needle aspiration sample, a ductal lavage sample, a blood sample, a serum sample, a plasma sample, a lymph sample or an interstitial fluid sample. For example, the presence of an oncofetal fibronectin indicating molecule can be detected in ductal lavage collected from subjects with breast cancer. In one embodiment, detection of any oncofetal fibronectin indicating molecule in a sample can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous breast cells. In another embodiment, detection of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous breast cells. An exemplary threshold value for oncofetal fibronectin protein in a ductal lavage sample (diluted from 250 μ l to 1000 μ l or about 250 μ l to about 1000 μ l) as an indicator of breast cancer is 5 ng/ml, 8 ng/ml, 10 ng/ml, 15 ng/ml or 20 ng/ml, or about 5 ng/ml, about 8 ng/ml, about 10 ng/ml, about 15 ng/ml or about 20 ng/ml. All forms of breast cancer can be indicated using the methods provided herein. Exemplary forms of breast cancer include infiltrating ductal carcinoma, invasive ductal carcinoma, other forms of ductal carcinoma, lobular carcinoma, nipple carcinoma and undifferentiated breast carcinoma.

[0511] In one embodiment, characterization of an oncofetal fibronectin indicating molecule in a sample can indicate whether or not the oncofetal fibronectin indicating molecule in the sample was produced by cancerous breast cells. In some cases, one or more forms of oncofetal fibronectin indicating molecule (varying, e.g., at EDA, EDB, IIICS and/or in post-translational modifications) present in a sample such as ductal lavage or breast tissue are known to be produced by cancerous breast cells. For example, oncofetal fibronectin protein in breast tissue samples from subjects with invasive ductal carcinoma contained EDB and O-glycosylated threonine 33 in the IIICS region of fibronectin in the tissue samples (Kaczmarek et al., *Int. J. Cancer* 59:11-16 (1994)), and cancerous breast tissue samples have also been found to contain EDA+ oncofetal fibronectin (Matsumoto et al., *Jpn. J. Cancer Res.* 90:320-325 (1999)). In other cases, one or more forms of oncofetal fibronectin indicating molecule, although present in a sample such as ductal lavage or breast tissue, are not produced by cancerous breast cells, but instead are produced by a different tissue or organ source. The methods herein can be used to characterize the oncofetal fibronectin indicating molecule in a sample, and such characterization can indicate whether or not the oncofetal fibronectin indicating molecule observed in the sample is an

oncofetal fibronectin indicating molecule form observed in breast cancer. When a sample contains an oncofetal fibronectin indicating molecule observed in breast cancer, the presence of the oncofetal fibronectin indicating molecule is consistent with a subject having breast cancer. When a sample contains an oncofetal fibronectin indicating molecule not observed in breast cancer, the presence of the oncofetal fibronectin indicating molecule is not consistent with a subject having breast cancer.

[0512] c. Cervical Cancer

[0513] The methods provided herein include use of an oncofetal fibronectin indicating molecule as a marker for cervical cancer. An oncofetal fibronectin indicating molecule can be present in subjects that have cervical cancer. Oncofetal fibronectin indicating molecule measurements can be used as a marker for cervical cancer in any of the cancer (tumor, metastasis or malignant neoplastic) indications provided herein. For example, the oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has cancerous (malignant neoplastic or metastatic) cervical cells. The methods include determining the presence or amount of an oncofetal fibronectin indicating molecule in a sample, and characterizing the sample according to the presence or absence of an oncofetal fibronectin indicating molecule in the sample or according to the amount of an oncofetal fibronectin indicating molecule in the sample, where presence or an amount of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate that a subject has cancerous cervical cells, and absence or an amount of an oncofetal fibronectin indicating molecule below a threshold level can indicate that a subject does not have cancerous cervical cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the risk or lack of risk of cervical cells such as normal, abnormal, dysplastic or hyperplastic cervical cells, becoming cancerous. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the development or lack of development of cervical cells such as normal, abnormal, dysplastic or hyperplastic cervical cells, into cancerous cervical cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for indicating the progression of a cervical cancer. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for distinguishing between aggressive and non-aggressive cervical tumors. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has or is likely to have a recurrence of cancerous cervical cells. The oncofetal fibronectin indicating molecule detection methods provided herein can indicate the likely or actual efficacy or lack of efficacy of cervical cancer treatment.

[0514] A sample collected for cervical cancer determination can be taken from any source, as provided herein. Exemplary sources for samples include a cervical or vaginal tissue sample, urine, lymph, lymphatic fluid, blood, serum, plasma, interstitial fluid and cervicovaginal fluid. Cervicovaginal fluid can be collected by any of a variety of methods, such as by swab, from any of a variety of cervicovaginal locations. Exemplary cervicovaginal swab samples include a swab of the point of a possible cervicovaginal lesion, the

cervical canal, the cervical os, the ectocervix, the transition zone on the cervix between squamous and columnar cells (i.e., the squamocolumnar junction), the vagina, the posterior fornix, the portion of the vagina below the posterior fornix such as the lower third of the vagina, the labia, or combinations thereof. Cervicovaginal samples also can be collected as cervicovaginal fluid leakage from the vagina. In one example, the presence of an oncofetal fibronectin indicating molecule can be detected in a swab of the cervical os in subjects with cervical cancer. Detection of any oncofetal fibronectin indicating molecule in a sample can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous cervical cells. In another embodiment, detection of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous cervical cells. An exemplary threshold value for an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, in a cervical os swab buffer-treated sample as an indicator of cervical cancer is 1 ng/ml, 3 ng/ml, 5 ng/ml, 8 ng/ml, 10 ng/ml, 15 ng/ml, 20 ng/ml, or 25 ng/ml, or about 1 ng/ml, about 3 ng/ml, about 5 ng/ml, about 8 ng/ml, about 10 ng/ml, about 15 ng/ml, about 20 ng/ml, or about 25 ng/ml. An exemplary threshold value for an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, in a cervical os swab untreated sample as an indicator of cervical cancer is 5 ng/ml, 10 ng/ml, 20 ng/ml, 30 ng/ml, 40 ng/ml, 50 ng/ml, 60 ng/ml, or 75 ng/ml, or about 5 ng/ml, about 10 ng/ml, about 20 ng/ml, about 30 ng/ml, about 40 ng/ml, about 50 ng/ml, about 60 ng/ml, or about 75 ng/ml. An exemplary threshold value for an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, in a lower vaginal swab buffer-treated sample as an indicator of cervical cancer is 1 ng/ml, 2 ng/ml, 3 ng/ml, 5 ng/ml, 8 ng/ml, 10 ng/ml, 15 ng/ml, or 25 ng/ml, or about 1 ng/ml, about 2 ng/ml, about 3 ng/ml, about 5 ng/ml, about 8 ng/ml, about 10 ng/ml, about 15 ng/ml, or about 25 ng/ml. An exemplary threshold value for an oncofetal fibronectin indicating molecule, such as oncofetal fibronectin protein, in a cervical os swab untreated sample as an indicator of cervical cancer is 2 ng/ml, 5 ng/ml, 10 ng/ml, 15 ng/ml, 20 ng/ml, 30 ng/ml, 40 ng/ml, or 50 ng/ml, or about 2 ng/ml, about 5 ng/ml, about 10 ng/ml, about 15 ng/ml, about 20 ng/ml, about 30 ng/ml, about 40 ng/ml, or about 50 ng/ml. All forms of cervical cancer can be indicated using the methods known in the art or provided herein. Exemplary forms of cervical cancer include squamous cell carcinoma and adenocarcinoma.

[0515] In one embodiment, characterization of an oncofetal fibronectin indicating molecule in a sample can indicate whether or not the oncofetal fibronectin indicating molecule in the sample was produced by cancerous cervical cells. In some cases, one or more forms of oncofetal fibronectin indicating molecule (varying, e.g., at EDA, EDB, IIICS and/or in post-translational modifications) present in a sample such as cervicovaginal fluid or cervical tissue are known to be produced by cancerous cervical cells. For example, as provided herein, oncofetal fibronectin protein in cervical os swab samples from subjects with cervical cancer contained O-glycosylated threonine 33 in the IIICS region of fibronectin in the samples. In other cases, one or more forms of oncofetal fibronectin indicating molecule, although

present in a sample such as cervicovaginal fluid or cervical tissue, are not produced by cancerous cervical cells, but instead are produced by a different tissue or organ source. The methods herein can be used to characterize the oncofetal fibronectin indicating molecule in a sample, and such characterization can indicate whether or not the oncofetal fibronectin indicating molecule observed in the sample is an oncofetal fibronectin indicating molecule form observed in cervical cancer. When a sample contains an oncofetal fibronectin indicating molecule observed in cervical cancer, the presence of the oncofetal fibronectin indicating molecule is consistent with a subject having cervical cancer. When a sample contains an oncofetal fibronectin indicating molecule not observed in cervical cancer, the presence of the oncofetal fibronectin indicating molecule is not consistent with a subject having cervical cancer.

[0516] d. Ovarian Cancer

[0517] The methods provided herein include use of an oncofetal fibronectin indicating molecule as a marker for ovarian cancer. An oncofetal fibronectin indicating molecule can be present in subjects that have ovarian cancer. Oncofetal fibronectin indicating molecule measurements can be used as a marker for ovarian cancer in any of the cancer (tumor, metastasis or malignant neoplastic) indications provided herein. For example, the oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has cancerous (malignant neoplastic or metastatic) ovarian cells. The methods include determining the presence or amount of an oncofetal fibronectin indicating molecule in a sample, and characterizing the sample according to the presence or absence of an oncofetal fibronectin indicating molecule in the sample or according to the amount of an oncofetal fibronectin indicating molecule in the sample, where presence or an amount of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate that a subject has cancerous ovarian cells, and absence or an amount of an oncofetal fibronectin indicating molecule below a threshold level can indicate that a subject does not have cancerous ovarian cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the risk or lack of risk of ovarian cells such as normal, abnormal, dysplastic or hyperplastic ovarian cells, becoming cancerous. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the development or lack of development of ovarian cells such as normal, abnormal, dysplastic or hyperplastic ovarian cells, into cancerous ovarian cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for indicating the progression of an ovarian cancer (tumor, metastasis or malignant neoplasia). The oncofetal fibronectin indicating molecule detection methods provided herein can be used for distinguishing between aggressive and non-aggressive ovarian tumors. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has or is likely to have a recurrence of cancerous ovarian cells. The oncofetal fibronectin indicating molecule detection methods provided herein can indicate the likely or actual efficacy or lack of efficacy of ovarian cancer treatment.

[0518] A sample collected for ovarian cancer determination can be taken from any source, as provided herein.

Exemplary sources for samples include an ovarian tissue sample, ascitic fluid, peritoneal fluid, urine, stool, plasma, blood, serum, lymph, lymphatic fluid and interstitial fluid samples. For example, the presence of an oncofetal fibronectin indicating molecule can be detected in peritoneal fluid collected from subjects with ovarian cancer. Detection of any oncofetal fibronectin indicating molecule in a sample can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous ovarian cells. In another embodiment, detection of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous ovarian cells. An exemplary threshold value for an oncofetal fibronectin indicating molecule in a peritoneal sample as an indicator of ovarian cancer is 3 ng/ml, 5 ng/ml, 8 ng/ml, 10 ng/ml, 15 ng/ml, 20 ng/ml, 25 ng/ml, or 30 ng/ml, or about 3 ng/ml, about 5 ng/ml, about 8 ng/ml, about 10 ng/ml, about 15 ng/ml, about 20 ng/ml, about 25 ng/ml, or about 30 ng/ml. All forms of ovarian cancer can be indicated using the methods known in the art or provided herein. Exemplary forms of ovarian cancer include serous cystoma, mucinous cystoma, endometriod tumor, mesonephroid tumor, dysgerminoma, endodermal sinus tumor, embryonal carcinoma, polyembroma, choriocarcinoma and teratoma.

[0519] In one embodiment, characterization of an oncofetal fibronectin indicating molecule in a sample can indicate whether or not the oncofetal fibronectin indicating molecule in the sample was produced by cancerous ovarian cells. In some cases, one or more forms of oncofetal fibronectin indicating molecule (varying, e.g., at EDA, EDB, IIICS and/or in post-translational modifications) present in a sample such as peritoneal fluid or ovarian tissue are known to be produced by cancerous ovarian cells. For example, oncofetal fibronectin protein in peritoneal samples from subjects with ovarian cancer contained O-glycosylated threonine 33 in the IIICS region of fibronectin and at least another portion of the IIICS region of fibronectin in the samples (see, e.g., U.S. Pat. No. 5,523,229; Menzin et al., *Cancer* 82:152-158 (1998)). In other cases, one or more forms of oncofetal fibronectin indicating molecule, although present in a sample such as peritoneal fluid or ovarian tissue, are not produced by cancerous ovarian cells, but instead are produced by a different tissue or organ source. The methods herein can be used to characterize the oncofetal fibronectin indicating molecule in a sample, and such characterization can indicate whether or not the oncofetal fibronectin indicating molecule observed in the sample is an oncofetal fibronectin indicating molecule form observed in ovarian cancer. When a sample contains an oncofetal fibronectin indicating molecule observed in ovarian cancer, the presence of the oncofetal fibronectin indicating molecule is consistent with a subject having ovarian cancer. When a sample contains an oncofetal fibronectin indicating molecule not observed in ovarian cancer, the presence of the oncofetal fibronectin indicating molecule is not consistent with a subject having ovarian cancer.

[0520] e. Prostate Cancer

[0521] The methods provided herein include use of an oncofetal fibronectin indicating molecule as a marker for prostate cancer. An oncofetal fibronectin indicating molecule can be present in subjects that have prostate cancer.

Oncofetal fibronectin indicating molecule measurements can be used as a marker for prostate cancer in any of the cancer (tumor, metastasis or malignant neoplastic) indications provided herein. For example, the oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has cancerous (malignant neoplastic or metastatic) prostate cells. The methods include determining the presence or amount of an oncofetal fibronectin indicating molecule in a sample, and characterizing the sample according to the presence or absence of an oncofetal fibronectin indicating molecule in the sample or according to the amount of an oncofetal fibronectin indicating molecule in the sample, where presence or an amount of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate that a subject has cancerous prostate cells, and absence or an amount of an oncofetal fibronectin indicating molecule below a threshold level can indicate that a subject does not have cancerous prostate cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the risk or lack of risk of prostate cells such as normal, abnormal, dysplastic or hyperplastic prostate cells, becoming cancerous. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the development or lack of development of prostate cells such as normal, abnormal, dysplastic or hyperplastic prostate cells, into cancerous prostate cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for indicating the progression of a prostate cancer. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for distinguishing between aggressive and non-aggressive prostate tumors. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has or is likely to have a recurrence of cancerous prostate cells. The oncofetal fibronectin indicating molecule detection methods provided herein can indicate the likely or actual efficacy or lack of efficacy of prostate cancer treatment.

[0522] A sample collected for prostate cancer determination can be taken from any source, as provided herein. Exemplary sources for samples include a prostate tissue sample, semen, urine, stool, plasma, blood, serum, lymph, lymphatic fluid and interstitial fluid samples. For example, the presence of an oncofetal fibronectin indicating molecule can be detected in prostate tissue samples collected from subjects with prostate cancer. Detection of any oncofetal fibronectin indicating molecule in a sample can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous prostate cells. In another embodiment, detection of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous prostate cells. An exemplary threshold value for an oncofetal fibronectin indicating molecule in a fine needle aspirate sample as an indicator of prostate cancer is 3 ng/ml, 5 ng/ml, 8 ng/ml, 10 ng/ml, 15 ng/ml, 20 ng/ml, 25 ng/ml, or 30 ng/ml, or about 3 ng/ml, about 5 ng/ml, about 8 ng/ml, about 10 ng/ml, about 15 ng/ml, about 20 ng/ml, about 25 ng/ml, or about 30 ng/ml. All forms of prostate cancer can be indicated using

the methods known in the art or provided herein. An exemplary form of prostate cancer includes prostate adenocarcinoma.

[0523] In one embodiment, characterization of an oncofetal fibronectin indicating molecule in a sample can indicate whether or not the oncofetal fibronectin indicating molecule in the sample was produced by cancerous prostate cells. In some cases, one or more forms of oncofetal fibronectin indicating molecule (varying, e.g., at EDA, EDB, IIICS and/or in post-translational modifications) present in a sample such as fine needle aspirate or prostate tissue are known to be produced by cancerous prostate cells. For example, an oncofetal fibronectin indicating molecule in tissue samples from subjects with prostate cancer contained the ED-B region of fibronectin in the samples (see, e.g., Albrecht et al., *Histochem. Cell. Biol.* 112:51-61 (1999)). In other cases, one or more forms of oncofetal fibronectin indicating molecule, although present in a sample such as fine needle aspirate or prostate tissue, are not produced by cancerous prostate cells, but instead are produced by a different tissue or organ source. The methods herein can be used to characterize the oncofetal fibronectin indicating molecule in a sample, and such characterization can indicate whether or not the oncofetal fibronectin indicating molecule observed in the sample is an oncofetal fibronectin indicating molecule form observed in prostate cancer. When a sample contains an oncofetal fibronectin indicating molecule observed in prostate cancer, the presence of the oncofetal fibronectin indicating molecule is consistent with a subject having prostate cancer. When a sample contains an oncofetal fibronectin indicating molecule not observed in prostate cancer, the presence of the oncofetal fibronectin indicating molecule is not consistent with a subject having prostate cancer.

[0524] f. Lung Cancer

[0525] The methods provided herein include use of an oncofetal fibronectin indicating molecule as a marker for lung cancer. An oncofetal fibronectin indicating molecule can be present in subjects that have lung cancer. Oncofetal fibronectin indicating molecule measurements can be used as a marker for lung cancer in any of the cancer (tumor, metastasis or malignant neoplastic) indications provided herein. For example, the oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has cancerous (malignant neoplastic or metastatic) lung cells. The methods include determining the presence or amount of an oncofetal fibronectin indicating molecule in a sample, and characterizing the sample according to the presence or absence of an oncofetal fibronectin indicating molecule in the sample or according to the amount of an oncofetal fibronectin indicating molecule in the sample, where presence or an amount of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate that a subject has cancerous lung cells, and absence or an amount of an oncofetal fibronectin indicating molecule below a threshold level can indicate that a subject does not have cancerous lung cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the risk or lack of risk of lung cells such as normal, abnormal, dysplastic or hyperplastic lung cells, becoming cancerous. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the development or

lack of development of lung cells such as normal, abnormal, dysplastic or hyperplastic lung cells, into cancerous lung cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for indicating the progression of a lung cancer. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for distinguishing between aggressive and non-aggressive lung tumors. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has or is likely to have a recurrence of cancerous lung cells. The oncofetal fibronectin indicating molecule detection methods provided herein can indicate the likely or actual efficacy or lack of efficacy of lung cancer treatment.

[0526] A sample collected for lung cancer determination can be taken from any source, as provided herein. Exemplary sources for samples include a lung tissue sample, sputum, blood, serum and plasma samples. For example, the presence of an oncofetal fibronectin indicating molecule can be detected in sputum collected from subjects with lung cancer. Detection of any oncofetal fibronectin indicating molecule in a sample can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous lung cells. In another embodiment, detection of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous lung cells. An exemplary threshold value for an oncofetal fibronectin indicating molecule in a sputum sample as an indicator of lung cancer is 3 ng/ml, 5 ng/ml, 8 ng/ml, 10 ng/ml, 15 ng/ml, 20 ng/ml, 25 ng/ml, or 30 ng/ml, or about 3 ng/ml, about 5 ng/ml, about 8 ng/ml, about 10 ng/ml, about 15 ng/ml, about 20 ng/ml, about 25 ng/ml, or about 30 ng/ml. All forms of lung cancer can be indicated using the methods known in the art or provided herein. Exemplary forms of lung cancer include small cell carcinoma, adenocarcinoma, squamous cell carcinoma and large cell carcinoma.

[0527] In one embodiment, characterization of an oncofetal fibronectin indicating molecule in a sample can indicate whether or not the oncofetal fibronectin indicating molecule in the sample was produced by cancerous lung cells. In some cases, one or more forms of oncofetal fibronectin indicating molecule (varying, e.g., at EDA, EDB, IIICS and/or in post-translational modifications) present in a sample such as sputum or lung tissue are known to be produced by cancerous lung cells. For example, oncofetal fibronectin in tissue samples from subjects with lung cancer contained the EDB region of fibronectin in the samples (Santimaria et al., *Clin Cancer Res* 9:571-579 (2003)). In other cases, one or more forms of oncofetal fibronectin indicating molecule, although present in a sample such as sputum or lung tissue, are not produced by cancerous lung cells, but instead are produced by a different tissue or organ source. The methods herein can be used to characterize the oncofetal fibronectin indicating molecule in a sample, and such characterization can indicate whether or not the oncofetal fibronectin indicating molecule observed in the sample is an oncofetal fibronectin indicating molecule form observed in lung cancer. When a sample contains an oncofetal fibronectin indicating molecule observed in lung cancer, the presence of the oncofetal fibronectin indicating molecule is consistent with a subject having lung cancer. When a sample contains an oncofetal

fibronectin indicating molecule not observed in lung cancer, the presence of the oncofetal fibronectin indicating molecule is not consistent with a subject having lung cancer.

[0528] g. Colorectal Cancer

[0529] The methods provided herein include use of an oncofetal fibronectin indicating molecule as a marker for colorectal cancer. An oncofetal fibronectin indicating molecule can be present in subjects that have colorectal cancer. Oncofetal fibronectin indicating molecule measurements can be used as a marker for colorectal cancer in any of the cancer (tumor, metastasis or malignant neoplastic) indications provided herein. For example, the oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has cancerous (malignant neoplastic or metastatic) colorectal cells. The methods include determining the presence or amount of an oncofetal fibronectin indicating molecule in a sample, and characterizing the sample according to the presence or absence of an oncofetal fibronectin indicating molecule in the sample or according to the amount of an oncofetal fibronectin indicating molecule in the sample, where presence or an amount of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate that a subject has cancerous colorectal cells, and absence or an amount of an oncofetal fibronectin indicating molecule below a threshold level can indicate that a subject does not have cancerous colorectal cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the risk or lack of risk of colorectal cells such as normal, abnormal, dysplastic or hyperplastic colorectal cells, becoming cancerous. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the development or lack of development of colorectal cells such as normal, abnormal, dysplastic or hyperplastic colorectal cells, into cancerous colorectal cells. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for indicating the progression of a colorectal cancer. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for distinguishing between aggressive and non-aggressive colorectal tumors. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has or is likely to have a recurrence of cancerous colorectal cells. The oncofetal fibronectin indicating molecule detection methods provided herein can indicate the likely or actual efficacy or lack of efficacy of colorectal cancer treatment.

[0530] A sample collected for colorectal cancer determination can be taken from any source, as provided herein. Exemplary sources for samples include a colorectal tissue sample, stool, plasma, blood, serum, lymph, lymphatic fluid and interstitial fluid samples. For example, the presence of an oncofetal fibronectin indicating molecule can be detected in stool collected from subjects diagnosed with colorectal cancer. Detection of any oncofetal fibronectin indicating molecule in a sample can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in treatment of cancerous colorectal cells. In another embodiment, detection of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate the presence of, risk of, development of, progression of, aggressiveness of, recurrence of, or efficacy in

treatment of cancerous colorectal cells. An exemplary threshold value for an oncofetal fibronectin indicating molecule in a stool sample as an indicator of colorectal cancer is 3 ng/ml, 5 ng/ml, 8 ng/ml, 10 ng/ml, 15 ng/ml, 20 ng/ml, 25 ng/ml, or 30 ng/ml, or about 3 ng/ml, about 5 ng/ml, about 8 ng/ml, about 10 ng/ml, about 15 ng/ml, about 20 ng/ml, about 25 ng/ml, or about 30 ng/ml. All forms of colorectal cancer can be indicated using the methods known in the art or provided herein. Exemplary forms of colorectal cancer include mucinous (colloid) adenocarcinoma, signet ring adenocarcinoma, scirrhous tumor, carcinoid tumor, squamous cell tumor, leiomyosarcoma.

[0531] In one embodiment, characterization of an oncofetal fibronectin indicating molecule in a sample can indicate whether or not the oncofetal fibronectin indicating molecule in the sample was produced by cancerous colorectal cells. In some cases, one or more forms of oncofetal fibronectin indicating molecule (varying, e.g., at EDA, EDB, IIICS and/or in post-translational modifications) present in a sample such as stool or colorectal tissue are known to be produced by cancerous colorectal cells. For example, an oncofetal fibronectin indicating molecule in tissue samples from subjects with colorectal cancer contained the EDB region of fibronectin (see, e.g., Midulla et al., *Cancer Res.* 60:164-169 (2000)).

[0532] In other cases, one or more forms of oncofetal fibronectin indicating molecule, although present in a sample such as stool or colorectal tissue, are not produced by cancerous colorectal cells, but instead are produced by a different tissue or organ source. The methods herein can be used to characterize the oncofetal fibronectin indicating molecule in a sample, and such characterization can indicate whether or not the oncofetal fibronectin indicating molecule observed in the sample is an oncofetal fibronectin indicating molecule form observed in colorectal cancer. When a sample contains an oncofetal fibronectin indicating molecule observed in colorectal cancer, the presence of the oncofetal fibronectin indicating molecule is consistent with a subject having colorectal cancer. When a sample contains an oncofetal fibronectin indicating molecule not observed in colorectal cancer, the presence of the oncofetal fibronectin indicating molecule is not consistent with a subject having colorectal cancer.

[0533] h. Additional Cancers

[0534] Methods provided herein can be used to provide information regarding any of a variety of additional cancers. A non-limiting group of exemplary cancer types and samples that can be used to detect an oncofetal fibronectin indicating molecule in providing information regarding the respective cancer types include: renal cancers, as indicated using urine, lymph, lymphatic fluid, blood, serum, plasma, interstitial fluid or tissue samples; thyroid cancers, as indicated using tissue such as fine needle aspirate, lymph, lymphatic fluid, blood, serum, plasma and interstitial fluid; skin cancers, as indicated using interstitial fluid, lymph, lymphatic fluid, blood, serum, plasma and tissue samples; oropharyngeal cancer, as indicated using oropharyngeal swab, blood, serum, plasma and tissue samples; lymphomas, as indicated using lymph, blood, serum, plasma and tissue samples such as lymph node samples; and leukemias, as indicated using plasma, blood or serum.

[0535] Thus, the oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicat-

ing whether or not a subject has cancerous (malignant neoplastic or metastatic) cells of these tissue types. The methods include determining the presence or amount of an oncofetal fibronectin indicating molecule in a sample, and characterizing the sample according to the presence or absence of an oncofetal fibronectin indicating molecule in the sample or according to the amount of an oncofetal fibronectin indicating molecule in the sample, where presence or an amount of an oncofetal fibronectin indicating molecule at or above a threshold level can indicate that a subject has cancerous cells of these tissue types, and absence or an amount of an oncofetal fibronectin indicating molecule below a threshold level can indicate that a subject does not have neoplastic, malignant or metastatic cells of these tissue types. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the risk or lack of risk of cells such as normal, abnormal, dysplastic or hyperplastic cells of these tissue types, of becoming cancerous. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for determining the development or lack of development of cells such as normal, abnormal, dysplastic or hyperplastic cells of these tissue types, into cancerous cells of these tissue types. The oncofetal fibronectin indicating molecule detection methods provided herein also can be used for indicating the progression of cancer in these tissue types. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for distinguishing between aggressive and non-aggressive tumors of these tissue types. The oncofetal fibronectin indicating molecule detection methods provided herein can be used for indicating whether or not a subject has or is likely to have a recurrence of neoplastic, malignant or metastatic cells of these tissue types. The oncofetal fibronectin indicating molecule detection methods provided herein can indicate the likely or actual efficacy or lack of efficacy of cancer treatment of these tissue types.

[0536] 3. Health State Assessment

[0537] The presence or absence of an oncofetal fibronectin indicating molecule in a sample can indicate the health state of an individual. The presence of an oncofetal fibronectin indicating molecule in a sample relative to one or more threshold levels can indicate the severity of the health state of an individual. Detection of an oncofetal fibronectin indicating molecule in a body tissue or fluid sample can be an indicator of a variety of health problems or risk therefor. The presence of, or elevated levels of, an oncofetal fibronectin indicating molecule in a cell, tissue and/or fluid sample does not necessarily indicate that the health problem is caused by an oncofetal fibronectin indicating molecule, but that elevated levels of an oncofetal fibronectin indicating molecule are observed in cell, tissue and/or fluid samples. For example, an oncofetal fibronectin indicating molecule can serve as an indicator of cancer, can serve as an indicator of pre-term or imminent delivery and also can serve as an indicator of arthritis (Kriegsman et al., *Rheumatol Int.* 24:25-33 (2004)), diabetic retinopathy (Khan et al., *Invest. Ophthalmol. Vis. Sci.* 45:287-295 (2004)), renal disease, and Dupuytren's contracture (Howard et al., *J. Surg. Res.* 117:232-238 (2004)). Detection of an oncofetal fibronectin indicating molecule in a body tissue or fluid sample at or above one or more thresholds or at a level above a baseline for a particular individual can be an indicator of a variety of health problems or risk therefor. Similarly, its absence or

presence below one or more thresholds or at a level below a baseline for a particular individual can be indicative of the absence of any of these variety of diseases and disorders.

[0538] Provided herein are methods of screening subjects for the presence of an oncofetal fibronectin indicating molecule in a sample and, if an oncofetal fibronectin indicating molecule is not present in the sample, or is present below a threshold level, concluding that the subject does not have a health problem associated with the presence of an oncofetal fibronectin indicating molecule or with elevated levels of an oncofetal fibronectin indicating molecule. Similarly, provided herein are methods of screening subjects for the presence of an oncofetal fibronectin indicating molecule in a sample and, if an oncofetal fibronectin indicating molecule is present in the sample, or is present at or above a threshold level, determining that the subject has a health problem associated with the presence of an oncofetal fibronectin indicating molecule or with elevated levels of an oncofetal fibronectin indicating molecule.

[0539] In one embodiment, a method is provided for indicating that a subject is free of a health problem associated with oncofetal fibronectin, by testing a sample for the presence or absence of an oncofetal fibronectin indicating molecule, where absence (or presence below a threshold) of an oncofetal fibronectin indicating molecule indicates that the subject is free of a health problem associated with oncofetal fibronectin. In another embodiment, a method is provided for screening a subject for a health problem associated with oncofetal fibronectin, by testing a sample from a subject for the presence or absence of an oncofetal fibronectin indicating molecule, wherein presence of an oncofetal fibronectin indicating molecule indicates that the subject has a health problem associated with oncofetal fibronectin.

[0540] Similarly, general health, or presence or absence of a health problem associated with oncofetal fibronectin can be indicated by an increased rate of change in an amount of an oncofetal fibronectin indicating molecule; for example, increasing amounts of an oncofetal fibronectin indicating molecule can indicate a health problem associated with oncofetal fibronectin, and decreasing amounts of an oncofetal fibronectin indicating molecule can indicate absence of a health problem associated with oncofetal fibronectin. The amount of an oncofetal fibronectin indicating molecule can be compared to one or more thresholds, where increasing thresholds can indicate increased likelihood or increased severity of a health problem associated with oncofetal fibronectin, and decreasing thresholds can indicate decreased likelihood or decreased severity of a health problem associated with oncofetal fibronectin.

[0541] The methods provided herein can be performed in any of a variety of settings or for a variety of purposes, including during a routine physical examination, or as a general diagnostic tool to attempt to determine an unidentified malady or illness of a subject. Baseline levels can be established based on averages in a population or in a particular individual. Deviation from the average or from the baseline in the individual can indicate a change or risk of change in the health status of the individual.

[0542] In some embodiments, prior to performing the test for an oncofetal fibronectin indicating molecule, the subject tested is not diagnosed as having a health problem, or is not

diagnosed as having a health problem associated with oncofetal fibronectin. In other embodiments, a subject can have an unidentified health problem and testing for an oncofetal fibronectin indicating molecule can be used to as a screen to indicate or diagnose the health problem. Thus, provided herein are methods that include performing routine tests of subjects where the tests include determining the presence or absence and/or amount of an oncofetal fibronectin indicating molecule in subjects' samples and determining the health state of the tested subjects according to the presence or absence (or in comparison to one or more thresholds) and/or amount of oncofetal fibronectin indicating molecule detected.

[0543] In one embodiment, detection of an oncofetal fibronectin indicating molecule can be performed in conjunction with one or more additional diagnostic tests, including routine diagnostic tests such as blood pressure, pulse, body weight, health history, family history or sample tests. In another embodiment, if an oncofetal fibronectin indicating molecule is present or is at or above a threshold level, one or more additional diagnostic tests can be conducted to diagnose the health problem of the subject. Such diagnostic tests can be conducted prior to, at the same time as, or subsequent to, testing for an oncofetal fibronectin indicating molecule. In one example, a subject can be identified as having an oncofetal fibronectin indicating molecule present or at or above a threshold level and the subject can then be further tested in one or more subsequent diagnostic tests to identify the health problem of the subject.

[0544] In one embodiment, a sample can be tested for the presence or absence (or in comparison to one or more thresholds) and/or amount of an oncofetal fibronectin indicating molecule and also can be tested for one or more additional properties, including the presence of one or more additional sample components. Collection of samples from a subject and screening the sample for properties such as the presence or absence of a plurality of components such as ions or molecules in the sample or for the level of components such as ions or molecules in the sample is well known in the art. For example, blood sample collection can be used to determine ion content such as sodium ion content, lipid content such as LDL and HDL content and urine samples can be tested for the presence of metabolites or sugar and saliva can be tested for hormones. Any of a variety of samples can be used to measure the presence and/or amount of any of a variety ions or molecules, as is known in the art.

[0545] In one example blood can be tested for an oncofetal fibronectin indicating molecule and one or more additional properties. Exemplary properties of blood that are routinely tested include, but are not limited to, red blood cell count, white blood cell count (including count of neutrophils, lymphocytes, T cells, B cells, monocytes, eosinophils and basophils), platelet count, hematocrit, hemoglobin, blood type, Rh factor, glucose, lactose dehydrogenase, creatine phosphokinase, blood urea nitrogen, creatinine, carbon dioxide, sodium, potassium, chloride, calcium, phosphorus, alkaline phosphatase, alanine amino transferase, aspartate amino transferase, albumin, gamma-glutamyl transpeptidase, serum glutamate pyruvate transaminase, total protein, fibrinogen, prothrombin, cholesterol, globulin, bilirubin, high density lipoproteins, low density lipoproteins, very low density lipoproteins, free testosterone, total testosterone,

dehydroepiandrosterone, prostate-specific antigen, estradiol, progesterone, homocysteine, C-reactive protein, uric acid, amylase and lipase.

[0546] In another example urine can be tested for an oncofetal fibronectin indicating molecule and one or more additional properties. Exemplary properties of urine that are routinely tested include, but are not limited to, color, appearance, specific gravity, pH, total protein, glucose, ketones, hemoglobin, bile, urobilinogen, nitrate, uroglobin, white blood cells, leukocytes, red blood cells, epithelial cells, bacteria, crystals, mucus and casts. In another example saliva can be tested for an oncofetal fibronectin indicating molecule and one or more additional properties. Exemplary properties of saliva that are routinely tested include, but are not limited to, estradiol, testosterone, DHEA-S, cortisol, sodium, potassium, chloride.

[0547] In one embodiment, an oncofetal fibronectin indicating molecule is determined as a component of a sample panel. For example, an oncofetal fibronectin indicating molecule can be determined as a component of a blood panel, urine panel or saliva panel.

[0548] 4. Other Health Problems

[0549] An oncofetal fibronectin indicating molecule also can be present in subjects with other health problems such as arthritis, diabetic retinopathy and Dupuytren's contracture. In some embodiments, the presence of an oncofetal fibronectin indicating molecule can indicate the presence of health problems such as arthritis, diabetic retinopathy and Dupuytren's contracture. In other embodiments, the presence of an oncofetal fibronectin indicating molecule can indicate the risk of developing health problems such as arthritis, diabetic retinopathy and Dupuytren's contracture. In yet other embodiments, the presence of an oncofetal fibronectin indicating molecule can indicate the severity of health problems such as arthritis, diabetic retinopathy and Dupuytren's contracture. In addition to the presence of an oncofetal fibronectin indicating molecule indicating presence, risk of developing or severity of such health problems, an amount of an oncofetal fibronectin indicating molecule in a sample can be greater than, equal to, or less than one or more thresholds, where each increasing threshold indicates an increased likelihood of the presence, an increased risk of developing, or increased severity of such health problems, relative to each lower threshold. The rate of increase or decrease of an oncofetal fibronectin indicating molecule in a sample can indicate the degree of likelihood of the presence, an degree of the risk of developing, or the degree of the severity of such health problems, where larger increases represent more likely or more severe health problems relative to smaller increases or decreases.

[0550] a. Arthritis

[0551] An oncofetal fibronectin indicating molecule can be present in arthritic joints. An oncofetal fibronectin indicating molecule can be present in the synovial fluid, synovial tissue and/or cartilage of individuals with arthritis. An oncofetal fibronectin indicating molecule can be present at higher levels in subjects having rheumatoid arthritis relative to a control sample having an amount of oncofetal fibronectin indicating molecule below a threshold. An oncofetal fibronectin indicating molecule can be present at higher levels in subjects having osteoarthritis relative to a control

sample having an amount of oncofetal fibronectin indicating molecule below a threshold. An oncofetal fibronectin indicating molecule can be present at higher levels in subjects with rheumatoid arthritis relative to the levels of oncofetal fibronectin indicating molecule present in subjects with osteoarthritis.

[0552] Arthritic subjects can have elevated levels of an oncofetal fibronectin indicating molecule in the affected areas. Thus, in one embodiment, presence of arthritis can be determined by detecting an oncofetal fibronectin indicating molecule in synovial fluid, synovial tissue or cartilage of a subject. Arthritic subjects also can have elevated levels of an oncofetal fibronectin indicating molecule in their bloodstream. Thus, presence of arthritis also can be determined by detecting an oncofetal fibronectin indicating molecule in blood, serum or plasma.

[0553] Presence of an oncofetal fibronectin indicating molecule also can be indicative of the risk of a subject developing arthritis. Thus, provided herein are methods for determining the risk of a subject developing arthritis by testing for the presence of an oncofetal fibronectin indicating molecule in a sample, where presence of an oncofetal fibronectin indicating molecule indicates an increased risk of developing arthritis. Presence of an oncofetal fibronectin indicating molecule also can indicate the severity of the arthritis of a subject. Thus, provided herein are methods for determining the severity of arthritis in a subject by testing for the presence of an oncofetal fibronectin indicating molecule in a sample, where presence of an oncofetal fibronectin indicating molecule at or above a threshold indicates an increased severity of arthritis in the subject relative to a sample that is oncofetal fibronectin negative (or below the threshold).

[0554] Presence of an oncofetal fibronectin indicating molecule can be measured by determining the presence of EDA+, EDB+ and/or IIICS+ oncofetal fibronectin indicating molecule in a sample. In some embodiments, the presence of EDB in synovial tissue indicates the presence of rheumatoid arthritis in a subject. In other embodiments, presence of a splice variant of IIICS, such as a IIICS splice variant that binds FDC-6, indicates the presence of synovial hyperplasia.

[0555] b. Diabetic Retinopathy

[0556] An oncofetal fibronectin indicating molecule can be present in diabetic retinopathy. An oncofetal fibronectin indicating molecule can be present in the aqueous humor, vitreous humor, or various tissues of the eye. Subjects with diabetic retinopathy can have elevated levels of EDA+ oncofetal fibronectin indicating molecule relative to subjects without diabetic retinopathy. Subjects with diabetic retinopathy can have elevated levels of EDB+ oncofetal fibronectin indicating molecule relative to subjects without diabetic retinopathy. Subjects with diabetic retinopathy can have elevated levels of IIICS+ oncofetal fibronectin indicating molecule relative to subjects without diabetic retinopathy. Thus, methods for detecting an oncofetal fibronectin indicating molecule associated with diabetic retinopathy can include detecting EDA+, EDB+ and/or IIICS+ oncofetal fibronectin indicating molecule.

[0557] In one embodiment, presence of diabetic retinopathy can be determined by detecting an oncofetal fibronectin indicating molecule in the aqueous humor, vitreous humor,

or eye tissue sample of a subject. Subjects with diabetic retinopathy also can have elevated levels of an oncofetal fibronectin indicating molecule in their bloodstream. Thus, presence of diabetic retinopathy also can be determined by detecting an oncofetal fibronectin indicating molecule in blood, serum or plasma.

[0558] Presence of an oncofetal fibronectin indicating molecule also can be indicative of the risk of a subject developing diabetic retinopathy. Thus, provided herein are methods for determining the risk of a subject developing diabetic retinopathy by testing for the presence of an oncofetal fibronectin indicating molecule in a sample, where presence of an oncofetal fibronectin indicating molecule indicates an increased risk of developing diabetic retinopathy. Presence at or above one or more thresholds of an oncofetal fibronectin indicating molecule also can indicate the severity of the diabetic retinopathy of a subject. Thus, provided herein are methods for determining the severity of diabetic retinopathy in a subject by testing for the presence of an oncofetal fibronectin indicating molecule in a sample, where presence at or above one or more thresholds of an oncofetal fibronectin indicating molecule indicates an increased severity of diabetic retinopathy the subject relative to a sample that is oncofetal fibronectin negative (or below the threshold).

[0559] c. Dupuytren's Contracture

[0560] An oncofetal fibronectin indicating molecule can be present in Dupuytren's contracture. An oncofetal fibronectin indicating molecule can be present in the blood, serum, plasma or tissue sample of subjects with Dupuytren's contracture. Subjects with Dupuytren's contracture can have elevated levels of IIICS+ oncofetal fibronectin indicating molecule relative to subjects without Dupuytren's contracture. Thus, methods for detecting an oncofetal fibronectin indicating molecule associated with Dupuytren's contracture can include detecting a IIICS+ oncofetal fibronectin indicating molecule.

[0561] In one embodiment, presence of Dupuytren's contracture can be determined by detecting an oncofetal fibronectin indicating molecule in tissue samples from an area suspected of having Dupuytren's contracture. Subjects with Dupuytren's contracture also can have elevated levels of an oncofetal fibronectin indicating molecule in their bloodstream. Thus, presence of Dupuytren's contracture also can be determined by detecting an oncofetal fibronectin indicating molecule in blood, serum or plasma.

[0562] Presence of an oncofetal fibronectin indicating molecule also can be indicative of the risk of a subject developing Dupuytren's contracture. Thus, provided herein are methods for determining the risk of a subject developing Dupuytren's contracture by testing for the presence of an oncofetal fibronectin indicating molecule in a sample, where presence of an oncofetal fibronectin indicating molecule indicates an increased risk of developing Dupuytren's contracture. Presence of an oncofetal fibronectin indicating molecule at or above one or more thresholds can also indicate the severity of the Dupuytren's contracture of a subject. Thus, provided herein are methods for determining the severity of Dupuytren's contracture in a subject by testing for the presence of an oncofetal fibronectin indicating molecule in a sample, where presence of an oncofetal fibronectin indicating molecule at or above one or more

thresholds indicates an increased severity of Dupuytren's contracture in the subject relative to a sample that is oncofetal fibronectin negative (or below the threshold).

E. Collection of Samples

[0563] In accordance with the methods disclosed herein, an oncofetal fibronectin indicating molecule can be detected in any of a variety of types of samples. For example, the sample can include urine, lymph, blood, plasma, serum, saliva, cervical fluid, cervicovaginal fluid, vaginal fluid, breast fluid, breast milk, synovial fluid, semen, seminal fluid, stool, sputum, cerebral spinal fluid, tears, mucus, interstitial fluid, follicular fluid, amniotic fluid, aqueous humor, vitreous humor, peritoneal fluid, ascites, sweat, lymphatic fluid, lung sputum and lavage. In addition, the sample can contain tissues specimens such as a biopsy. When a sample contains solid material, such as a tissue biopsy, the sample can be homogenized in order to bring into solution or otherwise increase the accessibility of sample components for use in the methods provided herein or otherwise known in the art. Exemplary tissue biopsy samples include cervicovaginal tissue and breast tissue biopsy samples.

[0564] The sample can be collected by any of a variety of techniques. The particular technique used for a given procedure will depend, at least in part, upon the type of sample to be analyzed. In general, tissue samples can be collected using aspiration (e.g., fine needle aspiration), lavage (e.g., ductal lavage), biopsy, swabbing (using, e.g., a fibrous tipped swab such as a cytobrush, polyester swab, rayon swab or cotton swab), suction, transcutaneous or transdermal extraction and other methods. Liquid samples can be collected by suction, needle-mediated withdrawal, swabbing (using, e.g. a fibrous tipped swab such as a cytobrush, polyester swab, rayon swab or cotton swab) and other methods. When the sample is collected with a cotton swab, the methods provided herein are conducted on the swab itself. As will be recognized by one skilled in the art, depending on the sample, sample collection can be performed by a medical professional, an untrained individual and/or the subject from whom the sample is to be collected. For example, a biopsy or fine needle aspirate sample is likely to be collected by a medical professional. In another example, a urine sample or a vaginal swab sample can be collected by an untrained individual, such as a family member, or by the subject from whom the sample is to be collected. Samples that can be collected by an untrained individual or by the subject from whom the sample is to be collected, can be collected at a site other than a clinical setting, such as the home. For example, a sample can be collected as part of a home testing procedure. Home testing procedures can be performed, for example, using a home testing kit, such as a kit provided herein.

[0565] Any of the sample collection techniques provided herein can be used in conjunction with any of the oncofetal fibronectin indicating molecule detection methods provided herein or otherwise known in the art for any of the diagnostic uses or other uses of detection of an oncofetal fibronectin indicating molecule provided herein or otherwise known in the art. The following are exemplary collection methods and sources.

[0566] 1. Swab and Cervicovaginal Samples

[0567] A swab sample can be collected from a subject and tested for the presence of an oncofetal fibronectin indicating

molecule. Swab samples can contain body fluids of the subject, cells of the subject, or body fluids and cells. Swab samples can be collected from any of a variety of regions of the subject, including, but not limited to, oral, aural, nasal, anal, urethral, cervicovaginal, ocular, skin, alimentary canal such as esophageal, gastric, intestinal, colon, or any other surface accessible to a swab, or lesions of any of the above.

[0568] Swab samples can be collected by a medical professional, an untrained individual such as a family member, or the subject who is providing the sample, according to the sample to be collected and the oncofetal fibronectin indicating molecule test to be performed. For example, a swab of the cervical os to be tested for the presence of an oncofetal fibronectin indicating molecule by mass spectrometry is typically collected by a medical professional, whereas a vaginal and/or labial swab sample to be tested for the presence of an oncofetal fibronectin indicating molecule by a test strip assay can be collected by an untrained individual or by the subject providing the sample and can be used in, for example, a home testing method. Devices that can be used in collecting swab samples can be any swab sample collection device known in the art, including, but not limited to, fibrous tipped swabs such as a cytobrush, polyester swab, rayon swab or cotton swab, configured to facilitate sample collection from the targeted body region. In one embodiment, the swab sample is a cervicovaginal swab sample.

[0569] Cervicovaginal samples, including cervicovaginal swab samples, can include samples from any of a variety of cervicovaginal regions and combinations thereof. Cervicovaginal samples can contain cervicovaginal fluid and can optionally contain cells from the cervicovaginal cavity. Cervicovaginal samples such as cervicovaginal fluid can be collected by any of a variety of methods, including cervicovaginal swabbing, or collecting cervicovaginal fluid leakage, for example, using an absorptive collection vehicle such as an absorptive pad. Exemplary cervicovaginal swab samples include, but are not limited to, a swab of the point of a possible cervicovaginal lesion, the cervical canal, the cervical os, the ectocervix, the transition zone on the cervix between squamous and columnar cells (i.e., the squamocolumnar junction), the vagina, the posterior fornix, the portion of the vagina below the posterior fornix such as the lower third of the vagina, the labia, or combinations thereof. In the case of a vaginal swab sample, the sample can be a swab of any portion of the vagina, including the posterior fornix or the portion of the vagina below the posterior fornix, such as, for example, the lower third of the vagina. In the case of a labial swab, the swab can be collected from the labia minora or labia majora and typically includes a swab of the labia minora.

[0570] With respect to cervicovaginal samples in general, a tissue or liquid sample to be assayed can be removed in the vicinity of the point of a possible cervicovaginal lesion, the cervical canal, the cervical os, the ectocervix, the transition zone on the cervix between squamous and columnar cells (i.e., the squamocolumnar junction), the vagina, the posterior fornix, the portion of the vagina below the posterior fornix such as the lower third of the vagina, the labia, or combinations thereof. Cervicovaginal samples also can include culdocentesis samples. Cervicovaginal samples include samples collected by passive collection methods. Passive collection methods include collecting cervicovaginal fluid and, optionally particulate matter such as cells by

placing a sample collection vehicle in a position that will contact and typically absorb the fluid and optionally particulate matter. Exemplary passive collection vehicles can include a device for collecting a sample that is inserted into the cervicovaginal cavity (e.g., a tampon-like device that can collect a cervicovaginal sample) and a device for collecting a sample that can collect the sample as it exits the cervicovaginal cavity (e.g., an absorbent pad-like device such as a sanitary napkin-like device). Use of passive collection devices such as modified sanitary napkins, for diagnostic purposes is known in the art, as exemplified in Alary et al., *J. Clin. Microbiol.* 39:2508-2512 (2001).

[0571] In some embodiments, as provided herein, the amount of an oncofetal fibronectin indicating molecule in a cervicovaginal swab sample collected from the portion of the vagina below or inferior to the posterior fornix, such as the lower third of the vagina, can be one-third or about one-third the amount of oncofetal fibronectin indicating molecule in a cervicovaginal swab of the posterior fornix collected from the same subject. Accordingly, in methods provided herein in which the level of an oncofetal fibronectin indicating molecule in a sample is compared to a threshold level, the threshold level for a swab of the lower portion of the vagina, such as the lower third of the vagina, can be one-third or about one-third of the threshold level for a swab of the posterior fornix. For example, when the threshold level for a buffer-treated swab of the posterior fornix is 60 ng/ml (or 600 ng/ml for an untreated sample), or about 60 ng/ml (or about 600 ng/ml for an untreated sample), the threshold level of a buffer-treated swab of the lower portion of the vagina such as the lower third of the vagina can be 20 ng/ml (or 200 ng/ml for an untreated sample) or about 20 ng/ml (or about 200 ng/ml for an untreated sample). Similarly, when the threshold level for a buffer-treated swab of the posterior fornix is 50 ng/ml (or 500 ng/ml for an untreated sample) or about 50 ng/ml (or about 500 ng/ml for an untreated sample), 30 ng/ml (or 300 ng/ml for an untreated sample) or about 30 ng/ml (or about 300 ng/ml for an untreated sample), 15 ng/ml (or 150 ng/ml for an untreated sample) or about 15 ng/ml (or about 150 ng/ml for an untreated sample), or 10 ng/ml (or 100 ng/ml for an untreated sample) or about 10 ng/ml (or about 100 ng/ml for an untreated sample), the threshold level of a buffer-treated swab of the lower portion of the vagina such as the lower third of the vagina can respectively be 15-20 ng/ml (or 150-200 ng/ml for an untreated sample) or about 15-20 ng/ml (or about 150-200 ng/ml for an untreated sample), 10 ng/ml (or 100 ng/ml for an untreated sample) or about 10 ng/ml (or about 100 ng/ml for an untreated sample), 5 ng/ml (or 50 ng/ml for an untreated sample) or about 5 ng/ml (or about 50 ng/ml for an untreated sample), or 3-4 ng/ml (or 30-40 ng/ml for an untreated sample) or about 3-4 ng/ml (or about 30-40 ng/ml for an untreated sample).

[0572] A cervicovaginal sample generally includes fluid and particulate solids and can contain vaginal or cervical mucus, other vaginal or cervical secretions, cells or cell debris, amniotic fluid, or other fetal or maternal materials.

[0573] In some of the methods provided herein, the sample essentially does not contain blood. For example, when the method is an immunoassay such as an ELISA assay or lateral flow, the samples essentially do not contain blood. The sample contains 5% or less, 2% or less, 1% or less, 0.5% or less, or 0.1% or less, or about 5% or less, about

2% or less, about 1% or less, about 0.5% or less, or about 0.1% or less blood. The sample can be removed using any of a variety of techniques including, but not limited to, use of a fibrous tipped swab such as a cytobrush, polyester swab, rayon swab or cotton swab (see, e.g., WO 91/16855, WO 89/10724, U.S. Pat. Nos. 4,759,376, 4,762,133 and 4,700,713), aspirator, suction device, lavage device, needle, or other devices known in the art. In other methods, such as immunoprecipitation, Western blots, dot blots, etc., the assay method is not affected by the presence of blood in the sample. One of ordinary skill in the art can empirically determine whether or not blood would be a contaminant based on the assay method.

[0574] Cervicovaginal sample collection can be performed according to the cervicovaginal region to be sampled. For example, a swab of transition zone between squamous and columnar cells of cervix can be performed by a medical professional with the aid of a vaginal speculum. In another example, a vaginal swab, such as a swab of the lower third of the vagina and/or a swab of the labia, can be performed by the subject herself, by an untrained individual such as a family member, or by a medical professional.

[0575] In some embodiments directed to vaginal samples, vaginal samples collected at the same location in the vagina can increase the reproducibility of sample collection, and can increase the reliability of the results of oncofetal fibronectin indicating molecule measurements. As provided herein, the concentration of oncofetal fibronectin indicating molecule in the vagina can vary at different locations in the vagina. Accordingly, two or more samples collected from the same location in the vagina have an increased likelihood of containing the same or substantially the same concentration of oncofetal fibronectin indicating molecule relative to two or more samples collected from different locations in the vagina. Thus, provided herein are methods for collecting vaginal samples or methods for determining the presence and/or amount of oncofetal fibronectin indicating molecule in a vaginal sample, where the methods include collecting two or more vaginal samples (from the same or different subjects), where the samples are collected from the same location in the vagina. Two or more samples collected from the same location in the vagina and collected from the same subject on different occasions (e.g., on different days or weeks) can more reliably indicate the change in presence and/or amount of oncofetal fibronectin indicating molecule in the subject over time relative to two or more samples collected from different locations in the vagina of the same subject. Accordingly, the methods provided herein include methods for increasing the reliability of a measured change in presence and/or amount of oncofetal fibronectin indicating molecule in a subject over time by collecting two or more samples from the same location in the vagina of the same subject. Also provided herein, a sample collected from a particular location in the vagina can be compared to one or more thresholds that specifically relate to samples collected at that location; such a sample equal to or greater than such one or more thresholds can more reliably reflect the health condition or likely outcome of the subject relative to samples and threshold that do not relate to a particular location in the vagina. Accordingly, the methods provided herein include methods for increasing the reliability of an indicated health condition or likely health condition outcome by collecting a vaginal sample from the same location in the vagina as the location the location specifically related

to by one or more thresholds. Also provided herein, collecting samples from the same location in the vagina can reduce the variability of oncofetal fibronectin indicating molecule amounts present in a sample due to variations in sample collection procedures (e.g., variation between individuals collecting the sample) or sample collection techniques (e.g., improper or careless sample collection techniques). Reduced variability of oncofetal fibronectin indicating molecule amounts due to sample collection can increase the correlation of presence/absence and/or amount of oncofetal fibronectin indicating molecule with a health condition or likely health condition outcome. Accordingly, provided herein are methods of correlating a health condition or health outcome in a subject with presence/absence and/or amount of oncofetal fibronectin indicating molecule in samples by collecting samples from the same location in the vagina of two or more subjects and correlating a health condition or likely health condition outcome with the presence/absence and/or amount of oncofetal fibronectin indicating molecule in the samples. Methods for collecting a vaginal sample from the same location in the vagina are known in the art, and can be accomplished, for example using a swab sample collection device with an over-insertion preventing device attached thereto, where the over-insertion preventing device standardizes the location in the vagina at which the sample is collected.

[0576] Following collection, the sample can be transferred to a container for storage and transport to a testing laboratory. The test sample is optionally dispersed in a liquid that preserves biomolecule analytes such as proteins or nucleic acids that can be unstable in the sampled composition. The storage and transfer medium minimizes decline in the protein analyte level during storage and transport. For example, the storage and transfer medium can contain reagents or conditions (e.g., pH, ionic strength or ionic composition) that decrease, inhibit or prevent degradative enzyme activity such as protease or nuclease activity. An exemplary preserving solution for storage and transfer contains of 0.05 M Tris buffer, pH 7.4, 150 mM NaCl, 0.02% NaN_3 , 1% BSA, 5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin, and 0.1% Triton X-100, as known in the art and exemplified in U.S. Pat. No. 4,919,889, issued Apr. 24, 1990. The solution can be used, for example, when detecting an oncofetal fibronectin indicating molecule. Calculations to account for any additional dilution of the samples collected using liquids can be performed as part of the interpretation of the assay procedure.

[0577] In one embodiment, home and office use devices for immediate processing of the sample can be used. If used, the sample can be placed directly in the device and testing can be performed within minutes of sample collection. In such cases, the need to stabilize the analyte is minimized and any solution that facilitates performing the assay and is not detrimental to analyte stability or user safety can be used. An exemplary solution for home or office use in immediate processing contains of 0.05 M Tris-HCl, pH 7.4; 0.15 M NaCl, 1% BSA and 5 mM EDTA.

[0578] In one embodiment, a kit for home testing of a cervicovaginal sample is provided. The kit can contain a sample collection device, such as a swab or a passive sample collection vehicle and optionally a solution for mixing with the sample and typically contains one or more fibronectin or

oncofetal fibronectin binding partners and instructions for use and/or interpretation of results.

[0579] A variety of diagnostic systems and kits are provided herein and are known in the art, such as those exemplified in U.S. Pat. Nos. 6,394,952 and 6,267,722. Such diagnostic systems and kits can be used to determine the level of an oncofetal fibronectin indicating molecule in the sample, in accordance with the methods provided herein or otherwise known in the art and can be used for any of the diagnostic purposes provided herein or otherwise known in the art.

[0580] 2. Lavage Samples

[0581] An oncofetal fibronectin indicating molecule can be present in or on a variety of regions in a subject's body. Samples of fluids, cells, or other matter from the subject that can contain an oncofetal fibronectin indicating molecule can be gathered from a variety of regions, including cavities and ducts, using lavage methods. Thus, as provided herein, lavage samples can be collected from a subject, where the lavage sample can be tested for any oncofetal fibronectin indicating molecule therein. Any of a variety of body surfaces, cavities and/or ducts can be used for collection of a lavage sample. Exemplary lavage samples include peritoneal, ductal, bronchial, bronchoalveolar, oral, nasal, ear, eye, bladder, colonic, gastric, cervicovaginal lavage samples. Any of a variety of lavage methods and apparatuses known in the art can be used for collecting a lavage sample. Lavage methods generally include contacting a region of the subject's body with a fluid and collecting the fluid. The methods also can include a step of washing or moving the fluid over the region. The methods also can include, but do not require, step of aspirating the fluid or applying a vacuum to collect the fluid.

[0582] a. Sample Collection

[0583] Lavage samples can be collected using known methods. Generally, a lavage tool is used to probe and/or collect a lavage sample from a region of the subject's body. Once the region of interest is located, a lavage fluid carried in the lavage tool can be contacted with the region. At least a portion of the lavage fluid can then be collected to obtain a sample.

[0584] In one embodiment, the access tool contains a double lumen catheter. The lavage fluid can be contacted with the region of the subject's body through one of the catheter lumens, and lavage fluid can be removed through the second catheter lumen, and collected. Collection of lavage fluid can further include application of suction to the second catheter lumen to facilitate withdrawal of the lavage fluid from the region of the subject's body and accumulation of the sample in an assayable form. Suction can be applied using any device for creating suction in a lumen. For example, suction can be applied using a syringe or other suction device operatively coupled to the lumen through which lavage fluid will be withdrawn the region of the subject's body. The suction can be applied for a brief period of time or for an extended period of time, according to the procedure intended by one skilled in the art. Introduction of the lavage fluid optionally continues even after the initial portions of the fluid begin to emerge from the second catheter lumen.

[0585] The volume of lavage fluid used can be any amount in which an oncofetal fibronectin indicating molecule can be

collected, and can vary according to a variety of factors, including, but not limited to, the region of the subject to be contacted, the measurement method, and any sample manipulation methods to be used, as is understood by those of skill in the art. Typically the volume will be at least enough to carry fluid or cells or other components contacted by the fluid that can contain an oncofetal fibronectin indicating molecule and be removed from the subject for oncofetal fibronectin indicating molecule measurement. Typically the volume will not be greater than an amount that can dilute any oncofetal fibronectin indicating molecule in a sample such that the oncofetal fibronectin indicating molecule cannot be detected by the selected oncofetal fibronectin indicating molecule measurement method. Exemplary volumes are at least 0.5 mL or about 0.5 mL, and as much as 25 mL or about 25 mL.

[0586] The lavage fluid can be introduced onto the surface of the region, or into a body cavity at a pressure low enough that the lavage method will not disrupt tissues, organs or membranes of the body region, but at a pressure high enough that fluid, cells and/or other material can be separated from the body region and carried by the lavage fluid. For example, the fluid can be introduced at a rate in the range of between 0.1 mL/s and 5 mL/s or about 0.1 mL/s and about 5 mL/s.

[0587] b. Lavage Fluid

[0588] The lavage fluid can contain any of a variety of components known in the art. The lavage fluid can contain H₂O, alcohol, or other liquid compatible with contacting a subject and detecting a oncofetal fibronectin indicating molecule. For example, the sample can be aqueous and contain saline and one or more optional ingredients such as, for example, an anesthetic, an oncotic agent, an osmotic agent, hormone, cytokine, chemokine, an antiseptic agent, an orifice dilating agent, a vasodilator, a vasoconstrictor, a muscle relaxant, a muscle constrictor, an anti-ischemic agent, a beta-blocker, a calcium channel blocker, or a dye or stain. The lavage fluid also can optionally contain one or more gases (e.g., air and/or nitrogen). The presence of gas can serve to increase retrieval of cells and fluid. The gas can be introduced into the lavage fluid by any of a variety of standard methods, including introduction of the gas from a pressurized container.

[0589] The anesthetic is any anesthetic agent capable of anesthetizing a region of the subject. The anesthetic can act topically, systemically, locally, or any combination thereof. The anesthetic can include, but is not limited to, the following: lidocaine, procaine, prevericaine, or marcaine. The anesthetic also can be a combination or mixture of anesthetic agents.

[0590] The oncotic agent includes, but is not limited to, existing commercially available sterile oncotic solutions, such as preparations of high molecular weight hydroxyethyl starch (e.g., Hespan (DuPont)) and low molecular weight hydroxyethyl starch (e.g., Pentaspan (DuPont)). Other polysaccharide derivatives, including hydroxymethyl alpha substituted (1-4) or (1-6) polymers and cyclodextrins, including hydroxypropyl substituted β or γ cyclodextrin, also can be used as oncotic agents.

[0591] Osmotic agents are known in the art. Exemplary osmotic agents include, but are not limited to, osmopolymers and osmagents. A variety of oncotic agents are known in the art, as exemplified in U.S. Pat. No. 5,413,572.

[0592] The antiseptic agent is any agent that can reduce an opportunity for sepsis at the region of the subject contacted with the lavage fluid. The antiseptic agent can serve a prophylactic purpose in preventing or forestalling sepsis. The antiseptic agent can be, but is not limited to, one or more of the following: a medicinal alcohol (e.g., ethyl alcohol or isopropyl alcohol), a topical antibiotic (e.g., Neosporin or bacteriomycin) and combinations thereof.

[0593] The orifice dilating agent is an agent that promotes dilation of an orifice such as a ductal orifice. The orifice dilating agent can include, but is not limited to, one or more of the following: an agent from the red pepper family of plants (genus *Capsicum*, where the agent can be, for example, capsaicin), a hormone capable of prompt or delayed reaction at the orifice (e.g., prolactin or oxytocin) and combinations thereof.

[0594] The vasodilator is any agent that encourages vasodilation, or opening of blood vessels to increase blood flow to and within the region contacted. The vasodilator can include, but is not limited to, one or more of the following: a vasodilator typically used in cardiac contexts; any vasodilator that can work at the surface or orifice; and combinations thereof.

[0595] The muscle relaxant is any agent that can cause relaxation of muscles in or near the body region to be sampled, such as a sphincter. The muscle relaxant can include, but is not limited to, one or more of the following: a smooth muscle relaxing agent, a calcium channel blocker (e.g., nifedipine), an antispasmodic (e.g., ditropan (oxybutinin), urospas, or terbutyline)) and combinations thereof. For example, the muscle relaxant can include a sphincter relaxer that is effective in relaxing sphincter muscle (e.g., breast duct sphincter muscle). The muscle constricting agent is any agent that can cause constriction of muscles found in or near the body region to be sampled, such as a sphincter.

[0596] The lactation stimulating agent is any agent that can stimulate lactation in a lactating woman. For example, an agent, as applied to a nipple surface and breast of a non-lactating woman is believed to act to increase the ductal fluid collectable from the breast duct. The lactation stimulating agent can include, but is not limited to, one or more of the following: oxytocin, prolactin and combinations thereof.

[0597] The secretion stimulating agent is any agent that can stimulate secretion of fluids and/or materials from the body region to be sampled. The secretion stimulating agent can include, but is not limited to, one or more of the following: oxytocin, prolactin and combinations thereof.

[0598] The anti-ischemic agent is any agent that can prevent or reducing ischemia. The anti-ischemic agent can work in a variety of ways to achieve the anti-ischemic effect and use of the agent is not limited by its mode of action. The anti-ischemic agent can act to increase blood and oxygen flow to the body region to be sampled.

[0599] The beta-blocker is any beta-blocker that can act effectively on a body region to be sampled (e.g., a breast) to increase blood and oxygen flow to the body region. The calcium channel blocker is any calcium channel blocker that can act effectively on a body region to be sampled (e.g., a breast) to increase blood and oxygen flow to the body region.

[0600] The dye or stain is any agent that can be used to identify a body region contacted by the lavage fluid.

[0601] The ductal lavage sample obtained can contain epithelial cells from the body region to be sampled, normally secreted and non-secreted fluids present in the region and proteins, peptides, nucleic acid molecules, antibodies, and other chemical species which can be secreted or otherwise present in the body region to be sampled.

[0602] c. Applying a Label with Lavage

[0603] Optionally, lavage can be performed in detection and/or therapeutic methods. Identification of an oncofetal fibronectin indicating molecule on body regions and treatment of body regions containing an oncofetal fibronectin indicating molecule can be facilitated by including in the lavage fluid a detectable and/or therapeutic fibronectin or oncofetal fibronectin binding partner conjugate. The surface of the region of the subject's body can be contacted with the lavage fluid, and the conjugate can be detected and/or can have a therapeutic effect on the region to which the conjugate bound. In some embodiments, the body region can be optionally washed with a solution that blocks or unblocks the body region to facilitate or permit specific binding of the binding partner conjugate to an oncofetal fibronectin indicating molecule in the body region. For example, the orifice can become plugged with keratin-containing materials and washing with a keratinolytic solution, such as acetic acid (e.g., 5% to 50% or about 5% to about 50% by weight) admixed in a pharmaceutical delivery vehicle, can expose sufficient oncofetal fibronectin indicating molecule sites to permit binding partner binding. The detectable or therapeutic conjugates can be formulated as liquids (e.g., aqueous solutions) using any of a variety of conventional techniques. For example, a fibronectin or oncofetal fibronectin binding partner conjugate can be in an aqueous solution. A variety of detectable or therapeutic conjugates containing a fibronectin or oncofetal fibronectin binding partner or a fibronectin or oncofetal fibronectin binding partner are provided herein or are known in the art.

[0604] d. Ductal Lavage

[0605] For exemplary purposes, ductal lavage methods are provided herein. Ductal lavage can be used to collect samples from a duct, or to apply a label to a duct. One skilled in the art will select the ductal lavage method according to the selected an oncofetal fibronectin indicating molecule detection method and guidance provided by the teachings herein.

[0606] i. Sample Collection

[0607] Samples can be collected from a duct using known methods. Generally, a ductal access tool is used to probe a surface in search of a ductal orifice. Once a ductal orifice is located, a lavage fluid carried in the ductal access tool can be infused into the duct. The lavage fluid can prepare the ductal orifice and duct system for access and fluid and/or can be used for material collection from the duct. In one embodiment, the lavage fluid is introduced into the duct so that it passes substantially throughout the entire ductal network. At least a portion of the lavage fluid can then be collected from the duct to obtain a sample or specimen. In some cases, it can be elected to collect specimens from only a single ductal network. The steps can be repeated in order to identify the presence of an oncofetal fibronectin indicating molecule in

two or more ductal networks. For example, between 6 and 9 ducts are typically present in female human breasts, each of which can be sampled individually or at the same time. A variety of ductal lavage techniques are known in the art. Exemplary ductal lavage techniques are described in U.S. Pat. No. 6,168,779 (issued Jan. 2, 2001) and U.S. Patent Application Nos. 2002/0019017 (published Feb. 14, 2002) and 2002/0037265 (published Mar. 28, 2002).

[0608] In one embodiment, the ductal access tool contains a double lumen catheter. The lavage fluid is introduced into the ductal system through one of the catheter lumens. The lavage fluid can be introduced into the ductal system by a syringe operatively connected to the catheter lumen. Once the ductal system is filled with lavage fluid, excess fluid flows outwardly through the second catheter lumen, from which it is collected.

[0609] Collection of fluid can further include application of suction to the second catheter lumen to facilitate withdrawal of the lavage fluid from the duct system. Suction can be applied using any device capable of creating suction in a lumen. For example, suction can be applied using a syringe or other suction device operatively coupled to the lumen through which lavage fluid will be withdrawn from the ductal system. The suction can be applied for a brief period of time (e.g., a period of time sufficient to establish flow of the ductal fluid from inside to outside the duct). Alternatively, suction can be applied for an extended period of time (e.g., during a corresponding continuous infusion of lavage fluid).

[0610] The ductal system is optionally first accessed with a guide wire, such as a conventional 0.014 in (0.036 cm) guide wire. After the guide wire is inserted past the ductal orifice (e.g., a distance from between 0.25 cm and 2.5 cm or about 0.25 cm and about 2.5 cm past the orifice), the ductal access tool is introduced over the guide wire and into the ductal orifice. After the ductal access tool is in place within the ductal orifice, the guide wire is optionally withdrawn.

[0611] When collecting a breast duct sample, for example, external pressure is optionally applied to the breast to facilitate sample collection. The application of external pressure can be manual or mechanical. The pressure is used to more effectively mix fluid, cells and other ductal contents together in the duct. The external pressure can be applied beginning at the base of the breast and working up to the areola and nipple. The pressure can be applied to the breast periodically, continuously, or cyclically during and/or after infusion of lavage fluid.

[0612] With respect to breast duct samples, for example, the volume of lavage fluid introduced into the ductal system will typically be at least 5 mL or about 5 mL, more typically between 5 mL and 25 mL or about 5 mL and about 25 mL and often 10 mL or about 10 mL. The lavage fluid is generally introduced into the ductal system at a low pressure (i.e., a pressure which will not rupture the ductal network). For example, the fluid can be introduced at a rate in the range of between 0.1 mL/s and 5 mL/s or about 0.1 mL/s and about 5 mL/s and often between 0.5 mL/s and 1 mL/s or about 0.5 mL/s and about 1 mL/s. In addition, the lavage fluid is generally introduced for a relatively short period of time (e.g., between 1 min and 5 min or about 1 min and about 5 min). Introduction of the lavage fluid optionally continues even after the initial portions of the fluid begin to emerge from the second catheter lumen.

[0613] ii. Applying a Label to a Duct

[0614] Identification of ductal orifices on the surface is optionally facilitated by first labeling some or all of the ductal orifices. Methods for labeling ductal orifices are known (see, e.g., U.S. Pat. No. 6,168,779). Briefly, a portion of the epithelial lining exposed at the ductal orifice is labeled with a visible or otherwise detectable marker, which allows the treating professional to identify the orifice for each of the ductal networks. Accordingly, one or more tissue markers at the ductal orifice are specifically labeled with a detectable compound that can preferentially bind to the ductal orifice region. In one embodiment, for example, binding of the label to the ductal orifice region is at least 2-fold, generally, 5-, 10-, 50-, 100-, or more fold, or about 2-fold, generally, about 5-, 10-, 50-, 100-, or more fold stronger than binding of the label to other regions. As such, binding of the label to the orifice will provide a discernable indication of the location of the orifice.

[0615] In certain embodiments, the surface is optionally washed with a solution capable of unblocking the orifice to facilitate or permit binding of the labeling agent to the tissue marker(s) at the ductal orifice. For example, the orifice can become plugged with keratin-containing materials and washing with a keratinolytic solution, such as acetic acid (e.g., 5% to 50% or about 5% to about 50% by weight) admixed in a pharmaceutical delivery vehicle, will expose sufficient marker sites to permit labeling of the ductal orifices.

[0616] The labeling reagents can be formulated as liquids (e.g., aqueous solutions) using any of a variety of conventional techniques. For example, a fibronectin or oncofetal fibronectin binding partner can be in an aqueous solution. The binding partner can be optionally coupled to one member of a signal-producing system capable of generating a detectable visual or other change on the tissue surface. Signal-producing systems include, but are not limited to, fluorescent systems, color-generating systems, luminescent systems, magnetic resonance detection systems, radionuclide systems and ultrasound imaging systems. For example, fluorescent systems that contain a single fluorescent label can be used. Alternatively, other systems that contain two or more components, including enzymes, substrates, catalysts and enhancers, also can be employed. At least one component of the signal-producing system is attached to the binding partner. Alternatively, primary antibodies specific for the tissue marker and labeled secondary antibodies can be employed to indirectly bind to the label to the tissue marker. For example, the primary antibody can be mouse IgG and the labeled secondary antibody can be FITC goat anti-mouse IgG (Zymed).

[0617] In particular embodiments, the tissue marker or markers are antigenic or epitopic sites characteristic of the epithelial lining of the breast ducts. The epithelial lining typically extends sufficiently far into the orifice region of the duct to permit successful labeling using generally conventional immunocytochemical labeling reagents and techniques. Exemplary tissue markers include, but are not limited to antigens and epitopes defined by an oncofetal fibronectin indicating molecule. Other breast epithelial tissue markers include cytokeratins present in the epithelial cytoplasmic lining, such as cytokeratin 8 and cytokeratin 18; and by molecules present in the membrane lining, such as E

cadherin and epithelial membrane antigen (EMA), and others described, for example, in Moll et al., *Cell* 30:11-19 (1982); Gown and Vogel, *Am. J. Pathol.*, 114:309-321 (1984); and Johnson, *Cancer Metastasis Rev.*, 10:11-22 (1991).

[0618] iii. Lavage Fluid

[0619] The lavage fluid typically contains saline and one or more optional ingredients such as, for example, an anesthetic, an oncotic agent, an osmotic agent, oxytocin, prolactin, an antiseptic agent, a ductal orifice dilating agent, a vasodilator, a vasoconstrictor, a muscle relaxant, a muscle constrictor, an anti-ischemic agent, a beta-blocker, a calcium channel blocker, a dye or stain to mark the surface excluding ductal orifices, a dye or stain to mark a perimeter of a ductal orifice and a dye or stain to mark a ductal orifice. The lavage fluid also can optionally contain one or more gases (e.g., air and/or nitrogen). It is expected that the presence of gas serves to increase retrieval of cells and fluid. The gas can be introduced into the lavage fluid by any of a variety of standard methods, including introduction of the gas from a pressurized container.

[0620] The lactation stimulating agent is any agent capable of stimulating lactation in a lactating woman. The agent, as applied to a nipple surface and breast of a non-lactating woman is believed to act to increase the ductal fluid collectable from the breast duct. The lactation stimulating agent can include, but is not limited to, one or more of the following: oxytocin, prolactin and combinations thereof.

[0621] The dye or stain to mark non-ductal orifice regions of the surface is any agent capable of identifying the non-ductal orifice regions on the surface, to the exclusion of the ductal orifices.

[0622] The dye or stain to mark the perimeter regions of ductal orifices is any agent capable of identifying a ring or region surrounding one or more ductal orifices.

[0623] The dye or stain to mark a ductal orifice is any agent capable of marking a ductal orifice to the exclusion of other regions of the surface. The dye or stain to mark a ductal orifice can include, but is not limited to, one or more keratin ligands having a fluorescent tag. In operation, the keratin ligand binds to a keratin plug at a ductal orifice and the fluorescent tag is observed at the ductal orifice, but not at non-keratinized regions of the surface.

[0624] The ductal lavage sample obtained can contain epithelial cells from the lining of the duct, normally secreted and non-secreted fluids present in the ducts and proteins, peptides, nucleic acid molecules, and other chemical species that can be secreted or otherwise released into a duct in response to a disease or other problem to be identified. In connection with breast ducts, material from the terminal ductal lobular unit also can be collected in a lavage procedure, as well as materials residing deep within the ductal passages that access the portion of the breast duct close to the nipple surface, depending upon the depth of penetration of the lavage fluid and the extent to which the fluid that is introduced into the ductal system is retrieved after mixing with fluid and material in the ductal system. The ductal system includes the terminal ductal lobular unit and any tributary ductal passages that connect with or feed into the ductal system leading to the main breast duct that is accessed at the nipple surface.

[0625] 3. Collection of Urine Samples

[0626] Urine can be examined according to the oncofetal fibronectin indicating molecule detection methods known in the art or disclosed herein to screen for the presence of analytes such as an oncofetal fibronectin indicating molecule, which can be used as an indicator for health problems such as overall health, cancer or delivery.

[0627] a. Sample Handling

[0628] Urine samples can be obtained using any of a variety of techniques known in the art. For example, a sample can be collected by a subject urinating into a sample container, a sample can be collected from a catheter that has been introduced into the bladder, or a sample can be obtained by a urine stream contacting a test device, without the need to be collected into a vessel.

[0629] The urine sample can be a pooled collection from a sample container or can be a sample from a urine stream. Either the pooled or the stream samples can contain urine from an entire urination, or only a portion of the urination. Portions of urination can be separated by methods known in the art for different testing purposes, as is known in the art. For example, the first catch of the urination includes the initial portion of the urination. A sample can include the first catch or can exclude the first catch, according to the desired purposes of one skilled in the art.

[0630] The urine sample can be collected from a subject at any time. Typically, a sample is collected at least 1 hour or about 1 hour after the most recent urination. A sample can be collected at least 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, or about 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, or more, after the most recent urination. In one example, a sample collected is a first morning void. Samples collected from a catheter can be stored drainage collected from the drainage bag, or can be fresh drainage collected from the catheter as the drainage flows from the bladder. The sample volume can be varied according to the type and number of tests to be performed and can range from as little as 100 microliters or less, to 50 milliliters or more.

[0631] The sample can be stored in any of a variety of containers known in the art. Typically, the container will be a plastic (e.g., polypropylene or polyethylene) container capable of forming a fluid-impermeable seal that prevents sample leakage and sample contamination and can optionally be sterile. In some embodiments, the container is formed from a material to which an oncofetal fibronectin indicating molecule does not adhere, such as polypropylene or polyethylene. In other embodiments, the container can be formed from a material to which an oncofetal fibronectin indicating molecule does adhere, such as glass, polycarbonate or polystyrene; in such embodiments, assays for an oncofetal fibronectin indicating molecule can be performed in the container. In one exemplary container, a sample can be stored in a container that permits use of a fraction of a sample without contaminating the remainder of the sample, such as that disclosed in U.S. Pat. No. 5,897,840.

[0632] The sample can be stored at room temperature, temperatures below room temperature (such as 4° C.), or can be frozen and stored at temperatures below freezing (such as -20° C. or -70° C.).

[0633] Typically, urine samples used in vertical flow assay methods as described herein are tested immediately after

collection, within 0.5 hours, within 1 hour, 2 hours, 4 hours, 8 hours or 12 hours after collection, or about 0.5 hours, 1 hour, 2 hours, 4 hours, 8 hours or 12 hours after collection. Generally, urine samples used in other assay methods, including those described herein (e.g., dot blot, lateral flow and western blot), that are stored at room temperature can be tested immediately after collection, within 0.5 hours or about 0.5 hours after collection, within 1 hour, 2 hours, 4 hours, 8 hours or 12 hours, or about 1 hour, 2 hours, 4 hours, 8 hours or 12 hours after collection, or later, such as 1 day after collection. Samples that are stored below room temperature (such as 4° C.) can be tested immediately after collection, within 0.5 hours or about 0.5 hours after collection, within 1 hour, 2 hours, 4 hours, 8 hours or 12 hours, or about 1 hour, 2 hours, 4 hours, 8 hours or 12 hours after collection, or later, such as 1 day after collection, 3 days after collection, 1 week after collection, or more. One of ordinary skill in the art can empirically determine, based on the assay method, whether the urine samples should be tested in a narrower time frame (e.g., within 12 hours after collection).

[0634] If a sample is to be frozen, the sample can be frozen by any method known in the art for freezing a liquid. For example, samples can be frozen by placing the sample in a container at -5° C. or less for 1 hour or more. Typically, the sample will be placed in a container, such as a freezer, at a temperature of -20° C., -50° C., -70° C., or less. One skilled in the art will understand that, in order to freeze the sample, the length of time needed for a sample to freeze decreases with a decrease in the temperature of the container in which the sample is placed. After freezing, the sample can be stored at or below a temperature in which the urine sample remains frozen. Typically the storage temperature will be -5° C., -20° C., -50° C., -70° C., or about -5° C., -20° C., -50° C., -70° C., or less. Frozen samples can be stored for 1 week after collection, 2 weeks after collection, 1 month after collection, 2 months after collection, 3 months after collection, 4 months after collection, 5 months after collection, 6 months after collection, or about 1 week after collection, 2 weeks after collection, 1 month after collection, 2 months after collection, 3 months after collection, 4 months after collection, 5 months after collection, 6 months after collection, or more. Frozen samples can be thawed at a variety of temperatures, including room temperature or cooler, such as 4° C. A frozen sample is typically fully thawed prior to analysis of the sample for presence of an oncofetal fibronectin indicating molecule. A frozen sample will usually be subjected to 3 freeze/thaw cycles or less. Typically, a frozen sample is subjected to only one freeze/thaw cycle (i.e., a frozen sample is thawed and tested, not thawed, refrozen, re-thawed and tested).

[0635] Typically, a urine sample used in other assay methods as described herein (lateral flow, dot blot, western blot, etc.), can be used

[0636] b. Sample Condition Modification

[0637] The urine sample can be used "neat" (i.e., without addition of further reagents) or can have added thereto one or more reagents such as preservatives or compounds that inhibit sample degradation such as protease or nuclease inhibitors as is known in the art (e.g., by dilution with a buffer, such as anti-protease buffer (APB) containing 0.05 M Tris buffer, pH 7.4, 150 mM NaCl, 0.02% NaN₃, 1% BSA,

5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin, and 0.1% Triton X-100). Adding one or more reagents can occur by any of a variety of condition modification methods, including, but not limited to, direct mixing of the sample and reagent, dialysis, dilution, filtration and buffer exchange. If the sample conditions are modified, modification can be performed at the time of sample collection, at the time of sample analysis, or any time in between. When a sample is to be frozen, sample condition modification can be performed before freezing the sample, while the sample is being frozen, while thawing the sample, or after the sample is thawed.

[0638] Urine samples can vary from subject to subject, or from sample to sample for the same subject. Sample condition modification can address these variations. To address sample variations, sample modification can be conducted using any of a variety of reagents and methods known in the art. Typically, the diluent or liquid for condition exchange (e.g., dialysis) is distilled water or an aqueous solution of one or more compounds. In one example, the diluent or liquid for condition exchange contains APB buffer: 0.05 M Tris buffer, pH 7.4, 150 mM NaCl, 0.02% NaN_3 , 1% BSA, 5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin, and 0.1% Triton X-100.

[0639] i. Ionic Strength

[0640] Sample condition modification with a buffer such as APB can result in a change in the ionic strength of the sample, depending on the ionic strength of the unmodified sample and the APB buffer. With increasing amounts of condition modification, the ionic strength of the sample will increasingly approximate the ionic strength of the diluent or liquid exchange buffer.

[0641] A sample tested for the presence of an oncofetal fibronectin indicating molecule by contacting the sample with a binding partner specific for an oncofetal fibronectin indicating molecule can yield different test results at different ionic strengths. For example, at low ionic strength, an oncofetal fibronectin indicating molecule can readily bind to a binding partner specific for the oncofetal fibronectin indicating molecule, but other background material such as non-oncofetal fibronectin molecules also can bind to such a binding partner at low ionic strength, thus potentially yielding false positive results. At high ionic strength, non-oncofetal fibronectin molecules will not readily bind to a binding partner specific for an oncofetal fibronectin indicating molecule, but the oncofetal fibronectin indicating molecule also may not strongly bind (detectably bind) to such a binding partner, thus potentially yielding false negative results. A mid-range ionic strength can be selected that permits specific binding between an oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner, while at the same time suppresses binding of background material such as non-oncofetal fibronectin molecules to the fibronectin or oncofetal fibronectin binding partner.

[0642] Urine samples can vary in ionic strength. In such methods, use of neat samples without sample condition modification can result in either false positive or false negative signals. By modifying the conditions of urine samples with a buffer such as APB, the ionic strength from

broadly varying urine samples can be approximated to a mid-range ionic strength determined to permit specific binding partner binding (decreasing false negative results) while suppressing non-specific binding (decreasing false positive results). Thus, modifying the conditions of urine samples with the appropriate buffer can increase the reliability of oncofetal fibronectin indicating molecule detection methods.

[0643] A sample can be modified by any of a variety of ratios of sample to modifying substance. When a buffer is added to a sample, the buffer can be added such that a desired ionic strength is achieved without unduly diluting the sample components such as an oncofetal fibronectin indicating molecule. One skilled in the art can determine a desirable dilution ratio, according to the desired ionic strength or range thereof, the ionic strength of the sample, the reagent to be added to the sample and the concentration of sample components. In some embodiments, a minimum amount of reagent is added to arrive at a desired ionic strength or ionic strength range, thus resulting in a minimum amount of sample dilution. In other embodiments, a dilution ratio and reagent that can be used to adjust most or all urine samples to a desired ionic strength or ionic strength range can be selected. Typical dilution ratios include 1:1.5 (volume before dilution:volume after dilution), 1:2, 1:3, 1:4, 1:5, 1:6, 1:7, 1:8, 1:9, 1:10, 1:12, 1:15, 1:20, about 1:1.5, 1:2, 1:3, 1:4, 1:5, 1:6, 1:7, 1:8, 1:9, 1:10, 1:12, 1:15 and 1:20. A typical dilution ratio is 1:4.

[0644] In one embodiment, the reagent added to the urine sample is a buffer solution such as a modified APB buffer. Typically, the principal contributor to the ionic strength of a buffer solution can be a salt, including a monovalent salt such as NaCl, KCl, NaBr, KBr and a variety of other salts known in the art. The salt concentrations can be any of a variety of concentrations that achieves the desired ionic strength when added to urine. Exemplary salt concentration ranges can include 50 mM to 350 mM, 100 mM to 250 mM, or 150 mM. Additional exemplary salt concentration ranges can include about 50 mM to about 350 mM, about 100 mM to about 250 mM, or about 150 mM. Such buffer solutions can contain a variety of additional components, including, but not limited to, a buffering compound (e.g., Tris or phosphate), a chelator (e.g., EDTA or EGTA), a protease inhibitor (e.g., PMSF or aprotinin), detergent (e.g., Tween or Triton X-100), other stabilizers (e.g., PEG or BSA), or combinations thereof. In one example, a buffer solution can contain 0.05 M Tris buffer, pH 7.4, 150 mM NaCl, 0.02% NaN_3 , 1% BSA, 5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin, and 0.1% Triton X-100.

[0645] ii. Ionic Strength Testing

[0646] Typically, a major ionic component in urine is sodium chloride. The concentrations of either sodium, chloride, or both, or total ionic strength, can be determined using a variety of testing methods known in the art including use of test strips, glass membrane electrodes, conductivity measurement, atomic absorption, laser induced fluorescence or x-ray fluorescence. The results of the ion composition test can either directly measure or approximate the ionic strength of the urine sample. If the ionic strength of the urine sample is higher than a maximum allowable amount, it is possible to use the ionic strength measurement to calculate the

minimum amount of dilution necessary to achieve a selected ionic strength in the sample prior to analysis, or the minimum amount of liquid exchange (e.g., dialysis) necessary to achieve a selected ionic strength in the sample prior to analysis. If the ionic strength of the urine sample is lower than a minimum allowable amount, it is possible to use the ionic strength measurement to calculate the minimum amount of salt solution or solid salt to add to the urine to achieve a selected ionic strength in the sample prior to analysis, or the minimum amount of liquid exchange to achieve a selected ionic strength in the sample prior to analysis. As one skilled in the art will recognize, the selected ionic strength for a binding partner to bind to an oncofetal fibronectin indicating molecule can include a range of ionic strengths and can vary based on a variety of factors including the nature of the binding partner. Typically, the selected ionic strength for binding assays ranges from 50μ to 500μ , from 75μ to 400μ , from 100μ to 300μ and will often be 150μ to 250μ . Alternatively, the selected ionic strength for binding assays ranges from about 50μ to about 500μ , from about 75μ to about 400μ , from about 100μ to about 300μ and will often be about 150μ to about 250μ .

[0647] iii. Normalization

[0648] The concentration of a component of urine can vary from urine sample to urine sample as a function of a variety of factors including condition modification (e.g., dilution or dialysis) of a urine sample and frequency of urination by the subject. One of several methods can be used to estimate a sample variation-independent concentration of an analyte such as an oncofetal fibronectin indicating molecule, in a urine sample, where the estimated concentration is less sensitive to variation between samples. Such a method of estimating sample variation-independent concentrations is termed normalization.

[0649] Normalization can be carried out using any of a variety of methods known in the art or provided herein. One method of normalization of a urine sample is to measure the concentrations of the analyte of interest (an oncofetal fibronectin indicating molecule) and a second analyte that typically enters urine at a constant rate.

[0650] A variety of constantly-entering analytes are known in the art. One example of such an analyte is creatinine, which enters urine at a steady state. The concentration of creatinine can be used to normalize the concentration of an oncofetal fibronectin indicating molecule in urine. A variety of different normalization methods are known in the art, as exemplified in U.S. Pat. Nos. 5,804,452 and 6,436,721.

[0651] c. Sample Treatment

[0652] i. Non-Specific Binding

[0653] The urine sample can be contacted with one or more non-specific binding compounds or one or more non-specific binders prior to contacting the urine sample with a molecule that can detect the presence of an oncofetal fibronectin indicating molecule.

[0654] Non-specific binding compounds are compounds that bind to at least a portion of background material in a sample without binding more than a small amount (e.g., less than 10%) of oncofetal fibronectin indicating molecule in the sample. Typically, non-specific binding compounds bind

to background material more readily than the non-specific binding compounds bind to an oncofetal fibronectin indicating molecule. Non-specific binding compounds that can be used include non-specific binding proteins. Non-specific binding proteins are typically water-soluble proteins including albumins such as bovine serum albumin (BSA), human, rabbit, goat, sheep and horse serum albumins; and other proteins such as ovalbumin, fibrinogen, thrombin, transferrin, glycoproteins, casein, antibodies not specific for an oncofetal fibronectin indicating molecule and other proteins. Non-specific binding proteins also can include water-soluble polyamino acids such as, for example, polymers of one or more amino acids such as lysine, glutamic acid, alanine, histidine, methionine and proline. Exemplary proteins that can be used for a non-specific binding surface include BSA, methylated BSA or antibodies such as mouse anti-MHC-1 antibody (e.g., ATCC No. W632) or mouse IgG. Non-specific binding compounds also can be protein-containing compositions including serum such as fetal calf serum, gelatin and dried milk.

[0655] Non-specific binders can include non-specific binding surfaces, which are solid structures that can contain one or more components, where the non-specific binding surface binds to at least a portion of background material in a sample while not binding more than a small amount (e.g., less than 10%) of oncofetal fibronectin indicating molecule in the sample. Typically, non-specific binding surfaces bind to background material more readily than the non-specific binding surfaces bind to an oncofetal fibronectin indicating molecule. Possible solid supports for non-specific binding surfaces include paper and cellulose derivatives, such as cellulose esters and ethers, natural and synthetic polymeric materials, such as latex, vinyl polymers, polypropylene, polyethylene and partially hydrolyzed derivatives, polycondensates, copolymers and inorganic materials. In one embodiment, a non-specific binding surface is a porous or fibulous member capable of transporting a liquid sample along a test strip. Non-specific binding surfaces that can be used include solid supports having immobilized thereon one or more non-specific binding proteins such as, but not limited to, albumin (including bovine serum albumin, or BSA), antibodies not specific for an oncofetal fibronectin indicating molecule and others provided herein or known in the art. Exemplary proteins that can be used for a non-specific binding surface include BSA, methylated BSA or antibodies such as W632 or mouse IgG. In one example, a non-specific binding surface can be a nitrocellulose membrane having methylated BSA immobilized thereon.

[0656] ii. Filtration

[0657] The urine sample also can be filtered to remove at least a portion of the background materials prior to detecting the presence of an oncofetal fibronectin indicating molecule. Filters that can be used are low protein binding filters that bind to no more than a small amount (less than 10%) of the oncofetal fibronectin indicating molecule present in the urine sample. Typically, filters used in the methods provided herein bind to background material more readily than the non-specific binding compounds bind to an oncofetal fibronectin indicating molecule. Exemplary low protein binding filters include polyester, polyurethane, fiberglass, polyacetate, polyvinylidene fluoride, polycarbonate, nylon, polyethersulfone, polysulfone, cellulose, cellulose acetate, cellulose mixed esters and hydrophilic modifications

thereof. An exemplary low protein binding filter is cellulose acetate. The filter pore size can be large enough to permit passage of an oncofetal fibronectin indicating molecule but small enough to prevent passage of at least a portion of the background materials. Filter pore sizes can range from 20 μm to 0.01 μm , 10 μm to 0.02 μm , 5 μm to 0.05 μm , 1 μm to 0.1 μm and from 0.5 μm to 0.2 μm . Exemplary filters have a pore size of 0.2 μm .

[0658] 4. Interstitial Fluid

[0659] Interstitial fluid also can be collected and tested for the presence of an oncofetal fibronectin indicating molecule. Interstitial fluid is the fluid in the spaces between tissue cells and can constitute about 16% of the weight of the body. Interstitial fluid can contain, as one of its components, an oncofetal fibronectin indicating molecule. Normal interstitial fluid either does not contain oncofetal fibronectin, or contains only low amounts of oncofetal fibronectin; therefore, presence of an oncofetal fibronectin indicating molecule in an interstitial fluid sample can indicate presence of a health problem associated with oncofetal fibronectin in the subject. Interstitial fluid can be collected from a variety of locations throughout the body, as is well known in the art. In one embodiment, interstitial fluid can be collected from the tissue or organ under examination, or from tissue adjacent thereto. For example, when a region or organ in or near the alimentary canal is under examination (e.g., large intestine, prostate, stomach, or gall bladder), interstitial fluid can be collected from the epithelium of that region or organ or tissue adjacent to the region or organ (e.g., in order to monitor the prostate, interstitial fluid can be collected from the region of the colon adjacent the prostate). In another embodiment, interstitial fluid can be collected from skin, regardless of whether or not the skin or some distal region or organ is under examination. For example, interstitial fluid in skin can contain an oncofetal fibronectin indicating molecule that indicates neoplasia in the breast or cervix or thyroid. The dermal layer of skin contains collagen fibers, cells and interstitial fluid in the space between the fibers and cells. Interstitial fluid collected from the dermal layer, or from other locations in the body of a subject, can be used to test for the presence of an oncofetal fibronectin indicating molecule in a subject.

[0660] Interstitial fluid samples can be collected by a variety of methods known in the art such as, but not limited to, needle aspiration including dermal needle aspiration (see, e.g., U.S. Pat. Nos. 6,702,791, 5,823,973 and 5,582,184), microporation (see, e.g., U.S. Pat. No. 6,508,785), ultrasound extraction (see, e.g., U.S. Pat. No. 6,589,173), transdermal extraction (see, e.g., U.S. Pat. No. 4,595,011), iontophoresis (see, e.g., U.S. Pat. Nos. 6,496,728; 5,989,409; 5,362,307 and 5,279,543), blister collection such as microblister collection (see, e.g., U.S. Pat. No. 6,334,851), microblade or microneedle array extraction (see, e.g., U.S. Pat. No. 6,562,014), enhanced cell permeation collection (see, e.g., U.S. Pat. No. 6,503,198) and a variety of other methods known in the art.

[0661] In one embodiment, an interstitial fluid sample can be collected by penetrating the outer layer, typically the epithelium, of the region or organ from which the interstitial fluid sample is to be collected, with an instrument such as a hypodermic needle, into which interstitial fluid can enter. The vesicle, such as a hypodermic needle, can pass into

and/or through the outermost portion of the region or organ, such as the epithelial cells, to a portion of the region or organ in which interstitial fluid is present. The depth of penetration will vary according to the tissue type and location of the region or organ to be examined, as will be known to those skilled in the art. The vesicle that penetrates the region or organ of interest will have an outer dimension that is small enough to pass with relative ease into the tissue or organ while maintaining physical integrity of the vesicle and an inner dimension that is large enough to permit interstitial fluid to pass into the vesicle. For example, needles of a size of 28 to 32 gauge or about 28 to about 32 gauge (360 micron outer diameter to 230 micron outer diameter) can be used; typically, a needle of 29 or 30 gauge is used. Optionally, the vesicle can be operably attached to a negative pressure device, such as a vacuum apparatus or a syringe in which a plunger can be manually or mechanically withdrawn. A variety of apparatuses and methods for collecting interstitial fluid by penetrating the region or organ of interest are known in the art, as exemplified in U.S. Pat. Nos. 6,702,791, 6,624,882, 5,823,973 and 5,582,184.

[0662] In one example of collecting interstitial fluid with a vesicle, interstitial fluid can be collected from skin. The vesicle, such as a needle, can pass through the outer layer of the epidermis (known as the stratum corneum) without passing through the dermal layer of the skin. After passing through the stratum corneum, interstitial fluid can enter or be drawn into the vesicle. The depth to which the vesicle, such as a needle, is inserted into the skin will be deep enough to penetrate the stratum corneum, but typically not deep enough to pass through the dermal layer and penetrate the subcutaneous layer. The stratum corneum is typically 10-15 microns or about 10-15 microns thick and the remainder of the epidermis is typically 80 microns or about 80 microns thick. The dermis is about 2,000-3,000 microns thick. As will be appreciated by one skilled in the art, such dimensions will vary somewhat from individual to individual and depending on the body location of the skin from which the sample is to be collected. A vesicle such as a needle, will typically be inserted into the skin at a depth of 50-2,500 microns or about 50-2,500 microns, often 700-1,500 microns or about 700-1,500 microns. The vesicle that penetrates the skin will have an outer dimension that is small enough to pass with relative ease through the stratum corneum with minimal pain while maintaining physical integrity of the vesicle and an inner dimension that is large enough to permit interstitial fluid to pass into the vesicle.

[0663] In another embodiment, an interstitial fluid sample can be collected by penetrating into or through the region or organ of interest, such as the epithelium of an organ or the stratum corneum of skin, with a microarray of blades or needles. Such microarrays can be applied to the tissue or organ of interest to provide a path in the tissue or organ, or to provide a vesicle, through which interstitial fluid can pass. For example, a blade microarray can be applied to the skin to provide a path in the stratum corneum through which interstitial fluid can pass. Microblade or microneedle arrays can provide a plurality of locations on the region or organ of interest for collection of interstitial fluid. The depth to which each microblade or microneedle is inserted will be deep enough to penetrate into the region or organ of interest to a region containing interstitial fluid. For example, when an interstitial fluid sample is collected from skin, the depth to which each microblade or microneedle is inserted into the

skin will be deep enough to penetrate the stratum corneum, but typically not deep enough to pass through the dermal layer and penetrate the subcutaneous layer. In an exemplary configuration, each microblade or needle can have an outer dimension of 1-50 microns or about 1-50 microns, a length of 50-500 microns or about 50-500 microns and be separated from neighboring microneedles or microblades by 50-1000 microns or about 50-1000 microns. The microblades or microneedles can optionally be operably coupled with an apparatus or compound for increasing interstitial fluid passage through the penetrated tissue. For example, the microneedle or microblade array can be coupled with a negative pressure device, such as a vacuum apparatus. In another example, a microblade array or microneedle array can be coupled with an absorbent pad that can draw interstitial fluid from the subject by capillarity and/or by osmotic pressure. A variety of microblade and microneedle arrays and methods for using such arrays for collecting interstitial fluid are known in the art, as exemplified in U.S. Pat. Nos. 6,663,612, 6,562,014 and 6,312,612.

[0664] In another embodiment, an interstitial fluid sample can be collected by blister formation and harvesting the fluid in the blister. Interstitial fluid can be a major component of a blister. Blisters can be formed by any of a variety of methods known in the art, including suction and heating methods. For example, when a negative pressure of 200 mm Hg or about 200 mm Hg is applied to skin for 1 hour or about 1 hour, a blister will form. The fluid in the suction blister can be collected using any of a variety of methods, such as, but not limited to, aspiration with a hypodermic needle. In another example, heat can be used to form a blister. For example, a laser energy absorbing substance can be placed onto the skin of a subject and exposure of the skin to laser energy can result in blister formation. Placement of the absorbing substance and the amount of laser energy applied can be controlled such that only microblisters form. The fluid in the heat blister can be collected using any of a variety of methods, such as, but not limited to, aspiration with a hypodermic needle. A variety of apparatuses and methods for blister formation and collection of interstitial fluid from the blister are known in the art, as exemplified in U.S. Pat. Nos. 6,409,679, 6,387,059 and 6,334,851.

[0665] In another embodiment, an interstitial fluid sample can be collected by ultrasound extraction. Treating a tissue such as skin with ultrasound can permeabilize the tissue and the treated area can remain permeable from 30 minutes or about 30 minutes after treatment up to 10 hours or about 10 hours after treatment. Interstitial fluid can be collected from the treated, permeable area by any of a variety of chemical and/or physical methods, including, but not limited to, applying negative pressure, contacting the area with a surfactant or organic solvent, contacting the area with a composition that creates osmotic pressure, contacting the area with an ultrasound coupling medium and applying electrical current. The interstitial fluid can be collected in a buffer, salve, gel, or other composition that contacts the treated, permeable area. Ultrasound can be applied directly to the region or organ under examination, such as the large intestine, or can be applied to a region adjacent the region or organ under examination, such as the region of the large intestine that is adjacent to the prostate when the prostate is under examination. Ultrasound also can be applied to skin. A variety of apparatuses and methods for collecting inter-

stitial fluid using ultrasound are known in the art, as exemplified in U.S. Pat. Nos. 6,620,123, 6,508,785 and 6,190,315.

[0666] In another embodiment, an oncofetal fibronectin indicating molecule in an interstitial fluid sample can be collected by electrical harvesting. In electrical harvesting, a substance such as an oncofetal fibronectin indicating molecule, diffuses through a membrane as a result of applying an electric field. Electrical harvesting can be used to collect substances within interstitial fluid. Electrical harvesting can be performed by conducting electrical current through the tissue to extract the substance into one or more sampling vesicles or into collection compositions (e.g., a buffer, a gel, salve or cream that can absorb components of interstitial fluid). For example, an oncofetal fibronectin indicating molecule sample can be collected by applying electrical energy of sufficient strength and duration to the tissue surface of a subject in order to transport a substance or a metabolite from beneath the tissue surface at a collection site on the surface of the tissue into a defined collection area or into a collection vesicle. These methods include the method known as iontophoresis, which is a method of treatment to drive uncharged non-ionic materials and positive or negative ions out of an organism through tissue. In conventional iontophoresis two electrodes are placed in contact with the tissue. One or both of the electrodes is in a sampling chamber/collection reservoir to collect the substance extracted and a voltage is applied between the two electrodes. The sampling chamber/collection reservoir is provided at the tissue surface to serve as a collector of material transported. Any method known in the art such as iontophoresis that results in collection of components of interstitial fluid by, for example, electrophoresis, electroosmosis, or electroporation, can be used to collect a sample containing an oncofetal fibronectin indicating molecule. The electrical harvesting is typically performed by applying an electrical field to the region or organ under examination, or to a region adjacent thereto. For example, the electrical field can be applied to skin. The electrical field applied to the tissue from which the sample is to be harvested can be any electrical field along which an analyte such as an oncofetal fibronectin indicating molecule can migrate. Exemplary electrical fields include direct current, pulsed direct current and current with alternating polarity such as AC current. A variety of apparatuses and methods for collecting an oncofetal fibronectin indicating molecule from interstitial fluid by electrical harvesting are known in the art, as exemplified in U.S. Pat. Nos. 6,496,728, 6,023,629 and 5,989,409.

[0667] In another embodiment, an interstitial fluid sample can be collected using a permeation enhancing compound. A permeation-enhancing compound can serve to increase the permeability of a membrane, cell or tissue and permit interstitial fluid, or selected components thereof such as an oncofetal fibronectin indicating molecule, to pass through the membrane, cell or tissue, to a location at which the interstitial fluid or components can be collected. A permeation enhancing compound typically functions by contacting the surface for which permeation is to be enhanced and causing one or more components of interstitial fluid to pass through the contacted surface. A variety of permeation enhancing compounds are known in the art and include, but are not limited to, aqueous hypertonic solutions, solutions containing organic solvents such as propylene glycol, dimethylsulfoxide or isopropyl alcohol, organic salts such as

bile salts including sodium cholate, surfactants such as detergents, including cationic detergents, anionic detergents, non-ionic detergents and zwittergents, organic compounds such as pyrrolidones including N-methyl-pyrrolidone. The permeation compound can be a liquid or a solid, but when the permeation compound is a solid, the solid is typically dissolved in a solvent. The permeation enhancing compounds can be applied directly to the surface of interest, or can be applied in a composition such as a salve, cream, gel, or in an absorbent pad. In one example, the permeation compound can be a component of a transdermal patch that can be affixed to the skin of a subject from which the sample is to be collected. A variety of apparatuses and methods for collecting components of interstitial fluid using permeation enhancing compounds are known in the art, as exemplified in U.S. Pat. Nos. 6,503,198, 6,492,180, 5,438,984, 5,139,023, 4,960,467, 4,746,508 and 4,595,011.

[0668] In another embodiment, an interstitial fluid sample can be collected by electromagnetic radiation mediated permeation. Electromagnetic irradiation of a membrane, cell or tissue can result in ablation or formation of micropores in the irradiated area. Electromagnetic irradiation can cause ablation or micropore formation by a variety of mechanisms including photochemical, photothermal and photomechanical ablation. Photochemical ablation occurs by dissociation and/or formation of atomic bonds that results in disruption of a membrane or tissue or extracellular matrix. Photothermal ablation occurs by absorption of heat, which can result in denaturation of a membrane or tissue or extracellular matrix, or in water vapor formation within a membrane or tissue or extracellular matrix, which results in pressure formation that causes fracturing of a membrane, tissue or extracellular matrix. Photomechanical ablation occurs by absorption of electromagnetic energy in such a way that mechanical stress is induced onto the absorbing surface, for example, by delivering radiation pulses at time lengths shorter than thermal diffusion time, resulting in mechanical stress caused by compression, expansion and/or recoil of the treated surface. By controlling the variables of electromagnetic irradiation such as the wavelength, intensity, pulse duration, pulse frequency and beam size, one skilled in the art can control the amount and type of energy delivered and the location and area of the irradiated surface. Such control also permits the skilled artisan to select the degree to which ablation or microchannel formation will be accomplished by photochemical, photothermal or photomechanical mechanisms, or combinations thereof. Such control also permits one skilled in the art to determine the size and depth of the micropores or ablations resultant from exposure to electromagnetic radiation. Electromagnetic radiation can be delivered by, for example, one of a variety of lasers which can be selected according to the properties of the particular laser, such as intensity and wavelength, or other considerations such as portability or expense. In addition, one or more electromagnetic energy absorbing compounds such as dyes can be used to aid or further control energy transfer to the surface, as is known in the art. Micropore formation or ablation by magnetic radiation can permit interstitial fluid to pass through the treated region, membrane or tissue. Any of a variety of tissues or organs throughout the body can be treated with electromagnetic radiation to form micropores or ablated tissue. In one example, the stratum corneum of the skin can have formed therein micropores through which interstitial fluid can pass; similarly, the micropores can be

formed to pass through the stratum corneum and epidermis and into the dermis. Typically, micropore formation or ablation will be performed such that a sufficient channel is formed to permit interstitial fluid to pass therethrough, without resulting in significant damage to surrounding tissue or discomfort to the subject being treated. Arrays of micropores or ablated tissue also can be readily formed using such methods. Micropore formation or ablation by electromagnetic irradiation can be coupled with any of a variety of methods for harvesting interstitial fluid including, for example, coupling with a negative pressure device such as a vacuum. A variety of apparatuses and methods for collecting interstitial fluid using electromagnetic radiation are known in the art, as exemplified in U.S. Pat. Nos. 6,685,699, 6,508,785 and 6,387,059.

[0669] In a similar embodiment, an interstitial fluid sample can be collected by micropore formation as a result of plasma formation or bombardment with microparticles. Plasma, in the present context, refers to any of a variety of charged molecules, atoms or subatomic particles such as electrons. Plasma can be formed on the surface of the target region or organ, such as the surface of skin, by, for example, pulses from an intense laser. The charged particles can be highly reactive and can cause ablation or micropore formation at the surface location of the plasma. A variety of apparatuses and methods for collecting interstitial fluid plasma for micropore formation or ablation are known in the art, as exemplified in U.S. Pat. Nos. 6,387,059 and 5,586,981. Micropores or ablation also can be caused by bombardment with microparticles. Microparticles are solid particles that can range from 0.1 micron in diameter to 100 microns in diameter or about 0.1 micron in diameter to about 100 microns in diameter. Microparticles can be composed of any of a variety of solid compounds or compositions known in the art, ranging from solid water (water ice) or solid carbon dioxide (dry ice) to insoluble compounds such as insoluble inorganic compounds such as aluminum oxide or titanium oxide, to soluble compounds such as sugars, starch or salts, or metals such as gold, platinum or tungsten. Microparticles can form micropores by being accelerated toward the surface of a region or organ under examination. Microparticles can be accelerated toward a surface using any of a variety of methods known in the art including, but not limited to, compressed gas, electric discharge, expansion of a liquid to a gas such as liquid helium expanding to a gas at room temperature, negative pressure acceleration, or momentum transfer by contact with a moving solid surface. The depth and size of the micropores formed can be a function of the size and weight of the microparticles used and the amount of acceleration of the microparticles. Typically, the micropores-formed will be sufficiently large to form a channel that permits interstitial fluid to pass therethrough, while not resulting in significant damage to surrounding tissue or discomfort to the subject being treated. For example, the micropores formed by microparticles can range from 10 microns to 1000 microns or about 10 microns to about 1000 microns in diameter. The depth of the micropore can depend on the tissue being sampled; for example, a micropore in skin can be 50 to 2000 microns or about 50 to about 2000 microns in depth. A variety of apparatuses and methods for collecting interstitial fluid using microparticle bombardment are known in the art, as exemplified in U.S. Pat. Nos. 6,706,032 and 6,482,604. Arrays of micropores also can be readily formed using

plasma ablation or microparticle bombardment. Micropore formation by plasma or microparticle bombardment can be coupled with any of a variety of methods for harvesting interstitial fluid including, for example, coupling with a negative pressure device such as a vacuum.

[0670] The above embodiments can be coupled together in collecting interstitial fluid or components thereof, such as an oncofetal fibronectin indicating molecule. For example, microparticle mediated microporation can be coupled with a permeation enhancing compound in collecting the sample. In another example, different microporation methods can be coupled together, such as, for example, ultrasound and iontophoresis. A large variety of additional combinations of the above methods are known in the art, as described in the patents related to each embodiment and in U.S. Pat. No. 6,692,456. In particular, the methods for enhancing passage of interstitial fluid or components thereof to a collectable region can be coupled with any of a variety of methods for collecting the fluid or components thereof, including applying negative pressure such as a vacuum, applying a liquid, gel, cream, salve, solid support such as a patch, or any other interstitial fluid collection vesicle or compound known in the art.

[0671] Methods and apparatuses for collecting interstitial fluid or components thereof, such as an oncofetal fibronectin indicating molecule, can be performed in conjunction with one or more additional steps of sample treatment and/or oncofetal fibronectin indicating molecule detection. In one embodiment, the sample collection device can be coupled with a composition containing a non-specific binding partner. For example, a sample can be collected by coupling a vacuum to a microneedle array and the microneedle array can be in fluid communication with a composition that contains a non-specific binding partner. For example, the microneedle array can be in fluid communication with a buffer containing a non-specific binding partner such as BSA. In another example, the microneedle array can be coupled to a solid support or a gel containing a non-specific binding partner. Any of a variety of formats for the non-specific binding partner, as provided elsewhere herein, can be used with any of the variety of interstitial fluid sampling methods and combinations thereof provided herein.

[0672] In another embodiment, the sample collection device can be coupled with a composition containing a fibronectin binding partner or an oncofetal fibronectin binding partner. For example, a sample can be collected by microporation followed by application of a transdermal patch containing a permeation enhancing compound and the transdermal patch can further contain a composition that contains a fibronectin binding partner or an oncofetal fibronectin binding partner. For example, the transdermal patch can contain a buffer containing a colloidal gold fibronectin binding partner conjugate. In another example, the transdermal patch can contain a solid support or containing a fibronectin or oncofetal fibronectin binding partner bound to the solid support. Any of a variety of formats for the fibronectin binding partner or oncofetal fibronectin binding partner, as provided elsewhere herein, can be used with any of the variety of interstitial fluid sampling methods and combinations thereof provided herein.

F. Methods of Detecting Oncofetal Fibronectin

[0673] Provided herein are methods of detecting an oncofetal fibronectin indicating molecule. Typically, the

methods are used to detect an oncofetal fibronectin indicating molecule in a sample, such as a sample from a subject. An oncofetal fibronectin indicating molecule can be, for example, an oncofetal fibronectin protein, a nucleic acid molecule encoding oncofetal fibronectin or a complement thereto, or an autoantibody that specifically binds oncofetal fibronectin protein or a nucleic acid molecule encoding oncofetal fibronectin, or a fragment thereof. Methods of detecting an oncofetal fibronectin indicating molecule in a sample can be used to determine the presence of an oncofetal fibronectin indicating molecule in the sample, can be used to determine the amount or concentration of an oncofetal fibronectin indicating molecule in a sample, can be used to determine whether or not a positive result is a false positive and can be used to determine the regions or composition of an oncofetal fibronectin indicating molecule present in the detected an oncofetal fibronectin indicating molecule.

[0674] For example, when the particular type of sample collected from a subject typically does not contain detectable amounts of an oncofetal fibronectin indicating molecule when collected from normal subjects, detecting any amount of an oncofetal fibronectin indicating molecule in such a sample can indicate the presence of a health problem associated with oncofetal fibronectin in the subject. In another example, when the type of sample collected from a subject typically contains a baseline amount of an oncofetal fibronectin indicating molecule when collected from normal subjects, detecting an amount of an oncofetal fibronectin indicating molecule greater than the baseline amount can indicate the presence of a health problem associated with oncofetal fibronectin in the subject.

[0675] Assays intended for use in the systems and methods disclosed herein include, but are not limited to: protein detection, including immunoassay or other antigen-binding-based detection; nucleic acid detection, including methods using amplification and non-amplification protocols; any assay that includes colorimetric or spectrometric detection, such as fluorometric and luminescent detection; mass spectrometric analysis; or any assay that includes binding of an oncofetal fibronectin indicating molecule to a fibronectin or oncofetal fibronectin binding partner. Any test that produces a signal, or from which a signal can be generated, or that can be detected by a detector, such as a photodetector, a gamma counter or a mass spectrometer, is intended for use in the methods provided herein. Any wavelength is intended to be included.

[0676] Any of the methods for detecting oncofetal fibronectin provided herein can be used in conjunction with any of the sample collection methods provided herein or known in the art, to provide information for any of the indications or other uses of detection of oncofetal fibronectin provided herein or otherwise known in the art.

[0677] Binding assays, including competitive binding assays and sandwich assays, are among those that can be used in the methods provided herein. The methods and systems provided herein have broad applicability to a variety of sample types for a variety of different indications, as will be apparent to one skilled in the art. A number of different types of binding assays using a variety of protocols and labels are well known. Binding assays can be performed in a single liquid phase, or a binding partner can be immobilized to a solid support upon which the assay is performed.

Sandwich assays can be performed. Competitive assays can be performed. The reaction steps can be performed simultaneously or sequentially, as will be known to those skilled in the art. Threshold assays can be performed, where only analyte levels of above a specified level or rate yield a positive result. Assay formats include, but are not limited to, for example, assays performed in test tubes, on membranes, in wells, in multi-well plates, on microarrays, on chromatographic test strips, as well as dipstick, lateral flow, vertical flow, or migratory format assays.

[0678] Assay methods also can include mass measurement, where mass of an oncofetal fibronectin indicating molecule, a fragment thereof, or another compound indicative of the presence of an oncofetal fibronectin indicating molecule can indicate the presence of the oncofetal fibronectin indicating molecule in the sample. An exemplary mass measurement method is mass spectrometry.

[0679] The assay methods provided herein also include nucleic acid molecule amplification methods, where a primer can hybridize with a nucleic acid molecule encoding oncofetal fibronectin (e.g., mRNA), and can serve as a substrate for nucleic acid synthesis methods, such as, for example, RT-PCR.

[0680] The detection methods provided herein also can include one or more sample manipulation methods. In one example, the sample can have one or more components separated or removed, for example, in a method of increasing the relative amount of oncofetal fibronectin indicating molecule present in the sample. In another example, the sample can be contacted with a fragmentation compound that can fragment oncofetal fibronectin indicating molecule present in the sample.

[0681] In one embodiment, detecting an oncofetal fibronectin indicating molecule with two or more different binding partners or with two or more different detection methods can be used to confirm the presence and/or amount of oncofetal fibronectin in the sample. Detecting an oncofetal fibronectin indicating molecule with two or more different binding partners or with two or more different detection methods also can be used to determine the regions present and/or not present in the oncofetal fibronectin indicating molecule. Determination of the regions or composition of an oncofetal fibronectin indicating molecule present in a sample can be used for a variety of purposes, including to identify the cell or tissue type(s) from which an oncofetal fibronectin indicating molecule could have been produced and/or the cell or tissue type(s) from which an oncofetal fibronectin indicating molecule was likely to not have been produced, to indicate or identify the binding or biochemical activity of the oncofetal fibronectin present in the sample (e.g., when EDA is present in oncofetal fibronectin, the oncofetal fibronectin can bind to $\alpha_9\beta_1$ integrin and can have improved cell spreading and migration properties (see, e.g., Manabe et al., *J. Cell. Biol.* 139:295-307 (1997) and Liao et al., *J. Biol. Chem.* 277:14467-14474 (2002)) and to rule out the possibility of a false positive result for the presence or amount of an oncofetal fibronectin indicating molecule in the sample.

[0682] A variety of cell types, tissue types and tumor types are known to contain oncofetal fibronectin proteins that include one or more of EDA, EDB and one of the splice variants of IIICS. In one case, the oncofetal fibronectin

protein can be characterized as the EDA portion of an oncofetal fibronectin protein, the EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, the portion of an autoantibody that binds to EDA of oncofetal fibronectin protein, and the portion of an autoantibody that binds to the EDA-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin. In another example, the fibronectin or oncofetal fibronectin protein: can be characterized as the EDB portion of an oncofetal fibronectin protein, the EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, the portion of an autoantibody that binds to EDB of oncofetal fibronectin protein, and the portion of an autoantibody that binds to the EDB-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin. In another example, the fibronectin or oncofetal fibronectin protein can be characterized as the IIICS portion of an oncofetal fibronectin protein, the IIICS-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin, the portion of an autoantibody that binds to IIICS of oncofetal fibronectin protein, and the portion of an autoantibody that binds to the IIICS-encoding portion of a nucleic acid molecule encoding oncofetal fibronectin. In another example, an oncofetal fibronectin protein can be characterized as containing the IIICS splice variant V64, V89, V95 or V120. In another example, an oncofetal fibronectin protein can be characterized as containing one or more post-translational modifications such as O-glycosylation of threonine 33 of IIICS. A variety of cell types, tissue types and tumor types are known to contain oncofetal fibronectin proteins that do not include one or more of EDA, EDB and one of the splice variants of IIICS. In one case, an oncofetal fibronectin protein can be characterized as lacking EDA, EDB or IIICS. In another example, IIICS is identified as lacking amino acids 1-25 of IIICS, or 90-120 of IIICS, or both. Oncofetal fibronectin variants and their association with various cell types, tissue types and tumor types are exemplified herein and are known in the art. Many samples for which the presence of oncofetal fibronectin can be determined, such as a sample collected from a specific organ or tissue, or from a body fluid sample such as urine, lymph, blood, plasma, serum, saliva, cervical fluid, cervicovaginal fluid, vaginal fluid, breast fluid, breast milk, synovial fluid, semen, seminal fluid, stool, sputum, cerebral spinal fluid, tears, mucus, interstitial fluid, follicular fluid, amniotic fluid, aqueous humor, vitreous humor, peritoneal fluid, ascites, sweat, lymphatic fluid, lung sputum and lavage, can contain components whose sources are two or more cell types, two or more tissue types, or two or more organs. As a result, the cell, tissue or organ source of the oncofetal fibronectin in a sample can be ambiguous. Using the methods provided herein, characterization of the regions or composition of oncofetal fibronectin present in the sample can be used to identify the likely cell source, tissue source or organ source of an oncofetal fibronectin indicating molecule, or can be used to identify the unlikely cell source, tissue source or organ source of an oncofetal fibronectin indicating molecule. Such methods can be used, for example, to identify the type of tumor or neoplastic cell likely present in a subject, or to identify the type of tumor or neoplastic cell unlikely present in a subject.

[0683] To illustrate, a sample can contain two or more different oncofetal fibronectin proteins. For example, a sample can contain an oncofetal fibronectin protein that contains EDA and V120 but does not contain EDB, while the

same sample also can contain an oncofetal fibronectin that contains EDA, EDB and V120. Such a sample can further be used to identify the likely cell source, tissue source or organ source of the oncofetal fibronectin, or can be used to identify the unlikely cell source, tissue source or organ source of the oncofetal fibronectins. As is known in the art, different cell and tissue types are known to produce two or more different oncofetal fibronectins. The different oncofetal fibronectin indicating molecules present in the sample can be compared to the oncofetal fibronectins known to be produced by one or more cell or tissue types to identify the likely or unlikely cell or tissue source(s) of an oncofetal fibronectin indicating molecule. Similarly, the relative levels of different oncofetal fibronectin indicating molecules present in the sample can be used to identify the likely cell source, tissue source or organ source of an oncofetal fibronectin indicating molecule, or can be used to identify the unlikely cell source, tissue source or organ source of an oncofetal fibronectin indicating molecule. Different cell and tissue types produce known ratios of two or more different oncofetal fibronectins. The relative amount of different oncofetal fibronectin indicating molecules present in the sample can be compared to the relative amount of oncofetal fibronectins known to be produced by one or more cell or tissue types to identify the likely or unlikely cell or tissue source(s) of an oncofetal fibronectin indicating molecule.

[0684] In another embodiment, characterizing an oncofetal fibronectin indicating molecule or measuring two or more different oncofetal fibronectin indicating molecules in a sample can be used to indicate the presence of a health problem associated with oncofetal fibronectin. For example, when the type of sample collected from a subject typically does not contain detectable amounts of a particular type of an oncofetal fibronectin indicating molecule when collected from normal subjects, detecting any amount of that particular type of an oncofetal fibronectin indicating molecule in such a sample can indicate the presence of a health problem associated with oncofetal fibronectin in the subject; this can occur even when a sample does typically contain a baseline amount of oncofetal fibronectin, provided that the baseline amount is of a different oncofetal fibronectin. In another example, when the type of sample collected from a subject typically contains a baseline amount of a particular type of oncofetal fibronectin indicating molecule when collected from normal subjects, detecting an amount of that type of oncofetal fibronectin indicating molecule greater than the baseline amount can indicate the presence of a health problem associated with oncofetal fibronectin in the subject. In another example, when the type of sample collected from a subject typically contains a ratio of two or more particular types of oncofetal fibronectin indicating molecules when collected from normal subjects, detecting a ratio of those types of oncofetal fibronectin indicating molecules that is different than the normal ratio can indicate the presence of a health problem associated with oncofetal fibronectin in the subject.

[0685] 1. Compounds and Compositions in Detecting Oncofetal Fibronectin

[0686] Any known assay procedure, including those that can be adapted for use in combination with fibronectin or oncofetal fibronectin binding partners, such as, for example, lateral flow devices, can be used in the systems and methods provided herein. Examples of assay procedures are protein

binding assays (e.g., dot blot analysis and Western blot analysis) and nucleic acid molecule hybridization assays (e.g., Northern blot analysis, Southern blot analysis and FISH). These can be practiced with any suitable format, including, for example, immunological methods, such as lateral flow and dip stick formats, mass spectrometry formats and others.

[0687] Binding assays, including competitive and non-competitive binding assays, are among those that can be used for determination of the presence or amount of analyte in a subject sample and are exemplified herein. It is understood that binding assays are provided for exemplification and that the methods and systems provided herein have broad applicability to a variety of sample types for a variety of different indications.

[0688] A number of different types of binding assays using a variety of protocols and labels are well known. Binding assays can be performed in a single liquid phase, or a binding partner can be immobilized to a solid support upon which the assay is performed. Sandwich assays can be performed. Competitive assays can be performed. The reaction steps can be performed simultaneously or sequentially, as will be known to those skilled in the art. Threshold assays can be performed, where only analyte levels of above a specified level or rate yield a positive result. Assay formats include, but are not limited to, for example, assays performed in test tubes, on membranes, in wells, in multi-well plates, on microarrays, on chromatographic test strips, as well as dipstick, lateral flow, vertical flow, or migratory format assays.

[0689] a. Molecules that Indicate the Presence of Oncofetal Fibronectin

[0690] Molecules detected in the methods provided herein can indicate the presence of oncofetal fibronectin in a subject. Molecules that can indicate the presence of oncofetal fibronectin in a subject include the oncofetal fibronectin protein, a fragment of the oncofetal fibronectin protein, mRNA encoding oncofetal fibronectin, a fragment of mRNA encoding oncofetal fibronectin, or an amplified nucleic acid molecule complementary to all or a portion of mRNA encoding oncofetal fibronectin (e.g., cDNA). For the detection methods described below, methods for detecting oncofetal fibronectin can apply to any of the oncofetal fibronectin indicating molecules provided herein, whether or not the particular detection method explicitly so indicates, subject to limits clear to one skilled in the art (e.g., use of PCR methods to amplify a molecule that indicates the presence of oncofetal fibronectin is not used to amplify the oncofetal fibronectin protein, protein fragments or autoantibodies).

[0691] In addition to the detection of an oncofetal fibronectin protein or nucleic acid molecule, or as an alternative therefor, one or more autoantibodies of oncofetal fibronectin, or antibody fragments thereof, can be detected using the methods provided herein, as will be understood by one skilled in the art. The presence of an autoantibody for oncofetal fibronectin can evidence the presence of oncofetal fibronectin in a subject. Thus, detection of the autoantibody can be used as an indicator of the presence of oncofetal fibronectin in a subject. Any of the protein detection methods described herein in relation to detection of oncofetal

fibronectin protein also can be used to detect an autoantibody to oncofetal fibronectin or to a nucleic acid encoding oncofetal fibronectin.

[0692] b. Fibronectin Portions Indicative of Oncofetal Fibronectin

[0693] The method provided herein can employ detection of a fibronectin region indicative of oncofetal fibronectin, or a fragment thereof. Regions indicative of oncofetal fibronectin include, but are not limited to, EDA, EDB, IIICS, splice variants of IIICS such as V64, V89, V95 and V120 and fragments of IIICS such as CS1, CS2, CS3, CS4, CS5 and CS6. The sequence of nucleotides of nucleic acid encoding human fibronectin EDA, EDB and IIICS regions and oncofetal fibronectins are known in the art and are available in public databases. Exemplary sequences are set forth in SEQ ID Nos:1-8 and 14-35. The amino acid and nucleic acid molecules for a variety of additional species including rat, mouse, chicken, cow and *Xenopus laevis* also are known and readily available in public databases. Allelic variants and other variants are known and/or can be identified.

[0694] Detection of a fibronectin region indicative of oncofetal fibronectin can be performed by a variety of methods known in the art or disclosed herein. For example, a fibronectin region indicative of oncofetal fibronectin can be detected using a protein that binds to an oncofetal fibronectin indicating region of a fibronectin protein. For example, a variety of anti-oncofetal fibronectin antibodies are known in the art, including IST-9, DH1, BC-1, L19, ME4C, H10, A134, C6, FDC-6, 5C10, X18A4, X20C4 and X8E3.

[0695] A fibronectin region indicative of oncofetal fibronectin also can bind to an integrin. For example, EDA can bind to $\alpha_4\beta_1$ integrin and $\alpha_9\beta_1$ integrin. The amino acid sequence EDGIHEL of EDA (EDA amino acids 40-46) can bind to $\alpha_4\beta_1$ integrin and $\alpha_9\beta_1$ integrin. IIICS can bind to $\alpha_4\beta_1$ integrin, $\alpha_4\beta_7$ integrin and heparin. The V95 splice variant of IIICS can bind to heparin. CS1 and CS5 can bind to $\alpha_4\beta_1$ integrin and $\alpha_4\beta_7$ integrin. The IIICS amino acid sequence LDV (IIICS amino acids 20-22) can bind to $\alpha_4\beta_1$ integrin and $\alpha_4\beta_7$ integrin. The IIICS amino acid sequence REDV (IIICS amino acids 100-103) can bind to $\alpha_4\beta_1$ integrin and $\alpha_4\beta_7$ integrin.

[0696] An oncofetal fibronectin region can be identified by detecting a glycosylation indicative of oncofetal fibronectin. For example, EDB can contain one or more N-linked glycosylation sites. IIICS can contain one or more N-linked glycosylation sites and from 1 to 6 or about 6 O-linked glycosylation sites, particularly IIICS can contain an O-glycosylation at threonine 33 of IIICS.

[0697] Detection of a fibronectin region indicative of oncofetal fibronectin can be performed by detecting one or more proteolysis fragments. For example, oncofetal fibronectin can yield trypsin fragments of 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa and/or 55 kDa. Typically, these six trypsin fragments can bind to the antibody FDC-6. In one example of an oncofetal fibronectin, trypsin fragments from a oncofetal fibronectin can be 200 kDa, 120 kDa or 55 kDa, where each smaller fragment represents a product of further trypsin cleavage of a larger fragment. In another example, trypsin fragments from an oncofetal fibronectin can be 235 kDa, 160 kDa or 65 kDa, where each smaller fragment represents a product of further trypsin cleavage of a larger fragment.

[0698] Oncofetal fibronectin can yield cathepsin D fragments of 110 kDa and/or 85 kDa. Typically these two cathepsin D fragments can bind to the antibody FDC-6. Oncofetal fibronectin also can be a fibronectin that yields thermolysin fragments of 120 kDa, 85 kDa and/or 35 kDa. Typically the 120 kDa and 85 kDa can bind to the antibody BC-1 and the 85 kDa fragment represents a product of further thermolysin cleavage of the 120 kDa fragment. Oncofetal fibronectin can be a fibronectin that yields an *Achromobacter* protease I fragment of 14 kDa, where this fragment typically can bind to the antibody FDC-6.

[0699] Nucleic acid molecules can indicate the presence of oncofetal fibronectin in a subject. Nucleic acid molecules indicative of the presence of oncofetal fibronectin or complements thereto also can be detected using the methods provided herein or known in the art.

[0700] A nucleic acid molecule indicative of the presence of oncofetal fibronectin or of the complement thereof, or of a fragment thereof can be detected herein. A nucleic acid molecule or fragment or complement thereof that can indicate the presence of oncofetal fibronectin indicates the presence of a fibronectin polypeptide region indicative of oncofetal fibronectin. For example, a detection of a nucleic acid molecule or fragment thereof can indicate the presence of a EDA, EDB or IIICS region, including the V0, V64, V89, V95 and V120 splice variants of the IIICS region. See, e.g., SEQ ID Nos. 28-35. Any of a variety of methods for detecting nucleic acid molecules or fragments thereof can be employed to detect a fibronectin polypeptide region indicative of oncofetal fibronectin. For example, a method can be used where the presence of any nucleic acid molecule indicates the presence of oncofetal fibronectin (e.g., where primers are designed to hybridize with the EDA, EDB or IIICS region, presence of any amplified nucleic acid molecules can indicate the presence of oncofetal fibronectin). In another example, presence of a nucleic acid molecule of an expected size can indicate the presence of oncofetal fibronectin (e.g., when a primer is located near EDA, EDB or IIICS encoding regions, small amplified nucleic acid molecules can indicate fibronectin without the EDA, EDB or IIICS encoding regions and large amplified nucleic acid molecules can indicate the presence of oncofetal fibronectin). In another example, presence of a nucleic acid molecule having an expected nucleotide sequence or sequence complementary thereto can indicate the presence of oncofetal fibronectin (e.g., when a Southern blot probe complementary to the EDA, EDB or IIICS encoding regions is used, binding of the probe to a sample band can indicate the presence of oncofetal fibronectin).

[0701] c. Autoantibody to Oncofetal Fibronectin or to a Nucleic Acid Molecule Encoding Oncofetal Fibronectin

[0702] Autoantibodies to oncofetal fibronectin or to a nucleic acid encoding oncofetal fibronectin, can indicate the presence of oncofetal fibronectin in a subject. Autoantibodies can be detected using the methods provided herein or known in the art. Autoantibodies are produced by a subject in response to a subject's own antigen. In the present methods, autoantibodies are produced in response to oncofetal fibronectin or a nucleic acid molecule encoding oncofetal fibronectin. Autoantibodies include endogenous antibodies that specifically bind to oncofetal fibronectin at one or more regions indicative of oncofetal fibronectin, including EDA,

EDB, IIICS, splice variants of IIICS such as V64, V89, V95 and V120 and peptide fragments of IIICS such as CS1, CS2, CS3, CS4, CS5 and CS6. Autoantibodies also can bind specifically to oncofetal fibronectin at one or more post-translational modification sites indicative of oncofetal fibronectin, such as one or more N-linked glycosylation sites in EDB or one or more N-linked glycosylation sites or one or more O-linked glycosylation sites in IIICS, particularly the O-glycosylation at threonine 33 of IIICS. Autoantibodies also can bind specifically to nucleic acid molecules encoding oncofetal fibronectin, such as a nucleic acid molecule encoding EDA, EDB, or a splice variant of IIICS such as V64, V89, V95 or V120.

[0703] Autoantibodies can be detected in any of a variety of samples, such as a tissue sample or a body fluid sample, including urine, lymph, blood, plasma, serum, saliva, cervical fluid, cervicovaginal fluid, vaginal fluid, breast fluid, breast milk, synovial fluid, semen, seminal fluid, stool, sputum, cerebral spinal fluid, tears, mucus, interstitial fluid, follicular fluid, amniotic fluid, aqueous humor, vitreous humor, peritoneal fluid, ascites, sweat, lymphatic fluid, lung sputum and lavage or purified fractions thereof. Autoantibodies can be detected by any of a variety of methods, including, for example, methods that include binding the autoantibody with a binding partner, such as described herein.

[0704] d. Binding Partner

[0705] Assays for detection of an oncofetal fibronectin indicating molecule can employ binding partners to aid in or permit detection. Binding partners can be used in a variety of roles in the methods provided herein. For example, binding partners can be used to bind an oncofetal fibronectin indicating molecule to a solid support, to increase the relative purity of an oncofetal fibronectin indicating molecule in a sample, to provide a detectable signal that indicates the presence of an oncofetal fibronectin indicating molecule in a sample, to provide a binding surface to which a detectable label can bind to indicate the presence of an oncofetal fibronectin indicating molecule in a sample, to amplify oncofetal fibronectin encoding nucleic acid molecules or complements thereto, and/or to indicate the presence of a specific domain of an oncofetal fibronectin indicating molecule in a sample.

[0706] In one embodiment, a binding partner can specifically bind to a fibronectin indicating molecule; thus, the binding partner can specifically bind to a fibronectin indicating molecule with increased a specificity relative to other molecules present in a sample. In another embodiment, a binding partner can specifically bind to an oncofetal fibronectin indicating molecule; thus, the binding partner can specifically bind to an oncofetal fibronectin indicating molecule with increased a specificity relative to a non-oncofetal fibronectin indicating molecule and/or other molecules present in a sample.

[0707] Exemplary binding partners include, but are not limited to, antibodies, antibody fragments, enzymes, metal ions, proteins, peptides, immunoglobulins, nucleic acid molecules, nucleic acid analogs, organic compounds, carbohydrates, lectins, dyes, reducing agents, energy absorbing molecules, affinity capture moieties, photolabile attachment molecules and combinations thereof.

[0708] A binding partner can be any of a variety of compounds that specifically bind to fibronectin protein.

Examples of compounds that bind to fibronectin protein include integrins such as integrin $\alpha_3\beta_1$, integrin $\alpha_4\beta_1$, integrin $\alpha_4\beta_7$, integrin $\alpha_5\beta_1$, integrin $\alpha_8\beta_1$, integrin $\alpha_9\beta_1$, integrin $\alpha_v\beta_1$, integrin $\alpha_v\beta_3$, integrin $\alpha_v\beta_5$, integrin $\alpha_v\beta_6$ and integrin $\alpha_{IIIb}\beta_3$; heparin; fibrin; collagen; gelatin; and antibodies such as IST-4 and 3E3.

[0709] A binding partner also can include compounds that bind specifically to oncofetal fibronectin protein. An oncofetal fibronectin binding partner can bind to a portion of oncofetal fibronectin that is not present or is not accessible in non-oncofetal fibronectin. Exemplary portions of oncofetal fibronectin not present in non-oncofetal fibronectin include amino acid regions such as EDA, EDB and V64, V89, V95 and V120 of IIICS. Additional exemplary portions of oncofetal fibronectin not present in non-oncofetal fibronectin include post-translationally modified regions such as O-glycosylated regions. Exemplary portions of oncofetal fibronectin not accessible in non-oncofetal fibronectin include amino acid regions present in oncofetal and non-oncofetal fibronectin, but that are conformationally different, for example, as a result of expression of amino acids not present in non-oncofetal fibronectin or as a result of post-translational modification not present in non-oncofetal fibronectin. Examples of compounds that bind preferentially to oncofetal fibronectin or antibodies therefor include antibodies, but are not limited to BC-1, FDC-6, L19, ME4C, X18A4, 5C10, IST-9 and DH-1.

[0710] i. Antibodies

[0711] Any antibody, including polyclonal or monoclonal antibodies, or any antigen-binding fragment thereof, include, but are not limited to, a Fab fragment or an scFv, that binds to an oncofetal fibronectin indicating molecule can be employed. For example, a mobile, labeled mouse monoclonal anti-oncofetal fibronectin antibody can be used to bind to oncofetal fibronectin protein in a sample and a polyclonal goat anti-fibronectin antibody immobilized on a test strip can be used to bind to oncofetal fibronectin protein to form a detectable sandwich complex indicating the presence of oncofetal fibronectin. In another example, a mouse monoclonal anti-oncofetal fibronectin antibody immobilized on a test strip can be used to bind to oncofetal fibronectin protein in a sample and a labeled goat polyclonal anti-oncofetal-fibronectin antibody conjugate can be used to bind to oncofetal fibronectin protein to form a detectable sandwich complex indicating the presence of oncofetal fibronectin protein in the sample. In another example, a membrane can be used to non-specifically bind to and immobilize oncofetal fibronectin protein and the immobilized oncofetal fibronectin protein can be exposed to a labeled anti-oncofetal fibronectin antibody conjugate, which can be detected to indicate presence of the oncofetal fibronectin protein in the sample. In another example, immobilized oncofetal fibronectin protein can be exposed to, for example, mouse anti-oncofetal fibronectin antibody and then exposed to a labeled polyclonal goat anti-mouse antibody complex to indicate the presence of oncofetal fibronectin protein in the sample.

[0712] Specifically binding antibody fragments for use in the methods described herein can be made from the respective monoclonal or polyclonal antibodies by conventional enzyme or chemical fragmentation procedures or can be made by routine recombinant methods. Various procedures

are known (see, e.g., Tijssen, P. *Laboratory Techniques in Biochemistry and Molecular Biology: Practice and Theories of Enzyme Immunoassays*, Elsevier, New York (1985)). For example, monoclonal antibody FDC-6 (ATCC Accession Number ATCC HB 9018) can be exposed to a protease such as papain or pepsin, respectively, to form Fab and F(ab)₂ fragments, as is known in the art. In addition, the VL and VH of FDC-6 can be joined using recombinant methods using a synthetic linker that permits them to be expressed as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see e.g., Bird et al., *Science*, 242:423-426 (1988); and Huston et al., *Proc. Natl. Acad. Sci. USA*, 85:5879-5883 (1988)). Bivalent, bispecific antibodies (i.e., diabodies) can be formed in which FDC-6 VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see e.g., Holliger, P., et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993); Poljak, R. J., et al., *Structure* 2:1121-1123 (1994)).

[0713] The antibodies can be raised and purified using methods known to those of skill in the art or obtained from publicly available sources. For example, methods for raising and purifying antibodies can be similar to the methods used in raising and purifying monoclonal antibody FDC-6 (deposited at the American Type Culture Collection as accession number ATCC HB 9018; see U.S. Pat. No. 4,894,326; see, also, Matsuura et al., *Proc. Natl. Acad. Sci. USA*, 82:6517-6521 (1985); see, also, U.S. Pat. Nos. 4,919,889; 5,096,830; 5,185,270; 5,223,440; 5,236,846; 5,281,522; 5,468,619 and 5,516,702). Anti-fibronectin antibodies and anti-oncofetal fibronectin antibodies can be formed into conjugates or immobilized on a solid support using methods disclosed herein or known in the art.

[0714] a. Antibodies for Oncofetal Fibronectin

[0715] In one embodiment, antibodies used herein can specifically bind to oncofetal fibronectin protein. Antibodies used herein can specifically bind an epitope present in oncofetal fibronectin protein. Polyclonal antibodies typically bind to two or more epitopes present in oncofetal fibronectin protein. A monoclonal antibody typically binds a single epitope present in oncofetal fibronectin protein.

[0716] In another example, an epitope present in oncofetal fibronectin can include an epitope that can be present in oncofetal fibronectin, such as an epitope that includes EDA, EDB or IIICS domains of oncofetal fibronectin. Antibodies that bind to epitopes present in oncofetal fibronectins include FDC-6 (ATCC Accession Number ATCC HB 9018), IST-9 (Carnemolla et al., *FEBS Lett.* 215:269-273 1987; Accurate Chemical and Scientific Corp., Westbury, N.Y.), DH1 (Vartio et al., *J. Cell Sci.* 88:419-430 1987), BC-1 (Carnemolla et al., *J. Cell Biol.* 108:1139-1148 1989), L19 (U.S. Pat. App. No. 20030176663), ME4C (Giovannoni et al., *Nucleic Acids Res.* 29:e27 (2001); the ME4C scFv nucleic acid and amino acid sequences are provided in SEQ ID Nos:9 and 10, and are available at GenBank accession no. AJ297960), H10 (U.S. Pat. App. No. 20030176663), A134 (Islami et al., *Eur. J. Obstet. Gynecol. Reprod. Biol.*, 97:40-45 2001), 5C10 (Mandel et al., *APMIS* 100:817-826 1992)

and X18A4, X20C4 and X8E3 (U.S. Pat. No. 5,523,229; ATCC accession numbers HB-11587, HB-11589 and HB-11588, respectively).

[0717] b. Conjugation of the Antibody to a Label

[0718] In one embodiment, an antibody that binds to an analyte of interest can be conjugated to a detectable label. The detectable label used in the antibody conjugate can be any physical or chemical label that can be detected and can be distinguished from other compounds and materials in the assay. For example, an antibody conjugate can be detected on a solid support using a reader, such as a reflectance reader. In one example, a mouse monoclonal anti-onfFN antibody (see, e.g., U.S. Pat. No. 5,281,522), can be conjugated to latex particles containing a blue dye or other spectrometrically detectable label. In another example, a goat polyclonal antibody to human fibronectin can be conjugated to a colloidal gold label.

[0719] Various antibody labels are well known to those of skill in the art. The labels include, but are not limited to enzyme-substrate combinations that produce color upon reaction, colored particles, such as latex particles, quantum dots, colloidal metal or metal or carbon sol labels, fluorescent labels and liposome or polymer sacs, which are detected due to aggregation of the label. In one particular embodiment, the label is a colored latex particle. In an alternative embodiment, colloidal gold is used in the labeled antibody conjugate.

[0720] The label can be derivatized for linking to antibodies, such as by attaching functional groups, such as carboxyl groups to the surface of a particle to permit covalent attachment of antibodies. Antibodies can be conjugated to the label using well known coupling methods. Coupling agents such as glutaraldehyde or carbodiimide can be used. The labels can be bonded or coupled to the antibodies by chemical or physical bonding. In an exemplary embodiment, a carbodiimide coupling reagent, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDAC), is used to link antibodies to latex particles.

[0721] ii. Nucleic Acid Molecules

[0722] A binding partner also can be a molecule that specifically binds a fibronectin-encoding nucleic acid molecule, a molecule that specifically binds a sequence complementary to a fibronectin-encoding nucleic acid molecule, amplicates thereof, or fragments thereof (the group of which is herein referred to as an oncofetal fibronectin nucleic acid molecule). In one embodiment, at least one binding partner binds specifically to a nucleic acid molecule encoding fibronectin or a complement thereto. In another embodiment, two or more binding partners can bind specifically or preferentially to an oncofetal fibronectin encoding nucleic acid molecule or a complement thereto. Binding partners can include single stranded or double stranded nucleic acid molecules, oligonucleotides, primers, deoxyribonucleic acid molecules (DNA), ribonucleic acid molecules (RNA), nucleic acid homologs such as peptide nucleic acids and hybrids thereof. Binding partners also can include antibodies, antibody fragments, enzymes, metal ions, proteins, peptides, immunoglobulins, carbohydrates, lectins, dyes, reducing agents, energy absorbing molecules, affinity capture moieties, photolabile attachment molecules and combinations thereof.

[0723] A binding partner that binds a fibronectin encoding nucleic acid molecule or complement thereto can specifically bind to a fibronectin-encoding nucleic acid molecule or an oncofetal fibronectin-encoding nucleic acid molecule or a complement thereto. A binding partner that binds an oncofetal fibronectin encoding nucleic acid molecule or complement thereto can specifically bind to an oncofetal fibronectin-encoding nucleic acid molecule or a complement thereto. Typically, such a binding partner will be complementary to nucleic acid encoding a fibronectin or oncofetal fibronectin nucleic acid molecule or complement thereof, but not present in other (non-fibronectin-encoding) nucleic acid molecules.

[0724] An oncofetal fibronectin binding partner can preferentially bind to an oncofetal fibronectin-encoding nucleic acid molecule or complement thereto relative to binding a non-oncofetal fibronectin-encoding nucleic acid molecule or complement thereto. Typically, such a binding partner will be complementary to a nucleic acid molecule that encodes oncofetal fibronectin or complement thereto. For example, a binding partner that preferentially binds an oncofetal fibronectin-encoding nucleic acid can bind to a nucleic acid sequence encoding the EDA, EDB or IIICS region of oncofetal fibronectin, including, but not limited to, a nucleic acid molecule encoding V120, V95, V89 or V64.

[0725] Binding partners used to detect the presence of an oncofetal fibronectin nucleic acid molecule or complement thereto can be conjugated with a moiety useful for binding, isolation, or detection. A moiety can include a bindable moiety such as biotin, nickel, magnetic bead, or other composition or compound used for binding or isolation. For example, a binding partner can be conjugated with a biotin moiety which can be used for binding or isolation when used in conjunction with avidin or streptavidin. Such a moiety also can be a detectable moiety such as a fluorescent compound, a compound containing a radionuclide, a quantum dot, colloidal metal, or any other moiety that can be used for detection by methods including, but not limited to, fluorimetry, absorption, reflection, visible inspection and scintillation. Binding partners also can be immobilized on a solid substrate. Methods for conjugating or immobilizing binding partners such as nucleic acids and nucleic acid analogs are known in the art and can be used in the methods herein.

[0726] Binding partners also can serve as nucleotide synthesis primers for amplification of an oncofetal fibronectin encoding nucleic acid molecule or complement thereto. Such binding partners are typically nucleic acid molecules or nucleic acid analogs that contain a 3' hydroxy moiety accessible for phosphodiester bond formation or other nucleotide synthesis steps. Methods for preparing primers for nucleotide synthesis reactions are well known in the art and can be used in the methods herein.

[0727] Nucleic acid molecule binding partners can bind to oncofetal fibronectin-encoding nucleic acid molecules or nucleic acid molecules complementary thereto. For example, methods provided herein include nucleic acid molecule synthesis methods in which nucleic acid molecules complementary to oncofetal fibronectin-encoding nucleic acid molecules are formed. Methods for detecting oncofetal fibronectin-encoding nucleic acid molecules are therefore

intended to also detect nucleic acid molecules complementary to oncofetal fibronectin-encoding nucleic acid molecules.

[0728] In one embodiment, binding partners can be nucleic acid molecules. Typically nucleic acid molecule binding partners are large enough in length to specifically bind to oncofetal fibronectin-encoding nucleic acid molecules or complements thereto, without also specifically binding other non-fibronectin-encoding nucleic acid molecules. In one example, nucleic acid molecule binding partners preferentially bind to oncofetal fibronectin-encoding nucleic acid molecules or complements thereto, without also specifically binding other non-oncofetal fibronectin-encoding nucleic acid molecules or complements thereto. For example, a nucleic acid molecule binding partner can specifically bind to a nucleic acid molecule encoding EDA, EDB or IIICS or complement thereto, or a fragment thereof.

[0729] Nucleic acid molecule binding partners can vary in length and can contain extra nucleotides in addition to the oncofetal-fibronectin encoding (or complement thereto) portion of the nucleic acid molecule binding partner (e.g., a transcriptional start site or a chip hybridization sequence). Exemplary lengths of the oncofetal fibronectin-encoding portion (or complement thereto) of a nucleic acid molecule binding partner are at least 5 nucleotides, at least 6 nucleotides, at least 7 nucleotides, at least 8 nucleotides, at least 10 nucleotides, at least 12 nucleotides, at least 15 nucleotides, at least 18 nucleotides, at least 20 nucleotides, at least 25 nucleotides, at least 35 nucleotides, at least 40 nucleotides, at least 50 nucleotides, or at least about 5 nucleotides, at least about 6 nucleotides, at least about 7 nucleotides, at least about 8 nucleotides, at least about 10 nucleotides, at least about 12 nucleotides, at least about 15 nucleotides, at least about 18 nucleotides, at least about 20 nucleotides, at least about 25 nucleotides, at least about 35 nucleotides, at least about 40 nucleotides, at least about 50 nucleotides, or more.

[0730] Similarly, the oncofetal fibronectin-encoding nucleic acid molecule or complement thereto need not be bound by the binding partner along the entire length of the oncofetal fibronectin-encoding sequence. In some embodiments, the oncofetal fibronectin-encoding nucleic acid molecule or complement thereto bound by the binding partner is the full length nucleic acid molecule, where the binding partner specifically binds a portion of the oncofetal fibronectin-encoding nucleic acid molecule or complement thereto, without also specifically binding other non-fibronectin-encoding nucleic acid molecules. In one example, a nucleic acid molecule binding partner preferentially binds a portion of an oncofetal fibronectin-encoding nucleic acid molecule or complement thereto, without also specifically binding other non-oncofetal fibronectin-encoding nucleic acid molecules or complements thereto. For example, a nucleic acid molecule binding partner can specifically bind to the portion of the oncofetal fibronectin-encoding nucleic acid molecule encoding EDA, EDB or IIICS or complements thereto.

[0731] In other embodiments, the oncofetal fibronectin-encoding nucleic acid molecule or complement thereto bound by the binding partner is a fragment of the full length nucleic acid molecule, where the binding partner specifically binds the fragment of the oncofetal fibronectin-encoding nucleic acid molecule or complements thereto, without also

specifically binding other non-fibronectin-encoding nucleic acid molecules. In one example, a nucleic acid molecule binding partner preferentially binds to a fragment of an oncofetal fibronectin-encoding nucleic acid molecule or complement thereto, without also specifically binding other non-oncofetal fibronectin-encoding nucleic acid molecules or complements thereto. For example, a nucleic acid molecule binding partner can specifically bind to a fragment of the oncofetal fibronectin-encoding nucleic acid molecule encoding at least a portion of EDA, EDB or IIICS or complement thereto. Fragments of an oncofetal fibronectin-encoding nucleic acid molecule or complement thereto can be generated by any of a variety of methods provided herein or otherwise known in the art, including enzymatic, chemical or physical cleavage of nucleic acid molecules and also including nucleic acid synthesis methods such as PCR or RT-PCR, where only a portion of the oncofetal fibronectin-encoding nucleic acid molecule or complement thereto is amplified. In some instances, the fragment will contain only nucleotides encoding EDA, EDB or IIICS or complement thereto, where the fragment can contain the entire splice region or less than the entire splice region or complement thereto. In regard to IIICS, a fragment can contain only nucleotides encoding V120, V95, V89 or V64 or complement thereto; or only the splice regions encoding amino acids 1-25, amino acids 26-89, or amino acids 90-102 or complement thereto. Fragments also can include regions adjacent to any of the aforementioned splice regions or complements thereto. For example, a fragment can contain 10 nucleotides 3' and 10 nucleotides 5' of the EDA splice region or complement thereto. Such fragments can be used, for example, in mass spectrometric detection methods to distinguish between nucleic acid molecules containing an oncofetal fibronectin splice region and nucleic acid molecules lacking the oncofetal fibronectin splice region. For example, a fragment can contain 10 nucleotides 3' and 10 nucleotides 5' of the EDA splice region or complement thereto, and nucleic acid molecules containing an oncofetal fibronectin splice region (110 nucleotide in length) can be distinguished from nucleic acid molecules lacking the oncofetal fibronectin splice region (20 nucleotides in length).

[0732] Fragments can vary in length according to any of a variety of factors known to one skilled in the art, including region(s) of oncofetal fibronectin to be detected, detection method and sample. Fragments, such as fragments formed by nucleic acid synthesis reactions, can contain extra nucleotides in addition to the oncofetal fibronectin-encoding portion of the nucleic acid molecule (e.g., a transcriptional start site or a non-oncofetal fibronectin encoding hybridization sequence). Exemplary lengths of the oncofetal fibronectin-encoding portion or complement thereto of a fragment (or complement thereto) are at least 5 nucleotides, at least 6 nucleotides, at least 7 nucleotides, at least 8 nucleotides, at least 10 nucleotides, at least 12 nucleotides, at least 15 nucleotides, at least 18 nucleotides, at least 20 nucleotides, at least 25 nucleotides, at least 35 nucleotides, at least 40 nucleotides, at least 50 nucleotides, or at least about 5 nucleotides, at least about 6 nucleotides, at least about 7 nucleotides, at least about 8 nucleotides, at least about 10 nucleotides, at least about 12 nucleotides, at least about 15 nucleotides, at least about 18 nucleotides, at least about 20

nucleotides, at least about 25 nucleotides, at least about 35 nucleotides, at least about 40 nucleotides, at least about 50 nucleotides, or more.

[0733] iii. Binding Partners to Autoantibodies

[0734] A binding partner can specifically bind to an autoantibody to oncofetal fibronectin or a nucleic acid molecule encoding oncofetal fibronectin. Such a binding partner can include antibodies, antibody fragments, enzymes, metal ions, proteins, peptides, immunoglobulins, carbohydrates, lectins, dyes, reducing agents, energy absorbing molecules, affinity capture moieties, photolabile attachment molecules and combinations thereof. Binding partners also can include single stranded or double stranded nucleic acids, oligonucleotides, primers, deoxyribonucleic acid molecules (DNA), ribonucleic acid molecules (RNA), nucleic acid homologs such as peptide nucleic acids and hybrids thereof.

[0735] A binding partner to an autoantibody also can include oncofetal fibronectin protein or a fragment thereof, or a nucleic acid molecule encoding oncofetal fibronectin or a fragment thereof. The autoantibodies can specifically bind to oncofetal fibronectin protein at one or more regions indicative of oncofetal fibronectin, including EDA, EDB, IIICS, splice variants of IIICS such as V64, V89, V95 and V120 and fragments of IIICS such as CS1, CS2, CS3, CS4, CS5 and CS6. Autoantibodies also can specifically bind to oncofetal fibronectin protein at one or more post-translational modification sites indicative of oncofetal fibronectin, such as one or more N-linked glycosylation sites in EDB or one or more N-linked glycosylation sites or one or more O-linked glycosylation sites in IIICS, particularly the O-glycosylation at threonine 33 of IIICS. Autoantibodies also can specifically bind to nucleic acid molecules encoding oncofetal fibronectin, such as a nucleic acid molecule encoding EDA, EDB, or a splice variant of IIICS such as V64, V89, V95 or V120. Accordingly, a protein or nucleic acid molecule that contains any of the regions indicative of oncofetal fibronectin, post-translation modifications indicative of oncofetal fibronectin, or nucleic acid molecule regions encoding a polypeptide region indicative of oncofetal fibronectin, can be used as a binding partner for an autoantibody in accordance with the methods provided herein. Fragments of a protein or nucleic acid molecule, such as a polypeptide or oligonucleotide, that contains any of the regions indicative of oncofetal fibronectin, post-translation modifications indicative of oncofetal fibronectin, or nucleic acid molecule regions encoding a polypeptide region indicative of oncofetal fibronectin, can be used as a binding partner for an autoantibody. In one example, a fragment such as a peptide or oligonucleotide, contains only a region indicative of oncofetal fibronectin, optionally also containing post-translation modifications indicative of oncofetal fibronectin, or nucleic acid molecule regions encoding a polypeptide region indicative of oncofetal fibronectin. For example, a polypeptide can contain only EDA, EDB, IIICS, splice variants of IIICS such as V64, V89, V95 and V120, peptide fragments of IIICS such as CS1, CS2, CS3, CS4, CS5 and CS6, or fragments thereof, which optionally contain one or more post-translational modification sites indicative of oncofetal fibronectin, such as one or more N-linked glycosylation sites in EDB or one or more N-linked glycosylation sites or one or more O-linked glycosylation sites in IIICS, particularly the O-glycosylation at threonine 33 of IIICS. In

another example, an oligonucleotide can contain only nucleotides encoding EDA, EDB, or a splice variant of IIICS such as V64, V89, V95 or V120. Exemplary polypeptide or oligonucleotide fragment sizes can range from 5 to 30 residues, 7 to 20 residues, or 9 to 15 residues, or about 5 to about 30 residues, about 7 to about 20 residues, or about 9 to about 15 residues.

[0736] Methods for detecting autoantibodies also can employ antibody binding partners that are not specific for anti-oncofetal fibronectin autoantibodies. For example, antibody binding partners such as anti-human antibodies, including anti-IgG, anti-IgA, anti-IgD, anti-IgE or anti-IgM antibodies, or fragments thereof, can be used in methods for detecting an anti-oncofetal fibronectin autoantibody. Antibody binding partners or fragments thereof can be polyclonal or monoclonal.

[0737] Binding partners for autoantibodies used in accordance with the methods provided herein can be used in the same manner as described herein for the use of binding partners for fibronectin or oncofetal fibronectin protein or binding partners for a nucleic acid molecule encoding oncofetal fibronectin. For example, such binding partners can be formed as conjugates or can be immobilized to a solid support. General methods and binding partners for detecting autoantibodies are known in the art and can be used in conjunction with the compounds, compositions and methods provided herein. Such methods and binding partners are exemplified in U.S. Pat. Nos. 20030232399 and 20040048320 and in WO/03 020115.

[0738] iv. Additional Binding Partners

[0739] A variety of additional compounds can be used as fibronectin binding partners of oncofetal fibronectin protein or oncofetal fibronectin encoding nucleic acids, including compounds that bind to fibronectin protein such as peptides and non-peptide organic compounds. Compounds that bind to fibronectin protein are known in the art and include, but are not limited to, small peptides such as GGWSHW (SEQ ID NO: 36) and related peptides and variants (e.g., cyclic peptides) thereof, as provided, for example, in U.S. Pat. No. 5,491,130. Also included are compounds that specifically bind to oncofetal fibronectin protein, including, but not limited to, diaryl alkylamines such as 2,2-diphenylethylamine, 2,2-diphenylpropylamine, 3,3-diphenylpropylamine, 2-naphthyl,2-phenylethylamine, 2-naphthyl,2-(2,6-dichlorophenyl)ethylamine, 2-phenyl,2-(4-trifluoromethylphenyl)ethylamine, 2-phenyl,2-(3,4-methylenedioxybenzyl)ethylamine, 2-phenyl,2-thienylethylamine, and derivatives thereof, which are known to bind to the EDB domain of oncofetal fibronectin. Such diaryl alkylamines are known in the art, as exemplified in the doctoral thesis of Jörg Scheuermann, submitted to the Swiss Federal Institute of Technology, Zurich, December 2002.

[0740] v. Binding Partners that Bind a Region of Oncofetal Fibronectin

[0741] As described herein, binding partners can be used to detect the presence of an oncofetal fibronectin indicating molecule. Binding partners also can be used to detect the presence of regions indicative of oncofetal fibronectin. For example, binding partners can be used to detect the presence of EDA, EDB, IIICS and any of the splice variants of IIICS including V0, V64, V89, V95 and V120, in an oncofetal

fibronectin indicating molecule. Binding partners can be used to detect one or more regions indicative of oncofetal fibronectin, two or more regions indicative of oncofetal fibronectin, or three or more regions indicative of oncofetal fibronectin.

[0742] Binding partners also can be used in the methods herein to detect a post translational modification indicative of oncofetal fibronectin. Post translational modifications include glycosylation such as O-linked and N-linked glycosylation, disulfide bridge formation, proteolysis, lipidylation and other known post translational modifications. Binding partners can be used to detect one or more post translational modifications indicative of oncofetal fibronectin, two or more post translational modifications indicative of oncofetal fibronectin, or three or more post translational modifications indicative of oncofetal fibronectin. Binding partners also can be used to detect combinations of regions indicative of oncofetal fibronectin and post translational modifications indicative of oncofetal fibronectin.

[0743] e. Non-Specific Binder

[0744] In order to improve detection of an oncofetal fibronectin indicating molecule, background signals can be reduced by, for example, removing, extracting, or decreasing binding of background material from a sample that can non-specifically bind to a binding partner. For example, a urine sample can be contacted with a non specific binder such as a non-specific binding compound or a non-specific binding surface prior to the sample contacting an analyte-specific antibody. For example, when using a lateral flow device, a zone containing a ligand, such as bovine serum albumin (BSA), can be included. This zone can remove background material from a sample solution prior to the sample reaching the zone where presence of an oncofetal fibronectin indicating molecule is to be detected.

[0745] Exemplary compounds for reducing non-specific binding include casein, albumin, substituted albumin (e.g., methylated albumin), IgG, antibodies that do not specifically bind to fibronectin or oncofetal fibronectin, and other proteins known to block non-specific protein interactions. In the case of nucleic acid molecule detection, compounds for reducing non-specific binding include salmon sperm DNA, herring sperm DNA, poly(dI-dC)-poly(dI-dC) and other nucleic acid molecules that block non-specific nucleic acid hybridization.

[0746] 2. Assays for Detecting Oncofetal Fibronectin Complexed with a Binding Partner

[0747] A variety of different assay procedures for detecting an oncofetal fibronectin indicating molecule, such as an oncofetal fibronectin protein, a nucleic acid molecule encoding oncofetal fibronectin or complement thereto, an autoantibody to oncofetal fibronectin protein or a nucleic acid molecule encoding oncofetal fibronectin, or a fragment thereof. Typically, assay procedures for detecting the presence of oncofetal fibronectin include, for example, a step of contacting a sample with a fibronectin or oncofetal fibronectin binding partner. Exemplary assay procedures include, but are not limited to, flow cytometry, Western blot analysis, dot blot analysis, Southern blot analysis, Northern blot analysis, sandwich assay, fluorescence polarization, test strip analysis, mass spectrometry and PCR-based methods. Assays can be performed with the oncofetal fibronectin indicating mol-

ecule immobilized on a solid support, with a fibronectin or oncofetal fibronectin binding partner immobilized on a solid support, or with no molecules immobilized on a solid support. In embodiments that include a solid support, any solid support provided herein or otherwise known in the art can be used including, but not limited to a microtiter plate, a microarray, a test strip, a mass spectrometry substrate, and a nitrocellulose membrane.

[0748] In conducting the assay, a subject sample is obtained. The samples that can be used include any sample provided herein or otherwise known in the art. The samples can contain, for example, fluid and particulate solids and, can be filtered prior to measuring an oncofetal fibronectin indicating molecule, for example, prior to application to an assay test strip. The sample can be removed from the subject using any method known in the art or provided herein, including using a swab having a fibrous tip, an aspirator, suction or lavage device, syringe, or any other known method of removing a bodily sample, including passive methods for collecting urine or saliva. The sample, particularly samples in or attached to an insoluble medium, can be extracted into a buffer solution and optionally filtered. In one embodiment, where oncofetal fibronectin is to be detected in a sample, the sample is obtained from in the vicinity of the point of a possible cervicovaginal lesion, the cervical canal, the cervical os, the ectocervix, the transition zone on the cervix between squamous and columnar cells (i.e., the squamocolumnar junction), the vagina, the posterior fornix, the portion of the vagina below the posterior fornix such as the lower third of the vagina, the labia, or combinations thereof, where the sample can be collected as a swab sample, using, for example, a swab having at its tip a fibrous material such as polyester, rayon, or cotton. When the sample is collected with a cotton swab, the assay methods are conducted on the swab and reagents are added to the swab.

[0749] The assay methods provided herein can generally be used for detecting oncofetal fibronectin indicating molecules that are proteins or for detecting oncofetal fibronectin indicating molecules that are nucleic acid molecules. Selection of the reagents and particular methodologies can vary according to the oncofetal fibronectin indicating molecule to be detected, as will be understood by one skilled in the art.

[0750] The assay methods provided herein can be used to detect binding of an oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner, and also can be used to detect competitive inhibition, where an oncofetal fibronectin indicating molecule in the sample competes with a known amount of oncofetal fibronectin indicating molecule or analog thereof for a predetermined amount of fibronectin or oncofetal fibronectin binding partner. For example, in an assay for oncofetal fibronectin protein, any oncofetal fibronectin protein present in the sample can compete with a known amount of labeled oncofetal fibronectin protein or a labeled analogue thereof for the fibronectin or oncofetal fibronectin binding partner. The amount of labeled fibronectin affixed to the solid phase or remaining in solution can be measured, and the measurement can be used to determine the amount of oncofetal fibronectin indicating molecule in the sample, using methods known in the art.

[0751] a. Solution Detection

[0752] Presence of an oncofetal fibronectin indicating molecule can be detected in solution. A sample can be

contacted with one or more fibronectin or oncofetal fibronectin binding partners and any complex formed between an oncofetal fibronectin indicating molecule and the binding partner can be detected. Detection of the complex can indicate the presence of an oncofetal fibronectin indicating molecule in a sample. Detection of complexes formed in solution are known in the art, such as detection of a signal indicative of complex formation, or detection of a substance with a molecular weight corresponding to complex formation. In some embodiments, detection of competition can be performed, where loss of a signal indicative of a complex can indicate the presence of an oncofetal fibronectin indicating molecule. Solution detection methods can be performed when detecting oncofetal fibronectin proteins or oncofetal fibronectin encoding nucleic acid molecules of complements thereto, as is understood in the art.

[0753] i. Signal Indicative of Complex Formation

[0754] Complex formation between an oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner can result in a signal indicative of complex formation. For example, complex formation can result in a change in a signal, a unique signal, an increasing signal, a decreasing signal, or a combination of signals. The signal indicative of complex formation can be detected by any of a variety of methods for detecting signals including, fluorescence polarization, fluorimetry, absorption, scintillation detection and agglutination, and can be configured in any of a number of experimental systems, such as a fluorimeter, flow cytometer, or microscope such as a confocal microscope. Exemplary signals indicative of complex formation can be agglutination of reaction components, a change in fluorescence polarization, a change in the intensity or wavelength of fluorescence, a change in the intensity or wavelength of absorption, a unique signal resulting from energy transfer such as FRET (fluorescence resonance energy transfer), or spatial proximity of two or more signals. Such signals indicative of complex formation can arise from a change resulting from a binding partner complexing with an oncofetal fibronectin indicating molecule (e.g., a conformational change in a binding partner that changes the signal of a detectable moiety), or from two or more binding partners binding to the same oncofetal fibronectin indicating molecule (e.g., a signal resulting from FRET between or among binding partners). Additionally, the above methods can be performed as competition assays, using, for example, a labeled oncofetal fibronectin indicating molecule or analog thereof, where disappearance of a signal as a result of competition from an oncofetal fibronectin indicating molecule in the sample can indicate the presence of the oncofetal fibronectin indicating molecule in the sample.

[0755] A variety of methods are known for identifying the presence and/or amount of an analyte in a solution without performing steps of separating the analyte from solution. Exemplified below are particular methods for performing such non-separation assays, however, a variety of additional methods are known in the art, as exemplified in Hussa, "The clinical marker hCG," Praeger Publishers (1987), pp 38-40.

[0756] In one embodiment, binding of one binding partner to an oncofetal fibronectin indicating molecule can provide a detectable signal that indicates the presence and/or level of the oncofetal fibronectin indicating molecule in the sample. For example, a binding partner labeled with a fluorescent

probe can provide a different fluorescence polarization signal when bound to an oncofetal fibronectin indicating molecule relative to the unbound signal. In another example, a binding partner can be bound by a molecule labeled with a fluorescent probe, where the molecule can be, for example, an oncofetal fibronectin indicating molecule or analog thereof, can provide a different fluorescence polarization signal when bound to oncofetal fibronectin indicating molecule from a sample relative to the signal when sample oncofetal fibronectin indicating molecule is not bound. Fluorescence polarization signals can vary according to the molecular weight of the dye-containing complex; when a fluorescent dye-labeled binding partner binds to an oncofetal fibronectin indicating molecule, the signal can be more polarized relative to the unbound form. Analogously, when a fluorescent dye-labeled oncofetal fibronectin indicating molecule or analog thereof binds to a binding partner, the signal can be more polarized relative to the unbound form, and when an oncofetal fibronectin indicating molecule in a sample displaces the fluorescently labeled oncofetal fibronectin indicating molecule or analog thereof, the signal can be less polarized relative to the bound form. Any of a variety of known dyes can be used in performing such methods, including fluorescein dyes, cyanine dyes, dansyl dyes, and polyazaindacene dyes, such as 4,4-difluoro-4-bora-3a,4a-diaza-s-indacene (BODIPY) dyes (Molecular Probes, Eugene, Oreg.; see, e.g., U.S. Pat. Nos. 6,323,186 and 6,005,113). Fluorescence polarization assays are known in the art and selection of dyes and assay conditions can be determined according to the assay design.

[0757] In some fluorescence polarization assays, the fibronectin or oncofetal fibronectin binding partner can be an antibody or an antibody fragment, for example FDC-6 or ME4C, labeled with a fluorescent dye. In embodiments where a larger mass difference between the oncofetal fibronectin indicating molecule and the antibody binding partner is desirable, the antibody binding partner can be an antibody fragment such as, for example, scFv, sc(Fv)₂, Fab, Fv, SIP, or another antigen-binding fragment provided herein or otherwise known in the art.

[0758] In other fluorescence polarization assays, an oncofetal fibronectin indicating molecule or analog thereof can be attached to a fluorescent dye, and presence of an oncofetal fibronectin indicating molecule in a sample can be determined by competition for binding to a fibronectin or oncofetal fibronectin binding partner. Exemplary analogs of an oncofetal fibronectin indicating molecule include oligopeptides or oligonucleotides containing all or a portion of EDA, EDB or IIICS, where any portion of such a region is selected according to its ability to specifically bind to a fibronectin or oncofetal fibronectin binding partner. For example, an oncofetal fibronectin protein analog can be a hexapeptide of the IIICS region, where the hexapeptide contains the sequence Val-Thr-His-Pro-Gly-Tyr (SEQ ID NO: 39) and having an O-glycosylation at the threonine residue, and can optionally contain a linker, and the analog can be attached to a fluorescent dye, either directly or through a linker, using methods known in the art.

[0759] In one embodiment, when two or more binding partners bind to the same oncofetal fibronectin indicating molecule, a first binding partner can transfer energy to a second binding partner to produce a signal that arises only when the first and second binding partners are spatially

proximal. For example, a chromophore on a first binding partner can, by fluorescence resonance energy transfer (FRET), transfer a quantum of energy to a proximal fluorophore of a second binding partner, whereupon the fluorophore of the second binding partner fluoresces. Such experiments can be configured such that fluorescence of the fluorophore of the second binding partner can only arise as a result of FRET, which can only arise if the chromophore and fluorophore are spatially close to one another. Detection of fluorescence from the fluorophore of the second binding partner can indicate the two binding partners are spatially proximal, which can indicate the presence of an oncofetal fibronectin indicating molecule. Methods of using FRET are well known in the art, and any of a variety of these methods can be used herein (see, e.g., Gaits et al., *Science STKE* 2003:PE3).

[0760] In another embodiment, spatial proximity of two different signals can indicate the presence of an oncofetal fibronectin indicating molecule in a sample. Binding of two differently labeled binding partners to one oncofetal fibronectin indicating molecule can result in a complex that provides two different signals, for example, fluorescence at two different wavelengths. Detection methods such as flow cytometry or confocal microscopy can be used to examine particular portions of a sample or small volumes of sample at any instant in time. Experimental conditions can be adapted such that the detection of two or more different signals at the same instance can indicate that two differently labeled binding partners are spatially proximate, which occurs when the two or more binding partners are bound to the same molecule, an oncofetal fibronectin indicating molecule. Thus, detection of two or more different signals at the same instance can indicate the presence of an oncofetal fibronectin indicating molecule in a sample.

[0761] In another embodiment, a polydentate binding partner, such as a bidentate binding partner can be used in agglutination assays that indicate the presence of an oncofetal fibronectin indicating molecule in a sample. A polydentate binding partner is able to bind two oncofetal fibronectin indicating molecules, which can result in linking together binding partners and oncofetal fibronectin indicating molecules into large complexes. Examples of polydentate binding partners include antibodies, which contain two binding sites and are, therefore, bidentate, and fusion proteins containing two or more regions that bind the same or different regions of oncofetal fibronectin protein (including a bidentate binding partner containing a first region that contains an oncofetal fibronectin fibronectin binding partner and a second region that contains a fibronectin or oncofetal fibronectin binding partner). Such large complexes can be detected by any of a variety of methods, such as, for example, spectrophotometric detection and visual inspection. An exemplary agglutination assay includes the use of gold sol particles coated with two or more of the same binding partner or two or more different binding partners, and color change by visual or spectrophotometric detection can indicate the presence and/or amount of an oncofetal fibronectin indicating molecule in a sample.

[0762] ii. Molecular Weight Corresponding to Complex

[0763] When an oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner form a complex, the mass of the complex is larger than the

masses of the oncofetal fibronectin indicating molecule alone or the binding partner alone. Detection of a mass corresponding to such a complex can indicate the presence of an oncofetal fibronectin indicating molecule in a sample. When a binding partner contains a moiety that can generate a detectable signal, presence of a compound having a molecular weight corresponding to an oncofetal fibronectin indicating molecule-binding partner complex, together with presence of the detectable signal from such a compound, can indicate the presence of an oncofetal fibronectin indicating molecule in a sample.

[0764] A variety of methods for determining the mass of a complex are known in the art, including, electrophoretic and chromatographic methods, such as gel electrophoresis (under conditions that do not disrupt the complex) and size exclusion chromatography and mass spectrometry. In one embodiment, a sample can be contacted with a fibronectin or oncofetal fibronectin binding partner conjugated with a fluorescent dye and then loaded onto an analytical gel filtration column. Elution can be monitored in terms of volume and fluorescence, where volume eluted is calibrated to molecular weight. A fluorescence signal at a volume corresponding to a molecular weight of an oncofetal fibronectin indicating molecule-binding partner complex can indicate the presence of an oncofetal fibronectin indicating molecule in a sample.

[0765] b. Immobilized Sample

[0766] A sample suspected of containing an oncofetal fibronectin indicating molecule can be examined for presence of the oncofetal fibronectin indicating molecule by immobilizing the sample on a solid support and probing the immobilized sample using a fibronectin or oncofetal fibronectin binding partner, where detection of binding of the binding partner to the solid support is indicative of the presence of an oncofetal fibronectin indicating molecule in the sample. Any oncofetal fibronectin indicating molecule, such as an oncofetal fibronectin protein, an oncofetal fibronectin-encoding nucleic acid molecule or a complement thereto, or an autoantibody to oncofetal fibronectin protein or a nucleic acid molecule encoding oncofetal fibronectin, can be immobilized to a solid support in the methods provided herein. A binding partner can be detected directly or indirectly. Direct detection of the binding partner can be performed using a binding partner conjugated to a detectable moiety or conjugated to a bindable moiety. Indirect detection of a fibronectin or an oncofetal fibronectin binding partner can be performed using a binding partner that can bind to the fibronectin or oncofetal fibronectin binding partner and can generate a detectable signal or can be bound by another binding partner. Competition assays using labeled and unlabeled (or differently labeled) binding partners also can be performed.

[0767] Solid supports used in the present methods can be any solid support to which, upon contact with a sample, an oncofetal fibronectin indicating molecule can be immobilized. Exemplary solid supports include microplates, microarrays, or membranes such as nitrocellulose, polyvinylidene fluoride (PVDF) or nylon membranes. Methods for immobilizing a sample on solid supports are known in the art and can be used in the methods herein. When sample is immobilized in two or more discrete locations, such as in a dot blot, a microplate or microarray, each sample can be

independently treated, where all samples can be differently treated, or some samples can be equally treated while others are differently treated. Exemplary treatments include contacting with different binding partners, including binding partners that bind to different regions of an oncofetal fibronectin indicating molecule such as EDA, EDB and IICS, contacting under different buffer conditions, and contacting with different concentrations of binding partner. Such methods are known to those skilled in the art.

[0768] Detection of a signal also can be used to quantitate the amount of an oncofetal fibronectin indicating molecule present in the sample, using any of a variety of known methods. For example, the intensity of the signal corresponding to an oncofetal fibronectin indicating molecule can be measured using any of a variety of methods known in the art, including, for example, fluorescence or absorption spectrometry, or phosphor imager measurement. In one example, known concentrations of a standard also can be included and the intensity of one or more sample signals can be qualitatively or quantitatively compared to the standards using known methods to provide an estimate or calculation of the amount of an oncofetal fibronectin indicating molecule in the sample. In another example, multiple dilutions of the sample can be detected, and the signal measured at each dilution can be used to estimate the amount of an oncofetal fibronectin indicating molecule using known methods.

[0769] i. Dot Blot Analysis

[0770] Dot blot assays can be used to detect the presence of an oncofetal fibronectin indicating molecule in a sample. Dot blot can be performed by first preparing a solid support, typically a membrane such as a nitrocellulose membrane or a PVDF membrane, such as by wetting the membrane in a solvent such as methanol or distilled water. In other examples, no preparation of the membrane is required. A sample can then be added to the membrane. In a dot blot, one or more aliquots of sample can be added at discrete loci on a membrane to form one or more "dots" on the membrane. When multiple sample aliquots are used, the aliquots can be identical or different, where the different aliquots can be at different levels of sample dilution or can be aliquots which have been treated with at least one different reagent. For example, different aliquots can be contacted with different non-specific binding compounds or surfaces; or one or more first aliquots can be untreated and one or more second aliquots can be treated, for example, using a non-specific binding compound. Control or reference samples also can be added at one or more discrete loci on the membrane. In one embodiment, samples applied to a membrane can be drawn through the membrane by a suction pump, or other similar device. The oncofetal fibronectin can be immobilized onto the membrane by any known methods, including, for example, drying the membrane or exposing the membrane to ultraviolet radiation. Such a membrane also can be washed to remove any substances that were not immobilized to the membrane. Wash solutions are known in the art and can contain detergent, such as a nonionic detergent such as Tween-20.

[0771] Membranes to which sample has been added can be contacted with a fibronectin or oncofetal fibronectin binding partner. In one embodiment, after adding the sample, but prior to adding a fibronectin or oncofetal fibronectin binding partner, the membrane can be washed with a solution that

suppresses additional non-specific binding to the membrane. For example, in the case of protein (including autoantibody) detection, a membrane can be washed with a solution containing, for example, BSA or casein, which can bind to all remaining surfaces of the membrane that can non-specifically bind to proteins. In the case of nucleic acid detection, a membrane can be washed with a solution containing, for example, salmon sperm DNA, which can bind to all remaining surfaces of the membrane that can non-specifically bind to nucleic acids. After washing with a solution that suppresses non-specific binding, a fibronectin or oncofetal fibronectin binding partner can be added. The presence of the fibronectin or oncofetal fibronectin binding partner on the membrane can be detected directly or indirectly. In one example, the binding partner can be conjugated to a moiety that can produce a directly detectable signal.

[0772] If the fibronectin or oncofetal fibronectin binding partner is not conjugated with a detectable moiety, the fibronectin or oncofetal fibronectin binding partner can be detected indirectly using a binding partner that binds to a fibronectin or oncofetal fibronectin binding partner, where the binding partner that binds to a fibronectin or oncofetal fibronectin binding partner can be conjugated with a moiety that can produce a detectable signal, or can itself be bound by a binding partner. For example, the fibronectin or oncofetal fibronectin binding partner can be a mouse monoclonal anti-oncofetal fibronectin antibody and the binding partner that binds to the mouse monoclonal anti-oncofetal fibronectin antibody can be a horseradish peroxidase-conjugated goat anti-mouse IgG antibody.

[0773] Moieties that can produce a detectable signal include radionuclides, fluorescent molecules, quantum dots, colloidal metal and proteins including green fluorescence protein, horseradish peroxidase and chloramphenicol acetyl transferase. Moieties can be detected directly, such as by detecting fluorescence of a fluorescent molecule. Moieties also can be detected indirectly such as by detecting chemiluminescence catalyzed by horseradish peroxidase. Detection of a moiety at a discrete location on the membrane can indicate presence of an oncofetal fibronectin indicating molecule in the sample that was applied to that discrete location. In one embodiment, the amount of an oncofetal fibronectin indicating molecule present in the sample can be quantitated using known methods, for example, using multiple dilutions of a sample at two or more dot loci and detecting binding partner(s) present at each locus.

[0774] ii. Western Blot Analysis

[0775] Western blot analysis can be used to detect the presence of oncofetal fibronectin protein or autoantibody in a sample. Western blotting is performed by first loading a protein sample in a gel, such as a sodium dodecyl sulfate polyacrylamide gel and electrophoresing the sample, using known methods. One sample can be loaded on each lane of the gel, which can contain a plurality of lanes. In one embodiment, a gel can contain multiple sample aliquots. When multiple sample aliquots are used, the aliquots can be identical or different, where the different aliquots can be at different levels of sample dilution or can be aliquots which have been treated with at least one different reagent. For example, different aliquots can be contacted with different non-specific binding compounds or surfaces; or one or more first aliquots can be untreated and one or more second aliquots can be treated, for example, using a non-specific binding compound. Control or reference samples also can be added to the gel. The gel can then be electrophoresed according to known methods.

aliquots can be treated, for example, using a non-specific binding compound. Control or reference samples also can be added to the gel. The gel can then be electrophoresed according to known methods.

[0776] The electrophoresed gel can then be placed onto a prepared membrane (as known in the art and described in regard to dot blot analysis) and the protein of the gel can be electroblotted onto the membrane using known methods. The membrane can then be washed, blocked and a fibronectin or oncofetal fibronectin binding partner can be added as described in regard to dot blot analysis. As with dot blot analysis, the fibronectin or oncofetal fibronectin binding partner can be detected directly, for example, using a fibronectin or oncofetal fibronectin binding partner conjugated with a moiety that can produce a detectable signal, or indirectly, for example, using a binding partner conjugate that binds the fibronectin or oncofetal fibronectin binding partner.

[0777] Detection of a signal from a moiety in one lane of the membrane can indicate the presence of oncofetal fibronectin protein or autoantibody in the sample loaded in that lane. Further, the location of the signal in the lane can be used to determine the molecular weight of the sample component to which the fibronectin or oncofetal fibronectin binding partner bound. This information can be used, for example, to eliminate false positive signals if a detected signal does not correspond to an expected molecular weight of oncofetal fibronectin protein or autoantibody (or a fragment thereof). This information also can be used, for example, to characterize the type of oncofetal fibronectin protein variant present, for example, to distinguish between a V0, V64, V89, V95 or V120 splice variant. In one embodiment, the amount of oncofetal fibronectin protein or autoantibody present in the sample can be quantitated using known methods.

[0778] iii. Southern and Northern Blot Analyses

[0779] Southern and Northern blot analysis can be used to detect the presence of a nucleic acid encoding oncofetal fibronectin or complementary thereto in a sample. A nucleic acid encoding oncofetal fibronectin can include mRNA encoding oncofetal fibronectin or amplicates thereof (e.g., cDNA) and fragments thereof. Southern and Northern blot analysis are performed by first loading a DNA (Southern) or RNA (Northern) sample in a gel, such as an agarose gel or polyacrylamide gel, and electrophoresing the sample under denaturing conditions, using known methods. One sample can be loaded on each lane of the gel, which can contain a plurality of lanes. In one embodiment, a gel can contain multiple sample aliquots. When multiple sample aliquots are used, the aliquots can be identical or different, where the different aliquots can be at different levels of sample dilution or can be aliquots which have been treated with at least one different reagent. For example, different aliquots can be contacted with different non-specific binding compounds or surfaces; or one or more first aliquots can be untreated and one or more second aliquots can be treated, for example, using a non-specific binding compound. Control or reference samples also can be added to the gel. The gel can then be electrophoresed according to known methods.

[0780] The electrophoresed gel can then be placed onto a prepared membrane (as known in the art or described herein), such as a nylon membrane and the nucleic acid

molecules of the gel can be blotted onto the membrane using known methods, including electroblotting, vacuum blotting, semi-dry electroblotting, bidirection transfer and positive pressure, as known in the art.

[0781] The membrane can then be washed, blocked and a fibronectin or oncofetal fibronectin binding partner can be added as described above. As with dot blot analysis, the fibronectin or oncofetal fibronectin binding partner can be conjugated with a moiety that can produce a detectable signal, or a binding partner conjugate that binds the fibronectin or oncofetal fibronectin binding partner can be used.

[0782] Detection of a signal from a moiety in one lane of the membrane can indicate the presence of an oncofetal fibronectin encoding nucleic acid or complement thereto or fragment thereof in the sample loaded in that lane. Further, the location of the signal in the lane can be used to determine the molecular weight of the sample component to which the fibronectin or oncofetal fibronectin binding partner bound. This information can be used, for example, to eliminate false positive signals if a detected signal does not correspond to an expected molecular weight of an oncofetal fibronectin encoding nucleic acid or complement thereto (or a fragment thereof). This information also can be used, for example, to characterize the oncofetal fibronectin as containing or not containing EDA, EDB and/or the splice variant of IIICS or complement thereto.

[0783] Detection of a signal also can be used to quantitate the amount of oncofetal fibronectin encoding nucleic acid molecule or complement thereto present in the sample. For example, the intensity of the band corresponding to oncofetal fibronectin can be measured using any of a variety of methods known in the art, including, for example, phosphor imager measurement. In one example, known concentrations of a standard also can be included in the blot and the intensity of one or more sample bands can be qualitatively or quantitatively compared to the standards using known methods to provide an estimate or calculation of the amount of oncofetal fibronectin encoding nucleic acid molecule or complement thereto present in the sample.

[0784] iv. In Situ Analysis

[0785] Presence of an oncofetal fibronectin indicating molecule can be determined in situ. An oncofetal fibronectin indicating molecule can be detected in tissue or organ samples, or can be detected in a subject. For example, a fibronectin or oncofetal fibronectin binding partner conjugated to a fluorophore can be applied to a tissue sample and detection of fluorescence on the tissue sample can indicate the presence of an oncofetal fibronectin indicating molecule in the tissue. In another example, a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging agent can be administered to a subject and detection of a localized imaging agent signal in the subject can indicate the presence and location of an oncofetal fibronectin indicating molecule in the subject. In another example, a fibronectin or oncofetal fibronectin binding partner, optionally conjugated to a therapeutic agent can be administered to a subject in treating the subject.

[0786] Methods of administering a fibronectin or oncofetal fibronectin binding partner conjugate to a subject for detecting an oncofetal fibronectin indicating molecule in a

subject also can be used for treatment of the subject. For example, a detectable fetal fibronectin or oncofetal fibronectin binding partner conjugate, such as a radionuclide-containing conjugate, can be detected by known methods and also can be used for treating the subject. Conjugates that can be used for both detection and therapeutic methods are known in the art.

[0787] In one embodiment, anti-fibronectin or anti-oncofetal fibronectin autoantibodies can be used according to the use of the fibronectin or oncofetal fibronectin binding partners. For example, anti-fibronectin or anti-oncofetal fibronectin autoantibodies can be removed from a subject and have conjugated thereto a detectable and/or therapeutic agent, and the autoantibody can then be administered back to the subject, where the autoantibody conjugate can be detectable, therapeutic, or both.

[0788] a. Tissue or Organ Samples

[0789] Presence of an oncofetal fibronectin indicating molecule can be determined in situ in a tissue or organ sample. A tissue or organ sample includes biological matter removed from a subject, where typically the biological matter contains cells or is solid. An exemplary tissue or organ sample is a biopsy or a paraffin-embedded tissue sample. In one embodiment, a fibronectin or oncofetal fibronectin binding partner conjugate can be applied to a tissue sample and detection of the conjugate bound to the tissue sample can indicate the presence of an oncofetal fibronectin indicating molecule in the tissue. A variety of conjugates useful for such methods are provided herein or known in the art and include conjugates containing fluorescent moieties, radionuclides, chromophores, latex microspheres, quantum dots, colloidal metal, or an enzyme capable of producing a detectable signal such as horseradish peroxidase or luciferase. In another embodiment, a fibronectin or oncofetal fibronectin binding partner can be applied to a tissue sample and detection of the fibronectin or oncofetal fibronectin binding partner bound to the tissue sample can indicate the presence of an oncofetal fibronectin indicating molecule in the tissue. Presence of the fibronectin or oncofetal fibronectin binding partner can be detected using any of a variety of methods, including contacting the tissue or organ sample with a detectable compound that specifically binds a fibronectin or oncofetal fibronectin binding partner, including, but not limited to, an antibody or an oligonucleotide. For example, when the sample is contacted with an anti-oncofetal fibronectin mouse monoclonal antibody, the sample can be subsequently contacted with a goat anti-mouse IgG antibody conjugated to, for example, an enzyme such as luciferase and presence of a luminescent signal can indicate the presence of anti-oncofetal fibronectin mouse monoclonal antibody bound to the sample and thereby indicate the presence of an oncofetal fibronectin indicating molecule in the sample.

[0790] Any of a variety of known histochemical methods can be used for determining the presence of an oncofetal fibronectin indicating molecule in a tissue or organ sample, according to the sample to be used, the oncofetal fibronectin indicating molecule to be detected, and the binding partner and binding partner detection method to be used. Generally, such techniques typically include steps of sample preparation or fixation, any post-fixation and/or antigen retrieval steps appropriate according to the sample preparation and an

oncofetal fibronectin indicating molecule detection methods, blocking non-specific binding and other false positive blocking steps such as endogenous peroxidase blocking, hybridization with a fibronectin or oncofetal fibronectin binding partner, washing and detection of the fibronectin or oncofetal fibronectin binding partner bound to the sample either directly or indirectly through a secondary compound. Any method known in the art for tissue or organ sample preparation can be used, including, but not limited to, fixing the sample with formaldehyde and embedding the sample in paraffin, fixing the sample in acetone and embedding the sample in paraffin and fixing the sample in paraformaldehyde and flash freezing the sample in liquid nitrogen. An oncofetal fibronectin indicating molecule detection methods can be performed according to any of a variety of well known methods, according to the oncofetal fibronectin indicating molecule to be detected. In one example, when an oncofetal fibronectin protein or fragment thereof, is to be detected, standard immunohistochemical methods employing a fibronectin or oncofetal fibronectin binding partner such as an anti-oncofetal fibronectin antibody such as FDC-6 or BC-1 can be performed. In another example, when an oncofetal fibronectin encoding nucleic acid molecule or fragment thereof, is to be detected, standard nucleotide in situ hybridization methods (e.g., FISH), employing an oligonucleotide complementary to an oncofetal fibronectin encoding nucleic acid molecule can be performed. Various known imaging methods can be used to detect the presence of an oncofetal fibronectin indicating molecule in the sample. In one example, confocal microscopy can be used to detect the presence of an oncofetal fibronectin indicating molecule in the sample.

[0791] The fibronectin or oncofetal fibronectin binding partner (e.g., antibody or oligonucleotide probe) can be selected according to the characteristics of the oncofetal fibronectin indicating molecule to be determined. For example, if presence of a IIICS-containing oncofetal fibronectin is to be determined, an antibody such as FDC-6 or X18A4 can be used, or an oligonucleotide probe complementary to at least a portion of the IIICS region can be used. Similar selections of oncofetal fibronectin binding partners can be used to determine the presence of EDA or EDB containing an oncofetal fibronectin indicating molecule in a sample and also to determine the presence of various splice variants of IIICS in an oncofetal fibronectin indicating molecule present in the sample. Oncofetal fibronectin binding partners also can bind oncofetal fibronectin protein or autoantibody, to indicate post-translational modifications of oncofetal fibronectin protein in the subject. Combinations of two or more oncofetal fibronectin binding partners that can, for example, detect the presence of two or more different regions of an oncofetal fibronectin indicating molecule can be used. In one example, multiple oligonucleotides can be used for detecting a nucleic acid molecule encoding oncofetal fibronectin. For example, five oligonucleotide probes can be used to determine the presence of oncofetal fibronectin containing EDA, oncofetal fibronectin containing EDB, oncofetal fibronectin containing the first, 25 amino acid splice region of IIICS, oncofetal fibronectin containing the second, 64 amino acid splice region of IIICS and oncofetal fibronectin containing the third, 31 amino acid splice region of IIICS; alternatively, 10 amplification primers can be used to determine the presence of these 5 regions using standard nucleic acid amplification methods. Oncofetal fibronectin

binding partners disclosed herein or known in the art can be used alone or in combinations, for example, of 2 or more oncofetal fibronectin binding partners, 3 or more oncofetal fibronectin binding partners, 4 or more oncofetal fibronectin binding partners, or 5 or more oncofetal fibronectin binding partners. In one embodiment, each different oncofetal fibronectin binding partner is differently detectable; for example, each different oncofetal fibronectin binding partner can contain a fluorescent moiety that fluoresces at peak wavelength different from the other fluorescent moieties.

[0792] A variety of methods for detecting an oncofetal fibronectin indicating molecule in a tissue sample are known in the art. For example, antibodies can be used for detecting oncofetal fibronectin protein in a tissue sample. Among antibodies that can be used is FDC-6, which can be used for detecting a variety of cancers (neoplastic diseases) in a subject, including carcinomas of the lung, breast, ovary, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, prostate, thyroid, pituitary, eye, brain, oral, skin, head and neck cancer, lymphoma, leukemia, squamous cell carcinomas, adenocarcinomas, small cell carcinomas, melanomas, gliomas, sarcomas and neuroblastomas. In one example, FDC-6 can be used to detect glycosylated IIICS-containing oncofetal fibronectin protein in cancerous tissue samples such as breast cancer (Kaczmarek et al., *Int. J. Cancer* 59:11-16 (1994)). Other antibodies that bind oncofetal fibronectin protein also can be used. For example, BC-1 can be used to detect the presence of an EDB-containing oncofetal fibronectin protein in cancerous tissue samples such as brain meningiomas and lung carcinoma (Carnemolla et al., *J. Cell. Biol.* 108:1139-1148 (1989)), L19 can be used to detect the presence of EDB-containing oncofetal fibronectin protein in tumor tissue in mice (Borsi et al., *Int. J. Cancer* 102:75-85 (2002)), 5C10 can be used to detect glycosylated IIICS-containing oncofetal fibronectin protein in cancerous tissue samples such as oral squamous cell carcinoma (Lyons et al., *Br. J. Oral Maxillofac. Surg.* 39:471-477 (2001)), X18A4 can be used to detect the presence of IIICS-containing oncofetal fibronectin protein in cancerous tissue samples such as ovarian cancer (Menzin et al., *Cancer* 82:152-158 (1998)) and IST-9 can be used to detect the presence of EDA-containing oncofetal fibronectin protein in cancerous tissue samples such as papillary carcinoma of the thyroid (Scarpino et al., *J. Pathol.* 188:163-167 (1999)). In another example, IST-9 and BC-1 can be used to detect the presence of EDA and EDB in oncofetal fibronectin protein, respectively, in hepatocellular carcinoma (Oyama et al., *Cancer Res.* 53:2005-2011 (1993)). Additional combinations of particular neoplastic diseases and oncofetal fibronectin protein binding partners can be determined empirically.

[0793] Nucleic acid probes also can be used for in situ detection of a nucleic acid encoding oncofetal fibronectin. For example, a nucleic acid probe can be used to detect the presence of mRNA encoding IIICS-containing oncofetal fibronectin in cancerous tissue samples such as thyroid papillary and anaplastic carcinoma (Takano et al., *Br. J. Cancer* 78:221-224 (1998)). In another example, a nucleic acid probe can be used to detect the presence of mRNA encoding EDB-containing oncofetal fibronectin in cancerous tissue such as oral squamous cell carcinoma (Kosmehl et al., *Br. J. Cancer* 81:1071-1079 (1999)). Additional combinations of particular neoplastic diseases and oncofetal

fibronectin encoding nucleic acid molecule binding partners can be determined empirically.

[0794] b. Detection in a Subject

[0795] Presence of an oncofetal fibronectin indicating molecule can be determined in vivo in a subject. A variety of tumorous tissues, such as cervical tumor tissue (as provided herein), express oncofetal fibronectin protein. In addition, as provided herein, cells or tissues expressing oncofetal fibronectin or expressing increased amounts of oncofetal fibronectin protein relative to normal cells or tissues can be developing into neoplastic cells or tissues or can have an increased likelihood of developing into neoplastic cells or tissues. Accordingly, by administering to a subject a fibronectin or oncofetal fibronectin binding partner or conjugate thereof, tumorous tissues or tissues with increased likelihood of developing into tumorous tissues can be specifically targeted. In accordance with this embodiment, a fibronectin or oncofetal fibronectin binding partner conjugate can be administered to a subject and detection of the conjugate at a location in the subject can indicate the presence of oncofetal fibronectin expression in that location. For example, tumorous tissue in a subject can be imaged by administering to a subject a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety, whereby the conjugate localizes to an oncofetal fibronectin indicating molecule at or near tumorous tissue and then detecting the localization of the conjugate within the subject to thereby image tumorous tissue in the subject. In another example, tissue developing or likely to develop into tumorous tissue can be imaged by administering to a subject a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety, whereby the conjugate localizes to an oncofetal fibronectin indicating molecule at or near tissue developing into or likely to develop into tumorous tissue and then detecting the localization of the conjugate within the subject to thereby image tissue developing or likely to develop into tumorous tissue in the subject. In such cases, the conjugate can localize to the tumorous or pre-tumorous tissue itself, or to extracellular matrix or vasculature adjacent the tumorous or pre-tumorous tissue. Further, a fibronectin or oncofetal fibronectin binding partner conjugate can be administered to a subject and detection of the conjugate at a location in the subject can indicate the presence of cancerous (tumorous or neoplastic) cells, or cells with increased likelihood of developing into cancerous cells in that location. For example, cancerous (tumorous or neoplastic) cells in a subject can be imaged by administering to a subject a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety, whereby the conjugate localizes to cancerous (tumorous or neoplastic) cells containing an oncofetal fibronectin indicating molecule and then detecting the localization of the conjugate within the subject to thereby image cancerous (tumorous or neoplastic) cells in the subject. In another example, cells developing into or having an increased likelihood of developing into cancerous (tumorous or neoplastic) cells can be imaged by administering to a subject a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety, whereby the conjugate localizes to cells developing into or with increased likelihood of developing into cancerous (tumorous or neoplastic) cells containing an oncofetal fibronectin indicating molecule and then detecting the localization of the conjugate within the subject to thereby image cells developing into or with increased likelihood of devel-

oping into cancerous (tumorous or neoplastic) cells in the subject. Methods also are provided herein for detecting an oncofetal fibronectin indicating molecule in a subject, by administering to a subject a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety, whereby the conjugate localizes to regions in the subject containing an oncofetal fibronectin indicating molecule and detecting the localization of the conjugate within the subject, thereby detecting an oncofetal fibronectin indicating molecule in the subject.

[0796] A variety of non-invasive imaging methods are known in the art, including magnetic resonance imaging or other resonance methods, ultrasonic imaging, fluorescence imaging, scintigraphy, or tomography methods such as computed tomography, computerized axial tomography, positron emission tomography, single photon emission computed tomography, ultrasound tomography or x-ray tomography. For example, BC-1 labeled with technetium(Tc)-99m can be used to indicate the presence of EDB containing oncofetal fibronectin protein in brain tumors in a subject (Calcagno et al., *Cancer* 80:2484-2489 (1997)) and ¹²³I-labeled-L19 can be used to indicate the presence of EDB containing oncofetal fibronectin protein in lung cancer or colorectal cancer in a subject (Santimaria et al., *Clin. Cancer Res.* 9:571-579 (2003)). In another example, L19 labeled with the fluorophore Cy5 can be used for in vivo fluorescence indication of the presence of EDB containing oncofetal fibronectin protein in angiogenesis in the eye (U.S. Pat. No. 20030045681).

[0797] Compounds that can be used in the imaging methods provided herein or known in the art typically contain a fibronectin or oncofetal fibronectin binding partner conjugated to an imaging moiety. For example, FDC-6 can be conjugated to ¹⁸F, which can be detected by positron emission tomography. Any of a variety of fibronectin or oncofetal fibronectin binding partners disclosed herein or otherwise known in the art can be used in the treatment methods provided herein, including fibronectin or oncofetal fibronectin binding partner proteins, fibronectin or oncofetal fibronectin binding partner nucleic acid molecules and fibronectin or oncofetal fibronectin binding partner organic molecules. The fibronectin or oncofetal fibronectin binding partner also can be a binding partner that specifically binds the EDA, EDB, IIICS regions of oncofetal fibronectin, or a particular splice variant of IIICS such as V64, V89, V95 or V120, or a particular post-translational modification of oncofetal fibronectin protein such as O-glycosylation of threonine 33 of IIICS, or combinations thereof. Typically the fibronectin or oncofetal fibronectin binding partner will be a binding partner that can be present in a subject for a length of time sufficient to localize to one or more sites where oncofetal fibronectin is present within the subject (e.g., the prostate, lungs, brain, breast, ovary, thyroid, cervix or bladder) and sufficient to be imaged by one or more of the imaging methods provided herein or known in the art. The fibronectin or oncofetal fibronectin binding partner will also typically preferentially bind to an oncofetal fibronectin indicating molecule such that imaging methods can determine the location of an oncofetal fibronectin indicating molecule within the subject. An exemplary oncofetal fibronectin binding partner that can be used for in vivo imaging is an antibody such as FDC-6, BC-1, ME4C or L19.

[0798] In one embodiment, the anti-oncofetal fibronectin antibody can be a “humanized” or chimeric antibody bearing human constant and/or variable domains, or can be an antibody otherwise derived from a human antibody source or human antibody sequence, such as L19. Methods of deriving antibodies from a human antibody source or human antibody sequence using, for example, phage display or filter selection of antibody-expressing bacteria are known in the art, as exemplified in WO 97/45544 and WO 02/46455. Methods for deriving antibodies from a human antibody sequence using, for example transgenic animals expressing human antibody sequences are known in the art, as exemplified in U.S. Pat. Nos. 6,632,976 and 6,713,610.

[0799] The conjugate used for in vivo detection methods typically contains a moiety that permits imaging. A variety of imaging moieties are known in the art, including fluorescent moieties, radionuclides, magnetically detectable isotopes or compounds, sonographic imaging agents, chromophores, latex microspheres, or quantum dots. The moiety can be selected according to the imaging method to be used. For example, a variety of ultrasound imaging agents are known in the art, including, but not limited to, microspheres containing perfluorocarbon such as perfluorocarbon-filled albumin microspheres (see, e.g., U.S. Pat. No. 6,174,287) or phospholipid coated microbubbles filled with perfluorocarbon gas (see, e.g., U.S. Pat. No. 6,146,657) and also including other imaging agents such as simethicone-coated cellulose (see, e.g., U.S. Pat. No. 6,024,939) or galactose and palmitic acid microparticles (see, e.g., U.S. Pat. No. 5,380,411). In another example, a variety of computed tomography imaging agents are known in the art, including, but not limited to, compounds containing ^{131}I , ^{18}F , ^{123}I , ^{201}Tl , ^{111}In and Tc-99m. In another example, a variety of magnetic resonance imaging agents are known in the art, including, but not limited to, metal complexes such as iron, manganese or gadolinium-containing complexes, or other compounds containing atoms that are low in natural abundance and have a nuclear magnetic moment, for example, compounds containing ^{13}C , ^{15}N , ^{17}O , ^{19}F , ^{29}Si , or ^{31}P . In another example, any of a variety of luminescent or fluorescent compounds can be used including bioluminescent enzymes (e.g., luciferase), fluorescent proteins (e.g., green fluorescence protein), fluorophores, dyes, latex microspheres and quantum dots. In another example, any of a variety of x-ray contrast agents can be used, including, but not limited to, barium compounds such as barium sulfate and iodinated compounds including ionic compounds such as iodamide, iodipamide and ioglycamic acid and non-ionic iodinated compounds such as metrizamide, iopamidol, iohexol, iopromide, iobitridol, iomeprol, iopentol, ioversol, ioxilan, iodixanol, iotrolan.

[0800] A fibronectin or oncofetal fibronectin binding partner conjugated to a detectable moiety can be administered to a subject in any of a variety of manners known in the art for delivering compounds to a subject, according a variety of factors known to those skilled in the art, including, but not limited to, the nature of the fibronectin or oncofetal fibronectin binding partner and the detectable moiety, the regions of the subject's body to be imaged, the selected speed of administration and the likelihood of unintended clearance. In one embodiment, the conjugate is administered intravenously to a subject. In another embodiment, the conjugate is administered topically, for example, as a component of a lavage composition, or as a cream, salve or gel. The conjugate also can be administered topically to a variety of

additional surfaces including, but not limited to, dermal, oral, aural, nasal, anal, urethral, ocular, breast, cervicovaginal, alimentary canal such as esophageal, gastric, intestinal, or colon surfaces. Exemplary topical applications include contacting the vaginal and uterine cavities with a conjugate, or contacting one or more breast ducts with a conjugate.

[0801] The methods provided herein also can include imaging cells or tissues containing a particular oncofetal fibronectin indicating molecule variant. For example, imaging can be performed using an oncofetal fibronectin binding partner that specifically binds the EDA, EDB, IIICS regions of an oncofetal fibronectin indicating molecule, or a particular splice variant of IIICS such as V64, V89, V95 or V120, or a particular post-translational modification of oncofetal fibronectin protein such as O-glycosylation of threonine 33 of IIICS, or combinations thereof. Accordingly cells or tissues can be specifically imaged according to whether or not the cells or tissues contain an oncofetal fibronectin indicating molecule containing EDA, EDB, or IIICS, or a particular splice variant of IIICS such as V64, V89, V95 or V120, or a particular post-translational modification of oncofetal fibronectin protein such as O-glycosylation of threonine 33 of IIICS, or combinations thereof.

[0802] c. Treatment in a Subject

[0803] Presence of an oncofetal fibronectin indicating molecule also can be used for in vivo treatment of a subject. A variety of tumorous tissues are known to express, shed and/or secrete oncofetal fibronectin. Accordingly, by administering to a subject a fibronectin or oncofetal fibronectin binding partner or conjugate thereof, cancerous (tumorous) tissues can be specifically targeted. As provided herein, tissues developing into cancerous tissues or with increased risk of developing into cancerous tissues express oncofetal fibronectin. Accordingly, by administering to a subject a fibronectin or oncofetal fibronectin binding partner or conjugate thereof, tissues developing into cancerous tissues or with increased risk of developing into cancerous tissues can be specifically targeted. In accordance with this embodiment, a fibronectin or oncofetal fibronectin binding partner or a fibronectin or oncofetal fibronectin binding partner conjugate can be administered to a subject and the binding partner or conjugate can accumulate in a location in the subject that has an oncofetal fibronectin indicating molecule, such as a location at or near tumorous tissue or tissue developing into neoplastic tissue or with increased risk of developing into cancerous tissue. For example, the conjugate can localize to the tumorous tissue itself, or to extracellular matrix or vasculature adjacent the tumorous tissue. In another example, the conjugate can localize to tissue developing into cancerous tissues or with increased risk of developing into cancerous tissues itself, or to extracellular matrix or vasculature adjacent the tissue developing into cancerous tissues or with increased risk of developing into cancerous tissues. Further, a fibronectin or oncofetal fibronectin binding partner or conjugate thereof can be administered to a subject and the binding partner or conjugate can accumulate in a location in the subject that has cancerous cells. Additionally, a fibronectin or oncofetal fibronectin binding partner or conjugate thereof can be administered to a subject and the binding partner or conjugate can accumulate in a location in the subject that has cells developing into cancerous cells or with increased risk of developing into cancerous cells. The localized binding part-

ner or conjugate can then, directly or indirectly, inhibit cell growth in that location or cause cell death in that location or inhibit development of cells into cancerous cells in that location.

[0804] Cell growth can be inhibited by any of a variety of methods, including but not limited to, reducing the amount of neovascularization in a location, or down-regulating cell growth-promoting cellular processes. Cell death can be caused by any of a variety of methods including increasing a subject's immune response, generating apoptosis signal or otherwise initiating apoptosis, or applying a toxin or toxic substance such as diphtheria toxin or a radionuclide or chemotherapeutic substance.

[0805] A fibronectin or oncofetal fibronectin binding partner alone can cause tumor inhibition or cell death response or a fibronectin or oncofetal fibronectin binding partner conjugate with a moiety that inhibits cell growth or promotes cell death can cause tumor inhibition or cell death response. Any of a variety of fibronectin or oncofetal fibronectin binding partners disclosed herein or otherwise known in the art can be used in the treatment methods provided herein, including fibronectin or oncofetal fibronectin binding partners that are proteins, fibronectin or oncofetal fibronectin binding partners that are nucleic acid molecules and fibronectin or oncofetal fibronectin binding partners that are organic molecules. The oncofetal fibronectin binding partner also can be a binding partner that specifically binds the EDA, EDB, IIICS regions of an oncofetal fibronectin indicating molecule, or a particular splice variant of IIICS such as V64, V89, V95 or V120, or a particular post-translational modification of oncofetal fibronectin protein such as O-glycosylation of threonine 33 of IIICS, or combinations thereof. In one example, the oncofetal fibronectin binding partner can be an anti-oncofetal fibronectin antibody and the anti-oncofetal fibronectin antibody can bind to oncofetal fibronectin protein located at or near the targeted cell or the targeted tissue and elicit an immune response by the subject's immune system which can result in death of at least a portion of the cells in the location to which the fibronectin or oncofetal fibronectin binding partner localizes. In another example, a fibronectin or oncofetal fibronectin binding partner can bind an oncofetal fibronectin indicating molecule and inhibit one or more activities of the oncofetal fibronectin indicating molecule, resulting in inhibition of cell proliferation. For example, a fibronectin or oncofetal fibronectin binding partner that binds an oncofetal fibronectin encoding nucleic acid molecule can inhibit expression of oncofetal fibronectin, resulting in inhibition of cell proliferation. In another example, a fibronectin or oncofetal fibronectin binding partner that binds oncofetal fibronectin protein can alter, inhibit or modulate the binding properties of oncofetal fibronectin, resulting in inhibition of cell proliferation. For present purposes, a fibronectin or oncofetal fibronectin binding partner conjugated to a moiety that does not inhibit cell growth or cause cell death can have the same cell growth inhibitory or cell death causing effect as the fibronectin or oncofetal fibronectin binding partner alone. Typically the fibronectin or oncofetal fibronectin binding partner will be a binding partner that can be present in a subject for a length of time sufficient to localize to one or more sites where oncofetal fibronectin is present within the subject (e.g., the prostate, lungs, brain, breast, ovary, thyroid, colon or rectum, cervix or bladder) and sufficient to inhibit cell growth

or cause cell death, for example, by eliciting an immune response in the subject. An exemplary fibronectin or oncofetal fibronectin binding partner that can be used for in vivo treatment is an antibody such as FDC-6, BC-1, ME4C or L19. In one embodiment, the antibody can be a "humanized" or chimeric antibody bearing human constant and/or variable domains, or can be an antibody otherwise derived from a human antibody source or human antibody sequence, such as L19. Methods of deriving antibodies from a human antibody source using, for example, phage display or filter selection of antibody-expressing bacteria are known in the art. Methods for deriving antibodies from a human antibody sequence using, for example transgenic animals expressing human antibody sequences are known in the art.

[0806] The fibronectin or oncofetal fibronectin binding partner also can be administered as a conjugate. The conjugate typically contains a therapeutic moiety that inhibits cell growth or promotes cell death, that can be activated to inhibit cell growth or promote cell death, or that can activate a compound to inhibit cell growth or promote cell death. Optionally, the moiety also can have one or more additional properties such as acting as an imaging agent, as described elsewhere herein. A variety of therapeutic moieties are known in the art, including, but not limited to, biological toxins, cytokines, photosensitizing agents, toxins, anticancer antibiotics, chemotherapeutic compounds, radionuclides, binding partners and bioluminescent compounds. For example, a therapeutic moiety can be a biological toxin such as, but not limited to, psuedomonas exotoxin, diphtheria toxin, ricin, cholera toxin, gelonin, shigella toxin, pokeweed anti-viral protein, exotoxin A, abrin toxin or saporin (see, e.g., U.S. Pub. No. 2004/0009551). For example, a therapeutic moiety can be a signaling compound such as a cytokine or growth factor such as an interleukin including interleukin-1, interleukin-2, interleukin-6 and interleukin-12, a tumor necrosis factor such as tumor necrosis factor alpha (TNF- α), an interferon such as interferon gamma (IFN- γ), granulocyte macrophage colony stimulating factor (GM-CSF), angiogenin, or tissue factor (see, e.g., U.S. Pub. No. 20030232010). For example, a therapeutic moiety can be an anticancer antibiotic such as, but not limited to, porfirimycin, doxorubicin, dactinomycin, plicamycin, mitomycin, bleomycin, actinomycin, and daunorubicin (see, e.g., U.S. Pub. No. 20040054014). For example, a therapeutic moiety can be a photosensitizing agent such as, but not limited to, indocyanine green, toluidine blue, aminolevulinic acid, texaphyrins, benzoporphyrins, phenothiazines, phthalocyanines, porphyrins such as sodium porphyrin, chlorins such as tetra(m-hydroxyphenyl)chlorin or tin(IV) chlorin e6, purpurins such as tin ethyl etiopurpurin, purpurinimides, bacteriochlorins, pheophorbides, pyropheophorbides and cationic dyes (see, e.g., U.S. Pub. Nos. 20040019032 and 20030114434). For example, a therapeutic moiety can be a high energy radionuclide such as, but not limited to, a compound containing ³²Phosphate, ⁶⁰Cobalt, ⁹⁰Yttrium, ⁹⁹Technetium, ¹⁰³Palladium, ¹⁰⁶Ruthenium, ¹¹¹Indium, ¹¹⁷Lutetium, ¹²⁵Iodine, ¹³¹Iodine, ¹³⁷Cesium, ¹⁵³Samarium, ¹⁸⁶Rhenium, ¹⁸⁸Rhenium, ¹⁹²Iridium, ¹⁹⁸Gold, ²¹¹Astatine, ²¹²Bismuth and ²¹³Bismuth.

[0807] A therapeutic moiety can include a variety of other toxins including, but not limited to, any of a large number of chemotherapeutic compounds such as, but not limited to, 5-fluorouridine, calicheamicin and maytansine (see, e.g., U.S. Pub. No. 20020039557). Additional chemotherapeutic

compounds include alkylating agents such as thiotepa and cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolmelamine nitrogen mustards such as chiorambucil, chlomaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as acclinomycins, actinomycin, anthramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitiostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguanzone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; polysaccharide-K; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; cytosine arabinoside; cyclophosphamide; thiotepa; taxoids, e.g. paclitaxel and doxetaxel; chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone and toremifene (Fareston); and antiandrogens such as flutamide, nilutamide, bicalutamide, leuprolide and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Such chemotherapeutic compounds that can be used

herein include compounds whose toxicities preclude use of the compound in general systemic chemotherapeutic methods.

[0808] Therapeutic moieties also can be binding partners such as an antibody or fragment thereof, receptor or fragment thereof or ligand for an antibody or receptor, where such moieties can bind to any of a variety of substances or compounds for inhibiting cell growth or causing cell death. Binding partners can bind, for example, toxins, chemotherapeutic compounds, or compounds containing a radionuclide. Binding partners also can bind, for example, cell surface proteins such as a cell surface protein on a leukocyte or other cell related to an immune response. For example, a binding partner can bind to a lymphocyte such as a B cell, T cell or NK cell; an exemplary binding partner is an IgG that can bind to the Fc receptor of an NK cell. Therapeutic moieties also can activate a compound to inhibit cell growth or promote cell death. For example, a therapeutic moiety can be bioluminescent, such as luciferase and can, upon administration to the subject of an appropriate substrate such as luciferin and a photosensitizing agent, activate the photosensitizing agent in the area of localization of the conjugate to inhibit cell growth or cause cell death in that location.

[0809] Therapeutic moieties can be cleavable from the binding partner portion of the conjugate. A variety of cleavable linkages are known in the art including photocleavable linkages, chemically cleavable linkages, thermally cleavable linkages, enzymatically cleavable linkages. For example, the linkage can be through a disulfide bond, by a photocleavable biotin derivative, or protease-sensitive peptides (see, e.g., U.S. Pat. No. 6,416,758). In one embodiment a moiety attached to the binding partner via a cleavable linkage is a moiety that directly inhibits cell growth or causes cell death.

[0810] A fibronectin or oncofetal fibronectin binding partner or fibronectin or oncofetal fibronectin binding partner conjugate can be administered to a subject in any of a variety of manners known in the art for delivering compounds to a subject, according to a variety of factors known to those skilled in the art, including, but not limited to, the nature of the fibronectin or oncofetal fibronectin binding partner and the nature of the therapeutic moiety in a fibronectin or oncofetal fibronectin binding partner conjugate, the regions of the subject's body to be treated, the selected speed of administration and the likelihood of unintended clearance. In one embodiment, the binding partner or conjugate is administered intravenously to a subject. In another embodiment, the binding partner or conjugate is administered topically, for example, as a component of a lavage composition, or as a cream, salve or gel. The binding partner or conjugate also can be administered topically to a variety of additional surfaces including, but not limited to, dermal, oral, aural, nasal, anal, urethral, ocular, breast, cervicovaginal, alimentary canal such as esophageal, gastric, intestinal, or colon surfaces. Exemplary topical applications include contacting the vaginal and uterine cavities with a binding partner or conjugate, or contacting one or more breast ducts with a binding partner or conjugate.

[0811] The methods provided herein also can include treating cells or tissues containing a particular oncofetal fibronectin indicating molecule variant. For example, treatment methods can be performed using an oncofetal fibronectin binding partner that specifically binds the EDA, EDB,

IIICS regions of an oncofetal fibronectin indicating molecule, or a particular splice variant of IIICS such as V64, V89, V95 or V120, or a particular post-translational modification of oncofetal fibronectin protein such as O-glycosylation of threonine 33 of IIICS, or combinations thereof. Accordingly cells or tissues can be specifically treated by containing an oncofetal fibronectin indicating molecule containing EDA, EDB, or IIICS, or a particular splice variant of IIICS such as V64, V89, V95 or V120, or a particular post-translational modification of oncofetal fibronectin protein such as O-glycosylation of threonine 33 of IIICS, or combinations thereof.

[0812] The methods provided herein also can include method for inhibiting the recurrence of neoplastic disease in a subject. Such methods can include treating a subject for a neoplastic disease and administering to the subject a fibronectin or oncofetal fibronectin binding partner, whereby recurrence of neoplastic disease is inhibited. Treatment of the subject can be performed by any of a variety of methods known in the art, including chemotherapy, radiation therapy, administration of a bacteria or virus, administration of a tumor-specific compound and combinations thereof, including administration of a fibronectin or oncofetal fibronectin binding partner. In addition to the treatment for neoplastic disease, the subject can have administered thereto a fibronectin or oncofetal fibronectin binding partner which can serve to inhibit recurrence or metastasis of a neoplastic disease. Such administration can be concomitantly, subsequently, or intermittently or mixtures thereof with the treatment of the neoplastic disease.

[0813] c. Immobilized Binding Partner

[0814] Presence of an oncofetal fibronectin indicating molecule in a sample can be detected by contacting a sample with a solid support, to which a fibronectin binding partner or an oncofetal fibronectin binding partner is immobilized. When an oncofetal fibronectin indicating molecule is present in the sample, complex formation of the immobilized binding partner and oncofetal fibronectin indicating molecule can be detected. Detection of a complex between the immobilized binding partner and oncofetal fibronectin indicating molecule can indicate the presence of the oncofetal fibronectin indicating molecule in a sample. Complex detection can be achieved by any of a variety of methods known in the art. In one example, complex formation between a fibronectin or oncofetal fibronectin binding partner and an oncofetal fibronectin indicating molecule can be detected by detecting the presence of a second binding partner (either fibronectin or oncofetal fibronectin binding partner) in the complex, forming a "sandwich" complex. A binding partner can be detected directly or indirectly. Direct detection of the binding partner can be performed using a binding partner conjugated to a detectable moiety or conjugated to a bindable moiety. Indirect detection of a fibronectin or an oncofetal fibronectin binding partner can be performed using a binding partner that can bind to the fibronectin or oncofetal fibronectin binding partner and can generate a detectable signal or can be bound by another binding partner. In another example, complex formation between an oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner can be detected by competition assay. For example, a fibronectin or oncofetal fibronectin binding partner can be contacted with a labeled oncofetal fibronectin indicating molecule or analog thereof, and then

contacted with the sample; presence of oncofetal fibronectin indicating molecule in the sample will result in a decrease in the presence of labeled oncofetal fibronectin indicating molecule or analog thereof in complex with the binding partner. Thus, presence of an oncofetal fibronectin indicating molecule can be detected by disappearance of a signal in a competition assay.

[0815] A fibronectin binding partner or an oncofetal fibronectin binding partner can be immobilized on a solid support by known methods for use in the methods provided herein. A solid support on which a binding partner can be immobilized can be any of a variety of supports to which the binding partner can bind directly or can be bound via a linker or coating. Exemplary solid supports include microplates, microarrays, or membranes such as nitrocellulose, polyvinylidene fluoride (PVDF) or nylon membranes. For example, a binding partner can be immobilized on an untreated microplate or a treated microplate, including a microplate coated with a compound for binding the binding partner. Methods for immobilizing a sample on solid supports are known in the art and can be used in the methods herein.

[0816] A solid support containing a fibronectin or oncofetal fibronectin binding partner immobilized thereon can be subjected to one or more treatment steps prior to contacting the solid support with a sample. Such treatment steps include blocking steps to prohibit the surface of the solid support from non-specifically binding one or more components of a sample. Any of a variety of blocking steps known in the art can be applied to the solid support; also, a step of contacting the solid support with a non-specific binding compound such as a non-specific binding protein for oncofetal fibronectin protein assays, and a non-specific binding nucleic acid molecule for assays of a nucleic acid molecule encoding oncofetal fibronectin or complement thereto.

[0817] Prior to contacting the sample to the solid support containing fibronectin or oncofetal fibronectin binding partner, a sample can be treated by one or more steps as provided herein or otherwise known in the art. Exemplary treatment steps include filtering the sample to remove particulate matter, contacting the sample with a non-specific binding compound or surface to reduce background material binding to immobilized binding partner, adding a soluble or mobile fibronectin or oncofetal fibronectin binding partner or conjugate thereof to the sample solution, and addition of one or more buffers or reagents to modify ambient conditions such as pH or ionic strength.

[0818] Sample, treated or untreated, can be contacted with the solid support under conditions in which the fibronectin or oncofetal fibronectin binding partner can specifically bind to an oncofetal fibronectin indicating molecule. In one embodiment, the solid support is contacted with a sample under conditions in which background materials do not significantly interfere with the solid support binding oncofetal fibronectin (i.e., 10% or about 10% or less of the binding partners on the solid support bind to background materials or 90% or about 90% or more of an oncofetal fibronectin indicating molecule in the sample is bound by binding partners on the solid support). Exemplary conditions include 0.05 M Tris buffer, pH 7.4, 150 mM NaCl, 0.02% NaN₃, 1% BSA, 5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin, and 0.1% Triton X-100. The solid support can optionally be washed to

remove any background material that can be non-specifically binding to the solid support.

[0819] When a fibronectin or oncofetal fibronectin binding partner is immobilized in two or more discrete locations, such as in a dot blot, a microplate or microarray, each discrete location can be independently treated, where all locations can be differently treated, or some locations can be equally treated while others are differently treated. Exemplary treatments include contacting with different dilutions of sample, contacting with different soluble or mobile binding partners, including binding partners that bind to different regions of an oncofetal fibronectin indicating molecule such as EDA, EDB and IICS, and contacting under different buffer conditions. Such methods are known to those skilled in the art.

[0820] Complex formation of the binding partner and an oncofetal fibronectin indicating molecule can be determined in a number of ways. Complex formation can be determined by use of soluble or mobile fibronectin or oncofetal fibronectin binding partners. The soluble or mobile fibronectin or oncofetal fibronectin binding partners can be detected directly using, for example, a detectable moiety conjugated with the binding partner, or indirectly using binding partners that bind fibronectin or oncofetal fibronectin binding partners. The assay can be quantitative, for example, can be an enzyme-linked immunosorbent assay (ELISA) in which at least a first fibronectin or oncofetal fibronectin binding partner is immobilized to a solid support and at least a second fibronectin or oncofetal fibronectin binding partner is soluble or mobile. An oncofetal fibronectin assay can be based on competitive inhibition, where an oncofetal fibronectin indicating molecule in the sample competes with a known amount of oncofetal fibronectin indicating molecule or analog thereof (typically labeled) for a fibronectin or oncofetal fibronectin binding partner. For example, in an assay for oncofetal fibronectin protein, any oncofetal fibronectin protein present in the sample can compete with a known amount of labeled oncofetal fibronectin protein or a labeled analog thereof for the fibronectin or oncofetal fibronectin binding partner. The amount of labeled oncofetal fibronectin indicating molecule affixed to the solid phase or remaining in solution can be measured, and the measurement can be used to determine the amount of oncofetal fibronectin indicating molecule in the sample, using methods known in the art.

[0821] Detection of a signal also can be used to quantitate the amount of an oncofetal fibronectin indicating molecule present in the sample, using any of a variety of known methods. For example, the intensity of the signal corresponding to an oncofetal fibronectin indicating molecule can be measured using any of a variety of methods known in the art, including, for example, fluorescence or absorption spectrometry, or phosphor imager measurement. In one example, known concentrations of a standard also can be included and the intensity of one or more sample signals can be qualitatively or quantitatively compared to the standards using known methods to provide an estimate or calculation of the amount of an oncofetal fibronectin indicating molecule in the sample. In another example, multiple dilutions of the sample can be detected, and the signal measured at each dilution can be used to estimate the amount of an oncofetal fibronectin indicating molecule using known methods.

[0822] i. Sandwich Assay

[0823] An oncofetal fibronectin indicating molecule can be detected when bound to two or more fibronectin or oncofetal fibronectin binding partners, where such a complex can produce a signal indicative of complex formation. For example, presence of an oncofetal fibronectin indicating molecule bound to an immobilized fibronectin or oncofetal fibronectin binding partner can be detected by detecting presence of a soluble or mobile fibronectin or oncofetal fibronectin binding partner or conjugate thereof, bound to a solid support, as a result of a complex with an oncofetal fibronectin indicating molecule. Detection can be performed immediately after contacting the solid support with a sample or after one or more subsequent steps such as washing steps. In assays in which no label soluble or mobile fibronectin or oncofetal fibronectin binding partner conjugate is added, the solid support can optionally be washed and the solid support can be contacted with one or more additional reagents, such as a binding partner that binds a fibronectin or oncofetal fibronectin binding partner.

[0824] In some instances, the mobile or soluble fibronectin or oncofetal fibronectin binding partner will be detected, either directly or indirectly. Direct detection can include, for example, detecting a detectable moiety conjugated to the mobile or soluble fibronectin or oncofetal fibronectin binding partner. Indirect detection can include, for example, detecting a binding partner that binds the mobile or soluble fibronectin or oncofetal fibronectin binding partner. For example, indirect detection of a soluble mouse anti-oncofetal fibronectin antibody can be accomplished by contacting a solid support with a goat anti-mouse IgG antibody conjugated to horseradish peroxidase and measuring light formation upon using the appropriate substrate. In any of the above assays, presence of a detectable signal on the solid support can indicate that the soluble binding partner is bound to an oncofetal fibronectin indicating molecule that is bound to an immobilized binding partner. Thus, presence of a detectable signal on the solid support can indicate the presence of an oncofetal fibronectin indicating molecule in a sample.

[0825] The immobilized fibronectin or oncofetal fibronectin binding partner, sample, and soluble or mobile fibronectin or oncofetal fibronectin binding partner can be added to each other in any order, as will be understood by one skilled in the art. For example, a sample can be treated with a soluble or mobile fibronectin or oncofetal fibronectin binding partner prior to contacting the sample with the solid support (which contains immobilized fibronectin or oncofetal fibronectin binding partner), and subsequent steps can include contacting the solid support with the sample that contains the soluble or mobile fibronectin or oncofetal fibronectin binding partner. In another example, a sample can be contacted with a solid support containing immobilized fibronectin or oncofetal fibronectin binding partner prior to contacting the sample with a soluble or mobile fibronectin or oncofetal fibronectin binding partner, and subsequent steps can include contacting the solid support with the soluble or mobile fibronectin or oncofetal fibronectin binding partner. In another example, a solid support containing immobilized fibronectin or oncofetal fibronectin binding partner can be contacted with a soluble or mobile fibronectin or oncofetal fibronectin binding partner prior to

either binding partner contacting the sample, and subsequent steps can include contacting the solid support with the sample.

[0826] ii. Test Device

[0827] A test device can contain an immobilized fibronectin or oncofetal fibronectin binding partner. Any test device that can be used to produce a signal, or from which a signal can be generated, is intended for use as part of the methods, combinations and kits provided herein (e.g., lateral flow formats and vertical flow formats). A test device can contain a test strip and one or more additional components used for determining the presence of an oncofetal fibronectin indicating molecule.

[0828] a. Test Strip

[0829] A test strip can be used in the test device to indicate the presence of an oncofetal fibronectin indicating molecule in a sample. Any test strip that can provide a detectable signal, for example, visually inspectable or reader-adapted test strip can be used in the methods, combinations and kits provided herein. Such test strip devices and methods for use as are known to those skilled in the art can be used in systems described herein (see, e.g., U.S. Pat. Nos. 6,394,952, 6,267,722, 5,658,801, 5,656,503, 5,656,502, 5,654,162, 5,622,871, 5,591,645, 5,578,577, 5,500,375, 5,270,166, 5,252,459, 5,209,904, 5,149,622, 5,132,097, 5,120,643, 5,073,484, 4,960,691 and 4,956,302).

[0830] A test strip generally can accommodate flow of a liquid sample and contains a region containing a fibronectin or oncofetal fibronectin binding partner and/or a fibronectin or oncofetal fibronectin binding partner conjugate immobilized thereto. A test strip also can contain one or more additional regions such as a region for applying the sample, a region for removing particulate or solid or undissolved matter from the sample, a region for removing background material that can non-specifically bind to a fibronectin or oncofetal fibronectin binding partner, a region containing a non-specific binding compound, a region containing a fibronectin or oncofetal fibronectin binding partner or conjugate thereof that is mobilizable upon contact with the sample, a region containing a binding partner that can specifically bind to a fibronectin or oncofetal fibronectin binding partner or conjugate thereof, and combinations thereof.

[0831] A test strip can include a membrane system that defines a liquid flow pathway. Exemplary test strips and systems include the Rapid fFN Cassette and the fFN membrane immunoassay. The test strips can be visually inspected, or used in conjunction with a test strip reader, such as the point of care device described in U.S. Pat. Nos. 6,267,722 and 6,394,952.

[0832] When a reader is used, one or more measurements can be made by the reader and the one or more measurements can be subjected to further analysis, such as image reconstruction or otherwise classifying an image. Methods for processing of reflectance data and methods of classifying an image are known in the art, as exemplified in U.S. Pat. No. 6,267,722.

[0833] The results of the test strip measurement and/or the results from classifying an image can be used alone, or in conjunction with other information input into a decision

support system, such as a neural network that can analyze a variety of data or information to guide further testing or treatment of a subject. Such neural nets generally analyze patient data or information, typically patient history or clinical data (see, e.g., U.S. Pat. Nos. 6,678,669 and 6,267,722).

[0834] Lateral flow test immunoassay devices are among those that can be employed in the methods herein. In such devices, a membrane system forms a single fluid flow pathway along the test strip. The membrane system includes components that act as a solid support for immunoreactions. For example, porous or bibulous or absorbent materials can be placed on a strip such that they partially overlap, or a single material can be used, in order to conduct liquid along the strip. The membrane materials can be supported on a backing, such as a plastic backing. In an exemplary embodiment, the test strip includes a glass fiber pad, a nitrocellulose strip and an absorbent cellulose paper strip supported on a plastic backing.

[0835] Fibronectin or oncofetal fibronectin binding partners and/or conjugates thereof can be immobilized on the solid support. The binding partners or conjugates can be bound to the test strip by adsorption, ionic binding, van der Waals adsorption, electrostatic binding, or by covalent binding, by using a coupling agent, such as glutaraldehyde. For example, the binding partners or conjugates can be applied to the conjugate pad and nitrocellulose strip using standard dispensing methods, such as a syringe pump, air brush, ceramic piston pump or drop-on-demand dispenser. In one embodiment, a volumetric ceramic piston pump dispenser is used to stripe binding partners that bind to the analyte of interest, including a labeled binding partner conjugate, onto a glass fiber conjugate pad and a nitrocellulose strip.

[0836] The test strip can be otherwise treated, for example, with sugar to facilitate mobilization of reagents including a fibronectin or oncofetal fibronectin binding partner or conjugate thereof, or with proteins, such as albumins, including bovine (BSA), immunoproteins, other animal proteins, water-soluble polyamino acids, or casein to block non-specific binding sites on the test strip. In one embodiment, a binding partner that can bind to a labeled fibronectin or oncofetal fibronectin binding partner conjugate is immobilized on the test strip; such a binding partner can bind to a labeled fibronectin or oncofetal fibronectin binding partner conjugate that is not complexed with an oncofetal fibronectin indicating molecule and thereby can be used as a control. For example, where the labeled conjugate includes a mouse monoclonal anti-oncofetal fibronectin antibody, a polyclonal goat anti-mouse IgG antibody can be used to bind the conjugate.

[0837] In test strips using a defined liquid sample flow pathway, a test strip can contain two or more separate regions, where the sample contacts a first region prior to contacting a second region. Typically, all regions can accommodate a liquid sample such that at least a portion of the liquid sample can pass through the region and, if applicable, interact with a binding partner immobilized thereto. For example, a test strip can contain a region for applying the sample and a region containing immobilized fibronectin or oncofetal fibronectin binding partner, where the sample contacts the sample application region prior to contacting the binding partner region.

[0838] A test strip can contain a region for applying the sample. This region can be referred to as the sample application region. This region is the region first contacted by the sample. This region can be formed from any of a variety of substances to form a solid structure capable of accommodating a liquid sample. This region also can be the site for one or more additional regions such as a filter region, non-specific surface region, or other region described herein.

[0839] A test strip can contain a region containing a fibronectin or oncofetal fibronectin binding partner or a conjugate thereof, immobilized onto the test strip. This region can be referred to as a fibronectin or oncofetal fibronectin binding region. Typically, the immobilized binding partner or conjugate will be in a region that can accommodate a liquid sample such that the liquid sample can interact with the immobilized binding partner or conjugate. An immobilized fibronectin or oncofetal fibronectin binding partner can react with the sample to specifically bind a fibronectin or oncofetal fibronectin indicating molecule, respectively. Accordingly, this region of the test strip can be a region where an oncofetal fibronectin indicating molecule in a sample forms a complex with one or more fibronectin and/or oncofetal fibronectin binding partners. Exemplary binding partners include polyclonal anti-fibronectin antibodies, monoclonal anti-oncofetal fibronectin antibodies, heparin, collagen, an integrin, fibrin, or a nucleic acid complementary to an oncofetal fibronectin nucleotide sequence. An additional exemplary binding partner is oncofetal fibronectin protein or an oncofetal fibronectin encoding nucleic acid molecule or fragment thereof, which can bind to an autoantibody to oncofetal fibronectin protein or nucleic acid present in the sample.

[0840] A test strip also can contain a filter region. A filter region can be a region in which particulate, solid or undissolved matter present in a sample that are greater than a cutoff size are physically blocked from advancing through the test strip. Exemplary matter that can be filtered include cells, mucus, debris, and insoluble matter. Cutoff sizes can be any of a variety of sizes, according to the selected substances to be removed from the sample. For example, a cutoff filter size can be 10 mm, 5 mm, 1 mm, 500 μm , 200 μm , 100 μm , 50 μm , 20 μm , 10 μm , 5 μm , 2 μm , 1 μm , 0.5 μm , 0.2 μm or 0.1 μm , or about 10 mm, 5 mm, 1 mm, 500 μm , 200 μm , 100 μm , 50 μm , 20 μm , 10 μm , 5 μm , 2 μm , 1 μm , 0.5 μm , 0.2 μm or 0.1 μm . Filter regions can be formed from any of a variety of substances used for filtration of particulate matter, including glass (e.g., glass wool), cellulose, nylon, polyether sulfone, polyvinyl chloride, teflon and any other substance having the selected cutoff size and properties. In one embodiment, the filter substance is a low protein binding substance. The filtering region can be located at the region of sample application to the test strip, or can be located downstream of the sample application region. Typically, the filter region is located upstream of a fibronectin or oncofetal fibronectin binding region.

[0841] A test strip also can contain a region containing a non-specific binder. This region can be referred to as a non-specific binding region. Non-specific binder can be present on solid structures, where the non-specific binder binds to at least a portion of background material in a sample while not binding more than a small amount (e.g., less than 10%) of an oncofetal fibronectin indicating molecule in the sample. Possible solid supports for non-specific binders

include paper and cellulose derivatives, such as cellulose esters and ethers, natural and synthetic polymeric materials, such as latex, vinyl polymers, polypropylene, polyethylene and partially hydrolyzed derivatives, polycondensates, copolymers and inorganic materials. In one embodiment, a non-specific binder is a porous or bibulous member capable of transporting a liquid sample along a test strip. Non-specific binders that can be used include solid supports having immobilized thereon one or more non-specific binding proteins such as albumin (including bovine serum albumin, or BSA), antibodies not specific for an oncofetal fibronectin indicating molecule and other surfaces known in the art or disclosed herein. Exemplary proteins that can be used for a non-specific binder include BSA, methylated BSA or antibodies such as W632 or mouse IgG. In one example, a non-specific binder can be a nitrocellulose membrane having BSA immobilized thereon. A non-specific binder can be at the same region as a sample application region, or downstream of a sample application region. A non-specific binder can be at the same region as a filter region, if present, or upstream or downstream of a filter region, if present. A non-specific binder is typically upstream of a fibronectin or oncofetal fibronectin binding region.

[0842] A test strip also can contain a region having immobilized thereto a binding partner that can bind to a fibronectin or oncofetal fibronectin binding partner. This region can be referred to as a control region. Such a region can act as a positive or negative control, according to the design of the test strip. Typically, the immobilization will be in a region that can accommodate a liquid sample such that the liquid sample can interact with the immobilized binding partner that binds to a fibronectin or oncofetal fibronectin binding partner. An immobilized binding partner of a fibronectin or oncofetal fibronectin binding partner can specifically bind to a fibronectin or oncofetal fibronectin binding partner. In one example, the immobilized binding partner of a fibronectin or oncofetal fibronectin binding partner can bind to a fibronectin or oncofetal fibronectin binding partner to which an oncofetal fibronectin indicating molecule is not bound. Exemplary binding partners include polyclonal anti-mouse IgG antibodies or other antibodies that specifically bind to a protein or nucleic acid or other compound that binds to a fibronectin or oncofetal fibronectin indicating molecule, or a nucleic acid molecule complementary to a nucleic acid molecule encoding a fibronectin or oncofetal fibronectin. A control region is typically downstream from a sample application region. A control region can be at the same region as a filter region, if present, or can be downstream of a filter region, if present. A control region can be upstream or downstream of a fibronectin or oncofetal fibronectin binding region, or upstream and downstream (i.e., two or more control regions can be present) of a fibronectin or oncofetal fibronectin binding region. A control region can be at the same region, upstream or downstream of a non-specific binding region (if present), or combinations thereof.

[0843] A test strip also can contain a region containing a mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate thereof. Such a region can be termed a mobilization region. A mobilizable fibronectin or oncofetal fibronectin binding partner can specifically bind to a fibronectin or oncofetal fibronectin indicating molecule. Exemplary fibronectin or oncofetal fibronectin binding partners include polyclonal anti-fibronectin antibodies, monoclonal anti-oncofetal fibronectin antibodies, heparin, col-

lagen, an integrin, fibrin, or a nucleic acid molecule complementary to a fibronectin or oncofetal fibronectin encoding nucleotide sequence. An additional exemplary binding partner is oncofetal fibronectin protein or nucleic acid molecule encoding oncofetal fibronectin or fragment thereof, which can bind to an autoantibody to oncofetal fibronectin protein or nucleic acid present in the sample. A mobilizable binding partner or conjugate can be a compound which, upon contact with a liquid sample, is mobilized such that the mobilizable compound can interact with solutes of the liquid sample and the mobilizable compound can migrate along the test strip as the liquid sample migrates along the test strip. For example, a mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate thereof can be a water soluble compound affixed to the surface of the test strip in a non-aqueous format (e.g., in the absence of any solvent, or in a non-aqueous solvent). Accordingly, a mobilizable compound that is a fibronectin or oncofetal fibronectin binding partner can, upon mobilization, bind or react with an oncofetal fibronectin indicating molecule present in the sample. A mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate thereof can be located in the sample application region or downstream of the sample application region. A mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate thereof can be located at the fibronectin or oncofetal fibronectin binding region or upstream of the fibronectin or oncofetal fibronectin binding region. A mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate thereof can be located upstream, at the same region, or downstream of a filter region (if present), or combinations thereof. A mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate thereof can be located upstream, at the same region, or downstream of a non-specific binding region (if present), or combinations thereof. A mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate thereof is typically located upstream of a control region, if present.

[0844] The above description of a test strip is meant to exemplify different possible regions that can be present and arrangements of possible regions; the above description is not meant to limit possible test strip combinations to those described above, since a variety of combinations will be apparent to one skilled in the art according to the selected assay configuration. A test strip can contain various combinations of the regions described herein, including two or more of the same type of regions in different locations along the test strip. A test strip also can contain two or more different compositions in the same region (e.g., a non-specific binding region can be at the same location as the filter region).

[0845] b. Test Strip Housing

[0846] The test strip optionally can be contained within a housing. Such a housing can serve any of a variety of purposes, including facilitating handling of the test strip or for insertion into a reflectance reader. A variety of test strip housings are known in the art, as exemplified in U.S. Pat. No. 6,267,722.

[0847] In an exemplary embodiment, the test strip housing includes a symbology, such as a bar code that can be associated with data related to the assay device, subject data and/or test run. For example, information associated with the device, such as lot number, expiration date, analyte and

intensity value, or information related to the test run, such as date, calibration data, reflectance value or other such information, can be encoded and associated, such as in a database with a bar code imprinted on the device. Any bar code system that provides the appropriate line thickness and spacing can be used. Code 39 and Code 128 are among the known bar code systems.

[0848] In a particular embodiment, Code 39 is used. The bar code is made up of 11 alphanumeric characters, including 2 alphabetic and 9 numeric characters. The first and last characters are asterisks (*), as is standard in the Code 39 system. The lot number is stored as 1 alpha and 4 numeric codes so that product complaints or questions can be traced to a particular lot number. In the exemplified embodiment, the first character represents the month of production, the second is a digit representing the year of production and the last three are an index value indicating the lot number. Thus, the lot number "A8001" represents the first device in a lot produced in January, 1998. The next two characters ("01") represent the identity of the analyte as 2 numerics (00-99). This permits the use of up to 100 different analytes with the system. The reflectance intensity value (00-99) is stored as the next two numeric characters ("01"). The intensity value sets the reference threshold for which controls and subject samples can be compared. This eliminates the need to run liquid reference samples on a daily basis. Finally, the cassette expiration date is stored as 1 alpha and 1 numeric code to prevent the use of expired devices. In the example given, an expiration code of "A9" represents an expiration date of January, 1999.

[0849] c. Analysis with a Test Device

[0850] A volume of sample can be delivered to a test strip using any known device for transporting a sample, for example, a standard plastic pipet. In one embodiment, for example, when the sample is liquid, the neat (e.g., undiluted or without added reagents) sample can be applied to the test strip, which can include direct application to the test strip (e.g., contacting the test strip with a urine stream). In another embodiment, any an oncofetal fibronectin indicating molecule present in the sample can bind to a labeled mobilizable fibronectin or oncofetal fibronectin binding partner conjugate (e.g., labeled anti-oncofetal fibronectin antibody conjugate) and the resulting complex migrates along the test strip. Alternatively, the sample can be pre-mixed with a labeled conjugate prior to applying the mixture to the test strip. When the labeled complex encounters a fibronectin or oncofetal fibronectin binding region of the test strip, an immobilized fibronectin or oncofetal fibronectin binding partner (e.g., anti-fibronectin antibody) therein can bind to the complexed an oncofetal fibronectin indicating molecule to form a sandwich complex, thereby resulting in a region in which the label of the mobilizable fibronectin or oncofetal fibronectin binding partner conjugate accumulates and can be detected.

[0851] In one embodiment, prior to contacting the fibronectin or oncofetal fibronectin binding region of a test strip, the sample can be contacted with a non-specific binder such as a non-specific binding compound or a non-specific binding surface. For example, a sample can be added to a test strip that is configured to have the sample flow through a non-specific binding region prior to flowing through a fibronectin or oncofetal fibronectin binding region. By first

contacting the non-specific binder such as a non-specific binding compound or non-specific binding surface, background materials in the sample, which might otherwise non-specifically bind to a fibronectin or oncofetal fibronectin binding partner or conjugate thereof, instead bind to the non-specific binding compound or surface such that the background materials are at least partially prohibited from binding to a fibronectin or oncofetal fibronectin binding partner or conjugate thereof.

[0852] In another embodiment, prior to contacting a non-specific binder such as a non-specific binding compound or a non-specific binding surface, the sample can be contacted with a mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate thereof. For example, a sample can be added to a test strip that is configured to have the sample flow through a mobilization region containing a fibronectin or oncofetal fibronectin binding partner conjugate prior to flowing through a non-specific binding region. By first contacting the mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate, an oncofetal fibronectin indicating molecule that might otherwise attach to the non-specific binder such as the non-specific binding compound or non-specific binding surface, instead bind to the mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate such that a larger amount of an oncofetal fibronectin indicating molecule can be bound to a binding partner.

[0853] In another embodiment, prior to contacting a mobilizable fibronectin or oncofetal fibronectin binding partner or conjugate thereof, the sample can be contacted with a non-specific binder such as a non-specific binding compound or a non-specific binding surface. For example, a sample can be added to a test strip that is configured to have the sample flow through a non-specific binding region prior to flowing through a mobilization region containing a fibronectin or oncofetal fibronectin binding partner conjugate. By first contacting the non-specific binder such as a non-specific binding compound or non-specific binding surface, background materials in the sample, which might otherwise non-specifically bind to a fibronectin or oncofetal fibronectin binding partner or conjugate, instead bind to the non-specific binding compound or surface such that the background materials are at least partially prohibited from binding to a fibronectin or oncofetal fibronectin binding partner or conjugate.

[0854] As the sample passes through the fibronectin or oncofetal fibronectin binding region, any unbound binding partner continues to migrate into a control zone where it can be captured by an immobilized binding partner that can bind to the fibronectin or oncofetal fibronectin binding partner. For example, a goat anti-mouse IgG antibody can be located in a control region and can bind to a mouse anti-oncofetal fibronectin antibody conjugate. The complex formed between the immobilized binding partner that binds to an fibronectin or oncofetal fibronectin binding partner and the fibronectin or oncofetal fibronectin binding partner can form a detectable signal in this region, such as a colored stripe, that reflects the aggregation of the labeled conjugate. Presence of a detectable signal in this region can indicate that the assay run has completed and also can serve as a positive control.

[0855] The results of the assay can be assessed using a reader and associated software. Use of the point of care

device described in U.S. Pat. Nos. 6,267,722 and 6,394,952 provides, at the very least, the same clinically relevant information as an onfFN ELISA (an enzyme linked immunosorbent sandwich assay (ELISA); see, e.g., U.S. Pat. No. 5,281,522), but in significantly less time and at the point of care. This oncofetal fibronectin immunoassay allows the user to test a cervicovaginal swab sample in 20 minutes or about 20 minutes. When comparing the 20 minute rapid onfFN test to the data from the onfFN ELISA, a Kappa coefficient of 0.81 was found with a 95% confidence interval [0.75, 0.88] and an overall concordance of at least 94.9%. These data were obtained using a system including an immunoassay test strip in combination with a reflectance reader and data processing software employing data reduction and curve fitting algorithms or neural networks, as known in the art.

[0856] iii. Quantitation

[0857] Immobilized fibronectin or oncofetal fibronectin binding partner also can be used in a format amenable to quantitation of the amount of oncofetal fibronectin indicating molecule in a sample. For example, a fibronectin or oncofetal fibronectin binding partner can be immobilized on a solid support that can be used in spectrophotometric measurements.

[0858] In one example, the amount of oncofetal fibronectin protein in a sample can be quantitated using an enzyme-linked immunosorbent assay (ELISA). An exemplary ELISA method can be performed by coating one or more wells of a reaction vessel such as a microtiter plate or microtiter strip with a fibronectin or oncofetal fibronectin binding partner, and incubating a sample in such wells. The incubated wells can be washed and then reacted with a mobile or soluble fibronectin or oncofetal fibronectin binding partner or conjugate thereof, and then washed again. The amount of mobile or soluble fibronectin or oncofetal fibronectin binding partner bound to the wells can be indirectly or directly measured spectrophotometrically. Methods for preparing ELISA plates and reagents and performing ELISA are known in the art, and can be used for the oncofetal fibronectin indicating molecule detection methods provided herein.

[0859] iv. Affinity-Based Isolation of Oncofetal Fibronectin

[0860] A fibronectin or oncofetal fibronectin binding partner can be used to specifically isolate an oncofetal fibronectin indicating molecule. A fibronectin or oncofetal fibronectin binding partner can be used to increase the concentration of an oncofetal fibronectin indicating molecule in a sample. A fibronectin or oncofetal fibronectin binding partner can be used to increase the concentration of an oncofetal fibronectin indicating molecule in a sample relative to the concentration of one or more background components of the sample, thereby increasing the purity of the oncofetal fibronectin indicating molecule in the sample. In accordance with other uses provided herein or otherwise known in the art, a fibronectin or oncofetal fibronectin binding partner can be affixed to a solid support and contacted with a sample. Any oncofetal fibronectin indicating molecule present in the sample can bind to the binding partner and one or more subsequent steps (e.g., washing and elution steps) can be used to separate the oncofetal fibronectin indicating mol-

ecule from background material and/or decrease the volume in which the oncofetal fibronectin indicating molecule is present.

[0861] A fibronectin or oncofetal fibronectin binding partner can be affixed to any solid support as described herein or known in the art. For example, a fibronectin or oncofetal fibronectin can be affixed to beads such as beads used in liquid chromatography, magnetic beads, or any beads that can be isolated by physical methods (e.g., centrifugation). The solid support can be in any form, including, but not limited to, in the form of a liquid chromatographic column or a slurry of beads.

[0862] The solid support can be contacted with a sample under conditions in which the fibronectin or oncofetal fibronectin binding partner can specifically bind to an oncofetal fibronectin indicating molecule. In one embodiment, the solid support is contacted with a sample under conditions in which background materials do not significantly interfere with the binding partners on the solid support binding an oncofetal fibronectin indicating molecule (i.e., 10% or about 10% or less of the binding partners on the solid support bind to background materials or 90% or about 90% or more of oncofetal fibronectin indicating molecule in the sample is bound by binding partners on the solid support). Exemplary conditions include standard phosphate-buffered saline (PBS) conditions (e.g., 137 mM NaCl, 2.7 mM KCl, 10 mM phosphate buffer pH 7.4).

[0863] In one embodiment, the solid support is washed with a solution that removes background material from the solid support in preference to removal of an oncofetal fibronectin indicating molecule from the solid support. Such conditions will typically remove 50% or about 50% or more background material while removing 10% or about 10% or less oncofetal fibronectin indicating molecule from the solid support. The conditions can be the same as initial conditions for contacting the sample and solid support, or can be different. Exemplary conditions include increased salt or detergent concentrations, or different pH, relative to the initial conditions for contacting the sample and solid support. Methods for determining conditions are known in the art.

[0864] An oncofetal fibronectin indicating molecule can be released from the solid support by any of a variety of methods known in the art. For example, an oncofetal fibronectin indicating molecule can be released from the solid support by a change in pH or ionic strength of the ambient buffer. In another example, an oncofetal fibronectin indicating molecule can be released from the solid support by subjecting the oncofetal fibronectin indicating molecule to denaturing conditions including denaturing, salt, pH, urea, detergent or temperature conditions. An exemplary condition for release of an oncofetal fibronectin indicating molecule from the solid support is 100 mM glycine, pH 2.5-3.0.

[0865] After release of the oncofetal fibronectin indicating molecule from the solid support, the solution containing the oncofetal fibronectin indicating molecule can be used directly for detection of the oncofetal fibronectin indicating molecule or can be treated prior to detection methods. Exemplary treatment includes adding a concentration solution of APB in order to achieve a final, diluted concentration of 1×APB or 0.05 M Tris buffer, pH 7.4, 150 mM NaCl, 0.02% NaN₃, 1% BSA, 5 mM EDTA, 1 mM phenylmeth-

ylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin, and 0.1% Triton X-100.

[0866] Presence of an oncofetal fibronectin indicating molecule can be detected using any of a variety of methods, such as the methods provided herein or detection methods known in the art. For example, an oncofetal fibronectin indicating molecule can be detected by gel electrophoresis, including SDS-PAGE and a band having a molecular weight corresponding to an oncofetal fibronectin indicating molecule can indicate the presence of an oncofetal fibronectin indicating molecule in the sample. An oncofetal fibronectin indicating molecule also can be detected using any of the methods described herein that use a binding partner conjugate to detect the presence of an oncofetal fibronectin indicating molecule, including, but not limited to, sandwich assays and blot analyses.

[0867] d. Detection of Regions of Oncofetal Fibronectin

[0868] Binding partners can be used to detect the presence of particular regions in an oncofetal fibronectin indicating molecule. Detection of the presence of particular regions in an oncofetal fibronectin indicating molecule can serve a variety of purposes, including identifying the likely cell or tissue or organ source of the oncofetal fibronectin indicating molecule, identifying the unlikely cell or tissue or organ source of the oncofetal fibronectin indicating molecule, or identifying a health problem associated with a particular form of oncofetal fibronectin. In one example, the antibody L19 can be used to detect the presence of EDB in an oncofetal fibronectin protein. Binding partners that bind oncofetal fibronectin proteins or autoantibodies to oncofetal fibronectin proteins also can be used to detect the presence of one or more post translational modifications in an oncofetal fibronectin protein. For example, the antibody FDC-6 can be used to detect the presence of O-glycosylation at threonine 33 of the IIICS region of oncofetal fibronectin protein. Binding partners also can be used to detect a IIICS splice variant (e.g., V0, V64, V89, V95 or V120 splice variants of IIICS) or to detect the presence of one or more splice regions of IIICS (e.g., aa1-25, aa26-89 or aa90-120 splice regions of IIICS).

[0869] Detection of particular oncofetal fibronectin regions, IIICS splice regions and post-translational modifications, can serve to characterize the oncofetal fibronectin indicating molecule in the sample. For example, binding partners can be used to characterize an oncofetal fibronectin indicating molecule present in a sample as containing or lacking the EDA, EDB or IIICS regions. Binding partners can be used to characterize oncofetal fibronectin protein present in a sample as containing or lacking one or more particular post-translational modifications, such as O-glycosylation at threonine 33 of IIICS. Binding partners can be used to characterize an oncofetal fibronectin indicating molecule present in a sample as containing or lacking a particular splice variant of IIICS such as V0, V64, V89, V95 or V120.

[0870] In using the binding methods disclosed herein to characterize an oncofetal fibronectin indicating molecule, two or more binding partners can be used in the same or different assays performed on a sample. Each binding partner can provide information regarding the composition of the oncofetal fibronectin indicating molecule in a sample. For example, if an oncofetal fibronectin protein binds

FDC-6 and IST-9, but not L19, the oncofetal fibronectin protein can be characterized as containing EDA and IIICS and containing an O-glycosylation at threonine 33 of IIICS, but not containing EDB. Thus, provided herein are methods for characterizing an oncofetal fibronectin indicating molecule in a sample by identifying one or more binding partners to which the oncofetal fibronectin indicating molecule is bound and by identifying any binding partners which are not bound to the oncofetal fibronectin indicating molecule. The bound and unbound binding partners (e.g., a binding profile) can be indicative of a region of oncofetal fibronectin or indicative of a particular oncofetal fibronectin variant. In one embodiment, such methods can be performed by comparing bound and unbound binding partners in an assay of a sample to bound and unbound binding partners in an assay of a reference (e.g., bound and unbound binding partners contacted with a known oncofetal fibronectin indicating molecule) or by comparing a binding profile to a calculated binding profile.

[0871] 3. Detection of Oncofetal Fibronectin by Mass Spectrometry

[0872] Disclosed herein are methods of detecting an oncofetal fibronectin indicating molecule using mass spectrometric formats. Using mass spectrometry, an atom, molecule or molecule fragment, such as a fragment of oncofetal fibronectin protein or oncofetal fibronectin-encoding nucleic acid or complement thereto, can be detected using mass spectrometry. The presence of that atom, molecule or molecule fragment, can indicate the presence of an oncofetal fibronectin indicating molecule in a sample. Oncofetal fibronectin indicating molecules detected using mass spectrometric methods described herein, include, but are not limited to, an oncofetal fibronectin protein, an autoantibody of oncofetal fibronectin protein or nucleic acid encoding oncofetal fibronectin, mRNA encoding oncofetal fibronectin, amplicates of the aforementioned mRNA, and fragments thereof.

[0873] A variety of mass spectrometric techniques can be used to perform the oncofetal fibronectin indicating molecule detection methods provided herein. Mass spectrometric techniques generally include desorption and detection methods. Any known desorption method can be used herein, including, for example, ultraviolet (UV) and infrared (IR) Matrix-Assisted Laser Desorption/Ionization (MALDI; see, e.g., published International PCT Application No. WO 99/57318 and U.S. Pat. No. 5,118,937) and electrospray (ES). Selection of the particular desorption method to be used can be made by one skilled in the art according to the selected mass measurement to be performed. Any known detection method can be used herein, including, for example, time-of-flight (TOF), Fourier transform and magnetic sector/magnetic deflection instruments in single or triple quadrupole mode. Selection of the particular detection method to be used can be made by one skilled in the art according to the selected mass measurement to be performed.

[0874] In some embodiments, prior to detection by mass spectrometry by the methods disclosed herein, a sample can be manipulated in one or more steps, which can include, for example, isolation of an oncofetal fibronectin indicating molecule, fragmentation of an oncofetal fibronectin indicating molecule and sample conditioning. For example, a

sample containing an oncofetal fibronectin indicating molecule can be first treated in a step of isolating an oncofetal fibronectin indicating molecule, then second treated in a step of contacting the sample with a fragmenting compound and third treated in a step of measuring the fragment masses using mass spectrometry.

[0875] In one embodiment, the mass spectrometry is used to measure the molecular weights of a sample treated with a protease, such as trypsin or cathepsin D. Measurement of one or more protease fragments within defined mass ranges can indicate the presence of oncofetal fibronectin protein in a sample.

[0876] The mass spectrometric methods provided herein can be used to detect both proteins and nucleic acid molecules that indicate the presence of oncofetal fibronectin in a subject. As is understood by one skilled in the art methods for protein detection by mass spectrometry (and sample treatment methods prior to detection) can differ from nucleic acid molecule detection methods. The methods provided herein can be modified by routine methods to detect the appropriate analyte.

[0877] a. Sample Manipulation

[0878] Prior to mass measurement using mass spectrometry, a sample can be manipulated and/or treated in one or more steps. Exemplary manipulation steps include, but are not limited to, contacting the sample with an ionic surface, contacting the sample with a hydrophobic surface, contacting a sample with a fibronectin or oncofetal fibronectin binding partner and contacting a sample with a fragmentation compound. Such manipulation also can include conditioning, which includes any procedures that improve resolution of a mass spectrum. Such manipulation steps also can include one or more rinsing steps where oncofetal fibronectin is at least partially separated from background material in the sample.

[0879] The sample manipulation steps can occur in any reaction vessel and can be performed immediately before mass measurement, or one or more hours or one or more days before mass measurement or concurrent therewith. In one embodiment, at least one sample manipulation step is performed on a mass spectrometry substrate. In another embodiment, two or more sample manipulation steps are performed on a mass spectrometry substrate.

[0880] Such manipulation steps include any of a variety of surface-enhanced laser desorption ionization (SELDI) mass spectrometric methods known in the art. For example, a mass spectrometry substrate can be coated with a substance or compound for sample manipulation, such as a reverse phase substance, ion exchange substance, binding partner, metal affinity substance, or other substances or compounds known in the art. A substrate can contain such a substance or compound at separate discrete locations. A substrate also have combinations of such substances or compounds at the same location or at separate discrete locations. When a substrate contains two or more discrete locations, sample can be applied to each discrete location, and each discrete location can have added thereto the same solution or different solutions. When a particular substance or compound or combination is present at multiple discrete locations, each location can have added thereto different solutions to result in variation in the compounds adhered to each discrete

location; for example, discrete locations can be treated with different solutions to form a gradient ranging from lower specificity to higher specificity. These and other SELDI methods are known in the art, as exemplified in U.S. Pat. Nos. 5,719,060, 5,894,063, 6,124,137 and 6,225,047.

[0881] i. Contact with Binding Partner

[0882] A sample can be contacted with a binding partner, such as an anti-oncofetal fibronectin antibody, or other moiety that binds to a fibronectin or oncofetal fibronectin indicating molecule with greater affinity than to other components in the sample. A binding partner can be used, as disclosed herein, to specifically and/or preferentially bind to an oncofetal fibronectin indicating molecule. For example, by contacting a sample with a binding partner and subsequently detecting the molecular weights of sample components that bind to the binding partner, an oncofetal fibronectin indicating molecule can be more readily detected among the components in the sample. Such a contacting step can achieve a reduction in the number of different masses measured and can resolve and enrich masses corresponding to an oncofetal fibronectin indicating molecule relative to masses of sample components that do not correspond to the oncofetal fibronectin indicating molecule.

[0883] For example, a binding partner immobilized on a solid support can be contacted with a sample under conditions in which an oncofetal fibronectin indicating molecule in the sample can bind to the binding partner. After contacting the sample with the solid support, the solid support can optionally be washed. Sample components bound to the solid support can be measured by mass spectrometry by desorbing the sample components using mass spectrometric desorption methods such as MALDI or ES, or sample components can be removed from the solid support using solvent conditions in which an oncofetal fibronectin indicating molecule no longer binds to the fibronectin or oncofetal fibronectin binding partner. Sample components removed from the solid support by solvent conditions optionally can be subjected to one or more subsequent sample manipulation steps prior to mass measurement of the sample components, including one or more fragmentation steps.

[0884] When a sample contains an oncofetal fibronectin indicating molecule, a step of contacting the sample with a fibronectin or oncofetal fibronectin binding partner bound to a solid support can be used to increase the relative concentration of the oncofetal fibronectin indicating molecule in the sample, thus facilitating detection of the oncofetal fibronectin indicating molecule using the mass measurement methods provided herein.

[0885] ii. Contact with a Fragmentation Compound

[0886] A sample can be contacted with a fragmentation compound. A fragmentation compound can be used to fragment an oncofetal fibronectin indicating molecule at specific sites (i.e., specifically fragment an oncofetal fibronectin indicating molecule), or can be used to fragment an oncofetal fibronectin indicating molecule at random sites (i.e., non-specifically fragment an oncofetal fibronectin indicating molecule), where random fragmentation refers to fragmentation where no particular site is more than two-fold more frequently cleaved than any other site, but random fragmentation does not require pure randomness in fragmentation.

[0887] A fragmentation compound can be a protein, peptide, oligonucleotide, or other compound that can be used to fragment proteins or nucleic acid molecules. In one embodiment, a fragmentation compound can be a protease or other compound that can be used to fragment an oncofetal fibronectin protein. Exemplary compounds for fragmenting oncofetal fibronectin protein include cathepsin D, trypsin, thermolysin, 2-nitro-5-thiocyanobenzoic acid (for S-cyanylation), *Achromobacter* protease 1, *S. aureus* V8 protease and hydroxylamine. In another embodiment, a fragmentation compound can be nuclease, ribozyme, DNAzyme, or other compound that can be used to fragment an oncofetal fibronectin encoding nucleic acid molecule or complement thereto. Exemplary compounds for fragmenting nucleic acid molecules include restriction endonucleases, hammerhead ribozymes and RNases.

[0888] Fragmentation methods can be performed prior to other steps of sample treatment, such as contacting the sample with a fibronectin or oncofetal fibronectin binding partner (e.g., a solid support to which a fibronectin or oncofetal fibronectin binding partner is immobilized), or conditioning the sample. Fragmentation methods can be performed subsequent to other steps of sample treatment. Fragmentation methods can coincide with other steps of sample treatment; for example, fragmentation methods can be performed while sample components are bound to a solid support.

[0889] a. Trypsin Proteolysis

[0890] In one embodiment, a sample can be treated with trypsin. When digested with trypsin, a human oncofetal fibronectin protein can yield proteolytic fragments that are 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa and/or 55 kDa in mass. Typically, each of these six trypsin fragments specifically binds to the antibody FDC-6. Thus, measurement of one or more trypsin fragments that are 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa or 55 kDa in mass and that can specifically bind to FDC-6 can indicate the presence of oncofetal fibronectin protein in a sample.

[0891] In an exemplary trypsin digest, trypsin fragments of a human oncofetal fibronectin protein can be 200 kDa, 120 kDa and/or 55 kDa, where each smaller fragment represents a product of further trypsin cleavage of a larger fragment. In another exemplary trypsin digest, trypsin fragments of oncofetal fibronectin protein can be 235 kDa, 160 kDa and/or 65 kDa, where each smaller fragment represents a product of further trypsin cleavage of a larger fragment.

[0892] In an exemplary method of oncofetal fibronectin protein detection, a sample can be contacted with a solid surface to which FDC-6 is immobilized, rinsed and then contacted with a dissociation solution that causes oncofetal fibronectin protein to no longer bind to FDC-6. The dissociation solution eluate can then be contacted by trypsin and the trypsin-treated eluate can be analyzed by MALDI-TOF mass spectrometry. A mass spectrum from such a treated sample having one or more measured masses of 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa or 55 kDa, or about 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa or 55 kDa can be identified as a sample containing oncofetal fibronectin protein. In another exemplary method, a sample can be contacted with trypsin and then can be contacted with a solid surface to which FDC-6 is immobilized. The solid surface can then be rinsed and then treated under conditions that

cause oncofetal fibronectin protein to no longer bind to FDC-6. A mass spectrum from such a treated sample having one or more measured masses of 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa or 55 kDa, or about 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa or 55 kDa can be identified as a sample containing oncofetal fibronectin protein. In another exemplary method, a sample can be contacted with a solid surface to which FDC-6 is immobilized and then the solid surface can be contacted with trypsin. The solid surface can then be rinsed and then treated under conditions that cause oncofetal fibronectin protein to no longer bind to FDC-6. A mass spectrum from such a treated sample having one or more measured masses of 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa or 55 kDa, or about 235 kDa, 200 kDa, 160 kDa, 120 kDa, 65 kDa or 55 kDa can be identified as a sample containing oncofetal fibronectin protein. In one exemplary case, a mass spectrum that identifies a sample as containing oncofetal fibronectin protein will contain all three 200 kDa, 120 kDa and 55 kDa masses. In another exemplary case, a mass spectrum that identifies a sample as containing oncofetal fibronectin protein will contain all three 235 kDa, 160 kDa and 65 kDa masses.

[0893] b. Cathepsin D Proteolysis

[0894] In another embodiment, a sample can be treated with cathepsin D. When digested with cathepsin D, oncofetal fibronectin protein can yield fragments that are 110 kDa and/or 85 kDa in mass. Typically these two cathepsin D fragments specifically bind to the antibody FDC-6. Thus, measurement of one or more cathepsin D fragments that are 110 kDa or 85 kDa in mass and that can specifically bind to FDC-6 can indicate the presence of oncofetal fibronectin protein in a sample.

[0895] In an exemplary method of oncofetal fibronectin protein detection, a sample can be contacted with a solid surface to which FDC-6 is immobilized, rinsed and then contacted with a dissociation solution that causes oncofetal fibronectin protein to no longer bind to FDC-6. The dissociation solution eluate can then be contacted by cathepsin D and the cathepsin D-treated eluate can be analyzed by MALDI-TOF mass spectrometry. A mass spectrum from such a treated sample having one or more measured masses of 110 kDa or 85 kDa, or about 110 kDa or 85 kDa can be identified as a sample containing oncofetal fibronectin protein. In other examples, the sample can be first contacted with cathepsin D and subsequently contacted with a solid support, or the sample can be contacted with a solid support and contacted with cathepsin D prior to eluting the sample from the solid support.

[0896] c. Thermolysin Proteolysis

[0897] In another embodiment, a sample can be treated with thermolysin. When digested with thermolysin, oncofetal fibronectin protein can yield fragments that are 120 kDa, 85 kDa and/or 35 kDa in mass. Typically the 120 kDa and 85 kDa can bind to the antibody BC-1. Thus, measurement of one or more thermolysin fragments that are 120 kDa or 85 kDa in mass and that can specifically bind to BC-1 can indicate the presence of oncofetal fibronectin protein in a sample. In an exemplary thermolysin digest, the 35 kDa and 85 kDa fragments represent products of further thermolysin cleavage of a larger fragment.

[0898] In an exemplary method of oncofetal fibronectin protein detection, a sample can be contacted with a solid

surface to which BC-1 is immobilized, rinsed and then contacted with a dissociation solution that causes oncofetal fibronectin protein to no longer bind to BC-1. The dissociation solution eluate can then be contacted by thermolysin and the thermolysin-treated eluate can be analyzed by MALDI-TOF mass spectrometry. A mass spectrum from such a treated sample having one or more measured masses of 120 kDa, 85 kDa or 35 kDa, or about 120 kDa, 85 kDa or 35 kDa can be identified as a sample containing oncofetal fibronectin protein. In other examples, the sample can be first contacted with thermolysin and subsequently contacted with a solid support, or the sample can be contacted with a solid support and contacted with thermolysin prior to eluting the sample from the solid support. In one exemplary case, a mass spectrum that identifies a sample as containing oncofetal fibronectin protein will contain all three 120 kDa, 85 kDa and 35 kDa masses.

[0899] d. *Acrhomobacter* Protease I Proteolysis

[0900] In another embodiment, a sample can be treated with *Acrhomobacter* protease I. When digested with *Acrhomobacter* protease I, oncofetal fibronectin protein can yield a fragment that is 14 kDa in mass. Typically, this 14 kDa fragment can bind to the antibody FDC-6. Thus, measurement of a *Acrhomobacter* protease I fragment that is 14 kDa and that can specifically bind to FDC-6 can indicate the presence of oncofetal fibronectin protein in a sample.

[0901] In an exemplary method of oncofetal fibronectin protein detection, a sample can be contacted with a solid surface to which FDC-6 is immobilized, rinsed and then contacted with a dissociation solution that causes oncofetal fibronectin protein to no longer bind to FDC-6. The dissociation solution eluate can then be contacted by *Acrhomobacter* protease I and the *Acrhomobacter* protease I-treated eluate can be analyzed by MALDI-TOF mass spectrometry. A mass spectrum from such a treated sample having a measured mass of 14 kDa or about 14 kDa can be identified as a sample containing oncofetal fibronectin protein. In other examples, the sample can be first contacted with *Acrhomobacter* protease I and subsequently contacted with a solid support, or the sample can be contacted with a solid support and contacted with *Acrhomobacter* protease I prior to eluting the sample from the solid support.

[0902] iii. Solid Support

[0903] A fibronectin or oncofetal fibronectin binding partner, a fragmentation compound, or both, can be immobilized on one or more solid supports for use in the methods provided herein. A solid support on which a binding partner or fragmentation compound can be immobilized can be any of a variety of supports to which the binding partner or fragmentation compound can bind to directly or can be bound using a linker. For example, a binding partner or fragmentation compound can be immobilized on a membrane such as a nitrocellulose membrane. A binding partner or fragmentation compound also can be immobilized on a microplate, such as a microplate coated with a compound for binding the binding partner. A binding partner or fragmentation compound also can be immobilized on a probe, pipette tip, or a conical, needle-shaped, or similarly shaped structure.

[0904] iv. Conditioning

[0905] An oncofetal fibronectin indicating molecule or fragment thereof, or an oncofetal fibronectin indicating

molecule or fragment thereof bound to a fibronectin or oncofetal fibronectin binding partner, can be optionally "conditioned". Conditioning is performed prior to mass spectrometric analysis and typically subsequent to or simultaneous with, one or more binding partner contacting or fragmentation steps. Conditioning can be performed, for example, in order to decrease the laser energy required for volatilization and/or to minimize unintended fragmentation. Conditioning can be performed before adding the oncofetal fibronectin indicating molecule, or fragment or complex thereof, to a mass spectrometry substrate. Conditioning can be performed while the oncofetal fibronectin indicating molecule, or fragment or complex thereof is bound or immobilized on the substrate. Conditioning can be performed after the oncofetal fibronectin indicating molecule, or fragment or complex thereof dissociates or is otherwise no longer bound to the substrate. Methods for conditioning are known in the art and include use of cation exchange resins and use of acetonitrile solutions.

[0906] v. Combinations of Sample Manipulation Steps

[0907] Sample manipulation steps that include different permutations of binding partners and fragmentation compounds can be employed. For example, a sample can be first contacted with an immobilized binding partner and then can be released from the solid support and contacted with an immobilized fragmentation compound. In another example, a sample can be first contacted with mobile or soluble fragmentation compound and then contacted with a solid support containing binding partner immobilized thereto. In another example, a sample can be first contacted with a solid support containing binding partner immobilized thereto and, while the sample is still exposed to the solid support, mobile or soluble fragmentation compound can be added to the sample. In another example, sample can first be contacted with a solid support containing binding partner immobilized thereto and then can be released from the solid support and contacted with a soluble or mobile fragmentation compound. In another example, a sample can be contacted with a solid support containing binding partner and fragmentation compound immobilized thereto. In another example, a sample can be first contacted with an immobilized fragmentation compound and then can be released from the solid support and contacted with an immobilized binding partner. Additional combinations also can be performed, as will be clear to one skilled in the art.

[0908] The sample manipulation steps prior to mass measurement can be in any order. For example, a sample can first be contacted with a fragmentation compound and then subsequently contacted with a binding partner. In another example, a sample can be first contacted with a binding partner and then subsequently contacted with a fragmentation compound. In another example, a sample can be contacted with a fragmentation compound while the sample is in contact with a binding partner. The binding partner, the fragmentation compound, or both, can be immobilized on a support. Where the fragmentation compound and binding partner are immobilized, they can be immobilized on the same or different solid supports. A solid support containing immobilized binding partner or immobilized fragmentation compound, or both, can serve as a substrate for mass spectrometric analysis.

[0909] In one example, a sample is first contacted by a first solid support to which a fibronectin or oncofetal fibronectin

binding partner, such as FDC-6, is immobilized, under conditions in which an oncofetal fibronectin indicating molecule can bind to the fibronectin or oncofetal fibronectin binding partner. Exemplary buffer conditions include 1% bovine serum albumin in phosphate-buffered saline (PBS; typically 10 mM phosphate, pH 7.4, 150 mM NaCl and 3 mM KCl). The contact between the sample and the first solid support can extend for any duration of time that results in binding of sufficient oncofetal fibronectin indicating molecule for carrying out subsequent steps such that one or more measurable fragments can be measured by mass spectrometry. For example, the first solid support can contact the sample for 10 minutes, 30 minutes, 1 hour, 2 hours, 5 hours, 10 hours, 24 hours, or about 10 minutes, 30 minutes, 1 hour, 2 hours, 5 hours, 10 hours, 24 hours, or more. The first solid support can then be separated from the sample and, in some embodiments, subjected to one or more washing steps. For example, the excess sample can be removed by washing the substrate with PBS and 0.05% Tween-20. The first solid support can then be exposed to an analyte release solution which causes bound sample compounds (including an oncofetal fibronectin indicating molecule, if present) to be released from the binding partner on the first solid support. Buffer conditions for releasing an oncofetal fibronectin indicating molecule from the binding partner generally include low pH solutions, high ionic strength solutions, or solutions containing a compound that displaces oncofetal fibronectin from the oncofetal fibronectin binding partner. Exemplary buffer conditions include buffers with a pH at or below 3.0 or about 3.0, buffers containing NaCl concentrations at or above about 1 M, or, buffers containing chaotropic agents such as 4 M NaSCN or 6 M urea. Released sample compounds (including oncofetal fibronectin, if present) can be, directly (e.g., immediately upon release from the solid support) or indirectly (e.g., after storage in a vessel, buffer exchange, freezing, centrifugation, filtration, etc.), added to a second solid support to which has been immobilized a fragmentation compound, such as trypsin. In one embodiment, the second solid support can serve as the substrate used for desorption of analytes to be measured by mass spectrometry. Released sample compounds (including an oncofetal fibronectin indicating molecule, if present) then are exposed to the fragmentation compound under conditions in which the fragmentation compound can cleave an oncofetal fibronectin indicating molecule in at least one site. The incubation of the released sample compounds (including oncofetal fibronectin, if present) and the second solid support can extend for any duration of time that results in fragmentation sufficient for carrying out subsequent steps such that one or more measurable fragments can be measured by mass spectrometry. For example, the second solid support can contact the released compounds for 10 seconds, 20 seconds, 30 seconds, 45 seconds, 1 minute, 2 minutes, 5 minutes, 10 minutes, 30 minutes, 1 hour, 2 hours, 5 hours, 10 hours, 24 hours, or about 10 seconds, 20 seconds, 30 seconds, 45 seconds, 1 minute, 2 minutes, 5 minutes, 10 minutes, 30 minutes, 1 hour, 2 hours, 5 hours, 10 hours, 24 hours, or more.

[0910] After fragmentation, the sample compounds (including oncofetal fibronectin fragments, if present) on the second solid support can be transferred to a substrate for desorption in mass spectrometry, or can be directly desorbed from the second solid support. Mass spectrometric methods can then be used to measure the charge to mass ratio of the

sample compounds and/or fragments (including oncofetal fibronectin fragments, if present), to detect the molecular weights of the sample compounds (including oncofetal fibronectin fragments, if present). Sample compound and fragment masses can then be used for further determinations of the presence of oncofetal fibronectin in a sample, as described elsewhere herein.

[0911] Exemplary binding partners, fragmentation compounds, solid supports to which binding partners or fragmentation compounds are immobilized, substrates for mass spectrometric analysis and methods including fragmentation, specific binding of analyte and mass spectrometric measurement of fragments and systems for carrying out one or more such steps are disclosed herein and known in the art (see, e.g., U.S. Pat. Nos. 6,498,039, 6,316,266, 6,093,541, 6,004,770, 5,955,729 and 5,719,060; and in U.S. Pat. App. Nos. 20030027216, 20020164818, 20020110904, 20020094566, 20020042075, 20010021535 and 20010019829). Any of a variety of methods of mass spectrometric analysis and analysis of mass spectrometric results known in the art or described herein can be used in conjunction with the above sample manipulation methods.

[0912] b. Substrate for Mass Spectrometry

[0913] The substrate for mass spectrometry can be any of a variety of materials. In particular, the substrate can be fabricated from virtually any insoluble or solid material that can be used in a particular mass spectrometry format.

[0914] MALDI substrates that can be used in the methods provided herein include those that are commercially available. For example, substrates can be metals such as gold, silicon, silver, copper, aluminum, or steel; non-metals such as glass or quartz; and polymers such as hydrocarbon polymers, polysilanes, PTFE, PTE, PE, PFA, perfluoroalkyals and methacrylates, polyethylene, polypropylene, polyamide, polyurethane, polyvinylidene difluoride, polyvinylidene fluoride, polysiloxanes optionally substituted with fluoroalkyl groups, perfluorodecyltrichlorosilanes, octadecyltrichlorosilanes, fluoropolymers, silicones, graphite, graphite filed polymers, or polysilanes. MALDI substrates can have any of a variety of shapes including beads, capillaries, flat surfaces (including flat surfaces containing pits such as an array of pits), microarrays, or pins (such as an array of pins). The substrate can be porous or non-porous, contain pits or wells and can have features (e.g., pits, wells, pins, etc.) organized in an array. MALDI substrates also can be coated with a monolayer, thin film, or thick film, using methods known in the art. A variety of compositions, shapes and coatings of substrates are commercially available and known in the art, as exemplified in U.S. Pat. App. No. 20030138823, U.S. Pat. Nos. 5,808,300, 6,265,715, 6,287,872 and 6,465,778.

[0915] The substrate can be the solid support to which the binding partner is immobilized, the solid support to which the fragmentation compound is immobilized, or the solid support to which the binding partner and the fragmentation compound are immobilized. The substrate also can have neither the binding partner nor the fragmentation compound immobilized thereto.

[0916] In some embodiments, the substrate can contain or have immobilized thereto, one or more energy absorbing compounds, such as matrix compounds, which can promote

analyte desorption and ionization. For example, an energy-absorbing compounds can be attached to a substrate by covalent attachment or physisorption, as is known in the art, and exemplified in U.S. Pat. No. 5,894,063. In another example, the substrate can have attached thereto a substance or compound to which an oncofetal fibronectin indicating molecule binds (including, e.g., a hydrophobic substance, ionic substance, or binding partner), where the substance or compound binds to an oncofetal fibronectin indicating molecule via a photolabile attachment or the substance or compound is energy absorbing and can promote analyte desorption and ionization. Such substrates are known in the art, as exemplified in U.S. Pat. No. 6,124,137.

[0917] c. Mass Spectrometric Analysis

[0918] Mass spectrometry can be used to detect presence of an oncofetal fibronectin indicating molecule. Mass spectrometric detection can be a direct detection of oncofetal fibronectin protein, a nucleic acid molecule encoding oncofetal fibronectin or complement thereto, an autoantibody for oncofetal fibronectin or a nucleic acid molecule encoding oncofetal fibronectin, or a fragment thereof, or can be a detection of a mass marker or other atom or molecule indicating presence of oncofetal fibronectin protein, nucleic acid molecule encoding oncofetal fibronectin or complement thereto, autoantibody therefor, or a fragment thereof. Mass spectrometry can be used following complex formation between a binding partner and oncofetal fibronectin protein, a nucleic acid molecule encoding oncofetal fibronectin or complement thereto, an autoantibody therefor, or a fragment thereof.

[0919] A variety of mass spectrometric detection formats are known to those skilled in the art. These include, for example, MALDI-TOF mass spectrometry, electrospray ionization mass spectrometry, inductively coupled plasma mass spectrometry, fast atom bombardment mass spectrometry, fourier transform mass spectrometry, electron impact mass spectrometry, chemical ionization mass spectrometry, ion cyclotron resonance mass spectrometry and combinations thereof, such methods for ionization and detection are known in the art, as exemplified in U.S. Pat. No. 6,657,191. Any known format can be adapted for detection of an oncofetal fibronectin indicating molecule.

[0920] i. Formation of Ions in the Gas Phase

[0921] In one step of mass spectrometry, gas phase ions are formed from sample material. Formation of gas phase ions of the sample can be accomplished using any of a variety of techniques. For example, a fragment of an oncofetal fibronectin indicating molecule can be desorbed and ionized using ultraviolet Matrix-Assisted Laser Desorption/Ionization, infrared Matrix-Assisted Laser Desorption/Ionization, electrospray, ion cyclotron resonance and/or inductively coupled plasma. In one example, if the sample material is sufficiently volatile, ions can be formed by electron impact (EI) or chemical ionization (CI) of the gas phase sample molecules. For solid samples, ions can be formed by desorption and ionization of sample molecules by bombardment with high energy photons or particles. Secondary ion mass spectrometry (SIMS), for example, uses keV ions to desorb and ionize sample material. In the SIMS process, a large amount of energy is deposited in the analyte molecules. As a result, fragile molecules will be fragmented. This fragmentation is not typically used when information

regarding the original composition of the sample, e.g., the molecular weight of sample molecules, will be lost.

[0922] In another example, for more labile or fragile molecules, other ionization methods now exist. The plasma desorption (PD) process results in the desorption of larger, more labile species—e.g., insulin and other protein molecules. Lasers can be used in a similar manner to induce desorption of biological or other labile molecules, as exemplified in Van Breeman et al., *Int. J. Mass Spectrom. Ion Phys.* 49:35-50 (1983); Tabet et al., *Anal. Chem.* 56:1662 (1984); Olthoff et al., *Anal. Instrument.* 16:93 (1987). The plasma or laser desorption and ionization of labile molecules relies on the deposition of little or no energy in the analyte molecules of interest. One use of lasers to desorb and ionize labile molecules intact is termed matrix assisted laser desorption ionization (MALDI) (see, e.g., Tanaka et al., *Rapid Commun. Mass Spectrom.* 2:151 (1988) and Karas et al., *Anal. Chem.* 60:2299 (1988)). In an example of the MALDI process, an analyte is dissolved in a solid, organic matrix. Laser light of a wavelength that is absorbed by the solid matrix but not by the analyte is used to excite the sample. The matrix is excited directly by the laser and the excited matrix sublimates into the gas phase carrying with it the analyte molecules. The analyte molecules then are ionized by proton, electron, or cation transfer from the matrix molecules to the analyte molecules. The MALDI process is typically used in conjunction with time-of-flight mass spectrometry (TOF-MS) and can be used to measure the molecular weights of proteins in excess of 100,000 daltons.

[0923] In another example, atmospheric pressure ionization (API) methods also can be used. Typically, analyte ions are produced from liquid solution at atmospheric pressure. One such method, is known as electrospray ionization (ESI) (see, e.g., Dole et al., *J. Chem. Phys.* 49:2240 (1968)). In the electrospray technique, analyte is dissolved in a liquid solution and sprayed from a needle. The spray is induced by the application of a potential difference between the needle and a counter electrode. The spray results in the formation of fine, charged droplets of solution containing analyte molecules. In the gas phase, the solvent evaporates leaving behind charged, gas phase, analyte ions. Very large ions can be formed by this method. Ions as large as 1 MDa have been detected by ESI in conjunction with mass spectrometry (ESMS). A variety of electrospray methods are known in the art, as exemplified by Wilm et al., *Int. J. Mass Spectrom. Ion Processes* 136:167 (1994), which teaches use of a small diameter needle in a method termed nano electrospray MS.

[0924] In one embodiment, when an oncofetal fibronectin indicating molecule or a fragment thereof is complexed with a fibronectin or oncofetal fibronectin binding partner that is immobilized on the substrate, the oncofetal fibronectin indicating molecule or fragment thereof is desorbed from the substrate. The desorption method can cause the oncofetal fibronectin indicating molecule or fragment thereof to desorb from the substrate without causing the binding partner to desorb from the substrate. In other embodiments, desorption methods can cause the oncofetal fibronectin indicating molecule or fragment thereof and its binding partner to desorb from the substrate.

[0925] For desorption methods such as MALDI, the sample is mixed with, or provided with, a matrix material that absorbs the laser light used in the MALDI method,

sufficiently for matrix and sample to desorb from the substrate. Many types of matrix materials are known in the art, including, for example, nicotinic acid, 3'-hydroxypicolinic acid, 2,5-dihydroxybenzoic acid, sinapinic acid, succinic acid, glycerol, urea and Tris-HCl, pH 7.3 or about 7.3. The matrix material can be applied simultaneously with the sample by, for example, mixing the matrix material with the sample. Alternatively, the matrix material can be present on a derivatized substrate prior to application of the sample, or introduced after application of the sample.

[0926] The gas phase ions of oncofetal fibronectin proteins, nucleic acids, fragments or complexes, then are detected using mass spectrometry.

[0927] ii. Detection

[0928] Gas phase ions can be detected by any of a variety of mass analyzers, such as a time-of-flight (TOF) mass analyzer. Mass analysis of gas phase ions can be performed using, for example, a magnetic or electrostatic analyzer, or both. Ions passing through a magnetic or electrostatic field will follow a curved path. In a magnetic field the curvature of the path will be indicative of the momentum-to-charge ratio of the ion. In an electrostatic field, the curvature of the path will be indicative of the energy-to-charge ratio of the ion. If magnetic and electrostatic analyzers are used consecutively, then the momentum-to-charge and energy-to-charge ratios of the ions can be detected and the mass of the ion can thereby be determined.

[0929] Another exemplary mass analyzer is a time-of-flight (TOF) mass analyzer. Typical TOF instruments take advantage of pulsed ionization as occurs with methods such as laser desorption methods including MALDI. In TOF methods, the ions are accelerated by a potential of 10-30 kV or about 10-30 kV and then allowed to drift down a field free region 1-2 m or about 1-2 m in length. Due to different velocities resulting from differences in mass, the ions arrive at the detector separated in time, permitting the mass to be determined. Reflector TOF mass analyzers also can be used in mass determination.

[0930] In another example, quadrupole mass analyzers can be used. In these mass analyzers, ions are accelerated electrically (5-15V) and passed along a long central axis of four rods arranged symmetrically. By applying combined DC and oscillating RF potentials, the ions drift along irregular flight paths along the rod axis. The DC/RF ratio is held constant and the absolute values of DC and RF are varied. Only ions with a particular m/z value have stable trajectories for a given value of DC and RF. If DC is set to 0, then all ions have stable trajectories. Quadrupole mass analyzers also can be used in conjunction with ion traps.

[0931] Ion cyclotron resonance (ICR) mass analyzers also can be used. In ICR mass analyzers, a range of rf components are used to excite a sample. By placing the ion trap within a superconducting magnet, the trapped ions undergo cyclotron gyration and are radially confined. The frequency of the cyclotron radiation is inversely proportional to the m/z ratio for an ion and directly proportional to the magnetic field. If an ion is excited at its natural cyclotron frequency, it moves to a higher energy level. The ion clouds then induce an image current at two or more detection electrodes. The resulting signal when subjected to Fourier transform analysis yields an extremely precise measure of ion cyclotron frequencies and hence m/z values and molecular weights.

[0932] iii. Use of Mass Spectrometry for Detecting Oncofetal Fibronectin in a Sample

[0933] Mass spectrometric methods can be used to detect the presence of an oncofetal fibronectin indicating molecule in a sample by detecting the molecular weight of a molecule, molecule fragment or atom indicative of the presence of an oncofetal fibronectin indicating molecule in a sample. In one embodiment, the molecule, molecule fragment or atom indicative of the presence of an oncofetal fibronectin indicating molecule in a sample includes an oncofetal fibronectin protein fragment, or a fragment of an oncofetal fibronectin encoding nucleic acid molecule or complement thereto. As described further hereinbelow, mass spectrometry can be used to detect the presence of particular regions in an oncofetal fibronectin indicating molecule. For example, a protein or nucleic acid fragment having a particular detected molecular weight can indicate the presence of EDA, EDB or IIICS in an oncofetal fibronectin protein or nucleic acid as described herein. Mass spectrometry also can be used to detect the presence of one or more post translational modifications in an oncofetal fibronectin protein. Measurement of an oncofetal fibronectin protein or nucleic acid fragment can be used to detect the presence of an oncofetal fibronectin protein or nucleic acid molecule in a sample and also to characterize structural elements present in the oncofetal fibronectin protein or nucleic acid molecule. Further in accordance with this embodiment, the method can include detecting the molecular weights of two or more oncofetal fibronectin protein fragments, or fragments of an oncofetal fibronectin encoding nucleic acid molecule or complement thereto; three or more oncofetal fibronectin protein fragments, or fragments of an oncofetal fibronectin encoding nucleic acid molecule or complement thereto; four or more oncofetal fibronectin protein fragments, or fragments of an oncofetal fibronectin encoding nucleic acid molecule or complement thereto; or more.

[0934] Mass spectrometry can be used to detect compounds according to their molecular weights. Detected signals can be used in measuring the molecular weights of compounds, and the measured molecular weights can be compared to one or more expected molecular weights, such as the molecular weight of an oncofetal fibronectin indicating molecule or fragment thereof. Detected mass spectrometric signals also can be compared to one or more references, such as reference mass spectra or reference signals of one or more mass spectra. A reference, such as a reference mass spectra or signal, can be a mass spectrum or signal corresponding to an oncofetal fibronectin indicating molecule or fragment thereof, or corresponding to a reference sample that contained an oncofetal fibronectin indicating molecule or fragment thereof. When detected mass spectrometric signals are compared to one or more references, calculation of the molecular weight of any particular peak is optional. For example, the mass spectrum from a sample can be compared to one or more references, where a detected signal that matches a reference signal can be identified as a detected signal indicative of the presence in the sample of an oncofetal fibronectin indicating molecule or fragment thereof.

[0935] In another embodiment, the molecule, molecule fragment or atom indicative of the presence of an oncofetal fibronectin indicating molecule in a sample will be detected by mass spectrometry after a binding event between an

oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner. A molecule, molecule fragment or atom that signals a binding event between an oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner can take any of a variety of forms. For example, a signalling molecule, molecule fragment or atom can be an oncofetal fibronectin protein, nucleic acid, autoantibody, or fragment thereof. In another example, a signalling molecule, molecule fragment or atom can be a fibronectin or oncofetal fibronectin binding partner or fragment thereof. In another example, a signalling molecule, molecule fragment or atom can be a detectable moiety bound to an oncofetal fibronectin protein, nucleic acid, autoantibody, binding partner, or fragment thereof. Such a detectable moiety can be any moiety that can be measured by a mass spectrometric method, typically a moiety having a mass or mass to charge ratio in a mass spectrometer that can be resolved from other detected masses or mass to charge ratios in the mass spectrum. Exemplary moieties include, for example, mass labels, as known in the art and described herein.

[0936] In one example of detecting a binding event between an oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner, a first oncofetal fibronectin binding partner can be fixed on a solid support and an oncofetal fibronectin indicating molecule-containing sample can be applied to the solid support to form a complex between the oncofetal fibronectin indicating molecule and the first oncofetal fibronectin binding partner. Following complex formation, a second fibronectin or oncofetal fibronectin binding partner can be applied to the solid support to form a complex between the oncofetal fibronectin indicating molecule and the first and second oncofetal fibronectin binding partners (i.e., a sandwich complex). The second fibronectin or oncofetal fibronectin binding partner can contain a detectable moiety, such as a photocleavable mass label. The detectible moiety can be measured directly using mass spectrometry, or the signal of the detectible moiety can be enhanced prior to measurement by increasing the number of detectible moieties present when an oncofetal fibronectin indicating molecule-oncofetal fibronectin binding partner complex forms.

[0937] a. Direct Measurement

[0938] Presence of an oncofetal fibronectin indicating molecule such as oncofetal fibronectin protein, a nucleic acid encoding oncofetal fibronectin or complement thereto, or an oncofetal fibronectin autoantibody, can be determined by directly detecting the oncofetal fibronectin indicating molecule, or a fragment thereof. For example, an oncofetal fibronectin indicating molecule can be detected using mass spectrometry by selectively binding an oncofetal fibronectin indicating molecule with a fibronectin or oncofetal fibronectin binding partner attached to a solid support, then subjecting the oncofetal fibronectin indicating molecule to conditions under which the oncofetal fibronectin indicating molecule or a fragment thereof no longer binds to the fibronectin or oncofetal fibronectin binding partner and then detecting oncofetal fibronectin indicating molecule or a fragment thereof using mass spectrometry. Conditions under which an oncofetal fibronectin indicating molecule or fragment thereof no longer binds to a binding partner can include, for example, low-binding buffer conditions, where a buffer contains substances such as high salt, low pH,

surfactants and denaturants; proteolysis; competitive displacement with a mass-distinguishable analog; or ionization or desorption, optionally coupled with fragmentation.

[0939] Presence of an oncofetal fibronectin indicating molecule also can be determined by directly detecting an oncofetal fibronectin indicating molecule in complex with a corresponding binding partner. For example, an oncofetal fibronectin indicating molecule/binding partner complex can be detected using mass spectrometry by selectively binding an oncofetal fibronectin indicating molecule with a fibronectin or oncofetal fibronectin binding partner attached to a solid support, then subjecting the complex to conditions under which the fibronectin or oncofetal fibronectin binding partner or a fragment thereof no longer is attached to solid substrate and then detecting oncofetal fibronectin indicating molecule/binding partner or a fragment thereof complex using mass spectrometry. Conditions under which a fibronectin or oncofetal fibronectin binding partner or fragment thereof no longer is attached to a solid substrate are known in the art and include, for example, ionization, desorption, proteolysis, or cleavage of a linker linking the binding partner to the substrate.

[0940] In yet another embodiment, presence of an oncofetal fibronectin indicating molecule or fragment thereof can be detected using a cleavable indicator atom or molecule, such as a cleavable mass label. A variety of cleavable mass labels can be selected and attached to a fibronectin or oncofetal fibronectin binding partner via appropriate cleavable linking chemistries. In accordance with the present method, upon binding or after binding of the binding partner to an oncofetal fibronectin indicating molecule or a fragment thereof, the mass label can be released and detected, to thereby indicate the presence of the oncofetal fibronectin indicating molecule or a particular fragment thereof.

[0941] A variety of mass labels and cleavable linking chemistries are known in the art, as exemplified in Pat. App. Nos. U.S. 200301947171, WO 98/31830, WO 98/26095, WO 97/27327 and U.S. Pat. Nos. 5,770,367, 6,558,902. Such mass labels can be detectable by mass spectrometry. Mass labels can include a vast array of different types of compounds including biopolymers and synthetic polymers. In one example of mass labels, biological monomer units can be used, either singly or in polymeric form, including amino acids, non-natural amino acids, nucleic acids, saccharides, carbohydrates, peptide mimics and nucleic acid mimics. Amino acids include those with simple aliphatic side chains (e.g., glycine, alanine, valine, leucine and isoleucine), amino acids with aromatic side chains (e.g., phenylalanine, tryptophan, tyrosine and histidine), amino acids with oxygen and sulfur containing side chains (e.g., serine, threonine, methionine and cysteine), amino acids with side chains containing carboxylic or amide groups (e.g., aspartic acid, glutamic acid, asparagine and glutamine) and amino acids with side chains containing strongly basic groups (e.g., lysine and arginine) and proline. Derivatives of the above described amino acids are monomer units. An amino acid derivative includes any compound that contains within its structure the basic amino acid core of an amino-substituted carboxylic acid, with representative examples including, but not limited to, azaserine, fluoroalanine, GABA, ornithine, norleucine and cycloserine. Polypeptides derived from the above described amino acids also can be used as monomer units. Representative examples include naturally occurring

and synthetic polypeptides with molecular weight above 500 Daltons or about 500 Daltons.

[0942] Representative examples of saccharides include ribose, arabinose, xylose, glucose, galactose and other sugar derivatives composed of chains from 2-7 carbons. Representative polysaccharides include combinations of the saccharide units listed above linked via a glycosidic bond. Mass labels also can be composed of nucleobase compounds, which include any moiety having within its structure a purine, a pyrimidine, a nucleic acid, nucleoside, nucleotide or derivative of any of these, such as a protected nucleobase, purine analog, pyrimidine analog, folinic acid analog, methyl phosphonate derivatives, phosphotriester derivatives, borano phosphate derivatives or phosphorothioate derivatives.

[0943] Mass labels also can include any organic or inorganic polymer that has a defined mass value, remains water soluble during bioassays and is detectable by mass spectrometry. Representative synthetic monomer units that can be used as mass units in polymeric form include polyethylene glycols, polyvinyl phenols, polymethyl methacrylates, polypropylene glycol, polypyrroles and derivatives thereof. The polymers can be composed of a single type of monomer unit or combinations of monomer units to create a mixed polymer. The sequence of the polymeric units within any one mass label is not critical; the total mass is the key feature of the label. For nonvolatile mass labels having mass below 500 Da or about 500 Da, usually significant ionic character is required; representative examples include polyethylene glycol oligomers of quaternary ammonium salts (e.g., $R-(O-CH_2-CH_2)_n-N(CH_3)_3^+/Cl^-$) and polyethylene glycol oligomers of carboxylic acids and salts (e.g., $R-(O-CH_2-CH_2)_n-CO_2^-Na^+$). Examples of involatile mass labels typically include small oligomers of polyethylene glycol and small peptides (natural or modified) less than 500 Da or about 500 Da in molecular weight. In these instances, as for all of the cases considered herein, mass analysis is not by electron attachment. Exemplary mass labels include a variety of nonvolatile and involatile organic compounds which are nonpolymeric. Representative examples of nonvolatile organic compounds include heme groups, dyes, organometallic compounds, steroids, fullerenes, retinoids, carotenoids and polyaromatic hydrocarbons.

[0944] A variety of cleavable linkers known in the art can be used to link the mass label to the fibronectin or oncofetal fibronectin binding partner. Different linker chemistries will confer cleavability under different specific physical or chemical conditions. Examples of conditions which serve to cleave various designs of linker include acid, base, oxidation, reduction, fluoride, thiol exchange, photolysis, ionization and enzymatic conditions. Examples of cleavable linkers that satisfy the general criteria for linkers listed above will be well known to those in the art and include, for example, those found in the catalog available from Pierce (Rockford, Ill.). Examples include: ethylene glycobis succinimidylsuccinate (EGS), an amine reactive cross-linking reagent which is cleavable by hydroxylamine (1 M at 37° C. for 3-6 hours); disuccinimidyl tartarate (DST) and sulfodST, which are amine reactive cross-linking reagents, cleavable by 0.015 M sodium periodate; bis[2-(succinimidyl-oxycarbonyloxy)ethyl]sulfone (BSOCOES) and sulfodBSOCOES, which are amine reactive cross-linking

reagents, cleavable by base (pH 11.6); 1,4-di-[3'-(2'-pyridyldithio(propionamido))butane (DPDPB), a pyridyldithiol crosslinker which is cleavable by thiol exchange or reduction; a silyl linking group that can be cleaved by fluoride or under acidic conditions; and a 3-, 4-, 5-, or 6-substituted-2-nitrobenzyloxy or 2-, 3-, 5- or 6-substituted-4-nitrobenzyloxy linking group that can be cleaved by a photon source (photolysis).

[0945] b. With Signal Enhancement

[0946] Upon formation of a complex between an oncofetal fibronectin indicating molecule and a fibronectin or oncofetal fibronectin binding partner, a second fibronectin or oncofetal fibronectin binding partner can be introduced that has attached thereto an amplifiable signalling nucleic acid or an amplifiable signalling nucleic acid attachment site. Amplifiable signalling nucleic acid attachment sites include a moiety such as biotin or poly-histidine, which can bind with specificity to a compound bound to an amplifiable signalling nucleic acid or to an intermediary binding partner that can bind with specificity to a compound bound to an amplifiable signalling nucleic acid, such as avidin to which a biotin-conjugated amplifiable signalling nucleic acid is bound.

[0947] Upon binding of the amplifiable signalling nucleic acid to the complex, the signalling nucleic acid can be amplified using methods known in the art. For example, a signalling nucleic acid can be amplified by transcription, PCR, ligase chain reaction, strand displacement amplification, rolling circle amplification, or other amplification reactions known in the art. In one example, the signalling nucleic acid is amplified using PCR.

[0948] In one embodiment, labeled anti-oncofetal fibronectin antibody (e.g., a biotinylated anti-oncofetal fibronectin antibody) is bound to a complex of oncofetal fibronectin protein and a fibronectin or oncofetal fibronectin binding partner, resulting in the formation of a sandwich complex. Streptavidin or avidin can then be added to the complex and can bind specifically to the labeled antibody. Biotinylated signalling nucleic acid, such as biotinylated linear DNA can then be added to the complex and can bind specifically to the streptavidin or avidin. The bound signalling nucleic acid can then be amplified in a multi-cycle (e.g., 30 cycles or about 30 cycles) polymerase chain reaction procedure. Generally, for example, each cycle includes a 1 minute denaturation step at 94° C., a 1 minute annealing step at 58° C. and a 1 minute primer extension step at 72° C., or about a 1 minute denaturation step at about 94° C., about a 1 minute annealing step at about 58° C. and about a 1 minute primer extension step at about 72° C. Amplification factors of about 10⁶ can be obtained.

[0949] Following amplification of the signalling nucleic acid, the signalling nucleic acid can be detected using mass spectrometry to indicate the presence of an oncofetal fibronectin indicating molecule in the sample.

[0950] iv. Detection of Regions of Oncofetal Fibronectin

[0951] Mass spectrometry can be used to detect the presence of particular regions in a fibronectin to thereby identify and/or characterize the oncofetal fibronectins and species thereof. Detection of the presence of particular regions in a fibronectin can serve a variety of purposes, including identifying the likely cell or tissue or organ source of the

oncofetal fibronectin, identifying the unlikely cell or tissue or organ source of the oncofetal fibronectin, or identifying a health problem associated with a particular form of oncofetal fibronectin. In one example, a fragment having a particular detected molecular weight can indicate the presence of an oncofetal fibronectin indication molecule, such as, but not limited to, EDA, EDB or IIICS in an oncofetal fibronectin protein or oncofetal fibronectin encoding nucleic acid molecule or complement thereto as described herein. Mass spectrometry also can be used to detect the presence of one or more post translational modifications in an oncofetal fibronectin protein. For example, a fragment having a particular detected molecular weight can indicate the presence of O-glycosylation at threonine 33 of the IIICS region of oncofetal fibronectin protein. Mass spectrometry also can be used to detect the presence of one or more splice regions of IIICS. For example, a fragment having a particular detected molecular weight can indicate the presence of the amino acid (aa) 90-120 splice region of an oncofetal fibronectin indicating molecule.

[0952] Detection of particular oncofetal fibronectin regions, IIICS splice regions and post-translational modifications, can serve to characterize an oncofetal fibronectin indicating molecule in the sample. For example, mass spectrometry can be used to characterize an oncofetal fibronectin indicating molecule present in a sample as containing or lacking all or part of the EDA, EDB or IIICS regions. Mass spectrometry can be used to characterize oncofetal fibronectin protein present in a sample as containing or lacking one or more particular post-translational modifications, such as O-glycosylation at threonine 33 of IIICS. Mass spectrometry can be used to characterize the oncofetal fibronectin indicating molecule present in a sample as containing or lacking a particular splice variant of IIICS such as V0, V64, V89, V95 or V120.

[0953] In using the mass spectrometry methods disclosed herein to characterize an oncofetal fibronectin indicating molecule, it is not necessary to use all detected molecular weights to characterize the oncofetal fibronectin indicating molecule. It also is not necessary to identify the composition of fragments whose mass is measured in order to characterize the oncofetal fibronectin. For example, measurement of one mass and comparison of that mass to one or more reference masses can indicate the presence of V120 of IIICS containing O-glycosylated threonine 33 in oncofetal fibronectin protein. In another example, measurement of two or more masses and comparison of those masses to two or more reference masses can indicate the presence of V120 of IIICS containing O-glycosylated threonine 33 in oncofetal fibronectin protein. In another example, measurement of two or more masses and comparison of those masses to two or more reference masses can indicate the presence of V120 of IIICS and EDB in an oncofetal fibronectin indicating molecule. Thus, provided herein are methods for characterizing an oncofetal fibronectin indicating molecule in a sample by identifying one or more masses indicative of a region of oncofetal fibronectin. In one embodiment, such methods can be performed by comparing one or more masses of a mass spectrum to masses of a reference mass spectrum (e.g., a mass spectrum collected from a known oncofetal fibronectin indicating molecule) or by comparing one or more masses of a mass spectrum to one or more reference masses (calculated or experimentally determined).

[0954] In another embodiment, one or more detected molecular weights can be used to distinguish or characterize different oncofetal fibronectin indicating molecules. For example, two oncofetal fibronectin indicating molecules having different compositions can yield different mass patterns when subjected to identical sample treatment and mass spectrometric methods. Accordingly, mass patterns for different oncofetal fibronectin indicating molecules can differ by one or more detected molecular weights and such different masses can be used to distinguish or characterize the different oncofetal fibronectin indicating molecules. A mass pattern can be indicative of a particular oncofetal fibronectin indicating molecule structure (e.g., an oncofetal fibronectin protein where the presence or absence of EDA, EDB and IIICS (and splice variants of IIICS), as well as the presence or absence of one or more post-translational modifications are known; or an oncofetal fibronectin nucleic acid or complement thereto known to encode or not encode a fibronectin protein containing EDA, EDB and IIICS (and splice variants thereof)) or complement thereto. Thus, provided herein are methods for characterizing an oncofetal fibronectin indicating molecule in a sample by identifying one or more masses indicative of a particular oncofetal fibronectin indicating molecule. In one embodiment, such methods can be performed by comparing one or more masses of a mass spectrum to masses of a reference mass spectrum (e.g., a mass spectrum collected from a known oncofetal fibronectin indicating molecule) or by comparing one or more masses of a mass spectrum to one or more reference masses (calculated or experimentally determined).

[0955] v. Quantitation of Oncofetal Fibronectin

[0956] Mass spectrometry can be used to determine the relative concentrations of components of a sample. Based on such methods, the amount of an oncofetal fibronectin indicating molecule or the amount of different types of oncofetal fibronectin indicating molecules in a sample, can be quantitated. In one example, a sample from a subject can have added thereto a known amount of a reference molecule, prior to mass spectrometric analysis. Comparison of the peak intensity of the reference molecule to the intensity of a peak representative of an oncofetal fibronectin indicating molecule can yield the ratio of oncofetal fibronectin indicating molecule present in the sample relative to the reference molecule; and knowledge of the amount of reference molecule present in the sample can then yield the concentration of the oncofetal fibronectin indicating molecule present in the sample. In one embodiment, a plurality of samples containing different concentrations of the reference molecule relative to the concentration of analytes in the sample, can be used to establish a standard curve against which the oncofetal fibronectin indicating molecule-associated peaks can be compared for determination of the amount of the oncofetal fibronectin indicating molecule present in the sample. Methods for quantitating analytes in a sample using mass spectrometry are known in the art, as exemplified in U.S. Pub. No. 20030027216.

[0957] 4. Detecting Nucleic Acid Molecules

[0958] Presence of oncofetal fibronectin in a subject can be indicated by detection of an oncofetal fibronectin-encoding nucleic acid in a sample. Oncofetal fibronectin-encoding nucleic acids include a nucleic acid molecule encoding oncofetal fibronectin such as transcribed mRNA encoding

oncofetal fibronectin. Encompassed within the scope of an oncofetal fibronectin-encoding nucleic acid are truncations, splice variants and fragments of an oncofetal fibronectin encoding nucleic acid. Full-length, truncations, variants and fragments of oncofetal fibronectin encoding nucleic acids can be detected using the methods provided herein. Typically, detection of an oncofetal fibronectin-encoding nucleic acid includes detection of a portion of a nucleic acid containing a sequence of nucleotides unique to oncofetal fibronectin or complement thereto.

[0959] a. Detection Methods

[0960] Amplified nucleic acid molecules from RT-PCR can be measured by a variety of methods for detecting nucleic acid molecules in a sample, including those provided herein and those known in the art. For example, nucleic acid molecules can be detected using gel electrophoresis, Southern blot analysis, Northern blot analysis, mass spectrometry, dot blot analysis, microarray or chip hybridization methods and other methods provided herein or known in the art. Oncofetal fibronectin-encoding nucleic acids or complements thereto can be detected directly or can be detected indirectly, for example, oncofetal fibronectin-associated nucleic acids or complements thereto can be detected indirectly by detecting nucleic acid molecules after nucleic acid amplification. A variety of amplification methods are known in the art and include, but are not limited to, PCR, rolling circle amplification, transcription, reverse transcription and reverse transcription PCR.

[0961] In accord with methods provided herein, detection of an amplified nucleic acid molecule corresponding to an oncofetal fibronectin-encoding nucleic acid molecule or complement thereto can indicate the presence of oncofetal fibronectin encoding nucleic acid molecule in a sample. Such an amplified nucleic acid molecule can contain all or a portion of the nucleotide sequence of the oncofetal fibronectin-encoding nucleic acid molecule or a complement thereto. For example, an amplified nucleic acid molecule can contain all or a portion of the nucleotide sequence encoding the EDA, EDB or IIICS regions of fibronectin or complement thereto. In one embodiment, an amplified nucleic acid molecule contains a nucleotide sequence encoding an amino acid region not present in non-oncofetal fibronectin protein, such as EDA, EDB or IIICS or complement thereto. In such an example, primers for nucleotide synthesis reactions can be designed to be complementary to a nucleotide sequence encoding an amino acid region not present in non-oncofetal fibronectin protein or complement thereto. Primers also can be designed to form amplified nucleic acid molecules which contain all or a portion of the EDA, EDB or IIICS encoding regions or complements thereto, if present, in the template nucleic acid molecule. Depending on the primer design, detection of the presence of any amplified nucleic acid, detection of an amplified nucleic acid having an expected size, or detection of an amplified nucleic acid containing an expected nucleotide sequence can indicate the presence of oncofetal fibronectin in a subject. For example, when the primers are complementary to regions that encode fibronectin portions occurring in oncofetal fibronectin and non-oncofetal fibronectin or complements thereto, but flanking a region that encodes a fibronectin portion occurring only in oncofetal fibronectin or complement thereto, the size of the fragment can indicate the presence or absence of oncofetal fibronectin in the sample, where a smaller fragment contains

nucleotides encoding non-oncofetal fibronectin and therefore is not indicative of oncofetal fibronectin in a sample and a larger fragment contains nucleotides encoding oncofetal fibronectin and therefore is indicative of oncofetal fibronectin in a sample. In another example, when one or more primers are complementary to regions encoding fibronectin portions that occur only in oncofetal fibronectin or complements thereto, presence of any amplicate can indicate the presence of oncofetal fibronectin in the sample.

[0962] b. Detection of RNA

[0963] In one embodiment, an oncofetal fibronectin-encoding nucleic acid is amplified using reverse transcription PCR (RT-PCR). Generally, reverse transcription PCR contains two types of reactions that can be performed in a single step or in separate steps. In the first type of reaction, RNA from a sample is reverse transcribed to complementary DNA (cDNA). In the second type of reaction, the cDNA is amplified using traditional PCR methods.

[0964] i. Reverse Transcription

[0965] Reverse transcription can be performed by contacting an RNA-containing sample with reverse transcriptase and primer, where the primer can be complementary to an oncofetal fibronectin-encoding RNA. Prior to contacting the sample with reverse transcriptase, the sample can be treated to remove DNA from the sample, using, for example, physical, chemical or enzymatic methods known in the art. Reverse transcription methods using a primer complementary to oncofetal fibronectin-encoding RNA can selectively yield oncofetal fibronectin-encoding cDNA.

[0966] The process of converting mRNA to cDNA typically uses a type of enzyme termed a reverse transcriptase, or a related enzyme with reverse transcriptase activity. A reverse transcriptase is an RNA-dependent DNA polymerase. All known reverse transcriptases require a primer to synthesize a DNA transcript from an RNA template. The reverse transcriptase can be obtained from eukaryotic cells which are infected with retrovirus, or from a number of plasmids which contain either a portion of, or the entire retrovirus genome. In addition, messenger RNA-like RNA which contains the RT gene can be obtained from retroviruses. Examples of sources for RT include Moloney murine leukemia virus (M-MLV); human T-cell leukemia virus type I (HTLV-I); bovine leukemia virus (BLV); Rous Sarcoma Virus (RSV); human immunodeficiency virus (HIV); yeast, including *Saccharomyces*, *Neurospora*, *Drosophila*; primates; and rodents. See, for example, Weiss et al., U.S. Pat. No. 4,663,290 (1987); Gerard, G. R., *DNA* 5(4):271-279 (1986); Kotewicz, M. L., et al., *Gene* 35:249-258 (1985); Tanese, N., et al., *Proc. Natl. Acad. Sci. USA* 82(15):4944-4948 (1985); Roth, M. J., et al., *J. Biol. Chem.* 260:9326-9335 (1985); Michel, F., et al., *Nature* 316:641-643 (1985); Akins, R. A., et al., *Cell* 47:505-516 (1986), *EMBO J.* 4:1267-1275 (1985); and Fawcett, D. F., *Cell* 47:1007-1015 (1986). M-MLV reverse transcriptase substantially lacking in RNase H activity has also been described. See, e.g., U.S. Pat. No. 5,244,797.

[0967] ii. cDNA Amplification

[0968] The methods also can use one or more DNA polymerases, including thermostable DNA polymerases, for amplifying the reverse-transcribed DNA. DNA polymerases can be isolated from natural or recombinant sources, by

techniques that are well-known in the art (See WO 92/06200, U.S. Pat. Nos. 5,455,170 and 5,466,591), from a variety of thermophilic bacteria that are available commercially (for example, from American Type Culture Collection, Rockville, Md.). A variety of sources of thermostable polymerases or the genes thereof for expression in recombinant systems are available, including the thermophilic bacteria *Thermus aquaticus*, *Thermus thermophilus*, *Thermococcus litoralis*, *Pyrococcus furiosus*, *Pyrococcus woosii* and other species of the *Pyrococcus* genus, *Bacillus sterothermophilus*, *Sulfolobus acidocaldarius*, *Thermoplasma acidophilum*, *Thermus flavus*, *Thermus ruber*, *Thermus brockianus*, *Thermotoga neapolitana*, *Thermotoga maritima* and other species of the *Thermotoga* genus and *Methanobacterium thermoautotrophicum* and mutants, variants or derivatives thereof. As an alternative to isolation, thermostable DNA polymerases are available commercially from, for example, Life Technologies, Inc. (Rockville, Md.), New England Biolabs (Beverly, Mass.), Finnzymes Oy (Espoo, Finland), Stratagene (La Jolla, Calif.), Boehringer Mannheim Biochemicals (Indianapolis, Ind.) and Perkin Elmer Cetus (Norwalk, Conn.). Exemplary thermostable DNA polymerases for use in the compositions and methods provided herein include, but are not limited to, Taq, Tne, Tma, Tli/VENT™, DEEPVENT™, Pfu, Pwo, Tfi or Tth DNA polymerases, or mutants or derivatives thereof. Taq DNA polymerase is commercially available, for example from Life Technologies, Inc. (Rockville, Md.), or can be isolated from its natural source, the thermophilic bacterium *Thermus aquaticus*, as described previously (U.S. Pat. Nos. 4,889,818 and 4,965,188). Tne DNA polymerase can be isolated from its natural source, the thermophilic bacterium *Thermotoga neapolitana* (See U.S. Pat. No. 5,939,301) and Tma DNA polymerase from its natural source, the thermophilic bacterium *Thermotoga maritima* (See U.S. Pat. No. 5,374,553). Methods for producing mutants and derivatives of thermophilic DNA polymerases, particularly of Tne and Tma polymerases, are disclosed in U.S. Pat. Nos. 5,948,614 and 6,015,668. Tfi, Tli/VENT™ and DEEPVENT™ are available commercially (e.g., from New England Biolabs; Beverly, Mass.), or can be produced as previously described (Bej, A. K. and Mahbubani, M. H., in: *PCR Technology: Current Innovations*, Griffin, H. G. and Griffin, A. M., eds., CRC Press, pp. 219-237 (1994) for Tli/VENT™; Flaman, J. M., et al., *Nucl. Acids Res.* 22 (15):3259-3260 (1994) for DEEPVENT™). Thermostable DNA polymerases can be added to the present compositions at a final concentration in solution of 0.1-200 units per milliliter, 0.1-50 units per milliliter, 0.1-40 units per milliliter, 0.1-36 units per milliliter, 0.1-34 units per milliliter, 0.1-32 units per milliliter, 0.1-30 units per milliliter, or 0.1-20 units per milliliter and most typically at a concentration of 20 units per milliliter, or about 0.1-200 units per milliliter, about 0.1-50 units per milliliter, about 0.1-40 units per milliliter, about 0.1-36 units per milliliter, about 0.1-34 units per milliliter, about 0.1-32 units per milliliter, about 0.1-30 units per milliliter, or about 0.1-20 units per milliliter and most typically at a concentration of about 20 units per milliliter.

[0969] In single-step reaction embodiments, the concentration of DNA polymerases can be determined as a ratio of the concentration of the enzymes having reverse transcriptase activity. Thus, compositions can have a unit ratio of the reverse transcriptase enzymes to the DNA polymerase enzymes ranging from 0.2:2 to 500:2, 0.5:2 to 250:2, or

greater than 3:2, or about 0.2:2 to about 500:2, about 0.5:2 to about 250:2, or greater than about 3:2.

[0970] iii. Additional Components

[0971] The compositions used herein include one or more nucleotides (e.g., deoxynucleoside triphosphates (dNTPs)). The nucleotide components of the present compositions serve as the "building blocks" for newly synthesized nucleic acids, being incorporated therein by the action of the reverse transcriptases or DNA polymerases. Examples of nucleotides for use in the present compositions include dUTP, dATP, dTTP, dCTP, dGTP, dITP, 7-deaza-dGTP, α -thio-dATP, α -thio-dTTP, α -thio-dGTP, α -thio-dCTP or derivatives thereof, all of which are available commercially from sources including Life Technologies, Inc. (Rockville, Md.), New England BioLabs (Beverly, Mass.) and Sigma Chemical Company (Saint Louis, Mo.). The dNTPs can be unlabeled, or they can be detectably labeled by coupling them by methods known in the art, such as, for example, with mass labels detectable by mass spectrometry, spectroscopically detectable labels, magnetic beads, radioisotopes (e.g., ^3H , ^{14}C , ^{32}P or ^{35}S), vitamins (e.g., biotin), fluorescent moieties (e.g., fluorescein, rhodamine, Texas Red, or phycoerythrin), chemiluminescent labels and dioxigenin. Labeled dNTPs also can be obtained commercially, for example from Life Technologies, Inc. (Rockville, Md.) or Sigma Chemical Company (Saint Louis, Mo.). The dNTPs can be added, for example, to give a working concentration of each dNTP of 10-1000 micromolar, 10-500 micromolar, 10-250 micromolar, 10-100 micromolar, or 100 micromolar, or about 10-1000 micromolar, about 10-500 micromolar, about 10-250 micromolar, about 10-100 micromolar, or about 100 micromolar.

[0972] In addition to nucleotides, the compositions include one or more primers which facilitate the synthesis of a first DNA molecule complementary to all or a portion of an RNA template (e.g., a single-stranded cDNA molecule). Primers also can be used to synthesize a DNA molecule complementary to all or a portion of the first DNA molecule, thereby forming a double-stranded cDNA molecule. Additionally, these primers can be used in amplifying nucleic acid molecules in accordance with the methods known in the art or provided herein. Such primers include, but are not limited to, target-specific primers (which are typically gene-specific primers such as oncofetal fibronectin-specific primers), oligo(dT) primers, random primers or arbitrary primers. Methods directed to specific detection of oncofetal fibronectin-encoding mRNA in a sample can use primers that bind specifically to oncofetal fibronectin-encoding mRNA and a reverse transcriptase to thereby selectively reverse transcribe oncofetal fibronectin-encoding mRNA to form oncofetal fibronectin-encoding cDNA.

[0973] Compositions for performing RT-PCR also can include one or more RNase inhibitors. Since RNA is the substrate of the reverse transcription reaction, fidelity of the RNA in the sample can be important in detecting RNA encoding oncofetal fibronectin. Any of a variety of known RNase inhibitors can be used in the RT-PCR methods provided herein, including, for example, human placental RNase inhibitor.

[0974] iv. Nucleic Acid Synthesis Steps

[0975] The reverse transcription reaction is performed by adding to an RNA sample reverse transcriptase, primer and

all four deoxynucleoside triphosphates under the appropriate reaction conditions. Typically, the reverse transcription reaction is performed under conditions that prevent or reduce degradation of the mRNA, which could result in incomplete cDNA synthesis. Exemplary reverse transcription reaction conditions include 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3 mM MgCl_2 , 10 mM dithiothreitol, 0.5 mM each dATP, dGTP and dTTP, 0.5 mM [^3H]dCTP (200 cpm/pmol), 50 $\mu\text{g/ml}$ (dT) $_{12-18}$, 20 $\mu\text{g/ml}$ 2.3 kb RNA and 4,000 units/ml RT and incubated at 37° C.

[0976] In one embodiment, DNA is removed from a sample prior to the reverse transcription reaction. DNA can be removed from a sample by any of a variety of methods, including physical, chemical and enzymatic methods. For example, DNA can be removed from a sample using a DNase, such as DNase I. DNA also can be removed by extraction using acid phenol:chloroform followed by precipitation of RNA using, for example, 0.5 M ammonium acetate and ethanol. DNA also can be separated from RNA by precipitation methods; for example, RNA can be precipitated from a sample using 7.5 M LiCl while the DNA remains in the supernatant, which can be discarded.

[0977] In another embodiment, DNA is not removed prior to reverse transcription. This embodiment can include a step of separating reverse transcribed DNA from other DNA in a sample prior to PCR or other amplification methods. For example, reverse transcribed DNA can be separated based on sequence composition, including a sequence specific to an oncofetal fibronectin-encoding sequence, or a sequence specific to RNA, such as a polyA sequence. In other embodiments, DNA is not removed prior to amplification, but PCR primers are designed to selectively amplify reverse transcribed DNA. Exemplary selective primers include primers that bind with sequence specificity to an oncofetal fibronectin-encoding sequence and primers that bind with sequence specificity to DNA products of reverse transcription, including DNA containing a poly(dT) sequence.

[0978] The RNA-DNA hybrid that results from reverse transcription can be subsequently treated, for example, with alkali or RNase H to selectively hydrolyze the RNA to leave cDNA that can be converted to double-stranded form in a second DNA amplification reaction catalyzed by reverse transcriptase or other DNA polymerase. See Old, R. W., et al., *Principals of Gene Manipulation, second edition, Studies in Microbiology*, Vol. 2, University of California Press, p. 26 (1981).

[0979] In the "uncoupled" RT-PCR procedure (e.g., two-step RT-PCR), reverse transcription is performed as an independent step using buffer conditions optimal for reverse transcriptase activity, such as 50 mM Tris-HCl (pH 8.3), 50 mM KCl, 10 mM MgCl_2 , 15 mM dithiothreitol, 0.1 mg/ml actinomycin D, at 37° C. Following cDNA synthesis, the reaction can be diluted to decrease MgCl_2 and deoxyribonucleotide triphosphate (dNTP) concentrations to conditions acceptable for Taq DNA Polymerase activity and PCR can be performed according to standard conditions (see, for example, U.S. Pat. Nos. 4,683,195 and 4,683,202).

[0980] In "coupled" RT-PCR methods, a common or compromised buffer is used for reverse transcriptase and Taq DNA Polymerase activities. In one version, the annealing of reverse primer is a separate step preceding the addition of enzymes, which then are added to the single reaction vessel.

In another version, the reverse transcriptase activity is a component of the thermostable Tth DNA polymerase. Annealing and cDNA synthesis can be performed in the presence of Mn^{++} , then PCR can be performed in the presence of Mg^{++} after the removal of Mn^{++} by a chelating agent.

[0981] A "continuous" method (e.g., one-step RT-PCR) can be used which integrates annealing, reverse transcription and PCR steps into a single continuous reaction that avoids the opening of the reaction tube for component or enzyme addition. Continuous RT-PCR can be performed as a single enzyme system using the reverse transcriptase activity of thermostable Taq DNA Polymerase and Tth polymerase, or as a two-enzyme system using AMV-RT and Taq DNA Polymerase where the initial temperature is 65° C.

[0982] U.S. Pat. Pub. No. 20030113712 describes compositions and methods useful for one-step/one-tube RT-PCR, using M-MLV-RT, or its RNase H-deficient derivatives, in combination with one or more DNA polymerases. The methods can be performed in the presence of sulfur-containing molecules or acetate-containing molecules (or combinations of sulfur-containing molecules and acetate-containing molecules) to relieve the inhibition of PCR when using compositions containing two or more enzymes having reverse transcriptase activity.

[0983] In the RT-PCR reaction, the reaction mixtures can be incubated at a temperature sufficient to synthesize a DNA molecule complementary to all or a portion of the RNA template. Such conditions will depend on the enzyme used and can range from 20° C. to 75° C., 35° C. to 60° C., or 45° C. to 55° C., or about 20° C. to about 75° C., about 35° C. to about 60° C., or about 45° C. to about 55° C. After the reverse transcription reaction, the reaction can be incubated at a temperature sufficient to amplify the synthesized DNA molecule. In one embodiment, the amplification is accomplished via one or more polymerase chain reactions (PCRs). Conditions for amplification can include thermocycling, such as alternating heating and cooling of the mixture sufficient to amplify the DNA molecule and which can include alternating from a first temperature range of from 90° C. to 100° C., to a second temperature range of from 45° C. to 75° C., 50° C. to 75° C., 55° C. to 75° C., or 65° C. to 75° C., or from a first temperature range from about 90° C. to about 100° C., to a second temperature range of from about 45° C. to about 75° C., about 50° C. to about 75° C., about 55° C. to about 75° C., or about 65° C. to about 75° C. The thermocycling can be performed any number of times, typically from 5 to 80 times or about 5 to about 80 times, greater than 10 times or about 10 times, or greater than 20 times or about 20 times.

[0984] U.S. Pat. Pub. No. 20030157550 provides methods for using oligonucleotide-immobilized microplates to which oligonucleotides are securely immobilized and which can be subjected to thermal cycles of PCR (PCR microplates), capturing of mRNA and reverse transcription-polymerase chain reaction (RT-PCR) can be conducted on the same plates. Using PCR microplates such as those made of polypropylene, polyolefin, or polycarbonate, because of their fluorescent characteristics, immobilized oligonucleotide, hybridized mRNA and synthesized cDNA can be quantitated fluorometrically by using nucleic acid stain or with the aid of a protein by producing fluorescence or

chemiluminescence. The PCR microplates also can capture mRNA from crude cell lysates without purification of RNA or mRNA.

[0985] Also provided herein are kits for carrying out the reverse transcription PCR methods described herein. Such kits can include a carrier being compartmentalized to receive a close confinement therein, one or more containers, such as vials and tubes, each of the containers including one of the separate elements of the method used to prepare cDNA from RNA. For example, there can be provided a containers containing reverse transcriptase having DNA polymerase activity and substantially no RNase H activity, in solution. Further containers can contain buffers, substrates for DNA synthesis such as the deoxynucleoside triphosphate, oligo(dT) primer and control RNA for use as a standard.

[0986] v. Detection

[0987] Amplified nucleic acid molecules from RT-PCR can be measured by a variety of methods for detecting nucleic acid molecules in a sample such as mass spectrometry or chip hybridization and any of a variety of detection results can indicate the presence or absence of oncofetal fibronectin encoding nucleic acid molecule in a sample, such as the presence or absence of an amplified nucleic acid molecule, or the size of the detected nucleic acid molecule. Presence of RNA encoding oncofetal fibronectin also can be quantitated using a variety of methods such as relative RT-PCR, competitive RT-PCR, comparative RT-PCR and real time RT-PCR.

[0988] a. DNA Detection Methods

[0989] Amplified nucleic acid molecules from RT-PCR can be measured by a variety of methods for detecting nucleic acid in a sample, including those provided herein or those known in the art. For example, DNA can be detected using gel electrophoresis, Southern blot analysis, mass spectrometry, dot blot analysis and chip hybridization methods.

[0990] In accordance with the methods provided herein, detection of a nucleic acid molecule such as an amplified deoxyribonucleic acid molecule corresponding to an oncofetal fibronectin-encoding nucleic acid molecule or complement thereto can indicate the presence of an oncofetal fibronectin encoding nucleic acid molecule in a sample. Such an amplified nucleic acid molecule can contain all or a portion of the nucleic acid molecule (or complement thereof) encoding the oncofetal fibronectin-encoding nucleic acid molecule. For example, an amplified nucleic acid molecule detected using the methods provided herein can contain all or a portion of the nucleic acid molecule encoding the EDA, EDB or IIICS regions of fibronectin or complement thereto. An amplified nucleic acid molecule also can contain a nucleotide sequence encoding an amino acid region not present in a non-oncofetal fibronectin protein or complement thereto. Primers used for reverse transcription or for amplification for nucleotide synthesis reactions can be designed to be complementary to a nucleic acid molecule encoding an amino acid region not present in non-oncofetal fibronectin protein or complement thereto. Primers also can be designed to form amplified nucleic acid molecules which contain all or a portion of the EDA, EDB or IIICS encoding regions or complement thereto, if present, in the template nucleic acid molecule. According to the primer design, detection of the presence of any amplified

nucleic acid, detection of an amplified nucleic acid having a particular size, or detection of an amplified nucleic acid containing a particular nucleotide sequence, can indicate the presence of oncofetal fibronectin in a subject.

[0991] b. Quantitation

[0992] Presence of RNA encoding oncofetal fibronectin can be quantitated using any of a variety of methods known in the art including relative RT-PCR, competitive RT-PCR, comparative RT-PCR and real time RT-PCR. In relative RT-PCR, amplification of the target nucleic acid molecule (i.e., mRNA encoding oncofetal fibronectin) is compared to amplification of a control sequence, such as the sequence of a housekeeping gene such as β -actin mRNA, GAPDH mRNA, or 18S rRNA. This method is performed for two or more samples in order to normalize the signal between samples, thus permitting quantitative calculation of the amount of the target nucleic acid molecule, such as an oncofetal fibronectin encoding nucleic acid molecule or complement thereto.

[0993] Competitive RT-PCR can use reverse transcription of a target nucleic acid molecule, followed by simultaneous PCR amplification of a target nucleic acid molecule and a competitive template for the target nucleic acid molecule, where the competitive template is a designed mutant of the target nucleic acid molecule. Exemplary mutants can arise from point mutations, insertions or deletions. The mutants can be detected as different from the wild type sequences; for example, the mutant can have a deletion and thus provide a shorter nucleic acid molecule than the wild type target nucleic acid molecule. Different amounts of competitive mutant template can be used in several samples in order to construct a standard curve used to quantify the amount of target nucleic acid molecule present. An exemplary competitive method is standardized RT-PCR, which uses standardized, competitor templates to allow for comparison between experiments. U.S. Pat. No. 5,876,978, for example, provides a standardized competitive RT-PCR process including reverse transcription of at least a target nucleic acid molecule and a housekeeping gene, followed by simultaneous PCR amplification of a target nucleic acid molecule, a housekeeping gene and competitive templates for each of these nucleic acids. The method simultaneously uses primers for a target nucleic acid molecule, primers for a housekeeping gene and two internal standard competitive templates that include designed mutants of the target nucleic acid molecule and housekeeping gene. Such methods also can be performed in multiplexed format, as described in WO 03/083051.

[0994] Another method for quantitating RT-PCR results is comparative RT-PCR. In this method, reverse transcription on two or more target nucleic acid molecule samples is performed using a unique probe for each sample, where the uniqueness is not in the RNA-recognition portion of the probe, resulting in a unique cDNA for every sample. After reverse transcription, the different target nucleic acid molecule sequences are combined and PCR amplification is performed where the different cDNAs compete with each other (as with competitive RT-PCR, described herein or known in the art). The relative amounts of target nucleic acid molecule in each sample can then be compared by determining the relative amount of each unique cDNA that was amplified.

[0995] RT-PCR also can be quantitated after every amplification cycle using "real time" RT-PCR. Real time RT-PCR is based on the ability to detect formation of double-stranded DNA. One exemplary method for real time RT-PCR quantitation includes use of probe nucleotides at least partially complementary to the target nucleic acid molecule and having a fluorescent dye and a quenching dye. The probe nucleotide is designed such that when the probe nucleotide does not interact (e.g., hybridize) with the target nucleic acid molecule, the fluorescent dye is quenched and little or no fluorescent signal arises and when the probe nucleotide does interact with the target nucleic acid molecule, the fluorescent dye is no longer quenched and fluorescent signal arises. As the target nucleic acid molecule is amplified, the probe nucleotide interacts with (e.g., hybridizes to) the target nucleic acid molecule, resulting in a fluorescent signal that increases proportionally to the amount of target nucleic acid molecule present. A variety of probe nucleotides are available, including TaqMan probes, molecular beacons and scorpions, as described, for example, in Bustin, *J. Mol. Endocrinology* 29:23-39 (2002) and Thelwell et al., *Nucleic Acids Res.* 28:3752-61 (2000). Real-time quantitation also can be performed using a dye that emits only upon binding to double-stranded DNA, such as SYBR green (exemplary kit made by Qiagen, Inc. (Valencia, Calif.); see, for example, Moullisseaux et al., *J. Virol. Methods* 111(2):121-127 (2003)). Additional reagents such as peptide nucleic acids (PNAs) or minor groove binding dyes also can be used (see Bustin, *J. Mol. Endocrinology* 29:23-39 (2002)). Results from real time RT-PCR measurements can be compared to a standardized curve describing signal intensity as a function of nucleic acid concentration, or can be compared to the signal intensity of one or more reference nucleic acids. These methods, such as those using a fluorescent dye/fluorescence quencher probe nucleotides, also can be used in multiplexing reactions.

[0996] c. Detection of Regions of Oncofetal Fibronectin

[0997] Nucleic acid detection methods can be used to detect the presence of particular regions in oncofetal fibronectin. Detection of the presence of particular regions in oncofetal fibronectin can serve a variety of purposes, including identifying the likely cell or tissue or organ source of the oncofetal fibronectin, identifying the unlikely cell or tissue or organ source of the oncofetal fibronectin, or identifying a health problem associated with a particular form of oncofetal fibronectin. In one example, detection of a particular nucleic acid molecule can indicate the presence of EDA, EDB or IIICS (or a particular splice variant thereof) in an oncofetal fibronectin protein. Nucleic acid detection methods also can be used to detect the presence of one or more splice regions of IIICS. For example, detection of a particular nucleic acid molecule can indicate the presence of the amino acid (aa) 90-120 splice region of oncofetal fibronectin.

[0998] Detection of particular oncofetal fibronectin regions and IIICS splice regions can serve to characterize the oncofetal fibronectin in the sample. For example, nucleic acid detection methods can be used to characterize oncofetal fibronectin present in a sample as containing or lacking the EDA, EDB or IIICS regions. Nucleic acid detection methods can be used to characterize oncofetal fibronectin present in a sample as containing or lacking a particular splice variant of IIICS such as V0, V64, V89, V95 or V120.

[0999] A variety of methods disclosed herein or known in the art can be used to characterize one or more oncofetal fibronectin nucleic acid molecules or complements thereto present in a sample, including but not limited to gel electrophoresis, mass spectrometry, Southern blot analysis, Northern blot analysis, dot blot analysis, mass spectrometry and chip array hybridization. In one embodiment, an oncofetal fibronectin encoding nucleic acid molecule or complement thereto can be characterized according to the molecular weight of the oncofetal fibronectin nucleic acid molecule or fragment thereof. For example, a nucleic acid molecule encoding oncofetal fibronectin containing EDA, EDB and IIICS V120 or complement thereto can have a greater mass than an oncofetal fibronectin encoding only EDB and IIICS V120 or complement thereto; thus, measurement of mass of the nucleic acid molecules can be used to characterize a sample as containing an oncofetal fibronectin nucleic acid molecule encoding EDA, EDB and IIICS V120 or complement thereto, or encoding EDB and IIICS V120 or complement thereto, or both.

[1000] In another embodiment, an oncofetal fibronectin encoding nucleic acid molecule or complement thereto can be characterized by hybridization of a sample nucleic acid molecule to a probe oligonucleotide. For example, a nucleic acid molecule encoding oncofetal fibronectin containing EDA, EDB and IIICS V120 can hybridize to oligonucleotide probes complementary to the EDA, EDB and IIICS V120 splice regions, while a nucleic acid molecule encoding oncofetal fibronectin containing EDB and IIICS V120 can hybridize to oligonucleotide probes complementary to the EDB and IIICS V120 splice regions, but not to an oligonucleotide probe complementary to the EDA splice region.

[1001] In using the nucleic acid detection methods disclosed herein to characterize an oncofetal fibronectin encoding nucleic acid molecule, one or more nucleic acid molecules such as probes or primers can be used to characterize the oncofetal fibronectin encoding nucleic acid molecule. For example, two or more nucleic acid molecules complementary to different oncofetal fibronectin regions or complements thereto can be used to distinguish or characterize an oncofetal fibronectin encoding nucleic acid molecule or complement thereto. Thus, provided herein are methods for characterizing oncofetal encoding fibronectin or complement thereto in a sample by identifying one or more nucleic acids indicative of a region of oncofetal fibronectin or indicative of a particular oncofetal fibronectin variant. Depending on the oncofetal fibronectin-encoding nucleic acid molecules or complements thereto present in the sample, some, but not all probe or primer nucleic acid molecules can be detected and/or amplified in characterizing the oncofetal fibronectin encoding nucleic acid molecule or complement thereto. For example, when a sample contains an oncofetal fibronectin-encoding nucleic acid molecule containing a nucleotide sequence that encodes EDA but not EDB, probes or primers complementary to both and to an EDA-encoding nucleic acid molecule and an EDB-encoding nucleic acid molecule can be added to the sample, but only the probes or primers complementary to the EDA-encoding nucleic acid molecule will be amplified and/or detected. It is possible, but not necessary, to identify the nucleic acid molecules or nucleotide compositions of the oncofetal fibronectin-encoding nucleic acid molecules or complements thereto of a sample in order to characterize the oncofetal fibronectin encoding nucleic acid molecule or

complement thereto. For example, two oncofetal fibronectin-encoding molecules or complements thereto having different nucleotide compositions (e.g., one encodes EDA and EDB while another encodes EDB but not EDA) can yield different bands on a gel or blot when subjected to identical sample treatment and nucleic acid detection methods, thus permitting characterization of one or both oncofetal fibronectin-encoding molecules or complements thereto without determining the nucleotide sequence or composition of the molecules. In one embodiment, such methods can be performed by comparing measurement of one or more nucleic acid molecules of a sample to measurement of nucleic acid molecules of a reference (e.g., a known oncofetal fibronectin molecule) or by comparing one or more nucleic acid molecules to one or more reference nucleic acid molecules (calculated or experimentally determined).

[1002] A variety of methods for using oligonucleotides to detect regions of oncofetal fibronectin in a sample are known in the art, including methods for using oligonucleotides to detect regions of oncofetal fibronectin or complements thereto in a tumorous tissue sample. For example, oligonucleotides can be used to detect the presence of mRNA encoding EDA, EDB and V0, V64 and V89 of IIICS in thyroid papillary carcinomas (Takano et al., *J. Endocrinol. Invest.* 22:18-22 (1999)). In another example, oligonucleotides can be used to detect the presence of EDA and EDB in hepatocellular carcinoma (Oyama et al., *Cancer Res.* 53:2005-2011 (1993)).

[1003] 5. Detection of Autoantibodies to Oncofetal Fibronectin

[1004] Detection of autoantibodies reactive against oncofetal fibronectin or a nucleic acid molecule encoding oncofetal fibronectin or a fragment thereof can indicate the presence of oncofetal fibronectin in a subject. Detection of autoantibodies reactive against oncofetal fibronectin or a nucleic acid molecule encoding oncofetal fibronectin or a fragment thereof also can indicate a variety of health conditions and/or health problems including overall health, cancer, or pregnancy-related conditions. Additionally, the monitoring of autoantibody levels can be used to stage the progression of the disease, to determine a disease or disorder outcome, to determine the likelihood of success of a particular treatment of the disease or disorder and/or to determine a treatment regimen. The detection of autoantibodies in a sample such as a serum sample from a subject can be accomplished by any of a number of methods. Any of a variety of methods known for detecting antibodies, including those provided herein, can be used to detect autoantibodies to oncofetal fibronectin or a nucleic acid molecule encoding oncofetal fibronectin. Such methods include immunoassays which include, but are not limited to, assay systems using techniques such as Western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, dipstick immunoassays, lateral flow immunoassays, vertical flow immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, flow cytometry assays and agglutination assays such as latex bead agglutination assays. Methods and compositions for detecting autoantibodies in a sample are known in the art, as

exemplified in U.S. Pat. No. 6,788,128, U.S. Pub. Nos. 20030232399 and 20040048320 and WO 03/020115.

[1005] An exemplary immunoassay is carried out by contacting a sample such as a serum sample of a subject with oncofetal fibronectin protein or a fragment thereof under conditions in which a specific antigen-antibody binding complex can form and detecting or measuring the amount of any complex. The presence of any anti-oncofetal fibronectin autoantibodies can indicate the presence of oncofetal fibronectin in a subject, or the presence of an oncofetal fibronectin associated disease or disorder in the subject. In another example, the levels of autoantibodies in a sample can be compared to the levels present in an analogous sample from a subject not having the disease or disorder, or to the levels present in a population of subjects not having the disease or disorder. The presence of anti-oncofetal fibronectin autoantibody levels higher than levels in subjects not having the disease or disorder can indicate an oncofetal fibronectin positive result and can indicate the presence of an oncofetal fibronectin associated disease or disorder in the subject.

[1006] In another exemplary immunoassay a sample is contacted with an anti-(human antibody) antibody, such as an anti-human IgG antibody, or a fragment thereof under conditions in which a specific antigen-antibody binding complex can form and detecting or measuring the amount of any complex. The presence of any anti-oncofetal fibronectin autoantibodies can indicate the presence of oncofetal fibronectin or the presence of an oncofetal fibronectin associated disease or disorder in the subject. In another example, the levels of autoantibodies in a sample can be compared to the levels present in an analogous sample from a subject not having the disease or disorder, or to the levels present in a population of subjects not having the disease or disorder. The presence of anti-oncofetal fibronectin autoantibody levels higher than levels in subjects not having the disease or disorder can indicate an oncofetal fibronectin positive result and can indicate the presence of an oncofetal fibronectin associated disease or disorder in the subject.

[1007] The immunoassays can be performed in a variety of ways. For example, oncofetal fibronectin protein or oncofetal fibronectin encoding nucleic acid molecule, or a fragment thereof, can be immobilized onto a solid support and anti-oncofetal fibronectin antibodies specifically bound thereto can be detected. Oncofetal fibronectin proteins or oncofetal fibronectin encoding nucleic acid molecules used in the assays provided herein can be prepared via recombinant DNA techniques well known in the art. For example, a DNA molecule encoding oncofetal fibronectin or an antigenic fragment thereof can be genetically engineered into an appropriate expression vector for large scale preparation of oncofetal fibronectin protein or nucleic acid. Fusion proteins can be expressed that facilitate labeling, immobilization or detection of oncofetal fibronectin protein. See, for example, the techniques described in Sambrook et al., 1989, *Molecular Cloning: A laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. Alternatively, oncofetal fibronectin protein or nucleic acid can be purified from natural sources, e.g., purified from cells, using purification and separation techniques known in the art. Such purification techniques can include, but are not limited to, molecular sieve chromatography and/or ion exchange chromatography. In one embodiment, microtitre plates can be used as the solid

support for oncofetal fibronectin protein or nucleic acid. The surfaces can be prepared in advance and stored. Anti-oncofetal fibronectin antibodies bound to the solid support can be detected by, for example, contacting the solid support with a detectable antibody such as an anti-(human antibody) antibody, such as an anti-human IgE antibody, where the detectable antibody can be linked to a detectable or bindable moiety or can be detected by a specifically binding partner containing a detectable or bindable moiety, such as an anti-human IgE-horseradish peroxidase conjugate.

[1008] In another example, an anti-(human antibody) antibody, such as an anti-human IgE antibody or an anti-human IgG antibody, or a fragment thereof, can be immobilized onto a solid support and anti-oncofetal fibronectin antibodies specifically bound thereto can be detected. Anti-(human antibody) antibodies used in the assays provided herein can be prepared via a variety of methods provided herein or otherwise known in the art. In one example, microtitre plates can be used as the solid support for anti-(human antibody) antibodies. The surfaces can be prepared in advance and stored. Anti-oncofetal fibronectin antibodies bound to the solid support can be detected by, for example, contacting the solid support with a binding partner such as oncofetal fibronectin protein or nucleic acid or a fragment thereof, where the binding partner can be linked to a detectable or bindable moiety or can be specifically bound by a compound containing a detectable or bindable moiety, such as an anti-human fibronectin-horseradish peroxidase conjugate.

[1009] In another example, anti-oncofetal fibronectin autoantibodies can be detected by contacting a sample with oncofetal fibronectin protein or nucleic acid or a fragment thereof and separating sample components bound to oncofetal fibronectin protein or nucleic acid or fragment thereof from the remainder of the sample components. Such separation methods can be accomplished in any of a variety of ways known in the art, such as, but not limited to, by contacting a sample with oncofetal fibronectin protein or nucleic acid or fragment thereof immobilized to a solid support and then separating the solid support from the sample (e.g., using a washing buffer), or by contacting a sample with oncofetal fibronectin protein or nucleic acid or fragment thereof conjugated to a bindable moiety such as a magnetic bead or biotin and separating the conjugate from the sample (e.g., using a magnet or solid support containing streptavidin to remove the conjugate from the sample). After separating bound sample components from the remainder of the sample, sample components or fragments thereof, can be dissociated from oncofetal fibronectin protein or nucleic acid or fragment thereof and the molecular weights of bound sample components or fragments thereof can be detected. One or more masses indicative of an antibody can indicate the presence of anti-oncofetal fibronectin autoantibodies in the sample. Optionally, prior to measuring masses of sample components, sample components bound to oncofetal fibronectin protein or nucleic acid or fragment thereof can be released from oncofetal fibronectin protein or nucleic acid or fragment thereof and contacted with an anti-(human antibody) antibody under conditions in which human antibodies will specifically bind to the anti-(human antibody) antibody. Sample components or fragments thereof bound to the anti-(human antibody) antibody can be separated from the remainder of the sample components and after such separation, masses of the bound sample components, or fragments thereof, can be measured. As one skilled in the art will

recognize, when two binding and separating events are performed prior to mass measurement, the sequential order of the autoantibody binding partner is readily interchangeable (e.g., sample can be bound first to anti-(human antibody) antibody and second to oncofetal fibronectin protein or nucleic acid, or vice versa).

[1010] Mass measurement of sample components or fragments thereof can be performed by, for example, mass spectrometry. When mass spectrometry is used to detect the molecular weights of sample components or fragments thereof, antibodies or fragments thereof can be identified by detected molecular weights. Proteolysis of antibodies can yield antibody fragments with known masses. Thus, proteolysis of sample components while bound to an autoantibody binding partner or after release from an autoantibody binding partner, can result in fragments having a mass indicative of an antibody fragment. Mass spectrometry can accurately measure masses to determine whether or not a fragment is present that has a mass indicative of an antibody fragment. Detection of a proteolysis fragment having a mass indicative of an antibody fragment can identify the sample as containing anti-oncofetal fibronectin autoantibodies and can thereby indicate the presence of oncofetal fibronectin in the subject.

[1011] Also provided herein are methods for determining the amino acid sequence of one or more autoantibodies that bind to oncofetal fibronectin protein or nucleic acid. Antibodies and antibody fragments measured by mass spectrometry can be further examined to determine all or a portion of the amino acid sequence of the antibody or antibody fragment. A variety of methods of using mass spectrometry to determine the amino acid sequence of proteins or protein fragments are known in the art, including, for example, tandem mass spectrometry measurement methods and computational methods such as SEQUEST® (Thermo Electron Corp., Woburn Mass.). Examples of the variety of known mass spectrometry sequencing methods are provided in U.S. Pat. Nos. 5,538,897, 6,017,693, 6,489,121, 6,670,194 and 6,716,636 and U.S. Pub. No. 20030104483. In an exemplary method, the amino acid sequence of one or more autoantibodies that bind to oncofetal fibronectin protein or nucleic acid can be determined by contacting a sample with an anti-oncofetal fibronectin autoantibody binding partner, separating sample components not bound to the binding partner, optionally releasing bound components and contacting released components with a second anti-oncofetal fibronectin autoantibody binding partner, optionally fragmenting bound components and detecting the molecular weight of bound components or fragments thereof by mass spectrometry, whereby the amino acid sequence of the components or fragments thereof is determined.

[1012] Also provided herein are methods for detecting anti-oncofetal fibronectin autoantibodies in conjunction with detection of one or more additional anti-tumor marker autoantibodies. Such detection methods can be carried out according to the same principles as described herein for detection of anti-oncofetal fibronectin autoantibodies, where the additional tumor marker(s) or fragment thereof can be used as an autoantibody binding partner. For example, a sample can be contacted with an anti-(human antibody) antibody immobilized to a solid support and the solid support also can be contacted with oncofetal fibronectin protein or nucleic acid or fragment thereof and one or more

additional tumor markers or fragment thereof, where oncofetal fibronectin protein or nucleic acid and each tumor marker can be conjugated to a distinguishably detectable or bindable moiety (e.g., fluorophores of different emission wavelength, or one conjugated to biotin and another conjugated to a magnetic bead), whereby distinguishable signals or separate binding events can independently identify the presence of autoantibodies to oncofetal fibronectin protein or nucleic acid and/or one or more additional tumor markers in the sample. In another example, a sample can be contacted with oncofetal fibronectin protein or nucleic acid or fragment thereof and one or more additional tumor markers immobilized to a solid support, where oncofetal fibronectin protein or nucleic acid and each tumor marker can be immobilized to different regions of the solid support and the solid support also can be contacted with an anti-(human antibody) antibody optionally conjugated to a detectable or bindable moiety, where presence of autoantibody in one or more regions can independently identify the presence of autoantibodies to oncofetal fibronectin protein or nucleic acid and/or one or more additional tumor markers in the sample. Methods and apparatuses for performing such procedures can be performed using, for example microarray chips, whose general use is known in the art.

[1013] A variety of additional tumor markers, for which autoantibodies can be detected, are known in the art, including, but not limited to, p53, c-erbB2, c-myc, MUC1, BRCA1, BRCA2, Her-2/neu, bcl-2, bax, PSA, CYFRA 21-1, PTH-RP, CA125, CEA gene family members, progastrin, gastrin G17, gastrin G34, CA 19-9, CA 15-3, CA 27-29, CA 72-4, APC, SCC, HPV subtypes, TK, alphaFP, p62, Kallikrein, ras, vasopressin, gastrin releasing peptide, annexin I, annexin II, Hu and KOC (see, e.g., U.S. Pub. No. 200030232399).

[1014] 6. Measurement of Other Analytes

[1015] In addition to oncofetal fibronectin, analytes can include, but are not limited to: hormones, such as steroidal hormones, including estriol; proteins or peptides, such as human immunodeficiency virus (HIV) antigens, antigens indicative of infectious organisms, such as *Salmonella*, *E. coli*, yeast or parasitic infections, apolipoprotein(a) and lipoprotein(a), environmental antigens, human chorionic gonadotropin (hCG), E-3-G, interleukins and other cytokines and immunomodulatory proteins, such as IL-6 and interferon, small nuclear ribonuclear particles (snRNP) antigens, insulin-like growth factor binding protein one (IGFBP-1), a marker of hypochlorous acid such as 3-chlorotyrosine, tumor markers, or other indicators of cancer or pregnancy related conditions; nucleic acids encoding one of the above proteins or peptides or complements thereto, methylated nucleic acids, or other nucleic acids associated with cancer or pregnancy related conditions; creatinine, sample specific gravity or other analytes that can be used to normalize oncofetal fibronectin indicating molecule measurements.

[1016] a. Normalization

[1017] A neat sample can contain any of a wide range of solute concentrations and also can have added thereto one or more reagents including a dilution buffer. Because of the possible variability of solute concentrations in a sample, it may be ambiguous whether or not the amount of oncofetal fibronectin indicating molecule present in the sample indicates an amount above a threshold level. If necessary, in

order to determine whether the amount of oncofetal fibronectin indicating molecule is at or above a threshold level, the concentration of solutes in a sample can be normalized. For example, the concentration of components in a urine sample can vary depending on a variety of factors including time between urinating, food and liquid intake by the subject and kidney function of the subject. In such instances, analyte concentrations can be normalized by, for example, comparing the analyte concentration to the concentration of a constantly produced urine component to yield a normalized analyte concentration that is less affected by, or independent of factors that cause solute concentrations to vary.

[1018] In one embodiment, one or more components or other characteristics of a sample can be measured in order to normalize the measured amount of the oncofetal fibronectin indicating molecule to the sample. Measured properties and/or components include pH, specific gravity, ionic strength, concentration of a compound produced at a constant rate and concentration of an administered compound that enters into the sample at a constant rate. In the case of urine, components of urine produced by the subject at a constant rate include creatinine, IgA, IgG, albumin, urea, cystatin-C, DTPA (diethylenetriaminepentaacetic acid), EDTA (ethylenediaminetetraacetic acid), iodinated aromatics, metal complexes and organic dyes (see, e.g., Achilefu et al., *Topics in Current Chemistry*, 222:31-72 (2002)). Methods for normalizing urine components using these and other compounds are known in the art, as exemplified in U.S. Pat. No. 6,436,721 for normalizing with creatinine or inulin and in U.S. Pat. No. 6,368,873 for normalizing with IgA, albumin or IgG. Devices for measuring compounds for normalization as well as analytes are known in the art, as exemplified in U.S. Pat. Nos. 5,804,452, 6,368,873 and 6,436,721.

[1019] In one example, the concentration of an oncofetal fibronectin indicating molecule in a urine sample can be normalized by comparing the measured concentration of oncofetal fibronectin indicating molecule in the urine sample to the measured concentration of creatinine in the urine sample. In determining the concentration of an oncofetal fibronectin indicating molecule in a sample, the concentration of oncofetal fibronectin indicating molecule and creatinine in the sample can be measured. The concentration of oncofetal fibronectin indicating molecule can be normalized by determining the ratio of the concentration of the oncofetal fibronectin indicating molecule to the concentration of creatinine in the urine sample. This normalized oncofetal fibronectin indicating molecule concentration can be used to more accurately determine the typical concentration of oncofetal fibronectin indicating molecule in the urine of the subject from whom the sample was collected and/or can be compared to a reference ratio that can be used to identify a subject as positive or negative for oncofetal fibronectin. The normalized oncofetal fibronectin indicating molecule concentration also can be used to more accurately determine a threshold above which a subject has a higher likelihood of disease or delivery. This normalized oncofetal fibronectin indicating molecule concentration also increases the ease of manipulating the urine sample (e.g., concentrating and/or diluting) without changing the experimental results.

[1020] b. Combination with Other Markers

[1021] In one embodiment, detection of an oncofetal fibronectin indicating molecule can be combined with detection of one or more additional markers to determine

increased or decreased likelihood of delivery by a pregnant woman, date of delivery of a pregnant woman, efficacy of induction procedures, presence or absence of a neoplastic disease, efficacy of treatment of a neoplastic disease, or tendency of a subject toward neoplastic disease. For example, an oncofetal fibronectin indicating molecule can be detected in combination with salivary estriol, whereby detection of the oncofetal fibronectin indicating molecule and salivary estriol can indicate increased likelihood of delivery by a pregnant woman, date of delivery of a pregnant woman, or efficacy of induction procedures.

[1022] In another example, detection of an oncofetal fibronectin indicating molecule can be combined with detection of one or more additional tumor markers to determine presence of neoplastic cells. For example, an oncofetal fibronectin indicating molecule can be detected in combination with interleukin-6 (IL-6), interleukin-2 (IL-2), interleukin-12 (L-12) and/or N-acetylgalactosaminyltransferase-T3 (GalNAc-T3). When an oncofetal fibronectin indicating molecule is detected in combination with IL-6, for example, detection of the oncofetal fibronectin indicating molecule and IL-6 can indicate the presence of neoplastic cells, such as neoplastic breast cells, neoplastic bladder cells, neoplastic cervical cells or neoplastic ovarian cells. In another example, tumor markers, such as particular expressed genes, antigens or proteins, or variants thereof, that are indicative of a tumor or neoplastic condition, can be detected in conjunction with the oncofetal fibronectin indicating molecule detection methods provided herein, to indicate the presence or absence of tumorous or neoplastic cells in a subject. For example, oncofetal fibronectin indicating molecule detection can be combined with Her/Neu detection methods in breast cancer diagnostic methods.

[1023] i. Indicators of Membrane Rupture

[1024] In one embodiment, measurement of an oncofetal fibronectin indicating molecule can be accompanied by measurement of an indicator of membrane rupture. Typically, when the oncofetal fibronectin assay is positive, an assay of an indicator of membrane rupture such as insulin-like growth factor binding protein one (IGFBP-1) or a marker of hypochlorous acid such as 3-chlorotyrosine, can be performed on a sample from the subject to determine whether the membranes are intact. The cervicovaginal sample can be the same or different from the sample used to assay for the oncofetal fibronectin indicating molecule. Methods for performing the IGFBP-1 assay are known in the art, as exemplified in international publication No. WO 94/17405; methods for detecting the presence of hypochlorous acid are known in the art, for example, by detecting 3-chlorotyrosine using one of a variety of methods known in the art and exemplified in WO 04/003555. If the membrane rupture indicator assay is negative, the membranes have not ruptured and the subject can be administered a tocolytic agent, as described herein. If, however, the membrane rupture indicator assay is positive, indicating that the membranes have ruptured, the test indicates that delivery cannot be delayed and the tocolytic agent typically is not administered.

[1025] a. Insulin-Like Growth Factor Binding Protein

[1026] IGFBP-1 is assayed by any quantitative or semi-quantitative procedure that can either determine the amount

of IGFBP-1 in the sample or that the amount of IGFBP-1 is at or above a threshold level that indicates rupture of membranes.

[1027] Anti-IGFBP-1 antibodies can be produced by a number of methods. Polyclonal antibodies can be induced by administering an immunogenic composition containing human IGFBP-1 to a host animal. Alternatively, amniotic fluid or another source of high levels of IGFBP-1 can be used as the immunogen and antibodies of the selected specificity can be identified.

[1028] Preparation of immunogenic compositions of IGFBP-1 can vary depending on the host animal and is well known. For example, IGFBP-1 or an antigenic portion thereof can be conjugated to an immunogenic substance such as KLH or BSA, or provided in an adjuvant or the like. The induced antibodies can be tested to determine whether the composition is IGFBP-1-specific. If a polyclonal antibody composition does not provide sufficient specificity, the antibodies can be purified to enhance specificity by a variety of conventional methods. For example, the composition can be purified to reduce binding to other substances by contacting the composition with IGFBP-1 affixed to a solid substrate. Those antibodies which bind to the substrate are retained. Purification techniques using antigens affixed to a variety of solid substrates such as affinity chromatography materials including Sephadex, Sepharose are well known.

[1029] Monoclonal IGFBP-1-specific antibodies also can be prepared by conventional methods. A mouse can be injected with an immunogenic composition containing IGFBP-1 and spleen cells obtained. Those spleen cells can be fused with a fusion partner to prepare hybridomas. Antibodies secreted by the hybridomas can be screened to select a hybridoma where the antibodies react with IGFBP-1 and exhibit substantially no reaction with the other proteins which can be present in a sample. Hybridomas that produce antibodies of selected specificity can be cultured by standard techniques. Hybridoma preparation techniques and culture methods are known in the art.

[1030] The assay conditions and reagents can be any of a variety of methods and conditions known in the art or disclosed herein. The assay can be heterogeneous or homogeneous, conveniently a sandwich assay. The assay usually employs solid phase-affixed anti-IGFBP-1 antibodies. The antibodies can be polyclonal or monoclonal or antibody fragments or other binding moieties. The solid phase-affixed antibodies are combined with the sample. Binding between the antibodies and sample can be determined in a number of ways. Complex formation can be determined by use of soluble antibodies specific for IGFBP-1. The antibodies can be labeled directly or can be detected using labeled second antibodies specific for the species of the soluble antibodies. Various labels include radionuclides, enzymes, fluorescent compounds, colloidal metals or the like. Conveniently, the assay will be a quantitative enzyme-linked immunosorbent assay (ELISA) in which antibodies specific for IGFBP-1 are used as the solid phase-affixed and enzyme-labeled, soluble antibodies. Alternatively, the assay can be based on competitive inhibition, where IGFBP-1 in the sample competes with a known amount of IGFBP-1 for a predetermined amount of anti-IGFBP-1 antibody. For example, any IGFBP-1 present in the sample can compete with a known amount of the labeled IGFBP-1 or IGFBP-1 analogue for

antibody binding sites. The amount of labeled IGFBP-1 affixed to the solid phase or remaining in solution can be determined.

[1031] Appropriate dilution of the conjugate can be performed to detect the selected threshold level of IGFBP-1 which is above background values for the assay as a positive sample.

[1032] The assay results can be interpreted as follows. IGFBP-1 levels below 20-50 ng/mL can be considered background and are negative. The cutoff of choice for the background level depends upon whether a high sensitivity or high specificity test is to be selected. For example, 24 cervicovaginal secretion specimens that exhibited a positive oncofetal fibronectin test results (>50 ng/mL for buffer-treated swab samples) for impending delivery, and exhibited negative ferning, pooling and nitrazine test results for rupture of membranes were tested for IGFBP-1. One of these specimens demonstrated 42 ng/mL IGFBP-1, and the remainder exhibited less than 20 ng/mL IGFBP-1. If a cut-off of 20 ng/mL were to be used, the demonstrated specificity of the test to rule out rupture would be 97%. On the other hand, if 50 ng/mL were to be used, the rule out specificity of the test would be 100%. In most cases, high rule-out specificity can be used, since subjects with ruptured membranes are in greater danger of infection than those who do not have rupture, so a cutoff can range between 20-50 ng/ml or about 20-50 ng/mL.

[1033] Other assays using different reagents can have different cutoff values. For example, IGFBP-1 antibodies which differ in their antigen binding characteristics can produce assay results with different optimal cutoff values. One of skill in the art understands that background values can vary when different reagents are used and will understand how to determine the proper background level for the selected specificity and sensitivity for a selected assay. In addition, different optimal cutoffs can be used for different applications. For example, when using oncofetal fibronectin as a predictor of delivery date, a cutoff for a buffer-treated swab sample of 50 ng/ml or about 50 ng/mL oncofetal fibronectin protein can be used.

[1034] The presence of IGFBP-1 in a cervicovaginal secretion sample from a subject who is positive for a marker that indicates increased risk of delivery indicates that the membranes have ruptured. If IGFBP-1 is less than 20-50 ng/mL or undetectable (background for the assay), the membranes remain intact. When IGFBP-1 is positive (>20-50 ng/mL) and the delivery marker (e.g., oncofetal fibronectin) is negative, then amniotic membranes can have ruptured, although most subjects who have ruptured membranes will exhibit positive IGFBP-1 and the delivery marker simultaneously. When IGFBP-1 is negative, the IGFBP-1 test can be repeated, typically daily, until the sample is positive for IGFBP-1.

[1035] b. Hypochlorous Acid

[1036] Presence of hypochlorous acid can be assayed by any of a variety of known methods, as known in the art, as exemplified in Winterbourn et al., *Free Radic. Biol. Med.* 29:403-409 (2000) and Chapman et al., *Arch. Biochem. Biophys.* 377:95-100 (2000). A typical method for determining the presence of hypochlorous acid is performed by detecting 3-chlorotyrosine and/or other chlorotyrosines in a

sample. Methods for detecting 3-chlorotyrosine and other chlorotyrosines including dichlorotyrosine, are known in the art, as exemplified in WO 04/003555 to Woods et al., Kettle, A.J., *Methods Enzymol.* 300:111-120 (1999) and Hazen et al., *J. Clin. Invest.* 99:2075-2081 (1997). Typical methods for detecting chlorotyrosines such as 3-chlorotyrosine include gas chromatography/mass spectrometry methods (see, e.g., Hazen et al., *J. Clin. Invest.* 99:2075-2081 (1997)) and binding partner-based methods such as immunoassays including ELISA (see, e.g., WO 04/003555 to Woods et al.).

[1037] In one embodiment, the presence of any indicator of hypochlorous acid can indicate membrane rupture. In other embodiments, the presence of an amount of indicator of hypochlorous acid at or above a threshold level can indicate membrane rupture. Typically, the amount of indicator of hypochlorous acid that can indicate membrane rupture is an amount greater than the average amount, such as mean or median amount, present in pregnant women without membrane rupture. The degree to which a sample can reflect membrane rupture can be determined according to the number of standard deviations above the mean amount present in pregnant women without membrane rupture, according to the diagnosis goal. For example, one standard deviation can be used for an assay that can include false positives which can be selected for further evaluation, or for subjects known to have normal or low estriol values and relatively little variation between samples, or for high risk individuals. Two standard deviations above the mean or three standard deviations above the mean, are typically used.

[1038] Anti-chlorotyrosine antibodies can be produced by a number of methods known in the art. Exemplary methods and antigens for raising such antibodies are demonstrated in WO 04/003555. For example, polyclonal antibodies can be induced by administering to a host animal an immunogenic composition containing, for example, 3-(3-chloro-4-hydroxy-benzyl)-6-mercaptopethylpiperazine-2,5-dione linked to a carrier protein, N-acetyl-3-chlorotyrosine linked to a carrier protein, or N-acetyl-3,5-chlorotyrosine linked to a carrier protein.

[1039] Preparation of immunogenic compositions can vary depending on the host animal and is well known. For example, a chlorotyrosine-containing compound can be conjugated to an immunogenic substance such as KLH or BSA, or provided in an adjuvant. The induced antibodies can be tested to determine whether the composition is chlorotyrosine-specific. If a polyclonal antibody composition does not provide sufficient specificity, the antibodies can be purified to enhance specificity by a variety of conventional methods. For example, the composition can be purified to reduce binding to other substances by contacting the composition with chlorotyrosine affixed to a solid substrate. Those antibodies which bind to the substrate are retained. Purification techniques using antigens affixed to a variety of solid substrates such as affinity chromatography materials including Sephadex, and Sepharose are well known.

[1040] Monoclonal chlorotyrosine-specific antibodies also can be prepared by conventional methods. A mouse can be injected with an immunogenic composition containing chlorotyrosine and spleen cells obtained. Those spleen cells can be fused with a fusion partner to prepare hybridomas. Antibodies secreted by the hybridomas can be screened to select a hybridoma where the antibodies react with chloro-

tyrosine and exhibit substantially no reaction with the other proteins which can be present in a sample. Hybridomas that produce antibodies of selected specificity can be cultured by standard techniques. Hybridoma preparation techniques and culture methods are known in the art.

[1041] The assay conditions and reagents can be any of a variety of methods and conditions known in the art or disclosed herein. The assay can be heterogeneous or homogeneous, conveniently a sandwich assay. The assay can employ solid phase-affixed anti-chlorotyrosine antibodies. The antibodies can be polyclonal or monoclonal or antibody fragments or other binding moieties. The solid phase-affixed antibodies are combined with the sample. Binding between the antibodies and sample can be determined in a number of ways. Complex formation can be determined by use of soluble antibodies specific for chlorotyrosine. The antibodies can be labeled directly or can be detected using labeled second antibodies specific for the species of the soluble antibodies. Various labels include radionuclides, enzymes, fluorescers, colloidal metals or the like. Conveniently, the assay will be a quantitative enzyme-linked immunosorbent assay (ELISA) in which antibodies specific for chlorotyrosine are used as the solid phase-affixed and enzyme-labeled, soluble antibodies. Alternatively, the assay can be based on competitive inhibition, where chlorotyrosine in the sample competes with a known amount of chlorotyrosine or compounds containing chlorotyrosine for a predetermined amount of anti-chlorotyrosine antibody and competition can be measured by methods known in the art.

[1042] Gas chromatography/Mass spectrometry methods for detecting chlorotyrosine can be performed as known in the art and exemplified in Hazen et al., *J. Clin. Invest.* 99:2075-2081 (1997). Briefly, a sample can be treated in one or more separation steps, including liquid chromatography or gas chromatography steps. The chromatographed sample can then be ionized, by, for example, chemical ionization and the mass and/or mass to charge ratio of the ions measured by mass spectrometry.

[1043] ii. Estriol

[1044] In addition to, or as an alternative to, the IGFBP-1 assay, the concentration of estriol can be determined in a sample obtained from the subject. There are no limitations on the type of assay used to measure estriol. Any assay for estriol can be employed.

[1045] A variety of examples of estriol assays are known in the art, as exemplified in U.S. Pat. No. 5,480,776, issued Jan. 2, 1996. Briefly, the assay can be performed on any sample of body fluid, such as blood (or a blood fraction, especially serum or plasma), urine, cervical or vaginal secretions, sweat, saliva or other fluid. Estriol is sufficiently soluble in water so that it is distributed in fluids throughout the body. For simplicity of sampling and because, unlike in urine, detection is not complicated by the presence of estrogen conjugates, saliva can be used.

[1046] Assays are generally directed to detection of free estriol, since conjugated estriol has reduced biological activity. In saliva about 92% of estriol is in the free form, while most estriol in urine is present as a conjugate. As will be clear to those familiar with steroid metabolism, an estriol conjugate is a compound formed by formation of a covalent linkage of a non-steroidal compound to estriol. Linkage is

typically through a hydroxyl group of the steroidal ring system. The non-steroidal component can be inorganic (e.g., a sulfate group) or organic (e.g., a glucuronide group).

[1047] There are no limitations on the collection and handling of samples as long as consistency is maintained. With some body fluids, such as saliva and plasma, there is little diurnal variation in estriol levels. For other fluids, notably urine, variations occur. In one embodiment variations are eliminated to the extent possible, for example by taking samples at the same time of day. Other techniques can be used to ensure consistency of measurement of analytes in clinical fluids. For example, creatinine can be measured concurrently with estriol in urine to normalize the concentration of estriol. Creatinine is produced at a constant rate in the kidneys and measurement of creatinine concentration allows correction of volume errors in urine samples, as is well known in the art.

[1048] Optionally and depending on the source of the fluid being tested, free estriol can be separated from estriol conjugates. Techniques for such separations are known in the art. See, for example, Evan, N. *Z. Med. Lab. Tech.* 33:86 (1979), which describes such separations as well as two radioimmunoassays useful for measuring plasma estriol. These separations are generally difficult and assays that do not require separation, either because of the use of specific antibodies or other binding partners that differentiate between free and conjugated estriol, or because the sample is obtained from a source containing mostly free estriol, such as saliva, can be used.

[1049] The concentration of estriol in the fluid assayed is correlated with a standard value to determine when labor is imminent. The standard is usually (1) a predetermined range of estriol concentrations for the same body fluid in normal pregnant humans in the general population, either at the corresponding time in the pregnancy or a specific time relative to normal termination of pregnancy, or (2) a previously measured estriol concentration of the same body fluid of the same pregnant human. A measured higher concentration of estriol relative to the standard value is an indication of potential onset of pre-term labor. The methods herein do not require the measurement of any other substance, such as the progesterone concentration in the body fluid, or require the measurement of total estriol production for a time interval. Optionally, measurements of total estriol for a given time period, such as 24 hours, can be used with urine.

[1050] The first general standard set out above, namely a predetermined range of estriol concentrations for the same body fluid in normal pregnant humans in general, is typically obtained by the same assay technique that will be used in the application of the method to an individual being tested, in order to ensure the highest correlation. Sufficient measurements are made in a normal population of pregnant women to produce a statistically significant range of normal values for the value to which a comparison will be made, which typically is at preselected time intervals during normal pregnancy. While comparison to a time immediately prior to normal delivery (38 to 40 weeks) is often used, other time periods can be used. For example, estriol levels during a given week of a individual pregnancy (i.e., that of the subject) can be compared to the normal range of concentrations for the same time period (e.g., the 20th week). Generally, the minimum concentration indicative of possible

onset of labor is considered to be at least 1, generally at least 2, typically at least 3 or at least 4, standard deviations above the mean estriol concentration determined just prior to the onset of labor for normal pregnant humans for any given body fluid.

[1051] It will be recognized by those familiar with statistics that the number of standard deviations used as an indication of pregnancy complications will be selected with an appropriate diagnosis goal in mind. For example, one standard deviation would encompass 68% or about 68% of normal samples; that is, 32% of normal samples would be expected to fall outside the lower and upper limits set by one standard deviation from the mean (16% would thus be expected to be above the selection limit). Thus, one standard deviation above the normal mean is not used for routine analysis, as it would include too many false positives. One standard deviation is appropriate for an assay that is selected to sweep in for further evaluation all possible candidates who might be predisposed toward pre-term labor, or this limit can be selected for subjects known to have normal or low estriol values and relatively little variation between samples. One standard deviation also can be selected for a subject known to have problems with pre-term labor in order to determine when to more closely monitor the subject under controlled conditions (such as by having a subject admitted to a hospital for constant monitoring). Two standard deviations from the mean would encompass 95% or about 95% of normal samples; three standard deviations, 99% or about 99%; four standard deviations, more than 99%. These levels are more appropriate generally, especially for subjects whose levels of estriol are known to be normal or slightly above normal or to vary from sample to sample as well as for assays with a high coefficient of variation.

[1052] It is not necessary to express the lower limit of the indication of labor (upper limit of the normal range) in standard deviations. Any other system that can be used to provide a statistically significant indication of probable onset of labor can be used. For example, the limit can be set to be a concentration that is at least as high as the 95th percentile concentration for normal subjects for the same body fluid for a normal pregnancy. In any case, a normal limit from the 38-42 week period, typically 40 weeks or about 40 weeks, can be selected for normal pregnancies. Monitoring the concentration can be initiated at a selected time, such as 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 weeks or earlier, and continued until delivery.

[1053] Because of the many different possible clinical goals, the actual estriol level indicative of probable onset of pre-term labor is best selected by the attending physician after collecting data from several samples during the initial portion of the pregnancy and taking into consideration the time at which the measurement is being made. For example, in a normal pregnancy at week 30, the change expected in the estriol concentration prior to the onset of labor is smaller than 2 standard deviations from the mean concentration of estriol at 30 weeks. Thus, while assays in the first portion of a pregnancy (prior to 30 weeks) can use 3 or 4 standard deviations as an indication of onset of labor, two, one and a half, or even one standard deviation would be more appropriate in the later portion of a pregnancy (e.g., after 30 weeks) depending on the condition of the subject, other clinical indications in the mother known to the attending physician and the health of the fetus. It is the earlier stages

of a pregnancy that require greater attention to avoiding pre-term labor, because of the lack of fetal development at these stages and the higher risk of infant death post partum. Pre-term labor is generally considered to be any labor prior to the end of a normal 40-week term of pregnancy.

[1054] The methods herein can be used for pregnancies during weeks 20 to 36, when prolonging pregnancy for even a short time is most efficacious in reducing the effects of premature birth. The assay, particularly when used to a detect rate of increase, can be employed in pregnancies terminated by labor and delivery after the end of 40 weeks and measurements can be made during this time period. When employed at weeks 38 and higher, the methods provided herein typically are practiced using the "self-comparison" method discussed elsewhere herein; i.e., by comparing the measurement at a given time with a measurement made earlier with the same subject.

[1055] In a similar manner, subject to the same constraints discussed above, an assay concentration of at least 1, generally at least 2, typically at least 3 or at least 4, standard deviations above the mean normal concentration for the same stage of pregnancy also can be used as an indication of an abnormal pregnancy and thus as an indication of possible onset of labor, although the probability is lower if the measured amount does not reach the levels considered normal for weeks 38-42.

[1056] Standard values will vary with the specific body fluid whose concentration is being measured and with the specific assay being used (although to a lesser extent). Typical minimum indicative levels of labor onset in an assay that measures unconjugated estriol are as follows for the indicated body fluids (all concentrations are in nM): saliva, at least 3, typically at least 5 or at least 6 or at least 7; serum, 30, at least 35 or at least 45.

[1057] As an alternative to comparing estriol concentrations to those present in a normal population, a previously measured estriol concentration of the same body fluid of the same pregnant human can be used as a standard for comparison. In this case, what is being determined is usually the rate of increase in estriol concentration in the fluid being tested. A positive assay (i.e., indication of imminent onset of labor) is considered to be present when the measured concentration exceeds a previously measured estriol concentration made in the same body fluid in the same pregnant human female by 50%, generally 75%, typically 100%, within one week. Again the selection of a particular rate of increase to label as the lower limit of labor onset can be selected by the attending physician for the particular reason selected. For example a screening test that is intended to collect potential problem subjects into the hospital for further observation and study could select the 50% increase as its limit in order to avoid false negative results, while accepting the problems caused by including a relatively large number of false positives. Higher percentage increases as the minimum positive indication are more acceptable for home assays and point of care assays, in the same manner as described above for standard deviations from the normal population mean. Increases in estriol concentration that meet the standards of this paragraph and additionally reach levels previously indicated to be indicative of the onset of labor in normal populations of subjects are particularly likely to indicate imminent onset of labor.

[1058] Many assays can be used. For example, in U.S. Pat. No. 5,480,776, an enzyme-label component (here a labeled estriol molecule or derivative thereof) is used in a competitive binding assay for estriol. The assay is a non-instrumented enzyme immunoassay that provides present/not-present or "threshold" (\pm) analysis results at a preselected cut-off value and thus can be used herein.

[1059] In a typical assay using this technique, the enzyme-labeled, competitive binding component contains estriol (or the portion thereof used to generate the antibody used in the assay) bound to the immunogen that is used to produce the antibody of the assay. An enzyme label is bound to this moiety, such as through a bulky linker such as an avidin-biotin complex. The use of such a competitive binding partner allows antibodies to be used without attempting to manipulate affinity of binding of antibody to competitor while still providing the steep competitive binding curve required for a \pm analysis.

[1060] In a typical such assay, antibody is attached to a solid surface, such as a microtiter plate well, a test tube, or a porous reagent strip (such as cellulose or glass fibers). The antibody-coated solid surface then is contacted simultaneously with a sample and with a competitive binding partner. By providing fewer antibody binding sites than are present in the combined total of analyte and competitive binding partner, only a fraction of the molecules in solution will bind to the solid surface. If there are no analyte molecules present, all of the binding sites will be taken up by the competitive binding partners so that a maximum amount of enzyme is attached to the solid surface. When a substrate for the enzyme is contacted with the solid surface after the sample is washed away, reaction of the enzyme with the substrate provides a detectable signal (usually formation of a color) that indicates to the user the absence of analyte in the sample (a negative result). If analyte is present in the sample, analyte competes for binding sites so that less of the enzyme-label competitor can bind. By using a bulky binding composition, which binds less rapidly to the antibody than does the analyte and by properly selecting the number of binding sites relative to the amount of sample added (which is a standard technique to one of skill in the art), analyte present at a concentration above a preselected minimum level will exclude binding of the competitive binding composition and thus binding of the enzyme to the solid substrate. An example of such a selection process to provide different threshold levels is found in U.S. Pat. No. 5,480,776. Thus, if sufficient analyte is present in the sample, after reaction, no enzyme is present to produce a color change and the reaction mixture stays the same (thus a positive reaction using this reaction scheme).

[1061] Other reaction schemes can be used in which the formation of color is indicative of the presence of the analyte. The previous example is merely one of many types of competitive binding assays in which estriol can be measured.

[1062] Antibody production for use in an assay for estriol is conventional and is not described here in detail. A brief discussion of general techniques for the production of antibodies specific for steroids follows.

[1063] An animal is injected with a composition containing estriol covalently attached to an immunogen, usually a protein, prepared as described above. Multiple injections or

the use of an adjuvant will ensure maximum stimulation of the immune system and production of antibodies. If polyclonal antibodies are selected, they can be prepared by simply collecting blood from the immunized animal and separating the antibodies from other blood components by standard techniques. To obtain monoclonal antibodies, the spleen or lymphocytes from the immunized animal are removed and immortalized or used to prepare hybridomas by cell-fusion methods known to those skilled in the art. Antibodies secreted by the immortalized cells are screened to determine the clones that secrete antibodies of the selected specificity. For monoclonal anti-estriol antibodies, the antibodies must bind to estriol. Cells producing antibodies of the selected specificity are selected, cloned and grown to produce the monoclonal antibodies of selected specificity.

[1064] Antibody can be attached to a solid surface for use in an assay using known techniques for attaching protein material to solid support materials. The solid support can include plastic surfaces of test tubes or microtiter plates, polymeric beads, dip sticks, or filter materials. The attachment methods include non-specific adsorption of the protein to the support and covalent attachment of the protein, typically through a free amino group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group.

[1065] iii. Other Tumor Indicators

[1066] Also provided herein are methods for detecting an oncofetal fibronectin indicating molecule in conjunction with detection of one or more additional tumor markers. Such detection methods can be carried out according to the same principles as described herein for detection of an oncofetal fibronectin indicating molecule, including protein detection methods, nucleic acid molecule detection methods and autoantibody detections, such as, but not limited to, mass spectrometric methods, sandwich assays, test strip-based assays and in situ assays.

[1067] For example, a sample can be contacted with a solid support having immobilized thereto an anti-oncofetal fibronectin polyclonal antibody and an anti-(tumor marker) polyclonal antibody and the solid support also can be contacted with a monoclonal anti-oncofetal fibronectin antibody or fragment thereof and a monoclonal anti-(tumor marker) antibody or fragment thereof, where the monoclonal antibodies are conjugated to distinguishably detectable or bindable moieties (e.g., fluorophores of different emission wavelength, or one conjugated to biotin and another conjugated to a magnetic bead), whereby distinguishable signals or separate binding events can independently identify the presence of an oncofetal fibronectin indicating molecule and/or one or more additional tumor markers in the sample. A variety of different combinations, such as those discussed for the various methods provided herein, also can be used for detection of an oncofetal fibronectin indicating molecule and one or more additional tumor markers. A variety of additional detectable tumor markers are known in the art, including, but not limited to, AE1/AE3, BCA-225, Cathepsin D, E-Cadherin, Epidermal Growth Factor Receptor (EGFR), Estrogen receptor (ER), Gross Cystic Disease Fluid Protein 15 (GCDFP-15), HOX-B3, Ki-67, p65, Progesterone Receptor (PR), Retinoblastoma (Rb) and Transglutaminase K (TGK), p21, DCC, NF-1, NF-2, BRCA-3, p16, FHIT, WT-1, MEN-I, MEN-IIa, MEN-IIb, VHL, FCC, MCC, raf, erb, src,

fms, jun, trk, ret, gsp, hst, bcr/ab1, p53, c-erbB2, c-myc, MUC1, BRCA1, BRCA2, Her-2/neu, bcl-2, bax, PSA, CYFRA 21-1, PTH-RP, CA125, CEA gene family members, pro-gastrin, gastrin G17, gastrin G34, CA 19-9, CA 15-3, CA 27-29, CA 72-4, APC, SCC, HPV subtypes, TK, alphaFP, p62, Kallikrein, ras, vasopressin, gastrin releasing peptide, annexin I, annexin II, Hu and KOC. Binding partners for such tumor markers, methods of detecting such tumor markers and tumors correlated with such tumor markers are known in the art, as exemplified in U.S. Pub. Nos. 20030190602 and 20040059519.

G. Analysis of Detection Measurements

[1068] Oncofetal fibronectin indicating molecule measurements made by any of the methods provided herein or known in the art can be analyzed to provide information about the subject from which the sample was collected, including information about the general health state of the subject, the propensity to childbirth of the subject or information about a tumor or neoplastic disease in the subject.

[1069] 1. Quantitation

[1070] The amount of an oncofetal fibronectin indicating molecule in a sample can be quantitated. Methods provided herein for measurement of oncofetal fibronectin indicating molecules can provide quantitative measurements. For example, mass spectrometry can be used to provide quantitative results about an analyte in a sample by a variety of methods including by use of an internal standard or by labeling components to be measured (see, e.g., Bucknall et al., *J. Am. Soc. Mass Spectrom.*, 13:1015-1027 (2002); Ross et al., *Biotechniques* 2000:620-629; Amexis et al., *Proc. Natl. Acad. Sci. USA* 98:12097-12102 (2001); Griffin et al., *Anal. Chem.* 73:978-986 (2001)). RT-PCR and related methods can yield quantitative results using procedures such as competitive RT-PCR, real time RT-PCR and other methods, as described herein or known in the art. Quantitative binding assays including competitive binding assays also are known in the art and can be used for quantitation of oncofetal fibronectin. A variety of spectroscopic or reflectance measurements can be used to determine the concentration of an oncofetal fibronectin indicating molecule in a sample according to the intensity of the signal that represents the presence of the oncofetal fibronectin indicating molecule. For example, a test strip can be used having a region containing a fibronectin or oncofetal fibronectin binding partner immobilized thereto and the amount of light reflected from the test strip at that region can indicate the amount of oncofetal fibronectin indicating molecule in the sample and the amount of light can be measured by a reflectance reader. Exemplary quantitative methods include ELISA methods for quantitating the level of an oncofetal fibronectin indicating molecule in a cervicovaginal swab sample and use of a lateral flow test strip device and a test strip reader for quantitating the level of an oncofetal fibronectin indicating molecule in a urine sample.

[1071] 2. Thresholds

[1072] Presence of an oncofetal fibronectin indicating molecule in a sample can be an indicator of any of a variety of biological or health conditions, including overall health state, pregnancy or delivery related conditions, pre-cancerous or cancer (e.g., neoplastic disease) conditions in a subject. In some cases a measurement is considered positive

for oncofetal fibronectin when any oncofetal fibronectin indicating molecule is detected in a sample. In other cases, a measurement is considered positive for oncofetal fibronectin when the presence of an oncofetal fibronectin indicating molecule in a sample is equal to or above one or more threshold levels. In one example, a threshold level of oncofetal fibronectin protein in a buffer-treated cervicovaginal sample assayed using a test strip can be 50 ng/mL. In another example, a threshold level for oncofetal fibronectin protein in a buffer-treated cervicovaginal sample assayed using a test strip can be 150 ng/mL.

[1073] In embodiments that compare the level of oncofetal fibronectin in a sample to a threshold level, the threshold level can be the amount of an oncofetal fibronectin indicating molecule present in an unmodified sample, or the threshold level can be the amount of an oncofetal fibronectin indicating molecule present in a modified sample (e.g., the concentration of an oncofetal fibronectin indicating molecule of a cervicovaginal swab sample after mixture with a buffer solution). Reference herein to the level of an oncofetal fibronectin indicating molecule in a sample or the threshold level of an oncofetal fibronectin indicating molecule typically refers to the level of an oncofetal fibronectin indicating molecule in a modified sample. For example, some oncofetal fibronectin indicating molecule measurements, such as measurement of an oncofetal fibronectin indicating molecule in a cervicovaginal swab sample, are known in the art according to the sample-modified form; thus, oncofetal fibronectin indicating molecule levels and threshold levels for a cervicovaginal swab sample typically refer to the sample modified level.

[1074] In some embodiments, the measured amount of an oncofetal fibronectin indicating molecule can be compared to one or more thresholds. Provided herein one or more thresholds is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more thresholds. Typically, an oncofetal fibronectin indicating molecule concentration in the sample equal to or above a threshold level indicates that the sample is oncofetal fibronectin positive. In one embodiment, an oncofetal fibronectin indicating molecule concentration in a buffer-treated cervicovaginal swab sample of 50 ng/ml or more (or 500 ng/ml untreated swab sample or more), or about 50 ng/ml or more (or about 500 ng/ml untreated swab sample or more) indicates that the sample is oncofetal fibronectin positive. Typically, an oncofetal fibronectin indicating molecule concentration in the sample below a threshold level indicates that the sample is oncofetal fibronectin negative. In one embodiment, an oncofetal fibronectin indicating molecule concentration in a buffer-treated cervicovaginal swab sample of less than 50 ng/ml (or less than 500 ng/ml untreated swab sample), or about 50 ng/ml (or less than about 500 ng/ml untreated swab sample) indicates that the sample is oncofetal fibronectin negative.

[1075] Different sample types can have different threshold levels. Provided herein, different sample types also can have related threshold levels. For example, the amount of an oncofetal fibronectin indicating molecule in a cervicovaginal swab sample collected from the portion of the vagina below the posterior fornix, such as the lower third of the vagina, can be one-third or about one-third the amount of an oncofetal fibronectin indicating molecule in a cervicovaginal swab of the posterior fornix collected from the same subject. In another example, the amount of an oncofetal

fibronectin indicating molecule in a urine sample can be one-tenth or about one-tenth the amount of an oncofetal fibronectin indicating molecule in a cervicovaginal swab of the posterior fornix collected from the same subject.

[1076] Exemplary threshold values for buffer-treated samples that can indicate different likelihoods of imminent or pre-term delivery include 50 ng/ml, 150 ng/ml, 200 ng/ml, 300 ng/ml, 500 ng/ml, 750 ng/ml and 1000 ng/ml, or about 50 ng/ml, about 150 ng/ml, about 200 ng/ml, about 300 ng/ml, about 500 ng/ml, about 750 ng/ml and about 1000 ng/ml. Exemplary threshold values for untreated samples that can indicate different likelihoods of imminent or pre-term delivery include 500 ng/ml, 1500 ng/ml, 2000 ng/ml, 3000 ng/ml, 5000 ng/ml, 7500 ng/ml and 10000 ng/ml, or about 500 ng/ml, about 1500 ng/ml, about 2000 ng/ml, about 3000 ng/ml, about 5000 ng/ml, about 7500 ng/ml and about 10000 ng/ml. In some embodiments, the oncofetal fibronectin indicating molecule is oncofetal fibronectin protein.

[1077] In one embodiment, where serum or plasma is the sample, a threshold level can be 3 $\mu\text{g/mL}$ to 12 $\mu\text{g/mL}$, or about 3 $\mu\text{g/mL}$ to about 12 $\mu\text{g/mL}$ of FDC-6 positive oncofetal fibronectin protein. A serum or plasma sample containing more than the threshold level can be indicative of a tumorous or neoplastic disease state in the subject, or indicative of an increased risk of pre-term or imminent delivery, or indicative of some other problem with the health state of the individual. In one example, subjects with preeclampsia exhibited oncofetal fibronectin protein plasma levels ranging from 11.5 $\mu\text{g/mL}$ to 38 $\mu\text{g/mL}$, while normal subject exhibited oncofetal fibronectin protein levels ranging from 4 $\mu\text{g/mL}$ to 12 $\mu\text{g/mL}$, as measured by ELISA test using FDC-6 and goat polyclonal anti-human fibronectin as oncofetal fibronectin protein binding partners (Kupferminc et al., *Am. J. Obstet. Gynecol.* 172:649-652 (1995)). Accordingly, detection of FDC-6 reactive oncofetal fibronectin protein levels above 12 $\mu\text{g/mL}$ or about 12 $\mu\text{g/mL}$ in a serum sample can indicate that the subject exhibits oncofetal fibronectin protein levels above the threshold and, therefore, is positive for oncofetal fibronectin; detection of FDC-6 reactive oncofetal fibronectin protein levels in serum between 8 $\mu\text{g/mL}$ and 12 $\mu\text{g/mL}$ or between about 8 $\mu\text{g/mL}$ and about 12 $\mu\text{g/mL}$ can indicate that the subject may or may not be positive for oncofetal fibronectin; and detection of FDC-6 reactive oncofetal fibronectin protein levels in serum below 8 $\mu\text{g/mL}$ or about 8 $\mu\text{g/mL}$ can indicate that the subject is negative for oncofetal fibronectin.

[1078] In other cases, multi-tiered thresholds can be applied to the oncofetal fibronectin measurement, where multi-tiered thresholds include two or more threshold levels, where each larger threshold level indicates a separate health state categorization; for example each larger threshold level can indicate a more severe health problem, an increased likelihood of imminent delivery, increased certainty of delivery date, or increased aggressiveness of a cancer. An exemplary multi-tiered threshold is a two-tiered threshold for oncofetal fibronectin protein, where the lower threshold is 50 ng/mL and the higher threshold is 150 ng/mL for buffer-treated samples. A sample can be categorized according to the thresholds below a measured amount of an oncofetal fibronectin indicating molecule in the sample. For example, a pregnant woman at week 24 of her pregnancy having an amount of an oncofetal fibronectin indicating

molecule that is higher than a first threshold but not higher than a second threshold, can be categorized as having an increased likelihood of pre-term delivery, whereas a pregnant woman at week 24 of her pregnancy having an amount of an oncofetal fibronectin indicating molecule that is higher than the first and second threshold levels can be categorized as having a higher risk of imminent delivery.

[1079] In some instances, one or more threshold levels or one or more threshold curves applied to a measured amount in a subject's sample can be determined according to any of a variety of subject-specific factors. In one example, a subject-specific factor can be the measured amount of one or more samples from a subject.

[1080] In another embodiment, the threshold level can vary according to one or more additional factors. Such factors can include, but are not limited to, biological state of the subject, such as the term of pregnancy, the presence of one or more other markers, anatomical factors, previous health history and genetic factors, the progression of disease, or the age of the subject. Factors also can include the change in the biological state of the subject over time, including the rate of increase in oncofetal fibronectin indicating molecule measurements over time. The varying threshold level can be expressed as a threshold curve where the threshold level of an oncofetal fibronectin indicating molecule varies as a function of time (e.g., week of pregnancy term). A threshold level can decrease with increasing time, or a threshold level can increase with increasing time.

[1081] In some instances, the rate of change of the amount of an oncofetal fibronectin indicating molecule in a particular sample type (e.g., cervicovaginal swab) from a subject can be used to identify a sample as oncofetal fibronectin positive or negative, or to categorize the sample into two or more populations. The rate of change of the amount of an oncofetal fibronectin indicating molecule in a type of sample can indicate a stable, increasing or decreasing amount of the oncofetal fibronectin indicating molecule in the sample.

[1082] 3. Identification of Tissue Source

[1083] An oncofetal fibronectin indicating molecule can contain any of a variety of different splice regions (e.g., EDA, EDB or IIICS) and oncofetal fibronectin protein can contain post-translational modifications (e.g., O-glycosylation). Thus, a variety of non-identical proteins can be termed oncofetal fibronectin, and a variety of non-identical nucleic acid molecules can encode oncofetal fibronectin. In studies of samples, a variety of different fibronectin proteins containing different combinations of these splice regions and post-translational modifications have been observed. The methods provided herein or known in the art can be used to identify the presence of an oncofetal fibronectin indicating molecule in a sample and also can be used to identify one or more splice regions and/or post-translational modifications present in an oncofetal fibronectin indicating molecule in the sample. Identification of the presence or absence of one or more splice regions and/or post-translational modifications present in an oncofetal fibronectin indicating molecule is referred to herein as characterization of oncofetal fibronectin.

[1084] In one embodiment, characterization of oncofetal fibronectin can be used to determine the biological source of the oncofetal fibronectin indicating molecule. For example,

characterization of oncofetal fibronectin can be used to identify the tissue source of an oncofetal fibronectin indicating molecule in a sample. For example, oncofetal fibronectin protein containing EDB and IIICS with threonine 33 O-glycosylated (referred to herein as EDB+, FDC-6+ onffFN) has been observed in liver cirrhosis, liver metastases, dilated cardiomyopathy, fibromatosis, rheumatoid arthritis, nodular palmar fibromatosis, pituitary adenoma, breast carcinoma, invasive ductal carcinoma of the breast, oral squamous cell carcinoma, colon carcinoma and renal carcinoma; an oncofetal fibronectin indicating molecule containing EDB (EDB+ onffFN) has been found in extracellular matrix containing newly forming blood vessels, in brain tumor, prostate carcinoma, benign prostatic hyperplasia, stomach adenocarcinoma, kidney clear cell carcinoma, urinary bladder carcinoma, skin carcinoma, skin and ocular melanoma, lung carcinoma and colon carcinoma; an oncofetal fibronectin indicating molecule containing EDB and EDA (EDB+, EDA+ onffFN) has been observed in fibroblasts and macrophages in healing wounds, glomerulonephritis, nodular palmar fibromatosis and thyroid tumor; an oncofetal fibronectin indicating molecule containing EDA (EDA+ onffFN) has been observed in psoriasis and liver tumor; an oncofetal fibronectin indicating molecule containing CS1 (IIICS/CS1+ onffFN) has been observed in liver tumor.

[1085] The methods provided herein and known in the art can be used to characterize an oncofetal fibronectin indicating molecule in a sample according to the presence of one or more domains and/or post-translational modifications. After such determination, the characterized an oncofetal fibronectin indicating molecule of the sample can be compared to tissues, tumors and other biological sources known to contain oncofetal fibronectin. A tissue, tumor or other source that is known to contain the same oncofetal fibronectin variant as the characterized oncofetal fibronectin indicating molecule from a sample, can be identified as a potential tissue, tumor or other biological source of the oncofetal fibronectin indicating molecule. For example, if a sample contains EDB+, FDC-6+ onffFN protein, the source of the oncofetal fibronectin protein in the sample can be liver, vascular tissue, pituitary tissue, breast tissue, oral squamous cells, colon or kidney.

[1086] Also provided herein are methods for identifying a sample as containing a biological source other than the tissue of the sample. As described above, the methods provided herein can be used to identify the tissue, tumor or other source of an oncofetal fibronectin indicating molecule in a sample. Such methods can be used to identify the oncofetal fibronectin indicating molecule as arising from a biological source other than the tissue of the sample. For example, pancreatic carcinoma typically does not contain EDB+ oncofetal fibronectin protein; thus, if an EDB+ oncofetal fibronectin indicating molecule is identified in a pancreas tissue sample, the sample can be identified as containing a biological material other than pancreatic tissue. In such cases, the biological material other than the tissue of the sample can include, for example metastasized tumor cells. For example, a pancreatic tissue sample containing metastasized liver cells can be a pancreatic tissue sample containing an EDB+ oncofetal fibronectin indicating molecule and thus can be identified using the methods provided herein as a sample containing metastasized tumor cells and not tumorous pancreatic cells or tissue. Thus, methods

provided herein that include characterizing an oncofetal fibronectin indicating molecule in a sample can be used in methods for determining the presence of metastasized tumor cells in a sample and methods for distinguishing between a sample containing metastasized tumor cells and non-metastasized tumor cells.

[1087] Methods provided herein also can be used to characterize the composition of an oncofetal fibronectin indicating molecule of a particular tissue, tumor or biological source. For example, extracellular matrix containing newly formed or forming vasculature can contain oncofetal fibronectin protein. The methods provided herein can be used to characterize the oncofetal fibronectin protein in the extracellular matrix as containing or not containing EDA, EDB, IIICS (and splice variants thereof) and one or more post translational modifications such as O-glycosylation of threonine 33 of IIICS. The methods provided herein can be used to characterize an oncofetal fibronectin indicating molecule present in any tissue of a subject. Similarly, tissue that is benign, hyperplastic, neoplastic, fetal, male, female, or combinations thereof (e.g., neoplastic male liver tissue, neoplastic female liver tissue), can be used as samples in the methods provided herein and any oncofetal fibronectin indicating molecule present in the sample can be characterized as containing or not containing EDA, EDB, IIICS (and splice variants thereof) and one or more post translational modifications such as O-glycosylation of threonine 33 of IIICS.

H. Combinations, Probes, Conjugates and Kits

[1088] Combinations and kits containing the combinations also are provided. The combinations include one or more fibronectin or oncofetal fibronectin binding partners; and, optionally, reagents for detecting at least one fibronectin or oncofetal fibronectin binding partner. The combination can include one or more fibronectin or oncofetal fibronectin binding partners immobilized to a solid support, such as a microtiter plate, a microarray, a membrane or a test strip. The combinations also include one or more solutions (e.g., buffer solution) with which the sample can be mixed. The combinations also can include one or more filters for removing particulate or solid or undissolved matter from a liquid sample. The combinations also can include a non-specific binding compound, dry or in solution and/or a solid support containing a non-specific binder. The combinations also can include a structure for immobilizing or manipulating sample components, such as an electrophoresis gel for manipulation of a sample, or a microtiter plate, a microarray, or membrane for immobilizing sample components. When used in connection with pre-term delivery, the combination also optionally contains a tocolytic agent and, optionally, a device for administering the tocolytic agent. When used in connection with inducing delivery, the combination optionally contains an inducing agent and, optionally, a device for administering the inducing agent. The combinations and kits optionally include instructions for collecting the sample and/or performing the assay. Any of a variety of combinations and kits known in the art that can be adapted for use in the methods provided herein, using techniques known to those skilled in the art, are contemplated herein, as exemplified in U.S. Pat. Nos. 5,281,522, 6,394,952, and 6,267,722. The combinations and kits can be used for detection of oncofetal fibronectin in a sample, for determining the amount of an oncofetal fibronectin indicating molecule in a sample and/or

for characterizing one or more oncofetal fibronectin indicating molecules in a sample.

[1089] The combinations also can include a device or solution for collecting or contacting a sample, such as, but not limited to, a polyester swab, a urine specimen cup, a lavage fluid, a dipstick, a passive sample collection device, or a transdermal patch. Exemplary swabs of the present combinations can include swabs long enough to insert into the vagina, but not long enough to contact the cervix; for example, the length of the swab can be 15 cm or less, 13 cm or less, 12 cm or less, 11 cm or less, 10 cm or less, 9 cm or less, 8 cm or less, 7 cm or less, 6 cm or less, 5 cm or less, 4 cm or less, 3 cm or less, or 2 cm or less. Swabs also can be a tissue or pad (e.g., a gauze pad) that can be used to wipe the vaginal vestibule and/or the labia, and/or can be used to wipe portions of the lower third of the vagina. Exemplary passive sample collection devices include absorbent devices, devices that can be inserted into the vagina, such as into the portion of the vagina below the posterior fornix including the lower third of the vagina, devices that contact the labia and/or vaginal orifice, devices that can be located between the labia and/or vaginal orifice and the subject's undergarment. Sample collection devices also can have immobilized thereto one or more fibronectin or oncofetal fibronectin binding partners.

[1090] Swabs, passive collection devices, and other sample collection devices that are inserted into the vagina in collecting the sample can optionally have attached thereto an overinsertion preventing device. An overinsertion preventing device can limit the distance into the vagina that the sample collection device can be inserted, and thereby ensure that the sample collection device can be readily withdrawn and/or ensure that the sample does not contact the cervix or other portion of the cervicovaginal cavity that could be damaged by unintended contact with a sample collection device. An overinsertion preventing device also can serve to standardize the location in the vagina at which the sample is collected, by ensuring that the sample is collected no further than a particular distance into the vagina, and, optionally, by demarking the location in the vagina at which some or most of the sample is collected. For example, an overinsertion preventing device that is 5 cm from the polyester tip of a swab can be used to ensure that the polyester tip collects fluid no further than 5 cm into the vagina, and, optionally, that most of the cervicovaginal fluid absorbed by the polyester tip is cervicovaginal fluid located about 5 cm from the vaginal vestibule or vaginal orifice. In one example of an overinsertion device, a long narrow swab, such as a swab with a solid shaft and a polyester tip at one end, can have attached thereto a broad shield that is sufficiently large to not enter the vagina and thereby prevent the swab from being further inserted into the vagina. Any of a variety of overinsertion preventing devices can be used, and typically an overinsertion preventing device portion of a sample collection device is larger than the portion of the sample collection device that is inserted into the vagina. Exemplary overinsertion preventing devices include, but are not limited to, a shield, a handle, or a three-dimensional structure (e.g., a sphere or cube). In some embodiments, an overinsertion preventing device that standardizes the location in the vagina at which the sample is collected can increase the reproducibility of sample collection.

[1091] Sample collection devices also can be selected for their ease of use. In some embodiments, the sample collection device can be a device that can be operated by the subject from whom the sample is to be collected or a non-medical professional such as a relative of the subject. Such sample collection device also can be accompanied by instructions for use that instruct the unskilled person on the methods for collecting the sample.

[1092] The combinations can include test strips or other devices containing one or more fibronectin or oncofetal fibronectin binding partners, where such devices can be used to detect the presence of an oncofetal fibronectin indicating molecule in a sample or to indicate the amount of an oncofetal fibronectin indicating molecule in a sample. In some embodiments, such a device can be configured to indicate a positive result when the amount of an oncofetal fibronectin indicating molecule is above a defined threshold level. Exemplary threshold levels include 1 ng/ml, 3 ng/ml, 5 ng/ml, 10 ng/ml, 15 ng/ml, 25 ng/ml, 35 ng/ml and 50 ng/ml, or about 1 ng/ml, about 3 ng/ml, about 5 ng/ml, about 10 ng/ml, about 15 ng/ml, about 25 ng/ml, about 35 ng/ml and about 50 ng/ml, for untreated samples, samples before treatment, or undiluted samples. Other exemplary threshold levels include 1 ng/ml, 2 ng/ml, 3 ng/ml, 5 ng/ml, 7 ng/ml, 10 ng/ml, 15 ng/ml and 20 ng/ml, or about 1 ng/ml, about 2 ng/ml, about 3 ng/ml, about 5 ng/ml, about 7 ng/ml, about 10 ng/ml, about 15 ng/ml and about 20 ng/ml, for buffer-treated samples. Methods for configuring devices such as test strips to return a positive result upon the presence of a threshold level of a sample component are known in the art. The test strip can be readable by a machine, such as a test strip reader, or can be configured to be read by an individual and not by a machine.

[1093] The combinations can include test strip readers or other devices such as absorbance devices, fluorescence devices, that can be used to determine the amount of an oncofetal fibronectin indicating molecule in a sample. In some embodiments, such a device can be configured to indicate a positive result when the amount of an oncofetal fibronectin indicating molecule is above a defined threshold level. Exemplary threshold levels include 1 ng/ml, 3 ng/ml, 5 ng/ml, 10 ng/ml, 15 ng/ml, 25 ng/ml, 35 ng/ml and 50 ng/ml, or about 1 ng/ml, about 3 ng/ml, about 5 ng/ml, about 10 ng/ml, about 15 ng/ml, about 25 ng/ml, about 35 ng/ml and about 50 ng/ml, for untreated samples or samples before treatment. Other exemplary threshold levels include 1 ng/ml, 2 ng/ml, 3 ng/ml, 5 ng/ml, 7 ng/ml, 10 ng/ml, 15 ng/ml and 20 ng/ml, or about 1 ng/ml, about 2 ng/ml, about 3 ng/ml, about 5 ng/ml, about 7 ng/ml, about 10 ng/ml, about 15 ng/ml and about 20 ng/ml, for buffer-treated samples.

[1094] Kits are packaged in combinations that optionally include other reagents or devices. For example, a kit optionally includes one or more devices for obtaining and manipulating a sample from the subject (e.g., a vaginal accessing tool or a ductal accessing tool). A kit also optionally includes one or more devices for transferring or mixing a sample, such as a dropper or pipette. In one embodiment, a kit includes a test strip that contains one or more fibronectin or oncofetal fibronectin binding partners located thereon. A test strip can contain a fibronectin or oncofetal fibronectin binding partner immobilized on the test strip, or mobilizable upon contact with the sample, or can contain an immobilized oncofetal fibronectin binding partner and a mobilizable

oncofetal fibronectin binding partner. Optionally, a test strip-containing kit can include a fibronectin or oncofetal fibronectin binding partner in solution or mobile form, with which the sample can be mixed prior to applying the sample to the test strip. One or more of the fibronectin or oncofetal fibronectin binding partners can be conjugated to a moiety for detection of the conjugate. A kit containing a test strip also can include a non-specific binder such as a non-specific binding compound or a non-specific binding surface, either as a component of a test strip or as a different structure, composition or solution. In one example, a kit contains all necessary components for performing a home test for the presence of an oncofetal fibronectin indicating molecule in a sample, such as, for example, a sample collection device or vesicle, a test strip, instructions for sample collection, use of the test strip and interpretation of the test strip results and optionally one or more compounds, compositions, buffers or solutions for mixing with the sample or for use in conjunction with detection of an oncofetal fibronectin indicating molecule. Typically, a kit that can be used for home testing does not include a swab long enough to contact the cervix when held by an individual.

[1095] The combinations provided herein also include a combination that has been contacted by a sample. The combinations can include one or more fibronectin or oncofetal fibronectin binding partners that have been contacted by a sample; and, optionally, reagents for detecting at least one fibronectin or oncofetal fibronectin binding partner, including reagents for detecting fibronectin or oncofetal fibronectin binding partner that have been contacted with a test device and/or fibronectin or oncofetal fibronectin binding partner. The combination can include a solid support, such as a microtiter plate, a microarray, a membrane or a test strip including, for example, a solid support having one or more fibronectin or oncofetal fibronectin binding partners immobilized thereto, where the solid support has been contacted with a sample. The combinations also can include a sample that has been mixed with one or more solutions (e.g., buffer solution). The combinations also can include one or more filters that has been contacted with a liquid sample. The combinations also can include a non-specific binding compound, dry or in solution and/or a solid support containing a non-specific binder that has been contacted by a sample. The combinations also can include a structure having immobilized thereto, or containing, one or more sample components, such as an electrophoresis gel containing sample components, or a microtiter plate, a microarray, or membrane having immobilized thereto sample components. An exemplary combination can be a combination that indicates the presence of an oncofetal fibronectin indicating molecule in the sample, or an amount of an oncofetal fibronectin indicating molecule in the sample, or an amount of an oncofetal fibronectin indicating molecule in the sample at or above a threshold level. Another exemplary combination can be a combination that indicates the absence of an oncofetal fibronectin indicating molecule in the sample, or an amount of an oncofetal fibronectin indicating molecule in the sample below a threshold level.

[1096] Also provided herein are kits that contain the combinations provided herein and one or more devices such as a thermal cycler, an apparatus for sample preparation such as a tissue homogenizer, as known in the art or provided herein. A kit also can include the appropriate buffers and

solutions for performing the oncofetal fibronectin indicating molecule detection methods described herein.

[1097] Also provided herein are systems including one or more fibronectin or oncofetal fibronectin binding partners and one or more devices for detecting the presence of an oncofetal fibronectin indicating molecule in a sample. The binding partners can be mobile in solution, mobilizable on a solid support, or immobilized on a solid support such as a test strip, a mass spectrometry substrate or a DNA array chip. Devices for detecting the presence of an oncofetal fibronectin indicating molecule in a sample can be any of a variety of detection devices including mass spectrometer, absorbance spectrometer, fluorescence spectrometer, reflectance reader, flow cytometer, or electrophoretic gel scanner. In one embodiment, the device for detecting the presence of an oncofetal fibronectin indicating molecule is designed to receive a solid support containing a sample or sample components and, optionally also containing a fibronectin or oncofetal fibronectin binding partner.

[1098] The packaging material used in the kit can be one or more physical structures used to house the contents of the kit and can be constructed by well known methods, typically to provide a sterile, contaminant-free environment. The packaging material can have a label which indicates the components of the kit. In addition, the packaging material contains instructions indicating how the materials within the kit are employed to determine the presence of an oncofetal fibronectin indicating molecule, determine the amount of an oncofetal fibronectin indicating molecule, or characterize an oncofetal fibronectin indicating molecule, present in the sample. Instructions typically include a tangible expression describing the reagent concentration or at least one assay method parameter, such as the relative amounts of reagent and sample to be admixed, maintenance time periods for reagent/sample admixtures, temperature, buffer conditions and other parameters. The kit can include one or more containers capable of holding within fixed limits a primer, enzyme or other reactant or buffer solution used in the methods of oncofetal fibronectin indicating molecule determination. For example, a kit can include a glass vial used to contain milligram quantities of a fibronectin or oncofetal fibronectin binding partner. A kit also can include substrates, supports or containers for performing the oncofetal fibronectin indicating molecule determination methods, including vials or tubes, or a mass spectrometry substrate.

[1099] Also provided in the kits herein are systems for classifying the sample according to one or more threshold levels. Samples having amounts of oncofetal fibronectin equal to or greater than one or more threshold(s) are considered positive for oncofetal fibronectin. For instance, a sample having equal to or greater than 25 ng/ml is considered positive for oncofetal fibronectin. On the other hand, samples having amounts of oncofetal fibronectin less than one or more threshold(s) are considered negative for oncofetal fibronectin. For instance, a sample having less than 25 ng/ml is considered positive for oncofetal fibronectin.

[1100] Probes for detection and/or quantitation of oncofetal fibronectin and indicating molecules thereof also are provided herein. The probes include, for example, a mass spectrometry substrate and a fibronectin or oncofetal fibronectin binding partner immobilized on the mass spectrometry substrate for detecting an oncofetal fibronectin indicating molecule. Mass spectrometry substrates are substances such as, but not limited to, glass, metals, ceramics, Teflon-coated magnetic materials, organic polymers,

biopolymers and inorganic polymers. Probes as provided herein can be used to identify oncofetal fibronectin indicating molecules by detecting a particular weight fragment by mass spectrometry.

[1101] Conjugates also are provided herein. A conjugate can include, for example, a fibronectin or oncofetal fibronectin binding partner linked directly or indirectly via a linker to a therapeutic agent or an imaging agent or detecting agent. Therapeutic agents include, but are not limited to a cytokine, a photosensitizing agent, a toxin, an anticancer antibiotic, a chemotherapeutic compound, a radionuclide, and a bioluminescent compound or to a detectable moiety such as, for example, a fluorescent moiety, a radionuclide, a magnetically detectable isotope or compound, a sonographic imaging agent, a chromophore, a latex microsphere, or a quantum dot. In certain cases, the therapeutic agent is an angiogenesis inhibitor. In a particular case, a therapeutic agent can be pseudomonas exotoxin, diphtheria toxin, ricin, cholera toxin, gelonin, shigella toxin, pokeweed antiviral protein, exotoxin A, abrin toxin, saporin, an interleukin, a tumor necrosis factor, an interferon, granulocyte macrophage colony stimulating factor, angiogenin, tissue factor, porfirimycin, doxorubicin, dactinomycin, plicamycin, mitomycin, bleomycin, actinomycin, daunorubicin, indocyanine green, toluidine blue, aminolevulinic acid, texaphyrin, benzoporphyrin, a phenothiazine, a phthalocyanine, a porphyrin, a chlorin, a purpurin, a purpurinimide, a bacteriochlorin, a pheophorbide, a pyropheophorbide, a cationic dye, ³²Phosphate, ⁶⁰Cobalt, ⁹⁰Yttrium, ⁹⁹Technetium, ¹⁰³Palladium, ¹⁰⁶Ruthenium, ¹¹¹Indium, ¹¹⁷Lutetium, ¹²⁵Iodine, ¹³¹Iodine, ¹³⁷Cesium, ¹⁵³Samarium, ¹⁸⁶Rhenium, ¹⁸⁸Rhenium, ¹⁹²Iridium, ¹⁹⁸Gold, ²¹¹Astatine, ²¹²Bismuth, ²¹³Bismuth, 5-fluorouridine, calicheamicin or maytansine. Therapeutic agents can also be signaling modulators, such as, for example, an inhibitor of MIF (macrophage inhibitory factor), a toll-like receptor agonist, or a stat 3 inhibitor. Binding partners of the conjugates can be antibodies, such as, for example, FDC-6, BC-1, ME4C or L19.

[1102] Provided herein is the use of any of the products provided herein for the preparation of a medicament for any of the methods provided herein, including, but not limited to, diagnosis, imaging, and/or treatment of a health disorder characterized by the presence of, or elevated levels of, an oncofetal fibronectin indicating molecule.

[1103] The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

I. EXAMPLES

Example 1

Polyclonal Anti-Oncofetal Fibronectin Antibody

[1104] Oncofetal fibronectin can be purified from amniotic fluid as described by Engvall and Ruoslahti, *Int. J. Cancer*, 20:1-5 (1977).

[1105] The anti-(oncofetal fibronectin) antibodies are elicited in rabbits using the immunization techniques and schedules described in the literature, e.g., Stollar, *Meth. Enzymol.*, 70(A):70-85 (1980), immunizing the rabbits with the oncofetal fibronectin antigen. The antiserum is screened in a solid phase assay similar to that used for monoclonal antibodies, such as described by Lange et al., *Clin. Exp. Immunol.* 25(2):191-198 (1976) and Pisetsky et al., *J. Immun. Meth.* 41(2):187-200 (1981).

[1106] The IgG fraction of the antisera is purified further by affinity chromatography using CNBr-Sepharose 4B (Pharmacia Fine Chemicals) to which has been coupled oncofetal fibronectin. The method used for coupling is that recommended by the gel manufacturer, AFFINITY CHROMATOGRAPHY, *Pharmacia Fine Chemicals*, pp. 15-18.

[1107] Briefly, the column is equilibrated with from 2 to 3 volumes of buffer (0.01 M PBS, pH 7.2) and the anti-(oncofetal fibronectin) antibody containing solution then is applied to the column. The absorbency of the eluate is monitored at 280 nm until protein no longer passes from the column. The column then is washed with 0.1 M glycine buffer, pH 2.5, to desorb the immunoaffinity bound anti-(oncofetal fibronectin) antibody. Peak protein fractions are collected, pooled and dialyzed against 0.01 M PBS, pH 7.2, for 24-36 hr at 4° C. with multiple buffer changes.

[1108] If a purity of preferentially binding antibodies is to be achieved, the affinity purified IgG can be passed through an adult plasma fibronectin bound affinity column by the procedure described above to remove antibodies that would cross-react with non-oncofetal fibronectin.

Example 2

Monoclonal Anti-Oncofetal Fibronectin Antibody

[1109] Using the purified oncofetal fibronectin obtained by the procedure of Example 1, mouse monoclonal antibodies to the oncofetal fibronectin are obtained using standard procedures of Galfre and Milstein, *Meth. Enzymol.*, 73(Pt.B):3-46 (1981); and Matsuura, H. and Hakomori, S. et. al. *Proc. Natl. Acad. Sci. USA*, 82:6517-6521 (1985), using oncofetal fibronectin as the antigen for immunizing the mice. The monoclonal antibodies are screened using a modification of the techniques described in the literature, e.g., Lange et al., *Clin. Exp. Immunol.*, 25(2):191-198 (1976); and Pisetsky et al., *J. Immun. Meth.*, 41(2):187-200 (1981).

[1110] Mouse monoclonal antibody is purified from ascites fluid or from hybridoma culture supernatants using Protein-A coupled Sepharose-4B (Pharmacia Fine Chemicals) according to the procedure of Tijssen, *Practice and Theory of Enzyme Immunoassays*, Elsevier Science Publishers, pp. 105-107 (1985).

Example 3

Polyclonal Anti-Oncofetal Fibronectin Antibody-Coated Microtiter Plate

[1111] Rabbit anti-(oncofetal fibronectin) prepared and further purified to remove non-oncofetal fibronectin cross-reactivity as described in Example 1 is diluted to 10 µg/mL in 0.05 M carbonate buffer, pH 9.6. 100 µL is dispersed into each well of an IMMULON II microtiter plate (Dynatech). The plate is covered and incubated 4 hr at room temperature or 4° C. overnight. The plate is washed 4 times with Wash Buffer (0.02 M Tris HCl, 0.015 M NaCl, 0.05% TWEEN-20), filling and emptying the wells completely with each use. The plate then is blocked by dispersing into each well 200 µL of a blocking solution (0.01 M PBS, 1% BSA, 0.02% NaN₃, pH 7.4) and incubating for 1 hr at room temperature. The wells then are washed 4 times with Wash Buffer, as described above. The plate is now ready for immunoassay of samples.

Example 4

Polyclonal Anti-Human Fibronectin Antibody

[1112] Human plasma fibronectin was purified from human plasma as described by Engvall and Ruoslahti, *Int. J. Cancer*, 20:1-5 (1977).

[1113] The anti-human plasma fibronectin antibodies were elicited in goats using the immunization techniques and schedules described in the literature, e.g., Stollar, *Meth. Enzymol.*, 70(A):70-85 (1980), immunizing the goats with the human plasma fibronectin antigen. The antiserum was screened in a solid phase assay similar to that used for monoclonal antibodies, e.g., as described by Lange et al., *Clin. Exp. Immunol.*, 25(2):191-198 (1976) and Pisetsky et al., *J. Immun. Meth.*, 41(2):187-200 (1981).

[1114] The IgG fraction of the antiserum was purified further by affinity chromatography using CNBr—Sepharose 4B (Pharmacia Fine Chemicals) to which has been coupled human plasma fibronectin according to the method recommended by the manufacturer (AFFINITY CHROMATOGRAPHY, *Pharmacia Fine Chemicals Catalogue*, 1990, pp. 15-18).

[1115] Briefly, the column was equilibrated with from 2 to 3 volumes of buffer (0.01 M PBS, pH 7.2) and the anti-human fibronectin antibody-containing solution was then applied to the column. The absorbency of the effluent was monitored at 280 nm until protein no longer passed from the column. The column was then washed with equilibration buffer until a baseline absorbance at 280 nm was obtained.

[1116] The immunoaffinity bound anti-human plasma fibronectin antibody was eluted with 0.1 M glycine buffer, pH 2.5. Peak protein fractions were collected, pooled and dialyzed against 0.01 M PBS, pH 7.2, for 24-36 hr at 4° C. with multiple buffer changes. The above procedure was repeated to immunize rabbits with human plasma fibronectin and to purify the resultant polyclonal anti-human fibronectin antibodies.

Example 5

Polyclonal Anti-Fibronectin Antibody-Coated Microtiter Plate

[1117] Goat anti-human plasma fibronectin prepared as described in Example 4 is diluted to 10 µg/mL in 0.05 M carbonate buffer, pH 9.6. 100 µL is dispersed into each well of a polystyrene microtiter plate such as supplied by Costar, Nunc, or Dynatech. The plate is covered and incubated 2 to 4 hr at room temperature or 4° C. overnight. The plate is washed 3 to 4 times with Wash Buffer (0.02 M Tris HCl, 0.015 M NaCl, 0.05% TWEEN-20), filling and emptying the wells completely with each use. The plate then is blocked by dispersing into each well 200 µL of a blocking/stabilizing solution (4% sucrose, 1% mannitol, 0.01 M PBS, 1% BSA, 0.02% NaN₃, pH 7.4) and incubated for 30 minutes to 2 hrs at room temperature. The wells then are aspirated to dryness, the plate is packaged in an air-tight container with a desiccant pouch and stored at 4° C. until needed.

Example 6

Monoclonal Antibodies from Hybridoma HB 9018

[1118] Preparation of the Hybridoma deposited at the American Type Culture Collection and given the accession number ATCC HB 9018 is described U.S. Pat. No. 4,894,

326 issued Jan. 16, 1990 to Matsuura et al. The hybridoma was cultured by growth in RPMI 1640 tissue culture medium supplemented with 10% fetal bovine serum. Additionally, the hybridoma was cultured in mice by the injection of the hybrid cells according to the procedure of Mishell and Shiigi, *Selected Methods in Cellular Immunology*, W.H. Freeman & Co, San Francisco, page 368, (1980).

[1119] The monoclonal antibody, designated FDC-6, was produced by the hybridoma was prepared for use in an immunoassay by the following procedure. The IgG fraction of the culture supernatant or the ascites was precipitated by ammonium sulfate fractionation. The antibody was redissolved and dialyzed into the appropriate buffer for purification by affinity chromatography on Protein-G Fast Flow (Pharmacia Fine Chemicals) according to the manufacturer's directions.

Example 7

Monoclonal Antibody-Coated Microtiter Plate

[1120] Microtiter plates were coated with FDC-6 monoclonal antibody by following the procedure described below.

[1121] Monoclonal antibody FDC-6 prepared as described in Example 6 was diluted to 10 $\mu\text{g}/\text{mL}$ in phosphate buffer, pH 7.2 and 100 $\mu\text{l}/\text{well}$ was dispersed into a polystyrene microtiter plate (Costar). The plates were incubated for 2 hours at room temperature or overnight at 4° C. The contents of the wells were aspirated and the wells washed 3 to 4 times with wash buffer (0.02 M Tris HCl, 0.015 M NaCl and 0.05% TWEEN-20) as described in Example 5. 200 $\mu\text{l}/\text{well}$ of blocking/stabilizing solution (4% sucrose, 1% mannitol, 0.5% casein and 0.01 M PBS) was then added to the wells and incubated for 30 minutes to 4 hours at room temperature. The wells were then aspirated to dryness and the plate was packaged in an air-tight container with a desiccant pouch and stored at 4° C. until needed.

[1122] The above procedure was repeated using microtiter plates from Nunc and Dynatech and gave equivalent results.

Example 8

Enzyme Labeled Anti-Fibronectin Antibody

[1123] Anti-human plasma fibronectin antibody prepared according to Example 4 was conjugated with calf intestine alkaline phosphatase following the one-step glutaraldehyde procedure of Avrameas, *Immunochem.* 6:43 (1969).

Example 9

Oncofetal Fibronectin Assay Kit and Method

[1124] An assay kit for oncofetal fibronectin included the following reagents:

[1125] 1. a microtiter plate coated with affinity-purified murine monoclonal anti-oncofetal fibronectin antibody,

[1126] 2. calf intestine alkaline phosphatase-conjugated, affinity purified, polyclonal, goat anti-fibronectin antibodies,

[1127] 3. enzyme substrate,

[1128] 4. a negative control,

[1129] 5. a positive control,

[1130] 6. rinse buffer concentrate (50 \times), and

[1131] 7. stopping solution.

[1132] The microtiter plate coated with murine monoclonal anti-oncofetal fibronectin antibody and the alkaline phosphatase-conjugated, affinity purified, polyclonal, goat anti-fibronectin antibodies were prepared as described in Examples 7 and 8, respectively. The microtiter plate was packaged as 12 strips of eight wells each in sealed plastic bags containing desiccant and stored at 2° C. to 8° C.

[1133] The stock antibody conjugate was appropriately diluted in conjugate diluent (0.05 M Tris Buffer pH 7.2, 2% D-Sorbitol, 2% BSA, 0.1% Sodium Azide, 0.01% Tween-20, 1 mM Magnesium Chloride and 0.1% Zinc Chloride) and 10 mL placed in a polyethylene dropper bottle container.

[1134] The enzyme substrate (10 mL in a polyethylene dropper bottle container) was phenolphthalein monophosphate (1 mg/mL) dissolved in 0.4 M aminomethylpropanediol buffer, pH 10 with 0.1 mM magnesium chloride and 0.2% sodium azide.

[1135] The positive control (3.4 mL in a polyethylene dropper bottle container) was amniotic fluid containing oncofetal fibronectin diluted to a concentration of oncofetal fibronectin of 50 ng/mL in sample diluent solution (0.05 M Tris buffer pH 7.4, 1% bovine serum albumin (BSA), 0.15 M sodium chloride, 0.02% Sodium Azide, 5 mM ethylenediamine tetraacetic acid (EDTA), 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/mL of Aprotinin and 0.1% Triton X-100). This sample diluent solution is described in U.S. Pat. No. 4,919,889 to Jones et al., issued Apr. 24, 1990.

[1136] The negative control (2.5 mL in a polyethylene dropper bottle container) was the sample diluent solution used for the positive control without oncofetal fibronectin.

[1137] The rinse buffer (20 mL in a polyethylene dropper bottle container) was a 50 \times concentrate containing 1.0 M Tris buffer pH 7.4, 4.0 M sodium chloride, 2.5% Tween-20 and 1% sodium azide. The rinse buffer was diluted with distilled or deionized water to a final concentration of 0.02 M Tris, 0.08 M sodium chloride, 0.05% Tween-20 and 0.02% sodium azide for use in the assay. The stopping solution (10 mL in a polyethylene dropper bottle container) contained 50 mM EDTA and 50 mM sodium phosphate.

[1138] In addition, 5 μm pore size polyethylene sample filters (Porex Technologies, Fairburn, Ga.), a microtiter strip holder, a microtiter plate cover, storage tubes, and an instruction sheet. All of the dropper bottles in the kit were polyethylene bottles designed to dispense approximately 50 μL drops of the reagent. All of the assay steps performed following sample collection used the reagents and materials in the kit. All samples, positive and negative controls were tested at the same time using kit reagents from the same kit.

[1139] The assay was performed as follows. All samples were collected in the vicinity of the posterior fornix or cervical os using polyester swabs. Swab samples were immersed in 1.0 mL of sample diluent in a collection vial.

The sample diluent solution is described above. The swabs were removed from the solution leaving as much liquid as possible in the collection tube. All samples and control reagents were pre-warmed to 37° C. for 20 minutes in an incubator or 10 minutes in a 37° C. water bath prior to the assay, either before or after filtration. A sample filter was snapped in place on each sample tube and the filter pushed to the bottom until the entire sample is filtered. For samples of which filtering were not effective, samples were centrifuged at 550×g at room temperature for 5 minutes and test supernatant used in assay. The 8-well strips were snapped into place in a strip holder. The holder had the alphanumeric indications of the 12 columns and eight rows of standard microtiter plates. Duplicate 100 μ L aliquots of each sample and the pre-warmed positive and negative controls were placed in separate wells of the microtiter strip and incubated for 1 hour at room temperature, covered.

[1140] Following incubation, samples and controls were aspirated from the wells. Wells were washed three times with diluted wash buffer (1×), being sure wells were completely filled each time. Following washing, 100 μ L of enzyme-antibody conjugate was added to each well and incubated for 30 minutes at room temperature, covered. The wells were aspirated and washed as described above. Following washing, 100 μ L of enzyme substrate was added to each well and incubated for 30 minutes at room temperature, covered. After the 30 minute incubation, 50 μ L of stopping solution was added to each well.

[1141] Plates were gently agitated by hand or with an orbital shaker to mix the well contents. The frame of strips was placed in an ELISA plate reader. The absorbance of each well at 550 nm was determined. The average absorbance of the duplicate wells for each sample and control was calculated. If the absorbance of the subject sample was less than the absorbance of the positive control, the sample was negative, indicating an undetectable level of oncofetal fibronectin in the sample. If the sample absorbance is greater than or equal to the absorbance of the positive control, the sample was positive, indicating that oncofetal fibronectin was present in the sample. In any assay if the average absorbance of the positive control was not at least 0.02 absorbance units greater than the average absorbance of the negative control, the results were discarded and the assay procedure was repeated.

Example 10

Pre-Term Labor Sandwich Immunoassay

[1142] The procedure of Example 9 was repeated with test samples obtained during weeks 20-36 of pregnancy. Studies were conducted at three perinatal referral clinics in the United States. Women were evaluated for admission to the hospital for either suspected pre-term rupture of membranes or suspected pre-term labor with intact membranes.

[1143] Confirmation of rupture of membranes was made by visual examination of the vagina for gross pooling of amniotic fluid, microscopic examination of dried vaginal secretions for ferning, presence of alkaline vaginal secretions using nitrazine paper and ultrasound diagnosis of oligohydramnios. Rupture of membranes was defined by the presence of any two of these four diagnostic criteria. One hundred-seventeen women with intact amniotic membranes

pregnant between 23 weeks and 36 weeks, 6 days of gestation based on last known menstrual period and expected date of confinement confirmed by first trimester pelvic examination and ultrasonography <28 weeks gestation are subsequently described. Women were determined by the attending physician to be at risk of pre-term labor and subsequent delivery based on medical history and clinical examination including recording of uterine contractions and examination of the cervix. Since the clinical definition of pre-term labor is sometimes difficult to establish, data establishing the clinical utility of oncofetal fibronectin were analyzed using pre-term delivery as the outcome variable.

[1144] To assess the potential for cervicovaginal contamination by maternal plasma fibronectin, maternal blood specimens were obtained from 52 women with apparently healthy pregnancies during second or third trimester. Amniotic fluid specimens were obtained from 92 subjects undergoing amniocentesis for genetic diagnosis in early second trimester and 8 subjects undergoing amniocentesis for evaluation of fetal lung maturity prior to elective repeat, cesarean section in third trimester. The assay results indicated that the concentration of oncofetal fibronectin in amniotic fluid in second trimester was $87.1 \pm 4.8 \mu\text{g/mL}$ (n=92) and $27.1 \pm 17.3 \mu\text{g/mL}$ (n=8) in third trimester. The concentration of oncofetal fibronectin in maternal plasma in the second trimester was $1.48 \pm 0.11 \mu\text{g/mL}$ (n=20) and $3.19 \pm 0.30 \mu\text{g/mL}$ (n=32) in the third trimester.

[1145] For the 117 subjects with suspected pre-term labor and intact amniotic membranes, 49 of 59 (sensitivity=83.1%) women delivering prematurely (PTD) had oncofetal fibronectin in their cervicovaginal secretions compared to 11 of 58 women (specificity=81.0%) delivering at term (TD) (p<0.01). Similarly, those subjects with oncofetal fibronectin in their cervicovaginal secretions were far more likely to deliver prematurely (positive predictive value=81.7%) than those women not expressing cervicovaginal oncofetal fibronectin (negative predictive value=82.5%).

[1146] The presence of cervicovaginal oncofetal fibronectin was a sensitive and specific predictor of the risk of pre-term delivery in these women with suspected pre-term labor. The presence of oncofetal fibronectin in these subjects was strongly associated with risk of pre-term delivery with a logistic regression odds ratio of 3.79 (95% CI:2.33, 6.15; p<0.01).

[1147] To evaluate for potential confounding by oncofetal fibronectin of maternal origin, the data were analyzed after exclusion of 31 samples contaminated with blood. As shown below, similar proportions of subjects had oncofetal fibronectin in their cervicovaginal secretions and delivered prematurely. Furthermore, inclusion of the presence or absence of vaginal bloody show into the stepwise logistic regression model gave an odds ratio of 1.70 (95% CI:0.91, 3.18; p=0.1) demonstrating that bloody show was not an independent predictor of pre-term delivery after oncofetal fibronectin was introduced into the model. It was clear, however, from univariate analysis that detection of oncofetal fibronectin in cervicovaginal secretions contaminated with blood is an indicator of imminent delivery.

[1148] The utility of oncofetal fibronectin for identifying women at risk of PTD was maintained even when women in pre-term contractions with intact membranes with cervical dilation exceeding 2 cm were eliminated from the analysis.

The logistic regression odds ratio of 3.18 (95% CI:1.8, 5.6, $p < 0.01$) confirmed the predictive value of oncofetal fibronectin in this clinically discrete population.

Example 11

Oncofetal Fibronectin Assay Kit and Method

[1149] An assay kit for oncofetal fibronectin included the following components. This kit was designed to be used to perform a rapid, bedside assay.

[1150] 1. an assay device having a plastic housing and containing:

[1151] (a) a porous nylon membrane to which is bound a monoclonal anti-oncofetal fibronectin antibody;

[1152] (b) a flow control membrane system; and

[1153] (c) an absorbent layer

[1154] 2. a colloidal gold-labeled goat anti-fibronectin antibody conjugate in a protein matrix

[1155] 3. conjugate reconstitution buffer

[1156] 4. a wash solution

[1157] 5. a sterile, polyester sample collection swab

[1158] The membrane device was prepared by the following procedure. Approximately 2 μL of the murine monoclonal antibody FDC-6 prepared as described in Example 6 is applied to a membrane surface (1.2 μm nylon, Biodyne-A, Pall) in a pH 6, 0.01 M phosphate buffered saline (PBS), 0.1 M citrate buffer containing 0.5 mg/mL BSA. A procedural control containing human plasma fibronectin purified as described in Example 4 in the same buffer also is applied to a discrete region of the membrane. After the membrane has air dried, a blocking reagent of PBS-buffered, 0.5% nonfat dry milk is added to the membrane. The excess blocking reagent is removed after at least about 20 minutes.

[1159] The membrane-holding device (Target Device, V-Tech, Pomona, Calif.) is assembled with a second porous layer (0.45 μm low protein-binding nylon, LoProdyne, Pall) beneath the antibody-bearing membrane (in the direction of sample application) for controlling the flow of sample solution from the assay membrane to the absorbent layer. The two porous membranes then are placed over an absorbent porous polyethylene layer having a capacity of greater than 1.5 mL (Chromex, Brooklyn, N.Y.) and enclosed in the device. The device is packaged individually in a sealed plastic bag containing desiccant.

[1160] The colloidal gold is prepared by the reduction of 0.01% tetrachloroauric acid with 0.16% sodium citrate in a manner which produces approximately 30 nm particles. Briefly, the two solutions are heated separately to 90° C. The reducing solution is added to the gold solution while vigorously stirring. The combined solution is boiled (100° C.) for at least 10 minutes.

[1161] Affinity purified goat anti-fibronectin antibody (prepared as described in Example 4) was bound to the colloidal gold by adsorption. Briefly, the colloidal gold solution prepared above was combined with the antibody (5-10 $\mu\text{g}/\text{mL}$) in water. Following conjugation, the conjugate

solution was stabilized by the addition of 5% BSA and 5% polyvinylpyrrolidone (final concentration).

[1162] The stock conjugate was concentrated approximately 10- to 12-fold by ultrafiltration using a hollow fiber filter. The concentrated conjugate was diluted to an appropriate level in 15 mM Tris, 2% BSA, 0.1% Tween 20, 0.2% polyethylene glycol, 8% polyvinylpyrrolidone and 0.04% thimerosal. An appropriate concentration was determined by using a range of dilutions in a sample assay procedure as described below and determining the dilution which produces the best result.

[1163] The selected conjugate dilution is placed in polyethylene sample collection tubes and lyophilized. The tubes are fitted with 2 μm pore size polyethylene sample filters (Porex Technologies, Fairburn, Ga.) during the lyophilization process. The lyophilized conjugate is individually packaged in a foil pouch with desiccant.

[1164] The conjugate reconstitution buffer is 100 mM sodium acetate. This buffer is packaged as a unit dose in a 1 mL disposable tube.

[1165] The wash solution is water packaged as a unit dose in a disposable tube.

[1166] The kit additionally contains an individually packaged sterile polyester swab and a procedural summary card.

[1167] The assay was performed as follows:

[1168] 1. Before collecting the sample, remove the plastic tube containing gold conjugate from the foil pouch, remove the dropper tip and add the entire contents of the tube containing the conjugate reconstitution buffer.

[1169] 2. Collect the sample with the swab provided. During a sterile speculum examination, insert the swab into the posterior fornix of the vagina, swirl for approximately 10 seconds to absorb fluid. Immediately proceed to perform the test. Place the swab in the gold conjugate solution and mix rapidly with an up and down motion for 10 to 15 seconds.

[1170] 3. Remove as much liquid as possible from the swab by rolling the tip on the inside of the tube. Dispose of the swab in a manner consistent with handling potentially infectious materials.

[1171] 4. Replace the dropper tip on the plastic tube and immediately dispense the entire volume of diluted filtered sample onto the surface of the membrane device.

[1172] 5. After the sample liquid has been absorbed into the membrane surface, add a few drops of wash solution and observe the results.

[1173] 6. A negative result is indicated by a red color in the procedural control area of the membrane only. A positive result is indicated by a pink or red spot in the test zone of the membrane as well as in the control zone.

Example 12

Detection of Oncofetal Fibronectin in Cervicovaginal Samples as a Marker for Cervical Cancer

[1174] Swab samples of the cervical os of 15 subjects diagnosed with cervical cancer were collected. Specifically,

a polyester swab (Adeza Specimen Collection Kit, Adeza Biomedical, Inc., Sunnyvale, Calif.) was used to swab either a cervical lesion at the cervical os, or the transition zone at the cervical os and some transition zone swabs also included a swab of the ectocervix. To extract material from each swab into a buffer, each swab was placed into separate vessels containing 1 ml antiprotease buffer (APB) containing 0.05 M Tris buffer, pH 7.4, 150 mM NaCl, 0.02% NaN₃, 1% BSA, 5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin, and 0.1% Triton X-100.

[1175] Lateral flow was performed for each sample by adding 200 μ l of each swab-released APB sample to a test strip containing an absorbent pad which draws the liquid sample to the test strip. The sample first flowed through a mobilization region containing a conjugate of FDC-6 conjugated to blue microspheres (Adeza Biomedical, Inc.). The conjugate was mobilized by the flow of the sample. The sample and conjugate then flowed through a fibronectin or oncofetal fibronectin binding region containing goat polyclonal anti-fibronectin antibodies, such as those described in Example 4 (also referred to herein as A120 antibodies) non-diffusibly bound to the test strip. Finally, the sample flowed through a control region containing goat polyclonal anti-mouse IgG antibody (which selectively binds the FDC-6 blue latex microsphere conjugate) non-diffusibly bound to the test strip. The test strip was then placed in a reader device which measured light reflectance in the detection and control regions.

[1176] Of samples from subjects having cervical cancer, 100% had detectable levels of oncofetal fibronectin in this test (15 out of 15 samples). Similar results were obtained from ELISA assays (see Example 9) of these samples.

Example 13

Detection of Oncofetal Fibronectin in Urine Samples as a Marker for Bladder Cancer

[1177] Urine samples were collected from 44 bladder cancer subjects. Among these, 23 subjects were diagnosed with T1 stage bladder cancer, 7 subjects were diagnosed with T2 stage bladder cancer, 6 subjects were diagnosed with T3 stage bladder cancer, 2 subjects were diagnosed with T4 stage bladder cancer and 6 subjects had an unknown stage bladder cancer. Control samples from 41 subjects free of bladder cancer also were collected. The samples were frozen and stored at -80° C. Before testing, the samples were thawed. Samples were tested neat (i.e., without adding buffer or reagents) or were diluted, as provided below.

[1178] Testing was performed by three different methods: dot blot analysis, dipstick/lateral flow and vertical flow.

[1179] A. Dot Blot

[1180] For this analysis, 49 urine samples were tested: 29 urine samples from subjects with bladder cancer and 20 urine samples from control subjects (positive and negative). Blot assays were performed by incubating the mouse monoclonal anti-oncofetal fibronectin FDC-6 antibody and a horseradish peroxidase-conjugated anti-mouse IgG antibody with a nitrocellulose membrane to which thawed frozen

urine samples were applied. Briefly, frozen urine samples were thawed and were added to a nitrocellulose membrane by applying 5 μ l of each sample to a discrete position on the nitrocellulose membrane. Liquid samples applied to the nitrocellulose membrane were air dried. Non-specific protein binding was blocked by incubating the nitrocellulose membrane in a solution containing 20 mM Tris-HCl, 150 mM NaCl, pH 7.5, 0.05% Tween 20 and 5% BSA (BSA/TBS-T), for 1 hour, after which the BSA/TBS-T solution was removed. Next, primary antibody solution containing 6 μ g/ml FDC-6 in BSA/TBS-T was incubated with the nitrocellulose membrane for 30 minutes. The nitrocellulose membrane was then rinsed three times in a solution containing 20 mM Tris-HCl, 150 mM NaCl, pH 7.5 and 0.05% Tween 20 (TBS-T). Next, secondary antibody solution containing anti-mouse IgG conjugated with horseradish peroxidase (Jackson Immunologicals, West Grove, Pa.) in BSA/TBS-T was incubated for 30 minutes, followed by three rinses with TBS-T, followed by two rinses in a solution containing a solution containing 20 mM Tris-HCl and 150 mM NaCl, pH 7.5 (TBS). The nitrocellulose membrane was then briefly incubated with equal amounts of enzyme chemiluminescence solutions 1 and 2 (Amersham ECL Western Blotting Detection Reagents, Cat. No. RPN2109; Amersham Biosciences Corp., Piscataway, N.J.); the excess liquid was removed and the nitrocellulose membrane was wrapped in plastic wrap, placed in a film cassette and exposed to film for 1 minute. The film was developed and scanned for exposure to chemiluminescence. Confirmation that the signals were indicative of oncofetal fibronectin was performed by Western blot analysis.

[1181] Of samples from subjects having bladder cancer, 79% were positive for oncofetal fibronectin in this test (23 out of 29 samples). Of samples from subjects not having bladder cancer, 90% were negative for oncofetal fibronectin in this test (18 out of 20 samples). Thus, presence of oncofetal fibronectin had a 92% positive predictive value for the presence of bladder cancer and absence of oncofetal fibronectin had a negative predictive value of 75% for the absence of bladder cancer. Determination of being positive for oncofetal fibronectin was made according to the presence of any signal above background.

[1182] B. Dipstick/Lateral Flow

[1183] For this analysis, 59 urine samples were tested: 35 urine samples from subjects with bladder cancer and 24 urine samples from control (positive and negative) subjects. Lateral flow was performed by adding 100 μ l of urine sample to 300 μ l of APB (0.05 M Tris buffer, pH 7.4, 150 mM NaCl, 0.02% NaN₃, 1% BSA, 5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin and 0.1% Triton X-100), and then by applying 200 μ l of this diluted urine sample mixture to a test strip containing an absorbent pad which draws the sample to the test strip. The sample first flowed through a mobilization region containing a conjugate of FDC-6 conjugated to blue microspheres (Adeza Biomedical, Inc.) diffusibly bound to the test strip. The conjugate was mobilized

by the flow of the sample. Next, the sample and conjugate flowed through a non-specific binding trap region containing methylated BSA attached to the test strip. The sample and conjugate then flowed through a region containing goat polyclonal anti-fibronectin antibodies, such as those described in Example 4 (also referred to herein as A120 antibodies), non-diffusably bound to the test strip. Finally, the sample and conjugate flowed through a control region containing goat polyclonal anti-mouse IgG antibody (which selectively binds the FDC-6 conjugate) non-diffusably bound to the test strip. The test strip was then visually inspected for the presence or absence of blue color in the detection and control regions.

[1184] Of samples from subjects having bladder cancer, 66% were positive for oncofetal fibronectin in this test (23 out of 35 samples). Of samples from subjects not having bladder cancer, 88% were negative for oncofetal fibronectin in this test (21 out of 24 samples). Thus, presence of oncofetal fibronectin had a 89% positive predictive value for the presence of bladder cancer and absence of oncofetal fibronectin had a negative predictive value of 64% for the absence of bladder cancer. Determination of being positive for oncofetal fibronectin was made according to the presence of any signal above background.

[1185] C. Vertical Flow

[1186] Vertical flow analysis was used to examine 43 urine samples of subjects with bladder cancer and 42 urine samples from control subjects. Vertical flow was performed by dissolving a lyophilized polyclonal goat anti-human fibronectin antibody/colloidal gold conjugate in a reconstitution buffer containing 100 mM NaOAc and 3 mM Na₂S₂O₃. Six drops (about 150-300 μ l) of urine sample was added to 1 ml of the reconstituted conjugate solution. The sample mixture was added to the surface of a membrane containing beneath the membrane an absorbent pad which draws the liquid sample through the membrane. In the center of the test strip is a fibronectin or oncofetal fibronectin binding region containing FDC-6 (ATCC HB-9018) non-diffusably bound to the test strip. Separate from and encircling the FDC-6 region is a ring-shaped control region containing fibronectin that can specifically bind to the polyclonal goat anti-human fibronectin antibody/colloidal gold conjugate. After the sample passed through the membrane, presence of a ring along the periphery detected by visual inspection indicated that the test was complete and that the anti-fibronectin/colloidal gold conjugate migrated along the test strip. Presence of a pink or purple spot within the ring detected by visual inspection indicated that the test was positive for oncofetal fibronectin.

[1187] Of samples from subjects having bladder cancer, 54% were positive for oncofetal fibronectin in this test (23 out of 43 samples). Of samples from subjects not having bladder cancer, 67% were positive for oncofetal fibronectin in this test (28 out of 42 samples). Thus, presence of oncofetal fibronectin had a 62% positive predictive value for the presence of bladder cancer and absence of oncofetal fibronectin had a negative predictive value of 58% for the

absence of bladder cancer. Determination of being positive for oncofetal fibronectin was made according to the presence of any signal above background.

[1188] D. BTA Stat

[1189] For this analysis, 68 urine samples were tested: 26 urine samples from subjects with bladder cancer and 42 urine samples from control (positive and negative) subjects. Blot assays were performed by adding 5 drops of urine to a BTA-stat test strip (Polymedco, Inc., Cortlandt Manor, N.Y.). The BTA-stat test strip contains a mobilizable anti-BTA antibody conjugated to a detectable moiety and the mixture migrates to a region containing an antibody that binds to BTA, which region, when sufficient BTA is present, form a visible line that indicates that the sample is positive for BTA.

[1190] Of samples from subjects having bladder cancer, 65% were positive for BTA in this test (17 out of 26 samples). Of samples from subjects not having bladder cancer, 21% were positive for BTA in this test (9 out of 42 samples) and 79% were negative for BTA in this test (33 out of 42 samples). Thus, presence of BTA had a 65% positive predictive value for the presence of bladder cancer and absence of BTA had a negative predictive value of 79% for the absence of bladder cancer.

[1191] E. OnfFN and BTA-Stat in Combination

[1192] Samples for which BTA analysis and onfFN measurements were performed were compared relative to the cancerous state of the subject; 48 urine samples were tested: 23 urine samples from subjects with bladder cancer and 25 urine samples from control (positive and negative) subjects.

[1193] All 15 subjects that provided samples positive for BTA and onfFN had bladder cancer. For subjects that provided samples negative for BTA and onfFN, 91% did not have bladder cancer (20 of 22). For subjects that provided samples negative for BTA and positive for onfFN, 71% had bladder cancer (5 of 7). For subjects that provided samples positive for BTA and negative for onfFN, 25% had bladder cancer (1 of 4).

[1194] F. Immunoprecipitation and Western Blot

[1195] For this analysis, urine samples were tested. Immunoprecipitation was performed by incubating the mouse monoclonal anti-oncofetal fibronectin A120 antibody (0.5-1.0 ng/well) or antibody A137 (2.5-5.0 ng/well) diluted 1:10 with APB buffer. 30 μ l A137 coated beads was added to 800 μ l of sample. Acid from the precipitation samples was neutralized by washing with PBS. An internal standard was set: (25 μ g/ml in APB=2 ng/well): <50% of 25 ng/ml in APB as negative (-); 50-60% of 25 ng/ml in APB as positive (+); 60-70% of 25 ng/ml in APB as positive (++); 70-80% of 25 ng/ml in APB as positive (+++); and 80-100% of 25 ng/ml in APB as positive (++++). Western blots were performed as described above.

[1196] G. Combined Results

[1197] Abbreviations are as follows: BC=bladder cancer, PC=prostate cancer; BPH=benign prostate hyperplasia, KC=kidney cancer, hydronephrosis (HN), vasectomy (VS) and incontinence (IN)=symptomatic urinary condition controls, stages and grades are as described above, pos=positive, and neg=negative.

TABLE 2

Urinary oncofetal fibronectin levels as an indication of bladder cancer								
Subject	Condition	Grade	Stage	Dot blot	Lateral flow	BTA Stat	Vertical Flow	IP/WB
1	BC	3	T3A	3	4	Pos		++++
2	BC	2	T1A			Pos		
3	BC	3	T1B	3		Pos		++++
4	BC	1	T1A			Neg		
5	BC	3	T1B	1	0	Pos	1	
6	BC	3	T3AN1	0.5	0	Pos	1	-
7	BC	3	T2B	0	1	Neg	1	
8	BC	3	T2B	2	0	Neg	1	++++
9	BC	3	T3A	1	2	Pos	0	++++
10	BC	3	T2			Pos		
11	BC	3	T1B	1	3	Pos	0	++++
12	BC	3	T1B			Neg		
13	BC	2	T1B	1	0.5	Pos	1	-
14	BC	3	T2B	2	1	Pos	1	++++
15	BC	1	T1A	0	0	Neg	0	
16	BC	1	T1A	1	1	Pos	0	++++
17	BC	1	T1A	0	0.5	Pos	0	-
18	BC	3		3	2	Pos		
19	BC	3	T4	3	3	Pos	1	++++
20	BC	3				Pos		
21	BC		None	0.5	0	Neg	1	-
22	BC	2	T1	3	1	Pos	1	++++
23	BC	1	T1A		0	Pos		+++
24	BC	2	T1	3	3	Pos	0	++++
25	BC	1	T1	0	0	Neg	1	-
26	BC		None	3	3			++++
27	BC	2	T1	0.5	0	Neg	0	-
28	BC	3	T1	1	1	Neg	0	-
29	BC	1	T1					++
30	BC		None	3	3			++++
31	BC		None	0	0.5			-
32	BC	1	T1	0	0			1
33	BC	3	T1	3	1			++++
34	BC	3	T3	0.5	0			++++
35	BC	3	T2	3	2			++++
36	BC	3	T2	3	2			++++
37	BC	1	T1					-
38	BC	3	T4		0			
39	BC	3	T2		0			-
40	BC	3	T3		1			
41	BC	2	T1		1			
42	BC	2	T1					+
43	BC	3	T3		0			
44	BC	3	T1		1			++++
45	PC					Neg		
46	PC				0	Neg		++++
47	PC					Pos		++++
48	PC					Neg		-
49	PC			0	0	Neg		-
50	BPH			0	0	Neg		
51	BPH				0	Neg		+++
52	BPH				0	Neg		-
53	BPH				0	Neg		-
54	BPH				0	Neg		-
55	BPH				0	Pos		++++
56	BPH			0	0	Neg		-
57	BPH					Pos		++++
58	BPH					Neg		-
59	BPH			0		Neg		-
60	BPH					Neg		-
61	KC			0	0	Neg		-
62	KC			0	0	Neg	1	
63	HN					Neg		-
64	VS					Pos		-
65	IN			0	0	Pos	0	-
66	Control			3	0.5	Pos		
67	Control			0	0	Neg		
68	Control			0	0	Neg		-
69	Control			0	0	Neg		-
70	Control			0	0.5	Neg		
71	Control			0	0	Neg		-

TABLE 2-continued

Urinary oncofetal fibronectin levels as an indication of bladder cancer								
Subject	Condition	Grade	Stage	Dot blot	Lateral flow	BTA Stat	Vertical Flow	IP/WB
72	Control			3	3	Neg		-
73	Control					Neg		-
74	Control					Neg		-
75	Control					Neg		-
76	Control			0	0	Neg	0	-
77	Control					Neg		-
78	Control					Neg		-
79	Control					Pos		-
80	Control			0	0	Neg		-
81	Control			0	0	Neg		-
82	Control			0		Neg	0	-
83	Control			0	0	Pos	1	-
84	Control			0	0	Neg	0	-
85	Control					Neg		-

[1198] Samples for which onfFN measurements were performed were compared relative to the cancerous state of the subject; 85 urine samples were tested: 44 urine samples from subjects with bladder cancer, 5 urine samples from subjects with prostate cancer, 11 urine samples from subjects with benign prostate hyperplasia, 2 urine samples from subjects with kidney cancer, 2 urine samples from symptomatic controls and 20 urine samples from control subjects.

[1199] Of samples from subjects having bladder cancer tested by lateral flow, 71% were positive for onfFN in this test (22 out of 31 samples). Of negative control samples from subjects not having bladder cancer tested by lateral flow, 94% were negative for onfFN in this test (15 out of 16 samples).

[1200] Of samples from subjects having bladder cancer as confirmed by immuno-precipitation and western blot, 65% were positive for onfFN in this test (20 out of 31 samples). Of negative control samples from subjects not having bladder cancer as determined by immunoprecipitation and western blot, 100% were negative for onfFN in this test (16 out of 16 samples). When a threshold of >50% of 25 ng/ml of onfFN was set for a positive outcome, 59% of stage T1 bladder cancer subjects were positive for onfFN in this test (10 out of 17 samples), 83% of stage T2 bladder cancer subjects were positive for onfFN in this test (5 out of 6 samples), 75% of stage T3 bladder cancer subjects were positive for onfFN in this test (3 out of 4 samples), 100% of stage T4 bladder cancer subjects were positive for onfFN in this test (1 out of 1 samples), 30% of benign prostate hyperplasia subjects were positive for onfFN in this test (3 out of 10 samples), 50% of prostate subjects were positive for onfFN in this test (2 out of 4 samples), 100% of subjects having symptomatic conditions (i.e., negative controls) were negative for onfFN in this test (10 out of 10 samples) and 100% of negative control subjects were negative for onfFN in this test (3 out of 3 samples).

[1201] Using immunoprecipitation and western blot analysis of the urine samples, it was possible to differentiate between invasive and non-invasive bladder cancer. Samples were tested for onfFN as described above, and a threshold of >50% of 25 ng/ml of onfFN was set for a positive outcome. Of the patients who had non-invasive bladder cancer, 54% tested at or above threshold, 80% of the patients identified

as having invasive bladder cancer tested positive. In contrast, 100% of negative controls that had a level of onfFN less than 25 ng/ml (16 out of 16 samples).

Example 14

Treatment of Urine Samples to Detect Oncofetal Fibronectin Indicating the Presence of Bladder Cancer

[1202] Urine samples were diluted 4-fold with antiprotease buffer (APB) containing 0.05 M Tris buffer, pH 7.4, 0.02% NaN₃, 1% BSA, 5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin, 150 mM NaCl and 0.1% Triton X-100.

[1203] Lateral flow was performed by adding 100 μ l of urine sample to 300 μ l of APB (0.05 M Tris buffer, pH 7.4, 150 mM NaCl, 0.02% NaN₃, 1% BSA, 5 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 500 Kallikrein Units/ml of Aprotinin and 0.1% Triton X-100), and then by applying 200 μ l of this diluted urine sample mixture to a test strip containing an absorbent pad which draws the sample to the test strip. The sample first flowed through a mobilization region containing a conjugate of FDC-6 conjugated to blue microspheres (Adeza Biomedical, Inc.) diffusibly bound to the test strip. The conjugate was mobilized by the flow of the sample. Next, the sample and conjugate flowed through a non-specific binding trap region containing methylated BSA attached to the test strip. The sample and conjugate then flowed through a region containing goat polyclonal anti-fibronectin antibodies, such as those described in Example 4 (also referred to herein as A120 antibodies) non-diffusibly bound to the test strip. Finally, the sample and conjugate flowed through a control region containing goat polyclonal anti-mouse IgG antibody (which selectively binds the FDC-6 conjugate) non-diffusibly bound to the test strip. The test strip was then visually inspected for the presence or absence of blue color in the detection and control regions. Variations of the lateral flow method were performed using BSA, W632 (anti-MHC Class I) and mouse IgG in the non-specific binding trap region upstream of the mobilization region.

Example 15

Immunoassay Test Strip

[1204] The test strip **100** of FIGS. 2 and 3 includes a membrane system including three components: a porous or bibulous member **102**; a conjugate pad **108**; and an absorbent pad **110**. The membrane system can be mounted on a substrate or backing **112**, with the conjugate pad **108** and the absorbent pad **110** slightly overlapping the porous or bibulous member **102**, which is interposed thereinbetween. As can be seen, the conjugate pad **108** overlaps the porous or bibulous member **102** so that a fluid sample placed onto the conjugate pad **108** is communicated from the conjugate pad **108** to the porous or bibulous member **102**. Similarly, the absorbent pad **110** overlaps with the porous or bibulous member **102** so that fluid samples introduced into the porous or bibulous member **102** from the conjugate pad **108** can then be transmitted to the absorbent pad **110**. Thus, the conjugate pad **108**, the absorbent pad **110** and the porous or bibulous member **102** are all in fluid communication with one another, making any fluid sample placed on the conjugate pad **108** able to propagate through the conjugate pad **108** to the porous or bibulous member **110** and then to the absorbent pad **110**.

[1205] The membrane system includes a conjugate pad **108**, which serves as a sample application component and which includes an antibody to the analyte, which is conjugated to a detectable label. The conjugate pad is in fluid communication with the porous or bibulous member **102**. The labeled antibody conjugate is diffusively bound to the conjugate pad and becomes mobile upon application of the liquid sample and moves along the test strip. The conjugate pad is made of a porous material, such as glass fiber. The conjugate pad also can act as a prefilter for the sample. The conjugate pad also can contain a non-specific binder such as a non-specific binding surface or a non-specific binding compound immobilized thereto.

[1206] The porous or bibulous member **102** is capable of transporting a liquid sample along the test strip and serves as the solid support upon which the immunoreactions occur. Antibodies which react with the target analyte and/or label are immobilized on the solid support. Possible solid supports include paper and cellulose derivatives, such as cellulose esters and ethers, natural and synthetic polymeric materials, such as vinyl polymers and partially hydrolyzed derivatives, polycondensates, copolymers and inorganic materials. An exemplary solid support is a nitrocellulose membrane.

[1207] The porous or bibulous member **102** contains at least two distinct zones, an analyte binding zone **104** and a control zone **106**, at which different antibodies are immobilized. The analyte zone contains an immobilized analyte binding partner such as an antibody that binds the analyte of interest, whereas the control zone contains an immobilized antibody or other component, such as an antigen, that binds labeled antibody conjugate which has not bound to analyte.

[1208] In addition, the porous member can contain a non-specific binder. A non-specific binder can be located along any portion of the conjugate pad **108** or any portion of the porous or bibulous member **102** located between the analyte binding zone **104** and the conjugate pad **108**.

[1209] The membrane system also can include an absorbent pad **112**, which also is in fluid communication with the

porous or bibulous member and which serves to draw liquid continuously through the device. The absorbent strip can be made of a material such as cellulose paper or other material known to those of skill in the art.

[1210] Referring to FIG. 4, which depicts an exemplary immunoassay device, including a test strip and housing assembly **300**, the housing **302** generally surrounds the test strip **100** (FIGS. 2 and 3) and includes an opening through which test sample is applied **304**, as well as an aperture above the detection and control zones **306** that permits measurement of the amount of label by the reader, which is correlated with the amount of analyte in the test sample. The housing **302** includes at its upper surface **308** a fattened end **310**, used for gripping the housing **302** and an application window **304** (or sample window) through which a sample is applied to a conjugate pad **108** of an immunoassay test strip within the housing **302**. The housing **302** also includes a test window **314** through which the test result of the immunoassay is viewed. In accordance with the embodiments shown, no window material is mounted within the test window **314** (or the sample window **312**). Thus, an optical path from outside the housing **302** through the test window **314** to the immunoassay test strip is unobscured by any material. Other alternative embodiments can include an optically transparent material (transparent at wavelengths emitted by light emitted from devices described herein).

Example 16

Oncofetal Fibronectin as an Indicator of Induction Outcome

A. Experimental Design

[1211] A.1 Objectives

[1212] A multicenter clinical trial was conducted to prospectively evaluate the utility of the onfFN test and outcomes associated with induction of labor in pregnant women with a gestational age of ≤ 36 0/7 and < 42 0/7 weeks. For the purposes of this investigation, favorable outcomes of induction of labor include a reduced rate of cesarean section, an increased rate of vaginal delivery within 24 and 48 hours, a shortened time to delivery following administration of cervical ripening agents, a reduction in the number of cervical ripening agent administrations and a reduction in time interval from oxytocin initiation until delivery.

[1213] This study compared:

[1214] 1. The rate of vaginal delivery within 24 hours and 48 hours of start of induction (including cervical ripening agent administration) in women testing onfFN positive versus onfFN negative.

[1215] 2. The time interval from first dose of cervical ripening agent to delivery in women testing onfFN positive versus onfFN negative.

[1216] 3. The number of cervical ripening agent administrations in women testing onfFN positive versus onfFN negative.

[1217] 4. The time interval from oxytocin initiation to delivery in women testing onfFN positive versus onfFN negative.

[1218] 5. The rate of cesarean section in nulliparous women with unfavorable cervixes and intact amniotic membranes testing onfFN positive versus onfFN negative.

[1219] A.2 Study Design

[1220] The study was conducted prospectively and clinicians managing patients were blinded to results of the onfFN test. Patient management was not based on the clinician's knowledge of the onfFN test result. Enrolled women were followed from the time the informed consent was signed through delivery. The sample size used was estimated to provide 80-90% power to test each of the above-listed five clinical outcomes. Each outcome was tested at the 2-sided 5% significance level. Nine hundred and one women were considered enrolled. Of these 901 enrolled women, onfFN test results were not available for 26 women due to incorrect specimen handling. Therefore, 875 women were enrolled and deemed evaluable for the safety and effectiveness data analysis.

[1221] A.3 Subjects

[1222] Women scheduled for an induction of labor were prescreened for inclusion in the study. Initial chart review determined if the woman was at least 18 years of age, nulliparous and with a singleton pregnancy in cephalic presentation with a gestational age between 36 0/7 and 42 0/7 weeks. An onfFN test specimen was obtained, as well as confirmation that amniotic fluid leakage had not occurred. Specimens for onfFN testing were collected from the posterior fornix of the vagina using the Adeza Biomedical Specimen Collection Kit (Sunnyvale, Calif.) before the clinician performed any other part of the pelvic exam. Testing was performed on the Rapid onfFN for the TLiIQ™ System (Adeza Biomedical, Sunnyvale, Calif.).

[1223] A.4 Data Collection and Analysis

[1224] Statistical analyses were performed using ANOVA for comparison of three or more groups. The 2-sample t-test was used to compare the means of continuous variables. The Fisher's Exact test, Chi-square Test or Wilcoxon 2-sample rank test was used to compare groups for discrete variables, as appropriate. The univariate and multivariate association of dichotomous or polychotomous variables to clinical outcome variables of interest was evaluated using logistic or general linear regression modeling methods, as appropriate. All tests were two-tailed and a p-value of less than 0.05 was required for statistical significance. Statistical evaluations were performed using the Statistical Analysis System (SAS Institute, Cary, N.C.).

B. Results

[1225] Of the 875 study women, 371 (42.4%) tested onfFN positive and 504 (57.6%) tested onfFN negative. The outcomes of induction for this population is presented below.

[1226] B.1 The Rate of Cesarean Section in Women Testing onfFN Positive Versus onfFN Negative

[1227] There was a difference in cesarean section rate in women testing onfFN positive and onfFN negative. In women testing onfFN negative, 187 delivered by cesarean section and 317 did not, resulting in an observed rate of cesarean section of 0.371. In women testing onfFN positive, 107 delivered by cesarean section and 264 did not, resulting in an observed rate of cesarean section of 0.288. The rate difference of 0.083 (standard error=0.032) was statistically significant with a p-value of 0.011 (Fisher's Exact test). Thus the cesarean section rate of women testing onfFN

negative was greater than the cesarean section rate for women testing onfFN positive.

[1228] The ratio of the rates is another way to quantify the difference in cesarean-section rates between women testing onfFN negative and women testing onfFN positive. The rate ratio (0.371/0.288) of 1.29 indicates that women testing onfFN negative prior to induction of labor were 29% [95% CI 6%, 57%] more likely to be delivered by cesarean section than women testing onfFN positive.

[1229] B.2 The Rate of Vaginal Delivery within 24 and 48 Hours of Start of Induction in Women Testing onfFN Positive Versus onfFN Negative

[1230] There was a difference in rate of vaginal delivery within 24 hours of initiation of cervical ripening agents or oxytocin among women testing onfFN positive and onfFN negative. In women testing onfFN positive, 229 delivered vaginally within 24 hours of initiating cervical ripening or oxytocin and 138 did not, resulting in an observed rate of vaginal delivery within 24 hours from initiation of cervical ripening or oxytocin in women testing onfFN positive of 0.624. In women testing onfFN negative, 225 delivered vaginally within 24 hours of initiating cervical ripening or oxytocin and 274 did not, resulting in an observed rate of vaginal delivery within 24 hours from initiation of cervical ripening or oxytocin in women testing onfFN negative of 0.451. Nine women were omitted either because they did not receive cervical ripening or oxytocin, or data were missing for them. The rate difference of 0.173 (standard error=0.034) was statistically significant with a p-value <0.0001 (Fisher's Exact test). Thus, the rate of vaginal delivery within 24 hours from initiation of cervical ripening or oxytocin in women testing onfFN positive was greater than the rate of vaginal delivery within 24 hours from initiation of cervical ripening or oxytocin in women testing onfFN negative.

[1231] The ratio of the rates of vaginal delivery within 24 hours from initiation of cervical ripening or oxytocin in women testing onfFN positive versus women testing onfFN negative (0.624/0.451) was 1.38. Thus, when results are expressed as a ratio of rates, women testing onfFN positive prior to induction of labor were 38% [95% CI 22%, 57%] more likely to have a vaginal delivery within 24 hours from initiation of cervical ripening or oxytocin than women testing onfFN negative.

[1232] In women testing onfFN positive, 253 delivered vaginally within 48 hours of initiating cervical ripening or oxytocin and 114 did not, resulting in an observed rate of vaginal delivery within 48 hours from initiation of cervical ripening or oxytocin in women testing onfFN positive of 0.689. In women testing onfFN negative, 299 delivered vaginally within 48 hours of initiating cervical ripening or oxytocin and 200 did not, resulting in an observed rate of vaginal delivery within 48 hours from initiation of cervical ripening or oxytocin in women testing onfFN negative of 0.599. Nine women were omitted either because they did not receive cervical ripening or oxytocin, or data were missing for them. The rate difference of 0.090 (standard error=0.033) was statistically significant with a p-value=0.007 (Fisher's Exact test). Thus, the rate of vaginal delivery within 48 hours from initiation of cervical ripening or oxytocin in women testing onfFN positive was greater than the rate of vaginal delivery within 48 hours from initiation of cervical ripening or oxytocin in women testing onfFN negative.

[1233] The ratio of the rates of vaginal delivery within 48 hours from initiation of cervical ripening or oxytocin in women testing onfFN positive versus women testing onfFN negative (0.689/0.599) was 1.15. Thus, when results are expressed as a ratio of rates, women testing onfFN positive prior to induction of labor were 15% [95% CI 4%, 27%] more likely to have a vaginal delivery within 48 hours from initiation of cervical ripening or oxytocin than women testing onfFN negative.

[1234] B.3 The Time Interval from First Dose of Cervical Ripening Agent or Oxytocin to Delivery in Women Testing onfFN Positive Versus onfFN Negative

[1235] The mean observed time from the first administration of a cervical ripening agent or oxytocin to delivery was 9.6 hours longer in women testing onfFN negative (28.7 hours) than for women testing onfFN positive (19.1 hours). The standard error of this difference was 1.92 and the 2-sample t-test p-value was <0.0001. Thus the mean number of hours from initiation of cervical ripening or oxytocin to delivery was fewer among women testing onfFN positive than among women testing onfFN negative.

[1236] B.4 The Number of Cervical Ripening Agent Administrations in Women Testing onfFN Positive Versus onfFN Negative

[1237] The mean observed number of cervical ripening agent administrations was 0.60 greater in women testing onfFN negative (1.43) than for women testing onfFN positive (0.83). The standard error of this difference was 0.08 and the 2-sample t-test p-value was <0.0001. Thus, the mean number of cervical ripening administrations was fewer among women testing onfFN positive than among women testing onfFN negative.

[1238] B.5 The Time Interval from Oxytocin Initiation to Delivery in Women Testing onfFN Positive Versus onfFN Negative

[1239] The mean observed time from oxytocin administration to delivery was 4.4 hours longer in women testing onfFN negative (16.8 hours) than for women testing onfFN positive (12.4 hours). The standard error of this difference was 1.07 and the 2-sample t-test p-value was <0.0001. Thus, the mean number of hours from oxytocin initiation to delivery was fewer among women testing onfFN positive than among women testing onfFN negative.

C. Summary of Association of onfFN Test Results with Outcome

[1240] The association of the onfFN test with induction of labor outcomes for differences in rates (cesarean section and vaginal delivery within 24 and 48 hours) and for differences in means (time interval from first dose of cervical ripening agent, number of cervical ripening agent administrations and time interval from oxytocin initiation to delivery) is shown in Tables 3 and 4, respectively. Compared to women testing onfFN negative, those testing onfFN positive are less likely to deliver by cesarean section and are more unlikely to have a vaginal delivery within 24 hours, a shortened time to delivery, a reduction in the number of cervical ripening agent administrations and a shortened time from oxytocin initiation until delivery.

TABLE 3

	Rate Difference	Standard Error	p-value
Cesarean Section	0.083	0.032	0.011
Vaginal Delivery <24 Hours	0.173	0.035	<0.0001
Vaginal Delivery >48 Hours	0.090	0.033	0.007

[1241]

TABLE 4

	Mean Difference	Standard Error	p-value
Interval From Cervical Ripening agent or Oxytocin to Delivery (hours)	9.6	1.92	<0.0001
Number of Cervical Ripening Agent Administrations	0.603	0.085	<0.0001
Interval from Oxytocin Initiation to Delivery (hours)	4.4	1.07	<0.0001

Example 17

Differential Outcomes at Different Levels of Oncofetal Fibronectin

[1242] Cervicovaginal swab samples collected from women in their 24th week of pregnancy were measured for concentrations of oncofetal fibronectin. The pregnant women were then followed and their date of delivery recorded. The pregnant women were pooled into three populations: (a) women with oncofetal fibronectin measurements below 60 ng/ml, (b) women with oncofetal fibronectin measurements between 60 ng/ml and 150 ng/ml and (c) women with oncofetal fibronectin measurements equal to or above 150 ng/ml. The percent of women from each population that remained undelivered was then plotted against the date of delivery (after the 24th week).

[1243] Each of the three populations showed different delivery outcomes. The lowest oncofetal fibronectin group had the highest percent undelivered throughout the remainder of the pregnancy term and the highest oncofetal fibronectin group had the lowest percent undelivered throughout the remainder of the pregnancy term. The intermediate group remained similar in outcome to the lowest oncofetal fibronectin group until about week 30 and then showed a delivery rate of about 5% per week for the following 6 weeks (30th week through 36th week). At the 36th week, about 65% of this population remained undelivered. The highest oncofetal fibronectin group showed a delivery rate of about 5% per week for the first four weeks (24th week through 28th week) and about 5% per two weeks for the following eight weeks (28th week through 36th week). Less than 60% of this population remained undelivered at the 36th week.

Example 18

Oncofetal Fibronectin Measurements in Swab Samples Collected from the Lower Third of the Vagina

[1244] Cervicovaginal swab samples were collected from 259 pregnant subjects in two formats. The first cervicovagi-

nal swab sample format was a swab of the posterior fornix. The second cervicovaginal swab sample format was a swab of the lower third of the vagina.

[1245] All samples were applied to a test strip using the lateral flow method described in Example 12, and the concentration of oncofetal fibronectin in the sample was determined according to reflectance intensity measured with a TLiQ™ reflectance reader. Cervicovaginal swab samples of the posterior fornix were categorized as oncofetal fibronectin positive when the reflectance signal indicated an oncofetal fibronectin concentration of 50 ng/ml or more (for the lot of test strips used, the corresponding reflectance signal indicating 50 ng/ml was 0.315). Cervicovaginal swab samples of the lower third of the vagina were categorized as oncofetal fibronectin positive when the reflectance signal was 0.1 or greater. Since the reflectance signal for this test strip is a linear function of oncofetal fibronectin at these conditions, cervicovaginal swab samples of the lower third of the vagina were categorized as oncofetal fibronectin positive when the amount of oncofetal fibronectin was measured to be 16 ng/ml or more.

[1246] Of the two sets of 259 samples, 239 sets of samples were in agreement. Specifically, of 239 sets of samples, for 205 sets both were negative and for 34 sets both were positive. Of the 20 sets of samples that differed, only 3 samples were negative in the lower third of the vagina while being positive in the posterior fornix. The remaining 17 samples were posterior fornix negative and lower third positive. Thus, a positive measurement of the posterior fornix is accompanied by a positive measurement of the lower third of the vagina in 91.9% of the measurements and a negative measurement of the posterior fornix is accompanied by a negative measurement of the lower third of the vagina in 92.3% of the measurements. A positive measurement of the lower third of the vagina is accompanied by a positive measurement for the posterior fornix in 66.7% of the measurements and a negative measurement of the lower third of the vagina is accompanied by a negative measurement for the posterior fornix in 98.6% of the measurements.

[1247] Since modifications will be apparent to those of skill in this art, it is intended that this invention be limited only by the scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 39

<210> SEQ ID NO 1

<211> LENGTH: 7388

<212> TYPE: DNA

<213> ORGANISM: Homo Sapiens

<220> FEATURE:

<223> OTHER INFORMATION: Fibronectin

<400> SEQUENCE: 1

```

gccccgcgcg gctgtgctgc acagggggag gagagggaac cccaggcgcg agcgggaaga      60
ggggacctgc agccacaact tctctgttcc tctgcatccc ttctgtccct ccaccctcc      120
ccttccccac cctctggccc ccagcctcct ccctatggcc actgtgtcac agacagtgg      180
gtggtctact ctgtggggat gcagtggctg aagacacaag gaaataagca aatgctttgc      240
acgtgcctgg gcaacggagt cagctgcaa gagacagctg taaccagac ttacggtggc      300
aactcaaatg gagagccatg tgtcttacca ttcacctaca atggcaggac gttctactcc      360
tgcaccacag aagggcgaca ggacggacat ctttggtgca gcacaacttc gaattatgag      420
caggaccaga aataactcttt ctgcacagac cacactgttt tggttcagac tcgaggagga      480
aattccaatg gtgccttctg ccacttcccc ttcctataca acaaccacaa ttactctgat      540
tgcacttctg agggcagaag agacaacatg aagtgggtg ggaccacaca gaactatgat      600
gccgaccaga agtttgggtt ctgccccatg gctgcccacg aggaaatctg cacaaccaat      660
gaaggggtca tgtaccgcat tggagatcag tgggataaag agcatgacat gggtcacatg      720
atgaggtgca cgtgtgttgg gaatggctgt ggggaatgga catgcattgc ctactcgcag      780
cttcgagatc agtgcattgt tgatgacatc acttacaatg tgaacgacac attccacaag      840
cgtcatgaag aggggcacat gctgaactgt acatgcttcg gtcagggtcg gggcaggtgg      900
aagtgtgatc ccgtcgacca atgccaggat tcagagactg ggacgtttta tcaaattgga      960
gattcatggg agaagtatgt gcatggtgtc agataccagt gctactgcta tggccgtggc     1020

```

-continued

attggggagt ggcattgcc	acctttacag acctatccaa gctcaagtgg tctgtcga	1080
gtatttatca ctgagactcc	gagtcagccc aactcccacc ccatccagtg gaatgcacca	1140
cagccatctc acatttccaa	gtacattctc aggtggagac ctaaaaattc tgtaggccgt	1200
tggaaggaag ctaccatacc	agggcactta aactcctaca ccatcaaagg cctgaagcct	1260
ggtgtggtat acgagggcca	gctcatcagc atccagcagt acggccacca agaagtgact	1320
cgctttgact tcaccaccac	cagcaccagc acacctgtga ccagcaacac cgtgacagga	1380
gagacgactc cttttctcc	tcttgtggcc acttctgaat ctgtgaccga aatcacagcc	1440
agtagctttg tggctctctg	ggtctcagct tccgacaccg tgtcgggatt cgggtggaa	1500
tatgagctga gtgagaggg	agatgagcca cagtacctgg atcttccaag cacagccact	1560
tctgtgaaca tccctgacct	gcttctctggc cgaaaataca ttgtaatgt ctatcagata	1620
tctgaggatg gggagcagag	tttgatcctg tctacttcac aaacaacagc gcctgatgcc	1680
ccacctgacc cgactgtgga	ccaagttgat gacacctcaa ttgttgttcg ctggagcaga	1740
ccccaggctc ccatcacagg	gtacagaata gtctattcgc catcagtaga aggtagcagc	1800
acagaactca accttctcga	aactgcaaac tccgtcacc tcagtgaact gcaacctggt	1860
gttcagtata acatcactat	ctatgctgtg gaagaaaatc aagaaagtac acctgttgc	1920
attcaacaag aaacctctg	caccccacgc tcagatacag tgcctctcc cagggacctg	1980
cagtttgtgg aagtgcagag	cgtgaagtc accatcatgt ggacaccgcc tgagagtga	2040
gtgaccggct accgtgtgga	tgtgatcccc gtcaacctgc ctggcgagca cgggcagagg	2100
ctgccatca gcaggaacac	ctttgcagaa gtcaccgggc tgtcccctgg ggtcacctat	2160
tacttcaaag tctttgcagt	gagccatggg agggagagca agcctctgac tgcacaacag	2220
acaacaaaac tggatgctc	cactaacctc cagtttgc atgaaactga ttctactgtc	2280
ctggtgagat ggactccacc	tcgggcccag ataacaggat accgactgac cgtgggcctt	2340
acccgaagag gacagcccag	gcagtacaat gtgggtccct ctgtctccaa gtaccactg	2400
aggaatctgc agcctgcac	tgagtacacc gtatccctcg tggccataaa gggcaaccaa	2460
gagagcccca aagccactg	agtctttacc aactgcagc ctgggagctc tattccacct	2520
tacaacaccg aggtgactga	gaccaccatt gtgatcacat ggacgcctgc tccaagaatt	2580
ggttttaagc tgggtgtacg	accaagccag ggaggagagg caccacgaga agtgacttca	2640
gactcaggaa gcatcgttgt	gtccggcttg actccaggag tagaatacgt ctacaccatc	2700
caagtcctga gagatggaca	ggaaagagat gcgccaattg taaacaaagt ggtgacacca	2760
ttgtctccac caacaaact	gcatctggag gcaaacctg aactggagt gtcacagtc	2820
tcctgggaga ggagcaccac	cccagacatt actggttata gaattaccac aaccctaca	2880
aacggccagc agggaaatc	tttgaagaa gtgtccatg ctgatcagag ctctgcact	2940
tttgataacc tgagtcccg	cctggagtac aatgtcagtg tttacactgt caaggatgac	3000
aaggaaagtg tcctatctc	tgataaccac atcccagagg tgcccact cactgacct	3060
agctttgttg atataaccga	ttcaagcatc ggctgagggt ggacccgct aaactcttc	3120
accattattg ggtaccgcat	cacagtagtt gcggcaggag aaggtatccc tatttttgaa	3180
gattttgtgg actcctcagt	aggatactac acagtcacag ggtggagcc gggcattgac	3240
tatgatatca gcgttatcac	tctcattaat ggcgcgaga gtgccctac tacactgaca	3300

-continued

caacaaacgg ctgttctctc tcccactgac ctgctgattca ccaacattgg tccagacacc	3360
atgctgtca cctgggctcc accaccatcc attgatttaa ccaacttctt ggtgcgttac	3420
tcacctgtga aaaatgagga agatgttgca gagttgtcaa tttctccttc agacaatgca	3480
gtggtcttaa caaatctctc gcctggtaga gaatatgtag tgagtgtctc cagtgtctac	3540
gaacaacatg agagcacacc tcttagagga agacagaaaa caggctctga tcccccaact	3600
ggcattgact tttctgatat tactgccaac tcttttactg tgcactggat tgctcctcga	3660
gccaccatca ctggctacag gatccgccat catcccgagc acttcagtgg gagacctcga	3720
gaagatcggg tgccccactc tcggaattcc atcacctca ccaacctcacc tccaggcaca	3780
gagtatgtgg tcagcatcgt tgctcttaac ggacagagagg aaagtccctt attgattggc	3840
caacaatcaa cagtttctga tgttccgagg gacctggaag ttgtgtctgc gacccccacc	3900
agcctactga tcagctggga tgctcctgct gtcacagtga gatattacag gatcacttac	3960
ggagaaacag gaggaaatag ccctgtccag gagttcactg tgctggggag caagtctaca	4020
gctaccatca gcggccttaa acctggagtt gattatacca tcaactgtgca tgctgtcact	4080
ggcctgtggg acagccccgc aagcagcaag ccaatttcca ttaattaccg aacagaaatt	4140
gacaaacatc ccagatgca agtgaccgat gttcaggaca acagcattag tgtcaagtgg	4200
ctgcttcaa gttccccctg tactggttac agagtaacca ccaactccaa aaatggacca	4260
ggaccaacaa aactaaaac tgcaggtcca gatcaaacag aaatgactat tgaaggcttg	4320
cagcccacag tggagtacgt ggttagtgc tatgctcaga atccaagcgg agagagtacg	4380
cctctggttc agactgcagt aaccaacatt gatcgcccta aaggactggc attcactgat	4440
gtggatgtcg attccatcaa aattgcttgg gaaagccac aggggcaagt tccaggtac	4500
agggtagcct actcgagccc tgaggatgga atccatgagc tattccctgc acctgatggt	4560
gaagaagaca ctgcagagct gcaaggcctc agaccgggtt ctgagtacac agtcagtgtg	4620
gttgcttgc acgatgatat ggagagccag cccctgattg gaaccagtc cacagctatt	4680
cctgcaccaa ctgacctgaa gttcactcag gtcacacca caagcctgag cgcccagtgg	4740
acaccacca atgttcagct cactggatat cgagtgcggg tgacccccaa ggagaagacc	4800
ggaccaatga aagaaatcaa ccttgcctct gacagctcat ccgtggtgt atcaggactt	4860
atggtggcca ccaaatatga agtgagtgc tatgctctta aggacacttt gacaagcaga	4920
ccagctcagg gtgtgtcacc cactctggag aatgtcggcc caccaagaag ggctcgtgtg	4980
acagatgcta ctgagaccac catcaccatt agctggagaa ccaagactga gacgatcact	5040
ggcttcaag ttgatgccgt tccagccaat ggccagactc caatccagag aacctcaag	5100
ccagatgtca gaagctacac catcacaggt ttacaaccag gcaactgact caagatctac	5160
ctgtacacct tgaatgacaa tgctcggagc tcccctgtgg tcatcgacgc ctccactgcc	5220
attgatgac catccaacct gcgtttcctg gccaccacac ccaattcctt gctggtatca	5280
tggcagccgc cactgcccag gattaaccgc tacatcatca agtatgagaa gcctgggtct	5340
cctcccagag aagtgttccc tcggccccgc cctggtgtca cagaggctac tattactggc	5400
ctggaaccgg gaaccgaata tacaatttat gtcattgccc tgaagaataa tcagaagagc	5460
gagcccctga ttggaaggaa aaagacagac gagcttcccc aactggtaac ccttccacac	5520
cccaatcttc atggaccaga gatcttgat gttccttcca cagttcaaaa gaccctttc	5580

-continued

```

gtcaccacc ctgggtatga cactggaaat ggtattcagc ttcctggcac ttctggtcag 5640
caaccacagt ttgggcaaca aatgatcttt gaggaacatg gtttaggcg gaccacaccg 5700
ccccaacggy ccacccccat aaggcatagg ccaagacat acccgccgaa tgtaggtgag 5760
gaaatccaaa ttggtcacat tcccagggaa gatgtagact atcacctgta cccacacggt 5820
ccgggactca atccaaatgc ctctacacaa gaagctctct ctcagacaac catctcatgg 5880
gccccattcc aggacacttc tgagtacatc atttcatgtc atcctgttgg cactgatgaa 5940
gaacccttac agttcagggt tcctggaact tctaccagtg cactctgac aggctcacc 6000
agaggtgcca cctacaacat catagtggag gcaactgaa accagcagag gcataagggt 6060
cgggaagagg ttgttaccgt gggcaactct gtcaacgaag gcttgaacca acctacggat 6120
gactcgtgct ttgacccta cacagtttcc cattatgccg ttggagatga gtgggaacga 6180
atgtctgaat caggctttaa actgttgtgc cagtgccttag gcttggaaag tggtcatttc 6240
agatgtgatt catctagatg gtgccatgac aatggtgtga actacaagat tggagagaag 6300
tgggaccgtc agggagaaaa tggccagatg atgagctgca catgtcttgg gaacggaaaa 6360
ggagaattca agtgtgaccc tcatgaggca acgtgttatg atgatgggaa gacataccac 6420
gtaggagaac agtggcagaa ggaatatctc ggtgccattt gctcctgcac atgctttgga 6480
ggccagcggg gctggcgtg tgacaactgc cgcagacctg ggggtgaacc cactcccgaa 6540
ggcactactg gccagtccta caaccagtat tctcagagat accatcagag acaaacact 6600
aatgttaatt gcccaattga gtgcttcatg cctttagatg tacaggctga cagagaagat 6660
tcccagagat aatcatctt tccaatccag aggaacaagc atgtctctct gccaatgacc 6720
atctaaactg gagtgatgtt agcagaccca gcttagagtt cttcttctt tcttaagccc 6780
tttgctctg aggaagtctt ccagcttcag ctcaactcac agcttctcca agcatcacc 6840
tgagggtttc ctgaggggtt tctcataaat gagggctgca cattgcctgt tctgcttcga 6900
agtattcaat accgctcagt attttaaatg aagtgattct aagatttggg ttgggatcaa 6960
taggaaagca tatgcagcca accaagatgc aatgttttg aatgatatg accaaaattt 7020
taagtaggaa agtcaccaa acacttctgc tttcacttaa gtgtctggcc cgcaatactg 7080
taggaacaag catgatcttg ttactgtgat attttaata tccacagtac tcaacttttc 7140
caaatgatcc tagtaattgc ctagaatat ctttctctta cctgttattt atcaattttt 7200
cccagtattt ttatacggaa aaaattgtat tgaaaacact tagtatgcag ttgataagag 7260
gaatttggta taattatggt ggggtgattat tttttatact gtatgtgcca aagctttact 7320
actgtggaaa gacaactggt ttaataaaaag atttacattc cacaacttga aaaaaaaaaa 7380
aaaaaaaaa 7388

```

```

<210> SEQ ID NO 2
<211> LENGTH: 2223
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens

```

```

<400> SEQUENCE: 2

```

```

Ala Arg Ala Gly Cys Ala Ala Gln Gly Glu Glu Arg Glu Pro Gln Ala
 1           5           10           15
Arg Ala Gly Arg Gly Asp Leu Gln Pro Gln Leu Leu Trp Ser Ser Ala
          20           25           30

```

-continued

Ser Leu Leu Ser Leu His Pro Ser Pro Ser Pro Pro Ser Gly Pro Gln
 35 40 45

Pro Pro Pro Tyr Gly His Cys Val Thr Asp Ser Gly Val Val Tyr Ser
 50 55 60

Val Gly Met Gln Trp Leu Lys Thr Gln Gly Asn Lys Gln Met Leu Cys
 65 70 75 80

Thr Cys Leu Gly Asn Gly Val Ser Cys Gln Glu Thr Ala Val Thr Gln
 85 90 95

Thr Tyr Gly Gly Asn Ser Asn Gly Glu Pro Cys Val Leu Pro Phe Thr
 100 105 110

Tyr Asn Gly Arg Thr Phe Tyr Ser Cys Thr Thr Glu Gly Arg Gln Asp
 115 120 125

Gly His Leu Trp Cys Ser Thr Thr Ser Asn Tyr Glu Gln Asp Gln Lys
 130 135 140

Tyr Ser Phe Cys Thr Asp His Thr Val Leu Val Gln Thr Arg Gly Gly
 145 150 155 160

Asn Ser Asn Gly Ala Leu Cys His Phe Pro Phe Leu Tyr Asn Asn His
 165 170 175

Asn Tyr Thr Asp Cys Thr Ser Glu Gly Arg Arg Asp Asn Met Lys Trp
 180 185 190

Cys Gly Thr Thr Gln Asn Tyr Asp Ala Asp Gln Lys Phe Gly Phe Cys
 195 200 205

Pro Met Ala Ala His Glu Glu Ile Cys Thr Thr Asn Glu Gly Val Met
 210 215 220

Tyr Arg Ile Gly Asp Gln Trp Asp Lys Gln His Asp Met Gly His Met
 225 230 235 240

Met Arg Cys Thr Cys Val Gly Asn Gly Arg Gly Glu Trp Thr Cys Ile
 245 250 255

Ala Tyr Ser Gln Leu Arg Asp Gln Cys Ile Val Asp Asp Ile Thr Tyr
 260 265 270

Asn Val Asn Asp Thr Phe His Lys Arg His Glu Glu Gly His Met Leu
 275 280 285

Asn Cys Thr Cys Phe Gly Gln Gly Arg Gly Arg Trp Lys Cys Asp Pro
 290 295 300

Val Asp Gln Cys Gln Asp Ser Glu Thr Gly Thr Phe Tyr Gln Ile Gly
 305 310 315 320

Asp Ser Trp Glu Lys Tyr Val His Gly Val Arg Tyr Gln Cys Tyr Cys
 325 330 335

Tyr Gly Arg Gly Ile Gly Glu Trp His Cys Gln Pro Leu Gln Thr Tyr
 340 345 350

Pro Ser Ser Ser Gly Pro Val Glu Val Phe Ile Thr Glu Thr Pro Ser
 355 360 365

Gln Pro Asn Ser His Pro Ile Gln Trp Asn Ala Pro Gln Pro Ser His
 370 375 380

Ile Ser Lys Tyr Ile Leu Arg Trp Arg Pro Lys Asn Ser Val Gly Arg
 385 390 395 400

Trp Lys Glu Ala Thr Ile Pro Gly His Leu Asn Ser Tyr Thr Ile Lys
 405 410 415

Gly Leu Lys Pro Gly Val Val Tyr Glu Gly Gln Leu Ile Ser Ile Gln
 420 425 430

Gln Tyr Gly His Gln Glu Val Thr Arg Phe Asp Phe Thr Thr Thr Ser

-continued

435				440				445							
Thr	Ser	Thr	Pro	Val	Thr	Ser	Asn	Thr	Val	Thr	Gly	Glu	Thr	Thr	Pro
450						455					460				
Phe	Ser	Pro	Leu	Val	Ala	Thr	Ser	Glu	Ser	Val	Thr	Glu	Ile	Thr	Ala
465					470					475					480
Ser	Ser	Phe	Val	Val	Ser	Trp	Val	Ser	Ala	Ser	Asp	Thr	Val	Ser	Gly
				485					490					495	
Phe	Arg	Val	Glu	Tyr	Glu	Leu	Ser	Glu	Glu	Gly	Asp	Glu	Pro	Gln	Tyr
			500					505					510		
Leu	Asp	Leu	Pro	Ser	Thr	Ala	Thr	Ser	Val	Asn	Ile	Pro	Asp	Leu	Leu
		515					520					525			
Pro	Gly	Arg	Lys	Tyr	Ile	Val	Asn	Val	Tyr	Gln	Ile	Ser	Glu	Asp	Gly
	530					535					540				
Glu	Gln	Ser	Leu	Ile	Leu	Ser	Thr	Ser	Gln	Thr	Thr	Ala	Pro	Asp	Ala
545					550					555					560
Pro	Pro	Asp	Pro	Thr	Val	Asp	Gln	Val	Asp	Asp	Thr	Ser	Ile	Val	Val
				565					570					575	
Arg	Trp	Ser	Arg	Pro	Gln	Ala	Pro	Ile	Thr	Gly	Tyr	Arg	Ile	Val	Tyr
			580					585					590		
Ser	Pro	Ser	Val	Glu	Gly	Ser	Ser	Thr	Glu	Leu	Asn	Leu	Pro	Glu	Thr
		595					600					605			
Ala	Asn	Ser	Val	Thr	Leu	Ser	Asp	Leu	Gln	Pro	Gly	Val	Gln	Tyr	Asn
	610					615					620				
Ile	Thr	Ile	Tyr	Ala	Val	Glu	Glu	Asn	Gln	Glu	Ser	Thr	Pro	Val	Val
625					630					635					640
Ile	Gln	Gln	Glu	Thr	Thr	Gly	Thr	Pro	Arg	Ser	Asp	Thr	Val	Pro	Ser
				645					650					655	
Pro	Arg	Asp	Leu	Gln	Phe	Val	Glu	Val	Thr	Asp	Val	Lys	Val	Thr	Ile
			660					665					670		
Met	Trp	Thr	Pro	Pro	Glu	Ser	Ala	Val	Thr	Gly	Tyr	Arg	Val	Asp	Val
		675					680					685			
Ile	Pro	Val	Asn	Leu	Pro	Gly	Glu	His	Gly	Gln	Arg	Leu	Pro	Ile	Ser
	690					695					700				
Arg	Asn	Thr	Phe	Ala	Glu	Val	Thr	Gly	Leu	Ser	Pro	Gly	Val	Thr	Tyr
705					710					715					720
Tyr	Phe	Lys	Val	Phe	Ala	Val	Ser	His	Gly	Arg	Glu	Ser	Lys	Pro	Leu
				725					730					735	
Thr	Ala	Gln	Gln	Thr	Thr	Lys	Leu	Asp	Ala	Pro	Thr	Asn	Leu	Gln	Phe
			740					745					750		
Val	Asn	Glu	Thr	Asp	Ser	Thr	Val	Leu	Val	Arg	Trp	Thr	Pro	Pro	Arg
		755					760					765			
Ala	Gln	Ile	Thr	Gly	Tyr	Arg	Leu	Thr	Val	Gly	Leu	Thr	Arg	Arg	Gly
	770					775					780				
Gln	Pro	Arg	Gln	Tyr	Asn	Val	Gly	Pro	Ser	Val	Ser	Lys	Tyr	Pro	Leu
785					790					795					800
Arg	Asn	Leu	Gln	Pro	Ala	Ser	Glu	Tyr	Thr	Val	Ser	Leu	Val	Ala	Ile
				805					810					815	
Lys	Gly	Asn	Gln	Glu	Ser	Pro	Lys	Ala	Thr	Gly	Val	Phe	Thr	Thr	Leu
			820					825					830		
Gln	Pro	Gly	Ser	Ser	Ile	Pro	Pro	Tyr	Asn	Thr	Glu	Val	Thr	Glu	Thr
		835					840					845			

-continued

Thr Ile Val Ile Thr Trp Thr Pro Ala Pro Arg Ile Gly Phe Lys Leu
 850 855 860

Gly Val Arg Pro Ser Gln Gly Gly Glu Ala Pro Arg Glu Val Thr Ser
 865 870 875 880

Asp Ser Gly Ser Ile Val Val Ser Gly Leu Thr Pro Gly Val Glu Tyr
 885 890 895

Val Tyr Thr Ile Gln Val Leu Arg Asp Gly Gln Glu Arg Asp Ala Pro
 900 905 910

Ile Val Asn Lys Val Val Thr Pro Leu Ser Pro Pro Thr Asn Leu His
 915 920 925

Leu Glu Ala Asn Pro Asp Thr Gly Val Leu Thr Val Ser Trp Glu Arg
 930 935 940

Ser Thr Thr Pro Asp Ile Thr Gly Tyr Arg Ile Thr Thr Thr Pro Thr
 945 950 955 960

Asn Gly Gln Gln Gly Asn Ser Leu Glu Glu Val Val His Ala Asp Gln
 965 970 975

Ser Ser Cys Thr Phe Asp Asn Leu Ser Pro Gly Leu Glu Tyr Asn Val
 980 985 990

Ser Val Tyr Thr Val Lys Asp Asp Lys Glu Ser Val Pro Ile Ser Asp
 995 1000 1005

Thr Ile Ile Pro Glu Val Pro Gln Leu Thr Asp Leu Ser Phe Val Asp
 1010 1015 1020

Ile Thr Asp Ser Ser Ile Gly Leu Arg Trp Thr Pro Leu Asn Ser Ser
 1025 1030 1035 1040

Thr Ile Ile Gly Tyr Arg Ile Thr Val Val Ala Ala Gly Glu Gly Ile
 1045 1050 1055

Pro Ile Phe Glu Asp Phe Val Asp Ser Ser Val Gly Tyr Tyr Thr Val
 1060 1065 1070

Thr Gly Leu Glu Pro Gly Ile Asp Tyr Asp Ile Ser Val Ile Thr Leu
 1075 1080 1085

Ile Asn Gly Gly Glu Ser Ala Pro Thr Thr Leu Thr Gln Gln Thr Ala
 1090 1095 1100

Val Pro Pro Pro Thr Asp Leu Arg Phe Thr Asn Ile Gly Pro Asp Thr
 1105 1110 1115 1120

Met Arg Val Thr Trp Ala Pro Pro Pro Ser Ile Asp Leu Thr Asn Phe
 1125 1130 1135

Leu Val Arg Tyr Ser Pro Val Lys Asn Glu Glu Asp Val Ala Glu Leu
 1140 1145 1150

Ser Ile Ser Pro Ser Asp Asn Ala Val Leu Thr Asn Leu Leu Pro
 1155 1160 1165

Gly Thr Glu Tyr Val Val Ser Val Ser Ser Val Tyr Glu Gln His Glu
 1170 1175 1180

Ser Thr Pro Leu Arg Gly Arg Gln Lys Thr Gly Leu Asp Ser Pro Thr
 1185 1190 1195 1200

Gly Ile Asp Phe Ser Asp Ile Thr Ala Asn Ser Phe Thr Val His Trp
 1205 1210 1215

Ile Ala Pro Arg Ala Thr Ile Thr Gly Tyr Arg Ile Arg His His Pro
 1220 1225 1230

Glu His Phe Ser Gly Arg Pro Arg Glu Asp Arg Val Pro His Ser Arg
 1235 1240 1245

-continued

Asn	Ser	Ile	Thr	Leu	Thr	Asn	Leu	Thr	Pro	Gly	Thr	Glu	Tyr	Val	Val
1250						1255					1260				
Ser	Ile	Val	Ala	Leu	Asn	Gly	Arg	Glu	Glu	Ser	Pro	Leu	Leu	Ile	Gly
1265					1270					1275					1280
Gln	Gln	Ser	Thr	Val	Ser	Asp	Val	Pro	Arg	Asp	Leu	Glu	Val	Val	Ala
				1285					1290						1295
Ala	Thr	Pro	Thr	Ser	Leu	Leu	Ile	Ser	Trp	Asp	Ala	Pro	Ala	Val	Thr
			1300					1305						1310	
Val	Arg	Tyr	Tyr	Arg	Ile	Thr	Tyr	Gly	Glu	Thr	Gly	Gly	Asn	Ser	Pro
		1315					1320					1325			
Val	Gln	Glu	Phe	Thr	Val	Pro	Gly	Ser	Lys	Ser	Thr	Ala	Thr	Ile	Ser
	1330					1335					1340				
Gly	Leu	Lys	Pro	Gly	Val	Asp	Tyr	Thr	Ile	Thr	Val	His	Ala	Val	Thr
1345					1350					1355					1360
Gly	Arg	Gly	Asp	Ser	Pro	Ala	Ser	Ser	Lys	Pro	Ile	Ser	Ile	Asn	Tyr
				1365					1370					1375	
Arg	Thr	Glu	Ile	Asp	Lys	Pro	Ser	Gln	Met	Gln	Val	Thr	Asp	Val	Gln
			1380					1385						1390	
Asp	Asn	Ser	Ile	Ser	Val	Lys	Trp	Leu	Pro	Ser	Ser	Ser	Pro	Val	Thr
		1395					1400						1405		
Gly	Tyr	Arg	Val	Thr	Thr	Thr	Pro	Lys	Asn	Gly	Pro	Gly	Pro	Thr	Lys
	1410					1415						1420			
Thr	Lys	Thr	Ala	Gly	Pro	Asp	Gln	Thr	Glu	Met	Thr	Ile	Glu	Gly	Leu
1425					1430					1435					1440
Gln	Pro	Thr	Val	Glu	Tyr	Val	Val	Ser	Val	Tyr	Ala	Gln	Asn	Pro	Ser
				1445					1450					1455	
Gly	Glu	Ser	Gln	Pro	Leu	Val	Gln	Thr	Ala	Val	Thr	Asn	Ile	Asp	Arg
			1460					1465					1470		
Pro	Lys	Gly	Leu	Ala	Phe	Thr	Asp	Val	Asp	Val	Asp	Ser	Ile	Lys	Ile
		1475					1480					1485			
Ala	Trp	Glu	Ser	Pro	Gln	Gly	Gln	Val	Ser	Arg	Tyr	Arg	Val	Thr	Tyr
	1490					1495					1500				
Ser	Ser	Pro	Glu	Asp	Gly	Ile	His	Glu	Leu	Phe	Pro	Ala	Pro	Asp	Gly
1505					1510					1515					1520
Glu	Glu	Asp	Thr	Ala	Glu	Leu	Gln	Gly	Leu	Arg	Pro	Gly	Ser	Glu	Tyr
				1525					1530					1535	
Thr	Val	Ser	Val	Val	Ala	Leu	His	Asp	Asp	Met	Glu	Ser	Gln	Pro	Leu
			1540					1545						1550	
Ile	Gly	Thr	Gln	Ser	Thr	Ala	Ile	Pro	Ala	Pro	Thr	Asp	Leu	Lys	Phe
		1555					1560						1565		
Thr	Gln	Val	Thr	Pro	Thr	Ser	Leu	Ser	Ala	Gln	Trp	Thr	Pro	Pro	Asn
	1570					1575					1580				
Val	Gln	Leu	Thr	Gly	Tyr	Arg	Val	Arg	Val	Thr	Pro	Lys	Glu	Lys	Thr
1585					1590					1595					1600
Gly	Pro	Met	Lys	Glu	Ile	Asn	Leu	Ala	Pro	Asp	Ser	Ser	Ser	Val	Val
				1605					1610					1615	
Val	Ser	Gly	Leu	Met	Val	Ala	Thr	Lys	Tyr	Glu	Val	Ser	Val	Tyr	Ala
			1620					1625					1630		
Leu	Lys	Asp	Thr	Leu	Thr	Ser	Arg	Pro	Ala	Gln	Gly	Val	Val	Thr	Thr
		1635					1640					1645			
Leu	Glu	Asn	Val	Gly	Pro	Pro	Arg	Arg	Ala	Arg	Val	Thr	Asp	Ala	Thr

-continued

Gly Phe Lys Leu Leu Cys Gln Cys Leu Gly Phe Gly Ser Gly His Phe
2065 2070 2075 2080

Arg Cys Asp Ser Ser Arg Trp Cys His Asp Asn Gly Val Asn Tyr Lys
2085 2090 2095

Ile Gly Glu Lys Trp Asp Arg Gln Gly Glu Asn Gly Gln Met Met Ser
2100 2105 2110

Cys Thr Cys Leu Gly Asn Gly Lys Gly Glu Phe Lys Cys Asp Pro His
2115 2120 2125

Glu Ala Thr Cys Tyr Asp Asp Gly Lys Thr Tyr His Val Gly Glu Gln
2130 2135 2140

Trp Gln Lys Glu Tyr Leu Gly Ala Ile Cys Ser Cys Thr Cys Phe Gly
2145 2150 2155 2160

Gly Gln Arg Gly Trp Arg Cys Asp Asn Cys Arg Arg Pro Gly Gly Glu
2165 2170 2175

Pro Thr Pro Glu Gly Thr Thr Gly Gln Ser Tyr Asn Gln Tyr Ser Gln
2180 2185 2190

Arg Tyr His Gln Arg Thr Asn Thr Asn Val Asn Cys Pro Ile Glu Cys
2195 2200 2205

Phe Met Pro Leu Asp Val Gln Ala Asp Arg Glu Asp Ser Arg Glu
2210 2215 2220

<210> SEQ ID NO 3
<211> LENGTH: 3185
<212> TYPE: DNA
<213> ORGANISM: Homo Sapiens
<220> FEATURE:
<223> OTHER INFORMATION: Fibronectin ED-A region
<300> PUBLICATION INFORMATION:
<308> DATABASE ACCESSION NUMBER: GenBank X07718
<309> DATABASE ENTRY DATE: 2005-04-18

<400> SEQUENCE: 3

```

atcaaacaga aatgactatt gaaggcttgc agcccacagt ggagtatgtg gttagtgtct    60
atgctcagaa tccaagcgga gagagtcagc ctctgggtca gactgcagta accagtacgt    120
aaccactgct tggtttccat tttcaaagtc aaatthttgtt cttgggtgtc tgaatgcca    180
cgacatgtct tttgcaatta cacatagga aagtgaactt gttggtagt ttatgtcttg    240
agctgagccc tttacgaaca tcttttttcc ttctcagtgc caagcgagga atttacagag    300
aaagaagttg tgaaccacc atagttagtt gctgtgcttt gaatttcttt ttgctcaaat    360
ggcctcagcg aaatcttatt tgcctatagc aaatctacaa aaaatthtcc tagaccgtct    420
ttctacaac tggatggtaa agttgattga agtgtgcctc atgtagcttt atgthttggg    480
catttgaag gctatggctg gaccagagtg taatataaat gcttaataga gaggggaaaa    540
gaagagtgtg agaaccatta tagggctggg ctacgcctg taatcccagc atthttgggag    600
gctgagggcg atcacagagt caggagttcg agaccagcct gaccaacatg gtgaaacccc    660
atctctacta aaaatacaaa aattagccag tgcggtggc acgtgcctgt aatcccagct    720
actcacggag gctgagcgag aagaatcact tggaccagg aggcagaagt tgcagtgagc    780
caagatcatg cctctgcacc ccagcctagg tgatagagtg agactccatc tcaaaaaaaaa    840
acaaaacaaa acaattataa caatttgaat ctgaaccata tgcaaatcag ctttaccact    900
tccaaggtat aagaaaaatcc aggtctatga gactaacatc acattgtaaa aatcaaatcg    960

```

-continued

tggtagaata cctttaaatt aatataaata catccccatt gtggggacat tttgcagggt	1020
atctgcttat ctacataca ccatgtttta ataagtgatg caacattgca ttttttctaa	1080
accaagaaaa attaagcaag tgtttaagtg atttttcctt tgatagtggy ttaattggac	1140
ttcatcaaaag aaaatggtat ctgcaaaaact gctttgcatg ttataaaaat gcttatttca	1200
caacttgctt toacataacc tcttaccatt aatttgccta acagacattg atcgccctaa	1260
aggactggca ttcaactgatg tggatgtcga ttccatcaaa attgcttggg aaagcccaca	1320
ggggcaagtt tccaggtaca ggggtaccta ctcgagccct gaggatggaa tccatgagct	1380
attccctgca cctgatggtg aagaagacac tgcagagctg caaggcctca gaccgggttc	1440
tgagtacaca gtcagtgtgg ttgccttgca cgatgatatg gagagccagc ccctgattgg	1500
aaccagctcc acaggtatat ggttaattgc acacaggtgc catgggagca gcggtttat	1560
gcctactgaa tgaattatgc ttcactgggc tattgattcc cgtgtaaggg tgaanaagaa	1620
ttattaggaa agatcctctt taaagaggaa tggtaagaaa caataaaact taggtgatat	1680
taaaggaac aagtctgatt aaaagaaatt ttggagtatc ctggcttata cacaagacca	1740
taaagcaaga ctttgaaga ggatactaaa gttgtgatt atttcctaag ctctgactcc	1800
ctgtgattac cctcactatg tataaagaaa agaagtttgg cattacagag cttacttata	1860
aaaaggaacc caaactggg catttcatag cagcatgatt ctgagcacac gtgggtaaga	1920
cctttcttct ctggttagat atcatatgct ggtgtataat tagcttaaat gattgtgatt	1980
tagacaccta ggaaataatc aatagggcaa ttgctttcca taatacttta tcttcttggt	2040
ctttatttct gaagcagagt agaagctaa agatgtatcc tagtgacagc ataaacccta	2100
gaggtgacag tctgtattat tgcttttcgc ttctcttttc tgcttctggt gggagccagt	2160
tttcttctta cgccgatta cagagagaac gtcaaattta gcaagccata tctgccatag	2220
gtccaaaata agagacaata aaaattatct tctctttttt ggatggaata ctgcgtgaaa	2280
tggttatcca tacaaagata ctttatgtag aatagaaaaa ggaggccggg tgcagtggct	2340
cacacatgta atcctagtgc ttggggagc taagccggga gcaactgattg aggccaggag	2400
ttcatgatca gcctgggcaa tgaagtgaga ccccgctctc acaaaaaaat atgaaaaaat	2460
tagcgaggty tggtagacaca tgcctgtagt cccagctact caagaggctg aggtagagga	2520
tcaactgagc ctacgagttc aaggctgcag tgagctatga taactccact gcaactgctc	2580
ctggatgaca cagagagacc gtttctaata taattaatta acaattttaa gaaagaaaaa	2640
gggccattgc ttatttttcc atacaaaagt aaaataaatc ataatggcca ataagccaat	2700
gtaacttttt tttttaaggg aaagcaaac ttgtaaaacc taaaatctct tagagttttg	2760
gcatttacc aaatgttttc agtgattctg agaattggtg gatataaac acatttctca	2820
gcaaacactt tcttcatttt gcatccctta ctgtactttc ttgtactgaa tctttgcttg	2880
accagggaac ccacctagcc caacaagaac aatccattct acttcttggg actacgttta	2940
tttctctttt ccccatctc ctataagata acctctaacc aatgacaatc tgcacagcta	3000
ttctgcacc aactgacctg aagttcactc aggtcacacc cacaagcctg agcgcccagt	3060
ggacaccacc caatgttcag ctcaactggat atcgagtgcg ggtgaccccc aaggagaaga	3120
ccggaccaat gaaagaagtc aacctgtctc ctgacagctc atccgtggtt gtatcaggac	3180
ttatc	3185

-continued

<210> SEQ ID NO 4
 <211> LENGTH: 190
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank X07718
 <309> DATABASE ENTRY DATE: 2005-04-18

<400> SEQUENCE: 4

Gln Thr Glu Met Thr Ile Glu Gly Leu Gln Pro Thr Val Glu Tyr Val
 1 5 10 15
 Val Ser Val Tyr Ala Gln Asn Pro Ser Gly Glu Ser Gln Pro Leu Val
 20 25 30
 Gln Thr Ala Val Thr Asn Ile Asp Arg Pro Lys Gly Leu Ala Phe Thr
 35 40 45
 Asp Val Asp Val Asp Ser Ile Lys Ile Ala Trp Glu Ser Pro Gln Gly
 50 55 60
 Gln Val Ser Arg Tyr Arg Val Thr Tyr Ser Ser Pro Glu Asp Gly Ile
 65 70 75 80
 His Glu Leu Phe Pro Ala Pro Asp Gly Glu Glu Asp Thr Ala Glu Leu
 85 90 95
 Gln Gly Leu Arg Pro Gly Ser Glu Tyr Thr Val Ser Val Val Ala Leu
 100 105 110
 His Asp Asp Met Glu Ser Gln Pro Leu Ile Gly Thr Gln Ser Thr Ala
 115 120 125
 Ile Pro Ala Pro Thr Asp Leu Lys Phe Thr Gln Val Thr Pro Thr Ser
 130 135 140
 Leu Ser Ala Gln Trp Thr Pro Pro Asn Val Gln Leu Thr Gly Tyr Arg
 145 150 155 160
 Val Arg Val Thr Pro Lys Glu Lys Thr Gly Pro Met Lys Glu Val Asn
 165 170 175
 Leu Ala Pro Asp Ser Ser Ser Val Val Val Ser Gly Leu Ile
 180 185 190

<210> SEQ ID NO 5
 <211> LENGTH: 2823
 <212> TYPE: DNA
 <213> ORGANISM: Homo Sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Fibronectin ED-B region
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank X07717
 <309> DATABASE ENTRY DATE: 2004-04-18

<400> SEQUENCE: 5

ctgcactttt gataacctga gtcccgccct ggagtacaat gtcagtgttt acaactgtcaa 60
 ggatgacaag gaaagtgtcc ctatctctga taccatcatc ccaggtaata gaaaataagc 120
 tgctatcctg agagtgcacat tccaataaga gtggggatta gcctcttaat ccccagatgc 180
 ttaagggtgt caactatatt tgggatttaa ttccgatctc ccagctgcac tttccaaaac 240
 caagaagtca aagcagcgat ttggacaaaa tgcttgctgt taacactgct ttactgtctg 300
 tgcttcaact ggatgctgtg tgttgacgag agtatgtaat ggagtggcag ccatggcttt 360
 aactctgtat tgtctgctca catggaagta tgactaaaac actgtcacgt gctctgtactc 420
 agtactgata ggctcaaagt aatatggtaa atgcatccca tcagtacatt tctgcccgat 480

-continued

tttacaatcc	atatcaatth	ccaacagctg	cctatthcat	cttgagttt	caaatccttc	540
ttttgaaa	ttgatttta	aaaaaagt	aagtaaaagt	cacacctca	gggttgtct	600
ttctgtggc	ctgaaagac	aacattgcaa	aggcctgtcc	taaggatagg	ctgtttgtc	660
cattgggta	taacataatg	aaagcattgg	acagatcgtg	tcccccttg	gactcttcag	720
tagaatgctt	ttactaacgc	taattacatg	ttttgattat	gaatgaacct	aaaatagtg	780
caatggcctt	aacctaggcc	tgtctttcct	cagcctgaat	gtgctttga	atggcacatt	840
tcacaccata	cattcataat	gcattagcgt	tatggccatg	atgttgcac	gagttttgta	900
tgggagaaa	aaaatcaatt	tatcacccat	ttattattht	ttccggtgt	tcatgcaagc	960
ttatthtcta	ctaaaacagt	tttggaaata	ttaaaagcat	tgctgatact	tacttcagat	1020
attatgtcta	ggctctaaga	atggtttcga	catcctaaac	agccatatga	tttttaggaa	1080
tctgaacagt	tcaaattgta	ccctttaagg	atgttttcaa	aatgtaaaaa	atatatatat	1140
atatatatat	tccttaaaag	aatattcctg	tttattcttc	tagggaagca	aactgttcat	1200
gatgcttagg	aagtctthtc	agagaattta	aaacagattg	catattacca	tcattgcttt	1260
aacattccac	caatthtact	actagtaacc	tgatatacac	tgctthtatt	ttcctcttt	1320
ttttcctct	atthtcttht	tgctcctccc	tcctthtct	ttgtaactca	atagaggtgc	1380
cccaactcac	tgacctaaag	tttgttgata	taaccgattc	aagcatcggc	ctgaggtgga	1440
ccccgtaaa	ctctccacc	attattgggt	accgcatcac	agtagttgcg	gcaggagaag	1500
gtatccctat	ttttgaagat	tttgtggact	cctcagtagg	atactacaca	gtcacagggc	1560
tggagccggg	cattgactat	gatatcagcg	ttatcactct	cattaatggc	ggcgagagt	1620
cccctactac	actgacacaa	caaacgggtg	aatthtgaaa	actctgctg	ttgagacata	1680
gatggtgtg	catgctgcca	ccagttaact	cggttaaata	tgatgtttc	atgggggaag	1740
tcagcaattg	gccaaagatt	cagataggtg	gaattggggg	gataaggaat	caaatgcac	1800
tgctaaactg	attggagaaa	aacacatgca	atatcttcag	tacactctca	tttaaaccac	1860
aagtagatat	aaagcctaga	gaaatacaga	tgtctgctct	gttaaatata	aaatagcaaa	1920
tgttcattca	atthgaagac	ctagaattht	tctcttaaa	taccaaacac	gaataccaaa	1980
ttgctgaagt	accaattgat	aagaatata	cacaaaatg	taccatcatg	ctcttccttc	2040
taccctthga	taaactctac	catgctcctt	ctthttagct	aaaaaccat	caaaatttag	2100
ggtagagtgg	atgggcattg	ttttgaggta	ggagaaaagt	aaacttggga	ccattctagg	2160
ttttgtgct	gtcactagg	aaagaacac	ctcttaacc	acagtctggg	gacaagcatg	2220
caacatthta	aaggttctct	gctgtgcatg	ggaaaagaaa	catgctgaga	accaatthgc	2280
atgaacatgt	tcacttgtaa	gtagaattca	ctgaatggaa	ctgtagctct	agatatctca	2340
catgggggga	agthtaggac	cctctgtct	ttttgtctgt	gtgcatgtat	ttctthgtaa	2400
agtactgcta	tgtthctct	tgctgtgtgg	caacttaagc	ctcttcggcc	tgggataaaa	2460
taatctgcag	tggatataat	aatgtacata	aagtcaacat	atthgaaagt	agattaaaat	2520
ctththtaaa	tatatcaatg	atggcaaaaa	ggttaaagg	ggcctaacag	tactgtgtgt	2580
agtgtthtat	thtaaacagt	agtacactat	aacttaaaat	agacttagat	tagactgttt	2640
gcatgattat	gattctgttt	cctthtatgca	tgaaatattg	atthtacctt	tcagctact	2700
tcgttagctt	taatthtaaa	atacattaac	tgagtcttcc	ttctgtctcg	aaaccagctg	2760

-continued

ttctctctcc cactgacctg cgattcacca acattggtcc agacaccatg cgtgtcacct 2820
 ggg 2823

<210> SEQ ID NO 6
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank X07717
 <309> DATABASE ENTRY DATE: 2005-04-18

<400> SEQUENCE: 6

Cys Thr Phe Asp Asn Leu Ser Pro Gly Leu Glu Tyr Asn Val Ser Val
 1 5 10 15
 Tyr Thr Val Lys Asp Asp Lys Glu Ser Val Pro Ile Ser Asp Thr Ile
 20 25 30
 Ile Pro Glu Val Pro Gln Leu Thr Asp Leu Ser Phe Val Asp Ile Thr
 35 40 45
 Asp Ser Ser Ile Gly Leu Arg Trp Thr Pro Leu Asn Ser Ser Thr Ile
 50 55 60
 Ile Gly Tyr Arg Ile Thr Val Val Ala Ala Gly Glu Gly Ile Pro Ile
 65 70 75 80
 Phe Glu Asp Phe Val Asp Ser Ser Val Gly Tyr Tyr Thr Val Thr Gly
 85 90 95
 Leu Glu Pro Gly Ile Asp Tyr Asp Ile Ser Val Ile Thr Leu Ile Asn
 100 105 110
 Gly Gly Glu Ser Ala Pro Thr Thr Leu Thr Gln Gln Thr Ala Val Pro
 115 120 125
 Pro Pro Thr Asp Leu Arg Phe Thr Asn Ile Gly Pro Asp Thr Met Arg
 130 135 140
 Val Thr Trp
 145

<210> SEQ ID NO 7
 <211> LENGTH: 360
 <212> TYPE: DNA
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 7

gacgagcttc cccaactggt aacccttcca cacccaatc ttcattggacc agagatcttg 60
 gatgttcctt ccacagtcca aaagaccctt ttcgtcacc accctgggta tgacactgga 120
 aatggtattc agcttctctg cacttctggt cagcaaccca gtgttgggca acaaatgatc 180
 tttgaggaac atggttttag gcggaccaca cggccacaaa cggccacccc cataaggcat 240
 aggccaagac cataccggcc gaatgtaggt gaggaaatcc aaattggtca cattcccagg 300
 gaagatgtag actatcacct gtaccacac ggtccgggac tcaatccaaa tgcctctaca 360

<210> SEQ ID NO 8
 <211> LENGTH: 120
 <212> TYPE: PRT
 <213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 8

Asp Glu Leu Pro Gln Leu Val Thr Leu Pro His Pro Asn Leu His Gly
 1 5 10 15

-continued

Pro Glu Ile Leu Asp Val Pro Ser Thr Val Gln Lys Thr Pro Phe Val
 20 25 30

Thr His Pro Gly Tyr Asp Thr Gly Asn Gly Ile Gln Leu Pro Gly Thr
 35 40 45

Ser Gly Gln Gln Pro Ser Val Gly Gln Gln Met Ile Phe Glu Glu His
 50 55 60

Gly Phe Arg Arg Thr Thr Pro Pro Thr Thr Ala Thr Pro Ile Arg His
 65 70 75 80

Arg Pro Arg Pro Tyr Pro Pro Asn Val Gly Glu Glu Ile Gln Ile Gly
 85 90 95

His Ile Pro Arg Glu Asp Val Asp Tyr His Leu Tyr Pro His Gly Pro
 100 105 110

Gly Leu Asn Pro Asn Ala Ser Thr
 115 120

<210> SEQ ID NO 9
 <211> LENGTH: 714
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: ME4C
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank AJ297960
 <309> DATABASE ENTRY DATE: 2001-03-01

<400> SEQUENCE: 9

```
gaggtgcagc tgttgagtc tgggggagc ttggtacagc ctggggggtc cctgagactc   60
tcctgtgcag cctctggatt cacctttagc agtttttcga tgagctgggt ccgccaggct   120
ccaggaagg ggctggagt ggtctcatct attagtggta gttcgggtac cacatactac   180
gcagactccg tgaaggccg gttcaccatc tccagagaca attccaagaa cacgctgtat   240
ctgcaaatga acagcctgag agccgaggac acggccgtat attactgtgc gaaacagaag   300
agtgcgccgt ttgactactg gggccagga accctggtca cctctcagag tggcgatggg   360
tccagtggcg gtagcggggg cgcgtcgtct tctgagctga ctcaggaccc tgctgtgtct   420
gtggccttgg gacagacagt caggatcaca tgccaaggag acagcctcag aagctattat   480
gcaagctggt accagcagaa gccaggacag gccctgtac ttgtcatcta tggtaaaaac   540
aaccggccct cagggatccc agaccgattc tctggctcca gctcaggaaa cacagcttcc   600
ttgaccatca ctggggctca ggcggaagat gaggtgact attactgtaa ctctctgctg   660
cccgttagta atagggtggt attcggcgga gggaccaagc tgaccgtcct aggc       714
```

<210> SEQ ID NO 10
 <211> LENGTH: 238
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: ME4C anti-(ED-B) scFv recombinant antibody
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank CAC33765
 <309> DATABASE ENTRY DATE: 2000-03-01

<400> SEQUENCE: 10

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Phe
 20 25 30

-continued

Ser Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Ser Ile Ser Gly Ser Ser Gly Thr Thr Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Gln Lys Ser Ala Pro Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ser Gly Asp Gly Ser Ser Gly Gly Ser Gly Gly Ala
115 120 125

Ser Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly
130 135 140

Gln Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg Ser Tyr Tyr
145 150 155 160

Ala Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile
165 170 175

Tyr Gly Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly
180 185 190

Ser Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly Ala Gln Ala
195 200 205

Glu Asp Glu Ala Asp Tyr Tyr Cys Asn Ser Ser Ala Pro Val Ser Asn
210 215 220

Arg Val Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
225 230 235

<210> SEQ ID NO 11

<211> LENGTH: 1975

<212> TYPE: DNA

<213> ORGANISM: Homo Sapiens

<220> FEATURE:

<223> OTHER INFORMATION: GalNAc-T3 gene, 3' UTR

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank Y10345

<309> DATABASE ENTRY DATE: 2000-10-27

<400> SEQUENCE: 11

```

taagtgttcc ttaaaattaa gttgaaaaag gaaatattct ttctcataaa actgtgacta    60
ggcatacaact gtagtttttg aaaattatgc aaaagcagct aaatgtaact tattccaagt    120
gcatttttct tatttatatc tttatgtagc actactacag aaattctgca agttttctgtt    180
tcaaagcaca ataactagta ataccaaaga ctatttcaaa atgtccagat gtaggggaag    240
agatgtttac agtatgatga aaataathtt ccaagtaaag tgatgtttgt gtgttttgta    300
cacttaggga tatatatata tagctacatt cacacactca caatttaaaa tatttcccct    360
agttttttgg ggggatagga agaaagattt gttactgtat ttttttaact acataaaaat    420
agatcaataa atgtcagcat tggcctctgt gtacaaacca agagctttta cagatccaga    480
atttattagt ttaaaatgca ggtgaacttt tttttgcgtt tggtttactt gtctgtcaaa    540
tgtttcctta aacatgaaac tgaataagga gaagagtatt ttaaacactt aaatttcttg    600
gcaaatttta aacatttttt tagtctgtaa taaactccac ttgaagcact taagtcttcc    660
ttaaagact tttcttaagt aatgatactg tgtgttttcc caaagcactt ttaaaaaaat    720

```


-continued

```

ttttataaat tactatctgt tgaaaagggt tccttttcct ttcttctagt attttttttc 780
ttacccaaaat tcactaatct tgaatgtttg tgatattaaa tttcaaatgc agaatacttg 840
actcatttaa agctaaatth tgttactgat tcaattataa ttgtaatgga tttttgactt 900
tgtaatggat tcttttcctc aaaaagcctt attatttttt atctatgtgg aaaacacaat 960
aaaaaatcct caacactatt gtaatcattt ggttaagtgc ttattcctct tttgggtaaa 1020
atctgtaatt gataataggt gggggaaaat gaattttgta tgctgaattt ctaagcgctt 1080
attgtttgta aaacatcag atatttctta tggcacaaaa aatgaggaat agcaaaattc 1140
ctgtgttcaa tatttagaaa attttgattt aatttctgat aaagtccctt aagcatctga 1200
tagaatgatg ttttaaaaa atttgacgct tgcttaggag atttaccact tttttttttt 1260
gtttttcgtc attttatatt tagatctcct gtattcctgt tcccgaagta aaatacgatc 1320
ggtttcatat tttaaatctg gcagagcctc agctgtacga aaaagagcat atactgggta 1380
ttgaccctat cttctcattg tttgtttgta agtttgaatt tgtattaaaa agcctgcatt 1440
ctgagctgga catggtggct cagcttctaa tcccagcact ttggtaggca aagggtggag 1500
gatcatttga gctcaggagt tccagaccag cctgggcaac atagcaaaat ctcatctcta 1560
caaaaagtaa aaatataaaa atgaaattaa aaataaaatt acctaggtgt ggtggcacgc 1620
atctgtagtt ccagctatac aggaagggtg gcagaagca ttgcttgagc ttgggagatc 1680
gaggctacag tgagctatga ttacaccact gcacttcagt ctgtgtgact gagcaagact 1740
ctttcaaaaa aaaaaaaag cctacattct ccagttgatt atttccaact aatgtgtatt 1800
atgtgcctaa ttttctatca gaagttgat taagcccgtt ttcacactgc tgttaaagac 1860
atacctgaga ctgggtaatt tataaagaaa aataggttca atggaccac aggtcccgtt 1920
ggctggggaa gcttcacaat catggcggaa ggtgaaagca tgtcttacgt ggaag 1975

```

<210> SEQ ID NO 12

<211> LENGTH: 1902

<212> TYPE: DNA

<213> ORGANISM: Homo Sapiens

<220> FEATURE:

<223> OTHER INFORMATION: UDP-GalNAc:polypeptide N-acetylgalactosaminyl transferase

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank X92689

<309> DATABASE ENTRY DATE: 2000-01-07

<400> SEQUENCE: 12

```

atggctcacc taaagcgact agtaaaatta cacattaaaa gacattacca taaaaagttc 60
tggaagcttg gtgcagtaat ttttttcttt ataatagttt tggttttaat gcaaagagaa 120
gtaagtgttc aatattccaa agaggaatca aggatggaaa ggaacatgaa aaacaaaaac 180
aagatgttgg atttaagtct agaagctgta aacaatatta aggatgccat gccaaaaatg 240
caaataggag cacctgtcag gcaaaacatt gatgctggty agagacctg tttgcaagga 300
tattatacag cagcagaatt gaagcctgtc cttgaccgtc cacctcagga ttcaaatgca 360
cctgtgtcct ctggtaaagc attcaagaca accaatttaa gtgttgaaga gcaaaaggaa 420
aaggaacgtg gggaagctaa aactgtcttt aatgctttcg caagtgcag gattttctttg 480
caccgagatc ttggaccaga cactcgcact cctgaatgta ttgaacaaaa atttaagcgc 540
tgccctcccc tgcccaccac cagtgtcata atagtttttc ataatgaagc gtggtccacg 600

```

-continued

```

ttgcttagaa ctgtccacag tgtgctctat tcttcacctg caatactgct gaaggaaatc 660
atitttggtgg atgatgctag tgtagatgag tactttacatg ataaactaga tgaatatgta 720
aaacaatttt ctatagtaaa aatagtcaga caaagagaaa gaaaaggctct gatcactgct 780
cggttgctag gagcaacagt cgcaacagct gaaacgctca catttttaga tgctcactgt 840
gagtgtttct atggttggtct agaacctctg ttggccagaa tagctgagaa ctacacggct 900
gtcgttaagtc cagatattgc atccatagat ctgaacacgt ttgaattcaa caaaccttct 960
ccttatggaa gtaaccataa ccgtaggaaat tttgactgga gtctttcatt tggctgggag 1020
tcgcttcctg atcatgagaa gcaaagaagg aaagatgaaa cctaccaaat taaaacaccc 1080
acttttgtag gaggactttt ttccatatca aaagaatatt ttgagtatat tggaaactat 1140
gatgaagaaa tggaaatctg gggaggtgaa aatatagaaa tgcctttcag agtatggcaa 1200
tgtgtggggc agttggagat tatgccttgc tctgttgttg gacatgtttt tcgcagcaaa 1260
agccctcata gctttccaaa aggcactcag gtgattgcta gaaaccaagt tcgccttgca 1320
gaaagtctga tggatgaata caaggaaata ttttatagga gaaatacaga tgcagcaaaa 1380
attgttaaac aaaaagcatt tgggtgatctt tcaaaaagat ttgaaataaa acaccgtctt 1440
cggtgtaaaa attttacatg gtatctgaac aacatttatac cagaggtgta tgtgccagac 1500
cttaatcctg ttatatctgg atacattaaa agcgttggtc agcctctatg tctggatggt 1560
ggagaaaaca atcaaggagg caaacatta attatgtata catgtcatgg acttggggga 1620
aaccagtact ttgaatactc tgctcaacat gaaattcggc acaacatcca gaaggaatta 1680
tgtcttcag ctgctcaagg tctcgttcag ctgaaggcat gtacctaca aggtcacaag 1740
acagttgtca ctggagagca gatatgggag atccagaagg atcaacttct atacaatcca 1800
ttcttaaaaa tgtgcctttc agcaaatgga gagcatccaa gtttagtgtc atgcaacca 1860
tcagatccac tccaaaaatg gatacttagc caaaatgatt aa 1902

```

<210> SEQ ID NO 13

<211> LENGTH: 633

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank CAA63371

<309> DATABASE ENTRY DATE: 2000-01-07

<400> SEQUENCE: 13

```

Met Ala His Leu Lys Arg Leu Val Lys Leu His Ile Lys Arg His Tyr
 1             5             10             15

His Lys Lys Phe Trp Lys Leu Gly Ala Val Ile Phe Phe Phe Ile Ile
      20             25             30

Val Leu Val Leu Met Gln Arg Glu Val Ser Val Gln Tyr Ser Lys Glu
      35             40             45

Glu Ser Arg Met Glu Arg Asn Met Lys Asn Lys Asn Lys Met Leu Asp
      50             55             60

Leu Met Leu Glu Ala Val Asn Asn Ile Lys Asp Ala Met Pro Lys Met
      65             70             75             80

Gln Ile Gly Ala Pro Val Arg Gln Asn Ile Asp Ala Gly Glu Arg Pro
      85             90             95

Cys Leu Gln Gly Tyr Tyr Thr Ala Ala Glu Leu Lys Pro Val Leu Asp
      100            105            110

```

-continued

Arg Pro Pro Gln Asp Ser Asn Ala Pro Gly Ala Ser Gly Lys Ala Phe
 115 120 125
 Lys Thr Thr Asn Leu Ser Val Glu Glu Gln Lys Glu Lys Glu Arg Gly
 130 135 140
 Glu Ala Lys His Cys Phe Asn Ala Phe Ala Ser Asp Arg Ile Ser Leu
 145 150 155 160
 His Arg Asp Leu Gly Pro Asp Thr Arg Pro Pro Glu Cys Ile Glu Gln
 165 170 175
 Lys Phe Lys Arg Cys Pro Pro Leu Pro Thr Thr Ser Val Ile Ile Val
 180 185 190
 Phe His Asn Glu Ala Trp Ser Thr Leu Leu Arg Thr Val His Ser Val
 195 200 205
 Leu Tyr Ser Ser Pro Ala Ile Leu Leu Lys Glu Ile Ile Leu Val Asp
 210 215 220
 Asp Ala Ser Val Asp Glu Tyr Leu His Asp Lys Leu Asp Glu Tyr Val
 225 230 235 240
 Lys Gln Phe Ser Ile Val Lys Ile Val Arg Gln Arg Glu Arg Lys Gly
 245 250 255
 Leu Ile Thr Ala Arg Leu Leu Gly Ala Thr Val Ala Thr Ala Glu Thr
 260 265 270
 Leu Thr Phe Leu Asp Ala His Cys Glu Cys Phe Tyr Gly Trp Leu Glu
 275 280 285
 Pro Leu Leu Ala Arg Ile Ala Glu Asn Tyr Thr Ala Val Val Ser Pro
 290 295 300
 Asp Ile Ala Ser Ile Asp Leu Asn Thr Phe Glu Phe Asn Lys Pro Ser
 305 310 315 320
 Pro Tyr Gly Ser Asn His Asn Arg Gly Asn Phe Asp Trp Ser Leu Ser
 325 330 335
 Phe Gly Trp Glu Ser Leu Pro Asp His Glu Lys Gln Arg Arg Lys Asp
 340 345 350
 Glu Thr Tyr Pro Ile Lys Thr Pro Thr Phe Ala Gly Gly Leu Phe Ser
 355 360 365
 Ile Ser Lys Glu Tyr Phe Glu Tyr Ile Gly Ser Tyr Asp Glu Glu Met
 370 375 380
 Glu Ile Trp Gly Gly Glu Asn Ile Glu Met Ser Phe Arg Val Trp Gln
 385 390 395 400
 Cys Gly Gly Gln Leu Glu Ile Met Pro Cys Ser Val Val Gly His Val
 405 410 415
 Phe Arg Ser Lys Ser Pro His Ser Phe Pro Lys Gly Thr Gln Val Ile
 420 425 430
 Ala Arg Asn Gln Val Arg Leu Ala Glu Val Trp Met Asp Glu Tyr Lys
 435 440 445
 Glu Ile Phe Tyr Arg Arg Asn Thr Asp Ala Ala Lys Ile Val Lys Gln
 450 455 460
 Lys Ala Phe Gly Asp Leu Ser Lys Arg Phe Glu Ile Lys His Arg Leu
 465 470 475 480
 Arg Cys Lys Asn Phe Thr Trp Tyr Leu Asn Asn Ile Tyr Pro Glu Val
 485 490 495
 Tyr Val Pro Asp Leu Asn Pro Val Ile Ser Gly Tyr Ile Lys Ser Val
 500 505 510

-continued

Gly Gln Pro Leu Cys Leu Asp Val Gly Glu Asn Asn Gln Gly Gly Lys
 515 520 525

Pro Leu Ile Met Tyr Thr Cys His Gly Leu Gly Gly Asn Gln Tyr Phe
 530 535 540

Glu Tyr Ser Ala Gln His Glu Ile Arg His Asn Ile Gln Lys Glu Leu
 545 550 555 560

Cys Leu His Ala Ala Gln Gly Leu Val Gln Leu Lys Ala Cys Thr Tyr
 565 570 575

Lys Gly His Lys Thr Val Val Thr Gly Glu Gln Ile Trp Glu Ile Gln
 580 585 590

Lys Asp Gln Leu Leu Tyr Asn Pro Phe Leu Lys Met Cys Leu Ser Ala
 595 600 605

Asn Gly Glu His Pro Ser Leu Val Ser Cys Asn Pro Ser Asp Pro Leu
 610 615 620

Gln Lys Trp Ile Leu Ser Gln Asn Asp
 625 630

<210> SEQ ID NO 14
 <211> LENGTH: 8815
 <212> TYPE: DNA
 <213> ORGANISM: Homo Sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Oncofetal Fibronectin variant 1
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank NM 212482
 <309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 14

```

gcccgcgccg gctgtgctgc acagggggag gagagggaac cccaggcgcg agcgggaaga      60
ggggacctgc agccacaact tctctgtgcc tctgcatccc ttctgtccct ccacccgtcc      120
ccttccccac cctctggccc ccaccttctt ggaggcgaca acccccggga ggcattagaa      180
gggatttttc ccgcaggttg cgaagggag caaacttggg ggcaacttgc ctcccgggtg      240
ggggctctct cccccaccgt ctcaaatgct ttaggggtcc ggggccggg ctgctgctgc      300
tggccgtcca gtgcctgggg acagcgggtg cctccacggg agcctcgaag agcaagaggc      360
aggctcagca aatggttcag ccccagtccc cggtggctgt cagtcaaagc aagcccggtt      420
gttatgacaa tggaaaacac tatcagataa atcaacagtg ggagcggacc tacctaggca      480
atgctgtggt ttgtacttgt tatggaggaa gccgaggttt taactgcgag agtaaacctg      540
aagctgaaga gacttgcttt gacaagtaca ctgggaacac ttaccgagtg ggtgacactt      600
atgagcgtcc taaagactcc atgatctggg actgtacctg catcggggct gggcggagga      660
gaataagctg taccatcgca aaccgctgcc atgaaggggg tcagtcctac aagattggtg      720
acacctggag gagaccacat gagactggtg gttacatggt agagtgtgtg tgtcttggtg      780
atggaaaagg agaatggacc tgcaagccca tagctgagaa gtgttttgat catgctgctg      840
ggacttccta tgtggtcgga gaaacgtggg agaagcccta ccaaggctgg atgatggtag      900
attgtacttg cctgggagaa ggcagcggac gcatcacttg cacttctaga aatagatgca      960
acgatcagga cacaaggaca tcctatagaa ttggagacac ctggagcaag aaggataatc     1020
gaggaaacct gctccagtgc atctgcacag gcaacggccg aggagagtgg aagtgtgaga     1080
ggcacacctc tgtgcagacc acatcgagcg gatctggccc cttcaccgat gttcgtgcag     1140
ctgtttacca accgcagcct caccocagc ctccctcccta tggccactgt gtcacagaca     1200
    
```

-continued

gtggtgtggt ctactctgtg gggatgcagt ggctgaagac acaaggaat aagcaaatgc	1260
tttgacacgtg cctgggcaac ggagtcagct gccaaagagac agctgtaacc cagacttacg	1320
gtggcaactc aaatggagag ccatgtgtct taccattcac ctacaatggc aggacgttct	1380
actcctgcac cacagaaggc cgacaggacg gacatccttg gtgcagcaca acttcgaatt	1440
atgagcagga ccagaaatac tctttctgca cagaccacac tgttttggtt cagactcgag	1500
gaggaaattc caatggtgcc ttgtgccact tccccttctc atacaacaac cacaattaca	1560
ctgattgcac ttctgagggc agaagagaca acatgaagtg gtgtgggacc acacagaact	1620
atgatgccga ccagaagttt gggttctgcc ccatggctgc ccacgaggaa atctgcacaa	1680
ccaatgaagg ggtcatgtac cgcattggag atcagtggga taagcagcat gacatgggtc	1740
acatgatgag gtgcacgtgt gttgggaatg gtcgtgggga atggacatgc attgcctact	1800
cgagccttcg agatcagtgc attgttgatg acatcactta caatgtgaac gacacattcc	1860
acaagcgtca tgaagagggg cacatgctga actgtacatg cttcggtcag ggtcggggca	1920
ggtggaagtg tgatcccgtc gaccaatgcc aggattcaga gactgggacg ttttatcaaa	1980
ttggagattc atgggagaag tatgtgcatg gtgtcagata ccagtgtac tgctatggcc	2040
gtggcattgg ggagtggcat tgccaacctt tacagaccta tccaagctca agtggctctg	2100
tcgaagtatt tatcactgag actccagtc agcccaactc ccacccatc cagtggaatg	2160
caccacagcc atctcacatt tccaagtaca ttctcagtg gagacctaaa aattctgtag	2220
gccgttgga ggaagctacc ataccaggcc acttaaaactc ctacaccatc aaaggcctga	2280
agcctggtgt ggtatacagag gccacgtca tcagcatcca gcagtacggc caccaagaag	2340
tgactcgctt tgacttcacc accaccagca ccagcacacc tgtgaccagc aacaccgtga	2400
caggagagac gactcccttt tctcctcttg tggccacttc tgaatctgtg accgaaatca	2460
cagccagtag ctttgtggtc tcttgggtct cagcttccga caccgtgtcg ggattccggg	2520
tggaatatga gctgagtgag gaggagatg agccacagta cctggatcct ccaagcacag	2580
ccacttctgt gaacatccct gacctgcttc ctggccgaaa atacattgta aatgtctatc	2640
agatatctga ggatggggag cagagtttga tctgtctac ttcacaaaca acagcgctg	2700
atgccctcc tgacccgact gtggaccaag ttgatgacac ctcaattgtt gttcgtgga	2760
gcagacccca ggtcccatc acagggtaca gaatagtcta ttcgcatca gtagaaggta	2820
gcagacaga actcaacctt cctgaaactg caaactccgt caccctcagt gacttgcaac	2880
ctggtgttca gtataacatc actatctatg ctgtggaaga aaatcaagaa agtacacctg	2940
ttgtcattca acaagaaacc actggcacc caccctcaga tacagtgcc tctcccaggg	3000
acctgcagtt tgtggaagt acagacgtga aggtcacat catgtggaca ccgcctgaga	3060
gtgcagtgac cggctaccgt gtggatgta tcccctcaa cctgcctggc gagcacgggc	3120
agaggctgcc catcagcag aacaccttg cagaagtac cgggctgtcc cctggggtca	3180
cctattactt caaagtctt gcagtgagc atgggagga gagcaagcct ctgactgctc	3240
aacagacaac caaactggat gctcccacta acctccagtt tgtcaatgaa actgattcta	3300
ctgtcctggt gagatggact ccacctcgg cccagataac aggataccga ctgacctgg	3360
gccttaccg aagagagac cccaggcagt acaatgtggg tccctctgtc tccaagtacc	3420
cactgaggaa tctgcagcct gcatctgagt acaccgtatc cctcgtggcc ataagggca	3480

-continued

accaagagag	ccccaaagcc	actggagtct	ttaccacact	gcagcctggg	agetctattc	3540
caccttacia	caccgagggt	actgagacca	ccattgtgat	cacatggacg	cctgctccaa	3600
gaattggttt	taagctgggt	gtacgaccaa	gccagggagg	agaggcacca	cgagaagtga	3660
cttcagactc	aggaagcatc	gttgtgtccg	gcttgactcc	aggagtagaa	tacgtctaca	3720
ccatccaagt	cctgagagat	ggacagggaa	gagatgcgcc	aattgtaaac	aaagtgggta	3780
caccattgtc	tccaccaaca	aacttgcata	tggaggcaaa	ccctgacact	ggagtgtctc	3840
cagtctcctg	ggagaggagc	accaccccag	acattactgg	ttagagaatt	accacaaccc	3900
ctacaaacgg	ccagcaggga	aattcttttg	aagaagtggg	ccatgctgat	cagagctcct	3960
gcacttttga	taacctgagt	cccggcctgg	agtacaatgt	cagtgtttac	actgtcaagg	4020
atgacaagga	aagtgtccct	atctctgata	ccatcatccc	agaggtgccc	caactcactg	4080
acctaagctt	tgttgatata	accgattcaa	gcacggccct	gaggtggacc	ccgctaaact	4140
cttccaccat	tattgggtac	cgcatcacag	tagttgcggc	aggagaaggt	atccctatct	4200
ttgaagatct	tgtggactcc	tcagtaggat	actacacagt	cacaggctg	gagccgggca	4260
ttgactatga	tatcagcgtt	atcactctca	ttaatggcgg	cgagagtgcc	cctactacac	4320
tgacacaaca	aacggctggt	cctcctccca	ctgacctgcg	attcaccaac	attggtccag	4380
acaccatgcg	tgtcacctgg	gctccacccc	catccattga	tttaaccaac	ttcctgggtg	4440
ggtactcacc	tgtgaaaaat	gaggaagatg	ttgcagagtt	gtcaatttct	ccttcagaca	4500
atgcagtggt	cttaacaaat	ctcctgcctg	gtacagaata	tgtagttagt	gtctccagtg	4560
tctacgaaca	acatgagagc	acacctctta	gaggaagaca	gaaaaacaggt	cctgattccc	4620
caactggcat	tgacttttct	gatattactg	ccaactcttt	tactgtgcac	tggattgtct	4680
ctcgagccac	catcactggc	tacaggatcc	gccatcatcc	cgagcacttc	agtgggagac	4740
ctcgagaaga	tcgggtgccc	cactctcgga	attccatcac	cctcaccaac	ctcactccag	4800
gcacagagta	tgtggctcag	atcgttgctc	ttaatggcag	agaggaaagt	cccttattga	4860
ttggccaaca	atcaacagtt	tctgatgttc	cgagggacct	ggaagttggt	gctgcgaccc	4920
ccaccagcct	actgatcagc	tgggatgtct	ctgctgtcac	agtgagatat	tacaggatca	4980
cttacggaga	gacaggagga	aatagccctg	tccaggagtt	cactgtgcct	gggagcaagt	5040
ctacagctac	catcagcggc	cttaaacctg	gagttgatta	taccatcact	gtgtatgctg	5100
tcactggccg	tggagacagc	cccgcaagca	gcaagccaat	ttccattaat	taccgaacag	5160
aaattgacaa	accatcccag	atgcaagtga	cogatgttca	ggacaacagc	attagtgtca	5220
agtggctgcc	ttcaagttcc	cctgttactg	gttacagagt	aaccaccact	ccccaaaatg	5280
gaccaggacc	aacaaaaact	aaaactgcag	gtccagatca	aacagaaatg	actattgaag	5340
gcttgacgcc	cacagtggag	tatgtggtta	gtgtctatgc	tcagaatcca	agcggagaga	5400
gtcagcctct	ggttcagact	gcagtaacca	acattgatcg	ccctaaagga	ctggcattca	5460
ctgatgtgga	tgtcgattcc	atcaaaaattg	cttgggaaag	cccacagggg	caagtttcca	5520
ggtacagggg	gacctactcg	agccctgagg	atggaatcca	tgagctattc	cctgcacctg	5580
atggtgaaga	agacactgca	gagctgcaag	gcctcagacc	gggttctgag	tacacagtca	5640
gtgtggttgc	cttgacagat	gatatggaga	gccagccct	gattggaacc	cagtccacag	5700
ctattcctgc	accaactgac	ctgaagttca	ctcaggtcac	accacaagc	ctgagcggcc	5760

-continued

agtggacacc	acccaatggt	cagctcactg	gatatcgagt	gcgggtgacc	cccaaggaga	5820
agaccggacc	aatgaaagaa	atcaaccttg	ctcctgacag	ctcatccgtg	gttgtatcag	5880
gacttatggt	ggccacaaa	tatgaagtga	gtgtctatgc	tcttaaggac	actttgacaa	5940
gcagaccagc	tcaggaggtt	gtcaccactc	tggagaatgt	cagcccacca	agaagggtc	6000
gtgtgacaga	tgctactgag	accaccatca	ccattagctg	gagaaccaag	actgagacga	6060
tcactggctt	ccaagttgat	gccgttccag	ccaatggcca	gactccaatc	cagagaacca	6120
tcaagccaga	tgtcagaagc	tacaccatca	caggtttaca	accaggcact	gactacaaga	6180
tctacctgta	caccttgaat	gacaatgctc	ggagctcccc	tgtggtcatc	gacgcctcca	6240
ctgccattga	tgaccatcc	aacctgcgtt	tcctggccac	cacaccaat	tccttgctgg	6300
tatcatggca	gccgccacct	gccaggatta	ccggctacat	catcaagtat	gagaagcctg	6360
ggtctcctcc	cagagaagtg	gtccctcggc	cccgccctgg	tgtcacagag	gctactatta	6420
ctggcctgga	accgggaacc	gaatatacaa	tttatgtcat	tgccctgaag	aataatcaga	6480
agagcgagcc	cctgattgga	aggaaaaaga	cagacgagct	tccccaactg	gtaacccttc	6540
cacaccccaa	tcttcatgga	ccagagatct	tggatgttcc	ttccacagtt	caaaagacct	6600
ctttcgtcac	ccaccctggg	tatgacactg	gaaatggtat	tcagcttctc	ggcacttctg	6660
gtcagcaacc	cagtgttggg	caacaaatga	tctttgagga	acatggtttt	aggcggacca	6720
caccgccacc	aacggccacc	cccataaggc	ataggccaag	accatacccg	ccgaatgtag	6780
gtgaggaat	caaattggt	cacatcccca	gggaagatgt	agactatcac	ctgtaccac	6840
acggtccggg	actcaatcca	aatgcctcta	caggacaaga	agctctctct	cagacaacca	6900
tctcatgggc	cccattccag	gacacttctg	agtacatcat	ttcatgtcat	cctgttggca	6960
ctgatgaaga	acccttacag	ttcagggttc	ctggaacttc	taccagtgcc	actctgacag	7020
gcctcaccag	aggtgccacc	tacaacatca	tagtggaggc	actgaaagac	cagcagaggc	7080
ataaggttcg	ggaagaggtt	gttacogtgg	gcaactctgt	caacgaaggc	ttgaaccaac	7140
ctacggatga	ctcgtgcttt	gacccttaca	cagtttccca	ttatgccgtt	ggagatgagt	7200
gggaacgaat	gtctgaatca	ggctttaaac	tgttgtgcca	gtgcttaggc	tttgggaagt	7260
gtcatttcag	atgtgattca	tctagatggt	gccatgacaa	tggtgtgaac	tacaagattg	7320
gagagaagtg	ggaccgtcag	ggagaaaatg	gccagatgat	gagctgcaca	tgtcttgggg	7380
acggaaaagg	agaattcaag	tgtgaccctc	atgaggcaac	gtgttatgat	gatgggaaga	7440
cataccacgt	aggagaacag	tggcagaagg	aatatctcgg	tgccatttgc	tcctgcacat	7500
gctttggagg	ccagcggggc	tggcgctgtg	acaactgccg	cagacctggg	ggtgaacca	7560
gtcccgaagg	cactactggc	cagtcctaca	accagtattc	tcagagatac	catcagagaa	7620
caaacactaa	tgtaatttgc	ccaattgagt	gottcatgcc	tttagatgta	caggctgaca	7680
gagaagattc	ccgagagtaa	atcatcttcc	caatccagag	gaacaagcat	gtctctctgc	7740
caagatccat	ctaaactgga	gtgatgttag	cagaccagc	ttagagttct	tctttcttcc	7800
taaagcctt	tgctctggag	gaagttctcc	agcttcagct	caactcacag	cttctccaag	7860
catcacccctg	ggagtttctc	gagggtttcc	tcataaatga	gggctgcaca	ttgcctgttc	7920
tgcttcgaag	tattcaatcc	cgctcagtat	tttaaatgaa	gtgattctaa	gattttggtt	7980
gggatcaata	ggaaagcata	tcagoccaac	caagatgcaa	atgttttgaa	atgatatgac	8040

-continued

```

caaaatttta agtaggaaag tcacccaac acttctgctt tcacttaagt gtctggcccg 8100
caatactgta ggaacaagca tgatcttggt actgtgatat tttaaatata cacagtactc 8160
actttttcca aatgatccta gtaattgcct agaaatatct ttctcttacc tgttatttat 8220
caatttttcc cagtattttt atacggaaaa aattgtattg aaaacactta gtagcagtt 8280
gataagagga atttgggata attatgggtg gtgattattt tttatactgt atgtgccaaa 8340
gctttactac tgtggaaaga caactgtttt aataaaagat ttacattcca caacttgaag 8400
ttcatctatt tgatataaga caccttcggg ggaaataatt cctgtgaata ttctttttca 8460
attcagcaaa catttgaaaa tctatgatgt gcaagtctaa ttgttgattt cagtacaaga 8520
ttttctaaat cagttgctac aaaaactgat tggtttttgt cacttcatct ctcoactaat 8580
ggagatagct ttacactttc tgctttaata gatttaagtg gacccaata tttattaaaa 8640
ttgctagttt accgttcaga agtataatag aaataatctt tagttgctct tttctaacca 8700
ttgtaattct tcccttcttc cctccacett tccttcattg aataaacctc tgttcaaaga 8760
gattgctgac aagggaata aaaatgacta agatattaaa aaaaaaaaaa aaaaa 8815

```

<210> SEQ ID NO 15

<211> LENGTH: 2477

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank NM 212482

<309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 15

```

Met Leu Arg Gly Pro Gly Pro Gly Leu Leu Leu Leu Ala Val Gln Cys
 1                    5                    10                    15

Leu Gly Thr Ala Val Pro Ser Thr Gly Ala Ser Lys Ser Lys Arg Gln
                20                    25                    30

Ala Gln Gln Met Val Gln Pro Gln Ser Pro Val Ala Val Ser Gln Ser
                35                    40                    45

Lys Pro Gly Cys Tyr Asp Asn Gly Lys His Tyr Gln Ile Asn Gln Gln
 50                    55                    60

Trp Glu Arg Thr Tyr Leu Gly Asn Ala Leu Val Cys Thr Cys Tyr Gly
65                    70                    75                    80

Gly Ser Arg Gly Phe Asn Cys Glu Ser Lys Pro Glu Ala Glu Glu Thr
                85                    90                    95

Cys Phe Asp Lys Tyr Thr Gly Asn Thr Tyr Arg Val Gly Asp Thr Tyr
                100                    105                    110

Glu Arg Pro Lys Asp Ser Met Ile Trp Asp Cys Thr Cys Ile Gly Ala
                115                    120                    125

Gly Arg Gly Arg Ile Ser Cys Thr Ile Ala Asn Arg Cys His Glu Gly
                130                    135                    140

Gly Gln Ser Tyr Lys Ile Gly Asp Thr Trp Arg Arg Pro His Glu Thr
145                    150                    155                    160

Gly Gly Tyr Met Leu Glu Cys Val Cys Leu Gly Asn Gly Lys Gly Glu
                165                    170                    175

Trp Thr Cys Lys Pro Ile Ala Glu Lys Cys Phe Asp His Ala Ala Gly
                180                    185                    190

Thr Ser Tyr Val Val Gly Glu Thr Trp Glu Lys Pro Tyr Gln Gly Trp
                195                    200                    205

```


-continued

Met Met Val Asp Cys Thr Cys Leu Gly Glu Gly Ser Gly Arg Ile Thr
 210 215 220

Cys Thr Ser Arg Asn Arg Cys Asn Asp Gln Asp Thr Arg Thr Ser Tyr
 225 230 235 240

Arg Ile Gly Asp Thr Trp Ser Lys Lys Asp Asn Arg Gly Asn Leu Leu
 245 250 255

Gln Cys Ile Cys Thr Gly Asn Gly Arg Gly Glu Trp Lys Cys Glu Arg
 260 265 270

His Thr Ser Val Gln Thr Thr Ser Ser Gly Ser Gly Pro Phe Thr Asp
 275 280 285

Val Arg Ala Ala Val Tyr Gln Pro Gln Pro His Pro Gln Pro Pro Pro
 290 295 300

Tyr Gly His Cys Val Thr Asp Ser Gly Val Val Tyr Ser Val Gly Met
 305 310 315 320

Gln Trp Leu Lys Thr Gln Gly Asn Lys Gln Met Leu Cys Thr Cys Leu
 325 330 335

Gly Asn Gly Val Ser Cys Gln Glu Thr Ala Val Thr Gln Thr Tyr Gly
 340 345 350

Gly Asn Ser Asn Gly Glu Pro Cys Val Leu Pro Phe Thr Tyr Asn Gly
 355 360 365

Arg Thr Phe Tyr Ser Cys Thr Thr Glu Gly Arg Gln Asp Gly His Leu
 370 375 380

Trp Cys Ser Thr Thr Ser Asn Tyr Glu Gln Asp Gln Lys Tyr Ser Phe
 385 390 395 400

Cys Thr Asp His Thr Val Leu Val Gln Thr Arg Gly Gly Asn Ser Asn
 405 410 415

Gly Ala Leu Cys His Phe Pro Phe Leu Tyr Asn Asn His Asn Tyr Thr
 420 425 430

Asp Cys Thr Ser Glu Gly Arg Arg Asp Asn Met Lys Trp Cys Gly Thr
 435 440 445

Thr Gln Asn Tyr Asp Ala Asp Gln Lys Phe Gly Phe Cys Pro Met Ala
 450 455 460

Ala His Glu Glu Ile Cys Thr Thr Asn Glu Gly Val Met Tyr Arg Ile
 465 470 475 480

Gly Asp Gln Trp Asp Lys Gln His Asp Met Gly His Met Met Arg Cys
 485 490 495

Thr Cys Val Gly Asn Gly Arg Gly Glu Trp Thr Cys Ile Ala Tyr Ser
 500 505 510

Gln Leu Arg Asp Gln Cys Ile Val Asp Asp Ile Thr Tyr Asn Val Asn
 515 520 525

Asp Thr Phe His Lys Arg His Glu Glu Gly His Met Leu Asn Cys Thr
 530 535 540

Cys Phe Gly Gln Gly Arg Gly Arg Trp Lys Cys Asp Pro Val Asp Gln
 545 550 555 560

Cys Gln Asp Ser Glu Thr Gly Thr Phe Tyr Gln Ile Gly Asp Ser Trp
 565 570 575

Glu Lys Tyr Val His Gly Val Arg Tyr Gln Cys Tyr Cys Tyr Gly Arg
 580 585 590

Gly Ile Gly Glu Trp His Cys Gln Pro Leu Gln Thr Tyr Pro Ser Ser
 595 600 605

-continued

Ser Gly Pro Val Glu Val Phe Ile Thr Glu Thr Pro Ser Gln Pro Asn
 610 615 620

Ser His Pro Ile Gln Trp Asn Ala Pro Gln Pro Ser His Ile Ser Lys
 625 630 635 640

Tyr Ile Leu Arg Trp Arg Pro Lys Asn Ser Val Gly Arg Trp Lys Glu
 645 650 655

Ala Thr Ile Pro Gly His Leu Asn Ser Tyr Thr Ile Lys Gly Leu Lys
 660 665 670

Pro Gly Val Val Tyr Glu Gly Gln Leu Ile Ser Ile Gln Gln Tyr Gly
 675 680 685

His Gln Glu Val Thr Arg Phe Asp Phe Thr Thr Thr Ser Thr Ser Thr
 690 695 700

Pro Val Thr Ser Asn Thr Val Thr Gly Glu Thr Thr Pro Phe Ser Pro
 705 710 715 720

Leu Val Ala Thr Ser Glu Ser Val Thr Glu Ile Thr Ala Ser Ser Phe
 725 730 735

Val Val Ser Trp Val Ser Ala Ser Asp Thr Val Ser Gly Phe Arg Val
 740 745 750

Glu Tyr Glu Leu Ser Glu Glu Gly Asp Glu Pro Gln Tyr Leu Asp Leu
 755 760 765

Pro Ser Thr Ala Thr Ser Val Asn Ile Pro Asp Leu Leu Pro Gly Arg
 770 775 780

Lys Tyr Ile Val Asn Val Tyr Gln Ile Ser Glu Asp Gly Glu Gln Ser
 785 790 795 800

Leu Ile Leu Ser Thr Ser Gln Thr Thr Ala Pro Asp Ala Pro Pro Asp
 805 810 815

Pro Thr Val Asp Gln Val Asp Asp Thr Ser Ile Val Val Arg Trp Ser
 820 825 830

Arg Pro Gln Ala Pro Ile Thr Gly Tyr Arg Ile Val Tyr Ser Pro Ser
 835 840 845

Val Glu Gly Ser Ser Thr Glu Leu Asn Leu Pro Glu Thr Ala Asn Ser
 850 855 860

Val Thr Leu Ser Asp Leu Gln Pro Gly Val Gln Tyr Asn Ile Thr Ile
 865 870 875 880

Tyr Ala Val Glu Glu Asn Gln Glu Ser Thr Pro Val Val Ile Gln Gln
 885 890 895

Glu Thr Thr Gly Thr Pro Arg Ser Asp Thr Val Pro Ser Pro Arg Asp
 900 905 910

Leu Gln Phe Val Glu Val Thr Asp Val Lys Val Thr Ile Met Trp Thr
 915 920 925

Pro Pro Glu Ser Ala Val Thr Gly Tyr Arg Val Asp Val Ile Pro Val
 930 935 940

Asn Leu Pro Gly Glu His Gly Gln Arg Leu Pro Ile Ser Arg Asn Thr
 945 950 955 960

Phe Ala Glu Val Thr Gly Leu Ser Pro Gly Val Thr Tyr Tyr Phe Lys
 965 970 975

Val Phe Ala Val Ser His Gly Arg Glu Ser Lys Pro Leu Thr Ala Gln
 980 985 990

Gln Thr Thr Lys Leu Asp Ala Pro Thr Asn Leu Gln Phe Val Asn Glu
 995 1000 1005

Thr Asp Ser Thr Val Leu Val Arg Trp Thr Pro Pro Arg Ala Gln Ile

-continued

1010				1015				1020							
Thr	Gly	Tyr	Arg	Leu	Thr	Val	Gly	Leu	Thr	Arg	Arg	Gly	Gln	Pro	Arg
1025						1030					1035				1040
Gln	Tyr	Asn	Val	Gly	Pro	Ser	Val	Ser	Lys	Tyr	Pro	Leu	Arg	Asn	Leu
					1045					1050					1055
Gln	Pro	Ala	Ser	Glu	Tyr	Thr	Val	Ser	Leu	Val	Ala	Ile	Lys	Gly	Asn
					1060				1065					1070	
Gln	Glu	Ser	Pro	Lys	Ala	Thr	Gly	Val	Phe	Thr	Thr	Leu	Gln	Pro	Gly
							1080						1085		
Ser	Ser	Ile	Pro	Pro	Tyr	Asn	Thr	Glu	Val	Thr	Glu	Thr	Thr	Ile	Val
						1095					1100				
Ile	Thr	Trp	Thr	Pro	Ala	Pro	Arg	Ile	Gly	Phe	Lys	Leu	Gly	Val	Arg
1105					1110					1115					1120
Pro	Ser	Gln	Gly	Gly	Glu	Ala	Pro	Arg	Glu	Val	Thr	Ser	Asp	Ser	Gly
					1125					1130					1135
Ser	Ile	Val	Val	Ser	Gly	Leu	Thr	Pro	Gly	Val	Glu	Tyr	Val	Tyr	Thr
					1140				1145				1150		
Ile	Gln	Val	Leu	Arg	Asp	Gly	Gln	Glu	Arg	Asp	Ala	Pro	Ile	Val	Asn
							1160						1165		
Lys	Val	Val	Thr	Pro	Leu	Ser	Pro	Pro	Thr	Asn	Leu	His	Leu	Glu	Ala
							1175				1180				
Asn	Pro	Asp	Thr	Gly	Val	Leu	Thr	Val	Ser	Trp	Glu	Arg	Ser	Thr	Thr
1185					1190					1195					1200
Pro	Asp	Ile	Thr	Gly	Tyr	Arg	Ile	Thr	Thr	Thr	Pro	Thr	Asn	Gly	Gln
					1205					1210					1215
Gln	Gly	Asn	Ser	Leu	Glu	Glu	Val	Val	His	Ala	Asp	Gln	Ser	Ser	Cys
					1220				1225						1230
Thr	Phe	Asp	Asn	Leu	Ser	Pro	Gly	Leu	Glu	Tyr	Asn	Val	Ser	Val	Tyr
							1240						1245		
Thr	Val	Lys	Asp	Asp	Lys	Glu	Ser	Val	Pro	Ile	Ser	Asp	Thr	Ile	Ile
						1255					1260				
Pro	Glu	Val	Pro	Gln	Leu	Thr	Asp	Leu	Ser	Phe	Val	Asp	Ile	Thr	Asp
1265					1270					1275					1280
Ser	Ser	Ile	Gly	Leu	Arg	Trp	Thr	Pro	Leu	Asn	Ser	Ser	Thr	Ile	Ile
					1285					1290					1295
Gly	Tyr	Arg	Ile	Thr	Val	Val	Ala	Ala	Gly	Glu	Gly	Ile	Pro	Ile	Phe
					1300				1305				1310		
Glu	Asp	Phe	Val	Asp	Ser	Ser	Val	Gly	Tyr	Tyr	Thr	Val	Thr	Gly	Leu
					1315				1320				1325		
Glu	Pro	Gly	Ile	Asp	Tyr	Asp	Ile	Ser	Val	Ile	Thr	Leu	Ile	Asn	Gly
						1335					1340				
Gly	Glu	Ser	Ala	Pro	Thr	Thr	Leu	Thr	Gln	Gln	Thr	Ala	Val	Pro	Pro
1345					1350					1355					1360
Pro	Thr	Asp	Leu	Arg	Phe	Thr	Asn	Ile	Gly	Pro	Asp	Thr	Met	Arg	Val
					1365					1370					1375
Thr	Trp	Ala	Pro	Pro	Pro	Ser	Ile	Asp	Leu	Thr	Asn	Phe	Leu	Val	Arg
					1380				1385				1390		
Tyr	Ser	Pro	Val	Lys	Asn	Glu	Glu	Asp	Val	Ala	Glu	Leu	Ser	Ile	Ser
					1395				1400				1405		
Pro	Ser	Asp	Asn	Ala	Val	Val	Leu	Thr	Asn	Leu	Leu	Pro	Gly	Thr	Glu
						1415					1420				

-continued

Tyr Val Val Ser Val Ser Ser Val Tyr Glu Gln His Glu Ser Thr Pro
 1425 1430 1435 1440
 Leu Arg Gly Arg Gln Lys Thr Gly Leu Asp Ser Pro Thr Gly Ile Asp
 1445 1450 1455
 Phe Ser Asp Ile Thr Ala Asn Ser Phe Thr Val His Trp Ile Ala Pro
 1460 1465 1470
 Arg Ala Thr Ile Thr Gly Tyr Arg Ile Arg His His Pro Glu His Phe
 1475 1480 1485
 Ser Gly Arg Pro Arg Glu Asp Arg Val Pro His Ser Arg Asn Ser Ile
 1490 1495 1500
 Thr Leu Thr Asn Leu Thr Pro Gly Thr Glu Tyr Val Val Ser Ile Val
 1505 1510 1515
 Ala Leu Asn Gly Arg Glu Glu Ser Pro Leu Leu Ile Gly Gln Gln Ser
 1525 1530 1535
 Thr Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro
 1540 1545 1550
 Thr Ser Leu Leu Ile Ser Trp Asp Ala Pro Ala Val Thr Val Arg Tyr
 1555 1560 1565
 Tyr Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu
 1570 1575 1580
 Phe Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys
 1585 1590 1595
 Pro Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Gly Arg Gly
 1605 1610 1615
 Asp Ser Pro Ala Ser Ser Lys Pro Ile Ser Ile Asn Tyr Arg Thr Glu
 1620 1625 1630
 Ile Asp Lys Pro Ser Gln Met Gln Val Thr Asp Val Gln Asp Asn Ser
 1635 1640 1645
 Ile Ser Val Lys Trp Leu Pro Ser Ser Ser Pro Val Thr Gly Tyr Arg
 1650 1655 1660
 Val Thr Thr Thr Pro Lys Asn Gly Pro Gly Pro Thr Lys Thr Lys Thr
 1665 1670 1675
 Ala Gly Pro Asp Gln Thr Glu Met Thr Ile Glu Gly Leu Gln Pro Thr
 1685 1690 1695
 Val Glu Tyr Val Val Ser Val Tyr Ala Gln Asn Pro Ser Gly Glu Ser
 1700 1705 1710
 Gln Pro Leu Val Gln Thr Ala Val Thr Asn Ile Asp Arg Pro Lys Gly
 1715 1720 1725
 Leu Ala Phe Thr Asp Val Asp Val Asp Ser Ile Lys Ile Ala Trp Glu
 1730 1735 1740
 Ser Pro Gln Gly Gln Val Ser Arg Tyr Arg Val Thr Tyr Ser Ser Pro
 1745 1750 1755
 Glu Asp Gly Ile His Glu Leu Phe Pro Ala Pro Asp Gly Glu Glu Asp
 1765 1770 1775
 Thr Ala Glu Leu Gln Gly Leu Arg Pro Gly Ser Glu Tyr Thr Val Ser
 1780 1785 1790
 Val Val Ala Leu His Asp Asp Met Glu Ser Gln Pro Leu Ile Gly Thr
 1795 1800 1805
 Gln Ser Thr Ala Ile Pro Ala Pro Thr Asp Leu Lys Phe Thr Gln Val
 1810 1815 1820

-continued

Thr Pro Thr Ser Leu Ser Ala Gln Trp Thr Pro Pro Asn Val Gln Leu
 1825 1830 1835 1840
 Thr Gly Tyr Arg Val Arg Val Thr Pro Lys Glu Lys Thr Gly Pro Met
 1845 1850 1855
 Lys Glu Ile Asn Leu Ala Pro Asp Ser Ser Ser Val Val Val Ser Gly
 1860 1865 1870
 Leu Met Val Ala Thr Lys Tyr Glu Val Ser Val Tyr Ala Leu Lys Asp
 1875 1880 1885
 Thr Leu Thr Ser Arg Pro Ala Gln Gly Val Val Thr Thr Leu Glu Asn
 1890 1895 1900
 Val Ser Pro Pro Arg Arg Ala Arg Val Thr Asp Ala Thr Glu Thr Thr
 1905 1910 1915 1920
 Ile Thr Ile Ser Trp Arg Thr Lys Thr Glu Thr Ile Thr Gly Phe Gln
 1925 1930 1935
 Val Asp Ala Val Pro Ala Asn Gly Gln Thr Pro Ile Gln Arg Thr Ile
 1940 1945 1950
 Lys Pro Asp Val Arg Ser Tyr Thr Ile Thr Gly Leu Gln Pro Gly Thr
 1955 1960 1965
 Asp Tyr Lys Ile Tyr Leu Tyr Thr Leu Asn Asp Asn Ala Arg Ser Ser
 1970 1975 1980
 Pro Val Val Ile Asp Ala Ser Thr Ala Ile Asp Ala Pro Ser Asn Leu
 1985 1990 1995 2000
 Arg Phe Leu Ala Thr Thr Pro Asn Ser Leu Leu Val Ser Trp Gln Pro
 2005 2010 2015
 Pro Arg Ala Arg Ile Thr Gly Tyr Ile Ile Lys Tyr Glu Lys Pro Gly
 2020 2025 2030
 Ser Pro Pro Arg Glu Val Val Pro Arg Pro Arg Pro Gly Val Thr Glu
 2035 2040 2045
 Ala Thr Ile Thr Gly Leu Glu Pro Gly Thr Glu Tyr Thr Ile Tyr Val
 2050 2055 2060
 Ile Ala Leu Lys Asn Asn Gln Lys Ser Glu Pro Leu Ile Gly Arg Lys
 2065 2070 2075 2080
 Lys Thr Asp Glu Leu Pro Gln Leu Val Thr Leu Pro His Pro Asn Leu
 2085 2090 2095
 His Gly Pro Glu Ile Leu Asp Val Pro Ser Thr Val Gln Lys Thr Pro
 2100 2105 2110
 Phe Val Thr His Pro Gly Tyr Asp Thr Gly Asn Gly Ile Gln Leu Pro
 2115 2120 2125
 Gly Thr Ser Gly Gln Gln Pro Ser Val Gly Gln Gln Met Ile Phe Glu
 2130 2135 2140
 Glu His Gly Phe Arg Arg Thr Thr Pro Pro Thr Thr Ala Thr Pro Ile
 2145 2150 2155 2160
 Arg His Arg Pro Arg Pro Tyr Pro Pro Asn Val Gly Glu Glu Ile Gln
 2165 2170 2175
 Ile Gly His Ile Pro Arg Glu Asp Val Asp Tyr His Leu Tyr Pro His
 2180 2185 2190
 Gly Pro Gly Leu Asn Pro Asn Ala Ser Thr Gly Gln Glu Ala Leu Ser
 2195 2200 2205
 Gln Thr Thr Ile Ser Trp Ala Pro Phe Gln Asp Thr Ser Glu Tyr Ile
 2210 2215 2220
 Ile Ser Cys His Pro Val Gly Thr Asp Glu Glu Pro Leu Gln Phe Arg

-continued

2225	2230	2235	2240
Val Pro Gly Thr Ser Thr Ser Ala Thr Leu Thr Gly Leu Thr Arg Gly	2245	2250	2255
Ala Thr Tyr Asn Ile Ile Val Glu Ala Leu Lys Asp Gln Gln Arg His	2260	2265	2270
Lys Val Arg Glu Glu Val Val Thr Val Gly Asn Ser Val Asn Glu Gly	2275	2280	2285
Leu Asn Gln Pro Thr Asp Asp Ser Cys Phe Asp Pro Tyr Thr Val Ser	2290	2295	2300
His Tyr Ala Val Gly Asp Glu Trp Glu Arg Met Ser Glu Ser Gly Phe	2305	2310	2315
Lys Leu Leu Cys Gln Cys Leu Gly Phe Gly Ser Gly His Phe Arg Cys	2325	2330	2335
Asp Ser Ser Arg Trp Cys His Asp Asn Gly Val Asn Tyr Lys Ile Gly	2340	2345	2350
Glu Lys Trp Asp Arg Gln Gly Glu Asn Gly Gln Met Met Ser Cys Thr	2355	2360	2365
Cys Leu Gly Asn Gly Lys Gly Glu Phe Lys Cys Asp Pro His Glu Ala	2370	2375	2380
Thr Cys Tyr Asp Asp Gly Lys Thr Tyr His Val Gly Glu Gln Trp Gln	2385	2390	2395
Lys Glu Tyr Leu Gly Ala Ile Cys Ser Cys Thr Cys Phe Gly Gly Gln	2405	2410	2415
Arg Gly Trp Arg Cys Asp Asn Cys Arg Arg Pro Gly Gly Glu Pro Ser	2420	2425	2430
Pro Glu Gly Thr Thr Gly Gln Ser Tyr Asn Gln Tyr Ser Gln Arg Tyr	2435	2440	2445
His Gln Arg Thr Asn Thr Asn Val Asn Cys Pro Ile Glu Cys Phe Met	2450	2455	2460
Pro Leu Asp Val Gln Ala Asp Arg Glu Asp Ser Arg Glu	2465	2470	2475

<210> SEQ ID NO 16
 <211> LENGTH: 8647
 <212> TYPE: DNA
 <213> ORGANISM: Homo Sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Oncofetal Fibronectin variant 2
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank NM 212475
 <309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 16

```

gcccgccgccc gctgtgctgc acagggggag gagaggaac cccaggcgcg agcgggaaga      60
ggggacctgc agccacaact tctctgtgcc tctgcatccc tctgtccct ccaccctcc      120
ccttccccac cctctggccc ccaccttctt ggaggcgaca acccccggga ggcattagaa      180
gggatttttc ccgacaggtg cgaaggaag caaacttggg ggcaacttgc ctcccgggtg      240
gggcgtctct cccccaccgt ctcaacatgc ttaggggtcc gggcccggg ctgctgctgc      300
tggccgtcca gtgcctgggg acagcgggtg cctccacggg agcctcgaag agcaagaggg      360
aggctcagca aatggttcag cccaggtccc cgggtgctgt cagtcaaagc aagcccgtt      420
gttatgacaa tggaaaacac tatcagataa atcaacagtg ggagcggacc tacctaggca      480
    
```

-continued

atgcgttggt ttgtacttgt tatggaggaa gccgaggttt taactgcgag agtaaacctg	540
aagctgaaga gacttgcttt gacaagtaca ctgggaacac ttaccgagtg ggtgacactt	600
atgagcgtcc taaagactcc atgatctggg actgtacctg catcggggct gggcgaggga	660
gaataagctg taccatcgca aaccgctgcc atgaaggggg tcagtcctac aagattggtg	720
acacctggag gagaccacat gagactggtg gttacatggt agagtgtgtg tgtcttggtg	780
atggaaaagg agaatggacc tgcaagccca tagctgagaa gtgttttgat catgctgctg	840
ggacttccta tgtggtcggg gaaacgtggg agaagcccta ccaaggctgg atgatggtg	900
attgtacttg cctgggagaa ggcagcggac gcatcacttg cacttctaga aatagatgca	960
acgatcagga cacaaggaca tcctatagaa ttggagacac ctggagcaag aaggataatc	1020
gaggaaacct gctccagtgc atctgcacag gcaacggccg aggagagtgg aagtgtgaga	1080
ggcacacctc tgtgcagacc acatcgagcg gatctggccc cttcaccgat gttcgtgcag	1140
ctgtttacca accgcagcct cccccacgc ctctcccta tggccactgt gtcacagaca	1200
gtggtgtggt ctactctgtg gggatgcagt ggctgaagac acaaggaaat aagcaaatgc	1260
tttgcacgtg cctgggcaac ggagtcagct gccaaagac agctgtaacc cagacttacg	1320
gtggcaactc aaatggagag ccatgtgtct taccattcac ctacaatggc aggacgttct	1380
actcctgcac cacagaaggg cgacaggacg gacatctttg gtgcagcaca acttcgaatt	1440
atgagcagga ccagaaatac tctttctgca cagaccacac tgttttggtt cagactcgag	1500
gaggaaatc caatggtgcc ttgtgccact tccccttct atacaacaac cacaattaca	1560
ctgattgcac ttctgagggc agaagagaca acatgaagtg gtgtgggacc acacagaact	1620
atgatgccga ccagaagttt gggttctgcc ccatggctgc ccacgaggaa atctgcacaa	1680
ccaatgaag ggtcatgtac cgcattggag atcagtgagg taagcagcat gacatgggtc	1740
acatgatgag gtgcacgtgt gttgggaatg gtcgtgggga atggacatgc attgcctact	1800
cgcagcttcg agatcagtgc attgttgatg acatcactta caatgtgaac gacacattcc	1860
acaagcgtca tgaagagggg cacatgctga actgtacatg cttcggtcag ggtcggggca	1920
ggtggaagtg tgatcccgtc gaccaatgcc aggattcaga gactgggacg ttttatcaaa	1980
ttggagattc atgggagaag tatgtgcatg gtgtcagata ccagtgttac tgctatggcc	2040
gtggcattgg ggagtggcat tgccaacctt tacagaccta tccaagctca agtggctctg	2100
tcgaagtatt tatcactgag actccgagtc agcccaactc ccaccccatc cagtggaatg	2160
caccacagcc atctcacatt tccaagtaca ttctcaggty gagacctaaa aattctgtag	2220
gccgttgga ggaagctacc ataccaggcc acttaaacct ctacaccatc aaaggcctga	2280
agcctggtgt ggtatacagag gccacgtca tcagcatcca gcagtacggc caccaagaag	2340
tgactcgctt tgacttcacc accaccagca ccagcacacc tgtgaccagc aacaccgtga	2400
caggagagac gactcccttt tctcctcttg tggccacttc tgaatctgtg accgaaatca	2460
cagccagtag ctttgtggtc tcctgggtct cagcttccga caccgtgtcg ggattccggg	2520
tggaatatga gctgagttag gagggagatg agccacagta cctggatctt ccaagcacag	2580
ccacttctgt gaacatccct gacctgcttc ctggccgaaa atacattgta aatgtctatc	2640
agatatctga ggatggggag cagagtttga tcctgtctac ttcacaaaca acagcgctc	2700
atgcccctcc tgacccgact gtggaccaag ttgatgacac ctcaattggt gttcgtgga	2760

-continued

gcagacccca	ggctcccac	acagggata	gaatagtcta	ttcgccatca	gtagaaggta	2820
gcagcacaga	actcaacctt	cctgaaactg	caaaactccgt	caccctcagt	gacttgcaac	2880
ctggtgttca	gtataacatc	actatctatg	ctgtggaaga	aaatcaagaa	agtacacctg	2940
ttgtcattca	acaagaacc	actggcacc	cacgctcaga	tacagtgcc	tctcccagg	3000
acctgcagtt	tgtggaagt	acagacgtga	aggtcacat	catgtggaca	ccgctgaga	3060
gtgcagtgac	cggctaccgt	gtggatgtga	tccccgtaa	cctgcctggc	gagcacgggc	3120
agaggctgcc	catcagcag	aacacctttg	cagaagtca	cgggctgtcc	cctgggggtca	3180
cctattactt	caaagtcttt	gcagtgagcc	atgggaggga	gagcaagcct	ctgactgctc	3240
aacagacaac	caactggat	gctcccacta	acctccagtt	tgtcaatgaa	actgattcta	3300
ctgtcctggt	gagatggact	ccacctcggg	cccagataac	aggataccga	ctgacctggg	3360
gccttaccg	aagaggacag	cccaggcagt	acaatgtggg	tccctctgtc	tccaagtacc	3420
cactgaggaa	tctgcagcct	gcatctgagt	acaccgtatc	cctcgtggcc	ataaagggca	3480
accaagagag	ccccaaagcc	actggagtct	ttaccacact	gcagcctggg	agctctattc	3540
caccttacia	caccgaggtg	actgagacca	ccattgtgat	cacatggacg	cctgctccaa	3600
gaattggttt	taagctgggt	gtacgaccaa	gccagggagg	agaggcacca	cgagaagtga	3660
cttcagactc	aggaagcacc	gtgtgtccg	gcttgactcc	aggagtagaa	tacgtctaca	3720
ccatccaagt	cctgagagat	ggacagga	gagatgcgcc	aattgtaaac	aaagtgggtga	3780
caccattgtc	tccaccaaca	aacttgcatc	tggaggcaaa	ccctgacact	ggagtgtctca	3840
cagtctcctg	ggagaggagc	accaccccag	acattactgg	ttatagaatt	accacaaccc	3900
ctacaaacg	ccagcaggga	aattcttttg	aagaagtgg	ccatgctgat	cagagctcct	3960
gcacttttga	taactgag	cccggcctg	agtacaatgt	cagtgtttac	actgtcaagg	4020
atgacaagga	aagtgtccct	atctctgata	ccatcatccc	agaggtgccc	caactcactg	4080
acctaagcct	tgttgatata	accgattcaa	gcatcgccct	gaggtggacc	ccgctaaact	4140
cttcacccat	tattgggtac	cgcacacag	tagttgccc	aggagaagg	atccctattt	4200
ttgaagattt	tgtgactcc	tcagttagg	actacacagt	cacagggctg	gagccgggca	4260
ttgactatga	tatcagcgtt	atcactctca	ttaatggcgg	cgagagtgcc	cctactacac	4320
tgacacaaca	aacggctggt	cctcctccca	ctgacctg	attcaccaac	attgggtccag	4380
acaccatg	gtcacctgg	gctccacccc	catccattga	tttaaccaac	ttcctgggtg	4440
gttactcacc	tgtgaaaaat	gaggaagatg	ttgcagagtt	gtcaatttct	ccttcagaca	4500
atgcagtggt	cttaacaaat	ctcctgcctg	gtacagaata	tgtagttagt	gtctccagt	4560
tctacgaaca	acatgagagc	acacctctta	gaggaagaca	gaaaacaggt	cttgattccc	4620
caactggcat	tgacttttct	gatattactg	ccaactcttt	tactgtgcac	tggattgctc	4680
ctcgagccac	catcactggc	tacaggatcc	gccatcatcc	cgagcacttc	agtgaggagac	4740
ctcgagaaga	tcgggtgcc	cactctcgga	attccatcac	cctcaccaac	ctcactccag	4800
gcacagagta	tgtggtcagc	atcgttgctc	ttaatggcag	agaggaaagt	cccttattga	4860
ttggccaaca	atcaacagtt	tctgatgttc	cgagggacct	ggaagttggt	gctgcgacce	4920
ccaccagcct	actgatcagc	tgggatgctc	ctgctgtcac	agtgagatat	tacaggatca	4980
cttacggaga	gacaggagga	aatagccctg	tccaggagtt	cactgtgcct	gggagcaagt	5040

-continued

ctacagctac catcagcggc cttaaacctg gagttgatta taccatcact gtgtatgctg	5100
tcactggccg tggagacagc cccgcaagca gcaagccaat ttccattaat taccgaacag	5160
aaattgacaa accatcccag atgcaagtga ccgatgttca ggacaacagc attagtgtca	5220
agtggtctgc ttcaagtcc cctgttactg gttacagagt aaccaccact cccaaaaatg	5280
gaccaggacc acaaaaaact aaaactgcag gtccagatca aacagaaatg actattgaag	5340
gcttgacgcc cacagtggag tatgtgggta gtgtctatgc tcagaatcca agcggagaga	5400
gtcagcctct ggttcagact gcagtaacca acattgatcg ccctaaagga ctggcattca	5460
ctgatgtgga tgtcattcc atcaaaatg cttgggaaag cccacagggg caagtttcca	5520
ggtacagggg gacctactcg agccctgagg atggaatcca tgagctattc cctgcacctg	5580
atggtgaaga agacactgca gagctgcaag gcctcagacc gggttctgag tacacagtca	5640
gtgtggttgc cttgcacgat gatatggaga gccagcccct gattggaacc cagtccacag	5700
ctattcctgc accaactgac ctgaagtcca ctcaggtcac acccacaagc ctgagcgccc	5760
agtgacaccc acccaatggt cagctcactg gatatcgagt gcgggtgacc cccaaggaga	5820
agaccggacc aatgaaagaa atcaacctg ctcctgacag ctcacccgtg gttgtatcag	5880
gacttatggt ggccacacaa tatgaagtga gtgtctatgc tcttaaggac actttgacaa	5940
gcagaccagc tcaggagatt gtcaccactc tggagaatgt cagcccacca agaagggtc	6000
gtgtgacaga tgctactgag accaccatca ccattagctg gagaaccaag actgagacga	6060
tcactggctt ccaagttgat gccgttccag ccaatggcca gactccaatc cagagaacca	6120
tcaagccaga tgtcagaagc tacaccatca caggtttaca accaggcact gactacaaga	6180
tctacctgta caccttgaat gacaatgctc ggagctcccc tgtggtcacg gacgcctcca	6240
ctgccattga tgcaccatcc aacctgcgtt tcctggccac cacaccaat tccttgcctg	6300
tatcatggca gccgccactg gccaggatta ccggctacat catcaagtat gagaagcctg	6360
ggtctcctcc cagagaagtg gtcctcggc cccgccctgg tgcacagag gctactatta	6420
ctggcctgga accgggaacc gaatatacaa tttatgtcat tgccctgaag aataatcaga	6480
agagcgagcc cctgattgga aggaaaaaga cagttcaaaa gaccctttc gtcaccacc	6540
ctgggtatga cactggaaat ggtattcagc ttcctggcac ttctggtcag caaccagtg	6600
ttgggaaca aatgatcttt gaggaacatg gttttaggcg gaccacaccg cccacaacgg	6660
ccacccccat aaggcatagc ccaagaccat acccgccgaa ttaggacaa gaagctctct	6720
ctcagacaac catctcatgg gcccattcc aggacacttc tgagtacatc atttcatgtc	6780
atcctgttg cactgatgaa gaacccttac agttcagggt tcctggaact tctaccagtg	6840
ccactctgac aggcctcacc agaggtgcca cctacaacat catagtggag gcaactgaaag	6900
accagcagag gcataaggtt cgggaagagg ttgttaccgt gggcaactct gtcaaccgaag	6960
gcttgaacca acctacggat gactcgtgct ttgaccctca cacagtttcc cattatgccg	7020
ttggagatga gtgggaacga atgtctgaat caggctttaa actgttgtgc cagtgttag	7080
gctttggaag tggctatttc agatgtgatt catctagatg gtgccatgac aatgggtgta	7140
actacaagat tggagagaag tgggaccgtc agggagaaaa tggccagatg atgagctgca	7200
catgtcttgg gaacggaaaa ggagaattca agtgtgaccc tcatgaggca acgtgttatg	7260
atgatgggaa gacataccac gtaggagaac agtggcagaa ggaatatctc ggtgccattt	7320

-continued

```

gctcctgcac atgctttgga ggccagcggg gctggcgctg tgacaactgc cgcagacctg 7380
gggggtaacc cagtcccga ggcactactg gccagtccta caaccagtat tctcagagat 7440
accatcagag aacaaact aatgtaatt gcccaattga gtgcttcacg cctttagatg 7500
tacaggctga cagagaagat tcccagagat aatcatctt tccaatccag aggacaagc 7560
atgtctctct gccaaagatc atctaaactg gagtgatggt agcagaccca gcttagagtt 7620
cttctttctt tcttaagccc ttgctctgg aggaagttct ccagcttcag ctcaactcac 7680
agcttctcca agcatcacc tgaggagttc ctgagggttt tctcataaat gagggctgca 7740
cattgcctgt tctgcttcga agtattcaat accgctcagt attttaaag aagtattct 7800
aagatttggg ttgggatcaa taggaagca tatgcagcca accaagatgc aaatgtttg 7860
aaatgatatg accaaaatt taagtaggaa agtcacccaa acacttctgc tttcacttaa 7920
gtgtctggcc cgcaactag taggaacaag catgatcttg ttactgtgat attttaaata 7980
tccacagtac tactttttc caaatgatcc tagtaattgc ctgaaatat ctttctctta 8040
cctgttattt atcaattttt cccagtattt ttatcggaa aaaattgtat tgaaaacact 8100
tagtatgcag ttgataagag gaatttgta taattatggt ggggtattat tttttatact 8160
gtatgtgcca aagctttact actgtggaaa gacaactggt ttaataaaag atttacattc 8220
cacaactgga agttcatcta ttgatataa gacaccttcg ggggaaataa ttcctgtgaa 8280
tattcttttt caattcagca aacatttgaa aatctatgat gtgcaagtct aattgttgat 8340
ttcagtacaa gattttctaa atcagttgct acaaaaactg attggttttt gtcacttcat 8400
ctcttacta atggagatag ctttacactt tctgctttaa tagatttaag tggaccccaa 8460
tatttattaa aattgctagt ttaccgttca gaagtataat agaaataatc tttagttgct 8520
cttttctaac cattgtaatt ctcccttct tccctccacc tttccttcat tgaataaacc 8580
tctgttcaaa gagattgcct gcaagggaaa taaaatgac taagatatta aaaaaaaaaa 8640
aaaaaaa 8647

```

<210> SEQ ID NO 17

<211> LENGTH: 2421

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank NM 212475

<309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 17

```

Met Leu Arg Gly Pro Gly Pro Gly Leu Leu Leu Leu Ala Val Gln Cys
 1                5                10                15

Leu Gly Thr Ala Val Pro Ser Thr Gly Ala Ser Lys Ser Lys Arg Gln
 20                25                30

Ala Gln Gln Met Val Gln Pro Gln Ser Pro Val Ala Val Ser Gln Ser
 35                40                45

Lys Pro Gly Cys Tyr Asp Asn Gly Lys His Tyr Gln Ile Asn Gln Gln
 50                55                60

Trp Glu Arg Thr Tyr Leu Gly Asn Ala Leu Val Cys Thr Cys Tyr Gly
 65                70                75                80

Gly Ser Arg Gly Phe Asn Cys Glu Ser Lys Pro Glu Ala Glu Glu Thr
 85                90                95

Cys Phe Asp Lys Tyr Thr Gly Asn Thr Tyr Arg Val Gly Asp Thr Tyr

```

-continued

100					105					110					
Glu	Arg	Pro	Lys	Asp	Ser	Met	Ile	Trp	Asp	Cys	Thr	Cys	Ile	Gly	Ala
		115					120					125			
Gly	Arg	Gly	Arg	Ile	Ser	Cys	Thr	Ile	Ala	Asn	Arg	Cys	His	Glu	Gly
		130					135					140			
Gly	Gln	Ser	Tyr	Lys	Ile	Gly	Asp	Thr	Trp	Arg	Arg	Pro	His	Glu	Thr
		145					150					155			160
Gly	Gly	Tyr	Met	Leu	Glu	Cys	Val	Cys	Leu	Gly	Asn	Gly	Lys	Gly	Glu
				165					170					175	
Trp	Thr	Cys	Lys	Pro	Ile	Ala	Glu	Lys	Cys	Phe	Asp	His	Ala	Ala	Gly
			180						185					190	
Thr	Ser	Tyr	Val	Val	Gly	Glu	Thr	Trp	Glu	Lys	Pro	Tyr	Gln	Gly	Trp
		195					200					205			
Met	Met	Val	Asp	Cys	Thr	Cys	Leu	Gly	Glu	Gly	Ser	Gly	Arg	Ile	Thr
		210					215					220			
Cys	Thr	Ser	Arg	Asn	Arg	Cys	Asn	Asp	Gln	Asp	Thr	Arg	Thr	Ser	Tyr
		225					230					235			240
Arg	Ile	Gly	Asp	Thr	Trp	Ser	Lys	Lys	Asp	Asn	Arg	Gly	Asn	Leu	Leu
				245					250					255	
Gln	Cys	Ile	Cys	Thr	Gly	Asn	Gly	Arg	Gly	Glu	Trp	Lys	Cys	Glu	Arg
				260					265					270	
His	Thr	Ser	Val	Gln	Thr	Thr	Ser	Ser	Gly	Ser	Gly	Pro	Phe	Thr	Asp
		275					280					285			
Val	Arg	Ala	Ala	Val	Tyr	Gln	Pro	Gln	Pro	His	Pro	Gln	Pro	Pro	Pro
		290					295					300			
Tyr	Gly	His	Cys	Val	Thr	Asp	Ser	Gly	Val	Val	Tyr	Ser	Val	Gly	Met
		305					310					315			320
Gln	Trp	Leu	Lys	Thr	Gln	Gly	Asn	Lys	Gln	Met	Leu	Cys	Thr	Cys	Leu
				325					330					335	
Gly	Asn	Gly	Val	Ser	Cys	Gln	Glu	Thr	Ala	Val	Thr	Gln	Thr	Tyr	Gly
				340					345					350	
Gly	Asn	Ser	Asn	Gly	Glu	Pro	Cys	Val	Leu	Pro	Phe	Thr	Tyr	Asn	Gly
				355					360					365	
Arg	Thr	Phe	Tyr	Ser	Cys	Thr	Thr	Glu	Gly	Arg	Gln	Asp	Gly	His	Leu
				370			375					380			
Trp	Cys	Ser	Thr	Thr	Ser	Asn	Tyr	Glu	Gln	Asp	Gln	Lys	Tyr	Ser	Phe
				385			390					395			400
Cys	Thr	Asp	His	Thr	Val	Leu	Val	Gln	Thr	Arg	Gly	Gly	Asn	Ser	Asn
				405					410					415	
Gly	Ala	Leu	Cys	His	Phe	Pro	Phe	Leu	Tyr	Asn	Asn	His	Asn	Tyr	Thr
				420					425					430	
Asp	Cys	Thr	Ser	Glu	Gly	Arg	Arg	Asp	Asn	Met	Lys	Trp	Cys	Gly	Thr
				435				440					445		
Thr	Gln	Asn	Tyr	Asp	Ala	Asp	Gln	Lys	Phe	Gly	Phe	Cys	Pro	Met	Ala
				450			455					460			
Ala	His	Glu	Glu	Ile	Cys	Thr	Thr	Asn	Glu	Gly	Val	Met	Tyr	Arg	Ile
				465			470					475			480
Gly	Asp	Gln	Trp	Asp	Lys	Gln	His	Asp	Met	Gly	His	Met	Met	Arg	Cys
				485					490					495	
Thr	Cys	Val	Gly	Asn	Gly	Arg	Gly	Glu	Trp	Thr	Cys	Ile	Ala	Tyr	Ser
				500				505						510	

-continued

Gln Leu Arg Asp Gln Cys Ile Val Asp Asp Ile Thr Tyr Asn Val Asn
 515 520 525
 Asp Thr Phe His Lys Arg His Glu Glu Gly His Met Leu Asn Cys Thr
 530 535 540
 Cys Phe Gly Gln Gly Arg Gly Arg Trp Lys Cys Asp Pro Val Asp Gln
 545 550 555
 Cys Gln Asp Ser Glu Thr Gly Thr Phe Tyr Gln Ile Gly Asp Ser Trp
 565 570 575
 Glu Lys Tyr Val His Gly Val Arg Tyr Gln Cys Tyr Cys Tyr Gly Arg
 580 585 590
 Gly Ile Gly Glu Trp His Cys Gln Pro Leu Gln Thr Tyr Pro Ser Ser
 595 600 605
 Ser Gly Pro Val Glu Val Phe Ile Thr Glu Thr Pro Ser Gln Pro Asn
 610 615 620
 Ser His Pro Ile Gln Trp Asn Ala Pro Gln Pro Ser His Ile Ser Lys
 625 630 635 640
 Tyr Ile Leu Arg Trp Arg Pro Lys Asn Ser Val Gly Arg Trp Lys Glu
 645 650 655
 Ala Thr Ile Pro Gly His Leu Asn Ser Tyr Thr Ile Lys Gly Leu Lys
 660 665 670
 Pro Gly Val Val Tyr Glu Gly Gln Leu Ile Ser Ile Gln Gln Tyr Gly
 675 680 685
 His Gln Glu Val Thr Arg Phe Asp Phe Thr Thr Thr Ser Thr Ser Thr
 690 695 700
 Pro Val Thr Ser Asn Thr Val Thr Gly Glu Thr Thr Pro Phe Ser Pro
 705 710 715 720
 Leu Val Ala Thr Ser Glu Ser Val Thr Glu Ile Thr Ala Ser Ser Phe
 725 730 735
 Val Val Ser Trp Val Ser Ala Ser Asp Thr Val Ser Gly Phe Arg Val
 740 745 750
 Glu Tyr Glu Leu Ser Glu Glu Gly Asp Glu Pro Gln Tyr Leu Asp Leu
 755 760 765
 Pro Ser Thr Ala Thr Ser Val Asn Ile Pro Asp Leu Leu Pro Gly Arg
 770 775 780
 Lys Tyr Ile Val Asn Val Tyr Gln Ile Ser Glu Asp Gly Glu Gln Ser
 785 790 795 800
 Leu Ile Leu Ser Thr Ser Gln Thr Thr Ala Pro Asp Ala Pro Pro Asp
 805 810 815
 Pro Thr Val Asp Gln Val Asp Asp Thr Ser Ile Val Val Arg Trp Ser
 820 825 830
 Arg Pro Gln Ala Pro Ile Thr Gly Tyr Arg Ile Val Tyr Ser Pro Ser
 835 840 845
 Val Glu Gly Ser Ser Thr Glu Leu Asn Leu Pro Glu Thr Ala Asn Ser
 850 855 860
 Val Thr Leu Ser Asp Leu Gln Pro Gly Val Gln Tyr Asn Ile Thr Ile
 865 870 875 880
 Tyr Ala Val Glu Glu Asn Gln Glu Ser Thr Pro Val Val Ile Gln Gln
 885 890 895
 Glu Thr Thr Gly Thr Pro Arg Ser Asp Thr Val Pro Ser Pro Arg Asp
 900 905 910

-continued

1315					1320					1325					
Glu	Pro	Gly	Ile	Asp	Tyr	Asp	Ile	Ser	Val	Ile	Thr	Leu	Ile	Asn	Gly
1330					1335					1340					
Gly	Glu	Ser	Ala	Pro	Thr	Thr	Leu	Thr	Gln	Gln	Thr	Ala	Val	Pro	Pro
1345					1350					1355					1360
Pro	Thr	Asp	Leu	Arg	Phe	Thr	Asn	Ile	Gly	Pro	Asp	Thr	Met	Arg	Val
				1365					1370					1375	
Thr	Trp	Ala	Pro	Pro	Pro	Ser	Ile	Asp	Leu	Thr	Asn	Phe	Leu	Val	Arg
			1380						1385					1390	
Tyr	Ser	Pro	Val	Lys	Asn	Glu	Glu	Asp	Val	Ala	Glu	Leu	Ser	Ile	Ser
		1395					1400						1405		
Pro	Ser	Asp	Asn	Ala	Val	Val	Leu	Thr	Asn	Leu	Leu	Pro	Gly	Thr	Glu
		1410					1415					1420			
Tyr	Val	Val	Ser	Val	Ser	Ser	Val	Tyr	Glu	Gln	His	Glu	Ser	Thr	Pro
1425					1430					1435					1440
Leu	Arg	Gly	Arg	Gln	Lys	Thr	Gly	Leu	Asp	Ser	Pro	Thr	Gly	Ile	Asp
				1445					1450					1455	
Phe	Ser	Asp	Ile	Thr	Ala	Asn	Ser	Phe	Thr	Val	His	Trp	Ile	Ala	Pro
			1460						1465					1470	
Arg	Ala	Thr	Ile	Thr	Gly	Tyr	Arg	Ile	Arg	His	His	Pro	Glu	His	Phe
		1475					1480					1485			
Ser	Gly	Arg	Pro	Arg	Glu	Asp	Arg	Val	Pro	His	Ser	Arg	Asn	Ser	Ile
		1490					1495					1500			
Thr	Leu	Thr	Asn	Leu	Thr	Pro	Gly	Thr	Glu	Tyr	Val	Val	Ser	Ile	Val
1505					1510					1515					1520
Ala	Leu	Asn	Gly	Arg	Glu	Glu	Ser	Pro	Leu	Leu	Ile	Gly	Gln	Gln	Ser
				1525					1530					1535	
Thr	Val	Ser	Asp	Val	Pro	Arg	Asp	Leu	Glu	Val	Val	Ala	Ala	Thr	Pro
			1540						1545					1550	
Thr	Ser	Leu	Leu	Ile	Ser	Trp	Asp	Ala	Pro	Ala	Val	Thr	Val	Arg	Tyr
		1555					1560					1565			
Tyr	Arg	Ile	Thr	Tyr	Gly	Glu	Thr	Gly	Gly	Asn	Ser	Pro	Val	Gln	Glu
		1570					1575					1580			
Phe	Thr	Val	Pro	Gly	Ser	Lys	Ser	Thr	Ala	Thr	Ile	Ser	Gly	Leu	Lys
1585					1590					1595					1600
Pro	Gly	Val	Asp	Tyr	Thr	Ile	Thr	Val	Tyr	Ala	Val	Thr	Gly	Arg	Gly
				1605					1610					1615	
Asp	Ser	Pro	Ala	Ser	Ser	Lys	Pro	Ile	Ser	Ile	Asn	Tyr	Arg	Thr	Glu
			1620						1625					1630	
Ile	Asp	Lys	Pro	Ser	Gln	Met	Gln	Val	Thr	Asp	Val	Gln	Asp	Asn	Ser
		1635					1640					1645			
Ile	Ser	Val	Lys	Trp	Leu	Pro	Ser	Ser	Ser	Pro	Val	Thr	Gly	Tyr	Arg
		1650					1655					1660			
Val	Thr	Thr	Thr	Pro	Lys	Asn	Gly	Pro	Gly	Pro	Thr	Lys	Thr	Lys	Thr
1665					1670					1675					1680
Ala	Gly	Pro	Asp	Gln	Thr	Glu	Met	Thr	Ile	Glu	Gly	Leu	Gln	Pro	Thr
				1685					1690					1695	
Val	Glu	Tyr	Val	Val	Ser	Val	Tyr	Ala	Gln	Asn	Pro	Ser	Gly	Glu	Ser
			1700						1705					1710	
Gln	Pro	Leu	Val	Gln	Thr	Ala	Val	Thr	Asn	Ile	Asp	Arg	Pro	Lys	Gly
		1715					1720							1725	

-continued

Leu Ala Phe Thr Asp Val Asp Val Asp Ser Ile Lys Ile Ala Trp Glu
 1730 1735 1740

Ser Pro Gln Gly Gln Val Ser Arg Tyr Arg Val Thr Tyr Ser Ser Pro
 1745 1750 1755 1760

Glu Asp Gly Ile His Glu Leu Phe Pro Ala Pro Asp Gly Glu Glu Asp
 1765 1770 1775

Thr Ala Glu Leu Gln Gly Leu Arg Pro Gly Ser Glu Tyr Thr Val Ser
 1780 1785 1790

Val Val Ala Leu His Asp Asp Met Glu Ser Gln Pro Leu Ile Gly Thr
 1795 1800 1805

Gln Ser Thr Ala Ile Pro Ala Pro Thr Asp Leu Lys Phe Thr Gln Val
 1810 1815 1820

Thr Pro Thr Ser Leu Ser Ala Gln Trp Thr Pro Pro Asn Val Gln Leu
 1825 1830 1835 1840

Thr Gly Tyr Arg Val Arg Val Thr Pro Lys Glu Lys Thr Gly Pro Met
 1845 1850 1855

Lys Glu Ile Asn Leu Ala Pro Asp Ser Ser Ser Val Val Val Ser Gly
 1860 1865 1870

Leu Met Val Ala Thr Lys Tyr Glu Val Ser Val Tyr Ala Leu Lys Asp
 1875 1880 1885

Thr Leu Thr Ser Arg Pro Ala Gln Gly Val Val Thr Thr Leu Glu Asn
 1890 1895 1900

Val Ser Pro Pro Arg Arg Ala Arg Val Thr Asp Ala Thr Glu Thr Thr
 1905 1910 1915 1920

Ile Thr Ile Ser Trp Arg Thr Lys Thr Glu Thr Ile Thr Gly Phe Gln
 1925 1930 1935

Val Asp Ala Val Pro Ala Asn Gly Gln Thr Pro Ile Gln Arg Thr Ile
 1940 1945 1950

Lys Pro Asp Val Arg Ser Tyr Thr Ile Thr Gly Leu Gln Pro Gly Thr
 1955 1960 1965

Asp Tyr Lys Ile Tyr Leu Tyr Thr Leu Asn Asp Asn Ala Arg Ser Ser
 1970 1975 1980

Pro Val Val Ile Asp Ala Ser Thr Ala Ile Asp Ala Pro Ser Asn Leu
 1985 1990 1995 2000

Arg Phe Leu Ala Thr Thr Pro Asn Ser Leu Leu Val Ser Trp Gln Pro
 2005 2010 2015

Pro Arg Ala Arg Ile Thr Gly Tyr Ile Ile Lys Tyr Glu Lys Pro Gly
 2020 2025 2030

Ser Pro Pro Arg Glu Val Val Pro Arg Pro Arg Pro Gly Val Thr Glu
 2035 2040 2045

Ala Thr Ile Thr Gly Leu Glu Pro Gly Thr Glu Tyr Thr Ile Tyr Val
 2050 2055 2060

Ile Ala Leu Lys Asn Asn Gln Lys Ser Glu Pro Leu Ile Gly Arg Lys
 2065 2070 2075 2080

Lys Thr Val Gln Lys Thr Pro Phe Val Thr His Pro Gly Tyr Asp Thr
 2085 2090 2095

Gly Asn Gly Ile Gln Leu Pro Gly Thr Ser Gly Gln Gln Pro Ser Val
 2100 2105 2110

Gly Gln Gln Met Ile Phe Glu Glu His Gly Phe Arg Arg Thr Thr Pro
 2115 2120 2125

-continued

Pro Thr Thr Ala Thr Pro Ile Arg His Arg Pro Arg Pro Tyr Pro Pro
 2130 2135 2140
 Asn Val Gly Gln Glu Ala Leu Ser Gln Thr Thr Ile Ser Trp Ala Pro
 2145 2150 2155 2160
 Phe Gln Asp Thr Ser Glu Tyr Ile Ile Ser Cys His Pro Val Gly Thr
 2165 2170 2175
 Asp Glu Glu Pro Leu Gln Phe Arg Val Pro Gly Thr Ser Thr Ser Ala
 2180 2185 2190
 Thr Leu Thr Gly Leu Thr Arg Gly Ala Thr Tyr Asn Ile Ile Val Glu
 2195 2200 2205
 Ala Leu Lys Asp Gln Gln Arg His Lys Val Arg Glu Glu Val Val Thr
 2210 2215 2220
 Val Gly Asn Ser Val Asn Glu Gly Leu Asn Gln Pro Thr Asp Asp Ser
 2225 2230 2235 2240
 Cys Phe Asp Pro Tyr Thr Val Ser His Tyr Ala Val Gly Asp Glu Trp
 2245 2250 2255
 Glu Arg Met Ser Glu Ser Gly Phe Lys Leu Leu Cys Gln Cys Leu Gly
 2260 2265 2270
 Phe Gly Ser Gly His Phe Arg Cys Asp Ser Ser Arg Trp Cys His Asp
 2275 2280 2285
 Asn Gly Val Asn Tyr Lys Ile Gly Glu Lys Trp Asp Arg Gln Gly Glu
 2290 2295 2300
 Asn Gly Gln Met Met Ser Cys Thr Cys Leu Gly Asn Gly Lys Gly Glu
 2305 2310 2315 2320
 Phe Lys Cys Asp Pro His Glu Ala Thr Cys Tyr Asp Asp Gly Lys Thr
 2325 2330 2335
 Tyr His Val Gly Glu Gln Trp Gln Lys Glu Tyr Leu Gly Ala Ile Cys
 2340 2345 2350
 Ser Cys Thr Cys Phe Gly Gly Gln Arg Gly Trp Arg Cys Asp Asn Cys
 2355 2360 2365
 Arg Arg Pro Gly Gly Glu Pro Ser Pro Glu Gly Thr Thr Gly Gln Ser
 2370 2375 2380
 Tyr Asn Gln Tyr Ser Gln Arg Tyr His Gln Arg Thr Asn Thr Asn Val
 2385 2390 2395 2400
 Asn Cys Pro Ile Glu Cys Phe Met Pro Leu Asp Val Gln Ala Asp Arg
 2405 2410 2415
 Glu Asp Ser Arg Glu
 2420

<210> SEQ ID NO 18

<211> LENGTH: 8449

<212> TYPE: DNA

<213> ORGANISM: Homo Sapiens

<220> FEATURE:

<223> OTHER INFORMATION: Oncofetal Fibronectin variant 3

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank NM 002026

<309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 18

gccccgcgcg gctgtgctgc acagggggag gagagggaac cccagggcgcg agcgggaaga 60

ggggacctgc agccacaact tctctgtgcc tctgcatccc tctgtccct ccaccctcc 120

ccttccccac cctctggccc ccaccttctt ggaggcgaca acccccggga ggcattagaa 180

-continued

gggatttttc ccgcaggttg cgaaggggaag caaacttggg ggcaacttgc ctcccgggtgc	240
gggogtctct cccccaccgt ctcaacatgc ttaggggtcc ggggcccggg ctgctgctgc	300
tggccgtcca gtgcctgggg acagcgggtgc cctccacggg agcctcgaag agcaagaggc	360
aggctcagca aatggttcag ccccagtccc cgggtgctgt cagtcaaagc aagcccggtt	420
gttatgaaa tgaaaaaac tatcagataa atcaacagtg ggagcggacc tacctaggca	480
atgctgttgt ttgtacttgt tatggaggaa gccgaggttt taactgcgag agtaaacctg	540
aagctgaaga gacttgcctt gacaagtaca ctgggaacac ttaccgagtg ggtgacactt	600
atgagcgtcc taaagactcc atgatctggg actgtacctg catcggggct gggcgaggga	660
gaataagctg taccatcgca aaccgctgcc atgaaggggg tcagtcctac aagattgggtg	720
acacctggag gagaccacat gagactgggt gttacatggt agagtgtgtg tgtcttggtg	780
atggaaaagg agaatggacc tgcaagccca tagctgagaa gtgttttgat catgctgctg	840
ggacttccta tgtgtcggga gaaacgtggg agaagcccta ccaaggctgg atgatggtag	900
attgtacttg cctgggagaa ggcagcggac gcataccttg cacttctaga aatagatgca	960
acgatcagga cacaaggaca tcctatagaa ttggagacac ctggagcaag aaggataatc	1020
gaggaaacct gctccagtgc atctgcacag gcaacggccg aggagagtgg aagtgtgaga	1080
ggcacacctc tgtgcagacc acatcgagcg gatctggccc cttcaccgat gttcgtgcag	1140
ctgtttacca accgcagcct cccccccagc ctccctcccta tggccactgt gtcacagaca	1200
gtggtgttgt ctactctgtg gggatgcagt ggctgaagac acaagaaat aagcaaatgc	1260
ttgcacgtg cctgggcaac ggagtcagct gccaaagagac agctgtaacc cagacttacg	1320
gtggcaactc aaatggagag ccattgtgtc taccattcac ctacaatggc aggacgttct	1380
actctgcac cacagaaggg cgacaggagc gacatctttg gtgcagcaca acttcgaatt	1440
atgagcagga ccagaaatac tctttctgca cagaccacac tgttttggtt cagactcgag	1500
gaggaaattc caatggtgcc ttgtgccact tccccttctc atacaacaac cacaattaca	1560
ctgattgcac ttctgagggc agaagagaca acatgaagtg gtgtgggacc acacagaact	1620
atgatccga ccagaagttt gggttctgcc ccattgctgc ccacgaggaa atctgcacaa	1680
ccaatgaagg ggtcatgtac cgcattggag atcagtgga taagcagcat gacatgggtc	1740
acatgatgag gtgcacgtgt gttgggaatg gtcgtgggga atggacatgc attgcctact	1800
cgcagcttcg agatcagtgc attgttgatg acatcactta caatgtgaac gacacattcc	1860
acaagcgtca tgaagagggg cacatgctga actgtacatg cttcggtcag ggtcggggca	1920
ggtggaagtg tgatcccgtc gaccaatgcc aggattcaga gactgggacg ttttatcaaa	1980
ttggagattc atgggagaag tatgtgcatg gtgtcagata ccagtgtac tgctatggcc	2040
gtggcattgg ggagtggcat tgccaacctt tacagacctc tccaagctca agtggctctg	2100
tcgaagtatt tatcactgag actccagatc agcccaactc ccacccatc cagtggaaatg	2160
caccacagcc atctcacatt tccaagtaca ttctcaggtg gagacctaaa aattctgtag	2220
gccgttgga ggaagctacc ataccaggcc acttaaacctc ctacaccatc aaaggcctga	2280
agcctgggtg ggtatcacgag gccacgtca tcagcatcca gcagtacggc caccaagaag	2340
tgactcgctt tgacttcacc accaccagca ccagcacacc tgtgaccagc aacaccgtga	2400
caggagagac gactcccttt tctcctcttg tggccacttc tgaatctgtg accgaaatca	2460

-continued

cagccagtag	ctttgtggtc	tccctgggtct	cagcttccga	caccgtgtcg	ggattccggg	2520
tggaatatga	gctgagtgag	gagggagatg	agccacagta	cctggatcct	ccaagcacag	2580
ccacttctgt	gaacatccct	gacctgcttc	ctggccgaaa	atacattgta	aatgtctatc	2640
agatatctga	ggatggggag	cagagtttga	tccctgtctac	ttcacaaaaca	acagcgctctg	2700
atgcccctcc	tgaccgcact	gtggaccaag	ttgatgacac	ctcaattggt	gttcgctgga	2760
gcagacccca	ggctcccctc	acagggtaca	gaatagtcta	ttcgccatca	gtagaaggta	2820
gcagcacaga	actcaacctt	cctgaaactg	caaaactccgt	cacctctcagt	gacttgcaac	2880
ctggtgttca	gtataacatc	actatctatg	ctgtggaaga	aaatcaagaa	agtacacctg	2940
ttgtcattca	acaagaaacc	actggcacc	cacgctcaga	tacagtgcct	tctcccaggg	3000
acctgcagtt	tgtggaagt	acagagctga	aggtcacct	catgtggaca	ccgcttgaga	3060
gtgcagtgac	cggtaccctg	gtggatgtga	tccccgtcaa	cctgcctggc	gagcacgggc	3120
agaggctgcc	catcagcagg	aacacctttg	cagaagtcc	cgggctgtcc	cctgggggtca	3180
cctattactt	caaagtcttt	gcagtgagcc	atgggagggg	gagcaagcct	ctgactgctc	3240
aacagacaac	caaactggat	gctcccacta	acctccagtt	tgtcaatgaa	actgattcta	3300
ctgtcctggt	gagatggact	ccacctcggg	cccagataac	aggataccga	ctgacctggg	3360
gccttaccgg	aagaggacag	cccaggcagt	acaatgtggg	tccctctgtc	tccaagtacc	3420
cactgaggaa	tctgcagcct	gcatctgagt	acaccgtatc	cctcgtggcc	ataaagggca	3480
accaagagag	ccccaaagcc	actggagtct	ttaccacact	gcagcctggg	agctctattc	3540
caccttacia	caccgaggtg	actgagacca	ccattgtgat	cacatggacg	cctgctccaa	3600
gaattggttt	taagctgggt	gtacgaccaa	gccagggagg	agaggcacca	cgagaagtga	3660
cttcagactc	aggaagcatc	gttgtgtccg	gcttgactcc	aggagtagaa	tacgtctaca	3720
ccatccaagt	cctgagagat	ggacaggaaa	gagatgcgcc	aattgtaaac	aaagtgggtga	3780
caccattgtc	tccaccaaca	aacttgcatc	tggaggcaaa	ccctgacact	ggagtgtctca	3840
cagtctcctg	ggagaggagc	accaccccag	acattactgg	ttatagaatt	accacaaccc	3900
ctacaaaacg	ccagcaggga	aattcttttg	aagaagtggg	ccatgctgat	cagagctcct	3960
gcacttttga	taacctgagt	cccggcctgg	agtacaatgt	cagtgtttac	actgtcaagg	4020
atgacaagga	aagtgtccct	atctctgata	ccatcatccc	agctgttcc	cctcccactg	4080
acctgcgatt	caccaacatt	ggtccagaca	ccatgcgtgt	cacctgggct	ccaccccat	4140
ccattgattt	aaccaacttc	ctggtgcgtt	actcacctgt	gaaaaatgag	gaagatgttg	4200
cagagttgtc	aatttctcct	tcagacaatg	cagtggctct	aacaaatctc	ctgcctggta	4260
cagaatatgt	agtgagtgtc	tccagtgtct	acgaacaaca	tgagagcaca	cctcttagag	4320
gaagacagaa	aacaggtcct	gattccccaa	ctggcattga	ctttctgat	attactgcca	4380
actcttttac	tgtgactcct	attgtctctc	gagccacat	cactggctac	aggatccgcc	4440
atcatcccg	gcacttcagt	gggagacctc	gagaagatcg	ggtgcccac	tctoggaatt	4500
ccatcaccct	caccaacctc	actccaggca	cagagtatgt	ggtcagcatc	ggtgctctta	4560
atggcagaga	ggaaagtccc	ttattgattg	gccacaatc	aacagtttct	gatgttccga	4620
gggacctgga	agttgtgtct	gcgaccccca	ccagcctact	gatcagctgg	gatgctcctg	4680
ctgtcacagt	gagatattac	aggatcactt	acggagagac	aggaggaat	agccctgtcc	4740

-continued

aggagttcac	tgtgcctggg	agcaagtcta	cagctacat	cagcggcctt	aaacctggag	4800
ttgattatac	catcactgtg	tatgctgtca	ctggccgtgg	agacagcccc	gcaagcagca	4860
agccaatttc	cattaattac	cgaacagaaa	ttgacaaacc	atcccagatg	caagtgaccg	4920
atgttcagga	caacagcatt	agtgtcaagt	ggctgccttc	aagttcccct	gttactgggt	4980
acagagtaac	caccactccc	aaaaatggac	caggaccaac	aaaaactaaa	actgcaggtc	5040
cagatcaaac	agaaatgact	attgaaggct	tgacagccac	agtgaggatg	gtggttagtg	5100
tctatgctca	gaatccaagc	ggagagagtc	agcctctggt	tcagactgca	gtaaccaaca	5160
ttgatcgccc	taaaggactg	gcattcactg	atgtggatgt	cgattccatc	aaaattgctt	5220
gggaaagccc	acaggggcaa	gtttccaggt	acaggggtgac	ctactcgagc	cctgaggatg	5280
gaatccatga	gctattccct	gcacctgatg	gtgaagaaga	cactgcagag	ctgcaaggcc	5340
tcagaccggg	ttctgagtac	acagtcagtg	tggttgcctt	gcacgatgat	atggagagcc	5400
agcccctgat	tggaaccag	tccacagcta	ttcctgcacc	aactgacctg	aagttcactc	5460
aggtcacacc	cacaagcctg	agcggcccag	ggacaccacc	caatgttcag	ctcactggat	5520
atcgagtgcg	ggtgaccccc	aaggagaaga	ccggaccaat	gaaagaaatc	aaccttgctc	5580
ctgacagctc	atccgtggtt	gtatcaggac	ttatgggtgg	caccaaatac	gaagtgagtg	5640
tctatgctct	taaggacact	ttgacaagca	gaccagctca	gggagttgtc	accactctgg	5700
agaatgtcag	cccaccaaga	agggctcgtg	tgacagatgc	tactgagacc	accatcacca	5760
ttagctggag	aaccaagact	gagacgatca	ctggcttcca	agttgatgcc	gttccagcca	5820
atggccagac	tccaatccag	agaaccatca	agccagatgt	cagaagctac	accatcacag	5880
gtttacaacc	aggcactgac	tacaagatct	acctgtacac	cttgaatgac	aatgctcgga	5940
gctcccctgt	ggtcatcgac	gcctccactg	ccattgatgc	accatccaac	ctgcgtttcc	6000
tgccaccacc	acccaattcc	ttgctggtat	catggcagcc	gccacgtgcc	aggattaccg	6060
gctacatcat	caagtatgag	aagcctgggt	ctcctcccag	agaagtggtc	cctcggccccc	6120
gccttggtgt	cacagaggct	actattactg	gcctggaacc	gggaaccgaa	tatacaattt	6180
atgtcattgc	cctgaagaat	aatcagaaga	gogagcccct	gattggaagg	aaaagacag	6240
acgagcttcc	ccaactggta	acccttccac	accccaatct	tcatggacca	gagatcttgg	6300
atgttccttc	cacagttcaa	aagaccccct	tcgtcaccca	ccctgggtat	gacactggaa	6360
atggtattca	gcttcctggc	acttctggtc	agcaaccag	tggtgggcaa	caaatgatct	6420
ttgaggaaca	tggttttagg	cggaccacac	cgcccacaac	ggccaccccc	ataaggcata	6480
ggccaagacc	ataccgcgag	aatgtaggac	aagaagctct	ctctcagaca	accatctcat	6540
gggcccatt	ccaggacact	tctgagtaca	tcatttcatg	tcacctggtt	ggcactgatg	6600
aagaaccctt	acagttcag	gttcctgaa	cttctaccag	tgccactctg	acaggcctca	6660
ccagagggtc	cacctacaac	atcatagtgg	aggcactgaa	agaccagcag	aggcataagg	6720
ttcgggaaga	ggttgttacc	gtgggcaact	ctgtcaacga	aggcttgaac	caacctacgg	6780
atgactcgtg	ctttgacccc	tacacagttt	cccattatgc	cgttggagat	gagtgggaac	6840
gaatgtctga	atcaggcttt	aaactgttgt	gccagtgett	aggctttgga	agtggtcatt	6900
tcagatgtga	ttcatctaga	tggtgccatg	acaatggtgt	gaactacaag	attggagaga	6960
agtgggaccg	tcaggagaaa	aatggccaga	tgatgagctg	cacatgtctt	gggaaccgaa	7020

-continued

```

aaggagaatt caagtgtgac cctcatgagg caacgtgtta tgatgatggg aagacatacc 7080
acgtaggaga acagtggcag aaggaatadc tcggtgccat ttgctcctgc acatgctttg 7140
gaggccagcg gggctggcgc tgtgacaact gccgcagacc tgggggtgaa cccagtcccc 7200
aaggcactac tggccagtcc tacaaccagt attctcagag ataccatcag agaacaaca 7260
ctaatgttaa ttgccaatt gagtgcttca tgcctttaga tgtacaggct gacagagaag 7320
attcccgaga gtaaatcadc tttccaatcc agaggaacaa gcatgtctct ctgccaaagt 7380
ccatcctaac tggagtgatg ttagcagacc cagcttagag ttcttctttc tttcttaagc 7440
cctttgctct ggaggaagt ctccagcttc agctcaactc acagcttctc caagcatcac 7500
cctgggagtt tctgagggt tttctcataa atgagggctg cacattgcct gttctgcttc 7560
gaagtattca ataccgctca gtattttaa tgaagtgatt ctaagatttg gtttgggatc 7620
aataggaaag catatgcagc caaccaagat gcaaatgttt tgaatgata tgacaaaaat 7680
ttaagttagg aaagtacccc aaacacttct gctttcactt aagtgtctgg cccgaatac 7740
tgtaggaaca agcatgatct tgttactgtg atattttaa tatccacagt actcactttt 7800
tccaaatgat ctagtaatt gcctagaaat atctttctct tacctgttat ttatcaattt 7860
ttcccagtat ttttatacgg aaaaaattgt attgaaaaca cttagatgc agttgataag 7920
aggaatttgg tataattatg gtgggtgatt atttttata ctgtatgtgc caaagcttta 7980
ctactgtgga aagacaactg ttttaataaa agatttcat tccacaactt gaagttcatc 8040
tatttgatat aagacacctt cgggggaaat aattcctgtg aatattcttt ttcaattcag 8100
caaacatttg aaaatctatg atgtgcaagt ctaattgttg atttcagtac aagattttct 8160
aaatcagttg ctacaaaaac tgattgggtt ttgtcacttc atctcttcac taatggagat 8220
agctttacac tttctgcttt aatagattta agtggacccc aatatttatt aaaattgcta 8280
gtttaccggt cagaagtata atagaataa tctttagtgt ctcttttcta accattgtaa 8340
ttcttccctt ctccctcca ccttctctc attgaataaa cctctgttca aagagattgc 8400
ctgcaaggga aataaaaatg actaagatat taaaaaaaa aaaaaaaaa 8449

```

<210> SEQ ID NO 19

<211> LENGTH: 2355

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank NM 002026

<309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 19

```

Met Leu Arg Gly Pro Gly Pro Gly Leu Leu Leu Leu Ala Val Gln Cys
 1             5             10             15

Leu Gly Thr Ala Val Pro Ser Thr Gly Ala Ser Lys Ser Lys Arg Gln
      20             25             30

Ala Gln Gln Met Val Gln Pro Gln Ser Pro Val Ala Val Ser Gln Ser
      35             40             45

Lys Pro Gly Cys Tyr Asp Asn Gly Lys His Tyr Gln Ile Asn Gln Gln
      50             55             60

Trp Glu Arg Thr Tyr Leu Gly Asn Ala Leu Val Cys Thr Cys Tyr Gly
      65             70             75             80

Gly Ser Arg Gly Phe Asn Cys Glu Ser Lys Pro Glu Ala Glu Glu Thr
      85             90             95

```

-continued

Cys Phe Asp Lys Tyr Thr Gly Asn Thr Tyr Arg Val Gly Asp Thr Tyr
 100 105 110
 Glu Arg Pro Lys Asp Ser Met Ile Trp Asp Cys Thr Cys Ile Gly Ala
 115 120 125
 Gly Arg Gly Arg Ile Ser Cys Thr Ile Ala Asn Arg Cys His Glu Gly
 130 135 140
 Gly Gln Ser Tyr Lys Ile Gly Asp Thr Trp Arg Arg Pro His Glu Thr
 145 150 155
 Gly Gly Tyr Met Leu Glu Cys Val Cys Leu Gly Asn Gly Lys Gly Glu
 165 170 175
 Trp Thr Cys Lys Pro Ile Ala Glu Lys Cys Phe Asp His Ala Ala Gly
 180 185 190
 Thr Ser Tyr Val Val Gly Glu Thr Trp Glu Lys Pro Tyr Gln Gly Trp
 195 200 205
 Met Met Val Asp Cys Thr Cys Leu Gly Glu Gly Ser Gly Arg Ile Thr
 210 215 220
 Cys Thr Ser Arg Asn Arg Cys Asn Asp Gln Asp Thr Arg Thr Ser Tyr
 225 230 235 240
 Arg Ile Gly Asp Thr Trp Ser Lys Lys Asp Asn Arg Gly Asn Leu Leu
 245 250 255
 Gln Cys Ile Cys Thr Gly Asn Gly Arg Gly Glu Trp Lys Cys Glu Arg
 260 265 270
 His Thr Ser Val Gln Thr Thr Ser Ser Gly Ser Gly Pro Phe Thr Asp
 275 280 285
 Val Arg Ala Ala Val Tyr Gln Pro Gln Pro His Pro Gln Pro Pro Pro
 290 295 300
 Tyr Gly His Cys Val Thr Asp Ser Gly Val Val Tyr Ser Val Gly Met
 305 310 315 320
 Gln Trp Leu Lys Thr Gln Gly Asn Lys Gln Met Leu Cys Thr Cys Leu
 325 330 335
 Gly Asn Gly Val Ser Cys Gln Glu Thr Ala Val Thr Gln Thr Tyr Gly
 340 345 350
 Gly Asn Ser Asn Gly Glu Pro Cys Val Leu Pro Phe Thr Tyr Asn Gly
 355 360 365
 Arg Thr Phe Tyr Ser Cys Thr Thr Glu Gly Arg Gln Asp Gly His Leu
 370 375 380
 Trp Cys Ser Thr Thr Ser Asn Tyr Glu Gln Asp Gln Lys Tyr Ser Phe
 385 390 395 400
 Cys Thr Asp His Thr Val Leu Val Gln Thr Arg Gly Gly Asn Ser Asn
 405 410 415
 Gly Ala Leu Cys His Phe Pro Phe Leu Tyr Asn Asn His Asn Tyr Thr
 420 425 430
 Asp Cys Thr Ser Glu Gly Arg Arg Asp Asn Met Lys Trp Cys Gly Thr
 435 440 445
 Thr Gln Asn Tyr Asp Ala Asp Gln Lys Phe Gly Phe Cys Pro Met Ala
 450 455 460
 Ala His Glu Glu Ile Cys Thr Thr Asn Glu Gly Val Met Tyr Arg Ile
 465 470 475 480
 Gly Asp Gln Trp Asp Lys Gln His Asp Met Gly His Met Met Arg Cys
 485 490 495

-continued

Thr	Cys	Val	Gly	Asn	Gly	Arg	Gly	Glu	Trp	Thr	Cys	Ile	Ala	Tyr	Ser	500	505	510	
Gln	Leu	Arg	Asp	Gln	Cys	Ile	Val	Asp	Asp	Ile	Thr	Tyr	Asn	Val	Asn	515	520	525	
Asp	Thr	Phe	His	Lys	Arg	His	Glu	Glu	Gly	His	Met	Leu	Asn	Cys	Thr	530	535	540	
Cys	Phe	Gly	Gln	Gly	Arg	Gly	Arg	Trp	Lys	Cys	Asp	Pro	Val	Asp	Gln	545	550	555	560
Cys	Gln	Asp	Ser	Glu	Thr	Gly	Thr	Phe	Tyr	Gln	Ile	Gly	Asp	Ser	Trp	565	570	575	
Glu	Lys	Tyr	Val	His	Gly	Val	Arg	Tyr	Gln	Cys	Tyr	Cys	Tyr	Gly	Arg	580	585	590	
Gly	Ile	Gly	Glu	Trp	His	Cys	Gln	Pro	Leu	Gln	Thr	Tyr	Pro	Ser	Ser	595	600	605	
Ser	Gly	Pro	Val	Glu	Val	Phe	Ile	Thr	Glu	Thr	Pro	Ser	Gln	Pro	Asn	610	615	620	
Ser	His	Pro	Ile	Gln	Trp	Asn	Ala	Pro	Gln	Pro	Ser	His	Ile	Ser	Lys	625	630	635	640
Tyr	Ile	Leu	Arg	Trp	Arg	Pro	Lys	Asn	Ser	Val	Gly	Arg	Trp	Lys	Glu	645	650	655	
Ala	Thr	Ile	Pro	Gly	His	Leu	Asn	Ser	Tyr	Thr	Ile	Lys	Gly	Leu	Lys	660	665	670	
Pro	Gly	Val	Val	Tyr	Glu	Gly	Gln	Leu	Ile	Ser	Ile	Gln	Gln	Tyr	Gly	675	680	685	
His	Gln	Glu	Val	Thr	Arg	Phe	Asp	Phe	Thr	Thr	Thr	Ser	Thr	Ser	Thr	690	695	700	
Pro	Val	Thr	Ser	Asn	Thr	Val	Thr	Gly	Glu	Thr	Thr	Pro	Phe	Ser	Pro	705	710	715	720
Leu	Val	Ala	Thr	Ser	Glu	Ser	Val	Thr	Glu	Ile	Thr	Ala	Ser	Ser	Phe	725	730	735	
Val	Val	Ser	Trp	Val	Ser	Ala	Ser	Asp	Thr	Val	Ser	Gly	Phe	Arg	Val	740	745	750	
Glu	Tyr	Glu	Leu	Ser	Glu	Glu	Gly	Asp	Glu	Pro	Gln	Tyr	Leu	Asp	Leu	755	760	765	
Pro	Ser	Thr	Ala	Thr	Ser	Val	Asn	Ile	Pro	Asp	Leu	Leu	Pro	Gly	Arg	770	775	780	
Lys	Tyr	Ile	Val	Asn	Val	Tyr	Gln	Ile	Ser	Glu	Asp	Gly	Glu	Gln	Ser	785	790	795	800
Leu	Ile	Leu	Ser	Thr	Ser	Gln	Thr	Thr	Ala	Pro	Asp	Ala	Pro	Pro	Asp	805	810	815	
Pro	Thr	Val	Asp	Gln	Val	Asp	Asp	Thr	Ser	Ile	Val	Val	Arg	Trp	Ser	820	825	830	
Arg	Pro	Gln	Ala	Pro	Ile	Thr	Gly	Tyr	Arg	Ile	Val	Tyr	Ser	Pro	Ser	835	840	845	
Val	Glu	Gly	Ser	Ser	Thr	Glu	Leu	Asn	Leu	Pro	Glu	Thr	Ala	Asn	Ser	850	855	860	
Val	Thr	Leu	Ser	Asp	Leu	Gln	Pro	Gly	Val	Gln	Tyr	Asn	Ile	Thr	Ile	865	870	875	880
Tyr	Ala	Val	Glu	Glu	Asn	Gln	Glu	Ser	Thr	Pro	Val	Val	Ile	Gln	Gln	885	890	895	
Glu	Thr	Thr	Gly	Thr	Pro	Arg	Ser	Asp	Thr	Val	Pro	Ser	Pro	Arg	Asp				

-continued

900				905				910							
Leu	Gln	Phe	Val	Glu	Val	Thr	Asp	Val	Lys	Val	Thr	Ile	Met	Trp	Thr
	915						920					925			
Pro	Pro	Glu	Ser	Ala	Val	Thr	Gly	Tyr	Arg	Val	Asp	Val	Ile	Pro	Val
	930						935					940			
Asn	Leu	Pro	Gly	Glu	His	Gly	Gln	Arg	Leu	Pro	Ile	Ser	Arg	Asn	Thr
	945				950					955					960
Phe	Ala	Glu	Val	Thr	Gly	Leu	Ser	Pro	Gly	Val	Thr	Tyr	Tyr	Phe	Lys
					965					970					975
Val	Phe	Ala	Val	Ser	His	Gly	Arg	Glu	Ser	Lys	Pro	Leu	Thr	Ala	Gln
			980							985					990
Gln	Thr	Thr	Lys	Leu	Asp	Ala	Pro	Thr	Asn	Leu	Gln	Phe	Val	Asn	Glu
			995						1000					1005	
Thr	Asp	Ser	Thr	Val	Leu	Val	Arg	Trp	Thr	Pro	Pro	Arg	Ala	Gln	Ile
	1010						1015					1020			
Thr	Gly	Tyr	Arg	Leu	Thr	Val	Gly	Leu	Thr	Arg	Arg	Gly	Gln	Pro	Arg
	1025					1030					1035				1040
Gln	Tyr	Asn	Val	Gly	Pro	Ser	Val	Ser	Lys	Tyr	Pro	Leu	Arg	Asn	Leu
					1045					1050					1055
Gln	Pro	Ala	Ser	Glu	Tyr	Thr	Val	Ser	Leu	Val	Ala	Ile	Lys	Gly	Asn
			1060							1065				1070	
Gln	Glu	Ser	Pro	Lys	Ala	Thr	Gly	Val	Phe	Thr	Thr	Leu	Gln	Pro	Gly
			1075				1080							1085	
Ser	Ser	Ile	Pro	Pro	Tyr	Asn	Thr	Glu	Val	Thr	Glu	Thr	Thr	Ile	Val
		1090				1095						1100			
Ile	Thr	Trp	Thr	Pro	Ala	Pro	Arg	Ile	Gly	Phe	Lys	Leu	Gly	Val	Arg
	1105					1110					1115				1120
Pro	Ser	Gln	Gly	Gly	Glu	Ala	Pro	Arg	Glu	Val	Thr	Ser	Asp	Ser	Gly
					1125						1130				1135
Ser	Ile	Val	Val	Ser	Gly	Leu	Thr	Pro	Gly	Val	Glu	Tyr	Val	Tyr	Thr
			1140								1145				1150
Ile	Gln	Val	Leu	Arg	Asp	Gly	Gln	Glu	Arg	Asp	Ala	Pro	Ile	Val	Asn
		1155					1160							1165	
Lys	Val	Val	Thr	Pro	Leu	Ser	Pro	Pro	Thr	Asn	Leu	His	Leu	Glu	Ala
		1170					1175							1180	
Asn	Pro	Asp	Thr	Gly	Val	Leu	Thr	Val	Ser	Trp	Glu	Arg	Ser	Thr	Thr
			1185			1190					1195				1200
Pro	Asp	Ile	Thr	Gly	Tyr	Arg	Ile	Thr	Thr	Thr	Pro	Thr	Asn	Gly	Gln
					1205						1210				1215
Gln	Gly	Asn	Ser	Leu	Glu	Glu	Val	Val	His	Ala	Asp	Gln	Ser	Ser	Cys
			1220							1225				1230	
Thr	Phe	Asp	Asn	Leu	Ser	Pro	Gly	Leu	Glu	Tyr	Asn	Val	Ser	Val	Tyr
		1235					1240							1245	
Thr	Val	Lys	Asp	Asp	Lys	Glu	Ser	Val	Pro	Ile	Ser	Asp	Thr	Ile	Ile
		1250				1255						1260			
Pro	Ala	Val	Pro	Pro	Pro	Thr	Asp	Leu	Arg	Phe	Thr	Asn	Ile	Gly	Pro
			1265			1270					1275				1280
Asp	Thr	Met	Arg	Val	Thr	Trp	Ala	Pro	Pro	Pro	Ser	Ile	Asp	Leu	Thr
					1285						1290				1295
Asn	Phe	Leu	Val	Arg	Tyr	Ser	Pro	Val	Lys	Asn	Glu	Glu	Asp	Val	Ala
			1300							1305				1310	

-continued

Glu Leu Ser Ile Ser Pro Ser Asp Asn Ala Val Val Leu Thr Asn Leu
 1315 1320 1325
 Leu Pro Gly Thr Glu Tyr Val Val Ser Val Ser Val Tyr Glu Gln
 1330 1335 1340
 His Glu Ser Thr Pro Leu Arg Gly Arg Gln Lys Thr Gly Leu Asp Ser
 1345 1350 1355 1360
 Pro Thr Gly Ile Asp Phe Ser Asp Ile Thr Ala Asn Ser Phe Thr Val
 1365 1370 1375
 His Trp Ile Ala Pro Arg Ala Thr Ile Thr Gly Tyr Arg Ile Arg His
 1380 1385 1390
 His Pro Glu His Phe Ser Gly Arg Pro Arg Glu Asp Arg Val Pro His
 1395 1400 1405
 Ser Arg Asn Ser Ile Thr Leu Thr Asn Leu Thr Pro Gly Thr Glu Tyr
 1410 1415 1420
 Val Val Ser Ile Val Ala Leu Asn Gly Arg Glu Glu Ser Pro Leu Leu
 1425 1430 1435 1440
 Ile Gly Gln Gln Ser Thr Val Ser Asp Val Pro Arg Asp Leu Glu Val
 1445 1450 1455
 Val Ala Ala Thr Pro Thr Ser Leu Leu Ile Ser Trp Asp Ala Pro Ala
 1460 1465 1470
 Val Thr Val Arg Tyr Tyr Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn
 1475 1480 1485
 Ser Pro Val Gln Glu Phe Thr Val Pro Gly Ser Lys Ser Thr Ala Thr
 1490 1495 1500
 Ile Ser Gly Leu Lys Pro Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala
 1505 1510 1515 1520
 Val Thr Gly Arg Gly Asp Ser Pro Ala Ser Ser Lys Pro Ile Ser Ile
 1525 1530 1535
 Asn Tyr Arg Thr Glu Ile Asp Lys Pro Ser Gln Met Gln Val Thr Asp
 1540 1545 1550
 Val Gln Asp Asn Ser Ile Ser Val Lys Trp Leu Pro Ser Ser Ser Pro
 1555 1560 1565
 Val Thr Gly Tyr Arg Val Thr Thr Pro Lys Asn Gly Pro Gly Pro
 1570 1575 1580
 Thr Lys Thr Lys Thr Ala Gly Pro Asp Gln Thr Glu Met Thr Ile Glu
 1585 1590 1595 1600
 Gly Leu Gln Pro Thr Val Glu Tyr Val Val Ser Val Tyr Ala Gln Asn
 1605 1610 1615
 Pro Ser Gly Glu Ser Gln Pro Leu Val Gln Thr Ala Val Thr Asn Ile
 1620 1625 1630
 Asp Arg Pro Lys Gly Leu Ala Phe Thr Asp Val Asp Val Asp Ser Ile
 1635 1640 1645
 Lys Ile Ala Trp Glu Ser Pro Gln Gly Gln Val Ser Arg Tyr Arg Val
 1650 1655 1660
 Thr Tyr Ser Ser Pro Glu Asp Gly Ile His Glu Leu Phe Pro Ala Pro
 1665 1670 1675 1680
 Asp Gly Glu Glu Asp Thr Ala Glu Leu Gln Gly Leu Arg Pro Gly Ser
 1685 1690 1695
 Glu Tyr Thr Val Ser Val Val Ala Leu His Asp Asp Met Glu Ser Gln
 1700 1705 1710

-continued

Pro Leu Ile Gly Thr Gln Ser Thr Ala Ile Pro Ala Pro Thr Asp Leu
 1715 1720 1725
 Lys Phe Thr Gln Val Thr Pro Thr Ser Leu Ser Ala Gln Trp Thr Pro
 1730 1735 1740
 Pro Asn Val Gln Leu Thr Gly Tyr Arg Val Arg Val Thr Pro Lys Glu
 1745 1750 1755 1760
 Lys Thr Gly Pro Met Lys Glu Ile Asn Leu Ala Pro Asp Ser Ser Ser
 1765 1770 1775
 Val Val Val Ser Gly Leu Met Val Ala Thr Lys Tyr Glu Val Ser Val
 1780 1785 1790
 Tyr Ala Leu Lys Asp Thr Leu Thr Ser Arg Pro Ala Gln Gly Val Val
 1795 1800 1805
 Thr Thr Leu Glu Asn Val Ser Pro Pro Arg Arg Ala Arg Val Thr Asp
 1810 1815 1820
 Ala Thr Glu Thr Thr Ile Thr Ile Ser Trp Arg Thr Lys Thr Glu Thr
 1825 1830 1835 1840
 Ile Thr Gly Phe Gln Val Asp Ala Val Pro Ala Asn Gly Gln Thr Pro
 1845 1850 1855
 Ile Gln Arg Thr Ile Lys Pro Asp Val Arg Ser Tyr Thr Ile Thr Gly
 1860 1865 1870
 Leu Gln Pro Gly Thr Asp Tyr Lys Ile Tyr Leu Tyr Thr Leu Asn Asp
 1875 1880 1885
 Asn Ala Arg Ser Ser Pro Val Val Ile Asp Ala Ser Thr Ala Ile Asp
 1890 1895 1900
 Ala Pro Ser Asn Leu Arg Phe Leu Ala Thr Thr Pro Asn Ser Leu Leu
 1905 1910 1915 1920
 Val Ser Trp Gln Pro Pro Arg Ala Arg Ile Thr Gly Tyr Ile Ile Lys
 1925 1930 1935
 Tyr Glu Lys Pro Gly Ser Pro Pro Arg Glu Val Val Pro Arg Pro Arg
 1940 1945 1950
 Pro Gly Val Thr Glu Ala Thr Ile Thr Gly Leu Glu Pro Gly Thr Glu
 1955 1960 1965
 Tyr Thr Ile Tyr Val Ile Ala Leu Lys Asn Asn Gln Lys Ser Glu Pro
 1970 1975 1980
 Leu Ile Gly Arg Lys Lys Thr Asp Glu Leu Pro Gln Leu Val Thr Leu
 1985 1990 1995 2000
 Pro His Pro Asn Leu His Gly Pro Glu Ile Leu Asp Val Pro Ser Thr
 2005 2010 2015
 Val Gln Lys Thr Pro Phe Val Thr His Pro Gly Tyr Asp Thr Gly Asn
 2020 2025 2030
 Gly Ile Gln Leu Pro Gly Thr Ser Gly Gln Gln Pro Ser Val Gly Gln
 2035 2040 2045
 Gln Met Ile Phe Glu Glu His Gly Phe Arg Arg Thr Thr Pro Pro Thr
 2050 2055 2060
 Thr Ala Thr Pro Ile Arg His Arg Pro Arg Pro Tyr Pro Pro Asn Val
 2065 2070 2075 2080
 Gly Gln Glu Ala Leu Ser Gln Thr Thr Ile Ser Trp Ala Pro Phe Gln
 2085 2090 2095
 Asp Thr Ser Glu Tyr Ile Ile Ser Cys His Pro Val Gly Thr Asp Glu
 2100 2105 2110
 Glu Pro Leu Gln Phe Arg Val Pro Gly Thr Ser Thr Ser Ala Thr Leu

-continued

atgcgttggt ttgtacttgt tatggaggaa gccgaggttt taactgagag agtaaacctg	540
aagctgaaga gacttgcttt gacaagtaca ctgggaacac ttaccgagtg ggtgacactt	600
atgagcgtcc taaagactcc atgatctggg actgtacctg catcggggct gggcgaggga	660
gaataagctg taccatcgca aaccgctgcc atgaaggggg tcagtcctac aagattggtg	720
acacctggag gagaccacat gagactggtg gttacatggt agagtgtgtg tgtcttggtg	780
atggaaaagg agaattggacc tgcaagccca tagctgagaa gtgttttgat catgctgctg	840
ggacttccta tgtggtcggg gaaacgtggg agaagcccta ccaaggctgg atgatggtag	900
attgtacttg cctgggagaa ggcagcggac gcatcacttg cacttctaga aatagatgca	960
acgatcagga cacaaggaca tcctatagaa ttggagacac ctggagcaag aaggataatc	1020
gaggaaacct gctccagtgc atctgcacag gcaacggccg aggagagtgg aagtgtgaga	1080
ggcacacctc tgtgcagacc acatcgagcg gatctggccc cttcaccgat gttcgtgcag	1140
ctgtttacca accgcagcct cccccacgc ctctcccta tggccactgt gtcacagaca	1200
gtggtgtggt ctactctgtg gggatgcagt ggctgaagac acaaggaaat aagcaaagtc	1260
tttgcacgtg cctgggcaac ggagtcagct gccaaagac agctgtaacc cagacttacg	1320
gtggcaactc aaatggagag ccatgtgtct taccattcac ctacaatggc aggacgttct	1380
actcctgcac cacagaaggg cgacaggacg gacatctttg gtgcagcaca acttcgaatt	1440
atgagcagga ccagaaatac tctttctgca cagaccacac tgttttggtt cagactcgag	1500
gaggaaattc caatggtgcc ttgtgccact tccccttct atacaacaac cacaattaca	1560
ctgattgcac ttctgagggc agaagagaca acatgaagtg gtgtgggacc acacagaact	1620
atgatgccga ccagaagttt gggttctgcc ccatggctgc ccacgaggaa atctgcacaa	1680
ccaatgaag ggtcatgtac cgcattggag atcagtgagg taagcagcat gacatgggtc	1740
acatgatgag gtgcacgtgt gttgggaatg gtcgtgggga atggacatgc attgcctact	1800
cgcagcttcg agatcagtgc attgttgatg acatcactta caatgtgaac gacacattcc	1860
acaagcgtca tgaagagggg cacatgctga actgtacatg cttcggtcag ggtcggggca	1920
ggtggaagtg tgatcccgtc gaccaatgcc aggattcaga gactgggacg ttttatcaaa	1980
ttggagattc atgggagaag tatgtgcatg gtgtcagata ccagtgtac tgctatggcc	2040
gtggcattgg ggagtggcat tgccaacctt tacagaccta tccaagctca agtggctctg	2100
tcgaagtatt tatcactgag actccgagtc agcccaactc ccacccatc cagtggaatg	2160
caccacagcc atctcacatt tccaagtaca ttctcaggty gagacctaaa aattctgtag	2220
gccgttgga ggaagctacc ataccaggcc acttaaacct ctacaccatc aaaggcctga	2280
agcctgggtg ggtatacagag gccacgtca tcagcatcca gcagtacggc caccaagaag	2340
tgactcgctt tgacttcacc accaccagca ccagcacacc tgtgaccagc aacaccgtga	2400
caggagagac gactcccttt tctcctcttg tggccacttc tgaatctgtg accgaaatca	2460
cagccagtag ctttgtggtc tcctgggtct cagcttccga caccgtgtcg ggattccggg	2520
tggaatatga gctgagtgag gagggagatg agccacagta cctggatcctt ccaagcacag	2580
ccacttctgt gaacatccct gacctgcttc ctggccgaaa atacattgta aatgtctatc	2640
agatatctga ggatggggag cagagtttga tcctgtctac ttcacaaaca acagcgctctg	2700
atgcccctcc tgacccgact gtggaccaag ttgatgacac ctcaattggt gttcgtgga	2760

-continued

gcagacccca	ggctcccac	acagggata	gaatagtcta	ttcgccatca	gtagaaggta	2820
gcagcacaga	actcaacctt	cctgaaactg	caaaactccgt	caccctcagt	gacttgcaac	2880
ctggtgttca	gtataacatc	actatctatg	ctgtggaaga	aaatcaagaa	agtacacctg	2940
ttgtcattca	acaagaacc	actggcacc	cacgctcaga	tacagtgcc	tctcccagg	3000
acctgcagtt	tgtggaagt	acagacgtga	aggtcacat	catgtggaca	ccgctgaga	3060
gtgcagtgac	cggctaccgt	gtggatgtga	tccccgtcaa	cctgcctggc	gagcacgggc	3120
agaggtgcc	catcagcag	aacacctttg	cagaagtca	cgggctgtcc	cctgggggtca	3180
cctattactt	caaagtcttt	gcagtgagcc	atgggaggga	gagcaagcct	ctgactgctc	3240
aacagacaac	caactggat	gctcccacta	acctccagtt	tgcaatgaa	actgattcta	3300
ctgtcctggt	gagatggact	ccacctcggg	cccagataac	aggataccga	ctgacctggg	3360
gccttaccg	aagaggacag	cccaggcagt	acaatgtggg	tccctctgtc	tccaagtacc	3420
cactgaggaa	tctgcagcct	gcatctgagt	acaccgtatc	cctcgtggcc	ataaagggca	3480
accaagagag	ccccaaagcc	actggagtct	ttaccacact	gcagcctggg	agctctattc	3540
caccttacia	caccgaggtg	actgagacca	ccattgtgat	cacatggacg	cctgctccaa	3600
gaattggttt	taagctgggt	gtacgaccaa	gccagggagg	agaggcacca	cgagaagtga	3660
cttcagactc	aggaagcacc	gtgtgtccg	gcttgactcc	aggagtagaa	tacgtctaca	3720
ccatccaagt	cctgagagat	ggacagga	gagatgcgcc	aattgtaaac	aaagtgggta	3780
caccattgtc	tccaccaaca	aacttgcatc	tggaggcaaa	ccctgacact	ggagtgtctca	3840
cagtctcctg	ggagaggagc	accaccccag	acattactgg	ttatagaatt	accacaaccc	3900
ctacaaacgg	ccagcaggga	aattcttttg	aagaagtggg	ccatgctgat	cagagctcct	3960
gcacttttga	taacctgagt	cccggcctgg	agtacaatgt	cagtgtttac	actgtcaagg	4020
atgacaagga	aagtgtccct	atctctgata	ccatcatccc	agctgttccct	cctcccactg	4080
acctgcgatt	caccaacatt	ggtccagaca	ccatgcgtgt	cacctgggct	ccaccccat	4140
ccattgattt	aaccaacttc	ctggtgcgtt	actcacctgt	gaaaaatgag	gaagatggtg	4200
cagagttgtc	aatttctcct	tcagacaatg	cagtggctct	aacaaatctc	ctgcttggtg	4260
cagaatagt	agtgagtgtc	tccagtgtct	acgaacaaca	tgagagcaca	cctcttagag	4320
gaagacagaa	aacaggtcct	gattccocaa	ctggcattga	cttttctgat	attactgcca	4380
actcttttac	tgtgactg	attgtctctc	gagccaccat	cactggctac	aggatccgcc	4440
atcatcccga	gcacttcagt	gggagacctc	gagaagatcg	ggtgcccac	tctcggaatt	4500
ccatcaccct	caccaacctc	actccaggca	cagagtatgt	ggtcagcatc	gttgctctta	4560
atggcagaga	ggaaagtccc	ttattgattg	gccaacaatc	aacagtttct	gatgttccga	4620
gggacctgga	agttgttgct	gcgaccccca	ccagcctact	gatcagctgg	gatgctcctg	4680
ctgtcacagt	gagatattac	aggatcactt	acggagagac	aggagaaat	agccctgtcc	4740
aggagttcac	tgtgcctggg	agcaagtcta	cagctaccat	cagcggcctt	aaacctggag	4800
ttgattatac	catcactgtg	tatgtgttca	ctggccgtgg	agacagcccc	gcaagcagca	4860
agccaatttc	cattaattac	cgaacagaaa	ttgacaaacc	atcccagatg	caagtgaccg	4920
atgttcagga	caacagcatt	agtgtaagt	ggctgccttc	aagttccctt	gttactggtt	4980
acagagtaac	caccactccc	aaaaatggac	caggaccaac	aaaaactaaa	actgcaggtc	5040

-continued

cagatcaaac	agaaatgact	attgaaggct	tcagcccac	agtggagtat	gtggttagtg	5100
tctatgctca	gaatccaagc	ggagagagtc	agcctctggt	tcagactgca	gtaaccaaca	5160
ttgatcgccc	taaaggactg	gcattcactg	atgtggatgt	cgattccatc	aaaattgctt	5220
gggaaagccc	acaggggcaa	gtttccaggt	acagggtgac	ctactcgagc	cctgaggatg	5280
gaatccatga	gctattccct	gcacctgatg	gtgaagaaga	cactgcagag	ctgcaaggcc	5340
tcagaccggg	ttctgagtac	acagtcagtg	tggttgcctt	gcacgatgat	atggagagcc	5400
agcccctgat	tggaaccag	tccacagcta	ttcctgcacc	aactgacctg	aagtccactc	5460
aggtcacacc	cacaagcctg	agcgcccagt	ggacaccacc	caatgttcag	ctcactggat	5520
atcgagtgcg	ggtgaccccc	aaggagaaga	ccggaccaat	gaaagaaatc	aaccttgctc	5580
ctgacagctc	atccgtggtt	gtatcaggac	ttatggtggc	caccaaatac	gaagtgagtg	5640
tctatgctct	taaggacact	ttgacaagca	gaccagctca	gggagttgtc	accactctgg	5700
agaatgtcag	cccaccaaga	agggctcgtg	tgacagatgc	tactgagacc	accatcacca	5760
ttagctggag	aaccaagact	gagacgatca	ctggcttcca	agttgatgcc	gttccagcca	5820
atggccagac	tccaatccag	agaaccatca	agccagatgt	cagaagctac	accatcacag	5880
gtttacaacc	aggcactgac	tacaagatct	acctgtacac	cttgaatgac	aatgctcgga	5940
gctcccctgt	ggtcatcgac	gcctccactg	ccattgatgc	accatccaac	ctgctgttcc	6000
tggccaccac	acccaattcc	ttgctgggat	catggcagcc	gccacgtgcc	aggattaccg	6060
gtacatcat	caagtatgag	aagcctgggt	ctcctcccag	agaagtggtc	cctcgccccc	6120
gcccctggtg	cacagaggct	actattactg	gcctggaacc	gggaaccgaa	tatacaattt	6180
atgtcattgc	cctgaagaat	aatcagaaga	gcgagcccct	gattggaagg	aaaagacag	6240
ttcaaaagac	ccctttctgc	acccaccctg	ggtatgacac	tggaaatggt	attcagcttc	6300
ctggcacttc	tggtcagcaa	cccagtggtg	ggcaacaaat	gatctttgag	gaacatggtt	6360
ttaggcggac	cacaccgccc	acaacggcca	cccccataag	gcataggcca	agaccatacc	6420
cgccgaatgt	aggacaagaa	gctctctctc	agacaaccat	ctcatgggcc	ccattccagg	6480
acaacttctg	gtacatcatt	tcatgtcatc	ctggtggcac	tgatgaagaa	cccttacagt	6540
tcagggttcc	tggaacttct	accagtgcca	ctctgacagg	cctcaccaga	ggtgccacct	6600
acaacatcat	agtggaggca	ctgaaagacc	agcagaggca	taaggttcgg	gaagaggttg	6660
ttaccgtggg	caactctgtc	aacgaaggct	tgaaccaacc	tacggatgac	tcgtgctttg	6720
acccctacac	agtttcccat	tatgcccgtg	gagatgagtg	ggaacgaatg	tctgaatcag	6780
gctttaaact	gttgtgccag	tgcttaggct	ttggaagtgg	tcatttcaga	tgtgattcat	6840
ctagatgggt	ccatgacaat	ggtgtgaact	acaagattgg	agagaagtgg	gacogtcagg	6900
gagaaaatgg	ccagatgatg	agctgcacat	gtcttgggaa	cggaaaagga	gaattcaagt	6960
gtgaccctca	tgaggcaacg	tgttatgatg	atgggaagac	ataccacgta	ggagaacagt	7020
ggcagaagga	atatctcgtg	gccatttgct	cctgcacatg	ctttggaggc	cagcggggct	7080
ggcgtctgta	caactgccgc	agacctgggg	gtgaaccag	tcccgaaggc	actactggcc	7140
agtctacaa	ccagtattct	cagagatacc	atcagagaac	aaacactaat	gttaattgcc	7200
caattgagtg	cttcatgcct	ttagatgtac	aggctgacag	agaagattcc	cgagagtaaa	7260
tcactcttcc	aatccagag	aacaagcatg	tctctctgcc	aagatccatc	taaactggag	7320

-continued

```

tgatgtagc agaccagct tagagttctt ctttctttct taagcccttt gctctggagg 7380
aagttctcca gttcagctc aactcacagc ttctccaagc atcaccctgg gagtttctctg 7440
agggttttct cataaatgag ggctgcacat tgcctgttct gcttcgaagt attcaatacc 7500
gctcagtatt ttaaataga tgattctaag atttggtttg ggatcaatag gaaagcatat 7560
gcagccaacc aagatgcaaa tgttttgaaa tgatatgacc aaaattttaa gtaggaaagt 7620
caccCAAACA cttctgcttt cacttaagtg tctggcccgc aatactgtag gaacaagcat 7680
gatcttggtta ctgtgatatt ttaaataatcc acagtactca ctttttccaa atgatcctag 7740
taattgccta gaaatatctt tctcttacct gttatttata aatttttccc agtattttta 7800
tacggaaaaa attgtattga aaacacttag tatgcagttg ataagaggaa tttggtataa 7860
ttatggtggg tgattatttt ttatactgta tgtgccaaag ctttactact gtggaaagac 7920
aactgtttta ataaaagatt tacattccac aactggaagt tcatctattt gatataagac 7980
accttcgggg gaaataatc ctgtgaatat tctttttcaa ttcagcaaac attgaaaat 8040
ctatgatgtg caagtcta atgttgatttc agtacaagat tttctaaatc agttgctaca 8100
aaaactgatt ggtttttgtc acttcatctc ttcactaatg gagatagctt tacactttct 8160
gctttaatag atttaagtgg accccaatat ttattaaaat tgctagttaa ccgttcagaa 8220
gtataataga aataatcttt agttgctctt ttctaaccat tgtaattctt cccttcttcc 8280
ctccaccttt cttcattga ataaacctct gttcaaagag attgcctgca agggaaataa 8340
aaatgactaa gatattaaaa aaaaaaaaaa aaaa 8374

```

<210> SEQ ID NO 21

<211> LENGTH: 2330

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank NM 212478

<309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 21

```

Met Leu Arg Gly Pro Gly Pro Gly Leu Leu Leu Leu Ala Val Gln Cys
 1             5             10             15

Leu Gly Thr Ala Val Pro Ser Thr Gly Ala Ser Lys Ser Lys Arg Gln
      20             25             30

Ala Gln Gln Met Val Gln Pro Gln Ser Pro Val Ala Val Ser Gln Ser
      35             40             45

Lys Pro Gly Cys Tyr Asp Asn Gly Lys His Tyr Gln Ile Asn Gln Gln
 50             55             60

Trp Glu Arg Thr Tyr Leu Gly Asn Ala Leu Val Cys Thr Cys Tyr Gly
65             70             75             80

Gly Ser Arg Gly Phe Asn Cys Glu Ser Lys Pro Glu Ala Glu Glu Thr
      85             90             95

Cys Phe Asp Lys Tyr Thr Gly Asn Thr Tyr Arg Val Gly Asp Thr Tyr
      100            105            110

Glu Arg Pro Lys Asp Ser Met Ile Trp Asp Cys Thr Cys Ile Gly Ala
      115            120            125

Gly Arg Gly Arg Ile Ser Cys Thr Ile Ala Asn Arg Cys His Glu Gly
      130            135            140

Gly Gln Ser Tyr Lys Ile Gly Asp Thr Trp Arg Arg Pro His Glu Thr
      145            150            155            160

```

-continued

Gly Gly Tyr Met Leu Glu Cys Val Cys Leu Gly Asn Gly Lys Gly Glu
 165 170 175
 Trp Thr Cys Lys Pro Ile Ala Glu Lys Cys Phe Asp His Ala Ala Gly
 180 185 190
 Thr Ser Tyr Val Val Gly Glu Thr Trp Glu Lys Pro Tyr Gln Gly Trp
 195 200 205
 Met Met Val Asp Cys Thr Cys Leu Gly Glu Gly Ser Gly Arg Ile Thr
 210 215 220
 Cys Thr Ser Arg Asn Arg Cys Asn Asp Gln Asp Thr Arg Thr Ser Tyr
 225 230 235 240
 Arg Ile Gly Asp Thr Trp Ser Lys Lys Asp Asn Arg Gly Asn Leu Leu
 245 250 255
 Gln Cys Ile Cys Thr Gly Asn Gly Arg Gly Glu Trp Lys Cys Glu Arg
 260 265 270
 His Thr Ser Val Gln Thr Thr Ser Ser Gly Ser Gly Pro Phe Thr Asp
 275 280 285
 Val Arg Ala Ala Val Tyr Gln Pro Gln Pro His Pro Gln Pro Pro Pro
 290 295 300
 Tyr Gly His Cys Val Thr Asp Ser Gly Val Val Tyr Ser Val Gly Met
 305 310 315 320
 Gln Trp Leu Lys Thr Gln Gly Asn Lys Gln Met Leu Cys Thr Cys Leu
 325 330 335
 Gly Asn Gly Val Ser Cys Gln Glu Thr Ala Val Thr Gln Thr Tyr Gly
 340 345 350
 Gly Asn Ser Asn Gly Glu Pro Cys Val Leu Pro Phe Thr Tyr Asn Gly
 355 360 365
 Arg Thr Phe Tyr Ser Cys Thr Thr Glu Gly Arg Gln Asp Gly His Leu
 370 375 380
 Trp Cys Ser Thr Thr Ser Asn Tyr Glu Gln Asp Gln Lys Tyr Ser Phe
 385 390 395 400
 Cys Thr Asp His Thr Val Leu Val Gln Thr Arg Gly Gly Asn Ser Asn
 405 410 415
 Gly Ala Leu Cys His Phe Pro Phe Leu Tyr Asn Asn His Asn Tyr Thr
 420 425 430
 Asp Cys Thr Ser Glu Gly Arg Arg Asp Asn Met Lys Trp Cys Gly Thr
 435 440 445
 Thr Gln Asn Tyr Asp Ala Asp Gln Lys Phe Gly Phe Cys Pro Met Ala
 450 455 460
 Ala His Glu Glu Ile Cys Thr Thr Asn Glu Gly Val Met Tyr Arg Ile
 465 470 475 480
 Gly Asp Gln Trp Asp Lys Gln His Asp Met Gly His Met Met Arg Cys
 485 490 495
 Thr Cys Val Gly Asn Gly Arg Gly Glu Trp Thr Cys Ile Ala Tyr Ser
 500 505 510
 Gln Leu Arg Asp Gln Cys Ile Val Asp Asp Ile Thr Tyr Asn Val Asn
 515 520 525
 Asp Thr Phe His Lys Arg His Glu Glu Gly His Met Leu Asn Cys Thr
 530 535 540
 Cys Phe Gly Gln Gly Arg Gly Arg Trp Lys Cys Asp Pro Val Asp Gln
 545 550 555 560

-continued

Cys Gln Asp Ser Glu Thr Gly Thr Phe Tyr Gln Ile Gly Asp Ser Trp
 565 570 575
 Glu Lys Tyr Val His Gly Val Arg Tyr Gln Cys Tyr Cys Tyr Gly Arg
 580 585 590
 Gly Ile Gly Glu Trp His Cys Gln Pro Leu Gln Thr Tyr Pro Ser Ser
 595 600 605
 Ser Gly Pro Val Glu Val Phe Ile Thr Glu Thr Pro Ser Gln Pro Asn
 610 615 620
 Ser His Pro Ile Gln Trp Asn Ala Pro Gln Pro Ser His Ile Ser Lys
 625 630 635 640
 Tyr Ile Leu Arg Trp Arg Pro Lys Asn Ser Val Gly Arg Trp Lys Glu
 645 650 655
 Ala Thr Ile Pro Gly His Leu Asn Ser Tyr Thr Ile Lys Gly Leu Lys
 660 665 670
 Pro Gly Val Val Tyr Glu Gly Gln Leu Ile Ser Ile Gln Gln Tyr Gly
 675 680 685
 His Gln Glu Val Thr Arg Phe Asp Phe Thr Thr Thr Ser Thr Ser Thr
 690 695 700
 Pro Val Thr Ser Asn Thr Val Thr Gly Glu Thr Thr Pro Phe Ser Pro
 705 710 715 720
 Leu Val Ala Thr Ser Glu Ser Val Thr Glu Ile Thr Ala Ser Ser Phe
 725 730 735
 Val Val Ser Trp Val Ser Ala Ser Asp Thr Val Ser Gly Phe Arg Val
 740 745 750
 Glu Tyr Glu Leu Ser Glu Glu Gly Asp Glu Pro Gln Tyr Leu Asp Leu
 755 760 765
 Pro Ser Thr Ala Thr Ser Val Asn Ile Pro Asp Leu Leu Pro Gly Arg
 770 775 780
 Lys Tyr Ile Val Asn Val Tyr Gln Ile Ser Glu Asp Gly Glu Gln Ser
 785 790 795 800
 Leu Ile Leu Ser Thr Ser Gln Thr Thr Ala Pro Asp Ala Pro Pro Asp
 805 810 815
 Pro Thr Val Asp Gln Val Asp Asp Thr Ser Ile Val Val Arg Trp Ser
 820 825 830
 Arg Pro Gln Ala Pro Ile Thr Gly Tyr Arg Ile Val Tyr Ser Pro Ser
 835 840 845
 Val Glu Gly Ser Ser Thr Glu Leu Asn Leu Pro Glu Thr Ala Asn Ser
 850 855 860
 Val Thr Leu Ser Asp Leu Gln Pro Gly Val Gln Tyr Asn Ile Thr Ile
 865 870 875 880
 Tyr Ala Val Glu Glu Asn Gln Glu Ser Thr Pro Val Val Ile Gln Gln
 885 890 895
 Glu Thr Thr Gly Thr Pro Arg Ser Asp Thr Val Pro Ser Pro Arg Asp
 900 905 910
 Leu Gln Phe Val Glu Val Thr Asp Val Lys Val Thr Ile Met Trp Thr
 915 920 925
 Pro Pro Glu Ser Ala Val Thr Gly Tyr Arg Val Asp Val Ile Pro Val
 930 935 940
 Asn Leu Pro Gly Glu His Gly Gln Arg Leu Pro Ile Ser Arg Asn Thr
 945 950 955 960
 Phe Ala Glu Val Thr Gly Leu Ser Pro Gly Val Thr Tyr Tyr Phe Lys

-continued

His Trp Ile Ala Pro Arg Ala Thr Ile Thr Gly Tyr Arg Ile Arg His
 1380 1385 1390

His Pro Glu His Phe Ser Gly Arg Pro Arg Glu Asp Arg Val Pro His
 1395 1400 1405

Ser Arg Asn Ser Ile Thr Leu Thr Asn Leu Thr Pro Gly Thr Glu Tyr
 1410 1415 1420

Val Val Ser Ile Val Ala Leu Asn Gly Arg Glu Glu Ser Pro Leu Leu
 1425 1430 1435 1440

Ile Gly Gln Gln Ser Thr Val Ser Asp Val Pro Arg Asp Leu Glu Val
 1445 1450 1455

Val Ala Ala Thr Pro Thr Ser Leu Leu Ile Ser Trp Asp Ala Pro Ala
 1460 1465 1470

Val Thr Val Arg Tyr Tyr Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn
 1475 1480 1485

Ser Pro Val Gln Glu Phe Thr Val Pro Gly Ser Lys Ser Thr Ala Thr
 1490 1495 1500

Ile Ser Gly Leu Lys Pro Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala
 1505 1510 1515 1520

Val Thr Gly Arg Gly Asp Ser Pro Ala Ser Ser Lys Pro Ile Ser Ile
 1525 1530 1535

Asn Tyr Arg Thr Glu Ile Asp Lys Pro Ser Gln Met Gln Val Thr Asp
 1540 1545 1550

Val Gln Asp Asn Ser Ile Ser Val Lys Trp Leu Pro Ser Ser Ser Pro
 1555 1560 1565

Val Thr Gly Tyr Arg Val Thr Thr Pro Lys Asn Gly Pro Gly Pro
 1570 1575 1580

Thr Lys Thr Lys Thr Ala Gly Pro Asp Gln Thr Glu Met Thr Ile Glu
 1585 1590 1595 1600

Gly Leu Gln Pro Thr Val Glu Tyr Val Val Ser Val Tyr Ala Gln Asn
 1605 1610 1615

Pro Ser Gly Glu Ser Gln Pro Leu Val Gln Thr Ala Val Thr Asn Ile
 1620 1625 1630

Asp Arg Pro Lys Gly Leu Ala Phe Thr Asp Val Asp Val Asp Ser Ile
 1635 1640 1645

Lys Ile Ala Trp Glu Ser Pro Gln Gly Gln Val Ser Arg Tyr Arg Val
 1650 1655 1660

Thr Tyr Ser Ser Pro Glu Asp Gly Ile His Glu Leu Phe Pro Ala Pro
 1665 1670 1675 1680

Asp Gly Glu Glu Asp Thr Ala Glu Leu Gln Gly Leu Arg Pro Gly Ser
 1685 1690 1695

Glu Tyr Thr Val Ser Val Val Ala Leu His Asp Asp Met Glu Ser Gln
 1700 1705 1710

Pro Leu Ile Gly Thr Gln Ser Thr Ala Ile Pro Ala Pro Thr Asp Leu
 1715 1720 1725

Lys Phe Thr Gln Val Thr Pro Thr Ser Leu Ser Ala Gln Trp Thr Pro
 1730 1735 1740

Pro Asn Val Gln Leu Thr Gly Tyr Arg Val Arg Val Thr Pro Lys Glu
 1745 1750 1755 1760

Lys Thr Gly Pro Met Lys Glu Ile Asn Leu Ala Pro Asp Ser Ser Ser
 1765 1770 1775

-continued

Val Val Val Ser Gly Leu Met Val Ala Thr Lys Tyr Glu Val Ser Val
 1780 1785 1790
 Tyr Ala Leu Lys Asp Thr Leu Thr Ser Arg Pro Ala Gln Gly Val Val
 1795 1800 1805
 Thr Thr Leu Glu Asn Val Ser Pro Pro Arg Arg Ala Arg Val Thr Asp
 1810 1815 1820
 Ala Thr Glu Thr Thr Ile Thr Ile Ser Trp Arg Thr Lys Thr Glu Thr
 1825 1830 1835 1840
 Ile Thr Gly Phe Gln Val Asp Ala Val Pro Ala Asn Gly Gln Thr Pro
 1845 1850 1855
 Ile Gln Arg Thr Ile Lys Pro Asp Val Arg Ser Tyr Thr Ile Thr Gly
 1860 1865 1870
 Leu Gln Pro Gly Thr Asp Tyr Lys Ile Tyr Leu Tyr Thr Leu Asn Asp
 1875 1880 1885
 Asn Ala Arg Ser Ser Pro Val Val Ile Asp Ala Ser Thr Ala Ile Asp
 1890 1895 1900
 Ala Pro Ser Asn Leu Arg Phe Leu Ala Thr Thr Pro Asn Ser Leu Leu
 1905 1910 1915 1920
 Val Ser Trp Gln Pro Pro Arg Ala Arg Ile Thr Gly Tyr Ile Ile Lys
 1925 1930 1935
 Tyr Glu Lys Pro Gly Ser Pro Pro Arg Glu Val Val Pro Arg Pro Arg
 1940 1945 1950
 Pro Gly Val Thr Glu Ala Thr Ile Thr Gly Leu Glu Pro Gly Thr Glu
 1955 1960 1965
 Tyr Thr Ile Tyr Val Ile Ala Leu Lys Asn Asn Gln Lys Ser Glu Pro
 1970 1975 1980
 Leu Ile Gly Arg Lys Lys Thr Val Gln Lys Thr Pro Phe Val Thr His
 1985 1990 1995 2000
 Pro Gly Tyr Asp Thr Gly Asn Gly Ile Gln Leu Pro Gly Thr Ser Gly
 2005 2010 2015
 Gln Gln Pro Ser Val Gly Gln Gln Met Ile Phe Glu Glu His Gly Phe
 2020 2025 2030
 Arg Arg Thr Thr Pro Pro Thr Thr Ala Thr Pro Ile Arg His Arg Pro
 2035 2040 2045
 Arg Pro Tyr Pro Pro Asn Val Gly Gln Glu Ala Leu Ser Gln Thr Thr
 2050 2055 2060
 Ile Ser Trp Ala Pro Phe Gln Asp Thr Ser Glu Tyr Ile Ile Ser Cys
 2065 2070 2075 2080
 His Pro Val Gly Thr Asp Glu Glu Pro Leu Gln Phe Arg Val Pro Gly
 2085 2090 2095
 Thr Ser Thr Ser Ala Thr Leu Thr Gly Leu Thr Arg Gly Ala Thr Tyr
 2100 2105 2110
 Asn Ile Ile Val Glu Ala Leu Lys Asp Gln Gln Arg His Lys Val Arg
 2115 2120 2125
 Glu Glu Val Val Thr Val Gly Asn Ser Val Asn Glu Gly Leu Asn Gln
 2130 2135 2140
 Pro Thr Asp Asp Ser Cys Phe Asp Pro Tyr Thr Val Ser His Tyr Ala
 2145 2150 2155 2160
 Val Gly Asp Glu Trp Glu Arg Met Ser Glu Ser Gly Phe Lys Leu Leu
 2165 2170 2175
 Cys Gln Cys Leu Gly Phe Gly Ser Gly His Phe Arg Cys Asp Ser Ser

-continued

2180	2185	2190
Arg Trp Cys His Asp Asn Gly Val Asn Tyr Lys Ile Gly Glu Lys Trp		
2195	2200	2205
Asp Arg Gln Gly Glu Asn Gly Gln Met Met Ser Cys Thr Cys Leu Gly		
2210	2215	2220
Asn Gly Lys Gly Glu Phe Lys Cys Asp Pro His Glu Ala Thr Cys Tyr		
2225	2230	2235
Asp Asp Gly Lys Thr Tyr His Val Gly Glu Gln Trp Gln Lys Glu Tyr		
2245	2250	2255
Leu Gly Ala Ile Cys Ser Cys Thr Cys Phe Gly Gly Gln Arg Gly Trp		
2260	2265	2270
Arg Cys Asp Asn Cys Arg Arg Pro Gly Gly Glu Pro Ser Pro Glu Gly		
2275	2280	2285
Thr Thr Gly Gln Ser Tyr Asn Gln Tyr Ser Gln Arg Tyr His Gln Arg		
2290	2295	2300
Thr Asn Thr Asn Val Asn Cys Pro Ile Glu Cys Phe Met Pro Leu Asp		
2305	2310	2315
Val Gln Ala Asp Arg Glu Asp Ser Arg Glu		
2325	2330	

<210> SEQ ID NO 22
 <211> LENGTH: 8272
 <212> TYPE: DNA
 <213> ORGANISM: Homo Sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Oncofetal Fibronectin variant 5
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank NM 212476
 <309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 22

```

gccccgcgcc gctgtgctgc acagggggag gagaggaac cccagggcgc agcgggaaga      60
ggggacctgc agccacaact tctctggctc tctgcatccc ttctgtccct ccaccctcc      120
ccttccccac cctctggccc ccaccttctt ggaggcgaca acccccggga ggcattagaa      180
gggatttttc ccgcaggttg cgaaggaag caaacttggg ggcaacttgc ctcccgggtg      240
gggctctctc cccccaccgt ctcaacatgc ttaggggtcc ggggccggg ctgctgctgc      300
tggccgtcca gtgcctgggg acagcgggtg cctccacggg agcctcgaag agcaagaggc      360
aggctcagca aatggttcag cccagctccc cgggtgctgt cagtcaaagc aagcccgtt      420
gttatgacaa tggaaaacac tatcagataa atcaacagtg ggagcggacc tacctaggca      480
atgctgttgt ttgtacttgt tatggaggaa gccgaggttt taactgcgag agtaaacctg      540
aagctgaaga gacttgcctt gacaagtaca ctgggaacac ttaccgagtg ggtgacactt      600
atgagcgtcc taaagactcc atgatctggg actgtacctg catcggggct gggcgaggga      660
gaataagctg taccatcgca aaccgctgcc atgaaggggg tcagtcctac aagattggtg      720
acacctggag gagaccacat gagactggtg gttacatggt agagtgtgtg tgtcttgta      780
atggaaaagg agaatggacc tgcaagccca tagctgagaa gtgttttgat catgctgctg      840
ggacttccta tgtgtcgga gaaacgtggg agaagcccta ccaaggctgg atgatggtag      900
attgtacttg cctgggagaa ggcagcggac gcatcacttg cacttctaga aatagatgca      960
acgatcagga cacaaggaca tcctatagaa ttggagacac ctggagcaag aaggataatc     1020
    
```

-continued

gaggaaacct gctccagtgc atctgcacag gcaacggccg aggagagtgg aagtgtgaga	1080
ggcacacctc tgtgcagacc acatcgagcg gatctggccc cttcaccgat gttcgtgcag	1140
ctgtttacca accgcagcct cccccccagc ctcctccccta tggccactgt gtcacagaca	1200
gtggtgtggt ctactctgtg gggatgcagt ggctgaagac acaaggaaat aagcaaatgc	1260
tttgcacgtg cctgggcaac ggagtcaact gccaaagac agctgtaacc cagacttacg	1320
gtggcaactc aaatggagag ccatgtgtct taccattcac ctacaatggc aggacgttct	1380
actcctgcac cacagaaggg cgacaggacg gacatctttg gtgcagcaca acttcgaatt	1440
atgagcagga ccagaaatac tctttctgca cagaccacac tgttttggtt cagactcgag	1500
gaggaaattc caatggtgcc ttgtgccact tccccttccct atacaacaac cacaattaca	1560
ctgattgcac ttctgagggc agaagagaca acatgaagtg gtgtgggacc acacagaact	1620
atgatgccga ccagaagttt gggttctgcc ccatggctgc ccacgaggaa atctgcacaa	1680
ccaatgaagg ggtcatgtac cgcattggag atcagtggga taagcagcat gacatgggtc	1740
acatgatgag gtgcacgtgt gttgggaatg gtcgtgggga atggacatgc attgcctact	1800
cgcagttcg agatcagtgc attgttgatg acatcactta caatgtgaac gacacattcc	1860
acaagcgtca tgaagagggg cacatgctga actgtacatg cttcggtcag ggtcggggca	1920
ggtggaagtg tgatcccgtc gaccaatgcc aggattcaga gactgggacg ttttatcaaa	1980
ttggagattc atgggagaag tatgtgcatg gtgtcagata ccagtgtac tgctatggcc	2040
gtggcattgg ggagtggcat tgccaacctt tacagaccta tccaagctca agtggctctg	2100
tcgaagtatt tatcactgag actccgagtc agcccaactc ccaccccatc cagtggaatg	2160
caccacagcc atctcacatt tccaagtaca ttctcaggtg gagacctaaa aattctgtag	2220
gccgttgga ggaagctacc ataccaggcc acttaaaactc ctacaccatc aaaggcctga	2280
agcctggtgt ggtatacagag gccacgtca tcagcatcca gcagtacggc caccaagaag	2340
tgactcgctt tgacttcacc accaccagca ccagcacacc tgtgaccagc aacaccgtga	2400
caggagagac gactcccctt tctcctcttg tggccacttc tgaatctgtg accgaaatca	2460
cagccagtag ctttgtggtc tcctgggtct cagcttccga caccgtgtcg ggattccggg	2520
tggaatatga gctgagtgag gagggagatg agccacagta cctggatcct ccaagcacag	2580
ccacttctgt gaacatccct gacctgcttc ctggccgaaa atacattgta aatgtctatc	2640
agatatctga ggatggggag cagagtttga tcctgtctac ttcacaaaca acagcgctctg	2700
atgcccctcc tgacccgact gtggaccaag ttgatgacac ctcaattggt gttcgtgga	2760
gcagacccca ggctcccac acagggatca gaatagtcta ttcgcatca gtagaaggta	2820
gcagcacaga actcaacctt cctgaaactg caaactccgt caccctcagt gacttgcaac	2880
ctggtgttca gtataacatc actatctatg ctgtggaaga aaatcaagaa agtacacctg	2940
ttgtcattca acaagaaacc actggcacc cagctcaga tacagtgcc tctcccaggg	3000
acctgcagtt tgtggaagt acagacgtga aggtcaccat catgtggaca ccgctgaga	3060
gtgcagtgc cggtaccgt gtggatgtga tcccgtcaa cctgcctggc gagcacgggc	3120
agaggctgcc catcagcag aacacctttg cagaagtac cgggctgtcc cctgggggtca	3180
cctattactt caaagtctt gcagtgagcc atgggaggg gagcaagcct ctgactgctc	3240
aacagacaac caaactggat gctcccacta acctccagtt tgtcaatgaa actgattcta	3300

-continued

ctgtcctggt gagatggact ccacctcggg cccagataac aggataccga ctgaccgtgg	3360
gccttaccg aagaggacag cccaggcagt acaatgtggg tccctctgtc tccaagtacc	3420
cactgaggaa tctgcagcct gcacttgagt acaccgtatc cctcgtggcc ataaagggca	3480
accaagagag ccccaaagcc actggagtct ttaccacact gcagcctggg agctctattc	3540
cacottaaa caccgaggtg actgagacca ccattgtgat cacatggacg cctgctccaa	3600
gaattggttt taagctgggt gtacgaccaa gccagggagg agaggcacca cgagaagtga	3660
cttcagactc aggaagcacc gttgtgtccg gcttgactcc aggagtagaa tacgtctaca	3720
ccatccaagt cctgagagat ggacaggaag gagatgcgcc aattgtaaac aaagtgggta	3780
caccattgtc tccaccaaca aacttgcatc tggaggcaaa ccctgacact ggagtgtctc	3840
cagtctcctg ggagaggagc accaccccag acattactgg ttatagaatt accacaacc	3900
ctacaaacgg ccagcaggga aattcttttg aagaagtggg ccatgctgat cagagctcct	3960
gcacttttga taactgaggt cccggcctgg agtacaatgt cagtgtttac actgtcaagg	4020
atgacaagga aagtgtccct atctctgata ccatcatccc agctgttccct cctcccactg	4080
acctgcgatt caccaacatt ggtccagaca ccatgcgtgt cacctgggct ccaccccat	4140
ccattgattt aaccaacttc ctggtgcgtt actcacctgt gaaaaatgag gaagatggtg	4200
cagagttgtc aatttctcct tcagacaatg cagtggctctt aacaaatctc ctgcttggtg	4260
cagaatatgt agtgagtgtc tccagtgtct acgaacaaca tgagagcaca cctcttagag	4320
gaagacagaa aacaggtcct gattcccaa ctggcattga cttttctgat attactgcca	4380
actcttttac tgtgactgg attgtcctc gagccaccat cactggctac aggatccgcc	4440
atcatcccga gcacttcagt gggagacctc gagaagatcg ggtgccccac tctcggaatt	4500
ccatcaccct caccaacctc actccaggca cagagtatgt ggtcagcacc gttgctctta	4560
atggcagaga ggaaagtccc ttattgattg gccacaatc aacagtttct gatgttccga	4620
gggacctgga agttgttctc gcgaccccca ccagcctact gatcagctgg gatgctctc	4680
ctgtcacagt gagatattac aggatcactt acggagagac agggagaaat agccctgtcc	4740
aggagttcac tgtgcctggg agcaagtcta cagctaccat cagcggcctt aaacctggag	4800
ttgattatac catcactgtg tatgtgtca ctggccgtgg agacagcccc gcaagcagca	4860
agccaatttc cattaattac cgaacagaaa ttgacaaacc atcccagatg caagtgaccg	4920
atgttcagga caacagcatt agtgcgaagt ggctgccttc aagttccctt gttactggtt	4980
acagagtaac caccactccc aaaaatggac caggaccaac aaaaactaaa actgcaggtc	5040
cagatcaaac agaaatgact attgaaggct tgcagcccac agtggagtat gttggttagtg	5100
tctatgctca gaatccaagc ggagagagtc agcctctggt tcagactgca gtaaccacta	5160
ttctgcacc aactgacctg aagttcactc aggtcacacc cacaagcctg agcggccagt	5220
ggacaccacc caatgttcag ctcaactggat atcgagtgcg ggtgaccccc aaggagaaga	5280
ccggaccaat gaagaaatc aacctgtctc ctgacagctc atccgtggtt gtatcaggac	5340
ttatggtggc caccaaatat gaagtgagtg tctatgctct taaggacact ttgacaagca	5400
gaccagctca gggagttgtc accactctgg agaatgtcag cccaccaaga agggctcgtg	5460
tgacagatgc tactgagacc accatcacca ttatgctggag aaccaagact gagacgatca	5520
ctggcttcca agttgatgcc gttccagcca atggccagac tccaatccag agaaccatca	5580

-continued

agccagatgt cagaagctac accatcacag gtttacaacc aggcactgac tacaagatct	5640
acctgtacac cttgaatgac aatgctcggg gctcccctgt ggtcatcgac gcctccactg	5700
ccattgatgc accatccaac ctgcgtttcc tggccaccac acccaattcc ttgctgggat	5760
catggcagcc gccacgtgcc aggattaccg gctacatcat caagtatgag aagcctgggt	5820
ctcctcccag agaagtggtc cctcggcccc gccctggtgt cacagaggct actattactg	5880
gcctggaacc gggaaccgaa tatacaatth atgtcattgc cctgaagaat aatcagaaga	5940
gcgagcccc gatggaagc aaaaagacag acgagcttcc ccaactggta acccttccac	6000
acccaatct tcatggacca gagatcttgg atgttccttc cacagttaa aagaccctt	6060
tcgtcaccca cctgggtat gacactggaa atggtattca gcttctggc acttctggtc	6120
agcaaccagc tgttgggcaa caaatgatct ttgaggaaca tggttttagg cggaccacac	6180
cgcccacaac ggccaccccc ataaggcata ggccaagacc ataccgccc aatgtaggtg	6240
aggaaatcca aattggctac atccccagg aagatgtaga ctatcacctg taccacacg	6300
gtcggggact caatccaat gcctctacag gacaagaagc tctctctcag acaaccatct	6360
catgggcccc attccaggac acttctgagt acatcatttc atgtcatcct gttggcactg	6420
atgaagaacc cttacagttc agggttcctg gaacttctac cagtgccact ctgacaggcc	6480
tcaccagagg tcccacctac aacatcatag tggaggcact gaaagaccag cagaggcata	6540
aggttcggga agaggttgtt accgtgggca actctgtcaa cgaaggcttg aaccaaccta	6600
cgatgactc gtgctttgac ccctacacag tttcccatta tgcggttga gatgagtggg	6660
aacgaatgtc tgaatcagcc tttaaactgt tgtgccagt cttaggcttt ggaagtggtc	6720
atctcagatg tgattcatct agatggtgcc atgacaatgg tgtgaactac aagattggag	6780
agaagtggga ccgtcaggga gaaaatggcc agatgatgag ctgcacatgt cttgggaaag	6840
gaaaaggaga attcaagtgt gaccctcatg aggcacagt ttatgatgat gggaaagacat	6900
accacgtagg agaacagtgg cagaaggaat atctcgggtgc ctttctctcc tgcacatgct	6960
ttggaggcca gggggctg cgctgtgaca actgccgag acctgggggt gaaccagtc	7020
ccgaaggcac tactggccag tcctacaacc agtatttctc gagataccat cagagaacaa	7080
acactaatgt taattgcca attgagtgtc tcatgccttt agatgtacag gctgacagag	7140
aagattccc agagtaaatc atctttccaa tccagaggaa caagcatgct tctctgcaa	7200
gatccatcta aactggagt atgttagcag acccagctta gagttcttct tcttttctta	7260
agccctttgc tctggaggaa gttctccagc ttcagctcaa ctacagctt ctccaagcat	7320
caccctggga gtttctgag ggttttctca taaatgaggg ctgcacattg cctgttctgc	7380
ttcgaagtat tcaataccgc tcagtatttt aatgaagtg attctaagat ttggtttggg	7440
atcaatagga aagcatatgc agccaaccaa gatgcaaatg tttgaaatg atatgaccaa	7500
aattttaagt aggaaagtca cccaacact tctgctttca cttaagtgtc tggcccgcaa	7560
tactgtagga acaagcatga tctgttact gtgatatttt aaatatccac agtactcact	7620
ttttccaaat gatcctagta attgcctaga aatatcttcc tcttacctgt tattttatcaa	7680
tttttcccag tatttttata cggaaaaaat tgtattgaaa acacttagta tgcagttgat	7740
aagaggaatt tggataaatt atggtgggtg attatttttt atactgtatg tgcxaaagct	7800
ttactactgt gaaagacaa ctgttttaaat aaaagattta cattccacaa cttgaagttc	7860

-continued

```

atctatttga tataagacac cttcggggga aataattcct gtgaatattc tttttcaatt 7920
cagcaaacat ttgaaaatct atgatgtgca agtctaattg ttgatttcag tacaagattt 7980
tctaaatcag ttgtacaaa aactgattgg tttttgtcac ttcattctctt cactaatgga 8040
gatagcttta cactttctgc tttaatagat ttaagtggac cccaatattt attaaaattg 8100
ctagtttacc gttcagaagt ataatagaaa taatctttag ttgctctttt ctaaccattg 8160
taattcttcc cttcttcocct ccacctttcc ttcattgaat aaacctctgt tcaaagagat 8220
tgctctgcaag ggaataaaaa atgactaaga tattaaaaaa aaaaaaaaaa aa 8272

```

```

<210> SEQ ID NO 23
<211> LENGTH: 2296
<212> TYPE: PRT
<213> ORGANISM: Homo Sapiens
<300> PUBLICATION INFORMATION:
<308> DATABASE ACCESSION NUMBER: GenBank NM 212476
<309> DATABASE ENTRY DATE: 2005-06-10

```

```

<400> SEQUENCE: 23

```

```

Met Leu Arg Gly Pro Gly Pro Gly Leu Leu Leu Ala Val Gln Cys
 1           5           10           15
Leu Gly Thr Ala Val Pro Ser Thr Gly Ala Ser Lys Ser Lys Arg Gln
 20           25           30
Ala Gln Gln Met Val Gln Pro Gln Ser Pro Val Ala Val Ser Gln Ser
 35           40           45
Lys Pro Gly Cys Tyr Asp Asn Gly Lys His Tyr Gln Ile Asn Gln Gln
 50           55           60
Trp Glu Arg Thr Tyr Leu Gly Asn Ala Leu Val Cys Thr Cys Tyr Gly
 65           70           75
Gly Ser Arg Gly Phe Asn Cys Glu Ser Lys Pro Glu Ala Glu Glu Thr
 85           90           95
Cys Phe Asp Lys Tyr Thr Gly Asn Thr Tyr Arg Val Gly Asp Thr Tyr
 100          105          110
Glu Arg Pro Lys Asp Ser Met Ile Trp Asp Cys Thr Cys Ile Gly Ala
 115          120          125
Gly Arg Gly Arg Ile Ser Cys Thr Ile Ala Asn Arg Cys His Glu Gly
 130          135          140
Gly Gln Ser Tyr Lys Ile Gly Asp Thr Trp Arg Arg Pro His Glu Thr
 145          150          155
Gly Gly Tyr Met Leu Glu Cys Val Cys Leu Gly Asn Gly Lys Gly Glu
 165          170          175
Trp Thr Cys Lys Pro Ile Ala Glu Lys Cys Phe Asp His Ala Ala Gly
 180          185          190
Thr Ser Tyr Val Val Gly Glu Thr Trp Glu Lys Pro Tyr Gln Gly Trp
 195          200          205
Met Met Val Asp Cys Thr Cys Leu Gly Glu Gly Ser Gly Arg Ile Thr
 210          215          220
Cys Thr Ser Arg Asn Arg Cys Asn Asp Gln Asp Thr Arg Thr Ser Tyr
 225          230          235
Arg Ile Gly Asp Thr Trp Ser Lys Lys Asp Asn Arg Gly Asn Leu Leu
 245          250          255
Gln Cys Ile Cys Thr Gly Asn Gly Arg Gly Glu Trp Lys Cys Glu Arg
 260          265          270

```


-continued

His	Thr	Ser	Val	Gln	Thr	Thr	Ser	Ser	Gly	Ser	Gly	Pro	Phe	Thr	Asp
		275					280					285			
Val	Arg	Ala	Ala	Val	Tyr	Gln	Pro	Gln	Pro	His	Pro	Gln	Pro	Pro	Pro
	290					295					300				
Tyr	Gly	His	Cys	Val	Thr	Asp	Ser	Gly	Val	Val	Tyr	Ser	Val	Gly	Met
305					310					315					320
Gln	Trp	Leu	Lys	Thr	Gln	Gly	Asn	Lys	Gln	Met	Leu	Cys	Thr	Cys	Leu
				325					330					335	
Gly	Asn	Gly	Val	Ser	Cys	Gln	Glu	Thr	Ala	Val	Thr	Gln	Thr	Tyr	Gly
			340					345						350	
Gly	Asn	Ser	Asn	Gly	Glu	Pro	Cys	Val	Leu	Pro	Phe	Thr	Tyr	Asn	Gly
		355					360					365			
Arg	Thr	Phe	Tyr	Ser	Cys	Thr	Thr	Glu	Gly	Arg	Gln	Asp	Gly	His	Leu
	370					375					380				
Trp	Cys	Ser	Thr	Thr	Ser	Asn	Tyr	Glu	Gln	Asp	Gln	Lys	Tyr	Ser	Phe
385					390					395					400
Cys	Thr	Asp	His	Thr	Val	Leu	Val	Gln	Thr	Arg	Gly	Gly	Asn	Ser	Asn
				405					410					415	
Gly	Ala	Leu	Cys	His	Phe	Pro	Phe	Leu	Tyr	Asn	Asn	His	Asn	Tyr	Thr
			420					425						430	
Asp	Cys	Thr	Ser	Glu	Gly	Arg	Arg	Asp	Asn	Met	Lys	Trp	Cys	Gly	Thr
		435					440					445			
Thr	Gln	Asn	Tyr	Asp	Ala	Asp	Gln	Lys	Phe	Gly	Phe	Cys	Pro	Met	Ala
	450					455					460				
Ala	His	Glu	Glu	Ile	Cys	Thr	Thr	Asn	Glu	Gly	Val	Met	Tyr	Arg	Ile
465					470					475					480
Gly	Asp	Gln	Trp	Asp	Lys	Gln	His	Asp	Met	Gly	His	Met	Met	Arg	Cys
				485					490					495	
Thr	Cys	Val	Gly	Asn	Gly	Arg	Gly	Glu	Trp	Thr	Cys	Ile	Ala	Tyr	Ser
		500						505						510	
Gln	Leu	Arg	Asp	Gln	Cys	Ile	Val	Asp	Asp	Ile	Thr	Tyr	Asn	Val	Asn
		515					520					525			
Asp	Thr	Phe	His	Lys	Arg	His	Glu	Glu	Gly	His	Met	Leu	Asn	Cys	Thr
	530					535					540				
Cys	Phe	Gly	Gln	Gly	Arg	Gly	Arg	Trp	Lys	Cys	Asp	Pro	Val	Asp	Gln
545					550					555					560
Cys	Gln	Asp	Ser	Glu	Thr	Gly	Thr	Phe	Tyr	Gln	Ile	Gly	Asp	Ser	Trp
				565					570					575	
Glu	Lys	Tyr	Val	His	Gly	Val	Arg	Tyr	Gln	Cys	Tyr	Cys	Tyr	Gly	Arg
			580					585						590	
Gly	Ile	Gly	Glu	Trp	His	Cys	Gln	Pro	Leu	Gln	Thr	Tyr	Pro	Ser	Ser
		595					600						605		
Ser	Gly	Pro	Val	Glu	Val	Phe	Ile	Thr	Glu	Thr	Pro	Ser	Gln	Pro	Asn
	610					615					620				
Ser	His	Pro	Ile	Gln	Trp	Asn	Ala	Pro	Gln	Pro	Ser	His	Ile	Ser	Lys
625					630					635					640
Tyr	Ile	Leu	Arg	Trp	Arg	Pro	Lys	Asn	Ser	Val	Gly	Arg	Trp	Lys	Glu
				645					650					655	
Ala	Thr	Ile	Pro	Gly	His	Leu	Asn	Ser	Tyr	Thr	Ile	Lys	Gly	Leu	Lys
			660					665						670	
Pro	Gly	Val	Val	Tyr	Glu	Gly	Gln	Leu	Ile	Ser	Ile	Gln	Gln	Tyr	Gly

-continued

675				680				685							
His	Gln	Glu	Val	Thr	Arg	Phe	Asp	Phe	Thr	Thr	Thr	Ser	Thr	Ser	Thr
	690					695						700			
Pro	Val	Thr	Ser	Asn	Thr	Val	Thr	Gly	Glu	Thr	Thr	Pro	Phe	Ser	Pro
	705				710					715					720
Leu	Val	Ala	Thr	Ser	Glu	Ser	Val	Thr	Glu	Ile	Thr	Ala	Ser	Ser	Phe
					725				730					735	
Val	Val	Ser	Trp	Val	Ser	Ala	Ser	Asp	Thr	Val	Ser	Gly	Phe	Arg	Val
			740						745				750		
Glu	Tyr	Glu	Leu	Ser	Glu	Glu	Gly	Asp	Glu	Pro	Gln	Tyr	Leu	Asp	Leu
		755					760					765			
Pro	Ser	Thr	Ala	Thr	Ser	Val	Asn	Ile	Pro	Asp	Leu	Leu	Pro	Gly	Arg
		770				775					780				
Lys	Tyr	Ile	Val	Asn	Val	Tyr	Gln	Ile	Ser	Glu	Asp	Gly	Glu	Gln	Ser
		785			790					795					800
Leu	Ile	Leu	Ser	Thr	Ser	Gln	Thr	Thr	Ala	Pro	Asp	Ala	Pro	Pro	Asp
				805					810					815	
Pro	Thr	Val	Asp	Gln	Val	Asp	Asp	Thr	Ser	Ile	Val	Val	Arg	Trp	Ser
			820						825				830		
Arg	Pro	Gln	Ala	Pro	Ile	Thr	Gly	Tyr	Arg	Ile	Val	Tyr	Ser	Pro	Ser
		835					840					845			
Val	Glu	Gly	Ser	Ser	Thr	Glu	Leu	Asn	Leu	Pro	Glu	Thr	Ala	Asn	Ser
	850					855					860				
Val	Thr	Leu	Ser	Asp	Leu	Gln	Pro	Gly	Val	Gln	Tyr	Asn	Ile	Thr	Ile
	865				870					875					880
Tyr	Ala	Val	Glu	Glu	Asn	Gln	Glu	Ser	Thr	Pro	Val	Val	Ile	Gln	Gln
				885					890					895	
Glu	Thr	Thr	Gly	Thr	Pro	Arg	Ser	Asp	Thr	Val	Pro	Ser	Pro	Arg	Asp
			900						905				910		
Leu	Gln	Phe	Val	Glu	Val	Thr	Asp	Val	Lys	Val	Thr	Ile	Met	Trp	Thr
		915					920					925			
Pro	Pro	Glu	Ser	Ala	Val	Thr	Gly	Tyr	Arg	Val	Asp	Val	Ile	Pro	Val
	930					935					940				
Asn	Leu	Pro	Gly	Glu	His	Gly	Gln	Arg	Leu	Pro	Ile	Ser	Arg	Asn	Thr
	945				950					955					960
Phe	Ala	Glu	Val	Thr	Gly	Leu	Ser	Pro	Gly	Val	Thr	Tyr	Tyr	Phe	Lys
				965					970					975	
Val	Phe	Ala	Val	Ser	His	Gly	Arg	Glu	Ser	Lys	Pro	Leu	Thr	Ala	Gln
		980							985				990		
Gln	Thr	Thr	Lys	Leu	Asp	Ala	Pro	Thr	Asn	Leu	Gln	Phe	Val	Asn	Glu
		995					1000					1005			
Thr	Asp	Ser	Thr	Val	Leu	Val	Arg	Trp	Thr	Pro	Pro	Arg	Ala	Gln	Ile
	1010					1015					1020				
Thr	Gly	Tyr	Arg	Leu	Thr	Val	Gly	Leu	Thr	Arg	Arg	Gly	Gln	Pro	Arg
	1025				1030					1035					1040
Gln	Tyr	Asn	Val	Gly	Pro	Ser	Val	Ser	Lys	Tyr	Pro	Leu	Arg	Asn	Leu
				1045					1050					1055	
Gln	Pro	Ala	Ser	Glu	Tyr	Thr	Val	Ser	Leu	Val	Ala	Ile	Lys	Gly	Asn
		1060						1065					1070		
Gln	Glu	Ser	Pro	Lys	Ala	Thr	Gly	Val	Phe	Thr	Thr	Leu	Gln	Pro	Gly
	1075						1080					1085			

-continued

Ser Ser Ile Pro Pro Tyr Asn Thr Glu Val Thr Glu Thr Thr Ile Val
 1090 1095 1100

Ile Thr Trp Thr Pro Ala Pro Arg Ile Gly Phe Lys Leu Gly Val Arg
 1105 1110 1115 1120

Pro Ser Gln Gly Gly Glu Ala Pro Arg Glu Val Thr Ser Asp Ser Gly
 1125 1130 1135

Ser Ile Val Val Ser Gly Leu Thr Pro Gly Val Glu Tyr Val Tyr Thr
 1140 1145 1150

Ile Gln Val Leu Arg Asp Gly Gln Glu Arg Asp Ala Pro Ile Val Asn
 1155 1160 1165

Lys Val Val Thr Pro Leu Ser Pro Pro Thr Asn Leu His Leu Glu Ala
 1170 1175 1180

Asn Pro Asp Thr Gly Val Leu Thr Val Ser Trp Glu Arg Ser Thr Thr
 1185 1190 1195 1200

Pro Asp Ile Thr Gly Tyr Arg Ile Thr Thr Thr Pro Thr Asn Gly Gln
 1205 1210 1215

Gln Gly Asn Ser Leu Glu Glu Val Val His Ala Asp Gln Ser Ser Cys
 1220 1225 1230

Thr Phe Asp Asn Leu Ser Pro Gly Leu Glu Tyr Asn Val Ser Val Tyr
 1235 1240 1245

Thr Val Lys Asp Asp Lys Glu Ser Val Pro Ile Ser Asp Thr Ile Ile
 1250 1255 1260

Pro Ala Val Pro Pro Pro Thr Asp Leu Arg Phe Thr Asn Ile Gly Pro
 1265 1270 1275 1280

Asp Thr Met Arg Val Thr Trp Ala Pro Pro Ser Ile Asp Leu Thr
 1285 1290 1295

Asn Phe Leu Val Arg Tyr Ser Pro Val Lys Asn Glu Glu Asp Val Ala
 1300 1305 1310

Glu Leu Ser Ile Ser Pro Ser Asp Asn Ala Val Val Leu Thr Asn Leu
 1315 1320 1325

Leu Pro Gly Thr Glu Tyr Val Val Ser Val Ser Ser Val Tyr Glu Gln
 1330 1335 1340

His Glu Ser Thr Pro Leu Arg Gly Arg Gln Lys Thr Gly Leu Asp Ser
 1345 1350 1355 1360

Pro Thr Gly Ile Asp Phe Ser Asp Ile Thr Ala Asn Ser Phe Thr Val
 1365 1370 1375

His Trp Ile Ala Pro Arg Ala Thr Ile Thr Gly Tyr Arg Ile Arg His
 1380 1385 1390

His Pro Glu His Phe Ser Gly Arg Pro Arg Glu Asp Arg Val Pro His
 1395 1400 1405

Ser Arg Asn Ser Ile Thr Leu Thr Asn Leu Thr Pro Gly Thr Glu Tyr
 1410 1415 1420

Val Val Ser Ile Val Ala Leu Asn Gly Arg Glu Glu Ser Pro Leu Leu
 1425 1430 1435 1440

Ile Gly Gln Gln Ser Thr Val Ser Asp Val Pro Arg Asp Leu Glu Val
 1445 1450 1455

Val Ala Ala Thr Pro Thr Ser Leu Leu Ile Ser Trp Asp Ala Pro Ala
 1460 1465 1470

Val Thr Val Arg Tyr Tyr Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn
 1475 1480 1485

-continued

Ser Pro Val Gln Glu Phe Thr Val Pro Gly Ser Lys Ser Thr Ala Thr
 1490 1495 1500
 Ile Ser Gly Leu Lys Pro Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala
 1505 1510 1515 1520
 Val Thr Gly Arg Gly Asp Ser Pro Ala Ser Ser Lys Pro Ile Ser Ile
 1525 1530 1535
 Asn Tyr Arg Thr Glu Ile Asp Lys Pro Ser Gln Met Gln Val Thr Asp
 1540 1545 1550
 Val Gln Asp Asn Ser Ile Ser Val Lys Trp Leu Pro Ser Ser Ser Pro
 1555 1560 1565
 Val Thr Gly Tyr Arg Val Thr Thr Thr Pro Lys Asn Gly Pro Gly Pro
 1570 1575 1580
 Thr Lys Thr Lys Thr Ala Gly Pro Asp Gln Thr Glu Met Thr Ile Glu
 1585 1590 1595 1600
 Gly Leu Gln Pro Thr Val Glu Tyr Val Val Ser Val Tyr Ala Gln Asn
 1605 1610 1615
 Pro Ser Gly Glu Ser Gln Pro Leu Val Gln Thr Ala Val Thr Thr Ile
 1620 1625 1630
 Pro Ala Pro Thr Asp Leu Lys Phe Thr Gln Val Thr Pro Thr Ser Leu
 1635 1640 1645
 Ser Ala Gln Trp Thr Pro Pro Asn Val Gln Leu Thr Gly Tyr Arg Val
 1650 1655 1660
 Arg Val Thr Pro Lys Glu Lys Thr Gly Pro Met Lys Glu Ile Asn Leu
 1665 1670 1675 1680
 Ala Pro Asp Ser Ser Ser Val Val Val Ser Gly Leu Met Val Ala Thr
 1685 1690 1695
 Lys Tyr Glu Val Ser Val Tyr Ala Leu Lys Asp Thr Leu Thr Ser Arg
 1700 1705 1710
 Pro Ala Gln Gly Val Val Thr Thr Leu Glu Asn Val Ser Pro Pro Arg
 1715 1720 1725
 Arg Ala Arg Val Thr Asp Ala Thr Glu Thr Thr Ile Thr Ile Ser Trp
 1730 1735 1740
 Arg Thr Lys Thr Glu Thr Ile Thr Gly Phe Gln Val Asp Ala Val Pro
 1745 1750 1755 1760
 Ala Asn Gly Gln Thr Pro Ile Gln Arg Thr Ile Lys Pro Asp Val Arg
 1765 1770 1775
 Ser Tyr Thr Ile Thr Gly Leu Gln Pro Gly Thr Asp Tyr Lys Ile Tyr
 1780 1785 1790
 Leu Tyr Thr Leu Asn Asp Asn Ala Arg Ser Ser Pro Val Val Ile Asp
 1795 1800 1805
 Ala Ser Thr Ala Ile Asp Ala Pro Ser Asn Leu Arg Phe Leu Ala Thr
 1810 1815 1820
 Thr Pro Asn Ser Leu Leu Val Ser Trp Gln Pro Pro Arg Ala Arg Ile
 1825 1830 1835 1840
 Thr Gly Tyr Ile Ile Lys Tyr Glu Lys Pro Gly Ser Pro Pro Arg Glu
 1845 1850 1855
 Val Val Pro Arg Pro Arg Pro Gly Val Thr Glu Ala Thr Ile Thr Gly
 1860 1865 1870
 Leu Glu Pro Gly Thr Glu Tyr Thr Ile Tyr Val Ile Ala Leu Lys Asn
 1875 1880 1885
 Asn Gln Lys Ser Glu Pro Leu Ile Gly Arg Lys Lys Thr Asp Glu Leu

-continued

<210> SEQ ID NO 24
<211> LENGTH: 7912
<212> TYPE: DNA
<213> ORGANISM: Homo Sapiens
<220> FEATURE:
)
<223> OTHER INFORMATION: Oncofetal Fibronectin variant 6
<300> PUBLICATION INFORMATION:
<308> DATABASE ACCESSION NUMBER: GenBank NM 212474
<309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 24

gccccgcgcc gctgtgctgc acagggggag gagagggaac cccaggcgcg agcgggaaga 60
ggggacctgc agccacaact tctctggctc tctgcatccc ttctgtccct ccaccctcc 120
ccttccccac cctctggccc ccaccttctt ggaggcgaca acccccggga ggcattagaa 180
gggatttttc ccgcaggttg cgaagggaaag caaacttggg ggcaacttgc ctcccgggtg 240
gggctctctc cccccaccgt ctcaacatgc ttaggggtcc gggcccggg ctgctgctgc 300
tggccgtcca gtgcctgggg acagcgggtg cctccacggg agcctcgaag agcaagaggc 360
aggctcagca aatggttcag ccccagctcc cgggtgctgt cagtcaaagc aagcccgtt 420
gttatgacaa tggaaaacac tatcagataa atcaacagtg ggagcggacc tacctaggca 480
atgctgttgt ttgtacttgt tatggaggaa gccgaggttt taactgcgag agtaaacctg 540
aagctgaaga gacttgcctt gacaagtaca ctgggaacac ttaccgagtg ggtgacactt 600
atgagcgtcc taaagactcc atgatctggg actgtacctg catcggggct gggcgaggga 660
gaataagctg taccatcgca aaccgctgcc atgaaggggg tcagtcctac aagattggtg 720
acacctggag gagaccacat gagactggtg gttacatggt agagtgtgtg tgtcttgta 780
atggaaaagg agaatggacc tgcaagccca tagctgagaa gtgttttgat catgctgctg 840
ggacttccta tgtgtcgga gaaacgtggg agaagcccta ccaaggctgg atgatgtag 900
attgtacttg cctgggagaa ggcagcggac gcatcacttg cacttctaga aatagatgca 960
acgatcagga cacaaggaca tcctatagaa ttggagacac ctggagcaag aaggataatc 1020
gaggaaacct gctccagtgc atctgcacag gcaacggccg aggagagtgg aagtgtgaga 1080
ggcacacctc tgtcagacc acatcgagcg gatctggccc cttcaccgat gttcgtgag 1140
ctgtttacca accgcagcct cccccccagc ctccctcccta tggccactgt gtcacagaca 1200
gtggtgtggt ctactctgtg gggatgcagt ggctgaagac acaagaaat aagcaaatgc 1260
tttgacagt cctgggcaac ggagtcagct gccaaagagac agctgtaacc cagacttacg 1320
gtggcaactc aaatggagag ccattgtctt taccattcac ctacaatggc aggacgttct 1380
actcctgcac cacagaaggg cgacaggacg gacatctttg gtgcagcaca acttcgaatt 1440
atgagcagga ccagaataac tctttctgca cagaccacac tgttttggtt cagactcgag 1500
gaggaaattc caatggtgcc ttgtgccact tccccttct atacaacaac cacaattaca 1560
ctgattgcac ttctgagggc agaagagaca acatgaagtg gtgtgggacc acacagaact 1620
atgatgccga ccagaagttt gggttctgcc ccatggctgc ccacgaggaa atctgcacaa 1680
ccaatgaagg ggtcatgtac cgcattggag atcagtgagg taagcagcat gacatgggtc 1740
acatgatgag gtgcacgtgt gttgggaatg gtcgtgggga atggacatgc attgcctact 1800
cgcagcttcg agatcagtgc attgtgatg acatcactta caatgtgaac gacacattcc 1860

-continued

acaagcgtca tgaagagggg cacatgctga actgtacatg cttcggtcag ggtcggggca	1920
ggtggaagtg tgatcccgtc gaccaatgcc aggattcaga gactgggacg ttttatcaaa	1980
ttggagattc atgggagaag tatgtgcatg gtgtcagata ccagtgtac tgctatggcc	2040
gtggcattgg ggagtggcat tgccaacctt tacagaccta tccaagctca agtggctctg	2100
tcgaagtatt tatcactgag actccgagtc agcccaactc ccaccccatc cagtggaatg	2160
caccacagcc atctcacatt tccaagtaca ttctcagtg gagacctaaa aattctgtag	2220
gccgttgga ggaagtacc ataccaggcc acttaaacctc ctacaccatc aaaggcctga	2280
agcctgggtg ggtatacgag gccagctca tcagcatcca gcagtacggc caccaagaag	2340
tgactcgctt tgacttcacc accaccagca ccagcacacc tgtgaccagc aacaccgtga	2400
caggagagac gactcccctt tctcctcttg tggccacttc tgaatctgtg accgaaatca	2460
cagccagtag ctttgtggtc tcctgggtct cagcttccga caccgtgtcg ggattccggg	2520
tggaatatga gctgagtgag gagggagatg agccacagta cctggatctt ccaagcacag	2580
ccacttctgt gaacatccct gacctgcttc ctggccgaaa atacattgta aatgtctatc	2640
agatatctga ggatggggag cagagtttga tcctgtctac ttcacaaaaca acagcgcctg	2700
atgccccctc tgacccgact gtggaccaag ttgatgacac ctcaattggt gttcgtgga	2760
gcagacccca ggctcccctc acagggatca gaatagtcta ttcgccatca gtagaaggta	2820
gcagcacaga actcaacctt cctgaaactg caaactccgt caccctcagt gacttgcaac	2880
ctggtgttca gtataacatc actatctatg ctgtggaaga aaatcaagaa agtacacctg	2940
ttgtcattca acaagaaacc actggcacc caccctcaga tacagtgcc tctcccaggg	3000
acctgcagtt tgtggaagt acagacgtga aggtcaccat catgtggaca ccgctgaga	3060
gtgcagtgac cggctaccgt gtggatgtga tcccctcaa cctgcctggc gagcacgggc	3120
agaggctgcc catcagcagg aacaccttg cagaagtac cgggctgtcc cctggggctca	3180
cctattactt caaagtctt gcagtgagcc atgggagggg gagcaagcct ctgactgctc	3240
aacagacaac caaactggat gctcccacta acctccagtt tgtcaatgaa actgattcta	3300
ctgtcctggt gatgtggact ccacctcgg cccagataac aggataccga ctgacctgg	3360
gccttaccgg aagaggacag cccaggcagt acaatgtggg tccctctgtc tccaagtacc	3420
cactgaggaa tctgcagcct gcactgagt acaccgtatc cctcgtggcc ataaagggca	3480
accaagagag ccccaagacc actggagtct ttaccacact gcagcctggg agctctatc	3540
caccttaca caccgaggt actgagacca coattgtgat cacatggacg cctgctccaa	3600
gaattggtt taagctgggt gtacgaccaa gccagggagg agaggcacca cgagaagtga	3660
cttcagactc aggaagcatc gttgtgtccg gcttgactcc aggagtagaa tacgtctaca	3720
ccatccaagt cctgagagat ggacagggaa gagatgcgcc aattgtaaac aaagtgggta	3780
caccattgtc tccaccaaca aacttgcatc tggaggcaaa ccctgacact ggagtgtca	3840
cagtctcctg ggagaggagc accaccccag acattactgg ttatagaatt accacaacc	3900
ctacaaacgg ccagcaggga aattctttgg aagaagtggc ccatgctgat cagagctcct	3960
gcacttttga taacctgagt cccggcctgg agtacaatgt cagtgtttac actgtcaagg	4020
atgacaagga aagtgtccct atctctgata ccatcatccc agctgttctc cctcccactg	4080
acctgcgatt caccaacatt ggtccagaca ccatgcgtgt cacctgggct ccaccccat	4140

-continued

ccattgattt aaccaacttc ctggtgcgtt actcacctgt gaaaaatgag gaagatggtg 4200
cagagttgtc aattttctct tcagacaatg cagtggctctt aacaaatctc ctgcttggtgta 4260
cagaatatgt agtgagtgtc tccagtgtct acgaacaaca tgagagcaca cctcttagag 4320
gaagacagaa aacaggctctt gattcccca ctggcattga cttttctgat attactgcca 4380
actcttttac tgtgactggt attgtcctc gagccacat cactggctac aggatccgcc 4440
atcatcccga gcaactcagt gggagacctc gagaagatcg ggtgccccac tctcggaatt 4500
ccatcaccct caccaacctc actccaggca cagagtatgt ggtcagcatc gttgctctta 4560
atggcagaga ggaaagtccc ttattgattg gccacaatc aacagttct gatggtccga 4620
gggacctgga agttgtgtc gcgacccca ccagcctact gatcagctgg gatgctctg 4680
ctgtcacagt gagatattac aggatcactt acggagagac aggaggaat agccctgtcc 4740
aggagttcac tgtgcctggg agcaagtcta cagctacat cagcggcctt aaacctggag 4800
ttgattatac catcactgtg tatgtgtca ctggccgtgg agacagcccc gcaagcagca 4860
agccaatttc cattaattac cgaacagaaa ttgacaaacc atcccagatg caagtgaccg 4920
atgttcagga caacagcatt agtgtcaagt ggctgccttc aagttccct gttactggtt 4980
acagagtaac caccactccc aaaaatggac caggaccaac aaaaactaaa actgcaggtc 5040
cagatcaaac agaaatgact attgaaggct tgcagcccc agtggagtat gtggttagtg 5100
tctatgctca gaatccaagc ggagagagtc agcctctggt tcagactgca gtaaccacta 5160
ttcctgcacc aactgacctg aagttcactc aggtcacacc cacaagcctg agcgcaccgt 5220
ggacaccacc caatgttcag ctcaactggat atcgagtgcg ggtgaccccc aaggagaaga 5280
ccggaccaat gaaagaaatc aacctgtctc ctgacagctc atccgtggtt gtatcaggac 5340
ttatggtggc caccaaatat gaagtgagtg tctatgctct taaggacact ttgacaagca 5400
gaccagctca gggagtgtc accactctgg agaatgtcag cccaccaaga agggctctgtg 5460
tgacagatgc tactgagacc accatcacca ttagtggag aaccaagact gagacgatca 5520
ctggcttcca agttgatgcc gttccagcca atggccagac tccaatccag agaaccatca 5580
agccagatgt cagaagctac accatcacag gtttacaacc aggcaactgac tacaagatct 5640
acctgtacac cttgaatgac aatgctcgga gctcccctgt ggtcatcgac gcctccactg 5700
ccattgatgc accatccaac ctgctgttcc tggccaccac acccaattcc ttgctggtat 5760
catggcagcc gccacgtgcc aggattaccg gctacatcat caagtatgag aagcctgggt 5820
ctctcccag agaagtggct cctcggcccc gccctggtgt cacagaggct actattactg 5880
gcctggaacc gggaaccgaa tatacaattt atgtcattgc cctgaagaat aatcagaaga 5940
gcgagcccc gattggaagg aaaaagacag gacaagaagc tctctctcag acaaccatct 6000
catgggcccc attccaggac acttctgagt acatcattc atgtcatcct gttggcactg 6060
atgaagaacc cttacagttc agggttcctg gaacttctac cagtgccact ctgacaggcc 6120
tcaccagagg tgccacctac aacatcatag tggaggcact gaaagaccag cagaggcata 6180
aggttcggga agaggtgtt accgtgggca actctgtcaa cgaaggcttg aaccaacta 6240
cggatgactc gtgctttgac ccctacacag tttcccatta tgccgttggga gatgagtggg 6300
aacgaatgtc tgaatcagc tttaaactgt tgtgccagt cttaggcttt ggaagtggc 6360
atctcagatg tgattcatct agatgggtcc atgacaatg tgtgaactac aagattggag 6420

-continued

agaagtggga ccgtcagggg gaaaatggcc agatgatgag ctgcacatgt cttgggaacg 6480
 gaaaaggaga attcaagtgt gaccctcatg aggcaacgtg ttatgatgat gggagacat 6540
 accacgtagg agaacagtgg cagaaggaat atctcgggtgc catttgctcc tgcacatgct 6600
 ttggaggcca gcggggctgg cgctgtgaca actgccgag acctgggggt gaaccagtc 6660
 ccgaaggcac tactggccag tcctacaacc agtatttca gagataccat cagagaacaa 6720
 aactaatgt taattgccca attgagtgt tcatgccttt agatgtacag gctgacagag 6780
 aagattcccc agagtaaatc atctttccaa tccagaggaa caagcatgct tctctgccaa 6840
 gatccatcta aactggagtg atgttagcag acccagctta gagttcttct tcttttctta 6900
 agccctttgc tctggaggaa gttctccagc ttcagctcaa ctcacagctt ctccaagcat 6960
 caccctggga gtttctctag ggttttctca taaatgaggg ctgcacattg cctgttctgc 7020
 ttcgaagtat tcaataccgc tcagtatttt aaatgaagtg attctaagat ttggtttggg 7080
 atcaatagga aagcatatgc agccaaccaa gatgcaaag ttttgaagt atagaccaa 7140
 aattttaagt aggaaagtca cccaacact tctgctttca cttaagtgtc tggcccgcaa 7200
 tactgtagga acaagcatga tcttgttact gtgatatttt aaatatccac agtactcact 7260
 ttttccaaat gatcctagta attgcctaga aatatctttc tcttacctgt tatttatcaa 7320
 tttttccag tatttttata cggaaaaaat tgtattgaaa acacttagta tgcagttgat 7380
 aagaggaatt tggataaatt atggtgggtg attatttttt atactgtatg tgccaaagct 7440
 ttactactgt gaaagacaa ctgttttaaat aaaagattta cattccacaa cttgaagttc 7500
 atctatttga tataagacac ctccggggga aataattcct gtgaatattc tttttcaatt 7560
 cagcaaacat ttgaaatct atgatgtgca agtctaattg ttgatttcag tacaagattt 7620
 tctaaatcag ttgtacaaa aactgattgg tttttgtcac ttcattctct cactaatgga 7680
 gatagcttta cactttctgc ttaaatagat ttaagtggac cccaatattt attaaaattg 7740
 ctagtttacc gttcagaagt ataatagaaa taatcttttag ttgctctttt ctaaccattg 7800
 taattcttcc ctctctccct ccacctttcc ttcattgaat aaacctctgt tcaaagagat 7860
 tgctgcaag gaaataaaa atgactaaga tattaaaaaa aaaaaaaaaa aa 7912

<210> SEQ ID NO 25

<211> LENGTH: 2176

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank NM 212474

<309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 25

Met Leu Arg Gly Pro Gly Pro Gly Leu Leu Leu Leu Ala Val Gln Cys
 1 5 10 15
 Leu Gly Thr Ala Val Pro Ser Thr Gly Ala Ser Lys Ser Lys Arg Gln
 20 25 30
 Ala Gln Gln Met Val Gln Pro Gln Ser Pro Val Ala Val Ser Gln Ser
 35 40 45
 Lys Pro Gly Cys Tyr Asp Asn Gly Lys His Tyr Gln Ile Asn Gln Gln
 50 55 60
 Trp Glu Arg Thr Tyr Leu Gly Asn Ala Leu Val Cys Thr Cys Tyr Gly
 65 70 75 80

-continued

Gly Ser Arg Gly Phe Asn Cys Glu Ser Lys Pro Glu Ala Glu Glu Thr
 85 90 95
 Cys Phe Asp Lys Tyr Thr Gly Asn Thr Tyr Arg Val Gly Asp Thr Tyr
 100 105 110
 Glu Arg Pro Lys Asp Ser Met Ile Trp Asp Cys Thr Cys Ile Gly Ala
 115 120 125
 Gly Arg Gly Arg Ile Ser Cys Thr Ile Ala Asn Arg Cys His Glu Gly
 130 135 140
 Gly Gln Ser Tyr Lys Ile Gly Asp Thr Trp Arg Arg Pro His Glu Thr
 145 150 155
 Gly Gly Tyr Met Leu Glu Cys Val Cys Leu Gly Asn Gly Lys Gly Glu
 165 170 175
 Trp Thr Cys Lys Pro Ile Ala Glu Lys Cys Phe Asp His Ala Ala Gly
 180 185 190
 Thr Ser Tyr Val Val Gly Glu Thr Trp Glu Lys Pro Tyr Gln Gly Trp
 195 200 205
 Met Met Val Asp Cys Thr Cys Leu Gly Glu Gly Ser Gly Arg Ile Thr
 210 215 220
 Cys Thr Ser Arg Asn Arg Cys Asn Asp Gln Asp Thr Arg Thr Ser Tyr
 225 230 235 240
 Arg Ile Gly Asp Thr Trp Ser Lys Lys Asp Asn Arg Gly Asn Leu Leu
 245 250 255
 Gln Cys Ile Cys Thr Gly Asn Gly Arg Gly Glu Trp Lys Cys Glu Arg
 260 265 270
 His Thr Ser Val Gln Thr Thr Ser Ser Gly Ser Gly Pro Phe Thr Asp
 275 280 285
 Val Arg Ala Ala Val Tyr Gln Pro Gln Pro His Pro Gln Pro Pro Pro
 290 295 300
 Tyr Gly His Cys Val Thr Asp Ser Gly Val Val Tyr Ser Val Gly Met
 305 310 315 320
 Gln Trp Leu Lys Thr Gln Gly Asn Lys Gln Met Leu Cys Thr Cys Leu
 325 330 335
 Gly Asn Gly Val Ser Cys Gln Glu Thr Ala Val Thr Gln Thr Tyr Gly
 340 345 350
 Gly Asn Ser Asn Gly Glu Pro Cys Val Leu Pro Phe Thr Tyr Asn Gly
 355 360 365
 Arg Thr Phe Tyr Ser Cys Thr Thr Glu Gly Arg Gln Asp Gly His Leu
 370 375 380
 Trp Cys Ser Thr Thr Ser Asn Tyr Glu Gln Asp Gln Lys Tyr Ser Phe
 385 390 395 400
 Cys Thr Asp His Thr Val Leu Val Gln Thr Arg Gly Gly Asn Ser Asn
 405 410 415
 Gly Ala Leu Cys His Phe Pro Phe Leu Tyr Asn Asn His Asn Tyr Thr
 420 425 430
 Asp Cys Thr Ser Glu Gly Arg Arg Asp Asn Met Lys Trp Cys Gly Thr
 435 440 445
 Thr Gln Asn Tyr Asp Ala Asp Gln Lys Phe Gly Phe Cys Pro Met Ala
 450 455 460
 Ala His Glu Glu Ile Cys Thr Thr Asn Glu Gly Val Met Tyr Arg Ile
 465 470 475 480

-continued

Gly Asp Gln Trp Asp Lys Gln His Asp Met Gly His Met Met Arg Cys
 485 490 495
 Thr Cys Val Gly Asn Gly Arg Gly Glu Trp Thr Cys Ile Ala Tyr Ser
 500 505 510
 Gln Leu Arg Asp Gln Cys Ile Val Asp Asp Ile Thr Tyr Asn Val Asn
 515 520 525
 Asp Thr Phe His Lys Arg His Glu Glu Gly His Met Leu Asn Cys Thr
 530 535 540
 Cys Phe Gly Gln Gly Arg Gly Arg Trp Lys Cys Asp Pro Val Asp Gln
 545 550 555 560
 Cys Gln Asp Ser Glu Thr Gly Thr Phe Tyr Gln Ile Gly Asp Ser Trp
 565 570 575
 Glu Lys Tyr Val His Gly Val Arg Tyr Gln Cys Tyr Cys Tyr Gly Arg
 580 585 590
 Gly Ile Gly Glu Trp His Cys Gln Pro Leu Gln Thr Tyr Pro Ser Ser
 595 600 605
 Ser Gly Pro Val Glu Val Phe Ile Thr Glu Thr Pro Ser Gln Pro Asn
 610 615 620
 Ser His Pro Ile Gln Trp Asn Ala Pro Gln Pro Ser His Ile Ser Lys
 625 630 635 640
 Tyr Ile Leu Arg Trp Arg Pro Lys Asn Ser Val Gly Arg Trp Lys Glu
 645 650 655
 Ala Thr Ile Pro Gly His Leu Asn Ser Tyr Thr Ile Lys Gly Leu Lys
 660 665 670
 Pro Gly Val Val Tyr Glu Gly Gln Leu Ile Ser Ile Gln Gln Tyr Gly
 675 680 685
 His Gln Glu Val Thr Arg Phe Asp Phe Thr Thr Thr Ser Thr Ser Thr
 690 695 700
 Pro Val Thr Ser Asn Thr Val Thr Gly Glu Thr Thr Pro Phe Ser Pro
 705 710 715 720
 Leu Val Ala Thr Ser Glu Ser Val Thr Glu Ile Thr Ala Ser Ser Phe
 725 730 735
 Val Val Ser Trp Val Ser Ala Ser Asp Thr Val Ser Gly Phe Arg Val
 740 745 750
 Glu Tyr Glu Leu Ser Glu Glu Gly Asp Glu Pro Gln Tyr Leu Asp Leu
 755 760 765
 Pro Ser Thr Ala Thr Ser Val Asn Ile Pro Asp Leu Leu Pro Gly Arg
 770 775 780
 Lys Tyr Ile Val Asn Val Tyr Gln Ile Ser Glu Asp Gly Glu Gln Ser
 785 790 795 800
 Leu Ile Leu Ser Thr Ser Gln Thr Thr Ala Pro Asp Ala Pro Pro Asp
 805 810 815
 Pro Thr Val Asp Gln Val Asp Asp Thr Ser Ile Val Val Arg Trp Ser
 820 825 830
 Arg Pro Gln Ala Pro Ile Thr Gly Tyr Arg Ile Val Tyr Ser Pro Ser
 835 840 845
 Val Glu Gly Ser Ser Thr Glu Leu Asn Leu Pro Glu Thr Ala Asn Ser
 850 855 860
 Val Thr Leu Ser Asp Leu Gln Pro Gly Val Gln Tyr Asn Ile Thr Ile
 865 870 875 880
 Tyr Ala Val Glu Glu Asn Gln Glu Ser Thr Pro Val Val Ile Gln Gln

-continued

885					890					895					
Glu	Thr	Thr	Gly	Thr	Pro	Arg	Ser	Asp	Thr	Val	Pro	Ser	Pro	Arg	Asp
			900					905					910		
Leu	Gln	Phe	Val	Glu	Val	Thr	Asp	Val	Lys	Val	Thr	Ile	Met	Trp	Thr
		915					920					925			
Pro	Pro	Glu	Ser	Ala	Val	Thr	Gly	Tyr	Arg	Val	Asp	Val	Ile	Pro	Val
		930					935					940			
Asn	Leu	Pro	Gly	Glu	His	Gly	Gln	Arg	Leu	Pro	Ile	Ser	Arg	Asn	Thr
		945					950					955			960
Phe	Ala	Glu	Val	Thr	Gly	Leu	Ser	Pro	Gly	Val	Thr	Tyr	Tyr	Phe	Lys
				965					970					975	
Val	Phe	Ala	Val	Ser	His	Gly	Arg	Glu	Ser	Lys	Pro	Leu	Thr	Ala	Gln
			980						985					990	
Gln	Thr	Thr	Lys	Leu	Asp	Ala	Pro	Thr	Asn	Leu	Gln	Phe	Val	Asn	Glu
			995					1000					1005		
Thr	Asp	Ser	Thr	Val	Leu	Val	Arg	Trp	Thr	Pro	Pro	Arg	Ala	Gln	Ile
		1010					1015					1020			
Thr	Gly	Tyr	Arg	Leu	Thr	Val	Gly	Leu	Thr	Arg	Arg	Gly	Gln	Pro	Arg
		1025					1030					1035			1040
Gln	Tyr	Asn	Val	Gly	Pro	Ser	Val	Ser	Lys	Tyr	Pro	Leu	Arg	Asn	Leu
				1045					1050					1055	
Gln	Pro	Ala	Ser	Glu	Tyr	Thr	Val	Ser	Leu	Val	Ala	Ile	Lys	Gly	Asn
			1060						1065					1070	
Gln	Glu	Ser	Pro	Lys	Ala	Thr	Gly	Val	Phe	Thr	Thr	Leu	Gln	Pro	Gly
			1075					1080					1085		
Ser	Ser	Ile	Pro	Pro	Tyr	Asn	Thr	Glu	Val	Thr	Glu	Thr	Thr	Ile	Val
		1090					1095					1100			
Ile	Thr	Trp	Thr	Pro	Ala	Pro	Arg	Ile	Gly	Phe	Lys	Leu	Gly	Val	Arg
		1105					1110					1115			1120
Pro	Ser	Gln	Gly	Gly	Glu	Ala	Pro	Arg	Glu	Val	Thr	Ser	Asp	Ser	Gly
				1125					1130					1135	
Ser	Ile	Val	Val	Ser	Gly	Leu	Thr	Pro	Gly	Val	Glu	Tyr	Val	Tyr	Thr
			1140						1145					1150	
Ile	Gln	Val	Leu	Arg	Asp	Gly	Gln	Glu	Arg	Asp	Ala	Pro	Ile	Val	Asn
			1155				1160						1165		
Lys	Val	Val	Thr	Pro	Leu	Ser	Pro	Pro	Thr	Asn	Leu	His	Leu	Glu	Ala
			1170				1175					1180			
Asn	Pro	Asp	Thr	Gly	Val	Leu	Thr	Val	Ser	Trp	Glu	Arg	Ser	Thr	Thr
			1185				1190					1195			1200
Pro	Asp	Ile	Thr	Gly	Tyr	Arg	Ile	Thr	Thr	Thr	Pro	Thr	Asn	Gly	Gln
				1205					1210					1215	
Gln	Gly	Asn	Ser	Leu	Glu	Glu	Val	Val	His	Ala	Asp	Gln	Ser	Ser	Cys
			1220						1225					1230	
Thr	Phe	Asp	Asn	Leu	Ser	Pro	Gly	Leu	Glu	Tyr	Asn	Val	Ser	Val	Tyr
			1235					1240					1245		
Thr	Val	Lys	Asp	Asp	Lys	Glu	Ser	Val	Pro	Ile	Ser	Asp	Thr	Ile	Ile
			1250				1255					1260			
Pro	Ala	Val	Pro	Pro	Pro	Thr	Asp	Leu	Arg	Phe	Thr	Asn	Ile	Gly	Pro
			1265				1270					1275			1280
Asp	Thr	Met	Arg	Val	Thr	Trp	Ala	Pro	Pro	Pro	Ser	Ile	Asp	Leu	Thr
				1285					1290					1295	

-continued

Asn Phe Leu Val Arg Tyr Ser Pro Val Lys Asn Glu Glu Asp Val Ala
 1300 1305 1310

Glu Leu Ser Ile Ser Pro Ser Asp Asn Ala Val Val Leu Thr Asn Leu
 1315 1320 1325

Leu Pro Gly Thr Glu Tyr Val Val Ser Val Ser Ser Val Tyr Glu Gln
 1330 1335 1340

His Glu Ser Thr Pro Leu Arg Gly Arg Gln Lys Thr Gly Leu Asp Ser
 1345 1350 1355 1360

Pro Thr Gly Ile Asp Phe Ser Asp Ile Thr Ala Asn Ser Phe Thr Val
 1365 1370 1375

His Trp Ile Ala Pro Arg Ala Thr Ile Thr Gly Tyr Arg Ile Arg His
 1380 1385 1390

His Pro Glu His Phe Ser Gly Arg Pro Arg Glu Asp Arg Val Pro His
 1395 1400 1405

Ser Arg Asn Ser Ile Thr Leu Thr Asn Leu Thr Pro Gly Thr Glu Tyr
 1410 1415 1420

Val Val Ser Ile Val Ala Leu Asn Gly Arg Glu Glu Ser Pro Leu Leu
 1425 1430 1435 1440

Ile Gly Gln Gln Ser Thr Val Ser Asp Val Pro Arg Asp Leu Glu Val
 1445 1450 1455

Val Ala Ala Thr Pro Thr Ser Leu Leu Ile Ser Trp Asp Ala Pro Ala
 1460 1465 1470

Val Thr Val Arg Tyr Tyr Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn
 1475 1480 1485

Ser Pro Val Gln Glu Phe Thr Val Pro Gly Ser Lys Ser Thr Ala Thr
 1490 1495 1500

Ile Ser Gly Leu Lys Pro Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala
 1505 1510 1515 1520

Val Thr Gly Arg Gly Asp Ser Pro Ala Ser Ser Lys Pro Ile Ser Ile
 1525 1530 1535

Asn Tyr Arg Thr Glu Ile Asp Lys Pro Ser Gln Met Gln Val Thr Asp
 1540 1545 1550

Val Gln Asp Asn Ser Ile Ser Val Lys Trp Leu Pro Ser Ser Ser Pro
 1555 1560 1565

Val Thr Gly Tyr Arg Val Thr Thr Pro Lys Asn Gly Pro Gly Pro
 1570 1575 1580

Thr Lys Thr Lys Thr Ala Gly Pro Asp Gln Thr Glu Met Thr Ile Glu
 1585 1590 1595 1600

Gly Leu Gln Pro Thr Val Glu Tyr Val Val Ser Val Tyr Ala Gln Asn
 1605 1610 1615

Pro Ser Gly Glu Ser Gln Pro Leu Val Gln Thr Ala Val Thr Thr Ile
 1620 1625 1630

Pro Ala Pro Thr Asp Leu Lys Phe Thr Gln Val Thr Pro Thr Ser Leu
 1635 1640 1645

Ser Ala Gln Trp Thr Pro Pro Asn Val Gln Leu Thr Gly Tyr Arg Val
 1650 1655 1660

Arg Val Thr Pro Lys Glu Lys Thr Gly Pro Met Lys Glu Ile Asn Leu
 1665 1670 1675 1680

Ala Pro Asp Ser Ser Ser Val Val Val Ser Gly Leu Met Val Ala Thr
 1685 1690 1695

-continued

Lys Tyr Glu Val Ser Val Tyr Ala Leu Lys Asp Thr Leu Thr Ser Arg
 1700 1705 1710

Pro Ala Gln Gly Val Val Thr Thr Leu Glu Asn Val Ser Pro Pro Arg
 1715 1720 1725

Arg Ala Arg Val Thr Asp Ala Thr Glu Thr Thr Ile Thr Ile Ser Trp
 1730 1735 1740

Arg Thr Lys Thr Glu Thr Ile Thr Gly Phe Gln Val Asp Ala Val Pro
 1745 1750 1755 1760

Ala Asn Gly Gln Thr Pro Ile Gln Arg Thr Ile Lys Pro Asp Val Arg
 1765 1770 1775

Ser Tyr Thr Ile Thr Gly Leu Gln Pro Gly Thr Asp Tyr Lys Ile Tyr
 1780 1785 1790

Leu Tyr Thr Leu Asn Asp Asn Ala Arg Ser Ser Pro Val Val Ile Asp
 1795 1800 1805

Ala Ser Thr Ala Ile Asp Ala Pro Ser Asn Leu Arg Phe Leu Ala Thr
 1810 1815 1820

Thr Pro Asn Ser Leu Leu Val Ser Trp Gln Pro Pro Arg Ala Arg Ile
 1825 1830 1835 1840

Thr Gly Tyr Ile Ile Lys Tyr Glu Lys Pro Gly Ser Pro Pro Arg Glu
 1845 1850 1855

Val Val Pro Arg Pro Arg Pro Gly Val Thr Glu Ala Thr Ile Thr Gly
 1860 1865 1870

Leu Glu Pro Gly Thr Glu Tyr Thr Ile Tyr Val Ile Ala Leu Lys Asn
 1875 1880 1885

Asn Gln Lys Ser Glu Pro Leu Ile Gly Arg Lys Lys Thr Gly Gln Glu
 1890 1895 1900

Ala Leu Ser Gln Thr Thr Ile Ser Trp Ala Pro Phe Gln Asp Thr Ser
 1905 1910 1915 1920

Glu Tyr Ile Ile Ser Cys His Pro Val Gly Thr Asp Glu Glu Pro Leu
 1925 1930 1935

Gln Phe Arg Val Pro Gly Thr Ser Thr Ser Ala Thr Leu Thr Gly Leu
 1940 1945 1950

Thr Arg Gly Ala Thr Tyr Asn Ile Ile Val Glu Ala Leu Lys Asp Gln
 1955 1960 1965

Gln Arg His Lys Val Arg Glu Glu Val Val Thr Val Gly Asn Ser Val
 1970 1975 1980

Asn Glu Gly Leu Asn Gln Pro Thr Asp Asp Ser Cys Phe Asp Pro Tyr
 1985 1990 1995 2000

Thr Val Ser His Tyr Ala Val Gly Asp Glu Trp Glu Arg Met Ser Glu
 2005 2010 2015

Ser Gly Phe Lys Leu Leu Cys Gln Cys Leu Gly Phe Gly Ser Gly His
 2020 2025 2030

Phe Arg Cys Asp Ser Ser Arg Trp Cys His Asp Asn Gly Val Asn Tyr
 2035 2040 2045

Lys Ile Gly Glu Lys Trp Asp Arg Gln Gly Glu Asn Gly Gln Met Met
 2050 2055 2060

Ser Cys Thr Cys Leu Gly Asn Gly Lys Gly Glu Phe Lys Cys Asp Pro
 2065 2070 2075 2080

His Glu Ala Thr Cys Tyr Asp Asp Gly Lys Thr Tyr His Val Gly Glu
 2085 2090 2095

Gln Trp Gln Lys Glu Tyr Leu Gly Ala Ile Cys Ser Cys Thr Cys Phe

-continued

2100		2105		2110											
Gly	Gly	Gln	Arg	Gly	Trp	Arg	Cys	Asp	Asn	Cys	Arg	Arg	Pro	Gly	Gly
		2115					2120						2125		
Glu	Pro	Ser	Pro	Glu	Gly	Thr	Thr	Gly	Gln	Ser	Tyr	Asn	Gln	Tyr	Ser
		2130					2135					2140			
Gln	Arg	Tyr	His	Gln	Arg	Thr	Asn	Thr	Asn	Val	Asn	Cys	Pro	Ile	Glu
		2145				2150					2155				2160
Cys	Phe	Met	Pro	Leu	Asp	Val	Gln	Ala	Asp	Arg	Glu	Asp	Ser	Arg	Glu
				2165						2170					2175

<210> SEQ ID NO 26
 <211> LENGTH: 2402
 <212> TYPE: DNA
 <213> ORGANISM: Homo Sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Oncofetal Fibronectin variant 7
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank NM 054034
 <309> DATABASE ENTRY DATE: 2005-06-10
 <400> SEQUENCE: 26

```

gcccgccgccc gctgtgctgc acagggggag gagaggaac cccagggcgcg agcgggaaga    60
ggggacctgc agccacaact tctctggtcc tctgcatccc ttctgtccct ccaccctcc    120
ccttccccac cctctggccc ccaccttctt ggaggcgaca acccccggga ggcattagaa    180
gggatttttc ccgcaggttg cgaaggaag caaacttggg ggcaacttgc ctcccgggtg    240
gggctctctt cccccaccgt ctcaacatgc ttaggggtcc ggggccggg ctgctgctgc    300
tggcctcca gtgcctgggg acagcgggtg cctccacggg agcctcgaag agcaagaggc    360
aggctcagca aatggttcag cccagctccc cgggtgctgt cagtcaaagc aagcccgtt    420
gttatgacaa tggaaaacac tatcagataa atcaacagtg ggagcggacc tacctaggca    480
atgctgttgt ttgtacttgt tatggaggaa gccgaggtt taactgcgag agtaaacctg    540
aagctgaaga gacttctttt gacaagtaca ctgggaacac ttaccgagtg ggtgacactt    600
atgagcgtcc taaagactcc atgatctggg actgtacctg catcggggct gggcgaggga    660
gaataagctg taccatcgca aaccgctgcc atgaaggggg tcagtcctac aagattggtg    720
acacctggag gagaccacat gagactggtg gttacatggt agagtgtgtg tgtcttgta    780
atggaaaagg agaattgacc tgcaagccca tagctgagaa gtgttttgat catgctgctg    840
ggacttccta tgtggtcgga gaaacgtggg agaagcccta ccaaggctgg atgatgtag    900
attgtacttg cctgggagaa ggcagcggac gcatcacttg cacttctaga aatagatgca    960
acgatcagga cacaaggaca tcctatagaa ttggagacac ctggagcaag aaggataatc   1020
gaggaaacct gctccagtgc atctgcacag gcaacggccg aggagagtgg aagtgtgaga   1080
ggcacacctc tgtgcagacc acatcgagcg gatctggccc cttcaccgat gttcgtgcag   1140
ctgtttacca accgcagcct cccccccagc ctccctcccta tggccactgt gtcacagaca   1200
gtggtgtggt ctactctgtg gggatgcagt ggctgaagac acaaggaaat aagcaaatgc   1260
tttgacagtg cctgggcaac ggagtcagct gccaaagagac agctgtaacc cagacttacg   1320
gtggcaactc aaatggagag ccatgtgtct taccattcac ctacaatggc aggacgttct   1380
actcctgcac cacagaaggg cgacaggacg gacatctttg gtgcagcaca acttcgaatt   1440
atgagcagga ccagaataac tctttctgca cagaccacac tgttttggtt cagactcgag   1500
  
```

-continued

```

gaggaaattc caatggtgcc ttgtgccact tccccttccct atacaacaac cacaattaca 1560
ctgattgcaac ttctgagggc agaagagaca acatgaagtg gtgtgggacc acacagaact 1620
atgatgccga ccagaagttt gggttctgcc ccatggctgc ccacgaggaa atctgcacaa 1680
ccaatgaagg ggtcatgtac cgcattggag atcagtggga taagcagcat gacatgggtc 1740
acatgatgag gtgcacgtgt gttgggaatg gtcgtgggga atggacatgc attgcctact 1800
cgcagcttcg agatcagtgc attgttgatg acatcactta caatgtgaac gacacattcc 1860
acaagcgtca tgaagagggg cacatgctga actgtacatg cttcggtcag ggtcggggca 1920
ggtggaagtg tgatcccgtc gaccaatgcc aggattcaga gactgggacg ttttatcaaa 1980
ttggagattc atgggagaag tatgtgcatg gtgtcagata ccagtgtac tgctatggcc 2040
gtggcattgg ggagtggcat tgccaacctt tacagaccta tccaagctca agtggctctg 2100
tcgaagtatt tatcactgag actccgagtc agcccaactc ccaccccatc cagtggaatg 2160
caccacagcc atctcacatt tccaagtaca ttctcaggtg gagacctgtg agtatccac 2220
ccagaaacct tggatactga gtctcctaatt cttatcaatt ctgatggttt ctttttttcc 2280
cagcttttga gccacaact ctgattaact attcctatag catttactat atttgtttag 2340
tgaacaaaca atatgtggtc aattaaatg acttgtagac tgaaaaaaaaa aaaaaaaaaa 2400
aa 2402

```

<210> SEQ ID NO 27

<211> LENGTH: 657

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<300> PUBLICATION INFORMATION:

<308> DATABASE ACCESSION NUMBER: GenBank NM 054034

<309> DATABASE ENTRY DATE: 2005-06-10

<400> SEQUENCE: 27

```

Met Leu Arg Gly Pro Gly Pro Gly Leu Leu Leu Leu Ala Val Gln Cys
 1                    5                10                15

Leu Gly Thr Ala Val Pro Ser Thr Gly Ala Ser Lys Ser Lys Arg Gln
                20                25                30

Ala Gln Gln Met Val Gln Pro Gln Ser Pro Val Ala Val Ser Gln Ser
                35                40                45

Lys Pro Gly Cys Tyr Asp Asn Gly Lys His Tyr Gln Ile Asn Gln Gln
 50                55                60

Trp Glu Arg Thr Tyr Leu Gly Asn Ala Leu Val Cys Thr Cys Tyr Gly
65                70                75                80

Gly Ser Arg Gly Phe Asn Cys Glu Ser Lys Pro Glu Ala Glu Glu Thr
                85                90                95

Cys Phe Asp Lys Tyr Thr Gly Asn Thr Tyr Arg Val Gly Asp Thr Tyr
                100                105                110

Glu Arg Pro Lys Asp Ser Met Ile Trp Asp Cys Thr Cys Ile Gly Ala
                115                120                125

Gly Arg Gly Arg Ile Ser Cys Thr Ile Ala Asn Arg Cys His Glu Gly
                130                135                140

Gly Gln Ser Tyr Lys Ile Gly Asp Thr Trp Arg Arg Pro His Glu Thr
145                150                155                160

Gly Gly Tyr Met Leu Glu Cys Val Cys Leu Gly Asn Gly Lys Gly Glu
                165                170                175

```


-continued

Trp Thr Cys Lys Pro Ile Ala Glu Lys Cys Phe Asp His Ala Ala Gly
 180 185 190
 Thr Ser Tyr Val Val Gly Glu Thr Trp Glu Lys Pro Tyr Gln Gly Trp
 195 200 205
 Met Met Val Asp Cys Thr Cys Leu Gly Glu Gly Ser Gly Arg Ile Thr
 210 215 220
 Cys Thr Ser Arg Asn Arg Cys Asn Asp Gln Asp Thr Arg Thr Ser Tyr
 225 230 235 240
 Arg Ile Gly Asp Thr Trp Ser Lys Lys Asp Asn Arg Gly Asn Leu Leu
 245 250 255
 Gln Cys Ile Cys Thr Gly Asn Gly Arg Gly Glu Trp Lys Cys Glu Arg
 260 265 270
 His Thr Ser Val Gln Thr Thr Ser Ser Gly Ser Gly Pro Phe Thr Asp
 275 280 285
 Val Arg Ala Ala Val Tyr Gln Pro Gln Pro His Pro Gln Pro Pro Pro
 290 295 300
 Tyr Gly His Cys Val Thr Asp Ser Gly Val Val Tyr Ser Val Gly Met
 305 310 315 320
 Gln Trp Leu Lys Thr Gln Gly Asn Lys Gln Met Leu Cys Thr Cys Leu
 325 330 335
 Gly Asn Gly Val Ser Cys Gln Glu Thr Ala Val Thr Gln Thr Tyr Gly
 340 345 350
 Gly Asn Ser Asn Gly Glu Pro Cys Val Leu Pro Phe Thr Tyr Asn Gly
 355 360 365
 Arg Thr Phe Tyr Ser Cys Thr Thr Glu Gly Arg Gln Asp Gly His Leu
 370 375 380
 Trp Cys Ser Thr Thr Ser Asn Tyr Glu Gln Asp Gln Lys Tyr Ser Phe
 385 390 395 400
 Cys Thr Asp His Thr Val Leu Val Gln Thr Arg Gly Gly Asn Ser Asn
 405 410 415
 Gly Ala Leu Cys His Phe Pro Phe Leu Tyr Asn Asn His Asn Tyr Thr
 420 425 430
 Asp Cys Thr Ser Glu Gly Arg Arg Asp Asn Met Lys Trp Cys Gly Thr
 435 440 445
 Thr Gln Asn Tyr Asp Ala Asp Gln Lys Phe Gly Phe Cys Pro Met Ala
 450 455 460
 Ala His Glu Glu Ile Cys Thr Thr Asn Glu Gly Val Met Tyr Arg Ile
 465 470 475 480
 Gly Asp Gln Trp Asp Lys Gln His Asp Met Gly His Met Met Arg Cys
 485 490 495
 Thr Cys Val Gly Asn Gly Arg Gly Glu Trp Thr Cys Ile Ala Tyr Ser
 500 505 510
 Gln Leu Arg Asp Gln Cys Ile Val Asp Asp Ile Thr Tyr Asn Val Asn
 515 520 525
 Asp Thr Phe His Lys Arg His Glu Glu Gly His Met Leu Asn Cys Thr
 530 535 540
 Cys Phe Gly Gln Gly Arg Gly Arg Trp Lys Cys Asp Pro Val Asp Gln
 545 550 555 560
 Cys Gln Asp Ser Glu Thr Gly Thr Phe Tyr Gln Ile Gly Asp Ser Trp
 565 570 575

-continued

Glu Lys Tyr Val His Gly Val Arg Tyr Gln Cys Tyr Cys Tyr Gly Arg
 580 585 590

Gly Ile Gly Glu Trp His Cys Gln Pro Leu Gln Thr Tyr Pro Ser Ser
 595 600 605

Ser Gly Pro Val Glu Val Phe Ile Thr Glu Thr Pro Ser Gln Pro Asn
 610 615 620

Ser His Pro Ile Gln Trp Asn Ala Pro Gln Pro Ser His Ile Ser Lys
 625 630 635 640

Tyr Ile Leu Arg Trp Arg Pro Val Ser Ile Pro Pro Arg Asn Leu Gly
 645 650 655

Tyr

<210> SEQ ID NO 28
 <211> LENGTH: 360
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Fibronectin IIIICS region V120

<400> SEQUENCE: 28

```
gacgagcttc cccaactggt aacccttcca caccccaatc ttcattggacc agagatcttg      60
gatgttcctt ccacagttca aaagaccctt ttcgtcacc accctgggta tgacactgga     120
aatggtattc agcttctctg cacttctggt cagcaacca gtgttgggca acaaatgatc     180
tttgaggaac atggttttag gcggaccaca cggccacaaa cgccacccc cataaggcat     240
aggccaagac catacccgcc gaatgtaggt gaggaatcc aaattggta cattcccagg     300
gaagatgtag actatcacct gtaccacac  ggtccgggac tcaatccaaa tgcctctaca     360
```

<210> SEQ ID NO 29
 <211> LENGTH: 120
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

Asp Glu Leu Pro Gln Leu Val Thr Leu Pro His Pro Asn Leu His Gly
 1 5 10 15

Pro Glu Ile Leu Asp Val Pro Ser Thr Val Gln Lys Thr Pro Phe Val
 20 25 30

Thr His Pro Gly Tyr Asp Thr Gly Asn Gly Ile Gln Leu Pro Gly Thr
 35 40 45

Ser Gly Gln Gln Pro Ser Val Gly Gln Gln Met Ile Phe Glu Glu His
 50 55 60

Gly Phe Arg Arg Thr Thr Pro Pro Thr Thr Ala Thr Pro Ile Arg His
 65 70 75 80

Arg Pro Arg Pro Tyr Pro Pro Asn Val Gly Glu Glu Ile Gln Ile Gly
 85 90 95

His Ile Pro Arg Glu Asp Val Asp Tyr His Leu Tyr Pro His Gly Pro
 100 105 110

Gly Leu Asn Pro Asn Ala Ser Thr
 115 120

<210> SEQ ID NO 30
 <211> LENGTH: 285
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Fibronectin IIIICS region V95

<400> SEQUENCE: 30

```

acagttcaaa agaccctttt cgtcaccac cctgggtatg aactggaaa tggatttcag    60
cttctggca cttctggta gcaaccagt gttgggcaac aaatgatctt tgaggaacat    120
ggttttaggc ggaccacacc gccacacaag gccaccccca taaggcatag gccaagacca    180
taccgcgca atgtaggtga ggaaatccaa attggtcaca tcccaggga agatgtagac    240
tatcaactgt acccacacgg tccgggactc aatccaaatg cctct                    285

```

<210> SEQ ID NO 31

<211> LENGTH: 95

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

```

Thr Val Gln Lys Thr Pro Phe Val Thr His Pro Gly Tyr Asp Thr Gly
  1          5          10
Asn Gly Ile Gln Leu Pro Gly Thr Ser Gly Gln Gln Pro Ser Val Gly
      20          25          30
Gln Gln Met Ile Phe Glu Glu His Gly Phe Arg Arg Thr Thr Pro Pro
      35          40          45
Thr Thr Ala Thr Pro Ile Arg His Arg Pro Arg Pro Tyr Pro Pro Asn
  50          55          60
Val Gly Glu Glu Ile Gln Ile Gly His Ile Pro Arg Glu Asp Val Asp
  65          70          75          80
Tyr His Leu Tyr Pro His Gly Pro Gly Leu Asn Pro Asn Ala Ser
      85          90          95

```

<210> SEQ ID NO 32

<211> LENGTH: 267

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<223> OTHER INFORMATION: Fibronectin IIIICS region V89

<400> SEQUENCE: 32

```

gacgagcttc cccaactggt aacccttcca caccccaatc ttcattggacc agagatcttg    60
gatgttcctt ccacagtcca aaagaccctt ttcgtcacc accctgggta tgacactgga    120
aatggtatc agcttctctg cacttctggt cagcaacca gtgttgggca acaaatgatc    180
tttgaggaac atggttttag gcggaccaca cggccacaaa cggccacccc cataaggcat    240
aggccaagac catacccgcc gaatgta                    267

```

<210> SEQ ID NO 33

<211> LENGTH: 89

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

```

Asp Glu Leu Pro Gln Leu Val Thr Leu Pro His Pro Asn Leu His Gly
  1          5          10          15
Pro Glu Ile Leu Asp Val Pro Ser Thr Val Gln Lys Thr Pro Phe Val
      20          25          30
Thr His Pro Gly Tyr Asp Thr Gly Asn Gly Ile Gln Leu Pro Gly Thr

```

-continued

35	40	45	
Ser Gly Gln Gln Pro	Ser Val Gly Gln Gln Met	Ile Phe Glu Glu His	
50	55	60	
Gly Phe Arg Arg Thr	Thr Pro Pro Thr Thr Ala	Thr Pro Ile Arg His	
65	70	75	80
Arg Pro Arg Pro Tyr	Pro Pro Asn Val		
85			

<210> SEQ ID NO 34
 <211> LENGTH: 192
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Fibronectin IIIICS region V64

<400> SEQUENCE: 34

```
acagttcaaa agacccttt cgtcaccac cctgggtatg aactggaaa tggatttcag      60
cttctggca cttctggta gcaaccagt gttgggcaac aatgatctt tgaggaacat      120
ggttttaggc ggaccacacc gccacacaag gccaccccca taaggcatag gccaagacca      180
taccgccga at                                                              192
```

<210> SEQ ID NO 35
 <211> LENGTH: 64
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Fibronectin IIIICS region V64

<400> SEQUENCE: 35

Thr Val Gln Lys Thr	Pro Phe Val Thr	His Pro Gly Tyr Asp Thr Gly	
1	5	10	15
Asn Gly Ile Gln Leu	Pro Gly Thr Ser Gly Gln Gln	Pro Ser Val Gly	
20	25	30	
Gln Gln Met Ile Phe	Glu Glu His Gly Phe Arg Arg	Thr Thr Pro Pro	
35	40	45	
Thr Thr Ala Thr Pro	Ile Arg His Arg Pro Arg Pro	Tyr Pro Pro Asn	
50	55	60	

<210> SEQ ID NO 36
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 36

Gly Gly Trp Ser His Trp	
1	5

<210> SEQ ID NO 37
 <211> LENGTH: 143947
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <223> OTHER INFORMATION: Fibronectin
 <300> PUBLICATION INFORMATION:
 <308> DATABASE ACCESSION NUMBER: GenBank AC012462
 <309> DATABASE ENTRY DATE: 2005-04-05

<400> SEQUENCE: 37

```
gaattctcct tttccgttcc caagacatgt gcagctcatc atctggccat tttctcctg      60
```

-continued

acgggccac	ttctctcaa	tcttgtagt	cacaccattg	tcatggcacc	atcctgtagg	120
gggtgggaaa	gtcaggacca	aaaagttatt	tgtatattct	gactcacaag	acagtaggtg	180
aactgggacg	gtcacaggg	tcttcagaac	catgatctgg	aaaaatagca	agttaaagat	240
aaaaaccct	ataatgggc	atctatttta	tttatttatt	tactttttat	ttttttttt	300
ttttttgaga	caaagtctt	ctctgtgcc	cagactggag	tgaagtggg	caatctcggc	360
tcactgtaac	ctccatctcc	tgggttcatt	cagttatcct	gcctcagcct	cctgagtagc	420
tgggattaca	ggcactcccc	acagtgcct	ttttttttt	ttttttttt	ttgtattttt	480
tgtagagatg	gggttttacc	atgttgggta	ggctggcttc	aaactcgtga	gctcaggtga	540
tccaccgcgc	tcagcctcca	aaagtgcctg	gattacaggt	gtgagccacc	acacctggcc	600
aatgggcat	ttctttggt	gaattttaaa	atattatttt	ttatcattta	ccattttcta	660
gggcatttta	agaccaatt	tattctgcca	caatcatgtc	atcagaatag	tcaaatgaaa	720
tgactttcat	ttgaattctc	actattaaga	tttaaaattg	tggaaaacta	aagtgggatt	780
ggagtagact	gttaggaata	gatcctagat	gaaaaatcac	ttttggattt	gcagaaacct	840
tggatacaaa	gtagtccaat	taaccataaa	acaaggatca	gctgatctct	acacgccacc	900
acctgtttca	tgtacatggg	aagagtaaaa	aatgaacccg	gataaaataa	aacaaaagca	960
aacaaaatgc	taaatcattg	gttattatcc	acatcaaata	agtctgggtc	tgtggaatat	1020
ctaaaagtca	cagttttatg	cctttaacta	ctatacataa	gggatgactt	tttaacctcc	1080
agggcttata	caacaaaaca	cacctcagaa	gcttatataa	caatatacta	ctttttccat	1140
tttatcaaca	attcagcctg	ccttaagcta	caaagtaaaa	taattagaca	actgtgatat	1200
caaaacaaag	attatgtaag	ttgagccaaa	gagtcacata	aatgaacttg	acatccccgc	1260
tcccacaatt	gcacctaagt	gctcaaacag	gtaataaaca	tacaaactta	aacgtagttc	1320
gatttcagga	gttaaaaacc	atctacttga	agtcaaatat	taaaatttcc	agctcctaaa	1380
cagcatgtga	tgtttttatg	gtttttcatt	ttcagatggg	caaacatgga	agtctatatt	1440
gaaaacactg	gctggctttt	acaacttcag	tctggattaa	atgatttcaa	gtatacaagt	1500
taagtaaaat	actttgaaaa	tctcttttac	caagtcttat	attctaattc	cacaggatat	1560
gatttctgta	tgtctatag	aggcttattt	ataggcataa	cggtcaacac	ctataaaaat	1620
gaaccaattg	tcaataatca	tcacatctcc	taaacacaaa	gaataaatga	aaggtttata	1680
aattctaate	ctaatctcaa	agcataatag	atatttctca	ggcacatttt	aacttaaaaa	1740
atttgaaaat	taaaacgtat	tagaatttga	aatcaaaaca	tttatcaatt	caagttgatt	1800
tttaatgatc	taaaagcaga	ttatctgttt	tttctttcat	attctagca	acgttcaagc	1860
tacctgggct	atgtgtgtga	cgcaaaagtg	tgtgcatggg	tactatgata	tgtgcatggt	1920
tggtttatgt	tgaaggaggc	tgtttttgag	tctaaatggt	cacctctgat	ctcattcaac	1980
actgaagatt	aaaagagatt	tgagggtttt	ctgtgtctct	aatactctta	aatagacctg	2040
gttccaaaag	tctttatcca	tttgttttca	cagaaaatth	caaaaatcaa	attagcacca	2100
taatcttatg	tgtaattaat	agttgtattg	gcatacaacc	cttgaaatga	agcaatgtcc	2160
aacaagtaaa	atctccatc	atctttctat	gatgcaaaat	attacttoga	gtcaaacagt	2220
attctcttgt	ttgcttcatt	tttgaagaaa	cctgtgtctt	ccaacatgc	ttccttggca	2280
catgagtgca	tgcattgaa	ttgaggagac	aaacacggaa	gtggatggac	aaagcaacta	2340

-continued

ctcactagat gaatcacatc tgaaatgacc acttccaaag cctaagcact ggcacaacag	2400
tttaaagcct gattcagaca ttcgttccca ctcatctcca acggcataat gggaaactgt	2460
gtaggggtca aagcacgagt catccgtagg ttggttcaag ccttcgttga ctatgaagaa	2520
aaggaagaaa aagcaaaaag agacatctta ttaatcgatt tggaccataa gaagaaaatc	2580
gaatgactgt atacaatgac ttaatctaaa acaagaattc caggaggatt aagaggcatt	2640
catctgttct atcaaggcat taactgctct aaaaaacat ggtagatgt gacacgtgc	2700
acaggcacc gacaggaagc ccatcttcta tttttccctt gcttctaaga taattgccat	2760
ttctgttgaa actacttcat agaacatgga acagatcttg aggcaattga agctagtgga	2820
gaactttagt gggatgcaaa gactaatgat gcctctccat tggtagctga gtgacctgg	2880
gccagctatt taataccact ctgccttgct tccctcatct gcaatactag cgtaataata	2940
aaaaatctta tacctctctg ggaatcttgt gaagattata taaggccatg aattcaaac	3000
attgaccctt ttccaagact tagaaaatgt tcaactaaata aagaataatt attaaaggct	3060
gtcttttctt tataatgcaa aagctcttagc attttaatat accattataa tgtaattttc	3120
acttttatta ttgtaatttt tgcataatag tgcagaaagc ctgatttttg gacattgtta	3180
ttatagtggg gagacagctt tatcagttag gtcaactggt ccaataaaga aaatgccctt	3240
ggaatataga tgatttagga ataagtaagg gagtctcttc atgttaccta agatcttatc	3300
tctttatgat ttgaaaagc ttattttaa aaaatctatg caaacaagc atgcttcaa	3360
ttattcccta aaaattagat aaactaattt gacttaaat cccactgtgt atttttcttt	3420
ttacagtgca aaactcaaaa gctcaaaaagc agtagatatt tgagggcctt aatttttaa	3480
agggtgcacc cttaaagaaa atagcgtttt ggagaactgg taagtctgag acgtctcaa	3540
gctgggagaa cctggttaga atgtagaata ctgctgctgg gaataaggag tttaaaagaa	3600
aaatggaag aacttaaat gttactaggg gtttctatac tcaaaagagc caggagtca	3660
aattccatcc taactaaagt taaagagatg gaatctttaa aaaaaaaaa aaaaaatgac	3720
ttagaaaagc acaaagacct ttttgcctt gaaaattctg ttcacgtggc attctactag	3780
taaaaccaca aaaggcaaag ataagcatta ctgtggatat gtaatacaac aaacacaaaa	3840
taatctctc ccaacctctc tccagccaaa tgctcaactg cttaattcac aggaatttat	3900
acatgtaag tagcacagtc ttctatcaga tgaggggaagg ggctaaaatt aaacaactgg	3960
ccttaggaat aaaatctgtc acttggcata gacaggactt agcactctct gttggtgtgg	4020
agtagagaac tctcctgttg gaagattggg gattcacggt tttgattatc ccctatctca	4080
tgtgccttgg tgaattttca gctgctgcaa attgagcagc cagggtggac gctattgatt	4140
taaaaggatt tcccagggtc tacaagaag agttaatgat cttttaacct ctatagaagc	4200
ttttaacctg tatagtca aaaaaaatg gaaataacat tggaaaaaa tggaaataac	4260
attggaataa cattctctgc agacagaaat gcacaatggt ggtagcattt tatatcagga	4320
gcaaaagcca actgtcccca tgtctgcaga aggggtatct tctttatagg taatcgtttc	4380
agttttaat cactcaagag gacagcagca tggtagatg gagagggtct gggaaaagac	4440
atcaggaac ctgtgcctta tatcaactca gatactcaga agtaaggact cagagtgttg	4500
tactaaataa tatttgaaat cccttocaga tctataatca tatcaccatt tgagagtctg	4560
tcaattacgt atgcattttg tgaagacaac aggatattggc aggagggtgg ggagagatta	4620

-continued

atttaataata	aagttgctca	tttatttcat	gttgatgtgt	tgcatataat	cccaaagata	4680
aaaagacaat	ggcaacaaag	gtacagtcaa	cagaaaagcc	catctctggt	gtacagcaga	4740
ggggagcctg	tgtctaaata	gtacgtgttt	acataccaga	gttgccacg	gtaacaacct	4800
cttcccgaac	cttatgcctc	tgctggctct	tcagtgcctc	cactatgacg	ttgtagggtg	4860
cacctctggt	gaggcctgtc	agagtggcac	tggtagaagt	tccaggaacc	ctgaactgca	4920
attatcggta	catccaaagc	agagagaaag	cattatagtg	aggaaatgaca	cgggctctcc	4980
tcttaccaat	aaccctcact	gtttaacaa	agcaaaggaa	gacaaaaaac	aaaaaacaaa	5040
aaaaaaaaaa	agaggggagg	tatagtggtc	ctaaccagct	gaggagagaa	atggaacaag	5100
gataggggcc	atcgttcctg	ccagattttc	ctgccacaa	tctacctacg	tgctccgtag	5160
ttaacaggag	ccctcgacaa	tactagagac	ggcctgccat	tgatcttgct	tcagtcgcag	5220
gtctgcataa	gccctctgga	agccctggcc	atgtttgctt	cacataaaac	ctggacactg	5280
cgctatatgg	gagacacggt	catgctcctc	caggtgacat	ctgaccccag	tgtggagtcc	5340
ttaaaacagt	tgggcaaatc	ctcaaacctc	tgtttctaag	ctcataaatt	tcctacagtg	5400
ttgtcagttt	ttaaattctg	ccactttcta	aaagtaacaa	cacaataaaa	attgatactg	5460
tcagtgggcc	gggtgtagtg	ggtggctcat	gcttgtagtc	ccagcacttt	gggaggccaa	5520
ggcgggcgga	tcacgaggtc	aggagatcga	gaccatcctg	gccaacatcg	tgaacccccg	5580
tctctactaa	aaatacaaaa	attagctggg	tgcggtggca	catgcctgta	atcccagcta	5640
tttgggaggc	tgaggcagga	gaattgcttg	aacctgggag	gcagaggttg	cagtgagccg	5700
agatcacgcc	actgcactcc	agccagggcc	atagagcaag	actccatctc	ggggaaaaaa	5760
aaaaatggat	actgtcaaca	caaattaaat	aatatacgtt	ttagagaatc	aatcacttat	5820
cagctgctat	gcacagcttc	gcctgttttt	ttttttcttt	agatatgttg	tagatagtgg	5880
ggttcttcaa	ggacaatcgc	taaaggtagt	gttttagact	tctgcacaca	aatggaaatt	5940
caggtagaat	atctttcttt	tctagaatca	tctatcttac	tcaaaaagga	acctcaaatg	6000
tgtgagaata	agagaaatag	tttcattttc	tctatgaaat	aaaagctagt	gactctctg	6060
tatgtcctcc	aaagttaggc	catattggga	tttaaaaaaa	aaaaaaagca	gggcttaaat	6120
gtgccatctt	ctaagaggtt	ctaaaagtgg	agccactttt	tttttttttt	ttttttcaag	6180
agacgagggt	tcacatggtt	ggccaggctg	gtcttgaact	cctaacctca	aatgatccac	6240
ctgcctcggc	ctccc aaagt	gctgggatta	caggcgtgag	ccatcacacc	cggccatgaa	6300
gccacttttt	attcgagggc	tggtcacttc	agttacctaa	cttattcctt	tctgtgctgc	6360
cccatgagaa	gtgaagagaa	caattaatta	cctgtaaggg	ttcttcatca	gtgccaacag	6420
gatgacatga	aatgatgtac	tcagaagtgt	cctggaatgg	ggcccatgag	atggttgtct	6480
gagagagagc	ttcttgcctc	gtagaggcat	ttggattgag	tcccgaaccg	tgtgggtaca	6540
ggtgatagtc	tacatcttcc	ctggggatgt	gaccaatttg	gatttcctca	cctacattcg	6600
gcgggtatgg	tcttggccta	tgccctatgg	gggtggccgt	tgtgggcggg	gtgggtccgc	6660
taaaaccatg	ttctctaaag	atcatttggt	goccaacact	gggttgctga	ccagaagtgc	6720
caggaagctg	aataccattt	ccagtgtcat	acccagggtg	ggtgacgaaa	ggggtctttt	6780
gaactgtgga	aggaacatcc	aagatctctg	gtccatgaag	attgggggtg	ggaagggtta	6840
ccagttgggg	aagctcgtct	agccgagaga	ggttagagcc	aaaaaagcaa	agcgcattaa	6900

-continued

gctccaggaa	gtagcagcaa	tgataataat	cgattcaggc	aacaatgact	gttcatcaat	6960
ttgataaaa	ccaccagaaa	atgagcagtt	ctgagatcac	aaaacacttt	ttccagaaa	7020
aaatccttag	gatttttctc	aaaggtttct	ttatagttta	gaaagaaaa	agtgttagta	7080
tgaatttatac	atacacacag	ctttaaacgg	ctgagccccg	tagaagctgc	ctctgctcaa	7140
attagaccta	atggaactga	aatgtgcagt	ctttctcttt	ctctctttct	tttttaaatt	7200
tgagaagcaa	gttgtaggaa	gtagaatcca	gactaatcat	atcagtgta	taagtatgag	7260
aattacagag	gtttggctca	acagtttttc	atggtttgaa	atggtaaaa	tagataaaac	7320
acacatgttc	atttttgaaa	ttttcctata	tcaggaatac	acattttctt	gaaattaact	7380
gaaactggat	gcaaaagtaa	gacattatth	agctctaagt	tgtagaaacc	aaataaagat	7440
aaatgtgaga	ggtttctctc	cctcctttaa	aaaaggaac	attttaaaat	tcaacagaag	7500
gtataaaaga	agagcaatga	tagctcttct	tttacatth	aacatcttgt	ttgaattttt	7560
aaaaggtgaa	taacaaggct	ttatattgag	tggctgtagt	aaagaaaaa	aataaaacca	7620
aactctgcag	tgcattgtaa	attatthtct	ctattaaaga	atacaatata	tacactatgc	7680
tgttagataa	aaaaaatcac	aagaaatgca	tcaaaacatg	gagaaccttt	tcatattgca	7740
agattgtctca	tttgttatta	gttttgcct	atctttctcc	ttgttacctg	caagatactc	7800
ttactgtct	ttttccttcc	aatcaggggc	tcgctcttct	gattattctt	cagggcaatg	7860
acataaattg	tattattcgt	tcccggttcc	aggcctgaag	ggagaataga	accatcacat	7920
tatgtcaatg	ggctcagcta	gtcaagtgga	agtcggctct	accagcagac	gctactggga	7980
gcaggcactt	cctctccagc	tggctgttgg	cctcttgaag	gtaaaatcga	cttcaacttt	8040
cctccataaa	cacccaaata	tcctgcata	aaagtccgca	tcgcaacta	attatcttac	8100
tatcatatct	gacgcctgaa	tagacatth	ttggetaact	ttccagggg	attctttttt	8160
tttttttttt	tttttttttt	gagatggagt	ctggctctgt	caccagggct	ggagtgcagt	8220
gggtctatct	cggtcactg	caagctccgc	ttcccagggt	cagccattct	cctgcctcag	8280
ccaccagagt	agctgggact	acaggcacc	atcaccacgc	ctggctaatt	ttgtttttgt	8340
atthtttagta	gagatgggg	ttcacctgt	tagccaggat	ggtctcgatc	tctgacctt	8400
gtgatctgcc	tgctcggcc	tcccaaagt	ctgggattac	aggcgtgagc	caccgcgct	8460
ggcctttcca	gggtattctt	ttaacgtggt	cttatttgcc	ttttgaatt	taagaaaatc	8520
tatcagcctc	atataccacc	actggaatat	aaatttgaaa	gagagctctg	cagattatat	8580
acatgaatct	acttaggcct	aataoccaa	cagtcctcag	tggcagatca	atgaaaagt	8640
aaactaaag	caagtgaag	gtaggagaga	ttggccagt	tttgtgcagc	ctctgtggga	8700
agtgtctagt	tcaattcctc	tcatgtctgc	ttctattht	ctgttcccc	caattctaaa	8760
aagccagtgt	ctcttcttaa	tgtctgtatg	ttctgtggga	atthtaaagt	ccaggtgata	8820
ctctatacag	actgatattg	tttggctctg	tgtctgcac	caaatctcat	ctggaattgt	8880
aatccccatg	tattgaggag	aggaggtgat	tggatcatgg	ggtggttccc	ccaggctggt	8940
ctcatgatag	tgagttctca	caagatctga	tagttttgta	agtatctgga	agttcctcct	9000
gtgtgcttct	ctctcctgct	gccatgtaag	acatgcctgc	ttttccttcc	accatgattg	9060
tacatthtct	gaggcctccc	cagccatgag	gaactgtgag	tcaatthaac	ctctttatag	9120
cagtatgaaa	acagactaat	acagatagtt	taaatatth	atctcttggg	caaaaataat	9180

-continued

actagaggct tatagtcttt cccaccaagg tagacaactc ttattcccca ttccaggcat 9240
tcttccaatg tgggtgggtg ggactgctct gcctcaagac ctatcctgtg atagtgggtca 9300
acaaggtcaa tggcagggtg tgcactggca agtcctgcaa gattgccata tctatagaga 9360
gctgagtgta atgacagaat cacaatcagc tagtatgctt tgaaaaatat ctttctccag 9420
gactgaattg acaggtatct ggcaaaactg cagcgaagg aaaacatgag tctagtcatg 9480
ccattagtac agtcttgaaa ttgcatttct cttcactcat gccttttggg atgtattagg 9540
ctctaccact cctacacaga gaatgctttt tgaggttatc aggaacagag tttcgcattc 9600
tcatgacaca gtatcaaaga cgctgtggga gtttgctaag gacttcatgt gtcctaggtg 9660
gtaagaagga aatgacagc atggaagcag caataccagt aatagtagcc tctgtgacac 9720
cagggcgggg ccgagggacc acttctctgg gaggagacc aggcttctca tacttgatga 9780
tgtagccggt aatcctggca cgtggcggct gccatgatac cagcaaggaa ttgggtgtgg 9840
tggccaggaa acgcagggtg gatgggtgat caatggctga aagaaacaa aattaagtct 9900
ctaagaaggc aaataaccaa gaaaattact cccacacttt tctcccgagc cgttctaaac 9960
acacttaaaa gaatctgaga atactgtaa ccgaaacaa gattctctgg aatctataat 10020
acaaaaatat acgcctcaca tttgtttttt gagttcatga aactgattgc atacaagtca 10080
atggcatttc ctcahtagaa ggtatagtta ccagtggagg cgtcgatgac cacaggggag 10140
ctccgagcat tgtcattcaa ggtgtacag tagatcttgt agtcagtgcc tggttgtaaa 10200
cctgggattt gagaagagat gatttttaac agttcttgct ttttactagg agaattctca 10260
agctagcctg caatttttct gggttcttct atcatcccat atttatatag actaccaaag 10320
tgggtccacat ttgaacagta aagtagagca tgctatggat gttactggtt aacaagaaac 10380
atatctcgaa agtgaatcac tgatacagac aaggacactg gtaaagtaaa ttctggaatg 10440
caaaaatgct atgttggaag aggccctctt gcagaatggg ggtcagccac ttattgaggt 10500
tttctcttta ccgtaatcag atggcaata gttgagcctt ggaatcacia ttttaaaaag 10560
ctaaaaaaga cctcagatat tgaatattct tcggttacia ttttggcctt tattaagtgg 10620
tattgtttac ttttatatta actatgcttt ttaatgctaa gatctatata tcaactgtttt 10680
ttatttcttt tagtttttaa cttatgtta caatttttta aataagaaag ggcacaatga 10740
tggaagatca aacttggtga ttgacactta cgcaatttcc atgacaaata ttacacaaaa 10800
tcaggcattc aaaggacaaa actatttctg atcatagcat gaaaaaacg ttctaaaact 10860
gggttatgat gataacactg aaaataaatt ttatcagtgt cagtcgtaga acagtttaata 10920
agcccgttta cattgtgggt atttgttaac aaatgtgggt agtgcaatga gttccctgac 10980
ctgtgatggt gtagcttctg acatctggct tgatggttct ctggattgga gtctggccat 11040
tggctggaac ggcacaaact tggaaagcag tgatcgtctc agtcttggtt ctccagctaa 11100
tgggtgatggt ggtctcagta gcatctgtca cacgagccct tcttgggtggg ctgacatctg 11160
cacaggagca tagctagaga ttagtgcctg cttggctcat ccatgtaaag ttttggattt 11220
ttaaaacggt tccttggtat atatacaatt catccatttc aagaaatata taactagtaa 11280
tgtgtagatt atctcctttg gcattcagag agaaagtca actgtaatta gaaaatagga 11340
aaatatctac ctctcagaca ttttattttg aagagctgaa aagctttata gatactacca 11400
tggttccca gggctggctt tatagtatct acaaaactt caataaatat taaaattatt 11460

-continued

aatgtattat aatgtaaaaa tatgtgtggt aggagttaca ataacatggc cagctaacaa 11520
ttttgtgtgt gtgtgtgtgt atgtgtgtgt gagagagaga gagagtgtgt gtgtgtgtgt 11580
gtgtgtgtgt gtgtctaata ttacggcaag tttttgtat tttctctcaa caggtacaat 11640
gccatccctc ttttataaat gataaaactt tgacacttta cctgactttt gaaaaggcat 11700
acatcaaggt attgatataa cagaagacta gaatctaagc ctgatttggc ccaatgggta 11760
ctgtgctgtg atataatttg gatatttgc cctgtccaaa tctctgttga gttgtaactt 11820
ccaatgctgg aggtggggcc tgggtgagagg tgtttggatc atggggcag atcccttgtg 11880
gcttgggtgc ctctttatga cagtgtgttc tcaagagact gggtcattta aatgtataag 11940
gtcctcccc cccaactctc tctctctctc ggtcttgctt ttgccacagc catgtgatgt 12000
gcctgtctcc ccttggcctt ccaccatgag taagcttccc aaggcctccc tagaagccaa 12060
gaagatacca gcaccatgct tcctgtaaa cctgctggcc tgtgggcaa ttaagcctct 12120
ttcttttaga aattaccag tctccggtat ttctttatag taatgcaaga atggcctaata 12180
atgcattggt aagatatagc agaagccagt ggggtctccc caagcttagc tctatgaaaa 12240
caaattctaa caagtagagc ataaactaga tgtcagtcac agtccataat atcaacatta 12300
taagcttatg aagtgatcc attttatggg tcagtagacc cctttctgaa ggagtataaa 12360
agatgacagt catgctcaa aaaggcaatc aaattgattt tcaaatagg actcattcca 12420
ctgtaaacat ataaactcaa aaaattattt atttatttta tttattttt ttaagagac 12480
agggccttgc tatttggccg ggtaggtctc aaactcctgg cctcaagtga cgtcccggc 12540
tcagcttccc aaagtcttg gattttagac atgagccact gaacccact gatttaccat 12600
tctttaattt ttgtctatac agaaggttg tccatatgaa gacattttgt tacttactct 12660
ccagagtgtg gacaactccc tgagctggtc tgcttgctca agtgcctta agagcataga 12720
cactcacttc atatttggg gccacctaga gaaataagg catggtgagc tttagcaacg 12780
tcctcaactg aaaccaag agtttacta cagcattagg taaatgcaat ggttagaaaa 12840
taaaaaataa acaaaccagc aaaaccaca acctgaaaa ttcctagatt aatttcattc 12900
ctacagtgca ggtgataaat gtatttatca attttaaat gctcttagcg gttaaagctta 12960
agtaagccat agattatata cttaactaa gctggtaaaa aaaatgagaa aagtgggtat 13020
tagcgtattt ttctttttta tgatctgaca ggtgtaaactc ataattttt ccttatcata 13080
gagaggcaaa gagaatgttc cttttgtct ctgaacctc tttctgtagt ttcccctgaa 13140
attggcctta tgtaagagg tacatcagtc ccttaggttc ctgacagttt tctgcatgta 13200
tgtctggaag gattttaaaa tgttttaatg agggtaggct caaggtagac tggtaaaaac 13260
tgtaaaatat tctttaccaa ggtaacagag ctcagaaaat gcataactaa gtcgatttat 13320
gcattcaacc aatggatag taaattgccc cccacaacct tttaaaataa gcacaatata 13380
tctaaaagag ctgcacaaaa tccaagcta tttataaat tctgtcccaa tagtcatctg 13440
gaaaacttag gtcaacataa aggaattccg ttgatataaa attacaataa gattatttga 13500
tgcagaggaa aagaacagtt agatttatta tgatattata ttttcaccac cttagaaact 13560
gtgttagaga tgatctccat ctttattcca atgaacaacg gtcattgctt accataagtc 13620
ctgatacaac cacggatgag ctgtcaggag caaggttgat ttctttcatt ggtccggtct 13680
tctccttggg ggtcaccgac actcgatatc cagtgtgctg aacattgggt ggtgtccact 13740

-continued

gggcgctcag gcttgtgggt gtgacctgag tgaacttcag gtcagttggt gcaggaatag 13800
ctgtcgagat tgtcattggt tagaggttat cttatagtaa atgggggaaa aggaaaataa 13860
agtgagttcc aagaagtaga atggattggt cttgttgggc taggtggggt ccctgggtcaa 13920
gcaaagattc agtacaagaa agtacgtaca gtaagggatg caaaatgaag aaagtgtttg 13980
ctgagaaatg tgtttatata ccaccaattc tcagaatcac tgaaaacatt tgggtaaagt 14040
ccaaaactct aagagatttt aggttttaca agttttgctt tcccttaaaa aaaaaagtta 14100
cattggctta ttggccatta tgatttattt tacttttgta tggaaaaata agcaatggcc 14160
ctttttcttt cttaaaattg ttaattaatt aatttagaaa cggctctctc gtgtcatcca 14220
ggcggcagtg cagtggagtt atcatagctc actgcagcct tgaactcgtg ggctcaagtg 14280
atcctctacc tcagcctctt gagtagctgg gactacaggc atgtgtcacc acacctcgtc 14340
aattttttca tatttttttg tagagacggg gtctcacttc attgccaggc ctgatcatga 14400
actcctggcc tcaatcagtg ctcccggctt agcctcccaa agcactagga ttacatgtgt 14460
gagccactgc acccggcctc ctttttctat tctacataaa gtatccttgt atggataacc 14520
atttcacgca gtattccatc caaaaagag agaataatgt ttttattgtc tctttatttg 14580
gaccctatgg cagatatggt tgctaaattt gacgttctct ctgtaatgct gcgtaagaag 14640
aaaactggct cccaacagaa gcagaaaaga gaagcgaaaa gcaataatac agactgtcac 14700
ctctagggtt tatgctgtca ctaggataca tctttagcat tctactctgc ttcagaaata 14760
aagcacaaga agataaagta ttatggaaag caattgcctt attgattatt tcctagggtg 14820
ctaaatcaca atcatttaag ctaattatac accagcatat gatatactaac cagagaagaa 14880
aggtcttacc cacgtgtgct cagaatcatg ctgctatgaa atgcccgagt ttgggttctc 14940
ttttataagt aagctctgta atgccaaact tcttttcttt atacatagtg agggtaatca 15000
cagggagtca gagcttagga aataatccac aactttagta tcctcttcaa atgtcttgc 15060
ttatggtctt gtgtataagc caggatactc caaaatttct ttaaatcaga cttgtttcct 15120
taaatatcac ctaagtttta ttgtttctta ccattcctct ttaaagagga tctttcctaa 15180
taattctttt tcacccttac acgggaatca atagcccagt gaagcataat tcattcagta 15240
gggcataaag ccgctgtctc catgggcacc tgggtggtgca attaaccata tacctgtgga 15300
ctgggttcca atcaggggct ggctctccat atcatcgtgc aaggcaacca cactgactgt 15360
gtactcagaa cccggtctga gcccttgtag ctctgcagtg tcttcttcaac catcaggtgc 15420
agggatagc tcatggattc catcctcagg gctcagtag gtcaccctgt acctggaaac 15480
ttgcccctgt gggctttccc aagcaatttt gatggaatcg acatccacat cagtgaatgc 15540
cagtccttta gggcgatcaa tgtctgtag gcaaatatg ggtaagaggt tatgtgaaaa 15600
gcaagttgtg aaataagcat ttttataaca tgcaaagcag ttttgagat accattttct 15660
ttgatgaagt ccaattaacc cactatcaaa aggaaaaatc acttaaacac ttgcttaatt 15720
ttcttgggtt tagaaaaat gcaatgttgc atcacttatt aaaacatagg tgtatgtgag 15780
ataagcagat accctgcaaa atgtcccac aatggggatg tatttatatt aatttaaga 15840
tattctacca cgatttgatt tttacaatgt gatgttagtc tcatagacct ggattttcta 15900
taccttgaa gtggtaaagc tgatttgcaa tgtcagattc aaattgttat aattgtttg 15960
ttttgtttt ttttgagatg gactctcact ctatcaccta ggctggggtg cagaggcatg 16020

-continued

atcttggtc actgcaactt ctgcctcctg ggtccaagt attcttctgc ctgagcctcc 16080
tgagtagctg ggattacagc cacgtgccac cgcgactggc taatTTTTgt atTTTTagta 16140
gagatggggt ttcaccatgt tggtcaggct ggtctcgaac tcctgacctc gtgatccgcc 16200
tgctcagcc tcccaaatg ctgggattac aggcgtgagc ccagccctat aatggttctt 16260
acactcttct tttcccctct ctattaagca tttatattac actctggtcc agccatagcc 16320
cttcaaatgc cccaaacata aagctacatg aggcacactt caatcaactt taccatccag 16380
ttgtagaaaa gacggtctag gaaaatTTTT tgtagatttg ctataggcaa ataagatttc 16440
gctgaggcca tttgagcaaa agaaattcaa agcacagcaa ctaactatgg tggtttcaca 16500
acttctttct ctgtaaattc ctcgcttggc actgagaagg aaaaaagatg ttcgtaaagg 16560
gctcagctca agacataaac taaccaacaa gttcactttc cctatgtgta attgcaaaag 16620
acatgctgctg ggcattcaga cacccaagaa caaaatttga ctttgaaaat ggaaccaag 16680
cagtggttac gtactggtta ctgcagtctg aaccagaggc tgactctctc cgcttgatt 16740
ctgagcatag aactaacca catactccac tgtgggctgc aagcctcaa tagtcatttc 16800
tgtttgatct gcaaaggag tgaagcaaa atgcaacatc cacgtcatcc actataaagt 16860
cctcaagtgg cgtctagagt ttggtagtaa acataggtg gcctagaatt tttatttgca 16920
gaattgtaac aaaaatattg agagtcagac tttgatttcc tcttgacaa acctgttcat 16980
tgctaataca aattaccata tgggagttaa gtcacactca aggataagat ctaaaactta 17040
atatgatgcc cccaacaatt agtgatactg ctctcagaaa aatgaaaat taacttgact 17100
ctaataaatt gccttaagat cctaatttca caagtattct atttaaaac aaagacaaca 17160
tagctaaaaa gagctgaaaa agtcagttag gccatttctc tcttcttagt gtgaatgata 17220
aacaaaacca gaagggtgaa tctagtaatt taagatgtcc atatcactaa atagaactat 17280
ccatgtaaaa acagaaatgc acttaagact tctactgtga aaagggttga taatgatctt 17340
gagccctgaa ctcatgaatg gggacaatga tgacatagga caaaaaata catagtactc 17400
acaacaaggt agccatttat ataaaaaaga atattaggtg gtcctacccc atctatctg 17460
ttgtatatga gatgcagatt actctttttt ttttttttct cctgagatgg agtctctcac 17520
tctgtcacc aggctggagt gcagtggggc tatcttggt cactgcaacc tctgctcca 17580
gacttcaagc cattctcctg cctcagcctc ctgagtagct gggattacag gcacaagcca 17640
ccatgccagc ctaatttttt tgtatttttt agtacagacg gggtttcgcc atattggcca 17700
ggctggtctc gaactcctaa cctcaagtga tctgcccgca tcagcctccc aaagttctg 17760
gattatagga gtgagccacc gcacctggcc gagatgcaga tgattcttac ctggacctgc 17820
agttttagtt tttgttggtc ctggtocatt tttgggagtg gtggttactc tgtaaccagt 17880
aacaggggaa cttgaaggca gccacttgac actaatgctg ttgtcctgaa catoggtcac 17940
ttgcatctgg gatggtttgt caatttctac aaataaaagc agggagaac cagtgaagcc 18000
ccagtccttg gagacaagat tggacaagtc ctcctctcta ggtctgttac atattcgaat 18060
gatogcttac atgagaaagg tatttcttct cgaaagaata cagagagagc ttttagagg 18120
atggggcagc catctcaata ccatttctca tttttgagat ggttgcttca ttcacaaaga 18180
agctacattc cacaagcagc tgtctcttcc tactttatca actcatccat tcttttccca 18240
caaatactta ctgaacactc tctatgctg tcgtattgca gacatgagga atatatcgac 18300

-continued

aaaactgact gaacttctcg cccttggggt gcttccttct ttatctggct caaaactctg 18360
ttcgtctcag gtgctagctg caggttgttg ctcattaaaa aggaaaatct attctacaat 18420
tagaaaaata ccttgggaga acattgcaaa acagtatctg aataagtaaa agctgggtgc 18480
acccagagat agaagtttgt acctgttcgg taattaatgg aaattggcct gctgcttgcg 18540
ggcgtgtctc cacggccagt gacagcatac acagtgatgg tataatcaac tccaggttta 18600
aggccgctga tggtagctgt agacttgctc ccaggcacag tgaactcctg gacagggcta 18660
tttctcctg tatgaaaaag ggtagttca gagtggagg ggtagtagc tacttgggta 18720
ttactgatta attgaattac cacatttata gcagcatgta aatcacatct tcttgccttat 18780
tcccttttaa agagcgcgat cttgaacaga aaggggtatt tatgcatgta gcaaaccttg 18840
gtaagtgtat aaacattgag agaaaattac agttagcgt gttttaacat ttcagcttta 18900
aaatgtcact gagtagtaga gtagcattag aagtgggaga aggtaagccc tcaacctcag 18960
attgcatgca ttgtgcctt tttaaaactt ttaacagaat acattccttt taaataaaat 19020
ttggaaaata cagaaaagta aagagaagaa aaaataaaaa tcacttattt ccaccactga 19080
taatagtgtt gtgaatttac aatcctttta tttcctcagc tctgccgttc cccctctagt 19140
tgtagtataa aaacaacttt ttatttgggt tttagaacaa attatatcat ctatatctta 19200
cgtattttct tggaaaaaac aagggtccg ttgaatggac ttgccaaagt ttctttaatt 19260
ctttgttcac tgttgggctt tcaggttatc cacaacagaa tctgatttaa tcagagtgta 19320
aaatagcatt ttactgctgt acctgtctct ccgtaagtga tcctgtaata tctcactgtg 19380
acagcaggag catcccagct gatcagtagg ctgggtgggg tgcagcaac aacttccagg 19440
tcccctcgaa catcagaaac tagaaaaaaa agggaaactt ttcacatccg taattttcaa 19500
acaatattca gatttactgt ttgttcattt tccagataat tgccttaatg attcttccat 19560
acaatcatgt aaatactacg tgtaattagc agaattttat attatgacga attcaagata 19620
aggggttaca gagcccattc aaattgtgat aaaatgatta tttcaaattt ggggaagagt 19680
ttttttccct caacatttta tttggaaaaa tttcaaaact acagaaaagt taaaaaaaaa 19740
aaaaaaaaaa aaagtgtatc aaggccgggc gtgggtggctc acgctgtaa tcccagcact 19800
ttgggaggcc gaggcgggtg gatcatttga ggtcaggagt tcaaaaccag cctggccaac 19860
atggtaaaat cccatctcta ctaaaaatc aaaaattagc cgggctgggt ggcgggcacc 19920
tgtagtccca gctacttggg aggctgagcc aagagaattg ctcgagcctg ggaggcagag 19980
gttgcatgta gccaaagatc caccattgca ctccagcctg ggcgacagag caagactcca 20040
ttcaggaag aaaaaaaaaa aaaaaaaagt aatacaatga acacctatat acccttcctt 20100
taggtatttg tctctggcct gacgacttga gactaaaaata cagataacat gatgctactg 20160
ttcccgtaag tacttcagcc tctatctgct aagaacaatg aactattac cctaaaatta 20220
tatttttaaa gtacttttga aacagtgttc agttttccac cttttttttt tttttttttg 20280
agacagagtc tccctctgtc accaggctgg agtgcagtga cacaactctt gcttactgca 20340
acctccgctt cctgggttca agcaattctc ctgcctcagc ctctgagta gctgggacta 20400
caggtgcatg ccaccagcc cagctaattt ttgtattttt agtagagacg aggtttcatc 20460
atgttgcca ggatgtctc gatctcttga cctcgtgatc caccgcctt ggctcccaa 20520
agtgtggga ttacaagcgt gagccattgc gctggccct tttttttttt ttttttttga 20580

-continued

gacagggctc tgctctgtca cccaggttgg agtgcaatgg cctgatctca gctcactgca 20640
accttcgcct cccaggttca aagattctcc tgcctcagcc tccccagtag ctgggattac 20700
agggtgatgc caccacacct tgctaatttt tgtatattta gtagagatgg ggtttcacta 20760
tgttgccag gctggtcttg aacttctgac ctcaagtgat ctgcccattt cggcctccca 20820
aagtgtgag attactgacg tgagccaccc cgtccggcca attttctaca ttttcaaca 20880
aacttgtgtt attagaaatg cataaatacc caaatcaggt gagaaatttc ttgagaggaa 20940
aaaaattatc ttctttactg ccaccacgcc cgggcttctt taccatattc taaggaagat 21000
gtttctccac attttctcac ttccctctcc atgtaccatg acaatgatct attttttttt 21060
tttttttttt ttttttgag agctgatgac agacaacagc aagctacttt acagaatcta 21120
ccaactgggt aggaaagtct tctgagtttc tttgcagaca agaaaagtta cctggtgatt 21180
gttgccaat caataagga ctttctctc tgccattaag agcaacgatg ctgaccacat 21240
actctgtgcc tggagtgagg ttggtgagg tgatggaatt ccgagagtgg ggcaccgat 21300
cttctcgagg tctcccactg aagtgtctcg gatgatggcg gatcctgtag ccagtgatgg 21360
tggctcgagg agcaatccag tgcacagtaa aagagttggc agtaatatca gaaaagtcaa 21420
tgccagttgg ggaatcaaga cctgttttcc ccaccgggg gaggaagaga aaaaaaaaaag 21480
aaaaacacc accagtttag gaagtgagga aggtgtaggg gaaattaacg tacatccaac 21540
atctgttcc tgtctcatca ataccatgat ttgccataaa ccaagagta agatgtactg 21600
attctaagct acatatgaat tttaaaatct taaattgact caattcaaaa tagccagctc 21660
aagcattcat tttttcttg tgctaagtaa ttcaagcaga ttttaggggt tattacagta 21720
agtttctgtg tcacatttgg agataagtgt tcaagatctt ctgagaattg acattttcaa 21780
attcttcagt gttcaagat tctgtgttcc tatcctgtaa gtgtatcctt ccaagtaaaag 21840
caagatctgc cactctcaa agtaaagtgt tgactctcag agaagcctcc taagctcaag 21900
aggaaattta tatgtcaaca gagtggtaat gatgccaac acaaaacttt ctgaaactaa 21960
agacattatc ccagacaaat tacaatgctg ataattcctc gtgaattgac agttttgaaa 22020
aatgcatttt tcaaaaaatgc acaaaaatc agagtttgtt gcaaaattcaa atcacttagt 22080
ctctgggact tccagcaaat ttctgtttca gttacagcat ttactagttc ttttcaataa 22140
gaaaccaaac taagatatca taagtcttct ttggaagcaa gtggactata aataaattaa 22200
tgaatacact aaagtgtacc ttttgaaaga aaaaattaaa agcattcata ttttaagtatg 22260
taatatttat gagattcaat tcaacttctc tctagtaaat gcatatttta ttaagtaaaa 22320
agaaaatgta aacttgctgg cttagtaaag gaagacaaat agatgacaat gagtgttggg 22380
tcagccatta gacagagctg cctaatagaa agaatgtct gctaaatcta aatacaaaag 22440
atcttcttag aaaatgtttt ttaaaaggac agacaacaaa tttcttaatt tgttagtgac 22500
tgaaagtcta ggtgattgtt gagtaagta tcttttagtt tttgagacca gcgataaaat 22560
catactgtga tgacaagggt acacacacca tgacaacaaa aatggagat gtatttttag 22620
aagtagcatt tcatttgtta tacagaattg ataggcaaaa gctactggat agtcatactg 22680
ccaacaccac tcacctgttt tctgtcttcc tctaagagggt gtgctctcat gttgttcgta 22740
gacactggag aactcacta catattctgt accaggcagg agatctgtag gggcaaatgg 22800
ggcttatttt aaaactctgc tcaaaagcat gagaaactaa agcacgcca ggaactattg 22860

-continued

aactgatgaa atatccaaat gtccagtcct atagcctact tacacctaag tgaacactac 22920
agcgaagtgt tcagtgaaca gaccttgaga aggtctgtaa ttagatgata aggaagggtc 22980
accttgcccc atggcctctc gaagccctcc attccacctg tgagtgtgac tcagctggat 23040
ttcagatggg atgaagttgc tgcaacagct ggatatttct cgcacccacc tgagagtatg 23100
tctctggagg gcacatgata atccttgctg tgggtgctgtg gattgccatt tatgcctcac 23160
aacaaccctg gaaggtagtt tgctcctttt aaaagcaatt tcttgtcata caacgtactg 23220
catatcttta aagtgatggg tttttacata tgtgtgactt gggagacat caccgcattc 23280
aagaaatgaa caaacctatc acctctaaaa atttgattta ctacaattat ctccattttg 23340
taaatgaaaa aacaaaaaac aaaacaaaa aaaactgagg cacagagaag ttatcttctt 23400
taatttactt gccaggatc atactgggat tgaccaggc caggactagt ggtcttaata 23460
attaaccttt actgaaaaa agccacactt agaaagcaat cggttctgcc atattaaaaat 23520
gagagcagaa ttttttttaa aaaattaaat atacatccct gagtatcttg ttccattttt 23580
ttttattttt attatttttt ctttttgacg aggttttact ctgttgccca ggcgggagtg 23640
cagtgggtgt atcttggtc actgaaacct ctgcctcctg ggttcaagtg attctcctgc 23700
ctcagactcc tgagttagct ggattacagg tgcctgccac catgcctggc taatttttgt 23760
atttttaata gagacggggc ttcaccgtgt taaccaggct tggactggaa atcctgacct 23820
tgtgatccac ccgccttgcc ctctcaaatg gctgggatta caggcgtgag ccaactgtgc 23880
cagcttcttc ttccattttg atacacagtc aggcatagct taatggtgga gatacatttt 23940
gataagtgtg tgatttggg attttattgt gtaagcatca tagagtgtac ctacacaaac 24000
ctagatgggc ttgtaatccc agcacttttg gaggccgagg aggtggatc acttgaggtc 24060
aggagtttga aaccagcctg gccaacatgg tgaaacccct cgtctactaa aaatacaaaa 24120
aaatagctg gccgtgggg ctccagcctg taatcccagc tactcgggag gctgaggcag 24180
gagaatcact tgagccttgg tgggtggagg tgcagtgagc ccagatcgtg ccaactgcact 24240
ccatcctggg caacagaggg agactccgtc tcaaaaacaa aacaaaacaa aacaaaacaa 24300
aaaaacctga tggtagagcc ttgacacac ctaagctata agctatgtgg tctagcctat 24360
tgctcctagg ctatgagcct gtgcagcatg ttactgtact gaacactaga ggcaactgta 24420
atataaggta aatatttga tatctaaaag atacagagag tgaggtttct aaatttggaa 24480
ctctactca ctaaaccctc attccctgc acaccattg cccttcagcg tttctataaa 24540
gcccctaca aaagtttga aaagtccag aatggaggac cagatgcctg gattccattt 24600
ccaccatact agctggccag tggcaattta cttaactgct ggggacctcg acggcttcat 24660
gtttatttgt ttgtttattt ttttgagaca gggctcact ctgtcaccca ggtggagtg 24720
cagtggcagc atcatggctt gctgcagcct caacctcccc aggtcaggt gatcctccta 24780
cctcaacttc ctacatagct gagaccacag gtgtgcacca ccatgcccgg ctaatttttg 24840
tattttttgt agagacttgg tctcactgtg ttgcccaggc tggctcgaac ctctgggct 24900
caagcaattg gcccacctca gcctccocaa gtgctgggat tatacaggcg agagccactg 24960
cactggtttg cttcatcttt aaaagaggat aataatgact ttgctaggag ggagaggggc 25020
taccctgagg tcatctgtag cattaggatt ataataaggc cttagaacac atgattagca 25080
tgtgacaggt ctacggctaa gggacagaca ttgtgatgca gggccttga tactgggtga 25140

-continued

tgctgtaatt agtgattcca ggggtgctcta gaacagaagc aacagccctg ggggtggtgga 25200
tatgccaata taaagatatt ctgcccctca gccacaaagt aagaaccaa gattatggtc 25260
acgctgagct gctcttcttt gtgtttcatt atctaaatta tgtttgaaag gtagatgttt 25320
tcagactaat acttggaat gtacctaaat gtaagctatt tttcccttat gtagaatctt 25380
ttgctgttgt tatatatttg actattacag tgaaaataat aaaaagcact gtttatgtga 25440
aacacagcac catcttgag cctactgttt tattctttaa caaagatgt ctattctata 25500
atggttggga ctagaagag acaatttcca taaattgttg ttatttattt aatgtcagtt 25560
tttttctaa gctctcagtc ttctgagaaa tccttctaac agattctttt tgaagtacaa 25620
attaaacttc tgactctctt cacaacaaca ttctaacta gaccacaa tcagtcactt 25680
ttctaagtat tcaaatccta aatcaatcaa ctgctgacac tttttttata aaaatgagaa 25740
taaagcacat atggtgacac tattttcagt tctcattttt ttctctaaag atttagattt 25800
agtcataata tttctcttga tattcactta tttttacgtg cttatttaga agcagaattt 25860
gtttgaagtc atggaacttt cctgcagggtg tttttccact atgatttttt taaagctgta 25920
gatatacaaa gtatattaag atagcaacaa acattacatc tttcaaaatt aatggagctg 25980
tgcatggaat tcagagttha gttttagact tgggtgcagc tctctgtctc ttagctctc 26040
ctacttgaga atttagaatt catttgctat gctctaggtt aatatttatg gaacagatac 26100
attcaactgc ttacttgtha agaccactgc attgtctgaa ggagaaattg acaactctgc 26160
aacatcttcc tcatttttca caggtgagta acgcaccagg aagttggtha aatcaatgga 26220
tgggggtgga gccaggtgga cacgcatggt gtctggacca atggtggtga atcgcaggtc 26280
agtgggagga ggaacagctg gtttcgaaca agaaggaaga ctcagttaat gtaattttaa 26340
aattaagct aacgaagtag ctggaaggtt aaaatcaata tttcatgcat aaaggaaca 26400
gaatcataat catgcaaca gtctaacta agtctatttt aagttatagt gtactactgt 26460
taaaaataaa aactacaca cagtactgtt agggcccctt taaccttttt gccatcattg 26520
atataattaa aaaaaatttt aatctacttt caaatatggt gactttatgt acattattaa 26580
taccactgca gattatttta tcccaggcgg aagaggctta agttgccaca cagcaagag 26640
aaacatagca gtactttaca aagaaataca tgcacacaga caaaaagaca agagggtcct 26700
aaacttcccc ccatgtgaga tatctagagc tacagttcca ttcagtgaat tctacttaca 26760
agtgaacatg ttcatgcaaa ttggttctca gcatgtttct tttcccatgc acagcagaga 26820
acctttaaaa tgttgcatgc ttgtcccag actgtggtta aagaggtgtt tctttactta 26880
gtgacagcaa caaacctag aatgtcccga agtttacttt tctctactct caaaacaatg 26940
cccatcact ctaccctaaa ttttgatggg tttttagcta caaagaagga gcatggtaga 27000
gtttatcaaa gggtagaag aagatcatga tggatcattt tgggtatata tttottataa 27060
ttaattggta ctacgcaat ttggtattcg tgtttggtat ttaagacgaa aaattctagg 27120
tcttcaaat gaatgaacat ttgctatttt atatttaaca gagcagacat ctgtatttct 27180
ctagctttat atctacttgt ggtttaaatg agagtgtact gaagaatact tgcatgtgtt 27240
tttctcaat cagtttagca gatgcatttg attccttacc ccccaatcc accctatctg 27300
aatctttggc caattgctga cttccccat gaaacatcca tatttaaccg gagtaactgg 27360
tggcagcatg caacaccatc tatgtotcaa acgcagaagt tttcaaaatt caccgtttg 27420

-continued

ttgtgtcagt gtagtagggg cactctcgcc gccattaatg agagtgataa cgctgatatc 27480
atagtcaatg cccggctcca gccctgtgac tgtgtagtat cctactgagg agtccacaaa 27540
atcttcaaaa atagggatac cttctcctgc cgcaactact gtgatgcggg acccaataat 27600
ggtggaagag tttagcgggg tccacctcag gccgatgctt gaatcgggta tatcaacaaa 27660
gcttaggtca gtgagttggg gcacctctat tgagttacaa agcaaaggga gggggaggca 27720
aaaggaaaat agagggaaaa aaagaggaaa aaataaagca gtgtatatca ggttactagt 27780
agtaaaatg gtggaatggt aaagcaatga tggtaatatg caatctgttt taaattctct 27840
gaaaagactt cctaagcatc atgaacagtt tgcttcccta gaagaataaa caggaatatt 27900
cttttaggga atatatatat atatatattt tttacatttt gaaaacatcc ttaaagggta 27960
caatttgaac tgttcagatt cctaaaaatc atatggctgt ttaggatgtc aaaaccattc 28020
ttagagccta gacataatat ctgaagtaag taccagcaat gcttttaata attccaaaac 28080
tgttttagta gaaaataagc ttgcatgaag aagggtaaaa aataataaat gggtgataaa 28140
ttgatTTTTT ttctcccata caaaactcat gacaacatca tggccataac gctaattgcat 28200
tatgaatgta tgggtgtgaa tgtgccattc aaaagcacat tcaggctgag gaaagacagg 28260
cctaaggta aggccattgc cactatttta gttcattcat aatcaaaaca tgtaattagc 28320
gtagtaaaa gcattctact gaagagtcca aagggggaca cgactctgctc aatgctttca 28380
ttatgttata acccaatgga caaacagcc tacccttaga caggcctttg caatgttgtc 28440
tttcaagcca caagaagaa caccctgaag gtgtgacttt tacttctttt ttttaaaatc 28500
caattttcaa aaagaaggat ttgaaactgc aagatgaaaa actagatgtg tcaaccaa 28560
gctgtttcag tgctcacata cagggacaaa actattttat aggcagctgt tggaaattga 28620
tatggattgt aaaatcgggc agaaatgtac tgatgggatg catttaccat attactttga 28680
gcctatcagt actgagtaca gacacgtgac agtgttttag tcatacttcc atgtgagcag 28740
acaatacaga gttaaagcca tggctgccac tccattacat actcgtgca acacacagca 28800
tcccagtga gacagacag taaagcagtg ttaacagcaa gcatttgtcc aaatcgtgc 28860
tttgacttct tggttttgga aagtcagct gggagatcgg aattaaatcc caaatatagt 28920
tgacaccctt aagcatctgg ggattaagat gctaataccc actcttattg gaagtgtcac 28980
tctcaggata gcagcttatt tttctattac ctgggatgat ggatcagag ataggacac 29040
tttctgtgc atccttgaca gtgtaaacac tgacattgta ctccaggccg ggactcaggt 29100
tatcaaaagt gcaggagctc tgatcagcat ggaccactt tccaaagaa tttccctgct 29160
ggcgtttgt aggggttgtg gtaattctat aaccagtaat gtctggagaa aaaagaaaag 29220
ggaagtatt gcacagagga tctgtgagcc agagcatatt taactagact ccaatgatg 29280
attcacagg cggaatgcag gacttggcat ttactgagga ctccactgct actgttattc 29340
aatgcttga acaatccttt caggcagtac taagatcctc actttatttt tttattttta 29400
ttttttgaga cagtctccct ctatcaccca ggctggagtg cagcgggtgtg atcttggctc 29460
acttcaacct ccgcaagat cctcaactta aaggaaatct tgaaccaggg gagtattcc 29520
ccaaagcttt cttgttttcc aaatctcatg gtcctatcac tgtaccctgc ttttccttaa 29580
tgaatcattt gaagacagct gaggaatatt cctttctacc tgcctagtat agaatcgtc 29640
tccaccttaa gcacattccc cagagagctc tctaattta aaggctctga aactatggga 29700

-continued

tttataggaa gagaaccaag aaaggagaaa aggtgagaaa caaaagtgct gatacttatt 29760
ctagaaagag tattgattgg gaggttgagg caggcagatc acctgagggt gggagtctga 29820
gaccagcctg gccaacacgg taaaaccctg tctctactaa aaatctaaaa aatagctggg 29880
cttgggtggcg catgtctgta gtcccagcta cttggggagg ctgaggcacg agaatcgctt 29940
gagcccagga gacggagggt gcagtgagtt gagatggcgc cactgcactc caagcctggg 30000
tgacagagtg agactctgtc tcaaaaaaaaa aaacaaaaac aaacaaaaaa agagagtatt 30060
gattgttgaa gtcttcagga atcacagaaa ttgggcagta aaagagtga gagggatgag 30120
agacattcaa ataaaaacag tgatgactag agatctgtgg ggacagatta aatgagaaa 30180
gtatatataa ctttataaaa ggcaaggaga tgacagaata ttttgccatt aggaacaggg 30240
taaacactaa tgtggcaaat acaaggaaac ccaactgaat gagaaataaa tattgagatc 30300
attacacttg ctagtgtgca tccatattgt tgggctgggc cattgagacg agatattaat 30360
gatcctgctg ctaacaagca cgaagattgt caacttaag tttcaaattc agagcaactc 30420
agctggaaat tgtatagctt tggcttaaac aatgtttccg cagataatga aaactttacc 30480
aaaaactact tacttcagag ggagatataa agagatattc tgagctagtt tcattctgtg 30540
tgtttgcata cattaatcag atttagagat gatagcctta gctctgtgcc cggcaaaga 30600
taagaactat ataatacttt tttttaaaca aaaatttcac aagaatttta cagtaaaatt 30660
agaaatagct aaataataac ctaaacatat cccttaaatt aacaagtata tgaggtaaga 30720
atgcaatcaa cattaattgg aacttttatt tttgtttaag atttttctcc ataggtttgt 30780
tgagacttcc atgtggtttt ggcaaaagta atgggtatct aaaatttcct gttattgcta 30840
tagtacgtca tgctgtttga atattgttaa caactatctt tatacatttt agcattttat 30900
aaataatttt caaatatag tgaacaagga atttagacaa atgcttcatt taaagctcta 30960
aaaaatgtca ggaaaccctt taatcagaat tttcagttg catagaatat aattaaaaaa 31020
aaattacatt ccaaataatg gacattgggt gtgagaagga gtctacaaac ttaagttaca 31080
atcccagttt tgtaacattt aaaatgtgtg actatgagta ggccaatttt ttctaagtct 31140
cagttttctt attctgtaaa atgaaggat tctgccttt catcacctt aaatattgt 31200
ggtgcaacc atgacaaaac attcataaaa cttcagggaa taataaactt aggctctgta 31260
acaactcaca aaagggaaat tacgtaatat gactcaacc atttggttg caatttttaa 31320
tgcaactcagc aaaagccacc ttccacagtg cttttcttca gccctacagc tgtgtgagga 31380
tctgtcttgt ccccctagac tagtgtaagg ttgttctgct tcctgtgatt gaagtcaacg 31440
tgcatataag catcctgcag gctgtccat gatgtacatc atgtactgct atagttttta 31500
tggagagcta agcatttgta atttgaatcc cttctttctc tgaagaaac acagtttcta 31560
aaatctttgt cttgggaag tatggtgat acttgccctc tgatccatcc caaacttacc 31620
tgggtgggtg ctectctccc aggagactgt gagcactcca gtgtcagggt ttgctccag 31680
atgcaagttt gttggtggag acaatgctat gcagaaagaa cattttaaaa gtcaaagctg 31740
acacaaaagct cttgacctgt gtaatctttc ctctctccct acctccctct cttccatgct 31800
tcttccctcc ctccctccct cccataaaaa tataggttac attatgaaaa tatgatggac 31860
agcagtaatt ctctgacaaa gacttcttta tgattaatga ttttgcaata ctattgaaat 31920
tattttcatc ttgccagttc ataaatattc ttgacatgga aagaacattt ctgatgccat 31980

-continued

aaaatataaa gcatgaattg aacctgctac tccaaattca ggaatagcat taataatata 32040
agtaaggata agatatctcc cctgtgccat tctcataagt aaaaagtgat attgacacta 32100
gcttttttcg aaaagtagaa cagtttttaa aactaataga aaagggaaaa aattcttctt 32160
acgtgtcacc actttgttta caattggcgc atctctttcc tgtccatctc tcaggacttg 32220
gatggtgtag acgtattcta ctctctggagt caagccggac acaacgatgc ttctctgagtc 32280
tgaagtcact tctcgtggtg cctctcctcc ctggcttggc cgtacacca gctagaggaa 32340
ggaatgcaaa gtaaaccacca aggacaaata tttccagagg actgttactg ctgtgaggct 32400
atgattatgc tttaaatagt gtaaactctt cagaaactgc acagtgtact tttattggga 32460
aatgatttgg ggactctttt atatacttta gctggaaatt ccccaggtg ggggaggat 32520
cttccaaact gcattaaggt ggcttgaat ggaaggttcc agagaaatgt cattacatac 32580
tgttttgctt gataaagaat ctcttgttac aaaccagacg tatacagga ctgtgggtca 32640
catggtagtt catttctatt tttctgtgtg tgtgctagaa ggaatgtctg agctctccct 32700
ttaacagatt attcacatc aattaacctt ttagtgtcca tttaaaggca ccagcccctg 32760
tagcaactgt attgcaaaaa cacacctttt gaagagatat gtacatagct tcaactgcag 32820
gatgatacat tacaagagaa gaattagaat aactatgcc ataatttaa atccttttaa 32880
gattctatct caaaaactta ggtaacactg ttatccaata gtgttcaact ctagagagat 32940
ccttcacaca tcataaggct tctgaaagtc agcatgacat tattaaggga attaaaagg 33000
atactggcac tcttcagaaa gttcttctat atttgagtag cataatcata aatatggcca 33060
tttcaaaact ttatcttttg tgtatttttt taaacgatca aggggattga gtcatgattt 33120
ccagcgataa caaatatttc tctcttcaca ctaaagcca aagaatggaa ttctttttaa 33180
catattcatt tgtattcacc gcttccctaa tatcattact ttaagaaga tgaatccaga 33240
tatccacag gaaaccttaa tcttaacttc tattaagaga ttcaccctca gtgctttgaa 33300
cagttggaga tgaactgat tggatgaaaa tgcgattttt aagtactct gagtttaact 33360
ggtctcccta aactggcttt tgtgaactg ttcacactca tttagaacaa cacatcaaat 33420
aaagttttcc catgtttttt tgaatgtaa ttccctgag gaaaggatag agaaaataag 33480
ataacaataa gcagcagctt cgtgctaatt ttttatcctt tgcaaaacac tttgcagcc 33540
tccatctacc cttccaacaa ttcagtgagg taggcaggga aggactgtta ggccctgaca 33600
atgtagtcaa agatgtatgg acagaagcag gtgacaaaac ccaggctctc tgactcctga 33660
gcacctgtt tccactacat gggaaocaca aggacagctc agggctgtat cacagagatt 33720
aggaacatct gcagtttacc ttaaaccacaa ttcttgagc aggcgtccat gtgatcacia 33780
tgggtgtctc agtcacctg gtgttgaag gtggaataga gctcccaggc tgcactgtga 33840
gagagaatca atgcacatat tcaaaactca aactcacaga tgattgcata gcatatagat 33900
aacaagtatt tttatggaga acctcagttt aaccaagcag tggatagagg atgtaacata 33960
tttttctatt ggggtgagaat tgggagagga tcaggcttct caatctttaa tgtgccacac 34020
acatcatctg gggctctgctt aaaatgcaga ttctgtttct ggaggctcaa caaaaaacta 34080
gggagtttg atttctgaca aacacagata ataccaattc tgctggcccg tgaatgatac 34140
ctgagtaaca aggggttaga tgtttgctaa ggtggtcctt tcattgaaat tatataatct 34200
aattcatgtc tttttatgat tgttgggatt tgactttgat gacaataaat ttccctatat 34260

-continued

cgtcagtgta aaatccctta taaaatgtaa aactggccac cacaaactga ttatcatatt 34320
gtttcttaac cattaattta agaacaatgg taaattatag aaggtgatct aaaaacagct 34380
ttatgctggg cggggatcgg tggctcatga ctgtaatccc agcactttgg gaagccgagg 34440
tgggtggatc acttgaggtc aggagttcga gaccaggctg gcaaacatgg tgaaaccttg 34500
tctctattaa aaatagaaa aattaactgt catggtggct catgcctgta gtcccagctt 34560
cttgggaggc tgagacacaa gaatcgcttg aaccctggag atagaagttg cagtgaagccg 34620
agatcatgcc attgcactcc agcctgggtg acagaatgag actccgtctc aaaaaaagaa 34680
aaaaaaaaag ctttacgctg aattattcca caataatggt cagctatttt tggttgcaat 34740
aaaagatgct atataaaagt taaacaaaat ctgccacat ggatgcaact tgaagtctaa 34800
agatttgtag tttaaatagc tgattattct ttcagaataa tactgcaaaa atttagtatt 34860
ttatcaccag aatcaggaat tatacaaaa ttattttgca tttctagca tctgacatc 34920
actgttattt ttcttattga aaatattact ttggccaggc gcagtggctc atgcctgtaa 34980
tcccagcact ttgggaggct gatgtaggtg aatcacttga ggccaggagt tcaagaccaa 35040
cctggccaac atggtgaaac ctgtctctg ctaaaaatat aaaaattagc caggtgtgat 35100
ggtggatgcc tgtaatccta gctgcttggg aagctgaggc aggagaatca cctgaacctg 35160
ggagatggag gtttcagtga gccaaagatca caccactgca ctccagccag ggtgagagag 35220
caaggctcca tctcaaaaa aaaaaaaaa aaaaaaaga aagaaaataa aattttactt 35280
aaacaatttt ttttttgaga tggactcttg ctctgtcacc caggctggag tgcagtggcg 35340
caatctcatc tcaactgaac ctccagctcc tgggttcaag tgattctcct gcctcagact 35400
cccaagtagc tgggactaca gttgccacc accacatcca gctaattttt gtatttttag 35460
taaagatggg gtttcactat gttggccagg ccagtctcga actcctgacc ttaggtgatc 35520
taccacctc ggctcccaa agtgtggaa ttacaggcat gagccatcac acctggccaa 35580
aaaaaaaaa aatcttattt ttaaataaaa gatttgtttt gggttttagc accaaaaata 35640
agaaagaaag aaagagagag agagtgagag agaaagaaag aaaagaaaga aagaaagaaa 35700
gaaagaaaga aagaaagaaa gaaagaaaga aaggaagaaa gaaaggaagg aaaggaagg 35760
aaaggaagaa aagagagaga gagaaaggaa ggaaggaagg aaagagaaag aaagaaaggg 35820
aaaggaagg aaagaaaaag aaagaaagag gcatgtatat ataaagttaa gatagagatt 35880
tccaacaacc ttgcaaatct tattataaat gtgaaaagg taaatcgagc aaggccagat 35940
attttgagtt gggtaaagtt aagttcaagg gtatcaagtg tgttcatagc tctggtcaca 36000
tgtttgacat ctccacaatt cttcctctct ttacatctca atcaattttt gttccattg 36060
ggaaaaaacc tctgaatgca tttatactaa attatgaatg gctggcaaga caaagaatta 36120
gtctccagag ttatacaaac agggattaaa aagtgttcag actccagtct caactcaact 36180
ccagtcaatt ttttttctt tgccatgtgg gaatttagaa tcaacatcgc aattctcgca 36240
tttttttttc agtctggatt ctaaggagcc attttttatt tttattttta aatattgtg 36300
tgagaaaaag ccaaatatat gtgtgggcca cttcataatg atagctctaa tggotgaatt 36360
gaggtactac tggtttagtg ggctgtttca tctatgtatg ttattatttt gatggtgttt 36420
ggtggggggg ggggggacat gactgggatg ggtgaccgga tgccttttag tacctctgaa 36480
attatacacc acttgactgc tgggtactgt gccaaaatgc catggccagc cctgtgtgac 36540

-continued

catttttaat ggcccacaaa aaacaattca gagaacagtc agtttctttc actgaacatc 36600
tgtgaaaacc acaacttttt ttacaaatta tattcgcttc aactgtcctg ggctttaaga 36660
ctaggtatth tctattgctt gagccagcca cctagagtct tcccaagttt aagatgatgc 36720
tctctgcctg ggaagccat tatgggaaaa tcaccaaacc acaaacccac aaacgatgtc 36780
aaacaaagga tgtgattcta agtgtgagct atgggttctg gcaggctttg ttagggaaga 36840
acagcccagc tgttaaagaa ataaaattg ccctatcaat ataagactaa gctgaattht 36900
ccccagaaac agcttttttc cctaactttt ctcacatttg ctcttaagaa ttagcacia 36960
gtataaaac tcagaaaagt tactcccaga attcagctta gtaagatagt aaaaaactac 37020
agaaatttag taatttcaaa aacatgcaag ccttgggat gtttacagcc ttagggagtg 37080
cttaaagttc aatggagtt gtagaatacc ctgcttcat tagtttattt aaatataagc 37140
ttcatttagc tgtttttatt attattcctt tctcaattat ctattcacca aatagttttt 37200
gaaagcatac aatgttccag acactatgth aacctttatc taataaaatt agttttggct 37260
atcagtactt cagttaaaat cattatattt tataatgaatg tattatgcaa agcataaaat 37320
aatatatcca atatatcagt cttcagaagt acttgaaata tcttggccca ccaccttccc 37380
ttaattattg ttagcatcat taaaatgct tttctttta taagcactgt gttcaacaaa 37440
agaaactaca taaaatattg tctcatgtaa ggaatgaaca agacaactat agaattctac 37500
atthtagtaa aagacgacat ctactgaaag ggcgcattca ctgaaatctc cattctaatt 37560
cactgaagga catttataag aatgaatat ttacaagaaa tataaagaca ttaatttcat 37620
gagagtatag cttagactat cttagaacg tgtaaaatth tagccaaatt ttcaagtaag 37680
cttatttgca aattaaaagt gaattttact tgcttacgat taactattat ttaggaaaaa 37740
gccttcatga gccaaacttc atacagtaaa tttaatgga agtaagtttt tcttaaagta 37800
tgaattctta tggggaaaag tttgtattht ctttgggaaa tcaattccat agttatttac 37860
aaactaggct tgtatggact ttttgaatgc tttttaacat ccatactttt aaattagtctg 37920
aagagtaaca tttgttgaaa gctcttagaa aattgcaaac tagtccatgg ttagaatata 37980
tgctgaaaa gtacttatgt taggtgcttc tataattcaa aatgtcgagt tgaataaac 38040
aatgatgtht cagctagaaa taaggagtct cataattctc tgtaatgtaa atggaaatgt 38100
atthaatagg aatcggtht tgaatagagg ttgttattgt gatgtagata tgtgggagat 38160
ttaagaccac ttcctthtaa atgttgagca aaaggataga catgaccaa aattattgtht 38220
agthttttc tctctgagg aatttcaaga gtaatttaag aaaagagcga ttcogtctgc 38280
aaaagtactt taatcttctg cagccaacac tcataaagaa atcatttctt tggcttcatg 38340
ttcaattagt tggthccat tcaatocata gataththt tctgcattt atagthtacta 38400
ttcaatgccc aaagtaatag aatgtagaca taggtgctaa tttttcttg cgttaactag 38460
tgtttgtaa atthattaa tgtaatcгаа tttaattht ctaataacag actgctthaa 38520
acctatgtht ctcaaagtgc tggccctaga ggaagagcg agcttggcaa tgtgtgcaa 38580
ttaacacact gthtcttca tcaagaaat ctggcttctg aaaaattctc agctgtacta 38640
agcagtgtgc tttgtgattt acgttctgac acaaccgtht tcttgttgac agactagcag 38700
caacaatta acgggcccag agccagtcta tggagcact tthtagtact actgattthaa 38760
atthagtact aathththaa tgtthththt gththththt gththththt aaagcatgaa 38820

-continued

gaatagagaa tgtaaatata gttaagaaaa ggcacttaat tttcagctta cgtgtggtaa 38880
agactccagt ggctttgggg ctctcttggg tgcctttat ggcacagagg gatacgggtg 38940
actcagatgc aggctgcaga ttcctcagtg ggtacttggg gacagagga cccacattgt 39000
actgcctggg ctgtcctctt cgggtaaggg ccacggtcag tcggtatcct gttatctggg 39060
cccagggtgg agtccatctc accaggacag tagaatcagt ttcattgaca aactggaggt 39120
tagtgggagc atccagttct aggaaaaag atgaaacatg ccaagaaata tttagatcag 39180
taatgatcat aactcaagtc ctgaaacttg attgaatgtc taagttttct ctctcaagg 39240
ttgtaactat gtgaaagtca aaacctgga aaaactgagc cagtaagaga ttgagtgcta 39300
cacaaaactt tgccaaaact ctgccagtc tgagaaattg tggaaaccatt ttgcttgact 39360
gtgatccttt gggaagtcta attttccagc aagactttcg catcagctc agaaattata 39420
gttacattaa agcaatgga tttttaaaaa ggcaaaaca acacttctga agttcacagt 39480
tagctctgag ccaaacagat taaggaaaaa gggatatgcc tcctaaaatg acagcaagaa 39540
gaagctctct atgggtcaaa agcaatattc tgtgaatagc cctttgctag agaaataaaa 39600
gttattttca catggttcag attagacagt aattacagta atgataacgt atcctcagat 39660
tcatggtagg atactggtga gtttgacta tcagaatcac ataagttgtg gagtttattc 39720
ttgagaataa taaccagcag ttttcccacc gaaatttaa ttaccttca aatagaggaa 39780
atctttaaga ggcagctcag tataataatt gagagcatgg gcttcggaca taggttatag 39840
agccagctct atcccttct atctctgtga ccctacaagt tgcagaactt ttttaaagcc 39900
tcagtttctt taagtattca ataccggatc tagtaagacc ttccggctgg gcacgatggc 39960
tcacgcctgt aatcccagca ctttgggag ccaaagtggg tggatcactt gaggtcagga 40020
gtttcagacc agcttgcca acacagcga acccatttct actaaaaa caaaaattag 40080
cagggatgg tggcacacgc ctgtagtccc agctactcgg gaggtcaggg caggagaatt 40140
gcttgaacc aggtggtaga ggttccagtg agctgagatg gtgccactgt actccagccc 40200
tggcgacaga gcgagactcc atctcaaaaa aataaataa taaaataata ataataagac 40260
cttcttctc aaggtcaaaa caaagagtaa atatgagaat aatgcttga aaaactgagc 40320
agataagctg gtggctagta atactagaca cttgacaatt gctattgta aatgaattct 40380
actctaagtg ttatgatcct gcattttaac tttgatctag atatatttag ttacacgaaa 40440
aatgcaacta aattagattc tcttcagttc tccccttta gtaccctca gctggtacat 40500
gtattttaga aatgggttac cacttagcat coatggagct tgtctgagtt ttgtttcaa 40560
gtttaatgtg ctttgtgttc actatgcatt cccttgccgg ctgggtatth ccctctctct 40620
aagccataca cctctgagtg aatggtttac tgcactttct cacaggctgc aagtgagggt 40680
ggagaatttg gaggggagat agagatagga gaagacatac tggttgtctg ttgagcagtc 40740
agaggcttgc tctccctccc atggctcact gcaaagactt tgaagtaata ggtgacccca 40800
ggggacagcc cggtagcttc tgcaaaagtg ttcctgctga tgggcagcct ctgcccgtgc 40860
tcgcaggca ggttgacggg gatcacatcc acacggtagc cggctactgc actctcaggg 40920
ggtgtccaca tgatggtgac cttcacgtct gtcacttcca caaactgcag gtccttgga 40980
gagggcactg tatctgacag acaagagtc actggtcatt cacattctct gctgaagatt 41040
acttttaaga agccagtttc ttggatagtg taggcagttc attgagcctc tcattcgaga 41100

-continued

gttttgctaa gttttgtctc ttaatcagat cacatthtca tggccagtc atagtgccta 41160
aattatgaca tttgaacctt ttatctaaaa taggtaataa gtgagttacc agcatatgaa 41220
gtgtcatatg gtggtggggg gagaggggta taagaaaaga tactaaaaaa ttttttagaa 41280
aaagcctttg aaataactgc agaaaataat gtaacagttt ttgttttact gaatcatttt 41340
ataaacttta agaatacaaaa atgaaatgac ccttaagctt ctaggggaaa attttagatg 41400
ggacgttatg gtgtcagttt ttaatcagaa aggccagtag ccagcttga taagttactt 41460
atgacatttc agctaaatat cagaagtta gggaaaaact aatataaag atttcttgaa 41520
tgagaggact gttgacagag acaaaacaac cctccttcag gaacaatcct gataagataa 41580
cataggaagt gtcttcttaa aaaagttacc tgagcgtggg gtgccagtg tttctgttg 41640
aatgacaaca ggtgtacttt cttgattttc ttccacagca tagatagtga tgttatactg 41700
aacaccaggt tgcaagtca tgagggtgac ggagtttgca gtttcaggaa ggttgagttc 41760
tgtgctgcta ctttctactg atggcgaata gactattctg tacctcagc attcatgagc 41820
aaaagtaaga tgcaactgtt tctttgagac agaggcttag aaacaccata aatcctccct 41880
tttctaatat cccttagcat cttgcttgat tagtgaagag ccagtgatac tttttgactc 41940
tttgaatatt gccacttctt ctaataatga ctattcagca cgaagataaa ataagaacag 42000
tcatgcgaac ctttttaata ataaatgtca tccttgaaga cctgacatcc ttaataaatg 42060
tcaattagag atgccattag gtgaaatat tcatgttaag ctgatttttc ctctttgagc 42120
tccagacgag ggaagcaatt aatgaaccag cctaagcaga gttgtctagc actgtgtgtg 42180
tctttttgga acttcagtga tgaggcagtt gaatcaccta atgagctaag acgacagctc 42240
agctttctcc ttaaggataa atcttagagc aacgttttca ggaaattaca cctcttcaac 42300
cctgggaaca cttgctttaa taagccaatg agctggcttt acaggttgtg tgtatgttgt 42360
ggtgtgtgtg tataaggag ttggtgaaat ggtttcaagg ggaaatgtgc catagaaaat 42420
aactcccca cccccgccac acacacacac acacacacac acaaacctc ctccataaa 42480
ttatattgga gatthaaagt gatatgcagg tccgcagtca gaactctgag cctccctgct 42540
gatgtcatta agattaacat agcagccaaa gagggggagc cttagctgac ctgtgatggg 42600
agcctggggt ctgctccagc gaacaacaat tgagggtgca tcaacttggc ccacagtcgt 42660
gtcaggaggg gcatcaggcg ctaagaaaga aagaaagtgg ggcaaacagt caggaagtgc 42720
tactacagc ctgtgtttta cagaaaaaga ttcttttaac actatgtagc acacatgtac 42780
ctgtgtttg tgaagtagac aggatcaaac tctgctcccc atcctcagat atctgataga 42840
catttacaat gtattttcgg ccaggaagca ggtcagggat gttcacagaa gtggctgtgc 42900
ttggaagatc tttgaaatga aaagaaaagg taactaatca gagcaacta gtcctcaaaa 42960
tgactctaca gcttataaga aggatatttt aagaattata tattcatagg gaaaagcagc 43020
tagaaggtaa tgtgctaata gctaatacgt gattattatt attattttta cacagagtct 43080
tgctgtgttg cccaggttg agtgcaagtg cacaaattca gctcactgca tcctctacct 43140
ttcgggttca agcaattctt ctgctctggc ctcccagta gctgggatta caggtgcaca 43200
ccaccacacc cggctatttt ttttgattt ttattagaga tggaggttca ccatgttggc 43260
caggctgac tcgaactcct ggcctcaagt gatccaccag cctcagcttc ccgaagtgtc 43320
gagattatag gagttagcca ccgcaaccac ctaatctgtg attattaatg tactatggga 43380

-continued

aaacaccact tcttgtcttt ctcccttctc ctccctctcc ttcttccttt catttttcca 43440
actttttttc taaagacgta taatttttgt aatcgaatgc caaattttta aaaatagctc 43500
tctgcagtaa gagtcagggt agtgattatt ggcggaagat gatgggggta gggtagagag 43560
atattctttt ccttgatctg ggtgctggtc ttgtgggtg ttcactttat aagaattcaa 43620
attcactgaa ctcagtgctt acgatggcac acacttcagt gcgccaatga agttttttaa 43680
aggccccca ccataacaat ttgaaaagta tgagtgaccg tactgtgctc acattgtcag 43740
ctgcagctct ggtaattact tccaccaaga acacagcttt tgctgtagga acttgtgcac 43800
aggtagggcg atgatgcagc tggcatcagc gaccaggaac atgtgacagt ggcttgggaa 43860
aaacccggcc acttcttaga aagagtcagt ccttcccttc ctcatgccag aggggtggtt 43920
gttcagctct agctgtcaac cagttccaaa ccacagatac cacctgccta gaggccacc 43980
acccccaca ggagagttt ccagcagctg ccctgaacct gtcttgagcg acatattgag 44040
cttaccagc tactgtggct catctccctc ctcaactcagc tcatattcca cccggaatcc 44100
cgacacgggt tcggaagctg agaccagga gaccacaaag ctactggctg tgatttcggt 44160
cacagattca gaagtggcca caagaggaga aaagggagtc gtctctctg tcaagggtgt 44220
gcctagagag tcacagaaag gggaaagtca gccttcagtc atctagaaaa tttaccgtcc 44280
tcgtcgccaa catcatttta aaaaacgttg gtgcccctcc tggagaata tgaattgaga 44340
aggaaagcat tgctcagga gagctagct ctaggaaacct cgggggtagg aggcagtagg 44400
gtggttgtgt acatactggt cacaggtgtg ctggtgctgg tgggtgtgaa gtcaaagcga 44460
gtcacttctt ggtggccgta ctgctggatg ctgatgagct ggccctcgta taccacacca 44520
ggcttcagcg ctttgatggt gtaggagttt aagtggcctg gtatggtagc ttccttccaa 44580
cggcctacag aatttttctg aaaatttaaa ttaacacaca cacacacaca cacacgtgtt 44640
tacaaggatg aacatttgaa gatgatgttc agtaatcttc aaagaaaaat gacttaagca 44700
ggacttaggc agctccatc tagaggaaaa tcatgaaaa ctggattccc aatggttata 44760
tcgtagata tagttagata taatgctaac tattgttaac atcaacaaac tgctgcagtt 44820
tacttactga taccatact ctgtgctgtt tacctacatt ctccattta gtctccatat 44880
ccttctgatg ataaggcaat ggcattgttg ctcccgtttt acagacggaa aaacagagtg 44940
atatgtgata tgccccgggt gacatagttg cagacactgg gcacctcacc tttttttttt 45000
tttttttgag acagagtttt gctctgttg cccaggctag agtgacagtg ctcaatctca 45060
gctcactgca aactccacct cccaggttta agcgttctc ctgctcagc ctcccgagta 45120
gctgggatta gaggcgtgtg ccaccatgcc tggctaattt tataatttta gtagagatgg 45180
ggtttctcca tgttggtcag gctggtctg aactcccagc ctgaggtgat ccacccgcct 45240
cggactccca aagtgtctgg attacaagcg tgagccacca cggccacca caccttcaact 45300
tttaccaggt taattgtgtt atatcagtaa cataaatga tgaacctaa ctttgcctct 45360
ctcacttatt ttttattctc accacaaaca gatataattc tctctgaagt acagaaatgc 45420
caatgccata tattttatct ttaactcag tgggatacaa tcaagtttc ttgtcctgaa 45480
aaaaaaaagg cagaatgttg ctttgtgact catcatatac caggttcaac gagtctggtt 45540
ttgtgactca ttcttgata tgagctacc catcagaagt ggttacccaa ttttgacaa 45600
gactacagaa aaaatgcaat tttggtttat ttaatctttt gcttccagtt tatatttcac 45660

-continued

atcctgttca gcttcattaa tatgccatgg gtcacaaaac tcagtgcaat aaaatgtgta 45720
tgaagaaaac acccttcaga aaagatgaga ctctttcaag tgtaaatact ctaaactaat 45780
ataagtcaaa atatattttt tgtgcccagt gattttttaa aattaccag tcaaccattt 45840
cctcaataat tcaaatactc aagtgtccat ttatattttt ggaataagcg agagtgatcg 45900
tagtactctt cacagtgaaa ttttaactca aatacactgt ggaaaatttt gaaaccagtc 45960
attgcaatta gctcaaattt agtaatcatt tttcaacatg agcttgtttg gttagttatt 46020
acaagaaaat aacagaaaat aaagtctcag gtggcatcca ttgaaaaaa ccctgaattt 46080
ttgtaattgt taaaaatctc aaaaaattca tcaactgctgc tttttgaaga ctttccaatt 46140
caatgtactt tttttttttt ttttttgaga tggagtctcg ctcttggtgc ccaggctgga 46200
gtgcagtggc gtgatcctgg ctcaactgcaa cctccgcctt ctggtttcaa gcgattctcc 46260
tgcctcagct tcccagtag ctgggattac aggtatgctc caccatgccc agctaatttt 46320
tttgtatttt taatagagac ggggtttcac catgttgcc aggatgctct cgatctcttg 46380
acctcgtgat ccgccaccct tggcctccca aagtgtctgg attacaggcg tgagccactg 46440
cgcttgccc aaagtacatt ttataccact ttttaagtctc ccattgcaat ggtctcacat 46500
aatactttgg atatccttaa ttttaaagta aataaaatta atgatgtaat aagaagatga 46560
gagactagaa gatattggat gcaactgatgc tgttgatttc gatcaactgga catcaagttc 46620
tgcaaatggc cagattatgg acataggggtg aaagaagaaa aagaatgact tttctttttt 46680
ttttttcat tcaccatttg ttaagcagtg tgccaggtgc tgaagcattc tagactctac 46740
ggagactaca ttgagccaat gagaaagaca cgattcccct ctttaaggga atctgtttcc 46800
atttctatt attctgtctc tgccctcatg tagacaacat gcttacacct ctatttgttt 46860
catttgtaa caataagtca tatttacaga gtctttaata tgagacttga tgatttattt 46920
atatatttat ttatgtgac acagggtctt tctctgtctc ccaggctgga gtgcagtagt 46980
gccactcttg ctcaactgcaa cttccgcctc tcaagctcaa gcgatccttc cgcctcagcc 47040
tcccagtag ctgggactac aggtgcatgc taccatgccc ggctaatttt tagccactgt 47100
gccagacaa agttgaggat ttatgtcaat aataccacc cttccttcca gaatcttga 47160
cattcccagt gtctgacatt tgtatcatta tatttgttct tcttggtttt atatttgtaa 47220
tttaaagtat tatattaagt atcttttttt tttttttttt tttttacttc atcagctttt 47280
ggtaccatct ctcaattctc tattttgaaa gataaagctg ttatctctc tcttctccac 47340
cttccatate caccattttt catgtaaatg tttacttttt atttgtctt gaagttgcaa 47400
ccaagtttcc atgacacatc tatagaccga tcgcatcatg cattagaaat acattctttt 47460
caatgagtct aacatcatga ctgctgggtg gatcaaagtt tccagaatca aggttcaatg 47520
gaattattcc tctttgtttt tataccttaa tattgctgca ttttagtttg ctctgtatag 47580
gaattattgc cttataatac acttgggggt tttttgaaac agaactctgc tctgttgccc 47640
aggctaggg gcaatggcgc gatcttggct caactccctg caacctttgc ctctgggtt 47700
caagtgattc tcctacctca gctcctgagg agctgggact acaggcgtgc gccccaagc 47760
ctggctaatt tttgtatttt tagtagagac ggggtttcgc catgttgcc aagctgtctt 47820
cgaaactctg acctcaggta atctgcccgg cttggcctcc caaagtctg ggattacagg 47880
tgtgagccac tctaccagc cacatttggg tttattatac attcctccac ctgattgtca 47940

-continued

tgttgattatt ttaactgcat tatttgcccc tgtgattaat tatccacaag cctacttttt 48000
ccatgaagtc cctttttatt ctgggagacc tcaatccctg atccctcaat cccatcctca 48060
tctttctggt coactctgat tggctgtcct ctaagccagc tacgcagttg ttgcattgca 48120
aagttttcca cctctatcag gggctgatt tgggaagcact gtttgcctgga ccccaact 48180
tccttcataa ttcactcttg tttttctgga gtgtatctac aaataccctt aaagaaaagg 48240
atatatggaa gataaactgc ctaaaactg tctgaaaata ttattttgcc ctcatatttg 48300
attgataacc tggctgagta tagaagactt taaaagtttc tttttatcct tggaaactctg 48360
acacatcaaa gatgcgtcta gacgtggttc tttccactaa ttttctgggc acctaggagg 48420
cccttttaaa tctgaaaaat tctgtccctc cactttagaa atttagaaaa agaataattt 48480
taacagctta aatttttctc gaagagtta aatttaaccg cctgggttg tcaaattttg 48540
tttgtttggt tctttgtaat tgtagaattc aatagctatg atttaaaatg gacatttttt 48600
tctcctgtgt ttaaaacagt gtttctcaaa ttatgagcca tgaactttgg agttaaacag 48660
ggcaggcttg agagagtaag agaaactaaa gctaaattga ccaatcctc ccttattttg 48720
tttcttcaat ttctagtca tctctcattag gaaatcacag ggctattggt taaataaaac 48780
caacttccat cctttatcta aacttccctc ccatcatctc ttttcttct atctaaataa 48840
attcagtggt tcaaaagtaa agtttttaaa atgatcactt aaaaagttaa tcaaaaaaga 48900
aatatattaa tttggctttt tttttcaaat tatgaaaatg atgccagtaa ttttctttg 48960
atgaggttgc aagacatata aaaaccttag atttcttctc cacattgggc aagaacaatc 49020
ttaagcatta tcatttctaa gtccttgcaa aaaagatttc tgtaacataa agagctaaat 49080
tcttccaaat accagcccc ccaccgaaa aaaacaaaac ccaaaaccaa aaccaaaatc 49140
ccctcagtct acaagtcaat ttaattgacc acatattggt tgttcaactaa acaaatatag 49200
taaatgctat aggaatagtt aatcagagtt gttggctcaa aagctgggaa aaaaagaac 49260
catcagaatt gataagatta ggagactcag tatccaaggt tcttgggttg gataactcaca 49320
ggctccacc tgagaatgta cttggaaatg tgagatggct gtggtgcatt ccaactggatg 49380
gggtgggagt tgggtgact cggagtctca gtgataaata cttcgacag accacttgag 49440
cctgaaaatg aaatgtagc atttaattat cttgaatatt cactcacttc aacttaaac 49500
tgtgtacatg gacagtcac attataaaca gctttgcttg tctttaaag tgagtataat 49560
aaaattatct ggcattaatc ttttaaatat ttacgttgaa aatcaatgca gcaaatccat 49620
ttttagccga taattattat tcttcacatg agaaatgcaa atcatagttg tttatagcaa 49680
ataaaatata tttttaaaaa aaagtctctt ggcttgcttt tattttctct actatcaaaag 49740
taagaattgt gagaaaattg tgaacagatc ctacaaaaat aatgtaatac agaacacaga 49800
cttcatttac ttattgtatt gacgtctttt taaagcaaat tctaattttg gaaggttttc 49860
atcaagaatg aatctactac ctaagaagga aacaacattc caaaggccat atgagatgat 49920
gtaaaacaaa cacatttcta aatgggggta gaatcttaga caaatagcaa aggtaataat 49980
taaagaccta gttttatttc tcatcagtct aagcagtcta tcagtttggg aaaaacatca 50040
agttttattt ttattcaaga agtttactcc caaatcacac cattgatcca tcaattatt 50100
aagtgattaa atttattttt tttccaaaat cagttgtcaa cagaaatgac tttatgaagt 50160
gtatactaaa ttcagttatt ctacacctg atgtatatca ctgttatcaa caaaatttaa 50220

-continued

agtatggttt tggttctttt gtttaaaata aaaggaaaat aagggaaagta gcatcaattt 50280
ctaagtagtg ttaactccat ttaagataat aatgaaatag ctagaaaatt tatgcttgca 50340
gaacagaaaa ctatcttttc ttaaaacact atctttatga tgtcagtcaa tgtttaaatt 50400
actagaaata aaaattgtgt cttatagtaa ccttttcaag aatttctaac tgacactctg 50460
tcaaacagag caaaattttg tatgtgtaat aaagcacttc accagattat taacagattt 50520
aagttttctg aaagcttcta ggaaaaaaa tacattgatt tcaaaacttt attatctaag 50580
ccattaaatt ctcttggaac acaaagcttt ctgtagcaga aaaaatatat aacttggata 50640
tccatgtaga ttcagaaatt ctaaaacaat atgctcactt ccacaccagt aacaattcaa 50700
agttggttga agattagcaa atactttgaa tcttaattcc atttaatgca catatttcca 50760
agccacatga acacaaaat tcatgcataa aattatgagt ttttcatttt aaagttgtat 50820
gtatttcaca gaacacacag gatgtttgtt ttaatcaatt ggcataagtc aattccatgc 50880
ctatctttga attaccctt ctgtttatga agatccttac aaatgtaaac acatttttaa 50940
agacaactat tcaagaaaa ttttgagat gtgactattt aacgtacttc catgtaaatc 51000
aggaaagcta ggaatatcat gtccttaatt atttatgata attggcatag ggctatttaa 51060
atattgatcc tttaaataaa tcctactaca atttcctcca taaagagaca cctgataatac 51120
actttattta gtcttgctca tcatgatttt gaggggggac aagcttatga tggataaatg 51180
tctattaagt tttgaaaaga taacataaat gacctttact gttcacctct aagcctaatac 51240
atatttctat gtccacaagg acacaaggag gaggataaaa atgggcaggg aaaggaaaga 51300
aaaagggtac agcagtggtt tttactcaag ccaaagtga gtacaactac agaggtccct 51360
tttgctttca taagatttcg agcactggtg tgaattcca cgttttaaaa tctagccctg 51420
tggcatagcc cacctttaa ggaaatggac tgaacagcat ggaatttagc atgtttttgt 51480
tcaggaagta aagagacttt gtgtcagtta tttcttagat caattaaatt cctatatctt 51540
agcttatact acgtataatt actctagctt atactatgta tcctttaaag acagaaaagt 51600
taaaaattat gacatctttt acattattta gatttaaact tttgacagtc ctttttcgct 51660
actggtagtt ggtataaact tttaaattaa ctttaattggc atatactagc tatcgccatt 51720
gtgaaatcct tactcccctt cttattcaat atgaataaac taacaataaa atagttgaga 51780
tgataccaaa aaaaggagat agtacaatgt ctacactttt aaaattagtt tcaataaaag 51840
tctaaatgaa taatttccaa tagtagtgc cttggagaag ttaaacattt atgatttttt 51900
gactcagtc tttttttctc tatttatttt gotgagatct gtgcttttct agttgcttat 51960
tgacaaacca aatgcctctt atcttttgct agtttgaaca actctacca gttctctgta 52020
ggagaaacaa acaaggcag actcagagac aatctggttg taatctgctg ttccttatt 52080
ctcaccacac tcttaaagcc ctacgtgcta ggcacggag caaacagac atacagataa 52140
gcaccaaaga gcatatctga ggtcaaatc caatctcttc cctccgaaat cacctctatg 52200
ctcagtgga tcttgctggt ctccttttta aagagtagga gtaaggtoaa catgtaatac 52260
tccaatactc aaggattcct tcctttccag attttatctc caacatgtcc acctaaatgc 52320
ttccagcaag cagctatgtg gacagcctta tgttgagaga gacgaagatc tctctcaatc 52380
attcagaagg aatcattttt cagcaaatg caacaccaat aacaatcata aatatagaaa 52440
ccacagagca tttttttaga agtttctacc atttaagtat catttgcac atataaacia 52500

-continued

acaaaccatt catatactta tagtacttcc acaatcagat ttaaatatat attaagttac 52560
aaaaaattaa gtgattccta cacctaaatc ctccaagaaa tacaatttag gaggttggtc 52620
aatctgtgct ataagccttc aaaagtgaag tcaccttccc cagcactgac agtgtctgag 52680
ggttgggaaa agcagaatct tccaagttcc tagaaaaggt gcctctgggt gtttagtatt 52740
taggatggaa cattaaggt aatcagctac aatcttccca tattacagat gaggaaatgg 52800
aggaatgagg ctggctaac actaccttg gccaaagtgt tcccatagct tgttctctgt 52860
cacagtgaat cttttatcac agagttaatc ttaatttctg tcgcatgcag attctgacac 52920
cctgaaactc ttttgtttct gaatgtaggc cataatttca gtcaaacggt agtgaagact 52980
ttggagactt tgcacctcag cagttgctca aataactgat gccttttaat tagccttca 53040
tgttttatta acatgtttta ttataatcac tcagtgaata tttaaatgca attttcccc 53100
caaatgaact aattacacct acagctaggc atgatgtctt tgtgacctgg aaaaaccag 53160
caataaacat ttgagagtgt ttgaactgtg tgggcagaga gttgaccaag ttactcaacc 53220
cttggcccaa cactactgtgt tcctgggtggc aactgatcaa acaaatgaaa agtttgatgc 53280
aaagggagct tagatgacat gttgaatgaa cacgtacctt atgctgcatt tcatgccatc 53340
tctctatgca gcttctctgt taagaatcat aatataaagc cccaaacctt tactcccagc 53400
ataactcact tgggtggtttt tatttccatg tatggatata agatgaaatt agccagagtg 53460
tcttaacctc acagaatatt ttacagagtt aagaagtaaa attataaaca ggcacatttt 53520
catagccttc tctacctttt acttgataga ggaagagaga aaaattattt taatcaaaaa 53580
ttcaattgat tattagataa taacaagaga aaaggggtggg aaagagggga gacccccaac 53640
ttaggcata gaagcattaat aatcactgcc ctggtgttat aacccaattg gcagttctca 53700
acttgacgta atagagctac ttacttggat aggtctgtaa aggttgcaa tgccactccc 53760
caatgccacg gccatagcag tagcactggt atctgacacc atgcacatac ttctcccatg 53820
aatctccaat ttgataaaac gtcccagtct ctgaatcctg gcattggtct aaaatacatg 53880
gaaggaaaag ataaacagcc ttgaagactt ataactacga atttaatttc aggggtgctag 53940
agcatacatt tagctttaat agaaatttgc aattgttctt acaaatatat gcagttaaaa 54000
aaatattttg ttcacattct ttaccttaca ggacagtttg tagtaaaaaa aaaaccaaag 54060
tacctattag gaggaatatt tagttttcta aaaggtaata tttctaataa aaaatgtaaa 54120
gatttagatt tcagtgtata ttgataatct acacattaat tttaaatatt agttattatt 54180
tttaccata ttaatataac tggcttaaca tttaaatagc aagaaaaaaa tctgcaataa 54240
tttcattttc ctacttctt caaaatttaa agaaactggt tgacatttaa atgaatgaac 54300
tagactttac gagtatacct gctgagcagc ttctaagaaa acctctcagg cttctctctc 54360
agttctggta atctcacgct aaacaacgg atgttatcag tttgttttct cagtgaacaa 54420
cctgtctctt ttggtaaaga aatagtaatg gatcacctta aactttgtag tgcgtgtaac 54480
acagctcttt taaggtcag ccttactat ttgcttagtt ctattttgaa gggcggccaa 54540
aaaaaacct agagtgtttt ggtgggtgat atttacacc cttgtctatg ctgttacaca 54600
gctggaagg aagtgaagtt gagaccagcc tgaccacat ggagaaaccc catctctact 54660
aaagatacaa aattagctgg gcctgggtggc gcctgcctgt aatcccagct actcaggagg 54720
ctgaggcagc agaattgctt gaaccaggga agcagaggtt gcagtgaacc gagattgtgc 54780

-continued

catacactcc agcctgggca acaagagcga aactctctct caaaaaaaca aaaacaaaaa 54840
caaaaaaaca caaaaaaaca aggaaagaat actttaata gcatggatta tctttatgat 54900
tctaaacca agaataaatt ccattcaatt aaaataagcc taaatttgtc accaaattca 54960
gcaaaaacca gacgactacc aatatcgaga tattttggtc cagtattaga gagaagactt 55020
tgcaaagttc ttgcaagca actgatcaaa gtctggagcc aacttcagtc atgtgatttc 55080
acatagcttt tcttatctga aacttgctaa ccattcccc tgtgcaaact catctagggga 55140
aatagggcta ctaccgagc ggatcacact tccacctgcc ccgaccctga ccgaagcatg 55200
tacagttcag catgtgcccc tcttcatgac gcttgggaa tgtgtcgttc acattgtaag 55260
tgatgtcatc aacaatgcac tgatctgttt aggaaacagg tgggtgagtg agaaactttt 55320
taaagttcca ttgatgaaag aaaaaggtg tgcttctcaa agcatacaac tacttccact 55380
taattatcaa caccttccc taaaataact tttctacaag catagaataa catttttttc 55440
aaagtttggc cagtaataca atttttgcca aaaaattttt tttcttcca aaatgtgtaa 55500
cttttgcaga ggttacagta tgctcaagta aagaatgtgg taaaatacac tttcctgtga 55560
gaagccagta ttgaaacaaa tacatgtcct actttctgaa aatttcttgc ttagttgata 55620
atttcttta aatattctgt tgtaggata aataagaaga agattaaaca tatttatcta 55680
acaccagca tgatgtggtg gtctgtctc tggaataggg ggaaggggag gagatactta 55740
gatttgggaa ataaaatgtc aacagtattc acaacgctaa aataaattag gccaaacttg 55800
ttgctcactc ttgagagat cctgagaaaa ttactttttc ctttttgggt ctgagctcct 55860
tgatctatat agagtgggt ctctatgtgt tattaaggtc tctttcagtt ctgcaacacc 55920
acatgtaatt attactccc gatggtgta cccactggac accacaaaag gagaacaaat 55980
tcagaccaat atctacgcac tagccacata gattattaat agctcagggt tctcccgaag 56040
tattactttt catatcaaag aaatggaaaa ctaaacact tttaaaagt caatacttcc 56100
tcatattata tagctgggtg caaaggagaa aaactaatct gaacaaggaa agttagtccc 56160
ctatgtattt gtttcctgaa atctggatct gtaaggatct gattttagaa tagcaaggtc 56220
ttcatttctt tacagaaatt atcctatcct ttctagacat gaatccagtg gcatagccta 56280
aagtgtaatg gatcaaatgt ttcattctta cttttgaagc catctttttg gctcagccta 56340
agtacttaca tgctaggagt gtttcagaac attagcattg ctatgccgct tgtcttagca 56400
ggctgcctgc taagccttcc aggttatct taccatgcac agtagactaa tgaatgacgt 56460
ggcattccat atttcttccc ctgcttttgg catagtgcaa gttttcataa aaaaagtgt 56520
ccctttctgt accttttggc atctcaaata tttttgtaa tcagccagca tacctcgaag 56580
ctgagagtag gcaatgatg tccattcccc acgaccattc ccaacacacg tgcacctcat 56640
catgtgacct atgtcatgct gcttatccca ctgatctcca atgctgtaca tgaoccttc 56700
attggttgtg cagatttctc cgtgggctgt ttggaatgg ataagaaatg cacttgataa 56760
atgatcacca ggtctaaaca tgtatgtgtg caaaagatc agttcagttt ggtcatccta 56820
agaattctca cttgggaaga agctttgtgt goatcaaaga catctaagtg ctgtcaggtg 56880
aagggtaga atttttgagc cctggaggca gaaaacgtag tccaaacgtg tcatagaaaa 56940
tagttacagc aagggtagg gtaggttaaa ctaagataat atctacacat ccacacaaag 57000
ggttcttctg gttctagaag cctggattta actacagctt cataactttt cccagttcca 57060

-continued

gtttattctg gatcaataa aaactatggg tttggatata gaattacaag caaaagtggg 57120
ttttttcttc taaactcagc atcttctttt gttgaaagaa ataaaagtat aagcaatttc 57180
tttgtctcat agtcacactt gacagtgcct accgcccagg ctcttgtttc tacaactgga 57240
aaattaaag gaaactatgg ggttcttgaa atctgaaaac aagtaaaag tatttcgtttc 57300
catttgttcg ccatctatac aaactgggga cctttaaaaa aagagtgaga ttccctacct 57360
gttagaaaa attgagtttt agtcactttg gcaactatta ataagtata gagcttatgc 57420
ccatcaaac acaacctatg gtcaatattg tagctctgtg actaaagccg tcatacatat 57480
attaaatgtt gtaagctact cagttgtgtg ctctttacaa agacaaacct gtgcaaatc 57540
atatttccag acttacaat aagtaattat atttttgaca ttaaatcatg tcaatgaggt 57600
ttaaaaaata tataaataat tatctaatta ttcttcccaa tccttatgta tatgatgtaa 57660
catcacacac acacacacac acacacacac acacaaacac acttcacatg aagttttctg 57720
ttgtctcaaa attgtgttaa tgaatgaaat gggagtaaac tggacactag tctttagtct 57780
ctactcccta aattgtgtgc aaacaagaca acccacaagg gcttcatctt accagccatg 57840
ggcgagaacc caaacttctg gtcggcatca tagttctgtg tggcccaca ccacttcatg 57900
ttgtctcttc tgcctcaga agtgcaatca gtgtaattgt ggttggtgta taggaagggg 57960
aagtggcaca aggcaccatt ggaatttctt cctcgagtct gaacaaaaac tgccaggaac 58020
aatacacaac aaagaaggaa aagattaccg ctgagctttc caatgtacac agaattactg 58080
gtctgagaga gccaggttct tgcagctcat tcacagacag tctatctcag aataaaaaat 58140
gtctgataat atagtcttaa gcttcttagc atatttacct tcacaccttc aaaacagggt 58200
taaatccac tttactgtc atttccccag ccccccccc acaaaagatt aatgatagc 58260
tttgaataaa acattgtttt aactggaaaa taagetaagg cactaaaaca ttcattcttt 58320
gtattcctca gccctgtggg atcatctact gattccaaaa tacaagtatc tttaatcaca 58380
tcataatccc aatgtgttaa atcaaatgaa ctggaaggct actcccttgg tgcattttcc 58440
caacagtact ggggaagatg gaagaatcct cagctgttac gtgagcgtg cgatctcttg 58500
agtcaaaact gaacaaaagc gatgcttctt gggcgggttc aaaataaatg gctagaatgc 58560
tggcaataa aactagggca tgaaagtgag tttttctcac ttctgtggtt ccccttggg 58620
acactcacca gtgtggtctg tgcagaaaga gtatttctgg tcctgctcat aattcgaagt 58680
tgtgtgac caaagatgtc cgtctgtg cccttctgtg gtgcaggagt agaacgtcct 58740
gccattgtag gtgaatgta agacacatgg ctctccattt gagttgccac cgtaagtctg 58800
ggttacagct acaatcataa tcaaaagagt gtcagtaaac agagatgctt tatctccac 58860
cagcagctgc ttatttatgg cttaaatgag atactgagct tgtgccctca aaggaagaag 58920
ctatgagat agatgatcat ctctgtaatt tcagtcaaaa tgcattactc aggaatccat 58980
gtctactgtg gatagaacga cacctgagtt cttctgtgta tgccttcat gaccacttct 59040
gctgccatgc ctcaagttct ctattttaa agctatattt atacaggtac attaaggatg 59100
ctgottaagg atgttggat aatatttatg actatatcta tgcataaaa ttaatatatg 59160
agactattct ggggagaat tgtgtagcaa ttgttattgc catagtgact accatgtttg 59220
tagtgacaga tacatatatt aggttattgg aagatgttta catcacaag ctgctgaaat 59280
ttgagattta gaataaaaa cattgatgat tattcagaaa gtagctctct tgaacaaca 59340

-continued

ctttagacag ataaatata taaaatTTTT acccatgata tccatgacca gaaagaaaat 59400
acataaatta aaaagattgg aagaaactat ttcaaagtc aaccctaatt gttatcacag 59460
ggtggaaga ttacaggcct cttgtttgtt ttattgtttc acattgaaaa tgtattgtcc 59520
ttcaaaaact aaataataaa gtcacagtc ctattcttca aaaagataat gcatacctgt 59580
ctcttgccag ctgactccgt tgcccaggca cgtgcaaagc atttgcttat ttccttgtgt 59640
cttcagccac tgcatcccca cagagtagac cacaccactg tctgtgacac agtggccata 59700
gggaggaggc tgggggtgag gctgcggtg gtaaacagct gcacgaacat cggggaagg 59760
gccagatcct aatggcatga aaaggaatg tcacaaaact ggggagaga agacaattca 59820
ctactttctg cagtcaggat cttcaggatc tcacagagat cactctgaaa tctatgtcgc 59880
tactggcctg ggactggaaa aaaatgtccc ttattcagca gtacttttg catttagcct 59940
aggaaataat ccacaagtgt ctaccaagtg gtctccatcc tccagctcag ctggggctac 60000
tgggattata tttctttttg tttgtttgtt ttttggtttt ttgggtgtt tttgagatgg 60060
ggtctcactc tgttgcccag gctggagtgc agtggcgcgg tctcggctca ctgcaacctc 60120
tgctcctgg gttcaagcta ttctcctgcc tcaacctccc aagtagctgc aattacaggc 60180
acgcaccacc atgcccagct gatTTTTTTT gtatttttta gtagagacgg ggttttgcca 60240
tgttgcaaaa ctctgacct caaacctcgt tctcaaacctc ctgacctcag gggatccacc 60300
cgcttcggcc tcccaaagtg ctgggattac aggcagtagc cacctcgcct gccgggatt 60360
atattttctc atcacagaca tcatcctgcc actccatata gcagatacct tcaacacatc 60420
ctcactcttc tgagtaaccg agctcatcag aggggtgcct cagcagctgt cctggatttt 60480
ctggaggcca cctccacctt ccacatttta tttctcctct ggattatttg tgattctatt 60540
gtgttttttc taaaaccaat tgatcctctt aagacagctt tcctttcaga ataacacaac 60600
aggatgtgtt gatcttttct tttttctttt tttttttttt tttttttttg agacggagtc 60660
tcaactctgc acccaggctg gagtgcagtg ggcgaatctc ggctcactgc aagctctgcc 60720
tcccgggttc acgccattct cctgcctcag cctcccagct agctgggact acaggcacc 60780
gccaccacgc ccggctaatt ttttgatttt ttagtagaga cggggtttca cctgttagc 60840
caggatggtc tcgatctcct gacctgtga tccaccgcc ttggcctccc aaagtcttg 60900
gattacaggc gtgagccacc gcgccggcc gatcttttct tcaactttt agatattctgt 60960
tctcttgaag gccagggcc accgcagaac agaactcagg ctgaggctag accaaaggta 61020
cactgaattc agtgcctccc taagtctgc tgaacactag aaccacaaga tgttcacaga 61080
caaagaatcc caaggtcaaa cgtactaggg agacatggcg cccatcccct aactctacc 61140
atggagagta tgatgtaaag gaacacacag attctgcaga gcaggttcca tttgtttta 61200
cccattgctt ctcaactttt ttgaaacaat ggaatctttt cagtattggc aatattcata 61260
tatagctaac atttattgag tgtttatccc catttaattc ccacctacac ttttaattca 61320
ttttgtaga ctgacattta taattttttt tattttttat tttattttat tttttgaga 61380
tggggctctg ctgtatcacc caggctggag tgcagtgtg ogatctcggg tcaactgcaaa 61440
ctccacctcc cgggttcagc ccattctcct gcctcagcct cccagtagc tgggactaca 61500
ggcgcgccgc acaacgcctg gtttaattttt tgtattttta gtagagatgg ggtttcaccg 61560
tgtagccag gatggtctg atctcctgac ctctgatcc gcctgcctcg gcctcccaaa 61620

-continued

gtgctgggat tacagcgtg agccaccgag cgtggccgat gttataaatt ttttttggcc 61680
agtagcaatt ttattaatga taataaaata actcaaagat attatcaaatt ctaattctgt 61740
tcaccttcat aagtgtcaa agcaagacaa aaactcatca aaagaagaaa aattagaaca 61800
attaatattt ttaattgaag gatggctgag aactacgta atgatgttg cacatcatat 61860
agtaaaaagt taatcaaaac atattaaaaa taatgagagg agtgcacgtc cttgttggcc 61920
atatttctct cttcagattt ttatgaaaca tttatgacgt aaaaatcttt ttctcatgcc 61980
aaaaatgatt aaaactgtt ttttaaaaag aatgcttaga tatcatgtct ttcatggcca 62040
tctctaagtc ctctctgaat ttctaaaatt tcgaatgcc atacatgaag gaaataattg 62100
tgtgttactt tatagaatgt aacaaagtt gccagagcat agaaaccaca tataaaggat 62160
ggtttccact acaaaaagag ttcaatgtcg ggacaggaa gtatatttca ttcaaaaaca 62220
aacaacaaa aaatccaaat gcactttttt tgcctttcc tagcagaaga accaggtagt 62280
catataatta attataactc tcatttcctt tcaaatgaac tttggatatt caaaatcgt 62340
gtgtgtgtgc gtgtgtgcgt gatctatcca tggtaagttt tatgatgcta gcattcttta 62400
aaactcagtt ttatcttttg gagataaatc tggctttttt cccctaagg tttatttga 62460
tgtcattttt ctttttttct gaaaattta aaacttgaca ttaaaaaag ttttaacaac 62520
caataatgtg taaattatct taggttatta gctgaaacac tgcaattat cctcaaatgg 62580
catgtcaag gaaagatgga ttgctgaaa ttttcttga cctgcttccc catttcccgc 62640
ccctgctcgt cctgtgcctc accgctcgat gtggtctgca cagaggtgtg cctctcacac 62700
ttccactctc ctgcggcgtt gcctgtgcag atgcactgga gcaggtttcc tcgattatcc 62760
ttctgtctcc aggtgtctcc aattctatag gatgtccttg tgcctgatc gttgatctta 62820
tctgtgtcac aaaggaagca catacataca tcaggctgag agtcgcaagt ggtcaattca 62880
aacttaaaa agaatgcta tctatgcctg gtaatctatt tttggtaata tgttcatatt 62940
cagctctggt gctgcaaggt attacacttc tgaagatga aattacattt gttacagtgg 63000
aaaggcagct gaatggttta gtctagtctc atctctctgt caaataacat tttcatttat 63060
ctatttattt attgttttca cactgttctt ttgactaact tcattctctg gcattatatt 63120
tcatcttttt cttctcagc ctcttccagc atttttctta gtcactactg tttatggcct 63180
gcctttcaat agggaccttg ctttagaaaa ctaaaaacag gctaagcgcg gtggctcacg 63240
cctgtaatcc tagtacttta ggagggtgga tcacctgag tcaggagttc aagaccagcc 63300
tgggcaacat ggtgaaaccc ccgtctctac taaaaatacc aaaaaaatt agcoggtgt 63360
ggtggcgggc gcctgtaatc caagctactc ggaagctga ggcaagagaa tcgcttgaac 63420
ccgggaggtg gaggttgcaa tgagctgaga tcacaccatt gcactccagc ctgggcgaca 63480
gggtgagact gtctcaaaaa caaacaaaa caaacaaaa caaaaaaac aagcaaaaa 63540
aaaaagcctt ataacaatgg ggaggcaagc aggtatcaa ggtcttccac caatggctac 63600
aaagaagcta taagaaggtg gtagtatctc tttcttctt tttatgtaa aaaatgtaga 63660
ggatgcacaa tttctttgac taaaatatta ttttcccttc ccctataacc aaacatctac 63720
agtatttaaa aactactagc aaaatagccc caaattattc agtgtagacc ctccactgga 63780
agagtgggat caacataaaa tgataatttg taactaaatg accattagtc tatgcgtgca 63840
tttatctggt aaatagatta tttaaattat aatccaaatt tgctctgtgc cacgatgtga 63900

-continued

gtatagagga tctgatgtct ttgtgaattg gatattgttg caactcctgg gttagagaaa 63960
aaaaattaat agttgatgta ggttcaaaca aatacacacc cacatccatc taccacaaga 64020
atagaagttt gcatgaaggt ataattactg attagagctt tttcccaaat cccagcgaa 64080
aagtattatc atcccctgaa cctccatagc actactttca gtattttctgt tgaatccttt 64140
aatactgtgt ttgacagatg ataaatatgt ataagacttt ctcatctccc tgaccccccta 64200
agtactcttt gaagcgctgg ctaatgttca tttcatatct gcaccctat aataccttga 64260
acaaagtaag tccttataaa attttctgga ataaatgact tctctttgag aaacttttcc 64320
caacttgcta tttaaagagt tctaaataac agataaggtt tgaatcagac atctaatttg 64380
tcttaagcta aaatattgcc acaaagatct tggaagtctg atgatcacag taacatttta 64440
aatgcaaaat atctaataaa ccagcaagtt tctgggaaag ctttggtcaa aatggttacc 64500
acttttcatg agagacatct ttttgtagc aaagaatttg aatttgatg atggctatgt 64560
taaaggtagt taaatgggtt cagctagata aataaacag gttcaaatta tttggatttg 64620
attttaact tatttacaac accagtagtt aatggctggt ttacttcaac aaactctcta 64680
gctgttatta gtttacaata tcaaaaaagg agccttcaa cccaaaggca acaattaaga 64740
ccccaagaca atcttagaag gagcagtaaa gacaccagag gccagcttct ttacagcact 64800
aatcaaatgc gtttagtcag agacactatc cctaactcca gagtagataa atcattttta 64860
tttatgggat aaattctttt taattttatc aatttagcct gtgaatgtgc cctattttaa 64920
tttcaaacat agaagaaaac ctaaaacttt tgttcaactc tatgtcagtg gcagtgagct 64980
gagattgaac cttatcaaag tgatatgttt ttctatttat gtagtcttct ctgaagatgc 65040
ggaataagtt ggttatgaaa gttttcttga cttccagaag ttaccaaagc atgctgatat 65100
actggatgtg ttctgagtaa catagtatta cccttctctc taggaatcca gaaaacaata 65160
cctggcttaa tctttaacat aaagatgtaa aacatacttc tagaagtga agtgatgcgt 65220
ccgctgcctt ctcccaggca agtacaatct accatcatcc agccttggtg gggcttctcc 65280
cacgtttctc cgaccacata ggaagtccca gcagcatgat caaaacactt ctgagctgta 65340
gggaaatttg gaagaaaaac aggaaaaaaa ggttattttg aattgtgcaa gctccctgta 65400
atttaaagt tagtgtgtaa gcaaaaatc agccacattt tttttaagtg gcaactctgtt 65460
cagaaagaat gacacacca gcatctaagc acacagtaag tagccaaggc aaaagaaagc 65520
tcaagctgag ttggcaggca gaaattccat tctcctaata agtacctgga ctacatattt 65580
tcagcgagcc ccaaacttcc cagcagcatt tccagtagca aggatctatg tatgtatcta 65640
gtctgtggga gactacatat gcaaaattcc aatcttaatt ataaatgacc aattatgaat 65700
tttgagtggc ctaaatacaa tgacaaagaa agctattttt aaatattgga aagcaaaaca 65760
atacaaaaca agacaaaaac ctctaaatta gaagacaagt aaacagagta aaatttatag 65820
ccagaaactt tcaggtttgt acttcaaaa ttcaactgag gattgagatc caaagaagct 65880
caacaacttt taatttgaaa agaaatgagc agagtgtgac aagcacacac tatttcatgt 65940
ttcggccag gacattccat atagaatgt gtccaaagt tcattttttt ttctttttaa 66000
taaattatat cttgacgaag gaggaatctg agatcagcta ttaggtaaaa gaaatgactc 66060
cattctaaat ctttcaatc taccttatgc ttttgaagca aagttaagac ttctatgtgt 66120
agtcagtact ttttgatcta actcctgtaa gaagactatg gattatataa taattcatta 66180

-continued

tttcaaacaa aattcccatt tctttattgg ttttcaactgg aattcaggtc tatgacctat 66240
gtagatgtga ttctgggtcca accccacatt agaactaagc atcccagctc ttgctcagcc 66300
ctaagactca cacacctatg ggcttgacagg tccatttctcc ttttccatta ccaagacaca 66360
cacactctaa catgtaacca ccagtctcat gtggtctcct ccagggtgca ccaatcttgt 66420
aggactgacc cccttcatgg cagcgggtctg ttgaagatac aacgaaaatg ttaggagagg 66480
gcagaacaga tttttttttt tggctcctata ttccaccat ggtaggtact taggttggt 66540
aaagtagaga cacagacaac agctttccca tcagagacag gggaaacgtt aacagggtctg 66600
gtcacttgga gtcaattcag ttaagtgtct tgcttagaac acaatggaaa accacagaga 66660
aatctggatg gcaaggaaga ggtcatgaat cctgctttcc cagagagatc acaggaaatt 66720
gggtcctaag cagaaaagaa aggctgggag gcttgggtctt cactcataca cagcaggcta 66780
agactgagac taattcagtg aaaagccac atcaggcaca gactccagc ttttgggagc 66840
atggagagcc tgtattttac cctaggatag aaatgtcatt ttggtggaaa gcaaacagac 66900
atatgcataa aaagtgtgga aatcctggta gttttctaag ttgtaagcc caatggcaca 66960
ttcaaatgct gaactagatt gtaaacacta tggacccttg cctcttgcca tttgcctgtg 67020
aacttgata aaccctaaca tgaaatatgg gaaagttga ttggcctat tcacacggga 67080
cttgttcttt tgaatagaag tctttaagac gacttatatt taaagaatgg attaaaaaac 67140
ttaaatacta tcttaataga atctattggt aaccagtgta atcactactg aaagagtgcc 67200
ttagctaaac tagagttgag aatcagcccg gttcccaaag ttccttagca aactctattg 67260
ggttaataat tttgtattat ccctaggaat agtgacatct gcttcctggg tataacctac 67320
tggtcaccaa gaaaccagac tgtttctgtg aggtgcaaga ggtctctgca ttgcaaacat 67380
gaagaccaga caacacatgg ctggttccact cttcatgaac accttcccag tcaactgcaga 67440
aaagtctcat gattttgtcta gcttgaagat atccagggtga tttaaaatgg gcaactcccc 67500
taagcatgct ttactgaaat ggaacccaag aggtctcttg gttctactat gtacagaaac 67560
acacacactc ataaatgcac acacactcat aaacgcacac acatcaaaca ctgattccat 67620
ctgtgctcat acccgtcttt aagaaacaca ggaaccagtt gggtgaccac ttaagagtac 67680
ctggtcacca ggggcaaacc tcaaagttcg gaatatccag tcattggataa cagaaaatga 67740
cagtgtggt aacagagaaa ttcataattg atattttggc atcattttac acaatctctt 67800
ccttacttgc gatggtacag cttatttctcc ctcgccagc cccgatgcag gtacagtccc 67860
agatcatgga gtcttttaga cgctcataag tgtcaccac tcggttaagtg ttcccaggtg 67920
acttgtcaaa gcaagtctct tcagctgagg gaaaaggaa agtccatgtg agcctcactt 67980
aggtaacaag ttttctagtt cactaatccc ctctaaagga aaaccactt ctctattcca 68040
caagtaacat ggtcacctc aaagaataaa atgaagttag agatcacagat ttttatccta 68100
agctctaata aggccatggt ttttgacctt gacatattgt caaattttct gtaaattatt 68160
catattgtag aaaaatctca ggctcttagg aaaaggccat ataaagtact aacttatgac 68220
agagctagat ttgtgaaggg tttcacattc caatctctga ctcagaaagt caaagcaagg 68280
ttcacataaa aagcaaacga ttggccgggc acagtggctc acgcctgtaa tcccagcact 68340
ttgggaggct gaggcgggtg gatcaactga ggtcaggaat tcgagaccac cctgaccaat 68400
acggtgaaac cccgtctcta ctaaaaatat gaaaattagc tgggtgtggt ggcattgtgc 68460

-continued

tgtagtccca gctactcagg aggctgagc aggagaattg cttgaaccct ggaggcagag 68520
gttgacagtga gctgagatag tgctactgca ctccagcctg gtgacagagc aagactctgt 68580
ctcaaaaaca aaacaaaaca aaaacaaaaca aacaattcga agatccccac ctcccacaag 68640
gccttgacag ttgttaatcc tatgatctcc attttcccct gatcagtgct tctctgacag 68700
agttgctata atttatgaga aaaatatcat tggtcctaaa ggaatgagaa ctttgaagtg 68760
gctaattggtt ttgtgttaac tgccgctgac aaatcatttt acgatgtaag tcccaaaagt 68820
gttttctggc tcataaacta acagtttaag atttgagact gctttctaataa aaattaaaat 68880
ttataagca attttttttg tcggaggaca cacaaaatct ttcatttttc tcatcaaaaa 68940
ctttgggaga gtaaagcaaa aagaagactc taggaactaa tttgggaaag attcgtttct 69000
ttgttttaaa ggtaaagca tgtgaaaagc aaatataagc agtttttaaa cttaagagga 69060
aatgacaggt aattctatta gaaaaatggt gtatgtatct ttacttacta atgcaaagtt 69120
taacaattac cacttcatgt attaaaagat actaactatg gggcttggtg tcacttacct 69180
tcaggtttac tctcgcagtt aaaacctcgg cttcctccat aacaagtaca aaccaacgca 69240
ttgcctaggt aggtccgctc cactggtga tttatctgat agtgttttcc attgctataa 69300
caaccggctg caaataattg aaggaaaacg ttacatttgc atttctcctt tcccaaaat 69360
tatggaattt tcttcatgtg aatattgacg tacatatctt tttcctttta actttgctaa 69420
aagtatatata caatgatgct atttccatct gccaggggt ctacataaaa aagttggaaa 69480
cattttggaa ggcagacaat tgaagtcaca tttttaaac atgcaaggaa cttaaatatt 69540
cttttgaggt aatagcaaa aaggacaact ttaagcagct tctcaaaagc aaaaggaggg 69600
tggtctttg aaaaacggaa agaattccaa cctctgttct tggtagaagta tcctgagcgg 69660
ccctaaagag gtctgaacca attacatctc ttccaagtta gttagaaca agacagctat 69720
ttcaatacc ctttccacc cccatgttga aaatatttga ccaattattt aacagaagga 69780
aataagtagc agtaaggacc aactaatcc cagacggaaa atttatagac atgcatcatt 69840
ttctcttta aaggcaagtt ttggttaatt cctagaagga caatctgaac gtgcagaatt 69900
tattagaaca cgggaaagtc gcagagctct cgtcctcaa tgcacggtct tatcatcca 69960
ggcctcaga accagcgtgc tcaaaactcg gtccttttgt gtgcacagct ggtttctctc 70020
agtaaagcgc gcacacactc gcacacacgc gcgcgcacaa aacttcagcc ccaactttgg 70080
tcggcttag ggtccatcc ctgaggcagc ctgtttcagc ccgcggtcag tactcacgct 70140
tgctttgact gacagccacc ggggactggg gctgaaccat ttgctgagcc tgcctcttgc 70200
tcttcgagc tcccgtggag ggcacgcgtg tcccaggca ctggacggcc agcagcagca 70260
gcccgggccc cggaccctta agcatgttga gacggtgggg gagagacgcc cgcaccggga 70320
ggcaagttgc caccaagttt gcttcccttc gcaacctcgc gaaaaatcc cttctaagtc 70380
ctcccggggg ttgtcgcctc caagaaggtg ggggcccagag ggtggggaag gggacgggtg 70440
gaggacaga aggatgcagc aggaccagag aagttgtggc tgcaggtccc ctcttcccgc 70500
tcgocctcg ggttccctct cctcccctg tgcagcacag ccggcgcggg cgtccgagcg 70560
ccgggagccg gggcttatat gggacggtcc cctcccggcc ctctggaggg ccggggcgcg 70620
ggggaggaga gaccgcggc gctgtcccgc ccccgcccca cccaccgcgc ccgcccggcc 70680
gccggccgcc gcagccgacc gcgcgocgat tggcccgggc tccgggtgac gtcacggggg 70740

-continued

actgtgggtt cgcagcgaac aaaagagatg ctgatggccc gccaggactg gggcaagaca 70800
actttttttt ggttttcttt gcggtcotca aacttttttag gggatggggg aagggatggg 70860
aagcggctgg gaggaagg agtggttggc cttgtgtgaa gcgaaggggt tgcaagacgc 70920
tcccccttc cttttctttt ttattggggg tgggtgtagt gtttgaggac attgcgtcac 70980
ctctcttcgg ggtggagggg gggaatctct gcttatagga gtgtattttg tgaggaagaa 71040
aaccataaa atatccacga ccttgcaact gttaagaatc cagacttttc agtcccaggg 71100
cggggctgcc tttccccca tcccgcctcc tttctttgga ctaaagggta ctgactcggg 71160
actcccttat ttgtcttca cagctccctg ttcggacttc tccccttctc agcgtctgtc 71220
ctgtctggct ctgagaaagg gccagcgggc gaacggggag caaaggggaa cagagacctt 71280
taatgagctt ctactaagta ccgggagttc cgcggggcgc tttgcagatc ccggggaagt 71340
tcagagaatg aagttttctt ttgcaaggac aaagtaacag tttacttttt tacacatgca 71400
ggaaagcctt cctttcctgg ggatgggacg cccgtttcca ctgaaagtga aatcaactgtg 71460
aacaacctga aaattggaaa gttggaccag ctgtggggag agaaaagtct tctgtctcctt 71520
tcccacgttc accgtgccag ttacacacaa agcagagatt tttttgcttg aatcattaaa 71580
agccgggtgg gtggggggca ccctatgtac tgtcttgccc tccttcggcc cctttctcac 71640
atttatttgc cttttctttt ttgctgacaa gttccattac agaatgcccc ctccatcccc 71700
cgccgccaga tttttatata ttcaattggt atttactcca agagtgtgga aagttctcct 71760
ttaagattcc cccgccttga actctaaact cagagtggaa agggaatctg aaaagtctct 71820
catttagcaa cacatcacat cagtcctttt tgtcagataa acacaggcac aattgctttt 71880
aaagatcttc agaaggagga gtccaccact tccacagtag atccagagag acccgaactg 71940
aatgctttgc ctctccttg tgttaggagt tcagaggttc catttgcttt aaaacacaag 72000
aagctactaa taaaatata ttatttcgpc aaggtaccta tttctccaa aagtttcacc 72060
ttcagtaatt gccacaggag taattagaac tctgtgttct ataaaatgcc tgacacatta 72120
aagacacaaa attagcattt gtttttattc atgcaaggat ttaaaaccaa accaaaacaa 72180
agcaaaacac ccagcaata tcagtttctt acaaatagtg taaaaaaccc tcttgaggaa 72240
tctttgtgac ccacactcca aattgaaaga gtgccatgga tttccccaaa tttagaatcc 72300
tgtctggggt agaggatttt tttggaattt ctcatcactt gacaaaacag ttttctcttt 72360
tcaaataaa aagagctcct tgtaaaatgc ttggtaaatt tctgggaagc cgagtgtttc 72420
ttccagaggg aaggatcctt ggaatccgg atgtaagcaa gataaacctg tagctaaagc 72480
tttcttaaaa ttcattgatt ctgggggaag gaaataagtg attagaaaag tatttgggtca 72540
accaaaacca tgatagtttg atatatattc tgcactgaaa atacactaga atcatatgca 72600
tagtggagct tgaagtggaa agcccactt acattttgga agaaactttt atatttcaat 72660
cccatagata tctaatttat ttaaggaaac aggatattct tgtctcataa gaaaaaaat 72720
aacatctgac tgatttttat aggaatgctt gggagtttac tcttttactt gaaaggtttt 72780
tttttttttt gccttttttt ttaaacttta gaaacagagg cgtttcttag ttaagataac 72840
agggcttact acagtattct ttaattatca aggtcccttg taactctaag tctgttatct 72900
ctaattggag agataattaa atcttaaaaga ccaactctgac tctaattcaa ccaacgtgtg 72960
tggtgtccag gatcgttggc cttccttaag aattctattt tatccagggt ggttcttctt 73020

-continued

cagtaaattg tattctccta ttggtaggaa atactccttg atttccttct ttctttcttt 73080
ttctttcttt ctttctttct ttctttcttt ctttctttct ttctttttcc ttctttctct 73140
ctttctttct aaaactgtat tcagagagag aaatgtgaca aaaaaagga aatcaataga 73200
gacagaaagt agattcatgg gtacccaagt ttatgggctg ggaacaggaa ttaactgtaa 73260
aagggcacag ggaatcttat tgaggtgatg gaaatctgtg ttctaaaact acattgcagt 73320
gatgactgta caatttggtg aaagcttcaa aattctgggg attatacaca aaatgagtga 73380
attttacggt atgtaaatga tgcttcaata aagttgtttt tttcaaaagt tgattttaaa 73440
aattagtatt tgttcttttt tactcttaac catctcctga ttttggtttc aaagacttgc 73500
tctcaggtag cagcaacata cacaaggcgt cctcatttct tcatttttct gagaggaagt 73560
ttgccttga ttagtagtg gatttcttct acaccagtaa ctccttctga aacctactat 73620
ttgtgtgaag gcatatatgt gtagagaca gaacaaatct tcatttagga ctgtgccttg 73680
tttcttcttg gaattgccat agtgatagaa ttttccaga gcctcaggca atttaaactt 73740
tttgaacaa aaagaagcac atatattttc gggaaactagg ggggtgtggg tggcgggggg 73800
caatggtatt gggaaatatt attattccag aaggtaagg aggagcatga gcaattatgg 73860
aaatctgaat taaattgatt tcatttggtt tgaagattag caatggagac atttgattaa 73920
gcacaaaaa taattgatca gatccagtca tttagaagga ctgaaataaa aacaggctag 73980
gtgtggtgat tcattcctgt aatcccagca ctttgggagg ccagtgtgga agggctactt 74040
aagcccagga gttcaagtcc agcctgggca atatagttag acactcttaa taataatgat 74100
aataataacc agataggtg acaggcacct ctcgtcccag ctattcagga gtctgacatg 74160
agtctgggg gatgactgga gcccagaaga tcgagagatc aaagctgcag ggagctgtga 74220
tcacaccact gactccaga cccagcctag atgacagagc aagatcctgt ctcaaaaatt 74280
taacatttaa cattgaaatg ttaaagtta aaggttcat ttaaatatgt cacatttaga 74340
tgttacattt caaatgttac atttaggtgt aacatttaat tgatgggctg tttttcctga 74400
aacctttggc ttttatcctg ttattaatth cttccagttc tgctagagag gaggccattg 74460
actctggaga gctcgaact tcatttcatc ttcttctata ggtagattag agggagataa 74520
attagggtag tttttttttt ttcttatttt gaattttttt tttttttttg gcataaaatg 74580
cttcccttct tccttgaaaa tggccacggt tacaaaaatg ttactcatta accacagggtg 74640
tgagaagtta aaaataaact tccctgctgg cagccctcgt aagtagaatg tctgaaaagt 74700
ttgtcaagaa ttgacagggt gcctgcaacc agggagaaaa tactggataa gggctttaa 74760
ttatgccatg aaagctaacc agttttcctg taataggcat tttgtttaca ttttaagtga 74820
ctttattttt taatcagcac tttttaaaag ccttaactta tgctttttgc ttcttaacag 74880
gaatattctt ttctcattag agtgatgtaa ctccaacag gcattggcaa ggtgtttaac 74940
ttatccagta aaatgggaac taactactgt gtagctggaa catgtaacca aaactgtcct 75000
ggagaacatt tcatctttcc aagcaagta gcagcttctt gtaaaaatth taattttgta 75060
tgttatttca tgttgctggc attaaagagt tttttgttgt tgttgaaagg ctgaatttaa 75120
aaaaaaatg aagttttgct atatttaaat tattttctat cctttttgca ttaacatact 75180
ttaaacatct catatgcagc taatcacaaa agttagttag gccaaatgct taaaaataac 75240
cacaacatt atctagtaaa taacaattta tgccaacaat agactattht gatatgagct 75300

-continued

gtcaatattt atgcctacct gtctaattag tgaattaatt agttaaata aattaagaat 75360
acattcaatc gaattgtaaa gatgccatat acatgttatt gtaaatttat attgaactat 75420
tttctagata agtttgatat attctttcat taagtatgac ctttttactg agataaaatg 75480
gacatttaaa atcattacta ttttaaacat taaaaactaa ggcactgctg gccgggctg 75540
gtggttcatg cctgtaatcc cagcactttg ggagaccgag gcgggctgat cacgaggtca 75600
ggagatcgag accatcctgg ctaacatggt gaaacccctg ctctactaaa aaaatacaaa 75660
aaaaaaaaa aaaattagcc aggcattggt gcaggcgcct gtagtcccag ctactcagga 75720
ggctgaggca ggagaatgac gtgaacccgg gaggcggagc ttgcagtgat ccgagatggc 75780
cccgtgcac tccagcctgg gtgacagagc gagactctgc ctaaaaaaaaa aaaaaaaaaa 75840
aaaaaaaaa aaaagaaac taaggcactg cttacatgca tgcacattta agtagtcacg 75900
aagtcaaaa gatttcagca cctagataga aataaatagc tacattttat taaactctct 75960
aaaccagatc cagtactaag tgtttcatat gaattttttt catctaacca tcaacaaact 76020
ttaaaggtaa gcactatcat cctctcctct tacagatgag gatataaagc ttagaaaaac 76080
aatgtcaaaa aatgctactc attaagcaca gatgtgaaaa ggaaaaagaa aatgccctga 76140
aggcagctcg catgagtata atgtctgaga agtttgttga taattgacaa agaatgaagt 76200
tttctcaatc acaagccttg tagaaattgt ggcgtcaagt tgtgtcactg tcaataccag 76260
gtgattagtc aagtgattac ataatggtct ccctcagaac caaaggatg ggcaagacaa 76320
ctggctaaag aattttgtgg gtgacttctt tcacatgaga ttaatttatc aaaagatgtc 76380
acacaggact ctgccaaactg acacgctacc agtttccatc tgtttagacc aaggccattc 76440
ttctgggaac caatttcctt caaacagcct ggctgtgttt cagcattttg aatcaaatgt 76500
ggaagattgt taccatcgac tctccactgg gctggaagt ttgctggggc agaggctctg 76560
tcttgtttcc tctgtatgcc cagcactagg taccaggcac ccttgataa gtacttgga 76620
aatgtcaaat gaatgaatgg aaggagagaa ggcagatga caaacactct gagattcaca 76680
ttcagtgact atccagccag gcacagtct aagtgtctct catatgctat tttattttat 76740
cttcaagct actgccagaa gtagatatca atagtccat tttactctct gaagctcgga 76800
gctgaaagat ggtcctagag cttatgctgt aagaagggg cagagctgaa attgaaatca 76860
cggctcctct gacacaaaat tcttgcctct accgaagaca tacatggctt tccccaaaga 76920
gactttgcc tatatgttaa aagaacgttg gaaacactct caggcagtt ctttttatct 76980
tcctctagat cgttctgcct tgcttctggg cctgggcct gacctgtga actgcgggc 77040
tgcttctct gtttctgga ggacagagga cggagatgag agttgttct gcccactccc 77100
tcctgtttct ggcttcagt ccagagttct agtggcagct gcctttccct aggacctgca 77160
ttatcaatgg ggcaacagt ccccaagagg gtaaaagtgg gttcttgggg tagaagggga 77220
gtggttaaaa aatttcaacc ctacctaata aaaattttat tccttagttg tgaatttaac 77280
actttttttt tttttttttt gacagaatct tgctctgtca ccaggctgg agtgcaggga 77340
caagattttg gctcactgca acctctgcct cccgggttca agcagttctc ctgctcagc 77400
ttccaagta gctggaacta catacatgag ccaccatgct gggctaattt ttgtattttt 77460
aatagagacg gggtttcatt atgttagcca ggctggtctt gaactcctga cctcaggtga 77520
tccgcccgc tcggcctccc aaagagctgg gattacaggc atgagccacc atgctggcc 77580

-continued

atgaatztat catattagat agactttaaa tttcatttct ttcttgagg gttgtataag 77640
gtagggcatt aatacataaa agaaaaaggg agaaacattg ctctaggaca gcagtcctcc 77700
ccaggctttg gtctccaag gctccagcta tcgctggctc cttccccatg atggcagccg 77760
taggagtgc aaactgccta cattgctagt ccctgggaac ctcagcatcc cttttttatt 77820
ccttggtcat gccacactg gggtaacagt gctttcacia aattctccc atggaaatca 77880
ttttaattcc atttctgct aaaaccccgg ggatccagat tcttttttt tcttttttt 77940
aagatggagt ttccccttg ttcccaggc tggagtgcaa cggcgtgact tggctcactg 78000
caacttccac ctcccgggt caagcgattc tctgcctca gccttacagg tatgcgccac 78060
cacgactagc taatttttgt attttagta gagatgagg ttcaccactt taaccaggct 78120
ggtcttgaac acctgacctc aggtgattca cccgcctcgg cctccaagg tgctgggatt 78180
acaggcatga gccaccatgc ccagcgacga tccagattct aaccatcaa tgatgacatg 78240
agcagtcacc cccatagttt taagtcttaa attatagcag aatagactta ggacacacga 78300
attcttttcc aaacttagca actagtatta atcatgctgt gatttctttg tatttcccat 78360
gacttacatg gtcttatttt tttctctcta ctgaatgcat gcgcttttaa cactctatct 78420
ttcccaaagt aaaacgctca gatgatagaa gcaaatgaca tagtcatcat acaatgacgg 78480
agaaaagaaa ccagaathtt atatttaaat ttatgatatt ttagacaggg tctcactttg 78540
tcgccagat tgtagtgcag tggcaagatc ttggctcact gcaaaactcca cctctcaggg 78600
tcaggcaatc ctctacctc agcctccgga gtaggtggga ttataggagg tgtacaccac 78660
catgccgggc aaagttttg tattttttgt agagatgggg ttttgccatg ttgcacagga 78720
tggctctcaa tttccgggct tgagtgatcc tctgccttg gcctccaaa gtgctgggat 78780
tacaggtgtg agccaccacg catggtcatg tcgacgtctt aactgtcttg gaaataatht 78840
gattttgtcc aggtaathtg gttcacgcaa agaagttgag atcggggcac atcagaaaga 78900
gattcatctt ttaaagtgtc tcttaaatct acaaatgcaa tatatgtaa agctaaaggt 78960
ttttatata ttttcacaca tgtaaaaatg taagttttgc ctttaaattc attttagata 79020
tgaaataatc tagctgagag catgtaaagt cgaggaaagc ttattcagat ccgctctagt 79080
tcagcatgtg gtaaattcct aggtgttcac aatattagt tacacaaatt ccttctcaag 79140
aaagaaaacc ttggacagac ctggtagact agtatcactt atgccttata gagactttct 79200
gtcaaaaacca acaaaaatc tgcataatct ttattcatag aataaaatgt gatattaggg 79260
aacacaactg tttctggaat gcattgaaac tgatagtatt ctatagattt ttctacttaa 79320
caattactta ttattactat tattattatt gttgttttg agactgagtc tcaactctatc 79380
accaggctg gagtgcagtg gcatgatctc agctcactgc agcctgcacc tcccgggctc 79440
aagcaattct cctgcctcag cctcccagc agctgggatt aaaggtgcc accccaacgc 79500
ccagctaatt tttgathttt tagtagagat gggtttcacc atgttgcca ggctgtctc 79560
gaactcttga cctaaatgat ctgcccgct tggcctccga aagtgtctgg attacaagtg 79620
tgagccactg cactggcct caacttacat tacttatttt tcaaaaacta gagaatccat 79680
taathtttgc ttgtttgaat ttaaagtga ttataatatt tttgttagtt atatatgagc 79740
agtatttcc tacagctcat cttaaagata tttgataatt tatcttctat ttttacagc 79800
aaatggtcac atttttgtca gttgacaata aattcttttt ttttttttt ttttttttt 79860

-continued

ttttgtagag ttggggtttc gctgtgtttg gccaggctgg tcttgaactc ctggcctcaa 79920
gtgaccaccc cggctcagcc tctcaaagtg ctgggattac agatgtgagc caccacacct 79980
ggcagacaat aaattcttga aagattcctg atactttacc aacttacaca ttaagttaat 80040
cttcccattt tctttgtcca attaagtttt aaagcaaaat aatataaata tttatagacc 80100
caaatgtaa acacatttca gttttgaaga acaatataaa gtaaatatac tatggaatta 80160
caattagaaa accaagtttg cactgatatt tttaaaactt ctctggggaa cttagttttt 80220
caacttaag acacaagtat taaattagaa ggtattttcc ttcactaacg aaaatttttg 80280
tttttaatt atctattctt gtctttcaaa ttttaagtttt aatggcggac caattaatgt 80340
tgcttaaagt tacaacttag acaaacagtt aaaatagctt agcaaattgt ggttctgggt 80400
gttctcctat aggtaattaa gagggcaatg actcaagatt taaaaatta aataacatgt 80460
ccacacaatt ttctctgaaa tgttcccttc agatttttaa aacaggttta ttgagatata 80520
attgccatct cataagcccc acatacctgt atacaatttg acaagtattg acatactttt 80580
gacaattttc tattactgtac attttatcctt ttcttttttc ttttttttac accaccctc 80640
tccacttatt acatatattt tgactgttgt tttaatattt ccttttagct tatatagcat 80700
taaatccttt tctatcagtc tgagcttttg aaagaaattt ttttttttg gaacaagata 80760
tcaaagagta aaattgaaga aaagttaat ctcaaactca tgggggaaaa gacatcacag 80820
gaaataaatg ctaaaacct ggctttttta tgtgttgaat taaatagtta ttaagttaat 80880
tggtagataa ttatgagatc tattatcatt taatataatt agtatgtaa taaggtgata 80940
taaatatgt tctgtgagaa gttgtcactg atggtgattg ttatctctgt gtaaaagtga 81000
gaaaaccaac agaaagatga tgattattag tggcaagatt aatgatcatg ggccttttct 81060
aattcttatg gcccaatctg gtatttttct tctcctcatt ttgactttac ctatactttt 81120
ttttttttcc acacaggcca gcatattatt atagctcttt caaaaatttt atattgattc 81180
ttttttcccc aacagtactg taaactcttt gattgtaggc actaaattaa ttccttttag 81240
aaagactttt ttttttcttt ctaccatttg aagtttttta agttgtgtgg tggagactgc 81300
tagttgccca cacagatcca ttctcttctt tcacagttag tccattgta cacagttagg 81360
ctaaatttta cagtctcctt tgtagttagg catggcaata gaaatagttc ctcatcagtg 81420
gactgtgaga agaagcagtc tatcccaact ctggttctgt gccttagttt aataacgtca 81480
ctctccaga ttccctgtgt cattccact agctgaaata cacaacacag aggacacca 81540
gcttctaacc atgtagatga tcaaaatgcc ttgggaaaat ttgggacctg aattatcatg 81600
aagagcagac cttgactatt gacatgagcc attcactttg agactgttgc atgagaagta 81660
gacaccctcc tttattgtag aaggcctgt gttactgagc agctttgtta cagcagccca 81720
gctttcatag taaccagtat gatgcactac cttccttaca tgcctagttg ctacagctgt 81780
ctgttcctgc tgactgcatg tgatttgttt actagctttg ccctggtatt tgcaattcta 81840
ctttcgagtt tcctctcttt tcctagcttt ccataaaactc cacgcagaat tttagcttct 81900
aagctaaact ccataaata tttaccattt gtttttaaat aataaattca agtcttccgt 81960
taaatattct tgagcacctg ctaagagaac tgttatgggt tgaatgatgt ccctcccaaa 82020
ttcagctgg accccagtc tttcagaatg taacctgatt tggaaatagg atctttaaag 82080
aggtcatcaa gttaaaacac gtacatcagg gtggatccca atccaatag actgctgtcc 82140

-continued

tataagagag gaaacagac acacgggcag ggagaatgcc aggaaagatg aagataaaga 82200
tcagggtag gcatctacaa gccaaaggaga aacgaagatt taccgcaaac caccagaagc 82260
tgaggagag gcatggaata gattcttctt cacaaccttc agaaggaacc cacactgctg 82320
acatcttgat ttggacttc tggttttgag aagtgtaga caatgcattt ctgtggttta 82380
aggcctccat tctatggtac ttgtattac cctagcaaat gaatccaagc acctggagta 82440
ttaatgataa caagagaatg agtgataaat ttaggttcat agatagaact ccagaggcat 82500
gacataatat gcccaaaggc aaatgcctgg ttaactgaag ggctgaattt gttgattctt 82560
gctctttctt acagatggtt ttgtgtcaat cattgtgtgt aggagacgtt gatagataag 82620
atgtaggctc tacttttaag gaatgtaaaa tgtactagtc agttgacaga gataaaaagg 82680
ttgatttttg tacactattg gcaagtgtgt cagttgaaa aatgtatata tgtgcacaga 82740
tttttaggag tgaaaatga ggtaatcttt taaaaggaca ttttgcaat atttaaatg 82800
cacgaaccta agacacagta gactgaacac aagcatttcc tctacacct cctgaaatca 82860
cacctatagt tcagagatga aaacaattta aacctacaag aataaagaga ataagaaagg 82920
agacagtaga tgagagacaa caacattttg gaaaaaggaa cacaggtgga gaagtgatac 82980
ctaaggtcac agaattaatg cctgcaagcc aaaaaaaaa aaagaagctg attgatattc 83040
aagaaacccg gaaagcctca atcatgggag gcatcaggta cctctgaagg gagggcagga 83100
aagtgaggct gagaggtgtt tgaaagtatc tgcataaaga ataactggac gcccttgta 83160
ctcctgccc tacttctacc agccatggac tgatccactc ctcttgccc agtaggacct 83220
aggaaatttg ctgtgtgat agattgaacc agaaaaacc caaactcagt ggcaggagg 83280
gtgatagtg tgaagcacag ggctaaaaa aatggagagt taagtgaagg tatacatgct 83340
tactctccat ttggcagatt gaaaatgttc tctctggact tagtctcttt ggagacaata 83400
atctgtgat ctgacactgg ggagtcctgt agtgaaatag ccagattcta accccatcac 83460
ctacagtgtt gccaatcagt cactaagctc cactcctctg tgcagtttcc aatcagatct 83520
ttagtgtccc gattctaaac atgaatgat gtcaaaggat cagtcaccaa accttgaga 83580
aaagcctctg acacaaaagg ttaaagtga tttatagaat acaaaacaaa agaataagag 83640
gcaatgcaa aatagaaga aagtgaaaa taaaaacaaa ctaaaccta aatcatgac 83700
tgtagaaaga gaagatattg tgagaaggaa acaagaatag tgatctatac atgtatgaat 83760
atttagagac caagaccca atcctataag attagcatcc ttattagaag agacaccaga 83820
gacatctccc ccacctgta cgccttctg accccaggca agcacagagg aaaaggatg 83880
taacaggaaa agagcccga ccagaacga accctgccag acctgatct tgactaccag 83940
cctccagaac tgtgagaaaa tgaatttcca ttatttaagc ccctaattt gtggtatctg 84000
ttatggcagc caaggagac tagcataatg taccacattt aataaaatta aatgaacact 84060
aaaaacagaa tgcatttgcc ttttgaggca gcaattccaa atttctctct gttctttaag 84120
atatgtaact taacctagga gtcagcaaac ccaggggcca aatccagcct acctttggt 84180
tttggcagc ctgtgagcta agattgattt ttagaattta tgcagttgaa aaatttaaaa 84240
ataaatatatt atgacatgag aatgttatat gagattcaaa tttcagtggt cataaatgaa 84300
gtttaaaaat atatagccac atctatattt aaaaatgtat catccctggc tacttttctg 84360
ctacagtggc acagttgagc agttgtgaca gggacccat tgcactacaaa actgaaata 84420

-continued

tttactatgt gttcatttag gaaaaagttt gctgactgct ggtgtagacc atctgagtga 84480
aatttccttc cagattccta taatattttc tccaatcttt tatagttgtc gtgcccaaac 84540
tgaattaaca gcaatatttt aaatatctca attttattgc gtctttcca agtaattcaa 84600
attggttttc ttaggaattg ctctctttca acattttgat tttaaagata atatggtttg 84660
gatattcacc ccctccaaat ctcatgttga aatgtgatcc ccaattgtat tagtccattg 84720
tcatactgct atgaagaaat acctgagact gagtaattta taaagaaaa gagatttaat 84780
gactgggaag gcctcacagt catgaaggaa ggcaaggag cagcaaaaggc acattttaca 84840
tgggtgcag caagacagca tgtacagggg aattgctgtt tataaaacca tcagatctca 84900
tgagactcac tcaactcac aagaacagca cggaaagcc tacctccatg attcaattac 84960
ctcgaacttg gtccctctca tgacacatgg ggataatggg agctacaatt caagatgaga 85020
tttgggtggg gacacagcca aacctgtca ccaatgttg aggtgtggtc tagcggaggt 85080
gtttgggtaa tagggcagat ccctcatcag tagcctggtg ccaacctcat ggtaatgagt 85140
gagttctcac tctattagtt ctggggagag ttgattgtta aaaggagtct ggcacttcct 85200
cccttttctc tctttcccc tttctctctc catgtgacat acctgctccc cctgcacctt 85260
ctgccatcag taaaagcttc ctgaggcctc cccagaagcc aagcagatgc tgtatattgc 85320
catgcttgta cagcctgcag aacctgagc caaataaacc tcttttctta accaattgcc 85380
cagcctcaag tttttcttta tagcaatgca aacagatga atacaaaagg caaaaacagc 85440
ttaattacat aattatcaaa tccattacaa tggcaataaa cagtttgta cccaacaact 85500
gacttctggt aaacatataa ctcaaattat aagagcacia gagcagtgac ctgctagggt 85560
ggaatcatga gctttgagta taccttgac atcaccagca cattttatgg agggaaagca 85620
gagcacagtt gggaagtggg ttgcatggaa gttgcgtaag agattttatg ttgtaattag 85680
cctagttatt tcaggtatct gatttttttt ctcgtccca ttaagaata aatctcctaa 85740
gagcagggat aatattttga tttcttttgt attcttcttt caagcatgga gcaccaagtt 85800
gggcacatga caaatgttga ataaatccct cttgaaagg tagatgatga agcttctaca 85860
tgccagagat gaccattctt ggggtaagtc agctctcagc cacctagaaa ggttttctca 85920
accgctgatt ttagactctc aatctgtttt gatttgcgat gctgtattct tctttgccat 85980
tctcccaaat gacacgtcag ctttcatgat tcatttttat gttttcttgt ctatcgctc 86040
aatgttggat ttctgcctaa gcagcaaat tcaagaaat ctcttctggt ggaattgctg 86100
atgtgaaaga agcccatgag gttcaatga cctcatagtg cataaaacaa tgtgagtga 86160
tatggggagt actatttatt gttcaagaca aaaatgagta tgttgtgaaa agcttgctca 86220
gtttatggta agatgaaatt gctgaccgaa gactgcctaa aactcaagtt ctgaaatgctc 86280
attaatatgg acaataaaaa ggaactgtct tagaaagttt attgttacac ataattgtaca 86340
taattcttgc aataactatt ttgattttta gctaccag agtatgatca agaactgtga 86400
gtagtacaat ctagttaag ttgccttgtt ggttctaaga gaaaaattct ttggggttga 86460
aaattagccc attttgaac acatctctgt gtacaattac tgttaaaagt tgaatatgtg 86520
aaatctggty gcatagtaag cactgataga aattaggttt tttattaatt tattattgat 86580
ttgccttatg attgtagaaa gggatgtat ttgtataatt taatgactta attactctca 86640
ttgtaaaagc tgataaaat tatatctggt ctattcacat cattagtata ttgcaaagat 86700

-continued

gcttgacagt ggcaactgatg gctttgggtg gtatgatatt tatcaattat aatttttata 86760
taattgacat tccagttcct gttctgtcag tataatatctg tgtgaacttg attataactta 86820
aatctttgtg cctcagtttc cctatttata aaacaaatat gttgaattag attaattcctt 86880
attccttggtg ttcaaaaatt acataactgt aaaaaagctt ttgtcatta taatttttta 86940
acctcaccta aaaatgccca aatataaaat aatgatatat gtgcacttcc taaacattca 87000
tgtgcacatt ttgcatatac ccagaatatt ggagcacaca tcttcgcaag gaatatagat 87060
ttatatccta aatttgcttt tgtttctaag agaagagatg aagagctcat accttaccac 87120
cagcaagaat aaaagctttg tttctctcct ctgtttaagt aattaaata tctttgctcc 87180
tttactttat gatttcccat tttgagggtc cctaggggtg agaggagcgc tcattaaaca 87240
actcatctct ttcctgcctg aattccacat agagaactaa ctgatggggg ctaaggtgaa 87300
ggaagtggct cggaagaga gtgaaggag cctgcctagt gtaccagag ttagggagtc 87360
tcatctatcc tctgcagtcc ctgctggcca ggagtggccc acacagggca cattgtgttt 87420
ccagtaaagc caactgcacg gacaaacaga ggagcagcag agatcgcagt gacagcctc 87480
cctaccagg agagaaaag agagcagagt tgatcaggct ggtctggata ttttcccaa 87540
atttacaag ccagtgcgtc acttcacttc tccagaggag aaacaaatg ggaatttg 87600
cccagaggag caccaagact gtgcttacg aaaatcacca ggaatggga aatgtgttc 87660
cctctaaacc tagaatcctc gtcactaata gctgtgttct ctataataa acttttgaa 87720
attaaaaag tataaatgc caagatatgt cgtgtaaaac aatttaagt aactttcaa 87780
gtagacttgt ctgtaatgag accactcttg gttttaaatt tgcagatgct ctttgcaaag 87840
aaagtagcat gtgtttaaca tttaccttca gtattaactg tgttaatact accattgctt 87900
attttgacat ttttagtttt aacatttata tttggaggaa ttaaaaacg acttgatttc 87960
aaaggaggta taactcattt tcacaatctg agttcatgtg aatgttttag aagaaatttc 88020
agaaatggct ttgaagtat ctgagaaaat gtttacaatt atgccatttg ggaattaca 88080
ttctatgtgt tggaattggt tctatggaga tacaattgc tgtacaatgt agaatgtagt 88140
ttgatagaat gagaatgta gaattttcac agtgcctaaa gactggtaat taaactataa 88200
ctagcaaggt aagaacaaaa taaccacatc ttaagtacag tttgactttc atgactaaag 88260
ataaaaatta caggagggta ggattttatt tcaactataa gatgaacttc ctaacggtaa 88320
gagctatcta agattgaaat gggctgccat agttaacaa aaaacataa attacttggg 88380
gatagtattt gaaggggagt cactcatgct tgtattgtac tgcatgcctt ccaagatctc 88440
ttccaattct aatacttgta ttctataaga gcaacagaat catgcaatga tgaaacctt 88500
ttaccaatt ttttctttg tattagttaa tagacaatac taagagtaca cagcatgcct 88560
ctgottgcaa taatagaaaa gaaccaagaa catttaaaat gggctattga aggagaccat 88620
ttgaaataa tttttacaaa ttaagggtgt aatcatatc tttttctaa gcagattttt 88680
aaaaaagatc cttgcaggat cttttttact gctagtattc ctttgttagt tctgaaaaga 88740
gattctgtgc attactctgt aaatagtact gtatattact aaaattttat tggtaaaatt 88800
atttgctga ttctaaggtt tagaagaaaa aactgaaca aaatttccca aagaacggac 88860
tgtattaatt tgaaagcaca gtcctgggga gttcccaagt ggtgggttc tagtgacctg 88920
gcaattattc atcaattaac cacagocact tattatctgt atgaatggaa caaagttgat 88980

-continued

acacagacct tgaggcaaca atgatcagga tatatggcta gttttcttat tccatctcag 89040
agttacactt cagaaacaaa tttccttaat atttgatatt aagtgtcatg ccctcaatga 89100
caactaggag gagctgtttg gtctcgaactc tgcctcagaa ctctttccat gggatcaaac 89160
attgcacctt gtactgaatt aaccttttgt gtcttttgac cacatacgtc aaaggtaaat 89220
aaaaatata gtattttaag ctctgttaaa aaaaaaacac aattgctggc ctcttgata 89280
attaagagtt tataacaatg aagattgctt tgccttttgt tctcaagatg ccctcatggg 89340
ctgttttaga aggatgacct gttcaatfff cacctcggtc atcatctctg agtctctgtt 89400
ttgcacatgg ggaacttcag tgcctcttac ctcttctttt agtttgaana taaagggttt 89460
taaagtctca atctagggtc tgcactgaaa aacgtctttt ctctctagtg gttatgctaa 89520
tgcaagtctc ccaatgttac tctgcataac tgaatcacta actctgcatt tcaaagtgcc 89580
aacactggta aacctgaaaa tttattaatg taacaacat tgtgctggaa acaaaaaatat 89640
ggtttctgat ctcaaacatt ttaagatcac attggagaaa agacacgaac tgtaatatct 89700
tcacaattgg taatagcagc agttagaaac aacagggttag gcagtagagg acattttctc 89760
tgcacagaa ctgagaaatt gtcaacatat atactttttt tgggtgtgtc attttgggga 89820
gtaaagaaca tggggatgag taaaatttat aaacctttat ttttttgcgt ctctctaaaa 89880
tgcaagtctc ttaagaacct atcagcaaat aatattagct tgatcatcct caaataatgt 89940
tagattgagt caaatgaagt aattatttct gtaggtcaca acagtcaata ttggcaattt 90000
catatgggtc acctaatttt aaaaagtggt gcattagttt tctttttttg agacagagtc 90060
tcactctgtc acccaggctg gagtgcaatg gcacaatctc ggctcactgc aacctctgct 90120
tcctgggttc aagtgttctc cctgtttcag ccttccaagt agctgggatt acagggcgac 90180
accaccatgc tcagctaatt cctgtacttt tagtagagat ggggttttgc catgttggcc 90240
aggcctctgc caccatgtct ggtaattttt tgtatttttg gtatagacga gggttcacca 90300
tgttgccagc gctggtctca aactcctaac ctcaagtgat ccgccctcct cagcctgcca 90360
aagtgtctgg attataggtg tgagccactg catctggcct tgggttagtt ttctggggct 90420
gtagaacaaa tgaccacaaa tctagcccgt taaaacaaca ctcatattt gtctcacagt 90480
tcagtaggtc agaaatctgg gtgggcttga ctgggttttc tgcttagggc ctcatgaggc 90540
tgaaatcaag gctgggctcc tatctgaagg ctctggagaa gagtctgcct ccaaaactcag 90600
atggattggt gccagaatc agttcctggt ggttgttgc agcgggtggt ttcagaaact 90660
agaagcagct ctccagctct tgcacgtggt cccctccatc ttcaggtagc agtggccttg 90720
tgctttgaat ctctctgaca tttcctctgt caccagtctg agaaagctct ttgcttttaa 90780
aaggttcatg tgattagatt aggccacca gataatctgc ttttaattaa cacaaaaata 90840
actaattaat aaccttaatt acatctgcaa aatcgctttt gccacataac ataacataat 90900
caaaagtgca gtgagaggtt gtgtagtaaa gtggctaagg gcatgggctt gagaatcaga 90960
ctatctgggt tcaaaagtcat gctctatcac taaattaggt gacactggac aaagcttttt 91020
agcttctccg tgcttctgca aaacaagcat catctttaat aacctgcctt gttggttggg 91080
ttttaagggt atcattctta acaaatattt agcagagtgc ctggcacata atagttgctc 91140
aatgagtaac atttagtatt attattgtag aaaaagaaaa tgagaaagaa aaagtgagag 91200
aaaagaaga agtgaagaa aattgttga gacagagtga tcaaggaagg atttataaag 91260

-continued

gaggtgaggc ttgagttgga ttaaggttaa ctgaatTTTT gtaagcatga agaggggcaa 91320
tgtattcatt ctaaattata acaaatttac caataaaagg tttgtatTTT gtagaaatga 91380
gaaactgatt acatctgctt atttggacag tgaatggaac cacatattct gcctgttggc 91440
aaacacaaa tctgtatagt ttatTTTctt caaaggtaag taatgagaaa attaaactgac 91500
aaggcacact ggatcaacag gttactatgc tataTTTgat aaataggctt attcctgata 91560
cctcagactt actccaggat tTTTTTTTT tTTTtaatag agtctcgtc tgtcggccag 91620
gctggagtgc agtggcatga tctgggctca ctgcaacctc tgccttcag gTTtaagtgg 91680
ttctactact tcagcctccc agtagctggg attacagtg cctgctacct cacgtggcta 91740
atTTTtgat tTTtagtaga gacagtgtt tgccatgtt gccagactgg tcttgaactt 91800
ctgacctcaa gtgatTTTc cgcctcggct tcccaaagt ctgaaattac agtcatgaac 91860
cacccgcct gcccaggaat ttggtaagcc atgtttacat atacagtga ccaggtagga 91920
accctaaata tgctTTTcct tggattgtc caaatgcctt ggtcattctc tcccagcata 91980
atccaagcaa agataaaaa tgaccaatga ataaggctca gtagtcaatc ggttgaatgg 92040
gaaagcacca cagccactta gatcataaaa agcccaagtc tgcagccact cagtcagtcc 92100
attctgctag caaatttctg tttatatac ttctctcaa gaaggggtc tcccaaagct 92160
ggccctagat agttcctcag ccgatcggg aattaatagt caattaacta agtattaggt 92220
tgatgtgaaa gtaatcacgg ttttgccatt gaatatgtta cattgaatat aaaggcaaaa 92280
attgcatca ctttcatacc aacctaatat taccacctct ttttctaata tgtgacctg 92340
tgataagaga aaacccccg attttcctct aaagacctat tcattctcca ctttcagtgc 92400
atgcttttca ggtcctgctg atctcagggt gggattttga ttcaggctcc agtaatcaga 92460
gcactgcagt acctgattac caacagcctt tactatttgt tcatggctgg gcttgcagtc 92520
caaaactaggc caatgaaaac ccttcctgag aatttttaa gaattaatta ataatacac 92580
ttcogtttg aattgctaaa gtggtagaat gtaggtgtga tgctactggt gattactctg 92640
cctctgtgtg ggaacagcct tcctgagaat gaaggcacc caaagaaaag tgagctctga 92700
gatggaagg ctcagctctg aagatgtcat ttgagcatct atacggtgct gcattgtaca 92760
tatoatgct ctggacttct aagttattga gccaataagt tattatattt tgtttaaagca 92820
catttgagtt tgcatttctg tcattgtcaa ccaagagttc taacagtcgg ctcacagagc 92880
ctccgatgta acgttcccac tgtaaacatt cagattgatt aattaattg aatcccagct 92940
atagaacct gatagagaga atctctacc tctgatcaa tcagacaggc tcacagagca 93000
caaacatacc tgttgactc tgggtgggtg gtgagggggg atagtatca tggaaagga 93060
acagagatac ataacaaaag ccatttggt caaatgtgtg tgacttttat aatcagacaa 93120
taagaatatt tttttgaaat aatcaactaa aacaatttaa gtagattga ataactatt 93180
aacaatttt tgcTTTTt ttacattcag tggctgcaa gagTTTTta tcaactcaagt 93240
gattttagtt aacaattaat aatgaaaac taagcaaaa aaatgttcc tctaagtcatt 93300
ttgacataag agaactagt ctcagatggt acacggagaa ccatttttc atttccacgc 93360
ttcaaatct cacactctt tgccaaatc ctacaaccag tccttaatga cctagtctca 93420
gtctcaagt acctctctc ctaaggcatc ctgtctccc agaaacagga atgtattgct 93480
aagtggacct tacagattat ttaggcagc tttcatatga gatgaaaagt tcaaaagag 93540

-continued

tcaataatag tgaaacattt ttttaaaagg ttgggggggg gattctaatt ttattgaaat 93600
cccccaaaga acttgatatt acaggtgaca tcagttccag ctcagactgt agctagtgtga 93660
gtcctctggt ccacagattt tattcctaag tagatccttg cctttagaag tgagtgtcaa 93720
gcctattcag taatttcaag ctggaagaga ccttgggaat tttctagtcc atatctgtca 93780
ttttatTTTT atTTTTattt tttttgagac tgagtcttgc tctgttgccc aggtgggagt 93840
gcagtgggtc tatctcgggt cactgcaacc tctgcctcct gggttcaagg gattctcctg 93900
cctcagcctc ccgagtagct gggattacag gcatgtgcca ccacgcctga ctgatttttg 93960
tatttttagt agagatgggt tttcactgtg ttggtcaggc tggctttaac tcctgacctc 94020
tggggatcca cctaccttgg cttcccaaag tgctgggatt acaggcacga gccactgcac 94080
ccggtccata ccattattt taaagaggag gaaactcagg ctcagaaaat atggaccaat 94140
aatcaaagca gggagacacg tacatcccag gaaggaaaaa tggctggaac aaaggcgtg 94200
agtcaagaaa agatatactc agggacaaaa ttttgaaaat gaatttggat gtgactttag 94260
ttgcttaatg ggttgactgt aaacttgaag ggtaattat taggtccaaa ttatagaagg 94320
ttttgaatgt taaggtaga ggttgttctg ttgtttttaa ctttataaac acataaaaagt 94380
agggaaagtt gaataatgaa agccatgtgg ttaccaccca ttttctacaa tgatcaactc 94440
atgatcagtt atattccttc tatgactcta cttcctcct ccttctcttc tgtctcccag 94500
attattttga agcaaatctt tgtcataata tagctttata aatatttcaa catatttcaa 94560
gccccaaact aaagacttag agggtttcta agtattctgt agataagaag aagcctttga 94620
atattttcaa ccaggggagc aacatgttca gaactgcaga aggctcttcc ggcagttcca 94680
tatgggatg attggagtag gaagtggcag ggtagtgtaa gcaactgata tgccacattg 94740
tccttgtcac agaagcgttg tttcctattt tccaagttta acgtatgctc tatagtttcc 94800
cttctggta ttaagggta gaaataattt atagattctt ggccaatgga gatatttccg 94860
ttttagaggt tttgtcattc tgacatttca gtttcacgct ttgtagtctt tttatgcatt 94920
gaggtgtatg gcaataatgt aagtgtgaat ttttctattt ctccttatgt attaaatatt 94980
caccttttat tttattatat tgcctaatga ggacattctg tgtactagtt tgatgtttat 95040
ggaagaaaaa caaaagcttc cctagctcta aaaacttgcc ttttaagttc ctcatctcta 95100
taagtagcaa aacataacac ataaattatt caggcaatac aaaatgaaaa taaatacatt 95160
tcaattggct atctaagaga ttatttctct gtgcattgag tcttctttta tgtagtcaact 95220
gcaaaccat gaagattga aatattttct gaatgagaaa tctttcaaca tgaagcagca 95280
cacacagcac tgtgagcaga aactaaaata aacctgtct ggcctcaagg ttcattgttg 95340
atatttagag ttggaaatac tcatttcagg gtacattttt ctttaacca gtcaccccag 95400
agtggaaaca caacatagag aatagataag attgtctgtc tggctagctg aacacactgt 95460
ccttcacact gctgagaaat ttgcagagta tggtagaggt ttttagatct tgaactgta 95520
caaggcactc ttctgctttt aatttatgct ttgaagattt atcaaaaata gaaagaccaa 95580
ataactcact ttccatttaa agtgctcacg ttttccatct ggtacattgt tatgaataac 95640
atgtctgcat gacataacag aaaatttttc ataccagtgc gggctgcca tcaatttata 95700
atcagaaatg tcacatgacg tattgaaaag aaatcagaat tacacacggg tgetgcattt 95760
gccttcttat atttcttacc agaagocctt ttccttcttt tcagctgocct ctgtgaattg 95820

-continued

tgtggaatca ttagagaaaa acccaagtag agagctagca attttcaaaa actgtaaaaa 95880
agttccttga ataaccataa caatgggata ttttcagtag ttttcctggt atctttttgg 95940
ttgcattttt aaatcaaaac acaaaattga tagatgacag ataataaaaa atttttataa 96000
ccagatctct tctaattgctt gaaatttcag gtgaatcaac agaagcaat atgagcatga 96060
ttatgctgct ataattgggtc agaccagtac ctcatgctaa ctaaaagtct gtctatgact 96120
gtggcagtca tgtgtatttt ctgaaaaaga attctagctt ctctacagga tttcagactt 96180
agaaagttag agacctgccc aaagctttcc caaacttcta attaacagag ggccaaggca 96240
gagaacttga gttctcatta ctgatctccc actaatggga tgtgccacct taggtgtatt 96300
gttaatttct gtttacagtg tttcactttc tttatttcca gataagaaaa gattctaatt 96360
acaaaaacaa ttttaaaaga gtctacacat agaagataat aattgctatt atttttactt 96420
atctaaatct agggaatcat gattgccaac aatcactcag caccaacaat ttactgaatg 96480
tttatctgcc tgcattcttc ttgtggctgg aaatgtagca gtgaaccaga ccgagaccct 96540
gatccgctgg gtatatagct gggggagtca gacattcatg aaaatgactt ctagtgccac 96600
agatatttca aaagatgtag taaagagttc tgtgggtctg aatgatattg tgtgctaacc 96660
aaatatgtga aggtggcgaa ggtctcctag aagaattggt gcaaaacttg agtctcaaag 96720
gatgagctgg gtgaggagtt gggcacagta ttctgggagc tgggaacatc atggacagaa 96780
gcctgaatgc agcggagaga ctgactcact tgtgggcatg aaggagatca gcatgactgg 96840
attggagctc acttcctggt gttttcacct cattgcatgt aagcatcaat ggacctacc 96900
ctgcttctgg ccatatcaaa cagttcaata gtttggcatc tttgtgatca gcttacagaa 96960
gtgctgtggg ctttgtcaaa ccagtttaga agctctagct tactggctga gagaacttgg 97020
catgagagga tgctgacgag ttagacagag gtctgattgg aaggggctgg gtgctgagcg 97080
ggagttggga aaggaggagt tagtgacgac aacacattca agatcattga aggttagtga 97140
tatggtttgg ctgtgttctt atgcaaatct catctcaaat tataatcccc ataaccctcg 97200
cgtatcgagg aagaaacttg gtgggaggtg actggatcat ggggcccgtt tccctcatgc 97260
agttctcctg aaagtgaata agttcggatg ggatctgatg gttttataag tgtgtgacag 97320
ttctccttc atgtgttctc tctctctctc acctgctgag atataagtcg tggctgcttc 97380
tccttctgcc atgattgtaa gtttccggag gcctcccagc catgtggaat tgagtcaatt 97440
aaacctctt tctttataaa ttaccagtc ttgggaggtt ctttaaagaa gtgtgagaac 97500
agattaatac agtcagagag gcctttctga gcataagttc ttgactatac taagggcaat 97560
tgagagccat ggaaaatggt taagcagag aaaagtttgt ctgagaaggg gaggaagaga 97620
ttctgaggtg cttttaagaa atccacttg aggggcccag cgcgggtggt tacgcctgta 97680
atcgagcac tttgggaggt tgaggcagcg gatcacgagg tcaggagacc gagacctcc 97740
tggctaaaa acgtgaaaccc cgtctctact aaaaatacaa aaaattagcc gggcatggtg 97800
gcggtgctct gtagtcccag ctactcggga ggctgaggca ggagaatggc gtgaatacgg 97860
gagggggagg ttgcagttag ctgagattgt gccactgcac tccagtctgg gccacagaga 97920
gagactacat ctcaaaaaaa aaaaagaaaa gaaaaagaa atccacttgt agccgctaag 97980
gtctggagct cagagaagag tctgaccca agcccagttc tccagagtca ttttgattct 98040
aagaaacctt gatattctt caataogttc cttttttttt ttttgtccac catagccctg 98100

-continued

gttggtttct gttgcttgca atgaaagaac ccttacaaat gcaataatgc agaagtaaca 98160
atgaagatga cactgattca cacaagattht aatttggtac atgagtaaat aaacaaagca 98220
aaatgaacag aatttactat actgcatccc tgctgggagt atgttgagct atctggcctg 98280
gaaggccttt gtgacctgtt gaaagaatat acatggattht acttcaggag aaaaataaag 98340
tgttatttta aaaatggatc atgttcattg taattaagaa taaaggttgt aaaaactatg 98400
gattgtggcc gggcttggtg cctcacgcct gtaatcccag cactttggga ggcgagggca 98460
ggtgaatcac ctgaggctcag gatttcaaga ccagcctgac caacatggtg aaaccccatc 98520
tctactaaaa atacaaaaat tggctgggca tgggtggtggg cacctgtaat cccagctact 98580
tgggaggctg aggcaggaga attgcttgaa cctgggaggc agaggttgca gtgagccaag 98640
atcacaccac tgcaactccag cctgggcaac agagggagac tttgtctcaa agaataataa 98700
taaaaaaaat tgtgggttgt atagaaaaa attacagtgg aagcattata tgtgaacttc 98760
attggagaaa tatttattat gacatataga gttatatata tgagacataa ggttaaaaag 98820
gtaccttact ttttgcata cttgtatagt cagtgaagg cttttttctg tatacattgg 98880
aaaactaatt tttaaattha atatttaaag catggccggg cacggtggct cacacctgta 98940
atcccagcac tttgggaggc cgtgactggt gaatcacaag gtcaggagat caagaccatc 99000
ctggctaaca cagtgaacc ctgtctctac taaaaatata aaaaattagc caggcatggt 99060
ggcagggccc tgtgtccca gctactcagg aggctcaggc aggagaatgg cgtgaacccg 99120
ggaggcggag ctgacagtaa gccgagatcg ggccaccgca ctccagcctg ggcgacagag 99180
caagactccg tctcaaaaa ataaataaat aataaataa ataaataaaa taaagcagat 99240
agcagttgat atatatgat tgggcacaaa tttagctagac tatcttccca gggcaaatga 99300
tgtcatthta tttgactcct tttctgthtt agctgagthtt tttccagtha cthtaataaa 99360
gaattaatga taaatthttt ctctacttgt attacttcag cagattaatt tthcatttht 99420
thccagthga gattaagatg gtagcatccc tatacaaaat cthtattacc atattcactt 99480
ggaaaaatat catgaaaagg thttactaaa atgaatgagg tgaagccatt cagthttgac 99540
tgtaagthtc agaggctgtt tcatcaatgc gtaagtagcc taaagacttc actggagtht 99600
tataagaatg caaaagagaa aagcaaacct acgaagacgc ataaaaatta tgaactctgg 99660
gagthtattha atthacacat ggtthcttht cataaattca ggcagaagcg taacctatth 99720
ctcaccccta gthtttgaa cthcttccct tthctatgct tggthacgth gthattcggth 99780
ctagacatac catccathaa gcaaccaatt thcacgthaa gthgagthgth tgtgththga 99840
thtctgtht cctctthaaa cctattgth tthcatctct gthgathcatca aattatatac 99900
atcatgggth tctatththa gagatathac ccagthgth tcatctthac tctcatctta 99960
gthgcatctc atctthctc ctgthgagat thccaccctt thctgththca catgaththt 100020
thttgcctta gagcaattgt ctcaatgthc aattgagcct thgataaggaa gthctthcagga 100080
aattctthga thgathacag catgththgta thctthacca ththaatgag acaaacathga 100140
aatcagththt aaaacctthaa thctgththaa ththaatthaa thgathththt cthctgathg 100200
ctctthcaag gaagcataat taattcttht thcacaththt thctagthct cathaacctc 100260
ctccccaaaa gaagthccac thththctaaa atthcaata thgactactg aagthattgt 100320
cctaaagath ththaattha aatathattht caagthgthaa gaaaththaac thctthataca 100380

-continued

aagctcagaa acaagagata acatgatgta gtcactctacc aagcagcact tctcttattt 100440
gggaaagtaa cattctattc ctcagtcaca cccatccatc tgccccctcc aggtgtctga 100500
caagagcatc tctaaaggta gtggggctgg gaggagctac taactttttc tgaaccaggc 100560
aactaaagt taaacctcat gccagtggtt tacaactttt ttaacttaca ttttatctaa 100620
ttgagaaaca aattaaag ccagatggtt ctgtgactga ggttcaaaaa attagcccat 100680
ccctaaaca tactggtcaa gtctctttca attgcaagca tccaaaatct cactctaact 100740
ggcctgagca aatgagttaa catgttggtt cctataattg aaaagttcac aggcatacaa 100800
cacagctgga tgcagatggt taatatgtgt catcagaact tggactcttt ttcaatactc 100860
ctggccctgc cttctccaca ttggcttgtt ttcaggcagg actcctccca tccacaaata 100920
tacagtgata agatgtctag cagcagttcc aggcctttat tctcctaatt tagcagaggg 100980
aagaatgctt ctttctccat tgatgcactc tttcagagac tcagtaaaca gcatctgtct 101040
ttgtgctgta gatttggaac cattgagaaa taagaaaata aggagcccaa ataccaccct 101100
tattactgat aaagttggcc ttagatgtca gttggcctca gatggacttc ttaatatgaa 101160
ccatataacc aggcagactt tctactcagt cctggactgg gtctcccaaa ggggcctctg 101220
tgagcaagct ggccatgta acgtgcccc tttattcttt tcaaatttac caattagata 101280
agtcattctt ccttgaagac tgggaggagg ggttgagaaa ggccagagct gttgcatcta 101340
gccaagctgg agatgtcatg aatgtggaga aagttctgac ctagtatctc ctggctcttt 101400
ttttcttaa cccctgaaat taactattaa acccatgagc agccagaaat gtggctgcct 101460
ccctccaag acagttttat tccctccttt atagggccac cccaatagt gaaaaccaact 101520
tcattgagga tccctctgga tcttcagtgg tcaccagag ctctgaagtg ggttagagca 101580
aaacaggcat aactcagaga tattgtgggt ttaattctag acgaccctaa taaagccaag 101640
attgcaatat agcaagttgc aaaaactttt tggtttctca gtggatataa aaatgatatt 101700
tctactacac cgtagtcttt taagtgcgca atagcattct gtctaaagaa acaatgtaca 101760
taccttagtt aaaaatgcct tattgcagcc aggcattggt gctcacgcct gtgatcccag 101820
cactatggga ggcccagggt ggccgatcac gaggtgagga gttcaagacc agcctgacca 101880
acatggtgaa accccgcctc tactaaaaat acaaaaatta gccaggcacg gtgggtgtgca 101940
cctgtaatcc cagctactca ggaggctgac acaggagaat agcttgaacc tgggaggtgg 102000
aggttgagct gagccagat cgtggcactg cactccagcc tgggtgacag agggagactc 102060
tgtcccaaaa acaaaacaaa caaaaacca aaaaaccca atcttattgt taataaatgc 102120
caatgatgat catctgagcc ttaaacatgt tgtaatat tctgtgtggt agggctcttc 102180
ctctatgtta atgactgctg actgatcagg gtgggtggtg ctgaagtttc ttagaataag 102240
acatcaatga agtttgccac atcaattgac ttttcctttc aaaaagttt tctctgtagc 102300
ctgtgatact gtttgaagc attttaccg cagtagaact tctttcaaaa ttggaataaa 102360
tcccctgaaa ctctaccact gttttatctg ctatgtttat ggaatagtct aatgaagctg 102420
tcatttcaac atgttcacag catcttcaca aggtgtagat tccatctcaa gaaaccactt 102480
tctttgttct tccataagaa gcaactcctc atctgttaaa gtttgattaa gagattgcag 102540
caatccactt acatcttcag gatccacttc taattctagt tccctaacta tttctaccat 102600
atctgcagtg acttccctca ctgaagcttt ggatccctca aagtcacca tgaaaattag 102660

-continued

aataaatttc ttccaaactt ctgттаатgt tgaaaatgtg atctcctccc gtgaatcatg 102720
aatgctctta atggcatctg gaatggtaaa acctttccag gttttcaatg tactttgccc 102780
agatccatca gaggaatcac tatctatggc agctatagcc ttaccaaatg tattttctta 102840
accataagac ttgagactgg gcgccgtggc tcacgcctgt aatcccagca ctttgggagg 102900
ccgaggtggg cggatcacga ggtcaggaaa tcaagacat cctggctaac attgtgaaac 102960
cccgtctcta ctaaaaatac aaaaaatta gccaggcatg gtggctttcg cctgtagtcc 103020
cagctacttg ggaggctgag gcaggagaat cgcttgaacc cgggaggtgg ggttgccggtg 103080
agccgagatc gtgccactgc actccagcct ggagacagag caatactccg tctcaaaaaa 103140
aaaaaaaaa ccgaaaaaca aaaaacaaaa cttgaaagtc aaacttagtc cttgatccat 103200
ttgctgcaga atgaatgctg tgттагсааg catgacagta acattcatct ccttgtagat 103260
cactagagct ctttggtgac gaggtgcatt gттаатgagc agtaaatatt tgaaaaaac 103320
aaacaaaaa aacttttgct ttgagcagta gttctcaaca gtgcacttaa aatagtcagt 103380
aaacatgcc atgctgtaaa taaatgtgct gtcacccagg ccttggtgtc ctacaaccag 103440
agtagattta gcataattct taaggccctt aggattttca gagtataaat gagcattggc 103500
ttcaacttaa gtaatcaatt gcattagaac ctaacaagag agtcagcctg ccttttgaag 103560
ctttgaagcc aggcattgac ttctccttct tagctatgaa ggtcctcaat ggcattctct 103620
tccaatagaa gactgcttca tccacatgga aaatctgtgg tttactgcag caactttttt 103680
ttttttttt ttttttttt ttttttttt gtgagacagt ctactctgt cacctaggct 103740
agagtgcagt ggcaccatct cagctcactg aaacctctgc ctctgggtt caagcaattc 103800
tcttgctca gcctcctgag tagctgggac tacaggtgct tgcagccatg cccagctaat 103860
ttttgtagtt ttagtagaga tggggtttcg ctatgttggc caggctggtc tcgaattccc 103920
gatctcaggt gatccgccca cctcggcctc ccaaagtgt gggattacag gtgtgagcca 103980
ccatgcccg tctactgcag tgactttatc aattatctta ggtatagatt tttctggat 104040
aattttctac atcagcactt gctgcttctt cttgctcttt tatgtaatga agatggcttc 104100
tttcttaaa cccatgaa caacctctgt tagcttcaa gttttcttcc acagctgctt 104160
cacctctctc agctttcata gaattgaaga gagttaggac ctactctgga ttaggctttg 104220
gcttggtgga atgttgtagc cagtttgatc tatctagatg gctaaaactt tatatcagca 104280
agaaagcagt ttgctctct tatcattcgt gtgttactg gagtagcatt tttcatttcc 104340
ttcgagaact tttcattgc ttttacaacc tggataactg gcacaggagg cctagcattc 104400
agcttatctt ggcttttttc atgcctttct caccaagctt aatcctttat agcttttgat 104460
ttgaagtggg agatgtgtaa ctctttcttc acttgaacgc ttagaggcca ttgtaggggt 104520
ataaattggc ttgatttcaa tattgttggtg tctcagggaa tgggaaggcc caagaaaaga 104580
gagagagaga gagagaaaga aagaacaac tggtcagtgg agcagtcaga acacacaaca 104640
tttatcaggt tctcctccat cttgtatggg cacagtttgt gttgctccca aatagttaga 104700
atagtaacat caaagatcac tgatcacaga tcaccataag agatataata ataataaagt 104760
ttaaaatag agaattgcca aaatgtgaca caaagacaaa aagtgagcac atactgttga 104820
aaaaatagtc caatagattt gttcaatgca gagttgccac aaatctgcaa tttgtaaaaa 104880
atgcagtatc tgtgaagtgc aataaaaca agcacaataa aacaagttat gggccgggag 104940

-continued

ccgtggctca cgctgtaat cccagcactt tgggaggcca aggccgccag atcacctgag 105000
gtcgggagtt caagaccagc ctgaccaaca tggagaaact ccatctctac taaaaatata 105060
aaattagctg ggcatggtg cacgtgcctg taattccagc tacttgggag gctgaggcag 105120
gagaaccgct tgaacccagg aggcggaggt tgtggtgagt cgagatcatg ccattgcact 105180
ccagcctggg caataagagc aaaactccat ctcaaaaaca acaaaaaca aaaacaaaca 105240
accgcccccc cccccaaaa aaaaaaaca agttatgcct ggacattaat tcaaccaact 105300
tcaaagccgg tttgtatcaa agactgaata ttcacttctt attctctttc ccatcccacc 105360
tgatgtcctg gatcatccat tatagtcttt cagcgggtcaa atgaatcat aaaaacactc 105420
cactaatgaa atgcatagaa atttcagtca tacattccaa caatttagag atccccaaa 105480
tgtaaccata acagtttttt aatgatcaaa gtcctgggaa gaaattataa agtccataat 105540
aaagctccct tttaaattat aagccacaga tattaaagtc tagagaatat tatattctat 105600
aaataaatgt cacgattcct gtgcacggaa aggtcaaagt ccaaatgtct gatgttaaag 105660
ttcaaaagcc attctccaaa atgtactatt cagtagctca gtcttattgt gtgaggctgt 105720
gggtatgttg gaagagggaa gcgggtggtt ctttctggac ctcagaaatt atactagttc 105780
cacacccaat ctctcacttt ctaaagaacc cagagagtta acaggacaca cagatttctg 105840
tagcactggc atcacactcc tcatttttag aaaacaagtt aataccaac cttctaaagc 105900
agccatcctt agcaaaatta tgtaccaaga actgagtatt ttatgctgcc aaaaagcgtt 105960
acagacatat ggaggtgttt gaagagagga tagtaggtga aaatcttttt tttttaatct 106020
accaattttt tttttttttt tacaattggg gtctcacagc ctggtacggt ggtcacacc 106080
tgtaatccca gacttttggg atgctgaggt ggggtgatca cctgaggcca ggagttcaag 106140
accagcttgg ccaacatggt gaaacctgt ctctactaaa aatacaaaaa ttagctgggt 106200
gtggtggtat acacctgtaa tcccagtgc ttgggaggct ggggcagaag aatcacttga 106260
accggggagg cagagattgc agtgagctga gatcgtgcc ttgcaactcca gcctgggtga 106320
caagagcaaa actccttctc aaaaaatagt aataataata ataataaat ggggtcttgc 106380
tatgttgctc aagcttgttg tgaactcctg ggctcaagca atcctcctgc ttcagcctcc 106440
caaatgctg ggactacag catgggccac tgtgcctggc ctcccccaat gctagttgcc 106500
tggactccat gcctataaa cagatattca agtagaagaa gattctcaga aagaaagaag 106560
ttcattctg cactttgtat ttaccataat ttgggagaaa catccatttc tgcccactct 106620
ctccattcct ggagagtggc atttgagaca ttgctgtctt cttgctcagt ttgcaaagt 106680
gcaacaacag ccgagctcac taagtcttgt ttaagattcc tgtttttctt ctcatactgg 106740
aaaaacttta attgctgttt tctcttgatc cctgtttttc tggcaactct gggctgcctt 106800
cagggtaggt tcagcctaag gagatagaat ctaccctttc tgggaacag tatatagatt 106860
cctgaggaac aagaaatctg cctccttaga gaaatactat ttccaattca gagccaaatt 106920
ttgactagat tttcagcgt ggggttggg accattggg ggaaggagtt acggataatt 106980
cttgagcctg tgataaacc ttgcaaat tctctccatc actgctctct ccctgaaaaa 107040
gcagagtgga atttgcat tgcgcacacct agcacaccag acttaggcc aaaggaaaca 107100
cattaaatct aaaaactccc agggctctgt agtgctgag acggagttgc ctggtgattt 107160
aaaagaagac aaatattatt ttgcctatca atgctcaatg gtccttaaag acacctttc 107220

-continued

cctgtcatca attcctggga tgagttgttc ctacttagga aaaggtgttt ccgcttgca 107280
aattggttct ggatttccca tcagcctagc aagattggaa cttagacctg gtgtacatta 107340
aattattcag aaattatgtc cttacctaata acaaaaaaaaa agcaatcttt aaatttctgg 107400
ccaaatatca atttctctgaa gtcacgtaac tccaaatcta aggatctcaa gtcttctagt 107460
gagaattcca gaatatgttt cttatagatt tgaggaagta aatcatctta gaataactgt 107520
ctcccttctc ccctctctcc ctggtaccac atagtctttt ctccattgca aatagaggaa 107580
atgacaaaat gtgctatgtt ctaattttta tgaaatctgc catgatgtat tttatttatt 107640
atgaccctga gtcccagggt gggtttatgg tattctactt tctgcctatg caattgacga 107700
gtattatgtt tcctctctct tttgaaaagt gatatgtttt ggctgtgtcc ccacccaaat 107760
ctcatcttga attgtagtcc ctacaatccc cacgtgttgt ggaagggacc tgggtggagg 107820
taattgaatc atggggcgag gtttttccca tgctgttctc atggtagtga ataagtctca 107880
aaagatctga tggttttata aatgggagtt cctctgaaca agttctcttg cctgatgcca 107940
tgtaagatgt gactttactc ccccctatc ttccaccatg attgtgaggc ctccccagcc 108000
atgtggaact gtgagcfaat taaacctctt tctttataaa ttactcagtc tctggtatat 108060
ttttattagc agcatgagaa tgaactaata cataaagttt gtaagtagcg ctatgtgcta 108120
tgtggtcatt ctgctccaag gctcatgggt ttgtcatgaa ccctgcatgt tacacaatgc 108180
taaagaagca attgtctgca cagggtgtctc ttaaattaac ttaagaatat ctatcaagca 108240
cctattttgt gcaaggtggg agggatacaa agaccacagt ctctaattat cttaagcttt 108300
tagtctagtg aatatcttcc ctcttacttt taagtgcca agtgcagaa cttggtgctg 108360
tttgttgaat tccaagaggc aatggagaaa atgtaggaat aaactacatt ttcttctaca 108420
tttaagatgt aaaggaaca ttttcttagt agaaaatcct tacttttcag aaaataactt 108480
cttaggtaaa gcacatcatt ttaacccttt ttgcagaaaa aaaaaataca atgcttctgg 108540
ttcaaggtgg caatttgagc acatatgttt gaatcctcct ctttcccaaa taacctgaa 108600
aatgcaggag acacttttta gcattgctga aaccacaaca gggtgacatc cacatgttca 108660
aactgagtg aattcatgga ggatataatgc agatagaaga agttgagtg tgggcaagac 108720
gaccaatggt tttctgcaca agaagaagcc agtcagagaa gtagggtgaa gcaccttcta 108780
ccccagagag ggagctggag gaaaggggtg gctttggggg ggtctttgag gtgcaaat 108840
ctgtgacact ggtcatagct cacatagaga gcgattgctg ccacggctct taaattattt 108900
gaaactcacac attacgtagc tggctattcc taggcaaaaa gccagggctt cctttgctga 108960
atcctaacga ggcagttatg ccaggtagga aaggaagtgc ctgcccagc tagctagatg 109020
ccgcaacaga ggaggcaagg aattcctcca aagagagagg cacacttggg tgtgatggcc 109080
tttggtttga agctggagtt ccttcttact cccaagtgc atgggcccacc agcctgagaa 109140
agcagtgaaa atgctttcct ttgtctgttt tcttgttttt gttttatca aagattttaa 109200
caggtacact tgcaatttct gatgaaagca ttttaaccaa catagataca cctctatctg 109260
tttttgcaag ggtaggtaaa tagaggtgaa tggcaggaca gataagtgag aacccccagc 109320
tgtcaaaag agaggaccat tatgagaaaa ggtgttcttg tgagacagtg attataaaag 109380
aggactaaaa tgtctaaaag ctcatcttcc aaaattgatt aaataaagt cctcatcgcc 109440
ataaaatggc agccattaga atgatatgat tagaaacatt tattgcatga aaggattttc 109500

-continued

acaatatatt gcaaagctta aaaaaagag agcaagaaaa caaatgaaat aatagcagaa 109560
tgaatgcatt gtgtgtatgg atgggaggaa ggcgggagtg agggttggca ataataaggaa 109620
gatcaatact aaaaaattat acacccaaat agttaataag gcaaaaagtt aatataagtc 109680
tggaattga tttcaatgat tttataatct ttttgatgat gtatttttaa ttttttcaca 109740
caaatgaac ataaaggat atgttttct taaagtaaat attgtttctc agaaagctta 109800
cttcccagaa agacactaga ctgaaactat attatggatg cgttcttaac acttaacaaa 109860
gagaggggta atttcaatc tatttaaaat attcttcaat gcaggaagaa atggaaagta 109920
taccaatca ttttatgaaa ctatcatac tctggatc aaagcctaac aaaggaggt 109980
ggaaaaaat caagcaggag ccaatttcg ttgtgaatat gatgcaaatt tttgaaaatg 110040
ataatagaat tgtttttagt attttggtaa gtaataatgc tatggccaag tagcgtttca 110100
tctcaaagt cgaacatatt ttaatactag caagcttatt aacaaaattg ttaagcacat 110160
cagaaagata agctaaggct cttagatgct ttatatgtt gacacacaaa gcaagagaaa 110220
atggctttca tttcaatctg gcactttgct gcgatggtac attcaacaga gaaagagggc 110280
cctgtcacag gtgaccccca acaaggatag ccttgccagt tacaggtggc aggctgggag 110340
gttgccacaa cctagcgagg agaagactga tggctcagaa tgcgtactc actctctatt 110400
gaagaagttt ggtcctctaa aacgactaaa caaacaagc aacaacaaca actacaacca 110460
cctaagtttt accccactct ccccttcgga ccctgaaaga cttgtgcagg gatttaaacc 110520
atctctgaca aaatccccta tcaaacagac aatgaagaaa actttaaaaa ataggtcaat 110580
cattttattc ttcagtaatt atataaatag gtaattcatt ttattcttca gtaataaaat 110640
gaattaccta tttatataat tcaataataa attacctatt caaagaaaca gaagcaaac 110700
taaaaaataa gtagttcatt atactactga aaattgtgag ggcatgtgat ttagatcaag 110760
atgtgaggtc aggaaacgct tcctaggggt gaaaacaagt tatttaaaaa atgtgatgga 110820
agcagtaaat tgcagatgta agaggcagaa ataaaatcgg caaattggaa gacaacttgg 110880
aattctccca gcattaagag aaaaataaaa gaatcaatat gatgagaaga aacaacagac 110940
atgaatgagc gatcctgata gttcagttta tgcagtaaga cttttagaag atgacagaaa 111000
cagatctgag tcagtgaaga tgtgatcttc ttagcaaaag ggctaagaaa atgccaggca 111060
cattaacaaa cacattccat acatagtcat aaacataaaa tttctgaatt tcagagacaa 111120
aggagaatct ttaaattgaa agttaaagg atcctgacat acagctgcag agacagacac 111180
tgtattgcct cggggagaac tccagcccca aaggtgagta ggtgatggag aagaagcaga 111240
gtttgacatg tacaaagcta aagtgagtct ctaaattttt caaatattcc aaatattcct 111300
ctctctgtga aagaaaatag atagtgtctt tccaaaatat gacaaggatc caaagaatt 111360
atgtttcatt accaataaga aggggtgaaa tgaaaagtag tttttgttac tgtcaataat 111420
aaaaaacaca ttttgatcac taggctatag aaattctctt acacgtctct gtatagataa 111480
aggcattaca aaatttttgt catttggggg gtgattaag aatatgcaac aaaaaatgta 111540
gaaaaatag tataatattg cagaagtata ccagacaatt aattaatata atgctatat 111600
ggttttctg gactttctga gttttgtcgt atgttctgat atttgaagat tgtgttgcc 111660
aggtgtggtg gctcgtgcat gtaatccag gactttggga ggcccaggca agtggatccc 111720
ttgagctcag gagtttgaga ccagactggg caacatggtg aaacctcatc tctttgaaa 111780

-continued

taaaataaaa ataaaatcca aagctcatac tccttctgca cactccattg acacataata 111840
tcttgtatatt gggaggggta cccttgaggt tttcaggttc aagggctcag ttgaatatcc 111900
ctcacaaggg cctagtgctc ttagaacttg gaagtttcaa aaacaaaaaa taaaaaaga 111960
aagaaaaat tttaaaaatt agaaaaaaa gaaattaaaa taaatattta aaaagattgt 112020
atgtgtgtt ctttctcctt ttaaagtcaa attcatattc atacctaat ttaaaatcaa 112080
agtattgttt atatacagta aaataagtca ttataattgt tttttttttt cttaaagtta 112140
acccctaaaa ttgtctaagt ttcaggcatc caaaaccatc ctgtacaaaa cccagggtgcc 112200
ctttctgcat ttctttatcc aaccaagaaa taaatcaaca taaggaagta aaaccttgaa 112260
aaggggatta agaactgaaa ggattctggg tgatatcaca attggaaaat atttttaatg 112320
tctacataat tttttttttt tttctgagat ggagtcctgc tctgtctccc aggctggagt 112380
ccagtggcca atctcagctc actgcaagct ctgcctcttg ggttcacgcc attctcctgc 112440
ctcagcctcc agagtagctg ggaccacagg cgcccaccac cacgcccggc taattttttg 112500
tagttttagt agagatgggg tttcaccgtg tttagccagga tgatctcgat ctctgactt 112560
cgtgatccgc ccacctggc ctcccaaagt gctgggatta cagaagtgag ccactgtgcc 112620
tggcctacat aattgttaat atagttataa aacctagtgt aagtacaaaa agtaattttt 112680
gaaagagaaa aatactataa aaatttatta ttatataatcc gaaatattgg atcattttat 112740
aagatgtggt atggagtgg gggagatag aagaggacta tatttcccat agtcataag 112800
ctttgtatatt agagtctggt gaggatggta tgtggttttg tcgtttaacc taacattggt 112860
gtaatatctt acatgtgata acttgtgtat catgggtatt cacaaatcaga atggacattg 112920
gttggaaaca cccacatagc catcctagag catactgaaa caaaacatta tcttagatgt 112980
aaataatcac tggtagaaaa ttgcaattct aaatcattag attttttaaa aaaagcaaat 113040
tctaaaaatct tcaattaata tccagttatt cttcagttag aatgttggat ttgggaatac 113100
aataactatt tagtggatca aaatattcct atcttatcct tgacatcatt aagccaaggc 113160
aaatagagat acagggacac attagctcct atactagatc attagaatac cagtggtaaa 113220
aacataacag ctacaagct tttggtgct tcaatgttgg tttcaaagag agcaattgca 113280
gtaaagtct gtgaactttt gaagggtga tctcactttc ttttctccac tccaggaatg 113340
aaagcaaac cttttgaaa aaatgttatt catcatctat cagtgccttg actttgttct 113400
tcccattaca ggttcattgg tttaaaaagt atttgctaaa ataggcaaa gcccctttc 113460
cccctcaaaa gcaataaaa acccccacac caaaaacaac aataaaattt tctgtgaagt 113520
gctaactaat ttctgcattt gttatataga aaatgttgaa attataattg gaaaagttagt 113580
gaaactcaga cttggaactc tgaaacagtg attctttcct gtacaagggt gatattttta 113640
atcttttaga cttttaagtt ctatataata tttttttgtt agttgttata gaaaacacaa 113700
ctctagaaat acatcctacc ttcctcttat cctgaaatct caggtaata cttgatcact 113760
tgtgggttat ggcatagaga tggcattcca attattcatc cgcctgaaca tggcaggagt 113820
attgttccca ccccctttt tttacattac tgtaaaactgt tgctactatg tttcaattcg 113880
tttctggct ataaatgtgt atgtgtgtgt gtatgtgtgt cctatctggt cctgtatata 113940
aaattttcta gtaactcaag attgtgatac caaggcaaat gtttcagca tcaaaaatga 114000
atatttcagt ggctgtcat ttttaatttt tattcttttg gctttcattt aggaatagtt 114060

-continued

tagttcagtt gcttgagtac agttgattga atacagttgc agcaactaat taaaatctgt 114120
tataagttag gggcacttta aatctgtgct cttttgtctc cactatgacg cattatttta 114180
agcccagagt ttagaacctg tacaaattcc ctctcagtct gaagctttga tgaatgatga 114240
agaggatgcg gtattactca ctttgaactg aaaaaagtt tgttaaaata taatgttcta 114300
taaaaaggtg gcagaaacag attctttctc ttttcttgtt gaaaaatagg aaggctccct 114360
tgccattggg tttctgtttt aatggttggg ttttctctgt aaaggtttct ccctgcagga 114420
agtcggagct tcagaggagt gcaaggctac ggtgtttgtt atcccactg cctgtctgtc 114480
aagtcgctga ggattggctt tgttctctca ccaaaggcca cagcccctgc cagggaacct 114540
cttgtccagc ctgccatctg ggtctattct ttcctccctg tatgtttcca gacctagggc 114600
tggcagcagc tcccttctat tggtagctc gaggtactgt gcctgccatg tcttcttget 114660
tcatctaaac cctgcacatt cctttgtaaa aagtggtttt aattaaactc acctcaaatt 114720
ttccagtgtg agtgtgcctt ctgttttgtt ttcaggctct tgggtaaggc agtcatttaa 114780
ataaattagg cttttgggc aatatttatg ataagggaga gagaataggg agccacagag 114840
tctctgatga cattctctgg caattctctc cctttcaact ccattgattg ccctaattgg 114900
ctcttccttc ttctggcccc cagtttctc atgtctgact aaagccttc atttcccttc 114960
tccatgaaat catctctact tcagattaag tccagtactc tagatataat gctctctctc 115020
atctcctgaa cttctcataa actttttgtc tcctatgcct cattctcttg aacaccgatg 115080
tcaattaatt gtaagttagg aaagcaattt tgacatacta acctatgta aggcacaaaag 115140
gtctactaat gccttaataca tgtgtagcac accatggagt actgacttct gaaaaataaa 115200
atgacttatc ccaggttaca tagcatgaac aaaagagtga ctctctaata atgaaaatat 115260
caccctgaaa ggcaggaggt gttattagaa agatattatg aagcaacaag cgcaacctaa 115320
agtcacacaa taggagctctg tcagctcaac ttaaaagtaa ggtttgact gagacctcag 115380
tggtaagttt aatgttcaag actatgcaa aatgagatca tttctcttcc tagtaataag 115440
aacaaaagta gaatggatat tggaaaactt gatacaagaa atgagcacct actataataa 115500
ttatgtctag atgtgagcaa tctccattta gatcttgggt gggtaacaact taacgtgta 115560
aaacttaagt agaaatctga agagtaagga attggacata ctttgataa agttaagacc 115620
ttccgaagga taaggaccat tttcagagcc atgttgcttt ttattctctc gatataaaaa 115680
gtttggggtg tcattaatac ttgtagtcta atttctgatg gtttgatta ttatgtattt 115740
gacgccatta acgtcagcac tcagatacgc tttacataca atatcttaag agcttctgtt 115800
ggtagtgcca tccaatgtag cccagaaatg aatgaaaaaa tctcactttg tcatatgcac 115860
tttatgcagt ttgattaggg catgcatgta tgttgcatag caggatttta ttaaaatact 115920
gcatatcaat tcatttgcac gggggaggcc agtaatgtcc ctccagcatc tacatatgat 115980
ttctagcacc ccaaactaga aatcatttgg ctcaaaaata gaacttgag aacagctag 116040
tttcaacatc agagacagag acctggggat aggagagtgt gaaccagag gagatttgca 116100
ggagaaaatc ttttagaaa ttcaatcctt tgaaaactct ccctaaaact tccctaaagc 116160
cctcatatac tgaccattga ggtttccatt gaatatttta cactgtttat tttaccattc 116220
accattcatc cagaagatgc ctaccaagca ccagatgcca ctctgggtgc tgcggatgca 116280
ttggtgagaa aaaaatagat aaactacttc tacctttagg ggagtgggg gcagggaaat 116340

-continued

agaccataag caatacatat aataaatcat taggccatat agtgggctag aagatgataa 116400
gtgttggtta aaaagaata gagctgggta agggggctcag gaatttggga agttatcatt 116460
tatcaagtga cttattgag caggtgacat ttgagccaag acttgaagga ggtaagaaaa 116520
tgagccacac agctacctct ttaacactga ccgttaaatc agcctgttaa agcagctctg 116580
cgctgggccc ggtggctcac gcccgtaatc ccagcacttt gggaggccga ggcaggtgga 116640
ttgcctgagc tcaggagttc gtgaccagcc tgggcaacac ggtgaaacc cgtctctatt 116700
aaaaatacaa aaattagcca ggtgtggcgg tgtactcttg tagtcccagc tactcgggag 116760
gctgaggcag gagaatcact tgaaccacag aggcggaggt tgcagtgagc tgagatcacg 116820
ccattgcact ccagcctggg tgacagagtg agactccatt aaaaaaaaa aaaaaaaaa 116880
aaaaaaaa gctctgtctg gttgaaatag aatgcaaata gcataggtca ttttaagattt 116940
tacagtagtc acattttaa aagtaaaaag aaataggtaa atttatttta acaaaatatt 117000
ttatctaacc tattctattc aaaatatttc agcatgaaat caaaattcta gaaaaaatt 117060
agatttttgc attttgtttt tcatactaag tcttagacat gtgatgtgta ttttccgatt 117120
acagcacatc tcaatttggg ctagccacat atcacatgct cagtagcaca tatggtcagt 117180
ggataccata ttgggtggta tagccttaaa gagggttgaa aaagtgtcac ttttgaagag 117240
catactcaaa aggagtgagt ggtccctctg ctttcccac cctccaggct aggcaggggg 117300
tgtttgtggg cagtggagag gaggcaatct taatttgggg aaggatgaat ttcctgcaaa 117360
caaagcatta gggagttttc tttctatttt agaccaataa gggtagctat ctcatatcat 117420
cccccttgaa tcagcagtgt gatatgaagg gtcattgtca attctcaggc ctttgtctac 117480
atcttcacc aaagccttcc tctcctttag gaattgtctc actgagagac tggtttggcc 117540
gaattatcca ttcaagatca agcctttatt tgtggtattt tgtatgtttg ttggagatgg 117600
agacgatagc cttaactag aggttaagaa tgtaaaatag tggctttgaa ggatgcaatt 117660
gtcctgtaga tgccagggtt ctaatattac gaccatttat aagtatcaa gttcagcaag 117720
aatgaggaag aaacaacca tatattatag ccatttttgt agttcctagc aatcaactat 117780
gaaaaacagc tgttattagt ttttctttc ccaactagat tataaatcca caaaaatagg 117840
agcoatgtat tcatgacatt ctcttttctt ttgtatcctc catagagggt agtctgagc 117900
tgctatttaa taaatattta tagatcactc caggtgttgg caaacctttt ctgtaaggac 117960
cagataataa atatttcagt ctttgagaac tatactgttt ctgtcacaac taccctacct 118020
tgctgttgca gcacaaaac aagcatagat aatatataac tgatgggcat agatgcatgc 118080
caataaaact ttattgatat atacaagttt aaacatcatg caatttttac atgttacaag 118140
gtattgttct gcttttgttt ttttcaattg tttaaaaatg cgaaaaaaca ctgttagctc 118200
agctgtgtga aaacaggaag tgagctggat taggtctaag agttatcgtg taccaatgcc 118260
tggtttccat tattctcatt ttttgcctag ccattttcgt ttcttttttt cctgtgcctc 118320
tttattacct aattctctct ccctccttcc tttttgactt tccttcatct cactgctgat 118380
tctctatat ctctcttctt tcccattctt cttatcttt tcttctttct tcaaacccca 118440
tagccatttt ctctatttg cctaagcact gtcactctag gtggagcggg ttccagaaac 118500
acagcttcc tttcaggaag aaatagatgg gattaaaata atatttaaaa catggagtcc 118560
taaagttatt atattatgcc ctcccatttc ccttcttctc ctctctctct tctttctatt 118620

-continued

tctccatata ccaagtacaa tggtaaaaga ataaaagtga cttacatgtg tatatggaca 118680
gaagtctcat tactaacatg tgcattgatg agagtgggta atggttatgt cagttgtgct 118740
atttgttttg tgttaacttg tagtaattga gccttttggc tgttcattgct ttcacatttt 118800
tctgcaaaat caggcactta tttgagaaaa ttttcttgac ctgctaacat atttagtttg 118860
gctgaaagta atttttatta ttcccaacca cttttaattg atagctacta attataaata 118920
cgacctatt caagtctttt tttacaacac tttcacaagc aatttgcctc aaatctatca 118980
ggtttgataa acagtcttta taattcagtt accagggat tctgaaagt acagacacac 119040
aaagcaaaaa gcaaacaaaa caacctccac aaaaacaata acaacaaaaat ccctttgggt 119100
cttgtcaaca ctgctcagag catgtctcct ttgaaacaag accatttcag caattaaaaa 119160
aaaaatcacg cagattttcc cttcttgctt ctttccccct ttttttaaat aaaattgcca 119220
acttaaaaaat attataacat tttaccattg tcagcagtta tacagaattt tgctcacaaa 119280
tccaagatgg ttccgccttt ccttattttt ttttaaggac aactgatta ggtattagaa 119340
tcaatacccc cctgcctttg atagaattct ttttctttt acaagtaac tctgaattaa 119400
aggttaggca atcgatttct tcagctattt ttaccattgt cttagtcaca gagatctgat 119460
ctaagacaga ccataatttt tccccttaat tcagaatatt cgatagaata tgatagtaca 119520
aatagcataa tctagactga agcaaatgc ttacaaaagg catgaaagct tcataatcat 119580
tctgtcttca aatgaacaca gagtattgat ttgaaatatt gttccagctc ctgtcctctt 119640
gaggagtata ctgacaatc actgaatatt acaaaagcat ctcagttgtg tggattgcag 119700
aaaaatgctg actatgataa catctgtgat ttgttattac aaattgaaag ataataaaat 119760
aattatctct agttgcatag ttactgattt tagcattgca accatattgc acttaatacc 119820
ttgataatgt ttataacatc ccacagatgt gagtgaaaaa tacaataacc agtattttag 119880
acaagaaagc ataggatttc agcaagtgtt ttgtaataaa atttgcacag tattataaag 119940
cfaatcataa atattaaagt ttagtttaca tactgtttac atttctctat ttgttcatta 120000
acaaaaatta atctgaagac tacaacata aaaagggttac tgtctataaa acacacaact 120060
ttgcattca gatgtcctaa agtctttcat atactttata tgtagtttta ccaatgtag 120120
agcaagagaa aaagacaacc ttctctttaa atccagaaaa gagcgagaaa aatagttttg 120180
tgggcatact caaagaaaca gaggtggtaa ttacggcagc ccactctaag agtgacattt 120240
ctctgtttca gccactaaa aaaatgtgca aagaaacaag taggccattg tggctgtatt 120300
ttcttttttc tttttttttt ttgagacgga gtctcacact gttgctcagc ctggagtgcg 120360
gtggcatatt cttgcaatct ccacctccc ggttcaagtg attctcctgc ctcaacctcc 120420
tgagttagctg agattacagc tgcacaccac cacgccagc taatttttgt atcttttagta 120480
gagacaggtt ttcaccatgt tggctaggct ggtctcgaac tcctgacctc aagtgatcca 120540
cccgccttgg catccccaaa tgctgagatt acaggtgtga gccaccgagc ccagcctgtg 120600
gctgtatttt ctaaagcttg ttgcaccaat atatatctca ttcacatgc tcttgttaca 120660
gtgtgtcatc aacatgcctc cattgagctg tgggtctctgt tttctcccct tgaacctggg 120720
tggagctttc taacagcctc aagcaataga acgtgatgaa tatgacatta caggacttat 120780
gagactagga cataaagga catataatct ctatctggtt ctttctttct cactcagggc 120840
acttgcccc tggagctcag ccaccatgtt atgaggaagt ctgggacaca tgaagaggcc 120900

-continued

acatgcactt attctgtcta aaagccctag ctaattgcaa ggctgacagc cagcagcaac 120960
tatcattttt gtgaataaat aagtcttcag atgattccaa atcccagcct ttgagtcacc 121020
ccagctgatg tcaagtagag caaagatgag ctatcctcac cgaactccat ctaaattgct 121080
gattcctgag caaaataaat gtcattgttc taagtctcta aatcttgag tcatttatta 121140
tgtagccata gataacagga atggccttca acaagatagt tctttaacac tgtaaaaatt 121200
atcttgctca tgtaataggg gttccactgg atctccagaa accaatgctg aaccatatca 121260
attcctagca tacttaactc atttcccat ttgttgagta tagagttccc tttgtgatt 121320
gccttgaggaa tggaaacaag agagatcact tcagaacaga aagagaccag ggaattactt 121380
ctctatgggt tttgtgagtc cccagagcta gaaaaaaat taattggctt aactaaaat 121440
gttaactta gttctgttct tggaggaaca cctttacggc tagcgacatg atagacattg 121500
ccactcttgt gtacttatta tgaaacaaga atgatctatc tccttaattg tagcaaaaat 121560
aacttttgag atgcgggtaa gtaagcagta gggacgggca gagcagtaac taagttgtgt 121620
tcataggtat tgtcaagtt cttaaagtag ctggaagcac atggttatgg caagtgtgga 121680
gttatttccc catttttaac tcagcaggat gactcctggt gttgacctaa gaatcacagg 121740
atgagaggaa gaaacctgcc aaggcatgtc ggccaactcc tcagagaaat actacagtta 121800
acttagaggt cagtgtcacc atttcagcgt ggctttttaa aatgttgca cctttcaaag 121860
attcctgctca acaaacact taaactttat taacataagt tgtttacagc atctatatgg 121920
aacaactagc tgatgtggct gaaaagtta tatttgggggt taattccagg tgttttgagg 121980
actccatgtc aataacatag ttcttccctg gatgagatcc caacaaattc atatggatta 122040
ttgggccaac ataacttita tagaacagca agaggcttgt atgcagccat gtaataact 122100
atggtaacat ctgaatcaat tgcagttcag tagacaaaac ttcgagtc aa agcttttatg 122160
caaatattct ttaaaactgc catgatgag tggccaagac aatataaaat ccatttcccta 122220
gaagaagaaa aaggtagaga gtagagtatt taatagtata gactctggag ccagactgtc 122280
tgggttcaaa ttctgtctct gacactggtt tcatgatcct aggtggaggt atgtagtctt 122340
tcttagcctc agttcactca tttaaaaat aaaaatgatg gagttgtagg cagaaagaaa 122400
ttgattcatg cacagagcat acgacagtgc ttgttttgcg gtaagtgtc agtaattggt 122460
acctattatt attattttgt gagtgaacta tagagtatag tgaataactc aaaaatgtcc 122520
cctctctcc cattatggag tatttttagg ttctgggag ttaattattc agcttctttg 122580
gccagaaaa gatagcctcc aagcacattc tataacatcg acactttctc tagaagtggc 122640
tcagtcgctc taggaatctg ctgtaataac cctaacacc aacattagat actattcttt 122700
taatttcacc cttggtaagt agagagcaca tcatgaccag tgagtgtgtg gtttcaatgc 122760
tgcatagtga gataatgaga aagccttgca aaaagagaaa acaaggcttg tattgtagtt 122820
gtgaaaattg agaaaggag tgggagactc ctggcttcaa aaggaattca ggctatagat 122880
gatattcttg aagcacttaa tgtttccgag ctagccatca ttatttttct cactatgtat 122940
agctcccttc ctataagctt aagaataaat gatagaaat tgtggtttat tctccctgaa 123000
tgccatcaa agatcgactg gataaagaaa atgtggtaca tataccat ggcatatctat 123060
gcagccataa aaaagaatga gatcatgtcc tttgcaggaa catgggtgga gctggaggcc 123120
attatactga gaaactaat gcaggaagag aaaacaaat accagttct cacttataag 123180

-continued

tgagagctaa gtggtgagaa cacatggaca catagagggg aacgacacac actggggcct 123240
attgcagggg ggaggctgga aggagggaaa cgatcaggaa aaataactaa tgagtactag 123300
atttaatacg gggagaaaat aatttgtgaa caaaccccca cgacacagtt ttacctatat 123360
aacaacttg cacatgtaca cttgaactaa aaagttaaaa caaaacaaac aaaaaatttc 123420
atattttaat tgtatgctgt taatctgtgg aaatacaatt gattttttaa aataaaaacc 123480
ctacactcag aaaaaatgt ggtttattct caacataaga aagtcatttt gaccgagaag 123540
caaatgacca tgtaacatcc cggaaagtaa tactgtgctg agatcagctc cattgaatgg 123600
ttttccagt aaaatgtgaa gtaacctgac aatggcagac ttttgaggca gaaaaacagt 123660
aggaagtga gagcccaag atagttcagt tttccttaac tttagatcat atccggtgtt 123720
gcacatatgt atctcagttg gctaaaagct gatagcagag atagagaaac ccaagaaggt 123780
gccagtgaa gttctcattc caaaaactcc ctcatcattt ggatgccac tatattctac 123840
ttttatgga ctgtcaactt tttttttcc ttccttgaaa agacattagt gttttaccat 123900
tatcagcagt tatgagaaac tttgtgaaga cacgattttt cctctctcca tagttctgag 123960
ttttctgtg aattcagcgt ttttatggat gccataatga acagattggt gaagtatgag 124020
caccttttgt cagtcacttc tcagttttgg tgtcagatcc agtctctgcc cttctcgata 124080
atgggagagc aacacttccc tgtttcccct gcactctgac ttttgatca ctctgaccaa 124140
tagaaggcac aagtgagaca ctggagggca gaaggagggg agaagacaaa ctaattctct 124200
ctctctgctt tggtaaatat ctctaacaga tgtgtctcct ctgtggctcc agcccccatt 124260
gggcagatgg aaatggttcc actctgcatt gggcttgggc tttagaactt ctgtaacatc 124320
tacttaacac attttctata ttaagggtact tctatggaaa acatagactt atttctgttt 124380
tctgactgga ctctgactga tacacacttg tagaatcttt atgaatatct ctgaaagaaa 124440
acattagaat gctggaaaat tctaagattg taggggtaat agaaattcta aataaataaa 124500
ttctggatca ttgatgatgc tttgggtatc gacaatcatg ccattaatga tcagatatta 124560
ttaggcatag acaagctaata ctttgtgaag acctatcct aggtaaggat agtgagaagt 124620
ttattaactc taaatactac atatggggat ctcttaaggg ctttctcag agtcttttta 124680
tagtctctat ggagttttaa caaaagcatc agtcacagta tgattagctt agtctctagc 124740
atgcgtaatt tattcaatag taccttagtt gactagaagc taaaatatct cttatatatt 124800
ttttccttaa attaaaaagc ttatttctga cttggagcaa attgctgcaa tagctggatg 124860
tagacagaat gatataggca tttctttctg cagtttttagg tttacagaat aattaaccaa 124920
aaagtatata taacttcatat ataccocctc actctacccc attcccaca cctcagtttc 124980
ccctagtatt aacatcttgc attagtgtgg tacatttatt agaatagatg agccaatatt 125040
gatacattat tattaagtaa agtctatagt ttacattggg gttcactcta tgtgtgcaca 125100
ttctatgagt tttgacaaat atatgatgat acgtattcaa ttacaatatt atagaaaata 125160
gttccacagc ccccaaaaat cccatgctct acctcctcat cccgctgtgt tgaaccctg 125220
acaataatag ctctactgtc tgcaaatttt ttccttttcc agaatgtcat atagttagaa 125280
tcacacagta tgtggccttt ttagattggc ttcttccact tagaaatag catttaagtt 125340
tcctccatgt attttcaagt tttgataatg tacttctttt tatcattgaa taccattcag 125400
tgatttggaa taccattgta tggatgatc acatttgttt actcattaac ctgttgcaag 125460

-continued

acatctgtt tgctccagg tttggcaat tatgaataaa gctgttataa gcatttgtat 125520
gcaggttctt gtgtagatgt aagtcttttag ctcatittgaa tatataccaa ggaccacatt 125580
gtaagagaac gtttagtttt gtaagaaatt gctaaagtgc cttcaaaagt gactgtatca 125640
tgttgcatga ccactagcaa tgaacgagat tcgtgttgtt ccacatctctg gccagcttcc 125700
gggtgtgtca atgttttga ttttagccat tctaataagg gtacagtgg accacattgt 125760
tgttttaatc tgtgattccc taatgtgaca tataatgttg agtatcttgt catatacttg 125820
tttgccattt gtatagcttc tcttttgagg tgtctgcoat atcttttacc cactttttca 125880
agggactgtt tgctttcta ctgttgagtt ttaagagttt tccatatatt ttgtatatca 125940
aaccattatc agatatgtgt ttgcaaata tttctccca gtagtctgtg gctgtctttt 126000
tcattctctt aacaggaaga ctttttagtg taatttttcc caatgtctct ggaattttat 126060
taccatcttg ttgaagctag ggtatttcta tggttttgaa tacctgggaa attaatcct 126120
aatgttttga gatagtaact cccctccaag atatttttat gatcagtgga aatgtatttt 126180
atagcccaca acataacttc tagtatcagt tgtgtgtgat cattataatc ttgaagaatt 126240
acgattagtg gtcatagcag tggtagaaa gggcttgag aaaatatcac tgaatggtct 126300
caatcatact tttagaccat ttctacagtt cacttcccat ttactctgca agcagtaaag 126360
cagtctggct cccagccctc tactcccacc ctttaacctc tctattgaa gtgtcttgag 126420
aaaccaatca gtgacctgtg atgttaactc caatcccttt ttcttgaaa tctctttact 126480
ataatcttgg cttttgtaac actactttat cctagagaca ggtaggtagg aattttacta 126540
tcttttttcc tgaactttgt aatggcctcc taagtatttt cattgcctct tctttttcca 126600
atatatcctt cacattgtca cagcacagaa ggcaacatgg tacaatgaga agatcacagc 126660
ccttgaagcc aataagacct aggtttacat ctacagctctg ccacataagt gactgaaatgc 126720
tgacctcaa gttcaacatg acatccctta actgtggtaa tttatgactg gagttgtttt 126780
tgaatcaaa cataataata cctataaagt ggcctgaaca tggataaag tcatttcaat 126840
ttctgaagaa cgaacccagt tatgaacaca tctctcttcc attagttata caataaattc 126900
cagactcctc agcctgggat gcctggcctc gcacaggtga ctgcttcttt tcagtcttag 126960
cttctatgcc atcatacatg ctctgaaaac acagatgggc ttgtttcctt ccacatgtgg 127020
cctcccagct atgcctatgt gtgtgtatat agtgctctca ctgtaataata ttctgtctctc 127080
ctgtcggaat ggcagacttc caatttattc ttcagggccc taatctaata ttacctcca 127140
gtcctttgcc ctcatctaa tagattttga acaacttatg ctattgaagt gatttcctat 127200
ccttgatgat ttggcccttc acatcctgga atttgccaac atgtgaaatct gtggctccat 127260
ttctcaagaa gtagataaac tataggacca gctcactttc cttaggggct gtttacctg 127320
ctttcaaaaag catgatattt cattttaaga acataaactt cagttaaaac tcattgtgat 127380
aattcttggc actgggtgaag atgtggtgaag ataagcattc catagcactt atgatgaaa 127440
tgcaaatctt agtatctttg gggaatccag tctgtcaata tatactctgag tagatactta 127500
aaatacttat accttttgat ccagtgtttt tttttttaa aataaactta aagaaaatat 127560
atztatatat aaacatgttt atcatgtata gtaaaaaaat tgaatggaat ttaataatca 127620
aaacagtttt ttagttacat aaattatctc cataccatag aataataaac agtgattaaa 127680
atgtacatgt ataaagagtt ttaaatgaga gaaaaacctt atgatataat attaaatca 127740

-continued

aaattcgaat acaaagacag actttcaata tgaccattac tgtattcaat atgtacctag 127800
aaaagagact gaaaagaaat atactgaaat gtttgtgatt ctattagtca gggttgacta 127860
cctgcttaat aaacaacctc tgagtctctg tggcttaaca ctaaaggta atttcttggt 127920
catgtcacag tacaatgcaa ggttggaagg tggctcttca tgtaattatc tgaggaccta 127980
gactccttcc ttcttttaaa gactgactta ctttatgtat tcaactggga agtgaaggaa 128040
ggaggtttca gaggcattgc ctggaattga gacacatcac ttccatccac ataccactgg 128100
gcagcagagc tcagtcacat gaacccaact aactgcacaa tattgagaac tatattccca 128160
acagaaagaa aaggaacttg agtattgaca gcactagcat tgtcttctc atctctaaga 128220
ggaaatctaa taattactgt ttcatttttc tgcattttcc aattattccc cagtaactatg 128280
tattacaaga gaaaaatgta tatctagaag ccctcaaagt tttaaaagag ggaactcctt 128340
taatttccac cgtgtaaatt tacactgcct aaagtatttc cttggacatt caaaaggta 128400
aatttggttc aagcaaaagt tgctggtgta agatttcaca aactgcacaa aagaaattca 128460
atgcagtatg aacagccttt tgattcagct aatgatatta aaatatgtaa atacatggag 128520
aagcacattg attatcccca cattcatcaa atgtttttaa taaaataggg tgttgtaaat 128580
ctaactttaa attatgtttt tagtagaaag agatgagagg tggagatgtc tgagatccac 128640
attggtggac gttaaactct tccctgtagg caccctttct atagtttata gctcccaaat 128700
tgtctttatt gttgatcaca tcagcactct tcttggttgt ggatagagaa taagtagaga 128760
gagaggatgt tggagaccta catggccact ttcttcagcc ctttccttca atgtcctgag 128820
gccagcatat gttctggata gtattaaaaa tcttggtgcc cagactgtca ggggtctggc 128880
aatatttatt tatttatttc ctctcttctc tcttctctc cctctctctc tccctcttcc 128940
ctctctctc cacttctctc ctctcttcca cccaatcacc tatgcaagag acattgatta 129000
aaaaactaca ctgaccacag aactacacaa tataactcta tttttgttc tctattttta 129060
tttgttacgt gatgaatgaa agtctttctc gatatgtata ttggagagcc agtatggtga 129120
caaaaattct tcctatttgc tctagaacag ccacactggg gaaagttgca ctaaaaagtt 129180
tgccggttaa aagaatgaaa agatgagcca cagactggga gaaaaatatt gcaaaacaca 129240
tatctgataa tggtttgata tacaaaatat ataaagaact cttaaaactc aacagtaaga 129300
aagcaaacag tcccgtgaaa aagtgggtaa gagatatgac cagacacccc acaagagaag 129360
ctatacagat ggcaaacag tatatgacaa gatactcaac attatatgtc acattagga 129420
atcacagatt aaaacaacaa tgtggtgcca ctgtacacct attagaatgg ctaaaatcca 129480
aaacattgac aacaccaaag gctagccagg atgtggaaca acaggaatct cattcattgc 129540
tgggtgggat gcaaaacggt acagtctctt ttgaagacac tttagcaatt tcttataaaa 129600
ctaaacattg ttgatcacat cagcactctt ctgatcctgc tcagactgct gagtggaggt 129660
aacattcttt tgcttattat tttatagat ttgaggatag acctactctg acagatgggt 129720
gctttccctt ttatgcacac atctgctcaaac aacctcttat tctcggggct gaaaaaactg 129780
tagcataata gttttttaa aaatcaagct aaaactagca ggtctaacac aaggtaatta 129840
cttctcaaaa caacacacct tgcaactagg aagtctcaga aagactctaa acagccatag 129900
ccagccatt acctcaatg tatttgggga taaaagagcc tcagtaaata cagggcgtat 129960
gagccattga ctttgccatc ccatgctgt caggctggtg gagtattgac agaggagctg 130020

-continued

cagccagtga tgttattgtg ctacgagaag aactgggcat gtcctcatgc tgaccaacc 130080
atgatctttc tacacttatt agaactctgct ttcacagtag ttcacaggta tccttcagtc 130140
ccagattggt aaaggaaaat gtgtctagaa caaaagtcct aaagaagaaa ttgaattaac 130200
aactcatttc attctataat actccagtta aaatgaagaa actctcattt tttttgccct 130260
tctcagttta tcacattgag aaatacagat tgctagtcta taccaacctt gggttattag 130320
catttcacaa tcattaacat ttcattacc ttttttcttc tcagtacaaa atgtcactgg 130380
aaatatatgt gaaatattta ggaactctgt ataaaatata taaaagaaca tgaaaaatat 130440
gaacatttaa ggataagagg agcaaagcac ttctcaaaat gtcttgtag tttcttattt 130500
gaattctaaa ctcttggtta gatgcttaag tcaatcctgc ctaactgaat gtgattttcc 130560
acctctccat cctgcccag tattgcata ctgctatcta cattcacata tttatgtagt 130620
tggctgtgtc tgcagaacct ggagattctg gacataggat tttattatat gtgtttggga 130680
ttattgaatc acccactcca cattatatta aagtcaagaa cattggattt tctaaattgg 130740
ttttgaaata gataaaattt tctaggtatt tcaatgacct tctcctttgt ctaggagact 130800
cagggatattg ctactagtt atgctgatgg ctttattgag tgcacacaag ttaagttatg 130860
cactatatga aaaagttact ttagattcag tccctgacca gcagtctcat agcctttatt 130920
taatactggc acttaaaagc aaacaaaatt aactgcactt ttactgaaat ataattgcta 130980
ttgtataatg gaaaggaaat tatgtttttt tccttttcta tcagcattgc ttgggtggtga 131040
taattttgac catgcagtga gagaaataaa agatgcctat tttatgaaag catttttcag 131100
taatacaagt atgtccattt taagtgagaa agtgattttt gggactagaa ttaaccacat 131160
gatatttgag ttgagatca atgccacta tttgaggcta ctgctttggt gcaggaagtc 131220
agggaccccg aatggaggga cgggtgaag ccatggcaga agaataaaa ttgtgaagat 131280
ttcatggaca tttattagat ccccaatta atactttcat aatttcttac acctgtcttt 131340
actgtaatct ctgaacataa attgtgaaga tttcacggac acttatcact tccccagtca 131400
atacccttgt gatttcttat gcctgtcttt actttaatct ctttaattccg tcattttcat 131460
aagctgagga ggatgtatgt tgcctcagga cctgtgatg attgcattaa ctgcacacat 131520
tgtttgtaga gcatgtgtgt ttgaacaata tgaaatctgg gcacctgaa aaaagaacag 131580
gataacagca acgttcaggg aacaagagag ataacctaa actctgaccg ccggtgagcc 131640
aggcagaaca gagccatatt tctcttcttt caaaagcaa tgggagaaat atcactgaat 131700
tctttttctc agcaaggaa atccctgaga aagagaatgc atccctgagg gtgggtctct 131760
aaaatggccc ccatgggtgc agccgtcttt tatggctgag ctgtaggat gaaataagcc 131820
ccagtttccc atagtctcc caggcttatt aggacgagga aattcccgcc taataaattt 131880
tggtcagaac ggttgtctgc tctcaaaccc tgtctcctga taagatgta tcaatgacaa 131940
tgtgtgcccc aaacttcatt agcaatttta atttctgcc ggtcctgtgg tcctgtgatc 132000
tcgccctgcc tccatttgcc ttgtgatatt ctattacett gtgaagcacg tgatctctgt 132060
gaccacacc ctattcgtac actccctacc cttttgaaat cactaataaa aacttgctgg 132120
ttttgatct tgtggggcat cacggaacct accgacatgt cataatctccc ctggacgccc 132180
agcttataaa tttctctctt ttgtactctg tccctttatt tctcaaacg gctgatgctt 132240
agggaaata gcaaagaac tacgtgaaat atcgggggtg aattttgcc gatatctggc 132300

-continued

tgaatttccc caatactgct tctcagtgag attcaagtta tgtgggagac caatatccaa 132360
tcatttactc caaaaaagtc tctgatctgc tcatagcatc attttctaata aacaatgtga 132420
gagatagatt tataaactgt tcaggaagac taacacgagt ccctggtgga gctgagagac 132480
tgggatagga ataactggca cttttttttt tttttctttt agcaatgttt taggttaaag 132540
ggcaatatta atcctccttt agaatcaggt gggtcatttg aggtcaggag ttcaagacca 132600
gcctggccaa catgatgaaa ccctgtctct actaaaaata caaaaattag ccagggtgtg 132660
tggctcacac ctgtagtctc agctatgtag gaggctgagg catgagaatc gctggagcct 132720
gggagtaaga ggctgcagtg agctgaaatc atgccattgc actccagtct gggtgacaaa 132780
gtgagattct gtccaaaaaa aaattatgca ttttcttatt ttctctgtat tctcttgaga 132840
cgacagctcc agtcattaga aaatgtaaac tcagctttcc tttttacctt tggttaaaaa 132900
aaataaaaaa gtattcacag ttctcttcac aactaggctg gcttactttg gcctgtgttt 132960
ttatttgctc ttttctcccc tatgttgctt tctttgtgcc ctgtttttcc ctttctgctt 133020
cttctgagtc acatgttgac gtaactcagt gtgaagcctg gggtttgaga agatgtggac 133080
gccaaagtcca ggtcacacaga gatTTTTgtc tgataacacc acatacactc cgcctgcaga 133140
cttctacat tggaaacatc gcattttcaa acgccaattt gccaaagcagg ttggctgtga 133200
atgtgcctgc cttctgctgc ttgtcagatg gaggctgact cgctccattc ctctaactct 133260
tgggcttcat tcgtgattta gccacactga taaatatagc ttctttttga agtagagagt 133320
ctagattttt tttctctcag gctttaaaaa aatggattgc tttattccaa actttatttt 133380
tgatgaaata caactcacia ctcatcttgg aaaaaaagc tcttcaaatt ttgaactggt 133440
gaaagtttga gtctgaggtg atatgacatg tttttgatca ttcaataact ttatttctct 133500
tcttctcgat tgcttccaaa gttggaagtt acatgcacct aataccactg ttcctatttg 133560
atgttctcag atgtgaaacg gggatgctc tcaatcctta attaaatctc ctcttgccca 133620
agctcacttc tacattttta ctaagttctt ttctattcta tttatagttt atgtatgtct 133680
gaattacata gaatgtatga gagggaaaga aacataatag agaaaataaa gagatacaaa 133740
ggaagaata aagtgaatga aaagacaatt caaagggggg gagaatctga gcaatgggta 133800
ggaattttta agaagaatg gaatagggtt tctacctaag aatatgaaat atttacttag 133860
gtgtagttag acaacttggc ttctttcatg aaacaaaaca taaaactggg atgttttcat 133920
aaagaaatgt tccatatttt aagactgcca aaagatgaca cccacacatg ccgggtggaat 133980
ttggtaacat ttaactgtgt gaaaacaaga ggtctcaaag ataatgattc agagtcggca 134040
gagtgtttat ttatctttga tttagaaaag acattgtctg ttatttggct gtagatatgt 134100
cttaaggact cttttagggt tggctctgct gaaaccaaga gcttaaggca aaaacagaca 134160
tgctttcccc tttttgatgt tcatgctgca ccaaaataca cctaacatct atggctctatg 134220
ctgggagacc ctccatgac acttaacttt aaatagacag atggataaat agatgtgact 134280
tgctcttact cagcactggt caaattttta caagaatgta ctttccagct tggagctcca 134340
catcttaaga gaactctaac atttgggaaa gattcagaag agactgtgaa tgttgatgag 134400
agggttggaa aatggaactt ataagaaaga gttaaagaaa ttggaattgc ttttcttga 134460
gaagagaagt aacctggtaa cagtcttccc atttttgag ttacataata tacatgatga 134520
ttgccaagtg ttttcttct ctatcaaaaga acatcagagg tcaagtgtc acactgcata 134580

-continued

tatcacatat aaggaaaact tgttgcttgt ggtggtggta gtgttaaaaa gataaactgg 134640
taaccattgg ctttgtaaaa acttttctgg aattacttaa atattggata gttttttggt 134700
ctcatatfff ctgaggattg gaagataata cttatgtcat agttcttaat gcatcatgat 134760
ttataagtct ataagaacaa aaatgtatff taaagaaata agtgtagata atggcccaag 134820
attcatagaa tctataatff actttccgff aaataagtga aaccaagaca ggagatgtgg 134880
tttccatagt gacttttaga gtaaaccatff tagtgtgata ctctgatttc ttaaaaaaac 134940
actgctgcta attgacataa ggaaaggtaa aatcaagtgt gtgtgtgtgt gtgtgtgtgt 135000
gtgtaaaata tacaatagat actatgtata ataatacaat atacaataga tatatgtatg 135060
ctaagtata tggccaccta agaacctcag ttagaattff atffttttaa aaaagagaaa 135120
aaccaccag ctctttctff tctttctff tctctttctt gctttctffc tctttctgfc 135180
tctcttctct cctcttctct ccttttttct ctgtcttctt ttttcttct ctcccttttc 135240
cttcttctct tcttctctff ctttcttctt ttctttcttc caactaagff tattcttcta 135300
gtgatcccca agtagatag ctgctatgca aaactccttg tgtgaaccaa tatactffta 135360
atattcctca tgttcccag tgttaaaata agctgatcaa tcaataata aaacaattg 135420
aatgtctatc tactctatag tcaactggtga tgttggttat ttggttacct caatgaagaa 135480
ataaaactca ttgaattctt tggcaaatff ctactttttg tagtgctagg atcatagaaa 135540
catagtctca tgattcatct gcattfftaag ctctaataga acacatcaca ttttaggggg 135600
aaaatcaatt ttcacctaaf gttagcaaaa atgagattaa atattgcaat tatcaaaaat 135660
agcattatgt gttttccatg atgcagggct atatttatgt tttggttttt aatcatacaa 135720
atgcattgta ttctgttttc agttccactc ttatcactag aagcatttct ttgtgctgag 135780
tagaactaca aatatacaca taattctcta tgagtaggca caaacagctt tgctttttaa 135840
atfttgagca caaatcatct gctatgaaaa catggtffta taatttctgg tggaaatata 135900
ttaataccac aggctttcag attaagtcat aaaaagaaag gtttagatff ttgtctttca 135960
cattttatff tagaaaaag acataccaca caggtcaca cctgtaatcc caacactffg 136020
ggaggccaag gcagggtgat cacaaggfca ggaaatcag actatcctgg ccaacatggt 136080
gaaaccccgf ctgtactaaa aatacaataa tttagctggg gtggtggcct gcaactgtgg 136140
tcctagctac ttgggaggct gaagcaggag aatggcgtga acctgggagg cggagcttgc 136200
agtgagccga gattgcacca ctgcaactca gcctcgtgac agagcgagac tccatctcac 136260
aaaaaaaa aacaaacaaa aagacataca gtgtttatff aatatagtgg ttttgcttag 136320
atgctcactt ttgaaagfca tffaattggg gggaaatgaa aatgtfftaa aactffattg 136380
tggacatff caaacattca gaaaagcaga taggatggtg taatgaaact ttatcaccca 136440
tcattcaaca tttatffaca ctctgcaatc ccaactffact taccctacc tctctccctt 136500
aaaacaaaca tacaacaaa caacaataa ctttactgga gcttaaatgc acttaactff 136560
cctttgggaa ttagtatata tttgtcagf gtctaaaatg taagcattff tattagtcaa 136620
tttattcata atffttact aacatfftg tacagaatac tgagtatatt aatffctctg 136680
gactgctgta acaaatgacc acaaacctag ttgcttacac caccaccaac ttattatctt 136740
atagttffg aggtcagaag tctgaaatgt gtctcagfgg gctaaaatca aggtgtcaac 136800
agcaatgtgt taacttctg aggccttagg gagaatctct tttcttgcct tttctggctt 136860

-continued

gtaaaggctg cctgaatttc atggtctcct ccatcttcaa aaccagcaat tgtcagttaa 136920
gtccttctca catagcatta cttaactctt tcctttctcg cctttctttt tctcttataa 136980
agacccttct gattacattg ggcccaactg gaaaaactca tctttaggtc agctaagag 137040
caacattaat tccatctgta actaccattt acttttgcca agtaaccta catagtcata 137100
ggttctgagc attagagatac cgacatcttt tggggccatt attttgccca ccacgtaggg 137160
catattttat attgatttgc aaatatacat ttatgtcttt catgtgaaat gtataatata 137220
gatttaacag tcatagagaa atagcaaaga agtattttac catattcttt gggttgagtc 137280
cacactaat tatgtggcct gattattctt taactcactg ttcaatctga acacacacag 137340
tagtaattat catctcatal ttgggagagt aaagtaaaaa aacagcaata ataacaaaaa 137400
taggccaggt gcagtggtc acacctgtaa tcccagcact ttgggaggca tgtggatcac 137460
ctgaggtcag gagtttgaaa ccagacaggc caacatggtg aaccgctgtc tctactaaaa 137520
atacaaaaaat tagccagggt tgggtggtgca cacctgtaat cccagctact cgggaggctg 137580
aggcaggaga attgcttgaa cccaggaggc ggaggttgc atgagctgag atggcaccac 137640
tgcactccag cctaggggac agagtggagc tgtctcaaaa aaacaaagca aaacaaaaca 137700
aaacaaaaat gaccacggaa cagatcaatt acagtagaat agtttacgca ggtggaaggg 137760
ccttctgacc aggaatctta acatcacaat ctatacattt atttactcta gagtatgtct 137820
gtagctctgt tacctaggat gcctcagcca actgaaatga taggtacttt ctctccggct 137880
tcacaacttt gtcttcaca gactgcccta ggccccatg ccatgccag acacttattt 137940
tcccttcatt gtgttccta atattatggt cagtggggag gcctcttaga tttgccttac 138000
agtgaagagg gatggtgtaa tttcactggt gccactagcc ataaagttgc agacagagag 138060
gctacagatg tggagcagat gtgcataatg atcatttaa aaattctttg tatagctctc 138120
ataaaaaagt gactgtgata aatgtttgtg tgtttggtgc tcacattcta ttctattatt 138180
taaacagaat cattaacttg tttgtacttt acaatctatt tttatttga ctaaaaaacta 138240
tttattacta cattatttct cttgagcttg ggaggatgac agctttgccc tccctcccaa 138300
tttaggatga attcatttac ctttctcttt ttctgttcta gacggagttg ttccaaagga 138360
tggggaaca ccttcttga aaggagcata tctcatggag gcagcatcag cattagaacc 138420
aggtgaaatc tgagcccaga cactcagatc agttaatttc tctgagcctt catttcttat 138480
ttgcaaatg ggactaaca tataatccat aaagtttctg tgagaaaatg cagggataat 138540
tatagaaagc aggtagtgtc gtatctgtgt gattaaaggt ctgttccctt cccttctcta 138600
tggaggagc catgccacta gaaaaacacc tttagacca caaacttttg cttttaaata 138660
aaattagttt cctatttctg ggtatcatga aaattcttag atagacttcc acaaaatttg 138720
gaaagtattg ttttaggaaga tctgatttag aatttgggtt atgtggcata tgataagaat 138780
cactttagga gatcctggga agcctcaaag aagtaggcag tcatcataca ggagctgaaa 138840
gtagggttga acaaggggta tgaataaacg tcatttagga gaggcctgtc atattcagta 138900
atatgccat agatggaaca aatggaaca ttggtttcca ataggtcctt aaattgaaag 138960
ccataagggt aggtgctttg caaagcaggt atgatttcta tccaaattac ctctcacatg 139020
ggcaccttgt ggatttctct ggacagaagt gcagcaggaa cctacttttt aatgaataat 139080
ggttattgag tcattcaaac atcccagaat agccagtggg acctggctgt tctcccttca 139140

-continued

agttcttcgc cacctgtaag tcacagagcc acagctgtgg ttcttctgca ggggccacag 139200
agcccttctct ctgcacttgg aaaaaaatgt catgtgctga ctccaagagt gagtcttctg 139260
ggggtacag agacacagaa acagaaactt tccaagaaaa acacactgac taaaccaaata 139320
gacagagcag aaaacatttt ctgataaggc aaagggtaat aacttgggta gaattacaca 139380
ggtaattaaa tagttgtttc caagggctag gaaaagacag gcacgcgtctg tgttgacag 139440
aaggggagtt ttggatttat taacatggat atgaggggtt agggttattt gtgtttgtgg 139500
taaagagctc agagaagaag tctttattta ctgcctaaag atttagtaag tcaatttgg 139560
ccctgccttc tatctcatcc acagacaata tcttcttaaa taggatacta gattgcatgc 139620
atgatgttaa aaaaactgag agtaggaaaa tcaccttatt gaaatagctc ttgaggtgtc 139680
cttaactcac aagaacttgg gaaaattttc agactttctc accaacagag gacataaata 139740
caaaggcaa gatgtaaagt tatgcaaacc tctgtggctg gaataaataat tttgtactta 139800
tgaaagtcc actagaagac cacttctgaa ttcagatctg tatcaattaa tggaaagaaa 139860
aaatgggttg gaggagccaa atggggaaatg ttacaagacg tattaatgta taatcaaaat 139920
ttagaaaatg aggtggttct gaattggaat tggctttaga taaaatacag aacaggaaaag 139980
aatgggagtc ctctgttcta gaagggtagg ctggctacaa tcgggtgaaat aagacaattt 140040
cataagaaaa cgttttgaaa gaatatgaga ggaaactgca agagcagaag aaagtaataa 140100
actcattgca caagaatttc agaagaacta gggtaaaaa tacagtcaca atctggcagc 140160
tttcaggagg aggtggggag tagataagga caaagagggtg ggaagggaga tagggccaag 140220
tccaaagtgc ctttcagact ctccctgagt cagcaaacct tgcttctttt tacttttact 140280
ttttctgtgg gctctgaatg gttcttttaa tctatattca ggcagagtaa caggcgtgat 140340
gaattataa tgtaaaaggt acactttggg aggctgaggc gggaggatca cgaagtcagg 140400
agatcgagac catcctggct aacacgggta aaccccgct ctactaaaga tacaaaaaat 140460
tagccgggagc tgggtggggg cgctgtagt cccagctact cgggaggctg aggcaggaga 140520
atggcgtgaa cccgggaggc ggagggtgca gtgagccgag atcacgccac tgactccag 140580
ctgggagcag agagtgagac tccgcttcaa aaaaaaaaa aaaaaaaaa agccagaaac 140640
gctcactagg caaaaaaaaa aacttctctt cgagcagcct ggtagggcct ctactggag 140700
aatattcagt tgtgttaag catcttagta gagcaccagt ggtatctgga agaaggtgcc 140760
cccaaaagcc tctatccctc atcagcaggt ttacacaatt catcgaacag atttaatagt 140820
tacagattac tctgttaact agattttgag aaaacagtgc agagtgttg cagagcaaag 140880
agagaaaatc aaattctgga gtgtggaatg tttaccctaa agaaatggca gtagcaggaa 140940
ctaggagcct gaaggggggt cttttgtgta aaaatgtttt tgatttctga tgctacagc 141000
tggattttta atgtagctca agtccgacag aatgcaggtg gaaaaagaa agcagcaatg 141060
aattgtttgc ctcttcttta aactctgttt ctgaaaataa aatttgttag tattttaaac 141120
ttaaaccctc tcatgattat tatatcctta gttcttatgt gtcactgata tactaaaata 141180
ggttgtgaat atgatagaaa gtgagatgaa agagagagag tgtggtaag ctgtgggtct 141240
gtgaaactcc aagggccagat gctgctcacc ttattatgga acaggatgtg tgattcaagc 141300
acagggtcaa aggaatgggc caggggtgact ctttttgtaa taatagttaa catgtgttac 141360
tcaactcatag actcaacctg ctgattaata tatgttatct cttgaatttg agaatgaagt 141420

-continued

tgtacaattc ccatttatag ttgtggaagc taaggctcag agatacacag taacttgctg 141480
tgtagccagt tgtggcaaaa aaaaaaaaaa aaaaaaaga atttgatattt atttctgtct 141540
gatcctaacc aggtgttttc caggatgca ccctcctgc tcacattttg attggttttg 141600
aattcttgtg tattattgac tgtctacttc actaacatca agtataaac aaaaaatgta 141660
ttgcagccta atgtttagct tctctactct tttatattgc ctatatttcc agttccttag 141720
atttgcaaaa gtatgaagt taaatataaa tcatacctgc aattgcatta gtatattggt 141780
atagtccctt tcattgagt ttgcaggaag tgagctgtct ttaagaatat ccagttgtag 141840
gaagttcctc tttttctttt ttgttactc tcaaaaagtc tttggtcttt cctttagta 141900
aattactaag cactctagtt tggctgatt ctagattgaa tattaaagaa ttctatgata 141960
aagcaaacctt taaaaatta caaacactga atatttttaa agtcaaagaa taaaatctgc 142020
agaggctgtc cattggatgt gattcagcca catgtgtttt tgggtcagat ttttctccct 142080
ttcagtgaaa atgtctttct attttcttga ctttgatcac tttggtgtca ggggtcttta 142140
tactcagtg tcatctgtca gccttttcat aagtcagttt cttgcatcca tctggggaat 142200
ggcaaaaagat ttgggttag gcaattgcat gcattactca aaagactgtg gagactttga 142260
gaagaaaga aggaaaatct tcattgcata cttttaactc tttttttttt ttaaattaac 142320
ttccccatca gagtgtttga aaaaacattt tgatatggca tgtgaaataa gaacaagagg 142380
aggtgagagg cttttgatct gtgtgctctc ctgaaatag catgaccagc tctaggataa 142440
agaggtgag aggcttgtgt ttgacacca catctctctt tccatcagtt cctcccactc 142500
agctgcgtgg tcttctgtt gctcagagcc ctaccttttg atctggctgc cagagctgca 142560
tttcttttgg gtaaactgtc cacactccag aatttgattt tctctgtttg ctctgccaac 142620
acataactgc cctgttctcc caaaatctaa ttaatggagt aatctgttaa actattcaat 142680
ctgttcaatg aattgtgtaa acctgctaag gagtgataga ggcttttggg ggctccttct 142740
tccagatacc actgggtgct tactaagatg ccctcacaca actgaatatt ctccagctag 142800
aagccccac ccaggctgct ggaagagaag ttttttttgc atagtggttg attttggcac 142860
cttttacgtc atacattcct catgctgtt actctgctg aatatagatt aaaacaacca 142920
ttcagagccc acagaaaaag tagacaagca actctgaaat tagggtatag ggctggaata 142980
tggaggatac aaaagagtct acattagccc caaatctggc catcacatgg gagaggcagc 143040
tgacacagat atctagaaaag caaagcagag gctcataact cctgcagatt tcgggagtg 143100
agtctgtag cctgacttcc aggaccagt tattcattta ctattattt tattctttca 143160
ttctttcaca tttctgtgac aactattatg ggtcagtc atgtggttaa atgogctaac 143220
agttctgtca ttttctgtt tggaaaagta gactcaattt acaaaaaaga acaaaactag 143280
ggtttctgaa ataaagattt gtaatttgct tttttcatgc aggtgcaaat tatcttaaac 143340
tgatataaaa aagaagaaac agaaatttta aaatatatgt tttttacaaa tgtgtaccat 143400
ctaattatca gctagaagtc ttattctgtg aagagctaac acaggttcta gtgtaataat 143460
aataatcagc ataacattta ttgagtattt actgtacatg ccaacaact gtgctgtgtg 143520
tttaacatga atttttgcatt ttaactctta ggaaaaattc tgtgagacag gtttgttatt 143580
accttcttt cccagataag caagcagatc aagaaatgtt gatcgcatag ctggaattta 143640
gagctgaaa ttgaaccag gcagctgtac tccagagtct gcagctttt gttgatgtta 143700

-continued

```

cctgtctcta ctccagccac aggtggttca cgacctatga tggttcaact taatgatttt 143760
ttgactttac aatggtacga aagcatttgg tagaaactgt acatcaagta ctcgtataac 143820
cattctgttt ttcactttca gtatagcatt caataaatta cataagatat tcaacacctt 143880
attatgaaat aggcctttgtg ttagatgatt ttgcccaact gtaggctaata aacatagttg 143940
ttctgag                                         143947

```

```

<210> SEQ ID NO 38
<211> LENGTH: 2217
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<300> PUBLICATION INFORMATION:
<308> DATABASE ACCESSION NUMBER: GenBank AC012462
<309> DATABASE ENTRY DATE: 2005-04-05

```

```

<400> SEQUENCE: 38

```

```

Met Leu Arg Gly Pro Gly Pro Gly Leu Leu Leu Ala Val Gln Cys
 1           5           10           15
Leu Gly Thr Ala Val Pro Ser Thr Gly Ala Ser Lys Ser Lys Arg Gln
 20           25           30
Ala Gln Gln Met Val Gln Pro Gln Ser Pro Val Ala Val Ser Gln Ser
 35           40           45
Lys Pro Gly Cys Tyr Asp Asn Gly Lys His Tyr Gln Ile Asn Gln Gln
 50           55           60
Trp Glu Arg Thr Tyr Leu Gly Asn Ala Leu Val Cys Thr Cys Tyr Gly
 65           70           75
Gly Ser Arg Gly Phe Asn Cys Glu Ser Lys Pro Glu Ala Glu Glu Thr
 85           90           95
Cys Phe Asp Lys Tyr Thr Gly Asn Thr Tyr Arg Val Gly Asp Thr Tyr
100          105          110
Glu Arg Pro Lys Asp Ser Met Ile Trp Asp Cys Thr Cys Ile Gly Ala
115          120          125
Gly Arg Gly Arg Ile Ser Cys Thr Ile Ala Asn Arg Cys His Glu Gly
130          135          140
Gly Gln Ser Tyr Lys Ile Gly Asp Thr Trp Arg Arg Pro His Glu Thr
145          150          155
Gly Gly Tyr Met Leu Glu Cys Val Cys Leu Gly Asn Gly Lys Gly Glu
165          170          175
Trp Thr Cys Lys Pro Ile Ala Glu Lys Cys Phe Asp His Ala Ala Gly
180          185          190
Thr Ser Tyr Val Val Gly Glu Thr Trp Glu Lys Pro Tyr Gln Gly Trp
195          200          205
Met Met Val Asp Cys Thr Cys Leu Gly Glu Gly Ser Gly Arg Ile Thr
210          215          220
Cys Thr Ser Arg Asn Arg Cys Asn Asp Gln Asp Thr Arg Thr Ser Tyr
225          230          235
Arg Ile Gly Asp Thr Trp Ser Lys Lys Asp Asn Arg Gly Asn Leu Leu
245          250          255
Gln Cys Ile Cys Thr Gly Asn Gly Arg Gly Glu Trp Lys Cys Glu Arg
260          265          270
His Thr Ser Val Gln Thr Thr Ser Ser Gly Ser Gly Pro Phe Thr Asp
275          280          285

```

-continued

Val Arg Ala Ala Val Tyr Gln Pro Gln Pro His Pro Gln Pro Pro Pro
 290 295 300
 Tyr Gly His Cys Val Thr Asp Ser Gly Val Val Tyr Ser Val Gly Met
 305 310 315 320
 Gln Trp Leu Lys Thr Gln Gly Asn Lys Gln Met Leu Cys Thr Cys Leu
 325 330 335
 Gly Asn Gly Val Ser Cys Gln Glu Thr Ala Val Thr Gln Thr Tyr Gly
 340 345 350
 Gly Asn Ser Asn Gly Glu Pro Cys Val Leu Pro Phe Thr Tyr Asn Gly
 355 360 365
 Arg Thr Phe Tyr Ser Cys Thr Thr Glu Gly Arg Gln Asp Gly His Leu
 370 375 380
 Trp Cys Ser Thr Thr Ser Asn Tyr Glu Gln Asp Gln Lys Tyr Ser Phe
 385 390 395 400
 Cys Thr Asp His Thr Val Leu Val Gln Thr Arg Gly Gly Asn Ser Asn
 405 410 415
 Gly Ala Leu Cys His Phe Pro Phe Leu Tyr Asn Asn His Asn Tyr Thr
 420 425 430
 Asp Cys Thr Ser Glu Gly Arg Arg Asp Asn Met Lys Trp Cys Gly Thr
 435 440 445
 Thr Gln Asn Tyr Asp Ala Asp Gln Lys Phe Gly Phe Cys Pro Met Ala
 450 455 460
 Ala His Glu Glu Ile Cys Thr Thr Asn Glu Gly Val Met Tyr Arg Ile
 465 470 475 480
 Gly Asp Gln Trp Asp Lys Gln His Asp Met Gly His Met Met Arg Cys
 485 490 495
 Thr Cys Val Gly Asn Gly Arg Gly Glu Trp Thr Cys Ile Ala Tyr Ser
 500 505 510
 Gln Leu Arg Asp Gln Cys Ile Val Asp Asp Ile Thr Tyr Asn Val Asn
 515 520 525
 Asp Thr Phe His Lys Arg His Glu Glu Gly His Met Leu Asn Cys Thr
 530 535 540
 Cys Phe Gly Gln Gly Arg Gly Arg Trp Lys Cys Asp Pro Val Asp Gln
 545 550 555 560
 Cys Gln Asp Ser Glu Thr Gly Thr Phe Tyr Gln Ile Gly Asp Ser Trp
 565 570 575
 Glu Lys Tyr Val His Gly Val Arg Tyr Gln Cys Tyr Cys Tyr Gly Arg
 580 585 590
 Gly Ile Gly Glu Trp His Cys Gln Pro Leu Gln Thr Tyr Pro Ser Ser
 595 600 605
 Ser Gly Pro Val Glu Val Phe Ile Thr Glu Thr Pro Ser Gln Pro Asn
 610 615 620
 Ser His Pro Ile Gln Trp Asn Ala Pro Gln Pro Ser His Ile Ser Lys
 625 630 635 640
 Tyr Ile Leu Arg Trp Arg Pro Lys Asn Ser Val Gly Arg Trp Lys Glu
 645 650 655
 Ala Thr Ile Pro Gly His Leu Asn Ser Tyr Thr Ile Lys Gly Leu Lys
 660 665 670
 Pro Gly Val Val Tyr Glu Gly Gln Leu Ile Ser Ile Gln Gln Tyr Gly
 675 680 685
 His Gln Glu Val Thr Arg Phe Asp Phe Thr Thr Thr Ser Thr Ser Thr

-continued

690				695				700							
Pro	Val	Thr	Ser	Asn	Thr	Val	Thr	Gly	Glu	Thr	Thr	Pro	Phe	Ser	Pro
705					710					715					720
Leu	Val	Ala	Thr	Ser	Glu	Ser	Val	Thr	Glu	Ile	Thr	Ala	Ser	Ser	Phe
				725					730					735	
Val	Val	Ser	Trp	Val	Ser	Ala	Ser	Asp	Thr	Val	Ser	Gly	Phe	Arg	Val
			740					745					750		
Glu	Tyr	Glu	Leu	Ser	Glu	Glu	Gly	Asp	Glu	Pro	Gln	Tyr	Leu	Asp	Leu
		755					760					765			
Pro	Ser	Thr	Ala	Thr	Ser	Val	Asn	Ile	Pro	Asp	Leu	Leu	Pro	Gly	Arg
		770				775					780				
Lys	Tyr	Ile	Val	Asn	Val	Tyr	Gln	Ile	Ser	Glu	Asp	Gly	Glu	Gln	Ser
785					790					795					800
Leu	Ile	Leu	Ser	Thr	Ser	Gln	Thr	Thr	Ala	Pro	Asp	Ala	Pro	Pro	Asp
				805					810					815	
Thr	Thr	Val	Asp	Gln	Val	Asp	Asp	Thr	Ser	Ile	Val	Val	Arg	Trp	Ser
			820					825					830		
Arg	Pro	Gln	Ala	Pro	Ile	Thr	Gly	Tyr	Arg	Ile	Val	Tyr	Ser	Pro	Ser
		835					840					845			
Val	Glu	Gly	Ser	Ser	Thr	Glu	Leu	Asn	Leu	Pro	Glu	Thr	Ala	Asn	Ser
		850				855					860				
Val	Thr	Leu	Ser	Asp	Leu	Gln	Pro	Gly	Val	Gln	Tyr	Asn	Ile	Thr	Ile
865					870					875					880
Tyr	Ala	Val	Glu	Glu	Asn	Gln	Glu	Ser	Thr	Pro	Val	Val	Ile	Gln	Gln
				885					890					895	
Glu	Thr	Thr	Gly	Thr	Pro	Arg	Ser	Asp	Thr	Val	Pro	Ser	Pro	Arg	Asp
			900					905					910		
Leu	Gln	Phe	Val	Glu	Val	Thr	Asp	Val	Lys	Val	Thr	Ile	Met	Trp	Thr
		915					920					925			
Pro	Pro	Glu	Ser	Ala	Val	Thr	Gly	Tyr	Arg	Val	Asp	Val	Ile	Pro	Val
		930				935					940				
Asn	Leu	Pro	Gly	Glu	His	Gly	Gln	Arg	Leu	Pro	Ile	Ser	Arg	Asn	Thr
945					950					955					960
Phe	Ala	Glu	Val	Thr	Gly	Leu	Ser	Pro	Gly	Val	Thr	Tyr	Tyr	Phe	Lys
				965					970					975	
Val	Phe	Ala	Val	Ser	His	Gly	Arg	Glu	Ser	Lys	Pro	Leu	Thr	Ala	Gln
			980					985					990		
Gln	Thr	Thr	Lys	Leu	Asp	Ala	Pro	Thr	Asn	Leu	Gln	Phe	Val	Asn	Glu
		995					1000					1005			
Thr	Asp	Ser	Thr	Val	Leu	Val	Arg	Trp	Thr	Pro	Pro	Arg	Ala	Gln	Ile
	1010					1015					1020				
Thr	Gly	Tyr	Arg	Leu	Thr	Val	Gly	Leu	Thr	Arg	Arg	Gly	Gln	Pro	Arg
1025					1030					1035					1040
Gln	Tyr	Asn	Val	Gly	Pro	Ser	Val	Ser	Lys	Tyr	Pro	Leu	Arg	Asn	Leu
				1045					1050					1055	
Gln	Pro	Ala	Ser	Glu	Tyr	Thr	Val	Ser	Leu	Val	Ala	Ile	Lys	Gly	Asn
			1060					1065					1070		
Gln	Glu	Ser	Pro	Lys	Ala	Thr	Gly	Val	Phe	Thr	Thr	Leu	Gln	Pro	Gly
		1075					1080					1085			
Ser	Ser	Ile	Pro	Pro	Tyr	Asn	Thr	Glu	Val	Thr	Glu	Thr	Thr	Ile	Val
	1090					1095					1100				

-continued

Ile Thr Trp Thr Pro Ala Pro Arg Ile Gly Phe Lys Leu Gly Val Arg
 1105 1110 1115 1120
 Pro Ser Gln Gly Gly Glu Ala Pro Arg Glu Val Thr Ser Asp Ser Gly
 1125 1130 1135
 Ser Ile Val Val Ser Gly Leu Thr Pro Gly Val Glu Tyr Val Tyr Thr
 1140 1145 1150
 Ile Gln Val Leu Arg Asp Gly Gln Glu Arg Asp Ala Pro Ile Val Asn
 1155 1160 1165
 Lys Val Val Thr Pro Leu Ser Pro Pro Thr Asn Leu His Leu Glu Ala
 1170 1175 1180
 Asn Pro Asp Thr Gly Val Leu Thr Val Ser Trp Glu Arg Ser Thr Thr
 1185 1190 1195 1200
 Pro Asp Ile Thr Gly Tyr Arg Ile Thr Thr Thr Pro Thr Asn Gly Gln
 1205 1210 1215
 Gln Gly Asn Ser Leu Glu Glu Val Val His Ala Asp Gln Ser Ser Cys
 1220 1225 1230
 Thr Phe Asp Asn Leu Ser Pro Gly Leu Glu Tyr Asn Val Ser Val Tyr
 1235 1240 1245
 Thr Val Lys Asp Asp Lys Glu Ser Val Pro Ile Ser Asp Thr Ile Ile
 1250 1255 1260
 Pro Ala Val Pro Pro Pro Thr Asp Leu Arg Phe Thr Asn Ile Gly Pro
 1265 1270 1275 1280
 Asp Thr Met Arg Val Thr Trp Ala Pro Pro Ser Ile Asp Leu Thr
 1285 1290 1295
 Asn Phe Leu Val Arg Tyr Ser Pro Val Lys Asn Glu Glu Asp Val Ala
 1300 1305 1310
 Glu Leu Ser Ile Ser Pro Ser Asp Asn Ala Val Val Leu Thr Asn Leu
 1315 1320 1325
 Leu Pro Gly Thr Glu Tyr Val Val Ser Val Ser Ser Val Tyr Glu Gln
 1330 1335 1340
 His Glu Ser Thr Pro Leu Arg Gly Arg Gln Lys Thr Gly Leu Asp Ser
 1345 1350 1355 1360
 Pro Thr Gly Ile Asp Phe Ser Asp Ile Thr Ala Asn Ser Phe Thr Val
 1365 1370 1375
 His Trp Ile Ala Pro Arg Ala Thr Ile Thr Gly Tyr Arg Ile Arg His
 1380 1385 1390
 His Pro Glu His Phe Ser Gly Arg Pro Arg Glu Asp Arg Val Pro His
 1395 1400 1405
 Ser Arg Asn Ser Ile Thr Leu Thr Asn Leu Thr Pro Gly Thr Glu Tyr
 1410 1415 1420
 Val Val Ser Ile Val Ala Leu Asn Gly Arg Glu Glu Ser Pro Leu Leu
 1425 1430 1435 1440
 Ile Gly Gln Gln Ser Thr Val Ser Asp Val Pro Arg Asp Leu Glu Val
 1445 1450 1455
 Val Ala Ala Thr Pro Thr Ser Leu Leu Ile Ser Trp Asp Ala Pro Ala
 1460 1465 1470
 Val Thr Val Arg Tyr Tyr Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn
 1475 1480 1485
 Ser Pro Val Gln Glu Phe Thr Val Pro Gly Ser Lys Ser Thr Ala Thr
 1490 1495 1500

-continued

1905	1910	1915	1920
Val Ser Trp Gln Pro	Pro Arg Ala Arg	Ile Thr Gly Tyr Ile	Ile Lys
	1925	1930	1935
Tyr Glu Lys Pro	Gly Ser Pro Pro	Arg Glu Val Val	Pro Arg Pro Arg
	1940	1945	1950
Pro Gly Val Thr	Glu Ala Thr Ile	Thr Gly Leu Glu	Pro Gly Thr Glu
	1955	1960	1965
Tyr Thr Ile Tyr	Val Ile Ala Leu	Lys Asn Asn Gln	Lys Ser Glu Pro
	1970	1975	1980
Leu Ile Gly Arg	Lys Lys Thr Asp	Glu Leu Pro Gln	Leu Val Thr Leu
1985	1990	1995	2000
Pro His Pro Asn	Leu His Gly Pro	Glu Ile Leu Asp	Val Pro Ser Thr
	2005	2010	2015
Val Gln Lys Thr	Pro Phe Val Thr	His Pro Gly Tyr	Asp Thr Gly Asn
	2020	2025	2030
Gly Ile Gln Leu	Pro Gly Thr Ser	Gly Gln Gln Pro	Ser Val Gly Gln
	2035	2040	2045
Gln Met Ile Phe	Glu Glu His Gly	Phe Arg Arg Thr	Thr Pro Pro Thr
	2050	2055	2060
Thr Ala Thr Pro	Ile Arg His Arg	Pro Arg Pro Tyr	Pro Pro Asn Val
2065	2070	2075	2080
Gly Gln Glu Ala	Leu Ser Gln Thr	Thr Ile Ser Trp	Ala Pro Phe Gln
	2085	2090	2095
Asp Thr Ser Glu	Tyr Ile Ile Ser	Cys His Pro Val	Gly Thr Asp Glu
	2100	2105	2110
Glu Pro Leu Gln	Phe Arg Val Pro	Gly Thr Ser Thr	Ser Ala Thr Leu
	2115	2120	2125
Thr Gly Leu Thr	Arg Gly Ala Thr	Tyr Asn Val Ile	Val Glu Ala Leu
	2130	2135	2140
Lys Asp Gln Gln	Arg His Lys Val	Arg Glu Glu Val	Val Thr Val Gly
2145	2150	2155	2160
Asn Ser Val Asn	Glu Gly Leu Asn	Gln Pro Thr Asp	Asp Ser Cys Phe
	2165	2170	2175
Asp Pro Tyr Thr	Val Ser His Tyr	Ala Val Gly Asp	Glu Trp Glu Arg
	2180	2185	2190
Met Ser Glu Ser	Gly Phe Lys Leu	Leu Cys Gln Cys	Leu Gly Phe Gly
	2195	2200	2205
Ser Gly His Phe	Arg Cys Asp Ser	Ser Ser	
	2210	2215	

<210> SEQ ID NO 39

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 39

Val Thr His Pro Gly Tyr

1

5

1. A method for assessing whether a subject has an increased likelihood of imminent or preterm delivery, comprising:

detecting an oncofetal fibronectin indicating molecule in a sample from a pregnant subject, wherein:

the sample is selected from among serum, plasma, blood, urine, a body tissue, lavage and cervical vaginal fluid sampled from among cervical canal, cervical os, ectocervix, transition zone on the cervix between squamous and columnar cells, posterior fornix, a portion of the vagina below the posterior fornix, lower third of the vagina, labia, cervical interstitial fluid; and

the presence of the oncofetal fibronectin indicating molecule in the sample indicates that the subject has an increased likelihood of imminent or preterm delivery.

2. The method of claim 1, wherein the sample is urine.

3. The method of claim 1, wherein the amount of oncofetal fibronectin detected is correlated with the likelihood of imminent or preterm delivery

4. The method of claim 1, wherein presence of an amount of oncofetal fibronectin indicating molecule at or above a threshold level indicates that the subject has an increased likelihood of imminent or preterm delivery.

5. The method of claim 1, wherein:

the amount of oncofetal fibronectin indicating molecule detected is compared to two or more thresholds; and

the sample is classified according to the highest threshold that is less than or equal to the detected amount of oncofetal fibronectin indicating molecule, whereby classification of oncofetal fibronectin indicating molecule in the sample according to the highest threshold indicates that the subject has an increased likelihood of imminent or preterm delivery.

6. The method of claim 1, wherein:

the oncofetal fibronectin indicating molecule is selected from among an oncofetal fibronectin protein or a fragment thereof, a nucleic acid molecule encoding oncofetal fibronectin or a fragment thereof, a nucleic acid molecule complementary to a nucleic acid molecule encoding oncofetal fibronectin or a fragment thereof, an autoantibody for oncofetal fibronectin protein or a fragment thereof and an autoantibody for a nucleic acid molecule encoding oncofetal fibronectin or a fragment thereof.

7. The method of claim 1, wherein:

detecting the oncofetal fibronectin indicating molecule further comprises:

contacting the sample with a first oncofetal fibronectin binding partner; and

detecting complexes of the binding partner and oncofetal fibronectin, whereby detection of a complex is indicative of the presence of oncofetal fibronectin indicating molecule in the sample.

8. The method of claim 6 further comprising contacting a sample from the subject with a non-specific binding compound.

9. The method of claim 7, wherein the binding partner is detectably labeled.

10. The method of claim 1, wherein:

detecting the oncofetal fibronectin indicating molecule comprises:

contacting the sample with a first fibronectin or oncofetal fibronectin binding partner;

contacting the sample with a second fibronectin or oncofetal fibronectin binding partner, wherein:

the second fibronectin or oncofetal fibronectin binding partner is conjugated to a detectable or bindable moiety, or the second fibronectin or oncofetal fibronectin binding partner is immobilized to a solid support.

11. The method of claim 7, wherein the binding partner is detectably labeled.

12. The method of claim 1, wherein the oncofetal fibronectin indicating molecule is detected by its molecular weight or the weight of a fragment thereof.

13. The method of claim 12, wherein the oncofetal fibronectin indicating molecule is detected by mass spectrometry or gel electrophoresis.

14. The method of claim 7, wherein the fibronectin or oncofetal fibronectin binding partner is immobilized on a test strip.

15. A method for determining whether a subject is a candidate for successful induction of labor, comprising:

determining the amount of an oncofetal fibronectin indicating molecule in a sample from a pregnant subject; wherein

if the amount of oncofetal fibronectin indicating molecule is equal to or above a threshold level, the subject is identified as one likely to have a successful induction.

16. The method of claim 15, wherein a subject who has a successful induction is an induction that results in vaginal delivery, shorter time to delivery or fewer administrations of induction or pre-induction agents compared to a subject that does not have a successful induction.

17. The method of claim 15, further comprising:

administering to the subject having an amount of oncofetal fibronectin indicating molecule equal to or above the threshold level a dose of an agent or procedure effective to induce labor.

18. The method of claim 17, further comprising:

evaluating the effectiveness of induction by determining the amount of an oncofetal fibronectin indicating molecule in a sample from a pregnant subject undergoing an induction procedure; and

if the amount of oncofetal fibronectin indicating molecule is equal to or above threshold level, identifying the subject as one who is likely to have a successful induction.

19. The method of claim 15, further comprising:

determining a second indicator of induction outcome for the subject, wherein:

if the amount of oncofetal fibronectin indicating molecule is above threshold level and the second indicator indicates favorable induction outcome, the subject is identified as one for whom induction is likely to be successful.

20. The method of claim 19, wherein the second indicator of induction outcome is selected from among measurements

or observations of the pregnant subject, a measurement or observation of the fetus(es), and medical history of the pregnant subject.

21. The method of claim 20, wherein the second indicator is selected from among cervical length, Bishop score, effacement, parity, cervical dilation, gestational age, body mass index, station, consistency, transvaginal ultrasound, and digital examination.

22. The method of claim 19, further comprising:

identifying the subject as one for whom induction is likely to be successful; and

administering an induction agent to the subject or subjecting the subject to a procedure.

23. The method of claim 15, wherein the sample is collected with a swab selected from among a polyester swab, a cotton swab and a rayon swab.

24. The method of claim 15, wherein the sample is contacted with a non-specific binding compound prior to determining the amount of the oncofetal fibronectin indicating molecule.

25. The method of claim 15, wherein the oncofetal fibronectin indicating molecule or a fragment thereof is detected by mass spectrometry or gel electrophoresis.

26. The method of claim 15, further comprising contacting the sample with a fibronectin or oncofetal fibronectin binding protein; and

detecting the complexes of the oncofetal fibronectin indicating molecule and the fibronectin or oncofetal fibronectin binding protein, wherein the complex is measured by detecting the fibronectin or oncofetal fibronectin binding partner.

27. The method of claim 26, wherein the fibronectin or oncofetal fibronectin binding partner is immobilized on a test strip.

28. The method of claim 1, wherein:

determining the amount of oncofetal fibronectin indicating molecule comprises:

contacting the sample with an immunoassay test strip.

29. The method of claim 28, wherein the test strip contains:

a mobilizable oncofetal fibronectin binding partner conjugated to a detectable moiety and a fibronectin or oncofetal fibronectin binding partner immobilized to the test strip; and

detecting complexes of the mobilizable binding partner, the oncofetal fibronectin indicating molecule, and the immobilized binding partner, whereby detection of a complex is indicative of the amount of oncofetal fibronectin indicating molecule in the sample.

30. A method for delivery date prediction, comprising:

measuring an oncofetal fibronectin indicating molecule in a sample from a pregnant subject, wherein:

an amount of oncofetal fibronectin indicating molecule in the sample at or above a threshold level indicates an increased likelihood that the subject will deliver within a particular time period.

31. The method of claim 30, wherein:

the particular time period is selected from among 3 weeks or less, 2 weeks or less, 10 days or less, 1 week or less, 6 days or less, 5 days or less, 4 days or less, 3 days or less, 2 days or less, or 1 day or less.

32. A combination, comprising:

a fibronectin or oncofetal fibronectin binding partner and a parturifacient.

33. The combination of claim 32, further comprising a non-specific binding compound.

34. A kit, comprising the combination of claim 32 and optionally instructions for use of the binding partner and parturifacient for assessing whether a subject has an increased likelihood of imminent or preterm delivery.

35. A kit, comprising:

a fibronectin or oncofetal fibronectin binding partner, a parturifacient, and

a system for classifying the sample according to one or more threshold levels.

* * * * *