

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2017/151746 A1

(43) International Publication Date
8 September 2017 (08.09.2017)

WIPO | PCT

(51) International Patent Classification:

C07D 498/10 (2006.01) A61P 7/02 (2006.01)
A61K 31/537 (2006.01)

(21) International Application Number:

PCT/US2017/020161

(22) International Filing Date:

1 March 2017 (01.03.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/302,220 2 March 2016 (02.03.2016) US

(71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; Route 206 and Province Line Road, Princeton, New Jersey 08543 (US).

(72) Inventors: SHI, Jun; c/o Bristol-Myers Squibb Drive, 311 Pennington-Rocky Hill Road, Pennington, New Jersey 08534 (US). EWING, William R.; c/o Bristol-Myers Squibb Company, Route 206 and Province Line Road, Princeton, New Jersey 08543 (US). PINTO, Donald J.P.; c/o Bristol-Myers Squibb Company, 311 Pennington-Rocky Hill Road, Princeton, New Jersey 08534 (US).

(74) Agents: LIU, Hong et al.; Bristol-Myers Squibb Company, Route 206 & Province Line Road, Princeton, New Jersey 08543 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))

(54) Title: DIAMIDE MACROCYCLES HAVING FACTOR XIA INHIBITING ACTIVITY



(I)

(57) Abstract: The present invention provides compounds of Formula (I): or stereoisomers, tautomers, or pharmaceutically acceptable salts thereof, wherein all the variables are as defined herein. These compounds are selective Factor XIa inhibitors or dual inhibitors of FXIa and plasma kallikrein. This invention also relates to pharmaceutical compositions comprising these compounds and methods of treating thromboembolic and/or inflammatory disorders using the same.

WO 2017/151746 A1

DIAMIDE MACROCYCLES HAVING FACTOR XIa INHIBITING ACTIVITY

CROSS REFERENCE TO RELATED APPLICATIONS

5 This application is entitled to priority pursuant to 35 U.S.C. §119(e) to U.S. provisional patent application No. 62/302,220, filed March 2, 2016, which is incorporated herein in its entirety.

FIELD OF THE INVENTION

10 The present invention relates generally to novel macrocyclic compounds, and their analogues thereof, which are inhibitors of Factor XIa and/or plasma kallikrein, compositions containing them, and methods of using them, for example, for the treatment or prophylaxis of thromboembolic disorders, or for the treatment of retinal vascular permeability associated with diabetic retinopathy and diabetic macular edema.

15 BACKGROUND OF THE INVENTION

Thromboembolic diseases remain the leading cause of death in developed countries despite the availability of anticoagulants such as warfarin (COUMADIN®), heparin, low molecular weight heparins (LMWH), and synthetic pentasaccharides and antiplatelet agents such as aspirin and clopidogrel (PLAVIX®). The oral anticoagulant warfarin, inhibits the post-translational maturation of coagulation Factors VII, IX, X and prothrombin, and has proven effective in both venous and arterial thrombosis. However, its usage is limited due to its narrow therapeutic index, slow onset of therapeutic effect, numerous dietary and drug interactions, and a need for monitoring and dose adjustment. Thus discovering and developing safe and efficacious oral anticoagulants for the prevention and treatment of a wide range of thromboembolic disorders has become increasingly important.

30 One approach is to inhibit thrombin generation by targeting the inhibition of coagulation Factor XIa (FXIa). Factor XIa is a plasma serine protease involved in the regulation of blood coagulation, which is initiated *in vivo* by the binding of tissue factor (TF) to Factor VII (FVII) to generate Factor VIIa (FVIIa). The resulting TF:FVIIa complex activates Factor IX (FIX) and Factor X (FX) that leads to the production of Factor Xa (FXa). The generated FXa catalyzes the transformation of prothrombin into

small amounts of thrombin before this pathway is shut down by tissue factor pathway inhibitor (TFPI). The process of coagulation is then further propagated via the feedback activation of Factors V, VIII and XI by catalytic amounts of thrombin. (Gailani, D. et al., *Arterioscler. Thromb. Vasc. Biol.*, 27:2507-2513 (2007).) The resulting burst of thrombin 5 converts fibrinogen to fibrin that polymerizes to form the structural framework of a blood clot, and activates platelets, which are a key cellular component of coagulation (Hoffman, M., *Blood Reviews*, 17:S1-S5 (2003)). Therefore, Factor XIa plays a key role in propagating this amplification loop and is thus an attractive target for anti-thrombotic therapy.

10 Plasma prekallikrein is a zymogen of a trypsin-like serine protease and is present in plasma at 35 to 50 µg/mL. The gene structure is similar to that of Factor XI. Overall, the amino acid sequence of plasma kallikrein has 58% homology to Factor XI. Plasma kallikrein is thought to play a role in a number of inflammatory disorders. The major inhibitor of plasma kallikrein is the serpin C1 esterase inhibitor. Patients who present with 15 a genetic deficiency in C1 esterase inhibitor suffer from hereditary angioedema (HAE) which results in intermittent swelling of face, hands, throat, gastro-intestinal tract and genitals. Blisters formed during acute episodes contain high levels of plasma kallikrein which cleaves high molecular weight kininogen liberating bradykinin leading to increased vascular permeability. Treatment with a large protein plasma kallikrein inhibitor has been 20 shown to effectively treat HAE by preventing the release of bradykinin which causes increased vascular permeability (Lehmann, A., "Ecallantide (DX-88), a plasma kallikrein inhibitor for the treatment of hereditary angioedema and the prevention of blood loss in on-pump cardiothoracic surgery", *Expert Opin. Biol. Ther.*, 8:1187-1199 (2008)).

25 The plasma kallikrein-kinin system is abnormally abundant in patients with advanced diabetic macular edema. It has been recently published that plasma kallikrein contributes to retinal vascular dysfunctions in diabetic rats (Clermont, A. et al., "Plasma kallikrein mediates retinal vascular dysfunction and induces retinal thickening in diabetic rats", *Diabetes*, 60:1590-1598 (2011)). Furthermore, administration of the plasma kallikrein inhibitor ASP-440 ameliorated both retinal vascular permeability and retinal 30 blood flow abnormalities in diabetic rats. Therefore, a plasma kallikrein inhibitor should have utility as a treatment to reduce retinal vascular permeability associated with diabetic retinopathy and diabetic macular edema. Other complications of diabetes such as cerebral

hemorrhage, nephropathy, cardiomyopathy and neuropathy, all of which have associations with plasma kallikrein may also be considered as targets for a plasma kallikrein inhibitor.

To date, no small molecule synthetic plasma kallikrein inhibitor has been 5 approved for medical use. The large protein plasma kallikrein inhibitors present risks of anaphylactic reactions, as has been reported for Ecallantide. Thus there remains a need for compounds that inhibit plasma kallikrein, that do not induce anaphylaxis and that are orally available. Furthermore, the molecules in the known art feature a highly polar and ionizable guanidine or amidine functionality. It is well known that such functionalities 10 may be limiting to gut permeability and therefore to oral availability.

SUMMARY OF THE INVENTION

The present invention provides novel macrocyclic compounds, their analogues, including stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof, 15 which are useful as selective inhibitors of serine protease enzymes, especially Factor XIa and/or plasma kallikrein.

The present invention also provides processes and intermediates for making the compounds of the present invention.

The present invention also provides pharmaceutical compositions comprising a 20 pharmaceutically acceptable carrier and at least one of the compounds of the present invention or stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof.

The compounds of the invention may be used in the treatment and/or prophylaxis of thromboembolic disorders.

25 The compounds of the invention may be used in the treatment of retinal vascular permeability associated with diabetic retinopathy and diabetic macular edema.

The compounds of the present invention may be used in therapy.

The compounds of the present invention may be used for the manufacture of a medicament for the treatment and/or prophylaxis of a thromboembolic disorder.

30 The compounds of the invention can be used alone, in combination with other compounds of the present invention, or in combination with one or more, preferably one to two other agent(s).


These and other features of the invention will be set forth in expanded form as the disclosure continues.

DETAILED DESCRIPTION OF THE INVENTION

5

I. COMPOUNDS OF THE INVENTION

In one aspect, the present invention provides, *inter alia*, compounds of Formula (I):



10 or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein:

---- is an optional bond;

ring A is independently selected from a mono or bicyclic heterocycle comprising: carbon atoms and 1-4 heteroatoms selected from NR², O, and S;

15 ring B is independently selected from aryl and a 5- to 10-membered heterocycle;

R¹ is, independently at each occurrence, selected from halogen, C₁₋₆ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, OH, =O, OCH₂F, OCHF₂, OCF₃, CF₃, CN, NH₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, CO₂(C₁₋₄ alkyl), CO(C₁₋₄ alkyl), -CH₂NH₂, -CONH₂, -CONH(C₁₋₄ alkyl), -OCH₂CO₂H, -NHCO(C₁₋₄ alkyl), -NHCO₂(C₁₋₄ alkyl), -NHSO₂(C₁₋₄ alkyl), -SO₂NH₂, and -C(=NH)NH₂;

20 R² is absent or independently selected from H and C₁₋₄alkyl;

R³ is independently selected from H, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, OH, =O, OCH₂F, OCHF₂, OCF₃, CN, NH₂, NH(C₁₋₄ alkyl), and N(C₁₋₄ alkyl)₂;

R⁴ is independently selected from H, C₁₋₄ alkyl, hydroxyl, F, CF₃, and C₃₋₆ cycloalkyl;

R⁵ is independently selected from H and C₁₋₄ alkyl;

R⁶ is independently selected from H, halogen, C(O)OH, and C(O)O(C₁₋₄ alkyl);


5 R⁷ is independently selected from H, C₁₋₄ alkyl, and CF₃;

alternatively, R⁶ and R⁷ together are =O; and

R⁸ is, independently at each occurrence, selected from H, halogen, NHCO₂C₁₋₄ alkyl, CN, OH, O-C₁₋₄ alkyl, CF₃, -OCF₃, -OCHF₂, -CHF₂, -CH₂F, CO₂H, CO₂(C₁₋₄ alkyl), -CH₂CO₂H, -(CH₂)₂CO₂H, -CH₂CO₂(C₁₋₄ alkyl), -(CH₂)₂CO₂(C₁₋₄ alkyl),

10 NH₂, -CH₂NH₂, -NHCO(C₁₋₄ alkyl), -NHCO₂(CH₂)₁₋₂O(C₁₋₄ alkyl), -NHCO₂(CH₂)₁₋₃O(C₁₋₄ alkyl), NHCO₂CH₂CH(C₁₋₄ alkyl)O(C₁₋₄ alkyl), -NHCO₂(CH₂)₁₋₂OH, -NHCO₂CH₂CO₂H, -CH₂NHCO₂(C₁₋₄ alkyl), -NHC(O)NH(C₁₋₄ alkyl), -NHC(O)N(C₁₋₄ alkyl)₂, NHC(O)NH(C₁₋₄ alkyl)N(5- to 6-membered heterocycle), -NHSO₂(C₁₋₄ alkyl), -CONH₂, -CONH(C₁₋₄ alkyl), -CON(C₁₋₄ alkyl)₂, and -CH₂CONH₂.

In another aspect, the present invention provides compounds of Formula (II):



or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs

20 thereof, wherein:

---- is an optional bond;

ring A is independently selected from a mono or bicyclic heterocycle comprising: carbon atoms and 1-4 heteroatoms selected from NR², O, and S;

R¹ is, independently at each occurrence, selected from halogen, C₁₋₆ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, OH, =O, OCH₂F, OCHF₂, OCF₃, CN, NH₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, CO₂(C₁₋₄ alkyl), CO(C₁₋₄ alkyl), -CH₂NH₂, -CONH₂, -CONH(C₁₋₄ alkyl), -OCH₂CO₂H, -NHCO(C₁₋₄ alkyl), -NHCO₂(C₁₋₄ alkyl), -NHSO₂(C₁₋₄ alkyl), -SO₂NH₂, and -C(=NH)NH₂;

5 R² is absent or independently selected from H and C₁₋₄alkyl;


R³ is independently selected from H and C₁₋₄ alkyl;

R⁴ is independently selected from H, C₁₋₄ alkyl, hydroxyl, and C₃₋₆ cycloalkyl;

R⁵ is independently selected from H and C₁₋₄ alkyl; and

10 R⁸ is independently selected from H, halogen, NHCO₂C₁₋₄ alkyl, CN, OH, O-C₁₋₄ alkyl, CF₃, -OCF₃, -OCHF₂, -CHF₂, -CH₂F, CO₂H, CO₂(C₁₋₄ alkyl), -CH₂CO₂H, -(CH₂)₂CO₂H, -CH₂CO₂(C₁₋₄ alkyl), -(CH₂)₂CO₂(C₁₋₄ alkyl), NH₂, -CH₂NH₂, -NHCO(C₁₋₄ alkyl), -NHCO₂(CH₂)₁₋₂O(C₁₋₄ alkyl), -NHCO₂(CH₂)₁₋₃O(C₁₋₄ alkyl), NHCO₂CH₂CH(C₁₋₄ alkyl)O(C₁₋₄ alkyl), -NHCO₂(CH₂)₁₋₂OH, -NHCO₂CH₂CO₂H, -CH₂NHCO₂(C₁₋₄ alkyl), -NHC(O)NH(C₁₋₄ alkyl), -NHC(O)N(C₁₋₄ alkyl)₂, NHC(O)NH(C₁₋₄ alkyl)N[5- to 15 6-membered heterocycle], -NHSO₂(C₁₋₄ alkyl), -CONH₂, -CONH(C₁₋₄ alkyl), -CON(C₁₋₄ alkyl)₂, and -CH₂CONH₂.

In another aspect, the present invention provides compounds of Formula (III):



20 or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein:

ring C is independently selected from 6-membered aryl and heteroaryl;

R¹ is, independently at each occurrence, selected from halogen, C₁₋₆ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, OH, OCH₂F, OCHF₂, OCF₃, CN, NH₂, NH(C₁₋₄ alkyl), and N(C₁₋₄ alkyl)₂;

5 R² is independently selected from H and C₁₋₄alkyl;


R⁴ is independently selected from H, C₁₋₄ alkyl, hydroxyl, and C₃₋₆ cycloalkyl;

R⁵ is independently selected from H and C₁₋₄ alkyl; and

10 R⁸ is independently selected from H, halogen, NHCO₂C₁₋₄ alkyl, CN, OH, O-C₁₋₄ alkyl, CF₃, -OCF₃, -OCHF₂, -CHF₂, -CH₂F, CO₂H, and CO₂(C₁₋₄ alkyl).

10

In another aspect, the present invention provides compounds of Formula (IV):



or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein:

15 ---- is an optional bond;

R¹ is independently selected from F and Cl;

R² is independently selected from H and C₁₋₄alkyl;

R⁴ is independently selected from H, C₁₋₄ alkyl, and hydroxyl;

R⁵ is independently selected from H and C₁₋₄ alkyl; and

20 R⁸ is independently selected from H, halogen, NHCO₂C₁₋₄ alkyl, CN, OH, O-C₁₋₄ alkyl, CF₃, -OCF₃, -OCHF₂, -CHF₂, -CH₂F, CO₂H, and CO₂(C₁₋₄ alkyl).

In another aspect, the present invention provides compounds of Formula (IV), or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein:

---- is an optional bond;

5 R¹ is independently selected from F and Cl;

R² is H;





R⁴ is independently selected from H, C₁₋₄ alkyl, hydroxyl, and C₃₋₆ cycloalkyl;

R⁵ is independently selected from H and C₁₋₄ alkyl; and

R⁸ is independently selected from H, halogen, and NHCO₂C₁₋₄ alkyl; and

10 other variables are as defined in Formula (IV) above.

In another aspect, the present invention provides compounds selected from:



15 or a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof.

In another aspect, the present invention provides a compound selected from any subset list of compounds exemplified in the present application.

In another embodiment, the compounds of the present invention have Factor XIa

20 or plasma kallikrein Ki values $\leq 10 \mu\text{M}$.

In another embodiment, the compounds of the present invention have Factor XIa or plasma kallikrein Ki values $\leq 1 \mu\text{M}$.

In another embodiment, the compounds of the present invention have Factor XIa or plasma kallikrein Ki values $\leq 0.5 \mu\text{M}$.

5 In another embodiment, the compounds of the present invention have Factor XIa or plasma kallikrein Ki values $\leq 0.1 \mu\text{M}$.

II. OTHER EMBODIMENTS OF THE INVENTION

In another embodiment, the present invention provides a composition comprising 10 at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

In another embodiment, the present invention provides a pharmaceutical 15 composition comprising a pharmaceutically acceptable carrier and at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate, thereof.

In another embodiment, the present invention provides a pharmaceutical 20 composition, comprising: a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

In another embodiment, the present invention provides a process for making a 25 compound of the present invention.

In another embodiment, the present invention provides an intermediate for making a compound of the present invention.

In another embodiment, the present invention provides a pharmaceutical 30 composition further comprising additional therapeutic agent(s). In a preferred embodiment, the present invention provides pharmaceutical composition, wherein the additional therapeutic agent(s) are an anti-platelet agent or a combination thereof.

Preferably, the anti-platelet agent(s) are clopidogrel and/or aspirin, or a combination thereof.

In another embodiment, the present invention provides a method for the treatment 35 and/or prophylaxis of a thromboembolic disorder comprising administering to a patient in need of such treatment and/or prophylaxis a therapeutically effective amount of at least

one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

In another embodiment, the present invention provides a compound of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, for use in therapy.

In another embodiment, the present invention provides a compound of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, for use in therapy for the treatment and/or prophylaxis of a thromboembolic disorder.

10 In another embodiment, the present invention also provides the use of a compound of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, for the manufacture of a medicament for the treatment and/or prophylaxis of a thromboembolic disorder.

In another embodiment, the present invention provides a method for treatment and/or prophylaxis of a thromboembolic disorder, comprising: administering to a patient in need thereof a therapeutically effective amount of a first and second therapeutic agent, wherein the first therapeutic agent is a compound of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, and the second therapeutic agent is at least one agent selected from a Factor Xa inhibitor such as apixaban, rivaroxaban, betrixaban, edoxaban, an anti-coagulant agent, an anti-platelet agent, a thrombin inhibiting agent such as dabigatran, a thrombolytic agent, and a fibrinolytic agent. Preferably, the second therapeutic agent is at least one agent selected from warfarin, unfractionated heparin, low molecular weight heparin, synthetic pentasaccharide, hirudin, argatroban, aspirin, ibuprofen, naproxen, sulindac, indomethacin, mefenamate, droxicam, diclofenac, sulfinpyrazone, piroxicam, ticlopidine, clopidogrel, tirofiban, eptifibatide, abciximab, melagatran, desulfatohirudin, tissue plasminogen activator, modified tissue plasminogen activator, anistreplase, urokinase, and streptokinase. Preferably, the second therapeutic agent is at least one anti-platelet agent. Preferably, the anti-platelet agent(s) are clopidogrel and/or aspirin, or a combination thereof.

The thromboembolic disorder includes arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, arterial cerebrovascular

thromboembolic disorders, and venous cerebrovascular thromboembolic disorders. Examples of the thromboembolic disorder include, but are not limited to, unstable angina, an acute coronary syndrome, atrial fibrillation, first myocardial infarction, recurrent myocardial infarction, ischemic sudden death, transient ischemic attack, stroke, 5 atherosclerosis, peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an artificial surface that promotes thrombosis.

10 In another embodiment, the present invention provides a method for the treatment and/or prophylaxis of an inflammatory disorder comprising: administering to a patient in need of such treatment and/or prophylaxis a therapeutically effective amount of at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof. Examples of the inflammatory 15 disorder include, but are not limited to, sepsis, acute respiratory distress syndrome, and systemic inflammatory response syndrome.

20 In another embodiment, the present invention provides a method for the prophylaxis of a disease or condition in which plasma kallikrein activity is implicated comprising administering to a patient in need of such treatment and/or prophylaxis a therapeutically effective amount of at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

25 The disease or condition in which plasma kallikrein activity is implicated includes, but not limited to, impaired visual acuity, diabetic retinopathy, diabetic macular edema, hereditary angioedema, diabetes, pancreatitis, nephropathy, cardio myopathy, neuropathy, inflammatory bowel disease, arthritis, inflammation, septic shock, hypotension, cancer, adult respiratory distress syndrome, disseminated intravascular coagulation, and cardiopulmonary bypass surgery.

30 In another embodiment, the present invention provides a combined preparation of a compound of the present invention and additional therapeutic agent(s) for simultaneous, separate or sequential use in therapy.

In another embodiment, the present invention provides a combined preparation of a compound of the present invention and additional therapeutic agent(s) for simultaneous, separate or sequential use in treatment and/or prophylaxis of a thromboembolic disorder.

The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof. This invention encompasses all combinations of preferred aspects of the invention noted herein. It is understood that any and all embodiments of the present invention may be taken in conjunction with any other embodiment or embodiments to describe additional embodiments. It is also to be understood that each individual element of the embodiments is its own independent embodiment. Furthermore, any element of an embodiment is meant to be combined with any and all other elements from any embodiment to describe an additional embodiment.

III. CHEMISTRY

Throughout the specification and the appended claims, a given chemical formula or name shall encompass all stereo and optical isomers and racemates thereof where such isomers exist. Unless otherwise indicated, all chiral (enantiomeric and diastereomeric) and racemic forms are within the scope of the invention. Many geometric isomers of C=C double bonds, C=N double bonds, ring systems, and the like can also be present in the compounds, and all such stable isomers are contemplated in the present invention.

Cis- and *trans-* (or *E*- and *Z*-) geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. The present compounds can be isolated in optically active or racemic forms. Optically active forms may be prepared by resolution of racemic forms or by synthesis from optically active starting materials. All processes used to prepare compounds of the present invention and intermediates made therein are considered to be part of the present invention. When enantiomeric or diastereomeric products are prepared, they may be separated by conventional methods, for example, by chromatography or fractional crystallization. Depending on the process conditions the end products of the present invention are obtained either in free (neutral) or salt form. Both the free form and the salts of these end products are within the scope of the invention. If so desired, one form of a compound may be converted into another form. A free base or acid may be converted into a salt; a salt may be converted into the free compound or

another salt; a mixture of isomeric compounds of the present invention may be separated into the individual isomers. Compounds of the present invention, free form and salts thereof, may exist in multiple tautomeric forms, in which hydrogen atoms are transposed to other parts of the molecules and the chemical bonds between the atoms of the 5 molecules are consequently rearranged. It should be understood that all tautomeric forms, insofar as they may exist, are included within the invention.

The term "stereoisomer" refers to isomers of identical constitution that differ in the arrangement of their atoms in space. Enantiomers and diastereomers are examples of stereoisomers. The term "enantiomer" refers to one of a pair of molecular species that are 10 mirror images of each other and are not superimposable. The term "diastereomer" refers to stereoisomers that are not mirror images. The term "racemate" or "racemic mixture" refers to a composition composed of equimolar quantities of two enantiomeric species, wherein the composition is devoid of optical activity.

The symbols "R" and "S" represent the configuration of substituents around a 15 chiral carbon atom(s). The isomeric descriptors "R" and "S" are used as described herein for indicating atom configuration(s) relative to a core molecule and are intended to be used as defined in the literature (IUPAC Recommendations 1996, *Pure and Applied Chemistry*, 68:2193-2222 (1996)).

The term "chiral" refers to the structural characteristic of a molecule that makes it 20 impossible to superimpose it on its mirror image. The term "homochiral" refers to a state of enantiomeric purity. The term "optical activity" refers to the degree to which a homochiral molecule or nonracemic mixture of chiral molecules rotates a plane of polarized light.

As used herein, the term "alkyl" or "alkylene" is intended to include both 25 branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, "C₁ to C₁₀ alkyl" or "C₁₋₁₀ alkyl" (or alkylene), is intended to include C₁, C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, and C₁₀ alkyl groups. Additionally, for example, "C₁ to C₆ alkyl" or "C_{1-C6} alkyl" denotes alkyl having 1 to 6 carbon atoms. Alkyl group can be unsubstituted or substituted with at least one hydrogen being replaced 30 by another chemical group. Example alkyl groups include, but are not limited to, methyl (Me), ethyl (Et), propyl (e.g., n-propyl and isopropyl), butyl (e.g., n-butyl, isobutyl, t-

butyl), and pentyl (e.g., n-pentyl, isopentyl, neopentyl). When "C₀ alkyl" or "C₀ alkylene" is used, it is intended to denote a direct bond.

"Alkynyl" or "alkynylene" is intended to include hydrocarbon chains of either straight or branched configuration having one or more, preferably one to three, carbon-5 carbon triple bonds that may occur in any stable point along the chain. For example, "C₂ to C₆ alkynyl" or "C₂₋₆ alkynyl" (or alkynylene), is intended to include C₂, C₃, C₄, C₅, and C₆ alkynyl groups; such as ethynyl, propynyl, butynyl, pentynyl, and hexynyl.

The term "alkoxy" or "alkyloxy" refers to an -O-alkyl group. "C₁ to C₆ alkoxy" or "C₁₋₆ alkoxy" (or alkyloxy), is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ alkoxy 10 groups. Example alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy (e.g., n-propoxy and isopropoxy), and *t*-butoxy. Similarly, "alkylthio" or "thioalkoxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge; for example, methyl-S- and ethyl-S-.

"Halo" or "halogen" includes fluoro, chloro, bromo, and iodo. "Haloalkyl" is 15 intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogens. Examples of haloalkyl include, but are not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, trichloromethyl, pentafluoroethyl, pentachloroethyl, 2,2,2-trifluoroethyl, heptafluoropropyl, and heptachloropropyl. Examples of haloalkyl also include 20 "fluoroalkyl" that is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more fluorine atoms.

"Haloalkoxy" or "haloalkyloxy" represents a haloalkyl group as defined above 25 with the indicated number of carbon atoms attached through an oxygen bridge. For example, "C₁ to C₆ haloalkoxy" or "C₁₋₆ haloalkoxy", is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ haloalkoxy groups. Examples of haloalkoxy include, but are not limited to, trifluoromethoxy, 2,2,2-trifluoroethoxy, and pentafluoroethoxy. Similarly, 30 "haloalkylthio" or "thiohaloalkoxy" represents a haloalkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge; for example, trifluoromethyl-S-, and pentafluoroethyl-S-.

The term "alkoxyalkyl", as used herein, refers to an alkyl group substituted with one, two, or three alkoxy groups.

The term "amino", as used herein, refers to -NH₂.

The term "substituted amino", as used herein, refers to the defined terms below having the suffix "amino" such as "aryl amino", "alkyl amino", "aryl amino", etc.

5 The term "alkoxyalkyl amino", as used herein, refers to -NHR wherein R is an alkoxyalkyl group.

The term "alkoxycarbonyl", as used herein, refers to an alkoxy group attached to the parent molecular moiety through a carbonyl group.

The term "alkoxycarbonyl amino", as used herein, refers to an -NHR wherein R is an alkoxy carbonyl group.

10 The term "alkyl amino", as used herein refers to -NHR, wherein R is an alkyl group.

The term "alkyl carbonyl", as used herein, refers to an alkyl group attached to the parent molecular moiety through a carbonyl group.

15 The term "alkyl carbonyl amino", as used herein, refers to -NHR wherein R is an alkyl carbonyl group.

The term "aminosulfonyl", as used herein, refers to -SO₂NH₂.

The term "aryl alkyl", as used herein, refers to an alkyl group substituted with one, two, or three aryl groups.

The term "aryl amino", as used herein, refers to -NHR wherein R is an aryl group.

20 The term "aryl carbonyl", as used herein, refers to an aryl group attached to the parent molecular moiety through a carbonyl group.

The term "aryl carbonyl amino", as used herein refers to -NHR wherein R is an aryl carbonyl group.

The term "carbonyl", as used herein, refers to -C(O)-.

25 The term "cyano", as used herein, refers to -CN.

The term "cycloalkyl amino", as used herein, refers to -NHR wherein R is a cycloalkyl group.

The term "cycloalkyl carbonyl", as used herein, refers to a cycloalkyl group attached to the parent molecular moiety through a carbonyl group.

30 The term "cycloalkyl carbonyl amino", as used herein, refers to -NHR wherein R is a cycloalkyl carbonyl group.

The term "cycloalkyloxy", as used herein, refers to a cycloalkyl group attached to the parent molecular moiety through an oxygen atom.

The term "dialkylamino", as used herein, refers to NR₂, wherein each R is an alkyl group. The two alkyl groups are the same or different.

5 The term "haloalkoxy", as used herein, refers to a haloalkyl group attached to the parent molecular moiety through an oxygen atom.

The term "haloalkyl", as used herein, refers to an alkyl group substituted by one, two, three, or four halogen atoms.

10 The term "haloalkylamino", as used herein, refers to -NHR wherein R is a haloalkyl group.

The term "carbonyl" refers to C(=O).

The term "carboxy" refers to C(=O)OH.

The term "haloalkylcarbonyl", as used herein, refers to a haloalkyl group attached to the parent molecular moiety through a carbonyl group.

15 The term "haloalkylcarbonylamino", as used herein, refers to -NHR wherein R is a haloalkylcarbonyl group.

The terms "alkylcarbonyl" refer to an alkyl or substituted alkyl bonded to a carbonyl.

20 The term "alkoxycarbonyl", as used herein, refers to an alkoxy group attached to the parent molecular moiety through a carbonyl group.

The term "hydroxy" or "hydroxyl" refers to OH.

25 The term "cycloalkyl" refers to cyclized alkyl groups, including mono-, bi- or poly-cyclic ring systems. "C₃ to C₇ cycloalkyl" or "C₃₋₇ cycloalkyl" is intended to include C₃, C₄, C₅, C₆, and C₇ cycloalkyl groups. Example cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and norbornyl. Branched cycloalkyl groups such as 1-methylcyclopropyl and 2-methylcyclopropyl are included in the definition of "cycloalkyl".

30 As used herein, "carbocycle" or "carbocyclic residue" is intended to mean any stable 3-, 4-, 5-, 6-, 7-, or 8-membered monocyclic or bicyclic or 7-, 8-, 9-, 10-, 11-, 12-, or 13-membered bicyclic or tricyclic hydrocarbon ring, any of which may be saturated, partially unsaturated, unsaturated or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclobutenyl, cyclopentyl, cyclopentenyl,

cyclohexyl, cycloheptenyl, cycloheptyl, cycloheptenyl, adamantlyl, cyclooctyl, cyclooctenyl, cyclooctadienyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantlyl, anthracenyl, and tetrahydronaphthyl (tetralin). As shown above, bridged rings 5 are also included in the definition of carbocycle (e.g., [2.2.2]bicyclooctane). Preferred carbocycles, unless otherwise specified, are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, and indanyl. When the term "carbocycle" is used, it is intended to include "aryl". A bridged ring occurs when one or more carbon atoms link two non-adjacent carbon atoms. Preferred bridges are one or two carbon atoms. It is noted that a 10 bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge.

As used herein, the term "bicyclic carbocycle" or "bicyclic carbocyclic group" is intended to mean a stable 9- or 10-membered carbocyclic ring system that contains two fused rings and consists of carbon atoms. Of the two fused rings, one ring is a benzo ring 15 fused to a second ring; and the second ring is a 5- or 6-membered carbon ring which is saturated, partially unsaturated, or unsaturated. The bicyclic carbocyclic group may be attached to its pendant group at any carbon atom which results in a stable structure. The bicyclic carbocyclic group described herein may be substituted on any carbon if the resulting compound is stable. Examples of a bicyclic carbocyclic group are, but not 20 limited to, naphthyl, 1,2-dihydronaphthyl, 1,2,3,4-tetrahydronaphthyl, and indanyl.

"Aryl" groups refer to monocyclic or polycyclic aromatic hydrocarbons, including, for example, phenyl, naphthyl, and phenanthranyl. Aryl moieties are well known and described, for example, in Lewis, R.J., ed., *Hawley's Condensed Chemical Dictionary*, 13th Edition, John Wiley & Sons, Inc., New York (1997). "C₆ or C₁₀ aryl" or 25 "C₆₋₁₀ aryl" refers to phenyl and naphthyl. Unless otherwise specified, "aryl", "C₆ or C₁₀ aryl" or "C₆₋₁₀ aryl" or "aromatic residue" may be unsubstituted or substituted with 1 to 5 groups, preferably 1 to 3 groups, OH, OCH₃, Cl, F, Br, I, CN, NO₂, NH₂, N(CH₃)H, N(CH₃)₂, CF₃, OCF₃, C(=O)CH₃, SCH₃, S(=O)CH₃, S(=O)₂CH₃, CH₃, CH₂CH₃, CO₂H, and CO₂CH₃.

30 The term "benzyl", as used herein, refers to a methyl group on which one of the hydrogen atoms is replaced by a phenyl group, wherein said phenyl group may optionally be substituted with 1 to 5 groups, preferably 1 to 3 groups, OH, OCH₃, Cl, F, Br, I, CN,

NO₂, NH₂, N(CH₃)H, N(CH₃)₂, CF₃, OCF₃, C(=O)CH₃, SCH₃, S(=O)CH₃, S(=O)₂CH₃, CH₃, CH₂CH₃, CO₂H, and CO₂CH₃.

As used herein, the term "heterocycle" or "heterocyclic ring" is intended to mean a stable 3-, 4-, 5-, 6-, or 7-membered monocyclic or bicyclic or 7-, 8-, 9-, 10-, 11-, 12-, 13-, 5 or 14-membered polycyclic heterocyclic ring that is saturated, partially unsaturated, or fully unsaturated, and that contains carbon atoms and 1, 2, 3 or 4 heteroatoms independently selected from the group consisting of N, O and S; and including any polycyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized (*i.e.*, 10 N→O and S(O)_p, wherein p is 0, 1 or 2). The nitrogen atom may be substituted or unsubstituted (*i.e.*, N or NR wherein R is H or another substituent, if defined). The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. A 15 nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. When the term "heterocycle" is used, it is intended to include heteroaryl.

20 Examples of heterocycles include, but are not limited to, acridinyl, azetidinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzoxazolinyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolinyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-25 dithiazinyl, dihydrofuro[2,3-*b*]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolinyl, imidazolyl, 1*H*-indazolyl, imidazolopyridinyl, indolenyl, indoliny, indolizinyl, indolyl, 3*H*-indolyl, isatinoyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isothiazolopyridinyl, isoxazolyl, isoxazolopyridinyl, methylenedioxophenyl, morpholinyl, naphthyridinyl, 30 octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolopyridinyl, oxazolidinylperimidinyl, oxindolyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl,

phenazinyl, phenothiazinyl, phenoxythiinyl, phenoazinyl, phthalazinyl, piperazinyl, piperidinyl, piperidonyl, 4-piperidonyl, piperonyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolopyridinyl, pyrazolyl, pyridazinyl, pyridoazazolyl, pyridoimidazolyl, pyridothiazolyl, pyridinyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2-pyrrolidonyl, 2H-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl, 4H-quinolizinyl, quinoxaliny, quinuclidinyl, tetrazolyl, tetrahydrofuran, tetrahydroisoquinolinyl, tetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thiazolopyridinyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

Examples of 5- to 10-membered heterocycles include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrazinyl, piperazinyl, piperidinyl, imidazolyl, imidazolidinyl, indolyl, tetrazolyl, isoxazolyl, morpholinyl, oxazolyl, 15 oxadiazolyl, oxazolidinyl, tetrahydrofuran, thiadiazinyl, thiadiazolyl, thiazolyl, triazinyl, triazolyl, benzimidazolyl, 1H-indazolyl, benzofuran, benzothiofuran, benzotetrazolyl, benzotriazolyl, benzisoxazolyl, benzoxazolyl, oxindolyl, benzoxazolinyl, benzthiazolyl, benzisothiazolyl, isatinoyl, isoquinolinyl, octahydroisoquinolinyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, isoxazolopyridinyl, quinazolinyl, 20 quinolinyl, isothiazolopyridinyl, thiazolopyridinyl, oxazolopyridinyl, imidazolopyridinyl, and pyrazolopyridinyl.

Examples of 5- to 6-membered heterocycles include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrazinyl, piperazinyl, piperidinyl, imidazolyl, imidazolidinyl, indolyl, tetrazolyl, isoxazolyl, morpholinyl, oxazolyl, 25 oxadiazolyl, oxazolidinyl, tetrahydrofuran, thiadiazinyl, thiadiazolyl, thiazolyl, triazinyl, and triazolyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

As used herein, the term "bicyclic heterocycle" or "bicyclic heterocyclic group" is intended to mean a stable 9- or 10-membered heterocyclic ring system which contains 30 two fused rings and consists of carbon atoms and 1, 2, 3, or 4 heteroatoms independently selected from the group consisting of N, O and S. Of the two fused rings, one ring is a 5- or 6-membered monocyclic aromatic ring comprising a 5-membered heteroaryl ring, a 6-

membered heteroaryl ring or a benzo ring, each fused to a second ring. The second ring is a 5- or 6-membered monocyclic ring which is saturated, partially unsaturated, or unsaturated, and comprises a 5-membered heterocycle, a 6-membered heterocycle or a carbocycle (provided the first ring is not benzo when the second ring is a carbocycle).

5 The bicyclic heterocyclic group may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The bicyclic heterocyclic group described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is
10 preferred that the total number of S and O atoms in the heterocycle is not more than 1.

Examples of a bicyclic heterocyclic group are, but not limited to, quinoliny1, isoquinoliny1, phthalaziny1, quinazoliny1, indolyl, isoindolyl, indoliny1, 1H-indazolyl, benzimidazolyl, 1,2,3,4-tetrahydroquinoliny1, 1,2,3,4-tetrahydroisoquinoliny1, 5,6,7,8-tetrahydro-quinoliny1, 2,3-dihydro-benzofuranyl, chromanyl, 1,2,3,4-tetrahydro-quinoxaliny1, and 1,2,3,4-tetrahydro-quinazoliny1.

15 As used herein, the term "aromatic heterocyclic group" or "heteroaryl" is intended to mean stable monocyclic and polycyclic aromatic hydrocarbons that include at least one heteroatom ring member such as sulfur, oxygen, or nitrogen. Heteroaryl groups include, without limitation, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl, quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuryl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 1,2,4-thiadiazolyl, isothiazolyl, purinyl, carbazolyl, benzimidazolyl, indoliny1, benzodioxolanyl, and benzodioxane. Heteroaryl groups are substituted or unsubstituted. The nitrogen atom is substituted or unsubstituted (*i.e.*, N or NR wherein R is H or another
20 substituent, if defined). The nitrogen and sulfur heteroatoms may optionally be oxidized (*i.e.*, N→O and S(O)_p, wherein p is 0, 1 or 2).

25 Bridged rings are also included in the definition of heterocycle. A bridged ring occurs when one or more atoms (*i.e.*, C, O, N, or S) link two non-adjacent carbon or nitrogen atoms. Examples of bridged rings include, but are not limited to, one carbon atom, two carbon atoms, one nitrogen atom, two nitrogen atoms, and a carbon-nitrogen group. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring.

When a ring is bridged, the substituents recited for the ring may also be present on the bridge.

The term "counterion" is used to represent a negatively charged species such as chloride, bromide, hydroxide, acetate, and sulfate.

5 When a dotted ring is used within a ring structure, this indicates that the ring structure may be saturated, partially saturated or unsaturated.

As referred to herein, the term "substituted" means that at least one hydrogen atom is replaced with a non-hydrogen group, provided that normal valencies are maintained and that the substitution results in a stable compound. When a substituent is keto (*i.e.*, 10 =O), then 2 hydrogens on the atom are replaced. Keto substituents are not present on aromatic moieties. When a ring system (*e.g.*, carbocyclic or heterocyclic) is said to be substituted with a carbonyl group or a double bond, it is intended that the carbonyl group or double bond be part (*i.e.*, within) of the ring. Ring double bonds, as used herein, are double bonds that are formed between two adjacent ring atoms (*e.g.*, C=C, C=N, or 15 N=N).

In cases wherein there are nitrogen atoms (*e.g.*, amines) on compounds of the present invention, these may be converted to N-oxides by treatment with an oxidizing agent (*e.g.*, mCPBA and/or hydrogen peroxides) to afford other compounds of this invention. Thus, shown and claimed nitrogen atoms are considered to cover both the 20 shown nitrogen and its N-oxide (N→O) derivative.

When any variable occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-3 R groups, then said group may optionally be substituted with up to three R groups, and at each 25 occurrence R is selected independently from the definition of R. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is 30 listed without indicating the atom in which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such

substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms that are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, and/or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic groups such as amines, and alkali or organic salts of acidic groups such as carboxylic acids. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, and nitric, and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, and isethionic.

The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 18th Edition, Mack Publishing Company, Easton, PA (1990), the disclosure of which is hereby incorporated by reference.

In addition, compounds of formula I may have prodrug forms. Any compound that will be converted *in vivo* to provide the bioactive agent (*i.e.*, a compound of formula

I) is a prodrug within the scope and spirit of the invention. Various forms of prodrugs are well known in the art. For examples of such prodrug derivatives, see:

- a) Bundgaard, H., ed., *Design of Prodrugs*, Elsevier (1985), and Widder, K. et al., eds., *Methods in Enzymology*, 112:309-396, Academic Press (1985);
- 5 b) Bundgaard, H., Chapter 5, "Design and Application of Prodrugs", *A Textbook of Drug Design and Development*, pp. 113-191, Krosgaard-Larsen, P. et al., eds., Harwood Academic Publishers (1991);
- c) Bundgaard, H., *Adv. Drug Deliv. Rev.*, 8:1-38 (1992);
- d) Bundgaard, H. et al., *J. Pharm. Sci.*, 77:285 (1988); and
- 10 e) Kakeya, N. et al., *Chem. Pharm. Bull.*, 32:692 (1984).

Compounds containing a carboxy group can form physiologically hydrolyzable esters that serve as prodrugs by being hydrolyzed in the body to yield formula I compounds *per se*. Such prodrugs are preferably administered orally since hydrolysis in many instances occurs principally under the influence of the digestive enzymes.

15 Parenteral administration may be used where the ester *per se* is active, or in those instances where hydrolysis occurs in the blood. Examples of physiologically hydrolyzable esters of compounds of formula I include C₁₋₆alkyl, C₁₋₆alkylbenzyl, 4-methoxybenzyl, indanyl, phthalyl, methoxymethyl, C₁₋₆ alkanoyloxy-C₁₋₆alkyl (e.g., acetoxyethyl, pivaloyloxymethyl or propionyloxymethyl), C₁₋₆alkoxycarbonyloxy-C₁₋₆alkyl (e.g., methoxycarbonyl-oxymethyl or ethoxycarbonyloxymethyl, glycyloxymethyl, phenylglycyloxymethyl, (5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl), and other well known physiologically hydrolyzable esters used, for example, in the penicillin and cephalosporin arts. Such esters may be prepared by conventional techniques known in the art.

20 Preparation of prodrugs is well known in the art and described in, for example, King, F.D., ed., *Medicinal Chemistry: Principles and Practice*, The Royal Society of Chemistry, Cambridge, UK (1994); Testa, B. et al., *Hydrolysis in Drug and Prodrug Metabolism. Chemistry, Biochemistry and Enzymology*, VCHA and Wiley-VCH, Zurich, Switzerland (2003); Wermuth, C.G., ed., *The Practice of Medicinal Chemistry*, Academic Press, San Diego, CA (1999).

The present invention is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include deuterium and tritium. Isotopes of carbon include ¹³C and ¹⁴C.

5 Isotopically-labeled compounds of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein, using an appropriate isotopically-labeled reagent in place of the non-labeled reagent otherwise employed. Such compounds have a variety of potential uses, *e.g.*, as standards and reagents in determining the ability of a potential 10 pharmaceutical compound to bind to target proteins or receptors, or for imaging compounds of this invention bound to biological receptors *in vivo* or *in vitro*.

"Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent. It is preferred that 15 compounds of the present invention do not contain a N-halo, S(O)₂H, or S(O)H group.

The term "solvate" means a physical association of a compound of this invention with one or more solvent molecules, whether organic or inorganic. This physical association includes hydrogen bonding. In certain instances the solvate will be capable of isolation, for example, when one or more solvent molecules are incorporated in the 20 crystal lattice of the crystalline solid. The solvent molecules in the solvate may be present in a regular arrangement and/or a non-ordered arrangement. The solvate may comprise either a stoichiometric or nonstoichiometric amount of the solvent molecules.

"Solvate" encompasses both solution-phase and isolable solvates. Exemplary solvates include, but are not limited to, hydrates, ethanolates, methanolates, and isopropanolates.

25 Methods of solvation are generally known in the art.

Abbreviations as used herein, are defined as follows: "1 x" for once, "2 x" for twice, "3 x" for thrice, "^oC" for degrees Celsius, "eq" for equivalent or equivalents, "g" for gram or grams, "mg" for milligram or milligrams, "L" for liter or liters, "mL" for milliliter or milliliters, " μ L" for microliter or microliters, "N" for normal, "M" for molar, "mmol" 30 for millimole or millimoles, "min" for minute or minutes, "h" for hour or hours, "rt" for room temperature, "RT" for retention time, "RBF" for round bottom flask, "atm" for atmosphere, "psi" for pounds per square inch, "conc." for concentrate, "RCM" for ring-

closing metathesis, "sat" or "sat'd" for saturated, "SFC" for supercritical fluid chromatography, "MW" for molecular weight, "mp" for melting point, "ee" for enantiomeric excess, "MS" or "Mass Spec" for mass spectrometry, "ESI" for electrospray ionization mass spectroscopy, "HR" for high resolution, "HRMS" for high resolution mass spectrometry, "LCMS" for liquid chromatography mass spectrometry, "HPLC" for high pressure liquid chromatography, "RP HPLC" for reverse phase HPLC, "TLC" or "tlc" for thin layer chromatography, "NMR" for nuclear magnetic resonance spectroscopy, "nOe" for nuclear Overhauser effect spectroscopy, "¹H" for proton, " δ " for delta, "s" for singlet, "d" for doublet, "t" for triplet, "q" for quartet, "m" for multiplet, "br" for broad, "Hz" for hertz, and " α ", " β ", "R", "S", "E", and "Z" are stereochemical designations familiar to one skilled in the art.

Me	methyl
Et	ethyl
Pr	propyl
<i>i</i> -Pr	isopropyl
Bu	butyl
<i>i</i> -Bu	isobutyl
<i>t</i> -Bu	<i>tert</i> -butyl
Ph	phenyl
Bn	benzyl
Boc	<i>tert</i> -butyloxycarbonyl
Boc ₂ O	di- <i>tert</i> -butyl dicarbonate
AcOH or HOAc	acetic acid
AlCl ₃	aluminum chloride
AIBN	Azobisisobutyronitrile
BBr ₃	boron tribromide
BCl ₃	boron trichloride
BEMP	2- <i>tert</i> -butylimino-2-diethylamino-1,3-dimethylperhydro-1,3,2-diazaphosphorine
BOP reagent	benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate

Burgess reagent	1-methoxy-N-triethylammoniosulfonyl-methanimide
CBz	carbobenzyloxy
DCM or CH ₂ Cl ₂	dichloromethane
CH ₃ CN or ACN	acetonitrile
CDCl ₃	deutero-chloroform
CHCl ₃	chloroform
mCPBA or m-CPBA	<i>meta</i> -chloroperbenzoic acid
Cs ₂ CO ₃	cesium carbonate
Cu(OAc) ₂	copper (II) acetate
Cy ₂ NMe	N-cyclohexyl-N-methylcyclohexanamine
DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
DCE	1,2 dichloroethane
DEA	diethylamine
Dess-Martin	1,1,1-tris(acetyloxy)-1,1-dihydro-1,2-beniziodoxol-3-(1H)-one
DIC or DIPCDI	diisopropylcarbodiimide
DIEA, DIPEA or Hunig's base	diisopropylethylamine
DMAP	4-dimethylaminopyridine
DME	1,2-dimethoxyethane
DMF	dimethyl formamide
DMSO	dimethyl sulfoxide
cDNA	complimentary DNA
Dppp	(<i>R</i>)-(+)1,2-bis(diphenylphosphino)propane
DuPhos	(+)-1,2-bis((2S,5S)-2,5-diethylphospholano)benzene
EDC	<i>N</i> -(3-dimethylaminopropyl)- <i>N'</i> -ethylcarbodiimide
EDCI	<i>N</i> -(3-dimethylaminopropyl)- <i>N'</i> -ethylcarbodiimide hydrochloride
EDTA	ethylenediaminetetraacetic acid
(<i>S,S</i>)-EtDuPhosRh(I)	(+)-1,2-bis((2S,5S)-2,5-diethylphospholano)benzene(1,5-cyclooctadiene)rhodium(I) trifluoromethanesulfonate
Et ₃ N or TEA	triethylamine
EtOAc	ethyl acetate
Et ₂ O	diethyl ether

EtOH	ethanol
GMF	glass microfiber filter
Grubbs II	(1,3-bis(2,4,6-trimethylphenyl)-2-imidazolidinylidene)dichloro (phenylmethylene)(tricyclohexylphosphine)ruthenium
HCl	hydrochloric acid
HATU	O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
HEPES	4-(2-hydroxyethyl)piperazine-1-ethanesulfonic acid
Hex	hexane
HOBt or HOBT	1-hydroxybenzotriazole
IBX	2-iodoxybenzoic acid
H ₂ SO ₄	sulfuric acid
Jones reagent	CrO ₃ in aqueous H ₂ SO ₄ , 2 M
K ₂ CO ₃	potassium carbonate
K ₂ HPO ₄	potassium phosphate dibasic
KOAc	potassium acetate
K ₃ PO ₄	potassium phosphate
LAH	lithium aluminum hydride
LG	leaving group
LiOH	lithium hydroxide
MeOH	methanol
MgSO ₄	magnesium sulfate
MsOH or MSA	methylsulfonic acid
NaCl	sodium chloride
NaH	sodium hydride
NaHCO ₃	sodium bicarbonate
Na ₂ CO ₃	sodium carbonate
NaOH	sodium hydroxide
Na ₂ SO ₃	sodium sulfite
Na ₂ SO ₄	sodium sulfate
NBS	N-bromosuccinimide
NCS	N-chlorosuccinimide

NH ₃	ammonia
NH ₄ Cl	ammonium chloride
NH ₄ OH	ammonium hydroxide
NH ₄ COOH	ammonium formate
OTf	triflate or trifluoromethanesulfonate
Pd ₂ (dba) ₃	tris(dibenzylideneacetone)dipalladium(0)
Pd(OAc) ₂	palladium(II) acetate
Pd/C	palladium on carbon
Pd(dppf)Cl ₂	[1,1'-bis(diphenylphosphino)-ferrocene]dichloropalladium(II)
Ph ₃ PCl ₂	triphenylphosphine dichloride
PG	protecting group
POCl ₃	phosphorus oxychloride
i-PrOH or IPA	isopropanol
PS	Polystyrene
SEM-Cl	2-(trimethylsilyl)ethoxymethyl chloride
SiO ₂	silica oxide
SnCl ₂	tin(II) chloride
TBAI	tetra- <i>n</i> -butylammonium iodide
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TMSCHN ₂	trimethylsilyldiazomethane
T3P®	propane phosphonic acid anhydride
TRIS	tris (hydroxymethyl) aminomethane
pTsOH	p-toluenesulfonic acid

The compounds of the present invention can be prepared in a number of ways known to one skilled in the art of organic synthesis.

5 IV. BIOLOGY

While blood coagulation is essential to the regulation of an organism's hemostasis, it is also involved in many pathological conditions. In thrombosis, a blood clot, or thrombus, may form and obstruct circulation locally, causing ischemia and organ damage.

Alternatively, in a process known as embolism, the clot may dislodge and subsequently become trapped in a distal vessel, where it again causes ischemia and organ damage. Diseases arising from pathological thrombus formation are collectively referred to as thromboembolic disorders and include acute coronary syndrome, unstable angina, 5 myocardial infarction, thrombosis in the cavity of the heart, ischemic stroke, deep vein thrombosis, peripheral occlusive arterial disease, transient ischemic attack, and pulmonary embolism. In addition, thrombosis occurs on artificial surfaces in contact with blood, including catheters, stents, artificial heart valves, and hemodialysis membranes.

Some conditions contribute to the risk of developing thrombosis. For example, 10 alterations of the vessel wall, changes in the flow of blood, and alterations in the composition of the vascular compartment. These risk factors are collectively known as Virchow's triad. (Colman, R.W. et al., eds., *Hemostasis and Thrombosis, Basic Principles and Clinical Practice*, Fifth Edition, p. 853, Lippincott Williams & Wilkins (2006)).

Antithrombotic agents are frequently given to patients at risk of developing 15 thromboembolic disease because of the presence of one or more predisposing risk factors from Virchow's triad to prevent formation of an occlusive thrombus (primary prevention). For example, in an orthopedic surgery setting (e.g., hip and knee replacement), an antithrombotic agent is frequently administered prior to a surgical procedure. The antithrombotic agent counterbalances the prothrombotic stimulus exerted by vascular 20 flow alterations (stasis), potential surgical vessel wall injury, as well as changes in the composition of the blood due to the acute phase response related to surgery. Another example of the use of an antithrombotic agent for primary prevention is dosing with aspirin, a platelet activation inhibitor, in patients at risk for developing thrombotic cardiovascular disease. Well recognized risk factors in this setting include age, male 25 gender, hypertension, diabetes mellitus, lipid alterations, and obesity.

Antithrombotic agents are also indicated for secondary prevention, following an 30 initial thrombotic episode. For example, patients with mutations in Factor V (also known as Factor V Leiden) and additional risk factors (e.g., pregnancy), are dosed with anticoagulants to prevent the reoccurrence of venous thrombosis. Another example entails secondary prevention of cardiovascular events in patients with a history of acute myocardial infarction or acute coronary syndrome. In a clinical setting, a combination of

aspirin and clopidogrel (or other thienopyridines) may be used to prevent a second thrombotic event.

Antithrombotic agents are also given to treat the disease state (*i.e.*, by arresting its development) after it has already started. For example, patients presenting with deep vein thrombosis are treated with anticoagulants (*i.e.*, heparin, warfarin, or LMWH) to prevent further growth of the venous occlusion. Over time, these agents also cause a regression of the disease state because the balance between prothrombotic factors and anticoagulant/profibrinolytic pathways is changed in favor of the latter. Examples on the arterial vascular bed include the treatment of patients with acute myocardial infarction or acute coronary syndrome with aspirin and clopidogrel to prevent further growth of vascular occlusions and eventually leading to a regression of thrombotic occlusions.

Thus, antithrombotic agents are used widely for primary and secondary prevention (*i.e.*, prophylaxis or risk reduction) of thromboembolic disorders, as well as treatment of an already existing thrombotic process. Drugs that inhibit blood coagulation, or anticoagulants, are "pivotal agents for prevention and treatment of thromboembolic disorders" (Hirsh, J. et al., *Blood*, 105:453-463 (2005)).

An alternative way of initiation of coagulation is operative when blood is exposed to artificial surfaces (*e.g.*, during hemodialysis, "on-pump" cardiovascular surgery, vessel grafts, bacterial sepsis), on cell surfaces, cellular receptors, cell debris, DNA, RNA, and extracellular matrices. This process is also termed contact activation. Surface absorption of Factor XII leads to a conformational change in the Factor XII molecule, thereby facilitating activation to proteolytic active Factor XII molecules (Factor XIIa and Factor XIIif). Factor XIIa (or XIIif) has a number of target proteins, including plasma prekallikrein and Factor XI. Active plasma kallikrein further activates Factor XII, leading to an amplification of contact activation. Alternatively, the serine protease prolylcarboxylpeptidase can activate plasma kallikrein complexed with high molecular weight kininogen in a multiprotein complex formed on the surface of cells and matrices (Shariat-Madar et al., *Blood*, 108:192-199 (2006)). Contact activation is a surface mediated process responsible in part for the regulation of thrombosis and inflammation, and is mediated, at least in part, by fibrinolytic-, complement-, kininogen/kinin-, and other humoral and cellular pathways (for review, Coleman, R., "Contact Activation Pathway", *Hemostasis and Thrombosis*, pp. 103-122, Lippincott Williams & Wilkins

(2001); Schmaier, A.H., "Contact Activation", *Thrombosis and Hemorrhage*, pp. 105-128 (1998)). The biological relevance of the contact activation system for thromboembolic diseases is supported by the phenotype of Factor XII deficient mice. More specifically, Factor XII deficient mice were protected from thrombotic vascular occlusion in several 5 thrombosis models as well as stroke models and the phenotype of the XII deficient mice was identical to XI deficient mice (Renne et al., *J. Exp. Med.*, 202:271-281 (2005); Kleinschmitz et al., *J. Exp. Med.*, 203:513-518 (2006)). The fact that Factor XI is downstream from Factor XIIa, combined with the identical phenotype of the XII and XI deficient mice suggest that the contact activation system could play a major role in Factor 10 XI activation *in vivo*.

Factor XI is a zymogen of a trypsin-like serine protease and is present in plasma at a relatively low concentration. Proteolytic activation at an internal R369-I370 bond yields a heavy chain (369 amino acids) and a light chain (238 amino acids). The latter contains a typical trypsin-like catalytic triad (H413, D464, and S557). Activation of 15 Factor XI by thrombin is believed to occur on negatively charged surfaces, most likely on the surface of activated platelets. Platelets contain high affinity (0.8 nM) specific sites (130-500/platelet) for activated Factor XI. After activation, Factor XIa remains surface bound and recognizes Factor IX as its normal macromolecular substrate. (Galiani, D., *Trends Cardiovasc. Med.*, 10:198-204 (2000)).

20 In addition to the feedback activation mechanisms described above, thrombin activates thrombin activated fibrinolysis inhibitor (TAFI), a plasma carboxypeptidase that cleaves C-terminal lysine and arginine residues on fibrin, reducing the ability of fibrin to enhance tissue-type plasminogen activator (tPA) dependent plasminogen activation. In the presence of antibodies to FXIa, clot lysis can occur more rapidly independent of 25 plasma TAFI concentration. (Bouma, B.N. et al., *Thromb. Res.*, 101:329-354 (2001).) Thus, inhibitors of Factor XIa are expected to be anticoagulant and profibrinolytic.

Further evidence for the anti-thromboembolic effects of targeting Factor XI is derived from mice deficient in Factor XI. It has been demonstrated that complete fXI deficiency protected mice from ferric chloride (FeCl₃)-induced carotid artery thrombosis 30 (Rosen et al., *Thromb. Haemost.*, 87:774-777 (2002); Wang et al., *J. Thromb. Haemost.*, 3:695-702 (2005)). Also, Factor XI deficiency rescues the perinatal lethal phenotype of complete protein C deficiency (Chan et al., *Amer. J. Pathol.*, 158:469-479 (2001)).

Furthermore, baboon cross-reactive, function blocking antibodies to human Factor XI protect against baboon arterial - venous shunt thrombosis (Gruber et al., *Blood*, 102:953-955 (2003)). Evidence for an antithrombotic effect of small molecule inhibitors of Factor XIa is also disclosed in published U.S. Publication No. 2004/0180855 A1. Taken 5 together, these studies suggest that targeting Factor XI will reduce the propensity for thrombotic and thromboembolic diseases.

Genetic evidence indicates that Factor XI is not required for normal homeostasis, implying a superior safety profile of the Factor XI mechanism compared to competing antithrombotic mechanisms. In contrast to hemophilia A (Factor VIII deficiency) or 10 hemophilia B (Factor IX deficiency), mutations of the Factor XI gene causing Factor XI deficiency (hemophilia C) result in only a mild to moderate bleeding diathesis characterized primarily by postoperative or posttraumatic, but rarely spontaneous hemorrhage. Postoperative bleeding occurs mostly in tissue with high concentrations of endogenous fibrinolytic activity (e.g., oral cavity, and urogenital system). The majority 15 of the cases are fortuitously identified by preoperative prolongation of aPTT (intrinsic system) without any prior bleeding history.

The increased safety of inhibition of XIa as an anticoagulation therapy is further supported by the fact that Factor XI knock-out mice, which have no detectable Factor XI protein, undergo normal development, and have a normal life span. No evidence for 20 spontaneous bleeding has been noted. The aPTT (intrinsic system) is prolonged in a gene dose-dependent fashion. Interestingly, even after severe stimulation of the coagulation system (tail transection), the bleeding time is not significantly prolonged compared to wild-type and heterozygous litter mates. (Gailani, D., *Frontiers in Bioscience*, 6:201-207 (2001); Gailani, D. et al., *Blood Coagulation and Fibrinolysis*, 8:134-144 (1997).) Taken 25 together, these observations suggest that high levels of inhibition of Factor XIa should be well tolerated. This is in contrast to gene targeting experiments with other coagulation factors, excluding Factor XII.

In vivo activation of Factor XI can be determined by complex formation with either C1 inhibitor or alpha 1 antitrypsin. In a study of 50 patients with acute myocardial 30 infarction (AMI), approximately 25% of the patients had values above the upper normal range of the complex ELISA. This study can be viewed as evidence that at least in a subpopulation of patients with AMI, Factor XI activation contributes to thrombin

formation (Minnema, M.C. et al., *Arterioscler. Thromb. Vasc. Biol.*, 20:2489-2493 (2000)). A second study establishes a positive correlation between the extent of coronary arteriosclerosis and Factor XIa in complex with alpha 1 antitrypsin (Murakami, T. et al., *Arterioscler. Thromb. Vasc. Biol.*, 15:1107-1113 (1995)). In another study, Factor XI 5 levels above the 90th percentile in patients were associated with a 2.2-fold increased risk for venous thrombosis (Meijers, J.C.M. et al., *N. Engl. J. Med.*, 342:696-701 (2000)).

Also, it is preferred to find new compounds with improved activity in *in vitro* clotting assays, compared with known serine protease inhibitors, such as the activated partial thromboplastin time (aPTT) or prothrombin time (PT) assay. (for a description of 10 the aPTT and PT assays see, Goodnight, S.H. et al., "Screening Tests of Hemostasis", *Disorders of Thrombosis and Hemostasis: A Clinical Guide*, Second Edition, pp. 41-51, McGraw-Hill, New York (2001)).

It is also desirable and preferable to find compounds with advantageous and 15 improved characteristics compared with known serine protease inhibitors, in one or more of the following categories that are given as examples, and are not intended to be limiting: (a) pharmacokinetic properties, including oral bioavailability, half life, and clearance; (b) pharmaceutical properties; (c) dosage requirements; (d) factors that decrease blood concentration peak-to-trough characteristics; (e) factors that increase the concentration of active drug at the receptor; (f) factors that decrease the liability for clinical drug-drug 20 interactions; (g) factors that decrease the potential for adverse side-effects, including selectivity versus other biological targets; and (h) factors that improve manufacturing costs or feasibility.

Pre-clinical studies demonstrated significant antithrombotic effects of small 25 molecule Factor XIa inhibitors in rabbit and rat model of arterial thrombosis, at doses that preserved hemostasis. (Wong P.C. et al., *American Heart Association Scientific Sessions*, Abstract No. 6118, November 12-15, 2006; Schumacher, W. et al., *J. Thromb. Haemost.*, 3(Suppl. 1):P1228 (2005); Schumacher, W.A. et al., *Eur. J. Pharmacol.*, 167-174 (2007)). Furthermore, it was observed that *in vitro* prolongation of the aPTT by specific XIa 30 inhibitors is a good predictor of efficacy in our thrombosis models. Thus, the *in vitro* aPTT test can be used as a surrogate for efficacy *in vivo*.

As used herein, the term "patient" encompasses all mammalian species.

As used herein, "treating" or "treatment" cover the treatment of a disease-state in a mammal, particularly in a human, and include: (a) inhibiting the disease-state, *i.e.*, arresting its development; and/or (b) relieving the disease-state, *i.e.*, causing regression of the disease state.

5 As used herein, "prophylaxis" or "prevention" cover the preventive treatment of a subclinical disease-state in a mammal, particularly in a human, aimed at reducing the probability of the occurrence of a clinical disease-state. Patients are selected for preventative therapy based on factors that are known to increase risk of suffering a clinical disease state compared to the general population. "Prophylaxis" therapies can be
10 divided into (a) primary prevention and (b) secondary prevention. Primary prevention is defined as treatment in a subject that has not yet presented with a clinical disease state, whereas secondary prevention is defined as preventing a second occurrence of the same or similar clinical disease state.

15 As used herein, "risk reduction" covers therapies that lower the incidence of development of a clinical disease state. As such, primary and secondary prevention therapies are examples of risk reduction.

20 "Therapeutically effective amount" is intended to include an amount of a compound of the present invention that is effective when administered alone or in combination to inhibit Factor XIa and/or plasma kallikrein and/or to prevent or treat the disorders listed herein. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the preventive or therapeutic effect, whether administered in combination, serially, or simultaneously.

25 The term "thrombosis", as used herein, refers to formation or presence of a thrombus (pl. thrombi); clotting within a blood vessel that may cause ischemia or infarction of tissues supplied by the vessel. The term "embolism", as used herein, refers to sudden blocking of an artery by a clot or foreign material that has been brought to its site of lodgment by the blood current. The term "thromboembolism", as used herein, refers to obstruction of a blood vessel with thrombotic material carried by the blood stream from the site of origin to plug another vessel. The term "thromboembolic
30 disorders" entails both "thrombotic" and "embolic" disorders (defined above).

The term "thromboembolic disorders" as used herein includes arterial cardiovascular thromboembolic disorders, venous cardiovascular or cerebrovascular

thromboembolic disorders, and thromboembolic disorders in the chambers of the heart or in the peripheral circulation. The term "thromboembolic disorders" as used herein also includes specific disorders selected from, but not limited to, unstable angina or other acute coronary syndromes, atrial fibrillation, first or recurrent myocardial infarction, 5 ischemic sudden death, transient ischemic attack, stroke, atherosclerosis, peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an artificial surface 10 that promotes thrombosis. The medical implants or devices include, but are not limited to: prosthetic valves, artificial valves, indwelling catheters, stents, blood oxygenators, shunts, vascular access ports, ventricular assist devices and artificial hearts or heart chambers, and vessel grafts. The procedures include, but are not limited to: cardiopulmonary bypass, percutaneous coronary intervention, and hemodialysis. In 15 another embodiment, the term "thromboembolic disorders" includes acute coronary syndrome, stroke, deep vein thrombosis, and pulmonary embolism.

In another embodiment, the present invention provides a method for the treatment of a thromboembolic disorder, wherein the thromboembolic disorder is selected from unstable angina, an acute coronary syndrome, atrial fibrillation, myocardial infarction, 20 transient ischemic attack, stroke, atherosclerosis, peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an artificial surface that promotes thrombosis. In 25 another embodiment, the present invention provides a method for the treatment of a thromboembolic disorder, wherein the thromboembolic disorder is selected from acute coronary syndrome, stroke, venous thrombosis, atrial fibrillation, and thrombosis resulting from medical implants and devices.

In another embodiment, the present invention provides a method for the primary prophylaxis of a thromboembolic disorder, wherein the thromboembolic disorder is 30 selected from unstable angina, an acute coronary syndrome, atrial fibrillation, myocardial infarction, ischemic sudden death, transient ischemic attack, stroke, atherosclerosis,

5 peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an
10 artificial surface that promotes thrombosis. In another embodiment, the present invention provides a method for the primary prophylaxis of a thromboembolic disorder, wherein the thromboembolic disorder is selected from acute coronary syndrome, stroke, venous thrombosis, and thrombosis resulting from medical implants and devices.

15 In another embodiment, the present invention provides a method for the secondary prophylaxis of a thromboembolic disorder, wherein the thromboembolic disorder is selected from unstable angina, an acute coronary syndrome, atrial fibrillation, recurrent myocardial infarction, transient ischemic attack, stroke, atherosclerosis, peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an artificial surface that promotes thrombosis. In another embodiment, the present invention provides a
20 method for the secondary prophylaxis of a thromboembolic disorder, wherein the thromboembolic disorder is selected from acute coronary syndrome, stroke, atrial fibrillation and venous thrombosis.

The term "stroke", as used herein, refers to embolic stroke or atherothrombotic stroke arising from occlusive thrombosis in the carotid communis, carotid interna, or intracerebral arteries.

25 It is noted that thrombosis includes vessel occlusion (*e.g.*, after a bypass) and reocclusion (*e.g.*, during or after percutaneous transluminal coronary angioplasty). The thromboembolic disorders may result from conditions including but not limited to atherosclerosis, surgery or surgical complications, prolonged immobilization, arterial fibrillation, congenital thrombophilia, cancer, diabetes, effects of medications or hormones, and complications of pregnancy.

30 Thromboembolic disorders are frequently associated with patients with atherosclerosis. Risk factors for atherosclerosis include but are not limited to male gender, age, hypertension, lipid disorders, and diabetes mellitus. Risk factors for

atherosclerosis are at the same time risk factors for complications of atherosclerosis, *i.e.*, thromboembolic disorders.

Similarly, arterial fibrillation is frequently associated with thromboembolic disorders. Risk factors for arterial fibrillation and subsequent thromboembolic disorders 5 include cardiovascular disease, rheumatic heart disease, nonrheumatic mitral valve disease, hypertensive cardiovascular disease, chronic lung disease, and a variety of miscellaneous cardiac abnormalities as well as thyrotoxicosis.

Diabetes mellitus is frequently associated with atherosclerosis and thromboembolic disorders. Risk factors for the more common type 2 include but are not 10 limited to are family history, obesity, physical inactivity, race/ethnicity, previously impaired fasting glucose or glucose tolerance test, history of gestational diabetes mellitus or delivery of a "big baby", hypertension, low HDL cholesterol, and polycystic ovary syndrome.

Risk factors for congenital thrombophilia include gain of function mutations in 15 coagulation factors or loss of function mutations in the anticoagulant- or fibrinolytic pathways.

Thrombosis has been associated with a variety of tumor types, *e.g.*, pancreatic cancer, breast cancer, brain tumors, lung cancer, ovarian cancer, prostate cancer, gastrointestinal malignancies, and Hodgkins or non-Hodgkins lymphoma. Recent studies 20 suggest that the frequency of cancer in patients with thrombosis reflects the frequency of a particular cancer type in the general population (Levitin, N. et al., *Medicine* (Baltimore), 78(5):285-291 (1999); Levine M. et al., *N. Engl. J. Med.*, 334(11):677-681 (1996); Blom, J.W. et al., *JAMA*, 293(6):715-722 (2005)). Hence, the most common cancers associated with thrombosis in men are prostate, colorectal, brain, and lung cancer, 25 and in women are breast, ovary, and lung cancer. The observed rate of venous thromboembolism (VTE) in cancer patients is significant. The varying rates of VTE between different tumor types are most likely related to the selection of the patient population. Cancer patients at risk for thrombosis may possess any or all of the following risk factors: (i) the stage of the cancer (*i.e.*, presence of metastases), (ii) the presence of 30 central vein catheters, (iii) surgery and anticancer therapies including chemotherapy, and (iv) hormones and antiangiogenic drugs. Thus, it is common clinical practice to dose patients having advanced tumors with heparin or low molecular heparin to prevent

thromboembolic disorders. A number of low molecular heparin preparations have been approved by the FDA for these indications.

There are three main clinical situations when considering the prevention of VTE in a medical cancer patient: (i) the patient is bedridden for prolonged periods of time; (ii) 5 the ambulatory patient is receiving chemotherapy or radiation; and (iii) the patient is with indwelling central vein catheters. Unfractionated heparin (UFH) and low molecular weight heparin (LMWH) are effective antithrombotic agents in cancer patients undergoing surgery. (Mismetti, P. et al., *Brit. J. Surg.*, 88:913-930 (2001).)

10 A. *In Vitro* Assays

The effectiveness of compounds of the present invention as inhibitors of the coagulation Factors XIa, VIIa, IXa, Xa, XIIa, plasma kallikrein or thrombin, can be determined using a relevant purified serine protease, respectively, and an appropriate synthetic substrate. The rate of hydrolysis of the chromogenic or fluorogenic substrate by 15 the relevant serine protease was measured both in the absence and presence of compounds of the present invention. Hydrolysis of the substrate resulted in the release of pNA (para nitroaniline), which was monitored spectrophotometrically by measuring the increase in absorbance at 405 nm, or the release of AMC (amino methylcoumarin), which was monitored spectrofluorometrically by measuring the increase in emission at 460 nm 20 with excitation at 380 nm. A decrease in the rate of absorbance or fluorescence change in the presence of inhibitor is indicative of enzyme inhibition. Such methods are known to one skilled in the art. The results of this assay are expressed as the inhibitory constant, K_i .

Factor XIa determinations were made in 50 mM HEPES buffer at pH 7.4 containing 145 mM NaCl, 5 mM KCl, and 0.1% PEG 8000 (polyethylene glycol; JT 25 Baker or Fisher Scientific). Determinations were made using purified human Factor XIa at a final concentration of 25-200 pM (Haematologic Technologies) and the synthetic substrate S-2366 (pyroGlu-Pro-Arg-pNA; CHROMOGENIX® or AnaSpec) at a concentration of 0.0002-0.001 M.

Factor VIIa determinations were made in 0.005 M calcium chloride, 0.15 M 30 sodium chloride, 0.05 M HEPES buffer containing 0.1% PEG 8000 at a pH of 7.5. Determinations were made using purified human Factor VIIa (Haematologic Technologies) or recombinant human Factor VIIa (Novo Nordisk) at a final assay

concentration of 0.5-10 nM, recombinant soluble tissue factor at a concentration of 10-40 nM and the synthetic substrate H-D-Ile-Pro-Arg-pNA (S-2288; CHROMOGENIX® or BMPM-2; AnaSpec) at a concentration of 0.001-0.0075 M.

Factor IXa determinations were made in 0.005 M calcium chloride, 0.1 M sodium chloride, 0.0000001 M Refludan (Berlex), 0.05 M TRIS base and 0.5% PEG 8000 at a pH of 7.4. Refludan was added to inhibit small amounts of thrombin in the commercial preparations of human Factor IXa. Determinations were made using purified human Factor IXa (Haematologic Technologies) at a final assay concentration of 20-100 nM and the synthetic substrate PCIXA2100-B (CenterChem) or Pefafluor IXa 3688 (H-D-Leu-10 Ph'Gly-Arg-AMC; CenterChem) at a concentration of 0.0004-0.0005 M.

Factor Xa determinations were made in 0.1 M sodium phosphate buffer at a pH of 7.5 containing 0.2 M sodium chloride and 0.5% PEG 8000. Determinations were made using purified human Factor Xa (Haematologic Technologies) at a final assay concentration of 150-1000 pM and the synthetic substrate S-2222 (Bz-Ile-Glu (gamma-15 OMe, 50%)-Gly-Arg-pNA; CHROMOGENIX®) at a concentration of 0.0002-0.00035 M.

Factor XIIa determinations were made in 0.05 M HEPES buffer at pH 7.4 containing 0.145 M NaCl, 0.05 M KCl, and 0.1% PEG 8000. Determinations were made using purified human Factor XIIa at a final concentration of 4 nM (American 20 Diagnostica) and the synthetic substrate SPECTROZYME® #312 (H-D-CHT-Gly-L-Arg-pNA.2AcOH; American Diagnostica) at a concentration of 0.00015 M.

Plasma kallikrein determinations were made in 0.1 M sodium phosphate buffer at a pH of 7.5 containing 0.1-0.2 M sodium chloride and 0.5% PEG 8000. Determinations were made using purified human plasma kallikrein (Enzyme Research Laboratories) at a 25 final assay concentration of 200 pM and the synthetic substrate S-2302 (H-(D)-Pro-Phe-Arg-pNA; CHROMOGENIX®) at a concentration of 0.00008-0.0004 M.

Thrombin determinations were made in 0.1 M sodium phosphate buffer at a pH of 7.5 containing 0.2 M sodium chloride and 0.5% PEG 8000. Determinations were made using purified human alpha thrombin (Haematologic Technologies or Enzyme Research 30 Laboratories) at a final assay concentration of 200-250 pM and the synthetic substrate S-2366 (pyroGlu-Pro-Arg-pNA; CHROMOGENIX® or AnaSpec) at a concentration of 0.0002-0.0004 M.

The Michaelis constant, K_m , for substrate hydrolysis by each protease, was determined at 25 °C or 37 °C in the absence of inhibitor. Values of K_i were determined by allowing the protease to react with the substrate in the presence of the inhibitor. Reactions were allowed to go for periods of 20-180 minutes (depending on the protease) and the velocities (rate of absorbance or fluorescence change versus time) were measured.

5 The following relationships were used to calculate K_i values:

$$(V_{max} \cdot S) / (K_m + S);$$

$$(v_0 - v_s) / v_s = I / (K_i (1 + S / K_m)) \text{ for a competitive inhibitor with one binding site; or}$$

$$v_s / v_0 = A + (B - A) / (1 + (I / IC_{50})^n); \text{ and}$$

$$10 K_i = IC_{50} / (1 + S / K_m) \text{ for a competitive inhibitor}$$

where:

v_0 is the velocity of the control in the absence of inhibitor;

v_s is the velocity in the presence of inhibitor;

V_{max} is the maximum reaction velocity;

15 I is the concentration of inhibitor;

A is the minimum activity remaining (usually locked at zero);

B is the maximum activity remaining (usually locked at 1.0);

n is the Hill coefficient, a measure of the number and cooperativity of potential inhibitor binding sites;

20 IC_{50} is the concentration of inhibitor that produces 50% inhibition under the assay conditions;

K_i is the dissociation constant of the enzyme: inhibitor complex;

S is the concentration of substrate; and

K_m is the Michaelis constant for the substrate.

25

The selectivity of a compound may be evaluated by taking the ratio of the K_i value for a given protease with the K_i value for the protease of interest (*i.e.*, selectivity for FXIa versus protease P = K_i for protease P / K_i for FXIa). Compounds with selectivity ratios >20 are considered selective.

30

The effectiveness of compounds of the present invention as inhibitors of coagulation can be determined using a standard or modified clotting assay. An increase in the plasma clotting time in the presence of inhibitor is indicative of anticoagulation.

Relative clotting time is the clotting time in the presence of an inhibitor divided by the clotting time in the absence of an inhibitor. The results of this assay may be expressed as IC1.5x or IC2x, the inhibitor concentration required to increase the clotting time by 50 or 100 percent, respectively. The IC1.5x or IC2x is found by linear interpolation from 5 relative clotting time versus inhibitor concentration plots using inhibitor concentration that spans the IC1.5x or IC2x.

Clotting times are determined using citrated normal human plasma as well as plasma obtained from a number of laboratory animal species (e.g., rat, or rabbit). A compound is diluted into plasma beginning with a 10 mM DMSO stock solution. The 10 final concentration of DMSO is less than 2%. Plasma clotting assays are performed in an automated coagulation analyzer (SYSMEX®, Dade-Behring, Illinois). Similarly, clotting times can be determined from laboratory animal species or humans dosed with compounds of the invention.

Activated Partial Thromboplastin Time (aPTT) is determined using ACTIN® FSL 15 (Dade-Behring, Illinois) following the directions in the package insert. Plasma (0.05 mL) is warmed to 37 °C for 1 minute. ACTIN® FSL (0.05 mL) is added to the plasma and incubated for an additional 2 to 5 minutes. Calcium chloride (25 mM, 0.05 mL) is added to the reaction to initiate coagulation. The clotting time is the time in seconds from the moment calcium chloride is added until a clot is detected.

20 Prothrombin Time (PT) is determined using thromboplastin (Thromboplastin C Plus or INNOVIN®, Dade-Behring, Illinois) following the directions in the package insert. Plasma (0.05 mL) is warmed to 37 °C for 1 minute. Thromboplastin (0.1 mL) is added to the plasma to initiate coagulation. The clotting time is the time in seconds from the moment thromboplastin is added until a clot is detected.

25 The exemplified Examples disclosed below were tested in the Factor XIa assay described above and found having Factor XIa inhibitory activity. A range of Factor XIa inhibitory activity (Ki values) of $\leq 10 \mu\text{M}$ (10000 nM) was observed. Table 1 below lists Factor XIa Ki values measured at 37 °C for the following Examples.

30

V. PHARMACEUTICAL COMPOSITIONS, FORMULATIONS AND COMBINATIONS

The compounds of this invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. They may also be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or 5 intramuscular form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. They can be administered alone, but generally will be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The term "pharmaceutical composition" means a composition comprising a 10 compound of the invention in combination with at least one additional pharmaceutically acceptable carrier. A "pharmaceutically acceptable carrier" refers to media generally accepted in the art for the delivery of biologically active agents to animals, in particular, mammals, including, *i.e.*, adjuvant, excipient or vehicle, such as diluents, preserving agents, fillers, flow regulating agents, disintegrating agents, wetting agents, emulsifying 15 agents, suspending agents, sweetening agents, flavoring agents, perfuming agents, antibacterial agents, antifungal agents, lubricating agents and dispensing agents, depending on the nature of the mode of administration and dosage forms.

Pharmaceutically acceptable carriers are formulated according to a number of factors well 20 within the purview of those of ordinary skill in the art. These include, without limitation: the type and nature of the active agent being formulated; the subject to which the agent-containing composition is to be administered; the intended route of administration of the composition; and the therapeutic indication being targeted. Pharmaceutically acceptable carriers include both aqueous and non-aqueous liquid media, as well as a variety of solid and semi-solid dosage forms. Such carriers can include a number of different ingredients 25 and additives in addition to the active agent, such additional ingredients being included in the formulation for a variety of reasons, *e.g.*, stabilization of the active agent, binders, etc., well known to those of ordinary skill in the art. Descriptions of suitable pharmaceutically acceptable carriers, and factors involved in their selection, are found in a variety of readily available sources such as, for example, *Remington's Pharmaceutical Sciences*, 18th Edition (1990).

The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the

particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired. A physician or 5 veterinarian can determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the thromboembolic disorder.

By way of general guidance, the daily oral dosage of each active ingredient, when used for the indicated effects, will range between about 0.001 to about 1000 mg/kg of body weight, preferably between about 0.01 to about 100 mg/kg of body weight per day, 10 and most preferably between about 0.1 to about 20 mg/kg/day. Intravenously, the most preferred doses will range from about 0.001 to about 10 mg/kg/minute during a constant rate infusion. Compounds of this invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

15 Compounds of this invention can also be administered by parenteral administration (*e.g.*, intra-venous, intra-arterial, intramuscularly, or subcutaneously). When administered intra-venous or intra-arterial, the dose can be given continuously or intermittent. Furthermore, formulation can be developed for intramuscularly and subcutaneous delivery that ensure a gradual release of the active pharmaceutical 20 ingredient. In one embodiment, the pharmaceutical composition is a solid formulation, *e.g.*, a spray-dried composition, which may be used as is, or whereto the physician or the patient adds solvents, and/or diluents prior to use.

Compounds of this invention can be administered in intranasal form via topical 25 use of suitable intranasal vehicles, or via transdermal routes, using transdermal skin patches. When administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

The compounds are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as 30 pharmaceutical carriers) suitably selected with respect to the intended form of administration, *e.g.*, oral tablets, capsules, elixirs, and syrups, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral 5 administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, 10 natural and synthetic gums such as acacia, tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

15 The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

Compounds of the present invention may also be coupled with soluble polymers 20 as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for 25 example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacylates, and crosslinked or amphipathic block copolymers of hydrogels. Solid dispersions are also called solid-state dispersions. In some embodiments, any compound described herein is formulated as a spray dried dispersion 30 (SDD). An SDD is a single phase amorphous molecular dispersion of a drug in a polymer matrix. It is a solid solution prepared by dissolving the drug and a polymer in a solvent (e.g., acetone, methanol or the like) and spray drying the solution. The solvent

rapidly evaporates from droplets which rapidly solidifies the polymer and drug mixture trapping the drug in amorphous form as an amorphous molecular dispersion.

5 Dosage forms (pharmaceutical compositions) suitable for administration may contain from about 1 milligram to about 1000 milligrams of active ingredient per dosage unit. In these pharmaceutical compositions the active ingredient will ordinarily be present in an amount of about 0.1-95% by weight based on the total weight of the composition.

Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like.

10 Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

15 Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably 20 contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben, and 25 chlorobutanol.

Suitable pharmaceutical carriers are described in *Remington's Pharmaceutical Sciences*, Mack Publishing Company, a standard reference text in this field.

Where the compounds of this invention are combined with other anticoagulant agents, for example, a daily dosage may be about 0.1 to about 100 milligrams of the 30 compound of the present invention and about 0.1 to about 100 milligrams per kilogram of patient body weight. For a tablet dosage form, the compounds of this invention generally

may be present in an amount of about 5 to about 300 milligrams per dosage unit, and the second anti-coagulant in an amount of about 1 to about 500 milligrams per dosage unit.

Where the compounds of the present invention are administered in combination with an anti-platelet agent, by way of general guidance, typically a daily dosage may be 5 about 0.01 to about 300 milligrams of the compound of the present invention and about 50 to about 150 milligrams of the anti-platelet agent, preferably about 0.1 to about 4 milligrams of the compound of the present invention and about 1 to about 3 milligrams of antiplatelet agents, per kilogram of patient body weight.

Where the compounds of the present invention are administered in combination 10 with thrombolytic agent, typically a daily dosage may be about 0.1 to about 100 milligrams of the compound of the present invention, per kilogram of patient body weight and, in the case of the thrombolytic agents, the usual dosage of the thrombolytic agent when administered alone may be reduced by about 50-80% when administered with a compound of the present invention.

15 Particularly when provided as a single dosage unit, the potential exists for a chemical interaction between the combined active ingredients. For this reason, when the compound of the present invention and a second therapeutic agent are combined in a single dosage unit they are formulated such that although the active ingredients are combined in a single dosage unit, the physical contact between the active ingredients is minimized (that is, reduced). For example, one active ingredient may be enteric coated. By enteric coating one of the active ingredients, it is possible not only to minimize the contact between the combined active ingredients, but also, it is possible to control the release of one of these components in the gastrointestinal tract such that one of these components is not released in the stomach but rather is released in the intestines. One of 20 the active ingredients may also be coated with a material that affects a sustained-release throughout the gastrointestinal tract and also serves to minimize physical contact between the combined active ingredients. Furthermore, the sustained-released component can be additionally enteric coated such that the release of this component occurs only in the intestine. Still another approach would involve the formulation of a combination product 25 in which the one component is coated with a sustained and/or enteric release polymer, and the other component is also coated with a polymer such as a low viscosity grade of hydroxypropyl methylcellulose (HPMC) or other appropriate materials as known in the 30

art, in order to further separate the active components. The polymer coating serves to form an additional barrier to interaction with the other component.

These as well as other ways of minimizing contact between the components of combination products of the present invention, whether administered in a single dosage form or administered in separate forms but at the same time by the same manner, will be readily apparent to those skilled in the art, once armed with the present disclosure.

In another embodiment, the present invention provides a pharmaceutical composition further comprising additional therapeutic agent(s) selected from potassium channel openers, potassium channel blockers, calcium channel blockers, sodium

10 hydrogen exchanger inhibitors, antiarrhythmic agents, antiatherosclerotic agents, anticoagulants, antithrombotic agents, prothrombolytic agents, fibrinogen antagonists, diuretics, antihypertensive agents, ATPase inhibitors, mineralocorticoid receptor antagonists, phosphodiesterase inhibitors, antidiabetic agents, anti-inflammatory agents, antioxidants, angiogenesis modulators, antiosteoporosis agents, hormone replacement

15 therapies, hormone receptor modulators, oral contraceptives, antiobesity agents, antidepressants, antianxiety agents, antipsychotic agents, antiproliferative agents, antitumor agents, antiulcer and gastroesophageal reflux disease agents, growth hormone agents and/or growth hormone secretagogues, thyroid mimetics, anti-infective agents, antiviral agents, antibacterial agents, antifungal agents, cholesterol/lipid lowering agents

20 and lipid profile therapies, and agents that mimic ischemic preconditioning and/or myocardial stunning, or a combination thereof.

In another embodiment, the present invention provides a pharmaceutical composition further comprising additional therapeutic agent(s) selected from an anti-arrhythmic agent, an anti-hypertensive agent, an anti-coagulant agent, an anti-platelet agent, a thrombin inhibiting agent, a thrombolytic agent, a fibrinolytic agent, a calcium channel blocker, a potassium channel blocker, a cholesterol/lipid lowering agent, or a combination thereof.

In another embodiment, the present invention provides a pharmaceutical composition further comprising additional therapeutic agent(s) selected from warfarin, unfractionated heparin, low molecular weight heparin, synthetic pentasaccharide, hirudin, argatroban, aspirin, ibuprofen, naproxen, sulindac, indomethacin, mefenamate, dipyridamol, droxicam, diclofenac, sulfinpyrazone, piroxicam, ticlopidine, clopidogrel,

tirofiban, eptifibatide, abciximab, melagatran, ximelagatran, disulfatohirudin, tissue plasminogen activator, modified tissue plasminogen activator, anistreplase, urokinase, and streptokinase, or a combination thereof.

In another embodiment, the present invention provides a pharmaceutical composition wherein the additional therapeutic agent is an antihypertensive agent selected from ACE inhibitors, AT-1 receptor antagonists, beta-adrenergic receptor antagonists, ETA receptor antagonists, dual ETA/AT-1 receptor antagonists, renin inhibitors (aliskiren) and vasopepsidase inhibitors, an antiarrhythmic agent selected from I_{Kur} inhibitors, an anticoagulant selected from thrombin inhibitors, antithrombin-III activators, heparin co-Factor II activators, other Factor XIa inhibitors, other kallikrein inhibitors, plasminogen activator inhibitor (PAI-1) antagonists, thrombin activatable fibrinolysis inhibitor (TAFI) inhibitors, Factor VIIa inhibitors, Factor IXa inhibitors, and Factor Xa inhibitors, or an antiplatelet agent selected from GPIIb/IIIa blockers, GP Ib/IX blockers, protease activated receptor 1 (PAR-1) antagonists, protease activated receptor4 (PAR-4) antagonists, prostaglandin E2 receptor EP3 antagonists, collagen receptor antagonists, phosphodiesterase-III inhibitors, P2Y₁ receptor antagonists, P2Y₁₂ antagonists, thromboxane receptor antagonists, cyclooxygenase-1 inhibitors, and aspirin, or a combination thereof.

In another embodiment, the present invention provides pharmaceutical composition, wherein the additional therapeutic agent(s) are an anti-platelet agent or a combination thereof.

In another embodiment, the present invention provides a pharmaceutical composition, wherein the additional therapeutic agent is the anti-platelet agent clopidogrel.

The compounds of the present invention can be administered alone or in combination with one or more additional therapeutic agents. By "administered in combination" or "combination therapy" it is meant that the compound of the present invention and one or more additional therapeutic agents are administered concurrently to the mammal being treated. When administered in combination, each component may be administered at the same time or sequentially in any order at different points in time. Thus, each component may be administered separately but sufficiently closely in time so as to provide the desired therapeutic effect.

Compounds that can be administered in combination with the compounds of the present invention include, but are not limited to, anticoagulants, anti-thrombin agents, anti-platelet agents, fibrinolytics, hypolipidemic agents, antihypertensive agents, and anti-ischemic agents.

5 Other anticoagulant agents (or coagulation inhibitory agents) that may be used in combination with the compounds of this invention include warfarin, heparin (either unfractionated heparin or any commercially available low molecular weight heparin, for example, LOVENOX®), synthetic pentasaccharide, direct acting thrombin inhibitors including hirudin and argatroban, as well as other Factor VIIa inhibitors, Factor IXa
10 inhibitors, Factor Xa inhibitors (e.g., ARIXTRA®, apixaban, rivaroxaban, LY-517717, DU-176b, DX-9065a, and those disclosed in WO 98/57951, WO 03/026652, WO 01/047919, and WO 00/076970), Factor XIa inhibitors, and inhibitors of activated TAFI and PAI-1 known in the art.

15 The term anti-platelet agents (or platelet inhibitory agents), as used herein, denotes agents that inhibit platelet function, for example, by inhibiting the aggregation, adhesion or granule-content secretion of platelets. Such agents include, but are not limited to, the various known non-steroidal anti-inflammatory drugs (NSAIDs) such as acetaminophen, aspirin, codeine, diclofenac, droxicam, fentanyl, ibuprofen, indomethacin, ketorolac, mefenamate, morphine, naproxen, phenacetin, piroxicam,
20 sufentanyl, sulfinpyrazone, sulindac, and pharmaceutically acceptable salts or prodrugs thereof. Of the NSAIDs, aspirin (acetylsalicylic acid or ASA) and piroxicam are preferred. Other suitable platelet inhibitory agents include glycoprotein IIb/IIIa antagonists (e.g., tirofiban, eptifibatide, abciximab, and integrilin), thromboxane-A2-receptor antagonists (e.g., ifetroban), thromboxane-A-synthetase inhibitors,
25 phosphodiesterase-III (PDE-III) inhibitors (e.g., dipyridamole, cilostazol), and PDE-V inhibitors (such as sildenafil), protease-activated receptor 1 (PAR-1) antagonists (e.g., E-5555, SCH-530348, SCH-203099, SCH-529153 and SCH-205831), and pharmaceutically acceptable salts or prodrugs thereof.

30 Other examples of suitable anti-platelet agents for use in combination with the compounds of the present invention, with or without aspirin, are ADP (adenosine diphosphate) receptor antagonists, preferably antagonists of the purinergic receptors P2Y₁ and P2Y₁₂, with P2Y₁₂ being even more preferred. Preferred P2Y₁₂ receptor antagonists

include clopidogrel, ticlopidine, prasugrel, ticagrelor, and cangrelor, and pharmaceutically acceptable salts or prodrugs thereof. Ticlopidine and clopidogrel are also preferred compounds since they are known to be more gentle than aspirin on the gastro-intestinal tract in use. Clopidogrel is an even more preferred agent.

5 A preferred example is a triple combination of a compound of the present invention, aspirin, and another anti-platelet agent. Preferably, the anti-platelet agent is clopidogrel or prasugrel, more preferably clopidogrel.

10 The term thrombin inhibitors (or anti-thrombin agents), as used herein, denotes inhibitors of the serine protease thrombin. By inhibiting thrombin, various thrombin-mediated processes, such as thrombin-mediated platelet activation (that is, for example, the aggregation of platelets, and/or the secretion of platelet granule contents including serotonin) and/or fibrin formation are disrupted. A number of thrombin inhibitors are known to one of skill in the art and these inhibitors are contemplated to be used in combination with the present compounds. Such inhibitors include, but are not limited to, 15 boroarginine derivatives, boropeptides, heparins, hirudin, argatroban, dabigatran, AZD-0837, and those disclosed in WO 98/37075 and WO 02/044145, and pharmaceutically acceptable salts and prodrugs thereof. Boroarginine derivatives and boropeptides include N-acetyl and peptide derivatives of boronic acid, such as C-terminal a-aminoboronic acid derivatives of lysine, ornithine, arginine, homoarginine and corresponding isothiouronium 20 analogs thereof. The term hirudin, as used herein, includes suitable derivatives or analogs of hirudin, referred to herein as hirulogs, such as disulfatohirudin.

25 The term thrombolytic (or fibrinolytic) agents (or thrombolytics or fibrinolytics), as used herein, denotes agents that lyse blood clots (thrombi). Such agents include tissue plasminogen activator (TPA, natural or recombinant) and modified forms thereof, anistreplase, urokinase, streptokinase, tenecteplase (TNK), lanoteplase (nPA), Factor VIIa inhibitors, thrombin inhibitors, inhibitors of Factors IXa, Xa, and XIa, PAI-I inhibitors (i.e., inactivators of tissue plasminogen activator inhibitors), inhibitors of activated TAFI, alpha-2-antiplasmin inhibitors, and anisoylated plasminogen streptokinase activator complex, including pharmaceutically acceptable salts or prodrugs thereof. The term 30 anistreplase, as used herein, refers to anisoylated plasminogen streptokinase activator complex, as described, for example, in European Patent Application No. 028,489, the disclosure of which is hereby incorporated herein by reference herein. The term

urokinase, as used herein, is intended to denote both dual and single chain urokinase, the latter also being referred to herein as prourokinase.

Examples of suitable cholesterol/lipid lowering agents and lipid profile therapies for use in combination with the compounds of the present invention include HMG-CoA 5 reductase inhibitors (*e.g.*, pravastatin, lovastatin, simvastatin, fluvastatin, atorvastatin, rosuvastatin, and other statins), low-density lipoprotein (LDL) receptor activity modulators (*e.g.*, HOE-402, PCSK9 inhibitors), bile acid sequestrants (*e.g.*, cholestyramine and colestipol), nicotinic acid or derivatives thereof (*e.g.*, NIASPAN®), GPR109B (nicotinic acid receptor) modulators, fenofibric acid derivatives (*e.g.*, 10 gemfibrozil, clofibrate, fenofibrate and benzafibrate) and other peroxisome proliferator-activated receptors (PPAR) alpha modulators, PPARdelta modulators (*e.g.*, GW-501516), PPARgamma modulators (*e.g.*, rosiglitazone), compounds that have multiple functionality for modulating the activities of various combinations of PPARalpha, PPARgamma and PPARdelta, probucol or derivatives thereof (*e.g.*, AGI-1067), 15 cholesterol absorption inhibitors and/or Niemann-Pick C1-like transporter inhibitors (*e.g.*, ezetimibe), cholesterol ester transfer protein inhibitors (*e.g.*, CP-529414), squalene synthase inhibitors and/or squalene epoxidase inhibitors or mixtures thereof, acyl coenzyme A: cholestryl acyltransferase (ACAT) 1 inhibitors, ACAT2 inhibitors, dual 20 ACAT1/2 inhibitors, ileal bile acid transport inhibitors (or apical sodium co-dependent bile acid transport inhibitors), microsomal triglyceride transfer protein inhibitors, liver-X-receptor (LXR) alpha modulators, LXRbeta modulators, LXR dual alpha/beta modulators, FXR modulators, omega 3 fatty acids (*e.g.*, 3-PUFA), plant stanols and/or fatty acid esters of plant stanols (*e.g.*, sitostanol ester used in BENECOL® margarine), endothelial lipase 25 inhibitors, and HDL functional mimetics which activate reverse cholesterol transport (*e.g.*, apoAI derivatives or apoAI peptide mimetics).

The compounds of the present invention are also useful as standard or reference compounds, for example, as a quality standard or control, in tests or assays involving the inhibition of thrombin, Factor VIIa, IXa, Xa, XIa, and/or plasma kallikrein. Such compounds may be provided in a commercial kit, for example, for use in pharmaceutical 30 research involving thrombin, Factor VIIa, IXa, Xa, XIa, and/or plasma kallikrein. XIa. For example, a compound of the present invention could be used as a reference in an assay to compare its known activity to a compound with an unknown activity. This

would ensure the experimenter that the assay was being performed properly and provide a basis for comparison, especially if the test compound was a derivative of the reference compound. When developing new assays or protocols, compounds according to the present invention could be used to test their effectiveness.

5 The compounds of the present invention may also be used in diagnostic assays involving thrombin, Factor VIIa, IXa, Xa, XIa, and/or plasma kallikrein. For example, the presence of thrombin, Factor VIIa, IXa, Xa XIa, and/or plasma kallikrein in an unknown sample could be determined by addition of the relevant chromogenic substrate, for example, S2366 for Factor XIa, to a series of solutions containing test sample and 10 optionally one of the compounds of the present invention. If production of pNA is observed in the solutions containing test sample, but not in the presence of a compound of the present invention, then one would conclude Factor XIa was present.

Extremely potent and selective compounds of the present invention, those having K_i values less than or equal to 0.001 μM against the target protease and greater than or 15 equal to 0.1 μM against the other proteases, may also be used in diagnostic assays involving the quantitation of thrombin, Factor VIIa, IXa, Xa, XIa, and/or plasma kallikrein in serum samples. For example, the amount of Factor XIa in serum samples could be determined by careful titration of protease activity in the presence of the relevant chromogenic substrate, S2366, with a potent Factor XIa inhibitor of the present invention.

20 The present invention also encompasses an article of manufacture. As used herein, article of manufacture is intended to include, but not be limited to, kits and packages. The article of manufacture of the present invention, comprises: (a) a first container; (b) a pharmaceutical composition located within the first container, wherein the composition, comprises: a first therapeutic agent, comprising: a compound of the present 25 invention or a pharmaceutically acceptable salt form thereof; and, (c) a package insert stating that the pharmaceutical composition can be used for the treatment of a thromboembolic and/or inflammatory disorder (as defined previously). In another embodiment, the package insert states that the pharmaceutical composition can be used in combination (as defined previously) with a second therapeutic agent to treat a 30 thromboembolic and/or inflammatory disorder. The article of manufacture can further comprise: (d) a second container, wherein components (a) and (b) are located within the second container and component (c) is located within or outside of the second container.

Located within the first and second containers means that the respective container holds the item within its boundaries.

The first container is a receptacle used to hold a pharmaceutical composition. This container can be for manufacturing, storing, shipping, and/or individual/bulk selling.

5 First container is intended to cover a bottle, jar, vial, flask, syringe, tube (e.g., for a cream preparation), or any other container used to manufacture, hold, store, or distribute a pharmaceutical product.

The second container is one used to hold the first container and, optionally, the package insert. Examples of the second container include, but are not limited to, boxes

10 (e.g., cardboard or plastic), crates, cartons, bags (e.g., paper or plastic bags), pouches, and sacks. The package insert can be physically attached to the outside of the first container via tape, glue, staple, or another method of attachment, or it can rest inside the second container without any physical means of attachment to the first container. Alternatively, the package insert is located on the outside of the second container. When located on the

15 outside of the second container, it is preferable that the package insert is physically attached via tape, glue, staple, or another method of attachment. Alternatively, it can be adjacent to or touching the outside of the second container without being physically attached.

The package insert is a label, tag, marker, etc. that recites information relating to

20 the pharmaceutical composition located within the first container. The information recited will usually be determined by the regulatory agency governing the area in which the article of manufacture is to be sold (e.g., the United States Food and Drug Administration). Preferably, the package insert specifically recites the indications for which the pharmaceutical composition has been approved. The package insert may be

25 made of any material on which a person can read information contained therein or thereon. Preferably, the package insert is a printable material (e.g., paper, plastic, cardboard, foil, adhesive-backed paper or plastic, etc.) on which the desired information has been formed (e.g., printed or applied).

Other features of the invention will become apparent in the course of the

30 following descriptions of exemplary embodiments that are given for illustration of the invention and are not intended to be limiting thereof. The following Examples have been prepared, isolated and characterized using the methods disclosed herein.

VI. GENERAL SYNTHESIS INCLUDING SCHEMES


The compounds of the present invention can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or by variations thereon as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below. The reactions are performed in a solvent or solvent mixture appropriate to the reagents and materials employed and suitable for the transformations being effected. It will be understood by those skilled in the art of organic synthesis that the functionality present on the molecule should be consistent with the transformations proposed. This will sometimes require a judgment to modify the order of the synthetic steps or to select one particular process scheme over another in order to obtain a desired compound of the invention. Example compounds are typically prepared as racemic mixtures. Preparation of homochiral examples may be carried out by techniques known to one skilled in the art. For example, homochiral compounds may be prepared by separation of racemic products by chiral phase preparative HPLC. Alternatively, the example compounds may be prepared by methods known to give enantiomerically enriched products. These include, but are not limited to, the incorporation of chiral auxiliary functionalities into racemic intermediates which serve to control the diastereoselectivity of transformations, providing enantio-enriched products upon cleavage of the chiral auxiliary.

It will also be recognized that another major consideration in the planning of any synthetic route in this field is the judicious choice of the protecting group used for protection of the reactive functional groups present in the compounds described in this invention. An authoritative account describing the many alternatives to the trained practitioner is Greene et al. (*Protective Groups in Organic Synthesis*, Fourth Edition, Wiley-Interscience (2006)).

Compounds of this invention can be derived from intermediates **1g**, the synthesis of which is described in Scheme 1. Esterification of amino acid **1a** gives amino ester **1b**. Amino ester **1b** can then be coupled with an appropriately substituted carboxylic acid **1h** using coupling agent, such as EDC/HOBt. Aniline **1c** can then be coupled with an appropriately substituted carboxylic acid **1j** using T3P® and a base, such as pyridine, to give the amide **1d**. Using a modified procedure described by Lovely (*Tetrahedron Lett.*,

44:1379 (2003)), **1d** can be cyclized via ring-closing metathesis using a catalyst, such as Grubbs (II), in a suitable solvent, such as DCM, DCE, or toluene at elevated temperature, to give the macrocycle **1e**. The alkene can be reduced with hydrogen over either palladium on carbon or platinum oxide to provide ester **1f**. Hydrolysis of ester **1f** gives acid **1g**. Compounds of the formula (I) can be synthesized from the coupling reaction of acid **1g** and amine **1i** by using coupling agents, such as HATU.

Scheme 1




10

Olefin containing macrocycles of this invention can also be prepared according to Scheme 2. Hydrolysis of ester **1e** gives acid **2a**. Compounds of the formula (I) can be synthesized from the coupling reaction of acid **2a** and amine **1i** by using coupling agents, such as HATU.


15

Scheme 2



WHAT IS CLAIMED IS:

1. A compound of Formula (I):



(I)

or a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein:

---- is an optional bond;

ring A is independently selected from a mono or bicyclic heterocycle comprising: carbon atoms and 1-4 heteroatoms selected from NR², O, and S;

10 ring B is independently selected from aryl and a 5- to 10-membered heterocycle;

R¹ is, independently at each occurrence, selected from halogen, C₁₋₆ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, OH, =O, OCH₂F, OCHF₂, OCF₃, CF₃, CN, NH₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, CO₂(C₁₋₄ alkyl), CO(C₁₋₄ alkyl), -CH₂NH₂, -CONH₂, -CONH(C₁₋₄ alkyl), -OCH₂CO₂H, -NHCO(C₁₋₄ alkyl), -NHCO₂(C₁₋₄ alkyl), -NHSO₂(C₁₋₄ alkyl), -SO₂NH₂, and -C(=NH)NH₂;

15 R² is absent or independently selected from H and C₁₋₄alkyl;

R³ is independently selected from H, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, OH, =O, OCH₂F, OCHF₂, OCF₃, CN, NH₂, NH(C₁₋₄ alkyl), and N(C₁₋₄ alkyl)₂;

R⁴ is independently selected from H, C₁₋₄ alkyl, hydroxyl, F, CF₃, and C₃₋₆

20 cycloalkyl;

R⁵ is independently selected from H and C₁₋₄ alkyl;

R⁶ is independently selected from H, halogen, C(O)OH, and C(O)O(C₁₋₄ alkyl);


R⁷ is independently selected from H, C₁₋₄ alkyl, and CF₃;

alternatively, R⁶ and R⁷ together are =O; and

R⁸ is, independently at each occurrence, selected from H, halogen, NHCO₂C₁₋₄ alkyl, CN, OH, O-C₁₋₄ alkyl, CF₃, -OCF₃, -OCHF₂, -CHF₂, -CH₂F, CO₂H, CO₂(C₁₋₄ alkyl), -CH₂CO₂H, -(CH₂)₂CO₂H, -CH₂CO₂(C₁₋₄ alkyl), -(CH₂)₂CO₂(C₁₋₄ alkyl), NH₂, -CH₂NH₂, -NHCO(C₁₋₄ alkyl), -NHCO₂(CH₂)₁₋₂O(C₁₋₄ alkyl), -NHCO₂(CH₂)₁₋₃O(C₁₋₄ alkyl), NHCO₂CH₂CH(C₁₋₄ alkyl)O(C₁₋₄ alkyl), -NHCO₂(CH₂)₁₋₂OH, -NHCO₂CH₂CO₂H, -CH₂NHCO₂(C₁₋₄ alkyl), -NHC(O)NH(C₁₋₄ alkyl), -NHC(O)N(C₁₋₄ alkyl)₂, NHC(O)NH(C₁₋₄ alkyl)N(5- to 6-membered heterocycle), -NHSO₂(C₁₋₄ alkyl), -CONH₂, -CONH(C₁₋₄ alkyl), -CON(C₁₋₄ alkyl)₂, and -CH₂CONH₂.

10

2. The compound of claim 1 having Formula (II):



or a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein:

---- is an optional bond;

15 ring A is independently selected from a mono or bicyclic heterocycle comprising: carbon atoms and 1-4 heteroatoms selected from NR², O, and S;

R¹ is, independently at each occurrence, selected from halogen, C₁₋₆ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, OH, =O, OCH₂F, OCHF₂, OCF₃, CN, NH₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, CO₂(C₁₋₄ alkyl), CO(C₁₋₄ alkyl), -CH₂NH₂, -CONH₂, -CONH(C₁₋₄ alkyl), -OCH₂CO₂H, -NHCO(C₁₋₄ alkyl), -NHCO₂(C₁₋₄ alkyl), -NHSO₂(C₁₋₄ alkyl), -SO₂NH₂, and -C(=NH)NH₂;

R² is absent or independently selected from H and C₁₋₄alkyl;

R³ is independently selected from H and C₁₋₄ alkyl;


R⁴ is independently selected from H, C₁₋₄ alkyl, hydroxyl, and C₃₋₆ cycloalkyl;

R^5 is independently selected from H and C_{1-4} alkyl; and

R^8 is independently selected from H, halogen, $NHCO_2C_{1-4}$ alkyl, CN, OH, $O-C_{1-4}$ alkyl, CF_3 , $-OCF_3$, $-OCHF_2$, $-CHF_2$, $-CH_2F$, CO_2H , $CO_2(C_{1-4}$ alkyl), $-CH_2CO_2H$, $-(CH_2)_2CO_2H$, $-CH_2CO_2(C_{1-4}$ alkyl), $-(CH_2)_2CO_2(C_{1-4}$ alkyl),
5 NH_2 , $-CH_2NH_2$, $-NHCO(C_{1-4}$ alkyl), $-NHCO_2(CH_2)_{1-2}O(C_{1-4}$ alkyl), $-NHCO_2(CH_2)_{1-3}O(C_{1-4}$ alkyl), $NHCO_2CH_2CH(C_{1-4}$ alkyl) $O(C_{1-4}$ alkyl), $-NHCO_2(CH_2)_{1-2}OH$, $-NHCO_2CH_2CO_2H$, $-CH_2NHCO_2(C_{1-4}$ alkyl), $-NHC(O)NH(C_{1-4}$ alkyl), $-NHC(O)N(C_{1-4}$ alkyl) $_2$, $NHC(O)NH(C_{1-4}$ alkyl) $N[5-$ to
6-membered heterocycle] $]$, $-NSO_2(C_{1-4}$ alkyl), $-CONH_2$, $-CONH(C_{1-4}$ alkyl), $-CON(C_{1-4}$ alkyl) $_2$, and $-CH_2CONH_2$.

10

3. The compound of claim 2 having Formula (III):



or a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein:

15

ring C is independently selected from 6-membered aryl and heteroaryl;

R^1 is, independently at each occurrence, selected from halogen, C_{1-6} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, C_{1-4} haloalkyl, OH, OCH_2F , $OCHF_2$, OCF_3 , CN, NH_2 , $NH(C_{1-4}$ alkyl), and $N(C_{1-4}$ alkyl) $_2$;

R^2 is independently selected from H and C_{1-4} alkyl;


20

R^4 is independently selected from H, C_{1-4} alkyl, hydroxyl, and C_{3-6} cycloalkyl;

R^5 is independently selected from H and C_{1-4} alkyl; and

R⁸ is independently selected from H, halogen, NHCO₂C₁₋₄ alkyl, CN, OH, O-C₁₋₄ alkyl, CF₃, -OCF₃, -OCHF₂, -CHF₂, -CH₂F, CO₂H, and CO₂(C₁₋₄ alkyl).

4. The compound of claim 3 having Formula (IV):



or a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein:

---- is an optional bond;

R¹ is independently selected from F and Cl;

R² is independently selected from H and C₁₋₄alkyl;

10 R⁴ is independently selected from H, C₁₋₄ alkyl, and hydroxyl;

R⁵ is independently selected from H and C₁₋₄ alkyl; and

R⁸ is independently selected from H, halogen, NHCO₂C₁₋₄ alkyl, CN, OH, O-C₁₋₄ alkyl, CF₃, -OCF₃, -OCHF₂, -CHF₂, -CH₂F, CO₂H, and CO₂(C₁₋₄ alkyl).

15 5. The compound of claim 4, or a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein:

---- is an optional bond;

R¹ is independently selected from F and Cl;





R² is H;

20 R⁴ is independently selected from H, C₁₋₄ alkyl, hydroxyl, and C₃₋₆ cycloalkyl;

R⁵ is independently selected from H and C₁₋₄ alkyl; and

R⁸ is independently selected from H, halogen, and NHCO₂C₁₋₄ alkyl.

6. The compound of claim 1 selected from:



or a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof.

5

7. A pharmaceutical composition comprising one or more compounds according to any one of claims 1-6 and a pharmaceutically acceptable carrier or diluent.

8. A method for the treatment and/or prophylaxis of a thromboembolic disorder, comprising: administering to a patient in need thereof a therapeutically effective amount of a compound of any one of claims 1-6, or a stereoisomer, a tautomer, or a pharmaceutically acceptable salt thereof.

9. A method according to claim 8, wherein the thromboembolic disorder is selected from the group consisting of arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, and thromboembolic disorders in the chambers of the heart or in the peripheral circulation.

10. A method according to claim 9, wherein the thromboembolic disorder is selected from unstable angina, an acute coronary syndrome, atrial fibrillation, myocardial

infarction, transient ischemic attack, stroke, atherosclerosis, peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants,
5 devices, or procedures in which blood is exposed to an artificial surface that promotes thrombosis.

11. A compound of any one of claims 1-6, or a stereoisomer, a tautomer, or a pharmaceutically acceptable salt thereof, for use in therapy.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/020161

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D498/10 A61K31/537 A61P7/02
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2014/022766 A1 (SQUIBB BRISTOL MYERS CO [US]) 6 February 2014 (2014-02-06) claims; examples -----	1-11
A	WO 2011/100401 A1 (SQUIBB BRISTOL MYERS CO [US]; CORTE JAMES R [US]; FANG TIANAN [US]; DE) 18 August 2011 (2011-08-18) claims; examples -----	1-11
Y	WO 2015/164308 A1 (MERCK SHARP & DOHME [US]; LIM YEON-HEE [US]; GUO ZHUYAN [US]; ALI AMJA) 29 October 2015 (2015-10-29) claims; examples -----	1-11
Y, P	WO 2016/036893 A1 (SQUIBB BRISTOL MYERS CO [US]) 10 March 2016 (2016-03-10) claims; examples -----	1-11



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

7 April 2017

20/04/2017

Name and mailing address of the ISA/
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Stroeter, Thomas

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/020161

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO 2014022766	A1 06-02-2014	AR 093759 A1 AU 2013296258 A1 CA 2880898 A1 CN 104520289 A DK 2882734 T3 EA 201590284 A1 EP 2882734 A1 ES 2605824 T3 HR P20161378 T1 JP 6082462 B2 JP 2015524443 A KR 20150038369 A LT 2882734 T SG 11201500270R A SI 2882734 T1 SM T201700003 B TW 201410666 A US 2014038969 A1 US 2015166550 A1 UY 34959 A WO 2014022766 A1		24-06-2015 19-03-2015 06-02-2014 15-04-2015 30-01-2017 29-05-2015 17-06-2015 16-03-2017 02-12-2016 15-02-2017 24-08-2015 08-04-2015 12-12-2016 30-03-2015 31-01-2017 08-03-2017 16-03-2014 06-02-2014 18-06-2015 31-01-2014 06-02-2014

WO 2011100401	A1 18-08-2011	AR 080167 A1 AU 2011215898 A1 CA 2789622 A1 CL 2012002229 A1 CN 102753555 A CN 102834397 A CO 6551666 A2 EA 201270716 A1 EP 2534152 A1 EP 2534154 A1 JP 5841547 B2 JP 5844287 B2 JP 2013519678 A JP 2013519679 A JP 2016102114 A KR 20120134122 A NZ 602266 A PE 18062012 A1 SG 183153 A1 TW 201130847 A TW 201605806 A US 2012041190 A1 US 2012309960 A1 US 2014343276 A1 US 2015112058 A1 US 2015266894 A1 US 2016362414 A1 WO 2011100401 A1 WO 2011100402 A1 ZA 201206758 B		21-03-2012 27-09-2012 18-08-2011 11-01-2013 24-10-2012 19-12-2012 31-10-2012 29-03-2013 19-12-2012 13-01-2016 13-01-2016 30-05-2013 30-05-2013 02-06-2016 11-12-2012 30-05-2014 02-01-2013 27-09-2012 16-09-2011 16-02-2016 16-02-2012 06-12-2012 20-11-2014 23-04-2015 24-09-2015 15-12-2016 18-08-2011 18-08-2011 26-02-2014

WO 2015164308	A1 29-10-2015	EP 3134408 A1 US 2017044183 A1 WO 2015164308 A1		01-03-2017 16-02-2017 29-10-2015

WO 2016036893	A1 10-03-2016	NONE		

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/US2017/020161

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
<hr/>			