



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : <b>C12N 15/12, C07K 14/47, A61K 38/17, C07K 16/18, G01N 33/50</b></p>	<p><b>A2</b></p>	<p>(11) International Publication Number: <b>WO 00/15797</b></p> <p>(43) International Publication Date: 23 March 2000 (23.03.00)</p>
<p>(21) International Application Number: PCT/US99/21547</p> <p>(22) International Filing Date: 15 September 1999 (15.09.99)</p> <p>(30) Priority Data: 60/100,858 17 September 1998 (17.09.98) US PCT/US98/19437 17 September 1998 (17.09.98) US</p> <p>(71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): FONG, Sherman [US/US]; 19 Basinside Way, Alameda, CA 94502 (US). GODDARD, Audrey [CA/US]; 110 Congo Street, San Francisco, CA 94131 (US). GURNEY, Austin, L. [US/US]; 1 Debbie Lane, Belmont, CA 94002 (US). TUMAS, Daniel [US/US]; 3 Rae Avenue, Orinda, CA 94563 (US). WOOD, William, I. [US/US]; 35 Southdown Court, Hillsborough, CA 94010 (US).</p> <p>(74) Agents: SCHWARTZ, Timothy, R. et al.; Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080-4990 (US).</p>	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i></p>	
<p>(54) Title: COMPOSITIONS AND METHODS FOR THE TREATMENT OF IMMUNE RELATED DISEASES</p>		
<p>(57) Abstract</p> <p>The present invention relates to a composition containing novel proteins and methods for the diagnosis and treatment of immune related diseases.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

<b>AL</b>	Albania	<b>ES</b>	Spain	<b>LS</b>	Lesotho	<b>SI</b>	Slovenia
<b>AM</b>	Armenia	<b>FI</b>	Finland	<b>LT</b>	Lithuania	<b>SK</b>	Slovakia
<b>AT</b>	Austria	<b>FR</b>	France	<b>LU</b>	Luxembourg	<b>SN</b>	Senegal
<b>AU</b>	Australia	<b>GA</b>	Gabon	<b>LV</b>	Latvia	<b>SZ</b>	Swaziland
<b>AZ</b>	Azerbaijan	<b>GB</b>	United Kingdom	<b>MC</b>	Monaco	<b>TD</b>	Chad
<b>BA</b>	Bosnia and Herzegovina	<b>GE</b>	Georgia	<b>MD</b>	Republic of Moldova	<b>TG</b>	Togo
<b>BB</b>	Barbados	<b>GH</b>	Ghana	<b>MG</b>	Madagascar	<b>TJ</b>	Tajikistan
<b>BE</b>	Belgium	<b>GN</b>	Guinea	<b>MK</b>	The former Yugoslav Republic of Macedonia	<b>TM</b>	Turkmenistan
<b>BF</b>	Burkina Faso	<b>GR</b>	Greece	<b>ML</b>	Mali	<b>TR</b>	Turkey
<b>BG</b>	Bulgaria	<b>HU</b>	Hungary	<b>MN</b>	Mongolia	<b>TT</b>	Trinidad and Tobago
<b>BJ</b>	Benin	<b>IE</b>	Ireland	<b>MR</b>	Mauritania	<b>UA</b>	Ukraine
<b>BR</b>	Brazil	<b>IL</b>	Israel	<b>MW</b>	Malawi	<b>UG</b>	Uganda
<b>BY</b>	Belarus	<b>IS</b>	Iceland	<b>MX</b>	Mexico	<b>US</b>	United States of America
<b>CA</b>	Canada	<b>IT</b>	Italy	<b>NE</b>	Niger	<b>UZ</b>	Uzbekistan
<b>CF</b>	Central African Republic	<b>JP</b>	Japan	<b>NL</b>	Netherlands	<b>VN</b>	Viet Nam
<b>CG</b>	Congo	<b>KE</b>	Kenya	<b>NO</b>	Norway	<b>YU</b>	Yugoslavia
<b>CH</b>	Switzerland	<b>KG</b>	Kyrgyzstan	<b>NZ</b>	New Zealand	<b>ZW</b>	Zimbabwe
<b>CI</b>	Côte d'Ivoire	<b>KP</b>	Democratic People's Republic of Korea	<b>PL</b>	Poland		
<b>CM</b>	Cameroon	<b>KR</b>	Republic of Korea	<b>PT</b>	Portugal		
<b>CN</b>	China	<b>KZ</b>	Kazakstan	<b>RO</b>	Romania		
<b>CU</b>	Cuba	<b>LC</b>	Saint Lucia	<b>RU</b>	Russian Federation		
<b>CZ</b>	Czech Republic	<b>LI</b>	Liechtenstein	<b>SD</b>	Sudan		
<b>DE</b>	Germany	<b>LK</b>	Sri Lanka	<b>SE</b>	Sweden		
<b>DK</b>	Denmark	<b>LR</b>	Liberia	<b>SG</b>	Singapore		
<b>EE</b>	Estonia						

## COMPOSITIONS AND METHODS FOR THE TREATMENT OF IMMUNE RELATED DISEASES

### Field of the Invention

The present invention relates to compositions and methods for the diagnosis and treatment of immune related diseases.

### Background of the Invention

Immune related and inflammatory diseases are the manifestation or consequence of fairly complex, often multiple interconnected biological pathways which in normal physiology are critical to respond to insult or injury, initiate repair from insult or injury, and mount innate and acquired defense against foreign organisms. Disease or pathology occurs when these normal physiological pathways cause additional insult or injury either as directly related to the intensity of the response, as a consequence of abnormal regulation or excessive stimulation, as a reaction to self, or as a combination of these.

Though the genesis of these diseases often involves multistep pathways and often multiple different biological systems/pathways, intervention at critical points in one or more of these pathways can have an ameliorative or therapeutic effect. Therapeutic intervention can occur by either antagonism of a detrimental process/pathway or stimulation of a beneficial process/pathway.

Many immune related diseases are known and have been extensively studied. Such diseases include immune-mediated inflammatory diseases, non-immune-mediated inflammatory diseases, infectious diseases, immunodeficiency diseases, neoplasia, etc.

T lymphocytes (T cells) are an important component of a mammalian immune response. T cells recognise antigens which are associated with a self-molecule encoded by genes within the major histocompatibility complex (MHC). The antigen may be displayed together with MHC molecules on the surface of antigen presenting cells, virus infected cells, cancer cells, grafts, etc. The T cell system eliminates these altered cells which pose a health threat to the host mammal. T cells include helper T cells and cytotoxic T cells. Helper T cells proliferate extensively following recognition of an antigen-MHC complex on an antigen presenting cell. Helper T cells also secrete a variety of cytokines, i.e. lymphokines, which play a central role in the activation of B cells, cytotoxic T cells and a variety of other cells which participate in the immune response.

A central event in both humoral and cell mediated immune responses is the activation and clonal expansion of helper T cells. Helper T cell activation is initiated by the interaction of the T cell receptor (TCR) - CD3 complex with an antigen-MHC on the surface of an antigen presenting cell.

This interaction mediates a cascade of biochemical events that induce the resting helper T cell to enter a cell cycle (the G<sub>0</sub> to G<sub>1</sub> transition) and results in the expression of a high affinity receptor for IL-2 and sometimes IL-4. The activated T cell progresses through the cycle proliferating and differentiating into memory cells or effector cells.

In addition to the signals mediated through the TCR, activation of T cells involves additional costimulation induced by cytokines released by the antigen presenting cell or through interactions with membrane bound molecules on the antigen presenting cell and the T cell. The cytokines IL-1 and IL-6 have been shown to provide a costimulatory signal. Also, the interaction between the B7 molecule expressed on the surface of an antigen presenting cell and CD28 and CTLA-4 molecules expressed on the T cell surface effect T cell activation. Activated T cells express an increased number of cellular adhesion molecules, such as ICAM-1, integrins, VLA-4, LFA-1, CD56, etc.

T-cell proliferation in a mixed lymphocyte culture or mixed lymphocyte reaction (MLR) is an established indication of the ability of a compound to stimulate the immune system. In many immune responses, inflammatory cells infiltrate the site of injury or infection. The migrating cells may be neutrophilic, eosinophilic, monocytic or lymphocytic. Histologic examination of the affected tissues provides evidence of an immune stimulating or inhibiting response Current Protocols in Immunology, ed. John E. Coligan, 1994, John Wiley & Sons, Inc.

Immune related diseases can be treated by suppressing the immune response. Using neutralizing antibodies that inhibit molecules having immune stimulatory activity would be beneficial in the treatment of immune-mediated and inflammatory diseases. Molecules which inhibit the immune response can be utilized (proteins directly or via the use of antibody agonists) to inhibit the immune response and thus ameliorate immune related disease.

#### Summary of the Invention

The present invention concerns compositions and methods for the diagnosis and treatment of immune related disease in mammals, including humans. The present invention is based on the identification of proteins (including agonist and antagonist antibodies) which either stimulate or inhibit the immune response in mammals. Immune related diseases can be treated by suppressing or enhancing the immune response. Molecules that enhance the immune response stimulate or potentiate the immune response to an antigen. Molecules which stimulate the immune response can be used therapeutically where enhancement of the immune response would be beneficial. Such stimulatory molecules can also be inhibited where suppression of the immune response would be of value. Neutralizing antibodies are examples of molecules that inhibit molecules having immune stimulatory activity and which would be beneficial in the treatment of immune related and inflammatory diseases. Molecules which inhibit the immune response can also be utilized (proteins directly or via the use of antibody agonists) to inhibit the immune response and thus ameliorate immune related disease.

Accordingly, the proteins of the invention encoded by the genes of the invention are useful for the diagnosis and/or treatment (including prevention) of immune related diseases. Antibodies which bind to stimulatory proteins are useful to suppress the immune system and the immune response. Antibodies which bind to inhibitory proteins are useful to stimulate the immune system and the immune response. The proteins and antibodies of the invention are also useful to prepare medicines and medicaments for the treatment of immune related and inflammatory diseases.

In one embodiment, the present invention concerns an isolated antibody which binds a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide. In one aspect, the antibody mimics the activity of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide (an agonist antibody) or conversely the antibody inhibits or neutralizes the activity of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide (an antagonist antibody). In another aspect, the antibody is a monoclonal antibody, which preferably has nonhuman complementarity determining region (CDR) residues and human framework region (FR) residues. The antibody may be labeled and may be immobilized on a solid support. In a further aspect, the antibody is an antibody fragment, a single-chain antibody, or an anti-idiotypic antibody.

In another embodiment, the invention concerns a composition containing a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide or an agonist or antagonist antibody which binds the polypeptide in admixture with a carrier or excipient. In one aspect, the composition contains a therapeutically effective amount of the peptide or antibody. In another aspect, when the composition contains an immune stimulating molecule, the composition is useful for: (a) increasing infiltration of inflammatory cells into a tissue of a mammal in need thereof, (b) stimulating or enhancing an immune response in a mammal in need thereof, or (c) increasing the proliferation of T-lymphocytes in a mammal in need thereof in response to an antigen. In a further aspect, when the composition contains an immune inhibiting molecule, the composition is useful for: (a) decreasing infiltration of inflammatory cells into a tissue of a mammal in need thereof, (b) inhibiting or reducing an immune response in a mammal in need thereof, or (c) decreasing the proliferation of T-lymphocytes in a mammal in need thereof in response to an antigen. In another aspect, the composition contains a further active ingredient, which may, for example, be a further antibody or a cytotoxic or chemotherapeutic agent. Preferably, the composition is sterile.

In another embodiment, the invention concerns the use of the polypeptides and antibodies of the invention to prepare a composition or medicament which has the uses described above.

In a further embodiment, the invention concerns nucleic acid encoding an anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody, and vectors and recombinant host cells comprising such nucleic acid. In a still further embodiment, the invention concerns a method for producing such an antibody by culturing a host cell transformed with nucleic acid encoding the antibody under conditions such that the antibody is expressed, and recovering the

antibody from the cell culture.

The invention further concerns antagonists and agonists of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide that inhibit one or more of the functions or activities of the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide.

In a further embodiment, the invention concerns isolated nucleic acid molecules that hybridize to the complement of the nucleic acid molecules encoding the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptides. The nucleic acid preferably is DNA, and hybridization preferably occurs under stringent conditions. Such nucleic acid molecules can act as antisense molecules of the amplified genes identified herein, which, in turn, can find use in the modulation of the respective amplified genes, or as antisense primers in amplification reactions. Furthermore, such sequences can be used as part of ribozyme and/or triple helix sequence which, in turn, may be used in regulation of the amplified genes.

In another embodiment, the invention concerns a method for determining the presence of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide comprising exposing a cell suspected of containing the polypeptide to an anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody and determining binding of the antibody to the cell.

In yet another embodiment, the present invention concerns a method of diagnosing an immune related disease in a mammal, comprising detecting the level of expression of a gene encoding a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide (a) in a test sample of tissue cells obtained from the mammal, and (b) in a control sample of known normal tissue cells of the same cell type, wherein a higher expression level in the test sample indicates the presence of immune related disease in the mammal from which the test tissue cells were obtained.

In another embodiment, the present invention concerns a method of diagnosing an immune disease in a mammal, comprising (a) contacting an anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody with a test sample of tissue cells obtained from the mammal, and (b) detecting the formation of a complex between the antibody and the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide in the test sample. The detection may be qualitative or quantitative, and may be performed in comparison with monitoring the complex formation in a control sample of known normal tissue cells of the same cell type. A larger quantity of complexes formed in the test sample indicates the presence of tumor in the mammal from which the test tissue cells were obtained. The antibody preferably carries a detectable label. Complex formation can be monitored, for example, by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. The test sample is usually obtained from an individual suspected of having a deficiency or abnormality of the immune system.

In another embodiment, the present invention concerns a diagnostic kit, containing an anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody and a carrier (e.g. a

buffer) in suitable packaging. The kit preferably contains instructions for using the antibody to detect the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide.

In a further embodiment, the invention concerns an article of manufacture, comprising:

a container;

a label on the container; and

a composition comprising an active agent contained within the container; wherein the composition is effective for stimulating or inhibiting an immune response in a mammal, the label on the container indicates that the composition can be used to treat an immune related disease, and the active agent in the composition is an agent stimulating or inhibiting the expression and/or activity of the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide. In a preferred aspect, the active agent is a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide or an anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody.

A further embodiment is a method for identifying a compound capable of inhibiting the expression and/or activity of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide by contacting a candidate compound with a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide under conditions and for a time sufficient to allow these two components to interact. In a specific aspect, either the candidate compound or the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide is immobilized on a solid support. In another aspect, the non-immobilized component carries a detectable label.

#### Brief Description of the Drawings

Figures 1A-D and Table 4 show hypothetical exemplifications for using the below described method to determine % amino acid sequence identity (Figures 1A-B) and % nucleic acid sequence identity (Figures 1C-D) using the ALIGN-2 sequence comparison computer program, wherein "PRO" represents the amino acid sequence of a hypothetical polypeptide of the invention of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, "PRO-DNA" represents a hypothetical "PRO"-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, "X", "Y" and "Z" each represent different hypothetical amino acid residues and "N", "L" and "V" each represent different hypothetical nucleotides.

Figures 2A-P and Table 5 provide the complete source code for the ALIGN-2 sequence comparison computer program. This source code may be routinely compiled for use on a UNIX operating system to provide the ALIGN-2 sequence comparison computer program.

Figure 3 shows the nucleotide sequence of a cDNA containing a nucleotide sequence encoding native sequence PRO245(UNQ219), wherein the nucleotide sequence (SEQ ID NO: 1) is a

clone designated herein as "DNA35638". Also presented in bold font and underlined are the positions of the respective start and stop codons.

Figure 4 and Table 6 show the amino acid sequence (SEQ ID NO: 2) of a native sequence PRO245 polypeptide as derived from the coding sequence of Figure 3. Also shown are the approximate locations of various other important polypeptide domains if known.

Figure 5 shows the nucleotide sequence of a cDNA containing a nucleotide sequence encoding native sequence PRO217(UNQ191), wherein the nucleotide sequence (SEQ ID NO: 3) is a clone designated herein as "DNA33094". Also presented in bold font and underlined are the positions of the respective start and stop codons.

Figure 6 and Table 7 show the amino acid sequence (SEQ ID NO: 4) of a native sequence PRO217 polypeptide as derived from the coding sequence of Figure 5. Also shown are the approximate locations of various other important polypeptide domains if known.

Figure 7 shows the nucleotide sequence of a cDNA containing a nucleotide sequence encoding native sequence PRO301(UNQ264), wherein the nucleotide sequence (SEQ ID NO: 5) is a clone designated herein as "DNA40628". Also presented in bold font and underlined are the positions of the respective start and stop codons.

Figure 8 and Table 8 show the amino acid sequence (SEQ ID NO: 6) of a native sequence PRO301 polypeptide as derived from the coding sequence of Figure 7. Also shown are the approximate locations of various other important polypeptide domains if known.

Figure 9 shows the nucleotide sequence of a cDNA containing a nucleotide sequence encoding native sequence PRO266 (UNQ233), wherein the nucleotide sequence (SEQ ID NO: 7) is a clone designated herein as "DNA37150". Also presented in bold font and underlined are the positions of the respective start and stop codons.

Figure 10 and Table 9 show the amino acid sequence (SEQ ID NO: 8) of a native sequence PRO266 polypeptide as derived from the coding sequence of Figure 9. Also shown are the approximate locations of various other important polypeptide domains if known.

Figure 11 shows the nucleotide sequence of a cDNA containing a nucleotide sequence encoding native sequence PRO335 (UNQ287V), wherein the nucleotide sequence (SEQ ID NO: 9) is a clone designated herein as "DNA41388". Also presented in bold font and underlined are the positions of the respective start and stop codons.

Figure 12 and Table 10 show the amino acid sequence (SEQ ID NO: 10) of a native sequence PRO335 polypeptide as derived from the coding sequence of Figure 11. Also shown are the approximate locations of various other important polypeptide domains if known.

Figure 13 shows the nucleotide sequence of a cDNA containing a nucleotide sequence encoding native sequence PRO331 (UNQ292), wherein the nucleotide sequence (SEQ ID NO: 11) is a

clone designated herein as "DNA40981". Also presented in bold font and underlined are the positions of the respective start and stop codons.

Figure 14 and Table 11 show the amino acid sequence (SEQ ID NO: 12) of a native sequence PRO331 polypeptide as derived from the coding sequence of Figure 13. Also shown are the approximate locations of various other important polypeptide domains if known.

Figure 15 shows the nucleotide sequence of a cDNA containing a nucleotide sequence encoding native sequence PRO326 (UNQ287), wherein the nucleotide sequence (SEQ ID NO: 13) is a clone designated herein as "DNA37140". Also presented in bold font and underlined are the positions of the respective start and stop codons.

Figure 16 and Table 12 show the amino acid sequence (SEQ ID NO: 14) of a native sequence PRO331 polypeptide as derived from the coding sequence of Figure 15. Also shown are the approximate locations of various other important polypeptide domains if known.

### Detailed Description of the Preferred Embodiments

#### I. Definitions

The term "immune related disease" means a disease in which a component of the immune system of a mammal causes, mediates or otherwise contributes to a morbidity in the mammal. Also included are diseases in which stimulation or intervention of the immune response has an ameliorative effect on progression of the disease. Included within this term are immune-mediated inflammatory diseases, non-immune-mediated inflammatory diseases, infectious diseases, immunodeficiency diseases, neoplasia, etc.

The term "T cell mediated" disease means a disease in which T cells directly or indirectly mediate or otherwise contribute to a morbidity in a mammal. The T cell mediated disease may be associated with cell mediated effects, lymphokine mediated effects, etc., and even effects associated with B cells if the B cells are stimulated, for example, by the lymphokines secreted by T cells.

Examples of immune-related and inflammatory diseases, some of which are immune or T cell mediated, which can be treated according to the invention include systemic lupus erythematosus, rheumatoid arthritis, juvenile chronic arthritis, spondyloarthropathies, systemic sclerosis (scleroderma), idiopathic inflammatory myopathies (dermatomyositis, polymyositis), Sjsgren's syndrome, systemic vasculitis, sarcoidosis, autoimmune hemolytic anemia (immune pancytopenia, paroxysmal nocturnal hemoglobinuria), autoimmune thrombocytopenia (idiopathic thrombocytopenic purpura, immune-mediated thrombocytopenia), thyroiditis (Grave's disease, Hashimoto's thyroiditis, juvenile lymphocytic thyroiditis, atrophic thyroiditis), diabetes mellitus, immune-mediated renal disease (glomerulonephritis, tubulointerstitial nephritis), demyelinating diseases of the central and peripheral nervous systems such as multiple sclerosis, idiopathic demyelinating polyneuropathy or

Guillain-Barré syndrome, and chronic inflammatory demyelinating polyneuropathy, hepatobiliary diseases such as infectious hepatitis (hepatitis A, B, C, D, E and other non-hepatotropic viruses), autoimmune chronic active hepatitis, primary biliary cirrhosis, granulomatous hepatitis, and sclerosing cholangitis, inflammatory and fibrotic lung diseases such as inflammatory bowel disease (ulcerative colitis: Crohn's disease), gluten-sensitive enteropathy, and Whipple's disease, autoimmune or immune-mediated skin diseases including bullous skin diseases, erythema multiforme and contact dermatitis, psoriasis, allergic diseases such as asthma, allergic rhinitis, atopic dermatitis, food hypersensitivity and urticaria, immunologic diseases of the lung such as eosinophilic pneumonias, idiopathic pulmonary fibrosis and hypersensitivity pneumonitis, transplantation associated diseases including graft rejection and graft -versus-host-disease. Infectious diseases include AIDS (HIV infection), hepatitis A, B, C, D, and E, bacterial infections, fungal infections, protozoal infections and parasitic infections.

"Treatment" is an intervention performed with the intention of preventing the development or altering the pathology of a disorder. Accordingly, "treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented. In treatment of an immune related disease, a therapeutic agent may directly decrease or increase the magnitude of response of a component of the immune response, or render the disease more susceptible to treatment by other therapeutic agents, e.g. antibiotics, antifungals, anti-inflammatory agents, chemotherapeutics, etc.

The "pathology" of an immune related disease includes all phenomena that compromise the well-being of the patient. This includes, without limitation, abnormal or uncontrollable cell growth (neutrophilic, eosinophilic, monocytic, lymphocytic cells), antibody production, auto-antibody production, complement production, interference with the normal functioning of neighboring cells, release of cytokines or other secretory products at abnormal levels, suppression or aggravation of any inflammatory or immunological response, infiltration of inflammatory cells (neutrophilic, eosinophilic, monocytic, lymphocytic) into cellular spaces, etc.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. "Intermittent" administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

"Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, polyethylene glycol (PEG), and PLURONICS™.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g. I<sup>131</sup>, I<sup>125</sup>, Y<sup>90</sup> and Re<sup>186</sup>), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant or animal origin, or fragments thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include adriamycin, doxorubicin, epirubicin, 5-fluorouracil, cytosine arabinoside ("Ara-C"), cyclophosphamide, thiopeta, busulfan, cytoxin, taxoids, e.g. paclitaxel (Taxol, Bristol-Myers Squibb Oncology, Princeton, NJ), and doxetaxel (Taxotere, Rhone-Poulenc Rorer, Antony, France), taxotere, methotrexate, cisplatin, melphalan, vinblastine, bleomycin, etoposide, ifosfamide, mitomycin C, mitoxantrone, vincristine, vinorelbine, carboplatin, teniposide, daunomycin, carminomycin, aminopterin, dactinomycin, mitomycins, esperamicins (see U.S. Pat. No. 4,675,187), melphalan and other related nitrogen mustards. Also included in this definition are hormonal agents that act to regulate or inhibit hormone action on tumors such as tamoxifen and onapristone.

A "growth inhibitory agent" when used herein refers to a compound or composition which inhibits growth of a cell, especially cancer cell overexpressing any of the genes identified herein, either *in vitro* or *in vivo*. Thus, the growth inhibitory agent is one which significantly reduces the percentage of cells overexpressing such genes in S phase. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxol, and topo II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in *The Molecular Basis of Cancer*, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogenes, and

antineoplastic drugs" by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13.

The term "cytokine" is a generic term for proteins released by one cell population which act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor-alpha and -beta; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors such as NGF-beta; platelet-growth factor; transforming growth factors (TGFs) such as TGF-alpha and TGF-beta; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon-alpha, -beta, and -gamma; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-11, IL-12; a tumor necrosis factor such as TNF-alpha or TNF-beta; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

As used herein, a "PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide" refers to a native sequence PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 having the same amino acid sequence as a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 derived from nature. Such native sequence PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can be isolated from nature or can be produced by recombinant and/or synthetic means. The term specifically encompasses naturally-occurring truncated or secreted forms (*e.g.*, an extracellular domain sequence), naturally-occurring variant forms (*e.g.*, alternatively spliced forms) and naturally-occurring allelic variants of the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326. In one embodiment of the invention, the native sequence PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 is a mature or full-length native sequence PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 comprising the amino acid sequences shown in Figures 4, 6, 8, 10, 12, 14 and 16.

The term "polypeptide of the invention" refers to each individual PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide. All disclosures in this specification which refer to the "polypeptide of the invention" or to "the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide" refer to each of the polypeptides individually as well as jointly. For example, descriptions of the preparation of, purification of, derivation of, formation of antibodies to or against, administration of, compositions containing, treatment of a disease with, etc.,

pertain to each polypeptide of the invention individually. The term "compound of the invention" includes the polypeptide of the invention, as well as agonist antibodies for and antagonist antibodies to these polypeptide, peptides or small molecules having agonist or antagonist activity developed from the polypeptide, etc.

The term "polypeptide of the invention" also includes variants of the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptides. A "variant" polypeptide means an active polypeptide as defined below having at least about 80% amino acid sequence identity with the amino acid sequence of the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptides. Such variant polypeptides include, for instance, polypeptides wherein one or more amino acid residues are added, or deleted, at the N- and/or C-terminus, as well as within one or more internal domains. Ordinarily, a variant polypeptide will have at least about 80% amino acid sequence identity, more preferably at least about 81% amino acid sequence identity, more preferably at least about 82% amino acid sequence identity, more preferably at least about 83% amino acid sequence identity, more preferably at least about 84% amino acid sequence identity, more preferably at least about 85% amino acid sequence identity, more preferably at least about 86% amino acid sequence identity, more preferably at least about 87% amino acid sequence identity, more preferably at least about 88% amino acid sequence identity, more preferably at least about 89% amino acid sequence identity, more preferably at least about 90% amino acid sequence identity, more preferably at least about 91% amino acid sequence identity, more preferably at least about 92% amino acid sequence identity, more preferably at least about 93% amino acid sequence identity, more preferably at least about 94% amino acid sequence identity, more preferably at least about 95% amino acid sequence identity, more preferably at least about 96% amino acid sequence identity, more preferably at least about 97% amino acid sequence identity, more preferably at least about 98% amino acid sequence identity and yet more preferably at least about 99% amino acid sequence identity with the amino acid sequence of the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptides. Variants do not encompass the native polypeptide sequence.

Ordinarily, variant polypeptides of the invention are at least about 10 amino acids in length, often at least about 20 amino acids in length, more often at least about 30 amino acids in length, more often at least about 40 amino acids in length, more often at least about 50 amino acids in length, more often at least about 60 amino acids in length, more often at least about 70 amino acids in length, more often at least about 80 amino acids in length, more often at least about 90 amino acids in length, more often at least about 100 amino acids in length, more often at least about 150 amino acids in length, more often at least about 200 amino acids in length, more often at least about 300 amino acids in length, or more.

"Percent (%) amino acid sequence identity" with respect to the polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are

identical with the amino acid residues in a sequence of the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptides, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Figures 2A-P. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Figures 2A-P has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Figures 2A-P. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

For purposes herein, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations, Figures 1A-B demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO".

Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-

BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

Also included within the term "polypeptides of the invention" are polypeptides which in the context of the amino acid sequence identity comparisons performed as described above, include amino acid residues in the sequences compared that are not only identical, but also those that have similar properties. These polypeptides are termed "positives". Amino acid residues that score a positive value to an amino acid residue of interest are those that are either identical to the amino acid residue of interest or are a preferred substitution (as defined in Table 1 below) of the amino acid residue of interest. For purposes herein, the % value of positives of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % positives to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scoring a positive value as defined above by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % positives of A to B will not equal the % positives of B to A.

An "invention variant polynucleotide" or "invention variant nucleic acid sequence" means a nucleic acid molecule which encodes an active polypeptide of the invention as defined herein and which has at least about 80% nucleic acid sequence identity with the nucleotide acid sequence of DNA35638, DNA33094, DNA40628, DNA37150, DNA41388, DNA40981, or DNA37140 or a specifically derived fragment thereof. Ordinarily, an invention variant polynucleotide will have at least about 80% nucleic acid sequence identity, more preferably at least about 81% nucleic acid sequence identity, more preferably at least about 82% nucleic acid sequence identity, more preferably at least about 83% nucleic acid sequence identity, more preferably at least about 84% nucleic acid sequence identity, more preferably at least about 85% nucleic acid sequence identity, more preferably at least about 86% nucleic acid sequence identity, more preferably at least about 87% nucleic acid sequence identity, more preferably at least about 88% nucleic acid sequence identity, more preferably at least about 89% nucleic acid sequence identity, more preferably at least about 90% nucleic acid sequence identity, more preferably at least about 91% nucleic acid sequence identity, more preferably at least about 92% nucleic acid sequence identity, more preferably at least about 93% nucleic acid sequence identity, more preferably at least about 94% nucleic acid sequence identity, more preferably at least about 95% nucleic acid sequence identity, more preferably at least about 96% nucleic acid sequence identity, more preferably at least about 97% nucleic acid sequence identity, more preferably at least about 98% nucleic acid sequence identity and yet more preferably at least about 99% nucleic acid sequence identity with the nucleic acid sequence of DNA35638, DNA33094, DNA40628, DNA37150, DNA41388, DNA40981, or DNA37140 or a derived fragment thereof. Variants do not encompass the native nucleotide sequence. In this regard, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of invention variant polynucleotides having at least about 80% nucleic acid sequence identity to nucleotides DNA35638, DNA33094, DNA40628, DNA37150, DNA41388, DNA40981, or DNA37140 will encode a polypeptide having an amino acid sequence which is identical to the amino acid sequence of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 or the amino acid sequence encoded by the clones deposited with the ATCC described below.

Ordinarily, invention variant polynucleotides are at least about 30 nucleotides in length, often at least about 60 nucleotides in length, more often at least about 90 nucleotides in length, more often at least about 120 nucleotides in length, more often at least about 150 nucleotides in length, more often at least about 180 nucleotides in length, more often at least about 210 nucleotides in length, more often at least about 240 nucleotides in length, more often at least about 270 nucleotides in length, more often at least about 300 nucleotides in length, more often at least about 450 nucleotides in length, more often at least about 600 nucleotides in length, more often at least about 900 nucleotides in length, or more.

"Percent (%) nucleic acid sequence identity" with respect to the invention polypeptide-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in an invention polypeptide-encoding sequence of interest, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % nucleic acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Figures 2A-P. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Figures 2A-P has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Figures 2A-P. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

For purposes herein, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Figures 1C-D demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA".

Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % nucleic acid sequence identity may also be determined using the sequence comparison

program NCBI-BLAST2 (Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

In other embodiments, invention variant polynucleotides are nucleic acid molecules that encode an active polypeptide of the invention and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding the full-length invention polypeptide. Invention variant polypeptides include those that are encoded by an invention variant polynucleotide.

An "isolated" nucleic acid molecule encoding a polypeptide of the invention is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide-encoding nucleic acid. An isolated polypeptide-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from the polypeptide-encoding nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule encoding a polypeptide of the invention includes polypeptide-encoding nucleic acid molecules contained in cells that ordinarily express a polypeptide of the invention where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a

ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel *et al.*, Current Protocols in Molecular Biology, Wiley Interscience Publishers, (1995).

"Stringent conditions" or "high stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 ug/ml), 0.1% SDS, and 10% dextran sulfate at 42C, with washes at 42C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55C.

"Moderately stringent conditions" may be identified as described by Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS)

less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/mL denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising a polypeptide of the invention fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

"Active" or "activity" in the context of variants of the polypeptide of the invention refers to form(s) of proteins of the invention which retain the biologic and/or immunologic activities of a native or naturally-occurring polypeptide of the invention.

"Biological activity" in the context of an antibody or another molecule that can be identified by the screening assays disclosed herein (e.g. an organic or inorganic small molecule, peptide, etc.) is used to refer to the ability of such molecules to induce or inhibit infiltration of inflammatory cells into a tissue, to stimulate or inhibit T-cell proliferation and to stimulate or inhibit lymphokine release by cells. Another preferred activity is increased vascular permeability or the inhibition thereof.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native polypeptide of the invention disclosed herein. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native polypeptide of the invention disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native polypeptides of the invention, peptides, small organic molecules, etc.

A "small molecule" is defined herein to have a molecular weight below about 600 daltons.

"Antibodies" (Abs) and "immunoglobulins" (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific antigen, immunoglobulins include both antibodies and other antibody-like molecules which lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas. The term "antibody" is used in the broadest sense and specifically covers, without limitation, intact monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g. bispecific antibodies) formed from at least two intact antibodies, and antibody

fragments so long as they exhibit the desired biological activity. An anti- PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody is an antibody which immunologically binds to a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide. The antibody may bind to any domain of the polypeptide of the invention which may be contacted by the antibody. For example, the antibody may bind to any extracellular domain of the polypeptide and when the entire polypeptide is secreted, to any domain on the polypeptide which is available to the antibody for binding.

"Native antibodies" and "native immunoglobulins" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain ( $V_H$ ) followed by a number of constant domains. Each light chain has a variable domain at one end ( $V_L$ ) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light- and heavy-chain variable domains.

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity-determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a beta-sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the beta-sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, *NIH Publ. No. 91-3242*, Vol. I, pages 647-669 (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

The term "hypervariable region" or "complementarity-determining regions" (CDRs) as used herein define a subregion within the variable region of extreme sequence variability of the antibody, which form the antigen-binding site and are the main determinants of antigen specificity. According to one definition, they can be, for example, residues (Kabat nomenclature) 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable region and residues (Kabat nomenclature) 31-35 (H1), 50-65

(H2), 95-102 (H3) in the heavy chain variable region. Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institute of Health, Bethesda, MD. [1991]. Alternatively, or in combination with the region defined by Kabat, the hypervariable region can be the "hypervariable loop", comprising, for example, residues (Chothia nomenclature) 26-32 (L1), 50-53 (L2), 91-96 (L3) in the light chain variable region and residue (Chothia nomenclature) 26-32 (H1), 53-55 (L2) and 96-101 (L3); Chothia and Lesk, *J. Mol. Biol.* 196: 901-917 [1987]. "Framework" or "FR" residues are those variable domain residues of relatively low sequence variability which lie in between the CDR regions.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments; diabodies; linear antibodies (Zapata *et al.*, *Protein Eng.* 8(10):1057-1062 [1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose PRO245, PRO217, PRO301, Pro266, pro335, pro331 or pro326 reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')<sub>2</sub> fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V<sub>H</sub>-V<sub>L</sub> dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')<sub>2</sub> antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains,

immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), *e.g.*, IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of immunoglobulins are called  $\alpha$ ,  $\delta$ ,  $\epsilon$ ,  $\gamma$ , and  $\mu$ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, Nature, 256:495 [1975], or may be made by recombinant DNA methods (see, *e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, Nature, 352:624-628 [1991] and Marks *et al.*, J. Mol. Biol., 222:581-597 (1991), for example. See also U.S. Patent Nos. 5,750,373, 5,571,698, 5,403,484 and 5,223,409 which describe the preparation of antibodies using phagemid and phage vectors.

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; Morrison *et al.*, Proc. Natl. Acad. Sci. USA, 81:6851-6855 [1984]).

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human

immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementarity-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and maximize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, Nature, 321:522-525 (1986); Reichmann *et al.*, Nature, 332:323-329 [1988]; and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992). The term "humanized antibody" includes a "primatized" antibody where the antigen-binding region of the antibody is derived from an antibody produced by immunizing macaque monkeys with the antigen of interest. Antibodies containing residues from Old World monkeys are also possible within the invention. See, for example, U.S. Patent Nos. 5,658,570; 5,693,780; 5,681,722; 5,750,105; and 5,756,096.

Antibodies and fragments thereof in this invention also include "affinity matured" antibodies in which an antibody is altered to change the amino acid sequence of one or more of the CDR regions and/or the framework regions to alter the affinity of the antibody or fragment thereof for the antigen to which it binds. Affinity maturation may result in an increase or in a decrease in the affinity of the matured antibody for the antigen relative to the starting antibody. Typically, the starting antibody will be a humanized, human, chimeric or murine antibody and the affinity matured antibody will have a higher affinity than the starting antibody. During the maturation process, one or more of the amino acid residues in the CDRs or in the framework regions are changed to a different residue using any standard method. Suitable methods include point mutations using well known cassette mutagenesis methods (Wells *et al.*, 1985, Gene, 34:315) or oligonucleotide mediated mutagenesis methods (Zoller *et al.*, 1987, Nucleic Acids Res., 10:6487-6504). Affinity maturation may also be performed using known selection methods in which many mutations are produced and mutants having the desired affinity are selected from a pool or library of mutants based on improved affinity for the antigen or ligand. Known phage display techniques can be conveniently used in this approach. See, for example, U.S. 5,750,373; U.S. 5,223,409, etc.

Human antibodies are also within the scope of the antibodies of the invention. Human antibodies can be produced using various techniques known in the art, including phage display

libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985); Boerner et al., *J. Immunol.*, 147(1):86-95 (1991); U. S. 5,750, 373]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks *et al.*, *Bio/Technology* 10, 779-783 (1992); Lonberg *et al.*, *Nature* 368 856-859 (1994); Morrison, *Nature* 368, 812-13 (1994); Fishwild *et al.*, *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

"Single-chain Fv" or "sFv" antibody fragments comprise the V<sub>H</sub> and V<sub>L</sub> domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V<sub>H</sub> and V<sub>L</sub> domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V<sub>H</sub>) connected to a light-chain variable domain (V<sub>L</sub>) in the same polypeptide chain (V<sub>H</sub> - V<sub>L</sub>). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the compound of the invention will be purified (1) to greater than 95% by weight of the compound as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated compound, e.g. antibody or

polypeptide, includes the compound *in situ* within recombinant cells since at least one component of the compound's natural environment will not be present. Ordinarily, however, isolated compound will be prepared by at least one purification step.

The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the compound, e.g. antibody or polypeptide, so as to generate a "labelled" compound. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

By "solid phase" is meant a non-aqueous matrix to which the compound of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Patent No. 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as the anti-ErbB2 antibodies disclosed herein and, optionally, a chemotherapeutic agent) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

## II. Compositions and Methods of the Invention

### 1. Preparation of the polypeptides of the invention

The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 (UNQ219, UNQ191, UNQ264, UNQ233, UNQ287V, UNQ292 or UNQ287 respectively). In particular, cDNA encoding a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide has been identified and isolated, as disclosed in further detail in the

Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by DNA35638, DNA33094, DNA40628, DNA37150, DNA41388, DNA40981 and DNA37140 as well as all further native homologues and variants included in the foregoing definition of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326, will be referred to as PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 or simply as "the polypeptide of the invention", regardless of their origin or mode of preparation.

The description below relates primarily to production of the polypeptide of the invention by culturing cells transformed or transfected with a vector containing nucleic acid which encodes of the polypeptide of the invention. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare of the polypeptide of the invention. For instance, the polypeptide sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., Solid-Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, J. Am. Chem. Soc., 85:2149-2154 (1963)]. *In vitro* protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the polypeptide of the invention may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length polypeptide.

In addition to the full-length native sequence polypeptides described herein, it is contemplated that variants can be prepared. Variants can be prepared by introducing appropriate nucleotide changes into the DNA, and/or by synthesis of the desired polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the polypeptide of the invention, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the native full-length sequence or in various domains of the polypeptide of the invention described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the polypeptide that results in a change in the amino acid sequence of the polypeptide as compared with the native sequence polypeptide sequence. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the polypeptide of the invention. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the polypeptide of the invention with that of homologous known protein molecules and minimizing the

number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

Polypeptide fragments of the polypeptides of the invention are also within the scope of the invention. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the invention polypeptide.

Polypeptide fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR. Preferably, polypeptide fragments share at least one biological and/or immunological activity with the native invention polypeptide.

In particular embodiments, conservative substitutions of interest are shown in Table 1 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 1, or as further described below in reference to amino acid classes, are introduced and the products screened.

Table 1

<u>Original Residue</u>	<u>Exemplary Substitutions</u>	<u>Preferred Substitutions</u>
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro; ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in function or immunological identity of the invention polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

(1) hydrophobic: norleucine, met, ala, val, leu, ile;

- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., Nucl. Acids Res., 13:4331 (1986); Zoller et al., Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells et al., Gene, 34:315 (1985)], restriction selection mutagenesis [Wells et al., Philos. Trans. R. Soc. London SerA, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

Covalent modifications of polypeptides of the invention are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of an invention polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the polypeptide. Derivatization with bifunctional agents is useful, for instance, for crosslinking the invention polypeptide to a water-insoluble support matrix or surface for use in the method for purifying anti-polypeptide antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimidate.

Other modifications include deamidation of glutamyl and asparagyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the invention polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence polypeptide (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Addition of glycosylation sites to the polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence polypeptide (for O-linked glycosylation sites). The amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the polypeptide of the invention is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

Removal of carbohydrate moieties present on the polypeptide of the invention may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., Arch. Biochem. Biophys., 259:52 (1987) and by Edge et al., Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., Meth. Enzymol., 138:350 (1987).

Another type of covalent modification comprises linking the invention polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The polypeptide of the present invention may also be modified in a way to form a chimeric molecule comprising the invention polypeptide fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the invention polypeptide with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the polypeptide of the invention. The presence of such epitope-tagged forms of the polypeptide of the invention can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the polypeptide of the invention to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., *Mol. Cell. Biol.*, **8**:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., *Molecular and Cellular Biology*, **5**:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., *Protein Engineering*, **3**(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., *BioTechnology*, **6**:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., *Science*, **255**:192-194 (1992)]; an  $\alpha$ -tubulin epitope peptide [Skinner et al., *J. Biol. Chem.*, **266**:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., *Proc. Natl. Acad. Sci. USA*, **87**:6393-6397 (1990)].

In an alternative embodiment, the chimeric molecule may comprise a fusion of the polypeptide of the invention with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of an invention polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also US Patent No. 5,428,130 issued June 27, 1995.

i. Isolation of DNA Encoding the Polypeptide of the Invention

DNA encoding the polypeptide of the invention may be obtained from a cDNA library prepared from tissue believed to possess the polypeptide mRNA and to express it at a detectable level. Accordingly, human DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The gene encoding the polypeptide of the invention may also be obtained from a genomic library or by oligonucleotide synthesis.

Libraries can be screened with probes (such as antibodies to the polypeptide of the invention

or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding the polypeptide of the invention is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., PCR Primer: A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1995)].

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like  $^{32}\text{P}$ -labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined through sequence alignment using computer software programs such as ALIGN, DNASTAR, and INHERIT which employ various algorithms to measure homology.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

#### ii. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for production of the polypeptides of the invention and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

Methods of transfection are known to the ordinarily skilled artisan, for example, CaPO<sub>4</sub> and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transformations have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact., 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635).

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for vectors encoding the polypeptides of the invention. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism.

Suitable host cells for the expression of glycosylated polypeptides of the invention are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen. Virol., 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

iii. Selection and Use of a Replicable Vector

The nucleic acid (*e.g.*, cDNA or genomic DNA) encoding the polypeptides of the invention may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, phagemid or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

The polypeptide of the invention may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the DNA encoding the polypeptide of the invention that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, *lpp*, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, *e.g.*, the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* alpha-factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2u plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, *e.g.*, the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the nucleic acid encoding the polypeptide of the invention,

such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., Nature, 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics, 85:12 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the nucleic acid sequence encoding the polypeptide of the invention to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the beta-lactamase and lactose promoter systems [Chang et al., Nature, 275:615 (1978); Goeddel et al., Nature, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, Nucleic Acids Res., 8:4057 (1980); EP 36,776], and hybrid promoters such as the *tac* promoter [deBoer et al., Proc. Natl. Acad. Sci. USA, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the polypeptide of the invention.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., J. Biol. Chem., 255:2073 (1980)] or other glycolytic enzymes [Hess et al., J. Adv. Enzyme Reg., 7:149 (1968); Holland, Biochemistry, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

Transcription of the polypeptide of the invention from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the polypeptide of the invention by higher eukaryotes may

be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, alpha-fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the coding sequence of the polypeptide of the invention, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding the polypeptide of the invention .

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of the polypeptide of the invention in recombinant vertebrate cell culture are described in Gething et al., Nature, 293:620-625 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

### iii. Detecting Gene Expression

Gene expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence of the inventive polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to DNA encoding the polypeptide of the invention and encoding a specific antibody epitope.

### iv. Purification of Polypeptide

Forms of the polypeptide of the invention may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (*e.g.* Triton-X 100) or by enzymatic cleavage. Cells employed in expression of the polypeptide of the invention can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It may be desired to purify the polypeptide of the invention from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the polypeptide of the invention. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular polypeptide of the invention produced.

## 2. Tissue Distribution

The location of tissues expressing the polypeptides of the invention can be identified by determining mRNA expression in various human tissues. The location of such genes provides information about which tissues are most likely to be affected by the stimulating and inhibiting activities of the polypeptides of the invention. The location of a gene in a specific tissue also provides sample tissue for the activity blocking assays discussed below.

As noted before, gene expression in various tissues may be measured by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 [1980]), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes.

Gene expression in various tissues, alternatively, may be measured by immunological methods, such as immunohistochemical staining of tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence of a polypeptide of the invention or against a synthetic peptide based on the DNA sequences encoding the polypeptide of the invention or against an exogenous sequence fused to a DNA encoding a polypeptide of the invention and encoding a specific antibody epitope. General techniques for

generating antibodies, and special protocols for Northern blotting and *in situ* hybridization are provided below.

### 3. Antibody Binding Studies

The activity of the polypeptides of the invention can be further verified by antibody binding studies, in which the ability of anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibodies to inhibit the effect of the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptides on tissue cells is tested. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies, the preparation of which will be described hereinbelow.

Antibody binding studies may be carried out in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, *Monoclonal Antibodies: A Manual of Techniques*, pp.147-158 (CRC Press, Inc., 1987).

Competitive binding assays rely on the ability of a labeled standard to compete with the test sample analyte for binding with a limited amount of antibody. The amount of target protein in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies preferably are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte which remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex. See, *e.g.*, U.S. Pat No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

For immunohistochemistry, the tissue sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin, for example.

### 4. Cell-Based Assays

Cell-based assays and animal models for immune related diseases can be used to further understand the relationship between the genes and polypeptides identified herein and the development and pathogenesis of immune related disease.

In a different approach, cells of a cell type known to be involved in a particular immune related disease are transfected with the cDNAs described herein, and the ability of these cDNAs to stimulate or inhibit immune function is analyzed. Suitable cells can be transfected with the desired gene, and monitored for immune function activity. Such transfected cell lines can then be used to test

the ability of poly- or monoclonal antibodies or antibody compositions to inhibit or stimulate immune function, for example to modulate T-cell proliferation or inflammatory cell infiltration. Cells transfected with the coding sequences of the genes identified herein can further be used to identify drug candidates for the treatment of immune related diseases.

In addition, primary cultures derived from transgenic animals (as described below) can be used in the cell-based assays herein, although stable cell lines are preferred. Techniques to derive continuous cell lines from transgenic animals are well known in the art (see, e.g. Small *et al.*, Mol. Cell. Biol. 5, 642-648 [1985]).

One suitable cell based assay is the mixed lymphocyte reaction (MLR). Current Protocols in Immunology, unit 3.12; edited by J. E. Coligan, A. M. Kruisbeek, D. H. Marglies, E. M. Shevach, W. Strober, National Institutes of Health, Published by John Wiley & Sons, Inc. In this assay, the ability of a test compound to stimulate the proliferation of activated T cells is assayed. A suspension of responder T cells is cultured with allogeneic stimulator cells and the proliferation of T cells is measured by uptake of tritiated thymidine. This assay is a general measure of T cell reactivity. Since the majority of T cells respond to and produce IL-2 upon activation, differences in responsiveness in this assay in part reflect differences in IL-2 production by the responding cells. The MLR results can be verified by a standard lymphokine (IL-2) detection assay. Current Protocols in Immunology, above, 3.15, 6.3.

A proliferative T cell response in an MLR assay may be due to direct mitogenic properties of an assayed molecule or to external antigen induced activation. Additional verification of the T cell stimulatory activity of the polypeptides of the invention can be obtained by a costimulation assay. T cell activation requires an antigen specific signal mediated through the T-cell receptor (TCR) and a costimulatory signal mediated through a second ligand binding interaction, for example, the B7(CD80, CD86)/CD28 binding interaction. CD28 crosslinking increases lymphokine secretion by activated T cells. T cell activation has both negative and positive controls through the binding of ligands which have a negative or positive effect. CD28 and CTLA-4 are related glycoproteins in the Ig superfamily which bind to B7. CD28 binding to B7 has a positive costimulation effect of T cell activation; conversely, CTLA-4 binding to B7 has a negative T cell deactivating effect. Chambers, C. A. and Allison, J. P., *Curr. Opin. Immunol.* (1997) 9:396. Schwartz, R. H., *Cell* (1992) 71:1065; Linsey, P. S. and Ledbetter, J. A., *Annu. Rev. Immunol.* (1993) 11:191; June, C. H. et al., *Immunol. Today* (1994) 15:321; Jenkins, M. K., *Immunity* (1994) 1:405. In a costimulation assay, the polypeptides of the invention are assayed for T cell costimulatory or inhibitory activity.

Polypeptides of the invention, as well as other compounds of the invention, which are stimulators (costimulators) of T cell proliferation and agonists, e.g. agonist antibodies, thereto as determined by MLR and costimulation assays, for example, are useful in treating immune related diseases characterized by poor, suboptimal or inadequate immune function. These diseases are treated

by stimulating the proliferation and activation of T cells (and T cell mediated immunity) and enhancing the immune response in a mammal through administration of a stimulatory compound, such as the stimulating polypeptides of the invention. The stimulating polypeptide may, for example, be a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide or an agonist antibody therefor.

Direct use of a stimulating compound as in the invention has been validated in experiments with 4-1BB glycoprotein, a member of the tumor necrosis factor receptor family, which binds to a ligand (4-1BBL) expressed on primed T cells and signals T cell activation and growth. Alderson, M. E. et al., *J. Immunol.* (1994) 24:2219.

The use of an agonist stimulating compound has also been validated experimentally. Activation of 4-1BB by treatment with an agonist anti-4-1BB antibody enhances eradication of tumors. Hellstrom, I. and Hellstrom, K. E., *Crit. Rev. Immunol.* (1998) 18:1. Immunoadjuvant therapy for treatment of tumors, described in more detail below, is another example of the use of the stimulating compounds of the invention.

An immune stimulating or enhancing effect can also be achieved by antagonizing or blocking the activity of a protein which has been found to be inhibiting in the MLR assay. Negating the inhibitory activity of the compound produces a net stimulatory effect. Suitable antagonists/blocking compounds are antibodies or fragments thereof which recognize and bind to the inhibitory protein, thereby blocking the effective interaction of the protein with its receptor and inhibiting signaling through the receptor. This effect has been validated in experiments using anti-CTLA-4 antibodies which enhance T cell proliferation, presumably by removal of the inhibitory signal caused by CTLA-4 binding. Walunas, T. L. et al., *Immunity* (1994) 1:405.

On the other hand, polypeptides of the invention, as well as other compounds of the invention, which are direct inhibitors of T cell proliferation/activation and/or lymphokine secretion, can be directly used to suppress the immune response. These compounds are useful to reduce the degree of the immune response and to treat immune related diseases characterized by a hyperactive, superoptimal, or autoimmune response. This use of the compounds of the invention has been validated by the experiments described above in which CTLA-4 binding to receptor B7 deactivates T cells. The direct inhibitory compounds of the invention function in an analogous manner.

Alternatively, compounds, e.g. antibodies, which bind to stimulating polypeptides of the invention and block the stimulating effect of these molecules produce a net inhibitory effect and can be used to suppress the T cell mediated immune response by inhibiting T cell proliferation/activation and/or lymphokine secretion. Blocking the stimulating effect of the polypeptides suppresses the immune response of the mammal. This use has been validated in experiments using an anti-IL2 antibody. In these experiments, the antibody binds to IL2 and blocks binding of IL2 to its receptor thereby achieving a T cell inhibitory effect.

## 5. Animal Models

The results of the cell based *in vitro* assays can be further verified using *in vivo* animal models and assays for T-cell function. A variety of well known animal models can be used to further understand the role of the genes identified herein in the development and pathogenesis of immune related disease, and to test the efficacy of candidate therapeutic agents, including antibodies, and other antagonists of the native polypeptides, including small molecule antagonists. The *in vivo* nature of such models makes them predictive of responses in human patients. Animal models of immune related diseases include both non-recombinant and recombinant (transgenic) animals. Non-recombinant animal models include, for example, rodent, e.g., murine models. Such models can be generated by introducing cells into syngeneic mice using standard techniques, e.g. subcutaneous injection, tail vein injection, spleen implantation, intraperitoneal implantation, implantation under the renal capsule, etc.

Graft-versus-host disease occurs when immunocompetent cells are transplanted into immunosuppressed or tolerant patients. The donor cells recognize and respond to host antigens. The response can vary from life threatening severe inflammation to mild cases of diarrhea and weight loss. Graft-versus-host disease models provide a means of assessing T cell reactivity against MHC antigens and minor transplant antigens. A suitable procedure is described in detail in Current Protocols in Immunology, above, unit 4.3.

An animal model for skin allograft rejection is a means of testing the ability of T cells to mediate *in vivo* tissue destruction and a measure of their role in transplant rejection. The most common and accepted models use murine tail-skin grafts. Repeated experiments have shown that skin allograft rejection is mediated by T cells, helper T cells and killer-effector T cells, and not antibodies. Auchincloss, H. Jr. and Sachs, D. H., Fundamental Immunology, 2nd ed., W. E. Paul ed., Raven Press, NY, 1989, 889-992. A suitable procedure is described in detail in Current Protocols in Immunology, above, unit 4.4. Other transplant rejection models which can be used to test the compounds of the invention are the allogeneic heart transplant models described by Tanabe, M. et al., Transplantation (1994) 58:23 and Tinubu, S. A. et al., J. Immunol. (1994) 4330-4338.

Animal models for delayed type hypersensitivity provides an assay of cell mediated immune function as well. Delayed type hypersensitivity reactions are a T cell mediated *in vivo* immune response characterized by inflammation which does not reach a peak until after a period of time has elapsed after challenge with an antigen. These reactions also occur in tissue specific autoimmune diseases such as multiple sclerosis (MS) and experimental autoimmune encephalomyelitis (EAE, a model for MS). A suitable procedure is described in detail in Current Protocols in Immunology, above, unit 4.5.

EAE is a T cell mediated autoimmune disease characterized by T cell and mononuclear cell inflammation and subsequent demyelination of axons in the central nervous system. EAE is generally

considered to be a relevant animal model for MS in humans. Bolton, C., *Multiple Sclerosis* (1995) 1:143. Both acute and relapsing-remitting models have been developed. The compounds of the invention can be tested for T cell stimulatory or inhibitory activity against immune mediated demyelinating disease using the protocol described in *Current Protocols in Immunology*, above, units 15.1 and 15.2. See also the models for myelin disease in which oligodendrocytes or Schwann cells are grafted into the central nervous system as described in Duncan, I. D. et al, *Molec. Med. Today* (1997) 554-561.

Contact hypersensitivity is a simple delayed type hypersensitivity in vivo assay of cell mediated immune function. In this procedure, cutaneous exposure to exogenous haptens which gives rise to a delayed type hypersensitivity reaction which is measured and quantitated. Contact sensitivity involves an initial sensitizing phase followed by an elicitation phase. The elicitation phase occurs when the T lymphocytes encounter an antigen to which they have had previous contact. Swelling and inflammation occur, making this an excellent model of human allergic contact dermatitis. A suitable procedure is described in detail in *Current Protocols in Immunology*, Eds. J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach and W. Strober, John Wiley & Sons, Inc., 1994, unit 4.2. See also Grabbe, S. and Schwarz, T., *Immun. Today* 19(1):37-44 (1998) .

An animal model for arthritis is collagen-induced arthritis. This model shares clinical, histological and immunological characteristics of human autoimmune rheumatoid arthritis and is an acceptable model for human autoimmune arthritis. Mouse and rat models are characterized by synovitis, erosion of cartilage and subchondral bone. The compounds of the invention can be tested for activity against autoimmune arthritis using the protocols described in *Current Protocols in Immunology*, above, units 15.5. See also the model using a monoclonal antibody to CD18 and VLA-4 integrins described in Issekutz, A. C. et al., *Immunology* (1996) 88:569.

A model of asthma has been described in which antigen-induced airway hyper-reactivity, pulmonary eosinophilia and inflammation are induced by sensitizing an animal with ovalbumin and then challenging the animal with the same protein delivered by aerosol. Several animal models (guinea pig, rat, non-human primate) show symptoms similar to atopic asthma in humans upon challenge with aerosol antigens. Murine models have many of the features of human asthma. Suitable procedures to test the compounds of the invention for activity and effectiveness in the treatment of asthma are described by Wolyniec, W. W. et al., *Am. J. Respir. Cell Mol. Biol.* (1998) 18:777 and the references cited therein.

Additionally, the compounds of the invention can be tested on animal models for psoriasis like diseases. Evidence suggests a T cell pathogenesis for psoriasis. The compounds of the invention can be tested in the scid/scid mouse model described by Schon, M. P. et al., *Nat. Med.* (1997) 3:183, in which the mice demonstrate histopathologic skin lesions resembling psoriasis. Another suitable

model is the human skin/scid mouse chimera prepared as described by Nickoloff, B. J. et al., *Am. J. Path.* (1995) 146:580.

Recombinant (transgenic) animal models can be engineered by introducing the coding portion of the genes identified herein into the genome of animals of interest, using standard techniques for producing transgenic animals. Animals that can serve as a target for transgenic manipulation include, without limitation, mice, rats, rabbits, guinea pigs, sheep, goats, pigs, and non-human primates, e.g. baboons, chimpanzees and monkeys. Techniques known in the art to introduce a transgene into such animals include pronucleic microinjection (Hoppe and Wanger, U.S. Patent No. 4,873,191); retrovirus-mediated gene transfer into germ lines (e.g., Van der Putten *et al.*, *Proc. Natl. Acad. Sci. USA* 82, 6148-615 [1985]); gene targeting in embryonic stem cells (Thompson *et al.*, *Cell* 56, 313-321 [1989]); electroporation of embryos (Lo, *Mol. Cell. Biol.* 3, 1803-1814 [1983]); sperm-mediated gene transfer (Lavitrano *et al.*, *Cell* 57, 717-73 [1989]). For review, see, for example, U.S. Patent No. 4,736,866.

For the purpose of the present invention, transgenic animals include those that carry the transgene only in part of their cells ("mosaic animals"). The transgene can be integrated either as a single transgene, or in concatamers, e.g., head-to-head or head-to-tail tandems. Selective introduction of a transgene into a particular cell type is also possible by following, for example, the technique of Lasko *et al.*, *Proc. Natl. Acad. Sci. USA* 89, 6232-636 (1992).

The expression of the transgene in transgenic animals can be monitored by standard techniques. For example, Southern blot analysis or PCR amplification can be used to verify the integration of the transgene. The level of mRNA expression can then be analyzed using techniques such as *in situ* hybridization, Northern blot analysis, PCR, or immunocytochemistry.

The animals may be further examined for signs of immune disease pathology, for example by histological examination to determine infiltration of immune cells into specific tissues. Blocking experiments can also be performed in which the transgenic animals are treated with the compounds of the invention to determine the extent of the T cell proliferation stimulation or inhibition of the compounds. In these experiments, blocking antibodies which bind to the polypeptide of the invention, prepared as described above, are administered to the animal and the effect on immune function is determined.

Alternatively, "knock out" animals can be constructed which have a defective or altered gene encoding a polypeptide identified herein, as a result of homologous recombination between the endogenous gene encoding the polypeptide and altered genomic DNA encoding the same polypeptide introduced into an embryonic cell of the animal. For example, cDNA encoding a particular polypeptide can be used to clone genomic DNA encoding that polypeptide in accordance with established techniques. A portion of the genomic DNA encoding a particular polypeptide can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used

to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the polypeptide.

#### 6. ImmunoAdjuvant Therapy

In one embodiment, the immunostimulating compounds of the invention can be used in immunoadjuvant therapy for the treatment of tumors (cancer). It is now well established that T cells recognize human tumor specific antigens. One group of tumor antigens, encoded by the MAGE, BAGE and GAGE families of genes, are silent in all adult normal tissues, but are expressed in significant amounts in tumors, such as melanomas, lung tumors, head and neck tumors, and bladder carcinomas. DeSmet, C. et al, (1996) Proc. Natl. Acad. Sci. USA, 93:7149. It has been shown that costimulation of T cells induces tumor regression and an antitumor response both in vitro and in vivo. Melero, I. et al., Nature Medicine (1997) 3:682; Kwon, E. D. et al., Proc. Natl. Acad. Sci. USA (1997) 94:8099; Lynch, D. H. et al., Nature Medicine (1997) 3:625; Finn, O. J. and Lotze, M. T., J. Immunol. (1998) 21:114. The stimulatory compounds of the invention can be administered as adjuvants, alone or together with a growth regulating agent, cytotoxic agent or chemotherapeutic agent, to stimulate T cell proliferation/activation and an antitumor response to tumor antigens. The growth regulating, cytotoxic, or chemotherapeutic agent may be administered in conventional amounts using known administration regimes. Immunostimulating activity by the compounds of the invention allows reduced amounts of the growth regulating, cytotoxic, or chemotherapeutic agents thereby potentially lowering the toxicity to the patient.

#### 7. Screening Assays for Drug Candidates

Screening assays for drug candidates are designed to identify compounds that bind or complex with the polypeptides encoded by the genes identified herein or a biologically active fragment thereof, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. Such screening assays will include assays amenable to high-throughput screening of

chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds, including peptides, preferably soluble peptides, (poly)peptide-immunoglobulin fusions, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

All assays are common in that they call for contacting the drug candidate with a polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, e.g. on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the polypeptide and drying. Alternatively, an immobilized antibody, e.g. a monoclonal antibody, specific for the polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, e.g. the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g. by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labelled antibody specifically binding the immobilized complex.

If the candidate compound interacts with but does not bind to a particular protein encoded by a gene identified herein, its interaction with that protein can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers [Fields and Song, *Nature (London)* 340, 245-246 (1989); Chien *et al.*, *Proc. Natl. Acad. Sci. USA* 88, 9578-9582 (1991)] as disclosed by Chevray and Nathans [*Proc. Natl. Acad. Sci. USA* 89, 5789-5793 (1991)]. Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, while the other one functioning as the transcription activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of

this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for beta-galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

In order to find compounds that interfere with the interaction of a gene identified herein and other intra- or extracellular components can be tested, a reaction mixture is usually prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a test compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described above. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

#### 8. Compositions and Methods for the Treatment of Immune Related Diseases

The compositions useful in the treatment of immune related diseases include, without limitation, antibodies, small organic and inorganic molecules, peptides, phosphopeptides, antisense and ribozyme molecules, triple helix molecules, etc. that inhibit or stimulate immune function, for example, T cell proliferation/activation, lymphokine release, or immune cell infiltration.

For example, antisense RNA and RNA molecule act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation initiation site, e.g. between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, e.g. Rossi, *Current Biology* 4, 469-471 (1994), and PCT publication No. WO 97/33551 (published September 18, 1997).

Nucleic acid molecules in triple helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple helix formation via Hoogsteen base pairing rules, which

generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details see, e.g. PCT publication No. WO 97/33551, *supra*.

These molecules can be identified by any or any combination of the screening assays discussed above and/or by any other screening techniques well known for those skilled in the art.

#### 9. Antibodies

Some of the most promising drug candidates according to the present invention are antibodies and antibody fragments which may inhibit (antagonists) or stimulate (agonists) T cell proliferation, eosinophil infiltration, etc.

##### i. Polyclonal Antibodies

Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the polypeptide of the invention or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

##### ii. Monoclonal Antibodies

Antibodies which recognize and bind to the polypeptides of the invention or which act as agonist therefor may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

The immunizing agent will typically include the polypeptide of the invention or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture

medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Rockville, Maryland. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications*, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the polypeptide of the invention or having similar activity as the polypeptide of the invention. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, *Anal. Biochem.*, 107:220 (1980).

After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, *supra*]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as ascites in a mammal.

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxyapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise

produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Patent No. 4,816,567; Morrison et al., *supra*] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies are preferably monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

*In vitro* methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

### iii. Human and Humanized Antibodies

The antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art. Generally, a

humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and coworkers [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeyen et al., Science, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985); Boerner et al., J. Immunol., 147(1):86-95 (1991); U. S. 5,750, 373]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks *et al.*, Bio/Technology 10, 779-783 (1992); Lonberg *et al.*, Nature 368 856-859 (1994); Morrison, Nature 368, 812-13 (1994); Fishwild *et al.*, Nature Biotechnology 14, 845-51 (1996); Neuberger, Nature Biotechnology 14, 826 (1996); Lonberg and Huszar, Intern. Rev. Immunol. 13 65-93 (1995).

#### iv. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities may be for the polypeptide of the invention, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the coexpression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, Nature, 305:537-539 [1983]). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct

molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker et al., EMBO J., 10:3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are cotransfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology, 121:210 (1986).

v. Heteroconjugate Antibodies

Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Patent No. 4,676,980], and for treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

vi. Effector function engineering

It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance the effectiveness of the antibody in treating an immune related disease, for example. For example cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, J. Exp Med. 176:1191-1195 (1992) and Shopes, B. J. Immunol. 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.* Cancer Research 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.*, Anti-Cancer Drug Design 3:219-230 (1989).

vii. Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (*e.g.* an enzymatically active toxin of bacterial, fungal, plant or animal origin, or fragments thereof), or a radioactive isotope (*i.e.* a

radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$  and  $^{186}\text{Re}$ .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutarealdehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, Science 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyl-diethylene triamine-pentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tissue pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (*e.g.* avidin) which is conjugated to a cytotoxic agent (*e.g.* a radionucleotide).

#### viii. Immunoliposomes

The proteins, antibodies, etc. disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82:3688 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, 77:4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.*, J. Biol. Chem. 257: 286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent (such as doxorubicin) may be optionally

contained within the liposome. See Gabizon *et al.*, J. National Cancer Inst. 81(19)1484 (1989).

#### 10. Pharmaceutical Compositions

The active molecules of the invention, polypeptides and antibodies, as well as other molecules identified by the screening assays disclosed above, can be administered for the treatment of immune related diseases, in the form of pharmaceutical compositions.

Therapeutic formulations of the active molecule, preferably a polypeptide or antibody of the invention, are prepared for storage by mixing the active molecule having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. [1980]), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as TWEEN<sup>TM</sup>, PLURONICS<sup>TM</sup> or polyethylene glycol (PEG).

Compounds identified by the screening assays of the present invention can be formulated in an analogous manner, using standard techniques well known in the art.

Lipofections or liposomes can also be used to deliver the polypeptide, antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment which specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable region sequences of an antibody, peptide molecules can be designed which retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology (see, *e.g.* Marasco *et al.*, Proc. Natl. Acad. Sci. USA 90, 7889-7893 [1993]).

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise a cytotoxic agent, cytokine or growth inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active molecules may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980).

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.* films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

#### 11. Methods of Treatment

It is contemplated that the polypeptides, antibodies and other active compounds of the present invention may be used to treat various immune related diseases and conditions, such as T cell mediated diseases, including those characterized by infiltration of inflammatory cells into a tissue, stimulation of T-cell proliferation, inhibition of T-cell proliferation, increased or decreased vascular permeability or the inhibition thereof.

Exemplary conditions or disorders to be treated with the polypeptides, antibodies and other compounds of the invention, include, but are not limited to systemic lupus erythematosus, rheumatoid arthritis, juvenile chronic arthritis, osteoarthritis, spondyloarthropathies, systemic sclerosis (scleroderma), idiopathic inflammatory myopathies (dermatomyositis, polymyositis), Sjogren's syndrome, systemic vasculitis, sarcoidosis, autoimmune hemolytic anemia (immune pancytopenia,

paroxysmal nocturnal hemoglobinuria), autoimmune thrombocytopenia (idiopathic thrombocytopenic purpura, immune-mediated thrombocytopenia), thyroiditis (Grave's disease, Hashimoto's thyroiditis, juvenile lymphocytic thyroiditis, atrophic thyroiditis), diabetes mellitus, immune-mediated renal disease (glomerulonephritis, tubulointerstitial nephritis), demyelinating diseases of the central and peripheral nervous systems such as multiple sclerosis, idiopathic demyelinating polyneuropathy or Guillain-Barré syndrome, and chronic inflammatory demyelinating polyneuropathy, hepatobiliary diseases such as infectious hepatitis (hepatitis A, B, C, D, E and other non-hepatotropic viruses), autoimmune chronic active hepatitis, primary biliary cirrhosis, granulomatous hepatitis, and sclerosing cholangitis, inflammatory bowel disease (ulcerative colitis: Crohn's disease), gluten-sensitive enteropathy, and Whipple's disease, autoimmune or immune-mediated skin diseases including bullous skin diseases, erythema multiforme and contact dermatitis, psoriasis, allergic diseases such as asthma, allergic rhinitis, atopic dermatitis, food hypersensitivity and urticaria, immunologic diseases of the lung such as eosinophilic pneumonias, idiopathic pulmonary fibrosis and hypersensitivity pneumonitis, transplantation associated diseases including graft rejection and graft-versus-host-disease.

In systemic lupus erythematosus, the central mediator of disease is the production of auto-reactive antibodies to self proteins/tissues and the subsequent generation of immune-mediated inflammation. Antibodies either directly or indirectly mediate tissue injury. Though T lymphocytes have not been shown to be directly involved in tissue damage, T lymphocytes are required for the development of auto-reactive antibodies. The genesis of the disease is thus T lymphocyte dependent. Multiple organs and systems are affected clinically including kidney, lung, musculoskeletal system, mucocutaneous, eye, central nervous system, cardiovascular system, gastrointestinal tract, bone marrow and blood.

Rheumatoid arthritis (RA) is a chronic systemic autoimmune inflammatory disease that mainly involves the synovial membrane of multiple joints with resultant injury to the articular cartilage. The pathogenesis is T lymphocyte dependent and is associated with the production of rheumatoid factors, auto-antibodies directed against self IgG, with the resultant formation of immune complexes that attain high levels in joint fluid and blood. These complexes in the joint may induce the marked infiltrate of lymphocytes and monocytes into the synovium and subsequent marked synovial changes; the joint space/fluid is infiltrated by similar cells with the addition of numerous neutrophils. Tissues affected are primarily the joints, often in symmetrical pattern. However, extra-articular disease also occurs in two major forms. One form is the development of extra-articular lesions with ongoing progressive joint disease and typical lesions of pulmonary fibrosis, vasculitis, and cutaneous ulcers. The second form of extra-articular disease is the so called Felty's syndrome which occurs late in the RA disease course, sometimes after joint disease has become quiescent, and

involves the presence of neutropenia, thrombocytopenia and splenomegaly. This can be accompanied by vasculitis in multiple organs with formations of infarcts, skin ulcers and gangrene. Patients often also develop rheumatoid nodules in the subcutis tissue overlying affected joints; the nodules late stage have necrotic centers surrounded by a mixed inflammatory cell infiltrate. Other manifestations which can occur in RA include: pericarditis, pleuritis, coronary arteritis, interstitial pneumonitis with pulmonary fibrosis, keratoconjunctivitis sicca, and rheumatoid nodules.

Juvenile chronic arthritis is a chronic idiopathic inflammatory disease which begins often at less than 16 years of age. Its phenotype has some similarities to RA; some patients which are rheumatoid factor positive are classified as juvenile rheumatoid arthritis. The disease is sub-classified into three major categories: pauciarticular, polyarticular, and systemic. The arthritis can be severe and is typically destructive and leads to joint ankylosis and retarded growth. Other manifestations can include chronic anterior uveitis and systemic amyloidosis.

Spondyloarthropathies are a group of disorders with some common clinical features and the common association with the expression of HLA-B27 gene product. The disorders include: ankylosing spondylitis, Reiter's syndrome (reactive arthritis), arthritis associated with inflammatory bowel disease, spondylitis associated with psoriasis, juvenile onset spondyloarthropathy and undifferentiated spondyloarthropathy. Distinguishing features include sacroileitis with or without spondylitis; inflammatory asymmetric arthritis; association with HLA-B27 (a serologically defined allele of the HLA-B locus of class I MHC); ocular inflammation, and absence of autoantibodies associated with other rheumatoid disease. The cell most implicated as key to induction of the disease is the CD8+ T lymphocyte, a cell which targets antigen presented by class I MHC molecules. CD8+ T cells may react against the class I MHC allele HLA-B27 as if it were a foreign peptide expressed by MHC class I molecules. It has been hypothesized that an epitope of HLA-B27 may mimic a bacterial or other microbial antigenic epitope and thus induce a CD8+ T cells response.

Systemic sclerosis (scleroderma) has an unknown etiology. A hallmark of the disease is induration of the skin; likely this is induced by an active inflammatory process. Scleroderma can be localized or systemic; vascular lesions are common and endothelial cell injury in the microvasculature is an early and important event in the development of systemic sclerosis; the vascular injury may be immune mediated. An immunologic basis is implied by the presence of mononuclear cell infiltrates in the cutaneous lesions and the presence of anti-nuclear antibodies in many patients. ICAM-1 is often upregulated on the cell surface of fibroblasts in skin lesions suggesting that T cell interaction with these cells may have a role in the pathogenesis of the disease. Other organs involved include: the gastrointestinal tract: smooth muscle atrophy and fibrosis resulting in abnormal peristalsis/motility; kidney: concentric subendothelial intimal proliferation affecting small arcuate and interlobular arteries with resultant reduced renal cortical blood flow, results in proteinuria, azotemia and hypertension;

skeletal muscle: atrophy, interstitial fibrosis; inflammation; lung: interstitial pneumonitis and interstitial fibrosis; and heart: contraction band necrosis, scarring/fibrosis.

Idiopathic inflammatory myopathies including dermatomyositis, polymyositis and others are disorders of chronic muscle inflammation of unknown etiology resulting in muscle weakness. Muscle injury/inflammation is often symmetric and progressive. Autoantibodies are associated with most forms. These myositis-specific autoantibodies are directed against and inhibit the function of components, proteins and RNA's, involved in protein synthesis.

Sjogren's syndrome is due to immune-mediated inflammation and subsequent functional destruction of the tear glands and salivary glands. The disease can be associated with or accompanied by inflammatory connective tissue diseases. The disease is associated with autoantibody production against Ro and La antigens, both of which are small RNA-protein complexes. Lesions result in keratoconjunctivitis sicca, xerostomia, with other manifestations or associations including biliary cirrhosis, peripheral or sensory neuropathy, and palpable purpura.

Systemic vasculitis are diseases in which the primary lesion is inflammation and subsequent damage to blood vessels which results in ischemia/necrosis/degeneration to tissues supplied by the affected vessels and eventual end-organ dysfunction in some cases. Vasculitides can also occur as a secondary lesion or sequelae to other immune-inflammatory mediated diseases such as rheumatoid arthritis, systemic sclerosis, etc., particularly in diseases also associated with the formation of immune complexes. Diseases in the primary systemic vasculitis group include: systemic necrotizing vasculitis: polyarteritis nodosa, allergic angiitis and granulomatosis, polyangiitis; Wegener's granulomatosis; lymphomatoid granulomatosis; and giant cell arteritis. Miscellaneous vasculitides include: mucocutaneous lymph node syndrome (MLNS or Kawasaki's disease), isolated CNS vasculitis, Behet's disease, thromboangiitis obliterans (Buerger's disease) and cutaneous necrotizing vasculitis. The pathogenic mechanism of most of the types of vasculitis listed is believed to be primarily due to the deposition of immunoglobulin complexes in the vessel wall and subsequent induction of an inflammatory response either via ADCC, complement activation, or both.

Sarcoidosis is a condition of unknown etiology which is characterized by the presence of epithelioid granulomas in nearly any tissue in the body; involvement of the lung is most common. The pathogenesis involves the persistence of activated macrophages and lymphoid cells at sites of the disease with subsequent chronic sequelae resultant from the release of locally and systemically active products released by these cell types.

Autoimmune hemolytic anemia including autoimmune hemolytic anemia, immune pancytopenia, and paroxysmal nocturnal hemoglobinuria is a result of production of antibodies that react with antigens expressed on the surface of red blood cells (and in some cases other blood cells including platelets as well) and is a reflection of the removal of those antibody coated cells via complement mediated lysis and/or ADCC/Fc-receptor-mediated mechanisms.

In autoimmune thrombocytopenia including thrombocytopenic purpura, and immune-mediated thrombocytopenia in other clinical settings, platelet destruction/removal occurs as a result of either antibody or complement attaching to platelets and subsequent removal by complement lysis, ADCC or FC-receptor mediated mechanisms.

Thyroiditis including Grave's disease, Hashimoto's thyroiditis, juvenile lymphocytic thyroiditis, and atrophic thyroiditis, are the result of an autoimmune response against thyroid antigens with production of antibodies that react with proteins present in and often specific for the thyroid gland. Experimental models exist including spontaneous models: rats (BUF and BB rats) and chickens (obese chicken strain); inducible models: immunization of animals with either thyroglobulin, thyroid microsomal antigen (thyroid peroxidase).

Type I diabetes mellitus or insulin-dependent diabetes is the autoimmune destruction of pancreatic islet  $\beta$  cells; this destruction is mediated by auto-antibodies and auto-reactive T cells. Antibodies to insulin or the insulin receptor can also produce the phenotype of insulin-non-responsiveness.

Immune mediated renal diseases, including glomerulonephritis and tubulointerstitial nephritis, are the result of antibody or T lymphocyte mediated injury to renal tissue either directly as a result of the production of autoreactive antibodies or T cells against renal antigens or indirectly as a result of the deposition of antibodies and/or immune complexes in the kidney that are reactive against other, non-renal antigens. Thus other immune-mediated diseases that result in the formation of immune-complexes can also induce immune mediated renal disease as an indirect sequelae. Both direct and indirect immune mechanisms result in inflammatory response that produces/induces lesion development in renal tissues with resultant organ function impairment and in some cases progression to renal failure. Both humoral and cellular immune mechanisms can be involved in the pathogenesis of lesions.

Demyelinating diseases of the central and peripheral nervous systems, including Multiple Sclerosis; idiopathic demyelinating polyneuropathy or Guillain-Barr syndrome; and Chronic Inflammatory Demyelinating Polyneuropathy, are believed to have an autoimmune basis and result in nerve demyelination as a result of damage caused to oligodendrocytes or to myelin directly. In MS there is evidence to suggest that disease induction and progression is dependent on T lymphocytes. Multiple Sclerosis is a demyelinating disease that is T lymphocyte-dependent and has either a relapsing-remitting course or a chronic progressive course. The etiology is unknown; however, viral infections, genetic predisposition, environment, and autoimmunity all contribute. Lesions contain infiltrates of predominantly T lymphocyte mediated, microglial cells and infiltrating macrophages; CD4+T lymphocytes are the predominant cell type at lesions. The mechanism of oligodendrocyte cell death and subsequent demyelination is not known but is likely T lymphocyte driven.

Inflammatory and Fibrotic Lung Disease, including Eosinophilic Pneumonias; Idiopathic Pulmonary Fibrosis, and Hypersensitivity Pneumonitis may involve a dysregulated immune-inflammatory response. Inhibition of that response would be of therapeutic benefit.

Autoimmune or Immune-mediated Skin Disease including Bullous Skin Diseases, Erythema Multiforme, and Contact Dermatitis are mediated by auto-antibodies, the genesis of which is T lymphocyte-dependent.

Psoriasis is a T lymphocyte-mediated inflammatory disease. Lesions contain infiltrates of T lymphocytes, macrophages and antigen processing cells, and some neutrophils.

Allergic diseases, including asthma; allergic rhinitis; atopic dermatitis; food hypersensitivity; and urticaria are T lymphocyte dependent. These diseases are predominantly mediated by T lymphocyte induced inflammation, IgE mediated-inflammation or a combination of both.

Transplantation associated diseases, including Graft rejection and Graft-Versus-Host-Disease (GVHD) are T lymphocyte-dependent; inhibition of T lymphocyte function is ameliorative.

Other diseases in which intervention of the immune and/or inflammatory response have benefit are infectious disease including but not limited to viral infection (including but not limited to AIDS, hepatitis A, B, C, D, E and herpes) bacterial infection, fungal infections, and protozoal and parasitic infections (molecules (or derivatives/agonists) which stimulate the MLR can be utilized therapeutically to enhance the immune response to infectious agents), diseases of immunodeficiency (molecules/derivatives/agonists) which stimulate the MLR can be utilized therapeutically to enhance the immune response for conditions of inherited, acquired, infectious induced (as in HIV infection), or iatrogenic (i.e. as from chemotherapy) immunodeficiency), and neoplasia.

It has been demonstrated that some human cancer patients develop an antibody and/or T lymphocyte response to antigens on neoplastic cells. It has also been shown in animal models of neoplasia that enhancement of the immune response can result in rejection or regression of that particular neoplasm. Molecules that enhance the T lymphocyte response in the MLR have utility in vivo in enhancing the immune response against neoplasia. Molecules which enhance the T lymphocyte proliferative response in the MLR (or small molecule agonists or antibodies that affected the same receptor in an agonistic fashion) can be used therapeutically to treat cancer. Molecules that inhibit the lymphocyte response in the MLR also function in vivo during neoplasia to suppress the immune response to a neoplasm; such molecules can either be expressed by the neoplastic cells themselves or their expression can be induced by the neoplasm in other cells. Antagonism of such inhibitory molecules (either with antibody, small molecule antagonists or other means) enhances immune-mediated tumor rejection.

Additionally, inhibition of molecules with proinflammatory properties may have therapeutic benefit in reperfusion injury; stroke; myocardial infarction; atherosclerosis; acute lung injury;

hemorrhagic shock; burn; sepsis/septic shock; acute tubular necrosis; endometriosis; degenerative joint disease and pancreatitis.

The compounds of the present invention, e.g. polypeptides or antibodies, are administered to a mammal, preferably a human, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation (intranasal, intrapulmonary) routes. Intravenous or inhaled administration of polypeptides and antibodies is preferred.

In immunoadjuvant therapy, other therapeutic regimens, such administration of an anti-cancer agent, may be combined with the administration of the proteins, antibodies or compounds of the instant invention. For example, the patient to be treated with a the immunoadjuvant of the invention may also receive an anti-cancer agent (chemotherapeutic agent) or radiation therapy. Preparation and dosing schedules for such chemotherapeutic agents may be used according to manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in *Chemotherapy Service* Ed., M.C. Perry, Williams & Wilkins, Baltimore, MD (1992). The chemotherapeutic agent may precede, or follow administration of the immunoadjuvant or may be given simultaneously therewith. Additionally, an anti-oestrogen compound such as tamoxifen or an anti-progesterone such as onapristone (see, EP 616812) may be given in dosages known for such molecules.

It may be desirable to also administer antibodies against other immune disease associated or tumor associated antigens, such as antibodies which bind to CD20, CD11a, CD18, ErbB2, EGFR, ErbB3, ErbB4, or vascular endothelial factor (VEGF). Alternatively, or in addition, two or more antibodies binding the same or two or more different antigens disclosed herein may be coadministered to the patient. Sometimes, it may be beneficial to also administer one or more cytokines to the patient. In one embodiment, the polypeptides of the invention are coadministered with a growth inhibitory agent. For example, the growth inhibitory agent may be administered first, followed by a polypeptide of the invention. However, simultaneous administration or administration first is also contemplated. Suitable dosages for the growth inhibitory agent are those presently used and may be lowered due to the combined action (synergy) of the growth inhibitory agent and the polypeptide of the invention.

For the treatment or reduction in the severity of immune related disease, the appropriate dosage of an a compound of the invention will depend on the type of disease to be treated, as defined above, the severity and course of the disease, whether the agent is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the compound, and the discretion of the attending physician. The compound is suitably administered to the patient at one time or over a series of treatments.

For example, depending on the type and severity of the disease, about 1 ug/kg to 15 mg/kg (e.g. 0.1-20mg/kg) of polypeptide or antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily dosage might range from about 1 ug/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

#### 12. Articles of Manufacture

In another embodiment of the invention, an article of manufacture containing materials useful for the diagnosis or treatment of the disorders described above is provided. The article of manufacture comprises a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for diagnosing or treating the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agent in the composition is usually a polypeptide or an antibody of the invention. The label on, or associated with, the container indicates that the composition is used for diagnosing or treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

#### 13. Diagnosis and Prognosis of Immune Related Disease

Cell surface proteins, such as proteins which are overexpressed in certain immune related diseases, are excellent targets for drug candidates or disease treatment. The same proteins along with secreted proteins encoded by the genes amplified in immune related disease states find additional use in the diagnosis and prognosis of these diseases. For example, antibodies directed against the protein products of genes amplified in multiple sclerosis, rheumatoid arthritis, or another immune related disease, can be used as diagnostics or prognostics.

For example, antibodies, including antibody fragments, can be used to qualitatively or quantitatively detect the expression of proteins encoded by amplified or overexpressed genes ("marker gene products"). The antibody preferably is equipped with a detectable, e.g. fluorescent label, and binding can be monitored by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. These techniques are particularly suitable, if the overexpressed gene encodes a cell surface protein. Such binding assays are performed essentially as described above.

*In situ* detection of antibody binding to the marker gene products can be performed, for example, by immunofluorescence or immunoelectron microscopy. For this purpose, a histological specimen is removed from the patient, and a labeled antibody is applied to it, preferably by overlaying the antibody on a biological sample. This procedure also allows for determining the distribution of the marker gene product in the tissue examined. It will be apparent for those skilled in the art that a wide variety of histological methods are readily available for *in situ* detection.

The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

### EXAMPLES

Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA. Unless otherwise noted, the present invention uses standard procedures of recombinant DNA technology, such as those described hereinabove and in the following textbooks: Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press N.Y., 1989; Ausubel *et al.*, *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y., 1989; Innis *et al.*, *PCR Protocols: A Guide to Methods and Applications*, Academic Press, inc., N.Y., 1990; Harlow *et al.*, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, 1988; Gait, M.J., *Oligonucleotide Synthesis*, IRL Press, Oxford, 1984; R.I. Freshney, *Animal Cell Culture*, 1987; Coligan *et al.*, *Current Protocols in Immunology*, 1991.

#### EXAMPLE 1

Isolation of cDNA clones Encoding Human PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326

##### I. Isolation of cDNA Clones Encoding Human PRO245

The extracellular domain (ECD) sequences (including the secretion signal, if any) of from about 950 known secreted proteins from the Swiss-Prot public protein database were used to search expressed sequence tag (EST) databases. The EST databases included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ<sup>TM</sup>, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the computer program BLAST or BLAST2 (Altshul *et al.*, *Methods in Enzymology* 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6

frame translation of the EST sequence. Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Washington).

A consensus DNA sequence encoding PRO245 was assembled relative to the other identified EST sequences, where the consensus sequence was designated herein as DNA30954, and the polypeptide showed some structural homology to transmembrane protein receptor tyrosine kinase proteins.

Based on the DNA30954 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO245.

A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-ATCGTTGTGAAGTTAGTGCCCC-3' (SEQ ID NO: 15)

reverse PCR primer 5'-ACCTGCGATATCCAACAGAATTG-3' (SEQ ID NO: 16)

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30954 sequence which had the following nucleotide sequence:

hybridization probe

5'-GGAAGAGGATACAGTCACTCTGGAAGTATTAGTGGCTCCAGCAGTTCC-3' (SEQ ID NO: 17)

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO245 gene using the probe oligonucleotide and one of the PCR primers.

RNA for construction of the cDNA libraries was isolated from human fetal liver tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to Sall hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO245 [herein designated as UNQ219 (DNA35638)] and the derived protein sequence for PRO245.

The entire nucleotide sequence of UNQ219 (DNA35638) is shown in Figure 3 (SEQ ID NO: 1). Clone UNQ219 (DNA35638) contains a single open reading frame with an apparent translational

initiation site at nucleotide positions 89-91 [Kozak et al., *supra*] and ending at the stop codon at nucleotide positions 1025-1027 (Fig. 3). The predicted polypeptide precursor is 312 amino acids long (Figure. 4; PRO245; SEQ ID NO: 2). Clone UNQ219 (DNA35638) has been deposited with ATCC on September 17, 1997 and is assigned ATCC Deposit No. 209265.

Analysis of the amino acid sequence of the full-length PRO245 suggests that a portion of it possesses 60% amino acid identity with the human c-myc protein and, therefore, may be a new member of the transmembrane protein receptor tyrosine kinase family.

## II. Isolation of cDNA clones Encoding PRO217

The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public databases (e.g., Dayhof, GenBank), and proprietary databases (e.g. LIFESEQ<sup>TM</sup>, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the computer program BLAST or BLAST2 (Altschul, SF and Gish (1996), *Methods in Enzymology* 266: 460-80 (1996); <http://blast.wustl.edu/blast/README.html>) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons with a Blast score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, WA; (<http://bozeman.mbt.washington.edu/phrap.docs/phrap.html>)).

Consensus DNA sequences encoding EGF-like homologues were assembled (DNA28726, DNA28730 and DNA28760) using phrap. In some cases, the consensus DNA sequence was extended using repeated cycles of blast and phrap to extend the consensus sequence as far as possible using the three sources of EST sequences listed above.

Based on this consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence. The pair of forward and reverse PCR primers (notated as \*.f and \*.r, respectively) may range from 20 to 30 nucleotides (typically 24), and are designed to give a PCR product of 100-1000 bp in length. The probe sequences (notated as \*.p) are typically 40-55 bp (typically 50) in length. In some cases additional oligonucleotides are synthesized when the consensus sequence is greater than 1-1.5 kbp. In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, *Current Protocols in Molecular Biology*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest by the *in vivo* cloning procedure using the probe oligonucleotide and one of the PCR primers. This library was used to isolate DNA32279, DNA32292 and DNA33094 was fetal kidney, fetal lung and fetal lung, respectively.

RNA for the construction of the cDNA libraries was isolated using standard isolation protocols, e.g., Ausubel *et al.*, *Current Protocols in Molecular Biology*, from tissue or cell line sources or it was purchased from commercial sources (e.g., Clontech). The cDNA libraries used to isolate the cDNA clones were constructed by standard methods (e.g., Ausubel *et al.*) using commercially available reagents (e.g., Invitrogen). The cDNA was primed with oligo dT containing a NotI site, linked with blunt to Sall hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation in a suitable cloning vector (pRK5B or pRK5D) in the unique XhoI and NotI sites.

A cDNA clone was sequenced in its entirety. The entire nucleotide sequence of EGF-like homologue PRO217 is shown in Figure 5 (SEQ ID NO: 3). The predicted polypeptide is 379 (PRO217; Figure 6; SEQ ID NO: 4) amino acids in length with a molecule weight of approximately 41.52 kDa.

The oligonucleotide sequences used in the above procedure were the following:

28726.p (OLI500) (SEQ ID NO: 18)

GGGTACACCTGCTCCTGCACCGACGGATATTGGCTTCTGGAAGGCC

28726.f (OLI 502) (SEQ ID NO: 19)

ACAGATTCCCACCAAGTGCAACC

28726.r (OLI 503) (SEQ ID NO: 20)

CACACTCGTTCACATCTTGGC

28730.p (OLI 516) (SEQ ID NO: 21)

AGGGAGCACGGACAGTGTGCAGATGTGGACGAGTGCTCACTAGCA

28730.f (OLI 517) (SEQ ID NO: 22)

AGAGTGTATCTCTGGCTACGC

28730.r (OLI 518) (SEQ ID NO: 23)

TAAGTCCGGCACATTACAGGTC

28760.p (OLI 617) (SEQ ID NO: 24)

CCCACGATGTATGAATGGTGGACTTTGTGTGACTCCTGGTTTCTGCATC

28760.f (OLI 618) (SEQ ID NO: 25)

AAAGACGCATCTGCGAGTGTC

28760.r (OLI 619) (SEQ ID NO: 26)

TGCTGATTCACACTGCTCTCCC

### III. Isolation of cDNA clones Encoding Human PRO301

The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public EST databases (e.g., GenBank), a proprietary EST database (LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the

computer program BLAST or BLAST2 [Altschul et al., *Methods in Enzymology*, 266:460-480 (1996); <http://blast.wustl.edu/blast/README.html>] as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons resulting in a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Washington; <http://bozeman.mbt.washington.edu/phrap.docs/phrap.html>).

A consensus DNA sequence encoding DNA35936 was assembled using phrap. In some cases, the consensus DNA sequence was extended using repeated cycles of blast and phrap to extend the consensus sequence as far as possible using the three sources of EST sequences listed above.

Based on this consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence. Forward and reverse PCR primers (notated as \*.f and \*.r, respectively) may range from 20 to 30 nucleotides (typically about 24), and are designed to give a PCR product of 100-1000 bp in length. The probe sequences (notated as \*.p) are typically 40-55 bp (typically about 50) in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than 1-1.5 kbp. In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, *Current Protocols in Molecular Biology*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest by the in vivo cloning procedure using the probe oligonucleotide and one of the PCR primers.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO301 gene using the probe oligonucleotide and one of the PCR primers.

RNA for construction of the cDNA libraries was isolated from human fetal kidney. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents (e.g., Invitrogen, San Diego, CA; Clontech, etc.) The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

A cDNA clone was sequenced in its entirety. The full length nucleotide sequence of native sequence PRO301 is shown in Figure 7 (SEQ ID NO: 5). Clone DNA40628 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 52-54 [Kozak et al., *supra*] (Fig. 7). The predicted polypeptide (PRO301; Figure 8; SEQ ID NO: 6) is 299 amino acids

long with a predicted molecular weight of 32583 daltons and pI of 8.29. Clone DNA40628 has been deposited with ATCC and is assigned ATCC deposit No. 209432.

Based on a BLAST and FastA sequence alignment analysis of the full-length sequence, PRO301 shows amino acid sequence identity to A33 antigen precursor (30%) and coxsackie and adenovirus receptor protein (29%).

The oligonucleotide sequences used in the above procedure were the following:

OLI2162 (35936.f1) (SEQ ID NO: 27)  
TCGCGGAGCTGTGTTCTGTTTCCC

OLI2163 (35936.p1) (SEQ ID NO: 28)  
TGATCGCGATGGGGACAAAGGCGCAAGCTCGAGAGGAACTGTTGTGCCT

OLI2164 (35936.f2) (SEQ ID NO: 29)  
ACACCTGGTTCAAAGATGGG

OLI2165 (35936.r1) (SEQ ID NO: 30)  
TAGGAAGAGTTGCTGAAGGCACGG

OLI2166 (35936.f3) (SEQ ID NO: 31)  
TTGCCTTACTCAGGTGCTAC

OLI2167 (35936.r2) (SEQ ID NO: 32)  
ACTCAGCAGTGGTAGGAAAG

#### IV. Isolation of cDNA Clones Encoding Human PRO266

The extracellular domain (ECD) sequences (including the secretion signal, if any) of from about 950 known secreted proteins from the Swiss-Prot public protein database were used to search expressed sequence tag (EST) databases. The EST databases included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ<sup>TM</sup>, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the computer program BLAST or BLAST2 (Altschul et al., Methods in Enzymology 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequence. Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Washington; <http://bozeman.mbt.washington.edu/phrap.docs/phrap.html>).

Based on an expression sequence tag oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO266. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5kbp. In

order to screen several libraries for a full-length clone. DNA from the libraries was screened by PCR amplification, as per Ausubel et al., Current Protocols in Molecular Biology, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest by the *in vivo* cloning procedure using the probe oligonucleotide and one of the primer pairs.

A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-GTTGGATCTGGGCAACAATAAC-3' (SEQ ID NO: 33)

reverse PCR primer 5'-ATTGTTGTGCAGGCTGAGTTTAAG-3' (SEQ ID NO: 34)

Additionally, a synthetic oligonucleotide hybridization probe was constructed which had the following nucleotide sequence:

hybridization probe

5'-GGTGGCTATACATGGATAGCAATTACCTGGACACGCTGTCCCGGG-3' (SEQ ID NO: 35)

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO266 gene using the probe oligonucleotide and one of the PCR primers.

RNA for construction of the cDNA libraries was isolated from human fetal brain tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO266 [herein designated as UNQ233 (DNA37150)] and the derived protein sequence for PRO266.

The entire nucleotide sequence of UNQ233 (DNA37150) is shown in Figure 9 (SEQ ID NO: 7). Clone UNQ233 (DNA37150) contains a single open reading frame with an apparent translational initiation site at nucleotide positions 1-3 [Kozak et al., supra] and ending at the stop codon after nucleotide position 2088. The predicted polypeptide precursor is 696 amino acids long (Figure 10; PRO266: SEQ ID NO: 8). Clone UNQ233 (DNA37150) has been deposited with ATCC and is assigned ATCC deposit no. 209401.

Analysis of the amino acid sequence of the full-length PRO266 polypeptide suggests that portions of it possess significant homology to a SLIT protein, thereby indicating that PRO266 may be a novel leucine rich repeat protein.

#### V. Isolation of cDNA Clones Encoding Human PRO335, PRO331 or PRO326

The extracellular domain (ECD) sequences (including the secretion signal, if any) of from about 950 known secreted proteins from the Swiss-Prot public protein database were used to search expressed sequence tag (EST) databases. The EST databases included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the computer program BLAST or BLAST2 (Altschul et al., Methods in Enzymology 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequence. Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Washington; <http://bozeman.mbt.washington.edu/phrap.docs/phrap.html>).

A consensus DNA sequence was assembled relative to other EST sequences using phrap. Based on the consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO335, PRO331 or PRO326. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., Current Protocols in Molecular Biology, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest by the *in vivo* clonin procedure using the probe oligonucleotide and one of the primer pairs.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO335, PRO331 or PRO326 gene using the probe oligonucleotide and one of the PCR primers.

RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue (PRO335 and PRO326) and human fetal brain (PRO331). The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO335 (Figure 11; SEQ ID NO: 9), PRO331 (Figure 13; SEQ ID NO: 11) or PRO326

(Figure 15; SEQ ID NO: 13) and the derived protein sequence for PRO335 (Figure 12; SEQ ID NO: 10), PRO331 (Figure 14; SEQ ID NO: 12) or PRO326 (Figure 16; SEQ ID NO: 14). The nucleic acid encoding PRO335 was deposited with the ATCC on 2 June 1998 and is assigned ATCC Accession No. 209927; the nucleic acid encoding PRO331 was deposited with the ATCC on 7 November 1997 and is assigned ATCC Accession No. 209439; and the nucleic acid encoding PRO326 was deposited with the ATCC on 21 November 1997 and is assigned ATCC Accession No. 209489.

Analysis of the amino acid sequence of the full-length PRO335, PRO331 or PRO326 polypeptide suggests that portions of it possess significant homology to the *LIG-1* protein, thereby indicating that PRO335, PRO331 and PRO326 may be a novel *LIG-1*-related protein.

## EXAMPLE 2

### Stimulatory Activity in Mixed Lymphocyte Reaction (MLR) Assay (No. 24)

This example shows that the polypeptides of the invention are active as a stimulator of the proliferation of stimulated T-lymphocytes. Compounds which stimulate proliferation of lymphocytes are useful therapeutically where enhancement of an immune response is beneficial. Compounds which inhibit proliferation of lymphocytes are useful therapeutically where suppression of an immune response is beneficial. A therapeutic agent may take the form of antagonists of the polypeptide of the invention, for example, murine-human chimeric, humanized or human antibodies against the polypeptide.

The basic protocol for this assay is described in *Current Protocols in Immunology*, unit 3.12; edited by J. E. Coligan, A. M. Kruisbeek, D. H. Marglies, E. M. Shevach, W. Strober, National Institutes of Health, Published by John Wiley & Sons, Inc.

More specifically, in one assay variant, peripheral blood mononuclear cells (PBMC) are isolated from mammalian individuals, for example a human volunteer, by leukopheresis (one donor will supply stimulator PBMCs, the other donor will supply responder PBMCs). If desired, the cells are frozen in fetal bovine serum and DMSO after isolation. Frozen cells may be thawed overnight in assay media (37°C, 5% CO<sub>2</sub>) and then washed and resuspended to 3 x 10<sup>6</sup> cells/ml of assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate).

The stimulator PBMCs are prepared by irradiating the cells (about 3000 Rads). The assay is prepared by plating in triplicate wells a mixture of:

100µl of test sample diluted to 1% or to 0.1%

50 µl of irradiated stimulator cells and

50 µl of responder PBMC cells.

100 microliters of cell culture media or 100 microliter of CD4-IgG is used as the control. The wells are then incubated at 37°C, 5% CO<sub>2</sub> for 4 days. On day 5 and each well is pulsed with tritiated thymidine (i.0 mC/well; Amersham). After 6 hours the cells are washed 3 times and then the uptake of the label is evaluated.

In another variant of this assay, PBMCs are isolated from the spleens of Balb/c mice and C57B6 mice. The cells are teased from freshly harvested spleens in assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate) and the PBMCs are isolated by overlaying these cells over Lympholyte M (Organon Teknika), centrifuging at 2000 rpm for 20 minutes, collecting and washing the mononuclear cell layer in assay media and resuspending the cells to  $1 \times 10^7$  cells/ml of assay media. The assay is then conducted as described above using a sample having a PRO concentration obtained by diluting a stock solution. The results of this assay for compounds of the invention are shown below. Positive increases over control are considered positive with increases of greater than or equal to 180% being preferred. However, any value greater than control indicates a stimulatory effect for the test protein.

Table 2

<u>PRO</u>	<u>PRO Concentration</u>	<u>Percent Increase Over Control</u>
PRO245	0.1%	189.7
"	0.1%	193.7
"	1.0%	212.5
"	1.0%	300.5
PRO217	0.1%	74.5
"	1.0%	89.5
"	0.99 nM	97.0
"	9.9 nM	122.3
"	0.25 nM	144.8
"	2.5 nM	126.9
PRO301	50.0%	109.4
"	70.0 nM	133.7
"	700.0 nM	83.6
	0.1%	58.7
PRO301	1.0%	127.7
"	0.1%	181.7
"	1.0%	187.3
"	0.1%	127.5
"	1.0%	108.3

PRO266	0.1%	136.4
"	0.1%	139.2
"	1.0%	189.8
"	1.0%	245.1
PRO335	50.0%	91.0
"	50.0%	103.8
"	0.1%	130.0
"	1.0%	180.2
PRO331	50.0%	155.5
"	0.1%	169.3
"	1.0%	128.1
"	0.1%	129.3
"	1.0%	162.5
PRO326	50.0%	91.0
"	50.0%	103.8
"	0.1%	130.0
"	1.0%	180.2

---

### EXAMPLE 3

#### Skin Vascular Permeability Assay (No. 64)

This assay shows that certain polypeptides of the invention stimulate an immune response and induce inflammation by inducing mononuclear cell, eosinophil and PMN infiltration at the site of injection of the animal. This skin vascular permeability assay is conducted as follows. Hairless guinea pigs weighing 350 grams or more are anesthetized with ketamine (75-80 mg/Kg) and 5 mg/Kg xylazine intramuscularly (IM). A sample of purified polypeptide of the invention or a conditioned media test sample is injected intradermally onto the backs of the test animals with 100  $\mu$ L per injection site. It is possible to have about 10-30, preferably about 16-24, injection sites per animal. One mL of Evans blue dye (1% in physiologic buffered saline) is injected intracardially. Blemishes at the injection sites are then measured (mm diameter) at 1hr and 6 hr post injection. Animals were sacrificed at 6 hrs after injection. Each skin injection site was biopsied and fixed in formalin. The skins were then prepared for histopathologic evaluation. Each site was evaluated for inflammatory cell infiltration into the skin. Sites with visible inflammatory cell inflammation were scored as

positive. Inflammatory cells can be neutrophilic, eosinophilic, monocytic or lymphocytic. The results of this test for compounds of the invention is shown below.

In the Table below, at least a minimal perivascular infiltrate at the injection site is scored as positive, no infiltrate at the site of injection is scored as negative.

Table 3

<u>PRO</u>	<u>Hours Post Injection</u>	<u>Infiltrate Designation</u>
PRO245	24 hr	positive
PRO217	24 hr	positive
PRO301	24 hr	positive
PRO266	24 hr	positive
PRO335	24 hr	positive
PRO331	24 hr	positive
PRO326	24 hr	positive

#### EXAMPLE 4

##### Human Co-Stimulation Assay

In addition to the activation signal mediated by the T cell receptor, T cell activation requires a costimulatory signal. One costimulatory signal is generated by the interaction of B7 (CD3) with CD28. In this assay, 96 well plates are coated with CD3 with or without CD28 and then human peripheral blood lymphocytes followed by a test protein, are added. Proliferation of the lymphocytes is determined by tritiated thymidine uptake. A positive assay indicates that the test protein provided a stimulatory signal for lymphocyte proliferation.

##### Material:

- 1) Hyclone D-PBS without Calcium, Magnesium
- 2) Nunc 96 well certified plates #4-39454
- 3) Anti-human CD3 Amac 0178 200 µg/ml stock
- 4) Anti-human CD28 Biodesign P42235M
- 5) Media: Gibco RPMI 1640 + 10 % Intergen #1020-90 FBS, 1% Glu, 1% P/S, 50 µg/ml Gentamycin, 10 mM Hepes. Fresh for each assay.
- 6) Tritiated Thymidine
- 7) Frozen human peripheral blood lymphocytes (PBL) collected via a leukaphoresis procedure

Plates are prepared by coating 96 well flat bottom plates with anti-CD3 antibody (Amac) or anti-CD28 antibody (Biodesign) or both in Hyclone D-PBS without calcium and magnesium. Anti - CD3 antibody is used at 10 ng/well (50 $\mu$ l of 200 ng/ml) and anti -CD28 antibody at 25 ng/well (50  $\mu$ l of 0.5  $\mu$ g/ml) in 100  $\mu$ l total volume.

PBLs are isolated from human donors using standard leukophoresis methods. The cell preparations are frozen in 50% fetal bovine serum and 50% DMSO until the assay is conducted. Cells are prepared by thawing and washing PBLs in media, resuspending PBLs in 25 mls of media and incubating at 37°C, 5% CO<sub>2</sub> overnight.

In the assay procedure, the coated plate is washed twice with PBS and the PBLs are washed and resuspended to 1 x 10<sup>6</sup> cells/ml using 100  $\mu$ L /well. 100  $\mu$ l of a test protein or control media are added to the plate making a total volume per well of 200  $\mu$ L. The plate is incubated for 72 hours. The plate is then pulsed for 6 hours with tritiated thymidine (1 mC/well; Amersham) and the PBLs are harvested from the plates and evaluated for uptake of the tritiated thymidine.

## EXAMPLE 5

### In situ Hybridization

*In situ* hybridization is a powerful and versatile technique for the detection and localization of nucleic acid sequences within cell or tissue preparations. It may be useful, for example, to identify sites of gene expression, analyze the tissue distribution of transcription, identify and localize viral infection, follow changes in specific mRNA synthesis and aid in chromosome mapping.

*In situ* hybridization was performed following an optimized version of the protocol by Lu and Gillett, Cell Vision 1: 169-176 (1994), using PCR-generated <sup>33</sup>P-labeled riboprobes. Briefly, formalin-fixed, paraffin-embedded human tissues were sectioned, deparaffinized, deproteinated in proteinase K (20 g/ml) for 15 minutes at 37°C, and further processed for *in situ* hybridization as described by Lu and Gillett, *supra*. A [<sup>33</sup>P] UTP-labeled antisense riboprobe was generated from a PCR product and hybridized at 55C overnight. The slides were dipped in Kodak NTB2 nuclear track emulsion and exposed for 4 weeks.

#### <sup>33</sup>P-Riboprobe synthesis

6.0  $\mu$ l (125 mCi) of <sup>33</sup>P-UTP (Amersham BF 1002, SA<2000 Ci/mmol) were speed vac dried. To each tube containing dried <sup>33</sup>P-UTP, the following ingredients were added:

2.0  $\mu$ l 5x transcription buffer

1.0  $\mu$ l DTT (100 mM)

2.0  $\mu$ l NTP mix (2.5 mM : 10 $\mu$  l; each of 10 mM GTP, CTP & ATP + 10 $\mu$  l H<sub>2</sub>O)

1.0  $\mu$ l UTP (50  $\mu$ M)

- 1.0  $\mu$ l Rnasin
- 1.0  $\mu$ l DNA template (1 $\mu$ g)
- 1.0  $\mu$ l H<sub>2</sub>O

The tubes were incubated at 37°C for one hour. 1.0  $\mu$ L RQ1 DNase were added, followed by incubation at 37°C for 15 minutes. 90  $\mu$ L TE (10 mM Tris pH 7.6/1mM EDTA pH 8.0) were added, and the mixture was pipetted onto DE81 paper. The remaining solution was loaded in a Microcon-50 ultrafiltration unit, and spun using program 10 (6 minutes). The filtration unit was inverted over a second tube and spun using program 2 (3 minutes). After the final recovery spin, 100  $\mu$ L TE were added. 1  $\mu$ L of the final product was pipetted on DE81 paper and counted in 6 ml of Biofluor II.

The probe was run on a TBE/urea gel. 1-3  $\mu$ L of the probe or 5  $\mu$ L of RNA Mrk III were added to 3  $\mu$ L of loading buffer. After heating on a 95C heat block for three minutes, the gel was immediately placed on ice. The wells of gel were flushed, the sample loaded, and run at 180-250 volts for 45 minutes. The gel was wrapped in saran wrap and exposed to XAR film with an intensifying screen in -70C freezer one hour to overnight.

#### <sup>33</sup>P-Hybridization

*Pretreatment of frozen sections* The slides were removed from the freezer, placed on aluminium trays and thawed at room temperature for 5 minutes. The trays were placed in 55C incubator for five minutes to reduce condensation. The slides were fixed for 10 minutes in 4% paraformaldehyde on ice in the fume hood, and washed in 0.5 x SSC for 5 minutes, at room temperature (25 ml 20 x SSC + 975 ml SQ H<sub>2</sub>O). After deproteination in 0.5  $\mu$ g/ml proteinase K for 10 minutes at 37°C (12.5 $\mu$ L of 10 mg/ml stock in 250 ml prewarmed RNase-free RNase buffer), the sections were washed in 0.5 x SSC for 10 minutes at room temperature. The sections were dehydrated in 70%, 95%, 100% ethanol, 2 minutes each.

*Pretreatment of paraffin-embedded sections* The slides were deparaffinized, placed in SQ H<sub>2</sub>O, and rinsed twice in 2 x SSC at room temperature, for 5 minutes each time. The sections were deproteinated in 20  $\mu$ g/ml proteinase K (500  $\mu$ L of 10 mg/ml in 250 ml RNase-free RNase buffer; 37C, 15 minutes) - human embryo, or 8 x proteinase K (100  $\mu$ L in 250 ml Rnase buffer, 37C, 30 minutes) - formalin tissues. Subsequent rinsing in 0.5 x SSC and dehydration were performed as described above.

*Prehybridization* The slides were laid out in plastic box lined with Box buffer (4 x SSC, 50% formamide) - saturated filter paper. The tissue was covered with 50  $\mu$ L of hybridization buffer (3.75g Dextran Sulfate + 6 ml SQ H<sub>2</sub>O), vortexed and heated in the microwave for 2 minutes with the cap loosened. After cooling on ice, 18.75 ml formamide, 3.75 ml 20 x SSC and 9 ml SQ H<sub>2</sub>O were added, the tissue was vortexed well, and incubated at 42C for 1-4 hours.

*Hybridization* 1.0 x 10<sup>6</sup> cpm probe and 1.0 µL tRNA (50 mg/ml stock) per slide were heated at 95C for 3 minutes. The slides were cooled on ice, and 48 µL hybridization buffer were added per slide. After vortexing, 50 µL <sup>33</sup>P mix were added to 50 µL prehybridization on slide. The slides were incubated overnight at 55C.

*Washes* Washing was done 2x10 minutes with 2xSSC, EDTA at room temperature (400 ml 20 x SSC + 16 ml 0.25M EDTA, V<sub>f</sub>=4L), followed by RNaseA treatment at 37C for 30 minutes (500 µL of 10 mg/ml in 250 ml Rnase buffer = 20 ug/ml), The slides were washed 2x10 minutes with 2 x SSC, EDTA at room temperature. The stringency wash conditions were as follows: 2 hours at 55C, 0.1 x SSC, EDTA (20 ml 20 x SSC + 16 ml EDTA, V<sub>f</sub>=4L).

#### DNA 35638 (1 TM receptor)

Expression was observed in the endothelium lining of a subset of fetal and placental vessels. Endothelial expression was confined to these tissue blocks. Expression was also observed over intermediate trophoblast cells of placenta.

Oligo C-120N: (SEQ ID NO: 36)

GGA TTC TAA TAC GAC TCA CTA TAG GGC TGC GGC GGC TCA GGT CTT CAG TT

Oligo c-120P (SEQ ID NO: 37)

CTA TGA AAT TAA CCC TCA CTA AAG GGA GCA TGG GAT GGG GAG GGA TAC GG

#### DNA 33094 (EGF Homolog)

A highly distinctive expression pattern was observed. In the human embryo expression was observed in outer smooth muscle layer of the GI tract, respiratory cartilage, branching respiratory epithelium, osteoblasts, tendons, gonad, in the optic nerve head and developing dermis. In the adult, expression was observed in the epidermal pegs of the chimp tongue, the basal epithelial / myoepithelial cells of the prostate and urinary bladder. Expression was also found in the alveolar lining cells of the adult lung, mesenchymal cells juxtaposed to erectile tissue in the penis and the cerebral cortex (probably glial cells). In the kidney, expression was only seen in disease, in cells surrounding thyroidized renal tubules.

Oligo D-200V (SEQ ID NO: 38)

CTA TGA AAT TAA CCC TCA CTA AAG GGA ATA GCA GGC CAT CCC AGG ACA

Oligo D-200Z (SEQ ID NO: 39)

CTA TGA AAT TAA CCC TCA CTA AAG GGA TGT CTT CCA TGC CAA CCT TC

## EXAMPLE 6

### *In situ* Hybridization in Cells and Diseased Tissues

The *in situ* hybridization method of Example 5 is used to determine gene expression, analyze the tissue distribution of transcription, and follow changes in specific mRNA synthesis for the genes/DNAs and the proteins of the invention in diseased tissues isolated from human individuals suffering from a specific disease. These results show more specifically where in diseased tissues the genes of the invention are expressed and are more predictive of the particular localization of the therapeutic effect of the inhibitory or stimulatory compounds of the invention (and agonists or antagonists thereof) in a disease. Hybridization is performed according to the method of Example 5 using one or more of the following tissue and cell samples:

(a) lymphocytes and antigen presenting cells (dendritic cells, langherhans cells, macrophages and monocytes, NK cells);

(b) lymphoid tissues: normal and reactive lymph node, thymus, Bronchial Associated Lymphoid Tissues, (BALT), Mucosal Associated Lymphoid Tissues (MALT);

(c) human disease tissues

- Synovium and joint of patients with Arthritis and Degenerative Joint Disease
- Colon from patients with Inflammatory Bowel Disease including Ulcerative Colitis and Crohns' disease
- Skin lesions from Psoriasis and other forms of dermatitis
- Lung tissue including BALT and tissue lymph nodes from Chronic and acute bronchitis, pneumonia, pneumonitis, pleuritis
- Lung tissue including BALT and tissue lymph nodes from Asthma
- nasal and sinus tissue from patients with rhinitis or sinusitis
- Brain and Spinal cord from Multiple Sclerosis. Alzheimer's Disease and Stroke
- Kidney from Nephritis, Glomerulonephritis and Systemic Lupus Erythematosis
- Liver from Infectious and non-infectious Hepatitis
- Tissues from Neoplasms/Cancer.

Expression is observed in one or more cell or tissue samples indicating localization of the therapeutic effect of the compounds of the invention (and agonists or antagonists thereof) in the

disease associated with the cell or tissue sample.

DNA 35638 (PRO245) was found to be expressed in inflamed human tissues (psoriasis, inflammatory bowel disease (IBD), inflamed kidney, inflamed lung, hepatitis (liver block), normal tonsil, adult and chimp (multiblocks). Expression was present in the endothelium/intima of large vessels in the lung afflicted with chronic inflammation, in the superficial dermal vessels of the psoriatic skin, in arterioles in a specimen of chronic sclerosing nephritis, and in capillaries including the perifollicular sinuses of the tonsil. These results indicate that PRO245 is immunostimulatory (enhances T lymphocyte proliferation in the MLR and costimulation) and has proinflammatory properties (induces a neutrophil infiltrate *in vivo*).

#### EXAMPLE 7

##### Use of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 as a hybridization probe

The following method describes use of a nucleotide sequence encoding PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 as a hybridization probe.

DNA comprising the coding sequence of full-length or mature PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 (as shown in Figures 4, 6, 8, 10, 12, 14 and 16) is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326) in human tissue cDNA libraries or human tissue genomic libraries.

Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326-derived probe to the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.

DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can then be identified using standard techniques known in the art.

#### EXAMPLE 8

##### Expression of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 in *E. coli*

This example illustrates preparation of an unglycosylated form of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 by recombinant expression in *E. coli*.

The DNA sequence encoding PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar et al., Gene, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 coding region, lambda transcriptional terminator, and an argU gene.

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., supra. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

PRO245, PRO217, PRO301 and PRO266 were expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO245, PRO217, PRO301 and PRO266 was initially amplified using selected PCR primers. The primers contained restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences were then ligated into an expression vector, which was used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) lon galE rpoHts(htpRts) clpP(lacIq)). Transformants were first grown in LB containing 50 mg/ml carbenicillin at 30C with shaking until an O.D.600 of 3-5 was reached. Cultures were then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.71 g sodium citrate.2H<sub>2</sub>O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield

hycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO<sub>4</sub>) and grown for approximately 20-30 hours at 30C with shaking. Samples were removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets were frozen until purification and refolding.

*E. coli* paste from 0.5 to 1 L fermentations (6-10 g pellets) was resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution was stirred overnight at 4C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution was centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant was diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. Depending the clarified extract was loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column was washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol grade), pH 7.4. The protein was eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein were pooled and stored at 4C. Protein concentration was estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

The proteins were refolded by diluting sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes were chosen so that the final protein concentration was between 50 to 100 micrograms/ml. The refolding solution was stirred gently at 4C for 12-36 hours. The refolding reaction was quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution was filtered through a 0.22 micron filter and acetonitrile was added to 2-10% final concentration. The refolded protein was chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A280 absorbance were analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein were pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

Fractions containing the desired folded PRO245, PRO217, PRO301 and PRO266 proteins, respectively, were pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins were formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium

chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

#### EXAMPLE 9

##### Expression of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 in mammalian cells

This example illustrates preparation of a potentially glycosylated form of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 by recombinant expression in mammalian cells.

The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 DNA using ligation methods such as described in Sambrook et al., *supra*. The resulting vector is called pRK5-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326.

In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 ug pRK5-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 DNA is mixed with about 1 ug DNA encoding the VA RNA gene [Thimmappaya et al., *Cell*, 31:543 (1982)] and dissolved in 500 uL of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl<sub>2</sub>. To this mixture is added, dropwise, 500 uL of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO<sub>4</sub>, and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 uCi/ml <sup>35</sup>S-cysteine and 200 uCi/ml <sup>35</sup>S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

In an alternative technique, PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 may be introduced into 293 cells transiently using the dextran sulfate method described by Sompariyac et al., *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 ug pRK5-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or

PRO326 DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5  $\mu\text{g/ml}$  bovine insulin and 0.1  $\mu\text{g/ml}$  bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

In another embodiment, PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can be expressed in CHO cells. The pRK5-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can be transfected into CHO cells using known reagents such as  $\text{CaPO}_4$  or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as  $^{35}\text{S}$ -methionine. After determining the presence of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can then be concentrated and purified by any selected method.

Epitope-tagged PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 may also be expressed in host CHO cells. The PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can then be concentrated and purified by any selected method, such as by  $\text{Ni}^{2+}$ -chelate affinity chromatography.

PRO245, PRO217 and PRO301 were expressed in CHO cells by both a transient and a stable expression procedure.

Stable expression in CHO cells was performed using the following procedure. The proteins were expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins were fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

Following PCR amplification, the respective DNAs were subcloned in a CHO expression

vector using standard techniques as described in Ausubel et al., *Current Protocols of Molecular Biology*, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNAs. The vector used expression in CHO cells is as described in Lucas *et al.*, *Nucl. Acids Res.* 24: 9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

Twelve micrograms of the desired plasmid DNA were introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect (Quiagen), Dosper or Fugene (Boehringer Mannheim). The cells were grown and described in Lucas *et al.*, *supra*. Approximately  $3 \times 10^7$  cells are frozen in an ampule for further growth and production as described below.

The ampules containing the plasmid DNA were thawed by placement into water bath and mixed by vortexing. The contents were pipetted into a centrifuge tube containing 10 mLs of media and centrifuged at 1000 rpm for 5 minutes. The supernatant was aspirated and the cells were resuspended in 10 mL of selective media (0.2  $\mu\text{m}$  filtered PS20 with 5% 0.2  $\mu\text{m}$  diafiltered fetal bovine serum). The cells were then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells were transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37C. After another 2-3 days, a 250 mL, 500 mL and 2000 mL spinners were seeded with  $3 \times 10^5$  cells/mL. The cell media was exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Patent No. 5,122,469, issued June 16, 1992 was actually used. 3L production spinner is seeded at  $1.2 \times 10^6$  cells/mL. On day 0, the cell number pH were determined. On day 1, the spinner was sampled and sparging with filtered air was commenced. On day 2, the spinner was sampled, the temperature shifted to 33C, and 30 mL of 500 g/L glucose and 0.6 mL of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion). Throughout the production, pH was adjusted as necessary to keep at around 7.2. After 10 days, or until viability dropped below 70%, the cell culture was harvested by centrifugation and filtering through a 0.22  $\mu\text{m}$  filter. The filtrate was either stored at 4C or immediately loaded onto columns for purification.

For the poly-His tagged constructs, the proteins were purified using a Ni-NTA column (Quiagen). Before purification, imidazole was added to the conditioned media to a concentration of 5 mM. The conditioned media was pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4C. After loading, the column was washed with additional equilibration buffer and the protein eluted with

equilibration buffer containing 0.25 M imidazole. The highly purified protein was subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80C.

Immunoadhesin (Fc containing) constructs of were purified from the conditioned media as follows. The conditioned medium was pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column was washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein was immediately neutralized by collecting 1 ml fractions into tubes containing 275  $\mu$ L of 1 M Tris buffer, pH 9. The highly purified protein was subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity was assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

PRO326 was also produced by transient expression in COS cells.

#### EXAMPLE 10

##### Expression of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 in Yeast

The following method describes recombinant expression of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 from the ADH2/GAPDH promoter. DNA encoding PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326. For secretion, DNA encoding PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 may further be

purified using selected column chromatography resins.

#### EXAMPLE 11

##### Expression of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 in Baculovirus-Infected Insect Cells

The following method describes recombinant expression of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 in Baculovirus-infected insect cells.

The sequence coding for PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 or the desired portion of the coding sequence of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 [such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular] is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4 - 5 days of incubation at 28°C, the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilley et al., Baculovirus expression vectors: A Laboratory Manual, Oxford: Oxford University Press (1994).

Expressed poly-his tagged PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can then be purified, for example, by Ni<sup>2+</sup>-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., Nature, 362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl<sub>2</sub>; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 um filter. A Ni<sup>2+</sup>-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A<sub>280</sub> with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH

6.0), which elutes nonspecifically bound protein. After reaching  $A_{280}$  baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with  $Ni^{2+}$ -NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His<sub>10</sub>-tagged PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG tagged (or Fc tagged) PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

PRO245, PRO301 and PRO266 were expressed in baculovirus infected Sf9 insect cells. While the expression was actually performed in a 0.5-2 L scale, it can be readily scaled up for larger (e.g. 8 L) preparations. The proteins were expressed as an IgG construct (immunoadhesin), in which the protein extracellular region was fused to an IgG1 constant region sequence containing the hinge, CH2 and CH3 domains and/or in poly-His tagged forms.

Following PCR amplification, the respective coding sequences were subcloned into a baculovirus expression vector (pb.PH.IgG for IgG fusions and pb.PH.His.c for poly-His tagged proteins), and the vector and Baculogold baculovirus DNA (Pharmingen) were co-transfected into 105 *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711), using Lipofectin (Gibco BRL). pb.PH.IgG and pb.PH.His are modifications of the commercially available baculovirus expression vector pVL1393 (Pharmingen), with modified polylinker regions to include the His or Fc tag sequences. The cells were grown in Hink's TNM-FH medium supplemented with 10% FBS (Hyclone). Cells were incubated for 5 days at 28C. The supernatant was harvested and subsequently used for the first viral amplification by infecting Sf9 cells in Hink's TNM-FH medium supplemented with 10% FBS at an approximate multiplicity of infection (MOI) of 10. Cells were incubated for 3 days at 28C. The supernatant was harvested and the expression of the constructs in the baculovirus expression vector was determined by batch binding of 1 ml of supernatant to 25 mL of Ni-NTA beads (QIAGEN) for histidine tagged proteins or Protein-A Sepharose CL-4B beads (Pharmacia) for IgG tagged proteins followed by SDS-PAGE analysis comparing to a known concentration of protein standard by Coomassie blue staining.

The first viral amplification supernatant was used to infect a spinner culture (500 ml) of Sf9 cells grown in ESF-921 medium (Expression Systems LLC) at an approximate MOI of 0.1. Cells were incubated for 3 days at 28C. The supernatant was harvested and filtered. Batch binding and SDS-PAGE analysis was repeated, as necessary, until expression of the spinner culture was confirmed.

The conditioned medium from the transfected cells (0.5 to 3 L) was harvested by centrifugation to remove the cells and filtered through 0.22 micron filters. For the poly-His tagged

constructs, the protein construct were purified using a Ni-NTA column (Qiagen). Before purification, imidazole was added to the conditioned media to a concentration of 5 mM. The conditioned media were pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4C. After loading, the column was washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein was subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80C.

Immunoadhesin (Fc containing) constructs of proteins were purified from the conditioned media as follows. The conditioned media were pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column was washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein was immediately neutralized by collecting 1 ml fractions into tubes containing 275 mL of 1 M Tris buffer, pH 9. The highly purified protein was subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity of the proteins was verified by SDS polyacrylamide gel (PAGE) electrophoresis and N-terminal amino acid sequencing by Edman degradation.

PRO245, PRO217, PRO301, PRO266, PRO331 and PRO326 were also expressed in baculovirus infected High-5 cells using an analogous procedure.

## EXAMPLE 12

### Preparation of Antibodies that Bind PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326

This example illustrates preparation of monoclonal antibodies which can specifically bind PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326.

Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326, fusion proteins containing PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326, and cells expressing recombinant PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

Mice, such as Balb/c, are immunized with the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days

later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 is within the skill in the art.

The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

#### Deposit of Material

The following materials have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, USA (ATCC):

<u>Material</u>	<u>ATCC Dep. No.</u>	<u>Deposit Date</u>
DNA40981	209439	7 November 1997
DNA37140	209489	21 November 1997
DNA41388	209927	2 June 1998
DNA35638	209265	17 September 1997
DNA37150	209401	17 October 1997

DNA33094	209256	16 September 1997
DNA32292	209258	16 September 1997
DNA32279	209259	16 September 1997
DNA40628	209432	7 November 1997

This deposit was made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposit will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC 122 and the Commissioner's rules pursuant thereto (including 37 CFR 1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct deposited, since the deposited embodiment is intended as a single illustration of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

TABLE 4

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXXXYYYYYYY	(Length = 12 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

5 divided by 15 = 33.3%

PRO	XXXXXXXXXX	(Length = 10 amino acids)
Comparison Protein	XXXXXXXXYYYYYYZZYZ	(Length = 15 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

5 divided by 10 = 50%

PRO-DNA	NNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLLLL	(Length = 16 nucleotides)

% nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

6 divided by 14 = 42.9%

PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLLTV	(Length = 9 nucleotides)

% nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

4 divided by 12 = 33.3%

TABLE 5

```

/*
 *
 * C-C increased from 12 to 15
 * Z is average of EQ
 * B is average of ND
 * match with stop is _M; stop-stop = 0; J (joker) match = 0
 */
#define _M -8 /* value of a match with a stop */

int _day[26][26] = {
/* A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */ {2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */ {0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */ {-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
/* D */ {0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */ {0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */ {-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */ {1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ {-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */ {-1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */ {-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */ {-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
/* M */ {-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */ {0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ {_M, _M, 0, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ {1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ {0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */ {-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */ {1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */ {1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ {0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */ {-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */ {-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ {0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4}
};

```

## TABLE 5 (cont.)

```

/*
*/
#include <stdio.h>
#include <ctype.h>

#define MAXJMP 16      /* max jumps in a diag */
#define MAXGAP 24     /* don't continue to penalize gaps larger than this */
#define JMPS 1024     /* max jmps in an path */
#define MX 4          /* save if there's at least MX-1 bases since last jmp */

#define DMAT 3        /* value of matching bases */
#define DMIS 0        /* penalty for mismatched bases */
#define DINS0 8       /* penalty for a gap */
#define DINS1 1       /* penalty per base */
#define PINS0 8       /* penalty for a gap */
#define PINS1 4       /* penalty per residue */

struct jmp {
    short      n[MAXJMP];      /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
};

struct diag {
    int      score;          /* score at last jmp */
    long     offset;        /* offset of prev block */
    short    jmp;           /* current jmp index */
    struct jmp jp;         /* list of jmps */
};

struct path {
    int      spc;           /* number of leading spaces */
    short    n[JMPS];      /* size of jmp (gap) */
    int      x[JMPS];      /* loc of jmp (last elem before gap) */
};

char        *ofile;        /* output file name */
char        *namex[2];     /* seq names: getseqs() */
char        *prog;         /* prog name for err msgs */
char        *seqx[2];      /* seqs: getseqs() */
int         dmax;          /* best diag: nw() */
int         dmax0;         /* final diag */
int         dna;           /* set if dna: main() */
int         endgaps;       /* set if penalizing end gaps */
int         gapx, gapy;     /* total gaps in seqs */
int         len0, len1;    /* seq lens */
int         ngapx, ngapy;  /* total size of gaps */
int         smax;          /* max score: nw() */
int         *xbm;          /* bitmap for matching */
long        offset;        /* current offset in jmp file */
struct      diag           /* holds diagonals */
*dx;
struct      path           /* holds path for seqs */
pp[2];

char        *calloc(), *malloc(), *index(), *strcpy();
char        *getseq(), *g_calloc();

```

TABLE 5 (cont.)

```

/* Needleman-Wunsch alignment program
*
* usage: prog file1 file2
* where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
#include "day.h"

static  _dbval[26] = {
1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};

static  _pbval[26] = {
1, 2|(1<<('D':'A'))|(1<<('N':'A')), 4, 8, 16, 32, 64,
128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
1<<23, 1<<24, 1<<25|(1<<('E':'A'))|(1<<('Q':'A'))
};

main(ac, av)      main
int              ac;
char             *av[];
{
prog = av[0];
if (ac != 3) {
fprintf(stderr, "usage: %s file1 file2\n", prog);
fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
fprintf(stderr, "The sequences can be in upper- or lower-case\n");
fprintf(stderr, "Any lines beginning with ';', '>' or '<' are ignored\n");
fprintf(stderr, "Output is in the file \"align.out\"\n");
exit(1);
}
namex[0] = av[1];
namex[1] = av[2];
seqx[0] = getseq(namex[0], &len0);
seqx[1] = getseq(namex[1], &len1);
xbm = (dna)? _dbval : _pbval;

endgaps = 0;                      /* 1 to penalize endgaps */
ofile = "align.out";              /* output file */

nw();                             /* fill in the matrix. get the possible jmps */
readjmps();                        /* get the actual jmps */
print();                           /* print stats. alignment */

cleanup(0);                        /* unlink any tmp files */
}

```

## TABLE 5 (cont.)

```

/* do the alignment, return best score: main()
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
* pro: PAM 250 values
* When scores are equal, we prefer mismatches to any gap, prefer
* a new gap to extending an ongoing gap, and prefer a gap in seqx
* to a gap in seq y.
*/
nw()    nw
{
    char          *px, *py;          /* seqs and ptrs */
    int           *ndely, *dely;     /* keep track of dely */
    int           ndelx, delx;       /* keep track of delx */
    int           *tmp;              /* for swapping row0, row1 */
    int           mis;               /* score for each type */
    int           ins0, ins1;        /* insertion penalties */
    register      id;                 /* diagonal index */
    register      ij;                 /* jmp index */
    register      *col0, *col1;      /* score for curr, last row */
    register      xx, yy;            /* index into seqs */

    dx = (struct diag *)g_malloc("to get diags", len0+len1+1, sizeof(struct diag));

    ndely = (int *)g_malloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_malloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_malloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_malloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;

    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;          /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
    */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
        */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0] - ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }
}

```

TABLE 5 (cont.)

...nw

```

for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
  mis = col0[yy-1];
  if (dna)
    mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
  else
    mis += _day[*px-'A'][*py-'A'];

  /* update penalty for del in x seq;
   * favor new del over ongong del
   * ignore MAXGAP if weighting endgaps
   */
  if (endgaps || ndely[yy] < MAXGAP) {
    if (col0[yy] - ins0 >= dely[yy]) {
      dely[yy] = col0[yy] - (ins0+ins1);
      ndely[yy] = 1;
    } else {
      dely[yy] -= ins1;
      ndely[yy]++;
    }
  } else {
    if (col0[yy] - (ins0+ins1) >= dely[yy]) {
      dely[yy] = col0[yy] - (ins0+ins1);
      ndely[yy] = 1;
    } else
      ndely[yy]++;
  }

  /* update penalty for del in y seq;
   * favor new del over ongong del
   */
  if (endgaps || ndelx < MAXGAP) {
    if (col1[yy-1] - ins0 >= delx) {
      delx = col1[yy-1] - (ins0+ins1);
      ndelx = 1;
    } else {
      delx -= ins1;
      ndelx++;
    }
  } else {
    if (col1[yy-1] - (ins0+ins1) >= delx) {
      delx = col1[yy-1] - (ins0+ins1);
      ndelx = 1;
    } else
      ndelx++;
  }

  /* pick the maximum score: we're favoring
   * mis over any del and delx over dely
   */

```

TABLE 5 (cont.)

...nw

```

id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    coll[yy] = mis;
else if (delx >= dely[yy]) {
    coll[yy] = delx;
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejumps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndelx;
    dx[id].jp.x[ij] = xx;
    dx[id].score = delx;
}
else {
    coll[yy] = dely[yy];
    ij = dx[id].ijmp;
if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejumps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = -ndely[yy];
    dx[id].jp.x[ij] = xx;
    dx[id].score = dely[yy];
}
if (xx == len0 && yy < len1) {
    /* last col
    */
    if (endgaps)
        coll[yy] -= ins0+ins1*(len1-yy);
    if (coll[yy] > smax) {
        smax = coll[yy];
        dmax = id;
    }
}
}
if (endgaps && xx < len0)
    coll[yy-1] -= ins0+ins1*(len0-xx);
if (coll[yy-1] > smax) {
    smax = coll[yy-1];
    dmax = id;
}
tmp = col0; col0 = col1; col1 = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)col1);
}

```

TABLE 5 (cont.)

```

/*
 *
 * print() -- only routine visible outside this module
 *
 * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
 * nums() -- put out a number line: dumpblock()
 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */

#include "nw.h"

#define SPC 3
#define P_LINE 256 /* maximum output line */
#define P_SPC 3 /* space between name or num and seq */

extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */

print() print
{
    int lx, ly, firstgap, lastgap; /* overlap */

    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}

```

TABLE 5 (cont.)

```

/*
 * trace back the best path, count matches
 */
static
getmat(lx, ly, firstgap, lastgap)  getmat
    int      lx, ly;                /* "core" (minus endgaps) */
    int      firstgap, lastgap;     /* leading trailing overlap */
{
    int      nm, i0, i1, siz0, siz1;
    char     outx[32];
    double   pct;
    register n0, n1;
    register char *p0, *p1;

    /* get total matches, score
     */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;

    nm = 0;
    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
        else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
        else {
            if (xbm[*p0-'A']&xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
    }

    /* pct homology:
     * if penalizing endgaps, base is the shorter seq
     * else, knock off overhangs and take shorter core
     */
    if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
        lx = (lx < ly)? lx : ly;
    pct = 100.*(double)nm/(double)lx;
    fprintf(fx, "\n");
    fprintf(fx, "< %d match%s in an overlap of %d: %.2f percent similarity\n",
            nm, (nm == 1)? "" : "es", lx, pct);
}

```

TABLE 5 (cont.)

```

fprintf(fx, "<gaps in first sequence: %d", gapx);      ...getmat
if (gapx) {
    (void) sprintf(outx, " (%d %s%s)",
        ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outx);

    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, " (%d %s%s)",
            ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
    if (dna)
        fprintf(fx,
            "\n<score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINS0, DINS1);
    else
        fprintf(fx,
            "\n<score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINS0, PINS1);
    if (endgaps)
        fprintf(fx,
            "<endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
            lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
    else
        fprintf(fx, "<endgaps not penalized\n");
}

static      nm;          /* matches in core -- for checking */
static      lmax;       /* lengths of stripped file names */
static      ij[2];      /* jmp index for a path */
static      nc[2];      /* number at start of current line */
static      ni[2];      /* current elem number -- for gapping */
static      siz[2];
static char *ps[2];     /* ptr to current element */
static char *po[2];     /* ptr to next output char slot */
static char out[2][P_LINE]; /* output line */
static char star[P_LINE]; /* set by stars() */

/*
 * print alignment of described in struct path pp[]
 */
static
pr_align() pr_align
{
    int      nn;         /* char count */
    int      more;
    register i;

    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(namex[i]);
        if (nn > lmax)
            lmax = nn;

        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
    }
}

```

TABLE 5 (cont.)

```

for (nn = nm = 0, more = 1; more; ) {      ...pr_align
    for (i = more = 0; i < 2; i++) {
        /*
         * do we have more of this sequence?
         */
        if (!*ps[i])
            continue;

        more++;

        if (pp[i].spc) {                    /* leading space */
            *po[i]++ = ' ';
            pp[i].spc--;
        }
        else if (siz[i]) {                  /* in a gap */
            *po[i]++ = '-';
            siz[i]--;
        }
        else {                              /* we're putting a seq element
         */
            *po[i] = *ps[i];
            if (islower(*ps[i]))
                *ps[i] = toupper(*ps[i]);

            po[i]++;
            ps[i]++;

            /*
             * are we at next gap for this seq?
             */
            if (ni[i] == pp[i].x[ij[i]]) {
                /*
                 * we need to merge all gaps
                 * at this location
                 */
                siz[i] = pp[i].n[ij[i]++];
                while (ni[i] == pp[i].x[ij[i]])
                    siz[i] += pp[i].n[ij[i]++];
            }
            ni[i]++;
        }
    }
    if (++nn == olen || !more && nn) {
        dumpblock();
        for (i = 0; i < 2; i++)
            po[i] = out[i];
        nn = 0;
    }
}

/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
static
dumpblock()      dumpblock
{
    register i;

    for (i = 0; i < 2; i++)
        *po[i]-- = '^0';
}

```

TABLE 5 (cont.)

...dumpblock

```

(void) putc('\n', fx);
for (i = 0; i < 2; i++) {
    if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
        if (i == 0)
            nums(i);
        if (i == 0 && *out[1])
            stars();
        putline(i);
        if (i == 0 && *out[1])
            fprintf(fx, star);
        if (i == 1)
            nums(i);
    }
}

/*
 * put out a number line: dumpblock()
 */
static
nums(ix)  nums
int      ix;      /* index in out[] holding seq line */
{
    char      nline[P_LINE];
    register  i, j;
    register  char *pn, *px, *py;

    for (pn = nline, i = 0; i < lmax+P_SPC; i++, pn++)
        *pn = ' ';
    for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
        if (*py == ' ' || *py == '-')
            *pn = ' ';
        else {
            if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                j = (i < 0)? -i : i;
                for (px = pn; j /= 10, px--)
                    *px = j%10 + '0';
                if (i < 0)
                    *px = '-';
            }
            else
                *pn = ' ';
            i++;
        }
    }
    *pn = '\0';
    nc[ix] = i;
    for (pn = nline; *pn; pn++)
        (void) putc(*pn, fx);
    (void) putc('\n', fx);
}

/*
 * put out a line (name, [num], seq, [num]): dumpblock()
 */
static
putline(ix)  putline
int         ix;
{

```

Page 5 of nwprint.c

TABLE 5 (cont.)

...putline

```

int          i;
register char *px;

for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
    (void) putc(*px, fx);
for (; i < lmax+P_SPC; i++)
    (void) putc(' ', fx);

/* these count from 1:
 * ni[] is current element (from 1)
 * nc[] is number at start of current line
 */
for (px = out[ix]; *px; px++)
    (void) putc(*px&0x7F, fx);
(void) putc('\n', fx);
}

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock()
 */
static
stars()
{
    int          i;
    register char *p0, *p1, cx, *px;

    if (!*out[0] || (*out[0] == ' ' && *(po[0]) == ' ') ||
        !*out[1] || (*out[1] == ' ' && *(po[1]) == ' '))
        return;
    px = star;
    for (i = lmax+P_SPC; i; i--)
        *px++ = ' ';

    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
        if (isalpha(*p0) && isalpha(*p1)) {
            if (xbrm[*p0-'A']&xbrm[*p1-'A']) {
                cx = '*';
                nm++;
            }
            else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                cx = '!';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
    *px = '\0';
}

```

TABLE 5 (cont.)

```
/*
 * strip path or prefix from pn, return len: pr_align()
 */
static
stripname(pn)      stripname
char      *pn;     /* file name (may be path) */
{
    register char    *px, *py;

    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
            py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
}
```

TABLE 5 (cont.)

```

/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
 * g_alloc() -- calloc() with error checkin
 * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>

char    *jname = "/tmp/homgXXXXXX";          /* tmp file for jumps */
FILE    *fj;

int     cleanup();                          /* cleanup tmp file */
long    lseek();

/*
 * remove any tmp file if we blow
 */
cleanup(i) cleanup
    int    i;
{
    if (fj)
        (void) unlink(jname);
    exit(i);
}

/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char    *
getseq(file, len)    getseq
    char    *file;    /* file name */
    int     *len;     /* seq len */
{
    char    line[1024], *pseq;
    register char    *px, *py;
    int     natgc, tlen;
    FILE    *fp;

    if ((fp = fopen(file, "r")) == 0) {
        fprintf(stderr, "%s: can't read %s\n", prog, file);
        exit(1);
    }
    tlen = natgc = 0;
    while (fgets(line, 1024, fp)) {
        if (*line == ';' || *line == '<' || *line == '>')
            continue;
        for (px = line; *px != '\n'; px++)
            if (isupper(*px) || islower(*px))
                tlen++;
    }
    if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
        fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
        exit(1);
    }
    pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';

```

TABLE 5 (cont.)

...getseq

```

py = pseq + 4;
*len = tlen;
rewind(fp);

while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
            natgc++;
    }
    *py++ = '\0';
    *py = '\0';
    (void) fclose(fp);
    dna = natgc > (tlen/3);
    return(pseq+4);
}

char *
g_alloc(msg, nx, sz) g_alloc
char *msg;          /* program, calling routine */
int nx, sz;         /* number and size of elements */
{
    char *px, *calloc();

    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g_alloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}

/*
 * get final jmps from dx[] or tmp file, set pp[], reset dmax: main()
 */
readjmps() readjmps
{
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;

    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; ; i++) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
                ;

```

Page 2 of nwsubr.c

TABLE 5 (cont.)

```

...readjumps
    if (j < 0 && dx[dmax].offset && fj) {
        (void) lseek(fd, dx[dmax].offset, 0);
        (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
        (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
        dx[dmax].ijmp = MAXJMP-1;
    }
    else
        break;
}
if (i >= JMPS) {
    fprintf(stderr, "%s: too many gaps in alignment\n", prog);
    cleanup(1);
}
if (j >= 0) {
    siz = dx[dmax].jp.n[j];
    xx = dx[dmax].jp.x[j];
    dmax += siz;
    if (siz < 0) { /* gap in second seq */
        pp[1].n[i1] = -siz;
        xx += siz;
        /* id = xx - yy + len1 - 1
        */
        pp[1].x[i1] = xx - dmax + len1 - 1;
        gapy++;
        ngapy -= siz;
/* ignore MAXGAP when doing endgaps */
        siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
        i1++;
    }
    else if (siz > 0) { /* gap in first seq */
        pp[0].n[i0] = siz;
        pp[0].x[i0] = xx;
        gapx++;
        ngapx += siz;
/* ignore MAXGAP when doing endgaps */
        siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
        i0++;
    }
}
else
    break;
}
/* reverse the order of jumps
*/
for (j = 0, i0--; j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--; j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}
}

```

TABLE 5 (cont.)

```

5  /*
   * write a filled jmp struct offset of the prev one (if any): nw()
   */
writejumps(ix)      writejumps
   int             ix;
{
10     char          *mktemp();

   if (!fj) {
15         if (mktemp(jname) < 0) {
               fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
               cleanup(1);
           }
           if ((fj = fopen(jname, "w")) == 0) {
               fprintf(stderr, "%s: can't write %s\n", prog, jname);
               exit(1);
           }
20     }
   (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
   (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}

```

TABLE 6

MARRSRHRLLLLLLRYLVVALGYHKAYGFSAPKDQQVVTAVEYQEAILACKTPKKTVSS  
 RLEWKKLGRSVSFVYYQQTLOGDFKNRAEMIDFNIRIKNVTRSDAGKYRCEVSAPSEQG  
 5 QNLEEDTVTLEVLVAPAVPSCEVPSSALSGTVVELRCQDKEGNPAPEYTWFKDGIKLLLE  
 NPRLGSQSTNSSYTMNTKTGTLQFNVTVSKLDTGEYSCEARNSVGYRRCPGKRMQVDDLN  
 ISGIIAAVVVVALVISVCGLGVCYAQRKGYFSKETSFOKSNSSSKATTMSENVQWLTPV  
 IPALWAAAAGGSRGQEF

10 N-glycosylation siteat residues:

98-102

187-191

236-240

277-281

15

Casein kinase II phosphorylation siteat residues:

39-43

59-63

100-104

20

149-153

205-209

284-288

N-myristoylation siteat residues:

25

182-188

239-245

255-261

257-263

305-311

30

Amidation site at residues:

226-230

TABLE 7

MARRSAFPAAALWLWSILLCLLALRAEAGPPQEESELYLWIDAHQARVLIGFEEDILIVS  
 EGKMAPFTHDFRKAQORMPAIPVNIHSMNFTWQAAGQAEYFYEFSLRSLDKGIMADPT  
 5 VNVPLLGTVPHKASVVQVGFPCLGKQDGVAAFEVDVIVMNSEGNTILQTPQNAIFFKTC  
 QQAECPPGGCRNGGFCNERRICECPDGFHGHCEKALCTPRCMNGGLCVTPGFCICPPGF  
 YGVNCDKANCSTTCFNGGTCFYPGKICPPGLEGEQCEISKCPQPCRNGGKCIGKSKCK  
 CSKGYQGDLCSPKPVCEPGCGAHGTCHPNKCQCQEGWHGRHCNKRYEASLIHALRPAGA  
 10 QLRQHTPSLKKAEERRDPPESNYIW

N-glycosylation site at residues:

88-92

245-249

15 Casein kinase II phosphorylation site at residues:

319-323

Tyrosine kinase phosphorylation site at residues:

370-378

20 N-myristoylation site at residues:

184-190

185-191

189-195

25 315-321

ATP/GTP-binding site motif A (P-loop) at residues:

285-293

30 EGF-like domain cysteine pattern signature at residues:

198-210

230-242

262-274

294-306

35 326-338

TABLE 8

MGTKAQVERKLLCLFILAILLCSLALGSVTVHSSEPEVRI PENNPVKLSCAYSGFSSPR  
 VEWKFDQGD TTRLVLCYNNKITASYEDRVTF LPTGITFKSVTREDTGTYT CMVSEEGGNS  
 5 YGEVKVKLIVLVPPSKPTVNI PSSATIGNRAVLTCSEQDGSPPSEYTWFKDGIVMPTNP  
 KSTRAFSNSSYVLNPTTGELVFDPLSASDTGEYSCEARNGYGT PMTSNAVRMEAVERNV  
 GVIVA AVLVT LILLGILVFGIWFAYS RGHFDRTKKGTSSKKVIYSQPSARSEGEFKQTS  
 SFLV

10 N-glycosylation site at residues:  
 185-189

cAMP- and cGMP-dependent protein kinase phosphorylation site at  
 residues:  
 15 270-274

Casein kinase II phosphorylation site at residues:  
 34-38  
 82-86  
 20 100-108  
 118-122  
 152-156  
 154-158  
 193-197  
 25 203-207  
 287-291

N-myristoylation site at residues:  
 105-111  
 30 116-122  
 158-164  
 219-225  
 237-243  
 256-262

TABLE 9

5 MLLWILLLETSLCFAAGNVTGDVCKEKICSCNEIEGDLHVDCEKKGFTSLQRFTAPTSQ  
 FYHLFLHGNSLTRLFPNEFANFYNAVSLHMENGLHEIVPGAFLGLQLVKRLHINNKI  
 KSRKQTFGLGDDLEYLQADFNLLRDIDPGAFQDLNKLEVLILNDNLISTLPANVFQYV  
 PITHLDLRGNRLKTLPEEVELEQIPGIAEILLEDNPWDCTCDLLSLKEWLENIIPKNALI  
 GRVVCEAPTRLQGKDLNETTEQDLCPLKNRVDSSLPAPPAQEETFAPGPLPTPFKTNGQ  
 EDHATPGSAPNGGTKIPGNWQIKIRPTAAIATGSSRNKPLANSLPCPGGCSCDHIPGSG  
 LKMNCNNRNVSADLKPKLSNVQELFLRDNKIHSIRKSHFVDYKNLILLDLGNNNIAT  
 10 VENNTFKNLLDLRWLYMDSNYLDTLSREKFAGLQNLLEYLNVEYNAIQILPGTFNAMPK  
 LRILILNLLRSLPVDVVFAGVSLSKLSLHNNYFMYLPVAGVLDQLTSIIQIDLHGNPW  
 ECSCTIVPFKQWAERLGSEVLMSDLKCETPVNFFRKDFMLLSNDEICPQLYARISPTLT  
 SHSKNSTGLAETGTHSNSYLDTSRVSISVLVPGLLLVFVTSFTVVGMLVFILRNKRKRS  
 KRRDANSSASEINSLQTVCDSSYWHNGPYNADGAHRVYDCGSHSLSD

15 N-glycosylation site at residues:

- 18-22
- 253-257
- 363-367
- 20 416-420
- 595-599
- 655-659

25 cAMP- and cGMP-dependent protein kinase phosphorylation site at residues:

- 122-126
- 646-650

30 Casein kinase II phosphorylation site at residues:

- 30-34
- 180-184
- 222-226
- 256-260
- 366-370
- 35 573-577
- 608-612
- 657-661
- 666-670

693-697

N-myristoylation site at residues:

5           17-23  
            67-73  
            100-106  
            302-308  
            328-334  
            343-349  
10          354-360  
            465-471  
            493-499  
            598-604  
            603-609

15

Prokaryotic membrane lipoprotein lipid attachment site at residues:

337-348

TABLE 10

MVDVLLLFSLCLLFHISRPDLSHNRLSFIKASSMSHLQSLREVKLNNELETIPNLGPV  
 SANITLLSLAGNRIVEILPEHLKEFQSLETLDLSSNNISELQTAFPALQLKYLYLNSNR  
 VTSMEPGYFDNLANTLLVLKLNRRNRISAIPPKMFKLPQLQHLELNRNKIKNVDGLTFQG  
 5 LGALKSLKMQRNGVTKLMDGAFWGLSNMEILQLDHNNLTEITKGWLYGLLMLQELHLSQ  
 NAINRISPDAWEFCQKLSLDELTFNHL SRLDDSSFLGSLLLNTLHIGNNRVSYIADCAF  
 RGLSSLKTLDLKNNEISWTIEDMNGAFSGLDKLRRLILQGNRIRSITKKAFTGLDALEH  
 LDLSDNAIMSLOGNAFSQMKKLOQLHLNTSSLLCDCQLKWLPOQVAENNFQSFVNASCA  
 HPQLLKGRSIFAVSPDGFVCDDFPKPQITVQPETQSAIKGSNLSFICSAASSSDSPMTF  
 10 AWKKDNELLHDAEMENYAHLRAQGGVMEYTTILRLREVEFASEGKYQCVISNHFGSSY  
 SVKAKLTVNMLPSFTKTPMDLTIRAGAMARLECAAVGHPAPQIAWQKDGGTDFPAARER  
 RMHVMPEDDVFFIVDVKIEDIGVYSCTAQN SAGSISANATLTVLETSPFLRPLLDRTVT  
 KGETAVLQCIAGGSPPPKLNWTKDDSPVVTERHFFAAGNQLLIIVDSVSDAGKYTCE  
 MSNTLGTGERGNVRLSVIPTPTCDSPQMTAPSLDDDGWATVGVVIIAVVCCVVGTSLVVW  
 15 VIIYHTRRRNEDCSITNTDETNLPADIPSYLSSQGTADRQDGYVSSESGSHHQFVTSS  
 GAGFFLPQHDSSTCHIDNSSEADVEAATDLFLCPFLGSTGPMYKGNVYGSDFPETYH  
 TGCSPPDRPTVLMDHYEPSYIKKKECYPCSHPSEESCERSFSNISWPSHVRKLLNTSYSH  
 NEGPGMKNLCLNKSSLD FSANPEPASVASSNSFMGTFGKALRRPHLDAYS SFGQPSDCQ  
 PRAFYLKAHSSPDLDSGSEEDGKERTDFQEEHNICTFKQTLNRYRTPNFQSYDLDT

20

N-glycosylation site at residues:

62-66

96-100

214-218

25

382-386

409-413

455-459

628-632

669-673

30

845-849

927-931

939-943

956-960

35

Glycosaminoglycan attachment site at residues:

826-830

Casein kinase II phosphorylation site at residues:

17-21

39-43  
120-124  
203-207  
254-258  
5 264-268  
314-318  
323-327  
347-351  
464-468  
10 548-552  
632-636  
649-653  
671-675  
739-743  
15 783-787  
803-807  
847-851  
943-947  
958-962  
20 1013-1017  
1019-1023  
1021-1025

Tyrosine kinase phosphorylation site at residues:

25 607-615

N-myristoylation site at residues:

179-185  
197-203  
30 320-326  
367-373  
453-459  
528-534  
612-618  
35 623-629  
714-720  
873-879

TABLE 11

MLNKMTLHPQQIMIGPRFNRALFDPLLVLVLLALQLLVVAGLVRAQTCPSVCSNCFQFSK  
 VICVRKNLREVPDGI STNTRLLNLHENQIQI IKVNSFKHLRHLEILQLSRNHIRTIEIG  
 5 AFNGLANLNTLELFDNRLTTIPNGAFVYLSKLELWLRNPNIESIPSYAFNRIPSLRRL  
 DLGELKRLSYI SEGAFEGLSNLRYLNLAMCNLREIPNL TPLIKLDELDSLGNHLSAIRP  
 GSFOGLMHLQKLWMIQSQIQVIERNAFDNLQSLVEINLAHNNLTLLPHDLFTPLHHLER  
 IHLHHPWNCNCDILWLSWWIKDMAPSNTACCARCNTPPNLKGRYIGELDQNYFTCYAP  
 VIVEPPADLNVTEGMAAELKCRAS TSLTSVSWITPNGTVMTHGAYKVR IAVLSDGTLNF  
 10 TNVTVQDTGMYTCMVSN SVGN TTASATLNVTAATTT PFSYFSTVTVETMEPSQDEARTT  
 DNNVGPTPVVDWETTNTTSLTPQSTRST EKTFTIPVTDINSGIPGIDEVMKTTKIIIG  
 CFVAITLMAAVMLVIFYKMRKQHRQNH HAPTRTVEI INV DDEITGDTPMESHLPMPAI  
 EHEHLNHYNSYKSPFNHTTTVNTINSI HSSVHEPLLIRMNSKDNVQETQI

N-glycosylation site at residues:

- 278-282
- 364-368
- 390-394
- 412-416
- 415-419
- 434-438
- 442-446
- 488-492
- 606-610

cAMP- and cGMP-dependent protein kinase phosphorylation site at residues:

- 183-187

Casein kinase II phosphorylation site at residues:

- 268-272
- 417-421
- 465-469
- 579-583
- 620-624

N-myristoylation site at residues:

- 40-46
- 73-79

118-124

191-197

228-234

237-243

5

391-397

422-428

433-439

531-537

TABLE 12

MSAPSLRARAAGLGLLLCAVLGRAGRSDSGRGELGQPSGVAAERPCPTTCRCLGDLDD  
 CSRKRLARLPEPLPSWVARLDLSHNRLSFIKASSMSHLQSLREVKLNNNELETIPNLGP  
 5 VSANITLLSLAGNRIVEILPEHLKEFQSLETLDLSSNNISELQTAFFPALQLKYLYLNSN  
 RVTSMEPGYFDNLANTLLVLKLNRRNRISAIPPKMFKLPQLQHLELNRNKIKNVDGLTFQ  
 GLGALKSLKMQRNGVTKLMDGAFWGLSNMEILQLDHNNLTEITKGWLYGLLMLQELHLS  
 QNAINRISPDWEFCQKLSELDLTFNHLSRLDDSSFLGLSLLNLTHIGNNRVSYIADCA  
 FRGLSSKLTLDLKNNEISWTIEDMNGAFSGLDKLRRLILQGNRIRSIKKAFTGLDALE  
 10 HLDLSDNAIMSLQGNAFSQMKKLQQLHLNLTSSLLCDCQLKWLPOVAENNFQSFVNASC  
 AHPQLLKGRSIFAVSPDGFVCDDFPKPQITVQPETQSAIKGSNLSFICSAASSSDSPMT  
 FAWKKNELLHDAEMENYAHLRAQGGVMEYTTILRLREVEFASEGKYQCVISNHFSS  
 YSVKAKLTVNMLPSFTKTPMDLTIRAGAMARLECAAVGHPAPQIAWQKDGTDFFPAARE  
 RRMHVMPEDDVFFIVDVKIEDIGVYSCTAONSAGSISANATLTVLETSPFLRPLLDRTV  
 15 TKGETAVLQCIAGGSPPKLNWTKDSDPLVVTERHFFAAGNQLLIIVDSDVSDAGKYTC  
 EMSNTLGTERRGNVRLSVIPTPTCDSPQMTAPSLDDDGWATVGVVIAVCCVVGTSLVW  
 VVIIYHTRRRNEDCSITNTDETNLPADIPSYLSSQGTLADRQDGYVSSSESGSHHQFVTS  
 SGAGFFLPQHDSSGTCHIDNSSEADVEAATDLFLCPFLGSTGPMYKGNVYGSDDPFETY  
 HTGCSDPRTVLMDHYEPSYIKKKECYPCHPSEESCERSFSNISWPSHVRKLLNTSYS  
 20 HNEGPGMKNLCLNKSSLDIFSANPEPASVASSNSFMGTFGKALRRPHLDAYSSFGQPSDC  
 QPRAFYLKAHSSPDLDGSEEDGKERTDFQEENHICTFKQTLNRYRTPNFQSYDLDT

N-glycosylation site at residues:

- 122-126
- 25 156-160
- 274-278
- 442-446
- 469-473
- 515-519
- 30 688-692
- 729-733
- 905-909
- 987-991
- 999-1003
- 35 1016-1020

Glycosaminoglycan attachment site at residues:

- 886-890

## Casein kinase II phosphorylation site at residues:

99-103  
180-184  
5 263-267  
314-318  
324-328  
374-378  
383-387  
10 407-411  
524-528  
608-612  
692-696  
709-713  
15 731-735  
799-803  
843-847  
863-867  
907-911  
20 1003-1007  
1018-1022  
1073-1077  
1079-1083  
1081-1085

25

## Tyrosine kinase phosphorylation site at residue:

667-675

## N-myristoylation site at residues:

30 14-20  
36-42  
239-245  
257-263  
380-386  
35 427-433  
513-519  
588-594  
672-678  
683-689

774-780

933-999

Leucine zipper pattern at residues:

5

58-80

65-87

What is claimed:

1. A composition, comprising a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide, agonist or fragment thereof and a carrier or excipient, useful for:
  - (a) increasing infiltration of inflammatory cells into a tissue of a mammal in need thereof,
  - (b) stimulating or enhancing an immune response in a mammal in need thereof, or
  - (c) increasing the proliferation of T-lymphocytes in a mammal in need thereof in response to an antigen.
  
2. Use of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide, agonist or a fragment thereof to prepare a composition useful for:
  - (a) increasing infiltration of inflammatory cells into a tissue of a mammal in need thereof,
  - (b) stimulating or enhancing an immune response in a mammal in need thereof, or
  - (c) increasing the proliferation of T-lymphocytes in a mammal in need thereof in response to an antigen.
  
3. A composition, comprising a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide, antagonist or a fragment thereof and a carrier or excipient, useful for:
  - (a) decreasing infiltration of inflammatory cells into a tissue of a mammal in need thereof,
  - (b) inhibiting or reducing an immune response in a mammal in need thereof, or
  - (c) decreasing the proliferation of T-lymphocytes in a mammal in need thereof in response to an antigen.
  
4. Use of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide, antagonist or a fragment thereof to prepare a composition useful for:
  - (a) decreasing infiltration of inflammatory cells into a tissue of a mammal in need thereof,
  - (b) inhibiting or reducing an immune response in a mammal in need thereof, or
  - (c) decreasing the proliferation of T-lymphocytes in a mammal in need thereof in response to an antigen.
  
5. A method of treating an immune related disorder, such as a T cell mediated

disorder, in a mammal in need thereof, comprising administering to the mammal an effective amount of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide, an agonist antibody thereof, an antagonist antibody thereto, or a fragment thereof.

6. The method of claim 5, wherein the disorder is selected from systemic lupus erythematosus, rheumatoid arthritis, juvenile chronic arthritis, spondyloarthropathies, systemic sclerosis (scleroderma), idiopathic inflammatory myopathies (dermatomyositis, polymyositis), Sjogren's syndrome, systemic vasculitis, sarcoidosis, autoimmune hemolytic anemia (immune pancytopenia, paroxysmal nocturnal hemoglobinuria), autoimmune thrombocytopenia (idiopathic thrombocytopenic purpura, immune-mediated thrombocytopenia), thyroiditis (Grave's disease, Hashimoto's thyroiditis, juvenile lymphocytic thyroiditis, atrophic thyroiditis), diabetes mellitus, immune-mediated renal disease (glomerulonephritis, tubulointerstitial nephritis), demyelinating diseases of the central and peripheral nervous systems such as multiple sclerosis, idiopathic demyelinating polyneuropathy or Guillain-Barré syndrome, and chronic inflammatory demyelinating polyneuropathy, hepatobiliary diseases such as infectious hepatitis (hepatitis A, B, C, D, E and other non-hepatotropic viruses), autoimmune chronic active hepatitis, primary biliary cirrhosis, granulomatous hepatitis, and sclerosing cholangitis, inflammatory and fibrotic lung diseases such as inflammatory bowel disease (ulcerative colitis: Crohn's disease), gluten-sensitive enteropathy, and Whipple's disease, autoimmune or immune-mediated skin diseases including bullous skin diseases, erythema multiforme and contact dermatitis, psoriasis, allergic diseases such as asthma, allergic rhinitis, atopic dermatitis, food hypersensitivity and urticaria, immunologic diseases of the lung such as eosinophilic pneumonias, idiopathic pulmonary fibrosis and hypersensitivity pneumonitis, transplantation associated diseases including graft rejection and graft -versus-host-disease.

7. The composition or use of any of the preceding claims, wherein the antibody is a monoclonal antibody.

8. The composition or use of any of the preceding claims, wherein the antibody is an antibody fragment or a single-chain antibody.

9. The composition or use of any of the preceding claims, wherein the antibody has nonhuman complementarity determining region (CDR) residues and human framework region (FR) residues.

10. A method for determining the presence of a PRO245, PRO217, PRO301,

PRO266, PRO335, PRO331 or PRO326 polypeptide, comprising exposing a cell suspected of containing the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide to an anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody and determining binding of the antibody to the cell.

11. A method of diagnosing an immune related disease in a mammal, comprising detecting the level of expression of a gene encoding a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide (a) in a test sample of tissue cells obtained from the mammal, and (b) in a control sample of known normal tissue cells of the same cell type, wherein a higher expression level in the test sample indicates the presence of immune related disease in the mammal from which the test tissue cells were obtained.

12. A method of diagnosing an immune related disease in a mammal, comprising (a) contacting an anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody with a test sample of tissue cells obtained from the mammal, and (b) detecting the formation of a complex between the antibody and the polypeptide in the test sample.

13. An immune related disease diagnostic kit, comprising an anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody or fragment thereof and a carrier in suitable packaging.

14. The kit of claim 13, further comprising instructions for using the antibody to detect a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide.

15. An article of manufacture, comprising:  
a container;  
a label on the container; and  
a composition comprising an active agent contained within the container; wherein the composition is effective for stimulating or enhancing an immune response in a mammal. the label on the container indicates that the composition can be used for treating an immune related disease, and the active agent in the composition is an agent inhibiting the expression and/or activity of a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide.

16. The article of manufacture of claim 21 wherein said active agent is an anti-PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 antibody.

17. A method for identifying a compound capable of inhibiting the expression or activity of a PRO245 polypeptide, comprising contacting a candidate compound with a PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide under conditions and for a time sufficient to allow these two components to interact.

18. The method of claim 17, wherein the candidate compound or the PRO245, PRO217, PRO301, PRO266, PRO335, PRO331 or PRO326 polypeptide is immobilized on a solid support.

19. The method of claim 18, wherein the non-immobilized component carries a detectable label.

FIGURE 1A

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXYYYYYYY	(Length = 12 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

5 divided by 15 = 33.3%



FIGURE 1C

PRO-DNA	NNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLL	(Length = 16 nucleotides)

% nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

6 divided by 14 = 42.9%

FIGURE 1D

PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLLVV	(Length = 9 nucleotides)

% nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

4 divided by 12 = 33.3%

FIGURE 2A

```

/*
 *
 * C-C increased from 12 to 15
 * Z is average of EQ
 * B is average of ND
 * match with stop is _M; stop-stop = 0; J (joker) match = 0
 */
#define _M -8 /* value of a match with a stop */

int _day[26][26] = {
/* A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */ {2,0,-2,0,0,-4,1,-1,-1,0,-1,-2,-1,0,_M,1,0,-2,1,1,0,0,-6,0,-3,0},
/* B */ {0,3,-4,3,2,-5,0,1,-2,0,0,-3,-2,2,_M,-1,1,0,0,0,-2,-5,0,-3,1},
/* C */ {-2,-4,15,-5,-5,-4,-3,-3,-2,0,-5,-6,-5,-4,_M,-3,-5,-4,0,-2,0,-2,-8,0,0,-5},
/* D */ {0,3,-5,4,3,-6,1,1,-2,0,0,-4,-3,2,_M,-1,2,-1,0,0,0,-2,-7,0,-4,2},
/* E */ {0,2,-5,3,4,-5,0,1,-2,0,0,-3,-2,1,_M,-1,2,-1,0,0,0,-2,-7,0,-4,3},
/* F */ {-4,-5,-4,-6,-5,9,-5,-2,1,0,-5,2,0,-4,_M,-5,-5,-4,-3,0,-1,0,0,7,-5},
/* G */ {1,0,-3,1,0,-5,5,-2,-3,0,-2,-4,-3,0,_M,-1,-1,-3,1,0,0,-1,-7,0,-5,0},
/* H */ {-1,1,-3,1,1,-2,-2,6,-2,0,0,-2,-2,2,_M,0,3,2,-1,-1,0,-2,-3,0,0,2},
/* I */ {-1,-2,-2,-2,-2,1,-3,-2,5,0,-2,2,2,-2,_M,-2,-2,-2,-1,0,0,4,-5,0,-1,-2},
/* J */ {0,0,0,0,0,0,0,0,0,0,0,0,0,0,_M,0,0,0,0,0,0,0,0,0,0},
/* K */ {-1,0,-5,0,0,-5,-2,0,-2,0,5,-3,0,1,_M,-1,1,3,0,0,0,-2,-3,0,-4,0},
/* L */ {-2,-3,-6,-4,-3,2,-4,-2,2,0,-3,6,4,-3,_M,-3,-2,-3,-3,-1,0,2,-2,0,-1,-2},
/* M */ {-1,-2,-5,-3,-2,0,-3,-2,2,0,0,4,6,-2,_M,-2,-1,0,-2,-1,0,2,-4,0,-2,-1},
/* N */ {0,2,-4,2,1,-4,0,2,-2,0,1,-3,-2,2,_M,-1,1,0,1,0,0,-2,-4,0,-2,1},
/* O */ {_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,_M,0,_M,_M,_M,_M,_M,_M,_M,_M,_M},
/* P */ {1,-1,-3,-1,-1,-5,-1,0,-2,0,-1,-3,-2,-1,_M,6,0,0,1,0,0,-1,-6,0,-5,0},
/* Q */ {0,1,-5,2,2,-5,-1,3,-2,0,1,-2,-1,1,_M,0,4,1,-1,-1,0,-2,-5,0,-4,3},
/* R */ {-2,0,-4,-1,-1,-4,-3,2,-2,0,3,-3,0,0,_M,0,1,6,0,-1,0,-2,2,0,-4,0},
/* S */ {1,0,0,0,0,-3,1,-1,-1,0,0,-3,-2,1,_M,1,-1,0,2,1,0,-1,-2,0,-3,0},
/* T */ {1,0,-2,0,0,-3,0,-1,0,0,0,-1,-1,0,_M,0,-1,-1,1,3,0,0,-5,0,-3,0},
/* U */ {0,0,0,0,0,0,0,0,0,0,0,0,0,0,_M,0,0,0,0,0,0,0,0,0,0},
/* V */ {0,-2,-2,-2,-2,-1,-1,-2,4,0,-2,2,2,-2,_M,-1,-2,-2,-1,0,0,4,-6,0,-2,-2},
/* W */ {-6,-5,-8,-7,-7,0,-7,-3,-5,0,-3,-2,-4,-4,_M,-6,-5,2,-2,-5,0,-6,17,0,0,-6},
/* X */ {0,0,0,0,0,0,0,0,0,0,0,0,0,0,_M,0,0,0,0,0,0,0,0,0,0},
/* Y */ {-3,-3,0,-4,-4,7,-5,0,-1,0,-4,-1,-2,-2,_M,-5,-4,-4,-3,-3,0,-2,0,0,10,-4},
/* Z */ {0,1,-5,2,3,-5,0,2,-2,0,0,-2,-1,1,_M,0,3,0,0,0,0,-2,-6,0,-4,4}
};

```

## FIGURE 2B

```

/*
*/
#include <stdio.h>
#include <ctype.h>

#define MAXJMP 16 /* max jumps in a diag */
#define MAXGAP 24 /* don't continue to penalize gaps larger than this */
#define JMPS 1024 /* max jmps in a path */
#define MX 4 /* save if there's at least MX-1 bases since last jmp */

#define DMAT 3 /* value of matching bases */
#define DMIS 0 /* penalty for mismatched bases */
#define DINS0 8 /* penalty for a gap */
#define DINS1 1 /* penalty per base */
#define PINS0 8 /* penalty for a gap */
#define PINS1 4 /* penalty per residue */

struct jmp {
    short n[MAXJMP]; /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
}; /* limits seq to 2^16 -1 */

struct diag {
    int score; /* score at last jmp */
    long offset; /* offset of prev block */
    short ijmp; /* current jmp index */
    struct jmp jp; /* list of jmps */
};

struct path {
    int spc; /* number of leading spaces */
    short n[JMPS]; /* size of jmp (gap) */
    int x[JMPS]; /* loc of jmp (last elem before gap) */
};

char *ofile; /* output file name */
char *namex[2]; /* seq names: getseqs() */
char *prog; /* prog name for err msgs */
char *seqx[2]; /* seqs: getseqs() */
int dmax; /* best diag: nw() */
int dmax0; /* final diag */
int dna; /* set if dna: main() */
int endgaps; /* set if penalizing end gaps */
int gapx, gapy; /* total gaps in seqs */
int len0, len1; /* seq lens */
int ngapx, ngapy; /* total size of gaps */
int smax; /* max score: nw() */
int *xbm; /* bitmap for matching */
long offset; /* current offset in jmp file */
struct diag *dx; /* holds diagonals */
struct path pp[2]; /* holds path for seqs */

char *calloc(), *malloc(), *index(), *strcpy();
char *getseq(), *g_calloc();

```

## FIGURE 2C

```

/* Needleman-Wunsch alignment program
*
* usage: progs file1 file2
* where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
#include "day.h"

static  _dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};

static  _pbval[26] = {
    1, 2|(1<<('D'-'A'))|(1<<('N'-'A')), 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
    1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
};

main(ac, av)          main
    int               ac;
    char              *av[];
{
    prog = av[0];
    if (ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';', '>' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)?_dbval : _pbval;

    endgaps = 0;                               /* 1 to penalize endgaps */
    ofile = "align.out";                       /* output file */

    nw();                                       /* fill in the matrix, get the possible jmps */
    readjmps();                                /* get the actual jmps */
    print();                                   /* print stats, alignment */

    cleanup(0);                                /* unlink any tmp files */
}

```

FIGURE 2D

```

/* do the alignment, return best score: main()
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
* pro: PAM 250 values
* When scores are equal, we prefer mismatches to any gap, prefer
* a new gap to extending an ongoing gap, and prefer a gap in seqx
* to a gap in seq y.
*/
nw()    nw
{
    char          *px, *py;          /* seqs and ptrs */
    int           *ndely, *dely;     /* keep track of dely */
    int           ndelx, delx;       /* keep track of delx */
    int           *tmp;              /* for swapping row0, row1 */
    int           mis;               /* score for each type */
    int           ins0, ins1; /* insertion penalties */
    register      id;                /* diagonal index */
    register      ij;                /* jmp index */
    register      *col0, *col1;      /* score for curr, last row */
    register      xx, yy;            /* index into seqs */

    dx = (struct diag *)g_malloc("to get diags", len0+len1+1, sizeof(struct diag));

    ndely = (int *)g_malloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_malloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_malloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_malloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;

    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;          /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
    */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
        */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0] - ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }
}

```

## FIGURE 2E

...nw

```

for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
  mis = col0[yy-1];
  if (dna)
    mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
  else
    mis += _day[*px-'A'][*py-'A'];

  /* update penalty for del in x seq;
   * favor new del over ongong del
   * ignore MAXGAP if weighting endgaps
   */
  if (endgaps || ndely[yy] < MAXGAP) {
    if (col0[yy] - ins0 >= dely[yy]) {
      dely[yy] = col0[yy] - (ins0+ins1);
      ndely[yy] = 1;
    } else {
      dely[yy] = ins1;
      ndely[yy]++;
    }
  } else {
    if (col0[yy] - (ins0+ins1) >= dely[yy]) {
      dely[yy] = col0[yy] - (ins0+ins1);
      ndely[yy] = 1;
    } else
      ndely[yy]++;
  }

  /* update penalty for del in y seq;
   * favor new del over ongong del
   */
  if (endgaps || ndelx < MAXGAP) {
    if (col1[yy-1] - ins0 >= delx) {
      delx = col1[yy-1] - (ins0+ins1);
      ndelx = 1;
    } else {
      delx = ins1;
      ndelx++;
    }
  } else {
    if (col1[yy-1] - (ins0+ins1) >= delx) {
      delx = col1[yy-1] - (ins0+ins1);
      ndelx = 1;
    } else
      ndelx++;
  }

  /* pick the maximum score; we're favoring
   * mis over any del and delx over dely
   */
}

```

FIGURE 2F

```

...nw
id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    coll[yy] = mis;
else if (delx >= dely[yy]) {
    coll[yy] = delx;
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejmps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
        dx[id].jp.n[ij] = ndelx;
        dx[id].jp.x[ij] = xx;
        dx[id].score = delx;
    }
} else {
    coll[yy] = dely[yy];
    ij = dx[id].ijmp;
if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejmps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
        dx[id].jp.n[ij] = -ndely[yy];
        dx[id].jp.x[ij] = xx;
        dx[id].score = dely[yy];
    }
}
if (xx == len0 && yy < len1) {
    /* last col
    */
    if (endgaps)
        coll[yy] = ins0+ins1*(len1-yy);
    if (coll[yy] > smax) {
        smax = coll[yy];
        dmax = id;
    }
}
}
if (endgaps && xx < len0)
    coll[yy-1] = ins0+ins1*(len0-xx);
if (coll[yy-1] > smax) {
    smax = coll[yy-1];
    dmax = id;
}
}
tmp = col0; col0 = coll; coll = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)coll); }

```

FIGURE 2G

```

/*
 *
 * print() -- only routine visible outside this module
 *
 * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
 * nums() -- put out a number line: dumpblock()
 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */

#include "nw.h"

#define SPC 3
#define P_LINE 256 /* maximum output line */
#define P_SPC 3 /* space between name or num and seq */

extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */

print() print
{
    int lx, ly, firstgap, lastgap; /* overlap */

    if((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}

```

FIGURE 2H

```

/*
 * trace back the best path, count matches
 */
static
getmat(lx, ly, firstgap, lastgap)  getmat
    int      lx, ly;                /* "core" (minus endgaps) */
    int      firstgap, lastgap;     /* leading trailing overlap */
{
    int      nm, i0, i1, siz0, siz1;
    char     outx[32];
    double   pct;
    register n0, n1;
    register char *p0, *p1;

    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;

    nm = 0;
    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
        else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
        else {
            if (xbrm[*p0-'A']&xbrm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
    }

    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */
    if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
        lx = (lx < ly)? lx : ly;
    pct = 100.*((double)nm)/(double)lx;
    fprintf(fx, "\n");
    fprintf(fx, "< %d match%s in an overlap of %d: %.2f percent similarity\n",
            nm, (nm == 1)? "" : "es", lx, pct);
}

```

FIGURE 2I

```

fprintf(fx, "<gaps in first sequence: %d", gapx);      ...getmat
if (gapx) {
    (void) sprintf(outr, "(%d %s%s)",
        ngapx, (dna)? "base":"residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outr);

    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outr, "(%d %s%s)",
            ngapy, (dna)? "base":"residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outr);
    }
    if (dna)
        fprintf(fx,
            "\n<score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINS0, DINS1);
    else
        fprintf(fx,
            "\n<score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINS0, PINS1);
    if (endgaps)
        fprintf(fx,
            "<endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
            lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
    else
        fprintf(fx, "<endgaps not penalized\n");
}

static      nm;           /* matches in core -- for checking */
static      lmax;        /* lengths of stripped file names */
static      ij[2];       /* jmp index for a path */
static      nc[2];       /* number at start of current line */
static      ni[2];       /* current elem number -- for gapping */
static      siz[2];
static char *ps[2];      /* ptr to current element */
static char *po[2];      /* ptr to next output char slot */
static char out[2][P_LINE]; /* output line */
static char star[P_LINE]; /* set by stars() */

/*
 * print alignment of described in struct path pp[]
 */
static
pr_align() pr_align
{
    int      nn;          /* char count */
    int      more;
    register i;

    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(namex[i]);
        if (nn > lmax)
            lmax = nn;

        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
    }
}

```

FIGURE 2J

```

for (nn = nm = 0, more = 1; more; ) {      ...pr_align
    for (i = more = 0; i < 2; i++) {
        /*
         * do we have more of this sequence?
         */
        if (!*ps[i])
            continue;

        more++;

        if (pp[i].spc) { /* leading space */
            *po[i]++ = ' ';
            pp[i].spc--;
        }
        else if (siz[i]) { /* in a gap */
            *po[i]++ = '-';
            siz[i]--;
        }
        else { /* we're putting a seq element
            */
            *po[i] = *ps[i];
            if (islower(*ps[i]))
                *ps[i] = toupper(*ps[i]);
            po[i]++;
            ps[i]++;

            /*
             * are we at next gap for this seq?
             */
            if (ni[i] == pp[i].x[ij[i]]) {
                /*
                 * we need to merge all gaps
                 * at this location
                 */
                siz[i] = pp[i].n[ij[i]++];
                while (ni[i] == pp[i].x[ij[i]])
                    siz[i] += pp[i].n[ij[i]++];
            }
            ni[i]++;
        }
    }
    if (++nn == olen || !more && nn) {
        dumpblock();
        for (i = 0; i < 2; i++)
            po[i] = out[i];
        nn = 0;
    }
}

/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
static
dumpblock()    dumpblock
{
    register    i;

    for (i = 0; i < 2; i++)
        *po[i]-- = '\0';
}

```

FIGURE 2K

...dumpblock

```

(void) putc('\n', fx);
for (i = 0; i < 2; i++) {
    if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
        if (i == 0)
            nums(i);
        if (i == 0 && *out[1])
            stars();
        putline(i);
        if (i == 0 && *out[1])
            fprintf(fx, star);
        if (i == 1)
            nums(i);
    }
}

/*
 * put out a number line: dumpblock()
 */
static
nums(ix)  nums
int      ix;      /* index in out[] holding seq line */
{
    char      nline[P_LINE];
    register  i, j;
    register char *pn, *px, *py;

    for (pn = nline, i = 0; i < lmax+P_SPC; i++, pn++)
        *pn = ' ';
    for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
        if (*py == ' ' || *py == '-')
            *pn = ' ';
        else {
            if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                j = (i < 0)? -i : i;
                for (px = pn; j; j /= 10, px--)
                    *px = j%10 + '0';
                if (i < 0)
                    *px = '-';
            }
            else
                *pn = ' ';
            i++;
        }
    }
    *pn = '\0';
    nc[ix] = i;
    for (pn = nline; *pn; pn++)
        (void) putc(*pn, fx);
    (void) putc('\n', fx);
}

/*
 * put out a line (name, [num], seq, [num]): dumpblock()
 */
static
putline(ix)  putline
int          ix;
{

```

FIGURE 2L

```

...putline

int          i;
register char *px;

for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
    (void) putc(*px, fx);
for (; i < lmax+P_SPC; i++)
    (void) putc(' ', fx);

/* these count from 1:
 * ni[] is current element (from 1)
 * nc[] is number at start of current line
 */
for (px = out[ix]; *px; px++)
    (void) putc(*px&0x7F, fx);
(void) putc('\n', fx);
}

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock()
 */
static
stars()
{
    int          i;
    register char *p0, *p1, cx, *px;

    if (!*out[0] || (*out[0] == ' ' && *(po[0]) == ' ') ||
        !*out[1] || (*out[1] == ' ' && *(po[1]) == ' '))
        return;
    px = star;
    for (i = lmax+P_SPC; i; i--)
        *px++ = ' ';

    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
        if (isalpha(*p0) && isalpha(*p1)) {
            if (xbm[*p0-'A']&xbm[*p1-'A']) {
                cx = '*';
                nm++;
            }
            else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                cx = '!';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
    *px = '\0';
}

```

FIGURE 2M

```
/*
 * strip path or prefix from pn, return len: pr_align()
 */
static
stripname(pn)      stripname
char      *pn;     /* file name (may be path) */
{
    register char    *px, *py;

    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
            py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
}
```

FIGURE 2N

```

/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
 * g_alloc() -- calloc() with error checkin
 * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>

char    *jname = "/tmp/homgXXXXXX";          /* tmp file for jumps */
FILE    *fj;

int      cleanup();                          /* cleanup tmp file */
long     lseek();

/*
 * remove any tmp file if we blow
 */
cleanup(i) cleanup
int      i;
{
    if (fj)
        (void) unlink(jname);
    exit(i);
}

/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char     *
getseq(file, len)
char     *file;    /* file name */
int      *len;     /* seq len */
{
    char          line[1024], *pseq;
    register char *px, *py;
    int          natgc, tlen;
    FILE         *fp;

    if ((fp = fopen(file, "r")) == 0) {
        fprintf(stderr, "%s: can't read %s\n", prog, file);
        exit(1);
    }
    tlen = natgc = 0;
    while (fgets(line, 1024, fp)) {
        if (*line == ';' || *line == '<' || *line == '>')
            continue;
        for (px = line; *px != '\n'; px++)
            if (isupper(*px) || islower(*px))
                tlen++;
    }
    if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
        fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
        exit(1);
    }
    pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';

```

FIGURE 20

```

...getseq

py = pseq + 4;
*len = tlen;
rewind(fp);

while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
            natgc++;
    }
    *py++ = '\0';
    *py = '\0';
    (void) fclose(fp);
    dna = natgc > (tlen/3);
    return(pseq+4);
}

char *
g_alloc(msg, nx, sz) g_alloc
char *msg;          /* program, calling routine */
int nx, sz;         /* number and size of elements */
{
    char *px, *calloc();

    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g_alloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}

/*
 * get final jmps from dx[] or tmp file, set pp[], reset dmax: main()
 */
readjmps() readjmps
{
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;

    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = ii = 0, dmax0 = dmax, xx = len0; ; i++) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)

```

FIGURE 2P

```

...readjumps
if (j < 0 && dx[dmax].offset && fj) {
    (void) lseek(fd, dx[dmax].offset, 0);
    (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
    (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
    dx[dmax].ijmp = MAXJMP-1;
}
else
    break;
}
if (i >= JMPS) {
    fprintf(stderr, "%s: too many gaps in alignment\n", prog);
    cleanup(1);
}
if (j >= 0) {
    siz = dx[dmax].jp.n[j];
    xx = dx[dmax].jp.x[j];
    dmax += siz;
    if (siz < 0) { /* gap in second seq */
        pp[1].n[i1] = -siz;
        xx += siz;
        /* id = xx - yy + len1 - 1
        */
        pp[1].x[i1] = xx - dmax + len1 - 1;
        gapy++;
        ngapy -= siz;
/* ignore MAXGAP when doing endgaps */
        siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
        i1++;
    }
    else if (siz > 0) { /* gap in first seq */
        pp[0].n[i0] = siz;
        pp[0].x[i0] = xx;
        gapx++;
        ngapx += siz;
/* ignore MAXGAP when doing endgaps */
        siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
        i0++;
    }
}
else
    break;
}
/* reverse the order of jumps
*/
for (j = 0, i0--; j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--; j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}

```

FIGURE 2Q

```
/*
 * write a filled jmp struct offset of the prev one (if any): nw()
 */
writejumps(ix)    writejumps
                  int      ix;
{
    char          *mktemp();

    if (!fj) {
        if (mktemp(jname) < 0) {
            fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}
```

Figure 3

CCCAGAAGTTCAAGGGCCCCCGCCTCCTGCGCTCCTGCCGCCGGGACCCTCGACCTCCT  
CAGAGCAGCCGGCTGCCGCCCGGGAAAGATGGCGAGGAGGAGCCGCCACCGCCTCCTCCT  
GCTGCTGCTGCGCTACCTGGTGGTCGCCCTGGGCTATCATAAGGCCATATGGGTTTTCTGC  
CCAAAAGACCAACAAGTAGTCACAGCAGTAGAGTACCAAGAGGCTATTTTAGCCTGCAA  
AACCCCAAAGAAGACTGTTTTCTCCAGATTAGAGTGGAAGAAACTGGGTCGGAGTGTCTC  
CTTTGTCTACTATCAACAGACTCTTCAAGGTGATTTTAAAAATCGAGCTGAGATGATAGA  
TTTCAATATCCGGATCAAAAATGTGACAAGAAGTGATGCCGGGAAATATCGTTGTGAAGT  
TAGTGCCCCATCTGAGCAAGGCCAAAACCTGGAAGAGGATACAGTCACTCTGGAAGTATT  
AGTGGCTCCAGCAGTTCATCATGTGAAGTACCCTCTTCTGCTCTGAGTGGAAGTGTGGT  
AGAGCTACGATGTCAAGACAAAGAAGGGAATCCAGCTCCTGAATACACATGGTTTAAAGGA  
TGGCATCCGTTTGTAGAAAATCCAGACTTGGCTCCCAAAGCACCAACAGCTCATAAC  
AATGAATACAAAACCTGGAAGTCTGCAATTTAATACTGTTTCCAAAACCTGGACACTGGAGA  
ATATTCCTGTGAAGCCCGCAATTCTGTGGATATCGCAGGTGTCTGGGAAACGAATGCA  
AGTAGATGATCTCAACATAAGTGGCATCATAGCAGCCGTAGTAGTTGTGGCCTTAGTGAT  
TTCCGTTTGTGGCCTTGGTGTATGCTATGCTCAGAGGAAAGGCTACTTTTCAAAGAAAC  
CTCCTTCCAGAAGAGTAATTCTTCATCTAAAAGCCACGACAATGAGTGAAAATGTGCAGTG  
GCTCACGCCTGTAATCCCAGCACTTTGGAAGGCCGCGGCGGGCGGATCACGAGGTCAGGA  
GTTCTAGACCAGTCTGGCCAATATGGTAAAACCCATCTCTACTAAAATACAAAATTAG  
CTGGGCATGGTGGCATGTGCCTGCAGTTCAGCTGCTTGGGAGACAGGAGAATCACTTGA  
ACCCGGGAGGCGGAGGTTGCAGTGAGCTGAGATCACGCCACTGCAGTCCAGCCTGGGTAA  
CAGAGCAAGATTCCATCTCAAAAATAAAAATAAAATAAAATAAAATAAAATACTGGTTTTTACC  
TGTAATTCCTTACAATAAATATAGCTTGATATTC

Figure 4

MARRSRHRLLLLLLRYLVVALGYHKAYGFSAPKDQQVVTAVEYQEAILACKTPKKTVSS  
RLEWKKLGRSVSFVYYQOTLQGDFKNRAEMIDFNIRIKNVTRSDAGKYRCEVSAPSEQG  
QNLEEDTVTLEVLVAPAVPSCEVPSSALSGTVVELRCQDKEGNPAPEYTWFKDGIRLLE  
NPRLGSQSTNSSYTMNTKTGTLQFNTVSKLDTGEYSCEARNSVGYRRCPGKRMQVDDLN  
ISGIIAAVVVALVISVCGLGVCYAQRKGYFSKETSFQKSNSSSKATTMSENVQLTPV  
IPALWCAAAGGSRGQEF

N-glycosylation siteat residues:

98-102  
187-191  
236-240  
277-281

Casein kinase II phosphorylation siteat residues:

39-43  
59-63  
100-104  
149-153  
205-209  
284-288

N-myristoylation siteat residues:

182-188  
239-245  
255-261  
257-263  
305-311

Amidation site at residues:

226-230

Figure 5

CCAGGCCGGGAGGCGACGCGCCAGCCGTCTAAACGGGAACAGCCCTGGCTGAGGGAGCT  
 GCAGCGCAGCAGAGTATCTGACGGCGCCAGGTTGCGTAGGTGCGGCACGAGGAGTTTTCC  
 CGGCAGCGAGGAGGTCCTGAGCAGCATGGCCCCGGAGGAGCGCCTTCCCTGCCGCCGCGCT  
 CTGGCTCTGGAGCATCCTCCTGTGCCTGTGGCACTGCGGGCGGAGGCCGGGCCGCCGA  
 GGAGGAGAGCCTGTACCTATGGATCGATGCTCACCAGGCAAGAGTACTCATAGGATTTGA  
 AGAAGATATCCTGATTGTTTCAGAGGGGAAAATGGCACCTTTTACACATGATTTTCAGAAA  
 AGCGCAACAGAGAATGCCAGCTATTCTGTCAATATCCATTCCATGAATTTTACCTGGCA  
 AGCTGCAGGGCAGGCAGAATACTTCTATGAATTCCTGTCTTGGCTCCCTGGATAAAGG  
 CATCATGGCAGATCCAACCGTCAATGTCCCTCTGCTGGGAACAGTGCCTCACAAGGCATC  
 AGTTGTTCAAGTTGGTTTCCCATGTCTTGGAAAACAGGATGGGGTGGCAGCATTTTGAAGT  
 GGATGTGATTGTTATGAATTCGAAGGCAACACCATTCTCCAAACACCTCAAAAATGCTAT  
 CTTCTTTAAAAACATGTCAACAAGCTGAGTGCCAGGCGGGTCCGAAAATGGAGGCTTTTG  
 TAATGAAAGACGCATCTGCGAGTGTCTGATGGGTTCCACGGACCTCACTGTGAGAAAGC  
 CCTTGTACCCACGATGTATGAATGGTGGACTTTGTGTGACTCCTGGTTTTCTGCATCTG  
 CCCACCTGGATTCTATGGAGTGAACCTGTGACAAAGCAAACCTGCTCAACCACCTGCTTTAA  
 TGGAGGGACCTGTTTCTACCCTGGAAAATGTATTTGCCCTCCAGGACTAGAGGGAGAGCA  
 GTGTGAAATCAGCAAATGCCACAACCCTGTGAAAATGGAGGTAAATGCATTGGTAAAG  
 CAAATGTAAGTGTTCAAAAGTTACCAGGGAGACCTCTGTTCAAAGCCTGTCTGCGAGCC  
 TGGCTGTGGTGCACATGGAACCTGCCATGAACCCAACAAATGCCAATGTCAAGAAGGTTG  
 GCATGGAAGACACTGCAATAAAAAGGTACGAAAGCCAGCCTCATACATGCCCTGAGGCCAGC  
 AGGCGCCAGCTCAGGCAGCACACGCCTTCACTTAAAAAGGCCGAGGAGCGGGGGATCC  
 ACCTGAATCCAATTACATCTGGTGAACCTCCGACATCTGAAACGTTTTTAAGTTACACCAAG  
 TTCATAGCCTTTGTTAACCTTTCATGTGTTGAATGTTCAAATAATGTTTATTACACTTAA  
 GAATACTGGCCTGAATTTATTAGCTTCATTATAAATCACTGAGCTGATATTTACTCTTC  
 CTTTTAAGTTTTCTAAGTACGTCTGTAGCATGATGGTATAGATTTTCTGTTTTAGTGCT  
 TTGGGACAGATTTTATATTATGTCAATTGATCAGGTTAAAATTTTTCAGTGTGTAGTTGGC  
 AGATATTTTCAAATTACAATGCATTTATGGTGTCTGGGGCAGGGGAACATCAGAAAGG  
 TTAAATTGGGCAAAAATGCGTAAGTCACAAGAATTTGGATGGTGCAGTTAATGTTGAAGT  
 TACAGCATTTCAGATTTTATTGTGAGATATTTAGATGTTTGTACATTTTTTAAAAATTGC  
 TCTTAATTTTTTAACTCTCAATACAATATATTTTGACCTTACCATTATTCCAGAGATTCA  
 GTATTAAAAAATTAATACTGTGGTAGTGGCATTTAAACAATATAATATATTCTA  
 AACACAATGAAATAGGGAATATAATGTATGAACTTTTTGCATTGGCTTGAAGCAATATAA  
 TATATTGTAACAACACAGCTCTTACCTAATAAACATTTTATACTGTTTGTATGTATA  
 AAATAAAGTGCTGCTTTAGTTTTTTTGGAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Figure 6

MARRSAFPAAALWLWSILLCLLALRAEAGPPQEESLYLWIDAHQARVLIGFEEDILIVS  
EGKMAPFTHDFRKAQQRMPAIPVNIHSMNFTWQAAGQAEYFYEFSLRSLDKGIMADPT  
VNVPLLGTVPHKASVVQVGFPCLGKQDGVAAFEVDVIVMNSEGNTILQTPQNAIFFKTC  
QQAECPPGGCRNGGFCNERRICECPDGFHGHCEKALCTPRCMNGGLCVTPGFICPPGF  
YGVNCDKANCSTTCFNGGTCFYPGKICPPGLEGEQCEISKCPQPCRNGGKCIGKSKCK  
CSKGYQGDLCSPVCEPGCGAHGTCHEPNKCQCQEGWHGRHCNKRYEASLIHALRPAGA  
QLRQHTPSLKKAERRDPPESNYIW

N-glycosylation site at residues:

88-92

245-249

Casein kinase II phosphorylation site at residues:

319-323

Tyrosine kinase phosphorylation site at residues:

370-378

N-myristoylation site at residues:

184-190

185-191

189-195

315-321

ATP/GTP-binding site motif A (P-loop) at residues:

285-293

EGF-like domain cysteine pattern signature at residues:

198-210

230-242

262-274

294-306

326-338

Figure 7

GTCTGTTCCCAGGAGTCCTTCGGCGGCTGTTGTGTGTCAGTGGCCTGATCGCGATGGGGACA  
AAGGCGCAAGTCGAGAGGAACTGTTGTGCCTCTTCATATTGGCGATCCTGTTGTGCTCC  
CTGGCATTGGGCAGTGTTACAGTGCACCTCTTCTGAACCTGAAGTCAGAATTCCTGAGAAT  
AATCCTGTGAAGTTGTCTGTGCCTACTCGGGCTTTTCTTCTCCCCGTGTGGAGTGGAAAG  
TTTGACCAAGGAGACACCACCAGACTCGTTTGCTATAATAACAAGATCACAGCTTCCTAT  
GAGGACCGGGTGACCTTCTTGCCAACCTGGTATCACCTTCAAGTCCGTGACACGGGAAGAC  
ACTGGGACATACTTGTATGGTCTCTGAGGAAGGCGGCAACAGCTATGGGGAGGTCAAG  
GTCAAGCTCATCGTGCTTGTGCCTCCATCCAAGCCTACAGTTAACATCCCCCTCCTTGCC  
ACCATTTGGGAACCGGGCAGTGCTGACATGCTCAGAACAAGATGGTTCCCCACCTTCTGAA  
TACACCTGGTTCAAAGATGGGATAGTGATGCCTACGAATCCCAAAGCACCCGTGCCTTC  
AGCAACTCTTCTATGTCCTGAATCCCACAACAGGAGAGCTGGTCTTTGATCCCCGTCA  
GCCTCTGATACTGGAGAATACAGCTGTGAGGCACGGAATGGGTATGGGACACCCATGACT  
TCAAATGCTGTGCGCATGGAAGCTGTGGAGCGGAATGTGGGGTTCATCGTGGCAGCCGTC  
CTTGTAACCCTGATTCTCCTGGGAATCTTGGTTTTTGGCATCTGGTTTGCCTATAGCCGA  
GGCCACTTTGACAGAACAAAGAAAGGGACTTCGAGTAAGAAGGTGATTTACAGCCAGCCT  
AGTGCCCGAAGTGAAGGAGAATTCAAACAGACCTCGTCATTCTGGTGTGAGCCTGGTCG  
GCTCACCGCCTATCATCTGCATTTGCCTTACTCAGGTGCTACCGGACTCTGGCCCCGTAT  
GTCTGTAGTTTACAGGATGCCTTATTTGTCTTCTACACCCACAGGGCCCCCTACTTCT  
TCGGATGTGTTTTTAATAATGTCAGCTATGTGCCCATCCTCCTTCATGCCCTCCCTCCC  
TTTCTACCCTGCTGAGTGGCCTGGAACCTGTTTTAAAGTGTATTATCCCCATTTCTTTG  
AGGGATCAGGAAGGAATCCTGGGTATGCCATTGACTTCCCTTCTAAGTAGACAGCAAAAA  
TGGCGGGGGTTCGAGGAATCTGCACTCACTGCCACCTGGCTGGCAGGGATCTTTGAAT  
AGGTATCTTGAGCTTGGTTCTGGGCTCTTCTTGTGTACTGACGACCAGGGCCAGCTGT  
TCTAGAGCGGGAATTAGAGGCTAGAGCGGCTGAAATGGTTGTTTGGTGTATGACTGGGG  
TCCTTCCATCTCTGGGGCCACTCTCTTCTGTCTTCCCATGGGAAGTGCCACTGGGATCC  
CTCTGCCCTGTCCTCCTGAATACAAGCTGACTGACATTGACTGTGTCTGTGGAAAATGGG  
AGCTCTTGTTGTGGAGAGCATAGTAAATTTTTCAGAGAACTTGAAGCCAAAAGGATTTAAA  
ACCGCTGCTCTAAAGAAAAGAAAACCTGGAGGCTGGGCGCAGTGGCTCACGCCTGTAATCC  
CAGAGGCTGAGGCAGGCGGATCACCTGAGGTCTGGGAGTTCTGGGATCAGCCTGACCAACAT  
GGAGAAACCCTACTGGAATACAAAGTTAGCCAGGCATGGTGGTGCATGCCTGTAGTCCC  
AGCTGCTCAGGAGCCTGGCAACAAGAGCAAACTCCAGCTCAAAAAAAAAAAAAAAAAA

Figure 8

MGTKAQVERKLLCLFILAILLCSLALGSVTVHSSEPEVRIPENNPVKLSCAYSGFSSPR  
VEWKFDQGDTRRLVCYNNKITASYEDRVTFLLPTGITFKSVTREDTGTYTTCMVSEEGNS  
YGEVKVKLIVLVPPSKPTVNIPSSATIGNRAVLTCSEQDGSPPSEYTWFKDGIVMPTNP  
KSTRAFSNSSYVLNPTTGELVFDPLSASDTGEYSCEARNGYGTTPMTSNAVRMEAVERNV  
GVIVA AVLVTLLILGILVFGIWFAYS RGHFDR TKKGTSSKKVIYSQPSARSEGEFKQTS  
SFLV

N-glycosylation site at residues:

185-189

cAMP- and cGMP-dependent protein kinase phosphorylation site at  
residues:

270-274

Casein kinase II phosphorylation site at residues:

34-38

82-86

100-108

118-122

152-156

154-158

193-197

203-207

287-291

N-myristoylation site at residues:

105-111

116-122

158-164

219-225

237-243

256-262

Figure 9

GGGGTTAGGGAGGAAGGAATCCACCCCCACCCCCCAAACCCCTTTTCTTCTCCTTTCT  
GGCTTCGGACATTGGAGCACTAAATGAACTTGAATTGTGTCTGTGGCGAGCAGGATGGTC  
GCTGTTACTTTGTGATGAGATCGGGGATGAATTGCTCGCTTTAAAAATGCTGCTTTGGAT  
TCTGTTGCTGGAGACGTCTCTTTGTTTTGCGCTGGAAACGTTACAGGGGACGTTTGCAA  
AGAGAAGATCTGTTCCCTGCAATGAGATAGAAGGGGACCTACACGTAGACTGTGAAAAAAA  
GGGCTTCACAAGTCTGCAGCGTTTCACTGCCCCGACTTCCCAGTTTTACCATTTATTTCT  
GCATGGCAATCCCTCACTCGACTTTTCCCTAATGAGTTTCGTAACCTTTTATAATGCGGT  
TAGTTTGACATGGAAAAAATGGCTTGCATGAAATCGTTCCGGGGGCTTTTCTGGGGCT  
GCAGCTGGTGAAGGCTGCACATCAACAACAAGATCAAGTCTTTTCGAAAGCAGAC  
TTTTCTGGGGCTGGACGATCTGGAATATCTCCAGGCTGATTTTAAATTTATTACGAGATAT  
AGACCCGGGGCCCTCCAGGACTTGAACAAGCTGGAGGTGCTCATTTTAAATGACAATCT  
CATCAGCACCTACCTGCCAACGTGTTCCAGTATGTGCCCATCACCCACCTCGACCTCCG  
GGTAAACAGGCTGAAAACGCTGCCCTATGAGGAGGTCTTGAGCAAATCCCTGGTATTGC  
GGAGATCCTGCTAGAGGATAACCCTTGGGACTGCACCTGTGATCTGCTCTCCCTGAAAGA  
ATGGCTGGAAAAATCCCAAGAATGCCCTGATCGGCCGAGTGGTCTGCGAAGCCCCCAC  
CAGACTGCAGGGTAAAGACCTCAATGAAACCACCGAACAGGACTTGTGTCCTTTGAAAAA  
CCGAGTGGATTCTAGTCTCCCGCGCCCCCTGCCCAAGAAGAGACCTTTGCTCCTGGACC  
CCTGCCAACTCCTTTCAAGACAAATGGGCAAGAGGATCATGCCACACCAGGGTCTGCTCC  
AAACGGAGGTACAAAGATCCCAGGCAACTGGCAGATCAAAATCAGACCCACAGCAGCGAT  
AGCGACGGGTAGCTCCAGGAACAAACCTTAGCTAACAGTTTACCCTGCCCTGGGGGCTG  
CAGCTGCGACCACATCCCAGGGTCGGGTTTAAAGATGAACTGCAACAACAGGAACGTGAG  
CAGCTTGGCTGATTTGAAGCCCAAGCTCTCTAACGTGCAGGAGCTTTTCTACGAGATAA  
CAAGATCCACAGCATCCGAAAATCGCACTTTGTGGATTACAAGAACCTCATTCTGTTGGA  
TCTGGGCAACAATAACATCGCTACTGTAGAGAACAACACTTTCAAGAACCTTTTGGACCT  
CAGGTGGCTATACATGGATAGCAATTACCTGGACACGCTGTCCCGGGAGAAATTCGCGGG  
GCTGCAAAACCTAGAGTACCTGAACGTGGAGTACAACGCTATCCAGCTCATCCTCCCGGG  
CACTTTCAATGCCATGCCCAAACCTGAGGATCCTCATTCTCAACAACAACCTGCTGAGGTC  
CCTGCCTGTGGACGTGTTTCGCTGGGGTCTCGCTCTCTAAACTCAGCCTGCACAACAATTA  
CTTCATGTACCTCCCGGTGGCAGGGGTGCTGGACCAGTTAACCTCCATCATCCAGATAGA  
CCTCCACGGAAACCCCTGGGAGTGCTCCTGCACAATTGTGCCTTTCAAGCAGTGGGCAGA  
ACGCTTGGGTTCCGAAGTGCTGATGAGCGACCTCAAGTGTGAGACGCCGGTGAACCTTCTT  
TAGAAAGGATTTTCATGCTCCTCTCCAATGACGAGATCTGCCCTCAGCTGTACGCTAGGAT  
CTCGCCCACGTTAACTTCGCACAGTAAAAACAGCACTGGGTTGGCGGAGACCGGGACGCA  
CTCCAACCTCCTACCTAGACACCAGCAGGGTGTCCATCTCGGTGTTGGTCCCGGGACTGCT  
GCTGGTGTGTTGTCACCTCCGCCTCACCGTGGTGGGCATGCTCGTGTATTATCCTGAGGAA  
CCGAAAGCGGTCCAAGAGACGAGATGCCAACTCCTCCGCGTCCGAGATTAATTCCTTACA

GACAGTCTGTGACTCTTCCTACTGGCACAATGGGCCTTACAACGCAGATGGGGCCACAG  
AGTGTATGACTGTGGCTCTCACTCGCTCTCAGACTTAAGACCCCAACCCCAATAGGGGAGG  
GCAGAGGGAAGGCGATACATCCTTCCCCACCGCAGGCACCCCGGGGGCTGGAGGGGCGTG  
TACCCAAATCCCCGCGCCATCAGCCTGGATGGGCATAAGTAGATAAAATAACTGTGAGCTC  
GCACAACCGAAAGGGCCTGACCCCTTACTTAGCTCCCTCCTTGAAACAAAGAGCAGACTG  
TGGAGAGCTGGGAGAGCGCAGCCAGCTCGCTCTTTGCTGAGAGCCCCTTTTGACAGAAAG  
CCCAGCACGACCCTGCTGGAAGAAGTACAGTGCCCTCGCCCTCGGCCCGGGCCCTGTG  
GGGTTGGATGCCGCGGTTCTATACATATATACATATATCCACATCTATATAGAGAGATAG  
ATATCTATTTTTCCCTGTGGATTAGCCCCGTGATGGCTCCCTGTTGGCTACGCAGGGAT  
GGGCAGTTGCACGAAGGCATGAATGTATTGTAAATAAGTAACTTTGACTTCTGAC

Figure 10

MLLWILLETSLCFAAGNVTGDVCKEKICSCNEIEGDLHVDCEKKGFTSLQRFTAPTSQ  
 FYHLFLHGNSLTRLFPNEFANFYNAVSLHMENGLHEIVPGAFLGLQLVKRLHINNNKI  
 KFRKQTFGLDDLEYLQADFNLLRDIDPGAQDLNKLEVLILNDNLISTLPANVFQYV  
 PITHLDRGNRLKTLPYEEVLEQIPGIAEILLEDNPWDCTCDLLSLKEWLENIPKNALI  
 GRVVCEAPTRLQGKDLNETTEQDLCPLKNRVDSLPAPPAQEETFAPGPLPTPFKTNGQ  
 EDHATPGSAPNGGTKIPGNWQIKIRPTAAIATGSSRNKPLANSRPCGGCSCDHI PGSG  
 LKMNCNRRNVSSLADLKPKLSNVQELFLRDNKIHSIRKSHFVDYKNLILLDLGNNNIAT  
 VENNTFKNLLDLRWLYMDSNYLDTLSREKFAQLNLEYLNVEYNAIQILPGTFNAMPK  
 LRILILNNLLRSLPVDVVFAGVSLSKLSLHNNYFMYLPVAGVLDQLTSIIQIDLHGNPW  
 ECSCTIVPFKQWAERLGSEVLMSDLKCEPVMNFRKDFMLLSNDEICPOLYARISPTLT  
 SHSKNSTGLAETGTHSNSYLDTSRVSISVLVPGLLLVFVTSFTVVGMLVFILRNKRKRS  
 KRRDANSSASEINSLQTVCDSSYWHNGPYNADGAHRVYDCGSHSLSD

N-glycosylation site at residues:

18-22  
 253-257  
 363-367  
 416-420  
 595-599  
 655-659

cAMP- and cGMP-dependent protein kinase phosphorylation site at residues:

122-126  
 646-650

Casein kinase II phosphorylation site at residues:

30-34  
 180-184  
 222-226  
 256-260  
 366-370  
 573-577  
 608-612  
 657-661  
 666-670  
 693-697

N-myristoylation site at residues:

17-23  
67-73  
100-106  
302-308  
328-334  
343-349  
354-360  
465-471  
493-499  
598-604  
603-609

Prokaryotic membrane lipoprotein lipid attachment site at residues:

337-348

Figure 11

GTAAGTGAAGTCAGGCTTTTCATTTGGGAAGCCCCCTCAACAGAATTCGGTCATTCTCCA  
AGTTAIGGTGGACGTACTTCTGTTGTTCTCCCTCTGCTTGCTTTTTTCACATTAGCAGACC  
GGACTTAAGTCACAACAGATTATCTTTTCATCAAGGCAAGTTCCATGAGCCACCTTCAAAG  
CCTTCGAGAAGTGAACTGAACAACAATGAATTGGAGACCATTCCAAATCTGGGACCAGT  
CTCGGCAAATATTACACTTCTCTCCTTGGCTGGAAACAGGATTGTTGAAATACTCCCTGA  
ACATCTGAAAGAGTTTCAGTCCCTTGAAACTTTGGACCTTAGCAGCAACAATATTTTCAGA  
GCTCCAAACTGCATTTCCAGCCCTACAGCTCAAATATCTGTATCTCAACAGCAACCGAGT  
CACATCAATGGAACCTGGGTATTTTGACAATTTGGCCAACACACTCCTTGTGTTAAAGCT  
GAACAGGAACCGAATCTCAGCTATCCACCCAAAGATGTTTAAACTGCCCAACTGCAACA  
TCTCGAATTGAACCGAAACAAGATTAATAATGTAGATGGACTGACATTCCAAGGCCTTGG  
TGCTCTGAAGTCTCTGAAAATGCAAAGAAATGGAGTAACGAAACTTATGGATGGAGCTTT  
TTGGGGGCTGAGCAACATGGAAATTTTGCAGCTGGACCATAACAACCTAACAGAGATTAC  
CAAAGGCTGGCTTTACGGCTTGCTGATGCTGCAGGAACTTCATCTCAGCCAAAATGCCAT  
CAACAGGATCAGCCCTGATGCCTGGGAGTTCTGCCAGAAGCTCAGTGAGCTGGACCTAAC  
TTTCAATCACTTATCAAGGTTAGATGATTCAAGCTTCCTTGGCCTAAGCTTACTAAATAC  
ACTGCACATTGGGAACAACAGAGTCAGCTACATTGCTGATTGTGCCTTCCGGGGGCTTTC  
CAGTTTAAAGACTTTGGATCTGAAGAACAATGAAATTTCTGGACTATTGAAGACATGAA  
TGGTGTCTTCTCTGGGCTTGACAAAATGAGGGCAGTACTCAAGGAAATCGGATCCG  
TTCTATTACTAAAAAGCCTTCACTGGTTTTGGATGCATTGGAGCATCTAGACCTGAGTGA  
CAACGCAATCATGTCTTTACAAGGCAATGCATTTTCACAAATGAAGAACTGCAACAATT  
GCATTTAAATACATCAAGCCTTTTGTGCGATTGCCAGCTAAAATGGCTCCACAGTGGGT  
GGCGGAAAACAACCTTTCAGAGCTTTGTAAAATGCCAGTTGTGCCCATCTCAGCTGCTAAA  
AGGAAGAAGCATTTTTGCTGTTAGCCAGATGGCTTTGTGTGTGATGATTTTTCCCAAACC  
CCAGATCACGGTTCAGCCAGAAACACAGTCGGCAATAAAAGGTTCCAATTTGAGTTTCAT  
CTGCTCAGCTGCCAGCAGCAGTGATTCCCCAATGACTTTTGTCTTGAAAAAAGACAATGA  
ACTACTGCATGATGCTGAAATGGAAAATTATGCACACCTCCGGGCCCCAAGGTGGCGAGGT  
GATGGAGTATACCACCATCCTTCGGCTGCGCGAGGTGGAATTTGCCAGTGAGGGGAAATA  
TCAGTGTGTCATCTCCAATCACTTTGGTTCATCCTACTCTGTCAAAGCCAAGCTTACAGT  
AAATATGCTTCCCTCATTCACCAAGACCCCCATGGATCTCACCATCCGAGCTGGGGCCAT  
GGCACGCTTGGAGTGTGCTGCTGTGGGGCACCCAGCCCCCAGATAGCCTGGCAGAAGGA  
TGGGGGCACAGACTTCCCAGCTGCACGGGAGAGACGCATGCATGTGATGCCCGAGGATGA  
CGTGTCTTTATCGTGGATGTGAAGATAGAGGACATTGGGGTATACAGCTGCACAGCTCA  
GAACAGTGCAGGAAGTATTTTCAGCAAATGCAACTCTGACTGTCTTAGAAACACCATCATT  
TTTGCGGCCACTGTTGGACCGAAGTGAACCAAGGGAGAAACAGCCGTCTACAGTGCAT  
TGCTGGAGGAAGCCCTCCCCCTAAACTGAACTGGACCAAAGATGATAGCCCATTTGGTGGT  
AACCGAGAGGCACTTTTTTGTCAGCAGGCAATCAGCTTCTGATTATTGTGGACTCAGATGT

CAGTGATGCTGGGAAATACACATGTGAGATGTCTAACACCCTTGGCACTGAGAGAGGAAA  
CGTGCGCCTCAGTGTGATCCCCACTCCAACCTGCGACTCCCCTCAGATGACAGCCCCATC  
GTTAGACGATGACGGATGGGCCACTGTGGGTGTCGTGATCATAGCCGTGGTTTGCTGTGT  
GGTGGGCACGTCACCTCGTGTGGGTGGTCATCATATACCACACAAGGCGGAGGAATGAAGA  
TTGCAGCATTACCAACACAGATGAGACCAACTTGCCAGCAGATATTCTAGTTATTTGTC  
ATCTCAGGGAACGTTAGCTGACAGGCAGGATGGGTACGTGTCTTCAGAAAGTGAAGCCA  
CCACCAGTTTGTACATCTTCAGGTGCTGGATTTTTCTTACCACAACATGACAGTAGTGG  
GACCTGCCATATTGACAATAGCAGTGAAGCTGATGTGGAAGCTGCCACAGATCTGTTCCCT  
TTGTCCGTTTTTGGGATCCACAGGCCCTATGTATTTGAAGGGAAATGTGTATGGCTCAGA  
TCCTTTTGAAACATATCATAAGGTTGCAGTCCTGACCCAAGAACAGTTTTAATGGACCA  
CTATGAGCCCAGTTACATAAAGAAAAAGGAGTGCTACCCATGTTCTCATCCTTCAGAAGA  
ATCCTGCGAACGGAGCTTCAGTAATATATCGTGGCCTTCACATGTGAGGAAGCTACTTAA  
CACTAGTTACTCTCACAATGAAGGACCTGGAATGAAAAATCTGTGTCTAAACAAGTCCTC  
TTTAGATTTTAGTGCAAATCCAGAGCCAGCGTCCGTTGCCTCGAGTAATCTTTTCATGGG  
TACCTTTGGAAAAGCTCTCAGGAGACCTCACCTAGATGCCTATTCAAGCTTTGGACAGCC  
ATCAGATTGTCAGCCAAGAGCCTTTTATTTGAAAGCTCATTCTTCCCCAGACTTGGACTC  
TGGGTGAGAGGAAGATGGGAAAGAAAGGACAGATTTTCAGGAAGAAAATCACATTTGTAC  
CTTTAAACAGACTTTAGAAAACCTACAGGACTCCAAATTTTCAGTCTTATGACTTGGACAC  
ATAGACTGAATGAGACCAAAGGAAAAGCTTAACATACTACCTCAAGTGAACCTTTTATTTA  
AAAGAGAGAGAATCTTATGTTTTTTAAATGGAGTTATGAATTTTAAAAGGATAAAAATGC  
TTTATTTATACAGATGAACCAAATTACAAAAGTTATGAAAATTTTATACTGGGAATG  
ATGCTCATATAAGAATACCTTTTTAAACTATTTTTTAACTTTGTTTTATGCAAAAAGTA  
TCTTACGTAAATTAATGATATAAATCATGATTATTTTATGTATTTTTATAATGCCAGATT  
TCTTTTTATGAAAATGAGTTACTAAAGCATTTTAAATAATACTGCCTTGTACCATTTT  
TTAAATAGAAGTTACTTCATTATATTTGCACATTATATTTAATAAAATGTGTCAATTTG  
AA

Figure 12

MVDVLLLFSLCLLFHISRPDLSHNRLSFIKASSMSHLQSLREVKLNNELETIPNLGPV  
 SANITLLSLAGNRIVEILPEHLKEFQSLETLDLSSNNISELQTAFPALQLKYLYLNSNR  
 VTSMEPGYFDNLANLTLVLKLNRRNRISAIPPKMFKLPQLQHLELNRNKIKNVDGLTFQG  
 LGALKSLKMQRNGVTKLMDGAFWGLSNMEILQLDHNNLTEITKGWLYGLLMLQELHLSQ  
 NAINRISPDAWEFCQKLELDLTFNHL SRLDDSSFLGSLNLTNLHIGNNRVSYIADCAF  
 RGLSSLKTLDLKNEISWTIEDMNGAFSGLDKLRRILIQGNRIRSITKKAFTGLDALEH  
 LDLSDNAIMSLQGNAFSQMKKLQQLHLNTSSLLCDCQLKWLPOQVAENNFQSFVNASCA  
 HPQLLKGRSIFAVSPDGFVCDDFPKPQITVQPETQSAIKGSNLSFICSAASSSDSPMTF  
 AWKKNELLHDAEMENYAHLRAQGGVMEYTTILRLREVEFASEGKYQCVISNHFGSSY  
 SVKAKLTVNMLPSFTKTPMDLTI RAGAMARLECAAVGHPAPQIAWQKDGTDFFAARER  
 RMHVMPEDDVFFIVDVKIEDIGVYSCTAQNSAGSISANATLTVLETSPFLRPLLDRTVT  
 KGETAVLQCIAGGSPPKLNWTKDDSPLVVTERHFFAAGNQLLIIVDSDVSDAGKYTCE  
 MSNTLGTTERGNVRLSVIPTPTCDS PQMTAPSLDDDGWATVGVV IIAVVCCVVGTSLVWV  
 V I IYHTRRRNEDCSITNTDETNL PADIPSYLSSQGT LADRQDGYVSSSESGSHHQFVTSS  
 GAGFFLPQHDSSTCHIDNSSEADVEAATDLFLCPFLGSTGPMYLGKNVYGSDFPETYH  
 TGCSPPDRTVLMDHYEPSYIKKKECYPCSHPSEESCERSFSNISWPSHVRKLLNTSYSH  
 NEGPGMKNLCLNKSSLD FSANPEPASVASSNSFMGTFGKALRRPHLDAYSSFGQPSDCQ  
 PRAFYLKAHSSPDLD SGSEEDGKERTDFQEENHICTFKQTL ENYRTPNFQSYDLDT

N-glycosylation site at residues:

- 62-66
- 96-100
- 214-218
- 382-386
- 409-413
- 455-459
- 628-632
- 669-673
- 845-849
- 927-931
- 939-943
- 956-960

Glycosaminoglycan attachment site at residues:

- 826-830

## Casein kinase II phosphorylation site at residues:

17-21  
39-43  
120-124  
203-207  
254-258  
264-268  
314-318  
323-327  
347-351  
464-468  
548-552  
632-636  
649-653  
671-675  
739-743  
783-787  
803-807  
847-851  
943-947  
958-962  
1013-1017  
1019-1023  
1021-1025

## Tyrosine kinase phosphorylation site at residues:

607-615

## N-myristoylation site at residues:

179-185  
197-203  
320-326  
367-373  
453-459  
528-534  
612-618  
623-629  
714-720  
873-879

Figure 13

GGGGAGAGGAATTGACCATGTAAAAGGAGACTTTTTTTTTTTGGTGGTGGTGGCTGTGGG  
 TGCCTTGCAAAAATGAAGGATGCAGGACGCAGCTTCTCCTGGAACCGAACGCAATGGAT  
 AAAGTATTGTGCAAGAGAGAAGGAAGAACGAAGCTTTTCTTGTGAGCCCTGGATCTTA  
 ACACAAATGTGTATATGTGCACACAGGGAGCATTCAAGAATGAAATAAACAGAGTTAGA  
 CCCGCGGGGGTTGGTGTGTTCTGACATAAATAAATAATCTTAAAGCAGCTGTTCCCTCC  
 CCACCCCAAAAAAAGGATGATTGGAAATGAAGAACCGAGGATTCACAAAGAAAAAGT  
 ATGTTCATTTTTCTCTATAAAGGAGAAAGTGAGCCAAGGAGATATTTTTGGAATGAAAAG  
 TTTGGGGCTTTTTTAGTAAAGTAAAGAAGCTGGTGTGGTGGTGTTTTCCTTTCTTTTTGAA  
 TTTCCACAAGAGGAGAGGAAATTAATAATACATCTGCAAAGAAATTTAGAGAAGAAAA  
 GTTGACCGCGCAGATTGAGGCATTGATTGGGGGAGAGAAACCAGCAGAGCACAGTTGGA  
 TTTGTGCCTATGTTGACTAAAATGACGGATAATGCAGTTGGATTTTTCTTCATCAACC  
 TCCTTTTTTTTAAATTTTTATTCTTTTGGTATCAAGATCATGCGTTTTCTCTTGTCTT  
 AACACCTGGATTTCCATCTGGATGTTGCTGTGATCAGTCTGAAATACAACGTTTGAAT  
 TCCAGAAGGACCAACACCAGATAAATTATGAATGTTGAACAAGATGACCTTACATCCACA  
 GCAGATAATGATAGGTCCTAGGTTTAAACAGGGCCCTATTTGACCCCTGCTTGTGGTGGT  
 GCTGGCTCTTCAACTTCTTGTGGTGGCTGGTCTGGTGCAGGGCTCAGACCTGCCCTCTGT  
 GTGCTCCTGCAGCAACCAGTTTCAAGCAAGGTGATTTGTGTTTCGAAAAACCTGCGTGAGGT  
 TCCGGATGGCATCTCCACCAACACACGGCTGCTGAACCTCCATGAGAACCAAATCCAGAT  
 CATCAAAGTGAACAGCTTCAAGCACTTGAGGCACTTGGAATCCTACAGTTGAGTAGGAA  
 CCATATCAGAACCATTGAAATGGGGCTTTCAATGGTCTGGCGAACCTCAACACTCTGGA  
 ACTCTTTGACAATCGTCTTACTACCATCCCGAATGGAGCTTTTGTATACCTTGTCTAAACT  
 GAAGGAGCTCTGGTTGCGAAACAACCCATTGAAAGCATCCCTTCTTATGCTTTTAAACAG  
 AATTCTTCTTTGCGCCGACTAGACTTAGGGGAATTGAAAAGACTTTCATACATCTCAGA  
 AGGTGCCTTTGAAGGTCTGTCCAACCTGAGGTATTTGAACCTTGCCATGTGCAACCTTCG  
 GGAAATCCCTAACCTCACACCGCTCATAAACTAGATGAGCTGGATCTTCTGGGAATCA  
 TTTATCTGCCATCAGGCCTGGCTCTTCCAGGGTTTGATGCACCTTCAAAAACCTGTGGAT  
 GATACAGTCCCAGATTCAAGTGATTGAACGGAATGCCCTTGACAACCTTCAGTCACTAGT  
 GGAGATCAACCTGGCACACAATAATCTAACATTACTGCCTCATGACCTCTTCACTCCCTT  
 GCATCATCTAGAGCGGATACATTTACATCACAACCCTTGGAACCTGTAACCTGTGACATACT  
 GTGGCTCAGCTGGTGGATAAAAGACATGGCCCCCTCGAACACAGCTTGTGTGCCCCGGTG  
 TAACACTCCTCCAATCTAAAGGGGAGGTACATTGGAGAGCTCGACCAGAATTACTTCCAC  
 ATGCTATGCTCCGGTGATTGTGGAGCCCCCTGCAGACCTCAATGTCAGTGAAGGCATGGC  
 AGCTGAGCTGAAATGTCGGGCCCTCCACATCCCTGACATCTGTATCTTGGATTACTCCAAA  
 TGGAACAGTCATGACACATGGGGCGTACAAAGTGGGATAGCTGTGCTCAGTGATGGTAC  
 GTTAAATTTACAAAATGTAACCTGTGCAAGATACAGGCATGTACACATGTATGGTGAAGTAA  
 TTCCGTTGGGAATACTACTGCTTCCAGCCACCCTGAATGTTACTGCAGCAACCACTACTCC

TTTCTCTTACTTTTCAACCGTCACAGTAGAGACTATGGAACCGTCTCAGGATGAGGCACG  
GACCACAGATAACAATGTGGGTCCCCTCCAGTGGTCGACTGGGAGACCACCAATGTGAC  
CACCTCTCTCACACCACAGAGCACAAGGTCGACAGAGAAAAACCTTCACCATCCCAGTGAC  
TGATATAAACAGTGGGATCCCAGGAATTGATGAGGTCATGAAGACTACCAAAATCATCAT  
TGGGTGTTTTGTGGCCATCACACTCATGGCTGCAGTGATGCTGGTCATTTTCTACAAGAT  
GAGGAAGCAGCACCATCGGCAAAACCATCACGCCCAACAAGGACTGTTGAAATTATTAA  
TGTGGATGATGAGATTACGGGAGACACCCCATGGAAAAGCCACCTGCCCATGCCTGCTAT  
CGAGCATGAGCACCTAAATCACTATAACTCATACAAATCTCCCTTCAACCACACAACAAC  
AGTTAACACAATAAATTC AATACACAGTTCAGTGCATGAACCGTTATTGATCCGAATGAA  
CTCTAAAGACAATGTACAAGAGACTCAAATCTAA AACATTTACAGAGTTACAAAAACAA  
ACAATCAAAAAAAGACAGTTTATTAAAAATGACACAAATGACTGGGCTAAATCTACTG  
TTTCAAAAAAGTGTCTTTACAAAAAACA AAAAGAAAATTTATTTATTA AAAATT  
CTATTGTGATCTAAAGCAGACAAAAA

Figure 14

MLNKMTLHPQQIMIGPRFNRALFDPELLVLLALQLLVAVGLVRAQTCPSVCSNSQFSK  
 VICVRKNLREVPDGI STNTRLLNLHENQIQI IKVNSFKHLRHLEILQLSRNHIRTIEIG  
 AFNGLANLNTLELFDNRLTTIPNGAFVYLSKLELWLRNPNIESIPSYAFNRIPSLRRL  
 DLGELKRLSYISEGAFEGLSNLRYLNLAMCNLREIPNLTPLIKLELDLDSGNHLSAIRP  
 GSFQGLMHLQKLWMIQSQIQVIERNAFDNLQSLVEINLAHNNLTLPHDLFTPLHHLER  
 IHLHNPWNCNCDILWLSWWIKDMAPSNTACCARCNTPPNLKGRYIGELDQNYFTCYAP  
 VIVEPPADLNVTEGMAAELKCRASSTLSVSWITPNGTVMTHGAYKVR IAVLSDGTLNF  
 TNVTVQDTGMYTCMVSNSVGNNTASATLNVTAATTT PFSYFSTVTVETMEPSQDEARTT  
 DNNVGPTPVVDWETTNTVTTSLTPQSTRSTEKTF TTPVTDINSGIPGIDEVMKTTKIIIG  
 CFVAITLMAAVMLVIFYKMRKQHHRQNH HAPTRTVEI INVDEITGDTPMESHLPMPAI  
 EHEHLNHYSYKSPFNHTTTVNTINSIHSSVHEPLLIRMNSKDNVQETQI

N-glycosylation site at residues:

- 278-282
- 364-368
- 390-394
- 412-416
- 415-419
- 434-438
- 442-446
- 488-492
- 606-610

cAMP- and cGMP-dependent protein kinase phosphorylation site at residues:

- 183-187

Casein kinase II phosphorylation site at residues:

- 268-272
- 417-421
- 465-469
- 579-583
- 620-624

N-myristoylation site at residues:

- 40-46

73-79  
118-124  
191-197  
228-234  
237-243  
391-397  
422-428  
433-439  
531-537

Figure 15

AGCCGACGCTGCTCAAGCTGCAACTCTGTTGCAGTTGGCAGTTCTTTTCGGTTTCCCTCC  
TGCTGTTTGGGGGCATGAAAGGGCTTCGCCGCCGGGAGTAAAAGAAGGAATTGACCGGGC  
AGCGCGAGGGAGGAGCGCGCACGCGACC CGGAGGGCGGGCGTGCACCCTCGGCTGGAAGT  
TTGTGCCGGGCCCCGAGCGCGCGCCGGCTGGGAGCTTCGGGTAGAGACCTAGGCCGCTGG  
ACCGCGATGAGCGCGCCGAGCCTCCGTGCGCGCGCCGCGGGGTGGGGCTGCTGCTGTGC  
GCGGTGCTGGGGCGCGCTGGCCGGTCCGACAGCGCGGTGCGGGGAACTCGGGCAGCCC  
TCTGGGTAGCCCGGAGCGCCATGCCCCACTACCTGCCGCTGCCTCGGGGACCTGCTG  
GACTGCAGTCGTAAGCGGCTAGCGCGTCTTCCCGAGCCACTCCCGTCTGGGTGCTCGG  
CTGGACTTAAGTCAACAACAGATTATCTTTCATCAAGGCAAGTTCCATGAGCCACCTTCAA  
AGCCTTCGAGAAGTGAAACTGAACAACAATGAATTGGAGACCATTCCAAATCTGGGACCA  
GTCTCGGCAAATATTACACTTCTCTCCTTGGCTGGAACAGGATTGTTGAAATACTCCCT  
GAACATCTGAAAAGAGTTTCAGTCCCTTGAAACTTTGGACCTTAGCAGCAACAATATTTCA  
GAGTCCAAAAGTGCATTTCCAGCCCTACAGCTCAAATATCTGTATCTCAACAGCAACCGA  
GTCACATCAATGGAACCTGGGTATTTTGACAATTTGGCCAACACACTCCTTGTGTTAAAG  
CTGAACAGGAACCGAATCTCAGCTATCCCACCCAAGATGTTTAAACTGCCCAACTGCAA  
CATCTCGAATTGAACCGAAAACAAGATTAAAAATGTAGATGGACTGACATTCCAAGGCCTT  
GGTGCTCTGAAGTCTCTGAAAATGCAAAGAAATGGAGTAACGAACTTATGGATGGAGCT  
TTTTGGGGCTGAGCAACATGGAATTTTGAGCTGGACCATAACAACCTAACAGAGATT  
ACCAAAGGCTGGCTTTACGGCTTGCTGATGCTGCAGGAACTTCATCTCAGCCAAAATGCC  
ATCAACAGGATCAGCCCTGATGCCTGGGAGTTCTGCCAGAAGCTCAGTGAGCTGGACCTA  
ACTTTCAATCACTTATCAAGGTTAGATGATTCAAGCTTCCTTGGCCTAAGCTTACTAAT  
ACACTGCACATTGGGAACAACAGAGTCAGCTACATTGCTGATTGTGCCTTCCGGGGCTT  
TCCAGTTTAAAGACTTTGGATCTGAAGAACAATGAAATTTCTGGACTATTGAAGACATG  
AATGGTGCTTTCTCTGGGCTTGACAACTGAGGCGACTGATACTCCAAGGAAATCGGATC  
CGTTCATTACTAAAAAGCCTTCACTGGTTTTGGATGCATTGGAGCATCTAGACCTGAGT  
GACAACGCAATCATGTCTTTACAAGGCAATGCATTTTCACAAATGAAGAACTGCAACAA  
TTGCATTTAAATACATCAAGCCTTTTGTGCGATTGCCAGCTAAAATGGCTCCCACAGTGG  
GTGGCGGAAAACAACCTTTCAGAGCTTTGTAAATGCCAGTTGTGCCCATCCTCAGCTGCTA  
AAAGGAAGAAGCATTTTTGCTGTTAGCCCAGATGGCTTTGTGTGTGATGATTTTCCAAA  
CCCCAGATCACGGTTCAGCCAGAAACACAGTCGGCAATAAAAGGTTCCAATTTGAGTTTC  
ATCTGCTCAGCTGCCAGCAGAGTATTCCCCAATGACTTTTGTCTGGAAAAAGACAAT  
GAACTACTGCATGATGCTGAAATGGAAAATTATGCACACCTCCGGGCCAAGGTGGCGAG  
GTGATGGAGTATAACCACCATCCTTCGGCTGCGCGAGGTGGAATTTGCCAGTGAGGGGAAA  
TATCAGTGTGTCACTCTCCAATCACTTTGGTTCATCCTACTCTGTCAAAGCCAAGCTTACA  
GTAAATATGCTTCCCTCATTACCAAGACCCCCATGGATCTCACCATCCGAGCTGGGGCC  
ATGGCACGCTTGAGTGTGCTGCTGTGGGGCACCCAGCCCCCAGATAGCCTGGCAGAAG

GATGGGGGCACAGACTTCCCAGCTGCACGGGAGAGACGCATGCATGTGATGCCCGAGGAT  
GACGTGTTCTTTATCGTGGATGTGAAGATAGAGGACATTGGGGTATACAGCTGCACAGCT  
CAGAACAGTGCAGGAAGTATTTTCAGCAAATGCAACTCTGACTGTCCTAGAAACACCATCA  
TTTTTGGCGCCACTGTTGGACCGAACTGTAACCAAGGGAGAAACAGCCGTCCTACAGTGC  
ATTGCTGGAGGAAGCCCTCCCCCTAAACTGAACTGGACCAAAGATGATAGCCCATTTGGTG  
GTAACCGAGAGGCACTTTTTTTGCAGCAGGCAATCAGCTTCTGATTATTGTGGACTCAGAT  
GTCAGTGATGCTGGGAAATACACATGTGAGATGTCTAACACCCTTGGCACTGAGAGAGGA  
AACGTGCGCCTCAGTGTGATCCCCACTCCAACCTGCGACTCCCCCTCAGATGACAGCCCCA  
TCGTTAGACGATGACGGATGGGCCACTGTGGGTGTCGTGATCATAGCCGTGGTTTGCTGT  
GTGGTGGGCACGTCACCTCGTGTGGTGGTTCATCATATACCACACAAGGCGGAGGAATGAA  
GATTGCAGCATTACCAACACAGATGAGACCAACTTGCCAGCAGATATTCCTAGTTATTTG  
TCATCTCAGGGAACGTTAGCTGACAGGCAGGATGGGTACGTGTCTTCAGAAAAGTGGAAAGC  
CACCACCAGTTTGTACATCTTCAGGTGCTGGATTTTTCTTACCACAACATGACAGTAGT  
GGGACCTGCCATATTGACAATAGCAGTGAAGCTGATGTGGAAGCTGCCACAGATCTGTTC  
CTTTGTCCGTTTTTTGGGATCCACAGGCCCTATGTATTTGAAGGAAATGTGTATGGCTCA  
GATCCTTTTTGAAACATATCATAACAGTTGTCAGTCCCTGACCCAAGAACAGTTTTAATGGAC  
CACTATGAGCCCAGTTACATAAAGAAAAAGGAGTGCTACCCATGTTCTCATCCTTCAGAA  
GAATCCTGCGAACGGAGCTTCAGTAATATATCGTGGCCTTCACATGTGAGGAAGCTACTT  
AACACTAGTTACTCTCAATGAAGGACCTGGAATGAAAAATCTGTGTCTAAACAAGTCC  
TCTTTAGATTTTAGTGCAAATCCAGAGCCAGCGTTCGGTTGCCTCGAGTAATTCTTTTCATG  
GGTACCTTTGGAAAAGCTCTCAGGAGACCTCACCTAGATGCCATTTCAAGCTTTGGACAG  
CCATCAGATTGTCAGCCAAGAGCCTTTTATTTGAAAGCTCATTCTTCCCAGACTTGGAC  
TCTGGGTGAGAGGAAGATGGGAAAGAAAGGACAGATTTTCAGGAAGAAAATCACATTTGT  
ACCTTTAAACAGACTTTAGAAAACCTACAGGACTCCAAATTTTCAGTCTTATGACTTGGAC  
ACATAGACTGAATGAGACCAAAGGAAAAGCTTAACATACTACCTCAAGTGAACCTTTTATT  
TAAAAGAGAGAGAATCTTATGTTTTTTAAATGGAGTTATGAATTTTAAAAGGATAAAAAAT  
GCTTTATTTATACAGATGAACCAAATTACAAAAAGTTATGAAAATTTTATACTGGGAA  
TGATGCTCATATAAGAATACCTTTTTTAACTATTTTTTAACTTTGTTTTATGCAAAAAAG  
TATCTTACGTAAATTAATGATATAAATCATGATTATTTTATGTATTTTATAATGCCAGA  
TTTCTTTTTATGGAAAATGAGTTACTAAAGCATTTTAAATAATACCTGCCTTGTACCATT  
TTTTAAATAGAAGTTACTTCATTATATTTTGCACATTATATTTAATAAAAATGTGTCAATT  
TGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Figure 16

MSAPSLRARAAGLGLLLCAVLGRAGRSDSGRGELGQPSGVAERPCPTTCRCLGDLDD  
 CSRKRLARLPEPLPSWVARLDLSDHNRLSFIKASSMSHLQSLREVKLNNNELETI PNLGP  
 VSANITLLSLAGNRIVEILPEHLKEFQSLETLDLSSNNISELQTAFPALQLKYLYLNSN  
 RVTSMEPGYFDNLANTLLVLKLNRRNRISAIPPKMFKLPQLQHLELNRNKIKNVDGLTFQ  
 GLGALKSLKMQRNGVTKLMDGAFWGLSNMEILQLDHNNLTEITKGWLYGLLMLQELHLS  
 QNAINRISPDWEFCQKLSSELDLTFNHLRSLDDSSFLGLSLLNLT LHIGNNRVSYIADCA  
 FRGLSSLKTLDLKNNEISWTIEDMNGAFSGLDKLRRLLIQGNRIRSITKKAFTGLDALE  
 HLDLSDNAIMSLQGNAFSQMKKLOQLHLNLTSSLLCDCQLKWLQWVAENNFQSFVNASC  
 AHPQLLKGRSIFAVSPDGFVCDDFPKQITVQPETQSAIKGSNLSFICSAASSSDSPMT  
 FAWKKNELLHDAEMENYAHLRAQGGEVMEYTTILRLREVEFASEGKYQCVISNHFSS  
 YSVKAKLTVNMLPSFTKTPMDLTI RAGAMARLECAAVGHPAQIAWQKDGGTDFPAARE  
 RRMHVMPEDDVFFIVDVKIEDIGVYSCTAONSAGSISANATLTVLETSPFLRPLLDRTV  
 TKGETAVLQCIAGGSPPKLNWTKDSDPLVVTERHFFAAGNQLLIIVSDVSDAGKYTC  
 EMSNTLGTGERGNVRLSVIPTPTCDSPQMTAPSLDDDGWATVGVVIAVCCVVGTSLVW  
 VVIYHTRRRNEDCSITNTDETNLPA DIPSYLSSQGLADRQDGYVSSSESGSHHQFVTS  
 SGAGFFLPQHDSSTCHIDNSSEADVEAATDLFLCPFLGSTGPMYKGNVYGSDFPFETY  
 HTGCSPDPRTVLMDHYEPSYIKKKECYPCSHPSEESCERSFSNISWPSHVRKLLNTSYS  
 HNEGPGMKNLCLNKSSLD FSANPEPASVASSNSFMGTFGKALRRPHLDAYSSFGQPSDC  
 QPRAFYLKAHSSPDLDSGSEEDGKERTDFQEENHICTFKQTLNRYRTPNFQSYDLDT

N-glycosylation site at residues:

- 122-126
- 156-160
- 274-278
- 442-446
- 469-473
- 515-519
- 688-692
- 729-733
- 905-909
- 987-991
- 999-1003
- 1016-1020

Glycosaminoglycan attachment site at residues:

886-890

Casein kinase II phosphorylation site at residues:

99-103  
180-184  
263-267  
314-318  
324-328  
374-378  
383-387  
407-411  
524-528  
608-612  
692-696  
709-713  
731-735  
799-803  
843-847  
863-867  
907-911  
1003-1007  
1018-1022  
1073-1077  
1079-1083  
1081-1085

Tyrosine kinase phosphorylation site at residue:

667-675

N-myristoylation site at residues:

14-20  
36-42  
239-245  
257-263  
380-386  
427-433  
513-519  
588-594  
672-678  
683-689

774-780

933-999

Leucine zipper pattern at residues:

58-80

65-87

## Sequence Listing

<110> Genentech, Inc.  
 Fong, Sherman  
 Audrey Goddard  
 Gurney, Austin L.  
 Tumas, Daniel  
 Wood, William I.

<120> COMPOSITIONS AND METHODS FOR THE TREATMENT OF IMMUNE  
 RELATED DISEASES

<130> P1624R2

<141> 1999-09-14

<150> US 60/100,858  
 <151> 1998-09-17

<160> 39

<210> 1  
 <211> 1295  
 <212> DNA  
 <213> artificial sequence

<400> 1  
 cccagaagtt caagggcccc cggcctcctg cgctcctgcc gccgggaccc 50  
 tcgacctcct cagagcagcc ggctgccgcc ccgggaagat ggcgaggagg 100  
 agccgccacc gcctcctcct gctgctgctg cgctacctgg tggtcgcctt 150  
 gggctatcat aaggcctatg ggttttctgc cccaaaagac caacaagtag 200  
 tcacagcagt agagtaccaa gaggctatct tagcctgcaa aaccccaaag 250  
 aagactgttt cctccagatt agagtggaag aaactgggtc ggagtgtctc 300  
 ctttgtctac tatcaacaga ctcttcaagg tgattttaa aatcgagctg 350  
 agatgataga tttcaatata cggatcaaaa atgtgacaag aagtgatgag 400  
 gggaaatata gttgtgaagt tagtgcccca tctgagcaag gccaaaacct 450  
 ggaagaggat acagtcactc tggaagtatt agtggctcca gcagttccat 500  
 catgtgaagt accctcttct gctctgagtg gaactgtggt agagctacga 550  
 tgtcaagaca aagaaggaa tccagctcct gaatacacat ggtttaagga 600  
 tggcatccgt ttgctagaaa atcccagact tggctcccaa agcaccaaca 650  
 gctcatacac aatgaataca aaaactggaa ctctgcaatt taatactgtt 700  
 tccaaactgg aactggaga atattcctgt gaagcccga attctgttgg 750  
 atatcgagcagg tgcctgggaa aacgaatgca agtagatgat ctcaacataa 800  
 gtggcatcat agcagccgta gtagttgtgg ccttagtgat ctccgtttgt 850

ggccttggtg tatgctatgc tcagaggaaa ggctactttt caaaagaaac 900  
 ctccttccag aagagtaatt cttcatctaa agccacgaca atgagtgaaa 950  
 atgtgcagtg gctcacgcct gtaatcccag cactttggaa ggccgcgggcg 1000  
 ggcggatcac gaggtcagga gttctagacc agtctggcca atatggtgaa 1050  
 accccatctc tactaaaata caaaaattag ctgggcatgg tggcatgtgc 1100  
 ctgcagttcc agctgcttgg gagacaggag aatcacttga acccgggagg 1150  
 cggaggttgc agtgagctga gatcacgcca ctgcagtcca gcctgggtaa 1200  
 cagagcaaga ttccatctca aaaaataaaa taaataaata aataaatact 1250  
 ggtttttacc tgtagaattc ttacaataaa tatagcttga tattc 1295

<210> 2

<211> 312

<212> PRT

<213> artificial sequence

<400> 2

Met	Ala	Arg	Arg	Ser	Arg	His	Arg	Leu	Leu	Leu	Leu	Leu	Leu	Arg	1	5	10	15
Tyr	Leu	Val	Val	Ala	Leu	Gly	Tyr	His	Lys	Ala	Tyr	Gly	Phe	Ser	20	25	30	
Ala	Pro	Lys	Asp	Gln	Gln	Val	Val	Thr	Ala	Val	Glu	Tyr	Gln	Glu	35	40	45	
Ala	Ile	Leu	Ala	Cys	Lys	Thr	Pro	Lys	Lys	Thr	Val	Ser	Ser	Arg	50	55	60	
Leu	Glu	Trp	Lys	Lys	Leu	Gly	Arg	Ser	Val	Ser	Phe	Val	Tyr	Tyr	65	70	75	
Gln	Gln	Thr	Leu	Gln	Gly	Asp	Phe	Lys	Asn	Arg	Ala	Glu	Met	Ile	80	85	90	
Asp	Phe	Asn	Ile	Arg	Ile	Lys	Asn	Val	Thr	Arg	Ser	Asp	Ala	Gly	95	100	105	
Lys	Tyr	Arg	Cys	Glu	Val	Ser	Ala	Pro	Ser	Glu	Gln	Gly	Gln	Asn	110	115	120	
Leu	Glu	Glu	Asp	Thr	Val	Thr	Leu	Glu	Val	Leu	Val	Ala	Pro	Ala	125	130	135	
Val	Pro	Ser	Cys	Glu	Val	Pro	Ser	Ser	Ala	Leu	Ser	Gly	Thr	Val	140	145	150	
Val	Glu	Leu	Arg	Cys	Gln	Asp	Lys	Glu	Gly	Asn	Pro	Ala	Pro	Glu	155	160	165	
Tyr	Thr	Trp	Phe	Lys	Asp	Gly	Ile	Arg	Leu	Leu	Glu	Asn	Pro	Arg	170	175	180	
Leu	Gly	Ser	Gln	Ser	Thr	Asn	Ser	Ser	Tyr	Thr	Met	Asn	Thr	Lys				

	185		190		195
Thr Gly Thr Leu	Gln Phe Asn Thr Val	Ser Lys Leu Asp Thr Gly			
	200	205			210
Glu Tyr Ser Cys	Glu Ala Arg Asn Ser	Val Gly Tyr Arg Arg Cys			
	215	220			225
Pro Gly Lys Arg	Met Gln Val Asp Asp	Leu Asn Ile Ser Gly Ile			
	230	235			240
Ile Ala Ala Val	Val Val Val Ala Leu	Val Ile Ser Val Cys Gly			
	245	250			255
Leu Gly Val Cys	Tyr Ala Gln Arg Lys	Gly Tyr Phe Ser Lys Glu			
	260	265			270
Thr Ser Phe Gln	Lys Ser Asn Ser Ser	Ser Lys Ala Thr Thr Met			
	275	280			285
Ser Glu Asn Val	Gln Trp Leu Thr Pro	Val Ile Pro Ala Leu Trp			
	290	295			300
Lys Ala Ala Ala	Gly Gly Ser Arg Gly	Gln Glu Phe			
	305	310	312		

<210> 3  
 <211> 2033  
 <212> DNA  
 <213> artificial sequence

<400> 3  
 ccaggccggg aggcgacgcg cccagccgtc taaacgggaa cagccctggc 50  
 tgagggagct gcagcgcagc agagtatctg acggcgccag gttgcgtagg 100  
 tgccgcacga ggagttttcc cggcagcgag gaggtcctga gcagcatggc 150  
 ccggaggagc gccttcctcg ccgcccgcgt ctggctctgg agcatcctcc 200  
 tgtgcctgct ggcactgcgg gcggaggccg ggccgcccga ggaggagagc 250  
 ctgtacctat ggatcgatgc tcaccaggca agagtactca taggatttga 300  
 agaagatatc ctgattgttt cagaggggaa aatggcacct tttacacatg 350  
 atttcagaaa agcgcAACag agaatgccag ctattcctgt caatatccat 400  
 tccatgaatt ttacctggca agctgcaggg caggcagaat acttctatga 450  
 attcctgtcc ttgcgctccc tggataaagg catcatggca gatccaaccg 500  
 tcaatgtccc tctgctggga acagtgcctc acaaggcatc agttgttcaa 550  
 gttggtttcc catgtcttgg aaaacaggat ggggtggcag catttgaagt 600  
 ggatgtgatt gttatgaatt ctgaaggcaa caccattctc caaacacctc 650  
 aaaatgctat cttctttaaA acatgtcaac aagctgagtg cccaggcggg 700  
 tgccgaaatg gaggcttttg taatgaaaga cgcacctgcg agtgcctga 750

tggggtccac ggacctcact gtgagaaagc cttttgtacc ccacgatgta 800  
 tgaatggtgg actttgtgtg actcctggtt tctgcatctg cccacctgga 850  
 ttctatggag tgaactgtga caaagcaaac tgctcaacca cctgctttaa 900  
 tggagggacc tgtttctacc ctggaaaatg tatttgcctt ccaggactag 950  
 agggagagca gtgtgaaatc agcaaagtc cacaaccctg tcgaaatgga 1000  
 ggtaaagca ttggtaaaag caaatgtaag tgttccaaag gttaccaggg 1050  
 agacctctgt tcaaagcctg tctgagagcc tggctgtggt gcacatggaa 1100  
 cctgccatga acccaacaaa tgccaatgtc aagaaggttg gcatggaaga 1150  
 cactgcaata aaaggtacga agccagcctc atacatgccc tgaggccagc 1200  
 aggcgcccag ctcaggcagc acacgccttc acttaaaaag gccgaggagc 1250  
 ggcgggatcc acctgaatcc aattacatct ggtgaaactcc gacatctgaa 1300  
 acgttttaag ttacaccaag ttcatagcct ttgttaacct ttcatgtggt 1350  
 gaatgttcaa ataatgttca ttacacttaa gaatactggc ctgaatttta 1400  
 ttagcttcat tataaatcac tgagctgata tttactcttc cttttaagtt 1450  
 ttctaagtac gtctgtagca tgatggtata gattttcttg tttcagtgtt 1500  
 ttgggacaga ttttatatta tgtcaattga tcagggttaa attttcagtg 1550  
 tgtagttggc agatattttc aaaattacaa tgcatttatg gtgtctgggg 1600  
 gcaggggaac atcagaaagg ttaaattggg caaaaatgcy taagtcacaa 1650  
 gaatttgat ggtgcagtta atgttgaagt tacagcattt cagattttat 1700  
 tgtcagatat ttagatgttt gttacatttt taaaaattgc tcttaatttt 1750  
 taaactctca atacaatata ttttgacctt accattatc cagagattca 1800  
 gtattaaaaa aaaaaaaatt aactgtggt agtggcattt aaacaatata 1850  
 atatattcta aacacaatga aatagggat ataatgtatg aactttttgc 1900  
 attggcttga agcaatataa tatattgtaa acaaacaca gctcttacct 1950  
 aataaacatt ttatactggt tgtatgtata aaataaagg gctgctttag 2000  
 ttttttgaa aaaaaaaaaa aaaaaaaaaa aaa 2033

<210> 4

<211> 379

<212> PRT

<213> artificial sequence

<400> 4

Met Ala Arg Arg Ser Ala Phe Pro Ala Ala Ala Leu Trp Leu Trp  
 1 5 10 15

Ser Ile Leu Leu Cys Leu Leu Ala Leu Arg Ala Glu Ala Gly Pro  
 20 25 30  
 Pro Gln Glu Glu Ser Leu Tyr Leu Trp Ile Asp Ala His Gln Ala  
 35 40 45  
 Arg Val Leu Ile Gly Phe Glu Glu Asp Ile Leu Ile Val Ser Glu  
 50 55 60  
 Gly Lys Met Ala Pro Phe Thr His Asp Phe Arg Lys Ala Gln Gln  
 65 70 75  
 Arg Met Pro Ala Ile Pro Val Asn Ile His Ser Met Asn Phe Thr  
 80 85 90  
 Trp Gln Ala Ala Gly Gln Ala Glu Tyr Phe Tyr Glu Phe Leu Ser  
 95 100 105  
 Leu Arg Ser Leu Asp Lys Gly Ile Met Ala Asp Pro Thr Val Asn  
 110 115 120  
 Val Pro Leu Leu Gly Thr Val Pro His Lys Ala Ser Val Val Gln  
 125 130 135  
 Val Gly Phe Pro Cys Leu Gly Lys Gln Asp Gly Val Ala Ala Phe  
 140 145 150  
 Glu Val Asp Val Ile Val Met Asn Ser Glu Gly Asn Thr Ile Leu  
 155 160 165  
 Gln Thr Pro Gln Asn Ala Ile Phe Phe Lys Thr Cys Gln Gln Ala  
 170 175 180  
 Glu Cys Pro Gly Gly Cys Arg Asn Gly Gly Phe Cys Asn Glu Arg  
 185 190 195  
 Arg Ile Cys Glu Cys Pro Asp Gly Phe His Gly Pro His Cys Glu  
 200 205 210  
 Lys Ala Leu Cys Thr Pro Arg Cys Met Asn Gly Gly Leu Cys Val  
 215 220 225  
 Thr Pro Gly Phe Cys Ile Cys Pro Pro Gly Phe Tyr Gly Val Asn  
 230 235 240  
 Cys Asp Lys Ala Asn Cys Ser Thr Thr Cys Phe Asn Gly Gly Thr  
 245 250 255  
 Cys Phe Tyr Pro Gly Lys Cys Ile Cys Pro Pro Gly Leu Glu Gly  
 260 265 270  
 Glu Gln Cys Glu Ile Ser Lys Cys Pro Gln Pro Cys Arg Asn Gly  
 275 280 285  
 Gly Lys Cys Ile Gly Lys Ser Lys Cys Lys Cys Ser Lys Gly Tyr  
 290 295 300  
 Gln Gly Asp Leu Cys Ser Lys Pro Val Cys Glu Pro Gly Cys Gly  
 305 310 315

Ala His Gly Thr Cys His Glu Pro Asn Lys Cys Gln Cys Gln Glu  
 320 325 330

Gly Trp His Gly Arg His Cys Asn Lys Arg Tyr Glu Ala Ser Leu  
 335 340 345

Ile His Ala Leu Arg Pro Ala Gly Ala Gln Leu Arg Gln His Thr  
 350 355 360

Pro Ser Leu Lys Lys Ala Glu Glu Arg Arg Asp Pro Pro Glu Ser  
 365 370 375

Asn Tyr Ile Trp  
 379

<210> 5  
 <211> 1857  
 <212> DNA  
 <213> artificial sequence

<400> 5  
 gtctgttccc aggagtcctt cggcggctgt tgtgtcagtg gcctgatcgc 50  
 gatggggaca aaggcgcaag tcgagaggaa actgttgtgc ctcttcatat 100  
 tggcgatcct gttgtgctcc ctggcattgg gcagtgttac agtgactct 150  
 tctgaacctg aagtcagaat tcttgagaat aatcctgtga agttgtcctg 200  
 tgcctactcg ggcttttctt ctccccgtgt ggagtggaag tttgaccaag 250  
 gagacaccac cagactcgtt tgctataata acaagatcac agcttcctat 300  
 gaggaccggg tgaccttctt gccaaactggg atcaccttca agtccgtgac 350  
 acgggaagac actgggacat aacttgtat ggtctctgag gaaggcggca 400  
 acagctatgg ggaggtcaag gtcaagctca tcgtgcttgt gcctccatcc 450  
 aagcctacag ttaacatccc ctctctgcc accattggga accgggcagt 500  
 gctgacatgc tcagaacaag atggttcccc accttctgaa tacacctggt 550  
 tcaaagatgg gatagtgatg cctacgaatc caaaagcac ccgtgccttc 600  
 agcaactctt cctatgtcct gaatcccaca acaggagagc tggctcttga 650  
 tcccctgtca gcctctgata ctggagaata cagctgtgag gcacggaatg 700  
 ggtatgggac acctatgact tcaaatgctg tgcgcatgga agctgtggag 750  
 cggaatgtgg gggcatcgt gccagccgtc cttgtaacct tgattctcct 800  
 gggaatcttg gtttttggca tctggtttgc ctatagccga ggccactttg 850  
 acagaacaaa gaaagggact tcgagtaaga aggtgattta cagccagcct 900  
 agtgcccga gtaaggaga attcaaacag acctcgtcat tcttgggtgtg 950  
 agcctggctg gctcaccgcc tatcatctgc atttgcctta ctcaggtgct 1000

accggactct ggcccctgat gtctgtagtt tcacaggatg ccttatttgt 1050  
 cttctacacc ccacagggcc ccctacttct tcggatgtgt ttttaataat 1100  
 gtcagctatg tgccccatcc tccttcatgc cctccctccc tttcctacca 1150  
 ctgctgagtg gcoctggaact tgtttaaagt gtttattccc catttctttg 1200  
 agggatcagg aaggaatcct gggatgcca ttgacttccc ttctaagtag 1250  
 acagcaaaaa tggcgggggt cgcaggaatc tgactcaac tgcccacctg 1300  
 gctggcaggg atctttgaat aggtatcttg agcttggttc tgggctcttt 1350  
 ccttgtgtac tgacgaccag ggccagctgt tctagagcgg gaattagagg 1400  
 ctagagcggc tgaaatggtt gtttggatgac gacactgggg tccttccatc 1450  
 tctggggccc actctcttct gtcttcccat ggaagtgcc actgggatcc 1500  
 ctctgccttg tcctcctgaa tacaagctga ctgacattga ctgtgtctgt 1550  
 ggaaaatggg agctcttgtt gtggagagca tagtaaattt tcagagaact 1600  
 tgaagccaaa aggatttaaa accgctgctc taaagaaaag aaaactggag 1650  
 gctgggcgca gtggctcacg cctgtaatcc cagaggctga ggcaggcggg 1700  
 tcacctgagg tcgggagttc gggatcagcc tgaccaacat ggagaaacct 1750  
 tactggaaat acaaagttag ccaggcatgg tggatgatgc ctgtagtccc 1800  
 agctgctcag gagcctggca acaagagcaa aactccagct caaaaaaaaaa 1850  
 aaaaaaa 1857

<210> 6  
 <211> 299  
 <212> PRT  
 <213> artificial sequence

<400> 6  
 Met Gly Thr Lys Ala Gln Val Glu Arg Lys Leu Leu Cys Leu Phe  
 1 5 10 15  
 Ile Leu Ala Ile Leu Leu Cys Ser Leu Ala Leu Gly Ser Val Thr  
 20 25 30  
 Val His Ser Ser Glu Pro Glu Val Arg Ile Pro Glu Asn Asn Pro  
 35 40 45  
 Val Lys Leu Ser Cys Ala Tyr Ser Gly Phe Ser Ser Pro Arg Val  
 50 55 60  
 Glu Trp Lys Phe Asp Gln Gly Asp Thr Thr Arg Leu Val Cys Tyr  
 65 70 75  
 Asn Asn Lys Ile Thr Ala Ser Tyr Glu Asp Arg Val Thr Phe Leu  
 80 85 90  
 Pro Thr Gly Ile Thr Phe Lys Ser Val Thr Arg Glu Asp Thr Gly

	95		100		105
Thr Tyr Thr Cys Met Val Ser Glu Glu Gly Gly Asn Ser Tyr Gly	110		115		120
Glu Val Lys Val Lys Leu Ile Val Leu Val Pro Pro Ser Lys Pro	125		130		135
Thr Val Asn Ile Pro Ser Ser Ala Thr Ile Gly Asn Arg Ala Val	140		145		150
Leu Thr Cys Ser Glu Gln Asp Gly Ser Pro Pro Ser Glu Tyr Thr	155		160		165
Trp Phe Lys Asp Gly Ile Val Met Pro Thr Asn Pro Lys Ser Thr	170		175		180
Arg Ala Phe Ser Asn Ser Ser Tyr Val Leu Asn Pro Thr Thr Gly	185		190		195
Glu Leu Val Phe Asp Pro Leu Ser Ala Ser Asp Thr Gly Glu Tyr	200		205		210
Ser Cys Glu Ala Arg Asn Gly Tyr Gly Thr Pro Met Thr Ser Asn	215		220		225
Ala Val Arg Met Glu Ala Val Glu Arg Asn Val Gly Val Ile Val	230		235		240
Ala Ala Val Leu Val Thr Leu Ile Leu Leu Gly Ile Leu Val Phe	245		250		255
Gly Ile Trp Phe Ala Tyr Ser Arg Gly His Phe Asp Arg Thr Lys	260		265		270
Lys Gly Thr Ser Ser Lys Lys Val Ile Tyr Ser Gln Pro Ser Ala	275		280		285
Arg Ser Glu Gly Glu Phe Lys Gln Thr Ser Ser Phe Leu Val	290		295		299

<210> 7

<211> 2755

<212> DNA

<213> artificial sequence

<400> 7

```

gggggtagg gaggaaggaa tccaccccca ccccccaaa cccttttctt 50
ctcctttcct ggcttcggac attggagcac taaatgaact tgaattgtgt 100
ctgtggcgag caggatggtc gctgttactt tgtgatgaga tcggggatga 150
attgctcgct ttaaaaatgc tgctttggat tctgttgctg gagacgtctc 200
tttgttttgc cgctggaaac gttacagggg acgtttgcaa agagaagatc 250
tgttcctgca atgagataga aggggacctt cacgtagact gtgaaaaaaaa 300
gggcttcaca agtctgcagc gtttactgac cccgacttcc cagttttacc 350

```

atattatttct gcatggcaat tccctcactc gacttttccc taatgagttc 400  
gctaactttt ataatgcggt tagtttgac atggaaaaca atggcttgca 450  
tgaaatcggt ccgggggctt ttctggggct gcagctggtg aaaaggctgc 500  
acatcaacaa caacaagatc aagtcttttc gaaagcagac ttttctgggg 550  
ctggacgatc tggaatatct ccaggctgat ttttaatttat tacgagatat 600  
agaccggggg gccttccagg acttgaacaa gctggaggtg ctcattttaa 650  
atgacaatct catcagcacc ctacctgcca acgtgttcca gtatgtgcc 700  
atcaccacc tcgacctcg gggtaacagg ctgaaaacgc tgcctatga 750  
ggaggtcttg gagcaaatcc ctggattgic ggagatcctg ctagaggata 800  
acccttggga ctgcacctgt gatctgctct ccctgaaaga atggctggaa 850  
aacattccca agaatgcctt gatcggcga gtggtctgcg aagccccac 900  
cagactgcag ggtaaagacc tcaatgaaac caccgaacag gacttgtgtc 950  
ctttgaaaaa ccgagtggat tctagtctcc cggcgcccc tgccaagaa 1000  
gagacctttg ctctggacc cctgccaaact ctttcaaga caaatgggca 1050  
agaggatcat gccacaccag ggtctgctcc aaacggaggt acaaagatcc 1100  
caggcaactg gcagatcaaa atcagacca cagcagcgt agcgacgggt 1150  
agctccagga acaaaccctt agctaacagt ttaccctgcc ctgggggctg 1200  
cagctgcgac cacatcccag ggtcgggttt aaagatgaac tgcaacaaca 1250  
ggaacgtgag cagcttggtt gatttgaagc ccaagctctc taactgtcag 1300  
gagcttttcc tacgagataa caagatccac agcatccgaa aatcgactt 1350  
tgtggattac aagaacctca ttctgttga tctgggcaac aataacatcg 1400  
ctactgtaga gaacaacact ttcaagaacc ttttggacct caggtggcta 1450  
tacatggata gcaattacct ggacacgctg tcccgggaga aattcgcggg 1500  
gctgcaaac ctagagtacc tgaacgtgga gtacaacgct atccagctca 1550  
tcctcccggg cactttcaat gccatgcca aactgaggat cctcattctc 1600  
aacaacaacc tgctgaggtc cctgcctgtg gacgtgttcg ctggggtctc 1650  
gctctctaaa ctcagcctgc acaacaatta cttcatgtac ctcccgttg 1700  
caggggtgct ggaccagtta acctccatca tccagataga cctccacgga 1750  
aaccctggg agtgctcctg cacaattgtg ctttcaagc agtgggcaga 1800  
acgcttgggt tccgaagtgc tgatgagcga cctcaagtgt gagacgccgg 1850  
tgaacttctt tagaaaggat ttcattgctcc tctccaatga cgagatctgc 1900

cctcagctgt acgctaggat ctgccccacg ttaacttcgc acagtaaaaa 1950  
 cagcactggg ttggcggaga ccgggacgca ctccaactcc tacctagaca 2000  
 ccagcagggg gtccatctcg gtgttggtcc cgggactgct gctggtgttt 2050  
 gtcacctccg ccttcaccgt ggtgggcatg ctctgtgtta tcttgaggaa 2100  
 ccgaaagcgg tccaagagac gagatgcca ctcctccgcg tccgagatta 2150  
 attcoctaca gacagtctgt gactcttctt actggcacia tgggccttac 2200  
 aacgcagatg gggccccacag agtgtatgac tgtggctctc actcgctctc 2250  
 agactaagac cccaaccca ataggggagg gcagagggaa ggcgatacat 2300  
 ccttccccac cgcaggcacc ccgggggctg gaggggcgtg taccxaaatc 2350  
 cccgcgccat cagcctggat gggcataagt agataaataa ctgtgagctc 2400  
 gcacaaccga aagggcctga ccccttactt agtccctcc ttgaaacaaa 2450  
 gagcagactg tggagagctg ggagagcgca gccagctcgc tctttgctga 2500  
 gagccccttt tgacagaaag ccagcacga cctgctgga agaactgaca 2550  
 gtgccctcgc cctcggcccc ggggcctgtg gggttggatg ccgcggttct 2600  
 atacatatac acatatatcc acatctatat agagagatag atatctattt 2650  
 ttcccctgtg gattagcccc gtgatggctc cctgttggtc acgcagggat 2700  
 gggcagttgc acgaaggcat gaatgtattg taaataagta actttgactt 2750  
 ctgac 2755

<210> 8  
 <211> 696  
 <212> PRT  
 <213> artificial sequence

<400> 8  
 Met Leu Leu Trp Ile Leu Leu Leu Glu Thr Ser Leu Cys Phe Ala  
 1 5 10 15  
 Ala Gly Asn Val Thr Gly Asp Val Cys Lys Glu Lys Ile Cys Ser  
 20 25 30  
 Cys Asn Glu Ile Glu Gly Asp Leu His Val Asp Cys Glu Lys Lys  
 35 40 45  
 Gly Phe Thr Ser Leu Gln Arg Phe Thr Ala Pro Thr Ser Gln Phe  
 50 55 60  
 Tyr His Leu Phe Leu His Gly Asn Ser Leu Thr Arg Leu Phe Pro  
 65 70 75  
 Asn Glu Phe Ala Asn Phe Tyr Asn Ala Val Ser Leu His Met Glu  
 80 85 90

Asn Asn Gly Leu His Glu Ile Val Pro Gly Ala Phe Leu Gly Leu  
 95 100 105

Gln Leu Val Lys Arg Leu His Ile Asn Asn Asn Lys Ile Lys Ser  
 110 115 120

Phe Arg Lys Gln Thr Phe Leu Gly Leu Asp Asp Leu Glu Tyr Leu  
 125 130 135

Gln Ala Asp Phe Asn Leu Leu Arg Asp Ile Asp Pro Gly Ala Phe  
 140 145 150

Gln Asp Leu Asn Lys Leu Glu Val Leu Ile Leu Asn Asp Asn Leu  
 155 160 165

Ile Ser Thr Leu Pro Ala Asn Val Phe Gln Tyr Val Pro Ile Thr  
 170 175 180

His Leu Asp Leu Arg Gly Asn Arg Leu Lys Thr Leu Pro Tyr Glu  
 185 190 195

Glu Val Leu Glu Gln Ile Pro Gly Ile Ala Glu Ile Leu Leu Glu  
 200 205 210

Asp Asn Pro Trp Asp Cys Thr Cys Asp Leu Leu Ser Leu Lys Glu  
 215 220 225

Trp Leu Glu Asn Ile Pro Lys Asn Ala Leu Ile Gly Arg Val Val  
 230 235 240

Cys Glu Ala Pro Thr Arg Leu Gln Gly Lys Asp Leu Asn Glu Thr  
 245 250 255

Thr Glu Gln Asp Leu Cys Pro Leu Lys Asn Arg Val Asp Ser Ser  
 260 265 270

Leu Pro Ala Pro Pro Ala Gln Glu Glu Thr Phe Ala Pro Gly Pro  
 275 280 285

Leu Pro Thr Pro Phe Lys Thr Asn Gly Gln Glu Asp His Ala Thr  
 290 295 300

Pro Gly Ser Ala Pro Asn Gly Gly Thr Lys Ile Pro Gly Asn Trp  
 305 310 315

Gln Ile Lys Ile Arg Pro Thr Ala Ala Ile Ala Thr Gly Ser Ser  
 320 325 330

Arg Asn Lys Pro Leu Ala Asn Ser Leu Pro Cys Pro Gly Gly Cys  
 335 340 345

Ser Cys Asp His Ile Pro Gly Ser Gly Leu Lys Met Asn Cys Asn  
 350 355 360

Asn Arg Asn Val Ser Ser Leu Ala Asp Leu Lys Pro Lys Leu Ser  
 365 370 375

Asn Val Gln Glu Leu Phe Leu Arg Asp Asn Lys Ile His Ser Ile  
 380 385 390

Arg Lys Ser His Phe Val Asp Tyr Lys Asn Leu Ile Leu Leu Asp

	395		400		405
Leu Gly Asn Asn Asn Ile Ala Thr Val	Glu Asn Asn Thr Phe Lys				
	410		415		420
Asn Leu Leu Asp Leu Arg Trp Leu Tyr	Met Asp Ser Asn Tyr Leu				
	425		430		435
Asp Thr Leu Ser Arg Glu Lys Phe Ala	Gly Leu Gln Asn Leu Glu				
	440		445		450
Tyr Leu Asn Val Glu Tyr Asn Ala Ile	Gln Leu Ile Leu Pro Gly				
	455		460		465
Thr Phe Asn Ala Met Pro Lys Leu Arg	Ile Leu Ile Leu Asn Asn				
	470		475		480
Asn Leu Leu Arg Ser Leu Pro Val Asp	Val Phe Ala Gly Val Ser				
	485		490		495
Leu Ser Lys Leu Ser Leu His Asn Asn	Tyr Phe Met Tyr Leu Pro				
	500		505		510
Val Ala Gly Val Leu Asp Gln Leu Thr	Ser Ile Ile Gln Ile Asp				
	515		520		525
Leu His Gly Asn Pro Trp Glu Cys Ser	Cys Thr Ile Val Pro Phe				
	530		535		540
Lys Gln Trp Ala Glu Arg Leu Gly Ser	Glu Val Leu Met Ser Asp				
	545		550		555
Leu Lys Cys Glu Thr Pro Val Asn Phe	Phe Arg Lys Asp Phe Met				
	560		565		570
Leu Leu Ser Asn Asp Glu Ile Cys Pro	Gln Leu Tyr Ala Arg Ile				
	575		580		585
Ser Pro Thr Leu Thr Ser His Ser Lys	Asn Ser Thr Gly Leu Ala				
	590		595		600
Glu Thr Gly Thr His Ser Asn Ser Tyr	Leu Asp Thr Ser Arg Val				
	605		610		615
Ser Ile Ser Val Leu Val Pro Gly Leu	Leu Leu Val Phe Val Thr				
	620		625		630
Ser Ala Phe Thr Val Val Gly Met Leu	Val Phe Ile Leu Arg Asn				
	635		640		645
Arg Lys Arg Ser Lys Arg Arg Asp Ala	Asn Ser Ser Ala Ser Glu				
	650		655		660
Ile Asn Ser Leu Gln Thr Val Cys Asp	Ser Ser Tyr Trp His Asn				
	665		670		675
Gly Pro Tyr Asn Ala Asp Gly Ala His	Arg Val Tyr Asp Cys Gly				
	680		685		690
Ser His Ser Leu Ser Asp					
	695		696		

<210> 9

<211> 3659

<212> DNA

<213> artificial sequence

<400> 9

gtaactgaag tcaggctttt catttgggaa gccccctcaa cagaattcgg 50  
tcattctcca agttatggtg gacgtacttc tgttgttctc cctctgcttg 100  
ctttttcaca ttagcagacc ggacttaagt cacaacagat tatctttcat 150  
caaggcaagt tccatgagcc accttcaaag ccttcgagaa gtgaaactga 200  
acaacaatga attggagacc attccaaatc tgggaccagt ctcggcaaat 250  
attacacttc tctccttggc tggaaacagg attggtgaaa tactccctga 300  
acatctgaaa gagtttcagt cccttgaaac tttggacctt agcagcaaca 350  
atatttcaga gctccaaact gcatttccag ccctacagct caaatatctg 400  
tatctcaaca gcaaccgagt cacatcaatg gaacctgggt attttgacia 450  
tttggccaac acactccttg tgttaaagct gaacaggaac cgaatctcag 500  
ctatcccacc caagatgttt aaactgcccc aactgcaaca tctcgaattg 550  
aaccgaaaca agattaaaaa tgtagatgga ctgacattcc aaggccttgg 600  
tgctctgaag tctctgaaaa tgcaaagaaa tggagtaacg aaacttatgg 650  
atggagcttt ttgggggctg agcaacatgg aaattttgca gctggacat 700  
aacaacctaa cagagattac caaaggctgg ctttacggct tgctgatgct 750  
gcaggaactt catctcagcc aaaatgccat caacaggatc agccctgatg 800  
cctgggagtt ctgccagaag ctcagtgagc tggacctaac tttcaatcac 850  
ttatcaagggt tagatgattc aagcttcctt ggcttaagct tactaaatac 900  
ctgcacattg ggaacaacag agtcagctac attgctgatt gtgccttccg 950  
ggggctttcc agtttaaaga ctttgatct gaagaacaat gaaatttcct 1000  
ggactattga agacatgaat ggtgctttct ctgggcttga caaactgagg 1050  
cgactgatac tccaaggaaa tggatccgt tctattacta aaaaagcctt 1100  
cactggtttg gatgcattgg agcatctaga cctgagtgac aacgcaatca 1150  
tgtctttaca aggcaatgca ttttcacaaa tgaagaaact gcaacaattg 1200  
catttaata catcaagcct tttgtgcat tgccagctaa aatggctccc 1250  
acagtgggtg gcggaaaaca actttcagag ctttgtaaat gccagttgtg 1300  
cccatcctca gctgctaaaa ggaagaagca tttttgctgt tagcccagat 1350

ggctttgtgt gtgatgattt tcccaaacc cagatcacgg ttcagccaga 1400  
aacacagtcg gcaataaaaag gttccaattt gagtttcatc tgctcagctg 1450  
ccagcagcag tgattcccca atgacttttg cttggaaaaa agacaatgaa 1500  
ctactgcatg atgctgaaat ggaaaattat gcacacctcc gggcccaagg 1550  
tggcgagggt atggagtata ccaccatcct tcggctgcgc gaggtggaat 1600  
ttgccagtga ggggaaatat cagtgtgtca tctccaatca ctttggttca 1650  
tcctactctg tcaaagcaa gcttacagta aatatgcttc cctcattcac 1700  
caagaccccc atggatctca ccatccgagc tggggccatg gcacgcttgg 1750  
agtgtgctgc tgtggggcac ccagcccccc agatagcctg gcagaaggat 1800  
gggggcacag acttcccagc tgcacgggag agacgcatgc atgtgatgcc 1850  
cgaggatgac gtgttcttta tcgtggatgt gaagatagag gacattgggg 1900  
tatacagctg cacagctcag aacagtgcag gaagtatttc agcaaatgca 1950  
actctgactg tcttagaaac accatcattt ttgcgggcac tgttgaccg 2000  
aactgtaacc aaggagaaa cagccgtcct acagtgcatt gctggaggaa 2050  
gccctcccc taaactgaac tggaccaaaag atgatagccc attggtggta 2100  
accgagaggc acttttttgc agcaggcaat cagcttctga ttattgtgga 2150  
ctcagatgtc agtgatgctg ggaaatacac atgtgagatg tctaaccacc 2200  
ttggcactga gagaggaaac gtgcgcctca gtgtgatccc cactccaacc 2250  
tgcgactccc ctcagatgac agccccatcg ttagacgatg acggatgggc 2300  
cactgtgggt gtctgatca tagccgtgggt ttgctgtgtg gtgggcacgt 2350  
cactcgtgtg ggtggtcac atataccaca caaggcggag gaatgaagat 2400  
tgcagcatta ccaacacaga tgagaccaac ttgccagcag atattcctag 2450  
ttatttgtca tctcagggaa cgttagctga caggcaggat gggtagctgt 2500  
cttcagaaag tggaagccac caccagtttg tcacatcttc aggtgctgga 2550  
tttttcttac cacaacatga cagtagtggg acctgccata ttgacaatag 2600  
cagtgaagct gatgtggaag ctgccacaga tctgttcctt tgtccgtttt 2650  
tgggatccac aggcctatg tatttgaagg gaaatgtgta tggctcagat 2700  
ccttttgaaa catatcatac aggttgcaat cctgacccaa gaacagtttt 2750  
aatggaccac atgagcccag ttacataaag aaaaaggagt gctacccatg 2800  
ttctcatcct tcagaagaat cctgcgaacg gagcttcagt aatatatcgt 2850  
ggccttcaca tgtgaggaag ctacttaaca ctagttactc tcacaatgaa 2900

ggacctggaa tgaaaaatct gtgtctaaac aagtcctctt tagatttttag 2950  
 tgcaaatcca gagccagcgt cggttgcctc gagtaattct ttcattggta 3000  
 cctttggaaa agctctcagg agacctcacc tagatgccta ttcaagcttt 3050  
 ggacagccat cagattgtca gccaaagagcc ttttatttga aagctcattc 3100  
 ttccccagac ttggactctg ggtcagagga agatgggaaa gaaaggacag 3150  
 attttcagga agaaaatcac atttgtacct taaacagact ttagaaaact 3200  
 acaggactcc aaattttcag tcttatgact tggacacata gactgaatga 3250  
 gaccaaagga aaagcttaac atactacctc aagtgaactt ttatttataa 3300  
 gagagagaat cttatgtttt ttaaatggag ttatgaattt taaaaggata 3350  
 aaaatgcttt atttatacag atgaaccaa attacaaaaa gttatgaaaa 3400  
 tttttatact gggaatgatg ctcatataag aatacctttt taaactattt 3450  
 tttaactttg ttttatgcaa aaaagtatct tacgtaaatt aatgatataa 3500  
 atcatgatta ttttatgtat ttttataatg ccagatttct ttttatggaa 3550  
 aatgagttac taaagcattt taaataatac ctgccttgta ccatttttta 3600  
 aatagaagtt acttcattat attttgcaca ttatatttaa taaaatgtgt 3650  
 caatttgaa 3659

<210> 10  
 <211> 1059  
 <212> PRT  
 <213> artificial sequence

<400> 10  
 Met Val Asp Val Leu Leu Leu Phe Ser Leu Cys Leu Leu Phe His  
 1 5 10 15  
 Ile Ser Arg Pro Asp Leu Ser His Asn Arg Leu Ser Phe Ile Lys  
 20 25 30  
 Ala Ser Ser Met Ser His Leu Gln Ser Leu Arg Glu Val Lys Leu  
 35 40 45  
 Asn Asn Asn Glu Leu Glu Thr Ile Pro Asn Leu Gly Pro Val Ser  
 50 55 60  
 Ala Asn Ile Thr Leu Leu Ser Leu Ala Gly Asn Arg Ile Val Glu  
 65 70 75  
 Ile Leu Pro Glu His Leu Lys Glu Phe Gln Ser Leu Glu Thr Leu  
 80 85 90  
 Asp Leu Ser Ser Asn Asn Ile Ser Glu Leu Gln Thr Ala Phe Pro  
 95 100 105  
 Ala Leu Gln Leu Lys Tyr Leu Tyr Leu Asn Ser Asn Arg Val Thr

	110		115		120
Ser Met Glu Pro Gly Tyr Phe Asp Asn Leu Ala Asn Thr Leu Leu					
	125		130		135
Val Leu Lys Leu Asn Arg Asn Arg Ile Ser Ala Ile Pro Pro Lys					
	140		145		150
Met Phe Lys Leu Pro Gln Leu Gln His Leu Glu Leu Asn Arg Asn					
	155		160		165
Lys Ile Lys Asn Val Asp Gly Leu Thr Phe Gln Gly Leu Gly Ala					
	170		175		180
Leu Lys Ser Leu Lys Met Gln Arg Asn Gly Val Thr Lys Leu Met					
	185		190		195
Asp Gly Ala Phe Trp Gly Leu Ser Asn Met Glu Ile Leu Gln Leu					
	200		205		210
Asp His Asn Asn Leu Thr Glu Ile Thr Lys Gly Trp Leu Tyr Gly					
	215		220		225
Leu Leu Met Leu Gln Glu Leu His Leu Ser Gln Asn Ala Ile Asn					
	230		235		240
Arg Ile Ser Pro Asp Ala Trp Glu Phe Cys Gln Lys Leu Ser Glu					
	245		250		255
Leu Asp Leu Thr Phe Asn His Leu Ser Arg Leu Asp Asp Ser Ser					
	260		265		270
Phe Leu Gly Leu Ser Leu Leu Asn Thr Leu His Ile Gly Asn Asn					
	275		280		285
Arg Val Ser Tyr Ile Ala Asp Cys Ala Phe Arg Gly Leu Ser Ser					
	290		295		300
Leu Lys Thr Leu Asp Leu Lys Asn Asn Glu Ile Ser Trp Thr Ile					
	305		310		315
Glu Asp Met Asn Gly Ala Phe Ser Gly Leu Asp Lys Leu Arg Arg					
	320		325		330
Leu Ile Leu Gln Gly Asn Arg Ile Arg Ser Ile Thr Lys Lys Ala					
	335		340		345
Phe Thr Gly Leu Asp Ala Leu Glu His Leu Asp Leu Ser Asp Asn					
	350		355		360
Ala Ile Met Ser Leu Gln Gly Asn Ala Phe Ser Gln Met Lys Lys					
	365		370		375
Leu Gln Gln Leu His Leu Asn Thr Ser Ser Leu Leu Cys Asp Cys					
	380		385		390
Gln Leu Lys Trp Leu Pro Gln Trp Val Ala Glu Asn Asn Phe Gln					
	395		400		405
Ser Phe Val Asn Ala Ser Cys Ala His Pro Gln Leu Leu Lys Gly					
	410		415		420

Arg Ser Ile Phe Ala Val Ser Pro Asp Gly Phe Val Cys Asp Asp	425	430	435
Phe Pro Lys Pro Gln Ile Thr Val Gln Pro Glu Thr Gln Ser Ala	440	445	450
Ile Lys Gly Ser Asn Leu Ser Phe Ile Cys Ser Ala Ala Ser Ser	455	460	465
Ser Asp Ser Pro Met Thr Phe Ala Trp Lys Lys Asp Asn Glu Leu	470	475	480
Leu His Asp Ala Glu Met Glu Asn Tyr Ala His Leu Arg Ala Gln	485	490	495
Gly Gly Glu Val Met Glu Tyr Thr Thr Ile Leu Arg Leu Arg Glu	500	505	510
Val Glu Phe Ala Ser Glu Gly Lys Tyr Gln Cys Val Ile Ser Asn	515	520	525
His Phe Gly Ser Ser Tyr Ser Val Lys Ala Lys Leu Thr Val Asn	530	535	540
Met Leu Pro Ser Phe Thr Lys Thr Pro Met Asp Leu Thr Ile Arg	545	550	555
Ala Gly Ala Met Ala Arg Leu Glu Cys Ala Ala Val Gly His Pro	560	565	570
Ala Pro Gln Ile Ala Trp Gln Lys Asp Gly Gly Thr Asp Phe Pro	575	580	585
Ala Ala Arg Glu Arg Arg Met His Val Met Pro Glu Asp Asp Val	590	595	600
Phe Phe Ile Val Asp Val Lys Ile Glu Asp Ile Gly Val Tyr Ser	605	610	615
Cys Thr Ala Gln Asn Ser Ala Gly Ser Ile Ser Ala Asn Ala Thr	620	625	630
Leu Thr Val Leu Glu Thr Pro Ser Phe Leu Arg Pro Leu Leu Asp	635	640	645
Arg Thr Val Thr Lys Gly Glu Thr Ala Val Leu Gln Cys Ile Ala	650	655	660
Gly Gly Ser Pro Pro Pro Lys Leu Asn Trp Thr Lys Asp Asp Ser	665	670	675
Pro Leu Val Val Thr Glu Arg His Phe Phe Ala Ala Gly Asn Gln	680	685	690
Leu Leu Ile Ile Val Asp Ser Asp Val Ser Asp Ala Gly Lys Tyr	695	700	705
Thr Cys Glu Met Ser Asn Thr Leu Gly Thr Glu Arg Gly Asn Val	710	715	720

Arg Leu Ser Val Ile Pro Thr Pro Thr Cys Asp Ser Pro Gln Met  
 725 730 735  
 Thr Ala Pro Ser Leu Asp Asp Asp Gly Trp Ala Thr Val Gly Val  
 740 745 750  
 Val Ile Ile Ala Val Val Cys Cys Val Val Gly Thr Ser Leu Val  
 755 760 765  
 Trp Val Val Ile Ile Tyr His Thr Arg Arg Arg Asn Glu Asp Cys  
 770 775 780  
 Ser Ile Thr Asn Thr Asp Glu Thr Asn Leu Pro Ala Asp Ile Pro  
 785 790 795  
 Ser Tyr Leu Ser Ser Gln Gly Thr Leu Ala Asp Arg Gln Asp Gly  
 800 805 810  
 Tyr Val Ser Ser Glu Ser Gly Ser His His Gln Phe Val Thr Ser  
 815 820 825  
 Ser Gly Ala Gly Phe Phe Leu Pro Gln His Asp Ser Ser Gly Thr  
 830 835 840  
 Cys His Ile Asp Asn Ser Ser Glu Ala Asp Val Glu Ala Ala Thr  
 845 850 855  
 Asp Leu Phe Leu Cys Pro Phe Leu Gly Ser Thr Gly Pro Met Tyr  
 860 865 870  
 Leu Lys Gly Asn Val Tyr Gly Ser Asp Pro Phe Glu Thr Tyr His  
 875 880 885  
 Thr Gly Cys Ser Pro Asp Pro Arg Thr Val Leu Met Asp His Tyr  
 890 895 900  
 Glu Pro Ser Tyr Ile Lys Lys Lys Glu Cys Tyr Pro Cys Ser His  
 905 910 915  
 Pro Ser Glu Glu Ser Cys Glu Arg Ser Phe Ser Asn Ile Ser Trp  
 920 925 930  
 Pro Ser His Val Arg Lys Leu Leu Asn Thr Ser Tyr Ser His Asn  
 935 940 945  
 Glu Gly Pro Gly Met Lys Asn Leu Cys Leu Asn Lys Ser Ser Leu  
 950 955 960  
 Asp Phe Ser Ala Asn Pro Glu Pro Ala Ser Val Ala Ser Ser Asn  
 965 970 975  
 Ser Phe Met Gly Thr Phe Gly Lys Ala Leu Arg Arg Pro His Leu  
 980 985 990  
 Asp Ala Tyr Ser Ser Phe Gly Gln Pro Ser Asp Cys Gln Pro Arg  
 995 1000 1005  
 Ala Phe Tyr Leu Lys Ala His Ser Ser Pro Asp Leu Asp Ser Gly  
 1010 1015 1020  
 Ser Glu Glu Asp Gly Lys Glu Arg Thr Asp Phe Gln Glu Glu Asn

	1025		1030		1035									
His	Ile	Cys	Thr	Phe	Lys	Gln	Thr	Leu	Glu	Asn	Tyr	Arg	Thr	Pro
				1040					1045					1050
Asn	Phe	Gln	Ser	Tyr	Asp	Leu	Asp	Thr						
				1055				1059						

<210> 11  
 <211> 2906  
 <212> DNA  
 <213> artificial sequence

<400> 11  
 ggggagagga attgaccatg taaaaggaga cttttttttt tgggtggtggt 50  
 ggctgttggg tgccttgcaa aaatgaagga tgcaggacgc agctttctcc 100  
 tggaaccgaa cgcaatggat aaactgattg tgcaagagag aaggaagaac 150  
 gaagcttttt cttgtgagcc ctggatctta acacaaatgt gtatatgtgc 200  
 acacagggag cattcaagaa tgaaataaac cagagttaga cccgcggggg 250  
 ttggtgtggt ctgacataaa taaataatct taaagcagct gttcccctcc 300  
 ccacccccaa aaaaaaggat gattggaaat gaagaaccga ggattcacia 350  
 agaaaaaagt atgttcattt ttctctataa aggagaaagt gagccaagga 400  
 gatatttttg gaatgaaaag tttggggctt ttttagtaaa gtaaagaact 450  
 ggtgtggtgg tgttttcctt tctttttgaa tttcccacia gaggagagga 500  
 aattaataat acatctgcaa agaaatttca gagaagaaaa gttgaccgcg 550  
 gcagattgag gcattgattg ggggagagaa accagcagag cacagttgga 600  
 tttgtgccta tgttgactaa aattgacgga taattgcagt tggatttttc 650  
 ttcacaaacc tccttttttt taaattttta ttccttttgg tatcaagatc 700  
 atgcggtttc tcttgttctt aaccacctgg atttccatct ggatgttgct 750  
 gtgatcagtc tgaaatacaa ctgtttgaat tccagaagga ccaacaccag 800  
 ataaattatg aatggtgaac aagatgacct tacatccaca gcagataatg 850  
 ataggtccta ggtttaacag ggcctattt gacccctgc ttgtggtgct 900  
 gctggctctt caacttcttg tgggtggctgg tctggtgcgg gctcagacct 950  
 gcccttctgt gtgctcctgc agcaaccagt tcagcaaggt gatttgtggt 1000  
 cggaaaaacc tgcgtgaggt tccggatggc atctccacca acacacggct 1050  
 gctgaacctc catgagaacc aaatccagat catcaaagtg aacagcttca 1100  
 agcacttgag gcacttgaa atcctacagt tgagtaggaa ccatatcaga 1150  
 accattgaaa ttggggcttt caatggtctg gcgaacctca acactctgga 1200

actctttgac aatcgtctta ctaccatccc gaatggagct tttgtatact 1250  
tgtctaaact gaaggagctc tggttgcaa acaaccccat tgaaagcatc 1300  
ccttcttatg cttttaacag aattccttct ttgcgccgac tagacttagg 1350  
ggaattgaaa agactttcat acatctcaga aggtgccttt gaaggtctgt 1400  
ccaacttgag gtatgtgaac ctgccatgt gcaaccttcg ggaaatccct 1450  
aacctcacac cgctcataaa actagatgag ctggatcttt ctgggaatca 1500  
tttatctgcc atcaggcctg gctctttcca gggtttgatg caccttcaaa 1550  
aactgtggat gatacagtcc cagattcaag tgattgaacg gaatgccttt 1600  
gacaaccttc agtcactagt ggagatcaac ctggcacaca ataatctaac 1650  
attactgcct catgacctct tcactccctt gcatcatcta gagcggatac 1700  
attacatca caacccttgg aactgtaact gtgacatact gtggctcagc 1750  
tggtgataa aagacatggc ccctcgaac acagcttggt gtgcccggtg 1800  
taacactcct cccaatctaa aggggaggtta cattggagag ctcgaccaga 1850  
attacttcac atgctatgct ccggtgattg tggagcccc tgcagacctc 1900  
aatgtcactg aaggcatggc agctgagctg aaatgtcggg cctccacatc 1950  
cctgacatct gtatcttggg ttactccaaa tggaacagtc atgacacatg 2000  
ggcgctaaa agtgccgata gctgtgctca gtgatggtac gttaaatttc 2050  
acaaatgtaa ctgtgcaaga tacaggcatg tacacatgta tggtgagtaa 2100  
ttccgttggg aatactactg cttcagccac cctgaatggt actgcagcaa 2150  
ccactactcc tttctcttac ttttcaaccg tcacagtaga gactatggaa 2200  
ccgtctcagg atgaggcacg gaccacagat aacaatgtgg gtccactcc 2250  
agtggtcgac tgggagacca ccaatgtgac cacctctctc acaccacaga 2300  
gcacaaggtc gacagagaaa accttcacca tcccagtgac tgatataaac 2350  
agtgggatcc caggaattga tgaggatcag aagactacca aatcatcat 2400  
tgggtgtttt gtggccatca cactcatggc tgcagtgatg ctggtcattt 2450  
tctacaagat gaggaagcag caccatcggc aaaacatca cgccccaaaca 2500  
aggactgttg aaattattaa tgtggatgat gagattacgg gagacacacc 2550  
catggaaagc cacctgcccc tgcoctgctat cgagcatgag cacctaaatc 2600  
actataactc atacaaatct cccttcaacc acacaacaac agttaacaca 2650  
ataaattcaa tacacagttc agtgcgatgaa ccgttattga tccgaatgaa 2700

ctctaaagac aatgtacaag agactcaaatt ctaaaacatt tacagagtta 2750  
 caaaaaacaa acaatcaaaa aaaaagacag tttattaata atgacacaaa 2800  
 tgactgggct aaatctactg tttcaaaaaa gtgtctttac aaaaaaacia 2850  
 aaaagaaaag aaatattatt attaaaaatt ctattgtgat ctaaagcaga 2900  
 caaaaa 2906

<210> 12  
 <211> 640  
 <212> PRT  
 <213> artificial sequence

<400> 12  
 Met Leu Asn Lys Met Thr Leu His Pro Gln Gln Ile Met Ile Gly  
 1 5 10 15  
 Pro Arg Phe Asn Arg Ala Leu Phe Asp Pro Leu Leu Val Val Leu  
 20 25 30  
 Leu Ala Leu Gln Leu Leu Val Val Ala Gly Leu Val Arg Ala Gln  
 35 40 45  
 Thr Cys Pro Ser Val Cys Ser Cys Ser Asn Gln Phe Ser Lys Val  
 50 55 60  
 Ile Cys Val Arg Lys Asn Leu Arg Glu Val Pro Asp Gly Ile Ser  
 65 70 75  
 Thr Asn Thr Arg Leu Leu Asn Leu His Glu Asn Gln Ile Gln Ile  
 80 85 90  
 Ile Lys Val Asn Ser Phe Lys His Leu Arg His Leu Glu Ile Leu  
 95 100 105  
 Gln Leu Ser Arg Asn His Ile Arg Thr Ile Glu Ile Gly Ala Phe  
 110 115 120  
 Asn Gly Leu Ala Asn Leu Asn Thr Leu Glu Leu Phe Asp Asn Arg  
 125 130 135  
 Leu Thr Thr Ile Pro Asn Gly Ala Phe Val Tyr Leu Ser Lys Leu  
 140 145 150  
 Lys Glu Leu Trp Leu Arg Asn Asn Pro Ile Glu Ser Ile Pro Ser  
 155 160 165  
 Tyr Ala Phe Asn Arg Ile Pro Ser Leu Arg Arg Leu Asp Leu Gly  
 170 175 180  
 Glu Leu Lys Arg Leu Ser Tyr Ile Ser Glu Gly Ala Phe Glu Gly  
 185 190 195  
 Leu Ser Asn Leu Arg Tyr Leu Asn Leu Ala Met Cys Asn Leu Arg  
 200 205 210  
 Glu Ile Pro Asn Leu Thr Pro Leu Ile Lys Leu Asp Glu Leu Asp  
 215 220 225

Leu Ser Gly Asn His Leu Ser Ala Ile Arg Pro Gly Ser Phe Gln  
 230 235 240  
 Gly Leu Met His Leu Gln Lys Leu Trp Met Ile Gln Ser Gln Ile  
 245 250 255  
 Gln Val Ile Glu Arg Asn Ala Phe Asp Asn Leu Gln Ser Leu Val  
 260 265 270  
 Glu Ile Asn Leu Ala His Asn Asn Leu Thr Leu Leu Pro His Asp  
 275 280 285  
 Leu Phe Thr Pro Leu His His Leu Glu Arg Ile His Leu His His  
 290 295 300  
 Asn Pro Trp Asn Cys Asn Cys Asp Ile Leu Trp Leu Ser Trp Trp  
 305 310 315  
 Ile Lys Asp Met Ala Pro Ser Asn Thr Ala Cys Cys Ala Arg Cys  
 320 325 330  
 Asn Thr Pro Pro Asn Leu Lys Gly Arg Tyr Ile Gly Glu Leu Asp  
 335 340 345  
 Gln Asn Tyr Phe Thr Cys Tyr Ala Pro Val Ile Val Glu Pro Pro  
 350 355 360  
 Ala Asp Leu Asn Val Thr Glu Gly Met Ala Ala Glu Leu Lys Cys  
 365 370 375  
 Arg Ala Ser Thr Ser Leu Thr Ser Val Ser Trp Ile Thr Pro Asn  
 380 385 390  
 Gly Thr Val Met Thr His Gly Ala Tyr Lys Val Arg Ile Ala Val  
 395 400 405  
 Leu Ser Asp Gly Thr Leu Asn Phe Thr Asn Val Thr Val Gln Asp  
 410 415 420  
 Thr Gly Met Tyr Thr Cys Met Val Ser Asn Ser Val Gly Asn Thr  
 425 430 435  
 Thr Ala Ser Ala Thr Leu Asn Val Thr Ala Ala Thr Thr Thr Pro  
 440 445 450  
 Phe Ser Tyr Phe Ser Thr Val Thr Val Glu Thr Met Glu Pro Ser  
 455 460 465  
 Gln Asp Glu Ala Arg Thr Thr Asp Asn Asn Val Gly Pro Thr Pro  
 470 475 480  
 Val Val Asp Trp Glu Thr Thr Asn Val Thr Thr Ser Leu Thr Pro  
 485 490 495  
 Gln Ser Thr Arg Ser Thr Glu Lys Thr Phe Thr Ile Pro Val Thr  
 500 505 510  
 Asp Ile Asn Ser Gly Ile Pro Gly Ile Asp Glu Val Met Lys Thr  
 515 520 525  
 Thr Lys Ile Ile Ile Gly Cys Phe Val Ala Ile Thr Leu Met Ala

	530		535		540
Ala Val Met Leu	Val Ile Phe Tyr Lys	Met Arg Lys Gln His His			
	545		550		555
Arg Gln Asn His	His Ala Pro Thr Arg	Thr Val Glu Ile Ile Asn			
	560		565		570
Val Asp Asp Glu	Ile Thr Gly Asp Thr	Pro Met Glu Ser His Leu			
	575		580		585
Pro Met Pro Ala	Ile Glu His Glu His	Leu Asn His Tyr Asn Ser			
	590		595		600
Tyr Lys Ser Pro	Phe Asn His Thr Thr	Thr Val Asn Thr Ile Asn			
	605		610		615
Ser Ile His Ser	Ser Val His Glu Pro	Leu Leu Ile Arg Met Asn			
	620		625		630
Ser Lys Asp Asn	Val Gln Glu Thr Gln Ile				
	635		640		

<210> 13

<211> 4051

<212> DNA

<213> artificial sequence

<400> 13

```

agccgacgct gctcaagctg caactctggt gcagttggca gttcttttcg 50
gtttccctcc tgctgtttgg gggcatgaaa gggcttcgcc gccgggagta 100
aaagaaggaa ttgaccgggc agcgcgaggg aggagcgcgc acgcgaccgc 150
gagggcgggc gtgcaccctc ggctggaagt ttgtgccggg ccccgagcgc 200
gcgcccggctg ggagcttcgg gtagagacct aggccgctgg accgcgatga 250
gcgcgcccag cctccgtgcg cgcgcccgcg ggttggggct gctgctgtgc 300
gcggtgctgg ggcgcgctgg ccggtccgac agcggcggtc gcggggaact 350
cgggcagccc tctggggtag ccgccgagcg cccatgcccc actacctgcc 400
gctgcctcgg ggacctgctg gactgcagtc gtaagcggct agcgcgtctt 450
cccgagccac tcccgtcctg ggtcgcctcg ctggacttaa gtcacaacag 500
attatctttc atcaaggcaa gttccatgag ccaccttcaa agccttcgag 550
aagtgaaact gaacaacaat gaattggaga ccattccaaa tctgggacca 600
gtctcgcaa atattacact tctctccttg gctggaacaa ggattgttga 650
aatactccct gaacatctga aagagtttca gtccttgaa actttggacc 700
ttagcagcaa caatatttca gagctccaaa ctgcatttcc agccctacag 750
ctcaaatatc tgtatctcaa cagcaaccga gtcacatcaa tggaacctgg 800

```

gtatTTTgac aattTggcca acacactcct tgtgtTaaag ctgaacagga 850  
accgaatctc agctatccca cccaagatgt ttaaactgcc ccaactgcaa 900  
catctcgaat tgaaccgaaa caagattaaa aatgtagatg gactgacatt 950  
ccaaggcctt ggtgctctga agtctctgaa aatgcaaaga aatggagtaa 1000  
cgaaacttat ggatggagct ttttgggggc tgagcaacat ggaaatTTTg 1050  
cagctggacc ataacaacct aacagagatt accaaaggct ggctTTTcgg 1100  
cttgctgatg ctgcaggaac ttcactctcag caaaatgcc atcaacagga 1150  
tcagccctga tgccctgggag ttctgccaga agctcagtga gctggacct 1200  
actTTcaatc acttatcaag gttagatgat tcaagcttcc ttggcctaag 1250  
cttactaaat aactgcaca ttgggaacaa cagagtcagc tacattgctg 1300  
attgtgcctt ccgggggctt tccagTTTaa agactTTgga tctgaagaac 1350  
aatgaaattt cctggactat tgaagacatg aatggTgctt tctctgggct 1400  
tgacaaaactg aggcgactga tactccaagg aaatcggatc cgTTctatta 1450  
ctaaaaaagc cttcactggt ttggatgcat tggagcatct agacctgagt 1500  
acaacgcaat catgtctTTa caaggcaatg cattTTcaca aatgaagaaa 1550  
ctgcaacaat tgcattTaaa tacatcaagc cTTTtTgtgcg attgccagct 1600  
aaaatggctc ccacagtggg tggcggaaaa caactTTcag agctTTgtaa 1650  
atgccagttg tgccatcctc cagctgctaa aaggagaag cattTTTgct 1700  
gttagcccag atggctTTgt gtgtgatgat tttcccaaac cccagatcac 1750  
ggttcagcca gaaacacagt cggcaataaa aggttccaat ttgagTTTca 1800  
tctgctcagc tgccagcagc agtgattccc caatgacttt tgctTgaaa 1850  
aaagacaatg aactactgca tgatgctgaa atggaaaatt atgcacacct 1900  
ccgggccccaa ggtggcgagg tgatggagta taccaccatc cttcggctgc 1950  
gcgaggtgga atttgccagt gaggggaaat atcagtgtgt catctccaat 2000  
cactTTggtt catcctactc tgtcaaagcc aagctTtacag taaatatgct 2050  
tcctcattc accaagacct ccatggatct caccatccga gctggggcca 2100  
tggcacgctt ggagtgtgct gctgtggggc acccagcccc ccagatagcc 2150  
tggcagaagg atgggggcac agactTcca gctgcacggg agagacgcat 2200  
gcatgtgatg cccgaggatg acgtgttctt tatcgtggat gtgaagatag 2250  
aggacattgg ggtatacagc tgcacagctc agaacagtgc aggaagtatt 2300  
tcagcaaatg caactctgac tgtcctagaa acaccatcat tttTcggcc 2350

actggttgac cgaactgtaa ccaagggaga aacagccgtc ctacagtgca 2400  
ttgctggagg aagccctccc cctaaactga actggaccaa agatgatagc 2450  
ccattggtgg taaccgagag gcactttttt gcagcaggca atcagcttct 2500  
gattattgtg gactcagatg tcagtgatgc tgggaaatac acatgtgaga 2550  
tgtctaacac ccttggcact gagagaggaa acgtgcgctt cagtgtgatc 2600  
cccactccaa cctgcgactc ccctcagatg acagcccat cgtagacga 2650  
tgacggatgg gccactgtgg gtgtcgtgat catagccgtg gtttgctgtg 2700  
tgggtggcac gtcactcgtg tgggtggtca tcatatacca cacaaggcgg 2750  
aggaatgaag attgcagcat taccaacaca gatgagacca acttgccagc 2800  
agatattcct agttatttgt catctcaggg aacgttagct gacaggcagg 2850  
atgggtacgt gtcttcagaa agtgggaagcc accaccagtt tgtcacatct 2900  
tcaggtgctg gatttttctt accacaacat gacagtagtg ggacctgcca 2950  
tattgacaat agcagtgaag ctgatgtgga agctgccaca gatctgttcc 3000  
ttgtccgttt ttgggatcca caggccctat gtatttgaag ggaaatgtgt 3050  
atggctcaga tccttttgaa acatatcata caggttgcag tcctgacca 3100  
agaacagttt taatggacca ctatgagccc agttacataa agaaaaagga 3150  
gtgctacca tgttctcatc cttcagaaga atcctgcgaa cggagcttca 3200  
gtaatatac gtggccttca catgtgagga agctacttaa cactagttac 3250  
tctcacaatg aaggacctgg aatgaaaaat ctgtgtctaa acaagtcctc 3300  
tttagatttt agtgcaaac cagagccagc gtcggttgcc tcgagtaatt 3350  
ctttcatggg tacctttgga aaagctctca ggagacctca cctagatgcc 3400  
tattcaagct ttggacagcc atcagattgt cagccaagag ccttttattt 3450  
gaaagctcat tcttccccag acttggactc tgggtcagag gaagatggga 3500  
aagaaaggac agattttcag gaagaaaatc acatttgtac ctttaaacag 3550  
actttagaaa actacaggac tccaaatttt cagtcttatg acttggacac 3600  
atagactgaa tgagaccaa ggaaaagctt aacatactac ctcaagtga 3650  
cttttattta aaagagagag aatcttatgt tttttaaag gagttatgaa 3700  
ttttaaaggg ataaaaatgc tttatttata cagatgaacc aaaattaca 3750  
aaagttatga aaatttttat actgggaatg atgctcatat aagaatacct 3800  
ttttaaacta ttttttaact ttgttttatg caaaaaagta tcttacgtaa 3850

attaatgata taaatcatga ttattttatg tatttttata atgccagatt 3900  
 tctttttatg gaaaatgagt tactaaagca ttttaaataa tacctgcctt 3950  
 gtaccatttt ttaaatagaa gttacttcat tatattttgc acattatatt 4000  
 taataaaatg tgtcaatttg aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 4050

a 4051

<210> 14  
 <211> 1119  
 <212> PRT  
 <213> artificial sequence

<400> 14  
 Met Ser Ala Pro Ser Leu Arg Ala Arg Ala Ala Gly Leu Gly Leu  
 1 5 10 15  
 Leu Leu Cys Ala Val Leu Gly Arg Ala Gly Arg Ser Asp Ser Gly  
 20 25 30  
 Gly Arg Gly Glu Leu Gly Gln Pro Ser Gly Val Ala Ala Glu Arg  
 35 40 45  
 Pro Cys Pro Thr Thr Cys Arg Cys Leu Gly Asp Leu Leu Asp Cys  
 50 55 60  
 Ser Arg Lys Arg Leu Ala Arg Leu Pro Glu Pro Leu Pro Ser Trp  
 65 70 75  
 Val Ala Arg Leu Asp Leu Ser His Asn Arg Leu Ser Phe Ile Lys  
 80 85 90  
 Ala Ser Ser Met Ser His Leu Gln Ser Leu Arg Glu Val Lys Leu  
 95 100 105  
 Asn Asn Asn Glu Leu Glu Thr Ile Pro Asn Leu Gly Pro Val Ser  
 110 115 120  
 Ala Asn Ile Thr Leu Leu Ser Leu Ala Gly Asn Arg Ile Val Glu  
 125 130 135  
 Ile Leu Pro Glu His Leu Lys Glu Phe Gln Ser Leu Glu Thr Leu  
 140 145 150  
 Asp Leu Ser Ser Asn Asn Ile Ser Glu Leu Gln Thr Ala Phe Pro  
 155 160 165  
 Ala Leu Gln Leu Lys Tyr Leu Tyr Leu Asn Ser Asn Arg Val Thr  
 170 175 180  
 Ser Met Glu Pro Gly Tyr Phe Asp Asn Leu Ala Asn Thr Leu Leu  
 185 190 195  
 Val Leu Lys Leu Asn Arg Asn Arg Ile Ser Ala Ile Pro Pro Lys  
 200 205 210  
 Met Phe Lys Leu Pro Gln Leu Gln His Leu Glu Leu Asn Arg Asn  
 215 220 225

Lys Ile Lys Asn Val Asp Gly Leu Thr Phe Gln Gly Leu Gly Ala  
 230 235 240  
 Leu Lys Ser Leu Lys Met Gln Arg Asn Gly Val Thr Lys Leu Met  
 245 250 255  
 Asp Gly Ala Phe Trp Gly Leu Ser Asn Met Glu Ile Leu Gln Leu  
 260 265 270  
 Asp His Asn Asn Leu Thr Glu Ile Thr Lys Gly Trp Leu Tyr Gly  
 275 280 285  
 Leu Leu Met Leu Gln Glu Leu His Leu Ser Gln Asn Ala Ile Asn  
 290 295 300  
 Arg Ile Ser Pro Asp Ala Trp Glu Phe Cys Gln Lys Leu Ser Glu  
 305 310 315  
 Leu Asp Leu Thr Phe Asn His Leu Ser Arg Leu Asp Asp Ser Ser  
 320 325 330  
 Phe Leu Gly Leu Ser Leu Leu Asn Thr Leu His Ile Gly Asn Asn  
 335 340 345  
 Arg Val Ser Tyr Ile Ala Asp Cys Ala Phe Arg Gly Leu Ser Ser  
 350 355 360  
 Leu Lys Thr Leu Asp Leu Lys Asn Asn Glu Ile Ser Trp Thr Ile  
 365 370 375  
 Glu Asp Met Asn Gly Ala Phe Ser Gly Leu Asp Lys Leu Arg Arg  
 380 385 390  
 Leu Ile Leu Gln Gly Asn Arg Ile Arg Ser Ile Thr Lys Lys Ala  
 395 400 405  
 Phe Thr Gly Leu Asp Ala Leu Glu His Leu Asp Leu Ser Asp Asn  
 410 415 420  
 Ala Ile Met Ser Leu Gln Gly Asn Ala Phe Ser Gln Met Lys Lys  
 425 430 435  
 Leu Gln Gln Leu His Leu Asn Thr Ser Ser Leu Leu Cys Asp Cys  
 440 445 450  
 Gln Leu Lys Trp Leu Pro Gln Trp Val Ala Glu Asn Asn Phe Gln  
 455 460 465  
 Ser Phe Val Asn Ala Ser Cys Ala His Pro Gln Leu Leu Lys Gly  
 470 475 480  
 Arg Ser Ile Phe Ala Val Ser Pro Asp Gly Phe Val Cys Asp Asp  
 485 490 495  
 Phe Pro Lys Pro Gln Ile Thr Val Gln Pro Glu Thr Gln Ser Ala  
 500 505 510  
 Ile Lys Gly Ser Asn Leu Ser Phe Ile Cys Ser Ala Ala Ser Ser  
 515 520 525  
 Ser Asp Ser Pro Met Thr Phe Ala Trp Lys Lys Asp Asn Glu Leu

	530		535		540
Leu His Asp Ala	Glu Met Glu Asn Tyr	Ala His Leu Arg Ala	Gln		
	545		550		555
Gly Gly Glu Val	Met Glu Tyr Thr Thr	Ile Leu Arg Leu Arg	Glu		
	560		565		570
Val Glu Phe Ala	Ser Glu Gly Lys Tyr	Gln Cys Val Ile Ser	Asn		
	575		580		585
His Phe Gly Ser	Ser Tyr Ser Val Lys	Ala Lys Leu Thr Val	Asn		
	590		595		600
Met Leu Pro Ser	Phe Thr Lys Thr Pro	Met Asp Leu Thr Ile	Arg		
	605		610		615
Ala Gly Ala Met	Ala Arg Leu Glu Cys	Ala Ala Val Gly His	Pro		
	620		625		630
Ala Pro Gln Ile	Ala Trp Gln Lys Asp	Gly Gly Thr Asp Phe	Pro		
	635		640		645
Ala Ala Arg Glu	Arg Arg Met His Val	Met Pro Glu Asp Asp	Val		
	650		655		660
Phe Phe Ile Val	Asp Val Lys Ile Glu	Asp Ile Gly Val Tyr	Ser		
	665		670		675
Cys Thr Ala Gln	Asn Ser Ala Gly Ser	Ile Ser Ala Asn Ala	Thr		
	680		685		690
Leu Thr Val Leu	Glu Thr Pro Ser Phe	Leu Arg Pro Leu Leu	Asp		
	695		700		705
Arg Thr Val Thr	Lys Gly Glu Thr Ala	Val Leu Gln Cys Ile	Ala		
	710		715		720
Gly Gly Ser Pro	Pro Pro Lys Leu Asn	Trp Thr Lys Asp Asp	Ser		
	725		730		735
Pro Leu Val Val	Thr Glu Arg His Phe	Phe Ala Ala Gly Asn	Gln		
	740		745		750
Leu Leu Ile Ile	Val Asp Ser Asp Val	Ser Asp Ala Gly Lys	Tyr		
	755		760		765
Thr Cys Glu Met	Ser Asn Thr Leu Gly	Thr Glu Arg Gly Asn	Val		
	770		775		780
Arg Leu Ser Val	Ile Pro Thr Pro Thr	Cys Asp Ser Pro Gln	Met		
	785		790		795
Thr Ala Pro Ser	Leu Asp Asp Asp Gly	Trp Ala Thr Val Gly	Val		
	800		805		810
Val Ile Ile Ala	Val Val Cys Cys Val	Val Gly Thr Ser Leu	Val		
	815		820		825
Trp Val Val Ile	Ile Tyr His Thr Arg	Arg Arg Asn Glu Asp	Cys		
	830		835		840

Ser Ile Thr Asn Thr Asp Glu Thr Asn Leu Pro Ala Asp Ile Pro  
 845 850 855  
 Ser Tyr Leu Ser Ser Gln Gly Thr Leu Ala Asp Arg Gln Asp Gly  
 860 865 870  
 Tyr Val Ser Ser Glu Ser Gly Ser His His Gln Phe Val Thr Ser  
 875 880 885  
 Ser Gly Ala Gly Phe Phe Leu Pro Gln His Asp Ser Ser Gly Thr  
 890 895 900  
 Cys His Ile Asp Asn Ser Ser Glu Ala Asp Val Glu Ala Ala Thr  
 905 910 915  
 Asp Leu Phe Leu Cys Pro Phe Leu Gly Ser Thr Gly Pro Met Tyr  
 920 925 930  
 Leu Lys Gly Asn Val Tyr Gly Ser Asp Pro Phe Glu Thr Tyr His  
 935 940 945  
 Thr Gly Cys Ser Pro Asp Pro Arg Thr Val Leu Met Asp His Tyr  
 950 955 960  
 Glu Pro Ser Tyr Ile Lys Lys Lys Glu Cys Tyr Pro Cys Ser His  
 965 970 975  
 Pro Ser Glu Glu Ser Cys Glu Arg Ser Phe Ser Asn Ile Ser Trp  
 980 985 990  
 Pro Ser His Val Arg Lys Leu Leu Asn Thr Ser Tyr Ser His Asn  
 995 1000 1005  
 Glu Gly Pro Gly Met Lys Asn Leu Cys Leu Asn Lys Ser Ser Leu  
 1010 1015 1020  
 Asp Phe Ser Ala Asn Pro Glu Pro Ala Ser Val Ala Ser Ser Asn  
 1025 1030 1035  
 Ser Phe Met Gly Thr Phe Gly Lys Ala Leu Arg Arg Pro His Leu  
 1040 1045 1050  
 Asp Ala Tyr Ser Ser Phe Gly Gln Pro Ser Asp Cys Gln Pro Arg  
 1055 1060 1065  
 Ala Phe Tyr Leu Lys Ala His Ser Ser Pro Asp Leu Asp Ser Gly  
 1070 1075 1080  
 Ser Glu Glu Asp Gly Lys Glu Arg Thr Asp Phe Gln Glu Glu Asn  
 1085 1090 1095  
 His Ile Cys Thr Phe Lys Gln Thr Leu Glu Asn Tyr Arg Thr Pro  
 1100 1105 1110  
 Asn Phe Gln Ser Tyr Asp Leu Asp Thr  
 1115 1119

<210> 15  
 <211> 22  
 <212> DNA

<213> artificial sequence

<400> 15

atcgttgatga agttagtgcc cc 22

<210> 16

<211> 23

<212> DNA

<213> artificial sequence

<400> 16

acctgcgata tccaacagaa ttg 23

<210> 17

<211> 48

<212> DNA

<213> artificial sequence

<400> 17

ggaagaggat acagtcactc tggaagtatt agtggctcca gcagttcc 48

<210> 18

<211> 46

<212> DNA

<213> artificial sequence

<400> 18

gggtacacct gtcctgcac cgacggatat tggcttctgg aaggcc 46

<210> 19

<211> 22

<212> DNA

<213> artificial sequence

<400> 19

acagattccc accagtgcaa cc 22

<210> 20

<211> 21

<212> DNA

<213> artificial sequence

<400> 20

cacactcgtt cacatcttgg c 21

<210> 21

<211> 45

<212> DNA

<213> artificial sequence

<400> 21

agggagcacg gacagtgtgc agatgtggac gaggctcac tagca 45

<210> 22

<211> 21

<212> DNA

<213> artificial sequence

<400> 22

agagtgtatc tctggctacg c 21

<210> 23  
<211> 22  
<212> DNA  
<213> artificial sequence

<400> 23  
taagtccggc acattacagg tc 22

<210> 24  
<211> 49  
<212> DNA  
<213> artificial sequence

<400> 24  
cccacgatgt atgaatggtg gactttgtgt gactcctggt ttctgcatc 49

<210> 25  
<211> 22  
<212> DNA  
<213> artificial sequence

<400> 25  
aaagacgcat ctgcgagtgt cc 22

<210> 26  
<211> 23  
<212> DNA  
<213> artificial sequence

<400> 26  
tgctgatttc aactgctct ccc 23

<210> 27  
<211> 24  
<212> DNA  
<213> artificial sequence

<400> 27  
tcgcgagct gtgttctggt tccc 24

<210> 28  
<211> 50  
<212> DNA  
<213> artificial sequence

<400> 28  
tgatcgcgat ggggacaaag gcgcaagctc gagaggaaac tgttgtgcct 50

<210> 29  
<211> 20  
<212> DNA  
<213> artificial sequence

<400> 29  
acacctgggt caaagatggg 20

<210> 30  
<211> 24  
<212> DNA  
<213> artificial sequence

<400> 30  
taggaagagt tgctgaaggc acgg 24

<210> 31  
<211> 20  
<212> DNA  
<213> artificial sequence

<400> 31  
ttgccttact caggtgctac 20

<210> 32  
<211> 20  
<212> DNA  
<213> artificial sequence

<400> 32  
actcagcagt ggtaggaaag 20

<210> 33  
<211> 22  
<212> DNA  
<213> artificial sequence

<400> 33  
gttggatctg ggcaacaata ac 22

<210> 34  
<211> 24  
<212> DNA  
<213> artificial sequence

<400> 34  
attggtgtgc aggctgagtt taag 24

<210> 35  
<211> 45  
<212> DNA  
<213> artificial sequence

<400> 35  
ggtggctata catggatagc aattacctgg acacgctgtc ccggg 45

<210> 36  
<211> 50  
<212> DNA  
<213> artificial sequence

<400> 36  
ggattctaatac gactcact atagggctgc ggcggctcag gtcttcagtt 50

<210> 37  
<211> 50  
<212> DNA  
<213> artificial sequence

<400> 37  
ctatgaaatt aaccctcact aaaggagca tgggatgggg agggatacgg 50

<210> 38  
<211> 48

<212> DNA  
<213> artificial sequence

<400> 38  
ctatgaaatt aaccctcact aaaggaata gcaggccatc ccaggaca 48

<210> 39  
<211> 47  
<212> DNA  
<213> artificial sequence

<400> 39  
ctatgaaatt aaccctcact aaagggatgt cttccatgcc aaccttc 47

1

1