



HU000029457T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 029 457**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 11 839071**(22) A bejelentés napja: **2011. 11. 11.**

(96) Az európai bejelentés bejelentési száma:

EP 20110839071

(97) Az európai bejelentés közzétételi adatai:

EP 2637690 A2 **2012. 05. 18.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2637690 B1 **2016. 09. 21.**(51) Int. Cl.: **A61K 393/95** (2006.01)

(86) A nemzetközi (PCT) bejelentési szám:

PCT/US 11/060388

(87) A nemzetközi közzétételi szám:

WO 12065072

(30) Elsőbbségi adatok:

413960 P **2010. 11. 15.** **US****412728 P** **2010. 11. 11.** **US**

(72) Feltalálók(k):

NEU, Michael, 68535 Edingen-Neckarhausen (DE)**TSCHOEPE, Markus, 67454 Hassloch (DE)****WEBER, Carsten, 67133 Maxdorf (DE)****REDDEN, Laura, Glenview, IL 60025 (US)****FRAUNHOFER, Wolfgang, Gurnee, IL 60031 (US)****GASTENS, Martin, 67157 Wachenheim (DE)****FEICK, Alexander, 4123 Allschwil (CH)****PAULSON, Susan K., Downers Grove, IL 60515 (US)****ZHU, Tong, Riverwoods, IL 60015 (US)**

(73) Jogosult(ak):

AbbVie Biotechnology Ltd, HM 11 Hamilton
(BM)

(74) Képviselő:

Danubia Szabadalmi és Jogi Iroda Kft.,
Budapest(54) **Folyékony készítmények, amelyek nagy koncentrációban tartalmaznak ANTI-TNF-alfa antitestet**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 637 690 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
21.09.2016 Bulletin 2016/38

(21) Application number: **11839071.5**

(22) Date of filing: **11.11.2011**

(51) Int Cl.:
A61K 39/395 (2006.01)

(86) International application number:
PCT/US2011/060388

(87) International publication number:
WO 2012/065072 (18.05.2012 Gazette 2012/20)

(54) **HIGH CONCENTRATION ANTI-TNFalpha ANTIBODY LIQUID FORMULATIONS**

ANTI-TNF-ALPHA-ANTIKÖRPER-FLÜSSIGFORMULIERUNGEN MIT HOHER KONZENTRATION
FORMULATION LIQUIDE D'ANTICORPS ANTI-TNFalpha HAUTEMENT CONCENTRÉE

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**
Designated Extension States:
BA ME

(30) Priority: **15.11.2010 US 413960 P**
11.11.2010 US 412728 P

(43) Date of publication of application:
18.09.2013 Bulletin 2013/38

(73) Proprietor: **AbbVie Biotechnology Ltd**
HM 11 Hamilton (BM)

(72) Inventors:

- **NEU, Michael**
68535 Edingen-Neckarhausen (DE)
- **TSCHOEPE, Markus**
67454 Hassloch (DE)
- **WEBER, Carsten**
67133 Maxdorf (DE)
- **REDDEN, Laura**
Glenview, IL 60025 (US)
- **FRAUNHOFER, Wolfgang**
Gurnee, IL 60031 (US)
- **GASTENS, Martin**
67157 Wachenheim (DE)
- **FEICK, Alexander**
4123 Allschwil (CH)
- **PAULSON, Susan K.**
Downers Grove, IL 60515 (US)
- **ZHU, Tong**
Riverwoods, IL 60015 (US)

(74) Representative: **Adams, Harvey Vaughan John et al**
Mathys & Squire LLP
The Shard
32 London Bridge Street
London SE1 9SG (GB)

(56) References cited:
WO-A2-2004/016286 WO-A2-2009/073569
US-A1- 2004 038 878 US-A1- 2010 278 822
US-A1- 2010 278 822

- Emea: "Assessment Report for Simponi EMEA/H/C/000992", , 3 March 2008 (2008-03-03), pages 1-70, XP055109044, UK Retrieved from the Internet:
URL:http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Public_assessment_report/human/000992/WC500052372.pdf [retrieved on 2014-03-20]
- ABBOTT BIOTECHNOLOGY LTD: "Additional Experimental Results for EP 03 748 438", ABBOTT BIOTECHNOLOGY LTD, 1 May 2009 (2009-05-01), pages 1-26, XP055057827,
- Abbott: "HUMIRA", US, 31 January 2003 (2003-01-31), pages 1-13, XP055109061, USA Retrieved from the Internet:
URL:http://www.fda.gov/ohrms/dockets/ac/03/briefing/3930B1_02_B-Abbott-Humira_Prescribing_Info.pdf [retrieved on 2014-03-20]

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 637 690 B1

- DAUGHERTY A L ET AL: "Formulation and delivery issues for monoclonal antibody therapeutics", ADVANCED DRUG DELIVERY REVIEWS, ELSEVIER, AMSTERDAM, NL, vol. 58, no. 5-6, 7 August 2006 (2006-08-07), pages 686-706, XP024892149, ISSN: 0169-409X, DOI: 10.1016/J.ADDR.2006.03.011 [retrieved on 2006-08-07]

Description

BACKGROUND OF THE INVENTION

5 [0001] The formulation of therapeutic proteins, such as antibodies, is often a challenge given the numerous desirable properties that the formulation must have to be economically and therapeutically successful, *e.g.*, stability, suitability for administration, concentration. During manufacturing, storage, and delivery, therapeutic proteins have been known to undergo physical and chemical degradations. These instabilities can reduce the potency of the protein and increase the risk of adverse events in patients, and, therefore, significantly impact regulatory approval (see, *e.g.*, Wang et al. J. Pharm. Sci. 96:1, 2007). As such, a stable protein formulation is essential to the success of a therapeutic protein.

10 [0002] To be effective, many therapeutic proteins require the administration of high doses, which, ideally, are formulated in high concentration formulations. High protein concentration formulations are desirable as they can impact the mode (*e.g.*, intravenous vs. subcutaneous) and frequency of administration of the drug to a subject.

15 [0003] Despite the benefits of high protein concentration formulations, formulating high concentration therapeutic proteins presents numerous challenges. For example, increasing protein concentration often negatively impacts protein aggregation, solubility, stability, and viscosity (see, *e.g.*, Shire et al. J. Pharm. Sci. 93:1390,2004). Increased viscosity, which is a very common challenge for high protein solutions, can have negative ramifications on administration of the formulation, *e.g.*, felt pain and burning syndromes and limitations in manufacturing, processing, fill-finish and drug delivery device options (see, *e.g.*, Shire et al. J. Pharm. Sci. 93:1390,2004). Even for therapeutic proteins having common structural features, *e.g.*, antibodies, approved formulations to date have had varying ingredients and ranges of concentrations. For example, the anti-CD20 antibody Rituxan is formulated for intravenous administration at a concentration of 10 mg/mL, while the anti-RSV antibody Synagis is formulated for intramuscular administration at a concentration of 100 mg/mL. Thus, high protein formulations, especially antibody formulations, which can be used for therapeutic purposes remain a challenge.

25 [0004] Another challenge associated with therapeutic proteins, such as antibodies, is drug delivery. While self-administering devices allow patients to avoid unnecessary trips to medical facilities to receive treatments, patients' self-awareness and fear of the pain associated with self-administration may frequently impact self-administered drug delivery. Moreover, formulations having high concentrations of protein may have high viscosity resulting in increased pain upon delivery, particularly for subcutaneous administration. Thus, there is especially a need for high concentration formulations that reduce pain associated with drug delivery (*e.g.*, self-injection).

30 [0005] Accordingly, there is a need for stable, high concentration protein formulations that provide dosing and administrative advantages, particularly with respect to a decrease in pain for the patient and/or improved bioavailability.

SUMMARY OF THE INVENTION

35 [0006] The present invention is based, at least in part, on the discovery of new high-concentration formulations for therapeutic antibodies (including human anti-TNF- α antibodies, or antigen-binding fragments thereof, *e.g.*, adalimumab). The formulations of the invention provide a number of surprising characteristics given the high concentration of the therapeutic antibody. Specifically, the present invention provides pharmaceutical formulations comprising human anti-TNF α antibodies which surprisingly have improved bioavailability or decreased pain upon subcutaneous injection.

40 [0007] In particular, the present invention is based, at least partly, on the unexpected and surprising discovery that a formulation having a high antibody concentration, a surfactant, and a polyol, provides dramatically reduced pain to the patient during drug delivery, particularly subcutaneous administration of the antibody through, for example, self-injection. The formulations of the invention are established, at least in part, on the surprising finding that a therapeutic protein (*e.g.*, an anti-TNF- α antibody, or antigen-binding portion thereof), can remain soluble at a high protein concentration (*e.g.*, at least about 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 96, 100, 105, 110 mg/mL or more) and maintain a viscosity suitable for injection (*e.g.*, subcutaneous administration). The formulation of the present invention is further surprising, in that the formulation does not contain a buffer or a salt, yet has a high concentration of antibody. Notably, the formulation of the invention reduces pain associated with injection in a patient by at least about 50% (*e.g.*, at least about 60%, 65%, 70%, 75%, 80%, 85%, 90% or more) when compared to injecting an otherwise identical formulation comprising at least one salt and/or at least one buffer.

50 [0008] Thus, in one instance the disclosure provides a liquid aqueous formulation comprising an anti-TNF α antibody, or antigen-binding portion thereof; a surfactant; and, a polyol; wherein the formulation does not contain a buffer or a salt, and reduces pain associated with injection in a patient by at least about 50% (*e.g.*, at least about 60%, 65%, 70%, 75%, 80%, 85%, 90% or more) when compared to injecting an otherwise identical formulation comprising at least one salt and/or at least one buffer.

55 [0009] In another instance, the disclosure provides a liquid aqueous formulation comprising an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, a surfactant, and less than 50 mg/mL of a polyol, wherein injection

of the formulation into a human subject results in a Pain Visual Analog Scale (VAS) score of less than 1.0. In one instance, the disclosure provides a liquid aqueous formulation consisting essentially of an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, a surfactant, and less than 50 mg/mL of a polyol, wherein injection of the formulation into a human subject results in a Pain Visual Analog Scale (VAS) score of less than 1.0. In one instance, the VAS scale is from 0 (no pain) to 10 (excruciating pain)

[0010] In a further instance, the disclosure provides a liquid aqueous formulation comprising an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, a surfactant, and less than 50 mg/ml of a polyol, wherein the formulation does not contain a buffer and a salt, and wherein injection of the formulation reduces pain associated with the injection in a human subject by at least about 50% when compared to injection of an otherwise identical formulation that comprises a salt and/or a buffer. In one instance, the otherwise identical formulation comprises a citrate and phosphate buffer and sodium chloride.

[0011] The disclosure further provides a liquid aqueous formulation comprising an anti-TNF α antibody or antigen-binding portion thereof, at a concentration of at least about 50 mg/mL; a surfactant; and, a polyol, wherein the formulation has a conductivity of less than about 2 mS/cm. In one instance, the formulation has a conductivity of less than 1 mS/cm. In another instance, the formulation has a conductivity of less than 0.9 mS/cm.

[0012] The disclosure also provides, in another instance, a liquid aqueous formulation comprising an anti-TNF α antibody or antigen-binding portion thereof, at a concentration of at least about 50 mg/mL; a surfactant; and, a polyol, wherein the antibody, or antigen-binding portion thereof, has a hydrodynamic diameter of less than 4 nm in the formulation. In one instance, the antibody or antigen-binding portion thereof, has a hydrodynamic diameter of less than 3 nm in the formulation.

[0013] The present disclosure also provides a liquid aqueous formulation comprising an isolated human anti-TNF α antibody, or an antigen-binding portion thereof; a surfactant; and, less than 50 mg/ml of a polyol; wherein the formulation has a characteristic selected from the group consisting of a conductivity of less than about 2 mS/cm; a hydrodynamic diameter (D_h) which is at least about 50% less than the D_h of the protein in a buffered solution at a given concentration; and a hydrodynamic diameter (D_h) of less than about 4 nm. In one instance, the formulation has a conductivity of less than about 1 mS/cm. In another instance, the formulation has a conductivity of less than about 0.9 mS/cm. In one instance, the antibody or antigen-binding portion thereof, has a hydrodynamic diameter of less than about 3 nm in the formulation. In another instance, the antibody or antigen-binding portion thereof, has a hydrodynamic diameter of less than about 2 nm in the formulation.

[0014] The disclosure also provides a liquid aqueous formulation consisting essentially of an anti-TNF α antibody or antigen-binding portion thereof; a surfactant; and, a polyol; wherein the concentration of the anti-TNF α antibody or antigen-binding portion thereof is at least about 50 mg/mL, 75 mg/mL, 100 mg/mL, or greater than 100 mg/mL.

[0015] In a particular instance, the disclosure provides a liquid aqueous formulation consisting essentially of a concentration of 90-110 mg/ml of an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, having a light chain variable region (LCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 3, or modified from SEQ ID NO: 3 by a single alanine substitution at position 1, 4, 5, 7 or 8, a CDR2 domain comprising the amino acid sequence of SEQ ID NO: 5, and a CDR1 domain comprising the amino acid sequence of SEQ ID NO: 7; and having a heavy chain variable region (HCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 4, or modified from SEQ ID NO: 4 by a single alanine substitution at position 2, 3, 4, 5, 6, 8, 9, 10 or 11, a CDR2 domain comprising the amino acid sequence of SEQ ID NO: 6, and a CDR1 domain comprising the amino acid sequence of SEQ ID NO: 8; a polysorbate, e.g., polysorbate 80; and, about 38-46 mg/ml of a polyol, e.g., mannitol.

[0016] In another instance, the present disclosure provides a liquid aqueous formulation comprising an isolated human anti-TNF α antibody, or an antigen-binding portion thereof; a surfactant; and, less than 50 mg/ml of a polyol; wherein the formulation is stable up to about 30 degrees C for at least about 6 days, about 10, days, or about 14 days, or is stable at about 28 degrees C for up to about 24 months.

[0017] In another instance, the disclosure provides a method of administering an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, to a subject such that injection pain is reduced upon administration, said method comprising subcutaneously administering to the subject a formulation comprising the antibody, or antigen-binding portion thereof, such that injection pain is reduced upon administration, wherein the formulation comprises more than 50 mg/ml of the antibody, or antigen-binding portion thereof; a surfactant; and less than 50 mg/ml of a polyol. In one instance, the injection pain is determined to be less than 1.0 according to a Pain Visual Analog Scale (VAS).

[0018] In certain instances, pain associated with injection is assessed using a pain visual analog scale (VAS). In one instance, the VAS scale is from 0 (no pain) to 10 (excruciating pain)

[0019] In certain instances, the pain associated with injection is assessed after injection (e.g., immediately, no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 minutes, or no more than 15 minutes after injection).

[0020] In certain instances, the formulation reduces pain associated with injection in the patient by at least about 60%, 70%, 80% or more, when compared to injecting the otherwise identical formulation comprising the at least one salt and/or at least one buffer.

[0021] The disclosure further provides a liquid aqueous formulation comprising an anti-TNF α antibody or antigen-binding portion thereof, at a concentration of at least about 50, 75, 100 mg/mL, or greater than 100 mg/mL; a surfactant; and, a polyol; wherein the formulation does not contain a buffer and a salt.

[0022] In another instance, the disclosure provides a liquid aqueous formulation comprising an isolated human anti-TNF α antibody, or an antigen-binding portion thereof; a surfactant; and, less than 50 mg/ml of a polyol; wherein the formulation is stable for up to about 30 degrees C for at least about 6 days. In one instance, the formulation is stable at room temperature for at least about 7 days. In one instance, the formulation is stable at room temperature for at least about 8 days. In one instance, the formulation is stable at room temperature for at least about 9 days. In one instance, the formulation is stable at room temperature for at least about 10 days. In one instance, the formulation is stable at room temperature for at least about 11 days. In one instance, the formulation is stable at room temperature for at least about 12 days. In one instance, the formulation is stable at room temperature for at least about 13 days. In one instance, the formulation is stable at room temperature for at least about 14 days. In one instance, the formulation is stable at room temperature for at least about 15 days.

[0023] In one instance, the polyol used in the formulation of the disclosure is mannitol or sorbitol.

[0024] In one instance, the formulation of the disclosure contains about 20-60 mg/mL mannitol, or, alternatively, about 30-50 mg/mL. In one instance, the formulation contains about 38-46 mg/ml of mannitol.

[0025] The present invention is also based, at least in part, on the unexpected and surprising discovery that a formulation having a high antibody concentration and a surfactant provides notably higher bioavailability than similar formulations containing additional excipients, such as a buffer, a polyol and/or a salt.

[0026] Thus, in one instance, the disclosure provides a liquid aqueous formulation comprising a surfactant and 30-90 mg of an isolated human anti-TNF α antibody or antigen-binding portion, wherein the formulation has an antibody concentration of 90-110 mg/ml, and wherein the formulation provides increased bioavailability of the antibody, or antigen-binding portion thereof, to a human subject upon subcutaneous injection of the formulation relative to a formulation comprising a citrate phosphate buffer, sodium chloride, and mannitol.

[0027] In one instance, the disclosure provides a liquid aqueous formulation consisting essentially of a surfactant and 30-90 mg of an isolated human anti-TNF α antibody or antigen-binding portion, wherein the concentration of the antibody, or antigen-binding portion thereof, is 90-110 mg/ml.

[0028] In another instance, the disclosure provides a liquid aqueous formulation comprising a surfactant and 30-90 mg of an isolated human anti-TNF α antibody, or an antigen-binding portion, wherein the formulation has an antibody concentration of 90-110 mg/ml, and wherein the formulation provides increased bioavailability of the antibody, or antigen-binding portion thereof, in a human subject upon subcutaneous injection of the formulation, such that the antibody or antigen-binding portion thereof, has an AUC₀₋₃₆₀ greater than about 1300 $\mu\text{g} \cdot \text{hr} / \text{ml}$.

[0029] In another instance, the disclosure provides a method for improving the bioavailability of an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, in a human subject, said method comprising administering a formulation comprising an effective amount of the antibody, or antigen-binding portion thereof, and a surfactant to the subject such that the bioavailability of the antibody, or antigen-binding portion thereof, is improved, wherein the formulation does not contain a buffer, a polyol, or a salt.

[0030] In a further instance, the disclosure provides a method of improving the bioavailability of an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, in a subject, said method comprising administering a formulation comprising an effective amount of the antibody, or antigen-binding portion thereof, and a surfactant to the subject such that the bioavailability of the antibody, or antigen-binding portion thereof, in the subject is improved, at least about 15% over a second formulation, wherein the formulation does not contain a buffer, a polyol, or a salt, and wherein the second formulation comprises a buffer, a polyol, and a salt. In one instance, the bioavailability of the antibody, or antigen-binding portion thereof, is improved at least about 30% over the second formulation. In one instance, the bioavailability of the antibody, or antigen-binding portion thereof, is improved at least about 40% over the second formulation.

[0031] The disclosure further provides a method of improving the bioavailability of an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, in a human subject, said method comprising administering a formulation comprising a surfactant and an effective amount of the antibody, or antigen-binding portion thereof, to the subject such that the bioavailability of the antibody, or antigen-binding portion thereof, is improved, wherein the formulation has a characteristic selected from the group consisting of a conductivity of less than about 2 mS/cm; the antibody, or antigen-binding portion thereof, has a hydrodynamic diameter (D_h) which is at least about 50% less than the D_h of the antibody, or antigen-binding portion thereof, in a buffered solution at the given concentration; and the antibody, or antigen-binding portion thereof, has a hydrodynamic diameter (D_h) of less than about 4 nm. In one instance, the formulation has a conductivity of less than about 1 mS/cm. In another instance, the formulation has a conductivity of less than about 0.9 mS/cm. In one instance, the antibody or antigen-binding portion thereof, has a hydrodynamic diameter of less than about 3 nm in the formulation.

[0032] In one instance, the bioavailability is determined according to either an AUC level or a C_{max}. In one instance, the bioavailability is determined according to either an AUC₀₋₃₆₀ or an AUC₀₋₁₃₄₄. In one instance, the bioavailability of

the antibody, or antigen-binding portion thereof, is an AUC₀₋₃₆₀ greater than about 1300 µg * hr / ml when subcutaneously injected into the human subject.

[0033] In certain instances, the anti-TNF α antibody is an isolated human antibody (e.g., a human IgG1 kappa antibody), a humanized antibody, a chimeric antibody, or a murine antibody. For example, the chimeric antibody may be infliximab or a biosimilar thereof, and the human antibody may be golimumab or adalimumab, or a biosimilar thereof.

[0034] In one instance, the human anti-TNF α antibody, or an antigen-binding portion thereof, is an IgG1 or an IgG4.

[0035] In one instance, human anti-TNF α antibody, or an antigen-binding portion thereof, dissociates from human TNF α with a K_d of 1 x 10⁻⁸ M or less and has a k_{off} rate constant of 1 x 10⁻³ s⁻¹ or less, both determined by surface plasmon resonance. In certain instances, the human anti-TNF α antibody, or an antigen-binding portion thereof, dissociates from human TNF α with a K_d of 1 x 10⁻⁸ M or less and a k_{off} rate constant of 1 x 10⁻³ s⁻¹ or less, both determined by surface plasmon resonance, and neutralizes human TNF α cytotoxicity in a standard *in vitro* L929 assay with an IC₅₀ of 1 x 10⁻⁷ M or less.

[0036] In certain instances, the human anti-TNF α antibody, or an antigen-binding portion thereof, has the following characteristics: dissociates from human TNF α with a k_{off} rate constant of 1 x 10⁻³ s⁻¹ or less, as determined by surface plasmon resonance; has a light chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 3, or modified from SEQ ID NO: 3 by a single alanine substitution at position 1, 4, 5, 7 or 8 or by one to five conservative amino acid substitutions at positions 1, 3, 4, 6, 7, 8 and/or 9; and, (c) has a heavy chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 4, or modified from SEQ ID NO: 4 by a single alanine substitution at position 2, 3, 4, 5, 6, 8, 9, 10 or 11 or by one to five conservative amino acid substitutions at positions 2, 3, 4, 5, 6, 8, 9, 10, 11 and/or 12.

[0037] In certain instances, the human anti-TNF α antibody, or an antigen-binding portion thereof, has a light chain variable region (LCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 3, or modified from SEQ ID NO: 3 by a single alanine substitution at position 1, 4, 5, 7 or 8, and with a heavy chain variable region (HCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 4, or modified from SEQ ID NO: 4 by a single alanine substitution at position 2, 3, 4, 5, 6, 8, 9, 10 or 11.

[0038] In certain instances, the human anti-TNF α antibody, or an antigen-binding portion thereof, has a light chain variable region (LCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 3, or modified from SEQ ID NO: 3 by a single alanine substitution at position 1, 4, 5, 7 or 8, a CDR2 domain comprising the amino acid sequence of SEQ ID NO: 5, and a CDR1 domain comprising the amino acid sequence of SEQ ID NO: 7; and has a heavy chain variable region (HCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 4, or modified from SEQ ID NO: 4 by a single alanine substitution at position 2, 3, 4, 5, 6, 8, 9, 10 or 11, a CDR2 domain comprising the amino acid sequence of SEQ ID NO: 6, and a CDR1 domain comprising the amino acid sequence of SEQ ID NO: 8.

[0039] In certain instances, the human anti-TNF α antibody, or an antigen-binding portion thereof, has a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO: 1 and a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO: 2.

[0040] In one instance, the human anti-TNF α antibody, or an antigen-binding portion thereof, comprises the CDRs corresponding to adalimumab.

[0041] In one instance, the human anti-TNF α antibody, or an antigen-binding portion thereof, is adalimumab or golimumab, or a biosimilar thereof.

[0042] In certain instances, the concentration of the human anti-TNF α antibody, or antigen-binding portion thereof, in the formulation is at least about 50 mg/mL, about 75 mg/mL, about 100 mg/mL, or greater than 100 mg/mL. In one instance, the concentration of the human anti-TNF α antibody, or antigen-binding portion thereof, in the formulation of the disclosure is 90-110 mg/ml. In one instance, the concentration of the human anti-TNF α antibody, or antigen-binding portion thereof, in the formulation of the disclosure is 95-105 mg/ml. In one instance, the formulation comprises more than 75 mg/ml of the antibody, or antigen-binding portion thereof. In one instance, the disclosure provides a stable, liquid aqueous formulation comprising a high concentration, e.g., 75-125 mg/mL, of a human anti-hTNF α antibody.

[0043] In certain instances, the surfactant used in the formulation of the disclosure is a polysorbate. In one instance, the concentration of polysorbate is about 0.1-1.5 mg/ml, about 0.2-1.4 mg/ml, about 0.3-1.3 mg/ml, about 0.4-1.2 mg/ml, about 0.5-1.1 mg/ml, about 0.6-1.0 mg/ml, about 0.6-1.1 mg/ml, about 0.7-1.1 mg/ml, about 0.8-1.1 mg/ml, or about 0.9-1.1 mg/ml. In certain instances, the polysorbate is at a concentration of about 0.1-10 mg/mL, about 0.5-5 mg/mL, about 0.1-2 mg/mL, or about 1 mg/mL. In one instance, the surfactant is polysorbate 80.

[0044] In certain instances, the patient is human, or a non-human mammal.

[0045] In certain instances, the formulation is Formulation 3 or Formulation 4 described in the Examples.

[0046] In certain instances, the otherwise identical formulation is the commercially available adalimumab formulation containing adalimumab, sodium chloride, monobasic sodium phosphate dihydrate, dibasic sodium phosphate dihydrate, sodium citrate, citric acid monohydrate, mannitol polysorbate 80, and water for Injection..

[0047] In one instance, the otherwise identical formulation contains a buffer and a salt. In certain instances, the salt is a neutral salt, or a salt from a base (e.g., NaOH) used for pH adjustment. In certain instances, the buffer comprises

a phosphate buffer and/or a citrate buffer. For example, the phosphate buffer may contain about 1.35-1.75 mg/mL or about 1.50-1.56 mg/mL of $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$, and about 0.75-0.95 mg/mL or about 0.83-0.89 mg/mL of $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$. The citrate buffer may contain about 1.15-1.45 mg/mL or about 1.30-1.31 mg/mL of citric acid- H_2O , and about 0.2-0.4 mg/mL or about 0.30-0.31 mg/mL of sodium citrate dehydrate. The at least one salt may be a neutral salt, such as a neutral sodium salt (e.g., NaCl).

[0048] In one embodiment, the formulation of the invention is a pharmaceutical formulation.

[0049] In certain embodiments, the formulation of the invention is suitable for subcutaneous injection. In one embodiment, the formulation of the invention is suitable for subcutaneous self-administration by a subject.

[0050] In certain embodiments, the volume of the aqueous formulation is no more than 1.5 mL, 1.0 mL, 0.8 mL, 0.5 mL, or 0.4 mL.

[0051] In certain instances, the formulation comprises a dose of about 30-90 mg of the antibody, or antigen binding portion thereof. In one instance, the formulation comprises about 40 mg of the anti-TNF α antibody, or antigen binding portion thereof. In one instance, the formulation comprises about 50 mg of the anti-TNF α antibody, or antigen binding portion thereof. In one instance, the formulation comprises about 60 mg of the anti-TNF α antibody, or antigen binding portion thereof. In one instance, the formulation comprises about 70 mg of the anti-TNF α antibody, or antigen binding portion thereof. In one instance, the formulation comprises about 80 mg of the anti-TNF α antibody, or antigen binding portion thereof. In one instance, the formulation comprises about 90 mg of the anti-TNF α antibody, or antigen binding portion thereof. In one instance, the formulation comprises 60-85 mg. In another instance, the formulation comprises 70-90 mg. In yet a further instance, the formulation contains 30-110 mg. In one instance, the formulation contains 70-110 mg.

[0052] Another instance of the disclosure provides a pre-filled syringe or autoinjector device, comprising any of the subject formulations described herein. In certain instances, the aqueous formulation stored in the pre-filled syringe or autoinjector device contains about 40 mg of adalimumab, or biosimilar thereof. In certain instances, the aqueous formulation stored in the pre-filled syringe or autoinjector device contains about 80 mg of adalimumab, or biosimilar thereof.

[0053] Another instance of the disclosure provides a method of treating a disorder associated with detrimental TNF α activity in a patient, comprising administering to the patient any one of the formulations described herein.

[0054] In one embodiment, the formulation of the invention is used to treat a subject having rheumatoid arthritis. In one embodiment, the formulation of the invention is used to treat a subject having Crohn's disease. In one embodiment, the formulation of the invention is used to treat a subject having psoriatic arthritis. In one embodiment, the formulation of the invention is used to treat a subject having psoriasis. In one embodiment, the formulation of the invention is used to treat a subject having juvenile idiopathic arthritis (JIA). In one embodiment, the formulation of the invention is used to treat a subject having ankylosing spondylitis. In one embodiment, the formulation of the invention is used to treat a subject having ulcerative colitis. In one embodiment, the formulation of the invention is used to treat a subject having hidradenitis suppurativa. In one embodiment, the formulation of the invention is used to treat a subject having diabetic retinopathy. In one embodiment, the formulation of the invention is used to treat a subject having giant cell arteritis. In one embodiment, the formulation of the invention is used to treat a subject having Behcet's disease. In one embodiment, the formulation of the invention is used to treat a subject having sarcoidosis, e.g. cutaneous sarcoidosis. In one embodiment, the formulation of the invention is used to treat a subject having axial spondyloarthritis. In one embodiment, the formulation of the invention is used to treat a subject having uveitis.

[0055] In one embodiment, the formulation is administered to the subject according to a periodicity selected from the group consisting of weekly, biweekly, every three weeks, and monthly. In one instance, the formulation of the disclosure contains 30-90 mg of a human anti-TNF α antibody, or antigen-binding portion thereof, and is administered on a biweekly dosing regimen. In another instance, the formulation of the disclosure contains 30-90 mg of a human anti-TNF α antibody, or antigen-binding portion thereof, and is administered according to a monthly dosing regimen. In one instance, the formulation of the disclosure contains 60-85 mg of a human anti-TNF α antibody, or antigen-binding portion thereof, and is administered on a biweekly dosing regimen. In another instance, the formulation of the disclosure contains 60-85 mg of a human anti-TNF α antibody, or antigen-binding portion thereof, and is administered according to a monthly dosing regimen.

[0056] In certain embodiments, the administration of the formulation of the invention to a subject is via self-administration.

[0057] Based on the disclosure that is contained herein, the present invention provides a liquid aqueous formulation consisting essentially of 100 mg/ml of adalimumab, 1 mg/ml of polysorbate-80, 42 mg/ml of mannitol, and water, wherein the formulation does not contain a buffer.

[0058] In a related aspect, the present invention provides a pre-filled syringe or autoinjector device comprising the formulation of the present invention.

[0059] In a related aspect, the present invention provides a formulation for use in treating rheumatoid arthritis, Crohn's disease, psoriatic arthritis, psoriasis, juvenile idiopathic arthritis, ankylosing spondylitis, ulcerative colitis, or hidradenitis suppurativa.

[0060] The present invention and embodiments thereof are set out in the appended claims.

[0061] It is contemplated that any one embodiment described herein can be combined with one or more other embodiments of the invention, including embodiments described only under one aspect of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0062]

Figure 1 is a panel of graphs that show administration of high concentration formulations 1 (F1) and 2 (F2) resulted in a significant decrease in pain assessment at all time points after injection (immediately, 15 minutes, and 30 minutes), compared to the other treatment groups (F4 and the current commercial formulation).

Figure 2 shows, on a linear scale, the means and standard deviations of adalimumab serum concentrations over a time period of 56 days following a single 40 mg SC dose of adalimumab.

Figures 3A and 3B are graphs that show the stability of the various adalimumab formulations assessed by the number of sum aggregates in the formulations (3A) or the sum aggregates (3B) over a range of polysorbate or a range of polyol.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

[0063] In order that the present invention may be more readily understood, certain terms are first defined. In addition, it should be noted that whenever a value or range of values of a parameter are recited, it is intended that values and ranges intermediate to the recited values are also intended to be part of this invention.

[0064] The term "pain associated with injection (in a patient)," as used herein, refers to the pain associated with the injection of drug into the patient's or subject's tissue. In one embodiment, the pain is separate from the pain caused by the injection device (if any), such as the injection needle stick. In one embodiment, the pain associated with injection may originate from the drug formulation being injected into patient's tissue.

[0065] The pain associated with injection may be evaluated using a number of art-recognized means, such as the Pain Visual Analog Scale (VAS). The pain measurement is, in one embodiment, quantifiable, such that a percentage pain scale reduction / increase can be directly compared using statistical methods. For example, when the Pain Visual Analog Scale is used, a numeric pain value (e.g., average $\pm$ SD) can be assigned to each treatment group, such that a percentage increase or reduction can be calculated.

[0066] In general, a Visual Analogue Scale (VAS) is a measurement instrument that measures a characteristic or attitude that is believed to range across a continuum of values (see, e.g., Singer and Thods (1998) Academic Emergency Medicine 5:1007). For example, the amount of pain that a patient feels ranges across a continuum from none (a score of, for example, 0) to an extreme amount of pain (a score of, for example, 10). From the patient's perspective this spectrum appears continuous - their pain does not take discrete jumps, as a categorization of none, mild, moderate and severe would suggest. Operationally, a VAS is usually a horizontal line, 100 mm in length, anchored by word descriptors at each end, such as "no pain" at one end, and "extreme pain" (or some variation thereof) on the other end. The patient marks on the line at a point (for example, a score of 0-10) that they feel represents their perception of their current state. The VAS score may determined by measuring in millimeters from the left hand end of the line to the point that the patient marks.

[0067] There are various ways in which VAS have been presented, including vertical lines and lines with extra descriptors. See Wewers & Lowe ("A critical review of visual analogue scales in the measurement of clinical phenomena." Research in Nursing and Health 13: 227-236, 1990) provide an informative discussion of the benefits and shortcomings of different styles of VAS.

[0068] The term "liquid formulation" refers to a formulation in a liquid state and is not intended to refer to resuspended lyophilized formulations. A liquid formulation of the invention is stable upon storage, and does not rely upon lyophilization (or other state change methods, e.g., spray drying) for stability.

[0069] The term "liquid aqueous formulation" refers to a liquid formulation using water as a solvent. In one embodiment, a liquid aqueous formulation is a formulation that maintains stability (e.g., chemical and/or physical stability/and/or biological activity) without the need for lyophilization, spray-drying, and/or freezing.

[0070] The term "pharmaceutical," as used herein, refers to a composition, e.g., an aqueous formulation, that it is useful for treating a disease or disorder.

[0071] The term "subject" or "patient" is intended to include mammalian organisms. Examples of subjects / patients

include humans and non-human mammals, *e.g.*, non-human primates, dogs, cows, horses, pigs, sheep, goats, cats, mice, rabbits, rats, and transgenic non-human animals. In specific embodiments of the invention, the subject is a human.

[0072] The term "excipient" refers to an agent which may be added to a formulation to provide a desired characteristic, *e.g.*, consistency, improving stability, and/or to adjust osmolality. Examples of commonly used excipients include, but are not limited to, sugars, polyols, amino acids, surfactants, and polymers.

[0073] A commonly used excipient is a polyol. As used herein, a "polyol" is a substance with multiple hydroxyl groups, and includes sugars (reducing and nonreducing sugars), sugar alcohols and sugar acids. Non-limiting examples of polyols are fructose, mannose, maltose, lactose, arabinose, xylose, ribose, rhamnose, galactose, glucose, sucrose, trehalose, sorbose, melezitose, raffinose, mannitol, xylitol erythritol, threitol, sorbitol, glycerol, L-gluconate and metallic salts thereof. In one instance, the polyol used in the formulation or methods of the disclosure is mannitol. In one instance, the polyol used in the formulation or methods of the disclosure is sorbitol.

[0074] A "therapeutically active antibody" or "therapeutic antibody" refers to an antibody which may be used for therapeutic purposes, *i.e.*, for the treatment of a disorder in a subject. It should be noted that while therapeutic proteins may be used for treatment purposes, the invention is not limited to such use, as said proteins may also be used for *in vitro* studies.

[0075] As used herein, "buffer" is an agent(s) in a solution that allows the solution to resist changes in pH by the action of its acid-base conjugate components. Examples of buffers include acetate (*e.g.* sodium acetate), succinate (such as sodium succinate), gluconate, histidine, methionine, citrate, phosphate, citrate / phosphate, imidazole, combinations thereof, and other organic acid buffers. In one embodiment, a buffer is not a protein. A buffer may provide a solution with a pH in the range from about 4 to about 8; from about 4.5 to about 7; or from about 5.0 to about 6.5.

[0076] Although the formulations of the invention do not contain a buffer(s), otherwise identical formulations containing one or more buffers may be used for pain or bioavailability comparison purposes. Examples of such buffers include phosphate, acetate (*e.g.*, sodium acetate), succinate (such as sodium succinate), gluconate, glutamate, histidine, citrate and other organic acid buffers. In one embodiment, a representative buffer in the otherwise identical formulation comprises a citrate buffer and/or a phosphate buffer.

[0077] As used herein, the term "surfactant" generally includes an agent that protects the protein, *e.g.*, antibody, from air/solution interface-induced stresses, solution/surface induced-stresses, to reduce aggregation of the antibody, or to minimize the formation of particulates in the formulation. Exemplary surfactants include, but are not limited to, nonionic surfactants such as polysorbates (*e.g.* polysorbates 20 and 80) or poloxamers (*e.g.* poloxamer 188). The term "surfactant" or "detergent" includes nonionic surfactants such as, but not limited to, polysorbates. In one embodiment, a surfactant includes poloxamers, *e.g.*, Poloxamer 188, Poloxamer 407; polyoxyethylene alkyl ethers, *e.g.*, Brij 35®, Cremophor A25, Sympatens ALM/230; and polysorbates/Tweens, *e.g.*, Polysorbate 20 (Tween 20), Polysorbate 80 (Tween 80), Mirj, and Poloxamers, *e.g.*, Poloxamer 188.

[0078] A "stable" formulation is one in which the antibody therein essentially retains its physical stability and/or chemical stability and/or biological activity during the manufacturing process and/or upon storage. Various analytical techniques for measuring protein stability are available in the art and are reviewed in Peptide and Protein Drug Delivery 247-301, Vincent Lee Ed., Marcel Dekker, Inc., New York, N.Y., Pubs. (1991); and Jones, A. (1993) Adv. Drug Delivery Rev. 10: 29-90). For example, in one embodiment, the stability of a protein is determined according to the percentage of monomer protein in the solution, with a low percentage of degraded (*e.g.*, fragmented) and/or aggregated protein. In one embodiment, the formulation may be stable at room temperature, at about 25-30°C, or at 40°C for at least 1 month and/or stable at about 2-8°C for at least 1 month, 1 year, or, alternatively, for at least 2 years. In another embodiment, the formulation may be stable up to about 30 degrees C for at least about 6 days, about 10, days, or about 14 days, or is stable at about 28 degrees C for up to about 24 months. In one embodiment, the formulation may be stable following freezing (to, *e.g.*, -70°C) and thawing of the formulation, hereinafter referred to as a "freeze/thaw cycle."

[0079] An antibody "retains its physical stability" in a pharmaceutical formulation if it shows substantially no signs of, *e.g.*, aggregation, precipitation and/or denaturation upon visual examination of color and/or clarity, or as measured by UV light scattering or by size exclusion chromatography. Aggregation is a process whereby individual molecules or complexes associate covalently or non-covalently to form aggregates. Aggregation can proceed to the extent that a visible precipitate is formed.

[0080] Stability, such as physical stability of a formulation, may be assessed by methods well-known in the art, including measurement of a sample's apparent attenuation of light (absorbance, or optical density). Such a measurement of light attenuation relates to the turbidity of a formulation. The turbidity of a formulation is partially an intrinsic property of the protein dissolved in solution and is commonly determined by nephelometry, and measured in Nephelometric Turbidity Units (NTU).

[0081] The degree of turbidity, *e.g.*, as a function of the concentration of one or more of the components in the solution, *e.g.*, protein and/or salt concentration, is also referred to as the "opalescence" or "opalescent appearance" of a formulation. The degree of turbidity can be calculated by reference to a standard curve generated using suspensions of known turbidity. Reference standards for determining the degree of turbidity for pharmaceutical compositions can be based on

the European Pharmacopeia criteria (European Pharmacopoeia, Fourth Ed., Directorate for the Quality of Medicine of the Council of Europe (EDQM), Strasbourg, France). According to the European Pharmacopeia criteria, a clear solution is defined as one with a turbidity less than or equal to a reference suspension which has a turbidity of approximately 3 according to European Pharmacopeia standards. Nephelometric turbidity measurements can detect Rayleigh scatter, which typically changes linearly with concentration, in the absence of association or nonideality effects. Other methods for assessing physical stability are well-known in the art.

[0082] An antibody "retains its chemical stability" in a pharmaceutical formulation, if the chemical stability at a given time is such that the antibody is considered to still retain its biological activity as defined below. Chemical stability can be assessed by, e.g., detecting and quantifying chemically altered forms of the antibody. Chemical alteration may involve size modification (e.g. clipping) which can be evaluated using size exclusion chromatography, SDS-PAGE and/or matrix-assisted laser desorption ionization/time-of-flight mass spectrometry (MALDI/TOF MS), for example. Other types of chemical alteration include charge alteration (e.g. occurring as a result of deamidation or oxidation) which can be evaluated by ion-exchange chromatography, for example.

[0083] An antibody "retains its biological activity" in a pharmaceutical formulation, if the antibody in a pharmaceutical formulation is biologically active for its intended purpose. For example, biological activity is retained if the biological activity of the antibody in the pharmaceutical formulation is within about 30%, about 20%, or about 10% (within the errors of the assay) of the biological activity exhibited at the time the pharmaceutical formulation was prepared (e.g., as determined in an antigen binding assay).

[0084] In a pharmacological sense, in the context of the present invention, a "therapeutically effective amount" or "effective amount" of an antibody refers to an amount effective in the prevention or treatment or alleviation of a symptom of a disorder for the treatment of which the antibody is effective.

[0085] The term "human TNF-alpha" (abbreviated herein as hTNF-alpha, TNF α , or simply hTNF), as used herein, is intended to refer to a human cytokine that exists as a 17 kDa secreted form and a 26 kDa membrane associated form, the biologically active form of which is composed of a trimer of noncovalently bound 17 kDa molecules. The structure of hTNF-alpha is described further in, for example, Pennica, D., et al. (1984) *Nature* 312:724-729; Davis, J. M., et al. (1987) *Biochem* 26:1322-1326; and Jones, E. Y., et al. (1989) *Nature* 338:225-228. The term human TNF-alpha is intended to include recombinant human TNF-alpha (rhTNF-alpha), which can be prepared by standard recombinant expression methods or purchased commercially (R & D Systems, Catalog No. 210-TA, Minneapolis, Minn.).

[0086] The term "antibody," as used herein, is intended to refer to immunoglobulin molecules comprised of four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Other naturally occurring antibodies of altered structure, such as, for example, camelid antibodies, are also included in this definition. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. In one instance of the disclosure, the formulation contains an antibody with CDR1, CDR2, and CDR3 sequences like those described in U.S. Patent Nos. 6,090,382 and 6,258,562. In certain instances, the formulation contains an antibody as claimed in U.S. Patent Nos. 6,090,382 and 6,258,562.

[0087] As used herein, the term "CDR" refers to the complementarity determining region within an antibody variable sequence. There are three CDRs in each of the variable regions of the heavy chain and the light chain, which are designated CDR1, CDR2 and CDR3, for each of the heavy and light chain variable regions. The exact boundaries of these CDRs have been defined differently according to different systems. The system described by Kabat (Id.) not only provides an unambiguous residue numbering system applicable to any variable region of an antibody, but also provides precise residue boundaries defining the three CDRs. These CDRs may be referred to as Kabat CDRs. Chothia *et al.* found that certain sub-portions within Kabat CDRs adopt nearly identical peptide backbone conformations, despite having great diversity at the level of amino acid sequence (Chothia et al. (1987) *Mol. Biol.* 196:901-917; Chothia et al. (1989) *Nature* 342:877-883) These sub-portions were designated as L1, L2 and L3 or H1, H2 and H3 where the "L" and the "H" designates the light chain and the heavy chains regions, respectively. These regions may be referred to as Chothia CDRs, which have boundaries that overlap with Kabat CDRs. Other boundaries defining CDRs overlapping with the Kabat CDRs have been described by Padlan (1995) *FASEB J.* 9:133-139 and MacCallum (1996) *J. Mol. Biol.* 262(5):732-45. Still other CDR boundary definitions may not strictly follow one of the herein described systems, but will nonetheless overlap with the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. The methods used herein may utilize CDRs defined according to any of these systems although certain embodiments use Kabat or Chothia defined CDRs. In one instance, the antibody used in the methods and compositions

of the disclosure includes the six CDRs from the antibody adalimumab.

[0088] The term "antigen-binding portion" of an antibody (or simply "antibody portion"), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (e.g., hTNF- α). It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward et al., (1989) Nature 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see e.g., Bird et al. (1988) Science 242:423-426; and Huston et al. (1988) Proc. Natl. Acad. Sci. USA 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. Other forms of single chain antibodies, such as diabodies are also encompassed. Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see e.g., Holliger, P., et al. (1993) Proc. Natl. Acad. Sci. USA 90:6444-6448; Poljak, R.J., et. al. (1994) Structure 2:1121-1123). In one instance of the disclosure, the formulation contains an antigen-binding portions described in U.S. Patent Nos. 6,090,382 and 6,258,562.

[0089] The phrase "recombinant antibody" refers to antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell, antibodies isolated from a recombinant, combinatorial antibody library, antibodies isolated from an animal (e.g., a mouse) that is transgenic for human immunoglobulin genes (see e.g., Taylor et al. (1992) Nucl. Acids Res. 20:6287-6295) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of particular immunoglobulin gene sequences (such as human immunoglobulin gene sequences) to other DNA sequences. Examples of recombinant antibodies include recombinant human, chimeric, CDR-grafted and humanized antibodies.

[0090] The term "human antibody," as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies used in the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs and in particular CDR3. However, the term "human antibody," as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

[0091] The term "chimeric antibody" refers to antibodies which comprise heavy and light chain variable region sequences from one species and constant region sequences from another species, such as antibodies having murine heavy and light chain variable regions linked to human constant regions.

[0092] The term "CDR-grafted antibody" refers to antibodies which comprise heavy and light chain variable region sequences from one species but in which the sequences of one or more of the CDR regions of VH and/or VL are replaced with CDR sequences of another species, such as antibodies having murine heavy and light chain variable regions in which one or more of the murine CDRs (e.g., CDR3) has been replaced with human CDR sequences.

[0093] An "isolated antibody," as used herein, is intended to refer to an antibody that is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds hTNF- α is substantially free of antibodies that specifically bind antigens other than hTNF- α). An isolated antibody that specifically binds hTNF- α may, however, have cross-reactivity to other antigens, such as TNF- α molecules from other species. Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals.

[0094] A "neutralizing antibody," as used herein (or an "antibody that neutralized hTNF- α activity"), is intended to refer to an antibody whose binding to hTNF- α results in inhibition of the biological activity of hTNF- α . This inhibition of the biological activity of hTNF- α can be assessed by measuring one or more indicators of hTNF- α biological activity, such as hTNF- α -induced cytotoxicity (either *in vitro* or *in vivo*), hTNF- α -induced cellular activation and hTNF- α binding to hTNF- α receptors. These indicators of hTNF- α biological activity can be assessed by one or more of several standard *in vitro* or *in vivo* assays known in the art, and described in U.S. Pat. Nos. 6,090,382 and 6,258,562.

[0095] In one embodiment, the ability of an antibody to neutralize hTNF- α activity is assessed by inhibition of hTNF- α -induced cytotoxicity of L929 cells. As an additional or alternative parameter of hTNF- α activity, the ability of an antibody to inhibit hTNF- α -induced expression of ELAM-1 on HUVEC, as a measure of hTNF- α -induced cellular activation, can be assessed.

[0096] The term "surface plasmon resonance," as used herein, refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore system (Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, N.J.). For further descriptions, see Jonsson, U., et al. (1993) *Ann. Biol. Clin.* 51:19-26; Jonsson, U., et al. (1991) *Biotechniques* 11:620-627; 5 Johnson, B., et al. (1995) *J. Mol. Recognit.* 8:125-131; and Johnson, B., et al. (1991) *Anal. Biochem.* 198:268-277.

[0097] The term " k_{on} ," as used herein, is intended to refer to the on rate constant for association of a binding protein (e.g., an antibody) to the antigen to form the, e.g., antibody/antigen complex as is known in the art.

[0098] The term " k_{off} ," as used herein, is intended to refer to the off rate constant for dissociation of an antibody from the antibody/antigen complex.

[0099] The term " k_d ," as used herein, is intended to refer to the dissociation constant of a particular antibody-antigen interaction and refers to the value obtained in a titration measurement at equilibrium, or by dividing the dissociation rate constant (k_{off}) by the association rate constant (k_{on}).

[0100] As used herein, "biosimilar" (of an approved reference product / biological drug, such as a protein therapeutic, antibody, etc.) refers to a biologic product that is similar to the reference product based upon data derived from (a) 15 analytical studies that demonstrate that the biological product is highly similar to the reference product notwithstanding minor differences in clinically inactive components; (b) animal studies (including the assessment of toxicity); and/or (c) a clinical study or studies (including the assessment of immunogenicity and pharmacokinetics or pharmacodynamics) that are sufficient to demonstrate safety, purity, and potency in one or more appropriate conditions of use for which the reference product is licensed and intended to be used and for which licensure is sought for the biological product. In 20 one embodiment, the biosimilar biological product and reference product utilize the same mechanism or mechanisms of action for the condition or conditions of use prescribed, recommended, or suggested in the proposed labeling, but only to the extent the mechanism or mechanisms of action are known for the reference product. In one embodiment, the condition or conditions of use prescribed, recommended, or suggested in the labeling proposed for the biological product have been previously approved for the reference product. In one embodiment, the route of administration, the 25 dosage form, and/or the strength of the biological product are the same as those of the reference product. In one embodiment, the facility in which the biological product is manufactured, processed, packed, or held meets standards designed to assure that the biological product continues to be safe, pure, and potent. The reference product may be approved in at least one of the U.S., Europe, or Japan.

[0101] The term "dosing", as used herein, refers to the administration of a substance (e.g., an anti-TNF α antibody) to 30 achieve a therapeutic objective (e.g., the treatment of a TNF α -associated disorder).

[0102] The terms "weekly dosing regimen", "weekly dosing" and "weekly administration" as used herein, refer to a certain time course (or periodicity) of administering a substance (e.g., an anti-TNF α antibody) to a subject to achieve a therapeutic objective (e.g., the treatment of a TNF α -associated disorder). In one instance, the antibody, or antigen-binding portion thereof, is administered every 6-8 days, or, alternatively, every 7 days.

[0103] The terms "biweekly dosing regimen", "biweekly dosing", and "biweekly administration", as used herein, refer to a certain time course (or periodicity) of administering a substance (e.g., an anti-TNF α antibody) to a subject to achieve a therapeutic objective (e.g., the treatment of a TNF α -associated disorder). The biweekly dosing regimen is not intended to include a weekly dosing regimen. In one instance, the antibody, or antigen-binding portion thereof, is administered every 9-19 days, more preferably, every 11-17 days, even more preferably, every 13-15 days, and most preferably, 40 every 14 days.

[0104] The terms "monthly dosing regimen", "monthly dosing", and "monthly administration", as used herein, refer to a certain time course (or periodicity) of administering a substance (e.g., an anti-TNF α antibody) to a subject to achieve a therapeutic objective (e.g., the treatment of a TNF α -associated disorder). In one instance, a monthly dosing regimen means that the antibody, or antigen-binding portion thereof, is administered every 28-31 days. In another instance, a 45 monthly dosing regimen means that the antibody, or antigen-binding portion thereof, is administered once a month, e.g. on the same day each month, such as, for example, the first day of each month.

[0105] AUC, Cmax, and Tmax are pharmacokinetic parameters that may be used to characterize the pharmacokinetic responses of a particular drug product in an animal or human subject. The term "AUC" refers to the "area under the curve" that represents changes in blood, serum, or plasma concentrations of a substance, e.g., a human anti-TNF α 50 antibody, over time. As used herein, the term "Cmax" refers to the maximum or peak blood, serum, or plasma concentration of substance observed in a subject after its administration. The term "Tmax" refers to the time at which the Cmax occurred, as measured from the time point of administration."

[0106] The term "hydrodynamic diameter" or " D_h " of a particle refers to the diameter of a sphere that has the density of water and the same velocity as the particle. Thus, the term "hydrodynamic diameter of an antibody" as used herein 55 refers to a size determination for an antibody, or an antigen-binding portion thereof, e.g., a human anti-TNF α antibody, or antigen-binding fragment thereof, in solution using dynamic light scattering (DLS). A DLS-measuring instrument measures the time-dependent fluctuation in the intensity of light scattered from the antibody, or antigen-binding fragment thereof, in solution at a fixed scattering angle. D_h is determined from the intensity autocorrelation function of the time-

dependent fluctuation in intensity. Scattering intensity data are processed using DLS instrument software to determine the value for the hydrodynamic diameter and the size distribution of the scattering molecules, e.g. the human anti-TNF α antibody, or antigen-binding fragment thereof, specimen.

[0107] The term "conductivity," as used herein, refers to the ability of an aqueous solution to conduct an electric current between two electrodes. Generally, electrical conductivity or specific conductivity is a measure of a material's ability to conduct an electric current. In solution, the current flows by ion transport. Therefore, with an increasing amount of ions present in the aqueous solution, the solution will have a higher conductivity. The unit of measurement for conductivity is mmhos (mS/cm), and can be measured using a conductivity meter sold, e.g., by Orion Research, Inc. (Beverly, MA). The conductivity of a solution may be altered by changing the concentration of ions therein. For example, the concentration of buffer and/or salt, in the solution may be altered in order to achieve the desired conductivity.

[0108] Conductivity of a solution is measured according to methods known in the art. Conductivity meters and cells may be used to determine the conductivity of the aqueous formulation, and should be calibrated to a standard solution before use. Examples of conductivity meters available in the art include MYRONL Digital (Cole Parmer®), Conductometer (Metrohm AG), and Series 3105/3115 Integrated Conductivity Analyzers (Kemotron).

[0109] Conductivity measurements may be taken with any commercially available conductivity meter suitable for conductivity analysis in protein solutions, e.g. conductivity meter Model SevenMulti, with expansion capacity for broad pH range (Mettler Toledo, Schwerzenbach, Switzerland). The instrument is operated according to the manufacturers instructions (e.g., if the conductivity sensor is changed in the Mettler Toledo instrument, calibration must be performed again, as each sensor has a different cell constant; refer to Operating Instructions of Model SevenMulti conductivity meter). If the instructions are followed, conductivity measurements can be taken by directly immersing the measuring probe into the sample solution.

[0110] Various aspects of the invention are described in further detail in the following subsections.

II. Formulation and Methods of the disclosure

[0111] The present disclosure features stable, liquid aqueous pharmaceutical formulations comprising an anti-TNF α antibody, or an antigen binding portion thereof, having improved properties as compared to art-recognized formulations. While high concentration formulations containing human anti-TNF α antibodies are known in the art (see, for example, US20060153846 and US20100278822), the instant disclosure provides high concentration formulations having unexpected characteristics, *i.e.*, significantly decreased pain or increased bioavailability. The formulations of the disclosure are based, at least in part, on the combination of only one or two excipients, *i.e.*, a surfactant and a polyol or, alternatively, a surfactant alone. Despite having few excipients, the formulations of the disclosure contain a high concentration of an antibody, e.g. 90-110 mg/ml, and are stable.

[0112] As described in the working examples below, a formulation containing an antibody concentration of more than 50 mg/ml of an isolated human anti-TNF α antibody, less than 50 mg/ml of a polyol, (such as mannitol), and a surfactant, (such as a polysorbate), was shown to have dramatically reduced pain upon injection relative to other high concentration formulations, including the commercial adalimumab formulation described in US20060153846, and the formulation described in USA20100278822. Thus, in one embodiment, the formulations of the invention are associated with a reduction of pain, despite having a high antibody concentration (e.g., 100 mg/mL) and having no buffer or salt. The low-pain formulations described herein are based, at least in part, on the surprising finding that by removing or excluding salt (e.g., NaCl) and/or a buffer (e.g., a phosphate / citrate buffer) the concentration of a human anti-TNF alpha antibody in a formulation can be increased, e.g., to about 100 mg/mL, while decreasing pain upon delivery to a patient.

[0113] In one embodiment, the formulation of the invention is surprising, in that the formulation does not contain a buffer or a salt, and reduces pain associated with injection in a patient by at least about 50% when compared to injecting an otherwise identical formulation comprising at least one salt and/or at least one buffer. In one embodiment, the formulation reduces pain associated with the injection in a human subject by at least about 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% (e.g., about 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, or 80%) when compared to the injection of an otherwise identical formulation that further comprises a salt and/or a buffer.

[0114] In one embodiment, the otherwise identical formulation used for pain comparison assay comprises at least one buffer, such as a citrate buffer and a phosphate buffer, and/or a salt, e.g., NaCl. For example, the buffer (excluded from the formulation of the invention and present in the reference formulation for pain comparisons) may include citric acid monohydrate, sodium citrate, disodium phosphate dihydrate, and/or sodium dihydrogen phosphate dihydrate. The buffer may include about 1.15-1.45 mg/ml of citric acid (e.g., about 1.15, 1.20, 1.25, 1.30, 1.35, 1.40, or 1.45), about 0.2-0.4 mg/mL of sodium citrate dehydrate (e.g., about 0.2, 0.25, 0.3, 0.35, or 0.4), about 1.35-1.75 mg/mL of disodium phosphate dehydrate (e.g., about 1.35, 1.40, 1.45, 1.50, 1.55, 1.60, 1.65, 1.70, or 1.75), about 0.75-0.95 mg/mL of sodium dihydrogen phosphate dehydrate (e.g., about 0.75, 0.80, 0.85, 0.9, or 0.95). Values and ranges intermediate to the aforementioned

concentrations are also intended to be part of this invention. In addition, ranges of values using a combination of any of the above-recited values as upper and/or lower limits are intended to be included, e.g., 0.1 to 0.5 mg/mL or 1.20-1.40 mg/mL. In one embodiment, the pH of the formulation is adjusted with sodium hydroxide.

[0115] In one instance, the formulation of the disclosure includes high concentrations of human anti-TNF α antibodies, or antigen binding portions thereof, e.g., 90-110 mg/ml, a polyol at a concentration less than 50 mg/ml, and a surfactant, such that the formulation is suitable for administration without significant pain as determined by a visual analog scale (VAS) score. In one instance, the formulation and methods of the disclosure include high concentrations of anti-TNF α antibodies, or antigen binding portions thereof, and no buffer or salt, such that they are suitable for, administration, e.g., subcutaneous administration, without significant felt pain as determined by a visual analog scale (VAS) score. For example, the formulation of the disclosure may result in a VAS score of less than 1 on a scale of 0 (no pain) to 10 (worst imaginable pain) following subcutaneous injection. As described in Example 1, a formulation having 100 mg/ml of adalimumab, polysorbate 80, and mannitol (less than 50 mg/ml) resulted in a VAS score of less than 1, e.g., 0.56, whereas other high antibody concentration formulations resulted in VAS scores ranging from 1.79 to 4.12.

[0116] In one instance, the disclosure provides a liquid aqueous formulation comprising an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, a surfactant, and less than 50 mg/ml of a polyol, wherein subcutaneous injection of the formulation results in a Pain Visual Analog Scale score of less than 1.0 following injection. In one instance, the formulation does not contain a buffer and a salt, and results in a reduction of pain of at least about 50% upon subcutaneous injection when compared to an injection of an otherwise identical formulation that further comprises a salt and/or a buffer(s).

[0117] Thus, in one embodiment of the invention, liquid formulations of the invention have advantageous tolerability properties in that the formulations produce less pain relative to formulations containing a buffer and a salt. In certain embodiments, the formulation reduces pain associated with injection (or any other form of administration) in a subject. In some embodiments, pain associated with injection is reduced by at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% (e.g., at least about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, or 95%). In one embodiment, pain is reduced by at least about 50%.

[0118] Pain may be evaluated using any type of pain assessment known in the art, including, for example, visual analog scales, qualitative assessments of pain, or needle pain assessments. For example, subject-perceived injection site pain may be assessed using the Pain Visual Analog Scale (VAS). A VAS is a measurement instrument that measures pain as it ranges across a continuum of values, e.g., from none to an extreme amount of pain. Operationally a VAS is a horizontal line, about 100 mm in length, anchored by numerical and/or word descriptors, e.g., 0 or 10, or "no pain" or "excruciating pain," optionally with additional word or numeric descriptors between the extremes, e.g., mild, moderate, and severe; or 1 through 9) (see, e.g., Lee JS, et al. (2000) Acad Emerg Med 7:550, or Singer and Thods (1998) Academic Emergency Medicine 5:1007). Pain may be assessed at a single time or at various times following administration of a formulation of the invention such as, for example, immediately after injection, at about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, or 45 minutes after injection.

[0119] In a certain embodiment of the invention, injection of the formulation into a subject results in a Pain Visual Analog Scale score of less than 0.6, 0.7, 0.8, 0.9, 1.0, 2.0, 3.0, 4.0, or 5.0 on a scale of 0 (no pain) to 10 (excruciating pain).

[0120] Other tools for pain assessment are known in the art, including, for example, the Numerical Rating Scale, the Verbal Rating Scale, and the Brief Pain Inventory. Such tools could also be used to assess pain in accordance with the invention.

[0121] Additional indices for skin irritation may be used, including, e.g., the Draize Scale (hemorrhage, petechiae, erythema, edema, pruritus).

[0122] Formulations of the disclosure containing a polyol preferably contain less than about 50 mg of the polyol. In one instance, the formulations contain less than about 45 mg/mL of the polyol. In another instance, the formulations of the disclosure contain about 38-46 mg/mL of the polyol (e.g., mannitol), e.g., about 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, or 55 mg/mL of the polyol. In addition, ranges of values using a combination of any of the above recited values as upper and/or lower limits are intended to be included, e.g., 39-45 mg/ml, 40-44 mg/ml, or 37-47 mg/ml. In one instance, the formulations of the disclosure contain about 12-72 mg/ml of polyol, e.g., mannitol. In one instance, suitable polyols for use in the formulations and methods of the disclosure are mannitol or sorbitol.

[0123] In one instance, the formulation of the disclosure contains adalimumab (or a biosimilar thereof), polysorbate 80, mannitol, and water for injection. In one embodiment, the formulation contains 80 mg of adalimumab, water for injection, 42 mg/ml of Mannitol, and 1 mg/ml of polysorbate 80. In one embodiment, the formulation may contain 20-110 mg, alternatively 20-90 mg of adalimumab or, alternatively, 30-80 mg of the antibody. In one embodiment, the formulation

contains 30 mg, 31 mg, 32 mg, 33 mg, 34 mg, 35 mg, 36 mg, 37 mg, 38 mg, 39 mg, 40 mg, 41 mg, 42 mg, 43 mg, 44 mg, 45 mg, 46 mg, 47 mg, 48 mg, 49 mg, 50 mg, 51 mg, 52 mg, 53 mg, 54 mg, 55 mg, 56 mg, 57 mg, 58 mg, 59 mg, 60 mg, 61 mg, 62 mg, 63 mg, 64 mg, 65 mg, 66 mg, 67 mg, 68 mg, 69 mg, 70 mg, 71 mg, 72 mg, 73 mg, 74 mg, 75 mg, 76 mg, 77 mg, 78 mg, 79 mg, 80 mg, 81 mg, 82 mg, 83 mg, 84 mg, 85 mg, 86 mg, 87 mg, 88 mg, 89 mg, 90 mg, 91 mg, 92 mg, 93 mg, 94 mg, 95 mg, 96 mg, 97 mg, 98 mg, 99 mg, 100 mg, 101 mg, 102 mg, 103 mg, 104 mg, 105 mg, 106 mg, 107 mg, 108 mg, 109 mg, or 110 mg of the antibody. Ranges including the aforementioned numbers are also included in the invention, e.g., 70-90 mg, 65-95, or 60-85 mg.

[0124] The present invention is also based, at least in part on the surprising discovery that a liquid aqueous pharmaceutical formulation having a high concentration of a human anti-TNF α antibody, or antigen binding portion thereof, and a surfactant (*i.e.*, in the absence of additional excipients), has greater bioavailability than other high concentration formulations having additional excipients. As described in the working examples below, a formulation containing more than 50 mg/ml of an isolated human anti-TNF α antibody, and a polysorbate was shown to have increased bioavailability relative to other high concentration formulations, including the commercial adalimumab formulation described in US20060153846.

[0125] As described in Example 2 below, bioavailability of an anti-TNF α antibody can be increased by combining the antibody with a surfactant, e.g., polysorbate 80. The increase in bioavailability is based on the combination of the antibody and surfactant and the omission or removal of other excipients, including a buffer, polyol, and salt. The increase in bioavailability results in an AUC₀₋₃₆₀ of the anti-TNF α antibody, or an antigen-binding portion thereof, of greater than about 1300 $\mu\text{g} \cdot \text{hr} / \text{ml}$ or an AUC₀₋₁₃₄₄ of the anti-TNF α antibody, or an antigen-binding portion thereof, of greater than about 2600 $\mu\text{g} \cdot \text{hr} / \text{ml}$, when subcutaneously injected into a human subject.

[0126] Accordingly, the present disclosure provides methods for improving the bioavailability of an isolated anti-TNF α antibody, or an antigen-binding portion thereof, in a pharmaceutical formulation. The methods include combining a therapeutically effective amount of the anti-TNF α antibody, or antigen-binding portion thereof, with a surfactant and excluding or removing other excipients, e.g., a buffer(s), salt, and polyol, or combinations thereof, such that the bioavailability of the antibody, or antigen-binding portion thereof, is improved. In one instance, the formulation is injected subcutaneously into a human subject. The methods may improve the bioavailability by providing an AUC₀₋₃₆₀ of the anti-TNF α antibody, or an antigen-binding portion thereof, of greater than about 1100, 1125, 1150, 1175, 1200, 1225, 1250, 1275, 1300, 1325, 1350, 1375, 1400, 1425, 1450, 1475, or about 1500 $\mu\text{g} \cdot \text{hr} / \text{ml}$ when subcutaneously injected into a human subject.

[0127] The disclosure further provides a method of improving the bioavailability of an isolated human anti-TNF α antibody, or an antigen-binding portion thereof, in a subject, said method comprising administering a formulation comprising a surfactant and an effective amount of the antibody, or antigen-binding portion thereof, to the subject such that the bioavailability of the antibody, or antigen-binding portion thereof, in the subject is improved at least about 15% over a second formulation. In one instance, the formulation of the disclosure does not contain a buffer, a polyol, or a salt, and the second formulation comprises a buffer, a polyol, and a salt. In one instance, the bioavailability of the antibody, or antigen-binding portion thereof, is improved at least about 30% over the second formulation. In one instance, the bioavailability of the antibody, or antigen-binding portion thereof, is improved at least about 40% over the second formulation. In one instance, the bioavailability may be determined according to either an AUC level, e.g., AUC₀₋₃₆₀ or an AUC₀₋₁₃₄₄, or a C_{max}.

[0128] In one instance, the present disclosure provides a liquid aqueous formulation which includes a surfactant and about 30-90 mg of an isolated human anti-TNF α antibody or antigen-binding portion, wherein the formulation has an antibody concentration of about 90-110 mg/ml, and wherein the formulation provides increased bioavailability of the antibody, or antigen-binding portion thereof, to a human subject upon subcutaneous injection of the formulation relative to a formulation comprising citrate phosphate buffer, sodium chloride, and mannitol.

[0129] In one instance, the present disclosure provides liquid aqueous formulations which include a surfactant and 30-90 mg of an isolated human anti-TNF α antibody, or an antigen-binding portion, wherein the formulation has an antibody concentration of 90-110 mg/ml, and wherein the formulation provides increased bioavailability of the antibody, or antigen-binding portion thereof, to a human subject upon subcutaneous injection of the formulation, such that the antibody or antigen-binding portion thereof, has an AUC₀₋₃₆₀ greater than about 1100, 1125, 1150, 1175, 1200, 1225, 1250, 1275, 1300, 1325, 1350, 1375, 1400, 1425, 1450, 1475, or about 1500 $\mu\text{g} \cdot \text{hr} / \text{ml}$.

[0130] In one instance, the formulation of the disclosure contains adalimumab (or a biosimilar thereof), polysorbate 80, and water for injection. In one instance, the formulation contains 80 mg of adalimumab, water for injection, and 1 mg/ml polysorbate 80. The formulation may contain 20-110 mg, alternatively 20-90 mg of adalimumab or, alternatively, 30-80 mg of the antibody. In one instance, the formulation contains 30 mg, 31 mg, 32 mg, 33 mg, 34 mg, 35 mg, 36 mg, 37 mg, 38 mg, 39 mg, 40 mg, 41 mg, 42 mg, 43 mg, 44 mg, 45 mg, 46 mg, 47 mg, 48 mg, 49 mg, 50 mg, 51 mg, 52 mg, 53 mg, 54 mg, 55 mg, 56 mg, 57 mg, 58 mg, 59 mg, 60 mg, 61 mg, 62 mg, 63 mg, 64 mg, 65 mg, 66 mg, 67 mg, 68 mg, 69 mg, 70 mg, 71 mg, 72 mg, 73 mg, 74 mg, 75 mg, 76 mg, 77 mg, 78 mg, 79 mg, 80 mg, 81 mg, 82 mg, 83 mg, 84 mg, 85 mg, 86 mg, 87 mg, 88 mg, 89 mg, 90 mg, 91 mg, 92 mg, 93 mg, 94 mg, 95 mg, 96 mg, 97 mg, 98 mg, 99

mg, 100 mg, 101 mg, 102 mg, 103 mg, 104 mg, 105 mg, 106 mg, 107 mg, 108 mg, 109 mg, or 110 mg of the antibody. Ranges including the aforementioned numbers are also included in the disclosure, e.g., 70-90 mg, 65-95 mg, or 60-85 mg.

[0131] Thus, the high antibody formulations and methods of the disclosure not only overcome a number of known challenges for pharmaceutical formulations, including high concentrations in a stable formulation, but also possesses the added benefit of producing improved bioavailability or providing significantly low levels of pain when injected into patients.

[0132] Another obstacle overcome by the formulations of the invention is the ability to remain stable at room temperature (at about 25 degree C or up to about 30 degrees C). Such stability provides advantages for the user of the antibody, providing for more flexible storage options, as the constant need for refrigeration is unnecessary. Both the decreased pain formulation and the increased bioavailability formulation (exemplified by formulations F3 and F4, respectively, in the Examples below) are stable for at least 6 days at about 25 degrees C or up to about 30 degrees C. As described in more detail in the Examples, the formulations of the invention are stable at up to 30 degrees C for at least 6 days, at least 7 days, at least 8 days, at least 9 days, at least 10 days, at least 11 days, at least 12 days, at least 13 days, and at least 14 days. Thus, the invention further provides formulations having extended (*i.e.*, at least 6 days, 10 days or 14 days) shelf life at room temperature (or about 25 degrees C or up to about 30 degrees C). In one embodiment, the formulation of the invention is stable at 20 to 32 degrees C for at least 6 days. Temperatures intermediate to the above recited concentrations are also intended to be part of this invention, *i.e.*, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 and 32 degrees Celsius (C). Ranges including the aforementioned temperatures are also included in the invention, *e.g.*, 22-26 degrees C, 25-30 degrees C, etc.

[0133] The formulations of the disclosure contain a high antibody concentration, including, for example, an antibody concentration of about 50 mg/mL, 55 mg/mL, 60 mg/mL, 65 mg/mL, 70 mg/mL, 75 mg/mL, 80 mg/mL, 85 mg/mL, 90 mg/mL, 95 mg/mL, 100 mg/mL, 105 mg/mL, 110 mg/mL, 115 mg/mL (or higher) of a human anti-TNF-alpha antibody or antigen-binding fragment thereof. Accordingly, as described in the examples below, in one instance of the disclosure the liquid pharmaceutical formulations of the disclosure contain a human anti-TNF alpha antibody concentration of 50-100 mg/mL or greater. In one instance, the formulations of the disclosure may comprise an antibody concentration between about 1 mg/mL - 150 mg/mL or about 40 mg/mL-125 mg/mL. In one instance, the antibody concentration of the formulation is 50-150 mg/mL, 55-150 mg/mL, 60-150 mg/mL, 65-150 mg/mL, 70-150 mg/mL, 75-150 mg/mL, 80-150 mg/mL, 85-150 mg/mL, 90-150 mg/mL, 90-110 mg/mL, 95-105 mg/mL, 95-150 mg/mL, 100-150 mg/mL, 105-150 mg/mL, 110-150 mg/mL, 115-150 mg/mL, 120-150 mg/mL, 125-150 mg/mL, 50-130 mg/mL, 75-125 mg/mL, etc. Concentrations and ranges intermediate to the above recited concentrations are also intended to be part of this disclosure (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150 mg/mL).

[0134] The formulations of the invention may contain an effective amount of the antibody. In one instance, an effective amount is about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or about 100 mg of the human anti-TNF α antibody, or antigen-binding portion thereof. In one instance, the formulations and methods of the disclosure comprise about 20-100, about 20-90, about 30-90, about 30-100, about 60-100, about 70-90, about 40-90, about 60-85 mg, or about 40-100 mg of a human anti-TNF α antibody, or antigen-binding portion thereof. In one instance, the formulation contains 30 mg, 31 mg, 32 mg, 33 mg, 34 mg, 35 mg, 36 mg, 37 mg, 38 mg, 39 mg, 40 mg, 41 mg, 42 mg, 43 mg, 44 mg, 45 mg, 46 mg, 47 mg, 48 mg, 49 mg, 50 mg, 51 mg, 52 mg, 53 mg, 54 mg, 55 mg, 56 mg, 57 mg, 58 mg, 59 mg, 60 mg, 61 mg, 62 mg, 63 mg, 64 mg, 65 mg, 66 mg, 67 mg, 68 mg, 69 mg, 70 mg, 71 mg, 72 mg, 73 mg, 74 mg, 75 mg, 76 mg, 77 mg, 78 mg, 79 mg, 80 mg, 81 mg, 82 mg, 83 mg, 84 mg, 85 mg, 86 mg, 87 mg, 88 mg, 89 mg, or 90 mg of the antibody. Ranges including the aforementioned numbers are also included in the disclosure, e.g., 70-90 or 75-85 mg or 60-85 mg.

[0135] An important aspect of the formulations and methods of the disclosure is the omission of a buffer and salt. Thus, in one instance, the formulations and methods of the disclosure do not contain any buffer(s) (*e.g.*, citrate and phosphate) and salts. It should be noted, however, that although the preferred formulations of the invention do not contain buffers or salts (*e.g.*, NaCl), a small amount of buffer and/or salt may be present in the formulations. Thus, in one embodiment, the formulations of the invention do not contain detectable levels of a buffer(s) and/or a salt.

[0136] In one embodiment, the buffer(s) omitted from the formulations of the invention (or those formulations for comparison which include a buffer(s)) may include citric acid (*e.g.*, about 1.3-1.31 mg/mL or 1.305 mg/mL). In another embodiment, the buffer system includes sodium citrate dehydrate (*e.g.*, about 0.27-0.33 mg/mL or about 0.305 mg/mL). In one embodiment, the buffer system includes disodium phosphate dehydrate (*e.g.*, about 1.5-1.56 mg/mL or about 1.53 mg/mL). In another embodiment, the buffer system includes sodium dihydrogen phosphate dihydrate (*e.g.*, about 0.83-0.89 mg/mL or about 0.86 mg/mL).

[0137] In one embodiment of the invention, the conductivity of the formulation may be used to determine if a formulation has a buffer and/or salt. Both Formulation F3 and F4 (described in the working examples below) have been determined to have a conductivity of less than about 2 mS/cm, *e.g.*, about 0.7 μ S/cm. Thus, in one embodiment, the reduced pain and increased bioavailability formulations of the invention have a conductivity of less than about 2 mS/cm. In another embodiment, the formulations of the invention have a conductivity of less than about 1 mS/cm.

[0138] In one instance, the formulation of the disclosure contains a human anti-TNF alpha antibody, or antigen binding portion thereof, at a concentration of about 100 mg/mL (or 75-125 mg/mL), a surfactant (*e.g.*, polysorbate 80), a polyol (*e.g.*, sorbitol or mannitol), and has a conductivity of less than 2 mS/cm. In one instance, the formulation of the disclosure contains a human anti-TNF alpha antibody, or antigen binding portion thereof, at a concentration of about 100 mg/mL (or 75-125 mg/mL), about 0.8-1.3 mg/ml of a surfactant (*e.g.*, polysorbate 80), less than about 50 mg/ml of a polyol (*e.g.*, sorbitol or mannitol), and has a conductivity of less than 2 mS/cm.

[0139] In one instance, the formulation of the disclosure contains a human anti-TNF alpha antibody, or antigen binding portion thereof, at a concentration of about 100 mg/mL (or 75-125 mg/mL), a surfactant (*e.g.*, polysorbate 80), and has a conductivity of less than 2 mS/cm. In one instance the formulation of the disclosure contains a human anti-TNF alpha antibody, or antigen binding portion thereof, at a concentration of about 100 mg/mL (or 75-125 mg/mL), about 0.8-1.3 mg/ml of a surfactant (*e.g.*, polysorbate 80), and has a conductivity of less than 2 mS/cm.

[0140] In another instance, the disclosure provides a stable formulation having a high concentration antibody, or antigen-binding portion thereof, wherein the antibody, or antigen has a hydrodynamic diameter (z-average) of less than about 4 nm or wherein the antibody, or antigen has a hydrodynamic diameter (z-average) which is at least about 50% less than the hydrodynamic diameter of a buffered solution at the same antibody concentration. In one instance, the antibody, or antigen has a hydrodynamic diameter (z-average) of less than about 3 nm.

[0141] In one instance, the formulation of the disclosure contains a human anti-TNF alpha antibody, or antigen binding portion thereof, at a concentration of about 100 mg/mL (or 75-125 mg/mL), a surfactant (*e.g.*, polysorbate 80), a polyol (*e.g.*, sorbitol or mannitol), and has a hydrodynamic diameter of less than 4 nm. In one instance, the formulation of the disclosure contains a human anti-TNF alpha antibody, or antigen binding portion thereof, at a concentration of about 100 mg/mL (or 75-125 mg/mL), about 0.8-1.3 mg/ml of a surfactant (*e.g.*, polysorbate 80), less than about 50 mg/ml of a polyol (*e.g.*, sorbitol or mannitol), and has a hydrodynamic diameter of less than 4 nm.

[0142] In one instance, the formulation of the disclosure contains a human anti-TNF alpha antibody, or antigen binding portion thereof, at a concentration of about 100 mg/mL (or 75-125 mg/mL), a surfactant (*e.g.*, polysorbate 80), and has a hydrodynamic diameter of less than 4 nm. In one instance, the formulation of the disclosure contains a human anti-TNF alpha antibody, or antigen binding portion thereof, at a concentration of about 100 mg/mL (or 75-125 mg/mL), about 0.8-1.3 mg/ml of a surfactant (*e.g.*, polysorbate 80), and has a hydrodynamic diameter of less than 4 nm.

[0143] A detergent or surfactant is included in the antibody formulation of the disclosure. Exemplary detergents include nonionic detergents such as polysorbates (*e.g.* polysorbates 20, 80. *etc.*) or poloxamers (*e.g.* poloxamer 188). The amount of detergent added is such that it reduces aggregation of the formulated antibody and/or minimizes the formation of particulates in the formulation and/or reduces adsorption. In a preferred instance of the disclosure, the formulation includes a surfactant which is a polysorbate. In another preferred instance of the disclosure, the formulation contains the detergent polysorbate 80. In one instance, the formulation contains between about 0.1 and about 2.0 mg/mL of surfactant (*e.g.*, polysorbate), *e.g.*, about 1 mg/mL. Other ranges of polysorbate that may be included in the formulations of the disclosure include 0.1 to 1.5 mg/ml, alternatively 0.2-1.4 mg/ml, 0.3-1.3 mg/ml, 0.4-1.2 mg/ml, 0.5-1.1 mg/ml, 0.6-1.0 mg/ml, 0.6-1.1 mg/ml, 0.7-1.1 mg/ml, 0.8-1.1 mg/ml, or 0.9-1.1 mg/ml. Values and ranges intermediate to the above recited concentrations are also intended to be part of this disclosure, *e.g.*, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9. In addition, ranges of values using a combination of any of the above-recited values as upper and/or lower limits are intended to be included, *e.g.*, 0.3 to 1.1 mg/mL.

[0144] In one instance, the formulation of the disclosure consists essentially of a human anti-TNF alpha antibody, or antigen binding portion thereof, at a concentration of about 100 mg/mL (or 75-125 mg/mL), a surfactant (*e.g.*, polysorbate 80), a polyol (*e.g.*, sorbitol or mannitol), does not contain a buffer(s) (*e.g.*, citric acid monohydrate, sodium citrate, disodium phosphate dihydrate, and/or sodium dihydrogen phosphate dihydrate), and does not contain a salt (*e.g.*, NaCl).

[0145] In certain embodiments, the otherwise identical formulation to which the formulation of the invention is compared for pain or bioavailability purposes is a formulation containing adalimumab, sodium chloride, monobasic sodium phosphate dihydrate, dibasic sodium phosphate dihydrate, sodium citrate, citric acid monohydrate, mannitol, polysorbate 80, and Water for Injection.

[0146] The formulation herein may also be combined with one or more other therapeutic agents as necessary for the particular indication being treated. In one embodiment, those with complementary activities that do not adversely affect the antibody of the formulation. Such therapeutic agents are suitably present in combination in amounts that are effective for the purpose intended. Additional therapeutic agents which can be combined with the formulation of the invention are further described in U.S. Pat. Nos. 6,090,382 and 6,258,562.

[0147] All formulations described herein may be used in the methods of the disclosure as well.

III. Antibodies for Use in the Formulations and Methods of the disclosure

[0148] The formulations and methods of the disclosure include an antibody, or antigen binding portion thereof, particularly an anti-TNF α antibody, or antigen binding portion or fragment thereof. Examples of antibodies that may be used in the disclosure include chimeric antibodies, non-human antibodies, isolated human antibodies, humanized antibodies, and domain antibodies (dAbs). All antibodies described herein may be used in the methods of the disclosure as well.

[0149] In one instance, the formulations of the disclosure comprises an antibody, or antigen-binding portion thereof, which binds human TNF α , including, for example, adalimumab (also referred to as Humira, adalimumab, or D2E7; Abbott Laboratories). In a further instance, the formulation comprises an antibody that binds the same epitope as adalimumab, such as, but not limited to, an adalimumab biosimilar antibody. In one instance, the antibody is a human IgG1 antibody having six CDRs corresponding to those of the light and heavy chain of adalimumab.

[0150] In one instance, the disclosure features an isolated human antibody, or antigen-binding portion thereof, that binds to human TNF-alpha with high affinity and a low off rate, and also has a high neutralizing capacity. In one instance, the human antibodies used in the disclosure are recombinant, neutralizing human anti-hTNF-alpha antibodies.

[0151] In one instance, the disclosure pertains to adalimumab antibodies and antibody portions, adalimumab-related antibodies and antibody portions, and other human antibodies and antibody portions with equivalent properties to adalimumab, such as high affinity binding to hTNF α , with low dissociation kinetics and high neutralizing capacity. In one instance, the antibody, or antigen-binding fragment thereof, is defined according to dissociation and binding characteristics similar to adalimumab. For example, the formulation may include a human antibody that dissociates from human TNF α with a K_d of 1×10^{-8} M or less, and a k_{off} rate constant of 1×10^{-3} s $^{-1}$ or less, both determined by surface plasmon resonance. In another instance, the human antibody that dissociates from human TNF α with a K_d of 1×10^{-9} M or less.

[0152] In one instance, the antibody, or antigen-binding fragment thereof, is a human antibody that dissociates from human TNF α with a K_d of 1×10^{-8} M or less, and a k_{off} rate constant of 1×10^{-3} s $^{-1}$ or less, both determined by surface plasmon resonance, and neutralizes human TNF α cytotoxicity in a standard *in vitro* L929 assay with an IC $_{50}$ of 1×10^{-7} M or less. Examples and methods for making human, neutralizing antibodies which have a high affinity for human TNF α , including sequences of the antibodies, are described in U.S. Patent No. 6,090,382 (referred to as D2E7).

[0153] The antibody used in the formulation of the invention is D2E7, also referred to as HUMIRA[™] or adalimumab (the amino acid sequence of the D2E7 VL region is shown in SEQ ID NO: 1; the amino acid sequence of the D2E7 VH region is shown in SEQ ID NO: 2). The properties of D2E7 (adalimumab / HUMIRA[®]) have been described in Salfeld et al., U.S. Patent Nos. 6,090,382, 6,258,562, and 6,509,015.

[0154] In one instance, the human TNF-alpha, or an antigen-binding portion thereof, dissociates from human TNF-alpha with a K_d of 1×10^{-8} M or less and a k_{off} rate constant of 1×10^{-3} s $^{-1}$ or less, both determined by surface plasmon resonance, and neutralizes human TNF-alpha cytotoxicity in a standard *in vitro* L929 assay with an IC $_{50}$ of 1×10^{-7} M or less. In one instance, the isolated human antibody, or antigen-binding portion thereof, dissociates from human TNF-alpha with a k_{off} of 5×10^{-4} s $^{-1}$ or less; or, in one instance, with a k_{off} of 1×10^{-4} s $^{-1}$ or less. In one instance, the isolated human antibody, or antigen-binding portion thereof, neutralizes human TNF-alpha cytotoxicity in a standard *in vitro* L929 assay with an IC $_{50}$ of 1×10^{-8} M or less; or, in one instance, with an IC $_{50}$ of 1×10^{-9} M or less; or, in one instance, with an IC $_{50}$ of 1×10^{-10} M or less. In one instance, the antibody is an isolated human recombinant antibody, or an antigen-binding portion thereof.

[0155] It is well known in the art that antibody heavy and light chain CDR3 domains play an important role in the binding specificity/affinity of an antibody for an antigen. Accordingly, in another instance, the antibody used in the formulation of the disclosure has slow dissociation kinetics for association with hTNF-alpha and has light and heavy chain CDR3 domains that structurally are identical to or related to those of adalimumab. Position 9 of the adalimumab VL CDR3 can be occupied by Ala or Thr without substantially affecting the k_{off} . Accordingly, a consensus motif for the adalimumab VL CDR3 comprises the amino acid sequence: Q-R-Y-N-R-A-P-Y-(T/A) (SEQ ID NO: 3). Additionally, position 12 of the adalimumab VH CDR3 can be occupied by Tyr or Asn, without substantially affecting the k_{off} . Accordingly, a consensus motif for the adalimumab VH CDR3 comprises the amino acid sequence: V-S-Y-L-S-T-A-S-S-L-D-(Y/N) (SEQ ID NO: 4). Moreover, as demonstrated in Example 2 of U.S. Patent No. 6,090,382, the CDR3 domain of the adalimumab heavy and light chains is amenable to substitution with a single alanine residue (at position 1, 4, 5, 7 or 8 within the VL CDR3 or at position 2, 3, 4, 5, 6, 8, 9, 10 or 11 within the VH CDR3) without substantially affecting the k_{off} . Still further, the skilled artisan will appreciate that, given the amenability of the adalimumab VL and VH CDR3 domains to substitutions by alanine, substitution of other amino acids within the CDR3 domains may be possible while still retaining the low off rate constant of the antibody, in particular substitutions with conservative amino acids. In one instance, no more than one to five conservative amino acid substitutions are made within the adalimumab VL and/or VH CDR3 domains. In one instance, no more than one to three conservative amino acid substitutions are made within the adalimumab VL and/or VH CDR3 domains. Additionally, conservative amino acid substitutions should not be made at amino acid positions critical for binding to hTNF alpha. Positions 2 and 5 of the adalimumab VL CDR3 and positions 1 and 7 of the adalimumab VH CDR3 appear to be critical for interaction with hTNF alpha, and thus, conservative amino acid substitutions preferably

are not made at these positions (although an alanine substitution at position 5 of the adalimumab VL CDR3 is acceptable, as described above) (see U.S. Patent No. 6,090,382).

[0156] Accordingly, in one instance, the antibody or antigen-binding portion thereof, used in the formulation of the disclosure contains the following characteristics:

a) dissociates from human TNF α with a k_{off} rate constant of $1 \times 10^{-3} \text{ s}^{-1}$ or less, as determined by surface plasmon resonance;

b) has a light chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 3, or modified from SEQ ID NO: 3 by a single alanine substitution at position 1, 4, 5, 7 or 8 or by one to five conservative amino acid substitutions at positions 1, 3, 4, 6, 7, 8 and/or 9;

c) has a heavy chain CDR3 domain comprising the amino acid sequence of SEQ ID NO: 4, or modified from SEQ ID NO: 4 by a single alanine substitution at position 2, 3, 4, 5, 6, 8, 9, 10 or 11 or by one to five conservative amino acid substitutions at positions 2, 3, 4, 5, 6, 8, 9, 10, 11 and/or 12.

[0157] In certain instances, the antibody or antigen-binding portion thereof, dissociates from human TNF-alpha with a k_{off} of $5 \times 10^{-4} \text{ s}^{-1}$ or less. In certain instances, the antibody or antigen-binding portion thereof, dissociates from human TNF-alpha with a k_{off} of $1 \times 10^{-4} \text{ s}^{-1}$ or less.

[0158] In yet another instance, the antibody or antigen-binding portion thereof contains a light chain variable region (LCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 3, or modified from SEQ ID NO: 3 by a single alanine substitution at position 1, 4, 5, 7 or 8, and with a heavy chain variable region (HCVR) having a CDR3 domain comprising the amino acid sequence of SEQ ID NO: 4, or modified from SEQ ID NO: 4 by a single alanine substitution at position 2, 3, 4, 5, 6, 8, 9, 10 or 11. In one instance, the LCVR further has a CDR2 domain comprising the amino acid sequence of SEQ ID NO: 5 (*i.e.*, the D2E7 VL CDR2) and the HCVR further has a CDR2 domain comprising the amino acid sequence of SEQ ID NO: 6 (*i.e.*, the D2E7 VH CDR2). In one instance, the LCVR further has CDR1 domain comprising the amino acid sequence of SEQ ID NO: 7 (*i.e.*, the D2E7 VL CDR1) and the HCVR has a CDR1 domain comprising the amino acid sequence of SEQ ID NO: 8 (*i.e.*, the D2E7 VH CDR1). The framework regions for VL may be from the VkI human germline family, or from the A20 human germline Vk gene, or from the adalimumab VL framework sequences shown in Figures 1A and 1B of U.S. Patent No. 6,090,382. The framework regions for VH may be from the VH3 human germline family, or from the DP-31 human germline VH gene, or from the D2E7 VH framework sequences shown in Figures 2A and 2B of U.S. Patent No. 6,090,382. Nucleic acid sequences corresponding to the adalimumab light and heavy variable regions are described in SEQ ID NOs: 36 and 37, respectively.

[0159] Accordingly, in another instance, the antibody or antigen-binding portion thereof contains a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO: 1 (*i.e.*, the adalimumab VL) and a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO: 2 (*i.e.*, the adalimumab VH). In certain instances, the antibody comprises a heavy chain constant region, such as an IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM or IgD constant region. In certain instances, the heavy chain constant region is an IgG1 heavy chain constant region or an IgG4 heavy chain constant region. Furthermore, the antibody can comprise a light chain constant region, either a kappa light chain constant region or a lambda light chain constant region. In one instance, the antibody comprises a kappa light chain constant region. Alternatively, the antibody portion can be, for example, a Fab fragment or a single chain Fv fragment.

[0160] In still other instances, the disclosure includes uses of an isolated human antibody, or an antigen-binding portion thereof, containing adalimumab-related VL and VH CDR3 domains. For example, antibodies or antigen-binding portions thereof may have a light chain variable region (LCVR) having a CDR3 domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 3, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25 and SEQ ID NO: 26 or with a heavy chain variable region (HCVR) having a CDR3 domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34 and SEQ ID NO: 35.

[0161] In one instance, the TNF α antibody used in the disclosure includes the chimeric antibody infliximab (Remicade®, Johnson and Johnson; described in U.S. Patent No. 5,656,272), CDP571 (a humanized monoclonal anti-TNF-alpha IgG4 antibody), CDP 870 (a humanized monoclonal anti-TNF-alpha antibody fragment), an anti-TNF dAb (Peptech), or CNTO 148 (golimumab; Medarex and Centocor, see WO 02/12502). Additional TNF antibodies which may be used in the disclosure are described in U.S. Patent Nos. 6,593,458; 6,498,237; 6,451,983; and 6,448,380.

[0162] An antibody, or antibody portion, used in the methods and compositions of the disclosure, can be prepared by recombinant expression of immunoglobulin light and heavy chain genes in a host cell. To express an antibody

recombinantly, a host cell is transfected with one or more recombinant expression vectors carrying DNA fragments encoding the immunoglobulin light and heavy chains of the antibody such that the light and heavy chains are expressed in the host cell and, preferably, secreted into the medium in which the host cells are cultured, from which medium the antibodies can be recovered. Standard recombinant DNA methodologies are used to obtain antibody heavy and light chain genes, incorporate these genes into recombinant expression vectors and introduce the vectors into host cells, such as those described in Sambrook, Fritsch and Maniatis (eds), *Molecular Cloning; A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y., (1989), Ausubel, F.M. et al. (eds.) *Current Protocols in Molecular Biology*, Greene Publishing Associates, (1989) and in U.S. Patent No. 4,816,397 by Boss et al.

[0163] To express an anti-TNF α antibody, e.g., adalimumab (D2E7) or an adalimumab (D2E7)-related antibody, DNA fragments encoding the light and heavy chain variable regions are first obtained. These DNAs can be obtained by amplification and modification of germline light and heavy chain variable sequences using the polymerase chain reaction (PCR). Germline DNA sequences for human heavy and light chain variable region genes are known in the art (see e.g., the "Vbase" human germline sequence database; see also Kabat, E.A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Tomlinson, I.M., et al. (1992) "The Repertoire of Human Germline VH Sequences Reveals about Fifty Groups of VH Segments with Different Hypervariable Loops" *J. Mol. Biol.* 227:776-798; and Cox, J.P.L. et al. (1994) "A Directory of Human Germ-line V78 Segments Reveals a Strong Bias in their Usage" *Eur. J. Immunol.* 24:827-836). For example, to obtain a DNA fragment encoding the heavy chain variable region of D2E7, or a D2E7-related antibody, a member of the VH3 family of human Germline VH genes is amplified by standard PCR. In certain embodiments, the DP-31 VH-germline sequence is amplified. To obtain a DNA fragment encoding the light chain variable region of D2E7, or a D2E7-related antibody, a member of the V κ L family of human germline VL genes is amplified by standard PCR. In certain embodiments, the A20 VL germline sequence is amplified. PCR primers suitable for use in amplifying the DP-31 germline VH and A20 Germline VL sequences can be designed based on the nucleotide sequences disclosed in the references cited supra, using standard methods.

[0164] Once the germline VH and VL fragments are obtained, these sequences can be mutated to encode the anti-TNF α antibody amino acid sequences disclosed herein. The amino acid sequences encoded by the Germline VH and VL DNA sequences are first compared to the anti-TNF α antibody VH and VL amino acid sequences to identify amino acid residues in the anti-TNF α antibody sequence that differ from germline. Then, the appropriate nucleotides of the germline DNA sequences are mutated such that the mutated germline sequence encodes the anti-TNF α antibody amino acid sequence, using the genetic code to determine which nucleotide changes should be made. Mutagenesis of the germline sequences is carried out by standard methods, such as PCR-mediated mutagenesis (in which the mutated nucleotides are incorporated into the PCR primers such that the PCR product contains the mutations) or site-directed mutagenesis.

[0165] Moreover, it should be noted that if the "germline" sequences obtained by PCR amplification encode amino acid differences in the framework regions from the true germline configuration (*i.e.*, differences in the amplified sequence as compared to the true germline sequence, for example as a result of somatic mutation), it may be desirable to change these amino acid differences back to the true germline sequences (*i.e.*, "backmutation" of framework residues to the germline configuration).

[0166] Once DNA fragments encoding the anti-TNF α antibody VH and VL segments are obtained (e.g., by amplification and mutagenesis of germline VH and VL genes, as described above), these DNA fragments can be further manipulated by standard recombinant DNA techniques, for example to convert the variable region genes to full-length antibody chain genes, to Fab fragment genes or to a scFv gene. In these manipulations, a VL- or VH-encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker. The term "operatively linked," as used in this context, is intended to mean that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame.

[0167] The isolated DNA encoding the VH region can be converted to a full-length heavy chain gene by operatively linking the VH-encoding DNA to another DNA molecule encoding heavy chain constant regions (CH1, CH2 and CH3). The sequences of human heavy chain constant region genes are known in the art (see e.g., Kabat, E.A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM or IgD constant region, but most preferably is an IgG1 or IgG4 constant region. For a Fab fragment heavy chain gene, the VH-encoding DNA can be operatively linked to another DNA molecule encoding only the heavy chain CH1 constant region.

[0168] The isolated DNA encoding the VL region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the VL-encoding DNA to another DNA molecule encoding the light chain constant region, CL. The sequences of human light chain constant region genes are known in the art (see e.g., Kabat, E.A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR

amplification. The light chain constant region can be a kappa or lambda constant region. In one instance, the light chain constant region is a kappa constant region.

[0169] To create a scFv gene, the VH- and VL-encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, e.g., encoding the amino acid sequences (Gly₄-Ser)₃, such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see e.g., Bird et al. (1988) Science 242:423-426; Huston et. al. (1988) Proc. Natl. Acad. Sci. USA 85:5879-5883; McCafferty et al., Nature (1990) 348:552-554).

[0170] To express the antibodies, or antibody portions used in the disclosure, DNAs encoding partial or full-length light and heavy chains, obtained as described above, are inserted into expression vectors such that the genes are operatively linked to transcriptional and translational control sequences. In this context, the term "operatively linked" is intended to mean that an antibody gene is ligated into a vector such that transcriptional and translational control sequences within the vector serve their intended function of regulating the transcription and translation of the antibody gene. The expression vector and expression control sequences are chosen to be compatible with the expression host cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vector or, more typically, both genes are inserted into the same expression vector. The antibody genes are inserted into the expression vector by standard methods (e.g., ligation of complementary restriction sites on the antibody gene fragment and vector, or blunt end ligation if no restriction sites are present) Prior to insertion of the anti-TNFα antibody light or heavy chain sequences, the expression vector may already carry antibody constant region sequences. For example, one approach to converting the anti-TNFα antibody VH and VL sequences to full-length antibody genes is to insert them into expression vectors already encoding heavy chain constant and light chain constant regions, respectively, such that the VH segment is operatively linked to the CH segment(s) within the vector and the VL segment is operatively linked to the CL segment within the vector. Additionally or alternatively, the recombinant expression vector can encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The antibody chain gene can be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the antibody chain gene. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide (i.e., a signal peptide from a non-immunoglobulin protein).

[0171] In addition to the antibody chain genes, the recombinant expression vectors of the disclosure carry regulatory sequences that control the expression of the antibody chain genes in a host cell. The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals) that control the transcription or translation of the antibody chain genes. Such regulatory sequences are described, for example, in Goeddel; Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990). It will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. Preferred regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from cytomegalovirus (CMV) (such as the CMV promoter/enhancer), Simian Virus 40 (SV40) (such as the SV40 promoter/enhancer), adenovirus, (e.g., the adenovirus major late promoter (AdMLP)) and polyoma. For further description of viral regulatory elements, and sequences thereof, see e.g., U.S. Patent No. 5,168,062 by Stinski, U.S. Patent No. 4,510,245 by Bell et al. and U.S. Patent No. 4,968,615 by Schaffner et al.

[0172] In addition to the antibody chain genes and regulatory sequences, the recombinant expression vectors used in the disclosure may carry additional sequences, such as sequences that regulate replication of the vector in host cells (e.g., origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see e.g., U.S. Patents Nos. 4,399,216, 4,634,665 and 5,179,017, all by Axel et al.). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Preferred selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in dhfr- host cells with methotrexate selection/amplification) and the neo gene (for G418 selection).

[0173] For expression of the light and heavy chains, the expression vector(s) encoding the heavy and light chains is transfected into a host cell by standard techniques. The various forms of the term "transfection" are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or eukaryotic host cell, e.g., electroporation, calcium-phosphate precipitation, DEAE-dextran transfection and the like. Although it is theoretically possible to express the antibodies of the invention in either prokaryotic or eukaryotic host cells, expression of antibodies is preferably in eukaryotic cells. In one embodiment, mammalian host cells, is the most preferred because such eukaryotic cells, and in particular mammalian cells, are more likely than prokaryotic cells to assemble and secrete a properly folded and immunologically active antibody. Prokaryotic expression of antibody genes has been reported to be ineffective for production of high yields of active antibody (Boss, M.A. and Wood, C. R. (1985) Immunology Today 6:12-13).

[0174] Preferred mammalian host cells for expressing the recombinant antibodies of the invention include Chinese

Hamster Ovary (CHO cells) (including dhfr- CHO cells, described in Urlaub and Chasin, (1980) Proc. Natl. Acad. Sci. USA 77:4216-4220, used with a DHFR selectable marker, e.g., as described in R.J. Kaufman and P.A. Sharp (1982) Mol. Biol. 159:601-621), NS0 myeloma cells, COS cells and SP2 cells. When recombinant expression vectors encoding antibody genes are introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more, in one embodiment, secretion of the antibody into the culture medium in which the host cells are grown. Antibodies can be recovered from the culture medium using standard protein purification methods.

[0175] Host cells can also be used to produce portions of intact antibodies, such as Fab fragments or scFv molecules. It is understood that variations on the above procedure are within the scope of the present disclosure. For example, it may be desirable to transfect a host cell with DNA encoding either the light chain or the heavy chain (but not both) of an antibody of this disclosure. Recombinant DNA technology may also be used to remove some or all of the DNA encoding either or both of the light and heavy chains that is not necessary for binding to hTNF alpha. The molecules expressed from such truncated DNA molecules are also encompassed by the antibodies of the disclosure. In addition, bifunctional antibodies may be produced in which one heavy and one light chain are an antibody of the disclosure and the other heavy and light chain are specific for an antigen other than hTNF alpha by crosslinking an antibody of the invention to a second antibody by standard chemical crosslinking methods.

[0176] In a preferred system for recombinant expression of an antibody, or antigen-binding portion thereof, of the disclosure, a recombinant expression vector encoding both the antibody heavy chain and the antibody light chain is introduced into dhfr-CHO cells by calcium phosphate-mediated transfection. Within the recombinant expression vector, the antibody heavy and light chain genes are each operatively linked to CMV enhancer/AdMLP promoter regulatory elements to drive high levels of transcription of the genes. The recombinant expression vector also carries a DHFR gene, which allows for selection of CHO cells that have been transfected with the vector using methotrexate selection/amplification. The selected transformant host cells are culture to allow for expression of the antibody heavy and light chains and intact antibody is recovered from the culture medium. Standard molecular biology techniques are used to prepare the recombinant expression vector, transfect the host cells, select for transformants, culture the host cells and recover the antibody from the culture medium.

[0177] In view of the foregoing, nucleic acid, vector and host cell compositions that can be used for recombinant expression of the antibodies and antibody portions used in the disclosure include nucleic acids, and vectors comprising said nucleic acids, comprising the human TNF alpha antibody adalimumab (D2E7). The nucleotide sequence encoding the D2E7 light chain variable region is shown in SEQ ID NO: 36. The CDR1 domain of the LCVR encompasses nucleotides 70-102, the CDR2 domain encompasses nucleotides 148-168 and the CDR3 domain encompasses nucleotides 265-291. The nucleotide sequence encoding the D2E7 heavy chain variable region is shown in SEQ ID NO: 37. The CDR1 domain of the HCVR encompasses nucleotides 91-105, the CDR2 domain encompasses nucleotides 148-198 and the CDR3 domain encompasses nucleotides 295-330. It will be appreciated by the skilled artisan that nucleotide sequences encoding D2E7-related antibodies, or portions thereof (e.g., a CDR domain, such as a CDR3 domain), can be derived from the nucleotide sequences encoding the D2E7 LCVR and HCVR using the genetic code and standard molecular biology techniques.

[0178] In one instance, the liquid pharmaceutical formulation comprises a human TNF alpha antibody, or antigen-binding portion thereof that is a bioequivalent or biosimilar to the antibody adalimumab. In one instance, a biosimilar antibody is an antibody which shows no clinically meaningful difference when compared to a reference antibody, e.g., adalimumab. A biosimilar antibody has equivalent safety, purity, and potency as a reference antibody, e.g., adalimumab.

IV. Administration of the Formulations of the Invention For Treatment of TNFa-Related Disorders

[0179] An advantage of the formulations of the invention is that they may be used to deliver a high concentration of an anti-TNF alpha antibody, or antigen-binding portion, (e.g., adalimumab) to a subject subcutaneously such that either pain upon injection is decreased or the bioavailability of the antibody is improved. Thus, in one embodiment, the formulation of the invention is delivered to a subject subcutaneously. In one embodiment, the subject administers the formulation to himself/herself (self-administration).

[0180] In one embodiment, an effective amount of the formulation is administered. An example of an effective amount of the formulation is an amount sufficient to inhibit detrimental TNF-alpha activity or treat a disorder in which TNF alpha activity is detrimental.

[0181] As used herein, the term "a disorder in which TNF-alpha activity is detrimental" is intended to include diseases and other disorders in which the presence of TNF-alpha in a subject suffering from the disorder has been shown to be or is suspected of being either responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Accordingly, a disorder in which TNF-alpha activity is detrimental is a disorder in which inhibition of TNF-alpha activity is expected to alleviate the symptoms and/or progression of the disorder. Such disorders may be evidenced, for example, by an increase in the concentration of TNF-alpha in a biological fluid of a subject

suffering from the disorder (e.g., an increase in the concentration of TNF-alpha. in serum, plasma, synovial fluid, *etc.* of the subject), which can be detected, for example, using an anti-TNF-alpha. antibody.

[0182] In one embodiment, the effective amount of antibody may be determined according to a strictly weight based dosing scheme (e.g., mg/kg) or may be a total body dose also referred to as a fixed dose) which is independent of weight.

In one instance, an effective amount of the antibody is about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or about 100 mg of the human anti-TNF α antibody, or antigen-binding portion thereof. In one embodiment, an effective amount of the antibody is about 20-100, about 20-90, about 30-90, about 30-100, about 60-100, about 70-90, about 40-90, about 60-85 mg, or about 40-100 mg. In one embodiment, the formulation contains an effective amount of the antibody of 30 mg, 31 mg, 32 mg, 33 mg, 34 mg, 35 mg, 36 mg, 37 mg, 38 mg, 39 mg, 40 mg, 41 mg, 42 mg, 43 mg, 44 mg, 45 mg, 46 mg, 47 mg, 48 mg, 49 mg, 50 mg, 51 mg, 52 mg, 53 mg, 54 mg, 55 mg, 56 mg, 57 mg, 58 mg, 59 mg, 60 mg, 61 mg, 62 mg, 63 mg, 64 mg, 65 mg, 66 mg, 67 mg, 68 mg, 69 mg, 70 mg, 71 mg, 72 mg, 73 mg, 74 mg, 75 mg, 76 mg, 77 mg, 78 mg, 79 mg, 80 mg, 81 mg, 82 mg, 83 mg, 84 mg, 85 mg, 86 mg, 87 mg, 88 mg, 89 mg, 90 mg, 91 mg, 92 mg, 93 mg, 94 mg, 95 mg, 96 mg, 97 mg, 98 mg, 99 mg, or 100 mg of the antibody. Ranges including the aforementioned numbers are also included in the invention, e.g., 70-90 or 75-85 mg or 60-85 mg.

[0183] In one example, an effective amount of the formulation is 0.4 mL or 0.8 mL of the formulation containing a total body dose of about 80 mg of antibody (*i.e.*, 0.8 mL of a 100 mg/mL antibody formulation of the invention). In another example, an effective amount of the formulation is 0.4 mL of the formulation of the invention containing a total body dose of about 40 mg of antibody (*i.e.*, 0.4 mL of a 100 mg/mL antibody formulation of the invention). In yet another example, an effective amount of the formulation is twice 0.8 mL of the formulation containing a total body dose of about 160 mg of antibody (*i.e.*, two units containing 0.8 mL each of a 100 mg/mL antibody formulation of the invention). In a further example, an effective amount of the formulation is 0.2 mL of the formulation of the invention containing total body dose of about 20 mg of antibody (*i.e.*, 0.2 mL of a 100 mg/mL antibody formulation of the invention). Alternatively, an effective amount may be determined according to a weight-based fixed dosing regimen (see, e.g., WO 2008/154543).

[0184] In one embodiment, the TNF-alpha is human TNF-alpha and the subject is a human subject. Alternatively, the subject can be a mammal expressing a TNF-alpha with which an antibody of the invention cross-reacts. Still further the subject can be a mammal into which has been introduced hTNF-alpha (e.g., by administration of hTNF-alpha or by expression of an hTNF-alpha transgene).

[0185] A formulation of the invention may be administered to a human subject for therapeutic purposes (discussed further below). In one embodiment of the invention, the liquid pharmaceutical formulation is easily administrable, which includes, for example, a formulation which is self-administered by the patient. In one embodiment, the formulation of the invention is administered through subcutaneous injection, such as single use subcutaneous injection. Moreover, a formulation of the invention can be administered to a non-human mammal expressing a TNF-alpha with which the antibody cross-reacts (e.g., a primate, pig or mouse) for veterinary purposes or as an animal model of human disease. Regarding the latter, such animal models may be useful for evaluating the therapeutic efficacy of antibodies of the invention (e.g., testing of dosages and time courses of administration).

[0186] The formulations of the invention may be administered according to a certain dosing schedule. For example, the formulations may be administered according to a weekly, biweekly, or monthly dosing regimen. Alternatively, the formulation may be administered once every three weeks. In one instance, the formulations and methods comprise administration to the subject of a human anti-TNF α antibody according to a periodicity selected from the group consisting of weekly, biweekly, every three weeks, and monthly.

[0187] In one embodiment, the liquid aqueous formulation of the invention may be administered to a subject via, for example, a prefilled syringe, an autoinjector pen, or a needle-free administration device. Thus, the invention also features an autoinjector pen, a prefilled syringe, or a needle-free administration device comprising the liquid aqueous formulation of the invention. In one instance, the disclosure features a delivery device comprising a dose of the formulation comprising 100 mg/mL a human TNF alpha antibody, or antigen-binding portion thereof, e.g., an autoinjector pen or prefilled syringe comprises a dose of about 19 mg, 20 mg, 21 mg, 22 mg, 23 mg, 24 mg, 25 mg, 26 mg, 27 mg, 28 mg, 29 mg, 30 mg, 31 mg, 32 mg, 33 mg, 34 mg, 35 mg, 36 mg, 37 mg, 38 mg, 39 mg, 40 mg, 41 mg, 42 mg, 43 mg, 44 mg, 45 mg, 46 mg, 47 mg, 48 mg, 49 mg, 50 mg, 51 mg, 52 mg, 53 mg, 54 mg, 55 mg, 56 mg, 57 mg, 58 mg, 59 mg, 60 mg, 61 mg, 62 mg, 63 mg, 64 mg, 65 mg, 66 mg, 67 mg, 68 mg, 69 mg, 70 mg, 71 mg, 72 mg, 73 mg, 74 mg, 75 mg, 76 mg, 77 mg, 78 mg, 79 mg, 80 mg, 81 mg, 82 mg, 83 mg, 84 mg, 85 mg, 86 mg, 87 mg, 88 mg, 89 mg, 90 mg, 91 mg, 92 mg, 93 mg, 94 mg, 95 mg, 96 mg, 97 mg, 98 mg, 99 mg, 100 mg, 101 mg, 102 mg, 103 mg, 104 mg, 105 mg, *etc.* of the formulation. In one embodiment, the syringe or autoinjector contains 60-100 mg, 70-90 mg, or about 80 mg of the antibody.

[0188] In one embodiment, the formulations of the invention may be self administered using, e.g., a preloaded syringe or an automatic injection device. Automatic injection devices offer an alternative to manually-operated syringes for delivering therapeutic agents into patients' bodies and allowing patients to self-administer injections. Automatic injection devices are described, for example, in the following publications, WO 2008/005315, WO 2010/127146, WO 2006/000785, WO 2011/075524, WO 2005/113039, WO 2011/075524.

[0189] Accordingly, in one embodiment, the present disclosure provides pre-filled syringes or autoinjector devices

containing the formulations of the disclosure, as well as use of pre-filled syringes or autoinjector devices comprising the formulations described herein in the methods of the disclosure.

[0190] In one embodiment, the formulation of the invention is used to treat disorders in which TNF alpha activity is detrimental. As used herein, the term "a disorder in which TNF-alpha activity is detrimental" is intended to include diseases and other disorders in which the presence of TNF-alpha in a subject suffering from the disorder has been shown to be or is suspected of being either responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Accordingly, a disorder in which TNF-alpha activity is detrimental is a disorder in which inhibition of TNF-alpha activity is expected to alleviate the symptoms and/or progression of the disorder. Such disorders may be evidenced, for example, by an increase in the concentration of TNF-alpha in a biological fluid of a subject suffering from the disorder (e.g., an increase in the concentration of TNF-alpha in serum, plasma, synovial fluid, etc. of the subject), which can be detected, for example, using an anti-TNF-alpha antibody as described above.

[0191] There are numerous examples of disorders in which TNF-alpha activity is detrimental. Examples in which TNF-alpha activity is detrimental are also described in U.S. Patent Nos. 6,015,557; 6,177,077; 6,379,666; 6,419,934; 6,419,944; 6,423,321; 6,428,787; and 6,537,549; and PCT Publication Nos. WO 00/50079 and WO 01/49321. The formulations of the invention may also be used to treat disorders in which TNF alpha activity is detrimental as described in U.S. Pat. Nos. 6,090,382, 6,258,562 and U.S. Patent Application Publication No. US20040126372.

[0192] The use of the formulations of the invention in the treatment of specific exemplary disorders is discussed further below:

A. Sepsis

[0193] The formulations the invention may be used to treat subjects having sepsis. Tumor necrosis factor has an established role in the pathophysiology of sepsis, with biological effects that include hypotension, myocardial suppression, vascular leakage syndrome, organ necrosis, stimulation of the release of toxic secondary mediators and activation of the clotting cascade (see e.g., Tracey, K. J. and Cerami, A. (1994) *Annu. Rev. Med.* 45:491-503; Russell, D and Thompson, R. C. (1993) *Curr. Opin. Biotech.* 4:714-721). Accordingly, the formulation of the invention can be used to treat sepsis in any of its clinical settings, including septic shock, endotoxic shock, gram negative sepsis and toxic shock syndrome.

[0194] Furthermore, to treat sepsis, the formulation of the invention can be coadministered with one or more additional therapeutic agents that may further alleviate sepsis, such as an interleukin-1 inhibitor (such as those described in PCT Publication Nos. WO 92/16221 and WO 92/17583), the cytokine interleukin-6 (see e.g., PCT Publication No. WO 93/11793) or an antagonist of platelet activating factor (see e.g., European Patent Application Publication No. EP 374 510).

[0195] Additionally, in one embodiment, the formulation of the invention is administered to a human subject within a subgroup of sepsis patients having a serum or plasma concentration of 1L-6 above 500 pg/ml; or, in one embodiment, 1000 pg/ml, at the time of treatment (see PCT Publication No. WO 95/20978).

B. Autoimmune Diseases

[0196] The formulations of the invention may be used to treat subjects having an autoimmune disease. Tumor necrosis factor has been implicated in playing a role in the pathophysiology of a variety of autoimmune diseases. For example, TNF-alpha has been implicated in activating tissue inflammation and causing joint destruction in rheumatoid arthritis (see e.g., Tracey and Cerami, supra; Arend, W. P. and Dayer, J-M. (1995) *Arth. Rheum.* 38:151-160; Fava, R. A., et al. (1993) *Clin. Exp. Immunol.* 94:261-266). TNF-alpha also has been implicated in promoting the death of islet cells and in mediating insulin resistance in diabetes (see e.g., Tracey and Cerami, supra; PCT Publication No. WO 94/08609). TNF-alpha also has been implicated in mediating cytotoxicity to oligodendrocytes and induction of inflammatory plaques in multiple sclerosis (see e.g., Tracey and Cerami, supra). Also included in autoimmune diseases that may be treated using the formulations of the invention is juvenile idiopathic arthritis (JIA) (also referred to as juvenile rheumatoid arthritis) (see Grom et al. (1996) *Arthritis Rheum.* 39:1703; Mangge et al. (1995) *Arthritis Rheum.* 8:211).

[0197] The formulation of the invention can be used to treat autoimmune diseases, in particular those associated with inflammation, including rheumatoid arthritis, rheumatoid spondylitis (also referred to as ankylosing spondylitis), osteoarthritis and gouty arthritis, allergy, multiple sclerosis, autoimmune diabetes, autoimmune uveitis, juvenile idiopathic arthritis (also referred to as juvenile rheumatoid arthritis), and nephrotic syndrome.

C. Infectious Diseases

[0198] The formulations of the invention may be used to treat subjects having an infectious disease. Tumor necrosis factor has been implicated in mediating biological effects observed in a variety of infectious diseases. For example, TNF-alpha has been implicated in mediating brain inflammation and capillary thrombosis and infarction in malaria (see e.g.,

Tracey and Cerami, supra). TNF-alpha also has been implicated in mediating brain inflammation, inducing breakdown of the blood-brain barrier, triggering septic shock syndrome and activating venous infarction in meningitis (see e.g., Tracey and Cerami, supra). TNF-alpha also has been implicated in inducing cachexia, stimulating viral proliferation and mediating central nervous system injury in acquired immune deficiency syndrome (AIDS) (see e.g., Tracey and Cerami, supra): Accordingly, the antibodies, and antibody portions, of the disclosure, can be used in the treatment of infectious diseases, including bacterial meningitis (see e.g., European Patent Application Publication No. EP 585 705), cerebral malaria, AIDS and AIDS-related complex (ARC) (see e.g., European Patent Application Publication No. EP 230 574), as well as cytomegalovirus infection secondary to transplantation (see e.g., Fietze, E., et al. (1994) Transplantation 58:675-680). The formulation of the invention, also can be used to alleviate symptoms associated with infectious diseases, including fever and myalgias due to infection (such as influenza) and cachexia secondary to infection (e.g., secondary to AIDS or ARC).

D. Transplantation

[0199] The formulations of the invention may be used to treat subjects having a transplantation. Tumor necrosis factor has been implicated as a key mediator of allograft rejection and graft versus host disease (GVHD) and in mediating an adverse reaction that has been observed when the rat antibody OKT3, directed against the T cell receptor CD3 complex, is used to inhibit rejection of renal transplants (see e.g., Tracey and Cerami, supra; Eason, J. D., et al. (1995) Transplantation 59:300-305; Suthanthiran, M. and Strom, T. B. (1994) New Engl. J. Med. 331:365-375). Accordingly, the formulations of the invention can be used to inhibit transplant rejection, including rejections of allografts and xenografts and to inhibit GVHD. Although the antibody or antibody portion may be used alone, it can be used in combination with one or more other agents that inhibit the immune response against the allograft or inhibit GVHD. For example, in one embodiment, the formulations of the invention are used in combination with OKT3 to inhibit OKT3-induced reactions. In another embodiment, the formulation of the invention is used in combination with one or more antibodies directed at other targets involved in regulating immune responses, such as the cell surface molecules CD25 (interleukin-2 receptor-.alpha.), CD11a (LFA-1), CD54 (ICAM-1), CD4, CD45, CD28/CTLA4, CD80 (B7-1) and/or CD86 (B7-2). In yet another embodiment, the formulation of the invention is used in combination with one or more general immunosuppressive agents, such as cyclosporin A or FK506.

E. Malignancy

[0200] The formulations of the invention may be used to treat subjects having cancer or a malignant tumor. Tumor necrosis factor has been implicated in inducing cachexia, stimulating tumor growth, enhancing metastatic potential and mediating cytotoxicity in malignancies (see e.g., Tracey and Cerami, supra). Accordingly, the formulations of the invention can be used in the treatment of malignancies, to inhibit tumor growth or metastasis and/or to alleviate cachexia secondary to malignancy. The formulation of the invention may be administered systemically or locally to the tumor site.

F. Pulmonary Disorders

[0201] The formulations of the invention may be used to treat subjects having a pulmonary disease. Tumor necrosis factor has been implicated in the pathophysiology of adult respiratory distress syndrome, including stimulating leukocyte-endothelial activation, directing cytotoxicity to pneumocytes and inducing vascular leakage syndrome (see e.g., Tracey and Cerami, supra). Accordingly, the formulations of the invention can be used to treat various pulmonary disorders, including adult respiratory distress syndrome (see e.g., PCT Publication No. WO 91/04054), shock lung, chronic pulmonary inflammatory disease, pulmonary sarcoidosis, pulmonary fibrosis and silicosis. The formulation of the invention may be administered systemically or locally to the lung surface, for example as an aerosol.

G. Intestinal Disorders

[0202] The formulations of the invention may be used to treat subjects having an intestinal disorder. Tumor necrosis factor has been implicated in the pathophysiology of inflammatory bowel disorders (see e.g., Tracy, K. J., et al. (1986) Science 234:470-474; Sun, X-M., et al. (1988) J. Clin. Invest. 81:1328-1331; MacDonald, T. T., et al. (1990) Clin. Exp. Immunol. 81:301-305). Chimeric murine anti-hTNF-alpha antibodies have undergone clinical testing for treatment of Crohn's disease (van Dulleman, H. M., et al. (1995) Gastroenterology 109:129-135). The formulation of the invention, also can be used to treat intestinal disorders, such as idiopathic inflammatory bowel disease, which includes two syndromes, Crohn's disease and ulcerative colitis. In one embodiment, the formulation of the invention is used to treat Crohn's disease. In one embodiment, the formulation of the invention is used to treat ulcerative colitis.

H. Cardiac Disorders

[0203] The formulations of the invention, also can be used to treat various cardiac disorders, including ischemia of the heart (see e.g., European Patent Application, Publication No. EP 453 898) and heart insufficiency (weakness of the heart muscle)(see e.g., PCT Publication No. WO 94/20139).

I. Spondyloarthropathies

[0204] The formulations of the invention may also be used to treat subjects who have a spondyloarthropathy, including, for example, an axial spondyloarthropathy. $\text{TNF}\alpha$ has been implicated in the pathophysiology of a wide variety of disorders, including inflammatory diseases such as spondyloarthropathies (see e.g., Moeller, A., et al. (1990) Cytokine 2:162-169; U.S. Patent No. 5,231,024 to Moeller et al.; European Patent Publication No. 260 610 B1 by Moeller, A). In one embodiment, the spondyloarthropathy is an axial spondyloarthropathy. Other examples of spondyloarthropathies which can be treated with the $\text{TNF}\alpha$ antibody of the invention are described below:

1. Psoriatic arthritis

[0205] The formulations of the invention may also be used to treat subjects who have psoriatic arthritis. Tumor necrosis factor has been implicated in the pathophysiology of psoriatic arthritis (Parsch et al. (1998) Ann Rheum Dis. 57:691; Ritchlin et al. (1998) J Rheumatol. 25:1544). As referred to herein, psoriatic arthritis (PsA) or psoriasis associated with the skin, refers to chronic inflammatory arthritis which is associated with psoriasis. Psoriasis is a common chronic skin condition that causes red patches on the body. About 1 in 20 individuals with psoriasis will develop arthritis along with the skin condition, and in about 75% of cases, psoriasis precedes the arthritis. PsA exhibits itself in a variety of ways, ranging from mild to severe arthritis, wherein the arthritis usually affects the fingers and the spine. When the spine is affected, the symptoms are similar to those of ankylosing spondylitis, as described above.

[0206] PsA is sometimes associated with arthritis mutilans. Arthritis mutilans refers to a disorder which is characterized by excessive bone erosion resulting in a gross, erosive deformity which mutilates the joint. In one embodiment, formulations of the invention can be used to treat arthritis mutilans.

2. Reactive arthritis / Reiter's syndrome

[0207] The formulations of the invention may also be used to treat subjects who have Reiter's syndrome or reactive arthritis. Tumor necrosis factor has been implicated in the pathophysiology of reactive arthritis, which is also referred to as Reiter's syndrome (Braun et al. (1999) Arthritis Rheum. 42(10):2039). Reactive arthritis (ReA) refers to arthritis which complicates an infection elsewhere in the body, often following enteric or urogenital infections. ReA is often characterized by certain clinical symptoms, including inflammation of the joints (arthritis), urethritis, conjunctivitis, and lesions of the skin and mucous membranes. In addition, ReA can occur following infection with a sexually transmitted disease or dysenteric infection, including chlamydia, campylobacter, salmonella, or yersinia.

3. Undifferentiated spondyloarthropathies

[0208] The formulations of the invention may also be used to treat subjects who have an undifferentiated spondyloarthropathy (see Zeidler et al. (1992) Rheum Dis Clin North Am. 18:187). Other terms used to describe undifferentiated spondyloarthropathies include seronegative oligoarthritis and undifferentiated oligoarthritis. Undifferentiated spondyloarthropathies, as used herein, refers to a disorder wherein the subject demonstrates only some of the symptoms associated with a spondyloarthropathy. This condition is usually observed in young adults who do not have IBD, psoriasis, or the classic symptoms of AS or Reiter's syndrome. In some instances, undifferentiated spondyloarthropathies may be an early indication of AS.

J. Skin and Nail Disorders

[0209] In one embodiment, the formulations of the invention are used to treat a skin and/or a nail disorder. As used herein, the term "skin and nail disorder in which $\text{TNF}\alpha$ activity is detrimental" is intended to include skin and/or nail disorders and other disorders in which the presence of $\text{TNF}\alpha$ in a subject suffering from the disorder has been shown to be or is suspected of being either responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder, e.g., psoriasis. An example of a skin disorder which may be treated using the formulation of the invention is psoriasis. In one embodiment, the formulation of the invention is used to treat plaque psoriasis. Tumor necrosis factor has been implicated in the pathophysiology of psoriasis (Takematsu et al. (1989) Arch Dermatol Res.

281:398; Victor and Gottlieb (2002) J Drugs Dermatol. 1(3):264).

1. Psoriasis

[0210] The formulations of the invention may be used to treat subjects having psoriasis, including subjects having plaque psoriasis. Tumor necrosis factor has been implicated in the pathophysiology of psoriasis (Takematsu et al. (1989) Arch Dermatol Res. 281:398; Victor and Gottlieb (2002) J Drugs Dermatol. 1(3):264). Psoriasis is described as a skin inflammation (irritation and redness) characterized by frequent episodes of redness, itching, and thick, dry, silvery scales on the skin. In particular, lesions are formed which involve primary and secondary alterations in epidermal proliferation, inflammatory responses of the skin, and an expression of regulatory molecules such as lymphokines and inflammatory factors. Psoriatic skin is morphologically characterized by an increased turnover of epidermal cells, thickened epidermis, abnormal keratinization, inflammatory cell infiltrates into the epidermis and polymorphonuclear leukocyte and lymphocyte infiltration into the epidermis layer resulting in an increase in the basal cell cycle. Psoriasis often involves the nails, which frequently exhibit pitting, separation of the nail, thickening, and discoloration. Psoriasis is often associated with other inflammatory disorders, for example arthritis, including rheumatoid arthritis, inflammatory bowel disease (IBD), and Crohn's disease.

[0211] Evidence of psoriasis is most commonly seen on the trunk, elbows, knees, scalp, skin folds, or fingernails, but it may affect any or all parts of the skin. Normally, it takes about a month for new skin cells to move up from the lower layers to the surface. In psoriasis, this process takes only a few days, resulting in a build-up of dead skin cells and formation of thick scales. Symptoms of psoriasis include: skin patches, that are dry or red, covered with silvery scales, raised patches of skin, accompanied by red borders, that may crack and become painful, and that are usually located on the elbows, knees, trunk, scalp, and hands; skin lesions, including pustules, cracking of the skin, and skin redness; joint pain or aching which may be associated with arthritis, e.g., psoriatic arthritis.

[0212] Treatment for psoriasis often includes a topical corticosteroids, vitamin D analogs, and topical or oral retinoids, or combinations thereof. In one embodiment, the TNFalpha inhibitor of the invention is administered in combination with or the presence of one of these common treatments.

[0213] The diagnosis of psoriasis is usually based on the appearance of the skin. Additionally a skin biopsy, or scraping and culture of skin patches may be needed to rule out other skin disorders. An x-ray may be used to check for psoriatic arthritis if joint pain is present and persistent.

[0214] In one embodiment of the invention, a TNFalpha inhibitor is used to treat psoriasis, including chronic plaque psoriasis, guttate psoriasis, inverse psoriasis, pustular psoriasis, pemphigus vulgaris, erythrodermic psoriasis, psoriasis associated with inflammatory bowel disease (IBD), and psoriasis associated with rheumatoid arthritis (RA). Specific types of psoriasis included in the treatment methods of the invention are described in detail below:

a. Chronic Plaque Psoriasis

[0215] The formulations of the invention may be used to treat subjects having chronic plaque psoriasis. Tumor necrosis factor has been implicated in the pathophysiology of chronic plaque psoriasis (Asadullah et al. (1999) Br J Dermatol. 141:94). Chronic plaque psoriasis (also referred to as psoriasis vulgaris) is the most common form of psoriasis. Chronic plaque psoriasis is characterized by raised reddened patches of skin, ranging from coin-sized to much larger. In chronic plaque psoriasis, the plaques may be single or multiple, they may vary in size from a few millimeters to several centimeters. The plaques are usually red with a scaly surface, and reflect light when gently scratched, creating a "silvery" effect. Lesions (which are often symmetrical) from chronic plaque psoriasis occur all over body, but with predilection for extensor surfaces, including the knees, elbows, lumbosacral regions, scalp, and nails. Occasionally chronic plaque psoriasis can occur on the penis, vulva and flexures, but scaling is usually absent. Diagnosis of patients with chronic plaque psoriasis is usually based on the clinical features described above. In particular, the distribution, color and typical silvery scaling of the lesions in chronic plaque psoriasis are characteristic of chronic plaque psoriasis.

b. Guttate Psoriasis

[0216] The formulations of the invention may be used to treat subjects having guttate psoriasis. Guttate psoriasis refers to a form of psoriasis with characteristic water drop shaped scaly plaques. Flares of guttate psoriasis generally follow an infection, most notably a streptococcal throat infection. Diagnosis of guttate psoriasis is usually based on the appearance of the skin, and the fact that there is often a history of recent sore throat.

c. Inverse Psoriasis

[0217] The formulations of the invention may be used to treat subjects having inverse psoriasis. Inverse psoriasis is

a form of psoriasis in which the patient has smooth, usually moist areas of skin that are red and inflamed, which is unlike the scaling associated with plaque psoriasis. Inverse psoriasis is also referred to as intertriginous psoriasis or flexural psoriasis. Inverse psoriasis occurs mostly in the armpits, groin, under the breasts and in other skin folds around the genitals and buttocks, and, as a result of the locations of presentation, rubbing and sweating can irritate the affected areas.

d. Pustular Psoriasis

[0218] The formulations of the invention may be used to treat subjects having pustular psoriasis. Pustular psoriasis is a form of psoriasis that causes pus-filled blisters that vary in size and location, but often occur on the hands and feet. The blisters may be localized, or spread over large areas of the body. Pustular psoriasis can be both tender and painful, can cause fevers.

e. Other Psoriasis Disorders

[0219] Other examples of psoriatic disorders which can be treated with the formulations of the invention include erythrodermic psoriasis, vulgaris, psoriasis associated with IBD, and psoriasis associated with arthritis, including rheumatoid arthritis.

2. Pemphigus Vulgaris

[0220] The formulations of the invention may be used to treat subjects having pemphigus vulgaris. Pemphigus vulgaris is a serious autoimmune systemic dermatologic disease that often affects the oral mucous membrane and skin. The pathogenesis of pemphigus vulgaris is thought to be an autoimmune process that is directed at skin and oral mucous membrane desmosomes. Consequentially, cells do not adhere to each other. The disorder manifests as large fluid-filled, rupture-prone bullae, and has a distinctive histologic appearance. Anti-inflammatory agents are the only effective therapy for this disease which has a high mortality rate. Complications that arise in patients suffering from pemphigus vulgaris are intractable pain, interference with nutrition and fluid loss, and infections.

3. Atopic Dermatitis/Eczema

[0221] The formulations of the invention may be used to treat subjects having atopic dermatitis. Atopic dermatitis (also referred to as eczema) is a chronic skin disorder categorized by scaly and itching plaques. People with eczema often have a family history of allergic conditions like asthma, hay fever, or eczema. Atopic dermatitis is a hypersensitivity reaction (similar to an allergy) which occurs in the skin, causing chronic inflammation. The inflammation causes the skin to become itchy and scaly. Chronic irritation and scratching can cause the skin to thicken and become leathery-textured. Exposure to environmental irritants can worsen symptoms, as can dryness of the skin, exposure to water, temperature changes, and stress.

[0222] Subjects with atopic dermatitis can be identified by certain symptoms, which often include intense itching, blisters with oozing and crusting, skin redness or inflammation around the blisters, rash, dry, leathery skin areas, raw areas of the skin from scratching, and ear discharges/bleeding.

4. Sarcoidosis

[0223] The formulations of the invention may be used to treat subjects having sarcoidosis. Sarcoidosis is a disease in which granulomatous inflammation occurs in the lymph nodes, lungs, liver, eyes, skin, and/or other tissues. Sarcoidosis includes cutaneous sarcoidosis (sarcoidosis of the skin) and nodular sarcoidosis (sarcoidosis of the lymph nodes). Patients with sarcoidosis can be identified by the symptoms, which often include general discomfort, uneasiness, or an ill feeling; fever; skin lesions.

5. Erythema Nodosum

[0224] The formulations of the invention may be used to treat subjects having erythema nodosum. Erythema nodosum refers to an inflammatory disorder that is characterized by tender, red nodules under the skin, typically on the anterior lower legs. Lesions associated with erythema nodosum often begin as flat, but firm, hot red painful lumps (approximately an inch across). Within a few days the lesions may become purplish, and then over several weeks fade to a brownish flat patch.

[0225] In some instances, erythema nodosum may be associated with infections including, streptococcus, coccidioidomycosis, tuberculosis, hepatitis B, syphilis, cat scratch disease, tularemia, yersinia, leptospirosis psittacosis, histo-

plasmosis, mononucleosis (EBV). In other instances, erythema nodosum may be associated with sensitivity to certain medications including, oral contraceptives, penicillin, sulfonamides, sulfones, barbiturates, hydantoin, phenacetin, salicylates, iodides, and progestin. Erythema nodosum is often associated with other disorders including, leukemia, sarcoidosis, rheumatic fever, and ulcerative colitis.

[0226] Symptoms of erythema nodosum usually present themselves on the shins, but lesions may also occur on other areas of the body, including the buttocks, calves, ankles, thighs and upper extremities. Other symptoms in subjects with erythema nodosum can include fever and malaise.

6. Hidradenitis Suppurativa

[0227] The formulations of the invention may be used to treat subjects having hidradenitis suppurativa. Hidradenitis suppurativa refers to a skin disorder in which swollen, painful, inflamed lesions or lumps develop in the groin and sometimes under the arms and under the breasts. Hidradenitis suppurativa occurs when apocrine gland outlets become blocked by perspiration or are unable to drain normally because of incomplete gland development. Secretions trapped in the glands force perspiration and bacteria into surrounding tissue, causing subcutaneous induration, inflammation, and infection. Hidradenitis suppurativa is confined to areas of the body that contain apocrine glands. These areas are the axillae, areola of the nipple, groin, perineum, circumanal, and periumbilical regions.

7. Lichen Planus

[0228] The formulations of the invention may be used to treat subjects having lichen planus. Tumor necrosis factor has been implicated in the pathophysiology of lichen planus (Sklavounou et al. (2000) J Oral Pathol Med. 29:370). Lichen planus refers to a disorder of the skin and the mucous membranes resulting in inflammation, itching, and distinctive skin lesions. Lichen planus may be associated with hepatitis C or certain medications.

8. Sweet's Syndrome

[0229] The formulations of the invention may be used to treat subjects having Sweet's syndrome. Inflammatory cytokines, including tumor necrosis factor, have been implicated in the pathophysiology of Sweet's syndrome (Reuss-Borst et al. (1993) Br J Haematol. 84:356). Sweet's syndrome, which was described by R. D. Sweet in 1964, is characterized by the sudden onset of fever, leukocytosis, and cutaneous eruption. The eruption consists of tender, erythematous, well-demarcated papules and plaques which show dense neutrophilic infiltrates microscopically. The lesions may appear anywhere, but favor the upper body including the face. The individual lesions are often described as pseudovesicular or pseudopustular, but may be frankly pustular, bullous, or ulcerative. Oral and eye involvement (conjunctivitis or episcleritis) have also been frequently reported in patients with Sweet's syndrome. Leukemia has also been associated with Sweet's syndrome.

9. Vitiligo

[0230] The formulations of the invention may be used to treat subjects having vitiligo. Vitiligo refers to a skin condition in which there is loss of pigment from areas of skin resulting in irregular white patches with normal skin texture. Lesions characteristic of vitiligo appear as flat depigmented areas. The edges of the lesions are sharply defined but irregular. Frequently affected areas in subjects with vitiligo include the face, elbows and knees, hands and feet, and genitalia.

10. Scleroderma

[0231] The formulations of the invention may be used to treat subjects having scleroderma. Tumor necrosis factor has been implicated in the pathophysiology of scleroderma (Tutuncu Z et al. (2002) Clin Exp Rheumatol. 20(6 Suppl 28):S 146-51; Mackiewicz Z et al. (2003) Clin Exp Rheumatol. 21(1):41-8; Murota H et al. (2003) Arthritis Rheum. 48(4):1117-25). Scleroderma refers to a diffuse connective tissue disease characterized by changes in the skin, blood vessels, skeletal muscles, and internal organs. Scleroderma is also referred to as CREST syndrome or Progressive systemic sclerosis, and usually affects people between the ages 30-50. Women are affected more often than men.

[0232] The cause of scleroderma is unknown. The disease may produce local or systemic symptoms. The course and severity of the disease varies widely in those affected. Excess collagen deposits in the skin and other organs produce the symptoms. Damage to small blood vessels within the skin and affected organs also occurs. In the skin, ulceration, calcification, and changes in pigmentation may occur. Systemic features may include fibrosis and degeneration of the heart, lungs, kidneys and gastrointestinal tract.

[0233] Patients suffering from scleroderma exhibit certain clinical features, including, blanching, blueness, or redness

of fingers and toes in response to heat and cold (Raynaud's phenomenon), pain, stiffness, and swelling of fingers and joints, skin thickening and shiny hands and forearm, esophageal reflux or heartburn, difficulty swallowing, and shortness of breath. Other clinical symptoms used to diagnose scleroderma include, an elevated erythrocyte sedimentation rate (ESR), an elevated rheumatoid factor (RF), a positive antinuclear antibody test, urinalysis that shows protein and microscopic blood, a chest X-ray that may show fibrosis, and pulmonary function studies that show restrictive lung disease.

11. Nail Disorders

[0234] The formulations of the invention may be used to treat subjects having a nail disorder. Nail disorders include any abnormality of the nail. Specific nail disorders include, but are not limited to, pitting, koilonychia, Beau's lines, spoon nails, onycholysis, yellow nails, pterygium (seen in lichen planus), and leukonychia. Pitting is characterized by the presence of small depressions on the nail surface. Ridges or linear elevations can develop along the nail occurring in a "lengthwise" or "crosswise" direction. Beau's lines are linear depressions that occur "crosswise" (transverse) in the fingernail. Leukonychia describes white streaks or spots on the nails. Koilonychia is an abnormal shape of the fingernail where the nail has raised ridges and is thin and concave. Koilonychia is often associated with iron deficiency.

[0235] Nail disorders which can be treated with the TNF α antibody of the invention also include psoriatic nails. Psoriatic nails include changes in nails which are attributable to psoriasis. In some instances psoriasis may occur only in the nails and nowhere else on the body. Psoriatic changes in nails range from mild to severe, generally reflecting the extent of psoriatic involvement of the nail plate, nail matrix, i.e., tissue from which the nail grows, nail bed, i.e., tissue under the nail, and skin at the base of the nail. Damage to the nail bed by the pustular type of psoriasis can result in loss of the nail. Nail changes in psoriasis fall into general categories that may occur singly or all together. In one category of psoriatic nails, the nail plate is deeply pitted, probably due to defects in nail growth caused by psoriasis. In another category, the nail has a yellow to yellow-pink discoloration, probably due to psoriatic involvement of the nail bed. A third subtype of psoriatic nails are characterized by white areas which appear under the nail plate. The white areas are actually air bubbles marking spots where the nail plate is becoming detached from the nail bed. There may also be reddened skin around the nail. A fourth category is evidenced by the nail plate crumbling in yellowish patches, i.e., onychodystrophy, probably due to psoriatic involvement in the nail matrix. A fifth category is characterized by the loss of the nail in its entirety due to psoriatic involvement of the nail matrix and nail bed.

[0236] The formulations of the invention may also be used to treat nail disorders often associated with lichen planus. Nails in subjects with lichen planus often show thinning and surface roughness of the nail plate with longitudinal ridges or pterygium.

[0237] The formulations of the invention may be used to treat nail disorders, such as those described herein. Often nail disorders are associated with skin disorders. In one instance, the disclosure includes a method of treatment for nail disorders with a TNF α antibody. In an embodiment, the nail disorder is associated with another disorder, including a skin disorder such as psoriasis. In another embodiment, the disorder associated with a nail disorder is arthritis, including psoriatic arthritis.

[0238] 12. Other Skin and Nail Disorders

[0239] The formulations of the invention may be used to treat other skin and nail disorders, such as chronic actinic dermatitis, bullous pemphigoid, and alopecia areata. Chronic actinic dermatitis (CAD) is also referred to as photosensitivity dermatitis/actinic reticuloid syndrome (PD/AR). CAD is a condition in which the skin becomes inflamed, particularly in areas that have been exposed to sunlight or artificial light. Commonly, CAD patients have allergies to certain substances that come into contact with their skin, particularly various flowers, woods, perfumes, sunscreens and rubber compounds. Bullous pemphigoid refers to a skin disorder characterized by the formation of large blisters on the trunk and extremities. Alopecia areata refers to hair loss characterized by round patches of complete baldness in the scalp or beard.

K. Metabolic Disorders

[0240] The formulations of the invention may be used to treat a metabolic disease. TNF α has been implicated in the pathophysiology of a wide variety of disorders, including metabolic disorders, such as diabetes and obesity (Spiegelman and Hotamisligil (1993) Cell 73:625; Chu et al. (2000) Int J Obes Relat Metab Disord. 24:1085; Ishii et al. (2000) Metabolism. 49:1616).

[0241] Metabolic disorders affect how the body processes substances needed to carry out physiological functions. A number of metabolic disorders share certain characteristics, i.e. they are associated with the insulin resistance, lack of ability to regulate blood sugar, weight gain, and increase in body mass index. Examples of metabolic disorders include diabetes and obesity. Examples of diabetes include type 1 diabetes mellitus, type 2 diabetes mellitus, diabetic neuropathy, peripheral neuropathy, diabetic retinopathy, diabetic ulcerations, retinopathy ulcerations, diabetic macrovasculopathy, and obesity. Examples of metabolic disorders which can be treated with the formulations of the invention are described in more detail below:

1. Diabetes

[0242] The formulations of the invention may be used to treat diabetes. Tumor necrosis factor has been implicated in the pathophysiology of diabetes. (see e.g., Navarro J.F., Mora C., Maca, Am J Kidney Dis. 2003 Jul;42(1):53-61; Daimon M et al., Diabetes Care. 2003 Jul;26(7):2015-20; Zhang M et. al., J Tongji Med Univ. 1999;19(3):203-5, Barbieri M et al., Am J Hypertens. 2003 Jul;16(7):537-43.) For example, $\text{TNF}\alpha$ is implicated in the pathophysiology for insulin resistance. It has been found that serum TNF levels in patients with gastrointestinal cancer correlates with insulin resistance (see e.g., McCall, J. et al. Br. J. Surg. 1992; 79: 1361-3).

[0243] Diabetes includes the two most common types of the disorder, namely type I diabetes and type II diabetes, which both result from the body's inability to regulate insulin. Insulin is a hormone released by the pancreas in response to increased levels of blood sugar (glucose) in the blood.

[0244] The term "type 1 diabetes," as used herein, refers to a chronic disease that occurs when the pancreas produces too little insulin to regulate blood sugar levels appropriately. Type 1 diabetes is also referred to as insulin-dependent diabetes mellitus, IDDM, juvenile onset diabetes, and diabetes - type I. Type 1 diabetes represents is the result of a progressive autoimmune destruction of the pancreatic β -cells with subsequent insulin deficiency.

[0245] The term "type 2 diabetes," refers to a chronic disease that occurs when the pancreas does not make enough insulin to keep blood glucose levels normal, often because the body does not respond well to the insulin. Type 2 diabetes is also referred to as noninsulin-dependent diabetes mellitus, NDDM, and diabetes - type II

[0246] Diabetes is can be diagnosed by the administration. of a glucose tolerance test. Clinically, diabetes is often divided into several basic categories. Primary examples of these categories include, autoimmune diabetes mellitus, non-insulin-dependent diabetes mellitus (type 1 NDDM), insulin-dependant diabetes mellitus (type 2 IDDM), non-autoimmune diabetes mellitus, non-insulin-dependant diabetes mellitus (type 2 NIDDM), and maturity-onset diabetes of the young (MODY). A further category, often referred to as secondary, refers to diabetes brought about by some identifiable condition which causes or allows a diabetic syndrome to develop. Examples of secondary categories include, diabetes caused by pancreatic disease, hormonal abnormalities, drug-or chemical-induced diabetes, diabetes caused by insulin receptor abnormalities, diabetes associated with genetic syndromes, and diabetes of other causes. (see e.g., Harrison's (1996) 14thed., New York, McGraw-Hill).

[0247] Diabetes manifests itself in the foregoing categories and can cause several complications that are discussed in the following sections. Accordingly, the antibody, or antigen-binding fragment thereof, of the disclosure can be used to treat diabetes. In one instance, the $\text{TNF}\alpha$ antibody, or antigen-binding fragment thereof, of the disclosure is used to treat diabetes associated with the above identified catagores.

[0248] Diabetes is often treated with diet, insulin dosages, and various medications described herein. Accordingly, the formulations of the invention may also be administered in combination with agents commonly used to treat metabolic disorders and pain commonly associated with diabetes.

[0249] Diabetes manifests itself in many complications and conditions associated with diabetes, including the following catagories:

a. Diabetic Neuropathy and Peripheral Neuropathy

[0250] The formulations of the invention may be used to treat diabetic neuropathy or peripheral neuropathy. Tumor necrosis factor has been implicated in the pathophysiology of diabetic neuropathy and peripheral neuropathy. (See Benjafeld et al. (2001) Diabetes Care. 24:753; Qiang, X. et al. (1998) Diabetologia.41:1321-6; Pfeiffer et al (1997) Horm Metab Res. 29:111).

[0251] The term "neuropathy," also referred to as nerve damage-diabetic, as used herein, refers to a common complication of diabetes in which nerves are damaged as a result of hyperglycemia (high blood sugar levels). A variety of diabetic neuropathies are recognized, such as distal sensorimotor polyneuropathy, focal motor neuropathy, and autonomic neuropathy.

[0252] The term "peripheral neuropathy," also known as peripheral neuritis and diabetic neuropathy, as used herein, refers to the failure of the nerves to carry information to and from the brain and spinal cord. Peripheral neuropathy produces symptoms such as pain, loss of sensation, and the inability to control muscles. In some cases, the failure of nerves to control blood vessels, intestinal function, and other organs results in abnormal blood pressure, digestion, and loss of other basic involuntary processes. Peripheral neuropathy may involve damage to a single nerve or nerve group (mononeuropathy) or may affect multiple nerves (polyneuropathy).

[0253] Neuropathies that affect small myelinated and unmyelinated fibers of the sympathetic and parasympathetic nerves are known as "peripheral neuropathies." Furthermore, the related disorder of peripheral neuropathy, also known as peripheral neuritis and diabetic neuropathy, refers to the failure of the nerves to carry information to and from the brain and spinal cord. This produces symptoms such as pain, loss of sensation, and the inability to control muscles. In some cases, failure of nerves controlling blood vessels, intestinal function, and other organs results in abnormal blood

pressure, digestion, and loss of other basic involuntary processes. Peripheral neuropathy may involve damage to a single nerve or nerve group (mononeuropathy) or may affect multiple nerves (polyneuropathy).

[0254] The term "diabetic neuropathy" refers to a common complication of diabetes in which nerves are damaged as a result of hyperglycemia (high blood sugar levels). Diabetic neuropathy is also referred to as neuropathy and nerve damage-diabetic. A variety of diabetic neuropathies are recognized, such as distal sensorimotor polyneuropathy, focal motor neuropathy, and autonomic neuropathy.

b. Diabetic Retinopathy

[0255] The formulations of the invention may be used to treat diabetic retinopathy. Tumor necrosis factor has been implicated in the pathophysiology of diabetic retinopathy (Scholz et al. (2003) Trends Microbiol. 11:171). The term "diabetic retinopathy" as used herein, refers to progressive damage to the eye's retina caused by long-term diabetes. Diabetic retinopathy, includes proliferative retinopathy. Proliferative neuropathy in turn includes includes neovascularization, peritinal hemorrhage and retinal detachment.

[0256] In advanced retinopathy, small vessels proliferate on the surface of the retina. These blood vessels are fragile, tend to bleed and can cause peritinal hemorrhages. The hemorrhage can obscure vision, and as the hemorrhage is resorbed fibrous tissue forms predisposing to retinal detachments and loss of vision. In addition, diabetic retinopathy includes proliferative retinopathy which includes neovascularization, peritinal hemorrhage and retinal detachment. Daibetic retinopathy also includes "background retinopathy" which involves changes occurring with the layers of the retina.

c. Diabetic Ulcerations and Retinopathy Ulceration

[0257] The formulations of the invention may be used to treat diabetic ulcerations or retinopathy ulcerations. Tumor necrosis factor has been implicated in the pathophysiology of diabetic ulcerations, (see Lee et al. (2003) Hum Immunol. 64:614; Navarro et al. (2003) Am J Kidney Dis. 42:53; Daimon et al (2003) Diabetes Care. 26:2015; Zhang et al: (1999) J Tongji Med Univ. 19:203; Barbieri et al. (2003) Am J Hypertens. 16:537; Venn et al. (1993) Arthritis Rheum. 36:819; Westacott et al. (1994) J Rheumatol. 21:1710).

[0258] The term "diabetic ulcerations," as used herein, refers to an ulcer which results as a complication of diabetes. An ulcer is a crater-like lesion on the skin or mucous membrane caused by an inflammatory, infectious, malignant condition, or metabolic disorder. Typically diabetic ulcers can be found on limbs and extremities, more typically the feet. These ulcers, caused by diabetic conditions, such as neurapthy and a vacualr insufficiency, can lead to ischemia and poor wound healing. More, extensive ulcerations may progress to osteomyelitis. Once osteomyelitis develops, it may be difficile to eradicate with antibiotics alonda nd amputation maybe necessary..

[0259] The term "retinopathy ulcerations," as used herein refers to an ulcer which causes or results in damages to the eye and the eye's retina. Retinopathy ulcerations may include conditions such has retinoathic hemorrhages.

d. Diabetic Macrovasculopathy

[0260] The formulations of the invention may be used to treat diabetic macrovasculopathy. Tumor necrosis factor has been implicated in the pathophysiology of diabetic macrovasculopathy (Devaraj et al. (2000) Circulation. 102:191; Hattori Y et al. (2000) Cardioasc Res. 46:188; Clausell N et al. (1999) Cardioasc Pathol.8:145). The term "diabetic macrovasculopathy;" also referred to as "macrovascular disease," as used herein, refers to a disease of the blood vessels that results from diabetes. Diabetic macrovasculopathy complication occurs when, for example, fat and blood clots build up in the large blood vessels and stick to the vessel walls. Diabetic macrovasculopathies. include diseases such as coronary disease, cerebrovascular disease, and peripheral vascular disease, hyperglycaemia and cardiovascular disease, and strokes.

2. Obesity

[0261] The formulations of the invention may be used to treat obesity. Tumor necrosis factor has been implicated in the pathophysiology of obesity (see e.g., Pihlajamaki J et al. (2003) Obes Res.11:912; Barbieri et al. (2003) Am J Hypertens. 16:537; Tsuda et al. (2003) J Nutr. 133:2125). Obesity increases a person's risk of illness and death due to diabetes, stroke, coronary artery disease, hypertension, high cholesterol, and kidney and gallbladder disorders. Obesity may also increase the risk for some types of cancer, and may be a risk factor for the development of osteoarthritis and sleep apnea. Obesity can be treated with the antibody of the invention alone or in combination with other metabolic disorders, including diabetes.

L. Vasculitides

[0262] The formulations of the invention may be used to treat a subject having a vasculitis. $\text{TNF}\alpha$ has been implicated in the pathophysiology of a variety of vasculitides, (see e.g., Deguchi et al. (1989) *Lancet*. 2:745). As used herein, the term "a vasculitis in which $\text{TNF}\alpha$ activity is detrimental" is intended to include vasculitis in which the presence of $\text{TNF}\alpha$ in a subject suffering from the disorder has been shown to be or is suspected of being either responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Such disorders may be evidenced, for example, by an increase in the concentration of $\text{TNF}\alpha$ in a biological fluid of a subject suffering from the disorder (e.g., an increase in the concentration of $\text{TNF}\alpha$ in serum, plasma, synovial fluid, etc. of the subject), which can be detected, for example, using an anti- $\text{TNF}\alpha$ antibody as described above.

[0263] There are numerous examples of vasculitides in which $\text{TNF}\alpha$ activity is detrimental, including Behcet's disease. The use of the formulations of the invention in the treatment of specific vasculitides are discussed further below. In certain instances, the antibody, or antibody portion, is administered to the subject in combination with another therapeutic agent, as described below

[0264] The formulations of the invention be used to treat vasculitis in which $\text{TNF}\alpha$ activity is detrimental, wherein inhibition of $\text{TNF}\alpha$ activity is expected to alleviate the symptoms and/or progression of the vasculitis or to prevent the vasculitis. Subjects suffering from or at risk of developing vasculitis can be identified through clinical symptoms and tests. For example, subjects with vasculitides often develop antibodies to certain proteins in the cytoplasm of neutrophils, antineutrophil cytoplasmic antibodies (ANCA). Thus, in some instances, vasculitides may be evidenced by tests (e.g., ELISA), which measure ANCA presence.

[0265] Vasculitis and its consequences may be the sole manifestation of disease or it may be a secondary component of another primary disease. Vasculitis may be confined to a single organ or it may simultaneously affect several organs, and depending on the syndrome, arteries and veins of all sizes can be affected. Vasculitis can affect any organ in the body.

[0266] In vasculitis, the vessel lumen is usually compromised, which is associated with ischemia of the tissues supplied by the involved vessel. The broad range of disorders that may result from this process is due to the fact that any type, size and location of vessel (e.g., artery, vein, arteriole, venule, capillary) can be involved. Vasculitides are generally classified according to the size of the affected vessels, as described below. It should be noted that some small and large vessel vasculitides may involve medium-sized arteries; but large and medium-sized vessel vasculitides do not involve vessels smaller than arteries. Large vessel disease includes, but is not limited to, giant cell arteritis, also known as temporal arteritis or cranial arteritis, polymyalgia rheumatica, and Takayasu's disease or arteritis, which is also known as aortic arch syndrome, young female arteritis and Pulseless disease. Medium vessel disease includes, but is not limited to, classic polyarteritis nodosa and Kawasaki's disease, also known as mucocutaneous lymph node syndrome. Non-limiting examples of small vessel disease are Behcet's Syndrome, Wegner's granulomatosis, microscopic polyangiitis, hypersensitivity vasculitis, also known as cutaneous vasculitis, small vessel vasculitis, Henoch-Schonlein purpura, allergic granulomatosis and vasculitis, also known as Churg Strauss syndrome. Other vasculitides include, but are not limited to, isolated central nervous system vasculitis, and thromboangitis obliterans, also known as Buerger's disease. Classic Polyarteritis nodosa (PAN), microscopic PAN, and allergic granulomatosis are also often grouped together and are called the systemic necrotizing vasculitides. A further description of vasculitis is described below:

1. Large vessel vasculitis

[0267] In one embodiment, the formulations of the invention are used to treat subjects who have large vessel vasculitis. The term "large vessel(s)" as used herein, refers to the aorta and the largest branches directed toward major body regions. Large vessels include, for example, the aorta, and its branches and corresponding veins, e.g., the subclavian artery; the brachiocephalic artery; the common carotid artery; the innominate vein; internal and external jugular veins; the pulmonary arteries and veins; the venae cavae; the renal arteries and veins; the femoral arteries and veins; and the carotid arteries. Examples of large vessel vasculitides are described below.

a. Giant cell arteritis (GCA)

[0268] The formulations of the invention may be used to treat giant cell arteritis. Tumor necrosis factor has been implicated in the pathophysiology of giant cell arteritis (Sneller, M.C. (2002) *Cleve. Clin. J. Med.* 69:SI140-3; Schett, G., et al. (2002) *Ann. Rheum. Dis.* 61:463). Giant cell arteritis (GCA), refers to a vasculitis involving inflammation and damage to blood vessels, particularly the large or medium arteries that branch from the external carotid artery of the neck. GCA is also referred to as temporal arteritis or cranial arteritis, and is the most common primary vasculitis in the elderly. It almost exclusively affects individuals over 50 years of age, however, there are well-documented cases of patients 40 years and younger. GCA usually affects extracranial arteries. GCA can affect the branches of the carotid arteries, including the temporal artery. GCA is also a systemic disease which can involve arteries in multiple locations.

[0269] Histopathologically, GCA is a panarteritis with inflammatory mononuclear cell infiltrates within the vessel wall with frequent Langhans type giant cell formation. There is proliferation of the intima, granulomatous inflammation and fragmentation of the internal elastic lamina. The pathological findings in organs is the result of ischemia related to the involved vessels.

[0270] Patients suffering from GCA exhibit certain clinical symptoms, including fever, headache, anemia and high erythrocyte sedimentation rate (ESR). Other typical indications of GCA include jaw or tongue claudication, scalp tenderness, constitutional symptoms, pale optic disc edema (particularly 'chalky white' disc edema), and vision disturbances. The diagnosis is confirmed by temporal artery biopsy.

b. Polymyalgia rheumatica

[0271] The formulations of the invention may be used to treat polymyalgia rheumatica. Tumor necrosis factor has been implicated in the pathophysiology of polymyalgia rheumatica (Straub, R.H., et al. (2002) *Rheumatology* (Oxford) 41:423; Uddhammar, A., et al. (1998) *Br. J. Rheumatol.* 37:766). Polymyalgia rheumatica refers to a rheumatic disorder that is associated with moderate to severe muscle pain and stiffness in the neck, shoulder, and hip, most noticeable in the morning. IL-6 and IL-1 β expression has also been detected in a majority of the circulating monocytes in patients with the polymyalgia rheumatica. Polymyalgia rheumatica may occur independently, or it may coexist with or precede GCA, which is an inflammation of blood vessels.

c. Takayasu's Arteritis

[0272] The formulations of the invention may be used to treat Takayasu's arteritis. Tumor necrosis factor has been implicated in the pathophysiology of Takayasu's arteritis (Kobayashi, Y. and Numano, F. (2002) *Intern. Med.* 41:44; Fraga, A. and Medina F. (2002) *Curr. Rheumatol. Rep.* 4:30). Takayasu's arteritis refers to a vasculitis characterized by an inflammation of the aorta and its major branches. Takayasu's arteritis (also known as Aortic arch syndrome, young female arteritis and Pulseless disease) affects the thoracic and abdominal aorta and its main branches or the pulmonary arteries. Fibrotic thickening of the aortic wall and its branches (e.g., carotid, innominate, and subclavian arteries) can lead to reduction of lumen size of vessels that arise from the aortic arch. This condition also typically affects the renal arteries.

[0273] Takayasu's arteritis primarily affects young women, usually aged 20-40 years old, particularly of Asian descent, and may be manifested by malaise, arthralgias and the gradual onset of extremity claudication. Most patients have asymmetrically reduced pulses, usually along with a blood pressure differential in the arms. Coronary and/or renal artery stenosis may occur.

[0274] The clinical features of Takayasu's arteritis may be divided into the features of the early inflammatory disease and the features of the later disease. The clinical features of the early inflammatory stage of Takayasu's disease are: malaise, low grade fever, weight loss, myalgia, arthralgia, and erythema multiforme. Later stages of Takayasu's disease are characterised by fibrotic stenosis of arteries and thrombosis. The main resulting clinical features are ischaemic phenomena, e.g. weak and asymmetrical arterial pulses, blood pressure discrepancy between the arms, visual disturbance, e.g. scotomata and hemianopia, other neurological features including vertigo and syncope, hemiparesis or stroke. The clinical features result from ischaemia due to arterial stenosis and thrombosis.

2. Medium Vessel Disease

[0275] The formulations of the invention may be used to treat subjects who have medium vessel vasculitis. The term "medium vessel(s)" is used to refer to those blood vessels which are the main visceral arteries. Examples of medium vessels include the mesenteric arteries and veins, the iliac arteries and veins, and the maxillary arteries and veins. Examples of medium vessel vasculitides are described below.

a. Polyarteritis Nodosa

[0276] The formulations of the invention may be used to treat polyarteritis nodosa. Tumor necrosis factor has been implicated in the pathophysiology of polyarteritis nodosa (DiGirolamo, N., et al. (1997) *J. Leukoc. Biol.* 61:667). Polyarteritis nodosa, or periarteritis nodosa refers to vasculitis which is a serious blood vessel disease in which small and medium-sized arteries become swollen and damaged because they are attacked by rogue immune cells. Polyarteritis nodosa usually affects adults more frequently than children. It damages the tissues supplied by the affected arteries because they don't receive enough oxygen and nourishment without a proper blood supply.

[0277] Symptoms which are exhibited in patients with polyarteritis nodosa generally result from damage to affected organs, often the skin, heart, kidneys, and nervous system. Generalized symptoms of polyarteritis nodosa include fever, fatigue, weakness, loss of appetite, and weight loss. Muscle aches (myalgia) and joint aches (arthralgia) are common.

The skin of subjects with polyarteritis nodosa may also show rashes, swelling, ulcers, and lumps (nodular lesions).

[02778] Classic PAN (polyarteritis nodosa) is a systemic arteritis of small to medium muscular arteritis in which involvement of renal and visceral arteries is common. Abdominal vessels have aneurysms or occlusions in 50% of PAN patients. Classic PAN does not involve the pulmonary arteries although the bronchial vessels may be involved. Granulomas, significant eosinophilia and an allergic diathesis are not part of the syndrome. Although any organ system may be involved, the most common manifestations include peripheral neuropathy, mononeuritis multiplex, intestinal ischemia, renal ischemia, testicular pain and livedo reticularis.

b. Kawasaki's Disease

[02799] The formulations of the invention may be used to treat Kawasaki's disease. Tumor necrosis factor has been implicated in the pathophysiology of Kawasaki's disease (Sundel, R.P. (2002) Curr. Rheumatol. Rep. 4:474; Gedalia, A.(2002) Curr. Rheumatol. Rep. 4:25). Although the cause of Kawasaki's disease is unknown, it is associated with acute inflammation of the coronary arteries, suggesting that the tissue damage associated with this disease may be mediated by proinflammatory agents such as $TNF\alpha$. Kawasaki's disease refers to a vasculitis that affects the mucus membranes, lymph nodes, lining of the blood vessels, and the heart. Kawasaki's disease is also often referred to as mucocutaneous lymph node syndrome, mucocutaneous lymph node disease, and infantile polyarteritis. Subjects afflicted with Kawasaki's disease develop vasculitis often involving the coronary arteries which can lead to myocarditis and pericarditis. Often as the acute inflammation diminishes, the coronary arteries may develop aneurysm, thrombosis, and lead to myocardial infarction.

[02800] Kawasaki's disease is a febrile systemic vasculitis associated with edema in the palms and the soles of the feet, with enlargement of cervical lymph nodes, cracked lips and "strawberry tongue". Although the inflammatory response is found in vessels throughout the body, the most common site of end-organ damage is the coronary arteries. Kawasaki's Disease predominantly affects children under the age of 5. The highest incidence is in Japan but is becoming increasingly recognized in the West and is now the leading cause of acquired heart disease in US children. The most serious complication of Kawasaki disease is coronary arteritis and aneurysm formation that occurs in a third of untreated patients.

3. Small vessel disease

[02811] The formulations of the invention may be used to treat small vessel disease. In one embodiment, the $TNF\alpha$ antibody of the invention is used to treat subjects who have small vessel vasculitis. The term "small vessel(s)" is used to refer to arterioles, venules and capillaries. Arterioles are arteries that contain only 1 or 2 layers of smooth muscle cells and are terminal to and continuous with the capillary network. Venules carry blood from the capillary network to veins and capillaries connect arterioles and venules. Examples of small vessel vasculitides are described below.

a. Behcet's Disease

[02821] The formulations of the invention may be used to treat Behcet's disease. Tumor necrosis factor has been implicated in the pathophysiology of Behcet's disease (Sfikakis, P.P. (2002) Ann. Rheum. Dis. 61:ii51-3; Dogan, D. and Farah, C. (2002) Oftalmologia. 52:23). Behcet's disease is a chronic disorder that involves inflammation of blood vessels throughout the body. Behcet's disease may also cause various types of skin lesions, arthritis, bowel inflammation, and meningitis (inflammation of the membranes of the brain and spinal cord). As a result of Behcet's disease, the subject with the disorder may have inflammation in tissues and organs throughout the body, including the gastrointestinal tract, central nervous system, vascular system, lungs, and kidneys. Behcet's disease is three times more common in males than females and is more common in the east Mediterranean and Japan.

b. Wegener's granulomatosis

[02831] The formulations of the invention may be used to treat Wegener's granulomatosis. Tumor necrosis factor has been implicated in the pathophysiology of Wegener's granulomatosis (Marquez, J., et al. (2003) Curr. Rheumatol. Rep. 5:128; Harman, L.E. and Margo, C.E. (1998) Surv. Ophthalmol. 42:458). Wegener's granulomatosis refers to a vasculitis that causes inflammation of blood vessels in the upper respiratory tract (nose, sinuses, ears), lungs, and kidneys. Wegener's granulomatosis is also referred to as midline granulomatosis. Wegener's granulomatosis includes a granulomatous inflammation involving the respiratory tract, and necrotizing vasculitis affecting small to medium-sized vessels. Subjects who have Wegener's granulomatosis often also have arthritis (joint inflammation). Glomerulonephritis may also be present in affected subjects, but virtually any organ may be involved.

c. Churg-Strauss syndrome

[0284] The formulations of the invention may be used to treat Churg-Strauss syndrome. Tumor necrosis factor has been implicated in the pathophysiology of Churg-Strauss syndrome (Gross, W.L (2002) Curr. Opin. Rheumatol. 14:11; Churg, W.A.(2001) Mod. Pathol. 14:1284). Churg-Strauss syndrome refers to a vasculitis that is systemic and shows early manifestation signs of asthma and eosinophilia. Churg-Strauss syndrome is also referred to as allergic granulomatosis and angiitis, and occurs in the setting of allergic rhinitis, asthma and eosinophilia. Sinusitis and pulmonary infiltrates also occur in Churg-Strauss syndrome, primarily affecting the lung and heart. Peripheral neuropathy, coronary arteritis and gastrointestinal involvement are common.

M. Other diseases

[0285] The formulations of the invention may be used to treat various other disorders in which TNF α activity is detrimental. Examples of other diseases and disorders in which TNF α activity has been implicated in the pathophysiology, and thus which can be treated using an antibody, or antibody portion, of the invention, include inflammatory bone disorders and bone resorption disease (see e.g., Bertolini, D. R., et al. (1986) Nature 319:516-518; Konig, A., et al. (1988) J. Bone Miner. Res. 3:621-627; Lerner, U. H. and Ohlin, A. (1993) J. Bone Miner. Res. 8:147-155; and Shanlar, G. and Stem, P. H. (1993) Bone 14:871-876), hepatitis, including alcoholic hepatitis (see e.g., McClain, C. J. and Cohen, D. A. (1989) Hepatology 9:349-351; Felver, M. E., et al. (1990) Alcohol. Clin. Exp. Res. 14:255-259; and Hansen, J., et al. (1994) Hepatology 20:461-474), viral hepatitis (Sheron, N., et al. (1991) J. Hepatol. 12:241-245; and Hussain, M. J., et al. (1994) J. Clin. Pathol. 47:1112-1115), and fulminant hepatitis; coagulation disturbances (see e.g., van der Poll, T., et al. (1990) N. Engl. J. Med. 322:1622-1627; and van der Poll, T., et al. (1991) Prog. Clin. Biol. Res. 367:55-60), burns (see e.g., Giroir, B. P., et al. (1994) Am. J. Physiol. 267:H 118-124; and Liu, X. S., et al. (1994) Burns 20:40-44), reperfusion injury (see e.g., Scales, W. E., et al. (1994) Am. J. Physiol. 267:G1122-1127; Serrick, C., et al. (1994) Transplantation 58:1158-1162; and Yao, Y. M., et al. (1995) Resuscitation 29:157-168), keloid formation (see e.g., McCauley, R. L., et al. (1992) J. Clin. Immunol. 12:300-308), scar tissue formation; pyrexia; periodontal disease; obesity and radiation toxicity.

[0286] Examples of other disorders that may be treated with the formulations of the invention are described in US20040126372 and US6258562.

[0287] In one embodiment, the formulation of the invention are used to treat rheumatoid arthritis, psoriatic arthritis, or ankylosing spondylitis. The formulation of the disclosure comprising an isolated human TNF α antibody, or antigen-binding portion thereof, (e.g., adalimumab), maybe administered to a human subject according to a dosing scheme and dose amount effective for treating rheumatoid arthritis, psoriatic arthritis, or ankylosing spondylitis. In one instance, a dose of about 40 mg of a human TNF α antibody, or antigen-binding portion thereof, (e.g., adalimumab) (e.g., 0.4 mL of a 100 mg/mL formulation of the disclosure) in the formulation of the disclosure is administered to a human subject every other week for the treatment of rheumatoid arthritis, psoriatic arthritis, or ankylosing spondylitis. In one instance, a dose of about 80 mg of a human TNF α antibody, or antigen-binding portion thereof, (e.g., adalimumab) (e.g., 0.8 mL of a 100 mg/mL formulation of the disclosure) in the formulation of the disclosure is administered to a human subject monthly for the treatment of rheumatoid arthritis, psoriatic arthritis, or ankylosing spondylitis. In one embodiment, the formulation is administered subcutaneously, every other week (also referred to as biweekly, see methods of administration described in US20030235585) for the treatment of rheumatoid arthritis, ankylosing spondylitis, or psoriatic arthritis. In one embodiment, the formulation is administered subcutaneously, monthly for the treatment of rheumatoid arthritis, ankylosing spondylitis, or psoriatic arthritis.

[0288] In one embodiment, the formulation of the invention is used to treat Crohn's disease or ulcerative colitis. The formulation of the disclosure comprising an isolated human TNF α antibody, or antigen-binding portion thereof, (e.g., adalimumab), may be administered to a human subject according to a dosing scheme and dose amount effective for treating Crohn's disease. In one instance, a dose of about 160 mg of a human TNF α antibody, or antigen-binding portion thereof, (e.g., adalimumab) (e.g., 1.6 mL of a 100 mg/mL formulation of the disclosure) in the formulation of the disclosure is administered to a human subject initially at about day 1, followed by a subsequent dose of 80 mg of the antibody (e.g., 0.8 mL of a 100 mg/mL formulation of the disclosure) two weeks later, followed by administration of about 40 mg (e.g., 0.4 mL of a 100 mg/mL formulation of the disclosure) every other week for the treatment of Crohn's disease. In one embodiment, the formulation is administered subcutaneously, according to a multiple variable dose regimen comprising an induction dose(s) and maintenance dose(s) (see, for example, U.S. Patent Publication Nos. US20060009385 and US20090317399) for the treatment of Crohn's disease or ulcerative colitis. In one embodiment, the formulation is administered subcutaneously, biweekly or monthly for the treatment of Crohn's disease or ulcerative colitis. In one instance, a dose of about 80 mg of a human TNF α antibody, or antigen-binding portion thereof, (e.g., adalimumab) (e.g., 0.8 mL of a 100 mg/mL formulation of the disclosure) in the formulation of the disclosure is administered to a human subject monthly for the treatment of Crohn's disease or ulcerative colitis.

[0289] In one embodiment, the formulation of the invention is used to treat psoriasis. The formulation of the disclosure

comprising an isolated human TNF alpha antibody, or antigen-binding portion thereof, (e.g., adalimumab), may be administered to a human subject according to a dosing scheme and dose amount effective for treating psoriasis. In one instance, an initial dose of about 80 mg of a human TNF alpha antibody, or antigen-binding portion thereof, (e.g., adalimumab) (e.g., 0.8 mL of a 100 mg/mL formulation of the disclosure) in the formulation of the disclosure is administered to a human subject, followed by a subsequent dose of 40 mg of the antibody (e.g., 0.4 mL of a 100 mg/mL formulation of the disclosure) every other week starting one week after the initial dose. In one embodiment, the formulation is administered subcutaneously, according to a multiple variable dose regimen comprising an induction dose(s) and maintenance dose(s) (see, for example, US 20060009385 and WO 2007/120823) for the treatment of psoriasis. In one embodiment, the formulation is administered subcutaneously, biweekly or monthly for the treatment of psoriasis. In one instance, a dose of about 80 mg of a human TNF alpha antibody, or antigen-binding portion thereof, (e.g., adalimumab) (e.g., 0.8 mL of a 100 mg/mL formulation of the disclosure) in the formulation of the disclosure is administered to a human subject monthly for the treatment of psoriasis.

[0290] In one embodiment, the formulation of the invention is used to treat juvenile idiopathic arthritis (JIA). The formulation of the disclosure comprising an isolated human TNF alpha antibody, or antigen-binding portion thereof, (e.g., adalimumab), may be administered to a human subject according to a dosing scheme and dose amount effective for treating JIA. In one instance, 20 mg of a human TNF alpha antibody, or antigen-binding portion thereof, in the formulation of the disclosure (e.g., 0.2 mL of a 100 mg/mL formulation of the disclosure) is administered to a subject weighing 15 kg (about 33 lbs) to less than 30 kg (66 lbs) every other week for the treatment of JIA. In another instance, 40 mg of a human TNF alpha antibody, or antigen-binding portion thereof, in the formulation of the disclosure (e.g., 0.4 mL of a 100 mg/mL formulation of the disclosure) is administered to a subject weighing more than or equal to 30 kg (66 lbs) every other week for the treatment of JIA. In one embodiment, the formulation is administered subcutaneously, according to a weight-based fixed dose (see, for example, U.S. Patent Publication No. 20090271164) for the treatment of JIA. In one embodiment, the formulation is administered subcutaneously biweekly or monthly for the treatment of JIA.

[0291] In one instance, an isolated human TNF alpha antibody, or antigen-binding portion thereof, (e.g., adalimumab), may be administered to a human subject for treatment of a disorder associated with detrimental TNFa activity according to a monthly dosing schedule, whereby the antibody is administered once every month Or once every four weeks. As described above, examples of disorders that may be treated according to a monthly dosing schedule using the formulations of the invention include, but are not limited to, rheumatoid arthritis, ankylosing spondylitis, JIA, psoriasis, Crohn's disease, ulcerative colitis, hidradenitis suppurativa, giant cell arteritis, Behcet's disease, sarcoidosis, diabetic retinopathy, or psoriatic arthritis. Thus, the formulation of the disclosure comprising an isolated human TNF alpha antibody, or antigen-binding portion thereof, (e.g., adalimumab), may be administered to a human subject for treatment of a disorder associated with detrimental TNFa activity according to a monthly dosing schedule. In one instance, 80 mg of a human TNF alpha antibody, or antigen-binding portion thereof, in the formulation of the disclosure (e.g., 0.8 mL of a 100 mg/mL formulation of the disclosure) is administered to a subject having a disorder associated with detrimental TNFa activity. In one instance, mg of a human TNF alpha antibody, or antigen-binding portion thereof, in the formulation of the disclosure (e.g., 0.8 mL of a 100 mg/mL formulation of the disclosure) is administered monthly or biweekly to a subject for the treatment of a disorder associated with detrimental TNFa activity.

[0292] Dose amounts described herein may be delivered as a single dose (e.g., a single dose of 40 mg in 0.4 mL or 80 mg dose in 0.8mL), or, alternatively may be delivered as multiple doses (e.g., four 40 mg doses or two 80 mg doses for delivery of a 160 mg dose).

[0293] The formulation of the disclosure comprising an isolated human TNF alpha antibody, or antigen-binding portion thereof, (e.g., adalimumab) may also be administered to a subject in combination with an additional therapeutic agent. In one embodiment, the formulation is administered to a human subject for treatment of rheumatoid arthritis in combination with methotrexate or other disease-modifying anti-rheumatic drugs (DMARDs). In another embodiment, the formulation is administered to a human subject for treatment of JIA in combination with methotrexate or other disease-modifying anti-rheumatic drugs (DMARDs). Additional combination therapies are described in U.S. Patent Nos. 6,258,562 and 7,541,031; and U.S. Patent Publication No. US20040126372.

[0294] The formulation of the disclosure comprising a human TNF alpha antibody, or antigen-binding portion thereof, may also be used to treat a subject who has failed previous TNF inhibitor therapy, e.g., a subject who has lost response to or is intolerant to infliximab.

[0295] The invention is further illustrated in the following examples, which should not be construed as further limiting.

EXAMPLES

EXAMPLE 1: HIGH CONCENTRATION ANTI-TNF α ANTIBODY FORMULATION REDUCES INJECTION PAIN

[0296] There have been reports of pain associated with the subcutaneous administration of a human anti-TNF α antibody, e.g., adalimumab. In placebo-controlled trials, 20% of patients treated with adalimumab developed injection

site reactions (erythema and/or itching, haemorrhage, pain or swelling), compared to 14% of patients receiving placebo. Most injection site reactions are mild and do not generally necessitate drug discontinuation.

[0297] There are two main components of the injection pain associated with adalimumab: the pain associated with the needle stick, and the pain associated with the injection of drug into the tissue. The injection-related pain may be related to the Adalimumab formulation and/or to the volume of medication. The following study examined whether various formulations have an impact on injection pain following subcutaneous delivery of adalimumab.

Materials and Methods

Study Design

[0298] The primary objectives of this study were to compare injection-related pain of three high concentration (100 mg/mL) adalimumab formulations in the PHYSIOLIS™ pre-filled syringe with the current (50 mg/mL) adalimumab commercial formulation in the current pre-filled syringe; and to assess the bioavailability of three high concentration (100 mg/mL) adalimumab formulations in comparison to the current (50 mg/mL) adalimumab commercial formulation. The secondary objective of this study was to assess the safety and tolerability of all four adalimumab formulations.

[0299] 200 healthy adult male and female subjects who fulfilled the study eligibility criteria were recruited to participate in the study. Generally, the study was conducted according to a randomized parallel-group design. Pain assessment data was preferably obtained from all 200 subjects. Evaluation of pharmacokinetics (PK) was done only for the first 100 or so subjects.

[0300] Subjects from each treatment group were scheduled to receive one subcutaneous injection of adalimumab 40 mg via a pre-filled syringe. There were four treatment groups, one for each of the four formulations as set forth in Table 1 below. After meeting the selection criteria, subjects were randomly assigned in roughly equal numbers to one of the four treatment groups shown in Table 1.

[0301] The three high concentration formulations (F1, F3, and F4), each contained 40 mg of adalimumab in 0.4 mL of solution in the PHYSIOLIS™ pre-filled syringe. F1, F3, and F4 were compared to the current adalimumab commercial formulation of 40 mg of adalimumab in 0.8 mL of solution in the current pre-filled syringe. The ingredients for each of the formulations is described below in Table 1. The formulations described in Table 1 also refer to the formulations described in Examples 2-7 below.

Table 1 Treatment Groups

Treatment Group	No. of Subjects	Study Day 1 SC Injection	Formulation
A	50	High Conc. Formulation 1 (F1) (40 mg/0.4 mL in the PHYSIOLIS™ pre-filled syringe)	adalimumab, Mannitol, Citric acid monohydrate, Sodium citrate, Disodium phosphate dihydrate, Polysorbate 80, Water for injection, Sodium Hydroxide added as necessary to adjust pH.
B	50	High Conc. Formulation 3 (F3) (40 mg/0.4 mL in the PHYSIOLIS™ pre-filled syringe)	adalimumab, Mannitol, polysorbate 80, Water for injection
C	50	High Conc. Formulation 4 (F4) (40 mg/0.4 mL in the PHYSIOLIS™ pre-filled syringe)	adalimumab, Polysorbate 80, Water for injection
D	50	Current Commercial Formulation (40 mg/0.8 mL in the current syringe)	adalimumab, Mannitol, Citric acid monohydrate, Sodium citrate, Disodium phosphate dihydrate, Sodium dihydrogen phosphate dihydrate, Sodium chloride, Polysorbate 80, Water for injection, Sodium Hydroxide added as necessary to adjust pH

[0302] The first about 100 subjects to fulfill all entry criteria and enroll in the study were randomized to the four treatment groups in roughly equal numbers in each group and participated as either Cohort 1 or Cohort 2. The second about 100 subjects to fulfill all entry criteria and enroll in the study were randomized to the four treatment groups, in roughly equal numbers in each group, and participated as Cohorts 3-5. It is the Cohort number that specifies if a subject has pharma-

cokinetic (PK) and pain assessments or only pain assessments as described in Table 2 below.

Table 2 Assignment of Study Subjects

Cohort	Total N	Assessments	No. of Subjects			
			A	B	C	D
1	50	PK and Pain	13	12	13	12
2	44	PK and Pain	11	12	10	11
3	38	Pain	9	9	10	10
4	39	Pain	10	10	10	9
5	29	Pain	7	7	7	8

[0303] Pharmacokinetic sample collection and pain assessments was done for all subjects in the first two cohorts of about 100 patients (Cohorts 1 and 2). Subjects in Cohorts 3-5 only participated in pain assessments, and no pharmacokinetic samples were collected for these subjects. Safety and tolerability were to be assessed in all subjects of all 5 cohorts. Each subject was randomly assigned to receive one injection of adalimumab on Study Day 1. Each dose of study drug was to be administered subcutaneously by an appropriate site staff member via a pre-filled syringe in accordance with the proper injection method. The injection was given subcutaneously in the abdomen 2 inches to the right of the navel. Questionnaires were administered by a different study staff member than the individual administering the injection, as often as possible.

[0304] Subjects in Cohorts 1 and 2 (pharmacokinetic and pain assessments) were confined to the study site and supervised for approximately 10 days (9 nights). Confinement for each subject began on Study Day -1 (1 day prior to the dosing day) and ended after the collection of the 192 hour blood samples and scheduled study procedures on Study Day 9. Serial blood samples were collected through Study Day 57 after dosing with subjects returning for outpatient visits. Safety and tolerability were assessed throughout the study. Subjects in Cohorts 3-5 (pain assessments only) were confined to the study site and supervised for approximately 3 days (2 nights). Confinement for each subject began on Study Day -1 (1 day prior to the dosing day) and ended after the completed study procedures on Study Day 2. Safety and tolerability were assessed throughout the study.

[0305] In addition to bioavailability and AAA assays, tolerability was preferably assessed as follows:

- 1) Immediately following the injection on Study Day 1: Pain Assessment Module was completed by the subject.
- 2) Approximately 10 minutes following the injection on Study Day 1: Draize Scale (hemorrhage, petechiae, erythema, edema, and pruritus) was evaluated by a qualified site staff member.
- 3) Approximately 15 minutes following the injection on Study Day 1: Pain Assessment Module was completed by the subject.
- 4) Approximately 30 minutes following the injection on Study Day 1: Pain Assessment Module and Draize Scale evaluation was completed by the subject and a qualified site staff member, respectively.

[0306] The demographics of subjects in the treatment groups are as follows, shown in Table 3, below.

Table 3 Patient Demographics

Variable	High Conc. Formulation 1 (N=50)	High Conc. Formulation 3 (N=50)	High Conc. Formulation 4 (N=50)	Commercial Formulation (N=50)
Age (yrs)	29.6 ± 8.7	29.5 ± 9.4	30.0 ± 8.9	30.3 ± 9.7
Weight (kg)	68.3 ± 13.9	69.6 ± 9.6	67.0 ± 8.4	68.5 ± 10.0
Sex	31 F (62%), 19 M (38%)	25 F (50%), 25 M (50%)	31 F (62%), 19 M (38%)	30 F (60%), 20 M (40%)

EP 2 637 690 B1

(continued)

Variable	High Conc. Formulation 1 (N=50)	High Conc. Formulation 3 (N=50)	High Conc. Formulation 4 (N=50)	Commercial Formulation (N=50)
Race	37 white (74%), 9 black (18%), 4 other (8%)	45 white (90%), 3 black (6%), 2 other (4%)	36 white (72%), 10 black (20%), 4 other (8%)	40 white (80%), 8 black (16%), 2 other (4%)

Formulations

[0307] Three new high concentration formulations (referred to herein as Formulation 1, 3, or 4; or F1, F3, or F4, respectively) were studied relative to the commercial 50 mg/mL adalimumab formulation. The compositions of each of these formulations are listed below in Tables 4 - 7.

Table 4: Formulation 1 (F1)

COMPOSITION OF THE BULK SOLUTION 1 mL bulk solution contains	
Name of ingredient	Concentration [mg]
Active Substance	
adalimumab (A-765865)*	100.00
Excipients	
Mannitol	42.00
Citric acid monohydrate	1.31
Sodium citrate	0.31
Disodium phosphate dihydrate	1.53
Sodium dihydrogen phosphate dihydrate	0.86
Polysorbate 80	1.00
Sodium Hydroxide	q.s.
Water for injections	ad 1,041.00
Nitrogen	---
Density of the solution: 1.041 g/mL * Used as concentrate.	

Table 5: Formulation 3 (F3)

COMPOSITION OF THE BULK SOLUTION 1 mL bulk solution contains	
Name of ingredient	Concentration [mg]
Active Substance	
adalimumab (A-765865)*	100
Excipients	
Mannitol	42.00
Polysorbate 80	1.00
Water for injections	ad 1,040.00

EP 2 637 690 B1

(continued)

Excipients	
Nitrogen	---
* Used as concentrate. Density of the solution: 1.040 g/mL	

Table 6: Formulation 4 (F4)

COMPOSITION OF THE BULK SOLUTION 1 mL bulk solution contains	
Name of ingredient	Concentration [mg]
Active Substance	
adalimumab (A-765865)*	100.00
Excipients	
Polysorbate 80	1.00
Water for injections	ad 1,026.00
Nitrogen	---
* Used as concentrate. Density of the solution: 1.026 g/mL	

Table 7: Commercial 50 mg/mL adalimumab formulation

COMPOSITION OF THE BULK SOLUTION 1 mL bulk solution contains	
Name of ingredient	Concentration [mg]
Active Substance	
adalimumab (A-765865)*	50.00
Excipients	
Mannitol	12.00
Citric acid monohydrate	1.30
Sodium citrate	0.30
Disodium phosphate dihydrate	1.53
Sodium dihydrogen phosphate dihydrate	0.86
Polysorbate 80	1.00
Sodium Hydroxide	q.s.
Water for injections	ad about 1,000
Sodium Chloride	6.16
Nitrogen	---

Study Drug Administration

[0308] Study drug (adalimumab) in the various formulations was administered in the morning at Hour 0 on Study Day 1. The four treatment groups are set forth in Table 1 above as Groups A, B; C and D. Subjects in each treatment group were subcutaneously injected with only a single adalimumab formulation via a pre-filled syringe.

[0309] Two types of syringes were used in this study; the currently commercially available glass pre-filled syringe ("current pre-filled syringe") was used for the reference current adalimumab commercial formulation (40 mg of adalimumab in 0.8 mL of solution), and the PHYSIOLIS™ pre-filled syringe for the three high concentration test formulations (40 mg of adalimumab in 0.4 mL of solution). The PHYSIOLIS™ pre-filled syringe has a 29 gauge needle (the current pre-filled syringe has a 27 gauge 1/2 inch length fixed needle), a latex-free needle shield, and a plunger stopper which is coated to minimize leachables.

Pain Testing: Pain Scale

[0310] The Pain Visual Analog Scale was used to quantitatively assess pain sensation. The following instructions were followed to assess the Pain Visual Analog Scale (VAS):

[03111] The pain scale was administered to the subject at three different times after the injection: immediately after the injection, at 15 minutes after the injection, and at 30 minutes after the injection on Study Day 1. The pain scale was shown and read to the subject, who was asked to place one straight vertical mark along the line in the pain scale to indicate their current level of pain at the injection site (for example see below). An indication of 0 meant no pain, while the highest score (10) indicated "the worst imaginable pain." An illustrative pain scale used in the study is shown below: What is your current level of pain at the injection site?

[illegible]

Qualitative Assessment of Pain

[0312] Following the completion of the pain scale, the qualitative assessment of pain was administered three times after the injection: immediately following the injection, at 15 minutes post the injection, and 30 minutes following the injection on Study Day 1. An exemplary qualitative assessment of pain used in the study is shown below:

[0313] Select all that describe your current level of pain at the injection site:

Shooting pain
Sharp pain
Stinging Pain
Dull
Uncomfortable
Pressure
Aching
Soreness
Localized burning
Other
OR
I currently have no

Needle Pain Assessment

[0314] After the qualitative assessment of pain was completed, a needle pain assessment was administered immediately following the injection. An exemplary needle pain assessment used in the study is shown below:

[Q0315] Were you able to tell the difference between the pain from the needle entering your skin and the pain from the solution that was injected?

Yes

No

a. If yes, was most of your pain caused by the needle entering your skin or was most of your pain caused by the solution that was injected?

Most of my pain was caused by the needle entry

Most of my pain was caused by the solution that was injected

Draize Scale

[0316] Qualified site personnel completed this assessment for each subject approximately 10 minutes and 30 minutes after the injection on Study Day 1.

[0317] Hemorrhage/Petechiae at injection site:

- 0: None
- 1: Isolated; up to 5 petechiae
- 2: Isolated but > 5 petechiae
- 3: Many petechiae, some coalescence
- 4: Spots of blood on surface
- 5: Frank bleeding

[0318] Erythema at injection site:

- 0: No erythema
- 1: Very slight (barely perceptible) erythema
- 2: Well-defined erythema
- 3: Moderate to severe erythema
- 4: Severe erythema (beet redness)

[0319] Edema at injection site:

- 0: No edema
- 1: Very slight (barely perceptible) edema
- 2: Slight edema (edges of area well defined by slight raising)
- 3: Moderate edema (raised ~1mm)
- 4: Severe edema (raised > 1 mm, extending beyond area of injection)

[0320] Pruritis at injection site:

- 0: No pruritis
- 1: Occasional pruritis
- 2: Constant pruritis

Results

[0321] To determine whether delivery of adalimumab could be improved, new high concentration formulations were developed. Formulations F1, F3, and F4, as shown below, have half of the volume (*i.e.*, 0.4 mL vs. 0.8 mL) and twice the protein concentration (100 mg/mL vs. 50 mg/mL) compared to the current commercial adalimumab formulation, and they also have different excipient compositions. Experiments described herein were designed to assess whether any of the new formulations are superior to the current commercial adalimumab formulation.

[0322] The pain visual analog scale was chosen to assess injection site pain, and was used to evaluate impact of formulation composition on pain sensations. In addition, tolerability of various new adalimumab 100 mg/mL formulations were compared to the current commercial formulation (50 mg/mL adalimumab formulation). Data in this example supports the surprising finding that the new formulations, especially Formulation 3 (F3), decreases pain significantly relative to the current commercial formulation. Surprisingly, F3 also decreased pain significantly relative to formulations F1 and F4.

[0323] Specifically, Figure 1 shows that administration of high concentration Formulations 1 and 3 resulted in a significant decrease in pain assessment at all time points after injection (immediately, 15 minutes, and 30 minutes), compared to the other treatment groups (F4 and the current commercial formulation). Table 8, shown below, summarizes the individual data and shows a comparison of the F1, F3, and F4 formulations with the 0.8 mL, 50 mg/mL commercial formulation.

[0324] As described in Table 8, immediately after injection, subjects who received the current Humira formulation reported a mean (SD) pain score of 3.29 (2.57) cm. The mean pain scores for Formulation 1 and Formulation 3 were statistically significantly lower than that for the current Humira formulation ($p < 0.001$). The estimated differences from the current Humira formulation were -1.50 (95% CI: -2.31 - -0.69 cm) for Formulation F1, and -2.70 (95% CI: -3.52 - -1.89 cm) for Formulation F3. Thus, Treatments A and B (high concentration Formulations 1 and 3) resulted in 45.6% and 82.7% reductions in injection site pain, respectively. Statistical tests were not performed for the pain scores assessed

at 15 minutes and 30 minutes after injection because a majority of the subjects reported no pain at these time points. As described in Figure 1, the minimum / maximum VAS scores immediately after injection were as follows: Formulation F1, 0.00-8.3; Formulation F3, 0.00-2.20; Formulation F4, 0.20-8.70; and current commercial formulation, 0.00-10.00.

[0325] It was apparent that the pain associated with the injection of Formulation 3 was dramatically reduced compared to the same for the current commercial formulation. Specifically, mean pain value, as assessed by the pain visual analog scale (VAS) immediately after injection, decreased from a mean of 3.29 in the current commercial formulation to 0.56 in Formulation 3, a stark 83% reduction. In fact, the pain reduction associated with Formulation 3 was so significant, it was 69% reduced from the level of the next best formulation (in terms of pain) - Formulation 1 (1.79).

[0326] Similarly, the mean pain scale for Formulation 1 reduced to 1.79, a 45% reduction from the 3.29 pain scale associated with the current commercial formulation.

Table 8 Pain Visual Analog Scales (VAS)

Immediately After Injection						
Treatment	N	Mean (SD)	LS Mean&	Comparisons with Current Formulation&		
				Estimate	P-value#	95% CI
High Conc. Formulation 1	50	1.79 (2.08)	1.79	-1.50	<0.001	-2.31,-0.69
High Conc. Formulation 3	50	0.56 (0.56)	0.58	-2.71	<0.001	-3.52, - 1.90
High Conc. Formulation 4	50	4.12 (2.50)	4.11	0.82	0.976	0.01, 1.63
Current Formulation	50	3.29 (2.57)	3.29			
# From a one-sided test with the null hypothesis being that the mean for the test formulation is $\geq$ the mean for the current formulation & Based on ANOVA.						

[0327] A Qualitative Assessment of Pain was also administered to subjects immediately after injection, 15 minutes after injection and at 30 minutes after injection for all four adalimumab treatments. Immediately after injection, an assessment of "no discomfort" was reported with greatest frequency by 31 subjects (31/50, 62.0%) who had received Formulation 3, followed by 19 subjects (19/50, 38.0%) who had received Formulation 1, 7 subjects (7/50, 14.0%) who had received the current Humira formulation, and one subject (1/50, 2.0%) who had received Formulation 4. Of those subjects who reported discomfort immediately after injection, "stinging pain" was the most frequently reported sensation with 30 subjects (30/50, 60%) for each of the current formulation and formulation 4, 16 subjects (16/50, 32.0%) for Formulation 1, and 4 subjects (4/50, 8.0%) for Formulation 3. At 15 minutes after injection, a large majority of subjects receiving each treatment reported "no discomfort" at the injection site.

[0328] Study site staff also utilized the Draize Scale to evaluate hemorrhage/petechiae, Erythema, edema and pruritis at the injection site of each subject. Ten minutes after injection the majority of subjects in all treatment groups had no observed injection-site hemorrhages or petechiae, edema or pruritis.

[0329] All four formulations were well tolerated during the study. A summary of preliminary adverse events (AE) data is shown below in Table 9.

Table 9 Preliminary Adverse Events (AE)

	High Conc. Formulation 1 (N=50)	High Conc. Formulation 3 (N=50)	High Conc. Formulation 4 (N=50)	Current Formulation (N=50)
Any AE	7 (14%)	7 (14%)	6 (12%)	3 (6%)
Any AE at least possibly drug related	3 (6%)	3 (6%)	2 (4%)	1 (2%)
Any severe AE	0	0	0	0
Any serious AE	0	0	0	0
Any AE leading to study discontinuation	0	0	0	0
Deaths	0	0	0	0

[0330] Accordingly, the data demonstrates that the new 100 mg/mL formulations, especially formulations 1 and 3, are well tolerated, and were effective in reducing injection site pain after subcutaneous injection of similar therapeutic doses as compared to the currently marketed adalimumab formulation. Formulation F3 had a particularly low VAS score relative to the other formulations tested.

[0331] The reduction in pain using the VAS score was not related to the difference in needle size (a 27G needle was used to administer the current adalimumab commercial formulation and a 29G needle was used to administer formulations F1, F3, and F4). In particular, a needle prick accounts for an immediate pain response, whereas the pain response measured by the VAS scale indicated a prolonged persistent pain over several minutes, demonstrating that the injected solution itself contributes to the majority of the response. In addition, all of the test formulations (F1, F3, and F4) were injected using the same size needle, yet F1, F3, and F4 had very different VAS scores. This result further demonstrates that it was the formulation contributing to the pain effect and that this can be separated from the size of the needle used to administer the formulations.

EXAMPLE 2: HIGH CONCENTRATION ANTI-TNFA ANTIBODY FORMULATIONS INCREASE BIOAVAILABILITY IN HUMANS

[0332] The following example describes a Phase 1, single-blind, single-dose, parallel-group design, randomized study in healthy volunteers (same study described above in Example 1). The primary objectives of this study were to compare injection-related pain of three high concentration (100 mg/mL) adalimumab formulations in the PhysioliS PFS with the current (50 mg/mL) adalimumab (Humira) formulation in the current PFS (see Example 1), and to assess the bioavailability of three high concentration (100 mg/mL) adalimumab formulations in comparison to the current commercial (50 mg/mL) adalimumab (Humira) formulation. The secondary objective of this study was to assess the safety and tolerability of all four adalimumab formulations.

Study Design

[0333] Two hundred healthy volunteers were enrolled in this study (Table 10). Pain assessment data were obtained from all 200 subjects. Adalimumab pharmacokinetics were evaluated in the first 100 subjects. A description of the formulations is provided above in Table 1.

Table 10. Treatment Groups

Treatment Group	Number of Subjects for Study	Number of Subjects for Pharmacokinetic Data	Study Day 1
			SC Injection
A	50	24	High Concentration Adalimumab Formulation No. 1 (40 mg/0.4 mL in the PhysioliS PFS)
B	50	24	High Concentration Adalimumab Formulation No. 3 (40 mg/0.4 mL in the PhysioliS PFS)
C	50	23	High Concentration Adalimumab Formulation No. 4 (40 mg/0.4 mL in the PhysioliS PFS)
D	50	23	Current Commercial Humira Formulation (40 mg/0.8 mL in the current PFS)
*See Tables 4-7 for formulation compositions.			

[0334] Subjects from each treatment group received one subcutaneous injection of 40 mg adalimumab *via* PFS on Study Day 1. Each dose of study drug was administered subcutaneously by an appropriate site staff member in accordance with the proper injection method as described in the protocol. The injection was given subcutaneously in the abdomen 2 inches to the right of the navel. Questionnaires were administered by a different study staff member than the individual administering the injection, as often as possible.

Results

Pharmacokinetic Results and Conclusions

[0335] Following a single subcutaneous dose of adalimumab, the central values of the pharmacokinetic parameters,

T_{max} , C_{max} , AUC_{0-360} and AUC_{0-1344} were similar between Treatments A, B (high concentration adalimumab Formulations 1 and 3, respectively) and D (current commercial Humira formulation). Following the single dose administration, mean T_{max} was earlier for Treatment C (high concentration adalimumab Formulation 4) relative to Treatment D (Figures 2 and 3). The central values of C_{max} and AUC_{0-360} values were greater ($p < 0.05$) for Treatment C versus Treatment D.

Bioavailability/Bioequivalence for Treatments A, B and C (Adalimumab High Concentration Formulations) Relative to Treatment D (Commercial Humira Formulation)

[0336] For Treatment Group A versus D, the point estimates for the ratios of C_{max} , AUC_{0-360} , and AUC_{0-1344} central values for Treatments A and B were near unity, and the 90% confidence intervals were within the 0.80 to 1.25 range. For Treatment B versus D, the point estimates for the ratios of C_{max} and AUC_{0-360} central values were near unity and the 90% confidence intervals were within the 0.80 to 1.25 range. For AUC_{0-1344} , the upper bound of the 90% confidence interval for Treatments B versus D was above 1.25. For Treatment C versus D, the point estimates for the ratio of C_{max} , AUC_{0-360} and AUC_{0-1344} central values were 1.429, 1.309, and 1.170 respectively, indicating that the relative bioavailability of Treatment C (Formulation 4) was greater.

Table 11. Relative Bioavailability and 90% Confidence Intervals for the Bioequivalence Assessment

Treatments [£] Test vs. Reference	PK Parameter	Central Value		Relative Bioavailability	
		Test	Reference	Point Estimate	90% Confidence Interval
A vs. D	C_{max}	4.47	4.39	1.018	0.859-1.207
	AUC_{0-360}	1192.14	1192.23	1.000	0.860-1.163
	AUC_{0-1344}	2306.91	2387.28	0.966	0.814-1.147
B vs. D	C_{max}	4.52	4.39	1.029	0.868-1.219
	AUC_{0-360}	1222.24	1192.23	1.025	0.882-1.192
	AUC_{0-1344}	2547.95	2387.28	1.067	0.899-1.266
C vs. D	C_{max}	6.28	4.39	1.429	1.202-1.699
	AUC_{0-360}	1561.05	1192.23	1.309	1.123-1.527
	AUC_{0-1344}	2794.29	2387.28	1.170	0.983-1.394

[£] Treatments A, B or C: a single dose of high concentration adalimumab Formulation 1, 3 or 4, respectively, administered as a single sc injection using a Physiolis PFS (40 mg/0.4 mL).

Treatment D: a single dose of the current Humira formulation administered as a single sc injection using the currently available glass PFS (40 mg/0.8 mL). PK = Pharmacokinetic.

Pharmacokinetic Conclusions.

[0337] Based on the pharmacokinetic results, the relative bioavailability of Treatments A and B were similar to Treatment D, the currently marketed Humira formulation. The relative bioavailability of Treatment C was greater when compared to Treatment D. The unexpected increase in bioavailability for Treatment C suggests that the effective dose amount administered to a subject may be reduced.

Immunogenicity Conclusions

[0338] Twelve subjects had positive AAA samples during any time in the study, with only two subjects determined as AAA positive according to the pre defined definition. Because of the small sample size and relatively similar numbers of AAA positive samples, no conclusions can be made of the immunogenicity between the treatments.

Safety Conclusions

[0339] The treatments tested were generally well tolerated by the subjects. No clinically significant vital signs, ECG or laboratory measurements were observed during the course of the study. The majority of adverse events were assessed by the investigator as probably not or not related to study drug and mild in severity. No adverse events were assessed as severe.

[0340] No deaths, serious adverse events or discontinuations due to adverse events occurred during the study.

[0341] Results of other safety analyses, including individual subject changes and potentially clinically significant values for vital signs, ECG and laboratory measurements, were unremarkable for all treatment groups.

5 Tolerability

[0342] The tolerability assessments that were conducted included completion of a Pain Assessment Module (Pain Visual Analog Scale [VAS]), Qualitative Assessment of Pain and Needle Pain Assessment) and the Draize Scale (see Example 1).

10 EXAMPLE 3: HIGH CONCENTRATION ANTI-TNFA ANTIBODY FORMULATION INCREASE BIOAVAILABILITY IN PRE-CLINICAL MODEL

[0343] The objective of the following study was to evaluate the pharmacokinetic profiles of adalimumab formulation F4 in contrast to the adalimumab commercial formulation (see Table 7 above for a description of the formulation).

[0344] The pharmacokinetic profiles of HUMIRA (Adalimumab) were studied in male and female Beagle dogs (2/sex/s.c. administration and 2 males/i.v. administration, Marshall Bio Resources USA, Inc., North Rose, NY 14516) after a single subcutaneous (s.c.) injection of the HUMIRA commercial formulation (adalimumab) and a HUMIRA test formulation corresponding to formulation F4 of the previous examples (adalimumab), as well as an intravenous (i.v.) injection of the HUMIRA commercial formulation as a control. The administered dose was 40 mg/dog (at 100 mg/mL for F4 and 50 mg/mL of the commercial formulation).

[0345] For the determination of Adalimumab serum exposure levels, blood samples were collected at 0.083, 4, 24, 48, 96, 168, 240, 312, 384, 456, 528 and 864 hours post administration (p.a.). Examined parameters were clinical signs (twice weekly) and mortality.

[0346] Apart from mucous feces in one male animal of the control group, no relevant clinical signs were observed. The incidences of clinical signs are summarized in Tables 14 and 15 below.

[0347] The pharmacokinetic results (described in Table 12 below) of this study indicated that the bioavailability after s.c. dosing was about 80 % and the exposure levels seemed to be higher in females than in males after s.c. dosing. There was a trend for higher exposure levels following s.c. dosing of the test formulation compared with s.c. dosing of the commercial formulation in males.

Table 12: Pharmacokinetic results

Treatment	Gender	Animal Number	AUC _{0-528h} ($\mu\text{g}\cdot\text{Hours/mL}$)	AUC/Dose ($\mu\text{g}\cdot\text{Hours/mL}/\text{mg/kg}$)	Vdss (mL)	T _{1/2} (Hours)
Test formulation, s.c.	male	1001	9020	226	708	39.3
		1003	11400	286	870	187.2
		Mean	10200 $\pm$ 1680	256 $\pm$ 42.4	789 $\pm$ 115	113.3 $\pm$ 104.6
	female	1002	15400	384	388	55.5
		1004	15800	395	469	54.5
		Mean	15600 $\pm$ 283	390 $\pm$ 7.78	429 $\pm$ 57.3	55 $\pm$ 0.7
Commercial formulation, s.c.	male	2001	8010	200	692	21.4
		2003	8230	206	695	72.7
		Mean	8120 $\pm$ 56	203 $\pm$ 4.24	694 $\pm$ 2.12	47.1 $\pm$ 6.3
	female	2002	12700	319	385	34.0
		2004	17200	431	477	119.0
		Mean	15000 $\pm$ 3180	375 $\pm$ 79.2	431 $\pm$ 65.1	76.5 $\pm$ 60.1

EP 2 637 690 B1

(continued)

Treatment	Gender	Animal Number	AUC _{0-528h} ($\mu\text{g}\cdot\text{Hours}/\text{mL}$)	AUC/Dose ($\mu\text{g}\cdot\text{Hours}/\text{mL}/\text{mg}/\text{kg}$)	Vdss (mL)	T _{1/2} (Hours)
Commercial formulation, i.v.	male	3001	9360	234	548	45.5
		3003	11900	298	407	22.2
		Mean	10600 $\pm$ 1800	266 $\pm$ 45.3	478 $\pm$ 99.7	33.9 $\pm$ 16.5

Table 13: Animal identification

Animal No.	Tattoo-No.	Sex
1001	1246730	Male
1003	1230230	Male
1002	1282302	Female
1004	1288689	Female
2001	1284879	Male
2003	1298951	Male
2002	1297237	Female
2004	1280491	Female
3001	1285514	Male
3003	1290143	Male

Table 14: Summary of Clinical Observations in Males

Dosage Group:	2	3	1	
Animals Examined:	4	4	2	
Number Normal:	3	3	2	
Category, Observation	a	b	a	b
Excretion, feces	1	4	0	0

Note: a = Number of animals with observation
b = Number of days observation seen

Table 15: Summary of Clinical Observations in Females

Dosage Group:	2	3	1
Animals Examined:	2	2	2
Number Normal:	2	2	2
Category, Observation			
Note: a = Number of animals with observation b = Number of days observation seen			

EXAMPLE 4: STABILITY OF HIGH CONCENTRATION ANTI-TNFA ANTIBODY FORMULATIONS AGAINST FREEZE/THAW STRESS

[0348] The following example compares the stability of high concentration formulations F1, F3, and F4 with the commercial adalimumab formulation. Stability was examined using freeze/thaw tests.

Experimental Setup

[0349] High concentration human anti-TNF α antibody formulations were prepared as described in Example 1, Table 1 above.

[0350] The compounded solutions were sterile filtered and aliquoted in 8x 30 mL PETG bottles at 20 mL, respectively. The solutions were practically free from particles in visual inspection.

[0351] The samples for T0 were directly placed into a 2-8°C refrigerator. The other bottles were put into the -80°C cube to freeze.

[0352] The next day the bottles were thawed in water baths with a temperature of 25°C or 37°C, respectively.

[0353] The Freeze /Thaw cycles were repeated 5 times. At T0 (before any freeze-thaw cycles), T1 (after one freeze-thaw cycle), T3 (after three freeze-thaw cycles) and T5 (after five freeze-thaw cycles) samples were taken for analysis and stored in a 2-8°C fridge.

→ n = 1 per pullpoint from 4 samples

→ Sample volume: 20 mL

→ Freeze/Thaw: -80°C / 25°C+37°C

→ Freeze/thaw cycles: 5

[0354] After the cycling the samples were analyzed in the lab using each of the following measures: Optical appearance (at each time point); absorption at 340 nm; subvisible particles (at GGDDA); Photon-correlation-spectroscopy (PCS); Size Exclusion Chromatography (SEC); and Ion Exchange Chromatography (IEC).

Subvisible particles

[0355] The measurement of subvisible particles was made at the Klotz particle measurement device. The results are shown in Table 16.

Table 16: Counts of particles $\geq 1 \mu\text{m}$, $\geq 10 \mu\text{m}$, and $\geq 25 \mu\text{m}$

Timepoint	Sample	Temperature	Charge	Particles $\geq$		
				1 μm	10 μm	25 μm
T0	HC F1		(25 °C) E161118001CL	9	1	0
T0	HC F1	(37°C)	E161118001CL	7	2	1
T1	HC F1	25°C	E161118001CL	3	0	0
T1	HC F1	37°C	E161118001CL	33	1	0
T3	HC F1	25°C	E161118001CL	3	0	0
T3	HC F1	37°C	E161118001CL	20	1	0
T5	HC F1	25°C	E161118001CL	4	0	0
T5	HC F1	37°C	E161118001CL	94	0	0
T0	HC F3	(25°C)	E161119001CL	6	3	1
T0	HC F3	(37°C)	E161119001CL	12	2	0
T1	HC F3	25°C	E161119001CL	4	1	0
T1	HC F3	37°C	E161119001CL	7	2	0
T3	HC F3	37°C	E161119001CL	3	1	0
T3	HC F3	37°C	E161119001CL	9	2	1
T5	HC F3	25°C	E161119001CL	7	0	0
T5	HC F3	37°C	E161119001CL	5	0	0

EP 2 637 690 B1

(continued)

5					Particles >=		
	T0	HC F4	(25°C)	E161120001CL	5	1	1
10	T0	HC F4	(37°C)	E161120001CL	7	1	0
	T1	HC F4	25°C	E161120001CL	6	1	0
	T1	HC F4	37°C	E161120001CL	5	1	0
	T3	HC F4	25°C	E161120001CL	12	1	1
15	T3	HC F4	37°9C	E161120001CL	60	0	0
	T5	HC F4	25 °C	E161120001CL	13	0	0
	T5	HC F4	37°C	E161120001CL	22	1	0
	T0	commercial	(25°C)	E161121001CL	464	2	1
20	T0	commercial	(37°C)	E161121001CL	198	0	0
	T1	commercial	25°C	E161121001CL	143	1	0
	T1	commercial	37°C	E161121001CL	285	0	0
	T3	commercial	25°C	E161121001CL	108	0	0
25	T3	commercial	37°C	E161121001CL	224	0	0
	T5	commercial	25°C	E161121001CL	39	0	0
	T5	commercial	37°C	E161121001CL	151	0	0

[0356] The >=1 µm particle data showed a clear trend to a higher particle load in Humira commercial and high concentration (HC) F1 at T5, reflecting a characteristic behavior of buffer salt or sodium chloride containing adalimumab formulations.

Sample name		Sum Aggregates	Monomer	Sum Fragments
35	T0, HCF1, 25°C	0,42	99,50	0,09
	T0, HCF1, 37°C	0,43	99,46	0,11
	T0, HC F3, 25°C	0,39	99,54	0,07
	T0, HCF3, 37°C	0,41	99,50	0,09
40	T0, HCF4, 25°C	0,43	99,46	0,11
	T0, HCF4, 37°C	0,42	99,48	0,11
	T0, commercial, 25°C	0,36	99,55	0,09
	T0, commercial, 37°C	0,35	99,56	0,09
45	T1, HC F1, 25°C	0,43	99,47	0,10
	T1, HC F1, 37°C	0,44	99,48	0,08
	T1, HC F3, 25°C	0,38	99,53	0,09
50	T1, HC F3, 37°C	0,37	99,54	0,09
	T1, HC F4, 25°C	0,44	99,47	0,09
	T, HC F4, 37°C	0,44	99,46	0,10
	T1, commercial, 25°C	0,35	99,56	0,08
55	T1, commercial, 37°C	0,35	99,56	0,10
	T3, HC F1, 25°C	0,42	99,47	0,10

(continued)

Sample name	Sum Aggregates	Monomer	Sum Fragments
T3, HC F1, 37°C	0,42	99,48	0,11
T3, HC F3, 25°C	0,40	99,48	0,12
T3, HC F3, 37°C	0,40	99,52	0,08
T3, HC F4, 25°C	0,48	99,41	0,11
T3, HC F4, 37°C	0,44	99,48	0,08
T3, commercial, 25°C	0,36	99,54	0,10
T3, commercial, 37°C	0,34	99,55	0,11
T5, HC F1, 25°C	0,43	99,48	0,09
T5, HC F1, 37°C	0,45	99,45	0,10
T5, HC F3, 25°C	0,41	99,48	0,11
T5, HC F3, 37°C	0,39	99,48	0,13
T5, HC F4, 25°C	0,47	99,43	0,10
T5, HC F4, 37°C	0,49	99,40	0,11
T5, commercial, 25°C	0,36	99,56	0,08
T5, commercial, 37°C	0,40	99,47	0,13

[0357] Subvisible particle counts for $\geq 10 \mu\text{m}$ and ≥ 25 both were very low. Freeze/thaw cycling did not lead to an increased number of subvisible particles, indicating that the tested formulations had favorable stability.

Size Exclusion Chromatography (SEC)

[0358] The SEC results are shown in Table 17. Table 17 indicates the percentages of SEC aggregates, monomers, and fragments in each of the solutions at T0 (before any freeze-thaw cycles), T1 (after one freeze-thaw cycle), T3 (after three freeze-thaw cycles) and T5 (after five freeze-thaw cycles). These results indicate that each of formulations 1, 3, and 4 show stabilities similar to that of the commercial formulation.

Table 17: Percentages of aggregates, monomers, and fragments before and after freeze-thaw cycles as assessed by SEC

Ion Exchange Chromatography (IEC)

[0359] Ion exchange chromatography did not reveal different sensitivity of the tested solutions. No significant degradation could be observed.

[0360] However, with increasing number of freeze/thaw cycles, the samples that were thawed at 25°C showed a higher amount in the 2nd acidic region after 5 cycles.

[0361] IEC results are shown in Table 18.

Table 18: IEC measurements before and after freeze-thaw cycles

Sample name	Sum Lysin Peaks	1 ac area	2 ac area	Lysin 0	Lysin 1	Peak between	Lysin 2
T0, HC F1, 25°C	86,74	2,09	10,51	68,63	16,72	0,99	0,40
T0, HC F3, 25°C	87,29	1,89	10,22	66,35	16,16	0,91	3,87
T0, HC F4, 25°C	87,35	1,85	10,20	66,46	16,15	0,89	3,85
T0, commercial, 25°C	87,18	1,93	10,22	66,25	16,12	0,94	3,88
T1, HC F1, 25°C	87,17	1,98	10,18	66,24	16,11	0,94	3,87
T1, HC F3, 25°C	87,30	1,81	10,24	66,39	16,13	0,90	3,88

EP 2 637 690 B1

(continued)

	Sample name	Sum Lysin Peaks	1 ac area	2 ac area	Lysin 0	Lysin 1	Peak between	Lysin 2
5	T1, HC F4, 25°C	87,21	1,88	10,25	66,31	16,11	0,90	3,88
	T1, commercial, 25°C	87,19	2,01	10,20	66,27	16,11	0,93	3,88
	T3, HC F1, 25°C	87,23	1,94	10,20	66,34	16,12	0,91	3,87
	T3, HC F3, 25°C	87,27	1,86	10,25	66,37	16,14	0,88	3,88
10	T3, HC F4, 25°C	87,26	1,82	10,27	66,34	16,15	0,88	3,88
	T3, commercial, 25°C	87,20	1,88	10,28	66,29	16,11	0,91	3,89
	T5, HC F1, 25°C	87,39	1,74	10,21	66,43	16,18	0,89	3,88
15	T5, HC F3, 25 °C	87,27	1,79	10,32	66,42	16,15	0,84	3,86
	T5, HC F4, 25°C	87,33	1,69	10,32	66,49	16,14	0,85	3,85
	T5, commercial, 25°C	87,05	1,95	10,38	66,17	16,10	0,89	3,88
	T0, HC F1, 37°C	87,25	1,94	10,19	66,36	16,13	0,90	3,86
20	T0, HC F3, 37°C	87,42	1,82	10,19	66,54	16,16	0,85	3,87
	T0, HC F4, 37°C	87,38	1,89	10,11	66,52	16,14	0,86	3,86
	T0, commercial, 37°C	87,25	1,86	10,24	66,34	16,12	0,91	3,88
25	T1, HC F1, 37°C	87,21	1,98	10,17	66,27	16,11	0,95	3,88
	T1, HC F3, 37°C	87,32	1,91	10,12	66,40	16,13	0,90	3,88
	T1, HC F4, 37°C	87,27	1,98	10,13	66,36	16,14	0,91	3,86
	T1, commercial, 37°C	87,28	1,97	10,14	66,34	16,12	0,93	3,88
30	T3, HC F1, 37°C	87,27	1,95	10,15	66,36	16,11	0,94	3,86
	T3, HC F3, 37°C	87,18	2,03	10,11	66,29	16,13	0,90	3,87
	T3, HC F4, 37°C	87,21	1,95	10,15	66,35	16,09	0,92	3,85
35	T3, commercial, 37°C	87,31	1,95	10,21	66,50	16,09	0,91	3,81
	T5, HC F1, 37°C	87,29	2,01	10,06	66,40	16,11	0,93	3,85
	T5, HC F3, 37°C	87,25	2,07	10,06	66,37	16,10	0,92	3,87
	T5, HC F4, 37°C	87,28	2,04	10,02	66,42	16,11	0,93	3,83
40	T5, commercial, 37°C	87,53	1,91	10,02	66,72	16,11	0,88	3,83

EXAMPLE 5: STABILITY OF HIGH CONCENTRATION ANTI-TNFA ANTIBODY FORMULATIONS AGAINST STIR STRESS

[0362] The following example describes a study which examined the stability of the F1, F3, and F4 formulations using the stir-stress test. Each formulation was tested in a range of pH levels.

Materials

- Humira HC F1 pH 4.2
100 mg/mL pH 4.7
 pH 5.7
 pH 6.2
- Humira HC F3 pH 4.2
100 mg/mL pH 4.7
 pH 5.7

(continued)

pH 6.2

- 5
- Humira HC F4 pH 4.2
100 mg/mL pH 4.7
pH 5.7
pH 6.2

10 *Procedure*

[0363] The vials, stir bars, and stoppers were steam sterilized prior to use.

[0364] The stirring experiment was performed with the following experimental set-up:

- 15
- Protein solutions: Humira HC F1, F3, F4, each at pH 4.2, 4.7, 5.7, 6.2 100mg/mL, Humira HC F3 pH 5.2 100 mg/mL, Humira from Vetter 50 mg/mL,
 - 5 mL filling volume per 6R vial
 - n = 3 → 2x stirred (with 7x2 mm magnetic bar), 1 unstirred control (without magnetic bar)
 - magnetic stirrer multipoint HP: 550rpm
 - ambient temperature
 - sample pull: t = 0, t = 1 h, t = 4 h, t = 24 h, t = 48 h
- 20

[0365] Three 6R vials were filled with 5 mL for each protein solution and closed with stoppers. Two of them were equipped with a magnetic stir bar.

25 [0366] The vials were kept at 5°C over night. The next morning the samples (one per protein solution, because in the beginning they were all the same), were measured with the turbidity meter. The measured solutions were filled back in the vials prior start of the experiment. After 1, 4, 24, and 48 h samples were taken and the turbidity was determined.

[0367] The unstirred samples were only measured at the time points 0 and 48 h.

30 [0368] For Humira HC F3 pH 5.2 also subvisible particles were determined for all time points.

Results

Turbidity

35 [0369] The turbidity results for samples subjected to stir stress for 0, 1, 4, 24, or 48 hours, as well as a 48 hour unstirred control, are shown in Table 19.

40 **Table 19: Turbidity (NTU) of samples subjected to stir stress**

sample	0 h	1 h	4 h	24 h	48 h
Commercial Adalimumab	20.90	23.90	31.20	98.05	176.00
Humira HC F3 pH 5.2	6.13	6.69	8.92	18.05	29.50
Humira HC F1 4.2	8.62	8.89	9.40	6.48	15.05
Humira HC F1 4.7	14.00	15.50	20.05	10.88	81.40
Humira HC F1 5.7	30.70	33.25	36.40	23.40	100.40

Humira HC F1 6.2	38.00	40.95	52.60	32.65	168.00
Humira HC F3 4.2	3.20	3.35	3.72	4.88	6.69
Humira HC F3 4.7	4.81	5.20	6.09	9.54	18.70
Humira HC F3 5.7	8.75	10.03	11.30	25.90	46.10
Humira HC F3 6.2	9.24	-	13.05	22.60	37.30
Humira HC F4 4.2	3.44	3.74	3.80	6.48	9.79
Humira HC F4 4.7	5.13	5.67	6.60	10.88	17.00
Humira HC F4 5.7	9.23	10.15	12.50	23.40	32.20
Humira HC F4 6.2	10.30	11.65	15.55	32.65	56.75

[0370] Increased pH correlated with increased turbidity for all tested formulations, both TO/unstirred and stirred samples. This effect was most pronounced for formulation 1. Also, formulation 1 showed the highest increase of turbidity after 48h at all pH values except 4.2. Formulation 3 and 4 showed similar behavior and turbidity values were comparable at all time points.

[0371] Humira HC (100 mg/mL), F3, pH 5.2 showed only a slight increase of the turbidity over the time. In contrast, the commercial Humira solution showed both a significantly higher starting value and increase in turbidity over the time. Thus, formulation 3 showed lower turbidity than the commercial Humira formulation.

[0372] The stirred samples showed a higher turbidity compared with the unstirred controls. The turbidity of the unstirred controls remained almost constant in comparison to the 0 h samples, indicating that running the experiment at room temperature did not bias the results.

Subvisible Particles

[0373] Table 20 shows the results for the numbers of subvisible particles.

Table 20: Counts of subvisible particles before and after stirring stress

	subvisible particles		
Humira HC F3 pH 5.2	$\geq 1 \mu\text{m}$	$\geq 10 \mu\text{m}$	$\geq 25 \mu\text{m}$
0 h	103	3	1
1 h	194	4	0
4h	202	4	0
24 h	262	2	0
48 h	192	3	0
48 h unstirred	80	1	0

Particles $\geq 1 \mu\text{m}$

[0374] Stirring induced a slight increase in sub-visible particle counts $\geq 1 \mu\text{m}$. The unstirred control was comparable to the 0 h sample.

Particles $\geq 10 \mu\text{m}$

[0375] Stirring had no significant effect on the particle counts $\geq 10 \mu\text{m}$. The unstirred control was comparable to the 0 h sample.

Particles $\geq 25 \mu\text{m}$

[0376] Stirring had no significant effect on the particle counts $\geq 25 \mu\text{m}$. The unstirred control was comparable to the 0 h sample.

[0377] Overall, the results of the experiments presented in Example 5 showed that formulation 3, when subjected to stirring stress, was surprisingly stable compared with commercial Humira solution. Formulation 3 was robust to stirring

stress according to the turbidity measure, and stirring of formulation 3 also had little or no effect on formation of subvisible particles.

EXAMPLE 6: LONG-TERM STORAGE STABILITY OF HIGH CONCENTRATION ANTI-TNFA ANTIBODY

[0378] The following example describes a study which examined the long-term storage stability (up to 24 months) of the F1, F3, and F4 formulations.

[0379] Formulations F1, F3, and F4 were tested prior to long term storage (Initial), and after 3, 6, 9, 12, 18, and 24 months of storage. The following storage conditions were used: (1) 5°C, (2) 25°C/60% relative humidity (R.H.), and (3) 40°C/75% R.H.. During storage, the samples were packaged in a 1 ml pre-filled syringe (colorless, glass type I, Ph.Eur.); BD Hypak Syringe BD 260 with a grey DB Hypak 4023/50 Fluorotec stopper. The following measures were used to assess storage stability: particulate contamination: visible particles; clarity and opalescence; color of solution (visual); *in vitro* TNF α -neutralization; cation exchange chromatography (CEX-HPLC), size exclusion chromatography (SE-HPLC); particulate contamination - sub-visible particles; container closure integrity; pH; and microbial quality.

[0380] All formulations tested were stable under the tested storage conditions of 2-8°C for up

RESULTS

[0381] The results for Formulation F1 are presented in Table 21.

Table 21: Stability Summary Report for Formulation F1

\Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/ % r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
Particulate contamination: visible particles	Visible particles	NMT 4.5	Initial	0.0	0.0	0.0
			3 months	0.0	0.0	0.0
			6 months	0.0	0.1	-
			9 months	0.0	-	-
			12 months	0.0	-	-
			18 months	0.0	-	-
			24 months	0.0	-	-
Clarity and Opalescence	Assessment	Not more opalescent than reference suspension IV	Initial	<= RS III	<= RS III	<= RS III
			3 months	<= RS III	<= RS III	<= RS IV
			6 months	<= RS III	<= RS III	<= RS IV
			9 months	<= RS IV	-	-
			12 months	<= RS III	-	-
			18 months	RS III	-	-
			24 months	<= RS III	-	-
Color of solution (visual)	BY-Scale	Report value	Initial	<= BY 7	<= BY 7	<= BY 7
			3 months	<= BY 7	<= BY 7	<= BY 7
			6 months	<= BY 7	<= BY 7	<= BY 6
			9 months	<= BY 7	-	-

EP 2 637 690 B1

(continued)

5	\\Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/ % r.H.]		
					5°C	25°C/60% R.H.	40°C/75% R.H.
10				12 months	<= BY 7	-	-
				18 months	<= BY 7	-	-
				24 months	<= BY 7	-	-
15	In vitro TNF-Neutralization	(cytotoxicity test) [%]	80 % to 125 % of the neutralization capacity of the reference standard	Initial	99	99	99
				3 months	97	110	97
				6 months	87	81	68
				9 months	88	-	-
				12 months	110	-	-
				18 months	97	-	-
				24 months	111	-	-
		Fiducial Limit of error (p=0.95) lower Limit [%]	NLT 64	Initial	96.1	96.1	196.1
				3 months	91.1	104.7	93.2
				6 months	84.2	76.3	64.4
				9 months	84.3	-	-
				12 months	105.2	-	-
				18 months	95.3	-	-
				24 months	108.9	-	-
		Fiducial Limit of error (p=0.95) upper Limit [%]	NMT 156	Initial	101.3	101.3	101.3
				3 months	101.9	115.5	100.1
				6 months	90.8	85.9	-
				9 months	90.8	-	71.2
				12 months	114.2	-	-
				18 months	98.8	-	-
				24 months	113.5	-	-
55	Cation Exchange Chromatography (CEX-HPLC)	Sum of lysine variants [%]	NLT 75	Initial	86.8	86.8	86.8
				3 months	86.2	74.7	26.0
				6 months	85.9	65.0	11.9
				9 months	85.2	-	-

EP 2 637 690 B1

(continued)

\\Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/ % r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
		24	12 months	85.2	-	-
			18 months	84.1	-	-
			24 months	83.9	-	-
Size exclusion chromatography (SE-HPLC)	Principal peak (monomer) [%]	NLT 98	Initial	99.6	99.6	99.6
			3 months	99.5	99.0	96.4
			6 months	99.4	98.5	92.9
			9 months	99.4	-	-
			12 months	99.4	-	-
			18 months	99.3	-	-
			24 months	99.3	-	-
Particulate contamination - Sub-visible Particles*	Particles >= 10 µm [/container]	NMT 6000	Initial	11	11	11
			3 months	8	37	55
			6 months	33	102	98
			9 months	32	-	-
			12 months	58	-	-
			18 months	44	-	-
	Particles >= 25 µm [/container]	NMT 600	24 months	11		-
			Initial	0	0	0
			3 months	0	0	1
			6 months	0	2	2
			9 months	0	-	-
			12 months	0	-	-
			18 months	1	-	-
			24 months	0	-	-
Container Closure Integrity	Assessment	Tight	Initial	Complies	Complies	Complies
			6 months	Complies	Complies	Complies
			12 months	Complies	-	-

EP 2 637 690 B1

(continued)

\Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/ % r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
			18 months	Complies	-	-
			24 months	Complies	-	-
pH	Single values	4.7 to 5.7	Initial	5.3	5.3	5.3
			3 months	5.2	5.2	5.2
			6 months	5.3	5.3	5.3
			9 months	5.3	-	-
			12 months	5.3	-	-
			18 months	5.3	-	-
			24 months	5.3	-	-
Microbial quality	Sterility drug product	No evidence of microbial growth is found	Initial	Complies	Complies	Complies

[0382] The results described above show that when stored for 24 months at 5°C, Formulation F1 showed no visible particulate contamination, clarity and opalescence ≤ RS III, and visual color ≤ BY7(brown yellow 7). Formulation F1 also demonstrated 111% of the TNF-neutralization capacity of the reference standard, 83.9% lysine variants, 99.3% monomers, 11 particles ≥10 μm, and no particles ≥25 μm. Furthermore, F1 maintained a stable pH of 5.3 and showed no evidence of Microbial growth. When stored for 6 months at 25°C/60% R.H., Formulation F1 showed 0.1 visible particles, clarity and opalescence ≤ RS III, visual color ≤ BY7, 81% of the TNF-neutralization capacity of the reference standard, 65% lysine variants, 98.5% monomers, 102 particles ≥10 μm, 2 particles ≥25 μm, a stable pH of 5.3 and no evidence of microbial growth. When stored for 6 months at 45°C/75% R.H., Formulation F1 showed no visible particles, clarity and opalescence ≤ RS IV, visual color ≤ BY6, 68% of the TNF-neutralization capacity of the reference standard, 11.9% lysine variants, 92.9% monomer, 98 particles ≥10 μm, 2 particles ≥25 μm, and no evidence of microbial growth.

[0383] The results for Formulation F3 are presented in Table 22.

Table 22: Stability Summary Report for Formulation F3

Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/% r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
Particulate contamination: visible particles	Visible particles	NMT 4.5	Initial	0.0	0.0	0.0
			3 months	0.0	0.0	0.2
			6 months	0.2	0.1	0.0
			9 months	0.0	-	-
			12 months	0.0	-	-
			18 months	0.0	-	-

EP 2 637 690 B1

(continued)

Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/% r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
			24 months	0.0	-	-
Clarity and Opalescence	Assessment	Not more opalescent than reference suspension IV	Initial	<= RS II	<= RS II	<= RS II
			3 months	<= RS II	<= RS II	<= RS II
			6 months	<= RS II	<= RS II	<= RS II
			9 months	<= RS II	-	-
			12 months	<= RS II	-	-
			18 months	<= RS II	-	-
			24 months	<= RS II	-	-
Color of solution (visual)	BY-Scale	Report value	Initial	<= BY 7	<= BY 7	<= BY 7
			3 months	<= BY 7	<= BY 7	<= BY 7
			6 months	<= BY 7	<= BY 7	<= BY 6
			9 months	<= BY 7	-	-
			12 months	<= BY 7	-	-
			18 months	<= BY 7	-	-
			24 months	<= BY 7	-	-
In vitro TNF-Neutralization	(cytotoxicity test) [%]	80 % to 125 % of the neutralization capacity of the reference standard	Initial	87	87	87
			3 months	101	106	89
			6 months	100	101	90
			9 months	98	-	-
			12 months	96	-	-
			18 months	96	-	-
			24 months	98	-	-
	Fiducial Limit of error (p=0.95) lower Limit [%]	NLT 64	Initial	85.4	85.4	85.4
			3 months	92.9	88.1	80.6
			6 months	98.3	97.4	86.5
			9 months	97.0	-	-
			12 months	93.3	-	-
			18 months	93.9	-	-

EP 2 637 690 B1

(continued)

Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/% r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
	Fiducial Umit of error (p=0.95) upper Limit [%]	NMT 156	24 months	96.7	-	-
			Initial	88.5	88.5	88.5
			3 months	110.5	122.2	97.9
			6 months	101.7	103.8	92.6
			9 months	99.8	-	-
			12 months	99.2	-	-
			18 months	98.1	-	-
			24 months	99.6	-	-
Cation Exchange Chromatography (CEX-HPLC)	Sum of lysine variants [%]	NLT 75	Initial	86.8	86.8	86.8
			3 months	86.6	77.8	32.8
			6 months	86.4	70.1	16.5
			9 months	86.0	-	-
			12 months	86.2	-	-
			18 months	85.2	-	-
Size exclusion chromatography (SE-HPLC)	Principal peak (monomer) [%]	NLT 98	24 months	85.1	-	-
			Initial	99.7	99.7	99.7
			3 months	99.6	99.2	96.9
			6 months	99.5	98.8	93.8
			9 months		-	-
			12 months	99.5	-	-
			18 months	99.4	-	-
			24 months	99.4	-	-
Particulate contamination - Sub-visible Particles*	Particles >= 10 µm [/container]	NMT 6000	Initial	10	10	10
			3 months	12	45	173
			6 months	22	157	275
			9 months	50	-	-
			12 months	54	-	-
			18 months	45	-	-

EP 2 637 690 B1

(continued)

Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/% r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
	Particles >= 25 µm [/container]	NMT 600	24 months	14	-	-
			Initial	0	0	
			3 months	0	0	1
			6 months	0	2	9
			9 months	0	-	-
			months	1	-	
			12 18 months	0	-	- -
			24 months	0	-	
			Initial	Complies	Complies	Complies
			6 months	Complies	Complies	Complies
			24 months	Complies	-	-
			Initial	5.2	5.2	5.2
	Single values	4.7 to 5.7	3 months	5.3	5.3	5.3
			6 months	5.2	5.2	5.3
			9 months	5.3	-	-
			12 months	5.4	-	-
			18 months	5.2	-	-
			24 months	5.1	-	-
Microbial quality	Sterility drug product	No evidence of microbial growth is found	Initial	Complies	Complies	Complies

[0384] The results provided in Table 22 indicate that when stored for 24 months at 5°C, Formulation F3 showed no visible particulate contamination, clarity and opalescence <= RS II, and visual color <= BY7. Formulation F3 showed 98% of the TNF-neutralization capacity of the reference standard, 85.1% lysine variants, 99.4% monomers, 14 particles >=10 µm, and no particles >=25 µm. The pH showed little change and there was no evidence of microbial growth.

[0385] When stored for 6 months at 25°C/60% R.H., Formulation F3 showed no visible particles, clarity and opalescence <= RS II, and visual color <= BY7. Also, formulation F3 showed 101% of the TNF-neutralization capacity of the reference standard, 97.4% lysine variants, 70.1% monomers, 157 particles >=10 µm, and 2 particles >=25 µm. The pH was stable and there was no evidence of microbial growth.

[0386] When stored for 6 months at 45°C/75% R.H., Formulation F3 showed no visible particles, clarity and opalescence <= RS II, and visual color <= BY6. Also, formulation F3 showed 90% of the TNF-neutralization capacity of the reference standard, 16.5% lysine variants, 93.8% monomers, 275 particles >=10 µm, and 9 particles >=25 µm. The pH was quite stable, and there was no evidence of microbial growth.

[0387] The results for Formulation F4 are presented in Table 23.

Table 23: Stability Summary Report for Formulation F4

Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/ % r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
Appearance	Visual particles	NMT 4.5	Initial	0.0	0.0	0.0
			3 months	0.0	0.0	0.0
			6 months	0.0	0.0	0.0
			9 months	0.0	-	-
			12 months	0.0	-	-
			18 months	0.0	-	-
Clarity	Assessment	Not more opalescent than reference suspension IV	Initial	<= RS II	<= RS II	<= RS II
			3 months	<= RS II	<= RS II	<= RS II
			6 months	<= RS II	<= RS II	<= RS II
			9 months	<= RS II	-	-
			12 months	<= RS II	-	-
			18 months	<= RS II	-	-
Color	BY-Scale		Initial	<= BY 7	<= BY 7	<= BY 7
			3 months	<= BY 6	<= BY 6	<= BY 6
			6 months	<= BY 7	<= BY 7	<= BY 6
			9 months	<=	BY 7	
			12 months	<= BY 7	-	-
			18 months	<= BY 7	-	-
In vitro TNF-Neutralisation (Cytotoxizitätstest)	(cytotoxicity test) [%]	80% to 125% of the neutralization capacity of the reference standard	Initial	111	111	111
			3 months	105	101	80
			6 months	97	101	76
			9 months	112	-	-
			12 months	97	-	-
			18 months	104	-	-

EP 2 637 690 B1

(continued)

Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/ % r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
Fiducial Limit of error (p=0.95) lower Limit [%]	NLT 64		Initial	105.2	105.2	105.2
			3 months	103.2	100.1	79.2
			6 months	92.9	97.5	74.7
			9 months	109.3	-	-
			12 months	90.2	-	-
			18 months	101.2	-	-
	Fiducial Limit of error (p=0.95) upper Limit [%]	NMT 156	Initial	116.3	116.3	116.3
			3 months	106.2	102.7	80.4
			6 months	101.1	103.5	78.1
			9 months	113.9	-	-
			12 months	104.9	-	-
			18 months	107.5	-	-
Cation Exchange Chromatography (CEX-HPLC)	Sum of lysine variants [%]	NLT 75	Initial	85.5	85.5	85.5
			3 months	85.8	76.8	31.6
			6 months	85.4	68.7	15.7
			9 months	85.2	-	-
			12 months	84.5	-	-
			18 months	84.4	-	-
Size exclusion chromatography (SE-HPLC)	Principal peak (monomer) [%]	NLT 98	Initial	99.7	99.7	99.7
			3 months	99.6	99.1	96.5
			6 months	99.6		93.1
			9 months	99.5	-	-
			12 months	99.5	-	-
			18 months	99.4	-	-
Particulate contamination - Sub-visible Particles	Particles >= 10 µm [/container]	NMT 6000	Initial	17	17	17
			3 months	51	174	207
			6 months	39	144	218
			9 months	82	-	
			12 months	57	-	-

(continued)

Test Item	Component	Specification	Duration of Testing	Storage Conditions [°C/ % r.H.]		
				5°C	25°C/60% R.H.	40°C/75% R.H.
	Particles $\geq 25 \mu\text{m}$ [/container]	NMT 600	Initial	0	0	0.5
			3 months	0	1	
			6 months	0	1	1
			9 months	1	-	-
			12 months	2	-	-
Container closure Integrity	Assessment	Must comply (no blue coloration)	Initial	Complies	1 Complies	Complies
			6 months	Complies	Complies	Complies
pH	Single values	4.7 to 5.7	Initial	5.1	5.1	5.1
			3 months	5.2	5.2	5.1
			6 months	5.2	5.1	5.2
			9 months	5.2	-	
			months	5.2	-	-
			12 months	5.1	-	-
Microbial quality	Sterility drug product	No evidence of microbial growth is found	Initial	Complies	Complies	Complies

[0388] The results provided in Table 23 indicate that when stored for 18 months at 5°C, Formulation F4 showed no visible particulate contamination, clarity and opalescence $\leq$ RS II, and visual color $\leq$ BY7. Formulation F4 showed 104% of the TNF-neutralization capacity of the reference standard, 84.4% lysine variants, and 99.4% monomers. Furthermore, the pH was stable and there was no evidence of microbial growth.

[0389] When stored for 6 months at 25°C/60% R.H., Formulation F4 showed no visible particles, clarity and opalescence $\leq$ RS II, and visual color $\leq$ BY7. Formulation F4 showed 101% of the TNF-neutralization capacity of the reference standard, 68.7% lysine variants, 98.8% monomers, 144 particles $\geq 10 \mu\text{m}$, and 1 particle $\geq 25 \mu\text{m}$. Furthermore, the pH was quite stable and there was no evidence of microbial growth.

[0390] When stored for 6 months at 45°C/75% R.H., Formulation F4 showed no visible particles, clarity and opalescence $\leq$ RS II, and visual color $\leq$ BY6. Formulation F4 showed 76% of the TNF-neutralization capacity of the reference standard, 15.7% lysine variants, 93.1% monomers, 218 particles $\geq 10 \mu\text{m}$, and 1 particle $\geq 25 \mu\text{m}$. Furthermore, the pH was quite stable and there was no evidence of microbial growth.

[0391] In summary, the results of the long-term stability experiments, as presented in Tables 21-23, show that high concentration formulations F1, F3, and F4 were surprisingly stable when subjected to long term storage. The stability of these formulations was similar to the commercial formulation. Formulations F1 and F3 showed stability similar to the commercial formulation after long term storage for at least 24 months. Formulation F4 showed stability similar to the commercial formulation after long term storage for at least 18 months.

EXAMPLE 7: ROOM TEMPERATURE STORAGE STABILITY OF HIGH CONCENTRATION ANTI-TNFA ANTIBODY

[0392] Liquid pharmaceutical products containing therapeutic antibodies often require storage at 2-8°C until end-of-shelf-life. Cooling is therefore also required by patients between purchasing of the medicines until use. Depending on the proposed dosing regimen, this can result in storage times under patient's responsibility in the case of self-administration drugs for up to several weeks.

[0393] Therefore, drugs that do not require storage under refrigerated conditions display both a significant increase in patient convenience for home care products and reduction of drug quality concerns in case of improper storage,

EP 2 637 690 B1

thereby reducing complaint rates and temperature excursion investigations.

[0394] The currently marketed Adalimumab containing product (Humira) was successfully reformulated at a higher protein concentration as Formulation F3, as described above in Examples 1-6. The following stability data for Formulation F3 resulted in findings of improved stability against protein degradation. The resulting degradation kinetics measured at 25 °C complied with requirements for ambient storage for up to 3 months.

[0395] For general long-term stability data related to storage at 25°C ofr Formulation F3, see Example 6 above.

[0396] The following data describes long-term storage characteristics for Formulation F3. The data shows that even after 18 months and 24 months of long-term storage at 2-5 °C, additional storage at 25 °C / 30°C is acceptable.

Table 24: 24m storage of F3 at 2-8 °C, followed by 7 days / 14 days at accelerated conditions (25°C, 30°C)

Test criterion	Specification	Characteristic	10 (24h 5°C)	+ 7 Days		+ 14 Days	
				25°C	30°C / 65% R.H.	25°C	30°C / 65% R.H.
Appearance	colourless to slightly yellow solution		complies	complies	complies	complies	complies
Visible particles*	single vial ≤ 2=> practically free from visible particulate matter 1 vial > 10=>		0.0	Value per analyst: 3 x 0; single vial ≤ 2=> practically free from visible particulate matter	Value per analyst: 3 x 0; single vial ≤ 2=> practically free from visible particulate matter	Value per analyst: 3 x 0; single vial ≤ 2=> practically free from visible particulate matter	Value per analyst: 3 x 0; single vial ≤ 2=> practically free from visible particulate matter
Size exclusion HPLC	Purity% monomer NLT 98%	aggregate	0.4	0.4	0.4	0.4	0.4
		monomer	99.4	99.3	99.3	99.3	99.2
		fragments	0.2	0.2	0.3	0.3	0.3
Cation exchange HPLC	NMT 8%	first acid region	2.7	2.8	3.0	2.9	3.3
	NMT 16%	second acid region	10.5	10.9	11.4	11.6	12.7
	NLT 75%	sum Lysine variants	85.1	84.2	83.4	83.0	83.2
	NMT 4%	peak between Lysine 1 and Lysine 2	1.5	1.7	1.9	1.6	1.8
PCS	report value [%]	peaks after Lysine 2	0.2	0.4	0.4	0.9	1.0
	report value [%]	Z-Average	1.390	1.365	1.395	1.397	1.420
		PdI	0.193	0.176	0.188	0.175	0.210

Table 25: 18m storage at 2-8 °C of F3, followed by 7 days / 14 days at accelerated conditions (25°C, 30°C)

Test criterion	Specification	Characteristic	ID (10115 °C)	+ 7 Days		+ 14 Days	
				25 °C	30 °C / 85% R.H.	25 °C	30 °C / 85% R.H.
Appearance	colourless to slightly yellow solution			complies	complies	complies	complies
Visible particles*	single vial ≤2⇒ practically free from visible particulate matter 1 vial > 10⇒ inform lab manager			Value per analyst:3 x 0; single vial ≤2⇒ practically free from visible particulate matter	Value per analyst:3 x 0; single vial ≤2⇒ practically free from visible particulate matter	Value per analyst:3 x 0; single vial ≤2⇒ practically free from visible particulate matter	Value per analyst:3 x 0; single vial ≤2⇒ practically free from visible particulate matter
Particulate contamination, subvisible particles*	particles ≤ 10 µm: ≤ 6000	≤ 1 µm / 1ml		2250	5623	9355	10252
	particles/container particles ≤ 25 µm: ≤ 600	≤ 10 µm / 1ml		6	7	39	45
	particles/container	≤ 25 µm / 1ml		0	0	0	1
PCS	report value [%]	Z-Average		2.378	2.344	2.353	2.358
		PdI		0.102	0.077	0.077	0.077

EXAMPLE 8: CONDUCTIVITY OF HIGH CONCENTRATION ANTI-TNFA ANTIBODY FORMULATIONS

[0397] The conductivity of the high concentration anti-TNF α antibody formulations F3 and F4 (see Examples 1-6, *supra*) was determined using an InoLab Cond Level2 WTW device normalized to 25°C. Table 26 shows the influence of non-ionic excipients on the conductivity of the F3 and F4 adalimumab formulations.

Table 26. Conductivity of Formulations F3 and F4

Sample	Temperature [°C]	Conductivity [μ S/cm]
Adalimumab DP F3		
1	22.4	663
2	22.4	651
3	23.8	660
4	21.4	715
5	21.7	691
6	23.1	680
7	23.3	644
8	22.9	647
Adalimumab DP F4		
1	22.0	797
2	22.9	746

[0398] As described above in Table 26, average conductivity for both formulations F3 and F4 was less than 2 mS/cm.

EXAMPLE 9: DYNAMIC LIGHT SCATTERING (DLS) OF HIGH CONCENTRATION ANTI-TNFA ANTIBODY FORMULATIONS

[0399] Dynamic light scattering analysis of diluted solutions was used to assess the hydrodynamic diameter (reported as the mean or Z-average size, calculated by cumulants analysis of the DLS measured intensity autocorrelation function and polydispersity index, PDI, of the size distribution of particles). DLS measurements were specifically used to detect low amounts of higher molecular weight species, e.g. aggregates, in a size distribution, since those species possess higher scattering intensity (proportional to d^6) and, therefore, will influence the Z-average and Polydispersity Index (PDI) as an indicator of the Z-average size distribution significantly.

[0400] A 150 μ L sample of each of formulations F3 and F4 (see examples 1-6 above) was measured to analyze the average size of the particles (Z-average) and the Polydispersity Index (PDI), an indicator of the "broadness" of the particle size distribution using DLS. The results are shown below. DLS data did not show any signs of the development of higher molecular weight aggregates, since the polydispersity index, a sensitive indicator for low levels of higher molecular weight sub-populations did not increase significantly.

<u>Formulation F3</u>		
Sample No.	ZAve (nm)	PDI
1	2.4	n.a.
2	2.3	0.08
3	2.3	0.14
4	2.3	0.09

<u>Formulation F4</u>		
Sample No.	ZAve (nm)	PDI
1	1.3	n.a.
2	2.5	n.a.

[0401] As described above, the z-average measurement for both F3 and F4 was less than 4 nm. This low hydrodynamic diameter is representative of the fact that both formulations F3 and F4 do not contain additional excipients other than a polysorbate and a polyol or a polysorbate.

EXAMPLE 10: FACTORS INFLUENCING THE STABILITY OF HIGH CONCENTRATION ANTI-TNFA ANTIBODY FORMULATIONS

[0402] The effect of varying mannitol concentrations and polysorbate concentrations on the stability of adalimumab in water was examined.

[0403] Formulations containing 100 mg/ml of adalimumab in water were prepared. Subsequently, various concentrations of either mannitol or polysorbate were added in a concentration range to determine the impact of each excipient on the stability of the formulation, as measured by aggregation and fragmentation. The concentrations of polysorbate and mannitol ranged from 0.1 to 1.0 mg/ml and 0-72 mg/ml, respectively, as shown in Figures 3A and 3B. As shown in Figure 3A, varying the concentration of mannitol from about 12 to about 72 mg/ml had a minimal effect on the stability of adalimumab. Similarly, varying the concentration of polysorbate-80 from about 0.1 to about 1.0 mg/ml had no effect on the stability of adalimumab.

[0404] The practice of the present invention will employ, unless otherwise indicated, conventional techniques of protein formulations, which are well known in the art.

EQUIVALENTS

[0405] The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting of the invention described herein. Scope of the invention is thus indicated by the appended claims rather than by the foregoing description.

SEQUENCE LISTING

[0406]

<110> Abbott Biotechnology Ltd. et al.

<120> Improved High Concentration Anti-TNF-alpha Antibody Liquid Formulation

<130> 117813-01020

<140> Concurrently filed herewith

<141> 2011-11-11

<150> US 61/412728

<151> 2010-11-11

<150> US 61/413960

<151> 2010-11-15

<160> 53

<170> PatentIn Ver. 2.0

<210> 1

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> human antibody

<400> 1

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asn Tyr
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Ala Ala Ser Thr Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Val Ala Thr Tyr Tyr Cys Gln Arg Tyr Asn Arg Ala Pro Tyr
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

15 <210> 2
 <211> 121
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <223> human antibody

<400> 2

25 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30
 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 30 35 40 45
 Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
 50 55 60
 Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
 100 105 110
 40 Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

45 <210> 3
 <211> 9
 <212> PRT
 <213> Artificial Sequence

50 <220>
 <223> human antibody

55 <220>
 <221> VARIANT
 <222> (9)
 <223> Xaa = Thr or Ala
 <400> 3

EP 2 637 690 B1

Gln Arg Tyr Asn Arg Ala Pro Tyr Xaa
1 5

5 <210> 4
<211> 12
<212> PRT
<213> Artificial Sequence

10 <220>
<223> human antibody

<220>
<221> VARIANT
<222> (12)
15 <223> Xaa = Tyr or Asn

<400> 4

20 Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Xaa
1 5 10

<210> 5
<211> 7
<212> PRT
25 <213> Artificial Sequence

<220>
<223> human antibody

30 <400> 5

Ala Ala Ser Thr Leu Gln Ser
1 5

35 <210> 6
<211> 17
<212> PRT
<213> Artificial Sequence

40 <220>
<223> human antibody

45 <400> 6

Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val Glu
1 5 10 15
Gly

50 <210> 7
<211> 11
<212> PRT
<213> Artificial Sequence

55 <220>
<223> human antibody

<400> 7

EP 2 637 690 B1

Arg Ala Ser Gln Gly Ile Arg Asn Tyr Leu Ala
1 5 10

5 <210> 8
<211> 5
<212> PRT
<213> Artificial Sequence

10 <220>
<223> human antibody

<400> 8

Asp Tyr Ala Met His
1 5

15 <210> 9
<211> 107
<212> PRT
20 <213> Artificial Sequence

<220>
<223> 2SD4 light chain variable region

25 <400> 9

30 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Ile Gly
1 5 10
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asn Tyr
20 25 30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45
Tyr Ala Ala Ser Thr Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
35 65 70 75 80
Glu Asp Val Ala Thr Tyr Tyr Cys Gln Lys Tyr Asn Ser Ala Pro Tyr
85 90 95
Ala Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
100 105

40 <210> 10
<211> 121
<212> PRT
<213> Artificial Sequence

45 <220>
<223> 2SD4 heavy chain variable region

50 <400> 10

55

EP 2 637 690 B1

5 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30
 10 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Asp Trp Val
 35 40 45
 Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
 50 55 60
 Glu Gly Arg Phe Ala Val Ser Arg Asp Asn Ala Lys Asn Ala Leu Tyr
 65 70 75 80
 15 Leu Gln Met Asn Ser Leu Arg Pro Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Thr Lys Ala Ser Tyr Leu Ser Thr Ser Ser Ser Leu Asp Asn Trp Gly
 100 105 110
 Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 11

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> 2SD4 light chain variable region CDR3

<400> 11

Gln Lys Tyr Asn Ser Ala Pro Tyr Ala
 1 5

<210> 12

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> EP B12 light chain variable region CDR3

<400> 12

Gln Lys Tyr Asn Arg Ala Pro Tyr Ala
 1 5

<210> 13

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> VL10E4 light chain variable region CDR3

<400> 13

Gln Lys Tyr Gln Arg Ala Pro Tyr Thr
 1 5

<210> 14

<211> 9

<212> PRT

EP 2 637 690 B1

<213> Artificial Sequence

<220>

<223> VL100A9 light chain variable region CDR3

5

<400> 14

Gln Lys Tyr Ser Ser Ala Pro Tyr Thr
1 5

10

<210> 15

<211> 9

<212> PRT

<213> Artificial Sequence

15

<220>

<223> VLL100D2 light chain variable region CDR3

<400> 15

20

Gln Lys Tyr Asn Ser Ala Pro Tyr Thr
1 5

<210> 16

25

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

30

<223> VLL0F9 light chain variable region CDR3

<400> 16

Gln Lys Tyr Asn Arg Ala Pro Tyr Thr
1 5

35

<210> 17

<211> 9

<212> PRT

40

<213> Artificial Sequence

<220>

<223> LOE5 light chain variable region CDR3

45

<400> 17

Gln Lys Tyr Asn Ser Ala Pro Tyr Tyr
1 5

50

<210> 18

<211> 9

<212> PRT

<213> Artificial Sequence

55

<220>

<223> VLLOG7 light chain variable region CDR3

<400> 18

EP 2 637 690 B1

Gln Lys Tyr Asn Ser Ala Pro Tyr Asn
1 5

5 <210> 19
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 <220>
 10 <223> VLLOG9 light chain variable region CDR3
 <400> 19

Gln Lys Tyr Thr Ser Ala Pro Tyr Thr
1 5

15 <210> 20
 <211> 9
 <212> PRT
 20 <213> Artificial Sequence
 <220>
 <223> VLLOH1 light chain variable region CDR3
 25 <400> 20

Gln Lys Tyr Asn Arg Ala Pro Tyr Asn
1 5

30 <210> 21
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 35 <220>
 <223> VLLOH10 light chain variable region CDR3
 <400> 21

Gln Lys Tyr Asn Ser Ala Ala Tyr Ser
1 5

40 <210> 22
 <211> 9
 45 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> VL1B7 light chain variable region CDR3
 50 <400> 22

Gln Gln Tyr Asn Ser Ala Pro Asp Thr
1 5

55 <210> 23
 <211> 9
 <212> PRT

<213> Artificial Sequence

<220>

<223> VL1C1 light chain variable region CDR3

5

<400> 23

Gln Lys Tyr Asn Ser Asp Pro Tyr Thr
1 5

10

<210> 24

<211> 9

<212> PRT

<213> Artificial Sequence

15

<220>

<223> VL0.1F4 light chain variable region CDR3

<400> 24

20

Gln Lys Tyr Ile Ser Ala Pro Tyr Thr
1 5

<210> 25

25

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

30

<223> VL0.1H8 light chain variable region CDR3

<400> 25

Gln Lys Tyr Asn Arg Pro Pro Tyr Thr
1 5

35

<210> 26

<211> 9

40

<212> PRT

<213> Artificial Sequence

<220>

<223> LOE7.A light chain variable region CDR3

45

<400> 26

Gln Arg Tyr Asn Arg Ala Pro Tyr Ala
1 5

50

<210> 27

<211> 12

<212> PRT

<213> Artificial Sequence

55

<220>

<223> 2SD4 heavy chain variable region CDR3

<400> 27

Ala Ser Tyr Leu Ser Thr Ser Ser Ser Leu Asp Asn
1 5 10

5

<210> 28

<211> 12

<212> PRT

<213> Artificial Sequence

10

<220>

<223> VH1B11 heavy chain variable region CDR3

<400> 28

15

Ala Ser Tyr Leu Ser Thr Ser Ser Ser Leu Asp Lys
1 5 10

<210> 29

20

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

25

<223> VH1D8 heavy chain variable region CDR3

<400> 29

Ala Ser Tyr Leu Ser Thr Ser Ser Ser Leu Asp Tyr
1 5 10

30

<210> 30

<211> 12

<212> PRT

35

<213> Artificial Sequence

<220>

<223> VH1A11 heavy chain variable region CDR3

40

<400> 30

Ala Ser Tyr Leu Ser Thr Ser Ser Ser Leu Asp Asp
1 5 10

45

<210> 31

<211> 12

<212> PRT

<213> Artificial Sequence

50

<220>

<223> VH1B12 heavy chain variable region CDR3

<400> 31

Ala Ser Tyr Leu Ser Thr Ser Phe Ser Leu Asp Tyr
1 5 10

55

<210> 32

EP 2 637 690 B1

<211> 12
<212> PRT
<213> Artificial Sequence

5 <220>
 <223> VH1E4 heavy chain variable region CDR3

 <400> 32

10 Ala. Ser Tyr Leu Ser Thr Ser Ser Ser Leu His Tyr
1 5 10

<210> 33
<211> 12
<212> PRT
<213> Artificial Sequence

20

<220>
<223> VH1F6 heavy chain variable region CDR3
<400> 33

Ala Ser Phe Leu Ser Thr Ser Ser Ser Leu Glu Tyr
1 5 10

<210> 34
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> 3C-H2 heavy chain variable region CDR3
<400> 34

Ala Ser Tyr Leu Ser Thr Ala Ser Ser Leu Glu Tyr
1 5 10

<210> 35
<211> 12
<212> PRT
<213> Artificial Sequence

45 <220>
<223> VH1-D2.N heavy chain variable region CDR3
<400> 35

50 Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Asn
1 5 10

<210> 36
<211> 321
<212> DNA
<213> Artificial Sequence

<220>
<223> adalimumab light chain variable region

<400> 36

5

gacatccaga	tgacccagtc	tccatcctcc	ctgtctgcat	ctgtagggga	cagagtcacc	60
atcacttgtc	gggcaagtca	gggcatcaga	aattacttag	cctgggtatca	gcaaaaacca	120
gggaaagccc	ctaagctcct	gatctatgct	gcattcactt	tgcaatcagg	gggtcccatct	180
cggttcagtg	gcagtggatc	tgggacagat	ttcactctca	ccatcagcag	cctacagcct	240
gaagatgttg	caacttatta	ctgtcaaagg	tataaccgtg	caccgtatac	ttttggccag	300
gggaccaagg	tggaaatcaa	a				321

10

<210> 37
<211> 363
<212> DNA
<213> Artificial Sequence

15 <220>
<223> adalimumab heavy chain variable region

<400> 37

20	gaggtgcagc	tggtggagtc	tgggggaggc	ttggtacagc	ccggcaggtc	cctgagactc	60
	tcctgtgcgg	cctctggatt	cacctttgat	gattatgcc	tgcactgggt	ccggcaagct	120
	ccaggaagg	gcctggaatg	ggtctcagct	atcacttgga	atagtgggtca	catagactat	180
	gcggactctg	tggaggcccg	attcaccatc	tccagagaca	acgccaaaga	ctccctgtat	240
	ctgcaaata	acagttctgag	agctgaggat	acggccgtat	attactgtgc	gaaagtctcg	300
25	taccttagca	ccgcgtcctc	ccttgactat	tggggccaaag	gtaccctggt	caccgtctcg	360
	agt						363

30

<210>	38
<211>	6
<212>	PRT
<213>	Homo sapiens

<400> 38

35 His Gly Ser His Asp Asn
1 5

40

<210>	39
<211>	12
<212>	PRT
<213>	Homo sapiens

<400> 39

45 Gln Ser Tyr Asp Arg Tyr Thr His Pro Ala Leu Leu
 1 5 10

50

<210>	40
<211>	17
<212>	PRT
<213>	Homo sapiens

<400> 40

55 Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15
Gly

<210> 41
 <211> 7
 <212> PRT
 <213> Homo sapiens

5

<400> 41

Tyr Asn Asp Gln Arg Pro Ser
 1 5

10

<210> 42
 <211> 9
 <212> PRT
 <213> Homo sapiens

15

<400> 42

Phe Thr Phe Ser Ser Tyr Gly Met His
 1 5

20

<210> 43
 <211> 13
 <212> PRT
 <213> Homo sapiens

25

<400> 43

Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn Thr Val Lys
 1 5 10

30

<210> 44
 <211> 115
 <212> PRT
 <213> Homo sapiens

35

<400> 44

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15

40

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

45

Ala Phe Ile Arg Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

50

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Lys Thr His Gly Ser His Asp Asn Trp Gly Gln Gly Thr Met Val Thr
 100 105 110

55

Val Ser Ser
 115

EP 2 637 690 B1

<210> 45
 <211> 112
 <212> PRT
 <213> Homo sapiens

5

<400> 45

10

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Arg Ser Asn Ile Gly Ser Asn
 20 25 30

15

Thr Val Lys Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

Ile Tyr Tyr Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

20

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln
 65 70 75 80

Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Arg Tyr Thr
 85 90 95

25

His Pro Ala Leu Leu Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
 100 105 110

<210> 46
 <211> 16
 <212> PRT
 <213> Homo Sapiens

30

<400> 46

35

Ser Ile His Asn Arg Gly Thr Ile Phe Tyr Leu Asp Ser Val Lys Gly
 1 5 10 15

<210> 47
 <211> 10
 <212> PRT
 <213> Homo Sapiens

40

<400> 47

45

Gly Arg Ser Asn Ser Tyr Ala Met Asp Tyr
 1 5 10

<210> 48
 <211> 16
 <212> PRT
 <213> Homo Sapiens

50

<400> 48

55

Arg Ser Thr Gln Thr Leu Val His Arg Asn Gly Asp Thr Tyr Leu Glu
 1 5 10 15

<210> 49
 <211> 7
 <212> PRT
 <213> Homo Sapiens
 5
 <400> 49

 Lys Val Ser Asn Arg Phe Ser
 1 5
 10
 <210> 50
 <211> 9
 <212> PRT
 <213> Homo Sapiens
 15
 <400> 50

 Phe Gln Gly Ser His Val Pro Tyr Thr
 1 5
 20
 <210> 51
 <211> 30
 <212> PRT
 <213> Homo Sapiens
 25
 <400> 51

 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala Ser Val Lys
 Val
 1 5 10 15
 20
 Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr
 25 30
 35
 <210> 52
 <211> 13
 <212> PRT
 <213> Homo Sapiens
 40
 <400> 52

 Trp Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met Gly
 1 5 10
 45
 <210> 53
 <211> 32
 <212> PRT
 <213> Homo Sapiens
 50
 <400> 53

 55

Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr Met Glu Leu Ser Ser
 Leu
 1 5 10 15
 20
 5 Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
 25 30

Claims

1. A liquid aqueous formulation consisting essentially of:

- (1) 100 mg/ml of adalimumab;
- (2) 1 mg/ml of polysorbate-80;
- (3) 42 mg/ml of mannitol; and
- (4) water,

wherein the formulation does not contain a buffer.

2. The liquid aqueous formulation of claim 1, wherein the formulation consists of:

- (1) 100 mg/ml of adalimumab;
- (2) 1 mg/ml of polysorbate-80;
- (3) 42 mg/ml of mannitol; and
- (4) water.

3. The liquid aqueous formulation of claim 1 or 2, wherein the formulation is stable up to about 30°C for at least 6 days.

4. The liquid aqueous formulation of any one of claims 1-3, wherein the formulation is stable up to about 30°C for at least 10 days.

5. The liquid aqueous formulation of any one of claims 1-4 having a characteristic selected from the group consisting of:

- a) a conductivity of less than about 2 mS/cm;
- b) a hydrodynamic diameter (D_h) which is at least about 50% less than the D_h of the protein in a buffered solution at a given concentration; and
- c) a hydrodynamic diameter (D_h) of less than about 4 nm.

6. A pre-filled syringe or autoinjector device, comprising the formulation of any one of claims 1-5.

7. The formulation of any one of claims 1-5 for use in treating rheumatoid arthritis, Crohn's disease, psoriatic arthritis, psoriasis, juvenile idiopathic arthritis, ankylosing spondylitis, ulcerative colitis, or hidradenitis suppurativa.

Patentansprüche

1. Flüssige wässrige Formulierung, die im Wesentlichen aus Folgendem besteht:

- (1) 100 mg/ml Adalimumab;
- (2) 1 mg/ml Polysorbat-80;
- (3) 42 mg/ml Mannit und
- (4) Wasser,

wobei die Formulierung keinen Puffer enthält.

2. Flüssige wässrige Formulierung nach Anspruch 1, wobei die Formulierung aus Folgendem besteht:

- (1) 100 mg/ml Adalimumab;
- (2) 1 mg/ml Polysorbat-80;
- (3) 42 mg/ml Mannit und
- (4) Wasser.

3. Flüssige wässrige Formulierung nach Anspruch 1 oder 2, wobei die Formulierung für mindestens 6 Tage bis zu etwa 30° C stabil ist.
4. Flüssige wässrige Formulierung nach einem der Ansprüche 1 bis 3, wobei die Formulierung für mindestens 10 Tage bis zu etwa 30° C stabil ist.
5. Flüssige wässrige Formulierung nach einem der Ansprüche 1 bis 4, die ein Merkmal aufweist, das aus der Gruppe ausgewählt ist, die aus Folgendem besteht:
 - a) einer Leitfähigkeit von weniger als etwa 2 mS/cm;
 - b) einem hydrodynamischen Durchmesser (D_h), der mindestens etwa 50 % geringer ist als der D_h des Proteins in einer gepufferten Lösung bei einer gegebenen Konzentration; und
 - c) einem hydrodynamischen Durchmesser (D_h) von weniger als etwa 4 nm.
6. Vorgefüllte Spritze oder Autoinjektorvorrichtung, die die Formulierung nach einem der Ansprüche 1 bis 5 umfasst.
7. Formulierung nach einem der Ansprüche 1 bis 5, zur Verwendung bei der Behandlung von rheumatoider Arthritis, Morbus Crohn, Psoriasis-Arthritis, Psoriasis, juveniler idiopathischer Arthritis, Spondylitis ankylosans, Colitis ulcerosa oder Acne inversa.

Revendications

1. Formulation aqueuse liquide constituée essentiellement de :

- (1) 100 mg/ml d'adalimumab ;
- (2) 1 mg/ml de polysorbate-80 ;
- (3) 42 mg/ml de mannitol ; et
- (4) d'eau,

la formulation ne contenant pas de tampon.

2. Formulation aqueuse liquide selon la revendication 1, la formulation étant constituée de :

- (1) 100 mg/ml d'adalimumab ;
- (2) 1 mg/ml de polysorbate-80 ;
- (3) 42 mg/ml de mannitol ; et
- (4) d'eau.

3. Formulation aqueuse liquide selon la revendication 1 ou 2, la formulation étant stable jusqu'à environ 30 °C pendant au moins 6 jours.
4. Formulation aqueuse liquide selon l'une quelconque des revendications 1 à 3, la formulation étant stable jusqu'à environ 30 °C pendant au moins 10 jours.
5. Formulation aqueuse liquide selon l'une quelconque des revendications 1 à 4, ayant une caractéristique sélectionnée dans le groupe constitué de :

- a) une conductivité inférieure à environ 2 mS/cm ;
- b) un diamètre hydrodynamique (D_h) qui est au moins environ 50 % inférieur au D_h de la protéine dans une solution tamponnée à une concentration donnée ; et
- c) un diamètre hydrodynamique (D_h) inférieur à environ 4 nm.

EP 2 637 690 B1

6. Seringue pré-remplie ou dispositif auto-injecteur, comprenant la formulation selon l'une quelconque des revendications 1 à 5.

5 7. Formulation selon l'une quelconque des revendications 1 à 5, destinée à une utilisation dans le traitement de la polyarthrite rhumatoïde, de la maladie de Crohn, de l'arthrite psoriasique, du psoriasis, de l'arthrite juvénile idiopathique, de la spondylarthrite ankylosante, de la colite ulcéreuse, ou de l'hydrosadénite.

10

15

20

25

30

35


40

45

50

55

Figure 1



A = High Conc. Formulation 1
B = High Conc. Formulation 3
C = High Conc. Formulation 4
D = Commercial Formulation

Figure 2




Figure 3A





Figure 3B



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 6090382 A [0086] [0088] [0094] [0146] [0152] [0153] [0155] [0158] [0191]
- US 6258562 B [0086] [0088] [0094] [0146] [0153] [0191] [0286] [0293]
- US 20060153846 A [0111] [0112] [0124]
- US 20100278822 A [0111] [0112]
- US 6509015 B [0153]
- US 5656272 A [0161]
- WO 0212502 A [0161]
- US 6593458 B [0161]
- US 6498237 B [0161]
- US 6451983 B [0161]
- US 6448380 B [0161]
- US 4816397 A, Boss [0162]
- US 5168062 A, Stinski [0171]
- US 4510245 A, Bell [0171]
- US 4968615 A, Schaffner [0171]
- US 4399216 A [0172]
- US 4634665 A [0172]
- US 5179017 A [0172]
- WO 2008154543 A [0183]
- WO 2008005315 A [0188]
- WO 2010127146 A [0188]
- WO 2006000785 A [0188]
- WO 2011075524 A [0188]
- WO 2005113039 A [0188]
- US 6015557 A [0191]
- US 6177077 B [0191]
- US 6379666 B [0191]
- US 6419934 B [0191]
- US 6419944 B [0191]
- US 6423321 B [0191]
- US 6428787 B [0191]
- US 6537549 B [0191]
- WO 0050079 A [0191]
- WO 0149321 A [0191]
- US 20040126372 A [0191] [0286] [0293]
- WO 9216221 A [0194]
- WO 9217583 A [0194]
- WO 9311793 A [0194]
- EP 374510 A [0194]
- WO 9520978 A [0195]
- WO 9408609 A [0196]
- EP 585705 A [0198]
- EP 230574 A [0198]
- WO 9104054 A [0201]
- EP 453898 A [0203]
- WO 9420139 A [0203]
- US 5231024 A, Moeller [0204]
- US 260610 B1, Moeller, A [0204]
- US 20030235585 A [0287]
- US 20060009385 A [0288] [0289]
- US 20090317399 A [0288]
- WO 2007120823 A [0289]
- US 20090271164 A [0290]
- US 7541031 B [0293]
- US 61412728 B [0406]
- US 61413960 B [0406]

Non-patent literature cited in the description

- **WANG et al.** *J. Pharm. Sci.*, 2007, vol. 96, 1 [0001]
- **SHIRE et al.** *J. Pharm. Sci.*, 2004, vol. 93, 1390 [0003]
- **SINGER ; THODS.** *Academic Emergency Medicine*, 1998, vol. 5, 1007 [0066] [0118]
- **WEWERS ; LOWE.** A critical review of visual analogue scales in the measurement of clinical phenomena. *Research in Nursing and Health*, 1990, vol. 13, 227-236 [0067]
- **Peptide and Protein Drug Delivery.** Marcel Dekker, Inc, 1991, 247-301 [0078]
- **JONES, A.** *Adv. Drug Delivery Rev.*, 1993, vol. 10, 29-90 [0078]
- **PENNICA, D. et al.** *Nature*, 1984, vol. 312, 724-729 [0085]
- **DAVIS, J. M. et al.** *Biochem*, 1987, vol. 26, 1322-1326 [0085]
- **JONES, E. Y. et al.** *Nature*, 1989, vol. 338, 225-228 [0085]
- **CHOTHIA et al.** *Mol. Biol.*, 1987, vol. 196, 901-917 [0087]
- **CHOTHIA et al.** *Nature*, 1989, vol. 342, 877-883 [0087]
- **PADLAN.** *FASEB J.*, 1995, vol. 9, 133-139 [0087]
- **MACCALLUM.** *J. Mol. Biol.*, 1996, vol. 262 (5), 732-45 [0087]
- **WARD et al.** *Nature*, 1989, vol. 341, 544-546 [0088]
- **BIRD et al.** *Science*, 1988, vol. 242, 423-426 [0088] [0169]
- **HUSTON et al.** *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 5879-5883 [0088]

- **HOLLIGER, P. et al.** *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 6444-6448 [0088]
- **POLJAK, R.J.** *Structure*, 1994, vol. 2, 1121-1123 [0088]
- **TAYLOR et al.** *Nucl. Acids Res.*, 1992, vol. 20, 6287-6295 [0089]
- **JONSSON, U. et al.** *Ann. Biol. Clin.*, 1993, vol. 51, 19-26 [0096]
- **JONSSON, U. et al.** *Biotechniques*, 1991, vol. 11, 620-627 [0096]
- **JOHNSON, B. et al.** *J. Mol. Recognit.*, 1995, vol. 8, 125-131 [0096]
- **JOHNSON, B. et al.** *Anal. Biochem.*, 1991, vol. 198, 268-277 [0096]
- **LEE JS et al.** *Acad Emerg Med*, 2000, vol. 7, 550 [0118]
- **Molecular Cloning; A Laboratory Manual.** Cold Spring Harbor, 1989 [0162]
- **Current Protocols in Molecular Biology.** Greene Publishing Associates, 1989 [0162]
- **KABAT, E.A. et al.** *Sequences of Proteins of Immunological Interest.* U.S. Department of Health and Human Services, 1991 [0163] [0167] [0168]
- **TOMLINSON, I.M. et al.** *The Repertoire of Human Germline VH Sequences Reveals about Fifty Groups of VH Segments with Different Hypervariable Loops.* *J. Mol. Biol.*, 1992, vol. 227, 776-798 [0163]
- **COX, J.P.L. et al.** *A Directory of Human Germ-line V78 Segments Reveals a Strong Bias in their Usage.* *Eur. J. Immunol.*, 1994, vol. 24, 827-836 [0163]
- **HUSTON.** *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 5879-5883 [0169]
- **MCCAFFERTY et al.** *Nature*, 1990, vol. 348, 552-554 [0169]
- **GOEDDEL.** *Gene Expression Technology: Methods in Enzymology.* Academic Press, 1990, vol. 185 [0171]
- **BOSS, M.A. ; WOOD, C. R.** *Immunology Today*, 1985, vol. 6, 12-13 [0173]
- **URLAUB ; CHASIN.** *Proc. Natl. Acad. Sci. USA*, 1980, vol. 77, 4216-4220 [0174]
- **R.J. KAUFMAN ; P.A. SHARP.** *Mol. Biol.*, 1982, vol. 159, 601-621 [0174]
- **TRACEY, K. J. ; CERAMI, A.** *Annu. Rev. Med.*, 1994, vol. 45, 491-503 [0193]
- **RUSSELL, D ; THOMPSON, R. C.** *Curr. Opin. Biotech.*, 1993, vol. 4, 714-721 [0193]
- **AREND, W. P. ; DAYER, J-M.** *Arth. Rheum.*, 1995, vol. 38, 151-160 [0196]
- **FAVA, R. A. et al.** *Clin. Exp. Immunol.*, 1993, vol. 94, 261-266 [0196]
- **GROM et al.** *Arthritis Rheum.*, 1996, vol. 39, 1703 [0196]
- **MANGGE et al.** *Arthritis Rheum.*, 1995, vol. 8, 211 [0196]
- **FIETZE, E. et al.** *Transplantation*, 1994, vol. 58, 675-680 [0198]
- **EASON, J. D. et al.** *Transplantation*, 1995, vol. 59, 300-305 [0199]
- **SUTHANTHIRAN, M. ; STROM, T. B.** *New Engl. J. Med.*, 1994, vol. 331, 365-375 [0199]
- **TRACY, K. J. et al.** *Science*, 1986, vol. 234, 470-474 [0202]
- **SUN, X-M. et al.** *J. Clin. Invest.*, 1988, vol. 81, 1328-1331 [0202]
- **MACDONALD, T. T. et al.** *Clin. Exp. Immunol.*, 1990, vol. 81, 301-305 [0202]
- **VAN DULLEMEN, H. M. et al.** *Gastroenterology*, 1995, vol. 109, 129-135 [0202]
- **MOELLER, A. et al.** *Cytokine*, 1990, vol. 2, 162-169 [0204]
- **PARTSCH et al.** *Ann Rheum Dis.*, 1998, vol. 57, 691 [0205]
- **RITCHLIN et al.** *J Rheumatol.*, 1998, vol. 25, 1544 [0205]
- **BRAUN et al.** *Arthritis Rheum.*, 1999, vol. 42 (10), 2039 [0207]
- **ZEIDLER et al.** *Rheum Dis Clin North Am.*, 1992, vol. 18, 187 [0208]
- **TAKEMATSU et al.** *Arch Dermatol Res.*, 1989, vol. 281, 398 [0209] [0210]
- **VICTOR ; GOTTLIEB.** *J Drugs Dermatol.*, 2002, vol. 1 (3), 264 [0209] [0210]
- **ASADULLAH et al.** *Br J Dermatol.*, 1999, vol. 141, 94 [0215]
- **SKLAVOUNOU et al.** *J Oral Pathol Med.*, 2000, vol. 29, 370 [0228]
- **REUSS-BORST et al.** *Br J Haematol.*, 1993, vol. 84, 356 [0229]
- **TUTUNCU Z et al.** *Clin Exp Rheumatol.*, 2002, vol. 20 (28), 146-51 [0231]
- **MACKIEWICZ Z et al.** *Clin Exp Rheumatol.*, 2003, vol. 21 (1), 41-8 [0231]
- **MUROTA H et al.** *Arthritis Rheum.*, 2003, vol. 48 (4), 1117-25 [0231]
- **SPIEGELMAN ; HOTAMISLIGIL.** *Cell*, 1993, vol. 73, 625 [0240]
- **CHU et al.** *Int J Obes Relat Metab Disord.*, 2000, vol. 24, 1085 [0240]
- **ISHII et al.** *Metabolism.*, 2000, vol. 49, 1616 [0240]
- **NAVARRO J.F. ; MORA C. ; MACA.** *Am J Kidney Dis.*, July 2003, vol. 42 (1), 53-61 [0242]
- **DAIMON M et al.** *Diabetes Care.*, July 2003, vol. 26 (7), 2015-20 [0242]
- **ZHANG M.** *J Tongji Med Univ.*, 1999, vol. 19 (3), 203-5 [0242]
- **BARBIERI M et al.** *Am J Hypertens.*, July 2003, vol. 16 (7), 537-43 [0242]
- **MCCALL, J. et al.** *Br. J. Surg.*, 1992, vol. 79, 1361-3 [0242]
- **BENJAFIELD et al.** *Diabetes Care*, 2001, vol. 24, 753 [0250]
- **QIANG, X. et al.** *Diabetologia*, 1998, vol. 41, 1321-6 [0250]

- PFEIFFER. *Horm Metab Res.*, 1997, vol. 29, 111 [0250]
- SCHOLZ et al. *Trends Microbiol.*, 2003, vol. 11, 171 [0255]
- LEE et al. *Hum Immunol.*, 2003, vol. 64, 614 [0257]
- NAVARRO et al. *Am J Kidney Dis.*, 2003, vol. 42, 53 [0257]
- DAIMON et al. *Diabetes Care*, 2003, vol. 26, 2015 [0257]
- ZHANG et al. *J Tongji Med Univ.*, 1999, vol. 19, 203 [0257]
- BARBIERI et al. *Am J Hypertens.*, 2003, vol. 16, 537 [0257]
- VENN et al. *Arthritis Rheum.*, 1993, vol. 36, 819 [0257]
- WESTACOTT et al. *J Rheumatol.*, 1994, vol. 21, 1710 [0257]
- DEVARAJ et al. *Circulation*, 2000, vol. 102, 191 [0260]
- HATTORI Y et al. *Cardiovasc Res.*, 2000, vol. 46, 188 [0260]
- CLAUSELL N et al. *Cardiovasc Pathol.*, 1999, vol. 8, 145 [0260]
- PIHLAJAMAKI J et al. *Obes Res.*, 2003, vol. 11, 912 [0261]
- BARBIERI et al. *Am J Hypertens*, 2003, vol. 16, 537 [0261]
- TSUDA et al. *J Nutr.*, 2003, vol. 133, 2125 [0261]
- DEGUCHI et al. *Lancet*, 1989, vol. 2, 745 [0262]
- SNELLER, M.C. *Cleve. Clin. J. Med.*, 2002, vol. 69, 40-3 [0268]
- SCHETT, G. et al. *Ann. Rheum. Dis.*, 2002, vol. 61, 463 [0268]
- STRAUB, R.H. et al. *Rheumatology*, 2002, vol. 41, 423 [0271]
- UDDHAMMAR, A. et al. *Br. J. Rheumatol.*, 1998, vol. 37, 766 [0271]
- KOBAYASHI, Y. ; NUMANO, F. *Intern. Med.*, 2002, vol. 41, 44 [0272]
- FRAGA, A. ; MEDINA F. *Curr. Rheumatol. Rep.*, 2002, vol. 4, 30 [0272]
- DIGIROLAMO, N. et al. *J. Leukoc. Biol.*, 1997, vol. 61, 667 [0276]
- SUNDEL, R.P. *Curr. Rheumatol. Rep.*, 2002, vol. 4, 474 [0279]
- GEDALIA, A. *Curr. Rheumatol. Rep.*, 2002, vol. 4, 25 [0279]
- SFIKAKIS, P.P. *Ann. Rheum. Dis.*, 2002, vol. 61, 51-3 [0282]
- DOGAN, D. ; FARAH, C. *Oftalmologia*, 2002, vol. 52, 23 [0282]
- MARQUEZ, J. et al. *Curr. Rheumatol. Rep.*, 2003, vol. 5, 128 [0283]
- HARMAN, L.E. ; MARGO, C.E. *Surv. Ophthalmol.*, 1998, vol. 42, 458 [0283]
- GROSS, W.L. *Curr. Opin. Rheumatol.*, 2002, vol. 14, 11 [0284]
- CHURG, W.A. *Mod. Pathol.*, 2001, vol. 14, 1284 [0284]
- BERTOLINI. D. R. et al. *Nature*, 1986, vol. 319, 516-518 [0285]
- KONIG, A. et al. *J. Bone Miner. Res.*, 1988, vol. 3, 621-627 [0285]
- LERNER, U. H. ; OHLIN, A. *J. Bone Miner. Res.*, 1993, vol. 8, 147-155 [0285]
- SHANLAR. G. ; STEM, P. H. *Bone*, 1993, vol. 14, 871-876 [0285]
- MCCLAIN, C. J. ; COHEN, D. A. *Hepatology*, 1989, vol. 9, 349-351 [0285]
- FELVER, M. E. *Alcohol. Clin. Exp. Res.*, 1990, vol. 14, 255-259 [0285]
- HANSEN, J. *Hepatology*, 1994, vol. 20, 461-474 [0285]
- SHERON, N. et al. *J. Hepatol.*, 1991, vol. 12, 241-245 [0285]
- HUSSAIN, M. J. et al. *J. Clin. Pathol.*, 1994, vol. 47, 1112-1115 [0285]
- VAN DER POLL, T. N. *Engl. J. Med.*, 1990, vol. 322, 1622-1627 [0285]
- VANDER POLL, T. et al. *Prog. Clin. Biol. Res.*, 1991, vol. 367, 55-60 [0285]
- GIROIR, B. P. *Am. J. Physiol.*, 1994, vol. 267, 118-124 [0285]
- LIU. X. S. *Bums*, 1994, vol. 20, 40-44 [0285]
- SCALES. W. E. et al. *Am. J Physiol.*, 1994, vol. 267, 1122-1127 [0285]
- SERRICK. C. *Transplantation*, 1994, vol. 58, 1158-1162 [0285]
- YAO, Y. M. et al. *Resuscitation*, 1995, vol. 29, 157-168 [0285]
- MCCAULEY, R. L. et al. *J. Clin. Immunol.*, 1992, vol. 12, 300-308 [0285]

Szabadalmi igénypontok

1. Folyékony vizes készítmény, amely alapvetően a következőkből áll:

- (1) 100 mg/ml adalimumab;
- (2) 1 mg/ml poliszorbát-80;
- (3) 42 mg/ml mannit; és
- (4) víz,

ahol a készítmény nem tartalmaz puffert.



2. Az 1. igénypont szerinti folyékony vizes készítmény, ahol a készítmény a következőkből áll:

- (1) 100 mg/ml adalimumab;
- (2) 1 mg/ml poliszorbát-80;
- (3) 42 mg/ml mannit; és
- (4) víz.

3. Az 1. vagy 2. igénypont szerinti folyékony vizes készítmény, ahol a készítmény 30°C-ig terjedő hőmérsékleten legalább 6 napig stabil.

4. Az 1-3. igénypontok bármelyike szerinti folyékony vizes készítmény, ahol a készítmény 30°C-ig terjedő hőmérsékleten legalább 10 napig stabil.

5. Az 1-4. igénypontok bármelyike szerinti folyékony vizes készítmény, amely a következő csoportból választott jellemzőkkel bír:

- a) a vezetőképesség kisebb, mint kb. 2 mS/cm;
- b) a hidrodinamikai átmérő (D_h) legalább kb. 50%-kal kisebb, mint a fehérje D_h értéke egy puffer oldatban egy adott koncentrációnál, és
- c) a hidrodinamikai átmérő (D_h) kisebb, mint kb. 4 nm.

6. Előretöltött fecskendő vagy autoinjektáló berendezés, amely tartalmazza az 1-5. igénypontok bármelyike szerinti készítményt.

7. Az 1-5. igénypontok bármelyike szerinti készítmény a következő kezelésekben történő alkalmazásra: reumás ízületi gyulladás, Crohn-betegség, pszoriázisos ízületi gyulladás, pszoriázis, ifjúkori idiopátiás ízületi gyulladás, ízületi merevséggel járó csigolyagyulladás, fekélyes vastagbélgyulladás vagy genyves verejtékmirigy-gyulladás.