




(86) Date de dépôt PCT/PCT Filing Date: 2009/07/21  
(87) Date publication PCT/PCT Publication Date: 2010/01/28  
(45) Date de délivrance/Issue Date: 2016/06/28  
(85) Entrée phase nationale/National Entry: 2011/01/07  
(86) N° demande PCT/PCT Application No.: US 2009/051217  
(87) N° publication PCT/PCT Publication No.: 2010/011630  
(30) Priorité/Priority: 2008/07/21 (US61/082,422)

(51) Cl.Int./Int.Cl. *C07C 43/23* (2006.01),  
*A23L 2/02* (2006.01), *A61K 8/33* (2006.01),  
*A61K 8/34* (2006.01), *A61K 8/49* (2006.01),  
*A61Q 19/00* (2006.01), *A61Q 19/02* (2006.01),  
*C07C 39/15* (2006.01), *C07D 213/30* (2006.01),  
*C07D 307/12* (2006.01), *C07D 307/44* (2006.01),  
*C07D 317/54* (2006.01)

(72) Inventeurs/Inventors:  
NANDY, SANDIP K., US;  
LIU, JIYUN, US;  
NESTEROV, ALEXANDRE MIKHAILOVITCH, US;  
HERTEL, CARMEN, US;  
PADMAPRIYA, ABEYSINGHE A., US

(73) Propriétaire/Owner:

(54) Titre : FAMILLE DE COMPOSES PERMETTANT LE BLANCHIMENT (ECLAIRCISSEMENT) DE LA PEAU  
(54) Title: SERIES OF SKIN-WHITENING (LIGHTENING) COMPOUNDS



Formula I

(57) Abrégé/Abstract:


The present invention provides compounds of formula I: (see formula I) or a pharmaceutically acceptable salt thereof, wherein: R is unsubstituted phenyl, substituted or unsubstituted biphenyl, substituted or unsubstituted pyrid-3-yl, substituted or unsubstituted pyrid-4-yl or substituted or unsubstituted furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group.

(73) Propriétaires(suite)/Owners(continued):UNIGEN, INC., US

(74) Agent: NORTON ROSE FULBRIGHT CANADA LLP/S.E.N.C.R.L., S.R.L.

**ABSTRACT**

The present invention provides compounds of formula I:



Formula I

or a pharmaceutically acceptable salt thereof, wherein: R is unsubstituted phenyl, substituted or unsubstituted biphenyl, substituted or unsubstituted pyrid-3-yl, substituted or unsubstituted pyrid-4-yl or substituted or unsubstituted furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group.

## SERIES OF SKIN-WHITENING (LIGHTENING) COMPOUNDS

### FIELD OF THE INVENTION

[0001] The present invention is directed to novel tyrosinase inhibitors, pharmaceutical, dermatologic and cosmetic compositions comprising such inhibitors, and methods of making and using the same.

### BACKGROUND OF THE INVENTION

[0002] The color of human skin is determined by the amount and the type of melanin produced by specialized cells, melanocytes, which are located in the basal layers of skin. Melanin, one of the most widely distributed natural pigments, is a homogeneous polyphenol-like biopolymer with a complex structure and color varying from brown to black (eumelanin) and red to yellow (pheomelanin) (Prota G. *Med. Res. Rev.* **1988**, 8, 525-556). In human skin, melanin is believed to act as a protective agent against ultraviolet radiation. As such, people living close to equator have darker skin than those living in areas away from the equator.

[0003] Overproduction of melanin can causes different types of abnormal skin color, hair color, and other dermatological disorders, such as melasma, age spots and sites of actinic damage. Melanin is produced through a series of oxidative reactions and polymerization steps starting with the amino acid tyrosine. Tyrosinase (EC1.14.18.1), a type III, copper-containing enzyme catalyzes two initial reactions in the production of melanin: 1) the *ortho*-hydroxylation of L-tyrosine by monophenolase action, and 2) the oxidation of 3,4-dihydroxyphenylalanine (L-DOPA)  $\rightarrow$  *o*-dopaquinone by diphenolase action. The later oxidation step is much more rapid than the former, thus, the hydroxylation of tyrosine is considered to be the rate-determining step in melanin biosynthesis. Subsequent conversion of *o*-dopaquinone to melanin occurs through a series of enzymatic and non-enzymatic polymerization reactions. Other enzymes, such as dopachrome tautomerase (Tyrosinase Related Protein 2; TRP-2) and dihydroxyindole carboxylic acid (DICA) oxidase (Tyrosinase Related Protein 1; TRP-1) are also involved in the process of the biosynthesis of melanin. Since tyrosinase plays the key role in the process of melanin production, inhibitors of this enzyme are often used as skin-whitening agents (Mosher et al. In *Dermatology in General Medicine*, **1983**, 205-125, Fitzpatrick T. B., Eisen A. Z., Wolff K., Freedberg I. M., Austern K. F. (eds), Mc-Graw-Hill, New York.; Maeda K., Fukuda M, In vitro effectiveness

of several whitening cosmetic components in human melanocytes. *J. Soc. Cosmet. Chem.* **1991**, 42, 361-368).

**[0004]** A number of naturally occurring, as well as, synthetic tyrosinase inhibitors have been described in the literature. The majority of compounds comprise a phenol structure. These compounds act as metal chelating agents (Kojima et al. *Biol. Pharma. Bull.* **1995**, 18, 1076-1078.; Seo et al. *J. Agric. Food Chem.* **2003**, 51, 2837-2853.; Fu et al. *J. Agric. Food Chem.* **2003**, 53, 7408-7414.; Kim, Y.-J.; Uyama, H. *Cell. Mol. Life Sci.* **2005**, 62, 1707-1723 and references cited therein). A pharmaceutical product containing hydroquinone (2-4%) is moderately efficacious, but hydroquinone is considered to be cytotoxic to melanocytes and potentially mutagenic to mammalian cells. Unfortunately, several purportedly active agents, e.g., arbutin and kojic acid, among others have not been demonstrated yet to be clinically efficacious when critically analyzed in carefully controlled studies (Frenk, E. In *Melasma: New Approaches to Treatment*; Martin Dunitz: London, **1995**, 9-15.; Dooley, T. P. In: *Drug Discovery Approaches for Developing Cosmeceuticals: Advanced Skin Care and Cosmetic Products*; Hori, W., Ed.; International business communications: Southborough, MA, **1997**; Dooley, T. P. *J. Dermatol. Treat.* **1997**, 7, 188-200).

**[0005]** Many tyrosinase inhibitors are resorcinol derivatives or polyphenol derivatives of flavonoids or of *trans*-stilbene, such as resveratrol or its derivatives. These types of compounds are known to form strong chelates with metal ions. (Seo et al. *J. Agric. Food Chem.* **2003**, 51, 2837-2853.; Fu et al. *J. Agric. Food Chem.* **2003**, 53, 7408-7414.; Kim, Y.-J.; Uyama, H. *Cell. Mol. Life Sci.* **2005**, 62, 1707-1723 and references cited therein.; Lerch, K. In: *Metal ions in Biological Systems*, pp.143-186; Sigel, H., Ed.; Marcel Dekker, NY, **1981**.; Wilcox et al. In: *Substrate analogue binding to the coupled binuclear copper active site in tyrosinase*, *J. Am. Chem. Soc.* **1985**, 107, 4015-4027.; Sanchez-Ferrer et al. *Biochim. Biophys. Acta* **1995**, 1247, 1-11.; Decker et al. *Angew. Chem. Int. Ed.* **2000**, 39, 1591-1595.; Decker et al. *Angew. Chem. Int. Ed.* **2006**, 45, 4546-4550.; Briganti et al. *Pigment Cell Res.* **2003**, 16, 101-110). Several compounds based on the resorcinol moiety have been used as tyrosinase inhibitors, see: JP 2008-056651 A; JP 2000-095721 A; US 2005/0267047A1; US 5,339,785; US 6,093,836; US 2008/0032938 A1; US 7,282,592 B2; US 7,339,076 B1; US 5,880,314; US 6,852,310 B2; US 6,077,503; US 2005/0271608 A1; US 5,523,421; US 2007/0098655 A1; US 2005/0267047 A1.

**[0006]** Although many compounds have been reported as potent tyrosinase inhibitors, very few of them have shown skin-whitener properties. Additionally, most of these agents

were found either to be toxic, or shown to have adverse side effects in humans. As such, the search for new natural products or synthetic compounds having potent tyrosinase inhibitory activity with low cytotoxicity continues.

## SUMMARY OF THE INVENTION

**[0007]** The instant application provides a new series of novel depigmenting agents having excellent inhibitory activity against tyrosinase and very low cytotoxicity. The compounds are stable and easily synthesized from the commonly available starting materials. More specifically, the present invention provides novel 2,4-dihydroxybenzene derivatives, which are useful as tyrosinase inhibitors. The invention encompasses pharmaceutically acceptable salts of these tyrosinase inhibitors. Also encompassed by the invention are pharmaceutical compositions comprised of at least one tyrosinase inhibitor of the invention and at least one pharmaceutically acceptable carrier. The compositions of the present invention can be prepared in any suitable pharmaceutically acceptable dosage form.

**[0008]** The present invention also provides a method for inhibiting the enzyme tyrosinase in a subject in need thereof, said method comprising administering an effective amount of a composition comprising at least one 2,4-dihydroxybenzene tyrosinase inhibitor of the instant invention.

**[0009]** The present invention further provides a method for the prevention and treatment of diseases and conditions related to the activity of the enzyme tyrosinase. The method of prevention and treatment according to this invention comprises administering internally or topically to a subject in need thereof a therapeutically effective amount of at least one novel 2,4-dihydroxybenzene tyrosinase inhibitor of the instant invention.

**[0010]** The present invention also provides a method for inhibiting the synthesis of melanin in a subject in need thereof. Such a method comprises administering an effective amount of a composition comprising at least one 2,4-dihydroxybenzene compound of the instant invention.

**[0011]** The present invention further provides methods for the prevention and treatment of diseases and conditions related to the overproduction or uneven distribution of melanin, said method comprising administering internally or topically to a subject in need thereof a therapeutically effective amount of at least one at least one 2,4-dihydroxybenzene compound of the instant invention. Diseases and conditions related to the overproduction or uneven distribution of melanin include, but not limited to hyper pigmentation spots caused

by skin aging, melasma, liver diseases, thermal burns and topical wounds, skin pigmentation due to inflammatory conditions caused by fungal, microbial and viral infections, vitilago, carcinoma, melanoma, as well as other mammalian skin conditions.

**[0012]** The method can also be used for preventing and treating skin darkening and damage resulting from exposure to the sun, ultraviolet (UV) radiation, chemicals, heat, wind and dry environments. Finally, the method can be used for preventing and treating wrinkles, saggy skin, lines and dark circles around the eyes, soothing sensitive skin and preventing and treating dermatitis and other allergy related conditions of the skin.


**[0013]** In specific embodiments, the dose of the 2,4-dihydroxybenzene tyrosinase inhibitor(s) of the instant invention administered to the subject in need thereof is an efficacious, nontoxic quantity generally selected from the range of 0.001% to 100% based on total weight of the final formulation, and/or 0.01 mg to 200 mg per kilogram based on the body weight of the subject. Persons skilled in the art using routine clinical testing are able to determine optimum doses for the particular ailment being treated. The compositions of this invention can be administered by any method known to one of ordinary skill in the art. The modes of administration include, but are not limited to, enteral (oral) administration, parenteral (intravenous, subcutaneous, and intramuscular) administration and topical application. The method of treatment according to this invention comprises administering internally or topically to a patient in need thereof a therapeutically effective amount of at least one 2,4-dihydroxybenzene of the instant invention. In a preferred embodiment the composition is administered topically.

**[0014]** Included in the present invention are novel methods for the synthesis of the 2,4-dihydroxybenzene tyrosinase inhibitor(s) of the invention as described in Examples 1-5. Thus, the present invention provides commercially viable options for the synthesis, and/or isolation, purification and formulation of the 2,4-dihydroxybenzenes of the instant invention to yield a composition of matter having the desirable physiological activity.

**[0015]** In yet another embodiment, the 2,4-dihydroxybenzenes can be used in the food industry to inhibit the browning and color changes in fruits, vegetables, juices and other food products.

**[0016]** Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below.

**[0016a]** According to one aspect of the present invention, there is provided a compound of formula I:




Formula I

or a pharmaceutically acceptable salt thereof, wherein:

R is unsubstituted phenyl, substituted or unsubstituted biphenyl, substituted or unsubstituted pyrid-3-yl, substituted or unsubstituted pyrid-4-yl or substituted or unsubstituted furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group.

**[0016b]** According to another aspect of the present invention, there is provided a compound of formula II:



Formula II


or a pharmaceutically acceptable salt thereof, wherein:

X is H, or OH; and

R is substituted or unsubstituted: phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl or furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group.

**[0016c]** According to still another aspect of the present invention, there is provided a compound selected from the group consisting of 4-(furan-2-ylmethyl)benzene-1,3-diol (5), 4-(furan-3-ylmethyl)benzene-1,3-diol (6), 4-(pyridin-ylmethyl)benzene-1,3-diol (7), 4-((2-methoxypyridin-3-yl)methyl)benzene-1,3-diol (8), 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9), 4-(benzo[d][1,3]dioxol-5-ylmethyl)benzene-1,3-diol (10) and 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11).

**[0016d]** According to yet another aspect of the present invention, there is provided a compound of Formula III:




or a pharmaceutically acceptable salt thereof, wherein:

$R_1$  is H or  $C_1$ - $C_{10}$  alkyl; and

$R_2$  is selected from the group consisting of 4-hydroxyphen-1-yl, 4-methoxyphen-1-yl, 2,4-dimethoxyphen-1-yl, 3,5-dimethoxyphen-1-yl, 2,4-dimethoxy-3-methylphen-1-yl, 2,4,6-trimethoxyphen-1-yl, furan-3-yl, tetrahydrofuran-3-yl, furan-2-yl, tetrahydrofuran-2-yl, 5-ethylfuran-2-yl and pyridine-3-yl.

**[0016e]** According to a further aspect of the present invention, there is provided a compound selected from the group consisting of 4-(4-hydroxyphenethylbenzene)-1,3-diol (13), 4-(4-methoxyphenethyl)benzene-1,3-diol (14), 4-(2,4-dimethoxyphenethyl)benzene-1,3-diol (17), 4-(3,5-dimethoxyphenethyl)benzene-1,3-diol (18), 4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol (19), 4-(2,4,6-trimethoxyphenethyl)benzene-1,3-diol (20), 4-(2-(furan-3-yl)ethyl)benzene-1,3-diol (22), 4-(2-(tetrahydrofuran-3-yl)ethyl)benzene-1,3-diol (23), 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol (24), 4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol (25), 4-(2-(5-ethylfuran-2-yl)ethyl)benzene-1,3-diol (26) and 4-(2-(pyridin-3-yl)ethyl)benzene-1,3-diol (27).

**[0016f]** According to yet a further aspect of the present invention, there is provided a compound of formula IV:



wherein  $\equiv$  represents a double or single bond;

$R_1$  is H or  $C_1$ - $C_{10}$  alkyl;

$R_4$  is H, OH or O; and

$R_5$  is a substituted or unsubstituted: phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl, furan-3-yl or tetrahydrofuran-2-yl; wherein, when substituted,  $R_5$  is

substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group with the following provisos:

proviso 1: when R<sub>1</sub> and R<sub>4</sub> are H; R<sub>5</sub> cannot be 1'-(2,4-dimethoxy-3-methyl)phenyl;


proviso 2: when R<sub>1</sub> and R<sub>4</sub> are H; R<sub>5</sub> cannot be 1'-(2,4-dihydroxy)phenyl; 1'-(2,4-dialkoxy)phenyl; 1'-(2,4-dialkenoxy)phenyl or 1'-(2,4-dibenzyloxy)phenyl;

proviso 3: R<sub>5</sub> cannot be 1'-(4-hydroxy)phenyl;

proviso 5: when R<sub>1</sub> and R<sub>4</sub> are H; R<sub>5</sub> cannot be a 1'-(2,4-disubstituted)phenyl; wherein said substituents are selected from -OR<sup>2</sup> and -OR<sup>3</sup>, wherein R<sup>2</sup> and R<sup>3</sup> are independently selected from H and a (C<sub>1</sub>-C<sub>10</sub>)-alkyl; and

proviso 6: when R<sub>1</sub> is H and R<sub>4</sub> is OH; R<sub>5</sub> cannot be 1'-(3,4-dihydroxy)phenyl; 1'-(3,4-dimethoxy)phenyl; 1'-(2-hydroxy)phenyl; 1'-(2-methoxy)phenyl; 1'-(4-methoxy)phenyl; phenyl; 1'-(3-methoxy,4-hydroxy)phenyl.

[0016g] According to still a further aspect of the present invention, there is provided a composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound of formula I:




Formula I

or a pharmaceutically acceptable salt thereof, wherein:

R is substituted or unsubstituted phenyl, substituted or unsubstituted biphenyl, substituted or unsubstituted pyrid-3-yl, substituted or unsubstituted pyrid-4-yl or substituted or unsubstituted furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group.

**[0016h]** According to another aspect of the present invention, there is provided a composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound of formula II:




Formula II

or a pharmaceutically acceptable salt thereof, wherein:

X is H, or OH; and

R is substituted or unsubstituted: phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl or furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group.

**[0016i]** According to yet another aspect of the present invention, there is provided a composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound of formula III:




Formula III

or a pharmaceutically acceptable salt thereof, wherein:

R<sub>1</sub> is H or C<sub>1</sub>-C<sub>10</sub> alkyl; and

R<sub>2</sub> is selected from the group consisting of phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl, furan-3-yl and tetrahydrofuran-2-yl; wherein R<sub>2</sub> is optionally substituted with 1 to 3 moieties (R', R'', R''') independently selected from the group consisting of a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group.

**[0016j]** According to still another aspect of the present invention, there is provided a composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound of formula IV:



Formula IV

wherein:

$\equiv$  represents a double or single bond;

R<sub>1</sub> is H or C<sub>1</sub>-C<sub>10</sub> alkyl;

R<sub>4</sub> is H, OH or O; and

R<sub>5</sub> is a substituted or unsubstituted: phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl, furan-3-yl or tetrahydrofuran-2-yl, and when substituted, R<sub>5</sub> is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group and a hydroxyl group;

with the following provisos:

proviso 1: when R<sub>1</sub> and R<sub>4</sub> are H; R<sub>5</sub> cannot be 1'-(2,4-dimethoxy-3-methyl)phenyl;

proviso 2: when R<sub>1</sub> and R<sub>4</sub> are H; R<sub>5</sub> cannot be 1'-(2,4-dihydroxy)phenyl; 1'-(2,4-dialkoxy)phenyl; 1'-(2,4-dialkenoxy)phenyl or 1'-(2,4-dibenzyloxy)phenyl;

proviso 3: R<sub>5</sub> cannot be 1'-(4-hydroxy)phenyl;

proviso 5: when R<sub>1</sub> and R<sub>4</sub> are H and R<sub>5</sub> cannot be a 1'-(2,4-disubstituted)phenyl;

wherein said substituents are selected from -OR<sup>2</sup> and -OR<sup>3</sup>, wherein R<sup>2</sup> and R<sup>3</sup> are independently selected from H and a (C<sub>1</sub>-C<sub>10</sub>)-alkyl; and

proviso 6: when R<sub>1</sub> is H and R<sub>4</sub> is OH; R<sub>5</sub> cannot be 1'-(3,4-dihydroxy)phenyl; 1'-(3,4-dimethoxy)phenyl; 1'-(2-hydroxy)phenyl; 1'-(2-methoxy)phenyl; 1'-(4-methoxy)phenyl; phenyl; or 1'-(3-methoxy,4-hydroxy)phenyl.

**[0016k]** According to a further aspect of the present invention, there is provided use of the composition described herein for inhibiting the activity of a tyrosinase.


**[0016l]** According to yet a further aspect of the present invention, there is provided use of the composition described herein for inhibiting the production of melanin.

**[0016m]** According to still a further aspect of the present invention, there is provided use of the composition described herein for preventing or treating diseases of conditions related to the overproduction or uneven distribution of melanin.

**[0016n]** According to another aspect of the present invention, there is provided use of the composition described herein for at least whitening or lightening skin.

[00160] According to yet another aspect of the present invention, there is provided use of a compound selected from the group consisting of:

a) a compound of formula I:




Formula I

or a pharmaceutically acceptable salt thereof, wherein:

R is substituted or unsubstituted phenyl, substituted or unsubstituted biphenyl, substituted or unsubstituted pyrid-3-yl, substituted or unsubstituted pyrid-4-yl or substituted or unsubstituted furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group;

b) a compound of formula II:




Formula II

or a pharmaceutically acceptable salt thereof, wherein:

X is H, or OH; and

R is substituted or unsubstituted: phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl or furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group,

c) a compound of formula III:



Formula III


or a pharmaceutically acceptable salt thereof, wherein:

R<sub>1</sub> is H or C<sub>1</sub>-C<sub>10</sub> alkyl; and

R<sub>2</sub> is phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl, furan-3-yl or tetrahydrofuran-2-yl; wherein R<sub>2</sub> is optionally substituted with 1 to 3 moieties (R', R'', R''') independently selected from the group consisting of a C1-C10 alkyl group, a C1-

C10 alkoxy group or a hydroxyl group, and

d) a compound of formula IV:



wherein:

$\equiv$  represents a double or single bond;

$R_1$  is H or  $C_1$ - $C_{10}$  alkyl;

$R_4$  is H, OH or O; and

$R_5$  is a substituted or unsubstituted: phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl, furan-3-yl or tetrahydrofuran-2-yl, and when substituted,  $R_5$  is substituted with 1 to 3 moieties ( $R'$ ,  $R''$ ,  $R'''$ ) independently selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group and a hydroxyl group;

with the following provisos:

proviso 1: when  $R_1$  and  $R_4$  are H;  $R_5$  cannot be 1'-(2,4-dimethoxy-3-methyl)phenyl;

proviso 2: when  $R_1$  and  $R_4$  are H;  $R_5$  cannot be 1'-(2,4-dihydroxy)phenyl; 1'-(2,4-dialkoxy)phenyl; 1'-(2,4-dialkenoxy)phenyl or 1'-(2,4-dibenzyloxy)phenyl;

proviso 3:  $R_5$  cannot be 1'-(4-hydroxy)phenyl;

proviso 5: when  $R_1$  and  $R_4$  are H and  $R_5$  cannot be a 1'-(2,4-disubstituted)phenyl;

wherein said substituents are selected from  $-OR^2$  and  $-OR^3$ , wherein  $R^2$  and  $R^3$  are independently selected from H and a  $(C_1-C_{10})$ -alkyl; and

proviso 6: when  $R_1$  is H and  $R_4$  is OH;  $R_5$  cannot be 1'-(3,4-dihydroxy)phenyl; 1'-(3,4-dimethoxy)phenyl; 1'-(2-hydroxy)phenyl; 1'-(2-methoxy)phenyl; 1'-(4-methoxy)phenyl; phenyl; or 1'-(3-methoxy,4-hydroxy)phenyl,

for inhibiting browning and color changes in fruits, vegetables, juices and food products.

[0017] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention as claimed. Other objects, advantages, and novel features will be readily apparent to those skilled in the art from the following detailed description.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0018] Figure 1 illustrates graphically a profile of the inhibition in the production of melanin (●) by 4-(pyridin-ylmethyl)benzene-1,3-diol (**7**) as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu\text{M}$ ). The  $\text{IC}_{50}$  of compound **7** was determined to be  $14 \mu\text{M}$ . Figure 1 also illustrates graphically cell viability ( $\text{LD}_{50}$ ) (x) at varying concentrations of compound **7**.

[0019] Figure 2 illustrates graphically a profile of the inhibition in the production of melanin (●) by 4-(2-(benzo[d][1,3]dioxol-5-ylethyl)benzene-1,3-diol (**15**) as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu\text{M}$ ). The  $\text{IC}_{50}$  of compound **15** was determined to be  $3.3 \mu\text{M}$ . Figure 2 also illustrates graphically cell viability ( $\text{LD}_{50}$ ) (x) at varying concentrations of compound **15**. The  $\text{LD}_{50}$  for compound **15** was determined to be  $66 \mu\text{M}$ .

[0020] Figure 3 illustrates graphically a profile of the inhibition in the production of melanin (●) by 4,4'-(ethane-1,2diyl)dibenzene-1,3-diol (**16**) as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu\text{M}$ ). The  $\text{IC}_{50}$  of compound **16** was determined to be  $1.7 \mu\text{M}$ . Figure 3 also illustrates graphically cell viability ( $\text{LD}_{50}$ ) (x) at varying concentrations of compound **16**. The  $\text{LD}_{50}$  for compound **16** was determined to be  $260 \mu\text{M}$ .

[0021] Figure 4 illustrates graphically a profile of the inhibition in the production of melanin (●) by 4-(2,4-dimethoxyphenethyl)benzene-1,3-diol (**17**) as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu\text{M}$ ). The  $\text{IC}_{50}$  of compound **17** was determined to be  $11.9 \mu\text{M}$ . Figure 4 also illustrates graphically cell viability ( $\text{LD}_{50}$ ) (x) at varying concentrations of compound **17**. The  $\text{LD}_{50}$  for compound **17** was determined to be  $157.8 \mu\text{M}$ .

[0022] Figure 5 illustrates graphically a profile of the inhibition in the production of melanin (●) by 4-(3,5-dimethoxyphenethyl)benzene-1,3-diol (**18**) as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu\text{M}$ ). The  $\text{IC}_{50}$  of compound **18** was determined to be  $2.97 \mu\text{M}$ . Figure 5 also illustrates

graphically cell viability ( $LD_{50}$ ) (x) at varying concentrations of compound **18**. The  $LD_{50}$  for compound **18** was determined to be 222.2  $\mu$ M.

[0023] Figure 6 illustrates graphically a profile of the inhibition in the production of melanin (●) by 4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol (**19**) as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu$ M). The  $IC_{50}$  of compound **19** was determined to be 24.7  $\mu$ M. Figure 6 also illustrates graphically cell viability ( $LD_{50}$ ) (x) at varying concentrations of compound **19**. The  $LD_{50}$  for compound **19** was determined to be 130.3  $\mu$ M.

[0024] Figure 7 illustrates graphically a profile of the inhibition in the production of melanin (●) by 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol (**24**) as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu$ M). The  $IC_{50}$  of compound **24** was determined to be 1.6  $\mu$ M. Figure 7 also illustrates graphically cell viability ( $LD_{50}$ ) (x) at varying concentrations of compound **24**. The  $LD_{50}$  for compound **24** was determined to be 345.9  $\mu$ M.

[0025] Figure 8 illustrates graphically a profile of the inhibition in the production of melanin (●) by 4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol (**25**) as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu$ M). The  $IC_{50}$  of compound **25** was determined to be 5  $\mu$ M. Figure 8 also illustrates graphically cell viability ( $LD_{50}$ ) (x) at varying concentrations of compound **25**. The  $LD_{50}$  for compound **25** was determined to be >1000  $\mu$ M.

[0026] Figure 9 illustrates graphically a profile of the inhibition in the production of melanin (●) by compound **30** as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu$ M). The  $IC_{50}$  of compound **30** was determined to be 156  $\mu$ M. Figure 9 also illustrates graphically cell viability ( $LD_{50}$ ) (x) at varying concentrations of compound **25**. The  $LD_{50}$  for compound **30** was determined to be 187.8  $\mu$ M.


[0027] Figure 10 illustrates graphically a profile of the inhibition in the production of melanin (●) by kojic acid as described in Example 7. The data is presented as melanin content (% of untreated control) vs. inhibitor concentration ( $\mu$ M). The  $IC_{50}$  of kojic acid in the murine assay was determined to be 303.5  $\mu$ M. Figure 10 also illustrates graphically cell viability ( $LD_{50}$ ) (x) at varying concentrations of kojic acid. The  $LD_{50}$  for kojic acid was determined to be > 1000  $\mu$ M.

[0028] Figure 11A depicts the reconstructed skin prepared as described in Example 8. The reconstructed skin was grown at the air-liquid interface, making it possible to mimic topical application of skin whitening agents.


[0029] Figure 11B depicts photographically the results of the reconstructed skin studies as described in Example 8. Photographs of skin specimen taken after 15 days of the experiment show significant dose-dependent whitening effects on melanocytes, which appear on photographs as dark dendritic cells.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT


[0030] The present disclosure provides pharmaceutical agents that are potent inhibitors of the tyrosinase enzyme. In particular, provided are substituted 2,4-dihydroxybenzene derivatives that are inhibitors of tyrosinase having the structures depicted in Formulas I-IV, below or pharmaceutically acceptable salts thereof.




Formula I



Formula II



Formula III



Formula IV

These compounds are shown to have excellent tyrosinase inhibitory activity and low cytotoxicity. Illustrated is the inhibitory activity of representative compounds against mushroom tyrosinase, as well as, the inhibition of melanin production by murine B16-F1 melanoma cells. The skin whitening properties of representative compounds were evaluated by means of a reconstructed human skin model as described in Example 8. These compounds or their pharmaceutically acceptable salts thereof are shown to be useful for both cosmetic and medical applications as detailed below.

[0031] As used in this context, the term "derivative" or "analog" refers to a compound having similar chemical structure or function as the compounds of Formula I-IV that retains the core 2,4-dihydroxybenzene ring.

[0032] Various terms are used herein to refer to aspects of the present invention. To aid in the clarification of the description of the components of this invention, the following definitions are provided. Unless defined otherwise all technical and scientific terms used herein have the meaning commonly understood by one of ordinary skill in the art to which this invention belongs.

[0033] It is to be noted that as used herein the term "a" or "an" entity refers to one or more of that entity; for example, a tyrosinase inhibitor refers to one or more tyrosinase inhibitors. As such, the terms "a" or "an", "one or more" and "at least one" are used interchangeably herein.

[0034] As used herein, "about" will be understood by persons of ordinary skill in the art and will vary to some extent on the context in which it is used. If there are uses of the term which are not clear to persons of ordinary skill in the art given the context in which it is used, "about" will mean up to plus or minus 10% of the particular term.

[0035] The term "alkyl" as used herein refers to a straight or branched chain, saturated hydrocarbon having the indicated number of carbon atoms. For example, (C<sub>1</sub>-C<sub>10</sub>) alkyl is meant to include a straight or branched chain hydrocarbon having one to ten carbon atoms. An alkyl group can be unsubstituted or optionally substituted with one or more substituents as described herein.

[0036] The term "alkoxy" as used herein refers to an -O-alkyl group having the indicated number of carbon atoms. For example, a (C<sub>1</sub>-C<sub>10</sub>) alkoxy group includes -OCH<sub>3</sub>, -OCH<sub>2</sub>CH<sub>3</sub>, etc. up to ten carbon atoms.

[0037] As used herein, the term "heteroatom" is meant to include oxygen (O), nitrogen (N) and sulfur (S).

[0038] The term "heteroaromatic" as used herein refers to an aromatic heterocyclic ring of 5 to 14 members and having at least one heteroatom selected from nitrogen, oxygen and sulfur, and containing at least 1 carbon atom. Representative heteroaromatics include pyridyl, furyl, thienyl, pyrrolyl and imidazolyl etc. The heteroaryl group can be attached via any heteroatom or carbon atom, where chemically acceptable. A heteroaryl group can be unsubstituted or optionally substituted with one or more substituents as described herein.

[0039] As used herein, the term "heterocycle" refers to non-aromatic 5 to 14-membered ring systems which are either saturated, unsaturated and which contain from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulfur, and wherein the nitrogen and sulfur heteroatoms can be optionally oxidized, and the nitrogen heteroatom can be optionally quaternized. The heterocycle can be attached via any heteroatom or carbon atom, where chemically acceptable. Representative examples of non-aromatic heterocycles include, but are not limited to tetrahydrofuranyl, tetrahydropyrrolyl, pyranlyl and tetrahydropyranlyl etc. A heterocycle group can be unsubstituted or optionally substituted with one or more substituents as described herein.

**[0040]** The term "aromatic" as used herein refers to any compound which includes or consists of one or more hydrocarbon aromatic rings. The rings may be mono or polycyclic ring systems. Examples of suitable rings include, but are not limited to benzene, biphenyl, terphenyl, naphthalene etc.

**[0041]** The term "hydroxyalkyl" as used herein, refers to an alkyl group having the indicated number of carbon atoms wherein one or more of the hydrogen atoms in the alkyl group is replaced with an -OH group. Representative examples of hydroxyalkyl groups include, but are not limited to, -CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, and branched versions thereof. Typically, the hydroxylalkyl is a compound of the formula -(C<sub>1</sub>-C<sub>10</sub>)alkyl-OH.

**[0042]** "Therapeutic" as used herein, includes prevention, treatment and/or prophylaxis. When used therapeutic refers to humans as well as other animals.

**[0043]** "Pharmaceutically or therapeutically effective dose or amount" refers to a dosage level sufficient to induce a desired biological result. That result may be the alleviation of the signs, symptoms or causes of a disease or any other alteration of a biological system that is desired. The precise dosage will vary according to a variety of factors, including but not limited to the age and size of the subject, the disease and the treatment being effected.

**[0044]** A "host" or "patient" or "subject" is a living mammal, human or animal, for whom therapy is desired. The "host," "patient" or "subject" generally refers to the recipient of the therapy to be practiced according to the method of the invention. It should be noted that the invention described herein may be used for veterinary as well as human applications and that the term "host" should not be construed in a limiting manner. In the case of veterinary applications, the dosage ranges can be determined as described below, taking into account the body weight of the animal.

**[0045]** As used herein, the term "pharmaceutically acceptable" means approved by a regulatory agency of a federal or a state government or listed in the U.S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals and, more particularly, in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered and includes, but is not limited to such sterile liquids as water and oils.


**[0046]** A "pharmaceutically acceptable salt" or "salt" of a tyrosinase inhibitor is a product of the disclosed compound that contains an ionic bond, and is typically produced by reacting the disclosed compound with either an acid or a base, suitable for administering to a

subject. A pharmaceutically acceptable salt can include, but is not limited to, acid addition salts including hydrochlorides, hydrobromides, phosphates, sulphates, hydrogen sulphates, alkylsulphonates, arylsulphonates, arylalkylsulfonates, acetates, benzoates, citrates, maleates, fumarates, succinates, lactates, and tartrates; alkali metal cations such as Li, Na, K, alkali earth metal salts such as Mg or Ca, or organic amine salts.

**[0047]** A "pharmaceutical composition" is a formulation comprising the disclosed compounds in a form suitable for administration to a subject. A pharmaceutical composition of the invention is preferably formulated to be compatible with its intended route of administration. Examples of routes of administration include, but are not limited to, oral and parenteral, *e.g.*, intravenous, intradermal, subcutaneous, inhalation, topical, transdermal, transmucosal, and rectal administration.


**[0048]** The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (*i.e.*, =O), then 2 hydrogens on the atom are replaced. Examples of substituents include, but are not limited to C<sub>1</sub>-C<sub>10</sub> alkyl, hydroxy (-OH); C<sub>1</sub>-C<sub>10</sub> alkoxy groups. Typically an aromatic, heteroaromatic or heterocyclic ring will have from 1-3 substituents.

**[0049]** In one of its aspects the present invention provides a compound having a structure shown in Formula I, or a pharmaceutically acceptable salt thereof:



wherein R is selected from a substituted or unsubstituted aromatic, heteroaromatic or heterocyclic ring. In one embodiment R is selected from the group consisting of a substituted or unsubstituted: phenyl, biphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furanyl, 3-furanyl. In one embodiment R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from the group consisting of a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group or a hydroxyl group. In other embodiments R is selected from the group consisting of 1'-(4'-methoxy)phenyl; 1'-(2,4-dihydroxy)phenyl; 1'-(3-pyridyl); 1'-(biphenyl-4-ol).


**[0050]** In one of its aspects the present invention provides a compound having a structure shown in Formula II or a pharmaceutically acceptable salt thereof.



Formula II

wherein X = H or -OH and R is selected from a substituted or unsubstituted aromatic, heteroaromatic or heterocyclic ring. In one embodiment R is selected from the group consisting of a substituted or unsubstituted: phenyl, biphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furanyl, 3-furanyl. In one embodiment, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from the group consisting of a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group or a hydroxyl group. In other embodiments R is selected from the group consisting of 2'-furyl; 3'-furyl; 3'-pyridyl; 3'-(2-methoxypyridyl); 1'-(2,4,6-trimethoxy phenyl); 1'-(3,4-dioxalane phenyl) and when X = OH, R is 3'-pyridyl.


**[0051]** In one of its aspects the present invention provides a compound having a structure shown in Formula III or a pharmaceutically acceptable salt thereof.



Formula III

wherein R<sub>1</sub> is selected from the group consisting of H, or C<sub>1</sub>-C<sub>10</sub> alkyl; and R<sub>2</sub> is selected from a substituted or unsubstituted: aromatic ring, heteroaromatic ring or heterocyclic ring. In particular embodiments R<sub>1</sub> is selected from H or -CH<sub>3</sub> and R<sub>2</sub> is selected from a substituted or unsubstituted: aromatic ring, heteroaromatic ring or heterocyclic ring. In yet other embodiments, R<sub>2</sub> is substituted with 1 to 3 moieties (R', R'', R''') independently selected from the group consisting of a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group or a hydroxyl group. In other embodiments R<sub>2</sub> is selected from the group consisting of a substituted or unsubstituted: phenyl, biphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furanyl, 3-furanyl or 2-tetrahydrofuryl. In specific embodiments R<sub>2</sub> is selected from the group consisting of R<sub>1</sub> is H and R<sub>2</sub> is 1'-phenyl; 1'-(4-hydroxyphenyl); 1'-(4-methoxyphenyl); 1'-(3,4-dioxalane)phenyl; 1'-(2,4-dihydroxyphenyl); 1'-(2,4-dimethoxyphenyl); 1'-(3,5-dimethoxyphenyl); 1'-(2,4-dimethoxy-3-methylphenyl); 1'-(2,4,6-trimethoxyphenyl); 3-furyl; 3-tetrahydrofuryl; 2'-furyl, 2'-tetrahydrofuryl; 2'-(5'-ethyl)-furyl; 3'-pyridyl and also R<sup>1</sup> = Me, R<sub>2</sub> = 1'-phenyl

**[0052]** In one of its embodiments the present invention provides a compound having a structure shown in Formula II or a pharmaceutically acceptable salt thereof.



wherein  $\equiv$  represents double or single bond,

$R_1$  is selected from the group consisting of H or  $C_1$ - $C_{10}$  alkyl;

$R_4$  is selected from the group consisting of = H, OH or O; and

$R_5$  is selected from the group consisting of a substituted or unsubstituted: aromatic ring, heteroaromatic ring, heterocyclic ring or a hydroxylalkyl moiety. In specific embodiments  $R_1$  is H or  $CH_3$  and  $R_5$  is selected from a substituted or unsubstituted: aromatic ring, heteroaromatic ring or a heterocyclic ring or a  $C_1$ - $C_{10}$  hydroxylalkyl ( $-C_1$ - $C_{10}$ )-OH). In yet other embodiments,  $R_5$  is substituted with 1 to 3 moieties ( $R'$ ,  $R''$ ,  $R'''$ ) independently selected from the group consisting of a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group or a hydroxyl group. In other embodiments  $R_5$  is selected from the group consisting of a substituted or unsubstituted: phenyl, biphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furanyl, 3-furanyl or 2-tetrahydrofuranyl. In specific embodiments  $R_1 = R_4 = H$  and  $R_5$  is selected from 1'-(3',5'-dimethoxy)phenyl; 1'-(3',5'-dihydroxy)phenyl; 1-(2',4',6-trimethoxyphenyl);  $R_1 = Me$ ,  $R_4 = OH$ ,  $R_5$  is 1'-(2',4'-dihydroxy)phenyl;  $R_1 = Me$ ,  $R_4 = O$ ,  $R_5$  is 1'-(2',4'-dihydroxy)phenyl; and  $R_1 = R_4 = H$ ,  $R_5$  - $CH_2CH_2OH$ .

with the following provisos:

proviso 1: when  $R_1 = R_4 = H$ ;  $R_5$  cannot be 1'-(2,4-dimethoxy-3-methyl)phenyl.

proviso 2: when  $R_1 = R_4 = H$ ;  $R_5$  cannot be 1'-(2,4-dihydroxy)phenyl; 1'-(2,4-dialkoxy)phenyl; 1'-(2,4-dialkenoxy)phenyl or 1'-(2,4-dibenzyloxy)phenyl.

proviso 3:  $R_5$  cannot be 1'-(4-hydroxy)phenyl.

proviso 4: when  $R_1 = R_4 = H$ ;  $R_5$  cannot be 1'-(2,4-di(CO) $R^1$ ), wherein  $R^1$  is a ( $C_1$ - $C_{20}$ )-alkyl group.

proviso 5: when  $R_1 = R_4 = H$  and  $R_5$  cannot be a 1'-(2,4-disubstituted)phenyl; wherein said substituents are selected from  $-OR^2$  and  $-OR^3$ , wherein  $R^2$  and  $R^3$  are independently selected from the group consisting of H or a  $(C_1-C_{20})$ -alkyl.

proviso 6: when  $R_1 = H$  and  $R_4 = OH$ ;  $R_5$  cannot be 1'-(3,4-dihydroxy)phenyl; 1'-(3,4-dimethoxy)phenyl; 1'-(2-hydroxy)phenyl; 1'-(2-methoxy)phenyl; 1'-(4-methoxy)phenyl; phenyl; 1'-(3-methoxy,4-hydroxy)phenyl; .

**[0053]** Table 1 lists representative novel 2,4-dihydroxybenzene analogs of Formula I which are useful as tyrosinase inhibitors. The synthetic methods that can be used to prepare each compound, identified in Table 1 are described in detail in Example 1. Supporting  $^1H$ - and  $^{13}C$ -NMR data is provided for each compound synthesized. In general, the compounds of Formula I can be synthesized from readily available materials using standard organic synthesis techniques. Further preparation routes may be found in the literature and relevant art.  $IC_{50}$  values for these compounds as determined by the tyrosinase assay as described in Example 6 are also set forth in Table 1. For select compounds of Formula I, Table 5 provides  $IC_{50}$  values as determined by the murine melanoma cell-based assays described in Example 7 and cell viability ( $LD_{50}$ ) as determined by the methods described in Example 8.

**[0054]** Table 2 lists representative novel 2,4-dihydroxybenzene derivatives of Formula II which are useful as tyrosinase inhibitors. The synthetic methods that can be used to prepare each compound identified in Table 2 are described in detail in Example 2. Supporting  $^1H$ - and  $^{13}C$ -NMR data is provided for each compound synthesized. In general, the compounds of Formula II can be synthesized from readily available materials using standard organic synthesis techniques. Further preparation routes may be found in the literature and relevant art.  $IC_{50}$  values for these compounds as determined by the tyrosinase assay as described in Example 6 are also set forth in Table 2. For select compounds of Formula II, Table 5 provides  $IC_{50}$  values as determined by the murine melanoma cell-based assay described in Example 7 and cell viability ( $LD_{50}$ ) as determined by the methods described in Example 8.

**[0055]** Table 3 lists representative novel 2,4-dihydroxybenzene analogs of Formula III which are useful as tyrosinase inhibitors. The synthetic methods that can be used to prepare each compound identified in Table 3 are described in detail in Example 3. Supporting  $^1H$ - and  $^{13}C$ -NMR data is provided for each compound synthesized. In general, the compounds of Formula III can be synthesized from readily available materials using

standard organic synthesis techniques. Further preparation routes may be found in the literature and relevant art.  $IC_{50}$  values for these compounds as determined by the tyrosinase assay as described in Example 6 are also set forth in Table 3. For select compounds of Formula III, Table 5 provides  $IC_{50}$  values as determined by the murine melanoma cell-based assay described in Example 7 and cell viability ( $LD_{50}$ ) as determined by the methods described in Example 8.

**[0056]** Table 4 lists representative novel 2,4-dihydroxybenzene analogs of Formula IV which are useful as tyrosinase inhibitors. The synthetic methods that can be used to prepare each compound identified in Table 4 are described in detail in Example 4 and Example 5 (compound 34). Supporting  $^1H$ - and  $^{13}C$ -NMR data is provided for each compound synthesized. In general, the compounds of Formula IV can be synthesized from readily available materials using standard organic synthesis techniques. Further preparation routes may be found in the literature and relevant art.  $IC_{50}$  values for these compounds as determined by the tyrosinase assay as described in Example 6 are also set forth in Table 4. For select compounds of Formula IV, Table 5 provides  $IC_{50}$  values as determined by the murine melanoma cell-based assay described in Example 7 and cell viability ( $LD_{50}$ ) as determined by the methods described in Example 8.

**[0057]** In summary, the compounds set forth shown in Tables 1-4 exhibited excellent inhibition of the mushroom tyrosinase enzyme, as well as, melanin production by B16-F1 cells (Table 5, Figures 1-9). Cytotoxicity was observed only at high concentrations. All compounds exemplified in Table 5 (i.e., compounds 7, 15, 16, 17, 18, 19, 24, 25, 30) were much more potent than kojic acid, a commonly used skin whitener as illustrated in Figure 10. The  $IC_{50}$  of kojic acid (tyrosinase assay) is 20  $\mu M$ . The  $IC_{50}$  of kojic acid (murine assay) is 303.5  $\mu M$ . Additionally, two of the compounds tested (compounds 16 and 25) exhibited strong skin-whitening effects in a reconstructed skin model without any detectable cytotoxicity as described in Example 8.

**[0058]** The present invention encompasses pharmaceutical compositions comprising at least one tyrosinase inhibitor described herein. The compositions of the present invention can be formulated as pharmaceutical compositions, which include other components such as a pharmaceutically and/or cosmetically acceptable excipient, an adjuvant, and/or a carrier. For example, compositions of the present invention can be formulated in an excipient that the host to be treated can tolerate. An excipient is an inert substance used as a diluent or vehicle for a therapeutic agent. Examples of such excipients include, but are not limited to water, buffers, saline, Ringer's solution, dextrose solution, mannitol, Hank's solution, preservatives

and other aqueous physiologically balanced salt solutions. Nonaqueous vehicles, such as fixed oils, sesame oil, ethyl oleate, or triglycerides may also be used. Other useful formulations include suspensions containing viscosity-enhancing agents, such as sodium carboxymethylcellulose, sorbitol or dextran. Excipients can also contain minor amounts of additives, such as substances that enhance isotonicity and chemical stability. Examples of buffers include phosphate buffer, bicarbonate buffer, tris buffer, histidine, citrate, and glycine, or mixtures thereof, while examples of preservatives include, but are not limited to EDTA, disodium EDTA, BHA, BHT, vitamin C, vitamin E, sodium bisulfite,  $\text{SnCl}_2$ , thimerosal, m- or o-cresol, formalin and benzyl alcohol. Standard formulations can be either liquid or solids, which can be taken up in a suitable liquid as a suspension or solution for administration. Thus, in a non-liquid formulation, the excipient can comprise dextrose, human serum albumin, preservatives, etc., to which sterile water or saline can be added prior to administration.

**[0059]** In one embodiment of the present invention, the composition can also include an adjuvant or a carrier. Adjuvants are typically substances that generally enhance the biological response of a host to a specific bioactive agent. Suitable adjuvants include, but are not limited to, Freund's adjuvant, other bacterial cell wall components, aluminum, magnesium, copper, zinc, iron, calcium, and other metal ion based salts, silica, polynucleotides, toxoids, serum proteins, viral coat proteins, other bacterial-derived preparations, gamma interferon; block copolymer adjuvants; such as Hunter's Titermax adjuvant (Vaxcel.TM., Inc. Norcross, Ga.), Ribi adjuvants (available from Ribi ImmunoChem Research, Inc., Hamilton, Mont.); and saponins and their derivatives, such as Quil A (available from Superfos Biosector A/S, Denmark). Carriers are typically compounds that increase the half-life of a therapeutic composition in the treated host. Suitable carriers include, but are not limited to, polymeric controlled release formulations, biodegradable implants, liposomes, bacteria, viruses, oils, esters, and glycols. Suitable carriers are described in "Remington: The Science and Practice, Twentieth Edition," published by Lippincott Williams & Wilkins, which is incorporated herein by reference.

**[0060]** In one embodiment, the composition is prepared as a controlled release formulation, which slowly releases the composition of the present invention into the host. As used herein, a controlled release formulation comprises a composition of the present invention in a controlled release vehicle. Suitable controlled release vehicles will be known to those skilled in the art. Preferred controlled release formulations are biodegradable (i.e., bioerodible).

**[0061]** The compositions of this invention can be administered by any method known to one of ordinary skill in the art. The modes of administration include, but are not limited to, enteral (oral) administration, parenteral (intravenous, subcutaneous, and intramuscular) administration and topical application. The method of treatment according to this invention comprises administering internally or topically to a patient in need thereof a therapeutically effective amount of at least one of the 2,4-dihydroxybenzene analogs of the instant invention.

**[0062]** In one embodiment, the therapeutic agents of the instant invention are administered topically by any suitable means, known to those of skill in the art for topically administering therapeutic compositions including, but not limited to as an ointment, gel, lotion, or cream base, or as a toothpaste, mouth-wash, or coated on dental flossing materials or as an emulsion, as a patch, dressing or mask, a nonsticking gauze, a bandage, a swab or a cloth wipe.

**[0063]** A therapeutic composition can be administered in a variety of unit dosage forms depending upon the method of administration. For particular modes of delivery, a therapeutic composition of the present invention can be formulated in an excipient of the present invention. A therapeutic reagent of the present invention can be administered to any host, preferably to mammals, and more preferably to humans. The particular mode of administration will depend on the condition to be treated.

**[0064]** In one embodiment, a suitable ointment is comprised of the desired concentration of at least one 2,4-dihydroxybenzene analog of the instant invention, that is an efficacious, nontoxic quantity generally selected from the range of 0.001% to 100% based on the total weight of the topical formulation, from 65 to 100% (preferably 75 to 96%) of white soft paraffin, from 0 to 15% of liquid paraffin, and from 0 to 7% (preferably 3 to 7%) of lanolin or a derivative or synthetic equivalent thereof. In another embodiment the ointment may comprise a polyethylene-liquid paraffin matrix.

**[0065]** In one embodiment, a suitable cream is comprised of an emulsifying system together with the desired concentration of at least one 2,4-dihydroxybenzene analog of the instant invention as provided above. The emulsifying system is preferably comprised of from 2 to 10% of polyoxyethylene alcohols (e.g. the mixture available under the trademark Cetomacrogol<sup>TM</sup>1000), from 10 to 25% of stearyl alcohol, from 20 to 60% of liquid paraffin, and from 10 to 65% of water; together with one or more preservatives, for example from 0.1 to 1% of N,N'-methylenebis[N'-[3-(hydroxymethyl)-2,5-dioxo-4-imidazolidinyl]urea] (available under the name Imidurea USNF), from 0.1 to 1% of alkyl 4-hydroxybenzoates

(for example the mixture available from Nipa Laboratories under the trade mark Nipastat), from 0.01 to 0.1% of sodium butyl 4-hydroxybenzoate (available from Nipa Laboratories under the trade mark Nipabutyl sodium), and from 0.1 to 2% of phenoxyethanol.

**[0066]** In one embodiment, a suitable gel is comprised of a semi-solid system in which a liquid phase is constrained within a three dimensional polymeric matrix with a high degree of cross-linking. The liquid phase may be comprised of water, together with the desired amount of at least one 2,4-dihydroxybenzene analog of the instant invention, from 0 to 20% of water-miscible additives, for example glycerol, polyethylene glycol, or propylene glycol, and from 0.1 to 10%, preferably from 0.5 to 2%, of a thickening agent, which may be a natural product, selected from the group including, but not limited to tragacanth, pectin, carrageen, agar and alginic acid, or a synthetic or semi-synthetic compound, selected from the group including, but not limited to methylcellulose and carboxypolymethylene (carbopol); together with one or more preservatives, selected from the group including, but not limited to for example from 0.1 to 2% of methyl 4-hydroxybenzoate (methyl paraben) or phenoxyethanol-differential. Another suitable base, is comprised of the desired amount of at least one 2,4-dihydroxybenzene analog of the instant invention, together with from 70 to 90% of polyethylene glycol (for example, polyethylene glycol ointment containing 40% of polyethylene glycol 3350 and 60% of polyethylene glycol 400, prepared in accordance with the U.S. National Formulary (USNF)), from 5 to 20% of water, from 0.02 to 0.25% of an anti-oxidant (for example butylated hydroxytoluene), and from 0.005 to 0.1% of a chelating agent (for example ethylenediamine tetraacetic acid (EDTA)).

**[0067]** The term soft paraffin as used above encompasses the cream or ointment bases white soft paraffin and yellow soft paraffin. The term lanolin encompasses native wool fat and purified wool fat. Derivatives of lanolin include in particular lanolins which have been chemically modified in order to alter their physical or chemical properties and synthetic equivalents of lanolin include in particular synthetic or semisynthetic compounds and mixtures which are known and used in the pharmaceutical and cosmetic arts as alternatives to lanolin and may, for example, be referred to as lanolin substitutes.

**[0068]** One suitable synthetic equivalent of lanolin that may be used is the material available under the trademark Softisan<sup>TM</sup> known as Softisan 649. Softisan 649, available from Dynamit Nobel Aktiengesellschaft, is a glycerine ester of natural vegetable fatty acids, of isostearic acid and of adipic acid; its properties are discussed by H. Hermsdorf in Fette, Seifen, Anstrichmittel, Issue No. 84, No.3 (1982), pp. 3-6.

[0069] The other substances mentioned hereinabove as constituents of suitable ointment or cream bases and their properties are discussed in standard reference works, for example pharmacopoeia. Cetomacrogol 1000 has the formula  $\text{CH}_3(\text{CH}_2)_m(\text{OCH}_2\text{CH}_2)_n\text{OH}$ , wherein  $m$  may be 15 or 17 and  $n$  may be 20 to 24. Butylated hydroxytoluene is 2,6-di-tert-butyl-p-cresol. Nipastat is a mixture of methyl, ethyl, propyl and butyl 4-hydroxybenzoates.

[0070] The compositions of the invention may be produced by conventional pharmaceutical techniques. Thus the aforementioned compositions, for example, may conveniently be prepared by mixing together at an elevated temperature, preferably 60-70°C, the soft paraffin, liquid paraffin if present, and lanolin or derivative or synthetic equivalent thereof. The mixture may then be cooled to room temperature, and, after addition of the hydrated crystalline calcium salt of mupirocin, together with the corticosteroid and any other ingredients, stirred to ensure adequate dispersion.

[0071] Regardless of the manner of administration, the specific dose is calculated according to the approximate body weight of the host. Further refinement of the calculations necessary to determine the appropriate dosage for treatment involving each of the above mentioned formulations is routinely made by those of ordinary skill in the art and is within the scope of tasks routinely performed by them without undue experimentation, especially in light of the dosage information and assays disclosed herein. These dosages may be ascertained through use of the established assays for determining dosages utilized in conjunction with appropriate dose-response data.

[0072] The tyrosinase inhibitors of the invention can readily be synthesized using known synthetic methodologies or via a modification of known synthetic methodologies. As would be readily recognized by a skilled artisan, the methodologies described below allow the synthesis of 2,4-dihydroxybenzene analogs having a variety of substituents. Included in this invention is a method of synthesizing the 2,4-dihydroxybenzen analogs described herein. Exemplary synthetic methods are described in Examples 1-5 below.

[0073] The invention encompasses methods of preventing or treating (*e.g.*, alleviating one or more symptoms of) medical conditions through use of one or more of the disclosed compounds. The methods of preventing or treating comprise administering a therapeutically effective amount of at least one tyrosinase inhibitor of the instant invention to a patient in need thereof. The compositions of the invention can also be used for prophylactic therapy.

[0074] The present invention provides a method of treating a subject afflicted with a disorder ameliorated by the inhibition of the activity tyrosinase or the production of excess

amounts of melanin. Such a method comprises administering to a subject a therapeutically effective amount of a tyrosinase inhibitor as described herein.

**[0075]** As used herein, "treating" describes the management and care of a patient for the purpose of combating a disease, condition, or disorder and includes the administration of a compound of the present invention to prevent the onset of the symptoms or complications, alleviating the symptoms or complications, or eliminating the disease, condition or disorder. More specifically, "treating" includes reversing, alleviating, minimizing, suppressing or halting at least one deleterious symptom or effect of a disease (disorder) state, disease progression, or other abnormal condition. Treatment is continued as long as symptoms and/or pathology ameliorate.

**[0076]** More specifically the present invention also provides a method for inhibiting the synthesis of melanin in a subject in need thereof. Such a method comprises administering an effective amount of a composition comprising at least one of the 2,4-dihydroxybenzene analogs of the instant invention.

**[0077]** The present invention further provides methods for the prevention and treatment of diseases and conditions related to the overproduction or uneven distribution of melanin, said method comprising administering internally or topically to a subject in need thereof a therapeutically effective amount of at least one at least one 2,4-dihydroxybenzene compound of the instant invention. Diseases and conditions related to the overproduction or uneven distribution of melanin include, but not limited to hyperpigmentation spots caused by skin aging, melasma, liver diseases, thermal burns and topical wounds, skin pigmentation due to inflammatory conditions caused by fungal, microbial and viral infections, vitiligo, carcinoma, melanoma, as well as other mammalian skin conditions.

**[0078]** The method can also be used for preventing and treating skin darkening and damage resulting from exposure to the sun, ultraviolet (UV) radiation, chemicals, heat, wind and dry environments. Finally, the method can be used for preventing and treating wrinkles, saggy skin, lines and dark circles around the eyes, soothing sensitive skin and preventing and treating dermatitis and other allergy related conditions of the skin.


**[0079]** In addition to their use for the prevention and treatment of the above described diseases and conditions of the skin, the therapeutic compositions described herein provide an efficacious composition that yields the benefit of smooth and youthful skin appearance with improved skin color, enhanced elasticity, reduced and delayed aging, enhanced youthful appearance and texture, and increased flexibility, firmness, smoothness and suppleness.

[0080] In yet another embodiment, the 2,4-dihydroxybenzene derivatives of the instant invention can be used in the food industry to inhibit the browning and color changes in fruits, vegetables, juices and other food products.

[0081] The following examples are given to illustrate the present invention. It should be understood, however, that the invention is not to be limited to the specific conditions or details described in these examples.

## EXAMPLES


### Example 1. General and specific methods for the synthesis of compounds of Formula I



**Formula I**

[0082] Compounds of Formula I were prepared generally as set forth in Scheme 1, using a substituted bromobenzene as R for purposes of illustration. Briefly, the biphenyl compounds of Formula I were prepared by Kumada coupling reaction (Tamao, K.; Sumitani, K.; Kumada, M, *Journal of the American Chemical Society*. **1972**, *94*. 4374-4376) using Ni(dppp)Cl<sub>2</sub> catalyst from the corresponding Grignard reagents with appropriate aryl bromide followed by reductive debenzylization reaction of benzyl group.

Scheme 1

**Representative procedure for Scheme 1: Synthesis of 4'-Methoxybiphenyl-2,4-diol (1)**

**[0083]** 4-Bromoanisole (2.80 g, 15 mmol) was treated with magnesium turnings (480 mg, 20 mmol) and iodine (a pinch) in THF to make the corresponding Grignard reagent.

The Grignard reagent was then added to a stirred solution of benzyl protected 4-bromoresorcinol (3.7 g, 10 mmol) and Ni(dppp)Cl<sub>2</sub> (2.7 g, 5 mmol) in THF at 0°C. The reaction mixture was allowed to stir at room temperature for 30 minutes after which it was heated to reflux for 4h. The reaction mixture was then cooled down and quenched with ammonium chloride, extracted with ethyl acetate, washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in rotary evaporator and purified by column chromatography to yield (1.98 g, 50%) of the coupled product as an off white solid.

**[0084]** The resulting coupled product (1.5 g, 3.7 mmol) was then treated with a catalytic amount of 10% Pd on activated carbon (150 mg) and an excess amount of ammonium formate (1.9 g, 29.6 mmol) in 15 mL THF/MeOH (1:2) solution at reflux for 3 h to remove the benzyl protection group. After filtration through Celite\* to remove the catalyst, the crude product was purified by column chromatography to yield (720 mg, 90%) of compound 1 as an off-white solid. <sup>1</sup>H-NMR (MeOD, 500MHz): δ 7.152 (d, 2H, *J* = 8.5Hz), 6.718-6.762 (m, 3H), 6.443(d, 1H, *J* = 2.5Hz), 6.241(dd, 1H, *J* = 2.5 & 8.0Hz), 3.663(s, 3H).

**[0085]** Representative compounds **2-4** of Formula I (see Table 1) were synthesized according to the same protocol using appropriate starting materials. Overall yields varied from 40-50%.

**Biphenyl-2,2',4,4'-tetraol (2)**

**[0086]** <sup>1</sup>H-NMR (MeOD, 300MHz): δ 7.007-6.953(m, 2H), 6.378-6.327(m, 6H).

<sup>13</sup>C-NMR (MeOD, 75MHz): δ 157.971(6C), 130.246(2CH), 107.134(2CH), 102.589(2CH).

**4-(Pyridin-3yl)benzene-1,3-diol (3)**

\*Trade-mark

[0087]  $^1\text{H-NMR}$ : ( $\text{CDCl}_3$ , 300MHz):  $\delta$  8.829(s, 1H), 8.539(d, 1H,  $J=3.6\text{Hz}$ ), 7.885(d, 1H,  $J=7.5\text{Hz}$ ), 7.605-7.261(m, 1H), 6.753(s, 1H), 6.711(d, 1H,  $J=8.4\text{Hz}$ ).  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ , 75MHz):  $\delta$  157.036(C), 150.534(CH), 137.420(CH), 136.926(CH), 134.400(C), 131.618(CH), 129.237(C), 127.931(CH), 120.872(C), 106.821(CH), 101.769(CH).


#### 4-(4'-Hydroxybiphenyl)benzene-1,3-diol (4)

[0088]  $^1\text{H-NMR}$  (MeOD, 300MHz):  $\delta$  7.511-7.183(m, 9H), 6.88-6.83(m, 2H).  $^{13}\text{C-NMR}$  (MeOD, 75MHz):  $\delta$  156.949 (3 C), 141.204(2 C), 132.709(2 C), 128.557(2 CH), 127.893(3 CH), 126.256(3 CH), 115.472(3 CH).

Table 1. Compounds Representative of Formula I

| Compound                                   | Structure | Mushroom Tyrosinase Assay            |
|--------------------------------------------|-----------|--------------------------------------|
| 4'-Methoxybiphenyl-2,4-diol (1)            |           | $\text{IC}_{50}: = 12.8 \mu\text{M}$ |
| Biphenyl-2,2',4,4'-tetraol (2)             |           | $\text{IC}_{50}: = 60.0 \mu\text{M}$ |
| 4-(Pyridin-3-yl)benzene-1,3-diol (3)       |           | $\text{IC}_{50}: = 0.5 \mu\text{M}$  |
| 4-(4'-Hydroxybiphenyl)benzene-1,3-diol (4) |           | $\text{IC}_{50}: = 70.0 \mu\text{M}$ |

#### Example 2. General and specific methods for the synthesis of compounds of Formula II




Formula II

[0089] Compounds of Formula II were prepared as illustrated in Scheme 2. Briefly, with reference to Scheme 2, the diphenylmethane compounds illustrative of Formula II were

prepared by Grignard addition reaction with an aldehyde (RCHO, wherein R=aromatic or heteroaromatic ring), followed by dehydroxylation of resulting benzylic hydroxyl group and debenzoylation of benzyloxy protecting groups in situ.

**Scheme 2**



**Representative procedure for Scheme 2: Synthesis of 4-(Furan-2-ylmethyl)benzene-1,3-diol (5)**

**[0090]** Benzyl protected 4-bromoresorcinol (2.80 g, 15 mmol) was treated with magnesium turnings (480 mg, 20 mmol) and iodine (a pinch) in THF to provide the corresponding Grignard reagent, which was then added to a stirred solution of 2-furaldehyde (3.97 g, 10 mmol) in THF at 0°C. The reaction mixture was allowed to warm up to room temperature and stirred at room temperature for 3h. The mixture was then quenched with ammonium chloride, extracted with ethyl acetate, washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, concentrated in a rotary evaporator and purified by column chromatography to provide (3.0 g, 80%) of the coupled product.

**[0091]** The coupled product (2.5 g, 6.5 mmol) was then treated with a catalytic amount of 10% Pd on activated carbon (250 mg) and an excess amount of ammonium formate (3.33 g, 52 mmol) in THF/MeOH (18 mL) at reflux to remove the benzyl protecting groups. Partial dehydroxylation of benzyl hydroxy group also took place during the debenzoylation reaction. Upon completion, the reaction mixture was filtered through Celite to remove the catalyst and concentrated to provide the crude product, which was then purified by column chromatography to yield (740 mg, 60%) of compound 5 as an off-white solid, as well as, the hydroxylated compound (X=OH) (270 mg, 20%). <sup>1</sup>H-NMR (MeOD, 500MHz): δ 7.306 (dd, 1H, *J*=1.0 & 2.0Hz), 6.8565(d, 1H, *J*= 8.5Hz), 6.8085(d, 1H, *J*= 8.5Hz), 6.3155(d, 1H, *J*= 2.5Hz), 6.2965(d, 1H, *J*= 2.5Hz), 6.255-6.223(m, 2H). <sup>13</sup>C-NMR(MeOD, 125MHz): δ 156.604(C), 156.504(C), 156.064(C), 155.556(C), 155.290(C), 140.575(CH), 131.443(CH), 130.325(CH), 116.098(C), 116.973(C), 109.761(CH), 106.303(CH), 106.111(CH), 105.156(CH), 102.513(CH), 102.061(CH), 79.946(CH), 67.489(CH<sub>2</sub>), 35.257(CH), 29.875(CH<sub>2</sub>), 27.102(CH<sub>2</sub>), 25.218(CH<sub>2</sub>).

**[0092]** Representative compounds **6-11** of Formula II (see Table 2) were synthesized according to the same general reaction scheme using the appropriate starting materials.

Overall yields varied from 50-60%.

**4-(Furan-3-ylmethyl)benzene-1,3-diol (6)**

**[0093]** <sup>1</sup>H-NMR (MeOD, 500 MHz): δ 8.39(d, 1H, *J* = 2Hz), 8.28(dd, 1H, *J* = 5 & 1.5 Hz), 6.63-6.69(m, 1H), 7.25-7.28(m, 1H), 6.87(d, 1H, *J* = 8.5 Hz), 6.31(d, 1H, *J* = 2.5 Hz), 6.24 (dd, 1H, *J* = 8 & 2.5 Hz), 3.25(s, 2H). <sup>13</sup>C-NMR (MeOD, 125 MHz): δ 156.897(C), 155.797 (C), 148.727(CH), 145.584(CH), 138.672(C), 137.021(CH), 130.536(CH), 123.476(CH), 117.446(C), 106.221(CH), 102.283(CH), 32.196(CH<sub>2</sub>).

**4-(Pyridin-ylmethyl)benzene-1,3-diol (7)**

**[0094]** <sup>1</sup>H-NMR (MeOD, 500 MHz): δ 8.39(d, 1H, *J* = 2Hz), 8.28(dd, 1H, *J* = 5 & 1.5 Hz), 6.63-6.69(m, 1H), 7.25-7.28(m, 1H), 6.87(d, 1H, *J* = 8.5 Hz), 6.31(d, 1H, *J* = 2.5 Hz), 6.24 (dd, 1H, *J* = 8 & 2.5 Hz), 3.25(s, 2H). <sup>13</sup>C-NMR (MeOD, 125 MHz): δ 156.897(C), 155.797 (C), 148.727(CH), 145.584(CH), 138.672(C), 137.021(CH), 130.536(CH), 123.476(CH), 117.446(C), 106.221(CH), 102.283(CH), 32.196(CH<sub>2</sub>).

**[0095]** Hydroxylated product (20 %) was also obtained and was successfully separated by column chromatography (See compound **11** below).

**4-((2-Methoxypyridin-3-yl)methyl)benzene-1,3-diol (8)**

**[0096]** <sup>1</sup>H-NMR (MeOD, 500 MHz): δ 7.9015 (dd, 1H, *J* = 1.5 & 5.0Hz), 7.277(dt, 1H, *J* = 1.0 & 7.0Hz), 6.778-6.823 (m, 2H), 6.313(dd, 1H, *J* = 2.5 & 6.5Hz), 6.239(dt, 1H, *J* = 2.0 & 8.5Hz), 3.935(s, 3H), 3.749(s, 2H). <sup>13</sup>C-NMR (MeOD, 125 MHz): δ 161.927(C), 157.766(C), 156.578(C), 155.941(C), 143.115(CH), 137.880(CH), 130.828(CH), 124.794(C), 116.498(CH), 106.133(CH), 102.098(CH), 52.482(CH<sub>3</sub>), 28.505(CH<sub>2</sub>).

**4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9)**

**[0097]** <sup>1</sup>H-NMR (MeOD, 500Hz): δ 6.587(d, 1H, *J* = 8.5Hz), 6.242 (s, 1H), 6.239(s, 2H), 6.106(dd, 1H, *J* = 2.5 & 8.0Hz), 3.798(s, 3H), 3.775(s, 6H), 3.703(s, 2H). <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 125Hz): δ 160.150(C), 158.100(2 C), 155.986(C), 155.509(C), 132.723(CH), 119.535(C), 109.882(C), 107.232(CH), 103.475(CH), 91.531(2CH), 56.258(2 CH<sub>3</sub>), 55.802(CH<sub>3</sub>), 23.157(CH<sub>2</sub>).







**4-(Benzo[d][1,3]dioxol-5-ylmethyl)benzene-1,3-diol (10)**


[0098]  $^1\text{H-NMR}$  (MeOD, 500 MHz):  $\delta$  6.775(d, 1H,  $J = 8.5\text{Hz}$ ), 6.62-6.68 (m, 3H), 6.295(d, 1H,  $J = 2.0\text{Hz}$ ), 6.202-6.223(m, 1H), 5.842(s, 2H), 3.733(s, 2H).  $^{13}\text{C-NMR}$  (MeOD, 125 MHz):  $\delta$  155.840(C), 155.124(C), 148.362(C), 146.548(C), 134.234(C), 131.868(CH), 121.597(CH), 119.754(C), 109.431(CH), 108.687(CH), 108.154(CH), 103.761(CH), 101.299(CH<sub>2</sub>), 35.949(CH<sub>2</sub>).

**4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11)**


[0099]  $^1\text{H-NMR}$  (MeOD, 300MHz) :  $\delta$  8.528(d, 1H,  $J = 2.1\text{Hz}$ ), 8.355(dd, 1H,  $J = 3.6$  & 4.8Hz), 7.820(td,  $J = 1.5$  & 7.5Hz), 7.353(t, 1H,  $J = 5.4\text{Hz}$ ), 7.086(d, 1H,  $J = 7.8\text{Hz}$ ), 6.315-6.267(m, 2H), 6.067(s, 1H); Minor partial: 8.497(d, 1H,  $J = 2.1\text{Hz}$ ), 7.018(d, 1H,  $J = 9.3\text{Hz}$ ), 5.661(s, 1H).  $^{13}\text{C-NMR}$ (MeOD, 125MHz):  $\delta$  157.659(C), 155.283(C), 147.120(CH), 146.694(CH), 141.308(C), 135.144(CH), 127.604(CH), 123.495(CH), 120.885(C), 106.466(CH), 102.231(CH), 68.285(CH).

Table 2. Compounds Representative of Formula II

| Compound                                                         | Structure                                                                             | Mushroom Tyrosinase Assay               |
|------------------------------------------------------------------|---------------------------------------------------------------------------------------|-----------------------------------------|
| 4-(Furan-2-ylmethyl)benzene-1,3-diol ( <b>5</b> )                |  | IC <sub>50</sub> : = 1.8 $\mu\text{M}$  |
| 4-(Furan-3-ylmethyl)benzene-1,3-diol ( <b>6</b> )                |  | IC <sub>50</sub> : = 3.0 $\mu\text{M}$  |
| 4-(Pyridin-ylmethyl)benzene-1,3-diol ( <b>7</b> )                |  | IC <sub>50</sub> : = 2.0 $\mu\text{M}$  |
| 4-((2-Methoxypyridin-3-yl)methyl)benzene-1,3-diol ( <b>8</b> )   |  | IC <sub>50</sub> : = 50.0 $\mu\text{M}$ |
| 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol ( <b>9</b> )          |  | IC <sub>50</sub> : = 30.0 $\mu\text{M}$ |
| 4-(Benzo[d][1,3]dioxol-5-ylmethyl)benzene-1,3-diol ( <b>10</b> ) |  | IC <sub>50</sub> : = 0.85 $\mu\text{M}$ |

|                                                                 |                                                                                     |                             |
|-----------------------------------------------------------------|-------------------------------------------------------------------------------------|-----------------------------|
| 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol ( <b>11</b> ) |  | IC <sub>50</sub> : = 4.9 μM |
|-----------------------------------------------------------------|-------------------------------------------------------------------------------------|-----------------------------|


**Example 3. General and specific methods for the synthesis of compounds of Formula III**



Formula III

**[00100]** Compounds of Formula III were prepared as illustrated in Scheme 3, below.

Briefly, with reference to Scheme 3, the 1,2-diphenylethene compounds of Formula III were prepared by Wittig reaction between the corresponding Wittig salt and aldehyde followed by hydrogenation with Pd on activated carbon and ammonium formate (AMF).



**Representative procedure for Scheme 3: 4-Phenethylbenzene-1,3-diol (**12**)**

**[00101]** To a solution of the phosphonium salt of benzyl bromide (5.2 g, 12 mmol) in THF (25 mL), nBu-Li in hexane (6.9 mL, 11 mmol) was added at 0°C. The reaction mixture was allowed to stir at room temperature for 1 h, during which time the mixture developed a red color indicating the formation of ylide. A solution of 2,4-bis(benzoyloxy)benzaldehyde (3.2 g, 10 mmol) in THF (10 mL) was added to the ylide solution at -80°C and allowed to stir overnight without removing the low temperature bath. During that period the red color was disappeared and a whitish suspension was formed which was diluted with hexane and filtered. The filtrate was concentrated and the residue was passed through a short bed of silica gel eluting with ethyl acetate-hexane (5:95) to yield the coupled product (3.6 g, 92%) in a ratio of (*E/Z* : 20/80) as a pale yellow oil.

**[00102]** The coupled product (3 g, 7.7 mmol) was then dissolved in 24 mL THF/MeOH (1:2) followed by treatment with 10% Pd on activated carbon (300 mg) and ammonium formate (3.92 g, 62 mmol) at reflux for 4 h. After cooling, the reaction mixture was filtered through celite to remove the catalyst. The crude product was purified by column chromatography eluting with hexane and ethyl acetate (75:25) to yield compound **12** (1.5 g, 95%) as off-white solid. <sup>1</sup>H-NMR (MeOD, 500 MHz): δ 7.19-7.09 (m, 5H), 7.77(d, 1H, *J*=8Hz), 6.35 (t, 1H, *J*=2.5Hz), 6.22 (s, 1H), 2.80 (d, 2H, *J* = 6Hz), 2.78 (d, 2H, *J* = 4Hz). <sup>13</sup>C-NMR (MeOD, 125 MHz): δ 155.911(C), 155.604(C), 142.535(C), 130.302(CH), 128.177(CH), 128.170(CH), 127.815(CH), 127.800(CH), 125.238(CH), 119.561(C), 106.029(CH), 102.150(CH), 36.274(CH<sub>2</sub>), 31.803(CH<sub>2</sub>).

**[00103]** Representative compounds **13-27** of Formula III (see Table 3) were synthesized according to the same general reaction scheme using the appropriate starting materials. Overall yields varied from 35-55%. In some cases, the reductive high pressure hydrogenation reaction was successfully performed using Parr apparatus instead of CTH methods to obtain the final compounds.

#### **4-(4-hydroxyphenethylbenzene)-1,3-diol (13)**

**[00104]** <sup>1</sup>H-NMR (MeOD, 500MHz): δ 6.985 (dd, 2H, *J*=3 & 5Hz), 6.764 (d, 1H, *J* = 8Hz), 6.716-6.617 (m, 2H), 6.326 (d, 1H, *J* = 2.5Hz), 6.215 (dd, 1H, *J* = 2.5 & 8.5Hz), 2.719(s, 2H), 2.675(d, 2H, *J* = 8.5Hz). <sup>13</sup>C-NMR (MeOD, 125MHz): δ 155.756(C), 155.500(C), 154.560(C), 133.707(C), 130.354(CH), 129.125(2CH), 119.742(C), 114.586(2CH), 106.048CH), 102.127(CH), 35.393(CH<sub>2</sub>), 31.988(CH<sub>2</sub>).

#### **4-(4-methoxyphenethyl)benzene-1,3-diol (14)**

**[00105]** <sup>1</sup>H-NMR (MeOD, 500MHz): δ 7.052 (d, 2H, *J*=8.5Hz), 6.748-6.772 (m, 3H), 6.343(d, 1H, *J* = 2.5Hz), 6.222(dd, 1H, *J* = 2.5 & 8.0Hz), 3.683(s, 3H), 2.745(t, 4H, *J* = 6.0Hz). <sup>13</sup>C-NMR (MeOD, 125MHz): δ 157.700(C), 155.893(C), 155.601(C), 134.660(C), 130.317(CH), 129.073(2CH), 119.608(C), 113.255(2CH), 106.007(CH), 102.139(CH), 54.303(CH<sub>3</sub>), 35.350(CH<sub>2</sub>), 31.944(CH<sub>2</sub>).

#### **4-(2-(benzo[d][1,3]dioxol-5-ylethyl)benzene-1,3-diol (15)**

**[00106]** <sup>1</sup>H-NMR (MeOD, 500 MHz): δ 6.75(d, 1H, *J* = 8Hz), 6.67(d, 1H, *J* = 8 Hz), 6.66(d, 1H, *J* = 1.5Hz), 6.61(dd, 1H, *J* = 8 & 1.5Hz), 6.27(d, 1H, *J* = 2.5 Hz), 6.17(dd, 1H, *J*

= 8 & 2.5 Hz), 5.87 (s, 2H), 2.71-2.72(m, 4H).  $^{13}\text{C}$ -NMR (MeOD, 125 MHz):  $\delta$  156.01(C), 155.01 (C), 147.41(C), 145.47(C), 136.52(C), 130.14(CH), 120.87(CH), 119.23(C), 108.48(CH), 107.39(CH), 105.82(CH), 102.03(CH), 100.45(CH<sub>2</sub>), 35.95(CH<sub>2</sub>), 32.03(CH<sub>2</sub>).

**4,4'-(ethane-1,2diyl)dibenzene-1,3-diol (16)**

[00107]  $^1\text{H}$ -NMR (MeOD, 500 MHz):  $\delta$  6.81(d, 2H,  $J = 2\text{Hz}$ ), 6.29 (s, 2H), 6.20 (dd, 2H,  $J = 2$  & 2 Hz), 2.68 (s, 4H).  $^{13}\text{C}$ -NMR (MeOD, 125 MHz):  $\delta$  155.83 (2 C), 155.47 (2 C), 130.04 (2 CH), 145.47(C), 120.01 (2 C), 105.95(2 CH), 102.05(2 CH), 30.04(2 CH<sub>2</sub>).

**4-(2,4-dimethoxyphenethyl)benzene-1,3-diol (17)**

[00108]  $^1\text{H}$ -NMR (MeOD, 500 MHz):  $\delta$  6.93(d, 1H,  $J = 8\text{Hz}$ ), 6.73(d, 1H,  $J = 8$  Hz), 6.46(d, 1H,  $J = 2$  Hz), 6.37(dd, 1H,  $J = 8$  & 2 Hz), 6.28(d, 1H,  $J = 2$  Hz), 6.17(dd, 1H,  $J = 8$  & 2Hz), 3.77 (s, 3H), 3.74 (s, 3H), 2.67-2.73(m, 4H).  $^{13}\text{C}$ -NMR (MeOD, 125 MHz):  $\delta$  159.177 (C), 158.307 (C), 155.801(C), 155.564(C), 155.564 (C), 130.047 (CH), 129.758 (CH), 122.988(C), 119.900 (C), 105.796 (CH), 103.771 (CH), 101.991 (CH), 97.867 (CH), 54.392 (CH<sub>3</sub>), 54.289 (CH<sub>3</sub>), 29.901(CH<sub>2</sub>), 29.834(CH<sub>2</sub>).

**4-(3,5-dimethoxyphenethyl)benzene-1,3-diol (18)**

[00109]  $^1\text{H}$ -NMR (MeOD, 500 MHz):  $\delta$  6.77(d, 1H,  $J = 10\text{Hz}$ ), 6.34(d, 2H,  $J = 5$  Hz), 6.32(d, 1H,  $J = 5$  Hz), 6.25(t, 1H,  $J = 5$  Hz), 6.20(dd, 1H,  $J = 10$  & 5 Hz), 3.68 (s, 6H), 2.75 (s, 4H).  $^{13}\text{C}$ -NMR (MeOD, 125 MHz):  $\delta$  160.67 (C, 2C), 155.98 (C), 155.60 (C), 144.91 (C), 130.32 (CH), 119.38 (C), 106.20 (CH, 2C), 105.97 (CH), 102.08 (CH), 97.50 (CH), 54.25 (2 CH<sub>3</sub>), 36.48 (CH<sub>2</sub>), 31.47 (CH<sub>2</sub>).

**4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol (19)**

[00110]  $^1\text{H}$ -NMR (MeOD, 500 MHz):  $\delta$  6.972 (d, 1H,  $J=8.5\text{Hz}$ ), 6.624(d, 1H,  $J=9.0$  Hz), 6.285(dd, 1H,  $J=1$  & 2.5Hz), 6.184 (ddd, 1H,  $J = 1.5, 2.5$  & 8 Hz), 3.783 (s, 3H), 3.694 (s, 3H), 2.723-2.762 (m, 4H), 2.114(s, 3H).  $^{13}\text{C}$ -NMR (MeOD, 125 MHz):  $\delta$  157.034 (C), 156.915 (C), 155.953 (C), 155.664 (C), 130.066 (CH), 127.197 (C), 126.967 (CH), 119.834 (C), 118.705 (C), 105.855 (CH), 105.841 (CH), 102.046 (CH), 59.749 (CH<sub>3</sub>), 54.637 (CH<sub>3</sub>), 31.008 (CH<sub>2</sub>), 29.916 (CH<sub>2</sub>), 7.941 (CH<sub>3</sub>).

**4-(2,4,6-trimethoxyphenethyl)benzene-1,3-diol (20)**

[00111]  $^1\text{H}$ -NMR (MeOD, 500 MHz):  $\delta$  6.698. (dd, 1H,  $J = 1.5$  & 8Hz), 6.262 (t, 1H,  $J = 2.5\text{Hz}$ ), 6.135-6.157 (m, 3H), 3.775 (s, 3H), 3.730 (s, 3H), 3.723(s, 3H), 2.750 (t, 2H,  $J = 8.0\text{Hz}$ ), 2.563 (t, 2H,  $J = 8.0\text{Hz}$ ).  $^{13}\text{C}$ -NMR (MeOD, 125 MHz):  $\delta$  159.369 (C), 158.799 (2C), 155.638 (C), 155.556 (C), 129.840 (CH), 120.359 (C), 111.082 (C), 105.696 (CH), 101.906 (CH), 90.367 (2 CH), 54.744 (2 CH<sub>3</sub>), 54.281 (CH<sub>3</sub>), 28.790 (CH<sub>2</sub>), 22.653 (CH<sub>2</sub>).

#### 4-(1-phenylpropan-2-yl)benzene-1,3-diol (21)

[00112]  $^1\text{H}$ -NMR (MeOD, 500Hz):  $\delta$  6.928 (d, 2H,  $J = 8.5\text{Hz}$ ), 6.8541(d, 1H,  $J = 8.0\text{Hz}$ ), 6.631(d, 2H,  $J = 8.0\text{Hz}$ ), 6.259(d, 1H,  $J = 2.0\text{Hz}$ ), 6.219(dd, 1H,  $J = 2.5$  & 8.0Hz), 3.231(q, 1H,  $J = 6.5\text{Hz}$ ), 2.854(dd, 1H,  $J = 5.5$  & 13.0Hz), 2.519(dd, 1H,  $J = 9.0$  & 13.5Hz), 1.087(d, 3H,  $J = 7.0\text{Hz}$ ).  $^{13}\text{C}$ -NMR(MeOD, 500Hz):  $\delta$  155.482(C), 155.093(C), 153.428(C), 154.716(C), 132.452(CH), 129.673(2CH), 124.583(C), 114.242(2CH), 105.870(CH), 102.016(CH), 42.284(CH<sub>2</sub>), 33.835(CH), 18.575(CH<sub>3</sub>).

#### 4-(2-(furan-3-yl)ethyl)benzene-1,3-diol (22)

[00113]  $^1\text{H}$ -NMR (CDCl<sub>3</sub>, 300MHz):  $\delta$  7.348(s, 2H), 7.187(s, 1H), 6.342 (d, 1H,  $J = 8.1$  Hz), 6.282 (d, 2H,  $J = 10.2$  Hz), 2.795-2.663 (m, 4H).  $^{13}\text{C}$ -NMR (CDCl<sub>3</sub>, 75 MHz):  $\delta$  154.903(C), 154.594(C), 143.043(CH), 139.307(CH), 131.236(CH), 124.802(C), 120.509(C), 111.259(CH), 107.970(CH), 103.249(CH), 30.310(CH<sub>2</sub>), 25.544(CH<sub>2</sub>).

#### 4-(2-(tetrahydrofuran-3-yl)ethyl)benzene-1,3-diol (23)

[00114]  $^1\text{H}$ -NMR (CDCl<sub>3</sub>, 300MHz):  $\delta$  6.933 (d, 1H,  $J = 8.4\text{Hz}$ ), 6.359-6.283 (m, 2H), 3.951-3.709 (m, 3H), 3.388 (t, 1H,  $J = 7.5\text{Hz}$ ), 2.602-2.492 (m, 2H), 2.262-2.163 (m, 2H), 1.707-1.585 (m, 3H).  $^{13}\text{C}$ -NMR (CDCl<sub>3</sub>, 75 MHz):  $\delta$  155.056, 154.544, 132.595, 120.639, 110.008, 107.810, 103.074, 68.188, 64.108, 33.905, 32.718, 28.432.

#### 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol (24)

[00115]  $^1\text{H}$ -NMR (MeOD, 500 MHz):  $\delta$  7.31(d, 1H,  $J = 2\text{Hz}$ ), 6.78(d, 1H,  $J = 8$  Hz), 6.28(d, 1H,  $J = 2$  Hz), 6.24(dd, 1H,  $J = 3$  & 2 Hz), 6.18(dd, 1H,  $J = 8$  & 2.5 Hz), 5.95(dd, 1H,  $J = 3$  Hz), 2.78-2.82(m, 4H).  $^{13}\text{C}$ -NMR (MeOD, 125 MHz):  $\delta$  156.145 (C), 156.104 (C), 155.675(C), 140.334(CH), 129.995 (CH), 118.790 (C), 109.576 (CH), 105.867(CH), 104.315 (CH), 102.020 (CH), 28.209(CH<sub>2</sub>), 28.150(CH<sub>2</sub>).

**4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol (25)**

[00116]  $^1\text{H-NMR}$  (MeOD, 500 MHz):  $\delta$  6.83(d, 1H,  $J = 8\text{Hz}$ ), 6.26(d, 1H,  $J = 2.5\text{ Hz}$ ), 6.21(dd, 1H,  $J = 8 \text{ \& } 2.5\text{ Hz}$ ), 3.79-3.85 (m, 2H), 3.70-3.73 (m, 1H), 2.57-2.60 (m, 1H), 2.51-2.56 (m, 1H), 1.98-2.01 (m, 1H), 1.86-1.90 (m, 2H), 1.77-1.80 (m, 1H), 1.70-1.71 (m, 1H), 1.48-1.54 (m, 1H).  $^{13}\text{C-NMR}$  (MeOD, 125 MHz):  $\delta$  156.01 (C), 155.619 (C), 155.675(C), 129.958 (CH), 119.326 (C), 105.855 (CH), 102.130 (CH), 79.187(CH), 67.104 (CH<sub>2</sub>), 35.728 (CH<sub>2</sub>), 30.771 (CH<sub>2</sub>), 25.951(CH<sub>2</sub>), 25.218 (CH<sub>2</sub>).



**4-(2-(5-ethylofuran-2-yl)ethyl)benzene-1,3-diol (26)**












[00117]  $^1\text{H-NMR}$  (MeOD, 500 MHz):  $\delta$  6.786 (d, 1H,  $J = 8.0\text{Hz}$ ), 6.278(d, 1H,  $J = 2.5\text{Hz}$ ), 6.1845(dd, 1H,  $J = 2.5 \text{ \& } 8.0\text{Hz}$ ), 5.189(s, 2H), 2.766(s, 4H), 2.578(q, 2H,  $J = 7.5\text{Hz}$ ), 1.194(td, 3H,  $J = 7.5 \text{ \& } 1.0\text{Hz}$ ).  $^{13}\text{C-NMR}$  (MeOD, 125 MHz):  $\delta$  156.090(C), 155.649(C), 155.401(C), 154.142(C), 129.999(CH), 118.938(C), 105.852(CH), 104.734(CH), 103.779(CH), 102.006(CH), 28.309(CH<sub>2</sub>), 28.268(CH<sub>2</sub>), 20.828(CH<sub>2</sub>), 11.443(CH<sub>3</sub>).




**4-(2-(pyridin-3-yl)ethyl)benzene-1,3-diol (27)**

[00118]  $^1\text{H-NMR}$  (MeOD, 300 MHz):  $\delta$  8.293 (dd, 1H,  $J = 4.8 \text{ \& } 1.5\text{Hz}$ ), 8.250(d, 1H,  $J = 1.8\text{Hz}$ ), 7.605(dt, 1H,  $J = 8.1 \text{ \& } 1.5\text{Hz}$ ), 7.291(ddd, 1H,  $J = 7.5, 4.5 \text{ \& } 0.6\text{Hz}$ ), 6.673 (d, 1H,  $J = 8.4\text{Hz}$ ), 6.270 (d, 1H,  $J = 2.4\text{Hz}$ ), 6.139 (dd, 1H,  $J = 8.4 \text{ \& } 2.1\text{Hz}$ ), 2.880(td, 2H,  $J = 6.6 \text{ \& } 1.8\text{Hz}$ ), 2.778(td, 2H,  $J = 6.6 \text{ \& } 1.2\text{Hz}$ ).  $^{13}\text{C-NMR}$  (MeOD, 125 MHz):  $\delta$  156.06(C), 156.04(C), 149.00(CH), 146.00(CH), 138.90(C), 137.34(CH), 130.57(CH), 123.72(CH), 118.31(C), 106.03(CH), 102.25(CH), 33.14(CH<sub>2</sub>), 31.48(CH<sub>2</sub>).


Table 3. Compounds Representative of Formula III

| Compound                                             | Structure                                                                             | Mushroom Tyrosinase Assay              |
|------------------------------------------------------|---------------------------------------------------------------------------------------|----------------------------------------|
| 4-Phenethylbenzene-1,3-diol ( <b>12</b> )            |  | IC <sub>50</sub> : = 2.8 $\mu\text{M}$ |
| 4-(4-hydroxyphenethylbenzene)-1,3-diol ( <b>13</b> ) |   | IC <sub>50</sub> : = 2.9 $\mu\text{M}$ |

|                                                                    |                                                                                       |                              |
|--------------------------------------------------------------------|---------------------------------------------------------------------------------------|------------------------------|
| 4-(4-methoxyphenethyl)benzene-1,3-diol ( <b>14</b> )               |     | IC <sub>50</sub> : = 1.2 μM  |
| 4-(2-(benzo[d][1,3]dioxol-5-ylethyl)benzene-1,3-diol ( <b>15</b> ) |     | IC <sub>50</sub> : = 0.25 μM |
| 4,4'-(ethane-1,2diyl)dibenzene-1,3-diol ( <b>16</b> )              |     | IC <sub>50</sub> : = 0.35 μM |
| 4-(2,4-dimethoxyphenethyl)benzene-1,3-diol ( <b>17</b> )           |    | IC <sub>50</sub> : = 3.0 μM  |
| 4-(3,5-dimethoxyphenethyl)benzene-1,3-diol ( <b>18</b> )           |   | IC <sub>50</sub> : = 2.8 μM  |
| 4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol ( <b>19</b> )  |   | IC <sub>50</sub> : = 0.80 μM |
| 4-(2,4,6-trimethoxyphenethyl)benzene-1,3-diol ( <b>20</b> )        |   | IC <sub>50</sub> : = 40.0 μM |
| 4-(1-phenylpropan-2-yl)benzene-1,3-diol ( <b>21</b> )              |  | IC <sub>50</sub> : = 8.4 μM  |
| 4-(2-(furan-3-yl)ethyl)benzene-1,3-diol ( <b>22</b> )              |  | IC <sub>50</sub> : = 1.0 μM  |
| 4-(2-(tetrahydrofuran-3-yl)ethyl)benzene-1,3-diol ( <b>23</b> )    |  | IC <sub>50</sub> : = 5.8 μM  |
| 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol ( <b>24</b> )              |  | IC <sub>50</sub> : = 0.2 μM  |

|                                                                 |                                                                                     |                              |
|-----------------------------------------------------------------|-------------------------------------------------------------------------------------|------------------------------|
| 4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol ( <b>25</b> ) |  | IC <sub>50</sub> : = 4.5 μM  |
| 4-(2-(5-ethylfuran-2-yl)ethyl)benzene-1,3-diol ( <b>26</b> )    |  | IC <sub>50</sub> : = 0.86 μM |
| 4-(2-(pyridin-3-yl)ethyl)benzene-1,3-diol ( <b>27</b> )         |  | IC <sub>50</sub> : = 1.8 μM  |


**Example 4. General and specific methods for the synthesis of compounds of Formula IV**



Formula IV

**[00119]** Compounds of Formula IV were prepared as described in Scheme 4. Briefly, the 1,3-diphenylpropanes illustrative of Formula IV were prepared by base catalyzed Aldol condensation to provide the respective chalcones, which were then reduced by catalytic transfer of hydrogen (CTH) using ammonium formate (AMF) and a Pd-C catalyst.

**Scheme 4**



**Representative procedure for Scheme 4: Synthesis of Compounds 28, 29 and 30**

**General method for aldol condensation reaction**

**[00120]** To a solution containing 2,4-dibenzoyloxyacetophenone (7.67 g, 23.1 mmol) and 3,5-dimethoxybenzaldehyde (3.80 g, 23 mmol) in 100 mL anhydrous MeOH was added

KOH (1.4 g, 25 mmol) and the mixture was stirred for 5h at room temperature. A yellow solid was precipitated out which was filtered and washed with water. The crude product was finally crystallized from MeOH to afford a yellow crystalline solid (9.97 g; 87%).

#### General method for CTH reduction

**[00121]** The condensation product,  $\alpha,\beta$ -unsaturated ketone (3 g, 7.5 mmol) was dissolved in 24 mL glacial acetic acid followed by the addition of 10% Pd on activated carbon (300 mg) and ammonium formate (3.92 g, 62 mmol) and the reaction mixture was refluxed for 4 h. After cooling down the reaction mixture was filtered through Celite™ to remove the catalyst. The crude product was concentrated at rotary evaporator and purified by column chromatography eluting with hexane and ethyl acetate (75:25) to yielded compound **28** (1.1 g, 50%) as off-white solid & compound **29** (0.57 g, 25%) as thick yellow liquid.

#### General method for demethylation

**[00122]** To a stirred solution of **28** (1.4g, 4.9 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under argon at -80°C, BBr<sub>3</sub> (2.67g 10.7 mmol) was added and the mixture was slowly warmed to 0°C. The reaction was quenched by adding ice-cooled water and methanol was added. The solvent was then removed in a rotary evaporator and the crude product was purified by column chromatography eluting with ethyl acetate/hexanes (25:75) to provide compound **30** (1.14 g, 90%) as a solid.

#### 4-(3-(3,5-dimethoxyphenyl)propyl)benzene-1,3-diol (**28**)

<sup>1</sup>H-NMR (MeOD, 500 MHz):  $\delta$  6.828 (d, 1H,  $J$ =8.5Hz), 6.344 (d, 2H,  $J$ =2.5Hz), 6.288(d, 1H,  $J$  = 2.5Hz), 2.265(t, 1H,  $J$  = 2.0Hz), 6.226(dd, 1H,  $J$  = 2.5Hz & 8.0Hz), 3.723(s, 6H), 2.503-2.558 (m, 4H), 1.800-1.826 (m, 2H). <sup>13</sup>C-NMR (MeOD, 125 MHz):  $\delta$  160.772 (2C), 155.860(C), 155.597(C), 145.065(C), 130.080(CH), 119.774(C), 106.055(2 CH), 105.937(CH), 102.094(CH), 97.326(CH), 54.226(2 CH<sub>3</sub>), 35.587(CH<sub>2</sub>), 31.563(CH<sub>2</sub>), 28.924(CH<sub>2</sub>).

#### 4-(3-(3,5-dimethoxyphenyl)-1-hydroxypropyl)benzene-1,3-diol (**29**)

**[00123]** <sup>1</sup>H-NMR (MeOD, 500 MHz):  $\delta$  6.977 (d, 1H,  $J$ =8.0Hz), 6.316-6.337 (m, 3H), 6.309(t, 1H,  $J$ =2.5Hz), 6.259(t, 1H,  $J$ =2.5Hz), 4.781(dd, 1H,  $J$ =5.5Hz,  $J$ 2=7.5Hz), 3.704(s, 6H), 2.623(ddd, 1H,  $J$ =13.5Hz,  $J$ 2=10.0Hz,  $J$ 3=5.5Hz), 2.5222(ddd, 1H,  $J$ =20.0Hz,  $J$ 2=9.5Hz,  $J$ 3=6.5Hz), 2.031-1.908(m, 2H). <sup>13</sup>C-NMR (MeOD, 125 MHz):  $\delta$

160.806(2C), 157.356(C), 156.189(C), 144.503(C), 128.070(CH), 118.275(C), 106.640(CH), 106.155(2 CH), 102.239(CH), 97.515(CH), 78.440(CH), 54.185(CH<sub>3</sub>), 54.059(CH<sub>3</sub>), 37.786(CH<sub>2</sub>), 32.070(CH<sub>2</sub>).

**4-(3-(3,5-dihydroxyphenyl)propyl)benzene-1,3-diol (30)**

[00124] <sup>1</sup>H-NMR (MeOD, 500MHz): δ 6.827 (d, 1H, *J* = 8.0Hz), 6.267(d, 1H, *J* = 2.0Hz), 6.211(dd, 1H, *J* = 2.5 & 8.5Hz), 6.142 (d, 2H, *J* = 2.0Hz), 6.074 (t, 1H, *J* = 2.5Hz), 2.507(t, 2H, *J* = 8.0Hz), 2.455(t, 2H, *J* = 8.0Hz), 1.76-1.84(m, 2H). <sup>13</sup>C-NMR (MeOD, 125MHz): δ 157.848(2C), 155.849(C), 155.582(C), 145.065(C), 129.995(CH), 119.793(C), 106.559(2CH), 105.859(CH), 102.054(CH), 99.514(CH), 35.420(CH<sub>2</sub>), 31.593(CH<sub>2</sub>), 28.961(CH<sub>2</sub>).

**4-(3-(2,4,6-trimethoxyphenyl)propyl)benzene-1,3-diol (31)**

[00125] Compound **31** was synthesized following a similar procedure as described for compound **28** above using as starting materials 2,4-dibenzoyloxyacetophenone and 2,4,6-trimethoxybenzaldehyde. <sup>1</sup>H-NMR (MeOD, 500 MHz): δ 6.822 (d, 1H, *J* = 8.0Hz), 6.250(d, 1H, *J* = 2.5Hz), 6.197(dd, 1H, *J* = 2.5 & 8.0Hz), 6.165(s, 2H), 3.768(s, 3H), 3.764(s, 6H), 2.562(t, 2H, *J* = 7.5Hz), 2.461(t, 2H, *J* = 7.5Hz), 1.617-1.653(m, 2H). <sup>13</sup>C-NMR (MeOD, 125 MHz): δ 159.273(C), 158.729(2C), 155.556(C), 155.449(C), 129.655(2CH), 120.426(C), 111.382(C), 105.789(CH), 101.968(CH), 90.285(CH), 54.655(2CH<sub>3</sub>), 54.274(CH<sub>3</sub>), 29.716(CH<sub>2</sub>), 29.064(CH<sub>2</sub>), 22.142(CH<sub>2</sub>).

**4-(3-(2,4-dihydroxyphenyl)propyl)-2-methylbenzene-1,3-diol (32)**


[00126] Compound **32** was synthesized following a similar procedure as described for compound **30** using as starting materials 2,4-dibenzoyloxyacetophenone and 2,4-dimethoxy-3-methylbenzaldehyde. <sup>1</sup>H-NMR (MeOD, 500 MHz): δ 6.879 (d, 1H, *J* = 8.5Hz), 6.747(d, 1H, *J* = 8.0Hz), 6.352(d, 1H, *J* = 1.5Hz), 6.341(d, 1H, *J* = 4.0Hz), 6.285(dd, 1H, *J* = 8.0Hz & 2.5Hz), 2.567(dd, 4H, *J* = 8.0 & 7.5Hz), 2.133(s, 3H), 1.786-1.817 (m, 2H). <sup>13</sup>C-NMR (MeOD, 125 MHz): δ 155.486(C), 155.316(C), 153.428(C), 152.906(C), 130.295(CH), 126.523(CH), 120.933(C), 120.518(C), 111.556(C), 106.788(CH), 106.377(CH), 102.283 (CH), 30.767(CH<sub>2</sub>), 29.649(CH<sub>2</sub>), 29.127(CH<sub>2</sub>), 7.833(CH<sub>3</sub>).

**4-(3-(2,4-dihydroxy-3-methylphenyl)-3--(2,4-dihydroxyphenyl)propane-1-one (33)**

[00127] Compound **33** was synthesized following a similar procedure as described for compound **28** using as starting materials 2,4-dibenzoyloxyacetophenone and 2,4-dimethoxy-3-methylbenzaldehyde. The reduction with Pd-C/ ammonium formate was carried out at room temperature instead of refluxing. Demethylation reaction with borontribromide was performed using as similar protocol described as in the procedure for compound **30**. <sup>1</sup>H-NMR (MeOD, 500MHz):  $\delta$  7.628 (d, 1H,  $J$  = 9.0Hz), 6.868 (d, 1H,  $J$  = 8.0Hz), 6.349 (d, 1H,  $J$  = 9.0Hz), 6.283 (d, 1H,  $J$  = 2.5Hz), 6.203 (dd, 1H,  $J$  = 2.5 & 8.0Hz), 3.127 (t, 2H,  $J$  = 7.5Hz), 2.862 (t, 2H,  $J$  = 7.5Hz), 2.031 (s, 3H). <sup>13</sup>C-NMR (MeOD, 500Hz):  $\delta$  205.420 (C), 162.894 (C), 162.264 (C), 156.438 (C), 155.782 (C), 130.710 (CH), 129.510 (CH), 118.364 (C), 112.249 (C), 110.820 (C), 106.596 (CH), 105.970 (CH), 102.142 (CH), 47.089 (CH<sub>2</sub>), 25.810 (CH<sub>2</sub>), 6.361 (CH<sub>3</sub>).

Example 5. Synthesis of 4-(5-hydroxyphenyl)benzene-1,3-diol (**34**)

[00128] Compound **34** was synthesized as set forth in Scheme 5



[00129] To a solution of the phosphonium salt of 2,4-bis(benzoyloxy)benzyl bromide (7.72, 12 mmol) in THF (25 mL) was added nBu-Li in hexane (6.9 mL, 11 mmol) at 0°C. The reaction mixture was allowed to stir at room temperature for 1 h, during that period the mixture developed a red color indicating the formation of ylide. A solution of 4-(tetrahydropyranyloxy)butanal (1.72 g, 10 mmol) in THF (10 mL) was added to the ylide solution at -80°C and the mixture was allowed to stir overnight without removing the low temperature bath, during that period the red color disappeared and a whitish suspension was formed which was diluted with hexane and filtered. The filtrate was concentrated and residue was passed through a short bed of silica gel eluting with ethyl acetate-hexane (5:95) to give the coupled product (5.2 g, 92%) in a ratio of ( $E/Z$  : 20/80) as pale yellow oil.




**[00130]** The coupled product (3.6 g, 7.7 mmol) was then dissolved in 24 mL THF/MeOH (1:2) followed by treatment with 10% Pd on activated carbon (360 mg) and ammonium formate (3.92 g, 62 mmol) at reflux for 4 h. After cooling down, the reaction mixture was filtered through celite to remove the catalyst. The crude product, after concentration, was purified by column chromatography eluting with hexane and ethyl acetate (80:20) to yield the reduced product (2.15 g, 95%) as a colorless thick liquid.

**[00131]** To a stirred solution of the reduced product (1.2 g, 4 mmol) in ethanol (20 mL) was added pyridinium toluene-4-sulfonate (350mg, 1.39 mmol) at room temperature and the mixture was stirred for 24 h. Removal of the solvent in vacuum followed by column chromatography of the residue on silica gel eluting with ethyl acetate–hexane (10:90) gave the title compound (**34**) as off-white solid (690 mg, 88%). <sup>1</sup>H-NMR (MeOD, 500MHz):  $\delta$  8.825(d, 1H,  $J = 7.5$ Hz), 6.2705 (d, 1H,  $J = 0.5$ Hz), 6.215(d, 1H,  $J = 7.5$ Hz), 3.538(t, 2H,  $J = 6.5$ Hz), 2.491(t, 2H,  $J = 7.5$ Hz), 1.526-1.585(m, 4H), 1.358-1.40(m, 2H). <sup>13</sup>C-NMR (MeOD, 125MHz):  $\delta$  155.705(C), 155.482(C), 129.984(CH), 120.108(C), 105.926(CH), 102.061(CH), 61.722(CH<sub>2</sub>), 32.189(CH<sub>2</sub>), 29.182(CH<sub>2</sub>), 29.086(CH<sub>2</sub>), 25.322(CH<sub>2</sub>), 103.761(CH).

**[00132]**

Table 4. Compounds Representative of Formula IV

| Compound                                                                  | Structure | Mushroom Tyrosinase Assay          |
|---------------------------------------------------------------------------|-----------|------------------------------------|
| 4-(3-(3,5-dimethoxyphenyl)propyl)benzene-1,3-diol ( <b>28</b> )           |           | IC <sub>50</sub> : = 1.02 $\mu$ M  |
| 4-(3-(3,5-dimethoxyphenyl)-1-hydroxypropyl)benzene-1,3-diol ( <b>29</b> ) |           | IC <sub>50</sub> : = 5.15 $\mu$ M  |
| 4-(3-(3,5-dihydroxyphenyl)propyl)benzene-1,3-diol ( <b>30</b> )           |           | IC <sub>50</sub> : = 0.5 $\mu$ M   |
| 4-(3-(2,4,6-trimethoxyphenyl)propyl)benzene-1,3-diol ( <b>31</b> )        |           | IC <sub>50</sub> : = 0.165 $\mu$ M |

|                                                                                        |                                                                                     |                             |
|----------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-----------------------------|
| 4-(3-(2,4-dihydroxyphenyl)propyl)-2-methylbenzene-1,3-diol ( <b>32</b> )               |  | IC <sub>50</sub> : = 2.3 μM |
| 4-(3-(2,4-dihydroxy-3-methylphenyl)-3-(2,4-dihydroxyphenyl)propane-1-one ( <b>33</b> ) |  | IC <sub>50</sub> = 2.54 μM  |
| 4-(5-hydroxypentyl)benzene-1,3-diol ( <b>34</b> )                                      |  | IC <sub>50</sub> = 1.0 μM   |

### Example 6. Tyrosinase Assay

#### Reagents

**[00133]** Tyrosinase isolated from the mushroom species *Agaricus bisporus* was purchased from the Sigma-Aldrich Inc. (Cat# T3824-50KU). The enzyme was dissolved in the tyrosinase assay buffer (100 mM sodium phosphate, pH 6.8) to a concentration 10 U/μl, and stored at -70°C. For the experiments the enzyme was freshly diluted in the assay buffer to a concentration of 0.2 U/μl.

**[00134]** All test compounds were initially dissolved in 100% DMSO at a concentration of 400 mM. The compounds were further diluted in 100% DMSO to a concentration of 80 μM. 80 μM stock solutions of the compounds were then diluted ten times in the tyrosinase assay buffer to the concentration of 8 μM. The compounds were then serially diluted at three-fold increments in the assay buffer containing 10% DMSO, keeping the concentration of DMSO constant across all samples. These serially diluted compounds were subsequently used as two-fold concentrated stock solutions in the tyrosinase activity assays.

**[00135]** Tyrosinase substrate, L-DOPA (Sigma-Aldrich Inc, Cat # 37830) was dissolved in the tyrosinase assay buffer to a concentration of 4 mM. This solution was then used as the four-fold concentrated stock of substrate in the tyrosinase activity assays.

#### Assay conditions

**[00136]** Tyrosinase assays were performed in clear-bottom 96-well plates at room temperature. The final volume of the assays was 200 μl per well. 100 μl of two-fold concentrated test compounds were mixed with 50 μl of 4 mM L-DOPA. The reactions were initiated by adding 50 μl of mushroom tyrosinase (0.2 U/μl, 10 U per reaction), and allowed to proceed for 15 minutes. Accumulation of colored product was monitored by light absorption at 450 nm using Victor 2 plate reader (Perkin-Elmer Inc.).

## Results

[00137] The assays were performed in triplicate and covered a concentration range of test compounds from 4  $\mu$ M to 5.5 nM. Mean (n=3) absorption of wells containing no enzyme was subtracted as blanks. The data was computed as percentage activity of wells that contained the tyrosinase, but no test compounds. IC<sub>50</sub> values were computed from non-linear regression fits using GraphPad Prism software. The results for representative compounds of Formulas I-IV are set forth in Tables 1-4 below, and Figures 1-9.

### Example 7. Murine Melanoma cell-based assays

[00138] Selected compounds were then tested for the ability to suppress melanin production by the murine melanoma cells B16-F1 as detailed below.

## Materials

[00139] Murine melanoma cells B16-F1 were purchased from the ATCC (Cat # CRL-6323). CellTiter\*96 AqueousOne Solution was purchased from Promega (Cat# G3581). 0.2  $\mu$ m pore size, low protein binding filters were purchased from PALL Life Sciences. (Cat # PN4454). Alpha-MSH was purchased from the Bachem Inc. (Cat # H-1075.0005). All tissue culture reagents were purchased from the Invitrogen Inc.

[00140] The B16-F1 cells were maintained in cell growth media (DMEM/High Glucose supplemented with glutamine, sodium pyruvate, 10% dialyzed fetal calf serum, 1% non-essential amino acids, 50 units/mL penicillin, and 50  $\mu$ g/mL of streptomycin) at 37°C in a humidified atmosphere of 95% air and 5% CO<sub>2</sub>.

[00141] All test compounds were initially dissolved in 100% DMSO at a concentration of 400 mM. Then, 6  $\mu$ L of 400 mM test compounds were added to 1.2 mL of the cell growth media supplemented with 200 nM alpha-MSH, giving a final concentration of 2 mM for the test compound and concentration of 0.5% for DMSO. The compounds were centrifuged for 1 h at 20,000xg. Supernatants (1 mL) were collected and filtered through sterile 0.2  $\mu$ m filters. The compounds were serially diluted in two-fold increments in the sterile cell growth media supplemented with 0.5% DMSO and 200 nM alpha-MSH, thus keeping the concentration of alpha-MSH and DMSO constant for all samples. These serially diluted compounds were subsequently used as two-fold concentrated stock solutions in the melanin production and cell viability assays.

\*Trade-mark

## Methods

**[00142]** B16-F1 cells were seeded into the wells of clear-bottom 96-well plates at 40,000 cells per well, in 100  $\mu$ l of the cell growth media. On the following day, 100  $\mu$ l of freshly prepared two fold concentrated test compounds were added to the wells. The cells were maintained at 37°C in a humidified atmosphere of 95% air and 5% CO<sub>2</sub> for 72 hours. At the end of the incubation period melanin-containing conditioned media was removed from the cells and transferred to another plate. To measure melanin content the light absorbance of conditioned media was taken at 450 nm using a Victor 2 plate reader (Perkin-Elmer Inc.).











**[00143]** Viability of cells was measured using standard tetrazolium reduction assay based on redox potential of live cells. After aspiration of melanin-containing conditioned media, the cells were immediately replenished with 100  $\mu$ l of fresh media supplemented with 16% CellTiter96 AqueousOne solution containing tetrazolium salts. The cells were maintained at 37°C for additional 20-40 minutes. Conversion of tetrazolium was monitored by measuring absorbance of cell wells at 450 nm using a Victor 2 plate reader (Perkin-Elmer Inc.).

## Results

**[00144]** The results of the representative experiments are summarized in Table 5 and Figures 1-9. More than 90% percent of the melanin produced by cultured melanoma cells is found in extra-cellular media. Therefore, at the end of the experiment melanin-containing media was collected and relative amounts of melanin were determined by adsorption at 450 nm. Viability of cells was determined by a commonly used colorimetric procedure that is based on conversion of the tetrazolium compounds to colored formazan products (using Promega's CellTiter96 AqueousOne assay). Dehydrogenase enzymes in metabolically active cells accomplish this conversion, and the amount of formazan product is directly proportional to the number of living cells in culture.

**[00145]** The assays were performed in quadruplicate and covered a concentration range of test compounds from 2  $\mu$ M to 1000  $\mu$ M. Mean (n=4) absorption of wells containing no cells was subtracted as blanks. The results were computed as percent of wells that contained the cells but no test compounds. IC<sub>50</sub> values were computed from non-linear regression fits using GraphPad Prism software. The results are set forth in Table 5.

Table 5. Results of Mushroom Tyrosinase Assay, Melanin Production Assay and Cell Viability Studies for representative compounds of Formulas I-IV

| Compound          | Structure                                                                           | Mushroom Tyrosinase Assay     | Melanin production            | Cell viability                |
|-------------------|-------------------------------------------------------------------------------------|-------------------------------|-------------------------------|-------------------------------|
| <b>7</b>          |    | IC <sub>50</sub> : = 2.0 μM   | IC <sub>50</sub> : = 14 μM    |                               |
| <b>15</b>         |    | IC <sub>50</sub> : = 0.25 μM  | IC <sub>50</sub> : = 3.3 μM   | LD <sub>50</sub> : = 66 μM    |
| <b>16</b>         |    | IC <sub>50</sub> : = 0.35 μM  | IC <sub>50</sub> : = 1.7 μM   | LD <sub>50</sub> : = 260 μM   |
| <b>17</b>         |   | IC <sub>50</sub> : = 3.0 μM   | IC <sub>50</sub> : = 11.9 μM  | LD <sub>50</sub> : = 157.8 μM |
| <b>18</b>         |  | IC <sub>50</sub> : = 2.8 μM   | IC <sub>50</sub> : = 2.97 μM  | LD <sub>50</sub> : = 222.2 μM |
| <b>19</b>         |  | IC <sub>50</sub> : = 0.80 μM  | IC <sub>50</sub> : = 24.7 μM  | LD <sub>50</sub> : = 130.3 μM |
| <b>24</b>         |  | IC <sub>50</sub> : = 0.2 μM   | IC <sub>50</sub> : = 1.60 μM  | LD <sub>50</sub> : = 345.9 μM |
| <b>25</b>         |  | IC <sub>50</sub> : = 4.5 μM   | IC <sub>50</sub> : = 5 μM     | LD <sub>50</sub> : >1000 μM   |
| <b>31</b>         |  | IC <sub>50</sub> : = 0.165 μM | IC <sub>50</sub> : = 156.1 μM | LD <sub>50</sub> : = 187.8 μM |
| <b>kojic acid</b> |  | IC <sub>50</sub> : = 20 μM    | IC <sub>50</sub> : = 303.5 μM | LD <sub>50</sub> : = >1000 μM |

Example 8. Reconstructed Human Skin Studies**Materials and Methods**

**[00146]** The skin whitening effects of test compounds were studied using a reconstructed skin model, Melanoderm™ provided by MatTek Corp. (Ashland, MA)

according to the manufacturer's specifications. Briefly, normal human epidermal keratinocytes and normal human melanocytes derived from dark skin donors were co-cultured on a surface of collagen-coated membrane to form multi-layered, highly differentiated skin tissue (MEL-300-B). The tissues were maintained in the CO<sub>2</sub> incubator at 37°C. The apical surfaces of the reconstructed skin (9 mm in diameter) were exposed to air whereas the bottom surfaces remained in contact with 5 mL of maintenance medium, containing skin differentiating factors (EPI-100-NNM-113). Test compounds were formulated in 80% propylene glycol as follows: 10 mg of each test compound was dissolved overnight in 1 mL of propylene glycol (1,2-propanediol, Sigma-Aldrich). The compounds were then sterilized by passing through 0.2 µm filter and diluted in sterile water/propylene glycol to final concentrations 0.2%, 0.4% and 0.8%. The concentration of propylene glycol was kept 80% for all samples. In addition the following controls were used: sterile water, 80% propylene glycol, and 1% kojic acid in water.

**[00147]** Test compounds were applied to the apical surface of the tissues as follows: 10 µl of each test compound, 10 µl of 80% propylene glycol (vehicle control), 25 µl of sterile water (negative control) and 25 µl of 1% kojic acid (positive control). The samples were re-applied every other day for 15 days. All the samples were tested in duplicate. At the end of the experiment microscopic images were taken using photo-microscope equipped with 25X phase contrast objective and color CCD camera.

## Results


**[00148]** The potential skin-whitening properties of test compounds were further explored in a reconstructed skin model. The model consists of normal, human-derived epidermal keratinocytes and melanocytes, which have been co-cultured to form a multilayered, highly differentiated human epidermis. In this study the melanocytes were obtained from a highly pigmented donor. The reconstructed skin was grown at the air-liquid interface, making it possible to mimic topical application of skin whitening agents (Figure 10A).

**[00149]** Different concentrations of test compounds, 80% propylene glycol (the vehicle control), water (negative control) or 1% kojic acid (positive control) were repeatedly applied topically on the surface of the reconstructed skin for 15 days. Two of the test compounds, namely, compound # 16, and compound # 25 exhibited significant whitening effects on skin melanocytes, without causing any detectable alterations of cell morphology (Figure 10B). Of them, compound #16 exhibited the greatest effect with significant whitening of

melanocytes observed as early as 3 days after the beginning of the experiment (data not shown). Photographs of skin specimen taken after 15 days of the experiment show significant dose-dependent whitening effects on melanocytes, which appear on photographs as dark dendritic cells.

## CLAIMS

1. A compound of formula I:




Formula I

or a pharmaceutically acceptable salt thereof, wherein:

R is substituted or unsubstituted biphenyl, substituted or unsubstituted pyrid-3-yl, substituted or unsubstituted pyrid-4-yl, substituted or unsubstituted furan-3-yl or a substituted phenyl; wherein, when substituted biphenyl, substituted pyrid-3-yl, substituted pyrid-4-yl or substituted furan-3-yl, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group and when substituted phenyl, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group and a C1-C10 alkoxy group.

2. The compound of claim 1, wherein R is pyrid-3-yl or biphenyl-4-yl.
3. The compound of claim 1, wherein R is 1'-(4'-methoxy)phenyl or 1'-(2',4'-dihydroxy)phenyl.
4. A compound selected from the group consisting of 4'-methoxybiphenyl-2,4-diol (1), biphenyl-2,2',4,4'-tetraol (2), 4-(pyridin-3-yl)benzene-1,3-diol (3) and 4-(4'-hydroxybiphenyl)benzene-1,3-diol (4).

5. A compound of formula II:



Formula II

or a pharmaceutically acceptable salt thereof, wherein:


X is H, or OH; and

R is substituted phenyl or substituted or unsubstituted: biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, or furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group.

6. The compound of claim 5, wherein R is selected from the group consisting of 3'-furyl; 3'-pyridyl; 3'-(2-methoxypyridyl); and 1'-(2',4',6'-trimethoxy phenyl); or wherein X is OH and R is 3'-pyridyl.

7. A compound selected from the group consisting of 4-(furan-3-ylmethyl)benzene-1,3-diol (6), 4-(pyridin-ylmethyl)benzene-1,3-diol (7), 4-((2-methoxypyridin-3-yl)methyl)benzene-1,3-diol (8), 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9), and 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11).

8. A compound of formula III:



or a pharmaceutically acceptable salt thereof, wherein:

R<sub>1</sub> is H or C<sub>1</sub>-C<sub>10</sub> alkyl; and


R<sub>2</sub> is selected from the group consisting of 4-hydroxyphen-1-yl, 4-methoxyphen-1-yl, 2,4-dimethoxyphen-1-yl, 3,5-dimethoxyphen-1-yl, 2,4-dimethoxy-3-methylphen-1-yl, 2,4,6-trimethoxyphen-1-yl, furan-3-yl, tetrahydrofuran-3-yl, furan-2-yl, tetrahydrofuran-2-yl, 5-ethylfuran-2-yl and pyridine-3-yl.

9. The compound of claim 8, wherein R<sub>1</sub> is H.

10. The compound of claim 8, wherein  $R_1$  is Me.

11. A compound selected from the group consisting of 4-(4-hydroxyphenethylbenzene)-1,3-diol (13), 4-(4-methoxyphenethylbenzene)-1,3-diol (14), 4-(2,4-dimethoxyphenethylbenzene)-1,3-diol (17), 4-(3,5-dimethoxyphenethylbenzene)-1,3-diol (18), 4-(2,4-dimethoxy-3-methylphenethylbenzene)-1,3-diol (19), 4-(2,4,6-trimethoxyphenethylbenzene)-1,3-diol (20), 4-(2-(furan-3-yl)ethylbenzene)-1,3-diol (22), 4-(2-(tetrahydrofuran-3-yl)ethylbenzene)-1,3-diol (23), 4-(2-(furan-2-yl)ethylbenzene)-1,3-diol (24), 4-(2-(tetrahydrofuran-2-yl)ethylbenzene)-1,3-diol (25), 4-(2-(5-ethylofuran-2-yl)ethylbenzene)-1,3-diol (26), and 4-(2-(pyridin-3-yl)ethylbenzene)-1,3-diol (27).

12. A compound of formula IV:



wherein  $==$  represents a double or single bond;

$R_1$  is H or  $C_1$ - $C_{10}$  alkyl;

$R_4$  is H, OH or O; and

$R_5$  is a substituted or unsubstituted: phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl, furan-3-yl or tetrahydrofuran-2-yl; wherein, when substituted,  $R_5$  is substituted with 1 to 3 moieties ( $R'$ ,  $R''$ ,  $R'''$ ) independently selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group and a hydroxyl group;

with the following provisos:

proviso 1: when  $R_1$  and  $R_4$  are H;  $R_5$  cannot be 1'-(2,4-dimethoxy-3-methyl)phenyl;


proviso 2: when  $R_1$  and  $R_4$  are H;  $R_5$  cannot be 1'-(2,4-dihydroxy)phenyl; 1'-(2,4-dialkoxy)phenyl; 1'-(2,4-dialkenoxy)phenyl or 1'-(2,4-dibenzyloxy)phenyl;

proviso 3:  $R_5$  cannot be 1'-(4-hydroxy)phenyl;

proviso 5: when  $R_1$  and  $R_4$  are H;  $R_5$  cannot be a 1'-(2,4-disubstituted)phenyl; wherein said substituents are selected from  $-OR^2$  and  $-OR^3$ , wherein  $R^2$  and  $R^3$  are independently selected from H and a  $(C_1-C_{10})$ -alkyl; and

proviso 6: when  $R_1$  is H and  $R_4$  is OH;  $R_5$  cannot be 1'-(3,4-dihydroxy)phenyl; 1'-(3,4-dimethoxy)phenyl; 1'-(2-hydroxy)phenyl; 1'-(2-methoxy)phenyl; 1'-(4-methoxy)phenyl; phenyl; 1'-(3-methoxy,4-hydroxy)phenyl.

13. A composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound of formula I:



Formula I

or a pharmaceutically acceptable salt thereof, wherein:


$R$  is substituted or unsubstituted biphenyl, substituted or unsubstituted pyrid-3-yl, substituted or unsubstituted pyrid-4-yl, substituted or unsubstituted furan-3-yl or a substituted phenyl; wherein, when substituted biphenyl, substituted pyrid-3-yl, substituted pyrid-4-yl or substituted furan-3-yl,  $R$  is substituted with 1 to 3 moieties ( $R'$ ,  $R''$ ,  $R'''$ ) independently selected from a  $C_1-C_{10}$  alkyl group, a  $C_1-C_{10}$  alkoxy group and a hydroxyl group and when substituted phenyl,  $R$  is substituted with 1 to 3 moieties ( $R'$ ,  $R''$ ,  $R'''$ ) independently selected from a  $C_1-C_{10}$  alkyl group and a  $C_1-C_{10}$  alkoxy group.

14. The composition of claim 13, wherein  $R$  is selected from the group consisting of 1'-(3-pyridyl); 1'-(biphenyl-4-ol).

15. The composition of claim 13, wherein R is 1'-(4'-methoxy)phenyl or 1'-(2',4'-dihydroxy)phenyl.

16. A composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound, wherein the compound is 4'-methoxybiphenyl-2,4-diol (1), biphenyl-2,2',4,4'-tetraol (2), 4-(pyridin-3-yl)benzene-1,3-diol (3) or 4-(4'-hydroxybiphenyl)benzene-1,3-diol (4).

17. A composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound of formula II:



Formula II

or a pharmaceutically acceptable salt thereof, wherein:


X is H, or OH; and

R is substituted phenyl or substituted or unsubstituted: biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, or furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group.

18. The composition of claim 17, wherein R is 3'-furyl; 3'-pyridyl; 3'-(2-methoxypyridyl); or 1'-(2',4',6'-trimethoxy phenyl); or wherein X is OH and R is 3'-pyridyl.

19. The composition of claim 18, wherein the compound is selected from the group consisting of 4-(furan-3-ylmethyl)benzene-1,3-diol (6), 4-(pyridin-ylmethyl)benzene-1,3-diol (7), 4-((2-methoxypyridin-3-yl)methyl)benzene-1,3-diol (8), 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9), and 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11).

20. A composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound of formula III:



Formula III

or a pharmaceutically acceptable salt thereof, wherein:

$R_1$  is H or  $C_1$ - $C_{10}$  alkyl; and


$R_2$  is 4-hydroxyphen-1-yl, 4-methoxyphen-1-yl, 2,4-dimethoxyphen-1-yl, 3,5-dimethoxyphen-1-yl, 2,4-dimethoxy-3-methylphen-1-yl, 2,4,6-trimethoxyphen-1-yl, furan-3-yl, tetrahydrofuran-3-yl, furan-2-yl, tetrahydrofuran-2-yl, 5-ethylfuran-2-yl and pyridine-3-yl.

21. The composition of claim 20 wherein  $R_1$  is  $-CH_3$ .

22. The composition of claim 20, wherein  $R_1$  is H.

23. A composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound, wherein the compound is 4-(4-hydroxyphenethylbenzene)-1,3-diol (13), 4-(4-methoxyphenethylbenzene)-1,3-diol (14), 4-(2,4-dimethoxyphenethylbenzene)-1,3-diol (17), 4-(3,5-dimethoxyphenethylbenzene)-1,3-diol (18), 4-(2,4-dimethoxy-3-methylphenethylbenzene)-1,3-diol (19), 4-(2,4,6-trimethoxyphenethylbenzene)-1,3-diol (20), 4-(2-(furan-3-yl)ethylbenzene)-1,3-diol (22), 4-(2-(tetrahydrofuran-3-yl)ethylbenzene)-1,3-diol (23), 4-(2-(furan-2-yl)ethylbenzene)-1,3-diol (24), 4-(2-(5-ethylfuran-2-yl)ethylbenzene)-1,3-diol (26) or 4-(2-(pyridin-3-yl)ethylbenzene)-1,3-diol (27).

24. A composition comprising a pharmaceutically or dermatologically acceptable carrier and a compound of formula IV:



Formula IV

wherein:

$\equiv$  represents a double or single bond;

$R_1$  is H or  $C_1$ - $C_{10}$  alkyl;

$R_4$  is H, OH or O; and

$R_5$  is a substituted or unsubstituted: phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl, furan-3-yl or tetrahydrofuran-2-yl, and when substituted,  $R_5$  is substituted with 1 to 3 moieties ( $R'$ ,  $R''$ ,  $R'''$ ) independently selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group and a hydroxyl group;

with the following provisos:

proviso 1: when  $R_1$  and  $R_4$  are H;  $R_5$  cannot be 1'-(2,4-dimethoxy-3-methyl)phenyl;

proviso 2: when  $R_1$  and  $R_4$  are H;  $R_5$  cannot be 1'-(2,4-dihydroxy)phenyl; 1'-(2,4-dialkoxy)phenyl; 1'-(2,4-dialkenoxy)phenyl or 1'-(2,4-dibenzyloxy)phenyl;

proviso 3:  $R_5$  cannot be 1'-(4-hydroxy)phenyl;

proviso 5: when  $R_1$  and  $R_4$  are H and  $R_5$  cannot be a 1'-(2,4-disubstituted)phenyl; wherein said substituents are selected from  $-OR^2$  and  $-OR^3$ , wherein  $R^2$  and  $R^3$  are independently selected from H and a  $(C_1-C_{10})$ -alkyl; and

proviso 6: when  $R_1$  is H and  $R_4$  is OH;  $R_5$  cannot be 1'-(3,4-dihydroxy)phenyl; 1'-(3,4-dimethoxy)phenyl; 1'-(2-hydroxy)phenyl; 1'-(2-methoxy)phenyl; 1'-(4-methoxy)phenyl; phenyl; or 1'-(3-methoxy,4-hydroxy)phenyl.

25. The composition of claim 24, wherein  $R_1$  and  $R_4$  are H and  $R_5$  is selected from 1'-(3',5'-dimethoxy)phenyl; 1'-(3',5'-dihydroxy)phenyl and 1-(2',4',6-trimethoxyphenyl); or wherein  $R_1$  is Me,  $R_4$  is OH, and  $R_5$  is 1'-(2',4'-dihydroxy)phenyl;

or wherein  $R_1$  is Me,  $R_4$  is O, and  $R_5$  is 1'-(2',4'-dihydroxy)phenyl; or wherein  $R_1$  and  $R_4$  are H, and  $R_5$  is  $\text{CH}_2\text{CH}_2\text{OH}$ .

26. The composition of claim 24, wherein the compound is 4-(3-(3,5-dimethoxyphenyl)propyl)benzene-1,3-diol (28), 4-(3-(3,5-dimethoxyphenyl)-1-hydroxypropyl)benzene-1,3-diol (29), 4-(3-(3,5-dihydroxyphenyl)propyl)benzene-1,3-diol (30), 4-(3-(2,4,6-trimethoxyphenyl)propyl)benzene-1,3-diol (31), 4-(3-(2,4-dihydroxyphenyl)propyl)-2-methylbenzene-1,3-diol (32), 4-(3-(2,4-dihydroxy-3-methylphenyl)-3-(2,4-dihydroxyphenyl)propane-1-one (33), or 4-(5-hydroxypentyl)benzene-1,3-diol (34).

27. Use of the composition of any one of claims 13-26 for inhibiting the activity of tyrosinase.

28. Use of the composition of any one of claims 13-26 for inhibiting the production of melanin.


29. Use of the composition of any one of claims 13-26 for preventing or treating diseases of conditions related to the overproduction or uneven distribution of melanin.

30. The use of claim 29, wherein the diseases or conditions are selected from hyperpigmentation or damage resulting from exposure to the sun or ultraviolet (UV) radiation, suntan, hyper pigmentation spots caused by skin aging, melasma, liver diseases, thermal burns, topical wounds, skin pigmentation due to inflammatory conditions caused by fungal, microbial or viral infections, vitiligo, carcinoma and melanoma.

31. Use of the composition of any one of claims 13-26 for whitening or lightening skin, or both.

32. Use of a compound selected from:

a) a compound of formula I:




Formula I

or a pharmaceutically acceptable salt thereof, wherein:

R is substituted or unsubstituted biphenyl, substituted or unsubstituted pyrid-3-yl, substituted or unsubstituted pyrid-4-yl, substituted or unsubstituted furan-3-yl or a substituted phenyl; wherein, when substituted biphenyl, substituted pyrid-3-yl, substituted pyrid-4-yl or substituted furan-3-yl, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group and when substituted phenyl, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group and a C1-C10 alkoxy group;

b) a compound of formula II:




Formula II

or a pharmaceutically acceptable salt thereof, wherein:

X is H, or OH; and

R is substituted phenyl, or substituted or unsubstituted biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl or furan-3-yl; wherein, when substituted, R is substituted with 1 to 3 moieties (R', R'', R''') independently selected from a C1-C10 alkyl group, a C1-C10 alkoxy group and a hydroxyl group,

c) a compound of formula III:



Formula III


or a pharmaceutically acceptable salt thereof, wherein:

$R_1$  is H or  $C_1$ - $C_{10}$  alkyl; and

$R_2$  is phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl, furan-3-yl or tetrahydrofuran-2-yl; wherein  $R_2$  is optionally substituted with 1 to 3 moieties ( $R'$ ,  $R''$ ,  $R'''$ ) independently selected from the group consisting of a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group or a hydroxyl group,

and

d) a compound of formula IV:



Formula IV

wherein:

$\equiv$  represents a double or single bond;

$R_1$  is H or  $C_1$ - $C_{10}$  alkyl;

$R_4$  is H, OH or O; and

$R_5$  is a substituted or unsubstituted: phenyl, biphenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, furan-2-yl, furan-3-yl or tetrahydrofuran-2-yl, and when substituted,  $R_5$  is substituted with 1 to 3 moieties ( $R'$ ,  $R''$ ,  $R'''$ ) independently selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group and a hydroxyl group;

with the following provisos:

proviso 1: when  $R_1$  and  $R_4$  are H;  $R_5$  cannot be 1'-(2,4-dimethoxy-3-methyl)phenyl;

proviso 2: when  $R_1$  and  $R_4$  are H;  $R_5$  cannot be 1'-(2,4-dihydroxy)phenyl; 1'-(2,4-dialkoxy)phenyl; 1'-(2,4-dialkenoxy)phenyl or 1'-(2,4-dibenzyloxy)phenyl;

proviso 3: R<sub>5</sub> cannot be 1'-(4-hydroxy)phenyl;

proviso 5: when R<sub>1</sub> and R<sub>4</sub> are H and R<sub>5</sub> cannot be a 1'-(2,4-disubstituted)phenyl; wherein said substituents are selected from -OR<sup>2</sup> and -OR<sup>3</sup>, wherein R<sup>2</sup> and R<sup>3</sup> are independently selected from H and a (C<sub>1</sub>-C<sub>10</sub>)-alkyl; and

proviso 6: when R<sub>1</sub> is H and R<sub>4</sub> is OH; R<sub>5</sub> cannot be 1'-(3,4-dihydroxy)phenyl; 1'-(3,4-dimethoxy)phenyl; 1'-(2-hydroxy)phenyl; 1'-(2-methoxy)phenyl; 1'-(4-methoxy)phenyl; phenyl; or 1'-(3-methoxy,4-hydroxy)phenyl, for inhibiting browning and color changes in fruits, vegetables, juices and food products.

33. The use of claim 32, wherein the compound is selected from the group consisting of 4'-methoxybiphenyl-2,4-diol (1), biphenyl-2,2',4,4'-tetraol (2), 4-(pyridin-3-yl)benzene-1,3-diol (3), 4-(4'-hydroxybiphenyl)benzene-1,3-diol (4), 4-(furan-2-ylmethyl)benzene-1,3-diol (5), 4-(furan-3-ylmethyl)benzene-1,3-diol (6), 4-(pyridin-ylmethyl)benzene-1,3-diol (7), 4-((2-methoxypyridin-3-yl)methyl)benzene-1,3-diol (8), 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9), 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11), 4-phenethylbenzene-1,3-diol (12), 4-(4-hydroxyphenethylbenzene)-1,3-diol (13), 4-(4-methoxyphenethyl)benzene-1,3-diol (14), 4,4'-(ethane-1,2-diyl)dibenzene-1,3-diol (16), 4-(2,4-dimethoxyphenethyl)benzene-1,3-diol (17), 4-(3,5-dimethoxyphenethyl)benzene-1,3-diol (18), 4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol (19), 4-(2,4,6-trimethoxyphenethyl)benzene-1,3-diol (20), 4-(1-phenylpropan-2-yl)benzene-1,3-diol (21), 4-(2-(furan-3-yl)ethyl)benzene-1,3-diol (22), 4-(2-(tetrahydrofuran-3-yl)ethyl)benzene-1,3-diol (23), 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol (24), 4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol (25), 4-(2-(5-ethylfuran-2-yl)ethyl)benzene-1,3-diol (26), 4-(2-(pyridin-3-yl)ethyl)benzene-1,3-diol (27), 4-(3-(3,5-dimethoxyphenyl)propyl)benzene-1,3-diol (28), 4-(3-(3,5-dimethoxyphenyl)-1-hydroxypropyl)benzene-1,3-diol (29), 4-(3-(3,5-dihydroxyphenyl)propyl)benzene-1,3-diol (30), 4-(3-(2,4,6-trimethoxyphenyl)propyl)benzene-1,3-diol (31), 4-(3-(2,4-dihydroxyphenyl)propyl)-2-

methylbenzene-1,3-diol (32), 4-(3-(2,4-dihydroxy-3-methylphenyl)-3-(2,4-dihydroxyphenyl)propane-1-one (33) and 4-(5-hydroxypentyl)benzene-1,3-diol (34).

34. A composition for inhibiting the activity of tyrosinase, the composition comprising one or more compounds selected from the group consisting of 4'-methoxybiphenyl-2,4-diol (1), 4-(pyridin-3yl)benzene-1,3-diol (3), 4-(4'-hydroxybiphenyl)benzene-1,3-diol (4), 4-(furan-2-ylmethyl)benzene-1,3-diol (5), 4-(furan-3-ylmethyl)benzene-1,3-diol (6), 4-(pyridin-ylmethyl)benzene-1,3-diol (7), 4-((2-methoxypyridin-3-yl)methyl)benzene-1,3-diol (8), 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9), 4-(benzo[d][1,3]dioxol-5-ylmethyl)benzene-1,3-diol (10), 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11), 4-phenethylbenzene-1,3-diol (12), 4-(4-hydroxyphenethylbenzene)-1,3-diol (13), 4-(4-methoxyphenethyl)benzene-1,3-diol (14), 4-(2-(benzo[d][1,3]dioxol-5-ylethyl)benzene-1,3-diol (15), 4-(2,4-dimethoxyphenethyl)benzene-1,3-diol (17), 4-(3,5-dimethoxyphenethyl)benzene-1,3-diol (18), 4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol (19), 4-(2,4,6-trimethoxyphenethyl)benzene-1,3-diol (20), 4-(1-phenylpropan-2-yl)benzene-1,3-diol (21), 4-(2-(furan-3-yl)ethyl)benzene-1,3-diol (22), 4-(2-(tetrahydrofuran-3-yl)ethyl)benzene-1,3-diol (23), 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol (24), 4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol (25), 4-(2-(5-ethylofuran-2-yl)ethyl)benzene-1,3-diol (26), 4-(2-(pyridin-3yl)ethyl)benzene-1,3-diol (27), 4-(3-(3,5-dimethoxyphenyl)propyl)benzene-1,3-diol (28), 4-(3-(3,5-dimethoxyphenyl)-1-hydroxypropyl)benzene-1,3-diol (29), 4-(3-(3,5-dihydroxyphenyl)propyl)benzene-1,3-diol (30), 4-(3-(2,4,6-trimethoxyphenyl)propyl)benzene-1,3-diol (31), 4-(3-(2,4-dihydroxyphenyl)propyl)-2-methylbenzene-1,3-diol (32), 4-(3-(2,4-dihydroxy-3-methylphenyl)-3-(2,4-dihydroxyphenyl)propane-1-one (33), and 4-(5-hydroxypentyl)benzene-1,3-diol (34) and a pharmaceutically acceptable salt thereof, and a pharmaceutically or dermatologically acceptable carrier.

35. A composition for inhibiting the production of melanin, the composition comprising one or more compounds selected from 4'-methoxybiphenyl-2,4-diol (1), 4-(pyridin-3yl)benzene-1,3-diol (3), 4-(4'-hydroxybiphenyl)benzene-1,3-diol (4), 4-(furan-

2-ylmethyl)benzene-1,3-diol (5), 4-(furan-3-ylmethyl)benzene-1,3-diol (6), 4-(pyridin-ylmethyl)benzene-1,3-diol (7), 4-((2-methoxypyridin-3-yl)methyl)benzene-1,3-diol (8), 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9), 4-(benzo[d][1,3]dioxol-5-ylmethyl)benzene-1,3-diol (10), 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11), 4-phenethylbenzene-1,3-diol (12), 4-(4-hydroxyphenethylbenzene)-1,3-diol (13), 4-(4-methoxyphenethyl)benzene-1,3-diol (14), 4-(2-(benzo[d][1,3]dioxol-5-ylethyl)benzene-1,3-diol (15), 4-(2,4-dimethoxyphenethyl)benzene-1,3-diol (17), 4-(3,5-dimethoxyphenethyl)benzene-1,3-diol (18), 4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol (19), 4-(2,4,6-trimethoxyphenethyl)benzene-1,3-diol (20), 4-(1-phenylpropan-2-yl)benzene-1,3-diol (21), 4-(2-(furan-3-yl)ethyl)benzene-1,3-diol (22), 4-(2-(tetrahydrofuran-3-yl)ethyl)benzene-1,3-diol (23), 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol (24), 4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol (25), 4-(2-(5-ethylfuran-2-yl)ethyl)benzene-1,3-diol (26), 4-(2-(pyridin-3yl)ethyl)benzene-1,3-diol (27), 4-(3-(3,5-dimethoxyphenyl)propyl)benzene-1,3-diol (28), 4-(3-(3,5-dimethoxyphenyl)-1-hydroxypropyl)benzene-1,3-diol (29), 4-(3-(3,5-dihydroxyphenyl)propyl)benzene-1,3-diol (30), 4-(3-(2,4,6-trimethoxyphenyl)propyl)benzene-1,3-diol (31), 4-(3-(2,4-dihydroxyphenyl)propyl)-2-methylbenzene-1,3-diol (32), 4-(3-(2,4-dihydroxy-3-methylphenyl)-3-(2,4-dihydroxyphenyl)propane-1-one (33), and 4-(5-hydroxypentyl)benzene-1,3-diol (34) and a pharmaceutically acceptable salt thereof, and a pharmaceutically or dermatologically acceptable carrier.

36. A composition for preventing or treating diseases or conditions related to the overproduction or uneven distribution of melanin, the composition comprising one or more compounds selected from 4'-methoxybiphenyl-2,4-diol (1), 4-(pyridin-3yl)benzene-1,3-diol (3), 4-(4'-hydroxybiphenyl)benzene-1,3-diol (4), 4-(furan-2-ylmethyl)benzene-1,3-diol (5), 4-(furan-3-ylmethyl)benzene-1,3-diol (6), 4-(pyridin-ylmethyl)benzene-1,3-diol (7), 4-((2-methoxypyridin-3-yl)methyl)benzene-1,3-diol (8), 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9), 4-(benzo[d][1,3]dioxol-5-ylmethyl)benzene-1,3-diol (10), 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11), 4-phenethylbenzene-1,3-diol (12), 4-(4-hydroxyphenethylbenzene)-1,3-diol (13), 4-(4-

methoxyphenethyl)benzene-1,3-diol (14), 4-(2-(benzo[d][1,3]dioxol-5-ylethyl)benzene-1,3-diol (15), 4-(2,4-dimethoxyphenethyl)benzene-1,3-diol (17), 4-(3,5-dimethoxyphenethyl)benzene-1,3-diol (18), 4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol (19), 4-(2,4,6-trimethoxyphenethyl)benzene-1,3-diol (20), 4-(1-phenylpropan-2-yl)benzene-1,3-diol (21), 4-(2-(furan-3-yl)ethyl)benzene-1,3-diol (22), 4-(2-(tetrahydrofuran-3-yl)ethyl)benzene-1,3-diol (23), 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol (24), 4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol (25), 4-(2-(5-ethylfuran-2-yl)ethyl)benzene-1,3-diol (26), 4-(2-(pyridin-3yl)ethyl)benzene-1,3-diol (27), 4-(3-(3,5-dimethoxyphenyl)propyl)benzene-1,3-diol (28), 4-(3-(3,5-dimethoxyphenyl)-1-hydroxypropyl)benzene-1,3-diol (29), 4-(3-(3,5-dihydroxyphenyl)propyl)benzene-1,3-diol (30), 4-(3-(2,4,6-trimethoxyphenyl)propyl)benzene-1,3-diol (31), 4-(3-(2,4-dihydroxyphenyl)propyl)-2-methylbenzene-1,3-diol (32), 4-(3-(2,4-dihydroxy-3-methylphenyl)-3-(2,4-dihydroxyphenyl)propane-1-one (33), and 4-(5-hydroxypentyl)benzene-1,3-diol (34) and a pharmaceutically acceptable salt thereof, and a pharmaceutically or dermatologically acceptable carrier.

37. The composition of claim 36 wherein the diseases or conditions are selected from the group consisting of suntan, hyper pigmentation spots caused by skin aging, melasma, liver diseases, thermal burns and topical wounds, skin pigmentation due to inflammatory conditions caused by fungal, microbial and viral infections, vitilago, carcinoma, and melanoma.



38. A composition for whitening and/or lightening skin, the composition comprising one or more compounds selected from the group consisting of 4'-methoxybiphenyl-2,4-diol (1), 4-(pyridin-3yl)benzene-1,3-diol (3), 4-(4'-hydroxybiphenyl)benzene-1,3-diol (4), 4-(furan-2-ylmethyl)benzene-1,3-diol (5), 4-(furan-3-ylmethyl)benzene-1,3-diol (6), 4-(pyridin-ylmethyl)benzene-1,3-diol (7), 4-((2-methoxypyridin-3-yl)methyl)benzene-1,3-diol (8), 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9), 4-(benzo[d][1,3]dioxol-5-ylmethyl)benzene-1,3-diol (10), 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11), 4-phenethylbenzene-1,3-diol (12),

4-(4-hydroxyphenethylbenzene)-1,3-diol (13), 4-(4-methoxyphenethyl)benzene-1,3-diol (14), 4-(2-(benzo[d][1,3]dioxol-5-ylethyl)benzene-1,3-diol (15), 4-(2,4-dimethoxyphenethyl)benzene-1,3-diol (17), 4-(3,5-dimethoxyphenethyl)benzene-1,3-diol (18), 4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol (19), 4-(2,4,6-trimethoxyphenethyl)benzene-1,3-diol (20), 4-(1-phenylpropan-2-yl)benzene-1,3-diol (21), 4-(2-(furan-3-yl)ethyl)benzene-1,3-diol (22), 4-(2-(tetrahydrofuran-3-yl)ethyl)benzene-1,3-diol (23), 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol (24), 4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol (25), 4-(2-(5-ethylfuran-2-yl)ethyl)benzene-1,3-diol (26), 4-(2-(pyridin-3yl)ethyl)benzene-1,3-diol (27), 4-(3-(3,5-dimethoxyphenyl)propyl)benzene-1,3-diol (28), 4-(3-(3,5-dimethoxyphenyl)-1-hydroxypropyl)benzene-1,3-diol (29), 4-(3-(3,5-dihydroxyphenyl)propyl)benzene-1,3-diol (30), 4-(3-(2,4,6-trimethoxyphenyl)propyl)benzene-1,3-diol (31), 4-(3-(2,4-dihydroxyphenyl)propyl)-2-methylbenzene-1,3-diol (32), 4-(3-(2,4-dihydroxy-3-methylphenyl)-3-(2,4-dihydroxyphenyl)propane-1-one (33), and 4-(5-hydroxypentyl)benzene-1,3-diol (34) and a pharmaceutically acceptable salt thereof, and a pharmaceutically or dermatologically acceptable carrier.



39. A method for inhibiting browning and color changes in fruits or vegetables, juices, the method comprising contacting the fruits, vegetables, juices or other food products with a compound selected from the group consisting of 4'-methoxybiphenyl-2,4-diol (1), 4-(pyridin-3yl)benzene-1,3-diol (3), 4-(4'-hydroxybiphenyl)benzene-1,3-diol (4), 4-(furan-2-ylmethyl)benzene-1,3-diol (5), 4-(furan-3-ylmethyl)benzene-1,3-diol (6), 4-(pyridin-ylmethyl)benzene-1,3-diol (7), 4-((2-methoxypyridin-3-yl)methyl)benzene-1,3-diol (8), 4-(2,4,6-trimethoxybenzyl)benzene-1,3-diol (9), 4-(benzo[d][1,3]dioxol-5-ylmethyl)benzene-1,3-diol (10), 4-(hydroxyl(pyridine-3-yl)methyl)benzene-1,3-diol (11), 4-phenethylbenzene-1,3-diol (12), 4-(4-hydroxyphenethylbenzene)-1,3-diol (13), 4-(4-methoxyphenethyl)benzene-1,3-diol (14), 4-(2-(benzo[d][1,3]dioxol-5-ylethyl)benzene-1,3-diol (15), , 4-(2,4-dimethoxyphenethyl)benzene-1,3-diol (17), 4-(3,5-dimethoxyphenethyl)benzene-1,3-diol (18), 4-(2,4-dimethoxy-3-methylphenethyl)benzene-1,3-diol (19), 4-(2,4,6-trimethoxyphenethyl)benzene-1,3-diol (20), 4-(1-phenylpropan-2-yl)benzene-1,3-diol

(21), 4-(2-(furan-3-yl)ethyl)benzene-1,3-diol (22), 4-(2-(tetrahydrofuran-3-yl)ethyl)benzene-1,3-diol (23), 4-(2-(furan-2-yl)ethyl)benzene-1,3-diol (24), 4-(2-(tetrahydrofuran-2-yl)ethyl)benzene-1,3-diol (25), 4-(2-(5-ethylfuran-2-yl)ethyl)benzene-1,3-diol (26), 4-(2-(pyridin-3yl)ethyl)benzene-1,3-diol (27), 4-(3-(3,5-dimethoxyphenyl)propyl)benzene-1,3-diol (28), 4-(3-(3,5-dimethoxyphenyl)-1-hydroxypropyl)benzene-1,3-diol (29), 4-(3-(3,5-dihydroxyphenyl)propyl)benzene-1,3-diol (30), 4-(3-(2,4,6-trimethoxyphenyl)propyl)benzene-1,3-diol (31), 4-(3-(2,4-dihydroxyphenyl)propyl)-2-methylbenzene-1,3-diol (32), 4-(3-(2,4-dihydroxy-3-methylphenyl)-3-(2,4-dihydroxyphenyl)propane-1-one (33), and 4-(5-hydroxypentyl)benzene-1,3-diol (34).



1/7

**Fig. 1****Fig. 2**



2/7

**Fig. 3****Fig. 4**


3/7

**Fig. 5****Fig. 6**


4/7

**Fig. 7****Fig. 8**



5/7


**Fig. 9**

6/7

**Fig. 10**

7/7

**Fig. 11A****Fig. 11B**



Formula I