

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(10) International Publication Number

WO 2018/231937 A2

(43) International Publication Date
20 December 2018 (20.12.2018)

(51) International Patent Classification:

Not classified

(21) International Application Number:

PCT/US2018/037263

(22) International Filing Date:

13 June 2018 (13.06.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/519,087 13 June 2017 (13.06.2017) US

(72) Inventor; and

(71) Applicant: HSIA, Houn Simon [US/US]; 17201 Daimler Street, Irvine, California 92614 (US).

(74) Agent: BRILLHART, Kurt L et al.; FISH IP LAW, LLC, 2603 Main Street, Ste 1000, Irvine, California 92614 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: COMPOSITIONS AND METHODS FOR ENHANCING CANCER RADIOTHERAPY




FIG. 3A

(57) **Abstract:** A nutritional supplement containing fish oil and selenium has been identified that potentiates radiotherapy of tumors in a synergistic fashion. Cotherapy with such a supplement and radiotherapy results in substantially improved reductions in tumor volume of animal models relative to radiotherapy alone, and reduces common side effects such as weight loss, inflammation, and damage to the gastrointestinal tract. Such cotherapy was also found to reduce the incidence of metastasis from the primary implantation site, and to modulate expression of genes associated with apoptosis in tumor cells. A formulation for such a supplement that is both well tolerated and palatable is also provided.

COMPOSITIONS AND METHODS FOR ENHANCING CANCER RADIOTHERAPY

[0001] This application claims the benefit of United States Provisional Application No. 62/519,087 filed on June 13, 2017. These and all other referenced extrinsic materials are incorporated herein by reference in their entirety. Where a definition or use of a term in a reference that is incorporated by reference is inconsistent or contrary to the definition of that term provided herein, the definition of that term provided herein is deemed to be controlling.

Field of the Invention

[0002] The field of the invention is cancer radiotherapy.

Background

[0003] The background description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed invention, or that any publication specifically or implicitly referenced is prior art.

[0004] Radiotherapy and chemotherapy protocols utilized in the treatment of cancer can clearly benefit patients, but can be ineffective or less effective with some cancers. In addition both radiotherapy and chemotherapy are associated with significant side effects, including nausea, weight loss, hair loss, damage to the gastrointestinal tract, and skin irritation.

[0005] Attempts have been made to enhance the effectiveness of radiotherapy. For example, gold nanoparticles that have been modified to target tumor cells have been used to enhance radiotherapy (Yang et al, ACS Nano, 2014, 8(9):8992-9002). All publications herein are incorporated by reference to the same extent as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Where a definition or use of a term in an incorporated reference is inconsistent or contrary to the definition of that term provided herein, the definition of that term provided herein applies and the definition of that term in the reference does not apply. Similarly, COX-2 inhibitors have been used to selectively sensitize tumor cells to the effects of radiation (Choy and Milas, J. Natl Cancer Inst (2003) 95(19):1140-1452). Such approaches, however, can have issues with

selectivity and may not be effective against all tumor types. To date attempts to reduce the side effects of radiotherapy are primarily directed to partitioning the total radiation dose into a number of smaller radiation doses (leaving time in between to allow for recovery), targeting of tumor using shielding, and identification of the boundaries of the tumor and localization of radiotherapy to that site. Unfortunately, such approaches can fail to adequately treat all of the tumor cells.

[0006] Attempts have also been made to enhance the effects of chemotherapy. Some studies have suggested that consumption of fish oil can improve results from chemotherapy, however other research has suggested that fish oil can interfere (Daenen et al, JAMA Oncol (2015) 1(3):350-358). Formulation of chemotherapeutic agents as nanoparticles has also been attempted (Xu et al, Coll. Surf. B: Biointerfaces (2006) 48(1):50-57). It is unclear, however, if all chemotherapeutic drugs are suitable for such reformulation. Codelivery of chemotherapeutic drugs with siRNA designed to interfere with multi-drug resistance has also been explored. Such siRNAs, however, are sequence specific and may not be suitable for some tumors.

[0007] Mitigation of the side effects of chemotherapy are generally directed at providing symptomatic relief. For example, antiemetics can be used to reduce nausea, along with diet modification and eating small, frequent meals that avoid certain foods. Unfortunately such approaches are not always effective. In some instances chemotherapeutic agents are selected to have reduced toxicity in order to reduce side effects, however such agents may also have reduced effectiveness against tumor cells.

[0008] Thus, there is still a need for safe and effective compositions and methods to enhance the effectiveness and/or reduce the side effects of cancer radiotherapy and/or chemotherapy.

Summary of The Invention

[0009] The inventive subject matter provides compositions and methods for enhancing and improving radiotherapy of tumors in a synergistic manner through the use of a nutritional supplement containing fish oil and selenium.

[0010] One embodiment of the inventive concept is a method of treating a tumor by applying a radiotherapy protocol to a patient in need of treatment while providing the patient with a

nutritional supplement containing fish oil and selenium (for example, as shown in Table 1) in an amount to provide a synergistic effect in reducing tumor volume or weight. In some embodiments the nutritional supplement is provided to the patient prior to the initiation of radiotherapy. Such a nutritional supplement can be formulated such that two or more of the components of the supplement are provided in amounts as described in Table 1.

[0011] Another embodiment of the inventive concept is a method of reducing side while providing the patient with a nutritional supplement formulated that includes fish oil and selenium (for example, as in Table 1) in an amount to reduce a side effect of the radiotherapy protocol. In some embodiments the nutritional supplement is provided to the patient prior to the initiation of radiotherapy. Such a nutritional supplement can be formulated such that two or more of the components of the supplement are provided in amounts as described in Table 1.

[0012] Another embodiment of the inventive concept is a method of modulating gene (*e.g.* an angiogenesis-related gene, an apoptosis-related gene, *etc.*) expression in a tumor, by providing a nutritional supplement to a tumor or an animal having a tumor, wherein the nutritional supplement includes fish oil and selenium (for example, as in Table 1), and where the nutritional supplement is provided in an amount to modulate expression of a gene of the tumor. In some embodiments the nutritional supplement is provided to the patient prior to the initiation of radiotherapy and during radiotherapy; in other embodiments such a supplement is provided during or at the initiation of the application of radiotherapy. In preferred embodiments the nutritional supplement is formulated such that two or more components of the supplement are provided in amounts as described in Table 1.

[0013] Another embodiment of the inventive concept is a method of reducing metastasis from a tumor, by providing the patient having a metastatic tumor with a nutritional supplement that includes fish oil and selenium (such as in Table 1) where the nutritional supplement is provided in an amount to reduce metastatic activity of the tumor. In some embodiments such a nutritional supplement is provided either prior to or at the initiation of radiotherapy, and can be provided through the course of radiotherapy. In a preferred embodiment the nutritional supplement is formulated such that two or more of the supplement's components are provided in amounts as described in Table 1.

[0014] Another embodiment of the inventive concept is a method of reducing angiogenesis in a tumor, comprising providing the patient having a tumor with a nutritional supplement that includes fish oil and selenium (for example, as in Table 1) in combination with radiotherapy, where the nutritional supplement is provided in an amount to reduce an angiogenic activity of the tumor. In some embodiments such a nutritional supplement is provided concurrent with the initiation of radiotherapy, and can be provided to the patient during the application of radiotherapy. In preferred embodiments the nutritional supplement is formulated such that two or more of the supplement's components are provided in amounts as described in Table 1.

[0015] Another embodiment of the inventive concept is a method of reducing incidences of cancer stem cells in a tumor, by providing the patient having a tumor with a nutritional supplement that includes fish oil and selenium (such as in Table 1), wherein the nutritional supplement is provided in an amount to reduce the incidence of stem cells in the tumor. In a preferred embodiment the nutritional supplement is provided concurrently with the application radiotherapy. The nutritional supplement can be formulated such that two or more components of the supplement are provided in amounts as described in Table 1.

[0016] Various objects, features, aspects and advantages of the inventive subject matter will become more apparent from the following detailed description of preferred embodiments, along with the accompanying drawing figures in which like numerals represent like components.

Brief Description of The Drawings

[0017] FIG. 1: FIG. 1 depicts a treatment protocol in which nutritional supplementation is provided starting either 7 days prior to implementation of radiotherapy or simultaneously with tumor cell implantation. Mice were sacrificed 21 days after tumor cell implantation.

[0018] FIGs. 2A and 2B: FIG. 2A depicts changes in body weight over time for various treatment groups. FIG. 2B depicts change in body weight in various treatment groups over the course of the study. Note that body weight is reported following removal of the primary tumor mass.

[0019] FIGs. 3A to 3D: FIGs. 3A to 3D show typical synergistic effects on tumor volume and weight of co-treatment with radiotherapy and the supplement containing fish oil and selenium.

FIG. 3A shows changes in tumor volume over time for various treatment groups. FIG. 3B provides a detailed view of the first 11 days of treatment as shown in FIG. 3A. FIG. 3C provides a histogram of tumor weight for various treatment groups. FIG. 3D provides photographs of exemplary tumors from various treatment groups.

[0020] FIGs. 4A to 4G: FIG. 4A: Effect of treatment using radiotherapy, a supplement containing fish oil and selenium, and combined treatment on gastrocnemius muscle mass 21 days after tumor cell injection. FIG. 4B: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on soleus muscle mass 21 days after tumor cell injection. FIG. 4C: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on lung (inclusive of metastatic tumor) weight 21 days after tumor cell injection. FIG. 4D: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on liver (inclusive of metastatic tumor) weight 21 days after tumor cell injection. FIG. 4E: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on spleen (inclusive of metastatic tumor) weight 21 days after tumor cell injection. FIG. 4F provides quantitative results of studies of lung metastasis in animal subjects treated with a supplement containing fish oil and selenium and/or radiotherapy. FIG. 4G shows the results of studies of Ki-67 expression in primary tumor sites and metastatic sites in animal subjects treated with a supplement containing fish oil and selenium and/or radiotherapy.

[0021] FIGs. 5A to 5E: FIG. 5A: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on platelet count 21 days after tumor cell injection. FIG. 5B: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on white blood cell count 21 days after tumor cell injection. FIG. 5C: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on lymphocyte count 21 days after tumor cell injection. FIG. 5D: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on granulocyte count 21 days after tumor cell injection. FIG. 5E: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on neutrophil/lymphocyte percentage ratio 21 days after tumor cell injection.

[0022] FIGs. 6A and 6B: FIG. 6A: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on serum albumin 21 days after tumor cell injection. FIG.

6B: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on serum creatinine 21 days after tumor cell injection.

[0023] FIGS. 7A and 7B: FIG. 7A: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on serum IL-6 21 days after tumor cell injection. FIG. 7B: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on serum IL-1 β 21 days after tumor cell injection.

[0024] FIGs. 8A to 8G: FIG. 8A: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the VEGF gene in an implanted tumor 21 days after tumor cell injection. FIG. 8B: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the BAX gene in an implanted tumor 21 days after tumor cell injection. FIG. 8C: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the Bcl-2 gene in an implanted tumor 21 days after tumor cell injection. FIG. 8D: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the caspase 3 gene in an implanted tumor 21 days after tumor cell injection. FIG. 8E: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the BAX gene in lung 21 days after tumor cell injection. FIG. 8F: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the Bcl-2 gene in lung 21 days after tumor cell injection. FIG. 8G: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the caspase 3 gene in lung 21 days after tumor cell injection.

[0025] FIGs. 9A and 9B: FIG. 9A depicts a testing protocol in which mice receive radiotherapy on days 8, 10, and 12, with nutritional supplementation beginning with the initiation of radiotherapy. Mice are sacrificed at day 14 or at day 24 following implantation of tumor cells. FIG. 9B: Treatment groups derived from the protocol shown in FIG. 9A.

[0026] FIGs. 10A to 10C: FIG. 10A: Serum albumin concentration at 14 days and 24 days following tumor cell implantation in mice treated as shown in FIG. 9A. FIG. 10B: Lymphocyte counts at 14 days and 24 days following tumor cell implantation in mice treated as shown in FIG. 9A. FIG. 10C: N/L Ratio at 14 days and 24 days following tumor cell implantation in mice treated as shown in FIG. 9A.

[0027] FIG. 11A to 11D: FIG. 11A: VEGF expression within the tumor mass 24 days following tumor cell implantation in mice treated as shown in FIG. 9A. FITC represents VEGF-specific staining. FIG. 11B: VEGF expression within the lung (metastasis) 24 days following tumor cell implantation in mice treated as shown in FIG. 9A. FITC represents VEGF-specific staining. FIG. 11C: EGFR expression within the lung (metastasis) 24 days following tumor cell implantation in mice treated as shown in FIG. 9A. FITC represents EGFR-specific staining. FIG. 11D: EGFR expression within the tumor mass 24 days following tumor cell implantation in mice treated as shown in FIG. 9A. FITC represents EGFR-specific staining.

[0028] FIG. 12: CD31 (cancer stem cell marker) expression within the tumor mass and in lung tissue (metastasis) 24 days following tumor cell implantation in mice treated as shown in FIG. 9A. FITC represents CD31-specific staining.

[0029] FIGs. 13A and 13B: FIG. 13AA shows the presence of H1F1- α (hypoxia marker) protein within the tumor mass and in lung tissue (metastasis) 24 days following tumor cell implantation in mice treated as shown in FIG. 9A. FITC represents H1F1- α -specific staining. FIG. 13B shows results of gene expression studies of tumor samples from subjects treated with a nutritional supplement containing fish oil and selenium and/or radiotherapy.

[0030] FIG. 14: Expression of apoptosis markers 24 days following tumor cell implantation in mice treated as shown in FIG. 9A.

[0031] FIGs. 15A and 15B: FIG. 15A shows the results of studies characterizing PDL-1 gene expression in primary and metastatic tumor sites in animal models of human disease treated with a supplement containing fish oil and selenium and/or radiotherapy. FIG. 15B shows the results of studies characterizing PD-1 gene expression in primary and metastatic tumor sites in animal models of human disease treated with a supplement containing fish oil and selenium and/or radiotherapy.

[0032] FIGs. 16A and 16B: FIG. 16A depicts a testing protocol in which mice receive radiotherapy on days 8, 10, and 12, with nutritional supplementation provided prior to tumor cell implantation, at the time of tumor cell implantation, or at the initiation of radiotherapy. Mice are

sacrificed at day 21 following implantation of tumor cells. FIG. 16B: Treatment groups derived from the protocol shown in FIG. 16A.

[0033] FIG. 17: Photomicrographs of cross sections of the intestines of mice treated by the protocol described in FIG. 16A.

[0034] FIGs. 18A to 18F: FIG. 18A: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the VEGF gene 21 days after tumor cell injection. FIG. 18B: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the BAX gene 21 days after tumor cell injection. FIG. 18C: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the Bcl-2 gene 21 days after tumor cell injection. FIG. 18D: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the Caspase 3 gene 21 days after tumor cell injection. FIG. 18E: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the Bcl-2 gene 21 days in lung (i.e. metastasis) after tumor cell injection. FIG. 18F: Effect of treatment using radiotherapy, NutraWell supplement, and combined treatment on expression of the Caspase 3 gene 21 days in lung (i.e. metastasis) after tumor cell injection.

[0035] FIGs. 19A and 19B: FIG. 19A depicts a testing protocol in which mice receive radiotherapy on days 8, 10, and 12, with nutritional supplementation provided for 7 days prior to tumor cell implantation. Mice are sacrificed at day 24 following implantation of tumor cells. FIG. 19B: Treatment groups derived from the protocol shown in FIG. 19A.

[0036] FIGs. 20A and 20B: FIG. 20A: Effect of a nutritional supplement containing fish oil and selenium on body mass in mice receiving repeated radiotherapy following tumor cell implantation using the protocol shown in FIG. 19A. FIG. 20B: Effect of treatment using radiotherapy, a nutritional supplement containing fish oil and selenium, and combined treatment on gastrocnemius muscle mass 21 days after tumor cell injection using the protocol shown in FIG. 19A.

[0037] FIGs. 21A and 21B: FIG. 21A depicts a testing protocol mice in which mice receive radiotherapy on days 8, 10, and 12, with nutritional supplementation provided for 7 days prior to,

the day of, or 8 days after tumor cell implantation. Mice are sacrificed at day 21 following implantation of tumor cells. FIG. 21B: Treatment groups derived from the protocol shown in FIG. 21A.

[0038] FIG. 22: Effect of NutraWell supplementation on body mass in mice receiving repeated radiotherapy following tumor cell implantation using the protocol shown in FIG. 21A.

[0039] FIG. 23: Changes in tumor volume over time on repeated radiotherapy in combination with NutraWell supplement in mice treated as in the protocol shown in FIG. 21A.

[0040] FIGs. 24A and 24B: FIG. 24A: Effect of treatment using repeated radiotherapy in combination with NutraWell supplement, and combined treatment on serum TNF- α 21 days after tumor cell injection in mice treated using the protocol shown in FIG. 21A. FIG. 24B: Effect of treatment using repeated radiotherapy in combination with NutraWell supplement, and combined treatment on serum IL-6 21 days after tumor cell injection in mice treated using the protocol shown in FIG. 21A.

Detailed Description

[0041] The following description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed invention, or that any publication specifically or implicitly referenced is prior art.

[0042] The inventive subject matter provides compositions and methods in which a nutritional supplement that includes fish oil and selenium (for example, a supplement containing fish oil, selenium derived from selenium yeast, and certain vitamins, minerals, amino acids, and saccharides, e.g. “NutraWell”) is used in combination with radiotherapy. Combination therapy with radiation and such a supplement surprisingly provides a significant synergistic effect in reduction of tumor size. In addition, side effects of radiotherapy (e.g. neutropenia, loss of body mass, loss of muscle mass, damage to acyl cells of the gastrointestinal tract, etc.) are reduced and/or mitigated relative to application of radiotherapy without use of such a supplement. Surprisingly, expression of genes related to angiogenesis and apoptosis were also found to be modulated in tumor cells on use of a supplement containing fish oil and selenium, both with and

without the application of radiotherapy. In addition, metastasis is prevented and the growth and spread of cancer stem cells was found to be reduced.

[0043] One should appreciate that the disclosed techniques provide many advantageous technical effects including enhancing the effectiveness of current radiotherapeutic protocols used in the treatment of cancer while reducing the side effects associated with these approaches.

[0044] The following discussion provides many example embodiments of the inventive subject matter. Although each embodiment represents a single combination of inventive elements, the inventive subject matter is considered to include all possible combinations of the disclosed elements. Thus if one embodiment comprises elements A, B, and C, and a second embodiment comprises elements B and D, then the inventive subject matter is also considered to include other remaining combinations of A, B, C, or D, even if not explicitly disclosed.

[0045] In some embodiments, the numbers expressing quantities of ingredients, properties such as concentration, reaction conditions, and so forth, used to describe and claim certain embodiments of the invention are to be understood as being modified in some instances by the term “about.” Accordingly, in some embodiments, the numerical parameters set forth in the written description and attached claims are approximations that can vary depending upon the desired properties sought to be obtained by a particular embodiment. In some embodiments, the numerical parameters should be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of some embodiments of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as practicable. The numerical values presented in some embodiments of the invention may contain certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

[0046] In one embodiment of the inventive concept, a nutritional supplement composition as shown in Table 1 (“Nutrawell”) is provided to improve the results of radiotherapy.

Component	Minimum	Maximum	Unit
Maltodextrin	10000	50000	mg
Whey Protein Isolate	5000	60000	mg

Whey Protein Concentrate	1000	50000	mg
Fructooligosaccharides/Inulin	40	15000	mg
Granulated Honey	1000	9000	mg
Oat Fiber	500	15000	mg
Natural French Vanilla Flavor	500	20000	mg
Soy Protein	500	50000	mg
Brownulated Powdered Brown Sugar	500	10000	mg
Natural Vanilla Masking Flavor	500	5000	mg
Lecithin	200	10000	mg
Milk, Non-fat	50	5000	mg
Rice Protein Powder	50	5000	mg
Calcium Caseinate	50	2000	mg
Oils			
Flax Seed Oil	100	7000	mg
Canola Oil	100	7000	mg
Borage Oil	100	7000	mg
Olive Oil	100	7000	mg
Fish Oil	150	10,000	mg
Pure Lemon Oil	100	1000	mg
Pure Orange Oil	50	1000	mg
Mixed Tocopherols	0.5	200	mg
Vitamins/Minerals			
Potassium Phosphate	200	1500	mg
Calcium Carbonate	100	5000	mg
Choline Bitartrate	150	2500	mg
Sodium Chloride	100	2000	mg
Calcium Phosphate Tribasic	100	2000	mg
Ascorbic Acid	50	3000	mg
Potassium Chloride	50	2000	mg
Magnesium Oxide	50	500	mg
Selenium Yeast	30	4000	mcg
Chromium Yeast	30	3000	mcg
Molybdenum Yeast	30	2000	mcg
Inositol	10	5000	mg
Zinc Sulfate Monohydrate	5	200	mg
Dry Vitamin E Acetate	5	2000	IU
Niacinamide	5	500	mg
Ferric Orthophosphate	3	100	mg
Calcium Pantothenate	3	200	mg
Manganese Sulfate Monohydrate	3	100	mg
Beta Carotene	1	100	mg

Copper Gluconate	1	15	mg
Vitamin D3	25	5000	IU
Vitamin K2	2	1000	mcg
Pyridoxine HCl	0.5	200	mg
Potassium Iodide	0.5	1500	mg
Riboflavin	0.5	1000	mg
Thiamine Hydrochloride	0.5	2500	mg
Dry Vitamin K1	1	500	mcg
Vitamin A Acetate	500	100000	IU
Folic Acid	100	10000	mcg
d-Biotin	10	10000	mcg
Vitamin B12	1	3000	mcg
Amino Acids			
L-Carnitine	300	30000	mg
L-Glutamine	500	60000	mg
L-Arginine Base	500	30000	mg
Taurine	50	2000	mg
L-Lysine	50	2000	mg
Alpha Lipoic Acid	10	1000	mg
Resveratrol	15	1500	mg
Co-Enzyme Q10	10	5000	mg
Glycine	5	1000	mg
Proline	5	1000	mg
Bacterial Cultures			
Lact. Acidophilus (app. 10 billion total)	2	500	mg
Bifido Bifidum (app. 10 billion total)	2	500	mg
Lac. Bulgaricus (app. 10 billion total)	2	500	mg
Bifido Longum (app. 10 billion total)	2	500	mg
Strep. Thermophilus (app. 10 billion total)	2	500	mg
Enzymes			
Papain	5	100	mg
Pepsin	5	100	mg
Lipase	5	100	mg
Bromelain	5	100	mg
Pancreatin 4X	0.5	100	mg
Lactase	1	100	mg
Betaine HCl	3	100	mg
Plant Products			
Pineapple Juice Powder	2	500	mg
Papaya Fruit Powder	2	500	mg
Quercetin	30	3000	mg

EGCG	25	600	mg
OPC	15	500	mg
Anthocyanins	15	5000	mg
Ellagic Acid	10	300	mg
Astaxanthin	2	90	mg
Fucoidan	20	1500	mg
Mushroom Preparation			
Cordyceps	5	6000	mg
Ganoderma Lucidum	15	10000	mg
Shiitake	40	15000	mg
Maitake	30	15000	mg
Turkey Tail	30	15000	mg

Table 1

[0047] The composition shown in Table 1 includes components that have various physiological and biochemical effects, including anti-inflammatory activity, lowering of blood glucose levels, lowering of cholesterol, and anti-tumor activity. Other components provide supplementation of necessary vitamins, minerals, and amino acids at elevated levels. Other components (e.g. enzymes, lecithin) serve to aid in digestion and absorption of components of the composition when consumed. The combination of these complementary activities provides a synergistic effect that exceeds the simple additive effect of individual components. It should be appreciated that the composition shown in Table 1 also includes certain flavorants (e.g. brown sugar, honey, vanilla flavor and masking agent) that serve to improve palatability and acceptance. Certain components (e.g. honey, brown sugar, milk, rice protein, casein) can provide both flavor and caloric energy. The Inventor has found that the combination of flavorants described above is effective in providing compliance with consumption of the nutritional supplement in effective amounts. In some embodiments, such flavorants can be excluded without negatively impacting the effectiveness of the nutritional supplement.

[0048] Components shown in Table 1 can be provided as a single formulation (for example, as a pill, tablet, capsule, powder, liquid, suspension, etc.) or can be segregated into different formulations (for example, as pills, tablets, capsules, powders, liquids, suspensions, or combinations thereof). The amounts shown in Table 1 are exemplary, and represent typical daily

dosages provided to an adult of normal stature and otherwise normal health. These amounts can be adjusted to account for differences in body mass, gender, medical condition, etc. For example, a relatively small patient weighing 40 kilos or less may receive benefit from dosages provided at or below the low end of the ranges provided, whereas a relatively large patient weighing 100 kilograms or more may require dosages provided at the high end of the ranges noted (or more). In some embodiments such a daily dose can be distributed as multiple doses throughout the day. In some of such embodiments the composition of each of such distributed doses can be identical. In other embodiments the composition of such distributed doses can be different, provided the summation of such doses provides the required supplementation.

[0049] In an exemplary embodiment human tumor cells (following transplantation into nude mice) are treated with 1 gram per day of the nutritional supplement, radiotherapy, or 1 gram per day of the nutritional supplement and chemotherapy. The mice are weighed during treatment to characterize side effects such as nausea and loss of appetite. After several weeks the mice are sacrificed and the tumor characterized. Tumor volume is determined, and the impact of therapy on organ and muscle volume is determined. The degree of neutropenia is also characterized. A typical protocol is shown in Table 2.

			Scarfify (21 th day)
1	C	Control	N=6
2	T	Tumor	N=6
3	PTN	Tumor + Nutrawell (-7 day start)	N=6
4	TN	Tumor + Nutrawell (0 day start)	N=6
5	TR	Tumor + Radiotherapy (3 Gy x 3)	N=6
6	PTRN	Tumor + Radiotherapy (3 Gy x 3) + Nutrawell (-7 day start)	N=6
7	TRN	Tumor + Radiotherapy (3 Gy x 3) + Nutrawell (8 day start)	N=6

Table 2

[0050] Body Weight/Wasting

[0051] A typical treatment schedule is depicted schematically in FIG. 1. Some subjects received treatment with a supplement containing fish oil and selenium (PTN, PTRN) prior to tumor cell implantation, with a portion of these (PTRN) receiving radiotherapy. Other subject started treatment with such a supplement at the time of tumor cell implantation (TN, TRN), with some of these receiving radiotherapy (TRN). Results of body weight studies compared to control subjects (C) and subjects implanted with tumor cells and otherwise untreated (T) are shown in FIGs. 2A and 2B. As shown, mice receiving both radiotherapy and the supplement containing fish oil and selenium gained weight at a significantly greater rate than those receiving only radiotherapy, indicating a reduction in side effects normally associated with this treatment mode. Pre-treatment with a nutritional supplement containing fish oil and selenium had a particularly pronounced effect.

[0052] Tumor Size

[0053] FIGs. 3A to 3D show typical synergistic effects on tumor volume and weight of co-treatment with radiotherapy and a supplement containing fish oil and selenium for similar treatment groups in a murine model of human cancer. FIG. 3A shows the effect of various treatment protocols on tumor volume over the course of 3 weeks, with FIG. 3B providing a scaled view of the effect over the initial 11 days. As shown, treatment with a supplement containing fish oil and selenium alone provides an approximately 60% reduction in tumor volume. Treatment with radiation alone provides a similar reduction in tumor volume. In the absence of a synergistic effect one would therefore anticipate a reduction in tumor volume to approximately 25% of that of the untreated tumor. Surprisingly, what is observed is a greater than 90% reduction in tumor volume to approximately 7% of the untreated tumor- indicative of a significant synergistic effect. FIG. 3C shows that similar effects are found when tumor weight is characterized. FIG. 3D shows photographs typical examples of tumors excised from test animals following treatment; the effects of cotherapy with a nutritional supplement containing fish oil and selenium with radiotherapy is readily apparent on visual inspection. As such, it is apparent that cotherapy with a supplement containing fish oil and selenium and radiotherapy can provide a synergistic effect in reducing tumor volume and/or mass.

[0054] Muscle Wasting

[0055] Side effects of radiotherapy go beyond loss of appetite and weight loss, and can include damage to internal organs, loss of muscle mass, anemia, neutropenia, reduction in kidney function, *etc.* To determine the protective effects of cotherapy with a supplement containing fish oil and selenium in regards to such side effects on muscle mass and organ weight of treated mice were also characterized following therapy. The results are shown in FIGs. 4A to 4E. FIGs. 4A and 4B show the effects of treatment with a supplement containing fish oil and selenium, radiotherapy, and cotherapy with radiotherapy using such a supplement (with and without supplement pre-treatment) on the weight of gastrocnemius muscle and soleus muscle (respectively), with test groups designated as described above. As shown, cotherapy with such a supplement and radiotherapy provides almost complete retention of muscle weight.

[0056] Metastasis

[0057] FIGs. 4C, 4D, and 4E show the effects of treatment as described above on the weight lungs inclusive of any metastatic tumors (FIG. 4C), livers inclusive of any metastatic tumors (FIG. 4D), and spleens inclusive of any metastatic tumors (FIG. 4E). As shown lack of treatment leads to an increase in weight, due at least in part to the presence of metastatic tumors. This increase is not entirely mitigated by radiotherapy alone (TR), however subjects receiving treatment with a supplement containing fish oil and selenium had organ weights similar to those of control subjects, both with and without cotherapy with radiotherapy.

[0058] Similarly, quantitation of metastatic sites within the lungs of animal models of human disease treated with a nutritional supplement containing fish oil and selenium and/or radiotherapy. Typical metastatic sites are shown in the photographs in the left panel of FIG. 4F, and enumerated in a histogram in the right panel. As shown, treatment with either such a supplement (TN) or radiotherapy (TR) reduced the number of metastatic sites to some extent, however treatment with both the supplement and radiotherapy (TRN) resulted in a complete lack of apparent metastatic sites in most subjects.

[0059] Without wishing to be bound by theory, the Inventor believes that this effect on metastasis (and also on tumor size and mass, as shown above) can be due to a synergistic effect provided by the use of a nutritional supplement containing fish oil and selenium in combination with radiotherapy on tumor cell proliferation. As shown in FIG. 4G, the expression of the

proliferation marker Ki-67 is suppressed in the primary tumor site on treatment with either a supplement containing fish oil and selenium (TN), on treatment with radiotherapy (TR), or treatment using both modes (TRN). Metastatic sites (right panel) show a related trend, with treatment with the nutritional supplement or radiotherapy alone resulting in a moderate reduction in the expression of this proliferation marker, but surprisingly showing a synergistic effect in reducing expression of Ki-67 in metastatic sites. This indicates that use of a supplement containing fish oil and selenium can reduce tumor metastasis, including metastasis that is not prevented by radiotherapy, and can do so in a synergistic manner.

[0060] Neutropenia/Anemia

[0061] Anemia and neutropenia are often found in cancer and as side effects of radiotherapy, due at least in part to suppression of bone marrow activity. The effects of cotherapy with a supplement containing fish oil and selenium on various blood cell populations are shown in FIGs. 5A to 5E. FIG. 5B shows the effects of a nutritional supplement containing fish oil and selenium, radiotherapy, and cotherapy with such a supplement and radiotherapy on white blood cells as shown in FIG. 5B, and lymphocytes as shown in FIG. 5C. This indicates that a nutritional supplement containing fish oil and selenium can be useful to improve red blood cell, granulocyte, white blood cell, and/or lymphocyte suppression resulting from the presence of a tumor and from radiotherapy utilized in the treatment of such tumors.

[0062] As shown in FIG. 5A, platelets were found to be elevated in untreated animals bearing tumors (T). Platelet concentrations were reduced by treatment with either a supplement containing fish oil and selenium or radiotherapy, but were most pronounced in animals receiving cotherapy with such a supplement and radiotherapy. This effect was particularly pronounced in subjects receiving pre-treatment with a supplement containing fish oil and selenium, which reduced platelet concentrations to the values found for control animals. Granulocyte concentrations were similarly elevated in untreated tumor bearing animals (as shown in FIG. 5D), and were further elevated in such animals receiving only radiotherapy. Cotherapy with a nutritional supplement containing fish oil and selenium and radiotherapy, was found to reduce this effect. As shown in FIG. 5E, tumor bearing animals show an elevated neutrophil/lymphocyte ratio (NLR) relative to control animals, which is even more elevated in

similar subjects receiving only radiotherapy. Treatment with a supplement containing fish oil and selenium was found to be effective in shifting this ratio towards a more normal value, both as a monotherapy and as a part of cotherapy with radiotherapy. This indicates that a nutritional supplement containing fish oil and selenium can be useful in reducing platelet concentration, granulocyte concentration, and/or neutrophil/lymphocyte ratios that are elevated due to the presence of a tumor and due to radiotherapy used to treat such a tumor.

[0063] Serum proteins

[0064] The effects of cotherapy with a supplement containing fish oil and selenium on various serum biochemistry markers, which provide information regarding kidney function, liver function, and/or nutritional status are shown in FIGs. 6A and 6B. FIG. 6A shows the effects of a nutritional supplement containing fish oil and selenium, radiotherapy, and cotherapy with such a supplement and radiotherapy on an animal tumor model. As shown, serum albumin concentration (an indicator of nutritional status) is reduced in untreated tumor-bearing animals, and is only slightly improved by radiotherapy alone. Treatment with a nutritional supplement containing fish oil and selenium, either as a monotherapy or as cotherapy with radiotherapy, improves serum albumin concentration- particularly when such a supplement is provided as a pretreatment. This is improved by cotherapy with a nutritional supplement containing fish oil and selenium. This indicates that treatment with a nutritional supplement containing fish oil and selenium, either as a monotherapy or as cotherapy with radiotherapy, can improve the nutritional status of tumor-bearing subjects.

[0065] FIG. 6B shows the results of similar studies where serum creatinine (a measure of kidney function) is characterized. As shown, untreated tumor bearing subjects shown elevated creatinine concentrations indicative of kidney damage. This is marginally improved by radiotherapy alone. Cotherapy with a nutritional supplement containing fish oil and selenium and radiotherapy, however, shows a synergistic effect in reducing serum creatinine concentrations- particularly when such a nutritional supplement is provided as a pretreatment.

[0066] Cytokines

[0067] The growth and spread of tumors is associated with inflammation, as is the application of radiotherapy. Surprisingly, Inventors have found that cotherapy with a supplement containing fish oil and selenium is effective in reducing the concentration of pro-inflammatory cytokines, indicating that such co-treatment is effective in reducing inflammation associated with tumors, and with radiotherapy of tumors. The effects of NutraWell supplement on serum concentrations of pro-inflammatory cytokines is shown in FIGs. 7A and 7B. FIG. 7A shows the concentration of IL-6 in control animals, untreated tumor-bearing animals, and tumor bearing animals treated with a nutritional supplement containing fish oil and selenium, radiotherapy, or such a supplement and radiotherapy in combination. As shown, untreated tumor-bearing animals show a large increase in serum IL-6 concentration. This is only somewhat reduced by radiotherapy. Use of a nutritional supplement containing fish oil and selenium, either as a monotherapy or as cotherapy with radiotherapy, was found to reduce serum IL-6 concentrations in tumor-bearing animals. FIG. 7B shows the results of similar studies of IL-1 β . The results with IL-1 β are similar to those found for IL-6. This indicates that a nutritional supplement containing fish oil and selenium can effectively reduce serum concentrations of inflammation-associated cytokines in tumor-bearing subjects, either as a monotherapy or as cotherapy with radiotherapy. The Inventor believes that such reduction is accompanied by a reduction in inflammation in such animals.

[0068] Tumor gene expression

[0069] Surprisingly, Inventors have also found that treatment with a nutritional supplement containing fish oil and selenium can modify gene expression in tumor cells *in vivo*, and can provide a synergistic effect to such changes in gene expression resulting from radiotherapy. In some embodiments the genes are related to cytokines and/or are related to apoptosis. Examples of the effect of radiotherapy, treatment with a nutritional supplement containing fish oil and selenium, and cotherapy with radiotherapy on gene expression in implanted tumor cells *in vivo* are shown in FIG. 8A to 8G. FIG. 8A shows the results of such treatments on the expression of VEGF in tumor cells. As shown, monotherapy with a nutritional supplement containing fish oil and selenium and radiotherapy reduced VEGF expression. Cotherapy with a nutritional supplement containing fish oil and selenium and radiotherapy provided dramatically reduced expression of VEGF.

[0070] FIGs. 8B and 8D show the effects of treatment with a nutritional supplement containing fish oil and selenium, radiotherapy, and cotherapy with such a supplement and radiotherapy on tumor BAX expression and metastatic (lung) tumor BAX expression, respectively. BAX is considered a marker for apoptosis. As shown, BAX expression in untreated tumor is low, and is not impacted by radiotherapy. Monotherapy with a nutritional supplement containing fish oil and selenium resulted in dramatic increases in BAX expression, and elevated BAX expression when used as cotherapy with radiotherapy (particularly with the supplement was provided as a pretreatment). As shown in FIGs. 8C and 8E, Bcl-2 expression was found to be elevated in tumor cells and metastatic (lung) tumor cells (respectively), and was reduced by either treatment with a nutritional supplement containing fish oil and selenium or radiotherapy as monotherapies. Greater reductions in Bcl-2 expression were found when such a supplement and radiotherapy were used as cotherapies. Expression of caspase 3 (which is associated with apoptosis) is elevated in tumors (FIG. 8D) treated with radiotherapy or with a nutritional supplement containing fish oil and selenium alone, and is also elevated by cotherapy with such a supplement as radiotherapy (particularly when the supplement is provided as a pre-treatment). As shown in FIG. 8F, expression of caspase 3 is reduced in metastatic (lung) tumors relative to lung tissue of control subjects. As shown, caspase 3 expression in such tumors is increased by radiotherapy or treatment with a nutritional supplement containing fish oil and selenium, as well as cotherapy.

[0071] Repeated radiotherapy

[0072] Similar studies were performed using modified treatment protocols. One modified treatment protocol utilizing multiple rounds of radiotherapy, as is typical with human radiotherapy, is shown in FIGs. 9A and 9B.

[0073] Results from serum albumin and blood cell characterization studies following treatment with NutraWell, radiotherapy, and combined treatment with a nutritional supplement containing fish oil and selenium and radiotherapy using the protocol shown in FIG. 9A are shown in FIGs. 10A to 10C. As shown in FIG. 10A serum albumin concentration is reduced in tumor bearing animals, particularly at later time points. This is improved by treatment with a nutritional supplement containing fish oil and selenium or radiotherapy as monotherapies, and by cotherapy. As shown in FIG. 10B, tumor bearing animals showed suppressed lymphocyte counts relative to

control subjects, particularly at later time points. This was only marginally improved by radiotherapy alone, however treatment with a nutritional supplement containing fish oil and selenium (either as a monotherapy or in combination with radiotherapy) was effective in increasing lymphocyte concentration- particularly in later time points. FIG. 10C shows the results of similar studies wherein the neutrophil/lymphocyte ratio (NLR) was characterized. As shown, animals bearing tumors show a dramatic elevation in this value at later time points. This is reduced by treatment with a nutritional supplement containing fish oil and selenium, radiotherapy, and cotherapy with such a supplement and radiotherapy.

[0074] Surprisingly, treatment with a nutritional supplement containing fish oil and selenium, radiotherapy, and combined treatment with such a supplement and radiotherapy using the protocol shown in FIG. 9A also has an impact on both expression of tumor cell markers and tumor cell metastasis. In the following studies the tumor cells selected for implantation are derived from a lung tumor and have a strong tendency to metastasize to the lung from the implantation site. FIGs. 11A to 11D show the results of immunocytochemistry studies of different tissues from mice treated with the protocol shown in FIG. 9A. It is notable that NutraWell supplementation alone reduces or eliminates metastasis.

[0075] As shown in FIG. 11A, VEGF expression within the tumor of untreated mice (left panel, FITC staining) is evident. Treatment with either a nutritional supplement containing fish oil and selenium or radiotherapy alone dramatically reduced VEGF expression, as does cotherapy. Numerical results are provided in the right panel of FIG. 11A. Similar results are found in tumors that have metastasized to the lungs, as shown in FIG. 11B. As shown, VEGF expression is apparent in untreated metastatic tumors (left panel, FITC staining) and is sharply reduced in animals that received treatment with a nutritional supplement containing fish oil and selenium, radiotherapy, or both. Numerical results of these studies are provided in the right panel of FIG. 11B.

[0076] Similar results are found with the expression of EGFR, overexpression of which is associated with tumors. As shown in FIG. 11C, elevated expression of EGFR (left panel, FITC staining) is evident in untreated tumors, and is reduced in subjects treated with a nutritional supplement containing fish oil and selenium, radiotherapy, or cotherapy with such a supplement

and radiotherapy. Numerical results are shown in the right panel of FIG. 11C. Similar results are found for EGFR expression of tumors that have metastasized to the lungs, as shown in FIG. 11D. As shown, elevated EGFR expression (left panel, FITC staining) is evident in untreated metastatic sites, and is reduced in subjects treated with a nutritional supplement containing fish oil and selenium or radiotherapy as monotherapies, and with cotherapy using such a supplement and radiotherapy. This suggests that use of a nutritional supplement containing fish oil and selenium, radiotherapy, and/or a combination of treatment with such a supplement and radiotherapy can be effective in enhancing EGFR-directed treatment protocols in individuals receiving treatment for cancer.

[0077] Surprisingly, treatment with a nutritional supplement containing fish oil and selenium, radiotherapy, and combined treatment with such a supplement and radiotherapy using the protocol shown in FIG. 9A also has an impact on tumor stem cells. Such stem cells are associated with metastasis and the development of resistance to various cancer therapies. In the following studies the tumor cells selected for implantation are derived from a lung tumor and have a strong tendency to metastasize to the lung from the implantation site. FIG. 12 shows the results of immunocytochemistry studies of different tissues from mice treated with the protocol shown in FIG. 9A. The left panel of FIG. 12 shows the results of staining for CD31 (FITC staining), a stem cell marker, in tumor cells. The right panel shows similar results for metastatic cells in the lungs. Untreated subjects show numerous cells with elevated expression of CD31. Surprisingly, treatment with a nutritional supplement containing fish oil and selenium in the absence of radiotherapy reduces or eliminates the occurrence of cancer stem cells both in the tumor implantation site and at the lung metastatic site.

[0078] Treatment with a nutritional supplement containing fish oil and selenium, radiotherapy, and combined NutraWell supplementation and radiotherapy using the protocol shown in FIG. 9A also has an impact on hypoxia often found in or among tumor cells. HIF1- α is a marker associated with hypoxia. In the following studies the tumor cells selected for implantation are derived from a lung tumor and have a strong tendency to metastasize to the lung from the implantation site. FIG. 13A shows the results of immunocytochemistry studies of different tissues from mice treated with the protocol shown in FIG. 9A. The left panel shows the results of immunocytochemistry staining for HIF1- α (FITC staining) in tumor cells, while the right

panel shows the results of similar staining at lung metastatic sites. Untreated subjects show numerous cells with elevated levels of HIF1- α . Surprisingly, treatment with a nutritional supplement containing fish oil and selenium in the absence of radiotherapy reduces or eliminates the occurrence of such hypoxia markers both in the tumor implantation site and at the lung metastatic site. FIG. 13B shows typical numerical results from a similar study where gene expression in tumor samples was characterized. As shown, both treatment with a supplement containing fish oil and selenium (TN) and pre-treatment with a nutritional supplement containing fish oil and selenium (PTN) for one week reduce tumor expression of HIF1- α , as does radiotherapy (TR). When radiotherapy and such a nutritional supplement are used in combination (PTRN, TRN) a dramatic reduction in tumor HIF1- α expression is evident, particularly in pretreated subjects (PTRN) (which show a synergistic effect). It's evident that use of a nutritional supplement containing fish oil and selenium in combination with radiotherapy can reduce both HIF1- α protein content and gene expression in tumor sites, potentially rendering them more vulnerable to hypoxic conditions.

[0079] Treatment with a nutritional supplement containing fish oil and selenium, radiotherapy, and combined supplementation and radiotherapy using the protocol shown in FIG. 9A also has an impact on apoptic activity in tumor cells. Results of qPCR studies for expression of various apoptosis markers (Bax, Bcl-2, and caspase 3) at 24 days from tumor cell implantation in mice is shown in FIG. 14. As shown, the Bax/Bcl-2 expression ratio is low in tumor cells, and only marginally improved by radiotherapy. Surprisingly, treatment with a nutritional supplement containing fish oil and selenium alone provided a significant increase in this ratio. In addition, a pronounced synergistic effect was found on cotherapy with such a supplement and radiotherapy. Caspase 3 expression was actually somewhat suppressed by radiotherapy alone, but was dramatically increased by treatment with a nutritional supplement containing fish oil and selenium, both as a monotherapy and as cotherapy with radiotherapy.

[0080] Treatment with a nutritional supplement containing fish oil and selenium in combination with radiotherapy has also been found to modulate expression of PDL-1 and PD-1 in primary tumor sites and metastatic (lung) tumor sites in animal models of human disease. As shown in FIG. 15A, both primary (left panel) and metastatic (right panel) tumor cells show high levels of PDL-1 expression (T). Surprisingly, these are markedly reduced by treatment with a nutritional

supplement containing fish oil and selenium, both when pretreated (PTN) and treated at the time of implantation (TN). As such, the Inventor believes that such supplements can be used to enhance immunotherapeutic approaches to cancer treatment. Treatment with radiotherapy (TR) also shows a reduction in PD-L1 gene expression in both primary and metastatic sites, suggesting that radiotherapy alone can be used to enhance immunotherapeutic approaches to cancer treatment. Cotherapy with such a nutritional supplement and radiotherapy is effective in reducing PDL-1 expression in both primary tumor sites and metastatic sites, whether the supplement is provided prior to implantation (PTRN) or at the time of implantation (TRN). Surprisingly, the effects are more marked at the primary tumor site than at metastatic sites. This suggests that cotherapy with a nutritional supplement that includes fish oil and selenium in combination with radiotherapy can render tumor cells at both primary and metastatic sites more susceptible to the patient's immune system, and/or can enhance the effect of immunotherapy approaches to cancer treatment.

[0081] Similar studies were performed on PD-1 expression. As shown in FIG. 15B, PD-1 expression is reduced in primary site tumor cells treated with radiotherapy (TR) and enhanced in primary site tumor cells (left panel) in animals treated with a supplement containing fish oil and selenium, either prior to implantation (PTN) or upon implantation (TN). Cotherapy with the nutritional supplement and radiotherapy provides differential results depending on if the supplement is provided prior to implantation (PTRN) or at the time of implantation (TRN). As shown in the right panel of FIG. 15B, tumor cells at metastatic sites showed reduced PD-1 expression relative to samples taken from control animals. Surprisingly, radiotherapy resulted in an increase in PD-1 expression at metastatic sites (right panel), as did treatment with a supplement containing fish oil and selenium either prior to implantation or at the time of implantation. Cotherapy with radiotherapy and such a supplement provided higher levels of PD-1 expression than those observed with monotherapy. It is apparent that treatment with a nutritional supplement containing fish oil and selenium, particularly in combination with radiotherapy, can shift PD-1 expression away from reduced expression levels seen in tumor-bearing subjects, particularly in metastatic sites.

[0082] Preimplantation, preradiotherapy, and initiation of radiotherapy supplementation with repeated radiotherapy

[0083] Another treatment protocol combining both pretreatment with a nutritional supplement containing fish oil and selenium and multiple rounds of radiotherapy typical of clinical application is shown in FIGs. 16A and 16B. In this protocol treatment with the supplement was initiated prior to tumor cell implantation, at the time of tumor cell implantation, and at the initiation of radiotherapy.

[0084] Loss of intestinal absorption and the resulting malnutrition are a well known side effect of radiotherapy, particularly repeated radiotherapy. FIG. 17 shows photomicrographs that demonstrate the effect treatment with a nutritional supplement containing fish oil and selenium on the cellular architecture of the gut during radiotherapy. Cellular architecture of the gut is shown for untreated control subjects (top left panel), tumor implanted subjects treated with multiple rounds of radiotherapy (top right panel), tumor implanted subjects pretreated with a nutritional supplement containing fish oil and selenium and subsequently treated with radiotherapy (lower left panel), and tumor implanted subjects treated with a nutritional supplement containing fish oil and selenium at the time of initiation of radiotherapy (lower right panel). As shown, treatment with a nutritional supplement containing fish oil and selenium can maintain the intestinal brush border during radiotherapy, with pretreatment apparently enhancing the intestinal brush border. This indicates that treatment with such a supplement, particularly pretreatment, can effectively address side effects of radiotherapy.

[0085] Such a treatment protocol has been found to modify expression of certain genes in tumor cells *in vivo*. Results of qPCR studies of gene expression (e.g. angiogenic factor-related, apoptosis-related, etc.) in tumor cells from mice treated by the protocol shown in FIG. 16A are shown in FIGs. 18A to 18F. As shown in FIG. 18A, treatment with a nutritional supplement containing fish oil and selenium reduces VEGF expression in tumor cells, as does repeated radiotherapy. Combined therapy with such a supplement and radiotherapy provides an enhanced reduction in VEGF expression, particularly when a nutritional supplement containing fish oil and selenium is provided prior to initiation of radiotherapy.

[0086] FIG. 18B shows that BAX expression in tumor cells is increased by treatment with a nutritional supplement containing fish oil and selenium, and relatively unaffected by repeated radiotherapy alone. Pretreatment with such a supplement in combination with repeated

radiotherapy also increased BAX expression. Similar results are found for expression of another apoptosis-related gene (Caspase 3), as shown in FIG. 18D. Results of studies of Caspase-3 expression in metastatic (lung) tumors are shown in FIG. 18F. As shown, expression of Caspase-3 in such metastatic tumors is elevated by treatment with a nutritional supplement containing fish oil and selenium and by repeated radiotherapy, and is similar to that of control cells when the supplement is provided as cotherapy prior to initiation of radiotherapy.

[0087] Expression of another apoptosis-related gene, Bcl-2, in tumor cells is reduced by treatment with a nutritional supplement containing fish oil and selenium and by repeated radiotherapy alone (see FIG. 18C). This reduction in Bcl-2 expression is more pronounced on cotherapy with such a supplement and repeated radiotherapy, particularly when the supplement is provided prior to initiation of radiotherapy. FIG. 18E shows the results of similar studies performed on metastatic (lung) tumors. As shown, the reduction in Bcl-2 expression is reduced in a synergistic fashion on cotherapy with a nutritional supplement containing fish oil and selenium and repeated radiotherapy, particularly when the supplement is provided prior to initiation of radiotherapy.

[0088] Pre-implantation supplementation and repeated radiotherapy

[0089] FIGs. 19A and 19B show a treatment protocol and related test groups (respectively) where a nutritional supplement containing fish oil and selenium is provided 7 days prior to implantation of tumor cells, with radiotherapy taking place on days 8, 10, and 12 following implantation. Mice were sacrificed on day 24 following implantation.

[0090] A well known side effect of both cancer and repeated radiotherapy is weight loss. This can be due to wasting associated with disease and with side effects of radiotherapy. The effect of treatment with a nutritional supplement containing fish oil and selenium on loss of body mass and on muscle mass following repeated radiotherapy is shown in FIGs. 20A and 20B. It should be appreciated that body mass was characterized following removal of the tumor mass. As shown in FIG. 20A, when tumor mass is removed a significant decrease in remaining body mass is apparent relative to control subjects. This is improved by treatment with a nutritional supplement containing fish oil and selenium and by radiotherapy as monotherapies. Surprisingly, gains in body mass surpass those of control subjects when such a supplement and

repeated radiotherapy are used in combination. FIG. 20B shows the weight of gastrocnemius muscle in the various test groups. Loss of muscle mass is apparent in untreated tumor bearing animals. This is marginally improved by repeated radiotherapy alone. Surprisingly, treatment with a nutritional supplement containing fish oil and selenium provides significant retention in muscle mass. Cotherapy with a nutritional supplement containing fish oil and selenium and repeated radiotherapy provides a synergistic improvement in muscle mass relative to untreated tumor bearing subjects. It is apparent that pre-treatment with a nutritional supplement containing fish oil and selenium effectively reverses the loss of body mass (relative to control subjects) and muscle mass resulting from both the presence of the tumor and repeated radiotherapy.

[0091] Preimplantation supplementation and repeated radiotherapy

[0092] Another treatment protocol with related treatment groups is shown in FIGs. 21A and 21B, which is similar to the protocol shown in FIGs 16A and 16B. In this protocol radiotherapy is provided on days 8, 10, and 12 following implantation of tumor cells. Treatment with a nutritional supplement containing fish oil and selenium is provided either 7 days prior to implantation, the day of implantation, or on the initiation of radiotherapy. Mice were sacrificed on day 21 following tumor cell implantation.

[0093] A well known side effect of both cancer and radiotherapy is weight loss. The effect of treatment with a nutritional supplement containing fish oil and selenium on loss of body mass and loss of muscle mass when provided prior to and when provided coincident with the initiation of repeated radiotherapy is shown in FIG. 22. It should be appreciated that body mass was characterized following removal of the tumor mass. As shown, body weight gains over time of untreated tumor bearing subjects (following excision of the tumor) is dramatically reduced relative to control subjects. Similar losses are noted on repeated radiotherapy when used alone. Surprisingly, treatment with a nutritional supplement containing fish oil and selenium provides remarkably improved body weight gains over time, both as a monotherapy and when used in combination with repeated radiotherapy. This is particularly evident with such a supplement is used as a pretreatment.

[0094] Inventors have also found that pre-treatment with a nutritional supplement containing fish oil and selenium enhances the reduction in tumor volume seen on repeated radiotherapy, as

shown in FIG. 23. As shown, both repeated radiotherapy and treatment with a nutritional supplement containing fish oil and selenium have a moderate effect in reducing tumor volume relative to tumors of untreated tumor bearing animals. Surprisingly, tumor volume shows almost no change over time when repeated radiotherapy and a nutritional supplement containing fish oil and selenium are used as cotherapies, indicating a synergistic effect.

[0095] The presence of tumors and repeated radiotherapy can also result in the development of inflammation, which can be characterized by the presence of pro-inflammatory cytokines in serum. FIGs. 24A and 24B show the effect of administration of a nutritional supplement containing fish oil and selenium with repeated radiotherapy on the concentration of pro-inflammatory cytokines in mice treated as in the protocol shown in FIG. 21A. FIG. 24A shows values for serum TNF- α . It is apparent that untreated tumor bearing animals show highly elevated concentrations of TNF- α , which is reduced to some extent by repeated radiotherapy. Treatment with a nutritional supplement containing fish oil and selenium also resulted in a reduction in serum TNF- α , particularly when such a supplement was used in cotherapy with repeated radiotherapy. FIG. 24B shows the results of similar studies where the serum concentration of IL-6 was characterized, which show similar results.

[0096] It should be apparent to those skilled in the art that many more modifications besides those already described are possible without departing from the inventive concepts herein. The inventive subject matter, therefore, is not to be restricted except in the spirit of the appended claims. Moreover, in interpreting both the specification and the claims, all terms should be interpreted in the broadest possible manner consistent with the context. In particular, the terms “comprises” and “comprising” should be interpreted as referring to elements, components, or steps in a non-exclusive manner, indicating that the referenced elements, components, or steps may be present, or utilized, or combined with other elements, components, or steps that are not expressly referenced. Where the specification claims refers to at least one of something selected from the group consisting of A, B, C and N, the text should be interpreted as requiring only one element from the group, not A plus N, or B plus N, etc.

CLAIMS

What is claimed is:

1. A method of treating a tumor, comprising:
 - applying a radiotherapy protocol to a patient in need of treatment; and
 - providing the patient with a nutritional supplement comprising fish oil and selenium, in an amount to provide a synergistic effect in reducing tumor volume or weight.
2. The method of claim 1, wherein the nutritional supplement is provided to the patient prior to the initiation of radiotherapy.
3. The method of claim 1 or 2, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
4. The method of claim 1 or 2, wherein the supplement comprises three or more components are as described in Table 1.
5. A method of reducing side effects of radiotherapy, comprising:
 - applying a radiotherapy protocol to a patient in need of treatment; and
 - providing the patient with a nutritional supplement comprising fish oil and selenium, in an amount to reduce a side effect of the radiotherapy protocol.
6. The method of claim 5, wherein the nutritional supplement is provided to the patient prior to the initiation of radiotherapy.
7. The method of claim 5 or 6, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
8. The method of claim 5 or 6, wherein the supplement comprises three or more components are as described in Table 1.
9. A method of modulating gene expression in a tumor, comprising:
 - providing a nutritional supplement to a tumor or an animal having a tumor, wherein the nutritional supplement comprising fish oil and selenium, and wherein the nutritional supplement is provided in an amount to modulate expression of a gene of the tumor.

10. The method of claim 9, wherein the nutritional supplement is provided to the patient prior to the initiation of radiotherapy.
11. The method of claim 9 or 10, wherein the nutritional supplement is provided to the patient during the application of radiotherapy.
12. The method of one of claims 9 to 11, wherein the nutritional supplement is formulated such that the plurality of components of the nutritional supplement are provided in amounts as described in Table 1.
13. The method of one of claims 9 to 11, wherein the supplement comprises three or more components are as described in Table 1.
14. The method of one of claims 9 to 13, wherein the gene encodes for an angiogenesis factor.
15. The method of one of claims 9 to 13, wherein the gene encodes for an apoptosis factor.
16. A method of reducing metastasis from a tumor, comprising:
 - providing the patient having a metastatic tumor with a nutritional supplement comprising fish oil and selenium, and
 - applying a radiotherapy protocol to the patient,
 - wherein the nutritional supplement is provided in an amount to reduce metastatic activity of the tumor.
17. The method of claim 16, wherein the nutritional supplement is provided concurrent with the radiotherapy protocol.
18. The method of claim 16 or 17, wherein the nutritional supplement is formulated such that the plurality of components of the nutritional supplement are provided in amounts as described in Table 1.
19. The method of claim 16 or 17, wherein the supplement comprises three or more components are as described in Table 1.
20. A method of reducing angiogenesis in a tumor, comprising:

providing the patient having a tumor with a nutritional supplement comprising fish oil and selenium;
applying a radiotherapy protocol to the patient,
wherein the nutritional supplement is provided in an amount to reduce an angiogenic activity of the tumor.

21. The method of claim 20, wherein the nutritional supplement is provided concurrent with radiotherapy.
22. The method of claim 20 or 21, wherein the nutritional supplement is formulated such that the plurality of components of the nutritional supplement are provided in amounts as described in Table 1.
23. The method of claim 20 or 21, wherein the supplement comprises three or more components are as described in Table 1.
24. A method of reducing cancer stem cell of a tumor, comprising:
providing a patient having a tumor with a nutritional supplement comprising fish oil and selenium; and
applying a radiotherapy protocol,
wherein the nutritional supplement is provided in an amount effective to reduce occurrence of cancer stem cells in the tumor.
25. The method of claim 24, wherein the nutritional supplement is provided concurrent with the radiotherapy protocol.
26. The method of claim 24 or 25, wherein the nutritional supplement is formulated such that the plurality of components of the nutritional supplement are provided in amounts as described in Table 1.
27. The method of claim 24 or 25, wherein the nutritional supplement comprises three or more components are as described in Table 1.
28. A method of reducing PDL-1 expression in a cancer cell, comprising applying a radiotherapy protocol to the cancer cell.

29. The method of claim 28, further comprising the step of contacting the cancer cell with a nutritional supplement comprising fish oil and selenium.
30. The method of claim 29, wherein the nutritional supplement is provided prior to the initiation of radiotherapy.
31. The method of claim 29, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
32. The method of claim 29, wherein the supplement comprises three or more components are as described in Table 1.
33. A method of reducing PDL-1 expression in a cancer cell, comprising contacting the cancer cell with a nutritional supplement comprising fish oil and selenium.
34. The method of claim 33, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
35. The method of claim 33, wherein the nutritional supplement comprises three or more components are as described in Table 1.
36. A method of enhancing anti-tumor immunotherapy in an individual, comprising:
 - applying an anti-tumor immunotherapy protocol to the patient; and
 - applying a radiotherapy protocol to the patient.
37. The method of claim 36, further comprising the step of administering a nutritional supplement comprising fish oil and selenium to the patient.
38. The method of claim 37, wherein the nutritional supplement is provided prior to the initiation of radiotherapy.
39. The method of claim 37 or 38, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
40. The method of claim 37 or 38, wherein the supplement comprises three or more components are as described in Table 1.

41. A method of enhancing anti-tumor immunotherapy in an individual, comprising:
 - applying an anti-tumor immunotherapy protocol to the patient; and
 - providing the patient with a nutritional supplement comprising fish oil and selenium.
42. The method of claim 41, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
43. The method of claim 41, wherein the nutritional supplement comprises three or more components are as described in Table 1.
44. A method of reducing EGFR expression in a cancer cell, comprising applying a radiotherapy protocol to the cancer cell.
45. The method of claim 44, further comprising the step of contacting the cancer cell with a nutritional supplement comprising fish oil and selenium.
46. The method of claim 45, wherein the nutritional supplement is provided prior to the initiation of radiotherapy.
47. The method of claim 45 or 46, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
48. The method of claim 45 or 46, wherein the supplement comprises three or more components are as described in Table 1.
46. A method of reducing EGFR expression in a cancer cell, comprising contacting the cancer cell with a nutritional supplement comprising fish oil and selenium.
47. The method of claim 46, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
48. The method of claim 46, wherein the nutritional supplement comprises three or more components are as described in Table 1.
49. A method of enhancing anti-EGFR therapy in an individual with cancer, comprising:
 - applying an anti-EGFR therapy protocol to the patient; and

applying a radiotherapy protocol to the patient.

50. The method of claim 49, further comprising the step of administering a nutritional supplement comprising fish oil and selenium to the patient.
51. The method of claim 50, wherein the nutritional supplement is provided prior to the initiation of radiotherapy.
52. The method of claim 50 or 51, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
53. The method of claim 50 or 51, wherein the supplement comprises three or more components are as described in Table 1.
54. A method of enhancing anti-EGFR therapy in an individual with cancer, comprising:
 - applying an anti-EGFR therapy protocol to the patient; and
 - providing the patient with a nutritional supplement comprising fish oil and selenium.
55. The method of claim 54, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
56. The method of claim 54, wherein the nutritional supplement comprises three or more components are as described in Table 1.
57. Use of a nutritional supplement for manufacturing a medicament that improves radiotherapy in an individual with cancer, comprising fish oil and selenium, wherein fish oil and selenium are provided in amounts that provide a synergistic effect in reducing tumor volume or weight when used in combination with a radiotherapy protocol.
58. The use of claim 57, wherein the nutritional supplement is formulated for provision to the patient prior to the initiation of radiotherapy.
59. The use of claim 57 or 58, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.

60. The use of claim 57 or 58, wherein the supplement comprises three or more components are as described in Table 1.
61. Use of a nutritional supplement for manufacturing a medicament that reduces side effects associated with radiotherapy in an individual receiving radiotherapy for cancer, comprising fish oil and selenium, wherein fish oil and selenium are provided in amounts that provide a synergistic effect in reducing side effects of a radiotherapy protocol.
62. The use of claim 61, wherein the nutritional supplement is formulated to be effective when administered to the patient prior to initiation of radiotherapy.
63. The use of claim 61 or 62, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
64. The use of claim 61 or 62, wherein the supplement comprises three or more components are as described in Table 1.
65. Use of a nutritional supplement for manufacturing a medicament that modulates gene expression in a tumor of a patient receiving radiotherapy, comprising fish oil and selenium, wherein the nutritional supplement is provided in an amount to modulate expression of a gene of the tumor.
66. The use of claim 65, wherein the nutritional supplement is formulated to be provided to the patient prior to initiation of radiotherapy.
67. The use of claim 65, wherein the nutritional supplement is formulated to be provided to the patient during application of radiotherapy.
68. The use of one of claims 65 to 67, wherein the nutritional supplement is formulated such that the plurality of components of the nutritional supplement are provided in amounts as described in Table 1.
69. The use of claim one of claims 65 to 67, wherein the supplement comprises three or more components are as described in Table 1.

70. The use of one of claims 65 to 69, wherein the gene encodes for an angiogenesis factor.
71. The use of one of claims 65 to 69, wherein the gene encodes for an apoptosis factor.
72. Use of a nutritional supplement for manufacturing a medicament that reduces metastasis from a tumor of a patient receiving radiotherapy, comprising fish oil and selenium, wherein the nutritional supplement is provided in an amount to reduce metastatic activity of the tumor.
73. The use of claim 72, wherein the nutritional supplement is formulated to be provided concurrent with a radiotherapy protocol.
74. The use of claim 72 or 73, wherein the nutritional supplement is formulated such that the plurality of components of the nutritional supplement are provided in amounts as described in Table 1.
75. The use of claim 72 or 73, wherein the supplement comprises three or more components are as described in Table 1.
76. Use of a nutritional supplement for manufacturing a medicament that reduces angiogenesis in a tumor of a patient receiving radiotherapy, comprising fish oil and selenium, wherein the nutritional supplement is provided in an amount to reduce an angiogenic activity of the tumor.
77. The use of claim 76, wherein the nutritional supplement is provided concurrent with radiotherapy.
78. The use of claim 76 or 77, wherein the nutritional supplement is formulated such that the plurality of components of the nutritional supplement are provided in amounts as described in Table 1.
79. The use of claim 76 or 77, wherein the supplement comprises three or more components are as described in Table 1.
80. Use of a nutritional supplement for manufacturing a medicament that reduces cancer stem cell content of a tumor of a patient receiving radiotherapy, comprising fish oil and selenium,

wherein the nutritional supplement is provided in an amount effective to reduce occurrence of cancer stem cells in the tumor.

81. The use of claim 80, wherein the nutritional supplement is formulated to be provided concurrent with a radiotherapy protocol.

82. The use of claim 80 or 81, wherein the nutritional supplement is formulated such that the plurality of components of the nutritional supplement are provided in amounts as described in Table 1.

83. The use of claim 80 or 81, wherein the nutritional supplement comprises three or more components are as described in Table 1.

84. Use of a nutritional supplement for manufacturing a medicament that reduces PDL-1 expression in a cancer cell receiving radiotherapy, comprising fish oil and selenium, wherein the nutritional supplement is provided in an amount effective to reduce PDL-1 of the cancer cell.

85. The use of claim 84, wherein the nutritional supplement is formulated to be provided prior to the initiation of radiotherapy.

86. The use of claim 84 or 85, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.

87. The use of claim 84 or 85, wherein the nutritional supplement comprises three or more components as described in Table 1.

88. The use of one of claims 84 to 87, wherein the cancer cell is being exposed to radiotherapy.

89. Use of a nutritional supplement for manufacturing a medicament that enhances immunotherapy in an individual receiving immunotherapy for cancer, comprising fish oil and selenium, wherein the nutritional supplement is provided in an amount effective to reduce PDL-1 of the cancer cell.

90. The use of claim 89, wherein the nutritional supplement is provided prior to initiation of radiotherapy.

91. The use of claim 89 or 90, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
92. The use of claim 89 or 90, wherein the supplement comprises three or more components are as described in Table 1.
93. Use of a nutritional supplement for manufacturing a medicament that reduces EGFR expression in a cancer cell, comprising fish oil and selenium, wherein the nutritional supplement is provided in an amount effective to reduce EGFR of the cancer cell.
94. The use of claim 93, wherein the cancer cell is subjected to radiotherapy.
95. The use of claim 94, wherein the nutritional supplement is formulated to be provided prior to initiation of radiotherapy.
96. The use of one of claims 93 to 95, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
96. The use of one of claims 93 to 95, wherein the supplement comprises three or more components are as described in Table 1.
97. Use of a nutritional supplement for manufacturing a medicament that enhances anti-EGFR therapy in an individual with cancer, comprising fish oil and selenium, wherein the nutritional supplement is provided in an amount effective to reduce EGFR of a cancer cell.
98. The use of claim 97, wherein the individual is receiving radiotherapy.
99. The use of claim 94, wherein the nutritional supplement is formulated to be provided prior to initiation of radiotherapy.
100. The use of one of claims 97 to 99, wherein the nutritional supplement is formulated such that a plurality of components of the supplement are provided in amounts as described in Table 1.
101. The use of one of claims 97 to 97, wherein the supplement comprises three or more components are as described in Table 1.




FIG. 1




FIG. 2A




FIG. 2B




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D




FIG. 4E




FIG. 4F





FIG. 4G

FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E




FIG. 6A




FIG. 6B




FIG. 7A












FIG. 7B

**FIG. 8A****FIG. 8B**

**FIG. 8C****FIG. 8D**

**FIG. 8E****FIG. 8F**

**FIG. 8G**

**FIG. 9A**

			Scalify (14 th day)	Scalify (24 th day)
1	C	control	N=6	N=6
2	T	Tumor	N=6	N=6
3	TN	Tumor + Nutrawell	N=6	N=6
4	T3R	Tumor + Radiotherapy x3	N=6	N=6
5	T3RN	Tumor + Radiotherapy x3+ Nutrawell	N=6	N=6

FIG. 9B




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 11D

24D





FIG. 12






FIG. 13B

	Bax	Bcl-2	Bax/Bcl-2	caspase3
T	0.31±0.00	1.41±0.67	0.22±0.12	1.03±0.88
TN	5199.29±2321.78	241.89±7.44	21.49±10.19	47708.35±7808.87
TR	1.09±1.39	0.99±0.16	1.10±0.03	0.71±0.06
TRN	7858.24±1593.75	213.40±8.45	36.82±9.56	46977.1±5263.41

FIG. 14




FIG. 15A




FIG. 15B




FIG. 16A

			Scalify (21 th day)
1	C	Control	N=6
2	T	Tumor	N=6
3	PTN	Tumor + Nutrawell (-7 day start)	N=6
4	TN	Tumor + Nutrawell (0 day start)	N=6
5	TR	Tumor + Radiotherapy (3 Gy x 3)	N=6
6	PTRN	Tumor + Radiotherapy (3 Gy x 3) + Nutrawell (-7 day start)	N=6
7	TRN	Tumor + Radiotherapy (3 Gy x 3) + Nutrawell (8 day start)	N=6

FIG. 16B




FIG. 17





FIG. 18A

**FIG. 18B**




FIG. 18C




FIG. 18D





FIG. 18E

**FIG. 18F**




FIG. 19A

			Scalify (24 th day)
1	C	Control	N=6
2	T	Tumor	N=6
3	TN	Tumor + Nutrawell	N=6
4	TR	Tumor + Radioherap x3	N=6
5	TRN	Tumor + Radioherap x3 + Nutrawell	N=6

FIG. 19B




FIG. 20A




FIG. 20B




FIG. 21A

			Scalify (21 th day)
1	C	Control	N=6
2	T	Tumor	N=6
3	PTN	Tumor + Nutrawell (-7 day start)	N=6
4	TN	Tumor + Nutrawell (0 day start)	N=6
5	TR	Tumor + Radiotherapy (3 Gy x 3)	N=6
6	PTRN	Tumor + Radiotherapy (3 Gy x 3) + Nutrawell (-7 day start)	N=6
7	TRN	Tumor + Radiotherapy (3 Gy x 3) + Nutrawell (8 day start)	N=6

FIG. 21B




FIG. 22




FIG. 23




FIG. 24A




FIG. 24B