



(51) International Patent Classification:

C12Q 1/68 (2006.01) *A61K 39/395* (2006.01)
A61B 5/00 (2006.01) *A61P 35/00* (2006.01)
A61B 5/06 (2006.01) *C07H 21/04* (2006.01)
A61K 31/4439 (2006.01) *G01N 33/48* (2006.01)
A61K 31/7088 (2006.01) *G01N 33/50* (2006.01)
A61K 38/00 (2006.01)

(21) International Application Number:

PCT/CA2016/050846

(22) International Filing Date:

18 July 2016 (18.07.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/194,109 17 July 2015 (17.07.2015) US

(71) Applicant: **PACYLEX PHARMACEUTICALS INC.**
[CA/CA]; 2600 Manulife Place, 10180-101 Street, Edmon-
ton, Alberta T5J 3Y2 (CA).

(72) Inventors: **BERTHIAUME, Luc G.**; 10061, 90 Avenue,
Edmonton, Alberta T6E 4X2 (CA). **BEAUCHAMP, Er-
wan**; #302 10033, 89 Avenue, Edmonton, Alberta T6E
2S7 (CA).

(74) Agents: **VICKERS, Mark, F.** et al.; Borden Ladner Ger-
vais LLP, World Exchange Plaza, 100 Queen Street, Suite
1300, Ottawa, Ontario K1P 1J9 (CA).


(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: EPIGENETIC SILENCING OF NMT2

Figure 2 B -- Epigenetic regulation of NMT2 in lymphoma cell lines and tumours.



(57) Abstract: The finding that multiple cancers lack one of two NMTs, while stromal and normal tissues do not, enables the treatment of NMT2-deficient cancer cells with an NMT inhibitor. It is shown herein that NMT2 expression is reduced or eliminated in certain cancers, and in one example lymphomas, via an epigenetic mechanism(s). Reduction or elimination of NMT2 expression renders the cancer sensitive inhibitors of NMT.

WO 2017/011907 A1



Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

EPIGENETIC SILENCING OF NMT2**CROSS REFERENCE TO RELATED APPLICATION**

- 5 This application claims priority to U.S. provisional patent application number US 62/194,109, filed July 17, 2015, the entire contents of which are hereby incorporated by reference.

FIELD

- 10 [0001] The present disclosure relates generally to epigenetic methods for cancer diagnosis and treatment.

BACKGROUND

- [0002] Cancer is a leading cause of death in Canada. The Canadian Cancer Society estimate there will be approximately 170000 new cases of cancer in 2011, and approximately 75000 deaths as a result of cancer.

[0003] It has been estimated that in 2015, about 589430 Americans will die of cancer, or about 1,620 people per day. Cancer is the second most common cause of death in the US, and accounts for nearly 1 of every 4 deaths.

- 20 [0004] Hematological cancers account for approximately nine percent of new cancer cases and nine percent of cancer related deaths. One type of hematological cancer is lymphoma. Aggressive non Hodgkin Lymphomas such as Burkitt Lymphoma (BL) and Diffuse Large B Cell Lymphoma (DLBCL) are treatable, but current therapies are expensive, carry serious toxicities, and achieve widely varying responses, with the majority of patients ultimately experiencing disease relapse. In North America alone, there are 78,000 new NHL cases per year and 600,000 NHL survivors in remission but with ongoing risk of relapse. Improved treatments are required.

- [0005] N-myristoylation of proteins is a modification in which myristate (a 14-carbon saturated fatty acid) is covalently attached to the NH₂ terminal glycine of a variety of cellular, viral, and onco-proteins (e.g., oncogenic Src-related tyrosine kinases, heterotrimeric G alpha subunits, etc.).

- 30 [0006] Cellular myristoylated proteins have diverse biological functions in signal transduction and oncogenesis. Modification of proteins by myristoylation is required for the subcellular targeting, protein conformation and biological activity of many important proteins in eukaryotic cells, including those required for signal transduction and regulatory
- 35

functions important in cell growth. Tyrosine kinases of the Src family (proto-oncogenes) are among the most extensively studied myristoylated proteins.

[0007] Myristoylation of proteins is catalyzed by N-myristoyltransferase (NMT).

NMT is responsible for this activity in eukaryotic cells and works by modifying its

5 polypeptide substrate after the removal of the initiator methionine residue by methionyl aminopeptidase. This modification occurs primarily as a cotranslational process, although myristoylation can also occur post-translationally after proteolytic cleavage of proteins, typically during apoptosis. Two isozymes of the mammalian NMT enzymes have been cloned and are designated NMT1 and NMT2. NMTs play a pro-survival role in cells.

10 The two NMTs are present in all normal mammalian cells.

[0008] There remains a need for compounds, composition and method for the treatment of cancer.

[0009] This background information is provided for the purpose of making known information believed by the applicant to be of possible relevance to the present invention.

15 No admission is necessarily intended, nor should it be construed, that any of the preceding information constitutes prior art against the present invention.

SUMMARY

[0010] In one aspect of the present disclosure there is provided a method of
20 predicting the response in a cancer patient to treatment with an NMT inhibitor comprising, providing a biological sample of said cancer patient, and determining in said biological sample the methylation status of the *NMT2* gene, and predicting a positive clinical response to said treatment with said NMT inhibitor, if hypermethylation is determined in the *NMT2* gene.

25 **[0011]** In one aspect of the present disclosure there is provided a method for identifying and/or selecting a patient with a cancer, or treated for a cancer, suitable for treatment with an NMT inhibitor, comprising providing a biological sample of said cancer patient, and determining the methylation status of the *NMT2* gene, and identifying and/or selecting the cancer patient for treatment with said NMT inhibitor if hypermethylation is
30 determined in the *NMT2* gene.

[0012] In one aspect of the present disclosure there is provided a method for selecting a suitable treatment regimen for a cancer in a cancer patient comprising providing a biological sample of said cancer patient, and determining in said biological sample the methylation status of the *NMT2* gene, and selecting an NMT inhibitor for

treatment if hypermethylation of *NMT2* gene is determined and/or under expression is determined in the *NMT2* gene.

[0013] In one aspect of the present disclosure there is provided a method of treating a patient with cancer with an NMT inhibitor comprising providing a biological sample of said cancer patient, and determining in said biological sample the methylation status of *NMT2* gene, and selecting said NMT inhibitor for treatment if hypermethylation is determined in *NMT2* gene.

[0014] In one aspect of the present disclosure there is provided a method comprising: a) obtaining a sample from a subject having cancer, or suspected as having cancer; b) contacting the sample with a reagent to form a product (in some cases a complex) indicative of methylation status of *NMT2* gene present in the sample; c) measuring the product formed to determine a methylation status of *NMT2* gene in the sample; and d) determining the benefit of NMT inhibitor treatment of said cancer in in said subject, wherein the determination of benefit of NMT inhibitor treatment is determined by hypermethylation of *NMT2* gene in said sample.

[0015] In one aspect, wherein in step b) said reagent comprises primers to detect hypermethylation in the *NMT2* gene.

[0016] In one aspect said primers comprise SEQ ID NO:1 and/or SEQ ID NO: 2.

[0017] In one aspect of the present disclosure there is provided a method of treating a subject with cancer or suspect of having cancer, comprising administering an NMT inhibitor to said subject when the *NMT2* gene is hypermethylated in said sample, optionally as compared to a control.

[0018] In one aspect of the present disclosure there is provided a method for treating cancer in a subject, comprising: a) obtaining nucleic acid from a sample from the subject; b) performing bisulfite modification to the nucleic acid in step a); c) performing a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, on the bisulfite modified nucleic acid from step b) using PCR primers and/or probes specific for the promoter region of the *NMT2* gene, e) determining a promoter methylation level of the promoter region of *NMT2* in the sample, optionally compared to a control, g) treating said patient with an inhibitor to NMT when promoter methylation level of the promoter region of *NMT2* gene in the sample is determined to be hypermethylated.

[0019] In one aspect said PCR primers comprise SEQ ID NO:1 and/or SEQ ID NO: 2.

[0020] In one aspect determining the methylation status comprises, a) performing bisulfite modification to the nucleic acid in said sample, b) determining methylation of the promoter of the NMT2 gene using PCR primers and/or probes specific for the promoter region of the NMT2 gene from step a).

[0021] In one aspect step b) is performed by a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, the bisulfite modified nucleic acid from step a).

[0022] In one aspect detection of hypermethylation of the NMT2 gene is performed with primers comprising SEQ ID NO: 1 and/or SEQ ID NO: 2.

[0023] In one aspect said NMT inhibitor comprises a small molecule, an antibody, a peptide fragment, a nucleic acid, or combinations thereof.

[0024] In one aspect said small molecule comprises DDD85646, DDD86481, or a derivative thereof.

[0025] In one aspect said antibody is a monoclonal antibody or a polyclonal antibody.

[0026] In one aspect said nucleic acid comprises a dsRNA molecule, a RNAi molecule, a miRNA molecule, a ribozyme, a shRNA molecule, or a siRNA molecule.

[0027] In one aspect said cancer is lymphoma, B cell lymphoma, follicular lymphoma, diffuse large B-cell lymphoma, mantle cell lymphoma, B-CLL/SLL, immunocytoma/Waldenstrom's, MALT-type/monocytoid B 20 cell lymphoma, Burkitt's lymphoma, a pediatric lymphoma, anaplastic large cell lymphoma, acute myeloid leukemia, Blast Phase Chronic Myeloid Leukaemia, Burkitt's Lymphoma, Plasma Cell Myeloma, Intestinal Adenocarcinoma, Lung mixed Adenosquamous Carcinoma, Lung Small Cell Carcinoma, Lung, Oesophagus Squamous Cell Carcinoma, Bone, Breast Ductal Carcinoma, Stomach Diffuse Adenocarcinoma, Thyroid Medullary Carcinoma, urinary Tract Transitional Cell Carcinoma, myeloma, ovarian clear cell carcinoma, transition cell carcinoma (ureter and bladder cancer), chronic myelogenous leukemia (CML), lymphoma-CLL, breast carcinoma, colorectal adenocarcinoma, pancreas adenocarcinoma, ovarian carcinoma, non-small cell lung carcinoma, osteosarcoma, melanoma, gastric adenocarcinoma, endometrial adenocarcinoma, esophageal squamous carcinoma.

[0028] In one aspect said NMT inhibitor comprises DDD85646 or DDD86481.

[0029] In one aspect said treatment further comprises treatment with a histone acetyl transferase inhibitor, a DNA demethylase Inhibitor, and/or histone demethylase inhibitor.

[0030] In one aspect said histone acetyl transferase inhibitor is anacardic acid,
5 CPTH2, MB-3, and/or Curcumin.

[0031] In one aspect said histone demethylase inhibitor is Daminozide, GSK J1, GSK J2, GSK J4, GSK J5, GSK LSD 1 dihydrochloride, IOX 1, JIB 04, RN 1 dihydrochloride, TC-E 5002, and/or Tranylcypromine hydrochloride.

[0032] In one aspect, when said cancer is a lymphoma, the method further
10 comprising administering a treatment selected from CHOP (i.e., cyclophosphamide, doxorubicin, vincristine, and prednisone), GAP-BOP (i.e., cyclophosphamide, doxorubicin, procarbazine, bleomycin, vincristine, and prednisone), m-BACOD (i.e., methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine, dexamethasone, and leucovorin), ProMACE-MOPP (i.e., prednisone, methotrexate, doxorubicin,
15 cyclophosphamide, etoposide, leucovorin with standard MOPP), ProMACE-CytaBOM (prednisone, doxorubicin, cyclophosphamide, etoposide, cytarabine, bleomycin, vincristine, methotrexate, and leucovorin), and MACOP-B (methotrexate, doxorubicin, cyclophosphamide, vincristine, prednisone, bleomycin, and leucovorin). For relapsed aggressive non-Hodgkin's Lymphoma the following chemotherapy drug combinations may
20 be used with the compounds and compositions described herein: IMVP-16 (i.e., ifosfamide, methotrexate, and etoposide), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), DHAP (i.e., dexamethasone, - 16 high dose cytarabine, and cisplatin), ESHAP (i.e., etoposide, methylprednisone, high dosage cytarabine, and cisplatin), CEFF(B) (i.e., cyclophosphamide, etoposide, procarbazine, prednisone, and
25 bleomycin), or CAMP (i.e., lomustine, mitoxantrone, cytarabine, and prednisone).

[0033] In one aspect, when said cancer is Hodgkin's disease, the method further comprising administering a treatment selected from VABCD (i.e., vinblastine, doxorubicin, dacarbazine, lomustine and bleomycin), ABDIC (i.e., doxorubicin, bleomycin, dacarbazine, lomustine, and prednisone), CBVD (i.e., lomustine, bleomycin, vinblastine,
30 dexamethasone), PCVP (i.e., vinblastine, procarbazine, cyclophosphamide, and prednisone), CEP (i.e., lomustine, etoposide, and prednimustine), EVA (i.e., etoposide, vinblastine, and doxorubicin), MOPLACE (i.e., cyclophosphamide, etoposide, prednisone, methotrexate, cytarabine, and vincristine), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), MINE (i.e., mitoxantrone, ifosfamide, vinorelbine, and
35 etoposide), MTX-CHOP (i.e., methotrexate and CHOP), CEM (i.e., lomustine, etoposide,

and methotrexate), CEVD (i.e., lomustine, etoposide, vindesine, and dexamethasone), CAVP (i.e., lomustine, melphalan, etoposide, and prednisone), EVAP (i.e., etoposide, vinblastine, cytarabine, and cisplatin), and EPOCH (i.e., etoposide, vincristine, ; doxorubicin, cyclophosphamide, and prednisone).

- 5 **[0034]** In one aspect, when said cancer is breast cancer, the method further comprising administering a treatment selected from paclitaxel, docetaxel, nanoparticle albumin bound paclitaxel, capecitabine, doxorubicin, eribulin, vinorelbine, trastuzumab, carboplatin, cisplatin; endocrine therapies including tamoxifen, anastrozole, letrozole, exemestane, fulvestrant, and cell cycle inhibitors including palbociclib, ribociclib, LEE001, 10 and everolimus; immune checkpoint inhibiting drugs including nivolumab and pembrolizumab, either individually or in combination.

[0035] In one aspect, when said cancer is small cell lung cancer, the method further comprising administering a treatment selected from carboplatin, cisplatin, etoposide, irinotecan, either individually or in combination.

- 15 **[0036]** In one aspect, when said cancer is non-small cell lung cancer, the method further comprising administering a treatment selected from carboplatin, paclitaxel, docetaxel, gefitinib, erlotinib, afatinib, bevacizumab, ramucirumab, osimertinib, necitumumab, crizotinib, ceritinib, alectinib, and immune checkpoint inhibiting drugs including nivolumab, pembrolizumab, either individually or in combination.

- 20 **[0037]** In one aspect, when said cancer is bladder carcinoma, the method further comprising administering a treatment selected from methotrexate, vincblastine, doxorubicin, cisplatin; carboplatin and paclitaxel; docetaxel, gemcitabine and cisplatin, either individually or in combination.

- [0038]** In one aspect the determination of hypermethylation in the *NMT2* gene is 25 performed by a method selected from the group consisting of PCR, methylation-specific PCR, realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, and bisulfite sequencing.

- [0039]** In one aspect said biological sample comprises, a tissue, a cell specimen 30 or fluids includes detached tumor cells and/or free nucleic acids, a blood sample, a fractionated blood sample, a bone marrow sample, a biopsy sample, a frozen tissue sample, a fresh tissue specimen, a cell sample, and/or a paraffin embedded section.

[0040] In one aspect said subject is a human.

- [0041]** In one aspect of the present disclosure there is described an *in vitro* assay 35 for selecting a treatment for a patient with cancer, comprising: a) obtaining nucleic acid

from a sample from the subject; b) performing bisulfite modification to the nucleic acid in step a); c) performing a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, on the bisulfite modified nucleic acid from step b) using PCR primers and/or probes specific for the promoter region of the NMT2 gene, e) determining a promoter methylation level of the promoter region of NMT2 in the sample, optionally compared to a control, wherein the selected treatment is an NMT inhibitor when promoter region of *NMT2* gene in the sample is determined to be hypermethylated.

[0042] In one aspect said PCR primers comprise SEQ ID NO:1 and/or SEQ ID NO: 2.

[0043] In one aspect of the present disclosure there is described an *in vitro* assay of a sample from a patient for selecting a treatment for a patient with cancer, comprising: a) performing bisulfite modification to the nucleic acid in said sample, b) determining methylation of the promoter of the NMT2 gene using PCR primers and/or probes specific for the promoter region of the NMT2 gene from step a).

[0044] In one aspect step b) is performed by a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, the bisulfite modified nucleic acid from step a).

[0045] In one aspect detection of hypermethylation of the NMT2 gene is performed with primers comprising SEQ ID NO: 1 and/or SEQ ID NO: 2.

[0046] In one aspect the selected treatment is an NMT inhibitor when promoter region of *NMT2* gene in the sample is determined to be hypermethylated.

[0047] In one aspect said NMT inhibitor comprises a small molecule, an antibody, a peptide fragment, a nucleic acid, or combinations thereof.

[0048] In one aspect said small molecule comprises DDD85646, DDD86481, or a derivative thereof.

[0049] In one aspect said antibody is a monoclonal antibody or a polyclonal antibody.

[0050] In one aspect said nucleic acid comprises a dsRNA molecule, a RNAi molecule, a miRNA molecule, a ribozyme, a shRNA molecule, or a siRNA molecule.

[0051] In one aspect of the present disclosure there is described a kit for selecting a suitable treatment for a cancer in a cancer patient, comprising: a) one or more reagents for determining the methylation status of the *NMT2* gene from a sample of said cancer patient, wherein the selected treatment is an NMT inhibitor when promoter region of
 5 *NMT2* gene in the sample is determined to be hypermethylated, b) instructions for the use thereof.

[0052] In one aspect said reagents comprise PCR primers and/or probes specific for the promoter region of the *NMT2* gene.

[0053] In one aspect said PCR primers comprise SEQ ID NO:1 and/or SEQ ID
 10 NO: 2.

[0054] In one aspect, further comprising and NMT inhibitor.

[0055] In one aspect said NMT inhibitor comprises a small molecule, an antibody, a peptide fragment, a nucleic acid, or combinations thereof.

[0056] In one aspect said small molecule comprises DDD85646, DDD86481, or a
 15 derivative thereof.

[0057] In one aspect said antibody is a monoclonal antibody or a polyclonal antibody.

[0058] In one aspect said nucleic acid comprises a dsRNA molecule, a RNAi molecule, a miRNA molecule, a ribozyme, a shRNA molecule, or a siRNA molecule.

20 [0059] In one aspect further comprising a histone acetyl transferase inhibitor, a DNA demethylase Inhibitor, and/or histone demethylase inhibitor.

[0060] In one aspect said histone acetyl transferase inhibitor is anacardic acid, CPTH2, MB-3, and/or Curcumin.

[0061] In one aspect said histone demethylase inhibitor is Daminozide, GSK J1, GSK J2, GSK J4, GSK J5, GSK LSD 1 dihydrochloride, IOX 1, JIB 04, RN 1
 25 dihydrochloride, TC-E 5002, and/or Tranylcypromine hydrochloride.

[0062] In one aspect further comprising reagents for a treatment selected from CHOP (i.e., cyclophosphamide, doxorubicin, vincristine, and prednisone), GAP-BOP (i.e., cyclophosphamide, doxorubicin, procarbazine, bleomycin, vincristine, and prednisone),
 30 m-BACOD (i.e., methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine, dexamethasone, and leucovorin), ProMACE-MOPP (i.e., prednisone, methotrexate, doxorubicin, cyclophosphamide, etoposide, leucovorin with standard MOPP), ProMACE-CytaBOM (prednisone, doxorubicin, cyclophosphamide, etoposide, cytarabine, bleomycin, vincristine, methotrexate, and leucovorin), and MACOP-B (methotrexate,
 35 doxorubicin, cyclophosphamide, vincristine, prednisone, bleomycin, and leucovorin). For

relapsed aggressive non-Hodgkin's lymphoma the following chemotherapy drug combinations may be used with the compounds and compositions described herein: IMVP-16 (i.e., ifosfamide, methotrexate, and etoposide), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), DHAP (i.e., dexamethasone, - 16 high dose cytarabine, and cisplatin), ESHAP (i. e., etoposide, methylprednisone, high dosage cytarabine, and cisplatin), CEFF(B) (i.e., cyclophosphamide, etoposide, procarbazine, prednisone, and bleomycin), or CAMP (i.e., lomustine, mitoxantrone, cytarabine, and prednisone).

[0063] In one aspect further comprising reagents for a treatment for Hodgkin's disease selected from VABCD (i.e., vinblastine, doxorubicin, dacarbazine, lomustine and bleomycin), ABDIC (i.e., doxorubicin, bleomycin, dacarbazine, lomustine, and prednisone), CBVD (i.e., lomustine, bleomycin, vinblastine, dexamethasone), PCVP (i.e., vinblastine, procarbazine, cyclophosphamide, and prednisone), CEP (i.e., lomustine, etoposide, and prednimustine), EVA (i.e., etoposide, vinblastine, and doxorubicin), MOPLACE (i.e., cyclophosphamide, etoposide, prednisone, methotrexate, cytaravine, and vincristine), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), MINE (i.e., mitoquazone, ifosfamide, vinorelbine, and etoposide), MTX-CHOP (i.e., methotrexate and CHOP), CEM (i.e., lomustine, etoposide, and methotrexate), CEVD (i.e., lomustine, etoposide, vindesine, and dexamethasone), CAVP (i.e., lomustine, melphalan, etoposide, and prednisone), EVAP (i.e., etoposide, vinblastine, cytarabine, and cisplatin), and EPOCH (i.e., etoposide, vincristine, ; doxorubicin, cyclophosphamide, and prednisone).

[0064] In one aspect further comprising reagents for a treatment for breast cancer selected from paclitaxel, docetaxel, nanoparticle albumin bound paclitaxel, capecitabine, doxorubicin, eribulin, vinorelbine, trastuzumab, carboplatin, cisplatin; endocrine therapies including tamoxifen, anastrozole, letrozole, exemestane, fulvestrant, and cell cycle inhibitors including palbociclib, ribociclib, LEE001, and everolimus; immune checkpoint inhibiting drugs including nivolumab and pembrolizumab, either individually or in combination.

[0065] In one aspect further comprising reagents for a treatment for small cell lung cancer selected from carboplatin, cisplatin, etoposide, irinotecan, either individually or in combination.

[0066] In one aspect further comprising reagents for a treatment for non-small cell lung cancer selected from carboplatin, paclitaxel, docetaxel, gefitinib, erlotinib, afatinib, bevacizumab, ramucirumab, osimertinib, necitumumab, crizotinib, ceritinib, alectinib, and

immune checkpoint inhibiting drugs including nivolumab, pembrolizumab, either individually or in combination.

[0067] In one aspect further comprising reagents for a treatment for bladder carcinoma, the method further comprising administering a treatment selected from
5 methotrexate, vincblastine, doxorubicin, cisplatin; carboplatin and paclitaxel; docetaxel, gemcitabine and cisplatin, either individually or in combination.

[0068] In one aspect of the present disclosure there is described use of an NMT inhibitor for treating a subject with a cancer wherein the *NMT2* gene in a sample from said subject is determined to be hypermethylated.

10 [0069] In one aspect of the present disclosure there is described use of an NMT inhibitor for the manufacture of a medicament for treating a subject with a cancer wherein the *NMT2* gene in a sample from said subject is determined to be hypermethylated.

[0070] In one aspect determining the methylation status comprises, a) performing bisulfite modification to the nucleic acid in said sample, b) determining methylation of the
15 promoter of the *NMT2* gene using PCR primers and/or probes specific for the promoter region of the *NMT2* gene from step a).

[0071] In one aspect step b) is performed by a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein,
20 quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, the bisulfite modified nucleic acid from step a).

[0072] In one aspect detection of hypermethylation of the *NMT2* gene is performed with primers comprising SEQ ID NO: 1 and/or SEQ ID NO: 2.

[0073] In one aspect said NMT inhibitor comprises a small molecule, an antibody,
25 a peptide fragment, a nucleic acid, or combinations thereof.

[0074] In one aspect said small molecule comprises DDD85646, DDD86481, or a derivative thereof.

[0075] In one aspect said antibody is a monoclonal antibody or a polyclonal antibody.

30 [0076] In one aspect said nucleic acid comprises a dsRNA molecule, a RNAi molecule, a miRNA molecule, a ribozyme, a shRNA molecule, or a siRNA molecule.

[0077] In one aspect said cancer is lymphoma, B cell lymphoma, follicular lymphoma, diffuse large B-cell lymphoma, mantle cell lymphoma, B-CLL/SLL, immunocytoma/Waldenstrom's, MALT-type/monocytoid B 20 cell lymphoma, Burkitt's
35 lymphoma, a pediatric lymphoma, anaplastic large cell lymphoma, acute myeloid

- leukemia, Blast Phase Chronic Myeloid Leukaemia, Burkitt's Lymphoma, Plasma Cell Myeloma, Intestinal Adenocarcinoma, Lung mixed Adenosquamous Carcinoma, Lung Small Cell Carcinoma, Lung, Oesophagus Squamous Cell Carcinoma, Bone, Breast Ductal Carcinoma, Stomach Diffuse Adenocarcinoma, Thyroid Medullary Carcinoma,
- 5 urinary Tract Transitional Cell Carcinoma, myeloma, ovarian clear cell carcinoma, transition cell carcinoma (ureter and bladder cancer), chronic myelogenous leukemia (CML), lymphoma-CLL, breast carcinoma, colorectal adenocarcinoma, pancreas adenocarcinoma, ovarian carcinoma, non-small cell lung carcinoma, osteosarcoma, melanoma, gastric adenocarcinoma, endometrial adenocarcinoma, esophageal
- 10 squamous carcinoma.
- [0078]** In one aspect said NMT inhibitor comprises DDD85646 or DDD86481.
- [0079]** In one aspect further comprising use of a histone acetyl transferase inhibitor, a DNA demethylase Inhibitor, and/or histone demethylase inhibitor.
- [0080]** In one aspect said histone acetyl transferase inhibitor is anacardic acid,
- 15 CPTH2, MB-3, and/or Curcumin.
- [0081]** In one aspect said histone demethylase inhibitor is Daminozide, GSK J1, GSK J2, GSK J4, GSK J5, GSK LSD 1 dihydrochloride, IOX 1, JIB 04, RN 1 dihydrochloride, TC-E 5002, and/or Tranylcypromine hydrochloride.
- [0082]** In one aspect, wherein when said cancer is a lymphoma, the use further
- 20 comprising use of a treatment selected from CHOP (i.e., cyclophosphamide, doxorubicin, vincristine, and prednisone), GAP-BOP (i.e., cyclophosphamide, doxorubicin, procarbazine, bleomycin, vincristine, and prednisone), m-BACOD (i.e., methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine, dexamethasone, and leucovorin), ProMACE-MOPP (i.e., prednisone, methotrexate, doxorubicin, cyclophosphamide,
- 25 etoposide, leucovorin with standard MOPP), ProMACE-CytaBOM (prednisone, doxorubicin, cyclophosphamide, etoposide, cytarabine, bleomycin, vincristine, methotrexate, and leucovorin), and MACOP-B (methotrexate, doxorubicin, cyclophosphamide, vincristine, prednisone, bleomycin, and leucovorin). For relapsed aggressive non-Hodgkin's lymphoma the following chemotherapy drug combinations may
- 30 be used with the compounds and compositions described herein: IMVP-16 (i.e., ifosfamide, methotrexate, and etoposide), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), DHAP (i.e., dexamethasone, - 16 high dose cytarabine, and cisplatin), ESHAP (i. e., etoposide, methylprednisone, high dosage cytarabine, and cisplatin), CEFF(B) (i.e., cyclophosphamide, etoposide, procarbazine, prednisone, and
- 35 bleomycin), or CAMP (i.e., lomustine, mitoxantrone, cytarabine, and prednisone).

- [0083]** In one aspect, wherein when said cancer is Hodgkin's disease, the use further comprising use of a treatment selected from VABCD (i.e., vinblastine, doxorubicin, dacarbazine, lomustine and bleomycin), ABDIC (i.e., doxorubicin, bleomycin, dacarbazine, lomustine, and prednisone), CBVD (i.e., lomustine, bleomycin, vinblastine, dexamethasone), PCVP (i.e., vinblastine, procarbazine, cyclophosphamide, and prednisone), CEP (i.e., lomustine, etoposide, and prednimustine), EVA (i.e., etoposide, vinblastine, and doxorubicin), MOPLACE (i.e., cyclophosphamide, etoposide, prednisone, methotrexate, cytarabine, and vincristine), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), MINE (i.e., mitomycin, ifosfamide, vinorelbine, and etoposide), MTX-CHOP (i.e., methotrexate and CHOP), CEM (i.e., lomustine, etoposide, and methotrexate), CEVD (i.e., lomustine, etoposide, vindesine, and dexamethasone), CAVP (i.e., lomustine, melphalan, etoposide, and prednisone), EVAP (i.e., etoposide, vinblastine, cytarabine, and cisplatin), and EPOCH (i.e., etoposide, vincristine, ; doxorubicin, cyclophosphamide, and prednisone).
- [0084]** In one aspect, wherein when said cancer is breast cancer, the use further comprising use of a treatment selected from paclitaxel, docetaxel, nanoparticle albumin bound paclitaxel, capecitabine, doxorubicin, eribulin, vinorelbine, trastuzumab, carboplatin, cisplatin; endocrine therapies including tamoxifen, anastrozole, letrozole, exemestane, fulvestrant, and cell cycle inhibitors including palbociclib, ribociclib, LEE001, and everolimus; immune checkpoint inhibiting drugs including nivolumab and pembrolizumab, either individually or in combination.
- [0085]** In one aspect, wherein when said cancer is small cell lung cancer, the use further comprising a use of a treatment selected from carboplatin, cisplatin, etoposide, irinotecan, either individually or in combination.
- [0086]** In one aspect, wherein when said cancer is non-small cell lung cancer, the use further comprising use of a treatment selected from carboplatin, paclitaxel, docetaxel, gefitinib, erlotinib, afatinib, bevacizumab, ramucirumab, osimertinib, necitumumab, crizotinib, ceritinib, alectinib, and immune checkpoint inhibiting drugs including nivolumab, pembrolizumab, either individually or in combination.
- [0087]** In one aspect, wherein when said cancer is bladder carcinoma, the use further comprising a use of a treatment selected from methotrexate, vincblastine, doxorubicin, cisplatin; carboplatin and paclitaxel; docetaxel, gemcitabine and cisplatin, either individually or in combination.
- [0088]** In one aspect, wherein the determination of hypermethylation in the *NMT2* gene is performed by a method selected from the group consisting of PCR, methylation-

specific PCR, realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, and bisulfite sequencing.

[0089] In one aspect, wherein said biological sample comprises, a tissue, a cell
 5 specimen or fluids includes detached tumor cells and/or free nucleic acids, a blood sample, a fractionated blood sample, a bone marrow sample, a biopsy sample, a frozen tissue sample, a fresh tissue specimen, a cell sample, and/or a paraffin embedded section.

[0090] In one aspect, wherein said subject is a human.

10 [0091] Other aspects and features of the present disclosure will become apparent to those ordinarily skilled in the art upon review of the following description of specific embodiments in conjunction with the accompanying figures.

BRIEF DESCRIPTION OF THE DRAWINGS

15 [0092] Embodiments of the present disclosure will now be described, by way of example only, with reference to the attached Figures.

[0093] Fig. 1 depicts NMT2 expression is decreased in several cancer cell lines in the CCLE database and in tumours from TCGA database. *NMT2* (grey) has a broader range of mRNA expression than *NMT1* (black) in cancer cell lines (Panel A) and tumours (Panel B). Box plot of NMT2 mRNA expression in cancer-specific cell lines and tumour types. ***, $p < 0.0001$ vs. all tumours by t test, #, data not available (Panels C, D, E). NMT2 mRNA copies determined by droplet digital PCR in various lymphocytic cell lines (Panel F). Specific NMT activity in the cell lines in (Panel G) *, $p < 0.05$. Kaplan-Meier plots show the progression-free survival (PFS) of 470 DLBCL patients (GSE31312) with
 20 high (red; top curve) versus low (blue; bottom curve) NMT2 expression (Panel H). P value was determined by log-rank test. All values are mean $\pm$ s.e.m. of 3 independent experiments;

[0094] Fig. 2 depicts a CpG island at the NMT2 promoter (Panel A). Epigenetic regulation of *NMT2* in lymphoma cell lines and tumours (Panel B) NMT2 expression
 30 (RSEM normalized counts) in TCGA tumours correlates negatively with NMT2 methylation values (cg02268561 beta values) at position 15,212,006 in WSU-DLCL2 (highlighted in red in Panel B). $p = 0.0082$, Spearman correlation analysis, $n = 43$ from the TCGA cohort). Bisulfite sequencing of DAPK1 promoter region reveals a highly methylated CpG containing promoter region only in malignant B lymphocytes (Panel C);

- [0095]** Fig. 3 depicts the number of NMT2 mRNA copies measured by digital PCR in various lymphocytic cell lines after a 24-h treatments with DAC in the presence or absence of SAHA (Panels A-D). NMT2 protein levels assessed by western blotting in BL2 (Panel E) and IM9 (Panel F) (normalized to GAPDH level) after 24-h treatments as above. Values are mean $\pm$ s.e.m. of 3 independent experiments. (Panel G) NMT2 expression (RSEM normalized counts) in TCGA tumours correlates negatively with NMT2 methylation values (cg02268561 beta values) at position 15,212,006 in WSU-DLCL2 (highlighted in red in Figure 2B). $p = 0.0082$, Spearman correlation analysis, $n = 43$ from the TCGA cohort. depicts graphs and immunoblots;
- 10 **[0096]** Fig. 4 shows a box and whiskers plot showing the distribution of individual mice tumor volumes in each group graphs (Panel A). PCLX-001 dose-response curves for xenografts derived from cell lines DOHH2 (Panel B); Combination therapy PCLX-001 and DOX dose-response curves for xenografts derived from cell lines DOHH2 (Panel C); and
- 15 **[0097]** Fig 5 PCLX-001 dose-response curves for xenografts derived from cell line BL2 (Panel A). Combination therapy PCLX-001 and DOX dose-response curves for xenografts derived from cell line BL2 (Panel B). Total NMT specific activity in BL2 tumour sample in Panels A (Panel C). Patient selection strategy for precision medicine (Panel D). Identification of NMT2-deficiency in DLBCLs from three patients by
- 20 immunohistochemistry (IHC) and RNA in situ hybridization (ISH). Tumour from patient DLBCL3 is also shown after engraftment in mice (Panel E). PCLX-001 dose-response curves for the xenograft derived from patient 3 (Panel F). Representative tumours from mice with patient-derived xenografts (PDX) (Panel G). PCLX-001 treatment of mice with NMT2-deficient DLBCL3 PDX induces apoptosis (increase in cleaved caspase-3 staining)
- 25 (Panel G) and reduces cell proliferation (reduction of Ki-67 staining) (Panel H). Shown are representative stained tumour slices for each treatment (Panel I).

DETAILED DESCRIPTION

- [0098]** As will be described in more detail below, there is described herein compounds, composition and methods for the treatment of a subject with cancer. There are also described here methods for identifying subject with cancer that are suitable for treatment with the compounds, composition and methods are described herein.

- [0099]** Cells need at least one NMT to survive. The finding that multiple cancers lack one of two NMTs, while stromal and normal tissues do not, enables the treatment of NMT2-deficient cancer cells with an NMT inhibitor. See WO2013/013302 and

WO2014/067002, the entire contents of which are hereby incorporated by reference in their entirety.

[00100] It is shown herein that NMT2 expression is reduced or eliminated in certain cancers, and in one example lymphomas, via an epigenetic mechanism(s). Reduction or
5 elimination of NMT2 expression renders the cancer sensitive inhibitors of NMT.

[00101] In some examples, determination of the methylation status of *NMT2* gene may be used to identify cancers which will respond to treatment with NMT inhibitors.

[00102] **Cancers**

[00103] The term "cancer", as used herein, refers to a variety of conditions caused
10 by the abnormal, uncontrolled growth of cells. Cells capable of causing cancer, referred to as "cancer cells", possess characteristic properties such as uncontrolled proliferation, immortality, metastatic potential, rapid growth and proliferation rate, and/or certain typical morphological features. Cancer cells may be in the form of a tumour, but such cells may also exist alone within a subject, or may be a non-tumorigenic cancer cell. A cancer can
15 be detected in any of a number of ways, including, but not limited to, detecting the presence of a tumor or tumors (e.g., by clinical or radiological means), examining cells within a tumor or from another biological sample (e.g., from a tissue biopsy), measuring blood markers indicative of cancer, and detecting a genotype indicative of a cancer. However, a negative result in one or more of the above detection methods does not
20 necessarily indicate the absence of cancer, e.g., a patient who has exhibited a complete response to a cancer treatment may still have a cancer, as evidenced by a subsequent relapse.

[00104] In a specific example of the present disclosure, the cancer is lymphoma.

[00105] The term "lymphoma" as used herein refers to a malignant growth of B or T
25 cells in the lymphatic system. "Lymphoma" includes numerous types of malignant growths, including Hodgkin's Lymphoma and non-Hodgkin's lymphoma. The term "non-Hodgkin's Lymphoma" as used herein, refers to a malignant growth of B or T cells in the lymphatic system that is not a Hodgkin's Lymphoma (which is characterized, e.g., by the presence of Reed-Sternberg cells in the cancerous area). Non-Hodgkin's lymphomas
30 encompass over 29 types of lymphoma, the distinctions between which are based on the type of cancer cells.

[00106] In a more specific example of the present disclosure, the cancer is a B-lymphoma.

[00107] Thus, in one embodiment, the compounds, compositions and methods of
35 the disclosure are suitable for the treatment of a subject with B cell lymphoma.

[00108] Examples of B-cell lymphomas include, but are not limited to, for example, follicular lymphoma, diffuse large B-cell lymphoma, mantle cell lymphoma, B-CLL/SLL, immunocytoma/Waldenstrom's, and MALT-type/monocytoid B cell lymphoma. Also contemplated are the treatment of pediatric lymphomas such as Burkitt's lymphoma,
 5 diffuse large B-cell lymphoma, follicular lymphoma, precursor B-LBL, precursor T-LBL, and anaplastic large cell lymphoma.

[00109] In other embodiments, the cancer is lymphoma, B cell lymphoma, follicular lymphoma, diffuse large B-cell lymphoma, mantle cell lymphoma, B-CLL/SLL, immunocytoma/Waldenstrom's, MALT-type/monocytoid B cell lymphoma, Burkitt's
 10 lymphoma, a pediatric lymphoma, anaplastic large cell lymphoma, acute myeloid leukemia, Blast Phase Chronic Myeloid Leukaemia, Burkitt's Lymphoma, Plasma Cell Myeloma, Intestinal Adenocarcinoma, Lung mixed Adenosquamous Carcinoma, Lung Small Cell Carcinoma, Lung, Oesophagus Squamous Cell Carcinoma, Bone, Breast Ductal Carcinoma, Stomach Diffuse Adenocarcinoma, Thyroid Medullary Carcinoma,
 15 urinary Tract Transitional Cell Carcinoma, myeloma, ovarian clear cell carcinoma, transition cell carcinoma (ureter and bladder cancer), chronic myelogenous leukemia (CML), lymphoma-CLL, breast carcinoma, colorectal adenocarcinoma, pancreas adenocarcinoma, ovarian carcinoma, non-small cell lung carcinoma, osteosarcoma, melanoma, gastric adenocarcinoma, endometrial adenocarcinoma, or esophageal
 20 squamous carcinoma.

[00110] The term "subject" or "patient" as used herein, refers to an animal, and can include, for example, domesticated animals, such as cats, dogs, etc., livestock (e.g., cattle, horses, pigs, sheep, goats, etc.), laboratory animals (e.g., mouse, rabbit, rat, guinea pig, etc.), mammals, non-human mammals, primates, non-human primates,
 25 rodents, birds, reptiles, amphibians, fish, and any other animal. In a specific example, the subject is a human.

[00111] The term "treatment" or "treat" as used herein, refers to obtaining beneficial or desired results, including clinical results. Beneficial or desired clinical results can include, but are not limited to, alleviation or amelioration of one or more symptoms or
 30 conditions, diminishment of extent of disease, stabilized (i.e. not worsening) state of disease, preventing spread of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, diminishment of the reoccurrence of disease, and remission (whether partial or total), whether detectable or undetectable. "Treating" and "Treatment" can also mean prolonging survival as compared to expected
 35 survival if not receiving treatment. "Treating" and "treatment" as used herein also include

prophylactic treatment. For example, a subject with early cancer, for example an early stage lymphoma, can be treated to prevent progression or alternatively a subject in remission can be treated with a compound or composition described herein to prevent recurrence.

- 5 [00112] The term "sample" or "biological sample" as used herein refers to any sample from a subject, including but not limited to a fluid, cell or tissue sample that comprises cancer cells, or which is suspected of containing cancer cells, which can be assayed for epigenetic modification, gene expression levels, protein levels, enzymatic activity levels, and the like. In a specific example, the sample is assayed for epigenetic
10 modification. In another specific example, the epigenetic modification is methylation or acetylation. The sample may include, for example, a blood sample, a fractionated blood sample, a bone marrow sample, a biopsy sample, a frozen tissue sample, a fresh tissue specimen, a cell sample, and/or a paraffin embedded section, material from which DNA can be extracted in sufficient quantities and with adequate quality to permit measurement
15 of epigenetic modification(s).

[00113] **Inhibitors of NMT**

[00114] In a specific aspect, said NMT inhibitor comprises a small molecule, an antibody, a peptide fragment, a nucleic acid, or combinations thereof.

- [00115] In a specific aspect, said small molecule comprises DDD85646, the
20 pyrazole sulphonamide inhibitor of *T. brucei* NMT [J.A.Freearson et al (2010) Nature. 464.728-723)], or a derivative thereof.

[00116] In a specific aspect, said small molecule comprises Trs-DBA.

[00117] In a specific aspect, said small molecule comprises DDD86481 (also referred to as PCLX-001 and CCI-002 herein), or a derivative thereof.

- 25 [00118] DDD85646 and DDD86481 are shown in Table 1.

Table 1	DDD00085646	DDD00086481
Structure		

[00119] In a specific aspect, said antibody is a monoclonal antibody or a polyclonal antibody.

[00120] In a specific aspect, said nucleic acid comprises a dsRNA molecule, a RNAi molecule, a miRNA molecule, a ribozyme, a shRNA molecule, or a siRNA molecule.

[00121] Peptide fragments may be prepared wholly or partly by chemical synthesis that fit in the active site of NMT1. Peptide fragments can be prepared according to
5 established, standard liquid or solid-phase peptide synthesis methods, which will be known to the skilled worker.

[00122] Nucleic acid inhibitors, or the complements thereof, inhibit activity or function by down-regulating production of active polypeptide. This can be monitored using conventional methods well known in the art, for example by screening using real time
10 PCR as described in the examples.

[00123] Examples of nucleic acid inhibitors include anti-sense or RNAi technology, the use of which is to down-regulate gene expression is well established in the art. Anti-sense oligonucleotides may be designed to hybridize to the complementary sequence of nucleic acid, pre-mRNA or mature mRNA, interfering with the production of the base
15 excision repair pathway component so that its expression is reduced or completely or substantially completely prevented. In addition to targeting coding sequence, anti-sense techniques may be used to target control sequences of a gene, e.g. in the 5' flanking sequence, whereby the anti-sense oligonucleotides can interfere with expression control sequences.

[00124] An alternative to anti-sense is to use a copy of all or part of the target gene inserted in sense, that is the same, orientation as the target gene, to achieve reduction in expression of the target gene by co-suppression.

[00125] Additionally, double stranded RNA (dsRNA) silencing may be used. dsRNA mediated silencing is gene specific and is often termed RNA interference (RNAi).

[00126] In another example, nucleic acid is used which on transcription produces a ribozyme, able to cut nucleic acid at a specific site and therefore also useful in influencing
25 NMT.

[00127] In yet another example, small RNA molecules may be employed to regulate gene expression. These include targeted degradation of mRNAs by small
30 interfering RNAs (siRNAs), post transcriptional gene silencing (PTGs), developmentally regulated sequence-specific translational repression of mRNA by micro-RNAs (miRNAs) and targeted transcriptional gene silencing.

[00128] In yet another example, the expression of a short hairpin RNA molecule (shRNA) in the cell may be used. A shRNA consists of short inverted repeats separated
35 by a small loop sequence. One inverted repeat is complimentary to the gene target. In the

cell the shRNA is processed by DICER into a siRNA which degrades the target NMT gene mRNA and suppresses expression. In a preferred embodiment the shRNA is produced endogenously (within a cell) by transcription from a vector.

[00129] The term "inhibit" or "inhibitor" as used herein, refers to any method or technique which inhibits protein synthesis, levels, activity, or function, as well as methods of inhibiting the induction or stimulation of synthesis, levels, activity, or function of the protein of interest, for example NMT1. The term also refers to any metabolic or regulatory pathway, which can regulate the synthesis, levels, activity, or function of the protein of interest. The term includes binding with other molecules and complex formation. Therefore, the term "inhibitor" refers to any agent or compound, the application of which results in the inhibition of protein function or protein pathway function. However, the term does not imply that each and every one of these functions must be inhibited at the same time.

[00130] Administration/pharmaceutical compositions

[00131] In other examples, the compounds and/or compositions are provided in a pharmaceutically effect amount suitable for administration to a subject.

[00132] The term "pharmaceutically effective amount" as used herein refers to the amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by a researcher or clinician. This amount can be a therapeutically effective amount.

[00133] The compounds and compositions are provided in a pharmaceutically acceptable form.

[00134] The term "pharmaceutically acceptable" as used herein includes compounds, materials, compositions, and/or dosage forms (such as unit dosages) which are suitable for use in contact with the tissues of a subject without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. Each carrier, excipient, etc. is also be "acceptable" in the sense of being compatible with the other ingredients of the formulation.

[00135] The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, e.g. decisions on dosage etc., is within the responsibility of general practitioners and other medical doctors, and typically takes account of the disorder to be treated, the condition of the individual patient, the site of delivery, the method of administration and other factors known to practitioners.

[00136] A compound or composition may be administered alone or in combination with other treatments, either simultaneously or sequentially, dependent upon the condition to be treated.

[00137] The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. Such methods include the step of bringing the active compound into association with a carrier, which may constitute one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active compound with liquid carriers or finely divided solid carriers or both, and then if necessary shaping the product.

[00138] The compounds and compositions may be administered to a subject by any convenient route of administration, whether systemically/peripherally or at the site of desired action, including but not limited to, oral (e.g. by ingestion); topical (including e.g. transdermal, intranasal, ocular, buccal, and sublingual); pulmonary (e.g. by inhalation or insufflation therapy using, e.g. an aerosol, e.g. through mouth or nose); rectal; vaginal; parenteral, for example, by injection, including subcutaneous, intradermal, intramuscular, intravenous, intraarterial, intracardiac, intrathecal, intraspinal, intracapsular, subcapsular, intraorbital, intraperitoneal, intratracheal, subcuticular, intraarticular, subarachnoid, and intrasternal; by implant of a depot / for example, subcutaneously or intramuscularly.

[00139] Formulations suitable for oral administration (e.g., by ingestion) may be presented as discrete units such as capsules, cachets or tablets, each containing a predetermined amount of the active compound; as a powder or granules; as a solution or suspension in an aqueous or non-aqueous liquid; or as an oil-in- water liquid emulsion or a water- in-oil liquid emulsion; as a bolus; as an electuary; or as a paste.

[00140] Formulations suitable for parenteral administration (e.g., by injection, including cutaneous, subcutaneous, intramuscular, intravenous and intradermal), include aqueous and non-aqueous isotonic, pyrogen-free, sterile injection solutions which may contain anti-oxidants, buffers, preservatives, stabilisers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non- aqueous sterile suspensions which may include suspending agents and thickening agents, and liposomes or other microparticulate systems which are designed to target the compound to blood components or one or more organs. Examples of suitable isotonic vehicles for use in such formulations include Sodium Chloride Injection, Ringer's Solution, or Lactated Ringer's Injection.

[00141] The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and
5 suspensions may be prepared from sterile powders, granules, and tablets. Formulations may be in the form of liposomes or other microparticulate systems which are designed to target the active compound to blood components or one or more organs.

[00142] Compositions comprising compounds disclosed herein may be used in the methods described herein in combination with standard chemotherapeutic regimes or in
10 conjunction with radiotherapy.

[00143] **Combination therapy**

[00144] In the case of cancer, and for example lymphoma, in a patient, known treatments are dependent upon the subject being treated, the type of disease, and its stage. Existing treatment modalities for various cancer, including for example for
15 lymphoma, are known to the skilled worker. Accordingly, known treatments for cancer(s) may be used together with the NMT inhibitors disclosed herein.

[00145] Common drug combinations for use in treating lymphomas include, but are not limited, to CHOP (i.e., cyclophosphamide, doxorubicin, vincristine, and prednisone), GAP-BOP (i.e., cyclophosphamide, doxorubicin, procarbazine, bleomycin, vincristine, and
20 prednisone), m-BACOD (i.e., methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine, dexamethasone, and leucovorin), ProMACE-MOPP (i.e., prednisone, methotrexate, doxorubicin, cyclophosphamide, etoposide, leucovorin with standard MOPP), ProMACE-CytaBOM (prednisone, doxorubicin, cyclophosphamide, etoposide, cytarabine, bleomycin, vincristine, methotrexate, and leucovorin), and MACOP-B
25 (methotrexate, doxorubicin, cyclophosphamide, vincristine, prednisone, bleomycin, and leucovorin). For relapsed aggressive non-Hodgkin's lymphoma the following chemotherapy drug combinations may be used with the compounds and compositions described herein: IMVP-16 (i.e., ifosfamide, methotrexate, and etoposide), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), DHAP (i.e., dexamethasone, - 16
30 high dose cytarabine, and cisplatin), ESHAP (i. e., etoposide, methylprednisone, high dosage cytarabine, and cisplatin), CEFF(B) (i.e., cyclophosphamide, etoposide, procarbazine, prednisone, and bleomycin), and CAMP (i.e., lomustine, mitoxantrone, cytarabine, and prednisone).

[00146] Treatment for salvage chemotherapy used for certain lymphomas such as
35 for relapsed, resistant Hodgkin's Disease include but are not limited to VABCD (i.e.,

vinblastine, doxorubicin, dacarbazine, lomustine and bleomycin), ABDIC (i.e., doxorubicin, bleomycin, dacarbazine, lomustine, and prednisone), CBVD (i.e., lomustine, bleomycin, vinblastine, dexamethasone), PCVP (i.e., vinblastine, procarbazine, cyclophosphamide, and prednisone), CEP (i.e., lomustine, etoposide, and prednimustine), EVA (i.e., etoposide, vinblastine, and doxorubicin), MOPLACE (i.e., cyclophosphamide, etoposide, prednisone, methotrexate, cytarabine, and vincristine), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), MINE (i.e., mitozantrone, ifosfamide, vinorelbine, and etoposide), MTX-CHOP (i.e., methotrexate and CHOP), CEM (i.e., lomustine, etoposide, and methotrexate), CEVD (i.e., lomustine, etoposide, vindesine, and dexamethasone), CAVP (i.e., lomustine, melphalan, etoposide, and prednisone), EVAP (i.e., etoposide, vinblastine, cytarabine, and cisplatin), and EPOCH (i.e., etoposide, vincristine, ; doxorubicin, cyclophosphamide, and prednisone).

[00147] Treatment for breast cancer includes treatment with paclitaxel, docetaxel, nanoparticle albumin bound paclitaxel, capecitabine, doxorubicin, eribulin, vinorelbine, trastuzumab, carboplatin, cisplatin; endocrine therapies including tamoxifen, anastrozole, letrozole, exemestane, fulvestrant, and cell cycle inhibitors including palbociclib, ribociclib, LEE001, and everolimus; immune checkpoint inhibiting drugs including nivolumab and pembrolizumab, either individually or in combination.

[00148] Treatment for small cell lung cancer includes treatment with carboplatin, cisplatin, etoposide, irinotecan, either individually or in combination.

[00149] Treatment for non-small cell lung cancer includes treatment with carboplatin, paclitaxel, docetaxel, gefitinib, erlotinib, afatinib, bevacizumab, ramucirumab, osimertinib, necitumumab, crizotinib, ceritinib, alectinib, and immune checkpoint inhibiting drugs including nivolumab, pembrolizumab, either individually or in combination.

[00150] Treatment for bladder carcinoma includes treatment with methotrexate, vinblastine, doxorubicin, cisplatin; carboplatin and paclitaxel; docetaxel, gemcitabine and cisplatin, either individually or in combination.

[00151] Epigenetic mechanisms

[00152] As mentioned above, it is shown herein that *NMT2* expression (i.e., *NMT2* gene expression) is reduced or eliminated in certain cancers, and in one example lymphomas, via an epigenetic mechanism(s). Reduction or elimination of *NMT2* expression renders the cancer sensitive inhibitors of NMT.

[00153] The term "epigenetic" as used herein refers to modification(s) in gene expression that is independent of DNA sequence.

[00154] Epigenetic factors include modifications in gene expression that are controlled by changes in DNA methylation (hypermethylation or hypomethylation), and chromatin structure.

[00155] For example, DNA methylation patterns are known to correlate with gene expression. DNA methylation occurs at the 5-position of cytosines at CpG sites.

[00156] In other example, epigenetic changes are altered acetylation patterns or levels.

[00157] **Epigenetic mechanisms- methylation**

[00158] In some examples herein, determination of the methylation status of *NMT2* gene may be used to identify cancers, which will respond to treatment with NMT inhibitors.

[00159] The term "methylation status" as used herein, refers to the level of methylation of cytosine residues in CpG sites in a gene, for example *NMT2*. CpG sites include CpG pairs, CpG islands, and CpG island shores.

[00160] The term "CpG island" refers to a genomic DNA region that contains a high percentage of CpG sites relative to the average genomic CpG incidence

[00161] In the example of CpG pairs, the methylation status may be methylated or unmethylated. In the example of CpG islands or CpG island shores, or to any stretch of residues, the methylation status refers to the level of methylation, which is the relative or absolute concentration of methylated C at the particular CpG island or CpG island shore or stretch of residues in a biological sample.

[00162] Methylation of a CpG sites or CpG island at a promoter typically reduces or eliminates expression of the gene.

[00163] CpG sites or CpG islands may surround the 5' region of the coding region of the gene as well as the 3' region of the coding region. Accordingly, CpG sites or CpG islands can be found in multiple regions of a nucleic acid sequence including upstream of coding sequences in a regulatory region including a promoter region, in the coding regions (e.g., exons), downstream of coding regions in, for example, enhancer regions, and in introns. All of these regions can be assessed to determine their methylation status, as appropriate.

[00164] The levels of methylation of the *NMT2* gene, for example hypermethylation of *NMT2* or hypomethylation of *NMT2*, may be determined by any suitable means in order to reflect the expression level of *NMT2*.

[00165] The term "expression level" as used herein refers to the level of gene expression.

[00166] The term "gene" as used herein refers to a nucleic acid (for example DNA) sequence that comprises coding sequences necessary for the production of a polypeptide such as NMT2, RNA (e.g., including but not limited to, mRNA, tRNA and rRNA) or precursor. The polypeptide, RNA, or precursor may be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired activity or functional properties (e.g., enzymatic activity, ligand binding, signal transduction, etc.) of the full-length or fragment are retained. The term also encompasses the coding region of a structural gene and the including sequences located adjacent to the coding region on both the 5' and 3' ends for a distance of about 1 kb, and in some example greater than 1 kb, on either end such that the gene corresponds to the length of the full-length mRNA.

[00167] The term "hypermethylation" as used herein refers to the average methylation status corresponding to an increased presence of 5-methyl cytidine at one or a more of CpG CpG site or CpG island within a DNA sequence, for example the *NMT2* gene, of a test DNA sample, relative to the amount of 5-methyl cytidine at corresponding CpG site or CpG island within a "normal" and/or control DNA sample.

[00168] In some example, hypermethylation refers to an of about 1%, about 5%, about 10%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, or more than about 95% greater than the extent of methylation of cytosines typically expected or observed for the CpG site or sites being evaluated. It will be appreciated that methylation varies under different metabolic conditions, stress, etc, as would be know to a skilled worker. In some example, hypermethylation refers to methylation at a single cytidine.

[00169] The term "hypomethylation" as used herein refers to the average methylation status corresponding to a decreased presence of 5-methyl cytidine at one or a more of CpG CpG site or CpG island within a DNA sequence, for example the *NMT2* gene, of a test DNA sample, relative to the amount of 5-methyl cytidine at corresponding CpG site or CpG island within a "normal" and/or control DNA sample.

[00170] Typically CpG motifs reside near the transcription start site, for example, within about 3 kbp, within about 2.5 kbp, within about 2 kbp, within about 1.5 kbp, within about 1 kbp, within about 750 bp, or within about 500 bp

[00171] In some examples, CpG islands are less than 100 base pairs, between 100 and 200 base pairs, between 200 and 300 base pairs, between 300 and 500 base pairs, between 500 and 750 base pairs; between 750 and 1000 base pair; 100 or more base pairs in length.

[00172] In some examples, the methylation status of the *NMT2* gene is determined to be hypermethylation. In some examples, the methylation status of *NMT2* is determined to be hypomethylation.

[00173] In the example in which the methylation of the *NMT2* gene in a sample is determined to be hypermethylation, *NMT2* expression is reduced or eliminated.

[00174] In the example in which the methylation of the *NMT2* gene in a sample is determined to be hypomethylation, *NMT2* expression is not reduced or eliminated.

[00175] The term "reduced expression" as used herein refers to a lesser expression level of a gene in a biological sample, compared to expression in a control or reference sample.

[00176] The methylation status of the *NMT2* gene in a sample may be determined using a wide range of suitable devices and methods of methylation quantitation or detection, as would be known to the skilled worker.

[00177] Suitable devices include, but are not limited to lab-on-chip technology, microfluidic technologies, biomonitor technology, proton recognition technologies (e.g., Ion Torrent), and/or other highly parallel and/or deep sequencing methods.

[00178] Suitable methods of DNA methylation analysis include, but are not limited to, MALDI-TOFF, MassARRAY, Methy Light, Quantitative analysis of ethylated alleles (QAMA), enzymatic regional methylation assay (ERMA), HeavyMethyl, QBSUPT, MS-SNuPE, MethylQuant, Quantitative PCR sequencing, and Oligonucleotide-based microarray systems.

[00179] Further suitable method of DNA methylation analysis include, but are not limited to, methylation-specific PCR (MSP), bisulfite sequencing (also referred to as a sequencing of DNA treated with bisulfite to convert Methylated Cytosine to Uracil), pyrosequencing, methylation- sensitive DNA restriction enzyme analysis, quantitative bisulfite pyrosequencing, bisulfite genomic sequencing PCR and AQAMA.

[00180] In some examples, determining of the methylation status of the *NMT2* gene comprises determining the methylation frequency of the CpG island or CpG island shore. Determining of the level of a nucleic acid polymer with altered methylation is carried out in a variety of ways, and as noted above, including but not limited to, methylation-specific PCR, quantitative methylation-specific PCR, methylation- sensitive DNA restriction enzyme analysis, quantitative bisulfite pyrosequencing, next generation sequencing and bisulfite genomic sequencing PCR.

[00181] In some examples, methylation-sensitive restriction endonucleases may be used to detect methylated CpG dinucleotidemotifs. Such endonucleases may either

preferentially cleave methylated recognition sites relative to non-methylated recognition sites. Non limiting examples include are Aat II, Ace III, Ad I, Acl I, Age I, Alu I, Asc I, Ase I, AsiS I, Ban I, Bbe I, Bsa I, BsaH I, BsiE I, BsiWI, BsrVI, BssK 1, BstB I, BstN I, Bs I, Cla I, Eae I, Eag I, Fau I, FseI, Hha I, mPI I, Hinc II, Hpa 11, Npy99 I, HpyCAIV, Kas I, Mbo I, Mlu I, MapA 11, Msp I, Nae I, Nar I, Not 1, Pml I, PstI, Pvu I, Rsr II, Sac II, Sap I, Sau3A I, SfiI, Sfo I, SgrA I, Sma I, SnaB I, Tsc I, Xma I, and Zra I. In some example, endonucleases preferentially cleave non-methylated relative to methylated recognition sites. Non limiting examples include Ace II, Ava I, BssH II, BstU I, Hpa II, Not I, and Mho I.

[00182] In some examples, chemical reagents can be used that selectively modify either the methylated or non-methylated form of CpG dinucleotide motifs. Modified products can be detected directly, or after a further reaction which creates products that are easily distinguishable. Means which detect altered size and/or charge can be used to detect modified products, including but not limited to electrophoresis, chromatography, and mass spectrometry. Examples of such chemical reagents for selective modification include hydrazine and bisulfite ions. Hydrazine-modified DNA can be treated with piperidine to cleave it. Bisulfite ion-treated DNA can be treated with alkali. Other means for detection that are reliant on specific sequences can be used, including but not limited to hybridization, amplification, sequencing, and ligase chain reaction. Combinations of such techniques can be used as is desired.

[00183] *NMT2* epigenetic modification - a marker for one or more of diagnosis, prognosis, classifying, stratifying, or monitoring, of a cancer in a subject.

[00184] Determination of the methylation status of *NMT2* may be used to monitor efficacy of a therapeutic regimen, for example, a chemotherapeutic agent or a biological agent, such as an inhibitor of NMT.

[00185] Determination of the methylation status of *NMT2* may also be used to determine what therapeutic or preventive regimen to employ on a patient.

[00186] Determination of the methylation status of *NMT2* may also be used to stratify patients into groups for testing agents and determining their efficacy on various groups of patients. Such uses characterize the cancer into categories based on the genes, which are epigenetically silenced and/or the amount of silencing of the genes. In the case of a diagnosis or characterization, information comprising data or conclusions can be written or communicated electronically or orally. The identification may be assisted by a machine. Communication of the data or conclusions may be from a clinical laboratory to a clinical Office, from a clinician to a patient, or from a specialist to a

generalist, as examples. The form of communication of data or conclusions typically may involve a tangible medium or physical human acts.

5 [00187] In one example, a test sample obtainable from tissue or cell specimens or fluids includes detached tumor cells and/or free nucleic acids that are released from dead or damaged tumor cells.

[00188] Nucleic acids include RNA, genomic DNA, mitochondrial DNA, single or double stranded. Any nucleic acid specimen in purified or non-purified form obtained from such sample may be utilized. The test samples may contain cancer cells or pre-cancer cells or nucleic acids from them. In a specific example, genomic DNA is isolated.

10 [00189] Demethylating agents can be contacted with cells in vitro or in vivo for the purpose of restoring normal gene expression to the cell or for validation of methylation. Suitable demethylating agents include, but are not limited to 5-aza-2'-deoxycytidine, 5-azacytidine, Zebularine, procaine, and L-ethionine. This reaction may be used for diagnosis, for determining predisposition, and for determining suitable therapeutic regimes.

15 [00190] In one example, determination of the methylation status of NMT2 is for determining the condition(s) for the risk of developing cancer, metastasis, tumour relapse, prediction of the response to a treatment, the identification and/or selection of a cancer, such as lymphoma, the selection of a suitable treatment regimen, stratification of a patient, and the treatment of a patient.

20 [00191] In certain embodiments of the invention, the methylation status is determined in part or wholly on the basis of a comparison with a control. The control can be a value or set of values related to the extent and/or pattern of methylation in a control sample. In certain embodiments of the invention, such a value or values may be determined, for example, by calculations, using algorithms, and/or from previously acquired and/or archived data. In certain embodiments of the invention, the value or set of values for the control is derived from experiments performed on samples or using a subject. For example, control data can be derived from experiments on samples derived from comparable tissues or cells, such as normal tissue adjacent to tumors.

30 [00192] In certain embodiments of the invention, a control comprising DNA that is mostly or entirely demethylated, at one or more of the CpG sites being analyzed, is used. Such a control may be obtained, for example, from mutant tissues or cells lacking methyltransferase activity and/or from tissues or cells that have been chemically demethylated. For example, controls may be obtained from tissues or cells lacking activity

of methyltransferases. Agents such as 5-aza-2'-deoxycytidine may be used to chemically demethylate DNA.

[00193] In certain embodiments of the invention, a control comprising DNA that is mostly or entirely methylated, at one or more of the CpG sites being analyzed, is used.

- 5 Such a control may be obtained, for example, from cells or tissues that are known or expected to be mostly or entirely methylated at the CpG site or sites of interest. Such a control could also be obtained by cells or tissues in which methylation levels have been altered and/or manipulated.

- [00194]** In one example, described herein is a method of predicting the response in
10 a cancer patient, for example a lymphoma patient, to treatment of cancer. The drug for the treatment is an NMT inhibitor. In some examples, the NMT inhibitor is used in combination with at least one additional chemotherapy.

- [00195]** The presence of hypermethylation of *NMT2* indicates that the *NMT2* gene is methylated to a higher extent, which enables intervention with an NMT inhibitor.
15 Therefore, the presence of hypermethylation of *NMT2* indicates a more positive clinical response to said treatment, whereas the absence of methylation or a lower level of methylation compared to a control sample indicates an unsuccessful clinical response to the treatment with an NMT inhibitor. If a positive clinical response to treatment with an NMT inhibitor is determined, the patient is identified or selected for a treatment with said
20 NMT inhibitor. In other cases, the patient is not selected for treatment with an NMT inhibitor, and one or more alternative drug or medical interventions may be more beneficial for the treatment of the cancer patient.

- [00196]** Accordingly, there is provided the use of NMT2 epigenetic modification, for example methylation of NMT2, as a marker for one or more of diagnosis, prognosis,
25 classifying, or monitoring of cancer in a subject.

[00197] The term "prognosis" as used herein refers to the prediction of the likelihood of cancer-attributable death or progression, including recurrence, metastatic spread, and drug resistance, of a neoplastic disease, such as breast cancer.

- [00198]** The term "prognostic marker" as used herein refers to a marker that
30 informs about the outcome of a patient in the absence of systemic therapy or portends an outcome different from that of the patients without the marker, despite empiric (not targeted to the marker) systemic therapy.

[00199] The term "predictive marker" as used herein refers to a marker that predicts that differential efficacy (benefit) of a particular therapy based on marker status.

[00200] The term "diagnosis" as used herein, refers to the identification of a molecular and/or pathological state, disease or condition, such as the identification of breast cancer, or other type of cancer.

5 [00201] Testing can be performed diagnostically or in conjunction with a therapeutic regimen.

[00202] Therefore, in one aspect, there is provided a method of predicting the response in a cancer patient (for example a lymphoma patient) to treatment with an NMT inhibitor comprising providing a biological sample of said cancer patient (for example the lymphoma patient), and determining in said biological sample the methylation status of
10 the *NMT2* gene, and predicting a positive clinical response to said treatment, if hypermethylation is determined in the *NMT2* gene.

[00203] In another aspect, there is provided a method for identifying and/or selecting a patient with a cancer (for example a lymphoma), or treated for a cancer (for example a lymphoma), suitable for treatment with an NMT inhibitor, comprising providing
15 a biological sample of cancer patient, and determining in the methylation status of the *NMT2* gene, and identifying and/or selecting the cancer patient for treatment with said NMT inhibitor if hypermethylation is determined in the *NMT2* gene.

[00204] Further provided is a method for selecting a suitable treatment regimen for a cancer (for example a lymphoma) in a cancer patient comprising providing a biological
20 sample of said cancer patient, and determining in said biological sample the methylation status of the *NMT2* gene, and selecting an NMT inhibitor for treatment if hypermethylation and/or under expression is determined in the *NMT2* gene.

[00205] Further provided is a method of treating a patient with cancer (for example lymphoma) with an NMT inhibitor comprising providing a biological sample of said cancer
25 patient, and determining in said biological sample the methylation status of *NMT2*, and selecting said NMT inhibitor for treatment if hypermethylation is determined in *NMT2*.

[00206] It will be appreciated that in some circumstances, a patient who initially responds to NMT inhibitor treatment may relapse. Such a relapse can manifest in several ways, including but not limited to, reoccurrence of the primary tumour and
30 development of metastasis. In addition to, or alternatively, an additional distinct tumour can arise.

[00207] In accordance with one aspect of the present invention, there is provided a method comprising: a) obtaining a sample from a subject with, or suspected as having, cancer; b) contacting the sample a reagent to form a product (in some cases a complex)
35 indicative of methylation status of *NMT2* present in the sample; c) measuring the product

formed to determine an amount or a concentration of methylated *NMT2* in the sample; and d) determining the benefit of NMT inhibitor treatment of said cancer in said subject, wherein the determination of benefit of NMT inhibitor treatment is determined by hypermethylation of *NMT2* in said sample.

- 5 [00208] In a specific aspect, administering an NMT inhibitor to said subject is indicated when *NMT2* is hypermethylated in said sample, optionally as compared to a control.

- [00209] In accordance with one aspect, there is provided a method for treating cancer in a subject, comprising: a) obtaining nucleic acid from a sample from the subject;
10 b) performing bisulfite modification to the nucleic acid in step a); c) performing a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, on the bisulfite modified nucleic acid from step b) using PCR primers and/or
15 probes specific for the promoter region of the *NMT2* gene, e) determining a promoter methylation level of the promoter region of *NMT2* in the sample, optionally compared to a control, g) treating said patient with an inhibitor to NMT when promoter methylation level of the promoter region of *NMT2* gene in the sample is determined to be hypermethylated.

- [00210] In accordance with one aspect, there is provided an *in vitro* assay for
20 selecting a treatment for a patient with cancer, comprising: a) obtaining nucleic acid from a sample from the subject; b) performing bisulfite modification to the nucleic acid in step a); c) performing a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay,
25 pyrosequencing, or bisulfite sequencing, on the bisulfite modified nucleic acid from step b) using PCR primers and/or probes specific for the promoter region of the *NMT2* gene, e) determining a promoter methylation level of the promoter region of *NMT2* in the sample, optionally compared to a control, wherein the selected treatment is an NMT inhibitor when promoter region of *NMT2* gene in the sample is determined to be
30 hypermethylated.

- [00211] In accordance with one aspect, there is provided an *in vitro* assay of a sample from a patient for selecting a treatment for a patient with cancer, comprising: a) performing bisulfite modification to the nucleic acid in said sample, b) determining methylation of the promoter of the *NMT2* gene using PCR primers and/or probes specific
35 for the promoter region of the *NMT2* gene from step a).

[00212] In one aspect of the assay step b) is performed by a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, the bisulfite modified nucleic acid from step a).

[00213] In one aspect, detection of hypermethylation of the *NMT2* gene is performed with primers comprising SEQ ID NO: 1 and/or SEQ ID NO: 2.

[00214] In one aspect of the, there is provided a kit for selecting a suitable treatment for a cancer in a cancer patient, comprising: a) one or more reagents for determining the methylation status of the *NMT2* gene from a sample of said cancer patient, wherein the selected treatment is an NMT inhibitor when promoter region of *NMT2* gene in the sample is determined to be hypermethylated, b) instructions for the use thereof.

[00215] In one aspect, said reagents comprise PCR primers and/or probes specific for the promoter region of the *NMT2* gene.

[00216] In one aspect, there is provided a use of an NMT inhibitor for treating a subject with a cancer wherein the *NMT2* gene in a sample from said subject is determined to be hypermethylated.

[00217] In one aspect, there is provided a use of an NMT inhibitor for the manufacture of a medicament for treating a subject with a cancer wherein the *NMT2* gene in a sample from said subject is determined to be hypermethylated.

[00218] Epigenetic mechanisms- acetylation

[00219] As noted above, in some example, the epigenetic modification is histone acetylation.

[00220] An additional mechanism underlying gene silencing involves histone acetylation. Histone acetylation is mediated by histone acetyl transferases, while acetyl groups are removed by histone deacetylase(s) (HDAC).

[00221] It is shown herein that that lymphoma cells aberrantly increase methylation (hypermethylation) and decrease acetylation (hypoacetylation) to silence *NMT2*.

[00222] There are several structurally distinct classes of HDAC inhibitors including, but not limited to, valproic acid, valproates, VPA; butyrates, NaB; vorinostat, SAHA; romidepsin, depsipeptide; belinostat, PXD101; trichostatin A, TSA; apicidin; etinostat, MS275; mocetinostat, MGCD0103; panobinostat, LBH589; givinostat, ITF2357; pracinostat, SB939; CHR--3996I; chidamide, CS055, HBI--8000; AR--42; quisinostat, JNJ--26481585; abexinostat, PCI--24782; resminostat, 4SC--201, RAS2410; CG200745;

M344, ME--344; WJ25591; A248; NK--HDAC--1; MHY219; Ky--2; HDACi 4b; OBP--801, YM753; DWP0016; BRD8430; largazole; adamantine; trifluoromethyloxadiazole (TFMO) series, eg TMP195; YK--4--272, compound 2 (see Katrina J. Falkenberg¹ and Ricky W. Johnstone (2014) NATURE REVIEWS | DRUG DISCOVERY. Vol 13. 673- 691; and
 5 Andrew A. Lane and Bruce A. Chabner (2009) J Clin Oncol 27:5459-5468.) the contents all of which are hereby incorporated by reference, in their entirety.

[00223] Histone Acetyl Transferase inhibitors, DNA Demethylase Inhibitors, and/or histone demethylase inhibitors.

[00224] Histone acetyltransferases (HAT) are enzymes that acetylate conserved
 10 lysine amino acids on histone proteins by transferring an acetyl group from acetyl CoA to form e-N-acetyl lysine. Histone acetylation is generally linked to transcriptional activation. These are generally associated with euchromatin. Histone acetyltransferases (HATs) act as transcriptional coactivators. Histone acetylation plays an important role in regulating the chromatin structure and is tightly regulated by two classes of enzyme, histone
 15 acetyltransferases (HAT) and histone deacetylases (HDAC).

[00225] Non-limiting examples of HAT inhibitors include anacardic acid, CPTH2, MB-3, and Curcumin (Sigma-Aldrich).

[00226] DNA methyltransferases are a family of enzymes that catalyze the transfer of methyl group to DNA. There are five related DNA cytosine- 5-methyltransferases
 20 (DNMTs) that transfer a methyl group from S-adenosylmethionine (AdoMet, SAM) to the C-5 position of cytosine. Inhibitors have been shown to have anti-proliferative activity.

[00227] Non-limiting examples of DNA Methyltransferase Inhibitors include 5-Azacytidine, Zebularine, Caffeic acid purum, Chlorogenic acid, (-)-Epigallocatechin gallate, Hydralazine, Procainamide hydrochloride, Procaine hydrochloride, Psammoplan
 25 A, and RG108 (Sigma-Aldrich).

[00228] DNA demethylation can be achieved either passively or actively, by a replication-independent process.

[00229] While methylations are performed by Histone Methyl Transferases (HMTs) using the ubiquitous methyl donor S-adenosyl-methionine, two different oxidation
 30 mechanisms for Histone DeMethylases (HDMs) have been shown to lead to demethylation. Lysine Specific Demethylase 1 (LSD1/KDM1) was the first HDM to be discovered. LSD1 oxidises the lysine methyl ammonium group by catalysing the overall transfer of hydrogen from the methyl amino group to FAD generating a methyliminium ion. Other known HDMs are from the Jumonji protein family, characterized by containing
 35 the JmjC domain

[00230] Non-limiting examples of histone demethylase inhibitors include Daminozide, GSK J1, GSK J2, GSK J4, GSK J5, GSK LSD 1 dihydrochloride, IOX 1, JIB 04, RN 1 dihydrochloride, TC-E 5002, Tranylcypromine hydrochloride (Tocris - A Bio-Techne Brand).

5 [00231] In some examples herein, a subject with a subject with cancer or suspected of having cancer is treated with an NMT inhibitor, such as PCLX-001, as well as a histone acetyl transferase inhibitor, a DNA demethylase Inhibitor, and/or a histone demethylase inhibitor.

10 [00232] In some examples herein, a subject with a subject with a methylation status, which is hypermethylated, is treated with an NMT inhibitor, such as PCLX-001, as well as a histone acetyl transferase inhibitor, a DNA demethylase Inhibitor, and/or a histone demethylase inhibitor.

15 [00233] While not wishing to be bound by theory, it is believed that the histone acetyl transferase inhibitor, DNA demethylase Inhibitor, and/or histone demethylase inhibitor maintain the chromatin in a condensed form.

[00234] Methods of the invention are conveniently practiced by providing the reagents, compounds and/or compositions used in such method in the form of a kit. Such a kit preferably contains the composition. Such a kit preferably contains instructions for the use thereof.

20 [00235] To gain a better understanding of the invention described herein, the following examples are set forth. It should be understood that these example are for illustrative purposes only. Therefore, they should not limit the scope of this invention in any way.

[00236] **EXAMPLES**

25 [00237] In the following examples, standard methodologies were employed, as would be appreciated by the skilled worker.

[00238] **Example 1**

[00239] *NMT* expression levels in the 967 cancer cell lines encyclopedia (CCLE) database and 8178 primary tumours from The Cancer Genome Atlas (TCGA).

30 [00240] In Figure 1, *NMT2* shows a very large range of mRNA expression (log2 normalized microarray fluorescence~ 3-11) compared to *NMT1* (log2 microarray fluorescence ~6-9) (Figure 1A). *NMT1* (black) and *NMT2* (grey) expression values are log transformed RSEM normalized counts relative to TBP (TATA Binding Protein) (Figure 1B). *NMT2* mRNA expression of cancer cell lines from several cancer subtypes are plotted and compared to the plot of all 967 cancer cell lines (Figures 1C-E). Student T-

35

tests were performed comparing each group to all tumours. *** denotes P values < 0.001. Absolute quantification *NMT2* (Panel F) mRNA copies was measured by digital droplet PCR in Burkitt's lymphoma cell lines Ramos, and BL-2 and Diffuse Large B Cell lymphomas DOHH-2, WSU-DLCL2 compared to immortalized "normal" B lymphocytic cell line IM-9 (n=3). Myristoylation activity in various Burkitt's lymphoma cell lines Ramos, and BL-2 and Diffuse Large B Cell lymphomas DOHH-2, WSU-DLCL2 compared to immortalized "normal" B lymphocytic cell line IM-9 (n=3) (Figure 1G). *NMT2* expression in relation to progression-free survival in adult DLBCL patients (Figure 1H). Kaplan-Meier plots show the PFS of DLBC patients (GSE31312, n=470) with high (upper curve) versus low (lower curve) *NMT2* expression (Figure 1I). P values were determined by log-rank test,

[00241] A prominent mechanism underlying gene silencing involves histone acetylation together with DNA methylation, most prominently at CpG islands. *In silico* analyses revealed a CpG island in the *NMT2* 5' regulatory region, shown in Figure 2A. Bisulphite sequencing of DNA from lymphoma cell lines and immortalized IM9 B cells confirmed that the *NMT2* locus was methylated in DOHH2 and WSU-DLCL2 lymphoma cells, but not in BL2 and benign IM9 cells (Figure 2B). To validate the methylation status of the cell lines, we used DAPK1 as a control and found it to be highly methylated only in malignant B lymphocytes (Figure 2C)

[00242] Thus, it is shown herein that *NMT2* expression is commonly lost in lymphomas, and that this phenomenon is associated with more aggressive disease.

[00243] The loss of *NMT2* expression in lymphomas was almost complete, suggesting epigenetically silencing of *NMT2* in this cancer.

[00244] In order to establish a direct link between epigenetic modifications and the regulation of *NMT2* expression in human lymphoma cell lines and tumours, we next treated cells with inhibitors of DNA methylation and histone deacetylation, deoxyazacytidine (DAC) and suberoylanilide hydroxamic acid (SAHA), respectively. Each produced time- and concentration-dependent recovery of *NMT2* mRNA levels in lymphoma cells, and combined treatment with DAC and SAHA restored transcript levels in Ramos, BL2, DOHH2 and WSU-DLCL2 cells to those in untransformed IM9 cells (Figure 3A–D). These inhibitors acted synergistically when added together. Treating BL2 cells with combination of DAC and SAHA increased *NMT2* expression 6 fold resulting in *NMT2* levels approximating those seen in IM-9 cells.

[00245] Absolute quantification of *NMT2* mRNA copies were measured in BL2 (Figure 3, panel A) and IM9 (Figure 3, panel B) after 24h treatment with increasing

concentration of demethylation agent (DAC) in the presence of histone deacetylase inhibitor (SAHA) (n=3). WSU-DLCL2 is shown Figure 3C and 3D.

[00246] NMT2 protein level quantification in BL2 (Figure 3E) and IM9 (Figure 3F) (normalized to GAPDH level) after 24h treatment with increasing concentration of demethylation agent (DAC) in the presence of histone deacetylase inhibitor (SAHA). Graphic representation of 3 independent experiments with 1 representative western blot.

[00247] Using data extracted from patient samples, we found an inverse correlation between *NMT2* expression (RSEM normalized counts) and *NMT2* methylation (beta value) status (spearman coefficient -0.4; p=0.0082) (Figure 3G).

[00248] In BL2 cells, the increase in *NMT2* mRNA after combined treatment was associated with increased protein levels (Figure 3E-F). However, in DOHH2, and WSU-DLCL2 cells, *NMT2* was not detected by western blotting despite significant increases in *NMT2* mRNA levels upon treatment with DAC and/or SAHA. In patient samples, *NMT2* expression correlated inversely with *NMT2* methylation status (Figure 3G).

[00249] Together, these results suggested that lymphoma cells aberrantly increase methylation and decrease acetylation to silence *NMT2*, and that this loss of expression is associated with a more aggressive disease.

[00250] Example 2

[00251] We next investigated the therapeutic potential of DDD86481 *in vivo* to determine if it could mitigate tumour progression.

[00252] We tested the potential tumoricidal effects of DDD86481 monotherapy in two lymphoma cell line-derived mouse xenograft (CDX) models developed by injecting 1×10^7 BL-2 (BL) or DOHH-2 (DLBCL) cells subcutaneously in the flank of immunocompromised NODscid mice. DOHH-2 tumour ($125\text{--}200\text{mm}^3$) bearing mice (n=10 per group) were injected subcutaneously with PCLX-001 into the opposite flank at doses of 10, 20, 50 daily (QD) or 50 (every other day, QOD) mg per kg (mpk) or daily saline control.

[00253] We observed a dose-dependent tumoricidal effect, which gained statistical significance ($p < 0.05$) when PCLX-001 was given at 20mpk or 50 mpk QOD.

[00254] In mice bearing DOHH2 tumours ($125\text{--}200\text{ mm}^3$) (n = 10 mice per group), PCLX-001 injected subcutaneously into the opposite flank had a significant, tumoricidal effect at 20 mg/kg daily or 50 mg/kg every other day ($p < 0.001$) (Figure 4B). When injected daily at 50 mg/kg, PCLX-001 shrank tumours by 70% (average tumour size, $44.0 \pm 8.1\text{ mm}^3$ at day 7). However, this was accompanied by some weight loss, necessitating a 5-day interruption in treatment (not shown). However, upon treatment

resumption, a mean tumour growth inhibition (TGI) of 95% was observed by day 16. Importantly, tumour volume shrank in 6 of 10 mice between days 14 and 16, indicating that a longer treatment period might have proven beneficial.

[00255] Similarly, in mice bearing BL2 CDXs, PCLX-001 at 20 mg/kg led to 40% TGI by day 9 ($p < 0.05$) (Figure 5A). PCLX-001 given at 50 or 60 mg/kg for 13 consecutive days caused tumours to disappear in 9 of 9 and 7 of 7 mice, respectively. It is not known whether the four mice that did not survive the higher daily doses died from drug toxicity or tumour lysis syndrome. Nonetheless, some of the mice died, and dosing at 60 mg/kg daily had no apparent benefit. Therefore, the maximal tolerated dose seems to be 50 mg/kg.

[00256] These results suggest that DDD86481 can reduce or eliminate NMT2-deficient B cell lymphoma tumours in vivo.

[00257] PCLX-001 also reduced total NMT specific activity in BL2 tumours ($p < 0.05$) in a concentration-dependent manner (Figure 5C). These results in two independent lymphoma CDX models confirm that PCLX-001 can eradicate NMT2-deficient B-cell lymphomas in vivo.

[00258] Since CDXs lack the complexity of human tumours, a strategy was developed to identify patients with NMT2-deficient DLBCLs by immunohistochemistry or RNA in situ hybridization (Figures 5D and 5E). We then dissected and propagated one such tumour from patient "DLCL3" to establish a DLBCL patient-derived xenograft (PDX) model in NODscid mice. The mean initial tumour volume was 240 mm³ (Figure 5F).

[00259] Treatment regimens were assessed in groups of 8 mice each. At 20 mg/kg daily for 21 days, PCLX-001 caused a TGI of 60% ($p < 0.001$). At 50 mg/kg daily for two 9-day periods separated by a 3-day treatment suspension (to allow the mice to recover from the nearly 15% body weight loss), PCLX-001 caused the tumours to disappear in 6 of 7 surviving mice at day 13; one mouse with no detectable tumour died at day 7 (Figure 5F). In surgically removed control and PCLX-001-treated PDX tumours, we confirmed the concentration-dependent reduction of tumour size (Figure 5G). In the only remaining tumour from the 50-mg/kg treatment, PCLX-001 effectively induced apoptosis (Figure 5H) and reduced cell proliferation (Figure 5I). Thus, NMT2-deficient lymphomas can be identified and treated in a mouse PDX model. Moreover, the eradication of DLBCL PDX tumours in 7 of 8 mice suggests that the loss of NMT2 expression is an essential driver event in some DLBCL cases.

[00260] Example 3 - DLBCL xenograft Results

[00261] Seven groups of female SCID mice ($n = 10/\text{group}$) were dosed s.c. to completion in accordance with the DOHH-2-e225 protocol in Table 2.

[00262] Table 2 Mouse protocol – DLBCL

Group	Treatment
1	Control
2	CCI-002 at 10mpk SC daily
3	CCI-002 at 10mpk SC daily with doxorubicin 3mpk once a week IV starting on day 0
4	CCI-002 at 20mpk SC daily
5	CCI-002 at 50mpk SC every other day (QOD)
6	CCI-002 at 50mpk SC daily (QD) x 4, rest 5 days, then resume 50mpk SC daily (QD)
7	Doxorubicin 4mpk once a week IV starting on day 0

[00263] Figure 4A shows a box and whiskers plot showing the distribution of individual mice tumor volumes in each group.

[00264] Figure 4B shows median tumor growth in monotherapy.

[00265] Figure 4C shows median tumor growth in combination thereapy.

[00266] Efficacy

[00267] Tumor Growth in Control Mice (Group 1)

[00268] Group 1 mice received vehicle on a qd x 16 schedule starting on Day 1 and served as the control group for calculation of percent TGI and statistical comparisons. On Day 16, the median tumor volume (MTV) for Group 1 was 2213 mm³, with a range of 1764 to 2944 mm³ (Figure 4A). Tumors engrafted in all ten control mice.. Median tumor growth for Group 1 controls was progressive from the original tumor size 123.4 +/- 6.1 mm³.

[00269] Response to Treatment with CCI-002 as a daily monotherapy (Groups 2 and 4)

[00270] Groups 2 and 4 were treated with CCI-002 at active dosages of 10 mg/kg and 20 mg/kg, respectively, s.c. qd x 16 resulting in Day 16 MTVs of 1895 and 922 mm³. These median tumor volumes (MTVs) corresponded to a non-significant TGI of 14% for Group 2 (P > 0.05) and a significant TGI of 58% for Group 4 (P < 0.001), which left both groups below the potential therapeutic activity threshold. Tumors engrafted in all treated mice.

[00271] Response to Treatment with CCI-002 & doxorubicin combination

therapy (Group 3)

[00272] Group 3 was treated with a combination of CCI-002 at an active dosages of 10 mg/kg s.c. qd x 16 and doxorubicin i.v. qwk x 3, resulting in a Day 16 MTVs of 864 mm³. Group 3 had one animal that was euthanized on Day 11 due to excessive weight loss. That animal was sampled according to the protocol and classified as a NTR death and removed from all analyses. Group 3 had a MTV of 864 mm³ corresponding to a significant TGI of 61% ($P < 0.001$) which exceeded the potential therapeutic activity threshold (Figure 4A). The Group 3 combination was significantly more effective than the corresponding CCI-002 monotherapy (Group 3 vs. 2, $P < 0.001$) but not the corresponding doxorubicin treated group (Group 3 vs. 7, $P > 0.05$). Tumors engrafted in all treated mice.

[00273] Response to Treatment with HIGH dose CCI-002 Monotherapy (Groups 5 and 6)

[00274] Groups 5 and 6 were treated with CCI-002 at an active dosage of 50 mg/kg s.c. every other day qod x 8 or qd x 4/5 days off/qd x 14, respectively, resulting in Day 16 MTVs of 1018 and 226 mm³. These MTVs corresponded to significant TGIs of 54% for Group 5 ($P < 0.01$) and 90% for Group 6 ($P < 0.001$) with Group 6 above the potential therapeutic activity threshold (Figure 4A). Tumors engrafted in all treated mice.

[00275] Response to Treatment with doxorubicin monotherapy (Group 7)

[00276] Group 7 was treated with doxorubicin at 3 mg/kg i.v. qwk x 3 resulting in a Day 16 MTV of 1268 mm³ corresponding to a significant TGI of 43% ($P < 0.05$). Tumors engrafted in all treated mice.

[00277] Summary

[00278] This study evaluated CCI-002 activity when dosed as a monotherapy and in combination with doxorubicin. Tumors were measured three times per week through the final day of the study (Day 23). The TGI analysis was performed on Day 16.

[00279] All therapy except Group 2 were significantly superior to the vehicle-treated control group ($P < 0.05$). The CCI-002 at 50 mg/kg s.c. qd x 4/5 days off/qd x 14 treated group (Group 6) was the most effective regimen in this study with a MTV of 226 mm³ which corresponded to a 90% TGI. The next most effective therapies included the CCI-002 /doxorubicin combination group (Group 3), the CCI-002 monotherapy groups at 20 mg/kg and 50 mg/kg treated groups (Groups 4 and 5) which had respective TGI values of 61, 58 and 54% compared to control treated animals. There were no regressions recorded during the study.

[00280] All of the therapies tested in this DOHH-2 study were acceptably tolerated.

[00281] Example 4 - Burkitt lymphoma

[00282] Seven groups of female NOD SCID mice (n = 10/group) were dosed s.c. to completion in accordance with the DOHH-2-e225 protocol in Table 3 (below). The original treatment plan for this study can be found in Table 3. Figure 5 shows average (n=10 per group) tumor volume, from Day 0 (A. monotherapy and B combination therapy with doxorubicin).

[00283] Table 3—Mouse protocol –BL-2

Group	Treatment
1	Control
2	CCI-002 at 20mpk SC daily
3	Doxorubicin 3mpk once a week IV starting on day 0
4	CCI-002 at 10mpk SC daily and Doxorubicin once a week at 4mpk IV
5	CCI-002 at 20mpk SC daily and Doxorubicin once a week at 4mpk IV
6	CCI-002 at 50mpk SC daily
7	CCI-002 at 60mpk SC daily

[00284] Also, to reduce dehydration, on the first day of injection water bottles were removed and mice were provided with a rehydration solution made of NaCl: 4.5g/L (0.45%final), Glucose (Dextrose): 25g/L (2.5% final) and KCl: 0.75 g/L (0.075% final).

[00285] Results:**[00286] Tumor Growth in Control Mice (Group 1)**

[00287] Group 1 mice received vehicle on a qd x 9 schedule starting on Day 1 and served as the control group for calculation of percent TGI and statistical comparisons. On Day 9, the average tumor volume (ATV) for Group 1 was 1492 mm³, with a range of 685 to 2742 mm³. Tumors engrafted in all ten control mice. Median tumor growth for Group 1 controls was progressive from initial tumor volume of 255 mm³.

[00288] Response to treatment with CCI-002 as a daily monotherapy (Groups 2, 6 and 7)

[00289] Groups 2, 6 and 7 were treated with CCI-002 at active dosages of 20 mg/kg, 50 mg/kg and 60 mg/kg, respectively, s.c. qd x 16 resulting in Day 9 MTVs of 918, 74 and 72 mm³. These average tumor volumes (ATVs) corresponded to a significant relative tumor growth inhibition of 38% for Group 2 (P <####) and highly a significant relative tumor growth inhibition of 95% for Group 6 and 7 (P < 0.001), which left both groups 6 and 7 in the potential therapeutic activity threshold (TGI > 60%) (Figure 4A).

Tumors engrafted in all treated mice. Average tumor growth in these groups is illustrated in Figure 4A. Median tumor volume reached 0 mm³ by day 13 of the study therefore groups 6 and 7 treatments shows that CCI-002 causes 100% tumor regression.

[00290] Response to treatment with doxorubicin monotherapy (Group 3)

5 **[00291]** Group 3 was treated with doxorubicin at 4 mg/kg i.v. qwk x 3 resulting in a Day 9 MTV of 1061 mm³ corresponding to a significant TGI of 29% (P < 0.05). Tumors engrafted in all treated mice. Median tumor growth in these groups is illustrated in Figure 4A.

[00292] Response to treatment with CCI-002 & doxorubicin combination therapy (Groups 4 and 5)

10 **[00293]** Group 4 was treated with a combination of CCI-002 at an active dosage of 10 mg/kg s.c. qd x 16 and doxorubicin i.v. qwk x 3, resulting in a Day 9 average tumor volume (ATV) of 952 mm³ corresponding to a significant TGI of 36% (P < 0.001) (Figure 5). The Group 4 combination was not significantly more effective than the corresponding
15 doxorubicin monotherapy (Group 4 vs. 3, P > 0.05##). Group 5 was treated with a combination of CCI-002 at an active dosage of 20 mg/kg s.c. qd x 16 and doxorubicin i.v. qwk x 3, resulting in a Day 9 average tumor volume (ATV) of 471 mm³ corresponding to a significant TGI of 68% (P < 0.001) (Figure 3.4). The Group 5 combination was significantly more effective than the corresponding doxorubicin monotherapy (Group 5 vs.
20 3, P < 0.05##).

[00294] Tumors engrafted in all treated mice. Average tumor growth in these groups (n=10 mice/group) is illustrated in Figure 5.

[00295] Figure 5A. Average group (n=10/group) tumor volume for the duration of the treatment of NOD SCID mice injected with BL-2 cells (Initial average tumor size 125
25 mm³). A. CCI-002 monotherapy and B combination therapy of CCI-002 with doxorubicin.

[00296] Summary of results (BL)

[00297] Overall, the test article CCI-002 at 20 mg/kg, 50 mg/kg and 60 mg/kg treatment groups produced significant antitumor activities in treatment of the subcutaneous BL-2 Burkitt lymphoma subcutaneous xenograft model.

30 **[00298]** The test article CCI-002 showed an obvious dose response effect as monotherapy. When in combination with doxorubicin, the synergetic effect of antitumor activities was observed at 10 and 20 mg/kg CCI-002.

[00299] At the high doses of CCI-002 (50 and 60 mpk), treatment was suspended for various period of times to allow for recovery from body weight loss and "normal" body
35 weight was recovered.

[00300] Regarding the safety profile, the test article CCI-002 at 20 mg/kg showed good safety profile. However, a severe body weight loss (10-30%) was observed in the animals treated with CCI-002 at 50 and 60 mg/kg, doxorubicin 4 mg/kg or in combination treatment. Diarrhea in some mice of groups in Groups 4, 5, 6, and 7 was observed. 5 Importantly, after suspended for dosing and rehydration treatment (final concentration: NaCl: 2.5 g/L, Glucose: 30 g/L) for two days under the animal conditions, including diarrhea and body weight loss exceeding 15%, were improved above the 15% threshold.

[00301] Materials and Methods

[00302] Cell culture

10 **[00303]** L0, IM9, Ramos, BL2, Daudi, KMH2, Jurkat and CEM cells were purchased from the American Type Culture Collection (ATCC; Manassas, VA, U.S.A) and were maintained in RPMI medium supplemented with 10% FBS, 100 U/ml penicillin, 0.1 mg/ml streptomycin, 1 mM sodium pyruvate and 2mM L-Glutamine. All cells used were maintained at 37°C and 5% CO₂ in a humidified incubator.

15 **[00304] Cell Lysis**

[00305] Cells were washed in cold PBS, harvested, lysed in 0.1% SDS-RIPA buffer [50 mM Tris pH 8.0, 150 mM NaCl, 1% Igepal CA-630, 0.5% NaDC, 2 mM MgCl₂, 2 mM EDTA with 1x complete protease inhibitor (Roche Diagnostics)] by rocking for 15 min at 4°C. Cell lysates were then centrifuged at 16,000 g for 10 min at 4°C and the post- 20 nuclear supernatant was collected.

[00306] Viability of cells treated with NMT inhibitors

[00307] 2 x 10⁶ cells CEM (T cell leukemia), L0 ("normal" B cell), BL2 and Ramos] were grown in 6 well plates and incubated with increasing amounts (0, 1, 2 and 5 µg/ml) of Tris-DBA (25) for 24 hours. Tris-DBA was a kind gift from Dr. Jack Arbiser (Emory 25 University, GA, USA). 1 x 10⁵ cells [KMH2 (Hodgkins lymphoma), IM9 ("normal" B cell), BL2, Ramos] were plated in 96-well plates and treated with increasing amounts of DDD85646, DDD73228 and DDD86481 for 24, 48 and 72 hours. DDD85646, DDD73228 and DDD86481 were kind gifts from Drs. David Gray and Paul Wyatt at the University of Dundee, Scotland, UK.

30 **[00308] Measurement of cell viability**

[00309] The viability of cells treated with Tris-DBA, DDD73228, and DDD85646 was measured by incubating cells treated with Tris-DBA using TC10TM Trypan Blue Dye (Biorad, Hercules, CA, USA) according to manufacturer's instructions. Cell viability was then quantified using the TC10TM automated cell counter (Biorad). The viability of cells 35 treated with DDD86481 was measured using the CellTiter 96 AQueous Non-Radioactive

cell proliferation assay (MTS) from Promega (Madison, WI, USA) according to manufacturer's instructions.

[00310] qRT-PCR of B cell lymphoma cell lines

[00311] RNA was isolated from IM9, KMH2, Ramos and BL2 cells using TRIzol®

- 5 reagent according to manufacturer's protocols. Next, the High Capacity cDNA Reverse Transcription kit with a random primer scheme from Applied Biosciences was used to synthesize cDNA from the isolated RNA according to manufacturer's instructions. Quantitative real time PCR (qRT PCR) reactions were set-up using the TaqMan® Universal Master Mix II and NMT1, NMT2 and 18 Taqman® probes purchased from Life
- 10 Technologies (Carlsbad, CA) and three replicates of each reaction were set up according to supplier's guidelines. qRT PCR was performed using a Mastercycler® ep realplex thermocycler (Eppendorf) and results were analyzed using the Realplex software (Eppendorf).

[00312] Treatment of cells with SAHA

- 15 **[00313]** IM9, BL2 and Ramos cells were plated at 3x10⁶ cells per well in a 6-well dish and treated with 1 μ M suberoylanilide hydroxamic acid (SAHA) for 24 hours. Equal amount of DMSO was added to the control samples. Cells were lysed and subjected to SDS-PAGE. Western blotting was performed with antibodies against NMT1, NMT2, p21/WAF-1 and GAPDH.

- 20 **[00314]** Bisulphite sequencing

[00315] Chromatin DNA was isolated from the cells using the QIAamp DNA and Blood Mini kit from Qiagen. DNA (20 μ l; concentration, 1 ng to 2 μ g/ μ l) was converted with the EpiTech Bisulfite kit (Qiagen) and amplified with bisulphite-specific primers (Table 4).

- 25 **[00316]** **Table 4. Primers used for bisulphite sequencing.**

Primers	Primer Sequence	SEQ ID NO:	Tm (°C)	Product size (bp)
1F	GTTTTGTTTTTGTGTTGAGAT	1	52.2	344
1R	TCCTTTTATATTTTAACCCCATTTAC	2		
2F	TTTAATAAGGATGGTAGGGAGTAGAG	3	53	519
2R	ACCAACCAACAAAACCTAAACTAA	4		
3F	GATTTAATAAGGATGGTAGGGAGTAG	5	51	360

3R	ACTCCCAAACTAAAAATTCTTC	6		
4F	GGTTTTGTTTTGAAGAGTTTTAGG	7	53	232
4R	ACTCCCTACCATCCTTATTAAATCC	8		

[00317] The amplified PCR products were cleaned with QIAquick PCR Purification kit (Qiagen) and cloned with the TA Cloning Kit (Life Technologies) and pC 2.1 Vector.

5 The sequences were analyzed with QUMA.

[00318] All publications, patents and patent applications mentioned in this Specification are indicative of the level of skill those skilled in the art to which this invention pertains and are herein incorporated by reference to the same extent as if each individual publication patent, or patent application was specifically and individually

10 indicated to be incorporated by reference.

[00319] The above-described embodiments are intended to be examples only. Alterations, modifications and variations can be effected to the particular embodiments by those of skill in the art. The scope of the claims should not be limited by the particular embodiments set forth herein, but should be construed in a manner consistent with the

15 specification as a whole.

WHAT IS CLAIMED IS:

1. A method of predicting the response in a cancer patient to treatment with an NMT inhibitor comprising, providing a biological sample of said cancer patient, and determining in said biological sample the methylation status of the *NMT2* gene, and predicting a positive clinical response to said treatment with said NMT inhibitor, if hypermethylation is determined in the *NMT2* gene.
2. A method for identifying and/or selecting a patient with a cancer, or treated for a cancer, suitable for treatment with an NMT inhibitor, comprising providing a biological sample of said cancer patient, and determining the methylation status of the *NMT2* gene, and identifying and/or selecting the cancer patient for treatment with said NMT inhibitor if hypermethylation is determined in the *NMT2* gene.
3. A method for selecting a suitable treatment regimen for a cancer in a cancer patient comprising providing a biological sample of said cancer patient, and determining in said biological sample the methylation status of the *NMT2* gene, and selecting an NMT inhibitor for treatment if hypermethylation of *NMT2* gene is determined and/or under expression is determined in the *NMT2* gene.
4. A method of treating a patient with cancer with an NMT inhibitor comprising providing a biological sample of said cancer patient, and determining in said biological sample the methylation status of *NMT2* gene, and selecting said NMT inhibitor for treatment if hypermethylation is determined in *NMT2* gene.
5. A method comprising: a) obtaining a sample from a subject having cancer, or suspected as having cancer; b) contacting the sample with a reagent to form a product (in some cases a complex) indicative of methylation status of *NMT2* gene present in the sample; c) measuring the product formed to determine a methylation status of *NMT2* gene in the sample; and d) determining the benefit of NMT inhibitor treatment of said cancer in in said subject, wherein the determination of benefit of NMT inhibitor treatment is determined by hypermethylation of *NMT2* gene in said sample.

6. The method of claim 5, wherein in step b) said reagent comprises primers to detect hypermethylation in the *NMT2* gene.

7. The method of claim 6, wherein said primers comprise SEQ ID NO:1 and/or SEQ ID NO: 2.

8. A method of treating a subject with cancer or suspect of having cancer, comprising administering an NMT inhibitor to said subject when the *NMT2* gene is hypermethylated in said sample, optionally as compared to a control.

9. A method for treating cancer in a subject, comprising:
a) obtaining nucleic acid from a sample from the subject;
b) performing bisulfite modification to the nucleic acid in step a);
c) performing a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, on the bisulfite modified nucleic acid from step b)
d) using PCR primers and/or probes specific for the promoter region of the *NMT2* gene,
e) determining a promoter methylation level of the promoter region of *NMT2* in the sample, optionally compared to a control,
f) treating said patient with an inhibitor to NMT when promoter methylation level of the promoter region of *NMT2* gene in the sample is determined to be hypermethylated.

10. The method of claim 9, wherein said PCR primers comprise SEQ ID NO:1 and/or SEQ ID NO: 2.

11. The method of any one of claims 1-9, wherein determining the methylation status comprises, a) performing bisulfite modification to the nucleic acid in said sample, b) determining methylation of the promoter of the *NMT2* gene using PCR primers and/or probes specific for the promoter region of the *NMT2* gene from step a).

12. The method of claim 11, wherein step b) is performed by a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein,

quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, the bisulfite modified nucleic acid from step a).

13. The method of claim 11 or 12, wherein detection of hypermethylation of the NMT2 gene is performed with primers comprising SEQ ID NO: 1 and/or SEQ ID NO: 2.

14. The method of any one of claim 1 – 13, wherein said NMT inhibitor comprises a small molecule, an antibody, a peptide fragment, a nucleic acid, or combinations thereof.

15. The method of claim 14, wherein said small molecule comprises DDD85646, DDD86481, or a derivative thereof.

16. The method of claim 14, wherein said antibody is a monoclonal antibody or a polyclonal antibody.

17. The method of claim 14, wherein said nucleic acid comprises a dsRNA molecule, a RNAi molecule, a miRNA molecule, a ribozyme, a shRNA molecule, or a siRNA molecule.

18. The method of any one of claim 1 to 17, wherein said cancer is lymphoma, B cell lymphoma, follicular lymphoma, diffuse large B-cell lymphoma, mantle cell lymphoma, B-CLL/SLL, immunocytoma/Waldenstrom's, MALT-type/monocytoid B 20 cell lymphoma, Burkitt's lymphoma, a pediatric lymphoma, anaplastic large cell lymphoma, acute myeloid leukemia, Blast Phase Chronic Myeloid Leukaemia, Burkitt's Lymphoma, Plasma Cell Myeloma, Intestinal Adenocarcinoma, Lung mixed Adenosquamous Carcinoma, Lung Small Cell Carcinoma, Lung, Oesophagus Squamous Cell Carcinoma, Bone, Breast Ductal Carcinoma, Stomach Diffuse Adenocarcinoma, Thyroid Medullary Carcinoma, urinary Tract Transitional Cell Carcinoma, myeloma, ovarian clear cell carcinoma, transition cell carcinoma (ureter and bladder cancer), chronic myelogenous leukemia (CML), lymphoma-CLL, breast carcinoma, colorectal adenocarcinoma, pancreas adenocarcinoma, ovarian carcinoma, non-small cell lung carcinoma, osteosarcoma, melanoma, gastric adenocarcinoma, endometrial adenocarcinoma, esophageal squamous carcinoma.

19. The method of any one of claims 1-18, wherein said NMT inhibitor comprises DDD85646 or DDD86481.
20. The method of any one of claims 1 to 19, wherein said treatment further comprises treatment with a histone acetyl transferase inhibitor, a DNA demethylase Inhibitor, and/or histone demethylase inhibitor.
21. The method of claim 20, wherein said histone acetyl transferase inhibitor is anacardic acid, CPTH2, MB-3, and/or Curcumin.
22. The method of claim 20, wherein said histone demethylase inhibitor is Daminozide, GSK J1, GSK J2, GSK J4, GSK J5, GSK LSD 1 dihydrochloride, IOX 1, JIB 04, RN 1 dihydrochloride, TC-E 5002, and/or Tranylcypromine hydrochloride.
23. The method of any one of claims 1 – 22, wherein when said cancer is a lymphoma, the method further comprising administering a treatment selected from CHOP (i.e., cyclophosphamide, doxorubicin, vincristine, and prednisone), GAP-BOP (i.e., cyclophosphamide, doxorubicin, procarbazine, bleomycin, vincristine, and prednisone), m-BACOD (i.e., methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine, dexamethasone, and leucovorin), ProMACE-MOPP (i.e., prednisone, methotrexate, doxorubicin, cyclophosphamide, etoposide, leucovorin with standard MOPP), ProMACE-CytaBOM (prednisone, doxorubicin, cyclophosphamide, etoposide, cytarabine, bleomycin, vincristine, methotrexate, and leucovorin), and MACOP-B (methotrexate, doxorubicin, cyclophosphamide, vincristine, prednisone, bleomycin, and leucovorin). For relapsed aggressive non-Hodgkin's lymphoma the following chemotherapy drug combinations may be used with the compounds and compositions described herein: IMVP-16 (i.e., ifosfamide, methotrexate, and etoposide), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), DHAP (i.e., dexamethasone, - 16 high dose cytarabine, and cisplatin), ESHAP (i. e., etoposide, methylprednisone, high dosage cytarabine, and cisplatin), CEFF(B) (i.e., cyclophosphamide, etoposide, procarbazine, prednisone, and bleomycin), or CAMP (i.e., lomustine, mitoxantrone, cytarabine, and prednisone).

24. The method of any one of claims 1-22, wherein when said cancer is Hodgkin's disease, the method further comprising administering a treatment selected from VABCD (i.e., vinblastine, doxorubicin, dacarbazine, lomustine and bleomycin), ABDIC (i.e., doxorubicin, bleomycin, dacarbazine, lomustine, and prednisone), CBVD (i.e., lomustine, bleomycin, vinblastine, dexamethasone), PCVP (i.e., vinblastine, procarbazine, cyclophosphamide, and prednisone), CEP (i.e., lomustine, etoposide, and prednimustine), EVA (i.e., etoposide, vinblastine, and doxorubicin), MOPLACE (i.e., cyclophosphamide, etoposide, prednisone, methotrexate, cytarabine, and vincristine), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), MINE (i.e., mitoguazone, ifosfamide, vinorelbine, and etoposide), MTX-CHOP (i.e., methotrexate and CHOP), CEM (i.e., lomustine, etoposide, and methotrexate), CEVD (i.e., lomustine, etoposide, vindesine, and dexamethasone), CAVP (i.e., lomustine, melphalan, etoposide, and prednisone), EVAP (i.e., etoposide, vinblastine, cytarabine, and cisplatin), and EPOCH (i.e., etoposide, vincristine, ; doxorubicin, cyclophosphamide, and prednisone).

25. The method of any one of claims 1-22, wherein when said cancer is breast cancer, the method further comprising administering a treatment selected from paclitaxel, docetaxel, nanoparticle albumin bound paclitaxel, capecitabine, doxorubicin, eribulin, vinorelbine, trastuzumab, carboplatin, cisplatin; endocrine therapies including tamoxifen, anastrozole, letrozole, exemestane, fulvestrant, and cell cycle inhibitors including palbociclib, ribociclib, LEE001, and everolimus; immune checkpoint inhibiting drugs including nivolumab and pembrolizumab, either individually or in combination.

26. The method of any one of claims 1-22, wherein when said cancer is small cell lung cancer, the method further comprising administering a treatment selected from carboplatin, cisplatin, etoposide, irinotecan, either individually or in combination.

27. The method of any one of claims 1-22, wherein when said cancer is non-small cell lung cancer, the method further comprising administering a treatment selected from carboplatin, paclitaxel, docetaxel, gefitinib, erlotinib, afatinib, bevacizumab, ramucirumab, osimertinib, necitumumab, crizotinib, ceritinib, alectinib, and immune checkpoint inhibiting drugs including nivolumab, pembrolizumab, either individually or in combination.

28. The method of any one of claims 1-22, wherein when said cancer is bladder carcinoma, the method further comprising administering a treatment selected from methotrexate, vincblastine, doxorubicin, cisplatin; carboplatin and paclitaxel; docetaxel, gemcitabine and cisplatin, either individually or in combination.
29. The method of any one of claims 1-22, wherein the determination of hypermethylation in the *NMT2* gene is performed by a method selected from the group consisting of PCR, methylation-specific PCR, realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, and bisulfite sequencing.
30. The method of any one of claims 1-29, wherein said biological sample comprises, a tissue, a cell specimen or fluids includes detached tumor cells and/or free nucleic acids, a blood sample, a fractionated blood sample, a bone marrow sample, a biopsy sample, a frozen tissue sample, a fresh tissue specimen, a cell sample, and/or a paraffin embedded section.
31. The method of anyone of claims 1-30, wherein said subject is a human.
32. An *in vitro* assay for selecting a treatment for a patient with cancer, comprising:
a) obtaining nucleic acid from a sample from the subject;
b) performing bisulfite modification to the nucleic acid in step a);
c) performing a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, on the bisulfite modified nucleic acid from step b) using PCR primers and/or probes specific for the promoter region of the *NMT2* gene,
e) determining a promoter methylation level of the promoter region of *NMT2* in the sample, optionally compared to a control,
wherein the selected treatment is an NMT inhibitor when promoter region of *NMT2* gene in the sample is determined to be hypermethylated.

33. The method of claim 23, wherein said PCR primers comprise SEQ ID NO:1 and/or SEQ ID NO: 2.
34. An *in vitro* assay of a sample from a patient for selecting a treatment for a patient with cancer, comprising:
- a) performing bisulfite modification to the nucleic acid in said sample,
 - b) determining methylation of the promoter of the NMT2 gene using PCR primers and/or probes specific for the promoter region of the NMT2 gene from step a).
35. The assay of claim 34, wherein step b) is performed by a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, the bisulfite modified nucleic acid from step a).
36. The assay of claim 34 or 35, wherein detection of hypermethylation of the NMT2 gene is performed with primers comprising SEQ ID NO: 1 and/or SEQ ID NO: 2.
37. The assay of any one of claims 34 to 36, wherein the selected treatment is an NMT inhibitor when promoter region of *NMT2* gene in the sample is determined to be hypermethylated.
38. The assay of any one of claim 34 to 37, wherein said NMT inhibitor comprises a small molecule, an antibody, a peptide fragment, a nucleic acid, or combinations thereof.
39. The assay of claim 38, wherein said small molecule comprises DDD85646, DDD86481, or a derivative thereof.
40. The assay of claim 38, wherein said antibody is a monoclonal antibody or a polyclonal antibody.
41. The assay of claim 38, wherein said nucleic acid comprises a dsRNA molecule, a RNAi molecule, a miRNA molecule, a ribozyme, a shRNA molecule, or a siRNA molecule.

42. A kit for selecting a suitable treatment for a cancer in a cancer patient, comprising:
a) one or more reagents for determining the methylation status of the *NMT2* gene from a sample of said cancer patient, wherein the selected treatment is an NMT inhibitor when promoter region of *NMT2* gene in the sample is determined to be hypermethylated,
b) instructions for the use thereof.
43. The kit of claim 42, wherein said reagents comprise PCR primers and/or probes specific for the promoter region of the *NMT2* gene.
44. The kit of claim 43, wherein said PCR primers comprise SEQ ID NO:1 and/or SEQ ID NO: 2.
45. The kit of any one of claims 42-44, further comprising and NMT inhibitor.
46. The kit of claim 45, said NMT inhibitor comprises a small molecule, an antibody, a peptide fragment, a nucleic acid, or combinations thereof.
47. The kit of claim 46, wherein said small molecule comprises DDD85646, DDD86481, or a derivative thereof.
48. The kit of claim 46, wherein said antibody is a monoclonal antibody or a polyclonal antibody.
49. The kit of claim 46, wherein said nucleic acid comprises a dsRNA molecule, a RNAi molecule, a miRNA molecule, a ribozyme, a shRNA molecule, or a siRNA molecule.
50. The kit of any one of claims 42 to 49, further comprising a histone acetyl transferase inhibitor, a DNA demethylase Inhibitor, and/or histone demethylase inhibitor.
51. The kit of claim 50, wherein said histone acetyl transferase inhibitor is anacardic acid, CPTH2, MB-3, and/or Curcumin.

52. The kit of claim 50, wherein said histone demethylase inhibitor is Daminozide, GSK J1, GSK J2, GSK J4, GSK J5, GSK LSD 1 dihydrochloride, IOX 1, JIB 04, RN 1 dihydrochloride, TC-E 5002, and/or Tranylcypromine hydrochloride.

53. The kit of any one of claims 42 to 52, further comprising reagents for a treatment selected from CHOP (i.e., cyclophosphamide, doxorubicin, vincristine, and prednisone), GAP-BOP (i.e., cyclophosphamide, doxorubicin, procarbazine, bleomycin, vincristine, and prednisone), m-BACOD (i.e., methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine, dexamethasone, and leucovorin), ProMACE-MOPP (i.e., prednisone, methotrexate, doxorubicin, cyclophosphamide, etoposide, leucovorin with standard MOPP), ProMACE-CytaBOM (prednisone, doxorubicin, cyclophosphamide, etoposide, cytarabine, bleomycin, vincristine, methotrexate, and leucovorin), and MACOP-B (methotrexate, doxorubicin, cyclophosphamide, vincristine, prednisone, bleomycin, and leucovorin). For relapsed aggressive non-Hodgkin's lymphoma the following chemotherapy drug combinations may be used with the compounds and compositions described herein: IMVP-16 (i.e., ifosfamide, methotrexate, and etoposide), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), DHAP (i.e., dexamethasone, - 16 high dose cytarabine, and cisplatin), ESHAP (i.e., etoposide, methylprednisone, high dosage cytarabine, and cisplatin), CEFF(B) (i.e., cyclophosphamide, etoposide, procarbazine, prednisone, and bleomycin), or CAMP (i.e., lomustine, mitoxantrone, cytarabine, and prednisone).

54. The kit of any one of claims 42-52, further comprising reagents for a treatment for Hodgkin's disease selected from VABCD (i.e., vinblastine, doxorubicin, dacarbazine, lomustine and bleomycin), ABDIC (i.e., doxorubicin, bleomycin, dacarbazine, lomustine, and prednisone), CBVD (i.e., lomustine, bleomycin, vinblastine, dexamethasone), PCVP (i.e., vinblastine, procarbazine, cyclophosphamide, and prednisone), CEP (i.e., lomustine, etoposide, and prednimustine), EVA (i.e., etoposide, vinblastine, and doxorubicin), MOPLACE (i.e., cyclophosphamide, etoposide, prednisone, methotrexate, cytarabine, and vincristine), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), MINE (i.e., mitoquazone, ifosfamide, vinorelbine, and etoposide), MTX-CHOP (i.e., methotrexate and CHOP), CEM (i.e., lomustine, etoposide, and methotrexate), CEVD (i.e., lomustine, etoposide, vindesine, and dexamethasone), CAVP (i.e., lomustine, melphalan, etoposide, and prednisone), EVAP (i.e., etoposide, vinblastine, cytarabine,

and cisplatin), and EPOCH (i.e., etoposide, vincristine, ; doxorubicin, cyclophosphamide, and prednisone).

55. The kit of any one of claims 42-52, further comprising reagents for a treatment for breast cancer selected from paclitaxel, docetaxel, nanoparticle albumin bound paclitaxel, capecitabine, doxorubicin, eribulin, vinorelbine, trastuzumab, carboplatin, cisplatin; endocrine therapies including tamoxifen, anastrozole, letrozole, exemestane, fulvestrant, and cell cycle inhibitors including palbociclib, ribociclib, LEE001, and everolimus; immune checkpoint inhibiting drugs including nivolumab and pembrolizumab, either individually or in combination.

56. The kit of any one of claims 42-52, further comprising reagents for a treatment for small cell lung cancer selected from carboplatin, cisplatin, etoposide, irinotecan, either individually or in combination.

57. The kit of any one of claims 42-52, further comprising reagents for a treatment for non-small cell lung cancer selected from carboplatin, paclitaxel, docetaxel, gefitinib, erlotinib, afatinib, bevacizumab, ramucirumab, osimertinib, necitumumab, crizotinib, ceritinib, alectinib, and immune checkpoint inhibiting drugs including nivolumab, pembrolizumab, either individually or in combination.

58. The kit of any one of claims 42-52, further comprising reagents for a treatment for bladder carcinoma, the method further comprising administering a treatment selected from methotrexate, vincblastine, doxorubicin, cisplatin; carboplatin and paclitaxel; docetaxel, gemcitabine and cisplatin, either individually or in combination.

59. Use of an NMT inhibitor for treating a subject with a cancer wherein the *NMT2* gene in a sample from said subject is determined to be hypermethylated.

60. Use of an NMT inhibitor for the manufacture of a medicament for treating a subject with a cancer wherein the *NMT2* gene in a sample from said subject is determined to be hypermethylated.

61. The use of claim 59 or 60, wherein determining the methylation status comprises, a) performing bisulfite modification to the nucleic acid in said sample, b) determining methylation of the promoter of the *NMT2* gene using PCR primers and/or probes specific for the promoter region of the *NMT2* gene from step a).

62. The use of claim 61, wherein step b) is performed by a method selected from PCR, methylation-specific PCR, quantitative methylation specific PCR (QMSP), realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, or bisulfite sequencing, the bisulfite modified nucleic acid from step a).

63. The use of any one of claims claim 59 to 62, wherein detection of hypermethylation of the *NMT2* gene is performed with primers comprising SEQ ID NO: 1 and/or SEQ ID NO: 2.

64. The use of any one of claim 59 – 63, wherein said NMT inhibitor comprises a small molecule, an antibody, a peptide fragment, a nucleic acid, or combinations thereof.

65. The use of claim 64, wherein said small molecule comprises DDD85646, DDD86481, or a derivative thereof.

66. The use of claim 64, wherein said antibody is a monoclonal antibody or a polyclonal antibody.

67. The use of claim 64, wherein said nucleic acid comprises a dsRNA molecule, a RNAi molecule, a miRNA molecule, a ribozyme, a shRNA molecule, or a siRNA molecule.

68. The use of any one of claim 59 to 67, wherein said cancer is lymphoma, B cell lymphoma, follicular lymphoma, diffuse large B-cell lymphoma, mantle cell lymphoma, B-CLL/SLL, immunocytoma/Waldenstrom's, MALT-type/monocytoid B 20 cell lymphoma, Burkitt's lymphoma, a pediatric lymphoma, anaplastic large cell lymphoma, acute myeloid leukemia, Blast Phase Chronic Myeloid Leukaemia, Burkitt's Lymphoma, Plasma Cell Myeloma, Intestinal Adenocarcinoma, Lung mixed Adenosquamous Carcinoma, Lung Small Cell Carcinoma, Lung, Oesophagus Squamous Cell Carcinoma, Bone, Breast

Ductal Carcinoma, Stomach Diffuse Adenocarcinoma, Thyroid Medullary Carcinoma, urinary Tract Transitional Cell Carcinoma, myeloma, ovarian clear cell carcinoma, transition cell carcinoma (ureter and bladder cancer), chronic myelogenous leukemia (CML), lymphoma-CLL, breast carcinoma, colorectal adenocarcinoma, pancreas adenocarcinoma, ovarian carcinoma, non-small cell lung carcinoma, osteosarcoma, melanoma, gastric adenocarcinoma, endometrial adenocarcinoma, esophageal squamous carcinoma.

69. The use of any one of claims 59-68, wherein said NMT inhibitor comprises DDD85646 or DDD86481.

70. The use of any one of claims 59 to 69, further comprising use of a histone acetyl transferase inhibitor, a DNA demethylase Inhibitor, and/or histone demethylase inhibitor.

71. The use of claim 70, wherein said histone acetyl transferase inhibitor is anacardic acid, CPTH2, MB-3, and/or Curcumin.

72. The use of claim 70, wherein said histone demethylase inhibitor is Daminozide, GSK J1, GSK J2, GSK J4, GSK J5, GSK LSD 1 dihydrochloride, IOX 1, JIB 04, RN 1 dihydrochloride, TC-E 5002, and/or Tranylcypromine hydrochloride.

73. The use of any one of claims 59 – 72, wherein when said cancer is a lymphoma, the use further comprising use of a treatment selected from CHOP (i.e., cyclophosphamide, doxorubicin, vincristine, and prednisone), GAP-BOP (i.e., cyclophosphamide, doxorubicin, procarbazine, bleomycin, vincristine, and prednisone), m-BACOD (i.e., methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine, dexamethasone, and leucovorin), ProMACE-MOPP (i.e., prednisone, methotrexate, doxorubicin, cyclophosphamide, etoposide, leucovorin with standard MOPP), ProMACE-CytaBOM (prednisone, doxorubicin, cyclophosphamide, etoposide, cytarabine, bleomycin, vincristine, methotrexate, and leucovorin), and MACOP-B (methotrexate, doxorubicin, cyclophosphamide, vincristine, prednisone, bleomycin, and leucovorin). For relapsed aggressive non-Hodgkin's lymphoma the following chemotherapy drug combinations may be used with the compounds and compositions described herein: IMVP-16 (i.e., ifosfamide, methotrexate, and etoposide), MIME (i.e., methyl-gag,

ifosfamide, methotrexate, and etoposide), DHAP (i.e., dexamethasone, - 16 high dose cytarabine, and cisplatin), ESHAP (i.e., etoposide, methylprednisone, high dosage cytarabine, and cisplatin), CEFF(B) (i.e., cyclophosphamide, etoposide, procarbazine, prednisone, and bleomycin), or CAMP (i.e., lomustine, mitoxantrone, cytarabine, and prednisone).

74. The use of any one of claims 59-72, wherein when said cancer is Hodgkin's disease, the use further comprising use of a treatment selected from VABCD (i.e., vinblastine, doxorubicin, dacarbazine, lomustine and bleomycin), ABDIC (i.e., doxorubicin, bleomycin, dacarbazine, lomustine, and prednisone), CBVD (i.e., lomustine, bleomycin, vinblastine, dexamethasone), PCVP (i.e., vinblastine, procarbazine, cyclophosphamide, and prednisone), CEP (i.e., lomustine, etoposide, and prednimustine), EVA (i.e., etoposide, vinblastine, and doxorubicin), MOPLACE (i.e., cyclophosphamide, etoposide, prednisone, methotrexate, cytarabine, and vincristine), MIME (i.e., methyl-gag, ifosfamide, methotrexate, and etoposide), MINE (i.e., mitozantrone, ifosfamide, vinorelbine, and etoposide), MTX-CHOP (i.e., methotrexate and CHOP), CEM (i.e., lomustine, etoposide, and methotrexate), CEVD (i.e., lomustine, etoposide, vindesine, and dexamethasone), CAVP (i.e., lomustine, melphalan, etoposide, and prednisone), EVAP (i.e., etoposide, vinblastine, cytarabine, and cisplatin), and EPOCH (i.e., etoposide, vincristine, ; doxorubicin, cyclophosphamide, and prednisone).

75. The use of any one of claims 59-72, wherein when said cancer is breast cancer, the use further comprising use of a treatment selected from paclitaxel, docetaxel, nanoparticle albumin bound paclitaxel, capecitabine, doxorubicin, eribulin, vinorelbine, trastuzumab, carboplatin, cisplatin; endocrine therapies including tamoxifen, anastrozole, letrozole, exemestane, fulvestrant, and cell cycle inhibitors including palbociclib, ribociclib, LEE001, and everolimus; immune checkpoint inhibiting drugs including nivolumab and pembrolizumab, either individually or in combination.

76. The use of any one of claims 59-72, wherein when said cancer is small cell lung cancer, the use further comprising a use of a treatment selected from carboplatin, cisplatin, etoposide, irinotecan, either individually or in combination.

77. The use of any one of claims 59-72, wherein when said cancer is non-small cell lung cancer, the use further comprising use of a treatment selected from carboplatin, paclitaxel, docetaxel, gefitinib, erlotinib, afatinib, bevacizumab, ramucirumab, osimertinib, necitumumab, crizotinib, ceritinib, alectinib, and immune checkpoint inhibiting drugs including nivolumab, pembrolizumab, either individually or in combination.


78. The use of any one of claims 59-72, wherein when said cancer is bladder carcinoma, the use further comprising a use of a treatment selected from methotrexate, vincblastine, doxorubicin, cisplatin; carboplatin and paclitaxel; docetaxel, gemcitabine and cisplatin, either individually or in combination.

79. The use of any one of claims 59-72, wherein the determination of hypermethylation in the *NMT2* gene is performed by a method selected from the group consisting of PCR, methylation-specific PCR, realtime methylation-specific PCR, PCR assay using a methylation DNA-specific binding protein, quantitative PCR, DNA chip-based assay, pyrosequencing, and bisulfite sequencing.

80. The use of any one of claims 59-79, wherein said biological sample comprises, a tissue, a cell specimen or fluids includes detached tumor cells and/or free nucleic acids, a blood sample, a fractionated blood sample, a bone marrow sample, a biopsy sample, a frozen tissue sample, a fresh tissue specimen, a cell sample, and/or a paraffin embedded section.


81. The use of anyone of claims 59-80, wherein said subject is a human.

Figure 1 A



(Black curve above grey curve at intersection of axis.)

Figure 1 B



(Black curve is top curve; grey curve is bottom curve.)

Figure 1 C





Figure 1 D



3/20

Figure 1 E




Figure 1 F




Figure 1 G






Figure 1 H



“Low NMT2 Expressors” curve is bottom curve; “High NMT2 Expressors” is top curve.

Figure 2 A



6/20


Figure 2 B -- Epigenetic regulation of NMT2 in lymphoma cell lines and tumours.

Figure 2 C

Bisulfite sequencing of *DAPK1* promoter region reveals a highly methylated CpG containing promoter region only in malignant B lymphocytes




Figure 3 A




Figure 3 B




Figure 3 C





Figure 3 D



11/20

Figure 3 E




Figure 3 F






Figure 3 G




13/20

Figure 4 A



Statistical Significance for Kruskal-Wallis Dunn's test: ne = not evaluable; ns = not significant; * = $P < 0.05$, ** = $P < 0.01$, *** = $P < 0.001$, compared to Group 1
 TGI $\geq$ 60% indicates potential therapeutic activity

Figure 4B



15/20

Figure 4 C




Figure 5 A




Figure 5B




Figure 5 C




Figure 5 D




Figure 5 E





Figure 5 F



20/20

Figure 5 G




Figure 5 H





Figure 5 I



INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA2016/050846

A. CLASSIFICATION OF SUBJECT MATTER

IPC: **C12Q 1/68** (2006.01), **A61B 5/00** (2006.01), **A61B 5/06** (2006.01), **A61K 31/4439** (2006.01),
A61K 31/7088 (2006.01), **A61K 38/00** (2006.01) (more IPCs on the last page)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 Keywords used across the whole IPC

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used)

Databases: Questel-Orbit, CIPO Library Database, Scopus and Google

Keywords: NMT2, N-myristoyltransferase 2, cancer, methylation, methylated, DNA, promoter, gene, expression, CpG

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO2014/067002 A1 (BERTHIAUME, L.G. et al.) 8 May 2014 (08-05-2014) *Entire document*	1-3 and 32-81
Y	WO2013/013302 A1 (BERTHIAUME, L.G. et al.) 31 January 2013 (31-01-2013) *Entire document*	1-3 and 32-81
Y	KUANG, S-Q. et al., "Genome-wide identification of aberrantly methylated promoter associated CpG islands in acute lymphocytic leukemia". Leukemia, August 2008 (08-2008), Vol. 22(8), pp. 1529-1538. *Abstract, pages 1535-1537 containing Figure 3 and Table 3*	1-3 and 32-81

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* "A" "E" "L" "O" "P"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier application or patent but published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	"T" "X" "Y" "&"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
--------------------------------------	--	--------------------------	--

Date of the actual completion of the international search
09 September 2016 (09-09-2016)

Date of mailing of the international search report
14 September 2016 (14-09-2016)

Name and mailing address of the ISA/CA
 Canadian Intellectual Property Office
 Place du Portage I, C114 - 1st Floor, Box PCT
 50 Victoria Street
 Gatineau, Quebec K1A 0C9
 Facsimile No.: 819-953-2476

Authorized officer

Maria Mill (819) 639-6815

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA2016/050846**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of the first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claim Nos.: 4-31
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 4-31 are directed to a method for treatment of the human or animal body by surgery or therapy, which the International Searching Authority is not required to search under PCT Rule 39.1(iv). However, this Authority has carried out a search based on the alleged use of the methylation status of the NMT2 gene as a marker in methods for cancer patient treatment and treatment selection.
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claim Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CA2016/050846

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
WO2014067002A1	08 May 2014 (08-05-2014)	WO2014067002A1	08 May 2014 (08-05-2014)
		AU2013337569A1	07 May 2015 (07-05-2015)
		CA2890113A1	08 May 2014 (08-05-2014)
		CN104768550A	08 July 2015 (08-07-2015)
		EP2914261A1	09 September 2015 (09-09-2015)
		EP2914261A4	27 July 2016 (27-07-2016)
		IL238481D0	30 June 2015 (30-06-2015)
		JP2016506362A	03 March 2016 (03-03-2016)
		KR20150079949A	08 July 2015 (08-07-2015)
		MX2015005441A	21 July 2015 (21-07-2015)
		SG11201503187UA	28 May 2015 (28-05-2015)
		US2015275309A1	01 October 2015 (01-10-2015)
WO2013013302A1	31 January 2013 (31-01-2013)	WO2013013302A1	31 January 2013 (31-01-2013)
		WO2013013302A8	27 February 2014 (27-02-2014)
		AU2012286542A1	06 February 2014 (06-02-2014)
		CA2842443A1	31 January 2013 (31-01-2013)
		CN103826623A	28 May 2014 (28-05-2014)
		EP2734199A1	28 May 2014 (28-05-2014)
		EP2734199A4	01 April 2015 (01-04-2015)
		IL230575D0	31 March 2014 (31-03-2014)
		JP2014523440A	11 September 2014 (11-09-2014)
		KR20140072026A	12 June 2014 (12-06-2014)
		MX2014000661A	08 September 2014 (08-09-2014)
		RU2014101787A	27 August 2015 (27-08-2015)
		US2014227284A1	14 August 2014 (14-08-2014)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA2016/050846

A61K 39/395 (2006.01) , *A61P 35/00* (2006.01) , *C07H 21/04* (2006.01) , *G01N 33/48* (2006.01) ,
G01N 33/50 (2006.01)