



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION


(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/12/11
(87) Date publication PCT/PCT Publication Date: 2019/06/20
(85) Entrée phase nationale/National Entry: 2020/05/15
(86) N° demande PCT/PCT Application No.: US 2018/064879
(87) N° publication PCT/PCT Publication No.: 2019/118411
(30) Priorité/Priority: 2017/12/11 (US62/597,160)

(51) Cl.Int./Int.Cl. *A61K 31/5386* (2006.01),
A61K 45/06 (2006.01), *C07D 405/12* (2006.01),
C07D 498/18 (2006.01), *C07K 16/28* (2006.01)
(71) Demandeur/Applicant:
TRIPHASE ACCELERATOR U.S. CORPORATION, US
(72) Inventeur/Inventor:
MACLAREN, ANN, US
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : CONJUGUES ANTICORPS ANTI-CD22-MAYTANSINE, ASSOCIATIONS ET METHODES D'UTILISATION
CORRESPONDANTES
(54) Title: ANTI-CD22 ANTIBODY-MAYTANSINE CONJUGATES, COMBINATIONS, AND METHODS OF USE
THEREOF

FIG. 1



(57) **Abrégé/Abstract:**

The present disclosure provides methods for treating a cancer or resistant cancer with a combination of an anti-CD22 antibody-maytansine conjugate and one or more anti-cancer agents. The disclosure also encompasses methods for sensitizing a cancer with such combinations. Also provided are pharmaceutical compositions including such combinations.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

20 June 2019 (20.06.2019)



(10) International Publication Number

WO 2019/118411 A3

(51) International Patent Classification:

A61K 31/5386 (2006.01) C07D 498/18 (2006.01)

C07D 405/12 (2006.01) C07K 16/28 (2006.01)

(21) International Application Number:

PCT/US2018/064879

(22) International Filing Date:

11 December 2018 (11.12.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/597,160 11 December 2017 (11.12.2017) US

(71) Applicant: **TRIPHASE ACCELERATOR U.S. CORPORATION** [US/US]; 3366 N. Torrey Pines Court, #210, La Jolla, California 92037 (US).

(72) Inventor: **MACLAREN, Ann**; c/o Triphase Accelerator U.S. Corporation, 3366 N. Torrey Pines Court, #210, La Jolla, California 92037 (US).

(74) Agent: **FARMER, Dean** et al.; Cooley LLP, 1299 Pennsylvania Ave. Suite 700, Washington, District of Columbia 20004 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:


- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:

26 March 2020 (26.03.2020)

(54) Title: ANTI-CD22 ANTIBODY-MAYTANSINE CONJUGATES, COMBINATIONS, AND METHODS OF USE THEREOF

FIG. 1



(57) Abstract: The present disclosure provides methods for treating a cancer or resistant cancer with a combination of an anti-CD22 antibody-maytansine conjugate and one or more anti-cancer agents. The disclosure also encompasses methods for sensitizing a cancer with such combinations. Also provided are pharmaceutical compositions including such combinations.



WO 2019/118411 A3

ANTI-CD22 ANTIBODY-MAYTANSINE CONJUGATES, COMBINATIONS, AND METHODS OF USE THEREOF

RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application No. 62/597,160, filed on December 11, 2017, which is incorporated by reference herein in its entirety.

INCORPORATION OF SEQUENCE LISTING

[0002] The contents of the text file named “TRPS-04001WO_SeqList_ST25.txt”, which was created on December 10, 2018 and is 95 KB in size, are hereby incorporated by reference in their entirety.

INTRODUCTION

[0003] The field of protein-small molecule therapeutic conjugates has advanced greatly, providing a number of clinically beneficial drugs with the promise of providing more in the years to come. Protein-conjugate therapeutics can provide several advantages, due to, for example, specificity, multiplicity of functions and relatively low off-target activity, resulting in fewer side effects. Chemical modification of proteins may extend these advantages by rendering them more potent, stable, or multimodal.

[0004] A number of standard chemical transformations are commonly used to create and manipulate post-translational modifications on proteins. There are a number of methods where one is able to modify the side chains of certain amino acids selectively. For example, carboxylic acid side chains (aspartate and glutamate) may be targeted by initial activation with a water-soluble carbodiimide reagent and subsequent reaction with an amine. Similarly, lysine can be targeted through the use of activated esters or isothiocyanates, and cysteine thiols can be targeted with maleimides and α -halo-carbonyls.

[0005] One significant obstacle to the creation of a chemically altered protein therapeutic or reagent is the production of the protein in a biologically active, homogenous form. Conjugation of a drug or detectable label to a polypeptide can be difficult to control, resulting in a heterogeneous mixture of conjugates that differ in the number of drug molecules attached and in the position of chemical conjugation. In some instances, it may be desirable to control the site of conjugation and/or the drug or detectable label conjugated to the polypeptide using the tools

of synthetic organic chemistry to direct the precise and selective formation of chemical bonds on a polypeptide.

SUMMARY

[0005] The present disclosure provides methods for treating cancers and resistant cancers with an anti-CD22 antibody-maytansine conjugate in combination with one or more anti-cancer agents. The disclosure also encompasses methods for sensitizing cancers by treatment with such conjugates and anti-cancer agents.

[0006] In one aspect, the present disclosure provides a method for treating a cancer in a subject, the method comprising administering to the subject in need thereof a therapeutically effective amount of one or more anti-cancer agents, and a conjugate as described herein.

[0007] In one aspect, the present disclosure provides a method for treating a resistant cancer in a subject, the method comprising administering to the subject in need thereof a therapeutically effective amount of one or more anti-cancer agents, and a conjugate as described herein.

[0008] In one aspect, the present disclosure provides a method for sensitizing a cancer in a subject, the method comprising administering to the subject in need thereof a therapeutically effective amount of one or more anti-cancer agents, and a conjugate as described herein.

[0009] In one aspect, the present disclosure provides a pharmaceutical composition for treating a cancer, the pharmaceutical composition comprising one or more anti-cancer agents; a conjugate as described herein; and a pharmaceutically acceptable excipient. In one aspect, the present disclosure provides the use of the pharmaceutical composition for the manufacture of a medicament for treating a cancer. In one aspect, the present disclosure provides the use of the pharmaceutical composition for the manufacture of a medicament for treating a resistant cancer.

[0010] In one aspect, the present disclosure provides the use of the pharmaceutical composition for the manufacture of a medicament for sensitizing a cancer. In one aspect, the present disclosure provides the use of the pharmaceutical composition for treating a cancer.


[0011] In one aspect, the present disclosure provides the use of the pharmaceutical composition for treating a resistant cancer. In one aspect, the present disclosure provides the use of the pharmaceutical composition for sensitizing a cancer.

[0012] In one aspect, the present disclosure provides a method for treating a resistant cancer in a subject, the method comprising administering to the subject in need thereof a therapeutically effective amount of a conjugate as disclosed herein.

[0013] In one aspect, the present disclosure provides the use of a combination comprising one or more anti-cancer agents, and a conjugate as described herein in the manufacture of a medicament for treating a cancer in a subject in need thereof.

[0014] In one aspect, the present disclosure provides the use of a combination comprising one or more anti-cancer agents, and a conjugate as described herein for treating a cancer in a subject in need thereof.

[0015] In some embodiments of any of the above-aspects, the conjugate includes at least one modified amino acid residue of formula (I):



wherein

Z is CR⁴ or N;

R¹ is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl

, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R² and R³ are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R² and R³ are optionally cyclically linked to form a 5 or 6-membered heterocyclyl;

each R⁴ is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino,

substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

L is a linker comprising $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d-$, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4;

T^1 , T^2 , T^3 and T^4 are each independently selected from (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(EDA)_w$, $(PEG)_n$, $(AA)_p$, $-(CR^{13}OH)_h-$, piperidin-4-amino (4AP), an acetal group, a hydrazine, a disulfide, and an ester, wherein EDA is an ethylene diamine moiety, PEG is a polyethylene glycol or a modified polyethylene glycol, and AA is an amino acid residue, wherein w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12;

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$ and $-P(O)OH-$, wherein q is an integer from 1 to 6;

each R^{13} is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl;

each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

W^1 is a maytansinoid; and

W^2 is an anti-CD22 antibody.


[0016] In certain embodiments,

T^1 is selected from a (C_1-C_{12}) alkyl and a substituted (C_1-C_{12}) alkyl;

T^2 , T^3 and T^4 are each independently selected from $(EDA)_w$, $(PEG)_n$, (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(AA)_p$, $-(CR^{13}OH)_h-$, piperidin-4-amino (4AP), an acetal group, a hydrazine, and an ester; and


V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$, and $-P(O)OH-$;


wherein:

(PEG)_n is , where n is an integer from 1 to 30;

EDA is an ethylene diamine moiety having the following structure:

where y is an integer from 1 to 6 and r is 0 or 1;



piperidin-4-amino is ;

each R¹² and R¹⁵ is independently selected from hydrogen, an alkyl, a substituted alkyl, a polyethylene glycol moiety, an aryl and a substituted aryl, wherein any two adjacent R¹² groups may be cyclically linked to form a piperazinyl ring; and

R¹³ is selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl.


[0017] In certain embodiments, T¹, T², T³ and T⁴, and V¹, V², V³ and V⁴ are selected from the following table:


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-CO-	(EDA) _w	-	-	-


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-SO ₂ -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(PEG) _n	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(CR ¹³ OH) _h	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	substituted (C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-SO ₂ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-	(CR ¹³ OH) _h	- CONR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	- NR ¹⁵ -

T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	- P(O)OH -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	(C ₁ -C ₁₂)alkyl	- NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	-	-

[0018] In certain embodiments, L is selected from one of the following structures:







wherein

each f is independently 0 or an integer from 1 to 12;

each y is independently 0 or an integer from 1 to 20;

each n is independently 0 or an integer from 1 to 30;


each p is independently 0 or an integer from 1 to 20;

each h is independently 0 or an integer from 1 to 12;

each R is independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

each R' is independently H, a sidechain group of an amino acid, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.


[0019] In certain embodiments, the maytansinoid is of the formula:



where ~~~ indicates the point of attachment between the maytansinoid and L.

[0020] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is 4AP, V² is -CO-, T³ is (C₁-C₁₂)alkyl, V³ is -CO-, T⁴ is absent and V⁴ is absent.

[0021] In certain embodiments, the linker, L, includes the following structure:




wherein


each f is independently an integer from 1 to 12; and

n is an integer from 1 to 30.

[0022] In certain embodiments, the conjugate includes the following structure:




[0023] In certain embodiments, the conjugate includes the following structure:



[0024] In certain embodiments, the anti-CD22 antibody binds an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C.

[0025] In certain embodiments, the anti-CD22 antibody comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue;

Z^{30} is a basic amino acid or an aliphatic amino acid;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), with the proviso that when the sequence is at the N-terminus of the conjugate, X^1 is present; and


X^2 and X^3 are each independently any amino acid, (e.g., any naturally-occurring amino acid).

[0026] In certain embodiments, the sequence is **L(FGly')TPSR** (SEQ ID NO: 185).

[0027] In certain embodiments, Z^{30} is selected from R, K, H, A, G, L, V, I, and P; X^1 is selected from L, M, S, and V; and X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

[0028] In certain embodiments, the modified amino acid residue is positioned at a C-terminus of a heavy chain constant region of the anti-CD22 antibody.

[0029] In certain embodiments, the heavy chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, (e.g., any naturally-occurring amino acid), and


wherein the sequence is C-terminal to the amino acid sequence **SLSLSPG** (SEQ ID NO: 186).

[0030] In certain embodiments, the heavy chain constant region comprises the sequence **SPGSL(FGly')TPSRGS** (SEQ ID NO: 184).

[0031] In certain embodiments, Z³⁰ is selected from R, K, H, A, G, L, V, I, and P; X¹ is selected from L, M, S, and V; and X² and X³ are each independently selected from S, T, A, V, G, and C.

[0032] In certain embodiments, the modified amino acid residue is positioned in a light chain constant region of the anti-CD22 antibody.

[0033] In certain embodiments, the light chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X^2 and X^3 are each independently any amino acid, (e.g., any naturally-occurring amino acid), and


wherein the sequence C-terminal to the sequence **KVDNAL** (SEQ ID NO: 58), and/or is N-terminal to the sequence **QSGNSQ** (SEQ ID NO: 59).

[0034] In certain embodiments, the light chain constant region comprises the sequence **KVDNAL(FGly')TPSRQSGNSQ** (SEQ ID NO: 60).

[0035] In certain embodiments, Z^{30} is selected from R, K, H, A, G, L, V, I, and P; X^1 is selected from L, M, S, and V; and X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

[0036] In certain embodiments, the modified amino acid residue is positioned in a heavy chain CH1 region of the anti-CD22 antibody.

[0037] In certain embodiments, the heavy chain CH1 region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue;

Z^{30} is a basic amino acid or an aliphatic amino acid;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), with the proviso that when the sequence is at the N-terminus of the conjugate, X^1 is present; and

X^2 and X^3 are each independently any amino acid, (e.g., any naturally-occurring amino acid), and

wherein the sequence is C-terminal to the amino acid sequence **SWNSGA** (SEQ ID NO: 61) and/or is N-terminal to the amino acid sequence **GVHTEP** (SEQ ID NO: 62).

[0038] In certain embodiments, the heavy chain CH1 region comprises the sequence **SWNSGAL(FGly')TPSRGVHTEP** (SEQ ID NO: 63).

[0039] In certain embodiments, Z^{30} is selected from R, K, H, A, G, L, V, I, and P; X^1 is selected from L, M, S, and V; and X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

[0045] In some embodiments, the one or more anti-cancer agents is selected from cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“CHOP”);

cyclophosphamide, vincristine sulfate, procarbazine hydrochloride, and prednisone (“COPP”); cyclophosphamide, vincristine sulfate, and prednisone (“CVP”); etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

[0046] In some embodiments, the one or more anti-cancer agents comprises rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”).

[0047] In some embodiments, the cancer is affiliated with dysregulation of BCR signaling. In some embodiments, the cancer is affiliated with dysregulation of BCR signaling, and the cancer is responsive to B-cell depletion.

[0048] In some embodiments, the cancer is lymphoma. In some embodiments, the cancer is a B-cell lymphoma. In some embodiments, the cancer is selected from Burkitt’s lymphoma, diffuse large B-cell lymphoma, Hodgkin’s lymphoma, and non-Hodgkin’s lymphoma. In some embodiments, the cancer is non-Hodgkin’s lymphoma. In some embodiments, the cancer is selected from marginal zone lymphoma, mantle cell lymphoma, follicular lymphoma, and primary central nervous system lymphoma. In some embodiments, the cancer is mantle cell lymphoma. In some embodiments, the cancer is diffuse large B-cell lymphoma. In some embodiments, the cancer is follicular lymphoma. In some embodiments, the cancer is marginal zone lymphoma. In some embodiments, the cancer is leukemia.

[0049] In some embodiments, the cancer is selected from chronic myeloproliferative syndrome, acute myelogenous leukemia, chronic lymphocytic leukemia, small lymphocytic leukemia, hairy cell leukemia, and acute lymphoblastic leukemia.

[0050] In some embodiments, the cancer is resistant to treatment with one or more anti-cancer agents selected from abiraterone, methotrexate, brentuximab vedotin, copanlisib, copanlisib hydrochloride, chlorambucil, nelarabine, axicabtagene ciloleucel, carmustine, belinostat, bendamustine, bendamustine hydrochloride, tositumomab iodine-131, tositumomab,

bleomycin, bortezomib, acalabrutinib, cyclophosphamide, cytarabine, cytarabine liposome, denileukin diftitox, cytarabine liposome, dexamethasone, doxorubicin, doxorubicin hydrochloride, methotrexate, pralatrexate, ofatumamb, obinutuzumab, ocrelizumab, ibrutinib, tiuxetan, ibrutinib, idelalisib, recombinant interferon alfa-2b, romidespsin, lenalidomide, mechlorethamine hydrochloride, plerixafor, prednisone, rituximab, rituximab and hyaluronidase human, bortezomib, vinblastine, vinblastine sulfate, vincristine, vincristine sulfate, and vorinostat.

[0051] In some embodiments, the cancer is resistant to treatment with one or more anti-cancer agents selected from: cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“CHOP”); cyclophosphamide, vincristine sulfate, procarbazine hydrochloride, and prednisone (“COPP”); cyclophosphamide, vincristine sulfate, and prednisone (“CVP”); etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

[0052] In some embodiments, the cancer is resistant to treatment with rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”).

BRIEF DESCRIPTION OF THE DRAWINGS

[0053] FIG. 1, panel A, shows a formylglycine-generating enzyme (FGE) recognition sequence inserted at the desired location along the antibody backbone using standard molecular biology techniques. Upon expression, FGE, which is endogenous to eukaryotic cells, catalyzes the conversion of the Cys within the consensus sequence to a formylglycine residue (FGly). FIG. 1, panel B, shows antibodies carrying aldehyde moieties (2 per antibody) reacted with a Hydrazino-*iso*-Pictet-Spengler (HIPS) linker and payload to generate a site-specifically conjugated ADC. FIG. 1, panel C, shows HIPS chemistry, which proceeds through an

intermediate hydrazone ion followed by intramolecular alkylation with a nucleophilic indole to generate a stable C-C bond.

[0054] FIG. 2 shows a hydrophobic interaction column (HIC) trace of an aldehyde-tagged anti-CD22 antibody conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker, according to embodiments of the present disclosure.

[0055] FIG. 3 shows a HIC trace of an aldehyde-tagged anti-CD22 antibody conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker, according to embodiments of the present disclosure.

[0056] FIG. 4 shows a reversed phase chromatography (PLRP) trace of an aldehyde-tagged anti-CD22 antibody conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker, according to embodiments of the present disclosure.

[0057] FIG. 5 shows a graph of analytical size exclusion chromatography (SEC) analysis of an aldehyde-tagged anti-CD22 antibody conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker, according to embodiments of the present disclosure.

[0058] FIG. 6A shows a graph indicating the *in vitro* potency against WSU-DLCL2 cells (% viability vs. Log antibody-drug conjugate (ADC) concentration (nM)) for anti-CD22 ADCs conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker, according to embodiments of the present disclosure. FIG. 6B shows a graph of *in vitro* potency against Ramos cells (% viability vs. Log antibody-drug conjugate (ADC) concentration (nM)) for anti-CD22 ADCs conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker, according to embodiments of the present disclosure.

[0059] FIG. 7 shows a graph indicating the *in vivo* efficacy against a WSU-DLCL2 xenograft model (mean tumor volume (mm³) vs. days) for anti-CD22 ADCs conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker, according to embodiments of the present disclosure.

[0060] FIG. 8A-8C provide amino acid sequences of CD22 isoforms, i.e., isoform2 (SEQ ID NO: 1), isoform4 (SEQ ID NO: 2), isoform1 (SEQ ID NO: 3), and isoform3 (SEQ ID NO: 4).

[0061] FIG. 9A depicts a site map showing possible modification sites for generation of an aldehyde tagged Ig polypeptide. The upper sequence is the amino acid sequence of the conserved region of an IgG1 light chain polypeptide (SEQ ID NO: 5) and shows possible modification sites in an Ig light chain; the lower sequence is the amino acid sequence of the

conserved region of an Ig heavy chain polypeptide (SEQ ID NO: 6); GenBank Accession No. AAG00909) and shows possible modification sites in an Ig heavy chain. The heavy and light chain numbering is based on the full-length heavy and light chains.

[0062] FIG. 9B depicts an alignment of immunoglobulin heavy chain constant regions for IgG1 (SEQ ID NO: 7), IgG2 (SEQ ID NO: 8), IgG3 (SEQ ID NO: 9), IgG4 (SEQ ID NO: 10), and IgA (SEQ ID NO: 11), showing modification sites at which aldehyde tags can be provided in an immunoglobulin heavy chain. The heavy and light chain numbering is based on the full- heavy and light chains.

[0063] FIG. 9C depicts an alignment of immunoglobulin light chain constant regions, i.e., *homo sapiens kappa* (SEQ ID NO: 12), GenBank Accession No. CAA75031.1; *homo sapiens kappa* (SEQ ID NO: 13), GenBank Accession No. BAC0168.1; *homo sapiens lambda* (SEQ ID NO: 14), GenBank Accession No. CAA75033; *Mus musculus* (SEQ ID NO: 15), GenBank Accession No. AAB09710.1; *Rattus norvegicus* (SEQ ID NO: 16), GenBank Accession No. AAD10133, showing modification sites at which aldehyde tags can be provided in an immunoglobulin light chain.

[0064] FIG. 10 shows illustrations of ELISA formats for detection of various analytes, according to embodiments of the present disclosure.

[0065] FIG. 11 – an anti-CD22 ADC according to the present disclosure was highly monomeric, had a average DAR of 1.8, and included a single light and heavy chain species. The anti-CD22 ADC was analyzed by (FIG. 11, panel A) Size exclusion chromatography to assess percent monomer (99.2%), and by hydrophobic interaction (HIC; FIG. 11, panel B) and reversed-phase (PLRP) chromatography (FIG. 11, panel C) to assess the drug-to-antibody ratio (DAR), which was 1.8.

[0066] FIG. 12 – an anti-CD22 ADC according to the present disclosure bound to human CD22 protein equally well as the wild-type anti-CD22 antibody. A competitive ELISA was used to compare the binding of the anti-CD22 ADC to the wild-type (WT) anti-CD22 antibody. The data are presented as the mean $\pm$ S.D. (n = 4).

[0067] FIG. 13 – an anti-CD22 ADC according to the present disclosure mediated the internalization of CD22 similarly to the wild-type anti-CD22 antibody. The NHL cell lines, Ramos, Granta-519, and WSU-DLCL2 were used to compare the internalization of cell surface CD22 as mediated by binding to either WT anti-CD22 or CAT-02-106.

[0068] FIG. 14 – an anti-CD22 ADC according to the present disclosure was equally potent against parental and MDR1-expressing NHL tumor cells *in vitro*. Ramos and WSU-DLCL2 parental (WT) cells (FIG. 14, panel A and panel C) and variants of those lines that were engineered to express MDR1 (MDR1+, FIG. 14, panel B and panel D) were used as targets for *in vitro* cytotoxicity studies of anti-CD22 ADC activity. Free maytansine and an α CD22 ADC made with the CAT-02 antibody but conjugated to maytansine using a valine-citrulline cleavable linker were used as controls. In an additional control experiment, the MDR1 inhibitor, cyclosporin, was added to WT or MDR1+ WSU-DLCL2 cells (FIG. 14, panel E and panel F). The data are presented as the mean $\pm$ S.D. (n = 2).

[0069] FIG. 15 – an anti-CD22 ADC according to the present disclosure did not mediate off-target cytotoxicity. The gastric tumor cell line, NCI-N87, was incubated *in vitro* for 5 days in the presence of increasing concentrations of the anti-CD22 ADC. Then, cell viability was assessed using an MTS-based method. The data are presented as the mean $\pm$ S.D. (n = 2).

[0070] FIG. 16 – The anti-CD22 ADC-related ADC, anti-HER2 conjugated to a HIPS-4AP-maytansine linker payload, did not induce bystander killing. *In vitro* cytotoxicity studies were conducted using HER2+ NCI-N87 cells, HER2- Ramos cells, or a coculture of both cells as targets. Free maytansine (2 nM) and anti-HER2 conjugated to MMAE via a cleavable valine-citrulline (vc) linker (2 nM payload), were used as positive controls for bystander killing. Anti-HER2 ADC was dosed at 2 nM payload. The data are presented as the mean $\pm$ S.D. (n = 2).

[0071] FIG. 17 – an anti-CD22 ADC according to the present disclosure was efficacious *in vivo* against the NHL-derived WSU-DLCL2 and Ramos xenograft models. Female CB17 ICR SCID mice (8/group) bearing WSU-DLCL2 xenografts were treated with vehicle alone or with the anti-CD22 ADC as either a (FIG. 17, panel A) single 10 mg/kg dose or (FIG. 17, panel B) as multiple 10 mg/kg doses delivered every four days for a total of four doses (q4d x 4). Treatment was initiated when tumors reached an average size of 118 or 262 mm³ for the single or multidose studies, respectively. (FIG. 17, panel C) Female CB17 ICR SCID mice (12/group) bearing Ramos xenografts were treated with vehicle alone, or with 5 or 10 mg/kg CAT-02-106 q4d x 4. Dosing was initiated when tumors reached an average size of 246 mm³. The data are presented as the mean $\pm$ S.E.M.

[0072] FIG. 18 – Ramos and WSU-DLCL2 cells expressed different levels of cell surface CD22. Ramos and WSU-DLCL2 cells were incubated with a fluorescein-labeled anti-CD22

antibody and then analyzed by flow cytometry. The mean fluorescence intensity of the FL1 channel (detecting fluorescein) for each cell type is shown in the graph.

[0073] FIG. 19 - Mouse body weights were not affected by treatment with an anti-CD22 ADC according to the present disclosure. Mean body weights of mice in the xenograft efficacy studies are shown. (FIG. 19, panel A) Single dose WSU-DLCL2 study; (FIG. 19, panel B) Multidose WSU-DLCL2 study; (FIG. 19, panel C) Ramos study. Error bars indicate S.D.

[0074] FIG. 20 – an anti-CD22 ADC according to the present disclosure can be dosed in rats up to 60 mg/kg with minimal effects. Sprague-Dawley rats (5/group) received a 6, 20, 40, or 60 mg/kg dose of CAT-02-106 followed by a 12 day observation period. (FIG. 20, panel A) Body weight was monitored at the times indicated. (FIG. 20, panel B) Alanine aminotransferase (ALT), and (FIG. 20, panel C) platelet counts were assessed at 5 and 12 days post-dose. The data are presented as the mean $\pm$ S.D.

[0075] FIG. 21 – an anti-CD22 ADC according to the present disclosure bound specifically to cynomolgus monkey B cells. Cynomolgus peripheral blood lymphocytes were gated according to their forward and side scatter profiles (upper left). Cells were incubated with either fluorescein-isothiocyanate (FITC)-conjugated streptavidin (SA) alone (upper right), or with biotinylated anti-CD22 ADC followed by FITC SA. Coincubation with antibodies recognizing T cells (CD3, lower left) or B cells (CD20, lower right) demonstrated specificity of CAT-02-106 binding to a B-cell population.

[0076] FIG. 22 – an anti-CD22 ADC according to the present disclosure demonstrated B cell-specific reactivity in human and cynomolgus monkey tissues. The anti-CD22 ADC bound to B-cell rich regions of the spleen (top). Heart tissues were negative for staining (middle). Lung sections were negative with the exception of scattered leukocytes (bottom).

[0077] FIG. 23 – Cynomolgus monkeys display no observed adverse effects with a repeat 60 mg/kg dose of an anti-CD22 ADC according to the present disclosure. Cynomolgus monkeys (2/sex/group) were given 10, 30, or 60 mg/kg of the anti-CD22 ADC once every three weeks for a total of two doses followed by a 21 day observation period. (FIG. 23, panel A) Aspartate transaminase (AST), (FIG. 23, panel B) alanine aminotransferase (ALT), (FIG. 23, panel C) platelets, and (FIG. 23, panel D) monocytes were monitored at the times indicated. The data are presented as the mean $\pm$ S.D.

[0078] FIG. 24 (panel A and panel B) – Treatment with an anti-CD22 ADC according to the present disclosure reduced peripheral B cell populations in cynomolgus monkeys. Peripheral blood mononuclear cells from cynomolgus monkeys enrolled in the toxicity study were monitored by flow cytometry to detect the ratio of B cells (CD20+), T cells (CD3+), and NK cells (CD20-/CD3-) observed in animals pre-dose and at days 7, 14, 28, and 35. The data are presented as the mean $\pm$ S.D.

[0079] FIG. 25 – an anti-CD22 ADC according to the present disclosure displayed very high *in vivo* stability as shown by a rat pharmacokinetic study. Sprague-Dawley rats (3/group) were given a single i.v. bolus dose of 3 mg/kg anti-CD22 ADC. Plasma samples were collected at the designated times and were analyzed (as shown in FIG. 10) for total antibody, total conjugate, and total ADC concentrations.

[0080] FIG. 26 shows Table 3: summary of mean ($\pm$ SD) pharmacokinetic and toxicokinetic (TK) parameters of total ADC values in animals dosed with an anti-CD22 ADC according to embodiments of the present disclosure.

[0081] FIG. 27 is a graph depicting tumor regression in a Granta-519 xenograft model. The graph compares dosing regimens of an anti-CD22 ADC of the present invention and compares anti-CD22 ADC treatment to treatment with rituximab.

[0082] FIG. 28 is a graph depicting tumor regression in a Granta-519 xenograft model. The graph compares treatments with rituximab, an anti-CD22 ADC of the present invention, R-CHOP, and treatment with R-CHOP followed by treatment with the anti-CD22 ADC.

DETAILED DESCRIPTION

[0083] The present disclosure provides methods of treating a cancer such as a resistant cancer using an anti-CD22 antibody-maytansine conjugate, for example in combination with one or more anti-cancer agents. Embodiments of each are described in more detail in the sections below.

DEFINITIONS

[0084] The following terms have the following meanings unless otherwise indicated. Any undefined terms have their art recognized meanings.

[0085] “Alkyl” refers to monovalent saturated aliphatic hydrocarbyl groups having from 1 to 10 carbon atoms and such as 1 to 6 carbon atoms, or 1 to 5, or 1 to 4, or 1 to 3 carbon atoms. This term includes, by way of example, linear and branched hydrocarbyl groups such as methyl

(CH₃-), ethyl (CH₃CH₂-), n-propyl (CH₃CH₂CH₂-), isopropyl ((CH₃)₂CH-), n-butyl (CH₃CH₂CH₂CH₂-), isobutyl ((CH₃)₂CHCH₂-), sec-butyl ((CH₃)(CH₃CH₂)CH-), t-butyl ((CH₃)₃C-), n-pentyl (CH₃CH₂CH₂CH₂CH₂-), and neopentyl ((CH₃)₃CCH₂-).

[0086] The term “substituted alkyl” refers to an alkyl group as defined herein wherein one or more carbon atoms in the alkyl chain (except the C₁ carbon atom) have been optionally replaced with a heteroatom such as -O-, -N-, -S-, -S(O)_n- (where n is 0 to 2), -NR- (where R is hydrogen or alkyl) and having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, oxo, thioketo, carboxyl, carboxylalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-aryl, -SO₂-heteroaryl, and -NR^aR^b, wherein R^a and R^b may be the same or different and are chosen from hydrogen, optionally substituted alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, aryl, heteroaryl and heterocyclic.

[0087] “Alkylene” refers to divalent aliphatic hydrocarbyl groups preferably having from 1 to 6 and more preferably 1 to 3 carbon atoms that are either straight-chained or branched, and which are optionally interrupted with one or more groups selected from -O-, -NR¹⁰-, -NR¹⁰C(O)-, -C(O)NR¹⁰- and the like. This term includes, by way of example, methylene (-CH₂-), ethylene (-CH₂CH₂-), n-propylene (-CH₂CH₂CH₂-), iso-propylene (-CH₂CH(CH₃)-), (-C(CH₃)₂CH₂CH₂-), (-C(CH₃)₂CH₂C(O)-), (-C(CH₃)₂CH₂C(O)NH-), (-CH(CH₃)CH₂-), and the like.

[0088] “Substituted alkylene” refers to an alkylene group having from 1 to 3 hydrogens replaced with substituents as described for carbons in the definition of “substituted” below.

[0089] The term “alkane” refers to alkyl group and alkylene group, as defined herein.

[0090] The term “alkylaminoalkyl”, “alkylaminoalkenyl” and “alkylaminoalkynyl” refers to the groups R¹NHR² - where R¹ is alkyl group as defined herein and R² is alkylene, alkenylene or alkynylene group as defined herein.

[0091] The term “alkaryl” or “aralkyl” refers to the groups -alkylene-aryl and -substituted alkylene-aryl where alkylene, substituted alkylene and aryl are defined herein.

[0092] “Alkoxy” refers to the group -O-alkyl, wherein alkyl is as defined herein. Alkoxy includes, by way of example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, t-butoxy, sec-

butoxy, n-pentoxy, and the like. The term “alkoxy” also refers to the groups alkenyl-O-, cycloalkyl-O-, cycloalkenyl-O-, and alkynyl-O-, where alkenyl, cycloalkyl, cycloalkenyl, and alkynyl are as defined herein.

[0093] The term “substituted alkoxy” refers to the groups substituted alkyl-O-, substituted alkenyl-O-, substituted cycloalkyl-O-, substituted cycloalkenyl-O-, and substituted alkynyl-O- where substituted alkyl, substituted alkenyl, substituted cycloalkyl, substituted cycloalkenyl and substituted alkynyl are as defined herein.

[0094] The term “alkoxyamino” refers to the group –NH-alkoxy, wherein alkoxy is defined herein.

[0095] The term “haloalkoxy” refers to the groups alkyl-O- wherein one or more hydrogen atoms on the alkyl group have been substituted with a halo group and include, by way of examples, groups such as trifluoromethoxy, and the like.

[0096] The term “haloalkyl” refers to a substituted alkyl group as described above, wherein one or more hydrogen atoms on the alkyl group have been substituted with a halo group. Examples of such groups include, without limitation, fluoroalkyl groups, such as trifluoromethyl, difluoromethyl, trifluoroethyl and the like.

[0097] The term “alkylalkoxy” refers to the groups -alkylene-O-alkyl, alkylene-O-substituted alkyl, substituted alkylene-O-alkyl, and substituted alkylene-O-substituted alkyl wherein alkyl, substituted alkyl, alkylene and substituted alkylene are as defined herein.

[0098] The term “alkylthioalkoxy” refers to the group -alkylene-S-alkyl, alkylene-S-substituted alkyl, substituted alkylene-S-alkyl and substituted alkylene-S-substituted alkyl wherein alkyl, substituted alkyl, alkylene and substituted alkylene are as defined herein.

[0099] “Alkenyl” refers to straight chain or branched hydrocarbyl groups having from 2 to 6 carbon atoms and preferably 2 to 4 carbon atoms and having at least 1 and preferably from 1 to 2 sites of double bond unsaturation. This term includes, by way of example, bi-vinyl, allyl, and but-3-en-1-yl. Included within this term are the cis and trans isomers or mixtures of these isomers.

[00100] The term “substituted alkenyl” refers to an alkenyl group as defined herein having from 1 to 5 substituents, or from 1 to 3 substituents, selected from alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano,

halogen, hydroxyl, oxo, thioketo, carboxyl, carboxylalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl.

[00101] “Alkynyl” refers to straight or branched monovalent hydrocarbyl groups having from 2 to 6 carbon atoms and preferably 2 to 3 carbon atoms and having at least 1 and preferably from 1 to 2 sites of triple bond unsaturation. Examples of such alkynyl groups include acetylenyl (-C≡CH), and propargyl (-CH₂C≡CH).

[00102] The term “substituted alkynyl” refers to an alkynyl group as defined herein having from 1 to 5 substituents, or from 1 to 3 substituents, selected from alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, oxo, thioketo, carboxyl, carboxylalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, and -SO₂-heteroaryl.

[00103] “Alkynyloxy” refers to the group -O-alkynyl, wherein alkynyl is as defined herein. Alkynyloxy includes, by way of example, ethynyloxy, propynyloxy, and the like.

[00104] “Acyl” refers to the groups H-C(O)-, alkyl-C(O)-, substituted alkyl-C(O)-, alkenyl-C(O)-, substituted alkenyl-C(O)-, alkynyl-C(O)-, substituted alkynyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, cycloalkenyl-C(O)-, substituted cycloalkenyl-C(O)-, aryl-C(O)-, substituted aryl-C(O)-, heteroaryl-C(O)-, substituted heteroaryl-C(O)-, heterocyclyl-C(O)-, and substituted heterocyclyl-C(O)-, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein. For example, acyl includes the “acetyl” group CH₃C(O)-

[00105] “Acylamino” refers to the groups -NR²⁰C(O)alkyl, -NR²⁰C(O)substituted alkyl, N R²⁰C(O)cycloalkyl, -NR²⁰C(O)substituted cycloalkyl, -

NR²⁰C(O)cycloalkenyl, -NR²⁰C(O)substituted cycloalkenyl, -NR²⁰C(O)alkenyl, -NR²⁰C(O)substituted alkenyl, -NR²⁰C(O)alkynyl, -NR²⁰C(O)substituted alkynyl, -NR²⁰C(O)aryl, -NR²⁰C(O)substituted aryl, -NR²⁰C(O)heteroaryl, -NR²⁰C(O)substituted heteroaryl, -NR²⁰C(O)heterocyclic, and -NR²⁰C(O)substituted heterocyclic, wherein R²⁰ is hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

[00106] “Aminocarbonyl” or the term “aminoacyl” refers to the group -C(O)NR²¹R²², wherein R²¹ and R²² independently are selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R²¹ and R²² are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

[00107] “Aminocarbonylamino” refers to the group -NR²¹C(O)NR²²R²³ where R²¹, R²², and R²³ are independently selected from hydrogen, alkyl, aryl or cycloalkyl, or where two R groups are joined to form a heterocyclyl group.

[00108] The term “alkoxycarbonylamino” refers to the group -NRC(O)OR where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclyl wherein alkyl, substituted alkyl, aryl, heteroaryl, and heterocyclyl are as defined herein.

[00109] The term “acyloxy” refers to the groups alkyl-C(O)O-, substituted alkyl-C(O)O-, cycloalkyl-C(O)O-, substituted cycloalkyl-C(O)O-, aryl-C(O)O-, heteroaryl-C(O)O-, and heterocyclyl-C(O)O- wherein alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, heteroaryl, and heterocyclyl are as defined herein.

[00110] “Aminosulfonyl” refers to the group -SO₂NR²¹R²², wherein R²¹ and R²² independently are selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl,

substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and where R^{21} and R^{22} are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group and alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

[00111] “Sulfonylamino” refers to the group $-NR^{21}SO_2R^{22}$, wherein R^{21} and R^{22} independently are selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R^{21} and R^{22} are optionally joined together with the atoms bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

[00112] “Aryl” or “Ar” refers to a monovalent aromatic carbocyclic group of from 6 to 18 carbon atoms having a single ring (such as is present in a phenyl group) or a ring system having multiple condensed rings (examples of such aromatic ring systems include naphthyl, anthryl and indanyl) which condensed rings may or may not be aromatic, provided that the point of attachment is through an atom of an aromatic ring. This term includes, by way of example, phenyl and naphthyl. Unless otherwise constrained by the definition for the aryl substituent, such aryl groups can optionally be substituted with from 1 to 5 substituents, or from 1 to 3 substituents, selected from acyloxy, hydroxy, thiol, acyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido, carboxyl, carboxylalkyl, cyano, halogen, nitro, heteroaryl, heteroaryloxy, heterocyclyl, heterocycloxy, aminoacyloxy, oxyacylamino, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl and trihalomethyl.

[00113] “Aryloxy” refers to the group –O-aryl, wherein aryl is as defined herein, including, by way of example, phenoxy, naphthoxy, and the like, including optionally substituted aryl groups as also defined herein.

[00114] “Amino” refers to the group –NH₂.

[00115] The term “substituted amino” refers to the group –NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, alkenyl, substituted alkenyl, cycloalkenyl, substituted cycloalkenyl, alkynyl, substituted alkynyl, aryl, heteroaryl, and heterocyclyl provided that at least one R is not hydrogen.

[00116] The term “azido” refers to the group –N₃.

[00117] “Carboxyl,” “carboxy” or “carboxylate” refers to –CO₂H or salts thereof.

[00118] “Carboxyl ester” or “carboxy ester” or the terms “carboxyalkyl” or “carboxylalkyl” refers to the groups –C(O)O-alkyl, –C(O)O-substituted alkyl, –C(O)O-alkenyl, –C(O)O-substituted alkenyl, –C(O)O-alkynyl, –C(O)O-substituted alkynyl, –C(O)O-aryl, –C(O)O-substituted aryl, –C(O)O-cycloalkyl, –C(O)O-substituted cycloalkyl, –C(O)O-cycloalkenyl, –C(O)O-substituted cycloalkenyl, –C(O)O-heteroaryl, –C(O)O-substituted heteroaryl, –C(O)O-heterocyclic, and –C(O)O-substituted heterocyclic, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

[00119] “(Carboxyl ester)oxy” or “carbonate” refers to the groups –O-C(O)O-alkyl, –O-C(O)O-substituted alkyl, –O-C(O)O-alkenyl, –O-C(O)O-substituted alkenyl, –O-C(O)O-alkynyl, –O-C(O)O-substituted alkynyl, –O-C(O)O-aryl, –O-C(O)O-substituted aryl, –O-C(O)O-cycloalkyl, –O-C(O)O-substituted cycloalkyl, –O-C(O)O-cycloalkenyl, –O-C(O)O-substituted cycloalkenyl, –O-C(O)O-heteroaryl, –O-C(O)O-substituted heteroaryl, –O-C(O)O-heterocyclic, and –O-C(O)O-substituted heterocyclic, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

[00120] “Cyano” or “nitrile” refers to the group –CN.

[00121] “Cycloalkyl” refers to cyclic alkyl groups of from 3 to 10 carbon atoms having single or multiple cyclic rings including fused, bridged, and spiro ring systems. Examples of suitable cycloalkyl groups include, for instance, adamantyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl and the like. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, and the like, or multiple ring structures such as adamantanyl, and the like.

[00122] The term “substituted cycloalkyl” refers to cycloalkyl groups having from 1 to 5 substituents, or from 1 to 3 substituents, selected from alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, oxo, thioketo, carboxyl, carboxylalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl.

[00123] “Cycloalkenyl” refers to non-aromatic cyclic alkyl groups of from 3 to 10 carbon atoms having single or multiple rings and having at least one double bond and preferably from 1 to 2 double bonds.

[00124] The term “substituted cycloalkenyl” refers to cycloalkenyl groups having from 1 to 5 substituents, or from 1 to 3 substituents, selected from alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxyl, carboxylalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl.

[00125] “Cycloalkynyl” refers to non-aromatic cycloalkyl groups of from 5 to 10 carbon atoms having single or multiple rings and having at least one triple bond.

[00126] “Cycloalkoxy” refers to -O-cycloalkyl.

[00127] “Cycloalkenyloxy” refers to -O-cycloalkenyl.

[00128] “Halo” or “halogen” refers to fluoro, chloro, bromo, and iodo.

[00129] “Hydroxy” or “hydroxyl” refers to the group –OH.

[00130] “Heteroaryl” refers to an aromatic group of from 1 to 15 carbon atoms, such as from 1 to 10 carbon atoms and 1 to 10 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur within the ring. Such heteroaryl groups can have a single ring (such as, pyridinyl, imidazolyl or furyl) or multiple condensed rings in a ring system (for example as in groups such as, indoliziny, quinoliny, benzofuran, benzimidazolyl or benzothienyl), wherein at least one ring within the ring system is aromatic. To satisfy valence requirements, any heteroatoms in such heteroaryl rings may or may not be bonded to H or a substituent group, e.g., an alkyl group or other substituent as described herein. In certain embodiments, the nitrogen and/or sulfur ring atom(s) of the heteroaryl group are optionally oxidized to provide for the N-oxide (N→O), sulfinyl, or sulfonyl moieties. This term includes, by way of example, pyridinyl, pyrrolyl, indolyl, thiophenyl, and furanyl. Unless otherwise constrained by the definition for the heteroaryl substituent, such heteroaryl groups can be optionally substituted with 1 to 5 substituents, or from 1 to 3 substituents, selected from acyloxy, hydroxy, thiol, acyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, substituted cycloalkyl, substituted cycloalkenyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido, carboxyl, carboxylalkyl, cyano, halogen, nitro, heteroaryl, heteroaryloxy, heterocyclyl, heterocycloxy, aminoacyloxy, oxyacylamino, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl and -SO₂-heteroaryl, and trihalomethyl.

[00131] The term “heteroaralkyl” refers to the groups -alkylene-heteroaryl where alkylene and heteroaryl are defined herein. This term includes, by way of example, pyridylmethyl, pyridylethyl, indolylmethyl, and the like.

[00132] “Heteroaryloxy” refers to –O-heteroaryl.

[00133] “Heterocycle,” “heterocyclic,” “heterocycloalkyl,” and “heterocyclyl” refer to a saturated or unsaturated group having a single ring or multiple condensed rings, including fused bridged and spiro ring systems, and having from 3 to 20 ring atoms, including 1 to 10 hetero atoms. These ring atoms are selected from nitrogen, sulfur, or oxygen, where, in fused ring systems, one or more of the rings can be cycloalkyl, aryl, or heteroaryl, provided that the point of attachment is through the non-aromatic ring. In certain embodiments, the nitrogen and/or sulfur

atom(s) of the heterocyclic group are optionally oxidized to provide for the N-oxide, -S(O)-, or -SO₂- moieties. To satisfy valence requirements, any heteroatoms in such heterocyclic rings may or may not be bonded to one or more H or one or more substituent group(s), e.g., an alkyl group or other substituent as described herein.

[00134] Examples of heterocycles and heteroaryls include, but are not limited to, azetidine, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, dihydroindole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline, phthalimide, 1,2,3,4-tetrahydroisoquinoline, 4,5,6,7-tetrahydrobenzo[b]thiophene, thiazole, thiazolidine, thiophene, benzo[b]thiophene, morpholinyl, thiomorpholinyl (also referred to as thiamorpholinyl), 1,1-dioxothiomorpholinyl, piperidinyl, pyrrolidine, tetrahydrofuranyl, and the like.

[00135] Unless otherwise constrained by the definition for the heterocyclic substituent, such heterocyclic groups can be optionally substituted with 1 to 5, or from 1 to 3 substituents, selected from alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, oxo, thioketo, carboxyl, carboxylalkyl, thioaryloxy, thioheteroaryloxy, thioheterocycloxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclyl, heterocycloxy, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl, and fused heterocycle.

[00136] “Heterocyclyloxy” refers to the group -O-heterocyclyl.

[00137] The term “heterocyclylthio” refers to the group heterocyclic-S-.

[00138] The term “heterocyclene” refers to the diradical group formed from a heterocycle, as defined herein.

[00139] The term “hydroxyamino” refers to the group -NHOH.

[00140] “Nitro” refers to the group -NO₂.

[00141] “Oxo” refers to the atom (=O).

[00142] “Sulfonyl” refers to the group SO₂-alkyl, SO₂-substituted alkyl, SO₂-alkenyl, SO₂-substituted alkenyl, SO₂-cycloalkyl, SO₂-substituted cycloalkyl, SO₂-cycloalkenyl, SO₂-

substituted cycloalkenyl, SO₂-aryl, SO₂-substituted aryl, SO₂-heteroaryl, SO₂-substituted heteroaryl, SO₂-heterocyclic, and SO₂-substituted heterocyclic, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein. Sulfonyl includes, by way of example, methyl-SO₂-, phenyl-SO₂-, and 4-methylphenyl-SO₂-.

[00143] “Sulfonyloxy” refers to the group –OSO₂-alkyl, OSO₂-substituted alkyl, OSO₂-alkenyl, OSO₂-substituted alkenyl, OSO₂-cycloalkyl, OSO₂-substituted cycloalkyl, OSO₂-cycloalkenyl, OSO₂-substituted cycloalkenyl, OSO₂-aryl, OSO₂-substituted aryl, OSO₂-heteroaryl, OSO₂-substituted heteroaryl, OSO₂-heterocyclic, and OSO₂ substituted heterocyclic, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

[00144] The term “aminocarbonyloxy” refers to the group -OC(O)NRR where each R is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

[00145] “Thiol” refers to the group -SH.

[00146] “Thioxo” or the term “thioketo” refers to the atom (=S).

[00147] “Alkylthio” or the term “thioalkoxy” refers to the group -S-alkyl, wherein alkyl is as defined herein. In certain embodiments, sulfur may be oxidized to -S(O)-. The sulfoxide may exist as one or more stereoisomers.

[00148] The term “substituted thioalkoxy” refers to the group -S-substituted alkyl.

[00149] The term “thioaryloxy” refers to the group aryl-S- wherein the aryl group is as defined herein including optionally substituted aryl groups also defined herein.

[00150] The term “thioheteroaryloxy” refers to the group heteroaryl-S- wherein the heteroaryl group is as defined herein including optionally substituted aryl groups as also defined herein.

[00151] The term “thioheterocycloxy” refers to the group heterocyclyl-S- wherein the heterocyclyl group is as defined herein including optionally substituted heterocyclyl groups as also defined herein.

[00152] In addition to the disclosure herein, the term “substituted,” when used to modify a specified group or radical, can also mean that one or more hydrogen atoms of the specified group or radical are each, independently of one another, replaced with the same or different substituent groups as defined below.

[00153] In addition to the groups disclosed with respect to the individual terms herein, substituent groups for substituting for one or more hydrogens (any two hydrogens on a single carbon can be replaced with =O, =NR⁷⁰, =N-OR⁷⁰, =N₂ or =S) on saturated carbon atoms in the specified group or radical are, unless otherwise specified, -R⁶⁰, halo, =O, -OR⁷⁰, -SR⁷⁰, -NR⁸⁰R⁸⁰, trihalomethyl, -CN, -OCN, -SCN, -NO, -NO₂, =N₂, -N₃, -SO₂R⁷⁰, -SO₂O⁻M⁺, -SO₂OR⁷⁰, -OSO₂R⁷⁰, -OSO₂O⁻M⁺, -OSO₂OR⁷⁰, -P(O)(O⁻)₂(M⁺)₂, -P(O)(OR⁷⁰)O⁻M⁺, -P(O)(OR⁷⁰)₂, -C(O)R⁷⁰, -C(S)R⁷⁰, -C(NR⁷⁰)R⁷⁰, -C(O)O⁻M⁺, -C(O)OR⁷⁰, -C(S)OR⁷⁰, -C(O)NR⁸⁰R⁸⁰, -C(NR⁷⁰)NR⁸⁰R⁸⁰, -OC(O)R⁷⁰, -OC(S)R⁷⁰, -OC(O)O⁻M⁺, -OC(O)OR⁷⁰, -OC(S)OR⁷⁰, -NR⁷⁰C(O)R⁷⁰, -NR⁷⁰C(S)R⁷⁰, -NR⁷⁰CO₂⁻M⁺, -NR⁷⁰CO₂R⁷⁰, -NR⁷⁰C(S)OR⁷⁰, -NR⁷⁰C(O)NR⁸⁰R⁸⁰, -NR⁷⁰C(NR⁷⁰)R⁷⁰ and -NR⁷⁰C(NR⁷⁰)NR⁸⁰R⁸⁰, where R⁶⁰ is selected from the group consisting of optionally substituted alkyl, cycloalkyl, heteroalkyl, heterocycloalkylalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl and heteroarylalkyl, each R⁷⁰ is independently hydrogen or R⁶⁰; each R⁸⁰ is independently R⁷⁰ or alternatively, two R⁸⁰'s, taken together with the nitrogen atom to which they are bonded, form a 5-, 6- or 7-membered heterocycloalkyl which may optionally include from 1 to 4 of the same or different additional heteroatoms selected from the group consisting of O, N and S, of which N may have -H or C₁-C₃ alkyl substitution; and each M⁺ is a counter ion with a net single positive charge. Each M⁺ may independently be, for example, an alkali ion, such as K⁺, Na⁺, Li⁺; an ammonium ion, such as ⁺N(R⁶⁰)₄; or an alkaline earth ion, such as [Ca²⁺]_{0.5}, [Mg²⁺]_{0.5}, or [Ba²⁺]_{0.5} (“subscript 0.5 means that one of the counter ions for such divalent alkali earth ions can be an ionized form of a compound of the invention and the other a typical counter ion such as chloride, or two ionized compounds disclosed herein can serve as counter ions for such divalent alkali earth ions, or a doubly ionized compound of the invention can serve as the counter ion for such divalent alkali earth ions). As specific examples, -NR⁸⁰R⁸⁰ is meant to include -NH₂, -NH-alkyl, *N*-pyrrolidinyl, *N*-piperazinyl, 4*N*-methyl-piperazin-1-yl and *N*-morpholinyl.

[00154] In addition to the disclosure herein, substituent groups for hydrogens on unsaturated carbon atoms in “substituted” alkene, alkyne, aryl and heteroaryl groups are, unless otherwise specified, $-R^{60}$, halo, $-O^+M^+$, $-OR^{70}$, $-SR^{70}$, $-S^+M^+$, $-NR^{80}R^{80}$, trihalomethyl, $-CF_3$, $-CN$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $-N_3$, $-SO_2R^{70}$, $-SO_3^-M^+$, $-SO_3R^{70}$, $-OSO_2R^{70}$, $-OSO_3^-M^+$, $-OSO_3R^{70}$, $-PO_3^{-2}(M^+)_2$, $-P(O)(OR^{70})O^-M^+$, $-P(O)(OR^{70})_2$, $-C(O)R^{70}$, $-C(S)R^{70}$, $-C(NR^{70})R^{70}$, $-CO_2^-M^+$, $-CO_2R^{70}$, $-C(S)OR^{70}$, $-C(O)NR^{80}R^{80}$, $-C(NR^{70})NR^{80}R^{80}$, $-OC(O)R^{70}$, $-OC(S)R^{70}$, $-OCO_2^-M^+$, $-OCO_2R^{70}$, $-OC(S)OR^{70}$, $-NR^{70}C(O)R^{70}$, $-NR^{70}C(S)R^{70}$, $-NR^{70}CO_2^-M^+$, $-NR^{70}CO_2R^{70}$, $-NR^{70}C(S)OR^{70}$, $-NR^{70}C(O)NR^{80}R^{80}$, $-NR^{70}C(NR^{70})R^{70}$ and $-NR^{70}C(NR^{70})NR^{80}R^{80}$, where R^{60} , R^{70} , R^{80} and M^+ are as previously defined, provided that in case of substituted alkene or alkyne, the substituents are not $-O^+M^+$, $-OR^{70}$, $-SR^{70}$, or $-S^+M^+$.

[00155] In addition to the groups disclosed with respect to the individual terms herein, substituent groups for hydrogens on nitrogen atoms in “substituted” heteroalkyl and cycloheteroalkyl groups are, unless otherwise specified, $-R^{60}$, $-O^+M^+$, $-OR^{70}$, $-SR^{70}$, $-S^+M^+$, $-NR^{80}R^{80}$, trihalomethyl, $-CF_3$, $-CN$, $-NO$, $-NO_2$, $-S(O)_2R^{70}$, $-S(O)_2O^+M^+$, $-S(O)_2OR^{70}$, $-OS(O)_2R^{70}$, $-OS(O)_2O^+M^+$, $-OS(O)_2OR^{70}$, $-P(O)(O^-)_2(M^+)_2$, $-P(O)(OR^{70})O^+M^+$, $-P(O)(OR^{70})(OR^{70})$, $-C(O)R^{70}$, $-C(S)R^{70}$, $-C(NR^{70})R^{70}$, $-C(O)OR^{70}$, $-C(S)OR^{70}$, $-C(O)NR^{80}R^{80}$, $-C(NR^{70})NR^{80}R^{80}$, $-OC(O)R^{70}$, $-OC(S)R^{70}$, $-OC(O)OR^{70}$, $-OC(S)OR^{70}$, $-NR^{70}C(O)R^{70}$, $-NR^{70}C(S)R^{70}$, $-NR^{70}C(O)OR^{70}$, $-NR^{70}C(S)OR^{70}$, $-NR^{70}C(O)NR^{80}R^{80}$, $-NR^{70}C(NR^{70})R^{70}$ and $-NR^{70}C(NR^{70})NR^{80}R^{80}$, where R^{60} , R^{70} , R^{80} and M^+ are as previously defined.

[00156] In addition to the disclosure herein, in a certain embodiment, a group that is substituted has 1, 2, 3, or 4 substituents, 1, 2, or 3 substituents, 1 or 2 substituents, or 1 substituent.

[00157] It is understood that in all substituted groups defined above, polymers arrived at by defining substituents with further substituents to themselves (e.g., substituted aryl having a substituted aryl group as a substituent which is itself substituted with a substituted aryl group, which is further substituted by a substituted aryl group, etc.) are not intended for inclusion herein. In such cases, the maximum number of such substitutions is three. For example, serial substitutions of substituted aryl groups specifically contemplated herein are limited to substituted aryl-(substituted aryl)-substituted aryl.

[00158] Unless indicated otherwise, the nomenclature of substituents that are not explicitly defined herein are arrived at by naming the terminal portion of the functionality followed by the adjacent functionality toward the point of attachment. For example, the substituent “arylalkyloxycarbonyl” refers to the group (aryl)-(alkyl)-O-C(O)-.

[00159] As to any of the groups disclosed herein which contain one or more substituents, it is understood, of course, that such groups do not contain any substitution or substitution patterns which are sterically impractical and/or synthetically non-feasible. In addition, the subject compounds include all stereochemical isomers arising from the substitution of these compounds.

[00160] The term “pharmaceutically acceptable salt” means a salt which is acceptable for administration to a patient, such as a mammal (salts with counterions having acceptable mammalian safety for a given dosage regime). Such salts can be derived from pharmaceutically acceptable inorganic or organic bases and from pharmaceutically acceptable inorganic or organic acids. “Pharmaceutically acceptable salt” refers to pharmaceutically acceptable salts of a compound, which salts are derived from a variety of organic and inorganic counter ions well known in the art and include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, formate, tartrate, besylate, mesylate, acetate, maleate, oxalate, and the like.

[00161] The term “salt thereof” means a compound formed when a proton of an acid is replaced by a cation, such as a metal cation or an organic cation and the like. Where applicable, the salt is a pharmaceutically acceptable salt, although this is not required for salts of intermediate compounds that are not intended for administration to a patient. By way of example, salts of the present compounds include those wherein the compound is protonated by an inorganic or organic acid to form a cation, with the conjugate base of the inorganic or organic acid as the anionic component of the salt.

[00162] “Solvate” refers to a complex formed by combination of solvent molecules with molecules or ions of the solute. The solvent can be an organic compound, an inorganic compound, or a mixture of both. Some examples of solvents include, but are not limited to, methanol, *N,N*-dimethylformamide, tetrahydrofuran, dimethylsulfoxide, and water. When the solvent is water, the solvate formed is a hydrate.

[00163] “Stereoisomer” and “stereoisomers” refer to compounds that have same atomic connectivity but different atomic arrangement in space. Stereoisomers include cis-trans isomers, *E* and *Z* isomers, enantiomers, and diastereomers.

[00164] “Tautomer” refers to alternate forms of a molecule that differ only in electronic bonding of atoms and/or in the position of a proton, such as enol-keto and imine-enamine tautomers, or the tautomeric forms of heteroaryl groups containing a -N=C(H)-NH- ring atom arrangement, such as pyrazoles, imidazoles, benzimidazoles, triazoles, and tetrazoles. A person of ordinary skill in the art would recognize that other tautomeric ring atom arrangements are possible.

[00165] It will be appreciated that the term “or a salt or solvate or stereoisomer thereof” is intended to include all permutations of salts, solvates and stereoisomers, such as a solvate of a pharmaceutically acceptable salt of a stereoisomer of subject compound.

[00166] “Pharmaceutically effective amount” and “therapeutically effective amount” refer to an amount of a compound sufficient to treat a specified disorder or disease or one or more of its symptoms and/or to prevent the occurrence of the disease or disorder. In reference to tumorigenic proliferative disorders, a pharmaceutically or therapeutically effective amount comprises an amount sufficient to, among other things, cause the tumor to shrink or decrease the growth rate of the tumor.

[00167] “Patient” refers to human and non-human subjects, especially mammalian subjects.

[00168] The term “treating” or “treatment” as used herein means the treating or treatment of a disease or medical condition in a patient, such as a mammal (particularly a human) that includes: (a) preventing the disease or medical condition from occurring, such as, prophylactic treatment of a subject; (b) ameliorating the disease or medical condition, such as, eliminating or causing regression of the disease or medical condition in a patient; (c) suppressing the disease or medical condition, for example by, slowing or arresting the development of the disease or medical condition in a patient; or (d) alleviating a symptom of the disease or medical condition in a patient.

[00169] The terms “polypeptide,” “peptide,” and “protein” are used interchangeably herein to refer to a polymeric form of amino acids of any length. Unless specifically indicated otherwise, “polypeptide,” “peptide,” and “protein” can include genetically coded and non-coded amino acids, chemically or biochemically modified or derivatized amino acids, and polypeptides

having modified peptide backbones. The term includes fusion proteins, including, but not limited to, fusion proteins with a heterologous amino acid sequence, fusions with heterologous and homologous leader sequences, proteins which contain at least one N-terminal methionine residue (e.g., to facilitate production in a recombinant bacterial host cell); immunologically tagged proteins; and the like.

[00170] “Native amino acid sequence” or “parent amino acid sequence” are used interchangeably herein to refer to the amino acid sequence of a polypeptide prior to modification to include a modified amino acid residue.

[00171] The terms “amino acid analog,” “unnatural amino acid,” and the like may be used interchangeably, and include amino acid-like compounds that are similar in structure and/or overall shape to one or more amino acids commonly found in naturally occurring proteins (e.g., Ala or A, Cys or C, Asp or D, Glu or E, Phe or F, Gly or G, His or H, Ile or I, Lys or K, Leu or L, Met or M, Asn or N, Pro or P, Gln or Q, Arg or R, Ser or S, Thr or T, Val or V, Trp or W, Tyr or Y). Amino acid analogs also include natural amino acids with modified side chains or backbones. Amino acid analogs also include amino acid analogs with the same stereochemistry as in the naturally occurring D-form, as well as the L-form of amino acid analogs. In some instances, the amino acid analogs share backbone structures, and/or the side chain structures of one or more natural amino acids, with difference(s) being one or more modified groups in the molecule. Such modification may include, but is not limited to, substitution of an atom (such as N) for a related atom (such as S), addition of a group (such as methyl, or hydroxyl, etc.) or an atom (such as Cl or Br, etc.), deletion of a group, substitution of a covalent bond (single bond for double bond, etc.), or combinations thereof. For example, amino acid analogs may include α -hydroxy acids, and α -amino acids, and the like.

[00172] The terms “amino acid side chain” or “side chain of an amino acid” and the like may be used to refer to the substituent attached to the α -carbon of an amino acid residue, including natural amino acids, unnatural amino acids, and amino acid analogs. An amino acid side chain can also include an amino acid side chain as described in the context of the modified amino acids and/or conjugates described herein.

[00173] The term “carbohydrate” and the like may be used to refer to monomers units and/or polymers of monosaccharides, disaccharides, oligosaccharides, and polysaccharides. The term sugar may be used to refer to the smaller carbohydrates, such as monosaccharides,

disaccharides. The term “carbohydrate derivative” includes compounds where one or more functional groups of a carbohydrate of interest are substituted (replaced by any convenient substituent), modified (converted to another group using any convenient chemistry) or absent (e.g., eliminated or replaced by H). A variety of carbohydrates and carbohydrate derivatives are available and may be adapted for use in the subject compounds and conjugates.

[00174] The term “antibody” is used in the broadest sense and includes monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, and multispecific antibodies (e.g., bispecific antibodies), humanized antibodies, single-chain antibodies, chimeric antibodies, antibody fragments (e.g., Fab fragments), and the like. An antibody is capable of binding a target antigen. (Janeway, C., Travers, P., Walport, M., Shlomchik (2001) *Immuno Biology*, 5th Ed., Garland Publishing, New York). A target antigen can have one or more binding sites, also called epitopes, recognized by complementarity determining regions (CDRs) formed by one or more variable regions of an antibody.

[00175] The term “natural antibody” refers to an antibody in which the heavy and light chains of the antibody have been made and paired by the immune system of a multi-cellular organism. Spleen, lymph nodes, bone marrow and serum are examples of tissues that produce natural antibodies. For example, the antibodies produced by the antibody producing cells isolated from a first animal immunized with an antigen are natural antibodies.

[00176] The term “humanized antibody” or “humanized immunoglobulin” refers to a non-human (e.g., mouse or rabbit) antibody containing one or more amino acids (in a framework region, a constant region or a CDR, for example) that have been substituted with a correspondingly positioned amino acid from a human antibody. In general, humanized antibodies produce a reduced immune response in a human host, as compared to a non-humanized version of the same antibody. Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Pat. Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, *Molecular Immunology* 28(4/5):489-498 (1991); Studnicka et al., *Protein Engineering* 7(6):805-814 (1994); Roguska. et al., *PNAS* 91:969-973 (1994)), and chain shuffling (U.S. Pat. No. 5,565,332). In certain embodiments, framework substitutions are identified by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular

positions (see, e.g., U.S. Pat. No. 5,585,089; Riechmann et al., Nature 332:323 (1988)).

Additional methods for humanizing antibodies contemplated for use in the present invention are described in U.S. Pat. Nos. 5,750,078; 5,502,167; 5,705,154; 5,770,403; 5,698,417; 5,693,493; 5,558,864; 4,935,496; and 4,816,567, and PCT publications WO 98/45331 and WO 98/45332. In particular embodiments, a subject rabbit antibody may be humanized according to the methods set forth in US20040086979 and US20050033031. Accordingly, the antibodies described above may be humanized using methods that are well known in the art.

[00177] The term “chimeric antibodies” refer to antibodies whose light and heavy chain genes have been constructed, typically by genetic engineering, from antibody variable and constant region genes belonging to different species. For example, the variable segments of the genes from a mouse monoclonal antibody may be joined to human constant segments, such as gamma 1 and gamma 3. An example of a therapeutic chimeric antibody is a hybrid protein composed of the variable or antigen-binding domain from a mouse antibody and the constant or effector domain from a human antibody, although domains from other mammalian species may be used.

[00178] An immunoglobulin polypeptide immunoglobulin light or heavy chain variable region is composed of a framework region (FR) interrupted by three hypervariable regions, also called “complementarity determining regions” or “CDRs”. The extent of the framework region and CDRs have been defined (see, “Sequences of Proteins of Immunological Interest,” E. Kabat et al., U.S. Department of Health and Human Services, 1991). The framework region of an antibody, that is the combined framework regions of the constituent light and heavy chains, serves to position and align the CDRs. The CDRs are primarily responsible for binding to an epitope of an antigen.

[00179] Throughout the present disclosure, the numbering of the residues in an immunoglobulin heavy chain and in an immunoglobulin light chain is that as in Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991), expressly incorporated herein by reference.

[00180] A “parent Ig polypeptide” is a polypeptide comprising an amino acid sequence which lacks an aldehyde-tagged constant region as described herein. The parent polypeptide may comprise a native sequence constant region, or may comprise a constant region with pre-existing amino acid sequence modifications (such as additions, deletions and/or substitutions).

[00181] In the context of an Ig polypeptide, the term “constant region” is well understood in the art, and refers to a C-terminal region of an Ig heavy chain, or an Ig light chain. An Ig heavy chain constant region includes CH1, CH2, and CH3 domains (and CH4 domains, where the heavy chain is a μ or an ϵ heavy chain). In a native Ig heavy chain, the CH1, CH2, CH3 (and, if present, CH4) domains begin immediately after (C-terminal to) the heavy chain variable (VH) region, and are each from about 100 amino acids to about 130 amino acids in length. In a native Ig light chain, the constant region begins immediately after (C-terminal to) the light chain variable (VL) region, and is about 100 amino acids to 120 amino acids in length.

[00182] As used herein, the term “CDR” or “complementarity determining region” is intended to mean the non-contiguous antigen combining sites found within the variable region of both heavy and light chain polypeptides. CDRs have been described by Kabat et al., J. Biol. Chem. 252:6609-6616 (1977); Kabat et al., U.S. Dept. of Health and Human Services, “Sequences of proteins of immunological interest” (1991); by Chothia et al., J. Mol. Biol. 196:901-917 (1987); and MacCallum et al., J. Mol. Biol. 262:732-745 (1996), where the definitions include overlapping or subsets of amino acid residues when compared against each other. Nevertheless, application of either definition to refer to a CDR of an antibody or grafted antibodies or variants thereof is intended to be within the scope of the term as defined and used herein. The amino acid residues which encompass the CDRs as defined by each of the above cited references are set forth below in Table 1 as a comparison.

Table 1: CDR Definitions

	Kabat¹	Chothia²	MacCallum³
V _H CDR1	31-35	26-32	30-35
V _H CDR2	50-65	53-55	47-58
V _H CDR3	95-102	96-101	93-101
V _L CDR1	24-34	26-32	30-36
V _L CDR2	50-56	50-52	46-55
V _L CDR3	89-97	91-96	89-96

¹ Residue numbering follows the nomenclature of Kabat et al., *supra*

² Residue numbering follows the nomenclature of Chothia et al., *supra*

³ Residue numbering follows the nomenclature of MacCallum et al., *supra*

[00183] By “genetically-encodable” as used in reference to an amino acid sequence of polypeptide, peptide or protein means that the amino acid sequence is composed of amino acid residues that are capable of production by transcription and translation of a nucleic acid encoding

the amino acid sequence, where transcription and/or translation may occur in a cell or in a cell-free in vitro transcription/translation system.

[00184] The term “control sequences” refers to DNA sequences that facilitate expression of an operably linked coding sequence in a particular expression system, e.g. mammalian cell, bacterial cell, cell-free synthesis, etc. The control sequences that are suitable for prokaryote systems, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cell systems may utilize promoters, polyadenylation signals, and enhancers.

[00185] A nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate the initiation of translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. Linking is accomplished by ligation or through amplification reactions. Synthetic oligonucleotide adaptors or linkers may be used for linking sequences in accordance with conventional practice.

[00186] The term “expression cassette” as used herein refers to a segment of nucleic acid, usually DNA, that can be inserted into a nucleic acid (e.g., by use of restriction sites compatible with ligation into a construct of interest or by homologous recombination into a construct of interest or into a host cell genome). In general, the nucleic acid segment comprises a polynucleotide that encodes a polypeptide of interest, and the cassette and restriction sites are designed to facilitate insertion of the cassette in the proper reading frame for transcription and translation. Expression cassettes can also comprise elements that facilitate expression of a polynucleotide encoding a polypeptide of interest in a host cell. These elements may include, but are not limited to: a promoter, a minimal promoter, an enhancer, a response element, a terminator sequence, a polyadenylation sequence, and the like.

[00187] As used herein the term “isolated” is meant to describe a compound of interest that is in an environment different from that in which the compound naturally occurs. “Isolated” is meant to include compounds that are within samples that are substantially enriched for the

compound of interest and/or in which the compound of interest is partially or substantially purified.

[00188] As used herein, the term “substantially purified” refers to a compound that is removed from its natural environment and is at least 60% free, at least 75% free, at least 80% free, at least 85% free, at least 90% free, at least 95% free, at least 98% free, or more than 98% free, from other components with which it is naturally associated.

[00189] The term “physiological conditions” is meant to encompass those conditions compatible with living cells, *e.g.*, predominantly aqueous conditions of a temperature, pH, salinity, *etc.* that are compatible with living cells.

[00190] By “reactive partner” is meant a molecule or molecular moiety that specifically reacts with another reactive partner to produce a reaction product. Exemplary reactive partners include a cysteine or serine of a sulfatase motif and Formylglycine Generating Enzyme (FGE), which react to form a reaction product of a converted aldehyde tag containing a formylglycine (FGly) in lieu of cysteine or serine in the motif. Other exemplary reactive partners include an aldehyde of an fGly residue of a converted aldehyde tag (*e.g.*, a reactive aldehyde group) and an “aldehyde-reactive reactive partner”, which comprises an aldehyde-reactive group and a moiety of interest, and which reacts to form a reaction product of a modified aldehyde tagged polypeptide having the moiety of interest conjugated to the modified polypeptide through a modified fGly residue.

[00191] “N-terminus” refers to the terminal amino acid residue of a polypeptide having a free amine group, which amine group in non-N-terminus amino acid residues normally forms part of the covalent backbone of the polypeptide.

[00192] “C-terminus” refers to the terminal amino acid residue of a polypeptide having a free carboxyl group, which carboxyl group in non-C-terminus amino acid residues normally forms part of the covalent backbone of the polypeptide.

[00193] By “internal site” as used in referenced to a polypeptide or an amino acid sequence of a polypeptide means a region of the polypeptide that is not at the N-terminus or at the C-terminus.

[00194] As used herein, an “anti-cancer agent” refers to, for example, a chemotherapeutic agent and/or biologic therapeutic agent, *e.g.*, a small molecule compound, an antibody, an

antibody-drug conjugate, and the like, or compositions thereof that are effective in treating or preventing cancer in a subject.

[00195] As used herein, a “resistant cancer” refers to a cancer that is not responsive to, is no longer responsive to, or is less responsive to, a particular mode(s) of treatment. As used herein, a “resistant cancer” is used synonymously with “refractory cancer.” In some embodiments, a resistant cancer can be a relapsed cancer, e.g., a cancer in which a treatment initially provided positive results, but wherein the cancer has become resistant to the treatment.

[00196] As used herein, “sensitizing a cancer” or “sensitized cancer” refers to a resistant cancer that has become no longer resistant to, or has become less resistant to further treatment. In other words, a resistant cancer becomes responsive or regains responsiveness to treatment when it is sensitized.

[00197] As used herein, the term “anti-CD22 antibody conjugate” is used synonymously with “antibody-drug conjugate”, “ADC,” “conjugate,” and “polypeptide-drug conjugate.”

[00198] Before the present invention is further described, it is to be understood that this invention is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[00199] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[00200] It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable sub-combination. All combinations of the embodiments pertaining to the invention are

specifically embraced by the present invention and are disclosed herein just as if each and every combination was individually and explicitly disclosed, to the extent that such combinations embrace subject matter that are, for example, compounds that are stable compounds (i.e., compounds that can be made, isolated, characterized, and tested for biological activity). In addition, all sub-combinations of the various embodiments and elements thereof (e.g., elements of the chemical groups listed in the embodiments describing such variables) are also specifically embraced by the present invention and are disclosed herein just as if each and every such sub-combination was individually and explicitly disclosed herein.

[00201] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

[00202] It must be noted that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. It is further noted that the claims may be drafted to exclude any optional element. As such, this statement is intended to serve as antecedent basis for use of such exclusive terminology as “solely,” “only” and the like in connection with the recitation of claim elements, or use of a “negative” limitation.

[00203] It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable sub-combination.

[00204] The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

ANTIBODY-DRUG CONJUGATES

[00205] The present disclosure provides conjugates, e.g., antibody-drug conjugates. By “conjugate” is meant a first moiety (e.g., an antibody) is stably associated with a second moiety (e.g., a drug). For example, a maytansine conjugate includes a maytansine (e.g., a maytansine active agent moiety) stably associated with another moiety (e.g., the antibody). By “stably associated” is meant that a moiety is bound to another moiety or structure under standard conditions. In certain embodiments, the first and second moieties are bound to each other through one or more covalent bonds.

[00206] In certain embodiments, the conjugate is a polypeptide conjugate, which includes a polypeptide conjugated to a second moiety. In certain embodiments, the moiety conjugated to the polypeptide can be any of a variety of moieties of interest such as, but not limited to, a detectable label, a drug, a water-soluble polymer, or a moiety for immobilization of the polypeptide to a membrane or a surface. In certain embodiments, the conjugate is a maytansine conjugate, where a polypeptide is conjugated to a maytansine or a maytansine active agent moiety. “Maytansine”, “maytansine moiety”, “maytansine active agent moiety” and “maytansinoid” refer to a maytansine and analogs and derivatives thereof, and pharmaceutically active maytansine moieties and/or portions thereof. A maytansine conjugated to the polypeptide can be any of a variety of maytansinoid moieties such as, but not limited to, maytansine and analogs and derivatives thereof as described herein.

[00207] The moiety of interest can be conjugated to the polypeptide at any desired site of the polypeptide. Thus, the present disclosure provides, for example, a modified polypeptide having a moiety conjugated at a site at or near the C-terminus of the polypeptide. Other examples include a modified polypeptide having a moiety conjugated at a position at or near the N-terminus of the polypeptide. Examples also include a modified polypeptide having a moiety conjugated at a position between the C-terminus and the N-terminus of the polypeptide (e.g., at an internal site of the polypeptide). Combinations of the above are also possible where the modified polypeptide is conjugated to two or more moieties.

[00208] In certain embodiments, a conjugate of the present disclosure includes a maytansine conjugated to an amino acid residue of a polypeptide at the α -carbon of an amino acid residue. Stated another way, a maytansine conjugate includes a polypeptide where the side chain

of one or more amino acid residues in the polypeptide have been modified to be attached to a maytansine (e.g., attached to a maytansine through a linker as described herein). For example, a maytansine conjugate includes a polypeptide where the α -carbon of one or more amino acid residues in the polypeptide has been modified to be attached to a maytansine (e.g., attached to a maytansine through a linker as described herein).


[00209] Embodiments of the present disclosure include conjugates where a polypeptide is conjugated to one or more moieties, such as 2 moieties, 3 moieties, 4 moieties, 5 moieties, 6 moieties, 7 moieties, 8 moieties, 9 moieties, or 10 or more moieties. The moieties may be conjugated to the polypeptide at one or more sites in the polypeptide. For example, one or more moieties may be conjugated to a single amino acid residue of the polypeptide. In some cases, one moiety is conjugated to an amino acid residue of the polypeptide. In other embodiments, two moieties may be conjugated to the same amino acid residue of the polypeptide. In other embodiments, a first moiety is conjugated to a first amino acid residue of the polypeptide and a second moiety is conjugated to a second amino acid residue of the polypeptide. Combinations of the above are also possible, for example where a polypeptide is conjugated to a first moiety at a first amino acid residue and conjugated to two other moieties at a second amino acid residue. Other combinations are also possible, such as, but not limited to, a polypeptide conjugated to first and second moieties at a first amino acid residue and conjugated to third and fourth moieties at a second amino acid residue, etc.

[00210] The one or more amino acid residues of the polypeptide that are conjugated to the one or more moieties may be naturally occurring amino acids, unnatural amino acids, or combinations thereof. For instance, the conjugate may include a moiety conjugated to a naturally occurring amino acid residue of the polypeptide. In other instances, the conjugate may include a moiety conjugated to an unnatural amino acid residue of the polypeptide. One or more moieties may be conjugated to the polypeptide at a single natural or unnatural amino acid residue as described above. One or more natural or unnatural amino acid residues in the polypeptide may be conjugated to the moiety or moieties as described herein. For example, two (or more) amino acid residues (e.g., natural or unnatural amino acid residues) in the polypeptide may each be conjugated to one or two moieties, such that multiple sites in the polypeptide are modified.

[00211] As described herein, a polypeptide may be conjugated to one or more moieties. In certain embodiments, the moiety of interest is a chemical entity, such as a drug or a detectable

label. For example, a drug (e.g., maytansine) may be conjugated to the polypeptide, or in other embodiments, a detectable label may be conjugated to the polypeptide. Thus, for instance, embodiments of the present disclosure include, but are not limited to, the following: a conjugate of a polypeptide and a drug; a conjugate of a polypeptide and a detectable label; a conjugate of two or more drugs and a polypeptide; a conjugate of two or more detectable labels and a polypeptide; and the like.

[00212] In certain embodiments, the polypeptide and the moiety of interest are conjugated through a coupling moiety. For example, the polypeptide and the moiety of interest may each be bound (e.g., covalently bonded) to the coupling moiety, thus indirectly binding the polypeptide and the moiety of interest (e.g., a drug, such as maytansine) together through the coupling moiety. In some cases, the coupling moiety includes a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl compound, or a derivative of a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl compound. For instance, a general scheme for coupling a moiety of interest (e.g., a maytansine) to a polypeptide through a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety is shown in the general reaction scheme below. Hydrazinyl-indolyl and hydrazinyl-pyrrolo-pyridinyl coupling moiety are also referred to herein as a hydrazino-*iso*-Pictet-Spengler (HIPS) coupling moiety and an aza-hydrazino-*iso*-Pictet-Spengler (azaHIPS) coupling moiety, respectively.



[00213] In the reaction scheme above, R is the moiety of interest (e.g., maytansine) that is conjugated to the polypeptide. As shown in the reaction scheme above, a polypeptide that includes a 2-formylglycine residue (fGly) is reacted with a drug (e.g., maytansine) that has been modified to include a coupling moiety (e.g., a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety) to produce a polypeptide conjugate attached to the coupling moiety, thus attaching the maytansine to the polypeptide through the coupling moiety.

[00214] As described herein, the moiety can be any of a variety of moieties such as, but not limited to, chemical entity, such as a detectable label, or a drug (e.g., a maytansinoid). R' and R'' may each independently be any desired substituent, such as, but not limited to, hydrogen,

alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. Z may be CR¹¹, NR¹², N, O or S, where R¹¹ and R¹² are each independently selected from any of the substituents described for R' and R" above.

[00215] Other hydrazinyl-indolyl or hydrazinyl-pyrrolo-pyridinyl coupling moieties are also possible, as shown in the conjugates and compounds described herein. For example, the hydrazinyl-indolyl or hydrazinyl-pyrrolo-pyridinyl coupling moieties may be modified to be attached (e.g., covalently attached) to a linker. As such, embodiments of the present disclosure include a hydrazinyl-indolyl or hydrazinyl-pyrrolo-pyridinyl coupling moiety attached to a drug (e.g., maytansine) through a linker. Various embodiments of the linker that may couple the hydrazinyl-indolyl or hydrazinyl-pyrrolo-pyridinyl coupling moiety to the drug (e.g., maytansine) are described in detail herein.

[00216] In certain embodiments, the polypeptide may be conjugated to a moiety of interest, where the polypeptide is modified before conjugation to the moiety of interest. Modification of the polypeptide may produce a modified polypeptide that contains one or more reactive groups suitable for conjugation to the moiety of interest. In some cases, the polypeptide may be modified at one or more amino acid residues to provide one or more reactive groups suitable for conjugation to the moiety of interest (e.g., a moiety that includes a coupling moiety, such as a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety as described above). For example, the polypeptide may be modified to include a reactive aldehyde group (e.g., a reactive aldehyde). A reactive aldehyde may be included in an "aldehyde tag" or "ald-tag", which as used herein refers to an amino acid sequence derived from a sulfatase motif (e.g., L(C/S)TPSR) that has been converted by action of a formylglycine generating enzyme (FGE) to contain a 2-formylglycine residue (referred to herein as "FGly"). The FGly residue generated by an FGE may also be referred to as a "formylglycine". Stated differently, the term "aldehyde tag" is used herein to refer to an amino acid sequence that includes a "converted" sulfatase motif (i.e., a sulfatase motif in which a cysteine or serine residue has been converted to FGly by action of an FGE, e.g., L(FGly)TPSR). A converted sulfatase motif may be derived from an amino acid


sequence that includes an “unconverted” sulfatase motif (i.e., a sulfatase motif in which the cysteine or serine residue has not been converted to FGly by an FGE, but is capable of being converted, e.g., an unconverted sulfatase motif with the sequence: L(C/S)TPSR). By “conversion” as used in the context of action of a formylglycine generating enzyme (FGE) on a sulfatase motif refers to biochemical modification of a cysteine or serine residue in a sulfatase motif to a formylglycine (FGly) residue (e.g., Cys to FGly, or Ser to FGly). Additional aspects of aldehyde tags and uses thereof in site-specific protein modification are described in U.S. Patent No. 7,985,783 and U.S. Patent No. 8,729,232, the disclosures of each of which are incorporated herein by reference.

[00217] In some cases, the modified polypeptide containing the FGly residue may be conjugated to the moiety of interest by reaction of the FGly with a compound (e.g., a compound containing a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety, as described above). For example, an FGly-containing polypeptide may be contacted with a reactive partner-containing drug under conditions suitable to provide for conjugation of the drug to the polypeptide. In some instances, the reactive partner-containing drug may include a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety as described above. For example, a maytansine may be modified to include a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety. In some cases, the maytansine is attached to a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl, such as covalently attached to a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl through a linker, as described in detail herein.

[00218] In certain embodiments, a conjugate of the present disclosure includes a polypeptide (e.g., an antibody, such as an anti-CD22 antibody) having at least one modified amino acid residue. The modified amino acid residue of the polypeptide may be coupled to a drug (e.g., maytansine) containing a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety as described above. In certain embodiments, the modified amino acid residue of the polypeptide (e.g., anti-CD22 antibody) may be derived from a cysteine or serine residue that has been converted to an FGly residue as described above. In certain embodiments, the FGly residue is conjugated to a drug containing a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety as described above to provide a conjugate of the present disclosure where the drug is conjugated to the polypeptide through the hydrazinyl-indolyl or hydrazinyl-pyrrolo-pyridinyl coupling moiety. As used herein, the term FGly’ refers to the modified amino acid

residue of the polypeptide (e.g., anti-CD22 antibody) that is coupled to the moiety of interest (e.g., a drug, such as a maytansinoid).

[00219] In certain embodiments, the conjugate includes at least one modified amino acid residue of the formula (I) described herein. For instance, the conjugate may include at least one modified amino acid residue with a side chain of the formula (I):



wherein

Z is CR⁴ or N;

R¹ is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R² and R³ are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R² and R³ are optionally cyclically linked to form a 5 or 6-membered heterocyclyl;

each R⁴ is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

L is a linker comprising -(T¹-V¹)_a-(T²-V²)_b-(T³-V³)_c-(T⁴-V⁴)_d, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4;

T^1 , T^2 , T^3 and T^4 are each independently selected from (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(EDA)_w$, $(PEG)_n$, $(AA)_p$, $-(CR^{13}OH)_h$, piperidin-4-amino (4AP), an acetal group, a hydrazine, a disulfide, and an ester, wherein EDA is an ethylene diamine moiety, PEG is a polyethylene glycol or a modified polyethylene glycol, and AA is an amino acid residue, wherein w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12;

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$ and $-P(O)OH-$, wherein q is an integer from 1 to 6;

each R^{13} is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl;

each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

W^1 is a maytansinoid; and

W^2 is an anti-CD22 antibody.

[00220] In certain embodiments, Z is CR^4 or N . In certain embodiments, Z is CR^4 . In certain embodiments, Z is N .

[00221] In certain embodiments, R^1 is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In certain embodiments, R^1 is hydrogen. In certain embodiments, R^1 is alkyl or substituted alkyl, such as C_{1-6} alkyl or C_{1-6} substituted alkyl, or C_{1-4} alkyl or C_{1-4} substituted alkyl, or C_{1-3} alkyl or C_{1-3} substituted alkyl. In certain embodiments, R^1 is alkenyl or substituted alkenyl, such as C_{2-6} alkenyl or C_{2-6} substituted alkenyl, or C_{2-4} alkenyl or C_{2-4} substituted alkenyl, or C_{2-3} alkenyl or C_{2-3} substituted alkenyl. In certain embodiments, R^1 is alkynyl or substituted alkynyl, such as C_{2-6} alkynyl or C_{2-6} substituted alkynyl, or C_{2-4} alkynyl or C_{2-4} substituted alkynyl, or C_{2-3} alkynyl or C_{2-3} substituted alkynyl. In certain embodiments, R^1 is aryl or substituted aryl, such as C_{5-8} aryl or C_{5-8} substituted aryl, such as a C_5 aryl or C_5 substituted

aryl, or a C₆ aryl or C₆ substituted aryl. In certain embodiments, R¹ is heteroaryl or substituted heteroaryl, such as C₅₋₈ heteroaryl or C₅₋₈ substituted heteroaryl, such as a C₅ heteroaryl or C₅ substituted heteroaryl, or a C₆ heteroaryl or C₆ substituted heteroaryl. In certain embodiments, R¹ is cycloalkyl or substituted cycloalkyl, such as C₃₋₈ cycloalkyl or C₃₋₈ substituted cycloalkyl, such as a C₃₋₆ cycloalkyl or C₃₋₆ substituted cycloalkyl, or a C₃₋₅ cycloalkyl or C₃₋₅ substituted cycloalkyl. In certain embodiments, R¹ is heterocyclyl or substituted heterocyclyl, such as C₃₋₈ heterocyclyl or C₃₋₈ substituted heterocyclyl, such as a C₃₋₆ heterocyclyl or C₃₋₆ substituted heterocyclyl, or a C₃₋₅ heterocyclyl or C₃₋₅ substituted heterocyclyl.

[00222] In certain embodiments, R² and R³ are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R² and R³ are optionally cyclically linked to form a 5 or 6-membered heterocyclyl.

[00223] In certain embodiments, R² is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In certain embodiments, R² is hydrogen. In certain embodiments, R² is alkyl or substituted alkyl, such as C₁₋₆ alkyl or C₁₋₆ substituted alkyl, or C₁₋₄ alkyl or C₁₋₄ substituted alkyl, or C₁₋₃ alkyl or C₁₋₃ substituted alkyl. In certain embodiments, R² is alkenyl or substituted alkenyl, such as C₂₋₆ alkenyl or C₂₋₆ substituted alkenyl, or C₂₋₄ alkenyl or C₂₋₄ substituted alkenyl, or C₂₋₃ alkenyl or C₂₋₃ substituted alkenyl. In certain embodiments, R² is alkynyl or substituted alkynyl. In certain embodiments, R² is alkoxy or substituted alkoxy. In certain embodiments, R² is amino or substituted amino. In certain embodiments, R² is carboxyl or carboxyl ester. In certain embodiments, R² is acyl or acyloxy. In certain embodiments, R² is acyl amino or amino acyl. In certain embodiments, R² is alkylamide or substituted alkylamide. In certain embodiments, R² is sulfonyl. In certain embodiments, R² is thioalkoxy or substituted thioalkoxy. In certain embodiments, R² is aryl or substituted aryl, such as C₅₋₈ aryl or C₅₋₈

substituted aryl, such as a C₅ aryl or C₅ substituted aryl, or a C₆ aryl or C₆ substituted aryl. In certain embodiments, R² is heteroaryl or substituted heteroaryl, such as C₅₋₈ heteroaryl or C₅₋₈ substituted heteroaryl, such as a C₅ heteroaryl or C₅ substituted heteroaryl, or a C₆ heteroaryl or C₆ substituted heteroaryl. In certain embodiments, R² is cycloalkyl or substituted cycloalkyl, such as C₃₋₈ cycloalkyl or C₃₋₈ substituted cycloalkyl, such as a C₃₋₆ cycloalkyl or C₃₋₆ substituted cycloalkyl, or a C₃₋₅ cycloalkyl or C₃₋₅ substituted cycloalkyl. In certain embodiments, R² is heterocyclyl or substituted heterocyclyl, such as a C₃₋₆ heterocyclyl or C₃₋₆ substituted heterocyclyl, or a C₃₋₅ heterocyclyl or C₃₋₅ substituted heterocyclyl.

[00224] In certain embodiments, R³ is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In certain embodiments, R³ is hydrogen. In certain embodiments, R³ is alkyl or substituted alkyl, such as C₁₋₆ alkyl or C₁₋₆ substituted alkyl, or C₁₋₄ alkyl or C₁₋₄ substituted alkyl, or C₁₋₃ alkyl or C₁₋₃ substituted alkyl. In certain embodiments, R³ is alkenyl or substituted alkenyl, such as C₂₋₆ alkenyl or C₂₋₆ substituted alkenyl, or C₂₋₄ alkenyl or C₂₋₄ substituted alkenyl, or C₂₋₃ alkenyl or C₂₋₃ substituted alkenyl. In certain embodiments, R³ is alkynyl or substituted alkynyl. In certain embodiments, R³ is alkoxy or substituted alkoxy. In certain embodiments, R³ is amino or substituted amino. In certain embodiments, R³ is carboxyl or carboxyl ester. In certain embodiments, R³ is acyl or acyloxy. In certain embodiments, R³ is acyl amino or amino acyl. In certain embodiments, R³ is alkylamide or substituted alkylamide. In certain embodiments, R³ is sulfonyl. In certain embodiments, R³ is thioalkoxy or substituted thioalkoxy. In certain embodiments, R³ is aryl or substituted aryl, such as C₅₋₈ aryl or C₅₋₈ substituted aryl, such as a C₅ aryl or C₅ substituted aryl, or a C₆ aryl or C₆ substituted aryl. In certain embodiments, R³ is heteroaryl or substituted heteroaryl, such as C₅₋₈ heteroaryl or C₅₋₈ substituted heteroaryl, such as a C₅ heteroaryl or C₅ substituted heteroaryl, or a C₆ heteroaryl or C₆ substituted heteroaryl. In certain embodiments, R³ is cycloalkyl or substituted cycloalkyl, such as C₃₋₈ cycloalkyl or C₃₋₈ substituted cycloalkyl, such as a C₃₋₆ cycloalkyl or C₃₋₆ substituted cycloalkyl, or a C₃₋₅ cycloalkyl or C₃₋₅ substituted cycloalkyl. In certain embodiments, R³ is heterocyclyl or substituted heterocyclyl, such as C₃₋₈ heterocyclyl or C₃₋₈ substituted

heterocyclyl, such as a C₃₋₆ heterocyclyl or C₃₋₆ substituted heterocyclyl, or a C₃₋₅ heterocyclyl or C₃₋₅ substituted heterocyclyl.

[00225] In certain embodiments, R² and R³ are optionally cyclically linked to form a 5 or 6-membered heterocyclyl. In certain embodiments, R² and R³ are cyclically linked to form a 5 or 6-membered heterocyclyl. In certain embodiments, R² and R³ are cyclically linked to form a 5-membered heterocyclyl. In certain embodiments, R² and R³ are cyclically linked to form a 6-membered heterocyclyl.

[00226] In certain embodiments, each R⁴ is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

[00227] The various possibilities for each R⁴ are described in more detail as follows. In certain embodiments, R⁴ is hydrogen. In certain embodiments, each R⁴ is hydrogen. In certain embodiments, R⁴ is halogen, such as F, Cl, Br or I. In certain embodiments, R⁴ is F. In certain embodiments, R⁴ is Cl. In certain embodiments, R⁴ is Br. In certain embodiments, R⁴ is I. In certain embodiments, R⁴ is alkyl or substituted alkyl, such as C₁₋₆ alkyl or C₁₋₆ substituted alkyl, or C₁₋₄ alkyl or C₁₋₄ substituted alkyl, or C₁₋₃ alkyl or C₁₋₃ substituted alkyl. In certain embodiments, R⁴ is alkenyl or substituted alkenyl, such as C₂₋₆ alkenyl or C₂₋₆ substituted alkenyl, or C₂₋₄ alkenyl or C₂₋₄ substituted alkenyl, or C₂₋₃ alkenyl or C₂₋₃ substituted alkenyl. In certain embodiments, R⁴ is alkynyl or substituted alkynyl. In certain embodiments, R⁴ is alkoxy or substituted alkoxy. In certain embodiments, R⁴ is amino or substituted amino. In certain embodiments, R⁴ is carboxyl or carboxyl ester. In certain embodiments, R⁴ is acyl or acyloxy. In certain embodiments, R⁴ is acyl amino or amino acyl. In certain embodiments, R⁴ is alkylamide or substituted alkylamide. In certain embodiments, R⁴ is sulfonyl. In certain embodiments, R⁴ is thioalkoxy or substituted thioalkoxy. In certain embodiments, R⁴ is aryl or substituted aryl, such as C₅₋₈ aryl or C₅₋₈ substituted aryl, such as a C₅ aryl or C₅ substituted aryl, or a C₆ aryl or C₆ substituted aryl (e.g., phenyl or substituted phenyl). In certain embodiments, R⁴ is heteroaryl or substituted heteroaryl, such as C₅₋₈ heteroaryl or C₅₋₈ substituted heteroaryl, such as a C₅ heteroaryl or C₅ substituted heteroaryl, or a C₆ heteroaryl or C₆ substituted

heteroaryl. In certain embodiments, R⁴ is cycloalkyl or substituted cycloalkyl, such as C₃₋₈ cycloalkyl or C₃₋₈ substituted cycloalkyl, such as a C₃₋₆ cycloalkyl or C₃₋₆ substituted cycloalkyl, or a C₃₋₅ cycloalkyl or C₃₋₅ substituted cycloalkyl. In certain embodiments, R⁴ is heterocyclyl or substituted heterocyclyl, such as C₃₋₈ heterocyclyl or C₃₋₈ substituted heterocyclyl, such as a C₃₋₆ heterocyclyl or C₃₋₆ substituted heterocyclyl, or a C₃₋₅ heterocyclyl or C₃₋₅ substituted heterocyclyl.

[00228] In certain embodiments, W¹ is a maytansinoid. Further description of the maytansinoid is found in the disclosure herein.

[00229] In certain embodiments, W² is an anti-CD22 antibody. Further description of the anti-CD22 antibody is found in the disclosure herein.

[00230] In certain embodiments, the compounds of formula (I) include a linker, L. The linker may be utilized to bind a coupling moiety to one or more moieties of interest and/or one or more polypeptides. In some embodiments, the linker binds a coupling moiety to either a polypeptide or a chemical entity. The linker may be bound (e.g., covalently bonded) to the coupling moiety (e.g., as described herein) at any convenient position. For example, the linker may attach a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety to a drug (e.g., a maytansine). The hydrazinyl-indolyl or hydrazinyl-pyrrolo-pyridinyl coupling moiety may be used to conjugate the linker (and thus the drug, e.g., maytansine) to a polypeptide, such as an anti-CD22 antibody.

[00231] In certain embodiments, L attaches the coupling moiety to W¹, and thus the coupling moiety is indirectly bonded to W¹ through the linker L. As described above, W¹ is a maytansinoid, and thus L attaches the coupling moiety to a maytansinoid, e.g., the coupling moiety is indirectly bonded to the maytansinoid through the linker, L.

[00232] Any convenient linkers may be utilized in the subject conjugates and compounds. In certain embodiments, L includes a group selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl amino, alkylamide, substituted alkylamide, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In certain embodiments, L includes an alkyl or substituted alkyl group. In certain embodiments, L includes an alkenyl or substituted alkenyl group. In certain embodiments, L includes an alkynyl or substituted alkynyl group. In certain embodiments, L

includes an alkoxy or substituted alkoxy group. In certain embodiments, L includes an amino or substituted amino group. In certain embodiments, L includes a carboxyl or carboxyl ester group. In certain embodiments, L includes an acyl amino group. In certain embodiments, L includes an alkylamide or substituted alkylamide group. In certain embodiments, L includes an aryl or substituted aryl group. In certain embodiments, L includes a heteroaryl or substituted heteroaryl group. In certain embodiments, L includes a cycloalkyl or substituted cycloalkyl group. In certain embodiments, L includes a heterocyclyl or substituted heterocyclyl group.

[00233] In certain embodiments, L includes a polymer. For example, the polymer may include a polyalkylene glycol and derivatives thereof, including polyethylene glycol, methoxypolyethylene glycol, polyethylene glycol homopolymers, polypropylene glycol homopolymers, copolymers of ethylene glycol with propylene glycol (e.g., where the homopolymers and copolymers are unsubstituted or substituted at one end with an alkyl group), polyvinyl alcohol, polyvinyl ethyl ethers, polyvinylpyrrolidone, combinations thereof, and the like. In certain embodiments, the polymer is a polyalkylene glycol. In certain embodiments, the polymer is a polyethylene glycol. Other linkers are also possible, as shown in the conjugates and compounds described in more detail below.

[00234] In some embodiments, L is a linker described by the formula $-(L^1)_a-(L^2)_b-(L^3)_c-(L^4)_d-$, wherein L^1 , L^2 , L^3 and L^4 are each independently a linker unit, and a, b, c and d are each independently 0 or 1, wherein the sum of a, b, c and d is 1 to 4.

[00235] In certain embodiments, the sum of a, b, c and d is 1. In certain embodiments, the sum of a, b, c and d is 2. In certain embodiments, the sum of a, b, c and d is 3. In certain embodiments, the sum of a, b, c and d is 4. In certain embodiments, a, b, c and d are each 1. In certain embodiments, a, b and c are each 1 and d is 0. In certain embodiments, a and b are each 1 and c and d are each 0. In certain embodiments, a is 1 and b, c and d are each 0.

[00236] In certain embodiments, L^1 is attached to the hydrazinyl-indolyl or the hydrazinyl-pyrrolo-pyridinyl coupling moiety (e.g., as shown in formula (I) above). In certain embodiments, L^2 , if present, is attached to W^1 . In certain embodiments, L^3 , if present, is attached to W^1 . In certain embodiments, L^4 , if present, is attached to W^1 .

[00237] Any convenient linker units may be utilized in the subject linkers. Linker units of interest include, but are not limited to, units of polymers such as polyethylene glycols, polyethylenes and polyacrylates, amino acid residue(s), carbohydrate-based polymers or

carbohydrate residues and derivatives thereof, polynucleotides, alkyl groups, aryl groups, heterocyclic groups, combinations thereof, and substituted versions thereof. In some embodiments, each of L^1 , L^2 , L^3 and L^4 (if present) comprise one or more groups independently selected from a polyethylene glycol, a modified polyethylene glycol, an amino acid residue, an alkyl group, a substituted alkyl, an aryl group, a substituted aryl group, and a diamine (e.g., a linking group that includes an alkylene diamine).

[00238] In some embodiments, L^1 (if present) comprises a polyethylene glycol, a modified polyethylene glycol, an amino acid residue, an alkyl group, a substituted alkyl, an aryl group, a substituted aryl group, or a diamine. In some embodiments, L^1 comprises a polyethylene glycol. In some embodiments, L^1 comprises a modified polyethylene glycol. In some embodiments, L^1 comprises an amino acid residue. In some embodiments, L^1 comprises an alkyl group or a substituted alkyl. In some embodiments, L^1 comprises an aryl group or a substituted aryl group. In some embodiments, L^1 comprises a diamine (e.g., a linking group comprising an alkylene diamine).

[00239] In some embodiments, L^2 (if present) comprises a polyethylene glycol, a modified polyethylene glycol, an amino acid residue, an alkyl group, a substituted alkyl, an aryl group, a substituted aryl group, or a diamine. In some embodiments, L^2 comprises a polyethylene glycol. In some embodiments, L^2 comprises a modified polyethylene glycol. In some embodiments, L^2 comprises an amino acid residue. In some embodiments, L^2 comprises an alkyl group or a substituted alkyl. In some embodiments, L^2 comprises an aryl group or a substituted aryl group. In some embodiments, L^2 comprises a diamine (e.g., a linking group comprising an alkylene diamine).

[00240] In some embodiments, L^3 (if present) comprises a polyethylene glycol, a modified polyethylene glycol, an amino acid residue, an alkyl group, a substituted alkyl, an aryl group, a substituted aryl group, or a diamine. In some embodiments, L^3 comprises a polyethylene glycol. In some embodiments, L^3 comprises a modified polyethylene glycol. In some embodiments, L^3 comprises an amino acid residue. In some embodiments, L^3 comprises an alkyl group or a substituted alkyl. In some embodiments, L^3 comprises an aryl group or a substituted aryl group. In some embodiments, L^3 comprises a diamine (e.g., a linking group comprising an alkylene diamine).

[00241] In some embodiments, L^4 (if present) comprises a polyethylene glycol, a modified polyethylene glycol, an amino acid residue, an alkyl group, a substituted alkyl, an aryl group, a substituted aryl group, or a diamine. In some embodiments, L^4 comprises a polyethylene glycol. In some embodiments, L^4 comprises a modified polyethylene glycol. In some embodiments, L^4 comprises an amino acid residue. In some embodiments, L^4 comprises an alkyl group or a substituted alkyl. In some embodiments, L^4 comprises an aryl group or a substituted aryl group. In some embodiments, L^4 comprises a diamine (e.g., a linking group comprising an alkylene diamine).

[00242] In some embodiments, L is a linker comprising $-(L^1)_a-(L^2)_b-(L^3)_c-(L^4)_d-$, where:

$-(L^1)_a-$ is $-(T^1-V^1)_a-$;

$-(L^2)_b-$ is $-(T^2-V^2)_b-$;

$-(L^3)_c-$ is $-(T^3-V^3)_c-$; and

$-(L^4)_d-$ is $-(T^4-V^4)_d-$,

wherein T^1 , T^2 , T^3 and T^4 , if present, are tether groups;

V^1 , V^2 , V^3 and V^4 , if present, are covalent bonds or linking functional groups; and


a, b, c and d are each independently 0 or 1, wherein the sum of a, b, c and d is 1 to 4.

[00243] As described above, in certain embodiments, L^1 is attached to the hydrazinyl-indolyl or the hydrazinyl-pyrrolo-pyridinyl coupling moiety (e.g., as shown in formula (I) above). As such, in certain embodiments, T^1 is attached to the hydrazinyl-indolyl or the hydrazinyl-pyrrolo-pyridinyl coupling moiety (e.g., as shown in formula (I) above). In certain embodiments, V^1 is attached to W^1 (the maytansinoid). In certain embodiments, L^2 , if present, is attached to W^1 . As such, in certain embodiments, T^2 , if present, is attached to W^1 , or V^2 , if present, is attached to W^1 . In certain embodiments, L^3 , if present, is attached to W^1 . As such, in certain embodiments, T^3 , if present, is attached to W^1 , or V^3 , if present, is attached to W^1 . In certain embodiments, L^4 , if present, is attached to W^1 . As such, in certain embodiments, T^4 , if present, is attached to W^1 , or V^4 , if present, is attached to W^1 .

[00244] Regarding the tether groups, T^1 , T^2 , T^3 and T^4 , any convenient tether groups may be utilized in the subject linkers. In some embodiments, T^1 , T^2 , T^3 and T^4 each comprise one or more groups independently selected from a (C_1-C_{12}) alkyl, a substituted (C_1-C_{12}) alkyl, an $(EDA)_w$, $(PEG)_n$, $(AA)_p$, $-(CR^{13}OH)_h-$, piperidin-4-amino (4AP), an acetal group, a disulfide, a


hydrazine, and an ester, where w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12.

[00245] In certain embodiments, when the sum of a, b, c and d is 2 and one of T¹-V¹, T²-V², T³-V³, or T⁴-V⁴ is (PEG)_n-CO, then n is not 6. For example, in some instances, the linker may have the following structure:



where n is not 6.

[00246] In certain embodiments, when the sum of a, b, c and d is 2 and one of T¹-V¹, T²-V², T³-V³, or T⁴-V⁴ is (C₁-C₁₂)alkyl-NR¹⁵, then (C₁-C₁₂)alkyl is not a C₅-alkyl. For example, in some instances, the linker may have the following structure:




where g is not 4.

[00247] In certain embodiments, the tether group (e.g., T¹, T², T³ and/or T⁴) includes a (C₁-C₁₂)alkyl or a substituted (C₁-C₁₂)alkyl. In certain embodiments, (C₁-C₁₂)alkyl is a straight chain or branched alkyl group that includes from 1 to 12 carbon atoms, such as 1 to 10 carbon atoms, or 1 to 8 carbon atoms, or 1 to 6 carbon atoms, or 1 to 5 carbon atoms, or 1 to 4 carbon atoms, or 1 to 3 carbon atoms. In some instances, (C₁-C₁₂)alkyl may be an alkyl or substituted alkyl, such as C₁-C₁₂ alkyl, or C₁-C₁₀ alkyl, or C₁-C₆ alkyl, or C₁-C₃ alkyl. In some instances, (C₁-C₁₂)alkyl is a C₂-alkyl. For example, (C₁-C₁₂)alkyl may be an alkylene or substituted alkylene, such as C₁-C₁₂ alkylene, or C₁-C₁₀ alkylene, or C₁-C₆ alkylene, or C₁-C₃ alkylene. In some instances, (C₁-C₁₂)alkyl is a C₂-alkylene.

[00248] In certain embodiments, substituted (C₁-C₁₂)alkyl is a straight chain or branched substituted alkyl group that includes from 1 to 12 carbon atoms, such as 1 to 10 carbon atoms, or 1 to 8 carbon atoms, or 1 to 6 carbon atoms, or 1 to 5 carbon atoms, or 1 to 4 carbon atoms, or 1 to 3 carbon atoms. In some instances, substituted (C₁-C₁₂)alkyl may be a substituted alkyl, such as substituted C₁-C₁₂ alkyl, or substituted C₁-C₁₀ alkyl, or substituted C₁-C₆ alkyl, or substituted C₁-C₃ alkyl. In some instances, substituted (C₁-C₁₂)alkyl is a substituted C₂-alkyl. For example,

substituted (C₁-C₁₂)alkyl may be a substituted alkylene, such as substituted C₁-C₁₂ alkylene, or substituted C₁-C₁₀ alkylene, or substituted C₁-C₆ alkylene, or substituted C₁-C₃ alkylene. In some instances, substituted (C₁-C₁₂)alkyl is a substituted C₂-alkylene.


[00249] In certain embodiments, the tether group (e.g., T¹, T², T³ and/or T⁴) includes an ethylene diamine (EDA) moiety, e.g., an EDA containing tether. In certain embodiments, (EDA)_w includes one or more EDA moieties, such as where w is an integer from 1 to 50, such as from 1 to 40, from 1 to 30, from 1 to 20, from 1 to 12 or from 1 to 6, such as 1, 2, 3, 4, 5 or 6). The linked ethylene diamine (EDA) moieties may optionally be substituted at one or more convenient positions with any convenient substituents, e.g., with an alkyl, a substituted alkyl, an acyl, a substituted acyl, an aryl or a substituted aryl. In certain embodiments, the EDA moiety is described by the structure:



where y is an integer from 1 to 6, r is 0 or 1, and each R¹² is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In certain embodiments, y is 1, 2, 3, 4, 5 or 6. In certain embodiments, y is 1 and r is 0. In certain embodiments, y is 1 and r is 1. In certain embodiments, y is 2 and r is 0. In certain embodiments, y is 2 and r is 1. In certain embodiments, each R¹² is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl and a substituted aryl. In certain embodiments, any two adjacent R¹² groups of the EDA may be cyclically linked, e.g., to form a piperazinyl ring. In certain embodiments, y is 1 and the two adjacent R¹² groups are an alkyl group, cyclically linked to form a piperazinyl ring. In certain embodiments, y is 1 and the adjacent R¹² groups are selected from hydrogen, an alkyl (e.g., methyl) and a substituted alkyl (e.g., lower alkyl-OH, such as ethyl-OH or propyl-OH).


[00250] In certain embodiments, the tether group includes a 4-amino-piperidine (4AP) moiety (also referred to herein as piperidin-4-amino, P4A). The 4AP moiety may optionally be substituted at one or more convenient positions with any convenient substituents, e.g., with an

alkyl, a substituted alkyl, a polyethylene glycol moiety, an acyl, a substituted acyl, an aryl or a substituted aryl. In certain embodiments, the 4AP moiety is described by the structure:




where R^{12} is selected from hydrogen, alkyl, substituted alkyl, a polyethylene glycol moiety (e.g., a polyethylene glycol or a modified polyethylene glycol), alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In certain embodiments, R^{12} is a polyethylene glycol moiety. In certain embodiments, R^{12} is a carboxy modified polyethylene glycol.

[00251] In certain embodiments, R^{12} includes a polyethylene glycol moiety described by the formula: $(PEG)_k$, which may be represented by the structure:



where k is an integer from 1 to 20, such as from 1 to 18, or from 1 to 16, or from 1 to 14, or from 1 to 12, or from 1 to 10, or from 1 to 8, or from 1 to 6, or from 1 to 4, or 1 or 2, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20. In some instances, k is 2. In certain embodiments, R^{17} is selected from OH, COOH, or COOR, where R is selected from alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In certain embodiments, R^{17} is COOH.

[00252] In certain embodiments, a tether group (e.g., T^1 , T^2 , T^3 and/or T^4) includes $(PEG)_n$, where $(PEG)_n$ is a polyethylene glycol or a modified polyethylene glycol linking unit. In certain embodiments, $(PEG)_n$ is described by the structure:



where n is an integer from 1 to 50, such as from 1 to 40, from 1 to 30, from 1 to 20, from 1 to 12 or from 1 to 6, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20. In some instances, n is 2. In some instances, n is 3. In some instances, n is 6. In some instances, n is 12.

[00253] In certain embodiments, a tether group (e.g., T^1 , T^2 , T^3 and/or T^4) includes $(AA)_p$, where AA is an amino acid residue. Any convenient amino acids may be utilized. Amino acids of interest include but are not limited to, L- and D-amino acids, naturally occurring amino acids such as any of the 20 primary alpha-amino acids and beta-alanine, non-naturally occurring amino acids (e.g., amino acid analogs), such as a non-naturally occurring alpha-amino acid or a non-naturally occurring beta-amino acid, etc. In certain embodiments, p is an integer from 1 to 50, such as from 1 to 40, from 1 to 30, from 1 to 20, from 1 to 12 or from 1 to 6, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20. In certain embodiments, p is 1. In certain embodiments, p is 2.

[00254] In certain embodiments, a tether group (e.g., T^1 , T^2 , T^3 and/or T^4) includes a moiety described by the formula $-(CR^{13}OH)_h-$, where h is 0 or n is an integer from 1 to 50, such as from 1 to 40, from 1 to 30, from 1 to 20, from 1 to 12 or from 1 to 6, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12. In certain embodiments, h is 1. In certain embodiments, h is 2. In certain embodiments, R^{13} is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In certain embodiments, R^{13} is hydrogen. In certain embodiments, R^{13} is alkyl or substituted alkyl, such as C_{1-6} alkyl or C_{1-6} substituted alkyl, or C_{1-4} alkyl or C_{1-4} substituted alkyl, or C_{1-3} alkyl or C_{1-3} substituted alkyl. In certain embodiments, R^{13} is alkenyl or substituted alkenyl, such as C_{2-6} alkenyl or C_{2-6} substituted alkenyl, or C_{2-4} alkenyl or C_{2-4} substituted alkenyl, or C_{2-3} alkenyl or C_{2-3} substituted alkenyl. In certain embodiments, R^{13} is alkynyl or substituted alkynyl. In certain embodiments, R^{13} is alkoxy or substituted alkoxy. In certain embodiments, R^{13} is amino or substituted amino. In certain embodiments, R^{13} is carboxyl or carboxyl ester. In certain embodiments, R^{13} is acyl or acyloxy. In certain embodiments, R^{13} is acyl amino or amino acyl. In certain embodiments, R^{13} is alkylamide or substituted alkylamide. In certain embodiments, R^{13} is sulfonyl. In certain embodiments, R^{13} is thioalkoxy or substituted thioalkoxy. In certain embodiments, R^{13} is aryl or substituted aryl, such as C_{5-8} aryl or C_{5-8} substituted aryl, such as a C_5 aryl or C_5 substituted aryl, or a C_6 aryl or C_6 substituted aryl. In certain embodiments, R^{13} is heteroaryl or substituted heteroaryl, such as C_{5-8} heteroaryl or C_{5-8}

substituted heteroaryl, such as a C₅ heteroaryl or C₅ substituted heteroaryl, or a C₆ heteroaryl or C₆ substituted heteroaryl. In certain embodiments, R¹³ is cycloalkyl or substituted cycloalkyl, such as C₃₋₈ cycloalkyl or C₃₋₈ substituted cycloalkyl, such as a C₃₋₆ cycloalkyl or C₃₋₆ substituted cycloalkyl, or a C₃₋₅ cycloalkyl or C₃₋₅ substituted cycloalkyl. In certain embodiments, R¹³ is heterocyclyl or substituted heterocyclyl, such as C₃₋₈ heterocyclyl or C₃₋₈ substituted heterocyclyl, such as a C₃₋₆ heterocyclyl or C₃₋₆ substituted heterocyclyl, or a C₃₋₅ heterocyclyl or C₃₋₅ substituted heterocyclyl.

[00255] In certain embodiments, R¹³ is selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl. In these embodiments, alkyl, substituted alkyl, aryl, and substituted aryl are as described above for R¹³.

[00256] Regarding the linking functional groups, V¹, V², V³ and V⁴, any convenient linking functional groups may be utilized in the subject linkers. Linking functional groups of interest include, but are not limited to, amino, carbonyl, amido, oxycarbonyl, carboxy, sulfonyl, sulfoxide, sulfonylamino, aminosulfonyl, thio, oxy, phospho, phosphoramidate, thiophosphoraidate, and the like. In some embodiments, V¹, V², V³ and V⁴ are each independently selected from a covalent bond, -CO-, -NR¹⁵-, -NR¹⁵(CH₂)_q-, -NR¹⁵(C₆H₄)-, -CONR¹⁵-, -NR¹⁵CO-, -C(O)O-, -OC(O)-, -O-, -S-, -S(O)-, -SO₂-, -SO₂NR¹⁵-, -NR¹⁵SO₂- and -P(O)OH-, where q is an integer from 1 to 6. In certain embodiments, q is an integer from 1 to 6 (e.g., 1, 2, 3, 4, 5 or 6). In certain embodiments, q is 1. In certain embodiments, q is 2.

[00257] In some embodiments, each R¹⁵ is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.

[00258] The various possibilities for each R¹⁵ are described in more detail as follows. In certain embodiments, R¹⁵ is hydrogen. In certain embodiments, each R¹⁵ is hydrogen. In certain embodiments, R¹⁵ is alkyl or substituted alkyl, such as C₁₋₆ alkyl or C₁₋₆ substituted alkyl, or C₁₋₄ alkyl or C₁₋₄ substituted alkyl, or C₁₋₃ alkyl or C₁₋₃ substituted alkyl. In certain embodiments, R¹⁵ is alkenyl or substituted alkenyl, such as C₂₋₆ alkenyl or C₂₋₆ substituted alkenyl, or C₂₋₄ alkenyl or C₂₋₄ substituted alkenyl, or C₂₋₃ alkenyl or C₂₋₃ substituted alkenyl. In certain embodiments,

R^{15} is alkynyl or substituted alkynyl. In certain embodiments, R^{15} is alkoxy or substituted alkoxy. In certain embodiments, R^{15} is amino or substituted amino. In certain embodiments, R^{15} is carboxyl or carboxyl ester. In certain embodiments, R^{15} is acyl or acyloxy. In certain embodiments, R^{15} is acyl amino or amino acyl. In certain embodiments, R^{15} is alkylamide or substituted alkylamide. In certain embodiments, R^{15} is sulfonyl. In certain embodiments, R^{15} is thioalkoxy or substituted thioalkoxy. In certain embodiments, R^{15} is aryl or substituted aryl, such as C_{5-8} aryl or C_{5-8} substituted aryl, such as a C_5 aryl or C_5 substituted aryl, or a C_6 aryl or C_6 substituted aryl. In certain embodiments, R^{15} is heteroaryl or substituted heteroaryl, such as C_{5-8} heteroaryl or C_{5-8} substituted heteroaryl, such as a C_5 heteroaryl or C_5 substituted heteroaryl, or a C_6 heteroaryl or C_6 substituted heteroaryl. In certain embodiments, R^{15} is cycloalkyl or substituted cycloalkyl, such as C_{3-8} cycloalkyl or C_{3-8} substituted cycloalkyl, such as a C_{3-6} cycloalkyl or C_{3-6} substituted cycloalkyl, or a C_{3-5} cycloalkyl or C_{3-5} substituted cycloalkyl. In certain embodiments, R^{15} is heterocyclyl or substituted heterocyclyl, such as C_{3-8} heterocyclyl or C_{3-8} substituted heterocyclyl, such as a C_{3-6} heterocyclyl or C_{3-6} substituted heterocyclyl, or a C_{3-5} heterocyclyl or C_{3-5} substituted heterocyclyl.

[00259] In certain embodiments, each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In these embodiments, the hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl substituents are as described above for R^{15} .

[00260] In certain embodiments, the tether group includes an acetal group, a disulfide, a hydrazine, or an ester. In some embodiments, the tether group includes an acetal group. In some embodiments, the tether group includes a disulfide. In some embodiments, the tether group includes a hydrazine. In some embodiments, the tether group includes an ester.

[00261] As described above, in some embodiments, L is a linker comprising $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d$, where a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4.


[00262] In some embodiments, in the subject linker:

T¹ is selected from a (C₁-C₁₂)alkyl and a substituted (C₁-C₁₂)alkyl;


T², T³ and T⁴ are each independently selected from (C₁-C₁₂)alkyl, substituted (C₁-C₁₂)alkyl, (EDA)_w, (PEG)_n, (AA)_p, -(CR¹³OH)_h-, 4-amino-piperidine (4AP), an acetal group, a disulfide, a hydrazine, and an ester; and


V¹, V², V³ and V⁴ are each independently selected from a covalent bond, -CO-, -NR¹⁵-, -NR¹⁵(CH₂)_q-, -NR¹⁵(C₆H₄)-, -CONR¹⁵-, -NR¹⁵CO-, -C(O)O-, -OC(O)-, -O-, -S-, -S(O)-, -SO₂-, -SO₂NR¹⁵-, -NR¹⁵SO₂- and -P(O)OH-, wherein q is an integer from 1 to 6;

wherein:

(PEG)_n is , where n is an integer from 1 to 30;

EDA is an ethylene diamine moiety having the following structure:

, where y is an integer from 1 to 6 and r is 0 or 1;

4-amino-piperidine (4AP) is 

AA is an amino acid residue, where p is an integer from 1 to 20; and

each R¹⁵ and R¹² is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl and a substituted aryl, wherein any two adjacent R¹² groups may be cyclically linked to form a piperazinyll ring; and

R¹³ is selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl.

[00263] In certain embodiments, T¹, T², T³ and T⁴ and V¹, V², V³ and V⁴ are selected from the following table, e.g., one row of the following table:

T ¹	V ¹	V ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-	-	-	-	-

T ¹	V ¹	V ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-CO-	(EDA) _w	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CONR ¹⁵ - -	(C ₁ -C ₁₂)alkyl	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-SO ₂ -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CONR ¹⁵ - -	(PEG) _n	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(CR ¹³ OH) _h	-CO-	-	-	-	-

T ¹	V ¹	V ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	substituted (C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-SO ₂ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-	(CR ¹³ OH) _h	-CONR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-P(O)OH -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	-	-

[00264] In certain embodiments, L is a linker comprising -(L¹)_a-(L²)_b-(L³)_c-(L⁴)_d-, where -(L¹)_a- is -(T¹-V¹)_a-, -(L²)_b- is -(T²-V²)_b-, -(L³)_c- is -(T³-V³)_c-, and -(L⁴)_d- is -(T⁴-V⁴)_d-.

[00265] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (AA)_p, V² is -NR¹⁵-, T³ is (PEG)_n, V³ is -CO-, T⁴ is absent and V⁴ is absent.

[00266] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (EDA)_w, V² is -CO-, T³ is (CR¹³OH)_h, V³ is -CONR¹⁵-, T⁴ is (C₁-C₁₂)alkyl and V⁴ is -CO-.

[00267] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (AA)_p, V² is -NR¹⁵-, T³ is (C₁-C₁₂)alkyl, V³ is -CO-, T⁴ is absent and V⁴ is absent.

[00268] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CONR¹⁵-, T² is (PEG)_n, V² is -CO-, T³ is absent, V³ is absent, T⁴ is absent and V⁴ is absent.

[00269] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (AA)_p, V² is absent, T³ is absent, V³ is absent, T⁴ is absent and V⁴ is absent.

[00270] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CONR¹⁵-, T² is (PEG)_n, V² is -NR¹⁵-, T³ is absent, V³ is absent, T⁴ is absent and V⁴ is absent.

[00271] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (AA)_p, V² is -NR¹⁵-, T³ is (PEG)_n, V³ is -NR¹⁵-, T⁴ is absent and V⁴ is absent.

[00272] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (EDA)_w, V² is -CO-, T³ is absent, V³ is absent, T⁴ is absent and V⁴ is absent.

[00273] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CONR¹⁵-, T² is (C₁-C₁₂)alkyl, V² is -NR¹⁵-, T³ is absent, V³ is absent, T⁴ is absent and V⁴ is absent.

[00274] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CONR¹⁵-, T² is (PEG)_n, V² is -CO-, T³ is (EDA)_w, V³ is absent, T⁴ is absent and V⁴ is absent.

[00275] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (EDA)_w, V² is absent, T³ is absent, V³ is absent, T⁴ is absent and V⁴ is absent.

[00276] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CONR¹⁵-, T² is (PEG)_n, V² is -CO-, T³ is (AA)_p, V³ is absent, T⁴ is absent and V⁴ is absent.

[00277] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (EDA)_w, V² is -CO-, T³ is (CR¹³OH)_h, V³ is -CO-, T⁴ is (AA)_p and V⁴ is absent.

[00278] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (AA)_p, V² is -NR¹⁵-, T³ is (C₁-C₁₂)alkyl, V³ is -CO-, T⁴ is (AA)_p and V⁴ is absent.

[00279] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (AA)_p, V² is -NR¹⁵-, T³ is (PEG)_n, V³ is -CO-, T⁴ is (AA)_p and V⁴ is absent.

[00280] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (AA)_p, V² is -NR¹¹-, T³ is (PEG)_n, V³ is -SO₂-, T⁴ is (AA)_p and V⁴ is absent.

[00281] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (EDA)_w, V² is -CO-, T³ is (CR¹³OH)_h, V³ is -CONR¹⁵-, T⁴ is (PEG)_n and V⁴ is -CO-.

[00282] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (CR¹³OH)_h, V² is -CO-, T³ is absent, V³ is absent, T⁴ is absent and V⁴ is absent.

[00283] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CONR¹⁵-, T² is substituted (C₁-C₁₂)alkyl, V² is -NR¹⁵-, T³ is (PEG)_n, V³ is -CO-, T⁴ is absent and V⁴ is absent.

[00284] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -SO₂-, T² is (C₁-C₁₂)alkyl, V² is -CO-, T³ is absent, V³ is absent, T⁴ is absent and V⁴ is absent.

[00285] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CONR¹⁵-, T² is (C₁-C₁₂)alkyl, V² is absent, T³ is (CR¹³OH)_h, V³ is -CONR¹⁵-, T⁴ is absent and V⁴ is absent.

[00286] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (AA)_p, V² is -NR¹⁵-, T³ is (PEG)_n, V³ is -CO-, T⁴ is (AA)_p and V⁴ is -NR¹⁵-.

[00287] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (AA)_p, V² is -NR¹⁵-, T³ is (PEG)_n, V³ is -P(O)OH-, T⁴ is (AA)_p and V⁴ is absent.

[00288] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (EDA)_w, V² is absent, T³ is (AA)_p, V³ is absent, T⁴ is absent and V⁴ is absent.

[00289] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (EDA)_w, V² is -CO-, T³ is (CR¹³OH)_h, V³ is -CONR¹⁵-, T⁴ is (C₁-C₁₂)alkyl and V⁴ is -CO(AA)_p-.

[00290] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CONR¹⁵-, T² is (C₁-C₁₂)alkyl, V² is -NR¹⁵-, T³ is absent, V³ is -CO-, T⁴ is absent and V⁴ is absent.


[00291] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CONR¹⁵-, T² is (C₁-C₁₂)alkyl, V² is -NR¹⁵-, T³ is absent, V³ is -CO-, T⁴ is (C₁-C₁₂)alkyl and V⁴ is -NR¹⁵-.


[00292] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is (EDA)_w, V² is -CO-, T³ is (CR¹³OH)_h, V³ is -CONR¹⁵-, T⁴ is (PEG)_n and V⁴ is -CO(AA)_p-.


[00293] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is 4AP, V² is -CO-, T³ is (C₁-C₁₂)alkyl, V³ is -CO-, T⁴ is (AA)_p and V⁴ is absent.

[00294] In certain embodiments, T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is 4AP, V² is -CO-, T³ is (C₁-C₁₂)alkyl, V³ is -CO-, T⁴ is absent and V⁴ is absent.

[00295] In certain embodiments, the linker is described by one of the following structures:








[00296] In certain embodiments of the linker structures depicted above, each f is independently 0 or an integer from 1 to 12; each y is independently 0 or an integer from 1 to 20; each n is independently 0 or an integer from 1 to 30; each p is independently 0 or an integer from 1 to 20; each h is independently 0 or an integer from 1 to 12; each R is independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and each R' is independently H, a sidechain of an amino acid, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl,

cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl. In certain embodiments of the linker structures depicted above, each f is independently 0, 1, 2, 3, 4, 5 or 6; each y is independently 0, 1, 2, 3, 4, 5 or 6; each n is independently 0, 1, 2, 3, 4, 5 or 6; each p is independently 0, 1, 2, 3, 4, 5 or 6; and each h is independently 0, 1, 2, 3, 4, 5 or 6. In certain embodiments of the linker structures depicted above, each R is independently H, methyl or $-(CH_2)_m-OH$ where m is 1, 2, 3 or 4 (e.g., 2).


[00297] In certain embodiments of the linker, L , T^1 is (C_1-C_{12}) alkyl, V^1 is $-CO-$, T^2 is 4AP, V^2 is $-CO-$, T^3 is (C_1-C_{12}) alkyl, V^3 is $-CO-$, T^4 is absent and V^4 is absent. In certain embodiments, T^1 is ethylene, V^1 is $-CO-$, T^2 is 4AP, V^2 is $-CO-$, T^3 is ethylene, V^3 is $-CO-$, T^4 is absent and V^4 is absent. In certain embodiments, T^1 is ethylene, V^1 is $-CO-$, T^2 is 4AP, V^2 is $-CO-$, T^3 is ethylene, V^3 is $-CO-$, T^4 is absent and V^4 is absent, where T^2 (e.g., 4AP) has the following structure:



wherein

R^{12} is a polyethylene glycol moiety (e.g., a polyethylene glycol or a modified polyethylene glycol).


[00298] In certain embodiments, the linker, L , includes the following structure:



wherein

each f is independently an integer from 1 to 12; and
 n is an integer from 1 to 30.

[00301] In certain embodiments, the linker, L, includes the following structure:



n is an integer from 1 to 30.

[00306] Additional disclosure related to hydrazinyl-indolyl and hydrazinyl-pyrrolo-pyridinyl compounds and methods for producing a conjugate is found in U.S. Application Publication No. 2014/0141025, filed March 11, 2013, and U.S. Application Publication No. 2015/0157736, filed November 26, 2014, the disclosures of each of which are incorporated herein by reference.

ANTI-CD22 ANTIBODIES

[00307] As noted above, a subject conjugate can comprise, as substituent W² an anti-CD22 antibody, where the anti-CD22 antibody has been modified to include a 2-formylglycine (FGly) residue. As used herein, amino acids may be referred to by their standard name, their standard three letter abbreviation and/or their standard one letter abbreviation, such as: Alanine or Ala or A; Cysteine or Cys or C; Aspartic acid or Asp or D; Glutamic acid or Glu or E; Phenylalanine or Phe or F; Glycine or Gly or G; Histidine or His or H; Isoleucine or Ile or I; Lysine or Lys or K; Leucine or Leu or L; Methionine or Met or M; Asparagine or Asn or N; Proline or Pro or P; Glutamine or Gln or Q; Arginine or Arg or R; Serine or Ser or S; Threonine or Thr or T; Valine or Val or V; Tryptophan or Trp or W; and Tyrosine or Tyr or Y.

[00308] In some cases, a suitable anti-CD22 antibody specifically binds a CD22 polypeptide, where the epitope comprises amino acid residues within a CD22 antigen (e.g., within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C).

[00309] The CD22 epitope can be formed by a polypeptide having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity to a contiguous stretch of from about 500 amino acids to about 670 amino acids of the human CD22 isoform 4 amino acid sequence depicted in FIG. 8A-8C. The CD22 epitope can be formed by a polypeptide having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity to a contiguous stretch of from about 500 amino acids to about 751 amino acids of the human CD22 isoform 3 amino acid sequence depicted in FIG. 8A-8C. The CD22 epitope can be formed by a polypeptide having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity to a contiguous stretch of from about 500 amino acids to about 759 amino acids of the human CD22 isoform 2 amino acid sequence depicted in FIG. 8A-8C. The CD22 epitope can be formed by a polypeptide having at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity to a contiguous stretch of from about 500 amino acids to about 847 amino acids of the human CD22 isoform 1 amino acid sequence depicted in FIG. 8A-8C.

[00310] A “CD22 antigen” or “CD22 polypeptide” can comprises an amino acid sequence having at least about 75%, at least about 80%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%, amino acid sequence identity to a contiguous stretch of from about 500 amino acids (aa) to about 847 aa (isoform 1), to about 759 aa (isoform 2), to about 751 aa (isoform 3), or to about 670 aa (isoform 4) of a CD22 isoform 1, 2, 3, or 4 amino acid sequence depicted in FIG. 8A-8C.

[00311] In some cases, a suitable anti-CD22 antibody exhibits high affinity binding to CD22. For example, in some cases, a suitable anti-CD22 antibody binds to CD22 with an affinity of at least about 10^{-7} M, at least about 10^{-8} M, at least about 10^{-9} M, at least about 10^{-10} M, at least about 10^{-11} M, or at least about 10^{-12} M, or greater than 10^{-12} M. In some cases, a suitable anti-CD22 antibody binds to an epitope present on CD22 with an affinity of from about 10^{-7} M to about 10^{-8} M, from about 10^{-8} M to about 10^{-9} M, from about 10^{-9} M to about 10^{-10} M, from about 10^{-10} M to about 10^{-11} M, or from about 10^{-11} M to about 10^{-12} M, or greater than 10^{-12} M.

[00312] In some cases, a suitable anti-CD22 antibody competes for binding to an epitope within CD22 with a second anti-CD22 antibody and/or binds to the same epitope within CD22, as a second anti-CD22 antibody. In some cases, an anti-CD22 antibody that competes for binding to an epitope within CD22 with a second anti-CD22 antibody also binds to the epitope as the second anti-CD22 antibody. In some cases, an anti-CD22 antibody that competes for binding to an epitope within CD22 with a second anti-CD22 antibody binds to an epitope that is overlapping with the epitope bound by the second anti-CD22 antibody. In some cases, the anti-CD22 antibody is humanized.

[00313] In some cases, a suitable anti-CD22 antibody can induce apoptosis in a cell that expresses CD22 on its cell surface.

[00314] An anti-CD22 antibody suitable for use in a subject conjugate will in some cases inhibit the proliferation of human tumor cells that overexpress CD22, where the inhibition occurs *in vitro*, *in vivo*, or both *in vitro* and *in vivo*. For example, in some cases, an anti-CD22 antibody suitable for use in a subject conjugate inhibits proliferation of human tumor cells that overexpress CD22 by at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, or more than 80%, e.g., by at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or 100%.

[00315] In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope comprising amino acid residues within a CD22 antigen (e.g., within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody comprising a heavy chain complementarity determining region (CDR) selected from **IYDMS** (VH CDR1; SEQ ID NO: 17), **YISSGGGTTYYPDTVKG** (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAF** (VH CDR3; SEQ ID NO: 19). In some cases, the anti-CD22 antibody is humanized. In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope comprising amino acid residues within a CD22 antigen (e.g., within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody comprising a light-chain CDR selected from **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized.

[00316] In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope comprising amino acid residues within a CD22 antigen (e.g., within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody comprising VH CDRs **IYDMS** (VH CDR1; SEQ ID NO: 17), **YISSGGGTTYYPDTVKG** (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAF** (VH CDR3; SEQ ID NO: 19). In some cases, the anti-CD22 antibody is humanized. In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope comprising amino acid residues within a CD22 antigen (e.g., an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody comprising VL CDRs **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized. In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope comprising amino acid residues within a CD22 antigen (e.g., within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody that comprises VH CDRs **IYDMS** (VH CDR1; SEQ ID NO: 17),

YISSGGGTTYYPDTVKG (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAFAY** (VH CDR3; SEQ ID NO: 19) and VL CDRs **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized.

[00317] In some cases, a suitable anti-CD22 antibody comprises VH CDRs **IYDMS** (VH CDR1; SEQ ID NO: 17), **YISSGGGTTYYPDTVKG** (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAFAY** (VH CDR3; SEQ ID NO: 19). In some cases, the anti-CD22 antibody is humanized. In some cases, a suitable anti-CD22 antibody comprises VL CDRs **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized. In some cases, a suitable anti-CD22 antibody comprises VH CDRs **IYDMS** (VH CDR1; SEQ ID NO: 17), **YISSGGGTTYYPDTVKG** (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAFAY** (VH CDR3; SEQ ID NO: 19) and VL CDRs **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized.

[00318] In some cases, a suitable anti-CD22 antibody comprises VH CDRs present in an anti-CD22 VH region comprising the following amino acid sequence:

**EVQLVESGGGLVKPGGSLRLSCAASGFAFSIYDMSWVRQAPGKGLEWVAYISSGG
GTTYYPDTVKGKFTISRDNKNSLYLQMSSLRAEDTAMYYCARHSGYGSSYGVLFAFAYWGQGT**
LVTVSS (SEQ ID NO: 23). In some cases, the anti-CD22 antibody is humanized.

[00319] In some cases, a suitable anti-CD22 antibody comprises VL CDRs present in an anti-CD22 VL region comprising the following amino acid sequence:

**DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPGKAVKLLIYYTSILHSG
VPSRFSGSGSGTDYTLTISSLQQEDFATYFCQQGNTLPWTFGGGGTKVEIKR** (SEQ ID
NO: 24). In some cases, the anti-CD22 antibody is humanized.

[00320] In some cases, a suitable anti-CD22 antibody comprises VH CDRs present in **EVQLVESGGGLVKPGGSLRLSCAASGFAFSIYDMSWVRQAPGKGLEWVAYISSGG
GTTYYPDTVKGKFTISRDNKNSLYLQMSSLRAEDTAMYYCARHSGYGSSYGVLFAFAYWGQGT**
LVTVSS (SEQ ID NO: 23) and VL CDRs present in **DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPGKAVKLLIYYTSILHSG**

VPSRFSGSGSGTDYTLTISSLQQEDFATYFCQQGNTLPWTFGGGGTKVEIKR (SEQ ID NO: 24). In some cases, the anti-CD22 antibody is humanized.

[00321] In some cases, a suitable anti-CD22 antibody comprises: a) a heavy chain comprising a VH region having the amino acid sequence

**EVQLVESGGGLVKPGGSLX¹LSCAASGFAFSIYDMSWVRQAPGKGLEWVAYISSGG
GTTYYPDTVKGRTISRDNAX²LYLQMX³SLRAEDTAMYYCARHSGYGSYGV
LAYWGQGTLLTVSS** (SEQ ID NO: 25), where X¹ is K (Lys) or R (Arg); X² is S (Ser) or T (Thr); and X³ is N (Asn) or S (Ser); and b) an immunoglobulin light chain.

[00322] A light chain can have any suitable VL amino acid sequence, so long as the resulting antibody binds specifically to CD22.

[00323] Exemplary VL amino acid sequences include:

[00324] **DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPGKAVKLLI
YYTSILHSGVPSRFSGSGSGTDYTLTISSLQQEDFATYFCQQGNTLPWTFGGGGTKVE
IKR** (SEQ ID NO: 24; VK1);

[00325] **DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPGKAVKLLI
YYTSILHSGVPSRFSGSGSGTDYTLTISSLQPEDFATYFCQQGNTLPWTFGGGGTKVE
IKR** (SEQ ID NO: 26; VK2); and

[00326] **DIQMTQSPSSVSASVGDRVTITCRASQDISNYLNWYQQKPGKAPKLLI
YYTSILHSGVPSRFSGSGSGTDYTLTISSLQPEDFATYFCQQGNTLPWTFGGGGTKVE
IKR** (SEQ ID NO: 27; VK4).

[00327] Thus, e.g., a suitable anti-CD22 antibody can comprise: a) a heavy chain comprising a VH region having the amino acid sequence set forth in SEQ ID NO: 25; and a light chain comprising the VL region of VK1 (SEQ ID NO: 24). In other cases, a suitable anti-CD22 antibody can comprise: a) a heavy chain comprising a VH region having the amino acid sequence set forth in SEQ ID NO: 25; and a light chain comprising the VL region of VK2 (SEQ ID NO: 26). In still other cases, a subject anti-CD22 antibody can comprise: a) a heavy chain comprising a VH region having the amino acid sequence set forth in SEQ ID NO: 25; and a light chain comprising the VL region of VK4 (SEQ ID NO: 27).

[00328] In some instances, a suitable anti-CD22 antibody comprises: a) an immunoglobulin light chain comprising the amino acid sequence

DIQMTQSPSSX¹SASVGDRVTITCRASQDISNYLNWYQQKPGKAX²KLLIYYTSILHS

GVPSRFSGSGSGTDYTLTISSLQX³EDFATYFCQQGNTLPWTFGGGTKVEIK (SEQ ID NO: 28), where X¹ is L (Leu) or V (Val); X² is V (Val) or P (Pro); and X³ is Q (Gln) or P (Pro); and b) an immunoglobulin heavy chain. The heavy chain can comprise an amino acid sequence selected from:

**[00329] EVQLVESGGGLVKPGGSLKLSCAASGFAFSIYDMSWVRQAPGKGLE
WVAYISSGGGTYYPDYTVKGRFTISRDNKNTLYLQMSSLRAEDTAMYYCARHSG
YGSSYGVLFAYWGQGTLVTVSS** (SEQ ID NO: 29; VH3);

**[00330] EVQLVESGGGLVKPGGSLRLSCAASGFAFSIYDMSWVRQAPGKGLE
WVAYISSGGGTYYPDYTVKGRFTISRDNKNSLYLQMSSLRAEDTAMYYCARHSG
YGSSYGVLFAYWGQGTLVTVSS** (SEQ ID NO: 23; VH4);

**[00331] EVQLVESGGGLVKPGGSLKLSCAASGFAFSIYDMSWVRQAPGKGLE
WVAYISSGGGTYYPDYTVKGRFTISRDNKNSLYLQMNSLRAEDTAMYYCARHSG
YGSSYGVLFAYWGQGTLVTVSS** (SEQ ID NO: 30; VH5); and

**[00332] EVQLVESGGGLVKPGGSLKLSCAASGFAFSIYDMSWVRQAPGKGLE
WVAYISSGGGTYYPDYTVKGRFTISRDNKNSLYLQMSSLRAEDTAMYYCARHSG
YGSSYGVLFAYWGQGTLVTVSS** (SEQ ID NO: 31; VH6).

[00333] In some cases, a suitable anti-CD22 antibody comprises a VH region comprising the following amino acid sequence:

**EVQLVESGGGLVKPGGSLRLSCAASGFAFSIYDMSWVRQAPGKGLEWVAYISSGG
GTTYYPDYTVKGRFTISRDNKNSLYLQMSSLRAEDTAMYYCARHSGYGSSYGVLF
AYWGQGTLVTVSS** (SEQ ID NO: 23).

[00334] In some cases, a suitable anti-CD22 antibody comprises a VH region comprising the following amino acid sequence:

**EVQLVESGGGLVKPGGSLRLSCAASGFAFSIYDMSWVRQAPGKGLEWVAYISSGG
GTTYYPDYTVKGRFTISRDNKNSLYLQMSSLRAEDTAMYYCARHSGYGSSYGVLF
AYWGQGTLVTVSS** (SEQ ID NO: 23) and VL region comprising the following amino acid sequence:

**DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPKGKAVKLLIYYTSILHSG
VPSRFSGSGSGTDYTLTISSLQQEDFATYFCQQGNTLPWTFGGGTKVEIKR** (SEQ ID NO: 24).

Modified constant region sequences

[00335] As noted above, the amino acid sequence of an anti-CD22 antibody is modified to include a sulfatase motif that contains a serine or cysteine residue that is capable of being converted (oxidized) to a 2-formylglycine (FGly) residue by action of a formylglycine generating enzyme (FGE) either *in vivo* (e.g., at the time of translation of an ald tag-containing protein in a cell) or *in vitro* (e.g., by contacting an ald tag-containing protein with an FGE in a cell-free system). Such sulfatase motifs may also be referred to herein as an FGE-modification site.

Sulfatase motifs

[00336] A minimal sulfatase motif of an aldehyde tag is usually 5 or 6 amino acid residues in length, usually no more than 6 amino acid residues in length. Sulfatase motifs provided in an Ig polypeptide are at least 5 or 6 amino acid residues, and can be, for example, from 5 to 16, 6-16, 5-15, 6-15, 5-14, 6-14, 5-13, 6-13, 5-12, 6-12, 5-11, 6-11, 5-10, 6-10, 5-9, 6-9, 5-8, or 6-8 amino acid residues in length, so as to define a sulfatase motif of less than 16, 15, 14, 13, 12, 11, 10, 9, 8 or 7 amino acid residues in length.


[00337] In certain embodiments, polypeptides of interest include those where one or more amino acid residues, such as 2 or more, or 3 or more, or 4 or more, or 5 or more, or 6 or more, or 7 or more, or 8 or more, or 9 or more, or 10 or more, or 11 or more, or 12 or more, or 13 or more, or 14 or more, or 15 or more, or 16 or more, or 17 or more, or 18 or more, or 19 or more, or 20 or more amino acid residues have been inserted, deleted, substituted (replaced) relative to the native amino acid sequence to provide for a sequence of a sulfatase motif in the polypeptide. In certain embodiments, the polypeptide includes a modification (insertion, addition, deletion, and/or substitution/replacement) of less than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3 or 2 amino acid residues of the amino acid sequence relative to the native amino acid sequence of the polypeptide. Where an amino acid sequence native to the polypeptide (e.g., anti-CD22 antibody) contains one or more residues of the desired sulfatase motif, the total number of modifications of residues can be reduced, e.g., by site-specification modification (insertion, addition, deletion, substitution/replacement) of amino acid residues flanking the native amino acid residues to provide a sequence of the desired sulfatase motif. In certain embodiments, the extent of modification of the native amino acid sequence of the target anti-CD22 polypeptide is minimized, so as to minimize the number of amino acid residues that are inserted, deleted, substituted (replaced), or added (e.g., to the N- or C-terminus). Minimizing the extent of amino

acid sequence modification of the target anti-CD22 polypeptide may minimize the impact such modifications may have upon anti-CD22 function and/or structure.

[00338] It should be noted that while aldehyde tags of particular interest are those comprising at least a minimal sulfatase motif (also referred to a “consensus sulfatase motif”), it will be readily appreciated that longer aldehyde tags are both contemplated and encompassed by the present disclosure and can find use in the compositions and methods of the present disclosure. Aldehyde tags can thus comprise a minimal sulfatase motif of 5 or 6 residues, or can be longer and comprise a minimal sulfatase motif which can be flanked at the N- and/or C-terminal sides of the motif by additional amino acid residues. Aldehyde tags of, for example, 5 or 6 amino acid residues are contemplated, as well as longer amino acid sequences of more than 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more amino acid residues.

[00339] An aldehyde tag can be present at or near the C-terminus of an Ig heavy chain; e.g., an aldehyde tag can be present within 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids of the C-terminus of a native, wild-type Ig heavy chain. An aldehyde tag can be present within a CH1 domain of an Ig heavy chain. An aldehyde tag can be present within a CH2 domain of an Ig heavy chain. An aldehyde tag can be present within a CH3 domain of an Ig heavy chain. An aldehyde tag can be present in an Ig light chain constant region, e.g., in a kappa light chain constant region or a lambda light chain constant region.

[00340] In certain embodiments, the sulfatase motif used may be described by the formula (SEQ ID NOs: 191 – 192):



where

Z^{10} is cysteine or serine (which can also be represented by (C/S));

Z^{20} is either a proline or alanine residue (which can also be represented by (P/A));

Z^{30} is a basic amino acid (e.g., arginine (R), and may be lysine (K) or histidine (H), e.g., lysine), or an aliphatic amino acid (alanine (A), glycine (G), leucine (L), valine (V), isoleucine (I), or proline (P), e.g., A, G, L, V, or I;

X^1 is present (SEQ ID NO: 191) or absent (SEQ ID NO: 192) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid

or a charged amino acid), e.g., L, M, V, S or T, e.g., L, M, S or V, with the proviso that when the sulfatase motif is at the N-terminus of the target polypeptide, X^1 is present; and

X^2 and X^3 independently can be any amino acid, (e.g., any naturally-occurring amino acid), though usually an aliphatic amino acid, a polar, uncharged amino acid, or a sulfur containing amino acid (i.e., other than an aromatic amino acid or a charged amino acid), e.g., S, T, A, V, G or C, e.g., S, T, A, V or G.

[00341] The amino acid sequence of an anti-CD22 heavy and/or light chain can be modified to provide a sequence of at least 5 amino acids of the formula $X^1Z^{10}X^2Z^{20}X^3Z^{30}$, where

Z^{10} is cysteine or serine;

Z^{20} is a proline or alanine residue;

Z^{30} is an aliphatic amino acid or a basic amino acid;


X^1 is present (SEQ ID NO: 191) or absent (SEQ ID NO: 192) and, when present, is any amino acid, (e.g., any naturally-occurring amino acid), with the proviso that when the heterologous sulfatase motif is at an N-terminus of the polypeptide, X^1 is present;

X^2 and X^3 are each independently any amino acid, (e.g., any naturally-occurring amino acid),

where the sequence is within or adjacent a solvent-accessible loop region of the Ig constant region, and wherein the sequence is not at the C-terminus of the Ig heavy chain.

[00342] The sulfatase motif is generally selected so as to be capable of conversion by a selected FGE, e.g., an FGE present in a host cell in which the aldehyde tagged polypeptide is expressed or an FGE which is to be contacted with the aldehyde tagged polypeptide in a cell-free *in vitro* method.

[00343] For example, where the FGE is a eukaryotic FGE (e.g., a mammalian FGE, including a human FGE), the sulfatase motif can be of the formula (SEQ ID NOs: 193 – 194):



where

X^1 may be present (SEQ ID NO: 193) or absent (SEQ ID NO: 194) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., L, M, S or V, with the proviso that when the sulfatase motif is at the N-terminus of the target polypeptide, X^1 is present;


X^2 and X^3 independently can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., S, T, A, V, G, or C, e.g., S, T, A, V or G; and

Z^{30} is a basic amino acid (e.g., arginine (R), and may be lysine (K) or histidine (H), e.g., lysine), or an aliphatic amino acid (alanine (A), glycine (G), leucine (L), valine (V), isoleucine (I), or proline (P), e.g., A, G, L, V, or I.

[00344] Specific examples of sulfatase motifs include **LCTPSR** (SEQ ID NO: 32), **MCTPSR** (SEQ ID NO: 33), **VCTPSR** (SEQ ID NO: 34), **LCSPSR** (SEQ ID NO: 35), **LCAPSR** (SEQ ID NO: 36), **LCVPSR** (SEQ ID NO: 37), **LCGPSR** (SEQ ID NO: 38), **ICTPAR** (SEQ ID NO: 39), **LCTPSK** (SEQ ID NO: 40), **MCTPSK** (SEQ ID NO: 41), **VCTPSK** (SEQ ID NO: 42), **LCSPSK** (SEQ ID NO: 43), **LCAPSK** (SEQ ID NO: 44), **LCVPSK** (SEQ ID NO: 45), **LCGPSK** (SEQ ID NO: 46), **LCTPSA** (SEQ ID NO: 47), **ICTPAA** (SEQ ID NO: 48), **MCTPSA** (SEQ ID NO: 49), **VCTPSA** (SEQ ID NO: 50), **LCSPSA** (SEQ ID NO: 51), **LCAPSA** (SEQ ID NO: 52), **LCVPSA** (SEQ ID NO: 53), and **LCGPSA** (SEQ ID NO: 54).

FGly-containing sequences

[00345] Upon action of FGE on the modified anti-CD22 heavy and/or light chain, the serine or the cysteine in the sulfatase motif is modified to FGly. Thus, the FGly-containing sulfatase motif can be of the formula (SEQ ID NOs: 187 and 188):



where

FGly is the formylglycine residue;

Z^{20} is either a proline or alanine residue (which can also be represented by (P/A));


Z^{30} is a basic amino acid (e.g., arginine (R), and may be lysine (K) or histidine (H), usually lysine), or an aliphatic amino acid (alanine (A), glycine (G), leucine (L), valine (V), isoleucine (I), or proline (P), e.g., A, G, L, V, or I;

X^1 may be present (i.e., SEQ ID NO: 187) or absent (i.e., SEQ ID NO: 188) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., L, M, V, S or T, e.g., L, M or V, with the

proviso that when the sulfatase motif is at the N-terminus of the target polypeptide, X¹ is present; and

X² and X³ independently can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., S, T, A, V, G or C, e.g., S, T, A, V or G.

[00346] As described above, the modified polypeptide containing the FGly residue may be conjugated to a drug (e.g., a maytansinoid) by reaction of the FGly with the drug (e.g., a drug containing a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety, as described above) to produce an FGly'-containing sulfatase motif. As used herein, the term FGly' refers to the modified amino acid residue of the sulfatase motif that is coupled to the drug, such as a maytansinoid (e.g., the modified amino acid residue of formula (I)). Thus, the FGly'-containing sulfatase motif can be of the formula (SEQ ID NOs: 189 – 190):



where

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue (which can also be represented by (P/A));


Z³⁰ is a basic amino acid (e.g., arginine (R), and may be lysine (K) or histidine (H), usually lysine), or an aliphatic amino acid (alanine (A), glycine (G), leucine (L), valine (V), isoleucine (I), or proline (P), e.g., A, G, L, V, or I;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., L, M, V, S or T, e.g., L, M or V, with the proviso that when the sulfatase motif is at the N-terminus of the target polypeptide, X¹ is present; and

X² and X³ independently can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., S, T, A, V, G or C, e.g., S, T, A, V or G.

[00347] In certain embodiments, the modified amino acid residue of formula (I) is positioned at a C-terminus of a heavy chain constant region of the anti-CD22 antibody. In some

instances, the heavy chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue (which can also be represented by (P/A));

Z^{30} is a basic amino acid (e.g., arginine (R), and may be lysine (K) or histidine (H), usually lysine), or an aliphatic amino acid (alanine (A), glycine (G), leucine (L), valine (V), isoleucine (I), or proline (P), e.g., A, G, L, V, or I;


X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., L, M, V, S or T, e.g., L, M or V, with the proviso that when the sulfatase motif is at the N-terminus of the target polypeptide, X^1 is present;

X^2 and X^3 independently can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., S, T, A, V, G or C, e.g., S, T, A, V or G; and

wherein the sequence is C-terminal to the amino acid sequence **QKSLSLSPGK** (SEQ ID NO: 55), and where the sequence may include 1, 2, 3, 4, 5, or from 5 to 10, amino acids not present in a native, wild-type heavy Ig chain constant region.

[00348] In certain embodiments, the heavy chain constant region comprises the sequence **SLSLSPGSL(FGly')TPSRGS** (SEQ ID NO: 56) at the C-terminus of the Ig heavy chain, e.g., in place of a native **SLSLSPGK** (SEQ ID NO: 57) sequence.

[00349] In certain embodiments, the modified amino acid residue of formula (I) is positioned in a light chain constant region of the anti-CD22 antibody. In certain embodiments, the light chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue (which can also be represented by (P/A));

Z^{30} is a basic amino acid (e.g., arginine (R), and may be lysine (K) or histidine (H), usually lysine), or an aliphatic amino acid (alanine (A), glycine (G), leucine (L), valine (V), isoleucine (I), or proline (P), e.g., A, G, L, V, or I;


X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., L, M, V, S or T, e.g., L, M or V, with the proviso that when the sulfatase motif is at the N-terminus of the target polypeptide, X^1 is present;

X^2 and X^3 independently can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., S, T, A, V, G or C, e.g., S, T, A, V or G; and

wherein the sequence is C-terminal to the amino acid sequence **KVDNAL** (SEQ ID NO: 58) and/or is N-terminal to the amino acid sequence **QSGNSQ** (SEQ ID NO: 59).

[00350] In certain embodiments, the light chain constant region comprises the sequence **KVDNAL(FGly')TPSRQSGNSQ** (SEQ ID NO: 60).

[00351] In certain embodiments, the modified amino acid residue of formula (I) is positioned in a heavy chain CH1 region of the anti-CD22 antibody. In certain embodiments, the heavy chain CH1 region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue (which can also be represented by (P/A));

Z^{30} is a basic amino acid (e.g., arginine (R), and may be lysine (K) or histidine (H), usually lysine), or an aliphatic amino acid (alanine (A), glycine (G), leucine (L), valine (V), isoleucine (I), or proline (P), e.g., A, G, L, V, or I;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic

amino acid or a charged amino acid), e.g., L, M, V, S or T, e.g., L, M or V, with the proviso that when the sulfatase motif is at the N-terminus of the target polypeptide, X¹ is present;

X² and X³ independently can be any amino acid, (e.g., any naturally-occurring amino acid), e.g., an aliphatic amino acid, a sulfur-containing amino acid, or a polar, uncharged amino acid, (i.e., other than an aromatic amino acid or a charged amino acid), e.g., S, T, A, V, G or C, e.g., S, T, A, V or G; and

wherein the sequence is C-terminal to the amino acid sequence **SWNSGA** (SEQ ID NO: 61) and/or is N-terminal to the amino acid sequence **GVHTEP** (SEQ ID NO: 62).

[00352] In certain embodiments, the heavy chain CH1 region comprises the sequence **SWNSGAL(FGly')TPSRGVHTEP** (SEQ ID NO: 63).

Site of modification

[00353] As noted above, the amino acid sequence of an anti-CD22 antibody is modified to include a sulfatase motif that contains a serine or cysteine residue that is capable of being converted (oxidized) to an FGly residue by action of an FGE either *in vivo* (e.g., at the time of translation of an ald tag-containing protein in a cell) or *in vitro* (e.g., by contacting an ald tag-containing protein with an FGE in a cell-free system). The anti-CD22 polypeptides used to generate a conjugate of the present disclosure include at least an Ig constant region, e.g., an Ig heavy chain constant region (e.g., at least a CH1 domain; at least a CH1 and a CH2 domain; a CH1, a CH2, and a CH3 domain; or a CH1, a CH2, a CH3, and a CH4 domain), or an Ig light chain constant region. Such Ig polypeptides are referred to herein as “target Ig polypeptides” or “target anti-CD22 antibodies” or “target anti-CD22 Ig polypeptides.”

[00354] The site in an anti-CD22 antibody into which a sulfatase motif is introduced can be any convenient site. As noted above, in some instances, the extent of modification of the native amino acid sequence of the target anti-CD22 polypeptide is minimized, so as to minimize the number of amino acid residues that are inserted, deleted, substituted (replaced), and/or added (e.g., to the N- or C-terminus). Minimizing the extent of amino acid sequence modification of the target anti-CD22 polypeptide may minimize the impact such modifications may have upon anti-CD22 function and/or structure.

[00355] An anti-CD22 antibody heavy chain constant region can include Ig constant regions of any heavy chain isotype, non-naturally occurring Ig heavy chain constant regions (including consensus Ig heavy chain constant regions). An Ig constant region can be modified to

include an aldehyde tag, where the aldehyde tag is present in or adjacent a solvent-accessible loop region of the Ig constant region. An Ig constant region can be modified by insertion and/or substitution of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 amino acids, or more than 16 amino acids, to provide an amino acid sequence of a sulfatase motif as described above.

[00356] In some cases, an aldehyde-tagged anti-CD22 antibody comprises an aldehyde-tagged Ig heavy chain constant region (e.g., at least a CH1 domain; at least a CH1 and a CH2 domain; a CH1, a CH2, and a CH3 domain; or a CH1, a CH2, a CH3, and a CH4 domain). The aldehyde-tagged Ig heavy chain constant region can include heavy chain constant region sequences of an IgA, IgM, IgD, IgE, IgG1, IgG2, IgG3, or IgG4 isotype heavy chain or any allotypic variant of same, e.g., human heavy chain constant region sequences or mouse heavy chain constant region sequences, a hybrid heavy chain constant region, a synthetic heavy chain constant region, or a consensus heavy chain constant region sequence, etc., modified to include at least one sulfatase motif that can be modified by an FGE to generate an FGly-modified Ig polypeptide. Allotypic variants of Ig heavy chains are known in the art. See, e.g., Jefferis and Lefranc (2009) *MAbs* 1:4.

[00357] In some cases, an aldehyde-tagged anti-CD22 antibody comprises an aldehyde-tagged Ig light chain constant region. The aldehyde-tagged Ig light chain constant region can include constant region sequences of a kappa light chain, a lambda light chain, e.g., human kappa or lambda light chain constant regions, a hybrid light chain constant region, a synthetic light chain constant region, or a consensus light chain constant region sequence, etc., modified to include at least one sulfatase motif that can be modified by an FGE to generate an FGly-modified anti-CD22 antibody polypeptide. Exemplary constant regions include human gamma 1 and gamma 3 regions. With the exception of the sulfatase motif, a modified constant region may have a wild-type amino acid sequence, or it may have an amino acid sequence that is at least 70% identical (e.g., at least 80%, at least 90% or at least 95% identical) to a wild type amino acid sequence.

[00358] In some embodiments the sulfatase motif is at a position other than, or in addition to, the C-terminus of the Ig polypeptide heavy chain. As noted above, an isolated aldehyde-tagged anti-CD22 polypeptide can comprise a heavy chain constant region modified to include a sulfatase motif as described above, where the sulfatase motif is in or adjacent a surface-accessible loop region of the anti-CD22 polypeptide heavy chain constant region.

[00359] In some instances, a target anti-CD22 immunoglobulin is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 122-127; 2) amino acids 137-143; 3) amino acids 155-158; 4) amino acids 163-170; 5) amino acids 163-183; 6) amino acids 179-183; 7) amino acids 190-192; 8) amino acids 200-202; 9) amino acids 199-202; 10) amino acids 208-212; 11) amino acids 220-241; 12) amino acids 247-251; 13) amino acids 257-261; 14) amino acid 269-277; 15) amino acids 271-277; 16) amino acids 284-285; 17) amino acids 284-292; 18) amino acids 289-291; 19) amino acids 299-303; 20) amino acids 309-313; 21) amino acids 320-322; 22) amino acids 329-335; 23) amino acids 341-349; 24) amino acids 342-348; 25) amino acids 356-365; 26) amino acids 377-381; 27) amino acids 388-394; 28) amino acids 398-407; 29) amino acids 433-451; and 30) amino acids 446-451; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as depicted in Figure 9B.

[00360] In some instances, a target anti-CD22 immunoglobulin is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region; sequence depicted in Figure 9B).

[00361] Exemplary surface-accessible loop regions of an IgG1 heavy chain include: 1) **ASTKGP** (SEQ ID NO: 64); 2) **KSTSGGT** (SEQ ID NO: 65); 3) **PEPV** (SEQ ID NO: 66); 4) **NSGALTSG** (SEQ ID NO: 67); 5) **NSGALTSGVHTFPAVLQSSGL** (SEQ ID NO: 68); 6)

QSSGL (SEQ ID NO: 69); 7) **VTV** (SEQ ID NO: 70); 8) **QTY** (SEQ ID NO: 71); 9) **TQTY** (SEQ ID NO: 72); 10) **HKPSN** (SEQ ID NO: 73); 11) **EPKSCDKTHTCPPCPAPELLGG** (SEQ ID NO: 74); 12) **FPPKP** (SEQ ID NO: 75); 13) **ISRTP** (SEQ ID NO: 76); 14) **DVSHEDPEV** (SEQ ID NO: 77); 15) **SHEDPEV** (SEQ ID NO: 78); 16) **DG** (SEQ ID NO: 79); 17) **DGVEVHNAK** (SEQ ID NO: 80); 18) **HNA** (SEQ ID NO: 81); 19) **QYNST** (SEQ ID NO: 82); 20) **VLTVL** (SEQ ID NO: 83); 21) **GKE** (SEQ ID NO: 84); 22) **NKALPAP** (SEQ ID NO: 85); 23) **SKAKGQPRE** (SEQ ID NO: 86); 24) **KAKGQPR** (SEQ ID NO: 87); 25) **PPSRKELTKN** (SEQ ID NO: 88); 26) **YPSDI** (SEQ ID NO: 89); 27) **NGQPENN** (SEQ ID NO: 90); 28) **TPPVLDSGGS** (SEQ ID NO: 91); 29) **HEALHNHYTQKSLSLSPGK** (SEQ ID NO: 92); and 30) **SLSPGK** (SEQ ID NO: 93), as shown in Figures 9A and 9B.

[00362] In some instances, a target immunoglobulin is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG2 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6 ; 2) amino acids 13-24; 3) amino acids 33-37; 4) amino acids 43-54; 5) amino acids 58-63; 6) amino acids 69-71; 7) amino acids 78-80; 8) 87-89; 9) amino acids 95-96; 10) 114-118; 11) 122-126; 12) 134-136; 13) 144-152; 14) 159-167; 15) 175-176; 16) 184-188; 17) 195-197; 18) 204-210; 19) 216-224 ; 20) 231-233; 21) 237-241; 22) 252-256; 23) 263-269; 24) 273-282; 25) amino acids 299-302; where the amino acid numbering is based on the numbering of the amino acid sequence set forth in SEQ ID NO: 8 (human IgG2; also depicted in Figure 9B).

[00363] Exemplary surface-accessible loop regions of an IgG2 heavy chain include 1) **ASTKGP** (SEQ ID NO: 64); 2) **PCSRSTSESTAA** (SEQ ID NO: 94); 3) **FPEPV** (SEQ ID NO: 95); 4) **SGALTSGVHTFP** (SEQ ID NO: 96); 5) **QSSGLY** (SEQ ID NO: 97); 6) **VTV** (SEQ ID NO: 70); 7) **TQT** (SEQ ID NO: 98); 8) **HKP** (SEQ ID NO: 99); 9) **DK** (SEQ ID NO: 100); 10) **VAGPS** (SEQ ID NO: 101); 11) **FPPKP** (SEQ ID NO: 75); 12) **RTP** (SEQ ID NO: 102); 13) **DVSHEDPEV** (SEQ ID NO: 77); 14) **DGVEVHNAK** (SEQ ID NO: 80); 15) **FN** (SEQ ID NO: 103); 16) **VLTVV** (SEQ ID NO: 104); 17) **GKE** (SEQ ID NO: 84); 18) **NKGLPAP** (SEQ ID NO: 105); 19) **SKTKGQPRE** (SEQ ID NO: 106); 20) **PPS** (SEQ ID NO: 107); 21) **MTKNQ** (SEQ ID NO: 108); 22) **YPSDI** (SEQ ID NO: 89); 23) **NGQPENN** (SEQ ID NO: 90); 24) **TPPMLDSGGS** (SEQ ID NO: 109); 25) **GNVF** (SEQ ID NO: 110); and 26) **HEALHNHYTQKSLSLSPGK** (SEQ ID NO: 92), as shown in Figure 9B.

[00364] In some instances, a target immunoglobulin is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG3 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 13-22; 3) amino acids 33-37; 4) amino acids 43-61; 5) amino acid 71; 6) amino acids 78-80; 7) 87-91; 8) amino acids 97-106; 9) 111-115; 10) 147-167; 11) 173-177; 16) 185-187; 13) 195-203; 14) 210-218; 15) 226-227; 16) 238-239; 17) 246-248; 18) 255-261; 19) 267-275; 20) 282-291; 21) amino acids 303-307; 22) amino acids 313-320; 23) amino acids 324-333; 24) amino acids 350-352; 25) amino acids 359-365; and 26) amino acids 372-377; where the amino acid numbering is based on the numbering of the amino acid sequence set forth in SEQ ID NO: 9 (human IgG3; also depicted in Figure 9B).

[00365] Exemplary surface-accessible loop regions of an IgG3 heavy chain include 1) **ASTKGP** (SEQ ID NO: 64); 2) **PCSRSTSGGT** (SEQ ID NO: 111); 3) **FPEPV** (SEQ ID NO: 95); 4) **SGALTSGVHTFPAVLQSSG** (SEQ ID NO: 112); 5) **V** (SEQ ID NO: 113); 6) **TQT** (SEQ ID NO: 98); 7) **HKPSN** (SEQ ID NO: 73); 8) **RVELKTPLGD** (SEQ ID NO: 114); 9) **CPRCPKP** (SEQ ID NO: 115); 10) **PKSCDTPPPCPRCPAPELLGG** (SEQ ID NO: 116); 11) **FPPKP** (SEQ ID NO: 75); 12) **RTP** (SEQ ID NO: 102); 13) **DVSHEDPEV** (SEQ ID NO: 77); 14) **DGVEVHNAK** (SEQ ID NO: 80); 15) **YN** (SEQ ID NO: 117); 16) **VL** (SEQ ID NO: 118); 17) **GKE** (SEQ ID NO: 84); 18) **NKALPAP** (SEQ ID NO: 85); 19) **SKTKGQPRE** (SEQ ID NO: 119); 20) **PPSREEMTKN** (SEQ ID NO: 120); 21) **YPSDI** (SEQ ID NO: 89); 22) **SSGQPENN** (SEQ ID NO: 121); 23) **TPPMLDSDGS** (SEQ ID NO: 109); 24) **GNI** (SEQ ID NO: 122); 25) **HEALHNR** (SEQ ID NO: 123); and 26) **SLSPGK** (SEQ ID NO: 93), as shown in Figure 9B.

[00366] In some instances, a target immunoglobulin is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG4 heavy chain constant region corresponding to one or more of: 1) amino acids 1-5; 2) amino acids 12-23; 3) amino acids 32-36; 4) amino acids 42-53; 5) amino acids 57-62; 6) amino acids 68-70; 7) amino acids 77-79; 8) amino acids 86-88; 9) amino acids 94-95; 10) amino acids 101-102; 11) amino acids 108-118; 12) amino acids 122-126; 13) amino acids 134-136; 14) amino acids 144-152; 15) amino acids 159-167; 16) amino acids 175-176; 17)

amino acids 185-186; 18) amino acids 196-198; 19) amino acids 205-211; 20) amino acids 217-226; 21) amino acids 232-241; 22) amino acids 253-257; 23) amino acids 264-265; 24) 269-270; 25) amino acids 274-283; 26) amino acids 300-303; 27) amino acids 399-417; where the amino acid numbering is based on the numbering of the amino acid sequence set forth in SEQ ID NO: 10 (human IgG4; also depicted in Figure 9B).

[00367] Exemplary surface-accessible loop regions of an IgG4 heavy chain include 1) **STKGP** (SEQ ID NO: 124); 2) **PCSRSTSESTAA** (SEQ ID NO: 94); 3) **FPEPV** (SEQ ID NO: 95); 4) **SGALTSGVHTFP** (SEQ ID NO: 96); 5) **QSSGLY** (SEQ ID NO: 97); 6) **VTV** (SEQ ID NO: 70); 7) **TKT** (SEQ ID NO: 125); 8) **HKP** (SEQ ID NO: 99); 9) **DK** (SEQ ID NO: 100); 10) **YG** (SEQ ID NO: 126); 11) **CPAPEFLGGPS** (SEQ ID NO: 127); 12) **FPPKP** (SEQ ID NO: 75); 13) **RTP** (SEQ ID NO: 102); 14) **DVSQEDPEV** (SEQ ID NO: 128); 15) **DGVEVHNAK** (SEQ ID NO: 80); 16) **FN** (SEQ ID NO: 103); 17) **VL** (SEQ ID NO: 118); 18) **GKE** (SEQ ID NO: 84); 19) **NKGLPSS** (SEQ ID NO: 129); 20) **SKAKGQPREP** (SEQ ID NO: 130); 21) **PPSQEEMTKN** (SEQ ID NO: 131); 22) **YPSDI** (SEQ ID NO: 89); 23) **NG** (SEQ ID NO: 132); 24) **NN** (SEQ ID NO: 133); 25) **TPPVLDSDGS** (SEQ ID NO: 91); 26) **GNVF** (SEQ ID NO: 110); and 27) **HEALHNHYTQKSLSLSLGK** (SEQ ID NO: 134), as shown in Figure 9B.

[00368] In some instances, a target immunoglobulin is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgA heavy chain constant region corresponding to one or more of: 1) amino acids 1-13; 2) amino acids 17-21; 3) amino acids 28-32; 4) amino acids 44-54; 5) amino acids 60-66; 6) amino acids 73-76; 7) amino acids 80-82; 8) amino acids 90-91; 9) amino acids 123-125; 10) amino acids 130-133; 11) amino acids 138-142; 12) amino acids 151-158; 13) amino acids 165-174; 14) amino acids 181-184; 15) amino acids 192-195; 16) amino acid 199; 17) amino acids 209-210; 18) amino acids 222-245; 19) amino acids 252-256; 20) amino acids 266-276; 21) amino acids 293-294; 22) amino acids 301-304; 23) amino acids 317-320; 24) amino acids 329-353; where the amino acid numbering is based on the numbering of the amino acid sequence set forth in SEQ ID NO: 11 (human IgA; also depicted in Figure 9B).

[00369] Exemplary surface-accessible loop regions of an IgA heavy chain include 1) **ASPTSPKVFPLSL** (SEQ ID NO: 135); 2) **QPDGN** (SEQ ID NO: 136); 3) **VQGFFPQEPL** (SEQ ID NO: 137); 4) **SGQGVARNFP** (SEQ ID NO: 138); 5) **SGDLYTT** (SEQ ID NO:

139); 6) **PATQ** (SEQ ID NO: 140); 7) **GKS** (SEQ ID NO: 141); 8) **YT** (SEQ ID NO: 142); 9) **CHP** (SEQ ID NO: 143); 10) **HRPA** (SEQ ID NO: 144); 11) **LLGSE** (SEQ ID NO: 145); 12) **GLRDASGV** (SEQ ID NO: 146); 13) **SSGKSAVQGP** (SEQ ID NO: 147); 14) **GCYS** (SEQ ID NO: 148); 15) **CAEP** (SEQ ID NO: 149); 16) **PE** (SEQ ID NO: 150); 17) **SGNTFRPEVHLLPPPSEELALNEL** (SEQ ID NO: 151); 18) **ARGFS** (SEQ ID NO: 152); 19) **QGSQELPREKY** (SEQ ID NO: 153); 20) **AV** (SEQ ID NO: 154); 21) **AAED** (SEQ ID NO: 155); 22) **HEAL** (SEQ ID NO: 156); and 23) **IDRLAGKPTHVNVSVVMAEVDGTCY** (SEQ ID NO: 157), as shown in Figure 9B.

[00370] A sulfatase motif can be provided within or adjacent one or more of these amino acid sequences of such modification sites of an Ig heavy chain. For example, an Ig heavy chain polypeptide can be modified (e.g., where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions) at one or more of these amino acid sequences to provide a sulfatase motif adjacent and N-terminal and/or adjacent and C-terminal to these modification sites. Alternatively or in addition, an Ig heavy chain polypeptide can be modified (e.g., where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions) at one or more of these amino acid sequences to provide a sulfatase motif between any two residues of the Ig heavy chain modifications sites. In some embodiments, an Ig heavy chain polypeptide may be modified to include two motifs, which may be adjacent to one another, or which may be separated by one, two, three, four or more (e.g., from about 1 to about 25, from about 25 to about 50, or from about 50 to about 100, or more, amino acids). Alternatively or in addition, where a native amino acid sequence provides for one or more amino acid residues of a sulfatase motif sequence, selected amino acid residues of the modification sites of an Ig heavy chain polypeptide amino acid sequence can be modified (e.g., where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions) so as to provide a sulfatase motif at the modification site.

[00371] The amino acid sequence of a surface-accessible loop region can thus be modified to provide a sulfatase motif, where the modifications can include insertions, deletions, and/or substitutions. For example, where the modification is in a CH1 domain, the surface-accessible loop region can have the amino acid sequence **NSGALTSG** (SEQ ID NO: 67), and the aldehyde-tagged sequence can be, e.g., **NSGALCTPSRG** (SEQ ID NO: 158), e.g., where the “TS” residues of the **NSGALTSG** (SEQ ID NO: 67), sequence are replaced with “CTPSR,” ,

i.e., **NSGALCTPSRG** (SEQ ID NO: 159), such that the sulfatase motif has the sequence **LCTPSR** (SEQ ID NO: 32). As another example, where the modification is in a CH2 domain, the surface-accessible loop region can have the amino acid sequence **NKALPAP** (SEQ ID NO: 85), and the aldehyde-tagged sequence can be, e.g., **NLCTPSRAP** (SEQ ID NO: 160), e.g., where the “KAL” residues of the **NKALPAP** (SEQ ID NO: 85) sequence are replaced with “LCTPSR,” such that the sulfatase motif has the sequence **LCTPSR** (SEQ ID NO: 32). As another example, where the modification is in a CH2/CH3 domain, the surface-accessible loop region can have the amino acid sequence **KAKGQPR** (SEQ ID NO: 87), and the aldehyde-tagged sequence can be, e.g., **KAKGLCTPSR** (SEQ ID NO: 161), e.g., where the “GQP” residues of the **KAKGQPR** (SEQ ID NO: 87) sequence are replaced with “LCTPS,” such that the sulfatase motif has the sequence **LCTPSR** (SEQ ID NO: 32).

[00372] As noted above, an isolated aldehyde-tagged anti-CD22 Ig polypeptide can comprise a light chain constant region modified to include a sulfatase motif as described above, where the sulfatase motif is in or adjacent a surface-accessible loop region of the Ig polypeptide light chain constant region. Illustrative examples of surface-accessible loop regions of a light chain constant region are presented in Figures 9A and 9C.

[00373] In some instances, a target immunoglobulin is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an Ig light chain constant region corresponding to one or more of: 1) amino acids 130-135; 2) amino acids 141-143; 3) amino acid 150; 4) amino acids 162-166; 5) amino acids 163-166; 6) amino acids 173-180; 7) amino acids 186-194; 8) amino acids 211-212; 9) amino acids 220-225; 10) amino acids 233-236; wherein the amino acid numbering is based on the amino acid numbering of human kappa light chain as depicted in Figure 9C. In some instances, a target immunoglobulin is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an Ig light chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107;

where the amino acid numbering is based on SEQ ID NOs: 12 and 13 (human kappa light chain; amino acid sequences depicted in Figure 9C, i.e., seq1 and seq2, respectively).

[00374] Exemplary surface-accessible loop regions of an Ig light chain (e.g., a human kappa light chain) include: 1) **RTVAAP** (SEQ ID NO: 162); 2) **PPS** (SEQ ID NO: 107); 3) Gly (see, e.g., Gly at position 150 of the human kappa light chain sequence depicted in Figure 9C); 4) **YPREA** (SEQ ID NO: 163); 5) **PREA** (SEQ ID NO: 164); 6) **DNALQSGN** (SEQ ID NO: 165); 7) **TEQDSKDST** (SEQ ID NO: 166); 8) **HK** (SEQ ID NO: 167); 9) **HQGLSS** (SEQ ID NO: 168); and 10) **RGEC** (SEQ ID NO: 169), as shown in Figures 9A and 9C.

[00375] Exemplary surface-accessible loop regions of an Ig lambda light chain, e.g., seq3 (SEQ ID NO: 14) in Figure 9C, include **QPKAAP** (SEQ ID NO: 170), **PPS** (SEQ ID NO: 107), **NK** (SEQ ID NO: 171), **DFYPGAV** (SEQ ID NO: 172), **DSSPVKAG** (SEQ ID NO: 173), **TTP** (SEQ ID NO: 174), **SN** (SEQ ID NO: 175), **HKS** (SEQ ID NO: 176), **EG** (SEQ ID NO: 177), and **APTECS** (SEQ ID NO: 178), as shown in Figure 9C.

[00376] In some instances, a target immunoglobulin is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of a rat Ig light chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acids 121-22; 4) amino acids 31-37; 5) amino acids 44-51; 6) amino acids 55-57; 7) amino acids 61-62; 8) amino acids 81-83; 9) amino acids 91-92; 10) amino acids 102-105; wherein the amino acid numbering is based on the amino acid numbering of rat light chain as set forth in SEQ ID NO: 16 (seq5 depicted in Figure 9C).

[00377] In some cases, a sulfatase motif is introduced into the CH1 region of an anti-CD22 heavy chain constant region. In some cases, a sulfatase motif is introduced at or near (e.g., within 1 to 10 amino acids of) the C-terminus of an anti-CD22 heavy chain. In some cases, a sulfatase motif is introduced in the light-chain constant region.

[00378] In some cases, a sulfatase motif is introduced into the CH1 region of an anti-CD22 heavy chain constant region, e.g., within amino acids 121-219 of the IgG1 heavy chain amino acid sequence depicted in Figure 9A. For example, in some cases, a sulfatase motif is introduced into the amino acid sequence:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL
QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVE (SEQ ID NO: 179). For

example, in some of these embodiments, the amino acid sequence **GALTSGVH** (SEQ ID NO: 180) is modified to **GALCTPSRGVH** (SEQ ID NO: 181), where the sulfatase motif is **LCTPSR** (SEQ ID NO: 32).

[00379] In some cases, a sulfatase motif is introduced at or near the C-terminus of an anti-CD22 heavy chain, e.g., the sulfatase motifs introduced within 1 amino acid, 2 amino acids (aa), 3 aa, 4 aa, 5 aa, 6 aa, 7 aa, 8 aa, 9 aa, or 10 aa the C-terminus of an anti-CD22 heavy chain. As one non-limiting example, the C-terminal lysine residue of an anti-CD22 heavy chain can be replaced with the amino acid sequence **SLCTPSRGS** (SEQ ID NO: 182).

[00380] In some cases, a sulfatase motif is introduced into the constant region of a light chain of an anti-CD22 antibody. As one non-limiting example, in some cases, a sulfatase motif is introduced into the constant region of a light chain of an anti-CD22 antibody, where the sulfatase motif is C-terminal to **KVDNAL** (SEQ ID NO: 58), and/or is N-terminal to **QSGNSQ** (SEQ ID NO: 59). For example, in some cases, the sulfatase motif is **LCTPSR** (SEQ ID NO: 32), and the anti-CD22 light chain comprises the amino acid sequence **KVDNALLCTPSRQSGNSQ** (SEQ ID NO: 183).

Exemplary anti-CD22 antibodies

[00381] In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody comprising a heavy chain VH CDR selected from **IYDMS** (VH CDR1; SEQ ID NO: 17), **YISSGGGTTYYPDTVKG** (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAF** (VH CDR3; SEQ ID NO: 19). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21)

amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, i.e., seq1 or seq2, respectively).

[00382] In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody comprising a light-chain CDR selected from **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 122-127; 2) amino acids 137-143; 3) amino acids 155-158; 4) amino acids 163-170; 5) amino acids 163-183; 6) amino acids 179-183; 7) amino acids 190-192; 8) amino acids 200-202; 9) amino acids 199-202; 10) amino acids 208-212; 11) amino acids 220-241; 12) amino acids 247-251; 13) amino acids 257-261; 14) amino acid 269-277; 15) amino acids 271-277; 16) amino acids 284-285; 17) amino acids 284-292; 18) amino acids 289-291; 19) amino acids 299-303; 20) amino acids 309-313; 21) amino acids 320-322; 22) amino acids 329-335; 23) amino acids 341-349; 24) amino acids 342-348; 25) amino acids 356-365; 26) amino acids 377-381; 27) amino acids 388-394; 28) amino acids 398-407; 29) amino acids 433-451; and 30) amino acids 446-451; wherein the amino acid

numbering is based on the amino acid numbering of human IgG1 as depicted in Figure 9B. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, i.e., seq1 or seq2, respectively).

[00383] In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody comprising VH CDRs **IYDMS** (VH CDR1; SEQ ID NO: 17), **YISSGGGTTYYPDTVKG** (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAAY** (VH CDR3; SEQ ID NO: 19). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or

adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, seq1 or seq2, respectively).

[00384] In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody comprising VL CDRs **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where

the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, i.e., seq1 or seq2, respectively).

[00385] In some cases, a suitable anti-CD22 antibody competes for binding to a CD22 epitope (e.g., an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C) with an antibody that comprises VH CDRs **IYDMS** (VH CDR1; SEQ ID NO: 17), **YISSGGGTTYYPDTVKG** (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAAY** (VH CDR3; SEQ ID NO: 19) and VL CDRs **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, i.e., seq1 or seq2, respectively).

[00386] In some cases, a suitable anti-CD22 antibody comprises VH CDRs **IYDMS** (VH CDR1; SEQ ID NO: 17), **YISSGGGTTYYPDTVKG** (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAFAY** (VH CDR3; SEQ ID NO: 19). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B.) In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, i.e., seq1 or seq2, respectively).

[00387] In some cases, a suitable anti-CD22 antibody comprises VL CDRs **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent

to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, seq1 or seq2, respectively).

[00388] In some cases, a suitable anti-CD22 antibody comprises VH CDRs **IYDMS** (VH CDR1; SEQ ID NO: 17), **YISSGGGTTYYPDTVKG** (VH CDR2; SEQ ID NO: 18), and **HSGYGSSYGVLFAAY** (VH CDR3; SEQ ID NO: 19) and VL CDRs **RASQDISNYLN** (VL CDR1; SEQ ID NO: 20), **YTSILHS** (VL CDR2; SEQ ID NO: 21), and **QQGNTLPWT** (VL CDR3; SEQ ID NO: 22). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino

acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, seq1 or seq2, respectively).

[00389] In some cases, a suitable anti-CD22 antibody comprises VH CDRs present in an anti-CD22 VH region comprising the following amino acid sequence:

**EVQLVESGGGLVKPGGSLRLSCAASGFAFSIYDMSWVRQAPGKGLEWVAYISSGG
GTTYYPDTVKGRFTISRDNKNSLYLQMSSLRAEDTAMYYCARHSGYGSSYGVLF
AYWGQGTLVTVSS** (SEQ ID NO: 23). In some cases, the anti-CD22 antibody is humanized.

In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-

287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, i.e., seq1 or seq2, respectively).

[00390] In some cases, a suitable anti-CD22 antibody comprises VL CDRs present in an anti-CD22 VL region comprising the following amino acid sequence:

**DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPGKAVKLLIYYTSILHSG
VPSRFSGSGSGTDYTLTISSLQQEDFATYFCQQGNTLPWTFGGGGTKVEIKR** (SEQ ID

NO: 24). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions,

deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, seq1 or seq2, respectively).

[00391] In some cases, a suitable anti-CD22 antibody comprises VH CDRs present in **EVQLVESGGGLVKPGGSLRLSCAASGFAFSIYDMSWVRQAPGKGLEWVAYISSGG GTTYYPDTVKGRFTISRDNKNSLYLQMSSLRAEDTAMYYCARHSGYGSSYGVLF AYWGQGTLTVSS** (SEQ ID NO: 23) and VL CDRs present in **DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPGKAVKLLIYYTSILHSG VPSRFSGSGSGTDYTLTISSLQQEDFATYFCQQGNTLPWTFGGGGTKVEIKR** (SEQ ID NO: 24). In some cases, the anti-CD22 antibody is humanized. In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions; e.g., where the sulfatase motif is within, or adjacent to, a region of an Ig kappa constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 12-14; 3) amino acid 21; 4) amino acids 33-37; 5) amino acids 34-37; 6) amino acids 44-

51; 7) amino acids 57-65; 8) amino acids 83-83; 9) amino acids 91-96; 10) amino acids 104-107; where the amino acid numbering is based on SEQ ID NOs: 12 or 13 (human kappa light chains; amino acid sequences depicted in Figure 9C, seq1 or seq2, respectively).

[00392] In some cases, a suitable anti-CD22 antibody comprises the VH amino acid sequence

**EVQLVESGGGLVKPGGSLRLSCAASGFAFSIYDMSWVRQAPGKGLEWVAYISSGG
GTTYYPDTVKGRTISRDNKNSLYLQMSSLRAEDTAMYYCARHSGYGSSYGVLF
AYWGQGTLVTVSS** (SEQ ID NO: 23). In some cases, a suitable anti-CD22 antibody

comprises the VL amino acid sequence

**DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPGKAVKLLIYYTSILHSG
VPSRFGSGSGTDYTLTISSLQQEDFATYFCQQGNTLPWTFGGGGTKVEIKR** (SEQ ID

NO: 24). In some cases, a suitable anti-CD22 antibody comprises the VH amino acid sequence


**EVQLVESGGGLVKPGGSLRLSCAASGFAFSIYDMSWVRQAPGKGLEWVAYISSGG
GTTYYPDTVKGRTISRDNKNSLYLQMSSLRAEDTAMYYCARHSGYGSSYGVLF
AYWGQGTLVTVSS** (SEQ ID NO: 23); and the VL amino acid sequence

**DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPGKAVKLLIYYTSILHSG
VPSRFGSGSGTDYTLTISSLQQEDFATYFCQQGNTLPWTFGGGGTKVEIKR** (SEQ ID

NO: 24). In some instances, the anti-CD22 antibody is modified to include a sulfatase motif as described above, where the modification includes one or more amino acid residue insertions, deletions, and/or substitutions. In certain embodiments, the sulfatase motif is within, or adjacent to, a region of an IgG1 heavy chain constant region corresponding to one or more of: 1) amino acids 1-6; 2) amino acids 16-22; 3) amino acids 34-47; 4) amino acids 42-49; 5) amino acids 42-62; 6) amino acids 34-37; 7) amino acids 69-71; 8) amino acids 79-81; 9) amino acids 78-81; 10) amino acids 87-91; 11) amino acids 100-121; 12) amino acids 127-131; 13) amino acids 137-141; 14) amino acid 149-157; 15) amino acids 151-157; 16) amino acids 164-165; 17) amino acids 164-172; 18) amino acids 169-171; 19) amino acids 179-183; 20) amino acids 189-193; 21) amino acids 200-202; 22) amino acids 209-215; 23) amino acids 221-229; 24) amino acids 22-228; 25) amino acids 236-245; 26) amino acids 217-261; 27) amino acids 268-274; 28) amino acids 278-287; 29) amino acids 313-331; and 30) amino acids 324-331; wherein the amino acid numbering is based on the amino acid numbering of human IgG1 as set out in SEQ ID NO: 7 (human IgG1 constant region depicted in Figure 9B).

Drugs for Conjugation to a Polypeptide

[00393] The present disclosure provides drug-polypeptide conjugates. Examples of drugs include small molecule drugs, such as a cancer anti-cancer agent. For example, where the polypeptide is an antibody (or fragment thereof) that has specificity for a tumor cell, the antibody can be modified as described herein to include a modified amino acid, which can be subsequently conjugated to a cancer anti-cancer agent, such as a microtubule affecting agents. In certain embodiments, the drug is a microtubule affecting agent that has antiproliferative activity, such as a maytansinoid. In certain embodiments, the drug is a maytansinoid, which has the following structure:



where ~~~ indicates the point of attachment between the maytansinoid and the linker, L, in formula (I). By “point of attachment” is meant that the ~~~ symbol indicates the bond between the N of the maytansinoid and the linker, L, in formula (I). For example, in formula (I), W^1 is a maytansinoid, such as a maytansinoid of the structure above, where ~~~ indicates the point of attachment between the maytansinoid and the linker, L.

[00394] As described above, in certain embodiments, L is a linker described by the formula $-(L^1)_a-(L^2)_b-(L^3)_c-(L^4)_d-$, wherein L^1 , L^2 , L^3 and L^4 are each independently a linker unit. In certain embodiments, L^1 is attached to the coupling moiety, such as a hydrazinyl-indolyl or a hydrazinyl-pyrrolo-pyridinyl coupling moiety (e.g., as shown in formula (I) above). In certain embodiments, L^2 , if present, is attached to W^1 (the maytansinoid). In certain embodiments, L^3 , if present, is attached to W^1 (the maytansinoid). In certain embodiments, L^4 , if present, is attached to W^1 (the maytansinoid).

[00395] As described above, in certain embodiments, the linker $-(L^1)_a-(L^2)_b-(L^3)_c-(L^4)_d-$ is described by the formula $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d-$, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4. In certain embodiments, as

described above, L^1 is attached to the hydrazinyl-indolyl or the hydrazinyl-pyrrolo-pyridinyl coupling moiety (e.g., as shown in formula (I) above). As such, in certain embodiments, T^1 is attached to the hydrazinyl-indolyl or the hydrazinyl-pyrrolo-pyridinyl coupling moiety (e.g., as shown in formula (I) above). In certain embodiments, V^1 is attached to W^1 (the maytansinoid). In certain embodiments, as described above, L^2 , if present, is attached to W^1 (the maytansinoid). As such, in certain embodiments, T^2 , if present, is attached to W^1 (the maytansinoid), or V^2 , if present, is attached to W^1 (the maytansinoid). In certain embodiments, as described above, L^3 , if present, is attached to W^1 (the maytansinoid). As such, in certain embodiments, T^3 , if present, is attached to W^1 (the maytansinoid), or V^3 , if present, is attached to W^1 (the maytansinoid). In certain embodiments, as described above, L^4 , if present, is attached to W^1 (the maytansinoid). As such, in certain embodiments, T^4 , if present, is attached to W^1 (the maytansinoid), or V^4 , if present, is attached to W^1 (the maytansinoid).

[00396] Embodiments of the present disclosure include conjugates where a polypeptide (e.g., anti-CD22 antibody) is conjugated to one or more drug moieties (e.g., maytansinoid), such as 2 drug moieties, 3 drug moieties, 4 drug moieties, 5 drug moieties, 6 drug moieties, 7 drug moieties, 8 drug moieties, 9 drug moieties, or 10 or more drug moieties. The drug moieties may be conjugated to the polypeptide at one or more sites in the polypeptide, as described herein. In certain embodiments, the conjugates have an average drug-to-antibody ratio (DAR) (molar ratio) in the range of from 0.1 to 10, or from 0.5 to 10, or from 1 to 10, such as from 1 to 9, or from 1 to 8, or from 1 to 7, or from 1 to 6, or from 1 to 5, or from 1 to 4, or from 1 to 3, or from 1 to 2. In certain embodiments, the conjugates have an average DAR from 1 to 2, such as 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9 or 2. In certain embodiments, the conjugates have an average DAR of 1.6 to 1.9. In certain embodiments, the conjugates have an average DAR of 1.7. By average is meant the arithmetic mean.

ANTI-CANCER AGENTS

[00397] The anti-cancer agents of the present disclosure include any agent for treating a cancer, e.g., chemotherapeutic agents and/or biologic therapeutic agents. In some embodiments, the cancer to be treated by the methods disclosed (e.g., by treatment with an ADC as disclosed herein) are resistant to an anti-cancer agent (e.g., an anti-cancer agent set forth below). In some embodiments, the anti-cancer agents set forth below are used in combination with an ADC as

disclosed herein for the treatment of a cancer, including cancers that are resistant to the anti-cancer agents set forth below.

[00398] An anti-cancer agent may include an alkylating agent, e.g., DNA alkylating agent. For example, an alkylating agent may include bendamustine chlorambucil, carmustine, cyclophosphamide, mechlorethamine, or the like, and pharmaceutically acceptable salts and formulations thereof. An anti-cancer agent may include a DNA or RNA synthesis inhibitors, e.g., topoisomerase inhibitors, polymerase inhibitors, dihydrofolate reductase inhibitors, or the like. For example, a DNA or RNA synthesis inhibitor may include nelarabine, bleomycin, cyarabine, doxorubicin, pralatrexate, methotrexate, or the like, and pharmaceutically acceptable salts and formulations thereof. An anti-cancer agent may include a kinase inhibitor, for example, a Bruton's tyrosine kinase (BTK) inhibitor, e.g., acalabrutinib, ibrutinib, or the like, and pharmaceutically acceptable salts and formulations thereof. An anti-cancer agent may include a phosphoinositide-3-kinase (PI3K) inhibitor or an inhibitor of an isoform thereof, e.g., P110 δ inhibitor. For example, a PI3K inhibitor may include copanlisib, idelalisib, or the like, and pharmaceutically acceptable salts and formulations thereof. An anti-cancer agent may include chemokine inhibitor, for example, a chemokine CXCR4 receptor inhibitor. The chemokine inhibitor may include plerixafor, or the like, and pharmaceutically acceptable salts and formulations thereof. An anti-cancer agent may include histone deacetylase inhibitor, e.g., vorinostat, romidespsin, belinostat, or the like, and pharmaceutically acceptable salts and formulations thereof. The anti-cancer agent may include a proteasome inhibitor, such as bortezomib. The anti-cancer agent may include a corticosteroid, such as dexamethasone, prednisone, or the like. The anti-cancer agent may include immunosuppressive agents or antineoplastic agents. For example, the anti-cancer agent may include an interleukin-2 inhibitor, e.g., denileukin difitox, or the like. For example, the anti-cancer agent may include recombinant interferon alfa 2b. The anti-cancer agent may include a tubulin inhibitor, e.g., vincristine, vinblastine, or the like, and pharmaceutically acceptable salts and formulations thereof. The anti-cancer agent may include a monoclonal antibody or drug conjugates thereof, e.g., small molecule drug conjugates, radioactive drug conjugates, or the like. For example the anti-cancer agent may include antibodies of CD19, CD20, CD22, CD30, or the like. Examples of antibody-based anti-cancer agents include brentuximab vedotin, tositumomab, iodine-131 tositumomab, ibritumomab, tiuxetan, obinutuzumab, rituximab, rituximab and hyaluronidase human, or the

like. The anti-cancer agent may include ubiquitin E3 ligase inhibitor, e.g., lenalidomide, or the like. The anti-cancer agent may include adoptive cell transfer therapy, for example, with axicabtagene ciloleucel.

[00399] The present invention may include therapies (e.g., combination therapies) involving more than one anti-cancer agent (e.g., in combination with an ADC as set forth herein). In one embodiment, an anti-cancer agent includes cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“CHOP”). In one embodiment, an anti-cancer agent includes cyclophosphamide, vincristine sulfate, procarbazine hydrochloride, and prednisone (“COPP”). In one embodiment, an anti-cancer agent includes cyclophosphamide, vincristine sulfate, and prednisone (“CVP”). In one embodiment, an anti-cancer agent includes etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”). In one embodiment, an anti-cancer agent includes cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”). In one embodiment, an anti-cancer agent includes ifosfamide, carboplatin, and etoposide phosphate (“ICE”). In one embodiment, an anti-cancer agent includes rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”). In one embodiment, an anti-cancer agent includes rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”). In one embodiment, an anti-cancer agent includes rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”). In one embodiment, an anti-cancer agent includes rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

[00400] In other embodiments, an anti-cancer agent may include R-CHOP, R-CVP, R-EPOCH, or R-ICE, wherein rituximab is replaced with a different anti-CD20 antibody, e.g., ofatumumab, obinutuzumab, ocrelizumab, or the like. For example, R-CHOP with rituximab replaced with obinutuzumab may include obinutuzumab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone. For example, R-CVP with rituximab replaced with obinutuzumab, may include obinutuzumab, cyclophosphamide, vincristine sulfate, and prednisone. For example, R-EPOCH with rituximab replaced with ocrelizumab, may include ocrelizumab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride.

[00401] In one embodiment, an anti-cancer agent includes R-CHOP.

[00402] In one embodiment, an anti-cancer agent includes rituximab.

[00403] In one embodiment an anti-cancer agent includes bendamustine.

FORMULATIONS

[00404] The conjugates (including antibody conjugates) of the present disclosure can be formulated in a variety of different ways. In general, where the conjugate is a polypeptide-drug conjugate, the conjugate is formulated in a manner compatible with the drug conjugated to the polypeptide, the condition to be treated, and the route of administration to be used.

[00405] The conjugate (e.g., polypeptide-drug conjugate) can be provided in any suitable form, e.g., in the form of a pharmaceutically acceptable salt, and can be formulated for any suitable route of administration, e.g., oral, topical or parenteral administration. Where the conjugate is provided as a liquid injectable (such as in those embodiments where they are administered intravenously or directly into a tissue), the conjugate can be provided as a ready-to-use dosage form, or as a reconstitutable storage-stable powder or liquid composed of pharmaceutically acceptable carriers and excipients.

[00406] The anti-cancer agents of the present disclosure can be formulated in a variety of different ways. In general, the anti-cancer agent is formulated in a manner compatible with the condition to be treated and the route of administration to be used.

[00407] The anti-cancer agents can be provided in any suitable form, e.g., in the form of a pharmaceutically acceptable salt, and can be formulated for any suitable route of administration, e.g., oral, topical, or parenteral administration.

[00408] Methods for formulating conjugates and/or anti-cancer agents can be adapted from those readily available. For example, conjugates and/or anti-cancer agents can be provided in a pharmaceutical composition comprising a therapeutically effective amount of a conjugate and a pharmaceutically acceptable carrier (e.g., saline). The pharmaceutical composition may optionally include other additives (e.g., buffers, stabilizers, preservatives, and the like). In some embodiments, the formulations are suitable for administration to a mammal, such as those that are suitable for administration to a human.

[00409] In certain embodiments, where a condition requires combination therapy, i.e., treatment with a conjugate of the present invention and one or more anti-cancer agent, the conjugate and one or more anti-cancer agent may be formulated together for co-administration, or may be formulated separately for subsequent administration. Likewise, when more than one

anti-cancer agent is employed, one or more of the anti-cancer agents may be formulated together or may be formulated separately.

METHODS OF TREATMENT

[00410] The polypeptide-drug conjugates of the present disclosure find use in treatment of a condition or disease in a subject that is amenable to treatment by administration of the parent drug (i.e., the drug prior to conjugation to the polypeptide). By “treatment” is meant that at least an amelioration of the symptoms associated with the condition afflicting the host is achieved, where amelioration is used in a broad sense to refer to at least a reduction in the magnitude of a parameter, e.g. symptom, associated with the condition being treated. As such, treatment also includes situations where the pathological condition, or at least symptoms associated therewith, are completely inhibited, e.g., prevented from happening, or stopped, e.g. terminated, such that the host no longer suffers from the condition, or at least the symptoms that characterize the condition. Thus treatment includes: (i) prevention, that is, reducing the risk of development of clinical symptoms, including causing the clinical symptoms not to develop, e.g., preventing disease progression to a harmful state; (ii) inhibition, that is, arresting the development or further development of clinical symptoms, e.g., mitigating or completely inhibiting an active disease; and/or (iii) relief, that is, causing the regression of clinical symptoms.

[00411] In the context of cancer, the term “treating” includes any or all of: reducing growth of a solid tumor, inhibiting replication of cancer cells, reducing overall tumor burden, and ameliorating one or more symptoms associated with a cancer.

[00412] The subject to be treated can be one that is in need of therapy, where the host to be treated is one amenable to treatment using the parent drug. Accordingly, a variety of subjects may be amenable to treatment using the polypeptide-drug conjugates disclosed herein. Generally, such subjects are “mammals”, with humans being of interest. Other subjects can include domestic pets (e.g., dogs and cats), livestock (e.g., cows, pigs, goats, horses, and the like), rodents (e.g., mice, guinea pigs, and rats, e.g., as in animal models of disease), as well as non-human primates (e.g., chimpanzees, and monkeys).

[00413] The amount of polypeptide-drug conjugate administered can be initially determined based on guidance of a dose and/or dosage regimen of the parent drug. In general, the polypeptide-drug conjugates can provide for targeted delivery and/or enhanced serum half-life of

the bound drug, thus providing for at least one of reduced dose or reduced administrations in a dosage regimen. Thus, the polypeptide-drug conjugates can provide for reduced dose and/or reduced administration in a dosage regimen relative to the parent drug prior to being conjugated in an polypeptide-drug conjugate of the present disclosure.

[00414] Furthermore, as noted above, because the polypeptide-drug conjugates can provide for controlled stoichiometry of drug delivery, dosages of polypeptide-drug conjugates can be calculated based on the number of drug molecules provided on a per polypeptide-drug conjugate basis.

[00415] In some embodiments, multiple doses of a polypeptide-drug conjugate are administered. The frequency of administration of a polypeptide-drug conjugate can vary depending on any of a variety of factors, e.g., severity of the symptoms, condition of the subject, etc. For example, in some embodiments, a polypeptide-drug conjugate is administered once per month, twice per month, three times per month, every other week, once per week (qwk), twice per week, three times per week, four times per week, five times per week, six times per week, every other day, daily (qd/od), twice a day (bds/bid), or three times a day (tds/tid), etc.

[00416] The polypeptide-drug conjugates of the present disclosure find use in combination treatment with anti-cancer agents for conditions or diseases in a subject that are amenable to treatment by administration of the parent drug, e.g., maytansine, and/or amenable to treatment by, or formerly amenable to treatment by, administration of the anti-cancer agent. The combination treatment may have a synergistic treatment effect on the condition or disease. The combination treatment may serve to render the condition or disease more susceptible to treatment. For example, a resistant cancer, e.g., a cancer resistant to treatment by one or more anti-cancer agents, may become responsive to a combination therapy described herein.

[00417] In several embodiments, a dosage amount in mg/kg of a polypeptide-drug conjugate administered to a subject may include one or more of: 0.10, 0.25, 0.50, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, and 20, or a range between any of the preceding numbers, e.g., between about 3 and about 5, between about 5 and about 10, between about 9 and 10, or the like. The polypeptide-drug conjugate may be administered at one of the preceding dose values at a rate of once every week (qw), once every two weeks (q2w), once every three weeks (q3w), or once every month (qm). The polypeptide-drug conjugate may be administered at one of the preceding dose values at a rate of once every week (qw), once every two weeks (q2w), once

every three weeks (q3w), or once every month (qm), for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 weeks. In certain embodiments, the polypeptide-drug conjugate is administered once every three weeks (e.g., for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 weeks). Alternatively, the polypeptide-drug conjugate may be administered in a single dose at any of the preceding dose values. The polypeptide-drug conjugate may be administered at another preceding dose value at one of the preceding rates for one of the preceding periods. For example the polypeptide-drug conjugate may be administered at 10 mg/kg, once every three weeks, for a period of six weeks. Further, for example, the polypeptide-drug conjugate may later be administered at 3 mg/kg, once every week, for a period of 4 weeks, and so forth.

[00418] In several embodiments, one or more anti-cancer agents may be administered to a subject in a single or multiple dose. The frequency of administration of one or more anti-cancer agents can vary depending on any of a variety of factors, e.g., severity of the symptoms, condition of the subject, etc. For example, in some embodiments, one or more anti-cancer agents is administered once per month, twice per month, three times per month, every other week, once per week (qwk), once every two weeks (q2wk), once every three weeks (q3wk), twice per week, three times per week, four times per week, five times per week, six times per week, every other day, daily (qd/od), twice a day (bds/bid), or three times a day (tds/tid), etc. An anti-cancer agent may be administered at a dose value, at a dose rate, and for a period of time that has been approved for a particular anti-cancer agent by the U.S. Food and Drug Administration.

[00419] In certain embodiments, a peptide-drug conjugate and one or more anti-cancer agents is co-administered.

[00420] In some embodiments, a peptide-drug conjugate is administered prior to administering one or more anti-cancer agent. For example, the peptide-drug conjugate is administered as described herein, followed by administration of one or more anti-cancer agents as described herein. In certain embodiments, a period of time separates administration of the peptide-drug conjugate and the administration of one or more anti-cancer agents.

[00421] In some embodiments, a peptide-drug conjugate is administered for a period of time and later the peptide-drug conjugate is co-administered with one or more anti-cancer agents.

[00422] In some embodiments, one or more anti-cancer agent is administered prior to administering a peptide-drug conjugate. For example, one or more anti-cancer agent is

administered as described herein, followed by administration of the peptide-drug conjugate as described herein.

[00423] In some embodiments, one or more anti-cancer agents is administered for a period of time and later one or more anti-cancer agents is co-administered with a peptide-drug conjugate.

[00424] In certain embodiments, a cancer becomes resistant to administration with one or more anti-cancer agents. Administering a peptide-drug conjugate to a subject having a resistant cancer may treat the cancer and/or sensitize the cancer to further treatment with one or more anti-cancer agents.

Methods of treating cancer

[00425] The present disclosure provides methods for delivering one or more cancer anti-cancer agents and a peptide-drug conjugate to an individual having a cancer or a cancer described herein that has become resistant to one or more anti-cancer agents, e.g., R-CHOP. The methods are useful for treating a wide variety of cancers, including carcinomas, sarcomas, leukemias, and lymphomas. The methods are further useful for sensitizing a cancer described herein, i.e., relieving the resistance of a cancer toward one or more anti-cancer agents described herein, e.g., R-CHOP. In some embodiments, the peptide-drug conjugate can be administered to a cancer in the absence of other therapies (e.g., as a monotherapy). In some embodiments, the peptide-drug conjugate can be administered to a cancer in combination with other cancer therapies (e.g., in combination with R-CHOP).

[00426] The present disclosure provides methods for delivering one or more cancer anti-cancer agents and a peptide-drug conjugate to an individual having a cancer. The methods are useful for treating cancers associated with dysregulation of BCR signaling owing to B-cell overexpression and/or dysfunction. The methods are useful for treating cancers that are responsive to B-cell depletion therapies. The methods are also useful for treating cancers associated with dysregulation of BCR signaling that have become resistant to one or more anti-cancer agents.

[00427] Carcinomas that can be treated using a subject method include, but are not limited to, esophageal carcinoma, hepatocellular carcinoma, basal cell carcinoma (a form of skin cancer), squamous cell carcinoma (various tissues), bladder carcinoma, including transitional cell carcinoma (a malignant neoplasm of the bladder), bronchogenic carcinoma, colon carcinoma,

colorectal carcinoma, gastric carcinoma, lung carcinoma, including small cell carcinoma and non-small cell carcinoma of the lung, adrenocortical carcinoma, thyroid carcinoma, pancreatic carcinoma, breast carcinoma, ovarian carcinoma, prostate carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinoma, cystadenocarcinoma, medullary carcinoma, renal cell carcinoma, ductal carcinoma in situ or bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical carcinoma, uterine carcinoma, testicular carcinoma, osteogenic carcinoma, epithelial carcinoma, and nasopharyngeal carcinoma, etc.


[00428] Sarcomas that can be treated using a subject method include, but are not limited to, fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, chordoma, osteogenic sarcoma, osteosarcoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's sarcoma, leiomyosarcoma, rhabdomyosarcoma, and other soft tissue sarcomas.

[00429] Other solid tumors that can be treated using a subject method include, but are not limited to, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.


[00430] Leukemias that can be treated using a subject method include, but are not limited to, a) chronic myeloproliferative syndromes (neoplastic disorders of multipotential hematopoietic stem cells); b) acute myelogenous leukemias (neoplastic transformation of a multipotential hematopoietic stem cell or a hematopoietic cell of restricted lineage potential; c) chronic lymphocytic leukemias (CLL; clonal proliferation of immunologically immature and functionally incompetent small lymphocytes), including B-cell CLL, T-cell CLL prolymphocytic leukemia, small lymphocytic leukemia (SLL), and hairy cell leukemia; and d) acute lymphoblastic leukemias (characterized by accumulation of lymphoblasts).

[00431] Lymphomas that can be treated using a subject method include, but are not limited to, B-cell lymphomas (e.g., Burkitt's lymphoma, diffuse large B-cell lymphoma); Hodgkin's lymphoma; non-Hodgkin's B cell lymphoma (e.g., marginal zone lymphomas (MZL), mantle cell lymphoma (MCL), follicular lymphoma, primary central nervous system lymphoma); and the like.

[00432] In some embodiments, the present disclosure provides for the treatment of non-Hodgkin's lymphoma with an ADC as set forth herein. The method can comprise administering the ADC wherein the non-Hodgkin's lymphoma is resistant to R-CHOP (e.g., following R-CHOP escape). In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg. In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg.

[00433] In some embodiments, the present disclosure provides for the treatment of non-Hodgkin's lymphoma with an ADC as set forth herein in combination with R-CHOP. The method can comprise administering the ADC in combination with R-CHOP wherein the non-

Hodgkin's lymphoma is resistant to R-CHOP (e.g., following R-CHOP escape). In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg. In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg.

[00434] In some embodiments, the present disclosure provides for the treatment of diffuse large B cell lymphoma with an ADC as set forth herein. The method can comprise administering the ADC wherein the diffuse large B cell lymphoma is resistant to R-CHOP (e.g., following R-CHOP escape). In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg. In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg.

[00435] In some embodiments, the present disclosure provides for the treatment of diffuse large B cell lymphoma with an ADC as set forth herein in combination with R-CHOP. The method can comprise administering the ADC in combination with R-CHOP wherein the diffuse large B cell lymphoma is resistant to R-CHOP (e.g., following R-CHOP escape). In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg. In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg.

[00436] In some embodiments, the present disclosure provides for the treatment of follicular lymphoma with an ADC as set forth herein. The method can comprise administering the ADC wherein the follicular lymphoma is resistant to R-CHOP (e.g., following R-CHOP escape). In some embodiments, the ADC is:





and is administered at a dose of about 10 mg/kg. In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg.


[00437] In some embodiments, the present disclosure provides for the treatment of follicular lymphoma with an ADC as set forth herein in combination with R-CHOP. The method can comprise administering the ADC in combination with R-CHOP wherein the follicular lymphoma is resistant to R-CHOP (e.g., following R-CHOP escape). In some embodiments, the ADC is:



123




and is administered at a dose of about 10 mg/kg. In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg.

[00439] In some embodiments, the present disclosure provides for the treatment of mantle cell lymphoma with an ADC as set forth herein in combination with R-CHOP. The method can comprise administering the ADC in combination with R-CHOP wherein the mantle cell lymphoma is resistant to R-CHOP (e.g., following R-CHOP escape). In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg. In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg.

[00440] In some embodiments, the present disclosure provides for the treatment of marginal zone lymphoma with an ADC as set forth herein. The method can comprise administering the ADC wherein the marginal zone lymphoma is resistant to R-CHOP (e.g., following R-CHOP escape). In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg. In some embodiments, the ADC is:




and is administered at a dose of about 10 mg/kg.

[00441] In some embodiments, the present disclosure provides for the treatment of marginal zone lymphoma with an ADC as set forth herein in combination with R-CHOP. The method can comprise administering the ADC in combination with R-CHOP wherein the marginal zone lymphoma is resistant to R-CHOP (e.g., following R-CHOP escape). In some embodiments, the ADC is:



and is administered at a dose of about 10 mg/kg. In some embodiments, the ADC is:



and is administered at a dose of about 10 mg/kg.

EXAMPLES

[00442] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts,

temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Celsius, and pressure is at or near atmospheric. By “average” is meant the arithmetic mean. Standard abbreviations may be used, e.g., bp, base pair(s); kb, kilobase(s); pl, picoliter(s); s or sec, second(s); min, minute(s); h or hr, hour(s); aa, amino acid(s); kb, kilobase(s); bp, base pair(s); nt, nucleotide(s); i.m., intramuscular(ly); i.p., intraperitoneal(ly); s.c., subcutaneous(ly); and the like.

General Synthetic Procedures

[00443] Many general references providing commonly known chemical synthetic schemes and conditions useful for synthesizing the disclosed compounds are available (see, e.g., Smith and March, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, Fifth Edition, Wiley-Interscience, 2001; or Vogel, *A Textbook of Practical Organic Chemistry, Including Qualitative Organic Analysis*, Fourth Edition, New York: Longman, 1978).

[00444] Compounds as described herein can be purified by any purification protocol known in the art, including chromatography, such as HPLC, preparative thin layer chromatography, flash column chromatography and ion exchange chromatography. Any suitable stationary phase can be used, including normal and reversed phases as well as ionic resins. In certain embodiments, the disclosed compounds are purified via silica gel and/or alumina chromatography. See, e.g., *Introduction to Modern Liquid Chromatography*, 2nd Edition, ed. L. R. Snyder and J. J. Kirkland, John Wiley and Sons, 1979; and *Thin Layer Chromatography*, ed E. Stahl, Springer-Verlag, New York, 1969.

[00445] During any of the processes for preparation of the subject compounds, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups as described in standard works, such as J. F. W. McOmie, “Protective Groups in Organic Chemistry”, Plenum Press, London and New York 1973, in T. W. Greene and P. G. M. Wuts, “Protective Groups in Organic Synthesis”, Third edition, Wiley, New York 1999, in “The Peptides”, Volume 3 (editors: E. Gross and J. Meienhofer), Academic Press, London and New York 1981, in “Methoden der organischen Chemie”, Houben-Weyl, 4th edition, Vol. 15/1, Georg Thieme Verlag, Stuttgart 1974, in H.-D. Jakubke and H. Jescheit, “Aminosäuren, Peptide, Proteine”,


Verlag Chemie, Weinheim, Deerfield Beach, and Basel 1982, and/or in Jochen Lehmann, “Chemie der Kohlenhydrate: Monosaccharide and Derivate”, Georg Thieme Verlag, Stuttgart 1974. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

[00446] The subject compounds can be synthesized via a variety of different synthetic routes using commercially available starting materials and/or starting materials prepared by conventional synthetic methods. A variety of examples of synthetic routes that can be used to synthesize the compounds disclosed herein are described in the schemes below.

EXAMPLE 1

[00447] A linker containing a 4-amino-piperidine (4AP) group was synthesized according to Scheme 1, shown below.

Scheme 1

Synthesis of (9H-fluoren-9-yl)methyl 4-oxopiperidine-1-carboxylate (**200**)

[00448] To a 100 mL round-bottom flask containing a magnetic stir bar was added piperidin-4-one hydrochloride monohydrate (1.53 g, 10 mmol), Fmoc chloride (2.58 g, 10 mmol), sodium carbonate (3.18 g, 30 mmol), dioxane (20 mL), and water (2 mL). The reaction mixture was stirred at room temperature for 1 h. The mixture was diluted with EtOAc (100 mL) and extracted with water (1 x 100 mL). The organic layer was dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The resulting material was dried in vacuo to yield compound **200** as a white solid (3.05 g, 95% yield).

[00449] ^1H NMR (CDCl_3) δ 7.78 (d, 2H, $J = 7.6$), 7.59 (d, 2H, $J = 7.2$), 7.43 (t, 2H, $J = 7.2$), 7.37 (t, 2H, $J = 7.2$), 4.60 (d, 2H, $J = 6.0$), 4.28 (t, 2H, $J = 6.0$), 3.72 (br, 2H), 3.63 (br, 2H), 2.39 (br, 2H), 2.28 (br, 2H).

[00450] MS (ESI) m/z: $[M+H]^+$ Calcd for $C_{20}H_{20}NO_3$ 322.4; Found 322.2.

Synthesis of (9H-fluoren-9-yl)methyl 4-((2-(2-(3-(tert-butoxy)-3-oxopropoxy)ethoxy)ethyl)amino)piperidine-1-carboxylate (201)

[00451] To a dried scintillation vial containing a magnetic stir bar was added piperidinone **200** (642 mg, 2.0 mmol), $H_2N-PEG_2-CO_2t-Bu$ (560 mg, 2.4 mmol), 4 Å molecular sieves (activated powder, 500 mg), and 1,2-dichloroethane (5 mL). The mixture was stirred for 1 h at room temperature. To the reaction mixture was added sodium triacetoxyborohydride (845 mg, 4.0 mmol). The mixture was stirred for 5 days at room temperature. The resulting mixture was diluted with EtOAc. The organic layer was washed with saturated $NaHCO_3$ (1 x 50 mL), and brine (1 x 50 mL), dried over Na_2SO_4 , filtered, and concentrated under reduced pressure to yield compound **201** as an oil, which was carried forward without further purification.

Synthesis of 13-(1-(((9H-fluoren-9-yl)methoxy)carbonyl)piperidin-4-yl)-2,2-dimethyl-4,14-dioxo-3,7,10-trioxa-13-azaheptadecan-17-oic acid (202)

[00452] To a dried scintillation vial containing a magnetic stir bar was added *N*-Fmoc-piperidine-4-amino- PEG_2-CO_2t-Bu (**201**) from the previous step, succinic anhydride (270 mg, 2.7 mmol), and dichloromethane (5 mL). The mixture was stirred for 18 hours at room temperature. The reaction mixture was partitioned between EtOAc and saturated $NaHCO_3$. The aqueous layer was extracted with EtOAc (3x). The aqueous layer was acidified with HCl (1 M) until the pH ~3. The aqueous layer was extracted (3x) with DCM. The combined organic layers were dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The reaction mixture was purified by C18 flash chromatography (elute 10-100% MeCN/water with 0.1% acetic acid). Product-containing fractions were concentrated under reduced pressure and then azeotroped with toluene (3 x 50 mL) to remove residual acetic acid to afford 534 mg (42%, 2 steps) of compound **202** as a white solid.

[00453] ^1H NMR ($DMSO-d_6$) δ 11.96 (br, 1H), 7.89 (d, 2H, $J = 7.2$), 7.63 (d, 2H, $J = 7.2$), 7.42 (t, 2H, $J = 7.2$), 7.34 (t, 2H, $J = 7.2$), 4.25-4.55 (m, 3H), 3.70-4.35 (m, 3H), 3.59 (t, 2H, $J = 6.0$), 3.39 (m, 5H), 3.35 (m, 3H), 3.21 (br, 1H), 2.79 (br, 2H), 2.57 (m, 2H), 2.42 (q, 4H, $J = 6.0$), 1.49 (br, 3H), 1.37 (s, 9H).

[00454] MS (ESI) m/z: $[M+H]^+$ Calcd for $C_{35}H_{47}N_2O_9$ 639.3; Found 639.2.

Synthesis of (2*S*)-1-(((1⁴*S*,1⁶*S*,3³*S*,2*R*,4*S*,10*E*,12*E*,14*R*)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl)oxy)-2,3-dimethyl-1,4,7-trioxo-8-(piperidin-4-yl)-11,14-dioxa-3,8-diazaheptadecan-17-oic acid (203)

[00455] To a solution of ester **202** (227mg, 0.356 mmol), diisopropylethylamine (174 μ L, 1.065 mmol), *N*-deacetyl maytansine **124** (231 mg, 0.355 mmol) in 2 mL of DMF was added PyAOP (185 mg, 0.355 mmol). The solution was stirred for 30 min. Piperidine (0.5 mL) was added to the reaction mixture and stirred for an additional 20 min. The crude reaction mixture was purified by C18 reverse phase chromatography using a gradient of 0-100% acetonitrile:water affording 203.2 mg (55%, 2 steps) of compound **203**.

Synthesis of 17-(*tert*-butyl) 1-(((1⁴*S*,1⁶*S*,3³*S*,2*R*,4*S*,10*E*,12*E*,14*R*)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl) (2*S*)-8-(1-(3-(2-(((9*H*-fluoren-9-yl)methoxy)carbonyl)-1,2-dimethylhydrazinyl)methyl)-1*H*-indol-1-yl)propanoyl)piperidin-4-yl)-2,3-dimethyl-4,7-dioxo-11,14-dioxa-3,8-diazaheptadecanedioate (204)

[00456] A solution of piperidine **203** (203.2 mg, 0.194 mmol), ester **12** (126.5 mg, 0.194 mmol), 2,4,6-trimethylpyridine (77 μ L, 0.582 mmol), HOAT (26.4 mg, 0.194 mmol) in 1mL DMF was stirred 30 min. The crude reaction was purified by C18 reverse phase chromatography using a gradient of 0-100% acetonitrile:water with 0.1% formic acid affording 280.5 mg (97% yield) of compound **204**.

[00457] MS (ESI) *m/z*: [M+H]⁺ Calcd for C₈₁H₁₀₆ClN₈O₁₈ 1513.7; Found 1514.0.

Synthesis of (2*S*)-8-(1-(3-(2-(((9*H*-fluoren-9-yl)methoxy)carbonyl)-1,2-dimethylhydrazinyl)methyl)-1*H*-indol-1-yl)propanoyl)piperidin-4-yl)-1-(((1⁴*S*,1⁶*S*,3³*S*,2*R*,4*S*,10*E*,12*E*,14*R*)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-

benzenacyclotetradecaphane-10,12-dien-4-yl)oxy)-2,3-dimethyl-1,4,7-trioxo-11,14-dioxo-3,8-diazaheptadecan-17-oic acid (205)


[00458] To a solution of compound **204** (108 mg, 0.0714 mmol) in 500 μ L anhydrous DCM was added 357 μ L of a 1M solution of SnCl_4 in DCM. The heterogeneous mixture was stirred for 1 h and then purified by C18 reverse phase chromatography using a gradient of 0-100% acetonitrile:water with 0.1% formic acid affording 78.4 mg (75% yield) of compound **205**.

[00459] MS (ESI) m/z : $[\text{M}-\text{H}]^-$ Calcd for $\text{C}_{77}\text{H}_{96}\text{ClN}_8\text{O}_{18}$ 1455.7; Found 1455.9.

EXAMPLE 2

[00460] A linker containing a 4-amino-piperidine (4AP) group was synthesized according to Scheme 2, shown below.

Scheme 2



Synthesis of *tert*-butyl 4-oxopiperidine-1-carboxylate (210)

[00461] To a 100 mL round-bottom flask containing a magnetic stir bar was added piperidin-4-one hydrochloride monohydrate (1.53 g, 10 mmol), di-*tert*-butyl dicarbonate (2.39 g, 11 mmol), sodium carbonate (1.22 g, 11.5 mmol), dioxane (10 mL), and water (1 mL). The reaction mixture was stirred at room temperature for 1 h. The mixture was diluted with water (100 mL) and extracted with EtOAc (3 x 100 mL). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting material was dried in vacuo to yield 1.74 g (87%) of compound **210** as a white solid.

[00462] ¹H NMR (CDCl₃) δ 3.73 (t, 4H, *J* = 6.0), 2.46 (t, 4H, *J* = 6.0), 1.51 (s, 9H).

[00463] MS (ESI) *m/z*: [M+H]⁺ Calcd for C₁₀H₁₈NO₃ 200.3; Found 200.2.

Synthesis of *tert*-butyl 4-((2-(2-(3-(*tert*-butoxy)-3-oxopropoxy)ethoxy)ethyl)amino)piperidine-1-carboxylate (211)

[00464] To a dried scintillation vial containing a magnetic stir bar was added *tert*-butyl 4-oxopiperidine-1-carboxylate (399 mg, 2 mmol), H₂N-PEG₂-COO*t*-Bu (550 mg, 2.4 mmol), 4 Å molecular sieves (activated powder, 200 mg), and 1,2-dichloroethane (5 mL). The mixture was stirred for 1 h at room temperature. To the reaction mixture was added sodium triacetoxyborohydride (845 mg, 4 mmol). The mixture was stirred for 3 days at room temperature. The resulting mixture was partitioned between EtOAc and saturated aqueous NaHCO₃. The organic layer was washed with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure to afford 850 mg of compound **211** as a viscous oil.

[00465] MS (ESI) *m/z*: [M+H]⁺ Calcd for C₂₁H₄₁N₂O₆ 417.3; Found 417.2.

Synthesis of 13-(1-(*tert*-butoxycarbonyl)piperidin-4-yl)-2,2-dimethyl-4,14-dioxo-3,7,10-trioxa-13-azaheptadecan-17-oic acid (212)

[00466] To a dried scintillation vial containing a magnetic stir bar was added *tert*-butyl 4-((2-(2-(3-(*tert*-butoxy)-3-oxopropoxy)ethoxy)ethyl)amino)piperidine-1-carboxylate **211** (220 mg, 0.5 mmol), succinic anhydride (55 mg, 0.55 mmol), 4-(dimethylamino)pyridine (5 mg, 0.04 mmol), and dichloromethane (3 mL). The mixture was stirred for 24 h at room temperature. The reaction mixture was partially purified by flash chromatography (elute 50-100% EtOAc/hexanes)

to yield 117 mg of compound **212** as a clear oil, which was carried forward without further characterization.

[00467] MS (ESI) m/z: [M+H]⁺ Calcd for C₂₅H₄₅N₂O₉ 517.6; Found 517.5.

Synthesis of 17-(*tert*-butyl) 1-(((1⁴S,1⁶S,3³S,2R,4S,10E,12E,14R)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl) (2S)-8-(1-(*tert*-butoxycarbonyl)piperidin-4-yl)-2,3-dimethyl-4,7-dioxo-11,14-dioxa-3,8-diazaheptadecanedioate (213)

[00468] To a dried scintillation vial containing a magnetic stir bar was added 13-(1-(*tert*-butoxycarbonyl)piperidin-4-yl)-2,2-dimethyl-4,14-dioxo-3,7,10-trioxa-13-azaheptadecan-17-oic acid **212** (55 mg, 0.1 mmol), *N*-deacyl maytansine **124** (65 mg, 0.1 mmol), HATU (43 mg, 0.11 mmol), DMF (1 mL), and dichloromethane (0.5 mL). The mixture was stirred for 8 h at room temperature. The reaction mixture was directly purified by C18 flash chromatography (elute 5-100% MeCN/water) to give 18 mg (16%) of compound **213** as a white film.

[00469] MS (ESI) m/z: [M+H]⁺ Calcd for C₅₇H₈₇ClN₅O₁₇ 1148.6; Found 1148.7.

Synthesis of (2S)-1-(((1⁴S,1⁶S,3³S,2R,4S,10E,12E,14R)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl)oxy)-2,3-dimethyl-1,4,7-trioxa-8-(piperidin-4-yl)-11,14-dioxa-3,8-diazaheptadecan-17-oic acid (214)

[00470] To a dried scintillation vial containing a magnetic stir bar was added maytansinoid **213** (31 mg, 0.027 mmol) and dichloromethane (1 mL). The solution was cooled to 0 °C and tin(IV) tetrachloride (1.0 M solution in dichloromethane, 0.3 mL, 0.3 mmol) was added. The reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was directly purified by C18 flash chromatography (elute 5-100% MeCN/water) to yield 16 mg (60%) of compound **214** as a white solid (16 mg, 60% yield).

[00471] MS (ESI) m/z: [M+H]⁺ Calcd for C₄₈H₇₁ClN₅O₁₅ 992.5; Found 992.6.

Synthesis of (2S)-8-(1-(3-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)-1,2-dimethylhydrazinyl)methyl)-1H-indol-1-yl)propanoyl)piperidin-4-yl)-1-(((1⁴S,1⁶S,3³S,2R,4S,10E,12E,14R)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-

tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl)oxy)-2,3-dimethyl-1,4,7-trioxo-11,14-dioxa-3,8-diazaheptadecan-17-oic acid (215)

[00472] To a dried scintillation vial containing a magnetic stir bar was added maytansinoid **214** (16 mg, 0.016 mmol), (9*H*-fluoren-9-yl)methyl 1,2-dimethyl-2-((1-(3-oxo-3-(perfluorophenoxy)propyl)-1*H*-indol-2-yl)methyl)hydrazine-1-carboxylate (**5**) (13 mg, 0.02 mmol), DIPEA (8 μ L, 0.05 mmol), and DMF (1 mL). The solution was stirred for 18 h at room temperature. The reaction mixture was directly purified by C18 flash chromatography (elute 5-100% MeCN/water) to yield 18 mg (77%) of compound **215** as a white solid.

[00473] MS (ESI) *m/z*: [M+H]⁺ Calcd for C₇₇H₉₈ClN₈O₁₈ 1457.7; Found 1457.9.

Synthesis of (2*S*)-1-(((1⁴*S*,1⁶*S*,3³*S*,2*R*,4*S*,10*E*,12*E*,14*R*)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl)oxy)-8-(1-(3-(2-((1,2-dimethylhydrazinyl)methyl)-1*H*-indol-1-yl)propanoyl)piperidin-4-yl)-2,3-dimethyl-1,4,7-trioxo-11,14-dioxa-3,8-diazaheptadecan-17-oic acid (216)

[00474] To a dried scintillation vial containing a magnetic stir bar was added maytansinoid **215** (18 mg, 0.012 mmol), piperidine (20 μ L, 0.02 mmol), and DMF (1 mL). The solution was stirred for 20 minutes at room temperature. The reaction mixture was directly purified by C18 flash chromatography (elute 1-60% MeCN/water) to yield 15 mg (98%) of compound **216** (also referred to herein as HIPS-4AP-maytansine or HIPS-4-amino-piperidin-maytansine) as a white solid.

[00475] MS (ESI) *m/z*: [M+H]⁺ Calcd for C₆₂H₈₈ClN₈O₁₆ 1235.6; Found 1236.0.

EXAMPLE 3

Experimental procedures

General

[00476] Experiments were performed to create site-specifically conjugated antibody-drug conjugates (ADCs). Site-specific ADC production included the incorporation of formylglycine (FGly), a non-natural amino acid, into the protein sequence. To install FGly (FIG. 1), a short consensus sequence, CXPXR, where X is serine, threonine, alanine, or glycine, was inserted at

the desired location in the conserved regions of antibody heavy or light chains using standard molecular biology cloning techniques. This “tagged” construct was produced recombinantly in cells that coexpress the formylglycine-generating enzyme (FGE), which cotranslationally converted the cysteine within the tag into an FGly residue, generating an aldehyde functional group (also referred to herein as an aldehyde tag). The aldehyde functional group served as a chemical handle for bioorthogonal conjugation. A hydrazino-*iso*-Pictet-Spengler (HIPS) ligation was used to connect the payload (e.g., a drug, such as a cytotoxin (e.g., maytansine)) to FGly, resulting in the formation of a stable, covalent C-C bond between the cytotoxin payload and the antibody. This C-C bond was expected to be stable to physiologically-relevant conditions encountered by the ADC during circulation and FcRn recycling, e.g., proteases, low pH, and reducing reagents. Antibodies bearing the aldehyde tag may be produced at a variety of locations. Experiments were performed to test the effects of inserting the aldehyde tag at the heavy chain C-terminus (CT). Biophysical and functional characterization was performed on the resulting ADCs made by conjugation to maytansine payloads via a HIPS linker.

Cloning, expression, and purification of tagged antibodies

[00477] The aldehyde tag sequence was inserted at the heavy chain C-terminus (CT) using standard molecular biology techniques. For small-scale production, CHO-S cells were transfected with human FGE expression constructs and pools of FGE-overexpressing cells were used for the transient production of antibodies. For larger-scale production, GPEX technology (Catalent, Inc., Somerset, NJ) was used to generate a clonal cell line overexpressing human FGE (GPEX). Then, the FGE clone was used to generate bulk stable pools of antibody-expressing cells. Antibodies were purified from the conditioned medium using a Protein A chromatography (MabSelect, GE Healthcare Life Sciences, Pittsburgh, PA). Purified antibodies were flash frozen and stored at -80 °C until further use.

Bioconjugation, Purification, and HPLC Analytics

[00478] C-terminally aldehyde-tagged α CD22 antibody (15 mg/mL) was conjugated to HIPS-4AP-maytansine (8 mol. equivalents drug:antibody) for 72 h at 37 °C in 50 mM sodium citrate, 50 mM NaCl pH 5.5 containing 0.85% DMA. Unconjugated antibody was removed using preparative-scale hydrophobic interaction chromatography (HIC; GE Healthcare 17-5195-01)

with mobile phase A: 1.0 M ammonium sulfate, 25 mM sodium phosphate pH 7.0, and mobile phase B: 25% isopropanol, 18.75 mM sodium phosphate pH 7.0. An isocratic gradient of 33% B was used to elute unconjugated material, followed by a linear gradient of 41-95% B to elute mono- and diconjugated species. To determine the DAR of the final product, ADCs were examined by analytical HIC (Tosoh #14947, Grove City, OH) with mobile phase A: 1.5 M ammonium sulfate, 25 mM sodium phosphate pH 7.0, and mobile phase B: 25% isopropanol, 18.75 mM sodium phosphate pH 7.0. To determine aggregation, samples were analyzed using analytical size exclusion chromatography (SEC; Tosoh #08541) with a mobile phase of 300 mM NaCl, 25 mM sodium phosphate pH 6.8.

Results

[00479] α CD22 antibodies modified to contain the aldehyde tag at the heavy chain C-terminus (CT) were conjugated to a maytansine payload attached to a HIPS-4AP linker as described above. Upon completion of the conjugation reaction, the unconjugated antibody was removed by preparative HIC and remaining free drug was removed during buffer exchange by tangential flow filtration. The reactions were high yielding, with $\geq 84\%$ conjugation efficiency and $>70\%$ total yield. The resulting ADCs had drug-to-antibody ratios (DARs) of 1.6-1.9 and were predominately monomeric. FIGS. 2-5 show DARs from representative crude reactions and the purified ADCs as determined by HIC and reversed phase PLRP chromatography, and show the monomeric integrity as determined by SEC.

[00480] FIG. 2 shows shows a hydrophobic interaction column (HIC) trace of an aldehyde-tagged anti-CD22 antibody conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker. FIG. 2 indicates that the crude DAR was 1.68 as determined by HIC.

[00481] FIG. 3 shows a HIC trace of an aldehyde-tagged anti-CD22 antibody conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker. FIG. 3 indicates that the final DAR was 1.77 as determined by HIC.

[00482] FIG. 4 shows a reversed phase chromatography (PLRP) trace of an aldehyde-tagged anti-CD22 antibody conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker. FIG. 4 indicates that the final DAR was 1.81 as determined by PLRP.

[00483] FIG. 5 shows a graph of analytical size exclusion chromatography (SEC) analysis of an aldehyde-tagged anti-CD22 antibody conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker. As shown in FIG. 5, analytical SEC indicated 98.2% monomer for the final product.

***In vitro* cytotoxicity**

[00484] The CD22-positive B-cell lymphoma cell lines, Ramos and WSU-DLCL2, were obtained from the ATCC and DSMZ cell banks, respectively. The cells were maintained in RPMI-1640 medium (Cellgro, Manassas, VA) supplemented with 10% fetal bovine serum (Invitrogen, Grand Island, NY) and Glutamax (Invitrogen). 24 h prior to plating, cells were passaged to ensure log-phase growth. On the day of plating, 5000 cells/well were seeded onto 96-well plates in 90 μ L normal growth medium supplemented with 10 IU penicillin and 10 μ g/mL streptomycin (Cellgro). Cells were treated at various concentrations with 10 μ L of diluted analytes, and the plates were incubated at 37 °C in an atmosphere of 5% CO₂. After 5 d, 100 μ L/well of Cell Titer-Glo reagent (Promega, Madison, WI) was added, and luminescence was measured using a Molecular Devices SpectraMax M5 plate reader. GraphPad Prism software was used for data analysis.

Results

[00485] α CD22 CT HIPS-4AP-maytansine exhibited very potent activity against WSU-DLCL2 and Ramos cells *in vitro* as compared to free maytansine (FIG. 6). The IC₅₀ concentrations were 0.018 and 0.086 nM for the ADC and the free drug, respectively, against WSU-DLCL2 cells, and were 0.007 and 0.040 nM for the ADC and the free drug, respectively, against Ramos cells.

[00486] FIG. 6A shows a graph of *in vitro* potency against WSU-DLCL2 cells (% viability vs. Log antibody-drug conjugate (ADC) concentration (nM)) for anti-CD22 ADCs conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker. FIG. 6B shows a graph of *in vitro* potency against Ramos cells (% viability vs. Log antibody-drug conjugate (ADC) concentration (nM)) for anti-CD22 ADCs conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker.

Xenograft studies

[00487] Female ICR SCID mice (8/group) were inoculated subcutaneously with 5×10^6 WSU-DLCL2 cells. Treatment began when the tumors reached an average of 262 mm^3 , at which time the animals were dosed intravenously with vehicle alone or CT-tagged αCD22 HIPS-4AP-maytansine (10 mg/kg). Dosing proceeded every four days for a total of four doses (q4d x 4). The animals were monitored twice weekly for body weight and tumor size. Animals were euthanized when tumors reached 2000 mm^3 .

Results

[00488] The median time to endpoint for animals in the vehicle control groups was 16 days; therefore, tumor growth inhibition (TGI%) was calculated at that day. TGI% was defined by the following formula:

$$\text{TGI (\%)} = (\text{TV}_{\text{control group}} - \text{TV}_{\text{treated group}}) / \text{TV}_{\text{control}} \times 100$$

where TV is tumor volume.

[00489] The animals that were dosed with αCD22 HIPS-4AP-maytansine demonstrated 90% TGI at day 16, with 5 of the 8 tumors undergoing complete regression (FIG. 7). Three of these complete regressions were durable through the end of the study (day 58). FIG. 7 shows a graph indicating the *in vivo* efficacy against a WSU-DLCL2 xenograft model (mean tumor volume (mm^3) vs. days) for anti-CD22 ADCs conjugated at the C-terminus (CT) to a maytansine payload attached to a HIPS-4AP linker. The vertical arrows in FIG. 7 indicate dosing, which occurred every four days for a total of four doses (q4d x 4).

EXAMPLE 4

Introduction

[00490] Hematologically-derived tumors make up ~10% of all newly-diagnosed cancer cases in the U.S. Of these, the non-Hodgkin lymphoma (NHL) designation describes a diverse group of cancers that collectively rank among the top 10 most commonly diagnosed cancers worldwide. Although long-term survival trends are improving, there remains a significant unmet clinical need for treatments to help patients with relapsed or refractory disease, one cause of which is drug efflux through upregulation of xenobiotic pumps, such as MDR1. A site-

specifically-conjugated antibody-drug conjugate targeted against CD22 and bearing a noncleavable maytansine payload that was resistant to MDR1-mediated efflux was produced. The construct was efficacious against CD22+ NHL xenografts and can be repeatedly dosed in cynomolgus monkeys at 60 mg/kg with no observed adverse effects. Together, the data indicated that this drug has the potential to be used effectively in patients with CD22+ tumors that have developed MDR1-related resistance to prior therapies. CD22 is a clinically-validated target for the treatment of NHL and ALL. An anti-CD22 antibody-drug conjugate (ADC) according to the present disclosure can be used for the treatment of relapsed/refractory NHL and ALL patients.

Material and Methods

[00491] An anti-CD22 antibody was conjugated site-specifically, using aldehyde tag technology, to a non-cleavable maytansine-payload linker. The ADC was characterized both biophysically and functionally *in vitro*. Then, *in vivo* efficacy was determined in mice using two xenograft models and toxicity studies were performed in both rat and cynomolgus monkeys. Pharmacodynamic studies were conducted in monkeys, and pharmaco- and toxicokinetic studies compared total ADC exposure in the efficacy and toxicity studies.

Results

[00492] The ADC was very potent *in vivo*, even against cell lines that had been constructed to overexpress the efflux pump, MDR1. The construct was efficacious at 10 mg/kg x 4 doses against NHL xenograft tumor models, and in a cynomolgus toxicity study, the ADC was dosed twice at 60 mg/kg with no observed adverse effects. Exposure to total ADC at these doses (as assessed by AUC_{0-inf}) indicated that the exposure needed to achieve efficacy was below tolerable limits. Finally, an examination of the pharmacodynamic response in the treated monkeys demonstrated that the B-cell compartment was selectively depleted, indicating that the ADC eliminated targeted cells without notable off-target toxicity.

[00493] The results indicated that the ADC can be used effectively in patients with CD22+ tumors that have developed MDR1-related resistance to prior therapies.

EXAMPLE 5**Introduction**

[00494] Leukemias, lymphomas, and myelomas are highly prevalent in the population, accounting for ~10% of all newly diagnosed cancer cases in the U.S. during 2015. Of these cancers, B-cell derived malignancies make up a large and diverse group that includes non-Hodgkin lymphoma (NHL), chronic lymphocytic leukemia (CLL), and acute lymphoblastic leukemia (ALL). Similarly, as a category, NHL designates about 60 lymphoma subsets, of which about 85% are B-cell derived, including diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), and mantle cell lymphoma (MCL). Collectively, NHL diseases are among the most common cancer types observed, ranking as the 7th most common cancer in the U.S., and the 10th most common cancer diagnosed worldwide in 2012. While long-term trends show improvements in 5-year survival rates for most blood cancer diagnoses, there remains a significant unmet clinical need, with 16% of CLL, 30% of ALL, and 30% of NHL patients diagnosed from 2004 to 2010 failing to meet the 5-year survival endpoint.

[00495] CD22 is a B-cell lineage-restricted cell surface glycoprotein that is expressed on the majority of B-cell hematologic malignancies, but is not expressed on hematopoietic stem cells, memory B cells, or other normal non-hematopoietic tissues. Its expression pattern and rapid internalization kinetics make it a target for antibody-drug conjugate (ADC) therapies, and it has been validated as such in clinical trials against NHL and ALL.

[00496] In the experiments described herein, site-specific conjugation technology based upon the aldehyde tag and Hydrazino-*iso*-Pictet-Spengler (HIPS) chemistry was used to place a maytansine payload coupled through a noncleavable linker to the antibody heavy chain C-terminus. The genetically-encoded aldehyde tag incorporated the six amino acid sequence, **LCTPSR** (SEQ ID NO: 32). Cotranslationally, overexpressed formylglycine generating enzyme (FGE) converted the cysteine within the consensus sequence to a formylglycine residue, bearing an aldehyde functional group, which was reacted with a HIPS-linker-payload to generate an ADC. This approach afforded control over both payload placement and DAR, and yielded highly homogenous ADC preparations. Site-specifically conjugated ADCs displayed improved pharmacokinetics (PK) and efficacy relative to stochastic conjugates, likely due to the lack of under- and overconjugated species in the preparation, which can lead to ineffective or overly toxic molecules, respectively. Furthermore, the noncleavable linker-maytansine payload used on

the anti-CD22 ADC was resistant to efflux by MDR1 and did not mediate off-target or bystander killing. Together, these features contributed to the efficacy and safety of the anti-CD22 ADC observed in preclinical studies.

Materials and Methods

General

[00497] All animal studies were conducted in accordance with Institutional Animal Care and Use Committee guidelines and were performed at Charles River Laboratories, Aragen Bioscience, or Covance Laboratories. The murine anti-maytansine antibody was made by ProMab and validated in-house. The rabbit anti-AF488 antibody was purchased from Life Technologies. The horseradish peroxidase (HRP)-conjugated secondary antibodies were from Jackson ImmunoResearch. The antibodies used for pharmacodynamic studies were from BD Pharmingen. Cell lines were obtained from ATCC and DSMZ cell banks where they were authenticated by morphology, karyotyping, and PCR based approaches.

Cloning, expression, and purification of tagged antibodies

[00498] Antibodies were generated using standard cloning and purification techniques and GPEX® expression technology.

Bioconjugation, purification, and HPLC analytics

[00499] ADCs were made and characterized as described in Drake et al., *Bioconjugate Chem.*, 2014, 25, 1331-41.

Generation of MDR1+ cell lines

[00500] MDR1 (ABCB1) cDNA was obtained from Sino Biological and cloned into a pEF plasmid with a hygromycin selection marker. An AMAXA Nucleofector™ instrument was used to electroporate Ramos (ATCC CRL-1923) and WSU-DLCL2 (DSMZ ACC 575) cells according to the manufacturer's instructions. After selection with hygromycin (Invitrogen 10687010), the pools were enriched with paclitaxel treatment (25 nM for up to 10 days) to further select cells with functional MDR1. The resulting cells were maintained under

hygromycin selection in RPMI (Gibco 21870-092) supplemented with 10% fetal bovine serum (FBS) and 1X GlutaMax (Gibco 35050-079).

***In vitro* cytotoxicity assays**

[00501] Cell lines were plated in 96-well plates (Costar 3610) at a density of 5×10^4 cells/well in 100 μ L of growth media and allowed to rest for 5 h. Serial dilution of test samples was performed in RPMI at 6x the final concentration and 20 μ L was added to the cells. After incubation at 37 °C with 5% CO₂ for 5 days, viability was measured using a Promega CellTiter 96® AQueous One Solution Cell Proliferation Assay (G3581) according to the manufacturer's instructions. GI₅₀ curves were calculated in GraphPad Prism using the ADC's drug-to-antibody ratio (DAR) value to normalize the dose to the payload concentration.

Xenograft studies

[00502] Female CB17 ICR SCID mice were inoculated subcutaneously with either WSU-DLCL2 or Ramos cells in 50% Matrigel. Tumors were measured twice weekly and tumor volume was estimated according to the formula: tumor volume (mm^3) = $\frac{w^2 \times l}{2}$ where w = tumor width and l = tumor length. When tumors reached the desired mean volume, animals were randomized into groups of 8-12 mice and were dosed as described below. Animals were euthanized at the end of the study or when tumors reached 2000 mm^3 .

Rat toxicology study and toxicokinetic (TK) analysis

[00503] Male Sprague-Dawley rats (8-9 wk old at study start) were given a single intravenous dose of 6, 20, 40, or 60 mg/kg of the anti-CD22 ADC (5 animals/group). Animals were observed for 12 days post-dose. Body weights were recorded on days 0, 1, 4, 8, and 11. Blood was collected from all animals at 8 h and at 5, 9, and 12 d and was used for toxicokinetic analyses (all time points) and for clinical chemistry and hematology analyses (days 5 and 12). Toxicokinetic analyses were performed by ELISA, using the same conditions and reagents as described for the pharmacokinetic analyses.

Non-human primate toxicology and TK studies

[00504] Cynomolgus monkeys (2/sex/group) were given two doses (every 21 days) of 10, 30, or 60 mg/kg of the anti-CD22 ADC followed by a 21 day observation period. Body weights were assessed prior to dosing on day 1, and on days 8, 15, 22 (predose), 29, 36, and 42. Blood was collected for toxicokinetic, clinical chemistry, and hematology analyses according to the schedules presented in Table 2. Toxicokinetic analyses were performed by ELISA, using the same conditions and reagents as described for the pharmacokinetic analyses, except that CD22-His protein was used as the capture reagent for the total antibody and total ADC measurements.

Table 2. Summary of pharmacokinetic findings in rats dosed at 3 mg/kg with anti-CD22 ADC

Parameter, mean (SD)	Total Ab	Total ADC	Total Conjugate
AUC _{0-inf} (day $\square$ μ g/mL)	304 (40)	218 (18)	261 (26)
Clearance (mL/day/kg)	10.0 (1)	13.8 (1)	11.6 (1)
C _{0.04 d}	73.2 (5)	83.9 (16)	76.8 (6)
t _{1/2 effective} (days)*	9.48 (1)	6.13 (0.6)	7.22 (0.6)
V _{ss} (mL/kg)	41.1 (3)	36.7 (7)	39.2 (3)

Total antibody measures conjugated and unconjugated Ab; Total ADC is a DAR-sensitive measurement; Total conjugate measures all analytes with DAR ≥ 1 . SD, standard deviation; AUC_{0-inf}, area under the concentration versus time curve from time 0 to infinity; C_{0.04 d}, concentration observed at 1 h; t_{1/2 effective}, Effective half-life; V_{ss}, volume of distribution at steady state. *The uncertainty for half-life is given as standard error.

Non-human primate pharmacodynamic study

[00505] Whole blood samples from the cynomolgus monkeys enrolled in the anti-CD22 ADC toxicology study were analyzed by flow cytometry to assess CD3⁺, CD20⁺, and CD3⁻/CD20⁻ leukocyte populations. Briefly, to a 100 μ L aliquot of whole blood, either fluorescein and phycoerythrin-conjugated isotype control antibodies or fluorescein-conjugated anti-CD20 and phycoerythrin-conjugated anti-CD3 antibodies were added and incubated on ice for 30 min. Then, red blood cells were lysed with an ammonium chloride solution (Stem Cell Technologies),

and cells were washed twice in phosphate buffered saline + 1% FBS. Labeled cells were analyzed by flow cytometry on a FACSCanto™ instrument running FACSDiva™ software.

Pharmacokinetic (PK) study designs

[00506] For the mouse study, animals used in the Ramos xenograft experiment were sampled in groups of three at time points beginning at 1 h post-first dose and continuing across the observation period. For the rat study, male Sprague-Dawley rats (3 per group) were dosed intravenously with a single 3 mg/kg bolus of ADC. Plasma was collected at 1 h, 8 h and 24 h, and 2, 4, 6, 8, 10, 14, and 21 days post-dose. Plasma samples were stored at -80 °C until use.

PK and TK sample analysis

[00507] The concentrations of total antibody, total ADC (DAR-sensitive), and total conjugate (DAR ≥ 1) were quantified by ELISA as diagrammed in FIG. 10. For total antibody, conjugates were captured with an anti-human IgG-specific antibody and detected with an HRP-conjugated anti-human Fc-specific antibody. For total ADC, conjugates were captured with an anti-human Fab-specific antibody and detected with a mouse anti-maytansine primary antibody, followed by an HRP-conjugated anti-mouse IgG-subclass 1-specific secondary antibody. For total conjugate, conjugates were captured with an anti-maytansine antibody and detected with an HRP-conjugated anti-human Fc-specific antibody. Bound secondary antibody was detected using Ultra TMB One-Step ELISA substrate (Thermo Fisher). After quenching the reaction with sulfuric acid, signals were read by taking the absorbance at 450 nm on a Molecular Devices Spectra Max M5 plate reader equipped with SoftMax Pro software. Data were analyzed using GraphPad Prism and Microsoft Excel software.

Indirect ELISA CD22 antigen binding

[00508] Maxisorp 96-well plates (Nunc) were coated overnight at 4 °C with 1 μ g/mL of human CD22-His (Sino Biological) in PBS. The plate was blocked with casein buffer (ThermoFisher), and then the anti-CD22 wild-type antibody and ADCs were plated in an 11-step series of 2-fold dilutions starting at 200 ng/mL. The plate was incubated, shaking, at room temperature for 2 h. After washing in phosphate-buffered saline (PBS) 0.1% Tween-20, bound analyte was detected with a donkey anti-human Fc- γ -specific horseradish peroxidase (HRP)-

conjugated secondary antibody. Signals were visualized with Ultra TMB (Pierce) and quenched with 2 N H₂SO₄. Absorbance at 450 nm was determined using a Molecular Devices SpectraMax M5 plate reader and the data were analyzed using GraphPad Prism.


Anti-CD22 ADC mediated CD22 internalization on CD22+ NHL cell lines

[00509] Ramos, Granta-519, and WSU-DLCL2 cells (1e6/test) were incubated either in labeling buffer alone [PBS+ 1% fetal bovine serum (FBS)], or in labeling buffer with the anti-CD22 ADC (1 µg/test). Samples were placed at 4 or 37 °C for 2 h. Then, cells were incubated on ice for 20 min with fluorescein-labeled anti-CD22. After washing 2x in labeling buffer, cells were analyzed by flow cytometry on a FACSCanto™ instrument running FACSDiva™ software. The difference in fluorescence between cells at 4 and 37 °C ± ADC was interpreted as anti-CD22 ADC-mediated internalization.

Cynomolgus and human tissue cross-reactivity studies

[00510] Tissue cross-reactivity studies were performed by Ensigna Biosystems Inc. (Richmond, CA) using biotinylated anti-CD22 ADC and a biotinylated HIPS-4AP-maytansine linker payload-conjugated isotype antibody as a control. Tissue microarrays containing skin, heart, lung, kidney, liver, pancreas, stomach, small intestine, large intestine, and spleen (a positive control) were used. Primary antibody was detected using streptavidin conjugated to horseradish peroxidase followed by visualization with DAB substrate.

Synthesis of HIPS-4AP-maytansine linker payload




(9H-Fluoren-9-yl)methyl 1,2-dimethylhydrazine-1-carboxylate (2).

[00511] MeNHNHMe•2HCl (1) (5.0 g, 37.6 mmol) was dissolved in CH₃CN (80 mL). Et₃N (22 mL, 158 mmol) was added and the precipitate that formed was removed by filtration. To the remaining solution of MeNHNHMe, a solution of FmocCl (0.49 g, 18.9 mmol, 0.5 eq) was added dropwise over 2.5 h at -20 °C. The reaction mixture was then diluted with EtOAc,

washed with H₂O, brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by flash chromatography on silica (hexanes:EtOAc = 3:2) to give 3.6 g (34%) of compound **2**.

[00512] ¹H NMR (400 MHz, CDCl₃) δ 7.75-7.37 (m, 8 H), 4.48 (br s, 2H), 4.27 (t, *J* = 6.0 Hz, 1H), 3.05 (s, 3H), 2.55 (br s, 3H).



2-(((*tert*-Butyldimethylsilyl)oxy)methyl)-1*H*-indole (**4**)

[00513] An oven-dried flask was charged with indole-2-methanol, **3**, (1.581 g, 10.74 mmol), TBSCl (1.789 g, 11.87 mmol), and imidazole (2.197 g, 32.27 mmol), and this mixture was suspended in CH₂Cl₂ (40 mL, anhydrous). After 16 h, the reaction mixture was concentrated to an orange residue. The crude mixture was taken up in Et₂O (50 mL), washed with aqueous AcOH (5% v/v, 3 x 50 mL) and brine (25 mL). The combined organic layers were dried over Na₂SO₄ and concentrated to give 2.789 g (99%) of compound **4** as a crystalline solid which was used without further purification.

[00514] ^1H NMR (500 MHz, CDCl_3) δ 8.29 (s, 1H), 7.57 (d, $J = 7.7$ Hz, 1H), 7.37 (dd, $J = 8.1, 0.6$ Hz, 1H), 7.19 – 7.14 (m, 1H), 7.12 – 7.07 (m, 1H), 6.32 (d, $J = 1.0$ Hz, 1H), 4.89 (s, 2H), 0.95 (s, 9H), 0.12 (s, 6H).

[00515] ^{13}C NMR (101 MHz, CDCl_3) δ 138.3, 136.0, 128.6, 121.7, 120.5, 119.8, 110.9, 99.0, 59.4, 26.1, 18.5, -5.2.

[00516] HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{24}\text{NOSi}$ $[\text{M}+\text{H}]^+$: 262.1627; found: 262.1625.

Methyl 3-(2-(((*tert*-butyldimethylsilyl)oxy)methyl)-1*H*-indol-1-yl)propanoate (6)

[00517] To a solution of indole **4** (2.789 μg , 10.67 mmol) in CH_3CN (25 mL) was added methyl acrylate, **5**, (4.80 mL, 53.3 mmol) followed by 1,8-diazabicyclo[5.4.0]undec-7-ene (800 μL , 5.35 mmol), and the resulting mixture was refluxed. After 18 h, the solution was cooled and concentrated to an orange oil which was purified by silica gel chromatography (9:1 hexanes:EtOAc) to yield 3.543 g (96%) of compound **6** as a colorless oil.

[00518] ^1H NMR (400 MHz, CDCl_3) δ 7.58 (d, $J = 7.8$ Hz, 1H), 7.34 (d, $J = 8.2$ Hz, 1H), 7.23 – 7.18 (m, 1H), 7.12 – 7.07_{sep}^[1.1] (m, 1H), 6.38 (s, 1H), 4.84 (s, 2H), 4.54 – 4.49 (m, 2H), 2.89 – 2.84 (m, 2H), 0.91 (s, 9H), 0.10 (s, 6H).

[00519] ^{13}C NMR (101 MHz, CDCl_3) δ 172.0, 138.5, 137.1, 127.7, 122.0, 121.0, 119.8, 109.3, 101.8, 58.2, 51.9, 39.5, 34.6, 26.0, 18.4, -5.2.

[00520] HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{30}\text{NO}_3\text{Si}$ $[\text{M}+\text{H}]^+$: 348.1995; found: 348.1996.

Methyl 3-(2-(hydroxymethyl)-1*H*-indol-1-yl)propanoate (7)

[00521] To a solution of compound **6** (1.283 g, 3.692 mmol) in THF (20 mL) at 0 °C was added a 1.0 M solution of tetrabutylammonium fluoride in THF (3.90 mL, 3.90 mmol). After 15 minutes, the reaction mixture was diluted with Et_2O (20 mL) and washed with NaHCO_3 (sat. aq., 3 x 20 mL), and concentrated to a pale green oil. The oil was purified by silica gel chromatography (2:1 hexanes:EtOAc) to yield 822 mg (95%) of **7** as a white crystalline solid.

[00522] ^1H NMR (500 MHz, CDCl_3) δ 7.60 (d, $J = 7.8$ Hz, 1H), 7.34 (dd, $J = 8.2, 0.4$ Hz, 1H), 7.27 – 7.23 (m, 1H), 7.16 – 7.11 (m, 1H), 6.44 (s, 1H), 4.77 (s, 2H), 4.49 (t, $J = 7.3$ Hz, 2H), 3.66 (s, 3H), 2.87 (t, $J = 7.3$ Hz, 2H), 2.64 (s, 1H).

[00523] ^{13}C NMR (126 MHz, CDCl_3) δ 172.3, 138.5, 137.0, 127.6, 122.2, 121.1, 119.9, 109.3, 102.3, 57.1, 52.0, 39.1, 34.3.

[00524] HRMS (ESI) calcd for $C_{13}H_{15}NNaO_3$ $[M+Na]^+$: 256.0950; found: 256.0946.

Methyl 3-(2-formyl-1*H*-indol-1-yl)propanoate (8)

[00525] Dess-Martin periodinane (5.195 g, 12.25 mmol) was suspended in a mixture of CH_2Cl_2 (20 mL) and pyridine (2.70 mL, 33.5 mmol). After 5 min, the resulting white suspension was transferred to a solution of methyl 3-(2-(hydroxymethyl)-1*H*-indol-1-yl)propanoate (7; 2.611 g, 11.19 mmol) in CH_2Cl_2 (10 mL), resulting in a red-brown suspension. After 1 h, the reaction was quenched with sodium thiosulfate (10% aqueous solution, 5 mL) and $NaHCO_3$ (saturated aqueous solution, 5 mL). The aqueous layer was extracted with CH_2Cl_2 (3 x 20 mL); the combined extracts were dried over Na_2SO_4 , filtered, and concentrated to a brown oil. Purification by silica gel chromatography (5-50% EtOAc in hexanes) yielded 2.165 g (84%) of compound **8** as a colorless oil.

[00526] ^1H NMR (400 MHz, $CDCl_3$) δ 9.87 (s, 1H), 7.73 (dt, $J = 8.1, 1.0$ Hz, 1H), 7.51 (dd, $J = 8.6, 0.9$ Hz, 1H), 7.45 – 7.40 (m, 1H), 7.29 (d, $J = 0.9$ Hz, 1H), 7.18 (ddd, $J = 8.0, 6.9, 1.0$ Hz, 1H), 4.84 (t, $J = 7.2$ Hz, 2H), 3.62 (s, 3H), 2.83 (t, $J = 7.2$ Hz, 2H).

[00527] ^{13}C NMR (101 MHz, $CDCl_3$) δ 182.52, 171.75, 140.12, 135.10, 127.20, 126.39, 123.46, 121.18, 118.55, 110.62, 51.83, 40.56, 34.97.

[00528] HRMS (ESI) calcd for $C_{13}H_{13}NO_3Na$ $[M+Na]^+$: 254.0793; found: 254.0786.

3-(2-Formyl-1*H*-indol-1-yl)propanoic acid (9)

[00529] To a solution of indole **8** (2.369 g, 10.24 mmol) dissolved in dioxane (100 mL) was added LiOH (4 M aqueous solution, 7.68 mL, 30.73 mmol). A thick white precipitate gradually formed over the^[1]_{SEP} course of several hours. After 21 h, HCl (1 M aqueous solution, 30 mL) was added dropwise to give a solution with pH = 4. The solution was concentrated and the resulting pale brown oil was dissolved in EtOAc (50 mL) and washed with water (2 x 50 mL) and brine (20 mL). The organic layer was dried over Na_2SO_4 , filtered, and concentrated to an orange solid. Purification by silica gel chromatography (10-50% EtOAc in hexanes with 0.1% acetic acid) yielded 1.994 g (84%) of compound **9** as a pale yellow solid.

[00530] ^1H NMR (400 MHz, $CDCl_3$) δ 9.89 (s, 1H), 7.76 (dt, $J = 8.1, 0.9$ Hz, 1H), 7.53 (dd, $J = 8.6, 0.9$ Hz, 1H), 7.48 – 7.43 (m, 1H), 7.33 (d, $J = 0.8$ Hz, 1H), 7.21 (ddd, $J = 8.0, 6.9, 1.0$ Hz, 1H), 4.85 (t, $J = 7.2$ Hz, 2H), 2.91 (t, $J = 7.2$ Hz, 2H).

[00531] ^{13}C NMR (101 MHz, CDCl_3) δ 182.65, 176.96, 140.12, 135.02, 127.33, 126.42, 123.53, 121.27, 118.76, 110.55, 40.19, 34.82.

[00532] HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{10}\text{NO}_3$ $[\text{M}-\text{H}]^-$: 216.0666; found: 216.0665.

3-(2-((2-(((9H-Fluoren-9-yl)methoxy)carbonyl)-1,2-dimethylhydrazinyl)methyl)-1H-indol-1-yl)propanoic acid (10)

[00533] To a solution of compound **9** (1.193 g, 5.492 mmol) and (9H-fluoren-9-yl)methyl 1,2-dimethylhydrazinecarboxylate, **2**, (2.147 g, 7.604 mmol) in 1,2-dichloroethane (anhydrous, 25 mL) was added sodium triacetoxyborohydride (1.273 g, 6.006 mmol). The resulting yellow suspension was stirred for 2 h and then quenched with NaHCO_3 (saturated aqueous solution, 10 mL), followed by addition of HCl (1 M aqueous solution) to pH 4. The organic layer was separated, and the aqueous layer was extracted with CH_2Cl_2 (5 x 10 mL). The pooled organic extracts were dried over Na_2SO_4 , filtered, and concentrated to an orange oil. Purification by C18 silica gel chromatography (20-90% CH_3CN in water) yielded 1.656 g (62%) of compound **10** as a waxy pink solid.

[00534] ^1H NMR (400 MHz, CDCl_3) δ 7.76 (d, $J = 7.4$ Hz, 2H), 7.70 – 7.47 (br m, 3H), 7.42 – 7.16 (br m, 6H), 7.12 – 7.05 (m, 1H), 6.37 (s, 0.6H), 6.05 (s, 0.4H), 4.75 – 4.30 (br m, 4H), 4.23 (m, 1H), 4.10 (br s, 1H), 3.55 (br d, 1H), 3.11 – 2.69 (m, 5H), 2.57 (br s, 2H), 2.09 (br s, 1H).

[00535] ^{13}C NMR (101 MHz, CDCl_3) δ 174.90, 155.65, 143.81, 141.42, 136.98, 134.64, 127.75, 127.48, 127.12, 124.92, 122.00, 120.73, 120.01, 119.75, 109.19, 103.74, 67.33, 66.80, 51.39, 47.30, 39.58, 39.32, 35.23, 32.10.


[00536] HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{30}\text{N}_3\text{O}_4$ $[\text{M}+\text{H}]^+$: 484.2236; found: 484.2222.

(9H-Fluoren-9-yl)methyl 1,2-dimethyl-2-((1-(3-oxo-3-(perfluorophenoxy)propyl)-1H-indol-2-yl)methyl)hydrazine-1-carboxylate (RED-004).

[00537] Compound **10** (5.006 g, 10.4 mmol), was added to a dried 100 mL 2-neck round bottom flask containing a dried stir bar. Anhydrous EtOAc, 40 mL, was added by syringe and the solution stirred at 20 °C for 5 min. giving a clear, pale, yellow-green solution. The solution was cooled to 0 °C in an ice water bath and pentafluorophenol (2098.8 mg, 11.4 mmol), in 3 mL of anhydrous EtOAc, was added dropwise. The solution was stirred at 0 °C for 5 min. DCC

(2348.0 mg, 11.4 mmol), in 7 mL of anhydrous EtOAc, was added dropwise, slowly by syringe. The solution was stirred at 0 °C for 5 min, then removed from the bath and warmed to 20 °C. The reaction was stirred for 2 h, cooled to 0 °C, and filtered to give a clear, pale, yellow-green solution. The solution was diluted with 50 mL of EtOAc, and washed with 2 x 25 mL H₂O, 1 x 25 mL 5 M NaCl, and dried over Na₂SO₄. The solution was filtered, evaporated, and dried under high vacuum, giving 6552.5 mg (97%) of **RED-004** as a greenish-white solid.

[00538] ¹H NMR (400 MHz, CDCl₃) δ 780 (d, J = 7.2 Hz, 2H), 7.58 (m, 3H), 7.45-7.22 (m, 6H), 7.14 (dd(appt. t), J = 7.4 Hz, 1H), 6.42 & 6.10 (2 br s, 1H), 4.74 (dd(appt. t), J = 5.4 Hz, 2H), 3.65-3.18 (br, 3H), 3.08 & 2.65 (2 br s, 3H), 2.88 (s, 3H).




(S)-3,4-dimethyloxazolidine-2,5-dione (RED-194).

[00539] To a solution of *N*-Boc-Ala-OH (**11**) (0.005 mol) in methylene chloride (25 ml) at 0°C, was added under nitrogen 1.2 equivalent of phosphorous trichloride. The reaction mixture was stirred for 2 h at 0 °C, the solvent was removed under reduced pressure and the residue was washed with carbon tetrachloride (3 x 20 ml) to afford **RED-194**.

(1⁴S,1⁶S,3²R,3³R,2R,4S,10E,12E,14R)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl methyl-L-alaninate (RED-062).

[00540] Maytansinol (**RED-063**) (4.53g, 8 mmol) was dissolved in anhydrous DMF (11 mL) to give a clear, colorless solution that was transferred to a dried two neck round bottom flask under N₂. Anhydrous THF (44 mL) was added followed by DIPEA (8.4 mL, 48 mmol). A solution of **RED-194** (5.4 g, 42 mmol) was added to give a clear, colorless solution. Dessicated, finely ground Zn(OTf)₂ (8.7 g, 24 mmol) was added to the stirring solution and the reaction mixture was stirred at 20 °C for 2 days. The reaction was quenched by adding to a solution of 70 mL of 1.2 M NaHCO₃ and 70 mL EtOAc. Upon stirring the resulting mixture produced a white precipitate that was removed by filtration. The filtrate was extracted with EtOAc (5 x 70 mL), dried (Na₂SO₄) and concentrated to give a reddish orange oil. This was dissolved in CH₂Cl₂ (15 mL) and purified using a Biotage system (adsorbed on 2x Biotage Ultra 10 g samplets, purification on 2x Biotage Ultra 100 g cartridge with 0-20% gradient of MeOH in CH₂Cl₂) to produce 4.38 g of the **RED-062** as a pale peach solid (95% de, 93.7% desired diastereomer).



tert-butyl 4-oxopiperidine-1-carboxylate (**13**).

[00541] To a 100 mL round-bottom flask containing a magnetic stir bar was added piperidin-4-one hydrochloride monohydrate (**12**) (1.53 g, 10 mmol), di-*tert*-butyl dicarbonate (2.39 g, 11 mmol), sodium carbonate (1.22 g, 11.5 mmol), dioxane (10 mL), and water (1 mL). The reaction mixture was stirred at room temperature for 1 h. The mixture was diluted with water (100 mL) and extracted with EtOAc (3 x 100 mL). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The resulting material was dried in vacuo to yield 1.74 g (87%) of compound **13** as a white solid.

[00542] ¹H NMR (CDCl₃) δ 3.73 (t, 4H, *J* = 6.0), 2.46 (t, 4H, *J* = 6.0), 1.51 (s, 9H).

[00543] MS (ESI) *m/z*: [M+H]⁺ Calcd for C₁₀H₁₈NO₃ 200.3; Found 200.2.

***tert*-butyl 4-((2-(2-(3-(*tert*-butoxy)-3-oxopropoxy)ethoxy)ethyl)amino)piperidine-1-carboxylate (**14**).**

[00544] To a dried scintillation vial containing a magnetic stir bar was added compound **13** (399 mg, 2 mmol), H₂N-PEG₂-COO*t*-Bu (550 mg, 2.4 mmol), 4 Å molecular sieves (activated powder, 200 mg), and 1,2-dichloroethane (5 mL). The mixture was stirred for 1 h at room temperature. To the reaction mixture was added sodium triacetoxymethylborohydride (845 mg, 4 mmol). The mixture was stirred for 3 days at room temperature. The resulting mixture was partitioned between EtOAc and saturated aqueous NaHCO₃. The organic layer was washed with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure to afford 850 mg of compound **14** as a viscous oil.

[00545] MS (ESI) *m/z*: [M+H]⁺ Calcd for C₂₁H₄₁N₂O₆ 417.3; Found 417.2.

13-(1-(*tert*-butoxycarbonyl)piperidin-4-yl)-2,2-dimethyl-4,14-dioxo-3,7,10-trioxa-13-azaheptadecan-17-oic acid (RED-195**).**

[00546] To a dried scintillation vial containing a magnetic stir bar was added compound **14** (220 mg, 0.5 mmol), succinic anhydride (55 mg, 0.55 mmol), 4-(dimethylamino)pyridine (5 mg, 0.04 mmol), and dichloromethane (3 mL). The mixture was stirred for 24 h at room temperature. The reaction mixture was partially purified by flash chromatography (elute 50-100% EtOAc/hexanes) to yield 117 mg of compound **RED-195** as a clear oil, which was carried forward without further characterization.

[00547] MS (ESI) *m/z*: [M+H]⁺ Calcd for C₂₅H₄₅N₂O₉ 517.6; Found 517.5.

17-(*tert*-butyl) 1-((1⁴S,1⁶S,3³S,2R,4S,10E,12E,14R)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl) (2S)-8-(1-(*tert*-butoxycarbonyl)piperidin-4-yl)-2,3-dimethyl-4,7-dioxo-11,14-dioxa-3,8-diazaheptadecanedioate (RED-196**).**

[00548] To a dried scintillation vial containing a magnetic stir bar was added **RED-195** (445 mg, 0.86 mmol), HATU (320 mg, 0.84 mmol), DIPEA (311 mg, 2.42 mmol), and dichloromethane (6 mL). The reaction mixture was stirred at room temperature for 5 minutes. The resulting solution was added to **RED-062** (516 mg, 0.79 mmol) and the reaction mixture was stirred for an additional 30 minutes at room temperature. The reaction mixture was directly

purified by flash chromatography (elute 3-10% MeOH/DCM) to give 820 mg (90%) of **RED-196** as a light tan solid.

[00549] MS (ESI) m/z: [M+H]⁺ Calcd for C₅₇H₈₇ClN₅O₁₇ 1148.6; Found 1148.8.

(2S)-1-(((1⁴S,1⁶S,3³S,2R,4S,10E,12E,14R)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl)oxy)-2,3-dimethyl-1,4,7-trioxo-8-(piperidin-4-yl)-11,14-dioxa-3,8-diazaheptadecan-17-oic acid (RED-197).

[00550] To a dried scintillation vial containing a magnetic stir bar was added **RED-196** (31 mg, 0.027 mmol) and dichloromethane (1 mL). The solution was cooled to 0 °C and tin(IV) tetrachloride (1.0 M solution in dichloromethane, 0.3 mL, 0.3 mmol) was added. The reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was directly purified by C18 flash chromatography (elute 5-100% MeCN/water) to yield 16 mg (60%) of **RED-197** as a white solid (16 mg, 60% yield).

[00551] MS (ESI) m/z: [M+H]⁺ Calcd for C₄₈H₇₁ClN₅O₁₅ 992.5; Found 992.6.

(2S)-8-(1-(3-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)-1,2-dimethylhydrazinyl)methyl)-1H-indol-1-yl)propanoyl)piperidin-4-yl)-1-(((1⁴S,1⁶S,3³S,2R,4S,10E,12E,14R)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-benzenacyclotetradecaphane-10,12-dien-4-yl)oxy)-2,3-dimethyl-1,4,7-trioxo-11,14-dioxa-3,8-diazaheptadecan-17-oic acid (RED-198).

[00552] To a dried scintillation vial containing a magnetic stir bar was added **RED-197** (16 mg, 0.016 mmol), (9H-fluoren-9-yl)methyl 1,2-dimethyl-2-((1-(3-oxo-3-(perfluorophenoxy)propyl)-1H-indol-2-yl)methyl)hydrazine-1-carboxylate (**12**) (13 mg, 0.02 mmol), DIPEA (8 µL, 0.05 mmol), and DMF (1 mL). The solution was stirred for 18 h at room temperature. The reaction mixture was directly purified by C18 flash chromatography (elute 5-100% MeCN/water) to yield 18 mg (77%) of **RED-198** as a white solid.

[00553] MS (ESI) m/z: [M+H]⁺ Calcd for C₇₇H₉₈ClN₈O₁₈ 1457.7; Found 1457.9.

(2S)-1-(((1⁴S,1⁶S,3³S,2R,4S,10E,12E,14R)-8⁶-chloro-1⁴-hydroxy-8⁵,14-dimethoxy-3³,2,7,10-tetramethyl-1²,6-dioxo-7-aza-1(6,4)-oxazinana-3(2,3)-oxirana-8(1,3)-

benzenacyclotetradecaphane-10,12-dien-4-yl)oxy)-8-(1-(3-(2-((1,2-dimethylhydrazinyl)methyl)-1H-indol-1-yl)propanoyl)piperidin-4-yl)-2,3-dimethyl-1,4,7-trioxo-11,14-dioxa-3,8-diazaheptadecan-17-oic acid (RED-106).

[00554] To a dried scintillation vial containing a magnetic stir bar was added **RED-197** (18 mg, 0.012 mmol), piperidine (20 μ L, 0.02 mmol), and DMF (1 mL). The solution was stirred for 20 minutes at room temperature. The reaction mixture was directly purified by C18 flash chromatography (elute 1-60% MeCN/water) to yield 15 mg (98%) of compound **RED-106** as a white solid.

[00555] MS (ESI) m/z: $[M+H]^+$ Calcd for $C_{62}H_{88}ClN_8O_{16}$ 1235.6; Found 1236.0.

Results and Discussion

Production and initial characterization of anti-CD22 ADC

[00556] The anti-CD22 antibody that was used (CAT-02) was a humanized variant of the RFB4 antibody. C-terminally tagged anti-CD22 antibody was made using a GPEX® clonal cell line with bioreactor titers of 1.6 g/L and 97% conversion of cysteine to formylglycine. The HIPS-4AP-maytansine linker payload was synthesized (described above) and conjugated to the aldehyde-tagged antibody. The resulting ADC was characterized (FIG. 11) by size exclusion chromatography to assess percent monomer (99.2%), and by hydrophobic interaction (HIC) and reversed-phase (PLRP) chromatography to assess the drug-to-antibody ratio (DAR), which was 1.8. The ADC was compared to the wild-type (untagged) anti-CD22 antibody in terms of affinity for human CD22 protein and internalization on CD22+ cells using an ELISA-based method (FIG. 12) and a flow cytometric-based method (FIG. 13), respectively. For both functional measures, the ADC performed equally well as the wild-type antibody, indicating that conjugation had no effect on these parameters.

The anti-CD22 ADC was not a substrate for MDR1 and does not promote off-target or bystander killing

[00557] Potency of the anti-CD22 ADC was tested *in vitro* against the Ramos and WSU-DLCL2 HNL tumor cell lines. Activity was compared to that of free maytansine and a related ADC made with the CAT-02 anti-CD22 antibody conjugated to maytansine through a cleavable valine-citrulline dipeptide linker. Both ADCs showed subnanomolar activity against wild-type

Ramos and WSU-DLCL2 cells (FIG. 14, panel A and panel C). In variants of those cells engineered to express the xenobiotic efflux pump, MDR1, only the anti-CD22 ADC of the present disclosure retained its original potency (FIG. 14, panel B and panel D). By contrast, free maytansine was ~10-fold less efficacious, and the ADC bearing cleavable maytansine was essentially devoid of activity. In a control experiment, cotreatment of WSU-DLCL2 cells with cyclosporin, an MDR1 inhibitor, had no effect on wild-type cells but restored the original potency of free maytansine and the cleavable ADC in MDR1+ cells (FIG. 14, panel E and panel F). Together, these results indicated that the active metabolite of the anti-CD22 ADC of the present disclosure was not a substrate for MDR1 efflux. In related *in vitro* cytotoxicity studies, the anti-CD22 ADC of the present disclosure had no effect on the antigen-negative cell line, NCI-N87 (FIG. 15), indicating that it had no off-target activity over a 5-day cell culture period. Furthermore, an anti-HER2-based ADC conjugated to the HIPS-4AP-maytansine linker payload did not mediate bystander killing of antigen-negative cells in coculture with antigen-positive cells (FIG. 16), implying that the active metabolite of the anti-CD22 ADC of the present disclosure, which would be the same as that of the anti-HER2 ADC conjugate, would also not mediate bystander killing.

The anti-CD22 ADC was efficacious against NHL xenograft models

[00558] The *in vivo* efficacy of the anti-CD22 ADC was assessed against the WSU-DLCL2 and Ramos xenograft models (FIG. 17), which expressed relatively higher and lower amounts of CD22, respectively (FIG. 18). In a single dose study, mice bearing WSU-DLCL2 tumors were given 10 mg/kg of the anti-CD22 ADC or a vehicle control. Dosing was initiated when the tumors averaged 118 mm³. Of the animals that received the ADC, 25% (2 of 8) had a partial response, with tumors that had regressed to 4 mm³ by day 31. The anti-CD22 ADC-treated and vehicle control groups had mean tumor volumes of 415 and 1783 mm³, respectively, by day 31. Next, in a multidose study, mice bearing WSU-DLCL2 xenografts were treated with 10 mg/kg of the anti-CD22 ADC or a vehicle control every four days for a total of four doses. Dosing was initiated when the tumors averaged 262 mm³. Of the animals that received the ADC, 75% (6 of 8) showed a complete response, with 38% of these (3 of 8) durable to the end of the study (day 59), 43 days after the last dose. By contrast, the vehicle control group reached a mean tumor volume of 2191 mm³ by day 17. Finally, in a multidose study, mice bearing Ramos

xenografts were treated with either 5 or 10 mg/kg of the anti-CD22 ADC or a vehicle control every four days for a total of four doses. Dosing was initiated when the tumors averaged 246 mm³. As anticipated, a dose effect was observed with the groups receiving the 5 or 10 mg/kg dose demonstrating 63% or 87% tumor growth delay, respectively. Specifically, the median times to endpoint were 12, 19, and 22 days for the vehicle control, 5-, and 10 mg/kg dosing groups, respectively. In all three studies, no effect was observed on mouse body weight in the anti-CD22 ADC dosing groups (FIG. 19).

The anti-CD22 ADC was well tolerated at up to 60 mg/kg in rats and cynomolgus monkeys [00559]

The anti-CD22 ADC did not bind to rodent CD22, however, dosing the ADC in these animals provided information related to off-target toxicity and safety of the linker-payload. As mentioned above, in mouse xenograft studies no effect of dosing was observed on body weight or clinical observations. In an exploratory rat toxicity study (FIG. 20), animals (5 per group) were given a single intravenous dose of the anti-CD22 ADC at 6, 20, 40, or 60 mg/kg and observed for 12 days post-dose. All animals survived until the end of the study. Animals dosed at 60 mg/kg experienced a 10% decrease in body weight relative to the vehicle control group. Clinical chemistry changes compatible with minimal to mild hepatobiliary injury occurred on Day 5 in animals given ≥ 40 mg/kg and included increased activities of alanine aminotransferase (ALT), aspartate transaminase (AST), and alkaline phosphatase (ALP). Most changes had reversed by Day 12. With respect to hematology, moderately to markedly decreased platelet counts occurred on Day 5 in animals given ≥ 40 mg/kg and had completely reversed by Day 12. Changes compatible with inflammation occurred on Days 5 and 12 in animals given ≥ 40 mg/kg and included slightly to moderately increased neutrophil and monocyte counts, slightly increased globulin concentrations, and decreased albumin:globulin ratio.

[00560] The anti-CD22 ADC did bind to cynomolgus CD22 (FIG. 21) and had a similar tissue cross-reactivity profile in monkeys as compared to humans (FIG. 22). Therefore, cynomolgus monkeys represented an appropriate model in which to test both the on-target and off-target toxicities of this ADC. In an exploratory repeat dose study, monkeys (2/sex/group) were given 10, 30, or 60 mg/kg of the anti-CD22 ADC once every three weeks for a total of two doses followed by a 21 day observation period. All animals survived until study termination. No anti-CD22 ADC-related changes in clinical observations, body weights, or food consumption

occurred. Clinical pathology changes occurred mostly in animals given ≥ 30 mg/kg, and were consistent with minimal liver injury, increased platelet consumption and/or sequestration, and inflammation (FIG. 23). These changes were similar at 30 and 60 mg/kg and after the first and second dose, and were of a magnitude that would not be expected to be associated with microscopic changes or clinical effects. Changes compatible with minimal liver injury in animals given ≥ 30 mg/kg consisted of increased ALT, AST, and ALP activities that had partially reversed by days 21 and 42. Slightly to moderately decreased platelet counts observed within a week of dosing had mostly reversed by days 21 and 42. Changes compatible with inflammation consisted of minimally to moderately increased neutrophil and monocyte counts, slightly to moderately increased globulin concentrations, and minimally decreased albumin concentrations.

Administration of the anti-CD22 ADC led to B-cell depletion in cynomolgus monkeys

[00561] In order to assess the pharmacodynamic effects of the anti-CD22 ADC in a cross-reactive species, peripheral blood mononuclear cell populations was monitored in samples taken from cynomolgus monkeys enrolled in the repeat dose toxicity study. Specifically, flow cytometry was used to detect the ratio of B cells (CD20+), T cells (CD3+), and NK cells (CD20-/CD3-) observed in animals pre-dose and at days 7, 14, 28, and 35 (Figure 5). In pre-dose anti-CD22 ADC-treated animals, B cells included an average of 11.6% of total lymphocytes; this value dropped to an average of 3.8% by day 35, representing an average decrease of 68% in the measured B cell populations relative to baseline levels (FIG. 24). B cell depletion was similar across all dosing groups, from 10 to 60 mg/kg, indicating that the lowest dose was sufficient to obtain the effect. Meanwhile, B cells in vehicle control-treated animals, and T cells and NK cells (not shown) in all groups were largely unchanged over the course of the treatment. The results indicated that the anti-CD22 ADC was able to selectively mediate the depletion of cynomolgus CD22+ cells in vivo without leading to adverse off-target toxicities.

Pharmaco- and toxicokinetics of the anti-CD22 ADC in mice, rats, and cynomolgus monkeys

[00562] In order to evaluate the in vivo stability of the anti-CD22 ADC, a pharmacokinetic (PK) study in rats was conducted. The concentrations of total antibody, total ADC, and total conjugate was monitored in the peripheral blood of animals (3/group) for 21 days

after receiving a single 3 mg/kg dose of the anti-CD22 ADC (Table 2 and FIG. 25). As shown in FIG. 10, the total ADC and total conjugate assays employed DAR-sensitive and DAR-insensitive measurements, respectively. The PK parameters obtained for all three analytes were similar, indicating that the conjugate was largely stable in circulation. For example, the elimination half-lives of total antibody, total ADC, and total conjugate were 9.48, 6.13, and 7.22 days, respectively.

[00563] Next, the anti-CD22 ADC analyte concentrations was measured over time in the peripheral blood of mice from the Ramos multidose efficacy study described above. The purpose of this analysis was to determine the total ADC exposure level achieved at an efficacious dose in xenograft studies (FIG. 26). For this benchmark, recall that 10 mg/kg x 4 doses over 22 days led to an 87% tumor growth delay in the Ramos model, and that 10 mg/kg x 4 doses over 28 days led to 75% of the animals exhibiting a complete response (no palpable tumor remaining) in the WSU-DLCL2 model. The mean area under the concentration versus time curve from time 0 to infinity (AUC_{0-inf}) for the 10 mg/kg x 4 dose in the mouse was 2530 ± 131 (S.D.) day $\cdot \mu\text{g/mL}$.

[00564] Finally, the anti-CD22 ADC analyte concentrations in toxicokinetic plasma samples from animals dosed in the previously described rat and cynomolgus monkey toxicity studies was assessed (FIG. 26). The purpose of these analyses was to determine the total ADC exposure levels achieved at doses correlated to the presence or absence of observed toxicities. With respect to the rat study, the C_{max} and AUC_{0-inf} values were generally proportional to the dose. The mean AUC_{0-inf} for the 60 mg/kg dose was 5201 ± 273 day $\cdot \mu\text{g/mL}$. With respect to the monkey study, the C_{max} and AUC_{0-inf} values were generally proportional to the dose. The mean AUC_{0-inf} for the first 60 mg/kg dose was 6140 ± 667 day $\cdot \mu\text{g/mL}$. The antibody bound to antigen in the cynomolgus model, however, clearance (not shown) was similar among all dosing groups. This indicated that the low (10 mg/kg) dose was sufficient to saturate target-mediated clearance mechanisms, and therefore that antigen-mediated clearance did not significantly affect the results of this study. This observation was consistent with the pharmacodynamic effect of the anti-CD22 ADC treatment on B-cell depletion, the extent of which was similar across all dosing groups.

EXAMPLE 6**Mantle Cell Lymphoma: Tumor Regression Following R-CHOP Treatment in Granta-519 Xenograft Model**

[00565] Granta-519 cells (0.5×10^7) were implanted into the right flanks of CB17-SCID mice. The average tumor volume 180 mm³ eleven days post implantation. The mice were randomized into 8 groups of 8 mice per group. The ADC, rituximab, and R-CHOP were tested for tumor growth inhibition against a vehicle control. All mice were injected intravenously (i.v.) with the ADC on either a once weekly or a once every three-week schedule for a period of 6 weeks, as shown in Table 4 below. R-CHOP was administered intravenously in a single dose: rituximab 30 mg/kg; cyclophosphamide 30 mg/kg, doxorubicin 2.475 mg/kg; and vincristine 0.365 mg/kg. Prednisone (0.15 mg/kg) was orally administered once per day.

Table 4.

Gr.	n	Treatment	Dosage (mg/kg)	Route	Schedule
1	8	Vehicle	-	i.v.	QW x 3
2	8	Rituximab	30mg/kg	i.v.	QW x 4
3	8	ADC	1mg/kg	i.v.	QW x 4
4	8	ADC	3mg/kg	i.v.	QW x 6
5	8	ADC	10mg/kg	i.v.	QW x 6
6	8	ADC	10mg/kg	i.v.	Q3W x 2
7	8	R- CHOP (Day 0 dosing)	R: 30mg/kg Cyclo: 30mg/kg Dox: 2.475mg/kg Vincrist:0.365mg/kg Predn: 0.15mg/kg	i.v. i.v. i.v. i.v. p.o.	Single dose Single dose Single dose Single dose QD x5
8	8	R-CHOP (D0) + ADC (D14)	R: 30mg/kg Cyclo: 30mg/kg Dox: 2.475mg/kg Vincrist:0.365mg/kg Predn: 0.15mg/kg ADC,10mg/kg	i.v. i.v. i.v. i.v. p.o. i.v.	Single dose Single dose Single dose Single dose QD x5 QW from D14

[00566] The ADC dosed at 1 mg/kg, 3 mg/kg, and 10 mg/kg on a once weekly schedule resulted in 19%, 59.7%, and 87.3% tumor growth inhibition (TGI), respectively, which without wishing to be bound by theory suggests dose-dependent anti-tumor activity. See FIG. 27. It was also observed that the anti-tumor activity was substantially greater on a once per week schedule (87.3% TGI) versus a once per three-week schedule (70.5% TGI). Treatment with R-CHOP (days 0-5) resulted in transient tumor growth inhibition. The tumors rebounded, and by day 14, the tumors were equal to the tumor volume recorded prior to treatment. On day 14, the rebounded tumors were treated with the ADC (10 mg/kg) once per week. The tumor showed significant growth inhibition by day 38 ($P < 0.001$). See FIG. 28.

[00567] The ADC inhibited Granta-519 tumor growth in mice in a dose-dependent and dose-frequency dependent manner without affecting body weight. In addition, the ADC resumed tumor growth inhibition following R-CHOP escape.

Conclusions


[00568] A CD22-targeted ADC site-specifically conjugated to a maytansine payload that was resistant to efflux by MDR1-expressing cells was produced. The ADC had a DAR of 1.8, displayed good biophysical characteristics, and mediated efficacy ranging from significant (87%) tumor growth delay to complete response *in vivo* against two NHL xenograft models. This efficacy was achieved at exposure levels well below those associated with toxicity; indeed, in the repeat dose cynomolgus toxicity study, no observed adverse effects were noted even at the highest dose of 60 mg/kg, indicating that higher doses may be used. The anti-CD22 ADC had a combination of efficacy and safety. As an added advantage, a number of the underlying components, including the target antigen, parental antibody, and the maytansine-based cytotoxic payload have been used in humans and have been well-studied regarding safety and toxicity. Based on the cynomolgus monkey, which is a reasonable model for projecting human pharmacokinetic and toxicity profiles, the results of these studies indicated that the anti-CD22 ADC is of therapeutic use for NHL patients, such as those who have developed refractory disease due to the upregulation of MDR1.

[00569] While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope

of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

WHAT IS CLAIMED IS:

1. A method for treating a cancer in a subject, the method comprising:
administering to the subject in need thereof a therapeutically effective amount of:
one or more anti-cancer agents, and
a conjugate, the conjugate comprising:
at least one modified amino acid residue with a side chain of formula (I):



wherein:

Z is CR⁴ or N;

R¹ is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R² and R³ are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R² and R³ are optionally cyclically linked to form a 5 or 6-membered heterocyclyl;

each R⁴ is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted

aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

L is a linker comprising $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d-$, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4;

T^1 , T^2 , T^3 and T^4 are each independently selected from (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(EDA)_w$, $(PEG)_n$, $(AA)_p$, $-(CR^{13}OH)_h-$, piperidin-4-amino (4AP), an acetal group, a hydrazine, a disulfide, and an ester, wherein EDA is an ethylene diamine moiety, PEG is a polyethylene glycol or a modified polyethylene glycol, and AA is an amino acid residue, wherein w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12;

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$ and $-P(O)OH-$, wherein q is an integer from 1 to 6;

each R^{13} is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl;

each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

W^1 is a maytansinoid; and

W^2 is an anti-CD22 antibody;

wherein the administering is effective to treat the cancer in the subject.


2. The method of claim 1, wherein:

T^1 is selected from a (C_1-C_{12}) alkyl and a substituted (C_1-C_{12}) alkyl;


T^2 , T^3 and T^4 are each independently selected from $(EDA)_w$, $(PEG)_n$, (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(AA)_p$, $-(CR^{13}OH)_h-$, 4-amino-piperidine (4AP), an acetal group, a hydrazine, and an ester; and

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, -CO-, -NR¹⁵-, -NR¹⁵(CH₂)_q-, -NR¹⁵(C₆H₄)-, -CONR¹⁵-, -NR¹⁵CO-, -C(O)O-, -OC(O)-, -O-, -S-, -S(O)-, -SO₂-, -SO₂NR¹⁵-, -NR¹⁵SO₂-, and -P(O)OH-;


wherein:

(PEG)_n is , where n is an integer from 1 to 30;

EDA is an ethylene diamine moiety having the following structure:



where y is an integer from 1 to 6 and r is 0 or 1;

4-amino-piperidine (4AP) is ;

each R¹² and R¹⁵ is independently selected from hydrogen, an alkyl, a substituted alkyl, a polyethylene glycol moiety, an aryl and a substituted aryl, wherein any two adjacent R¹² groups may be cyclically linked to form a piperazinyl ring; and

R¹³ is selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl.


3. The method of claim 1, wherein T¹, T², T³ and T⁴, and V¹, V², V³ and V⁴ are selected from the following table:


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	-	-	-	-


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	(EDA) _w	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-SO ₂ -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(PEG) _n	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(CR ¹³ OH) _h	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	substituted (C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-SO ₂ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-	-	-


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-	(CR ¹³ OH) _h	-CONR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-P(O)OH-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	-	-

4. The method of claim 1, wherein the linker, L, is selected from one of the following structures:









wherein

each f is independently 0 or an integer from 1 to 12;

each y is independently 0 or an integer from 1 to 20;

each n is independently 0 or an integer from 1 to 30;


each p is independently 0 or an integer from 1 to 20;

each h is independently 0 or an integer from 1 to 12;

each R is independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

each R' is independently H, a sidechain group of an amino acid, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.


5. The method of claim 1, wherein the maytansinoid is of the formula:



where ~~~ indicates the point of attachment between the maytansinoid and L.

6. The method of claim 1, wherein T^1 is (C_1-C_{12}) alkyl, V^1 is $-CO-$, T^2 is 4AP, V^2 is $-CO-$, T^3 is (C_1-C_{12}) alkyl, V^3 is $-CO-$, T^4 is absent and V^4 is absent.

7. The method of claim 1, wherein the linker, L, comprises the following structure:




wherein

each f is independently an integer from 1 to 12; and
 n is an integer from 1 to 30.

8. The method of claim 1, wherein the anti-CD22 antibody binds an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C.

9. The method of claim 1, wherein the anti-CD22 antibody comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid.

10. The method of claim 9, wherein the sequence is **L(FGly')TPSR** (SEQ ID NO: 185).

11. The method of claim 9, wherein


Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

12. The method of claim 1, wherein the modified amino acid residue is positioned at a C-terminus of a heavy chain constant region of the anti-CD22 antibody.

13. The method of claim 12, wherein the heavy chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X^1 is present; and

X^2 and X^3 are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SLSLSPG** (SEQ ID NO: 186).

14. The method of claim 13, wherein the heavy chain constant region comprises the sequence **SPGSL(FGly')TPSRGS** (SEQ ID NO: 184).

15. The method of claim 13, wherein


Z^{30} is selected from R, K, H, A, G, L, V, I, and P;

X^1 is selected from L, M, S, and V; and

X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

16. The method of claim 1, wherein the modified amino acid residue is positioned in a light chain constant region of the anti-CD22 antibody.

17. The method of claim 16, wherein the light chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue;

Z^{30} is a basic amino acid or an aliphatic amino acid;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X^1 is present; and

X^2 and X^3 are each independently any amino acid, and


wherein the sequence C-terminal to the sequence **KVDNAL** (SEQ ID NO: 58), and/or is N-terminal to the sequence **QSGNSQ** (SEQ ID NO: 59).

18. The method of claim 17, wherein the light chain constant region comprises the sequence **KVDNAL(FGly')TPSRQSGNSQ** (SEQ ID NO: 60).

19. The method of claim 17, wherein
 Z^{30} is selected from R, K, H, A, G, L, V, I, and P;
 X^1 is selected from L, M, S, and V; and
 X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

20. The method of claim 1, wherein the modified amino acid residue is positioned in a heavy chain CH1 region of the anti-CD22 antibody.

21. The method of claim 20, wherein the heavy chain CH1 region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue;

Z^{30} is a basic amino acid or an aliphatic amino acid;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X^1 is present; and

X^2 and X^3 are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SWNSGA** (SEQ ID NO: 61)

and/or is N-terminal to the amino acid sequence **GVHTEP** (SEQ ID NO: 62).

22. The method of claim 21, wherein the heavy chain CH1 region comprises the sequence **SWNSGAL(FGly')TPSRGVHTEP** (SEQ ID NO: 63).

23. The method of claim 21, wherein
 Z^{30} is selected from R, K, H, A, G, L, V, I, and P;
 X^1 is selected from L, M, S, and V; and

X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

24. The method of claim 1, wherein the modified amino acid residue is positioned in a heavy chain CH2 region of the anti-CD22 antibody.

25. The method of claim 1, wherein the modified amino acid residue is positioned in a heavy chain CH3 region of the anti-CD22 antibody.

26. The method of claim 1, wherein the one or more anti-cancer agents is selected from abitrexate methotrexate, brentuximab vedotin, copanlisib, copanlisib hydrochloride, chlorambucil, nelarabine, axicabtagene ciloleucel, carmustine, belinostat, bendamustine, bendamustine hydrochloride, tositumomab iodine-131, tositumomab, bleomycin, bortezomib, acalabrutinib, cyclophosphamide, cytarabine, cytarabine liposome, denileukin diftitox, cytarabine liposome, dexamethasone, doxorubicin, doxorubicin hydrochloride, methotrexate, pralatrexate, ofatumumab, obinutuzumab, ocrelizumab, ibritumomab, tiuxetan, ibrutinib, idelalisib, recombinant interferon alfa-2b, romidespsin, lenalidomide, mechlorethamine hydrochloride, plerixafor, prednisone, rituximab, rituximab and hyaluronidase human, bortezomib, vinblastine, vinblastine sulfate, vincristine, vincristine sulfate, and vorinostat.

27. The method of claim 1, wherein the one or more anti-cancer agents is selected from a Bruton's tyrosine kinase inhibitor, an anti-CD30 antibody-drug conjugate, a PI3K inhibitor, a DNA alkylating agent, a DNA synthesis inhibitor, a histone deacetylase inhibitor, an anti-CD20 monoclonal antibody, a proteasome inhibitor, a DNA polymerase inhibitor, an RNA polymerase inhibitor, an interleukin-2 inhibitor, a corticosteroid, a topoisomerase II inhibitor, dihydrofolate reductase inhibitor, an anti-CD20 antibody-drug conjugate, an anti-CD20 antibody-radioactive drug conjugate, a P110 δ inhibitor, an ubiquitin E3 ligase inhibitor, a chemokine CXCR4 receptor inhibitor, a tubulin inhibitor, and an agent for adoptive cell transfer therapy.

28. The method of claim 1, wherein the one or more anti-cancer agents is selected from cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone ("CHOP"); cyclophosphamide, vincristine sulfate, procarbazine hydrochloride, and prednisone ("COPP");

cyclophosphamide, vincristine sulfate, and prednisone (“CVP”); etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

29. The method of claim 1, wherein the one or more anti-cancer agents comprises rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”).

30. The method of claim 1, wherein the cancer is afflicted with dysregulation of BCR signaling.

31. The method of claim 1, wherein the cancer is afflicted with dysregulation of BCR signaling, and the cancer is responsive to B-cell depletion.

32. The method of claim 1, wherein the cancer is lymphoma.

33. The method of claim 1, wherein the cancer is a B-cell lymphoma.


34. The method of claim 1, wherein the cancer is selected from Burkitt’s lymphoma, diffuse large B-cell lymphoma, Hodgkin’s lymphoma, and non-Hodgkin’s lymphoma.

35. The method of claim 1, wherein the cancer is non-Hodgkin’s lymphoma.

36. The method of claim 1, wherein the cancer is selected from marginal zone lymphoma, mantle cell lymphoma, follicular lymphoma, and primary central nervous system lymphoma.

etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

45. A method for treating a resistant cancer in a subject, the method comprising:
administering to the subject in need thereof a therapeutically effective amount of:
one or more anti-cancer agents, and
a conjugate, the conjugate comprising:
at least one modified amino acid residue with a side chain of formula (I):



wherein:

Z is $\mathbb{C}R^4$ or $\mathbb{N}$;

R¹ is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R² and R³ are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted

aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R^2 and R^3 are optionally cyclically linked to form a 5 or 6-membered heterocyclyl;

each R^4 is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

L is a linker comprising $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d-$, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4;

T^1 , T^2 , T^3 and T^4 are each independently selected from (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(EDA)_w$, $(PEG)_n$, $(AA)_p$, $-(CR^{13}OH)_h-$, piperidin-4-amino (4AP), an acetal group, a hydrazine, a disulfide, and an ester, wherein EDA is an ethylene diamine moiety, PEG is a polyethylene glycol or a modified polyethylene glycol, and AA is an amino acid residue, wherein w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12;

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$ and $-P(O)OH-$, wherein q is an integer from 1 to 6;

each R^{13} is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl;

each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

W^1 is a maytansinoid; and

W^2 is an anti-CD22 antibody;

wherein the administering is effective to treat the resistant cancer in the subject.


46. The method of claim 45, wherein:


T^1 is selected from a (C₁-C₁₂)alkyl and a substituted (C₁-C₁₂)alkyl;

T^2 , T^3 and T^4 are each independently selected from (EDA)_w, (PEG)_n, (C₁-C₁₂)alkyl, substituted (C₁-C₁₂)alkyl, (AA)_p, -(CR¹³OH)_h-, 4-amino-piperidine (4AP), an acetal group, a hydrazine, and an ester; and

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, -CO-, -NR¹⁵-, -NR¹⁵(CH₂)_q-, -NR¹⁵(C₆H₄)-, -CONR¹⁵-, -NR¹⁵CO-, -C(O)O-, -OC(O)-, -O-, -S-, -S(O)-, -SO₂-, -SO₂NR¹⁵-, -NR¹⁵SO₂-, and -P(O)OH-;


wherein:

(PEG)_n is , where n is an integer from 1 to 30;



EDA is an ethylene diamine moiety having the following structure:

where y is an integer from 1 to 6 and r is 0 or 1;

4-amino-piperidine (4AP) is ;

each R¹² and R¹⁵ is independently selected from hydrogen, an alkyl, a substituted alkyl, a polyethylene glycol moiety, an aryl and a substituted aryl, wherein any two adjacent R¹² groups may be cyclically linked to form a piperazinyl ring; and

R¹³ is selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl.


47. The method of claim 45, wherein T^1 , T^2 , T^3 and T^4 , and V^1 , V^2 , V^3 and V^4 are selected from the following table:


T^1	V^1	T^2	V^2	T^3	V^3	T^4	V^4
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	-	-


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-CO-	(EDA) _w	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-SO ₂ -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CONR ¹⁵ -	(PEG) _n	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(CR ¹³ OH) _h	-CO-	-	-	-	-

T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	substituted (C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-SO ₂ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-	(CR ¹³ OH) _h	-CONR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-P(O)OH -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	-	-

48. The method of claim 45, wherein the linker, L, is selected from one of the following structures:







wherein

each f is independently 0 or an integer from 1 to 12;

each y is independently 0 or an integer from 1 to 20;

each n is independently 0 or an integer from 1 to 30;


each p is independently 0 or an integer from 1 to 20;

each h is independently 0 or an integer from 1 to 12;

each R is independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

each R' is independently H, a sidechain group of an amino acid, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino,

49. The method of claim 45, wherein the maytansinoid is of the formula:




50. The method of claim 45, wherein T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is 4AP, V² is -CO-, T³ is (C₁-C₁₂)alkyl, V³ is -CO-, T⁴ is absent and V⁴ is absent.

each f is independently an integer from 1 to 12; and
 n is an integer from 1 to 30.

52. The method of claim 45, wherein the anti-CD22 antibody binds an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C.

53. The method of claim 45, wherein the anti-CD22 antibody comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid.

54. The method of claim 53, wherein the sequence is **L(FGly')TPSR** (SEQ ID NO: 185).

55. The method of claim 53, wherein


Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

56. The method of claim 45, wherein the modified amino acid residue is positioned at a C-terminus of a heavy chain constant region of the anti-CD22 antibody.

57. The method of claim 56, wherein the heavy chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SLSLSPG** (SEQ ID NO: 186).

58. The method of claim 57, wherein the heavy chain constant region comprises the sequence **SPGSL(FGly')TPSRGS** (SEQ ID NO: 184).

59. The method of claim 57, wherein


Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

60. The method of claim 45, wherein the modified amino acid residue is positioned in a light chain constant region of the anti-CD22 antibody.

61. The method of claim 60, wherein the light chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, and


wherein the sequence C-terminal to the sequence **KVDNAL** (SEQ ID NO: 58), and/or is N-terminal to the sequence **QSGNSQ** (SEQ ID NO: 59).

62. The method of claim 61, wherein the light chain constant region comprises the sequence **KVDNAL(FGly')TPSRQSGNSQ** (SEQ ID NO: 60).

63. The method of claim 61, wherein
 Z^{30} is selected from R, K, H, A, G, L, V, I, and P;
 X^1 is selected from L, M, S, and V; and
 X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

64. The method of claim 45, wherein the modified amino acid residue is positioned in a heavy chain CH1 region of the anti-CD22 antibody.

65. The method of claim 64, wherein the heavy chain CH1 region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue;

Z^{30} is a basic amino acid or an aliphatic amino acid;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X^1 is present; and

X^2 and X^3 are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SWNSGA** (SEQ ID NO: 61) and/or is N-terminal to the amino acid sequence **GVHTEP** (SEQ ID NO: 62).

66. The method of claim 65, wherein the heavy chain CH1 region comprises the sequence **SWNSGAL(FGly')TPSRGVHTEP** (SEQ ID NO: 63).

67. The method of claim 65, wherein

Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

68. The method of claim 45, wherein the modified amino acid residue is positioned in a heavy chain CH2 region of the anti-CD22 antibody.

69. The method of claim 45, wherein the modified amino acid residue is positioned in a heavy chain CH3 region of the anti-CD22 antibody.

70. The method of claim 45, wherein the one or more anti-cancer agents is selected from abitrexate, methotrexate, brentuximab vedotin, copanlisib, copanlisib hydrochloride, chlorambucil, nelarabine, axicabtagene ciloleucel, carmustine, belinostat, bendamustine, bendamustine hydrochloride, tositumomab, iodine-131, bleomycin, bortezomib, acalabrutinib, cyclophosphamide, cytarabine, cytarabine liposome, denileukin diftitox, cytarabine liposome, dexamethasone, doxorubicin, doxorubicin hydrochloride, methotrexate, pralatrexate, ofatumamb, obinutuzumab, ocrelizumab, ibritumomab, tiuxetan, ibrutinib, idelalisib, recombinant interferon alfa-2b, romidespsin, lenalidomide, mechlorethamine hydrochloride, plerixafor, prednisone, rituximab, rituximab and hyaluronidase human, bortezomib, vinblastine, vinblastine sulfate, vincristine, vincristine sulfate, and vorinostat.

71. The method of claim 45, wherein the one or more anti-cancer agents is selected from a Bruton's tyrosine kinase inhibitor, an anti-CD30 antibody-drug conjugate, a PI3K inhibitor, a DNA alkylating agent, a DNA synthesis inhibitor, a histone deacetylase inhibitor, an anti-CD20 monoclonal antibody, a proteasome inhibitor, a DNA polymerase inhibitor, an RNA polymerase inhibitor, an interleukin-2 inhibitor, a corticosteroid, a topoisomerase II inhibitor, dihydrofolate reductase inhibitor, an anti-CD20 antibody-drug conjugate, an anti-CD20 antibody-radioactive drug conjugate, a P110δ inhibitor, an ubiquitin E3 ligase inhibitor, a chemokine CXCR4 receptor inhibitor, a tubulin inhibitor, and an agent for adoptive cell transfer therapy.

72. The method of claim 45, wherein the one or more anti-cancer agents is selected from cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“CHOP”); cyclophosphamide, vincristine sulfate, procarbazine hydrochloride, and prednisone (“COPP”); cyclophosphamide, vincristine sulfate, and prednisone (“CVP”); etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

73. The method of claim 45, wherein the one or more anti-cancer agents comprises rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”).

74. The method of claim 45, wherein the resistant cancer is affiliated with dysregulation of BCR signaling.

75. The method of claim 45, wherein the resistant cancer is affiliated with dysregulation of BCR signaling, and the cancer is responsive to B-cell depletion.

76. The method of claim 45, wherein the resistant cancer is lymphoma.

77. The method of claim 45, wherein the resistant cancer is a B-cell lymphoma.

78. The method of claim 45, wherein the resistant cancer is selected from Burkitt’s lymphoma, diffuse large B-cell lymphoma, Hodgkin’s lymphoma, and non-Hodgkin’s lymphoma.


79. The method of claim 45, wherein the resistant cancer is non-Hodgkin's lymphoma.
80. The method of claim 45, wherein the resistant cancer is selected from marginal zone lymphoma, mantle cell lymphoma, follicular lymphoma, and primary central nervous system lymphoma.
81. The method of claim 45, wherein the resistant cancer is mantle cell lymphoma.
82. The method of claim 45, wherein the resistant cancer is diffuse large B-cell lymphoma.
83. The method of claim 45, wherein the resistant cancer is follicular lymphoma.
84. The method of claim 45, wherein the resistant cancer is marginal zone lymphoma.
85. The method of claim 45, wherein the resistant cancer is leukemia.
86. The method of claim 45, wherein the resistant cancer is selected from chronic myeloproliferative syndrome, acute myelogenous leukemia, chronic lymphocytic leukemia, small lymphocutic leukemia, hairy cell leukemia, and acute lymphoblastic leukemia.
87. The method of claim 45, wherein the cancer is resistant to treatment with one or more anti-cancer agents selected from abitrexate methotrexate, brentuximab vedotin, copanlisib, copanlisib hydrochloride, chlorambucil, nelarabine, axicabtagene ciloleuce, carmustine, belinostat, bendamustine, bendamustine hydrochloride, tosimumab, iodine-131, tosimumab, bleomycin, bortezomib, acalabrutinib, cyclophosphamide, cytarabine, cytarabine liposome, denileukin diftitox, cytarabine liposome, dexamethasone, doxorubicin, doxorubicin hydrochloride, methotrexate, pralatrexate, ofatumumab, obinituzumab, ocrelizumab, ibritumomab, tiuxetan, ibrutinib, idelalisib, recombinant interferon alfa-2b, romidespsin, lenalidomide, mechlorethamine hydrochloride, plerixafor, prednisone, rituximab, rituximab and

hyaluronidase human, bortezomib, vinblastine, vinblastine sulfate, vincristine, vincristine sulfate, and vorinostat.

88. The method of claim 45, wherein the cancer is resistant to treatment with one or more anti-cancer agents selected from: cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“CHOP”); cyclophosphamide, vincristine sulfate, procarbazine hydrochloride, and prednisone (“COPP”); cyclophosphamide, vincristine sulfate, and prednisone (“CVP”); etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

89. The method of claim 45, wherein the cancer is resistant to treatment with rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”).

90. A method for sensitizing a cancer in a subject, the method comprising:
administering to the subject in need thereof a therapeutically effective amount of:
one or more anti-cancer agents, and
a conjugate, the conjugate comprising:
at least one modified amino acid residue with a side chain of formula (I):



wherein:

Z is CR⁴ or N;

R¹ is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R² and R³ are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R² and R³ are optionally cyclically linked to form a 5 or 6-membered heterocyclyl;

each R⁴ is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

L is a linker comprising $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d$, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4;

T¹, T², T³ and T⁴ are each independently selected from (C₁-C₁₂)alkyl, substituted (C₁-C₁₂)alkyl, (EDA)_w, (PEG)_n, (AA)_p, -(CR¹³OH)_h, piperidin-4-amino (4AP), an acetal group, a hydrazine, a disulfide, and an ester, wherein EDA is an ethylene diamine moiety, PEG is a polyethylene glycol or a modified polyethylene glycol, and AA is an amino acid residue, wherein w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12;

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-\text{CO}-$, $-\text{NR}^{15}-$, $-\text{NR}^{15}(\text{CH}_2)_q-$, $-\text{NR}^{15}(\text{C}_6\text{H}_4)-$, $-\text{CONR}^{15}-$, $-\text{NR}^{15}\text{CO}-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{SO}_2-$, $-\text{SO}_2\text{NR}^{15}-$, $-\text{NR}^{15}\text{SO}_2-$ and $-\text{P}(\text{O})\text{OH}-$, wherein q is an integer from 1 to 6;

each R^{13} is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl;

each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

W^1 is a maytansinoid; and

W^2 is an anti-CD22 antibody;

wherein the administering is effective to sensitize a cancer in the subject.


91. The method of claim 90, wherein:

T^1 is selected from a $(\text{C}_1\text{-C}_{12})$ alkyl and a substituted $(\text{C}_1\text{-C}_{12})$ alkyl;

T^2 , T^3 and T^4 are each independently selected from $(\text{EDA})_w$, $(\text{PEG})_n$, $(\text{C}_1\text{-C}_{12})$ alkyl, substituted $(\text{C}_1\text{-C}_{12})$ alkyl, $(\text{AA})_p$, $-(\text{CR}^{13}\text{OH})_h-$, 4-amino-piperidine (4AP), an acetal group, a hydrazine, and an ester; and


V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-\text{CO}-$, $-\text{NR}^{15}-$, $-\text{NR}^{15}(\text{CH}_2)_q-$, $-\text{NR}^{15}(\text{C}_6\text{H}_4)-$, $-\text{CONR}^{15}-$, $-\text{NR}^{15}\text{CO}-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{S}(\text{O})-$, $-\text{SO}_2-$, $-\text{SO}_2\text{NR}^{15}-$, $-\text{NR}^{15}\text{SO}_2-$, and $-\text{P}(\text{O})\text{OH}-$;


wherein:

$(\text{PEG})_n$ is , where n is an integer from 1 to 30;

EDA is an ethylene diamine moiety having the following structure:

where y is an integer from 1 to 6 and r is 0 or 1;

4-amino-piperidine (4AP) is ;



each R^{12} and R^{15} is independently selected from hydrogen, an alkyl, a substituted alkyl, a polyethylene glycol moiety, an aryl and a substituted aryl, wherein any two adjacent R^{12} groups may be cyclically linked to form a piperazinyl ring; and

R^{13} is selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl.


92. The method of claim 90, wherein T^1 , T^2 , T^3 and T^4 , and V^1 , V^2 , V^3 and V^4 are selected from the following table:


T^1	V^1	T^2	V^2	T^3	V^3	T^4	V^4
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	(EDA) _w	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-	-	-


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-SO ₂ -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(PEG) _n	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(CR ¹³ OH) _h	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	substituted (C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-SO ₂ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-	(CR ¹³ OH) _h	- CONR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	- NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	- P(O)OH -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	(C ₁ -C ₁₂)alkyl	- NR ¹⁵ -

T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	-	-

93. The method of claim 90, wherein the linker, L, is selected from one of the following structures:







wherein

each f is independently 0 or an integer from 1 to 12;

each y is independently 0 or an integer from 1 to 20;

each n is independently 0 or an integer from 1 to 30;

each p is independently 0 or an integer from 1 to 20;


each h is independently 0 or an integer from 1 to 12;

each R is independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

each R' is independently H, a sidechain group of an amino acid, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino,

substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.


94. The method of claim 90, wherein the maytansinoid is of the formula:



where ~ indicates the point of attachment between the maytansinoid and L.

95. The method of claim 90, wherein T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is 4AP, V² is -CO-, T³ is (C₁-C₁₂)alkyl, V³ is -CO-, T⁴ is absent and V⁴ is absent.

96. The method of claim 90, wherein the linker, L, comprises the following structure:




wherein

each f is independently an integer from 1 to 12; and
n is an integer from 1 to 30.

97. The method of claim 90, wherein the anti-CD22 antibody binds an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C.

98. The method of claim 90, wherein the anti-CD22 antibody comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid.

99. The method of claim 98, wherein the sequence is **L(FGly')TPSR** (SEQ ID NO: 185).

100. The method of claim 99, wherein


Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

101. The method of claim 90, wherein the modified amino acid residue is positioned at a C-terminus of a heavy chain constant region of the anti-CD22 antibody.

102. The method of claim 101, wherein the heavy chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SLSLSPG** (SEQ ID NO: 186).

103. The method of claim 102, wherein the heavy chain constant region comprises the sequence **SPGSL(FGly')TPSRGS** (SEQ ID NO: 184).

104. The method of claim 102, wherein


Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

105. The method of claim 90, wherein the modified amino acid residue is positioned in a light chain constant region of the anti-CD22 antibody.

106. The method of claim 105, wherein the light chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, and


wherein the sequence C-terminal to the sequence **KVDNAL** (SEQ ID NO: 58), and/or is N-terminal to the sequence **QSGNSQ** (SEQ ID NO: 59).

107. The method of claim 106, wherein the light chain constant region comprises the sequence **KVDNAL(FGly')TPSRQSGNSQ** (SEQ ID NO: 60).

108. The method of claim 106, wherein
 Z^{30} is selected from R, K, H, A, G, L, V, I, and P;
 X^1 is selected from L, M, S, and V; and
 X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

109. The method of claim 90, wherein the modified amino acid residue is positioned in a heavy chain CH1 region of the anti-CD22 antibody.

110. The method of claim 109, wherein the heavy chain CH1 region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue;

Z^{30} is a basic amino acid or an aliphatic amino acid;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X^1 is present; and

X^2 and X^3 are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SWNSGA** (SEQ ID NO: 61) and/or is N-terminal to the amino acid sequence **GVHTEP** (SEQ ID NO: 62).

111. The method of claim 110, wherein the heavy chain CH1 region comprises the sequence **SWNSGAL(FGly')TPSRGVHTEP** (SEQ ID NO: 63).

112. The method of claim 110, wherein

Z^{30} is selected from R, K, H, A, G, L, V, I, and P;

X^1 is selected from L, M, S, and V; and

X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

113. The method of claim 90, wherein the modified amino acid residue is positioned in a heavy chain CH2 region of the anti-CD22 antibody.

114. The method of claim 90, wherein the modified amino acid residue is positioned in a heavy chain CH3 region of the anti-CD22 antibody.

115. The method of claim 90, wherein the one or more anti-cancer agents is selected from abitrexate methotrexate, brentuximab vedotin, copanlisib, copanlisib hydrochloride, chlorambucil, nelarabine, axicabtagene ciloleucel, carmustine, belinostat, bendamustine, bendamustine hydrochloride, tositumomab, iodine-131 tositumomab, bleomycin, bortezomib, acalabrutinib, cyclophosphamide, cytarabine, cytarabine liposome, denileukin diftitox, cytarabine liposome, dexamethasone, doxorubicin, doxorubicin hydrochloride, methotrexate, pralatrexate, ofatumamb, obinutuzumab, ocrelizumab, ibritumomab, tiuxetan, ibrutinib, idelalisib, recombinant interferon alfa-2b, romidespsin, lenalidomide, mechlorethamine hydrochloride, plerixafor, prednisone, rituximab, rituximab and hyaluronidase human, bortezomib, vinblastine, vinblastine sulfate, vincristine, vincristine sulfate, and vorinostat.

116. The method of claim 90, wherein the one or more anti-cancer agents is selected from a Bruton's tyrosine kinase inhibitor, an anti-CD30 antibody-drug conjugate, a PI3K inhibitor, a DNA alkylating agent, a DNA synthesis inhibitor, a histone deacetylase inhibitor, an anti-CD20 monoclonal antibody, a proteasome inhibitor, a DNA polymerase inhibitor, an RNA polymerase inhibitor, an interleukin-2 inhibitor, a corticosteroid, a topoisomerase II inhibitor, dihydrofolate reductase inhibitor, an anti-CD20 antibody-drug conjugate, an anti-CD20 antibody-radioactive drug conjugate, a P110 δ inhibitor, an ubiquitin E3 ligase inhibitor, a chemokine CXCR4 receptor inhibitor, a tubulin inhibitor, and an agent for adoptive cell transfer therapy.

117. The method of claim 90, wherein the one or more anti-cancer agents is selected from cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“CHOP”); cyclophosphamide, vincristine sulfate, procarbazine hydrochloride, and prednisone (“COPP”); cyclophosphamide, vincristine sulfate, and prednisone (“CVP”); etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

118. The method of claim 90, wherein the one or more anti-cancer agents comprises rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”).

119. The method of claim 90, wherein the cancer is affiliated with dysregulation of BCR signaling.

120. The method of claim 90, wherein the cancer is affiliated with dysregulation of BCR signaling, and the cancer is responsive to B-cell depletion.

121. The method of claim 90, wherein the cancer is lymphoma.

122. The method of claim 90, wherein the cancer is a B-cell lymphoma.

123. The method of claim 90, wherein the cancer is selected from Burkitt’s lymphoma, diffuse large B-cell lymphoma, Hodgkin’s lymphoma, and non-Hodgkin’s lymphoma.

124. The method of claim 90, wherein the cancer is non-Hodgkin’s lymphoma.

- 209


133. The method of claim 90, wherein the cancer is resistant to treatment with one or more anti-cancer agents selected from: cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“CHOP”); cyclophosphamide, vincristine sulfate, procarbazine hydrochloride, and prednisone (“COPP”); cyclophosphamide, vincristine sulfate, and prednisone (“CVP”); etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

134. The method of claim 90, wherein the cancer is resistant to treatment with rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”).

135. A pharmaceutical composition for treating a cancer, the pharmaceutical composition comprising:

- one or more anti-cancer agents;
- a conjugate; and
- a pharmaceutically acceptable excipient,

wherein the conjugate comprises at least one modified amino acid residue with a side chain of formula (I):



wherein:

Z is CR⁴ or N;

R^1 is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R^2 and R^3 are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R^2 and R^3 are optionally cyclically linked to form a 5 or 6-membered heterocyclyl;

each R^4 is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

L is a linker comprising $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d-$, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4; T^1 , T^2 , T^3 and T^4 are each independently selected from (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(EDA)_w$, $(PEG)_n$, $(AA)_p$, $-(CR^{13}OH)_h-$, piperidin-4-amino (4AP), an acetal group, a hydrazine, a disulfide, and an ester, wherein EDA is an ethylene diamine moiety, PEG is a polyethylene glycol or a modified polyethylene glycol, and AA is an amino acid residue, wherein w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12;

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$ and $-P(O)OH-$, wherein q is an integer from 1 to 6;

each R^{13} is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl;

each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

W^1 is a maytansinoid; and

W^2 is an anti-CD22 antibody.


136. The pharmaceutical composition of claim 135, wherein:

T^1 is selected from a (C_1-C_{12}) alkyl and a substituted (C_1-C_{12}) alkyl;

T^2 , T^3 and T^4 are each independently selected from $(EDA)_w$, $(PEG)_n$, (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(AA)_p$, $-(CR^{13}OH)_h$, 4-amino-piperidine (4AP), an acetal group, a hydrazine, and an ester; and


V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$, and $-P(O)OH-$;

wherein:

$(PEG)_n$ is , where n is an integer from 1 to 30;


EDA is an ethylene diamine moiety having the following structure:

where y is an integer from 1 to 6 and r is 0 or 1;

4-amino-piperidine (4AP) is ;

each R^{12} and R^{15} is independently selected from hydrogen, an alkyl, a substituted alkyl, a polyethylene glycol moiety, an aryl and a substituted aryl, wherein any two adjacent R^{12} groups may be cyclically linked to form a piperazinyl ring; and

R^{13} is selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl.





137. The pharmaceutical composition of claim 135, wherein T¹, T², T³ and T⁴, and V¹, V², V³ and V⁴ are selected from the following table:


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	(EDA) _w	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-SO ₂ -	(AA) _p	-

T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(PEG) _n	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(CR ¹³ OH) _h	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	substituted (C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-SO ₂ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-	(CR ¹³ OH) _h	- CONR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	- NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	- P(O)OH -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	(C ₁ -C ₁₂)alkyl	- NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	-	-

138. The pharmaceutical composition of claim 135, wherein the linker, L, is selected from one of the following structures:







wherein

each f is independently 0 or an integer from 1 to 12;

each y is independently 0 or an integer from 1 to 20;

each n is independently 0 or an integer from 1 to 30;

each p is independently 0 or an integer from 1 to 20;


each h is independently 0 or an integer from 1 to 12;

each R is independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

each R' is independently H, a sidechain group of an amino acid, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino,

substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.


139. The pharmaceutical composition of claim 135, wherein the maytansinoid is of the formula:



where ~ indicates the point of attachment between the maytansinoid and L.

140. The pharmaceutical composition of claim 135, wherein T^1 is (C_1-C_{12}) alkyl, V^1 is $-CO-$, T^2 is 4AP, V^2 is $-CO-$, T^3 is (C_1-C_{12}) alkyl, V^3 is $-CO-$, T^4 is absent and V^4 is absent.

141. The pharmaceutical composition of claim 135, wherein the linker, L, comprises the following structure:




wherein

each f is independently an integer from 1 to 12; and
 n is an integer from 1 to 30.

142. The pharmaceutical composition of claim 135, wherein the anti-CD22 antibody binds an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C.

143. The pharmaceutical composition of claim 135, wherein the anti-CD22 antibody comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue;

Z^{30} is a basic amino acid or an aliphatic amino acid;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X^1 is present; and

X^2 and X^3 are each independently any amino acid.

144. The pharmaceutical composition of claim 143, wherein the sequence is L(FGly')TPSR.

145. The pharmaceutical composition of claim 143, wherein


Z^{30} is selected from R, K, H, A, G, L, V, I, and P;

X^1 is selected from L, M, S, and V; and

X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

146. The pharmaceutical composition of claim 135, wherein the modified amino acid residue is positioned at a C-terminus of a heavy chain constant region of the anti-CD22 antibody.

147. The pharmaceutical composition of claim 146, wherein the heavy chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SLSLSPG** (SEQ ID NO: 186).

148. The pharmaceutical composition of claim 147, wherein the heavy chain constant region comprises the sequence **SPGSL(FGly')TPSRGS** (SEQ ID NO: 184) .

149. The pharmaceutical composition of claim 147, wherein


Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

150. The pharmaceutical composition of claim 135, wherein the modified amino acid residue is positioned in a light chain constant region of the anti-CD22 antibody.

151. The pharmaceutical composition of claim 150, wherein the light chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, and

wherein the sequence C-terminal to the sequence **KVDNAL** (SEQ ID NO: 58), and/or is N-terminal to the sequence **QSGNSQ** (SEQ ID NO: 59).

152. The pharmaceutical composition of claim 151, wherein the light chain constant region comprises the sequence **KVDNAL(FGly')TPSRQSGNSQ** (SEQ ID NO: 60).

153. The pharmaceutical composition of claim 151, wherein


Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

154. The pharmaceutical composition of claim 135, wherein the modified amino acid residue is positioned in a heavy chain CH1 region of the anti-CD22 antibody.

155. The pharmaceutical composition of claim 154, wherein the heavy chain CH1 region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SWNSGA** (SEQ ID NO: 61) and/or is N-terminal to the amino acid sequence **GVHTEP** (SEQ ID NO: 62).

156. The pharmaceutical composition of claim 155, wherein the heavy chain CH1 region comprises the sequence **SWNSGAL(FGly')TPSRGVHTFP** (SEQ ID NO: 63).

157. The pharmaceutical composition of claim 155, wherein
 Z^{30} is selected from R, K, H, A, G, L, V, I, and P;
 X^1 is selected from L, M, S, and V; and
 X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

158. The pharmaceutical composition of claim 135, wherein the modified amino acid residue is positioned in a heavy chain CH2 region of the anti-CD22 antibody.

159. The pharmaceutical composition of claim 135, wherein the modified amino acid residue is positioned in a heavy chain CH3 region of the anti-CD22 antibody.

160. The pharmaceutical composition of claim 135, wherein the one or more anti-cancer agents is selected from abitrexate methotrexate, brentuximab vedotin, copanlisib, copanlisib hydrochloride, chlorambucil, nelarabine, axicabtagene ciloleucel, carmustine, belinostat, bendamustine, bendamustine hydrochloride, tositumomab, iodine-131 tositumomab, bleomycin, bortezomib, acalabrutinib, cyclophosphamide, cytarabine, cytarabine liposome, denileukin diftitox, cytarabine liposome, dexamethasone, doxorubicin, doxorubicin hydrochloride, methotrexate, pralatrexate, ofatumamb, obinutuzumab, ocrelizumab, ibritumomab, tiuxetan, ibrutinib, idelalisib, recombinant interferon alfa-2b, romidespsin, lenalidomide, mechlorethamine hydrochloride, plerixafor, prednisone, rituximab, rituximab and hyaluronidase human, bortezomib, vinblastine, vinblastine sulfate, vincristine, vincristine sulfate, and vorinostat.

161. The pharmaceutical composition of claim 135, wherein the one or more anti-cancer agents is selected from a Bruton's tyrosine kinase inhibitor, an anti-CD30 antibody-drug conjugate, a PI3K inhibitor, a DNA alkylating agent, a DNA synthesis inhibitor, a histone deacetylase inhibitor, an anti-CD20 monoclonal antibody, a proteasome inhibitor, a DNA polymerase inhibitor, an RNA polymerase inhibitor, an interleukin-2 inhibitor, a corticosteroid, a

topoisomerase II inhibitor, dihydrofolate reductase inhibitor, an anti-CD20 antibody-drug conjugate, an anti-CD20 antibody-radioactive drug conjugate, a P110 δ inhibitor, an ubiquitin E3 ligase inhibitor, a chemokine CXCR4 receptor inhibitor, a tubulin inhibitor, and an agent for adoptive cell transfer therapy.

162. The pharmaceutical composition of claim 135, wherein the one or more anti-cancer agents is selected from cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“CHOP”); cyclophosphamide, vincristine sulfate, procarbazine hydrochloride, and prednisone (“COPP”); cyclophosphamide, vincristine sulfate, and prednisone (“CVP”); etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).

163. The pharmaceutical composition of claim 135, wherein the one or more anti-cancer agents comprises rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”).

164. Use of a pharmaceutical composition for the manufacture of a medicament for treating a cancer, the pharmaceutical composition comprising a pharmaceutical composition of any of claims 135-163.

165. Use of a pharmaceutical composition for the manufacture of a medicament for treating a resistant cancer, the pharmaceutical composition comprising a pharmaceutical composition of any of claims 135-163.


166. Use of a pharmaceutical composition for the manufacture of a medicament for sensitizing a cancer, the pharmaceutical composition comprising a pharmaceutical composition of any of claims 135-163.

167. Use of a pharmaceutical composition for treating a cancer, the pharmaceutical composition comprising a pharmaceutical composition of any of claims 135-163.

168. Use of a pharmaceutical composition for treating a resistant cancer, the pharmaceutical composition comprising a pharmaceutical composition of any of claims 135-163.

169. Use of a pharmaceutical composition for sensitizing a cancer, the pharmaceutical composition comprising a pharmaceutical composition of any of claims 135-163.

170. A method for treating a resistant cancer in a subject, the method comprising:
administering to the subject in need thereof a therapeutically effective amount of a conjugate,
the conjugate comprising:
at least one modified amino acid residue with a side chain of formula (I):



wherein:

Z is CR⁴ or N;

R¹ is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R² and R³ are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl

ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R^2 and R^3 are optionally cyclically linked to form a 5 or 6-membered heterocyclyl;

each R^4 is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

L is a linker comprising $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d-$, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4;

T^1 , T^2 , T^3 and T^4 are each independently selected from (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(EDA)_w$, $(PEG)_n$, $(AA)_p$, $-(CR^{13}OH)_h-$, piperidin-4-amino (4AP), an acetal group, a hydrazine, a disulfide, and an ester, wherein EDA is an ethylene diamine moiety, PEG is a polyethylene glycol or a modified polyethylene glycol, and AA is an amino acid residue, wherein w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12;

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$ and $-P(O)OH-$, wherein q is an integer from 1 to 6;

each R^{13} is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl;

each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

W¹ is a maytansinoid; and

W² is an anti-CD22 antibody;

wherein the administering is effective to treat the resistant cancer in the subject.


171. The method of claim 170, wherein:

T¹ is selected from a (C₁-C₁₂)alkyl and a substituted (C₁-C₁₂)alkyl;

T², T³ and T⁴ are each independently selected from (EDA)_w, (PEG)_n, (C₁-C₁₂)alkyl, substituted (C₁-C₁₂)alkyl, (AA)_p, -(CR¹³OH)_h-, 4-amino-piperidine (4AP), an acetal group, a hydrazine, and an ester; and


V¹, V², V³ and V⁴ are each independently selected from the group consisting of a covalent bond, -CO-, -NR¹⁵-, -NR¹⁵(CH₂)_q-, -NR¹⁵(C₆H₄)-, -CONR¹⁵-, -NR¹⁵CO-, -C(O)O-, -OC(O)-, -O-, -S-, -S(O)-, -SO₂-, -SO₂NR¹⁵-, -NR¹⁵SO₂-, and -P(O)OH-;

wherein:


(PEG)_n is , where n is an integer from 1 to 30;

EDA is an ethylene diamine moiety having the following structure:

where y is an integer from 1 to 6 and r is 0 or 1;



4-amino-piperidine (4AP) is



each R¹² and R¹⁵ is independently selected from hydrogen, an alkyl, a substituted alkyl, a polyethylene glycol moiety, an aryl and a substituted aryl, wherein any two adjacent R¹² groups may be cyclically linked to form a piperazinyl ring; and


R¹³ is selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl.

172. The method of claim 170, wherein T¹, T², T³ and T⁴, and V¹, V², V³ and V⁴ are selected from the following table:


T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-NR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	(EDA) _w	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	-	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-SO ₂ -	(AA) _p	-

T ¹	V ¹	T ²	V ²	T ³	V ³	T ⁴	V ⁴
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-CO-	(CR ¹³ OH) _h	- CONR ¹⁵ -	(PEG) _n	-CO-
(C ₁ -C ₁₂)alkyl	-CO-	(CR ¹³ OH) _h	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	substituted (C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	(PEG) _n	-CO-	-	-
(C ₁ -C ₁₂)alkyl	-SO ₂ -	(C ₁ -C ₁₂)alkyl	-CO-	-	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-	(CR ¹³ OH) _h	- CONR ¹⁵ -	-	-
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	-CO-	(AA) _p	- NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-NR ¹⁵ -	(PEG) _n	- P(O)OH -	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	(EDA) _w	-	(AA) _p	-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	-	-
(C ₁ -C ₁₂)alkyl	- CONR ¹⁵ -	(C ₁ -C ₁₂)alkyl	-NR ¹⁵ -	-	-CO-	(C ₁ -C ₁₂)alkyl	- NR ¹⁵ -
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	(AA) _p	-
(C ₁ -C ₁₂)alkyl	-CO-	4AP	-CO-	(C ₁ -C ₁₂)alkyl	-CO-	-	-

173. The method of claim 170, wherein the linker, L, is selected from one of the following structures:







wherein

each f is independently 0 or an integer from 1 to 12;

each y is independently 0 or an integer from 1 to 20;

each n is independently 0 or an integer from 1 to 30;

each p is independently 0 or an integer from 1 to 20;


each h is independently 0 or an integer from 1 to 12;

each R is independently hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl; and

each R' is independently H, a sidechain group of an amino acid, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino,

substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl.


174. The method of claim 170, wherein the maytansinoid is of the formula:



where ~ indicates the point of attachment between the maytansinoid and L.

175. The method of claim 170, wherein T¹ is (C₁-C₁₂)alkyl, V¹ is -CO-, T² is 4AP, V² is -CO-, T³ is (C₁-C₁₂)alkyl, V³ is -CO-, T⁴ is absent and V⁴ is absent.

176. The method of claim 170, wherein the linker, L, comprises the following structure:




wherein

each f is independently an integer from 1 to 12; and
n is an integer from 1 to 30.

177. The method of claim 170, wherein the anti-CD22 antibody binds an epitope within amino acids 1 to 847, within amino acids 1-759, within amino acids 1-751, or within amino acids 1-670, of a CD22 amino acid sequence depicted in FIG. 8A-8C.

178. The method of claim 170, wherein the anti-CD22 antibody comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid.

179. The method of claim 178, wherein the sequence is **L(FGly')TPSR** (SEQ ID NO: 185).

180. The method of claim 178, wherein


Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

181. The method of claim 170, wherein the modified amino acid residue is positioned at a C-terminus of a heavy chain constant region of the anti-CD22 antibody.

182. The method of claim 181, wherein the heavy chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SLSLSPG** (SEQ ID NO: 186).

183. The method of claim 182, wherein the heavy chain constant region comprises the sequence **SPGSL(FGly')TPSRGS** (SEQ ID NO: 184).

184. The method of claim 182, wherein


Z³⁰ is selected from R, K, H, A, G, L, V, I, and P;

X¹ is selected from L, M, S, and V; and

X² and X³ are each independently selected from S, T, A, V, G, and C.

185. The method of claim 170, wherein the modified amino acid residue is positioned in a light chain constant region of the anti-CD22 antibody.

186. The method of claim 185, wherein the light chain constant region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z²⁰ is either a proline or alanine residue;

Z³⁰ is a basic amino acid or an aliphatic amino acid;

X¹ may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X¹ is present; and

X² and X³ are each independently any amino acid, and


wherein the sequence C-terminal to the sequence **KVDNAL** (SEQ ID NO: 58), and/or is N-terminal to the sequence **QSGNSQ** (SEQ ID NO: 59).

187. The method of claim 186, wherein the light chain constant region comprises the sequence **KVDNAL(FGly')TPSRQSGNSQ** (SEQ ID NO: 60).

188. The method of claim 186, wherein
 Z^{30} is selected from R, K, H, A, G, L, V, I, and P;
 X^1 is selected from L, M, S, and V; and
 X^2 and X^3 are each independently selected from S, T, A, V, G, and C.

189. The method of claim 170, wherein the modified amino acid residue is positioned in a heavy chain CH1 region of the anti-CD22 antibody.

190. The method of claim 189, wherein the heavy chain CH1 region comprises a sequence of the formula (II) (SEQ ID NOs: 189 – 190):



wherein

FGly' is the modified amino acid residue of formula (I);

Z^{20} is either a proline or alanine residue;

Z^{30} is a basic amino acid or an aliphatic amino acid;

X^1 may be present (SEQ ID NO: 189) or absent (SEQ ID NO: 190) and, when present, can be any amino acid, with the proviso that when the sequence is at the N-terminus of the conjugate, X^1 is present; and

X^2 and X^3 are each independently any amino acid, and

wherein the sequence is C-terminal to the amino acid sequence **SWNSGA** (SEQ ID NO: 61) and/or is N-terminal to the amino acid sequence **GVHTEP** (SEQ ID NO: 62).

191. The method of claim 190, wherein the heavy chain CH1 region comprises the sequence **SWNSGAL(FGly')TPSRGVHTEP** (SEQ ID NO: 63).

192. The method of claim 190, wherein
 Z^{30} is selected from R, K, H, A, G, L, V, I, and P;
 X^1 is selected from L, M, S, and V; and
 X^2 and X^3 are each independently selected from S, T, A, V, G, and C.
193. The method of claim 170, wherein the modified amino acid residue is positioned in a heavy chain CH2 region of the anti-CD22 antibody.
194. The method of claim 170, wherein the modified amino acid residue is positioned in a heavy chain CH3 region of the anti-CD22 antibody.
195. The method of claim 170, wherein the resistant cancer is affiliated with dysregulation of BCR signaling.
196. The method of claim 170, wherein the resistant cancer is affiliated with dysregulation of BCR signaling, and the cancer is responsive to B-cell depletion.
197. The method of claim 170, wherein the resistant cancer is lymphoma.
198. The method of claim 170, wherein the resistant cancer is a B-cell lymphoma.
199. The method of claim 170, wherein the resistant cancer is selected from Burkitt's lymphoma, diffuse large B-cell lymphoma, Hodgkin's lymphoma, and non-Hodgkin's lymphoma.
200. The method of claim 170, wherein the resistant cancer is non-Hodgkin's lymphoma.
201. The method of claim 170, wherein the resistant cancer is selected from marginal zone lymphoma, mantle cell lymphoma, follicular lymphoma, and primary central nervous system lymphoma.

- 237

doxorubicin hydrochloride (“EPOCH”); cyclophosphamide, vincristine sulfate, doxorubicin hydrochloride, and dexamethasone (“hyper-CVAD”); ifosfamide, carboplatin, and etoposide phosphate (“ICE”); rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”); rituximab, cyclophosphamide, vincristine sulfate, and prednisone (“R-CVP”); rituximab, etoposide phosphate, prednisone, vincristine sulfate, rituximab, etoposide phosphate, prednisone, vincristine sulfate, cyclophosphamide, and doxorubicin hydrochloride (“R-EPOCH”); and rituximab, ifosfamide, carboplatin, and etoposide phosphate (“R-ICE”).


210. The method of claim 170, wherein the cancer is resistant to treatment with rituximab, cyclophosphamide, doxorubicin hydrochloride, vincristine sulfate, and prednisone (“R-CHOP”).

211. Use of a combination comprising:

one or more anti-cancer agents, and

a conjugate, the conjugate comprising:

at least one modified amino acid residue with a side chain of formula (I):



wherein:

Z is CR⁴ or N;

R¹ is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R² and R³ are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted

cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R^2 and R^3 are optionally cyclically linked to form a 5 or 6-membered heterocyclyl;

each R^4 is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

L is a linker comprising $-(T^1-V^1)_a-(T^2-V^2)_b-(T^3-V^3)_c-(T^4-V^4)_d-$, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4;

T^1 , T^2 , T^3 and T^4 are each independently selected from (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(EDA)_w$, $(PEG)_n$, $(AA)_p$, $-(CR^{13}OH)_h-$, piperidin-4-amino (4AP), an acetal group, a hydrazine, a disulfide, and an ester, wherein EDA is an ethylene diamine moiety, PEG is a polyethylene glycol or a modified polyethylene glycol, and AA is an amino acid residue, wherein w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12;

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$ and $-P(O)OH-$, wherein q is an integer from 1 to 6;

each R^{13} is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl;

each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

W^1 is a maytansinoid; and

W^2 is an anti-CD22 antibody;


in the manufacture of a medicament for treating a cancer in a subject in need thereof.

212. Use of a combination comprising:

one or more anti-cancer agents, and

a conjugate, the conjugate comprising:

at least one modified amino acid residue with a side chain of formula (I):



wherein:

Z is CR⁴ or N;

R¹ is selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

R² and R³ are each independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl, or R² and R³ are optionally cyclically linked to form a 5 or 6-membered heterocyclyl;

each R⁴ is independently selected from hydrogen, halogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, amino, substituted amino, carboxyl, carboxyl ester, acyl, acyloxy, acyl amino, amino acyl, alkylamide, substituted alkylamide, sulfonyl, thioalkoxy, substituted thioalkoxy, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

L is a linker comprising -(T¹-V¹)_a-(T²-V²)_b-(T³-V³)_c-(T⁴-V⁴)_d, wherein a, b, c and d are each independently 0 or 1, where the sum of a, b, c and d is 1 to 4;

T^1 , T^2 , T^3 and T^4 are each independently selected from (C_1-C_{12}) alkyl, substituted (C_1-C_{12}) alkyl, $(EDA)_w$, $(PEG)_n$, $(AA)_p$, $-(CR^{13}OH)_h-$, piperidin-4-amino (4AP), an acetal group, a hydrazine, a disulfide, and an ester, wherein EDA is an ethylene diamine moiety, PEG is a polyethylene glycol or a modified polyethylene glycol, and AA is an amino acid residue, wherein w is an integer from 1 to 20, n is an integer from 1 to 30, p is an integer from 1 to 20, and h is an integer from 1 to 12;

V^1 , V^2 , V^3 and V^4 are each independently selected from the group consisting of a covalent bond, $-CO-$, $-NR^{15}-$, $-NR^{15}(CH_2)_q-$, $-NR^{15}(C_6H_4)-$, $-CONR^{15}-$, $-NR^{15}CO-$, $-C(O)O-$, $-OC(O)-$, $-O-$, $-S-$, $-S(O)-$, $-SO_2-$, $-SO_2NR^{15}-$, $-NR^{15}SO_2-$ and $-P(O)OH-$, wherein q is an integer from 1 to 6;

each R^{13} is independently selected from hydrogen, an alkyl, a substituted alkyl, an aryl, and a substituted aryl;

each R^{15} is independently selected from hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, carboxyl, carboxyl ester, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, and substituted heterocyclyl;

W^1 is a maytansinoid; and

W^2 is an anti-CD22 antibody;

for treating a cancer in a subject in need thereof.






FIG. 1




#	Time	Area	Height	Width	Area%	Symmetry
1	4.272	90.2	2.8	0.5446	5.511	0.46
2	6.015	348.2	8.3	0.7005	21.267	1.757
3	7.102	1199	36.5	0.548	73.223	0.809

FIG. 2




#	Time	Area	Height	Width	Area%	Symmetry
1	5.855	151.4	3.6	0.7039	22.779	1.421
2	6.965	513.1	14.7	0.5819	77.221	0.65

FIG. 3



#	Time	Area	Height	Width	Area%	Symmetry
1	3.389	30.5	5.6	0.0913	9.281	1729.18
2	3.518	298.5	23.7	0.2102	90.719	0.54

FIG. 4



#	Time	Area	Height	Width	Area%	Symmetry
1	7.408	19	4E-1	0.7939	1.791	1.355
2	8.502	1043.7	18	0.9658	98.209	0.423

FIG. 5




FIG. 6A




FIG. 6B




FIG. 7

FIG. 8A

isoform2	MHLLGPWLLLLLVLEYLAFSDSSKWVFEHPETLYAWEGACVWIPCTYRALDGDLESFILFH	60
isoform4	MHLLGPWLLLLLVLEYLAFSDSSKWVFEHPETLYAWEGACVWIPCTYRALDGDLESFILFH	60
isoform1	MHLLGPWLLLLLVLEYLAFSDSSKWVFEHPETLYAWEGACVWIPCTYRALDGDLESFILFH	60
isoform3	MHLLGPWLLLLLVLEYLAFSDSSKWVFEHPETLYAWEGACVWIPCTYRALDGDLESFILFH	60

isoform2	NPEYNKNTSKFDGTRLYESTKDGKVPSEQKRVQFLGDKNKNCCTLSIHPVHLNDSGQLGLR	120
isoform4	NPEYNKNTSKFDGTRLYESTKDGKVPSEQKRVQFLGDKNKNCCTLSIHPVHLNDSGQLGLR	120
isoform1	NPEYNKNTSKFDGTRLYESTKDGKVPSEQKRVQFLGDKNKNCCTLSIHPVHLNDSGQLGLR	120
isoform3	NPEYNKNTSKFDGTRLYESTKDGKVPSEQKRVQFLGDKNKNCCTLSIHPVHLNDSGQLGLR	120

isoform2	MESKTEKWMERIHLNVSERPPPHIQLPPEIQESQEVTLTCLLNFSCYGYPIQLQWLLEG	180
isoform4	MESKTEKWMERIHLNVSERPPPHIQLPPEIQESQEVTLTCLLNFSCYGYPIQLQWLLEG	180
isoform1	MESKTEKWMERIHLNVSERPPPHIQLPPEIQESQEVTLTCLLNFSCYGYPIQLQWLLEG	180
isoform3	MESKTEKWMERIHLNVSERPPPHIQLPPEIQESQEVTLTCLLNFSCYGYPIQLQWLLEG	180

isoform2	VPMRQAAVTSTSLTIKSVFTRSELKFSQWSHHGKIIVTCQLQDADGKFLSNDTVQLNVKH	240
isoform4	VPMRQAAVTSTSLTIKSVFTRSELKFSQWSHHGKIIVTCQLQDADGKFLSNDTVQLNVKH	240
isoform1	VPMRQAAVTSTSLTIKSVFTRSELKFSQWSHHGKIIVTCQLQDADGKFLSNDTVQLNVKH	240
isoform3	VPMRQAAVTSTSLTIKSVFTRSELKFSQWSHHGKIIVTCQLQDADGKFLSNDTVQLNVKH	240

isoform2	TPKLEIKVTPSDAIVREGDSVTMTCEVSSSNPEYTTVSWLKDGTSLKKQNTFTLNLRVLT	300
isoform4	-----	
isoform1	TPKLEIKVTPSDAIVREGDSVTMTCEVSSSNPEYTTVSWLKDGTSLKKQNTFTLNLRVLT	300
isoform3	TPKLEIKVTPSDAIVREGDSVTMTCEVSSSNPEYTTVSWLKDGTSLKKQNTFTLNLRVLT	300

isoform2	KDQSGKYCCQVNSNDVGPGRSEEVFLQVQ-----	328
isoform4	-----	
isoform1	KDQSGKYCCQVNSNDVGPGRSEEVFLQVQYAPEPSTVQIILHSPAPEGSQVEFLCMLANPL	360
isoform3	KDQSGKYCCQVNSNDVGPGRSEEVFLQVQYAPEPSTVQIILHSPAPEGSQVEFLCMLANPL	360

FIG. 8B

isoform2	-----YPPK	332
isoform4	-----PPK	243
isoform1	PTNYTWYHNGKEMQGRTEEKVHIPKILPWHAGTYSVCAENILGTGQRGPGAELDVQYPPK	420
isoform3	PTNYTWYHNGKEMQGRTEEKVHIPKILPWHAGTYSVCAENILGTGQRGPGAELDVQYPPK	420

isoform2	KVTTVIQNPMPIREGDTVTILSCNVNNSNPSTRYEWKPHGAWEEPSTGLVKIQNVGWDNT	392
isoform4	KVTTVIQNPMPIREGDTVTILSCNVNNSNPSTRYEWKPHGAWEEPSTGLVKIQNVGWDNT	303
isoform1	KVTTVIQNPMPIREGDTVTILSCNVNNSNPSTRYEWKPHGAWEEPSTGLVKIQNVGWDNT	480
isoform3	KVTTVIQNPMPIREGDTVTILSCNVNNSNPSTRYEWKPHGAWEEPSTGLVKIQNVGWDNT	480

isoform2	TIACAACNSWCWSWASPVALNVQYAPRDVVRVKIKPLSEIHSGNSVSLQCDEFSSHPKEVQ	452
isoform4	TIACAACNSWCWSWASPVALNVQYAPRDVVRVKIKPLSEIHSGNSVSLQCDEFSSHPKEVQ	363
isoform1	TIACAACNSWCWSWASPVALNVQYAPRDVVRVKIKPLSEIHSGNSVSLQCDEFSSHPKEVQ	540
isoform3	TIACAACNSWCWSWASPVALNVQYAPRDVVRVKIKPLSEIHSGNSVSLQCDEFSSHPKEVQ	540

isoform2	FFWEKNGRLLGKESQLNFDSISPEDAGSYSCWVNNSIGQTASKAWTLEVLVAPRRLRVSM	512
isoform4	FFWEKNGRLLGKESQLNFDSISPEDAGSYSCWVNNSIGQTASKAWTLEVLVAPRRLRVSM	423
isoform1	FFWEKNGRLLGKESQLNFDSISPEDAGSYSCWVNNSIGQTASKAWTLEVLVAPRRLRVSM	600
isoform3	FFWEKNGRLLGKESQLNFDSISPEDAGSYSCWVNNSIGQTASKAWTLEVLVAPRRLRVSM	600

isoform2	SPGDQVMEGKSATLTCESDANPPVSHYTWFDWNNQSLPYHSQKLRLEFPVKVQHSWAYWCQ	572
isoform4	SPGDQVMEGKSATLTCESDANPPVSHYTWFDWNNQSLPYHSQKLRLEFPVKVQHSWAYWCQ	483
isoform1	SPGDQVMEGKSATLTCESDANPPVSHYTWFDWNNQSLPYHSQKLRLEFPVKVQHSWAYWCQ	660
isoform3	SPGDQVMEGKSATLTCESDANPPVSHYTWFDWNNQSLPYHSQKLRLEFPVKVQHSWAYWCQ	660

isoform2	GTNSVGKGRSPLSTLTVYSPETIGRRVAVGLGSCIAILILAI CGLKLRQRRWKRTQSQQG	632
isoform4	GTNSVGKGRSPLSTLTVYSPETIGRRVAVGLGSCIAILILAI CGLKLRQRRWKRTQSQQG	543
isoform1	GTNSVGKGRSPLSTLTVYSPETIGRRVAVGLGSCIAILILAI CGLKLRQRRWKRTQSQQG	720
isoform3	GTNSVGKGRSPLSTLTVYSPETIGRRVAVGLGSCIAILILAI CGLKLRQRRWKRTQSQQG	720

FIG. 8C

isoform2	LQENSSGQSFVVRNKKVRRAPLSEGP HSLGCYNPMMEDGISYTTLRFPEMNI	692
isoform4	LQENSSGQSFVVRNKKVRRAPLSEGP HSLGCYNPMMEDGISYTTLRFPEMNI	603
isoform1	LQENSSGQSFVVRNKKVRRAPLSEGP HSLGCYNPMMEDGISYTTLRFPEMNI	780
isoform3	LQENSSGQSFVVRNKKRORVLR-----DAET	746
	***** *.	***;
isoform2	SEMQRPPDCDDTVTYSALHKRQVG DYENVIPDFPEDEGIHSELIQFGVG	752
isoform4	SEMQRPPDCDDTVTYSALHKRQVG DYENVIPDFPEDEGIHSELIQFGVG	663
isoform1	SEMQRPPDCDDTVTYSALHKRQVG DYENVIPDFPEDEGIHSELIQFGVG	840
isoform3	SPGLR-----	751
	* *	
isoform2	DYVILKH	759
isoform4	DYVILKH	670
isoform1	DYVILK-	846
isoform3	-----	
	Isoform2 = SEQ ID NO: 1	
	Isoform4 = SEQ ID NO: 2	
	Isoform1 = SEQ ID NO: 3	
	Isoform3 = SEQ ID NO: 4	

Light chain conserved region: SEQ ID NO: 5

140 150 160 170 180 189
 SVFI FPPSDEQLKS GTASVVCILN NFY KVKQ WKV SQESV
 200 210 220 230 236
 YSLSS TLTLKADYE KHKVYACEVT PVTK SEN

Heavy chain conserved region: SEQ ID NO: 6

130 140 150 160 170 180
 SVFP LAPSSKTSQGAALGCLVK DYF TSVS WSNALISGV HTFPAVL
 190 200 210 220 230 240
 YSLSSVVT VPSSSLGT ICNVNHKPS NTKVDKKV
 250 260 270 280 290 300
 PSVFLFPKP KDTLAEVTCVVVDV KENW YVGVVEVHN KTKPRER
 310 320 330 340 350 360
 YRVVSVLT VLHQDWLNCK EYKCKVSI IENTIS KGGPRFQ VYTLPPSRE
 370 380 390 400 410 420
 MTKNQVSLT LVKCFYPSDI AVEWESNQQ ENYKTP LSSGFFLY SKLTVDKSRW
 430 440 450
 OCGNVFSCSV MHEALHNHYT QKSL

FIG. 9A


```

IgG1      LTVLHQDWLNGKEYKCKVSNKALPAPIEKTIISKAKGQPREPQVYTLPPSRDELTKNQVS-- 368
IgG3      LTVLHQDWLNGKEYKCKVSNKALPAPIEKTIISKAKGQPREPQVYTLPPSRDEMTKNQVS--
IgG2      LTVVHQDWLNGKEYKCKVSNKGLPAPIEKTIISKTKGQPREPQVYTLPPSREEMTKNQVS--
IgG4      LTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIISKAKGQPREPQVYTLPPSQEEMTKNQVS--
IgA       LSGCAEPWNHGTKFTCTAAYPESKTPLTATLSKS--GNTFERPEVHLLPPPSEELALNELVT
*:  : *  **  :*:  :*:  :*:  :*:  :*:  :*:  :*:  :*:  :*:  :*:  :*:  :*:
IgG1      LTCLVKGFYPSDIAVEWESNGQ--PENNYKTPPPVLDSDG--SFFLYSKLTVDKSRWQQ 423
IgG3      LTCLVKGFYPSDIAVEWESGQ--PENNYNTTPPMLDSDG--SFFLYSKLTVDKSRWQQ
IgG2      LTCLVKGFYPSDISVEWESNGQ--PENNYKTPPMLDSDG--SFFLYSKLTVDKSRWQQ
IgG4      LTCLVKGFYPSDIAVEWESNGQ--PENNYKTPPPVLDSDG--SFFLYSKLTVDKSRWQE
IgA       LTCLARGFSPPKDVLRWLQGSQELPREKYLTTWASRQEPSQGTTFAVTSILRVAAEDWKK
****:*** *: *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *
IgG1      GNVFSCSVMHEALHNHYTQKSLSLSPGK----- 451
IgG3      GNVFSCSVMHEALHNRYTQKSLSLSPGK-----
IgG2      GNVFSCSVMHEALHNHYTQKSLSLSPGK-----
IgG4      GNVFSCSVMHEALHNHYTQKSLSLSPGK-----
IgA       GDTFSCMVGHEALPLAFTQKTIIDRLAGKPTHVNVSVVMAEVDGTCY
*:  *** *  ****  :***:..  **

```

FIG. 9B, continued

```

IgG1 = SEQ ID NO: 7
IgG3 = SEQ ID NO: 9
IgG2 = SEQ ID NO: 8
IgG4 = SEQ ID NO: 10
IgG  = SEQ ID NO: 11

```

```
Seq1 = Homo sapiens kappa light chain constant region; GenBank CAA75031.1
Seq2 = Homo sapiens kappa light chain constant region; GenBank BAC0168.1
Seq3 = Homo sapiens lambda light chain constant region; GenBank CAA75033
Seq4 = Mus musculus light chain constant region; GenBank AAB09710.1
Seq5 = Rattus norvegicus light chain constant region; GenBank AAD10133

seq1 RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD 189
seq2 RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD
seq4 RADAAPTVISIFFPPSSEQLTSGGASVVCFLNFFPKDINVKKIDGSERQNGVLNSWTDQD
seq5 RADAAPTVISIFFPPSMEQLTSGGATVVCVFNFFPRDISVKWKIDGSEQFDGLDSVTDQD
seq3 QPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQ
      :.. ***:* : **** * : * : * : * : * : * : * : * : * : * : * : * :
      : .. ***:* : **** * : * : * : * : * : * : * : * : * : * : * : * :

seq1 SKDSTYSLSSLTILTSKADYEKHKVYACEVTHQGSLSPVTKSFNRGEC 236
seq2 SKDSTYSLSSLTILTSKADYEKHKLYACEVTHQGSLSPVTKSFNRGEC
seq4 SKDSTYSMSSTLTITLKDEYERHNSYTCEATHKTSTSPIVKSFNRGEC
seq5 SKDSTYSMSSTLSLTKEVEYERHNLYTCEVVKHTSSSPVKSFNRNEC
seq3 S--NNKYAASSYLSTPEQWKSHKSYSCQVTHEG--STVEKTVAPTECS
      * :... * : ** ***: : : : * : * : * : * : * : * : * : *
```




FIG. 10




FIG. 11




FIG. 12




FIG. 13

- ◆ Maytansine
- ◆ Anti-CD22 ADC
- ◆ α CD22 cleavable maytansine




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 20, continued




FIG. 21




FIG. 22




FIG. 23

- ◆ Vehicle
- ◆ 10 mg/kg
- ◆ 30 mg/kg
- ◆ 60 mg/kg




FIG. 24




FIG. 25

Table 3. Summary of mean ($\pm$ SD) pharmacokinetic (PK) and toxicokinetic (TK) parameters of total ADC values in animals dosed with Anti-CD22 ADC.

Dose (mg/kg)	Mouse (q4d x 4) ^a			Rat (single dose)		Cynomolgus monkey (q3w x 2) ^b	
	C _{max} first dose (μg/mL)	AUC _{0-inf} (day·μg/mL)	C _{max} (μg/mL)	AUC _{0-inf} (day·μg/mL)	C _{max} first dose (μg/mL)	AUC _{0-inf} (day·μg/mL)	
3			83.9 (16)	218 (18.4)			
5	74.4 (5.48)	1500 (45.1)					
6			110 (38.5)	660 (143)			
10	136 (4.57)	2530 (131)			318 (110)	1360 (556)	
20			382 (63.0)	2280 (325)			
30					1030 (57.4)	4200 (768)	
40			687 (52.8)	3740 (185)			
60			1020 (158)	5201 (273)	1630 (138)	6140 (667)	

^aAUC calculation includes all doses.

^bAUC calculation from the first dose only.

SD, standard deviation; AUC_{0-inf}, area under the concentration versus time curve from time 0 to infinity; C_{max}, highest concentration observed at the first sampling time point from each study as follows: mouse and rat PK, 1 h; rat TK, 8 h; cynomolgus TK, 5 min.

FIG. 26




FIG. 27




FIG. 28

FIG. 1

