(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
29 January 2004 (29.01.2004)

PCT

(10) International Publication Number
WO 2004/009582 A1

(51) International Patent Classification[7]: C07D 333/50, 333/80, A61P 29/00

(21) International Application Number: PCT/US2003/022768

(22) International Filing Date: 18 July 2003 (18.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/397,052 19 July 2002 (19.07.2002) US

(71) Applicant *(for all designated States except US)*: **PHARMACIA CORPORATION** [US/US]; Corporate Patent Department, P.O. Box 1027, Chesterfiels, MI 63006 (US).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: **HAGEN, Timothy, J.** [US/US]; 1920 Madison Avenue, Gurnee, IL 60031 (US). **WEIER, Richard, M.** [US/US]; 240 Hickory Court, Lake Bluff, IL 60044 (US). **XU, Xiangdong** [CN/US]; 7498 Abbey Road, Gurnee, IL 60031 (US). **HOUDEK, Stephen, C.** [US/US]; 1538 Van Buren Avenue, Des Plaines, IL 60018 (US). **CLARE, Michael** [GB/US]; 5154 West Brown Street, Skokie, IL 60077 (US).

(74) Agents: **BAUER, S., Christopher** et al.; Pharmacia Corporation, Corporate Patent Department, P.O. Box 1027, Chesterfield, MO 63006 (US).

(81) Designated States *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States *(regional)*: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2004/009582 A1

(54) Title: SUBSTITUTED THIOPHENE CARBOXAMIDE COMPOUNDS FOR THE TREATMENT OF INFLAMMATION

(57) Abstract: The present invention provides substituted thiophene carboxamide compounds having structural Formula I and isomers, tautomers, polymorphs, carriers, prodrugs, and pharmaceutically acceptable salts thereof, compositions comprising such, and methods for treating diseases associated with kinase activity. More specifically, the present invention provides methods of treatment of a variety of diseases associated with IKK2 including the treatment of inflammation, other inflammation-associated disorders, such as, as an analgesic in the treatment of pain and headaches, arthritis, including but not limited to rheumatoid arthritis, asthma, gastrointestinal conditions such as inflammatory bowel disease, vascular diseases, viral infections such as AIDS, and cancer.

SUBSTITUTED THIOPHENE CARBOXAMIDE COMPOUNDS FOR THE TREATMENT OF INFLAMMATION

FIELD OF THE INVENTION

The present invention in general is in the field of anti-inflammatory pharmaceutical agents and specifically relates to substituted thiophene carboxamide derivatives, compositions comprising such, and methods for treating cancer, inflammation, and inflammation-associated disorders, such as arthritis.

BACKGROUND OF THE INVENTION

The following description of the background of the invention is provided to aid in the understanding the invention, but is not admitted to be or describe prior art to the invention.

NF-κB is a ubiquitous transcription factor that plays a prominent role in the activation of the immune system and in stress responses by regulating the transcription of many early, inducible genes including proinflammatory cytokines, adhesion molecules, growth factors, enzymes, and receptors (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342). Specificity of gene expression is determined at a cellular level by a diverse array of external stimuli such as bacterial products including LPS, as well as cytokines, most importantly tumor necrosis factor-α (TNFα) and interleukin-β (IL1β). Through the synergistic interaction with other transcription factors, further specificity can be achieved while maintaining enormous potential to coordinately induce a large number of functionally related genes. NF-κB is composed of homo and heterodimers of the Rel protein family and is sequestered in an inactive form in the cytoplasm by members of the IκB family of inhibitory proteins (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342). IκBs mask the nuclear localization signal on NF-κB, preventing

nuclear translocation and hence DNA binding to the promoter regions of responsive genes. Stimulation of cells with an agonist that activates NF-κB leads to a series of biochemical signals, ultimately resulting in the phosphorylation, ubiquitinylation, and degradation of IκBs, thereby releasing NF-κB for nuclear translocation (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342). Recently, two IκB kinases (IKK1 or IKKα and IKK2 or IKKβ), which phosphorylate IκBs and thereby initiate their degradation, have been cloned and characterized by a number of laboratories (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342). The catalytic subunits, IKK1 and IKK2, are similar structurally as well as enzymatically and exist as a heterodimer in a large protein complex referred to as the IKK signalsome (Regnier, C., Song, H., Gao, X., Goeddel, D., Cao, Z. and Rothe, M. (1997) *Cell* **90**, 373-383; DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. (1997) *Nature* **388**, 548-554; Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. and Roa, A. (1997) *Science* **278**, 860-866; Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. And Goeddel, D.V. (1997) *Science* **278**, 866-869). A third protein, NEMO (IKKγ, IKKAP1), is a regulatory adapter protein necessary for IKK activation and kinase activity (Yamaoka, S., Courtois, G., Bessia, C., Whiteside, S. T., Weil, R., Agou, F., Kirk, H. E., Kay, R. J., and Ireal, A. (1998) *Cell* **93**, 1231-1240; Rothwarf, D. M., Zandi, E., Natoli, G., Karin, M. (1998) *Nature* **395**, 297; Mercurio, F., Murray, B. W., Shevchenko, A., Bennet, B. L., Young, D. B., Li, J. W., Pascual, G., Motiwala, A., Zhu, H., Mann, M and Manning, A. M. (1999) *Mol. Cell. Biol.* **2**, 1526-1538). IKK1 and IKK2 are co-expressed in most human adult tissues as well as in different developmental stages of mouse embryos (Regnier, C., Song, H., Gao, X., Goeddel, D., Cao, Z. and Rothe, M. (1997) *Cell* **90**, 373-383; DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. (1997) *Nature* **388**, 548-554; Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. and Roa, A. (1997) *Science* **278**, 860-866; Zandi, E.

Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. and Goeddel, D.V. (1997) *Science* **278**, 866-869; Hu, M. C. T., and Wang, Y. (1998) *Gene* **222**, 31-40). This kinase complex appears to represent a critical, common denominator in the activation of NF-κB in a number of signal transduction pathways stimulated by a variety of agonists including cytokines, such as TNFα and IL1β, microbial products such as LPS and viral proteins such as TAX, as well as phorbol esters, oxidizing agents and serine/tyrosine phosphatases (Ghosh S., May, M. J., and Kopp. E (1998) *Annu. Rev. Immunol.* **16**, 115-260; Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342).

IKK1 (also termed IKKα, Regnier, C., Song, H., Gao, X., Goeddel, D., Cao, Z. and Rothe, M. (1997) *Cell* **90**, 373-383; DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. (1997) *Nature* **388**, 548-554; Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. And Roa, A. (1997) *Science* **278**, 860-866) was cloned simultaneously by standard biochemical purification of the IκB kinase activity from TNFα stimulated HeLa S3 cells and by its interaction with the MAP3K, NF-κB inducing kinase (NIK), in a yeast two-hybrid screen. IKK1 was identified as the previously cloned serine-threonine kinase, CHUK (Connelly, M. and Marcu, K. (1995) *Cell. Mol. Biol. Res.* **41**, 537-549). IKK1 (also termed IKKα) is an 85 kDa, 745 amino acid protein that contains an N-terminal serine/threonine kinase catalytic domain, a leucine zipper-like amphipathic helix, and a C-terminal helix-loop-helix domain. IKK2 (also termed IKKβ) was also cloned by standard biochemical purification, copurifying with IKK1 from TNFα stimulated HeLa S3 cells as well as by being identified in the public database from an EST clone with sequence homology to IKK1 (Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. and Roa, A. (1997) *Science* **278**, 860-866; Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. And Goeddel, D.V. (1997) *Science* **278**, 866-869). IKK2 is an 87 kDa, 756 amino acid protein with the same over all topology as IKK1 except for the addition of an 11 amino acid extension at the C-terminus. IKK1 and IKK2 are 52% identical

overall with 65% identity in the kinase domain and 44% identity in the protein interaction domains in the C-terminus. Data obtained using transient mammalian expression analysis, by *in vitro* translation experiments and by coexpression in a baculoviral system reveals that IKK1 and IKK2 associate preferentially as a heterodimer through their leucine zipper motifs. Although homodimers have also been described in these systems, the heterodimer is thought to be the physiologic form of the kinase in mammalian cells (Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Li, J., Peet, G.W., Pullen, S.S., Schembri-King, J., Warren, T.C., Marcu, K.B., Kehry, M.R., Barton, R. and Jakes, S. (1998) *J. Biol. Chem.* **273**, 30736-30741). Finally, NEMO (also termed IKKγ) contains three α-helical regions including a leucine zipper, interacts preferentially with IKK2 and is required for activation of the heterodimeric kinase complex perhaps by bringing other proteins into the signalsome complex (Yamaoka, S., Courtois, G., Bessia, C., Whiteside, S. T., Weil, R., Agou, F., Kirk, H. E., Kay, R. J., and Ireal, A. (1998) *Cell* **93**, 1231-1240; Rothwarf, D. M., Zandi, E., Natoli, G., Karin, M. (1998) *Nature* **395**, 297; Mercurio, F., Murray, B. W., Shevchenko, A., Bennet, B. L., Young, D. B., Li, J. W., Pascual, G., Motiwala, A., Zhu, H., Mann, M and Manning, A. M. (1999) *Mol. Cell. Biol.* **2**, 1526-1538).

The kinase activities of IKK1 and IKK2 are regulated by phosphorylation and require an intact leucine zipper (LZ) for dimerization as well as an intact helix-loop-helix (HLH) domain, which can exert a positive regulatory effect on kinase activity even when it is expressed in trans with the remainder of the IKK protein (Regnier, C., Song, H., Gao, X., Goeddel, D., Cao, Z. and Rothe, M. (1997) *Cell* **90**, 373-383; DiDonato, J.A., Hayakawa, M., Rothwarf, D.M., Zandi, E. and Karin, M. (1997) *Nature* **388**, 548-554; Mercurio,F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. and Roa, A. (1997) *Science* **278**, 860-866; Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. and Goeddel, D.V. (1997) *Science* **278**, 866-869; Dehase, M., Hayakawa, M., Chen, Y., and Karin, M. (1999) *Science* **284**, 309-313). Both IKK subunits contain a canonical MAPKK activation loop motif near the N- terminus which is the target for phosphorylation and activation of kinase activity by

MAP3Ks such as NIK and MEKK1, although the physiologic regulation by these two upstream kinases awaits further characterization (Zandi, E., and Karin, M. (1999) *Mol. Cell. Biol.* **19**, 4547-4551; Karin, M. (1999) *J. Biol. Chem.* **274**, 27339-27342; Karin, M., and Delhase, M. (1998) *Proc. Natl. Acad. Sci.* USA **95**, 9067-9069). Finally, phosphorylation of serines in the C-terminus of IKK2 results in a decrease in IKK activity and it is postulated to be responsible for the transient kinase activity seen after stimulation of cells with an agonist (Dehase, M., Hayakawa, M., Chen, Y., and Karin, M. (1999) *Science* **284**, 309-313).

IKK2 demonstrates a more potent kinase activity compared to IKK1 using IκBα or IκBβ as a substrate (Mercurio, F., Zhu, H., Murray, B.W., Shevchenko, A., Bennett, B.L., Li, J.W., Young, D.B., Barbosa, M., Mann, M., Manning, A. and Roa, A. (1997) *Science* **278**, 860-866; Zandi, E. Rothwarf, D.M., Delhase, M., Hayadawa, M and Karin, M. (1997) *Cell* **91**, 243-252; Woronicz, J.D., Gao, X., Cao, Z., Rothe, M. and Goeddel, D.V. (1997) *Science* **278**, 866-869; Dehase, M., Hayakawa, M., Chen, Y., and Karin, M. (1999) *Science* **284**, 309-313). Mutations of the phospho-acceptor serine residues within the MAPKK activation loop alters IKK2 kinase activity; the serine to alanine substitutions result in decreased kinase activity whereas the serine to glutamic acid substitutions result in a constitutively active kinase. Similar alanine mutations in IKK1 do not result in a decreased stimulation of total IKK activity in response to TNFα or IL1β (Dehase, M., Hayakawa, M., Chen, Y., and Karin, M. (1999) *Science* **284**, 309-313). IKK2 being the dominant kinase activity within the IKK complex is further supported by the analysis of fibroblasts from mice deficient in IKK1 or IKK2. Fibroblasts lacking IKK1 retain full IKK activity in response to cytokines and could activate NF-κB. In contrast, fibroblasts lacking IKK2 do not exhibit IKK activity when stimulated with cytokines nor do they activate NF-κB. Furthermore, the phenotypes of each IKK knock out is unique with IKK1 deficiency resulting in skin and skeletal defects and IKK2 knock out being embryonic lethal due to hepatocyte apoptosis (Li, Q., Antwerp, D. V., Mercurio, F., Lee, K., and Verma, I. M. (1999) *Science* **284**, 321-325; Takeda, K., Tekeuchi, O., Tsujimura, T., Itami, S., Adachi, O., Kawai, T., Sanjo, H., Yoshikawa, K., Terada, N, and Akira, S. (1999) *Science* **284**, 313-316; Hu, Y., Baud, V., Delhase, M., Zhang, P., Deerinck, T., Ellisman, M., Johnson, R.,

and Karin, M. (1999) *Science* **284**, 315-320; Li, Q., Lu, Q., Hwang, J. Y., Buscher, D., Lee, K., Izpisua-Belmonte, J. C., and Verma, I. M. (1999) *Gene and Development* **13**, 1322-1328; Tanaka, M., Fuentes, M. E., Yamaguchi, K., Durnin, M. H., Dalrymple, S. A., Hardy, K. L., and Goeddel, D. V. (1999) *Immunity* **10**, 421-429).

It is well-known that NF-KB plays a key role in the regulated expression of a large number of pro-inflammatory mediators including cytokines such as IL-6 and IL-8, cell adhesion molecules, such as ICAM and VCAM, and inducible nitric oxide synthase (iNOS). Such mediators are known to play a role in the recruitment of leukocytes at sites of inflammation and in the case of iNOS, may lead to organ destruction in some inflammatory and autoimmune diseases. The importance of NF-κB in inflammatory disorders is further strengthened by studies of airway inflammation including asthma in which NF-κB has been shown to be activated. This activation may underlie the increased cytokine production and leukocyte infiltration characteristic of these disorders. In addition, inhaled steroids are known to reduce airway hyperresponsiveness and suppress the inflammatory response in asthmatic airways. In light of the recent findings with regard to glucocorticoid inhibition of NF-κB, one may speculate that these effects are mediated through an inhibition of NF-κB. Further evidence for a role of NF-κB in inflammatory disorders comes from studies of rheumatoid synovium. Although NF-κB is normally present as an inactive cytoplasmic complex, recent immunohistochemical studies have indicated that NF-κB is present in the nuclei, and hence active, in the cells comprising rheumatoid synovium. Furthermore, NF-κB has been shown to be activated in human synovial cells in response to stimulation with TNF-α. Such a distribution may be the underlying mechanism for the increased cytokine and eicosanoid production characteristic of this tissue. See Roshak, A. K., et al., J. Biol. Chem., 271, 31496-31501 (1996).

The NF-κB/Rel and IκB proteins are also likely to play a key role in neoplastic transformation. Family members are associated with cell transformation in vitro and in vivo because of overexpression, gene amplification, gene rearrangements, or translocations (Gilmore TD, *Trends Genet* 7:318-322, 1991; Gillmore TD,

Oncogene **18**:6925-6937, 1999; Rayet B. et al., *Oncogene* **18**: 6938-6947, 1991). In addition, rearrangement and/or amplification of the genes encoding these proteins are seen in 20-25% of certain human lymphoid tumors. In addition, a role for NF-κB in the regulation of apoptosis, cell cycle progression, invasion, and metastasis has been reported (Bours V. et al., *Biochemical Pharmacology* **60**:1085-1090, 2000) strengthening the role of this transcription factor in the control of cell proliferation. The inhibition of NF-κB has been shown to potentiate TNF- and cancer therapy through increased apoptosis (Wang C-Y et al., *Science* **274**:784-787, 1996; Wang C-Y et al., *Nat Med* **5**:412-417, 1999). It has also been shown that human T-cell leukemia virus type 1 (HTLV1) infected cells (the etiological agent of an aggressive malignancy of activated $CD4^+$ T lymphocytes), IKKα and IKKβ are expressed constitutively, which normally function in a transient manner (Chu Z-L et al., *J of Biological Chemistry* **273**:15891-15894, 1998). The HTLV1 transforming and transactivating protein (Tax) has been shown to bind MEKK1 and increases the activity of IKKβ to enhance phosphorylation of serine residues in IκBα that lead to its degradation.


US 4,999,436 discloses aryl and heterocyclic substituted thiophene derivatives as inhibitors of 5-lipoxygenase for treating inflammation.

US 4,797,414 discloses thienobenzothiopyran derivatives as respiratory enhancing agents.

US 5,468,750 discloses heterocycle-coupled substituted pyrrolo[3,2-C]pyridin-2-carbxylic acids as inhibitors of the biological effect of oxygenated free radicals.

DETAILED DESCRIPTION OF THE INVENTION

The present invention concerns novel pharmaceutical composition comprising tricyclic compounds of the Formula I



or isomers, tautomers, polymorphs, carriers, prodrugs, pharmaceutically acceptable salts thereof, wherein;

A is $(CH_2)_m$ or $(CH_2)_m$-W-$(CH_2)_n$, wherein A is optionally substituted with one or more substituent independently selected from the group consisting of sulfamyl, halo, alkyl, alkoxy, hydroxyl and haloalkyl, CF_3, $COCF_3$, CN, NO_2, hydrido, OR^3, $OCOOR^3$, CO_2H, CO_2R^3, $CONH_2$, $CONHR^3$, $CON(R^3)_2$, COR^3, SR^3, SOR^3, $SCOOR^3$, SO_2R^3, NH_2, NHR^3, NR^3R^3, NR^3COR^3, NR^3CONHR^3, $NR^3SO_2R^3$, $NR^3SO_2NHR^3$, SO_2NHR^3, and $SO_2N(R^3)_2$;

B is a 6-membered aromatic hydrocarbon ring, optionally substituted with one or more substituent independently selected from the group consisting of OR^3, SR^4, $SO_2N(R^4)_2$, NHR^4, $NHCOR^4$, NR^4COR^4, $NHCO(OR^4)$, $NR^4CO(OR^4)$, $NHCONHR^3$, NR^3CONHR^4, $NR^3SO_2R^4$, $NHSO_2R^4$, $NHSO_2N(R^4)_2$, $NHCONR^4R^4$, CO_2R^4, $CON(R^4)_2$, aryl, heteroaryl, heterocyclic, halo, alkylthio, alkylsulfinyl, alkylsulfonyl, cyano, carboxyl, alkoxycarbonyl, amido, N-monoalkylamido, N-monoarylamido, alkyl, alkenyl, alkynyl, N,N-dialkylamido, N-alkyl-N-arylamido, haloalkyl, hydrido, hydroxyl, alkoxy, hydroxyalkyl, haloalkoxy, sulfamyl, N-alkylsulfamyl, amino, alkylamino, and nitro;

W is S(O)p, O, CH=N, N(O)=CH, CH_2= CH_2, and NR^4;
m is 0 to 3, inclusive;
n is 0 to 3, inclusive;

p is 0 to 2, inclusive;

R^1 and R^2 are independently selected from the group consisting of : hydrido, cyano, nitro, hydroxyl, alkyl, hydroxyalkyl, haloalkyl, alkoxy, haloalkoxy, amidine, guanidine, $CONHR^5$, NHR^5, and $NHCXNHR^5$, wherein X is O, S, or NR^6;

R^3 is selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, alkenyl, alkynyl, and heteroalkyl;

R^4 is selected from the group consisting of: lower alkyl, aryl, heteroaryl, arylalkyl, heteroalkyl, haloalkyl, arylalkylamino, and heteroarylalkyl, wherein aryl, arylalkyl, heteroaryl, or heteroarylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

R^5 is selected from the group consisting of: hydrido, alkyl, aminoalkyl, hydroxyalkyl, haloalkyl, and acyl;

R^6 is selected from the group consisting of: hydrido, alkyl, cyano, and nitro.

With the proviso, when W is CH=N and m is 0, the nitrogen atom in CH=N has to be directly attached to the ring B.

DEFINITIONS

The present invention includes the use of all hydrates, solvates, complexes and prodrugs of the compounds of this invention. Prodrugs are any covalently bonded compounds, which release the active parent drug according to Formula I in vivo. If a chiral center or another form of an isomeric center is present in a compound of the present invention all forms of such isomer or isomers, including enantiomers and diastereomers, are intended to be covered herein. Compounds containing a chiral

center may be used as a racemic mixture, an enantiornerically enriched mixture, or the racemic mixture may be separated using well-known techniques and an individual enantiomer may be used alone. In cases in which compounds have unsaturated carbon-carbon double bonds, both the cis (Z) and trans (E) isomers are within the scope of this invention. In cases wherein compounds may exist in tautomeric forms, such as keto-enol tautomers, each tautomeric form is contemplated as being included within this invention whether existing in equilibrium or predominantly in one form.

The meaning of any substituent at any one occurrence in Formula I or any sub-formula thereof is independent of its meaning, or any other substituents meaning, at any other occurrence, unless specified otherwise.

The term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "alkylsulfonyl"; it embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about five carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isoamyl, hexyl, octyl and the, like. The term "hydrido" denotes a single hydrogen atom (H). This hydrido radical may be attached, for example, to an oxygen atom to form a hydroxyl radical or two hydrido radicals may be attached to a carbon atom to form a methylene ($-CH_2-$) radical. The term "halo" means halogens such as fluorine, chlorine, and bromine or iodine atoms. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl, and polyhaloalkyl radicals. A monohaloalkyl radical, for one example, may have a bromo, chloro, or a fluoro atom within the radical. Dihalo radicals may have two or more of the same halo atoms or a combination of different halo radicals and polyhaloalkyl radicals may have more than two of the same halo atoms or a combination of different halo radicals. The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more hydroxylradicals. The terms "alkoxy" and

"alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms, such as methoxy radical. The term "alkoxyalkyl" also embraces alkyl radicals having two or more alkoxy radicals attached to the alkyl radical, that is, to form monoalkoxyalkyl and diaikoxyalkyl radicals. The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro, or bromo, to provide "haloalkoxy" or "haloalkoxyalkyl" radicals. Examples of "alkoxy" radicals include methoxy, butoxy, and trifluoromethoxy. The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one, two, or three rings wherein such rings may be attached together in a pendent manner or may be fused. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, tetrahydronapthyl, indane, and biphenyl. The term "heterocyclic" embraces saturated, partially saturated, and unsaturated heteroatom-containing ring-shaped radicals, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. Examples of saturated heterocyclic radicals include pyrrolidyl and morpholinyl. The term "heteroaryl" embraces unsaturated heterocyclic radicals. Examples of unsaturated heterocyclic radicals, also termed "heteroaryl" radicals include thienyl, pyrrolyl, furyl, pyridyl, pyrimidyl, pyrazinyl, pyrazolyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl, and tetrazolyl. The term also embraces radicals where heterocyclic radicals are fused with aryl radicals. Examples of such fused bicyclic radicals include benzofuran, benzothiophene, and the like. The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively divalent radicals $-SO_2-$. "Alkylsulfonyl", embraces alkyl radicals attached to a sulfonyl radical, where alkyl is defined as above. The term "arylsulfonyl" embraces sulfonyl radicals substituted with an aryl radical. The terms "sulfamyl" or "sulfonamidyl", whether alone or used with terms such as "N-alkylsulfamyl", "N-arylsulfamyl", "N,N-dialkylsulfamyl" and "N-alkyl-N-arylsulfamyl", denotes a sulfonyl radical substituted with an amine radical, forming a sulfonamide ($-SO_2NH_2$). The terms "N-alkylsulfamyl" and "N,N-dialkylsulfamyl" denote sulfamyl radicals substituted, respectively, with one alkyl radical, a cycloalkyl ring, or two alkyl radicals. The terms "N-arylsulfamyl" and "N-alkyl-N-arylsulfamyl" denote sulfamyl radicals substituted, respectively, with one aryl radical, and one alkyl and one aryl radical. The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", denotes

$-CO_2H$. The term "carboxyalkyl" embraces radicals having a carboxyradical as defined above, attached to an alkyl radical. The term "carbonyl", whether used alone or with other terms, such as "alkylcarbonyl", denotes –C(=O)–. The term "alkylcarbonyl" embraces radicals having a carbonyl radical substituted with an alkyl radical. An example of an "alkylcarbonyl" radical is CH_3–C(=O)–. The term "alkylcarbonylalkyl" denotes an alkyl radical substituted with an "alkylcarbonyl" radical. The term "alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl (C=O) radical. Examples of such "alkoxycarbonyl" radicals include $(CH_3)_3CO$-C(=O) – and –(O=)C–OCH_3. The term "alkoxycarbonylalkyl" embraces radicals having "alkoxycarbonyl", as defined above substituted to an alkyl radical. Examples of such "alkoxycarbonylalkyl" radicals include $(CH_3)_3COC(=O)$-$(CH_2)_2$– and –$(CH_2)_2$ $(O=)COCH_3$. The term "amido" when used by itself or with other terms such as "amidoalkyl", "N-monoalkylamido", "N-monoarylamido", "N,N-dialkylamido", "N-alkyl-N-arylamido", "N-alkyl-N-hydroxyamido" and "N-alkyl-N-hydroxyamidoalkyl", embraces a carbonyl radical substituted with an amino radical. The terms "N-alkylamido" and "N,N-dialkylamido" denote amido groups which have been substituted with one alkyl radical and with two alkyl radicals, respectively. The terms "N-monoarylamido" and "N-alkyl-N-arylamido" denote amido radicals substituted, respectively, with one aryl radical, and one alkyl and one aryl radical. The term "N-alkyl-N-hydroxyamido" embraces amido radicals substituted with a hydroxyl radical and with an alkyl radical. The term "N-alkyl-N-hydroxyamidoalkyl" embraces alkyl radicals substituted with an N-alkyl-N-hydroxyamido radical. The term "amidoalkyl" embraces alkyl radicals substituted with amido radicals. The term "aminoalkyl" embraces alkyl radicals substituted with amino radicals. The term "alkylaminoalkyl" embraces aminoalkyl radicals having the nitrogen atom substituted with an alkyl radical. The term "amidino" denotes an –C(=NH)–NH_2 radical. The term "cyanoamidino" denotes an –C(=N)–CN)–NH_2 radical. The term "heterocycloalkyl" embraces heterocyclic-substituted alkyl radicals such as pyridylmethyl and thienylmethyl. The term "aralkyl" embraces aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenethyl, and diphenethyl. The terms benzyl and phenylmethyl are interchangeable. The term "cycloalkyl" embraces radicals having three to ten

carbon atoms, such as cyclopropyl cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl. The term "cycloalkenyl" embraces unsaturated radicals having three to ten carbon atoms, such as cylopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, and cycloheptenyl. The term "alkylthio" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent sulfur atom. An example of "alkylthio" is methylthio, (CH_3S–). The term "alkylsulfinyl" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent –S(O)– atom. The terms "N-alkylamino" and "N,N-dialkylamino" denote amino groups which have been substituted with one alkyl radical and with two alkyl radicals, respectively. The term "acyl", whether used alone, or within a term such as "acylamino", denotes a radical provided by the residue after removal of hydroxyl from an organic acid. The term "acylamino" embraces an amino radical substituted with an acyl group. An example of an "acylamino" radical is acetylamino ($CH_3C(=O)$–NH–).

Compounds of Formula I or would be useful for, but not limited to, the treatment of inflammation in a subject, and for treatment of other inflammation-associated disorders, such as, as an analgesic in the treatment of pain and headaches, or as an antipyretic for the treatment of fever. For example, compounds of Formula I would be useful to treat arthritis, including but not limited to rheumatoid arthritis, spondylo arthopathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus, and juvenile arthritis. Such compounds of Formula I would be useful in the treatment of asthma, bronchitis, menstrual cramps, tendinitis, bursitis, and skin related conditions such as psoriasis, eczema, burns, and dermatitis. Compounds of Formula I also would be useful to treat gastrointestinal conditions such as inflammatory bowel disease, Crohn's disease, gastritis, irritable bowel syndrome, and ulcerative colitis and for the prevention of colorectal cancer. Compounds of Formula I would be useful in treating inflammation in such diseases as vascular diseases such as vascularitus, migraine headaches, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, sclerodoma, rheumatic fever, type I diabetes, myasthenia gravis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, polymyositis, gingivitis, hypersensitivity, conjunctivitis, swelling occurring after injury, myocardial ischemia, and the like. The compounds of the present invention

may also be used for pain. The compounds are useful as antiinflammatory agents, such as for the treatment of arthritis, with the additional benefit of having significantly less harmful side effects. The compounds of Formula I are useful as agents for treating cancer or anticancer agents. The compounds of Formula I may be proapoptotic, antiapoptotic, anticell cycle progressive, antiinvasive, antiproliferative, antiangiogenic, and antimetastatic. The cancer may be colon, ovarian, breast, prostate, gastric, B-cell lymphoma, and multiple myeloma. The compounds of Formula I may be used as an anitviral agent. The compounds of this invention may act as inhibitors of protein kinases. The compounds of this invention may act as inhibitors of IKK1 and/or IKK2, IKKα/IKKβ heterodimer, TBK or IKK*i*. The present invention preferably includes compounds, which selectively inhibit IKK2 over IKK1. Preferably, the compounds have an IKK2 IC50 of less than 1 μM, and have a selectivity ratio of IKK2 inhibition over IKK1 inhibition of at least 50, and more preferably of at least 100. Even more preferably, the compounds have an IKK1 IC50 of greater than 10 μM, and more preferably of greater than 100 μM. The compounds of formula may also be used to treat angiogenesis associated cardiovascular, ophthalmology and osteoporosis disorders. The compounds of the present invention may also be used for treatment of knee injury such as sport injuries.

While it is possible for an active ingredient to be administered alone as the raw chemical, it is preferable to present it as a pharmaceutical formulation. The present invention comprises a pharmaceutical composition comprising a therapeutically effective amount of a compound of the present invention in association with at least one pharmaceutically acceptable carrier, adjuvant, or diluent. The present invention also comprises a method of treating inflammation or inflammation associated disorders in a subject, the method comprising administering to the subject having such inflammation or disorders a therapeutically effective amount of a compound of the present invention. Also included in the family of compounds of the present invention are the pharmaceutically acceptable salts thereof. The term "pharmaceutically acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically acceptable. Suitable

pharmaceutically acceptable acid addition salts of compounds of the present invention may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric, and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicyclic, salicyclic, phydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, stearic, cyclohexylaminosulfonic, algenic, β-hydroxybutyric, salicyclic, galactaric and galacturonic acid. Suitable pharmaceutically acceptable base addition salts of compounds of the present invention include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methyl-glucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of the present invention by reacting, for example, the appropriate acid or base with the compound of the present invention.

Also embraced within this invention are pharmaceutical compositions comprising one or more compounds of the present invention in association with one or more non-toxic, pharmaceutically acceptable carriers and/or diluents and/or adjuvants and/or excipient (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. Accordingly, the compounds of the present invention may be used in the manufacture of a medicament. Pharmaceutical compositions of the compounds of the present invention prepared as herein before described may be formulated as solutions or lyophilized powders for parenteral administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation may be a buffered, isotonic aqueous solution. The compounds of the present invention may be administered by any suitable route, preferably in the form of a

pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The compounds and composition may, for example, be administered intravascularly, intraperitoneally, intravenously, subcutaneously, intramuscularly, intramedullary, orally, or topically. For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension, or liquid. The active ingredient may also be administered by injection as a composition wherein, for example, normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate solution may be used as a suitable carrier. Such formulation is especially suitable for parenteral administration, but may also be used for oral administration or contained in a metered dose inhaler or nebulizer for insufflation. It may be desirable to add excipients such as polyvinylpyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride, or sodium citrate. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. The amount of therapeutically active compound that is administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the sub-ject, the severity of the disease, the route and frequency of administration, and the particular compound employed, and thus may vary widely. The pharmaceutical compositions may contain active ingredient in the range of about 0.1 to 2000 mg, preferably in the range of about 0.5 to 500 mg and most preferably between about 1 and 100 mg. A daily dose of about 0.01 to 100 mg/kg bodyweight, preferably between about 0.1 and about 50 mg/kg body weight and most preferably between about 1 to 20 mg/kg bodyweight, may be appropriate. The daily dose can be administered in one to four doses per day. For therapeutic purposes, the compounds of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered orally, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanoic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated

for convenient administration. Such capsules or tablets may contain a controlled release formulation as may be provided in a dispersion of active compound in a sustained release material such as glyceryl monostearate, glyceryl distearate, hydroxypropylmethyl cellulose alone or with a wax. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. The pharmaceutical preparations are made following the conventional techniques of pharmacy involving milling, mixing, granulating, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion, or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered orally or filled into a soft gelatin capsule. For rectal administration, the compounds of the present invention may also be combined with excipients such as cocoa butter, glycerin, gelatin, or polyethylene glycols and molded into a suppository. The methods of the present invention include topical administration of the compounds of the present invention. By topical administration is meant non-systemic administration, including the application of a compound of the invention externally to the epidermis, to the buccal cavity and instillation of such a compound into the ear, eye, and nose, wherein the compound does not significantly enter the blood stream. By systemic administration is meant oral, intravenous, intraperitoneal, and intramuscular administration. The amount of a compound of the present invention (hereinafter referred to as the active ingredient) required for therapeutic or prophylactic effect upon topical administration will, of course, vary with the compound chosen, the nature and severity of the condition being treated and the animal undergoing treatment, and is ultimately at the discretion of the physician.

The topical formulations of the present invention, both for veterinary and for human medical use, comprise an active ingredient together with one or more acceptable

carriers therefore, and optionally any other therapeutic ingredients. The carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of where treatment is required such as: liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.01 to 5.0 wt% of the formulation.

Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container, which is then sealed and sterilized by autoclaving, or maintaining at 90-100° C for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.00217c), benzalkonium chloride (0.0 1%) and chlorhexidine acetate (0.0 1%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol, and propylene glycol.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil. Creams, ointments, or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy basis. The basis may comprise


hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol or macrogols. The formulation may incorporate any suitable surface-active agent such as an anionic, cationic, or non-ionic surface-active agent such as sorbitan esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as silicaceous silicas, and other ingredients such as lanolin may also be included. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art. Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

GENERAL SYNTHETIC PROCEDURES

The starting materials used herein are commercially available or are prepared by routine methods well known to those of ordinary skill in the art and can be found in standard reference books, such as the COMPENDIUM OF ORGANIC SYNTHETIC METHODS, Vol. I-VI (published by Wiley-Interscience).


The compounds of the invention can be synthesized according to the following procedures of Scheme I, wherein the R^1-R^4 substituents are as defined for Formula I, above, except where further noted.

Scheme I



a) $POCl_3$, DMF 5°C – RT, 16h b) NH_2OH x HCl, NaOAc, EtOH (90%) reflux 2h
c) Acetic anhydride, pyridine, reflux d) 2-Mercaptoacetamide, NaOMe, reflux

Scheme II




$R^3 = CONH_2$ or CN, $R^4 = CH_3$ or NH_2

a) sulfur, amine base, EtOH reflux, b) sodium cyanate in AcOH

The complete content of all publications, patents, and patent applications cited in this disclosure are herein incorporated by reference as if each individual publication, patent, or patent application were specifically and individually indicated to incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for the purposes of clarity of understanding, it will be readily apparent to one skilled in the art in light of the teachings of this invention that changes and modifications can be made without departing from the spirit and scope of the present invention. The following examples are provided for exemplification purposes only and are not intended to limit the scope of the invention, which has been described in broad terms above.

EXAMPLES

Examples 1



3-amino-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide

1-chloro-3,4-dihydronaphthalene-2-carbaldehyde 1a: In a 3 neck flask fitted with a N_2 inlet tube, thermometer, and addition funnel, was placed DMF (8.4 g) and CH_2Cl_2 (40 mL). After cooling in an ice bath to 0-5°C, $POCl_3$ (14 g) was added dropwise. After the addition was complete the reaction was stirred at RT for 2 h. The mixture was then cooled to 0-5°C and a solution of α-tetralone (8.76 g, 60 mmol) in CH_2Cl_2 (50 mL) was added. After the addition was complete the reaction was allowed to stir at RT for 16 h. The reaction solution was poured onto ice and


$NaHCO_3$ (satd, 200 mL), added an additional CH_2Cl_2 and stirred until gas evolution ceased. The layers were separated, and the aqueous layer was extracted with CH_2Cl_2 (250 mL). The CH_2Cl_2 extracts were combined, washed with water and dried (Na_2SO_4), and concentrated in vacuo to yield the title compound as a red liquid (11.7 g., 96% yield).

1-chloro-3,4-dihydronaphthalene-2-carbaldehyde oxime 1b: A mixture of **1** (11.0 g), hydroxyl amine hydrochloride (5.2 g) and sodium acetate (11.75 g) in ethanol (90 mL, 90%) was refluxed for 2 h. The reaction was cooled to RT, then poured onto ice water (90 mL). The resulting precipitate was collected by filtration to yield the title compound as a tan solid (12 g, 100% yield).

1-chloro-3,4-dihydronaphthalene-2-carbonitrile 1c: A mixture of 2 (2.08 g) and pyridine (0.4 mL) in acetic anhydride was refluxed for 10 h. The reaction solution was cooled and poured onto ice water. The resulting precipitate was collected by filtration to yield the title compound as a tan solid (1.28 g, 68% yield)

3-amino-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide 1:
2-Mercaptoacetamide (480 mg, 5.27 mmol) was added to a freshly prepared solution of sodium methoxide (5.27 mmol) in methanol (20 mL). The resulting solution was stirred at RT for 15 minutes, followed by the addition of **3** (1.0 g, 5.27 mmol). The resulting solution was refluxed overnight. The solvent was removed in vacuo and the residue suspended in ethylacetate (50 mL) and HCl (1 N, 50 mL). The resulting solid was collected by filtration and recrystallized from methanol to yield the title compound as an off white solid (358 mg, 32%).The title compound 3-amino-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide was evaluated in the IKK-2 Resin assay and the IC_{50} was $\geq 10 \leq 20$ μM.

Example 2



3-amino-8-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide

1-chloro-7-methoxy-3,4-dihydronaphthalene-2-carbaldehyde 2a: This material was prepared by the method of example **1a** from 7-methoxy-α-tetralone (5 g) to yield the title compound as a red liquid. This material was used directly in the next step.


1-chloro-7-methoxy-3,4-dihydronaphthalene-2-carbaldehyde oxime 2b: A mixture of **2a** (6.2 g), hydroxyl amine hydrochloride (2.78 g) and sodium acetate (6.15 g) in ethanol (50 mL, 90%) was refluxed for 2.5 h. The reaction was cooled to RT, then poured onto ice water (50 mL). The resulting precipitate was collected by filtration to yield the title compound as a yellow solid (5.8 g, 86% yield from 7-methoxy-α-tetralone).

1-chloro-7-methoxy-3,4-dihydronaphthalene-2-carbonitrile 2c: A mixture of **2b** (5.8 g) and pyridine (0.5 mL) in acetic anhydride (50 mL) was refluxed for 8 h. The reaction solution was cooled and poured onto ice water (250 mL). The resulting precipitate was collected by filtration to yield the title compound as a brown solid (5.7 g, 68% yield).

3-amino-8-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide 2: 2-Mercaptoacetamide (2.09 g) was added to a freshly prepared solution of sodium methoxide (23 mmol) in methanol (50 mL). The resulting solution was stirred at RT for 15 minutes, followed by the addition of **2c** (5.0 g). The resulting solution was refluxed overnight. The solvent was removed in vacuo and the residue suspended in ethylacetate (50 mL) and HCl (1 N, 50 mL). The resulting solid (6.0 g) was collected by filtration and recrystallized from methanol to yield the title

compound as an off white solid (1.1 g, 17%). The title compound 3-amino-8-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide was evaluated in the IKK-2 Resin assay and the IC_{50} was ≥1 ≤10 μM.

Example 3




HBr

3-[(2-aminoethyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide hydrobromide

A solution of **1** (732 mg) and 2-bromoethylamine hydrobromide (1.89 g) in DMF was heated at 80°C for 16 h. The solvent was removed and the residue was purified by reverse phase HPLC to yield the title compound. The title compound 3-[(2-aminoethyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide hydrobromide was evaluated in the IKK-2 Resin assay and the IC_{50} was ≥1 ≤10 μM.

Example 4




HCl

3-[(3-aminopropyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide hydrochloride


A solution of **1** (732 mg) and 2-bromopropylamine hydrobromide (2.0 g) in DMF was heated at 80°C for 16 h. The solvent was removed and the residue was purified by reverse phase HPLC. The residue was lyophilized from 1N HCl to yield the title compound. The title compound of 3-[(3-aminopropyl)amino]-4,5-dihydronaphtho-[1,2-b]thiophene-2-carboxamide hydrochloride was evaluated in the IKK-2 Resin assay and the IC_{50} was ≥10 ≤20 μM.

Example 5



3-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide To a stirred solution of **1** (512 mg) in acetic acid (35 mL) and water (7 mL) was added sodium cyanate. The resulting solution was stirred for 16 h. The resulting solid was removed by filtration. The solid was suspended in ethanol (95%, 50 mL) and filtered and dried to yield the title compound. The title compound 3-[(amino-carbonyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide was evaluated in the IKK-2 Resin assay and the IC_{50} was ≥10 ≤20 μM.


Example 6



2-amino-4,5-dihydronaphtho[1,2-b]thiophene-3-carbonitrile

To a suspension of β-tetralone (22.6 g, 0.154 mol), malonitrile (20 mL, 0.16 mol) and sulfur (7.2 g, 0.22 mol) in ethanol (160 mL) was added morpholine (5 mL) dropwise. The reaction mixture was refluxed for 15 min and then another 150 mL of ethanol was added. The hot suspension as filtered and the solid was treated with a mixture of acetone and ether. The residual sulfur was removed by filtration and the filtrate was concentrated to give 13.3 g of desired product as a tan solid (33% yield). The title compound 2-amino-4,5-dihydronaphtho[1,2-b]thiophene-3-carbonitrile was evaluated in the IKK-2 Resin assay and the IC_{50} was ≥10 ≤20 μM.


Example 7



2-(acetylamino)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide

To a suspension of β-tetralone (11.3 g, 0.077mol), cyanoacetamide (5.34 g, 0.08 mol) and sulfur (3.6 g, 0.11 mol) in ethanol (40 mL) was added diethylamine (2.5 mL) dropwise at room temperature. The reaction mixture was refluxed for 3 h and then cooled. The precipitate was collected by filtration, washed with ether and air-dried to give 1.47 g of 2-amino-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide as a light yellow solid (8% yield). To a solution of this compound (0.13 g, 0.00053 mol) in AcOH (5 mL) was added a solution of sodium cyanate (0.07 g, 0.0011 mol) in water (2 mL). The mixture was stirred at room temperature overnight. This crude solid was isolated by filtration and purified by preparative HPLC to give 0.01 g of the desired product as a white solid. The title compound 2-(acetylamino)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide was evaluated in the IKK-2 Resin assay and the IC_{50} was ≥10 ≤20 μM.

Example 8




2-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide






To a solution of 2-amino-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide (0.5 g, 0.002 mol) in AcOH (10 mL) was added a solution of sodium cyanate (0.135 g, 0.004 mol) in water (4 mL). The mixture was stirred at room temperature overnight. Another 4 eq of sodium cyanate was added and the solution was stirred for another 18 h. The precipitate was collected by filtration and purified by preparative HPLC to give 0.19 g of the desired product as a tan solid. The title compound 2-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide was evaluated in the IKK-2 Resin assay and the IC_{50} was ≤ 1 μM.







The following compounds of the invention can be synthesized according to the Scheme III and to similar procedure as described in the preparation of example 8 using corresponding starting materials.

Scheme III















a) cyanoacetamide, sulfur, diethylamine b) sodium cyanate, AcOH.

Example number	Structure	Name
9.		2-[(aminocarbonyl)-amino]-7-methoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
10.		2-[(aminocarbonyl)-amino]-6-methoxy-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide
11.		2-[(aminocarbonyl)-amino]-6,7-dimethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
12.		2-[(aminocarbonyl)-amino]-7-isopropyl-8-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
13.		2-[(aminocarbonyl)-amino]-4,4-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide







14.		2-[(aminocarbonyl)-amino]-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
15.		2-[(aminocarbonyl)-amino]-7-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
16.		2-[(aminocarbonyl)amino]-8-fluoro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
17.		2-[(aminocarbonyl)-amino]-4-ethyl-4-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
18.		2-[(aminocarbonyl)-amino]-4,4-diethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
19.		2-[(aminocarbonyl)-amino]-8-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







5







20.		2-[(aminocarbonyl)-amino]-9-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
21.		2-[(aminocarbonyl)-amino]-7-methoxy-5,5-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
22.		2-[(aminocarbonyl)-amino]-9-ethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
23.		2-[(aminocarbonyl)-amino]-7,8-dimethoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
24.		2-[(aminocarbonyl)-amino]-6,9-dimethoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
25.		2-[(aminocarbonyl)-amino]-6-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







26.		2-[(aminocarbonyl)-amino]-9-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
27.		2-[(aminocarbonyl)-amino]-7-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
28.		2-[(aminocarbonyl)-amino]-8-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
29.		2-[(aminocarbonyl)-amino]-7,9-dimethoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
30.		2-[(aminocarbonyl)-amino]-6,7-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
31.		2-[(aminocarbonyl)-amino]-7-ethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







5







32.		2-[(aminocarbonyl)-amino]-7,8-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
33.		2-[(aminocarbonyl)-amino]-7-tert-butyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
34.		2-[(aminocarbonyl)-amino]-7-propyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
35.		2-[(aminocarbonyl)-amino]-6-ethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
36.		2-[(aminocarbonyl)-amino]-7-butyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
37.		2-[(aminocarbonyl)-amino]-9-propoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide

38.		2-[(aminocarbonyl)-amino]-9-isopropoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
39.		2-[(aminocarbonyl)-amino]-9-butoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
40.		2-[(aminocarbonyl)-amino]-9-isobutoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
41.		2-[(aminocarbonyl)-amino]-9-sec-butoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
42.		2-[(aminocarbonyl)-amino]-7-methoxy-6-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
43.		2-[(aminocarbonyl)-amino]-8-methoxy-9-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







44.		2-[(aminocarbonyl)-amino]-6-isopropyl-8-methoxy-9-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
45.		2-[(aminocarbonyl)-amino]-7-isopropyl-6-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
46.		2-[(aminocarbonyl)-amino]-7-isopropyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide
47.		2-[(aminocarbonyl)-amino]-7-bromo-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
48.		2-[(aminocarbonyl)-amino]-5,5,8-trimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
49.		2-[(aminocarbonyl)-amino]-6-chloro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







50.		2-[(aminocarbonyl)-amino]-7-chloro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
51.		2-[(aminocarbonyl)-amino]-8-chloro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
52.		2-[(aminocarbonyl)-amino]-6,8-dichloro-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
53.		2-[(aminocarbonyl)-amino]-7,8-dichloro-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
54.		2-[(aminocarbonyl)-amino]-5,5-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
55.		2-[(aminocarbonyl)-amino]-4,5,5-trimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide







56.		2-[(aminocarbonyl)-amino]-8-hexyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
57.		6-(acetylamino)-2-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
58.		2-[(aminocarbonyl)-amino]-9-bromo-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
59.		2-[(aminocarbonyl)-amino]-6-fluoro-9-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
60.		2-[(aminocarbonyl)-amino]-4-ethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
61.		2-[(aminocarbonyl)-amino]-6,9-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide







62.		2-[(aminocarbonyl)-amino]-7-ethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
63.		2-[(aminocarbonyl)-amino]-7-propoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
64.		2-[(aminocarbonyl)-amino]-8-ethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
65.		2-[(aminocarbonyl)-amino]-5-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
66.		2-[(aminocarbonyl)-amino]-7-hydroxy-8-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
67.		2-[(aminocarbonyl)-amino]-8-nitro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







5

68.		2-[(aminocarbonyl)-amino]-6,8-dimethoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
69.		2-[(aminocarbonyl)-amino]-6-fluoro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
70.		2-[(aminocarbonyl)-amino]-9-fluoro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
71.		2-[(aminocarbonyl)-amino]-6-methoxy-4-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
72.		2-[(aminocarbonyl)-amino]-8-methoxy-4-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
73.		2-[(aminocarbonyl)-amino]-6-chloro-7-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







74.		2-[(aminocarbonyl)-amino]-8-(hydroxy-methyl)-7-methoxy-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide
75.		2-[(aminocarbonyl)-amino]-6-bromo-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
76.		2-[(aminocarbonyl)-amino]-6-iodo-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
77.		2-[(aminocarbonyl)-amino]-8-bromo-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
78.		2-[(aminocarbonyl)-amino]-8-iodo-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
79.		2-[(aminocarbonyl)-amino]-6,8-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide

80.		2-[(aminocarbonyl)-amino]-6-(methylthio)-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
81.		2-[(aminocarbonyl)-amino]-6-(methyl-sulfonyl)-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
82.		2-[(aminocarbonyl)-amino]-8-ethynyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
83.		2-[(aminocarbonyl)-amino]-8-chloro-5-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
84.		2-[(aminocarbonyl)-amino]-8-(methylthio)-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
85.		2-[(aminocarbonyl)-amino]-8-(methyl-sulfonyl)-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







86.		2-[(aminocarbonyl)-amino]-7-chloro-5-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
87.		2-[(aminocarbonyl)-amino]-8-(ethylthio)-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
88.		2-[(aminocarbonyl)-amino]-8-(ethyl-sulfonyl)-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
89.		2-[(aminocarbonyl)-amino]-6-chloro-5-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
90.		2-[(aminocarbonyl)-amino]-6-(ethylthio)-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
91.		2-[(aminocarbonyl)-amino]-6-(ethyl-sulfonyl)-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide






92.		2-[(aminocarbonyl)-amino]-9-chloro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
93.		2-[(aminocarbonyl)-amino]-9-methoxy-4,4-dimethyl-4,5-dihydro-naphtho[1,2-b-]-thiophene-3-carboxamide
94.		2-[(aminocarbonyl)-amino]-8-methoxy-7-methyl-4,5-dihydron-aphtho[1,2-b]-thiophene-3-carboxamide
95.		2-[(aminocarbonyl)-amino]-7-methoxy-8-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
96.		2-[(aminocarbonyl)-amino]-6-methoxy-7-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
97.		2-[(aminocarbonyl)-amino]-9-methoxy-4-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide

5







98.		2-[(aminocarbonyl)-amino]-6,7-diethoxy-4,5-dihydronaphtho[-1,2-b]thiophene-3-carboxamide
99.		2-[(aminocarbonyl)-amino]-8-ethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
100.		2-[(aminocarbonyl)-amino]-8-isopropyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide
101.		2-[(aminocarbonyl)-amino]-7,9-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
102.		2-[(aminocarbonyl)-amino]-8-tert-butyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide
103.		2-[(aminocarbonyl)-amino]-7-methoxy-9-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







5







104.		2-[(aminocarbonyl)-amino]-6,8-difluoro-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide
105.		2-[(aminocarbonyl)-amino]-7-isopropoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
106.		2-[(aminocarbonyl)-amino]-6-chloro-9-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
107.		2-[(aminocarbonyl)-amino]-7,9-dichloro-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
108.		2-[(aminocarbonyl)-amino]-8-(trifluoro-methyl)-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
109.		2-[(aminocarbonyl)-amino]-9-methoxy-6-(methylthio)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







110.		2-[(aminocarbonyl)-amino]-9-methoxy-6-(methylsulfonyl)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
111.		2-[(aminocarbonyl)-amino]-6-ethoxy-9-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
112.		2-[(aminocarbonyl)-amino]-6,9-diethoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
113.		2-[(aminocarbonyl)-amino]-8,9-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
114.		2-[(aminocarbonyl)-amino]-7,8-dinitro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide








5







115.		2-[(aminocarbonyl)-amino]-7-nitro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
116.		2-[(aminocarbonyl)-amino]-8-methoxy-4,7-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
117.		2-[(aminocarbonyl)-amino]-6-methoxy-5-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
118.		2-[(aminocarbonyl)-amino]-7-iodo-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
119.		2-[(aminocarbonyl)-amino]-7-chloro-8-nitro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
120.		2-[(aminocarbonyl)-amino]-5,8-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide








121.		2-[(aminocarbonyl)-amino]-5-ethyl-8-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
122.		benzyl 3-(amino-carbonyl)-2-[(amino-carbonyl)amino]-4,5-dihydronaphtho[1,2-b]-thien-8-ylcarbamate
123.		8-amino-2-[(amino-carbonyl)amino]-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
124.		8-(acetylamino)-2-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
125.		2-[(aminocarbonyl)-amino]-5-isopropyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
126.		2-[(aminocarbonyl)-amino]-6,9-dichloro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







127.		2-[(aminocarbonyl)-amino]-9-tert-butyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
128.		2-[(aminocarbonyl)amino]-5,5,7-trimethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
129.		2-[(aminocarbonyl)-amino]-5-butyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
130.		2-[(aminocarbonyl)-amino]-5-butyl-7-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
131.		2-[(aminocarbonyl)-amino]-5-butyl-7-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
132.		2-[(aminocarbonyl)-amino]-5-butyl-7-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide

133.		2-[(aminocarbonyl)-amino]-5-butyl-7,8-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
134.		2-(acetylamino)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
135.		2-(acetylamino)-7-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
136.		2-(acetylamino)-6-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
137.		2-(acetylamino)-6,7-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
138.		2-(acetylamino)-7-isopropyl-8-methoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide







139.		2-(acetylamino)-4,4-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
140.		2-(acetylamino)-4-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
141.		2-(acetylamino)-7-fluoro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
142.		2-(acetylamino)-8-fluoro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
143.		2-(acetylamino)-4-ethyl-4-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
144.		2-(acetylamino)-4,4-diethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
145.		2-(acetylamino)-8-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide

146.		2-(acetylamino)-9-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
147.		2-(acetylamino)-7-methoxy-5,5-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
148.		2-(acetylamino)-9-ethoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
149.		2-(acetylamino)-7,8-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
150.		2-(acetylamino)-6,9-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
151.		2-(acetylamino)-6-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







152.		2-(acetylamino)-9-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
153.		2-(acetylamino)-7-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
154.		2-(acetylamino)-8-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
155.		2-(acetylamino)-7,9-dimethoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
156.		2-(acetylamino)-6,7-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
157.		2-(acetylamino)-7-ethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
158.		2-(acetylamino)-7,8-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







159.		2-(acetylamino)-7-tert-butyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
160.		2-(acetylamino)-7-propyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
161.		2-(acetylamino)-6-ethoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
162.		2-(acetylamino)-7-butyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
163.		2-(acetylamino)-9-propoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
164.		2-(acetylamino)-9-isopropoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide

5







165.		2-(acetylamino)-9-butoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
166.		2-(acetylamino)-9-isobutoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
167.		2-(acetylamino)-9-sec-butoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
168.		2-(acetylamino)-7-methoxy-6-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
169.		2-(acetylamino)-8-methoxy-9-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
170.		2-(acetylamino)-6-isopropyl-8-methoxy-9-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







5

171.		2-(acetylamino)-7-isopropyl-6-methoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
172.		2-(acetylamino)-7-isopropyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
173.		2-(acetylamino)-7-bromo-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
174.		2-(acetylamino)-5,5,8-trimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
175.		2-(acetylamino)-6-chloro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
176.		2-(acetylamino)-7-chloro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







177.		2-(acetylamino)-8-chloro-4,5-dihydro-naphtho-[1,2-b]thiophene-3-carboxamide
178.		2-(acetylamino)-6,8-dichloro-4,5-dihydro-naphtho[1,2-b]-hiophene-3-carboxamide
179.		2-(acetylamino)-7,8-dichloro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
180.		2-(acetylamino)-5,5-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
181.		2-(acetylamino)-4,5,5-trimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
182.		2-(acetylamino)-8-hexyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







5







183.		2,6-bis(acetylamino)-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
184.		2-(acetylamino)-9-bromo-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
185.		2-(acetylamino)-6-fluoro-9-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
186.		2-(acetylamino)-4-ethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
187.		2-(acetylamino)-6,9-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
188.		2-(acetylamino)-7-ethoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







189.		2-(acetylamino)-7-propoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
190.		2-(acetylamino)-8-ethoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
191.		2-(acetylamino)-5-methyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
192.		2-(acetylamino)-7-hydroxy-8-methoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
193.		2-(acetylamino)-8-nitro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
194.		2-(acetylamino)-6,8-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







5








195.		2-(acetylamino)-6-fluoro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
196.		2-(acetylamino)-9-fluoro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
197.		2-(acetylamino)-6-methoxy-4-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
198.		2-(acetylamino)-8-methoxy-4-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
199.		2-(acetylamino)-6-chloro-7-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
200.		2-(acetylamino)-8-(hydroxymethyl)-7-methoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







201.		2-(acetylamino)-6-bromo-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
202.		2-(acetylamino)-6-iodo-4,5-dihydro-naphtho-[1,2-b]thiophene-3-carboxamide
203.		2-(acetylamino)-8-bromo-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
204.		2-(acetylamino)-8-iodo-4,5-dihydro-naphtho-[1,2-b]-thiophene-3-carboxamide
205.		2-(acetylamino)-6,8-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
206.		2-(acetylamino)-6-(methylthio)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







207.		2-(acetylamino)-6-(methylsulfonyl)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
208.		2-(acetylamino)-8-ethynyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
209.		2-(acetylamino)-8-chloro-5-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
210.		2-(acetylamino)-8-(methylthio)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
211.		2-(acetylamino)-8-(methylsulfonyl)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
212.		2-(acetylamino)-7-chloro-5-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







213.		2-(acetylamino)-8-(ethylthio)-4,5-dihydro-naphtho[1,2b]-thiophene-3-carboxamide
214.		2-(acetylamino)-8-(ethylsulfonyl)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
215.		2-(acetylamino)-6-chloro-5-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
216.		2-(acetylamino)-6-(ethylthio)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
217.		2-(acetylamino)-6-(ethylsulfonyl)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
218.		2-(acetylamino)-9-chloro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide

219.		2-(acetylamino)-9-methoxy-4,4-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
220.		2-(acetylamino)-8-methoxy-7-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
221.		2-(acetylamino)-7-methoxy-8-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
222.		2-(acetylamino)-6-methoxy-7-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
223.		2-(acetylamino)-9-methoxy-4-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
224.		2-(acetylamino)-6,7-diethoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







225.		2-(acetylamino)-8-ethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
226.		2-(acetylamino)-8-isopropyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
227.		2-(acetylamino)-7,9-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
228.		2-(acetylamino)-8-tert-butyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
229.		2-(acetylamino)-7-methoxy-9-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
230.		2-(acetylamino)-6,8-difluoro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
231.		2-(acetylamino)-7-isopropoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide







232.		2-(acetylamino)-6-chloro-9-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
233.		2-(acetylamino)-7,9-dichloro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
234.		2-(acetylamino)-8-(trifluoromethyl)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
235.		2-(acetylamino)-9-methoxy-6-(methylthio)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
236.		2-(acetylamino)-9-methoxy-6-(methyl-sulfonyl)-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
237.		2-(acetylamino)-6-ethoxy-9-methoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide







238.		2-(acetylamino)-6,9-diethoxy-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
239.		2-(acetylamino)-8,9-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
240.		2-(acetylamino)-7,8-dinitro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
241.		2-(acetylamino)-7-nitro-4,5-dihydro-naphtho-[1,2-b]-thiophene-3-carboxamide
242.		2-(acetylamino)-8-methoxy-4,7-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
243.		2-(acetylamino)-6-methoxy-5-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide







244.		2-(acetylamino)-7-iodo-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
245.		2-(acetylamino)-7-chloro-8-nitro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
246.		2-(acetylamino)-5,8-dimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
247.		2-(acetylamino)-5-ethyl-8-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
248.		benzyl 2-(acetyl-amino)-3-(amino-carbonyl)-4,5-dihydro-naphtho[1,2-b]-thien-8-ylcarbamate
249.		2-(acetylamino)-8-amino-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







5







250.		2,8-bis(acetylamino)-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
251.		2-(acetylamino)-5-isopropyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
252.		2-(acetylamino)-6,9-dichloro-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
253.		2-(acetylamino)-9-tert-butyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
254.		2-(acetylamino)-5,5,7-trimethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide
255.		2-(acetylamino)-5-butyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide







256.		2-(acetylamino)-5-butyl-7-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
257.		2-(acetylamino)-5-butyl-7-fluoro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
258.		2-(acetylamino)-5-butyl-7-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide
259.		2-(acetylamino)-5-butyl-7,8-dimethoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide
260.		2-(acetylamino)-5,6-dihydro-4H-benzo-[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide
261.		2-(acetylamino)-9-chloro-5,6-dihydro-4H-benzo[6,7]cyclo-hepta[1,2-b]thiophene-3-carboxamide






262.		2-(acetylamino)-8,9-dichloro-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide
263.		2-(acetylamino)-8-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide
264.		2-(acetylamino)-8-methoxy-6,6-dimethyl-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide
265.		2-(acetylamino)-9-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide
266.		2-(acetylamino)-8,10-dimethoxy-5,6-dihydro-4H-benzo[6,7]cyclo-hepta[1,2-b]thiophene-3-carboxamide
267.		2-(acetylamino)-8-methyl-5,6-dihydro-4H-benzo-[6,7]cyclohepta-[1,2-b]-thiophene-3-carboxamide

268.		2-(acetylamino)-8,9-dimethoxy-5,6-dihydro-4H-benzo[6,7]cyclo-hepta[1,2-b]thiophene-3-carboxamide
269.		2-(acetylamino)-8-fluoro-9-methoxy-5,6-dihydro-4H-benzo[6,7]-cyclo-hepta[1,2-b]-thiophene-3-carboxamide
270.		2-(acetylamino)-7-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide
271.		2-(acetylamino)-6,9-dimethyl-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide
272.		2-(acetylamino)-10-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide
273.		2-(acetylamino)-9-bromo-6-methyl-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide







274.		2-(acetylamino)-7,8-dimethoxy-5,6-dihydro-4H-benzo[6,7]cyclo-hepta[1,2-b]thiophene-3-carboxamide
275.		2-(acetylamino)-6-methyl-5,6-dihydro-4H-benzo-[6,7]cyclohepta-[1,2-b]-thiophene-3-carboxamide
276.		2-(acetylamino)-9-chloro-6-methyl-5,6-dihydro-4H-benzo-[6,7]cyclohepta-[1,2-b]-thiophene-3-carboxamide
277.		2-(acetylamino)-9-fluoro-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide
278.		2-(acetylamino)-9-methyl-5,6-dihydro-4H-benzo-[6,7]cyclo-hepta[1,2-b]-thiophene-3-carboxamide
279.		2-(acetylamino)-9-amino-5,6-dihydro-4H-benzo-[6,7]cyclo-hepta[1,2-b]-thiophene-3-carboxamide


280.		2-(acetylamino)-9-nitro-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide
281.		2-(acetylamino)-9,10-dimethoxy-5,6-dihydro-4H-benzo[6,7]cyclo-hepta[1,2-b]thiophene-3-carboxamide
282.		2-[(aminocarbonyl)-amino]-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide
283.		2-[(aminocarbonyl)-amino]-9-chloro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]-thiophene-3-carboxamide
284.		2-[(aminocarbonyl)-amino]-8,9-dichloro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]-thiophene-3-carboxamide
285.		2-[(aminocarbonyl)-amino]-8-methoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]-thiophene-3-carboxamide

286.		2-[(aminocarbonyl)-amino]-8-methoxy-6,6-dimethyl-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide
287.		2-[(aminocarbonyl)-amino]-9-methoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]-thiophene-3-carboxamide
288.		2-[(aminocarbonyl)-amino]-8,10-dimethoxy-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide
289.		2-[(aminocarbonyl)-amino]-8-methyl-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]-thiophene-3-carboxamide
290.		2-[(aminocarbonyl)amino]-8,9-dimethoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]-thiophene-3-carboxamide
291.		2-[(aminocarbonyl)-amino]-8-fluoro-9-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide

292.		2-[(aminocarbonyl)-amino]-7-methoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]-thiophene-3-carboxamide
293.		2-[(aminocarbonyl)-amino]-6,9-dimethyl-5,6-dihydro-4H-benzo-[6,7]cyclo-hepta[1,2-b]-thiophene-3-carboxamide
294.		2-[(aminocarbonyl)-amino]-10-methoxy-5,6-dihydro-4H-benzo[6,7]-cyclo-hepta[1,2-b]-thiophene-3-carboxamide
295.		2-[(aminocarbonyl)-amino]-9-bromo-6-methyl-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide
296.		2-[(aminocarbonyl)-amino]-7,8-dimethoxy-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide

5

297.		2-[(aminocarbonyl)-amino]-6-methyl-5,6-dihydro-4H-benzo-[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide
298.		2-[(aminocarbonyl)-amino]-9-chloro-6-methyl-5,6-dihydro-4H-benzo-[6,7]cyclo-hepta[1,2-b]thiophene-3-carboxamide
299.		2-[(aminocarbonyl)-amino]-9-fluoro-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide
300.		2-[(aminocarbonyl)-amino]-9-methyl-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]-thiophene-3-carboxamide
301.		9-amino-2-[(amino-carbonyl)amino]-5,6-dihydro-4H-benzo-[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide
302.		2-[(aminocarbonyl)-amino]-9-nitro-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide

303.		2-[(aminocarbonyl)-amino]-9,10-dimethoxy-5,6-dihydro-4H-benzo-[6,7]cyclo-hepta[1,2-b]thiophene-3-carboxamide

BIOLOGICAL EVALUATION

Materials

SAM^{2} ™ 96 Biotin capture plates were from Promega. Anti-FLAG affinity resin, FLAG-peptide, NP-40 (Nonidet P-40), BSA, ATP, ADP, AMP, LPS (*E. coli* serotype 0111:B4), and dithiothreitol were obtained from Sigma Chemicals. Antibodies specific for NEMO (IKKγ) (FL-419), IKK1(H-744), IKK2(H-470) and IκBα(C–21) were purchased from Santa Cruz Biotechnology. Ni-NTA resin was purchased from Qiagen. Peptides were purchased from American Peptide Company. Protease inhibitor cocktail tablets were from Boehringer Mannheim. Sephacryl S-300 column was from Pharmacia LKB Biotechnology. Centriprep-10 concentrators with a molecular weight cutoff of 10 kDa and membranes with molecular weight cut-off of 30 kDa were obtained from Amicon. [ϒ-^{33}P] ATP (2500 Ci/mmol) and [ϒ-^{32}P] ATP (6000 Ci/mmol) were purchased from Amersham. The other reagents used were of the highest grade commercially available.

Cloning and Expression

cDNAs of human IKK1 and IKK2 were amplified by reverse transcriptase-polymerase chain reaction from human placental RNA (Clonetech). hIKK1 was subcloned into pFastBac HTa (Life Technologies) and expressed as N-terminal His_6-tagged fusion protein. The hIKK2 cDNA was amplified using a reverse oligonucleotide primer which incorporated the peptide sequence for a FLAG-epitope tag at the C-terminus of the IKK2 coding region (DYKDDDDKD). The

hIKK2:FLAG cDNA was subcloned into the baculovirus vector pFastBac. The rhIKK2 (S177S, E177E) mutant was constructed in the same vector used for wild type rhIKK2 using a QuikChange™ mutagenesis kit (Stratagene). Viral stocks of each construct were used to infect insect cells grown in 40L suspension culture. The cells were lysed at a time that maximal expression and rhIKK activity were demonstrated. Cell lysates were stored at -80 °C until purification of the recombinant proteins was undertaken as described below.

Enzyme Isolation

All purification procedures were carried out at 4 °C unless otherwise noted. Buffers used are: buffer A: 20 mM Tris-HCl, pH 7.6, containing 50 mM NaCl, 20 mM NaF, 20 mM β-Glycerophosphate, 500 uM sodiumortho-vanadate, 2.5 mM metabisulfite, 5 mM benzamidine, 1 mM EDTA, 0.5 mM EGTA, 10% glycerol, 1 mM DTT, 1X Complete™ protease inhibitors; buffer B: same as buffer A, except 150 mM NaCl, and buffer C: same as buffer A, except 500 mM NaCl.

Isolation of rhIKK1 homodimer

Cells from an 8-liter fermentation of baculovirus-expressed IKK1 tagged with His peptide were centrifuged and the cell pellet (MOI 0.1, I=72 hr) was re-suspended in 100 ml of buffer C. The cells were microfluidized and centrifuged at 100,000 X g for 45 min. The supernatant was collected, imidazole added to the final concentration of 10 mM and incubated with 25 ml of Ni-NTA resin for 2 hrs. The suspension was poured into a 25 ml column and washed with 250 ml of buffer C and then with 125 ml of 50 mM imidazole in buffer C. rhIKK1 homodimer was eluted using 300 mM imidazole in buffer C. BSA and NP-40 were added to the enzyme fractions to the final concentration of 0.1 %. The enzyme was dialyzed against buffer B, aliquoted and stored at -80 °C.

Isolation of rhIKK2 homodimer

A 10-liter culture of baculovirus-expressing IKK2 tagged with FLAG peptide was centrifuged and the cell pellet (MOI=0.1 and I=72 hrs) was re-suspended in buffer A. These cells were microfluidized, and centrifuged at 100,000 X g for 45 min.

Supernatant was passed over a G-25 column equilibrated with Buffer A. Protein peak was collected and incubated with anti-FLAG affinity resin on a rotator overnight in buffer B. The resin was washed in batch with 10-15 bed volumes of buffer C. Washed resin was poured into a column and rhIKK2 homodimer was eluted using 5 bed volumes of buffer B containing FLAG peptide. 5 mM DTT, 0.1% NP-40 and BSA (concentrated to 0.1% in final amount) was added to the eluted enzyme before concentrating in using an Amicon membrane with a molecular weight cut-off of 30 kDa. Enzyme was aliquoted and stored at -80 °C.

Isolation of rhIKK1/IKK2 heterodimer

The heterodimer enzyme was produced by coinfection in a baculovirus system (FLAG IKK2/IKK1 His; MOI=0.1 and I=72 hrs). Infected cells were centrifuged and the cell pellet (10.0 g) was suspended in 50 ml of buffer A. The protein suspension was microfluidized and centrifuged at 100,000 X g for 45 min. Imidazole was added to the supernatant to a final concentration of 10 mM. The protein was allowed to bind 25 ml of Ni-NTA resin by mixing for 2 hrs. The protein-resin slurry was poured into a 25 ml column and washed with 250 ml of buffer A containing 10 mM imidazole followed by 125 ml of buffer A containing 50 mM imidazole. Buffer A, containing 300 mM imidazole, was then used to elute the protein. A 75 ml pool was collected and NP-40 was added to a final concentration of 0.1%. The protein solution was then dialyzed against buffer B. The dialyzed heterodimer enzyme was then allowed to bind to 25 ml of anti-FLAG M2 agarose affinity gel overnight with constant mixing. The protein-resin slurry was then centrifuged for 5 min at 2,000 rpm. The supernatant was collected and the resin re-suspended in 100 ml of buffer C containing 0.1% NP-40. The resin was washed with 375 ml of buffer C containing 0.1 % NP-40. The protein-resin was poured into a 25 ml column and the enzyme eluted using buffer B containing FLAG peptide. Enzyme fractions (100 ml) were collected and concentrated to 20 ml using an Amicon membrane with molecular weight cut-off of 30 kDa. Bovine serum albumin was added to the concentrated enzyme to final concentration of 0.1 %. The enzyme was then aliquoted and stored at -80 °C.

Cell Culture

The wild type (wt) human pre-B cell line, 70Z/3, and its mutant, 1.3E2, were generously provided by Dr. Carol Sibley. Wt 70Z/3 and 1.3E2 cells were grown in RPMI 1640 (Gibco) supplemented with 7 % defined bovine serum (Hyclone) and 50 μM 2-mercaptoethanol. Human monocytic leukemia THP-1 cells, obtained from ATCC, were cultured in RPMI 1640 supplemented with 10% defined bovine serum, 10 mM HEPES, 1.0 mM sodium pyruvate and 50 μM 2-mercaptoethanol. For experiments, cells were plated in 6 well plates at $1x10^6$ cells/ml in fresh media. Pre-B cells were stimulated by the addition of 10 μg/ml LPS for varying lengths of time ranging from 0-4 hr. THP-1 cells were stimulated by the addition of 1 μg/ml LPS for 45 minutes. Cells were pelleted, washed with cold 50 mM sodium phosphate buffer, pH 7.4 containing 0.15 M NaCl and lysed at 4 °C in 20 mM Hepes buffer, pH 7.6 containing 50 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM sodium orthovanadate, 10 mM β-glycerophosphate, 1 mM NaF, 1 mM PMSF, 1 mM DTT and 0.5 % NP40 (lysis buffer). The cytosolic fractions obtained following centrifugation at 10,000 X g were stored at -80° C until used.

Immunoprecipitation and Western Blotting

SF9 cells paste containing rhIKKs were centrifuged (100,000 X g, 10 min) to remove debris. rhIKKs were immunoprecipitated (100 μg of cell paste) from the cell supernatant using 3 μg of anti-NEMO antibody (FL-419), followed by coupling to protein A sepharose beads. rhIKKs were also immunoprecipitated from affinity chromatography purified protein preparations (1 μg) using anti-FLAG, anti-His or anti-NEMO antibodies (1-4 μg) followed by protein A sepharose coupling. The native, human IKK complex was immunoprecipitated from THP-1 cell homogenates (300 μg/condition) using the anti-NEMO antibody. Immune complexes were pelleted and washed 3 times with 1 ml cold lysis buffer. Immunoprecipitated rhIKKs were chromatographed by SDS-PAGE (8% Tris-glycine) and transferred to nitrocellulose membranes (Novex) and detected by chemiluminescense (SuperSignal) using specific anti-IKK antibodies (IKK2 H-470, IKK1 H-744). Native IKK2, IκBα and NEMO proteins from cytosolic lysates (20-

80 μg) were separated by SDS-PAGE and visualized by chemiluminescense using specific antibodies.

Phosphatase Treatment

Immunoprecipitated rhIKKs were washed 2 times in 50 mM Tris-HCl, pH 8.2 containing 0.1 mM EDTA, 1 mM DTT, 1 mM PMSF and 2 mM $MnCl_2$ and resuspended in 50 μl. Phosphatase (λPPase, 1000 U) was pre-diluted in the same buffer and added to the IKK samples. Following incubation at room temperature for 30 minutes with intermittent mixing, cold lysis buffer was added to the tubes to stop the reaction. After several washes, 10 % of the beads were removed for Western analysis, and the remaining material was pelleted and resuspended in 100 μl of the buffer used for the *in vitro* kinase assay.

IKKα SAM Enzyme Assay

IKKα kinase activity was measured using a biotinylated IκBα peptide (Gly-Leu-Lys-Lys-Glu-Arg-Leu-Leu-Asp-Asp-Arg-His-Asp-Ser_{32}-Gly-Leu-Asp-Ser_{36}-Met-Lys-Asp-Glu-Glu), a SAM^{2} ™ 96 Biotin capture plate, and a vacuum system. The standard reaction mixture contained 5 μM biotinylated IκBα peptide, 1 μM [γ-^{33}P] ATP (about 1 X 10^5 cpm), 1 mM DTT, 50 mM KCl, 2 mM $MgCl_2$, 2 mM $MnCl_2$, 10 mM NaF, 25 mM Hepes buffer, pH. 7.6 and enzyme solution (1-10 μl) in a final volume of 50 μl. After incubation at 25 °C for 30 min, 25 μl of the reaction mixture was withdrawn and added to a SAM^{2} ™ 96 Biotin capture 96-well plate. Each well was then washed successively with 800 μl 2 M NaCl, 1.2 ml of NaCl containing 1% H_3PO_4, 400 μl H_2O, and 200 μl 95% ethanol. The plate was allowed to dry in a hood at 25 °C for 1 hr and then 25 μl of scintillation fluid (Microscint 20) was added to each well. Incorporation of [γ-^{33}P] ATP was measured using a Top-Count NXT (Packard). Under each assay condition, the degree of phosphorylation of IκBα peptide substrate was linear with time and concentration for all purified enzymes. Results from the biotinylated peptide assay were confirmed by SDS-PAGE analysis of kinase reaction utilizing a GST- $IκBα_{1-54}$ and [γ-^{32}P]ATP. The resulting radiolabeled substrate was quantitated by Phosphoimager

(Molecular Dynamics). An ion exchange resin assay was also employed using [γ-^{33}P] ATP and GST-IκBα_{1-54} fusion protein as the substrates. Each assay system yielded consistent results in regard to K_m and specific activities for each of the purified kinase isoforms. One unit of enzyme activity was defined as the amount required to catalyze the transfer of 1 nmole of phosphate from ATP to IκBα peptide per min. Specific activity was expressed as units per mg of protein. For experiments related to K_m determination of purified enzymes, various concentrations of ATP or IκBα peptide were used in the assay at either a fixed IκBα or ATP concentration. For IκBα peptide K_m, assays were carried out with 0.1 μg of enzyme, 5 μM ATP and IκBα peptide from 0.5 to 20 μM. For ATP K_m, assays were carried out with 0.1 μg of enzyme, 10 μM IκBα peptide and ATP from 0.1 to 10 μM. For K_m determination of rhIKK1 homodimer, due to its low activity and higher K_m for IκBα peptide, rhIKK1 homodimer (0.3 μg) was assayed with 125 μM IκBα peptide and a 5-fold higher specific activity of ATP (from 0.1 to 10 μM) for ATP K_m experiments and a 5-fold higher specific activity of 5 μM ATP and IκBα peptide (from 5 to 200 μM) for IκBα peptide K_m experiments.

IKKβ Resin Enzyme Assay


IKKβ kinase activity was measured using a biotinylated IκBα peptide (Gly-Leu-Lys-Lys-Glu-Arg-Leu-Leu-Asp-Asp-Arg-His-Asp-Ser$_{32}$-Gly-Leu-Asp-Ser$_{36}$-Met-Lys-Asp-Glu-Glu) (American Peptide Co.). 20 ul of the standard reaction mixture contained 5 μM biotinylated IκBα peptide, 0.1 μCi/reaction [γ-^{33}P] ATP (Amersham) (about 1 X 10^5 cpm), 1 μM ATP (Sigma), 1 mM DTT (Sigma), 2 mM $MgCl_2$ (Sigma), 2 mM $MnCl_2$ (Sigma), 10 mM NaF (Sigma), 25 mM Hepes (Sigma) buffer, pH 7.6 and 20 μl enzyme solution and 10 ul inhibitor in a final volume of 50 μl. After incubation at 25 °C for 30 min, 150 μl resin (Dowex anion-exchange resin AG1X8 200-400 mesh) in 900 mM formate, pH 3.0 was added to each well to stop the reaction. Resin was allowed to settle for one hour and 50 ul of supernatant was removed to a Micolite-2 flat bottom plate (Dynex). 150 μl of scintillation fluid (Microscint 40) (Packard) was added to each well. Incorporation of [γ-^{33}P] ATP was measured using a Top-Count NXT (Packard).

IKK heterodimer Resin Enzyme Assay

IKK heterodimer kinase activity was measured using a biotinylated IκBα peptide (Gly-Leu-Lys-Lys-Glu-Arg-Leu-Leu-Asp-Asp-Arg-His-Asp-Ser_{32}-Gly-Leu-Asp-Ser_{36}-Met-Lys-Asp-Glu-Glu) (American Peptide Co.). 20 ul of the standard reaction mixture contained 5 μM biotinylated IκBα peptide, 0.1 μCi/reaction [γ-^{33}P] ATP (Amersham) (about 1 X 10^5 cpm), 1 μM ATP (Sigma), 1 mM DTT (Sigma), 2 mM $MgCl_2$ (Sigma), 2 mM $MnCl_2$ (Sigma), 10 mM NaF (Sigma), 25 mM Hepes (Sigma) buffer, pH 7.6 and 20 μl enzyme solution and 10 μl inhibitor in a final volume of 50 μl. After incubation at 25 °C for 30 min, 150 μl resin (Dowex anion-exchange resin AG1X8 200-400 mesh) in 900 mM formate, pH 3.0 was added to each well to stop the reaction. Resin was allowed to settle for one hour and 50 μl of supernatant was removed to a Micolite-2 flat bottom plate (Dynex). 150 μl of scintillation fluid (Microscint 40) (Packard) was added to each well. Incorporation of [γ-^{33}P] ATP was measured using a Top-Count NXT (Packard).

What is claimed is:

1 A compound of formula I



I

or isomers, tautomers, polymorphs, carriers, prodrugs, pharmaceutically acceptable salts thereof, wherein;

A is $(CH_2)_m$ or $(CH_2)_m$-W-$(CH_2)_n$, wherein A is optionally substituted with one or more substituent independently selected from the group consisting of sulfamyl, halo, alkyl, alkoxy, hydroxyl and haloalkyl, CF_3, $COCF_3$, CN, NO_2, hydrido, OR^3, $OCOOR^3$, CO_2H, CO_2R^3, $CONH_2$, $CONHR^3$, $CON(R^3)_2$, COR^3, SR^3, SOR^3, $SCOOR^3$, SO_2R^3, NH_2, NHR^3, NR^3R^3, NR^3COR^3, NR^3CONHR^3, $NR^3SO_2R^3$, $NR^3SO_2NHR^3$, SO_2NHR^3, and $SO_2N(R^3)_2$;

B is a 6-membered aromatic hydrocarbon ring, optionally substituted with one or more substituent independently selected from the group consisting of OR^3, SR^4, $SO_2N(R^4)_2$, NHR^4, $NHCOR^4$, NR^4COR^4, $NHCO(OR^4)$, $NR^4CO(OR^4)$, $NHCONHR^3$, NR^3CONHR^4, $NR^3SO_2R^4$, $NHSO_2R^4$, $NHSO_2N(R^4)_2$, $NHCONR^4R^4$, CO_2R^4, $CON(R^4)_2$, aryl, heteroaryl, heterocyclic, halo, alkylthio, alkylsulfinyl, alkylsulfonyl, cyano, carboxyl, alkoxycarbonyl, amido, N-monoalkylamido, N-monoarylamido, alkyl, alkenyl, alkynyl, N,N-dialkylamido, N-alkyl-N-arylamido, haloalkyl hydrido, hydroxyl, alkoxy, hydroxyalkyl, haloalkoxy, sulfamyl, N-alkylsulfamyl, amino, alkylamino, and nitro;

W is S(O)p, O, CH=N, N(O)=CH, and NR^4;

m is 0 to 3, inclusive;

n is 0 to 3, inclusive;

p is 0 to 2, inclusive;

R^1 and R^2 are independently selected from the group consisting of : hydrido, cyano, nitro, hydroxyl, alkyl, hydroxyalkyl, haloalkyl, alkoxy, haloalkoxy, amidine, guanidine, $CONHR^5$, NHR^5, and $NHCXNHR^5$, wherein X is O, S, or NR^6;

R^3 is selected from the group consisting of: hydrido, aryl, heteroaryl, lower alkyl, alkenyl, alkynyl, and heteroalkyl;

R^4 is selected from the group consisting of: lower alkyl, aryl, heteroaryl, arylalkyl, heteroalkyl, haloalkyl, arylalkylamino, and heteroarylalkyl, wherein aryl, arylalkyl, heteroaryl, or heteroarylalkyl are optionally substituted with one or more radical selected from alkyl, alkoxy, halo, haloalkyl, cyano, haloalkoxy, acyl, carboxyl, hydroxy, hydroxyalkyloxy, phenoxy, benzyloxy, dialkylaminoalkyloxy, and heterocyclic;

R^5 is selected from the group consisting of: hydrido, alkyl, aminoalkyl, hydroxyalkyl, haloalkyl, and acyl;

R^6 is selected from the group consisting of: hydrido, alkyl, cyano, and nitro.

With the proviso, when W is CH=N and m is 0, the nitrogen atom in CH=N has to be directly attached to the ring B.

2 The compound of claim 1, wherein A is $(CH_2)_2$.

3 The compound of claim 1, wherein A is $(CH_2)_3$.

4 The compound of claim 2 or 3, wherein R^1 is NHR^5.

5 The compound of claim 4, wherein R^1 is $NHCOCH_3$.

6 The compound of claim 4 wherein R^1 is $NHCONHR^5$.

7 The compound of claim 6 wherein R^1 is $NHCONH_2$.

8 The compound of claim 2 or 3, wherein R^2 is $CONHR^5$.

9 The compound of claim 8 wherein R^2 is $CONH_2$.

10 The compound of claim 8 wherein R^1 is $NHCOCH_3$ or $NHCONH_2$.

11 The compound of claim 1 selected from the group consisting of:

3-amino-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide,
3-amino-8-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide,
3-[(2-aminoethyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide hydrobromide,
3-[(3-aminopropyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide hydrochloride,
3-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-2-carboxamide,
2-amino-4,5-dihydronaphtho[1,2-b]thiophene-3-carbonitrile,
2-(acetylamino)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide, and
2-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide.

12 The compound of claim 1 selected from the group consisting of:

2-[(aminocarbonyl)amino]-7-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6,7-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-isopropyl-8-methoxy-4,5-dihydro-naphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-4,4-dimethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-4-ethyl-4-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-4,4-diethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-methoxy-5,5-dimethyl-4,5-dihydro-naphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-ethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7,8-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6,9-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-6-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7,9-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6,7-dimethyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-ethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7,8-dimethyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-tert-butyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-propyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-ethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-butyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-propoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-isopropoxy-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-butoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-isobutoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-9-sec-butoxy-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-7-methoxy-6-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-8-methoxy-9-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-6-isopropyl-8-methoxy-9-methyl-4,5-dihydro-naphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-7-isopropyl-6-methoxy-4,5-dihydro-naphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-7-isopropyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-7-bromo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-5,5,8-trimethyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-6-chloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-7-chloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-8-chloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-6,8-dichloro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-7,8-dichloro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-5,5-dimethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)-mino]-4,5,5-trimethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-8-hexyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

6-(acetylamino)-2-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-bromo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-fluoro-9-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-4-ethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6,9-dimethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)-mino]-7-ethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-propoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-ethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-hydroxy-8-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-nitro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6,8-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-methoxy-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-methoxy-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-6-chloro-7-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-(hydroxy-methyl)-7-methoxy-4,5-dihydro-naphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-bromo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-iodo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-bromo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-iodo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6,8-dimethyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-(methylthio)-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-(methyl-sulfonyl)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-ethynyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-chloro-5-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-(methylthio)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-(methyl-sulfonyl)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-chloro-5-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-(ethylthio)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-(ethyl-sulfonyl)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-6-chloro-5-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-6-(ethylthio)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-6-(ethyl-sulfonyl)-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-9-chloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-9-methoxy-4,4-dimethyl-4,5-dihydronaphtho-[1,2-b-]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-8-methoxy-7-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-7-methoxy-8-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-6-methoxy-7-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-9-methoxy-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-6,7-diethoxy-4,5-dihydronaphtho[-1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-8-ethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-8-isopropyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-7,9-dimethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-8-tert-butyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-7-methoxy-9-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-6,8-difluoro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)-amino]-7-isopropoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)-amino]-6-chloro-9-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7,9-dichloro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-(trifluoro-methyl)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-methoxy-6-(methylthio)-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-methoxy-6-(methylsulfonyl)-4,5-dihydro-naphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-ethoxy-9-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6,9-diethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8,9-dimethyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7,8-dinitro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-nitro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-methoxy-4,7-dimethyl-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-methoxy-5-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-iodo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-chloro-8-nitro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5,8-dimethyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-5-ethyl-8-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
benzyl 3-(aminocarbonyl)-2-[(amino-carbonyl)amino]-4,5-dihydronaphtho-[1,2-b]thien-8-ylcarbamate,
8-amino-2-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
8-(acetylamino)-2-[(aminocarbonyl)amino]-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5-isopropyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6,9-dichloro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-tert-butyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5,5,7-trimethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5-butyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5-butyl-7-methyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5-butyl-7-fluoro-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5-butyl-7-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5-butyl-7,8-dimethoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,7-dimethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-(acetylamino)-7-isopropyl-8-methoxy-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-4,4-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-4-ethyl-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-4,4-diethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-methoxy-5,5-dimethyl-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-9-ethoxy-4,5-dihydronaphtho[1,2-b-thiophene-3-carboxamide,
2-(acetylamino)-7,8-dimethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,9-dimethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-(acetylamino)-8-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
2-(acetylamino)-7,9-dimethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,7-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-ethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7,8-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-tert-butyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-propyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-ethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-butyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-propoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-isopropoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-butoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-isobutoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-sec-butoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-methoxy-6-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-methoxy-9-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-(acetylamino)-6-isopropyl-8-methoxy-9-methyl-4,5-dihydronaphtho-[1,2b]thiophene-3-carboxamide,
2-(acetylamino)-7-isopropyl-6-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-7-isopropyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-bromo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5,5,8-trimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-chloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-chloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-chloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,8-dichloro-4,5-dihydronaphtho[1,2-b]hiophene-3-carboxamide,
2-(acetylamino)-7,8-dichloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5,5-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-4,5,5-trimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-hexyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2,6-bis(acetylamino)-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-bromo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-fluoro-9-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-4-ethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-(acetylamino)-6,9-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-ethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-propoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-ethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-hydroxy-8-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-8-nitro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,8-dimethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-methoxy-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
2-(acetylamino)-8-methoxy-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-chloro-7-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-(hydroxymethyl)-7-methoxy-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-6-bromo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-iodo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-(acetylamino)-8-bromo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-iodo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,8-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-(methylthio)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-(methylsulfonyl)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-ethynyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-chloro-5-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-(methylthio)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-(methylsulfonyl)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-(ethylthio)-4,5-dihydronaphtho[1,2b]thiophene-3-carboxamide,
2-(acetylamino)-8-(ethylsulfonyl)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-chloro-5-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-(ethylthio)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-(ethylsulfonyl)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-chloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-methoxy-4,4-dimethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,

2-(acetylamino)-8-methoxy-7-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-methoxy-8-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-methoxy-7-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-methoxy-4-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,7-diethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-ethyl-4,5-dihydro-naphtho[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-8-isopropyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7,9-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide
2-(acetylamino)-8-tert-butyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-methoxy-9-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,8-difluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-isopropoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-chloro-9-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7,9-dichloro-4,5-dihydronaphtho[1,2-b-thiophene-3-carboxamide,
2-(acetylamino)-8-(trifluoromethyl)-4,5-dihydronaphtho[1,2-b] thiophene-3-carboxamide,
2-(acetylamino)-9-methoxy-6-(methylthio)-4,5-dihydronaphtho[1,2-b] -thiophene-3-carboxamide,

2-(acetylamino)-9-methoxy-6-(methyl-sulfonyl)-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-ethoxy-9-methoxy-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,9-diethoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8,9-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7,8-dinitro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-nitro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-methoxy-4,7-dimethyl-4,5-dihydronaphtho[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-6-methoxy-5-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-iodo-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7-chloro-8-nitro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5,8-dimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5-ethyl-8-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
benzyl 2-(acetyl-amino)-3-(amino-carbonyl)-4,5-dihydronaphtho[1,2-b]-thien-8-ylcarbamate,
2-(acetylamino)-8-amino-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2,8-bis(acetylamino)-4,5-dihydronaphtho-[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5-isopropyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6,9-dichloro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,

2-(acetylamino)-9-tert-butyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5,5,7-trimethyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5-butyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5-butyl-7-methyl-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5-butyl-7-fluoro-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5-butyl-7-methoxy-4,5-dihydronaphtho[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-5-butyl-7,8-dimethoxy-4,5-dihydronaphtho-[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-5,6-dihydro-4H-benzo-[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-9-chloro-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-8,9-dichloro-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-8-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-8-methoxy-6,6-dimethyl-5,6-dihydro-4H-benzo[6,7]cyclo-hepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-9-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-8,10-dimethoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta-[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-8-methyl-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-8,9-dimethoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide,

2-(acetylamino)-8-fluoro-9-methoxy-5,6-dihydro-4H-benzo[6,7]cyclo-hepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-7-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-6,9-dimethyl-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-10-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-9-bromo-6-methyl-5,6-dihydro-4H-benzo[6,7]cyclohepta-[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-7,8-dimethoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide,
2-(acetylamino)-6-methyl-5,6-dihydro-4H-benzo-[6,7]cyclohepta-[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-9-chloro-6-methyl-5,6-dihydro-4H-benzo[6,7]cyclohepta-[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-9-fluoro-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-9-methyl-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-9-amino-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-9-nitro-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-(acetylamino)-9,10-dimethoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta-[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]-thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-chloro-5,6-dihydro-4H-benzo[6,7]cyclo-hepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8,9-dichloro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,

2-[(aminocarbonyl)amino]-8-methoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-methoxy-6,6-dimethyl-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-methoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8,10-dimethoxy-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-methyl-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8,9-dimethoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-8-fluoro-9-methoxy-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7-methoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6,9-dimethyl-5,6-dihydro-4H-benzo[6,7]cyclo-hepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-10-methoxy-5,6-dihydro-4H-benzo[6,7]-cyclo-hepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-bromo-6-methyl-5,6-dihydro-4H-benzo[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-7,8-dimethoxy-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-6-methyl-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-chloro-6-methyl-5,6-dihydro-4H-benzo-[6,7]cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-fluoro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-methyl-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,

9-amino-2-[(aminocarbonyl)amino]-5,6-dihydro-4H-benzo-[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide,
2-[(aminocarbonyl)amino]-9-nitro-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide, and
2-[(aminocarbonyl)amino]-9,10-dimethoxy-5,6-dihydro-4H-benzo[6,7]-cyclohepta[1,2-b]thiophene-3-carboxamide.

13 A composition comprising the compound of claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12, and at least one pharmaceutically acceptable carrier.

14 A method of treating cancer, inflammation or an inflammation associated disorder in a subject, said method comprising administering to the subject having or susceptible to such cancer, inflammation or inflammation associated disorder, a therapeutically-effective amount of a compound of claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12.

15 The method of claim 14 for use in the treatment of cancer.

16 The method of claim 14 for use in the treatment of inflammation.

17 The method of claim 14 for use in the treatment of an inflammation-associated disorder.

18 The method of claim 17 wherein the inflammation-associated disorder is arthritis.

19 The method of claim 17 wherein the inflammation-associated disorder is pain

20 The method of claim 17 wherein the inflammation-associated disorder is fever.

21 The method of claim 17 wherein the inflammation-associated disorder is asthma.

22 The method of claim 17 wherein the inflammation-associated disorder is inflammatory bowel disease.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 03/22768

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D333/50 C07D333/80 A61P29/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
PAJ, WPI Data, EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 318 066 A (MERCK & CO INC) 31 May 1989 (1989-05-31) page 3, line 26 -page 3, line 29; claims 5-9; examples	1-22
A	US 4 775 691 A (LEUTWILER ALBERT) 4 October 1988 (1988-10-04) page 1, line 29 -page 1, line 35; claims; examples	1-22

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
27 October 2003	04/11/2003

Name and mailing address of the ISA	Authorized officer
European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tx. 31 651 epo nl, Fax: (+31–70) 340–3016	Schmid, A

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 03/22768

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. [X] Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 14-22 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

2. [X] Claims Nos.: 1-3,13, 14 (all partly)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210

3. [ ] Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. [ ] As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. [ ] As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. [ ] As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. [ ] No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

[ ] The additional search fees were accompanied by the applicant's protest.

[ ] No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-3,13, 14 (all partly)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claim(s) may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claims 1-3 is impossible. Consequently, the search has been restricted to the examples as regards claims 1-3, 13 and 14 and closely related compounds and claims 4-12.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 03/22768

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0318066 A	31-05-1989	DE 3873278 D1	03-09-1992
		DE 3873278 T2	25-02-1993
		EP 0318066 A1	31-05-1989
		JP 1186881 A	26-07-1989
		US 4999436 A	12-03-1991
US 4775691 A	04-10-1988	AT 323986 A	15-09-1992
		AU 600243 B2	09-08-1990
		AU 6617586 A	11-06-1987
		CA 1276158 C	13-11-1990
		CH 672129 B	31-10-1989
		DE 3641907 A1	11-06-1987
		DK 589886 A	10-06-1987
		ES 2003176 A6	16-10-1988
		FI 864978 A	10-06-1987
		FR 2595359 A1	11-09-1987
		GB 2183648 A ,B	10-06-1987
		GR 862863 A1	09-04-1987
		HU 46903 A2	28-12-1988
		IL 80906 A	31-01-1991
		IT 1214756 B	18-01-1990
		JP 1737086 C	26-02-1993
		JP 4022912 B	20-04-1992
		JP 62149672 A	03-07-1987
		LU 86706 A1	24-07-1987
		NL 8603098 A	01-07-1987
		NZ 218547 A	27-03-1990
		PH 24024 A	09-02-1990
		SE 8605255 A	10-06-1987
		ZA 8609296 A	27-07-1988