

(19) **DANMARK**

(10) **DK/EP 2274415 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 N 1/20 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-04-04**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-01-13**
- (86) Europæisk ansøgning nr.: **09755567.6**
- (86) Europæisk indleveringsdag: **2009-04-17**
- (87) Den europæiske ansøgnings publiceringsdag: **2011-01-19**
- (86) International ansøgning nr.: **US2009040920**
- (87) Internationalt publikationsnr.: **WO2009146205**
- (30) Prioritet: **2008-04-17 US 45915 P**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **DuPont Nutrition Biosciences ApS, Langebrogade 1 , Postboks 17, 1001 København K, Danmark**
- (72) Opfinder: **BELLOT, Marianne, Cain, 2640 N 73rd Street, Wauwatosa,WI 53213, USA**
MERTZ, Keith, J., N2533 Jefferson Road, Neosho,WI 53059, USA
REHBERGER, Thomas, G., 8422 Jackson Park Blvd, Wauwatosa,WI 53226, USA
- (74) Fuldmægtig i Danmark: **Marks & Clerk (Luxembourg) LLP, 44 rue de la Vallée, B.P. 1775, L-1017 Luxembourg, Luxembourg**
- (54) Benævnelse: **ANVENDELIGE BACILLESTAMMER TIL BEKÆMPELSE AF DYRELUGT**
- (56) Fremdragne publikationer:
US-A- 5 507 250
US-A- 5 945 333
US-A- 5 964 187
US-A1- 2007 202 088
JANSTOVA ET AL.: 'HEAT RESISTANCE OF BACILLUS spp. SPORES ISOLATED FROM COW'S MILK AND FARM ENVIRONMENT.' ACTA VET. BRNO, [Online] vol. 70, 2001, pages 179 - 184, XP008133200 Retrieved from the Internet: <URL:http://vfuu-www.vfu.cz/acta-vet/vo170/p df/70_179.pdf> [retrieved on 2009-05-29]

DESCRIPTION

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority under 35 U.S.C. §119(e) to U.S. Provisional Patent Application No. 61/045,915, filed April 17, 2008.

FIELD OF THE INVENTION

[0002] The invention relates to controlling odors associated with animals. More particularly, the invention relates to *Bacillus* strains for controlling odors associated with the bedding or other environment of animals, such as horses, guinea pigs, and the like and methods of making and using the *Bacillus* strains.

BACKGROUND OF THE INVENTION

[0003] Animals, including, but not limited to, companion animals, livestock, laboratory animals, working animals, and sport animals, generate odors that most people find offensive, especially when the odors are strong. For animals kept outdoors, this can lead to problems with neighbors, staff that work with the animals, and owners of the animals. Odors from animals kept indoors can aggravate the animal's owners, those living in the same household of the indoor animal, and visitors to the household where the animal is kept.

[0004] Ammonia and other odors coming from farms, such as horse farms, can aggravate neighbors, especially in residential areas. The ability to effectively control and alleviate waste odors associated with equine in urban areas provides a healthier living environment and helps horse enthusiasts to maintain good relationships with their neighbors.

[0005] Ammonia negatively affects the health of stabled horses worldwide. Even long-term, low level exposure to ammonia can affect a horse's respiratory health and immune response. (Davis, M. S. Foster, W.M. "Inhalation Toxicology in the Equine Respiratory Tract." In: Equine Respiratory Disease, P. Lekeus. International Veterinary Information Service, 2002). Horse bedding consists of materials such as straw, sand and other material. Wood shavings are commonly used as bedding material to help absorb urine and provide a surface for easy clean up. The condition or quality of bedding in equine stalls can be affected by a number of factors including frequency of clean-out, ventilation, moisture and temperature. Due to the health problems associated with high levels of ammonia, proper barn management is crucial. Daily cleaning, good ventilation and complete clean-out of stalls when bedding is too soiled are essential to control ammonia exposure. Due to the cost of bedding materials, most horse owners clean stalls daily but prefer not to strip stalls on a regular basis. There is great value in extending the bedding life and the amount of time that bedding remains in the stall. By reducing the ammonia odors in the bedding, bedding life can be extended, resulting in a great cost savings.

[0006] There are more horses in the United States currently than there were in the 1800's, many of them kept in urban areas. In this setting, the manure and urine from horses can be offensive to non-horse owners and can be a problem for municipal landfills. Ammonia production can also cause performance loss in competitive horse events like horse racing.

[0007] Some products to treat horse waste material are available, but most are chemically based and only provide short term relief of odors and no increased digestion of solid waste. Some products control ammonia with absorbents, such as clay and zeolite. These products function by providing negatively charged exchange sites to attract ammonium ions. In this process, more weakly bound ions such as hydrogen and sodium are replaced by ammonium ions, reduction the total concentration of ammonia in solution. These products also bind water, reducing the microbial activity and therefore the breakdown of urea to ammonia. Other products control ammonia via yucca additives and fragrances that simply mask the odor. Ammonia negatively affects the health of caged small animal pets as well.

[0008] Guinea pigs, hamsters, rats and other small animals need soft clean bedding that is changed frequently in order for the animal to stay in top health. Bedding is replaced to keep down ammonia, and to keep small animals and the cage clean. Ammonia is a component of urine. In high concentrations not only does it smell offensive to humans, but it can eventually lead to respiratory problems for the small animals. Breathing concentrated ammonia will damage their lungs, burn their esophagus and create other

health problems. Therefore, reducing ammonia levels is important for small animals. There is also a great value and need in extending bedding life.

[0009] Ammonia concentrations in poorly ventilated horse barns and cages can rise to levels potentially harmful to the equine and small animal respiratory tract. High levels of ammonia have been associated with foal pneumonia (McMillan K: Foal pneumonia: An Illinois survey. *An Health and Nutrit* 1986; 34). High levels of ammonia may also predispose horses to chronic obstructive pulmonary disease. (Tanner MK, Swinker AM, Traub-Dargatz JL, Stiffler LA, McCue PM, Vanderwall DK, Johnson KE: Respiratory and environmental effects of recycled phone book paper versus sawdust as bedding for horses. *J Eq Vet Sci* 1998; 468-476). Ammonia and odor smells coming from horse farms can aggravate neighbors, especially in residential areas.

[0010] The ammonia present in equine facilities and small animal cages is the product of microbial decomposition of excreted nitrogenous compounds. This includes urea, nonabsorbed proteins, amino acids, and nonprotein nitrogen present in the diet.

[0011] In view of the foregoing, it would be desirable to provide one or more *Bacillus* strains to treat or prevent animal odors and to provide animal bedding including one or more of these strains, methods of using the *Bacillus* strains and animal bedding treated with the *Bacillus* strains.

SUMMARY OF THE INVENTION

[0012] The invention, which is defined by the claims set out at the end of this disclosure, is intended to solve at least some of the problems noted above. A composition is provided that includes *Bacillus subtilis* 2084 (NRRL B-50013), *B. subtilis* 27 (NRRL B-50105) and *B. licheniformis* 21 (NRRL B-50134).

[0013] In some embodiments of the composition, at least one carrier capable of binding ammonia is included with the *Bacillus* strains. In some embodiments of the composition, at least one water-soluble carrier is included with the *Bacillus* strains.

[0014] Animal bedding that includes the *Bacillus* strains is also provided, as is a method of making the animal bedding.

[0015] Also provided are methods of controlling odors from animal waste. In some embodiments of these methods, at least one carrier capable of binding ammonia and the *Bacillus* strains are used to control odors. In some embodiments of these methods, at least one water-soluble carrier and the *Bacillus* strains are used to control odors.

[0016] A method of making a composition is also provided. In it, each of *Bacillus subtilis* 2084 (NRRL B-50013), *B. subtilis* 27 (NRRL B-50105) and *B. licheniformis* 21 (NRRL B-50134) is grown in a broth. Each of the strains is separated from its broth to make the composition.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017] Preferred exemplary embodiments described herein are illustrated in the accompanying drawings, in which like reference numerals represent like parts throughout and in which:

FIG. 1 is a graph of ammonia concentration at 48 hours, with bars with different letters being significantly different.

FIG. 2 is a graph of the reduction in ammonia levels in bedding with FreshShield™ product or FreshShield™ stall treatment product.

FIG. 3 is a graph of smell panel results after 5, 23, and 29 hours, showing reduction in ammonia smell with FreshShield™ product treatment.

FIG. 4 is a graph showing the reduction in ammonia after one week of FreshShield™ product treatment.

FIG. 5 is a graph showing the difference in reduction of ammonia levels from used guinea pig bedding after 7 days of treatments at different concentrations of FreshShield™ product compared to after 7 days with no treatment.

FIG. 6 is a graph showing reductions in ammonia levels after 24 hours in manure treated with FreshShield stall treatment product, Stall Fresh stall treatment product, Stall DRY Plus stall treatment product, and Sweet PDZ stall treatment product.

[0018] Before explaining embodiments described herein in detail, it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the drawings. The invention is capable of other embodiments or being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

DETAILED DESCRIPTION

[0019] Provided herein are combinations of three *Bacillus* strains that can be used to form odor-controlling compositions for controlling ammonia and other odors from non-human animals. Laboratory studies have shown a 50-60% reduction in ammonia on wood shavings treated with one or more compositions provided herein. Additional benefits of using the compositions to treat or prevent animal odors can include one or more of the following.

Compositions:

[0020] *Bacillus* strains have many qualities that make them useful for treating and preventing animal odors. For example, *Bacillus* strains produce extracellular enzymes, such as proteases, amylases, and cellulase. In addition, *Bacillus* strains produce antimicrobial factors, such as gramicidin, subtilin, bacitracin, and polymyxin. Several *Bacillus* species also have GRAS status, i.e., they are generally recognized as safe by the US Food and Drug Administration and are also approved for use in animal feed by the Association of American Feed Control Officials (AAFCO). All *B. subtilis* strains are GRAS.

[0021] The *Bacillus* strains described herein are aerobic and facultative sporeformers and thus, are stable. *Bacillus* species are the only sporeformers that are considered GRAS. *Bacillus* strains preferably are used as spores. When rehydrated, with liquid containing nutrients, the *Bacillus* strains can go into the vegetative state. However, when again dehydrated or without enough nutrients, the *Bacillus* strains will go back to spores.

[0022] *Bacillus* strains found to be useful in the compositions provided herein include *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21. *B. subtilis* strains 2084 and 27 were deposited on March 8, 2007 and January 24, 2008, respectively, at the Agricultural Research Service Culture Collection (NRRL), 1815 North University Street, Peoria, Illinois, 61604 and given accession numbers NRRL B-50013 and NRRL B-50105, respectively. On April 15, 2008, *B. licheniformis* strain 21 was deposited at NRRL and given accession number NRRL B-50134. All of these deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure.

[0023] The compositions provided herein include *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21 in an effective amount to control odors. In at least one embodiment of the composition, substantially equal amounts (based on colony forming units (cfu)) of each strain are used. However, differing amounts can also be used. For example, each strain can be added from about 10% to about 45% (based on cfu) of the total amount of strain in the composition. In an exemplary embodiment, *B. subtilis* strain 27 is included at 10%, *B. subtilis* strain 2084 is included at 45%, and *B. licheniformis* strain 21 is included at 45%.

[0024] In at least one embodiment of the composition, the total cfu of the *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21 is from about 7.5×10^7 to about 7.5×10^8 in the final form of the composition. For example, about 2.5×10^8 total cfu of the *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21 can be included in the final form of the composition.

[0025] In at least some embodiments of the composition, one or more water-soluble carrier, which is useful for rehydrating the strains, is added to the strains. In one embodiment, carriers include Baker's sugar, maltodextrin M100, and baylith. This embodiment, i.e., the three strains, *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21, in substantially equal amounts (based on CFU) of each strain and the carriers, is referred to herein as FreshShield™ product. In at least some embodiments, FreshShield™ product is applied to animal bedding. This can be done during the manufacturing of bedding or at other times, as is explained in more detail below.

[0026] In another embodiment, carriers that bind ammonia are used. In one embodiment, these carriers include bentonite, Fuller's earth, mineral oil, and baylith. These carriers in combination with the three strains, *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21, in substantially equal amounts (based on CFU) of each strain, Baker's sugar, and baylith are referred to

herein as FreshShield™ stall treatment product. Other carriers, such as clintpolite, diamaceous earth, beolite clay, and limestone, also bind ammonia, and can also be used.

Making and Using the Compositions:

[0027] The *Bacillus* strains are grown in a liquid nutrient broth. In at least one embodiment, the *Bacillus* strains are grown to a level at which the highest number of spores are formed.

[0028] The *Bacillus* strains are produced by fermentation of the bacterial strains. Fermentation can be started by scaling-up a seed culture. This involves repeatedly and aseptically transferring the culture to a larger and larger volume to serve as the inoculum for the fermentation, which is carried out in large stainless steel fermentors in medium containing proteins, carbohydrates, and minerals necessary for optimal growth. A non-limiting exemplary medium is TSB. After the inoculum is added to the fermentation vessel, the temperature and agitation are controlled to allow maximum growth. Once the culture reaches a maximum population density, the culture is harvested by separating the cells from the fermentation medium. This is commonly done by centrifugation.

[0029] In one embodiment, to prepare the *Bacillus* strains, each *Bacillus* strain is fermented to a 5×10^8 CFU/ml to about 4×10^9 CFU/ml level. In at least one embodiment, a level of 2×10^9 CFU/ml is used. The bacteria are harvested by centrifugation, and the supernatant is removed. The pelleted bacteria can then be applied to the wood shavings. In at least some embodiments, the pelleted bacteria are freeze-dried and mixed with a carrier before they are applied to the wood shavings. However, it is not necessary to freeze-dry the *Bacillus* before using them. The strains can also be used with or without preservatives, and in concentrated, unconcentrated, or diluted form.

[0030] The count of the culture can then be determined. CFU or colony forming unit is the viable cell count of a sample resulting from standard microbiological plating methods. The term is derived from the fact that a single cell when plated on appropriate medium will grow and become a viable colony in the agar medium. Since multiple cells may give rise to one visible colony, the term colony forming unit is a more useful unit measurement than cell number.

[0031] To prepare compositions described herein, the cultures and carriers (where used) can be added to a ribbon or paddle mixer and mixed for about 15 minutes, although the timing can be increased or decreased. The components are blended such that a uniform mixture of the cultures and carriers result. The final product is preferably a dry, flowable powder.

[0032] In a non-limiting example of the FreshShield™ stall treatment product, the composition is applied to the surface on which an animal is kept, such as in an animal stall. For example, the composition is used as a topical treatment in a stall to treat wet spots. The FreshShield™ stall treatment product can be applied after wet bedding is removed. Clean bedding can be added on top of the FreshShield™ stall treatment product and mixed with the FreshShield™ stall treatment product. This treatment can be done on a daily basis, if desired, or more (or less) frequently.

[0033] In a non-limiting example of the FreshShield™ product, the compositions are added on to the bedding while the bedding is being manufactured, such as just prior to packaging. This provides the convenience of having bedding material that reduces odor and ammonia emissions and extends the bedding life.

[0034] In at least one embodiment of a method of making the FreshShield™ product, the bacteria are mixed into a liquid and applied onto bedding, such as wood shavings.

[0035] Different application rates of the three-strain combination for making the FreshShield™ product can be used. In at least one embodiment, the application rate for bedding for large animals (including, but not limited to, equine, bovine, ovine, and porcine) differs from the rate for bedding for small animals (including, but not limited to, hamster, rodents, guinea pig, rabbits, reptiles, dogs, and birds).

[0036] For example, for large animal bedding, a total of 2.9×10^{10} total cfu of bacteria per cubic foot of wood shavings is used. The bacteria are added along with the carriers, M100 and Baker's sugar, to the wood shavings. However, the application rate can vary from 2.9×10^8 to 2.9×10^{11} total cfu per cubic foot of wood shavings. In at least one embodiment, the *Bacillus* strains are at a concentration of 8.8×10^{10} total cfu per gram of bacteria (each strain being 1/3 of this). In an exemplary embodiment, the *Bacillus* strains are combined with the wood shavings by mixing 10 kg of *Bacillus* with 55 gallons of water to form a stock solution. A total of

7 ml of stock solution is diluted with 53 ml of water and applied via an auger to one cubic foot of wood shavings prior to the bagging of the wood shavings. This diluted solution treats 10,000 bags of wood shavings, with one bag of shavings being approximately 3.0 cubic foot compressed and 9 cubic feet uncompressed. A typical horse stall (10'x10') uses 4 bags (3 cubic foot each) of shavings, which delivers 3.5×10^{11} cfu/gram of *Bacillus* when applied as above.

[0037] In at least one embodiment, the application for small animal bedding is a total of 5.8×10^{10} total cfu per cubic foot of wood shavings. The bacteria are added along with the carriers, M100 and Baker's sugar, to the wood shavings. However, the application rate can vary from 5.8×10^9 to 5.8×10^{11} total cfu per cubic foot of wood shavings. In at least one embodiment, the *Bacillus* strains are at a concentration of 1.7×10^{11} total cfu per gram of bacteria (each strain being 1/3 of this). In at least some embodiments, the *Bacillus* combination is added to the wood shavings by mixing 10 kg of *Bacillus* with 55 gallons of water to form a stock solution. In an exemplary embodiment, 7 ml of stock solution is diluted with 53 ml of water and applied to one cubic foot of wood shavings as they are being conveyed via an auger prior to bagging of the wood shavings. This diluted solution treats the equivalent of 30,000-10,000 bags of wood shavings, with one bag of shavings being approximately 1-3 cubic foot compressed and 3-9 cubic feet uncompressed, respectively. Small animal bedding materials come in a variety of sizes. Any dilution that would achieve a desired cfu per cubic foot is suitable within the scope of this invention.

[0038] An exemplary composition of FreshShield™ product is shown below in Table 1.

Table 1.

Ingredient	Amount added	% composition by weight
<i>B. licheniformis</i> 21 at 3.8×10^{11} cfu/g	0.925 kg	2.70*
<i>B. subtilis</i> 27 at 3.0×10^{11} cfu/g	1.172 kg	9.50*
<i>B. subtilis</i> 2084 at 2.0×10^{11} cfu/g	1.764 kg	3.53*
Baker's sugar	3.069 kg	42.13
Maltodextrin M100	3.008 kg	41.29
Baylith	0.061 kg	0.85
Total	10 kg	100%
*This is for a specific lot of <i>Bacillus</i> . The amount added varies depending on the concentration of bacteria.		

[0039] An exemplary composition of FreshShield™ stall treatment product is shown below in Table 2.

Table 2.

Ingredient	Amount added	% composition by weight
<i>B. licheniformis</i> 21 at 5.0×10^{11} cfu/g	0.001 kg	0.02*
<i>B. subtilis</i> 27 at 5.0×10^{11} cfu/g	0.001 kg	0.02*
<i>B. subtilis</i> 2084 at 5.0×10^{11} cfu/g	0.001 kg	0.02*
Bentonite	3.176 kg	69.75
Fullers Earth	1.361 kg	29.886
Mineral Oil	0.009 kg	0.197
Baylith	0.005 kg	0.109
Total	4.554 kg	100%
*This is for a specific lot of <i>Bacillus</i> . The amount added varies depending on the concentration of bacteria.		

[0040] Additional uses of the compositions include the following. The composition can be added to as an ingredient to existing stall treatment products as an all-natural additive for further odor control. The compositions can also be used with alternative bedding products, e.g., corn cob bedding, paper bedding, straw, and wood pellets, used in the small and large animal market. For use with alternative bedding products, the compositions are applied as is described above for wood shavings or in other ways.

[0041] Further uses of the compositions include using them on an absorbent material, such as granulated clay, for covering the floor of an animal's cage or excretory box. For example, the compositions can be used as a litter additive for cat litter boxes and

litter boxes for other animals. In one embodiment of the litter additive, *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21 and carriers are combined. In an exemplary embodiment, bentonite, Fuller's earth, mineral oil, and baylith are the carriers. The cat litter additive can have the same formula as shown in Table 2 or it can have a different formula. The cat litter additive can be applied to cat litter as needed for odor control. The cat litter additive can also be used as an ingredient for litter manufacturing or as a topical treatment.

[0042] Routine use of the one or more compositions provided herein can dramatically reduce and even eliminate animal odors. The compositions are ecologically friendly products because instead of chemicals, bacteria are used to control animal odors. The compositions are considered "green," that is, something good for the environment. The compositions are all natural and safe. The compositions are highly stable, therefore, they have a long shelf life. In addition, the compositions extend bedding life. This is increasingly important as wood and other bedding sources are becoming more expensive and more difficult to obtain. When one or more compositions is added to animal bedding, the treated bedding is easy to use because nothing extra needs to be added to the bedding for odor control. The compositions also provide a healthier living environment for animals living on bedding or other surfaces treated with one or more composition. Animals do not like living in ammonia-rich environments. Ammonia burns their eyes and causes other problems such as respiratory problems. Therefore, reductions in ammonia and other odors improve the animal's quality of life. In addition, more efficient composting of animal manure is accomplished by using the compositions provided herein.

[0043] A healthier working environment is provided for workers that train and take care of animals. For instance, ammonia, which irritates the people's lungs and can cause respiratory and other problems, is reduced.

[0044] The compositions can be easily applied to wood shavings. It also provides a surface for the *Bacillus* bacteria to live. Unlike simply adding untreated wood shavings, wood shavings treated with a composition provided herein adds the waste digesting power of the *Bacillus* bacteria, which reduces the odor and ammonia.

[0045] Although not intended to be a limitation to the present disclosure, it is believed that inhibition of odors is accomplished via reduced volatilization of ammonia through controlled decomposition of nitrogen-containing compounds and utilization of the nitrogen by the *Bacillus* strains. Where used, the carriers limestone, clay, zeolite, diatomaceous earth, clintpolite, bentonite, and Fuller's earth, also bind ammonia and therefore reduce odors.

EXAMPLES

[0046] The following Examples are provided for illustrative purposes only. The Examples are included herein solely to aid in a more complete understanding of the presently described invention. The Examples do not limit the scope described herein described or claimed herein in any fashion.

Example 1

[0047] Overview: This research was conducted to determine the effectiveness of a horse waste product at lessening the odor associated with waste excretion in confined areas such as the common equine box stall.

[0048] Protocol: 12 horses housed in a barn were utilized to "soil" the stalls. Horses/stalls were blocked by age/ration/sex so that there were similar types of horses within each treatment. Horses were stalled for approximately 12 hours/day with daily turnout. The treatments were as follows: *Control:* Consisted of normal bedding practices per the protocols of the facility with 6 stalls and *Bedding Treatment:* In addition to normal bedding protocols, a top dress of a "horse waste product" applied to shavings was applied to selected stalls with 6 stalls. Bedding treatment was applied by spreading 1 (one) scoop (in this example, one (1) scoop equals 8.64×10^{10} total cfu of the *Bacillus* strains) of treatment material, which is prepared to contain 2.4×10^8 total cfu/g of *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21 bacteria provided in FreshShield™ product, that is, one scoop per 1 bag (2.0 cubic foot) of Marth Easy Pick™ shavings.

[0049] The trial initiation had a 3 week trial period. At the beginning of the trial all stalls were cleaned out of all waste and bedding material.

[0050] Control stalls were re-bedded with 7 bags each of untreated Marth Easy Pick Shavings. Treated stalls had 7 scoops of FreshShield™ product on 7 Marth Easy Pick™ 2.0 cubic foot bags of wood shavings, applied and thoroughly mixed in the

bedding. Treated stalls were uniquely marked so farm personnel could easily identify treatment stalls. For the remainder of the trial, bedding protocols were adhered to per the facility managers direction. One scoop of FreshShield™ product was applied per bag of fresh shaving material put down in the stall. Fresh FreshShield™ product was mixed in with fresh shavings at time of application. Individual stall records were maintained regarding the amount of fresh bedding put down in each stall on a daily basis. The amount of deodorizer, such as Sweet PDZ or Stall Dry, was applied daily was recorded.

[0051] Results: The FreshShield™ product reduced odors in the treated stalls of the barn. The test was done during the time of year that the barn was open. The FreshShield™ product reduced stall odors and ammonia smells, making the facility and environment better for the horses.

Example 2

[0052] Aim: To determine if FreshShield™ product will modulate the decomposition of manure mixed with Marth Easy Pick™ Bedding over time. This model utilized cow urine and cow fecal material.

[0053] Protocol: A waste *in vitro* model was prepared by placing 100g of Marth Easy Pick™ bedding into each of fifteen 6"x9"x3" Ziploc™ containers with a small hole drilled in one end. The hole was taped to reseal the container. 500 ml of freshly collected cow urine was added to each container followed by 50g of freshly collected cow fecal matter. Three treatments were designated across the 15 containers with 5 replicates/treatment. Treatments consisted of control, containing no FreshShield™ product, and FreshShield™ product added at a 1X dose and a 2X dose. The 1X dose contained 3.12×10^9 CFU of the three *Bacillus* strains, i.e., *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21, with the 2X dose containing twice that amount. After addition of all materials to the Ziploc containers, contents of the containers were thoroughly mixed using a plastic teaspoon, with different spoons used to mix each treatment. Ziploc containers were sealed by placing the lid on top and placed in the Bio-Cold environmental chamber set at 81°F and 55% humidity.

[0054] Ammonia measurements were taken from each container at 24 hours, 48 hours, and 7 days after placement in the Bio-Cold chamber. Measurements were taken using the Drager pump fitted with disposable ammonia tubes.

[0055] Results: Ammonia concentrations at 48 hours can be seen in FIG. 1. Results from ammonia concentration after 7 days were also observed. There was no significant difference between treatments at 24 hours. At 48 hours the 2X dose was significantly less than that of the control, but not significantly different for the 1X dose. The 1X dose was less than that of the control but not significant. There were no significant differences between treatment groups after seven days.

Example 3

[0056] Study Details: The following study was performed to show the effect of bedding containing the FreshShield™ product and having Baker's sugar, maltodextrin M100, and baylith as the carriers (formulated as shown in Table 1 above) and bedding treated with FreshShield™ stall treatment product having bentonite, Fuller's earth, mineral oil, and baylith as the carriers (formulated as shown in Table 2 above) to determine the effect of carriers on the ability of the product to control ammonia production. Both products included the three strains, *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21, as shown in Tables 1 and 2. Soiled horse bedding material collected from a stable was placed into sealed containers. Treatments were applied at a rate of 1.0×10^{10} total cfu of the three strains of the bacteria in the FreshShield™ product per 500 grams of soiled bedding and at a rate of 1.0×10^{10} total cfu of the three strains of bacteria in the FreshShield™ stall treatment product. Ammonia measurements were taken at time 0, 6, and 16 hours post-treatment using a Draeger ammonia detector.

[0057] Results: FIG. 2 shows an average ammonia reduction of 50% in bedding with FreshShield™ product and 62% in bedding with FreshShield™ stall treatment product

Example 4

[0058] Study Details: The following study was performed to show the effect of FreshShield™ product on ammonia production in soiled pea gravel. Soiled pea gravel was placed into sealed containers. The composition of the FreshShield™ product is shown in Table 3 below. Control samples received no FreshShield™ product. Treatments were applied as in Example 3 for the

FreshShield™ stall treatment product. A smell panel evaluated the ammonia odors over time. Ammonia measurements were taken at 1 week post-treatment using a Draeger ammonia detector.

Table 3.

Ingredient	Amount added	% composition by weight
<i>B. licheniformis</i> 21 at 5.0x10 ¹¹ cfu/g	0.001 kg	0.02*
<i>B. subtilis</i> 27 at 5.0x10 ¹¹ cfu/g	0.001 kg	0.02*
<i>B. subtilis</i> 2084 at 5.0x10 ¹¹ cfu/g	0.001 kg	0.02*
Rice hulls	4.546 kg	69.75
Baylith	0.005 kg	0.109
Total	4.554 kg	100%

[0059] Results: The smell panel noted a marked reduction in ammonia smell with FreshShield™ product, at least a 50% reduction at 5, 23, and 29 hours after initial treatment as compared to the control samples (FIG. 3). The reduction in measurable ammonia was seen after one week of treatment, and no noticeable ammonia odors were present. In comparison, after one week the control containers still had offensive levels of ammonia. FIG. 4 shows that after one week, the average ammonia levels were 225 ppm for the control and less than 20 ppm in the treated containers.

Example 5

[0060] Initial Visit: A commercial guinea pig farm tested the FreshShield™ product at rates of 1.45x10¹⁰ total cfu per cubic foot of bedding (0.5x), 2.9x10¹⁰ total cfu per cubic foot of bedding (1.0x) and 5.8x10¹⁰ total cfu per cubic foot of bedding (2.0x), to measure ammonia reduction within a guinea pig commercial facility. The FreshShield™ product included the three strains of bacteria, *B. subtilis* strains 2084 and 27 and *B. licheniformis* strain 21. During the initial visit, control samples were collected from pens that were to be treated and used within the experiment. For each of nine pens, one composite sample was formed by collecting the same quantity from a similar location in each pen and combining it into one sample bag.

[0061] Forty-eight hours after collection, ammonia readings were taken and recorded. After composite samples were collected, the appropriate concentration of product to be tested was applied to clean fresh bedding. 1.45x10¹⁰ total cfu per cubic foot of bedding (0.5x), 2.9x10¹⁰ total cfu per cubic foot of bedding (1.0x) and 5.8x10¹⁰ total cfu per cubic foot of bedding (2.0x), by spraying the concentrated liquid on to the clean bedding. Treatments included a control (no FreshShield™ product), 0.5X FreshShield™ product, 1.0X FreshShield™ product, and 2X FreshShield™ product. Guinea pigs were allowed to maintain normal living conditions on treated bedding.

[0062] Follow-up Visit: One week following the initial treatment of bedding treated with the *Bacillus* shaving product, used bedding samples were collected. Again, the same quantity of bedding from similar locations was collected from each pen. Three pens were combined into one composite sample. Three replicate composite samples were collected and combined for each treatment. Forty-eight hours after collection ammonia readings were taken and recorded.

[0063] Results: The results show that ammonia readings from used guinea pig bedding with no treatment were higher than ammonia readings from used guinea pig bedding containing 0.5X, 1X and 2X *Bacillus* treatment levels. The difference between non-treated and treated bedding taken from the same pens after 7 days of treatment were decreased compared to the control difference, as shown in FIG. 5.

Example 6

The Problem:

[0064] Ammonia negatively affects the health of stabled horses worldwide. Even long-term, low level exposure to ammonia can affect a horse's respiratory health, immune response, and the health of those people working in and around horses. FreshShield

is an all natural, product that proactively stops ammonia production. The active ingredient in FreshShield stall treatment product starts working when applied to urine and/or feces to reduce harmful ammonia emissions. There are a number of stall treatment products available, but most work by masking the ammonia only for a short time and not by attacking the underlying cause of high ammonia.

Study Details:

[0065] The following study was performed to compare the effect of FreshShield™ stall treatment product on ammonia production to three other stall treatment products: Stall Fresh, Stall DRY Plus, and Sweet PDZ stall treatment products. Soiled horse bedding collected from a stable was placed into sealed containers. The FreshShield™ stall treatment product was formulated as in Table 1 above.

[0066] Treatments of the FreshShield™ stall treatment product were applied by applying 4 oz. of the product daily. The other products were applied according to each of their label directions. Ammonia measurements were taken at time 0 and 24 hours post-treatment using a Draeger ammonia detector.

The Results:

[0067] As is shown in FIG. 6, after 24 hours, ammonia was reduced by 70% in manure treated with FreshShield stall treatment product, by 22.2% in manure treated with Stall Fresh stall treatment product, by 49.8% in manure treated with Stall DRY Plus stall treatment product, and by 38.9% in manure treated with Sweet PDZ stall treatment product. The other products worked initially, but then lost their effectiveness over time. FreshShield stall treatment product kept working long after the other products stop. FreshShield stall treatment product worked well with both high and low initial ammonia levels, and reduced the ammonia by the greatest amount overall.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US61045915A](#) [0001]

Non-patent literature cited in the description

- Inhalation Toxicology in the Equine Respiratory Tract **DAVIS, M. S.FOSTER, W.M.**Equine Respiratory DiseaseInternational Veterinary Information Service20020000 [0005]
- **MCMILLAN K**Foal pneumonia: An Illinois surveyAn Health and Nutrit, 1986, 34- [0009]
- **TANNER MKSWINKER AM TRAUB-DARGATZ JLSTIFFLER LAMCCUE PMVANDERWALL DKJOHNSON KE**Respiratory and environmental effects of recycled phone book paper versus sawdust as bedding for horsesJ Eq Vet Sci, 1998, 468-476 [0009]

ANVENDELIGE BACILLESTAMMER TIL BEKÆMPELSE AF DYRELUGT

PATENTKRAV

1. Sammensætning omfattende:
Bacillus subtilis 2084, der er deponeret som B-50013,
5 *B. subtilis* 27, der er deponeret som NRRL B-50105, og
B. licheniformis 21, der er deponeret som B-50134.
2. Sammensætning ifølge krav 1, hvor mængden af hver stamme omtrent er ens.
3. Sammensætning ifølge krav 1 eller krav 2, hvilken sammensætning endvidere omfatter mindst én bærer, der kan binde ammoniak, hvor bæreren fortrinsvis er udvalgt fra gruppen bestående af kalksten, ler,
10 zeolit, diatoméjord, clintpolit, bentonit og blegejord.
4. Sammensætning ifølge krav 1 eller krav 2, hvilken sammensætning endvidere omfatter bentonit, blegejord, mineralolie og baylith.
5. Sammensætning ifølge krav 1 eller krav 2, hvilken sammensætning endvidere omfatter mindst én vandopløselig bærer, hvor bæreren fortrinsvis er udvalgt fra gruppen bestående af superfint sukker,
15 maltodextrin M100 og baylith.
6. Sammensætning ifølge et hvilket som helst af de foregående krav, hvilken sammensætning endvidere omfatter et absorberende materiale, hvor det absorberende materiale fortrinsvis er kattegrus.
7. Sammensætning ifølge et hvilket som helst af kravene 1 til 5, hvilken sammensætning endvidere omfatter dyrestrøelse.
- 20 8. Sammensætning ifølge krav 7, hvor dyrestrøelsen er udvalgt fra gruppen bestående af høvlspåner, majscolbestrøelse, papirstrøelse, strå og træpiller.
9. Fremgangsmåde til bekæmpelse af lugte og ammoniak fra dyreaffald, hvilken fremgangsmåde omfatter påføring af sammensætningen ifølge krav 3 eller krav 7 på et område, hvor dyret er installeret, hvor påføring af sammensætningen bekæmper lugte og ammoniak i dyreaffaldet.
- 25 10. Fremgangsmåde til fremstilling af en sammensætning, hvilken fremgangsmåde omfatter:
dyrkning af hver af *Bacillus subtilis* 2084, der er deponeret som NRRL B-50013; *B. subtilis* 27, der er deponeret som NRRL B-50105, og *B. licheniformis* 21, der er deponeret som NRRL B-50134, i en bouillon og
separation af hver af stammerne fra deres bouillon for fremstilling af sammensætningen.
- 30 11. Fremgangsmåde ifølge krav 10, hvilken fremgangsmåde endvidere omfatter frysetørring af hver stamme; og eventuelt tilsætning af den frysetørrede stamme til mindst én vandopløselig bærer eller mindst én bærer, der kan binde ammoniak.
12. Fremgangsmåde ifølge krav 10, hvilken fremgangsmåde endvidere omfatter påføring af stammerne på dyrestrøelse.
- 35 13. Fremgangsmåde ifølge krav 12, hvilken fremgangsmåde endvidere omfatter:
tilsætning af en vandopløselig bærer til stammerne og
rehydrering af stammerne, hvor de rehydrerede stammer påføres dyrestrøelse.

-2-

14. Fremgangsmåde ifølge krav 12, hvor stammerne påføres for at tilvejebringe i alt væsentligt den samme cfi for hver stamme.
15. Fremgangsmåde ifølge krav 12, hvilken fremgangsmåde endvidere omfatter emballering af dyrestrøelsen, hvor stammerne påføres for emballering af dyrestrøelsen; fortrinsvis hvor dyrestrøelsen er
5 udvalgt fra gruppen bestående af høvlspåner, majscolbestrøelse, papirstrøelse, strå og træpiller.

DRAWINGS




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6