



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2606206 A1 2006/11/02

(21) **2 606 206**

(12) **DEMANDE DE BREVET CANADIEN  
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2006/04/24  
(87) Date publication PCT/PCT Publication Date: 2006/11/02  
(85) Entrée phase nationale/National Entry: 2007/10/16  
(86) N° demande PCT/PCT Application No.: EP 2006/003809  
(87) N° publication PCT/PCT Publication No.: 2006/114273  
(30) Priorités/Priorities: 2005/04/26 (US11/114,301);  
2005/06/14 (EPPCT/EP2005/006461);  
2005/12/16 (US11/367,601)

(51) Cl.Int./Int.Cl. *A61K 39/12* (2006.01)  
(71) Demandeur/Applicant:  
GLAXOSMITHKLINE BIOLOGICALS S.A., BE  
(72) Inventeurs/Inventors:  
DUBIN, GARY, US;  
INNIS, BRUCE, US;  
SLAOUI, MONCEF MOHAMMED, US;  
WETTENDORFF, MARTINE ANNE CECILE, BE  
(74) Agent: OGILVY RENAULT LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : VACCIN  
(54) Title: VACCINE

(57) **Abrégé/Abstract:**

This invention pertains to methods and vaccines for treating infections caused by human papillomaviruses. It has been determined that immunization with HPV16 and HPV 18 virus like particles provides cross-protection against other HPV types.



## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
2 November 2006 (02.11.2006)

PCT

(10) International Publication Number  
**WO 2006/114273 A3**

(51) International Patent Classification:  
*A61K 39/12* (2006.01)

(21) International Application Number:  
PCT/EP2006/003809

(22) International Filing Date: 24 April 2006 (24.04.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
11/114,301 26 April 2005 (26.04.2005) US  
PCT/EP2005/006461 14 June 2005 (14.06.2005) EP  
11/367,601 16 December 2005 (16.12.2005) US

(71) Applicant (for all designated States except US): **GLAXO-SMITHKLINE BIOLOGICALS S.A.** [BE/BE]; Rue De L'institut 89, B-1330 Rixensart (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DUBIN, Gary** [US/US]; GlaxoSmithKline, 2301 Renaissance Blvd., PO Box 61540, King Of Prussia, Pennsylvania 19406-2772 (US). **INNIS, Bruce** [US/US]; GlaxoSmithKline, 2301 Renaissance Blvd., PO Box 61540, King Of Prussia, PA 19406-2772 (US). **SLAOUI, Moncef Mohammed** [BE/US]; GlaxoSmithKline, 2301 Renaissance Blvd., PO Box 61540, King Of Prussia, Pennsylvania 19406-2772 (US). **WETTENDORFF, Martine Anne Cecile** [BE/BE]; GlaxoSmithKline Biologicals sa, Rue De L'institut 89, B-1330 Rixensart (BE).

(74) Agent: **STEPHEN, Robert**; GlaxoSmithKline, Corporate Intellectual Property (CN925.1), 980 Great West Road, Brentford Middlesex TW8 9GS (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

#### Published:

- with international search report

(88) Date of publication of the international search report:  
15 March 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: VACCINE

(57) Abstract: This invention pertains to methods and vaccines for treating infections caused by human papillomaviruses. It has been determined that immunization with HPV16 and HPV 18 virus like particles provides cross-protection against other HPV types.

WO 2006/114273 A3

## DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME            1    DE    2  
CONTENANT LES PAGES    1    À    33

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

## JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME            1    OF    2  
CONTAINING PAGES    1    TO    33

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:



## VACCINE

### Field of the Invention

5           The present invention relates to human papillomavirus (HPV) vaccines.

### Background of the Invention

Papillomaviruses are small DNA tumour viruses, which are highly species specific. So far, over 100 individual human papillomavirus (HPV) genotypes have  
10   been described. HPVs are generally specific either for the skin (e.g. HPV-1 and -2) or mucosal surfaces (e.g. HPV-6 and -11) and usually cause benign tumours (warts) that persist for several months or years. Such benign tumours may be distressing for the individuals concerned but tend not to be life threatening, with a few exceptions.

Some HPVs are also associated with cancers, known as oncogenic HPV types.  
15   The strongest positive association between an HPV and human cancer is that which exists between HPV-16 and HPV-18 and cervical carcinoma. Cervical cancer is the most common malignancy in developing countries, with about 500,000 new cases occurring in the world each year.

Other HPV types which can cause cancer are types 31, 33, 35, 39, 45, 51, 52,  
20   56, 58, 59, 66 and 68 (referred to as "oncogenic-HPV types"). Types 16 and 18 are those which have the highest association with cervical cancer. Types 31 and 45 are the types with the next highest association with a cancer risk (Munoz N, Bosch FX, de Sanjose S *et al.* International Agency for Research on Cancer Multicenter Cervical Cancer Study Group. *N Engl J Med* 2003; 348: 518–27.)

25           HPV virus like particles (VLPs) have been suggested as potential vaccines for treatment of HPV. Animal studies have shown that VLPs produce no cross protection against infection for other HPV types – see, for example Suzich, J. A., et al, *Proc Natl Acad Sci*, 92: 11553-11557, 1995, and Breitburd, *Seminars in Cancer Biology*, vol 9, 1999, pp 431 – 445.

30           WO2004/056389 discloses that an HPV 16, 18 VLP vaccine can provide cross protection against infection by HPV types other than 16 and 18. Statistically significant

protection was observed against certain groups of HPV types. However, the level of cross protection against individual types within groups was not disclosed.

There is still a need for a vaccine that protects against multiple HPV types.

5

### Summary of the invention

The present invention relates to a multivalent HPV vaccine, the vaccine comprising an L1 protein or immunogenic fragment thereof from at least 3 different oncogenic HPV types, those types including HPV 16 and HPV 18, wherein the vaccine does not comprise an L1 protein or immunogenic fragment thereof from an HPV type  
10 selected from the list consisting of HPV 31, HPV 45, HPV 52 or any combination thereof.

The present invention further relates to use of a composition comprising an L1 protein or immunogenic fragment thereof from HPV 16 and HPV 18 in the manufacture of a medicament for prevention of infection and/or disease by one or more  
15 of the group consisting of HPV 31, HPV 45 and HPV 52.

The present invention further relates to use of a composition comprising an L1 protein or immunogenic fragment thereof from HPV 16 and HPV 18 in the manufacture of a medicament for prevention of cytological abnormalities, and/or reduction of the frequency of cytological abnormalities, and/or prevention of CIN  
20 lesions (ASCUS, CIN 1, CIN 2, CIN , cervical cancer) in an individual, the abnormalities or lesions caused by at least one HPV type other than HPV 16 or HPV 18, suitably being caused by HPV type 31, or 45, or 52, or a combination thereof.

The invention further relates to a method of prevention and/or treatment of HPV infection and/or disease, the method comprising delivering to an individual in need  
25 thereof an effective amount of a composition comprising an L1 protein or immunogenic fragment thereof from at least 3 different oncogenic HPV types, those types including HPV 16 and HPV 18, wherein the vaccine does not comprise an L1 protein or immunogenic fragment thereof from an HPV type selected from the list consisting of HPV 31, HPV 45, HPV 52 or any combination thereof.

30 The invention further relates to use of a multivalent composition in the manufacture of a medicament for the prevention and/or treatment of HPV infection and/or disease, the multivalent composition comprising an L1 protein or immunogenic



fragment thereof from at least 3 different oncogenic HPV types, those types including HPV 16 and HPV 18, wherein the vaccine does not comprise an L1 protein or immunogenic fragment thereof from an HPV type selected from the list consisting of HPV 31, HPV 45, HPV 52 or any combination thereof, and wherein the medicament  
5 provides protection against infection and/or disease caused by the omitted HPV type.

The invention also relates to a method for manufacture of a vaccine, the method comprising combining an L1 protein or immunogenic fragment thereof from at least 3 different oncogenic HPV types, those types including types HPV 16 and HPV 18, wherein the vaccine does not comprise an L1 protein or immunogenic fragment thereof  
10 from an HPV type selected from the list consisting of HPV 31, HPV 45, HPV 52 or any combination thereof.

### Detailed description

The general existence of cross protection afforded by HPV 16 and HPV 18  
15 against both incident and persistent infection, as assessed in relation to certain groups of HPV types, has been disclosed in WO2004/056389.

We have surprisingly discovered that the cross protection against certain (non HPV16, HPV 18) HPV types ( as assessed by the efficacy of an HPV 16 and HPV 18 vaccine against those types), is higher than against certain other (non HPV16, HPV 18)  
20 HPV types. Cross protection may be considered as the protection afforded by a vaccine containing one HPV type against infection (incident or persistent) and/or disease caused by a different HPV type. Cross protection may be assessed by considering the vaccine efficacy (V.E.), wherein the V.E. is the % improvement in protection against infection or disease by the vaccine compared to a placebo group for a given type.

25 Infection may be incident or persistent infection. Disease may be abnormal cytology, ASCUS, CIN1, CIN2, CIN3 or cervical cancer related to HPV infection. Infection may be assessed by PCR, for example. Disease may be assessed by, for example, histological examination or analysis of biomarkers such as p16.

Such a finding has potential implications for vaccine design. For example, the  
30 level of cross protection afforded by HPV 16 and HPV 18 L1 containing vaccines against certain other HPV types, such as HPV 31, HPV 45 and HPV 52, allows L1 components from these HPV types to be omitted from a vaccine comprising HPV 16

and HPV18 while still providing a vaccine which provides some protection against incident and/or persistent infection and/or disease related to those omitted types.

After HPV types 16 (found in 53.5% of cervical cancer) and 18 (found in 17.2% of cervical cancer), types 45 (6.7%) and 31 (2.9%) are the next most significant in terms of their frequency in cervical cancers (Muñoz et al. *supra*). HPV 33 (2.6%) is next, followed by HPV 52 (2.3%). Thus, when designing a multivalent HPV vaccine against cervical cancer containing at least HPV 3 types then types 31 and 45 would generally be included by the skilled person after types 16 and 18 from a statistical perspective.

10 The ability to omit antigens from certain HPV types potentially allows inclusion of L1 protein from other HPV types, or indeed antigens from other viruses or pathogens, into a vaccine in a scenario where the total amount of antigen in a vaccine may be limited, for example by physical, chemical, regulatory or other constraints.

In particular other HPV types include oncogenic HPV types such as HPV 31,  
15 33, 35, 39, 45, 51, 52, 56, 58, 59, 66 and 68.

The present invention relates to a multivalent HPV vaccine, the vaccine comprising an L1 protein or immunogenic fragment thereof from at least 3 different oncogenic HPV types, those types including HPV 16 and HPV 18, wherein the vaccine  
20 does not comprise an L1 protein or immunogenic fragment thereof from an HPV type selected from the list consisting of HPV 31, HPV 45, HPV 52 or any combination thereof.

In one aspect of the invention the vaccine does not contain an L1 protein or  
25 immunogenic fragment thereof from HPV 31.

In one aspect of the invention the vaccine is capable of providing protection against incident and /or persistent HPV infection by HPV 31.

In one aspect of the invention the vaccine of the invention does not contain an L1 protein or immunogenic fragment thereof from HPV 45.

30 In one aspect of the invention the vaccine is capable of providing protection against incident and /or persistent HPV infection by HPV 45.



In one aspect of the invention the vaccine does not contain an L1 protein or immunogenic fragment thereof from HPV 52.

In one aspect of the invention the vaccine is capable of providing protection against incident and /or persistent HPV infection by HPV 52.

5 In one aspect of the invention the vaccine of the invention does not contain an L1 protein or immunogenic fragment thereof from HPV 31 and 45.

In one aspect of the invention the vaccine is capable of providing protection against incident and /or persistent HPV infection by both HPV 31 and 45.

10 In one aspect of the invention the vaccine does not contain an L1 protein or immunogenic fragment thereof from HPV 31 and 52.

In one aspect of the invention the vaccine is capable of providing protection against incident and /or persistent HPV infection by both HPV 31 and 52.

In one aspect of the invention the vaccine of the invention does not contain an L1 protein or immunogenic fragment thereof from HPV 45 and 52.

15 In one aspect of the invention the vaccine is capable of providing protection against incident and /or persistent HPV infection by both HPV 52 and 45.

In one aspect of the invention the vaccine is capable of providing protection against incident and /or persistent HPV infection by HPV 31 and HPV 45 and HPV52.

Suitably the vaccine is capable of protection against persistent infection.

20 Suitably the vaccine is capable of protection against incident infection.

Incident and persistent cervical infection are defined in Example 1.

We have also determined that a vaccine comprising HPV 16 L1 and HPV 18 L1 proteins (for example, as described in example 1) provides protection against  
25 cytological abnormalities caused by certain other oncogenic HPV types such as HPV 52, and is significantly protective with respect to such abnormalities caused by a group of HPV high risk types (defined as 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, and 68). Cytological abnormalities are suitably detected by the well known Pap smear technique.

30 Thus the invention further relates to use of a combination of an L1 protein or immunogenic fragment thereof from HPV 16 and HPV 18 in the preparation of a composition for the prevention of cytological abnormalities or reduction of the



frequency of cytological abnormalities in an individual caused by other (non HPV 16, HPV 18) HPV types, suitably oncogenic HPV types, and in the prevention of histologically-confirmed CIN lesions (CIN 1, CIN 2, CIN 3) and cervical cancer associated with infection by HPV types which are not HPV 16 or 18. Said use is in addition to the prevention or reduction of such events caused by the HPV types in the vaccine, HPV 16 and 18.

Suitably the prevention of cytological abnormalities, reduction of the frequency of cytological abnormalities or prevention of histological-confirmed CIN lesions is prevention against those abnormalities or lesions caused by types not included in the combination, suitably selected from the list of HPV 31, HPV 45 and HPV 52, or is prevention against those abnormalities or lesions caused by the group of 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, and 68. Said use is in addition to the prevention or reduction of such events caused by the HPV types in the vaccine, HPV 16 and 18.

Suitably the composition comprising HPV 16 and HPV 18 for use as above is the multivalent HPV vaccine of the invention, the vaccine comprising an L1 protein or immunogenic fragment thereof from at least 3 different oncogenic HPV types, those types including HPV 16 and HPV 18, wherein the vaccine does not comprise an L1 protein or immunogenic fragment thereof from an HPV type selected from the list consisting of HPV 31, HPV 45, HPV 52 or any combination thereof.

The vaccine of the invention comprises L1 or immunogenic fragment from HPV 16, HPV 18 and at least one other oncogenic HPV type. The oncogenic HPV types are those types associated with a risk of cervical cancer and those oncogenic types that might be included in the vaccine of the invention in addition to HPV 16 and HPV 18 include, but are not limited to, HPV 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66 and 68, with the proviso that the vaccine does not comprise all of HPV 31, 45 and 52.

The vaccine of the invention suitably comprises an HPV 33 L1 protein or immunogenic fragment thereof.

The vaccine of the invention suitably comprises an HPV 58 L1 protein or immunogenic fragment thereof.

The vaccine of the invention suitably comprises an HPV 59 L1 protein or immunogenic fragment thereof.

The vaccine of the invention suitably comprises an HPV 16 L1 protein or immunogenic fragment thereof, HPV 18 L1 protein or immunogenic fragment thereof, HPV 33 L1 protein or immunogenic fragment thereof and HPV 58 L1 protein or immunogenic fragment thereof.

5 L1 proteins or protein fragments from additional HPV types can be included in the vaccine of the invention, such as skin types (in particular HPV 5 and 8) and types associated with genital warts, such as HPV 6 and 11. Types 6 and 11 are not considered oncogenic types herein.

10 In one aspect of the invention the vaccine may include an HPV early antigen, for example an antigen selected from the list consisting of HPV E1, E2, E3, E4, E5, E6, E7, or E8. In an alternative aspect the vaccine may lack an HPV early antigen, for example an antigen selected from the list consisting of HPV E1, E2, E3, E4, E5, E6, E7, or E8.

15 In one aspect the vaccine of the invention is trivalent (contains an HPV L1 or fragment thereof from 3 different oncogenic HPV types). In a further aspect the vaccine is tetravalent. In a further aspect the vaccine is pentavalent. In a further aspect the vaccine is heptavalent. In a further aspect the vaccine is septavalent. In a further aspect the vaccine is octavalent. Higher order valencies are also contemplated herein. In further aspects the vaccine is at least tetravalent, pentavalent, heptavalent, 20 septavalent or octavalent with respect to oncogenic HPV types.

25 Preferably the combination of HPV components within the vaccine does not significantly impact the immunogenicity of any one HPV component. In particular it is preferred that there is no biologically relevant interference between HPV antigens in the combination of the invention, such that the combined vaccine of the invention is able to offer effective protection against infection or disease caused by each HPV genotype represented in the vaccine. Suitably the immune response against a given HPV type in the combination is at least 50 % of the immune response of that same HPV type when measured individually, preferably 100% or substantially 100%. For responses to the HPV 16 and HPV 18, the combined vaccine of the invention 30 preferably stimulates an immune response which is at least 50% of that provided by a combined HPV 16 / HPV 18 vaccine. Suitably the immune response generated by the vaccine of the invention is at a level in which the protective effect of each HPV type is



still seen. The immune response may suitably be measured, for example, by antibody responses, in either preclinical or human experiments. Measurement of antibody responses is well known in the art, and disclosed in (for example) WO03/077942.

- 5 We have determined that a vaccine comprising HPV 16 L1 and HPV 18 L1 proteins (e.g. see example 1) provides cross protection against infection or disease caused by certain HPV types. As well as providing novel compositions, this information allows new uses to be developed.

10 In particular, the invention relates to use of a composition comprising HPV 16 and HPV 18 L1 protein, or immunogenic fragment thereof, in the manufacture of a medicament for prevention of infection by HPV 31.

The invention further relates to use of a composition comprising HPV 16 and HPV 18 L1 protein, or immunogenic fragment thereof, in the manufacture of a medicament for prevention of infection by HPV 45.

15 The invention further relates to use of a composition comprising HPV 16 and HPV 18 L1 protein, or immunogenic fragment thereof in the manufacture of a medicament for prevention of infection by HPV 52.

In one aspect the invention relates to use of a vaccine comprising HPV 16 L1 proteins, or immunogenic fragment thereof, in the preparation of a medicament for the prevention of infection and/or disease caused by HPV 31, or HPV 52, or any combination thereof.

In one aspect the invention relates to use of a vaccine comprising HPV 18 L1 proteins, or immunogenic fragment thereof, in the preparation of a medicament for the prevention of infection and/or disease caused by HPV 45.

25 The composition for said use may lack an antigenic component from the HPV type for which cross protection is provided. Alternatively the composition for said use may comprise such an antigenic component, e.g. the L1 protein or fragment thereof from said cross protected type. In the latter case the use of the composition comprising HPV 16 and HPV 18 L1 protein, or immunogenic fragment thereof, provides both cross protection (e.g. against HPV 31, 45 and 52) and homologous protection (e.g. against HPV 16 and HPV 18).

The composition in one aspect is a multivalent composition comprising L1 proteins or immunogenic fragments thereof from HPV 16, HPV 18 and at least one other oncogenic HPV type, wherein an L1 protein or immunogenic fragment thereof from one or more HPV types selected from the group consisting of HPV 31, HPV 45, and HPV 52 is omitted from the vaccine and wherein the vaccine provides protection against infection caused by the omitted HPV type.

Where the vaccine or composition of the invention comprises an immunogenic fragment of L1, then suitable immunogenic fragments of HPV L1 include truncations, deletions, substitution, or insertion mutants of L1. Such immunogenic fragments are suitably capable of raising an immune response (if necessary, when adjuvanted), said immune response being capable of recognising an L1 protein such as a virus like particle, from the HPV type from which the L1 protein was derived.

A suitable immunogenic fragment of HPV 16 is capable of cross protection against at least one of HPV 31 and HPV 52, and in an aspect of the invention, capable of cross protection against both.

A suitable immunogenic fragment of HPV 18 is capable of cross protection against HPV 45.

Cross protection obtainable by immunogenic fragments of HPV 16 and/or HPV 18 can be assessed by trials in humans, for example as outlined in Example 1.

Similarly, different vaccines according to the present invention can be tested using standard techniques, for example as in Example 1, or in standard preclinical models, to confirm that the vaccine is immunogenic.

Suitable immunogenic L1 fragments include truncated L1 proteins. In one aspect the truncation removes a nuclear localisation signal. In another aspect the truncation is a C terminal truncation. In a further aspect the C terminal truncation removes fewer than 50 amino acids, such as fewer than 40 amino acids. Where the L1 is from HPV 16 then in another aspect the C terminal truncation removes 34 amino acids from HPV 16 L1. Where the L1 is from HPV 18 then in a further aspect the C terminal truncation removes 35 amino acids from HPV 18 L1. Truncated L1 Proteins are described in US 6,060,324, US 6,361,778, and US 6,599,508 incorporated herein by reference.



In one aspect the HPV 16 amino acid sequence is the following sequence: (SEQ ID NO: 1)

	MSLWLPSEATVYLPPVPVSKVVSTDEYVARTNIYYHAGTSRLLAVGHPYFPIKKPNNNKI	60
	LVPKVSGLQYRVFRIHLPDPNKFGLPDTSFYNPDTQRLVWACVGVEVGRGQPLGVGISGH	120
5	PLLNLDDTENASAYAANAGVDNRECISMDYKQTQLCLIGCKPPIGEHWGKGSPCTNVAV	180
	NPGDCPPLELINTVIQDGMVDTGFGAMDFTTLQANKSEVPLDICTSICKYPDYIKMVSE	240
	PYGDSLFFYLRRQMFVRHLFNRAVGENVPDDLVIKSGSGSTANLASSNYFPTPSGSMV	300
	TSDAQIFNKPYWLQRAQGHNNGICWGNQLFVTVDTRSTNMSLCAAISTSETTYKNTNF	360
	KEYLRHGEEYDLQFIFQLCKITLTADVMTYIHSMNSTILEDWNFGLQPPPGGTLEDTYRF	420
10	VTSQAIACQKHTPPAPKEDPLKKYTFWEVNLKEKFSADLDQFPLGRKFLQ	471

In another aspect the invention relates to virus like particles consisting only of HPV 16 L1 having the amino sequence above, and to compositions containing such VLPs.

15

The HPV 16 sequence may also be that disclosed in WO9405792 or US6649167, for example, suitably truncated. Suitable truncates are truncated at a position equivalent to that shown above, as assessed by sequence comparison.

20 In one aspect the HPV 18 amino acid sequence is the following sequence: (SEQ ID NO: 2)

	MALWRPSDNTVYLPSPSVARVVNTDDYVTRTSIFYHAGSSRLLTVGNPYFRVPAGGGNKQ	60
	DIPKVSAYQYRVFRVQLPDPNKFGLPDNSIYNPETQRLVWACVGVEIGRGQPLGVGLSGH	120
	PFYNKLDDTESSHAATSNVSEDVRDNVSVDYKQTQLCILGCAPAIGEHWAKGTACKSRPL	180
25	SQGDCPPLELKNTVLEDGDMVDTGYGAMDFSTLQDTKCEVPLDICQSICKYPDYLQMSAD	240
	PYGDSMFFCLRREQLFARHFWNRAGTMGDTVPPSLYIKGTGMRASPGSCVYSPSPSGSIV	300
	TSDSQLFNKPYWLHKAQGHNNGVCWHNQLFVTVDTRSTNLTICASTQSPVPGQYDATK	360
	FKQYSRHHVEEYDLQFIFQLCTITLTADVMSYIHSMNSSILEDWNFGVPPPPTTSLVDTYR	420
	FVQSVAITCQKDAAPAENKDPYDKLKFWNVDLKEKFSLDLDQYPLGRKFLVQ	472

30

In another aspect the invention relates to virus like particles consisting only of HPV 18 L1 having the amino sequence above, and to compositions containing such VLPs.

35 An alternative HPV 18 sequence is disclosed in WO9629413, which may be suitably truncated. Suitable truncates are truncated at a position equivalent to that shown above, as assessed by sequence comparison.

Other HPV 16 and HPV 18 sequences are well known in the art and may be suitable for use in the present invention.

Suitable truncations of HPV 31, HPV 45 and HPV 52 may also be made, suitably removing equivalent C terminal portions of the L1 protein to those described  
5 above as assessed by sequence alignment.

Truncated L1 proteins are disclosed in, for example, WO9611272 and US6066324, herein incorporated by reference.

The L1 protein or fragment of the invention may optionally be in the form of a fusion protein, such as the fusion of the L1 protein with L2 or an early protein.

10 The HPV L1 protein is suitably in the form of a capsomer or virus like particle (VLP). In one aspect HPV VLPs may be used in the present invention. HPV VLPs and methods for the production of VLPs are well known in the art. VLPs typically are constructed from the L1 and optionally L2 structural proteins of the virus, see for example WO9420137, US5985610, W09611272, US6599508B1, US6361778B1, EP  
15 595935. Any suitable HPV VLP may be used in the present invention which provides cross protection, such as an L1 or L1 + L2 VLP.

Suitably the VLP is an L1-only VLP.

In one aspect of the invention the vaccine comprises HPV 16 and HPV 18 L1 only VLPs, suitably in combination with an L1 VLP selected from HPV 31, 33, 35, 39,  
20 45, 51, 52, 56, 58, 59, 66 and 68, with the proviso that the vaccine does not comprise VLPs from all of HPV 31, 45 and 52.

VLP formation can be assessed by standard techniques such as, for example, electron microscopy and dynamic laser light scattering.

The VLP may comprise full length L1 protein. In one aspect the L1 protein  
25 used to form the VLP is a truncated L1 protein, as described above.

VLPs may be made in any suitable cell substrate such as yeast cells or insect cells e.g. baculovirus cells, and techniques for preparation of VLPs are well known in the art, such as WO9913056, US 6416945B1, US 6261765B1 and US6245568, and references therein, the entire contents of which are hereby incorporated by reference.

30 VLPS are suitably made by disassembly and reassembly techniques, which can provide for more stable and/or homogeneous papillomavirus VLPs. For example, McCarthy et al, 1998 "Quantitative Disassembly and Reassembly of Human



Papillomavirus Type 11 Virus like Particles in Vitro" J. Virology 72(1):33-41, describes the disassembly and reassembly of recombinant L1 HPV 11 VLPs purified from insect cells in order to obtain a homogeneous preparation of VLPs. WO9913056 and US6245568 also describe disassembly/reassembly processes for making HPV

5 VLPs.


In one aspect HPV VLPS are made as described WO9913056 or US6245568.

The HPV L1 the invention may be combined with an adjuvant or immunostimulant such as, but not limited to, detoxified lipid A from any source and non-

10 toxic derivatives of lipid A, saponins and other reagents capable of stimulating a TH1 type response.

It has long been known that enterobacterial lipopolysaccharide (LPS) is a potent stimulator of the immune system, although its use in adjuvants has been curtailed by its toxic effects. A non-toxic derivative of LPS, monophosphoryl lipid A (MPL), produced

15 by removal of the core carbohydrate group and the phosphate from the reducing-end glucosamine, has been described by Ribí et al (1986, Immunology and Immunopharmacology of bacterial endotoxins, Plenum Publ. Corp., NY, p407-419) and has the following structure:



A further detoxified version of MPL results from the removal of the acyl chain from the 3-position of the disaccharide backbone, and is called 3-O-Deacylated monophosphoryl lipid A (3D-MPL). It can be purified and prepared by the methods taught in GB 2122204B, which reference also discloses the preparation of diphosphoryl lipid A, and 3-O-deacylated variants thereof.

A suitable form of 3D-MPL is in the form of an emulsion having a small particle size less than 0.2µm in diameter, and its method of manufacture is disclosed in WO 94/21292. Aqueous formulations comprising monophosphoryl lipid A and a surfactant have been described in WO9843670A2.

The bacterial lipopolysaccharide derived adjuvants to be formulated in the compositions of the present invention may be purified and processed from bacterial sources, or alternatively they may be synthetic. For example, purified monophosphoryl lipid A is described in Ribi et al 1986 (supra), and 3-O-Deacylated monophosphoryl or diphosphoryl lipid A derived from *Salmonella sp.* is described in GB 2220211 and US 4912094. Other purified and synthetic lipopolysaccharides have been described (Hilgers *et al.*, 1986, *Int.Arch.Allergy.Immunol.*, 79(4):392-6; Hilgers *et al.*, 1987, *Immunology*, 60(1):141-6; and EP 0 549 074 B1). In one aspect the bacterial lipopolysaccharide adjuvant is 3D-MPL.

Accordingly, the LPS derivatives that may be used in the present invention are those immunostimulants that are similar in structure to that of LPS or MPL or 3D-MPL. In another aspect of the present invention the LPS derivatives may be an acylated monosaccharide, which is a sub-portion to the above structure of MPL.

Saponins are taught in: Lacaille-Dubois, M and Wagner H. (1996. A review of the biological and pharmacological activities of saponins. *Phytomedicine* vol 2 pp 363-386). Saponins are steroid or triterpene glycosides widely distributed in the plant and marine animal kingdoms. Saponins are noted for forming colloidal solutions in water which foam on shaking, and for precipitating cholesterol. When saponins are near cell membranes they create pore-like structures in the membrane which cause the membrane to burst. Haemolysis of erythrocytes is an example of this phenomenon, which is a property of certain, but not all, saponins.



Saponins are known as adjuvants in vaccines for systemic administration. The adjuvant and haemolytic activity of individual saponins has been extensively studied in the art (Lacaille-Dubois and Wagner, *supra*). For example, Quil A (derived from the bark of the South American tree *Quillaja Saponaria* Molina), and fractions thereof, are  
5 described in US 5,057,540 and "Saponins as vaccine adjuvants", Kensil, C. R., *Crit Rev Ther Drug Carrier Syst*, 1996, 12 (1-2):1-55; and EP 0 362 279 B1. Particulate structures, termed Immune Stimulating Complexes (ISCOMS), comprising fractions of Quil A are haemolytic and have been used in the manufacture of vaccines (Morein, B., EP 0 109 942 B1; WO 96/11711; WO 96/33739). The haemolytic saponins QS21 and  
10 QS17 (HPLC purified fractions of Quil A) have been described as potent systemic adjuvants, and the method of their production is disclosed in US Patent No.5,057,540 and EP 0 362 279 B1. Other saponins which have been used in systemic vaccination studies include those derived from other plant species such as *Gypsophila* and *Saponaria* (Bomford *et al.*, *Vaccine*, 10(9):572-577, 1992).

15 An enhanced system involves the combination of a non-toxic lipid A derivative and a saponin derivative particularly the combination of QS21 and 3D-MPL as disclosed in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol as disclosed in WO 96/33739.

A particularly potent adjuvant formulation involving QS21 and 3D-MPL in an  
20 oil in water emulsion is described in WO 95/17210 and use of this adjuvant forms an aspect of the invention.

Accordingly in one embodiment of the present invention there is provided a vaccine adjuvanted with detoxified lipid A or a non-toxic derivative of lipid A, more suitably adjuvanted with a monophosphoryl lipid A or derivative thereof.

25 In one aspect the vaccine additionally comprises a saponin, for example QS21.

In one aspect the vaccine formulation comprises an oil in water emulsion. The present invention also provides a method for producing a vaccine formulation comprising mixing an L1 peptide of the present invention together with a pharmaceutically acceptable excipient, such as 3D-MPL.

30 Additional components that may be included present in an vaccine formulation according to the invention include non-ionic detergents such as the octoxynols and polyoxyethylene esters as described herein, particularly t-octylphenoxy

polyethoxyethanol (Triton X-100) and polyoxyethylene sorbitan monooleate (Tween 80); and bile salts or cholic acid derivatives as described herein, in particular sodium deoxycholate or taurodeoxycholate. Thus, in one aspect of the invention a formulation comprises 3D-MPL, Triton X-100, Tween 80 and sodium deoxycholate, which may be  
5 combined with an L2 antigen preparation to provide a suitable vaccine.

In one embodiment of the present invention, the vaccine comprises a vesicular adjuvant formulation comprising cholesterol, a saponin and an LPS derivative. In this regard the adjuvant formulation suitably comprises a unilamellar vesicle comprising cholesterol, having a lipid bilayer suitably comprising dioleoyl phosphatidyl choline,  
10 wherein the saponin and the LPS derivative are associated with, or embedded within, the lipid bilayer. In one aspect these adjuvant formulations comprise QS21 as the saponin, and 3D-MPL as the LPS derivative, wherein the ratio of QS21:cholesterol is from 1:1 to 1:100 weight/weight, and in one aspect, a ratio of 1:5 weight/weight. Such adjuvant formulations are described in EP 0 822 831 B, the disclosure of which is  
15 incorporated herein by reference.

Suitably the vaccines of the invention are used in combination with aluminium, and are suitably adsorbed or partially adsorbed onto aluminium adjuvants. Suitably the adjuvant is an aluminium salt, which may be in combination with 3D MPL, such as aluminium phosphate and 3D MPL. Aluminium hydroxide, optionally in combination  
20 with 3D MPL is also suitable.

In another aspect of the present invention the vaccine comprises the combination of HPV VLPs with an aluminium salt or with an aluminium salt + 3D MPL. Aluminium hydroxide is suitable as the aluminium salt.

The vaccine may also comprise aluminium or an aluminium compound as a  
25 stabiliser.

In another aspect the adjuvant may be a combination of an oil-in-water emulsion adjuvant and 3D MPL. In one aspect the oil-in-water emulsion comprises a metabolisable oil, a sterol and an emulsifying agent.

The vaccines of the invention may be provided by any of a variety of routes  
30 such as oral delivery (e.g. see WO9961052 A2), topical, subcutaneous, mucosal (typically intravaginal), intravenous, intramuscular, intranasal, sublingual, intradermal and via suppository.



Optionally the vaccine may also be formulated or co-administered with other HPV antigens or non-HPV antigens. Suitably these non-HPV antigens can provide protection against other diseases, such as sexually transmitted diseases such as herpes simplex virus, EBV, chlamydia and HIV. We particularly prefer that the vaccine  
5 comprises gD or a truncate thereof from HSV. In this way the vaccine provides protection against both HPV and HSV.

The dosage of the vaccine components will vary with the condition, sex, age and weight of the individual, the administration route and HPV of the vaccine. The quantity may also be varied with the number of VLP types. Suitably the delivery is of  
10 an amount of vaccine suitable to generate an immunologically protective response. Suitably each vaccine dose comprises 1-100 µg of each VLP, in one aspect 5-80µg, in another aspect 5- 30 µg each VLP, in a further aspect 5-20 µg of each VLP, in a yet further aspect 5µg, 6µg, 10µg, 15 µg or 20µg.

In one aspect the vaccine can comprise HPV L1 protein components, preferably  
15 as virus like particles, in different amounts. In one aspect, HPV 16 and HPV 18 VLPs may be provided at a higher dose than other oncogenic types, such as HPV 33 or 58. In one aspect HPV 16 and HPV 18 L1 only VLPs are provided at 20µg per dose for human use. Other HPV VLPs may be used at a lower dose, such as 15 or 10µg per dose for human use.

20 For all vaccines of the invention, in one aspect the vaccine is used for the vaccination of adolescent girls aged 10-15, such as 10-13 years. However, older girls above 15 years old and adult women may also be vaccinated. The vaccine may also be administered to women following an abnormal pap smear or after surgery following removal of a lesion caused by HPV, or who are seronegative and DNA negative for  
25 HPV cancer types.

The vaccine of the invention may be used in men.

In one aspect the vaccine is delivered in a 2 or 3 dose regime, for example in a 0, 1 month regime or 0,1 and 6 month regime respectively. Suitably the vaccination regime incorporates a booster injection after 5 to 10 years, such as 10 years.

30 In one aspect the vaccine is a liquid vaccine formulation, although the vaccine may be lyophilised and reconstituted prior to administration.

The teaching of all references in the present application, including patent applications and granted patents, are herein fully incorporated by reference.

The vaccines of the invention comprise certain HPV components as laid out above. In a further aspect of the invention the vaccine consists essentially of, or  
5 consists of, said components.

The term 'vaccine', as used in the present invention, refers to a composition that comprises an immunogenic component capable of provoking an immune response in an individual, such as a human, optionally when suitably formulated or adjuvant. A vaccine suitably elicits a protective immune response against incident infection, or  
10 persistent infection, or cytological abnormality such as ASCUS, CINI, CIN2, CIN3, or cervical cancer caused by one or more HPV types.

The present invention is now described with respect to the following examples which serve to illustrate the invention.

#### 15 **Example 1**

Precise details of the experiment carried out are provided in Harper *et al*, the Lancet. 2004 Nov 13;364(9447):1757-65, incorporated herein by reference.

In summary, healthy women between the ages of 15 and 25 years were immunised with a mixture of HPV 16 and HPV 18 L1 VLPs. The women at enrolment  
20 were: 1) seronegative for HPV-16 and HPV-18; 2) negative for high risk HPV infection of the cervix (detected by HPV PCR); 3) had 6 or fewer lifetime sexual partners and 4) had normal PAP smears.

The mixture comprised, per 0.5 ml dose, 20 µg of HPV-16 L1 VLP, 20 µg of HPV-18 L1 VLP and was adjuvanted with 500 µg of aluminum hydroxide and 50 µg of  
25 3D MPL. The placebo group was injected with 500 µg of aluminum hydroxide alone.

The vaccine efficacy (V.E.) against certain cancer HPV types was assessed, wherein the V.E. is the % improvement in protection against infection or disease by the vaccine compared to a placebo group.

Cross protection was assessed by detecting the presence of nucleic acid specific  
30 for various oncogenic types in the vaccinees and control group. Detection was carried out using techniques as described in WO03014402, and references therein, particularly for non-specific amplification of HPV DNA and subsequent detection of DNA types



using a LiPA system as described in WO 99/14377, and in Kleter et al, [Journal of Clinical Microbiology (1999), 37 (8): 2508-2517], the whole contents of which are herein specifically incorporated by reference.

Any suitable method can, however, be used for the detection of HPV DNA in a sample, such as type specific PCR using primers specific for each HPV type of interest. Suitable primers are known to the skilled person, or can be easily constructed given that the sequences of the oncogenic HPV types are known.

In detail, the methods section of the Lancet paper is reproduced here below, for completeness (continues until section entitled **“Initial analysis and results”**)

The primary objective of this study was to assess vaccine efficacy in the prevention of infection with HPV-16, HPV-18, or both (HPV-16/18), between months 6 and 18 in participants who were initially shown to be seronegative for HPV-16/18 by ELISA and negative for HPV-16/18 DNA by PCR. Secondary objectives included: evaluation of vaccine efficacy in the prevention of persistent infection with HPV-16/18, and the evaluation of vaccine efficacy in the prevention of cytologically confirmed low-grade squamous intraepithelial lesions (LSIL), high-grade squamous intraepithelial lesions (HSIL), and histologically confirmed LSIL (CIN 1), HSIL (CIN 2 or 3) squamous cell cancer, or adenocarcinoma associated with HPV-16/18 infection between months 6 and 18, and months 6 and 27. The prevention of atypical squamous cells of undetermined significance (ASCUS) cytology associated with HPV-16/18 infection was added post-hoc to the outcome analyses.

We also did an exploratory analysis of the histopathological endpoints CIN 1 and 2 associated with HPV-16/18 DNA detected by PCR in lesional tissue. Other objectives included the assessment of vaccine immunogenicity, safety, and tolerability. Investigators in North America (Canada and the USA) and Brazil recruited women for this efficacy study through advertisements or previous participation in an HPV cross-sectional epidemiology study that took place between July and December, 2000. For each of the 32 study sites, an institutional review board approved the protocol, consent forms, and amendments. Women signed separate written consents for study

participation and colposcopy. For those under 18 years, parental consent and assent from the participant were obligatory.

There were two study phases: an initial phase for vaccination and follow-up that concluded at month 18; and a blinded follow-up extension phase that concluded at  
5 month 27.

Women eligible for the initial phase (months 0–18) included healthy women aged 15–25 years, who had had no more than six sexual partners, no history of an abnormal Pap test or ablative or excisional treatment of the cervix, and no ongoing treatment for external condylomata; and who were cytologically negative, seronegative  
10 for HPV-16 and HPV-18 antibodies by ELISA, and HPV-DNA-negative by PCR for 14 high-risk HPV types (16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, and 68) no more than 90 days before study entry.

Women who completed the initial phase of the study earliest, and who did not have ablative or excisional therapy of the cervix, or hysterectomy after enrolment, were  
15 eligible to participate in the extension phase of the study (months 18–27).

### Procedures

Each dose of the bivalent HPV-16/18 virus-like particle vaccine (GlaxoSmithKline Biologicals, Rixensart, Belgium) contained 20 µg of HPV-16 L1  
20 virus-like particle and 20 µg of HPV-18 L1 virus-like particle. Each type of virus-like particle was produced on *Spodoptera frugiperda* Sf-9 and *Trichoplusia ni* Hi-5 cell substrate with AS04 adjuvant containing 500 µg aluminum hydroxide and 50 µg 3-deacylated monophosphoryl lipid A (MPL, Corixa, Montana, USA) provided in a monodose vial. The placebo contained 500 µg of aluminum hydroxide per dose, and  
25 was identical in appearance to the HPV-16/18 vaccine. Every study participant received a 0.5 mL dose of vaccine or placebo at 0 months, 1 month, and 6 months.

Health-care providers obtained cervical specimens with a cervical brush and spatula (washed in PreservCyt, Cytyc Corporation, Boxborough, MA, USA) for cytology and HPV DNA testing at screening and months 6, 12, and 18. At months 0 and 6, and  
30 subsequently every 3 months, women self-obtained cervicovaginal samples with two sequential swabs (placed in PreservCyt) for HPV DNA testing.[ DM Harper, WW Noll, DR Belloni and BF. Cole, Randomized clinical trial of PCR-determined human



papillomavirus detection methods: self-sampling versus clinician-directed—biologic concordance and women's preferences. *Am J Obstet Gynecol* **186** (2002), pp. 365–373]

A central laboratory (Quest Diagnostics, Teterboro, NJ, USA) reported cytology results (ThinPrep, Cytyc Corporation) by use of the 1991 Bethesda classification system.

- 5 Protocol guidelines recommended colposcopy after two reports of ASCUS, or one report of atypical glandular cells of undetermined significance, LSIL or HSIL, squamous cell carcinoma, adenocarcinoma in situ, or adenocarcinoma. These guidelines also recommended biopsy for any suspected lesions.

- The central histology laboratory made an initial diagnosis from the formalin-fixed  
10 tissue specimens for clinical management. A panel of three pathologists made a subsequent consensus diagnosis for HPV-16 and HPV-18 associated lesions with the CIN system. This consensus diagnosis also included review of the sections taken at the time of microdissection for PCR detection of lesional HPV DNA.

- HPV DNA isolated from the cytology specimen (MagNaPure Total Nucleic Acid  
15 system, Roche Diagnostics, Almere, Netherlands) and from the cervical biopsy specimen (proteinase K extraction) was amplified from an aliquot of purified total DNA with the SPF10 broad-spectrum primers that amplify a 65 bp region of the L1 gene.[ B Kleter, LJ van Doorn, J ter Schegget *et al.*, Novel short-fragment PCR assay for highly sensitive broad-spectrum detection of anogenital human papillomaviruses.  
20 *Am J Pathol* **153** (1998), pp. 1731–1739: LJ van Doorn, W Quint, B Kleter *et al.*, Genotyping of human papillomavirus in liquid cytology cervical specimens by the PGMY line blot assay and the SPF(10) line probe assay. *J Clin Microbiol* **40** (2002), pp. 979–983 and WG Quint, G Scholte, LJ van Doorn, B Kleter, PH Smits and J. Lindeman, Comparative analysis of human papillomavirus infections in cervical  
25 scrapes and biopsy specimens by general SPF(10) PCR and HPV genotyping. *J Pathol* **194** (2001), pp. 51–58] The amplification products were detected by a DNA enzyme immunoassay. A line probe assay (LiPA Kit HPV INNO LiPA HPV genotyping assay, SPF-10 system version 1, Innogenetics, Gent, Belgium, manufactured by Labo Bio-  
30 medical Products, Rijswijk, Netherlands) detected 25 HPV genotypes (6, 11, 16, 18, 31, 33, 34, 35, 39, 40, 42, 43, 44, 45, 51, 52, 53, 56, 58, 59, 66, 68, 70, and 74). [B Kleter, LJ van Doorn, L Schrauwen *et al.*, Development and clinical evaluation of a highly sensitive PCR-reverse hybridization line probe assay for detection and identification of

anogenital human papillomavirus. *J Clin Microbiol* **37** (1999), pp. 2508–2517] Any specimen that was positive by DNA enzyme immunoassay was tested by type-specific HPV-16 and HPV-18 PCR. HPV-16 type-specific PCR primers amplified a 92 bp segment of the E6/E7 gene and HPV-18 type-specific PCR primers amplified a 126 bp  
5 segment of the L1 gene. [MF Baay, WG Quint, J Koudstaal *et al.*, Comprehensive study of several general and type-specific primer pairs for detection of human papillomavirus DNA by PCR in paraffin-embedded cervical carcinomas. *J Clin Microbiol* **34** (1996), pp. 745–747]

We defined incident cervical infection with HPV-16/18 as at least one positive PCR  
10 result for HPV-16 or HPV-18 during the trial, and persistent infection with HPV-16/18 as at least two positive HPV-DNA PCR assays for the same viral genotype separated by at least 6 months. [H Richardson, G Kelsall, P Tellier *et al.*, The natural history of type-specific human papillomavirus infections in female university students. *Cancer Epidemiol Biomarkers Prev* **12** (2003), pp. 485–490 and AB Moscicki, JH Ellenberg, S  
15 Farhat and J. Xu, Persistence of human papillomavirus infection in HIV-infected and -uninfected adolescent girls: risk factors and differences, by phylogenetic type. *J Infect Dis* **190** (2004), pp. 37–45] HPV-DNA test results were concealed from investigators during the study and cytological and histological diagnoses were only revealed for clinical management purposes. Analyses included HPV-16/18 DNA results for cervical  
20 specimens and combined cervical and self-obtained cervicovaginal specimens.

We collected serum from study participants at months 0, 1, 6, 7, 12, and 18 for assessment of immunogenicity. Serological testing for antibodies to HPV-16 and HPV-18 virus-like particles was by ELISA. Recombinant HPV-16 or HPV-18 virus-like particles were used as coating antigens for antibody detection (see webappendix  
25 <http://image.thelancet.com/extras/04art10103webappendix.pdf>). Seropositivity was defined as a titre greater than or equal to the assay cut-off titre established at 8 ELISA units/mL for HPV-16 and 7 ELISA units/mL for HPV-18. Typical natural titres were determined by use of blood samples obtained from women in the preceding epidemiology study who were found to be seropositive for HPV-16 or HPV-18 by  
30 ELISA.

Women recorded symptoms experienced during the first 7 days after vaccination on diary cards with a three-grade scale of symptom intensity. Additionally,



they reported to study personnel by interview all adverse events within the first 30 days after vaccination. Information on serious adverse events and pregnancies was collected throughout the study.

## 5 Statistical methods

Assuming a 6% cumulative incidence rate of both HPV-16 and HPV-18 type infections over 12 months, we estimated that 500 women per treatment group would provide 80% power to assess a lower limit of the 95% CI of the vaccine efficacy above zero. We assumed an 80% retention rate over 18 months. Interim analyses for efficacy,  
10 safety, and immunogenicity were done for future study planning purposes only; the O'Brien and Fleming method was used to adjust the  $\alpha$  value for the final analysis after interim analyses occurred (overall  $\alpha=0.05$ ; two-sided test). [PC O'Brien and TR. Fleming, A multiple testing procedure for clinical trials. *Biometrics* 35 (1979), pp. 549–556]

15 Stratified, block randomisation according to validated algorithms was centralised with an internet randomisation system. Stratification was according to age (15–17, 18–21, and 22–25 years) and region (North America and Brazil). Each vaccine dose was attributed a randomly chosen number based on specific participant information entered into the computerised randomisation system by study personnel. Treatment allocation  
20 remains concealed from investigators and the women participating in a long-term follow-up study.

The intention-to-treat and according-to-protocol cohorts are shown in the figure, in which the reasons for exclusion from analyses are listed in rank order; women who met more than one exclusion criterion were only counted once according to the highest  
25 ranking criterion. We refer to the sets of participants entered in the intention-to-treat and according-to-protocol analyses as cohorts, although the information used to restrict subject inclusion in the according-to-protocol was only known after follow-up. We did both according-to-protocol and intention-to-treat analyses for efficacy. Calculation of vaccine efficacy in the according-to-protocol 18-month analysis was  
30 based on the proportion of participants with HPV-16/18 infection in the vaccinated versus placebo groups. Vaccine efficacy was defined as 1 minus the ratio between these two proportions; 95% CIs measured the precision of the efficacy estimates. p values


were calculated with the two-sided Fisher's exact test. Corresponding rates were expressed as the numbers of cases with the outcome divided by the numbers of participants at risk. The according-to-protocol 18-month cohort included enrolled women who received three scheduled doses of vaccine and complied with the protocol  
5 as described in the figure.

Calculation of vaccine efficacy in the intention-to-treat and according-to-protocol 27-month analyses was based on the Cox proportional hazard model using the time-to-occurrence of cases with HPV-16/18 infection in the vaccinated versus placebo groups. This allowed controlling for the accrued person-time data in each group.  
10 Vaccine efficacy was calculated using 1 minus the hazard ratio and p values calculated using the log rank test. Corresponding rates were expressed as the number of cases divided by the total person-time. All enrolled women who received at least one dose of vaccine or placebo, were negative for high-risk HPV-DNA at month 0, and had any data available for outcome measurement were included in the intention-to-treat cohort.

15 The according-to-protocol 27-month cohort included outcome results from the according-to-protocol 18-month cohort and results that occurred during the extension phase (from 18 months to 27 months).

Calculation of p values for the safety analysis was performed using Fisher's exact test comparisons. The cohort for safety analysis included all enrolled women who  
20 received at least one dose of vaccine or placebo and complied with specified, minimal protocol requirements (see protocol below:)





Immunogenicity was assessed in a subset of the according-to-protocol safety cohort, which included women with serology results at months 0, 7, and 18, who received all  
5 three doses of study vaccine or placebo according to schedule, complied with the blood sampling schedule, and did not become positive for HPV-16/18-DNA during the trial. Seropositivity rates between the vaccine and placebo groups were compared with Fisher's exact test ( $p < 0.001$  judged significant). Geometric mean titres were compared with ANOVA and Kruskal-Wallis test.

10 Block randomisation and statistical analyses were done with SAS version 8.2 (SAS Institute, Cary, North Carolina).

### **Initial analysis and results**

Results of the initial analysis on cross protection are presented in patent  
15 application WO2004/056389, the whole contents of which herein incorporated by reference.

An initial analysis was carried out on an "ITT" (Intention To Treat cohort, representing all individuals who received at least one dose of vaccine). This data is shown in Table A.

20 The results presented in Tables B and C relate to the "ATP" (According To Protocol) group for those patients who complied with all the criteria of the trial. Table B is a midpoint analysis with data taken from all patients at the timepoint at which at least 50% of the cohort were 18 months after their first vaccination. Table C gives the final results, all data being from subjects at 18 months post first vaccination (month 0).  
25 In the ATP group all patients received 3 doses of vaccine at 0, 1 and 6 months and were seronegative at 6 months.

As demonstrated by the data presented in table A, immunization with a mixture of HPV16 and HPV18 VLPs provided apparent cross-protection against other HPV types. At this point the sample sizes are too small to provide for a rigorous statistical  
30 analysis, however the data demonstrate a positive trend and suggest that immunization with HPV16 and HPV18 VLPs will be efficacious against infection with other HPV types.



This was confirmed as the study progressed.

Table B demonstrates that HPV 16 and HPV 18 provide statistically significant cross protection against the group of high risk cancer types

5 31,33,35,39,45,51,52,56,58,59, 66 and 68.

Table C demonstrates that, except for the HPV-18 related types (which show a very strong trend), there is statistically significant cross-protection against the groups of: HPV 31, 35, 58; HPV 31, 33, 35, 52, 58; and the 12 high risk (non HPV-16/18) types evaluated.

10 Further analysis was carried out on the specific cross protection against specific types.

Vaccine efficacy was assessed against infections and diseases related to the 12 high risk cancer types 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66 and 68, HPV-16 phylogenetic-related types (the groups of; 31, 35, and 58; 31, 33, 35, 52 and 58) and  
15 HPV-18 phylogenetic related types (45 and 59).

An analysis was carried out on an“ATP” (According To Protocol) group for those patients who complied with all the criteria of the trial. In the ATP group all patients received 3 doses of vaccine at 0, 1 and 6 months and were seronegative at 6 months.

20 As demonstrated by the data presented in Table D, immunization with a mixture of HPV16 and HPV18 VLPs provided statistically significant cross protection against incident infection by HPV types 31, 52 and 45 compared to the control.

Statistically significant cross protection against incident infection was also observed against the group of all HPV 16 related types (HPV-31, 33, 35, 52 and 58)  
25 and the group of all high risk types, excluding 16 and 18 (HPV 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, and 68).

Statistically significant cross protection against persistent infection was also observed against types 31 and 52 and was also observed against the group of all HPV 16 related types (see Table E).

30 Statistically significant cross protection was observed against cytological abnormalities associated with HPV 52 and was also observed against cytological abnormalities associated with the group of all HPV 16 related types (HPV-31, 33, 35,

52, and 58) and the group of all high risk types, excluding 16 and 18 (31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, and 68) (Table F).

Table A

HPV types analysed	HPV 31, 35, 58	HPV 31, 33, 35, 52, 58	HPV 45, 59	HPV 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, 68.
Number of women infected (vaccine group)	5	17	3	27
% women infected (vaccine group) = A	1.1	3.8	0.7	6.3
Number of women infected (placebo group)	11	24	6	40
% women infected (placebo group) = B	2.4	5.4	1.3	9.4
% vaccine efficacy $1 - (A/B) \times 100$ , adjusted for relative size of vaccine and placebo group	55.1	30.3	50.6	34.6
95% confidence limits -lower limit	-29.1	-29.7	-97.7	-6.5
95% confidence limits -upper limit	84.4	62.6	87.6	59.9
P	0.127	0.252	0.309	0.086

5

Samples were taken at 9, 12, 15 and 18 months from patients and tested for HPV infection by the types specified above.



**Table B - vaccine efficacy after three doses in preventing incident heterologous infections.**

Vaccine efficacy against infection with HPV-16 phylogenetically related types, HPV-18 phylogenetically related types, HPV-16 and/or HPV-18 phylogenetically related types and all high-risk types exclusive of HPV-16 and HPV-18 – ATP cohort (month 6-18)

Infection Type	Attack rate						Vaccine efficacy		
	Vaccine			Placebo					
	N	n	AR	N	n	AR	%	95% CI	p-value
HPV-16 related	433	12	2.8	438	24	5.5	49.4	0.2 74.4	0.060
HPV-16 related*	423	29	6.9	423	46	10.9	37.0	1.6 59.6	0.052
HPV-18 related	442	9	2.0	449	16	3.6	42.9	-27.9 74.5	0.223
HPV-16/18 related	433	21	4.9	438	41	9.4	48.2	13.8 68.9	0.012
HPV-16/18 related*	423	34	8.0	423	56	13.2	39.3	9.0 59.5	0.019
High-risk**	385	53	13.8	386	88	22.8	39.6	17.7 55.7	0.001

N = number of subjects in specific cohort

n = number of subjects with incident HPV infection

AR = Attack rate =  $n / N$

10 95% CI = 95% confidence interval

lower limit =  $1 - \exp ( \log (arv / arp) + 1.96 * \sqrt{1/nv - 1/Nv + 1/np - 1/Np} )$

upper limit =  $1 - \exp ( \log (arv / arp) - 1.96 * \sqrt{1/nv - 1/Nv + 1/np - 1/Np} )$

when number of cases in vaccine = 0 :

15 lower limit\* =  $1 - \exp ( \log (arv^* / arp^*) + 1.96 * \sqrt{1/(nv+0.5) - 1/(Nv+0.5) + 1/(np+0.5) - 1/(Np+0.5)} )$

$$\text{upper limit}^* = 1 - \exp \left( \log \left( \text{arv}^* / \text{arp}^* \right) - 1.96 * \sqrt{1/(nv+0.5) - 1/(Nv+0.5) + 1/(np+0.5) - 1/(Np+0.5)} \right)$$

with: arv = attack rate in vaccine recipients

arp = attack rate in placebo recipients

5 nv = number of cases in vaccine recipients

Nv = number of cases and non-cases in vaccine recipients

np = number of cases in placebo recipients

Np = number of cases and non-cases in placebo recipients

10 HPV-16 related: HPV-16 phylogenetically related types 35, 31, 58 without considering other HPV types

HPV-16 related\*: HPV-16 phylogenetically related types 35, 31, 58, 33, 52 without considering other HPV types

HPV-18 related: HPV-18 phylogenetically related types 45, 59 without considering other HPV types

15 HPV-16 and/or HPV-18 related: HPV-16 and/or HPV-18 phylogenetically related types 35, 31, 58, 45, 59 without considering other HPV types

HPV-16 and/or HPV-18 related\*: HPV-16 and/or HPV-18 phylogenetically related types 35, 31, 58, 33, 52, 45, 59 without considering other HPV types

\*\* = High-risk types exclusive of HPV-16 and HPV-18



**Table C**

<b>HPV types analysed</b>	<b>HPV 31, 35, 58</b>	<b>HPV 31, 33, 35, 52, 58</b>	<b>HPV 45, 59</b>	<b>HPV 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, 68.</b>
Total number of number of subjects with information available per group	412	403	421	368
Number of women infected (vaccine group)	11	28	10	58
% women infected (vaccine group) = A	2.7	6.9	2.4	15.8
Number of women infected (placebo group).	26	48	15	90
% women infected (placebo group) = B	6.3	12.2	3.6	25.3
% vaccine efficacy $1 - (A/B) \times 100$ , adjusted for relative size of vaccine and placebo group	57.9	43.0	33.5	37.7
95% confidence limits -lower limit	15.9	11.0	-46.3	16.2
95% confidence limits -upper limit	78.9	63.5	69.8	53.6
P	0.012	0.015	0.319	0.002

Samples were taken at 18 months from patients and tested for HPV infection by the types specified above.

Table D

## Efficacy against Incident Infections with 16/18 Related Types\*

	HPV Type	Vaccine		Control		Vaccine Efficacy	
		N	AR	N	AR	%	P value
16 related {	HPV-31	1	0.2	10	2.4	90.0	0.006
	HPV-33	6	1.4	6	1.4	-0.2	1.000
	HPV-35	1	0.2	3	0.7	66.5	0.624
	HPV-52	6	1.4	16	3.9	63.0	0.031
	HPV-58	5	1.2	5	1.2	0.0	1.000
18 related {	HPV-45	0	0.0	5	1.2	100.0	0.031
	HPV-59	4	0.9	2	0.5	-100.5	0.448
	All 16 related	16	4.0	32	8.1	51.1	0.017
	All 18 related	4	1.0	7	1.7	43.0	0.384
	All HR (except 16/18)	32	9.0	53	15.6	42.3	0.011

\*Cervical samples; ATP cohort





**Table E**

## Efficacy against Persistent Infections with 16/18 Related Types\*

	HPV Type	Vaccine		Control		Vaccine Efficacy	
		N	AR	N	AR	%	P value
16 related {	HPV-31	2	0.48	9	2.15	78.5	0.030
	HPV-33	3	0.71	5	1.18	40.2	0.476
	HPV-35	1	0.24	1	0.24	0.4	0.998
	HPV-52	5	1.20	21	5.10	77.1	0.001
	HPV-58	4	0.95	6	1.42	34.1	0.515
18 related {	HPV-45	1	0.24	4	0.94	75.4	0.174
	HPV-59	3	0.71	0	0.00	-	0.083
	All 16 related	11	2.7	30	7.6	65.1	0.002
	All 18 related	4	1.0	4	1.0	1.0	0.989
	All HR (except 16/18)	36	10.1	46	13.5	27.1	0.155

\*All samples; ATP cohort



**Table F**

## Efficacy against Cytological Abnormalities ass. with 16/18 Related Types\*

	HPV Type	Vaccine		Control		Vaccine Efficacy	
		N	AR	N	AR	%	P value
16 related {	HPV-31	1	0.24	5	1.20	80.1	0.123
	HPV-33	2	0.47	4	0.94	49.9	0.686
	HPV-35	0	0.00	2	0.47	100	0.499
	HPV-52	1	0.24	11	2.67	91	0.003
	HPV-58	2	0.47	2	0.47	0.2	1.000
18 related {	HPV-45	0	0.00	2	0.47	100	0.249
	HPV-59	4	0.94	2	0.47	-101	0.451
All 16 related		5	1.2	18	4.6	72.8	0.005
All 18 related		4	1.0	4	1.0	0.2	1.000
All HR (except 16/18)		10	2.8	30	8.8	68.2	< 0.001

\*ATP cohort



In tables D, E and F,

5

N = number of subjects in specific cohort

AR = Attack rate = n (number of subjects with HPV either incident infection, persistent infection or

cytological abnormality, as appropriate for the table) / N

10

% Vaccine efficacy is  $1 - (A/B) \times 100$ , adjusted for relative size of vaccine and placebo group, wherein

A = % women in vaccine group with incident infection, persistent infection or cytological abnormality, as appropriate for the table

15

B = % women in placebo group with incident infection, persistent infection or cytological abnormality, as appropriate for the table.



## DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME           1    DE    2  
CONTENANT LES PAGES    1    À    33

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

## JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME           1    OF    2  
CONTAINING PAGES    1    TO    33

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

**Claims**

1. A multivalent HPV vaccine comprising L1 proteins or immunogenic fragments thereof from HPV 16, HPV 18 and at least one other oncogenic HPV type, wherein an L1 protein or immunogenic fragment thereof from one or more HPV types  
5 selected from the group consisting of HPV 31, HPV 45, and HPV 52 is omitted from the vaccine and wherein the vaccine provides protection against infection caused by the omitted HPV type.
2. The vaccine according to claim 1 wherein an L1 protein or immunogenic fragment thereof from HPV 31 is omitted from the vaccine.
- 10 3. The vaccine according to claim 1 wherein an L1 protein or immunogenic fragment thereof from HPV 45 is omitted from the vaccine.
4. The vaccine according to claim 1 wherein an L1 protein or immunogenic fragment thereof from HPV 52 is omitted from the vaccine.
5. The vaccine according to claim 1 wherein an L1 protein or  
15 immunogenic fragment thereof from HPV 31 and from HPV 45 are omitted from the vaccine.
6. The vaccine according to claim 1 wherein an L1 protein or immunogenic fragment thereof from HPV 31 and from HPV 52 are omitted from the vaccine.
- 20 7. The vaccine according to claim 1 wherein an L1 protein or immunogenic fragment thereof from HPV 45 and from HPV 52 are omitted from the vaccine.
8. The vaccine according to claim 1 wherein an L1 protein or immunogenic fragment thereof from HPV 31 and from HPV 45 and from HPV 52 are  
25 omitted from the vaccine.
9. The vaccine according to claim 1 wherein the vaccine protects against incident infection.
10. The vaccine according to claim 1 wherein the vaccine protects against persistent infection.
- 30 11. The vaccine according to claim 1 wherein the other oncogenic HPV type is HPV 33.



12. The vaccine according to claim 1 wherein the other oncogenic HPV type is HPV 58.
13. The vaccine according to claim 1 wherein the other oncogenic HPV type is HPV 59.
- 5 14. The vaccine according to claim 1 comprising HPV 16 L1 protein or immunogenic fragment thereof, HPV 18 L1 protein or immunogenic fragment thereof, HPV 33 L1 protein or immunogenic fragment thereof and HPV 58 L1 protein or immunogenic fragment thereof.
15. The vaccine according to claim 1 wherein at least one of the L1 proteins  
10 or fragments thereof is in the form of a virus like particle.
16. The vaccine according to claim 1 wherein at least one of the L1 proteins is a truncated L1 protein.
17. The vaccine according to claim 16 wherein the at least one L1 protein is a C terminally truncated L1 protein.
- 15 18. The vaccine according to claim 1 further comprising an adjuvant.
19. The vaccine according to claim 18 wherein the adjuvant is an aluminium salt.
20. The vaccine according to claim 19 wherein the adjuvant is aluminium hydroxide.
- 20 21. The vaccine according to claim 18 wherein the adjuvant is 3D MPL.
22. The vaccine according to claim 18 wherein the adjuvant is 3D MPL and aluminium hydroxide.
- 23 A vaccine according to claim 18 wherein the adjuvant is an oil in water emulsion.
- 25 24 A vaccine according to claim 23 wherein the adjuvant additionally comprises an aluminium salt.
25. A method to protect a patient against infection caused by HPV 16, HPV 18 and at least one other HPV type selected from the group consisting of HPV 31, HPV 45 and HPV 52, the method comprising administering the vaccine of claim 1 wherein  
30 the vaccine provides protection against infection caused by the omitted HPV type.
26. The method of claim 25 wherein the omitted HPV type is HPV 31.
27. The method of claim 25 wherein the omitted HPV type is HPV 45.

28. The method of claim 25 wherein the omitted HPV type is HPV 52.

29. A method to prevent or reduce the frequency of cytological abnormalities in a patient caused by HPV 16, HPV 18 and at least one other HPV type selected from the group consisting of HPV 31, HPV 45 and HPV 52, the method comprising administering the vaccine of claim 1 wherein the vaccine prevents or reduces the frequency of cytological abnormalities caused by the omitted HPV type.

30. The method of claim 29 wherein the omitted HPV type is HPV 52.

31. The method of claim 29 wherein the omitted HPV type is HPV 45.

32. The method of claim 29 wherein the omitted HPV type is HPV 31.

33. A method to prevent the formation of histologically-confirmed CIN lesions caused by HPV 16, HPV 18 and at least one other HPV type selected from the group consisting of HPV 31, HPV 45 and HPV 52, the method comprising administering the vaccine of claim 1 wherein the vaccine prevents the formation of histologically-confirmed CIN lesions caused by the omitted HPV type.

34. The method of claim 33 wherein the omitted HPV type is HPV 52.

35. The method of claim 33 wherein the omitted HPV type is HPV 45.

36. The method of claim 33 wherein the omitted HPV type is HPV 31.

37. A method to prevent or reduce the frequency of cytological abnormalities in a patient caused by oncogenic HPV types, the method comprising administering the vaccine of claim 1.

38. A method to prevent the formation of histologically-confirmed CIN lesions in a patient caused by oncogenic HPV types, the method comprising administering the vaccine of claim 1.

39. A method to manufacture the vaccine of claim 1, the method comprising combining L1 proteins or immunogenic fragments thereof from HPV 16, HPV 18 and at least one other oncogenic HPV type, wherein the vaccine does not comprise an L1 protein or immunogenic fragment thereof from one or more HPV types selected from the group consisting of HPV 31, HPV 45, and HPV 52.

40. Use of a composition comprising HPV 16 L1 and HPV 18 L1 proteins, or immunogenic fragment thereof, in the preparation of a medicament for prevention of infection and/or disease caused by one or more of: HPV 31, HPV 45 or HPV 52.



41        Use of a vaccine comprising an HPV 16 L1 protein, or immunogenic fragment thereof, in the preparation of a medicament for the prevention of infection and/or disease caused by HPV 31, or HPV 52, or a combination thereof.

42        Use of a vaccine composition comprising an HPV 18 L1 protein, or an  
5   immunogenic fragment thereof, in the preparation of a medicament for the prevention of infection and/or disease caused by HPV 45.

43        Use of a vaccine according to claim 40 in the manufacture of a medicament for the prevention of cytological abnormalities or reduction of the frequency of cytological abnormalities in an individual caused by a type other than  
10   HPV 16 and HPV 18.

44        Use of a vaccine or combination according to claim 40 in the manufacture of a medicament for the prevention of histologically-confirmed CIN lesions (CIN 1, CIN 2, CIN 3) caused by a type other than HPV 16 and HPV 18.

45        Use according to any of claims 40 – 44 wherein the composition  
15   comprising L1 proteins or immunogenic fragments thereof from HPV 16, HPV 18 and at least one other oncogenic HPV type, wherein an L1 protein or immunogenic fragment thereof from one or more HPV types selected from the group consisting of HPV 31, HPV 45, and HPV 52 is omitted from the vaccine and wherein the vaccine provides protection against infection and/or disease caused by the omitted HPV type.