

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3456334 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **A 61 K 35/747 (2015.01)** **A 23 L 33/135 (2016.01)** **A 61 K 31/59 (2006.01)**
A 61 P 19/08 (2006.01) **C 12 Q 1/68 (2018.01)**
- (45) Oversættelsen bekendtgjort den: **2021-01-25**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-11-11**
- (86) Europæisk ansøgning nr.: **18202594.0**
- (86) Europæisk indleveringsdag: **2013-06-04**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-03-20**
- (30) Prioritet: **2012-06-04 US 201261689338 P**
- (62) Stamansøgningsnr: **13800449.4**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Biogaia AB, P.O. Box 3242, 103 64 Stockholm, Sverige**
Michigan State University, 325 E. Grand River , Suite 350, East Lansing, Michigan 48823, USA
- (72) Opfinder: **BRITTON, Robert Allen, Michigan State University, 1209 Blanchette Drive, EAST LANSING, MI Michigan 48823, USA**
McCABE, Laura Rae, Michigan State University, 1330 Orlando Drive, HASLETT, MI Michigan 48840, USA
CONNOLLY, Eamonn, BioGaia AB, Motionsvägen 3, 181 29 LIDINGÖ, Sverige
- (74) Fuldmægtig i Danmark: **Synergion AB, Box 32, S-443 21 Lerum, Sverige**
- (54) Benævnelse: **MÆLKESYREBAKTERIER TIL ANVENDELSE I FOREBYGGELSE AF KNOGLETAB HOS PATTEDYR**
- (56) Fremdragne publikationer:
EP-A1- 1 918 373
WO-A1-2006/110088
US-A1- 2008 254 011
BRENDAN E BOYCE ET AL: "TNF.ALPHA. and pathologic bone resorption", KEIO JOURNAL OF MEDICINE., vol. 54, no. 3, 1 January 2005 (2005-01-01), pages 127-131, XP055270832, JP ISSN: 0022-9717, DOI: 10.2302/kjm.54.127
SCHOLZ-AHRENS K E ET AL: "Prebiotics, probiotics, and synbiotics affect mineral absorption, bone mineral content, and bone structure", THE JOURNAL OF NUTRITION, AMERICAN SOCIETY FOR NUTRITION, US, vol. 137, no. 3, Suppl. S, 1 March 2007 (2007-03-01), pages 838S-846S, XP002737535, ISSN: 0022-3166
CHING-YI LAI ET AL: "Preventing Bone Loss and Weight Gain with Combinations of Vitamin D and Phytochemicals", JOURNAL OF MEDICINAL FOOD, vol. 14, no. 11, 1 November 2011 (2011-11-01), pages 1352-1362, XP055190442, ISSN: 1096-620X, DOI: 10.1089/jmf.2010.0232
IYER CHANDRA ET AL: "Probiotic Lactobacillus reuteri promotes TNF-induced apoptosis in human myeloid leukemia-derived cells by modulation of NF-kappaB and MAPK signalling.", CELLULAR MICROBIOLOGY JUL 2008, vol. 10, no. 7, July 2008 (2008-07), pages 1442-1452, XP002757502, ISSN: 1462-5822
BRITTON ROBERT A ET AL: "Probiotic L. reuteri Treatment Prevents Bone Loss in a Menopausal

Fortsættes ...

Ovariectomized Mouse Model", JOURNAL OF CELLULAR PHYSIOLOGY, vol. 229, no. 11, November 2014 (2014-11), pages 1822-1830, XP002757503,

DESCRIPTION

FIELD OF THE INVENTION

[0001] The present invention relates generally to medicine, pharmacology and food supplements. More specifically the invention relates to lactic acid bacteria for use in the prevention of bone loss in mammals.

BACKGROUND OF THE INVENTION

[0002] Over 40 million Americans over the age of 50 (14 million of which are men) are afflicted with low bone density or osteoporosis and its associated increased risk of fractures. Individuals with osteoporotic fractures are prone to depression, dependency and increased mortality. While aging is a major cause of osteoporosis, disease, disuse, and certain drugs can also cause bone loss at any stage in life.

[0003] The skeleton is a highly organized system that supports the body's weight, houses mesenchymal and hematopoietic stem cells, and serves as a calcium reservoir. The structure of bone comprises an outer cortical dense shell and an inner trabecular bone meshwork. Exercise can increase trabecular bone mineral density (BMD), and bone volume fraction (BVF), trabeculi thickness, and cortical BMD and thickness. In contrast, disease, disuse, and certain drugs (such as glucocorticoids) can decrease these parameters and cause osteoporosis in both males and females. Osteoporosis is defined by a reduction in bone mass (more than 2.5 standard deviations (SD) below average) and altered bone micro-architecture (such as decreased trabeculi thickness). With decreasing bone mass there is an increased risk of bone fractures. Thus, at the point of being diagnosed as osteoporotic, a patient has a 16-fold increase in fracture risk compared to someone with normal bone density. Fractures are associated with depression, dependency, and increased mortality (greater than 25% within 12 months for the elderly) and hip fractures account for over 50,000 deaths annually (National Osteoporosis Foundation (NOF) statistic). While osteoporosis is less prevalent in men, over 30% of hip fractures occur in men and mortality rates are greater for males compared to females. Currently, over 20 billion dollars are spent in the US and 30 billion dollars in the European Union to cover the direct costs of osteoporosis. Of even greater concern, it is estimated that by 2020 more than 61 million men and women in the US, over the age of 50, will have low bone density or osteoporosis (NOF statistic), and finding effective novel treatments is therefore a priority. In fact, one in three women over the age of 50 will experience an osteoporosis related fracture in their lifetime. Along with its associated increase in fracture risk, bone loss may have negative effects on metabolism and insulin secretion. Despite all the available treatments on the market, the number of osteoporotic patients is on the rise in the U.S. and worldwide. There are several reasons for this, including a lack of awareness that one is at risk early in life, an increasing elderly population, and patient noncompliance due to

unwanted medication side effects. In addition, conventional bone loss treatments are not always effective. Currently there are no alternative or natural treatments that can be used in place of osteoporosis medicines for people with low bone density or osteoporosis. Therefore, doctors are looking for new approaches to increase bone density in their patients and companies are working to improve pharmacologic bone therapeutic drugs.

[0004] Certain people are more likely to develop osteoporosis than others, some risk factors are;

*Being female

*Older age

*Family history of osteoporosis or broken bones

*Being small and thin

*Certain race/ethnicities such as Caucasian, Asian, or Hispanic/Latino although African Americans are also at risk

*History of broken bones

*Low levels of sex hormones

*Low estrogen levels in women, including menopause

*Missing periods (amenorrhea)

*Low levels of testosterone and estrogen in men

*Diet

- Low calcium intake
- Low vitamin D intake
- Excessive intake of protein, sodium and caffeine

*Inactive lifestyle

*Smoking

*Alcohol abuse

*Certain medications such as steroid medications, some anticonvulsants and others

*Certain diseases and conditions such as anorexia nervosa, rheumatoid arthritis, gastrointestinal diseases and others

[0005] Menopausal women are prone to losing bone during menopause time due to

decreased estrogen levels. Even during perimenopause (the period of 2 to 8 years before menopause) estrogen levels may start to drop off. Over time, too much bone loss can first cause osteopenia (low bone mass) and then osteoporosis.

[0006] Diagnosis of type 1 diabetes (T1D) is increasing in children and adults. While medical advances are extending patient lifespan, maintaining euglycemia remains difficult, even under therapeutic vigilance. Thus, more T1D patients (males and females) are suffering from complications, including bone loss. This means that patients begin aging/menopause with an already increased fracture risk. Once fractures occur, they can be difficult to heal, require extended hospitalizations, reduce the quality of life and increase mortality. Poor bone health also negatively affects the entire body. Postmenopausal women with T1D diabetes have higher incidences of osteoporotic fractures than women without diabetes. Children with T1D have lower bone mineral density than children without diabetes. Thus, maintaining bone health is critical for the overall quality of life of T1D patients and important for maximizing therapeutic/curative treatments involving marrow immune/progenitor cells since marrow cells and bone cells communicate.

[0007] Type 2 Diabetes (T2D) patients are also at higher risk of osteoporotic fractures than non-diabetics.

[0008] The two key components to strengthening bone and preventing osteoporosis are 1) attaining maximum bone density and 2) preventing bone loss during adulthood and aging. Bone remodeling occurs because bone is dynamic and constantly adapts to environmental cues to form or resorb bone. Targeted bone remodeling through the activities of osteoblasts (bone forming cells) and osteoclasts (bone resorbing cells) maintains blood calcium levels within a critical range while keeping bone strong at sites where support is needed. When formation and resorption activities are in balance there is no net gain or loss of bone, however when formation is decreased and/or resorption is increased then bone loss ensues.

[0009] Increased osteoclast activity results in bone resorption. Osteoclasts are derived from hematopoietic stem cells. These cells give rise to cells of the monocyte/macrophage lineage which, under the right conditions, develop into osteoclast precursors. Further signaling through factors such as RANKL (located on osteoblast surfaces) stimulate osteoclast maturation. Mature osteoclasts express enzymes involved in bone matrix degradation (including cathepsin K and TRAP5b).

[0010] Increased osteoblast activity results in bone formation, which can be regulated at several levels including 1) lineage selection, 2) maturation and 3) death. Because bone marrow stromal cells (BMSC) give rise to osteoblasts, adipocytes and other cell types, selection of one lineage (adipocyte) could be at the cost of another (osteoblast). This is supported by the reciprocal relationship between bone adiposity and mineral density recognized with aging, limb unloading, cell culture models, and type I (T1) diabetes. Osteoblast activity can be further regulated through death/apoptosis. An increase in osteoblast death will result in fewer bone making cells and therefore bone loss. Examples include the rapid bone

adaptation to disuse/unloading, which results in bone loss, increased marrow adiposity, and increased bone cell death. Aging also increases bone cell apoptosis. Many factors contribute to modulating some or all aspects of osteoblast regulation (lineage, maturation, death) including: positive factors such as TGF β , bone morphogenic proteins (BMPs), parathyroid hormone (PTH), and Wnts and negative factors such as cytokines.

[0011] Bisphosphonates are one of the most common treatments for osteoporosis. These compounds incorporate into the bone mineral and inhibit bone catabolism by osteoclasts and are effective at reducing fractures. However, many of these compounds need to be taken on an empty stomach and can cause gastric reflux and nausea resulting in reduced patient compliance. There is also concern about the length of time that these compounds reside in bone and their long-term impact on bone remodeling and strength. Selective estrogen receptor modifiers (SERMS) are another therapeutic treatment, but they still carry some concerns with regard to cancer. Hormone replacement therapy has been studied as useful in preventing or slowing the occurrence of osteoporosis, but sustained use of hormone replacement over many years may increase women's risk of breast cancer, may increase incidence of venous thrombosis (blood clots), exacerbation of pre-existing liver diseases and an increased risk of endometrial cancer as well as hypertension. Amgen has a drug under development (that is similar to osteoprotegerin) that works by modifying the RANKL/RANK system and hence suppresses osteoclast activity. Intermittent PTH treatment is an anabolic treatment, but this intravenous treatment is expensive and only indicated for severe osteoporotic patients. Taken together, it is not surprising that many people diagnosed with low bone density are confused about what to do. Many people do not want to take medication for fear of long-term effects. While weight bearing exercise and adequate calcium intake are two natural approaches, they cannot always overcome effects of disease, medications, and aging.

[0012] Brendan E Boyce et al. ("TNF α and pathologic bone resorption", Keio Journal of Medicine, vol.54, no.3, 1 January 2005, p. 127-131) describes the involvement of the pro-inflammatory cytokine TNF α in osteoclastic bone resorption causing bone loss around affected joints and teeth which is associated with chronic inflammatory bone diseases. It is also stated that TNF α is not absolutely required for osteoclastogenesis, erosive arthritis or osteolysis, as all these events could occur in the absence of TNF α .

[0013] Scholz-Ahrens K et al. ("Prebiotics, probiotics, and synbiotics affect mineral absorption, bone mineral content and bone structure", The Journal of Nutrition, American Society for Nutrition, US, vol.137, no.3, Suppl. S, 1 March 2007, p.838S-846S.) discusses the involvement of prebiotics, probiotics and synbiotics on bone mineral content and bone structure. In this regard, it was concluded that "prebiotics" (i.e. nondigestible food ingredients that beneficially affect the host by selectively stimulating the growth and/or activity of a limited number of bacteria in the gastrointestinal tract) are the most promising but also best investigated substances with respect to a bone-health promoting potential, compared with probiotics and synbiotics.

[0014] US2008/254011 concerns the selection of lactic acid bacteria based on their capability

of increasing the BSH (Bile Salt Hydrolase)-activity and consequently lowering serum LDL-cholesterol and simultaneously decreasing the pro-inflammatory cytokine TNF- α levels for use in prophylaxis and/or treatment of atherosclerosis and other cardiovascular diseases.

[0015] Ching-Yi Lai et al. ("Preventing Bone Loss and Weight Gain with Combinations of Vitamin D and Phytochemicals", Journal of Medicinal Food, vol.14, no.11, 1 November 2011, p.1352-1362") describes synergistic effects of a combination of phytochemicals with vitamin D and that such combinations may be effective in reducing bone loss and weight gain after menopause.

[0016] WO2006/110088A1 concerns the selection and use of lactic acid bacteria for reducing inflammation in mammals. The selection of the lactic acid bacteria is based on determining a decrease in TNF α levels in a cell line obtained from a human source.

[0017] EP 1918373 A1 discloses an anti-obesity agent containing as an active ingredient a microorganism belonging to the *Lactobacillus reuteri* species which is capable of producing lipases having specific amino acid sequences. The anti-obesity agent is described as enabling that the patient can ingest a regular meal without fat being absorbed into the body.

SUMMARY OF THE INVENTION

[0018] Herein is described a method for how to find a lactic acid bacterial strain that may prevent bone loss, especially in menopausal women, in diabetics, in osteopenic that includes for example young men with large energy intake and low exercise frequency.

[0019] It is also described how to use products containing such strains in menopausal women to prevent bone loss.

[0020] It is also described how to use products containing such strains in women who have had a hysterectomy to prevent bone loss.

[0021] It is also described how to use products containing such strains in men, including but not limited to diabetic, young men with metabolic disturbance and osteopenic men, to prevent bone loss.

[0022] It is also described how to use such product in combination with therapies for bone loss or bone formation in order to reduce the dose of such drugs to be able to minimize side effects.

[0023] It is also described how to improve bone repair after fracture.

[0024] Herein it is described a method for selecting a lactic acid bacterial strain for use in preventing or treating bone loss, comprising selecting a lactic acid bacterial strain having at

least 95 % identity to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1), and harboring an identical nucleotide relative to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1) in at least one of the following four positions: C in base pair 271 391, G in base pair 453 538, G in base pair 529 228, and C in base pair 599 338.

[0025] The method may comprise selecting a lactic acid bacterial strain having at least 96%, such as 97%, such as 98%, such as 99% identity to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1), and harboring an identical nucleotide relative to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1) in at least one of the following four positions: C in base pair 271 391, G in base pair 453 538, G in base pair 529 228, and C in base pair 599 338.

[0026] Herein is also described a method for selecting a lactic acid bacterial strain, such as a *Lactobacillus reuteri* strain, for use in preventing or treating bone loss, comprising selecting a *Lactobacillus reuteri* harboring an identical nucleotide relative to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1) in at least one of the following four positions: C in base pair 271 391, G in base pair 453 538, G in base pair 529 228, and C in base pair 599 338.

[0027] The lactic acid bacterial strain may harbor at least two of said four nucleotides, such as at least three of said four nucleotides, such as all four of said nucleotides.

[0028] The present invention provides a lactic acid bacterial strain having at least 99 % identity to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1), and harboring an identical nucleotide relative to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1) in all of the following four positions: C in base pair 271 391, G in base pair 453 538, G in base pair 529 228, and C in base pair 599 338, for use in the prevention or treatment of bone loss.

[0029] It is also described herein a lactic acid bacterial strain that has at least 96%, such as 97%, such as 98%, such as 99% identity to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1), and harbors an identical nucleotide relative to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1) in at least one of the following four positions: C in base pair 271 391, G in base pair 453 538, G in base pair 529 228, and C in base pair 599 338.

[0030] The lactic acid bacterial strain may harbor at least two of said four nucleotides, such as at least three of said four nucleotides, such as all four of said nucleotides.

[0031] The present invention also provides a composition comprising a lactic acid bacterial strain having at least 99 % identity to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1), and harboring an identical nucleotide relative to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1) in all of the following four positions: C in base pair 271 391, G in base pair 453 538, G in base pair 529 228, and C in base pair 599 338.

[0032] It is also described herein a lactic acid bacterial strain that has at least 96%, such as 97%, such as 98%, such as 99% identity to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1), and harbors an identical nucleotide relative to the genome of *L reuteri* JCM 1112 (SEQ ID NO:

1) in at least one of the following four positions: C in base pair 271 391, G in base pair 453 538, G in base pair 529 228, and C in base pair 599 338.

[0033] It is also described a lactic acid bacterial strain that harbors at least two of said four nucleotides, such as at least three of said four nucleotides, such as all four of said nucleotides.

[0034] The present invention also provides the composition for use in preventing bone loss in menopausal women, women who have had hysterectomy, diabetics, osteopenic individuals, osteoporotic individuals, and individuals with metabolic disturbance.

[0035] The present invention also provides the composition for use in improving bone repair after fracture.

[0036] The present invention also provides the above-described composition in combination with vitamin D for use in preventing or treating bone loss.

[0037] The present invention also provides the above-described composition in combination with a hormone (for use in hormone replacement therapy) for use in preventing or treating bone loss.

[0038] According to the present invention, the above-described composition may be a pharmaceutical composition (optionally comprising at least one pharmaceutically acceptable excipient), or a food product or a food supplement (optionally comprising at least one food-grade excipient, as known to a person of ordinary skill in the art).

[0039] It is also described herein a method for the treatment or prevention of bone loss, comprising administering, to an individual, a lactic acid bacterial strain having at least 95 % identity to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1), and harboring an identical nucleotide relative to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1) in at least one of the following four positions: C in base pair 271 391, G in base pair 453 538, G in base pair 529 228, and C in base pair 599 338.

[0040] It is also described herein the lactic acid bacterial strain that has at least 96%, such as 97%, such as 98%, such as 99% identity to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1), and harbors an identical nucleotide relative to the genome of *L reuteri* JCM 1112 (SEQ ID NO: 1) in at least one of the following four positions: C in base pair 271 391, G in base pair 453 538, G in base pair 529 228, and C in base pair 599 338.

[0041] It is also described herein, the lactic acid bacterial strain that harbors at least two of said four nucleotides, such as at least three of said four nucleotides, such as all four of said nucleotides.

BRIEF DESCRIPTION OF THE DRAWINGS

[0042]

Figure 1 shows microbial community clustering in jejunum and ileum.

Figure 2 shows the suppression of bone loss by *L. reuteri* ATCC PTA 6475.

Figure 3 shows the effect on bone loss in different *L. reuteri* strains.

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS THEREOF

[0043] Chronic inflammatory diseases are frequently associated with systemic bone loss. In the NIH grant abstract, grant number 1R21AT005472-01A1, McCabe suggests that therapies which improve overall intestinal health have the potential to benefit bone health. McCabe and Britton found out that *L. reuteri* treatment decreased TNF levels in the ileum and increased bone volume in healthy male but not female mice and suggest that *L. reuteri* increases bone density in a gender dependent manner through suppression of intestinal inflammation and upregulation of bone formation. They suggest that they have a novel way of increasing bone mass by use of a probiotic bacterium that attenuates intestinal inflammation. This is a gender dependent up regulation of bone formation, not prevention of bone loss, associated to *L. reuteri*'s anti-inflammatory properties, unlike the present invention where certain specifically selected strains are used to prevent bone loss in both male and female.

[0044] Probiotics can increase chicken cortical bone thickness and reduce bone loss in aging mice. Narva et al. described in "Effects of bioactive peptide, valyl-prolyl-proline (VPP), and lactobacillus helveticus fermented milk containing VPP on bone loss in ovariectomized rats." that *L. helveticus* fermented milk prevent bone loss, the effect might be due to the peptide valyl-prolyl-proline. Narva et al. further described in "The effect of Lactobacillus helveticus fermented milk on acute changes in calcium metabolism in postmenopausal women" that fermentation of milk with *Lactobacillus helveticus* had a positive acute effect on calcium metabolism.

[0045] Yeo et al. suggest in "Angiotensin I-converting enzyme inhibitory activity and bioconversion of isoflavones by probiotics in soymilk supplemented with prebiotics" that probiotic incorporated into soymilk supplemented with prebiotic could potentially be used as a dietary therapy in for example osteoporosis.

[0046] Kim et al. showed in "Effects of a Lactobacillus casei 393 fermented milk product on bone metabolism in ovariectomised rats" that *L. casei* 393 FMP had a preventative effect on bone loss in ovariectomised rats.

[0047] However none of the above mentioned prior art neither alone nor in combination teaches how one can select specific probiotic strains that are effective for preventing bone loss.

[0048] It is described herein a method for selecting lactic acid bacterial strains effective for preventing bone loss in humans and strains that have been selected according to the presented method. Products such as foods, nutritional additives and formulations, pharmaceuticals or medical devices containing whole cells or components derived from these strains may be formulated as is known in the art, and generally include an ingestible support as known plus the lactic acid bacterial strain, or its derived component.

[0049] Based on prior art it would be natural to think that a strain's capability of preventing bone loss would be associated with its general effect on intestinal health or its anti-inflammatory properties, however the inventors have surprisingly found out that these properties are not predictive on the efficiency on preventing bone loss. *Lactobacillus reuteri* ATCC PTA 6475 and *Lactobacillus reuteri* ATCC PTA 4659 are two almost identical strains, which are both anti-inflammatory and improve overall intestinal health. It is natural to assume that these strains therefore would have the same effect on bone loss as well. However the inventors have shown that these strains do not have the same impact on preventing bone loss and based on this observation they have invented a novel way of selecting lactic acid bacterial strains, such as for example *Lactobacillus reuteri* that will be effective for treatment and/or prevention of bone loss.

[0050] Lactic acid bacteria specifically selected by the method presented herein may be administered to humans to prevent bone loss.

[0051] *L. reuteri* ATCC PTA 6475 and ATCC PTA 4659 differ in four SNPs, which are important for the bacteria's ability to prevent bone loss. These SNPs are shown in Walter et al. (Walter et al. Host-microbial symbiosis in the vertebrate gastrointestinal tract and the *Lactobacillus reuteri* paradigm; PNAS, vol. 108 p. 4645-4652). For the SNP analysis, sequencing results were mapped onto a reference genome (*L. reuteri* JCM 1112, GenBank accession no AP007281, SEQ ID NO: 1). Seven SNPs were found in *L. reuteri* ATCC PTA 4659, and three of them were also found in *L. reuteri* ATCC PTA 6475 (SNP 4 located at bp 567 368, SNP 6 located at bp 968 088, and SNP 8 located at bp 1 358 460, referring to the reference genome, *L. reuteri* JCM 1112, GenBank accession no AP007281, SEQ ID NO: 1). The remaining four unique SNPs (for the purpose of the present text, hereinafter called SNP 1, SNP 2, SNP 3 and SNP 5, respectively) constitute the genomic differences between *L. reuteri* ATCC PTA 6475 and *L. reuteri* ATCC PTA 4659. Said four SNPs are located at:

- bp 271 391 (SNP 1),
- bp 453 538 (SNP 2),
- bp 529 228 (SNP 3), and
- bp 599 338 (SNP 5),

(referring to the reference genome, *L. reuteri* JCM 1112, GenBank accession no AP007281;

SEQ ID NO: 1).

[0052] SNP 1 is located in a gene coding for a conserved hypothetical protein (*L. reuteri* JCM 1112: <http://www.ncbi.nlm.nih.gov/protein/183224225>), SNP 2 is located in a gene coding for a chloride channel protein (*L. reuteri* JCM 1112: <http://www.ncbi.nlm.nih.gov/protein/183224386>), SNP 3 is located in a gene coding for an ATP synthase gamma subunit (*L. reuteri* JCM 1112: <http://www.ncbi.nlm.nih.gov/protein/183224455>) and SNP 5 is located in a gene coding for a DNA mismatch repair protein HexB (*L. reuteri* JCM 1112: <http://www.ncbi.nlm.nih.gov/protein/183224511>). The SNPs involved in this invention are the ones that match *L. reuteri* ATCC PTA 6475, the sequence of which has identical nucleotides as *L. reuteri* JCM 1112 in the positions of SNP 1, SNP 2, SNP 3 and SNP 5). Listed below are the nucleotides that differ between the strains *L. reuteri* ATCC PTA 6475 and 4659:

SNP 1)

a gene coding for a hypothetical protein, where nucleotide 267 has been changed in ATCC PTA 4659 from a C (as in ATCC PTA 6475 and JCM 1112) to a T.

SNP 2)

the gene coding for the chloride channel protein, where nucleotide 373 has been changed in ATCC PTA 4659 from a G (as in ATCC PTA 6475 and JCM 1112) to an A.

SNP 3)

the gene coding for ATP synthase gamma subunit, where nucleotide 296 has been changed in ATCC PTA 4659 from a G (as in ATCC PTA 6475 and JCM 1112) to an A.

SNP 5)

the gene coding for the HexB protein, where nucleotide 1966 has been changed from a C (as in ATCC PTA 6475 and JCM 1112) to a T.

[0053] In the selection method, strains are sought that, in at least one of these SNPs, harbor the same nucleotides as *L. reuteri* ATCC PTA 6475 for the above mentioned SNPs.

[0054] The microbiota plays an important role in bone loss; many patients suffering from bone loss have a disturbed intestinal microbiota. Lactic acid bacteria that are able to reestablish the normal microbial community in the GI tract are surprisingly more effective in preventing bone loss.

[0055] It is described herein, a unique method of selection, selecting strains effective for preventing bone loss. The ability to reestablish the total gut microbial composition is surprisingly also important for the function in preventing bone loss. The inventors have found out that strains capable of reestablishing an altered microbial community to normal and/or harboring at least one of the four specific SNPs are effective for preventing bone loss.

[0056] The ability to prevent bone loss is unique for certain strains and is not at all general for all lactic acid bacteria. When selecting effective strains it is not sufficient to use anti-inflammatory capacity as selection criteria since the inventors clearly show that this effect is not

dependent on anti-inflammatory features. *L. reuteri* ATCC PTA 6475 and *L. reuteri* ATCC PTA 4659 are both anti-inflammatory strains, but *L. reuteri* ATCC PTA 6475 is much more effective when used for prevention of bone loss, *L. reuteri* ATCC PTA 4659 is not selected according to this invention. Specific lactic acid bacterial strains selected according to the present invention may be used for preventing bone loss in general and the embodiments below are not intended to limit the scope of this invention, but to exemplify preferred embodiments.

[0057] Vitamin D is crucial to bone health and people with low levels of vitamin D have lower bone density or bone mass. People that do not get enough vitamin D may lose bone, since vitamin D is required to absorb calcium. The inventors have seen that an altered microbiota will lead to vitamin D deficiency and bone loss, administration of lactic acid bacteria selected according to the present invention will reestablish the microbiota and thereby increase the intestinal vitamin D absorption and restore the levels of vitamin D. It is also an option to combine vitamin D with the selected strains in order to get an even more efficient method/product for preventing bone loss.

[0058] T1D patients suffer from complications such as bone loss. Patients suffering from T1D will as a result of the condition have an altered microbiota. Administration of lactic acid bacteria according to the present invention will reestablish the microbiota and prevent bone loss.

[0059] High bone density during youth and adulthood can help prevent diseases like osteoporosis later in life. This is due to the fact that high bone density will allow a higher degree of bone loss before reaching a bone density within the osteoporosis zone. Thus, it is an object of the present invention to prevent bone loss by administering lactic acid bacterial strains according to the present invention, to young and adult people, this will help individuals to obtain maximum bone density to prevent osteoporosis from occurring later in life. Specific lactic acid bacteria selected according to the present invention prevents bone loss in healthy recipient as well as those suffering from bone loss.

[0060] Administration of selected lactic acid bacteria may be combined with hormone replacement therapy. Such a combination would make it possible to reduce the amount of hormones and thereby reduce the side effects, such as reducing the risk of cancer.

[0061] Lactic acid bacteria selected for preventing bone loss would preferably be administered to menopausal women and osteopenic men who are prone to develop osteoporosis, and administration of selected lactic acid bacteria will prevent bone loss and thus preventing low bone density and osteoporosis.

[0062] The inventors have seen that estrogen depletion alters the gut microbiota, treatment with lactic acid bacteria according to the present invention will reestablish the microbiota in people suffering from decreased estrogen levels, including but not limited to menopausal women and women who had hysterectomy, consequently preventing bone loss.

[0063] Lactic acid bacterial strains according to the present invention may also be used to

improve fracture repair.

[0064] In order to reduce side effects of drugs, such as for example bisphosphonates and hormone replacement therapy used to treat bone loss it is possible to combine drugs with administration of lactic acid bacteria and thereby reduce the dose, which will minimize the side effects.

EXAMPLE 1

Study of *L. reuteri* ATCC PTA 6475's ability to reestablish altered microbial communities in ovx mice.

[0065] There are significant changes in the intestinal microbial communities of control (non-ovx), ovx and ovx fed by *L. reuteri*.

Experimental groups and tissue collection.

[0066] In order to measure the effects of ovariectomy (ovx) and *L. reuteri* 6475 treatment of ovx mice we compared three experimental groups of animals. Control mice were non-ovx mice that received a vehicle control gavage three times per week. Ovx mice received a vehicle control gavage three times per week. Ovx + *L. reuteri* 6475 were mice that received 300 µl of overnight *L. reuteri* 6475 three times per week for four weeks. At the end of the experiment mice were euthanized and tissue samples from the stomach, duodenum, jejunum, ileum, proximal and distal colons were isolated and saved for microbial ecology analysis.

DNA extraction

[0067] Murine intestinal tissue was placed in MoBio Ultra Clean Fecal DNA Bead Tubes (cat.# 12811-100-DBT) containing 360 µl Buffer ATL (Qiagen cat.#19076) and lysed on a Mini-Beadbeater-8 (BioSpec Products) for 1 minute at full speed. DNA was extracted from murine intestinal tissue using Qiagen DNeasy Blood and Tissue kit (cat.# 69504). The tissue was further disrupted by adding 40 µl proteinase K (Qiagen, cat.# 19133) and incubating at 55°C for 1 hour. DNA was extracted using the Qiagen DNeasy Blood and Tissue kit (cat.#69504). DNA yield was quantified using a Nanodrop 1000.

PCR amplification

[0068] Bacterial 16S sequences were amplified for 454 sequencing from murine intestinal tissue using the V3-V5 barcoded primer set and amplification protocol developed by the Broad Institute for the Human Microbiome Project. Barcoded forward primers were synthesized by IDT DNA Technologies and the reverse primer was synthesized by Sigma. Barcoded forward primers were diluted to a working concentration of 4 μ M in 96 well plates; the reverse primer was added to each well to a final concentration of 4 μ M. Triplicate reactions in a 25 μ l volume were prepared containing 400 μ g murine intestinal DNA, 2 μ l 4 μ M primers, and 0.15 μ l Accuprime HiFi Taq polymerase in IX Accuprime Buffer II (Invitrogen, cat.# 12346086). Reactions were amplified in an Eppendorf Pro aluminum plate thermal cycler with a 2 minute 95°C denaturation, followed by 30 cycles of 95°C x 20 sec, 50°C x 30 sec, 72°C x 5min.

Amplification product purification

[0069] 16S amplification products were purified using Ampure Agencourt XP beads (Beckman Coulter, cat# A63880). First, triplicate reactions for each sample were combined into 1.7ml microfuge tubes and Ampure XP beads were added at a 0.7X volume ratio. After vortexing, the mixed samples were incubated for 10 minutes at room temperature then placed on a magnetic stand to separate the beads (Invitrogen, cat.#123-21D). The beads were washed according to the manufacturer's protocol with 2 washes of 200 μ l of 70% ethanol. The beads were dried at 37°C for 5 minutes, and DNA was eluted with 20 μ l 10 μ M Tris buffer/0.1 μ M EDTA. The eluent was separated from the beads on the magnetic stand, transferred to a new 1.7ml microfuge tube, and quantified using the Quant-It dsDNA high-sensitivity assay kit (Invitrogen, cat# Q33120). Equal amounts of each sample were then pooled into one tube for 454 sequencing.

454 sequencing and sequence analysis

[0070] 454 sequencing was performed using the GS Junior (Roche) using Titanium chemistry. In addition to the standard filters utilized by the GS Jr. to identify passed reads we utilized a modified amplicon processing algorithm to reduce the number of incorrectly discarded sequences. 16S rRNA sequences were aligned by the Ribosomal Database Project staff at MSU to *E. coli* 16S sequences and trimmed at *E. coli* 16S nucleotide positions 617 to 900. Subsequent processing and analysis (including diversity metrics) were performed using MOTHUR v.1.21 (<http://www.mothur.org/wiki/>). ANOSIM (analysis of similarity) and principle coordinate analysis were performed using the software package PAST. The accompanying figures and table utilize the Bray-Curtis method for measuring the level of dissimilarity between two or more microbial communities. In these analyses we chose an operational taxonomic unit (OUT) cutoff of 0.03, which is considered to be viewing the communities at the species level. From these data we conclude that in that treatment of ovariectomized mice with *L. reuteri* ATCC PTA 6475 causes a significant shift in the both the ileal and jejunal microbial communities, which correlates with improved bone health. (ovx+lacto in the table 1).

Table 1: ANOSIM analysis at species level using Bray-Curtis dissimilarity matrix, ** indicates statistical significance.

Tissue	Comparison	R Value (p value)
Jejunum	wt-ovx-ovxl acto	0.3367 (0.0183)*
	wt-ovx	0.0443 (0.3633)
	wt-ovxlacto	0.6078 (0.0250)*
	ovx -ovxlacto	0.3297 (0.0712)
Ileum	wt-ovx-ovxl acto	0.2068 (0.0084)*
	wt-ovx	0.1710 (0.1180)
	wt-ovxlacto	0.2540 (0.0290)*
	ovx -ovxlacto	0.2209 (0.0206)*

[0071] Three-way comparison of the wild-type, ovx, and ovx treated with *L. reuteri* showed significant shifts in microbial communities (table 1). These differences are largely driven by substantial shifts in communities after *L. reuteri* treatment. Principle coordinate analysis of microbial communities from the wild-type control group (triangle Δ), ovx group (circle •) and the ovx group treated with *L. reuteri* (square ■) was used to visualize how communities clustered in the jejunum and the ileum. Figure 1 shows that ovx mice treated with *L. reuteri* form a cluster of communities that is distinct from wild-type and ovx communities in both the jejunum and ileum. Several OTUs that were classified as *Clostridiales* are the main groups of bacteria that are driving the separation of the *L. reuteri* treated Ovx communities from the other two groups.

EXAMPLE 2

Study of L. reuteri ATCC PTA 4659's ability to reestablish altered microbial communities in ovx mice.

[0072] The experiment is performed as in example 1, but *L. reuteri* ATCC PTA 4659 is used instead of *L. reuteri* ATCC PTA 6475.

[0073] *L. reuteri* ATCC PTA 4659 treatment is not able to restore ovx communities toward control.

EXAMPLE 3

Identification of certain SNPs

Illumina Sequencing of *L. reuteri* Genomes

[0074] *L. reuteri* strains used in this study were ATCC PTA 4659 and 6475 grown in MRS media (Difco) and genomic DNA prepared by using the Qiagen Genomic-Tip System. DNA was fragmented by 20 min sonication (130 W) to obtain an average fragment size of 500 bp, then further purified and concentrated with QIAquick PCR Purification Spin Columns (Qiagen). Treatment to remove 3' overhangs and fill in 5' overhangs resulted in blunt-ended genomic fragments. An adenine residue was added by terminal transferase to the 3' end, and the resulting fragments were ligated to Solexa adapters. The products were separated by agarose gel electrophoresis, and the band between 150 and 200 bp was excised from the gel. The DNA fragments were extracted from the agarose slice using a QIAquick Gel Extraction Kit (Qiagen). Adapter-modified DNA fragments were enriched by an 18-cycle PCR using Solexa universal adapter primers. The DNA fragment library was quantitated, and then diluted to a 10-nM working stock for cluster generation. Adapter-ligated fragments (2 nM) were denatured in 0.1 M NaOH for 5 min, then further diluted to a final 9 pM concentration in 1 mL of prechilled hybridization buffer, and introduced onto the Solexa flow cell using the Cluster Station. Following isothermal amplification, clusters were made single-stranded by 0.1 M NaOH denaturation, metered across the flow cell by the Solexa Cluster Station. A sequencing primer complementary to one Solexa adapter was added to prime the single strands of each cluster. Once hybridized and with excess primer removed by a wash, the flow cell was ready for sequencing. The Solexa Genome Analyzer II was programmed to provide up to 36 sequential flows of fluorescently labeled, 3'-OH blocked nucleotides and polymerase to the surface of the flow cell, thus producing a fixed 36-bp read length. After each base incorporation step, the flow cell surface was washed to remove reactants and then imaged by microscope objective. The experiments collected 300 tiled images ("tiles") per flow cell lane, each containing on average 30,000 clusters.

SNP Analysis

[0075] The two lanes' sequencing results were mapped onto the reference genome *L. reuteri* JCM 1112T (GenBank accession no AP007281) separately. The mapping software Maq version 0.6.6 (<http://maq.sourceforge.net/maq-man.shtml>) was used to perform the mapping (default parameters). SNPs were identified and validated by the MAQ software, and classified into coding SNP and intergenetic SNPs. Coding SNPs were identified as synonymous and nonsynonymous. The SNPs were finally verified by PCR amplification of the surrounding region, followed by Sanger sequence determination.

EXAMPLE 4

Method of selection of strains

[0076] The selection of strains effective for prevention of bone loss is based on the ability to restore altered microbial communities. Based on the results of examples 1 and 2, *L. reuteri* ATCC PTA 6475 is selected based on the fact that this strain has the ability to restore altered microbial communities. *L. reuteri* ATCC PTA 4659 is not selected based on the results of example 2.

EXAMPLE 5***Method of selection of strains***

[0077] The selection of strains effective for prevention of bone loss is based on the presence of certain SNPs. As a consequence of the results of example 3, *L. reuteri* ATCC PTA 6475 is selected, since it harbors all of the four sought SNPs. Due to the lack of these SNPs *L. reuteri* ATCC PTA 4659 is not selected.

EXAMPLE 6***Method of selection of strains***

[0078] The selection of strains effective for prevention of bone loss is based on example 3 and 4 and 5, strains harboring at least one of the four sought SNPS as well as the capacity to restore altered microbial communities is selected. Based on these criteria *L. reuteri* ATCC PTA 6475 is selected.

EXAMPLE 7***L. reuteri* ATCC PTA 6475 suppresses ovx induced bone loss**

[0079] In this study ovariectomized (ovx) BALB/c mice were used as a mouse model for bone loss. Mice (12 weeks old) were ovariectomized and divided into two groups were the first group was treated with *L. reuteri* ATCC PTA 6475 three times a week during four weeks. BALB/c that had not been ovariectomized were used as a control group. Distal femur bone volume fraction

(BV/TV) and bone TRAP5 RNA (relative to HPRT) were measured. Mice treated with *L. reuteri* ATCC PTA 6475 showed the same bone volume fraction as the control group. Further it was to be seen that TRAP5 (a marker of osteoclast function) is returned to baseline (control group) upon *L. reuteri* ATCC PTA 6475 treatment.

[0080] Figure 2 shows that the suppression of bone loss by *L. reuteri* ATCC PTA 6475 is nearly 100% and that the expression of TRAP5 is returned to baseline.

EXAMPLE 8

The selected *L. reuteri* ATCC PTA 6475 is superior to the non-selected *L. reuteri* ATCC PTA 4659 in suppressing bone loss.

[0081] In this experiment we gavaged animals three times per week with the *L. reuteri* ATCC PTA 6475 and *L. reuteri* ATCC PTA 4659 strains while also providing the strains continuously in the drinking water for 28 days. Distal femur bone volume fraction (BV/TV) was measured by μ CT. *L. reuteri* ATCC PTA 6475 suppressed bone loss and was indistinguishable from control mice (Fig.3). *L. reuteri* ATCC PTA 4659 did not suppress bone loss to a sufficient level that reached statistical significance ($p < .01$). *L. reuteri* ATCC PTA 4659 is not as effective as the selected strain *L. reuteri* ATCC PTA 6475.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US2008254011A [0014]
- WO2006110088A1 [0016]
- EP1918373A1 [0017]

Non-patent literature cited in the description

- **BRENDAN E BOYCE et al.** TNF α and pathologic bone resorption *Keio Journal of Medicine*, 2005, vol. 54, 3127-131 [0012]
- Prebiotics, probiotics, and synbiotics affect mineral absorption, bone mineral content and bone structure **SCHOLZ-AHRENS K et al.** *The Journal of Nutrition* American Society for Nutrition 2007 03 01 vol. 137, 838S-846S [0013]
- **CHING-YI LAI et al.** Preventing Bone Loss and Weight Gain with Combinations of Vitamin D and Phytochemicals *Journal of Medicinal Food*, 2011, vol. 14, 111352-1362 [0015]
- **NARVA et al.** Effects of bioactive peptide, valyl-prolyl-proline (VPP), and lactobacillus helveticus fermented milk containing VPP on bone loss in ovariectomized rats, [0044]
- The effect of *Lactobacillus helveticus* fermented milk on acute changes in calcium metabolism in postmenopausal women, [0044]
- **YEO et al.** Angiotensin I-converting enzyme inhibitory activity and bioconversion of isoflavones by probiotics in soymilk supplemented with prebiotics, [0045]
- **KIM et al.** Effects of a *Lactobacillus casei* 393 fermented milk product on bone metabolism in ovariectomised rats, [0046]
- **WALTER et al.** Host-microbial symbiosis in the vertebrate gastrointestinal tract and the *Lactobacillus reuteri* paradigm *PNAS*, vol. 108, 4645-4652 [0051]

PATENTKRAV

1. Mælkesyrebakteriestamme med mindst 99 % identitet med nukleinsyresekvensen af genomet af *L reuteri* JCM 1112 (SEQ ID NO: 1), og som har et identisk nukleotid i forhold til nukleinsyresekvensen af genomet af *L reuteri* JCM 1112 (SEQ ID NO: 1) i alle de følgende fire positioner: C i basepar 271 391, G i basepar 453 538, G i basepar 529 228 og C i basepar 599 338, til anvendelse i forebyggelse eller behandling af knogletab.
 2. Sammensætning, der omfatter en mælkesyrebakteriestamme med mindst 99 % identitet med nukleinsyresekvensen af genomet af *L reuteri* JCM 1112 (SEQ ID NO: 1), og som har et identisk nukleotid i forhold til nukleinsyresekvensen af genomet af *L reuteri* JCM 1112 (SEQ ID NO: 1) i alle de følgende fire positioner: C i basepar 271 391, G i basepar 453 538, G i basepar 529 228 og C i basepar 599 338, til anvendelse i forebyggelse eller behandling af knogletab.
 3. Sammensætning til anvendelse ifølge krav 2, hvor knogletabet forebygges hos menopausale kvinder, kvinder, der har fået foretaget hysterektomi, diabetikere, individer med osteopeni, individer med osteoporose og individer med metabolismeforstyrrelse.
 4. Sammensætning til anvendelse ifølge krav 2, hvor forebyggelsen eller behandlingen af knogletab forbedrer knoglereparation efter fraktur.
 5. Sammensætning til anvendelse ifølge krav 2, hvor sammensætningen anvendes i kombination med D-vitamin eller et hormon.
 6. Sammensætning til anvendelse ifølge et hvilket som helst af kravene 2-5, hvilken sammensætningen er en farmaceutisk sammensætning, et fødevareprodukt eller et kosttilskud.
 7. Mælkesyrebakteriestamme til anvendelse ifølge krav 1, hvor bakteriestammen er *L. reuteri* JCM 1112.
 8. Sammensætning til anvendelse ifølge krav 2, hvor bakteriestammen er *L. reuteri* JCM 1112.
-

DRAWINGS

Figure 1




Figure 2




Figure 3

