

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2005249566 B2

(54) Title
Method for treating multiple sclerosis

(51) International Patent Classification(s)
C07K 16/28 (2006.01) **A61P 25/00** (2006.01)
A61K 39/395 (2006.01)

(21) Application No: **2005249566** (22) Date of Filing: **2005.06.02**

(87) WIPO No: **WO05/117978**

(30) Priority Data

(31) Number
60/576,993 (32) Date
2004.06.04 (33) Country
US

(43) Publication Date: **2005.12.15**
(44) Accepted Journal Date: **2010.11.11**

(71) Applicant(s)
Genentech, Inc.

(72) Inventor(s)
Frohna, Paul A.

(74) Agent / Attorney
Griffith Hack, Level 3 509 St Kilda Road, Melbourne, VIC, 3004

(56) Related Art
WO 2004/035607 A1

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
15 December 2005 (15.12.2005)

PCT

(10) International Publication Number
WO 2005/117978 A3(51) International Patent Classification:
A61K 39/395 (2006.01) A61P 25/00 (2006.01)

CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2005/019641

(22) International Filing Date: 2 June 2005 (02.06.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/576,993 4 June 2004 (04.06.2004) US

(71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): FROHNA, Paul, A. [US/US]; 3880 19th Street, San Francisco, CA 94114 (US).

(74) Agents: LEE, Wendy, M. et al.; Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080-4990 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
22 June 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR TREATING MULTIPLE SCLEROSIS

Sequence Alignment of Variable Light-Chain Domain

	FR1		CDR1	
	10	20	30	40
2H7	QJYLSQSPRLAGSPQEKVINTC	[RASSSVS-YMM]	WYQOKP	
hu2H7.v15	DIGMTQSPSSLSASVGCRVITC	[RASSSVS-YMM]	WYQOKP	
hum KI	DIGMTQSPSSLSASVGCRVITC	[RASQKISIYCA]	WYQOKP	
	FR2	CDR2	FR3	
	50	60	70	80
2B7	GSSEPKPNIV [APSNLAV]	GVPARPSGSGSGTYSVSLTISVEA		
hu2H7.v16	GRAKPPLIV [APSNLAV]	GVPARPSGSGSGTDFTLTISVSLQP		
hum KI	GRAKPPLIV [AASSLAV]	GVPARPSGSGSGTDFTLTISVSLQP		
	CDR3	FR4		
	90	100		
2H7	EDPAPYYC [QQWSPNPP]	FGAGTAKLRLR		
hu2H7.v16	EDPAPYYC [QQWSPNPP]	FGAGTAKVDIR		
hum KI	EDPAPYYC [QQWSPNPP]	FGAGTAKVDIR		

Sequence Alignment of Variable Heavy-Chain Domain

	FR1		CDR1	
	10	20	30	40
2H7	QAYLQSGAELWVPGASVWMSCNAG	[GYTFTSYNGH]	WYQOKT	
hu2H7.v16	EVQIYEVSGGGLVQPGNSLRLSCAAS	[GYTFCSYNGH]	WYQOKA	
hum III	EVQIYEVSGGGLVQPGNSLRLSCAAS	[GYTFSSYAMS]	WYQQA	
	FR2	CDR2	FR3	
	50	60	70	80
2H7	PRQGLEWIG [AIYPONGOTSYNQPKG]	KATGTVKSSSTAIM		
hu2H7.v16	PRQGLEWIG [AIYPONGOTSYNQPKG]	RPTISVRSKNEYL		
hum III	PRQGLEWIG [VTSQDQGSTVYADQWKG]	RPTISRDNRSKNEYL		
	CDR3	FR4		
	90	100	110	
2H7	abc	100abcde	110	
hu2H7.v16	QISSELTSEDAVYCAR	[VYVYENQSYNFDV]	WQGQMTAVTUS	
hum III	QNSNLRAEDTAVYCAR	[GRVQYSLY--DY]	WQGQMTAVTUS	

A3

WO 2005/117978 A3

(57) Abstract: Methods for treating multiple sclerosis (MS) with a CD20 antibody using special dosing regimens and protocols are described. Articles of manufacture for use in such methods are also described.

METHOD FOR TREATING MULTIPLE SCLEROSISField of the Invention

The present invention concerns methods for treating multiple sclerosis (MS) in a subject
5 using special dosing regimens and protocols, and an article of manufacture with instructions for such
use.

Background of the Invention**Multiple Sclerosis**

10 Multiple Sclerosis (MS) is an inflammatory and demyelinating degenerative disease of the human central nervous system (CNS). It is a worldwide disease that affects approximately 300,000 persons in the United States; it is a disease of young adults, with 70%–80% having onset between 20 and 40 years old (Anderson *et al.* *Ann Neurology* 31(3):333–6 (1992); Noonan *et al.* *Neurology* 58:136–8 (2002)). MS is a heterogeneous disorder based on clinical course, magnetic resonance
15 imaging (MRI) scan assessment, and pathology analysis of biopsy and autopsy material (Lucchinetti *et al.* *Ann Neurol* 47:707–17 (2000)). The disease manifests itself in a large number of possible combinations of deficits, including spinal cord, brainstem, cranial nerve, cerebellar, cerebral, and cognitive syndromes. Progressive disability is the fate of most patients with MS, especially when a 25-year perspective is included. Half of MS patients require a cane to walk within 15 years of disease
20 onset. MS is a major cause of neurologic disability in young and middle-aged adults and, until the past decade, has had no known beneficial treatments. MS is difficult to diagnose because of the non-specific clinical findings, which led to the development of highly structured diagnostic criteria that include several technological advances, consisting of MRI scans, evoked potentials, and cerebrospinal fluid (CSF) studies. All diagnostic criteria rely upon the general principles of scattered
25 lesions in the central white matter occurring at different times and not explained by other etiologies such as infection, vascular disorder, or autoimmune disorder (McDonald *et al.* *Ann Neurol* 50:121–7 (2001)). MS has four patterns of disease: relapsing-remitting MS (RRMS; 80%–85% of cases at onset), primary progressive MS (PPMS; 10%–15% at onset), progressive relapsing MS (PRMS; 5% at onset); and secondary progressive MS (SPMS) (Kremenchutzky *et al.* *Brain* 122 (Pt 10):1941–50
30 (1999); Confavreux *et al.* *N Engl J Med* 343(20):1430–8 (2000)). An estimated 50% of patients with RRMS will develop SPMS in 10 years, and up to 90% of RRMS patients will eventually develop SPMS (Weinshenker *et al.* *Brain* 112(Pt 1):133–46 (1989)).

Currently, six drugs in four classes are approved in the United States for the treatment of RRMS, whereas no drugs have been approved for PPMS. The RRMS treatments include the
35 following: interferon class, IFN-beta-1a (REBIF® and AVONEX®) and IFN-beta-1b (BETASERON®); glatiramer acetate (COPAXONE®), a polypeptide; natalizumab (TYSABRI®); and mitoxantrone (NOVANTRONE®), a cytotoxic agent. Other drugs have been used with varying

degrees of success, including corticosteroids, methotrexate, cyclophosphamide, azathioprine, and intravenous (IV) immunoglobulin. The benefits of currently approved treatments are relatively modest (~30%) for relapse rate and prevention of disability in RRMS as suggested by two recent meta-analyses (Filippini *et al.* *Lancet* 361:545–52 (2003)).

5 Other clinical studies evaluated other immunomodulatory agents in MS, including tumor necrosis factor- α inhibitors and altered peptide ligands, which aggravated rather than improved MS (Lenercept Multiple Sclerosis Study Group and the University of British Columbia MS/MRI *Neurology* 53:457–65 (1999); Bielekova *et al.* *Nat Med* 2000;6:1167–75 (2000), erratum appears in *Nat Med* 6:1412 (2000)).

10 The predominant view of MS pathophysiology has held that inflammation is principally mediated by CD4 $^{+}$ Th1 T cells. Therapeutic approaches based on this theory such as IFN-beta and glatiramer acetate decrease, but do not fully prevent, occurrence of exacerbations or accumulation of disability.

15 The existence of a humoral component in human MS has been implicitly recognized for decades, as evidenced by inclusion of CSF oligoclonal bands and increased intrathecal IgG synthesis in diagnostic criteria for MS (Siden A. *J Neurol* 221:39–51(1979); McDonald *et al.* *Ann Neurol* 50:121–7 (2001); Andersson *et al.* *Eur J Neurol* 9:243–51 (2002); O'Connor, P. *Neurology* 59:S1–33 (2002)). The presence of oligoclonal bands, increased free light chains, and increased intrathecal IgM synthesis correlates with MS disease activity and may be a predictor of more severe outcomes 20 (Rudick *et al.* *Mult Scler* 1:150–5 (1995); Zeman *et al.* *Acta Cytol* 45:51–9 (2001); Izquierdo *et al.* *Acta Neurol Scand* 105:158–63 (2002); Wolinsky J. *J Neurol Sci* 206:145–52 (2003); Villar *et al.* *Ann Neurol* 53:222–6 (2003)).

25 Anti-myelin antibodies (myelin basic protein (MBP) and myelin oligodendrocyte glycoprotein (MOG)) have been detected in the serum of patients with progressive and relapsing forms of MS (Reindl *et al.* *Brain* 122:2047–56 (1999); Egg *et al.* *Mult Scler* 7(5):285–9 (2001)). Anti-myelin antibodies have also been detected in the CSF of MS patients (Reindl *et al.* *Brain* 122:2047–56 (1999); Egg *et al.* *Mult Scler* 7(5):285–9 (2001); Andersson *et al.* *Eur J Neurol* 9:243–51 (2002)). Additional types of antibodies such as anti-ganglioside antibodies or anti-neurofilament antibodies have been observed in patients with MS (Mata *et al.* *Mult Scler* 5:379–88 (1999); Sadatipour *et al.* *Ann Neurol* 44:980–3 (1998)). A recent report indicated that the presence of serum anti-MOG and anti-MBP antibodies was a strong predictor of progression from a clinically isolated demyelinating event to definite RRMS (Berger *et al.* *N Engl J Med* 349:139–45 (2003)). The adjusted hazard ratio for experiencing an exacerbation was 76.5 for patients who were seropositive for both antibodies and 31.6 for patients who were seropositive only for anti-MOG.

35 An international pathology consortium found that antibodies bound to myelin are present in the majority of patients with MS, with plasma cells and B cells also found in MS lesions, providing

additional evidence for a humoral role in MS (Prineas and Wright, *Lab Invest* 38:409-21 (1978); Esiri M. *Neuropathol Appl Neurobiol* 6:9-21 (1980); Genain *et al.* *Nat Med* 5:170-5 (1999); Lucchinetti *et al.* *Ann Neurol* 47:707-17 (2000); Wingerchuk *et al.* *Lab Invest* 81:263-81 (2001)). B cells are detectable in the CSF of patients with MS, and the presence of a relatively high proportion of B cells 5 may be predictive of more severe disability progression (Cepok *et al.* *Brain* 124(Pt 11):2169-76 (2001)).

In subjects with RRMS or opsoclonus-myoclonus syndrome, Rituximab reportedly depleted 10 peripheral B cells in all subjects and decreased the number of CSF B cells in some patients (Pranzatelli *et al.* *Neurology* 60(Suppl1) PO5.128:A395 (2003); Cross *et al.* "Preliminary Results from a Phase II Trial of Rituximab in MS" (abstract) *Eighth Annual Meeting of the Americas Committees for Research and Treatment in Multiple Sclerosis ACTRIMS* 20-1 (October, 2003)). See also Cree *et al.* "Tolerability and Effects of Rituximab "Anti-CD20 Antibody" in Neuromyelitis Optica and Rapidly Worsening Multiple Sclerosis" *Meeting of the Am. Acad. Neurol.* (April, 2004).

CD20 Antibodies and Therapy Therewith

15 Lymphocytes are one of many types of white blood cells produced in the bone marrow during the process of hematopoiesis. There are two major populations of lymphocytes: B lymphocytes (B cells) and T lymphocytes (T cells). The lymphocytes of particular interest herein are B cells.

B cells mature within the bone marrow and leave the marrow expressing an antigen-binding 20 antibody on their cell surface. When a naive B cell first encounters the antigen for which its membrane-bound antibody is specific, the cell begins to divide rapidly and its progeny differentiate into memory B cells and effector cells called "plasma cells". Memory B cells have a longer life span and continue to express membrane-bound antibody with the same specificity as the original parent cell. Plasma cells do not produce membrane-bound antibody but instead produce the antibody in a form that can be secreted. Secreted antibodies are the major effector molecule of humoral immunity.

25 The CD20 antigen (also called human B-lymphocyte-restricted differentiation antigen, Bp35) is a hydrophobic transmembrane protein with a molecular weight of approximately 35 kD located on pre-B and mature B lymphocytes (Valentine *et al.* *J. Biol. Chem.* 264(19):11282-11287 (1989); and Einfeld *et al.* *EMBO J.* 7(3):711-717 (1988)). The antigen is also expressed on greater than 90% of B-cell non-Hodgkin's lymphomas (NHL) (Anderson *et al.* *Blood* 63(6):1424-1433 (1984)), but is not 30 found on hematopoietic stem cells, pro-B cells, normal plasma cells or other normal tissues (Tedder *et al.* *J. Immunol.* 135(2):973-979 (1985)). CD20 regulates an early step(s) in the activation process for cell cycle initiation and differentiation (Tedder *et al.*, *supra*) and possibly functions as a calcium ion channel (Tedder *et al.* *J. Cell. Biochem.* 14D:195 (1990)).

Given the expression of CD20 in B-cell lymphomas, this antigen can serve as a candidate for 35 "targeting" of such lymphomas. In essence, such targeting can be generalized as follows: antibodies specific to the CD20 surface antigen of B cells are administered to a patient. These anti-CD20

antibodies specifically bind to the "CD20" antigen of (ostensibly) both normal and malignant B cells; the antibody bound to the CD20 surface antigen may lead to the destruction and depletion of neoplastic B cells. Additionally, chemical agents or radioactive labels having the potential to destroy the tumor can be conjugated to the anti-CD20 antibody such that the agent is specifically "delivered" 5 to the neoplastic B cells. Irrespective of the approach, a primary goal is to destroy the tumor; the specific approach can be determined by the particular anti-CD20 antibody that is utilized and, thus, the available approaches to targeting the CD20 antigen can vary considerably.

The Rituximab (RITUXAN®) antibody is a genetically engineered chimeric murine/human monoclonal antibody directed against the CD20 antigen. Rituximab is the antibody called "C2B8" in 10 US Patent No. 5,736,137 issued April 7, 1998 (Anderson *et al.*). RITUXAN® is indicated for the treatment of patients with relapsed or refractory low-grade or follicular, CD20-positive, B-cell non-Hodgkin's lymphoma. *In vitro* mechanism of action studies have demonstrated that RITUXAN® binds human complement and lyses lymphoid B-cell lines through complement-dependent cytotoxicity (CDC) (Reff *et al.* *Blood* 83(2):435-445 (1994)). Additionally, it has significant activity 15 in assays for antibody-dependent cellular cytotoxicity (ADCC). More recently, RITUXAN® has been shown to have anti-proliferative effects in tritiated thymidine incorporation assays and to induce apoptosis directly, while other anti-CD19 and CD20 antibodies do not (Maloney *et al.* *Blood* 88(10):637a (1996)). Synergy between RITUXAN® and chemotherapies and toxins has also been observed experimentally. In particular, RITUXAN® sensitizes drug-resistant human B-cell 20 lymphoma cell lines to the cytotoxic effects of doxorubicin, CDDP, VP-16, diphtheria toxin and ricin (Demidem *et al.* *Cancer Chemotherapy & Radiopharmaceuticals* 12(3):177-186 (1997)). *In vivo* preclinical studies have shown that RITUXAN® depletes B cells from the peripheral blood, lymph nodes, and bone marrow of cynomolgus monkeys, presumably through complement and cell-mediated processes (Reff *et al.* *Blood* 83(2):435-445 (1994)).

25 Rituximab was approved in the United States in November 1997 for the treatment of patients with relapsed or refractory low-grade or follicular CD20⁺ B-cell non-Hodgkin's lymphoma (NHL) at a dose of 375 mg/m² weekly for four doses. In April 2001, the Food and Drug Administration (FDA) approved additional claims for the treatment of low-grade NHL: retreatment (weekly for four doses) and an additional dosing regimen (weekly for eight doses). There have been more than 300,000 30 patient exposures to Rituximab either as monotherapy or in combination with immunosuppressant or chemotherapeutic drugs. Patients have also been treated with Rituximab as maintenance therapy for up to 2 years (Hainsworth *et al.* *J Clin Oncol* 21:1746-51 (2003); Hainsworth *et al.* *J Clin Oncol* 20:4261-7 (2002)).

Rituximab has also been studied in a variety of non-malignant autoimmune disorders, in 35 which B cells and autoantibodies appear to play a role in disease pathophysiology (Edwards *et al.* *Biochem Soc Trans* 30:824-8 (2002)). Rituximab has been reported to potentially relieve signs and symptoms of rheumatoid arthritis (RA) (Leandro *et al.* *Ann Rheum Dis.* 61:883-8 (2002); Emery *et*

al. *Arthritis Rheum* 48(9):S439 (2003)), lupus (Eisenberg R. *Arthritis Res Ther* 5:157–9 (2003); Leandro *et al. Arthritis Rheum* 46:2673–7 (2002)), immune thrombocytopenia (D’Arena *et al. Leuk Lymphoma* 44:561–2 (2003)), autoimmune anemia (Zaja *et al. Haematologica* 87:189–95 (2002) (erratum appears in *Haematologica* 87:336 (2002)), autoimmune neuropathy (Pestronk *et al. J Neurol Neurosurg Psychiatry* 74:485–9 (2003)), paraneoplastic opsoclonus-myoclonus syndrome (Pranzatelli *et al. Neurology* 60(Suppl1) PO5.128:A395 (2003)), and relapsing-remitting multiple sclerosis (RRMS) (Cross *et al.* (abstract) Eighth Annual Meeting of the Americas Committees for Research and Treatment in Multiple Sclerosis 20–1 (2003)).

A Phase II study (WA16291) has been conducted in patients with rheumatoid arthritis (RA), providing 48-week follow-up data on safety and efficacy of Rituximab (Emery *et al. Arthritis Rheum* 48(9):S439 (2003); Szczepanski *et al. Arthritis Rheum* 48(9):S121 (2003)). A total of 161 patients were evenly randomized to four treatment arms: methotrexate, Rituximab alone, Rituximab plus methotrexate, Rituximab plus cyclophosphamide (CTX). The treatment regimen of Rituximab was 1 g administered intravenously on Days 1 and 15. Infusions of Rituximab in most patients with RA were well tolerated by most patients, with 36% of patients experiencing at least one adverse event during their first infusion (compared with 30% of patients receiving placebo). Overall, the majority of adverse events were considered to be mild to moderate in severity and were well balanced across all treatment groups. There were a total of 19 serious adverse events across the four arms over the 48 weeks, which were slightly more frequent in the Rituximab/CTX group. The incidence of infections was well balanced across all groups. The mean rate of serious infection in this RA patient population was 4.6 6 per 100 patient-years, which is lower than the rate of infections requiring hospital admission in RA patients (9.57 per 100 patient-years) reported in a community-based epidemiologic study (Doran *et al. Arthritis Rheum* 46:2287–93 (2002)).

The reported safety profile of Rituximab in a small number of patients with neurologic disorders, including autoimmune neuropathy (Pestronk *et al. J Neurol Neurosurg Psychiatry* 74:485–9 (2003)), opsoclonus/myoclonus syndrome (Pranzatelli *et al. Neurology* 60(Suppl1) PO5.128:A395 (2003)), and RRMS (Cross *et al.* Preliminary results from a phase II trial of Rituximab in MS (abstract) Eighth Annual Meeting of the Americas Committees for Research and Treatment in Multiple Sclerosis 20–1 (2003)), was similar to that reported in oncology or RA. In an ongoing investigator-sponsored trial (IST) of Rituximab in combination with interferon-beta (IFN-beta) or glatiramer acetate in subjects with RRMS (Cross *et al., supra*), 1 of 10 treated subjects was admitted to the hospital for overnight observation after experiencing moderate fever and rigors following the first infusion of Rituximab, while the other 9 subjects completed the four-infusion regimen without any reported adverse events.

Patents and patent publications concerning CD20 antibodies include US Patent Nos. 5,776,456, 5,736,137, 5,843,439, 6,399,061, and 6,682,734, as well as US 2002/0197255, US 2003/0021781, US 2003/0082172, US 2003/0095963, US 2003/0147885 (Anderson *et al.*); US Patent

No. 6,455,043, US 2003/0026804, and WO 2000/09160 (Grillo-Lopez, A.); WO 2000/27428 (Grillo-Lopez and White); WO 2000/27433 and US 2004/0213784 (Grillo-Lopez and Leonard); WO 2000/44788 (Braslawsky *et al.*); WO 2001/10462 (Rastetter, W.); WO 2001/10461 (Rastetter and White); WO 2001/10460 (White and Grillo-Lopez); US 2001/0018041, US 2003/0180292, WO 2001/34194 (Hanna and Hariharan); US 2002/0006404 and WO 2002/04021 (Hanna and Hariharan); US 2002/0012665 and WO 2001/74388 (Hanna, N.); US 2002/0058029 (Hanna, N.); US 2003/0103971 (Hariharan and Hanna); US 2002/0009444 and WO 2001/80884 (Grillo-Lopez, A.); WO 2001/97858 (White, C.); US 2002/0128488 and WO 2002/34790 (Reff, M.); WO 2002/060955 (Braslawsky *et al.*); WO 2002/096948 (Braslawsky *et al.*); WO 2002/079255 (Reff and Davies); US Patent No. 6,171,586 and WO 1998/56418 (Lam *et al.*); WO 1998/58964 (Raju, S.); WO 1999/22764 (Raju, S.); WO 1999/51642, US Patent No. 6,194,551, US Patent No. 6,242,195, US Patent No. 6,528,624 and US Patent No. 6,538,124 (Idusogie *et al.*); WO 2000/42072 (Presta, L.); WO 2000/67796 (Curd *et al.*); WO 2001/03734 (Grillo-Lopez *et al.*); US 2002/0004587 and WO 2001/77342 (Miller and Presta); US 2002/0197256 (Grewal, I.); US 2003/0157108 (Presta, L.); WO 04/056312 (Lowman *et al.*); US 2004/0202658 and WO 2004/091657 (Benyunes, K.); WO 2005/000351 (Chan, A.); US 2005/0032130A1 (Beresini *et al.*); US 2005/0053602A1 (Brunetta, P.); US Patent Nos. 6,565,827, 6,090,365, 6,287,537, 6,015,542, 5,843,398, and 5,595,721, (Kaminski *et al.*); US Patent Nos. 5,500,362, 5,677,180, 5,721,108, 6,120,767, and 6,652,852 (Robinson *et al.*); US Pat No. 6,410,391 (Raubitschek *et al.*); US Patent No. 6,224,866 and WO00/20864 (Barbera-Guillem, E.); WO 2001/13945 (Barbera-Guillem, E.); US2005/0079174A1 (Barbera-Guillem *et al.*); WO 2000/67795 (Goldenberg); US 2003/0133930 and WO 2000/74718 (Goldenberg and Hansen); US 2003/0219433 and WO 2003/68821 (Hansen *et al.*); WO2004/058298 (Goldenberg and Hansen); WO 2000/76542 (Golay *et al.*); WO 2001/72333 (Wolin and Rosenblatt); US Patent No. 6,368,596 (Ghetie *et al.*); US Patent No. 6,306,393 and US 2002/0041847 (Goldenberg, D.); US 2003/0026801 (Weiner and Hartmann); WO 2002/102312 (Engleman, E.); US 2003/0068664 (Albitar *et al.*); WO 2003/002607 (Leung, S.); WO 2003/049694, US2002/0009427, and US 2003/0185796 (Wolin *et al.*); WO 2003/061694 (Sing and Siegall); US 2003/0219818 (Bohen *et al.*); US 2003/0219433 and WO 2003/068821 (Hansen *et al.*); US 2003/0219818 (Bohen *et al.*); US2002/0136719 (Shenoy *et al.*); WO 2004/032828 (Wahl *et al.*); and WO 2002/56910 (Hayden-Ledbetter). See also US Patent No. 5,849,898 and EP 330,191 (Seed *et al.*); EP332,865A2 (Meyer and Weiss); US Patent No. 4,861,579 (Meyer *et al.*); US2001/0056066 (Bugelski *et al.*); WO 1995/03770 (Bhat *et al.*); US 2003/0219433 A1 (Hansen *et al.*); WO 2004/035607 (Teeling *et al.*); US 2004/0093621 (Shitara *et al.*); WO 2004/103404 (Watkins *et al.*); WO 2005/000901 (Tedder *et al.*); US 2005/0025764 (Watkins *et al.*); WO2005/016969 and US 2005/0069545 A1 (Carr *et al.*); WO 2005/014618 (Chang *et al.*). Certain of these include, *inter alia*, treatment of multiple sclerosis.

Publications concerning therapy with Rituximab include: Perotta and Abuel “Response of chronic relapsing ITP of 10 years duration to Rituximab” Abstract # 3360 *Blood* 10(1)(part 1-2): p.

88B (1998); Stashi *et al.* "Rituximab chimeric anti-CD20 monoclonal antibody treatment for adults with chronic idiopathic thrombocytopenic purpura" *Blood* 98(4):952-957 (2001); Matthews, R. "Medical Heretics" *New Scientist* (7 April, 2001); Leandro *et al.* "Clinical outcome in 22 patients with rheumatoid arthritis treated with B lymphocyte depletion" *Ann Rheum Dis* 61:833-888 (2002); 5 Leandro *et al.* "Lymphocyte depletion in rheumatoid arthritis: early evidence for safety, efficacy and dose response. *Arthritis and Rheumatism* 44(9): S370 (2001); Leandro *et al.* "An open study of B lymphocyte depletion in systemic lupus erythematosus", *Arthritis & Rheumatism* 46(1):2673-2677 (2002); Edwards and Cambridge "Sustained improvement in rheumatoid arthritis following a protocol designed to deplete B lymphocytes" *Rheumatology* 40:205-211 (2001); Edwards *et al.* "B-lymphocyte depletion therapy in rheumatoid arthritis and other autoimmune disorders" *Biochem. Soc. Trans.* 30(4):824-828 (2002); Edwards *et al.* "Efficacy and safety of Rituximab, a B-cell targeted chimeric monoclonal antibody: A randomized, placebo controlled trial in patients with rheumatoid arthritis. *Arthritis and Rheumatism* 46(9): S197 (2002); Levine and Pestronk "IgM antibody-related polyneuropathies: B-cell depletion chemotherapy using Rituximab" *Neurology* 52: 1701-1704 (1999); 10 DeVita *et al.* "Efficacy of selective B cell blockade in the treatment of rheumatoid arthritis" *Arthritis & Rheum* 46:2029-2033 (2002); Hidashida *et al.* "Treatment of DMARD-Refractory rheumatoid arthritis with Rituximab." Presented at the *Annual Scientific Meeting of the American College of Rheumatology*; Oct 24-29; New Orleans, LA 2002; Tuscano, J. "Successful treatment of Infliximab-refractory rheumatoid arthritis with Rituximab" Presented at the *Annual Scientific Meeting of the American College of Rheumatology*; Oct 24-29; New Orleans, LA 2002; Specks *et al.* "Response of Wegener's granulomatosis to anti-CD20 chimeric monoclonal antibody therapy" *Arthritis & Rheumatism* 44(12):2836-2840 (2001); Anolik *et al.*, "B lymphocyte Depletion in the Treatment of Systemic Lupus (SLE): Phase I/II Trial of Rituximab (RITUXAN®) in SLE" *Arthritis And Rheumatism*, 46(9), S289-S289 Abstract 717 (October, 2002), and Albert *et al.*, "A Phase I Trial of 15 Rituximab (Anti-CD20) for Treatment of Systemic Lupus Erythematosus" *Arthritis And Rheumatism*, 48(12): 3659-3659, Abstract LB9 (December, 2003); Martin and Chan "Pathogenic Roles of B cells in Human Autoimmunity: Insights from the Clinic" *Immunity* 20:517-527 (2004). 20

All references, including any patents or patent applications, cited in this specification are hereby incorporated by reference. No admission is made that any reference constitutes prior art. The 25 discussion of the references states what their authors assert, and the applicants reserve the right to challenge the accuracy and pertinency of the cited documents. It will be clearly understood that, although a number of prior art publications are referred to herein, this reference does not constitute an admission that any of these documents forms part of the common general knowledge in the art, in Australia or in any other country.

Summary of the Invention

The present invention involves, at least in part, the selection of a dose for a CD20 antibody that provides a safe and active treatment regimen in subjects with multiple sclerosis, such as PPMS or RRMS.

5 Accordingly, a first aspect of the invention concerns a method of treating multiple sclerosis in a subject comprising administering an effective amount of a CD20 antibody to the subject to provide an initial antibody exposure of about 0.5 to 4 grams followed by a second antibody exposure of about 0.5 to 4 grams, the second exposure not being provided until from about 16 to 60 weeks from the initial exposure, and each of the antibody exposures is provided to the subject as one or two doses of antibody.

10 A second aspect provides an article of manufacture when used according to the method of the first aspect comprising a container comprising a CD20 antibody.

15 A third aspect provides use of a CD20 antibody in the manufacture of a medicament for treating multiple sclerosis in a subject, wherein the medicament is administered to the subject to provide an initial antibody exposure of about 0.5 to 4 grams followed by a second antibody exposure of about 0.5 to 4 grams, the second exposure not being provided until from about 16 to 60 weeks from the initial exposure, and each of the antibody exposures is provided to the subject as one or two doses of antibody.

20 In addition, disclosed herein is an article of manufacture comprising: (a) a container comprising a CD20 antibody; and (b) a package insert with instructions for treating multiple sclerosis in a subject, wherein the instructions indicate that an amount of the antibody is administered to the subject that is effective to provide an initial antibody exposure of about 0.5 to 4 grams followed by a second antibody exposure of about 0.5 to 4 grams, the second exposure not being administered until from about 16 to 60 weeks from the initial exposure, and each of the antibody exposures is provided to the subject as one or two doses of antibody.

Brief Description of the Drawings

25 FIG. 1A is a sequence alignment comparing the amino acid sequences of the light chain variable domain (V_L) of each of murine 2H7 (SEQ ID NO:1), humanized 2H7.v16 variant (SEQ ID NO:2), and the human kappa light chain subgroup I (SEQ ID NO:3). The CDRs of V_L of 2H7 and hu2H7.v16 are as follows: CDR1 (SEQ ID NO:4), CDR2 (SEQ ID NO:5), and CDR3 (SEQ ID NO:6).

30 FIG. 1B is a sequence alignment comparing the amino acid sequences of the heavy chain variable domain (V_H) of each of murine 2H7 (SEQ ID NO:7), humanized 2H7.v16 variant (SEQ ID NO:8), and the human consensus sequence of the heavy chain subgroup III (SEQ ID NO:9). The

CDRs of V_H of 2H7 and hu2H7.v16 are as follows: CDR1 (SEQ ID NO:10), CDR2 (SEQ ID NO:11), and CDR3 (SEQ ID NO:12).

In FIG. 1A and FIG. 1B, the CDR1, CDR2 and CDR3 in each chain are enclosed within brackets, flanked by the framework regions, FR1-FR4, as indicated. 2H7 refers to the murine 2H7 antibody. The asterisks in between two rows of sequences indicate the positions that are different between the two sequences. Residue numbering is according to Kabat *et al. Sequences of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991), with insertions shown as a, b, c, d, and e.

FIG. 2 shows the amino acid sequence of the mature 2H7.v16 light chain (SEQ ID NO:13)

FIG. 3 shows the amino acid sequence of the mature 2H7.v16 heavy chain (SEQ ID NO:14).

FIG. 4 shows the amino acid sequence of the mature 2H7.v31 heavy chain (SEQ ID NO:15).

The L chain of 2H7.v31 is the same as for 2H7.v16.

FIG. 5 shows an alignment of the mature 2H7.v16 and 2H7.v511 light chains (SEQ ID NOS.

13 and 16, respectively), with Kabat variable domain residue numbering and Eu constant domain

residue numbering.

FIG. 6 shows an alignment of the mature 2H7.v16 and 2H7.v511 heavy chains (SEQ ID NOS. 14 and 17, respectively), with Kabat variable domain residue numbering and Eu constant domain residue numbering.

5

Detailed Description of the Preferred Embodiments

I. Definitions

A "B-cell" is a lymphocyte that matures within the bone marrow, and includes a naive B cell, memory B cell, or effector B cell (plasma cells). The B-cell herein may be a normal or non-malignant B cell.

10 A "B-cell surface marker" or "B-cell surface antigen" herein is an antigen expressed on the surface of a B cell that can be targeted with an antibody that binds thereto. Exemplary B-cell surface markers include the CD10, CD19, CD20, CD21, CD22, CD23, CD24, CD37, CD40, CD53, CD72, CD73, CD74, CDw75, CDw76, CD77, CDw78, CD79a, CD79b, CD80, CD81, CD82, CD83, CDw84, CD85 and CD86 leukocyte surface markers (for descriptions, see *The Leukocyte Antigen Facts Book*, 2nd Edition. 1997, ed. Barclay *et al.* Academic Press, Harcourt Brace & Co., New York). Other B-cell surface markers include RP105, FcRH2, B-cell CR2, CCR6, P2X5, HLA-DOB, CXCR5, FCER2, BR3, Btig, NAG14, SLGC16270, FcRH1, IRTA2, ATWD578, FcRH3, IRTA1, FcRH6, BCMA, and 239287. The B-cell surface marker of particular interest herein is preferentially expressed on B cells compared to other non-B-cell tissues of a mammal and may be expressed on 15 both precursor B cells and mature B cells. The preferred B-cell surface marker herein is CD20.

20 The "CD20" antigen, or "CD20," is an about 35-kDa, non-glycosylated phosphoprotein found on the surface of greater than 90% of B cells from peripheral blood or lymphoid organs. CD20 is present on both normal B cells as well as malignant B cells, but is not expressed on stem cells. Other names for CD20 in the literature include "B-lymphocyte-restricted antigen" and "Bp35". The 25 CD20 antigen is described in Clark *et al.* *Proc. Natl. Acad. Sci. (USA)* 82:1766 (1985), for example.

20 An "antibody antagonist" herein is a antibody that, upon binding to a B cell surface marker on B cells, destroys or depletes B cells in a mammal and/or interferes with one or more B-cell functions, *e.g.* by reducing or preventing a humoral response elicited by the B cell. The antibody antagonist preferably is able to deplete B cells (*i.e.* reduce circulating B-cell levels) in a mammal treated therewith. Such depletion may be achieved via various mechanisms such antibody-dependent cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC), inhibition of B-cell proliferation and/or induction of B-cell death (*e.g.* via apoptosis).

35 "Antibody-dependent cell-mediated cytotoxicity" and "ADCC" refer to a cell-mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FcRs) (*e.g.* Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. The primary cells for mediating ADCC, NK cells, express Fc γ RIII only, whereas monocytes express Fc γ RI, Fc γ RII and Fc γ RIII. FcR expression on hematopoietic cells in

summarized is Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as that described in US Patent No. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or 5 additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, e.g., in a animal model such as that disclosed in Clynes *et al.* *PNAS (USA)* 95:652-656 (1998).

“Human effector cells” are leukocytes that express one or more FcRs and perform effector functions. Preferably, the cells express at least Fc γ RIII and carry out ADCC effector function. Examples of human leukocytes that mediate ADCC include peripheral blood mononuclear cells 10 (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred.

The terms “Fc receptor” or “FcR” are used to describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one that binds an IgG antibody (a gamma receptor) and includes receptors of the Fc γ RI, Fc γ RII, and 15 Fc γ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors. Fc γ RII receptors include Fc γ RIIA (an “activating receptor”) and Fc γ RIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor Fc γ RIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in 20 its cytoplasmic domain. Inhibiting receptor Fc γ RIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (see Daëron, *Annu. Rev. Immunol.* 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-92 (1991); Capel *et al.*, *Immunomethods* 4:25-34 (1994); and de Haas *et al.*, *J. Lab. Clin. Med.* 126:330-41 (1995). Other 25 FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer *et al.*, *J. Immunol.* 117:587 (1976) and Kim *et al.*, *J. Immunol.* 24:249 (1994)).

“Complement dependent cytotoxicity” or “CDC” refer to the ability of a molecule to lyse a target in the presence of complement. The complement activation pathway is initiated by the binding of the first component of the complement system (C1q) to a molecule (e.g. an antibody) complexed with a cognate antigen. To assess complement activation, a CDC assay, e.g. as described in Gazzano- 30 Santoro *et al.*, *J. Immunol. Methods* 202:163 (1996), may be performed.

“Growth inhibitory” antibodies are those that prevent or reduce proliferation of a cell expressing an antigen to which the antibody binds. For example, the antibody may prevent or reduce proliferation of B cells *in vitro* and/or *in vivo*.

Antibodies that “induce apoptosis” are those that induce programmed cell death, e.g. of a B 35 cell, as determined by standard apoptosis assays, such as binding of annexin V, fragmentation of DNA, cell shrinkage, dilation of endoplasmic reticulum, cell fragmentation, and/or formation of membrane vesicles (called apoptotic bodies).

The term "antibody" hereinafter is used in the broadest sense and specifically covers monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g. bispecific antibodies) formed from at least two intact antibodies, and antibody fragments so long as they exhibit the desired biological activity.

5 "Antibody fragments" comprise a portion of an intact antibody, preferably comprising the antigen binding region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

10 For the purposes herein, an "intact antibody" is one comprising heavy and light variable domains as well as an Fc region.

15 "Native antibodies" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface 20 between the light chain and heavy chain variable domains.

25 The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called hypervariable regions both in the light chain and the heavy chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FRs). The variable domains of native heavy and light chains each comprise four FRs, largely adopting a β -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the β -sheet structure. The hypervariable regions in each chain are held together 30 in close proximity by the FRs and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody dependent cellular 35 cytotoxicity (ADCC).

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects

its ability to crystallize readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen-binding sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment that contains a complete antigen-recognition and antigen-binding site. This region consists of a dimer of one heavy chain and one light chain variable domain in tight, non-covalent association. It is in this configuration that the three hypervariable regions of each variable domain interact to define an antigen-binding site on the surface of the V_H - V_L dimer. Collectively, the six hypervariable regions confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three hypervariable regions specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear at least one free thiol group. $F(ab')_2$ antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, antibodies can be assigned to different classes. There are five major classes of intact antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy chain constant domains that correspond to the different classes of antibodies are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

"Single-chain Fv" or "scFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains that enables the scFv to form the desired structure for antigen binding. For a review of scFv see Plückthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenburg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy chain variable domain (V_H) connected to a light chain variable domain (V_L) in the same polypeptide chain ($V_H - V_L$). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are

described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical and/or bind the same epitope, except for possible variants that may arise during production of the monoclonal antibody, such variants generally being present in minor amounts. In contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, *e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991), for example.

The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; Morrison *et al.*, *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)). Chimeric antibodies of interest herein include "primatized" antibodies comprising variable domain antigen-binding sequences derived from a non-human primate (*e.g.* Old World Monkey, such as baboon, rhesus or cynomolgus monkey) and human constant region sequences (US Pat No. 5,693,780).

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor

5 antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence, except for FR substitution(s) as noted above. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region, typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992).

10 The term "hypervariable region" when used herein refers to the amino acid residues of an antibody that are responsible for antigen binding. The hypervariable region comprises amino acid residues from a "complementarity determining region" or "CDR" (e.g. residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues 15 from a "hypervariable loop" (e.g. residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). "Framework" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined.

20 A "naked antibody" is an antibody (as herein defined) that is not conjugated to a heterologous molecule, such as a cytotoxic moiety or radiolabel.

25 Examples of antibodies that bind the CD20 antigen include: "C2B8," which is now called "Rituximab" ("RITUXAN®") (US Patent No. 5,736,137); the yttrium-[90]-labeled 2B8 murine antibody designated "Y2B8" or "Ibritumomab Tiuxetan" (ZEVALIN®) commercially available from IDEC Pharmaceuticals, Inc. (US Patent No. 5,736,137; 2B8 deposited with ATCC under accession no. HB11388 on June 22, 1993); murine IgG2a "B1," also called "Tositumomab," optionally labeled with ¹³¹I to generate the "¹³¹I-B1" or "iodine I131 tositumomab" antibody (BEXXAR™) 30 commercially available from Corixa (see, also, US Patent No. 5,595,721); murine monoclonal antibody "1F5" (Press *et al.* *Blood* 69(2):584-591 (1987) and variants thereof including "framework patched" or humanized 1F5 (WO03/002607, Leung, S.; ATCC deposit HB-96450); murine 2H7 and chimeric 2H7 antibody (US Patent No. 5,677,180); humanized 2H7; huMax-CD20 (Genmab, Denmark, WO2004/035607); AME-133 (Applied Molecular Evolution); A20 antibody or variants thereof such as chimeric or humanized A20 antibody (cA20, hA20, respectively) or IMMU-106 (US 2003/0219433, Immunomedics); CD20-binding antibodies, including epitope-depleted Leu-16, 1H4, or 2B8, optionally conjugated with IL-2, as in US 2005/0069545A1 and WO 2005/16969 (Carr *et al.*); bispecific antibody that binds CD22 and CD20, for example, hLL2xhA20 (WO2005/14618, Chang *et al.*); monoclonal antibodies L27, G28-2, 93-1B3, B-C1 or NU-B2 available from the International Leukocyte Typing Workshop (Valentine *et al.*, In: *Leukocyte Typing III* (McMichael,

Ed., p. 440, Oxford University Press (1987)); 1H4 (Haisma *et al.* *Blood* 92:184 (1998)); anti-CD20 auristatin E conjugate (Seattle Genetics); anti-CD20-IL2 (EMD/Biovation/City of Hope); anti-CD20 MAb therapy (EpiCyte); and anti-CD20 antibody TRU 015 (Trubion).

The terms "Rituximab" or "RITUXAN®" herein refer to the genetically engineered chimeric murine/human monoclonal antibody directed against the CD20 antigen and designated "C2B8" in US Patent No. 5,736,137, including fragments thereof that retain the ability to bind CD20. Rituximab is commercially available from Genentech.

Purely for the purposes herein and unless indicated otherwise, "humanized 2H7" refers to a humanized antibody that binds human CD20, or an antigen-binding fragment thereof, wherein the antibody is effective to deplete primate B cells *in vivo*, the antibody comprising in the H chain variable region (V_H) thereof at least a CDR H3 sequence of SEQ ID NO:12 (Fig. 1B) from an anti-human CD20 antibody and substantially the human consensus framework (FR) residues of the human heavy- chain subgroup III (V_HIII). In a preferred embodiment, this antibody further comprises the H chain CDR H1 sequence of SEQ ID NO:10 and CDR H2 sequence of SEQ ID NO:11, and more preferably further comprises the L chain CDR L1 sequence of SEQ ID NO:4, CDR L2 sequence of SEQ ID NO:5, CDR L3 sequence of SEQ ID NO:6 and substantially the human consensus framework (FR) residues of the human light chain subgroup I (VI), wherein the V_H region may be joined to a human IgG chain constant region, wherein the region may be, for example, IgG1 or IgG3. In a preferred embodiment, such antibody comprises the V_H sequence of SEQ ID NO:8 (v16, as shown in Fig. 1B), optionally also comprising the V_L sequence of SEQ ID NO:2 (v16, as shown in Fig. 1A), which may have the amino acid substitutions of D56A and N100A in the H chain and S92A in the L chain (v96). Preferably the antibody is an intact antibody comprising the light and heavy chain amino acid sequences of SEQ ID NOS:13 and 14, respectively, as shown in Figs. 2 and 3. Another preferred embodiment is where the antibody is 2H7.v31 comprising the light and heavy chain amino acid sequences of SEQ ID NOS:13 and 15, respectively, as shown in Figs. 2 and 4. The antibody herein may further comprise at least one amino acid substitution in the Fc region that improves ADCC and/or CDC activity, such as one wherein the amino acid substitutions are S298A/E333A/K334A, more preferably 2H7.v31 having the heavy chain amino acid sequence of SEQ ID NO:15 (as shown in Fig. 4). Any of these antibodies may further comprise at least one amino acid substitution in the Fc region that decreases CDC activity, for example, comprising at least the substitution K322A. See U.S. Patent No. 6,528,624B1 (Idusogie *et al.*).

A preferred humanized 2H7 is an intact antibody or antibody fragment comprising the variable light chain sequence:

DIQMTQSPSSLASAVGDRVTITCRASSSVSYMHWYQQKPGKAPKPLIYAPSNLASGVPSRFSG
35 SGSGTDFTLTISSLQPEDFATYYCQQWSFNPPTFGQGTKVEIKR (SEQ ID NO:2);

and the variable heavy chain sequence:

EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPGNGDTSY

NQKFKGRFTISVDKSKNTLYLQMNSLRAEDTAVYYCARVVYYSNSYWYFDVWGQGTLVTV
SS (SEQ ID NO:8).

Where the humanized 2H7 antibody is an intact antibody, preferably it comprises the light chain amino acid sequence:

5 DIQMTQSPSSLSASVGDRVTITCRASSSVSYMHWYQQKPGKAPKPLIYAPSNLASGVPSRFSG
SGSGTDFTLTISSLQPEDFATYYCQQWSFNPPTFQGQTKVEIKRTVAAPSVFIFPPSDEQLKSG
TASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSSTLTLKADYEKH
KVYACEVTHQGLSSPVTKSFRGEC (SEQ ID NO:13);

and the heavy chain amino acid sequence:

10 EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPNGDTSY
NQKFKGRFTISVDKSKNTLYLQMNSLRAEDTAVYYCARVVYYSNSYWYFDVWGQGTLVTV
SSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVWSNSGALTSGVHTFPAVLQSSG
LYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVF
LFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV
15 VSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTIASKAKGQPREPQVYTLPPSREEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCS
VMHEALHNHYTQKSLSLSPGK (SEQ ID NO:14)

or the heavy chain amino acid sequence:

EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPNGDTSY
20 NQKFKGRFTISVDKSKNTLYLQMNSLRAEDTAVYYCARVVYYSNSYWYFDVWGQGTLVTV
SSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVWSNSGALTSGVHTFPAVLQSSG
LYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVF
LFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNATYRV
VSVLTVLHQDWLNGKEYKCKVSNKALPAPIATISKAKGQPREPQVYTLPPSREEMTKNQVS
25 LTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCS
VMHEALHNHYTQKSLSLSPGK (SEQ ID NO:15).

In the preferred embodiment of the invention, the variable region of variants based on 2H7 version 16 will have the amino acid sequences of v16 except at the positions of amino acid substitutions that are indicated in the table below. Unless otherwise indicated, the 2H7 variants will have the same light chain as that of v16.

2H7 Version	Heavy chain (V _H) changes	Light chain (V _L) changes	Fc changes
31			S298A, E333A, K334A
96	D56A, N100A	S92A	
114	D56A, N10	M32L, S92A	S298A, E333A, K334A
115	D56A, N100A	M32L, S92A	S298A, E333A, K334A, E356D, M358L

An "isolated" antibody is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the antibody, and may include 5 enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing 10 conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

A "subject" herein is a human subject. Generally, the subject is eligible for treatment for 15 multiple sclerosis. For the purposes herein, such eligible subject is one who is experiencing, has experienced, or is likely to experience, one or more signs, symptoms or other indicators of multiple sclerosis; has been diagnosed with multiple sclerosis, whether, for example, newly diagnosed (with "new onset" MS), previously diagnosed with a new relapse or exacerbation, previously diagnosed and in remission, etc; and/or is at risk for developing multiple sclerosis. One suffering from or at risk for 20 suffering from multiple sclerosis may optionally be identified as one who has been screened for elevated levels of CD20-positive B cells in serum, cerebrospinal fluid (CSF) and/or MS lesion(s) and/or is screened for using an assay to detect autoantibodies, assessed qualitatively, and preferably quantitatively. Exemplary such autoantibodies associated with multiple sclerosis include anti-myelin basic protein (MBP), anti-myelin oligodendrocytic glycoprotein (MOG), anti-ganglioside and/or 25 anti-neurofilament antibodies. Such autoantibodies may be detected in the subject's serum, cerebrospinal fluid (CSF) and/or MS lesion. By "elevated" autoantibody or B cell level(s) herein is meant level(s) of such autoantibodies or B cells which significantly exceed the level(s) in an individual without MS.

"Treatment" of a subject herein refers to both therapeutic treatment and prophylactic or 30 preventative measures. Those in need of treatment include those already with the multiple sclerosis as well as those in which the multiple sclerosis is to be prevented. Hence, the subject may have been diagnosed as having the multiple sclerosis or may be predisposed or susceptible to the multiple sclerosis.

A "symptom" of MS is any morbid phenomenon or departure from the normal in structure, 35 function, or sensation, experienced by the subject and indicative of MS.

"Multiple sclerosis" refers to the chronic and often disabling disease of the central nervous system characterized by the progressive destruction of the myelin. There are four internationally

recognized forms of MS, namely, primary progressive multiple sclerosis (PPMS), relapsing-remitting multiple sclerosis (RRMS), secondary progressive multiple sclerosis (SPMS), and progressive relapsing multiple sclerosis (PRMS).

“Primary progressive multiple sclerosis” or “PPMS” is characterized by a gradual progression of the disease from its onset with no superimposed relapses and remissions at all. There may be periods of a leveling off of disease activity and there may be good and bad days or weeks. PPMS differs from RRMS and SPMS in that onset is typically in the late thirties or early forties, men are as likely women to develop it, and initial disease activity is often in the spinal cord and not in the brain. PPMS often migrates into the brain, but is less likely to damage brain areas than RRMS or SPMS; for example, people with PPMS are less likely to develop cognitive problems. PPMS is the sub-type of MS that is least likely to show inflammatory (gadolinium enhancing) lesions on MRI scans. The Primary Progressive form of the disease affects between 10 and 15% of all people with multiple sclerosis. PPMS may be defined according to the criteria in McDonald *et al. Ann Neurol* 50:121–7 (2001). The subject with PPMS treated herein is usually one with probable or definitive diagnosis of PPMS.

“Relapsing-remitting multiple sclerosis” or “RRMS” is characterized by relapses (also known as exacerbations) during which time new symptoms can appear and old ones resurface or worsen. The relapses are followed by periods of remission, during which time the person fully or partially recovers from the deficits acquired during the relapse. Relapses can last for days, weeks or months and recovery can be slow and gradual or almost instantaneous. The vast majority of people presenting with MS are first diagnosed with RRMS. This is typically when they are in their twenties or thirties, though diagnoses much earlier or later are known. Twice as many women as men present with this sub-type of MS. During relapses, myelin, a protective insulating sheath around the nerve fibers (neurons) in the white matter regions of the central nervous system (CNS), may be damaged in an inflammatory response by the body's own immune system. This causes a wide variety of neurological symptoms that vary considerably depending on which areas of the CNS are damaged. Immediately after a relapse, the inflammatory response dies down and a special type of glial cell in the CNS (called an oligodendrocyte) sponsors remyelination - a process whereby the myelin sheath around the axon may be repaired. It is this remyelination that may be responsible for the remission. Approximately 50% of patients with RRMS convert to SPMS within 10 years of disease onset. After 30 years, this figure rises to 90%. At any one time, the relapsing-remitting form of the disease accounts around 55% of all people with MS.

“Secondary progressive multiple sclerosis” or “SPMS” is characterized by a steady progression of clinical neurological damage with or without superimposed relapses and minor remissions and plateaux. People who develop SPMS will have previously experienced a period of RRMS which may have lasted anything from two to forty years or more. Any superimposed relapses and remissions there are, tend to tail off over time. From the onset of the secondary progressive phase

of the disease, disability starts advancing much quicker than it did during RRMS though the progress can still be quite slow in some individuals. After 10 years, 50% of people with RRMS will have developed SPMS. By 25 to 30 years, that figure will have risen to 90%. SPMS tends to be associated with lower levels of inflammatory lesion formation than in RRMS but the total burden of disease 5 continues to progress. At any one time, SPMS accounts around 30% of all people with multiple sclerosis.

“Progressive relapsing multiple sclerosis” refers to “PRMS” is characterized by a steady progression of clinical neurological damage with superimposed relapses and remissions. There is significant recovery immediately following a relapse but between relapses there is a gradual 10 worsening of symptoms. PRMS affects around 5% of all people with multiple sclerosis. Some neurologists believe PRMS is a variant of PPMS.

The expression “effective amount” refers to an amount of the antibody (or other drug) that is effective for preventing, ameliorating or treating the multiple sclerosis. Such an effective amount will generally result in an improvement in the signs, symptoms or other indicators of MS, such as reducing 15 relapse rate, preventing disability, reducing number and/or volume of brain MRI lesions, improving timed 25-foot walk, extending the time to disease progression (*e.g.* using Expanded Disability Status Scale, EDSS), etc.

“Antibody exposure” refers to contact with or exposure to the antibody herein in one or more doses administered over a period of time of about 1-20 days. The doses may be given at one time or 20 at fixed or irregular time intervals over this period of exposure. Initial and later (*e.g.* second or third) antibody exposures are separated in time from each other as described in detail herein.

The term “immunosuppressive agent” as used herein for adjunct therapy refers to substances that act to suppress or mask the immune system of the mammal being treated herein. This would include substances that suppress cytokine production, down-regulate or suppress self-antigen 25 expression, or mask the MHC antigens. Examples of such agents include 2-amino-6-aryl-5- substituted pyrimidines (see U.S. Pat. No. 4,665,077); nonsteroidal anti-inflammatory drugs (NSAIDs); ganciclovir, tacrolimus, glucocorticoids such as cortisol or aldosterone, anti-inflammatory agents such as a cyclooxygenase inhibitor, a 5-lipoxygenase inhibitor, or a leukotriene receptor antagonist; purine antagonists such as azathioprine or mycophenolate mofetil (MMF); alkylating 30 agents such as cyclophosphamide; bromocryptine; danazol; dapsone; glutaraldehyde (which masks the MHC antigens, as described in U.S. Pat. No. 4,120,649); anti-idiotypic antibodies for MHC antigens and MHC fragments; cyclosporin A; steroids such as corticosteroids or glucocorticosteroids or glucocorticoid analogs, *e.g.*, prednisone, methylprednisolone, and dexamethasone; dihydrofolate reductase inhibitors such as methotrexate (oral or subcutaneous); hydroxychloroquine; sulfasalazine; 35 leflunomide; cytokine or cytokine receptor antagonists including anti-interferon-alpha, -beta, or -gamma antibodies, anti-tumor necrosis factor-alpha antibodies (infliximab or adalimumab), anti-TNF-alpha immunoahesin (etanercept), anti-tumor necrosis factor-beta antibodies, anti-interleukin-2

antibodies and anti-IL-2 receptor antibodies; anti-LFA-1 antibodies, including anti-CD11a and anti-CD18 antibodies; anti-L3T4 antibodies; heterologous anti-lymphocyte globulin; pan-T antibodies, preferably anti-CD3 or anti-CD4/CD4a antibodies; soluble peptide containing a LFA-3 binding domain (WO 90/08187 published 7/26/90); streptokinase; TGF-beta; streptodornase; RNA or DNA from the host; FK506; RS-61443; deoxyspergualin; rapamycin; T-cell receptor (Cohen *et al.*, U.S. Pat. No. 5,114,721); T-cell receptor fragments (Offner *et al.*, *Science*, 251: 430-432 (1991); WO 90/11294; Ianeway, *Nature*, 341: 482 (1989); and WO 91/01133); and T cell receptor antibodies (EP 340,109) such as T10B9.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g. At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³² and radioactive isotopes of Lu), chemotherapeutic agents, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, or fragments thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylololmelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, chlophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e. g., calicheamicin, especially calicheamicin gamma1I and calicheamicin omega1I (see, e.g., Agnew, *Chem Intl. Ed. Engl.*, 33: 183-186 (1994)); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabicin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN® doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as

methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, 5 dromostanolone propionate, epitostanol, mepitiostane, testolactone; anti- adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as folinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestramycin; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; el fornithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and 10 ansamitocins; mitoguazone; mitoxantrone; moperanmol; niraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2- ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, OR); razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; trichothecenes (especially T-2 toxin, 15 verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g., TAXOL® paclitaxel (Bristol- Myers Squibb Oncology, Princeton, N.J.), ABRAXANE™ Cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumberg, Illinois), and TAXOTERE® doxetaxel (Rhône- Poulenc Rorer, Antony, France); chlorambucil; GEMZAR® gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum 20 analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; NAVELBINE® vinorelbine; novantrone; teniposide; edatraxate; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

25 Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX® tamoxifen), raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON toremifene; aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen 30 production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, MEGASE® megestrol acetate, AROMASIN® exemestane, formestane, fadrozole, RIVISOR® vorozole, FEMARA® letrozole, and ARIMIDEX® anastrozole; and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine analog); antisense oligonucleotides, particularly those that inhibit 35 expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Ral and H-Ras; vaccines such as gene therapy vaccines, for example, ALLOVECTIN® vaccine, LEUVECTIN® vaccine, and VAXID® vaccine; PROLEUKIN® rIL-2;

LURTOTECAN® topoisomerase I inhibitor; ABARELIX® rmRH; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

The term "cytokine" is a generic term for proteins released by one cell population that act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines; interleukins (ILs) such as IL-1, IL-1 α , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-11, IL-12, IL-15; a tumor necrosis factor such as TNF- α or TNF- β ; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines, including synthetically produced small-molecule entities and pharmaceutically acceptable derivatives and salts thereof.

The term "hormone" refers to polypeptide hormones, which are generally secreted by glandular organs with ducts. Included among the hormones are, for example, growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); prolactin, placental lactogen, mouse gonadotropin-associated peptide, inhibin; activin; mullerian-inhibiting substance; and thrombopoietin. As used herein, the term hormone includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence hormone, including synthetically produced small-molecule entities and pharmaceutically acceptable derivatives and salts thereof.

The term "growth factor" refers to proteins that promote growth, and include, for example, hepatic growth factor; fibroblast growth factor; vascular endothelial growth factor; nerve growth factors such as NGF- β ; platelet-derived growth factor; transforming growth factors (TGFs) such as TGF- α and TGF- β ; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon- α , - β , and - γ ; and colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF). As used herein, the term growth factor includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence growth factor, including synthetically produced small-molecule entities and pharmaceutically acceptable derivatives and salts thereof.

The term "integrin" refers to a receptor protein that allows cells both to bind to and to respond to the extracellular matrix and is involved in a variety of cellular functions such as wound healing, cell differentiation, homing of tumor cells and apoptosis. They are part of a large family of cell adhesion receptors that are involved in cell-extracellular matrix and cell-cell interactions. Functional integrins consist of two transmembrane glycoprotein subunits, called alpha and beta, that are non-covalently bound. The alpha subunits all share some homology to each other, as do the beta subunits. The receptors always contain one alpha chain and one beta chain. Examples include Alpha6beta1,

Alpha3beta1, Alpha7beta1, LFA-1, alpha 4 integrin etc. As used herein, the term integrin includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence integrin, including synthetically produced small-molecule entities and pharmaceutically acceptable derivatives and salts thereof.

5 Examples of “integrin antagonists or antibodies” herein include an LFA-1 antibody such as Efalizumab (RAPTIVA®) commercially available from Genentech; an alpha 4 integrin antibody such as natalizumab (TYSABRI®) available from Biogen; diazacyclic phenylalanine derivatives (WO 2003/89410); phenylalanine derivatives (WO 2003/70709, WO 2002/28830, WO 2002/16329 and WO 2003/53926); phenylpropionic acid derivatives (WO 2003/10135); enamine derivatives (WO 10 2001/79173); propanoic acid derivatives (WO 2000/37444); alkanoic acid derivatives (WO 2000/32575); substituted phenyl derivatives (US Pat. Nos. 6,677,339 and 6,348,463); aromatic amine derivatives (US Pat. No. 6,369,229); and ADAM disintegrin domain polypeptide (US2002/0042368), antibodies to alphavbeta3 integrin (EP 633945); aza-bridged bicyclic amino acid derivatives (WO 2002/02556) etc.

15 For the purposes herein, “tumor necrosis factor alpha (TNF-alpha)” refers to a human TNF-alpha molecule comprising the amino acid sequence as described in Pennica *et al.*, *Nature*, 312:721 (1984) or Aggarwal *et al.*, *JBC*, 260:2345 (1985).

20 A “TNF-alpha inhibitor” herein is an agent that inhibits, to some extent, a biological function of TNF-alpha, generally through binding to TNF-alpha and neutralizing its activity. Examples of TNF inhibitors specifically contemplated herein are Etanercept (ENBREL®), Infliximab (REMICADE®) and Adalimumab (HUMIRA™).

25 Examples of “disease-modifying anti-rheumatic drugs” or “DMARDs” include hydroxycchloroquine, sulfasalazine, methotrexate, leflunomide, etanercept, infliximab (plus oral and subcutaneous methotrexate), azathioprine, D-penicillamine, Gold (oral), Gold (intramuscular), minocycline, cyclosporine, Staphylococcal protein A immunoabsorption, including salts and derivatives thereof, etc.

Examples of “nonsteroidal anti-inflammatory drugs” or “NSAIDs” are acetylsalicylic acid, ibuprofen, naproxen, indomethacin, sulindac, tolmetin, including salts and derivatives thereof, etc.

30 “Corticosteroid” refers to any one of several synthetic or naturally occurring substances with the general chemical structure of steroids that mimic or augment the effects of the naturally occurring corticosteroids. Examples of synthetic corticosteroids include prednisone, prednisolone (including methylprednisolone), dexamethasone, glucocorticoid and betamethasone.

A “package insert” is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage,

administration, contraindications, other therapeutic products to be combined with the packaged product, and/or warnings concerning the use of such therapeutic products, etc.

II. Therapy

Disclosed herein is a method of treating multiple sclerosis in a subject suffering therefrom, comprising administering an effective amount of an antibody that binds to a B-cell surface marker (preferably a CD20 antibody) to the subject. The MS to be treated herein includes primary progressive multiple sclerosis (PPMS), relapsing-remitting multiple sclerosis (RRMS), secondary progressive multiple sclerosis (SPMS), and progressive relapsing multiple sclerosis (PRMS), but therapy of either PPMS or RRMS are the preferred embodiments herein.

According to a preferred dosing protocol, the method comprises administering an effective amount of a CD20 antibody to the MS subject to provide an initial antibody exposure of about 0.5 to 4 grams (preferably about 1.5 to 2.5 grams) followed by a second antibody exposure of about 0.5 to 4 grams (preferably about 1.5 to 2.5 grams), the second antibody exposure not being provided until from about 16 to 60 weeks from the initial antibody exposure. For purposes of this invention, the second antibody exposure is the next time the subject is treated with the CD20 antibody after the initial antibody exposure, there being no intervening CD20 antibody treatment or exposure between the initial and second exposures.

The interval between the initial and second or subsequent antibody exposures can be measured from either the first or second dose of the initial antibody exposure, but preferably from the first dose of the initial antibody exposure.

In the preferred embodiments herein, the antibody exposures are approximately 24 weeks or 6 months apart; or approximately 48 weeks or 12 months apart.

In one embodiment, the second antibody exposure is not provided until about 20 to 30 weeks from the initial exposure, optionally followed by a third antibody exposure of about 0.5 to 4 grams (preferably about 1.5 to 2.5 grams), the third exposure not being administered until from about 46 to 60 weeks (preferably from about 46 to 54 weeks) from the initial exposure, and then, preferably no further antibody exposure is provided until at least about 70-75 weeks from the initial exposure.

In an alternative embodiment, the second antibody exposure is not provided until about 46 to 60 weeks from the initial exposure, and subsequent antibody exposures, if any, are not provided until about 46 to 60 weeks from the previous antibody exposure.

Any one or more of the antibody exposures herein may be provided to the subject as a single dose of antibody, or as two separate doses of the antibody (*i.e.*, constituting a first and second dose). The particular number of doses (whether one or two) employed for each antibody exposure is dependent, for example, on the type of MS treated, the type of antibody employed, whether and what type of second medicament is employed, and the method and frequency of administration. Where two separate doses are administered, the second dose is preferably administered from about 3 to 17 days, more preferably from about 6 to 16 days, and most preferably from about 13 to 16 days from the

time the first dose was administered. Where two separate doses are administered, the first and second dose of the antibody is preferably about 0.5 to 1.5 grams, more preferably about 0.75 to 1.3 grams.

In one embodiment, the subject is provided at least about three, or at least four exposures of the antibody, for example, from about 3 to 60 exposures, and more particularly about 3 to 40 exposures, most particularly, about 3 to 20 exposures. Preferably, such exposures are administered at intervals each of approximately 24 weeks or 6 months, or 48 weeks or 12 months. In one embodiment, each antibody exposure is provided as a single dose of the antibody. In an alternative embodiment, each antibody exposure is provided as two separate doses of the antibody. However, not every antibody exposure need be provided as a single dose or as two separate doses.

10 The antibody may be a naked antibody or may be conjugated with another molecule such as a cytotoxic agent such as a radioactive compound. The preferred antibody herein is Rituximab, humanized 2H7 (e.g. comprising the variable domain sequences in SEQ ID NOS. 2 and 8) or humanized 2H7 comprising the variable domain sequences in SEQ ID NOS. 23 and 24, or huMax-CD20 (Genmab).

15 In one embodiment, the subject has never been previously treated with drug(s), such as immunosuppressive agent(s), to treat the multiple sclerosis and/or has never been previously treated with an antibody to a B-cell surface marker (e.g. never previously treated with a CD20 antibody).

The antibody is administered by any suitable means, including parenteral, topical, subcutaneous, intraperitoneal, intrapulmonary, intranasal, and/or intralesional administration. 20 Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. Intrathecal administration is also contemplated (see, e.g., US Patent Appln No. 2002/0009444, Grillo-Lopez, A concerning intrathecal delivery of a CD20 antibody). In addition, the antibody may suitably be administered by pulse infusion, e.g., with declining doses of the antibody. Preferably, the dosing is given intravenously, subcutaneously or intrathecally, most preferably by 25 intravenous infusion(s).

While the CD20 antibody may be the only drug administered to the subject to treat the multiple sclerosis, one may optionally administer a second medicament, such as a cytotoxic agent, chemotherapeutic agent, immunosuppressive agent, cytokine, cytokine antagonist or antibody, growth factor, hormone, integrin, integrin antagonist or antibody (e.g. an LFA-1 antibody such as efalizumab 30 (RAPTIVA®) commercially available from Genentech, or an alpha 4 integrin antibody such as natalizumab (TYSABRI®) available from Biogen) etc, with the antibody that binds a B cell surface marker (e.g. with the CD20 antibody).

In the preferred embodiment of combination therapy, the antibody is combined with an interferon class drug such as IFN-beta-1a (REBIF® and AVONEX®) or IFN-beta-1b 35 (BETASERON®); an oligopeptide such a glatiramer acetate (COPAXONE®); a cytotoxic agent such as mitoxantrone (NOVANTRONE®), methotrexate, cyclophosphamide, chlorambucil, azathioprine;

intravenous immunoglobulin (gamma globulin); lymphocyte-depleting therapy (e.g., mitoxantrone, cyclophosphamide, Campath, anti-CD4, cladribine, total body irradiation, bone marrow transplantation); corticosteroid (e.g. methylprednisolone, prednisone, dexamethasone, or glucocorticoid), including systemic corticosteroid therapy; non-lymphocyte-depleting immunosuppressive therapy (e.g., mycophenolate mofetil (MMF) or cyclosporine); cholesterol-lowering drug of the “statin” class, which includes cerivastatin (BAYCOL®), fluvastatin (LESCOL®), atorvastatin (LIPITOR®), lovastatin (MEVACOR®), pravastatin (PRAVACHOL®), Simvastatin (ZOCOR®); estradiol; testosterone (optionally at elevated dosages; Stuve *et al.*

Neurology 8:290–301 (2002)); hormone replacement therapy; treatment for symptoms secondary or related to MS (e.g., spasticity, incontinence, pain, fatigue); a TNF inhibitor; disease-modifying anti-rheumatic drug (DMARD); non-steroidal anti-inflammatory drug (NSAID); plasmapheresis; levothyroxine; cyclosporin A; somatostatin analogue; cytokine or cytokine receptor antagonist; anti-metabolite; immunosuppressive agent; rehabilitative surgery; radioiodine; thyroidectomy; another B-cell surface antagonist/antibody; etc.

The second medicament is administered with the initial exposure and/or later exposures of the CD20 antibody, such combined administration includes co-administration, using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents simultaneously exert their biological activities.

Aside from administration of antibodies to the subject the present application contemplates administration of antibodies by gene therapy. Such administration of nucleic acid encoding the antibody is encompassed by the expression administering an “effective amount” of an antibody. See, for example, WO96/07321 published March 14, 1996 concerning the use of gene therapy to generate intracellular antibodies.

There are two major approaches to getting the nucleic acid (optionally contained in a vector) into the subject's cells; *in vivo* and *ex vivo*. For *in vivo* delivery the nucleic acid is injected directly into the subject, usually at the site where the antibody is required. For *ex vivo* treatment, the subject's cells are removed, the nucleic acid is introduced into these isolated cells and the modified cells are administered to the subject either directly or, for example, encapsulated within porous membranes that are implanted into the subject (see, *e.g.* U.S. Patent Nos. 4,892,538 and 5,283,187). There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. A commonly used vector for *ex vivo* delivery of the gene is a retrovirus.

The currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral vectors (such as adenovirus, Herpes simplex I virus, or adeno-associated virus) and lipid-based systems (useful lipids for lipid-mediated transfer of the gene are DOTMA, DOPE and DC-Chol, for example). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins that bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, *e.g.* capsid proteins or fragments thereof specific for a particular cell type, antibodies for proteins that undergo internalization in cycling, and proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu *et al.*, *J. Biol. Chem.* 262:4429-4432 (1987); and Wagner *et al.*, *Proc. Natl. Acad. Sci. USA* 87:3410-3414 (1990). For review of the currently known gene marking and gene therapy protocols see Anderson *et al.*, *Science* 256:808-813 (1992). See also WO 93/25673 and the references cited therein.

15 III. Production of Antibodies

The methods and articles of manufacture of the present invention use, or incorporate, an antibody that binds to a B-cell surface marker, especially one that binds to CD20. Accordingly, methods for generating such antibodies will be described here.

The B cell surface marker to be used for production of, or screening for, antibodies may be, 20 *e.g.*, a soluble form of the marker or a portion thereof, containing the desired epitope. Alternatively, or additionally, cells expressing the marker at their cell surface can be used to generate, or screen for, antibodies. Other forms of the B cell surface marker useful for generating antibodies will be apparent to those skilled in the art.

A description follows as to exemplary techniques for the production of the antibodies used in 25 accordance with the present invention.

(i) Polyclonal antibodies

30 Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the relevant antigen to a protein that is immunogenic in the species to be immunized, *e.g.*, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl groups.

35 Animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, *e.g.*, 100 μg or 5 μg of the protein or conjugate (for rabbits or mice, respectively) with 3

5 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitably used to enhance the immune response.

10 (ii) *Monoclonal antibodies*

15 Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical and/or bind the same epitope except for possible variants that arise during production of the monoclonal antibody, such variants generally being present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete or polyclonal antibodies.

20 For example, the monoclonal antibodies may be made using the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

25 In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)).

30 The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

35 Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal

antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

5 Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson *et al.*, *Anal. Biochem.*, 107:220 (1980).

10 After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal.

15 The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

20 DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese Hamster Ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant 25 expression in bacteria of DNA encoding the antibody include Skerra *et al.*, *Curr. Opinion in Immunol.*, 5:256-262 (1993) and Plückthun, *Immunol. Revs.*, 130:151-188 (1992).

30 In a further embodiment, antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty *et al.*, *Nature*, 348:552-554 (1990). Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by 35 chain shuffling (Marks *et al.*, *Bio/Technology*, 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse *et al.*, *Nuc. Acids. Res.*, 21:2265-2266 (1993)). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

¶¶¶¶¶ The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, *et al.*, *Proc. Natl Acad. Sci. USA*, 81:6851 (1984)), or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

5 Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody, or they are substituted for the variable domains of one antigen-combining site of an antibody to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an antigen and another antigen-combining site having specificity for a different 10 antigen.

10 (iii) *Humanized antibodies*

Methods for humanizing non-human antibodies have been described in the art. Preferably, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human. These non-human amino acid residues are often referred to as "import" residues, which are 15 typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, *Nature*, 321:522-525 (1986); Riechmann *et al.*, *Nature*, 332:323-327 (1988); Verhoeyen *et al.*, *Science*, 239:1534-1536 (1988)), by substituting hypervariable region sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567) 20 wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some hypervariable region residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the 25 humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence that is closest to that of the rodent is then accepted as the human framework region (FR) for the humanized antibody (Sims *et al.*, *J. Immunol.*, 151:2296 (1993); Chothia *et al.*, *J. Mol. Biol.*, 196:901 (1987)). Another method uses a 30 particular framework region derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chain variable regions. The same framework may be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta *et al.*, *J. Immunol.*, 151:2623 (1993)).

It is further important that antibodies be humanized with retention of high affinity for the 35 antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and

various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available that illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

5 (iv) *Human antibodies*

As an alternative to humanization, human antibodies can be generated. For example, it is now possible to produce transgenic animals (*e.g.*, mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, *e.g.*, Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993); Jakobovits *et al.*, *Nature*, 362:255-258 (1993); Brugermann *et al.*, *Year in Immuno.*, 7:33 (1993); and US Patent Nos. 5,591,669, 5,589,369 and 5,545,807.

15 Alternatively, phage display technology (McCafferty *et al.*, *Nature* 348:552-553 (1990)) can be used to produce human antibodies and antibody fragments *in vitro*, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimics some of the properties of the B cell. Phage display can be performed in a variety of formats; for their review see, *e.g.*, Johnson, Kevin S. and Chiswell, David J., *Current Opinion in Structural Biology* 3:564-571 (1993). Several sources of V-gene segments can be used for phage display. Clackson *et al.*, *Nature*, 352:624-628 (1991) isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V genes derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by

Marks *et al.*, *J. Mol. Biol.* 222:581-597 (1991), or Griffith *et al.*, *EMBO J.* 12:725-734 (1993). See, also, US Patent Nos. 5,565,332 and 5,573,905.

Human antibodies may also be generated by *in vitro* activated B cells (see US Patents 5,567,610 and 5,229,275).

5 (v) *Antibody fragments*

Various techniques have been developed for the production of antibody fragments.

Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, *e.g.*, Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992) and Brennan *et al.*, *Science*, 229:81 (1985)). However, these fragments can now be produced directly by recombinant 10 host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, *Bio/Technology* 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell 15 culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). See WO 93/16185; US Patent No. 5,571,894; and US Patent No. 5,587,458. The antibody fragment may also be a “linear antibody”, *e.g.*, as described in US Patent 5,641,870 for example. Such linear antibody fragments may be monospecific or bispecific.

(vi) *Bispecific antibodies*

20 Bispecific antibodies are antibodies that have binding specificities for at least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of the B cell surface marker. Other such antibodies may bind the B cell surface marker and further bind a second different B-cell surface marker. Alternatively, an anti-B cell surface marker binding arm may be combined with an arm that binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (*e.g.* 25 CD2 or CD3), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the B cell. Bispecific antibodies may also be used to localize cytotoxic agents to the B cell. These antibodies possess a B cell surface marker-binding arm and an arm that binds the cytotoxic agent (*e.g.* saporin, anti-interferon- α , vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies can be prepared as 30 full length antibodies or antibody fragments (*e.g.* F(ab')₂ bispecific antibodies).

Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein *et al.*, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these 35 hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually

done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker *et al.*, *EMBO J.*, 10:3655-3659 (1991).

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986).

According to another approach described in US Patent No. 5,731,168, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers that are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089).

Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

Techniques for generating bispecific antibodies from antibody fragments have also been 5 described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science*, 229: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) 10 derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Various techniques for making and isolating bispecific antibody fragments directly from 15 recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelnik *et al.*, *J. Immunol.*, 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized 20 for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy chain variable domain (V_H) connected to a light chain variable domain (V_L) by a linker that is too short to allow pairing between 25 the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, *J. Immunol.*, 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.* *J. Immunol.* 147: 60 (1991).

30 IV. Conjugates and Other Modifications of the Antibody'

The antibody used in the methods or included in the articles of manufacture herein is optionally conjugated to a cytotoxic agent. For instance, the antibody may be conjugated to a drug as described in WO2004/032828.

35 Chemotherapeutic agents useful in the generation of such antibody-cytotoxic agent conjugates have been described above.

Conjugates of an antibody and one or more small molecule toxins, such as a calicheamicin, a maytansine (US Patent No. 5,208,020), a trichothene, and CC1065 are also contemplated herein. In one embodiment of the invention, the antibody is conjugated to one or more maytansine molecules (e.g. about 1 to about 10 maytansine molecules per antibody molecule). Maytansine may, for example, be converted to May-SS-Me, which may be reduced to May-SH3 and reacted with modified antibody (Chari *et al. Cancer Research* 52: 127-131 (1992)) to generate a maytansinoid-antibody conjugate.

Alternatively, the antibody is conjugated to one or more calicheamicin molecules. The calicheamicin family of antibiotics are capable of producing double-stranded DNA breaks at sub-picomolar concentrations. Structural analogues of calicheamicin that may be used include, but are not limited to, γ_1^1 , α_2^1 , α_3^1 , N-acetyl- γ_1^1 , PSAG and θ^1 (Hinman *et al. Cancer Research* 53: 3336-3342 (1993) and Lode *et al. Cancer Research* 58: 2925-2928 (1998)).

Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the trichothecenes. See, for example, WO 93/21232 published October 28, 1993.

The present invention further contemplates use of an antibody conjugated with a compound with nucleolytic activity (e.g. a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

A variety of radioactive isotopes are available for the production of radioconjugated antibodies. Examples include At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³² and radioactive isotopes of Lu.

Conjugates of the antibody and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate, iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al. Science* 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026. The linker may be a "cleavable linker" facilitating release of the cytotoxic drug in the cell. For example, an acid-labile linker,

peptidase-sensitive linker, dimethyl linker or disulfide-containing linker (Chari *et al. Cancer Research* 52: 127-131 (1992)) may be used.

Alternatively, a fusion protein comprising the antibody and cytotoxic agent may be made, *e.g.* by recombinant techniques or peptide synthesis.

5 In yet another embodiment, the antibody may be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the subject, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (*e.g.* avidin) that is conjugated to a cytotoxic agent (*e.g.* a radionucleotide).

10 The antibodies used in the method of the present invention may also be conjugated with a prodrug-activating enzyme that converts a prodrug (*e.g.* a peptidyl chemotherapeutic agent, see WO81/01145) to an active anti-cancer drug. See, for example, WO 88/07378 and U.S. Patent No. 4,975,278.

15 The enzyme component of such conjugates includes any enzyme capable of acting on a prodrug in such a way so as to convert it into its more active, cytotoxic form.

Enzymes that are useful in the method of this invention include, but are not limited to, alkaline phosphatase useful for converting phosphate-containing prodrugs into free drugs; arylsulfatase useful for converting sulfate-containing prodrugs into free drugs; cytosine deaminase useful for converting non-toxic 5-fluorocytosine into the anti-cancer drug, 5-fluorouracil; proteases, 20 such as serratia protease, thermolysin, subtilisin, carboxypeptidases and cathepsins (such as cathepsins B and L), that are useful for converting peptide-containing prodrugs into free drugs; D-alanylcarboxypeptidases, useful for converting prodrugs that contain D-amino acid substituents; carbohydrate-cleaving enzymes such as β -galactosidase and neuraminidase useful for converting glycosylated prodrugs into free drugs; β -lactamase useful for converting drugs derivatized with β -lactams into free drugs; and penicillin amidases, such as penicillin V amidase or penicillin G amidase, 25 useful for converting drugs derivatized at their amine nitrogens with phenoxyacetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies with enzymatic activity, also known in the art as "abzymes", can be used to convert the prodrugs of the invention into free active drugs (see, *e.g.*, Massey, *Nature* 328: 457-458 (1987)). Antibody-abzyme conjugates can be prepared as 30 described herein for delivery of the abzyme to a tumor cell population.

The enzymes that may be used in the method of this invention can be covalently bound to the antibody by techniques well known in the art such as the use of the heterobifunctional crosslinking reagents discussed above. Alternatively, fusion proteins comprising at least the antigen binding region of an antibody of the invention linked to at least a functionally active portion of an enzyme of 35 the invention can be constructed using recombinant DNA techniques well known in the art (see, *e.g.*, Neuberger *et al.*, *Nature*, 312: 604-608 (1984)).

Other modifications of the antibody are contemplated herein. For example, the antibody may be linked to one of a variety of nonproteinaceous polymers, *e.g.*, polyethylene glycol (PEG), polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol. Antibody fragments, such as Fab', linked to one or more PEG molecules are an especially preferred embodiment of the invention.

The antibodies disclosed herein may also be formulated as liposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, *Proc. Natl. Acad. Sci. USA*, 82:3688 (1985); Hwang *et al.*, *Proc. Natl. Acad. Sci. USA*, 77:4030 (1980); U.S. Pat. Nos. 4,485,045 and 4,544,545; and WO97/38731 published October 23, 1997. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of an antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.* *J. Biol. Chem.* 257: 286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent is optionally contained within the liposome. See Gabizon *et al.* *J. National Cancer Inst.* 81(19)1484 (1989).

Amino acid sequence modification(s) of the antibody are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of the antibody are prepared by introducing appropriate nucleotide changes into the antibody nucleic acid, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the antibody, such as changing the number or position of glycosylation sites.

A useful method for identification of certain residues or regions of the antibody that are preferred locations for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham and Wells *Science*, 244:1081-1085 (1989). Here, a residue or group of target residues are identified (*e.g.*, charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with antigen. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* need not be predetermined. For example, to analyze the

performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed antibody variants are screened for the desired activity.

Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as

5 intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue or the antibody fused to a cytotoxic polypeptide. Other insertional variants of the antibody molecule include the fusion to the N- or C- terminus of the antibody of an enzyme, or a polypeptide that increases the serum half-life of the antibody.

10 Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the antibody molecule replaced by different residue. The sites of greatest interest for substitutional mutagenesis of antibody antibodies include the hypervariable regions, but FR alterations are also contemplated. Conservative substitutions are shown in Table 1 under the heading of "preferred substitutions". If such substitutions result in a change in biological activity, 15 then more substantial changes, denominated "exemplary substitutions" in Table 1, or as further described below in reference to amino acid classes, may be introduced and the products screened.

Table 1

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu

Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, 5 (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Amino acids may be grouped according to similarities in the properties of their side chains (in A. L. Lehninger, in *Biochemistry*, second ed., pp. 73-75, Worth Publishers, New York (1975)):

- (1) non-polar: Ala (A), Val (V), Leu (L), Ile (I), Pro (P), Phe (F), Trp (W), Met (M)
- (2) uncharged polar: Gly (G), Ser (S), Thr (T), Cys (C), Tyr (Y), Asn (N), Gln (Q)
- 10 (3) acidic: Asp (D), Glu (E)
- (4) basic: Lys (K), Arg (R), His(H)

Alternatively, naturally occurring residues may be divided into groups based on common side-chain properties:

- (1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;
- 15 (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;
- (3) acidic: Asp, Glu;
- (4) basic: His, Lys, Arg;
- (5) residues that influence chain orientation: Gly, Pro;
- (6) aromatic: Trp, Tyr, Phe.

20 Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

Any cysteine residue not involved in maintaining the proper conformation of the antibody also may be substituted, generally with serine, to improve the oxidative stability of the molecule and 25 prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

A particularly preferred type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody. Generally, the resulting variant(s) selected for

further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants is affinity maturation using phage display. Briefly, several hypervariable region sites (e.g. 6-7 sites) are mutated to generate all possible amino substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (e.g. binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or in addition, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

Another type of amino acid variant of the antibody alters the original glycosylation pattern of the antibody. Such altering includes deleting one or more carbohydrate moieties found in the antibody, and/or adding one or more glycosylation sites that are not present in the antibody.

Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylglucosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

Where the antibody comprises an Fc region, the carbohydrate attached thereto may be altered. For example, antibodies with a mature carbohydrate structure that lacks fucose attached to an Fc region of the antibody are described in US Pat Appl No US 2003/0157108 A1, Presta, L. See also US 2004/0093621 A1 (Kyowa Hakko Kogyo Co., Ltd) concerning a CD20 antibody composition. Antibodies with a bisecting N-acetylglucosamine (GlcNAc) in the carbohydrate attached to an Fc

region of the antibody are referenced in WO03/011878, Jean-Mairet *et al.* and US Patent No. 6,602,684, Umana *et al.* Antibodies with at least one galactose residue in the oligosaccharide attached to an Fc region of the antibody are reported in WO97/30087, Patel *et al.* See, also, WO98/58964 (Raju, S.) and WO99/22764 (Raju, S.) concerning antibodies with altered carbohydrate attached to the Fc region thereof.

The preferred glycosylation variant herein comprises an Fc region, wherein a carbohydrate structure attached to the Fc region lacks fucose. Such variants have improved ADCC function. Optionally, the Fc region further comprises one or more amino acid substitutions therein which further improve ADCC, for example, substitutions at positions 298, 333, and/or 334 of the Fc region (Eu numbering of residues). Examples of publications related to "defucosylated" or "fucose-deficient" antibodies include: US Pat. Appl. No. US 2003/0157108 A1, Presta, L; WO 00/61739A1; WO01/29246A1; US2003/0115614A1; US2002/0164328A1; US2004/0093621A1; US2004/0132140A1; US2004/0110704A1; US2004/0110282A1; US2004/0109865A1; WO03/085119A1; WO03/084570A1; WO2005/035778; WO2005/035586 (describing RNA inhibition (RNAi) of fucosylation); Okazaki *et al.* *J. Mol. Biol.* 336:1239-1249 (2004); Yamane-Ohnuki *et al.* *Biotech. Bioeng.* 87: 614 (2004). Examples of cell lines producing defucosylated antibodies include Lec13 CHO cells deficient in protein fucosylation (Ripka *et al.* *Arch. Biochem. Biophys.* 249:533-545 (1986); US Pat Appl No US 2003/0157108 A1, Presta, L; and WO 2004/056312 A1, Adams *et al.*, especially at Example 11), and knockout cell lines, such as alpha-1,6-fucosyltransferase gene, *FUT8*, knockout CHO cells (Yamane-Ohnuki *et al.* *Biotech. Bioeng.* 87: 614 (2004)).

Nucleic acid molecules encoding amino acid sequence variants of the antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the antibody.

It may be desirable to modify the antibody used in the method of the invention with respect to effector function, *e.g.* so as to enhance antigen-dependent cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC) of the antibody. This may be achieved by introducing one or more amino acid substitutions in an Fc region of an antibody antibody. Alternatively or additionally, cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, *J. Exp Med.* 176:1191-1195 (1992) and Shope, B. *J. Immunol.* 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.* *Cancer Research* 53:2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may

thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.* *Anti-Cancer Drug Design* 3:219-230 (1989).

WO00/42072 (Presta, L.) describes antibodies with improved ADCC function in the presence of human effector cells, where the antibodies comprise amino acid substitutions in the Fc region thereof. Preferably, the antibody with improved ADCC comprises substitutions at positions 298, 10 333, and/or 334 of the Fc region. Preferably the altered Fc region is a human IgG1 Fc region comprising or consisting of substitutions at one, two or three of these positions.

Antibodies with altered C1q binding and/or complement dependent cytotoxicity (CDC) are described in WO99/51642, US Patent No. 6,194,551B1, US Patent No. 6,242,195B1, US Patent No. 10 6,528,624B1 and US Patent No. 6,538,124 (Idusogie *et al.*). The antibodies comprise an amino acid substitution at one or more of amino acid positions 270, 322, 326, 327, 329, 313, 333 and/or 334 of the Fc region thereof.

To increase the serum half life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment) as described in US Patent 15 5,739,277, for example. As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule. Antibodies with substitutions in an Fc region thereof and increased serum half-lives are also described in WO00/42072 (Presta, L.).

Engineered antibodies with three or more (preferably four) functional antigen binding sites 20 are also contemplated (US Appln No. US2002/0004587 A1, Miller *et al.*).

V. Pharmaceutical Formulations

Therapeutic formulations of the antibodies used in accordance with the present invention are prepared for storage by mixing an antibody having the desired degree of purity with optional pharmaceutical acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 25 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; 30 phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, 35 mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or

sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICSTM or polyethylene glycol (PEG).

Exemplary anti-CD20 antibody formulations are described in WO98/56418. This publication describes a liquid multidose formulation comprising 40 mg/mL Rituximab, 25 mM acetate, 150 mM trehalose, 0.9% benzyl alcohol, 0.02% polysorbate 20 at pH 5.0 that has a minimum shelf life of two years storage at 2-8°C. Another anti-CD20 formulation of interest comprises 10mg/mL Rituximab in 9.0 mg/mL sodium chloride, 7.35 mg/mL sodium citrate dihydrate, 0.7mg/mL polysorbate 80, and Sterile Water for Injection, pH 6.5.

Lyophilized formulations adapted for subcutaneous administration are described in US Pat 10 No. 6,267,958 (Andya *et al.*). Such lyophilized formulations may be reconstituted with a suitable diluent to a high protein concentration and the reconstituted formulation may be administered subcutaneously to the mammal to be treated herein.

Crystallized forms of the antibody or antibody are also contemplated. See, for example, US 2002/0136719A1 (Shenoy *et al.*).

15 The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide a cytotoxic agent; chemotherapeutic agent; immunosuppressive agent; cytokine; cytokine antagonist or antibody; growth factor; hormone; integrin; integrin antagonist or antibody (e.g. an LFA-1 antibody such as 20 efalizumab/RAPTIVA commercially available from Genentech, or an alpha 4 integrin antibody such as natalizumab/TYSABRI® available from Biogen); interferon class drug such as IFN-beta-1a (REBIF® and AVONEX®) or IFN-beta-1b (BETASERON®); an oligopeptide such a glatiramer acetate (COPAXONE®); a cytotoxic agent such as mitoxantrone (NOVANTRONE®), methotrexate, cyclophosphamide, chlorambucil, or azathioprine; intravenous immunoglobulin (gamma globulin); 25 lymphocyte-depleting drug (e.g., mitoxantrone, cyclophosphamide, Campath, anti-CD4, or cladribine); non-lymphocyte-depleting immunosuppressive drug (e.g., mycophenolate mofetil (MMF) or cyclosporine); cholesterol-lowering drug of the “statin” class; estradiol; testosterone; hormone replacement therapy; drug that treats symptoms secondary or related to MS (e.g., spasticity, incontinence, pain, fatigue); a TNF inhibitor; disease-modifying anti-rheumatic drug (DMARD); non- 30 steroidal anti-inflammatory drug (NSAID); corticosteroid (e.g. methylprednisolone, prednisone, dexamethasone, or glucocorticoid); levothyroxine; cyclosporin A; somatostatin analogue; cytokine antagonist; anti-metabolite; immunosuppressive agent; integrin antagonist or antibody (e.g. an LFA-1 antibody, such as efalizumab or an alpha 4 integrin antibody such as natalizumab); or another B-cell surface antagonist/antibody; etc in the formulation. The type and effective amounts of such other 35 agents depend, for example, on the amount of antibody present in the formulation, the type of multiple sclerosis being treated, and clinical parameters of the subjects. These are generally used in

the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the heretofore employed dosages.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or 5 gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980).

Sustained-release preparations may be prepared. Suitable examples of sustained-release 10 preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.* films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-15 glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

VI. Articles of Manufacture

An article of manufacture containing materials useful for the treatment of multiple sclerosis 20 described above is also disclosed. Preferably, the article of manufacture comprises:(a) a container comprising a composition comprising an antibody that binds to a B-cell surface marker (*e.g.* a CD20 antibody) and a pharmaceutically acceptable carrier or diluent within the container; and (b) a package insert with instructions for administering the composition to a subject suffering from multiple 25 sclerosis to the subject to provide an initial antibody exposure of about 0.5 to 4 grams followed by a second antibody exposure of about 0.5 to 4 grams, the second exposure not being provided until from about 16 to 60 weeks from the initial exposure; or instructions for administering the composition to a subject suffering from PPMS.

The article of manufacture may comprise a container and a label or package insert on or 30 associated with the container. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds or contains a composition that is effective for treating the multiple sclerosis and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is 35 the antibody. The label or package insert indicates that the composition is used for treating multiple sclerosis in a subject suffering therefrom with specific guidance regarding dosing amounts and

intervals of antibody and any other drug being provided. The article of manufacture may further comprise a second container comprising a pharmaceutically acceptable diluent buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. The article of manufacture may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

Optionally, the article of manufacture herein further comprises a container comprising an agent other than the antibody for treatment and further comprising instructions on treating the mammal with such agent, such agent preferably being a chemotherapeutic agent or immunosuppressive agent, interferon class drug such as IFN-beta-1a (REBIF® and AVONEX®) or IFN-beta-1b (BETASERON®); an oligopeptide such a glatiramer acetate (COPAXONE®); a cytotoxic agent such as mitoxantrone (NOVANTRONE®), methotrexate, cyclophosphamide, chlorambucil, or azathioprine; intravenous immunoglobulin (gamma globulin); lymphocyte-depleting drug (e.g., mitoxantrone, cyclophosphamide, Campath, anti-CD4, or cladribine); non-lymphocyte-depleting immunosuppressive drug (e.g., mycophenolate mofetil (MMF) or cyclosporine); cholesterol-lowering drug of the "statin" class; estradiol; hormone replacement therapy; drug that treats symptoms secondary or related to MS (e.g., spasticity, incontinence, pain, fatigue); a TNF inhibitor; disease-modifying anti-rheumatic drug (DMARD); non-steroidal anti-inflammatory drug (NSAID); corticosteroid (e.g. methylprednisolone, prednisone, dexamethasone, or glucocorticoid); levothyroxine; cyclosporin A; somatostatin analogue; cytokine or cytokine receptor antagonist; anti-metabolite; immunosuppressive agent; integrin antagonist or antibody (e.g. an LFA-1 antibody, such as efalizumab or an alpha 4 integrin antibody such as natalizumab); and another B-cell surface marker antibody; etc.

Further details of the invention are illustrated by the following non-limiting Examples. The disclosures of all citations in the specification are expressly incorporated herein by reference.

25

EXAMPLE 1

TREATMENT OF PRIMARY PROGRESSIVE MULTIPLE SCLEROSIS (PPMS)

A subject with diagnosis of PPMS as defined by McDonald *et al. Ann Neurol* 50:121-7 (2001) is treated with a CD20 antibody in this example.

30 Rituximab, commercially available from Genentech, is formulated for IV administration as a sterile product in 9.0 mg/mL sodium chloride, 0.7 mg/mL polysorbate 80, 7.35 mg/mL sodium citrate dehydrate, and Sterile Water for Injection (pH 6.5).

The first course of treatment will consist of a dose of 1 g intravenous (IV) Rituximab administered on each of Days 1 and 15. Subjects will receive acetaminophen (1 g) and diphenhydramine HCl (50 mg), or equivalent, by mouth 30–60 minutes prior to the start of each infusion.

5 Subsequent courses of treatment will be administered starting at Week 24 (Day 169), Week 48 (Day 337), and Week 72 (Day 505). The second infusion of the subsequent courses of treatment will be 14 ± 1 days after the first infusion.

10 Subjects who experience a first relapse may receive rescue treatment with IV or oral corticosteroids. Systemic corticosteroids may be administered using a regimen that does not exceed exposure or duration of treatment appropriate for an MS relapse. A relapse is defined as all of the following:

- An acute appearance of a neurologic abnormality that persists for at least 24 hours
- A change not attributable to fever, infection, trauma, concomitant medications, or other etiology
- 15 • An event with objective change on examination by the blinded examining investigator, including a minimum of 1-point change on one of the following FS scales: pyramidal, cranial nerves, cerebellar, sensory, vision, or gait

The following regimen of corticosteroids may be used: 1 g IV methylprednisolone daily for 3 days, followed by 60 mg prednisone daily for 5 days, and decreasing by 10-mg increments each day thereafter. If IV methylprednisolone is not available, then 150 mg IV dexamethasone daily for 3 days may be substituted. Only one course of corticosteroids should be administered for an exacerbation. Subsequent immunological studies and MRI scans should be obtained at least 4 weeks after completion of the corticosteroid regimen. Corticosteroid inhalers (oral and nasal) or intra-articular injections may be used.

25 Additional treatments to be optionally combined with the CD20 antibody include IFN-beta, glatiramer acetate, methotrexate, cyclophosphamide, or mitoxantrone.

5 The primary efficacy outcome measure is the time to confirmed disease progression. Disease progression is defined as an increase of ≥ 1.0 point from baseline Expanded Disability Status Scale (EDSS) (Kurtzke *J. Neurology* 33(11):1444-52 (1983)), if the baseline EDSS is between 2.0 and 5.5 points (inclusive), or an increase of ≥ 0.5 point if the baseline EDSS is ≥ 5.5 points (inclusive), for which change is not attributable to another etiology (e.g., fever, concurrent illness, MS relapse or exacerbation, or concomitant medication). Confirmation of disease progression may occur at a regularly scheduled visit that is at least 12 weeks (84 days) after the initial progression.

Secondary efficacy outcome measures include:

10

- Change from baseline to Week 96 in the total volume of T2 lesions on brain MRI scan
- Change from baseline to Week 96 in brain volume on brain MRI scan

Optionally, improvement in any one or more of:

15

- Multiple Sclerosis Functional Composite Scale (MSFCS)
- EDSS
- Proportion of subjects with confirmed disease progression at Week 96, as determined using EDSS
- Upper extremity function, as measured by the 9-Hole Peg Test (a subscale of the MSFCS)
- Ambulation, as measured by the Timed 25-Foot Walk (a subscale of the MSFCS)
- Cognition, as measured by the Paced Auditory Serial Addition Test (3 seconds only; a subscale of the MSFCS)

20

- Total volume of brain T2 lesions on MRI scans (Weeks 48 and 122)
- Total volume of brain T1 lesions on MRI scans
- Cross sectional area of the cervical spinal cord on MRI scans
- Brain volume on MRI scans (Weeks 48 and 122)

25 The subject treated with Rituximab as described herein will show improvement in the signs, symptoms or other indicators of PPMS according to any one or more of the above outcome measures.

EXAMPLE 2**TREATMENT OF RELAPSING-REMITTING MULTIPLE SCLEROSIS**

Subjects with RRMS as defined by McDonald *et al. Ann Neurol* 50:121–7 (2001) are treated 5 with a CD20 antibody in this example, where the antibody exposures are approximately 6 months apart.

Rituximab, commercially available from Genentech, is formulated for IV administration as a sterile product in 9.0 mg/mL sodium chloride, 0.7 mg/mL polysorbate 80, 7.35 mg/mL sodium citrate dehydrate, and Sterile Water for Injection (pH 6.5).

10 The first course of treatment will consist of a dose of 1 g intravenous (IV) Rituximab administered on each of Days 1 and 15. Subjects will receive acetaminophen (1 g) and diphenhydramine HCl (50 mg), or equivalent, by mouth 30–60 minutes prior to the start of each infusion.

15 Subsequent courses of treatment will be administered starting at Week 24 (Day 169), Week 48 (Day 337), and Week 72 (Day 505). The second infusion of the subsequent courses of treatment will be 14 ± 1 days after the first infusion.

Preferably Rituximab is the only agent administered to treat the RRMS. However, subjects may optionally receive IV or oral corticosteroids, IFN-beta, glatiramer acetate, methotrexate, cyclophosphamide, or mitoxantrone.

20 Subjects who experience a first relapse may receive rescue treatment with IV or oral corticosteroids. Systemic corticosteroids may be administered using a regimen that does not exceed exposure or duration of treatment appropriate for an MS relapse. A relapse in this Example is defined as:

The appearance of new or recurrent neurological symptoms consistent with MS lasting more than 48 hours in a subject who has been in a relatively stable or improving neurologic state for at least 30 days. The change in neurologic symptoms must be accompanied by objective neurologic worsening consistent with an increase of at least half a step on the EDSS, or 2 points on one of the appropriate functional system scores (FSS), or 1 point on two or more of the appropriate FSS. The change must be verified by the examining investigator and must affect the selected FS scales (i.e., pyramidal, gait, cerebellar, brainstem, sensory, or visual). Symptoms must persist for \geq 24 hours and should not be attributable to confounding clinical factors (e.g., fever, infection, injury, adverse reactions to concomitant medications). A single episode of a paroxysmal symptom (e.g., tonic spasm) is not a relapse, but the new onset of multiple occurrences of a paroxysmal symptom over at least 24 hours can be a relapse if accompanied by new, corresponding objective manifestations. Sensory symptoms with no change on clinical examination, fatigue, mood change, or bladder or bowel urgency or incontinence will not be sufficient to establish a relapse.

The primary efficacy outcome measure is the MRI endpoint of gadolinium-enhancing lesions, or time to confirmed disease progression (defined as an increase of \geq 1.0 point from baseline Expanded Disability Status Scale (EDSS); Kurtzke *J. Neurology* 33(11):1444-52 (1983)). The primary efficacy endpoint may be the total number of gadolinium-enhancing T1 lesions observed on serial MRI scans of the brain at Weeks 12, 16, 20, and 24.

Secondary efficacy outcome measures include frequency of relapse; change from baseline to Week 96 in the total volume of T2 lesions on brain MRI scan (e.g. change in total volume of T2 lesions on MRA scans of the brain from screening to weeks 24 and 36); change from baseline to Week 96 in brain volume on brain MRI scan; Multiple Sclerosis Functional Composite Scale (MSFCS) and its subscales; upper extremity function, as measured by the 9-Hole Peg Test (a subscale of the MSFCS); ambulation, as measured by the Timed 25-Foot Walk (a subscale of the MSFCS); cognition, as measured by the Paced Auditory Serial Addition Test (a subscale of the MSFCS); Multiple Sclerosis Quality of Life-54 (MSQOL-54) questionnaire; total volume of brain T1 lesions on MRI scans (e.g. total number of gadolinium-enhancing T1 lesions observed on serial MRA scans of the brains at weeks 20, 28, and 36); cross sectional area of the cervical spinal cord on MRI scans; proportion of subjects relapsing by weeks 24 (i.e. between week 0 and week 24) and 36 (i.e. between week 0 and week 36); the Combined Unique Activity Measure at week 24 and 36.

The patient treated with Rituximab as described above will display an improvement in any one or more of the above outcome measures.

EXAMPLE 3**TREATMENT OF RELAPSING-REMITTING MULTIPLE SCLEROSIS**

5 A subject with RRMS as defined by McDonald *et al. Ann Neurol* 50:121–7 (2001) is treated with a CD20 antibody herein. In this example, the antibody exposures are approximately 1 year apart.

Rituximab, commercially available from Genentech, is formulated for IV administration as a sterile product in 9.0 mg/mL sodium chloride, 0.7 mg/mL polysorbate 80, 7.35 mg/mL sodium citrate dehydrate, and Sterile Water for Injection (pH 6.5).

10 The first course of treatment will consist of a dose of 1 g intravenous (IV) Rituximab administered on each of Days 1 and 15. Subjects will receive acetaminophen (1 g) and diphenhydramine HCl (50 mg), or equivalent, by mouth 30–60 minutes prior to the start of each infusion.

Subsequent courses of treatment will be administered starting at Week 48, and Week 96. The second infusion of the subsequent courses of treatment will be 14 ± 1 days after the first infusion.

15 Preferably Rituximab is the only agent administered to treat the RRMS. However, subjects may optionally receive IV or oral corticosteroids, IFN-beta, glatiramer acetate, methotrexate, cyclophosphamide, or mitoxantrone.

20 Subjects who experience a first relapse may receive rescue treatment with IV or oral corticosteroids. Systemic corticosteroids may be administered using a regimen that does not exceed exposure or duration of treatment appropriate for an MS relapse. A relapse in this Example is defined as:

The appearance of new or recurrent neurological symptoms consistent with MS lasting more than 48 hours in a subject who has been in a relatively stable or improving neurologic state for at least 30 days. The change in neurologic symptoms must be accompanied by objective neurologic worsening consistent with an increase of at least half a step on the EDSS, or 2 points on one of the appropriate functional system scores (FSS), or 1 point on two or more of the appropriate FSS. The change must be verified by the examining investigator and must affect the selected FS scales (i.e., pyramidal, gait, cerebellar, brainstem, sensory, or visual). Symptoms must persist for ≥ 24 hours and should not be attributable to confounding clinical factors (e.g., fever, infection, injury, adverse reactions to concomitant medications). A single episode of a paroxysmal symptom (e.g., tonic spasm) is not a relapse, but the new onset of multiple occurrences of a paroxysmal symptom over at least 24 hours can be a relapse if accompanied by new, corresponding objective manifestations. Sensory symptoms with no change on clinical examination, fatigue, mood change, or bladder or bowel urgency or incontinence will not be sufficient to establish a relapse.

The primary efficacy outcome measure is the MRI endpoint of gadolinium-enhancing lesions, or time to confirmed disease progression (defined as an increase of ≥ 1.0 point from baseline Expanded Disability Status Scale (EDSS); Kurtzke J. *Neurology* 33(11):1444–52 (1983)). The primary efficacy endpoint may be the total number of gadolinium-enhancing T1 lesions observed on serial MRI scans of the brain at Weeks 12, 16, 20, and 24.

Secondary efficacy outcome measures include frequency of relapse; change from baseline to Week 96 in the total volume of T2 lesions on brain MRI scan (e.g. change in total volume of T2 lesions on MRA scans of the brain from screening to weeks 24 and 36); change from baseline to Week 96 in brain volume on brain MRI scan; Multiple Sclerosis Functional Composite Scale (MSFCS) and its subscales; upper extremity function, as measured by the 9-Hole Peg Test (a subscale of the MSFCS); ambulation, as measured by the Timed 25-Foot Walk (a subscale of the MSFCS); cognition, as measured by the Paced Auditory Serial Addition Test (a subscale of the MSFCS); Multiple Sclerosis Quality of Life-54 (MSQOL-54) questionnaire; total volume of brain T1 lesions on MRI scans (e.g. total number of gadolinium-enhancing T1 lesions observed on serial MRA scans of the brains at weeks 20, 28, and 36); cross sectional area of the cervical spinal cord on MRI scans; proportion of subjects relapsing by week 24 (i.e. between week 0 and week 24) and week 36 (i.e. between week 0 and week 36); the Combined Unique Activity Measure at week 24 and 36.

The patient treated with the above with Rituximab will display an improvement in any one or more of the above outcome measures.

EXAMPLE 4
HUMANIZED 2H7 VARIANTS

This example describes humanized 2H7 antibody variants for use in the methods disclosed
5 herein. The humanized 2H7 antibody preferably comprises one, two, three, four, five or six of the
following CDR sequences:

CDR L1 sequence RASSSVSYXH wherein X is M or L (SEQ ID NO. 18), for example SEQ ID
NO:4 (Fig. 1A),

CDR L2 sequence of SEQ ID NO:5 (Fig. 1A),

10 CDR L3 sequence QQWXFNPP wherein X is S or A (SEQ ID NO. 19), for example SEQ ID NO:6
(Fig. 1A),

CDR H1 sequence of SEQ ID NO:10 (Fig. 1B),

CDR H2 sequence of AIYPGNGXTSYNQFKKG wherein X is D or A (SEQ ID NO. 20), for
example SEQ ID NO:11 (Fig. 1B), and

15 CDR H3 sequence of VVYYSSXYWYFDV wherein the X at position 6 is N, A, Y, W or D, and the
X as position 7 is S or R (SEQ ID NO. 21), for example SEQ ID NO:12 (Fig. 1B).

The CDR sequences above are generally present within human variable light and variable
heavy framework sequences, such as substantially the human consensus FR residues of human light
chain kappa subgroup I (V_L6I), and substantially the human consensus FR residues of human heavy
20 chain subgroup III (V_HIII). See also WO 2004/056312 (Lowman *et al.*).

The variable heavy region may be joined to a human IgG chain constant region, wherein the
region may be, for example, IgG1 or IgG3, including native sequence and variant constant regions.

In a preferred embodiment, such antibody comprises the variable heavy domain sequence of
SEQ ID NO:8 (v16, as shown in Fig. 1B), optionally also comprising the variable light domain
25 sequence of SEQ ID NO:2 (v16, as shown in Fig. 1A), which optionally comprises one or more
amino acid substitution(s) at positions 56, 100, and/or 100a, *e.g.* D56A, N100A or N100Y, and/or
S100aR in the variable heavy domain and one or more amino acid substitution(s) at positions 32
and/or 92, *e.g.* M32L and/or S92A, in the variable light domain. Preferably, the antibody is an intact
30 antibody comprising the light chain amino acid sequences of SEQ ID NOs. 13 or 16, and heavy chain
amino acid sequences of SEQ ID NO. 14, 15, 17, 22 or 25.

A preferred humanized 2H7 antibody is ocrelizumab (Genentech).

The antibody herein may further comprise at least one amino acid substitution in the Fc
region that improves ADCC activity, such as one wherein the amino acid substitutions are at positions
298, 333, and 334, preferably S298A, E333A, and K334A, using Eu numbering of heavy chain
35 residues. See also US Patent No. 6,737,056B1, Presta.

Any of these antibodies may comprise at least one substitution in the Fc region that improves FcRn binding or serum half-life, for example a substitution at heavy chain position 434, such as N434W. See also US Patent No. 6,737,056B1, Presta.

5 Any of these antibodies may further comprise at least one amino acid substitution in the Fc region that increases CDC activity, for example, comprising at least a substitution at position 326, preferably K326A or K326W. See also US Patent No. 6,528,624B1 (Idusogie *et al.*).

Some preferred humanized 2H7 variants are those comprising the variable light domain of SEQ ID NO:2 and the variable heavy domain of SEQ ID NO:8, including those with or without substitutions in an Fc region (if present), and those comprising a variable heavy domain with 10 alteration N100A; or D56A and N100A; or D56A, N100Y, and S100aR; in SEQ ID NO:8 and a variable light domain with alteration M32L; or S92A; or M32L and S92A; in SEQ ID NO:2.

M34 in the variable heavy domain of 2H7.v16 has been identified as a potential source of antibody stability and is another potential candidate for substitution.

15 In a summary of some various preferred embodiments of the invention, the variable region of variants based on 2H7.v16 comprise the amino acid sequences of v16 except at the positions of amino acid substitutions that are indicated in the table below. Unless otherwise indicated, the 2H7 variants will have the same light chain as that of v16.

Exemplary Humanized 2H7 Antibody Variants

2H7 Version	Heavy chain (V _H) changes	Light chain (V _L) changes	Fc changes
16 for reference			-
31	-	-	S298A, E333A, K334A
73	N100A	M32L	
75	N100A	M32L	S298A, E333A, K334A
96	D56A, N100A	S92A	
114	D56A, N100A	M32L, S92A	S298A, E333A, K334A
115	D56A, N100A	M32L, S92A	S298A, E333A, K334A, E356D, M358L
116	D56A, N100A	M32L, S92A	S298A, K334A, K322A
138	D56A, N100A	M32L, S92A	S298A, E333A, K334A, K326A
477	D56A, N100A	M32L, S92A	S298A, E333A, K334A, K326A, N434W
375	-	-	K334L
588	-	-	S298A, E333A, K334A, K326A
511	D56A, N100Y, S100aR	M32L, S92A	S298A, E333A, K334A, K326A

One preferred humanized 2H7 comprises 2H7.v16 variable light domain sequence:

DIQMTQSPSSLSASVGDRVTITCRASSSVSYMHWYQQKPGKAPKPLIYAPSNLASGVPSRFSG
SGSGTDFTLTISSLQPEDFATYYCQQWSFNPPTFGQGTKVEIKR (SEQ ID NO:2);

5 and 2H7.v16 variable heavy domain sequence:

EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPNGDTSY
NQKFKGRFTISVDKSNTLYLQMNSLRAEDTAVYYCARVVYYSNSYWYFDVWGQGTLVTV
SS (SEQ ID NO:8).

Where the humanized 2H7.v16 antibody is an intact antibody, it may comprise the light chain
10 amino acid sequence:

DIQMTQSPSSLSASVGDRVTITCRASSSVSYMHWYQQKPGKAPKPLIYAPSNLASGVPSRFSG
SGSGTDFTLTISSLQPEDFATYYCQQWSFNPPTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSG
TASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSTTLSKADYEKH
KVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:13);

15 and the heavy chain amino acid sequence of SEQ ID NO. 14 or:

EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPNGDTSY
NQKFKGRFTISVDKSNTLYLQMNSLRAEDTAVYYCARVVYYSNSYWYFDVWGQGTLVTV
SSASTKGPSVFPLAPSSKSTSGGTAAAGCLVKDYFPEPVTVWSNSGALTSGVHTFPAVLQSSG
LYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVF
20 LFPPPKPKDTLMISRTPEVTCVVVDVSHEDPEVFKFNWYVDGVEVHNAKTKPREEQYNSTYRV
VSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTIASKAKGQPREPQVYTLPPSREEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCS
VMHEALHNHYTQKSLSLSPG (SEQ ID NO:22).

Another preferred humanized 2H7 antibody comprises 2H7.v511 variable light domain
25 sequence:

DIQMTQSPSSLSASVGDRVTITCRASSSVSYLHWYQQKPGKAPKPLIYAPSNLASGV
SRFSGSGSGTDFTLTISSLQPEDFATYYCQQWAFNPPTFGQGTKVEIKR (SEQ ID NO:23)

and 2H7.v511 variable heavy domain sequence:

EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPGN
30 GATSYNQKFKGRFTISVDKSNTLYLQMNSLRAEDTAVYYCARVVYYSYRYWYFDVWGQ
GTLTVVSS (SEQ ID NO. 24).

Where the humanized 2H7.v511 antibody is an intact antibody, it may comprise the light
chain amino acid sequence:

DIQMTQSPSSLSASVGDRVTITCRASSSVSYLHWYQQKPGKAPKPLIYAPSNLASGVPSRFSGS
35 GSGTDFTLTISSLQPEDFATYYCQQWAFNPPTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGT
ASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSTTLSKADYEKH
VYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:16)

and the heavy chain amino acid sequence of SEQ ID NO. 17 or:

EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPGNGATSY
NQKFKGRFTISVDKSNTLYLQMNSLRAEDTAVYYCARVYYYSYRYWYFDVWGQGTLVT
VSSASTKGPSVFPLAPSSKSTSGGTAAAGCLVKDYFPEPVTVSWNSGALTSGVHTFPALQSS
5 GLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSV
FLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTPREEQYNATYR
VVSVLTVLHQDWLNGKEYKCKVSNAALPAPIAATISKAKGQPREPQVYTLPPSREEMTKNQ
VSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFS
CSVMHEALHNHYTQKSLSLSPG (SEQ ID NO. 25).

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method of treating multiple sclerosis in a subject comprising administering an effective amount of a CD20 antibody to the subject to provide an initial antibody exposure of about 0.5 to 4 grams followed by a second antibody exposure of about 0.5 to 4 grams, the second exposure not being provided until from about 16 to 60 weeks from the initial exposure, and each of the antibody exposures is provided to the subject as one or two doses of antibody.

5 2. The method of claim 1 wherein the second exposure is not provided until from about 20 to 30 weeks from the initial exposure.

0 3. The method of claim 1 or claim 2 wherein the initial and second antibody exposures are each provided in amounts of about 1.5 to 2.5 grams.

4. The method of claim 2 or claim 3 additionally comprising administering to the subject an effective amount of the CD20 antibody to provide a third antibody exposure of about 0.5 to 4 grams, the third exposure not being administered until from about 46 to 60 weeks from the initial exposure.

5 5. The method of claim 4 wherein the third antibody exposure is provided in an amount of about 1.5 to 2.5 grams.

6. The method of claim 4 or claim 5 wherein the third exposure is not provided until from about 46 to 54 weeks from the initial exposure.

7. The method of any one of claims 4 to 6 wherein no further antibody exposure is provided until at least about 70-75 weeks from the initial exposure.

10 8. The method of claim 1 wherein the second exposure is not provided until from about 46 to 60 weeks from the initial exposure.

9. The method of any one of claims 1 to 8 wherein each of the antibody exposures is provided to the subject as one dose of antibody.

25 10. The method of any one of claims 1 to 8 wherein each of the antibody exposures is provided to the subject as two doses of antibody, wherein the two doses constitute a first dose and a second dose.

11. The method of claim 10 wherein the second dose is administered from about 3 to 17 days from the time the first dose was administered.

12. The method of claim 11 wherein the second dose is administered from about 6 to 16 days from the time the first dose was administered.

30 13. The method of claim 12 wherein the second dose is administered from about 13 to 16 days from the time the first dose was administered.

14. The method of any one of claims 10 to 13 wherein the first and second dose of antibody are each about 0.5 to 1.5 grams.

35 15. The method of any one of claims 10 to 14 wherein the first and second dose of antibody are each about 0.75 to 1.3 grams.

16. The method of any one of the claims 1 to 15 wherein three or more antibody exposures are administered to the subject.
17. The method of any one of claims 1 to 16 wherein four or more antibody exposures are administered to the subject.
- 5 18. The method of any one of claims 1 to 17 wherein a second medicament is administered with the initial exposure or later exposures, wherein the CD20 antibody is a first medicament.
19. The method of claim 18 wherein the second medicament is selected from the group consisting of an interferon, glatiramer acetate, a cytotoxic agent, chemotherapeutic agent, mitoxantrone, methotrexate, cyclophosphamide, chlorambucil, azathioprine, gamma globulin, Campath, anti-CD4, cladribine, corticosteroid, mycophenolate mofetil (MMF), cyclosporine, cholesterol-lowering drug of the statin class, estradiol, testosterone, hormone replacement drug, a TNF inhibitor, disease-modifying anti-rheumatic drug (DMARD), non-steroidal anti-inflammatory drug (NSAID), levothyroxine, cyclosporin A, somatostatin analogue, cytokine or cytokine receptor antagonist, anti-metabolite, immunosuppressive agent, integrin antagonist or antibody, LFA-1 antibody, efalizumab, alpha 4 integrin antibody, natalizumab, and another B-cell surface marker antibody.
- 15 20. The method of any one of claims 1 to 19 wherein the multiple sclerosis is relapsing-remitting multiple sclerosis (RRMS).
21. The method of any one of claims 1 to 19 wherein the multiple sclerosis is primary progressive multiple sclerosis (PPMS).
- 20 22. The method of any one of claims 1 to 21 wherein the subject has never been previously treated with a CD20 antibody.
23. The method of any one of claims 1 to 22 wherein the antibody is a naked antibody.
24. The method of any one of claims 1 to 22 wherein the antibody is conjugated with another molecule.
- 25 25. The method of claim 24 wherein the other molecule is a cytotoxic agent.
26. The method of any one of claims 1 to 25 wherein the antibody is administered intravenously.
27. The method of claim 26 wherein the antibody is administered intravenously for each antibody exposure.
28. The method of any one of claims 1 to 25 wherein the antibody is administered subcutaneously.
- 30 29. The method of any one of claims 1 to 25 wherein the antibody is administered intrathecally.
30. The method of any one of claims 1 to 29 wherein the CD20 antibody is the only medicament administered to the subject to treat the multiple sclerosis.
31. The method of any one of claims 1 to 30 wherein the antibody is Rituximab.
- 35 32. The method of any one of claims 1 to 30 wherein the antibody is humanized 2H7 comprising the variable domain sequences in SEQ ID NOS. 2 and 8.

33. The method of any one of claims 1 to 30 wherein the antibody is humanized 2H7 comprising the variable domain sequences in SEQ ID NOS. 23 and 24.
34. The method of any one of claims 1 to 33 wherein the subject has elevated anti-myelin basic protein (MBP), anti-myelin oligodendrocyte glycoprotein (MOG), anti-ganglioside and/or anti-neurofilament antibody level(s).
- 5 35. The method of any one of claims 1 to 34 wherein elevated levels of B cells are present in cerebrospinal fluid (CSF), multiple sclerosis lesion, or serum of the subject.
36. An article of manufacture when used according to the method of any one of claims 1 to 35 comprising a container comprising a CD20 antibody.
- 10 37. The article of claim 36 further comprising a container comprising a second medicament, wherein the CD20 antibody is a first medicament.
38. The article of claim 37 wherein the second medicament is selected from the group consisting of an interferon, glatiramer acetate, a cytotoxic agent, chemotherapeutic agent, mitoxantrone, methotrexate, cyclophosphamide, chlorambucil, azathioprine, gamma globulin, Campath, anti-CD4, 15 cladribine, corticosteroid, mycophenolate mofetil (MMF), cyclosporine, cholesterol-lowering drug of the statin class, estradiol, testosterone, hormone replacement drug, a TNF inhibitor, disease-modifying anti-rheumatic drug (DMARD), non-steroidal anti-inflammatory drug (NSAID), levothyroxine, cyclosporin A, somastatin analogue, cytokine or cytokine receptor antagonist, anti-metabolite, immunosuppressive agent, and another B-cell surface marker antibody.
- 20 39. Use of a CD20 antibody in the manufacture of a medicament for treating multiple sclerosis in a subject, wherein the medicament is administered to the subject to provide an initial antibody exposure of about 0.5 to 4 grams followed by a second antibody exposure of about 0.5 to 4 grams, the second exposure not being provided until from about 16 to 60 weeks from the initial exposure, and each of the antibody exposures is provided to the subject as one or two doses of antibody.
- 25 40. A method according to claim 1, an article of manufacture according to claim 36, or use according to claim 39, substantially as hereinbefore described with reference to any one of the examples or figures.

Sequence Alignment of Variable Light-Chain Domain

	FR1		CDR1		
	10	20	30	40	
2H7	Q1VLSQSPAILSASPGERVIMTC [RASSSVS-YMH] WYQQKP * * * * * * * * *				
hu2H7.v16	DIQMTQSPSSLSASVGDRVTITC [RASSSVS-YMH] WYQQKP * * * *				
hum KI	DIQMTQSPSSLSASVGDRVTITC [RASQSIISVLA] WYQQKP				
	FR2		CDR2	FR3	
	50	60	70	80	
2H7	GSSPKPKWIY [APSNLAS] GVPARPSGSGSGTYSLTISRVEA * * * * * * * * * * * * * *				
hu2H7.v16	GKAPKPLIY [APSNLAS] GVPSRPSGSGSGTDFTLTISSLQP * * * *				
hum KI	GKAPKLLIY [AASSLES] GVPSRPSGSGSGTDFTLTISSLQP				
	CDR3		FR4		
	90	100			
2H7	EDEATYYC [QQWSFNPPT] FGAGTKLEIKR * * * *				
hu2H7.v16	EDEATYYC [QQWSFNPPT] FGQGTTKVEIKR * * * *				
hum KI	EDEATYYC [QQVNSLPWT] FGQGTTKVEIKR				

Sequence Alignment of Variable Heavy-Chain Domain

	FR1		CDR1		
	10	20	30	40	
2H7	QAYLQQSGAELVVRPGASVMSCKAS [GYTFTSYNMH] WVKQD * * * * * * * * * * * * *				
hu2H7.v16	EVQLVESGGGLVQPGGSLRLSCAAS [GYTFTSYNMH] WVRQA * * * *				
hum III	EVQLVESGGGLVQPGGSLRLSCAAS [GFTFSSYAMS] WVRQA				
	FR2		CDR2	FR3	
	50	a	60	70	80
2H7	PRQGLEWIG [AIYPGNGDTSYNQKFKG] KATETVDKSSSTAYM * * * * * * * * * * * * *				
hu2H7.v16	PGKGLEWVG [AIYPGNGDTSYNQKFKG] RFTISVDKSKNTLYL * * * * * * * * * * * * *				
hum III	PGKGLEWVA [VISGDDGSTYVADSVKG] RPTISGRDNNSKNTLTL				
	CDR3		FR4		
	abc	90	100	abcde	110
2H7	Q1SSLTSEDSAVYPCAR [VVVYSNSYWYFDV] WGTGTTVTVSS * * * * * * * * *				
hu2H7.v16	QMNSLRAEDTAVYYCAR [VVVYSNSYWYFDV] WGQGTLVTVSS * * * * * * * * *				
hum III	QMNSLRAEDTAVYYCAR [GRVGYSLY---DY] WGQGTLVTVSS				

Sequence Alignment of Variable Heavy-Chain Domain

				FR1					CDR1							
				10	20					30	40					
2H7	QAYLQQSGAELVRPGASVKMSCKAS [GYTFTSYNMH] WVKQT															
	***	***	*	*	***	*				*	*					
hu2H7.v16	EVQLVESGGGLVQPGGSLRLSCAAS [GYTFTSYNMH] WVRQA															
							*	*	*	*						
hum III	EVQLVESGGGLVQPGGSLRLSCAAS [GFTFSSYAMS] WVRQA															
				FR2	CDR2				FR3							
				50	a	60					70	80				
2H7	PROQGLEWIG [AIYPGNGDTSYNQFKKG] KATLTVDKSSSTAYM															
	**	*					**	**		**	*					
hu2H7.v16	PGKGLEWVG [AIYPGNGDTSYNQFKKG] RFTISVDKSKNTLYL															
	*	*	****	*	*	****			*	*						
hum III	PGKGLEWVA [VISGDDGSTYYADSVKG] RFTISRDNSKNTLTL															
				FR4	CDR3				FR4							
				abc	90	100	abcde					110				
2H7	QLSSLTSEDSAVYFCAR [VVVYSNSYWYFDV] WGTGTTVTVSS															
	**	**	*	*					*	*						
hu2H7.v16	QMNSLRAEDTAVYYCAR [VVVYSNSYWYFDV] WGQGTLVTVSS															
	*****	*****	*	*												
hum III	QMNSLRAEDTAVYYCAR [GRVGYSLY---DY] WGQGTLVTVSS															

Humanized 2H7.v16 Light Chain

DIQMTQSPSSLSASVGDRVТИTCAASSVSYMHYQQKPGKAPKPLIYAPSNLASGVPSRFSG
SGSGTDFTLTISSLQPEDFATYYCQQWSFNPPTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKS
GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSLTLSKADYEKHK
VYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:13)

FIG._2**Humanized 2H7.v16 Heavy Chain**

EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPNGDTSYNQK
FKGRFTISVDKSNTLYLQMNSLRAEDTAVYYCARVYYYSNSYWYFDVWGQGTLTVSSASTK
GPSVFLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSS
VVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPCPAPELLGGPSVFLFPPKP
KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTPREEQYNSTYRVSVLTVLH
QDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREEMTKNQVSLTCLVKGFY
PSDIAVEWESNGQOPENNYKTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHY
TQKSLSLSPGK (SEQ ID NO:14)

FIG._3**Humanized 2H7.v31 Heavy Chain**

EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHWVRQAPGKGLEWVGAIYPNGDTSYNQK
FKGRFTISVDKSNTLYLQMNSLRAEDTAVYYCARVYYYSNSYWYFDVWGQGTLTVSSASTK
GPSVFLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSS
VVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPCPAPELLGGPSVFLFPPKP
KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTPREEQYNATYRVSVLTVLH
QDWLNGKEYKCKVSNKALPAPIATISKAKGQPREPVYTLPPSREEMTKNQVSLTCLVKGFY
PSDIAVEWESNGQOPENNYKTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHY
TQKSLSLSPGK (SEQ ID NO:15)

FIG._4

Light Chain Alignment

	1	32
hu2H7.v16	DIQMTQSPSSLSASVGDRVТИTCRASSSVSYMHYQQKPGKAPKPLIYAP	*****
hu2H7.v511	DIQMTQSPSSLSASVGDRVТИTCRASSSVSYLHWYQQKPGKAPKPLIYAP	*****
	52	
hu2H7.v16	SNLASGVPSRSGSGTDFTLTISSSLQPEDFATYYCQQWSFNPPFGQG	*****
hu2H7.v511	SNLASGVPSRSGSGTDFTLTISSSLQPEDFATYYCQQWAFNPPTFGQG	*****
	102	
hu2H7.v16	TKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVD	*****
hu2H7.v511	TKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVD	*****
	152	
hu2H7.v16	NALQSGNSQESVTEQDSKDSTYSLSSLTLSKADYEHKVYACEVTHQGL	*****
hu2H7.v511	NALQSGNSQESVTEQDSKDSTYSLSSLTLSKADYEHKVYACEVTHQGL	*****
	202	214
hu2H7.v16	SSPVTKSFNRGEC	*****
hu2H7.v511	SSPVTKSFNRGEC	

FIG._5

Heavy Chain Alignment

hu2H7.v16	1	EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHW *****	
hu2H7.v511		EVQLVESGGGLVQPGGSLRLSCAASGYTFTSYNMHW	
hu2H7.v16	37	52a	82abc
		VRQAPGKGLEWVGAIYPGNGDTSYNQFKGRFTISVDKSNTLYLQMNSL *****	*****
hu2H7.v511		VRQAPGKGLEWVGAIYPGNGATSYNQFKGRFTISVDKSNTLYLQMNSL	
hu2H7.v16	83	100abcde	113
		RAEDTAVYYCARVYYYSNSYWYFDVWGQGTLVTVSS *****	*****
hu2H7.v511		RAEDTAVYYCARVYYYSYRYWYFDVWGQGTLVTVSS	
hu2H7.v16	118	ASTKGPSVFPLAPS	

hu2H7.v511		ASTKGPSVFPLAPS	
hu2H7.v16	132	SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS *****	*****
hu2H7.v511		SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS	
hu2H7.v16	182	LSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPCPA *****	*****
hu2H7.v511		LSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPCPA	
hu2H7.v16	232	PELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG *****	*****
hu2H7.v511		PELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG	
hu2H7.v16	282	VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAP *****	*****
hu2H7.v511		VEVHNAKTKPREEQYNATYRVVSVLTVLHQDWLNGKEYKCKVSNALPAP	
hu2H7.v16	332	IEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEW *	*****
hu2H7.v511		IAATISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEW	
hu2H7.v16	382	ESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCVMHEA *****	*****
hu2H7.v511		ESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCVMHEA	
hu2H7.v16	432	447	
		LHNHYTQKSLSLSPGK *****	
hu2H7.v511		LHNHYTQKSLSLSPGK	

FIG.-6

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)
Sequence Listing

<110> GENENTECH, INC.
FROHNA, PAUL A.

<120> METHOD FOR TREATING MULTIPLE SCLEROSIS

<130> P2134R1 PCT

<140> PCT/US2005/019641
<141> 2005-06-02

<150> US 60/576,993
<151> 2004-06-04

<160> 25

<210> 1
<211> 107
<212> PRT
<213> *Mus musculus*

<400> 1
Gln Ile Val Leu Ser Gln Ser Pro Ala Ile Leu Ser Ala Ser Pro
1 5 10 15
Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser Ser Ser Val Ser
20 25 30
Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Ser Ser Pro Lys Pro
35 40 45
Trp Ile Tyr Ala Pro Ser Asn Leu Ala Ser Gly Val Pro Ala Arg
50 55 60
Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser
65 70 75
Arg Val Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp
80 85 90
Ser Phe Asn Pro Pro Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu
95 100 105
Lys Arg

<210> 2
<211> 107
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized.

<400> 2
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
1 5 10 15
Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Ser Val Ser
20 25 30
Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro
35 40 45
Leu Ile Tyr Ala Pro Ser Asn Leu Ala Ser Gly Val Pro Ser Arg
50 55 60

06 Feb 2009

2005249566

P62780.AU P2134R1(SEQLISTING REVISED)
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75
Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp
80 85 90
Ser Phe Asn Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
95 100 105
Lys Arg

<210> 3
<211> 108
<212> PRT
<213> Artificial Sequence

<220>
<223> Sequence is synthesized.

<400> 3
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
1 5 10 15

Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser
20 25 30

Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
35 40 45

Leu Leu Ile Tyr Ala Ala Ser Ser Leu Glu Ser Gly Val Pro Ser
50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
65 70 75

Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
80 85 90

Tyr Asn Ser Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu
95 100 105

Ile Lys Arg

<210> 4
<211> 10
<212> PRT
<213> Mus musculus

<400> 4
Arg Ala Ser Ser Ser Val Ser Tyr Met His
5 10

<210> 5
<211> 7
<212> PRT
<213> Mus musculus

<400> 5
Ala Pro Ser Asn Leu Ala Ser
5

<210> 6
<211> 9
<212> PRT
<213> Mus musculus

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)

<400> 6
Gln Gln Trp Ser Phe Asn Pro Pro Thr
5

<210> 7

<211> 122

<212> PRT

<213> Mus musculus

<400> 7

Gln Ala Tyr Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Pro Gly
1 5 10 15

Ala Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr
20 25 30

Ser Tyr Asn Met His Trp Val Lys Gln Thr Pro Arg Gln Gly Leu
35 40 45

Glu Trp Ile Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr
50 55 60

Asn Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser
65 70 75

Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp
80 85 90

Ser Ala Val Tyr Phe Cys Ala Arg Val Val Tyr Tyr Ser Asn Ser
95 100 105

Tyr Trp Tyr Phe Asp Val Trp Gly Thr Gly Thr Thr Val Thr Val
110 115 120

Ser Ser

<210> 8

<211> 122

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

<400> 8

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly
1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
20 25 30

Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45

Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr
50 55 60

Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Val Asp Lys Ser
65 70 75

Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90

Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr Tyr Ser Asn Ser
95 100 105

Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val

2005249566 06 Feb 2009

110 P62780.AU P2134R1(SEQLISTING REVISED)
115
120

Ser Ser

<210> 9
<211> 119

<212> PRT

<213> Artificial sequence

<220>
<223> Sequence is synthesized.

<400> 9

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly
1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
20 25 30

Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45

Glu Trp Val Ala Val Ile Ser Gly Asp Gly Gly Ser Thr Tyr Tyr
50 55 60

Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser
65 70 75

Lys Asn Thr Leu Thr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90

Thr Ala Val Tyr Tyr Cys Ala Arg Gly Arg Val Gly Tyr Ser Leu
95 100 105

Tyr Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
110 115

<210> 10

<211> 10

<212> PRT

<213> Mus musculus

<400> 10

Gly Tyr Thr Phe Thr Ser Tyr Asn Met His
5 10

<210> 11

<211> 17

<212> PRT

<213> Mus musculus

<400> 11

Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr Asn Gln Lys Phe
1 5 10 15

Lys Gly

<210> 12

<211> 13

<212> PRT

<213> Mus musculus

<400> 12

Val Val Tyr Tyr Ser Asn Ser Tyr Trp Tyr Phe Asp Val
5 10

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)

<210> 13
<211> 213
<212> PRT

<213> Artificial sequence

<220>
<223> Sequence is synthesized.

<400> 13
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
1 5 10 15
Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Ser Val Ser
20 25 30
Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro
35 40 45
Leu Ile Tyr Ala Pro Ser Asn Leu Ala Ser Gly Val Pro Ser Arg
50 55 60
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75
Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp
80 85 90
Ser Phe Asn Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
95 100 105
Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser
110 115 120
Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu
125 130 135
Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp
140 145 150
Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln
155 160 165
Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu
170 175 180
Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val
185 190 195
Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg
200 205 210
Gly Glu Cys

<210> 14
<211> 452
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized.

<400> 14
Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly
1 5 10 15
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
20 25 30

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)

Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45
Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr
50 55 60
Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Val Asp Lys Ser
65 70 75
Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90
Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr Tyr Ser Asn Ser
95 100 105
Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val
110 115 120
Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro
125 130 135
Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu
140 145 150
Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
155 160 165
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
170 175 180
Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
185 190 195
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
200 205 210
Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys
215 220 225
Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
230 235 240
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
245 250 255
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
260 265 270
Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
275 280 285
Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
290 295 300
Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
305 310 315
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
320 325 330
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys
335 340 345
Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
350 355 360
Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)
365 370 375

Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
380 385 390

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
395 400 405

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
410 415 420

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
425 430 435

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
440 445 450

Gly Lys

<210> 15

<211> 452

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

<400> 15

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly
1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
20 25 30

Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45

Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr
50 55 60

Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Val Asp Lys Ser
65 70 75

Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90

Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr Tyr Ser Asn Ser
95 100 105

Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val
110 115 120

Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro
125 130 135

Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu
140 145 150

Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
155 160 165

Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
170 175 180

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
185 190 195

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
200 205 210
Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys
215 220 225
Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
230 235 240
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
245 250 255
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
260 265 270
Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
275 280 285
Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
290 295 300
Tyr Asn Ala Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
305 310 315
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
320 325 330
Lys Ala Leu Pro Ala Pro Ile Ala Ala Thr Ile Ser Lys Ala Lys
335 340 345
Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
350 355 360
Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
365 370 375
Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
380 385 390
Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
395 400 405
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
410 415 420
Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
425 430 435
Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
440 445 450
Gly Lys

<210> 16
<211> 213
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized.

<400> 16
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
1 5 10 15
Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Ser Val Ser
20 25 30

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)

Tyr Leu His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro
35 40 45
Leu Ile Tyr Ala Pro Ser Asn Leu Ala Ser Gly Val Pro Ser Arg
50 55 60
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75
Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp
80 85 90
Ala Phe Asn Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
95 100 105
Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser
110 115 120
Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu
125 130 135
Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp
140 145 150
Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln
155 160 165
Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu
170 175 180
Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val
185 190 195
Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg
200 205 210
Gly Glu Cys

<210> 17

<211> 452

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

<400> 17

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly
1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
20 25 30

Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45

Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Ala Thr Ser Tyr
50 55 60

Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Val Asp Lys Ser
65 70 75

Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90

Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr Tyr Ser Tyr Arg

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)
95 100 105
Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val
110 115 120
Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro
125 130 135
Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu
140 145 150
Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
155 160 165
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
170 175 180
Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
185 190 195
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
200 205 210
Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys
215 220 225
Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
230 235 240
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
245 250 255
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
260 265 270
Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
275 280 285
Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
290 295 300
Tyr Asn Ala Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
305 310 315
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
320 325 330
Ala Ala Leu Pro Ala Pro Ile Ala Ala Thr Ile Ser Lys Ala Lys
335 340 345
Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
350 355 360
Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
365 370 375
Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
380 385 390
Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
395 400 405
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
410 415 420
Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
425 430 435

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
440 445 450

Gly Lys

<210> 18

<211> 10

<212> PRT

<213> Artificial sequence

<220>

<223> sequence is synthesized

<220>

<221> xaa

<222> 9

<223> xaa is M or L

<400> 18

Arg Ala Ser Ser Ser Val Ser Tyr Xaa His
5 10

<210> 19

<211> 9

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

<220>

<221> xaa

<222> 4

<223> xaa is S or A

<400> 19

Gln Gln Trp Xaa Phe Asn Pro Pro Thr
5

<210> 20

<211> 17

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

<220>

<221> xaa

<222> 8

<223> xaa is D or A

<400> 20

Ala Ile Tyr Pro Gly Asn Gly xaa Thr Ser Tyr Asn Gln Lys Phe
1 5 10 15

Lys Gly

<210> 21

<211> 13

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)

<220>
<221> xaa
<222> 6
<223> xaa is N, A, Y, W or D

<220>
<221> xaa
<222> 7
<223> xaa is S or R

<400> 21
Val Val Tyr Tyr Ser Xaa Xaa Tyr Trp Tyr Phe Asp Val
5 10

<210> 22
<211> 451
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized.

<400> 22
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
1 5 10 15
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
20 25 30
Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45
Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr
50 55 60
Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Val Asp Lys Ser
65 70 75
Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90
Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr Tyr Ser Asn Ser
95 100 105
Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val
110 115 120
Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro
125 130 135
Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu
140 145 150
Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
155 160 165
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
170 175 180
Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
185 190 195
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
200 205 210
Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys
215 220 225

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)
Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
230 235 240
Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
245 250 255
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
260 265 270
Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
275 280 285
Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
290 295 300
Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
305 310 315
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
320 325 330
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys
335 340 345
Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
350 355 360
Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
365 370 375
Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
380 385 390
Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
395 400 405
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
410 415 420
Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
425 430 435
Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
440 445 450
Gly

<210> 23
<211> 107
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized.

<400> 23
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
1 5 10 15
Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ser Ser Val Ser
20 25 30
Tyr Leu His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Pro
35 40 45
Leu Ile Tyr Ala Pro Ser Asn Leu Ala Ser Gly Val Pro Ser Arg
50 55 60

P62780.AU P2134R1(SEQLISTING REVISED)

Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75

Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Trp
80 85 90

Ala Phe Asn Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
95 100 105

Lys Arg

<210> 24

<211> 122

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

<400> 24

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly
1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
20 25 30

Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45

Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Ala Thr Ser Tyr
50 55 60

Asn Gln Lys Phe Lys Gly Arg Phe Thr Ile Ser Val Asp Lys Ser
65 70 75

Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90

Thr Ala Val Tyr Tyr Cys Ala Arg Val Val Tyr Tyr Ser Tyr Arg
95 100 105

Tyr Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val
110 115 120

Ser Ser

<210> 25

<211> 451

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

<400> 25

Glu Val Gln Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly
1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Thr Phe Thr
20 25 30

Ser Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45

Glu Trp Val Gly Ala Ile Tyr Pro Gly Asn Gly Ala Thr Ser Tyr
Page 14

P62780.AU P2134R1(SEQLISTING REVISED)

50	55	60												
Asn	Gln	Lys	Phe	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Val	Asp	Lys	Ser
65									70					75
Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp
80									85					90
Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Val	Val	Tyr	Tyr	Ser	Tyr	Arg
95									100					105
Tyr	Trp	Tyr	Phe	Asp	Val	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val
110									115					120
Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro
125									130					135
Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu
140									145					150
Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
155									160					165
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln
170									175					180
Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
185									190					195
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys
200									205					210
Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys
215									220					225
Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu
230									235					240
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr
245									250					255
Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp
260									265					270
Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp
275									280					285
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln
290									295					300
Tyr	Asn	Ala	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His
305									310					315
Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn
320									325					330
Ala	Ala	Leu	Pro	Ala	Pro	Ile	Ala	Ala	Thr	Ile	Ser	Lys	Ala	Lys
335									340					345
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg
350									355					360
Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys
365									370					375
Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly
380									385					390

2005249566 06 Feb 2009

P62780.AU P2134R1(SEQLISTING REVISED)
Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
395 400 405
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
410 415 420
Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
425 430 435
Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
440 445 450
Gly