

COMMONWEALTH OF AUSTRALIA

Patents Act 1952

599985

CONVENTION APPLICATION FOR A PATENT

X/We, MEIJI SEIKA KAISHA, LTD.,

of 4-16, Kyobashi 2-chome, Chuo-ku, Tokyo, Japan

hereby apply for the grant of a Patent
for an invention entitled:-

"NEW PROCESS FOR THE PRODUCTION OF L-2-AMINO-4-(HYDROXYMETHYL-
PHOSPHINYL)-BUTYRIC ACID"

which is described in the accompanying complete specification.

This application is a Convention application and is based on
the application numbered 131743/86 for a patent or
similar protection made in Japan

on 9th June, 1986, the application numbered 12513/87 for a
patent or similar protection made in Japan on 23rd January,
1987, the application numbered 74476/87 for a patent or
similar protection made in Japan on 30th March, 1987, and
the application numbered 10152/87 for a patent or similar
protection made in Japan on 25th April, 1987.

X/Our address for service is: T.G. AHEARN & CO. Patent
Attorneys, of 79 Eagle Street, Brisbane, in
the State of Queensland, Commonwealth of Australia.

Dated this 2nd day of June, 1987.

APPLICATION ACCEPTED AND AMENDMENTS

MEIJI SEIKA KAISHA, LTD.
By their Patent Attorneys

ALLOWED

18.5.90

G.R. CULLEN & CO.

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952

DECLARATION IN SUPPORT OF A

CONVENTION APPLICATION FOR PATENT

In support of the Convention Application made for a patent
for an invention entitled

NEW PROCESS FOR THE PRODUCTION OF L-2-AMINO-4-(HYDROXYMETHYL-
PHOSPHINYL)-BUTYRIC ACID

I, Akira SASAI, President,
of MEIJI SEIKA KAISHA, LTD, 4-16, Kyobashi 2-chome, Chuo-
ku, Tokyo, Japan,

do solemnly and sincerely declare as follows:-

1. We are authorised by MEIJI SEIKA KAISHA, LTD, the
applicant, for the patent to make this declaration on its
behalf.

2. The basic applications as defined by Section 141 of
the Act was made in JAPAN on 9 JUNE 1986, 23 JANUARY 1987, 30
MARCH 1987 and 25 APRIL 1987 by MEIJI SEIKA KAISHA, LTD.

3. SATOSHI IMAI, NOBUHIKO TAKANE, YACHIYO YOSHIZAWA,
TOSHINORI SAITO, HIROSHI OGAWA, HIDEHI TAKEBE, ATSUYUKI SATO,
SHUNZO FUKATSU and AKIRA OKADA, all C/o MEIJI SEIKA KAISHA, LTD,
Pharmaceutical Development Laboratories, 580 Horikawa-cho,
Saiwai-ku, Kawasaki-shi, Kanagawa-ken, Japan and TAKESHI
MURAKAMI, OSAMU HARA, SHINJI MIYADO, YOICHI KUMADA, HIROYUKI
ANZAI and KOZO NAGAOKA all C/o MEIJI SEIKA KAISHA, LTD,
Pharmaceutical Research Laboratories, 760 Morooka-cho, Kohhoku-
ku, Yokohama-shi, Kanagawa-ken, Japan, are the actual inventors
of the invention and the facts upon which the applicant is
entitled to make the application is as follows:

MEIJI SEIKA KAISHA, LTD is the assignee of the said
SATOSHI IMAI, NOBUHIKO TAKANE, YACHIYO YOSHIZAWA, TOSHINORI
SAITO, HIROSHI OGAWA, HIDEHI TAKEBE, ATSUYUKI SATO, SHUNZO
FUKATSU, AKIRA OKADA, TAKESHI MURAKAMI, OSAMU HARA, SHINJI
MIYADO, YOICHI KUMADA, HIROYUKI ANZAI and KOZO NAGAOKA

4. The basic application referred to in paragraph 2 of
this Declaration was the first application made in a Convention
country in respect of the invention the subject of this
application.

DECLARED at TOKYO this 13th day of March 1990.

MEIJI SEIKA KAISHA, LTD.


.....*Akira Sasai*.....
Signature of Declarant
Akira SASAI, President

TO: THE COMMISSIONER OF PATENTS
AUSTRALIA.


(12) PATENT ABRIDGMENT (11) Document No. AU-B-73750/87
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 599985

- (54) Title
PRODUCTION OF 2-AMINO-4-(HYDROXYMETHYLPHOSPHINYL)-BUTYRIC ACID
- International Patent Classification(s)
(51)⁴ C12P 013/04
- (21) Application No. : 73750/87 (22) Application Date : 02.06.87
- (30) Priority Data
- | (31) Number | (32) Date | (33) Country |
|-------------|-----------|--------------|
| 61-131743 | 09.06.86 | JP JAPAN |
| 62-12513 | 23.01.87 | JP JAPAN |
| 62-74476 | 30.03.87 | JP JAPAN |
| 62-101152 | 25.04.87 | JP JAPAN |
- (43) Publication Date : 10.12.87
- (44) Publication Date of Accepted Application : 02.08.90
- (71) Applicant(s)
MEIJI SEIKA KAISHA, LTD.
- (72) Inventor(s)
SATOSHI IMAI; TAKESHI MURAKAMI; OSAMU HARA; SHINJI MIYADO; YOICHI KUMADA; HIROYUKI ANZAI; NOBUHIKO TAKANE; YACHIYO YOSHIZAWA; TOSHINORI SAITO; HIROSHI OGAWA; HIDEHI TAKEBE; ATSUYUKI SATO; KOZO NAGAOKA; SHUNZO FUKATSU; AKIRA OKADA
- (74) Attorney or Agent
CULLEN & CO. BRISBANE
- (57) Claim

1. A process for the production of L-2-amino-4-(hydroxymethylphosphinyl)-butyric acid represented by the formula (I):



which comprises treating 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid represented by the formula (II):



with one or more transaminases or with one or more

(11) AU-B-73750/87

-2-

(10) 599985

microorganisms capable of producing one or more
transaminases, in the presence of one or more amino-
donors.

599985

This document contains the
amendments made under
Section 49 and is correct for
printing.

COMMONWEALTH OF AUSTRALIA

Patents Act 1952

Name of Applicant: MEIJI SEIKA KAISHA, LTD.

Address of Applicant: 4-16, Kyobashi 2-chome,
Chuo-ku,
Tokyo,
Japan.

Actual Inventors: Satoshi IMAI
Takeshi MURAKAMI
Osamu HARA
Shinji MIYADO
Yoichi KUMADA
Hiroyuki ANZAI
Nobuhiko TAKANE
Yachiyo YOSHIZAWA
Toshinori SAITO
Hiroshi OGAWA
Hidehi TAKEBE
Atsuyuki SATO
Kozo NAGAOKA
Shunzo FUKATSU
Akira OKADA

Address for Service: T.G. AHEARN & CO.,
Patent and Trade Mark Attorneys,
79 Eagle Street,
Brisbane, Queensland.

COMPLETE SPECIFICATION FOR THE INVENTION ENTITLED:-

"NEW PROCESS FOR THE PRODUCTION OF L-2-AMINO-4-(HYDROXYMETHYL-
PHOSPHINYL)-BUTYRIC ACID"


The following statement is a full description of
the invention including the best method of performing it
known to us:-

Summary of the Invention

This invention relates to a new process for the production of L-2-amino-4-(hydroxymethylphosphinyl)-butyric acid which exhibits herbicidal activities and is known to be useful as a herbicide (see Japanese patent publication No. 56210/86 specification or U.S. patent No. 4,265,654 specification).

Background of the Invention

The known methods for the production of L-2-amino-4-(hydroxymethylphosphinyl)-butyric acid (hereinafter abbreviated as "L-AMPB") represented by the formula (I):



include such method wherein an antibiotic substance SF-1293 namely, L-2-amino-4-(hydroxymethylphosphinyl)-butyryl-L-alanyl-L-alanine, which contains L-AMPB as a moiety of the SF-1293 molecule and has herbicidal activities (also called "bialaphos"; see Japanese Patent Publication No. 639/76 and U.S. patent No. 4,309,208) is subjected to hydrolysis (see Japanese Patent Application first publication "Kokai" No. 85538/73), and such method wherein the SF-1293 substance is decomposed with a microbial enzyme (see Japanese Patent Application first publication "Kokai" No. 31890/74). Besides, there has also been

known a further method wherein AMPB in the form of a racemic mixture is at first produced by a chemical synthetic process (see Japanese Patent Application first publication "Kokai" Nos. 91019/73 and 84529/79) and then
5 the racemic AMPB product is subjected to optical resolution with aid of a microbial enzyme to yield L-AMPB. In addition, there has been known another method as reported recently by the present inventors, wherein an L-AMPB producing strain of the genus Streptomyces is cultivated, followed
10 by recovering L-AMPB directly from the resulting culture broth (see Japanese Patent Application first publication "Kokai" No. 47485/82).

In order that a substance having herbicidal activities such as L-AMPB is produced, investigated and developed as
15 a herbicide and then put actually on the market as a commercial herbicide product, some investigation needs to be made indispensably to improve the process of producing the herbicidal substance so as to make said process suitable for the low-price and large-scale production, while making
20 considerable researches and development works for enhancing the safety and the herbicidal effects of said substance. The AMPB as produced by the above-described chemical synthetic process is a mixture of L-AMPB and D-AMPB. However, D-AMPB itself substantially lacks herbicidal
25 activities. Moreover, D-AMPB is a non-natural substance

and when applied to the soil, its decomposition by soil bacteria is slow so that it tends to remain in the soil and may cause environmental pollution in some instances.

When L-AMPB is to be produced by the synthetic process, AMPB can

5 be first produced in the form of a racemic mixture. It is hence necessary to isolate L-AMPB and D-AMPB separately from each other out of the racemic mixture. The synthetic process is therefore cumbersome and the yield of L-AMPB is low. In contrast, the process of producing L-AMPB which makes
10 use of the microorganism or microbial enzyme can provide exclusively L-AMPB which is a naturally occurring substance. The naturally-occurring L-AMPB is considered to be an ideal herbicide which is free of the danger of environmental pollution, since such portions of the L-AMPB which have
15 taken no part in effecting the herbicidal effects and are still remaining in the soil are easily decomposed and metabolized by soil bacteria and are hence not allowed to remain in the soil for long duration.


With the foregoing in view, we, the present inventors,
20 have made researches in an attempt to combine the merit of the chemical synthetic process of producing AMPB that AMPB can be produced in a large scale, with the advantage of the microbiological process of producing L-AMPB that L-AMPB can exclusively be produced, so that an improved
25 process for the production of L-AMPB capable of producing

L-AMPB selectively in a large scale will be established.


L-AMPB can be deemed as a sort of derivatives of α -amino acid. Regarding such usual α -amino acids which normally constitute proteins, it has been known that α -amino acids can generally be formed by transamination of their corresponding 2-oxo-acids under the action of specific transaminases. We, the present inventors, have therefore been interesting in a 2-oxo-acid corresponding to L-AMPB, and have made an extensive investigation on such 2-oxo-acid. As a result, it has unexpectedly been discovered that when 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid (hereinafter abbreviated as OMPB) is treated with a certain sort of transaminase or a microorganism capable of producing such transaminase, in the presence of at least one amino donor, OMPB can be converted into L-AMPB in a significant yield for a reasonable reaction time.

Detailed Description of the Invention

According to this invention, therefore, there is provided a process for the production of L-2-amino-4-(hydroxymethylphosphinyl)-butyric acid represented by the formula (I):



which comprises treating 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid represented by the formula (II):



with one or more transaminases or with one or more microorganisms capable of producing one or more transaminases, in the presence of one or more amino-donors.

5 Upon carrying out the process of this invention, the transaminase or the microorganism capable of producing transaminase is caused to react with OMPB and the amino-donor compound in an aqueous liquid reaction medium which contains OMPB and the compound capable of acting as the amino-donor
10 dissolved therein.

 In the process of this invention, it may generally be preferable to conduct the conversion reaction of OMPB into L-AMPB while adjusting the reaction medium at a pH in a range of 7.5 or higher, preferably, within a range of
15 8.0-9.0. The pH adjustment can be effected by adding sodium hydroxide or a suitable buffer solution. Desirable reaction conditions may be set so that the temperature and pH of the reaction medium fall within such temperature and pH ranges optimal for the action of the transaminase
20 or the transaminase-producing microorganism participating in the reaction. Usually, it is preferable to conduct the reaction at a temperature in a range of from room temperature to 60°C preferably of 25°C to 50°C.

The OMPB used as the starting compound is a known substance. The production and physicochemical properties of OMPB are described, for example, in Japanese Patent Application first publication "Kokai" No. 92897/81 or U.S. patent No. 4,399,287 specification. In the process of this invention, the starting OMPB compound is usually dissolved in the reaction medium at an initial OMPB concentration preferably within a range of 0.10-100 mg/ml. at the beginning of the reaction.

10 The microorganisms used in the process of this invention may include actinomycetes, bacteria, yeasts and fungi (molds). As exemplary actinomycetes useful in the present invention, may be mentioned Streptomyces albus, Streptomyces griseus, Streptomyces hygroscopicus,
15 Streptomyces lividans, Streptomyces viridochromogenes, Streptomyces morookaensis, Streptoverticillium cinnamoneum, Nocardia mediterranei, Nocardiosis dassonvillei, Saccharopolyspora hirsuta, Kitasatospora phosalacinea, Micromonospora carbonacea, Streptosporangium pseudovulgare, and
20 so on.

Illustrative examples of the bacteria useful in the process of this invention may include Bacillus subtilis, Micrococcus luteus, Staphylococcus aureus, Escherichia coli, Pseudomonas aeruginosa, Pseudomonas cepacia, Serratia marcescens, Corynebacterium glutamicum,
25 and others.

As exemplary yeasts useful in the process of the present invention, may be mentioned Saccharomyces cerevisiae, Candida albicans, Cryptococcus neoformans, Debaryomyces hansenii, Trigonopsis variabilis, Hansenula
5 schneggi, and the like.

As illustrative examples of the fungi useful in the process of this invention, may be mentioned Aspergillus flavus, Aspergillus terreus, Mucor
spinescens, Aureobasidium pullulans, Chaetomium
10 globosum, Penicillium funiculosum, Gliocladium vireus,
etc.

The above-mentioned species of actinomycetes, bacteria, fungi and yeasts may be their Type Culture strains which have been deposited and stored in the well-
15 known public depositories of microorganisms and are now
available freely from these depositories.

Preferable examples of the microorganism available in the process of the process of this invention may include Streptomyces hygroscopicus SF-1293 strain (FERM BP-130 or
20 ATCC 21705; see Japanese patent publication No. 639/76 or
U.S. Patent No. 3,832,394 specification), which has been
known as an SF-1293 substance-producing strain of
Streptomyces, and its mutant strain, Streptomyces
hygroscopicus NP-50 strain (FERM P-7804 or FERM BP-1368;
25 see Japanese Patent Application first publication "Kokai"

No. 58589/86 or European patent application publication No. 0 173 327) as well as Streptomyces lividans 66 strain (FERM BP-737; see Japanese Patent Application first publication "Kokai" No. 175889/84 or European patent application publication No. 0 196 375 specification).

In addition, any other microorganisms can also be used so long as they can each produce an enzyme having the transaminase activity capable of converting OMPB into L-AMPB in the presence of the amino-donor. Microbiological

characteristics of the Streptomyces hygrosopicus SF-1293 strain are described in Japanese Patent Publication No. 639/76 or U.S. Patent No. 3,832,394 specification. The Streptomyces hygrosopicus NP-50 strain (FERM BP-1368)

has the same microbiological characteristics as those of the aforesaid SF-1293 strain but the genetic character of the NP-50 strain is different from that of the SF-1293 strain in that the NP-50 strain is lacking the biosynthetic ability of producing the SF-1293 substance (see Japanese patent Application first publication "Kokai" No. 58589/86

or European patent application publication No. 0 173 327).

Further, the microbiological characteristics of Streptomyces lividans 66 strain (FERM BP-737) are described in the

Japanese patent application first publication "Kokai"

No. 175889/84. Among the above-mentioned microbial species,

such those having the FERM P-numbers have been deposited

and stored in a Japanese public depository, the "Fermentation Research Institute, Agency of Industrial Science & Technology, Ministry of International Trade and Industry", of 1-3, Higashi 1-chome, Yatabe-machi, Tsukuba-gun, Ibaraki-ken, Japan, and such those having the FERM BP-number have been deposited and stored in the same Japanese public depository under the Butapest Treaty.

The enzyme useful in the process of this invention, i.e., the transaminase may be any transaminase so long as it possesses the transaminase activity capable of converting OMPB into L-AMPB in the presence of the amino-donor.

Preferred examples of the transaminase useful in the process of this invention may include a commercially available glutamic acid-oxaloacetic acid-transaminase (usually abbreviated as GOT) (International Enzyme Classification Number EC 2,6,1,1) and a commercially available glutamic acid-pyruvic acid-transaminase (usually abbreviated as GPT) (International Enzyme Classification Number EC 2,6,1,2), as well as such enzymes having the transaminase activity capable of converting OMPB into L-AMPB in the presence of L-glutamic acid as the amino-donor. These transaminases may be used either singly or in combination of two or more of them. Of these, a preferred combination of the transaminases is a combination or system of a transaminase also having the enzymatic activity of

converting 2-ketoglutaric acid into glutamic acid in the presence of aspartic acid as the amino-donor, namely such transaminase also having so-called GOT activity, and a second enzyme having the transaminase activity of converting OMPB into L-AMPB in the presence of glutamic acid as the amino-donor. For instance, it is possible to use, as such transaminase also having so-called GOT activity, the commercially-available glutamic acid-oxaloacetic acid-transaminase (International Enzyme Classification Number: EC 2,6,1,1); or such transaminase also having the GOT activity which has been extracted by a conventional method from a microorganism having the GOT activity, for example, Streptomyces hygroscopicus SF-1293 strain (see Japanese Patent Application first publication "Kokai" No. 47485/82; FERM BP-130; ATCC 21705). These transaminases also having the GOT activity may again be used either singly or in combination. As the second transaminase which can convert OMPB into L-AMPB in the presence of L-glutamic acid as the amino-donor and which is used in combination with said transaminase also having the so-called GOT activity, any transaminase may be employed as long as it can convert OMPB into L-AMPB in the presence of L-glutamic acid.

As the amino-donor compounds useful in the process of this invention, any known amino-donors such as those disclosed in the "SEIKAGAKU JIKKEN KOZA", Volume 11

"Metabolism of Amino Acids and Bio-amines", compiled by The Japanese Biochemical Society and published from Tokyo Kagaku Dozin Company Limited or in the "Journal of Biological Chemistry", 247, 2486 (1972) may all be used.

5 Preferred examples of the amino-donor compounds available according to this invention include straight-chain aliphatic L- α -amino acids such as L-glutamic acid, L-aspartic acid, L-alanine, L-methionine, L-glutamine and the like; branched-chain aliphatic L- α -amino acids such as
10 L-leucine, L-isoleucine, L-valine and the like; basic amino acids such as L-lysine, L-ornithine, L-histidine and the like; as well as aromatic amino acids such as L-phenylalanine, L-tyrosine, L-tryptophan and the like, and an alkali metal salts such as sodium or potassium salt of
15 these amino acids. Besides, D-amino acids corresponding to the above-mentioned L-amino acids may equally be used as the amino-donor in the present process. These amino-donors may be used either singly or in combination. As the amino-donor, it is preferable to use glutamic acid
20 or a salt thereof, in combination with aspartic acid or a salt thereof. Molar ratio of the quantity of the amino donor to the quantity of OMPB present in the reaction medium may generally be in a range of 10:1 to 1:10 before the reaction is initiated.

25 When glutamic acid and/or aspartic acid is used

as the amino-donor(s) in the process of this invention, commercially available glutamic acid or aspartic acid may be used as such. In general, these amino acids may each be in the form of a mixture of the D-isomer and
5 L-isomer. Their L-isomers are however preferred. As salts of glutamic acid and aspartic acid, their alkali metal salts, especially, their sodium or potassium salts may be used.

In a preferred embodiment of the process of this
10 invention, glutamic acid (or its salt) and aspartic acid (or its salt) are used in combination. The respective concentrations of these amino-donor compounds and the ratio of the amino-donor compounds to OMPB may preferably be set as described below. When these amino-
15 donors are provided at higher ratios than OMPB so that a greater quantity of the amino donor is present than OMPB in the reaction medium, the reaction speed and the rate of the conversion of OMPB into L-AMPB can increase. Suitable ratio of the amino-donor compound to OMPB may,
20 however, be set from the economical standpoint. In general, molar ratio of the concentration (or amount to be added) of glutamic acid or its salt to the concentration (or amount to be added) of OMPB may preferably be in a range of 0.2:1 to 3.0:1. On the other
25 hand, the molar ratio of the concentration (or amount

to be added) of aspartic acid or its salt to the concentration (or amount to be added) of OMPB may desirably be in a range of 1.0:1 to 3.0:1.

When a transaminase-producing microorganism is
5 to be reacted with OMPB of the formula (II) and the amino-donor in the process of this invention, the present process may be carried out in such manner that said microorganism is at first cultured in a culture medium which contains nutrients useful in usual cultivation of microorganisms.
10 It is possible to use, as the nutrient sources, any conventional nutrient sources which have been utilized in the ordinary cultivation of microorganisms. As the nutrient carbon source may be used, for example, glucose, starch, glycerin, sucrose, thick malt syrup, molasses or
15 the like. They may be used either singly or in combination. As the nutrient nitrogen source may be used, for example, soybean meal, wheat germs, meat extract, peptone, dry yeast, corn steep liquor, ammonium sulfate, and the like, either singly or in combination. It is also feasible
20 to add to the culture medium one or more inorganic salts such as calcium carbonate, sodium chloride, potassium chloride and phosphates as needed. As the cultivation method for the microorganism, the liquid cultivation method, especially, the submerged cultivation method is
25 most suited. The cultivation of the microorganism may be

conducted under aerobic conditions. The temperature suitable for the cultivation may range from 25°C to 40°C. The cultivation may suitably be continued for 1-4 days for actinomycetes, 1-2 days for bacteria, 1-2 days for yeasts and 1-4 days for fungi (molds).

The resulting culture broth of the transaminase-producing microorganism so cultivated may be used as such. If desired, the microbial cells may be separated from the culture broth and then washed with water or physiological saline. The resulting washed, intact cells may then be suspended at a suitable cell concentration in a volume of water, physiological saline or a suitable aqueous buffered solution to provide a cell suspension ready for use. To the culture broth or aqueous liquid suspension containing the cells of the microorganism employed, OMPB (a first substrate) and the amino-donor compound(s) (a second substrate for the enzyme) are then added either simultaneously or successively. The resultant liquid mixture is thereafter maintained under such conditions that the microorganism is allowed to react with OMPB and the amino-donor to proceed with the conversion reaction of OMPB into L-AMPB. The cell concentration and the concentrations of the OMPB and amino-donor in the aqueous liquid reaction medium, where OMPB and amino-donor are treated with the microorganism in the above-

described manner to be converted into L-AMPB, as well as the reaction temperature and pH conditions may suitably be adjusted so as to maintain the reaction medium under such conditions that the conversion reaction of OMPB into
5 L-AMPB should proceed efficiently and the microorganism should be allowed to exhibit its function. The reaction time may also be adjusted to produce and accumulate a substantial amount of L-AMPB in the reaction mixture.

When OMPB and the amino-donor compound are to be
10 treated with transaminase(s) in the process of this invention on the other hand, OMPB and the amino-donor(s) are added and dissolved in an enzyme solution of suitable transaminase(s) in water or a buffered solution, followed by conducting the enzymatic reaction of OMPB. The
15 respective concentrations of the transaminase(s), OMPB and amino-donor(s) in the reaction system as well as the reaction temperature and pH conditions may suitably be adjusted to their optimal ranges in which the conversion reaction of OMPB into L-AMPB should proceed with good
20 efficiency.

As the transaminase may be used a transaminase which is available commercially in the form of an enzyme product. As an alternative, it is also feasible to use either such an aqueous, crude enzyme solution which has
25 been prepared directly from a culture broth of an

actinomycetes, bacterium, fungus or yeast as the trans-aminase-producing microorganism useful in the present invention by disintegrating the cells of the microorganism in said broth, or an aqueous solution of a crude enzyme
5 product in water. The aqueous crude enzyme solution may also be used in the form of an aqueous extract of the disintegrated cells of the microorganism.

Such a crude enzyme solution which has been prepared from the above-described microorganism or its
10 culture broth by a known method such as ultrasonic treatment or lysozyme treatment of the cells may also be used in the process of this invention, so long as said crude enzyme solution shows the ability to produce L-AMPB from OMPB in the presence of the amino-donor(s). Needless to
15 say, an aqueous solution of the enzyme in a purified form may also be used. It is known that the stability and operation efficiency of an enzyme or microorganism can be enhanced by immobilizing same with aid of an organic solvent, crosslinking agent and carrier or the like.

20 An immobilized transaminase or an immobilized, transaminase-producing microorganism, which has been treated by such a known immobilization method, may also be employed in the process of this invention as far as it has the ability to produce L-AMPB from OMPB.

25 In the process of this invention, the enzymatic

reaction for converting OMPB into L-AMPB with transaminase(s) may be carried out preferably at a pH of 7.5 or higher, notably, within a pH range of 8.0-9.0. Reaction conditions should properly be chosen within the temperature and pH
5 ranges optimal to the activity of the transaminase(s) which takes or take part in the enzymatic reaction.

In a preferred embodiment of the process of this invention, 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid, namely, OMPB is treated with one or more transaminases or
10 transaminase-producing microorganism(s) in the presence of both of L-glutamic acid or its salt and L-aspartic acid or its salt as the amino-donors. As the transaminase(s) employed in this embodiment, it is preferable to use an enzyme system which is composed of a combination of such
15 an enzyme having the transaminase activity to convert 2-ketoglutaric acid into glutamic acid in the presence of L-aspartic acid as amino-donor, namely, such an enzyme having the transaminase activity which falls within the category of the so-called GOT activity, and such second enzyme
20 having the transaminase activity to convert OMPB into L-AMPB in the presence of L-glutamic acid as amino-donor. Again, the microorganism to be employed in the above preferred embodiment of the present process may preferably be such a microorganism capable of producing an
25 enzyme system which has not only an enzymatic activity to

convert 2-ketoglutaric acid into glutamic acid in the presence of L-aspartic acid as amino-donor, i.e., the so-called GOT activity, but also a transaminase activity to convert OMPB into L-AMPB in the presence of L-glutamic acid as amino-donor, and said microorganism may be, for example, Streptomyces hygroscopicus SF-1293 strains (FERM BP-130 or ATCC 21705) or its mutant strain, Streptomyces hygroscopicus NP-50 strain (see Japanese Patent Application first publication "Kokai" No. 58589/86; FERM P-7804 or FERM BP-1368).

In the above preferred embodiment, it may be presumed that OMPB will receive the amino group from the glutamic acid (or its salt) as one of the amino-donors under the action of the above-mentioned transaminase system or microorganism so as to form L-AMPB with the glutamic acid turning to 2-ketoglutaric acid upon donation of its amino group, while the aspartic acid (or its salt) as the other amino-donor gives its amino group to said 2-ketoglutaric acid under the action of such transaminase also having the GOT activity, whereby 2-ketoglutaric acid can be regenerated into glutamic acid and aspartic acid (or its salt) itself is converted into oxaloacetic acid and finally into pyruvic acid.

In this way, the process of this invention provides a reaction solution containing L-AMPB as produced.

In a summary, therefore, the process of this invention may suitably be carried out in such a way that 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid is treated or reacted with at least one transaminase in the presence of at least one amino-donor compound, preferably in the presence of both L-glutamic acid and L-aspartic acid or both their sodium salts, in an aqueous liquid reaction medium in which 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid and the amino-donor compound(s) as well as the transaminase(s) have been dissolved, under alkaline conditions in a range of pH 7.5 to pH 9.0 and at a temperature in a range of room temperature to 60°C. Also, the process of this invention may suitably be carried out in such a way that 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid is treated or reacted with at least one microorganisms capable of producing at least one transaminase, in the presence of at least one amino-donor compound, preferably in the presence of both L-glutamic acid and L-aspartic acid or both their sodium salts, in an aqueous liquid reaction medium in which 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid and the amino-donor compound(s) have been dissolved and the cells of said microorganism have been suspended, under alkaline conditions of pH 7.5 to pH 9.0 and at a temperature in a range of room temperature to 60°C. When carrying out the present process using the cells of a trans-

aminase-producing microorganism, 4-(hydroxymethylphosphinyl)-
2-oxo-butyric acid and amino-donor compound(s) may be added
to such a culture broth of a microorganism capable of pro-
ducing the transaminase in which the intact cells of said
5 microorganism are suspended, and then the interaction bet-
ween 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid, the
amino-donor compound(s) and said microorganism is effected.

Again also, the process of this invention may suitab-
ly be carried out in such a way that 4-(hydroxymethyl-
10 phosphinyl)-2-oxo-butyric acid is treated with an extract
of a microorganism capable of producing at least one trans-
aminase and containing the transaminase, in the presence of
at least one amino-donor compound, preferably in the presence
of both L-glutamic acid and L-aspartic acid or both their
15 sodium salts, in an aqueous liquid reaction medium in which
4-(hydroxymethylphosphinyl)-2-oxo-butyric acid and the amino-
donor compound(s) as well as the transaminase-containing
extract of said microorganism have been dissolved, under
alkaline conditions of pH 7.5 to pH 9.0 and at a temperature
20 in a range of room temperature to 60°C. In whatever way
the process of this invention is to be carried out, 4-
(hydroxymethylphosphinyl)-2-oxo-butyric acid may be dis-
solved at its initial concentration of 0.1 to 100 mg/ml in
an aqueous liquid reaction medium before the enzymatic re-
25 action begins to proceed. And 4-(hydroxymethylphosphinyl)-

2-oxo-butyric acid and the amino-donor compound(s) may be present at a molar ratio in a range of 1:10 to 10:1 in an aqueous liquid reaction medium initially before the enzymatic reaction of converting 4-(hydroxymethylphosphinyl)-
5 2-oxo-butyric acid into L-2-amino-4-(hydroxymethylphosphinyl)-butyric acid takes place.

When the interaction between 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid (i.e, OMPB), the amino-donor compound(s) and the transaminase(s) or the cells of the trans-
10 aminase-producing microorganism has been effected according to the process of this invention, there is provided an aqueous reaction solution or mixture containing an amount of L-2-amino-4-(hydroxymethylphosphinyl)-butyric acid (i.e. L-AMPB) produced, an amount of OMPB remaining un-
15 reacted, an amount of the transaminase(s) used or the cells of the microorganism used. Before recovering L-AMPB from said reaction solution or mixture, it is preferable to centrifuge the reaction solution or mixture if this is containing the cells of the microorganisms
20 used. By this centrifugation, the microbial cells can be removed from the reaction solution or mixture, and there is obtained a supernatant solution containing L-AMPB but free from the microbial cells.

Now, the recovery of the L-AMPB product from its
25 aqueous solution is described below.

Recovery and purification of L-AMPB from its aqueous solution so obtained may be conducted in the same manner as in the known methods of recovering and purifying L-AMPB from the culture broth of the L-AMPB-producing micro-
5 organism which is obtained according to the known fermentative method of producing L-AMPB. Detailed procedures for recovery and purification of L-AMPB are disclosed in Japanese patent application first publication "Kokai" No. 47485/82. For instance, the recovery and purification
10 of L-AMPB from the reaction solution as obtained from the process of this invention may be achieved by passing the L-AMPB-containing reaction solution through a column of a cation-exchange resin such as Dowex 50 W (product of Rohm & Haas Co., Ltd., U.S.A.) to make L-AMPB adsorbed by
15 this resin, and then eluting the resin column with water or diluted aqueous ammonia solution to give such fractions of the eluate containing L-AMPB. The eluate fractions containing L-AMPB may then be collected together and concentrated under reduced pressure to afford a dry powder
20 of L-AMPB. The L-AMPB as produced according to the process of this invention shows the same physical and chemical properties as those of an authentic sample of L-AMPB which has been afforded by the known fermentative method as disclosed in the Japanese patent application first
25 publication "Kokai" No. 47485/82, and also it has been

observed that the L-AMPB as produced according to this invention shows the herbicidal activities same as those of the L-AMPB as produced by the known fermentative method tested by some herbicidal methods.

5 This invention is now illustrated with reference to the following Examples to which this invention is not limited.

Example 1

10 The production of L-AMPB was conducted by reaction of different kinds of commercially available transaminases with OMPB which was dissolved in a volume of a 50 mM phosphate buffered solution (pH 6) containing an amino-donor compound as added.

15 Thus, in the experiments where glutamic acid-oxaloacetic acid-transaminase (GOT) (a product of Boehringer Mannheim Co., Ltd.) was employed as the transaminase enzyme, L-aspartic acid (Asp) and L-glutamic acid (Glu) were both added as the amino-donors (as the substrates for the donation of amino group) to and dissolved
20 in the reaction medium. When glutamic acid-pyruvic acid-transaminase (GPT) (a product of Boehringer Mannheim Co., Ltd.) was used on the other hand, L-alanine (Ala) was added and dissolved as the amino-donor into the reaction medium. When glutamic acid dehydrogenase (GLDH) was employed for
25 a reference test, ammonium chloride and NADH were both

added as the amino-donor. For comparison purpose, production of glutamic acid (Glu) was also observed by reacting 2-ketoglutaric acid (2-KG) (as a control substrate) with the transaminase employed.

5 After conducting the enzymatic reaction at 30°C for 50 minutes in each experiment, the reaction mixture was heated at 100°C for 3 minutes to terminate the reaction. The reaction mixture or solution was adjusted to pH 2 by addition of diluted aqueous sulfuric and cen-
10 trifuged to obtain a supernatant solution. The amounts of L-AMPB and glutamic acid as formed and present in the supernatant solution were determined by an amino acid analyzer. Results of the experiments are shown in Table 1.

Table 1

Enzyme used and its amount added	Substrate com- pound and its concentration ($\mu\text{g/ml}$)		Amino-donor compound and its concentration ($\mu\text{g/ml}$)		Reaction product and its amount produced ($\mu\text{g/ml}$)	
GOT (10 units/ml)	OMP	100	Asp	1000	L-AMPB	2.8
	OMP	100	Glu	1000	L-AMPB	3.7
	2-KG (Control)	100	Asp	1000	Glu	109.3
GPT (4 units/ml)	OMP	100	Ala	1000	L-AMPB	0.9
	2-KG (Control)	100	Ala	1000	Glu	92.7
GLDH (60 units/ml)	OMP	100	NADH+NH ₄ Cl 1000		L-AMPB	1.9
	2-KG (Control)	100	NADH+NH ₄ Cl 1000		Glu	109.3

It is clear from the results of Table 1 that L-AMPB was produced from OMPB by the enzymatic reaction, although the yield of L-AMPB was significant but not so high as the yield of glutamic acid as produced from 2-ketoglutaric acid.

5 Example 2

Test bacterial strains indicated in Table 2 given below were separately inoculated in 40-ml portions of a culture medium comprising Nutrient broth (product of Difco Corporation), followed by productive cultivation
10 of the bacteria at 28°C for 6 hours. The resultant culture broths each were used as a seed culture, and it was inoculated at an inoculum size of 2% in 40-ml portions of a culture medium of the same composition as above and cultivated at 28°C overnight, and OMPB was added to
15 a concentration of 100 µg/ml in each of the resulting culture broths containing the bacterial cells.

To each of the resulting culture broths containing OMPB as added, there was further added sodium L-aspartate to a concentration of 100 µg/ml as the amino-donor. The
20 liquid mixtures comprising the culture broth, OMPB and sodium L-aspartate were thus provided. Thereafter, the enzymatic conversion of OMPB into L-AMPB was allowed to proceed at 28°C for 24 hours, and the reaction mixtures were adjusted to pH 2 with 25% sulfuric acid to terminate
25 the reaction. The cells were removed by centrifugation.

The amounts of L-AMPB as formed in the respective supernatant solutions obtained were determined using an amino acid analyzer. The test results are summarized in Table 2.

Table 2

<u>Tested bacterial strain</u>	<u>Amount of L-AMPB produced ($\mu\text{g/ml}$)</u>	<u>GOT activity (potency) of Crude enzyme solution ($\times 10^{-3}$ units/ml)</u>	<u>Specific GOT activity of enzyme ($\times 10^{-3}$ units/mg of protein)</u>
Bacillus subtilis ATCC 6633	2.4	22.0	46.7
Micrococcus luteus ATCC 9341	1.9	22.0	196.4
Staphylococcus aureus 209P (ATCC 6538)	26.6	17.5	182.3
Escherichia coli ATCC 10798	0.8	3.8	5.4
Pseudomonas aeruginosa ATCC 10145	8.3	342.0	686.7
Serratia marcescens ATCC 13880	5.0	45.6	300.0

5 It is apparent from Table 2 that AMPB can be produced from OMPB in significant yields.

10 Further, portions of the culture broths as obtained after the above-mentioned overnight cultivation at 28°C were separately subjected to an ultrasonic treatment to disintegrate the bacterial cells and obtain cell extracts. The cell extracts so prepared were separately centrifuged to produce aqueous crude enzyme solutions. The GOT activity levels (potency) of the respective crude enzyme solutions so obtained were also measured and are shown

in Table 2 for information. As shown in Table 2, significant correlation is observed between the amounts of L-AMPB produced and the measured value of the GOT activity of said crude enzyme solution. Besides, said crude enzyme solutions were each analysed for their protein contents according to the Bio-Rad Protein Assay method, and the specific GOT activity of the enzyme ($\times 10^{-3}$ units/mg of protein) present in said solution was evaluated and shown in Table 2 above.

10 Example 3

Streptomyces hygroscopicus SF-1293 strain (FERM BP-130) was inoculated in 10 ml of a preliminary culture medium (comprising 2.0% soluble starch, 1.0% polypeptone, 0.3% meat extract, 0.05% dipotassium hydrogen phosphate; pH 7.0). The above strain was shake-cultivated at 28°C for 24 hours. The resulting culture broth was used as a seed culture, and it was inoculated at an inoculum size of 2% into a productive culture medium (comprising 7.0% glucose, 4.4% bactosoyton, 0.327% potassium dihydrogen phosphate, 0.0852% disodium hydrogen phosphate, 1.15% dotite TES, namely N-tris(hydroxymethyl)methyl-2-aminoethanesulfonic acid, 0.0001% cobalt chloride; pH 6.0) and then the cultivation of the SF-1293 strain was made at 28°C under aeration and agitation. After the cultivation for 4 days, the microbial cells were collected by

centrifugation and washed with a 50mM phosphate buffered solution (pH 6.0). The cells were then disintegrated by ultrasonic treatment (using a device designated "KUBOTA INSONATOR"; 1.5A, for 1 minute), followed by centrifugation to obtain a crude enzyme solution.

OMPB was added to a concentration of 100 µg/ml and L-aspartic acid was added to a concentration of 200 µg/ml into said crude enzyme solution. A further control test was also made using 2-ketoglutaric acid (2-KG) as a control substrate for the transamination reaction. After allowing the enzymatic reaction to proceed at 30°C for 2 hours, the reaction solutions were heated at 100°C for 3 minutes to terminate the reaction. The reaction solution obtained was adjusted to pH 2 with diluted sulfuric acid and a supernatant solution was obtained by centrifugation of the reaction solution. The amounts of L-AMPB and the glutamic acid as formed from 2-KG which were present in the supernatant solutions were determined by an amino acid analyzer. Test results are shown in Table 3.

Table 3

Substrate	Product and its amount produced (µg/ml)	
OMPB	L-AMPB	2.8
2-KG (Control)	Glu	67.9

Example 4

(1) Streptomyces hygroscopicus NP-50 strain (FERM P-7804 or FERM BP-1368) was inoculated in 10 ml of a preliminary culture medium (comprising 2.0% soluble starch, 1.0% polypeptone, 0.3% meat extract, 0.05% di-
5 potassium hydrogen phosphate; pH 7.0). The above NP-50 strain was shake-cultured at 28°C for 24 hours. The resulting culture broth was used as a seed culture, and it was inoculated at an inoculum size of 2% into a pro-
10 ductive culture medium (comprising 7.0% glucose, 4.4% bactosoyton, 0.327% potassium dihydrogen phosphate, 0.0852% disodium hydrogen phosphate, 1.15% dotite TES, 0.0001% cobalt chloride; pH 6.0) and then the cultivation of the NP-50 strain was made at 28°C under aeration and
15 agitation.

(2) In a 250-ml Erlenmeyer flask, were combined 20 ml of the culture broth of Streptomyces hygroscopicus NP-50 strain cultivated for 3 days in the above productive culture medium, 30 ml of an aqueous solution of OMPB
20 (OMPB concentration: 87 mg/ml, adjusted to pH 7.0 in advance), 40 ml of an aqueous solution of commercial sodium L-glutamate (sodium L-glutamate concentration: 170 mg/ml) and 10 ml of 1M tris-HCl buffered solution (pH 8.5). The OMPB concentration in the combined liquids
25 amounted to about 26 mg/ml. While gently shaking the

resulting liquid mixture at 37°C in the flask, the enzymatic reaction was allowed to proceed for 24 hours. Thereafter, the reaction mixture so obtained was centrifuged to remove the microbial cells therefrom, and the resulting supernatant solution containing L-AMPB produced (100 ml) was analyzed by an amino acid analyzer to determine the amount of L-AMPB in the solution. It was found that 14 mg/ml of L-AMPB had been produced.

In addition, similar experiments were also conducted using L-alanine or sodium L-aspartate separately as the amino-donors (as added to a concentration of 170 mg/ml). It was observed that 2.8 mg/ml of L-AMPB was produced when L-alanine was used, and 1.5 mg/ml of L-AMPB when sodium L-aspartate was employed.

(3) The supernatant solution (100 ml) obtained by centrifugating the reaction mixture which had been formed upon the use of sodium L-glutamate as the amino-donor in the above procedure (2) was passed into a column of 400 ml of a cation exchange resin "Dowex 50 W x 2" (H^+ -form) (trade name; product of Rohm & Haas Co.), followed by development with dilute aqueous ammonia. The L-AMPB-containing fractions of the eluate were collected and concentrated. The concentrated solution was then subjected to chromatography on an anion-exchange resin column of 150 ml of "Dowex 1 x 2" (CH_3COO^- -form) (trade

name; product of Rohm & Haas Co.). After washing the resin column with water, the column was eluted with a 0.3N aqueous solution of acetic acid.

5 The L-AMPB-containing fractions of the eluate were concentrated to dryness under reduced pressure, followed by drying in vacuo to give 720 mg of L-AMPB as a white powder. The white powder was recrystallized from methanol. The crystals thus obtained were analyzed in a usual manner, i.e., by determining the elemental
10 analysis, specific optical rotation, melting point, infrared absorption spectrum, nuclear magnetic resonance spectrum and mass spectrum. It was confirmed that the crystals of L-AMPB obtained was fully indential to an authentic sample of L-AMPB.

15 Example 5

(1) The culture broth (20 ml) of Streptomyces
hygroscopicus NP-50 strain (FERM P-7804; FERM BP-1368), which had been cultivated in the productive culture medium used in the procedure (1) of Example 4, was placed
20 in a 30-ml centrifuge tube and then centrifuged at 3000 rpm. for 10 minutes. Precipitated cells of the NP-50 strain were suspended in 30 ml of 50mM tris-HCl buffered solution (pH 8.5) to prepare the cell suspension. In a 250-ml Erlenmeyer flask, were combined 20 ml of the cell
25 suspension, 30 ml of an aqueous solution of OMPB (OMP

concentration: 87 mg/ml, adjusted to pH 7.0 in advance),
40 ml of an aqueous solution of commercial sodium L-
glutamate (sodium L-glutamate concentration: 170 mg/ml)
and 10 ml of 1M tris-HCl buffered solution (pH 8.5).

5 The OMPB concentration in the combined liquids amounted
to about 26 mg/ml. While gently shaking the resulting
liquid mixture at 37°C, the enzymatic reaction was allowed
to proceed for 24 hours. Thereafter, the reaction mixture
so obtained was centrifuged to remove the microbial cells
10 therefrom, and the resulting supernatant solution con-
taining L-AMPB produced (100 ml) was analyzed by an amino
acid analyzer to determine the amount of L-AMPB in the
solution. It was found that 15 mg/ml of L-AMPB had been
produced.

15 (2) In the same manner as in the procedure (3)
of Example 4, L-AMPB-containing supernatant solution
(100 ml) obtained in the above procedure (1) was subjected
to chromatography on a column of 400 ml of a cation-
exchange resin "Dowex 50W x 2" (H⁺-form), followed by
20 development with a dilute aqueous ammonia. Upon post-
treatment of the L-AMPB-containing fractions of the
eluate in the same manner as in the procedure (3) of
Example 4, 750 mg of L-AMPB was obtained as a white
powder.

Example 6

(1) The culture broth (20 ml) of Streptomyces
hygroscopicus NP-50 strain (FERM P-7804; FERM BP-1368),
which had been cultured in the productive culture medium
5 used in the procedure (1) of Example 4, was placed in a
30-ml centrifuge tube and centrifuged at 3000 rpm. for
10 minutes. Precipitated cells of the NP-50 strain were
suspended in 30 ml of 50mM tris-HCl buffered solution
(pH 8.5) to prepare the cell suspension. After dis-
10 integrating the cells in the cell suspension for 10
minutes by an ultrasonic disintegrator, the resultant
suspension was centrifuged at 10000 rpm. for 10 minutes
to yield 20 ml of a crude enzyme solution as the super-
natant solution.

15 (2) In a 250-ml Erlenmeyer flask, were combined
20 ml of the supernatant solution (the crude enzyme
solution) obtained as above, 30 ml of an aqueous solution
of OMPB (OMPB concentration: 87 mg/ml, adjusted to pH
7.0 in advance), 40ml of an aqueous solution of commercial
20 sodium L-glutamate (sodium L-glutamate concentration:
170 mg/ml) and 10 ml of 1M tris-HCl buffered solution
(pH 8.5). The OMPB concentration in the combined liquids
amounted to about 26 mg/ml. While gently shaking the
resulting liquid mixture at 37°C, the enzymatic reaction
25 was allowed to proceed for 24 hours. Thereafter, the

resulting reaction mixture obtained (100 ml) was analyzed by an amino acid analyzer to determine the amount of L-AMPB formed in said reaction mixture. It was found that 16 mg/ml of L-AMPB had been produced.

5 (3) In the same manner as in the procedure (3) of Example 4, the reaction mixture (100 ml) obtained from the enzymatic reaction of the above procedure (2) was subjected to chromatography on an ion-exchange resin column of 400 ml of "Dowex 50W x 2" (H⁺-form), followed
10 by development with a dilute aqueous ammonia. Upon post-treatment of the L-AMPB-containing fractions of the eluate in the same manner as in the procedure (3) of Example 4, 740 mg of L-AMPB was obtained as a white powder.

15 Example 7

 (1) Streptomyces lividans 66 strain (FERM BP-737) was inoculated in 10 ml of a preliminary culture medium of the same composition as that employed in the procedure (1) of Example 4. The above microorganism was shake-
20 cultivated at 28°C for 24 hours. The resultant culture broth was used as a seed culture, and it was inoculated at an inoculum size of 2% in a productive culture medium of the same composition as that used in the procedure (1) of Example 4 and was then cultivated at 28°C under
25 aeration and agitation.

(2) In a 250-ml Erlenmeyer flask, were combined 20 ml of the culture broth of Streptomyces lividans 66 strain cultivated for 3 days in the above productive culture medium, 30 ml of an aqueous solution of OMPB (OMP concentration: 87 mg/ml, adjusted to pH 7.0 in advance), 40 ml of an aqueous solution of commercial sodium L-glutamate (sodium L-glutamate concentration: 170 mg/ml) and 10 ml of 1M tris-HCl buffered solution (pH 8.5). The OMPB concentration in the combined liquids amounted to about 26 mg/ml. While gently shaking the resulting liquid mixture at 37°C in the flask, the enzymatic reaction was allowed to proceed for 24 hours. Thereafter, the reaction mixture so obtained was centrifuged to remove the microbial cells therefrom, and the resulting supernatant solution containing L-AMPB produced (100 ml) was analyzed by an amino acid analyzer to determine the amount of L-AMPB in the supernatant solution. It was found that 6 mg/ml of L-AMPB had been produced.

Example 8

(1) Streptomyces hygroscopicus SF-1293 strain (FERM BP-130; ATCC 21705) was inoculated in 10 ml of a preliminary culture medium of the same composition as that employed in the procedure (1) of Example 4. The above SF-1293 strain was shake-cultivated at 28°C for 24 hours. The resultant culture broth was used as a seed

culture, and it was inoculated at an inoculum size of 2% in a productive culture medium of the same composition as that used in the procedure (1) of Example 4 and then the SF-1293 strain was cultivated at 28°C under aeration and agitation.

(2) In a 250-ml Erlenmeyer flask, were combined 20 ml of the culture broth of Streptomyces hygroscopicus SF-1293 strain cultured for 3 days in the above productive culture medium, 30 ml of an aqueous solution of OMPB (OMP concentration: 87 mg/ml, adjusted to pH 7.0 in advance), 40 ml of an aqueous solution of commercial sodium L-glutamate (sodium L-glutamate concentration: 170 mg/ml) and 10 ml of 1M tris-HCl buffered solution (pH 8.5). The OMPB concentration in the combined liquids amounted to about 26 mg/ml. While gently shaking the resulting liquid mixture at 37°C in the flask, the enzymatic reaction was allowed to proceed for 24 hours. Thereafter, the reaction mixture so obtained was centrifuged to remove the microbial cells therefrom, and the resulting supernatant solution containing L-AMPB produced (100 ml) was analyzed by an amino acid analyzer to determine the amount of L-AMPB produced in the supernatant solution. It was found that 7 mg/ml of L-AMPB had been produced.

(3) In the same manner as in the procedure (3)

of Example 4, the supernatant solution (100 ml) obtained in the above procedure (2) was subjected to chromatography on an ion-exchange resin column of 400 ml of "Dowex 50W x 2" (H^+ -form), followed by development with
5 a dilute aqueous ammonia. Upon post-treatment of the L-AMPB-containing fractions of the eluate in the same manner as in the procedure (3) of Example 4, 350 mg of L-AMPB was obtained as a white powder.

Example 9

10 Suspended in 100 ml of 50mM tris-HCl buffered solution (pH 8.5) was 20 g of a wet cake of commercial baker's yeast (Saccharomyces cerevisae; a product of Oriental Yeast Co., Japan). In a 250-ml Erlenmeyer flask, were combined 20 ml of the resulting cell sus-
15 pension of the baker's yeast, 30 ml of an aqueous solution of OMPB (OMPB concentration: 87 mg/ml, adjusted to pH 7.0 in advance), 40 ml of an aqueous solution of commercial sodium L-glutamate (sodium L-glutamate concentration: 170 mg/ml) and 10 ml of 1M tris-HCl buffered solution
20 (pH 8.5). The OMPB concentration in the combined liquids then amounted to about 26 mg/ml. While shaking the resulting liquid mixture at 37°C for 24 hours, the enzymatic reaction of OMPB was allowed to proceed. The reaction mixture so obtained was centrifuged to
25 remove the yeast cells therefrom, and the resulting

supernatant solution containing L-AMPB produced (100 ml) was analyzed by an amino acid analyzer to determine the amount of L-AMPB present in the supernatant solution. It was found that 6 mg/ml of L-AMPB had been produced.

5 Example 10

 In 50 mM tris-HCl buffered solution (pH 8.5) containing 400 µg/ml of OMPB and 600 µg/ml of L-glutamic acid added and dissolved therein, a transaminase was reacted with OMPB to produce L-AMPB. As this trans-
10 aminase, a commercial glutamic acid-oxaloacetic acid-transaminase (GOT) (a product of Boehringer Mannheim Co.) was used. The GOT concentration was set at 40 units/ml. After allowing the enzymatic reaction to proceed at 37°C for 24 hours, the resulting reaction solution was heated
15 at 100°C for 3 minutes to terminate the reaction. After this, the reaction solution was adjusted to pH 2 with diluted sulfuric acid and then centrifuged to remove the insoluble solids and give a supernatant solution. The amount of L-AMPB in the supernatant solution was deter-
20 mined by an amino acid analyzer. The amino acid analyzer was Model "MLC-703" manufactured by ATTO CORPORATION, Japan, and the retention time was set for 12 minutes. It was found that 100 µg/ml of L-AMPB had been produced in the supernatant solution.

Example 11

(1) Cultivation method:

Several species of microorganisms indicated in Tables 4-10 below were separately inoculated from their
5 respective seed culture slants into volumes (10 ml-
portions) of a liquid culture medium (comprising 0.5%
glucose, 0.3% yeast extract, 1.0% meat extract, 1.0%
peptone, 0.3% sodium chloride; pH 7.0) as charged in
large-capacity test tubes. A piece of stainless
10 steel-coil was also placed in such test tubes as
employed for the cultivation of the actinomycetes.
The inoculated microorganisms were then separately
cultivated at 28°C for 24 hours (or 48 hours) on tube
shakers.

15 (2) Preparation of cell samples:

After the cultivation, 1-ml portions of the
resulting culture broths (except 0.2-ml portions of the
fungi culture in the case of the fungi) were each with-
drawn into micro-test tubes (product of Eppendorf Co.,
20 Ltd.) and centrifuged at 12000 rpm. for 3 minutes. The
supernatants as formed were discarded and the remaining
cells deposited were each collected as the cell samples.

(3) Engymatic reactions:

25 Into each of the cell samples so packed in the
respective micro-test tubes, were then added 20 µl of

tris-HCl buffered solution (1 mole, pH 8.0), 20 μ l of an aqueous OMPB solution (OMPB concentration: 200 mg/ml, pH 8.0), 40 μ l of an aqueous solution of an amino-donor as indicated below (containing either one of sodium L-glutamate, sodium L-aspartate and L-alanine, at a concentration of 1 mol/ml) and 120 μ l of water. The resultant mixture was allowed to stand at 37°C for 24 hours to the effect the interaction between the microbial cells, OMPB and the amino-donor compound present. After the reaction, the reaction mixture was centrifuged at 12000 rpm. for 3 minutes, and the amount of L-AMPB present in the supernatant solution obtained was measured by determining the amount of L-AMPB with an amino acid analyzer. Test results are summarized in Tables 4-10.

Table 4

<u>Microorganisms used (Bacteria)</u>	<u>Amount of L-AMPB produced (mg/ml)</u>		
	<u>Glu</u>	<u>Asp</u>	<u>Ala</u>
Bacillus subtilis ATCC 6633	3.0	1.9	2.2
Micrococcus luteus ATCC 9341	2.6	0.2	0.3
Staphylococcus aureus 209P ATCC 6538	3.2	0.6	0.5
Escherichia coli ATCC 10798	9.8	4.2	3.3
Pseudomonas aeruginosa ATCC 10145	8.3	3.3	3.8
Pseudomonas cepacia ATCC 17759	9.6	2.4	3.0
Serratia marcescens ATCC 13880	9.5	1.8	1.6
Corynebacterium glutamicum ATCC 13032	2.5	0.3	0.4

Table 5

<u>Microorganisms used (Yeasts)</u>	<u>Amount of L-AMPB produced (mg/ml)</u>		
	<u>Glu</u>	<u>Asp</u>	<u>Ala</u>
Saccharomyces cerevisiae ATCC 9763	0.9	0.3	0.2
Candida albicans IAM 4888	1.2	0.9	0.4
Candida albicans IAM 4829	12.1	6.1	4.0
Cryptococcus neoformans IAM 4772	2.4	1.3	0.8
Debaryomyces hansenii IAM 4356	2.7	1.2	0.7
Trigonopsis variabilis IAM 4443	2.2	1.5	0.4
Hansenula schneggi IAM 4269	1.6	0.8	0.2

Table 6

<u>Microorganisms used (Actinomycetes)</u>	<u>Amount of L-AMPB produced (mg/ml)</u>		
	<u>Glu</u>	<u>Asp</u>	<u>Ala</u>
Streptomyces albus IFO 13014 (ATCC 3004)	7.4	1.0	0.8
Streptomyces griseus IFO 12875 (ATCC 23345)	8.3	1.7	1.9
Streptovericillium cinnamomeum IFO 12852 (ATCC 11874)	10.6	6.0	2.2
Streptomyces morookaensis IFO 13416 (ATCC 19166)	6.8	7.5	3.3
Nocardia mediterranei ATCC 21271	8.1	2.6	1.6
Nocardiopsis dassonvillei JCM 3237	6.3	1.5	1.5
Saccharopolyspora hirsuta JCM 3170 (ATCC 27875)	3.7	0.8	0.9

Table 7

<u>Microorganisms used (Actinomycetes)</u>	<u>Amount of L-AMPB produced (mg/ml)</u>		
	<u>Glu</u>	<u>Asp</u>	<u>Ala</u>
Streptomyces viridochromogenes IFO 13347 (ATCC 14925)	9.5	1.6	1.1
Streptomyces pilosus IFO 12807 (ATCC 19797)	1.4	0.1	0.1
Micromonospora carbonacea NRRL 2972 (ATCC 27114)	6.1	1.4	1.6
Streptosporangium pseudovulgare ATCC 27100	3.0	0.8	1.7

* Cultivation time: 48 hours

Table 8

<u>Microorganism used (Actinomycetes)</u>	<u>Amount of L-AMPB produced (mg/ml)</u>		
	<u>Glu</u>	<u>Asp</u>	<u>Ala</u>
Streptomyces viridochromogenes JCM 4977 *	8.8	5.6	3.6

* Bialaphos-producing actinomycetes

Table 9

<u>Microorganisms used (Fungi)</u>	<u>Amount of L-AMPB produced (mg/ml)</u>		
	<u>Glu</u>	<u>Asp</u>	<u>Ala</u>
Aspergillus flavus ATCC 9643	0.5	0.2	0.3
Aspergillus terreus IAM QM82J	0.3	0.2	0.1
Mucor spinescens IAM Mu3	5.3	0.9	1.1
Aureobasidium pullulans IAM F24	1.7	0.6	0.4
Chaetomium globosum ATCC 6205	0.3	0.1	0.2

Table 10

<u>Microorganisms used (Fungi)</u>	<u>Amount of L-AMPB produced (mg/ml)</u>		
	<u>Glu</u>	<u>Asp</u>	<u>Ala</u>
Penicillium funiculosum ATCC 9644	0.6	0.4	0.3
Gliocladium virens ATCC 9645	2.0	0.4	0.5

* Cultivation time: 48 hours

Amongst the species or strains of the microorganisms indicated in Table 2 and Tables 4-10 hereinbefore, the species or strains having the ATCC-numbers have been deposited and stored in the "American Type Culture Collection", Washington D.C., U.S.A.; the species or strains having the JCM-numbers have been deposited and stored in the "Japan Collection of Microorganism", Institute of Physical and Chemical Research at Wako City, Saitama-ken, Japan; the species or strains having the IAM-numbers have been deposited and stored in the "Institute of Applied Microbiology, University of Tokyo", at Tokyo, Japan; and the species or strains having the IFO-numbers have been deposited and stored in the "Institute for Fermentation, Osaka" at Osaka, Japan. All of the species or strain of the microorganisms having the access numbers indicated as above are available and distributable as the known Type Culture strains from the above-identified public depositories.

Example 12

- (1) Streptomyces hygroscopicus NP-50 strain (FERM P-7804 or FERM BP-1368) was inoculated in 10 ml of a preliminary culture medium (comprising 2.0% soluble starch, 1.0% polypeptone, 0.3% meat extract, 0.05% dipotassium hydrogen phosphate; pH 7.0). The above NP-50 strain was shake-cultured at 28°C for 24 hours.

The resultant culture broth was used as a seed culture, and it was inoculated at an inoculum size of 2% in a productive culture medium (comprising 7.0% glucose, 3.9% wheat germs, 2.5% soluble vegetative protein, 5 0.3% potassium dihydrogen phosphate, 0.0001% cobalt chloride; pH 6.8). The cultivation was then conducted at 28°C under aeration and stirring.

(2) To 20-ml portions of the culture broth obtained by cultivation of the NP-50 strain in the above mentioned productive culture medium for 3 days and containing cells of Streptomyces hygrosopicus NP-50 strain, were added OMPB and sodium L-glutamate and sodium L-aspartate as the amino-donors in such quantities so as to give their respective concentrations 15 in the resulting mixture as given in Table 11 below at the beginning of the respective transamination reaction. By further addition of 10 ml of 1M tris-HCl buffered solution (pH 8.5), each of the resulting mixtures was adjusted to pH 8.5 and the volume of 20 the resultant mixture was brought to 100 ml. Thus, the original volume of said culture broth had been diluted fivefold.

The enzymatic amination reaction of OMPB was then allowed to proceed at 37°C for 24 hours with 25 gently shaking the resultant solution containing

the cells suspended therein as well as OMPB and the amino-donor compounds dissolved therein.

After the reaction, 100-ml portions of the respective reaction mixtures each were subjected
5 to a heat treatment (sterilization and inactivation of the enzyme) so as to terminate the reaction. The reaction mixture each were filtered to remove the microbial cells, and the amounts of L-AMPB in the individual filtrates obtained were measured by an
10 amino acid analyzer ("MLC-703", trade name, manufactured by ATTO CORPORATION: retention time of 12 minutes). Test results are shown in Table 11.

Table 11

Test No.	Initial Concentration of OMPB ($\mu\text{g/ml}$)	Initial concentration of amino donor before start of reaction ($\mu\text{g/ml}$)		Final concentration of L-AMPB produced after end of reaction ($\mu\text{g/ml}$)
		Sodium L-glutamate	Sodium L-aspartate	
1	10000	10000	0	5930
2	10000	20000	0	7170
3	10000	0	10000	350
4	10000	10000	10000	8500
5	30000	30000	30000	28360

From the results of Table 11, it is clearly observed that the amount of L-AMPB produced from OMPB increased significantly when the enzymatic amination reaction of OMPB was effected in the presence of sodium L-glutamate and sodium L-aspartate in combination as the amino-donors, as compared to when the enzymatic reaction of OMPB was effected in the presence of either one of sodium L-glutamate and sodium L-aspartate.

10 Example 13

A culture broth of Streptomyces hygroscopicus NP-50 strain, which had been prepared under the same preliminary cultivation conditions as in Example 12, was used as a seed culture, and it was inoculated into a volume of the same productive culture medium as that

employed in Example 12 in a 3-liter jar fermentor, followed by conducting the cultivation at 28°C for 3 days.

Two liters of the culture broth so obtained was transferred into a reaction tank, followed by addition of 300 g of OMPB into the culture broth. In addition, 310 g of sodium L-glutamate (product of Wako Pure Chemical Industries, Ltd.) and 290 g of sodium L-aspartate (product of Nakarai Chemicals, Ltd.) were also added and dissolved in said culture broth. After adjusting the resultant mixture to pH 8.5 with aqueous NaOH, the volume of the mixture was brought to 10 liters by addition of water. Thus, the concentration of OMPB in the resulting liquid mixture was 30000 µg/ml at the start of the reaction. The respective concentrations of sodium L-glutamate and sodium L-aspartate were equal at this time to the molar concentration of OMPB (1/6 mole). The enzymatic amination reaction of OMPB was then allowed to proceed at 37°C for 24 hours with gently stirring the liquid mixture containing the cells of the NP-50 strain, OMPB and the amino-donor compounds, and with controlling the pH at 8.5. After the reaction, the reaction mixture obtained was adjusted to pH 3.0 with addition of 50% sulfuric acid. After addition of a filter aid, the reaction mixture was filtered to remove the cells, affording 9 g of the

filtrate.


The concentration of L-AMPB in the filtrate was analyzed in the same manner as in Example 12. It was found to be 29000 µg/ml. The filtrate was then chromatographed by passing through a column of 5 l of "Dowex 50 W" (H⁺-form) as a cation-exchange resin, followed by development with water. The L-AMPB containing fractions of the eluate were again subjected to the chromatography on a column of "Dowex 1 x 2" (CH₃COO⁻-form) as an anion-
10 ion exchange resin, which was washed with water and then eluted with a 0.3N aqueous solution of acetic acid.

The L-AMPB-containing fractions of the eluate were concentrated to dryness under reduced pressure, followed by drying in vacuo to give L-AMPB as white powder. This
15 white powder was recrystallized from methanol. The yield of L-AMPB was 183 g (about 70%).


The crystals of L-AMPB thus obtained were analyzed in a usual manner, i.e., by measuring the elemental analysis, specific optical rotation, melting point, infrared
20 absorption spectrum, nuclear magnetic resonance spectrum and mass spectrum. It was confirmed that the L-AMPB crystal obtained was fully identical to an authentic sample of L-AMPB.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A process for the production of L-2-amino-4-(hydroxymethylphosphinyl)-butyric acid represented by the formula (I):



which comprises treating 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid represented by the formula (II):



with one or more transaminases or with one or more microorganisms capable of producing one or more transaminases, in the presence of one or more amino-donors.

2. A process as claimed in Claim 1, wherein the microorganism used is an actinomycetes.
3. A process as claimed in Claim 1, wherein the microorganism used is a bacterium.
4. A process as claimed in Claim 1, wherein the microorganism used is a yeast.
5. A process as claimed in Claim 1, wherein the microorganism used is a fungus.

6. A process as claimed in claim 2, wherein the microorganism used is an actinomycetes selected from the genus Streptomyces, Streptoverticillium, Nocardia, Nocardiosis, Saccharopolyspora, Kitasatosporia, Micromonospora or Streptosporangium.

7. A process as claimed in Claim 3, wherein the microorganism used is a bacterium selected from the genus Bacillus, Micrococcus, Staphylococcus, Escherichia, Pseudomonas, Serratia or Corynebacterium.

8. A process as claimed in Claim 4, wherein the microorganism used is a yeast selected from the genus Saccharomyces, Candida, Cryptococcus, Debaryomyces, Trigonopsis or Hansenula.

9. A process as claimed in claim 5, wherein the microorganism used is a fungus selected from the genus Aspergillus, Mucor, Aureobasidium, Chaetomium, Penicillium or Gliocladium.

10. A process as claimed in Claim 1, wherein the transaminase used is ^{selected from} ~~any one of~~ a glutamic acid-oxaloacetic acid-transaminase, a glutamic acid-pyruvic acid-transaminase, ^{or} ~~and~~ a transaminase capable of converting 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid into L-2-amino-4-(hydroxymethylphosphinyl)-butyric acid in the presence of glutamic acid as an amino-donor,



or a combination of two or more of said transaminases.

11. A process as claimed in any one of Claims 1-10, wherein the amino-donor used is glutamic acid and/or aspartic acid and/or salts thereof.

12. A process as claimed in Claim 1, wherein 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid is treated or reacted with at least one transaminase in the presence of at least one amino-donor compound in an aqueous medium in which 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid and the amino-donor compound as well as the transaminase(s) have been dissolved, under alkaline conditions in a range of pH 7.5 to pH 9.0 and at a temperature in a range of room temperature to 60°C.

13. A process as claimed in Claim 1, wherein 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid is treated or reacted with at least one microorganisms capable of producing at least one transaminase, in the presence of at least one amino-donor compound in an aqueous medium in which 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid and the amino-donor compound have been dissolved and the cells of said microorganism have been suspended, under alkaline conditions of pH 7.5 to pH 9.0 and at a temperature in a range of room temperature to 60°C.



14. A process as claimed in Claim 1, wherein 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid is treated with an extract of a microorganism capable of producing at least one transaminase and containing the transaminase, in the presence of at least one amino-donor compound in an aqueous medium in which 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid and the amino-donor compound as well as the transaminase-containing extract of said microorganism have been dissolved, under alkaline conditions of pH 7.5 to pH 9.0 and at a temperature in a range of room temperature to 60°C.

15. A process as claimed in Claim 1, wherein 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid is dissolved at its initial concentration of 0.1 to 100 mg/ml in an aqueous medium.

16. A process as claimed in Claim 1, wherein 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid and amino-donor compound(s) are present at a molar ratio in a range of 1:10 to 10:1 in an aqueous medium initially before the conversion of 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid into L-2-amino-4-(hydroxymethylphosphinyl)-butyric acid takes place.

17. A process as claimed in Claim 1 c. 13, wherein 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid and amino-donor compound(s) are added to such a culture broth of a

microorganism capable of producing the transaminase in which the intact cells of said microorganism are suspended, and then the interaction between 4-(hydroxymethylphosphinyl)-2-oxo-butyric acid, the amino-donor compound(s) and said microorganism is effected.

18. A process as claimed in Claim 1, wherein the microorganism used which is capable of producing the transaminase is selected from Streptomyces hygroscopicus SF-1293 FERM BP-130 or ATCC 21705, Streptomyces hygroscopicus NP-50 FERM BP-1368, Streptomyces lividans 66 FERM BP-737, Streptomyces albus IFO 13014 (ATCC 3004), Streptomyces griseus IFO 12875 (ATCC 23345), Streptovericillium cinnamoneum IFO 12852 (ATCC 11874), Streptomyces morookaensis IFO 13416 (ATCC 19166), Nocardia mediterranei ATCC 21271, Nocardiopsis dassonvillei JCM 3237, Saccharopolyspora hirsuta JCM 3170 (ATCC 27875), Streptomyces viridochromogenes IFO 13347 (ATCC 14925), Streptomyces viridochromogenes JCM 4977, Streptomyces pilosus IFO 12807 (ATCC 19797), Micromonospora carbonaceae NRRL 2972 (ATCC 27114), Streptosporangium pseudovulgare ATCC 27100; Bacillus subtilis ATCC 6633, Micrococcus luteus ATCC 9341, Staphylococcus aureus 209P ATCC 6538, Escherichia coli ATCC 10798, Pseudomonas aeruginosa ATCC 10145, Pseudomonas cepacia ATCC 17759, Serratia marcescens ATCC 13880, Corynebacterium glutamicum ATCC 13032; Saccharomyces

cerevisiae ATCC 9763, Candida albicans IAM 4888, Candida albicans IAM 4829, Cryptococcus neoformans IAM 4772, Debaryomyces hansenii IAM 4356, Trigonopsis variabilis IAM 4443, Hansenula schneggi IAM 4269; Aspergillus flavus ATCC 9643, Aspergillus terreus IAM QM82J, Mucor spinescens IAM Mu3, Aureobasidium pullulans IAM F24, Chaetomium globosum ATCC 6205, Penicillium funiculosum ATCC 9644 and Gliocladium virens ATCC 9645.

DATED this 2nd day of June, 1987.

MEIJI SEIKA KAISHA, LTD.

By their Patent Attorneys

T.G. AHEARN & CO.