

**(12) PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 199742608 B2**  
**(10) Patent No. 734090**

(54) Title  
**CXCR3 chemokine receptor, antibodies, nucleic acids, and methods of use**

(51)<sup>7</sup> International Patent Classification(s)  
C12N 015/12 C07K 016/18  
A61K 038/17 C12N 005/10  
C07K 014/19 G01N 033/50  
C07K 014/715 G01N 033/53

(21) Application No: **199742608** (22) Application Date: **1997.09.10**

(87) WIPO No: **WO98/11218**

(30) Priority Data

(31) Number (32) Date (33) Country  
08/709838 1996.09.10 US  
08/829839 1997.03.31 US

(43) Publication Date : **1998.04.02**  
(43) Publication Journal Date : **1998.05.28**  
(44) Accepted Journal Date : **2001.06.07**

(71) Applicant(s)  
**Theodor-Kocher Institute; Millennium Pharmaceuticals, Inc.**

(72) Inventor(s)  
**Marcel Loetscher; Bernhard Moser; Shixin Qin; Charles R Mackay**

(74) Agent/Attorney  
**DAVIES COLLISON CAVE, 1 Little Collins Street, MELBOURNE VIC 3000**

(56) Related Art  
**A. MARCHESE ET AL. 1995 CLONING+CHROMOSOMAL MAPPING OF THREE NOVEL GENES GPR9,GPR10 GPR14 ENCODING RECEPTORS RELATED TO INTERLEUKIN 8, NEUROPEPTIDE Y ..., GENOMICS,29:335-344**



<b>(51) International Patent Classification<sup>6</sup> :</b> <b>C12N 15/12, C07K 14/715, C12N 5/10, C07K 14/19, 16/18, G01N 33/50, 33/53, A61K 38/17</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/11218</b> <b>(43) International Publication Date:</b> <b>19 March 1998 (19.03.98)</b>
<b>(21) International Application Number:</b> <b>PCT/US97/15915</b> <b>(22) International Filing Date:</b> <b>10 September 1997 (10.09.97)</b>  <b>(30) Priority Data:</b> 08/709,838 10 September 1996 (10.09.96) US 08/829,839 31 March 1997 (31.03.97) US			<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). <i>MILLENIUM PHARMA CENTRALS, Inc</i>
<b>(71) Applicants:</b> THEODOR-KOCHER INSTITUTE [CH/CH]; Freiestrasse 1, CH-3012 Bern (CH). <del>LEUKOSITE, INC.</del> [US/US]; 215 First Street, Cambridge, MA 02142 (US). 15 Sidney Street, 39			<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> <i>With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description.</i> <i>Date of receipt by the International Bureau:</i> 10 September 1997 (10.09.97)
<b>(72) Inventors:</b> LOETSCHER, Marcel; Sonnenweg 4, CH-3098 Koeniz (CH). MOSER, Bernhard; Kammstrasse 11, CH-3066 Stettlen (CH). QIN, Shixin; 14 Taft Avenue, Lexington, MA 02173 (US). MACKAY, Charles, R.; 126 Church Street, Watertown, MA 02172 (US).			
<b>(74) Agents:</b> BROOK, David, E. et al.; Hamilton, Brook, Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA 02173 (US).			

**(54) Title:** CXCR3 CHEMOKINE RECEPTOR, ANTIBODIES, NUCLEIC ACIDS, AND METHODS OF USE

**(57) Abstract**

The present invention relates to recombinant mammalian (e.g., human) IP-10/Mig receptor proteins designated CXC Chemokine Receptor 3 (CXCR3) and variants thereof, including those characterized by selective binding of one or more chemokines (e.g., IP-10 and/or Mig), and/or the ability to induce a cellular response (e.g., chemotaxis, exocytosis). Another aspect of the invention relates to antisense nucleic acid, recombinant nucleic acid constructs, such as plasmids or retroviral vectors, methods of identifying ligands, and inhibitors (e.g., antagonists) or promoters (e.g., agonists) of receptor function.

-1-

CXCR3 CHEMOKINE RECEPTOR, ANTIBODIES, NUCLEIC ACIDS, AND METHODS OF USE

Description

BACKGROUND

5       Chemokines constitute a family of small cytokines that are produced in inflammation and regulate leukocyte recruitment (Baggiolini, M. et al., *Adv. Immunol.* 55: 97-179 (1994); Springer, T.A., *Annu. Rev. Physiol.* 57: 827-872 (1995); and Schall, T.J. and K.B. Bacon, *Curr. Opin. Immunol.* 6: 865-873 (1994)). Chemokines are capable of selectively inducing chemotaxis of the formed elements of the blood (other than red blood cells), including leukocytes such as neutrophils, monocytes, macrophages, eosinophils, basophils, mast cells, and lymphocytes, such as T cells and B cells. In addition to stimulating chemotaxis, other changes can be selectively induced by chemokines in responsive cells, including changes in cell shape, transient rises in the concentration of intracellular free calcium ions ( $[Ca^{2+}]_i$ ), granule exocytosis, integrin upregulation, formation of bioactive lipids (e.g., leukotrienes) and respiratory burst, associated with leukocyte activation. Thus, the chemokines are early triggers of the inflammatory response, causing inflammatory mediator release, chemotaxis and extravasation to sites of infection or inflammation.

20       Two subfamilies of chemokines, designated as CXC and CC chemokines, are distinguished by the arrangement of the first two of four conserved cysteine residues, which are either separated by one amino acid (as in CXC chemokines IL-8,  $\gamma$ IP-10, Mig, PF4, ENA-78, GCP-2, GRO $\alpha$ , GRO $\beta$ , GRO $\gamma$ ,

-2-

NAP-2, NAP-4) or are adjacent residues (as in CC chemokines MIP-1 $\alpha$ , MIP-1 $\beta$ , RANTES, MCP-1, MCP-2, MCP-3, I-309). Most CXC chemokines attract neutrophil leukocytes. For example, the CXC chemokines interleukin 8 (IL-8), platelet factor 4 (PF4), and neutrophil-activating peptide 2 (NAP-2) are potent chemoattractants and activators of neutrophils. The CXC chemokines designated Mig (monokine induced by gamma interferon) and IP-10 ( $\gamma$ IP-10, interferon-gamma inducible 10 kDa protein) are particularly active in inducing 5 chemotaxis of activated peripheral blood lymphocytes. CC chemokines are generally less selective and can attract a variety of leukocyte cell types, including monocytes, eosinophils, basophils, T lymphocytes and natural killer cells. CC chemokines such as human monocyte chemotactic 10 proteins 1-3 (MCP-1, MCP-2 and MCP-3), RANTES (Regulated on Activation, Normal T Expressed and Secreted), and the 15 macrophage inflammatory proteins 1 $\alpha$  and 1 $\beta$  (MIP-1 $\alpha$  and MIP-1 $\beta$ ) have been characterized as chemoattractants and activators of monocytes or lymphocytes, but do not appear 20 to be chemoattractants for neutrophils.

CC and CXC chemokines act through receptors which belong to a superfamily of seven transmembrane spanning G protein-coupled receptors (Murphy, P.M., *Annu. Rev. Immunol.*, 12: 593-633 (1994); Gerard, C. and N.P. Gerard, 25 *Curr. Opin. Immunol.*, 6: 140-145 (1994)). This family of G-protein coupled (serpentine) receptors comprises a large group of integral membrane proteins, containing seven transmembrane-spanning regions. The receptors are coupled to G proteins, which are heterotrimeric regulatory proteins 30 capable of binding GTP and mediating signal transduction from coupled receptors, for example, by the production of intracellular mediators.

The chemokine receptors can be divided into two groups: CC chemokine receptors 1 through 5 (CCR1-5), which 35 bind CC chemokines, and CXC chemokine receptors 1 through 4

-3-

(CXCR1-4), which bind CXC chemokines. In general, the CC chemokine receptors occur on several types of leukocytes, and are important for the migration of monocytes, eosinophils, basophils, and T cells (Qin, S., et al., *Eur. J. Immunol.*, 26:640-647 (1996); Carr, M.W., et al., *Proc. Natl. Acad. Sci. USA*, 91(9):3652-3656 (1994); Taub, D.D., et al., *J. Clin. Invest.*, 95(3):1370-1376 (1995); Neote, K. et al., *Cell*, 72: 415-425 (1993); Gao, J.-L. et al., *J. Exp. Med.*, 177: 1421-1427 (1993); Charo, I.F. et al., *Proc. Natl. Acad. Sci. USA*, 91: 2752-2756 (1994); Myers, S.J., et al., *J. Biol. Chem.*, 270: 5786-5792 (1995); Combadiere, C. et al., *J. Biol. Chem.*, 270 (27): 16491-16494 (1995); and Correction, *J. Biol. Chem.*, 270: 30235 (1995); Ponath, P.D. et al., *J. Exp. Med.*, 183: 2437-15 2448 (1996); and Daugherty, B.L. et al., *J. Exp. Med.*, 183: 2349-2354 (1996); Power, C.A. et al., 1995, *J. Biol. Chem.*, 270: 19495-19500 (1995); Hoogewerf, A.J. et al., *Biochem. Biophys. Res. Commun.*, 218: 337-343 (1996); Samson, M. et al., *Biochemistry*, 35: 3362-3367 (1996)). In contrast, the 20 two IL-8 receptors, CXCR1 and CXCR2, are largely restricted to neutrophils and are important for the migration of neutrophils (Baggiolini, M., et al., *Adv. Immunol.*, 55:97-179 (1994)). The IL-8 receptors, CXCR1 (IL-8R1, interleukin-8 receptor type 1; Holmes, W.E. et al., 25 *Science*, 253: 1278-1280 (1991)) and CXCR2 (IL-8R2, interleukin-8 receptor type 2; Murphy, P.M. and H.L. Tiffany, *Science*, 253: 1280-1283 (1991)) recognize the NH<sub>2</sub>-terminal Glu-Leu-Arg (ELR) motif, an essential binding epitope observed in CXC chemokines that induce neutrophil 30 chemotaxis (Clark-Lewis, I. et al., *J. Biol. Chem.*, 266: 23128-23134 (1991); Hébert, C.A. et al., *J. Biol. Chem.*, 266: 18989-18994 (1991); and Clark-Lewis, I. et al., *Proc. Natl. Acad. Sci. USA*, 90: 3574-3577 (1993)). In contrast to monocytes and granulocytes, lymphocyte 35 responses to chemokines are not well understood. Notably,

- 4 -

none of the receptors of known specificity appear to be restricted to lymphocytes and the chemokines that recognize these receptors cannot, therefore, account for events such as the selective recruitment of T lymphocytes that is 5 observed in T cell-mediated inflammatory conditions. Moreover, although a number of proteins with significant sequence similarity and similar tissue and leukocyte subpopulation distribution to known chemokine receptors have been identified and cloned, the ligands for these 10 receptors remain undefined. Thus, these proteins are referred to as orphan receptors. The characterization of the ligand(s) of a receptor, is essential to an understanding of the interaction of chemokines with their target cells, the events stimulated by this interaction, 15 including chemotaxis and cellular activation of leukocytes, and the development of therapies based upon modulation of receptor function.

#### SUMMARY OF THE INVENTION

The present invention relates to proteins or 20 polypeptides, referred to herein as isolated and/or recombinant mammalian (e.g., a primate such as a human) IP-10/Mig receptor proteins designated CXC Chemokine Receptor 3 (CXCR3) and variants thereof. Recombinant CXCR3 proteins and variants can be produced in host cells as 25 described herein. In one embodiment, a CXCR3 protein or variant thereof is characterized by selective binding (e.g., high affinity binding) of one or more chemokines, such as IP-10 and/or Mig, and/or the ability to induce a (one or more) cellular response(s) (e.g., chemotaxis, 30 exocytosis, release of one or more inflammatory mediators).

Another aspect of the present invention relates to isolated and/or recombinant nucleic acids which encode a mammalian (e.g., a primate such as a human) CXCR3 protein or variant thereof. The invention further relates to

-5-

recombinant nucleic acid constructs, such as plasmids or retroviral vectors, which contain a nucleic acid which encodes a protein of the present invention or a variant thereof. The nucleic acids and constructs can be used to 5 produce recombinant receptor proteins and host cells comprising a construct. In another embodiment, the nucleic acid encodes an antisense nucleic acid which can hybridize with a second nucleic acid encoding a CXCR3 protein and which, when introduced into cells, can inhibit the 10 expression of receptor.

The invention further relates to antibodies reactive with CXCR3 receptors, which can be produced using the proteins or variants thereof (e.g., a peptide) or cells expressing receptor protein or variant as immunogen, for 15 example. Such antibodies or fragments thereof are useful in therapeutic, diagnostic and research applications, including the purification and study of the receptor proteins, identification of cells expressing surface receptor, and sorting or counting of cells. Thus, the 20 present invention encompasses use of an antibody or fragment thereof described herein (e.g., mAb 1C6 or an antigen-binding fragment thereof) in therapy (including prophylaxis) or diagnosis, and use of such antibodies or fragments for the manufacture of a medicament for use in 25 treatment of diseases or conditions as described herein.

Also encompassed by the present invention are methods of identifying ligands of the receptor, inhibitors (e.g., antagonists) or promoters (e.g., agonists) of receptor function. In one embodiment, suitable host cells which 30 have been engineered to express a receptor protein or variant encoded by a nucleic acid introduced into said cells are used in an assay to identify and assess the efficacy of ligands, inhibitors or promoters of receptor function. Such cells are also useful in assessing the 35 function of the expressed receptor protein or polypeptide.

- 6 -

According to the present invention, ligands, inhibitors and promoters of receptor function can be identified in a suitable assay, and further assessed for therapeutic effect. Inhibitors of receptor function can be 5 used to inhibit (reduce or prevent) receptor activity, and ligands and/or promoters can be used to induce (trigger or enhance) normal receptor function where indicated. Thus, the present invention provides a method of treating inflammatory diseases including autoimmune disease and 10 graft rejection, comprising administering an inhibitor of receptor function to an individual (e.g., a mammal). The present invention further provides a method of stimulating receptor function by administering a ligand or promoter to an individual, providing a new approach to selective 15 stimulation of leukocyte function, which is useful, for example, in the treatment of infectious diseases and cancer.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is an illustration of the nucleotide sequence 20 determined from the 1670 bp insert of a cDNA encoding a human IP-10/Mig receptor designated CXCR3, which was isolated from a human CD4<sup>+</sup> T cell (KT30) cDNA library (SEQ ID NO:1). An open reading frame (69-1175) encodes a predicted protein of 368 amino acids (SEQ ID NO:2). A 25 putative poly-A signal and poly A site are located at positions 1534-1539 and at 1624-1670, respectively.

Figure 2 is an illustration of the conceptual translation of the open reading frame of the sequence in Figure 1, which encodes a human IP-10/Mig receptor (SEQ ID 30 NO:2). Arrowheads indicate potential N-linked glycosylation sites and horizontal lines indicate the location of putative transmembrane domains (TM1-TM7).

Figures 3A-3C are graphs illustrating the responses induced by IP-10 and Mig in stably transfected cells

- 7 -

expressing IP-10/MigR. Figure 3A is a graph illustrating the concentration dependent  $[Ca^{2+}]_i$  changes in IP-10/MigR-transfected 300-19 cells. IP-10 or Mig were each added at 1, 10, and 100 nM to Fura-2/AM loaded cells (arrowhead), 5 and  $[Ca^{2+}]_i$ -dependent fluorescence changes were recorded. Non-transfected cells (lower tracings) were stimulated with IP-10 or Mig at 100 nM under identical conditions.

Figure 3B is a graph illustrating the results of studies assessing receptor desensitization and cross- 10 desensitization, in which IP-10/MigR expressing 300-19 cells were sequentially stimulated with 100 nM IP-10 or Mig, and with IP-10 followed by Mig or vice versa, and fluorescence changes were recorded.

Figure 3C is a graph illustrating the chemotaxis of 15 IP-10/MigR expressing Jurkat cells stimulated with IP-10 (filled circles) or Mig (filled squares). The lower panel shows the response of non-transfected Jurkat cells when stimulated with increasing amounts of IP-10 (open circles) or Mig (open squares). Mean numbers ( $\pm$  SD) of migrating 20 cells per five high power fields are presented.

Figures 4A-4B are graphs illustrating the responses of peripheral blood lymphocytes (PBL) to IP-10 and Mig. Freshly isolated PBL from donor blood buffy coats were used as such (lower tracings and open symbols), or were used 25 after culturing for 10 days in the presence of IL-2 (400 U/ml) (upper tracings and closed symbols). Figure 4A is a graph illustrating  $[Ca^{2+}]_i$  changes induced by IP-10 or Mig. IP-10 or Mig were each added at 1, 10, and 100 nM to Fura-2/AM loaded cultured cells (arrowhead), and  $[Ca^{2+}]_i$ - 30 dependent fluorescence changes were recorded (upper tracings). Freshly isolated cells (lower tracings) were stimulated with IP-10 or Mig at 100 nM under the same conditions. Figure 4B is a graph illustrating chemotaxis of PBL in response to increasing concentrations of IP-10 35 (filled circles) or Mig (filled squares) (mean numbers

- 8 -

( $\pm$  SD) of migrating cells per five high power fields are presented).

Figures 5A-5B are graphs illustrating the binding of radiolabeled IP-10 to either L1.2 cells transfected with 5 CXCR3 DNA (Figure 5A) and to activated T cells (Figure 5B). Cells were incubated with 0.05 nM  $^{125}$ I-labeled IP-10 in the presence of increasing concentrations of unlabeled IP-10. Scatchard analysis (inset) indicated 37,000 receptors per cell (Kd of 614 pM) for CXCR3 L1.2 transfectants, and 10 17,000 receptors per cell (Kd of 156 pM) for CD3 blasts.

Figure 6 is an illustration of the specificity of anti-CXCR3 antibody 1C6 as assessed by staining of stable L1.2 transfectants expressing either CCR1, CCR2b, CCR3, CCR4, CCR5, CXCR1, CXCR2, CXCR3 or CXCR4 by anti-CXCR3 15 peptide mAb 1C6. Negative control staining for all the L1.2 transfectants (not shown) resembled the staining shown for 1C6 on untransfected L1.2 cells (L1.2 wt).

Figures 7A-7C are fluorescence histograms illustrating the expression of CXCR3 on neutrophils (Figure 7A), 20 lymphocytes (Figure 7B), and activated T cells (Figure 7C). Leukocyte subsets were identified in whole blood by their forward angle and side scatter, and were gated accordingly. To generate CD3 blasts, PBMC were activated with anti-CD3 25 mAb for 3 days, and were then maintained in media containing IL-2 for 7 days. In each plot, the blackened profile represents staining with anti-CXCR3 mAb 1C6, and the unfilled profile represent staining with an isotype-matched control mAb.

Figure 8 is a series of plots illustrating CXCR3 30 expression on populations of blood lymphocytes. A two color staining protocol was used to assess expression of CXCR3 on T cells (CD3), B cells (CD20), and NK cells (CD56).

Figure 9 is a series of plots illustrating CXCR3 35 expression versus various markers on the CD3+ subset of

-9-

blood lymphocytes, as analyzed by three color analysis of immunofluorescence. Anti-CD3 Cy-Chrome was used to stain T cells, and these cells were gated electronically for analysis. Quadrants were set according to the staining of 5 control mAbs. The staining shown was representative of five donors analyzed.

Figure 10 is a histogram illustrating inhibition of IP-10- or MCP-1-mediated chemotaxis of activated T cells by a panel of anti-CXCR3 mAbs.  $1 \times 10^6$  human CD3 blasts were 10 placed in the top chamber of a transwell and chemokine (12.5 nM) was placed in the bottom chamber. Various anti-CXCR3 mAbs (in tissue culture supernatant, without FCS) were placed in the top well with cells at the beginning of 15 the assay. After 1.5 hours the cells migrating to the bottom chamber were counted using flow cytometry. The percentage inhibition of chemotaxis was calculated using the number of cells that migrated in the absence of mAb as 100%. The results are representative of at least four separate experiments.

20 Figure 11 is a graph illustrating the inhibition of IP-10 mediated chemotaxis by purified anti-CXCR3 mAb 1C6. Various concentrations of 1C6 mAb were placed in the top well, and the assay was performed as described for Figure 10. mAb 1C6 inhibited 50% total chemotaxis at a 25 concentration of 856 ng/ml ( $IC_{50} = 856$  ng/ml).

Figure 12 is a graph illustrating the inhibition of  $^{125}\text{I}$ -IP-10 binding to activated T cells by mAb 1C6. CD3 blasts were incubated with 0.05 nM  $^{125}\text{I}$ -IP-10 in the presence of increasing concentrations of 1C6 as indicated. 30 After 60 minutes at room temperature, cell pellets were washed and counted. Data was analyzed by KaleidaGraph, which gave an  $IC_{50}$  of 0.16  $\mu\text{g/ml}$ .

Figures 13A-13H illustrate the inhibition by mAb 1C6 of  $[\text{Ca}^{2+}]_i$  by human T cells in response to IP-10, but not 35 Mig. Anti-CD3 activated, IL-2-stimulated human T cells

-10-

were labeled with Fura-2, and were stimulated sequentially with the indicated chemokines (Figures 13A-13B), or with mAb followed 40 seconds later by the indicated chemokine (Figures 13C-13H).  $[Ca^{2+}]_i$  fluorescence changes were recorded using a spectrofluorimeter. The tracings were representative of five separate experiments. Antibody was used at a final concentration of either 50  $\mu$ g/ml (Figures 13C-13D); 25  $\mu$ g/ml (Figure 13E); 12.5  $\mu$ g/ml (Figure 13F); 6.125  $\mu$ g/ml (Figure 13G); or 3.0625  $\mu$ g/ml (Figure 13H).

5 Chemokines were used at 2 nM.

10 Figures 14A-14D are fluorescence histograms illustrating the results of a flow cytometry analysis in which CXCR3-expressing transfectants were stained with mAb 1C6 in the presence of P1 peptide (Figure 14B), P2 peptide (Figure 14C), P3 peptide (Figure 14D), or in the absence of peptide (Figure 14A). In each plot, the profile defined by the heavy line represents mAb 1C6 staining, and the profile defined by the dotted line represents staining with an isotype-matched irrelevant control mAb.

15

20 Figure 15 is a histogram illustrating the percent inhibition of binding of radiolabeled IP-10 to CXCR3 transfectants by 40 nM cold IP-10, mAb 1C6, monoclonal antibodies raised against CXCR3 transfectants (2F8, 3A12, 3E2, 4B4, 4D2, 5B12, 7B8, or 8D5), or by anti-CXCR2 mAb.

25 DETAILED DESCRIPTION OF THE INVENTION

As described herein, a nucleic acid encoding a novel chemokine receptor that is selective for the CXC chemokines IP-10 and Mig was cloned and characterized. The clone, which was isolated from a human CD4 $^{+}$  T cell library, was not detected in monocyte- or granulocyte-derived cDNA libraries. Sequence analysis of the clone revealed an open reading frame of 1104 base pairs (Figure 1, SEQ ID NO:1), encoding a predicted protein of 368 amino acids with a predicted molecular mass of 40,659 daltons (Figure 2, SEQ

30

-11-

ID NO:2). The amino acid sequence includes seven putative transmembrane segments which are characteristic of G-protein coupled receptors and are found in other chemoattractant receptors. Consistent with this 5 observation, the receptor mediates  $\text{Ca}^{2+}$  (calcium ion) mobilization and chemotaxis in response to IP-10 and Mig (Example 2). No significant response to the CXC chemokines IL-8, GRO $\alpha$ , NAP-2 (neutrophil-activating protein-2), GCP-2 (granulocyte chemotactic protein-2), ENA78 10 (epithelial-derived neutrophil-activating peptide 78), PF4 (platelet factor 4), or the CC chemokines MCP-1 (monocyte chemotactic protein-1), MCP-2, MCP-3, MCP-4, MIP-1 $\alpha$  (macrophage inflammatory protein-1 $\alpha$ ), MIP-1 $\beta$ , RANTES (regulated on activation, normal T cell expressed and 15 secreted), I309, eotaxin or lymphotactin was detected under similar conditions.

The restricted expression of human CXCR3 in activated T lymphocytes and the ligand selectivity of the receptor for IP-10 and Mig are noteworthy. The human receptor is 20 highly expressed in IL-2 activated T lymphocytes, but was not detected in resting T lymphocytes, B lymphocytes, monocytes or granulocytes under the conditions used in Example 2. Additional studies of receptor distribution indicate that it is mostly CD3+ cells that express CXCR3, 25 including cells which are CD95+, CD45RO+, and CD45RA<sup>low</sup>, a phenotype consistent with previous activation, although a proportion of CD20+ (B) cells and CD56+ (NK) cells also express this receptor. The selective expression in activated T lymphocytes is of interest, because other 30 receptors for chemokines which have been reported to attract lymphocytes (e.g., MCP-1, MCP-2, MCP-3, MIP-1 $\alpha$ , MIP-1 $\beta$  and RANTES) are also found in granulocytes, such as neutrophils, eosinophils and basophils, as well as monocytes. These results suggest that the IP-10/Mig

-12-

receptor designated CXCR3 is involved in the selective recruitment of effector T cells.

The receptor recognizes two unusual CXC chemokines, designated IP-10 and Mig. Although IP-10 and Mig both belong to the CXC subfamily, in contrast to IL-8 and other CXC chemokines which are potent chemoattractants for neutrophils, the primary targets of IP-10 and Mig are lymphocytes, particularly effector cells such as activated or stimulated T lymphocytes and natural killer (NK) cells

10 (Taub, D.D. et al., *J. Exp. Med.*, 177: 18090-1814 (1993); Taub, D.D. et al., *J. Immunol.*, 155: 3877-3888 (1995)). (NK cells are large granular lymphocytes, which lack a specific T cell receptor for antigen recognition, but possess cytolytic activity against cells such as tumor

15 cells and virally infected cells.) Consistently, IP-10 and Mig lack the ELR motif, an essential binding epitope in those CXC chemokines that efficiently induce neutrophil chemotaxis (Clark-Lewis, I. et al., *J. Biol. Chem.*, 266: 23128-23134 (1991); Hébert, C.A. et al., *J. Biol. Chem.*,

20 266: 18989-18994 (1991); and Clark-Lewis, I. et al., *Proc. Natl. Acad. Sci. USA*, 90: 3574-3577 (1993)). In addition, both recombinant human Mig and recombinant human IP-10 have been reported to induce calcium flux in tumor infiltrating lymphocytes (TIL) (Liao, F. et al., *J. Exp. Med.*, 182: 1301-1314 (1995)). While IP-10 has been reported to induce chemotaxis of monocytes *in vitro* (Taub, D.D. et al., *J. Exp. Med.*, 177: 1809-1814 (1993), the receptor responsible has not been identified), human Mig appears highly selective, and does not show such an effect (Liao, F. et

25 al., *J. Exp. Med.*, 182: 1301-1314 (1995)). IP-10 expression is induced in a variety of tissues in inflammatory conditions such as psoriasis, fixed DRUG eruptions, cutaneous delayed-type hypersensitivity responses, tuberculoid leprosy, and in experimental

30 glomerulonephritis, and experimental allergic

35

-13-

encephalomyelitis. IP-10 also has a potent *in vivo* antitumor effect that is T cell dependent, is reported to be an inhibitor of angiogenesis *in vivo*, and can induce chemotaxis and degranulation of NK cells *in vitro*,

5 suggesting a role as a mediator of NK cell recruitment and degranulation (in tumor cell destruction, for example) (Luster, A.D. and P. Leder, *J. Exp. Med.*, 178: 1057-1065 (1993); Luster, A.D. et al., *J. Exp. Med.* 182: 219-231 (1995); Angiolillo, A.L. et al., *J. Exp. Med.*, 182: 155-162 (1995); Taub, D.D. et al., *J. Immunol.*, 155: 3877-3888 (1995)). The expression patterns of IP-10 and Mig are also distinct in that expression of each is induced by interferon-gamma (IFN $\gamma$ ), while the expression of IL-8 is down-regulated by IFN $\gamma$  (Luster, A.D. et al., *Nature*, 315: 15 672-676 (1985); Farber, J.M., *Proc. Natl. Acad. Sci. USA*, 87: 5238-5242 (1990); Farber, J.M., *Biochem. Biophys. Res. Commun.*, 192 (1): 223-230 (1993), Liao, F. et al., *J. Exp. Med.*, 182: 1301-1314 (1995); Seitz, M. et al., *J. Clin. Invest.*, 87: 463-469 (1991); Galy, A.H.M. and H. Spits, *J. 20 Immunol.*, 147: 3823-3830 (1991)).

Chemokines have been recently recognized as the long-sought mediators for the recruitment of lymphocytes. Several CC chemokines were found to elicit lymphocyte chemotaxis (Loetscher, P. et al., *FASEB J.*, 8: 1055-1060 (1994)), but they are also active on granulocytes and monocytes (Uguzzioni, M. et al., *Eur. J. Immunol.*, 25: 25 64-68 (1995); Baggiolini, M. and C.A. Dahinden, *Immunol. Today*, 15: 127-133 (1994)). The situation is different for IP-10 and Mig, which are selective in their action on 30 lymphocytes, including activated T lymphocytes and NK cells, and which bind CXCR3, a receptor which does not recognize numerous other chemokines and which displays a selective pattern of expression (Example 2, Example 5).

In view of these observations, it is reasonable to 35 conclude that the formation of the characteristic

-14-

infiltrates in inflammatory lesions, such as delayed-type hypersensitivity lesions, sites of viral infection, and certain tumors is a process mediated by via CXCR3 and regulated by CXCR3 expression. Lymphocytes, particularly T 5 lymphocytes, bearing a CXCR3 receptor as a result of activation can be recruited into inflammatory lesions, sites of infection, or tumors by IP-10 and/or Mig, which can be induced locally by interferon-gamma. Thus, CXCR3 plays a role in the selective recruitment of lymphocytes, 10 particularly effector cells such as activated or stimulated T lymphocytes.

#### Proteins and Peptides

The present invention relates to isolated and/or recombinant (including, e.g., essentially pure) proteins or 15 polypeptides designated mammalian CXCR3 proteins and variants thereof. In a preferred embodiment, the isolated and/or recombinant proteins of the present invention have at least one property, activity or function characteristic of a mammalian CXCR3 protein (as defined herein), such as a 20 binding activity (e.g., ligand, inhibitor and/or promoter binding), a signalling activity (e.g., activation of a mammalian G protein, induction of rapid and transient increase in the concentration of cytosolic free calcium  $[Ca^{2+}]_i$ ), cellular response function (e.g., stimulation of 25 chemotaxis, exocytosis or inflammatory mediator release by leukocytes), and/or an immunological property as defined herein. For example, some proteins of the present invention can selectively bind to IP-10 and/or Mig, mediate cellular signalling and/or a response thereto *in vitro* 30 and/or *in vivo* (e.g., calcium flux, chemotaxis and/or degranulation especially of activated T lymphocytes). For example, as shown herein, a human CXCR3 protein, produced in mammalian cells by expression of a cDNA clone, can selectively bind to CXC chemokines IP-10 and/or Mig, and

-15-

mediate signalling, and a cellular response (e.g., chemotaxis). In one embodiment, proteins of the present invention can bind a CXC chemokine from the same or a different mammalian species (e.g., human IP-10, murine 5 IP-10, human Mig, murine Mig) (human IP-10, Luster, A.D. et al., *Nature*, 315: 672-676 (1985); murine IP-10 (also referred to as CRG-2), Vanguri, P. and J.M. Farber, *J. Biol. Chem.*, 265: 15049 (1990) and Luster, A.D. and P. Leder, *J. Exp. Med.*, 178: 1057-1065 (1993); murine Mig, 10 Farber, J.M., *Proc. Natl. Acad. Sci. USA*, 87: 5238-5242 (1990); human Mig, Farber, J.M., *Biochem. Biophys. Res. Commun.*, 192 (1): 223-230 (1993) and Liao, F. et al., *J. Exp. Med.*, 182: 1301-1314 (1995)).

Proteins or polypeptides referred to herein as 15 "isolated" are proteins or polypeptides purified to a state beyond that in which they exist in mammalian cells, and include proteins or polypeptides obtained by methods described herein, similar methods or other suitable methods, including essentially pure proteins or 20 polypeptides, proteins or polypeptides produced by chemical synthesis (e.g., synthetic peptides), or by combinations of biological and chemical methods, and recombinant proteins or polypeptides which are isolated. The proteins can be obtained in an isolated state of at least about 50 % by 25 weight, preferably at least about 75 % by weight, or in essentially pure form. Proteins or polypeptides referred to herein as "recombinant" are proteins or polypeptides produced by the expression of recombinant nucleic acids.

As used herein "mammalian CXCR3 protein" refers to 30 naturally occurring or endogenous mammalian CXCR3 proteins and to proteins having an amino acid sequence which is the same as that of a naturally occurring or endogenous corresponding mammalian CXCR3 protein (e.g., recombinant proteins). Accordingly, as defined herein, the term 35 "mammalian CXCR3 protein" includes mature protein,

-16-

polymorphic or allelic variants, and other isoforms of mammalian CXCR3 (e.g., produced by alternative splicing or other cellular processes), and modified or unmodified forms of the foregoing (e.g., glycosylated, unglycosylated, 5 phosphorylated or unphosphorylated CXCR3 proteins). Naturally occurring or endogenous mammalian CXCR3 proteins include wild type proteins such as mature CXCR3, polymorphic or allelic variants and other isoforms which occur naturally in mammals (e.g., humans, non-human 10 primates). Such proteins can be recovered from a source which naturally produces mammalian CXCR3, for example. These proteins and mammalian CXCR3 proteins having the same amino acid sequence as a naturally occurring or endogenous corresponding mammalian CXCR3, are referred to by the name 15 of the corresponding mammal. For example, where the corresponding mammal is a human, the protein is designated as a human CXCR3 protein (e.g., a recombinant human CXCR3 produced in a suitable host cell).

"Functional variants" of mammalian CXCR3 proteins 20 include functional fragments, functional mutant proteins, and/or functional fusion proteins (e.g., produced via mutagenesis and/or recombinant techniques). Generally, fragments or portions of mammalian CXCR3 proteins encompassed by the present invention include those having a 25 deletion (i.e., one or more deletions) of an amino acid (i.e., one or more amino acids) relative to the mature mammalian CXCR3 protein (such as N-terminal, C-terminal or internal deletions). Fragments or portions in which only contiguous amino acids have been deleted or in which 30 non-contiguous amino acids have been deleted relative to mature mammalian CXCR3 protein are also envisioned.

Generally, mutants or derivatives of mammalian CXCR3 proteins, encompassed by the present invention include natural or artificial variants differing by the addition, 35 deletion and/or substitution of one or more contiguous or

-17-

non-contiguous amino acid residues, or modified polypeptides in which one or more residues is modified, and mutants comprising one or more modified residues.

Preferred mutants are natural or artificial variants of 5 mammalian CXCR3 proteins differing by the addition, deletion and/or substitution of one or more contiguous or non-contiguous amino acid residues. Such mutations can be in a conserved region or nonconserved region (compared to other CXC and/or CC chemokine receptors), extracellular, 10 cytoplasmic, or transmembrane region, for example.

A "functional fragment or portion", "functional mutant" and/or "functional fusion protein" of a mammalian CXCR3 protein refers to an isolated and/or recombinant protein or oligopeptide which has at least one property, 15 activity or function characteristic of a mammalian CXCR3 receptor (as defined herein), such as a binding activity (e.g., ligand, inhibitor and/or promoter binding), a signalling activity (e.g., activation of a mammalian G protein, induction of rapid and transient increase in the 20 concentration of cytosolic free calcium  $[Ca^{2+}]_i$ ), cellular response function (e.g., stimulation of chemotaxis, exocytosis or inflammatory mediator release by leukocytes), and/or an immunological property as defined herein.

As used herein, a protein or polypeptide having "at 25 least one immunological property" of a mammalian CXCR3 protein is one which (a) is bound by at least one antibody of a selected epitopic specificity which binds to a naturally occurring or endogenous mammalian CXCR3 protein or to a protein having the same amino acid sequence as the 30 naturally occurring or endogenous mammalian CXCR3 protein (e.g., human CXCR3), and/or (b) is an immunogen capable of inducing the formation (e.g., when conjugated to a suitable carrier) in a suitable animal of an antibody of a selected epitopic specificity which binds to a naturally occurring 35 or endogenous mammalian CXCR3 or to a protein having the

-18-

same amino acid sequence as the naturally occurring or endogenous mammalian CXCR3. For example, a suitable fragment can cross-react with an antibody which is raised against and/or reactive with isolated mammalian CXCR3.

5        Suitable fragments or mutants can be identified by screening. For example, the N-terminal, C-terminal, or internal regions of the protein can be deleted in a step-wise fashion and the resulting protein or polypeptide can be screened using a suitable assay, such as an assay  
10      described herein (e.g., chemotaxis, calcium flux). Where the resulting protein displays activity in the assay, the resulting protein ("fragment") is functional. Information regarding the structure and function of mammalian G protein coupled receptors, including CXC chemokine and CC chemokine  
15      receptors, provides a basis for dividing mammalian CXCR3 proteins into functional domains (Murphy, P.M., *Annu. Rev. Immunol.*, 12: 593-633 (1994) and Gerard, C. and N.P. Gerard, *Curr. Opin. Immunol.*, 6: 140-145 (1994), and references cited therein).

20       The term variant also encompasses fusion proteins, comprising a mammalian CXCR3 protein (e.g., human CXCR3) as a first moiety, linked to a second moiety not occurring in the mammalian CXCR3 as found in nature. Thus, the second moiety can be an amino acid, oligopeptide or  
25      polypeptide. The first moiety can be in an N-terminal location, C-terminal location or internal to the fusion protein. In one embodiment, the fusion protein comprises an affinity ligand (e.g., an enzyme, an antigen, epitope tag) as the first moiety, and a second moiety comprising a  
30      linker sequence and human CXCR3 or portion thereof.

35       Examples of mammalian CXCR3 proteins include proteins encoded by a nucleic acid of the present invention, such as a protein having an amino acid sequence as set forth or substantially as set forth in Figure 2 (SEQ ID NO:2). In a preferred embodiment, a mammalian CXCR3 or variant (e.g., a

-19-

variant including the extracellular N-terminal segment) has an amino acid sequence which is at least about 50% identical, more preferably at least about 70% identical, and still more preferably at least about 80 % identical, to 5 the protein shown in Figure 2 (SEQ ID NO:2).

It will be appreciated that isolated and/or recombinant mammalian CXCR3 proteins and variants thereof can be modified, for example, by incorporation of or attachment (directly or indirectly (e.g., via a linker)) of 10 a detectable label such as a radioisotope, spin label, antigen (e.g., epitope label such as a FLAG tag) or enzyme label, flourescent or chemiluminescent group and the like, and such modified forms are included within the scope of the invention.

15 Nucleic Acids, Constructs and Vectors

The present invention relates to isolated and/or recombinant (including, e.g., essentially pure) nucleic acids having sequences which encode a mammalian (e.g., human) CXCR3 protein or variant thereof as described 20 herein. Nucleic acids referred to herein as "isolated" are nucleic acids separated away from the nucleic acids of the genomic DNA or cellular RNA of their source of origin (e.g., as it exists in cells or in a mixture of nucleic acids such as a library), and may have undergone further 25 processing. "Isolated" nucleic acids include nucleic acids obtained by methods described herein, similar methods or other suitable methods, including essentially pure nucleic acids, nucleic acids produced by chemical synthesis, by combinations of biological and chemical methods, and 30 recombinant nucleic acids which are isolated. Nucleic acids referred to herein as "recombinant" are nucleic acids which have been produced by recombinant DNA methodology, including those nucleic acids that are generated by procedures which rely upon a method of artificial

- 20 -

recombination, such as the polymerase chain reaction (PCR) and/or cloning into a vector using restriction enzymes.

"Recombinant" nucleic acids are also those that result from recombination events that occur through the natural

5 mechanisms of cells, but are selected for after the introduction to the cells of nucleic acids designed to allow and make probable a desired recombination event.

In one embodiment, the nucleic acid or portion thereof encodes a protein or polypeptide having at least one

10 function characteristic of a mammalian CXCR3 protein (e.g., a human CXCR3 receptor), such as a binding activity (e.g., ligand, inhibitor and/or promoter binding), a signalling activity (e.g., activation of a mammalian G protein, induction of rapid and transient increase in the

15 concentration of cytosolic free calcium  $[Ca^{2+}]_i$ ), and/or stimulation of a cellular response (e.g., stimulation of chemotaxis, exocytosis or inflammatory mediator release by leukocytes). The present invention also relates more specifically to isolated and/or recombinant nucleic acids

20 or a portion thereof comprising sequences which encode a mammalian CXCR3 receptor or a portion thereof. The present invention relates even more specifically to isolated and/or recombinant nucleic acids comprising sequences which encode a human CXCR3 protein.

25 The invention further relates to isolated and/or recombinant nucleic acids, including double or single stranded DNA or RNA, that are characterized by (1) their ability to hybridize to: (a) a nucleic acid having the sequence SEQ ID NO:1, (b) a nucleic acid having a sequence

30 which is complementary to SEQ ID NO:1, or (c) a portion of the foregoing comprising the open reading frame of SEQ ID NO:1 (a portion of the strand illustrated in Figure 1 or the corresponding portion of the complementary strand); and/or (2) by their ability to encode a polypeptide having

35 the amino acid sequence SEQ ID NO:2 or a functional

-21-

equivalent thereof (i.e., a polypeptide having ligand binding activity for one or more natural or physiological ligand(s) of the receptor and/or stimulatory function responsive to ligand binding, such that it can induce a 5 cellular response (e.g., induction (including triggering or stimulation) of chemotaxis, exocytosis or inflammatory mediator release by leukocytes); and/or (3) by both characteristics.

In one embodiment, the percent amino acid sequence 10 identity between SEQ ID NO:2 and functional equivalents thereof is at least about 60% ( $\geq 60\%$ ). In a preferred embodiment, functional equivalents of SEQ ID NO:2 share at least about 70% sequence identity with SEQ ID NOS:2. More preferably, the percent amino acid sequence identity 15 between SEQ ID NO:2 and functional equivalents thereof is at least about 80%, and still more preferably, at least about 90%.

Isolated and/or recombinant nucleic acids meeting 20 these criteria comprise nucleic acids having sequences identical to sequences of naturally occurring mammalian CXCR3 receptors and portions thereof, or variants of the naturally occurring sequences. Such variants include mutants differing by the addition, deletion or substitution of one or more residues, modified nucleic acids in which 25 one or more residues is modified (e.g., DNA or RNA analogs), and mutants comprising one or more modified residues. In one embodiment, the nucleic acid shares at least about 50% nucleotide sequence similarity, more preferably at least about 75% nucleotide sequence 30 similarity, and still more preferably at least about 90% nucleotide sequence similarity, with one strand of the sequence illustrated in SEQ ID NO:1 or to the coding region thereof. Preferred nucleic acids have lengths of at least about 40 nucleotides, more preferably at least about 50, 35 and still more preferably at least about 75 nucleotides.

-22-

Such nucleic acids can be detected and isolated by hybridization under high stringency conditions or moderate stringency conditions, for example. "High stringency conditions" and "moderate stringency conditions" for 5 nucleic acid hybridizations are explained on pages 2.10.1-2.10.16 (see particularly 2.10.8-11) and pages 6.3.1-6 in *Current Protocols in Molecular Biology* (Ausubel, F.M. et al., eds., Vol. 1, Suppl. 26, 1991), the teachings of which are incorporated herein by reference. Factors such as 10 probe length, base composition, percent mismatch between the hybridizing sequences, temperature and ionic strength influence the stability of nucleic acid hybrids. Thus, high or moderate stringency conditions can be determined empirically to achieve the desired selectivity.

15 Isolated and/or recombinant nucleic acids that are characterized by their ability to hybridize to a nucleic acid having the sequence SEQ ID NO:1 or the complement thereof (e.g., under high or moderate stringency conditions) may further encode a protein or polypeptide 20 having at least one function characteristic of a mammalian CXCR3 protein (e.g., a human CXCR3 protein), such as a binding activity (e.g., ligand, inhibitor and/or promoter binding), a signalling activity (e.g., activation of a mammalian G protein, induction of rapid and transient 25 increase in the concentration of cytosolic free calcium  $[Ca^{2+}]_i$ ), and/or stimulation of a cellular response (e.g., stimulation of chemotaxis, exocytosis or inflammatory mediator release by leukocytes).

The human CXCR3 nucleic acid described herein, or 30 sufficient portions thereof, whether isolated, recombinant and/or synthetic, including fragments produced by PCR, can be used as probes or primers to detect and/or recover nucleic acids (e.g., genomic DNA, allelic variants, cDNA) encoding CXCR3 receptors (homologs) or other related 35 receptor genes (e.g., novel CXC chemokine receptor genes)

-23-

from other mammalian species including, but not limited to primates (e.g., a primate other than a human, such as a monkey (e.g., cynomolgus monkey)), bovine, ovine, equine, canine, feline and rodent (e.g., guinea pig, murine species such as rat, mouse). This can be achieved using the procedures described herein or other suitable methods, including hybridization, PCR or other suitable techniques. Mammalian nucleic acids can be used to prepare constructs (e.g., vectors), receptor or fragments thereof, and host strains useful in the production and methods of use of receptor.

In one embodiment, a nucleic acid encoding a mammalian CXCR3 protein (or variant) is producible by methods such as PCR amplification. For example, appropriate primers (e.g., 15 a pair of primers or nested primers) can be designed which comprise a sequence which is complementary or substantially complementary to a portion of the human CXCR3 cDNA described herein. For instance, primers complementary to the 5'- or 3'-ends of the coding sequence and/or flanking 20 the coding sequence can be designed. Such primers can be used in a polymerase chain reaction with a suitable template nucleic acid to obtain nucleic acid encoding a mammalian CXCR3, for example. Suitable templates include e.g., constructs described herein (such as pcDNA3-Clone8), 25 a cDNA or genomic library or another suitable source of mammalian (e.g., a human, primate) cDNA or genomic DNA. Primers can contain portions complementary to flanking sequences of a construct selected as template as appropriate.

30 The binding function of a protein or polypeptide (e.g., encoded by hybridizing nucleic acid) can be detected in binding or binding inhibition assays, using membrane fractions containing receptor or cells expressing receptor, for example (see e.g., Van Riper et al., *J. Exp. Med.*, 177: 35 851-856 (1993); Sledziewski et al., U.S. Patent No.

- 24 -

5,284,746 (Feb. 8, 1994)). Thus, the ability of the encoded protein or polypeptide to bind a ligand, such as IP-10 or Mig, an inhibitor and/or promoter, can be assessed. The antigenic properties of proteins or 5 polypeptides encoded by nucleic acids of the present invention can be determined by immunological methods employing antibodies that bind to a mammalian CXCR3, such as immunoblotting, immunoprecipitation and immunoassay (e.g., radioimmunoassay, ELISA).

10 The signalling function of a protein or polypeptide (e.g., encoded by hybridizing nucleic acid) can be detected by enzymatic assays for G protein activity responsive to receptor binding (e.g., exchange of GTP for GDP on the G protein  $\alpha$  subunit, using membrane fractions). G protein 15 coupling can be further assessed, for example, using assays in which stimulation by G protein is blocked by treatment or pre-treatment of cells or a suitable cellular fraction (e.g., membranes) with specific inhibitors of G proteins, such as *Bordetella pertussis* toxin (Bischoff, S.C. et al., 20 *Eur. J. Immunol.*, 23: 761-767 (1993); Sozzani, S. et al., *J. Immunol.*, 147: 2215-2221 (1991)).

The stimulatory function of a protein or polypeptide (e.g., encoded by hybridizing nucleic acid) can be detected by standard assays for chemotaxis or mediator release, 25 using cells expressing the protein or polypeptide (e.g., assays which monitor chemotaxis, exocytosis (e.g., degranulation of enzymes, such as esterases (e.g., serine esterases), perforin, granzymes) or mediator release (e.g., histamine, leukotriene) in response to a ligand (e.g., a 30 chemokine such as IP-10 or Mig) or a promoter (see e.g., Taub, D.D. et al., *J. Immunol.*, 155: 3877-3888 (1995); Bagliolini, M. and C.A. Dahinden, *Immunology Today*, 15: 127-133 (1994) and references cited therein). Functions characteristic of a mammalian CXCR3 receptor may also be 35 assessed by other suitable methods.

-25-

These methods, alone or in combination with other suitable methods can also be used in procedures for the identification and/or isolation of nucleic acids which encode a polypeptide having the amino acid sequence SEQ ID

5 NO:2 or functional equivalents thereof, and having an activity detected by the assay. Portions of isolated nucleic acids which encode polypeptide portions of SEQ ID NO:2 having a certain function can be also identified and isolated in this manner.

10 Nucleic acids of the present invention can be used in the production of proteins or polypeptides. For example, a nucleic acid containing all or part of the coding sequence for a mammalian CXCR3 receptor, or DNA which hybridizes to the sequence SEQ ID NO:1, or the complement thereof, can be  
15 incorporated into a construct for further manipulation of sequences or for production of the encoded polypeptide in suitable host cells. Nucleic acids of the present invention can also be modified, for example, by incorporation of or attachment (directly or indirectly) of  
20 a detectable label such as a radioisotope, spin label, antigen or enzyme label, fluorescent or chemiluminescent group and the like, and such modified forms are included within the scope of the invention.

#### Antisense Constructs

25 In another embodiment, the nucleic acid is an antisense nucleic acid, which is complementary, in whole or in part, to a target molecule comprising a sense strand, and can hybridize with the target molecule. The target can be DNA, or its RNA counterpart (i.e., wherein T residues of  
30 the DNA are U residues in the RNA counterpart). When introduced into a cell using suitable methods, antisense nucleic acid can inhibit the expression of the gene encoded by the sense strand. Antisense nucleic acids can be produced by standard techniques.

-26-

In one embodiment, the antisense nucleic acid is wholly or partially complementary to and can hybridize with a target nucleic acid, wherein the target nucleic acid can hybridize to a nucleic acid having the sequence of the 5 complement of SEQ ID NO:1. For example, antisense nucleic acid can be complementary to a target nucleic acid having the sequence of SEQ ID NO:1 or a portion thereof sufficient to allow hybridization. In another embodiment, the antisense nucleic acid is wholly or partially complementary 10 to and can hybridize with a target nucleic acid which encodes a mammalian CXCR3 receptor (e.g., human IP-10/Mig receptor CXCR3).

Antisense nucleic acids are useful for a variety of purposes, including research and therapeutic applications. 15 For example, a construct comprising an antisense nucleic acid can be introduced into a suitable cell to inhibit receptor expression. Such a cell provides a valuable control cell, for instance in assessing the specificity of receptor-ligand interaction with the parent cell or other 20 related cell types. In another aspect, such a construct can be introduced into some or all of the cells of a mammal. The antisense nucleic acid inhibits receptor expression, and inflammatory processes mediated by CXCR3 receptors in the cells containing the construct can be 25 inhibited. Thus, an inflammatory disease or condition can be treated using an antisense nucleic acid of the present invention. Suitable laboratory animals comprising an antisense construct can also provide useful models for deficiencies of leukocyte function, and of activated T 30 lymphocyte deficiency in particular, and can provide further information regarding CXCR3 receptor function. Such animals can provide valuable models of infectious disease or cancer, useful for elucidating the role of leukocytes, such as T lymphocytes and NK cells, in host 35 defenses.

-27-

Method of Producing Recombinant Proteins

Another aspect of the invention relates to a method of producing a mammalian CXCR3 protein or variant (e.g., portion) thereof. Recombinant protein can be obtained, for 5 example, by the expression of a recombinant nucleic acid (e.g., DNA) molecule encoding a mammalian CXCR3 or variant thereof in a suitable host cell, for example.

Constructs (e.g., expression vectors) suitable for the expression of a mammalian CXCR3 protein or variant thereof 10 are also provided. The constructs can be introduced into a suitable host cell, and cells which express a recombinant mammalian CXCR3 protein or variant thereof can be produced and maintained in culture. Such cells are useful for a variety of purposes, including use in the production of 15 protein for characterization, isolation and/or purification, (e.g., affinity purification), use as immunogen, and in binding assays or other functional assays (e.g., to screen for ligands, inhibitors and/or promoters of receptor function), for instance. Suitable host cells 20 can be prokaryotic, including bacterial cells such as *E. coli*, *B. subtilis* and/or other suitable bacteria, or eucaryotic, such as fungal or yeast cells (e.g., *Pichia pastoris*, *Aspergillus* species, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Neurospora crassa*), or other 25 lower eucaryotic cells, and cells of higher eucaryotes such as those from insects (e.g., Sf9 insect cells (WO 94/26087, O'Connor, published November 24, 1994)) or mammals (e.g., Chinese hamster ovary cells (CHO), COS cells, HuT 78 cells, 293 cells). (See, e.g., Ausubel, F.M. et al., eds. *Current 30 Protocols in Molecular Biology*, Greene Publishing Associates and John Wiley & Sons Inc., (1993)).

Host cells which produce a recombinant mammalian CXCR3 protein or variant thereof can be produced as follows. For example, a nucleic acid encoding all or part of the coding 35 sequence for the desired protein can be inserted into a

-28-

nucleic acid vector, e.g., a DNA vector, such as a plasmid, virus or other suitable replicon for expression. A variety of vectors are available, including vectors which are maintained in single copy or multiple copy, or which become 5 integrated into the host cell chromosome.

Transcriptional and/or translational signals of a mammalian CXCR3 gene can be used to direct expression. Suitable expression vectors for the expression of a nucleic acid encoding all or part of the coding sequence of the 10 desired protein are also available. Suitable expression vectors can contain a number of components, including, but not limited to one or more of the following: an origin of replication; a selectable marker gene; one or more expression control elements, such as a transcriptional 15 control element (e.g., a promoter, an enhancer, terminator), and/or one or more translation signals; a signal sequence or leader sequence for membrane targeting in a selected host cell (e.g., of mammalian origin or from a heterologous mammalian or non-mammalian species). In a 20 construct, a signal sequence can be provided by the vector, the mammalian CXCR3 coding sequence, or other source. Sequences present at a site of integration can also provide these elements.

A promoter can be provided for expression in a 25 suitable host cell. Promoters can be constitutive or inducible. For example, a promoter can be operably linked to a nucleic acid encoding the mammalian CXCR3 protein or variant thereof, such that it directs expression of the encoded polypeptide. A variety of suitable promoters for 30 prokaryotic (e.g., lac, tac, T3, T7 promoters for *E. coli*) and eucaryotic (e.g., yeast alcohol dehydrogenase (ADH1), SV40, CMV) hosts are available.

In addition, the expression vectors typically comprise a selectable marker for selection of host cells carrying 35 the vector, and, in the case of replicable expression

-29-

vector, an origin or replication. Genes encoding products which confer antibiotic or drug resistance are common selectable markers and may be used in prokaryotic (e.g.,  $\beta$ -lactamase gene (ampicillin resistance), Tet gene for 5 tetracycline resistance) and eukaryotic cells (e.g., neomycin (G418 or geneticin), gpt (mycophenolic acid), ampicillin, or hygromycin resistance genes). Dihydrofolate reductase marker genes permit selection with methotrexate in a variety of hosts. Genes encoding the gene product of 10 auxotrophic markers of the host (e.g., LEU2, URA3, HIS3) are often used as selectable markers in yeast. Use of viral (e.g., baculovirus) or phage vectors, and vectors which are capable of integrating into the genome of the host cell, such as retroviral vectors, are also 15 contemplated. The present invention also relates to cells carrying these expression vectors.

For example, a nucleic acid encoding a mammalian CXCR3 protein or variant thereof, or a construct comprising such nucleic acid, can be introduced into a suitable host cell 20 by a method appropriate to the host cell selected (e.g., transformation, transfection, electroporation, infection), such that the nucleic acid is operably linked to one or more expression control elements (e.g., in a vector, in a construct created by processes in the cell, integrated into 25 the host cell genome). Host cells can be maintained under conditions suitable for expression (e.g., in the presence of inducer, suitable media supplemented with appropriate salts, growth factors, antibiotic, nutritional supplements, etc.), whereby the encoded polypeptide is produced. If 30 desired, the encoded protein (e.g., human CXCR3) can be isolated (e.g., from the host cells, medium, milk). It will be appreciated that the method encompasses expression in a host cell of a transgenic animal (see e.g., WO 92/03918, GenPharm International, published March 19, 35 1992).

-30-

Fusion proteins can also be produced in this manner. For example, some embodiments can be produced by the insertion of a mammalian CXCR3 protein cDNA or portion thereof into a suitable expression vector, such as

- 5 Bluescript®II SK +/- (Stratagene), pGEX-4T-2 (Pharmacia), pcDNA-3 (Invitrogen) or pET-15b (Novagen). The resulting construct can be introduced into a suitable host cell for expression. Upon expression, fusion protein can be isolated or purified from a cell lysate by means of a
- 10 suitable affinity matrix (see e.g., *Current Protocols in Molecular Biology* (Ausubel, F.M. et al., eds., Vol. 2, Suppl. 26, pp. 16.4.1-16.7.8 (1991)). In addition, affinity labels provide a means of detecting a fusion protein. For example, the cell surface expression or
- 15 presence in a particular cell fraction of a fusion protein comprising an antigen or epitope affinity label can be detected by means of an appropriate antibody.

#### Antibodies

The invention further relates to antibodies reactive with a mammalian CXCR3 protein or portion thereof. In a preferred embodiment, the antibodies specifically bind mammalian CXCR3 receptor(s) or a portion thereof. In one embodiment, antibodies are raised against an isolated and/or recombinant mammalian CXCR3 protein or portion thereof (e.g., a peptide) or against a host cell which expresses recombinant mammalian CXCR3.

Antibodies which can inhibit one or more functions characteristic of a mammalian (e.g., a primate such as a human) CXCR3 protein, such as a binding activity, a signalling activity, and/or stimulation of a cellular response, are also encompassed by the present invention. In one embodiment, antibodies of the present invention can inhibit binding of a ligand (i.e., one or more ligands) to a mammalian CXCR3 protein and/or can inhibit one or more

-31-

functions mediated by a mammalian CXCR3 protein in response to ligand binding. In a particularly preferred embodiment, the antibodies can inhibit (reduce or prevent) the interaction of receptor with a natural ligand, such as

5 IP-10 and/or Mig. For example, as shown herein, antibodies of the present invention can selectively inhibit the interaction of a human CXCR3 protein with IP-10 and/or selectively inhibit receptor functions in response thereto (e.g., signalling activity and/or a cellular response).

10 An antibody designated 1C6, which displays this selectivity, can inhibit binding of IP-10 to a human CXCR3 protein as well as calcium flux and chemotaxis induced by IP-10, but does not significantly inhibit calcium flux induced by Mig under the same conditions. As shown herein,

15 additional antibodies can inhibit IP-10 binding to CXCR3 (e.g., 3A8, 2F8, 3A12, 3E2, 4B4, 4D2, 5B12, 7B8 and 8D5) or IP-10-induced chemotaxis (e.g., 1A5, 3A8, 5F10, and 10C6), although these antibodies did not inhibit Mig-induced signalling under the conditions used, indicating that they

20 are also selective inhibitors of the interaction of a human CXCR3 protein with IP-10 and/or of receptor functions in response thereto.

In a particularly preferred embodiment, the antibodies of the present invention have specificity for a human CXCR3 protein, and even more preferably have an epitopic specificity which is the same as or similar to that of a murine monoclonal antibody (mAb) designated 1C6. Antibodies having an epitopic specificity which is the same as or similar to that of mAb 1C6 can be identified using

25 one or more suitable techniques for characterizing epitopic specificity. For example, antibodies having an epitopic specificity which is the same as or similar to that of mAb 1C6 can be identified by their ability to compete with murine mAb 1C6 for binding to a human CXCR3 protein or

30 portion thereof (e.g., to cells bearing human CXCR3,

- 32 -

including lymphocytes such as activated T cells, NK cells, or recombinant host cells comprising a nucleic acid of the present invention). In one embodiment, antibodies having an epitopic specificity which is the same as or similar to 5 that of mAb 1C6 are further characterized by the ability of a polypeptide having a sequence which is the same as that of residues 1-15 ("P1") of SEQ ID NO:2 to inhibit binding of the antibodies to a human CXCR3 protein in a suitable assay (see e.g., Example 8). In one aspect of this 10 embodiment, binding to a human CXCR3 protein by such antibodies is not significantly inhibited by a polypeptide having a sequence which is the same as that of residues 16-30 ("P2") or 31-45 ("P3") of SEQ ID NO:2.

Other antibodies encompassed by the present invention 15 include antibodies which can bind to a human CXCR3 protein, wherein said binding can be inhibited by a portion of SEQ ID NO:2 corresponding to the N-terminal extracellular segment or a portion thereof. Suitable portions of the N-terminal extracellular segment include N-terminal, 20 internal or C-terminal portions of that segment, such as a polypeptide comprising the N-terminus and having a sequence which is the same as that of residues 1-30, 1-45 or 1-58 of SEQ ID NO:2, for example, or a polypeptide having a sequence which is the same as that of residues 16-30 or 25 45-58 of SEQ ID NO:2. In a preferred embodiment, such portions have "at least one immunological property" of a mammalian CXCR3 protein as defined hereinabove, wherein the mammal is a human. For example, antibodies reactive with a 30 human CXCR3 protein for which binding can be inhibited by a polypeptide having a sequence which is the same as that of residues 16-30 of SEQ ID NO:2 have been obtained (Example 9).

The antibodies of the present invention can be 35 polyclonal or monoclonal, and the term antibody is intended to encompass both polyclonal and monoclonal antibodies.

-33-

The terms polyclonal and monoclonal refer to the degree of homogeneity of an antibody preparation, and are not intended to be limited to particular methods of production.

Antibodies of the present invention can be raised

5 against an appropriate immunogen, including proteins or polypeptides of the present invention, such as isolated and/or recombinant mammalian CXCR3 protein or portion thereof (including synthetic molecules, such as synthetic peptides). In addition, cells expressing recombinant

10 mammalian CXCR3, such as transfected cells, can be used as immunogens or in a screen for antibody which binds receptor. See for example, Chuntharapai et al., *J. Immunol.*, 152: 1783-1789 (1994); and Chuntharapai et al., U.S. Patent No. 5,440,021.

15 Preparation of immunizing antigen, and polyclonal and monoclonal antibody production can be performed using any suitable technique. A variety of methods have been described (see e.g., Kohler et al., *Nature*, 256: 495-497 (1975) and *Eur. J. Immunol.* 6: 511-519 (1976); Milstein et al., *Nature* 266: 550-552 (1977); Koprowski et al., U.S. Patent No. 4,172,124; Harlow, E. and D. Lane, 1988, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory: Cold Spring Harbor, NY); *Current Protocols In Molecular Biology*, Vol. 2 (Supplement 27, Summer '94),

20 Ausubel, F.M. et al., Eds., (John Wiley & Sons: New York, NY), Chapter 11, (1991)). Generally, a hybridoma can be produced by fusing a suitable immortal cell line (e.g., a myeloma cell line such as SP2/0) with antibody producing cells. The antibody producing cell, preferably those of

25 the spleen or lymph nodes, can be obtained from animals immunized with the antigen of interest. The fused cells (hybridomas) can be isolated using selective culture conditions, and cloned by limiting dilution. Cells which produce antibodies with the desired specificity can be

30 selected by a suitable assay (e.g., ELISA).

35

-34-

Other suitable methods of producing or isolating antibodies of the requisite specificity can be used, including, for example, methods which select recombinant antibody from a library, or which rely upon immunization of 5 transgenic animals (e.g., mice) capable of producing a full repertoire of human antibodies (see e.g., Jakobovits et al., *Proc. Natl. Acad. Sci. USA*, 90: 2551-2555 (1993); Jakobovits et al., *Nature*, 362: 255-258 (1993); Lonberg et al., U.S. Patent No. 5,545,806; Surani et al., U.S. Patent 10 No. 5,545,807).

Single chain antibodies, and chimeric, humanized or primatized (CDR-grafted), or veneered antibodies, as well as chimeric, CDR-grafted or veneered single chain antibodies, comprising portions derived from different 15 species, and the like are also encompassed by the present invention and the term "antibody". The various portions of these antibodies can be joined together chemically by conventional techniques, or a polypeptide(s) can be prepared as a contiguous protein using genetic engineering 20 techniques. The term "humanized antibody or immunoglobulin" as used herein refers to an antibody or immunoglobulin comprising portions of immunoglobulins of different origin, wherein at least one portion is of human origin. Accordingly, the present invention relates to a 25 humanized antibody (including antigen-binding fragments thereof) having binding specificity for a mammalian CXCR3 protein (e.g., a human CXCR3 protein), comprising antigen binding regions of nonhuman origin (e.g., rodent) and at least of portion of an immunoglobulin of human origin 30 (e.g., a human framework region, a human constant region, or portions thereof). For example, humanized antibodies can comprise portions derived from an immunoglobulin of nonhuman origin with the requisite specificity (e.g., a nonhuman mouse variable region) and from immunoglobulin 35 sequences of human origin (e.g., a human constant region or

portion thereof) joined together chemically or prepared as a contiguous polypeptide using genetic engineering techniques. Another example of a humanized antibody of the invention is an immunoglobulin comprising one or more 5 immunoglobulin chains, said immunoglobulin comprising a CDR of nonhuman origin (e.g., one or more CDRs derived from an antibody of nonhuman origin) and a framework region derived from a light and/or heavy chain of human origin (e.g., CDR-grafted antibodies with or without framework changes). In 10 one aspect of this embodiment, the immunoglobulin comprises the light chain CDRs (CDR1, CDR2 and CDR3) and heavy chain CDRs (CDR1, CDR2, CDR3) of a nonhuman immunoglobulin. In one embodiment, the humanized immunoglobulin has an epitopic specificity which is the same or similar to that 15 mAb 1C6. In a preferred embodiment, the antigen binding region of the humanized immunoglobulin is derived from mAb 1C6. The term humanized antibody also encompasses single chain antibodies. See, e.g., Cabilly et al., U.S. Patent No. 4,816,567; Cabilly et al., European Patent No. 20 0,125,023 B1; Boss et al., U.S. Patent No. 4,816,397; Boss et al., European Patent No. 0,120,694 B1; Neuberger, M.S. et al., WO 86/01533; Neuberger, M.S. et al., European Patent No. 0,194,276 B1; Winter, U.S. Patent No. 5,225,539; Winter, European Patent No. 0,239,400 B1; Queen et al., 25 European Patent No. 0 451 216 B1; and Padlan, E.A. et al., EP 0 519 596 A1. See also, Newman, R. et al., *BioTechnology*, 10: 1455-1460 (1992), regarding primatized antibody, and Ladner et al., U.S. Patent No. 4,946,778 and Bird, R.E. et al., *Science*, 242: 423-426 (1988)) regarding 30 single chain antibodies.

In addition, functional fragments of antibodies, including fragments of chimeric, humanized, primatized, veneered or single chain antibodies, can also be produced. Functional fragments of foregoing antibodies retain at 35 least one binding function and/or modulation function of

-36-

the full-length antibody from which they are derived. For example, antibody fragments capable of binding to a mammalian CXCR3 protein or portion thereof, including, but not limited to, Fv, Fab, Fab' and F(ab')<sub>2</sub>, fragments are 5 encompassed by the invention. Such fragments can be produced by enzymatic cleavage or by recombinant techniques. For instance, papain or pepsin cleavage can generate Fab or F(ab')<sub>2</sub>, fragments, respectively. Antibodies can also be produced in a variety of truncated 10 forms using antibody genes in which one or more stop codons has been introduced upstream of the natural stop site. For example, a chimeric gene encoding a F(ab')<sub>2</sub>, heavy chain portion can be designed to include DNA sequences encoding the CH<sub>1</sub> domain and hinge region of the heavy chain. 15 The antibodies of the present invention are useful in a variety of applications, including research, diagnostic and therapeutic applications. For instance, they can be used to isolate, purify and/or detect receptor or portions thereof, and to study receptor structure (e.g., 20 conformation) and function.

The antibodies of the present invention can also be used to modulate receptor function in research and therapeutic applications. For instance, antibodies can act as inhibitors to inhibit (reduce or prevent) (a) binding 25 (e.g., of a ligand, a second inhibitor or a promoter) to the receptor, (b) receptor signalling, and/or (c) a cellular response. Antibodies which act as inhibitors of receptor function can block ligand or promoter binding directly or indirectly (e.g., by causing a conformational 30 change in the receptor). For example, antibodies can inhibit receptor function by inhibiting binding of a ligand, or by desensitization (with or without inhibition of binding of a ligand). Antibodies of the present invention can be used as antagonists of effector cells such 35 as activated or stimulated T lymphocytes and natural killer

(NK) cells, and find use in methods of therapy (including prophylaxis) of diseases or conditions which can be treated with inhibitors of CXCR3 function as described herein.

Antibodies which can selectively inhibit the interaction of

5 a human CXCR3 protein with IP-10 and/or selectively inhibit receptor functions in response thereto (e.g., mAb 1C6, antibodies having an epitopic specificity similar to that of mAb 1C6) are particularly useful in the treatment of diseases or disorders mediated by IP-10-CXCR3 interaction.

10 For example, inflammatory conditions such as psoriasis, inflammatory bowel diseases, nephritis, and multiple sclerosis can be particularly amenable to therapy.

Surprisingly, mAb 1C6 inhibits T cell activation as assessed in a mixed lymphocyte reaction (MLR).

15 Accordingly, mAb 1C6 and other anti-CXCR3 antibodies which can inhibit T cell activation, can be used to inhibit T cell activation and function(s) associated with activation, such as cytokine production, cytotoxic T cell killing, and/or provision of T cell help. When used in methods of

20 therapy or prophylaxis as described herein, such antibodies can have the added advantage of inhibiting further T cell activation. Such antibodies are particularly attractive as therapeutic agents for treating graft rejection (e.g., in transplantation), including allograft rejection or graft-

25 versus-host disease, or other diseases or conditions in which inhibition of the activation of T cells is desired.

Antibodies which bind receptor can also act as agonists of receptor function, triggering or stimulating a receptor function, such as signalling and/or a cellular

30 response (e.g., calcium flux, chemotaxis, exocytosis or pro-inflammatory mediator release) upon binding to receptor. Accordingly, such antibodies can be used as agonists of effector cells such as activated or stimulated T lymphocytes and natural killer (NK) cells, and find use

35 in methods of therapy (including prophylaxis) of diseases

or conditions which can be treated with promoters of CXCR3 function as described herein.

In addition, the various antibodies of the present invention can be used to detect or measure the expression 5 of receptor, for example, on leukocytes such as activated T cells or natural killer cells (NK cells), or on cells transfected with a receptor gene. Thus, they also have utility in applications such as cell sorting (e.g., flow cytometry, fluorescence activated cell sorting), for 10 diagnostic or research purposes.

Anti-idiotypic antibodies are also provided. Anti-idiotypic antibodies recognize antigenic determinants associated with the antigen-binding site of another antibody. Anti-idiotypic antibodies can be prepared a 15 against a first antibody by immunizing an animal of the same species, and preferably of the same strain as the animal used to produce the first antibody, with said first antibody. See e.g., U.S. Patent No. 4,699,880.

In one embodiment, antibodies are raised against 20 receptor or a portion thereof, and these antibodies are used in turn as immunogen to produce an anti-idiotypic antibody. The anti-Id produced thereby can mimic receptor and bind compounds which bind receptor, such as ligands, inhibitors or promoters of receptor function, and can be 25 used in an immunoassay to detect, identify or quantitate such compounds. Such an anti-idiotypic antibody can also be an inhibitor of receptor function, although it does not bind receptor itself.

Anti-idiotypic (i.e., Anti-Id) antibody can itself be 30 used to raise an anti-idiotypic antibody (i.e., Anti-anti-Id). Such an antibody can be similar or identical in specificity to the original immunizing antibody. In one embodiment, antibody antagonists which block binding to receptor can be used to raise Anti-Id, and 35 the Anti-Id can be used to raise Anti-anti-Id, which can

- 39 -

have a specificity which is similar or identical to that of the antibody antagonist. These anti-anti-Id antibodies can be assessed for inhibitory effect on receptor function to determine if they are antagonists.

5 Single chain, and chimeric, humanized, primateized (CDR-grafted), veneered, as well as chimeric, CDR-grafted, or veneered single chain anti-idiotypic antibodies can be prepared, and are encompassed by the term anti-idiotypic antibody. Antibody fragments of such antibodies can also  
10 be prepared.

The antibodies and fragments of the present invention can be modified, for example, by incorporation of or attachment (directly or indirectly) of a detectable label such as a radioisotope, spin label, antigen or enzyme  
15 label, fluorescent or chemiluminescent group and the like, and such modified forms are included within the scope of the invention.

Identification of Ligands, Inhibitors or Promoters of Receptor Function

20 As used herein, a ligand is a substance which binds to a receptor protein. A ligand of a selected mammalian CXCR3 protein is a substance which binds to the selected mammalian CXCR3 protein. In a preferred embodiment, ligand binding of a mammalian CXCR3 protein occurs with high  
25 affinity. The term ligand refers to substances including, but not limited to, a natural ligand, whether isolated and/or purified, synthetic, and/or recombinant, a homolog of a natural ligand (e.g., from another mammal), antibodies, portions of such molecules, and other  
30 substances which bind receptor. A natural ligand of a selected mammalian receptor can bind to the receptor under physiological conditions, and is of a mammalian origin which is the same as that of the mammalian CXCR3 protein. The term ligand encompasses substances which are inhibitors

-40-

or promoters of receptor activity, as well as substances which selectively bind receptor, but lack inhibitor or promoter activity.

As used herein, an inhibitor is a substance which  
5 inhibits at least one function characteristic of a  
mammalian CXCR3 protein (e.g., a human CXCR3), such as a  
binding activity (e.g., ligand binding, promoter binding),  
a signalling activity (e.g., activation of a mammalian G  
protein, induction of rapid and transient increase in the  
10 concentration of cytosolic free calcium  $[Ca^{2+}]_i$ ), and/or  
cellular response function (e.g., stimulation of  
chemotaxis, exocytosis or inflammatory mediator release by  
leukocytes). The term inhibitor refers to substances  
including antagonists which bind receptor (e.g., an  
15 antibody, a mutant of a natural ligand, other competitive  
inhibitors of ligand binding), and substances which inhibit  
receptor function without binding thereto (e.g., an  
anti-idiotypic antibody).

As used herein, a promoter is a substance which  
20 promotes (induces or enhances) at least one function  
characteristic of a mammalian CXCR3 protein (e.g., a human  
CXCR3), such as a binding activity (e.g., ligand, inhibitor  
and/or promoter binding), a signalling activity (e.g.,  
activation of a mammalian G protein, induction of rapid and  
25 transient increase in the concentration of cytosolic free  
calcium  $[Ca^{2+}]_i$ ), and/or a cellular response function (e.g.,  
stimulation of chemotaxis, exocytosis or inflammatory  
mediator release by leukocytes). The term promoter refers  
to substances including agonists which bind receptor (e.g.,  
30 an antibody, a homolog of a natural ligand from another  
species), and substances which promote receptor function  
without binding thereto (e.g., by activating an associated  
protein). In a preferred embodiment, the agonist is other  
than a homolog of a natural ligand.

-41-

The assays described below, which rely upon the nucleic acids and proteins of the present invention, can be used, alone or in combination with each other or other suitable methods, to identify ligands, inhibitors or 5 promoters of a mammalian CXCR3 protein or variant. The in vitro methods of the present invention can be adapted for high-throughput screening in which large numbers of samples are processed (e.g., a 96 well format). Host cells comprising a nucleic acid of the present invention and 10 expressing recombinant mammalian CXCR3 (e.g., human CXCR3) at levels suitable for high-throughput screening can be used, and thus, are particularly valuable in the identification and/or isolation of ligands, inhibitors and 15 promoters of mammalian CXCR3 proteins. Expression of receptor can be monitored in a variety of ways. For instance, expression can be monitored using antibodies of the present invention which bind receptor or a portion thereof. Also, commercially available antibodies can be used to detect expression of an antigen- or epitope-tagged 20 fusion protein comprising a receptor protein or polypeptide (e.g., FLAG tagged receptors), and cells expressing the desired level can be selected.

Nucleic acid encoding a mammalian CXCR3 protein, the 25 can be incorporated into an expression system to produce a receptor protein or polypeptide as described above. An isolated and/or recombinant receptor protein or polypeptide, such as a receptor expressed in cells stably or transiently transfected with a construct comprising a nucleic acid of the present invention, or in a cell 30 fraction containing receptor (e.g., a membrane fraction from transfected cells, liposomes incorporating receptor), can be used in tests for receptor function. The receptor can be further purified if desired. Testing of receptor function can be carried out *in vitro* or *in vivo*.

-42-

An isolated and/or recombinant mammalian CXCR3 protein, such as a human CXCR3 as shown in Figure 2 (SEQ ID NO:2), can be used in the present method, in which the effect of a compound is assessed by monitoring receptor 5 function as described herein or using other suitable techniques. For example, stable or transient transfectants such as those described in Example 2 or other suitable cells (e.g., baculovirus infected Sf9 cells, stable transfectants of mouse L1-2 pre-B cells (derived from a pre- 10 B lymphoma, Dr. Eugene Butcher (Stanford University, Stanford, CA)), can be used in binding assays. Stable transfectants of Jurkat cells (Example 2) or of other suitable cells capable of chemotaxis can be used (e.g., mouse L1-2 pre-B cells) in chemotaxis assays, for example. 15 According to the method of the present invention, compounds can be individually screened or one or more compounds can be tested simultaneously according to the methods herein. Where a mixture of compounds is tested, the compounds selected by the processes described can be 20 separated (as appropriate) and identified by suitable methods (e.g., PCR, sequencing, chromatography). The presence of one or more compounds (e.g., a ligand, inhibitor, promoter) in a test sample can also be determined according to these methods. 25 Large combinatorial libraries of compounds (e.g., organic compounds, recombinant or synthetic peptides, "peptoids", nucleic acids) produced by combinatorial chemical synthesis or other methods can be tested (see e.g., Zuckerman, R.N. et al., *J. Med. Chem.*, 37: 2678-2685 30 (1994) and references cited therein; see also, Ohlmeyer, M.H.J. et al., *Proc. Natl. Acad. Sci. USA* 90:10922-10926 (1993) and DeWitt, S.H. et al., *Proc. Natl. Acad. Sci. USA* 90:6909-6913 (1993), relating to tagged compounds; Rutter, W.J. et al. U.S. Patent No. 5,010,175; Huebner, V.D. et 35 al., U.S. Patent No. 5,182,366; and Geysen, H.M., U.S.

-43-

Patent No. 4,833,092). Where compounds selected from a combinatorial library by the present method carry unique tags, identification of individual compounds by chromatographic methods is possible.

- 5 In one embodiment, phage display methodology is used. For example, receptor can be contacted with a phage (e.g., a phage or collection of phage such as a library) displaying a polypeptide under conditions appropriate for receptor binding (e.g., in a suitable binding buffer).
- 10 Phage bound to receptor can be selected using standard techniques or other suitable methods. Phage can be separated from receptor using a suitable elution buffer. For example, a change in the ionic strength or pH can lead to a release of phage. Alternatively, the elution buffer
- 15 can comprise a release component or components designed to disrupt binding of compounds (e.g., one or more compounds which can disrupt binding of the displayed peptide to the receptor, such as a ligand, inhibitor, and/or promoter which competitively inhibits binding). Optionally, the
- 20 selection process can be repeated or another selection step can be used to further enrich for phage which bind receptor. The displayed polypeptide can be characterized (e.g., by sequencing phage DNA). The polypeptides identified can be produced and further tested for ligand
- 25 binding, inhibitor and/or promoter function. Analogs of such peptides can be produced which will have increased stability or other desirable properties.

In one embodiment, phage expressing and displaying fusion proteins comprising a coat protein with an

- 30 N-terminal peptide encoded by random sequence nucleic acids can be produced. Suitable host cells expressing a receptor protein or polypeptide of the present invention are contacted with the phage, bound phage are selected, recovered and characterized. (See e.g., Doorbar, J. and G.

-44-

Winter, *J. Mol. Biol.*, 244: 361 (1994) discussing a phage display procedure used with a G protein-coupled receptor).

Other sources of potential ligands, inhibitors and/or promoters of mammalian CXCR3 proteins include, but are not limited to, variants of CXCR3 ligands, including naturally occurring, synthetic or recombinant variants of IP-10 or Mig, substances such as other chemoattractants or chemokines, variants thereof, other inhibitors and/or promoters (e.g., anti-CXCR3 antibodies, antagonists, agonists), other G-protein coupled receptor ligands, inhibitors and/or promoters (e.g., antagonists or agonists), and soluble portions of a mammalian CXCR3 receptor, such as a suitable receptor peptide or analog which can inhibit receptor function (see e.g., Murphy, R.B., WO 94/05695).

#### Binding Assays

The isolated and/or recombinant receptor proteins or functional variants thereof, including portions thereof or suitable fusion proteins, can be used in a method to select and identify agents which bind to a (one or more) mammalian CXCR3 protein, such as human CXCR3, and which are ligands, or potential inhibitors or promoters of receptor activity. Agents selected by the method, including ligands, inhibitors or promoters, can be further assessed for an inhibitory or stimulatory effect on receptor function and/or for therapeutic utility.

In one embodiment, an agent which binds to an active, isolated and/or recombinant mammalian CXCR3 protein or polypeptide is identified by the method. In this embodiment, the receptor protein or polypeptide used has at least one property, activity or function characteristic of a mammalian CXCR3 protein (as defined herein), such as a binding activity (e.g., ligand, inhibitor and/or promoter binding), a signalling activity (e.g., activation of a

-45-

mammalian G protein, induction of rapid and transient increase in the concentration of cytosolic free calcium  $[Ca^{2+}]_i$ , cellular response function (e.g., stimulation of chemotaxis, exocytosis or inflammatory mediator release by 5 leukocytes), and/or an immunological property as defined herein. In a preferred embodiment, the isolated and/or recombinant mammalian CXCR3 protein or variant has ligand binding function, and more preferably binds a natural ligand of the receptor. In a particularly preferred 10 embodiment, the isolated and/or recombinant protein is a human CXCR3 protein encoded by the nucleic acid illustrated Figure 1 (SEQ ID NO:1).

For example, a composition comprising an isolated and/or recombinant mammalian CXCR3 protein or variant 15 thereof can be maintained under conditions suitable for binding, the receptor can be contacted with an agent (e.g., a composition comprising one or more agent) to be tested, and binding is detected or measured. In one embodiment, a receptor protein can be expressed in cells stably or 20 transiently transfected with a construct comprising a nucleic acid sequence which encodes a receptor of the present invention. The cells can be maintained under conditions appropriate for expression of receptor. The cells are contacted with an agent under conditions suitable 25 for binding (e.g., in a suitable binding buffer), and binding can be detected by standard techniques. For example, the extent of binding can be determined relative to a suitable control (e.g., compared with background determined in the absence of agent, compared with binding 30 of a second agent (i.e., a standard), compared with binding of the agent to untransfected cells). Optionally, a cellular fraction, such as a membrane fraction, containing receptor can be used in lieu of whole cells.

Binding or complex formation can be detected directly 35 or indirectly. In one embodiment, the agent can be labeled

-46-

with a suitable label (e.g., fluorescent label, chemiluminescent label, isotope label, enzyme label), and binding can be determined by detection of the label. Specificity of binding can be assessed by competition or 5 displacement, for example, using unlabeled agent or a ligand (e.g., IP-10, Mig) as competitor.

Ligands of the mammalian receptor, including natural ligands from the same mammalian species or from another species, can be identified in this manner. The binding 10 activity of a promoter or inhibitor which binds receptor can also be assessed using such a ligand binding assay.

Binding inhibition assays can also be used to identify ligands, and inhibitors and promoters which bind receptor and inhibit binding of another agent such as a ligand. For 15 example, a binding assay can be conducted in which a reduction in the binding of a first agent (in the absence of a second agent), as compared with binding of the first agent in the presence of the second test agent, is detected or measured. The receptor can be contacted with the first 20 and second agents simultaneously, or one after the other, in either order. A reduction in the extent of binding of the first agent in the presence of the second test agent, is indicative of inhibition of binding by the second agent. For example, binding of the first agent could be decreased 25 or abolished.

In one embodiment, direct inhibition of the binding of a first agent (e.g., a chemokine such as IP-10, Mig) to a human CXCR3 by a second test agent is monitored. For example, the ability of an agent to inhibit the binding of 30  $^{125}\text{I}$ -labeled Mig to human CXCR3 can be monitored. Such an assay can be conducted using whole cells (e.g., a suitable cell line containing nucleic acid encoding a human CXCR3 receptor), or a membrane fraction from said cells, for instance.

- 47 -

Other methods of identifying the presence of an agent(s) which binds a receptor are available, such as methods which monitor events which are triggered by receptor binding, including signalling function and/or 5 stimulation of a cellular response.

It will be understood that the inhibitory effect of antibodies of the present invention can be assessed in a binding inhibition assay. Competition between antibodies for receptor binding can also be assessed in the method in 10 which the first agent in the assay is another antibody, under conditions suitable for antibody binding.

Ligands, receptor-binding inhibitors and promoters, which are identified in this manner, can be further assessed to determine whether, subsequent to binding, they 15 act to inhibit or activate other functions of CXCR3 receptors and/or to assess their therapeutic utility.

#### Signalling Assays

The binding of a G protein-coupled receptor (e.g., by an agonist) can result in signalling by the receptor, and 20 stimulation of the activity of G protein. The induction of signalling function by an agent can be monitored using any suitable method. For example, G protein activity, such as hydrolysis of GTP to GDP, or later signalling events triggered by receptor binding, such as induction of rapid 25 and transient increase in the concentration of intracellular (cytosolic) free calcium  $[Ca^{2+}]_i$ , can be assayed by methods known in the art or other suitable methods (Example 2; see also, Neote, K. et al., *Cell*, 72: 415-425 1993); Van Riper et al., *J. Exp. Med.*, 177: 851-856 30 (1993); Dahinden, C.A. et al., *J. Exp. Med.*, 179: 751-756 (1994)).

The functional assay of Sledziewski et al. using hybrid G protein coupled receptors can also be used to identify a ligand or promoter by its ability to activate a

-48-

hybrid G protein or to identify an inhibitor by its ability to inhibit such activation (Sledziewski et al., U.S. Patent No. 5,284,746, the teachings of which are incorporated herein by reference). In one embodiment, a biological response of the host cell (triggered by binding to hybrid receptor) can be monitored, detection of the response being indicative of the presence of ligand in the test sample. For example, a method of detecting the presence of a ligand in a test sample is described, wherein the ligand is an agent which is capable of being bound by the ligand-binding domain of a receptor. In one embodiment of the method, yeast host cells are transformed with a DNA construct capable of directing the expression of a biologically active hybrid G protein-coupled receptor (i.e., a fusion protein). The hybrid receptor comprises a mammalian G protein-coupled receptor having at least one domain other than the ligand-binding domain replaced with a corresponding domain of a yeast G protein-coupled receptor, such as a *STE2* gene product. The yeast host cells containing the construct are maintained under conditions in which the hybrid receptor is expressed, and the cells are contacted with a test sample under conditions suitable to permit binding of ligand to the hybrid receptor. A biological response of the host cell (triggered by binding to hybrid receptor) is monitored, detection of the response being indicative of a signalling function. For instance, binding to a hybrid receptor derived from *STE2* gene product can lead to induction of the *BAR1* promoter. Induction of the promoter can be measured by means of a reporter gene (e.g.,  $\beta$ -gal), which is linked to the *BAR1* promoter and introduced into host cells on a second construct. Expression of the reporter gene can be detected by an *in vitro* enzyme assay on cell lysates or by the presence of blue colonies on plates containing an indicator (e.g., X-gal) in the medium, for example.

- 49 -

In another embodiment, the assay can be used to identify potential inhibitors of receptor function. The inhibitory activity of an agent can be determined using a ligand or promoter in the assay, and assessing the ability 5 of the test agent to inhibit the activity induced by ligand or promoter.

Variants of known ligands can also be screened for reduced ability (decreased ability or no ability) to stimulate activity of a coupled G protein. In this 10 embodiment, although the agent has ligand binding activity (as determined by another method), engagement of the receptor does not trigger or only weakly triggers activity of a coupled G protein. Such agents are potential antagonists, and can be further assessed for inhibitory 15 activity.

#### Chemotaxis and Other Assays of Cellular Responses

Chemotaxis assays can also be used to assess receptor function. These assays are based on the functional migration of cells *in vitro* or *in vivo* induced by an agent, 20 and can be used to assess the binding and/or effect on chemotaxis of ligands, inhibitors, or promoters.

The use of an *in vitro* chemotaxis assay to assess the response of cells to IP-10 and Mig is described in Example 2. Springer et al. describe a transendothelial 25 lymphocyte chemotaxis assay (Springer et al., WO 94/20142, published September 15, 1994, the teachings of which are incorporated herein by reference; see also Berman et al., *Immunol Invest.*, 17: 625-677 (1988)). Migration across endothelium into collagen gels has also been described 30 (Kavanaugh et al., *J. Immunol.*, 146: 4149-4156 (1991)).

Generally, chemotaxis assays monitor the directional movement or migration of a suitable cell capable of chemotaxis, such as a leukocyte (e.g., T lymphocytes, NK cells, monocytes), stable transfectants of Jurkat cells,

-50-

mouse L1-2 pre-B cells or of other suitable host cells, for example, into or through a barrier (e.g., endothelium, a filter), toward increased levels of an agent, from a first surface of the barrier toward an opposite second surface.

5   Membranes or filters provide convenient barriers, such that the directional movement or migration of a suitable cell into or through a filter, toward increased levels of an agent, from a first surface of the filter toward an opposite second surface of the filter, is monitored. In  
10   some assays, the membrane is coated with a substance to facilitate adhesion, such as ICAM-1, fibronectin or collagen.

For example, one can detect or measure the migration of cells in a suitable container (a containing means), from  
15   a first chamber into or through a microporous membrane into a second chamber which contains an agent to be tested, and which is divided from the first chamber by the membrane. A suitable membrane, having a suitable pore size for monitoring specific migration in response to the agent,  
20   including, for example, nitrocellulose, polycarbonate, is selected. For example, pore sizes of about 3-8 microns, and preferably about 5-8 microns can be used. Pore size can be uniform on a filter or within a range of suitable pore sizes.

25   To assess migration, the distance of migration into the filter, the number of cells crossing the filter that remain adherent to the second surface of the filter, and/or the number of cells that accumulate in the second chamber can be determined using standard techniques (e.g., by  
30   microscopy). In one embodiment, the cells are labeled with a detectable label (e.g., radioisotope, fluorescent label, antigen or epitope label), and migration can be assessed by determining the presence of the label adherent to the membrane and/or present in the second chamber using an  
35   appropriate method (e.g., by detecting radioactivity,

-51-

fluorescence, immunoassay). The extent of migration induced by an agent can be determined relative to a suitable control (e.g., compared to background migration determined in the absence of the agent, to the extent of 5 migration induced by a second agent (i.e., a standard), compared with migration of untransfected cells induced by the agent).

Chambers can be formed from various solids, such as plastic, glass, polypropylene, polystyrene, etc. Membranes 10 which are detachable from the chambers, such as a Biocoat (Collaborative Biomedical Products) or Transwell (Costar, Cambridge, MA) culture insert, facilitate counting adherent cells. In the container, the filter can be situated so as to be in contact with fluid containing cells in the first 15 chamber, and the fluid in the second chamber. Other than the test agent or additional ligand, inhibitor, or promoter present for the purpose of the assay, the fluid on either side of the membrane is preferably the same or substantially similar. The fluid in the chambers can 20 comprise protein solutions (e.g., bovine serum albumin, fetal calf serum, human serum albumin) which may act to increase stability and inhibit nonspecific binding of cells, and/or culture media.

In one embodiment, transendothelial migration is 25 assessed. In addition to lower background (signal to noise ratio), transendothelial migration models *in vivo* conditions in which leukocytes emigrate from blood vessels toward chemoattractants present in the tissues at sites of inflammation by crossing the endothelial cell layer lining 30 the vessel wall. In this embodiment, transmigration through an endothelial cell layer assessed. To prepare the cell layer, endothelial cells can be cultured on a microporous filter or membrane, optionally coated with a substance such as collagen, fibronectin, or other 35 extracellular matrix proteins, to facilitate the attachment

-52-

of endothelial cells. Preferably, endothelial cells are cultured until a confluent monolayer is formed. A variety of mammalian endothelial cells can be available for monolayer formation, including for example, vein, artery or 5 microvascular endothelium, such as human umbilical vein endothelial cells (Clonetics Corp, San Diego, CA) or a suitable cell line, such as the ECV 304 cell line used (European Collection of Animal Cell Cultures, Porton Down, Salisbury, U.K.). To assay chemotaxis in response to a 10 particular mammalian receptor, endothelial cells of the same mammal are preferred; however endothelial cells from a heterologous mammalian species or genus can also be used.

Generally, the assay is performed by detecting the directional migration of cells into or through a membrane 15 or filter, in a direction toward increased levels of an agent, from a first surface of the filter toward an opposite second surface of the filter, wherein the filter contains an endothelial cell layer on a first surface.

Directional migration occurs from the area adjacent to the 20 first surface, into or through the membrane, towards an agent situated on the opposite side of the filter. The concentration of agent present in the area adjacent to the second surface, is greater than that in the area adjacent to the first surface.

25 In one embodiment, a chemotaxis assay is used to test for ligand or promoter activity of an agent, a composition comprising cells capable of migration and expressing a mammalian CXCR3 protein or functional variant thereof are placed in the first chamber, and a composition comprising 30 the agent (one or more agents) to be tested is placed in the second chamber, preferably in the absence of other ligands or promoters capable of inducing chemotaxis of the cells in the first chamber (having chemoattractant function). However, one or more ligands or promoters 35 having chemoattractant function may be present. The

-53-

ability of an agent to induce chemotaxis of the cells expressing a mammalian CXCR3 receptor in this assay is indicative that the agent is a ligand or promoter of receptor function.

5        In one embodiment used to test for an inhibitor, a composition comprising cells capable of migration and expressing a mammalian CXCR3 protein or functional variant are placed in the first chamber. A composition comprising a ligand or promoter (i.e., one or more ligands or

10      promoters) capable of inducing chemotaxis of the cells in the first chamber (having chemoattractant function) is placed in the second chamber. Before (preferably shortly before) the cells are placed in the first chamber, or simultaneously with the cells, a composition comprising the

15      agent to be tested is placed, preferably, in the first chamber. The ability of an agent to inhibit ligand- or promoter-induced chemotaxis of the cells expressing a mammalian CXCR3 protein in this assay is indicative that the agent is an inhibitor of receptor function (e.g., an

20      inhibitor of cellular response function). A reduction in the extent of migration induced by the ligand or promoter in the presence of the test agent, is indicative of inhibitory activity. Separate binding studies (see above) can be performed to determine whether inhibition is a

25      result of binding of the test agent to receptor or occurs via a different mechanism.

In vivo assays which monitor leukocyte infiltration of a tissue, in response to injection of an agent in the tissue, are described below. These models measure the

30      ability of cells to respond to a ligand or promoter by emigration and chemotaxis to a site of inflammation.

The effects of a ligand, inhibitor or promoter on the cellular response function of a CXCR3 receptor can be assessed by monitoring other cellular responses induced by

35      active receptor, using suitable host cells containing

receptor. Similarly, these assays can be used to determine the function of a receptor. For instance, exocytosis (e.g., degranulation of natural killer cells leading to release of one or more enzymes or other granule components, 5 such as esterases (e.g., serine esterases), perforin, and/or granzymes), inflammatory mediator release (such as release of bioactive lipids such as leukotrienes (e.g., leukotriene C<sub>4</sub>)), and respiratory burst, can be monitored by methods known in the art or other suitable methods.

10 (See e.g., Taub, D.D. et al., *J. Immunol.*, 155: 3877-3888 (1995), regarding assays for release of granule-derived serine esterases (the teachings of which are incorporated herein by reference) and Loetscher et al., *J. Immunol.*, 156: 322-327 (1996), regarding assays for enzyme and

15 granzyme release by NK cells and cytotoxic T lymphocytes (CTLs) (the teachings of which are incorporated herein by reference); Rot, A. et al., *J. Exp. Med.*, 176: 1489-1495 (1992) regarding respiratory burst; Bischoff, S.C. et al., *Eur. J. Immunol.*, 23: 761-767 (1993) and Baggliolini, M.

20 and C.A. Dahinden, *Immunology Today*, 15: 127-133 (1994)).

In one embodiment, a ligand, inhibitor and/or promoter is identified by monitoring the release of an enzyme upon degranulation or exocytosis by a cell capable of this function. Cells containing a nucleic acid of the present 25 invention, which encodes an active receptor protein capable of stimulating exocytosis or degranulation are maintained in a suitable medium under suitable conditions, whereby receptor is expressed and degranulation can be induced. The receptor is contacted with an agent to be tested, and

30 enzyme release is assessed. The release of an enzyme into the medium can be detected or measured using a suitable assay, such as in an immunological assay, or biochemical assay for enzyme activity.

The medium can be assayed directly, by introducing 35 components of the assay (e.g., substrate, co-factors,

-55-

antibody) into the medium (e.g., before, simultaneous with or after the cells and agent are combined). Alternatively, the assay can be performed on medium which has been separated from the cells or further processed (e.g., 5 fractionated) prior to assay. For example, convenient assays for are available for enzymes such serine esterases (see e.g., Taub, D.D. et al., *J. Immunol.*, 155: 3877-3888 (1995) regarding release of granule-derived serine esterases).

10       Stimulation of degranulation by an agent can be indicative that the agent is a ligand or promoter of a mammalian CXCR3 protein. In another embodiment, cells expressing receptor are combined with a ligand or promoter, and an agent to be tested is added before, after or 15 simultaneous therewith, and degranulation is assessed. Inhibition of ligand- or promoter-induced degranulation is indicative that the agent is an inhibitor of mammalian CXCR3 protein function.

Cellular adherence can also monitored by methods known 20 in the art or other suitable methods. Engagement of the chemokine receptors of a lymphocyte can cause integrin activation, and induction of adherence to adhesion molecules expressed in vasculature or the perivascular space. In one embodiment, a ligand, inhibitor and/or 25 promoter is identified by monitoring cellular adherence by a cell capable of adhesion. For example, an agent to be tested can be combined with (a) cells expressing receptor (preferably non-adherent cells which when transfected with receptor acquire adhesive ability), (b) a composition 30 comprising a suitable adhesion molecule (e.g., a substrate such as a culture well coated with an adhesion molecule, such as fibronectin), and (c) a ligand or promoter (e.g., agonist), and maintained under conditions suitable for ligand- or promoter-induced adhesion. Labeling of cells 35 with a fluorescent dye provides a convenient means of

-56-

detecting adherent cells. Nonadherent cells can be removed (e.g., by washing) and the number of adherent cells determined. The effect of the agent in inhibiting or enhancing ligand- or promoter-induced adhesion can be 5 indicative of inhibitor or promoter activity, respectively. Agents active in the assay include inhibitors and promoters of binding, signalling, and/or cellular responses. In another embodiment, an agent to be tested can be combined with cells expressing receptor and a composition comprising 10 a suitable adhesion molecule under conditions suitable for ligand- or promoter-induced adhesion, and adhesion is monitored. Increased adhesion relative to a suitable control is indicative of the presence of a ligand and/or promoter.

15 Models of Inflammation

A variety of *in vivo* models of inflammation are available, which can be used to assess the effects of ligands, inhibitors, or promoters *in vivo* as therapeutic agents, including a sheep model for asthma (see e.g., Weg, 20 V.B. et al., *J. Exp. Med.*, 177: 561 (1993), the teachings of which are incorporated herein by reference), a rat delayed type hypersensitivity model (Rand, M.L. et al., *Am. J. Pathol.*, 148: 855-864 (1996), the teachings of which are incorporated herein by reference), or other suitable 25 models. The activity of antibodies which cross-react with other mammalian CXCR3 proteins can be assessed in such mammals.

In addition, leukocyte infiltration upon intradermal injection of a compound into a suitable animal, such as 30 rabbit, rat, or guinea pig, can be monitored (see e.g., Van Damme J. et al., *J. Exp. Med.*, 176: 59-65 (1992); Zachariae, C.O.C. et al., *J. Exp. Med.*, 171: 2177-2182 (1990); Jose, P.J. et al., *J. Exp. Med.*, 179: 881-887 (1994)). In one embodiment, skin biopsies are assessed

-57-

histologically for infiltration of leukocytes (e.g., T lymphocytes, monocytes, natural killer cells). In another embodiment, labeled cells (e.g., cells expressing a mammalian CXCR3 protein which are labeled with  $^{111}\text{In}$ , for example) capable of chemotaxis and extravasation are administered to the animal. Infiltration of labelled cells in the vicinity of the site of injection of a test sample (e.g., a compound to be tested in a suitable buffer or physiological carrier) is indicative of the presence of a ligand or promoter, such as an agonist, in the sample. These assays can also be modified to identify inhibitors of chemotaxis and leukocyte extravasation. For example, an inhibitor can be administered, either before, simultaneously with or after ligand or agonist is administered to the test animal. A decrease of the extent of infiltration in the presence of inhibitor as compared with the extent of infiltration in the absence of inhibitor is indicative of inhibition.

#### Diagnostic Applications

The present invention has a variety of diagnostic applications. For example, a mutation(s) in a gene encoding a mammalian CXCR3 protein can cause a defect in at least one function of the encoded receptor, thereby reducing or enhancing receptor function. For instance, a mutation which produces a variant of receptor or alters the level of expression, can reduce or enhance receptor function, reducing or enhancing processes mediated by receptor (e.g., inflammatory processes). The presence of such a mutation can be determined using methods which detect or measure the presence of receptor or receptor function in cells (e.g., leukocytes, such as activated T lymphocytes) of an individual or in a receptor preparation isolated from such cells. In these assays, reduced or

- 58 -

enhanced levels of receptor and/or reduced or enhanced receptor function can be assessed.

The nucleic acids of the present invention provide reagents, such as probes and PCR primers, which can be used 5 to screen for, characterize and/or isolate a defective mammalian CXCR3 gene, which encodes a receptor having reduced or enhanced activity. Standard methods of screening for a defective gene can be employed, for instance. A defective gene can be isolated and expressed 10 in a suitable host cell for further assessment as described herein for mammalian CXCR3 proteins. A number of human diseases are associated with defects in the function of a G-protein coupled receptor (Clapham, D.E., *Cell*, 75: 1237-1239 (1993); Lefkowitz, R.J., *Nature*, 365: 603-04 (1993)).

15 The nucleic acids of the present invention provide reagents, such as probes and PCR primers, which can also be used to assess expression of receptor (e.g., by detecting transcription of mRNA) by cells in a sample (e.g., by Northern analysis, by in situ hybridization). For example, 20 expression in activated T lymphocytes or other cell types can be assessed.

The antibodies of the present invention have application in procedures in which receptor can be detected on the surface of cells. The receptor provides a marker of 25 the leukocyte cell types in which it is expressed, particularly of activated T cells. For example, antibodies raised against a receptor protein or peptide, such as the antibodies described herein (e.g., mAb 1C6), can be used to detect and/or quantify cells expressing receptor. In one 30 embodiment, the antibodies can be used to sort cells which express receptor from among a mixture of cells (e.g., to isolate activated T cells, such as CD4<sup>+</sup> T cells). Suitable methods for counting and/or sorting cells can be used for this purpose (e.g., flow cytometry, fluorescence activated 35 cell sorting). Cell counts can be used in the diagnosis of

-59-

diseases or conditions in which an increase or decrease in leukocyte cell types (e.g., activated T cells) is observed. The presence of an increased level of activated T cells in a sample obtained from an individual can be indicative of 5 infiltration due to an inflammatory disease or condition, such as a delayed type hypersensitivity reaction, allograft rejection, or a pathologic condition, including bacterial or viral infection.

Furthermore, the antibodies can be used to detect or 10 measure expression of receptor. For example, antibodies of the present invention can be used to detect or measure receptor in a sample (e.g., tissues or body fluids from an individual such as blood, serum, leukocytes (e.g., activated T lymphocytes), bronchoalveolar lavage fluid, 15 saliva, bowel fluid). For example, a sample (e.g., tissue and/or fluid) can be obtained from an individual and a suitable assay can be used to assess the presence or amount of CXCR3 protein. Suitable assays include immunological methods such as FACS analysis and enzyme-linked 20 immunosorbent assays (ELISA), including chemiluminescence assays, radioimmunoassay, and immunohistology. Generally, a sample and antibody of the present invention are combined under conditions suitable for the formation of an antibody-receptor complex, and the formation of antibody-receptor 25 complex is assessed (directly or indirectly).

The presence of an increased level of receptor reactivity in a sample obtained from an individual can be indicative of inflammation and/or leukocyte (e.g., activated T cell) infiltration and/or accumulation 30 associated with an inflammatory disease or condition, such as allograft rejection, delayed type hypersensitivity reaction, or an infection such as a viral or bacterial infection. The level of expression of a mammalian CXCR3 protein or variant can also be used to correlate increased 35 or decreased expression of a mammalian CXCR3 protein with a

- 60 -

particular disease or condition, and in the diagnosis of a disease or condition in which increased or decreased expression of a mammalian CXCR3 protein occurs (e.g., increased or decreased relative to a suitable control, such 5 as the level of expression in a normal individual). Similarly, the course of therapy can be monitored by assessing CXCR3 immunoreactivity in a sample from a patient. For example, antibodies of the present invention can be used to monitor the number of cells bearing CXCR3 in 10 a sample (e.g., blood, tissue) from a patient being treated with an anti-inflammatory or immunosuppressive agent.

Transgenic Animals

Transgenic animals, in which the genome of the animal host is altered using recombinant DNA techniques, can be 15 constructed. In one embodiment, the alteration is not heritable (e.g., somatic cells, such as progenitor cells in bone marrow, are altered). In another embodiment, the alteration is heritable (the germ line is altered). Transgenic animals can be constructed using standard 20 techniques or other suitable methods (see e.g., Cooke, M.P. et al., *Cell*, 65: 281-291 (1991) regarding alteration of T lymphocytes; Hanahan, D., *Science*, 246: 1265-1275, (1989); Anderson et al., U.S. Patent No. 5,399,346).

In one aspect, an endogenous mammalian CXCR3 gene can 25 be inactivated or disabled, in whole or in part, in a suitable animal host (e.g., by gene disruption techniques) to produce a transgenic animal. Nucleic acids of the present invention can be used to assess successful construction of a host containing an inactivated or 30 disabled CXCR3 gene (e.g., by Southern hybridization). In addition, successful construction of a host containing an inactivated or disabled CXCR3 gene can be assessed by suitable assays which monitor the function of the encoded receptor. Such animals can be used to assess the effect of

- 61 -

receptor inactivation on inflammation and host defenses against cancer and pathogens (e.g., a viral pathogen).

In another embodiment, a nucleic acid encoding a mammalian CXCR3 protein or polypeptide is introduced into a 5 suitable host to produce a transgenic animal. In a preferred embodiment, endogenous CXCR3 receptor genes present in the transgenic animals are inactivated (e.g., simultaneously with introduction of the nucleic acid by homologous recombination, which disrupts and replaces the 10 endogenous gene). For example, a transgenic animal (e.g., a mouse, guinea pig, sheep) capable of expressing a nucleic acid encoding a mammalian CXCR3 receptor of a different mammalian species (e.g., a human CXCR3 such as the CXCR3 encoded by SEQ ID NO:1) in leukocytes (such as lymphocytes 15 (e.g., activated T lymphocytes), natural killer cells) can be produced, and provides a convenient animal model for assessing the function of the introduced receptor. In addition, a test agent can be administered to the transgenic animal, and the effect of the agent on a 20 receptor-mediated process (e.g., inflammation) can be monitored as described herein or using other suitable assays. In this manner, agents which inhibit or promote receptor function can be identified or assessed for *in vivo* effect.

25 Methods of Therapy

Modulation of mammalian CXCR3 function according to the present invention, through the inhibition or promotion of at least one function characteristic of a mammalian CXCR3 protein, provides an effective and selective way of 30 inhibiting or promoting receptor-mediated functions. As CXC chemokine receptors selectively expressed on activated lymphocytes, responsive to chemokines such as IP-10 and Mig whose primary targets are lymphocytes, particularly effector cells such as activated or stimulated T

-62-

lymphocytes and NK cells, mammalian CXCR3 proteins provide a target for selectively interfering with or promoting lymphocyte function in a mammal, such as a human. Once lymphocytes are recruited to a site, other leukocyte types, 5 such as monocytes, may be recruited by secondary signals. Thus, agents which inhibit or promote CXCR3 function, including ligands, inhibitors (e.g., 1C6) and/or promoters, such as those identified as described herein, can be used to modulate leukocyte function (e.g., leukocyte 10 infiltration including recruitment and/or accumulation), particularly of lymphocytes, for therapeutic purposes.

In one aspect, the present invention provides a method of inhibiting or promoting an inflammatory response in an individual in need of such therapy, comprising 15 administering an agent which inhibits or promotes mammalian CXCR3 function to an individual in need of such therapy. In one embodiment, a compound which inhibits one or more functions of a mammalian CXCR3 protein (e.g., a human CXCR3) is administered to inhibit (i.e., reduce or prevent) 20 inflammation. For example, antibodies of the present invention, including mAb 1C6 can be used in the method. As a result, one or more inflammatory processes, such as leukocyte emigration, chemotaxis, exocytosis (e.g., of enzymes) or inflammatory mediator release, is inhibited. 25 For example, leukocytic infiltration of inflammatory sites (e.g., in a delayed-type hypersensitivity response) can be inhibited according to the present method.

In another embodiment, an agent (e.g., receptor agonist) which promotes one or more functions of a 30 mammalian CXCR3 protein (e.g., a human CXCR3) is administered to induce (trigger or enhance) an inflammatory response, such as leukocyte emigration, chemotaxis, exocytosis (e.g., of enzymes) or inflammatory mediator release, resulting in the beneficial stimulation of 35 inflammatory processes. For example, natural killer cells

-63-

can be recruited to combat viral infections or neoplastic disease.

The term "individual" is defined herein to include animals such as mammals, including, but not limited to, 5 primates (e.g., humans), cows, sheep, goats, horses, dogs, cats, rabbits, guinea pigs, rats, mice or other bovine, ovine, equine, canine, feline, rodent or murine species. Diseases and conditions associated with inflammation, 10 infection, and cancer can be treated using the method. In a preferred embodiment, the disease or condition is one in which the actions of lymphocytes, particularly effector 15 cells such as activated or stimulated T lymphocytes and natural killer (NK) cells, are to be inhibited or promoted for therapeutic (including prophylactic) purposes. In a particularly preferred embodiment, the inflammatory disease or condition is a T cell-mediated disease or condition.

Diseases or conditions, including chronic diseases, of 20 humans or other species which can be treated with inhibitors of CXCR3 function, include, but are not limited to:

- inflammatory or allergic diseases and conditions, including systemic anaphylaxis or hypersensitivity responses, drug allergies (e.g., to penicillin, cephalosporins), insect sting allergies; inflammatory bowel 25 diseases, such as Crohn's disease, ulcerative colitis, ileitis and enteritis; vaginitis; psoriasis and inflammatory dermatoses such as dermatitis, eczema, atopic dermatitis, allergic contact dermatitis, urticaria; vasculitis (e.g., necrotizing, cutaneous, and 30 hypersensitivity vasculitis); spondyloarthropathies; scleroderma; respiratory allergic diseases such as asthma, allergic rhinitis, hypersensitivity lung diseases, hypersensitivity pneumonitis, interstitial lung diseases

-64-

(ILD) (e.g., idiopathic pulmonary fibrosis, or ILD associated with rheumatoid arthritis, or other autoimmune conditions);

- autoimmune diseases, such as arthritis (e.g.,  
5 rheumatoid arthritis, psoriatic arthritis), multiple sclerosis, systemic lupus erythematosus, myasthenia gravis, diabetes, including diabetes mellitus and juvenile onset diabetes, glomerulonephritis and other nephritides, autoimmune thyroiditis, Behcet's disease;
- 10 • graft rejection (e.g., in transplantation), including allograft rejection or graft-versus-host disease;
- other diseases or conditions in which undesirable inflammatory responses are to be inhibited can be treated, including, but not limited to, atherosclerosis, cytokine-  
15 induced toxicity, myositis (including polymyositis, dermatomyositis).

Diseases or conditions of humans or other species which can be treated with promoters (e.g., an agonist) of CXCR3 function, include, but are not limited to:

- 20 • cancers, particularly those with leukocytic infiltration of the skin or organs such as cutaneous T cell lymphoma (e.g., mycosis fungoides);
- diseases in which angiogenesis or neovascularization plays a role, including neoplastic disease, retinopathy  
25 (e.g., diabetic retinopathy), and macular degeneration;
- infectious diseases, such as bacterial infections and tuberculoid leprosy, and especially viral infections;

- 65 -

- immunosuppression, such as that in individuals with immunodeficiency syndromes such as AIDS, individuals undergoing radiation therapy, chemotherapy, or other therapy which causes immunosuppression; immunosuppression 5 due congenital deficiency in receptor function or other causes. Promoters of CXCR3 function can also have protective effects useful to combat stem cell depletion during cancer chemotherapy (Sarris, A.H. et al., *J. Exp. Med.*, 178: 1127-1132 (1993)).

10

#### *Modes of Administration*

According to the method, one or more agents can be administered to the host by an appropriate route, either alone or in combination with another drug. An effective 15 amount of an agent (e.g., a receptor peptide which inhibits ligand binding, an anti-CXCR3 antibody or antigen-binding fragment thereof) is administered. An effective amount is an amount sufficient to achieve the desired therapeutic or prophylactic effect, under the conditions of 20 administration, such as an amount sufficient for inhibition or promotion of CXCR3 receptor function, and thereby, inhibition or promotion, respectively, of a receptor-mediated process (e.g., an inflammatory response).

A variety of routes of administration are possible 25 including, but not necessarily limited to oral, dietary, topical, parenteral (e.g., intravenous, intraarterial, intramuscular, subcutaneous injection), and inhalation (e.g., intrabronchial, intranasal or oral inhalation, intranasal drops) routes of administration, depending on 30 the agent and disease or condition to be treated. For respiratory allergic diseases such as asthma, inhalation is a preferred mode of administration.

Formulation of an agent to be administered will vary 35 according to the route of administration selected (e.g., solution, emulsion, capsule). An appropriate composition

- 66 -

comprising the agent to be administered can be prepared in a physiologically acceptable vehicle or carrier. For solutions or emulsions, suitable carriers include, for example, aqueous or alcoholic/aqueous solutions, emulsions 5 or suspensions, including saline and buffered media. Parenteral vehicles can include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils, for instance. Intravenous vehicles 10 can include various additives, preservatives, or fluid, nutrient or electrolyte replenishers and the like (See, generally, *Remington's Pharmaceutical Sciences*, 17th Edition, Mack Publishing Co., PA, 1985). For inhalation, the agent can be solubilized and loaded into a suitable 15 dispenser for administration (e.g., an atomizer, nebulizer or pressurized aerosol dispenser).

Furthermore, where the agent is a protein or peptide, the agent can be administered via *in vivo* expression of the recombinant protein. *In vivo* expression can be accomplished via somatic cell expression according to 20 suitable methods (see, e.g. U.S. Patent No. 5,399,346). In this embodiment, nucleic acid encoding the protein can be incorporated into a retroviral, adenoviral or other suitable vector (preferably, a replication deficient 25 infectious vector) for delivery, or can be introduced into a transfected or transformed host cell capable of expressing the protein for delivery. In the latter embodiment, the cells can be implanted (alone or in a barrier device), injected or otherwise introduced in an 30 amount effective to express the protein in a therapeutically effective amount.

-67-

EXEMPLIFICATION

The present invention will now be illustrated by the following Examples, which are not intended to be limiting in any way.

5 *Human chemokines (Examples 1-2)*

The CXC chemokines Mig, IL-8, GRO $\alpha$ , NAP-2, GCP-2, ENA78, PF4, the CC chemokines MCP-1, MCP-2, MCP-3, MCP-4, MIP-1 $\alpha$ , MIP-1 $\beta$ , RANTES, I309, eotaxin and the chemokine-related lymphotactin were chemically synthesized 10 according to established protocols (Clark-Lewis, I. et al., *Biochemistry* 30: 3128-3135 (1991)). The CXC chemokine IP-10 was purchased from PeproTech, Rocky Hill, NJ.

Example 1. Cloning of Receptor cDNA

Standard molecular biology techniques were used 15 (Sambrook, J. et al., 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

DNA fragments coding for putative T lymphocyte-restricted chemokine receptors were generated using the 20 polymerase chain reaction (PCR). Two degenerate oligonucleotide primers were designed based on conserved motifs of chemokine receptors. Primer design was based on the conserved nucleotide sequences within transmembrane domain 2 (TM2) and transmembrane domain 7 (TM7) of the 25 chemokine receptors IL-8R1 (CXCR1), IL-8R2 (CXCR2), CC-CKR1 (CCR1), CC-CKR2 (CCR2) and the orphan receptors EBI I, LESTR, and BLR1/MDR15 (EBI I, Birkenbach, M. et al., *J. Virol.*, 67: 2209-2220 (1993)); LESTR, Loetscher, M. et al., *J. Biol. Chem.*, 269: 232-237 (1994); and BLR1/MDR15, 30 Dobner, T. et al., *Eur. J. Immunol.*, 22: 2795-2799 (1992) and Barella, L. et al., *Biochem. J.*, 309: 773-779 (1995)).

The sequences of the primers were as follows:

- 68 -

SEQ ID NO:3:

5'-GGG CTG CAG CII T(T/G) (T/G) C(C/A)G AC(A/C) TIC  
TI(C/T) T-3'

SEQ ID NO:4:

5 5'-GGG TCT AGA IGG GTT IAI (G/A)CA (G/A)C(T/A)  
(G/A) (T/C)G-3'

(I = inosine). These primers were used in a polymerase chain reaction (PCR) to amplify DNA fragments using human genomic DNA isolated from human peripheral blood

10 lymphocytes as template as follows. A 100  $\mu$ l reaction mixture containing 2  $\mu$ g human genomic DNA, 1X DynaZyme buffer (Finnzymes OY, Espoo, Finland), 1.5 mM MgCl<sub>2</sub>, 500  $\mu$ M of each deoxynucleotide, 1  $\mu$ M of both primers, and 2.5 U of DynaZyme DNA polymerase was subjected to 30 cycles (94°C  
15 for 1 minute; 55°C for 1 minute; and 72°C for 2 minutes) on a DNA thermal cycler (Techne PHC-2, Brouwer, Switzerland). PCR products of the predicted size (approximately 700 bp) were cloned into the Gene Scribe-Z vectors pTZ18/19 U/R (USB, Cleveland, OH), were partially sequenced (Sanger, F.  
20 et al., Proc. Natl. Acad. Sci. USA, 74: 5463-5467 (1977)), and were evaluated for their similarity to known chemokine receptors and for expression of their corresponding mRNA in leukocytes. A DNA fragment designated 2MLC22 revealed 64% nucleotide sequence identity with IL-8R2. Fragment 2MLC22  
25 specifically hybridized to RNA from T cells, but not monocytes or neutrophils, as assessed by Northern blot analysis using a hybridization probe prepared by enzymatically labeling 2MLC22 with the radioactive isotope <sup>32</sup>P using Klenow fragment of DNA Polymerase I and a  
30 commercially available random-prime labeling kit.

Fragment 2MLC22 was enzymatically labeled with <sup>32</sup>P as described and used as a probe to screen a human tetanus toxoid-specific CD4<sup>+</sup> T cell (KT30) cDNA library, prepared

-69-

in lambda-ZAP Express (Stratagene, Zurich, Switzerland) (Loetscher, M. et al., *J. Biol. Chem.*, 269: 232-237 (1994)). A cDNA library was prepared in a  $\lambda$  ZAP Express system according to the manufacturer's protocol (Stratagene 5 GMBH, Zurich, Switzerland) using poly(A)<sup>+</sup> RNA from human tetanus toxoid-specific CD4<sup>+</sup> T cells (KT30). The resulting library contained about  $1.8 \times 10^6$  independent clones with an average insert size of approximately 1.1 kb. For plaque hybridization screening, about  $4 \times 10^5$  clones were 10 transferred onto Biodyne nylon membranes (PALL AG, Muttenz, Switzerland) and probed with 2MLC22 which had been labeled to a specific activity of  $1 \times 10^9$  dpm/ $\mu$ g DNA using the high prime DNA labeling kit (Boehringer Mannheim, Mannheim, Germany). Hybridization was carried out in 50% formamide, 15 6X SSC, 0.5% SDS, 100  $\mu$ g/ml denatured salmon sperm DNA at 42°C for 20 hours using  $1 \times 10^6$  dpm 2MLC22/ml hybridization solution. The membranes were washed once in 2X SSC, 0.1% SDS at room temperature for 10 minutes, twice in 1X SSC, 0.1% SDS at 65°C for 30 minutes, and finally once 20 in 0.5X SSC, 0.1% SDS at 65°C for 10 minutes. Twenty-three clones were isolated from hybridization positive lambda plaques following the high stringency washes, and the clone with the largest insert (1670 bp) was sequenced. CXCR3 cDNA was subcloned into commercially available plasmid 25 vectors for nucleotide sequencing, generation of hybridization probes, and construction of stably transfected mammalian cell clones expressing CXCR3, and these CXCR3 cDNA-containing constructs are maintained in *E. coli* strains.

30 Results

A cDNA was isolated from a human CD4<sup>+</sup> T cell library by searching for T lymphocyte-specific chemokine receptors (Figure 1, SEQ ID NO:1). This cDNA was not recovered in the course of searching a commonly used monocyte-derived

- 70 -

cDNA library or granulocyte (HL60)-derived cDNA library for novel chemokine receptor cDNAs; however, a direct search of the libraries specifically for CXCR3 cDNA has not been conducted. The CXCR3 cDNA, which was shown to encode an

5 IP-10/Mig receptor (see below), and has an open reading frame (ORF) of 1104 bp beginning at residue 69 which encodes a protein of 368 amino acids with a predicted molecular mass of 40,659 daltons. The amino acid sequence (Figure 2, SEQ ID NO:2) includes seven putative

10 transmembrane segments, which are characteristic of G-protein coupled receptors, and three potential N-glycosylation sites (Asn<sup>22</sup>, Asn<sup>32</sup>, and Asn<sup>199</sup>) (Figure 2). In addition, one threonine and nine serine residues, which are potential phosphorylation sites for receptor kinases

15 (Palczewski, K. and J.L. Benovic, *Trends Biochem. Sci.*, 16: 387-391 (1991); Chuang, T.T. et al., *J. Biol. Chem.*, 267: 6886-6892 (1992); and Giannini, E. et al., *J. Biol. Chem.*, 270: 19166-19172 (1995)), can be found in the intracellular COOH-terminal region (Figure 2).

20 The 368 amino acid sequence of the receptor (IP-10/MigR, Figure 2, SEQ ID NO:2) was aligned with the amino acid sequences of other human chemokine receptors, including IL-8R1 (CXCR1), IL-8R2 (CXCR2), CC-CKR1 (CCR1), CC-CKR2A (CCR2a), CC-CKR3 (CCR3) and CC-CKR4 (CCR4).

25 Multiple protein alignment was performed according to Higgins and Sharp (Higgins, D.G. and P.M. Sharp, "Description of the method used in CLUSTAL," *Gene*, 73: 237-244 (1988)). Residues framed in black in Figure 2 represent regions of identity between IP-10/MigR and at

30 least two other chemokine receptors. Hyphens indicate gaps in the alignment. The alignment revealed several conserved motifs, particularly in the transmembrane domains and the second intracellular loop. Significant sequence identity with CXC receptors IL-8R1 and IL-8R2, but not with the CC

- 71 -

chemokine receptors, was observed in the third and the sixth transmembrane domains (Figure 2).

The sequence shares 40.9% and 40.3% amino acid identity overall with the IL-8R1 and IL-8R2 receptors, 5 respectively, and 34.2 to 36.9% identity with the five known CC chemokine receptors (Table 1). A lower degree of similarity was found with seven-transmembrane-domain receptors that are expressed in T cells, but which do not bind chemokines, e.g., 27.2% identity with the thrombin 10 receptor (Vu, T.-K.H. et al., *Cell*, 64: 1057-1068 (1991)). A truncated clone of unidentified function, with an incomplete coding sequence which can be aligned with that of Figure 2, was previously isolated from a human genomic DNA library (Marchese, A. et al., *Genomics*, 29: 335-344 15 (1995)).

TABLE 1  
Amino Acid Sequence Comparison of IP-10/MigR with Human Chemokine Receptors

	IL-8R1 (CXCR1)	IL-8R2 (CXCR2)	CC-CKR1 (CCR1)	CC-CKR2A (CCR2a)	CC-CKR3 (CCR3)	CC-CKR4 (CCR4)	CC-CKR5 (CCR5)	ThrombR
IP-10/MigR	40.9 <sup>a</sup>	40.3	34.9	34.2	34.4	35.8	36.9	27.2
IL-8R1		77.1	33.7	32.9	34.3	39.7	34.3	29.1
IL-8R2			34.9	33.6	34.1	40.8	34.4	29.7
CC-CKR1				54.1	63.1	49.3	56.3	26.8
CC-CKR2A					50.7	46.1	68.8	24.6
CC-CKR3						46.5	52.3	27.3
CC-CKR4							50.0	29.2
CC-CKR5								23.6

<sup>a</sup> Numbers refer to percentage amino acid identity. Pairwise protein sequence alignments were carried out using the program PALIGN with an open gap cost and unit gap cost of 3 and 2, respectively.

- 73 -

Example 2. Biological Activity

*Expression in Activated T Lymphocytes*

In view of the observed chemokine selectivity, the occurrence of the IP-10/MigR in leukocytes and related cell lines was examined by Northern blot analysis. 10  $\mu$ g samples of total RNA were examined from freshly isolated human blood monocytes, neutrophils, lymphocytes (PBL), nylon-wool purified T cells, and from cultured cells including cloned human CD4 $^{+}$  T cells (KT30) and CD8 $^{+}$  T cells (ERCD8), cloned NK cells (ERNK57), and PBL cultured for 10 days (1-2.5  $\times$  10 $^{6}$  cells/ml in RPMI 1640 medium containing 2 mM glutamine, 1X non-essential amino acids, 1 mM sodium pyruvate, 100  $\mu$ g/ml kanamycin, 5  $\times$  10 $^{-5}$  M 2-mercaptoethanol, and 5% human serum) in the presence of 400 U/ml hrIL-2. (human recombinant IL-2 was a gift of Dr. A. Lanzavecchia, Basel Institute of Immunology, Basel, Switzerland). Agarose gels were stained with ethidium bromide to check the integrity and amount of total RNA on the gel prior to blotting. RNA samples were analyzed with  $^{32}$ P-labeled 5'-fragment of the IP-10/MigR DNA (10 $^9$  cpm/ $\mu$ g DNA) at 5  $\times$  10 $^6$  cpm/ml hybridization solution as described (Loetscher, M. et al., *J. Biol. Chem.*, 269: 232-237 (1994)). The 5'-fragment used as a Northern probe was prepared by digestion of CXCR3 cDNA in pBK-CMV vector (Stratagene GMBH, Zurich, Switzerland) with PstI yielding the 724 bp 5'-end of the CXCR3 cDNA (Figure 1).

Results

Abundant expression of mRNA of the expected size was found in the cloned CD4 $^{+}$  T cells, KT30, that were used for isolation of the receptor cDNA. Similar levels of expression were observed in the CD8 $^{+}$  T cell clone, ERCD8, and the NK cell clone, ERNK57. In contrast, in freshly isolated blood lymphocytes and nylon-wool purified T cells,

- 74 -

IL10/MigR transcripts were barely detectable. However, when these cells were cultured in the presence of IL-2, a strong upregulation was obtained, and the level of receptor mRNA approached that of T and NK cell clones. No

5 IP-10/MigR transcripts were detected under these conditions in freshly isolated blood monocytes, neutrophil leukocytes, or eosinophil leukocytes. Additional leukocyte-related cells that did not express IP-10/MigR mRNA include the mast cell line, HMC-1, the promyelocytic leukemia line, HL60, 10 the histiocytic lymphoma, U937, the chronic myelogenous leukemia line, K562, the acute T cell leukemia line, Jurkat, the acute lymphoblastic leukemia line, Molt, the B-lymphoblastic cell lines Daudi and Raji, lymphocytes from patients with chronic and acute B-lymphoid leukemia (B-CLL 15 and B-ALL), mature basophils from a patient with basophilic leukemia, and the erythroleukemia cell line, HEL. By contrast, the receptors for chemokines which have been shown previously to attract lymphocytes, i.e. MCP-1 MCP-2, MCP-3, MIP-1 $\alpha$ , MIP-1 $\beta$  and RANTES (Loetscher, P. et al., 20 FASEB J., 8: 1055-1060 (1994); Carr, M.W. et al., Proc. Natl. Acad. Sci. USA 91: 3652-3656 (1994); Taub, D.D. et al., Science, 260: 355-358 (1993); Schall, T.J. et al., J. Exp. Med., 177: 1821-1825 (1993); Schall, T.J. et al., Nature, 347: 669-672 (1990)), are also found in monocytes 25 and granulocytes. The restricted expression of IP-10/MigR in activated T lymphocytes and a natural killer cell line suggests that this novel receptor can mediate selective lymphocyte recruitment.

#### Stable Transfectants

30 CXCR3 cDNA was released from pBK-CMV (Stratagene GMBH, Zurich, Switzerland) by digestion with BamHI and XbaI, and was cloned into BamHI and XbaI sites of pcDNA3 (Invitrogen BV, WB Leek, Netherlands) to yield pcDNA3-Clone8, which is maintained and stored Escherichia coli (XL1Blue).

-75-

To generate stable transfectants,  $4 \times 10^6$  of either mouse pre-B cells (300-19) (Thelen, M. et al., *FASEB J.*, 2: 2702-2706 (1988)), human promyelocytic cells (GM-1) (Garotta, G. et al., *J. Leukocyte Biol.*, 49: 294-301 (1991)) or human acute T cell leukemia cells (Jurkat) (Loetscher, P. et al., *FEBS Lett.* 341: 187-192 (1994)), were transfected by electroporation with 20  $\mu$ g of receptor cDNA in pcDNA3 which was linearized with Bgl II as described previously (Moser, B. et al., *Biochem. J.*, 294: 10 285-292 (1993)).

IP-10/MigR transfected cells were cloned by limiting dilution under G-418 (Life Technologies, Inc.) selection (1.0 mg/ml G-418 for 300-19 and 0.8 mg/ml G-418 for Jurkat and GM-1 cells). G-418 resistant clones were screened for 15 receptor expression by RNA Dot-blot analysis.

#### $\text{Ca}^{2+}$ Flux

To determine whether the receptor was functional, clones of murine pre-B cells (300-19), human promyelocytic cells (GM-1), and human T cell leukemia cells (Jurkat) were 20 stably transfected with receptor cDNA as described above. Activation of chemokine receptors leads to a transient rise in the cytosolic free  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ), and this assay was used to monitor signalling in the transfected cells.

25 Changes in the cytosolic free  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) were measured in cells loaded with fura-2 by incubation for 30 minutes at 37°C with 0.1 nmol fura-2 acetoxyxymethyl ester per  $10^6$  cells in a buffer containing 136 mM NaCl, 4.8 mM KCl, 1 mM  $\text{CaCl}_2$ , 5 mM glucose, and 20 mM 30 HEPES, pH 7.4. After centrifugation, loaded cells were resuspended in the same buffer ( $10^6$  cells/ml), stimulated with the indicated chemokine at 37°C, and the  $[\text{Ca}^{2+}]_i$ -related fluorescence changes were recorded (von Tscharner, V. et al., *Nature*, 324: 69-372 (1986)).

Results

A rapid  $[Ca^{2+}]_i$  rise was observed in response to IP-10 and Mig. The chemokine IP-10 has been shown to be expressed in cutaneous delayed-type hypersensitivity reactions (Luster, A.D. et al., *Nature*, 315: 672-676 (1985); Kaplan, G. et al., *J. Exp. Med.*, 166: 1098-1108 (1987)). The chemokine designated Mig was recently identified (Farber, J.M., *Proc. Natl. Acad. Sci. USA*, 87: 5238-5242 (1990); Farber, J.M., *Biophys. Res. Commun.*, 192: 10 223-230 (1993)). Both chemokines have the CXC arrangement of the first two cysteines like IL-8, but are not chemotactic for neutrophil leukocytes. It was recently reported that IP-10 attracts T lymphocytes (Luster, A.D. and P. Leder, *J. Exp. Med.*, 178: 1057-1065 (1993); Taub, D.D. et al., *J. Exp. Med.*, 177: 1809-1814 (1993)), and that Mig is chemotactic for tumor-associated lymphocytes (Liao, F. et al., *J. Exp. Med.*, 182: 1301-1314 (1995)).

Figures 3A-3C summarize the effects of IP-10 and Mig on cells transfected with the cDNA and expressing the functional IP-10/MigR. As shown by the  $[Ca^{2+}]_i$  changes (Figure 3A), the action of IP-10 and Mig was concentration dependent and already detectable at 1 nM, indicating that both chemokines have high affinity for the novel receptor. The IP-10/MigR transfectants, by contrast, did not respond to any of 16 other potential agonists at concentrations up to 100 nM, including the CXC chemokines IL-8, GRO $\alpha$ , NAP-2, GCP-2, ENA78, PF4, the CC chemokines MCP-1, MCP-2, MCP-3, MCP-4, MIP-1 $\alpha$ , MIP-1 $\beta$ , RANTES, I309, eotaxin or the chemokine related lymphotactin (not shown). Identical results were obtained with the murine and the human transfected cells. These observations demonstrate that the novel receptor is highly selective for IP-10 and Mig. Accordingly, the receptor is referred to herein as an IP-10/Mig receptor (IP-10/MigR), or as "CXCR3", reflecting its specificity for CXC chemokines.

-77-

As shown in Figure 3B, repeated stimulation with IP-10 or Mig resulted in desensitization typical of chemokine receptors. Furthermore, cross-desensitization occurred when the cells were stimulated with IP-10 followed by Mig 5 or vice versa, confirming that the receptor has high affinity for both chemokines. At 100 nM concentration, it became evident that Mig was more potent in cross-desensitization than IP-10, suggesting higher affinity or binding stability of the IP-10/Mig receptor for 10 Mig.

While expression of functional IP-10/MigR was demonstrated, binding experiments using radioactive ligands revealed non-specific binding between 60 and 80% of the total, preventing determination of binding parameters. 15 Since IP-10 and Mig are highly cationic (pI values of 10.8 and 11.1), nonspecific interaction with cell surface proteoglycans may explain these results. Indeed, chemokine receptor-unrelated, heparinase-sensitive binding sites for IP-10 (and PF4) have been detected on a variety of blood 20 and tissue cells (Luster, A.D. et al., *J. Exp. Med.*, 182: 219-231 (1995)), and heparan sulfate binds IP-10 and Mig and prevents lymphocyte chemotaxis (not shown). The heparin binding site is probably not involved in CXCR3 receptor binding, and inclusion of a suitable heparin 25 derivative such as chondroitin sulfate in the reaction (e.g., in binding buffer) can be used to inhibit non-specific binding to cells through the heparin binding site.

#### Chemotaxis

30 PBL were freshly isolated from donor blood buffy coats. Donor blood buffy coats were provided by the Swiss Central Laboratory Blood Transfusion Service, SRK. Isolation of buffy coat PBL was performed as described in Colotta, F. et al., *J. Immunol.*, 132: 936-944 (1984).

- 78 -

Freshly isolated PBL from donor blood buffy coats were used without further processing, or were used after culturing for 10 days in the presence of IL-2 (1-2.5  $\times$  10<sup>6</sup> cells/ml in RPMI 1640 medium containing 2 mM glutamine, 1X 5 non-essential amino acids, 1 mM sodium pyruvate, 100  $\mu$ g/ml kanamycin, 5  $\times$  10<sup>-5</sup> M 2-mercaptoethanol, and 5% human serum in the presence of 400 U/ml hrIL-2).

Cell migration was assessed in 48-well chambers (Neuro Probe, Cabin John, MD, USA) using polyvinylpyrrolidone-free 10 polycarbonate membranes (Nucleopore) with 5- $\mu$ m pores for IP-10/MigR transfected cells (Loetscher, P. et al., FEBS Lett. 341: 187-192 (1994)) or with 3- $\mu$ m pores for human PBL (Loetscher, P. et al., FASEB J., 8: 1055-1060 (1994)). RPMI 1640 supplemented with 20 mM Hepes, pH 7.4, and 1% 15 pasteurized plasma protein solution (Swiss Red Cross Laboratory, Bern, Switzerland) was used to dissolve the chemokines (lower wells), and to dilute the cells (100,000 receptor transfectants or PBL in the upper well). After 60 minutes at 37°C, the membrane was removed, washed on the 20 upper side with PBS, fixed and stained. All assays were done in triplicate, and the migrated cells were counted in five randomly selected fields at 1,000-fold magnification. Spontaneous migration was determined in the absence of chemoattractant.

25 Results - Transfected Cells

Transfected cells expressing the IP-10/MigR readily migrated toward IP-10 or Mig, while the non-transfected, parental cells did not respond (Figure 3C). Both agonists showed a typically biphasic concentration dependence. 30 IP-10 induced migration at concentrations above 1 nM, whereas the response of Mig became detectable above 10 nM. The efficacy, which is measured by the maximum number of migrating cells, was about twice as high for Mig as for IP-10. These results demonstrate that the IP-10/MigR, like

-79-

all known chemokine receptors in leukocytes, mediates chemotaxis in response to ligand.

#### Results - Human Blood Leukocytes

In agreement with the cellular distribution of the IP-5 10/MigR, activated human T lymphocytes were found to be highly responsive to IP-10 and Mig (Figures 4A-4B). The activity of IP-10 and Mig as inducers of  $[Ca^{2+}]_i$  changes (Figure 4A) and *in vitro* chemotaxis (Figure 4B) was consistent with the effects observed using transfected 10 cells expressing the IP-10/MigR, with IP-10 being more potent but less efficacious than Mig. Activation of the T lymphocytes by culturing in the presence of IL-2 was required for induction of calcium flux and chemotaxis, and no response was observed with freshly isolated blood 15 lymphocytes under the conditions used.

#### Materials and Methods for Examples 3-9

The following materials and methods were used in Examples 3-9.

##### *Chemokines*

20 Recombinant human chemokines were obtained from Peprotech (Rocky Hill, NJ), except for eotaxin, described previously (Ponath, P.D., et al., *J. Clin. Invest.*, 97:604-612 (1996); see also Ponath et al., WO 97/00960, published 9 January 1997), which was a gift from Dr. Ian Clark-Lewis. 25  $^{125}I$ -labeled chemokines were obtained from Du Pont NEN (Boston, MA).

##### *Cells and Cell Lines*

Neutrophils and PBMCs were isolated as described (Ponath, P.D., et al., *J. Clin. Invest.*, 97:604-612 30 (1996)). To generate CD3 blasts,  $2 \times 10^6$  PBMC/ml in RPMI-1640 plus 10% FCS were added to tissue culture plates which

-80-

had been coated with the anti-CD3 antibody TR66. After 4-6 days blasts were removed to fresh media and supplemented with IL-2 (kindly provided by Antonio Lanzavecchia, Basel) at 50 units/ml.

5 Other cell lines used included transfectants of the L1.2 murine pre B cell lymphoma, expressing either CXCR3 (see below), IL-8 RA (Ponath, P.D., et al., *J. Exp. Med.*, 183:2437-2448 (1996)), IL-8 RB (Ponath, P.D., et al., *J. Exp. Med.*, 183:2437-2448 (1996)), CCR2b (G. LaRosa, 10 unpublished), CCR4, CCR5 (Wu, L., et al., *Nature*, 384:179-183 (1996)), or CCR1 (Campbell, J.J., et al., *J. Cell Biol.*, 134:255-266 (1996)).

#### *Preparation of CXCR3 Transfectants*

##### *Cells*

15 L1.2 cells were grown in RPMI medium 1640, 10% Fetal Clone (from Hyclone, Inc.), 50 U/ml Penicillin/Streptomycin, 1X L-Glutamine, 1 mM NaPyruvate, and 5.5 x 10<sup>-5</sup>M  $\beta$ -Mercaptoethanol. Media components were purchased from GibcoBRL, except for 10% Fetal Clone, which was 20 purchased from Hyclone, Inc. Two days prior to transfection, the L1.2 cells were diluted 1:5 into fresh medium. This resulted in 150 million cells in log phase growth at a concentration of about 1-3 million cells/ml.

##### *CXCR3 DNA and Transfection*

25 *E. coli* XL1Blue cells (Stratagene, Inc. (Cat# 200236)) were transformed with pcDNA3-Clone8 (Example 2; Loetscher, M., et al., *J. Exp. Med.*, 184:963-969 (1996)) according to the manufacturer's protocol. Transformants were grown at 37°C while shaking at 250 rpm in 500 ml of LB containing 30 100  $\mu$ g/ml Ampicillin. The culture was then collected by centrifugation at 8,000 X g, and the plasmid was purified using a Maxi plasmid purification column and protocol (Qiagen, Cat# 12162). Plasmid concentration and purity

-81-

were determined using a 1% agarose gel and OD260/280 ratios. Plasmid DNA was suspended in ddH<sub>2</sub>O, and stored at -20°C until use.

ScaI endonuclease was used to linearize the vector.

5 100 µg of DNA was digested with 10 µl of ScaI for 8 hours at 37°C following the manufacturer's protocol (GibcoBRL, Cat# 15436-017). 20 µg was used directly in stable transfection construction. 80 µg was cleaned of proteins and salts with a Phenol:Chloroform:IsoAmyl Alcohol 10 (25:24:1) extraction, 100% Ethanol precipitation (with 0.1 volume NH<sub>4</sub>COOH), and a 70% ethanol wash.

Stable transfectants of murine pre-B lymphoma cell line (L1.2) were prepared essentially as described (Ponath, P.D., et al., *J. Exp. Med.*, 183:2437-2448 (1996)).

15 25 million L1.2 cells in 0.8 ml of 1 x PBS were electroporated with 20 µg of linearized DNA, 20 µg linearized DNA that was then cleaned (see above under Linearization of DNA), or without DNA. Before electroporation, the L1.2 cells and the DNA were incubated 20 for 10 minutes in 50 ml conical tubes (Falcon Model 2070) with gentle mixing (swirling) every 2 minutes. The L1.2 cell-DNA mixture was transferred into Gene Pulser cuvettes (BioRad, Cat# 165-2088) with a 0.4 cm electrode gap. The mixture was then electroporated at 250V and 960 µF, with 25 the duration of shock and the actual voltage being measured. After electroporation, the cuvette was left undisturbed for 10 minutes at room temperature. All of the L1.2 cells-DNA mixture was then transferred to a T-25 flask (Costar), and grown for two days in 10 ml non-selective 30 medium.

#### Selection

L1.2 cells expressing CXCR3 were then subjected to selection for neomycin resistance. After two days of growth in non-selective medium, 10 ml of 1.6 g/L Geneticin

-82-

(GibcoBRL) was added for a final concentration of 0.8 g/L (The selective and maintenance concentration). This was then allowed to grow for 10 to 15 days, with fresh selective medium added when cells started to over-grow.

5 Fresh selective medium consisted of RPMI-1640 supplemented with 10% bovine serum and 800  $\mu$ g/ml G418.

The cell surface expression of CXCR3 was assessed by chemotaxis, and ligand binding and Scatchard analysis was also used to monitor surface expression. After G418  
10 selection, CXCR3 expressing L1.2 cells were selected based on chemotaxis ability. For each electroporation reaction, 30 ml (800,000 cells/ml) were collected, and suspended in 600  $\mu$ l selective medium. Selective medium, 600  $\mu$ l, containing 10 nM IP-10, was placed into the bottom chamber  
15 of BioCoat cell culture plates from Becton-Dickinson (Cat# 40575). 100  $\mu$ l/well of the L1.2 cells were added into the top chamber of the BioCoat plates. These cells were then left to chemotax overnight in a CO<sub>2</sub> incubator at 37°C. The top chambers with the non-chemotaxing cells were removed.  
20 The chemotaxed cells were collected, transferred into fresh medium, and allowed to grow in a 24 well plate. They were subsequently expanded into a T-25 and then a T-75 flask from Costar.

Transfectants expressing high level of receptors were  
25 cloned by limiting dilution. CXCR3 transfected cells were diluted to between 30-3 cells/ml in selection medium containing G418. Aliquots were added to 96-well tissue culture plates at 100  $\mu$ l/well. After 14 days at 37°C and 5% CO<sub>2</sub>, wells containing single colonies were identified  
30 under an inverted microscope. 50  $\mu$ l of the cells were then transferred and stained with anti-CXCR3 mAb as above and analyzed by flow cytometry. The level of receptor expression correlated with mean fluorescence intensity and high expressors were selected. Once a stable cell line was  
35 established, the line was expanded for use.

- 83 -

In addition to selection for antibiotic resistance and chemotaxis, CXCR3 transfected cells can be further selected by sorting for higher receptor expression by antibody staining, although this was not done here. For 5 staining, transfected cells can be resuspended at  $5 \times 10^6$ /ml in sterile PBS containing 1% bovine serum albumin. Isolated, sterile anti-CXCR3 mAb can be added to a final concentration of 3  $\mu$ g/ml and cells incubated on ice for 30 minutes. After washing with cold sterile PBS, the bound 10 mAb can be detected with FITC-conjugated anti-mouse IgG, sterilized by filtration through a 0.2  $\mu$ m filter. The cells can be washed again and sorted by flow cytometry. The top 5% positive cells can be collected and returned to tissue culture for expansion under selective conditions 15 (e.g., RPMI-1640 supplemented with 10% bovine serum and 800  $\mu$ g/ml G418).

Example 3. IP-10 Binds with High Affinity to a Receptor  
Expressed on L1.2 Cells Transfected with CXCR3 DNA and on  
Activated T Cells

20 Additional binding studies were performed using radiolabeled IP-10. Chemokine binding to target cells was carried out as described previously (Ponath, P.D., et al., J. Clin. Invest., 97:604-612 (1996); Van Riper, G., et al., J. Exp. Med., 177(3):851-856 (1993)). Cells were washed 25 once in PBS and resuspended in binding buffer (50 mM HEPES, pH 7.5, 1 mM CaCl<sub>2</sub>, 5 mM MgCl<sub>2</sub>, 0.5% BSA, and 0.05% azide) at a concentration of  $10^7$ /ml. Aliquots of 50  $\mu$ l ( $5 \times 10^5$  cells) were dispensed into microfuge tubes, followed by the addition of cold competitor (unlabeled IP-10) and 30 radiolabeled chemokine (0.05 nM <sup>125</sup>I-labeled IP-10). The final reaction volume was 200  $\mu$ l. Nonspecific binding was determined by incubating cells with radiolabeled chemokines in the presence of 250-500 nM of unlabeled chemokines. After a 60-minute incubation at room temperature, the cells

-84-

were washed three times with 1 ml of binding buffer containing 0.5 M NaCl. Cell pellets were then counted. The competition was presented as the percent specific binding as calculated by  $100 \times [(S-B)/(T-B)]$ , where S is 5 the radioactivity of the sample, B is background binding, and T is total binding without competitors. Background binding was obtained by incubating cells with radiolabeled chemokine and at least 400-fold excess of unlabeled chemokines. Duplicates were used throughout the 10 experiments and the standard deviations were always < 10% of the mean. All experiments were repeated at least three times. Curve fit and concentrations that inhibit 50% specific binding ( $IC_{50}$ ) were calculated by KaleidaGraph software (Synergy Software, Reading, PA).

15 Figures 5A-5B show that  $^{125}\text{I}$ -labeled IP-10 bound to L1.2 cells transfected with CXCR3 (Figure 5A) and to CD3-activated T cells (Figure 5B), and that this binding could be inhibited with increasing concentrations of cold IP-10. Scatchard analysis revealed that IP-10 bound to 20 L1.2 CXCR3 transfectants with a  $K_d$  of 614 pM, and that these transfectants expressed 37,000 receptors per cell (Figure 5A, inset). A similar analysis of anti-CD3 activated, IL-2 stimulated T cells revealed a  $K_d$  of 156 pM, and 17,000 receptors per cell (Figure 5B).  $^{125}\text{I}$ -labeled 25 IP-10 binding to activated T cells could be totally inhibited by cold Mig under the same conditions, although Mig was slightly less efficient at blocking  $^{125}\text{I}$ -labeled IP-10 binding than was cold IP-10 (not shown). IP-10 and Mig bind CXCR3 with high affinity ( $K_d$  ~150-600 pM).

30 Example 4. Production and Characterization of Monoclonal Antibodies (mAbs) Specific for CXCR3

To develop antagonists of CXCR3, and to study receptor expression and regulation, a panel of mAbs was produced by immunizing mice with a synthetic peptide corresponding to

- 85 -

the N-terminus of this receptor. These mAbs specifically recognized CXCR3 transfectants, but not a range of other receptor transfectants.

*mAb Production and Flow Cytometry*

5 mAbs reactive with CXCR3 were generated by immunizing Balb/C mice with 10 µg of 37-mer synthetic peptide corresponding to the first 37 N-terminal amino acids of CXCR3 (see also, Loetscher M., et al., *J. Exp. Med.*, 184:963-969 (1996)), five times over a period of 10 weeks.

10 This peptide was synthesized and coupled to purified protein derivative of tuberculin (Severn Biotech Ltd., Kidderminster, U.K.). The first immunization was intraperitoneal (IP) with Freund's Complete Adjuvant (FCA). The second, third and fourth immunizations were IP with

15 Freund's Incomplete Adjuvant (FIA), and the final immunization was with peptide conjugate alone (no adjuvant), and was administered intravenously (IV). Four days after the last immunization, the spleen was taken and cell fusion performed using the cell line SP2/0, as

20 described (Coligan, J.E., et al., *Current Protocols In Immunology* (John Wiley and Sons, New York), Unit 2.5.4 (1992)). mAbs were generated that reacted with the N-terminal 37-mer peptide as assessed by ELISA (Coligan, J.E., et al., *Current Protocols In Immunology* (John Wiley and Sons, New York), Unit 2.1.3 (1992)). mAbs reactive

25 with CXCR3 were identified using untransfected and CXCR3 transfected L1.2 cells or 300.19 cells (a murine B cell line, Loetscher, M. et al., *J. Exp. Med.*, 184: 963-969, (1996)), and immunofluorescent staining and analysis using

30 a FACScan® (Becton Dickinson & Co., Mountain View, CA).

Monoclonal antibodies specific for CXCR3 were generated. Eight mAbs were found to recognize surface expressed CXCR3, as judged by staining of CXCR3 transfected L1.2 cells, but not untransfected L1.2 cells

- 86 -

transfected with other receptor types. The FACS profile of one of these mAbs, 1C6, is shown in Figure 6. These mAbs also stained human T cells and T cell clones that had been activated *in vitro* with PHA or anti-CD3 (illustrated in 5 Figure 7C with 1C6). However, these anti-CXCR3 mAbs were unreactive with neutrophils (illustrated in Figure 7A with 1C6), monocytes, or eosinophils (not shown). This pattern of reactivity was consistent with the analysis by Northern blot. However, the phenotypic analysis unexpectedly 10 revealed the expression of CXCR3 on a large subset of circulating lymphocytes (Figure 7B). This expression pattern was observed in all individuals examined, indicating that CXCR3 is normally expressed on a subset of blood lymphocytes. CXCR3 was found to mark a population of 15 circulating T cells, which were contained within the CD45RO+ (memory) subset.

Murine hybridoma 1C6 (also referred to as LS77-1C6) was deposited on March 28, 1997 at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD, 20 20852, in accordance with the terms of the Budapest Treaty, under ATCC Accession Number HB-12330, on behalf of LeukoSite, Inc., 215 First Street, Cambridge, MA, 02142, U.S.A.

Example 5. CXCR3 is Expressed on Activated/memory T Cells

25 *Flow Cytometry*

mAbs to CXCR1, CXCR2, CXCR3, and CCR5 have been described (Qin, S., et al., *Eur. J. Immunol.*, 26:640-647 (1996); Heath, H., et al., *J. Clin. Invest.*, in press (1997)). Anti-CXCR4 mAb 12G5 (Endres, M.J., et al., *Cell*, 30 87:745-756 (1996)) was kindly provided by Jim Hoxie (Univ. Penn.). PE-conjugated mAbs to CD4, CD8, CD14, CD20, CD25, CD26, CD69, CD45RO, CD45RA, CD95, and anti-CD3 and anti-CD4 Cy-Chrome were supplied by PharMingen (La Jolla, CA).

To assess reactivity of mAbs against transfected cells or leukocytes, indirect immunofluorescence and flow cytometry were used. Cells were washed once with PBS, and resuspended in 100  $\mu$ l PBS containing 2% human serum and 5 0.1% sodium azide (staining buffer), 5  $\mu$ g/ml purified antibody, 5  $\mu$ g/ml IgG<sub>2a</sub> isotype matched control mAb (Sigma Chemical Co., St. Louis, MO) or 50  $\mu$ l hybridoma culture supernatant. After 20 min at 4°C, cells were washed twice with staining buffer, and resuspended in 50  $\mu$ l FITC-10 conjugated affinity purified F(ab')<sub>2</sub>, goat anti-mouse IgG (Jackson ImmunoResearch Laboratories). After incubating for 20 min at 4°C, cells were washed twice in staining buffer and analyzed on the FACScan® to determine the level of surface expression. Propidium iodide was used to 15 exclude dead cells.

A two color immunofluorescence analysis of lymphocytes showed that it was mostly CD3+ cells that expressed CXCR3, although a small proportion of CD20+ (B) cells and CD56+ (NK) cells also expressed this receptor (Figure 8). 20 A three color analysis of T cells, performed using anti-CD3 Cy-Chrome to label T cells, showed that a portion of the CD4+ cells and a portion of the CD8+ cells expressed CXCR3 (Figure 9). An analysis using markers of cellular activation, such as CD25 and CD69, revealed that activated 25 T cells generally expressed this receptor. CXCR3+ T cells were CD95+, CD45RO+, and CD45RA<sup>low</sup>, a phenotype consistent with previous activation. The expression of CXCR3 and CCR5, a chemokine receptor that is also biased in its expression to previously activated T cells (Wu, L., 30 personal communication), was also compared. Figure 9 shows that the CCR5+ cells in blood were contained within the CXCR3+ subset, and that CXCR3 was more widely expressed than CCR5. Unlike other T cell chemokine receptors, such as CCR5 or CXCR4, CXCR3 was expressed on the majority of 35 circulating, activated T cells.

-88-

Example 6. Anti-CXCR3 mAb Blocks IP-10 Binding and Chemotaxis

Chemotaxis of human leukocytes was assessed using a modification of a transendothelial assay (Carr, M.W., et 5 al., *Proc. Natl. Acad. Sci. USA*, 91(9):3652-3656 (1994)), which has been described previously (Ponath, P.D., et al., *J. Clin. Invest.*, 97:604-612 (1996)), using the ECV 304 endothelial cell line (European Collection of Animal Cell Cultures, Porton Down, Salisbury, U.K.). Cells that had 10 migrated to the bottom chamber were placed in a tube, and relative cell counts were obtained using the FACScan®.

The anti-peptide mAbs were also tested for their ability to inhibit the chemotaxis of CD3 activated T cell blasts. The results for mAbs designated 1A5, 1C6, 3A8, 15 5F10, 10C6 and 10G12 are illustrated in Figure 10. One mAb, 1C6, was superior to the other mAbs in its ability to block the chemotaxis of T cells to IP-10. mAb 1C6 was able to inhibit completely the chemotaxis of T cells to IP-10 in a dose-dependent manner, with an IC<sub>50</sub> of ~0.8 µg/ml (Figure 20 11). A concentration of 2-5 µg/ml was achieved 100% inhibition, using an optimal concentration of IP-10 (12.5 nM) in the bottom of the transwell. 1C6 was unable to significantly inhibit T cell chemotaxis to MCP-1 under the conditions used (Figure 10), which occurs through the 25 chemokine receptor CCR2b.

mAb 1C6 was also able to block completely <sup>125</sup>I-labeled IP-10 binding to activated T cells, with an IC<sub>50</sub> of 0.16 µg/ml (Figure 12). Between 1 and 10 µg/ml of antibody gave complete inhibition. The complete inhibition of both IP-10 30 binding and chemotaxis by mAb 1C6 indicates that activated T cells do not express another receptor that binds this chemokine.

Northern blot analysis indicated that CXCR3 was expressed in activated T cells. Using a panel of specific 35 mAbs, CXCR3 was found to be expressed on a subset of blood

- 89 -

T cells, as well as on other leukocyte types (B cells and NK cells). CXCR3 expressing T cells had a phenotype consistent with previous activation, i.e. CD45RO+, CD26+ (Example 5). Staining of T cells was markedly increased 5 when T cells were activated by CD3 and IL-2, which correlated with increased cell migration in response to IP-10 and radiolabeled ligand binding.

The poor responsiveness of blood T cells to IP-10 or Mig, at least in chemotaxis assays, appears anomalous. A 10 possible explanation is that factors other than receptor expression may determine cellular responsiveness to chemoattractants. Appropriate G protein coupling may be necessary for signalling. Another possibility is that the blood separation procedure disrupts this receptor, as has 15 been observed for IL-8 receptors on T cells and NK cells (C.R. Mackay). Injection of IP-10 into the skin of an appropriate experimental animal can address the significance of CXCR3 expression on blood T cells. On activated T cells, IP-10 is one of the most potent 20 chemoattractants. Activation of T cells may induce the receptor signalling molecules or coupling needed for signal transduction.

Example 7. mAb 1C6 Selectively Inhibits [Ca<sup>2+</sup>]<sub>i</sub> by T cells in Response to IP-10, but Not Mig

25 Intracellular calcium concentration ([Ca<sup>2+</sup>]<sub>i</sub>) was determined as follows. A stock solution of Fura-2 AM (Molecular Probes, Eugene, OR) was prepared by dissolving 50 µg of the dye in 44 µl of DMSO. Immediately prior to addition to cells, this stock was diluted 1:100 into HBSS 30 with Ca<sup>2+</sup> and Mg<sup>2+</sup> and 2% BSA. Fura-2 AM was added to cells at a final concentration of 0.2 moles/10<sup>6</sup> cells at 37°C for 30 minutes. Following labeling, excess dye was removed by centrifugation and cells were resuspended at a concentration of 10<sup>6</sup>/ml in 125 mM NaCl, 5 mM KCl, 1 mM

-90-

$MgCl_2$ , 1 mM  $CaCl_2$ , 0.5 mM glucose, 0.025% BSA and 20 mM HEPES, pH 7.4.  $[Ca^{2+}]_i$  was measured using excitation at 340 and 380 nm on a Hitachi F-2000 fluorescence spectrometer. Calibration was performed using 1% NP-40 for total release 5 and 25  $\mu M$  EGTA to chelate free  $Ca^{2+}$ .

IP-10 and Mig induce  $[Ca^{2+}]_i$  by human T cells, and each chemokine was able to completely desensitize responses to the other chemokine (Figures 13A-13B). A titration of the chemokines revealed that maximal  $[Ca^{2+}]_i$  was achieved with 10 as little as 2 nM IP-10 or 2 nM Mig. To examine the agonist/antagonist function of mAb 1C6, activated T cells were assessed for  $[Ca^{2+}]_i$  following injection of mAb 1C6, or an irrelevant isotype-matched control mAb. T cells injected with an irrelevant isotype-matched control mAb 15 showed a robust  $[Ca^{2+}]_i$  response to subsequent injection of IP-10 (not shown). However, T cells treated with mAb 1C6 showed no  $[Ca^{2+}]_i$  upon stimulation with 2 nM IP-10.  $[Ca^{2+}]_i$  in activated T cells in response to IP-10 was completely suppressed by 12.5 mg of 1C6. However, as much as 50  $\mu g/ml$  20 of 1C6 had no effect on the response of T cells to 2 nM Mig (Figure 13D), indicating that the two ligands were differentially affected by this mAb. As a control, mAb 1C6 was tested for its effects on the  $[Ca^{2+}]_i$  of T cells in response to MIP-1 $\alpha$  or RANTES, which occurs through 25 receptors other than CXCR3 (not shown). The mAb had no effect under the conditions used.

mAb 1C6, which inhibits IP-10 binding and IP-10-induced chemotaxis, also inhibited calcium flux by activated T cells, but did not inhibit Mig-induced calcium-flux under these conditions. These results suggest the 30 IP-10 and Mig bind and/or signal through different regions of CXCR3, and that mAb 1C6 was able to block the IP-10 binding site and subsequent signalling. Thus, it is possible to develop receptor antagonists which selectively

- 91 -

inhibit the effects of individual chemokines, illustrated here using mAb 1C6.

As discussed below, staining of CXCR3 transfectants can be inhibited by a peptide comprising the first 15 5 N-terminal amino acid residues of CXCR3. The inhibition of IP-10 binding and subsequent responses by mAb 1C6 suggests that this region is of particular functional importance for ligand binding. Based upon the inability of 1C6 to inhibit Mig-induced calcium flux under the conditions used, the 10 epitope recognized by mAb 1C6, which appears to be important for IP-10 binding and signalling, does not appear to be involved in Mig binding and/or signalling.

Example 8. Epitope Mapping

Peptides were ordered from Genemed, South San 15 Francisco, CA. The peptides, each 15 amino acids long, correspond to different portions of the first 45 N-terminal residues of CXCR3 protein:

P1: MVLEVSDHQVLNDAE (SEQ ID NO:2, residues 1-15)  
P2: VAALLENFSSSYDYG (SEQ ID NO:2, residues 16-30)  
20 P3: ENESDSCCTSPPCPQ (SEQ ID NO:2, residues 31-45)

The peptides were first dissolved in DMSO and diluted to 1 mg/ml in phosphate buffered saline (PBS). To test the ability of the peptides to block staining with 1C6 antibody, 1  $\mu$ g/ml of purified 1C6 antibody was incubated in 25 1X PBS, 5% Fetal Calf Serum (FCS) with CXCR3 L1.2 transfectants ( $10^5$  cells; see Materials and Methods for Examples 3-9) in the presence of 100  $\mu$ g/ml of each peptide for 30 minutes at 4°C. Final volume was 100  $\mu$ l. Positive staining was carried out in the absence of peptide. Bound 30 mAb was detected with anti-mouse IgG-FITC and the results were analyzed by flow cytometry.

- 92 -

Staining with mAb 1C6 was completely inhibited by P1, suggesting that the mAb recognizes an epitope in the first N-terminal 15 amino acids of the CXCR3 protein (Figures 14A-14D). Binding of another mAb designated 3A8 (also referred to as LS77-3A8), which was produced from the same fusion (LS-77) as mAb 1C6, was also inhibited by the P1 peptide.

Example 9. Production of Anti-CXCR3 Monoclonal Antibodies by Immunization with Transfected Cells

Additional anti-CXCR3 monoclonal antibodies were generated by immunization of mice with L1.2 cells transfected with a CXCR3 construct and expressing high levels of human CXCR3 (see Materials and Methods above). Immunization and generation of fusion hybridomas was performed as described (Qin, S. et al., *Eur. J. Immunol.*, 26: 640 (1996); and Heath, H. et al., *J. Clin. Invest.*, 99(2): 178 (1997)). Anti-CXCR3 mAbs were identified by positive staining of activated human T cells and CXCR3 transfectants. Eight anti-CXCR3 antibodies were obtained in one fusion (LS-104).

The ability of these antibodies to inhibit binding of IP-10 to CXCR3 was tested. Tissue culture supernatants (25  $\mu$ l) from the positive clones were incubated with CXCR3 L1.2 transfectant cells (Materials and Methods for Examples 3-9) and 0.05 nM of radiolabeled  $^{125}$ I-labeled IP-10 in 1X PBS, 5% Fetal Calf Serum (FCS) (100  $\mu$ l final volume) for 30 minutes at 4°C. Supernatants of mAb 1C6 and an anti-CXCR2 (IL-8 receptor B) mAb were also used as specific and non-specific mAb controls, respectively. Total binding was determined in the absence of antibodies. Background binding was obtained using 40 nM unlabeled IP-10 as the competitor and this value was used to calculate the percentage of inhibition by the mAbs. The results are shown in Figure 15. All antibodies from this fusion (LS-104) were able to

-93-

block IP-10 binding, with the percentage inhibition ranging from 50-70%.

These mAbs were also assessed for their ability to inhibit signalling (ability to induce  $Ca^{2+}$  flux)

5. essentially as described in Example 7, but none of the antibodies was able to block Mig-mediated  $Ca^{2+}$  signalling under the conditions used. Antibody inhibition of IP-10-mediated CXCR3 binding and signalling, but not Mig-mediated signalling via CXCR3, indicates that these antibodies can  
10 selectively inhibit CXCR3 functions mediated by IP-10.

Epitope mapping studies were also carried out using the antibodies from the LS-104 fusion essentially as described in Example 8. The results indicate that a variety of binding sites are recognized (Table 2). The  
15 results suggest that three of the antibodies recognize epitope(s) within the first 15 N-terminal amino acids of CXCR3 (residues 1-15 of SEQ ID NO:2), and two of the antibodies recognize epitope(s) within amino acids 16-30 of CXCR3 (residues 16-30 of SEQ ID NO:2). P3 peptide did not  
20 block staining with any of these antibodies, indicating that none of these antibodies bound the peptide representing amino acids 31-45 of CXCR3 (residues 31-45 of SEQ ID NO:2). Staining using the three remaining mAbs could not be significantly inhibited by any of the peptides  
25 under the conditions used, suggesting that these mAbs may bind epitopes comprising overlapping segments of the peptides, conformational epitopes displayed on the cell surface or epitopes on other parts of the receptor. These data also suggest that mAbs against various portions of  
30 CXCR3 can be obtained by immunizing mice with receptor transfectants.

-94-

Table 2  
Epitope mapping of anti-CXCR3 mAbs

	Fusion Number mAb Name	Peptides which inhibit mAb staining of CXCR3 transfectants	Binding Region <sup>3</sup>
5	LS-77 <sup>1</sup>		
	1C6	P1	AA 1-15
	3A8	P1	AA 1-15
10	LS-104 <sup>2</sup>		
	2F8	none	
	3A12	P1	AA 1-15
	3E2	P1	AA 1-15
	4B4	P2	AA 16-30
	4D2	none	
15	5B12	none	
	7B8	P2	AA 16-30
	8D5	P1	AA 1-15

<sup>1</sup> Antibodies obtained by peptide immunization

<sup>2</sup> Antibodies obtained by immunization with CXCR3 transfectants

<sup>3</sup> Amino acid residue position in SEQ ID NO:2

Example 10. Immunohistochemical Analysis of Normal and Inflamed Tissues Using mAb 1C6

25 *Tissues*

Human tissues (normal and inflamed) were obtained from the National Disease Research Institute, a service organization funded by the National Institutes of Health. Normal macaque (*Macaca mulatta*) tissues were obtained from the New England Regional Primate Research Center, Southboro, MA.

*Immunohistochemistry*

Alkaline Phosphatase Technique. Tissue was sectioned at a thickness of 4  $\mu$ m, desiccated, and then fixed in 2% paraformaldehyde/0.5X PBS for 10 minutes at 4°C. After PBS washing, nonspecific antibody binding sites were blocked with 10% normal goat serum/5% human AB serum/PBS for 30

-95-

minutes at room temperature. Next, the purified, anti-CXCR3 murine mAb 1C6, was diluted to a concentration of 10  $\mu$ g/ml in 0.3% Triton X 100/0.2% Tween 20/1% FCS/5% human AB serum, and 0.1% sodium azide, and applied to tissue  
5 sections which were incubated overnight at 4°C. An isotype-matched irrelevant monoclonal antibody was used as a negative control on step sections of tissues (IgG<sub>1</sub>, MOPC-21, Sigma, St. Louis, MO). Subsequently, biotinylated goat anti-mouse IgG (Vector, Burlingame, CA) and avidin-biotin-  
10 alkaline phosphatase complexes (Biogenex, San Ramon, CA) were added in sequence. Fast Red (Biogenex, San Ramon, CA), containing levamisole to block endogenous alkaline phosphatase activity, was used as the chromogen and Mayers hematoxylin as the counterstain.

15 **Results**

Human and macaque normal lymph node: In both species, staining was limited to 70-80% of lymphocytes within the paracortex and medullary cords, consistent with CXCR3 expression on T lymphocytes.

20 Human and macaque spleen: In both species, staining was limited to lymphocytes along the periphery of lymphoid follicles of white pulp and scattered lymphocytes within splenic sinusoids. This pattern is consistent with CXCR3 expression on T lymphocytes.

25 Human thymus: The thymic medulla contained scattered CXCR3 immunoreactive mononuclear cells morphologically consistent with lymphocytes.

30 This analysis revealed that macaque CXCR3 is recognized by mAb 1C6. Separate studies showed that CXCR3 is upregulated by culturing macaque cells T lymphocytes with concanavalin A and IL-2, macaque T cell blasts can

-96-

chemotax in response to human IP-10, mAb 1C6 can block this chemotaxis, and prior incubation with human IP-10 desensitizes macaque blasts as assessed by chemotaxis.

In the following discussion, variable numbers of CXCR3 5 immunoreactive "mononuclear cells" were identified in both normal and inflamed tissues. These mononuclear cells are most likely T lymphocytes for the following reasons: a) flow cytometry revealed that the majority of CXCR3<sup>+</sup> 10 cells are T lymphocytes, although CXCR3 has been detected on some B cells and NK cells, but not on other mononuclear cells; b) within the lymph node and spleen, lymphocytes in regions known to be populated by T cells are the only cells that are immunoreactive for CXCR3; and c) the CXCR3 15 immunoreactive mononuclear cells in the tissues listed below are morphologically consistent with lymphocytes.

Non-inflamed human tissues: In non-inflamed human tissues including heart, liver, kidney, lung, skin, breast, skeletal muscle, salivary gland, pancreas, vagina, uterus, and ovary, CXCR3 expression was limited to rare, scattered 20 interstitial mononuclear cells. In sections of small and large intestine, mononuclear cells expressing CXCR3 were observed within the lamina propria and Peyer's patches. In liver, periductal lymphocytes were also stained and hepatocytes were lightly stained.

25 Human brain: No staining was observed.

Inflamed human tissues: Several sections of chronically inflamed tissue characterized by interstitial and perivascular accumulation of mononuclear cells were examined for CXCR3 expression. In these tissues, including 30 tissue from human patients with interstitial nephritis (1 case, kidney tissue), ulcerative colitis (1 case, colon tissue), enteritis (1 case, small intestine) and chronic vaginitis (4 cases, vagina), about 50%-90% of mononuclear cells were immunoreactive for CXCR3. Thus, compared to

normal tissues, chronically inflamed tissues contained a greater number of interstitial mononuclear cells, and a greater percentage of these cells were immunoreactive for CXCR3.

5       Analysis of inflamed tissues revealed that about 50-90% of lymphocytes expressed CXCR3, whereas much lower percentages of lymphocytes in the corresponding normal tissue expressed CXCR3. These observations suggest specific recruitment of lymphocytes expressing CXCR3, most 10 likely T lymphocytes expressing CXCR3, to sites of chronic inflammation. CXCR3 appears to mark T cells with a predilection for homing or migration to inflammatory sites.

#### EQUIVALENTS

Those skilled in the art will be able to recognize, or 15 be able to ascertain, using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

20

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a 25 stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any 30 form of suggestion that that prior art forms part of the common general knowledge in Australia.



- 98 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Theodor-Kocher Institute
- (B) STREET: Freiestrasse 1
- (C) CITY: Bern
- (D) STATE/PROVINCE:
- (E) COUNTRY: Switzerland
- (F) POSTAL CODE/ZIP: CH-3012
- (G) TELEPHONE: 41-31-631-4141
- (I) TELEFAX: 41-31-631-3799

(i) APPLICANT:

- (A) NAME: LeukoSite, Inc.
- (B) STREET: 215 First Street
- (C) CITY: Cambridge
- (D) STATE/PROVINCE: MA
- (E) COUNTRY: USA
- (F) POSTAL CODE/ZIP: 02142
- (G) TELEPHONE: 617-621-9350
- (I) TELEFAX: 617-621-9349

(ii) TITLE OF INVENTION: CXCR3, ANTIBODIES, NUCLEIC ACIDS, AND METHODS OF USE

(iii) NUMBER OF SEQUENCES: 4

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: Hamilton, Brook, Smith & Reynolds, P.C.
- (B) STREET: Two Militia Drive
- (C) CITY: Lexington
- (D) STATE: MA
- (E) COUNTRY: USA
- (F) ZIP: 02173

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:
- (B) FILING DATE:
- (C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 08/829,839
- (B) FILING DATE: 31-MAR-1997

(viii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 08/709,838
- (B) FILING DATE: 10-SEP-1996

(ix) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Brook Esq., David E.
- (B) REGISTRATION NUMBER: 22,592
- (C) REFERENCE/DOCKET NUMBER: TKI96-01A2 PCT

- 99 -

(ix) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: (781) 861-6240  
(B) TELEFAX: (781) 861-9540

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1670 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 69..1172

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

- 100 -

GCC TGC ATC AGC TTT GAC CGC TAC CTG AAC ATA GTT CAT GCC ACC CAG Ala Cys Ile Ser Phe Asp Arg Tyr Leu Asn Ile Val His Ala Thr Gln 145 150 155	542
CTC TAC CGC CGG GGG CCC CCG GCC CGC GTG ACC CTC ACC TGC CTG GCT Leu Tyr Arg Arg Gly Pro Pro Ala Arg Val Thr Leu Thr Cys Leu Ala 160 165 170	590
GTC TGG GGG CTC TGC CTG CTT TTC GCC CTC CCA GAC TTC ATC TTC CTG Val Trp Gly Leu Cys Leu Leu Phe Ala Leu Pro Asp Phe Ile Phe Leu 175 180 185 190	638
TCG GCC CAC CAC GAC GAG CGC CTC AAC GCC ACC CAC TGC CAA TAC AAC Ser Ala His His Asp Glu Arg Leu Asn Ala Thr His Cys Gln Tyr Asn 195 200 205	686
TTC CCA CAG GTG GGC CGC ACG GCT CTG CGG GTG CTG CAG CTG GTG GCT Phe Pro Gln Val Gly Arg Thr Ala Leu Arg Val Leu Gln Leu Val Ala 210 215 220	734
GGC TTT CTG CTG CCC CTG CTG GTC ATG GCC TAC TGC TAT GCC CAC ATC Gly Phe Leu Leu Pro Leu Leu Val Met Ala Tyr Cys Tyr Ala His Ile 225 230 235	782
CTG GCC GTG CTG CTG GTT TCC AGG GGC CAG CGG CGC CTG CGG GCC ATG Leu Ala Val Leu Leu Val Ser Arg Gly Gln Arg Arg Leu Arg Ala Met 240 245 250	830
CGG CTG GTG GTG GTG GTC GTG GTG GCC TTT GCC CTC TGC TGG ACC CCC Arg Leu Val Val Val Val Val Ala Phe Ala Leu Cys Trp Thr Pro 255 260 265 270	878
TAT CAC CTG GTG GTG CTG GTG GAC ATC CTC ATG GAC CTG GGC GCT TTG Tyr His Leu Val Val Leu Val Asp Ile Leu Met Asp Leu Gly Ala Leu 275 280 285	926
GCC CGC AAC TGT GGC CGA GAA AGC AGG GTA GAC GTG GCC AAG TCG GTC Ala Arg Asn Cys Gly Arg Glu Ser Arg Val Asp Val Ala Lys Ser Val 290 295 300	974
ACC TCA GGC CTG GGC TAC ATG CAC TGC TGC CTC AAC CCG CTG CTC TAT Thr Ser Gly Leu Gly Tyr Met His Cys Cys Leu Asn Pro Leu Leu Tyr 305 310 315	1022
GCC TTT GTA GGG GTC AAG TTC CGG GAG CGG ATG TGG ATG CTG CTC TTG Ala Phe Val Gly Val Lys Phe Arg Glu Arg Met Trp Met Leu Leu Leu 320 325 330	1070
CGC CTG GGC TGC CCC AAC CAG AGA GGG CTC CAG AGG CAG CCA TCG TCT Arg Leu Gly Cys Pro Asn Gln Arg Gly Leu Gln Arg Gln Pro Ser Ser 335 340 345 350	1118
TCC CGC CGG GAT TCA TCC TGG TCT GAG ACC TCA GAG GCC TCC TAC TCG Ser Arg Arg Asp Ser Ser Trp Ser Glu Thr Ser Glu Ala Ser Tyr Ser 355 360 365	1166
GGC TTG TGAGGCCGGA ATCCGGGCTC CCCTTCGCC CACAGTCTGA CTTCCCCGCA Gly Leu	1222
TTCCAGGCTC CTCCCTCCCT CTGCCGGCTC TGGCTCTCCC CAATATCCTC GCTCCGGGA	1282

- 101 -

CTCACTGGCA GCCCCAGCAC CACCAAGGTCT CCCGGGAAGC CACCCCTCCCA GCTCTGAGGA	1342
CTGCACCATT GCTGCTCCTT AGCTGCCAAG CCCCATCCTG CCGCCCGAGG TGGCTGCCTG	1402
GAGCCCCACT GCCCTTCTCA TTTGGAAACT AAAACTTCAT CTICCCCAAG TGCGGGGAGT	1462
ACAAGGCATG GCGTAGAGGG TGCTGCCCA TGAAGCCACA GCCCAGGCCT CCAGCTCAGC	1522
AGTGAAGTGTG GCCATGGTCC CCAAGACCTC TATATTGCT CTTTTATTTT TATGTCTAAA	1582
ATCCTGCTTA AAACTTTCA ATAAAACAAGA TCGTCAGGAC CTTTTTTTTT TTTTTTTTT	1642
TTTTTTTTT TTTTTTTTTT TTTTTTTT	1670

**(2) INFORMATION FOR SEQ ID NO:2:**

**(i) SEQUENCE CHARACTERISTICS:**  
 (A) LENGTH: 368 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

**(ii) MOLECULE TYPE: protein**

**(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:**

Met Val Leu Glu Val Ser Asp His Gln Val Leu Asn Asp Ala Glu Val			
1	5	10	15
Ala Ala Leu Leu Glu Asn Phe Ser Ser Ser Tyr Asp Tyr Gly Glu Asn			
20	25	30	
Glu Ser Asp Ser Cys Cys Thr Ser Pro Pro Cys Pro Gln Asp Phe Ser			
35	40	45	
Leu Asn Phe Asp Arg Ala Phe Leu Pro Ala Leu Tyr Ser Leu Leu Phe			
50	55	60	
Leu Leu Gly Leu Leu Gly Asn Gly Ala Val Ala Ala Val Leu Leu Ser			
65	70	75	80
Arg Arg Thr Ala Leu Ser Ser Thr Asp Thr Phe Leu Leu His Leu Ala			
85	90	95	
Val Ala Asp Thr Leu Leu Val Leu Thr Leu Pro Leu Trp Ala Val Asp			
100	105	110	
Ala Ala Val Gln Trp Val Phe Gly Ser Gly Leu Cys Lys Val Ala Gly			
115	120	125	
Ala Leu Phe Asn Ile Asn Phe Tyr Ala Gly Ala Leu Leu Leu Ala Cys			
130	135	140	
Ile Ser Phe Asp Arg Tyr Leu Asn Ile Val His Ala Thr Gln Leu Tyr			
145	150	155	160
Arg Arg Gly Pro Pro Ala Arg Val Thr Leu Thr Cys Leu Ala Val Trp			
165	170	175	

-102-

Gly Leu Cys Leu Leu Phe Ala Leu Pro Asp Phe Ile Phe Leu Ser Ala  
180 185 190

His His Asp Glu Arg Leu Asn Ala Thr His Cys Gln Tyr Asn Phe Pro  
195 200 205

Gln Val Gly Arg Thr Ala Leu Arg Val Leu Gln Leu Val Ala Gly Phe  
210 215 220

Leu Leu Pro Leu Leu Val Met Ala Tyr Cys Tyr Ala His Ile Leu Ala  
225 230 235 240

Val Leu Leu Val Ser Arg Gly Gln Arg Arg Leu Arg Ala Met Arg Leu  
245 250 255

Val Val Val Val Val Ala Phe Ala Leu Cys Trp Thr Pro Tyr His  
260 265 270

Leu Val Val Leu Val Asp Ile Leu Met Asp Leu Gly Ala Leu Ala Arg  
275 280 285

Asn Cys Gly Arg Glu Ser Arg Val Asp Val Ala Lys Ser Val Thr Ser  
290 295 300

Gly Leu Gly Tyr Met His Cys Cys Leu Asn Pro Leu Leu Tyr Ala Phe  
305 310 315 320

Val Gly Val Lys Phe Arg Glu Arg Met Trp Met Leu Leu Leu Arg Leu  
325 330 335

Gly Cys Pro Asn Gln Arg Gly Leu Gln Arg Gln Pro Ser Ser Ser Arg  
340 345 350

Arg Asp Ser Ser Trp Ser Glu Thr Ser Glu Ala Ser Tyr Ser Gly Leu  
355 360 365

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ix) FEATURE:

- (A) NAME/KEY: modified\_base
- (B) LOCATION: 11
- (D) OTHER INFORMATION: /mod\_base= i

(ix) FEATURE:

- (A) NAME/KEY: modified\_base
- (B) LOCATION: 12
- (D) OTHER INFORMATION: /mod\_base= i

(ix) FEATURE:

- (A) NAME/KEY: modified\_base
- (B) LOCATION: 23
- (D) OTHER INFORMATION: /mod\_base= i

-103-

(ix) FEATURE:  
(A) NAME/KEY: modified\_base  
(B) LOCATION: 26  
(D) OTHER INFORMATION: /mod\_base= i

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGGCTGCAGC NNTKKCMGAC MTNCTNYT

28

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

(ix) FEATURE:  
(A) NAME/KEY: modified\_base  
(B) LOCATION: 10  
(D) OTHER INFORMATION: /mod\_base= i

(ix) FEATURE:  
(A) NAME/KEY: modified\_base  
(B) LOCATION: 16  
(D) OTHER INFORMATION: /mod\_base= i

(ix) FEATURE:  
(A) NAME/KEY: modified\_base  
(B) LOCATION: 18  
(D) OTHER INFORMATION: /mod\_base= i

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGGTCTAGAN GGGTTNANRC ARCWRYG

27

CLAIMS

We Claim:

1. A method of detecting or identifying an agent which binds a mammalian CXC Chemokine receptor 3 (CXCR3) protein or ligand binding variant thereof, comprising combining an agent to be tested with a composition comprising an 5 isolated and/or recombinant mammalian CXCR3 protein or ligand binding variant thereof under conditions suitable for binding of ligand thereto, and detecting or measuring the formation of a complex between said agent and said mammalian CXCR3 protein or variant, wherein said ligand binding variant comprises an amino acid sequence that has at least about 80% amino acid sequence identity with SEQ ID 10 NO:2.
2. The method of Claim 1 wherein said isolated and/or recombinant mammalian CXCR3 protein is human CXCR3 which has the amino acid sequence set forth in SEQ ID NO:2.
3. The method of Claim 1 wherein the amino acid sequence of said CXCR3 protein is 15 a sequence encoded by SEQ ID NO:1.



4. A method of detecting or identifying an agent which binds a mammalian CXC Chemokine Receptor 3 (CXCR3) protein or a ligand binding variant thereof comprising:

- a) combining an agent to be tested with a host cell expressing recombinant mammalian CXCR3 protein or a ligand binding variant thereof under conditions suitable for binding of ligand thereto; and
- b) detecting or measuring the formation of a complex between said agent and the mammalian CXCR3 protein or a ligand binding variant, wherein said ligand binding variant comprises an amino acid sequence that has at least 10 about 80% amino acid sequence identity with SEQ ID NO:2.

5

5. The method of Claim 4 wherein the host cell expresses a recombinant human CXCR3 protein having the amino acid sequence set forth in SEQ ID NO:2.

10

6. The method of Claim 4 wherein the host cell expresses a recombinant human CXCR3 protein, and the amino acid sequence of said human CXCR3 protein is a sequence encoded by SEQ ID NO:1.

15

7. The method of any one of Claims 1-6 wherein the agent is a ligand selected from the group consisting of human IP-10, human Mig, a mammalian homolog of IP-10, and a mammalian homolog of Mig.

20

8. The method of Claim 7, wherein the ligand is labeled with a label selected from the group consisting of a radioisotope, spin label, antigen label, enzyme label fluorescent group or chemiluminescent group.



-106-

9. The method of any one of Claims 1-6, wherein the method is a competitive binding assay, in which binding is determined in the presence of one or more ligands selected from the group consisting of human IP-10, human Mig, a mammalian homolog of IP-10, and a mammalian homolog of Mig.
- 5 10. The method of any one of Claims 1-6 wherein the method is a competitive binding assay, in which binding is determined in the presence of one or more ligands selected from the group consisting of human IP-10 and human Mig.
- 10 11. The method of Claim 4, 5 or 6 wherein the mammalian CXCR3 protein or ligand binding variant thereof can mediate cellular signalling and/or a cellular response, and the formation of a complex is monitored by detecting or measuring a signalling activity or cellular response of said CXCR3 protein or variant in response thereto.
- 15 12. A method of detecting or identifying an inhibitor of ligand binding to a mammalian CXC Chemokine Receptor 3 (CXCR3) protein or a ligand binding variant thereof comprising:
  - a) combining an agent to be tested with a ligand of said mammalian CXCR3 protein and a composition comprising isolated and/or recombinant mammalian CXCR3 protein or ligand binding variant thereof under conditions suitable for binding of ligand thereto; and
  - b) detecting or measuring the formation of a complex between said mammalian CXCR3 protein or variant and said ligand, whereby inhibition of complex formation by the agent is indicative that the agent is an inhibitor, wherein said ligand binding variant comprises an amino acid sequence that has at least about 80% amino acid sequence identity with
- 20 25 SEQ ID NO:2.



13. The method of Claim 12 wherein an agent to be tested is combined with a ligand and a composition comprising an isolated and/or recombinant human CXCR3 protein which has the amino acid sequence set forth in SEQ iD NO:2.

14. The method of Claim 12 wherein an agent to be tested is combined with a ligand and a composition comprising an isolated and/or recombinant human CXCR3 protein, and wherein the amino acid sequence of said human CXCR3 protein is a sequence encoded by SEQ ID NO:1.

15. The method of Claim 12, 13 or 14 wherein the composition comprising isolated and/or recombinant CXCR3 protein or a ligand binding variant thereof contains a host cell expressing said recombinant CXCR3 protein or ligand binding variant thereof.

16. The method of Claim 15 wherein the mammalian CXCR3 protein or ligand binding variant thereof can mediate cellular signalling and/or a cellular response, and the formation of a complex is monitored by detecting or measuring a signalling activity or cellular response of said CXCR3 protein or variant in response thereto.

17. A method of detecting or identifying an inhibitor of ligand binding to a human CXCR3 Chemokine Receptor 3 (CXCR3) protein comprising:

20 a) combining an agent to be tested with a ligand of said human CXCR3 protein and a host cell expressing a recombinant protein comprising SEQ ID NO:2 under conditions suitable for binding of ligand thereto; and  
b) detecting or measuring the formation of a complex between said protein and said ligand, whereby inhibition of complex formation by the agent is indicative that the agent is an inhibitor.

18. The method of Claim 17 wherein the agent is an antibody or antibody fragment.



19. A method of detecting or identifying an inhibitor of a mammalian CXCR3 protein or functional variant thereof comprising combining an agent to be tested with
  - 5 a) a host cell expressing a recombinant mammalian CXCR3 protein or functional variant thereof, and
  - b) a ligand or promoter thereof, under conditions suitable for detecting a ligand- or promoter-induced response, and assessing the ability of the test agent to inhibit said response, whereby inhibition of a ligand- or promoter-induced response by the agent is indicative that the agent is an inhibitor,
- 10 wherein said functional variant can selectively bind at least one chemokine selected from the group consisting of IP-10, Mig, a homolog of IP-10, and a homolog of Mig, and wherein said functional variant comprises an amino acid sequence that has at least about 80% amino acid sequence identity with SEQ ID NO:2.
- 15
20. The method of Claim 19 wherein said recombinant mammalian CXCR3 protein is recombinant human CXCR3 having the amino acid sequence set forth in SEQ ID NO:2.
21. The method of Claim 19 wherein the amino acid sequence of said CXCR3 protein is a sequence encoded by SEQ ID NO:1.
- 20
22. The method of Claim 19, 20 or 21 wherein the response is monitored by detecting or measuring a signalling activity or cellular response of said CXCR3 protein or variant thereof in response thereto.



-109-

23. The method of any one of Claims 12-22 wherein the ligand is selected from the group consisting of human IP-10, human Mig, a mammalian homolog of IP-10, and a mammalian homolog of Mig.

5 24. The method of Claim 23, wherein the ligand is labeled with a label selected from the group consisting of a radioisotope, spin label, antigen label, enzyme label, fluorescent group or chemiluminescent group.

25. The method of any one of Claims 12-22 wherein the ligand is human IP-10 and/or human Mig.

10 26. A method of detecting or identifying a promoter of a mammalian CXCR3 protein or functional variant thereof comprising combining an agent to be tested with a host cell expressing a recombinant mammalian CXCR3 protein or functional variant thereof under conditions suitable for detecting a receptor-mediated response, and detecting or measuring said response, whereby induction or stimulation of said response by the agent is indicative that the agent is a promoter, wherein said functional variant can selectively bind at least one chemokine selected from the group consisting of IP-10, Mig, a homolog of IP-10, and a homolog of Mig, and wherein said functional variant comprises an amino acid sequence that has at least about 80% amino acid sequence identity with SEQ ID NO:2.

15 27. The method of Claim 26 wherein said recombinant mammalian CXCR3 protein is recombinant human CXCR3 which has the amino acid sequence set forth in SEQ ID NO:2.

20 28. The method of Claim 26 wherein the amino acid sequence of said CXCR3 protein is a sequence encoded by SEQ ID NO:1.



29. A method of detecting or identifying an agent which binds a mammalian CXC  
Chemokine receptor 3 (CXCR3) protein, comprising combining an agent to be  
tested with a composition comprising an isolated and/or recombinant fusion protein  
comprising a human CXCR3 protein under conditions suitable for binding of ligand  
thereto, and detecting or measuring the formation of a complex between said agent  
and said fusion protein, wherein said fusion protein can selectively bind at least one  
chemokine selected from the group consisting of human IP-10 and human Mig.

30. The method of Claim 29 wherein said human CXCR3 protein has the amino acid  
sequence set forth in SEQ ID NO:2.

10 31. The method of Claim 29 wherein the amino acid sequence of said human CXCR3  
protein is a sequence encoded by SEQ ID NO:1.

32. A method of detecting or identifying an agent which binds a mammalian CXC  
Chemokine Receptor 3 (CXCR3) protein comprising:

- 15 a) combining an agent to be tested with a host cell expressing a recombinant  
fusion protein comprising human CXCR3 protein under conditions suitable  
for binding of ligand thereto; and
- b) detecting or measuring the formation of a complex between said agent and  
said fusion protein.

20 33. The method of Claim 32 wherein said human CXCR3 protein has the amino acid  
sequence set forth in SEQ ID NO:2.

34. The method of Claim 32 wherein the amino acid sequence of said human CXCR3  
protein is a sequence encoded by SEQ ID NO:1.



-111-

35. The method of any one of Claims 29-34 wherein the agent is a ligand selected from the group consisting of human IP-10, human Mig, a mammalian homolog of IP-10, and a mammalian homolog of Mig.

5 36. The method of Claim 35, wherein the ligand is labeled with a label selected from the group consisting of a radioisotope, spin label, antigen label, enzyme label fluorescent group or chemiluminescent group.

10 37. The method of any one of Claims 29-34 wherein the method is a competitive binding assay, in which binding is determined in the presence of one or more ligands selected from the group consisting of human IP-10, human Mig, a mammalian homolog of IP-10, and a mammalian homolog of Mig.

38. The method of any one of Claims 29-34 wherein the method is a competitive binding assay, in which binding is determined in the presence of one or more ligands selected from the group consisting of human IP-10 and human Mig.

15 39. The method of Claim 32, 33 or 34 wherein the mammalian CXCR3 protein can mediate cellular signalling and/or a cellular response, and the formation of a complex is monitored by detecting or measuring a signalling activity or cellular response of said CXCR3 protein in response thereto.



40. A method of detecting or identifying an inhibitor of ligand binding to human CXC Chemokine Receptor 3 (CXCR3) protein comprising:

- a) combining an agent to be tested with a ligand of said human CXCR3 protein and a composition comprising isolated and/or recombinant fusion protein comprising human CXCR3 under conditions suitable for binding of ligand thereto; and
- b) detecting or measuring the formation of a complex between said fusion protein and said ligand, whereby inhibition of complex formation by the agent is indicative that the agent is an inhibitor.

10 41. The method of Claim 40 wherein said human CXCR3 protein has the amino acid sequence set forth in SEQ ID NO:2.

42. The method of Claim 40 wherein the amino acid sequence of said human CXCR3 protein is a sequence encoded by SEQ ID NO:1.

15 43. A method of detecting or identifying an inhibitor of ligand binding to a human CXCR3 Chemokine Receptor 3 (CXCR3) protein comprising:

- a) combining an agent to be tested with a ligand of said human CXCR3 protein and a host cell expressing a fusion protein comprising a human CXCR3 protein under conditions suitable for binding of ligand thereto; and
- b) detecting or measuring the formation of a complex between said CXCR3 protein and said ligand, whereby inhibition of complex formation by the agent is indicative that the agent is an inhibitor.

20 44. The method of Claim 43 wherein said human CXCR3 protein has the amino acid sequence set forth in SEQ ID NO:2.



-113-

45. The method of Claim 43 wherein the amino acid sequence of said human CXCR3 protein is a sequence encoded by SEQ ID NO:1.
46. The method of Claim 43, 44 or 45 wherein the agent is an antibody or antibody fragment.
- 5 47. A method of detecting or identifying an inhibitor of a human CXCR3 protein comprising combining an agent to be tested with
  - a) a host cell expressing a fusion protein comprising human CXCR3 protein, and
  - b) a ligand or promoter thereof, under conditions suitable for detecting a ligand- or promoter-induced response, and assessing the ability of the test agent to inhibit said response, whereby inhibition of a ligand- or promoter-induced response by the agent is indicative that the agent is an inhibitor,  
wherein said fusion protein can selectively bind at least one chemokine selected from the group consisting of IP-10, Mig, a homolog of IP-10, and a homolog of Mig.
- 10 48. The method of Claim 47 wherein said human CXCR3 protein has the amino acid sequence set forth in SEQ ID NO:2.
- 15 49. The method of Claim 47 wherein the amino acid sequence of said human CXCR3 protein is a sequence encoded by SEQ ID NO:1.
- 20 50. The method of Claim 47, 48 or 49 wherein the response is monitored by detecting or measuring a signalling activity or cellular response of said protein in response thereto.



51. The method of any one of Claims 40-50 wherein the ligand is selected from the group consisting of human IP-10, human Mig, a mammalian homolog of IP-10, and a mammalian homolog of Mig.
52. The method of Claim 51 wherein the ligand is labeled with a label selected from the group consisting of a radioisotope, spin label, antigen label, enzyme label flourescent group or chemiluminescent group.
53. The method of any one of Claims 40-50 wherein the ligand is human IP-10 and/or human Mig.
- 10 54. The method of Claim 47, 48 or 49 wherein the human CXCR3 protein can mediate cellular signalling and/or a cellular response, and the formation of a complex is monitored by detecting or measuring a signalling activity or cellular response of said protein in response thereto.
- 15 55. A method of detecting or identifying a promoter of a mammalian CXCR3 protein comprising combining an agent to be tested with a host cell expressing a fusion protein comprising human CXCR3 protein under conditions suitable for detecting a receptor-mediated response, and detecting or measuring said response, whereby induction or stimulation of said response by the agent is indicative that the agent is a promoter, wherein said fusion protein can selectively bind at least one chemokine selected from the group consisting of IP-10, Mig, a homolog of IP-10, and a homolog of Mig.
- 20 56. The method of Claim 55 wherein an agent to be tested is combined with a host cell expressing a fusion protein comprising SEQ ID NO:2 under conditions suitable for detecting a receptor-mediated response



-115-

57. The method of Claim 55 wherein an agent to be tested is combined with a host cell expressing a fusion protein comprising human CXCR3 protein under conditions suitable for detecting a receptor-mediated response, and wherein the amino acid sequence of said human CXCR3 protein is a sequence encoded by SEQ ID NO:1.

58. An inhibitor of at least one function characteristic of a mammalian CXCR3 protein identified according to the method of Claim 15 or Claim 40, wherein said inhibitor is an antagonist.

59. An inhibitor of at least one function characteristic of a mammalian CXCR3 protein identified according to the method of Claim 12, 17 or 19, wherein said inhibitor is an antagonist.

10 60. An inhibitor of at least one function characteristic of a mammalian CXCR3 protein identified according to the method of Claim 40, 43 or 47, wherein said inhibitor is an antagonist.

15 61. A method of modulating at least one function of a mammalian CXC Chemokine Receptor 3 (CXCR3) protein, comprising the step of contacting said protein with an inhibitor or promoter of at least one function of said protein.

62. A method of modulating at least one function of a mammalian CXC Chemokine Receptor 3 (CXCR3) protein having the amino acid sequence set forth in SEQ ID 20 NO:2, comprising the step of contacting said protein with an inhibitor or promoter of at least one function of said protein.



63. A method of modulating at least one function of a mammalian CXC Chemokine Receptor 3 (CXCR3) protein comprising the step of contacting said protein with an inhibitor or promoter of at least one function of said protein, wherein the amino acid sequence of said CXCR3 protein is a sequence encoded by SEQ ID NO:1.

5 64. A method for treating an inflammatory disease or condition, comprising administering to a mammal a therapeutically effective amount of an inhibitor of a mammalian CXC Chemokine Receptor 3 (CXCR3) protein, whereby inflammation is reduced.

10 65. The method of Claim 64, wherein the inflammatory disease or condition is a T cell mediated disease or condition.

15 66. The method of Claim 64, wherein the inhibitor is an antibody or antigen binding fragment thereof which specifically binds a human CXC Chemokine Receptor 3 (CXCR3) protein and which can inhibit one or more functions of said human CXCR3 protein selected from the group consisting of binding of a ligand, signalling activity and cellular response function.

67. The method of Claim 66, wherein the antibody or fragment thereof can compete with monoclonal antibody 1C6 for binding to a human CXCR3 protein.

68. The method of Claim 67, wherein the antibody or fragment thereof is monoclonal antibody 1C6 or an antigen binding fragment thereof.

20 69. A method of antitumor therapy, comprising administering to a mammal a therapeutically effective amount of a promoter of a mammalian CXC Chemokine Receptor 3 (CXCR3) protein, other than a natural ligand of said protein.



70. A method of antiviral therapy, comprising administering to a mammal a therapeutically effective amount of a promoter of a mammalian CXC Chemokine Receptor 3 (CXCR3) protein, other than a natural ligand of said protein.

71. An antibody or antigen-binding fragment thereof which binds a human CXC Chemokine Receptor 3 (CXCR3) protein having the amino acid sequence of SEQ ID NO:2.

72. The antibody or antigen-binding fragment thereof of Claim 71, wherein said antibody or fragment can inhibit the interaction of a human CXCR3 protein with one or more ligands selected from the group consisting of IP-10 and Mig.

73. The antibody or antigen-binding fragment of Claim 72 wherein said ligand is human IP-10.

74. The antibody or antigen-binding fragment of Claim 72 wherein said ligand is human Mig.

75. The antibody or antigen-binding fragment thereof of Claim 72, wherein said antibody or fragment can compete with monoclonal antibody 1C6 for binding to a human CXCR3 protein.

76. An antibody or antigen-binding fragment thereof of Claim 71, wherein said binding can be inhibited by a portion of a human CXCR3 corresponding to the N-terminal extracellular segment of SEQ ID NO:2 or a portion thereof having at least one immunological property of a human CXCR3 protein.



77. An antibody or antigen-binding fragment thereof of Claim 71, wherein said binding can be inhibited by a polypeptide having a sequence which is the same as that of residues 1-15 of SEQ ID NO:2.

78. The antibody or antigen-binding fragment thereof of Claim 77, wherein said binding 5 is not inhibited by a polypeptide having a sequence which is the same as that of residues 16-30 of SEQ ID NO:2 or by a polypeptide having a sequence which is the same as that of residues 31-45 of SEQ ID NO:2.

79. An antibody or antigen-binding fragment thereof of Claim 71, wherein said binding can be inhibited by a polypeptide having a sequence which is the same as that of 10 residues 16-30 of SEQ ID NO:2.

80. An antibody or antigen-binding fragment thereof which binds a human CXC Chemokine Receptor 3 (CXCR3) protein having the amino acid sequence of SEQ ID NO:2, wherein said antibody is a chimeric or humanized antibody having the same or similar epitopic specificity as monoclonal antibody 1C6.

15 81. The antibody or antigen-binding fragment of Claim 71 wherein said antibody or fragment inhibits T cell activation.

82. An antibody or antigen-binding fragment thereof which binds a human CXC Chemokine Receptor 3 (CXCR3) protein having the amino acid sequence of SEQ ID NO:2, wherein said antibody is a humanized immunoglobulin comprising the 20 light chain CDRs (CDR1, CDR2 and CDR3) and the heavy chain CDRs (CDR1, CDR2 and CDR3) of monoclonal antibody 1C6 and a human framework region, and said humanized immunoglobulin has the same or similar epitopic specificity as monoclonal antibody 1C6.



83. An antibody or antigen-binding fragment thereof which binds a human CXC Chemokine Receptor 3 (CXCR3) protein having the amino acid sequence of SEQ ID NO:2, wherein said antibody or fragment can inhibit one or more functions of said human CXCR3 protein selected from the group consisting of binding of a ligand, signalling activity and cellular response function.

84. The antibody or antigen-binding fragment thereof of Claim 83, wherein the antibody or fragment inhibits binding of a ligand to said human CXCR3 protein.

85. The antibody or antigen-binding fragment of Claim 84 wherein said ligand is human Mig.

10 86. The antibody of antigen-binding fragment of Claim 84 wherein said ligand is human IP-10.

87. An antibody or antigen-binding fragment thereof which binds a human CXC Chemokine Receptor 3 (CXCR3) protein having the amino acid sequence of SEQ ID NO:2, wherein said antibody or fragment selectively inhibits the interaction of a 15 human CXCR3 protein with IP-10.

88. The antibody or antigen-binding fragment thereof of Claim 87, wherein said antibody or fragment inhibits a signalling activity or a cellular response function mediated by IP-10 interaction with said receptor.

89. Monoclonal antibody 1C6 or an antigen-binding fragment thereof which binds a 20 human CXC Chemokine Receptor 3 (CXCR3) protein having the amino acid sequence of SEQ ID NO:2.



90. An antibody or antigen-binding fragment thereof, wherein said antibody or fragment can inhibit binding of human IP-10 to a human CXC Chemokine Receptor 3 (CXCR3) protein and can inhibit one or more functions mediated by said CXCR3

5 protein in response to IP-10 binding selected from the group consisting of signalling activity and cellular response function, wherein said antibody or fragment can bind a human CXCR3 protein having the amino acid sequence of SEQ ID NO:2.

91. An antibody or antigen-binding fragment of Claim 90, wherein said antibody or fragment can inhibit one or more functions of said human CXC Chemokine

10 Receptor 3 (CXCR3) protein selected from the group consisting of IP-10-induced calcium flux or IP-10-induced chemotaxis.


92. Hybridoma cell line HB-12330.

93. An antibody produced by hybridoma cell line HB-12330 or an antigen-binding fragment thereof.

94. An antibody or antigen-binding fragment thereof which binds a human CXC

15 Chemokine Receptor 3 (CXCR3) protein having the amino acid sequence of SEQ ID NO:2, wherein said antibody or fragment can inhibit one or more functions of said human CXCR3 protein selected from the group consisting of binding of a ligand, signalling activity and cellular response function, and wherein the binding of said antibody or antigen-binding fragment to said human CXCR3 can be inhibited 20 by a polypeptide having a sequence which is the same as that of residues 1-15 of SEQ ID NO:2.

95. The antibody or antigen-binding fragment of Claim 94 wherein said ligand is human IP-10.



96. The antibody or antigen-binding fragment of Claim 94 wherein said ligand is human  
Mig.

97. An antibody or antigen-binding fragment thereof which binds a human CXC  
Chemokine Receptor 3 (CXCR3) protein having the amino acid sequence of SEQ  
5 ID NO:2, wherein said antibody or fragment can inhibit one or more functions of  
said human CXCR3 protein selected from the group consisting of binding of a  
ligand, signalling activity and cellular response function, and wherein the binding of  
said antibody or antigen-binding fragment to said human CXCR3 can be inhibited  
by a polypeptide having a sequence which is the same as that of residues 16-30 of  
10 SEQ ID NO:2.

98. The antibody or antigen-binding fragment of Claim 97 wherein said ligand is human  
IP-10.

99. The antibody or antigen-binding fragment of Claim 97 wherein said ligand is human  
Mig.

15 100. The antibody or antigen-binding fragment of Claim 71, 72, 83 or 90 wherein said  
antigen-binding fragment is selected from the group consisting of a Fab  
fragment, a Fab' fragment, a  $F(ab')_2$  fragment and a Fv fragment.


101. The antibody or antigen-binding fragment of Claim 71, 72, 83 or 90 wherein  
said antibody is a human, humanized or chimeric antibody.

20 102. A composition comprising the antibody or antigen-binding fragment of Claim  
71, 72, 83 or 90 and a physiologically acceptable vehicle or carrier.



-122-

103. The antibody or antigen-binding fragment of Claim 77, 82, 89, 93, 94 or 97 wherein said antigen-binding fragment is selected from the group consisting of a Fab fragment, a Fab' fragment, a F(ab')<sub>2</sub> fragment and a Fv fragment.
104. The antibody or antigen-binding fragment of Claim 75, 77, 94 or 97 wherein said antibody is a human, humanized or chimeric antibody.
105. A composition comprising the antibody or antigen-binding fragment of Claim 77, 82, 89, 93, 94 or 97 and a physiologically acceptable vehicle or carrier.
106. A method of detecting a human CXC Chemokine Receptor 3 (CXCR3) protein in a sample comprising:
  - 10 a) contacting a sample with an antibody or antigen-binding fragment thereof, wherein said antibody or fragment binds a human CXCR3 protein having the sequence of SEQ ID NO:2, under conditions suitable for specific binding of said antibody or fragment to said protein having the sequence of SEQ ID NO:2; and
  - 15 b) detecting antibody-CXCR3 or antibody fragment-CXCR3 complexes, wherein detection of complex is indicative of the presence of a human CXCR3 in said sample.
107. The method of detecting a human CXCR3 protein of Claim 106 wherein said antibody or fragment comprises a detectable label.
- 20 108. The method of detecting a human CXCR3 protein of Claim 106 wherein the sample is contacted with an antibody which binds said protein.



-123-

109. The method of detecting a human CXCR3 protein of Claim 106, wherein the sample is contacted with an antigen-binding fragment of an antibody which binds said protein.
110. The method of Claim 109, wherein said antigen-binding fragment is selected 5 from the group consisting of a Fab fragment, a Fab' fragment, a F(ab')<sub>2</sub> fragment and a Fv fragment.
111. A method of detecting a human CXC Chemokine Receptor 3 (CXCR3) protein in a sample comprising:
  - 10 a) contacting a sample with an antibody or antigen-binding fragment thereof, wherein said antibody or fragment binds a human CXCR3 protein having the sequence of SEQ ID NO:2 and can inhibit one or more functions of said human CXCR3 protein selected from the group consisting of binding of a ligand, signalling activity and cellular response function, under conditions suitable for specific binding of said antibody or fragment to said protein having the sequence of SEQ ID NO:2; and
  - 15 b) detecting antibody-CXCR3 or antibody fragment-CXCR3 complexes, wherein detection of complex is indicative of the presence of a human CXCR3 in said sample.
112. The method of detecting a human CXCR3 protein of Claim 111, wherein said 20 antibody or antigen-binding fragment inhibits binding of a ligand to said human CXCR3 protein.
113. The method of detecting a human CXCR3 protein of Claim 112, wherein said ligand is selected from the group consisting of IP-10 and Mig.



114. The method of detecting a human CXCR3 protein of Claim 113, wherein said ligand is human IP-10.

115. The method of detecting a human CXCR3 protein of Claim 113, wherein said ligand is human Mig.

5 116. The method of detecting a human CXCR3 protein of Claim 111, wherein said antibody or fragment comprises a detectable label.

117. The method of detecting a human CXCR3 protein of Claim 111, wherein said antigen-binding fragment is selected from the group consisting of a Fab fragment, a Fab' fragment, a F(ab')<sub>2</sub> fragment and a Fv fragment.

10 118. An isolated human CXC Chemokine Receptor 3 (CXCR3) protein or functional variant thereof, wherein said protein or variant can selectively bind one or more chemokines and can mediate cellular signalling and/or a cellular response in response thereto, and wherein said protein or variant is encoded by a nucleic acid which hybridizes under high stringency conditions to a second nucleic acid and the sequence of said second nucleic acid is selected from the group consisting of Figure 1 (SEQ ID NO:1) and a sequence complementary to Figure 1 (SEQ ID NO:1).

15 119. The isolated human CXCR3 protein or functional variant thereof of Claim 118, wherein the protein can selectively bind one or more chemokines selected from the group consisting of human IP-10 and human Mig.

20 120. An isolated human CXC Chemokine Receptor 3 (CXCR3) protein encoded by the nucleic acid illustrated in Figure 1 (SEQ ID NO:1).



121. An isolated human CXC Chemokine Receptor 3 (CXCR3) protein comprising an amino acid sequence as set forth in Figure 2 (SEQ ID NO:2).

122. A fusion protein comprising a human CXC Chemokine Receptor 3 (CXCR3) protein, wherein the amino acid sequence of said CXCR3 protein is a sequence encoded by the nucleic acid illustrated in Figure 1 (SEQ ID NO:1).

5

123. A fusion protein comprising a human CXC Chemokine Receptor 3 (CXCR3) protein wherein the amino acid sequence of said CXCR3 protein consists of the amino acid sequence of Figure 2 (SEQ ID NO:2).

124. A fusion protein comprising a human CXC Chemokine Receptor 3 (CXCR3) protein or functional variant thereof, wherein said CXCR3 protein or variant can selectively bind one or more chemokines and can mediate cellular signalling and/or a cellular response in response thereto, and wherein said CXCR3 protein or variant is encoded by a nucleic acid which hybridizes under high stringency conditions to a second nucleic acid and the sequence of said second nucleic acid is selected from the group consisting of Figure 1 (SEQ ID NO:1) and a sequence complementary to Figure 1 (SEQ ID NO:1).

10

15

125. The fusion protein of Claim 124, wherein said CXCR3 protein or variant can selectively bind one or more chemokines selected from the group consisting of human IP-10 and human Mig.



126. An isolated human CXC Chemokine Receptor 3 (CXCR3) protein or variant thereof, wherein the amino acid sequence of said CXCR3 protein or variant is at least about 80% identical to that of the protein shown in Figure 2 (SEQ ID NO:2), said protein or variant comprises the extracellular N-terminal segment of the protein shown in Figure 2 (SEQ ID NO:2), and said protein or variant can selectively bind one or more chemokines selected from the group consisting of IP-10 and Mig and mediate cellular signalling and/or a cellular response in response thereto.

5

127. A fusion protein comprising a human CXC Chemokine Receptor 3 (CXCR3) protein or variant thereof, wherein the amino acid sequence of said CXCR3 protein or variant is at least about 80% identical to that of the protein shown in Figure 2 (SEQ ID NO:2), said CXCR3 protein or variant comprises the extracellular N-terminal segment of the protein shown in Figure 2 (SEQ ID NO:2), and said CXCR3 protein or variant can selectively bind one or more chemokines selected from the group consisting of IP-10 and Mig and mediate cellular signalling and/or a cellular response in response thereto.

10

15

128. The isolated human CXCR3 protein or functional variant thereof of Claim 118, 119, 120, 121 or 126 wherein said protein is labeled with a detectable label.

129. The isolated human CXCR3 protein or functional variant thereof of Claim 128, wherein the label is a radioisotope, a spin label, an enzyme label, a fluorescent label, a chemiluminescent label, an antigen or epitope label.

20

130. The fusion protein of Claim 122, 123, 124, 125 or 127 wherein said fusion protein is labeled with a detectable label.



131. The fusion protein of Claim 130, wherein the label is a radioisotope, a spin label, an enzyme label, a fluorescent label, a chemiluminescent label, an antigen or epitope label.

132. An isolated nucleic acid encoding a human CXC Chemokine Receptor 3 (CXCR3) protein or functional variant thereof, wherein said protein or variant can selectively bind one or more chemokines and can mediate cellular signalling and/or a cellular response in response thereto, and wherein said nucleic acid hybridizes under high stringency conditions to a second nucleic acid and the sequence of said second nucleic acid is selected from the group consisting of Figure 1 (SEQ ID NO:1) and a sequence complementary to Figure 1 (SEQ ID NO:1).

133. The isolated nucleic acid of Claim 132, wherein the isolated nucleic acid is essentially pure.

134. The isolated nucleic acid of Claim 132, wherein the human CXCR3 protein or variant can selectively bind one or more chemokines selected from the group consisting of human IP-10 and human Mig.

135. The isolated nucleic acid of Claim 132 comprising SEQ ID NO:1, its complement, or a portion of SEQ ID NO:1 comprising a coding sequence or its complement.

136. An isolated nucleic acid encoding a protein comprising the amino acid sequence of Figure 2 (SEQ ID NO:2).



137. An isolated nucleic acid encoding a human CXC Chemokine Receptor 3 (CXCR3) protein, wherein the amino acid sequence of said CXCR3 protein is a sequence encoded by the nucleic acid of Figure 1 (SEQ ID NO:1).

138. An isolated nucleic acid encoding a human CXC Chemokine Receptor 3 (CXCR3) protein or variant thereof, wherein the amino acid sequence of said CXCR3 protein or variant is at least about 80% identical to that of the protein shown in Figure 2 (SEQ ID NO:2), said protein or variant comprises the extracellular N-terminal segment of the protein shown in Figure 2 (SEQ ID NO:2), and said protein or variant can selectively bind one or more chemokines selected from the group consisting of IP-10 and Mig and mediate cellular signalling and/or a cellular response in response thereto.

139. A recombinant nucleic acid encoding a fusion protein comprising a human CXC Chemokine Receptor 3 (CXCR3) protein or functional variant thereof, wherein said CXCR3 protein or variant can selectively bind one or more chemokines and can mediate cellular signalling and/or a cellular response in response thereto, and wherein said CXCR3 protein or variant is encoded by a nucleic acid which hybridizes under high stringency conditions to a second nucleic acid and the sequence of said second nucleic acid is selected from the group consisting of Figure 1 (SEQ ID NO:1) and a sequence complementary to Figure 1 (SEQ ID NO:1).

140. The recombinant nucleic acid of Claim 139, wherein said CXCR3 protein or variant can selectively bind one or more chemokines selected from the group consisting of human IP-10 and human Mig.



141. A recombinant nucleic acid encoding a fusion protein comprising a human CXC Chemokine Receptor 3 (CXCR3) protein, wherein the amino acid sequence of said CXCR3 protein is a sequence encoded by the nucleic acid of Figure 1 (SEQ ID NO:1).

5 142. A recombinant nucleic acid encoding a fusion protein comprising a human CXC Chemokine Receptor 3 (CXCR3) protein, wherein the amino acid sequence of said CXCR3 protein consists of the amino acid sequence of Figure 2 (SEQ ID NO:2).

10 143. A recombinant nucleic acid encoding a fusion protein comprising a human CXC Chemokine Receptor 3 (CXCR3) protein or variant thereof, wherein the amino acid sequence of said CXCR3 protein or variant is at least about 80% identical to that of the protein shown in Figure 2 (SEQ ID NO:2), said CXCR3 protein or variant comprises the extracellular N-terminal segment of the protein shown in Figure 2 (SEQ ID NO:2), and said CXCR3 protein or variant can selectively bind 15 one or more chemokines selected from the group consisting of IP-10 and Mig and mediate cellular signalling and/or a cellular response in response thereto.

20 144. An isolated nucleic acid construct comprising a nucleic acid encoding a human CXC Chemokine Receptor 3 (CXCR3) protein or functional variant thereof, wherein said protein or variant can selectively bind one or more chemokines and can mediate cellular signalling and/or a cellular response in response thereto, and wherein said nucleic acid hybridizes under high stringency conditions to a second nucleic acid and the sequence of said second nucleic acid is selected from the group consisting of Figure 1 (SEQ ID NO:1) and a sequence complementary to Figure 1 (SEQ ID NO:1).



145. The isolated nucleic acid construct of Claim 144, wherein said CXCR3 protein or variant can selectively bind one or more chemokines selected from the group consisting of human IP-10 and human Mig.

146. The isolated nucleic acid construct of Claim 144, wherein the nucleic acid is operably linked to an expression control sequence.

147. The isolated nucleic acid construct of Claim 144, wherein the nucleic acid comprises SEQ ID NO:1, its complement, or a portion of SEQ ID NO:1 comprising the coding sequence or its complement.

148. An isolated nucleic acid construct comprising a nucleic acid encoding a human CXC Chemokine Receptor 3 (CXCR3) protein, wherein the amino acid sequence of said CXCR3 protein is a sequence encoded by the nucleic acid of Figure 1 (SEQ ID NO:1).

149. An isolated nucleic acid construct comprising a nucleic acid encoding a protein comprising the amino acid sequence of Figure 2 (SEQ ID NO:2).

150. An isolated nucleic acid construct, comprising a nucleic acid encoding a human CXC Chemokine Receptor 3 (CXCR3) protein or variant thereof, wherein the amino acid sequence of said CXCR3 protein or variant is at least about 80% identical to that of the protein shown in Figure 2 (SEQ ID NO:2), said protein or variant comprises the extracellular N-terminal segment of the protein shown in Figure 2 (SEQ ID NO:2), and said protein or variant can selectively bind one or more chemokines selected from the group consisting of IP-10 and Mig and mediate cellular signalling and/or a cellular response in response thereto.



151. An isolated nucleic acid construct comprising a recombinant nucleic acid of any one of Claims 139-143.

152. A host cell comprising a recombinant nucleic acid encoding a human CXC Chemokine Receptor 3 (CXCR3) protein or functional variant thereof, wherein  
5 said protein or variant can selectively bind one or more chemokines and can mediate cellular signalling and/or a cellular response in response thereto, and wherein said nucleic acid hybridizes under high stringency conditions to a second nucleic acid and the sequence of said second nucleic acid is selected from the group consisting of Figure 1 (SEQ ID NO:1) and a sequence complementary  
10 to Figure 1 (SEQ ID NO:1).

153. The host cell of Claim 152, wherein said CXCR3 protein or variant can selectively bind one or more chemokines selected from the group consisting of human IP-10 and human Mig.

154. The host cell of Claim 152, wherein the nucleic acid is operably linked to an  
15 expression control sequence.

155. The host cell of Claim 152, wherein the nucleic acid comprises SEQ ID NO:1, its complement, or a portion of SEQ ID NO:1 comprising the coding sequence or its complement.

156. A host cell comprising a recombinant nucleic acid encoding a human CXC Chemokine Receptor 3 (CXCR3) protein, wherein the amino acid sequence of  
20 said CXCR3 protein is a sequence encoded by the nucleic acid of Figure 1 (SEQ ID NO:1).



-132-

157. A host cell comprising a recombinant nucleic acid encoding a protein comprising the amino acid sequence of Figure 2 (SEQ ID NO:2).

158. A host cell comprising a recombinant nucleic acid encoding a human CXC Chemokine Receptor 3 (CXCR3) protein or variant thereof, wherein the amino acid sequence of said CXCR3 protein or variant is at least about 80% identical to that of the protein shown in Figure 2 (SEQ ID NO:2), said protein or variant comprises the extracellular N-terminal segment of the protein shown in Figure 2 (SEQ ID NO:2), and said protein or variant can selectively bind one or more chemokines selected from the group consisting of IP-10 and Mig and mediate cellular signalling and/or a cellular response in response thereto.

159. A host cell comprising the recombinant nucleic acid encoding a fusion protein of any one of Claims 139-143.

160. A method for producing a human CXCR3 protein or variant thereof, wherein said protein or variant can selectively bind one or more chemokines and can mediate cellular signalling and/or a cellular response in response thereto, the method comprising:

(a) introducing into a host cell a nucleic acid encoding said human CXCR3 protein or variant thereof, wherein said nucleic acid hybridizes under high stringency conditions to a second nucleic acid and the sequence of said second nucleic acid is selected from the group consisting of Figure 1 (SEQ ID NO:1) and a sequence complementary to Figure 1 (SEQ ID NO:1), whereby a recombinant host cell is produced having said nucleic acid operably linked to an expression control sequence; and

(b) maintaining the host cells produced in step (a) under conditions whereby the nucleic acid is expressed.



-133-

161. The method of Claim 160, further comprising the step of isolating the human CXCR3 protein or variant thereof.
162. A method for producing a human CXCR3 protein or variant thereof comprising maintaining a host cell of Claim 152 under conditions suitable for expression of the nucleic acid.  
5
163. The method of Claim 162 further comprising the step of isolating the human CXCR3 protein or variant thereof.
164. The method of Claim 163, wherein the protein or variant thereof can selectively bind one or more chemokines selected from the group consisting of human IP-10 and human Mig.  
10
165. A non-human transgenic animal having genome that comprises a recombinant nucleic acid encoding a mammalian CXC Chemokine Receptor 3 (CXCR3) protein or ligand binding variant thereof, wherein said ligand binding variant comprises an amino acid sequence that has at least about 80% amino acid sequence identity with SEQ ID NO:2.  
15
166. The non-human transgenic animal of Claim 165 wherein said CXCR3 protein has the amino acid sequence of SEQ ID NO:2.
167. The non-human transgenic animal of Claim 165 wherein the amino acid sequence of said CXCR3 protein is a sequence encoded by SEQ ID NO:1.
- 20 168. The non-human transgenic animal of Claim 165 wherein said ligand binding variant comprises an amino acid sequence that has at least about 80% amino acid sequence identity with SEQ ID NO:2.



169. The non-human transgenic animal of Claim 165 wherein said transgenic animal is a mouse, guinea pig or sheep.

170. A non-human transgenic animal having a genome that comprises an inactivated or disabled CXC chemokine Receptor 3 (CXCR3) gene, wherein said inactivated or disabled CXCR3 gene results from insertion of a nucleic acid which disrupts the endogenous CXCR3 gene.

171. The non-human transgenic animal of Claim 170 wherein said insertion results from homologous recombination.

172. The non-human transgenic animal of Claim 170 wherein said animal is homozygous for said inactivated or disabled CXCR3 gene.

173. The non-human transgenic animal of Claim 170 wherein said transgenic animal is a mouse, guinea pig or sheep.

174. A promoter of at least one function characteristic of a mammalian CXCR3 protein identified according to the method of Claim 26 or 55, wherein said promoter is other than IP-10 or Mig.

DATED this 12<sup>th</sup> day of April 2001

20 MILLENNIUM PHARMACEUTICALS, INC. AND THEODOR-KOCHER INSTITUTE

by DAVIES COLLISON CAVE  
Patent Attorneys for the Applicants



1/15

## Complete nucleotide sequence of IP10/MigR (MLRA) cDNA

CCAAACACAA	GCACCAAAGC	AGAGGGCAG	GCAGCACACC	ACCCAGCAGC	CAGAGCACCA	60
GCCCAGCCAT	GGTCCTTGAG	GTGAGTGACC	ACCAAGTGCT	AAATGACGCC	GAGGTTGCCG	120
CCCTCCTGGA	GAACITCAGC	TCTTCCTATG	ACTATGGAGA	AAACGAGAGT	GACTCGTGCT	180
GTACCTCCCC	GCCCTGCCCA	CAGGACTTCA	GCCTGAACCT	CGACCGGGCC	TTCTGCCCC	240
CCCTCTACAG	CCTCCTCTTT	CTGCTGGGGC	TGCTGGGCAA	CGGCGCGGTG	GCAGCCGTGC	300
TGCTGAGCCG	GCGGACAGCC	CTGAGCAGCA	CCGACACCTT	CCTGCTCCAC	CTAGCTGTAG	360
CAGACACGCT	GCTGGTGCTG	ACACTGCCGC	TCTGGGAGT	GGACGCTGCC	GTCCAGTGGG	420
TCTTTGGCTC	TGGCCTCTGC	AAAGTGGCAG	GTGCCCTCTT	CAACATCAAC	TTCTACGCAG	480
GAGCCCTCCT	GCTGGCCTGC	ATCAGCTTTG	ACCGCTACCT	GAACATAGTT	CATGCCACCC	540
AGCTCTACCG	CCGGGGGGCCC	CCGGCCCGCG	TGACCCCTCAC	CTGCCTGGCT	GTCTGGGGC	600
TCTGCCTGCT	TTTCGCCCTC	CCAGACTTCA	TCTTCCTGTC	GGCCCACACAC	GACGAGCGCC	660
TCAACGCCAC	CCACTGCCAA	TACAACTTCC	CACAGGTGGG	CCGCACGGCT	CTGCGGGTGC	720
TGCAGCTGGT	GGCTGGCTTT	CTGCTGCCCG	TGCTGGTCAT	GGCCTACTGC	TATGCCACACA	780
TCCTGGCCGT	GCTGCTGGTT	TCCAGGGGCC	AGCGGCCCT	GCAGGCCATG	CGGCTGGTGG	840
TGGTGGTCGT	GGTGGCCTTT	GCCCTCTGCT	GGACCCCCCTA	TCACCTGGTG	TGCTGGTGG	900
ACATCCTCAT	GGACCTGGC	GCTTGGCCC	GCAACTGTGG	CCGAGAAAGC	AGGGTAGACG	960
TGGCCAAGTC	GGTCACCTCA	GGCCTGGCT	ACATGCACTG	CTGCCTCAAC	CCGCTGCTCT	1020
ATGCCTTGT	AGGGGTCAAG	TTCCGGGAGC	GGATGTGGAT	GCTGCTCTTG	CGCCTGGGCT	1080
GCCCCAACCA	GAGAGGGCTC	CAGAGGCAGC	CATCGTCTTC	CCGCCGGGAT	TCATCCTGGT	1140
CTGAGACCTC	AGAGGCCTCC	TACTCGGGCT	TGTGAGGGCG	GAATCCGGGC	TCCCCCTTCG	1200
CCCACAGTCT	GAATTCCCCG	CATTCCAGGC	TCCTCCCTCC	CTCTGCCGGC	TCTGGCTCTC	1260
CCCAATATCC	TCGCTCCCCG	GAATCACTGG	CAGCCCCAGC	ACCACCAAGT	CTCCCAGGAA	1320
GCCACCCCTCC	CAGCTCTGAG	GAATGCACCA	TTGCTGCTCC	TTAGCTGCCA	AGCCCCATCC	1380
TGCCGCCCGA	GGTGGCTGCC	TGGAGCCCCA	CTGCCCTCT	CATTGGAAA	CTAAAACCTTC	1440
ATCTTCCCCA	AGTGCAGGGGA	GTACAAGGCA	TGGCGTAGAG	GGTGTGCC	CATGAAGCCA	1500
CAGCCCCAGGC	CTCCAGCTCA	GCAGTGACTG	TGGCCATGGT	CCCCAAGACC	TCTATATTTG	1560
CTCTTTTATT	TTTATGTCTA	AAATCCTGCT	TAAAACTTT	CAATAAACAA	GATCGTCAGG	1620
ACCTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	1670

## FIGURE 1

SUBSTITUTE SHEET (RULE 26)

031000 42606

IP10/MigR	MVLEVSDHQVLNDAEVAALLEN <u>FSSSYD</u> YGENESDSCCTS <u>PPCPQDFSLNF</u> DRA <u>FLPALYSLL</u> <u>FLLG</u> <u>LLNG</u> AVA <u>AV</u> LLS <u>RRTA</u> LS <u>STD</u> T	90	
▼      ▼			
IP10/MigR	TM2	TM3	TM4
	<u>FLLHL</u> <u>LA</u> <u>VADT</u> <u>LLVL</u> <u>TLPL</u> <u>WAV</u> -DAA <u>VQW</u> <u>VFG</u> <u>SGL</u> <u>CKVAG</u> <u>ALF</u> <u>NINFYAG</u> <u>ALL</u> <u>LAC</u> <u>ISF</u> <u>DRYLN</u> <u>IVH</u> <u>ATQ</u> <u>LYR</u> <u>RGPP</u> <u>AV</u> <u>TLT</u> <u>CL</u> <u>AVW</u> <u>GLC</u>		
	179		
IP10/MigR	TM4	TM5	TM6
	<u>LLF</u> <u>ALP</u> <u>DFI</u> <u>FLSAHH</u> <u>DERLN</u> <u>NATH</u> <u>CQYNF</u> <u>PQVG</u> - - - - - <u>RTA</u> <u>LRL</u> <u>QLVAG</u> <u>FLL</u> <u>PLL</u> <u>VMA</u> <u>CYA</u> <u>HIL</u> <u>AV</u> <u>LLV</u> <u>SRG</u> <u>QRL</u> - <u>RAM</u> <u>RLV</u> <u>VV</u> <u>VV</u> <u>VV</u>		
	262		
IP10/MigR	TM6	TM7	
	<u>A</u> <u>FALC</u> <u>WT</u> <u>PYH</u> <u>L</u> <u>V</u> <u>V</u> <u>L</u> <u>V</u> <u>D</u> <u>I</u> <u>L</u> <u>M</u> <u>D</u> <u>L</u> <u>G</u> <u>A</u> <u>R</u> <u>N</u> <u>C</u> <u>G</u> <u>E</u> <u>R</u> <u>S</u> <u>R</u> <u>V</u> <u>D</u> <u>V</u> <u>A</u> <u>K</u> <u>S</u> <u>V</u> <u>T</u> <u>G</u> <u>L</u> <u>G</u> <u>Y</u> <u>M</u> <u>H</u> <u>C</u> <u>C</u> <u>L</u> <u>N</u> <u>P</u> <u>L</u> <u>Y</u> <u>A</u> <u>F</u> <u>V</u> <u>G</u> <u>V</u> <u>K</u> <u>F</u> <u>R</u> <u>E</u> <u>M</u> <u>W</u> <u>M</u> <u>L</u> <u>L</u> <u>R</u> - - - <u>L</u> <u>G</u> <u>C</u> <u>P</u> <u>N</u> <u>Q</u> <u>R</u> <u>G</u> <u>L</u> <u>Q</u> <u>R</u> <u>Q</u> <u>P</u> <u>S</u>		
	349		
IP10/MigR	<u>S</u> <u>S</u> <u>R</u> <u>R</u> <u>D</u> <u>S</u> <u>S</u> <u>W</u> <u>S</u> <u>E</u> <u>T</u> <u>S</u> <u>E</u> <u>A</u> <u>Y</u> <u>S</u> <u>G</u> <u>L</u>	368	

FIGURE 2

## SUBSTITUTE SHEET (RULE 26)




FIGURE 3A




FIGURE 3B




FIGURE 3C




FIGURE 4A




FIGURE 4B




FIGURE 5A




FIGURE 5B

Cold IP-10 (nM)

WO 98/11218

6/15




FIGURE 6

SUBSTITUTE SHEET (RULE 26)

7/15




FIGURE 7A




FIGURE 7B




FIGURE 7C

CXCR3 staining →




FIGURE 8C




FIGURE 8B

↑ CXCR3




FIGURE 8A




FIGURE 9A





FIGURE 9B



→ CXCR3

FIGURE 9C

10/15




FIGURE 9D





FIGURE 9E



→ CXCR3

FIGURE 9F

SUBSTITUTE SHEET (RULE 26)




FIGURE 9G





FIGURE 9H



→ CXCR3

FIGURE 9I

SUBSTITUTE SHEET (RULE 26)

12/15




FIGURE 10




FIGURE 11





FIGURE 12

SUBSTITUTE SHEET (RULE 26)

13/15

## SUBSTITUTE SHEET (RULE 26)



FIGURES 13A - 13H

14/15




FIGURE 14A




FIGURE 14B




FIGURE 14C




FIGURE 14D




FIGURE 15

SUBSTITUTE SHEET (RULE 26)