



(86) Date de dépôt PCT/PCT Filing Date: 2000/09/14
(87) Date publication PCT/PCT Publication Date: 2001/03/22
(45) Date de délivrance/Issue Date: 2013/11/19
(85) Entrée phase nationale/National Entry: 2002/03/13
(86) N° demande PCT/PCT Application No.: US 2000/021970
(87) N° publication PCT/PCT Publication No.: 2001/019955
(30) Priorité/Priority: 1999/09/14 (US60/153,831)

(51) Cl.Int./Int.Cl. *C12N 15/54* (2006.01),
A61K 38/46 (2006.01), *C07H 21/04* (2006.01),
C07K 14/00 (2006.01), *C12N 1/20* (2006.01),
C12N 15/00 (2006.01), *C12N 15/63* (2006.01),
C12N 5/10 (2006.01), *C12N 9/12* (2006.01),
C12N 9/14 (2006.01), *C12N 9/16* (2006.01),
C12P 21/00 (2006.01), *A61K 38/00* (2006.01)
(72) Inventeur/Inventor:
CANFIELD, WILLIAM M., US
(73) Propriétaire/Owner:
GENZYME CORPORATION, US
(74) Agent: NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : ENZYMES A VOIE DE CIBLAGE LYSOSOMIAL
(54) Title: LYSOSOMAL TARGETING PATHWAY ENZYMES

(57) **Abrégé/Abstract:**

The lysosomal targeting pathway enzymes GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase are recovered and used to produce highly phosphorylated lysosomal hydrolases that can be used to treat lysosomal storage diseases. Generally, the nucleic acid molecules coding for the enzymes are incorporated into expression vectors that are used to transfect host cells that express the enzymes. The expressed enzymes are recovered and used to treat lysosomal hydrolases having high mannose structures. The resulting lysosomal hydrolases have asparagine-linked oligosaccharides that are highly phosphorylated with mannose phosphate. The highly phosphorylated lysosomal hydrolases are readily taken into the cell and into the lysosome during enzyme replacement therapy procedures.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19955 A3

(51) International Patent Classification⁷: C12N 9/12, 9/14, 1/20, 15/00, C07H 21/04, A61K 38/44, 38/51, C07K 14/00

(21) International Application Number: PCT/US00/21970

(22) International Filing Date:
14 September 2000 (14.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/153,831 14 September 1999 (14.09.1999) US

(71) Applicant and

(72) Inventor: CANFIELD, William, M. [US/US]; 7108 Brentwood Avenue, Oklahoma City, OK 73116 (US).

(74) Agents: CHINN, Richard, L. et al.; Oblon, Spivak, McClelland, Maier & Neustadt, P.C., 4th Floor, Crystal Square Five, 1755 Jefferson Davis Highway, Arlington, VA 22202 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
4 October 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: LYSOSOMAL TARGETING PATHWAY ENZYMES

(57) Abstract: The lysosomal targeting pathway enzymes GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase are recovered and used to produce highly phosphorylated lysosomal hydrolases that can be used to treat lysosomal storage diseases. Generally, the nucleic acid molecules coding for the enzymes are incorporated into expression vectors that are used to transfect host cells that express the enzymes. The expressed enzymes are recovered and used to treat lysosomal hydrolases having high mannose structures. The resulting lysosomal hydrolases have asparagine-linked oligosaccharides that are highly phosphorylated with mannose phosphate. The highly phosphorylated lysosomal hydrolases are readily taken into the cell and into the lysosome during enzyme replacement therapy procedures.

WO 01/19955 A3



TITLE OF THE INVENTION

LYSOSOMAL TARGETING PATHWAY ENZYMES

BACKGROUND OF THE INVENTIONField of the Invention

This invention relates generally to enzymes involved in the lysosomal targeting pathway and particularly to isolated and purified GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase, nucleic acids encoding the enzymes, processes for production of recombinant GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase, and the use of the enzymes for the preparation of highly phosphorylated lysosomal enzymes that are useful for the treatment of lysosomal storage diseases.

Description of the Prior Art

Lysosomes and Lysosomal Storage Diseases

Lysosomes are organelles in eukaryotic cells that function in the degradation of macromolecules into component parts that can be reused in biosynthetic pathways or discharged by the cell as waste. Normally, these macromolecules are broken down by enzymes known as lysosomal enzymes or lysosomal hydrolases. However, when a lysosomal enzyme is not present in the lysosome or does not function properly, the enzymes specific macromolecular substrate accumulates in the lysosome as "storage material" causing a variety of diseases, collectively known as lysosomal storage diseases.

Lysosomal storage diseases can cause chronic illness and death in hundreds of

individuals each year. There are approximately 50 known lysosomal storage diseases, e.g., Pompe Disease, Hurler Syndrome, Fabry Disease, Maroteaux-Lamy Syndrome (mucopolysaccharidosis VI), Morquio Syndrome (mucopolysaccharidosis IV), Hunter Syndrome (mucopolysaccharidosis II), Farber Disease, Acid Lipase Deficiency, Krabbe Disease, and Sly Syndrome (mucopolysaccharidosis VII). In each of these diseases, lysosomes are unable to degrade a specific compound or group of compounds because the enzyme that catalyzes a specific degradation reaction is missing from the lysosome, is present in low concentrations in the lysosome, or is present at sufficient concentrations in the lysosome but is not functioning properly.

Lysosomal storage diseases have been studied extensively and the enzymes (or lack thereof) responsible for particular diseases have been identified. Most of the diseases are caused by a deficiency of the appropriate enzyme in the lysosome, often due to mutations or deletions in the structural gene for the enzyme. For some lysosomal storage diseases, the enzyme deficiency is caused by the inability of the cell to target and transport the enzymes to the lysosome, e.g., I-cell disease and pseudo-Hurler polydystrophy.

Lysosomal Storage diseases have been studied extensively and the enzymes (or lack thereof) responsible for particular diseases have been identified (Scriver, Beaudet, Sly, and Vale, eds., The Metabolic Basis of Inherited Disease, 6th Edition, 1989, Lysosomal Enzymes, Part 11, Chapters 61-72, pp. 1565-1839). Within each disease, the severity and the age at which the disease presents may be a function of the amount of residual lysosomal enzyme that exists in the patient.

Lysosomal Targeting Pathway

The lysosomal targeting pathways have been studied extensively and the process by

which lysosomal enzymes are synthesized and transported to the lysosome has been well described. Kornfeld, S. (1986). "Trafficking of lysosomal enzymes in normal and disease states." *Journal of Clinical Investigation* 77: 1-6 and Kornfeld, S. (1990). "Lysosomal enzyme targeting." *Biochem. Soc. Trans.* 18: 367-374. Generally, lysosomal enzymes are synthesized by membrane-bound polysomes in the rough endoplasmic reticulum ("RER") along with secretory glycoproteins. In the RER, lysosomal enzymes acquire N-linked oligosaccharides by the en-bloc transfer of a preformed oligosaccharide from dolichol phosphate containing 2 N-acetylglucosamine, 9-mannose and 3-glucose. Glycosylated lysosomal enzymes are then transported to the Golgi apparatus along with secretory proteins. In the cis-Golgi or intermediate compartment lysosomal enzymes are specifically and uniquely modified by the transfer of GlcNAc-phosphate to specific mannoses. In a second step, the GlcNAc is removed thereby exposing the mannose 6-phosphate ("M6P") targeting determinant. The lysosomal enzymes with the exposed M6P binds to M6P receptors in the trans-Golgi and is transported to the endosome and then to the lysosome. In the lysosome, the phosphates are rapidly removed by lysosomal phosphatases and the mannoses are removed by lysosomal mannosidases (Einstein, R. and Gabel, C.A. (1991). "Cell- and ligand-specific dephosphorylation of acid hydrolases: evidence that the mannose 6-phosphate is controlled by compartmentalization." *Journal of Cell Biology* 112: 81-94).

The synthesis of lysosomal enzymes having exposed M6P is catalyzed by two different enzymes, both of which are essential if the synthesis is to occur. The first enzyme is UDP-N-acetylglucosamine: lysosomal enzyme N-Acetylglucosamine-1-phosphotransferase ("GlcNAc-phosphotransferase") (E.C. 2.7.8.17). GlcNAc-phosphotransferase catalyzes the transfer of N-acetylglucosamine-1-phosphate from UDP-GlcNAc to the 6 position of α 1,2-linked mannoses on the lysosomal enzyme. The recognition and addition of N-

acetylglucosamine-1-phosphate to lysosomal hydrolases by GlcNAc-phosphotransferase is the critical and determining step in lysosomal targeting. The second step is catalyzed by N-acetylglucosamine-1-phosphodiester α -N-Acetylglucosaminidase ("phosphodiester α -GlcNAcase") (E.C. 3.1.4.45). Phosphodiester α -GlcNAcase catalyzes the removal of N-Acetylglucosamine from the GlcNAc-phosphate modified lysosomal enzyme to generate a terminal M6P on the lysosomal enzyme. Preliminary studies of these enzymes have been conducted. Bao *et al.*, in *The Journal of Biological Chemistry*, Vol. 271, Number 49, Issue of December 6, 1996, pp. 31437-31445, relates to a method for the purification of bovine UDP-N-acetylglucosamine: Lysosomal enzyme N-Acetylglucosamine-1-phosphotransferase and proposes a hypothetical subunit structure for the protein. Bao *et al.*, in *The Journal of Biological Chemistry*, Vol. 271, Number 49, Issue of December 6, 1996, pp. 31446-31451, relates to the enzymatic characterization and identification of the catalytic subunit for bovine UDP-N-acetylglucosamine: Lysosomal enzyme N-Acetylglucosamine-1-phosphotransferase. Kornfeld *et al.*, in *The Journal of Biological Chemistry*, Vol. 273, Number 36, Issue of September 4, 1998, pp. 23203-23210, relates to the purification and multimeric structure of bovine N-Acetylglucosamine-1-phosphodiester α -N-Acetylglucosaminidase. However, the proprietary monoclonal antibodies required to isolate these proteins have not been made available to others and the protein sequences for the enzymes used in these preliminary studies have not been disclosed.

Although the lysosomal targeting pathway is known and the naturally occurring enzymes involved in the pathway have been partially studied, the enzymes responsible for adding M6P in the lysosomal targeting pathway are difficult to isolate and purify and are poorly understood. A better understanding of the lysosomal targeting pathway enzymes and the molecular basis for their action is needed to assist with the development of effective

techniques for the utilization of these enzymes in methods for the treatment of lysosomal storage diseases, particularly in the area of targeted enzyme replacement therapy.

Treatment of Lysosomal Storage Diseases

Lysosomal storage diseases caused by the lack of enzymes can in theory be treated using enzyme replacement therapy, *i.e.*, by administering isolated and purified enzymes to the patient to treat the disease. However, to be effective, the lysosomal enzyme administered must be internalized by the cell and transported to the lysosome. Naturally occurring enzymes and their recombinant equivalents, however, have been of limited value in enzyme replacement therapy because the purified or recombinant lysosomal enzymes do not contain adequate amounts of exposed M6P, or contain undesirable oligosaccharides which mediate their destruction. Without sufficient M6P, the administered lysosomal enzyme cannot efficiently bind to M6P receptors and be transported to the lysosome. For example, human acid α -glucosidase purified from placenta contains oligomannose oligosaccharides which are not phosphorylated (Mutsaers, J. H. G. M., Van Halbeek, H., Vliegenthart, J. F. G., Tager, J. M., Reuser, A. J. J., Kroos, M., and Galjaard, H. (1987). "Determination of the structure of the carbohydrate chains of acid α -glucosidase from human placenta." *Biochimica et Biophysica Acta* 911: 244 - 251), and this glycoform of the enzyme is not efficiently internalized by cells (Reuser, A. J., Kroos, M. A., Ponne, N. J., Wolterman, R. A., Loonen, M. C., Busch, H. T., Visser, W. J., and Bolhuis, P. A. (1984). "Uptake and stability of human and bovine acid α -glucosidase in cultured fibroblasts and skeletal muscle cells from glycogenosis type II patients." *Experimental Cell Research* 155: 178-189). As a result of the inability to purify or synthesize lysosomal enzymes with the desired oligosaccharide

structures, these enzyme preparations are inefficiently targeted to affected cells and are of limited effectiveness in the treatment of these diseases. There exists, therefore, a need for enzymes that can be used in enzyme replacement therapy procedures, particularly highly phosphorylated enzymes that will be efficiently internalized by the cell and transported to the lysosome.

SUMMARY OF THE INVENTION

It is, therefore, an object of the present invention to provide biologically active GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase as isolated and purified polypeptides.

It is another object of the present invention to provide nucleic acid molecules encoding GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase.

It is another object of the present invention to provide expression vectors having DNA that encodes GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase.

It is a further object of the present invention to provide host cells that have been transfected with expression vectors having DNA that encodes GlcNAc-phosphotransferase or phosphodiester α -GlcNAcase.

It is another object of the present invention to provide methods for producing recombinant GlcNAc-phosphotransferase and recombinant phosphodiester α -GlcNAcase by culturing host cells that have been transfected or transformed with expression vectors having DNA that encodes GlcNAc-phosphotransferase or phosphodiester α -GlcNAcase.

It is another object of the present invention to provide isolated and purified recombinant GlcNAc-phosphotransferase and recombinant phosphodiester α -GlcNAcase.

It is another object of the present invention to provide methods for the preparation of highly phosphorylated lysosomal enzymes that are useful for the treatment of lysosomal storage diseases.

It is a further object of the present invention to provide highly phosphorylated lysosomal hydrolases that are useful for the treatment of lysosomal storage diseases.

It is still another object of the present invention to provide methods for the treatment of lysosomal storage diseases.

It is still another object of the present invention to provide monoclonal antibodies capable of selectively binding to bovine GlcNAc-phosphotransferase and to bovine phosphodiester α -GlcNAcase.

These and other objects are achieved by recovering isolated and purified biologically active GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase and using the enzymes to obtain nucleic acid molecules that encode for the enzymes. The nucleic acid molecules coding for either enzyme are incorporated into expression vectors that are used to transfect host cells that express the enzyme. The expressed enzyme is recovered and used to prepare highly phosphorylated lysosomal hydrolases useful for the treatment of lysosomal storage diseases. In particular, the enzymes are used to produce highly phosphorylated-lysosomal hydrolases that can be effectively used in enzyme replacement therapy procedures.

Lysosomal hydrolases having high mannose structures are treated with GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase resulting in the production of asparagine-linked oligosaccharides that are highly modified with mannose 6-phosphate ("M6P"). The treated hydrolase binds to M6P receptors on the cell membrane and is transported into the cell and delivered to the lysosome where it can perform its normal or a desired function.

Other aspects and advantages of the present invention will become apparent from the following more detailed description of the invention taken in conjunction with the accompanying drawings.

BRIEF OF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a model of the subunit structure of GlcNAc-phosphotransferase. The enzyme is a complex of six polypeptides. The α - and β -subunits are the product of a single gene. Following translation, the α - and β -subunits are separated by proteolytic cleavage between Lys⁹²⁹ and Asp⁹³⁰. The α -subunit is a type II membrane glycoprotein with a single amino terminal membrane spanning domain. The β -subunit is a type I membrane spanning glycoprotein with a single carboxyl terminal membrane spanning domain. The γ -subunit is the product of a second gene. The γ -subunit is a soluble protein with a cleaved signal peptide. The α -, β -, and γ -subunits are all tightly associated.

Figure 2 shows a model of the subunit structure of phosphodiester α -GlcNAcase. The enzyme is a tetramer composed of four identical subunits arranged as two non-covalently-associated dimers which are themselves disulfide-linked. The single subunit is a type I membrane protein containing a signal peptide, a pro region not present in the mature enzyme and a single carboxyl terminal membrane spanning domain.

Figure 3 shows a diagram of recombinant glycoprotein expression in CHO cells. In overexpressing CHO cells, the rh-GAA is processed along the pathways 1 and 2, depending on whether or not the enzyme is acted upon by GlcNAc-phosphotransferase (GnPT). Secreted GAA contains predominantly sialylated biantennary complex-type glycans and is not a substrate for GlcNAc-phosphotransferase. In the presence of the α 1,2-mannosidase inhibitors, 1-deoxymannojirimycin or kifunensine conversion of MAN9 to MAN5 structures

is blocked, resulting in secretion of GAA-bearing MAN7-9 structures which can be modified with GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase (UCE) generating phosphorylated species (pathway 3).

Figure 4 shows transient expression analysis of various plasmid constructs of the α/β and γ subunits of human GlcNAc-phosphotransferase. Plasmids containing the α/β and/or the γ subunits were transfected into 293T cells, the expressed protein was purified from the culture at 23, 44.5 and 70 hours after transfection and relative amounts of expression were assessed by measuring phosphotransferase activity using methyl- α -D-mannoside and [β - ^{32} P] UDP-GlcNAc as substrates.

DETAILED DESCRIPTION OF THE INVENTION

The term "GlcNAc-phosphotransferase" as used herein refers to enzymes that are capable of catalyzing the transfer of N-acetylglucosamine-1-phosphate from UDP-GlcNAc to the 6' position of α 1,2-linked mannoses on lysosomal enzymes.

The term "phosphodiester α -GlcNAcase" as used herein refers to enzymes that are capable of catalyzing the removal of N-Acetylglucosamine from GlcNAc-phosphate-mannose diester modified lysosomal enzymes to generate terminal M6P.

The terms "GlcNAc-phosphotransferase" and "phosphodiester α -GlcNAcase" as used herein refer to enzymes obtained from any eukaryotic species, particularly mammalian species such as bovine, porcine, murine, equine, and human, and from any source whether natural, synthetic, semi-synthetic, or recombinant. The terms encompass membrane-bound enzymes and soluble or truncated enzymes having less than the complete amino acid sequence and

biologically active variants and gene products.

The term "naturally occurring" as used herein means an endogenous or exogenous protein isolated and purified from animal tissue or cells.

The term "isolated and purified" as used herein means a protein that is essentially free of association with other proteins or polypeptides, *e.g.*, as a naturally occurring protein that has been separated from cellular and other contaminants by the use of antibodies or other methods or as a purification product of a recombinant host cell culture.

The term "biologically active" as used herein means an enzyme or protein having structural, regulatory, or biochemical functions of a naturally occurring molecule.

The term "nucleotide sequence" as used herein means a polynucleotide molecule in the form of a separate fragment or as a component of a larger nucleic acid construct that has been derived from DNA or RNA isolated at least once in substantially pure form (*i.e.*, free of contaminating endogenous materials) and in a quantity or concentration enabling identification, manipulation, and recovery of its component nucleotide sequences by standard biochemical methods. Such sequences are preferably provided in the form of an open reading frame uninterrupted by internal non-translated sequences, or introns that are typically present in eukaryotic genes. Sequences of non-translated DNA may be present 5' or 3' from an open reading frame where the same do not interfere with manipulation or expression of the coding region.

The term "nucleic acid molecule" as used herein means RNA or DNA, including cDNA, single or double stranded, and linear or covalently closed molecules. A nucleic acid molecule may also be genomic DNA corresponding to the entire gene or a substantial portion thereof to fragments and derivatives thereof. The nucleotide sequence may correspond to the naturally occurring nucleotide sequence or may contain single or multiple nucleotide

substitutions, deletions and/or additions including fragments thereof. All such variations in the nucleic acid molecule retain the ability to encode a biologically active enzyme when expressed in the appropriate host or an enzymatically active fragment thereof. The nucleic acid molecule of the present invention may comprise solely the nucleotide sequence encoding an enzyme or may be part of a larger nucleic acid molecule that extends to the gene for the enzyme. The non-enzyme encoding sequences in a larger nucleic acid molecule may include vector, promoter, terminator, enhancer, replication, signal sequences, or non-coding regions of the gene.

The term "variant" as used herein means a polypeptide substantially homologous to a naturally occurring protein but which has an amino acid sequence different from the naturally occurring protein (human, bovine, ovine, porcine, murine, equine, or other eukaryotic species) because of one or more deletions, insertions, derivations, or substitutions. The variant amino acid sequence preferably is at least 50% identical to a naturally occurring amino acid sequence but is most preferably at least 70% identical. Variants may comprise conservatively substituted sequences wherein a given amino acid residue is replaced by a residue having similar physiochemical characteristics. Conservative substitutions are well known in the art and include substitution of one aliphatic residue for another, such as Ile, Val, Leu, or Ala for one another, or substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Conventional procedures and methods can be used for making and using such variants. Other such conservative substitutions such as substitutions of entire regions having similar hydrophobicity characteristics are well known. Naturally occurring variants are also encompassed by the present invention. Examples of such variants are enzymes that result from alternate mRNA splicing events or from proteolytic cleavage of the enzyme that leave the enzyme biologically active and capable of

performing its catalytic function. Alternate splicing of mRNA may yield a truncated but biologically active protein such as a naturally occurring soluble form of the protein. Variations attributable to proteolysis include differences in the N- or C-termini upon expression in different types of host cells due to proteolytic removal of one or more terminal amino acids from the protein.

The term "substantially the same" as used herein means nucleic acid or amino acid sequences having sequence variations that do not materially affect the nature of the protein, i.e., the structure and/or biological activity of the protein. With particular reference to nucleic acid sequences, the term "substantially the same" is intended to refer to the coding region and to conserved sequences governing expression and refers primarily to degenerate codons encoding the same amino acid or alternate codons encoding conservative substitute amino acids in the encoded polypeptide. With reference to amino acid sequences, the term "substantially the same" refers generally to conservative substitutions and/or variations in regions of the polypeptide not involved in determination of structure or function.

The term "percent identity" as used herein means comparisons among amino acid sequences as defined in the UWGCG sequence analysis program available from the University of Wisconsin. (Devereaux *et al.*, Nucl. Acids Res. 12: 387-397 (1984)).

The term "highly phosphorylated lysosomal hydrolase" as used to herein means a level of phosphorylation on a purified lysosomal hydrolase which could not be obtained by only isolating the hydrolase from a natural source and without subsequent treatment with the GlcNAc-phosphotransferase and phosphodiester- α -GlcNAcase. In particular, "highly phosphorylated lysosomal hydrolase" means a lysosomal hydrolase that contains from about 6% to about 100% bis-phosphorylated oligosaccharides.

This invention is not limited to the particular methodology, protocols, cell lines,

vectors, and reagents described because these may vary. Further, the terminology used herein is for the purpose of describing particular embodiments only and is not intended to limit the scope of the present invention. As used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise, *e.g.*, reference to "a host cell") includes a plurality of such host cells.

Because of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding GlcNAc-phosphotransferase, phosphodiester α -GlcNAcase, or other sequences referred to herein may be produced. Some of these sequences will be highly homologous and some will be minimally homologous to the nucleotide sequences of any known and naturally occurring gene. The present invention contemplates each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring GlcNAc-phosphotransferase or phosphodiester α -GlcNAcase, and all such variations are to be considered as being specifically disclosed.

Unless defined otherwise, all technical and scientific terms and any acronyms used herein have the same meanings as commonly understood by one of ordinary skill in the art in the field of the invention. Although any methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred methods, devices, and materials are described herein.

However, nothing herein is to be construed as an admission that the invention is

not entitled to antedate such disclosure by virtue of prior invention.

The Invention

GlcNAc-phosphotransferase

In one aspect, the present invention provides isolated and purified biologically active GlcNAc-phosphotransferase, nucleic acid molecules encoding GlcNAc-phosphotransferase and its subunits, expression vectors having a DNA that encodes GlcNAc-phosphotransferase, host cells that have been transfected or transformed with expression vectors having DNA that encodes GlcNAc-phosphotransferase, methods for producing recombinant GlcNAc-phosphotransferase by culturing host cells that have been transfected or transformed with expression vectors having DNA that encodes GlcNAc-phosphotransferase, isolated and purified recombinant GlcNAc-phosphotransferase, and methods for using GlcNAc-phosphotransferase for the preparation of highly phosphorylated lysosomal enzymes that are useful for the treatment of lysosomal storage diseases.

To obtain isolated and purified GlcNAc-phosphotransferase and its subunits and the nucleic acid molecules encoding the enzyme according to the present invention, bovine GlcNAc phosphotransferase was obtained and analyzed as follows. Splenocytes from mice immunized with a partially purified preparation of bovine GlcNAc-phosphotransferase were fused with myeloma cells to generate a panel of hybridomas. Hybridomas secreting monoclonal antibodies specific for GlcNAc-phosphotransferase were identified by immunocapture assay. In this assay, antibodies which could capture GlcNAc-phosphotransferase from a crude source were identified by assay of immunoprecipitates with a specific GlcNAc-phosphotransferase enzymatic assay. Hybridomas were subcloned twice, antibody produced in ascites culture, coupled to a solid support and evaluated for

immunoaffinity chromatography. Monoclonal PT18-Emphaze was found to allow a single step purification of GlcNAc-phosphotransferase to homogeneity. Bao, *et.al.*, The Journal of Biological Chemistry, Vol. 271, Number 49, Issue of December 6, 1996, pp. 31437-31445 relates to a method for the purification of bovine UDP-N-acetylglucosamine:Lysosomal-enzyme N-Acetylglucosamine-l-phosphotransferase and proposes a hypothetical subunit structure for the protein. Bao, *et. al.*, *The Journal of Biological Chemistry*, Vol. 271, Number 49, Issue of December 6, 1996, pp. 31446-31451. Using this technique, the enzyme was purified 488,000-fold in 29% yield. The eluted GlcNAc-phosphotransferase has a specific activity of $>10^6$, preferably $>5 \times 10^6$, more preferably $>12 \times 10^6$ pmol/h/mg and is apparently a homogenous, multi-subunit enzyme based on silver-stained SDS-PAGE. The monoclonal antibody labeled PT18 was selected for use in further experiments. A hybridoma secreting monoclonal antibody PT 18 was deposited with the American Type Culture Collection, 10801 Univerisity Blvd., Manassas, VA 20110 on August 29, 2000 and assigned ATCC Accession No. PTA 2432.

GlcNAc-phosphotransferase was determined to be a complex of six polypeptides with a subunit structure $\alpha_2\beta_2\gamma_2$. Figure 1 shows a model of the subunit structure obtained from quantitative amino acid sequencing, immunoprecipitation with subunit-specific monoclonal antibodies, SDS-PAGE, and cDNA sequences. The evidence for the model is summarized below. The molecular mass of the complex estimated by gel filtration is 570,000 Daltons. The 166,000 Dalton α -subunit is found as a disulfide-linked homodimer. Likewise, the 51,000 Dalton γ -subunit is found as a disulfide-linked homodimer. Because both the α - and γ -subunits are found in disulfide-linked homodimers, each molecule must contain at least one α - and one γ homodimer. Although the 56,000 Dalton β -subunit is not found in a disulfide-linked homodimer, two independent lines of evidence strongly suggest each complex contains

two β -subunits as well. First, quantitative aminoterminal sequencing demonstrates a 1:1 molar ratio between the β - and γ -subunits. Secondly, since the α - and β -subunits are encoded by a single cDNA and divided by proteolytic processing, two β -subunits are produced for each α -subunit dimer. The predicted mass of the complex based on the composition $\alpha_2\beta_2\gamma_2$ is 546,000 Daltons ($2 \times 166,000 + 2 \times 56,000 + 2 \times 51,000$) in excellent agreement with the mass estimated by gel filtration.

GlcNAc-phosphotransferase was purified using an assay for the transfer of GlcNAc-1-Phosphate to the synthetic acceptor α -methylmannoside. However, the natural acceptors for GlcNAc-phosphotransferase are the high mannose oligosaccharides of lysosomal hydrolases. To evaluate the ability of the purified GlcNAc-phosphotransferase to utilize glycoproteins as acceptors, the transfer of GlcNAc-1-P to the lysosomal enzymes uteroferrin and cathepsin D, the nonlysosomal glycoprotein RNase B, and the lysosomal hydrolase β -glucocerebrosidase (which is trafficked by a M6P independent pathway), were investigated. Both uteroferrin and cathepsin D are effectively utilized as acceptors by purified GlcNAc-phosphotransferase with K_m s below 20 μ m. In contrast, neither RNase B nor β -glucocerebrosidase is an effective acceptor.

The ineffectiveness of RNase B, which contains a single high mannose oligosaccharide, as an acceptor is especially notable since the K_m was not reached at the solubility limit of the protein (at 600 μ m). This data clearly demonstrates the specific phosphorylation of Lysosomal hydrolases previously observed with crude preparations (Waheed, Pohlmann A., R., *et al.* (1982). "Deficiency of UDP-N-acetylglucosamine:lysosomal enzyme N-Acetylglucosamine-1-phosphotransferase in organs of I-Cell patients." *Biochemical and Biophysical Research Communications* 105(3): 1052-10580 is a property of the GlcNAc-phosphotransferase itself.

The α -subunit was identified as containing the UDP-GlcNAc binding site since this subunit was specifically photoaffinity-labeled with [β - ^{32} P]-5-azido-UDP-Glc.

The amino-terminal and internal (tryptic) protein sequence data was obtained for each subunit. N-terminal sequence was obtained from each subunit as follows. Individual subunits of GlcNAc-phosphotransferase were resolved by polyacrylamide gel electrophoresis in the presence of sodium dodecyl sulfate before and after disulfide bond reduction. Subunits were then transferred to a PVDF membrane by electroblotting, identified by Coomassie blue staining, excised, and subjected to N-terminal sequencing. To obtain internal sequence, GlcNAc-phosphotransferase was denatured, reduced, and alkylated, and individual subunits were resolved by gel filtration chromatography. Isolated subunits were then digested with trypsin and the tryptic peptides fractionated by reverse phase HPLC. Peaks which appeared to contain only a single peptide were analyzed for purity by MALDI and subjected to N-terminal amino acid sequencing.

The amino acid sequence for the human α -subunit is shown in amino acids 1-928 of SEQ ID NO: 1; the human β -subunit in amino acids 1-328 of SEQ ID NO:2; and the human γ -subunit in amino acids 25-305 of SEQ ID NO:3. The γ -subunit has a signal sequence shown in amino acids 1-24 of SEQ ID NO:3.

Comparison with the databases using the blast algorithms demonstrate these proteins have not been previously described although several EST sequences of the corresponding cDNAs are present.

Using these peptide sequences and a combination of library screening, RACE, PCR and Blast searching of expressed sequence tag ("EST") files, full-length human cDNAs encoding each subunit were cloned and sequenced.

The nucleotide sequence for the human α/β -subunit precursor cDNA is shown in

nucleotides 165-3932 of SEQ ID NO:4; the nucleotide sequence for the α -subunit is shown in nucleotides 165-2948 of SEQ ID NO:4; the nucleotide sequence for the β -subunit is shown in nucleotides 2949-3932 of SEQ ID NO:4; and the nucleotide sequence for the γ -subunit is shown in nucleotides 96-941 of SEQ ID NO:5. The nucleotide sequence for the γ -subunit signal peptide is shown in nucleotides 24-95 of SEQ ID NO:5.

For each subunit a N-terminal peptide and two internal peptide sequences have been identified in the respective cDNA sequence. Although the protein sequence data is from the bovine protein and the cDNA sequences are human, the sequences are highly homologous (identities: α -subunit 43/50; β -subunit 64/64; γ -subunit 30/32), confirming the cloned cDNAs represent the human homologs of the bovine GlcNAc-phosphotransferase subunits. The α - and β -subunits were found to be encoded by a single cDNA whose gene is on chromosome 12. The γ -subunit is the product of a second gene located on chromosome 16. The α/β -subunits precursor gene has been cloned and sequenced. The gene spans ~80 kb and contains 21 exons. The γ -subunit gene has also been identified in data reported from a genome sequencing effort. The γ -subunit gene is arranged as 11 exons spanning 12 kb of genomic DNA.

Using the human cDNAs, the homologous murine cDNAs for the α -, β - and γ -subunits were isolated and sequenced using standard techniques. The murine α - β -subunit precursor cDNA is shown in SEQ ID NO:16. The deduced amino acid sequence for the murine α -subunit is shown in SEQ ID NO: 15 and the β -subunit in SEQ ID NO:8.

The mouse γ -subunit cDNA was isolated from a mouse liver library in λ Zap II using the γ -human γ -subunit cDNA as a probe. The human γ -subunit cDNA was random hexamer-labeled with ^{32}P -dCTP and used to screen a mouse liver cDNA library in λ Zap II. The probe hybridized to three of 500,000 plaques screened. Each was subcloned to

homogeneity, the insert excised, cloned into pUC19, and sequenced using standard methods. Sarnbrook, J., Fritsch E. F., *et al.* (1989). *Molecular Cloning. A Laboratory Manual.* Cold Spring Harbor, Cold Spring Harbor Laboratory Press. The mouse γ -subunit cDNA sequence is shown in SEQ ID NO:10 and the deduced amino acid sequence for the mouse γ -subunit is shown in SEQ ID NO:9.

Comparison of the deduced amino acid sequences of the human and mouse α -, β -, and γ -subunits demonstrates that the proteins are highly homologous with about an 80 percent identity.

To confirm that these enzymes were substantially the same between species, a partial homologous rat cDNA for the α - and β -subunits was isolated and sequenced using standard techniques. The partial rat α - and β -subunit cDNA is shown in SEQ ID NO:12. The deduced amino acid sequence corresponding to the cDNA is shown in SEQ ID NO:11. Further, a partial homologous *Drosophila* cDNA for the α - and β -subunits was isolated and sequenced using standard techniques. The partial *Drosophila* α - and β -subunit cDNA is shown in SEQ ID NO:17. The deduced amino acid sequence corresponding to the cDNA is shown in SEQ ID NO:13. Comparisons of the deduced amino acid sequences of the partial human, rat, and *Drosophila* α - and β -subunits show that the proteins are highly homologous.

Phosphodiester α -GlcNAcase

In another aspect, the present invention provides isolated and purified biologically active phosphodiester α -GlcNAcase, nucleic acid molecules encoding phosphodiester α -GlcNAcase, expression vectors having a DNA that encodes phosphodiester α -GlcNAcase, host cells that have been transfected or transformed with expression vectors having DNA that encodes phosphodiester α -GlcNAcase, methods for producing recombinant phosphodiester α -

GlcNAcase by culturing host cells that have been transfected or transformed with expression vectors having DNA that encodes phosphodiester α -GlcNAcase, isolated and purified recombinant phosphodiester α -GlcNAcase, and methods for using phosphodiester α -GlcNAcase for the preparation of highly phosphorylated lysosomal enzymes that are useful for the treatment of lysosomal storage diseases.

To obtain isolated and purified phosphodiester α -GlcNAcase and the nucleic acid molecules encoding the enzyme according to the present invention, bovine phosphodiester α -GlcNAcase was obtained and analyzed as follows. Mice were immunized with a partially purified preparation of phosphodiester α -GlcNAcase and a functional screening strategy was utilized to identify and isolate a monoclonal antibody specific for phosphodiester α -GlcNAcase. Immunogen was prepared by partially purifying phosphodiester α -GlcNAcase ~6000-fold from a bovine pancreas membrane pellet using chromatography on DEAE-Sepharose, iminodiacetic acid Sepharose, and Superose 6. Two BALB/c mice were each injected intraperitoneally with 5 μ g partially purified phosphodiester α -GlcNAcase emulsified in Freund's complete adjuvant. On day 28, the mice were boosted intraperitoneally with 5 μ g phosphodiester α -GlcNAcase emulsified in Freund's incomplete adjuvant. On day 42 the mice were bled and an phosphodiester α -GlcNAcase specific immune response was documented by "capture assay." To perform the capture assay, serum (5 μ l) was incubated overnight with 1.2 units partially purified phosphodiester α -GlcNAcase. Mouse antibody was then captured on rabbit antimouse IgG bound to protein A-Ultralink™ resin. Following extensive washing, bound phosphodiester α -GlcNAcase was determined in the Ultralink pellet by assay of cleavage of [^{3} H]-GlcNAc-1-phosphomannose α -methyl.

Following a second intravenous boost with phosphodiester α -GlcNAcase, the spleen was removed and splenocytes fused with SP2/0 myeloma cells according to our modifications

(Bag, M., Booth J. L., *et al.* (1996). "Bovine UDP-N-acetylglucosamine: lysosomal enzyme N-acetylglucosamine-1-phosphotransferase. I. Purification and subunit structure." *Journal of Biological Chemistry* 271: 31437 - 31445) of standard techniques; Harlow, E. and Lane, D. (1988). *Antibodies: a laboratory manual*, Cold Spring Harbor Laboratory). The fusion was plated in eight 96-well plates in media supplemented with recombinant human IL-6 (Bazin, R. and Lemieux, R. (1989). "Increased proportion of B cell hybridomas secreting monoclonal antibodies of desired specificity in cultures containing macrophage-derived hybridoma growth factor (IL-6)." *Journal of Immunological Methods* 116: 245 - 249); and grown until hybridomas were just visible. Forty-eight pools of 16-wells were constructed and assayed for antiphosphodiester α -GlcNAcase activity using the capture assay. Four pools were positive. Subpools of 4-wells were then constructed from the wells present in the positive 16-well pools. Three of the four 16-well pools contained a single 4-well pool with anti-phosphodiester α -GlcNAcase activity. The 4 single wells making up the 4-well pools were then assayed individually identifying the well containing the anti-phosphodiester α -GlcNAcase secreting hybridomas. Using the capture assay, each hybridoma was subcloned twice and antibody prepared by ascites culture. Monoclonals UC2 and UC3 were found to be low affinity antibodies. UC1, a high affinity IgG monoclonal antibody, was prepared by ascites culture and immobilized on Emphaze for purification of phosphodiester α -GlcNAcase. The monoclonal antibody labeled UC1 was selected for use in further experiments. A hybridoma secreting monoclonal antibody UC1 was deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110 on August 29, 2000 and assigned ATCC Accession No. PTA 2431.

To purify phosphodiester α -GlcNAcase, a solubilized membrane fraction was prepared from bovine liver. Phosphodiester α -GlcNAcase was absorbed to monoclonal

antibody UC1 coupled to Emphaze resin by incubation overnight with gentle rotation. The UC1-Emphaze was then packed in a column, washed sequentially with EDTA and NaHCO₃ at pH 7.0, then phosphodiester α -GlcNAcase was eluted with NaHCO₃ at pH 10. Fractions containing phosphodiester α -GlcNAcase at specific activities >50,000 μ /mg were pooled and adjusted to pH 8.0 with 1/5th volume of 1 M Tris HCl, pH 7.4. Following chromatography on UCI-Emphaze the phosphodiester α -GlcNAcase was purified 92,500-fold in 32% yield.

The phosphodiester α -GlcNAcase from UC1-Emphaze was concentrated and chromatographed on Superose 6. Phosphodiester α -GlcNAcase eluted early in the chromatogram as a symmetric activity peak with a coincident protein peak. Following chromatography on Superose 6, the enzyme was purified ~715,000-fold in 24% yield. The purified enzyme catalyzed the cleavage of 472 μ mol/hr/mg [³H]-GlcNAc-1-phosphomannose- α -methyl, corresponding to a specific activity of 472,000 units/mg.

The purified phosphodiester α -GlcNAcase was subjected to SDS-PAGE and protein was detected by silver staining (Blum, H., Beier H., *et al.* (1987). "Improved silver staining of plant proteins, RNA and DNA in polyacrylamide gels." *Electrophoresis*: 93-99). A diffuse band was observed with a molecular mass of approximately 70 kDa whose intensity varies with the measured phosphodiester α -GlcNAcase activity. The diffuse appearance of the band suggests the protein may be heavily glycosylated. A faint band with a molecular mass of ~150,000, which does not correlate with activity, was also present.

A model for the subunit structure of phosphodiester α -GlcNAcase was determined by gel filtration chromatography and SDS-PAGE with and without disulfide bond reduction. The mass by gel filtration is about 300,000. SDS-PAGE without disulfide bond reduction is ~140,000. Following disulfide bond reduction, the apparent mass is 70,000. Together these data show phosphodiester α -GlcNAcase is a tetramer composed of disulfide linked

homodimers. Figure 2 shows a model of the subunit structure of phosphodiester α -GlcNAcase.

The amino terminal amino acid sequence of affinity purified, homogeneous bovine phosphodiester α -GlcNAcase was determined using standard methods (Matsudaira, P., Ed. (1993). *A Practical Guide to Protein and Peptide Purification for Microsequencing*. San Diego, Academic Press, Inc.). The pure enzyme was also subjected to trypsin digestion and HPLC to generate two internal tryptic peptides which were sequenced. The amino acid sequences of these three peptides are:

Peptide 1 - Amino Terminal DXTRVHAGRLEHESWPPAAQTAGAHGPSVRFTFV (SEQ ID NO:23);

Peptide 2 - Tryptic RDGTLVTGYLSEEEVLDTEN (SEQ ID NO:24); and

Peptide 3 - Tryptic GINLWEMAEFLK (SEQ ID NO:25).

The protein, nucleotide, and EST data bases were searched for sequences that matched these peptide sequences and several human and mouse ESTs were found that had the sequence of the third peptide at their amino termini. Three human infant brain EST clones and one mouse embryo clone were obtained from ATCC and sequenced. The three human clones were all identical except for total length at their 3' ends and virtually identical to the mouse clone, except that the mouse EST contained a 102 bp region that was absent from all three human brain ESTs. An EcoR I -Hind III fragment of about 700 bp was excised from the human cDNA clone (ATCC # 367524) and used to probe a human liver cDNA library directionally cloned in TriplEx vector (Clontech). Of the positive clones isolated from the library and converted to plasmids (pTriplEx), the largest (2200 bp) was represented by clone 6.5 which was used for the rest of the analysis.

The cDNA clone has been completely sequenced on both strands and is a novel

sequence that predicts a mature protein of about 50 kDa which is in agreement with the size of the deglycosylated mature bovine liver phosphodiester α -GlcNAcase.

There is a unique BamH I site at base #512 and a unique Hind ID site at base # 1581. All three bovine peptide sequences (peptides 1, 2, and 3) were found. Although the sequences of peptides 2 and 3 in the human are 100% identical to the bovine sequences, the amino-terminal peptide in humans is only 67% identical to the bovine sequence. The human liver clone contains the 102 base pair insert that has the characteristics of an alternatively spliced segment that was missing in the human brain EST. The hydrophilicity plot indicates the presence of a hydrophobic membrane spanning region from amino acids 448 to 474 and another hydrophobic region from amino acid 8 to 24 which fits the motif for a signal sequence and there is a likely signal sequence cleavage site between G24 and G25. There are six Asn-X-Ser/Thr potential N-linked glycosylation sites, one of which is within the 102 bp insert. All of these sites are amino terminal of the putative trans-membrane region. These features indicate that the phosphodiester α -GlcNAcase is a type I membrane spanning glycoprotein with the amino terminus in the lumen of the Golgi and the carboxyl terminus in the cytosol. This orientation is different from that of other glycosyltransferases and glycosidases involved in glycoprotein processing, which to date have been shown to be type II membrane spanning proteins.

The amino acid sequence for the phosphodiester α -GlcNAcase monomer is shown in amino acids 50-515 of SEQ ID NO:6. The signal peptide is shown in amino acids 1-24 of SEQ ID NO:6 and the pro segment is shown in amino acids 25-49 of SEQ ID NO:6. The human cDNA was cloned using the techniques described above. The nucleotide sequence for the monomer that associates to form the phosphodiester α -GlcNAcase tetramer is shown in nucleotides 151-1548 of SEQ ID NO:7. The nucleotide sequence for the signal sequence is

shown in nucleotides 1-72 of SEQ ID NO:7. The nucleotide sequence for the propeptide is shown in nucleotides 73-150 of SEQ ID NO:7.

The murine cDNA for phosphodiester α -GlcNAcase is shown in SEQ ID NO:18. The deduced amino acid sequence for the murine phosphodiester α -GlcNAcase is shown in SEQ ID NO:19. Comparison of the deduced amino acid sequences of the human and mouse enzymes demonstrates that the proteins are highly homologous with about an 80 percent identity. This is especially true in the region of the active site where identity exceeds 90%. The murine gene for phosphodiester α -GlcNAcase is shown in SEQ ID NO:14.

The human phosphodiester α -GlcNAcase gene has been identified by database searching. The sequence was determined during the sequencing of clone l65E7 from chromosome 16.13.3, GenBank AC007011.1, gi4371266. Interestingly, the phosphodiester α -GlcNAcase gene was not identified by the SCAN program used to annotate the sequence.

Because of the degeneracy of the genetic code, a DNA sequence may vary from that shown in SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:7 and still encode a GlcNAc phosphotransferase and a phosphodiester α -GlcNAcase enzyme having the amino acid sequence shown in SEQ ID NO: 1, SEQ ID NO:2, SEQ ID NO:3, and SEQ ID NO:6. Such variant DNA sequences may result from silent mutations, e.g., occurring during PCR amplification, or may be the product of deliberate mutagenesis of a native sequence. The invention, therefore, provides equivalent isolated DNA sequences encoding biologically active GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase selected from: (a) the coding region of a native mammalian GlcNAc-phosphotransferase gene and phosphodiester α -GlcNAcase gene; (b) cDNA comprising the nucleotide sequence presented in SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:7; (c) DNA capable of hybridization to the native mammalian GlcNAc-phosphotransferase gene and phosphodiester α -GlcNAcase gene under

moderately stringent conditions and which encodes biologically active GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase; and (d) DNA which is degenerate as a result of the genetic code to a DNA defined in (a), (b), or (c) and which encodes biologically active GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase. GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase proteins encoded by such DNA equivalent sequences are encompassed by the invention.

Those sequences which hybridize under stringent conditions and encode biologically functional GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase are preferably at least 50-100% homologous, which includes 55, 60, 65, 70, 75, 75, 80, 85, 90, 95, 99% and all values and subranges therebetween. Homology may be determined with the software UWCG as described above. Stringent hybridization conditions are known in the art and are meant to include those conditions which allow hybridization to those sequences with a specific homology to the target sequence. An example of such stringent conditions are hybridization at 65°C in a standard hybridization buffer and subsequent washing in 0.2 X concentrate SSC and 0.1% SDS at 42-65°C, preferably 60°C. This and other hybridization conditions are disclosed in Sambrook, J., Fritsch E. F., *et al.* (1989). *Molecular Cloning. A Laboratory Manual*. Cold Spring Harbor, Cold Spring Harbor Laboratory Press. Alternatively, the temperature for hybridization conditions may vary dependent on the percent GC content and the length of the nucleotide sequence, concentration of salt in the hybridization buffer and thus the hybridization conditions may be calculated by means known in the art.

Recombinant Expression for GlcNAc-phosphotransferase and Phosphodiester α -GlcNAcase Isolated and purified recombinant GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase enzymes are provided according to the present invention by

incorporating the DNA corresponding to the desired protein into expression vectors and expressing the DNA in a suitable host cell to produce the desired protein.

Expression Vectors

Recombinant expression vectors containing a nucleic acid sequence encoding the enzymes can be prepared using well known techniques. The expression vectors include a DNA sequence operably linked to suitable transcriptional or translational regulatory nucleotide sequences such as those derived from mammalian, microbial, viral, or insect genes. Examples of regulatory sequences include transcriptional promoters, operators, enhancers, mRNA ribosomal binding sites, and appropriate sequences which control transcription and translation initiation and termination. Nucleotide sequences are "operably linked" when the regulatory sequence functionally relates to the DNA sequence for the appropriate enzyme. Thus, a promoter nucleotide sequence is operably linked to a GlcNAc-phosphotransferase or phosphodiesterase α -GlcNAcase DNA sequence if the promoter nucleotide sequence controls the transcription of the appropriate DNA sequence.

The ability to replicate in the desired host cells, usually conferred by an origin of replication and a selection gene by which transformants are identified, may additionally be incorporated into the expression vector.

In addition, sequences encoding appropriate signal peptides that are not naturally associated with GlcNAc-phosphotransferase or phosphodiesterase α -GlcNAcase can be incorporated into expression vectors. For example, a DNA sequence for a signal peptide (secretory leader) may be fused in-frame to the enzyme sequence so that the enzyme is initially translated as a fusion protein comprising the signal peptide. A signal peptide that is functional in the intended host cells enhances extracellular secretion of the appropriate

polypeptide. The signal peptide may be cleaved from the polypeptide upon secretion of enzyme from the cell.

Host Cells

Suitable host cells for expression of GlcNAc-phosphotransferase and phosphodiester at α -GlcNAcase include prokaryotes, yeast, archae, and other eukaryotic cells. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are well known in the art, *e.g.*, Pouwels *et al.* Cloning Vectors: A Laboratory Manual, Elsevier, New York (1985). The vector may be a plasmid vector, a single or double-stranded phage vector, or a single or double-stranded RNA or DNA viral vector. Such vectors may be introduced into cells as polynucleotides, preferably DNA, by well known techniques for introducing DNA and RNA into cells. The vectors, in the case of phage and viral vectors also may be and preferably are introduced into cells as packaged or encapsulated virus by well known techniques for infection and transduction. Viral vectors may be replication competent or replication defective. In the latter case viral propagation generally will occur only in complementing host cells. Cell-free translation systems could also be employed to produce the enzymes using RNAs derived from the present DNA constructs.

Prokaryotes useful as host cells in the present invention include gram negative or gram positive organisms such as *E. coli* or *Bacilli*. In a prokaryotic host cell, a polypeptide may include a N-terminal methionine residue to facilitate expression of the recombinant polypeptide in the prokaryotic host cell. The N-terminal Met may be cleaved from the expressed recombinant GlcNAc-phosphotransferase or phosphodiester α -GlcNAcase polypeptide. Promoter sequences commonly used for recombinant prokaryotic host cell expression vectors include β -lactamase and the lactose promoter system.

Expression vectors for use in prokaryotic host cells generally comprise one or more phenotypic selectable marker genes. A phenotypic selectable marker gene is, for example, a gene encoding a protein that confers antibiotic resistance or that supplies an autotrophic requirement. Examples of useful expression vectors for prokaryotic host cells include those derived from commercially available plasmids such as the cloning vector pBR322 (ATCC 37017). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells. To construct an expression vector using pBR322, an appropriate promoter and a DNA sequence are inserted into the pBR322 vector.

Other commercially available vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, Wisconsin, USA).

Promoter sequences commonly used for recombinant prokaryotic host cell expression vectors include β -lactamase (penicillinase), lactose promoter system (Chang *et al.*, Nature 275:615, (1978); and Goeddel *et al.*, Nature 281:544, (1979)), tryptophan (trp) promoter system (Goeddel *et al.*, Nucl. Acids Res. 8:4057, (1980)), and tac promoter (Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, p. 412 (1982)).

Yeasts useful as host cells in the present invention include those from the genus *Saccharomyces*, *Pichia*, *K. Actinomyces* and *Kluyveromyces*. Yeast vectors will often contain an origin of replication sequence from a 2 μ yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Suitable promoter sequences for yeast vectors include, among others, promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman *et al.*, J. Biol. Chem. 255:2073, (1980)) or other glycolytic enzymes

(Holland *et al.*, Biochem. 17:4900, (1978)) such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Other suitable vectors and promoters for use in yeast expression are further described in Fleer *et al.*, Gene, 107:285-195 (1991). Other suitable promoters and vectors for yeast and yeast transformation protocols are well known in the art.

Yeast transformation protocols are known to those of skill in the art. One such protocol is described by Hinnen *et al.*, *Proceedings of the National Academy of Sciences USA*, 75:1929 (1978). The Hinnen protocol selects for Trp.^{sup.}+ transformants in a selective medium, wherein the selective medium consists of 0.67% yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10 µg/ml adenine, and 20 µg/ml uracil.

Mammalian or insect host cell culture systems well known in the art could also be employed to express recombinant GlcNAc-phosphotransferase or phosphodiester α -GlcNAcase polypeptides, *e.g.*, Baculovirus systems for production of heterologous proteins in insect cells (Luckow and Summers, Bio/Technology 6:47 (1988)) or Chinese hamster ovary (CHO) cells for mammalian expression may be used. Transcriptional and translational control sequences for mammalian host cell expression vectors may be excised from viral genomes. Commonly used promoter sequences and enhancer sequences are derived from Polyoma virus, Adenovirus 2, Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome may be used to provide other genetic elements for expression of a structural gene sequence in a mammalian host cell, *e.g.*, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites. Viral early and late promoters are particularly useful because both are easily obtained from a viral genome as

a fragment which may also contain a viral origin of replication. Exemplary expression vectors for use in mammalian host cells are well known in the art.

The enzymes of the present invention may, when beneficial, be expressed as a fusion protein that has the enzyme attached to a fusion segment. The fusion segment often aids in protein purification, *e.g.* by permitting the fusion protein to be isolated and purified by affinity chromatography. Fusion proteins can be produced by culturing a recombinant cell transformed with a fusion nucleic acid sequence that encodes a protein including the fusion segment attached to either the carboxyl and/or amino terminal end of the enzyme. Preferred fusion segments include, but are not limited to, glutathione-S-transferase, β -galactosidase, a poly-histidine segment capable of binding to a divalent metal ion, and maltose binding protein. In addition, the HPC-4 epitope purification system may be employed to facilitate purification of the enzymes of the present invention. The HPC-4 system is described in U.S. Patent No. 5,202,253.

Expression by gene activation technology

In addition to expression strategies involving transfection of a cloned cDNA sequence, the endogenous GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase genes can be expressed by altering the promoter.

Methods of producing the enzymes of the present invention can also be accomplished according to the methods of protein production as described in U.S. Patent No. 5,968,502, using the sequences for GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase as described herein.

Expression and Recovery

According to the present invention, isolated and purified GlcNAc-phosphotransferase or phosphodiester α -GlcNAcase enzymes may be produced by the recombinant expression systems described above. The method comprises culturing a host cell transformed with an expression vector comprising a DNA sequence that encodes the enzyme under conditions sufficient to promote expression of the enzyme. The enzyme is then recovered from culture medium or cell extracts, depending upon the expression system employed. As is known to the skilled artisan, procedures for purifying a recombinant protein will vary according to such factors as the type of host cells employed and whether or not the recombinant protein is secreted into the culture medium. When expression systems that secrete the recombinant protein are employed, the culture medium first may be concentrated. Following the concentration step, the concentrate can be applied to a purification matrix such as a gel filtration medium. Alternatively, an anion exchange resin can be employed, *e.g.*, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose, or other types commonly employed in protein purification. Also, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Further, one or more reversed-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media (*e.g.*, silica gel having pendant methyl or other aliphatic groups) can be employed to further purify the enzyme. Some or all of the foregoing purification steps, in various combinations, are well known in the art and can be employed to provide an isolated and purified recombinant protein.

Recombinant protein produced in bacterial culture is usually isolated by initial disruption of the host cells, centrifugation, extraction from cell pellets if an insoluble

polypeptide, or from the supernatant fluid if a soluble polypeptide, followed by one or more concentration, salting-out, ion exchange, affinity purification, or size exclusion chromatography steps. Finally, RP-HPLC can be employed for final purification steps. Microbial cells can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Preparation of Highly Phosphorylated Lysosomal Enzymes

In another aspect, the present invention provides highly phosphorylated lysosomal hydrolases and methods for the preparation of such hydrolases. The highly phosphorylated lysosomal hydrolases can be used in clinical applications for the treatment of lysosomal storage diseases.

The method comprises obtaining lysosomal hydrolases having asparagine-linked oligosaccharides with high mannose structures and modifying the α 1,2-linked or other outer mannoses by the addition of M6P *in vitro* to produce a hydrolase that can be used for the treatment of lysosomal storage diseases because it binds to cell membrane M6P receptors and is readily taken into the cell and into the lysosome. Typically, the high mannose structures consist of from six to nine molecules of mannose and two molecules of N-acetylglucosamine (GlcNAc). In the preferred embodiment, the high mannose structure is a characteristic MAN7(D₂D₃) isomer structure consisting of seven molecules of mannose and two molecules of N-acetylglucosamine (GlcNAc).

Highly phosphorylated Lysosomal hydrolases are produced by treating the high mannose hydrolases with GlcNAc-phosphotransferase which catalyzes the transfer of N-acetylglucosamine-1-phosphate from UDP-GlcNAc to the 6' position of α 1,2-linked or other outer mannoses on the hydrolase. This GlcNAc-phosphotransferase modified hydrolase is

then treated with phosphodiester α -GlcNAcase which catalyzes the removal of N-Acetylglucosamine to generate terminal M6P on the hydrolase.

In one embodiment of the invention, the GlcNAc-phosphotransferase treated hydrolase may be isolated and stored without any subsequent treatment. Subsequently, the GlcNAc-phosphotransferase treated hydrolase may be modified further by treating the hydrolase with a phosphodiester α -GlcNAcase.

Surprisingly, it has been found that the hydrolases containing M6P generated by this method are highly phosphorylated when compared to naturally occurring or known recombinant hydrolases. The highly phosphorylated lysosomal hydrolases of the present invention contain from about 6% to about 100% bis-phosphorylated oligosaccharides compared to less than about 5% bis-phosphorylated oligosaccharides on known naturally occurring or recombinant hydrolases.

These highly phosphorylated hydrolases have a higher affinity for the M6P receptor and are therefore more efficiently taken into the cell by plasma membrane receptors. (Reuser, A. J., Kroos, M. A., Ponne, N. J., Wolterman, R. A., Loonen, M. C., Busch, H. F., Visser, W. J., and Bolhuis, P. A. (1984). "Uptake and stability of human and bovine acid alpha-glucosidase in cultured fibroblasts and skeletal muscle cells from glycogenosis type II patients." *Experimental Cell Research* 155: 178-189).

The high-affinity ligand for the cation-independent M6P receptor is an oligosaccharide containing two M6P groups (*i.e.*, a bis-phosphorylated oligosaccharide). Since a bisphosphorylated oligosaccharides binds with an affinity 3500-fold higher than a monophosphorylated oligosaccharides, virtually all the high-affinity binding of a lysosomal enzyme to the M6P receptor will result from the content of bis-phosphorylated oligosaccharides (Tong, P. Y., Gregory, W., and Kornfeld, S. (1989)). "Ligand interactions

of the cation-independent mannose 6-phosphate receptor. The stoichiometry of mannose 6-phosphate binding." *Journal of Biological Chemistry* 264: 7962-7969). It is therefore appropriate to use the content of bis-phosphorylated oligosaccharides to compare the binding potential of different preparations of lysosomal enzymes.

The extent of mannose 6-phosphate modification of two different lysosomal enzymes has been published. The oligosaccharide composition of human α -galactosidase A secreted from Chinese hamster ovary cells has been published (Matsuura, F., Ohta, M., Ioannou, Y. A., and Desnick, R. I. (1998). "Human alpha-galactosidase A: characterization of the N-linked oligosaccharides on the intracellular and secreted glycoforms overexpressed by Chinese hamster ovary cells." *Glycobiology* 8(4): 329-39). Of all oligosaccharides on α -gal A released by hydrazinolysis, only 5.2% were bis-phosphorylated. Zhao *et al.* partially characterized the oligosaccharide structures on recombinant human α -iduronidase secreted by CHO cells (Zhao, K. W., Faull, K. F., Kakkis, E. D., and Neufeld, E. F. (1997). "Carbohydrate structures of recombinant human alpha-L-iduronidase secreted by Chinese hamster ovary cells." *J Biol Chem* 272(36): 22758-65) and demonstrated a minority of the oligosaccharides were bisphosphorylated. The qualitative techniques utilized precluded the determination of the fraction of oligosaccharides phosphorylated.

The production and secretion of human acid α -glucosidase by CHO cells has been reported (Van Hove, J. L., Yang, H. W., Wu, J. Y., Brady, R. O., and Chen, Y. T. (1996). "High level production of recombinant human lysosomal acid alpha-glucosidase in Chinese hamster ovary cells which targets to heart muscle and corrects glycogen accumulation in fibroblasts from patients with Pompe disease." *Proceedings of the National Academy of Sciences USA*, 93(1): 6570). The carbohydrate structures of this preparation were not characterized in this publication. However, this preparation was obtained and analyzed. The

results, given in the examples below, showed that less than 1% of the oligosaccharides contained any M6P and bis-phosphorylated oligosaccharides were not detectable. Together, these data show that known preparations of recombinant lysosomal enzymes contain no more than 5.2% phosphorylated oligosaccharides. It appears that the preparation of more highly phosphorylated lysosomal enzymes is unlikely to be achieved with known techniques. Naturally occurring human acid α -glucosidase purified from human placenta contains very low levels of M6P (Mutsaers, I. H. G. M., Van Halbeek, H., Vliegenthart, J. F. G., Tager, J. M., Reuser, A. J. J., Kroos, M., and Galjaard, H. (1987). "Determination of the structure of the carbohydrate chains of acid α -glucosidase from human placenta." *Biochimica et Biophysica Acta* 911: 244 - 251). The arrangement of the phosphates as either bis- or monophosphorylated oligosaccharides has not been determined, but less than 1% of the oligosaccharides contain any M6P.

The highly phosphorylated hydrolases of the present invention are useful in enzyme replacement therapy procedures because they are more readily taken into the cell and the lysosome. (Reuser, A. J., Kroos, M. A., Ponne, N. J., Wolterman, R. A., Loonen, M. C., Busch, H. F., Visser, W. J. and Bolhuis, P. A. (1984). "Uptake and stability of human and bovine acid alpha-glucosidase in cultured fibroblasts and skeletal muscle cells from glycogenosis type II patients." *Experimental Cell Research* 155: 178-189).

Any lysosomal enzyme that uses the M6P transport system can be treated according to the method of the present invention. Examples include α -glucosidase (Pompe Disease), α -L-iduronidase (Hurler Syndrome), α -galactosidase A (Fabry Disease), arylsulfatase (Maroteaux-Lamy Syndrome), N-acetylgalactosamine-6-sulfatase or β -galactosidase (Morquio Syndrome), iduronate 2-sulfatase (Hunter Syndrome), ceramidase (Farber Disease), galactocerebrosidase (Krabbe Disease), β -glucuronidase (Sly Syndrome), Heparan N-

sulfatase (Sanfilippo A), N-Acetyl- α -glucosaminidase (Sanfilippo B), Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase (Sanfilippo D), Galactose 6-sulfatase (Morquio A), Arylsulfatase A, B, and C (Multiple Sulfatase Deficiency), Arylsulfatase A Cerebroside (Metachromatic Leukodystrophy), Ganglioside (Mucopolidosis IV), Acid β -galactosidase G_{MI} Galgliside (G_{MI} Gangliosidosis), Acid β -galactosidase (Galactosialidosis), Hexosaminidase A (Tay-Sachs and Variants), Hexosaminidase B (Sandhoff), α -fucosidase (Fucosidosis), α -N-Acetyl galactosaminidase (Schindler Disease), Glycoprotein Neuraminidase (Sialidosis), Aspartylglucosamine amidase (Aspartylglucosaminuria), Acid Lipase (Wolman Disease), Acid Ceramidase (Farber Lipogranulomatosis), Lysosomal Sphingomyelinase and other Sphingomyelinase (Niemann-Pick).

Methods for treating any particular lysosomal hydrolase with the enzymes of the present invention are within the skill of the artisan. Generally, the lysosomal hydrolase at a concentration of about 10 mg/ml and GlcNAc-phosphotransferase at a concentration of about 100,000 units/mL are incubated at about 37°C for 2 hours in the presence of a buffer that maintains the pH at about 6-7 and any stabilizers or coenzymes required to facilitate the reaction. Then, phosphodiester α -GlcNAcase is added to the system to a concentration of about 1000 units/mL and the system is allowed to incubate for about 2 more hours. The modified lysosomal enzyme having highly phosphorylated oligosaccharides is then recovered by conventional means.

In a preferred embodiment, the lysosomal hydrolase at 10 mg/ml is incubated in 50 mM Tris-HCl, pH 6.7, 5 mM $MgCl_2$, 5 mM $MnCl_2$, 2 mM UDP-GlcNAc with GlcNAc phosphotransferase at 100,000 units/mL at 37°C for 2 hours. Phosphodiester α -GlcNAcase, 1000 units/mL, is then added and the incubation continued for another 2 hours. The modified

enzyme is then repurified by chromatography on Q-Sepharose and step elution with NaCl.

Methods for Obtaining High Mannose Lysosomal Hydrolases

High mannose lysosomal hydrolases for treatment according to the present invention can be obtained from any convenient source, e.g., by isolating and purifying naturally occurring enzymes or by recombinant techniques for the production of proteins. High mannose lysosomal hydrolases can be prepared by expressing the DNA encoding a particular hydrolase in any host cell system that generates a oligosaccharide modified protein having high mannose structures, e.g., yeast cells, insect cells, other eukaryotic cells, transformed Chinese Hamster Ovary (CHO) host cells, or other mammalian cells.

In one embodiment, high mannose lysosomal hydrolases are produced using mutant yeast that are capable of expressing peptides having high mannose structures. These yeast include the mutant *S. cerevisiae* $\Delta ochl$, $\Delta mnnl$ (Nakanishi-Shindo, Y., Nakayama, K. I., Tanaka, A., Toda, Y. and Jigami, Y. (1993). "Structure of the N-linked oligosaccharides that show the complete loss of α -1,6-polymannose outer chain from *ochl*, *ochl mnnl*, and *ochl mnnl alg3* mutants of *Saccharomyces cerevisiae*." *Journal of Biological Chemistry* 268: 26338 - 26345).

Preferably, high mannose lysosomal hydrolases are produced using over-expressing transformed insect, CHO, or other mammalian cells that are cultured in the presence of certain inhibitors. Normally, cells expressing lysosomal hydrolases secrete acid α -glucosidase that contains predominantly sialylated biantennary complex type glycans that do not serve as a substrate for GlcNAc-phosphotransferase and therefore cannot be modified to use the M6P receptor.

According to the present invention, a new method has been discovered for

manipulating transformed cells containing DNA that expresses a recombinant hydrolase so that the cells secrete high mannose hydrolases that can be modified according to the above method. In this method, transformed cells are cultured in the presence of α 1,2-mannosidase inhibitors and the high mannose recombinant hydrolases are recovered from the culture medium. Inhibiting alpha 1,2-mannosidase prevents the enzyme from trimming mannoses and forces the cells to secrete glycoproteins having the high mannose structure. High mannose hydrolases are recovered from the culture medium using known techniques and treated with GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase according to the method herein to produce hydrolases that have M6P and can therefore bind to membrane M6P receptors and be taken into the cell. Preferably, the cells are CHO cells and the hydrolases are secreted with the MAN7(D₂D₃) structure. Figure 3 shows the reaction scheme for this method.

In a preferred embodiment, recombinant human acid alpha glucosidase ("rh-GAA") is prepared by culturing CHO cells secreting rh-GAA in Iscove's Media modified by the addition of an alpha 1,2-mannosidase inhibitor. Immunoprecipitation of rh-GAA from the media followed by digestion with either N-glycanase or endoglycosidase-H demonstrates that in the presence of the alpha 1,2-mannosidase inhibitor the rh-GAA retains high mannose structures rather than the complex structures found on a preparation secreted in the absence of the inhibitor. The secreted rh-GAA bearing high mannose structures is then purified to homogeneity, preferably by chromatography beginning with ion exchange chromatography on ConA-Sepharose, Phenyl-Sepharose and affinity chromatography on Sephadex G-100. The purified rh-GAA is then treated *in vitro* with GlcNAc-phosphotransferase to convert specific mannoses to GlcNAc-phospho-mannose diesters. The GlcNAcphosphomannose diesters are then converted to M6P groups by treatment with phosphodiester α GlcNAcase. Experiments

show that 74% of the rh-GAA oligosaccharides were phosphorylated, 62% being bis-phosphorylated, and 12% monophosphorylated. Since each molecule of rh-GAA contains 7 N-linked oligosaccharides, 100% of the rh-GAA molecules are likely to contain the mannose-phosphate modification.

Any alpha 1,2-mannosidase inhibitor can function in the present invention. Preferably, the inhibitor is selected from the group consisting of deoxymannojirimycin (dMM), kifunensine, D-Mannonolactam amidrazone, and N-butyl-deoxymannojirimycin. Most preferably the inhibitor is deoxymannojimycin.

Treatment of Lysosomal Storage Diseases

In a further aspect, the present invention provides a method for the treatment of lysosomal storage diseases by administering a disease treating amount of the highly phosphorylated lysosomal hydrolases of the present invention to a patient suffering from the corresponding lysosomal storage disease. While dosages may vary depending on the disease and the patient, the enzyme is generally administered to the patient in amounts of from about 0.1 to about 1000 milligrams per 50 kg of patient per month, preferably from about 1 to about 500 milligrams per 50 kg of patient per month. The highly phosphorylated enzymes of the present invention are more efficiently taken into the cell and the lysosome than the naturally occurring or less phosphorylated enzymes and are therefore effective for the treatment of the disease. Within each disease, the severity and the age at which the disease presents may be a function of the amount of residual lysosomal enzyme that exists in the patient. As such, the present method of treating lysosomal storage diseases includes providing the highly phosphorylated lysosomal hydrolases at any or all stages of disease progression.

The lysosomal enzyme is administered by any convenient means. For example, the enzyme can be administered in the form of a pharmaceutical composition containing the

enzyme and any pharmaceutically acceptable carriers or by means of a delivery system such as a liposome or a controlled release pharmaceutical composition. The term "pharmaceutically acceptable" refers to molecules and compositions that are physiologically tolerable and do not typically produce an allergic or similar unwanted reaction such as gastric upset or dizziness when administered. Preferably, "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals, preferably humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the compound is administered. Such pharmaceutical carriers can be sterile liquids, such as saline solutions, dextrose solutions, glycerol solutions, water and oils emulsions such as those made with oils of petroleum, animal, vegetable, or synthetic origin (peanut oil, soybean oil, mineral oil, or sesame oil). Water, saline solutions, dextrose solutions, and glycerol solutions are preferably employed as carriers, particularly for injectable solutions.

The enzyme or the composition can be administered by any standard technique compatible with enzymes or their compositions. For example, the enzyme or composition can be administered parenterally, transdermally, or transmucosally, *e.g.*, orally or nasally. Preferably, the enzyme or composition is administered by intravenous injection.

The following Examples provide an illustration of embodiments of the invention and should not be construed to limit the scope of the invention which is set forth in the appended claims. In the following Examples, all methods described are conventional unless otherwise specified.

EXAMPLES

Materials and Methods

Lactating bovine udders were obtained from Mikkelson Beef, Inc. (Oklahoma City, OK). Ultrasphere ODS columns were obtained from Beckman Instruments. Microsorb MV-NH₂ columns were obtained from Rainin Instrument Co., Inc. (Woburn, MA). [γ ³²P]ATP (7000 Ci/mmol; end labeling grade), Na¹²⁵I, and Lubrol (C₁₆H₃₃(CH₂CH₂O)₂₃H) were obtained from ICN (Costa Mesa, CA). Superose 6 (prep grade), DEAE-Sepharose FF, QAE-Sephadex A-25, molecular mass standards for SDS-PAGE, HiTrap-protein G columns, and Mono Q columns were obtained from Pharmacia Biotech Inc. 3M-Emphaze Biosupport Medium AB1, IODO GEN iodination reagent, and the BCA protein assay reagent were obtained from Pierce. Glycerol, sucrose, α -methylmannoside, α -methylglucoside, reactive green 19-agarose, sodium deoxycholate, benzamidine, UDP-GlcNAc, phenylmethylsulfonyl fluoride, Tris, rabbit anti-mouse IgG, and mouse monoclonal antibody isotyping reagents were obtained from Sigma.

POROS 50 HQ was obtained from PerSeptive Biosystems (Cambridge, MA). ProBlott polyvinylidene difluoride membranes were obtained from Applied Biosystems Inc. (Foster City, CA). A Model QT12 rotary tumbler was obtained from LORTONE, Inc. (Seattle, WA). A mouse immunoglobulin standard panel was obtained from Southern Biotechnology Associates, Inc. (Birmingham, AL). Recombinant interleukin-6, porcine uteroferrin, and monoclonal antibody BP95 were gifts from colleagues. Other chemicals were reagent grade or better and were from standard suppliers.

Example 1

Preparation of monoclonal antibodies specific for bovine GlcNAc-phosphotransferase

Bovine GlcNAc-phosphotransferase was partially purified 30,000 fold as described (Bao, M., Booth J. L., *et al.* (1996). "Bovine UDP-N-acetylglucosamine: Lysosomal enzyme N-acetylglucosamine-1-phosphotransferase. I. Purification and subunit structure." *Journal of*

Biological Chemistry 271: 31437 - 31445) and used to immunize mice. Spleens of immune mice were removed and splenocytes fused with SP2/0 myeloma cells according to Harlow (Harrow, E. and Lane, D. (1988). *Antibodies: a laboratory manual*, Cold Spring Harbor Laboratory). The fusion was plated into 96 well plates and cultured in HAT media until hybridomas were visible.

Hybridomas secreting monoclonal antibodies capable of capturing GlcNAc-phosphotransferase from a crude sample were identified by incubation of hybridoma media (200 μ l) with 200 units. Partially purified GlcNAc-phosphotransferase and capturing the resulting immune complex on rabbit anti-mouse IgG bound to protein A coupled to UltralinkTM matrix. Immune complexes which contained monoclonal antibodies directed against GlcNAc-phosphotransferase were then identified by assay of the immune complex for GlcNAc-phosphotransferase activity. By this strategy, four monoclonals directed against GlcNAc-phosphotransferase were identified in the fifth fusion screened. The hybridomas identified were subcloned twice using the same assay and ascites was produced in BALBc mice according to standard techniques (Harlow, E. and Lane, D. (1988). *Antibodies: a laboratory manual*, Cold Spring Harbor Laboratory). The monoclonal antibody labeled PT18 was selected for use in further experiments.

Example 2

Purification of bovine GlcNAc-phosphotransferase

Lactating bovine mammary gland (6 kg) was collected at slaughter and immediately sliced into 10 cm thick slices and chilled in ice. Following homogenization in a Waring commercial blender, the post-nuclear supernatant fraction was prepared by centrifugation. Membrane fragments were collected by high speed centrifugation (39,000 x g, 45 minutes) and membrane proteins were solubilized in 4% Lubrol, 0.5 % deoxycholate. GlcNAc-

phosphotransferase was specifically adsorbed from the solubilized membrane fraction by incubation overnight with 10 ml of monoclonal antibody PT18 coupled to UltralinkTM matrix (substitution 5 mg/ml). The matrix was then collected by low speed centrifugation, washed with 0.025 M Tris-HCl, pH 7.4, 0.005 M MgCl₂, 0.3 % Lubrol buffer containing 1 M NaCl. The column was then washed with 2 column volumes of 0.01 M Tris-HCl, pH 7.4, 0.005 M MgCl₂, 0.3 % Lubrol buffer. GlcNAc-phosphotransferase was then eluted from the column with 0.10 M Tris-HCl, pH 10.0, 0.005 M MgCl₂, 0.3 % Lubrol and neutralized with 1/10th volume of 1 M Tris-HCl, pH 6.0. Recovery is typically 20-50% of the GlcNAc-phosphotransferase activity present in the homogenized tissue, and approximately 0.5 mg of enzyme is recovered per 10 kg of tissue processed.

Example 3

Amino acid sequencing of bovine GlcNAc-phosphotransferase

Example 3A

Reduction, alkylation and separation of individual subunits

Bovine GlcNAc-phosphotransferase, 1.9 mg was desalted on a column of Sephadex G-25 superfine equilibrated in 9% formic acid and lyophilized. The lyophilized protein was dissolved in 1 ml of 500 mM Tris-HCl, pH 8.6, 6 M guanidine-HCl, 10 mM EDTA, 2 mM DTT degassed by bubbling N₂ gas through the solution and incubated at 37°C for 1 hour. The solution was made 5 mM in iodoacetic acid and incubated at 37°C in the dark for a further 2 1/2 hours. The solution was then made 15 mM in β-mercaptoethanol and chromatographed on a column of Sephadex G-25 superfine equilibrated in 9% formic acid. The void fraction was pooled and lyophilized. The individual subunits were resolved by chromatography on a 1.0 x 30 cm column of Superose 12 equilibrated with 9% formic acid.

Example 3B

Amino terminal sequencing of individual subunits

Bovine GlcNAc-phosphotransferase, 0.5 mg was equilibrated with sodium dodecyl sulfate, electrophoresed on a 6% polyacrylamide gel in the presence of sodium dodecyl sulfate. The resolved subunits were then electro-transferred to a PVDF membrane and the protein bands detected by staining with Coomassie Blue. The bands corresponding to the individual subunits were then excised with a razor blade and subjected to amino-terminal sequencing in an Applied Biosystems Model 492 protein sequencer. The amino terminal sequence of the α -subunit was Met Leu Leu Lys Leu Leu Gln Arg Gln Arg Gln Thr Tyr (SEQ ID NO:26). The amino terminal sequence of the β Subunit is Asp Thr Phe Ala Asp Ser Leu Arg Tyr Val Asn Lys Ile Leu Asn Ser Lys Phe Gly Phe Thr Ser Arg Lys Val Pro Ala His (SEQ ID NO:27). The amino terminal sequence of the γ -subunit is Ala Lys Met Lys Val Val Glu Glu Pro Asn Thr Phe Gly Leu Asn Asn Pro Phe Leu Pro Gln (SEQ ID NO:28).

Example 3C

Internal amino acid sequence of the β - and γ -subunits

The resolved β - and γ -subunits from example 3B were treated with trypsin at a 1/40 mass ratio overnight at 37°C in 0.1 M Tris-HCl, pH 8.0. The tryptic fragments were then resolved by reverse phase chromatography on a C18 column equilibrated with 0.1% trifluoroacetic acid and developed with a linear gradient in acetonitrile. Well resolved peaks were then subjected to amino terminal sequencing as described in example 3B. The peptides sequenced from the β -subunit had the sequences Ile Leu Asn Ser Lys (SEQ ID NO:29), Thr Ser Phe His Lys (SEQ ID NO:30), Phe Gly Phe Thr Ser Arg (SEQ ID NO:31), and Ser Leu Val Thr Asn Cys Lys Pro Val Thr Asp Lys (SEQ ID NO:32). The peptide sequenced from the γ -subunit had the sequence Leu Ala His Val Ser Glu Pro Ser Thr Cys Val Tyr (SEQ ID NO:33). A second peptide sequence from the γ -subunit was obtained by

chymotryptic digestion with the sequence Asn Asn Pro Phe Leu Pro Gln Thr Ser Arg Leu Gln Pro (SEQ ID NO:34).

Example 3D

Internal amino acid sequence of the α -subunit

Internal peptide sequences of the α -subunit were obtained as follows. Bovine GlcNAc phosphotransferase was reduced, alkylated, electrophoresed and transferred to PVDF as previously described. The α -subunit band was excised and tryptic peptides generated by in situ digestion with trypsin, eluted with acetonitrile/trifluoroacetic acid and fractionated by reverse phase HPLC. Individual peaks were then examined by Matrix Associated Laser Desorption-Ionization-Mass Spectroscopy (MALDI-MS) and peaks containing a single mass were subjected to amino terminal sequencing as above. The peptide sequences determined from the α -subunit are Val Pro Met Leu Val Leu Asp Xaa Ala Xaa Pro Thr Xaa Val Xaa Leu Lys (SEQ ID NO:35) and Glu Leu Pro Ser Leu Tyr Pro Ser Phe Leu Ser Ala Ser Asp Val Phe Asn Val Ala Lys Pro Lys (SEQ ID NO:36).

Example 4

Cloning the human GlcNAc-phosphotransferase α/β -subunit cDNA

The amino-terminal protein sequence determined from the isolated bovine β -subunit was used to search the Expressed Sequence Tag (EST) data base using the program tblastn. Altschul, S. F., Gish W., *et al.* (1990). "Basic Local Alignment Search Tool." *Journal of Molecular Biology* 215: 403-410. This search identified a partial mouse cDNA previously identified during a positional cloning strategy. Cordes, S. P. and Barsh, G. S. (1994). "The mouse segmentation gene *kr* encodes a novel basic domain-leucine zipper transcription factor." *Cell* 79: 1025 - 11034.

A forward PCR primer was designed based on the mouse sequence and used with an

oligo dT reverse primer for RT-PCR amplification of a 1,848 bp product using mouse liver poly A RNA as template. The PCR product was cloned and sequenced and proved to contain all the determined β -subunit sequences, demonstrating it encoded the murine β -subunit.

The human β -subunit cDNA was cloned by screening a size selected human placental cDNA library (Fischman, K., Edman J. C., *et al.* (1990). "A murine *fer* testis-specific transcript (*ferT* encodes a truncated *fer* protein." *Molecular and Cellular Biology* 10: 146 - 153) obtained from ATCC with the random hexamer labeled murine β -subunit cDNA under conditions of reduced stringency (55°C, 2XSSC). The remaining portion of the α/β -subunit precursor cDNA was cloned by a combination of a walking strategy beginning with the portion of the cDNA encoding the human β -subunit and standard library screening strategies. Additionally, EST data base searches were used to identify clones containing portions of the human α/β cDNA, which were obtained from the corresponding repositories and sequenced. Together these strategies allowed the determination of the full length human α/β -subunits precursor cDNA sequence. A clone containing this sequence was assembled using the appropriate fragments and cloned into pUC19. The 5597 bp sequence is given in Sequence NO:4 and contains DNA sequences predicted to encode protein sequences homologous to all of the amino terminal and internal peptide sequences determined from the bovine α - and β -subunits.

Example 5

Cloning the human GlcNAc-phosphotransferase γ -subunit cDNA

The γ -subunit amino terminal and tryptic peptide sequences were used to search the Expressed Sequence Tag (EST) data base using the program tblastn. Altschul, S. F., Gish W., *et al.* (1990). "Basic Local Alignment Search Tool." *Journal of Molecular Biology* 215: 403-10. Three human EST sequences were identified which were highly homologous to the

determined bovine protein sequences. cDNA clone 48250 from which EST sequence 280314 was determined was obtained from Genome Systems and sequenced using standard techniques. This clone contained a 1191 bp insert which contained all the determined protein sequences and appeared to contain a signal sequence 5' of the determined amino terminal sequence. The clone however lacked an initiator methionine or any 5' non-coding sequence. The 5' portion of the cDNA was obtained by PCR. the reverse primer 5'-GCGAAGATGAAGGTGGTGGAGGACC-3' (SEQ ID NO:37) and a T7 promoter primer were used in a reaction along with template DNA from a human brain cDNA library in pCMV-SPORT(GIBCO). A 654 bp product was obtained, cloned in pCR2.1 and sequenced. The sequence demonstrated the amplified product contained 23 bp of 5' non-coding sequence, the initiator methionine and the signal peptide identified in EST 280314. A full length cDNA for the γ -subunit (pBC36) was assembled by ligating a 75 bp EcoRI-ApaI fragment from the cloned PCR product, an ApaI-NotI fragment from clone 48250 and EcoRI-NotI cut pcDNA3 (Invitrogen).

Example 6

Cloning the human GlcNAc-phosphotransferase α/β -subunit gene

Plasmid DNA was prepared from a human brain cDNA library (Life Technologies) according to the manufacturers protocol. This DNA was used as template for PCR using primers with the sequences 5'-TGCAGAGACAGACCTATACCTGCC-3' (SEQ ID NO:38) and 5' ACTCACCTCTCCGAACTG-GAAAG-3' (SEQ ID NO:39) using Taq DNA polymerase and buffer A from Fischer Scientific using 35 cycles of 94°C 1 minute, 55°C 1 minute, and 79°C 1 minute. A 106 bp product was obtained, purified by agarose gel electrophoresis, isolated by GeneClean (Biol01) and cloned into pCR2. DNA sequencing determined the resulting plasmid pAD39 contained a 106 bp insert which was excised by

digestion with EcoRI and submitted to Genome Systems for screening of a human genomic BAC library. Four human BACs were identified and BAC #14951 was sequenced. For sequencing BAC #14951 was submitted to a colleague's laboratory at the University of Oklahoma. The BAC was then fragmented by nebulization, and fragments cloned into pUC18 and shotgun sequenced. Contigs were generated by computer analysis and gaps closed by primer walking strategies. The sequence of the BAC spans 177,364 bp. The GlcNAc-phosphotransferase α/β -subunits precursor gene spans ~80 kb and is arranged as 21 exons.

Example 7

Cloning the human GlcNAc-phosphotransferase γ -subunit gene

The human γ -subunit gene was identified by blastn searching of the NCBI High Throughput Genomic Sequence (HGTS) database with the full length human Subunit cDNA sequence. The search identified a clone HS316G12(gi 4495019) derived from human chromosome 16 which contained the human γ -subunit gene. The human GlcNAc-phosphotransferase γ -subunit gene spans about 12 kb and is arranged as 11 exons. Exons 1-3 and 4-11 are separated by a large intron of about 9 kb.

Example 8

Preparation of modified expression plasmid for the human GlcNAc-phosphotransferase

α/β -subunits precursor cDNA

An expression vector for the GlcNAc-phosphotransferase α/β cDNA was constructed in pcDNA3.1(+) as follows. Two upstream ATG's in the 5'-noncoding sequence of the human GlcNAc-phosphotransferase cDNA were removed and the Kozak sequence were modified as follows. Two fragments from pAD98, which was the human GlcNAc-phosphotransferase ct/p cDNA cloned into pcDNA3.1(+), were excised. A 1068 bp XhoI-

PstI fragment and a 9746 bp NheI-XhoI fragment were ligated with oligonucleotides with sequences 5'-CTAGCCACCATGGGGTTCAAGCTCTTGCA-3' (SEQ ID NO:40) and 5'-AGAGCTTGAACCCCATGGTGG-3' (SEQ ID NO:41) generating pAD105. The poly A sequence near the 3' end of the cDNA clone was removed by ligating a NheI-BglII fragment from the cDNA with NheI-BamHI cut vector pcDNA3.1(+) generating pAD128.

Example 9

Preparation of an expression plasmids for the human GlcNAc-phosphotransferase α/β subunits precursor cDNA

DNA sequencing of pAD128 identified deletion of an A in an AAAAAA sequence (positions 2761-2765 shown in SEQ ID NO:4) that disrupted the coding sequence. Plasmid pAD130 was constructed in an attempt to correct this by ligating a 5929 bp NheI-MfeI fragment and a 2736 bp NheI-AgeI fragment (both from pAD128 with a 515 bp MfeI-AgeI fragment derived from pAD124). Plasmid pAD130 was then grown and subsequent sequencing of plasmid pAD130 demonstrated that the AAAAAA sequence had reverted to AAAA again indicating instability in the sequence at this point.

In order to eliminate this instability the first AAA (position 2761-2763 shown in SEQ ID NO:4) that codes for lysine was changed to AAG (also coding for lysine) so that the unstable AAAAAA sequence was changed to a stable AAGAA without altering the encoded amino acid. Plasmid pAD130 was corrected by removing a 214 bp MfeI-DraIII fragment and replacing it with a fragment with the correct sequence. The correct MfeI- DraIII fragment was prepared by PCR using pAD130 as a template with forward primer 5'-GAAGACACAATTGGCATACTTCACTGATAGCAAAGAAATACTGGGAGGC AACTAAAAGATAC-3' (SEQ ID NO:42) (oligo TTI 25 with desired AAGAA sequence as underlined) and reverse primer 5'-ACTGCATATCCTCAGAATGG-3' (SEQ ID NO:43)

(oligo TTI 24). The PCR fragment was subcloned into the EcoRV site of pBluescript KS II(+) (Stratagene) generating pMK16. The insert was sequenced for confirmation and the 215 bp MfeI- DraIII fragment was prepared. To avoid MfeI- DraIII sites on the vector pcDNA 3.1(+) (Invitrogen), the NheI-XbaI fragment was prepared from pAD130 and subcloned into the XbaI site of pUC19 (Life Technologies) to construct pMK15. pMK15 was cleaved with MfeI and DraIII and the 6317 bp fragment was purified and ligated with the MfeI- DraIII fragment from pMK16 to form pMK19 containing the desired stable sequence in pUC19.

The corrected cDNA for the α/β subunit was excised from pMK19 as a KpnI-XbaI fragment and subcloned between the KpnI and XbaI sites of pcDNA6/V5/His-A and designated pMK25. Plasmid pMK25 containing the cDNA as shown in SEQ ID NO:20 where the nucleotide sequence for the modified human α/β -subunit precursor cDNA is shown in nucleotides 1-3768. This sequence corresponds to and is a modification of the nucleotide sequence 165-3932 shown in SEQ ID NO:4.

Example 10

Construction of Expression vectors for soluble, human GlcNAc-phosphotransferase α/β subunits precursor cDNA

Plasmid pMK19 was digested with BglII (cutting at positions 255 and 2703 shown in SEQ ID NO:20) and self-ligated to reduce the length of the cDNA to be amplified from approx. 3.5 kb to 1 kb so that the 5' and 3' ends of the cDNA can be modified by PCR to remove the transmembrane domains of the α and β subunits of human GlcNAc-phosphotransferase and used to construct expression vectors to produce soluble GlcNAc-phosphotransferase. This plasmid was designated pMK21. The strategy is that the nucleotides encoding the first 44 amino acids containing the transmembrane domain of the α

subunit (nucleotides 1-132 of SEQ ID NO:20) are replaced with a HindIII site, and nucleotides encoding the last 47 amino acids containing the transmembrane domain of the β subunit (nucleotides 3628-3768 of SEQ ID NO:21) are replaced with a stop codon and a XbaI site.

Plasmid pMK21 was used as a template for PCR with the following primers: A forward primer (5'-TGGTTCTGAAAGCTTAGCCGAGATCAATACCATG-3' (SEQ ID NO:44), oligo TTI 76) containing a HindIII site (underlined) and a sequence complementary to nucleotides 133 to 151 of SEQ ID NO:20 (*italics*), which will produce the 5'-end of a PCR fragment that removes the coding sequence of the first 44 amino acids comprising the putative transmembrane domain of the α subunit. A reverse primer (5'-TAGTACACTCTAGActactaCTTCAATTTGTCTCGATAAG-3' (SEQ ID NO:45), oligo TTI 78) containing a XbaI site (underlined), two stop codons (lower case) and a sequence complementary to nucleotides 3608 to 3627 of SEQ ID NO:21 (*italics*), which will produce the 3'-end of a PCR fragment that removes the coding sequence of the last 47 amino acids comprising the putative transmembrane domain of the β subunit and replaces it with two stop codons. The resulting PCR fragment was subcloned into the EcoRV site of pBluescript KS II+ (Stratagene). This plasmid, designated pMK42, was sequenced to ensure no errors were introduced by PCR. The BglII- BglII fragment (positions 255-2703 shown in SEQ ID NO:20) which was previously removed was subcloned back into the BglII site of pMK42. The orientation of this fragment was determined to be correct and this plasmid was designated pMK49. Thus, plasmid pMK49 contained a cDNA comprising a 5' HindIII site and a 3' XbaI site flanking a coding region for the human GlcNAc-phosphotransferase α/β subunits precursor cDNA with the α subunit putative transmembrane domain deleted and the

putative transmembrane domain of the β subunit replaced with two stop codons (soluble α/β -cDNA).

This "soluble α/β -cDNA" can now be conveniently sub-cloned into vectors constructed to contain the HPC4 epitope (used for rapid purification of the soluble enzyme) and different secretion signal peptides. These pcDNA6/V5/His-A+tag) vectors were constructed as follows:

Synthetic oligonucleotide cassettes containing a 5'-NheI site and a 3'-HindIII site flanking nucleotide sequences coding for different secretion signal peptides and the nucleotide sequence coding for the HPC4 epitope were inserted into plasmid pcDNA6/V5/His-A cut with NheI and HindIII. The following plasmids were prepared with the indicated cassette:

1. pMK45 - mouse immunoglobulin Kappa chain signal peptide (sequence in italics) and HPC4 epitope (sequence underlined)

CTAGCCGCCACC *ATGGAGACAGACACACTC* CTGCTATGGGTACTGCTGCTC

GGCGGTGGTACC TC TGTCT GTGTGAGGACGATACCCATGACGACGAG
TGGGTTCC AGGT TC CACTGGTGA CGAAGATCAGGTAGATCCGCGGTT AATC
 ACCCAAGGTCCAAGGTGACCACTGCTTC TAGTCCAT CTAGGCGCCAATTAG
GACGGTA

CT GCCATTCGA (SEQ ID NO:46)

1. pMK44 - a transferrin signal peptide sequence (in italics) and HPC4 epitope (sequence

underlined)

CTAGCGGTACCATGAGATT AGCAGTAGGCGCC TT ATTAG TATGCGC AGTACT C
CGCCATGGTACTCTAATCGTCATCCGCGGAATAATCATACGCGTCATGAG
GGATTAT GTC TCGCAG AAGATCAGGTAGATCCGC GGTT AATCGACGGTA
CCTTATACAGAGCGTCTTCTAG TCCAT CTAGGCGCCAAT TAGCTGCCATTCGA
(SEQ ID NO:47)

1. pMK43 - a transferrin secretion peptide sequence modified to satisfy a Kozak's
sequence(sequence in italics) and HPC4 epitope (sequence underlined),

CTAGCCGCCACCATGGGATT AGCAGTAGGCGCCTT ATT AGT ATGCGC AGT
CGCCGGTGGTACCCTAATCGTCATCCGCGGAATAATCATACGCGTCA

ACT CGGATTAT GT C TCGCA GAAGATCAGGTAGATCCGC GGTTAATCGACG
TGAGCCTAATACAGAGCGTCTT CTAGT CCATCTAGGCGCCAAT TAGCTGC
GTA
CATTCGA (SEQ ID NO:48)

The cDNA encoding "soluble α/β subunits" can be obtained as a HindIII- XbaI
fragment from pMK49 and inserted into the plasmid pMK43 to form pMK50; pMK44 to
form pMK51, and into pMK45 to form pMK52, plasmids capable of encoding the α/β
subunits of human GlcNAc-phosphotransferase with putative transmembrane domains
deleted, with different signal peptides and all having the HPC4 epitope tag to facilitate

purification of the soluble, secreted enzyme.

Example 11

Construction of Expression vectors for the human GlcNAc-phosphotransferase γ subunit precursor cDNA

The human GlcNAc-phosphotransferase γ -subunit precursor cDNA was obtained from plasmid pAD133 in pAC5.1/V5-His by cutting with EcoRI. This cDNA was inserted into EcoRI digested pcDNA6/V5/His-A to form plasmid pMK17 containing cDNA as shown in SEQ ID NO:5. Plasmid pMK17 was digested with MluI (position 124-129 as shown in SEQ ID NO:5) and EcoRI (position 1103-1108 as shown in SEQ ID NO:5) and the 980 bp MluI- EcoRI fragment was then subcloned in pBluescriptKSII(+) with a synthetic double stranded cassette having an HindIII site and a MluI site flanking a nucleotide sequence including positions corresponding to 95-123 as shown in SEQ ID NO:5 thereby removing the nucleotide sequence encoding the amino terminal, 24-amino acid signal peptide in plasmid pMK26. Plasmid pMK26 was sequenced to ensure its sequence. The correct cDNA from pMK26, which encodes amino acids for the human GlcNAc-phosphotransferase γ subunit with the signal peptide removed, is then excised from pMK26 by HindIII and EcoRI digestion and placed into plasmids pMK43 to form pMK58; pMK44 to form pMK59, and into pMK45 to form pMK64, plasmids capable of encoding the γ subunit of human GlcNAc-phosphotransferase with its signal peptide deleted, with different signal peptides and all having the HPC4 epitope tag to facilitate purification of the soluble, γ subunit.

To evaluate the behavior of $\alpha/\beta/\gamma$ secreted products, the α/β subunit precursor and the γ subunit were co-expressed in the bi-cistronic vector pIRES (Clontech). This was accomplished by subcloning α/β and γ cDNAs expressing the desired subunit with a selected

signal peptide and the HPC4 Tag into NheI site (MCS-A) and XbaI site (MCS-B) of pIRES, respectively.

Example 12

Transient expression of the α/β and γ subunits of human GlcNAc-phosphotransferase in 293T Cells

Plasmids were transfected into 293T cells using Fugene6 (Roche) according to manufacturer's instructions. Culture media was collected 23 h, 44.5 h and 70 h after transfection. Aliquots of media containing expressed protein was captured on anti-HPC4 monoclonal antibody (U.S. Patent No. 5,202,253) conjugated with Ultralink beads (Pierce) by overnight incubation at 4 °C. The beads were washed to remove unbound protein and assayed directly for phosphotransferase activity as described previously (REF).

Plasmids used for expression all containing a sequence encoding for the HPC4 tag were as follows:

1. pMK50 -modified transferrin secretion peptide and α/β subunit in pcDNA6/V5/His-4
2. pMK51 -transferrin secretion peptide and α/β subunit in pcDNA6/V5/His-4
3. pMK52 - mouse immunoglobulin secretion peptide and α/β subunit in pcDNA6/V5/His-4
4. pMK75 - modified transferrin secretion peptide and α/β subunit and modified transferrin secretion peptide and γ subunit in pIRES
5. pMK81 -transferrin secretion peptide and α/β subunit and transferrin secretion peptide and γ subunit in pIRES
6. pMK76 - mouse immunoglobulin secretion peptide and α/β subunit and mouse

immunoglobulin secretion peptide and γ in pIRES

The relative amounts of expression detected by assay for phosphotransferase using methyl- α -D-mannoside and UDP- $[\beta$ - ^{32}P]-GlcNAc as substrates with cell transfected with pcDNA6/V5/His-4 as controls is shown in Figure 4.

Example 13

Expression and purification GlcNAc-phosphotransferase $\alpha/\beta/\gamma$

For expression and purification of the enzyme, a modified expression plasmid is constructed in a modified expression vector derived from pEE14. The plasmid directs the synthesis of a soluble epitope tagged GlcNAc-phosphotransferase molecule. The α/β -subunit precursor is modified as follows: The 5' portion of the cDNA which encodes the α -subunit cytoplasmic and transmembrane domain is deleted and replaced with nucleotides which encode the transferrin signal peptide followed by amino acids which encode the epitope for monoclonal antibody HPC4. The 3' portion of the cDNA is modified by the insertion of a stop codon before the β -subunit transmembrane segment. The vector pEE14.1 (Lonza Biologics) is modified by the insertion of a 850 bp MluI-NcoI fragment containing a modified vascular endothelial growth factor (VEGF) promoter at the unique MluI site in pEE14.1. This vector encoding the modified GlcNAc-phosphotransferase α/β -subunit precursor is co-transfected with a wild type γ -subunit construct containing the VEGF promoter in pEE14.1 into CHO-K1 cells using Fugene6 and plated into 96 well plates. Transfectants are selected in 25 μM methionine sulfoximine and the plasmid amplified by selection in 96 well plates with 50 μM , 100 μM , 250 μM , and 500 μM methionine sulfoximine. Clones are picked into duplicate 96 well plate and the highest expressing clones selected by dot blotting media and immuno-detection with monoclonal antibody HPC4. The highest expressing clone is

expanded into cell factories. The recombinant soluble epitope tagged GlcNAc-phosphotransferase is purified from the media by chromatography on monoclonal antibody HPC4 coupled to Ultralink in the presence of 5 mM MgCl₂ and 1 mM CaCl₂. The soluble epitope tagged GlcNAc-phosphotransferase is eluted with 5 mM EGTA and 5 mM MgCl₂.

Example 14

Preparation of monoclonal antibodies specific for bovine phosphodiester α -GlcNAcase

Murine monoclonal antibodies specific for bovine phosphodiester α -GlcNAcase were generated by immunization of mice with a partially purified preparation of phosphodiester α -GlcNAcase. Spleens were then removed from immune mice and fused with SP2/O myeloma cells according to standard techniques (Harrow, E. and Lane, D. (1988). *Antibodies: a laboratory manual*, Cold Spring Harbor Laboratory). Hybridomas were plated in eight 96-well plates and grown until hybridomas were visible. Hybridomas secreting antibodies to phosphodiester α -GlcNAcase were identified measuring phosphodiester α -GlcNAcase activity in immunoprecipitates prepared by incubation of a partially purified preparation of phosphodiester α -GlcNAcase with pooled hybridoma supernatants. Pools from 16 and 4 wells were assayed followed by individual wells. Monoclonal UC1 was identified by this protocol and coupled to Ultralink™ for use in purification of phosphodiester α -GlcNAcase.

Example 15

Purification of bovine phosphodiester α -GlcNAcase

Bovine calf liver (1 kg) was homogenized in 0.05 M Imidazole-HCl, pH 7.0, 0.15 M NaCl, 0.01 M EDTA and a washed post-nuclear supernatant was prepared. Membranes were collected by centrifugation at 30,000 x g for 30 minutes and washed three times with the above buffer. Membrane proteins were then solubilized in buffer containing 2% Triton X* 100, 0.05% deoxycholate and insoluble material removed by centrifugation, as before. The

solubilized membrane fraction was incubated with 20 ml of monoclonal antibody UC1 coupled to Ultralink™ (substitution 5 mg/ml) with constant rotation for 16 hours at 4°C. The UC1-Ultralink™ was collected by low speed centrifugation. packed into a column and washed with 0.025 M Tris-HCl, pH 7.4, 0.3% Lubrol, followed by two column volumes of 0.5 M NaHCO₃, pH 8.0, 0.3% Lubrol. Phosphodiester α -GlcNAcase was then eluted with 0.5 M NaHCO₃, pH 10.0, 0.3% Lubrol and collected in 1/10 volume of 1.0 M Tris-HCl, pH 5.5.

Example 16

Amino acid sequencing of bovine phosphodiester α -GlcNAcase

Example 16A

Amino-terminal sequence of bovine phosphodiester α -GlcNAcase

Bovine phosphodiester α -GlcNAcase was bound to a 0.25 ml column of POROS HQ and step-eluted with buffer containing 0.5 M NaCl. Fractions containing phosphodiester α -GlcNAcase activity were identified by phosphodiester α -GlcNAcase assay, pooled and absorbed to a ProSorb Sample Preparation Cartridge (Perkin Elmer) and subjected to amino acid sequencing in an Applied Biosystems Model 492 Protein Sequencer operated according to the manufacturer's instructions. The sequence Asp-Xaa-Thr-Arg-Val-His-Ala-Gly-Arg-Leu-Glu-His-Glu-Ser-Trp-Pro-Pro-Ala-Ala-Gln-Thr-Ala-Gly-Ala-His-Arg-Pro-Ser-Val-Arg-Thr-Phe-Val was obtained.

Example 16B

Internal sequence of bovine phosphodiester α -GlcNAcase

Bovine liver phosphodiester α -GlcNAcase was concentrated to 10 μ l in a Speed Vac, combined with 30 μ l 0.1 M Tris-HCl, pH 7.4, 8 M guanidine-HCl, and 2-4 μ l 25 mM DTT and incubated at 50°C for 1 hour. Iodoacetamide 2.4 μ l 50 μ M was then added and the

incubation was continued for 1 hour. The reaction mixture was then desalted on a column of Sephadex G25 superfine as described for GlcNAc-phosphotransferase and digested with trypsin. The peptides were fractionated by HPLC and sequenced as described for GlcNAc-phosphotransferase. The sequences determined are Arg Asp Gly Thr Leu Val Thr Gly Tyr Leu Ser Glu Glu Glu Val Leu Asp Thr Glu Asn and Gly Ile Asn Leu Trp Glu Met Ala Glu Phe Leu Leu Lys.

Example 17

Cloning the human phosphodiester α -GlcNAcase cDNA

The phosphodiester α -GlcNAcase tryptic peptide sequences were used to search the EST data bases as described for GlcNAc-phosphotransferase above. Three EST sequences were identified which contained the human phosphodiester α -GlcNAcase cDNA and clone ATCC #367524 was obtained and a ~700 bp EcoRI-NotI fragment was excised from this clone and used to probe a human liver cDNA library in the vector TriplEx. Several clones were identified and sequenced, one of which (clone 6.5) proved to contain a nearly full length cDNA for the human phosphodiester α -GlcNAcase. The genomic clone described in Example 18 demonstrated that clone 6.5 was missing only the initiator methionine.

Example 18

Cloning the human phosphodiester α -GlcNAcase gene

The human phosphodiester α -GlcNAcase gene was identified by searching the NCBI database nr with the human phosphodiester α -GlcNAcase cDNA using the program blastn. The genomic sequence was determined during the sequencing of a clone from chromosome 16p13.3 and deposited 06-Mar-1999 in GenBank as an unidentified sequence of 161264 bp with the accession number AC007011. The gene spans about 12 kb of genomic DNA on chromosome 16.13 and is arranged in 11 exons.

Example 19

Construction of an expression vector for human phosphodiester or α -GlcNAcase

An expression vector for human phosphodiester α -GlcNAcase was prepared as follows: The 5' end of the sequence of clone 6.5 was modified by PCR amplification of the 5' end of the cDNA with a forward primer with the sequence 5'-GGAATTCCACCATGGCGACCTCCACGGGTCG-3' (SEQ ID NO:49) and a reverse primer 5'-TGACCAGGGTCCCGTCGCG-3' (SEQ ID NO:49). This served to add a consensus Kozak sequence and initiator methionine to the sequence of clone 6.5. The ~500 bp PCR product was purified, digested with EcoRI and BamHI and ligated into pcDNA3.1(-) which was sequenced. This construct was then digested with BamHI and HindIII and ligated with a ~1600 bp BamHI-HindIII fragment containing the 3' portion of the cDNA from clone 6.5 generating the full length expression plasmid.

Example 20

Host cell preparation for human phosphodiester α -GlcNAcase

Cos cells were grown in 60 mm plates in Dulbeccos minimal essential media (DMEM) at 37°C in 5% CO₂ until they reached 50-80% confluence. The plates were then washed with OptiMEM I and the cells transfected with the expression vector described in Example 19 using Lipofectamine Plus (GIBCO BRL Life Technologies) according to the manufacturers instructions. Cells were harvested at 48 hours, a solubilized membrane fraction prepared and assayed for phosphodiester α -GlcNAcase activity.

Example 21

Expression and purification of soluble recombinant human phosphodiester α -GlcNAcase

For expression and purification of the enzyme, a modified expression plasmid is constructed in a modified expression vector derived from pEE14.1. The plasmid directs the

synthesis of a soluble epitope tagged phosphodiester α -GlcNAcase molecule. The phosphodiester α -GlcNAcase precursor is modified as follows: The 3' portion of the cDNA which encodes the phosphodiester α -GlcNAcase transmembrane and cytoplasmic domains is deleted and replaced with nucleotides which encode the epitope for monoclonal antibody HPC4 followed by a stop codon. The vector pEE14.1 (Lonza Biologics) is modified by the insertion of a 850 bp MluI-NcoI fragment containing a modified vascular endothelial growth factor (VEGF) promoter at the unique MluI site in pEE14.1. This vector encoding the epitope tagged soluble phosphodiester α -GlcNAcase precursor is transfected into CHO-K1 cells using Fugene6 and plated into 96 well plates. Transfectants are selected in 25 μ M methionine sulfoximine, and the plasmid amplified by selection in 96 well plates with 50 μ M, 100 μ M, 250 μ M, and 500 μ M methionine sulfoximine. Clones are picked into duplicate 96 well plate and the highest expressing clones selected by dot blotting media and immuno-detection with monoclonal antibody HPC4. Media from clones demonstrating the highest level of epitope tag expression is assayed for phosphodiester α -GlcNAcase activity. The highest expressing clone is expanded into cell factories. The recombinant soluble epitope tagged phosphodiester α -GlcNAcase is purified from the media by chromatography on monoclonal antibody HPC4 coupled to Ultralink™ in the presence of 5 mM MgCl₂ and 1 mM CaCl₂. The soluble epitope tagged phosphodiester α -GlcNAcase is eluted with 5 mM EGTA and 5 mM MgCl₂.

Example 22

Construction of an Expression Vector for Soluble, Human Phosphodiester α -GlcNAcase

For expression and purification of the enzyme, a modified expression plasmid is constructed in a modified expression vector derived from the pEE14.1 vector (Lonza Biologics). The plasmid directs the synthesis of a soluble epitope tagged phosphodiester

α -GlcNAcase molecule. The phosphodiester α -GlcNAcase precursor is modified as follows: The 3' portion of the cDNA (1342-1548 of SEQ ID NO: 7) which encodes the phosphodiester α -GlcNAcase transmembrane and cytoplasmic domains was deleted and replaced with nucleotide sequence GAGGACCAGGTGGACCCCAGGCTGATCCAC GGCAAGGAT (SEQ ID NO:51) that encodes the epitope for monoclonal antibody HPC4 (EDQVDPRLIDGKD (SEQ ID NO:52)) followed by a stop codon.

This expression vector was constructed by generating two intermediate plasmids and ligating a fragment from each into pEE14.1 vector (Lonza Biologics) to yield the final expression vector. The first intermediate plasmid designated pKB4 was constructed by ligating the 1034 bp FseI-Bsu36I fragment of phosphodiester α -GlcNAcase (lacking the C-terminal transmembrane and cytoplasmic domains) from clone 6.5, and a Bsu36I-XbaI oligonucleotide fragment that contains the HPC4 epitope into a modified pUC19 vector. The second intermediate plasmid designated pKB5, was constructed by ligating a 850 bp MluI-NcoI fragment containing a modified vascular endothelial growth factor (VEGF) promoter from pcDNA4/HisMax (Invitrogen), a 256 bp BseI-FseI fragment encoding the N-terminus of human phosphodiester α -GlcNAcase from clone 6.5, and an oligonucleotide linker into a modified pUC19 vector. The final expression vector designated pKB6 was constructed by ligating the MluI-FseI fragment from pKB5, and the FseI-HindIII fragment from pKB4 into a MluI/HindIII digested pEE14.1 vector. The plasmid pKB6 contains the nucleotide sequence shown in SEQ ID NO:22.

Expression and purification of soluble recombinant human phosphodiester α -GlcNAcase

Approximately 10^8 293T cells were plated in a cell factory using Dulbecco's modified

eagle's medium (DMEM) containing 10% fetal bovine serum in a humidified atmosphere at 37°C with 5% CO₂. These cells were transfected with approximately 700 g of pKB6 using 2 ml of transfection reagent Fugene-6 (Roche) for the transient expression of soluble human phosphodiester α -GlcNAcase. After three days of culturing the transfected cells, the medium containing soluble, epitope-tagged, human phosphodiester α -GlcNAcase was collected and applied in the presence of 1 mM CaCl₂ to a column of monoclonal antibody HPC4 coupled to Ultralink (Pierce). Affinity purified, epitope-tagged, human phosphodiester α -GlcNAcase (approximately 11 mg) was eluted with buffer containing 5 mM EDTA and stored at -20°C in 50 mM Tris, 150 mM NaCl, 2 mM CaCl₂, 50% glycerol, pH 7.2. The enzyme had a specific activity of 500,000 units/mg with [³H]GlcNAc-phosphomannose- α -methyl as a substrate (Kornfeld R, et al., JBC 273:23203-23210).

Example 23

CHO cells expressing recombinant human acid α -glucosidase

The human acid α -glucosidase cDNA was obtained from Dr. Frank Martinuk (Martiniuk, F., Mehler, M., Tzall, S., Meredith, G. and Hirschhorn, R. (1990). "Sequence of the cDNA and 5'-flanking region for human acid alpha-glucosidase, detection of an intron in the 5' untranslated leader sequence, definition of 18-bp polymorphisms, and differences with previous cDNA and amino acid sequences." *DNA Cell Biol* 9(2): 85-94) and cloned into the expression vector pEE14.1. This vector was used to transfect CHO-K1 cells using Fugene6 and plated into 96 well plates. Transfectants were selected in 25 μ M methionine sulfoximine, and clones picked and plated into 96 well plates. The plasmid was amplified by selection with 50 μ M, 100 μ M, 250 μ M, and 500 μ M methionine sulfoximine. Clones were picked

into duplicate 96 well plates and the highest expressing clones selected by assay of the media for acid α -glucosidase activity and the cells for DNA content. The highest expressing clone (Clone 3.49.13) based on acid α -glucosidase activity to DNA content ratio was then expanded into a cell factory. This clone was incubated at 37°C in 5% CO₂ and maintained in Glasgow Minimal Essential Media containing 20 mM TES, pH 7.2, 5% fetal bovine serum.

Example 24

Growth of CHO cells expressing recombinant human acid α -glucosidase
in the presence of α -1,2 mannosidase inhibitors

CHO cells expressing human acid α -glucosidase were cultured in Glasgow Modified Minimal Essential Media containing 5% Fetal Bovine Serum, 25 μ M methionine sulfoximine, 20 mM TES, pH 7.2, and 7.5 mM l-deoxymannojirimycin-HCl. Alternatively, the cells can be cultured in the above media containing 100 μ g/mL l-deoxymannojirimycin-HCl and 25 μ g/mL kifunensine.

Example 25

Isolation of recombinant human acid α -glucosidase

Recombinant human acid α -glucosidase was purified from spent tissue culture media as follows: Media was concentrated 10 fold by tangential ultrafiltration with a 30,000 dalton cutoff membrane and dialyzed into 50 mM sodium phosphate, pH 6.5, and applied to a column of ConA Sepharose (Pharmacia). Following a wash with the same buffer to remove the unbound proteins, acid α -glucosidase was eluted with 1.0 M α -methyl glucoside, pooled, concentrated and dialyzed as before. The acid α -glucosidase was then applied to a column of Sephadex G-200 equilibrated with 50 mM sodium phosphate, pH 6.5 and eluted isocratically with the same buffer.

Example 26

Treatment of recombinant human acid α -glucosidase with GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase

Human acid α -glucosidase at 10 mg/ml was incubated in 50 mM Tris-HCl, pH 6.7, 5 mM $MgCl_2$, 5 mM $MnCl_2$, 2 mM UDP-GlcNAc with GlcNAc-phosphotransferase at 100,000 u/mL at 37°C for 2 hours. Phosphodiester α -GlcNAcase, 1000 u/mL was then added and the incubation continued for another 2 hours. The acid α -glucosidase was then repurified by chromatography on Q-Sepharose, and step elution with NaCl.

Example 27

Characterization of the oligosaccharide structures on modified recombinant human acid α -glucosidase

Recombinant acid α -glucosidase treated or untreated with GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase was digested with N-glycanase (New England Biolabs) or endomannosidase H (New England Biolabs) according to the manufacturer's conditions. The released oligosaccharides were then labeled on the reducing terminus with 2-aminobenzamide and fractionated by HPLC with fluorescent detection according to the manufacturer's instructions (Oxford Glycosystems). Peaks were identified by comparison with standards chromatographed on the same system, and confirmed by digestion with linkage specific glycosidases and/or mass determination by MALDI. The results are shown in Table 1.

Table 1

Enzyme Preparation	M6	M7	M8	M9	1P-Gn	2P-Gn	1M6P	Complex
Rh-GAA (Secreted)	0	0	0	0	0	0	1	99
Rh-GAA (dMM/intracellular)	23	31	23	6	0	0	17	0
Rh-GAA (dMM/intracellular) Ptase-treated	6	11	7	2	12	62	0	0

Referring to Table 1, the data (given in mole percent) show that the Lysosomal enzymes prepared using the GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase of the present invention are highly phosphorylated. The data shows that the present invention produces lysosomal enzymes having about 5-10 M6P groups per enzyme compared to about 0-2 for untreated enzymes and enzymes known in the art. When compared to naturally occurring or recombinant lysosomal enzymes, the *in vitro*-modified preparation is very highly phosphorylated. In the most highly phosphorylated lysosomal enzyme known in the art, the α -galactosidase A described by Matsuura, F., Ohta, M., Ioannou, Y. A. and Desnick, R. J. (1998). "Human alpha-galactosidase A: characterization of the N-linked oligosaccharides on the intracellular and secreted glycoforms overexpressed by Chinese hamster ovary cells." *Glycobiology* 8(4): 329-39, 5.2% of the oligosaccharides are bis-phosphorylated. In marked contrast, 62% of the oligosaccharides on the *in vitro*-phosphorylated acid α -glucosidase, preparation described here contains bis-phosphorylated oligosaccharides. This represents about a 12 fold increase. When the *in vitro* phosphorylated preparation of rh-GAA shown in Table 1 is compared with GAA secreted from CHO cells by methods known in the art, an

even greater increase in phosphorylation is evident, about a 62 fold increase.

Thus, the *in vitro*-phosphorylated GAA is 12-62 fold more phosphorylated than any other described preparation of natural or recombinant lysosomal enzyme. This difference has a major influence on the rate and extent of internalization (Reuser, A. J., Kroos, M. A., Ponne, N. J., Wolterman, R. A., Loonen, M. C., Busch, H. F., Visser, W. J. and Bolhuis, P. A. (1984). "Uptake and stability of human and bovine acid alpha-glucosidase in cultured fibroblasts and skeletal muscle cells from glycogenosis type II patients." *Experimental Cell Research* 155: 178-189).

Example 28

Comparison of cell uptake of recombinant human acid α -glucosidase with or without modification by GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase

Human Pompe disease fibroblasts are obtained from ATCC and cultured in DMEM with 10% fetal bovine serum in 6 well plates and incubated at 37°C in 5% CO₂. Recombinant human acid α -glucosidase with different carbohydrate structures are compared for the rate and extent of internalization. Controls include each preparation incubated with 5 mM mannose 6-phosphate and incubations without added recombinant human acid α -glucosidase. The different preparations to be examined include acid α -glucosidase secreted from CHO cells, acid α -glucosidase secreted from CHO cells in the presence of α 1,2-mannosidase inhibitors, acid α -glucosidase secreted from CHO cells in the presence of α 1,2-mannosidase inhibitors treated with GlcNAc-phosphotransferase, and acid α -glucosidase secreted from CHO cells in the presence of α 1,2-mannosidase inhibitors treated with GlcNAc-phosphotransferase and phosphodiester α -GlcNAcase. Equal amounts of the four different preparations are added to each well and incubated at 37°C for periods varying from 5 minutes to 4 hours. At the end of each incubation period the cell monolayers are washed

with phosphate buffered saline containing 5 mM mannose 6-phosphate and the monolayer solubilized in 1% Triton X^{*}-100 and assayed for internalized acid α -glucosidase by enzymatic assay.

Applicant and the assignee acknowledge their responsibility to replace these cultures should they die before the end of the term of a patent issued hereon, 5 years after the last request for a culture, or 30 years, whichever is the longer, and their responsibility to notify the depository of the issuance of such a patent, at which time the deposit will be made irrevocably available to the public. Until that time the deposit will be made available to the Commissioner of Patents under the terms of 37 C.F.R. 1.14 and 35 U.S.C. 112.

While the preferred embodiments are shown to illustrate the invention, numerous changes to the materials and methods can be made by those skilled in the art. All such changes are encompassed within the spirit of the invention as defined by the appended claims.

This application claims priority to U.S. Provisional application No. 60/153,831 filed September 14, 1999.

SEQUENCE LISTING

<110> CANFIELD, William M

<120> LYSOSOMAL TARGETING PATHWAY ENZYMES

<130> 196351WO77

<140> PCT/US00/21970

<141> 2000-08-10

<160> 52

<170> PatentIn Ver. 2.1

<210> 1

<211> 928

<212> PRT

<213> Homo sapiens

<400> 1

Met	Leu	Phe	Lys	Leu	Leu	Gln	Arg	Gln	Thr	Tyr	Thr	Cys	Leu	Ser	His
1				5					10					15	
Arg	Tyr	Gly	Leu	Tyr	Val	Cys	Phe	Leu	Gly	Val	Val	Val	Thr	Ile	Val
			20					25						30	
Ser	Ala	Phe	Gln	Phe	Gly	Glu	Val	Val	Leu	Glu	Trp	Ser	Arg	Asp	Gln
		35					40					45			
Tyr	His	Val	Leu	Phe	Asp	Ser	Tyr	Arg	Asp	Asn	Ile	Ala	Gly	Lys	Ser
	50					55					60				
Phe	Gln	Asn	Arg	Leu	Cys	Leu	Pro	Met	Pro	Ile	Asp	Val	Val	Tyr	Thr
	65				70					75					80
Trp	Val	Asn	Gly	Thr	Asp	Leu	Glu	Leu	Leu	Lys	Glu	Leu	Gln	Gln	Val
				85				90						95	
Arg	Glu	Gln	Met	Glu	Glu	Glu	Gln	Lys	Ala	Met	Arg	Glu	Ile	Leu	Gly
			100					105					110		
Lys	Asn	Thr	Thr	Glu	Pro	Thr	Lys	Lys	Ser	Glu	Lys	Gln	Leu	Glu	Cys
		115					120					125			
Leu	Leu	Thr	His	Cys	Ile	Lys	Val	Pro	Met	Leu	Val	Leu	Asp	Pro	Ala
	130					135					140				
Leu	Pro	Ala	Asn	Ile	Thr	Leu	Lys	Asp	Val	Pro	Ser	Leu	Tyr	Pro	Ser
145					150					155					160
Phe	His	Ser	Ala	Ser	Asp	Ile	Phe	Asn	Val	Ala	Lys	Pro	Lys	Asn	Pro
				165					170					175	
Ser	Thr	Asn	Val	Ser	Val	Val	Val	Phe	Asp	Ser	Thr	Lys	Asp	Val	Glu
			180					185					190		
Asp	Ala	His	Ser	Gly	Leu	Leu	Lys	Gly	Asn	Ser	Arg	Gln	Thr	Val	Trp
		195					200					205			
Arg	Gly	Tyr	Leu	Thr	Thr	Asp	Lys	Glu	Val	Pro	Gly	Leu	Val	Leu	Met

210	215	220
Gln Asp Leu Ala Phe Leu Ser Gly Phe Pro Pro Thr Phe Lys Glu Thr 225 230 235 240		
Asn Gln Leu Lys Thr Lys Leu Pro Glu Asn Leu Ser Ser Lys Val Lys 245 250 255		
Leu Leu Gln Leu Tyr Ser Glu Ala Ser Val Ala Leu Leu Lys Leu Asn 260 265 270		
Asn Pro Lys Asp Phe Gln Glu Leu Asn Lys Gln Thr Lys Lys Asn Met 275 280 285		
Thr Ile Asp Gly Lys Glu Leu Thr Ile Ser Pro Ala Tyr Leu Leu Trp 290 295 300		
Asp Leu Ser Ala Ile Ser Gln Ser Lys Gln Asp Glu Asp Ile Ser Ala 305 310 315 320		
Ser Arg Phe Glu Asp Asn Glu Glu Leu Arg Tyr Ser Leu Arg Ser Ile 325 330 335		
Glu Arg His Ala Pro Trp Val Arg Asn Ile Phe Ile Val Thr Asn Gly 340 345 350		
Gln Ile Pro Ser Trp Leu Asn Leu Asp Asn Pro Arg Val Thr Ile Val 355 360 365		
Thr His Gln Asp Val Phe Arg Asn Leu Ser His Leu Pro Thr Phe Ser 370 375 380		
Ser Pro Ala Ile Glu Ser His Ile His Arg Ile Glu Gly Leu Ser Gln 385 390 395 400		
Lys Phe Ile Tyr Leu Asn Asp Asp Val Met Phe Gly Lys Asp Val Trp 405 410 415		
Pro Asp Asp Phe Tyr Ser His Ser Lys Gly Gln Lys Val Tyr Leu Thr 420 425 430		
Trp Pro Val Pro Asn Cys Ala Glu Gly Cys Pro Gly Ser Trp Ile Lys 435 440 445		
Asp Gly Tyr Cys Asp Lys Ala Cys Asn Asn Ser Ala Cys Asp Trp Asp 450 455 460		
Gly Gly Asp Cys Ser Gly Asn Ser Gly Gly Ser Arg Tyr Ile Ala Gly 465 470 475 480		
Gly Gly Gly Thr Gly Ser Ile Gly Val Gly His Pro Trp Gln Phe Gly 485 490 495		
Gly Gly Ile Asn Ser Val Ser Tyr Cys Asn Gln Gly Cys Ala Asn Ser 500 505 510		
Trp Leu Ala Asp Lys Phe Cys Asp Gln Ala Cys Asn Val Leu Ser Cys 515 520 525		
Gly Phe Asp Ala Gly Asp Cys Gly Gln Asp His Phe His Glu Leu Tyr 530 535 540		

Lys	Val	Ile	Leu	Leu	Pro	Asn	Gln	Thr	His	Tyr	Ile	Ile	Pro	Lys	Gly	545	550	555	560
Glu	Cys	Leu	Pro	Tyr	Phe	Ser	Phe	Ala	Glu	Val	Ala	Lys	Arg	Gly	Val	565	570	575	
Glu	Gly	Ala	Tyr	Ser	Asp	Asn	Pro	Ile	Ile	Arg	His	Ala	Ser	Ile	Ala	580	585	590	
Asn	Lys	Trp	Lys	Thr	Ile	His	Leu	Ile	Met	His	Ser	Gly	Met	Asn	Ala	595	600	605	
Thr	Thr	Ile	His	Phe	Asn	Leu	Thr	Phe	Gln	Asn	Thr	Asn	Asp	Glu	Glu	610	615	620	
Phe	Lys	Met	Gln	Ile	Thr	Val	Glu	Val	Asp	Thr	Arg	Glu	Gly	Pro	Lys	625	630	635	640
Leu	Asn	Ser	Thr	Ala	Gln	Lys	Gly	Tyr	Glu	Asn	Leu	Val	Ser	Pro	Ile	645	650	655	
Thr	Leu	Leu	Pro	Glu	Ala	Glu	Ile	Leu	Phe	Glu	Asp	Ile	Pro	Lys	Glu	660	665	670	
Lys	Arg	Phe	Pro	Lys	Phe	Lys	Arg	His	Asp	Val	Asn	Ser	Thr	Arg	Arg	675	680	685	
Ala	Gln	Glu	Glu	Val	Lys	Ile	Pro	Leu	Val	Asn	Ile	Ser	Leu	Leu	Pro	690	695	700	
Lys	Asp	Ala	Gln	Leu	Ser	Leu	Asn	Thr	Leu	Asp	Leu	Gln	Leu	Glu	His	705	710	715	720
Gly	Asp	Ile	Thr	Leu	Lys	Gly	Tyr	Asn	Leu	Ser	Lys	Ser	Ala	Leu	Leu	725	730	735	
Arg	Ser	Phe	Leu	Met	Asn	Ser	Gln	His	Ala	Lys	Ile	Lys	Asn	Gln	Ala	740	745	750	
Ile	Ile	Thr	Asp	Glu	Thr	Asn	Asp	Ser	Leu	Val	Ala	Pro	Gln	Glu	Lys	755	760	765	
Gln	Val	His	Lys	Ser	Ile	Leu	Pro	Asn	Ser	Leu	Gly	Val	Ser	Glu	Arg	770	775	780	
Leu	Gln	Arg	Leu	Thr	Phe	Pro	Ala	Val	Ser	Val	Lys	Val	Asn	Gly	His	785	790	795	800
Asp	Gln	Gly	Gln	Asn	Pro	Pro	Leu	Asp	Leu	Glu	Thr	Thr	Ala	Arg	Phe	805	810	815	
Arg	Val	Glu	Thr	His	Thr	Gln	Lys	Thr	Ile	Gly	Gly	Asn	Val	Thr	Lys	820	825	830	
Glu	Lys	Pro	Pro	Ser	Leu	Ile	Val	Pro	Leu	Glu	Ser	Gln	Met	Thr	Lys	835	840	845	
Glu	Lys	Lys	Ile	Thr	Gly	Lys	Glu	Lys	Glu	Asn	Ser	Arg	Met	Glu	Glu	850	855	860	

Asn Ala Glu Asn His Ile Gly Val Thr Glu Val Leu Leu Gly Arg Lys
 865 870 875 880
 Leu Gln His Tyr Thr Asp Ser Tyr Leu Gly Phe Leu Pro Trp Glu Lys
 885 890 895
 Lys Lys Tyr Phe Gln Asp Leu Leu Asp Glu Glu Glu Ser Leu Lys Thr
 900 905 910
 Gln Leu Ala Tyr Phe Thr Asp Ser Lys Asn Thr Gly Arg Gln Leu Lys
 915 920 925

<210> 2
 <211> 328
 <212> PRT
 <213> Homo sapiens

<400> 2
 Asp Thr Phe Ala Asp Ser Leu Arg Tyr Val Asn Lys Ile Leu Asn Ser
 1 5 10 15
 Lys Phe Gly Phe Thr Ser Arg Lys Val Pro Ala His Met Pro His Met
 20 25 30
 Ile Asp Arg Ile Val Met Gln Glu Leu Gln Asp Met Phe Pro Glu Glu
 35 40 45
 Phe Asp Lys Thr Ser Phe His Lys Val Arg His Ser Glu Asp Met Gln
 50 55 60
 Phe Ala Phe Ser Tyr Phe Tyr Tyr Leu Met Ser Ala Val Gln Pro Leu
 65 70 75 80
 Asn Ile Ser Gln Val Phe Asp Glu Val Asp Thr Asp Gln Ser Gly Val
 85 90 95
 Leu Ser Asp Arg Glu Ile Arg Thr Leu Ala Thr Arg Ile His Glu Leu
 100 105 110
 Pro Leu Ser Leu Gln Asp Leu Thr Gly Leu Glu His Met Leu Ile Asn
 115 120 125
 Cys Ser Lys Met Leu Pro Ala Asp Ile Thr Gln Leu Asn Asn Ile Pro
 130 135 140
 Pro Thr Gln Glu Ser Tyr Tyr Asp Pro Asn Leu Pro Pro Val Thr Lys
 145 150 155 160
 Ser Leu Val Thr Asn Cys Lys Pro Val Thr Asp Lys Ile His Lys Ala
 165 170 175
 Tyr Lys Asp Lys Asn Lys Tyr Arg Phe Glu Ile Met Gly Glu Glu Glu
 180 185 190
 Ile Ala Phe Lys Met Ile Arg Thr Asn Val Ser His Val Val Gly Gln
 195 200 205

Leu Asp Asp Ile Arg Lys Asn Pro Arg Lys Phe Val Cys Leu Asn Asp
 210 215 220
 Asn Ile Asp His Asn His Lys Asp Ala Gln Thr Val Lys Ala Val Leu
 225 230 235 240
 Arg Asp Phe Tyr Glu Ser Met Phe Pro Ile Pro Ser Gln Phe Glu Leu
 245 250 255
 Pro Arg Glu Tyr Arg Asn Arg Phe Leu His Met His Glu Leu Gln Glu
 260 265 270
 Trp Arg Ala Tyr Arg Asp Lys Leu Lys Phe Trp Thr His Cys Val Leu
 275 280 285
 Ala Thr Leu Ile Met Phe Thr Ile Phe Ser Phe Phe Ala Glu Gln Leu
 290 295 300
 Ile Ala Leu Lys Arg Lys Ile Phe Pro Arg Arg Arg Ile His Lys Glu
 305 310 315 320
 Ala Ser Pro Asn Arg Ile Arg Val
 325

<210> 3
 <211> 305
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SIGNAL
 <222> (1)..(24)

<400> 3
 Met Ala Ala Gly Leu Ala Arg Leu Leu Leu Leu Leu Gly Leu Ser Ala
 1 5 10 15
 Gly Gly Pro Ala Pro Ala Gly Ala Ala Lys Met Lys Val Val Glu Glu
 20 25 30
 Pro Asn Ala Phe Gly Val Asn Asn Pro Phe Leu Pro Gln Ala Ser Arg
 35 40 45
 Leu Gln Ala Lys Arg Asp Pro Ser Pro Val Ser Gly Pro Val His Leu
 50 55 60
 Phe Arg Leu Ser Gly Lys Cys Phe Ser Leu Val Glu Ser Thr Tyr Lys
 65 70 75 80
 Tyr Glu Phe Cys Pro Phe His Asn Val Thr Gln His Glu Gln Thr Phe
 85 90 95
 Arg Trp Asn Ala Tyr Ser Gly Ile Leu Gly Ile Trp His Glu Trp Glu
 100 105 110
 Ile Ala Asn Asn Thr Phe Thr Gly Met Trp Met Arg Asp Gly Asp Ala
 115 120 125
 Cys Arg Ser Arg Ser Arg Gln Ser Lys Val Glu Leu Ala Cys Gly Lys
 130 135 140

Ser Asn Arg Leu Ala His Val Ser Glu Pro Ser Thr Cys Val Tyr Ala
145 150 155 160

Leu Thr Phe Glu Thr Pro Leu Val Cys His Pro His Ala Leu Leu Val
165 170 175

Tyr Pro Thr Leu Pro Glu Ala Leu Gln Arg Gln Trp Asp Gln Val Glu
180 185 190

Gln Asp Leu Ala Asp Glu Leu Ile Thr Pro Gln Gly His Glu Lys Leu
195 200 205

Leu Arg Thr Leu Phe Glu Asp Ala Gly Tyr Leu Lys Thr Pro Glu Glu
210 215 220

Asn Glu Pro Thr Gln Leu Glu Gly Gly Pro Asp Ser Leu Gly Phe Glu
225 230 235 240

Thr Leu Glu Asn Cys Arg Lys Ala His Lys Glu Leu Ser Lys Glu Ile
245 250 255

Lys Arg Leu Lys Gly Leu Leu Thr Gln His Gly Ile Pro Tyr Thr Arg
260 265 270

Pro Thr Glu Thr Ser Asn Leu Glu His Leu Gly His Glu Thr Pro Arg
275 280 285

Ala Lys Ser Pro Glu Gln Leu Arg Gly Asp Pro Gly Leu Arg Gly Ser
290 295 300

Leu
305

<210> 4

<211> 5597

<212> DNA

<213> Homo sapiens

<400> 4

```
cggagccgag cgggcgtccg tcgccggagc tgcaatgagc ggcgcccggga ggctgtgacc 60
tgcgcgcggc ggcccgaccg gggcccctga atggcggctc gctgaggcgg cggcggcggc 120
ggcggctcag gctcctcggg gcgtggcgtg gcggtgaagg ggtgatgctg ttcaagctcc 180
tgcagagaca aacctatacc tgccctgtccc acaggtatgg gctctacgtg tgcttcttgg 240
gcgtcgttgt caccatcgtc tccgccttcc agttcggaga ggtgggttctg gaatggagcc 300
gagatcaata ccatgttttg tttgattcct atagagacaa tattgctgga aagtcctttc 360
agaatcggct ttgtctgccc atgccgattg acgttggtta cacctgggtg aatggcacag 420
atcttgaact actgaaggaa ctacagcagg tcagagaaca gatggaggag gagcagaaag 480
caatgagaga aatccttggg aaaaacacaa cggaacctac taagaagagt gagaagcagt 540
tagagtgttt gctaacacac tgcattaagg tgccaatgct tgtactggac ccagccctgc 600
cagccaacat caccctgaag gacgtgccat ctctttatcc ttcttttcat tctgccagtg 660
acattttcaa tgttgcaaaa ccaaaaaacc cttctaccaa tgtctcagtt gttgtttttg 720
acagtactaa ggatgttgaa gatgccact ctggactgct taaaggaaat agcagacaga 780
cagtatggag ggggtacttg acaacagata aagaagtccc tggattagtg ctaatgcaag 840
atttggtttt cctgagtgga tttccaccaa cattcaagga acaaatcaa ctaaaaacaa 900
aattgccaga aaatctttcc tctaaagtca aactgttgca gttgtattca gaggccagtg 960
tagcgcttct aaaactgaat aaccccaagg attttcaaga attgaataag caaactaaga 1020
agaacatgac cattgatgga aaagaactga ccataagtcc tgcatattta ttatgggatc 1080
tgagcgccat cagccagtct aagcaggatg aagacatctc tgccagtcgt tttgaagata 1140
acgaagaact gaggtactca ttgcgatcta tcgagaggca tgcacatgg gttcgggaata 1200
```

ttttcattgt	caccaacggg	cagattccat	cctggctgaa	ccttgacaat	cctcgagtga	1260
caatagtaac	acaccaggat	gttttttcgaa	atttgagcca	cttgccctacc	tttagttcac	1320
ctgctattga	aagtcacatt	catcgcatcg	aagggctgtc	ccagaagttt	atttacctaa	1380
atgatgatgt	catgtttggg	aaggatgtct	ggccagatga	tttttacagt	cactccaaag	1440
gccagaaggt	ttatttgaca	tggcctgtgc	caaactgtgc	cgagggctgc	ccaggttcct	1500
ggattaagga	tggctattgt	gacaaggcct	gtaataattc	agcctgcat	tgggatgggtg	1560
gggattgctc	tggaaacagt	ggagggagtc	gctatattgc	aggaggtgga	ggtactggga	1620
gtattggagt	tggacacccc	tggcagtttg	gtggaggaat	aaacagtgtc	tcttactgta	1680
atcagggatg	tgcgaattcc	tggctcgctg	ataagtctctg	tgaccaagca	tgcaatgtct	1740
tgtcctgtgg	gtttgatgct	ggcgactgtg	ggcaagatca	ttttcatgaa	ttgtataaag	1800
tgatecttct	cccaaaccag	actcactata	ttattccaaa	aggtgaatgc	ctgccttatt	1860
tcagctttgc	agaagtagcc	aaaagaggag	ttgaagggtgc	ctatagtgc	aatccaataa	1920
ttcgacatgc	ttctattgcc	aacaagtgga	aaaccatcca	cctcataatg	cacagtggaa	1980
tgaatgccac	cacaatacat	tttaattctca	cgtttcaaaa	tacaaacgat	gaagagttca	2040
aaatgcagat	aacagtggag	gtggacacaa	gggagggacc	aaaactgaat	tctacggccc	2100
agaaggggta	cgaaaattta	gttagtccca	taacacttct	tccagaggcg	gaaatccttt	2160
ttgaggatat	tcccaaagaa	aaacgcttcc	cgaagttaa	gagacatgat	gttaactcaa	2220
caaggagagc	ccaggaagag	gtgaaaattc	ccctggtaaa	tatttcactc	cttccaaaag	2280
acgcccagtt	gagtcctcaat	accttggatt	tgcaactgga	acatggagac	atcactttga	2340
aaggatacaa	tttgtccaag	tcagccttgc	tgagatcatt	tctgatgaac	tcacagcatg	2400
ctaaaataaa	aaatcaagct	ataataacag	atgaaacaaa	tgacagtttg	gtggctccac	2460
aggaaaaaca	ggttcataaa	agcatcttgc	caaacagctt	aggagtgtct	gaaagattgc	2520
agagggtgac	ttttcctgca	gtgagtgtaa	aagtgaatgg	tcatgaccag	ggtcagaatc	2580
caccctgga	cttggagacc	acagcaagat	ttagagtgga	aactcacacc	caaaaaacca	2640
taggcggaaa	tgtgacaaaa	gaaaagcccc	catctctgat	tgttccactg	gaaagccaga	2700
tgacaaaaga	aaagaaaatc	acagggaaag	aaaaagagaa	cagtagaatg	gaggaaaatg	2760
ctgaaaatca	cataggcggt	actgaagtgt	tacttggaa	aaagctgcag	cattacacag	2820
atagttactt	gggctttttg	ccatgggaga	aaaaaaagta	tttccaagat	cttctcgacg	2880
aagaagagtc	attgaagaca	caattggcat	acttcactga	tagcaaaaat	actgggaggc	2940
aactaaaaga	tacatttgca	gattccctca	gatatgtaaa	taaaattcta	aatagcaagt	3000
ttggattcac	atcgcgga	gtccctgctc	acatgcctca	catgattgac	cggattgtta	3060
tgcaagaact	gcaagatatg	ttccctgaag	aatttgacaa	gacgtcattt	cacaaagtgc	3120
gccattctga	ggatatgcag	tttgcccttct	cttattttta	ttatctcatg	agtgcagtgc	3180
agccactgaa	tatatctcaa	gtctttgatg	aagttgatac	agatcaatct	ggtgtcttgt	3240
ctgacagaga	aatccgaaca	ctggctacca	gaattcacga	actgccgtta	agtttgcagg	3300
at ttgacagg	tctggaacac	atgctaataa	attgctcaaa	aatgcttcct	gctgatataca	3360
cgcagctaaa	taatattcca	ccaactcagg	aatcctacta	tgatcccaac	ctgccaccgg	3420
tcactaaaag	tctagtaaca	aactgtaaac	cagtaactga	caaaatccac	aaagcatata	3480
aggacaaaaa	caaatatagg	tttgaaatca	tgggagaaga	agaaatcgct	tttaaaatga	3540
ttcgtaccaa	cgtttctcat	gtgggttgcc	agttggatga	cataagaaaa	aaccctagga	3600
agtttgtttg	cctgaatgac	aacattgacc	acaatcataa	agatgctcag	acagtgaagg	3660
ctgttctcag	ggacttctat	gaatccatgt	tccccatacc	ttcccaattt	gaactgccaa	3720
gagagtatcg	aaaccgtttc	cttcatatgc	atgagctgca	ggaatggagg	gcttatcgag	3780
acaaattgaa	gttttgagacc	cattgtgtac	tagcaacatt	gattatgttt	actatattct	3840
catttttttg	tgagcagtta	attgcactta	agcggaagat	atttcccaga	aggaggatac	3900
acaaagaagc	tagtcccaat	cgaatcagag	tatagaagat	cttcatttga	aaaccatcta	3960
cctcagcatt	tactgagcat	tttaaaactc	agcttcacag	agatgtcttt	gtgatgtgat	4020
gcttagcagt	ttggcccga	gaaggaaaat	atccagttacc	atgctgtttt	gtggcatgaa	4080
tatagcccac	tgactaggaa	ttatttaacc	aaccactga	aaacttgtgt	gtcgagcagc	4140
tctgaactga	ttttactttt	aaagaatttg	ctcatggacc	tgtcatcctt	tttataaaaa	4200
ggctcactga	caagagacag	ctgttaattt	cccacagcaa	tcattgcaga	ctaactttat	4260
taggagaagc	ctatgccagc	tgggagtgat	tgctaagagg	ctccagtcct	tgcatcccaa	4320
agccttttgc	taaagttttg	cacttttttt	ttttcatttc	ccatttttaa	gtagttacta	4380
agttaactag	ttattcttgc	ttctgagtat	aacgaattgg	gatgtctaaa	cctattttta	4440
tagatgttat	ttaaataatg	cagcaatatc	acctcttatt	gacaatacct	aaattatgag	4500
ttttattaat	atttaagact	gtaaatggtc	ttaaaccact	aactactgaa	gagctcaatg	4560
attgacatct	gaaatgcttt	gtaattattg	acttcagccc	ctaagaatgc	tatgatttca	4620
cgtgcaggtc	taatttcaac	aggctagagt	tagtactact	taccagatgt	aattatgttt	4680
tggaaatgta	catattcaaa	cagaagtgcc	tcatttttaga	aatgagtagt	gctgatggca	4740
ctggcacatt	acagtgggtg	cttgtttaat	actcattggg	atattccagt	agctatctct	4800
ctcagttggg	ttttgataga	acagaggcca	gcaaactttc	tttgtaaaag	gctgggttagt	4860

```

aaattattgc aggccacctg tgtctttgtc atacattctt cttgctgttg tttagtttgt 4920
tttttttcaa acaaccctct aaaaatgtaa aaaccatgtt tagcttgcag ctgtacaaaa 4980
actgcccacc agccagatgt gaccctcagg ccatcatttg ccaatcactg agaattattt 5040
ttgttggtgt tgttggtgtt gtttttgaga cagagtctct ctctgttgcc caggctggag 5100
tgcagtggcg caatctcagc tcaactgcaac ctccgcctcc cgggttcaag cagttctgtc 5160
tcagccttct gagtagctgg gactacaggt gcatgccacc acaccctgct aatttttgta 5220
tttttagtag agacgggggt tccaccatat tggtcaggct tatcttgaac tcctgacctc 5280
aggtgatcca cctgcctctg cctcccaaag tgctgagatt acaggcataa gccagtgcac 5340
ccagccgaga attagtattt ttatgtatgg ttaaaccctg gcgtctagcc atattttatg 5400
tcataataca atggatttgt gaagagcaga ttccatgagt aactctgaca ggtatttttag 5460
atcatgatct caacaatatt cctcccaaat ggcatacatc ttttgtacaa agaacttgaa 5520
atgtaaatac tgtgtttgtg ctgtaagagt tgtgtatttc aaaaactgaa atctcataaa 5580
aagttaaatt ttgaaaa 5597

```

```

<210> 5
<211> 1219
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> sig_peptide
<222> (24)..(95)

```

```

<400> 5
gtagagcgca ggtgcgcggc tcgatggcgg cggggctggc gcggctcctg ttgctcctcg 60
ggctctcggc cggcggggccc gcgccggcag gtgcagcgaa gatgaagggt gtggaggagc 120
ccaacgcgtt tgggggtgaac aaccgcgttct tgcctcaggc cagtcgcctc caggccaaga 180
gggatccttc acccgtgtct ggaccgcgtc atctcttccg actctcgggc aagtgccttca 240
gcctggtgga gtccacgtac aagtatgagt tctgcccgtt ccacaacgtg acccagcacg 300
agcagacctt ccgctggaac gcctacagtg ggatcctcgg catctggcac gagtgggaga 360
tcgccaaaca caccttcacg ggcattgtga tgagggacgg tgacgcctgc cgttcccgga 420
gccggcagag caaggtggag ctggcgtgtg gaaaaagcaa ccggtcggcc catgtgtccg 480
agccgagcac ctgcgtctat gcgctgacgt tcgagacccc cctcgtctgc ccccccacg 540
ccttgctagt gtaccaacc ctgccagagg ccctgcagcg gcagtgggac caggtagagc 600
aggacctggc cgatgagctg atcaccccc aggcccatga gaagttgctg aggacacttt 660
ttgaggatgc tggctactta aagaccccag aagaaaatga acccaccag ctggaggagg 720
gtcctgacag cttgggggtt gagaccctgg aaaactgcag gaaggctcat aaagaactct 780
caaaggagat caaaaggctg aaaggtttgc tcaccagca cggcatcccc tacacgaggc 840
ccacagaaac ttccaacttg gagcacttgg gccacgagac gccagagacc aagtctccag 900
agcagctgcg gggtgaccca ggactgcgtg ggagtttgtg acctgttgtt gggagagcag 960
aggtggacgc ggccgagagc cctacagaga agctggctgg taggaccgcg aggaccagct 1020
gaccaggctt gtgctcagag aagcagacaa aacaaagatt caaggtttta attaattccc 1080
atactgataa aaataactcc atgaattctg taaaccattg cataaatgct atagtgtaaa 1140
aaaattttaa caagtgttaa ctttaaacag ttcgctacaa gtaaatagatt ataaatacta 1200
aaaaaaaaaa aaaaaaaaaa 1219

```

```

<210> 6
<211> 515
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> SIGNAL
<222> (1)..(24)

```

```

<220>
<221> PROPEP
<222> (25)..(49)

```

<400> 6

```

Met Ala Thr Ser Thr Gly Arg Trp Leu Leu Leu Arg Leu Ala Leu Phe
 1           5           10           15

Gly Phe Leu Trp Glu Ala Ser Gly Gly Leu Asp Ser Gly Ala Ser Arg
 20           25           30

Asp Asp Asp Leu Leu Leu Pro Tyr Pro Arg Ala Arg Ala Arg Leu Pro
 35           40           45

Arg Asp Cys Thr Arg Val Arg Ala Gly Asn Arg Glu His Glu Ser Trp
 50           55           60

Pro Pro Pro Pro Ala Thr Pro Gly Ala Gly Gly Leu Ala Val Arg Thr
 65           70           75           80

Phe Val Ser His Phe Arg Asp Arg Ala Val Ala Gly His Leu Thr Arg
 85           90           95

Ala Val Glu Pro Leu Arg Thr Phe Ser Val Leu Glu Pro Gly Gly Pro
100           105           110

Gly Gly Cys Ala Ala Arg Arg Arg Ala Thr Val Glu Glu Thr Ala Arg
115           120           125

Ala Ala Asp Cys Arg Val Ala Gln Asn Gly Gly Phe Phe Arg Met Asn
130           135           140

Ser Gly Glu Cys Leu Gly Asn Val Val Ser Asp Glu Arg Arg Val Ser
145           150           155           160

Ser Ser Gly Gly Leu Gln Asn Ala Gln Phe Gly Ile Arg Arg Asp Gly
165           170           175

Thr Leu Val Thr Gly Tyr Leu Ser Glu Glu Glu Val Leu Asp Thr Glu
180           185           190

Asn Pro Phe Val Gln Leu Leu Ser Gly Val Val Trp Leu Ile Arg Asn
195           200           205

Gly Ser Ile Tyr Ile Asn Glu Ser Gln Ala Thr Glu Cys Asp Glu Thr
210           215           220

Gln Glu Thr Gly Ser Phe Ser Lys Phe Val Asn Val Ile Ser Ala Arg
225           230           235           240

Thr Ala Ile Gly His Asp Arg Lys Gly Gln Leu Val Leu Phe His Ala
245           250           255

Asp Gly His Thr Glu Gln Arg Gly Ile Asn Leu Trp Glu Met Ala Glu
260           265           270

Phe Leu Leu Lys Gln Asp Val Val Asn Ala Ile Asn Leu Asp Gly Gly
275           280           285

Gly Ser Ala Thr Phe Val Leu Asn Gly Thr Leu Ala Ser Tyr Pro Ser
290           295           300

Asp His Cys Gln Asp Asn Met Trp Arg Cys Pro Arg Gln Val Ser Thr
305           310           315           320

```

Val Val Cys Val His Glu Pro Arg Cys Gln Pro Pro Asp Cys His Gly
 325 330 335
 His Gly Thr Cys Val Asp Gly His Cys Gln Cys Thr Gly His Phe Trp
 340 345 350
 Arg Gly Pro Gly Cys Asp Glu Leu Asp Cys Gly Pro Ser Asn Cys Ser
 355 360 365
 Gln His Gly Leu Cys Thr Glu Thr Gly Cys Arg Cys Asp Ala Gly Trp
 370 375 380
 Thr Gly Ser Asn Cys Ser Glu Glu Cys Pro Leu Gly Trp His Gly Pro
 385 390 395 400
 Gly Cys Gln Arg Arg Cys Lys Cys Glu His His Cys Pro Cys Asp Pro
 405 410 415
 Lys Thr Gly Asn Cys Ser Val Ser Arg Val Lys Gln Cys Leu Gln Pro
 420 425 430
 Pro Glu Ala Thr Leu Arg Ala Gly Glu Leu Ser Phe Phe Thr Arg Thr
 435 440 445
 Ala Trp Leu Ala Leu Thr Leu Ala Leu Ala Phe Leu Leu Leu Ile Ser
 450 455 460
 Ile Ala Ala Asn Leu Ser Leu Leu Leu Ser Arg Ala Glu Arg Asn Arg
 465 470 475 480
 Arg Leu His Gly Asp Tyr Ala Tyr His Pro Leu Gln Glu Met Asn Gly
 485 490 495
 Glu Pro Leu Ala Ala Glu Lys Glu Gln Pro Gly Gly Ala His Asn Pro
 500 505 510
 Phe Lys Asp
 515

<210> 7
 <211> 2183
 <212> DNA
 <213> Homo sapiens

<400> 7
 atggcgacct ccacgggtcg ctggcttctc ctccggcttg cactattcgg cttcctctgg 60
 gaagcgctccg gcggcctcga ctcggggggcc tcccgcgacg acgacttgct actgccctat 120
 ccacgcgcgc gcgcgcgcct cccccgggac tgcacacggg tgcgcgccgg caaccgcgag 180
 cacgagagtt ggcctccgcc tcccgcgact cccggcgccg gcggtctggc cgtgcgcacc 240
 ttcgtgtcgc acttcaggga ccgcgcggtg gccggccacc tgacgcgggc cgttgagccc 300
 ctgcgcacct tctcgggtgct ggagcccggg ggacccggcg gctgcgcggc gagacgacgc 360
 gccaccgtgg aggagacggc gcgggcggcc gactgccgtg tcgcccagaa cggcggcttc 420
 ttccgcatga actcgggcga gtgcctgggg aacgtggtga gcgacgagcg gcgggtgagc 480
 agctccgggg ggctgcagaa cgcgcagttc gggatccgcc gcgacgggac cctggtcacc 540
 gggtagctgt ctgaggagga ggtgctggac actgagaacc catttgtagc gctgctgagt 600
 ggggtcgtgt ggctgattcg taatggaagc atctacatca acgagagcca agccacagag 660
 tgtgacgaga cacaggagac aggttccttt agcaaatttg tgaatgtgat atcagccagg 720
 acggccattg gccacgaccg gaaagggcag ctggtgctct ttcattgcaga cggccatacg 780
 gagcagcgtg gcatcaacct gtgggaaatg gcggagttcc tgctgaaaca ggacgtggtc 840
 aacgccatca acctggatgg ggggtggctct gccacctttg tgctcaacgg gaccttggcc 900

```

agttaccctgt cagatcactg ccaggacaac atgtggcgct gtccccgcca agtgtccacc 960
gtggtgtgtg tgcacgaacc ccgctgccag ccgcctgact gccacggcca cgggacctgc 1020
gtggacgggc actgccaatg caccggggcac ttctggcggg gtcccggctg tgatgagctg 1080
gactgtggcc cctctaactg cagccagcac ggactgtgca cggagaccgg ctgccgctgt 1140
gatgccggat ggaccgggtc caactgcagt gaagagtgtc cccttggctg gcatggggccg 1200
ggctgccaga ggcgttgtaa gtgtgagcac cattgtccct gtgaccccaa gactggcaac 1260
tgcagcgtct ccagagtaaa gcagtgtctc cagccacctg aagccaccct gagggcgga 1320
gaactctcct ttttcaccag gaccgcctgg ctagccctca ccctggcgct ggccttcctc 1380
ctgctgatca gcattgcagc aaacctgtcc ttgctcctgt ccagagcaga gaggaaccgg 1440
cgctgcatg gggactatgc ataccacccg ctgcaggaga tgaacgggga gcctctggcc 1500
gcagagaagg agcagccagg gggcgccac aacccttca aggactgaag cctcaagctg 1560
cccggggtgg cacgtcgca aagcttggtt cccacggtc tggcttctgc aggggaaatt 1620
tcaaggccac tggcgtggac catctgggtg tcctcaatgg ccctgtggg gcagccaagt 1680
tcctgatagc acttgtgcct cagccctca cctggccacc tgccaggga cctgcaacct 1740
tagcaatacc atgctcgctg gagaggctca gctgcctgct tctcgctgc ctgtgtctgc 1800
tgccgagaag ccctgcccc cgggagggct gccgcactgc caaagagtct ccctcctcct 1860
ggggaagggg ctgccaacga accagactca gtgaccacgt catgacagaa cagcacatcc 1920
tggccagcac ccctggctgg agtgggttaa agggacgagt ctgccttcct ggctgtgaca 1980
cgggaccct tttctacaga cctcatcact ggatttgcca actagaattc gatttcctgt 2040
cataggaagc tccttggaag aagggatggg gggatgaaat catgtttaca gacctgtttt 2100
gtcatcctgc tgccaagaag ttttttaatc acttgaataa attgatataa taaaaggagc 2160
caccaggtgg tgtgtggatt ctg 2183

```

<210> 8

<211> 328

<212> PRT

<213> Mus musculus

<400> 8

```

Asp Thr Phe Ala Asp Ser Leu Arg Tyr Val Asn Lys Ile Leu Asn Ser
  1             5             10             15

```

```

Lys Phe Gly Phe Thr Ser Arg Lys Val Pro Ala His Met Pro His Met
      20             25             30

```

```

Ile Asp Arg Ile Val Met Gln Glu Leu Gln Asp Met Phe Pro Glu Glu
      35             40             45

```

```

Phe Asp Lys Thr Ser Phe His Lys Val Arg His Ser Glu Asp Met Gln
      50             55             60

```

```

Phe Ala Phe Ser Tyr Phe Tyr Tyr Leu Met Ser Ala Val Gln Pro Leu
      65             70             75             80

```

```

Asn Ile Ser Gln Val Phe His Glu Val Asp Thr Asp Gln Ser Gly Val
      85             90             95

```

```

Leu Ser Asp Arg Glu Ile Arg Thr Leu Ala Thr Arg Ile His Asp Leu
      100            105            110

```

```

Pro Leu Ser Leu Gln Asp Leu Thr Gly Leu Glu His Met Leu Ile Asn
      115            120            125

```

```

Cys Ser Lys Met Leu Pro Ala Asn Ile Thr Gln Leu Asn Asn Ile Pro
      130            135            140

```

```

Pro Thr Gln Glu Ala Tyr Tyr Asp Pro Asn Leu Pro Pro Val Thr Lys
      145            150            155            160

```

```

Ser Leu Val Thr Asn Cys Lys Pro Val Thr Asp Lys Ile His Lys Ala

```

165					170					175					
Tyr	Lys	Asp	Lys	Asn	Lys	Tyr	Arg	Phe	Glu	Ile	Met	Gly	Glu	Glu	Glu
			180					185					190		
Ile	Ala	Phe	Lys	Met	Ile	Arg	Thr	Asn	Val	Ser	His	Val	Val	Gly	Gln
			195				200					205			
Leu	Asp	Asp	Ile	Arg	Lys	Asn	Pro	Arg	Lys	Phe	Val	Cys	Leu	Asn	Asp
	210					215					220				
Asn	Ile	Asp	His	Asn	His	Lys	Asp	Ala	Arg	Thr	Val	Lys	Ala	Val	Leu
225				230					235						240
Arg	Asp	Phe	Tyr	Glu	Ser	Met	Phe	Pro	Ile	Pro	Ser	Gln	Phe	Glu	Leu
				245					250					255	
Pro	Arg	Glu	Tyr	Arg	Asn	Arg	Phe	Leu	His	Met	His	Glu	Leu	Gln	Glu
			260					265					270		
Trp	Arg	Ala	Tyr	Arg	Asp	Lys	Leu	Lys	Phe	Trp	Thr	His	Cys	Val	Leu
		275					280					285			
Ala	Thr	Leu	Ile	Ile	Phe	Thr	Ile	Phe	Ser	Phe	Phe	Ala	Glu	Gln	Ile
	290					295					300				
Ile	Ala	Leu	Lys	Arg	Lys	Ile	Phe	Pro	Arg	Arg	Arg	Ile	His	Lys	Glu
305				310					315						320
Ala	Ser	Pro	Asp	Arg	Ile	Arg	Val								
				325											

<210> 9
 <211> 307
 <212> PRT
 <213> Mus musculus

<400> 9															
Met	Ala	Gly	Arg	Leu	Ala	Gly	Phe	Leu	Met	Leu	Leu	Gly	Leu	Ala	Ser
1				5				10					15		
Gln	Gly	Pro	Ala	Pro	Ala	Cys	Ala	Gly	Lys	Met	Lys	Val	Val	Glu	Glu
			20					25					30		
Pro	Asn	Thr	Phe	Gly	Leu	Asn	Asn	Pro	Phe	Leu	Pro	Gln	Ala	Ser	Arg
		35					40					45			
Leu	Gln	Pro	Lys	Arg	Glu	Pro	Ser	Ala	Val	Ser	Gly	Pro	Leu	His	Leu
	50					55					60				
Phe	Arg	Leu	Ala	Gly	Lys	Cys	Phe	Ser	Leu	Val	Glu	Ser	Thr	Tyr	Lys
65					70					75					80
Tyr	Glu	Phe	Cys	Pro	Phe	His	Asn	Val	Thr	Gln	His	Glu	Gln	Thr	Phe
				85					90					95	
Arg	Trp	Asn	Ala	Tyr	Ser	Gly	Ile	Leu	Gly	Ile	Trp	His	Glu	Trp	Glu
			100					105					110		
Ile	Ile	Asn	Asn	Thr	Phe	Lys	Gly	Met	Trp	Met	Thr	Asp	Gly	Asp	Ser

115					120					125								
Cys	His	Ser	Arg	Ser	Arg	Gln	Ser	Lys	Val	Glu	Leu	Thr	Cys	Gly	Lys			
130					135					140								
Ile	Asn	Arg	Leu	Ala	His	Val	Ser	Glu	Pro	Ser	Thr	Cys	Val	Tyr	Ala			
145					150					155					160			
Leu	Thr	Phe	Glu	Thr	Pro	Leu	Val	Cys	His	Pro	His	Ser	Leu	Leu	Val			
					165					170					175			
Tyr	Pro	Thr	Leu	Ser	Glu	Ala	Leu	Gln	Gln	Arg	Leu	Asp	Gln	Val	Glu			
					180					185					190			
Gln	Asp	Leu	Ala	Asp	Glu	Leu	Ile	Thr	Pro	Gln	Gly	Tyr	Glu	Lys	Leu			
					195					200					205			
Leu	Arg	Val	Leu	Phe	Glu	Asp	Ala	Gly	Tyr	Leu	Lys	Val	Pro	Gly	Glu			
					210					215					220			
Thr	His	Pro	Thr	Gln	Leu	Ala	Gly	Gly	Ser	Lys	Gly	Leu	Gly	Leu	Glu			
					225					230					235		240	
Thr	Leu	Asp	Asn	Cys	Arg	Lys	Ala	His	Ala	Glu	Leu	Ser	Gln	Glu	Val			
					245					250					255			
Gln	Arg	Leu	Thr	Ser	Leu	Leu	Gln	Gln	His	Gly	Ile	Pro	His	Thr	Gln			
					260					265					270			
Pro	Thr	Glu	Thr	Thr	His	Ser	Gln	His	Leu	Gly	Gln	Gln	Leu	Pro	Ile			
					275					280					285			
Gly	Ala	Ile	Ala	Ala	Glu	His	Leu	Arg	Ser	Asp	Pro	Gly	Leu	Arg	Gly			
					290					295					300			

Asn Ile Leu
305

<210> 10
<211> 2070
<212> DNA
<213> Mus musculus

<220>
<221> misc_feature
<222> (186)
<223> N IS A, G, T OR C

<400> 10
gtgagaccct aggagcaatg gccgggcggc tggctggctt cctgatgttg ctggggctcg 60
cgtcgcaggg gcccgcgccg gcatgtgccg ggaagatgaa ggtggtggag gaggcctaaca 120
cattcgggtg agcggatcac ggtcctgcgg cttggggacc gaggcctggct gggttcttctg 180
accttntcaa ttccataggc tgaataaccc gttcttgccc caggcaagcc gccttcagcc 240
caagagagag ccttcagctg tatcccgcga attaagagaa attaatttca aacgatttag 300
aaagtattct agccaggcga tgatggcgca cgcctttaat cccagcactt gggaggcaga 360
ggcaggcaga tttccgagtt caaggccatc agaactgact gtacatctta gtacagttta 420
gcatgtgatc agagatctga atcacaaagc tgggcctgcg tggtaaagca ggtcctttct 480
aataagggtg cagtttagat tttctttctt aactctttta ttctttgaga cagggtttct 540
caacagtggg tgtcctggaa ctcacttttg taaaccaggc tgcccttaaa ctcacaaagc 600
tctgtcagcc tctgcctcct gagtgctggg attaaaggtc cacaccctgt tcattcattt 660

```

ttaatttttg agactgggtc tcattatgtg gccctagaca gatactgaga gcctcctcca 720
caggaacaag catgggaatc ctgccacaga caaccagttc tgtgggtctgg agatgagttt 780
gtcagtccct aggagttagg tcagcctgcc tctgcattcc caataattta ggaaaggagc 840
ttggggcggt ctggccttga tggttagtgc cctcctgcca accttagctt ccagcttttag 900
gggtagcaga gtttataaccg atgctaaact gctgttgtgt tcttccccag ggcccctgca 960
tctcttcaga cttgctggca agtgcttttag cctagtggag tccacgtgag tgccaggctg 1020
gtgggtggag tgggcggagt ctgcagagct cctgatgtgc ctgtgtttcc caggtacaag 1080
tatgaattct gccctttcca caacgtcacc cagcacgagc agaccttccg ctggaatgcc 1140
tacagcggga tccttggcat ctggcatgag tgggaaatca tcaacaatac cttcaagggc 1200
atgtggatga ctgatgggga ctccctgccac tcccggagcc ggcagagcaa ggtggagctc 1260
acctgtggaa agatcaaccg actggcccac gtgtctgagc caagcacctg tgtctatgca 1320
ttgacattcg agacctctct tgtttgccat cccactctt tgttagtgtg tccaactctg 1380
tcagaagccc tgcagcagcc cttggaccag gtggaacagg acctggcaga tgaactgatc 1440
acaccacagg gctatgagaa gttgctaagg gtactttttg aggatgctgg ctacttaaag 1500
gtcccaggag aaacctatcc caccagctg gcaggaggtt ccaagggcct ggggcttgag 1560
actctggaca actgtagaaa ggcacatgca gagctgtcac aggaggtaca aagactgacg 1620
agtctgctgc aacagcatgg aatccccac actcagccca caggtcagtc tgctgccct 1680
ggtcagctgc cagccactcc ggggcctgca gcactggggc agatctttat tgctacccat 1740
tctggcagaa accactcact ctcagcacct gggtcagcag ctccccatag gtgcaatcgc 1800
agcagagcat ctgcggagtg acccaggact acgtgggaac atcctgtgag caaggtggcc 1860
acgaagaata gaaatatcct gagcttttag tgtcctttca cagagtgaac aaaactggtg 1920
tggtgtagac acggcttctt ttggcatatt ctagatcaga cagtgtcact gacaaacaag 1980
agggacctgc tggccagcct ttgttgtgcc caaagatcca gacaaaataa agattcaaag 2040
ttttaattaa aaaaaaaaaa aaaggaattc 2070

```

<210> 11
 <211> 113
 <212> PRT
 <213> Rattus rattus

<400> 11
 Phe Pro Pro Thr Phe Lys Glu Thr Ser Gln Leu Lys Thr Lys Leu Pro
 1 5 10 15
 Glu Asn Leu Ser Ser Lys Ile Lys Leu Leu Gln Leu Tyr Ser Glu Ala
 20 25 30
 Ser Val Ala Leu Leu Lys Leu Asn Asn Pro Lys Gly Phe Pro Glu Leu
 35 40 45
 Asn Lys Gln Thr Lys Lys Asn Met Ser Ile Ser Gly Lys Glu Leu Ala
 50 55 60
 Ile Ser Pro Ala Tyr Leu Leu Trp Asp Leu Ser Ala Ile Ser Gln Ser
 65 70 75 80
 Lys Gln Asp Glu Asp Val Ser Ala Ser Arg Phe Glu Asp Asn Glu Glu
 85 90 95
 Leu Arg Tyr Ser Leu Arg Ser Ile Glu Arg His Asp Ser Met Ser Pro
 100 105 110
 Leu

<210> 12
 <211> 460
 <212> DNA
 <213> Rattus rattus

<400> 12

```

attccccacca acattcaagg agacgagtca gctgaagaca aaactgccag aaaatctttc 60
ttctaaaata aaactgttgc agctgtactc ggaggccagc gtcgctcttc tgaaattgaa 120
taaccccaaaa ggtttccccg agctgaacaa gcagaccaag aagaacatga gcatcagtgg 180
gaaggaactg gccatcagcc ctgcctatct gctgtgggac ctgagcgcca tcagccagtc 240
caagcaggat gaagatgtgt ctgccagccg cttcgaggat aacgaagagc tgaggtactc 300
actgagatct atcgagagac atgattccat gagtccttta tgaattctgg ccatatcttc 360
aatcatgatc tcagtagtat tcctctgaaa tggcacacat ttttctaata agaacttgaa 420
atgtaaataat tgtgtttgtg ctgtaaattt tgtgtatttc 460

```

<210> 13

<211> 502

<212> PRT

<213> *Drosophila melanogaster*

<400> 13

```

Gly Thr Arg Arg Phe Asp Asp Lys Asn Glu Leu Arg Tyr Ser Leu Arg
 1              5              10              15

Ser Leu Glu Lys His Ala Ala Trp Ile Arg His Val Tyr Ile Val Thr
      20              25              30

Asn Gly Gln Ile Pro Ser Trp Leu Asp Leu Ser Tyr Glu Arg Val Thr
      35              40              45

Val Val Pro His Glu Val Leu Ala Pro Asp Pro Asp Gln Leu Pro Thr
      50              55              60

Phe Ser Ser Ser Ala Ile Glu Thr Phe Leu His Arg Ile Pro Lys Leu
65              70              75              80

Ser Lys Arg Phe Leu Tyr Leu Asn Asp Asp Ile Phe Leu Gly Ala Pro
      85              90              95

Leu Tyr Pro Glu Asp Leu Tyr Thr Glu Ala Glu Gly Val Arg Val Tyr
      100             105             110

Gln Ala Trp Met Val Pro Gly Cys Ala Leu Asp Cys Pro Trp Thr Tyr
      115             120             125

Ile Gly Asp Gly Ala Cys Asp Arg His Cys Asn Ile Asp Ala Cys Gln
      130             135             140

Phe Asp Gly Gly Asp Cys Ser Glu Thr Gly Pro Ala Ser Asp Ala His
145             150             155             160

Val Ile Pro Pro Ser Lys Glu Val Leu Glu Val Gln Pro Ala Ala Val
      165             170             175

Pro Gln Ser Arg Val His Arg Phe Pro Gln Met Gly Leu Gln Lys Leu
      180             185             190

Phe Arg Arg Ser Ser Ala Asn Phe Lys Asp Val Met Arg His Arg Asn
195             200             205

Val Ser Thr Leu Lys Glu Leu Arg Arg Ile Val Glu Arg Phe Asn Lys
210             215             220

Ala Lys Leu Met Ser Leu Asn Pro Glu Leu Glu Thr Ser Ser Ser Glu

```

225		230		235		240									
Pro	Gln	Thr	Thr	Gln	Arg	His	Gly	Leu	Arg	Lys	Glu	Asp	Phe	Lys	Ser
				245					250					255	
Ser	Thr	Asp	Ile	Tyr	Ser	His	Ser	Leu	Ile	Ala	Thr	Asn	Met	Leu	Leu
			260					265					270		
Asn	Arg	Ala	Tyr	Gly	Phe	Lys	Ala	Arg	His	Val	Leu	Ala	His	Val	Gly
		275					280					285			
Phe	Leu	Ile	Asp	Lys	Asp	Ile	Val	Glu	Ala	Met	Gln	Arg	Arg	Phe	His
	290					295					300				
Gln	Gln	Ile	Leu	Asp	Thr	Ala	His	Gln	Arg	Phe	Arg	Ala	Pro	Thr	Asp
305					310					315					320
Leu	Gln	Tyr	Ala	Phe	Ala	Tyr	Tyr	Ser	Phe	Leu	Met	Ser	Glu	Thr	Lys
			325						330					335	
Val	Met	Ser	Val	Glu	Glu	Ile	Phe	Asp	Glu	Phe	Asp	Thr	Asp	Gly	Ser
			340					345					350		
Ala	Thr	Trp	Ser	Asp	Arg	Glu	Val	Arg	Thr	Phe	Leu	Thr	Arg	Ile	Tyr
		355					360					365			
Gln	Pro	Pro	Leu	Asp	Trp	Ser	Ala	Met	Arg	Tyr	Phe	Glu	Glu	Val	Val
	370					375					380				
Gln	Asn	Cys	Thr	Arg	Asn	Leu	Gly	Met	His	Leu	Lys	Val	Asp	Thr	Val
385					390					395					400
Glu	His	Ser	Thr	Leu	Val	Tyr	Glu	Arg	Tyr	Glu	Asp	Ser	Asn	Leu	Pro
				405					410					415	
Thr	Ile	Thr	Arg	Asp	Leu	Val	Val	Arg	Cys	Pro	Leu	Leu	Ala	Glu	Ala
			420					425					430		
Leu	Ala	Ala	Asn	Phe	Ala	Val	Arg	Pro	Lys	Tyr	Asn	Phe	His	Val	Ser
		435					440					445			
Pro	Lys	Arg	Thr	Ser	His	Ser	Asn	Phe	Met	Met	Leu	Thr	Ser	Asn	Leu
	450					455					460				
Thr	Glu	Val	Val	Glu	Ser	Leu	Asp	Arg	Leu	Arg	Arg	Asn	Pro	Arg	Lys
465					470					475					480
Phe	Asn	Cys	Ile	Asn	Asp	Asn	Leu	Asp	Ala	Asn	Arg	Gly	Glu	Asp	Asn
				485					490					495	
Glu	Asp	Gly	Ala	Pro	Ser										
			500												

<210> 14

<211> 9792

<212> DNA

<213> Mus musculus

<400> 14

caggctcggg acttactata acacaggaca cttgtcacct gaaagcttga gtcagtcagt 60

tattatggtc	tgtgtgtgag	atacaagtgg	gtgcataggc	agtgggtgcac	acatgtagat	120
cagactttct	acagccaatt	ctcttcttcc	tcctctccat	gggttcaggg	tcttcatctc	180
aggttgacac	gcgagttcat	ttatgtgctg	tgccatctcg	ccagtcgttc	ctatatccta	240
gaggaaaact	agtttcttct	ggtcaagagg	aggaaagagt	ggagacctgt	cattctaaga	300
tacccaaaac	agggccaggt	tggggacctg	tgcctttaat	cccatcactt	ggggattagg	360
tagaagcaag	aggctctaga	ccagtctaca	cactgaattt	caagccagcc	tacctataaa	420
tcagagaccc	tgcttcaaaa	ataaaattaa	acaaaaacga	agataaacca	agctacccaa	480
aacacaagag	ttaatccagt	cagacaggtc	tagcaaatgc	taggatgaaa	ggtgtgcacc	540
accacgagtg	ggctgcaagc	ctctctctct	ctctctctct	ctctctctct	ctcgtttgtt	600
ttgtttttcg	agacaagggt	tctctgtgta	gccctggctg	tcctggaact	cactctgtag	660
accaggctgg	cctcgagctt	cactcttaaa	agttcctctt	cctcctcctc	catcttttcc	720
tcctcttacc	ccctaggctc	cttttctctt	tcttgtcttt	cagataaagt	ctcaagtagt	780
ccagactggg	ctcaaaactaa	ctaactagcc	aagaatagcc	aacctcttaa	cttccgattc	840
tcctgcctct	gctgaatgct	gggggtgtgg	cgtggggccac	cacttctggg	ttgtgcaaca	900
cagaaggaac	tagggcttta	agcacgagaa	gcaagtctctg	tacagactta	cacaggccca	960
gcatctgttc	ttgcaatttt	ctgtaagtth	gacataatat	gagaataaaa	agctatctat	1020
ctcccttcca	gccttaccct	ctctgatgga	attcgaatgc	gtaatcaaag	cacccaacag	1080
cctggcctga	aatcacgtgg	ggcaagccca	cgtgaccgga	gcaccaatcc	aatatggcgg	1140
cgcccagggg	gcccgggctg	ttcctcatac	ccgcgctgct	cggttactc	ggggtggcgt	1200
ggtgcagctt	aagcttcggg	tgagtgcagg	ccgcccgggc	cagcctggct	ggggtccacc	1260
tttcttgagc	gctctcaggc	acagccctcc	gacctcacga	tcgccccgtc	cctgcagggt	1320
ttcccgcgac	gatgacctgc	tgctgcctta	cccactagcg	cgcagacgtc	cctcgcgaga	1380
ctgcgcccgg	gtgcgctcag	gtagcccaga	gcaggagagc	tggcctccgc	cacctctggc	1440
caccacgaa	ccccgggcgc	caagccacca	cgcgggcctg	cgcaccttcg	tgctgcactt	1500
cgagggggcg	gcggtggccg	gccacctgac	gcgggtcgcc	gatccccctac	gcactttctc	1560
ggtgctggag	cccggaggag	ccgggggctg	cggcggcaga	agcgccgcgg	ctactgtgga	1620
ggacacagcc	gtccggggcg	gttgccgcac	cgctcagaac	ggtggcttct	tcgcatgag	1680
cactggcgag	tgcttgggga	acgtggtgag	cgacggggcg	ctggtgagca	gctcaggggg	1740
actgcagaac	gcgcagttcg	gtatccgacg	cgatggaacc	atagtcaccg	ggtgaggagg	1800
cagggagccc	cggggctgta	gagggcaaa	ggtctctgat	gttctttcag	agccatgcct	1860
ccgagtcag	gtccctaacc	aaacttctct	tctttcttct	tccgagtaat	gacgctgaca	1920
ccttccttcc	tttaagttha	ttcatgtgcc	actgaataat	ctgtgatcag	gccgtgtgtg	1980
gggacttggg	gagggcagcc	tgagcctgaa	cacagtthgt	gccctagtga	actttgtgta	2040
gtattagaga	aacatttctg	gttcaacgaa	gccatggaac	caattggaaa	tagtgtagag	2100
tttatggagc	agtcccagac	agctagctgg	aggccttttg	ctgtcctgat	aaaaatccag	2160
gttagacaag	gagcttggtg	agggcagcct	ttggaagttt	ctgtgtttct	tgaaatttga	2220
cagcagccag	agttgacagc	aggcaggcag	gagtagaagg	tagcgccatc	tggtgttcca	2280
gttctcttcc	aagggtccgt	tttttgccaa	ggctgggaag	tgggctttcc	ccaactcttc	2340
tcagcccttg	gttgcaattt	ctgggcctgc	ccatgtatct	ggttcttcat	ccttcaacat	2400
cagccagtgt	caccactggt	gatcttaggt	tttcacagat	cctaaaactt	ctgccagtga	2460
ccagcgcttg	cagtttctct	tccttggtct	tgtccttcaa	cctctctaca	ttccagccat	2520
ctccctagct	cctctcttgg	actccctttc	agacttggtg	tcattgatcac	tgtctcagaa	2580
cccctattgc	tcctttacaa	tggtccactg	acctgctcac	ctcctacttt	ttttttttta	2640
atgtgtgtgc	atctgtgtgt	gcctgagggg	agaccagagt	ttgatttcaa	atgtcttcta	2700
ttctcttttc	ctccatctta	ttttctaaca	caaaatctga	atctagagat	cactggttca	2760
gttaacctgg	ctggccggta	aaccccaggg	ccctcctgct	tccctctgtc	cacccacccc	2820
cagcactaag	gctacagtgt	gtgctgttcc	agccagcttt	ctcatgggtg	ctgaggatct	2880
gaacgcaggt	tcacatgtgt	gggtgggaagg	cttttaccca	atgctctgtc	tttccagccc	2940
atcctccctt	gttaactgcc	aaacagctgc	ctatcctgtc	catgtgtagc	tcactgctac	3000
ttcttttatt	atgaggtcag	cacatgttac	taaagatggc	aagagaagaa	ggttctttca	3060
ttgtgtcata	gctatagctc	aggaggaatt	ttatttctct	tgtaggcaca	caggagagca	3120
tcttccagct	cacactccaa	ctgaactaac	tgaacacctg	cctatatatc	caaagaaggg	3180
gtgtcagtgc	caatcacagc	acacctccag	tgcaaatgaa	ggtttgtgtt	tgcaccaatc	3240
acagccttgc	ctcttttagc	atgcatcaca	acaaagtcct	cctagactat	caggggatat	3300
gctctcttgg	ccaaggtagg	aatagttgca	gtgtcatctg	gcacaaacca	tttcaaacgg	3360
cctggctgag	gttatgcctt	cgggaacctg	aagtctttgt	gtggttgtct	ccaagtgtct	3420
gtggagctcc	aggcggctgg	tgctgacaga	cgctttgtct	agttggctgt	ttgacttttg	3480
cttaagcagc	cagggcagta	gagtctaaca	gatgctaatt	tcaggatcag	gaagactgta	3540
gaaaaatgag	catcaagaag	cccctggtag	ccaaagctgc	tcttgccaat	gagtgaacct	3600
ctgccttccc	gcttccaggt	cctgtcttga	agaagaggtt	ctggatcccg	tgaatccggt	3660
cgtgcagctg	ctgagcggag	tcgtgtggct	catccgcaat	ggaaacatct	acatcaacga	3720

gagccaagcc	atcgagtgtg	acgagacaca	ggagacaggt	caggaagcac	aggtgttctg	3780
ttttatttgt	attaggtttt	gatttgttta	ttttgtgcat	gcagcgggtg	catgcatgct	3840
cctttccttt	cgccatgtga	gtcctgagta	ttgaactcag	actgttaagt	gtgatgggag	3900
gcactttacc	cactgagcca	ctttcccagc	cctcagcatc	agctttcttc	agacccagga	3960
acagtgtgag	tgggttattc	tttagtggtc	ccaaacattt	actgagcagc	tatttactgt	4020
ttagcactat	ggtgagagtc	ctagggattc	agtcttatgt	agaatataga	aggagaatcc	4080
ttggcaataa	gctggaaaat	tgtgacaagt	gccaagaaag	aaacaggaga	aaggggaccg	4140
gtggggacca	gaagcacagg	tatgaggaaa	gtgcctgcag	atttgctgta	tgggtggcctc	4200
cacatggcct	aggagtgtgt	cataaatgca	gagccatgag	tccaccctcc	ctatacctcc	4260
catccagaaa	ccactggtta	aatcctaaca	acttgggtgt	gcaggcactc	ccttgggtgac	4320
tctgatggac	actcaaggtc	aagggccact	tggggatggg	ctgatgagtt	ggcttgggtca	4380
gtaaagtatt	tgccttgaaa	gtgtgaggac	ctgagttgga	gccccagaaa	gaaacattaa	4440
aagccaagtg	ctgggatgca	cacttgcatt	cccagggatg	gagctggaag	gcagggatag	4500
gcagatccac	ggccacacgg	tgatattcta	agctaacaag	agacctgtct	cacacagaaa	4560
gtgggtggca	cctgaggacc	aacacccagg	gttatcctct	gacgtacctc	cagagtggaa	4620
aatactgggg	tgggtgaaaa	ggacactttg	gtcctgggaa	tctggctatt	cagggtatag	4680
tgtagaggga	gagggagact	caagaggctg	tctttgagtc	aaaggaacaa	gctatcagaa	4740
gaactcaggg	cagaggcctg	tggttcccag	gctcagggca	gccttcaagg	ccctaggcag	4800
agagtagctg	ctgggtgaac	aagtacagaa	gtgaggcctg	gggcctcagg	caaggcctgt	4860
gaaatccttc	caccaacata	gaagtttctg	gagactgaga	tcacatgaag	tgcttctggc	4920
tgtggcatgg	aagctcactg	gaggtggagc	tgggatgtgg	ctcagtgatc	cagtgccttg	4980
cacacgtgca	cgagggaagg	agccatcaaa	agagagaaag	tcgggagacc	tgaggggtcc	5040
cctggagagc	tgggtaacca	ccccgggccc	ttctccttta	ggttctttta	gcaaatttgt	5100
gaatgtgatg	tcagccagga	cagccgtggg	tcatgaccgt	gaggggcagc	ttatcctctt	5160
ccatgctgat	ggacagacgg	aacagcgtgg	tgagtcccag	gaaccttggg	gctgtttgca	5220
cttcagccac	cctacctttc	cagtcgggtc	tggggatttg	gtgggacaag	acagctttcc	5280
ggccattttg	gaagtttcat	ctggaggcaa	tagcatttac	ctactagtga	aagaagccag	5340
ttaagccaga	gaccacaggg	gctcaagctg	cataccccct	ctgcacagcc	ttaacctatg	5400
ggagatggca	gagttcctgc	gtcaacaaga	tgtcgtcaat	gccatcaacc	tggatggagg	5460
cggttctgct	acttttgtgc	tcaatgggac	cctggccagt	tacccttcag	atcactggta	5520
agaacccttg	agccaccttt	gtggctctct	cagactgtct	cactcagtca	atactgagac	5580
cctgtttgtg	gccaggccct	gggtatccaa	aagtgagcag	aagagccgag	atctcttccc	5640
tcagggtgct	gcacagccca	tccttggaag	cctgagacag	gtcaggaaag	gcctccctga	5700
ggacagtga	gtaagacctg	aggagatggc	tggccggggg	tgagagagcc	tttaccggaa	5760
gacaaactgt	acgcaatggg	gaaatccgct	aagtggccca	gggagaggct	ggagctatag	5820
ctcaggagga	aaagtacttg	cctcgcaagc	gaaggacctg	agtttaaact	ccaaaaccca	5880
tataaaaagc	cagatacgag	caagtggcac	atgcttgacg	tcccagcctt	gttgaggaag	5940
agtcagggtg	atcctgaccc	tctggccagc	cagcctagcc	tacttttttg	caagggtccag	6000
gccagcgaga	aagataaata	aaataaagtt	ttaaatgaca	tgtatctaag	gttgtcctga	6060
ctccatatgc	gcacgcacgc	atgcacgcac	gcacaactgg	cagaatggaa	agggaggcaa	6120
actggacagc	ctttataggc	tgccggcaggg	accagcacca	aggcctagac	ctcgtctcac	6180
agtgaatccc	ccacagccag	gacaacatgt	ggcgctgtcc	ccgccaagtg	tccactgtgg	6240
tgtgtgtgca	tgaaccgcgc	tgccagccac	ccgactgcag	tggccatggg	acctgtgtgg	6300
atggccactg	tgaatgcacc	agccacttct	ggcgggggcg	ggcctgcagc	gagctggact	6360
gtggccccct	caactgcagc	cagcatgggc	tgtgcacaga	gagtgagtgg	ggagcccaca	6420
ggaggggtgg	gctctggcgg	gaccccagct	cgcccatgct	agactcccgc	ctgtgtcctt	6480
accagcctc	tgtggtcctg	ctttggtagc	tggctgccac	tgtgatgctg	ggtggacagg	6540
atccaactgc	agtgaagggt	agagctgcct	gcaaacactc	ctggagaggg	tggcctggct	6600
gcacgcagct	ggtatgacgc	cttcgtccct	ccttctggct	tggaaacttac	cttcagagcc	6660
ttttctcatt	tcgcatgtgg	ataccgatg	ttctacctac	tgaaagagcc	cacaagtagg	6720
aagccagatt	ttcagtattg	tcactcaact	ctaaggacca	atagcaaaaa	aacaaagtgg	6780
ccacgcccct	gagggagatc	caccaaagtc	cttaactcct	ggaaagcagc	tcctgggtgat	6840
cctaggcatg	ggtaggggtg	tttcagcatc	agctcagtg	agttcccatt	cataatttct	6900
tcatcctttt	aaggtcataa	gttctagagc	ccaccttaaa	tctaggcagt	attcttgggtg	6960
tttatctgag	acaaagtctt	atacagccca	cgcagttctc	taacttagta	tgtaaccgag	7020
aatggcctca	agcaacctgc	ttcctccttt	caagcgctgg	gattataggg	atagcaccaa	7080
cttatagggg	gctagaagtc	aaaccagggg	ccctatgtat	atgcagcaag	cactctagaa	7140
actggaacac	agccctgttt	gcagcccggg	taccttgagg	ggttgggtcc	cagggatctg	7200
agggcatctc	cttcagcatg	gccatgtgca	cacccaggag	ccaggctgtc	tgtgacagga	7260
gaccatgcca	cccaagggtg	gacctccctg	ccaccatctc	ctctccacag	agtgtcctct	7320
gggctgggtat	gggccagggt	gccagaggcc	ctgccagtgt	gagcaccagt	gtttctgtga	7380

```

cccgagact ggcaactgca gcatctccca aggtatgcgg ccttaaaggt tcttgagctg 7440
ggagcccttg gggcaggtct ggggtaggtg gactctcccc agcccttctt tctggtgtct 7500
tgcagtgagg cagtgtctcc agccaactga ggctacgccg agggcaggag agctggcctc 7560
tttcaccagg taagtgtttt agcaggcact gagcccctat gtctcatccg tgaggcacta 7620
gccaggccag gaggtcacag gttaccctct actttgcaag ctcagggaca gtcacaggta 7680
aaactggcat ccaggaaaga ccctgagcta cccagtggaa ctcaaaggta gcaggctatg 7740
ggtgtcatgc ctctggctgc agagactcca cttagatgct ggagcagggc catagagaca 7800
ggaaggactc accttatttc tgaactcttc cgtgtgttca ggctttgtgt tgttgttgct 7860
tcctttctgc tgtttcctgg gtttccagct ccatccccac agggctcatg gaaagaattg 7920
tgaagcaggg ggtgtggctc aattggcaga ttgattgcct ggcatgcaga aagccctagg 7980
ttcaatcccc agcatttcat atcataacce aggcattgtg gcatcatgtg cctgtaagtc 8040
cagcacttgg gaggtagaag cagaaaagcc acgagttaa gaatgttagg gagtcttagg 8100
ccaacctggg atacctaaga caagagatag atgtaggagg atagattgac agacagacag 8160
acagacagac agacagacag atcttgagct ggaccttctg gcacaagcct gtcaccttag 8220
ctattccagg aagctgaagc aggaagatag caaattcaag gccagcttaa gccacagatt 8280
gagttcaaga tcaacctgag caactttatg aaatcctatt ataacataaa aagtaggggt 8340
gggaggttag gctgtagctc agtggttagg tgattgccta gcacgcacaa gaccaggtt 8400
caattcccag tactgcaaaa aatatattag gaaccccta aaagcagtaa cattcacatt 8460
agatgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtttt 8520
ttgggtattht atttcattht catttccaat gctatcccaa aagtccccca catcctcccc 8580
caccaccac cttgtttttt tttttttttt tttttttttt tttgacctga aactcacagg 8640
ttaggttaga caagctgact ggtgagctcc aacttccaac gtaccatcat gcctggcttt 8700
tgthttggtg tctctgtgta accctggatg tcctggagct ctctctgtag accagcctgg 8760
ccttaaactc acagaaacce acctgtttct gcctcccatg tgctgggatt aaaggcgtgt 8820
gccacctcac ccagccctgc tggacttaaa ttgggtcttc attttataag acaagcatga 8880
gctaattccc cagttcctaa aatgttttta acatccttaa acatcagaga ctgtctgtgg 8940
tattccctcc atgtgtcttc agtataccta cccccctccc tgcctactgg gttcaacatg 9000
cccagtttgg gttctggctg cctgccccca ctcaagactc tcttttccat ctcaggacca 9060
cctggctagc cctcaccctg acactaattt tcctgctgct gatcagcact ggggtcaacg 9120
tgctccttgt cctgggctcc agggccgaga ggaaccggca cctcgacggg gactatgtgt 9180
atcaccctact gcaggaggtg aacggggaag cgctgactgc agagaaggag cacatggagg 9240
aaactagcaa ccccttcaag gactgaagag ctgccccaac ggcatgctcc agataatctt 9300
gtccctgctc ctcaactcca caggggacat tgtgaggcca ctggcatgga tgctatgcac 9360
cccacccttt gctggccata ttctcctgt ccccatgctg tggctcatgc caacctagca 9420
ataaggagct ctggagagcc tgcacctgcc tcccgtctgc ctatatctgc tgcccagagg 9480
cctgtctcgc acaggggtct cgccactgcc aaagactccc aggaagtcaa agactcccag 9540
taatccacta gcaaatggaa ctctgtaacg ccatcataac aagagtggcc actctccgcg 9600
tgcacaggta tgaaatataa atccttacac acacacacac acacaccctc ggctcagcca 9660
cggcactcgc cttttataca gcgtcatcgc tggacagcca actagaactc tgcacctgt 9720
cacaggaagc acctcataag aaggaatggg gaggggaaggc agtcgccttg ttttcagacc 9780
ttagccgaat tc 9792

```

<210> 15
 <211> 908
 <212> PRT
 <213> Mus musculus

<400> 15
 Met Leu Phe Lys Leu Leu Gln Arg Gln Thr Tyr Thr Cys Leu Ser His
 1 5 10 15
 Arg Tyr Gly Leu Tyr Val Cys Phe Val Gly Val Val Val Thr Ile Val
 20 25 30
 Ser Ala Phe Gln Phe Gly Glu Val Val Leu Glu Trp Ser Arg Asp Gln
 35 40 45
 Tyr His Val Leu Phe Asp Ser Tyr Arg Asp Asn Ile Ala Gly Lys Ser
 50 55 60

Phe Gln Asn Arg Leu Cys Leu Pro Met Pro Ile Asp Val Val Tyr Thr
 65 70 75 80
 Trp Val Asn Gly Thr Asp Leu Glu Leu Leu Lys Glu Leu Gln Gln Val
 85 90 95
 Arg Glu His Met Glu Glu Glu Gln Arg Ala Met Arg Glu Thr Leu Gly
 100 105 110
 Lys Asn Thr Thr Glu Pro Thr Lys Lys Ser Glu Lys Gln Leu Glu Cys
 115 120 125
 Leu Leu Thr His Cys Ile Lys Val Pro Met Leu Val Leu Asp Pro Ala
 130 135 140
 Leu Pro Ala Thr Ile Thr Leu Lys Asp Leu Pro Thr Leu Tyr Pro Ser
 145 150 155 160
 Phe His Ala Ser Ser Asp Met Phe Asn Val Ala Lys Pro Lys Asn Pro
 165 170 175
 Ser Thr Asn Val Pro Val Val Val Phe Asp Thr Thr Lys Asp Val Glu
 180 185 190
 Asp Ala His Ala Gly Pro Phe Lys Gly Gly Gln Gln Thr Asp Val Trp
 195 200 205
 Arg Ala Tyr Leu Thr Thr Asp Lys Asp Ala Pro Gly Leu Val Leu Ile
 210 215 220
 Gln Gly Leu Ala Phe Leu Ser Gly Phe Pro Pro Thr Phe Lys Glu Thr
 225 230 235 240
 Ser Gln Leu Lys Thr Lys Leu Pro Arg Lys Ala Phe Pro Leu Lys Ile
 245 250 255
 Lys Leu Leu Arg Leu Tyr Ser Glu Ala Ser Val Ala Leu Leu Lys Leu
 260 265 270
 Asn Asn Pro Lys Gly Phe Gln Glu Leu Asn Lys Gln Thr Lys Lys Asn
 275 280 285
 Met Thr Ile Asp Gly Lys Glu Leu Thr Ile Ser Pro Ala Tyr Leu Leu
 290 295 300
 Trp Asp Leu Ser Ala Ile Ser Gln Ser Lys Gln Asp Glu Asp Ala Ser
 305 310 315 320
 Ala Ser Arg Phe Glu Asp Asn Glu Glu Leu Arg Tyr Ser Leu Arg Ser
 325 330 335
 Ile Glu Arg His Ala Pro Trp Val Arg Asn Ile Phe Ile Val Thr Asn
 340 345 350
 Gly Gln Ile Pro Ser Trp Leu Asn Leu Asp Asn Pro Arg Val Thr Ile
 355 360 365
 Val Thr His Gln Asp Ile Phe Gln Asn Leu Ser His Leu Pro Thr Phe
 370 375 380
 Ser Ser Pro Ala Ile Glu Ser His Ile His Arg Ile Glu Gly Leu Ser

385		390		395		400
Gln Lys Phe Ile Tyr Leu Asn Asp Asp Val Met Phe Gly Lys Asp Val						
		405		410		415
Trp Pro Asp Asp Phe Tyr Ser His Ser Lys Gly Gln Lys Val Tyr Leu						
		420		425		430
Thr Trp Pro Val Pro Asn Cys Ala Glu Gly Cys Pro Gly Ser Trp Ile						
		435		440		445
Lys Asp Gly Tyr Cys Asp Lys Ala Cys Asn Thr Ser Pro Cys Asp Trp						
		450		455		460
Asp Gly Gly Asn Cys Ser Gly Asn Thr Ala Gly Asn Arg Phe Val Ala						
		465		470		475
Arg Gly Gly Gly Thr Gly Asn Ile Gly Ala Gly Gln His Trp Gln Phe						
		485		490		495
Gly Gly Gly Ile Asn Thr Ile Ser Tyr Cys Asn Gln Gly Cys Ala Asn						
		500		505		510
Ser Trp Leu Ala Asp Lys Phe Cys Asp Gln Ala Cys Asn Val Leu Ser						
		515		520		525
Cys Gly Phe Asp Ala Gly Asp Cys Gly Gln Asp His Phe His Glu Leu						
		530		535		540
Tyr Lys Val Thr Leu Leu Pro Asn Gln Thr His Tyr Val Val Pro Lys						
		545		550		555
Gly Glu Tyr Leu Ser Tyr Phe Ser Phe Ala Asn Ile Ala Arg Lys Arg						
		565		570		575
Ile Glu Gly Thr Tyr Ser Asp Asn Pro Ile Ile Arg His Ala Ser Ile						
		580		585		590
Ala Asn Lys Trp Lys Thr Leu His Leu Ile Met Pro Gly Gly Met Asn						
		595		600		605
Ala Thr Thr Ile Tyr Phe Asn Leu Thr Leu Gln Asn Ala Asn Asp Glu						
		610		615		620
Glu Phe Lys Ile Gln Ile Ala Val Glu Val Asp Thr Arg Glu Ala Pro						
		625		630		635
Lys Leu Asn Ser Thr Thr Gln Lys Ala Tyr Glu Ser Leu Val Ser Pro						
		645		650		655
Val Thr Pro Leu Pro Gln Ala Asp Val Pro Phe Glu Asp Val Pro Lys						
		660		665		670
Glu Lys Arg Phe Pro Lys Ile Arg Arg His Asp Val Asn Ala Thr Gly						
		675		680		685
Arg Phe Gln Glu Glu Val Lys Ile Pro Arg Val Asn Ile Ser Leu Leu						
		690		695		700
Pro Lys Glu Ala Gln Val Arg Leu Ser Asn Leu Asp Leu Gln Leu Glu						
		705		710		715
						720

Arg Gly Asp Ile Thr Leu Lys Gly Tyr Asn Leu Ser Lys Ser Ala Leu
 725 730 735
 Leu Arg Ser Phe Leu Gly Asn Ser Leu Asp Thr Lys Ile Lys Pro Gln
 740 745 750
 Ala Arg Thr Asp Glu Thr Lys Gly Asn Leu Glu Val Pro Gln Glu Asn
 755 760 765
 Pro Ser His Arg Arg Pro His Gly Phe Ala Gly Glu His Arg Ser Glu
 770 775 780
 Arg Trp Thr Ala Pro Ala Glu Thr Val Thr Val Lys Gly Arg Asp His
 785 790 795 800
 Ala Leu Asn Pro Pro Pro Val Leu Glu Thr Asn Ala Arg Leu Ala Gln
 805 810 815
 Pro Thr Leu Gly Val Thr Val Ser Lys Glu Asn Leu Ser Pro Leu Ile
 820 825 830
 Val Pro Pro Glu Ser His Leu Pro Lys Glu Glu Glu Ser Asp Arg Ala
 835 840 845
 Glu Gly Asn Ala Val Pro Val Lys Glu Leu Val Pro Gly Arg Arg Leu
 850 855 860
 Gln Gln Asn Tyr Pro Gly Phe Leu Pro Trp Glu Lys Lys Lys Tyr Phe
 865 870 875 880
 Gln Asp Leu Leu Asp Glu Glu Glu Ser Leu Lys Thr Gln Leu Ala Tyr
 885 890 895
 Phe Thr Asp Arg Lys His Thr Gly Arg Gln Leu Lys
 900 905

<210> 16
 <211> 5229
 <212> DNA
 <213> Mus musculus

<400> 16
 ggcggtgaag gggatgatgct gttcaagctc ctgcagagac agacctatac ctgcctatcc 60
 cacaggtatg ggctctacgt ctgcttcgtg ggcgtcgttg tcaccatcgt ctcggttttc 120
 cagttcggag aggtggttct ggaatggagc cgagatcagt accatgtttt gtttgattcc 180
 tacagagaca acattgctgg gaaatccttt cagaatcggc tctgtctgcc catgccaatc 240
 gacgtgggtt acacctgggt gaatggcact gaccttgaac tgctaaagga gctacagcag 300
 gtccgagagc acatggagga agagcagaga gccatgcggg aaaccctcgg gaagaacaca 360
 accgaaccga caaagaagag tgagaagcag ctggaatgtc tgctgacgca ctgcattaag 420
 gtgcccattg ttgttctgga cccggccctg ccagccacca tcaccctgaa ggatctgcca 480
 accctttacc catctttcca cgcgtccagc gacatgttca atgttgcgaa accaaaaaat 540
 ccgtctacaa atgtccccgt tgtcgttttt gacactacta aggatgttga agacgcccac 600
 gctggaccgt ttaagggagg ccagcaaaca gatgtttgga gagcctactt gacaacagac 660
 aaagacgccc ctggcttagt gctgatacaa ggcttggcgt tcctgagtgg attcccaccg 720
 accttcaagg agacgagtca actgaagaca aagctgccaa gaaaagcttt ccctctaaaa 780
 ataaagctgt tgcggctgta ctggagggcc agtgtcgtc ttctgaaatt gaataatccc 840
 aagggtttcc aagagctgaa caagcagacc aagaagaaca tgaccatcga tgggaaggaa 900
 ctgaccatca gccctgcgta tctgctgtgg gacctgagt ccatcagcca gtccaagcag 960
 gatgaggacg cgtctgccag ccgctttgag gataatgaag agctgaggta ctcgctgcga 1020

tctatcgaga	gacacgcgcc	atgggtacgg	aatatTTTTca	ttgtcaccaa	cgggcagatt	1080
ccatcctggc	tgaaccttga	caaccctcga	gtgaccatag	tgaccaccca	ggacattttc	1140
caaaatctga	gccacttgcc	tacttttcagt	tcccctgcta	ttgaaagtca	cattcaccgc	1200
atcgaagggc	tgtcccagaa	gtttatTTtat	ctaaatgacg	atgtcatgtt	cggtaaggac	1260
gtctggccgg	acgatttttta	cagccactcc	aaaggTcaaa	aggTttattt	gacatggcct	1320
gtgccaaact	gtgcagaggg	ctgcccgggc	tcttgataa	aggacggcta	ttgtgataag	1380
gcctgtaata	cctcacccctg	tgactgggat	ggcggaaact	gctctggtaa	tactgcaggg	1440
aaccggtttg	ttgcaagagg	tgggggtacc	gggaatattg	gagctggaca	gcactggcag	1500
tttggtggag	gaataaacac	catctcttac	tgtaaccaag	gatgtgcaaa	ctcctggctg	1560
gctgacaagt	tctgtgacca	agcctgtaac	gtcttatcct	gcgggtttga	tgctggtgac	1620
tgtggacaag	atcatttttca	tgaattgtat	aaagtaaacac	ttctcccaa	ccagactcac	1680
tatgttgtcc	ccaaagggtga	atacctgtct	tatttcagct	ttgcaaacad	agccagaaaa	1740
agaattgaag	ggacctacag	cgacaacccc	atcatccgcc	acgcgtccat	tgcaaacaag	1800
tggaaaaccc	tacacctgat	aatgcccggg	gggatgaacg	ccaccacgat	ctattttaac	1860
ctcactcttc	aaaacgccaa	cgacgaagag	ttcaagatcc	agatagcagt	agagggtggac	1920
acgagggagg	cgcccaaact	gaattctaca	acccagaagg	cctatgaaag	tttggttagc	1980
ccagtgcacac	ctcttcctca	ggctgacgtc	ccttttgaag	atgtcccaa	agagaaacgc	2040
ttccccaaga	tcaggagaca	tgatgtaaat	gcaacaggga	gattccaaga	ggagggtgaaa	2100
atcccccggg	taaatatTTc	actccttccc	aaagaggccc	aggTgaggct	gagcaacttg	2160
gatttgcaac	tagaacgtgg	agacatcact	ctgaaaggat	ataacttgtc	caagtcagcc	2220
ctgctaaggt	ctttcctggg	gaattcacta	gatactaaaa	taaaacctca	agctaggacc	2280
gatgaaacaa	aaggcaacct	ggaggTccca	caggaaaacc	cttctcacag	acgtccacat	2340
ggctttgctg	gtgaacacag	atcagagaga	tggactgccc	cagcagagac	agtgaccgtg	2400
aaaggccgtg	accacgcttt	gaatccaccc	ccggtgttgg	agaccaatgc	aagattggcc	2460
cagcctacac	taggcgtgac	tgtgtccaaa	gagaaccttt	caccgctgat	cgttccccc	2520
gaaagccact	tgccaaaaga	agaggagagt	gacagggcag	aaggcaatgc	tgtacctgta	2580
aaggagttag	tgccctggcag	acggttgca	cagaattatc	caggcttttt	gccctgggag	2640
aaaaaaaaagt	atttccaaga	ccttcttgat	gaggaagagt	cattgaagac	ccagttggcg	2700
tactttacag	accgcaaaca	taccgggagg	caactaaaag	atacatttgc	agactccctc	2760
cgatacgtca	ataaaattct	caacagcaag	tttgatttca	catccaggaa	agtccctgca	2820
cacatgccgc	acatgattga	caggatcggt	atgcaagaac	tccaagatat	gttccctgaa	2880
gaatttgaca	agacttcatt	tcacaagggt	cgtcactctg	aggacatgca	gtttgccttc	2940
tcctactttt	attacctcat	gagtgcagtt	cagccccctca	atatttccca	agtctttcat	3000
gaagtagaca	cagaccaatc	tggtgtcttg	tctgataggg	aaatccgaac	wctggccacg	3060
agaattcacg	acctaccttt	aagcttgca	gatttgacag	gtttggaaca	catgttaata	3120
aattgctcaa	aaatgctccc	cgctaataatc	actcaactca	acaacatccc	accgactcag	3180
gaagcatact	acgaccccaa	cctgcctccg	gtcactaaga	gtcttgtcac	caactgtaag	3240
ccagtaactg	acaagatcca	caaagcctat	aaagacaaga	acaaatacag	gtttgaaatc	3300
atgggagagg	aagaaatcgc	tttcaagatg	atacgaacca	atgtttctca	tgtggttggt	3360
cagttggatg	acatcagaaa	aaaccccagg	aagttcgttt	gtctgaatga	caacattgac	3420
cacaaccata	aagatgcccg	gacagtgaag	gctgtcctca	gggacttcta	tgagtccatg	3480
tttcccatat	cttcccagtt	tgagctgcca	agagagtatc	ggaaccgctt	tctgcacatg	3540
catgagctcc	aagaatggcg	ggcatatcga	gacaagctga	agttttggac	ccactgcgta	3600
ctagcaacgt	tgattatatt	tactatatct	tcattttttg	ctgaacagat	aattgctctg	3660
aagcgaaaga	tatttcccag	gaggaggata	cacaaagaag	ctagtccaga	ccgaatcagg	3720
gtgtagaaga	tcttcatttg	aaagtcacct	accttagcat	ctgtgaacat	ctccctcctc	3780
gacaccacag	cggagtccct	gtgatgtggc	acagaggcag	cctcgtgggg	agaagggaca	3840
tcgtgcagac	cgggttcttc	tgcaatggga	agagagccca	ctgacctgga	attattcagc	3900
acactaagaa	cctgtgtcaa	tagcttgtac	agcttgtact	tttaaaggat	ttgccgaagg	3960
acctgtcggc	ttgttgacaa	accctccctg	acaagctgct	ggtttcttcc	cccagttact	4020
gcagactgag	aaaccagtcc	atcttgaaag	caagtgcgga	ggggccccag	tctttgcatt	4080
ccaaagcttt	ccagcataat	ttctggcttg	tctcctcctt	tgatccattt	cccatttttt	4140
tttaaaaaaac	aataagtggc	tactaagtta	gtcatttctca	cttctcaaaa	taacaaatca	4200
ggatgtcaaa	acatttgtat	agatcttatt	taaataatat	agaacgatta	cttcttttagc	4260
ctatctaaat	tattgatttt	tattaacagt	caagtgggtct	tgaaccgcta	acaactactg	4320
aagagctcga	gattgacgtt	gaaagtgcct	tgagcttggt	taactcattc	cccaagaata	4380
ctgtgacctc	gtgtgcgggc	ctgattgcga	agggttagtg	tcacgtagca	gtgctgctca	4440
ccggatgtaa	ttatgtcgtg	gaaatgtaca	tacagacaaa	agtgcctcac	ttcagaaatg	4500
agtagtgctg	atggcaccag	cgagtgatgg	tgtccatttg	gaaacccatg	ataccttcca	4560
atgcccaccc	tgcttacttt	atacagagca	gggtttaacc	aacttctgtc	aaagaacagt	4620
aaagaacttg	agatacatcc	atctttgtca	aatagttttc	cttgctaaca	tttattattg	4680

```

ttggtgtttt gggagggttta ttttatttta ttgctttgtt atttttcaag acgggggattc 4740
tctgtgtagc tctggctgtt tggtaattca ctctaaagac caggctggcc ttgaacttag 4800
agattcacct gcttctgctt cctgaatggt aggacatgtg cccacattgc ctaccacccc 4860
cccttttggg ggggggtgagc aactcaataa aaagatgaaa acctgcttta gtttgcagct 4920
atacaaaaagc agcaggcctc agccagactt gacccccggg gccattgttg gcccacggga 4980
gaatcatttt tgacgtgggt aagcaaacc ctgatattgt catgctgtgt tatgtcatta 5040
tgtgggtggt ttgaattttg gaagatattt tcagtcatga tttcagtagt attcctccaa 5100
aatggcacac atttttgtaa taagaacttg aaatgtaaat attgtgtttg tgctgtaa at 5160
tttgtgtatt tcaaaaactg aagtttcata aaaaaacaca cttattggaa aaaaaaaaaa 5220
aaaaaaaaa 5229

```

```

<210> 17
<211> 1105
<212> DNA
<213> Drosophila melanogaster

```

```

<220>
<221> misc_feature
<222> (903)
<223> N IS A, G, C, OR T

```

```

<220>
<221> misc_feature
<222> (935)
<223> N IS A, G, C, OR T

```

```

<220>
<221> misc_feature
<222> (1023)
<223> N IS A, T, G, OR C

```

```

<220>
<221> misc_feature
<222> (1035)
<223> N IS A, G, T, OR C

```

```

<220>
<221> misc_feature
<222> (1071)
<223> N IS A, G, T, OR C

```

```

<220>
<221> misc_feature
<222> (1100)
<223> N IS A, T, G, OR C

```

```

<400> 17
ctgcaggaat tcggcacgag gcggttcgat gacaagaatg agctgcggtg ctctctgagg 60
tccctggaaa aacacgccgc atggatcagg catgtgtaca tagtaaccaa tggccagatt 120
ccaagtggc tggatctcag ctacgaaagg gtcacgggtg tgccccacga agtcctggct 180
cccgatccc accagctgcc caccttctcc agctcggcca tcgagacatt tctgcaccgc 240
atacaaaagc tgtccaagag gttectctac ctcaacgacg acatattcct gggagctccg 300
ctgtatccgg aggacttgta cactgaagcg gagggagttc gcgtgtacca ggcattggatg 360
gtgcccggct ggccttgga ttgcccctgg acgtacatag gtgatggagc ttgcatcg 420
cactgcaaca ttgatgcgtg ccaatttgat ggaggcgact gcagtgaaac tgggccagcg 480
agcgatgcc acgtcattcc accaagcaaa gaagtgtcgc aggtgcagcc tgccgctgtt 540
ccacaatcaa gagtccaccg atttcctcag atgggtctcc aaaagctgtt caggcgcagc 600
tctgccaatt ttaaggatgt tatgcggcac cgcaatgtgt ccacactcaa ggaactacgt 660
cgcattgtgg agcgttttaa caaggccaaa ctcatgtcgc tgaaccccgga actggagacc 720
tccagctccg agccacagac aactcagcgc cacgggctgc gcaaggagga ttttaagtct 780

```

```

tccaccgata tttactctca ctcgctgatt gccaccaata tggtgctgaa tagagcctat 840
ggctttaagg cagcccatgt cctggcgcac gtgggcttcc taattgacaa ggatattgtg 900
gangccatgc aacgacgttt taccagcgaa ttctngacac tggccattaa cgctttccga 960
gccccaacag atttgacgta cgcattcgct tactacttct ttctaattgag cgaaatccaa 1020
gtnatgagtg tagangaaat cttcgatgaa gtcgacaccg gacggtttgg ncacctggtc 1080
ggatccagaa gtgcgaaccn tttta 1105

```

<210> 18
 <211> 2005
 <212> DNA
 <213> Mus musculus

```

<400> 18
gtttcccgcg acgatgacct gctgctgcct taccactag cgcgcagacg tccctcgcga 60
gactgcgccc ggggtgcgctc aggtagccca gagcaggaga gctggcctcc gccacctctg 120
gccaccacag aaccccgggc gccaaagccac cagcggcccg tgcgcacctt cgtgtcgcac 180
ttcgagggggc gcgcgggtggc cggccacctg acgcgggtcg ccgatcccct acgcactttc 240
tcggtgctgg agcccggagg agccgggggc tgcggcggca gaagcgcgcg ggctactgtg 300
gaggacacag ccgtccgggc cggttgcgcg atcgctcaga acggtggctt cttccgcatg 360
agcactggcg agtgcttggg gaacgtgggtg agcgacgggc ggctgggtgag cagctcaggg 420
ggactgcaga acgcgcagtt cggtatccga cgcgatggaa ccatagtcac cgggtcctgt 480
cttgaagaag aggttctgga tcccgtgaat ccgttcgtgc agctgctgag cggagtcgtg 540
tggctcatcc gcaatggaaa catctacatc aacgagagcc aagccatcga gtgtgacgag 600
acacaggaga caggttcttt tagcaaattt gtgaatgtga tgtcagccag gacagccgtg 660
ggctcatgacc gtgaggggca gcttatcctc ttccatgctg atggacagac ggaacagcgt 720
ggccttaacc tatgggagat ggcagagttc ctgctcaac aagatgtcgt caatgccatc 780
aacctggatg gaggcgggtc tgctactttt gtgctcaatg ggaccctggc cagttaccct 840
tcagatcact gccaggacaa catgtggcgc tgtccccgcc aagtgtccac tgtggtgtgt 900
gtgcatgaac cgcgctgcc a gccaccgcac tgcagtggcc atgggacctg tgtggatggc 960
cactgtgaat gcaccagcca cttctggcgg ggcgaggcct gcagcgagct ggactgtggc 1020
ccctccaact gcagccagca tgggctgtgc acagctggct gccactgtga tgctgggtgg 1080
acaggatcca actgcagtga agagtgtcct ctgggctggg atggggccagg ttgccagagg 1140
ccctgccagt gtgagcacca gtgtttctgt gaccgcaga ctggcaactg cagcatctcc 1200
caagtgaggc agtgtctcca gccaaactgag gctacgccga gggcaggaga gctggcctct 1260
ttcaccagga ccacctggct agccctcacc ctgacactaa ttttcctgct gctgatcagc 1320
actggggtea acgtgtcctt gttcctgggc tccagggccg agaggaaccg gcacctcgac 1380
ggggactatg tgtatcaccc actgcaggag gtgaacgggg aagcgctgac tgcagagaag 1440
gagcacatgg aggaaactag caacccttc aaggactgaa gagctgcccc aacggcatgc 1500
tccagataat cttgtccctg ctctcactt ccacagggga cattgtgagg ccactggcat 1560
ggatgctatg caccaccacc tttgctggcc atattcctcc tgtccccatg ctgtggctca 1620
tgccaacct a gcaataagga gctctggaga gcctgcacct gcctcccgct cgcctataatc 1680
tgctgcccag aggctgtct cgcacagggg tctcgccact gccaaagact cccaggaagt 1740
caaagactcc cagtaatcca ctagcaaagt gaactctgta acgccatcat aacaagagt 1800
gccactctcc gcgtgcacag gtatgaaata taaatcctta cacacacaca cacacacacc 1860
ctcggtcag ccacggcact cgccttttat acagcgtcat cgctggacag ccaactagaa 1920
ctctgcaccc tgtcacagga agcacctcat aagaaggaaat ggggagggaa ggcagtcgcc 1980
ttgttttcag accttagccg aattc 2005

```

<210> 19
 <211> 492
 <212> PRT
 <213> Mus musculus

```

<400> 19
Val Ser Arg Asp Asp Asp Leu Leu Leu Pro Tyr Pro Leu Ala Arg Arg
  1                      5                      10          15
Arg Pro Ser Arg Asp Cys Ala Arg Val Arg Ser Gly Ser Pro Glu Gln
                20                25                30

```

Glu Ser Trp Pro Pro Pro Pro Leu Ala Thr His Glu Pro Arg Ala Pro
 35 40 45
 Ser His His Ala Ala Val Arg Thr Phe Val Ser His Phe Glu Gly Arg
 50 55 60
 Ala Val Ala Gly His Leu Thr Arg Val Ala Asp Pro Leu Arg Thr Phe
 65 70 75 80
 Ser Val Leu Glu Pro Gly Gly Ala Gly Gly Cys Gly Gly Arg Ser Ala
 85 90 95
 Ala Ala Thr Val Glu Asp Thr Ala Val Arg Ala Gly Cys Arg Ile Ala
 100 105 110
 Gln Asn Gly Gly Phe Phe Arg Met Ser Thr Gly Glu Cys Leu Gly Asn
 115 120 125
 Val Val Ser Asp Gly Arg Leu Val Ser Ser Ser Gly Gly Leu Gln Asn
 130 135 140
 Ala Gln Phe Gly Ile Arg Arg Asp Gly Thr Ile Val Thr Gly Ser Cys
 145 150 155 160
 Leu Glu Glu Glu Val Leu Asp Pro Val Asn Pro Phe Val Gln Leu Leu
 165 170 175
 Ser Gly Val Val Trp Leu Ile Arg Asn Gly Asn Ile Tyr Ile Asn Glu
 180 185 190
 Ser Gln Ala Ile Glu Cys Asp Glu Thr Gln Glu Thr Gly Ser Phe Ser
 195 200 205
 Lys Phe Val Asn Val Met Ser Ala Arg Thr Ala Val Gly His Asp Arg
 210 215 220
 Glu Gly Gln Leu Ile Leu Phe His Ala Asp Gly Gln Thr Glu Gln Arg
 225 230 235 240
 Gly Leu Asn Leu Trp Glu Met Ala Glu Phe Leu Arg Gln Gln Asp Val
 245 250 255
 Val Asn Ala Ile Asn Leu Asp Gly Gly Gly Ser Ala Thr Phe Val Leu
 260 265 270
 Asn Gly Thr Leu Ala Ser Tyr Pro Ser Asp His Cys Gln Asp Asn Met
 275 280 285
 Trp Arg Cys Pro Arg Gln Val Ser Thr Val Val Cys Val His Glu Pro
 290 295 300
 Arg Cys Gln Pro Pro Asp Cys Ser Gly His Gly Thr Cys Val Asp Gly
 305 310 315 320
 His Cys Glu Cys Thr Ser His Phe Trp Arg Gly Glu Ala Cys Ser Glu
 325 330 335
 Leu Asp Cys Gly Pro Ser Asn Cys Ser Gln His Gly Leu Cys Thr Ala
 340 345 350

Gly Cys His Cys Asp Ala Gly Trp Thr Gly Ser Asn Cys Ser Glu Glu
 355 360 365

Cys Pro Leu Gly Trp Tyr Gly Pro Gly Cys Gln Arg Pro Cys Gln Cys
 370 375 380

Glu His Gln Cys Phe Cys Asp Pro Gln Thr Gly Asn Cys Ser Ile Ser
 385 390 395 400

Gln Val Arg Gln Cys Leu Gln Pro Thr Glu Ala Thr Pro Arg Ala Gly
 405 410 415

Glu Leu Ala Ser Phe Thr Arg Thr Thr Trp Leu Ala Leu Thr Leu Thr
 420 425 430

Leu Ile Phe Leu Leu Leu Ile Ser Thr Gly Val Asn Val Ser Leu Phe
 435 440 445

Leu Gly Ser Arg Ala Glu Arg Asn Arg His Leu Asp Gly Asp Tyr Val
 450 455 460

Tyr His Pro Leu Gln Glu Val Asn Gly Glu Ala Leu Thr Ala Glu Lys
 465 470 475 480

Glu His Met Glu Glu Thr Ser Asn Pro Phe Lys Asp
 485 490

<210> 20

<211> 3783

<212> DNA

<213> Homo sapiens

<400> 20

```

gccaccatgg ggttcaagct cttgcagaga caaacctata cctgcctgtc ccacaggtat 60
gggctctacg tgtgcttctt gggcgctcgtt gtcaccatcg tctccgcctt ccagttcgga 120
gaggtggttc tggaatggag ccgagatcaa taccatgttt tgtttgattc ctatagagac 180
aatattgctg gaaagtcctt tcagaatcgg ctttgtctgc ccatgccgat tgacgttggt 240
tacacctggg tgaatggcac agatcctgaa ctactgaagg aactacagca ggtcagagaa 300
cagatggagg aggagcagaa agcaatgaga gaaatccttg ggaaaaacac aacggaacct 360
actaagaaga gtgagaagca gttagagtgt ttgctaacac actgcattaa ggtgccaatg 420
cttgtcctgg acccagccct gccagccaac atcacctga aggacctgcc atctctttat 480
ccttcttttc attctgccag tgacattttc aatgttgcaa aacaaaaaaa cccttctacc 540
aatgtctcag ttgttggttt tgacagtact aaggatgttg aagatgcca ctctggactg 600
cttaaaggaa atagcagaca gacagtatgg aggggctact tgacaacaga taaagaagtc 660
cctggattag tgctaattgca agatttggtt ttcctgagtg gatttccacc aacattcaag 720
gaaacaaatc aactaaaaac aaaattgcca gaaaatcttt cctctaaagt caaactgttg 780
cagttgtatt cagaggccag tgtagcgctt ctaaaactga ataaccctaa ggattttcaa 840
gaattgaata agcaaactaa gaagaacatg accattgatg gaaaagaact gaccataagt 900
cctgcatatt tattatggga tctgagcgcc atcagccagt ctaagcagga tgaagacatc 960
tctgccagtc gttttgaaga taacgaagaa ctgaggtact cattgcatc tatcgagagg 1020
catgcaccat gggttcggaa tattttcatt gtcaccaacg ggcagattcc atcctggctg 1080
aaccttgaca atcctcgagt gacaatagta acacaccagg atgttttttcg aaatttgagc 1140
cacttgccct ccttttagttc acctgctatt gaaagtcacg ttcatcgcat cgaagggctg 1200
tcccagaagt ttatttacct aaatgatgat gtcatgtttg ggaaggatgt ctggccagat 1260
gattttttaca gtcactccaa aggccagaag gtttatttga catggcctgt gccaaactgt 1320
gccgagggct gccaggttc ctggattaag gatggctatt gtgacaaggc ttgtaataat 1380
tcagcctgcg attgggatgg tggggattgc tctggaaaca gtggaggagg tcgctatatt 1440
gcaggagggt gaggtactgg gagtattgga gttggacagc cctggcagtt tgggtggagg 1500
ataaacagtg tctcttactg taatcagga tgtgcgaatt cctggctcgc tgataagttc 1560
tgtgaccaag catgcaatgt cttgtcctgt gggtttgatg ctggcgactg tgggcaagat 1620

```

```

cattttcatg aattgtataa agtgatcctt ctcccaaacc agactcacta tattattcca 1680
aaagggtgaat gcctgcctta tttcagcttt gcagaagtag ccaaaagagg agttgaagggt 1740
gcctatagtg acaatccaat aattcgacat gcttctattg ccaacaagtg gaaaaccatc 1800
cacctcataa tgcacagtgg aatgaatgcc accacaatac attttaatct cacgtttcaa 1860
aatacaaacg atgaagagtt caaaatgcag ataacagtgg aggtggacac aagggaggga 1920
ccaaaactga attctacggc ccagaagggt tacgaaaatt tagttagtcc cataacactt 1980
cttccagagg cggaaatcct ttttgaggat attcccaaag aaaaacgctt cccgaagttt 2040
aagagacatg atgttaactc aacaaggaga gccaggaag aggtgaaaat tcccctggta 2100
aatatttcac tccttccaaa agacgcccag ttgagtctca ataccttgga tttgcaactg 2160
gaacatggag acatcacttt gaaaggatac aatttgtcca agtcagcctt gctgagatca 2220
tttctgatga actcacagca tgctaaaata aaaaatcaag ctataataac agatgaaaca 2280
aatgacagtt tgggtggctcc acaggaaaaa caggttcata aaagcatcct gccaaacagc 2340
ttaggagtggt ctgaaagatt gcagagggtg acttttcctg cagtgagtggt aaaagtgaat 2400
ggtcgatgacc agggtcagaa tccacccctg gacttggaga ccacagcaag atttagagtg 2460
gaaactcaca cccaaaaaac cataggcgga aatgtgacaa aagaaaagcc cccatctctg 2520
attgttccac tggaaagcca gatgacaaaa gaaaagaaaa tcacagggaa agaaaaagag 2580
aacagtagaa tggaggaaaa tgctgaaaat cacataggcg ttactgaagt gttacttgga 2640
agaaagctgc agcattacac agatagttac ttgggctttt tgccatggga gaaaaaaaag 2700
tatttcctag atcttctcga cgaagaagag tcattgaaga cacaattggc atacttcact 2760
gatagcaaga atactgggag gcaactaaaa gatacatttg cagattccct cagatatgta 2820
aataaaattc taaatagcaa gtttggattc acatcgcgga aagtccctgc tcacatgcct 2880
cacatgattg accggattgt tatgcaagaa ctgcaagata tgttccctga agaatttgac 2940
aagacgtcat ttcacaaagt gcgccattct gaggatatgc agtttgcctt ctcttatttt 3000
tattatctca tgagtgcagt gcagccactg aatatatctc aagtctttga tgaagttgat 3060
acagatcaat ctggtgtctt gtctgacaga gaaatccgaa cactggctac cagaattcac 3120
gaactgccgt taagtttgca ggatttgaca ggtctggaac acatgctaataaattgctca 3180
aaaatgcttc ctgctgatat cacgcagcta aataatattc caccaactca ggaatcctac 3240
tatgatccca acctgccacc ggtcactaaa agtctagtaa caaactgtaa accagtaact 3300
gacaaaatcc acaaagcata taaggacaaa aacaaatata ggtttgaaat catgggagaa 3360
gaagaaatcg cttttaaaat gattcgtacc aacgtttctc atgtgggttg ccagttggat 3420
gacataagaa aaaaccctag gaagtttgtt tgcctgaatg acaacattga ccacaatcat 3480
aaagatgctc agacagtga ggctgttctc agggacttct atgaatccat gttccccata 3540
ccttcccaat ttgaactgcc aagagagtat cgaaaccgtt tccttcatat gcatgagctg 3600
caggaatgga gggcttatcg agacaaattg aagttttgga cccattgtgt actagcaaca 3660
ttgattatgt ttactatatt ctcatttttt gctgagcagt taattgcact taagcggaag 3720
atatttccca gaaggaggat acacaaagaa gctagtccca atcgaatcag agtatagaag 3780
atc 3783

```

<210> 21
 <211> 3621
 <212> DNA
 <213> Homo sapiens

```

<400> 21
ctagccgcca ccatggagac agacacactc ctgctatggg tactgctgct ctgggttcca 60
ggttccactg gtgacgaaga tcaggtagat ccgcggttaa tcgacggtaa gcttagccga 120
gatcaatacc atgttttggt tgattcctat agagacaata ttgctggaaa gtcctttcag 180
aatcggcttt gtctgcccac gccgattgac gttgtttaca cctgggtgaa tggcacagat 240
cttgaactac tgaaggaaact acagcaggtc agagaacaga tggaggagga gcagaaagca 300
atgagagaaa tccttgggaa aaacacaacg gaacctacta agaagagtga gaagcagtta 360
gagtgtttgc taacacactg cattaagggt ccaatgcttg tcctggaccc agccctgcc 420
gccaacatca ccctgaagga cctgccatct ctttatcctt cttttcattc tgccagtga 480
attttcaatg ttgcaaaacc aaaaaaccct tctaccaatg tctcagttgt tgtttttgac 540
agtactaagg atgttgaaga tgcccactct ggactgctta aaggaaatag cagacagaca 600
gatatggagg gctacttgac aacagataaa gaagtcctg gattagtgt aatgcaagat 660
ttggctttcc tgagtggatt tccaccaaca ttcaaggaaa caaatcaact aaaaacaaaa 720
ttgccagaaa atctttcctc taaagtcaaa ctgttgacgt tgtattcaga ggccagtga 780
gcgcttctaa aactgaataa cccaaggat tttcaagaat tgaataagca aactaagaag 840
aacatgacca ttgatggaaa agaactgacc ataagtcctg catatttatt atgggatctg 900
agcgccatca gccagtctaa gcaggatgaa gacatctctg ccagtcgttt tgaagataac 960

```

```

gaagaactga ggtactcatt gcgatctatc gagagggcatg caccatgggt tcggaatatt 1020
ttcattgtca ccaacgggca gattccatcc tggctgaacc ttgacaatcc tcgagtga 1080
atagtaacac accaggatgt ttttcgaaat ttgagccact tgcctacctt tagttcacct 1140
gctattgaaa gtcacgttca tcgcatcgaa gggctgtccc agaagtttat ttacctaaat 1200
gatgatgtca tgtttgggaa ggatgtctgg ccagatgatt tttacagtca ctccaaaggc 1260
cagaagggtt atttgacatg gcctgtgcc aactgtgccg agggctgccc aggttcctgg 1320
attaaggatg gctattgtga caaggcttgt aataattcag cctgcgattg ggatggagg 1380
gattgctctg gaaacagtgg agggagtgc tatattgcag gaggtggagg tactgggagt 1440
attggagttg gacagccctg gcagtttggg ggaggaataa acagtgtctc ttactgtaat 1500
cagggatgtg cgaattcctg gctcgtgat aagttctgtg accaagcatg caatgtcttg 1560
tcctgtgggt ttgatgctgg cgactgtggg caagatcatt ttcatgaatt gtataaagt 1620
atccttctcc caaaccagac tcactatatt attccaaaag gtgaatgcct gccttatttc 1680
agctttgcag aagtagccaa aagaggagtt gaaggtgcct atagtgaaca tccaataatt 1740
cgacatgctt ctattgccaa caagtggaaa accatccacc tcataatgca cagtggaatg 1800
aatgccacca caatacattt taatctcacg tttcaaaata caaacgatga agagttcaaa 1860
atgcagataa cagtggaggt ggacacaagg gagggacca aactgaattc tacggcccag 1920
aagggttacg aaaatttagt tagtcccata acacttcttc cagaggcgga aatccttttt 1980
gaggatattc ccaaagaaaa acgcttcccg aagtttaaga gacatgatgt taactcaaca 2040
aggagagccc aggaagaggt gaaaattccc ctggtaaata tttcactcct tccaaaagac 2100
gccagttga gtctcaatac cttggatttg caactggaac atggagacat cactttgaaa 2160
ggatacaatt tgtccaagtc agccttgctg agatcatttc tgatgaactc acagcatgct 2220
aaaaataaaa atcaagctat aataacagat gaaacaaatg acagtttggg ggctccacag 2280
gaaaaacagg ttcataaaaag catcttgcca aacagcttag gagtgtctga aagattgcag 2340
aggttgactt ttcctgcagt gagtgtaaaa gtgaatgggtc atgaccaggg tcagaatcca 2400
cccctggact tggagaccac agcaagattt agagtggaaa ctcacacca aaaaaccata 2460
ggcggaaatg tgacaaaaga aaagccccc tctctgattg ttccactgga aagccagatg 2520
acaaaagaaa agaaaatcac agggaaagaa aaagagaaca gtagaatgga ggaaaatgct 2580
gaaaatcaca taggcgttac tgaagtgtta cttggaagaa agctgcagca ttacacagat 2640
agttacttgg gctttttgcc atgggagaaa aaaaagtatt tcctagatct tctcgacgaa 2700
gaagagtcac tgaagacaca attggcatac ttcactgata gcaagaatac tgggaggcaa 2760
ctaaaagata catttgcaga ttccttcaga tatgtaaata aaattctaaa tagcaagttt 2820
ggattcacat cgcggaagat ccctgctcac atgcctcaca tgattgaccg gattgttatg 2880
caagaactgc aagatatgtt ccctgaagaa tttgacaaga cgtcatttca caaagtgcgc 2940
cattctgagg atatgcagtt tgccttctct tatttttatt atctcatgag tgcagtgcag 3000
ccactgaata tatctcaagt ctttgatgaa gttgatacag atcaatctgg tgtcttgtct 3060
gacagagaaa tccgaacact ggctaccaga attcacgaac tgccgttaag tttgcaggat 3120
ttgacaggtc tggaacacat gctaataaat tgctcaaaaa tgcttctctg tgatatcacg 3180
cagctaaata atattccacc aactcaggaa tcctactatg atcccaacct gccaccggtc 3240
actaaaagtc tagtaacaaa ctgtaaacca gtaactgaca aaatccacaa agcatataag 3300
gacaaaaaca aatatagggt tgaaatcatg ggagaagaag aaatcgcttt taaaatgatt 3360
cgtaccaacg tttctcatgt ggttggccag ttggatgaca taagaaaaaa ccctaggaag 3420
tttgtttgcc tgaatgacaa cattgaccac aatcataaag atgctcagac agtgaaggct 3480
gttctcaggg acttctatga atccatgttc ccatacctt cccaatttga actgccaaga 3540
gagtatcgaa accgtttcct tcatatgcat gagctgcagg aatggagggc ttatcgagac 3600
aaattgaagt agtagtctag a
3621

```

<210> 22
<211> 1383
<212> DNA
<213> Homo sapiens

```

<400> 22
atggcgacct ccacgggtcg ctggcttctc ctccggcttg cactattcgg cttcctctgg 60
gaagcgtccg gcggcctcga ctcgggggcc tcccgcgacg acgacttgct actgccctat 120
ccacgcgcgc gcgcgcgcct ccccggggac tgcacacggg tgcgcgcggg caaccgcgag 180
cacgagagtt ggctccgcc tcccgcgact cccggcgccg gcggtctggc cgtgcgcacc 240
ttcgtgtcgc acttcaggga ccgcgcggtg gccggccacc tgacgcgggc cgttgagccc 300
ctgcgcacct tctcgggtgct ggagcccggg ggacccggcg gctgcgcggc gagacgacgc 360
gccaccgtgg aggagacggc gcgggcggcc gactgccgtg tcgcccagaa cggcggcttc 420
ttccgcataa actcgggcga gtgcctgggg aacgtggtga gcgacgagcg gcgggtgagc 480

```

```

agctccgggg ggctgcagaa cgcgcagttc gggatccgcc gcgacgggac cctgggtcacc 540
gggtacctgt ctgaggagga ggtgctggac actgagaacc catttgtgca gctgctgagt 600
ggggtcgtgt ggctgattcg taatggaagc atctacatca acgagagcca agccacagag 660
tgtgacgaga cacaggagac aggttccttt agcaaatttg tgaatgtgat atcagccagg 720
acggccattg gccacgaccg gaaagggcag ctggtgctct ttcattgcaga cggccatacg 780
gagcagcgtg gcatcaacct gtgggaaatg gcggagttcc tgctgaaaca ggacgtggtc 840
aacgccatca acctggatgg ggggtggctct gccacctttg tgctcaacgg gaccttggcc 900
agttaccctg cagatcactg ccaggacaac atgtggcgct gtccccgcca agtgtccacc 960
gtggtgtgtg tgcacgaacc ccgctgccag ccgcctgact gccacggcca cgggacctgc 1020
gtggacgggc actgccaatg caccgggcac ttctggcggg gtcccggctg tgatgagctg 1080
gactgtggcc cctctaactg cagccagcac ggactgtgca cggagaccgg ctgccgctgt 1140
gatgccggat ggaccgggtc caactgcagt gaagagtgtc cccttggctg gcatgggccc 1200
ggctgccaga ggccttgtaa gtgtgagcac cattgtccct gtgaccccaa gactggcaac 1260
tgcagcgtct ccagagtaaa gcagtgtctc cagccacctg aagccaccct gagggcggga 1320
gaactctcct ttttcaccag ggaggaccag gtggacccca ggctgatcga cggcaaggat 1380
tga

```

<210> 23
 <211> 32
 <212> PRT
 <213> Bos taurus

<220>
 <223> Xaa is one of Ala, Arg, Asn, Asp, Cys, Gln, Glu,
 Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr,
 Trp, Tyr, or Val

<400> 23
 Asp Xaa Thr Arg Val His Ala Gly Arg Leu Glu His Glu Ser Trp Pro
 1 5 10 15
 Pro Ala Ala Gln Thr Ala Gly Ala His Arg Pro Ser Val Arg Thr Phe
 20 25 30

<210> 24
 <211> 20
 <212> PRT
 <213> Bos taurus

<400> 24
 Arg Asp Gly Thr Leu Val Thr Gly Tyr Leu Ser Glu Glu Glu Val Leu
 1 5 10 15

Asp Thr Glu Asn
 20

<210> 25
 <211> 13
 <212> PRT
 <213> Bos taurus

<400> 25
 Gly Ile Asn Leu Trp Glu Met Ala Glu Phe Leu Leu Lys
 1 5 10

<210> 26
 <211> 13
 <212> PRT
 <213> Bos taurus

<400> 26
 Met Leu Leu Lys Leu Leu Gln Arg Gln Arg Gln Thr Tyr
 1 5 10

<210> 27
 <211> 28
 <212> PRT
 <213> Bos taurus

<400> 27
 Asp Thr Phe Ala Asp Ser Leu Arg Tyr Val Asn Lys Ile Leu Asn Ser
 1 5 10 15

Lys Phe Gly Phe Thr Ser Arg Lys Val Pro Ala His
 20 25

<210> 28
 <211> 21
 <212> PRT
 <213> Bos taurus

<400> 28
 Ala Lys Met Lys Val Val Glu Glu Pro Asn Thr Phe Gly Leu Asn Asn
 1 5 10 15

Pro Phe Leu Pro Gln
 20

<210> 29
 <211> 5
 <212> PRT
 <213> Bos taurus

<400> 29
 Ile Leu Asn Ser Lys
 1 5

<210> 30
 <211> 5
 <212> PRT
 <213> Bos taurus

<400> 30
 Thr Ser Phe His Lys
 1 5

<210> 31
 <211> 5
 <212> PRT
 <213> Bos taurus

<400> 31

Phe Gly Phe Ser Arg
 1 5

<210> 32

<211> 12

<212> PRT

<213> Bos taurus

<400> 32

Ser Leu Val Thr Asn Cys Lys Pro Val Thr Asp Lys
 1 5 10

<210> 33

<211> 12

<212> PRT

<213> Bos taurus

<400> 33

Leu Ala His Val Ser Glu Pro Ser Thr Cys Val Tyr
 1 5 10

<210> 34

<211> 13

<212> PRT

<213> Bos taurus

<400> 34

Asn Asn Pro Phe Leu Pro Gln Thr Ser Arg Leu Gln Pro
 1 5 10

<210> 35

<211> 16

<212> PRT

<213> Bos taurus

<220>

<223> Xaa is one of Ala, Arg, Asn, Asp, Cys, Gln, Glu,
 Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr,
 Trp, Tyr or Val

<400> 35

Val Pro Met Leu Val Leu Asp Xaa Ala Pro Thr Xaa Val Xaa Leu Lys
 1 5 10 15

<210> 36

<211> 22

<212> PRT

<213> Bos taurus

<400> 36

Glu Leu Pro Ser Leu Tyr Pro Ser Phe Leu Ser Ala Ser Asp Val Phe
 1 5 10 15

Asn Val Ala Lys Pro Lys

20

<210> 37
<211> 25
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 37
gcgaagatga aggtggtgga ggacc

25

<210> 38
<211> 24
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 38
tgcagagaca gacctatacc tgcc

24

<210> 39
<211> 23
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 39
actcacctct ccgaactgga aag

23

<210> 40
<211> 29
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 40
ctagccacca tggggttcaa gctcttgca

29

<210> 41
<211> 21
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 41
agagcttgaa ccccatggtg g

21

<210> 42
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 42
gaagacacaa ttggcatact tcactgatag caagaatact gggaggcaac taaaagatac 60

<210> 43
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 43
actgcatatc ctcagaatgg 20

<210> 44
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 44
tggttctgaa gcttagccga gatcaatacc atg 33

<210> 45
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic DNA

<400> 45
tagtacactc tagactacta cttcaatttg tctcgataag 40

<210> 46
<211> 218
<212> DNA
<213> Unknown Organism

<220>
<223> Description of Unknown Organism:mouse/human hybrid
sequences

<400> 46
ctagccgcca ccatggagac agacacactc ctgctatggg tactgctgct cggcggtggt 60

acctctgtct gtgtgaggac gatacccatg acgacgagtg ggttccaggt tccactgggtg 120
 acgaagatca ggtagatccg cggttaatca cccaaggtcc aaggtgacca ctgcttctag 180
 tccatctagg cgccaattag gacggtactg ccattcga 218

<210> 47
 <211> 205
 <212> DNA
 <213> Unknown Organism

<220>
 <223> Description of Unknown Organism:mouse/human hybrid
 sequences

<400> 47
 ctagcggtag catgagatta gcagtaggcg ccttattagt atgcgcagta ctccgccatg 60
 gtactctaata cgtcatccgc ggaataatca tacgcgtcat gagggattat gtctcgcaga 120
 agatcaggta gatccgcggt taatcgacgg taccttatac agagcgtctt ctagtccatc 180
 taggcgcca ttagctgcca ttcga 205

<210> 48
 <211> 207
 <212> DNA
 <213> Unknown Organism

<220>
 <223> Description of Unknown Organism:mouse/human hybrid
 sequences

<400> 48
 ctagccgcca ccatgggatt agcagtaggc gccttattag tatgcgcagt cgccgggtggt 60
 accctaatac tcatccgcgg aataatcata cgcgtaact cggattatgt ctgcgagaag 120
 atcaggtaga tccgcggtta atcgacgtga gcctaataca gagcgtcttc tagtccatct 180
 aggcgccaat tagctgcgta cattcga 207

<210> 49
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 49
 ggaattccac catggcgacc tccacgggtc g 31

<210> 50
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 50
 tgaccagggt cccgtcgcg 19

<210> 51
 <211> 39
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic DNA

<400> 51
 gaggaccagg tggaccccag gctgatccac ggcaaggat 39

<210> 52
 <211> 13
 <212> PRT
 <213> Homo sapiens

<400> 52
 Glu Asp Gln Val Asp Pro Arg Leu Ile Asp Gly Lys Asp
 1 5 10

CLAIMS:

1. A method of modifying a human lysosomal hydrolase comprising contacting said human lysosomal hydrolase with an isolated N-acetylglucosamine-phosphotransferase to produce a modified human lysosomal hydrolase, wherein said N-acetylglucosamine-phosphotransferase comprises SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3, and purifying said modified human lysosomal hydrolase after said contacting.
2. The method of claim 1, wherein said human lysosomal hydrolase comprises an asparagine-linked oligosaccharide with a high mannose structure.
3. The method of claim 1, wherein said N-acetylglucosamine-phosphotransferase catalyzes the transfer of N-acetylglucosamine-1-phosphate from UDP-N-Acetylglucosamine to a mannose on the human lysosomal hydrolase.
4. The method of claim 1, wherein said human lysosomal hydrolase is a recombinant hydrolase.
5. The method of claim 1, wherein said N-acetylglucosamine-phosphotransferase comprises amino acids 1-928 of SEQ ID NO:1, amino acids 1-328 of SEQ ID NO:2, and amino acids 25-305 of SEQ ID NO:3.
6. The method of claim 1, wherein said lysosomal hydrolase is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylglucosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucuronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase G_{M1} Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.

7. The method of claim 6, wherein said human lysosomal hydrolase is α -glucosidase.
8. A phosphorylated purified human lysosomal hydrolase comprising a terminal mannose-6-phosphate.
9. A phosphorylated purified human lysosomal hydrolase comprising a mannose 6-phosphate, which comprises at least 6% bis-phosphorylated oligosaccharides.
10. The phosphorylated purified human lysosomal hydrolase of claim 8 or 9, which comprises up to 100% bis-phosphorylated oligosaccharides.
11. The phosphorylated purified human lysosomal hydrolase of claim 8 or 9, which comprises at least 5 mannose 6-phosphates.
12. The phosphorylated purified human lysosomal hydrolase of claim 8 or 9, which is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylgalactosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucuronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase G_{M1} Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.
13. The phosphorylated purified human lysosomal hydrolase of claim 12, which is α -glucosidase.
14. A method of preparing a phosphorylated human lysosomal hydrolase comprising contacting said human lysosomal hydrolase with an isolated phosphodiester α -N-Acetylglucosaminidase, wherein said lysosomal hydrolase comprises a terminal mannose 6-phosphate, and said phosphodiester α -N-Acetylglucosaminidase comprises the amino acid sequence in SEQ ID

- NO:6, and purifying the phosphorylated human lysosomal hydrolase after said contacting.
15. The method of claim 14, wherein said human lysosomal hydrolase is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylgalactosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucoronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase G_{M1} Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.
 16. The method of claim 14, wherein said phosphodiester α -N-Acetylglucosaminidase catalyzes the removal of N-acetylglucosamine from said modified human lysosomal hydrolase and generates a terminal mannose 6-phosphate on said hydrolase.
 17. The method of claim 14, wherein said phosphodiester α -N-Acetylglucosaminidase comprises amino acids 50-515 of SEQ ID NO:6.
 18. The method of claim 15, wherein said human lysosomal hydrolase is α -glucosidase.
 19. A method of preparing a phosphorylated human lysosomal hydrolase comprising: contacting said human lysosomal hydrolase with an isolated N-acetylglucosamine-phosphotransferase to produce a modified human lysosomal hydrolase, wherein said N-acetylglucosamine-phosphotransferase comprises SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3; contacting said modified human lysosomal hydrolase with an isolated phosphodiester α -N-Acetylglucosaminidase, wherein said N-phosphodiester α -N-Acetylglucosaminidase comprises the amino acid in SEQ ID NO:6; and

purifying said phosphorylated human lysosomal hydrolase prior to or after said contacting with the isolated phosphodiester α -N-Acetylglucosaminidase.

20. The method of claim 19, wherein said modified lysosomal hydrolase is purified prior to said contacting with the isolated phosphodiester α -N-Acetylglucosaminidase.
21. The method of claim 19, wherein said modified lysosomal hydrolase is purified after said contacting with the isolated phosphodiester α -N-Acetylglucosaminidase.
22. The method of claim 19, wherein said N-acetylglucosamine-phosphotransferase comprises amino acids 1-928 of SEQ ID NO:1, amino acids 1-328 of SEQ ID NO:2, and amino acids 25-305 of SEQ ID NO:3.
23. The method of claim 19, wherein said phosphodiester α -N-Acetylglucosaminidase comprises amino acids 50-515 of SEQ ID NO:6.
24. The method of claim 19, wherein said human lysosomal hydrolase is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylgalactosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucoronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase G_{M1} Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.
25. The method of claim 24, wherein said human lysosomal hydrolase is α -glucosidase.
26. A method of modifying a human lysosomal hydrolase comprising contacting said human lysosomal hydrolase with an isolated N-acetylglucosamine-phosphotransferase to produce a modified human lysosomal hydrolase, wherein

said N-acetylglucosamine-phosphotransferase comprises an amino acid sequence that is at least 70% identical to SEQ ID NO:1, an amino acid sequence that is at least 70% identical to SEQ ID NO:2, and an amino acid sequence that is at least 70% identical to SEQ ID NO:3, and said N-acetylglucosamine-phosphotransferase catalyzes the transfer of N-acetylglucosamine-1-phosphate from UDP-N-Acetylglucosamine to a mannose on the human lysosomal hydrolase; and purifying said modified human lysosomal hydrolase after said contacting.

27. The method of claim 26, wherein said human lysosomal hydrolase comprises an asparagine-linked oligosaccharide with a high mannose structure.
28. The method of claim 26, wherein said human lysosomal hydrolase is a recombinant hydrolase.
29. The method of claim 26, wherein said human lysosomal hydrolase is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylgalactosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucuronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase GM, Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.
30. The method of claim 29, wherein said lysosomal hydrolase is α -glucosidase.
31. The method of claim 26, further comprising contacting said modified lysosomal hydrolase with an isolated phosphodiester α -N-Acetylglucosaminidase, which comprises an amino acid sequence that is at least 70% identical to SEQ ID NO:6, and wherein said phosphodiester α -N-Acetylglucosaminidase catalyzes the removal of N-acetylglucosamine from said modified lysosomal hydrolases and generates a terminal mannose 6-phosphate on said hydrolase.

32. The method of claim 31, further comprising purifying said lysosomal hydrolase after contacting with the isolated phosphodiester α -N-Acetylglucosaminidase.
33. A method of modifying a human lysosomal hydrolase comprising contacting said human lysosomal hydrolase with an isolated N-acetylglucosamine-phosphotransferase, which has a specific activity of at least 10^6 pmol/h/mg to produce a modified human lysosomal hydrolase, and purifying said modified human lysosomal hydrolase after said contacting.
34. The method of claim 33, wherein said N-acetylglucosamine-phosphotransferase catalyzes the transfer of N-acetylglucosamine-1-phosphate from UDP-N-Acetylglucosamine to a mannose on the hydrolase.
35. The method of claim 33, wherein said human lysosomal hydrolase is a recombinant hydrolase.
36. The method of claim 33, wherein said human lysosomal hydrolase is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylgalactosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucuronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase G_{M1} Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.
37. The method of claim 1, further comprising contacting said human modified lysosomal hydrolase with an isolated phosphodiester α -N-Acetylglucosaminidase, which catalyzes the removal of N-acetylglucosamine from said modified lysosomal hydrolase and generates a terminal mannose 6-phosphate on said hydrolase.
38. The method of claim 33, wherein said N-acetylglucosamine-phosphotransferase has a specific activity of at least 5×10^6 pmol/h/mg.

39. The method of claim 33, wherein said N-acetylglucosamine-phosphotransferase has a specific activity of at least 12×10^6 pmol/h/mg.
40. The method of claim 37, wherein said isolated phosphodiester α -N-Acetylglucosaminidase has a specific activity of at least about 1000 units/mg.
41. The method of claim 37, wherein said isolated phosphodiester α -N-Acetylglucosaminidase has a specific activity of at least about 472,000 units/mg.
42. The method of claim 33, wherein the N-acetylglucosamine-phosphotransferase comprises an α subunit, a β subunit, and a γ subunit; and wherein the α and β subunits are encoded by a DNA molecule comprising SEQ ID NO:20; and the γ subunit is encoded by a DNA molecule comprising nucleotides 96 to 941 of SEQ ID NO:5.
43. The method of claim 33, wherein the N-acetylglucosamine-phosphotransferase comprises an amino acid sequence that is at least 70% identical to SEQ ID NO:1, an amino acid sequence that is at least 70% identical to SEQ ID NO:2, and an amino acid sequence that is at least 70% identical to SEQ ID NO: 3, and said N-acetylglucosamine-phosphotransferase catalyzes the transfer of N-acetylglucosamine-1-phosphate from UDP-N-Acetylglucosamine to a mannose on the lysosomal hydrolase.
44. The method of claim 33, wherein the lysosomal hydrolase is α -glucosidase.
45. The method of claim 33, wherein the lysosomal hydrolase is α -iduronidase.
46. The method of claim 33, wherein the lysosomal hydrolase is α -galactosidase A.
47. A method of preparing a phosphorylated human lysosomal hydrolase comprising contacting said human lysosomal hydrolase with an isolated phosphodiester α -N-Acetylglucosaminidase, which has a specific activity of at least about 472,000 units/mg and which catalyzes the removal of N-acetylglucosamine from said human lysosomal hydrolase and generates a terminal mannose 6-phosphate on said hydrolase, and wherein said human

- lysosomal hydrolase comprises a N-acetylglucosamine phosphomannose diester; and purifying the phosphorylated human lysosomal hydrolase.
48. The method of claim 47, wherein said human lysosomal hydrolase is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylgalactosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucoronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase G_{M1} Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.
 49. The method of claim 47, wherein said phosphodiester α -N-Acetylglucosaminidase catalyzes the removal of N-acetylglucosamine from N-acetylglucosamine phosphomannose diester to generate a terminal mannose 6-phosphate on said human lysosomal hydrolase.
 50. The method of claim 47, wherein the phosphodiester α -N-Acetylglucosaminidase is encoded by a DNA molecule comprising nucleotides 151 to 1548 of SEQ ID NO:7.
 51. The method of claim 47, wherein the phosphodiester α -N-Acetylglucosaminidase is encoded by a DNA molecule which hybridizes under stringent conditions to the complement of nucleotides 151 to 1548 of SEQ ID NO:7.
 52. The method of claim 47, wherein the human lysosomal hydrolase is α -glucosidase.
 53. The method of claim 47, wherein the human lysosomal hydrolase is α -iduronidase.

54. The method of claim 47, wherein the human lysosomal hydrolase is α -galactosidase A.
55. A composition comprising the human lysosomal hydrolase of any one of claims 8-13 and a carrier.
56. A method of preparing a phosphorylated human lysosomal hydrolase comprising: contacting said lysosomal hydrolase with an isolated N-acetylglucosamine-phosphotransferase, which has a specific activity of at least 10^6 pmol/h/mg to produce a modified human lysosomal hydrolase; and contacting said modified human lysosomal hydrolase with an isolated phosphodiester α -N-Acetylglucosaminidase, which catalyzes the removal of N-acetylglucosamine from said modified human lysosomal hydrolase and generates a terminal mannose 6-phosphate on said hydrolase.
57. The method of claim 56, further comprising purifying said phosphorylated human lysosomal hydrolase after said contacting with the isolated phosphodiester α -N-Acetylglucosaminidase.
58. The method of claim 56, further comprising purifying said modified human lysosomal hydrolase prior to said contacting with the isolated phosphodiester α -N-Acetylglucosaminidase.
59. The method of claim 56, wherein said human lysosomal hydrolase is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylgalactosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucuronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase G_{M1} Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.

60. The method of claim 56, wherein said N-acetylglucosamine-phosphotransferase has a specific activity of at least 5×10^6 pmol/h/mg.
61. The method of claim 56, wherein said N-acetylglucosamine-phosphotransferase has a specific activity of at least 12×10^6 pmol/h/mg.
62. The method of claim 56, wherein the phosphorylated human lysosomal hydrolase comprises at least 6% bis-phosphorylated oligosaccharides.
63. The method of claim 56, wherein the phosphorylated human lysosomal hydrolase comprises at least 100% bis-phosphorylated oligosaccharides.
64. The method of claim 56, wherein the phosphorylated human lysosomal hydrolase comprises at least 5 mannose 6-phosphates.
65. The method of claim 56, wherein the N-acetylglucosamine-phosphotransferase comprises an α subunit, a β subunit, and a γ subunit; and wherein the α and β subunits are encoded by a DNA molecule comprising SEQ ID NO:20; and the γ subunit is encoded by a DNA molecule comprising nucleotides 96 to 941 of SEQ ID NO:5.
66. The method of claim 56, wherein the phosphodiester α -N-Acetylglucosaminidase is encoded by a DNA molecule comprising nucleotides 151 to 1548 of SEQ ID NO:7.
67. The method of claim 56, wherein the N-acetylglucosamine-phosphotransferase comprises an amino acid sequence that is at least 70% identical to SEQ ID NO:1, an amino acid sequence that is at least 70% identical to SEQ ID NO:2, and an amino acid sequence that is at least 70% identical to SEQ ID NO: 3, and said N-acetylglucosamine-phosphotransferase catalyzes the transfer of N-acetylglucosamine-1-phosphate from UDP-N-Acetylglucosamine to a mannose on the lysosomal hydrolase.
68. The method of claim 56, wherein the N-acetylglucosamine-phosphodiester α -N-Acetylglucosaminidase comprises an amino acid sequence that is at least 70% identical to SEQ ID NO:6, and wherein said N-acetylglucosamine-phosphodiester α -N-Acetylglucosaminidase catalyzes the removal of N-

acetylglucosamine from said modified lysosomal hydrolases and generates a terminal mannose 6-phosphate on said hydrolase.

69. The method of claim 56, wherein the human lysosomal hydrolase is α -glucosidase.
70. The method of claim 56, wherein the human lysosomal hydrolase is α -iduronidase.
71. The method of claim 56, wherein the human lysosomal hydrolase is α -galactosidase A.
72. The method of claim 56, wherein said isolated phosphodiester α -N-Acetylglucosaminidase has a specific activity of at least about 1000 units/mg.
73. The method of claim 56, wherein said isolated phosphodiester α -N-Acetylglucosaminidase has a specific activity of at least about 472,000 units/mg.
74. A method of producing a modified human lysosomal hydrolase comprising culturing transformed cells comprising a recombinant polynucleotide encoding a human lysosomal hydrolase in the presence of an α 1,2-mannosidase inhibitor; recovering the human lysosomal hydrolase from the cultured transformed cells; and contacting the recovered human lysosomal hydrolase with an isolated N-acetylglucosamine-phosphotransferase to produce a modified human lysosomal hydrolase, wherein said N-acetylglucosamine-phosphotransferase comprises SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3.
75. The method of claim 74, further comprising purifying said modified human lysosomal hydrolase after said contacting.
76. The method of claim 74, wherein said N-acetylglucosamine-phosphotransferase catalyzes the transfer of N-acetylglucosamine-1-phosphate from UDP-N-Acetylglucosamine to a mannose on the human lysosomal hydrolase.
77. The method of claim 74, wherein said N-acetylglucosamine-phosphotransferase comprises amino acids 1-928 of SEQ ID NO:1, amino acids 1-328 of SEQ ID NO:2, and amino acids 25-305 of SEQ ID NO:3.


78. The method of claim 74, wherein said human lysosomal hydrolase is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylgalactosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucoronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase G_{M1} Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.
79. The method of claim 74, further comprising contacting said modified human lysosomal hydrolase with an isolated phosphodiester α -N-Acetylglucosaminidase, which comprises the amino acid sequence in SEQ ID NO:6.
80. The method of claim 79, wherein said phosphodiester α -N-Acetylglucosaminidase catalyzes the removal of N-acetylglucosamine from said modified human lysosomal hydrolase and generates a terminal mannose 6-phosphate on said hydrolase.
81. The method of claim 79, wherein said phosphodiester α -N-Acetylglucosaminidase comprises amino acids 50-515 of SEQ ID NO:6.
82. The method of claim 74, wherein said alpha 1,2-mannosidase inhibitor is selected from the group consisting of deoxymannojirimycin, kifunensine, D-Mannonolactam amidrazone and N-butyl-dexoymannojirimycin.
83. The method of claim 74, wherein the alpha 1,2-mannosidase inhibitor is kifunensine.
84. The method of claim 74, wherein the alpha 1,2-mannosidase inhibitor is deoxymannojirimycin.
85. The method of claim 79, wherein said lysosomal hydrolase is α -glucosidase.

86. A method of modifying a human lysosomal hydrolase comprising culturing transformed cells comprising a recombinant polynucleotide encoding a human lysosomal hydrolase in the presence of an α 1,2-mannosidase inhibitor; recovering the human lysosomal hydrolase from the cultured transformed cells; and contacting said recovered human lysosomal hydrolase with an isolated N-acetylglucosamine-phosphotransferase to produce a modified human lysosomal hydrolase, wherein said N-acetylglucosamine-phosphotransferase comprises an amino acid sequence that is at least 70% identical to SEQ ID NO:1, an amino acid sequence that is at least 70% identical to SEQ ID NO:2, and an amino acid sequence that is at least 70% identical to SEQ ID NO:3, and said N-acetylglucosamine-phosphotransferase catalyzes the transfer of N-acetylglucosamine-1-phosphate from UDP-N-Acetylglucosamine to a mannose on the human lysosomal hydrolase.
87. The method of claim 86, further comprising purifying said modified human lysosomal hydrolase after said contacting.
88. The method of claim 86, wherein said human lysosomal hydrolase is selected from the group consisting of α -glucosidase, α -iduronidase, α -galactosidase A, arylsulfatase, N-acetylgalactosamine-6-sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, B-glucoronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α -glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside, Acid β -galactosidase G.sub.MI Galglioside, Acid β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.
89. The method of claim 86, wherein said human lysosomal hydrolase is α -glucosidase.
90. The method of claim 86, further comprising contacting said modified human lysosomal hydrolase with an isolated phosphodiester

α -N-Acetylglucosaminidase which comprises an amino acid sequence that is at least 70% identical to SEQ ID NO:6, and wherein said phosphodiester α -N-Acetylglucosaminidase catalyzes the removal of N-acetylglucosamine from said modified lysosomal hydrolases and generates a terminal mannose 6-phosphate on said hydrolase.

91. The method of claim 90, further comprising purifying said human lysosomal hydrolase after contacting with the isolated phosphodiester α -N-Acetylglucosaminidase.
92. The method of claim 86, wherein said alpha 1,2-mannosidase inhibitor is selected from the group consisting of deoxymannojirimycin, kifunensine, D-Mannonolactam amidrazone and N-butyl-dexoymannojirimycin.
93. The method of claim 86, wherein the alpha 1,2-mannosidase inhibitor is kifunensine.
94. The method of claim 86, wherein the alpha 1,2-mannosidase inhibitor is deoxymannojirimycin.
95. An isolated nucleic acid which encodes a protein comprising the amino acid sequence of SEQ ID NO:1, SEQ ID NO:2, or SEQ ID NO:3.
96. An isolated nucleic acid which hybridizes under stringent conditions to the complement of a nucleic acid which encodes a protein having the amino acid sequence of SEQ ID NO:1, SEQ ID NO:2, or SEQ ID NO:3, wherein said stringent conditions comprise washing in 0.2x SSC and 0.1% SDS at 65 °C, and wherein the nucleic acid encodes a biologically active N-acetylglucosamine-phosphotransferase.
97. A vector comprising the isolated nucleic acid of claim 95 or 96.
98. A host cell comprising the isolated nucleic acid of claim 95 or 96 or the vector of claim 97.
99. A method of producing N-acetylglucosamine-phosphotransferase comprising culturing the cell of claim 98 under conditions suitable for expression of the isolated nucleic acid molecule; and recovering the N-acetylglucosamine-phosphotransferase from the cell.

100. A composition comprising the isolated nucleic acid of claim 95 or 96 and a carrier.
101. A composition comprising the vector of claim 97 and a carrier.
102. An isolated peptide encoded by any of the isolated nucleic acids of claim 95 or 96.
103. Use of a composition for treating Pompe's disease in a patient in need thereof, the composition comprising recombinant human α -glucosidase and a pharmaceutically acceptable carrier, wherein the composition has an average of at least 5 mannose 6-phosphates per recombinant human α -glucosidase.
104. Use of a composition for treating Pompe's disease in a patient in need thereof, the composition comprising recombinant human α -glucosidase and a physiologically acceptable carrier, wherein the composition has an average of about at least 6% bis-phosphorylated oligosaccharides per recombinant human α -glucosidase.
105. A pharmaceutical composition for the treatment of Pompe disease in a human patient, comprising recombinant human α -glucosidase having an average of at least 5 mannose-6-phosphates per recombinant human α -glucosidase and a pharmaceutically acceptable carrier.
106. A pharmaceutical composition for the treatment of Pompe disease in a human patient, comprising recombinant human α -glucosidase having an average of about at least 6% bis-phosphorylated oligosaccharides per recombinant human α -glucosidase and a pharmaceutically effective carrier.







FIG. 2

FIG. 3






FIG. 4