(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
16 June 2005 (16.06.2005)

PCT

(10) International Publication Number
WO 2005/053700 A1

(51) International Patent Classification[7]: A61K 31/5375, A61P 25/14

(21) International Application Number:
PCT/EP2004/013248

(22) International Filing Date:
19 November 2004 (19.11.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0327195.4 21 November 2003 (21.11.2003) GB

(71) Applicant *(for all designated States except US)*: **SMITHKLINE BEECHAM CORPORATION** [US/US]; One Franklin Plaza, P O Box 7929, Philadelphia, Pennsylvania 19101 (US).

(72) Inventor; and
(75) Inventor/Applicant *(for US only)*: **BARLAND, Jean-Christophe** [FR/GB]; GlaxoSmithKline, Stockley Park, Uxbridge Middlesex UB11 1BU (GB).

(74) Agent: **REED, Michael**; GlaxoSmithKline, Corporate Intellectual Property CN925.1, 980 Great West Road, Brentford Middlesex TW8 9GS (GB).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2005/053700 A1

(54) Title: USE OF (2S,3S)2-(3-CHLOROPHENYL)-3,5,5-TRIMETHYL-2-MORPHOLINOL


(57) Abstract: This invention relates to a novel use of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol, in particular its use in the treatment of Restless Legs Syndrome, or its use in the treatment of Periodic Limb Movement Disorder (PLMD).

USE OF (2S, 3S) 2-(3-CHLOROPHENYL)-3,5,5-TRIMETHYL-2-MORPHOLINOL

This invention relates to a novel use of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol, in particular its use in the treatment of Restless Legs Syndrome, or its use in the treatment of Periodic Limb Movement Disorder (PLMD).

Background of the Invention


Bupropion hydrochloride, (±)-1-(3-chlorophenyl)-2-[(1,1-dimethylethyl)-amino]-1-propanone hydrochloride, is the active ingredient of Wellbutrin® which is marketed in the United States for the treatment of depression. It is also the active ingredient of Zyban® which is marketed in the United States as an aid to smoking cessation. Bupropion is an inhibitor of the neuronal uptake of noradrenaline (NA), and dopamine (DA), does not inhibit monoamine oxidase and has a negligible effect on the neuronal uptake of seretonin. While the mechanism of action of bupropion, as with other antidepressants, is not fully understood, it is presumed that this action is mediated by noradrenergic and/or dopaminergic mechanisms. Initial clinical evidence suggested Wellbutrin® to be a selective inhibitor of noradrenaline (NA) at doses that are predictive of antidepressant activity in animal models (Ascher, J. A., *et al.*, *Journal of Clinical Psychiatry*, 56: p. 395-401,1995). A more recent analysis (Stahl, S. M. *et al.*, Prim. Care Companion, Journal of Clinical Psychiatry, 6(4), p 159-166, 2004) concludes that bupropion acts via dual inhibition of norepinephrine and dopamine reuptake, having slightly greater functional potency at the dopamine transporter.



Bupropion HCl

Bupropion is extensively metabolized in man as well as laboratory animals. Urinary and plasma metabolites include biotransformation products formed via hydroxylation of the tert-butyl group and/or reduction of the carbonyl group of bupropion. Four basic metabolites have been identified. They are the erythro- and threo-amino alcohols of bupropion, the erythro-amino diol of bupropion (found in urine but not in plasma), and a morpholinol metabolite.


The morpholinol metabolite (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol is believed to be formed from hydroxylation of the tert-butyl group of bupropion.



Morpholinol Metabolite of Bupropion

It was discovered that despite the (-) form of the morpholinol metabolite predominating significantly in human plasma samples, it was the (+) enantiomer, (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol in which the optimal monoamine reuptake inhibitory activity resides (WO 99/37305), hereinafter referred to as the compound of formula (I):

(+)-enantiomer



(I)

The compound of formula (I) and its salts and solvates have been disclosed as being of use in the treatment of depression (including major depressive disorder (MDD), bipolar depression (type I and II), major (unipolar) depression and depression with atypical features (eg. lethargy, over-eating/obesity, hypersomnia)), attention deficit hyperactivity disorder (ADHD), obesity, migraine, pain (including neuropathic pain, eg. diabetic neuropathy, sciatica, non-specific lower back pain, multiple sclerosis pain, fibromyalgia, HIV-related neuropathy, neuralgia such as as post-herpetic neuralgia and trigeminal neuralgia and pain resulting from physical trauma, amputation, cancer, toxins or chronic inflammatory conditions), sexual dysfunction (including inhibited sexual desire (low libido), inhibited sexual arousal or excitement, orgasm dysfunction, inhibited female orgasm, inhibited male orgasm, hypoactive sexual desire disorder (HSDD), female sexual desire disorder (FSDD) and sexual dysfunction side-effects induced by treatment with antidepressants of the SSRI-class), Parkinson's disease (including relief from the symptoms of Parkinson's disease which include, but are not limited to, locomotor deficits and/or motor disability, including slowly increasing disability in purposeful movement, tremors, bradykinesia, hyperkinesia (moderate and severe), akinesia, rigidity, disturbance of balance and co-ordination, and a disturbance of posture), Alzheimer's disease, or addiction to cocaine or nicotine-containing (especially tobacco) products (WO 99/37305 and US2003-0064988; both Glaxo Group Limited).

US2003-0032643 (Glaxo Group Limited) discloses the use of the compound

of formula (I) and its salts and solvates in the treatment of seasonal affective disorder, chronic fatigue, narcolepsy and cognitive impairment.

US2003-0083330 (Glaxo Group Limited) discloses the use of the compound of formula (I) and its salts and solvates in the treatment of addiction to alcohol.

WO 00/51546 and WO 01/62257 (both Separacor Inc) disclose the use of a bupropion metabolite in the treatment of a disorder that is ameliorated by the inhibition of neuronal monoamine reuptake, sexual dysfunction (including erectile dysfunction), an affective disorder (including depression, anxiety disorders, attention deficit hyperactivity disorder, bipolar and manic conditions, sexual dysfunction, psycho-sexual dysfunction, bulimia, obesity or weight gain, narcolepsy, chronic fatigue syndrome, seasonal affective disorder, premenstrual syndrome, and substance addiction or abuse), nicotine addicton, a cerebral function disorder (including senile dementia, Alzheimer's type dementia, memory loss, amnesia/amnestic syndrome, epilepsy, disturbances or consciousness, coma, lowering of attention, speech disorders, Parkinson's disease, Lennox syndrome, autistic disorder, autism, hyperkinetic syndrome, schizophrenia, cerebral infarction, cerebral bleeding, cerebral auteriosclerosis, cerebral venous thrombosis and head injury), epilepsy, smoking cessation and incontinence.

Dopaminergic agents such as L-Dopa, pergolide and agonists at the D_3 subtype of the D_2 receptor such as ropinerole and pramipexole are recommended by physicians as first choice treatments in RLS (Stiasny K., et al., *Restless legs syndrome and its treatment by dopamine agonists*, Parkinsonism and Related Disorders, 7, p21-25, 2001; Chesson A. L., *Practice parameters for the treatment of restless legs syndrome and periodic limb movement disorders,* Sleep, 22(7), p961-968, 1999). Reference is also made to a small retrospective study of bupropion SR in depressed patients (Nofzinger E. A. et al., *Bupropion SR reduces periodic limb movements associated with arousals from sleep in depressed patients with periodic limb movement disorder*, Journal of Clinical Psychiatry, 61, p858-862, 2000).

Summary of the Invention

The present invention provides the use of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof in the manufacture of a medicament for the treatment of Restless Legs Syndrome (RLS).

A further aspect of the invention provides a method of treating Restless Legs Syndrome (RLS) in a mammal (human or animal subject) comprising the administration to said subject of an effective amount of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof.

One further aspect of the present invention provides the use of enantiomerically pure (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof in the manufacture

of a medicament for the treatment of Restless Legs Syndrome (RLS).

A yet further aspect of the invention provides a method of treating Restless Legs Syndrome (RLS) in a mammal (human or animal subject) comprising the administration to said subject of an effective amount of enantiomerically pure (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof.

A further aspect of the present invention provides the use of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof in the manufacture of a medicament for the treatment of Periodic Limb Movement Disorder (PLMD).

A further aspect of the invention provides a method of treating Periodic Limb Movement Disorder (PLMD) in a mammal (human or animal subject) comprising the administration to said subject of an effective amount of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof.

One further aspect of the present invention provides the use of enantiomerically pure (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof in the manufacture of a medicament for the treatment of Periodic Limb Movement Disorder (PLMD).

A yet further aspect of the invention provides a method of treating Periodic Limb Movement Disorder (PLMD) in a mammal (human or animal subject) comprising the administration to said subject of an effective amount of enantiomerically pure (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof.

Detailed Description of the Invention

It will be appreciated that references herein to "treatment" extend to prophylaxis, prevention of recurrence and suppression or amelioration of symptoms (whether mild, moderate or severe) as well as the treatment of established conditions.

As used herein, "Restless Legs Syndrome (RLS)" is also known as the Ekbom syndrome and is a sensorimotor disorder with a general-population prevalence of 5–10%. RLS is an internationally recognised disorder listed in the diagnostic manuals ICD 10 (Chapter VI, G25.8; World Health Organisation, Geneva, 1994), DSM IV (Dyssomnia not otherwise specified 307.47) and also in the American Sleep Association International Classification of Sleep Disorders (ICSD) (Thorpy M.J, Chairman ICSD, Diagnostic Classification Steering Committee, Rochester Minnesota, 1990). RLS is characterised by stereotypical jerks of the lower limbs, typically during sleep (including periodic limb movements (PLMs)). RLS may also be characterised as an unpleasant twitching, burning or painful sensation, likened by sufferers to 'crawling ants' or 'writhing worms' in the muscles and bones which usually occurs during the evenings. The sensations usually occur in the calf,

sometimes in the thighs and feet and once they have begun, there is an irresistible urge to move the legs to release the feelings and general discomfort. Symptoms are worse or exclusively present at rest, in the evenings and at night, and are relieved by movement. The need to move occurs on average every 20 to 40 seconds and the movements last for about 1 to 5 seconds. For some patients, RLS is mild and causes little inconvenience, in others however, the impact on sleep is considerable, compromising work and social activities (Allen, R. P. and Earley, C. J. (2001) J Clin Neurophysiol;**18**:128–147 and Earley, C. J. (2003) N Engl J Med; **348**:2103–2109). A minority of RLS cases are secondary to a pre-existing condition (pregnancy, renal failure and iron-deficiency anaemia), and resolve with that underlying condition. Minimal criteria for the diagnosis of RLS were published by the International Restless Legs Syndrome Study Group (IRLSSG) in 1995 (Walters, A. S. (1995) Mov Disord;**10**:634–642) and updated in 2003 (Allen *et al.* (2003) Sleep Med;**4**:101–119).

The use of the compound of formula (I) or a salt or solvate thereof in the treatment of RLS may result in improvement in the subject's condition as determined by one or more of the following clinical measures, following administration:
PLMI (periodic limb movement index), PLMAI (periodic leg movement with arousal index), PLMW (periodic leg movements during wakefulness), and IRLS (International Restless Leg Syndrome) rating scale, although other measures may also be used as appropriate (for example Clinical Global Improvement score, domains of Medical Outcomes Study (MOS) Sleep Scale, St. Mary's Sleep Questionnaire Scale, and other measures of discomfort, sleep efficiency, sleep latency, or % sleep time in various stages in the sleep cycle).

In addition, the compound of formula (I) may be less prone to cause augmentation in the treatment of RLS than is the case with traditional agents such as L-Dopa.

Periodic Limb Movement Disorder (PLMD) is a condition related to RLS, also referenced in the American Sleep Association International Classification of Sleep Disorders (ICSD) (Thorpy M.J, Chairman ICSD, Diagnostic Classification Steering Committee, Rochester Minnesota, 1990). PLMD is characterized by repetitive stereotyped movements of the limbs during sleep. The movements, while most common in the legs, can also affect the arms. The sufferer may or may not notice the movements while sleeping. These movements typically occur every 20 to 40 seconds, and may be associated with repeated arousal, and severely fragmented sleep. Generally, to be diagnosed with the disorder, the periodic limb movements (PLMs) will occur five or more times during each hour of sleep. The PLMs are most common in the stage of sleep known as non-REM (Rapid Eye Movement) sleep, which usually occurs during the first half of the night. The disorder may cause poor sleep and/or subsequent daytime somnolence.

Although RLS and PLMD both affect the limbs - and both affect a person's ability to sleep at night and function normally during the day.- they are two different disorders. The movements of RLS occur most often when a person is awake and are

a voluntary response to uncomfortable or painful feelings in the legs. The movements of PLMD occur most often when a person is asleep and are involuntary (not consciously controlled). People with periodic limb movements are often not aware of these movements, although on rare occasions they may notice the involuntary movement of PLMD while they are still awake. It has been estimated that about 80% of RLS patients also have periodic limb movement disorder; however, patients with PLMD often do not have RLS.

The use of the compound of formula (I) or a salt or solvate thereof in the treatment of PLMD may result in improvement in the subject's condition as determined by one or more of the following clinical measures, following administration: PLMI (periodic limb movement index), and PLMAI (periodic leg movement with arousal index), although other measures may also be used as appropriate.

As used herein, "enantiomerically pure" means that the composition contains greater than about 90% of the desired enantiomer by weight, preferably greater than about 95% of the desired enantiomer by weight, more preferably greater than about 99% of the desired enantiomer by weight, most preferably greater than 99.5% of the desired enantiomer by weight, said weight percent based upon the total weight of the compound of formula (I).

Preferred for use according to the present invention are pharmaceutically acceptable salts or solvates of the compound of formula (I), particularly those disclosed in U.S. Patent No. 6,342,496 B1, U.S. Patent No. 6,337,328 B1, U.S. Patent No. 6,391,875 B1, U.S. Patent No. 6,274,579 B1, U.S. Patent Application Publication Nos. 2002/0052340 A1, 2002/0052341 A1, and 2003/0027827 A1, as well as WO 01/62257, WO 99/37305, WO 00/51546 and WO 01/62257. Suitable pharmaceutically acceptable salts can include, but are not limited to, hydrochloride salt, hydrogen sulfate salt and other sulfate salts, hydrogen phosphate salt and other phosphate salts, methanesulfonate salt, p-toluenesulfonate salt, citrate salt, fumarate salt, tartrate salt, and the like. Of these, (+)-(2S, 3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride is particularly preferred.

Preparation

The compound of formula (I) or a salt or solvate thereof may be prepared in isolated form, and preferably in an enantiomerically pure form, in accordance with the procedures set forth in WO 99/37305, US2003-0064988, US2003-0032643 and US2003-0027827 (all of Glaxo Group Limited) or WO 00/51546 and WO 01/62257 (both of Sepracor Inc.) the procedures of which are herein incorporated by reference.

Dosage and Formulation

The compound of formula (I) or a salt or solvate thereof is administered in isolated form, and is preferably administered in an enantiomerically pure form.

The amount of compound of formula (I) or a salt or solvate thereof required to

achieve the desired therapeutic effect will, of course depend on a number of factors, for example, the mode of administration and the recipient being treated. In general, the daily dose will be in the range of 0.02 to 5.0 mg/kg, more particularly 0.1 to 1.5mg/kg, or 0.15 to 1.2 mg/kg. More particular ranges include 0.02 to 2.5 mg/kg, 0.02 to 1.0 mg/kg, 0.1 to 1.5 mg/kg, 0.02 to 0.25 mg/kg, 0.02 to 0.15 mg/kg and 0.02 to 0.07 mg/kg given as a single once a day dose or as single or divided doses throughout the day. Preferably in the treatment of RLS, administration will be at appropriate time(s) of the day so that a peak of plasma concentration of the compound of formula (I) coincides with late evening or bedtime.

The compound of formula (I) or a salt or solvate thereof may be employed in the treatment of Restless Legs Syndrome (RLS) as the compound *per se*, but is preferably presented with one or more pharmaceutically acceptable carriers, diluents or excipients in the form of a pharmaceutical formulation. The carriers, diluents and exipients must, of course, be acceptable in the sense of being compatible with the other ingredients of the formulation and must not be deleterious to the recipient. The carrier may be a solid or a liquid, or both, and is preferably formulated with the agent as a unit-dose formulation, for example, a tablet containing 1mg, 2mg, 5mg, 10mg, 20mg, 40mg, 60mg, 80mg, 100mg, 120mg, 150mg and 200mg of the compound of formula (I) or a salt or solvate thereof, more preferably 10-80mg of the compound of formula (I) or a salt or solvate thereof. Suitable formulations for use in the present invention include sustained release solid-dosage formulations, optionally film-coated solid-dosage formulations, and especially tablet and caplet formulations, for oral administration of the compound of formula (I), particularly once-daily administration, for example those illustrated in Examples 1 to 5 below.

The formulations include those suitable for oral, rectal, topical, buccal (e.g. sub-lingual) and parenteral (e.g. subcutaneous, intramuscular, intradermal or intravenous) administration.

Formulations suitable for buccal (sub-lingual) administration include lozenges comprising a compound of formula (I) or a salt or solvate thereof in a flavoured base, usually sucrose and acacia or tragacanth, and pastilles comprising the agent in an inert base such as gelatin and glycerin or sucrose and acacia.

Formulations of the present invention suitable for parenteral administration conveniently comprise sterile aqueous preparations of a compound of formula (I) or a salt or solvate thereof, preferably isotonic with the blood of the intended recipient. These preparations are preferably administered intravenously, although administration may also be effected by means of subcutaneous, intramuscular, or intradermal injection. Such preparations may conveniently be prepared by admixing the agent with water and rendering the resulting solution sterile and isotonic with the blood.

Formulations suitable for rectal administration are preferably presented as unit-dose suppositories. These may be prepared by admixing a compound of formula (I) with one or more conventional solid carriers, for example, cocoa butter, and then

shaping the resulting mixture.

Formulations suitable for topical application to the skin preferably take the form of an ointment, cream, lotion, paste, gel, spray, transdermal patch, aerosol, or oil. Carriers which may be used include vaseline, lanolin, polyethylene glycols, alcohols, and combinations of two or more thereof.

It should be understood that in addition to the ingredients particularly mentioned above, the formulations may include other agents conventional in the art having regard to the type of formulation in question.

Formulation Examples

The following non-limiting examples illustrate suitable formulations for use in the present invention, particularly for once-daily administration.

Example 1

Component	Amount/unit (mg)
Tablet Core	
Formula (I).HCl	22.86(*)
Microcrystalline Cellulose (Avicel PH102) Ph.Eur /USNF	176.20
Hydroxypropylmethylcellulose (Methocel E4M CR) Ph.Eur/USNF	120.25
Sodium Bisulphate	3.25
Purified Water (removed during processing) Ph.Eur	qs
Magnesium Stearate Ph.Eur /USNF	2.44
TOTAL	325.00
Tablet coating	
Opadry White : OY-S-28876	13.00
TOTAL	**338.00**

(*) corresponding to 20mg of the compound of formula (I)

Example 2

Component	Amount/unit (mg)
Tablet Core	
Formula (I).HCl	45.72(*)
Microcrystalline Cellulose (Avicel PH102) Ph.Eur /USNF	159.84
Hydroxypropylmethylcellulose (Methocel E4M CR) Ph.Eur/USNF	113.75
Sodium Bisulphate	3.25
Purified Water (removed during processing) Ph.Eur	qs
Magnesium Stearate Ph.Eur /USNF	2.44
TOTAL	325.00
Tablet coating	**13.00**
Opadry White : OY-S-28876	
TOTAL	**338.00**

(*) corresponding to 40mg of the compound of formula (I)

Example 3

Component	Amount/unit (mg)
Tablet Core	
Formula (I).HCl	68.58(*)
Microcrystalline Cellulose (Avicel PH102) Ph.Eur /USNF	153.23
Hydroxypropylmethylcellulose (Methocel E4M CR) Ph.Eur/USNF	97.50
Sodium Bisulphate	3.25
Purified Water (removed during processing) Ph.Eur	qs
Magnesium Stearate Ph.Eur /USNF	2.44
TOTAL	325.00
Tablet coating	
Opadry White : OY-S-28876	13.00
TOTAL	**338.00**

(*) corresponding to 60mg of the compound of formula (I)

Example 4

Component	Amount/unit (mg)
Tablet Core	
Formula (I).HCl	91.44(*)
Microcrystalline Cellulose (Avicel PH102) Ph.Eur /USNF	130.37
Hydroxypropylmethylcellulose (Methocel E4M CR) Ph.Eur/USNF	97.50
Sodium Bisulphate	3.25
Purified Water (removed during processing) Ph.Eur	qs
Magnesium Stearate Ph.Eur /USNF	2.44
TOTAL	325.00
Tablet coating	
Opadry White : OY-S-28876	13.00
TOTAL	**338.00**

(*) corresponding to 80mg of the compound of formula (I)

Example 5

Component	Amount/unit (mg)
Tablet Core	
Formula (I).HCl	11.43(*)
Microcrystalline Cellulose (Avicel PH102) Ph.Eur /USNF	160.87
Hydroxypropylmethylcellulose (Methocel E4M CR) Ph.Eur/USNF	146.25
Sodium Bisulphate	3.25
Purified Water (removed during processing) Ph.Eur	qs
Magnesium Stearate Ph.Eur /USNF	3.25
TOTAL	325.05
Tablet coating	
Opadry White : YS-1R-7003	9.75
TOTAL	**334.80**

(*) corresponding to 10mg of the compound of formula (I)

Examples 1 to 5 above were prepared by a process similar to the following general process: The drug substance is blended and wet granulated with the pharmaceutically acceptable excipients described, including HPMC as the rate-controlling polymer. The acidic stabiliser (sodium bisulphate) is first dissolved in purified water to produce the granulation solution, and the granule is then produced by conventional processing techniques, for example either high shear or a fluid bed process, followed by drying, milling, blending, compression into a tablet, and finally aqueous film-coating.

Biological Data

In vitro Synaptosomal Uptake

In vitro uptake was determined, as reported previously, using synaptosomes prepared from rat caudoputamen (for dopamine uptake) and hypothalamus (for NA and serotonin uptake) using [^{3}H]-dopamine, [^{3}H]-NA and [^{3}H]-serotonin as transport substrates, respectively. *See* Eckhardt, S.B., R.A. Maxwell, and R.M. Ferris, A Structure-Activity Study of the Transport Sites for the Hypothalamic and Striatal Catecholamine Uptake Systems. Similarities and differences. *Molecular Pharmacology*, 21: p. 374-9,1982.

Synaptosomes for use in obtaining in vitro uptake data were prepared from hypothalamus or striatum by gently homogenizing the tissue in a 0.3 M sucrose/25 mM Tris pH 7.4 buffer containing iproniazid phosphate to inhibit monoamine oxidase. The homogenate was centrifuged at 1100 x g at 4°C for 10 min and the supernatant was used for uptake studies. The supernatant (~ 1 mg tissue protein) was incubated with Km concentrations of [^{3}H]-noradrenaline, [^{3}H]-dopamine or [^{3}H]-serotonin at 37°C for 5 minutes in Modified Krebs-Henseleit buffer (118 mM NaCl, 5 mM KCl, 25 mM $NaHCO_3$, 1.2 mM NaH_2PO_4, 1.2 mM $MgSO_4$, 11 mM Dextrose, 2.5 mM $CaCl_2$) in the absence and presence of drug. Under these conditions uptake was linear with respect to both for substrate and tissue (with <5% total substrate transported). Non-specific uptake was defined as uptake at 0°C. [^{3}H]-substrate, which had been transported into synaptosomes, was separated from free [^{3}H]-substrate by filtration over GF/B filters and washing with cold Krebs-Henseleit buffer. The filters were counted for tritum in a liquid scintillation spectrometer.

The data for *in vitro* synaptosomal uptake are presented below as Table 1. The compound of formula (I), inhibited noradrenaline (NA) uptake with an IC_{50} of 1.1 μM. On dopamine (DA) uptake, the compound of formula (I) had an IC_{50} of ~10 μM. The compound of formula (I) showed no inhibition of serotonin uptake at 30 μM.

Table 1

Compound	IC_{50} NA	IC_{50} DA	IC_{50} Serotonin
Formula (I)	1.1 ± 0.07	9.3 ± 0.41	>30

Uptake values are means ± SEM of 3 separate experiments. The IC_{50} values are concentrations (μM) required for 50% inhibition of uptake.

Functional reuptake inhibition on human monoamine transporters

Three separate cell-lines expressing human monoamine transporters for dopamine (hDAT) noradrenaline (hNET) and serotonin (hSERT) were used to measure the functional reuptake inhibiting properties of the compound of formula (I) (as its hydrochloride salt). The following methods were utilised.

Human noradrenaline transporter (hNET): MDCK/hNET (dog kidney) cells (4

x 10^4 cells/well) expressing the human norepinephrine transporter were plated on 96-well format one day before the assay. When the cells were 80% confluent, cell monolayers were washed and preincubated with test compound and/or vehicle in modified Tris-HEPES buffer pH 7.1 at 25°C for 20 minutes, then 25 nM [^{3}H]Norepinephrine was added to make the total volume to 200 µl and the cells were further incubated for 10 minutes. Cells in the well were then rinsed twice, solubilized with 1% SDS lysis buffer and the lysate was counted to determine [^{3}H]Norepinephrine uptake. Non-specific signal was determined in the presence of 10 µM desipramine. Reduction of [^{3}H]Norepinephrine uptake by 50 per cent or more (≥50%) relative to vehicle controls indicated significant inhibitory activity.

Human dopamine transporter (hDAT): CHO-K1/hDAT cells (8 x 10^4 cells/well) expressing the human dopamine transporter (hDAT) were plated on 96-well format one day before the assay. Cells were preincubated with test compound and/or vehicle in modified Tris-HEPES buffer pH 7.1 at 25°C for 20 minutes, then 50 nM [^{3}H]Dopamine was added to make the total volume to 200 µl and further incubated for 10 minutes. Cells in the well were then rinsed twice, solubilized with 1% SDS lysis buffer and the lysate was counted to determine [^{3}H]Dopamine uptake. Non-specific signal was determined in the presence of 10µM nomifensine. Reduction of [^{3}H]Dopamine uptake by 50 per cent or more (≥50%) relative to vehicle controls indicates significant inhibitory activity.

Human serotonin transporter (hSERT): HEK-293/hSERT cells (5 x 10^4 cells/tube) expressing the human serotonin transporter (hSERT) were added into the minitube on 96-tube holder prior to assay. Cells were preincubated with test compound or vehicle in modified Tris-HEPES buffer pH 7.1 at 25°C for 20 minutes, then 65 nM [^{3}H]Serotonin was added to make the total volume to 200 µl and further incubated for 10 minutes. Cells were then washed by filtration through cell harvester four times with PBS buffer containing 0.1% BSA and the GF/B filter was counted to determine [^{3}H]Serotonin uptake. Non-specific signal was determined in the presence of 10µM fluoxetine. Reduction of [^{3}H]Serotonin uptake by 50 percent or more (≥50%) relative to vehicle-control indicates significant inhibitory activity.

Compounds were screened at 10, 1, 0.1, 0.01 and 0.001 µM. These same concentrations were concurrently applied to a separate group of untreated cells and evaluated for possible compound-induced cytotoxicity only if significant inhibition of uptake was observed. Radioactivity retained on the filters was determined by scintillation counting overnight using a Packard scintillation counter.

The potencies for monoamine reuptake inhibition for the hydrochloride salt of the compound of formula (I) are expressed in Table 2 below as IC_{50} (in µM; mean ± SEM) following three separate experiments, each performed in duplicate (n=3). The compound demonstrated reuptake inhibition at both hDAT (pIC50=6.36) and hNET (pIC50=6.70) but reuptake inhibition was not observed on hSERT (pIC50 < 5) at the highest concentration tested (10µM). No cytotoxicity was observed at any of the concentrations causing reuptake inhibition.

Table 2

Compound	hNET	hDAT	hSERT
Formula (I).HCl	0.20 ± 0.05 (n=3)	0.44 ± 0.01 (n=3)	>10 (n=3)

Dopamine transporter (DAT) receptor occupancy study

The study was conducted in two parts. Part A utilized positron emission tomography (PET) to characterize the concentration-percent occupancy relationship with respect to the dopamine transporter (DAT) after dosing with intravenous compound of formula (I) to pseudo-steady-state. Part B utilized PET to evaluate the time course of occupancy at the dopamine transporter following steady-state dosing of the compound of formula (I) via a modified release oral formulation. The tracer used in this study to assess the occupancy at the DAT was ^{11}C-βCIT-FE.

Part A. Six healthy male volunteers each received a 20.6mg intravenous dose of the compound of formula (I) administered via a 2-hour loading infusion followed by a 2-hour maintenance infusion to achieve pseudo-steady-state. Each subject then received two additional doses (61.8mg and 91.2mg) of the compound of formula (I), with each dose separated by 3 weeks. A baseline PET scan was performed between 1 and 7 days prior to first dosing with formula (I), and then each post-dose PET scan performed 2.5 hours after the initiation of each of the three dosing regimens.

The concentrate solution for infusion was formulated to contain 10mg free base equivalent per mL as the hydrochloride salt of the compound of formula (I), and consisted of a clear, colorless solution, pH adjusted to approximately 4.5, buffered with 50mmol citrate buffer. The product was diluted to a total volume of 250mL with 0.9% weight-to-volume ratio (w/v) sodium chloride injection prior to being administered.

The study showed that the compound of formula (I) bound effectively to dopamine transporter in the striatum. A clear concentration-dependent transporter occupancy was observed, with receptor occupancy values of 14±0.38%, 37±8.0% and 47±6.8% being observed for the 20.6mg, 61.8mg, and 91.2mg doses respectively.

Part B. Six healthy male volunteers received 60mg of the compound of formula (I) via a modified release oral formulation once daily for 6 days in order to achieve steady-state by the last day. Four ^{11}C-βCIT-FE PET scans were performed for each subject, the first between 1 and 7 days prior to dosing and the remaining three at 6, 12 and 24 hours following the final dose on Day 6. The oral formulations were uncoated tablets, containing 20mg and 40mg of the compound of formula (I) as the hydrochloride salt; details of the formulations are set out below as Examples 6 and 7 (prepared by an analogous method to Examples 1 to 5 above).

This oral dosing of 60mg of the compound of formula (I) resulted in an average receptor occupancy of 29% (29±5.3%) at 6 hours after the last dose of the drug. Further PET investigations showed that the degree of receptor occupancy remained at almost the same level, on average 27% (27±11.3%), during the following 6 hours. The results from Part B of the study were in accordance with those from Part A.

Example	Components	Amount/unit (mg)
6	Formula (I).HCl Microcrystalline Cellulose (Avicel PH-102) EP/USP Hydroxypropyl Methylcellulose (Methocel E4M CR) EP/USP Sulfuric Acid EP/USP Purified Water (removed during processing) EP/USP Lactose monohydrate spray dried EP/USP Magnesium stearate EP/USP **TOTAL**	45.72(*) 70.30 81.26 2.72 qs 121.75 3.25 **325.00**
7	Formula (I).HCl Microcrystalline Cellulose (Avicel PH-102) EP/USP Hydroxypropyl Methylcellulose (Methocel E4M CR) EP/USP Sulfuric Acid EP/USP Purified Water (removed during processing) EP/USP Lactose monohydrate spray dried EP/USP Magnesium stearate EP/USP **TOTAL**	22.86(§) 93.16 81.26 2.72 qs 121.75 3.25 **325.00**

(*) corresponding to 40.00mg of the compound of formula (I)

(§) corresponding to 20.00mg of the compound of formula (I)

The biological data presented above demonstrates that the compound of formula (I) inhibits the dopamine transporter in human subjects. Thus the compound of formula (I) may, as a result of enhancing dopaminergic activity in patients suffering from RLS or from PLMD, be an effective treatment for RLS or for PLMD.

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the present invention, and all such modifications as would be obvious to one skilled in the art are intended to be included within the scope of the following claims.

CLAIMS

1. Use of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof in the manufacture of a medicament for the treatment of Restless Legs Syndrome.

2. Use according to claim 1 wherein said (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof is enantiomerically pure.

3. Use according to claim 1 or claim 2 wherein the salt is (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride.

4. Method of treating Restless Legs Syndrome (RLS) in a mammal comprising the administration to said subject of an effective amount of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof.

5. Method according to claim 4 wherein said (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof is enantiomerically pure.

6. Method according to claim 4 or claim 5 wherein the salt is (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride. 1.

7. Use of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof in the manufacture of a medicament for the treatment of Periodic Limb Movement Disorder.

8. Use according to claim 7 wherein said (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof is enantiomerically pure.

9. Use according to claim 7 or claim 8 wherein the salt is (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride.

10. Method of treating Periodic Limb Movement Disorder (PLMD) in a mammal comprising the administration to said subject of an effective amount of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof or pharmaceutical compositions thereof.

11. Method according to claim 10 wherein said (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a salt or solvate thereof is enantiomerically pure.

12. Method according to claim 10 or claim 11 wherein the salt is (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/013248

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K31/5375 A61P25/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, CHEM ABS Data, EMBASE, MEDLINE, SCISEARCH, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2003/064988 A1 (MORGAN PHILLIP FREDERICK ET AL) 3 April 2003 (2003-04-03) paragraph '0006! - paragraph '0010! paragraph '0023!	
A	US 6 337 328 B1 (FANG QUN KEVIN ET AL) 8 January 2002 (2002-01-08) column 4, line 40 - line 63	
A	WO 01/62257 A (SEPRACOR INC) 30 August 2001 (2001-08-30) page 4, line 29 - page 5, line 1 page 7, line 26 - line 31	

-/--

[X] Further documents are listed in the continuation of box C.

[X] Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
13 April 2005	19/04/2005

Name and mailing address of the ISA	Authorized officer
European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Bonzano, C

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99/37305 A (GLAXO GROUP LIMITED; MORGAN, PHILLIP, FREDERICK; MUSSO, DAVID, LEE; PA) 29 July 1999 (1999-07-29) cited in the application abstract page 2, line 5 - line 14	
Y	KELLEY JAMES L ET AL: "-2-(3,5-difluorophenyl)-3,5-dimethyl-2-morpholinol: A novel antidepressant agent and selective inhibitor of norepinephrine uptake" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 39, no. 2, 1996, pages 347-349, XP002164358 ISSN: 0022-2623 page 347, column 2, paragraph 3	1-12
Y	EP 1 336 406 A (SOLVAY PHARM BV 'NL!) 20 August 2003 (2003-08-20) paragraph '0013! paragraph '0001!	1-12

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2004/013248

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. [X] Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 10-12 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

2. [ ] Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. [ ] Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. [ ] As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. [ ] As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. [ ] As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. [ ] No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

[ ] The additional search fees were accompanied by the applicant's protest.

[ ] No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP2004/013248

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2003064988 A1	03-04-2003	US 2002019396 A1	14-02-2002
		US 6274579 B1	14-08-2001
		US 2003027827 A1	06-02-2003
		US 2003083330 A1	01-05-2003
		US 2003032643 A1	13-02-2003
US 6337328 B1	08-01-2002	US 6342496 B1	29-01-2002
		AU 6926800 A	03-09-2001
		CA 2400482 A1	30-08-2001
		CZ 20022857 A3	14-01-2004
		EP 1259243 A2	27-11-2002
		HU 0300030 A2	28-05-2003
		JP 2003529563 T	07-10-2003
		PL 357389 A1	26-07-2004
		WO 0162257 A2	30-08-2001
		US 2002052341 A1	02-05-2002
		CA 2362361 A1	08-09-2000
		EP 1165086 A2	02-01-2002
		JP 2004513061 T	30-04-2004
		US 2002052340 A1	02-05-2002
		AU 775642 B2	12-08-2004
		AU 3505500 A	21-09-2000
		WO 0051546 A2	08-09-2000
WO 0162257 A 30-08-2001		US 6342496 B1	29-01-2002
		US 6337328 B1	08-01-2002
		AU 6926800 A	03-09-2001
		CA 2362361 A1	08-09-2000
		CA 2400482 A1	30-08-2001
		CZ 20022857 A3	14-01-2004
		EP 1165086 A2	02-01-2002
		EP 1259243 A2	27-11-2002
		HU 0300030 A2	28-05-2003
		JP 2004513061 T	30-04-2004
		JP 2003529563 T	07-10-2003
		PL 357389 A1	26-07-2004
		WO 0162257 A2	30-08-2001
		US 2002052340 A1	02-05-2002
		US 2002052341 A1	02-05-2002
WO 9937305 A 29-07-1999		AP 1229 A	04-12-2003
		AU 755536 B2	12-12-2002
		AU 2328099 A	09-08-1999
		BR 9907203 A	17-10-2000
		CA 2318268 A1	29-07-1999
		CN 1294513 A	09-05-2001
		CN 1528753 A	15-09-2004
		EA 2410 B1	25-04-2002
		EE 200000438 A	17-12-2001
		EP 1047428 A1	02-11-2000
		HR 20000494 A1	31-12-2000
		HU 0100900 A2	29-05-2002
		ID 26334 A	14-12-2000
		JP 2002501025 T	15-01-2002
		NO 20003721 A	19-09-2000
		NZ 505809 A	27-09-2002
		NZ 520349 A	27-02-2004
		NZ 529316 A	28-05-2004

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP2004/013248

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO 9937305 A		PL	342012 A1	07-05-2001
		SK	10912000 A3	09-04-2001
		TR	200002126 T2	21-12-2000
		WO	9937305 A1	29-07-1999
		US	6274579 B1	14-08-2001
		US	2002019396 A1	14-02-2002
EP 1336406 A	20-08-2003	EP	1336406 A1	20-08-2003
		AU	2003208714 A1	04-09-2003
		BR	0306147 A	19-10-2004
		CA	2462688 A1	21-08-2003
		WO	03068207 A2	21-08-2003
		EP	1476148 A2	17-11-2004
		US	2005038015 A1	17-02-2005