



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12Q 1/70	A1	(11) International Publication Number: WO 93/06247 (43) International Publication Date: 1 April 1993 (01.04.93)
(21) International Application Number: PCT/US92/07813 (22) International Filing Date: 16 September 1992 (16.09.92) (30) Priority data: 760,292 16 September 1991 (16.09.91) US (71) Applicant: ABBOTT LABORATORIES [US/US]; Chad-0377/AP6D-2, One Abbott Park Road, Abbott Park, IL 60064-3500 (US). (72) Inventors: LESNIEWSKI, Richard, R. ; 8706 110th Avenue, Kenosha, WI 53142 (US). LEUNG, Tat, K. ; 2825 Grandville Court, Waukegan, IL 60085 (US). (74) Agents: GORMAN, Edward, Hoover, Jr. et al.; Abbott Laboratories, CHAD-0377, AP6D/2, One Abbott Park Road, Abbott Park, IL 60064-3500 (US).		(81) Designated States: AU, CA, JP, KR, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE). Published <i>With international search report.</i>
(54) Title: HEPATITIS C ASSAY (57) Abstract The present invention provides an improved assay for detecting the presence of an antibody to an HCV antigen in a sample by contacting the sample with at least one polypeptide containing at least one epitope of an HCV antigen. Preferred assay formats include a confirmatory assay, a combination assay, a synthetic polypeptide-based assay, an immunodot assay and a competition assay.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MN	Mongolia
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Faso	GN	Guinea	NO	Norway
BG	Bulgaria	GR	Greece	NZ	New Zealand
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	PT	Portugal
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CS	Czechoslovakia	LU	Luxembourg	SU	Soviet Union
CZ	Czech Republic	MC	Monaco	TD	Chad
DE	Germany	MG	Madagascar	TG	Togo
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain			US	United States of America

HEPATITIS C ASSAY

Background of the Invention

5 This invention relates generally to an assay for identifying the presence and/or amount in a test sample of an antibody which is immunologically reactive with a hepatitis C viral antigen and more specifically, to an assay for detecting a complex of an antibody and a polypeptide having at least one epitope of a hepatitis C viral antigen.

10

 Acute viral hepatitis is clinically diagnosed by a well-defined set of patient symptoms, including jaundice, hepatic tenderness, and an increase in the serum levels of alanine aminotransferase and aspartate aminotransferase. Additional serological immunoassays are generally performed to diagnose the specific type of viral causative agent. Historically, patients presenting clinical hepatitis symptoms and not otherwise infected by hepatitis A, hepatitis B, Epstein-Barr, or cytomegalovirus were clinically diagnosed as having non-A, non-B hepatitis (NANBH) by default. The disease may result in chronic liver damage.

20

 Each of the well-known, immunologically characterized hepatitis-inducing viruses, hepatitis A virus (HAV), hepatitis B virus (HBV), and hepatitis D virus (HDV) belongs to a separate family of viruses and has a distinctive viral organization, protein structure and mode of replication.

25

 Attempts to identify the NANBH virus by virtue of genomic similarity to one of the known hepatitis viruses have failed, suggesting that NANBH has a distinct organization and structure. Fowler et al., J. Med. Virol. 12:205-213 (1983) and Weiner et al., J. Med. Virol. 21:239-247 (1987).

30

 Progress in developing assays to detect antibodies specific for NANBH has been particularly hampered by difficulties in correctly identifying antigens associated with NANBH. See, for example, J. Wands et al., U.S. Patent 4,870,076, Wands et al., Proc. Nat'l. Acad. Sci. 83:6608-6612 (1986), Ohori et al., J. Med. Virol. 12:161-178 (1983), Bradley et al., Proc. Nat'l. Acad. Sci. 84:6277-6281 (1987), T. Akatsuka et al., J. Med.

35

Virology 20:43-56 (1986), B. Seto et al., U.S. Patent Application No. 07/234,641 (available from U.S. Department of Commerce National Technical Information Service, Springfield, Virginia No. 89138168), K. Takahashi et al., European Patent Application No. 0 293 274, published
5 November 30, 1988, and R. Seelig et al., PCT Application PCT/EP88/00123.

Recently, another hepatitis-inducing virus has been unequivocally identified as hepatitis C virus (HCV) by M. Houghton et al., European Patent
10 Application publication number 0 318 216, May 31, 1989. Related papers describing this virus include G. Kuo et al., Science 244:359-361 (1989) and Q. Choo et al., Science 244:362-364 (1989). M. Houghton et al. reported isolating cDNA sequences from HCV which encode antigens which react immunologically with antibodies present in patients infected with NANBH,
15 thus establishing that HCV is the viral agent causing NANBH.

The cDNA sequences associated with HCV were isolated from a cDNA library prepared from the RNA obtained from pooled serum from a chimpanzee with chronic HCV infection. The cDNA library contained cDNA
20 sequences of approximate mean size of about 200 base pairs. The cDNA library was screened for encoded epitopes expressed in clones that could bind to antibodies in sera from patients who had previously experienced NANBH.

25 In the European Patent Application, M. Houghton et al. also described the preparation of several superoxide dismutase fusion polypeptides (SOD) and the use of these SOD fusion polypeptides to develop an HCV screening assay. The most complex SOD fusion polypeptide described in the European Patent Application, designated
30 C100-3, was described as containing 154 amino acids of human SOD at the amino-terminus, 5 amino acid residues derived from the expression of a synthetic DNA adapter containing a restriction site, EcoRI, 363 amino acids derived from the expression of a cloned HCV cDNA fragment, and 5
35 carboxy terminal amino acids derived from an MS2 cloning vector nucleotide sequence. The DNA sequence encoding this polypeptide was transformed into yeast cells using a plasmid. The transformed cells were

cultured and expressed a 54,000 molecular weight polypeptide which was purified to about 80% purity by differential extraction.

Other SOD fusion polypeptides designated SOD-NANB5-1-1 and
5 SOD-NANBH81 were expressed in recombinant bacteria. The E. coli
fusion polypeptides were purified by differential extraction and by
chromatography using anion and cation exchange columns. The
purification procedures were able to produce SOD-NANB5-1-1 as about
80% pure and NANBH81 as about 50% pure.

10

The recombinant SOD fusion polypeptides described by M.
Houghton et al. were coated on microtiter wells or polystyrene beads and
used to assay serum samples. Briefly, coated microtiter wells were
incubated with a sample in a diluent. After incubation, the microtiter wells
15 were washed and then developed using either a radioactively labelled
sheep anti-human antibody or a mouse anti-human IgG-HRP (horseradish
peroxidase) conjugate. These assays were used to detect both post acute
phase and chronic phase of HCV infection. Due to the preparative
methods, assay specificity required adding yeast or E. coli extracts to the
20 samples in order to prevent undesired immunological reactions with any
yeast or E. coli antibodies present in serum samples.

Ortho Diagnostics Systems Inc. have developed a research
immunoenzyme assay to detect antibodies to HCV antigens. The Ortho
25 assay procedure is a three-stage test for serum/plasma carried out in a
microwell coated with the recombinant yeast/hepatitis C virus SOD fusion
polypeptide C100-3.

In the first stage, a test specimen is diluted directly in the test well
30 and incubated for a specified length of time. If antibodies to HCV antigens
are present in the specimen, antigen-antibody complexes will be formed on
the microwell surface. If no antibodies are present, complexes will not be
formed, and the unbound serum or plasma proteins will be removed in a
washing step.

35

In the second stage, anti-human IgG murine monoclonal antibody
horseradish peroxidase conjugate is added to the microwell. The

conjugate binds specifically to the antibody portion of the antigen-antibody complexes. If antigen-antibody complexes are not present, the unbound conjugate will also be removed by a washing step.

5 In the third stage, an enzyme detection system composed of o-phenylenediamine 2 HCl (OPD) and hydrogen peroxide is added to the test well. If bound conjugate is present, the OPD will be oxidized, resulting in a colored end product. After formation of the colored end product, dilute sulfuric acid is added to the microwell to stop the color-forming detection
10 reaction.

The intensity of the colored end product is measured with a microwell reader. The assay may be used to screen patient serum and plasma.

15 It is established that HCV may be transmitted by contaminated blood and blood products. In transfusion patients, as many as 10% will suffer from post-transfusion hepatitis. Of those, approximately 90% are the result of infections diagnosed as HCV. The prevention of transmission of HCV by
20 blood and blood products requires reliable, sensitive and specific diagnosis and prognostic tools to identify HCV carriers as well as contaminated blood and blood products. Thus, there exists a need for an HCV assay which uses reliable and efficient reagents and methods to accurately detect the presence of HCV antibodies in samples.

25

Summary of the Invention

The present invention provides an improved assay for detecting the presence of an antibody to an HCV antigen in a test sample by contacting
30 the sample with a polypeptide containing at least one epitope of an HCV antigen, wherein the polypeptides are selected from the group consisting of p380-JH1, p-380.LG, p380-J and p408.

35 One assay format according to the invention provides a confirmatory assay for unequivocally identifying the presence of an antibody that is immunologically reactive with an HCV antigen. Briefly, a fluid sample is used to prepare first and second aliquots. The aliquots then are contacted

with at least two polypeptides duplicative of a continuous amino acid sequence putatively contained in proteins expressed by clones containing HCV cDNA sequences containing at least one epitope of an HCV antigen under conditions suitable for complexing the antibody with the polypeptide.

- 5 Finally, the antibody-antigen complex is detected. The first aliquot and the second aliquot are contacted with at least one polypeptide selected from the group consisting of p380-JH1, p-380.LG, p380-J and p408J, with the proviso that the polypeptide(s) used for the first aliquot are not used for the second aliquot..

10

- Another assay format provides an assay for identifying the presence of an antibody that is immunologically reactive with an HCV antigen in a fluid sample comprising contacting the sample with a solid phase having bound thereto at least one polypeptide selected from the group consisting of p380JH1, p380J, p380.LG and p408J containing at least one epitope of an HCV antigen, under conditions suitable for complexing the antibody with the polypeptide and detecting the antibody-polypeptide complex.

15

- Another assay format provides an assay for identifying the presence of an antibody that is immunologically reactive with an HCV antigen in a fluid sample comprising contacting the sample with a polypeptide selected from the group consisting of p380JH1, p380J, p380.LG and p408J containing at least one epitope of an HCV antigen under conditions suitable for complexing the antibody with the polypeptide and detecting the antibody-polypeptide complex as an indication of chronic HCV infection.

25

- Another assay format provides an immunodot assay for identifying the presence of an antibody that is immunologically reactive with an HCV antigen by concurrently contacting a sample with at least two polypeptides selected from the group consisting of p380JH1, p380J, p380.LG and p408J each containing distinct epitopes of an HCV antigen under conditions suitable for complexing the antibody with at least one of the polypeptides and detecting the antibody-polypeptide complex by reacting the complex with color-producing reagents.

30

- Another assay format provides a competition assay directed to the confirmation that positive results are not false by identifying the presence of

35

an antibody that is immunologically reactive with an HCV antigen in a fluid sample where the sample is used to prepare first and second immunologically equivalent aliquots. The first aliquot is contacted with a solid support containing a bound polypeptide selected from the group consisting of p380JH1, p380J, p380.LG and p408J which contains at least one epitope of an HCV antigen under conditions suitable for complexing with the antibody to form a detectable antibody-polypeptide complex and the second aliquot is first contacted with unbound polypeptide selected from the group consisting of p380JH1, p380J, p380.LG and p408J and then contacted with the same solid support containing bound polypeptide selected from the group consisting of p380JH1, p380J, p380.LG and p408J.

In all of the assays, the sample, especially serum or plasma, preferably is diluted before contacting the polypeptide absorbed on a solid support. However, it is contemplated that when testing some test samples, an undiluted or even concentrated sample may be preferred. Samples may be obtained from different biological samples such as whole blood, serum, plasma, cerebrospinal fluid, urine, and lymphocytes or cell culture supernatants. Solid support materials may include cellulose materials, such as paper and nitrocellulose, natural and synthetic polymeric materials, such as polyacrylamide, polystyrene, and cotton, silicon chips, porous gels such as silica gel, agarose, dextran and gelatin, particles which are capable of forming a charge such as those used for ion capture assays and inorganic materials such as deactivated alumina, magnesium sulfate and glass. Suitable solid support materials may be used in assays in a variety of well known physical configurations, including microtiter wells, test tubes, beads, strips, membranes, and microparticles, either magnetic or non-magnetic. A preferred solid support for a non-immunodot assay is a polystyrene bead. A preferred solid support for an immunodot blot assay is nitrocellulose.

Suitable methods and reagents for detecting an antibody-antigen complex in an assay of the present invention are commercially available or known in the relevant art. Representative methods may employ detection reagents such as an enzymatic, radioisotopic, fluorescent, luminescent or chemiluminescent reagents. These reagents may be used to prepare hapten-labelled anti-hapten detection systems according to known

procedures, for example, a biotin-labelled anti-biotin system may be used to detect an antibody-antigen complex.

5 The present invention also encompasses assay kits including polypeptides which contain at least one epitope of an HCV antigen bound to a solid support as well as needed sample preparation reagents, wash reagents, detection reagents and signal producing reagents.

10 Other aspects and advantages of the invention will be apparent to those skilled in the art upon consideration of the following detailed description which provides illustrations of the invention in its presently preferred embodiments.

Brief Description of the Drawings

15

FIGURES 1a and 1b illustrate the HCV genome.

FIGURE 2 illustrates the use of antigenic polypeptides to identify the presence of antibodies in a chimpanzee inoculated with HCV.

20 FIGURES 3a and 3b illustrate the sensitivity increase using a combination assay format.

FIGURE 4 illustrates a test cartridge for an immunodot assay.

25 FIGURE 5 illustrates a seroconversion graph wherein the amount of anti-NS5 S/N antibody, shown as the solid line between closed circles, and the amount of anti-HCV 2.0 S/CO antibody shown as a solid line between open squares, is plotted against days post presentation.

Detailed Description of the Invention

30 The present invention is directed to an assay to detect an antibody to an HCV antigen in a test sample. In a preferred format, human serum or plasma is diluted in a sample diluent and incubated with a polystyrene bead coated with a polypeptide that includes an HCV antigenic epitope. If antibodies are present in the sample, they will form a complex with the antigenic polypeptide and become affixed to the polystyrene bead. After
35 the complex has formed, unbound materials and reagents are removed by washing the bead and the bead-antigen-antibody complex is reacted with a solution containing horseradish peroxidase labelled goat antibodies

directed against human antibodies. This peroxidase enzyme then binds to the antigen-antibody complex already fixed to the bead. In a final reaction, the horseradish peroxidase is contacted with o-phenylenediamine and hydrogen peroxide, which results in a yellow-orange color. The intensity of the color is proportional to the amount of antibody which initially binds to the antigen fixed to the bead.

The preferred polypeptides having HCV antigenic epitopes were selected from portions which possessed amino acid sequences similar to other known immunologically reactive agents and which were identified as having some immunological reactivity. The immunological reactivity of a polypeptide was initially identified by reacting the cellular extract of *E. coli* clones which had been transformed with cDNA fragments of the HCV genome with HCV infected serum. The clones presumably expressed polypeptides encoded by the incorporated cDNA which were immunologically reactive with serum known to contain antibody to HCV antigens. An analysis of a given amino acid sequence, however, only provides rough guides to predicting immunological reactivity. There is no invariably predictable way to ensure immunological activity short of preparing a given amino acid sequence and testing the suspected sequence in an assay. As illustrated in Table 1, some peptides which were expected to provide immunological reactivity were found to be unreactive when used in an actual assay.

TABLE 1

CHIRON UPDATE	GENOMIC REGION	SAMPLE DILN.	A00642 RESULT	SAMPLE DILN.	#401 RESULT	SAMPLE DILN.	#423 RESULT
1694-1735	C-100	1:800	POS.	1:40	POS.	1:20	POS.
1866-1930	C-100	1:100	POS.	-	-	-	-
1689-1805	C-100	1:500	POS.	1:40	POS.	1:20	POS.
1684-1750	C-100	1:500	POS.	1:40	POS.	1:20	POS.
1899-1930	C-100	1:50	POS.	1:40	POS.	1:20	NEG.
1192-1240	33C	1:800	POS.	1:40	POS.	1:20	POS.
1223-1240	33C	1:200	POS.	1:40	POS.	1:20	POS.
1-75	Put. Core	1:100	POS.	1:40	POS.	1:20	POS.
35-75	Put. Core	1:50	POS.	1:40	NEG.	1:20	POS.
99-126	Put. Core	1:50	NEG.	1:40	NEG.	1:20	POS.

1569-1593	C-100	1:25	NEG.	1:40	NEG.	1:20	NEG.
1357-1407	33C	1:25	NEG.	1:40	NEG.	1:20	NEG.
1418-1457	33C	1:25	NEG.	1:40	NEG.	1:20	NEG.
195-262	Put.Core/ Envelope	1:50	NEG.	1:40	NEG.	1:20	NEG.
230-262	Put.Core/ Envelope	1:25	NEG.	1:40	NEG.	1:20	NEG.

A sample value is considered positive if the absorbance at 492 nm \geq 4 X absorbance value of the negative control (S/N \geq 4.0)

- 5 A00642 Human plasma sample convalescent from NANB (HCV) Hepatitis. Patient was clinically diagnosed with NANB and negative for HBV and HAV markers.
- 10 #491 based Human paid plasma donor positive by screening assays on C-100. No known clinical history.
- #423 based Human paid plasma donor positive by screening assays on C100-3. No known clinical history.
- 15 The use of polypeptides having one or more than one epitope of an HCV antigen to detect the presence of an antibody to an HCV antigen is illustrated in Figure 2. The course of HCV infection in the chimpanzee, Melilot, was followed with one assay using recombinant C100-3 polypeptide and with another assay using p1689 polypeptide. Both assays
- 20 gave negative results before inoculation and both assays detected the presence of antibodies about 100 days after the animal had been infected with HCV.

25 There are several known methods using both synthetic and recombinant methodologies to prepare the polypeptides of the present invention which have been found to be immunologically reactive. Preferably, the polypeptides may be prepared using automated synthesizers. The synthesis of p1684 is provided below.

30 Synthesis of p1684:

H-GRVVLSGKPAIIPDREVLVREFDEMEECSQHLPLYIEQGMM-LAEQFKQKALGLLQTASRQAEVIAPAV-OH

The fully protected peptide was assembled on a phenylacetamidomethyl (PAM) resin by stepwise solid phase synthesis (starting the with carboxyl terminal residue) according to the general procedure described by G. Barany and R. B. Merrifield in The Peptides (E. Gross and R. Meinhoeffer, eds.) 2, 1-284 (1980) Academic Press. New York, NY. The C-terminal amino acid valine (Val) was coupled to the solid support via an oxymethylphenylacetamidomethyl (OMPA) linkage to yield PAM resin which ensured improved stability to prolonged treatment with trifluoroacetic acid (TFA). A BOC-Val-OCH₂-PAM-resin (0.78 mmol/g, 0.13 g) was transferred to the reaction vessel of an Applied Biosystems Peptide Synthesizer, model 430A. All subsequent amino acids starting from the carboxyl terminal to N-terminus were coupled in a step-wise manner using Applied Biosystems' small scale rapid cycle protocol,. Protected amino acids were coupled using preformed symmetric anhydride chemistry expect of asparagine, glutamine, arginine and histidine, which were double coupled using N-N'-dicyclohexylcarbodiimide (DCC)/1-hydroxybenzotriazole (HOBt) chemistry. In the first coupling, protected amino acids were coupled using preformed symmetric anhydrides dissolved in dimethylformamide (DMF). The symmetric anhydride of an individual amino acid was formed in methylene chloride followed by solvent exchange to DMF before transferring to the reaction vessel of the peptide synthesizer. The second coupling of symmetric anhydride was also conducted in DMF. The N-amino group of all amino acids used was protected by a t-butyloxycarbonyl (t-BOC) linkage. The side chain functional groups of various amino acids were protected by the following groups:

Arg-Tos	(Tosyl)
Lys-2Clz	(2-Chlorobenzyloxycarbonyl)
Thr,Ser-Bzl	(Benzyl)
30 Tyr-2BrZ	(2-Bromobenzyloxycarbonyl)
Cys-4MeBzl	(4-Methylbenzyl)
Asp,Glu-OBzl	(O-Benzyl)
His-DNP	Dinitrophenyl

35 The fully protected peptide-resin (0.2g) was allowed to swell in methylene chloride (CH₂Cl₂) for five minutes. The peptide-resin was transferred to a manual reaction vessel, treated twice with 5% thiophenol in

DMF for 20 minutes each followed by six CH₂Cl₂ washes for one minute each, and then transferred to the reaction vessel of the synthesizer. The t-BOC protecting group then was removed using 60% TFA/CH₂Cl₂ according to the manufacturer's protocol and the partially deprotected peptide-resin then was dried overnight under house vacuum at room temperature.

Partially deprotected peptide-resin then was treated with dimethyl sulfide (DMS [1 ml], p-cresol [1 ml], p-thiocresol [0.2 g] and HF [10 ml]) at 0°C for one hour to cleave the peptide from the resin support. The HF/DMS and other volatiles were distilled off in vacuo at 0°C. The cleaved peptide and resin were washed three times with 15 ml aliquots of diethyl ether, and the cleaved peptide was extracted by washing three times each with 10 ml aliquots of 40% aqueous acetic acid and 15% aqueous acetic acid, respectively. The aqueous extracts were combined and washed three times with 15 ml aliquots of diethyl ether and then lyophilized to yield a crude peptide.

The crude peptide was analyzed for purity using reversed-phase high performance liquid chromatography on a C₄, 4.6 x 30mm column (Brownlee, Applied Biosystems, Inc., Foster City, California), flow rate one ml/minute employing 0.1% aqueous TFA (A) and 100% acetonitrile (B) as the solvent system. The preferred solvent gradient employed for this peptide analysis started with 30% B solvent. The column was maintained at 30% B for one minute followed by an increase over 20 minutes using a linear gradient to 55% B and maintained for one minute. Finally, the column was brought back to 30% B over a two minute period. The presence of peptide in the effluent was monitored simultaneously at 225 nm and 280 nm. The composition of the purified peptide was determined by acid hydrolysis. After removal of the acid, the hydrolysate was analyzed on a Beckmar 6300 amino acid analyzer.

If increased quantities of purified polypeptide were desired, semi-preparative reversed phase high performance liquid chromatography was performed in a similar manner using a C₄, 10 x 100mm column (Brownlee, Applied Biosystems, Inc., Foster City, California), using the same aqueous 0.1% TFA (A) and 100% acetonitrile (B) solvent system described above.

The preferred solvent gradient for a semi-preparative run started with 27% B at 3 ml/minute for two minutes followed by an increase over 20 minutes using a linear gradient to 50% B. The concentration was maintained at 50% B for one minute and then reduced to 27% B within one minute.

5

Other peptides described herein were assembled on solid support in a manner analogous to the synthesis described above. The amino acids tryptophan and methionine, if present, were used without any side chain protection. Usually, after incorporating methionine during the chain assembly, ethanedithiol (0.1% v/v) was added to TFA for all subsequent removal of t-BOC groups. However, if histidine protected by DNP was present in the sequence, ethanedithiol was not added to TFA; instead, indole (1% v/v) was used. Also, after incorporating tryptophan, indole (1% v/v) was added to the TFA solution.

15

HF cleavage from the resin and purification of the peptides were achieved essentially as described above.

The peptides synthesized as described above were evaluated for their antigenic/immunogenic properties. A summary of the amino acid sequences, beginning with the amino terminus and ending with the carboxy terminus, of immunologically reactive peptides is presented in Tables 2 and 8.

25

TABLE 2

(Note: H signifies the amino terminus; OH signifies the carboxyl terminus)

p1	H - M-S-T-N-P-K-P-Q-K-K-N-K-R-N-T-N-R-R-P-Q -
(1-75)	D-V-K-F-P-G-G-G-Q-I-V-G-G-V-Y-L-L-P-R-R-
	G-P-R-L-G-V-R-A-T-R-K-T-S-E-R-S-Q-P-R-G-
30	R-R-Q-P-I-P-K-A-R-R-P-E-G-R-T- OH
p35	H- Y-L-L-P-R-R-G-P-R-L-G-V-R-A-T-R-K-T-S-E-R-S-Q-P-R-G-
(35-75)	R-R-Q-P-I-P-K-A-R-R-P-E-G-R-T - OH
35	p99 H -S-P-R-G-S-R-P-S-W-G-P-T-D-P-R-R-R-S-R-N-L-G-K-V-I-D-
(99-126)	T-L-OH

TABLE 2 (CONT.)(Note: **H** signifies the amino terminus; **OH** signifies the carboxyl terminus)

5	p195	H-R-N-S-T-G-L-Y-H-V-T-N-D-C-P-N-S-S-I-V-Y-E-A-A-D-A-I-L-
	(195-262)	H-T-P-G-C-V-P-C-V-R-E-G-N-A-S-R-C-W-V-A-M-T-P-T-V-A-T-R-D-G-K-L-P-A-T-Q-L-R-R-H-I-OH
10	p230	H-V-R-E-G-N-A-S-R-C-W-V-A-M-T-P-T-V-A-T-R-D-G-K-L-P-A-
	(230-262)	T-Q-L-R-R-H-I-OH
15	P1192	H-A-V-D-F-I-P-V-E-N-L-E-T-T-M-R-S-P-V-F-T-D-N-S-S-P-P-V-
	(1192-1240)	V-P-Q-S-F-Q-V-A-H-L-H-A-P-T-G-S-G-K-S-T-K-V-OH
20	p1223	H-F-Q-V-A-H-L-H-A-P-T-G-S-G-K-S-T-K-V-OH
	(1223-1240)	
25	p1357	H-Y-V-P-H-P-N-I-E-E-V-A-L-S-T-T-G-E-I-P-F-Y-G-K-A-I-P-L-E-
	(1357-1407)	V-I-K-G-G-R-H-L-I-F-C-H-S-K-K-K-C-D-E--L-A-A-K-L-OH
30	p1418	H-R-G-L-D-V-S-V-I-P-T-S-G-D-V-V-V-V-A-T-D-A-L-M-T-G-Y-T-
	(1418-1457)	G-D-F-D-S-V-I-D-C-N-T-C-OH
35	p1569	H-D-A-H-F-L-S-Q-T-K-Q-S-G-E-N-L-P-Y-L-V-A-Y-Q-A-T-V-OH
	(1569-1593)	
40	p1684	H-G-R-V-V-L-S-G-K-P-A-I-I-P-D-R-E-V-L-Y-R-E-F-D-E-M-E-E-
	(1684-1750)	C-S-Q-H-L-P-Y-I-E-Q-G-M-M-L-A-E-Q-F-K-Q-K-A-L-G-L-L-Q-T-A-S-R-Q-A-E-V-I-A-P-A-V-OH
45	P1689	H-S-G-K-P-A-I-I-P-D-R-E-V-L-Y-R-E-F-D-E-M-E-E-C-S-Q-H-L-
	(1689-1805)	P-Y-I-E-Q-G-M-M-L-A-E-Q-F-K-Q-K-A-L-G-L-L-Q-T-A-S-R-Q-A-E-V-I-A-P-A-V-Q-T-N-W-Q-K-L-E-T-F-W-A-K-H-M-W-N-F-I-S-G-I-Q-Y-L-A-G-L-S-T-L-P-G-N-P-A-I-A-S-L-M-A-F-T-A-A-V-T-S-P-L-T-T-S-Q-OH
50	p1694	H-I-I-P-D-R-E-V-L-Y-R-E-F-D-E-M-E-E-C-S-Q-H-L-P-Y-I-E-Q-
	(1694-1735)	G-M-M-L-A-E-Q-F-K-Q-K-A-L-G-L-OH

TABLE 2 (CONT.)

(Note: **H** signifies the amino terminus; **OH** signifies the carboxyl terminus)

5 P1866 H-F-K-I-M-S-G-E-V-P-S-T-E-D-L-V-N-L-L-P-A-I-L-S-P-G-A-L-
(1866-1930) V-V-G-V-V-C-A-A-I-L-R-R-H-V-G-P-G-E-G-A-V-Q-W-M-N-R-L-I-
A-F-A-S-R-G-N-H-V-S-**OH**

10 P1899 H-A-A-I-L-R-R-H-V-G-P-G-E-G-A-V-Q-W-M-N-R-L-I-A-F-A-S-R-
(1899-1930) G-N-H-V-S-**OH**

TABLE 8

(Note: **H** signifies the amino terminus; **OH** signifies the carboxyl terminus.
The two underlined Tyr residues are not part of the HCV sequence but are
engineered there for ease of iodinating the peptide at a later time)

15 p380 H-Gly-Val-Asp-Ala-Glu-Thr-His-Val-Thr-Gly-Gly-Ser-Ala-Gly-
(380-436) His-Thr-Val-Ser-Gly-Phe-Val-Ser-Leu-Leu-Ala-Pro-Gly-Ala-
Lys-Gln-Asn-Val-Gln-Leu-Ile-Asn-Thr-Asn-Gly-Ser-Trp-His-
Leu-Asn-Ser-Thr-Ala-Leu-Asn-Cys-Asn-Asp-Ser-Seu-Asn-
Thr-Gly-**OH**

20 p380.LG H-Gly-Val-Asp-Ala--Ala-Thr-Tyr-Thr-Thr-Gly-Gly-Ser-Val-Ala-
(380-436) Arg-Thr-Thr-His-Gly-Phe-Ser-Ser-Leu-Phe-Ser-Gln-Gly-Ala-
Lys-Gln-Asn-Ile-Gln-Leu-Ile-Asn-Thr-Asn-Gly-Ser-Trp-His-Ile-
Asn-Arg-Thr-Ala-Leu-Asn-Cys-Asn-Ala-Ser-Leu-Asp-Thr-Gly-
25 **OH**

p447 H-Phe-Asn-Ser-Ser-Gly-Cys-Pro-Glu-Arg-Leu-Ala-Ser-Cys-
(447-483) Arg-Pro-Leu-Thr-Asp-Phe-Asp-Gln-Gly-Trp-Gly-Pro-Ile-Ser-
Tyr-Ala-Asn-Gly-Ser-Gly-Pro-Asp-Gln-Arg-**OH**

30 p607 H-Cys-Leu-Val-Asp-Tyr-Pro-Tyr-Arg-Leu-Trp-His-Tyr-Pro-Cys-
(607-627) Thr-Ile-Asn-Tyr-Thr-Ile-Phe-**OH**

35 p643a H-Ala-Cys-Asn-Trp-Thr-Arg-Gly-Glu-Arg-Cys-Asp-Leu-Glu-
(643-663) Asp-Arg-Asp-Arg-Ser-Glu-Leu-Ser-Tyr-**OH**

TABLE 8

(Note: **H** signifies the amino terminus; **OH** signifies the carboxyl terminus. The two underlined Tyr residues are not part of the HCV sequence but are engineered there for ease of iodinating the peptide at a later time)

5	p643b (643-683)	H-Ala-Cys-Asn-Trp-Thr-Arg-Gly-Glu-Arg-Cys-Asp-Leu-Glu-Asp-Arg-Asp-Arg-Ser-Glu-Leu-Ser-Pro-Leu-Leu-Leu-Thr-Thr-Thr-Gln-Trp-Gln-Val-Leu-Pro-Cys-Ser-Phe-Thr-Thr-Leu-Pro- OH
10	p666 (666-683)	H-Leu-Leu-Thr-Thr-Thr-Gln-Trp-Gln-Val-Leu-Pro-Cys-Ser-Phe-Thr-Thr-Leu-Pro- <u>Tyr</u> - OH
	p691 (691-714)	H-His-Leu-His-Gln-Asn-Ile-Val-Asp-Val-Gln-Tyr-Leu-Tyr-Gly-Val-Gly-Ser-Ser-Ile-Ala-Ser-Trp-Ala-Ile- OH
15	p2302 (2302-2352)	H-Lys-Lys-Pro-Asp-Tyr-Gln-Pro-Pro-Val-Val-His-Gly-Cys-Pro-Leu-Pro-Pro-Pro-Lys-Ser-Pro-Pro-Val-Pro-Pro-Pro-Lys-Lys-Lys-Arg-Thr-Val-Val-Leu-Thr-Glu-Ser-Thr-Leu-Ser-Thr-Ala-Leu-Ala-Glu-Leu-Ala-Thr-Arg-Ser-Phe- OH
20	p380-JH1 (380-436)	H-Gly-Val-Asp-Gly-His-Thr-Arg-Val-Thr-Gly-Gly-Val-Gln-Gly-His-Val-Thr-Ser-Thr-Leu-Thr-Ser-Leu-Phe-Arg-Pro-Gly-Ala-Ser-Gln-Lys-Ile-Gln-Leu-Val-Asn-Thr-Asn-Gly-Ser-Trp-His-Ile-Asn-Arg-Thr-Ala-Leu-Asn-Cys-Asn-Asp-Ser-Leu-Gln-Thr-Gly- OH
25	p380J (380-436)	H-Gly-Val-Asp-Gly-His-Thr-His-Val-Thr-Gly-Gly-Arg-Val-Ala-Ser-Ser-Thr-Gln-Ser-Leu-Val-Ser-Trp-Leu-Ser-Gln-Gly-Pro-Ser-Gln-Lys-Ile-Gln-Leu-Val-Asn-Thr-Asn-Gly-Ser-Trp-His-Ile-Asn-Arg-Thr-Ala-Leu-Asn-Cys-Asn-Asp-Ser-Leu-Gln-Thr-Gly- OH
30	p408J (408-436)	H-Ser-Gln-Lys-Ile-Gln-Leu-Val-Asn-Thr-Asn-Gly-Ser-Trp-His-Ile-Asn-Arg-Thr-Ala-Leu-Asn-Cys-Asn-Asp-Ser-Leu-Gln-Thr-Gly- OH
35		

The polypeptides illustrated in Table 2 also may be prepared in a stepwise fashion or in a fragment coupling protocol using various side chain protection methodologies known to those skilled in the art. The polypeptides also may be prepared using enzymatic methodology.

5

Further, the polypeptides useful in the practice of this invention may be prepared using recombinant technologies. Briefly, DNA sequences which encode the desired polypeptides are preferably assembled from fragments of the total desired sequence. The fragments are generally prepared using well known automated processes and apparatus. After the complete sequence has been prepared, the desired sequence is incorporated into an expression vector which is transformed into a host cell. The DNA sequence then is expressed by the host cell to give the desired polypeptide which is harvested from the host cell or from the medium in which the host cell is cultured. In most cases, the manufactured DNA sequence is assembled using codons which are known to be best expressed in the host cell. When smaller peptides are to be made using recombinant technologies, it may be advantageous to prepare a single DNA sequence which encodes several copies of the desired polypeptide in a connected chain. The long chain then is isolated, and the chain is cleaved into the shorter, desired sequences.

The amino acid sequence for p1684 is reverse translated to give the codons listed in Table 3 which are optimized (where not inconsistent with assembly and synthesis of fragments) to facilitate high level expression in E. coli. Individual oligonucleotides are synthesized on Applied Biosystem 380A DNA synthesizer using methods and reagents recommended by the manufacturer. These purified oligonucleotides are annealed and ligated together to assemble the entire DNA sequence for digestion with BamHI and SalI, allowing ligation into pUC18. The resulting plasmid is suitably transformed into E. coli JM103 cells. Table 3 also lists preferred codons to express p1 and p1223.

In order to establish that a clone expresses the DNA sequence, it is grown at 37°C in 50 ml Luria Broth, in a 250 ml Erlenmeyer flask. When the culture reaches an OD600 of 0.3-0.5, IPTG is added to a final concentration of 1 mM to induce expression. Samples (1.5 ml) are removed at one hour

intervals, and the cells are pelleted and resuspended to an OD600 of 10.0 in 2X SDS/PAGE loading buffer. Aliquots (15 μ l) of the prepared samples are loaded on a 15% SDS/PAGE gel, the expressed polypeptides separated, and then electrophoretically transferred to nitrocellulose for immunoblotting. The nitrocellulose sheet containing the transferred proteins is incubated with a blocking solution for one hour and incubated overnight at 4°C with HCV patients' sera diluted in TBS containing 5% E. coli JM103 lysate. The nitrocellulose sheet is washed three times in TBS, then incubated with HRPO-labelled goat anti-human IgG, diluted in TBS containing 10% fetal calf sera. The nitrocellulose is washed three times with TBS and the color is developed in TBS containing 2 mg/ml 4-chloro-1-naphthol, 0.02% hydrogen peroxide and 17% methanol. Strong immunoreactive band formation with HCV patients' sera indicates that the synthetic polypeptide is expressed in E. coli in immunologically reactive form.

Preferred formats for assays using the polypeptides described above are provided in the following examples. However, other assay formats known to those skilled in the art can be used. These assays include ion capture assays, microparticle assays, and the use of scanning tunneling microscopy in which at least one polypeptide is attached to a solid phase, contacted with a test serum, and the surface of the solid phase is scanned for antigen-antibody complexes.

Example 1 describes a confirmatory assay. Example 2 describes a combination assay. Example 3 describes a synthetic polypeptide-based assay. Example 4 describes an immunodot assay. Example 5 describes a competition assay. Example 6 describes an EIA assay in which peptides 380-436 and 447-483, 643-683 and 2302-2352 are used. Example 7 describes an EIA utilizing peptide p380.LG. Example 8 describes an EIA utilizing peptide 2302 (NS-5) compared to an EIA utilizing antigens NS3 (CKS-33C, NS4 (C-100) or CORE (CKS-CORE). Example 9 describes the PEPSCAN protocol followed. Example 10 describes the preparation of synthetic peptides based on the sequences of four distinct HCV isolates. Example 11 describes the data obtained from studies on chronic patients and plasma donors using the peptides of Example 10. These examples are

meant to illustrate, but not to limit, the scope and spirit of the present invention.

EXAMPLES

5

Example 1. CONFIRMATORY ASSAY

The confirmatory assay uses at least two polypeptides containing HCV antigenic epitopes which are preferably prepared and isolated from different sources. One polypeptide is used to screen serum or plasma samples. The other polypeptide is used to confirm the presence of a HCV
10 antibody in a sample initially identified as containing a HCV antibody by the screening procedure.

In the presently preferred confirmatory assay, the screening procedures uses a recombinant C100-3 polypeptide. The C100-3
15 recombinant polypeptide is believed to contain multiple epitopes as well as an immunodominant region defined by the 1689-1806 amino acid sequence. The C100-3 polypeptide is expressed in recombinant yeast cells and isolated from the cell extract as described in EPA Publication Number 0 318 216. Other recombinant polypeptides containing amino acid
20 sequences essentially duplicative of C100-3 also may be used.

The other peptide used in the confirmatory assay is a synthetic peptide selected from the group consisting of p1, p35, p99, p1192, p1223, p1684, p1689, p1694, p1866, and p1899. Preferably the peptide is p1684
25 or p1866. These peptides were prepared following procedures described above. In the confirmatory assay, both C100-3 and the synthetic peptides, p1684, p1694 or p1866, were separately coated onto polystyrene beads. A combination of synthetic peptides coated on a polystyrene bead also may be used if desired.

30

The polystyrene beads are first washed with distilled water and propanol, then incubated with crude or purified HCV synthetic peptides diluted to 0.1-20.0 $\mu\text{g/ml}$ in a 0.1 M solution of an appropriate buffer containing about 0.4-0.5 M NaCl, about 0.0022% Triton X-100, and
35 adjusted to about pH 6.5-10.0. The following buffers, tris, $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, boric acid, and citrate buffers are preferred and are optimized for each peptide; preferred buffers, pH and coating concentration for the synthetic

peptides are listed in Table 4. Successful coatings also have been accomplished with lower or higher pH. The beads are incubated in the antigen solution for about two hours at 38-42°C, washed in phosphate buffer solution (PBS) and soaked in 0.1% Triton X-100 in PBS for 60 minutes at 38-42°C. The beads then are washed two times in PBS, overcoated with a solution of 5% (w/v) bovine serum albumin (BSA) in PBS for 60 minutes, and washed three times with PBS. Finally, the beads are overcoated with 5% (w/v) sucrose in PBS and dried under nitrogen or air.

10 The peptides are each individually coated onto polystyrene beads and used in an antibody capture format. Ten microliters of sample are added to the wells of a reaction tray along with 400 µl of a sample diluent and a peptide coated bead. The sample diluent consists of about 10% (v/v), or less, bovine serum and about 20% (v/v), or less, goat serum in 20 mM Tris phosphate buffer containing 0.20%, or less, (v/v) Triton X-100, 3% (W/v), or less, BSA. When the recombinant yeast C100-3 polypeptide is used, antibodies to yeast antigens which may be present in a test sample are reacted with yeast extracts which are added to the sample diluent (typically about 200 µg/ml). The addition of yeast extracts to the sample diluent is used to prevent false positive results. The final material is sterile filtered and filled in plastic bottles, and preserved with 0.1% sodium azide.

After one hour of incubation at 40°C, the beads are washed and 200 µl of conjugate is added to the wells of the reaction tray.

25

The preferred conjugate is goat anti-human IgG horseradish peroxidase conjugate. Concentrated conjugate is purchased from Kirkegaard and Perry Laboratories, Inc., Gaithersburg, Maryland, and is titrated to determine a working concentration. A 20-fold concentrate of the working conjugate solution is then prepared by diluting the concentrate in diluent. The conjugate diluent includes 10% (v/v) bovine serum, 10% (v/v) goat serum and 0.15% Triton X-100 in 20 mM Tris buffer, pH 7.5 with 0.01% gentamicin sulfate, pink dye and antifungal agents as preservatives. The conjugate is sterile filtered and filled in plastic bottles.

35

After one hour of incubation with the conjugate at 40°C, the beads are washed, exposed to the OPD substrate for 30 minutes at room

temperature, and the reaction terminated by the addition of 1 N H₂SO₄.
The absorbance is read at 492 nm.

5 Samples found to be repeatably reactive by a screening assay using
the polypeptide C100-3 are tested in duplicate using p1684 or p1689
coated beads. Reactive specimens are considered confirmed samples.
Samples not reacting with p1684 or p1689 are tested in duplicate with
p1694 and p1866 beads. Samples reacting with one or both of these
peptides are considered confirmed. Those specimens not reacting with any
10 of these peptides are considered nonconfirmed.

In order to maintain acceptable specificity, the cutoff for the assay
should be at least 5-15 standard deviations above the absorbance value of
the normal population mean. Consistent with these criteria, a cutoff for the
15 assay may be selected which clearly separated most of the presumed "true
negatives" from "true positive" specimens. A general cutoff value may be
calculated as about 2.1 to 8 times the negative control mean absorbance
value.

20 Confirmatory Assay Performance

1. Intravenous Drug User Samples

Samples were collected from a population of intravenous drug
users enrolled in an NIH-funded study. The population consisted of
individuals who were acknowledged users of intravenous drugs selected
25 over a two-year period from patients at the Edward Hines Jr. Veteran's
Administration Hospital in Maywood, Illinois by Dr. Connie Pachucki and
members of the Infectious Diseases staff.

As illustrated in Table 5, a total of 296 specimens, each obtained
30 from a single donor, were screened using recombinant yeast C100-3
polypeptide. A total of 271 of 296 (91.6%) specimens initially tested
positive; upon retesting, 269 of 271 (99.3%) were repeat-positives.

Confirmatory testing indicated that 263 of 269 (97.8%) of the repeat
35 positives were reactive with p1689, five specimens were non-reactive with
p1689, and one specimen was not tested with any of the confirmatory
polypeptides. Four of the five specimens which were non-reactive with

p1689 were reactive with p1866 only; one specimen which was non-reactive with p1689 was reactive with only p1694.

5 All specimens which were repeatably reactive were confirmed reactive in assays using the HCV synthetic peptides.

TABLE 5

INTRAVENEOUS DRUG USERS SAMPLES

10	Confirmatory Testing					No. of Repeat Positives Confirmed
	C100-3 Initial Pos.	C100-3 Repeat Pos.	p1689	p1866	p1694	
15	271/296	269/271	263/269	4/5	1/5	268/269

2. Chimpanzees Samples

Confirmatory assays were used to evaluate 92 samples from six chimpanzees. All were initially reactive with recombinant C100-3.
 20 (Duplicative, repeat testing of chimp sera was not done because of the rare nature of these specimens and their utility for serological studies with other HCV antigens). Eighty-three of 92 (90.2%) specimens were confirmed reactive using p1689. Confirmation of initial reactives improved to 96.7% (89 of 92) when repeat testing with p1694 and p1866 was done.

25

3. Chiron Corporation Non-A, Non-B Hepatitis Virus Proficiency Panel #2

A proficiency panel comprised of neat and diluted human plasma including specimens containing antibodies to HCV C100-3 was provided
 30 by scientists at the Chiron Corporation (12 specimens). This panel contains specimens ranging from low to high reactivity in other assays, non-reactive presumed "true negative" specimens, and reactive specimens diluted to give low-level or negative results.

35 Results using the confirmatory assay on the Chiron Corporation Non-A, Non-B Hepatitis Virus Proficiency Panel #2 indicated 9 of 9 (100%) of the

specimens reactive by the preliminary screening assay are confirmed by p1689. All negative specimens were non-reactive.

4. Confirmatory Testing on NANB Panel II

5 A panel of highly pedigreed human sera from Dr. H. Alter, NIH, Bethesda, MD, containing infectious HCV sera, negative sera and other disease controls were tested. A total of 44 specimens were present in the panel.

10 All specimens (16/16, 100%) reactive in the assay using C100-3 were confirmed by p1689, as shown in Table 6. Again, there were no nonspecific reactives as all pedigreed negative or "other disease" controls were non-reactive in the peptide assay.

15

TABLE 6

Comparison of p1689 Results with HCV Screening Assay Results on NANB Panel II (H. Alter, NIH)

SAMPLE	MANUAL S/CO	C-100 MACHINE S/CO	ORTHO C1003 S/CO	p1689 EIA S/CO
1	>5.88	>6.47	>6.38	>8.33
2	0.63	0.93	0.27	0.45
3	>5.88	>6.47	>6.38	>8.33
4	>5.88	>6.47	>6.38	>8.33
5	0.43	0.35	0.16	0.43
6	>5.88	>6.47	>6.38	>8.33
7	0.46	0.73	0.36	0.32
8	0.41	0.50	0.32	0.38
9	1.87	2.21	0.91	2.84
10	0.35	0.41	0.32	0.30
11	0.48	0.45	0.27	0.46
12	0.32	0.41	0.17	0.39
13	0.48	0.69	0.32	0.51
14	0.37	0.40	0.19	0.32
15	>5.88	>6.47	>6.38	>8.33
16	>5.88	>6.47	>6.38	>8.33
17	0.34	0.40	0.20	0.44
18	3.01	3.68	0.68	6.80
19	0.74	0.61	0.53	0.72
20	0.53	0.59	0.28	0.33
21	>5.88	>6.47	>6.38	>8.33
22	0.24	0.26	0.20	0.23
23	>5.88	>6.47	>6.38	>8.33
24	0.69	0.64	0.53	0.70

23

25	0.50	0.60	0.49	0.40
26	3.41	4.11	0.77	6.61
27	0.62	0.74	0.30	0.65
28	0.61	0.77	0.08	0.47
29	0.34	0.42	0.13	0.33
30	1.58	2.40	1.26	2.65
31	0.32	0.35	0.22	0.37
32	>5.88	>6.47	>6.38	>8.33
33	0.45	0.48	0.24	0.45
34	>5.88	>6.47	>6.38	>8.33
35	>5.88	>6.47	>6.38	>8.33
36	0.37	0.36	0.21	0.40
37	0.40	0.46	0.24	0.52
38	>5.88	>6.47	>6.38	>8.33
39	0.40	0.49	0.30	0.46
40	0.53	0.59	0.30	0.56
41	0.41	0.26	0.15	0.32
42	0.52	0.55	0.33	0.60
43	0.28	0.30	0.36	0.33
44	0.44	0.57	0.35	0.53

Data presented demonstrate the efficacy of the confirmatory assay for detection of antibodies to HCV antigens. The current assay is both sensitive and specific for detection of antibodies to HCV antigens.

5

The data further support the utility of the confirmatory strategy using synthetic peptides. The synthetic peptides serve as an independent source of antigen for use in immunoassays. The ability to confirm an average of 99% of repeatably reactive specimens in high risk or pedigreed positive

10 HCV panels, establishes the utility of this strategy.

Example 2. COMBINATION ASSAY

The combination assay uses more than one polypeptide antigen coated on the same bead. To prepare multiple polypeptide-containing
15 beads, the polystyrene beads described in Example 1 are contacted simultaneously with the polypeptide in appropriate buffer solutions. After the beads have been contacted with the polypeptides, the bead is treated further as described above.

20 For a polystyrene bead containing both C100-3 and p1694, the sensitivity of the assay increases. As graphically illustrated in Figure 3a, adding about 0.3, 0.95, and 3 micrograms of p1694 to the coating solution,

respectively, shows a significant increase in the signal when the detection procedures of Example 1 are utilized. Figure 3b graphically illustrates the data which show no corresponding increase in the signals (such as may attend non-specific binding) generated from negative human plasma.

5

Example 3. SYNTHETIC POLYPEPTIDE-BASED ASSAY

The use of synthetic polypeptides which contain epitopes of HCV antigens provide immunological assays which have increased sensitivity and may be more specific than HCV immunological assays using the SOD fusion polypeptide C100-3. The use of shorter amino acid sequences on polystyrene beads provides an increase in sensitivity.

The increased sensitivity of an assay employing synthetic polypeptide compared to recombinant C100-3 polypeptide was demonstrated in a serial dilution study. The serial dilution study employed 15 samples which were identified as having antibodies to HCV antigens using a recombinant C100-3 screening assay. Each positive sample was assayed using recombinant C100-3 polypeptide in one assay and p1689 polypeptide in a second assay, and the samples were then diluted twofold until the S/CO value was less than one. In 12 samples, the p1689 polypeptide gave increased sensitivity (larger S/CO values) at all dilutions. In two samples, the p1689 polypeptide and the recombinant yeast C100-3 polypeptides were essentially equivalent. In one sample, the p1689 polypeptide have a negative response to a positive sample at all dilutions.

25

Additional studies on samples from serial bleeds of three chimps which developed an acute resolved case of HCV infections and three chimps which developed chronic HCV infections showed different immunological responses believed to be due to both the type of infection and the polypeptide used in the assay. This study assayed serum from serial bleeds of six chimps inoculated with HCV. The assay protocols were similar to those described in Example 1 above with the following differences.

35

The antibodies, IgG, IgM, and IgA were detected using affinity purified goat antibodies to human IgG, IgM and IgA coupled to horseradish peroxidase (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, Maryland)

which were used at working concentrations of 0.2 µg/ml anti-IgG; 0.5 µg/ml anti-IgM and 0.2 µg/ml anti-IgA. Serum dilutions for each assay were 1:41 for IgG, 1:101 for IgM and 1:41 for IgA.

- 5 The polypeptides that were used in the study include C100-2, p1694, p1689 and p1866.

10 Briefly, beads containing the polypeptides were incubated with diluted serum for one hour at 40°C, the beads then were washed and incubated with the appropriate goat antibody for one hour at 40°C. The beads were washed again and the assay was developed by incubating the beads with OPD for 30 minutes at room temperature. The color development was quenched with 1 N sulfuric acid and the results read at 492 nm.

15 All chimps developed antibodies that were detected by C100-3, p1684, and p1866 within seven to 17 wells post-inoculation (WPI). Within each chimp, IgG antibody reacting with C100-3, p1684, and p1866 appeared at approximately the same time. The response to p1694 and 20 p1866 was variable within that time period with indications that antibody to these two peptides can be either undetectable or significantly delayed following HCV infection. These data suggest that, out of the five peptides tested, antibody to C100-3, p1684, or p1689 would be the earliest and most consistent serologic indicator of HCV infection.

25 IgM antibody was detected in only three of the six chimps studied. The response of each of the three animals to C100-3, p1684, p1689 and p1694 was detected in seven to ten WPI whereas IgM antibody to p1866 was undetectable in two chimps and delayed in the third. All IgM 30 responses were short lived with levels falling below positive (S/N less than 3.0) within two to 22 weeks.

35 The explanation and significance of finding IgM antibodies in three chimps with acute resolved disease while not detecting IgM antibodies in three chimps with chronic infection is unexpected. Preliminary experimental results indicated that false negative IgM results due to preferential IgM binding is an unlikely explanation. If the pattern observed

in these six chimps with the five peptides holds true, antibody assays will provide important HCV prognostic information.

5 A positive IgA response (S/N greater than 3.0) was detected in only two of the six chimps, and proved to be either biphasic or significantly later than the IgG or IgM response. Although these two chimps had chronic disease, no conclusions regarding the significance of IgA antibodies can be made since sera from the three resolved chimps is available only through 30 to 40 WPI.

10

The results show the polypeptides when used to assay for antibodies to HCV antigens are useful to follow the progression of HCV infection, and that the polypeptides exhibit unexpected sensitivity to different antibodies generated during the clinical progression of HCV infection.

15

Example 4. IMMUNODOT ASSAY

The immunodot assay system uses a panel of purified synthetic polypeptides placed in an array on a nitrocellulose solid support. The prepared solid support is contacted with a sample, and captures specific 20 antibodies to HCV antigens. The captured antibodies are detected by a conjugate-specific reaction. Preferably, the conjugate-specific reaction is quantified using a reflectance optics assembly within an instrument which has been described in U.S. Patent Application Serial No. 07/227,408, filed August 2, 1988. The related U.S. Patent Applications Serial No. 25 07/227,272, 07/227,586 and 07/227,590 further describe specific methods and apparatus useful to perform an immunodot assay. Briefly, a nitrocellulose-base test cartridge which may be used in an automated process for performing an immunodot assay described above is illustrate in Figure 4. Each polypeptide is contained within a specific reaction zone on 30 the test cartridge. After all the antigenic polypeptides have been placed on the nitrocellulose, excess binding sites on the nitrocellulose are blocked. The test cartridge then is contacted with a test sample such that each antigenic polypeptide in each reaction zone will react if the sample contains the appropriate antibody. After reaction, the test cartridge is washed and 35 any antigen-antibody reactions are identified using suitable well-known reagents.

As described in the patent applications listed above, the entire process is amenable to automation. The specifications of these applications related to the methods and apparatus for performing an immunodot assay, listed above, are incorporated herein by reference.

5

In a preferred immunodot assay, the synthetic polypeptides p1223, p1684, p1689 and p1866 were diluted into an aqueous buffered solution (polypeptide diluent: 0.03% Triton X-100 and 0.1% sodium azide in 50 mM Hepes buffer, pH 7.6) and applied to a preassembled nitrocellulose test cartridge at about 40 ng in each reaction zone. After drying the cartridge overnight at room temperature, the nonspecific binding capacity of the nitrocellulose phase was blocked. The blocking solution contained 1% porcine gelatin, 1% casein enzymatic hydrolysate, 5% Tween-20, 0.1% sodium azide, 0.5 M sodium chloride and 20 mM Tris, pH 7.5.

15

Test cartridges were incubated with samples 00642 and 423 (see Table 1) and ALT 27. The sample ALT 27 was obtained from a volunteer donor having elevated alanine aminotransferase levels. After sample incubation, sequential incubations with a biotin-conjugated goat anti-human immunoglobulin -specific antibody, an alkaline phosphatase-conjugated rabbit anti-biotin specific antibody, and 5-bromo-4-chloro-3-indolyl phosphate produced a colored product at the site of the reaction.

20

A detectable reaction is defined by the formation of a visually discernable product at the antigen site on the array; when quantified by the instrument, a reflectance density (D_r) value of greater than or equal to approximately 0.0150 above background is obtained. None of the tested polypeptides elicited a detectable reaction with a negative control serum that was previously demonstrated negative for antibodies to HCV antigens using a recombinant C100-3 polypeptide.

25

30

A reaction with each of the synthetic polypeptides p1684, p1689, p1694 and p1866 occurred when the prepared test cells were incubated with either sample 00642 (1:100 dilution in negative serum) or sample 423 (1:40 dilution in negative serum). Polypeptide p1223, in addition to polypeptides p1684, p1689, p1694 and p1866 demonstrated a significant reaction with the elevated ALT 27 specimen. In all specimens, highest

35

reactivity was obtained with p1689. Enhanced reactivity of polypeptide p1684 with sample 00642 was achieved through subtle modification of the antigen dilution (the modified polypeptide diluent was 0.5 M sodium chloride, 0.0022% Triton X-100 and 0.1 M Tris/HCl, pH 8.5).

5

The net reflectance (Dr) for a test cartridge containing the polypeptides p1223, p1684, p1689, p1694 and p1866 which indicate a positive or negative response is set out in Table 7.

10

TABLE 7
NET REFLECTANCE DENSITY (INTERPRETATION*)

Sample	p1223	p1684	p1689	p1694	p1866
Neg Ctrl	-.0051 (-)	-.0042 (-)	.0047 (-)	.0012 (-)	.0134 (-)
Sample	.0011 (-)	.0281 (+)	.4401 (+)	.0748 (+)	.1444 (+)
00642	.0002 (-)	.0368 (+)	.4552 (+)	.0853 (+)	.1564 (+)
(1:100)					
Sample 423	.0073 (-)	.1501 (+)	1.5389 (++)	.1922 (+)	.0323 (+)
(1:40)					
Sample	.3040 (+)	10.40 (++)	29.64 (+++)	13.00 (++)	7.450 (++)
ALT27	.1469 (+)	9.619 (++)	30.20 (+++)	13.26 (++)	7.887 (++)

* + / - Values Based on S/N

- Dr/Bkg < 2.5

+ 2.5 < Dr/Bkg < 100

++ 100 < Dr/Bkg < 1000

+++ Dr/Bkg > 1000

15

Example 5. COMPETITION ASSAY

The synthetic peptides containing antigenic HCV epitopes are useful for competition assays. To perform a neutralization assay, peptides representing epitopes within the C100-3 region such as p1694, p1684 or p1689 are 34 and mixed with a specimen diluent to a final concentration of 0.5-50 µg/ml. Ten microliters (µl) of specimen or diluted specimen is added to a reaction well, followed by 400 µl of the specimen diluent containing peptide and if desired, the mixture may be pre-incubated for about 15

25

minutes to two hours. A bead coated with C100-3 antigen of HCV then is added to the reaction well and incubated for one hour at 40°C. After washing, 200 µl of peroxidase labeled goat anti-human IgG in conjugate diluent is added and incubated for one hour at 40°C. After washing, OPD substrate is added and incubated at room temperature for 30 minutes. The reaction is terminated by the addition of 1 N sulfuric acid, and the absorbance read at 492 nm.

Samples containing antibodies to the C100-3 antigen generate a reduced signal caused by the competitive binding of the peptides to these antibodies in solution. The percentage of competitive binding may be calculated by comparing the absorbance value of the sample in the presence of a synthetic peptide to the absorbance value of the sample assayed in the absence of a synthetic peptide at the same dilution.

Example 6. EIA ASSAY

Beads were coated with either peptides 380-436 and 447-483, 643-686 or 2302-2353 according to the method described in Example 1, except that peptides 380-436 and 447-483 were coated simultaneously on the same solid phase, both sequences being from the putative envelope region of HCV. Either peptide alone had activity in this type of assay. EIA was performed using each bead configuration described herein. The EIA method performed was as is described in Example 1, with the cutoff set at four times the negative control value. Table 9 presents data obtained from these assays in which serum specimens from patients diagnosed with chronic NANBH were assayed.

TABLE 9
ANTIGEN
POS./NO. TESTED

	p380	p643b	p2302
	70/165 (42%)	62/165 (38%)	102/165 (62%)

Example 7. EIA UTILIZING p380.LG and p380

Beads were coated with either p380.LG or p380 according to Example 1. An EIA following the procedures of Example 1 was used to assay samples. As can be seen by the data presented in Table 10, the p380.LG peptide detected antibody in specimens that were negative to p380. The HCV sequence is highly variable in the region 380-436 a.a. Therefore, there is reasonable probability that differentiation between HCV "serotypes" based on reactivity of human specimens to one or the other of these envelope region peptide sequences is possible. The data of Table 10 suggest that p380.LG can detect chronically infected HCV patients who are negative to p380.

TABLE 10

SAMPLE	p380		p380.LG	
	OD	S/N	OD	S/N
#8	.155	2.12	.399	5.87
#28	.246	3.37	.950	13.97
#23	.114	1.56	.458	6.74

Example 8. EIA UTILIZING p2302

Beads were coated with either p2302 or recombinant HCV antigens according to the method of Example 1. The bead coated with recombinant antigens comprised antigens from the NS3 (CKS-33C), NS4 (C-100) and Core (CKS-CORE) regions of the HCV genome. A patient sample which exhibited seroconversion to p2302, but not to antigens of HCV 2.0, is shown in Figure 5. Thus, this peptide improves the ability to detect HCV infected individuals.

Example 9. PEPSCAN PROTOCOL

NS1 region of HCV genome from a.a. 600-720 was mapped with PEPSCAN analysis, which is serological analysis of series of overlapping peptides spanning the protein sequence to identify immunogenic domains. A total of 106 overlapping hexamer peptides were synthesized on polypropylene pins following the manufacturer's instructions (Cambridge Research Bioscience, Valley Stream, NY). Fab dimers of IgG purified from sera of individuals seropositive for HCV were tested with these peptides.

Based on the reactivity in EIA (performed as described by the manufacturer) four peptide sequences were selected as illustrated in Table 11.

TABLE 11

5 AMINO ACID SEQUENCE OF PEPTIDES SELECTED FROM THE
NS1 REGION OF HCV GENOME (A.A. 600-720) BASED ON
PEPSCAN ANALYSIS

10	A.A. No. of HCV Genome	Peptide Sequence
	607-627	CLVDYPYRLWHYPCTINYTIF
	643-663	ACNWTRGERCDLEDRDRSELSY
	666-683	LLTTTQWQVLPCSFTTPLY
15	691-714	HLHQNIVDVQYLYGVGSSIASWAI

Each of these peptides were synthesized by a stepwise solid phase synthesis, starting the the carboxy terminus residue. A panel of sera
20 positive for antibodies to C-100 protein of HCV was tested for their reactivity to NS peptide by microtiter EIA as described below.

EIA PROTOCOL

Wells of microtiter plates were coated with 100 µl of the peptide at 10 µg/ml in 0.02 M bicarbonate buffer, pH 9.5 at ambient temperatures for 12-
25 16 hours. After washing with phosphate buffered saline (PBS) which also contained 0.01% sodium dodecyl sulfate (SDS) and 0.05% Tween-20® (available from BioRad Laboratories, Richmond, CA), free sites were overcoated with 1% BSA in bicarbonate buffer pH 9.5. Plates were stored at 4°C following a final wash.

30

Sera from individuals seropositive for antibodies to HCV C-100 were serially diluted in 100 µl of a buffer containing 20 mM sodium phosphate, pH 7.4, 0.15 M NaCl, 20% normal goat serum, 10% fetal calf serum, 5 mM EDTA, 10 mM EGTA, 50 mM Tris, 0.2% Tween-20 with sodium azide as
35 preservative, pH 6.8. The diluted sera were reacted with peptides in microtiter wells for three hours at 37°C or overnight at ambient temperatures. The plates were washed and 100 µl of appropriately diluted

goat anti-mouse (HH) Horseradish Peroxidase (HRPO)-conjugated antibody (Jackson Immunochemicals, West Grove, PA) was added. The plates were incubated at 37°C for two hours. After a final wash, 100 µl of O-phenylenediamine 2HCl (OPD) color reagent was added. The reaction was carried out at room temperature in the dark for 20-25 minutes, and stopped by the addition of 100 µl of 1N H₂SO₄. The absorbance of the reaction mixture was recorded at 492 nm. A negative control which was previously confirmed to be negative for HCV infection was included with each plate in triplicate. The sample was considered reactive if the absorbance of the sample at a 1:2000 dilution was three times the absorbance of the negative control at the same dilution. Table 12 illustrates the reactivity of these samples with each of the peptides.

The legend for Table 12 is as follows:

+ = Sample showing A₄₉₂ 3X negative control

++ = Sample titering to 1:5000 dilution

+++ = Strong reactivity with sample titering to 1:10,000 dilution.

TABLE 12
REACTIVITY OF A PANEL OF HCV SEROPOSITIVE SAMPLES WITH NS1
PEPTIDES BY MICROTITER EIA
Peptides Selected From NS1 Region (A.A. Numbes)

Sample ID*	607-627	643-663	666-683	691-714
15	-	-	+	-
22	-	-	-	-
23	-	-	-	-
24	+/-	-	-	-
25	+/-	-	-	-
32	+/-	+	+	-
36	-	-	-	-
46	-	-	-	-
50	-	-	-	-
65	-	-	-	-
70	-	+	+	-
71	-	-	+	-
75	-	-	-	-
89	-	++	-	+
95	-	++	++	+

33

100	-	++	-	-
102	-	-	-	-
108	-	-	-	-
130	-	-	+	-
137	-	-	-	-
LG	+	++	+++	+
301060	-	++	+++	+
PB3178	++	+	+++	+
PB3180	++	+	+++	++
300423	++	-	+++	+
% Positive	16	36	44	27

* All samples showed the presence of antibodies to Hepatitis C Virus by EIA as well as Western Blot Analysis.

Example 10. SYNTHETIC PEPTIDES TO 5' END OF E2/NS1

- 5 The 5' end of the E2/NS1 putative structural region of HCV has been shown to be hypervariable with respect to both nucleic acid and amino acid content among multiple virus isolates studied throughout the world. It has been speculated but not proven, that this region is involved in a viral mechanism to defeat host immune surveillance. If this region were to
- 10 mutate under humoral immune selective pressures, then specific antibody responses to genotypic variants of the virus should be detectable in HCV infected patients. We demonstrated the presence of multiple epitopes within this hypervariable region, thus supporting the hypothesis that, an in flaviviruses and pestiviruses, this E2/NS1 region is exposed to the host's
- 15 immune system and the subsequent selective pressures toward antigenically distinct variants.

- Synthetic peptides were prepared based on the sequences of four distinct HCV isolates. The peptides represented the sequence from amino
- 20 acid 380 through amino acid 436 of the large open reading frame of HCV. A smaller peptide encoded by amino acids 408-436 also was synthesized. The sequences were obtained from HCV-1 (Chiron prototype), HCV G83 (p380.LG), HCV-JH-1 and HCV-J. Peptide HCV-JH-1 was as described by K. Takeuchi et al., Nucleic Acids Research 18:46226 (1989), while peptide
- 25 HCV-J was as described by N. Kato et al., Proc. Nat'l. Acad. Sci. 87:9524-9528 (December, 1990). Though encoded by the same genomic region,

each of these peptides are divergent from each other by a minimum of 25% (range 25-35%) based on amino acid sequence over the span of 57 amino acids. Peptide 408-436 was prepared based on the sequence of HCV-JH-1 and HCV-J which are identical in this region. Each of these peptides was coated onto a solid phase under the coating conditions listed in Table 4 and used as antigenic targets in separate enzyme immunoassays (EIAs), according to the methods described hereinabove for EIA. The specificity of the individual EIA's were demonstrated by testing populations of normal plasma donors. A cutoff value of 4 times the negative control sample was used for all analyses. Populations of chronic HCV patients from the U.S. and Italy, acute phase HCV patients from the U.S. and Italy, and HCV antibody positive plasma donors from the U.S. and Japan were tested.

TABLE 4

PEPTIDE	COATING CONC. µg/ml	COATING BUFFER	OTHER COMPONENTS IN COATING SOLUTION
1-75	2.0	0.1M NaPhosphate, pH 6.5	0.4M NaCl, .0022% Triton X-100
35-75	2.0	0.1M NaPhosphate, pH 6.5	0.4M NaCl, .0022% Triton X-100
99-126	2.0	0.1M NaPhosphate, pH 6.5	0.4M NaCl, .0022% Triton X-100
195-262	2.0	0.1M NaPhosphate, pH 6.5	0.4M NaCl, .0022% Triton X-100
230-262	2.0	0.1M NaPhosphate, pH 6.5	0.4M NaCl, .0022% Triton X-100
1357-1407	2.0	0.1M Boric Acid, pH 9.0	0.4M NaCl, .0022% Triton X-100
1418-1457	2.0	0.1M Boric Acid, pH 9.0	0.4M NaCl, .0022% Triton X-100
1569-1593	3.0	0.1M Tris/HCl, pH 8.5	0.5M NaCl, .0022% Triton X-100
1899-1930	2.0	0.1M Tris/HCl, pH 8.5	0.5M NaCl, .0022% Triton X-100
1192-1240	2.0	0.1M Boric Acid, pH 9.0	0.4M NaCl, .0022% Triton X-100
1223-1240	5.0	0.1M Boric Acid, pH 9.0	0.4M NaCl, .0022% Triton X-100
1684-1750	1.0	0.1M Boric Acid, pH 10.0	0.4M NaCl, .0022% Triton X-100
1689-1805	1.0	0.1M Boric Acid, pH 10.0	0.4M NaCl, .0022% Triton X-100
1694-1735	3.0	0.1M Tris/HCl, pH 8.5	0.5M NaCl, .0022% Triton X-100
1866-1930	0.75	0.1M Tris/HCl, pH 8.5	0.5M NaCl, .0022% Triton X-100
380-436	3.0	0.1M Tris/HCl, pH 8.5	0.9% NaCl
447-483	3.0		
643-683	3.0	0.1M Tris/HCl, pH 8.5	0.5M NaCl, .0022% Triton X-100
2302-2352	3.0	0.1M Borate	0.4M NaCl, .0022% Triton X-100
p380JH1	5.0	0.1M Borate, pH 10.0	0.4M NaCl, .0022% Triton X-100
p380J	5.0	0.1M Borate, pH 10.0	0.4M NaCl, .0022% Triton X-100
p408J	5.0	0.1M Borate, pH 10.0	0.9% NaCl

Data from studies on chronic patients and plasma donors is presented in Table 13. Antibodies were detected to each of the four distinct synthetic peptides in chronic HCV patients. The most commonly
 5 recognized serotype among U.S. patients was HCV-1 (43.5%), while the least reactive was HCV-J (12.9%). However, among Italian patients, the HCV-G83 was the most prevalent serotype (52.0%), and HCV-1 (12.0%) was least commonly seen in these patients. The most commonly
 10 recognized HCV variant in U.S. plasma donors positive for other HCV antibodies was HCV-1. Conversely, the HCV-1 variant was the least recognized variant among Japanese donors. The HCV-G83 variant was recognized most frequently in Japanese donors, and was nearly as reactive as HCV-1 in the U.S. donors as well.

15 Seroconversion to at least one of these serotypes has been observed in the acute phase of several HCV infected patients. Loss of antibody signal to one serotype and the concurrent seroconversion to a second serotype has been observed in one patient.

20 Among the group of U.S. plasma donors were three specimens which had detectable antibody to all four variant peptides used, suggesting the presence of at least one conserved epitope within the 380-436 region. Since the sequences within the second half (a.a. 408-436) of this region are more conserved among the four virus isolates, this peptide was
 25 prepared based on the HCV-JH-1 and HCV-J sequences which are identical in this region. All three of these specimens were also reactive to this smaller, more conserved region, which demonstrated the existence of a conserved epitope within these boundaries.

30
 TABLE 13
 HCV SEROTYPE ANALYSIS

Population	Reactivity to Peptide From Isolate:				TOTAL
	HCV-1	G83	JH-1	J	
35 US Chronics N=62	29/62 (43.5%)	20/62 (32.3%)	17/62 (27.4%)	8/62 (12.9)	36/62 (58.1%)

36

	Unique Reactives	1/36 (2.8%)	3/36 (8.3%)	4/36 (11.1%)	0/36	8/36 (22.2%)
<hr/>						
5	Italy Chronics N=50	6/50 (12.0%)	26/50 (52.0%)	19/50 (38.0%)	8/47* (17.0%)	34/60 (68.0%)
	Unique Reactives	1/34 (2.9%)	11/34 (32.4%)	6/34 (17.7%)	1/30* (3.3%)	19/34 (58.8%)
<hr/>						
10	U.S. HCV Positive Plasma Donors N=55	16/55 (29.1%)	12/55 (21.8%)	10/55 (18.2%)	5/48* (10.4%)	22/55 (40.0%)
15	Unique Reactives	6/22 (27.3%)	5/22 (22.7%)	1/22 (4.6%)	0/17* (0.0%)	12/22 (54.6%)
<hr/>						
	Japan HCV Positive Plasma Donors N=27	1/27 (3.7%)	13/27 (48.2%)	9/27 (33.3%)	4/27 (14.8%)	18/27 (68.7%)
20	Unique Reactants	0/18	5/18 (27.6%)	2/18 (11.1%)	3/18 (16.7%)	10/18 (55.6%)
<hr/>						

* Some samples depleted.

25

Our data thus demonstrates the existence of at least two epitopes within the hypervariable region of HCV E2/NS1. We conclude that genotypic variability in this region resulted in antigenically distinct HCV variants. Unique E2/NS1 antibody responses can be found in patients with acute and chronic phase HCV infections, as well as in plasma donors from different parts of the world. Our preliminary data thus points to the possibility of geographic variability with regard to antibody responses to HCV as well. It is likely that a high proportion of patients with chronic HCV infection have antibody to the hypervariable region of E2/NS1, since we detected 58% and 68% of U.S. and Italian chronic patients, respectively, as reactive to this region using sequences from only four HCV isolates. The implications for HCV vaccine development are significant if virus neutralizing epitopes exist within this hypervariable region.

35

It is envisioned that these peptides may be used for the development of unique polyclonal and monoclonal antibodies. The polypeptides of this invention, along with those described in U.S. Patent Application Serial Nos. 5 07/456,162 and 07/610,180 (both of which were previously incorporated herein by reference) can be used in combination to develop unique assays for the detection of various epitopes of HCV. This combination of polypeptides can be accomplished, for example, by coating a solid surface with more than one peptide, or by utilizing a mixture of solid supports, 10 wherein the mixture comprises a unique blend of two or more polypeptides to HCV. These techniques are known to the routineer. Other variations of applications and modifications of the specific embodiments of the invention as set forth herein will be apparent to those skilled in the art. Accordingly, the invention is intended to be limited only in accordance with the 15 appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: LESNIEWSKI, RICHARD R.
LEUNG, TAT K.
- (ii) TITLE OF INVENTION: HEPATITIS C ASSAY
- (iii) NUMBER OF SEQUENCES: 27
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: ABBOTT LABORATORIES CHAD377/AP6D
 - (B) STREET: ONE ABBOTT PARK ROAD
 - (C) CITY: ABBOTT PARK
 - (D) STATE: ILLINOIS
 - (E) COUNTRY: U.S.A.
 - (F) ZIP: 60064-3500
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: POREMBSKIP, PRISCILLA E.
 - (B) REGISTRATION NUMBER: 33,207
 - (C) REFERENCE/DOCKET NUMBER: 4767.P3.03
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 708-937-6365
 - (B) TELEFAX: 708-937-9556

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 75 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

39

Met	Ser	Thr	Asn	Pro	Lys	Pro	Gln	Lys	Lys	Asn	Lys	Arg	Asn	Thr	Asn
1				5				10						15	
Arg	Arg	Pro	Gln	Asp	Val	Lys	Phe	Pro	Gly	Gly	Gly	Gln	Ile	Val	Gly
			20					25					30		
Gly	Val	Tyr	Leu	Leu	Pro	Arg	Arg	Gly	Pro	Arg	Leu	Gly	Val	Arg	Ala
		35					40					45			
Thr	Arg	Lys	Thr	Ser	Glu	Arg	Ser	Gln	Pro	Arg	Gly	Arg	Arg	Gln	Pro
	50					55					60				
Ile	Pro	Lys	Ala	Arg	Arg	Pro	Glu	Gly	Arg	Thr					
65					70					75					

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 41 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Tyr	Leu	Leu	Pro	Arg	Arg	Gly	Pro	Arg	Leu	Gly	Val	Arg	Ala	Thr	Arg
1				5				10						15	
Lys	Thr	Ser	Glu	Arg	Ser	Gln	Pro	Arg	Gly	Arg	Arg	Gln	Pro	Ile	Pro
			20					25					30		
Lys	Ala	Arg	Arg	Pro	Glu	Gly	Arg	Thr							
		35					40								

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 28 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Ser	Pro	Arg	Gly	Ser	Arg	Pro	Ser	Trp	Gly	Pro	Thr	Asp	Pro	Arg	Arg
1				5				10						15	

40

Arg Ser Arg Asn Leu Gly Lys Val Ile Asp Thr Leu
 20 25

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 68 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Arg Asn Ser Thr Gly Leu Tyr His Val Thr Asn Asp Cys Pro Asn Ser
 1 5 10 15
 Ser Ile Val Tyr Glu Ala Ala Asp Ala Ile Leu His Thr Pro Gly Cys
 20 25 30
 Val Pro Cys Val Arg Glu Gly Asn Ala Ser Arg Cys Trp Val Ala Met
 35 40 45
 Thr Pro Thr Val Ala Thr Arg Asp Gly Lys Leu Pro Ala Thr Gln Leu
 50 55 60
 Arg Arg His Ile
 65

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Val Arg Glu Gly Asn Ala Ser Arg Cys Trp Val Ala Met Thr Pro Thr
 1 5 10 15
 Val Ala Thr Arg Asp Gly Lys Leu Pro Ala Thr Gln Leu Arg Arg His
 20 25 30
 Ile

41

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 49 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ala	Val	Asp	Phe	Ile	Pro	Val	Glu	Asn	Leu	Glu	Thr	Thr	Met	Arg	Ser
1				5					10					15	
Pro	Val	Phe	Thr	Asp	Asn	Ser	Ser	Pro	Pro	Val	Val	Pro	Gln	Ser	Phe
			20					25					30		
Gln	Val	Ala	His	Leu	His	Ala	Pro	Thr	Gly	Ser	Gly	Lys	Ser	Thr	Lys
		35					40					45			
Val															

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 18 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Phe	Gln	Val	Ala	His	Leu	His	Ala	Pro	Thr	Gly	Ser	Gly	Lys	Ser	Thr
1				5					10					15	
Lys	Val														

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 52 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

Tyr Val Pro His Pro Asn Ile Glu Glu Val Ala Leu Ser Thr Thr Gly
1           5           10           15
Glu Ile Pro Phe Tyr Gly Lys Ala Ile Pro Leu Glu Val Ile Lys Gly
          20           25           30
Gly Arg His Leu Ile Phe Cys His Ser Lys Lys Lys Cys Asp Glu Leu
          35           40           45
Ala Ala Lys Leu
          50

```

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 39 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

Arg Gly Leu Asp Val Ser Val Ile Pro Thr Ser Gly Asp Val Val Val
1           5           10           15
Val Ala Thr Asp Ala Leu Met Thr Gly Tyr Thr Gly Asp Phe Asp Ser
          20           25           30
Val Ile Asp Cys Asn Thr Cys
          35

```

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 25 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```

Asp Ala His Phe Leu Ser Gln Thr Lys Gln Ser Gly Glu Asn Leu Pro

```

43

1	5	10	15
Tyr	Leu	Val	Ala
		Tyr	Gln
		Ala	Thr
		Val	
	20	25	

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 67 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Gly	Arg	Val	Val	Leu	Ser	Gly	Lys	Pro	Ala	Ile	Ile	Pro	Asp	Arg	Glu
1				5				10					15		

Val	Leu	Tyr	Arg	Glu	Phe	Asp	Glu	Met	Glu	Glu	Cys	Ser	Gln	His	Leu
				20				25					30		

Pro	Tyr	Ile	Glu	Gln	Gly	Met	Met	Leu	Ala	Glu	Gln	Phe	Lys	Gln	Lys
				35				40					45		

Ala	Leu	Gly	Leu	Leu	Gln	Thr	Ala	Ser	Arg	Gln	Ala	Glu	Val	Ile	Ala
				50			55				60				

Pro	Ala	Val
65		

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 117 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Ser	Gly	Lys	Pro	Ala	Ile	Ile	Pro	Asp	Arg	Glu	Val	Leu	Tyr	Arg	Glu
1				5				10					15		

Phe	Asp	Glu	Met	Glu	Glu	Cys	Ser	Gln	His	Leu	Pro	Tyr	Ile	Glu	Gln
				20				25					30		

Gly	Met	Met	Leu	Ala	Glu	Gln	Phe	Lys	Gln	Lys	Ala	Leu	Gly	Leu	Leu
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

44

35	40	45
Gln Thr Ala Ser Arg Gln Ala Glu Val Ile Ala Pro Ala Val Gln Thr		
50	55	60
Asn Trp Gln Lys Leu Glu Thr Phe Trp Ala Lys His Met Trp Asn Phe		
65	70	75
Ile Ser Gly Ile Gln Tyr Leu Ala Gly Leu Ser Thr Leu Pro Gly Asn		
85	90	95
Pro Ala Ile Ala Ser Leu Met Ala Phe Thr Ala Ala Val Thr Ser Pro		
100	105	110
Leu Thr Thr Ser Gln		
115		

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 42 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Ile Ile Pro Asp Arg Glu Val Leu Tyr Arg Glu Phe Asp Glu Met Glu	
1	15
Glu Cys Ser Gln His Leu Pro Tyr Ile Glu Gln Gly Met Met Leu Ala	
20	30
Glu Gln Phe Lys Gln Lys Ala Leu Gly Leu	
35	40

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 65 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Phe Lys Ile Met Ser Gly Glu Val Pro Ser Thr Glu Asp Leu Val Asn

45

1	5	10	15												
Leu	Leu	Pro	Ala	Ile	Leu	Ser	Pro	Gly	Ala	Leu	Val	Val	Gly	Val	Val
		20						25					30		
Cys	Ala	Ala	Ile	Leu	Arg	Arg	His	Val	Gly	Pro	Gly	Glu	Gly	Ala	Val
		35					40					45			
Gln	Trp	Met	Asn	Arg	Leu	Ile	Ala	Phe	Ala	Ser	Arg	Gly	Asn	His	Val
		50					55				60				
Ser															
65															

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 32 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Ala	Ala	Ile	Leu	Arg	Arg	His	Val	Gly	Pro	Gly	Glu	Gly	Ala	Val	Gln
1				5					10					15	
Trp	Met	Asn	Arg	Leu	Ile	Ala	Phe	Ala	Ser	Arg	Gly	Asn	His	Val	Ser
			20					25					30		

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 57 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Gly	Val	Asp	Ala	Glu	Thr	His	Val	Thr	Gly	Gly	Ser	Ala	Gly	His	Thr
1				5					10					15	
Val	Ser	Gly	Phe	Val	Ser	Leu	Leu	Ala	Pro	Gly	Ala	Lys	Gln	Asn	Val
			20					25					30		

46

Gln Leu Ile Asn Thr Asn Gly Ser Trp His Leu Asn Ser Thr Ala Leu
 35 40 45

Asn Cys Asn Asp Ser Ser Asn Thr Gly
 50 55

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Gly Val Asp Ala Ala Thr Tyr Thr Thr Gly Gly Ser Val Ala Arg Thr
 1 5 10 15

Thr His Gly Phe Ser Ser Leu Phe Ser Gln Gly Ala Lys Gln Asn Ile
 20 25 30

Gln Leu Ile Asn Thr Asn Gly Ser Trp His Ile Asn Arg Thr Ala Leu
 35 40 45

Asn Cys Asn Ala Ser Leu Asp Thr Gly
 50 55

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 37 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Phe Asn Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Pro Leu
 1 5 10 15

Thr Asp Phe Asp Gln Gly Trp Gly Pro Ile Ser Tyr Ala Asn Gly Ser
 20 25 30

Gly Pro Asp Gln Arg
 35

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Cys	Leu	Val	Asp	Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr	Pro	Cys	Thr	Ile
1				5				10					15		
Asn Tyr Thr Ile Phe															
20															

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Ala	Cys	Asn	Trp	Thr	Arg	Gly	Glu	Arg	Cys	Asp	Leu	Glu	Asp	Arg	Asp
1				5				10					15		
Arg Ser Glu Leu Ser Tyr															
20															

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Ala Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp

48

1	5	10	15
Arg	Ser	Glu	Leu
Ser	Pro	Leu	Leu
Leu	Leu	Thr	Thr
Thr	Thr	Gln	Trp
Gln	Val		
20	25	30	
Leu	Pro	Cys	Ser
Phe	Thr	Thr	Leu
Pro			
35	40		

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 19 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Leu	Leu	Thr	Thr	Thr	Gln	Trp	Gln	Val	Leu	Pro	Cys	Ser	Phe	Thr	Thr
1			5				10							15	
Leu Pro Tyr															

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

His	Leu	His	Gln	Asn	Ile	Val	Asp	Val	Gln	Tyr	Leu	Tyr	Gly	Val	Gly
1			5						10					15	
Ser Ser Ile Ala Ser Trp Ala Ile															
20															

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 51 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Lys	Lys	Pro	Asp	Tyr	Gln	Pro	Pro	Val	Val	His	Gly	Cys	Pro	Leu	Pro
1				5					10					15	
Pro	Pro	Lys	Ser	Pro	Pro	Val	Pro	Pro	Pro	Lys	Lys	Lys	Arg	Thr	Val
			20					25					30		
Val	Leu	Thr	Glu	Ser	Thr	Leu	Ser	Thr	Ala	Leu	Ala	Glu	Leu	Ala	Thr
			35				40					45			
Arg	Ser	Phe													
			50												

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 57 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Gly	Val	Asp	Gly	His	Thr	Arg	Val	Thr	Gly	Gly	Val	Gln	Gly	His	Val
1				5					10					15	
Thr	Ser	Thr	Leu	Thr	Ser	Leu	Phe	Arg	Pro	Gly	Ala	Ser	Gln	Lys	Ile
			20					25					30		
Gln	Leu	Val	Asn	Thr	Asn	Gly	Ser	Thr	His	Ile	Asn	Arg	Thr	Ala	Leu
			35				40					45			
Asn	Ser	Asn	Asp	Ser	Leu	Gln	Thr	Gly							
			50				55								

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 57 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Gly	Val	Asp	Gly	His	Thr	His	Val	Thr	Gly	Gly	Arg	Val	Ala	Ser	Ser	
1				5					10					15		
Thr	Gln	Ser	Leu	Val	Ser	Trp	Leu	Ser	Gln	Gly	Pro	Ser	Gln	Lys	Ile	
			20					25					30			
Gln	Leu	Val	Asn	Thr	Asn	Gly	Ser	Gln	His	Ile	Asn	Arg	Thr	Ala	Leu	
			35				40					45				
Asn	Cys	Asn	Asp	Ser	Leu	Gln	Thr	Gly								
			50				55									

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Ser	Gln	Lys	Ile	Gln	Leu	Val	Asn	Thr	Asn	Gly	Ser	Trp	His	Ile	Asn	
1				5					10					15		
Arg	Thr	Ala	Leu	Asn	Cys	Asn	Asp	Ser	Leu	Gln	Thr	Gly				
			20					25								

CLAIMS

1. An assay for identifying the presence of an antibody immunologically reactive with an HCV antigen in a fluid sample comprising:
5 contacting the sample with a polypeptide containing at least one epitope of an HCV antigen selected from the group consisting of p380-JH1, p380-J, p-380.LG, and p408 under conditions suitable for complexing the antibody with the polypeptide; and detecting the antibody-polypeptide complex as an indication of chronic HCV infection.
10
- 2.. The assay of claim 1 wherein the polypeptide is bound to a solid support.
3. A combination assay for the presence of an antibody
15 immunologically reactive with an HCV antigen in a fluid sample comprising contacting the sample with a solid support having bound thereto at least one polypeptide selected from the group consisting of p380-JH1, p380-J, p-380.LG, and p408 containing at least one epitope of an HCV antigen, under conditions suitable for complexing the antibody with the
20 polypeptide; and detecting the antibody-polypeptide complex as an indication of chronic HCV infection.
4. The assay of claim 3 wherein the solid support is a
25 polystyrene bead.
5. In a confirmatory assay for identifying the presence of an antibody in a fluid sample immunologically reactive with an HCV antigen wherein the sample is used to prepare first and second aliquots and the first
30 aliquot is contacted with a first polypeptide selected from the group consisting of p380-JH1, p380-J, p-380.LG, and p408 which contains at least one epitope of an HCV antigen under conditions suitable for complexing the antibody with the polypeptide and wherein the first antibody-antigen complex is detected;
35 contacting the second aliquot with a second polypeptide selected from the group consisting of p380-JH1, p380-J, p-380.LG, and p408 under

conditions suitable to form a second antibody-antigen complex, wherein said second polypeptide is different from the first polypeptide; and detecting the second antibody-antigen complex.

5 6. The assay of claim 5 wherein the antigens are bound to a solid support.

 7. The assay of claim 6 wherein said solid support is a polystyrene bead.

10

 8. An immunodot assay for identifying the presence of an antibody immunologically reactive with an HCV antigen in a fluid sample comprising

 concurrently contacting the sample with at least two polypeptides
15 selected from the group consisting of p380-JH1, p380-J, p-380.LG, and p408, each containing distinct epitopes of an HCV antigen and coated onto a solid support, under conditions suitable for complexing the antibody with the polypeptides; and

 detecting the antibody-polypeptide by reacting the complex with
20 color-producing reagents.

 9. A competition assay for identifying the presence of an antibody immunologically reactive with an HCV antigen in a fluid sample wherein the sample is used to prepare first and second immunologically
25 equivalent aliquots comprising

 contacting a first aliquot of test sample with a solid support coated with a polypeptide that contains at least one epitope of an HCV antigen under conditions suitable for complexing with the antibody to form a detectable antibody-polypeptide complex;

30 contacting the second aliquot with unbound polypeptide and then contacting said second aliquot with a solid support coated with at least one polypeptide selected from the group consisting of p380-JH1, p380-J, p-380.LG, and p408;

 detecting the presence of bound or unbound polypeptide.

35

 10. An immunoassay kit comprising:

a polypeptide containing at least one epitope of an HCV antigen selected from the group consisting of p380-JH1, p380-J, p-380.LG, and p408;

- 5 one or more sample preparation reagents; and
 one or more detection and signal producing reagents.

11. The kit of claim 10 wherein the polypeptides are coated on a solid support.




FIG.1a

2/7




FIG.1b

3/7




FIG. 2

SUBSTITUTE SHEET




FIG.3a

5/7




FIG.3b

SUBSTITUTE SHEET




FIG. 4

7/7




FIG.5

INTERNATIONAL SEARCH REPORT

 International application No.
PCT/US92/07813

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C12Q 1/70

US CL : 435/5

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/5; 435/518, 820

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, Dialog

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP, A 0.419,182 (Miyamura et al.) 27 March 1991, see page 2, last paragraph; page 3, fourth paragraph; Figure 3.	1-11
Y	GB, A, 2,099,578 (Gordon et al.) 08 December 1982, see page 11, Example 2.	8
Y	Science, Volume 244, issued 21 April 1989, G. Kuo et al., "An Assay for Circulating Antibodies to a Major Etiologic Virus of Human Non-A, Non-B Hepatitis," pages 262-264, entire document.	1-11
Y	Biochemical and Biophysical Research Communications, Volume 175, No. 1, issued 28 February 1991, M. Hijikata et al., "Hypervariable Regions in the Putative Glycoprotein of Hepatitis C Virus, pages 220-228, especially page 224.	1-11
Y	Journal of General Virology, Volume 72, issued 1991, D. Kremsdorf et al., "Partial Nucleotide Sequence Analysis of a French Hepatitis C Virus: Implications for HCV Genetic Variability", pages 2557-2561, especially page 2559, Figure 2(b).	1-11

☐

Further documents are listed in the continuation of Box C.

☐

See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be part of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

08 December 1992

Date of mailing of the international search report

16 DEC 1992

 Name and mailing address of the ISA/
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

DONNA C. WORTMAN

Telephone No. (703) 308-0196