



(86) Date de dépôt PCT/PCT Filing Date: 1999/02/26
(87) Date publication PCT/PCT Publication Date: 1999/09/02
(45) Date de délivrance/Issue Date: 2014/07/15
(85) Entrée phase nationale/National Entry: 2000/08/25
(86) N° demande PCT/PCT Application No.: US 1999/004438
(87) N° publication PCT/PCT Publication No.: 1999/043711
(30) Priorité/Priority: 1998/02/27 (US60/076,105)


(51) Cl.Int./Int.Cl. *C07K 14/715* (2006.01),
A61K 38/19 (2006.01), *A61K 38/00* (2006.01)

(72) Inventeurs/Inventors:
TARASOVA, NADYA I., US;
MICHEJDA, CHRISTOPHER J., US

(73) Propriétaire/Owner:
THE GOVERNMENT OF THE UNITED STATES OF
AMERICA, REPRESENTED BY THE SECRETARY
OF THE DEPARTMENT OF HEALTH AND HUMAN
SERVICES, US

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : ANTAGONISTES DES RECEPTEURS COUPLES A LA PROTEINE G
(54) Title: G PROTEIN-COUPLED RECEPTOR ANTAGONISTS



(57) Abrégé/Abstract:

G-protein coupled receptors (GPCR) generally contain seven transmembrane helices. The present invention provides synthetic peptides derived from these transmembrane helices. The peptides inhibit GPCR function by disrupting GPCR structure. In certain embodiments, charged residues are added at one terminus to promote correct orientation of the peptide in the membrane.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/715, A61K 38/19	A1	(11) International Publication Number: WO 99/43711 (43) International Publication Date: 2 September 1999 (02.09.99)
---	-----------	---

(21) International Application Number: PCT/US99/04438

(22) International Filing Date: 26 February 1999 (26.02.99)

(30) Priority Data:
60/076,105 27 February 1998 (27.02.98) US

(71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Bethesda, MD 20892 (US).

(72) Inventors; and


(75) Inventors/Applicants (for US only): TARASOVA, Nadya I. [RU/US]; 103 Pipe Meadow Way, Frederick, MD 21702 (US). MICHEJDA, Christopher, J. [US/US]; 13814 Hidden Glen Lane, Potomac, MD 20878 (US).

(74) Agents: WEBER, Kenneth, A. et al.; Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111-3834 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published*With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

(54) Title: G PROTEIN-COUPLED RECEPTOR ANTAGONISTS



(57) Abstract

G-protein coupled receptors (GPCR) generally contain seven transmembrane helices. The present invention provides synthetic peptides derived from these transmembrane helices. The peptides inhibit GPCR function by disrupting GPCR structure. In certain embodiments, charged residues are added at one terminus to promote correct orientation of the peptide in the membrane.

5

G Protein-Coupled Receptor Antagonists

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 60/076,105 filed 2/27/98.

10

FIELD OF THE INVENTION

The present invention relates to modulating, especially inhibiting, biological activities of G protein coupled receptors (GPCRs) by exposing GPCRs to molecules which interfere with correct receptor assembly. In particular, the invention relates to synthetic, isolated and/or recombinant peptides, fragments and/or consensus peptides of the transmembrane domain of GPCRs that inhibit GPCR-mediated signal transduction.

15

BACKGROUND OF THE INVENTION

20

Many physiologically important events are mediated by the binding of guanine nucleotide-binding regulatory proteins (G proteins) to G protein-coupled receptors (GPCRs). These events include vasodilation, stimulation or decrease in heart rate, bronchodilation, stimulation of endocrine secretions and enhancement of gut peristalsis, development, mitogenesis, cell proliferation and oncogenesis.

25

G proteins are a diverse superfamily of guanine nucleotide-binding proteins that play a central role in signal transduction and regulation of cellular metabolism. They are generally comprised of three subunits: a guanyl-nucleotide binding alpha subunit; a beta subunit; and a gamma subunit. (For a review, see Conklin et al. *Cell* 73, 631-641, (1993)). G proteins commonly cycle between two forms, depending on whether GDP or GTP is bound to the alpha subunit. When GDP is bound, the G protein exists as a heterotrimer, the G alpha-beta-gamma complex. When an alpha-beta-gamma complex operatively associates with a ligand-activated GPCR in a cell membrane, the rate of exchange of GTP for bound GDP is increased and the G alpha subunit dissociates from the G beta-gamma complex. The

30

free G alpha subunit and G beta-gamma complex are capable of transmitting a signal to downstream elements of a variety of signal transduction pathways, for example by binding to and activating adenyl cyclase. This fundamental scheme of events forms the basis for a multiplicity of different cell signaling phenomena.

5 Recent studies have suggested that all members of the GPCR superfamily have a conserved structure. Comparisons of avian and mammalian beta-adrenergic receptor cDNA's (Yarden *et al.*, *Proc. Natl. Acad. Sci. USA* 83: 6795-6799, 1986; Dixon *et al.*, *Nature* 321:75-79, 1986; and Kobilka *et al.*, *Proc. Natl. Acad. Sci. USA* 84:46-50, 1987), a bovine rhodopsin cDNA (Nathans and
10 Hogness, *Cell* 34:807-814, 1983), an alpha 2-adrenergic receptor (Kobilka *et al.*, *Science* 238:650-656, 1987), an angiotensin receptor cDNA (Young *et al.*, *Cell* 45:711-719, 1986; Jackson *et al.*, *Nature* 335:437-439, 1988), a bovine substance K receptor (Masu *et al.*, *Nature* 329:836-838, 1987), and a muscarinic acetylcholine receptor cDNA (Kubo *et al.*, *Nature* 323:411-416, 1986) predict that all GPCR share
15 a highly conserved presence of seven hydrophobic transmembrane domains that are suggested to be transmembrane helices of 20-30 amino acids connected by extracellular or cytoplasmic loops. Kobilka *et al.*, *Science* 240: 1310 (1988); Maggio *et al.*, *FEBS Lett.* 319: 195 (1993); Maggio *et al.*, *Proc. Natl. Acad. Sci USA* 90: 3103 (1993); Ridge *et al.*, *Proc. Natl. Sci USA* 91, 3204 (1995); Schonenberg *et al.*, *J. Biol. Chem.* 270: 18000 (1995); Huang *et al.*, *J. Biol. Chem.* 256: 3802 (1981); Popot *et al.*, *J. Mol. Biol.* 198: 655 (1987); Kahn and Engelman, *Biochemistry* 31: 6144 (1992); Schoneberg *et al.* *EMBO J.* 15: 1283 (1996); Wong *et al.*, *J. Biol. Chem.* 265: 6219 (1990); Monnot *et al.*, *J. Biol. Chem.* 271: 1507 (1996); Gudermann *et al.*, *Annu. Rev. Neurosci.* 20: 399 (1997); Osuga *et al.*, *J. Biol. Chem.* 272: 25006 (1997); Lefkowitz *et al.*, *J. Biol. Chem.* 263:4993-4996, 1988; Panayotou and Waterfield, *Curr. Opinion Cell Biol.* 1:167-176, 1989. These transmembrane domains of G-protein coupled receptors are designated TM1, TM2, TM3, TM4, TM5, TM6 and TM7. TM4, TM5, TM6 and TM7 are the most highly conserved and are postulated to provide sequences which impart biological activity to GPCRs. TM3 is
20 also implicated in signal transduction.

 The coupling of GPCRs to intracellular signaling molecules such as adenylate cyclase (Anand-Srivastava *et al.*, *J. Biol. Chem.* 271: 19324-19329 (1996)) and G-proteins (Merkouris *et al.*, *Mol. Pharmacol.* 50: 985-993 (1996)) is reportedly

inhibited by peptides corresponding to the intracellular loops of the receptors. Those studies were conducted primarily to provide an understanding of molecular mechanisms of receptor function and could not be applied directly for drug design, because of the difficulties in intracellular delivery of the inhibitors.

5 WO 94/05695 and U.S. Patent No. 5,508,384 set forth sequences of transmembrane regions for 74 GPCRs. The WO 94/05695 patent publication describes and claims polypeptides corresponding to fragments or homologous sequences of GPCRs which can bind a GPCR ligand or which can modulate ligand binding. Both references disclose that a membrane spanning fragment of the third
10 TM domain of the dopamine D₂ receptor specifically bound a ligand of the intact receptor in a simple, small unilamellar vesicle model. The fragment used was terminated with a lysine (which is positively charged at physiological pH) at one end and with an aspartic acid (which is negatively charged at physiological pH) at the other. This peptide would not be expected to insert readily into a biological
15 membrane.

SUMMARY OF THE INVENTION

The invention generally comprises peptide or peptidomimetic compounds that modulate, and preferably inhibit the biological properties and
20 activities of GPCRs, by targeting the transmembrane portions of these receptors. The present invention specifically comprises methods for disrupting GPCR function by using these GPCR antagonists.

The present invention provides for the use of chemical or recombinant DNA technology to obtain GPCR polypeptides, which preferably are as small as
25 possible while still retaining sufficiently high affinity for binding to, or association with, GPCRs. Non-limiting examples of GPCR polypeptides include fragments of 10 to 50 amino acids corresponding to at least one transmembrane domain of domains 1-7. The following are nonlimiting examples of GPCR peptides with antagonist properties.

30 From the GPCR CXCR4

F-2-2: LLFVITLPFWAVDAVANWYFGNDD (SEQ. ID No.: 1)

F-2-5: LLFVITLPFWAVDAVANDD-OH (SEQ. ID No.: 2)

F-4-2: VYVGWIPALLLTIPDFIFANDD-OH (SEQ. ID No.: 3)

F-6-1: VILILAFFACWLPYYIGISID-OH (SEQ. ID No.: 4)

F-7-3: DDEALAFFHCCLNPILYAFL-NH₂ (SEQ. ID No.: 5)

F-7-4: DDSITEALAFFHCCLNPILYAFL-NH₂ (SEQ. ID No.: 6)

5

From the GPCR CCR5

CCR5-TM-2-2: LFFL LTVPFWAHYAAAQWDFGDD (SEQ. ID No.: 7)

CCR5-TM-4-1: FGVVTSVITWVAVFASLPGIIFTSSDD (SEQ. ID No.: 8)

CCR5-TM-6-1: LIFTIMIVYFLFWAPYNIVLLLNTFQED (SEQ. ID No.: 9)

10 CCR5-TM-7-1: DDQAMQVTETLGMTHCCINPIIYAFV (SEQ. ID No.: 10)

From the GPCR CCR2

CCR2-TM-2-1: IYLLNLAISDLLFLITLPLWADD-OH (SEQ. ID No.: 11)

CCR2-TM-2-2: LLFLITLPLWAH SAANEVWFGNDD-OH (SEQ. ID No.: 12)

15 CCR2-TM-4-1: FGVVTSVITWLVAVF ASVPGIIFTDD (SEQ. ID No.: 13)

CCR2-TM-6-1: VIFTIMIVYFLFWTPYN IVILLNTFQED (SEQ. ID No.: 14)

CCR2-TM-7-1: DDATQVT ETLGMTHCCINPIIYAFV (SEQ. ID No.: 15)

From the GPCR CCR3

20 CCR3-TM-2-1: LLFLVTLPFW IHYVRGHNWVFGDDD (SEQ. ID No.: 16)

CCR3-TM-4-1: FGVITSIVTWGLAVLAALPEFI FYETED (SEQ. ID No.: 17)

CCR3-TM-6-1: IFVIMAVFFI FWTPYNVAILLSSYQSDD (SEQ. ID No.: 18)

CCR3-TM-7-1: DDLVMLVTEVIAYSHCCMNPVIYAFV (SEQ. ID No.: 19)

25 From the GPCR CCKAR

CCKAR-TM-1-6: DDEWQSALQILLYSIIFLLSVLGNTLVITV (SEQ. ID No.: 20)

CCKAR-TM-2-1: FLLSLAVSDLMLCLFCMPFNLP (SEQ. ID No.: 21)

CCKAR-TM-2-2: FLLSLAVSDLMLCLFCM PFNLIDD (SEQ. ID No.: 22)

CCKAR-TM-6-4: IVVLFFLCWMPIFSANAWRAYDTVDD (SEQ. ID No.: 23)

30

One embodiment of the invention is an isolated G protein-coupled receptor (GPCR)-modulating molecule comprising a peptide or peptidomimetic that is a structural analog of a portion of a transmembrane domain of a GPCR, wherein said

molecule has a first end and a second end and said molecule has at said first end a negatively charged group and at said second end a neutral charge under physiological conditions; said molecule spontaneously inserts into a membrane in the same orientation as the transmembrane domain from which it is derived; and said molecule modulates a biological property or activity of said GPCR.

In a particular embodiment, the molecules contain a hydrophilic, negatively charged non-peptidic head group and an uncharged tail, which assures correct orientation of the molecule in the cell membrane. In another embodiment, the negatively charged head group is one or more acidic amino acids.

Another embodiment is an isolated GPCR-modulating molecule comprising a peptide or peptidomimetic that is a structural analog of a portion of a transmembrane domain of CXCR4, wherein said portion of said transmembrane domain has a sequence selected from the group of sequences consisting of:

LLFVITLPFWAVDAVANWYFGNDD, (SEQ. ID No.: 1)

LLFVITLPFWAVDAVANDD-OH, (SEQ. ID No.: 2)

VYVGWIPALLLTIPDFIFANDD-OH, (SEQ. ID No.: 3)

VILILAFFACWLPPYYIGISID-OH, (SEQ. ID No.: 4)

DDEALAFFHCCLNPILYAFL-NH₂, (SEQ. ID No.: 5)

DDSITEALAFFHCCLNPILYAFL-NH₂, (SEQ. ID No.: 6)

wherein said molecule modulates a biological activity of said CXCR4. The CXCR4 activity modulated by said peptide includes inhibition of CXCR4-mediated intracellular Ca²⁺ release and inhibition of CXCR4-mediated HIV infection.

The invention also comprises methods of modulating the biological activity of a target GPCR by contacting a cell that expresses said GPCR with a molecule of the invention. In one method, the target GPCR is CXCR4, CCR5 or CCR2, and the modulated biological activity is inhibition of GPCR-mediated HIV infection. In another method, the target GPCR is CXCR4 and the modulated biological activity is inhibition of CXCR4-mediated intracellular Ca²⁺ release.

Another embodiment is a method of inhibiting HIV-1 infection, comprising contacting a cell that expresses a GPCR that binds HIV-1 with a molecule that comprises a peptide or peptidomimetic that is a structural analog of a portion of

a transmembrane domain of said GPCR, wherein contacting the cell with said molecule inhibits HIV-1 infection. The peptide or peptidomimetic may be a structural analog of a portion of a transmembrane domain of CXCR4 or CCR5. Peptides corresponding to TM regions of CXCR4, a GPCR that functions as a co-receptor during the cell entry of HIV, were designed and tested in cells, and yielded potent inhibition of HIV entry without apparent toxicity to the cells.

The usefulness of the method is demonstrated by specifically targeting the CXCR4 that functions as a co-receptor during the cell entry of T-cell tropic strains of HIV-1. Peptides containing 20-25 amino acid residues inhibited receptor signaling and HIV-1 infection *in vitro* at concentration as low as 0.2 micromolar.

In one embodiment, the molecules of the present invention mimic a transmembrane domain of the chosen receptor and block self-assembly of that receptor, possibly by competitive inhibition with the native TM domain. They thereby block or inhibit signal transduction in the affected cell.

The invention also includes peptide analogs and peptidomimetics which possess beneficial properties such as increased half-life, lack of immunogenicity, and the ability to cross the blood-brain barrier.

The peptide analogs of the invention mediate the chemical and/or biological effects of hormone agonists/antagonists or other peptides. They are believed to be useful for the development of pharmaceutical, therapeutic, and diagnostic techniques. Accordingly, the invention also provides methods for producing a prophylactic or therapeutic response in a mammal by administering to the mammal a pharmaceutically effective amount of one or more peptide analogs of the invention. In preferred embodiments, the present invention provides methods for producing such responses by modulating the activity of at least one mammalian G-protein-linked receptor by administering an effective amount of one or more peptide analogs of the invention.

In another embodiment, a peptide of the invention may modulate the biological activity of more than one GPCR. In another embodiment, more than one peptide of the invention are administered as a cocktail to modulate the biological activity of more than one GPCR.

BRIEF DESCRIPTION OF FIGURES

Figure 1. Anti-HIV efficacy and toxicity of F-2-2 assay. CEM-SS cells were infected with the LAV strain of HIV-1.

5 **Figure 2.** Anti-HIV efficacy and toxicity of F-4-2 in cytoprotection. CEM-SS cells were infected with the RF strain of HIV-1, which causes cell death, if the inhibitor of infection is not present.

10 **Figure 3.** The proposed model of transmembrane antagonists action.

DEFINITIONS

A "G-protein" is any member of the superfamily of signal transducing guanine nucleotide binding proteins.

15 A "G-protein-coupled receptor" is any member of a superfamily of receptors that mediates signal transduction by coupling with a G protein. Examples of such receptors include, but are not limited to: CC chemokine receptor 5 (CCR5), CXC chemokine receptor (CXCR4) cholecystokinin type A receptor (CCKAR), adenosine receptors, somatostatin receptors, dopamine receptors, muscarinic cholinergic receptors, alpha-adrenergic receptors, beta-adrenergic receptors, opiate
20 receptors, cannabinoid receptors, growth hormone releasing factor, glucagon, cAMP receptors, serotonin receptors (5-HT), histamine H2 receptors, thrombin receptors, kinin receptors, follicle stimulating hormone receptors, opsins and rhodopsins, odorant receptors, cytomegalovirus GPCRs, histamine H2 receptors, octopamine
25 receptors, N-formyl receptors, anaphylatoxin receptors, thromboxane receptors, IL-8 receptors, platelet activating factor receptors, endothelin receptors, bombesin gastrin releasing peptide receptor, neuromedin B preferring bombesin receptors, vasoactive intestinal peptide receptors, neurotensin receptors, bradykinin receptors, thyrotropin-releasing hormone receptors, substance P receptors, neuromedin K
30 receptors, renal angiotensin II type I receptors, mas oncogene (angiotensin) receptors, lutropin-choriogonadotropin receptors, thyrotropin receptors, follicle stimulating hormone receptors, cannabinoid receptors, glucocorticoid-induced receptors, endothelial cell GPCRs, testis GPCRs, and thoracic aorta GPCRs, and homologs thereof having a homology of at least 80% with at least one of transmembrane

domains 1-7, as described herein. See, e.g., Probst et al, *DNA and Cell Biology* 11: 1-20 (1992).

The term further encompasses subtypes of the named receptors, and mutants and homologs thereof, along with the DNA sequences encoding the same.

5 The term "membrane" refers generally to a lipid bilayer. Preferably, the lipid bilayer is the plasma membrane that delimits a cell, but may be any cellular membrane. The term membrane also encompasses bilayer structures, such as artificial liposomes.

10 The term "GPCR polypeptide" includes polypeptides having an amino acid sequence which substantially corresponds to at least one 10 to 50 (e.g., 10, 20, 25 30 residues) amino acid fragment and/or homologous sequence of a known GPCR or group of GPCRs, wherein the GPCR polypeptide has homology of at least 80%, such as 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92 93, 94, 95, 96, 97, 98, 99 or 100% homology, while maintaining GPCR modulating activity, wherein a GPCR
15 polypeptide of the present invention is not naturally occurring or is naturally occurring but is in a purified or isolated form which does not occur in nature.

 Preferably, a GPCR polypeptide of the present invention substantially corresponds to a transmembrane domain of a GPCRs. Also preferred are GPCR polypeptides wherein the GPCR amino acid sequence is 4-10 to 50 amino acids in
20 length, such as 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 amino acids, or any range therein.

 The term "spontaneously inserts into a membrane" means that a peptide that is brought into contact with a membrane will, under physiological
25 conditions, arrange itself within the lipid bilayer such that the hydrophobic portion of the peptide is within the membrane, and any charged end is exposed to either surface of a membrane. Preferably, molecules of the present invention that have a net negative charge at one end will orient themselves so that the charged end faces the extracellular surface of the cell.

30 The term "tumor cell" or "cancer cell" or "neoplastic cell" denotes a cell that demonstrates inappropriate, unregulated proliferation. A cell line is said to be "malignant" if, when the cell line is injected into a host animal, the host animal develops tumors or cancers that are anaplastic, invasive, and/or metastatic. A

"human" tumor is comprised of cells that have human chromosomes. Such tumors include those in a human patient, and tumors resulting from the introduction of a human malignant cell line into a non-human host animal if cells from such tumors have human chromosomes.

5 The terms "treating cancer", "cancer therapy", and the like mean generally a treatment that causes any improvement in a mammal having a cancer wherein the improvement is due to treatment with a peptide of the invention. The improvement can be either subjective or objective. For example, if the mammal is human, the patient may note improved vigor or vitality or decreased pain as
10 subjective symptoms of improvement or response to therapy. Alternatively, the clinician may notice a decrease in tumor size or tumor burden based on physical exam, laboratory parameters, tumor markers, or radiographic findings.

 The phrase "inhibiting tumor [or cell] growth" generally means that the rate of increase in mass, size, number and/or the metabolism of treated cells and/or
15 tumors is slower as a result of treatment than that of nontreated cells and/or tumors. The growth of a cell line or tumor is said to be "inhibited" by a treatment if, when assayed by means such as radioisotope incorporation into the cells, the treated cells increase in number at a rate that is less than the proliferation rate of untreated control cells, and preferably less than about 50% of the untreated cell proliferation
20 rate. More preferably, the growth rate is inhibited by at least 80%. If growth is assayed by a means such as plating in methylcellulose, the growth of a cell line is said to be "inhibited" if the treated cells give rise to less than the number of colonies that grow from a like number of untreated cells. Preferably, the number of colonies from treated cells is less than about 70% of the number from untreated cells. More
25 preferably, the number of colonies is decreased by at least 50%. "Inhibition of cell growth" also encompasses zero growth and, most importantly, consequent death of the tumor cells and eradication of the tumor. When measured *in vivo*, "inhibition of tumor growth" encompasses fewer or smaller tumors (for example, smaller diameter) as compared to control animals or untreated patients. Progression of a tumor refers
30 to events other than growth, such as morphological and physiological changes, and changes in gene and protein expression.

 Inhibition can be evaluated by any accepted method of measuring whether growth or size of the tumor and/or increase in the number of cancerous or

tumor cells has been slowed, stopped, or reversed. This includes direct observation and indirect evaluation such as subjective symptoms or objective signs. The clinician may notice a decrease in tumor size or tumor burden (number of tumors) based on physical exam, laboratory parameters, tumor markers, or radiographic findings.

5 Alternatively, if the mammal is human, the patient may note improved vigor or vitality or decreased pain as subjective symptoms of improvement or response to therapy. Some laboratory signs that the clinician may observe for response to therapy include normalization of tests such as white blood cell count, red blood cell count, platelet count, erythrocyte sedimentation rate, and various enzyme levels such as
10 transaminases and hydrogenases. Additionally, the clinician may observe a decrease in a detectable tumor marker such as prostatic specific antigen (PSA) or chorio embryonic antigen (CEA). Alternatively, other tests can be used to evaluate objective improvement such as sonograms, computerized axial tomography scans, nuclear magnetic resonance scans and positron emission testing.

15 The term "GPCR transmembrane peptide" can include a GPCR transmembrane domain fragment and/or a homologous peptide thereof, of at least 4-50, and preferably 4-30, and preferably at least 10-30 amino acids in length, such as 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47,
20 48, 49, or 50 amino acids, or any range therein, or any corresponding sequences having conservative amino acid substitutions. Sample transmembrane peptides of the invention include, but are not limited to, the peptides listed in Table I of the present disclosure. A preferred transmembrane peptide of the present invention, when contacted with a cell or membrane structure (e.g., liposome) that contains a
25 biologically active GPCR, modulates the biological activity of said GPCR *in vitro*, *in vivo* or *in situ*. The concentration of the peptide in a solution that contacts the cell *in vivo* (e.g., blood plasma or interstitial fluid) or *in vitro* (e.g., culture medium) is between 1 nanomolar and 50 micromolar, preferably between 1 nanomolar and 1 micromolar, and most preferably less than 5 micromolar.

30 The term "residue" refers to an amino acid or amino acid mimetic incorporated in a oligopeptide by an amide bond or amide bond mimetic.

"Negatively charged" refers to those amino acids, amino acid derivatives, amino acid mimetics and chemical moieties that are negatively charged

at physiological pH. Negatively charged amino acids include, for example Asp and Glu. An "acidic" residue is a residue that is negatively charged at physiological pH.

"Positively charged" refers to those amino acids, amino acid derivatives, amino acid mimetics and chemical moieties that are positively charged at physiological pH. Positively charged amino acids include, for example, Lys and Arg. A "basic residue" is a residue that is positively charged at physiological pH.

"Neutral" refers to those amino acids, amino acid derivatives, amino acid mimetics and chemical moieties that are neither positively nor negatively charged at physiological pH.

"Consensus" sequence refers to peptides which are distinct from known GPCR sequences in critical structural features, but which are derived from consensus sequences of homologous GPCR transmembrane domains 1-7. Such consensus peptides may be derived by molecular modeling, optionally combined with hydrophobicity analysis and/or fitting to model helices, as non-limiting examples. Such modeling can be accomplished according to known method steps using known modeling algorithms, such as, but not limited to, ECEPP, INSIGHT, DISCOVER, CHEM-DRAW, AMBER, FRODO and CHEM-X. Such algorithms compare transmembrane domains between related G-protein coupled receptors, determine probable energy-minimized structures and define alternative consensus polypeptide fragments.

An amino acid or nucleic acid sequence of a GPCR polypeptide of the present invention is said to "substantially correspond" to another amino acid or nucleic acid sequence, respectively, if the sequence of amino acids or nucleic acid in both molecules provides polypeptides having biological activity that is substantially similar, qualitatively or quantitatively, to the corresponding fragment of at least one GPCR transmembrane domain, or which may be synergistic when two or more transmembrane domains, consensus sequences or homologs thereof are present.

Additionally or alternatively, such "substantially corresponding" sequences of GPCR polypeptides include conservative amino acid or nucleotide substitutions, or degenerate nucleotide codon substitutions wherein individual amino acid or nucleotide substitutions are well known in the art.

The term "modulates a biological property or activity" means that in the presence of a test transmembrane peptide a measurable biological parameter or event

is increased or decreased relative to a control in the absence of said peptide. Examples of biological property or activity include: the conformation of the GPCR, association of the GPCR with other molecules, signal transduction, extracellular secretion of cellular proteins, conformational changes in proteins, changes in enzymatic activity, changes in metabolic activity, changes in affinity for a ligand, changes in levels of viral infection, changes in vasodilation, modulation of heart rate, modulation of bronchodilation, modulation of endocrine secretions and modulation of gut peristalsis. Note that the GPCR biological activity need not be one that is limited to the precise *in vivo* role performed by the GPCR. The term also covers GPCR properties, such as viral protein binding, that are not part of the *in vivo* biological role of the GPCR. It further covers intrinsic properties of GPCRs that are only disclosed by experimental manipulation in the laboratory, such as the ability of GPCRs in artificial bilayers (e.g., liposomes) to interact with GPCR ligands.

"Signal transduction" is the process by which binding of a ligand to a receptor is translated into physiological change. In general, binding of a ligand to a receptor causes a change in a physical property of the receptor, for example a change in its conformation, or its orientation, or in its ability to bind other ligands. This change in a physical property can result, directly or indirectly, in increased or decreased ion fluxes, increased or decreased enzymatic activity, increased or decreased phosphorylation, increased or decreased translocation of the receptor or of any molecule (e.g., an inositol moiety or a G protein subunit) from one cellular compartment to another.

"GPCR ligands" refers to biological molecules that bind GPCRs *in vitro*, *in situ* or *in vivo*, and may include hormones, neurotransmitters, viruses or receptor binding domains thereof, G proteins, opsins, rhodopsins, nucleosides, nucleotides, coagulation cascade factors, odorants or pheromones, toxins, colony stimulating factors, platelet activating factors, neuroactive peptides, neurohumor, or any biologically active compounds, such as drugs or synthetic or naturally occurring compounds.

The phrase "inhibits HIV infection" means that a peptide of the invention inhibits binding of an HIV to a GPCR or inhibits a GPCR biological activity that mediates the entry and successful reproduction of an HIV virus into a GPCR-expressing cell.

The term "effective amount" means a dosage sufficient to produce a desired result. The desired result can be subjective or objective changes in the biological activity of a GPCR, especially signal transduction. Effective amounts of the GPCR polypeptide or composition, which may also include a functional derivative thereof, are from about 0.01 micrograms to about 100 mg/kg body weight, and preferably from about 10 micrograms to about 50 mg/kg body weight, such 0.05, 0.07, 0.09, 0.1, 0.5, 0.7, 0.9, 1, 2, 5, 10, 20, 25, 30, 40, 45, or 50 mg/kg.

A "conservative substitution", when describing a protein refers to a change in the amino acid composition of the protein that does not substantially alter the protein's activity. Thus, "conservatively modified variations" of a particular amino acid sequence refers to amino acid substitutions of those amino acids that are not critical for protein activity or substitution of amino acids with other amino acids having similar properties (e.g., acidic, basic, positively or negatively charged, polar or non-polar, etc.) such that the substitutions of even critical amino acids do not substantially alter activity. Conservative substitution tables providing functionally similar amino acids are well known in the art. Such substitutions preferably are made in accordance with the following list, which substitutions may be determined by routine experimentation provide modified structural and functional properties of a synthesized polypeptide molecule, while maintaining the receptor binding, or inhibiting or mimicking biological activity, as determined by known GPCR receptor activity assays.

	Original	Exemplary
	Residue	Substitution
25	Ala	Gly;Ser
	Arg	Lys
	Asn	Gln;His
	Original	Exemplary
	Residue	Substitution
30	Asp	Glu
	Cys	Se
	Gln	Asn
	Glu	Asp

WO 99/43711

PCT/US99/04438

14

	Gay	Ala;Pro
	His	Asn;Gln
	Ile	Leu;Val
	Leu	Ile;Val
5	Lys	Arg;Gln;Glu
	Met	Leu;Tyr;Ile
	Phe	Met;Leu;Tyr
	Se	Thr
	Original	Exemplary
10	Residue	Substitution
	Thr	Se
	Trp	Tyr
	Tyr	Trp;Phe
	Val	Ile;Leu

15

Put differently, the following six groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Serine (S), Threonine (T);
- 20 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

25 See also, Creighton (1984) PROTEINS, W.H. Freeman and Company. In addition, individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids in an encoded sequence are also "conservatively modified variations".

30 The term "substantial identity" or "substantial similarity" in the context of a polypeptide indicates that a polypeptides comprises a sequence which can have 40% sequence identity to a reference sequence, or preferably 70%, or more preferably 85% sequence identity to the reference sequence, or most preferably 90% identity over a comparison window of about 10-20 amino acid residues. "Percentage

amino acid identity" or "percentage amino acid sequence identity" refers to a comparison of the amino acids of two polypeptides which, when optimally aligned, have approximately the designated percentage of the same amino acids. For example, "95% amino acid identity" refers to a comparison of the amino acids of two polypeptides which when optimally aligned have 95% amino acid identity.

Preferably, residue positions which are not identical differ by conservative amino acid substitutions. Because the substituted amino acids have similar properties, the substitutions do not change the functional properties of the polypeptides. An indication that two polypeptide sequences are substantially identical is that one peptide is immunologically reactive with antibodies raised against the second peptide. Thus, a polypeptide is substantially identical to a second polypeptide, for example, where the two peptides differ only by a conservative substitution. An indication that two nucleic acid sequences are substantially identical is that the polypeptide which the first nucleic acid encodes is immunologically cross reactive with the polypeptide encoded by the second nucleic acid. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2: 482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48: 443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (see generally Ausubel et al., *supra*).

One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J. Mol. Evol.* 35: 351-360 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151-153 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. For example, a reference sequence can be compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps.

Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to

zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989))
5 alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences
10 (see, e.g., Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest
15 sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

The term "nucleic acid" refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form.
20 Unless specifically limited, the term encompasses nucleic acids containing known analogues of natural nucleotides which have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g. degenerate
25 codon substitutions) and complementary sequences and as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer et al., *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka et al., *J. Biol. Chem.* 260:2605-2608
30 (1985); and Cassol et al., 1992; Rossolini et al., *Mol. Cell. Probes* 8: 91-98 (1994)). The term nucleic acid is used interchangeably with gene, cDNA, and mRNA encoded by a gene.

The phrase "a nucleic acid sequence encoding" refers to a nucleic acid which contains sequence information for a structural RNA such as rRNA, a tRNA, or the primary amino acid sequence of a specific protein or peptide, or a binding site for a trans-acting regulatory agent. This phrase specifically encompasses degenerate
5 codons (*i.e.*, different codons which encode a single amino acid) of the native sequence or sequences which may be introduced to conform with codon preference in a specific host cell.

"Nucleic acid probes" may be DNA or RNA fragments. DNA fragments can be prepared, for example, by digesting plasmid DNA, or by use of
10 PCR, or synthesized by either the phosphoramidite method described by Beaucage and Carruthers, *Tetrahedron Lett.* 22:1859-1862 (1981), or by the triester method according to Matteucci, *et al.*, *J. Am. Chem. Soc.*, 103:3185 (1981).

A double stranded fragment may then be obtained, if desired, by annealing the chemically synthesized single strands together
15 under appropriate conditions or by synthesizing the complementary strand using DNA polymerase with an appropriate primer sequence. Where a specific sequence for a nucleic acid probe is given, it is understood that the complementary strand is also identified and included. The complementary strand will work equally well *in situations* where the target is a double-stranded nucleic acid.

20 The terms "substantial identity" or "substantial sequence identity" as applied to nucleic acid sequences and as used herein and denote a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, and more preferably at least 99 percent sequence identity as compared to a
25 reference sequence over a comparison window of at least 20 nucleotide positions, frequently over a window of at least 25-50 nucleotides, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the polynucleotide sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the window of comparison. The
30 reference sequence may be a subset of a larger sequence.

The phrase "specifically binds to an antibody" or "specifically immunoreactive with", when referring to a protein or peptide, refers to a binding reaction which is determinative of the presence of the protein in the presence of a

heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein and do not bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is
5 selected for its specificity for a particular protein. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See Harlow and Lane (1988) ANTIBODIES, A LABORATORY MANUAL, Cold Spring Harbor Publications,
10 New York, for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity.

DETAILED DESCRIPTION

15

20 The present invention is based partly on evidence that transmembrane domains (TM) of GPCRs interact in a specific way in the assembly of receptor molecules. These interactions do not lead to a rigid structure, because some flexibility is required to allow for conformational changes to be made following ligand binding in order to provide the ability of the molecule to signal from the cell
25 surface to the intracellular parts. It was also demonstrated for several GPCRs that the transmembrane domains are involved in ligand binding and thus contain openings that allow penetration of the ligands. Reports that expression of missing transmembrane domains rescues inactive truncated V2 vasopressin, beta-adrenergic and muscarinic M3 receptors (Schoneberg et al. EMBO J. 15: 1283 (1996); Wong et al., J. Biol. Chem. 265: 6219 (1990); Monnot et al., J. Biol. Chem. 271: 1507 (1996);
30 Gudermann et al., Annu. Rev. Neurosci. 20: 399 (1997); Osuga et al., J. Biol. Chem. 272: 25006 (1997)) suggested peptide derived from the sixth transmembrane domain of P2-adrenergic receptor was found to inhibit receptor activation and dimerization

(Hebert et al., *J. Biol. Chem.*, 271(27):16384-92 (1996)). All these observations suggested to us that targeting intramembrane interactions of GPCRs can specifically regulate GPCR function.

5 The hydrophobic nature of the transmembrane peptides makes their penetrations into the bilayer highly probable. Orientation inside the membrane can be controlled by addition of charged residues to the terminus that is supposed to be extracellular.

1. GPCR peptides

10 GPCR polypeptides of the present invention, or nucleic acids encoding therefor, include a finite set of substantially corresponding sequences as substitution peptides or polynucleotide which can be routinely obtained by one of ordinary skill in the art, without undue experimentation, based on the teachings and guidance presented herein. For a detailed description of protein chemistry and structure, see Schulz et al., *PRINCIPLES OF PROTEIN STRUCTURE*, Springer-Verlag, New York, 1978, and Creighton, T. E., *PROTEINS: STRUCTURE AND MOLECULAR PROPERTIES*, W. H. Freeman & Co., San Francisco, 1983.

15 For a presentation of nucleotide sequence substitutions, such as codon preferences, see Ausubel et al, *supra*, at sections A.1.1-A.1.24, and Sambrook et al., *supra*, at Appendices C and D.

20 GPCR polypeptides include homologous sequences and/or fragments of at least one of transmembrane domain 1-7 of one or more GPCRs or homologs thereof, which GPCR polypeptides do not occur naturally, and/or which are provided in an isolated and/or purified form not found in nature.

25 However, in the context of the present invention, GPCR polypeptides of greater than 15-20 amino acids are preferred such that the GPCR polypeptides are able to span the lipid bilayer.

It is particularly preferred that peptides of the invention be selected or modified so that one end is charged and the other is neutral under physiological conditions. This is so that the peptide spontaneously inserts into a membrane. It is of particular importance that the peptide insert in the same orientation as the transmembrane GPCR domain from which it is derived.

30 Peptides of the invention can be derived from any of the 7 TM domains. Non-limiting, illustrative examples of GPCR TM1 and TM2 transmembrane

domains that are used to generate molecules of the present invention include the following:

		101	TM1	150	SEQ. ID No.:
5	GPCRAelegans	HPCEDIMGYV	WLTVVVSFMVG	AVALVANLVV	ALVLLTSQ... ..RRLNV 24
	GRH	NLPTLTLSGK	IRVTVTFFLF	LLSATFNASF	LLKLOKWTQK KEKGKKLSRM 25
	TRH	RAVVALEYQV	VTILLVLIIC	GLGIVGNIMV	VLVVMR.... .TKHM RTP 26
	FSHprec	NPCEDIMGYN	ILRVLIWFIS	ILAITGNIIV	LVILTTSQYK LTV 27
	TSHprec	NPCEDIMGYK	FLRIVVWFVS	LLALLGNV FV	LLILLTSHYK LNV 28
10	LH_CGprec	NPCEDIMGYD	FLRVLIWLIN	ILAIMGNMTV	LFVLLTSRY. KLT V 29
	PGE_EP1	PNTSAVPPSG	ASPALPIFSM	TLGAVSNLLA	LALLAQAA.G RLRRRRSATT 30
	PGE_EP2	SASLSPDRLN	SPVTIPAVMF	IFGVVGNLVA	IVVLCKS... .RKEQKETT 31
	PGE_EP3	QWLPPGE...	.SPAISSVMF	SAGVLGNLIA	LALLARRW.R SAGRSSLSL 32
	PGF	SNTTCQ TENR	LSVFFSVIFM	TVGILSNSLA	IAILMKAY.Q RFRQSKA.S 33
15	PGI	CRNLTYVRGS	VGPATSTLMF	VAGVVGNGLA	LGILSARRPA RP.....SA 34
	TXA2	NITLEERRLI	ASPWFAASFC	VVGLASNLLA	LSVLAGA... RQGGSHTRSS 35
	PAF	HMDSEFRYTL	F.PIVYSIIF	VLGVIANGYV	LWVFARLY.P CKKFNEIK.. 36
	M2	...YKTFEVV	FIVLVAGSL	LVTIIGNILV	MVSI.KVN.R HLQT.....V 37
	M4	HNRYETVEMV	FIATVTGSL	LVTVVGNILV	MLSI.KVN.R QLQT.....V 38
20	M1	..GKGPWQVA	FIGITTGLLS	LATVTGNLLV	LISF.KVN.T ELKT.....V 39
	M3	LGGHTVWQVV	FIAFLTGILA	LVTIIGNILV	IVSF.KVN.K QLKT.....V 40
	M5	LERHRLWEVI	TIAAVTAVVS	LITIVGNVLV	MISF.KVN.S QLKT.....V 41
	H1	KTTMASPQLM	PLVVVLSTIC	LVTVGLNLLV	LYAVRSER.. KLHT.....V 42
	H2	FCLDSTACKI	TITVVLAVLI	LITVAGNVVV	CLAVGLNR.. RLRN.....L 43
25	5HT1A	ISDVTVSQV	ITSLLLGTLI	FCAVLGNACV	VAAIALER.. SLQN.....V 44
	5HT1B	QDSISLPWKV	LLVMLLALIT	LATTLSNAFV	IATVYRTR.. KLHT.....P 45
	5HT1D	DPRTLQALKI	SLAVVLSVIT	LATVLSNAFV	LTILLTR.. KLHT.....P 46
	5HT1E	IRPKTITEKM	LICMTLVVIT	TLTLLNLAV	IMAIGTTK.. KLHQ.....P 47
	5HT1F	ELNRMPSKI	LVSLTSLGLA	LMTTINSLV	IAAIVTR.. KLHH.....P 48
30	5HT2A	QEKN.....	WSALLTAVVI	ILTIAGNILV	IMAVSLEK.. KLQN.....A 49
	5HT2B	IVEEQGNKLH	WAALLILMVI	IPTIGGNTLV	ILAVSLEK.. KLOQ.....A 50
	5HT2C	..QN.....	WPALSIVIII	IMTIGGNILV	IMAVSMEK.. KLHN.....A 51
	5HT5A	SSPLLSVFGV	LILTLLGFLV	AATFAWNLLV	LATILRVR.. TFHR.....V 52
	5HT5Brat	REPPFSAFTV	LVVTLLVLLI	AATFLWNLLV	LVTILRVR.. AFHR.....V 53
35	5HT6rat	GPPAPGGSG	WVAAALCVVI	VLTAANSLL	IVLICTQP.. AVR.....T 54
	5HT7	QINYGRVEKV	VIGSILTLIT	LLTIAGNCLV	VISVCFVK.. KLRQ.....P 55
	alpha1A	GGLVVSAQGV	GVGVFLAAFI	LMAVAGNLLV	ILSVACNR.. HLQT.....V 56
	alpha1B	..QLDITRAI	SVGLVLGAFI	LFAIVGNILV	ILSVACNR.. HLRT.....P 57
	alpha1C	PAPVNISKAI	LLGVILGGLI	LFGVLCNILV	ILSVACHR.. HLHS.....V 58
40	alpha2A	...YSLQVTL	TLVCLAGLLM	LLTVFGNVLV	IIAVFTSR.. ALKA.....P 59
	alpha2B	QDPYSVQATA	AIAAAITFLI	LFTIFGNALV	ILAVLTSR.. SLRA.....P 60
	alpha2C1	RGQYSAGAVA	GLAAVVGFLI	VFTVVGNVLV	VIAVLT SR.. ALRA.....P 61
	alpha2C2	RGQYSAGAVA	GLAAVVGFLI	VFTVVGNVLV	VIAVLT SR.. ALRA.....P 61
	beta1	EPLSQQWTAG	M.GLLMALIV	LLIVAGNVLV	IVAIKTP.. RLQT.....L 62
45	beta2	QQRDEVWVG	M.GIVMSLIV	LAIVFGNVLV	ITAIKFE.. RLQT.....V 63
	beta3	GLPGVPWEAA	LAGALLALAV	LATVGGNLLV	IVAIWTP.. RLQT.....M 64
	beta4turkey	SWAAVLSRQW	AVGAALSITI	LVIVAGNLLV	IVAIKTP.. RLQT.....M 65
	D1A	VVERDFSURI	LTACFLSLLI	LSTLLGNTLV	CAAVIRFR.. HLRSK.....V 66
	D2	DGKADRP HYN	YYATLLTLLI	AVIVFGNVLV	CMASREK.. ALQT.....T 67
50	D3	TGASQARPHA	YYALSYCALI	LAIVFGNGLV	CMAVLKER.. ALQT.....T 68
	D4	ASAGLAGQGA	AALVGGVLLI	GAVLAGNSLV	CVSVATER.. ALQT.....P 69
	D5	GAPPLGPSQV	VTACLLTLLI	IWTLLGNVLV	CAAI VRSR.. HLRAN.....M 70
	A1	MPPSISAFQA	AYIGIEVLIA	LVSVPGNVLV	IWAVKVNQ.. ALRD.....A 71
	A2a	...MPIMGSS	VYITVELAIA	VLAILGNVLV	CWAVWLNS.. NLQN.....V 72
55	A2b	..MLLETQDA	LYVALELVIA	ALSVAGNVLV	CAAVGTAN.. TLQT.....P 73
	A3	NSTTSLANV	TYITMEIFIG	LCAIVGNVLV	ICVVKLNP.. SLQT.....T 74
	OCdrome	LAVPE.WEAL	LTALVLSVII	VLTIIIGNILV	ILSVFTYK.. PLRI.....V 75
	ACTH	RNNSDCPRVV	LPEEIFFTIS	IVGVLENLIV	LLAVFKNK.. NLQA.....P 76
	MSH	QTGARCLEVS	ISDGLFLSLG	LVSLVENALV	VATIAKNR.. NLHS.....P 77
60	MC3	SSSAFCEQVF	IKPEIFLSLG	IVSLLENILV	ILAVVRNG.. NLHS.....P 78
	MC4	SDGGCYEQLF	VSPEVFVTLG	VISLLENILV	IVAIKKNK.. NLHS.....P 79
	MC5	NKSSPCEDMG	IAVEVFLTLG	VISLLENILV	IGAIVKNK.. NLHS.....P 80
	melatonin	DGARPSWLAS	ALACVLIFTI	VVDILGNLLV	ILSVYRNKKL RNA 81
	oxytocin	R..RNEALAR	VEVAVLCLIL	LLALSGNACV	LLALRTTRQK HSR 82
65	conopressinLs	FHGVDEDLLK	IEIAVQATIL	YMTLFGNGIV	LLVLRRLRRQK LTR 83
	V1A	RDVRNEELAK	LEIAVLAVTF	AVAVLGNSSV	LLAL..... .HRTPRKTSR 84

	V1B	WLGRDEELAK	VEIGVLATVL	VLATGGNLAV	LLTLGQLGRK	RSR	85
	V2	LDTRDPLLAR	AELALLSIVF	VAVALSNGLV	LAALA.....	RRGRRGHWAP	86
	CCK_A	PRPSKEWQPA	VOILLYSLIF	LLSVLGNTLV	ITVLI.....	RNCRM..RTV	87
	CCK_B	GAGTRELELA	IRITLYAVIF	LMSVGGNMLI	IVVLGLS...	..RRL..RTV	88
5	NPY1	DCHLPLAMIF	TLALAYGAVI	ILGVSGNLAL	IIIIIL.....	KQKEM..RNV	89
	NTR	DVNTDIYSKV	LVTAVYLALF	VVGTVGNTVT	AFTLAR....	KKSLOSLOST	90
	NK1	QFVQPAWQIV	LWAAAYTVIV	VTSVVGNNVV	MWIILA....	.HKRM..RTV	91
	NK2	AFSMPSWQLA	LWAPAYLALV	LVAVTGNAIV	IWIILA....	.HRRM..RTV	92
	NK3	QFVQPSWRIA	LWSLAYGVVV	AVAVLGNLIV	IWIILA....	.HKRM..RTV	93
10	blueops	YHIAPVWAFY	LQAAFMGTVF	LIGFPLNAMV	LVATL.....	RYKKL..RQP	94
	greenops	YHIAPRWVYH	LTSVWMIFVV	IASVFTNGLV	LAATM.....	KFKKL..RHP	95
	redops	YHIAPRWVYH	LTSVWMIFVV	TASVFTNGLV	LAATM.....	KFKKL..RHP	96
	rhodopsin	YYLAEPWQFS	MLAAYMFLLI	VLGFPINFLT	LYVTVQ....	.HKKL..RTP	97
	violetopsGg	YHIAPPWAFY	LQTAFMGIVF	AVGTPLNAV	LWVTVRYKRL	RQP	98
15	opsin_crab	FPPMNPLWYS	ILGVAMIILG	IICVLGNMGV	IYLMMTTKSL	RTP	99
	ET_Aprec	QTKITSAPKY	INTVISCTIF	IVGMVGNATL	LRRIYQ....	.NKCM..RNG	100
	ET_Bprec	PIEIKETFKY	INTVVSCLVF	VLGIIGNSTL	LRRIYKNK..	...CM..RNG	101
	ET_Cfrog	RAKIRHAFKY	VTILSCVIF	LVGIVGNSTL	LRRIYKNK..	...CM..RNG	102
	galanin	PLFGIGVENF	VTLVVFGILF	ALGVLGNSLV	ITVLARSK..	...PGKPRST	103
20	NMB	GTTTELVIRC	VIPSLYLLII	TVGLLGNIML	VKIFITNS..	...AM..RSV	104
	GRP	DDWSHPGILY	VIPAVYGVII	LIGLIGNITL	IKIFCTVK..	...SM..RNV	105
	BRS3	DNSPGIEALC	AIYITYAVII	SVGILGNAIL	IKVFFKTK..	...SM..QTV	106
	deltaOP	GSASSLALAI	AITALYSAVC	AVGLLGNVLV	MFGIVRYT..	...KM..KTA	107
	kappaOP	PAHISPAIPV	IITAVYSVVF	VVGLVGNLSV	MFVIRRYT..	...KM..KTA	108
25	muOP	GSP.SMITAI	TIMALYSIVC	VVGLFGNFLV	MYVIVRYT..	...KM..KTA	109
	OPRX	GAFLPLGLKV	TIVGLYLAVC	VGGLLGNCLV	MYVILRHT..	...KM..KTA	110
	CB1	FMVLNPSQQL	AI AVLSTLTG	TFTVLENLLV	LCVILHSR..	SLRCR....P	111
	CB2	YMILSGPQKT	AVAVLCTLLG	LLSALENVAV	LYLILSSH..	QLRRK....P	112
30	SSTR1	TLSEGQGSAL	LISFIYSVVC	LVGLCGNSMV	IYVILRYA..	...KM..KTA	113
	SSTR2	EPYYDLTSNA	VLTFIYFVVC	IIGLCGNTLV	IYVILRYA..	...KM..KTI	114
	SSTR3	SPAGLAVSGV	LIPLVYLVC	VVGLLGNLSV	IYVILRHT..	...AS..PSV	115
	SSTR4	GDARAAG.MV	AIQCIYALVC	LVGLVGNALV	IFVILRYA..	...KM..KTA	116
	SSTR5	PAPSAGARAV	LVPVLYLLVC	AAGLGGNTLV	IYVILRFA..	...KM..KTV	117
35	IL8A	MLETETLNKY	VVIIAYALVF	LLSLLGNLSV	MLVILYSR..	...VG..RSV	118
	IL8B	EPESLEINKY	FVVIYALVF	LLSLLGNLSV	MLVILYSR..	...VG..RSV	119
	AT1a	KAGRHNYIFV	MIPTLYSIIF	VVGIFGNSLV	VIVIFYFYM..	...KL..KTV	120
	AT1brat	KAGRHNYIFV	MIPTLYSIIF	VVGIFGNSLV	VIVIFYFYM..	...KL..KTV	120
	AT2	QKPSDKH.LD	AIPILYIIF	VIGFLVNIVV	VTLFCCQK..	...GP..KKV	121
40	BK1	APEAWDLLHR	VLPTFIISIC	FFGLLGNLFV	LLVFLLP..	RQLNV	122
	BK2	QVEWLGLWNT	IQPPFLWVLF	VLATLENIFV	LSVFCLHK..	SSCTV	123
	P2Y7	PSLGVFELSL	LAIILLSVAL	AVGLPGNSFV	VWSILKRMQ..	KRSV	124
	P2Y6	CVYREDFKRL	LLPPVYSVVL	VVGLPLNVCV	IAQICASRR..	TLTR	125
	P2Y5	CSTEDSFKYT	LYGCVFSMVF	VLGLIANCVA	IYIFTFTLK..	VRNE	126
	P2Y4	CWFDEDFKFI	LLPVSYAVVF	VLGLGLNAPT	LWLFIFRLR..	PWDA	127
45	P2Y3chick	CTFHEEFKQV	LLPLVYSVVF	LLGLPLNAV	IGQIWLARK..	ALTR	128
	P2Y2	CRFNEDFKYV	LLPVSYGVVC	VLGLCLNAV	LYIFLCRLK..	TWNA	129
	P2Y1	ALTKTGFOFY	YLPVYILVF	IIGFLGNSVA	IWMFVFHMK..	PWSG	130
	THRprec	GYLTSSWLT	FVPSVYTGVE	VVSLPLNIMA	IVVFILKM..	...KV..KKP	131
	C5a	TSNTLRVPDI	LALVIFAVVF	LVGVLGNALV	VWVTAFEA..	...K...RTI	132
50	GP01mouse	AESEPELVVN	PWDIVLCSSG	TLICCENAVV	VLIIF.HSPS	LR.....AP	133
	R334rat	VESEPELVVN	PWDIVLCSSG	TLICCENAVV	VLIIF.HSPS	LR.....AP	134
	GP21mouse	GPATLLPSPR	AWDVLCISG	TLVSCENALV	VAIIV.GTPA	FR.....AP	135
	GCRCmouse	AESQNPVKA	LLIVAYSFTI	VFSLFGNVLV	CHVIFK.NQR	MHSA	136
	TXKR	...QPPWAVA	LWSLAYGAVV	AVAVLGNLVV	IWIVLA.HKR	MR.....TV	137
55	G10Drat	MELNENTKQV	VLFVFLAIF	VVGLVENVLV	IC.VNCRSG	R.....VGM	138
	RDC1	NMPNKSULLY	TLSEFIYIF	VIGMIANSV	VW.VNIQAKT	TGYDT.....	139
	BLR1	...MASFKAV	FVPVAYSLIF	LLGVIGNVLV	LV.ILERHRQ	TRSSTE....	140
	CL5	REENANFNKI	FLPTIYSIIF	LTGIVGNL	IL.VMGYQKK	LRSMTDKYR.	141
	LCR1	REENANFNKI	FLPTIYSIIF	LTGIVGNL	IL.VMGYQKK	LRSMTDKYR.	141
60	EBI1	KKDVRNFKAW	FLPIMYSIIC	FVGLLGNL	VL.TYIYFKR	LKTMTDTY..	142
	RBS1rat	LGDIVAFGTI	FLSIFYSLVF	TFGLVGNLLV	VL.ALTNSRK	SKSITDIY..	143
	EBI2	LYAHHSTARI	VMPLHYSLVF	IIGLVGNLLA	LV.VIVQNRK	KINSTTLY..	144
	GCRTchick	CSTEDSFKYT	LYGCVFSMVF	VLGLIANCVA	IY.IFTFTLK	VRNETTLY..	145
	APJ	EYTDWKSSGA	LIPAIYMLVF	LLGTTGNL	LWTVFRSSRE	KRRSAD....	146
65	RTArat	EQIATLPPPA	VTNYIFLLC	LCGLVGNL	LWFFGFSIK..	.RT.....P	147
	UHRrat	SLQLVHQLKG	LIVMLYSIVV	VVGLVGNCLL	VLVIARVR..	RLHNV	148
	FMRL1	EPAGHTVLWI	FSLLVHGVTF	VFGVLGNL	IWVA.GFR..	MTRTV	149
	FMRL2	ESAGYTVLRI	LPLVVLGVTF	VLGVLGNL	IWVA.GFR..	MTRTV	150

	fMLP	VSAGYLFIDI	ITYLVFAVTF	VLGVLGNGLV	IWVA.GFR..		MTHTV	151
	OLF1catfish	NGFYNIPTK	YYYAFLCIAY	AVTVLGNSFI	MCTIYLAR..		SLHTA	152
	OLF3catfish	TGLYNIPHAK	YYFLFCFVY	TVTFLGNSFI	MGTIYLAR..		SLHTA	153
	OLF8catfish	GFHDLGEWGP	ILSIPYLLMF	LLSSTSNTL	IYLIISQR..		ALHSP	154
5	OLF32Acatfish	SGFSGIPFSQ	YYFAFLIFIY	IISLCGNSIV	LFMILVDR..		TLHIP	155
	OLF32Bcatfish	SGFSGIPFSQ	YYFVFLIFIY	IISLCGNSIV	LFMILVDR..		TLHIP	156
	OLF32Ccatfish	SGFSGIPFSQ	YYFVFLIFIY	IISLCGNSIV	LFMILVDR..		TLHIP	156
	OLF32Dcatfish	SGFSGIPFSQ	YYFVFLIFIY	IISLCGNSIV	LFMILVDR..		TLHIP	156
	OLF47catfish	IAYNSLGNKN	YLILALGIY	LITLLCNFTL	LAILMNS..		SLQNP	157
10	OLF202catfish	FPGLPPNYYG	LVSVMFFVY	VCTLIGNCTF	FTLFLREK..		SLQKP	158
	OLFCOR1chicken	LTD.NPGLQM	PLFMVFLAIY	TITLLTNLGL	IALISVDL..		HLQTP	159
	OLFCOR2chicken	LTD.NPRLQM	PLFMVFLVIY	TTTLLTNLGL	IALIGMDL..		HLQTP	160
	OLFCOR3chicken	LTD.NPGLQM	PLFMVFLAIY	TITLLTNLGL	IRLISVDL..		HLQTP	161
	OLFCOR4chicken	LTD.NPGLQM	PLFMVFLAIY	TITLLTNLGL	IRLISVDL..		HLQTP	161
15	OLFCOR5chicken	LTD.NPRLQM	PLFMVFLAIY	TITLLANLGL	IALISVDF..		HLQTP	162
	OLFCOR6chicken	LTD.NPGLQM	PLFMVFLAIY	TITLLTNLGL	IALIRIDL..		QLQTP	163
	OLFdog	LPI.DPDQRD	LFYALFLAMY	VTTILGNLLI	IVLIQLDS..		HLHTP	164
	OLF07E	MSE.SPEQQQ	ILFWMFLSMY	LVTVVG NVLI	ILAISSDS..		RLHTP	165
	OLF07I	LPI.QPEQQN	LCYALFLAMY	LTTLLGNLLI	IVLIRLDS..		HLHTP	166
20	OLF07J	FSS.FHEQQI	TLFGVFLALY	ILTLAGNIII	VTIIRIDL..		HLHTP	167
	OLFOR3mouse	VSD.HPHLEI	IFFAVILASY	LLTLVGNLTI	ILLSRLDA..		RLHTP	168
	OLFrat	LTK.QPELLL	PLFFFLVLIY	VLTVVG NVLGM	ILLIIVSP..		LLHTP	169
	OLFF3rat	FVE.NKDLQP	LIYGLFLSMY	LVTVIGNISI	IVAIISDP..		CLHTP	170
	OLFF5rat	LSR.QPQQQQ	LLFLLFLIMY	LATVLGNLLI	ILAIGTDS..		RLHTP	171
25	OLFF6rat	FPG.PRSMRI	GLFLLFLVMY	LLTVVG NLAI	ISLVGAHR..		CLQTP	172
	OLFF12rat	FTE.NPQLHF	LIFALFLSMY	LVTVLGNLLI	IMAIITQS..		HLHTP	173
	OLFI3rat	LPI.PEEHQH	LFYALFLVMY	LTTILGNLLI	IVLVQLDS..		QLHTP	174
	OLFI7rat	FPA.PAPLRV	LLFFLSLLXY	VLVLTENMLI	IIAIRNHP..		TLHKP	175
	OLFI8rat	LPI.PPEHQQ	LFFALFLIMY	LTTFLGNLLI	VVLVQLDS..		HLHTP	176
30	OLFI9rat	LPF.PPEYQH	LFYALFLAMY	LTTLLGNLII	IILILLDS..		HLHTP	177
	OLFI14rat	LPI.PSEYHL	LFYALFLAMY	LTIILGNLLI	IVLVRLDS..		HLHMP	178
	OLFI15rat	LPI.PSEHQH	VFYALFLSMY	LTTVLGNLII	IILIHLDL..		HLHTP	179
	OLFOR17_40	LLE.APGLQP	VVFVLF LFAY	LVTVRGNLSI	LA AVLVEP..		KLHTP	180
	GUST27rat	...MILNCN	PFSGLFLSMY	LVTVLGNLLI	ILAVSSNSHL		HNL	181
35	RPE	PTGFGELEVL	AVGMVLLVEA	LSGLSLNTLT	IFSCKTPEL		RTP	182
	HHRF1	FTDVLNQSKP	VTLFLYGVVF	LFGSIGNFLV	IFTITWRRRI		QCS	183
	HHRF2	NSTEIYQLFE	YTRLGVWLMC	IVGTFLNVLV	ITTILYRRK	K.....	KSP	184
	HHRF3	MTGPLFAIRT	TEAVLNTFII	FVGGLNLAIV	LITQLLTNRV	LG.....	YST	185
	MCP-1A	..DVKQIGAQ	LLPPLYSLVF	IFGFVG NMLV	VLILINCKKL		KCL	186
40	MCP-1B	..DVKQIGAQ	LLPPLYSLVF	IFGFVG NMLV	VLILINCKKL		KCL	186
	PPR1bovine	..EVRKFAKV	FLPAFFTIAF	IIGLAGNSTV	VAIYAYYKKR		RTK	187

TM2

SEQ. ID No.:

45	GPCRAelegans	151	TRFLMCNLAF	ADFILGLYIF	ILTSVSAVTR	GDYHNYVQOW	QNGAGCKILG	188
	GRH		.KLLLKHLTL	ANLLETILVM	PLDGMWNITV	QWYA.....	.GELLCKVLS	189
	TRH		TNCYLVSLAV	ADLMVLVAAG	LPNITDSIYG	SWVYGYV...	GCLCIT	190
	FSHprec		PRFLMCNLAF	ADLCIGIYLL	LIASVDIHTK	SOYHNYAIDW	QTGAGCDAAG	191
50	TSHprec		PRFLMCNLAF	ADFCMGMYLL	LIASVDLYTH	SEYYNHAIDW	QTGPGCNTAG	192
	LH_CGprec		PRFLMCNLSF	ADFCMGLYLL	LIASVDSQTK	GQYYNHAIDW	QTGSGCSTAG	193
	PGE_EP1		FLLFVASLLA	TDLAGHVIPG	ALVLRLYTA.	GRA	PAGGACHFLG	194
	PGE_EP2		FYTLVCGLAV	TDLLGTLLVS	PVTIATYMKG	QWPG	GQP.LCEYST	195
	PGE_EP3		FHVLVTELVF	TDLLGTCLIS	PVVLASYARN	QT..LVALAP	ESR.ACTYFA	196
55	PGF		FLLLASGLVI	TDFFGHLING	AIASFVYASD	KE..WIRFDQ	.SNVLCISIFG	197
	PGI		FAVLVTGLAA	TDLLGTSFLS	PAVFVAYARN	SS..LLGLAR	GGPALCDAFA	198
	TXA2		FLTFLCGLVL	TDFLGLLVTG	TIVVSQHAAL	FE..WHAUDP	GCR.LCRFMG	199
	PAF		..IFMVNLTM	ADMLFLITLP	LWIVYYQ.NQ	GNWIL.....	PK.FLCNVAG	200
	M2		NNYFLFSLAC	ADLIIGVFSM	NLYTLYTVIG	YWPL	.GPVVC DLWL	201
60	M4		NNYFLFSLAC	ADLIIGAFSM	NLYTVYIIGK	YWPL	.GAVVC DLWL	202
	M1		NNYFLLSLAC	ADLIIGTFSM	NLYTTYLLMG	HWAL	.GTLACDLWL	203
	M3		NNYFLLSLAC	ADLIIGVISM	NLFTTYIIMN	RWAL	.GNLACDLWL	204
	M5		NNYLLSLAC	ADLIIGIFSM	NLYTTYILMG	RWAL	.GSLACDLWL	205
	H1		GNLYIVSLSV	ADLIVGAVVM	PMNILYLLMS	KWSL	.GRPLCLFWL	206
65	H2		TNCFIVSLAI	TDLLGLLVVL	PFSAIYQLSC	KWSF	G.KVFCNIYT	207
	5HT1A		ANYLIGSLAV	TDLMVSVLVL	PMAALYQVLN	KWTL	.GOVTCDLFI	208
	5HT1B		ANYLIASLAV	TDLLVSILVM	PISTMYTVTG	RWTL	.GOVVCDFWL	209
	5HT1D		ANYLIGSLAT	TDLLVSILVM	PISIAYTITH	TWNF	.GOILCDIWL	210

	5HT1E	ANYLICSLAV	TDLLVAVLVM	PLSIIYIVMD	RWKL	.GYFLCEVWL	211
	5HT1F	ANYLICSLAV	TDFLVAVLVM	PFSIVYIVRE	SWIM	.GOVVCIDIWL	212
	5HT2A	TNYFLMSLAI	ADMLLGFLVM	PVSMLTILYG	YRWPL	P.SKLCVWI	213
5	5HT2B	TNYFLMSLAV	ADLLVGLFVM	PIALLTIMFE	AMWPL	P.LVLCPAWL	214
	5HT2C	TNYFLMSLAI	ADMLVGLLVM	PLSLLAILYD	YVWPL	P.RYLCPVWI	215
	5HT5A	PHNLVASMAY	SDVLVAALVM	PLSLVHELSE	GRRWQL	.GRRLCQLWI	216
	5HT5Brat	PHNLVASTAV	SDVLVAALVM	PLSLVSELSA	GRRWQL	.GRSLCHVWI	217
	5HT6rat	SNFFLVSLFT	SDLMVGLVVM	PPAMLNALYG	RWVL	A.RGLCLLWT	218
	5HT7	SNYLIVSLAL	ADLSVAVAVM	PFVSVTDLIG	G.....KWIF	.GHFFCNVFI	219
10	alpha1A	TNYFIVNLAV	ADLLLSATVL	PFSATMEVLG	FWAF	G.RAFCDVWA	220
	alpha1B	TNYFIVNLAM	ADLLLSFTVL	PFSAALEVLG	YWVL	G.RIFCDIWA	221
	alpha1C	THYYIVNLAV	ADLLLTSTVL	PFSAlFEVLG	YWAF	G.RVFCNIWA	222
	alpha2A	QNLFLVSLAS	ADILVATLVI	PFSLANEVMG	.Y.....WYF	.GKAWCEIYL	223
	alpha2B	QNLFLVSLAA	ADILVATLII	PFSLANELLG	.Y.....WYF	R.RTWCEVYL	224
15	alpha2C1	QNLFLVSLAS	ADILVATLVM	PFSLANELMA	.Y.....WYF	.GOVWCGVYL	225
	alpha2C2	QNLFLVSLAS	ADILVATLVM	PFSLANELMA	.Y.....WYF	.GOVWCGVYL	225
	beta1	TNLFIMSLAS	ADLVMGLLVV	PFGATIVVWG	RWEY	GS.FFCELWT	226
	beta2	TNYFITSLAC	ADLVMGLAVV	PFGAAHILMK	MWTF	GN.FWCEFWT	227
	beta3	TNVFVTSLAA	ADLVMGLLVV	PPAATLALTG	HWPL	GA.TGCELWT	228
20	beta4turkey	TNVFVTSLAC	ADLVMGLLVV	PPGATILLSG	HWPY	GT.VVCELWT	229
	D1A	TNFFVISLAV	SDLLVAVLVM	PWKAVAEIAG	FWPF	GS..FCNIWV	230
	D2	TNYLIVSLAV	ADLLVATLVM	PWVVYLEVVG	E.....WKF	S.RIHCDIFV	231
	D3	TNYLVVSLAV	ADLLVATLVM	PWVVYLEVTG	GV.....WNF	S.RICCDVFW	232
	D4	TNSFIVSLAA	ADLLLALLVL	PLFVYSEVQG	GA.....WLL	SPRLC.DALM	233
25	D5	TNVFIVSLAV	SDLFVALLVM	PWKAVAEVAG	YWPF	GA..FCDVWV	234
	A1	TFCFIVSLAV	ADVAVGALVI	PLAILINIGP	QTYFHTCL..	MVA	235
	A2a	TNYFVVSLLA	ADIAVGVLAI	PFAITISTGF	CAACHGCL..	FIA	236
	A2b	TNYFLVSLAA	ADVAVGLFAI	PFAITISLGF	CTDFYGCL..	FLA	237
	A3	TFYFIVSLAL	ADIAVGVLVM	PLAIVVSLGI	TIHFYSCL..	FMT	238
30	OCdrome	QNFFIVSLAV	ADLTVALLV	PFNVAYSILG	R.....WEF	GI.HLCKLWL	239
	ACTH	MYFFICSLAI	SDMLGSLYKI	LENILILRN	MGYLKPRGSF	ET.TADDIID	240
	MSH	MYCFICCLAL	SDLLVSGTNV	LETAVILLE	AGALVARAAV	LQ.QLDNVID	241
	MC3	MYFFLCSLAV	ADMLVSVSNA	LETIMIAIVH	SDDYTFEDQF	IQ.HMDNIFD	242
	MC4	MYFFICSLAV	ADMLVSVSNG	SETIIITLLN	STD.TDAQSF	TV.NIDNVID	243
35	MC5	MYFFVCSLAV	ADMLVSMSSA	WETITTYLLN	NKHLVIADAF	V.RHIDNVFD	244
	melatonin	GNIFVVSLLAV	ADLVVAIYPY	PLVLMsIFNN	GWNLGYLH..	CQVSG	245
	oxytocin	LFFFMKHLAI	ADLVVAVFQV	LPQLLWDITF	RFYGP.....	..DLLCRLVK	246
	conopressinLs	MOWFIAHLAF	ADIFVGFFNI	LPQLISDVTI	VFHGDD....	...FTCRFIK	247
	V1A	MHLFIRHLAL	ADLAVAFFQV	LPQMCWDITY	RFRGPD....	...WLCRVVK	248
40	V1B	MHLFVLHLAL	TDLAVALFQV	LPQLLWDITY	RFQGP.....	..DLLCRAVK	249
	V2	IHFVFIGHLCL	ADLAVALFQV	LPQLAWKATD	RFRGPD....	...ALCRAVK	250
	CCK_A	TNIFLLSLAV	SDMLCLFCM	PFNLIPNLLK	DFIFGS....	...AVCKTTT	251
	CCK_B	TNAFLLSLAV	SDLLAVACM	PFTLLPNLMG	TFIFGT....	...VICKAVS	252
	NPY1	TNIVNLVSLF	SDLLVAIMCL	PFTFVYTLMD	HWVFG.....	...AMCKLNP	253
45	NTR	VHYHLGSLAL	SDLLTLLLAM	PVELYNFIWV	HHPWAF....	.GDAGCRGY	254
	NK1	TNYFLVNLAF	AEASMAAFNT	VVNFTYAVHN	EWYGL.....	...FYCKFHN	255
	NK2	TNYFIVNLAL	ADLCMAAFNA	AFNFVYASHN	IWYFGR....	...AFCYFQN	256
	NK3	TNYFLVNLAF	SDASMAAFNT	LVNFIYALHS	EWYFGA....	...NYCRFQN	257
50	blueops	LNILVNVVSF	GGFLLCIFSV	FPVFVASCNG	YFVFG.....	...HVCALLEG	258
	greenops	LNWILVNLAV	ADLAETVIAS	TISVNVQVYG	YFVLGH....	...PMCVLEG	259
	redops	LNWILVNLAV	ADLAETVIAS	TISIVNQVSG	YFVLGH....	...PMCVLEG	260
	rhodopsin	LNILVNLAV	ADLFMVLGGF	TSTLYTSLHG	YFVFG.....	...TGCNLEG	261
	violetopsGg	LNILVNLISA	SGFVSCVLSV	FVVFVASARG	YFVFG.....	...KRVCELEA	262
	opsin_crab	TNLLVNLAF	SDFCMMAFMM	PTMTSNCFAE	TWILG.....	...PFMCEVYG	263
55	ET_Aprec	PNALIASLAL	GDLIYVIDL	PINVFKLLAG	RWPFDH.NDF	GV.FLCKLFP	264
	ET_Bprec	PNILIASLAL	GDLLHIVIDI	PINVKLLAE	DWPFGE....	...MCKLVP	265
	ET_Cfrog	PNVLIASLAL	GDLFYILIAI	PIISISFWLS	TGH.....	...SEYIYQ	266
	galanin	TNLFILNLSI	ADLAYLLFCI	PFOATVYALP	TWVLGA....	...FICKFIH	267
	NMB	PNIFISNLAA	GDLLLLLTCV	PVDASRYFFD	EWMFGKVG..	CKLIP	268
60	GRP	PNLFISLAL	GDLLLLLITCA	PVDASRYLAD	RWLFGRIG..	CKLIP	269
	BRS3	PNIFITSLAF	GDLLLLLTCV	PVDATHYLAE	GWLFGRIG..	CKVLS	270
	deltaOP	TNIYIFNLAL	ADALATSTLP	FQSAKYLMET	.WPFGE....	...LLCKAVL	271
	kappaOP	TNIYIFNLAL	ADALVTSTMP	FQSTVYLMNS	.WPFGE....	...VLCKIVI	272
	muOP	TNIYIFNLAL	ADALATSTLP	FQSVNYLMGT	.WPFGE....	...ILCKIVI	273
65	OPRX	TNIYIFNLAL	ADTLVLLTLP	FQGTDILLGF	.WPFGE....	...ALCKTVI	274
	CB1	SYHFIGSLAV	ADLLGSVIFV	YSFIDFHVFH	RKD.....	.SRNVFLFKL	275
	CB2	SYLFIGSLAG	ADFLASVFA	CSFVNHFVFH	GVD.....	.SKAVFLFKI	276
	SSTR1	TNIYILNLAI	ADELLMLSVP	FLVTSTLLRH	.WPFGE....	...LLCRLVL	277

	SSTR2	TNIYILNLAI	ADELFLMLGLP	FLAMQVALVH	.WPFGK....	...AICRVVM	278
	SSTR3	TNVYILNLAL	ADELFLMLGLP	FLAAQNALSY	.WPFGS....	...LMCRLVM	279
	SSTR4	TNIYLLNLAV	ADELFLMLSV	FVASSAALRH	.WPFGS....	...VLCRAVL	280
5	SSTR5	TNIYILNLAV	ADVLVYMLGLP	FLATQNAASF	.WPFGP....	...VLCRLVM	281
	IL8A	TDVYLLNLAL	ADLLFALTLP	IWAA..SKVN	GWIFGT....	...FLCKVVS	282
	IL8B	TDVYLLNLAL	ADLLFALTLP	IWAA..SKVN	GWIFGT....	...FLCKVVS	282
	AT1a	ASVFLNLAL	ADLCFLLTLP	LWAVYTAMEY	RWPFGN....	...YLCKIAS	283
	AT1brat	ASVFLNLAL	ADLCFLLTLP	LWAVYTAMEY	RWPFGN....	...HLCKIAS	284
10	AT2	SSIYIFNLAV	ADLLLLATLP	LWATYYSYRY	DWLFPG....	...VMCKVFG	285
	BK1	AEIYLANLAA	SDLVFLVGLP	FWAENIWNQF	NWPFGA....	...LLCRVIN	286
	BK2	AEIYLGNLAA	ADLILACGLP	FWAITISNNF	DWLFGE....	...TLCRVVM	287
	P2Y7	TALMVNLAL	ADLAVLLTAP	FFLHFLAQGT	WSFGLA....	...GCRLCH	288
	P2Y6	SAVYTLNLAL	ADLLYACSLP	LLIYNYARGD	HWPFGD....	...LACRLVR	289
15	P2Y5	TTYMLNLAI	SDLLFVFTLP	FRIYYFVVRN	.WPFGE....	...VLCKISV	290
	P2Y4	TATYMFHLAL	SDTLYVVSPL	TLIYYAAHN	HWPFGT....	...EICKFVR	291
	P2Y3chick	TTYMLNLAM	ADLLYVCSLP	LLIYNYTQKD	YWPFGD....	...FTCKFVR	292
	P2Y2	STTYMFHLAV	SDALYAASLP	LLVYYARGD	HWPFGT....	...VLCKLVR	293
	P2Y1	ISVYMFNLAL	ADFLYVLTLP	ALIFYFVNKT	DWIFGD....	...AMCKLQR	294
20	THRprec	AVVYMLHLAT	ADVLFVSPL	FKISYYFSGS	DWQFGS....	...ELCRFVT	295
	C5a	NAIWFLNLAV	ADFLSCLALP	ILFTSIVQHH	WPFGE....	...ACSILP	296
	GP01mouse	MFLIGSLAL	ADLLAGLGLI	INFVFAYLLO	SE....	...ATKLVTI	297
	R334rat	MFLIGSLAL	ADLLAGLGLI	INFVFAYLLO	SE....	...ATKLVTI	297
	GP21mouse	MFLVGLSLAV	ADLLAGLGLV	LHFAADFCIG	SP....	...EMSLMLV	298
25	GCRCmouse	TSLFIVNLAV	ADIMITLLNT	PFTLVRFVNS	TWVFGK....	...GMCHVSR	299
	TXKR	TNSFLVNLAF	ADAAMAALNA	LVNFIYALHG	EWYFGA....	...NYCRFQN	300
	G10Drat	LNLYILNMAV	ADLGIILSLP	VWMLEVMLEY	TWLWGS....	...FSCRFIH	301
	RDC1	.HCYILNLAI	ADLWVLTIP	VWVSVLVQHN	QWPMGE....	...LTCKVTH	302
	BLR1	..TFLFHLAV	ADLLLVFILP	FAVAEGSV..	GWVLGT....	...FLCKTVI	303
30	CL5	LHLSV	ADLLFVITLP	FWAVDAVA..	NWYFGN....	...FLCKAVH	304
	LCR1	LHLSV	ADLLFVITLP	FWAVDAVA..	NWYFGN....	...FLCKAVH	304
	EBI1	LLNLAV	ADILFLLTLP	FWAYSAAK..	SWVFGV....	...HFCKLIF	305
	RBS1rat	LLNLAL	SDLLFVATLP	FWTHYLIS..	HEGLHN....	...AMCKLTT	306
	EBI2	STNLVI	SDILFTTALP	TRIAYYAMGF	DWRIGD....	...ALCRITA	307
35	GCRTchick	MLNLAI	SDLLFVFTLP	FRIYYFVVR.	NWPFGD....	...VLCKISV	308
	APJ	..IFIASLAV	ADLTFVVTLP	LWATYTYRDY	DWPFGT....	...FFCKLSS	309
	RTArat	FSIYFLHLAS	ADGIYLFASKA	VIALLNMGTF	LGSFPD....	...YVRRVSR	310
	UHRrat	TNFLIGNLAL	SDVLMCAACV	PLTLAYAFEP	RGWVFG....	...GGLCHLVF	311
	FMRL1	NTICYNLAL	ADFSFSAILP	FRMVSAMRE	KWPFAS....	...FLCKLVH	312
40	FMRL2	TTICYNLAL	ADFSFTATLP	FLIVSMAMGE	KWPFGE....	...FLCKLIH	313
	fMLP	TTISYLNALAV	ADFCFTSTLP	FFMVRKAMGG	HWPFGW....	...FLCKFLF	314
	OLF1catfish	KYITVFNALAL	SDLGSSALI	PKLIDTFLF.	ENQV	ISYEACLANM	315
	OLF3catfish	KYIAVFNALAL	SDLCGSSALI	PKLLDMLLF.	ENQS	ISYEACLSNM	316
	OLF8catfish	MCILIGLMAV	VDLSMPIFCV	PNMLLSFLF.	NWKG	ISLVGCLVQM	317
45	OLF32Acatfish	KYMGIFNLAL	SDFGETNVLI	PSLVKTLFF.	DSQY	ISYDACLANM	318
	OLF32Bcatfish	KYMGIFNLAL	SDFGETNALI	PSLVKTLFF.	DSQY	ISYDACLANM	319
	OLF32Ccatfish	KYMGIFNLAL	SDIGETNALI	PSLVKTLFF.	DSQY	ISYDACLTNM	320
	OLF32Dcatfish	KYMGIFNLAL	SDFGETNALI	PSLVKTLFF.	DSQY	ISYDACLANM	319
	OLF47catfish	KFLAVFNALAV	VDISINSVII	PQMVPVFVF.	NLNH	ISFESCFSQM	321
50	OLF202catfish	MYIIMNLALAA	SDVLFSTTTL	PKIARYWF.	GDGS	ISFVGCFIQM	322
	OLFCOR1chicken	MYIFLQNLSE	TDAAYSTVIT	PKMLATFL..	EERKT	ISYVGCILQY	323
	OLFCOR2chicken	MYIFLQNLSE	TDAAYSTVIT	PKMLATFL..	EERT	ISYVGCILQY	324
	OLFCOR3chicken	MYIFLQNLSE	TDAAYSTVIT	PKMLATFL..	EERKT	ISYVGCILQY	323
	OLFCOR4chicken	MYIFLQNLSE	TDAAYSTVIT	PKMLATFL..	EERKT	ISYVGCILQY	323
55	OLFCOR5chicken	MYIFLQNLSE	TDAAYSTVIT	PKMLATFL..	EERT	ISYVGCILQY	324
	OLFCOR6chicken	MYIFLQNLSE	TDAVYSTVIT	PKMLATFL..	EETKT	ISYVGCILQY	325
	OLFdog	MYLFLSNLSE	SDLCFSSVTM	PKLLQNMQ..	SQVPS	IPYAGCLTQM	326
	OLF07E	VYFFLANLSE	TDLFFVTNTI	PKMLVNLQ..	SHNKA	ISYAGCLTQL	327
	OLF07I	MYLFLSNLSE	SDLCFSSVTI	PKLLQNMQ..	NQDPS	IPYADCLTQM	328
	OLF07J	MYFFLSMLST	SETVYTLVIL	PRMLSSLV..	GMSQP	MSLAGCATQM	329
60	OLFOR3mouse	MYFFLSNLSS	LDLAFTTSSV	PQMLKNLW..	GPKDT	ISYGGCVTQL	330
	OLFrat	MYFFLSNLSE	VDLCYSTVIT	PKMLVNLQ..	GKKNF	ITYSECMAQF	331
	OLFF3rat	MYFFLSNLSE	VDICFISTTV	PKMLVNIQ..	TQNNV	ITYAGCITQI	332
	OLFF5rat	MYFFLSNLSE	VDVCFSSSTTV	PKVLANHI..	LGSQA	ISFSGCLTQL	333
65	OLFF6rat	MYFFLCNLSE	LEIWFTTACV	PKTLATF...	APRGGV	ISLAGCATQM	334
	OLFF12rat	MYFFLANLSE	VDICFTSTTI	PKMLVNIY..	TQSKS	ITYEDCISQM	335
	OLFI3rat	MYLFLSNLSE	SDLCFSSVTM	PKLLQNMNR..	SQDTS	IPYGGCLAQT	336
	OLFI7rat	MYFFLANMSF	LEIWYVTVTI	PKMLAGFIG.	..SKENHGQL	ISFEACMTQL	337
	OLFI8rat	MYLFLSNLSE	SDLCFSSVTM	LKLLQNIQ..	SQVPS	ISYAGCLTQI	338

	OLFI9rat	MYLFLSNLSF	ADLCFSSVTM	PKLLQNMQ...	...SQVPS	IPYAGCLAQI	339
	OLFI14rat	MYLFLSNLSF	SDLCFSSVTM	PKLLQNMQ...	...SQVPS	ISYTGCLTQL	340
	OLFI15rat	MYLFLSNLSF	SDLCFSSVTM	PKLLQNMQ...	...SQVPS	IPFAGCLTQL	341
	OLFOR17_40	MYFFLGNSLV	LDVGCISVTV	PSMLSRLL...	...SRKRA	VPCGACLTQL	342
5	GUST27rat	MYFFLSNLSF	VDICFISTTI	PKMLVNIH...	...SQTkd	ISYIECLSQV	343
	RPE	CHLLVLSLAL	ADSGISLNAL	VAATSSLLRR	WPYG.....	..SDGCQAHG	344
	HHRF1	GDVYFINLAA	ADLLFVCTLP	LWMQYLLDHN	SLA.....	..SVPCTLLT	345
	HHRF2	SDTYICNLAV	ADLLIVVGLP	FFLEYAKHHP	KLSR.....	..EVVCSGLN	346
	HHRF3	PTIYMTNLYS	TNFLTTLTVLP	FIVLSNQWLL	PAG.....	..VASCKFLS	347
10	MCP-1A	TDIYLLNLAI	SDLLFLITLP	LWAHSAANEW	VFG.....	..NAMCKLFT	348
	MCP-1B	TDIYLLNLAI	SDLLFLITLP	LWAHSAANEW	VFG.....	..NAMCKLFT	348
	PPR1bovine	TDVYILNLAV	ADLFLFLFTLP	FWAVNAVHGW	VLG.....	..KIMCKVTS	349

The above sequences were obtained from a public database. Examples of TM3 and
 15 TM5 transmembrane domain sequences are included in WO 94/05695.

Examples of TM4, TM6, and TM7 transmembrane domain
 sequences can similarly be obtained from public sources

2. Synthesis of peptides

The peptides or fragments of GPCRs may be isolated from a natural
 20 source, chemically synthesized or produced recombinantly, in order to provide
 GPCR polypeptides which mimic, modulate or inhibit binding of ligands to G-protein
 coupled receptors.

a. Chemical synthesis of GPCR transmembrane peptides

Transmembrane peptides of the present invention are be made using
 25 well known peptide synthesis procedures, as described in e.g., Merrifield, *Science*
 232: 341-347 (1986), Barany and Merrifield, *THE PEPTIDES*, Gross and Meienhofer,
 eds. (N.Y., Academic Press), pp. 1-284 (1979); and Stewart and Young, *SOLID PHASE*
PEPTIDE SYNTHESIS (Rockford, Ill., Pierce), 2d Ed. (1984).

30 The peptides were synthesized by a flow-through solid phase peptide
 synthesis on 432A Applied Biosystems Peptide Synthesizer utilizing Fmoc amino acid
 derivatives. To overcome aggregation that frequently occurs during the synthesis of
 hydrophobic peptides and leads to the blockage of the growing peptide chain,
 FmocHmb derivatives of Ala, Val and Leu were introduced into the different
 35 sequences, but not more than two derivatives of that type per peptide to prevent
 sterical hindrance during the synthesis. Coupling on the step after FmocHmb amino
 acid was prolonged to 90 min, since this protection group causes slowing of the next
 coupling step due to steric hindrance (T. Johnson, M. Quibell, *Tetrahedron Lett.*
 35:463 (1994). The purity of the peptides was assessed by reverse phase HPLC and
 40 the structures were confirmed by matrix-assisted laser-desorption mass spectrometry.

b. Recombinant production of GPCR transmembrane peptides

Nucleic acids that encode GPCR transmembrane peptides may be obtained by synthesizing, isolating or obtaining a nucleic acid sequence that encodes a GPCR protein, and subcloning a region of the sequence that encodes a desired transmembrane peptide.

i. Chemical synthesis of oligonucleotides

Oligonucleotides used in the present invention, including sequences that encode transmembrane peptides, are optionally chemically synthesized using the solid phase phosphoramidite triester method of Beaucage and Carruthers, *Tetrahedron Lett.*, 22(20): 1859-1862 (1981) using an automated synthesizer as described in Needham-VanDevanter *et al.*, *Nucleic Acids Res.*, 12: 6159-6168 (1984). The chemically synthesized oligonucleotides are then purified by native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson and Regnier, *J. Chrom.*, 255: 137-149 (1983). The sequence of the synthetic oligonucleotide is verified, for example by using the chemical degradation method of Maxam and Gilbert in Grossman, L. and Moldave, D., eds. Academic Press, New York, *Methods in Enzymology*, 65:499-560 (1980).

The DNA sequences of the present invention coding for GPCR transmembrane peptides protein can be modified (*i.e.*, mutated) to prepare various mutations. Such mutations may be either degenerate, *i.e.*, the mutation does not change the amino acid sequence encoded by the mutated codon, or non-degenerate, *i.e.*, the mutation changes the amino acid sequence encoded by the mutated codon. These modified DNA sequences may be prepared, for example, by mutating known sequences so that the mutation results in the deletion, substitution, insertion, inversion or addition of one or more amino acids in the encoded polypeptide using various methods known in the art. For example, the methods of site-directed mutagenesis described in Taylor *et al.*, *Nucl. Acids Res.* 13, 8749-8764 (1985) and Kunkel, *Proc. Natl. Acad. Sci. USA* 82, 482-492 (1985) may be employed. In addition, kits for site-directed mutagenesis may be purchased from commercial vendors. For example, a kit for performing site-directed mutagenesis may be purchased from Amersham Corp. (Arlington Heights, Ill.). Both degenerate and non-degenerate mutations may be advantageous in producing or using the polypeptides of the present invention. For example, these mutations may permit

higher levels of production, easier purification, or provide additional restriction endonuclease recognition sites. All such modified DNAs (and the encoded polypeptide molecules) are included within the scope of the present invention.

ii. Recombinant isolation of GPCR transmembrane peptide-encoding nucleic acids

Nucleic acids that encode GPCR can be isolated from genomic or cDNA libraries, subcloning the library into expression vectors, labelling probes, DNA hybridization, and the like, as described in Sambrook, et al., MOLECULAR CLONING - A LABORATORY MANUAL (2nd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989. This manual is hereinafter referred to as "Sambrook, et al."

Various methods of amplifying target sequences, such as the polymerase chain reaction (PCR), can also be used to prepare DNA encoding GPCR transmembrane peptides or a peptide fragment thereof. In PCR techniques, oligonucleotide primers complementary to the two 3' borders of the DNA region to be amplified are synthesized. The polymerase chain reaction is then carried out using the two primers. See PCR PROTOCOLS: A GUIDE TO METHODS AND APPLICATIONS. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.), Academic Press, San Diego (1990). Primers can be selected to amplify the entire regions encoding a full-length GPCR transmembrane peptides or to amplify smaller DNA segments as desired. Once selected sequences are PCR-amplified, oligonucleotide probes can be prepared from sequence obtained. These probes can then be used to isolate DNA's encoding GPCR transmembrane peptides or a peptide fragment thereof.

iii. Recombinant expression of transmembrane peptide-encoding nucleic acids

Once a nucleic acid encoding a GPCR transmembrane peptides or a peptide fragment thereof is isolated and cloned, the nucleic acid is expressed in a variety of recombinantly engineered cells to ascertain that the isolated nucleic acid indeed encodes the desired GPCR transmembrane peptides or a peptide fragment thereof. The expression of natural or synthetic nucleic acids is typically achieved by operably linking a nucleic acid of interest to a promoter (which is either constitutive or inducible), incorporating the construct into an expression vector, and introducing

the vector into a suitable host cell. Typical vectors contain transcription and translation terminators, transcription and translation initiation sequences, and promoters useful for regulation of the expression of the particular nucleic acid. The vectors optionally comprise generic expression cassettes containing at least one independent terminator sequence, sequences permitting replication of the cassette in eukaryotes, or prokaryotes, or both, (e.g., shuttle vectors) and selection markers for both prokaryotic and eukaryotic systems. Vectors are suitable for replication and integration in prokaryotes, eukaryotes, or preferably both. See, Gilman and Smith (1979), *Gene*, 8: 81-97; Roberts et al. (1987), *Nature*, 328:731-734; Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in Enzymology* 152, Academic Press, Inc., San Diego, CA (Berger); Sambrook et al. (1989), *MOLECULAR CLONING - A LABORATORY MANUAL* (2nd ed.) Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor Press, N.Y., (Sambrook); and F.M. Ausubel et al., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (1994 Supplement) (Ausubel). Product information from manufacturers of biological reagents and experimental equipment also provide information useful in known biological methods. Such manufacturers include the SIGMA chemical company (Saint Louis, MO), R&D systems (Minneapolis, MN), Pharmacia LKB Biotechnology (Piscataway, NJ), CLONTECH Laboratories, Inc. (Palo Alto, CA), Chem. Genes Corp., Aldrich Chemical Company (Milwaukee, WI), Glen Research, Inc., GIBCO BRL Life Technologies, Inc. (Gaithersburg, MD), Fluka Chemica-Biochemika Analytika (Fluka Chemie AG, Buchs, Switzerland), and Applied Biosystems (Foster City, CA), as well as many other commercial sources known to one of skill.

The nucleic acids (e.g., coding sequences, promoters and vectors) used in the present method can be isolated from natural sources, obtained from such sources as ATCC or GenBank libraries, or prepared by synthetic methods. Synthetic nucleic acids can be prepared by a variety of solution or solid phase methods. Detailed descriptions of the procedures for solid phase synthesis of nucleic acids by phosphite-triester, phosphotriester, and H-phosphonate chemistries are widely available. See, for example, Itakura, U.S. Pat. No. 4,401,796; Caruthers, et al., U.S. Pat. Nos. 4,458,066 and 4,500,707; Beaucage, et al., (1981) *Tetrahedron Lett.*, 22:1859-1862; Matteucci, (1981) et al., *J. Am. Chem. Soc.*, 103:3185-3191;

Caruthers, et al., (1982) *Genetic Engineering*, 4:1-17; Jones, chapter 2, Atkinson, et al., chapter 3, and Sproat, et al., chapter 4, in *OLIGONUCLEOTIDE SYNTHESIS: A PRACTICAL APPROACH*, Gait (ed.), IRL Press, Washington D.C. (1984); Froehler, et al., (1986) *Tetrahedron Lett.*, 27:469-472; Froehler, et al., (1986) *Nucleic Acids Res.*, 14:5399-5407; Sinha, et al. (1983) *Tetrahedron Lett.*, 24:5843-5846; and Sinha, et al., (1984) *Nucl. Acids Res.*, 12:4539-4557.

3. Derivatized Peptides and Peptidomimetics

The design of chemically modified peptides and peptide mimics which are resistant to degradation by proteolytic enzymes or have improved solubility or binding properties is well known.

Modified amino acids or chemical derivatives of GPCRs peptides according to the present invention may contain additional chemical moieties or modified amino acids not normally a part of the protein. Covalent modifications of the peptide are thus included within the scope of the present invention. Such modifications may be introduced into a GPCR polypeptide by reacting targeted amino acid residues of the polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues. The following examples of chemical derivatives are provided by way of illustration and not by way of limitation.

The design of peptide mimics which are resistant to degradation by proteolytic enzymes is well known, both for hormone agonist/antagonist and for enzyme inhibitor design. See e.g., Sawyer, in *STRUCTURE-BASED DRUG DESIGN*, P. Verapandia, Ed., NY 1997; U.S. Patent No. 5,552,534; and U.S. Patent No. 5,550,251.

Historically, the major focus of peptidomimetic design has evolved from receptor-targeted drug discovery research and has not been directly impacted by an experimentally-determined three-dimensional structure of the target protein. Nevertheless, a hierarchical approach of peptide→peptidomimetic molecular design and chemical modification has evolved over the past two decades, based on systematic transformation of a peptide ligand and iterative analysis of the structure-activity and structure-conformation relationships of "second generation" analogs. Such work has typically integrated biophysical techniques (x-ray crystallography

and/or NMR spectroscopy) and computer-assisted molecular modeling with biological testing to advance peptidomimetic drug design.

The three-dimensional structural properties of peptides are defined in terms of torsion angles (Ψ , ϕ , ω , χ) between the backbone amine nitrogen (N^α), backbone carbonyl carbon (C^1), backbone methionine carbon (C^α), and side chain hydrocarbon functionalization (e.g., C^β , C^γ , C^δ , C^ϵ of Lys) derived from the amino acid sequence. A Ramachandran plot (Ψ versus ϕ) may define the preferred combinations of torsion angles for ordered secondary structures (conformations), such as α helix, β turn, γ turn, or β sheet. Molecular flexibility is directly related to covalent and/or noncovalent bonding interactions within a particular peptide. Even modest chemical modifications by N^α -methyl, C^α -methyl or C^β -methyl can have significant consequences on the resultant conformation.

The N^α - C^α - C' scaffold may be transformed by introduction of olefin substitution (e.g., C^α - $C^\beta \rightarrow C=C$ or dehydroamino acid or insertion (e.g., C^α - $C' \rightarrow C^\alpha$ - $C=C$ - C' or vinylogous amino acid. Also the C^β carbon may be substituted to advance the design of so-called "chimeric" amino acids. Finally, with respect to N-substituted amides it is also noteworthy to mention the intriguing approach of replacing the traditional peptide scaffold by achiral N-substituted glycine building blocks. Overall, such N^α - C^α - C scaffold or C^α - C^β side chain modifications expand peptide-based molecular diversity (i.e., so-called "peptoid" libraries) as well as extend our 3-D structural knowledge of traditional ϕ - Ψ - χ space.

In one approach, such as disclosed by Sherman and Spatola, *J. Am. Chem. Soc.* 112: 433 (1990), one or more amide bonds are replaced in an essentially isosteric manner by a variety of chemical functional groups. For example, any amide linkage in any of the GPCR polypeptides can be replaced by a ketomethylene moiety, e.g. $(-C(=O)-CH_2-)$ for $(-C(=O)-NH-)$. A few of the known amide bond replacements include: aminomethylene or $\Psi[CH_2NH]$; ketomethylene or $\Psi[COCH_2]$; ethylene or $\Psi[CH_2CH_2]$; olefin or $\Psi[CH=CH]$; ether or $\Psi[CH_2O]$; thioether or $\Psi[CH_2S]$; tetrazole or $\Psi[CN_4]$; thiazole or $\Psi[thz]$; retroamide or $\Psi[NHCO]$; thioamide or $\Psi[CSNH]$; and ester or $\Psi[CO_2]$. These amide bond surrogates alter conformational and H-bonding properties that may be requisite for peptide molecular recognition and/or biological activity at receptor targets. Furthermore, such backbone replacements can impart metabolic stability towards peptidase cleavage

relative to the parent peptide. The discovery of yet other nonhydrolyzable amide bond isostere has particularly impacted the design of protease inhibitors, and these include: hydroxymethylene or $\Psi[\text{CH}(\text{OH})]$; hydroxyethylene or $\Psi[\text{CH}(\text{OH})\text{CH}_2]$ and $\Psi[\text{CH}_2\text{CH}(\text{OH})]$; dihydroxyethylene or $(\Psi[\text{CH}(\text{OH})\text{CH}(\text{OH})])$, hydroxyethylamine or $\Psi[\text{CH}(\text{OH})\text{CH}_2\text{N}]$, dihydroxyethylene and C_2 -symmetric hydroxymethylene. Such backbone modifications have been extremely effective, as they may represent transition state mimics or bioisosteres of the hypothetical tetrahedral intermediate (e.g., $\Psi[\text{C}(\text{OH})_2\text{NH}]$) for this class of proteolytic enzymes. Such derivatives are expected to have the property of increased stability to degradation by enzymes, and therefore possess advantages for the formulation of compounds which may have increased *in vivo* half lives, as administered by oral, intravenous, intramuscular, intraperitoneal, topical, rectal, intraocular, or other routes.

Both peptide backbone and side chain modifications may provide prototypic leads for the design of secondary structure mimicry, as typically suggested by the fact that substitution of D-amino acids, N^α -Me-amino acids, C_α -Me amino acids, and/or dehydroamino acids within a peptide lead may induce or stabilize regiospecific β -turn, γ -turn, β -sheet, or α -helix conformations. To date, a variety of secondary structure mimetics have been designed and incorporated in peptides or peptidomimetics. The β -turn has been of particular interest to the area of receptor-targeted peptidomimetic drug discovery. This secondary structural motif exists within a tetrapeptide sequence in which the first and fourth C_α atoms are $\leq 7 \text{ \AA}$ separated, and they are further characterized as to occur in a nonhelical region of the peptide sequence and to possess a ten-membered intramolecular H-bond between the i and $i+4$ amino acid residues. One of the initial approaches of significance to the design of β -turn mimetics was the monocyclic dipeptide-based template which employs side chain to backbone constraint at the $i+1$ and $i+2$ sites. Over the past decade a variety of other monocyclic or bicyclic templates have been developed as β -turn mimetics. Monocyclic β -turn mimetic has been described that illustrate the potential opportunity to design scaffolds that may incorporate each of the side chains (i , $i+1$, $i+2$ and $i+3$ positions), as well as five of the eight NH or C=O functionalities, within the parent tetrapeptide sequence, tetrapeptide sequence modeled in type I-IV β -turn conformations. Similarly, a benzodiazepine template has shown utility as a β -turn mimetic scaffold which also may be multisubstituted to simulate side chain

functionalization,, particularly at the i and $i+3$ positions of the corresponding tetrapeptide sequence modeled in type I-VI β -turn conformations. A recently reported γ -turn mimetic, illustrates an innovative approach to incorporate a retroamide surrogate between the i and $i+1$ amino acid residues with an ethylene bridge between the N^1 (i.e., nitrogen replacing the carbonyl C') and N atoms of the i and $i+2$ positions, and this template allows the possibility for all three side chains of the parent tripeptide sequence. Finally, the design of a β -sheet mimetic provides an attractive template to constrain the backbone of a peptide to that simulating an extended conformation. The β -sheet is of particular interest to the area of protease-targeted peptidomimetic drug discovery.

Aromatic amino acids may be replaced with D- or L-naphthylalanine, D- or L-phenylglycine, D- or L-2-thieneylalanine, D- or L-1-, 2-, 3- or 4-pyreneylalanine, D- or L-3-thieneylalanine, D- or L-(2-pyridinyl)-alanine, D- or L-(3-pyridinyl)-alanine, D- or L-(2-pyrazinyl)-alanine, D- or L-(4-isopropyl)-phenylglycine, D-(trifluoromethyl)-phenylglycine, D-(trifluoromethyl)-phenylalanine, D-p-fluorophenylalanine, D- or L-p-biphenylphenylalanine, D- or L-p-methoxybiphenylphenylalanine, D- or L-2-indole(alkyl)alanines, and D- or L-alkylainines where alkyl may be substituted or unsubstituted methyl, ethyl, propyl, hexyl, butyl, pentyl, isopropyl, iso-butyl, sec-isotyl, iso-pentyl, non-acidic amino acids, of C_1 - C_{20} .

Acidic amino acids can be substituted with non-carboxylate amino acids while maintaining a negative charge, and derivatives or analogs thereof, such as the non-limiting examples of (phosphono)alanine, glycine, leucine, isoleucine, threonine, or serine; or sulfated (e.g., $-SO_3H$) threonine, serine, tyrosine.

Other substitutions may include unnatural hydroxylated amino acids made by combining "alkyl" (as defined and exemplified herein) with any natural amino acid. Basic amino acids may be substituted with alkyl groups at any position of the naturally occurring amino acids lysine, arginine, ornithine, citrulline, or (guanidino)-acetic acid, or other (guanidino)alkyl-acetic acids, where "alkyl" is defined as above. Nitrile derivatives (e.g., containing the CN-moiety in place of $COOH$) may also be substituted for asparagine or glutamine, and methionine sulfoxide may be substituted for methionine. Methods of preparation of such peptide derivatives are well known to one skilled in the art.

In addition, any amino acid of said peptides can be replaced by the same amino acid but of the opposite chirality. Thus, any amino acid naturally occurring in the L-configuration (which may also be referred to as the R or S, depending upon the structure of the chemical entity) may be replaced with an amino acid of the same chemical structural type, but of the opposite caroled, generally referred to as the D- amino acid but which can additionally be referred to as the R- or the S-, depending upon its composition and chemical configuration. Such derivatives have the property of greatly increased stability co degradation by enzymes, and therefore are advantageous in the formulation of compounds which may have longer *in vivo* half lives, when administered by oral, intravenous, intramuscular, intraperitoneal, topical, rectal, intraocular, or other routes.

Additional amino acid modifications of amino acids of GPCR polypeptides of to the present invention may include the following: Cysteiny residues may be reacted with alpha-haloacetates (and corresponding amines), such as 2-chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteiny residues may also be derivatized by reaction with compounds such as bromotrifluoroacetone, alpha-bromo-beta-(5-imidozoyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

Histidyl residues may be derivatized by reaction with compounds such as diethylprocarbonate e.g., at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain, and para-bromophenacyl bromide may also be used; e.g., where the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0.

Lysiny and amino terminal residues may be reacted with compounds such as succinic or other carboxylic acid anhydrides. Derivatization with these agents is expected to have the effect of reversing the charge of the lysiny residues. Other suitable reagents for derivatizing alpha-amino-containing residues include compounds such as imidoesters/e.g., as methyl picolinimide; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4 pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginyl residues may be modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin according to known method steps. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

Tyrosyl residues may be modified by reaction with aromatic diazonium compounds or tetranitromethane. N-acetylimidizol and tetranitromethane may be used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively.

Carboxyl side groups (aspartyl or glutamyl) may be selectively modified by reaction with carbodiimides (R'-N-C-N-R') such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore aspartyl and glutamyl residues may be converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

Glutaminyl and asparaginyl residues may be frequently deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues may be deamidated under mildly acidic conditions. Either form of these residues falls within the scope of the present invention.

Derivatization with bifunctional agents is useful for cross-linking the peptide to certain chemical moieties. Commonly used cross-linking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propioimide yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440

, may be employed for protein immobilization.

Other modifications of GPCR polypeptides of the present invention may include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups

of seryl or threonyl residues, methylation of the alpha-amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, *Proteins: Structure and Molecule Properties*, W. H. Freeman & Co., San Francisco, pp. 79-86 (1983)), acetylation of the N-terminal amine, methylation of main chain amide residues (or substitution with
5 N-methyl amino acids) and, in some instances, amidation of the C-terminal carboxyl groups, according to known method steps.

Such derivatized moieties may improve the solubility, absorption, permeability across the blood brain barrier biological half life, and the like. Such moieties or modifications of GPCR polypeptides may alternatively eliminate or
10 attenuate any possible undesirable side effect of the protein and the like. Moieties capable of mediating such effects are disclosed, for example, in Remington's *Pharmaceutical Sciences*, 16th ed., Mack Publishing Co., Easton, Pa. (1980).

Such chemical derivatives of GPCR polypeptides also may provide attachment to solid supports, including but not limited to, agarose, cellulose, hollow
15 fibers, or other polymeric carbohydrates such as agarose, cellulose, such as for purification, generation of antibodies or cloning; or to provide altered physical properties, such as resistance to enzymatic degradation or increased binding affinity or modulation for GPCRs, which is desired for therapeutic compositions comprising GPCR polypeptides, antibodies thereto or fragments thereof. Such peptide derivatives
20 are well-known in the art, as well as method steps for making such derivatives using carbodiimides active esters of N-hydroxy succinimide, or mixed anhydrides, as non-limiting examples.

Variation upon the sequences of GPCR polypeptides of the present invention may also include: the addition of one or more (e.g., two, three, four, or
25 five) lysine, arginine or other basic residues or one, or more (e.g., two, three, four, or five) glutamate or aspartate or other acidic residues at one end of the peptide, where "acidic" and "basic" are as defined herein. Negative charges can also be introduced by the addition of carboxyl, phosphate, borate, sulfonate or sulfate groups. Such modifications may increase the alpha-helical content of the peptide by the "helix
30 dipole effect". They also can provide enhanced aqueous solubility of the peptide, and allow the correct insertion of peptides into a membrane structure.

In another approach, a variety of uncoded or modified amino acids such as D-amino acids and N-methyl amino acids have been used to modify

mammalian peptides. Alternatively, a presumed bioactive conformation has been stabilized by a covalent modification, such as cyclization or by incorporation of gamma-lactam or other types of bridges. See, e.g., Veber and Hirschmann, *et al.*, *Proc. Natl. Acad. Sci. USA*, 1978 75 2636 and Thorsett, *et al.*, *Biochem Biophys. Res. Comm.*, 1983 111 166. The primary purpose of such manipulations has not been to avoid metabolism or to enhance oral bioavailability but rather to constrain a bioactive conformation to enhance potency or to induce greater specificity for a receptor subtype.

The above examples of peptide scaffold- or nonpeptide template-based peptidomimetic agonists or antagonists illustrate various strategies to elaborate bioactive conformation and/or pharmacophore models of peptide ligands at their receptors. In many cases, receptor subtype selectivity has also been achieved by systematic structural modifications of prototypic leads of peptidomimetics. Thus, although the 3D structures of GPCRs remains elusive (except for models constructed from homology-based low-resolution 3D structures of bacteriorhodopsin or rhodopsin, see below) the development of pharmacophore models using the hierarchical approach in peptide → peptidomimetic structure-based drug design is promising.

4. Purification of GPCR transmembrane peptides

The polypeptides of this invention may be purified to substantial purity by standard techniques, including selective precipitation with such substances as ammonium sulfate, column chromatography, immunopurification methods, and others. See, for instance, R. Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag: New York (1982). For example, the GPCR transmembrane peptides proteins and polypeptides produced by recombinant DNA technology are purified by a combination of cell lysis (e.g., sonication) and affinity chromatography or immunoprecipitation with a specific antibody to GPCR transmembrane peptides or a peptide fragment thereof. For fusion products, subsequent digestion of the fusion protein with an appropriate proteolytic enzyme releases the desired polypeptide. The proteins may then be further purified by standard protein chemistry techniques. A purified protein preferably exhibits a single band on an electrophoretic gel. Those of skill are reminded that the methods should take into account the hydrophobic nature of the peptides.

5. Detection of GPCR transmembrane peptide gene products

GPCR transmembrane peptides or a peptide fragment thereof to may be detected or quantified by a variety of methods. Preferred methods involve the use of specific antibodies.

5 a. Detection of GPCR transmembrane peptides by Immunoassay i. Antibody Production

Methods of producing polyclonal and monoclonal antibodies are known to those of skill in the art. See, e.g., Coligan (1991), CURRENT PROTOCOLS IN IMMUNOLOGY, Wiley/Greene, NY; and Harlow and Lane (1989), ANTIBODIES: A LABORATORY MANUAL, Cold Spring Harbor Press, NY; Stites et al. (eds.) BASIC AND CLINICAL IMMUNOLOGY (4th ed.) Lange Medical Publications, Los Altos, CA, and references cited therein; Goding (1986), MONOCLONAL ANTIBODIES: PRINCIPLES AND PRACTICE (2d ed.) Academic Press, New York, NY; and Kohler and Milstein (1975), *Nature*, 256:495-497. Such techniques include antibody preparation by selection of antibodies from libraries of recombinant antibodies in phage or similar vectors. See, Huse et al. (1989), *Science*, 246:1275-1281; and Ward et al. (1989) *Nature*, 341:544-546. For example, in order to produce antisera for use in an immunoassay, a polypeptide is isolated as described herein. For example, recombinant protein is produced in a transformed cell line. An inbred strain of mice or rabbits is immunized with the peptide using a standard adjuvant, such as Freund's adjuvant, and a standard immunization protocol. Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used an immunogen.

A number of immunogens may be used to produce antibodies specifically reactive with GPCR transmembrane peptides or a peptide fragment thereof. Recombinant protein is the preferred immunogen for the production of monoclonal or polyclonal antibodies. Naturally occurring protein may also be used either in pure or impure form. Synthetic peptides made using the GPCR transmembrane peptides or a peptide fragment thereof sequences described herein may also used as an immunogen for the production of antibodies to the protein. Recombinant protein can be expressed in eukaryotic or prokaryotic cells as described above, and purified as generally described above. The product is then injected into an animal capable of producing antibodies. Either monoclonal or polyclonal

antibodies may be generated, for subsequent use in immunoassays to measure the protein.

Methods of production of polyclonal antibodies are known to those of skill in the art. In brief, an immunogen, preferably a purified protein such as GPCR
5 transmembrane peptides or a peptide fragment thereof is mixed with an adjuvant and injected into an animal of choice (e.g., a mouse, rat, rabbit, pig, goat, cow, horse, chicken, etc.) at intervals of 1-4 weeks. The immunogen may be conjugated to a carrier protein can be used an immunogen. The animal's immune response to the immunogen preparation is monitored by taking test bleeds and determining the titer
10 of reactivity to the GPCR transmembrane peptides or a peptide fragment thereof. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the protein can be done if desired. (See Harlow and Lane, *supra*).

15 Polyclonal sera are collected and titered against the immunogen protein in an immunoassay, for example, a solid phase immunoassay with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of 10^4 or greater are selected and tested for their cross reactivity against non-GPCR transmembrane peptides or even GPCR transmembrane peptides from other cell types or species or a
20 peptide fragment thereof, using a competitive binding immunoassay (see, e.g., Harlow and Lane, *supra*, at pages 570-573). Specific monoclonal and polyclonal antibodies and antisera will usually bind with a K_D of at least about 0.1 mM, more usually at least about 1 μ M, preferably at least about .1 μ M or better, and most preferably, .01 μ M or better.

25 Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (See, Kohler and Milstein, *Eur. J. Immunol.* 6:511-519 (1976), incorporated herein by reference). Alternative methods of immortalization include transformation with
30 Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various

techniques, including injection into the peritoneal cavity of a vertebrate host. Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according to the general protocol outlined by Huse, et al. (1989) *Science* 246:1275-1281.

ii. Immunoassays

A particular protein can be measured by a variety of immunoassay methods. For a review of immunological and immunoassay procedures in general, see BASIC AND CLINICAL IMMUNOLOGY, 7th Edition (D. Stites and A. Terr, eds.) 1991. Moreover, the immunoassays of the present invention can be performed in any of several configurations, which are reviewed extensively in ENZYME IMMUNOASSAY, E.T. Maggio, ed., CRC Press, Boca Raton, Florida (1980); "Practice and Theory of Enzyme Immunoassays," P. Tijssen, in LABORATORY TECHNIQUES IN BIOCHEMISTRY AND MOLECULAR BIOLOGY, Elsevier Science Publishers B.V. Amsterdam (1985); and, Harlow and Lane, ANTIBODIES, A LABORATORY MANUAL, *supra*.

Immunoassays to GPCR transmembrane peptides, peptidomimetics or subfragments thereof may use a polyclonal antiserum raised against a peptide or peptidomimetic of the invention. This antiserum is selected to have low cross-reactivity against other (other non-GPCR transmembrane peptides or other GPCR transmembrane peptides) peptides and any such cross-reactivity is removed by immunoabsorption prior to use in the immunoassay.

Immunoassays in the competitive binding format can be used for the crossreactivity determinations. For example, a reference peptide antigen of the invention can be immobilized to a solid support. The ability of other molecules (other GPCR transmembrane peptides, or non-GPCR transmembrane peptides, or unknowns) to compete with the binding of antisera which recognize the immobilized reference peptide antigen is measured. The ability of such molecules to compete with the binding of an antiserum or antibody to the immobilized reference peptide is compared to a standard molecule, such as the reference peptide antigen itself. The percent crossreactivity is calculated, using standard calculations. Antisera with less than 10% crossreactivity to crossreacting molecules are selected and pooled. Any

cross-reacting antibodies are optionally removed from the pooled antisera by immunoabsorption with cross-reacting molecules.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay to compare the binding of a second protein to that of the reference peptide antigen. In order to make this comparison, the two molecules are each assayed at a wide range of concentrations and the amount of each molecule required to inhibit 50% of the binding of the antisera to the immobilized reference peptide antigen is determined. If the amount of the second protein required is less than 10 times the amount of the reference peptide used to make the antibody, then the second protein is said to specifically bind to an antibody generated to the reference peptide antigen.

The presence of a desired polypeptide (including peptide, translation product, or enzymatic digestion product) in a sample may be detected and quantified using Western blot analysis. The technique generally comprises separating sample products by gel electrophoresis on the basis of molecular weight, transferring the separated proteins to a suitable solid support, (such as a nitrocellulose filter, a nylon filter, or derivatized nylon filter), and incubating the sample with labeling antibodies that specifically bind to the analyte protein. The labeling antibodies specifically bind to analyte on the solid support. These antibodies are directly labeled, or alternatively are subsequently detected using labeling agents such as antibodies (e.g., labeled sheep anti-mouse antibodies where the antibody to an analyte is a murine antibody) that specifically bind to the labeling antibody.

6. Detection of GPCR transmembrane peptide sequences, peptides and peptidomimetics that optimally inhibit GPCR biological properties and functions

Peptides or peptide variants of the invention that modulate biological activity of GPCRs are generally identified as follows. Peptide sequences are selected from the transmembrane domains of the GPCR to be targeted. The transmembrane domains are readily ascertained by the application of computer models to known sequences. Computer modeling and comparison with known transmembrane peptide sequences are also used to define the orientation of the peptide sequence in the membrane, thus allowing the determination of the end of the peptide sequence that is towards the extracellular aspect of the plasma membrane. The selection of a

preferred transmembrane domain to be targeted is largely empirical. We have found that peptides derived from transmembrane domain 2 are particularly effective inhibitors of GPCR function. Alternatively, peptide sequences selected from transmembrane domain 4 have also been effective antagonists.

5 Upon selection of a peptide sequence, a reference transmembrane sequence is synthesized and systematically modified to identify variants (or analogs) that have improved properties. The modifications introduce a negative charge at the extracellular end of the peptide sequence. Negative charges may be added in the form of acidic amino acid residues such as Asp or Glu. The number of acidic
10 residues that is added is typically from 1 to 3 depending upon the hydrophobicity of the peptide sequence and the subsequent necessity to increase the solubility of the peptide. Further, preferable peptides have a neutral charge at the end of the peptide that is oriented towards the intracellular aspect of the plasma membrane. Thus, the overall hydrophobic nature of such a transmembrane peptide will result in insertion
15 into a membrane and the negative charge at the extracellular end will result in the peptide having the same orientation as the transmembrane GPCR domain from which it is derived. Insertion into the membrane may be tested by fluorescent microscopy of labeled peptide analogs using methodology known to those of skill in the art as illustrated in Example 3 herein.

20 The ability of the peptide or peptide variants to modulate activity of the targeted GPCR is generally determined by testing the ability of the peptide to inhibit activation that is induced by a natural ligand of the targeted GPCR. Activation of most GPCRs results in an increase in cAMP or the release of intracellular calcium. Thus, if activation of the target GPCR increases cAMP, the inhibitory activity of the
25 peptide is determined by measuring cAMP levels using methods known to those in the art (see e.g., C. Nordstedt and B.B. Fredholm *Anal. Chem.* 189: 231-234 (1990)). Similarly, if activation of the target GPCR releases intracellular calcium, the inhibitory activity of the peptide is determined by measuring the intracellular calcium levels as illustrated in the examples below.

30 Peptides may be tested for other properties including the following:

- enhanced ability to modulate GPCR activity;
- increased resistance to proteolysis;
- improved solubility;

- longer or shorter half-life, particularly in culture medium or a biological fluid such as plasma or whole blood;
- improved ability to insert into a membrane compartment, especially in a particular orientation, by means known in the art.

5 Variant peptides may also be synthesized having any one or more of the following modifications:

- conservative or non-conservative substitution of any of the amino acid residues;
- deletion or addition of residues at any position;
- 10 - chemical modification at any residue;
- peptidomimetic analogs of the reference peptide.

Variant peptides can be rationally designed and/or screened for using high throughput screening methodologies applied to combinatorial libraries. Methods of generating combinatorial libraries and screening such libraries using high-throughput
15 methods are well known to those of skill in the art (see, e.g., Baum, *C&EN* (Feb. 7, 1994): 20-26 and references cited therein).

These variant peptides are also tested for the any of the above-listed properties. In general, a variant peptide is considered to have improved properties relative to the reference peptide if a given measurable property or parameter
20 associated with the peptide has a value that is at least 10%, preferably at least 30%, more preferably at least 75%, and most preferably at least 95% better than the value for the reference peptide.

The relative ability of the modified peptides (as compared to the reference peptide) to modulate a GPCR biological activity is tested as follows. A cell
25 line that expresses a GPCR and exhibits a GPCR-mediated biological activity is exposed to either the reference or the modified peptide under identical conditions, and the biological property of the GPCR is measured in the absence or presence of either peptide. Examples of cell lines, GPCRs expressed by the cell line, and GPCR-regulated properties measured include the following:

- 30 - any cell that stably expresses CXCR4, especially attached cells, , including cells that are genetically engineered to express CXCR4, including HeLa cells; CXCR4; stroma cell derived factor I-induced calcium flux;

- any cell that stably expresses CXCR4, especially attached cells, including cells that are genetically engineered to express CXCR4, including CM cells; CXCR4; HIV-1 infection;
- any cell that stably expresses CCKAR, especially attached cells, including cells that are genetically engineered to express CCKAR, such as CHO cells; CCKAR; cholecystokinin-induced calcium release;
- any cell that stably expresses human CCR5, especially attached cells, including cells that are genetically engineered to express CCR5, including HEK cells; CCR5; RANTES induced calcium release.

10 The inhibitory activity is measured by exposing GPCR-expressing cells to a range of concentrations of a test antagonist, and measuring a biological property or activity associated with that GPCR. The test concentrations can range from 1 nanomolar to 100 micromolar, depending on peptide solubility and affinity. Initial screening is performed using 10-fold dilutions, such as 50, 5, 0.5, 0.05 micromolar.

15 Then, the lowest active concentration is lowered in decrements of 10% to determine the lowest effective concentration. The property measured can be binding to a ligand (for example, binding of cholecystokinin octapeptide to CCKAR), or production of a measurable metabolic response (e.g., altered ion flux or translocation, altered phosphorylation, altered protein synthesis or degradation,

20 altered cellular morphology, altered secretion, altered production of particular components such as soluble inositol phosphates, binding of a virus and subsequent infection, tumor growth, chemotaxis, mitogenic response, cell growth activation, secretion, muscle contraction, vasopressing and vasodepressing activity, synaptic transmission, and release of intracellular calcium, etc.)

25 The following GPCRs have been reported to play a role in HIV infection:

STRL33 U.S. Application Publication No. 20030203450

CCRR5 U.S. Application Publication No. 20030203450

30 CCR2 *Proc. Natl. Acad. Sci. USA*: 2752-2756 (1994)

J. Biol. Chem. 270: 29671-29675 (1995)

CCR3: *J. Biol. Chem.* 270: 16491-16494 (1995)

CX3CR1: *DNA Cell Biol.* 14: 673-680 (1995)

The following is a list of transmembrane peptides that have GPCR antagonist properties:

From the GPCR CXCR4

F-2-2: LLFVITLPFWAVDAVANWYFGNDD (SEQ. ID No.: 1)

5 F-2-5: LLFVITLPFWAVDAVANDD-OH (SEQ. ID No.: 2)

F-4-2: VYVGWIPALLLTIPDFIFANDD-OH (SEQ. ID No.: 3)

F-6-1: VILILAFFACWLPYYIGISID-OH (SEQ. ID No.: 4)

F-7-3: DDEALAFFHCCLNPILYAFL-NH₂ (SEQ. ID No.: 5)

F-7-4: DDSITEALAFFHCCLNPILYAFL-NH₂ (SEQ. ID No.: 6)

10

From the GPCR CCR5

CCR5-TM-2-2: LFFL LTVPFWAHYAAAQWDFGDD (SEQ. ID No.: 7)

CCR5-TM-4-1: FGVVTSVITWVAVFASLPGIIFTSSDD (SEQ. ID No.: 8)

CCR5-TM-6-1: LIFTIMIVYFLFWAPYNIVLLLNTFQED (SEQ. ID No.: 9)

15 CCR5-TM-7-1: DDQAMQVTETLGMTHCCINPIIYAFV (SEQ. ID No.: 10)

From the GPCR CCR2

CCR2-TM-2-1: IYLLNLAISDLLFLITLPLWADD-OH (SEQ. ID No.: 11)

CCR2-TM-2-2: LLFLITLPLWAH SAANEWVFGNDD-OH (SEQ. ID No.: 12)

20 CCR2-TM-4-1: FGVVTSVITWLVAVF ASVPGIIFTDD (SEQ. ID No.: 13)

CCR2-TM-6-1: VIFTIMIVYFLFWTPYN IVILLNTFQED (SEQ. ID No.: 14)

CCR2-TM-7-1: DDATQVT ETLGMTHCCINPIIYAFV (SEQ. ID No.: 15)

From the GPCR CCR3

25 CCR3-TM-2-1: LLFLVTLPFW IHYVRGHNWVFGDDD (SEQ. ID No.: 16)

CCR3-TM-4-1: FGVITSIVTWGLAVLAALPEFI FYETED (SEQ. ID No.: 17)

CCR3-TM-6-1: IFVIMAVFFI FWTPYNVAILLSSYQSDD (SEQ. ID No.: 18)

CCR3-TM-7-1: DDLVMLVTEVIAYSHCCMNPVIYAFV (SEQ. ID No.: 19)

30 From the GPCR CCKAR

CCKAR-TM-1-6: DDEWQSALQILLYSIIFLLSVLGNTLVITV (SEQ. ID No.: 20)

CCKAR-TN-2-1: FLLSLAVSDLMLCLFCMPFNLP (SEQ. ID No.: 21)

CCKAR-TM-2-2: FLLSLAVSDLMLCLFCM PFNLIDD (SEQ. ID No.: 22)

CCKAR-TM-6-4: IVVLFFLCWMPIFSANAWRAYDTVDD (SEQ. ID No.: 23)

7. Treatment embodiments

The compositions containing the present GPCR transmembrane peptides, or a cocktail thereof (*i.e.*, with other molecules, including other peptides of the invention), can be administered for therapeutic treatments. The molecules of the present invention are used to protect a patient from pathologies associated with GPCR, by modulating the biological activities associated with the GPCR.

"Protection" from infection or disease as used herein is intended to encompass "prevention" or "treatment." "Treatment" involves administration of the protective composition to a patient exhibiting symptoms of a GPCR-associated pathology (for example, HIV-1 infection), so as to reduce or suppress the symptoms of the pathology. Other examples of GPCR-associated conditions that may be treated with the peptides of the invention include:

- cancer. For example, vasoactive intestinal peptide (VIP) receptor is known to be overexpressed in breast cancer and lung cancer, and VIP antagonists are known to inhibit cancer growth. Thus, a peptide of the invention is probably effective in inhibiting such cancers;
- antagonists of chemokine receptors as anti-inflammatory and anti asthma drugs;
- tissue rejection;
- neuropeptide Y receptor antagonists as anti-obesity drugs;
- dopamine receptor D4 antagonists as drugs for treatment of depression, attention deficit hyperactivity disorder and schizophrenia;
- antagonists of Corticotropin-Releasing Factor Receptor for the treatment of depression and anxiety related disorders;
- angiotensin receptor antagonists as a mean of blood pressure control;
- antagonists of gastrin-releasing peptide receptor, somatostatin and gastrin receptors as anti-neoplastic agents that slow down growth of endocrine tumors;
- antagonists of opiod receptors as pain killers.

a. Pharmaceutical compositions

The compositions for administration may be in the form of a solution, suspension, tablets, pill, capsule, powder, gel, cream, lotion, ointment, aerosol or the like. In a preferred embodiment, the compositions for administration comprise a

solution of the GPCR transmembrane peptides dissolved in a pharmaceutically acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers can be used, e.g., buffered saline and the like. These solutions are sterile and generally free of undesirable matter. These compositions may be sterilized by conventional, well known sterilization techniques. In certain embodiments, the GPCR transmembrane peptides are provided in powder form.

The GPCR transmembrane peptides and analogs may be combined with conventional excipient, such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium, carbonate, and the like. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents and the like, for example, sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate and the like. The concentration of the GPCR in these formulations can vary widely, and will be selected primarily based on fluid volumes, viscosities, body weight and the like in accordance with the particular mode of administration selected and the patient's needs.

b. Administration and dosage

The pharmaceutical composition or medium that comprises a GPCR transmembrane peptide is administered orally, parenterally, enterically, gastrically, topically, subcutaneously, rectally, locally or systemically. For example, the compounds can be injected into the bloodstream using a cannula or catheter; the vein or artery is selected to maximize delivery of cells to the affected tissue(s). Actual methods for preparing parenterally administrable compositions will be known or apparent to those skilled in the art and are described in more detail in such publications as *Remington's Pharmaceutical Science*, 15th ed., Mack Publishing Company, Easton, Pennsylvania (1980). It is recognized that the GPCR transmembrane peptides polypeptides and related compounds described above, when administered orally, must be protected from digestion. This is typically accomplished either by complexing the protein with a composition to render it resistant to acidic and enzymatic hydrolysis or by packaging the protein in an appropriately resistant carrier such as a liposome. Means of protecting proteins from digestion are well known in the art.

In therapeutic applications, compositions are administered to a patient suffering from a disease or condition that in an amount sufficient to cure or at least partially arrest symptoms of the disease or conditions and its complications. An amount adequate to accomplish this is defined as a "therapeutically effective dose." Amounts effective for this use will depend upon the severity of the condition to be treated and the general state of the patient's health.

Generally, the dosage to be administered is the amount necessary to modulate a GPCR biological activity. It is understood that the dosage of a GPCR polypeptide of the present invention will be dependent upon the age, sex, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired. The ranges of effective doses provided herein are not intended to limit the inventors and represent preferred dose ranges. The most preferred dosage will be tailored to the individual subject, as is understood and determinable by one of skill in the art, without undue experimentation. It is contemplated that the compounds will be administered under the guidance of a physician, who will determine the exact dosages, monitor the progress of the treatment, and determine whether a given administration is successful and sufficient, or whether subsequent administrations are needed.

The concentration of compounds to be administered at a given time and to a given patient will vary from 0.1 μg -100 mg and preferably 0.1-10 mg per day per patient. The dosage and mode of administration may be chosen to achieve and optionally maintain a local concentration in fluids that contact the target cells of about 0.001-50 $\mu\text{g/ml}$, preferably 0.1-10 $\mu\text{g/ml}$. Dosages from 0.1 up to about 100 mg per patient per day may be used, particularly when the drug is administered to a secluded site and not into the blood stream, such as into a body cavity or into a lumen of an organ. Substantially higher dosages are possible in topical administration.

Single or multiple administrations of the compositions may be necessary depending on the dosage and frequency as required and tolerated by the patient. In any event, the composition should provide a sufficient quantity of the peptides of this invention to effectively treat the patient.

c. **Gene therapy**

The present invention provides packageable GPCR transmembrane peptide-encoding nucleic acids for the transformation of cells *in vitro* and *in vivo*. These packageable nucleic acids can be inserted into any of a number of well known
5 vectors for the transfection and transformation of target cells and organisms.

The nucleic acids are transfected into cells, *ex vivo* or *in vivo*, through the interaction of the vector and the target cell. The GPCR transmembrane peptide-encoding nucleic acid, under the control of a promoter, then expresses the GPCR transmembrane peptide, thereby modulating the biological activity of a target GPCR.

10 Such gene therapy procedures have been used to correct acquired and inherited genetic defects, cancer, and viral infection in a number of contexts. The ability to express artificial genes in humans facilitates the prevention and/or cure of many important human diseases, including many diseases which are not amenable to treatment by other therapies. As an example, *in vivo* expression of
15 cholesterol-regulating genes, genes which selectively block the replication of HIV, and tumor-suppressing genes in human patients dramatically improves the treatment of heart disease, AIDS, and cancer, respectively. For a review of gene therapy procedures, see Anderson, *Science* (1992) 256:808-813; Nabel and Felgner (1993) *TIBTECH* 11: 211-217; Mitani and Caskey (1993) *TIBTECH* 11: 162-166; Mulligan (1993) *Science* 926-932; Dillon (1993) *TIBTECH* 11: 167-175; Miller (1992) *Nature* 357: 455-460; Van Brunt (1988) *Biotechnology* 6(10): 1149-1154; Vigne (1995) *Restorative Neurology and Neuroscience* 8: 35-36; Kremer and Perricaudet (1995) *British Medical Bulletin* 51(1) 31-44; Haddada et al. (1995) in *CURRENT TOPICS IN MICROBIOLOGY AND IMMUNOLOGY* Doerfler and Böhm (eds) Springer-Verlag,
20 Heidelberg Germany; and Yu et al., *GENE THERAPY* (1994) 1:13-26.

25 Delivery of the gene or genetic material into the cell is the first critical step in gene therapy treatment of disease. A large number of delivery methods are well known to those of skill in the art. Such methods include, for example liposome-based gene delivery (Debs and Zhu (1993) WO 93/24640; Mannino and
30 Gould-Fogerite (1988) *BioTechniques* 6(7): 682-691; Rose U.S. Pat No. 5,279,833; Brigham (1991) WO 91/06309; and Felgner et al. (1987) *Proc. Natl. Acad. Sci. USA* 84: 7413-7414), and replication-defective retroviral vectors harboring a therapeutic polynucleotide sequence as part of the retroviral genome (see, e.g., Miller et al.

(1990) *Mol. Cell. Biol.* 10:4239 (1990); Kolberg (1992) *J. NIH Res.* 4:43, and Cornetta et al. *Hum. Gene Ther.* 2:215 (1991)). Widely used retroviral vectors include those based upon murine leukemia virus (MuLV), gibbon ape leukemia virus (GaLV), Simian Immuno deficiency virus (SIV), human immuno deficiency virus (HIV), and combinations thereof. See, e.g., Buchscher et al. (1992) *J. Virol.* 66(5) 2731-2739; Johann et al. (1992) *J. Virol.* 66 (5):1635-1640 (1992); Sommerfelt et al., (1990) *Virol.* 176:58- 59; Wilson et al. (1989) *J. Virol.* 63:2374-2378; Miller et al., *J. Virol.* 65:2220-2224 (1991); Wong-Staal et al., PCT/US94/05700, and Rosenberg and Fauci (1993) in *Fundamental Immunology, Third Edition* Paul (ed) Raven Press, Ltd., New York and the references therein, and Y et al., *GENE THERAPY* (1994), *supra*.

AAV-based vectors are also used to transduce cells with target nucleic acids, e.g., in the *in vitro* production of nucleic acids and peptides, and in *in vivo* and *ex vivo* gene therapy procedures. See, West et al. (1987) *Virology* 160:38- 47; Carter et al. (1989) U.S. Patent No. 4,797,368; Carter et al. WO 93/24641 (1993); Kotin (1994) *Human Gene Therapy* 5:793- 801; Muzyczka (1994) *J. Clin. Invest.* 94:1351 and Samulski (*supra*) for an overview of AAV vectors. Construction of recombinant AAV vectors are described in a number of publications, including Lebkowski, U.S. Pat. No. 5,173,414; Tratschin et al. (1985) *Mol. Cell. Biol.* 5(11):3251-3260; Tratschin, et al. (1984) *Mol. Cell. Biol.*, 4:2072-2081; Hermonat and Muzyczka (1984) *Proc. Natl. Acad. Sci. USA*, 81:6466-6470; McLaughlin et al. (1988) and Samulski et al. (1989) *J. Virol.*, 63:03822-3828. Cell lines that can be transformed by rAAV include those described in Lebkowski et al. (1988) *Mol. Cell. Biol.*, 8:3988-3996.

i. in vitro gene transfer

It is expected that those of skill in the art are knowledgeable in the numerous expression systems available for expression of DNA encoding GPCR transmembrane peptides or a peptide fragment thereof. No attempt to describe in detail the various methods known for the expression of proteins in prokaryotes or eukaryotes is made here.

There are several well-known methods of introducing nucleic acids into bacterial and animal cells, any of which may be used in the present invention. These include: calcium phosphate precipitation, fusion of the recipient cells with bacterial protoplasts containing the DNA, treatment of the recipient cells with

liposomes containing the DNA, DEAE dextran, receptor-mediated endocytosis, electroporation, micro-injection of the DNA directly into the cells, infection with viral vectors, etc.

For *in vitro* applications, the delivery of nucleic acids can be to any
5 cell grown in culture, whether of bacterial, plant or animal origin, vertebrate or invertebrate, and of any tissue or type. Contact between the cells and the genetically engineered nucleic acid constructs, when carried out *in vitro*, takes place in a biologically compatible medium. The concentration of nucleic acid varies widely depending on the particular application, but is generally between about 1 μ mol and
10 about 10 mmol. Treatment of the cells with the nucleic acid is generally carried out at physiological temperatures (about 37° C) for periods of time of from about 1 to 48 hours, preferably of from about 2 to 4 hours.

In one group of embodiments, a nucleic acid is added to 60-80% confluent plated cells having a cell density of from about 10^3 to about 10^5 cells/mL, more preferably about 2×10^4 cells/mL. The concentration of the suspension added
15 to the cells is preferably of from about 0.01 to 0.2 μ g/mL, more preferably about 0.1 μ g/mL.

ii. *In vivo* gene transfer

Alternatively, the GPCR transmembrane peptide encoding nucleic acids
20 can also be introduced into target cells *in vivo*, using recombinant methods which are known to those of skill in the art. The insertion of genes into cells for the purpose of medicinal therapy is a rapidly growing field in medicine which has enormous clinical potential. Research in gene therapy has been on-going for several years, and has entered human clinical trials. Zhu, et al., *Science*, 261:209-211
25 (1993), incorporated herein by reference, describes the intravenous delivery of cytomegalovirus (CMV)-chloramphenicol acetyltransferase (CAT) expression plasmid using DOTMA-DOPE complexes. Hyde, et al., *Nature*, 362:250-256 (1993), describes the delivery of the cystic fibrosis transmembrane conductance regulator (CFTR) gene to epithelia of the airway and to
30 alveoli in the lung of mice, using liposomes. Brigham, et al., *Am. J. Med. Sci.*, 298:278-281 (1989), describes the *in vivo* transfection of lungs of mice with a functioning prokaryotic gene encoding the intracellular enzyme chloramphenicol acetyltransferase (CAT).

Formulations suitable for administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can
5 include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The formulations of packaged nucleic acid can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials. Injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described.

10 For *in vivo* administration, pharmaceutical compositions that comprise GPCR transmembrane peptide-encoding nucleic acids are preferably administered parenterally, *i.e.*, intraarticularly, intravenously, intraperitoneally, subcutaneously, or intramuscularly. More preferably, the pharmaceutical compositions are administered intravenously or intraperitoneally by a bolus injection. For example, see Stadler, et
15 *al.*, U.S. Patent No. 5,286,634.

Intracellular nucleic acid delivery has also been discussed in Straubinger, *et al.*, *Methods in Enzymology*, Academic Press, New York. 101:512-527 (1983); Mannino, *et al.*, *Biotechniques*, 6:682-690 (1988); Nicolau, *et al.*, *Crit. Rev. Ther. Drug Carrier Syst.*, 6:239-271 (1989), and Behr, *Acc. Chem. Res.*, 26:274-278 (1993). Still other
20 methods of administering therapeutics are described in, for example, Rahman *et al.*, U.S. Patent No. 3,993,754; Sears, U.S. Patent No. 4,145,410; Papahadjopoulos *et al.*, U.S. Patent No. 4,235,871; Schneider, U.S. Patent No. 4,224,179; Lenk *et al.*, U.S. Patent No. 4,522,803; and Fountain *et al.*, U.S. Patent No. 4,588,578.

In preferred embodiments, the pharmaceutical preparations may be
25 contacted with the target tissue by direct application of the preparation to the tissue. The application may be made by topical, "open" or "closed" procedures. By "topical", it is meant the direct application of the pharmaceutical preparation to a tissue exposed to the environment, such as the skin, oropharynx, external auditory canal, and the like. "Open" procedures are those procedures which include incising
30 the skin of a patient and directly visualizing the underlying tissue to which the pharmaceutical preparations are applied. This is generally accomplished by a surgical procedure, such as a thoracotomy to access the lungs, abdominal laparotomy to access abdominal viscera, or other direct surgical approach to the target tissue.

"Closed" procedures are invasive procedures in which the internal target tissues are not directly visualized, but accessed via inserting instruments through small wounds in the skin. For example, the preparations may be administered to the peritoneum by needle lavage. Likewise, the preparations may be administered through
5 endoscopic devices.

The nucleic acid can also be administered in an aerosol inhaled into the lungs (see, Brigham, *et al.*, *Am. J. Sci.*, 298(4):278-281 (1989)) or by direct injection at the site of disease (Culver, *Human Gene Therapy*, MaryAnn Liebert, Inc., Publishers, New York. pp.70-71 (1994)).

10 Effective doses of the compositions of the present invention will vary depending upon many different factors, including means of administration, target site, physiological state of the patient, and other medicants administered. Thus, treatment dosages will need to be titrated to optimize safety and efficacy. In determining the effective amount of the vector to be administered, the physician evaluates the
15 particular nucleic acid used, the disease state being diagnosed; the age, weight, and condition of the patient, circulating plasma levels, vector toxicities, progression of the disease, and the production of anti-vector antibodies. The size of the dose also will be determined by the existence, nature, and extent of any adverse side-effects that accompany the administration of a particular vector. Doses ranging from about 10
20 ng to 1 g, 100 ng to 100 mg, 1 μ g to 10 mg, or 30-300 μ g DNA per patient are typical. Doses generally range between about 0.01 and about 50 mg per kilogram of body weight; preferably between about 0.1 and about 5 mg/kg of body weight or about 10^8 - 10^{10} or 10^{12} particles per injection. In general, the dose equivalent of a naked nucleic acid from a vector is from about 1 μ g to 100 μ g for a typical 70
25 kilogram patient, and doses of vectors which include a retroviral particle are calculated to yield an equivalent amount of inhibitor nucleic acid.

Prior to infusion, blood samples are obtained and saved for analysis. Between 10^8 and 1×10^{12} vectors are infused intravenously over 60-200 minutes. Vital signs and oxygen saturation by pulse oximetry are closely monitored. Blood
30 samples are obtained 5 minutes and 1 hour following infusion and saved for subsequent analysis. At the physician's discretion, reinfusion is repeated every 2 to 3 months for a total of 4 to 6 treatments in a one year period. After the first treatment, infusions can be performed on a outpatient basis at the discretion of the clinician. If

the reinfusion is given as an outpatient, the participant is monitored for at least 4, and preferably 8 hours following the therapy.

If a patient undergoing infusion of a vector or transduced cell develops fevers, chills, or muscle aches, he/she receives the appropriate dose of aspirin, ibuprofen or acetaminophen. Patients who experience reactions to the infusion such as fever, muscle aches, and chills are premedicated 30 minutes prior to the future infusions with either aspirin, acetaminophen, or diphenhydramine. Meperidine is used for more severe chills and muscle aches that do not quickly respond to antipyretics and antihistamines. Vector infusion is slowed or discontinued depending upon the severity of the reaction.

In vivo gene transfer may be practiced in a variety of hosts. Preferred hosts include mammalian species, such as humans, non-human primates, dogs, cats, cattle, horses, sheep, and the like.

15

EXAMPLES

The following examples are simply embodiments of the invention and are not intended to limit the invention. A person of ordinary skill in the art can modify and/or adapt the invention for various applications without undue experimentation, without departing from the generic concept of the present invention. Therefore, such adaptations and modifications are within the scope and range of the present invention.

Example 1

Example 1 illustrates that peptides derived from transmembrane regions of CXCR4 inhibit CXCR4-mediated calcium fluxes.

Peptides having the selected sequences were synthesized by a flow-through solid phase peptide synthesis on 432A Applied Biosystems Peptide Synthesizer utilizing Fmoc amino acid derivatives. To overcome the aggregation that frequently occurs during the synthesis of hydrophobic peptides and leads to the blockage of the growing peptide chain, FmocHmb derivatives of Ala, Val and Leu were introduced into the difficult sequences. Charged residues were added to the peptide termini to assure a proper orientation of the peptides during penetration into

30

the cellular membrane, and to improve the solubility of the highly hydrophobic peptides.

The purity of the peptides was assessed by reverse phase HPLC and the structures were confirmed by matrix-assisted laser-desorption time-of-flight (MALDI-TOF) mass spectrometry (Tarasova et al. (1998), *Ad. Exp. Med. Biol.*, Plenum Press, NY, pp. 201-206.)

Peptides used in this example are listed in Tables 1, 2, and 3.

The effect of the peptides on CXCR4-mediated calcium fluxes in HeLa cells that naturally express the CXCR4 receptor and U87 cells stably expressing the CXCR4 receptor was tested as follows. Cells grown on Nunc cover glass chamber slides were incubated with 1 micromolar Fura-2/AM for 20 min in a CO₂-incubator, rinsed with PBS and mounted on the stage of a Zeiss Axiovert inverted microscope. [Ca²⁺]_i measurements were performed using an Attofluor digital imaging system (Atto Instruments, Rockville, MD). Fluorescence of Fura was excited at alternating wavelength of 340 and 380 nm. Fluorescence was monitored by an intensified CCD camera using a 505 cut-off filter. Calibrations of [Ca²⁺]_i signals were performed using Ca²⁺ standards containing 1 micromolar Fura. CXCR4 antagonists were tested on HeLa cells and U87 cells. Stromal cell-derived factor-1 α (SDF-1 α) was used as a specific CXCR4 agonist.

CCR5 antagonists, which were used in selectivity studies as described below, were tested on HEK (human kidney carcinoma) cells stably expressing the CCR5 receptor and RANTES was used as an agonist. The antagonist activity of the peptides was evaluated by measuring the inhibition of agonist-evoked intracellular Ca²⁺_i release. These measurements were carried out in Fura-2/AM-treated cells, utilizing an Attofluor digital imaging system as described above. The agonist was SDF-1 α .

In the preliminary screen, peptides corresponding to the second and sixth transmembrane domains were found to abolish SDF-1 α - induced signaling through CXCR4 receptor (Table 1). Further optimization and structure-activity studies allowed to obtain antagonists derived from all but the third and fifth transmembrane domains (Table 2).

Table 1. Activity of synthetic peptides corresponding to predicted transmembrane domains of CXCR4 in inhibition of SDF-1 α -induced intracellular calcium release.

5	Peptide	Concentration, required for complete inhibition of [Ca ²⁺] _i release
	F-1-5: DDIFLPTIYSIIFLTGIV-HN ₂ (SEQ. ID No.: 350)	> 30 μ M
10	F-2-1: LLFVITLPFWAVDAVANWYFGN-OH (SEQ. ID No.: 351)	5 μ M
	F-3-1: KAVHVIYTVNLYSSVLILAFISL-NH ₂ (SEQ. ID No.: 352)	> 50 μ M
15	F-4-1 KVYVGWIPALLLTIPDFIF-OH (SEQ. ID No.: 353)	> 50 μ M
	F-5-1 HIMVGLILPGIVILSCYCIII-NH ₂ (SEQ. ID No.: 354)	> 50 μ M
20	F-6-1 VILILAFFACWLPYYIGISID-OH (SEQ. ID No.: 4)	10 μ M
25	F-7-1 ALAFFHCCLNPILYAFLGAK-NH ₂ (SEQ. ID No.: 355)	> 100 μ M

Table 2. Biological activity of CXCR4 antagonists derived from different transmembrane domains. Anti signaling activity was determined in inhibition of SDF-1 α -induced intracellular calcium release. Anti-HIV-1 activity was assessed in cytoprotection assay utilizing CEM-SS cells infected with HIV-1_{RF}.

Peptide	Concentration, required for inhibition of signal transduction (μ M)	EC ₅₀ in anti-HIV-1 assay (μ M)
F-2-2 LLFVITLPFWAVDAVANWYFGNDD (SEQ. ID No.: 1)	0.2	2.27
F-4-2 VYVGWIPALLLTIPDFIFANDD-OH (SEQ. ID. No.: 3)	5	0.3
F-6-1 VILILAFFACWLPYYIGISID-OH (SEQ. ID No.: 4)	10	>50
F-7-3 DDEALAFFHCCLNPILYAFL-NH ₂ (SEQ. ID No.: 5)	25	3.27
F-6-1 + F-7-3	1	No data

To further understand the structural requirements for a successful antagonist, structure-activity studies were conducted on the peptides derived from the second transmembrane domain of CXCR4 (Table 3). The most potent antagonist, a 24 amino acid residue peptide F-2-2, completely blocked signal transduction at 0.2 micromolar. Addition of negatively charged residues to the termini appeared to be important for the activity. Elimination of the added negative charges provided by two C-terminal Asp residues (F-2-1) decreased antagonist potency more than ten-fold. Consistent with those findings, the substitution of negatively charged aspartate residues with positively charged lysines (F-2-3) resulted in 100-fold decrease in antagonist activity. Deletion of five residues preceding the C-terminal aspartates (F-2-4) reduced the potency 20 -fold. Truncation of the transmembrane portion by three N-terminal residues Leu-Leu-Phe rendered the peptide inactive.

Table 3. Structure-activity relationships in peptides derived from the second transmembrane domain of CXCR4 :

...HLSVADLLFVITLPFWAVDAVANWYFGNFLCK...(SEQ. ID No.: 356) (predicted intramembrane portion is underlined)

5

10

15

20

25

Peptide	Concentration, required for complete inhibition of [Ca ²⁺] release
F-2-1: LLFVITLPFWAVDAVANWYFGN-OH (SEQ. ID No.: 351)	5 μ M
F-2-2: LLFVITLPFWAVDAVANWYFGNDD-OH (SEQ. ID No.: 1)	0.2 μ M
F-2-8: LLFVITLPFWAVDAVANWYFGNKK-OH (SEQ. ID No.: 357)	20 μ M
F-2-4: VITLPFWAVDAVANWYFGNKK-OH (SEQ. ID No.: 358)	> 50 μ M
F-2-5: LLFVITLPFWAVDAVANDD-OH (SEQ. ID No.: 2)	10 μ M
AcF-2-5: AcLLFVITLPFWAVDAVANDD-OH (SEQ. ID No.: 359)	10 μ M
F-2-6: LSVADLLFVITLPFWAVDAVANDD-OH (SEQ. ID No.: 360)	20 μ M
Rhod - F-2: AcLLFVITLPFWAVDAVANWYFGNDDK(Rhod)D-OH (SEQ. ID No.: 361)	8 μ M

30

Similar results were also observed for peptides derived from additional transmembrane regions. For example, in the case of peptides corresponding to the fourth transmembrane domain, positioning of the charged residue at the intracellular end of the peptide (F-4-1, Table 1) instead of the extracellular end (F-4-2, Table 2) abolished the antagonist activity. Further, substitution of extracellular aspartates with lysines also abolished the antagonist activity (data not shown).

The specificity of the transmembrane domain interaction was demonstrated by the fact that all peptides derived from CXCR4 showed selectivity for that receptor and had no influence on signaling of the other chemokine receptor involved in HIV-1 entry, CCR5. Similarly, a peptide derived from the second transmembrane domain of CCR5, LFFLLTVPFWAHYAAAQWDFGDD (SEQ. ID No.: 7), completely abolished agonist induced signaling of the receptor in U87 cells at 500 nM concentrations, but had no effect on signaling of CXCR4.

It was further noted that an equimolar mixture of two peptides, F-6-1 and F-7-3, was an order of magnitude more potent than the most active of the two peptides. This synergistic effect produced by the derivatives of the sixth and seventh transmembrane regions may be a general phenomenon. Thus, pairs of TM analogs in optimized combinations may act as very potent antagonists.

Example 2

Example 2 illustrates that synthetic peptides corresponding to transmembrane domains of CXCR4 inhibit CXCR4-mediated HIV infection.

CCR5 and CXCR4 are believed to be the main co-receptors for HIV-1 cell entry (Broder et al. (1997), *J. Leukoc. Biol.* 62:2029; Doranz et al. (1997) *Immunol. Res.* 16: 1528; Premack and Schall (1996), *Nat. Med.* 2:1174-1178.), although other chemokine receptors appear to mediate infection as well (Michael et al. (1997) *Nat. Med.* 3(10):11602).

The ability of synthetic CXCR4-derived peptides of Table 1 to inhibit HIV-1 infection of CEM-SS cells was tested using an LAV strain of the virus that is known to utilize CXCR4 as a co-receptor. Anti-HIV-1 assay. Buckheit et al. (1993) *Antiviral Research* 21: 247. The CEM-SS cells were maintained in RPMI 1640 medium containing 10% fetal bovine serum. The cells were placed in each well of a 96-well microtiter plate to a density of 5×10^3 cells per well. The cells were infected with HIV-1 virus at a multiplicity of infection (MOI) previously determined to produce maximal level of viral production at 6 days post infection (MOI of 0.01).

Serial half-log dilutions of test compound were added to appropriate wells in triplicate to evaluate their ability to inhibit HIV-1 infection. AZT was used in parallel as a positive control. Following 6 days of incubation at 37° C, the presence and relative abundance of viral p24 protein was determined by ELISA in cell-free supernatants derived from each well of the microtiter plate. The p24 ELISA kit was purchased from the AIDS Vaccine Program, NCI, FCRDC (Frederick, MD) and the assay was performed according to the manufacturer's instructions.

Most of the peptides in Table 1 showed some antiviral activity (data not shown). However, the peptides corresponding to the second and sixth transmembrane domains were the most potent in inhibition of HIV entry. The F-2-2 compound completely inhibited infection at a 5 micromolar concentration (Fig. 1).

Peptides corresponding to transmembrane domains of the cholecystikinin type A receptor (CCKAR) were used as negative controls and did not effect CXCR4 function, thereby confirming the specificity of the effect.

The peptides showed no cell toxicity in concentrations up to 100 micromolar (higher concentrations could not be tested because of solubility problems).

The ability of synthetic peptides to inhibit HIV-1 infection was additionally tested by cytoprotection assay using the highly cytopathic HIV-1 strain RF (Rice, *et al.* (1995) *Adv. Pharmacol.* 33:389,) (Table 2). The most potent peptide, F-4-2, completely inhibited infection at 1 micromolar concentration (Fig. 2). The peptides used as negative controls, which correspond to transmembrane domains of the cholecystikinin receptor type A, did not effect chemokine receptors functions.

The above results generally demonstrate the ability of externally added molecules to compete for interaction between transmembrane domains of GPCRs and thereby to disrupt receptor function (Figure 3). In addition, it is important to note that the peptides of the invention inhibit HIV infection by targeting a cellular molecule and function rather than a viral molecule and function. Viral proteins have a relatively high mutation rate, which often allows viruses to become resistant to a given treatment. Because cellular proteins mutate at a far slower rate, the probability that a virus will be able to develop a resistance is greatly reduced.

Example 3

Example 3 shows that the peptides of the invention partition to the plasma membrane and other membrane compartments.

A fluorescent derivative of CXCR4 TM2, rhodamine-F-2, was synthesized by solid-phase synthesis on 432A Applied Biosystems Peptide Synthesizer utilizing Fmoc amino acid derivatives. Rhodamine B (Fluka) was loaded onto the amino acid column of the instrument. The purity of the peptide was assessed by reverse phase HPLC and the
5 structures were confirmed by matrix-assisted laser desorption mass spectrometry.

A chimeric protein consisting of the CXCR4 and the green fluorescent protein (GFP) was used for studying receptor localization, internalization, and recycling in live cells in real time. This construct was made and stably expressed in HeLa cells as described in Tarasova et al. (1997), *J. Biol. Chem.* 272: 14817-14824. Fusion of the C terminus of
10 the CXCR4 to the N terminus of the GFP did not appear to alter receptor ligand binding affinity, signal transduction, or the pattern of receptor surface expression and distribution.

Transfected CXCR4-GFP-expressing HeLa cells were grown in coated 50 mm cover glass bottom dishes (MatTek, MA) in medium without phenol red. The cells were then exposed to 1 micromolar peptide in DMEM medium for 30 min in a CO₂-incubator.
15 The distribution of fluorescent label was determined by confocal laser scanning microscopy on a Zeiss inverted LSM 410 laser scanning confocal microscope. Fluorescence of GFP was excited using a 488 nm argon/krypton laser; emitted fluorescence was detected with 515-540 nm bandpass filter. For rhodamine red a 568 nm helium/neon laser was used for excitation and fluorescence was detected with a 590-640 nm bandpass filter.

20 The results demonstrated that the rhodaminated peptide co-localized with the CXCR4-GFP and was present at the cellular membrane within minutes after application and saturated endosomes and the endoplasmic reticulum after 15 minutes of incubation. This confirmed the ability of the peptides to concentrate in the cellular membranes and suggested that the peptides interacted with receptor molecules.

25 **Example 4**

Example 4 shows that peptides corresponding to transmembrane domains of the cholecystinin type A receptor (CCKAR) inhibits agonist-evoked intracellular calcium release with a potency similar to CXCR4 compounds.

To further illustrate the present invention, we have synthesized peptides
30 derived from the transmembrane domains of the rat cholecystinin receptor type A (CCKAR). Although CCKAR belongs to the same rhodopsin family of GPCRs as CXCR4, its sequence is only 15% identical to that of CXCR4, when aligned using the Dialign 2

program (Morgenstern, et al., (1996) *Proc. Natl. Acad. Sci. USA* 93:12098) and the degree of identify in transmembrane regions is only 27%.

The activity of peptides from the CCKAR transmembrane domain were tested in transfected CHO cells that stably express rat CCKAR (Tarasova et al. (1997), *J. Biol. Chem.* 272: 14817-14824). Sulfated cholecystinin octapeptide was the CCKAR agonist. Determination of intracellular calcium release was performed as described in example 1 above. The results are shown in Table 4.

None of CCKAR-derived peptides served as antagonists of chemokine receptors and none could inhibit HIV-1 infection. The activity of the antagonists in inhibiting signaling through the receptor was compared to the ability to prevent agonist binding (Table 4). Inhibition of signaling was assessed in CCK-8 evoked intracellular calcium release in FURA-2/AM treated CHO cells stably transfected with rat CCKAR (Tarasova, et al. (1997) *J. Biol. Chem.* 272:14817). Inhibition of ligand binding was measured with the use of a fluorescent agonist, rhodamine green CCK-8 (RG-CCK-8) and quantitative confocal laser scanning microscopy (Tarasova, et al. (1997) *J. Biol. Chem.* 272:14817). Peptides derived from the first, second, and sixth transmembrane domains inhibited CCK-induced signaling through the CCKAR receptor. Peptides derived from the first and the second transmembrane domains, CCKAR-1-1 and CCKAR-2-1, had comparable potencies with respect to the inhibition of ligand signaling and binding. A peptide derived from the sixth domain, CCKAR-6-1, was active in inhibition of signaling, but had very low activity in inhibition of RG-CCK-8 binding.

Table 4. The activity of CCKAR-derived TM peptides in inhibition of CCK-8 -induced intracellular calcium release and RG-CCK-8 binding.

5	Peptide	Concentration, required for inhibition of signaling	IC ₅₀ in inhibition of RG-CCK-8 binding
	CCKAR-TM-1-6: DDEWQSALQILLYSIIFLLSVLGNTLVITV (SEQ ID No.: 20)	50 μ M	20 μ M
10	CCKAR-TM-2-1: FLLSLAVSDLMLCLFCMPFNLP (SEQ ID No.: 21)	2 μ M	0.5 μ M
	CCKAR-TM-4-2 (#71) VIAATWCLSFTIMTPYPIYSNLVPFTDD (SEQ ID No.: 362)	> 50 μ M	> 50 μ M
15	CCKAR-TM-5-3 (#45) DDQTFLLLILFLLPGIVMVVAYGL (SEQ ID No.: 363)	> 50 μ M	> 50 μ M
20	CCKAR-TM-6-4 (#77) IVVLFFLCWMPIFSANAWRAYDTVDD (SEQ ID No.: 23)	5 μ M	> 50 μ M

25

64A

SEQUENCE LISTING

<110> Tarasova, Nadya I.
 Michejda, Christopher J.
 The Government of the United States of America
 as represented by the Secretary of the
 Department of Health and Human Services

<120> G Protein-Coupled Receptor Antagonists

<130> 40330-1644

<140> WO PCT/US99/04438

<141> 1999-02-26

<150> US 60/076,105

<151> 1998-02-27

<160> 363

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 24

<212> PRT

<213> Artificial Sequence

<220>

<223> F-2-2 GPCR CXCR4

<400> 1

Leu	Leu	Phe	Val	Ile	Thr	Leu	Pro	Phe	Trp	Ala	Val	Asp	Ala	Val	Ala
1				5					10					15	
Asn	Trp	Tyr	Phe	Gly	Asn	Asp	Asp								
			20												

<210> 2

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> F-2-5 GPCR CXCR4

<400> 2

Leu	Leu	Phe	Val	Ile	Thr	Leu	Pro	Phe	Trp	Ala	Val	Asp	Ala	Val	Ala
1				5					10					15	
Asn	Asp	Asp													

<210> 3

<211> 23

<212> PRT

<213> Artificial Sequence

<220>

<223> F-4-2 GPCR CXCR4

64B

<400> 3
 Val Tyr Val Gly Val Trp Ile Pro Ala Leu Leu Leu Thr Ile Pro Asp
 1 5 10 15
 Phe Ile Phe Ala Asn Asp Asp
 20

<210> 4
 <211> 21
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> F-6-1 GPCR CXCR4

<400> 4
 Val Ile Leu Ile Leu Ala Phe Phe Ala Cys Trp Leu Pro Tyr Tyr Ile
 1 5 10 15
 Gly Ile Ser Ile Asp
 20

<210> 5
 <211> 20
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> F-7-3 GPCR CXCR4

<221> MOD_RES
 <222> (20)...(20)
 <223> Xaa = leucinamide

<400> 5
 Asp Asp Glu Ala Leu Ala Phe Phe His Cys Cys Leu Asn Pro Ile Leu
 1 5 10 15
 Tyr Ala Phe Xaa
 20

<210> 6
 <211> 23
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> F-7-4 GPCR CXCR4

<221> MOD_RES
 <222> (23)...(23)
 <223> Xaa = leucinamide

<400> 6
 Asp Asp Ser Ile Thr Glu Ala Leu Ala Phe Phe His Cys Cys Leu Asn
 1 5 10 15
 Pro Ile Leu Tyr Ala Phe Xaa
 20

<210> 7
 <211> 23

64C

<212> PRT
<213> Artificial Sequence

<220>
<223> CCR5-TM-2-2 GPCR CCR5

<400> 7
Leu Phe Phe Leu Leu Thr Val Pro Phe Trp Ala His Tyr Ala Ala Ala
1 5 10 15
Gln Trp Asp Phe Gly Asp Asp
20

<210> 8
<211> 28
<212> PRT
<213> Artificial Sequence

<220>
<223> CCR5-TM-4-1 GPCR CCR5

<400> 8
Phe Gly Val Val Thr Ser Val Ile Thr Trp Val Val Ala Val Phe Ala
1 5 10 15
Ser Leu Pro Gly Ile Ile Phe Thr Ser Ser Asp Asp
20 25

<210> 9
<211> 28
<212> PRT
<213> Artificial Sequence

<220>
<223> CCR5-TM-6-1 GPCR CCR5

<400> 9
Leu Ile Phe Thr Ile Met Ile Val Tyr Phe Leu Phe Trp Ala Pro Tyr
1 5 10 15
Asn Ile Val Leu Leu Leu Asn Thr Phe Gln Glu Asp
20 25

<210> 10
<211> 26
<212> PRT
<213> Artificial Sequence

<220>
<223> CCR5-TM-7-1 GPCR CCR5

<400> 10
Asp Asp Gln Ala Met Gln Val Thr Glu Thr Leu Gly Met Thr His Cys
1 5 10 15
Cys Ile Asn Pro Ile Ile Tyr Ala Phe Val
20 25

<210> 11
<211> 23
<212> PRT
<213> Artificial Sequence

64D

<220>

<223> CCR2-TM-2-1 GPCR CCR2

<400> 11

Ile	Tyr	Leu	Leu	Asn	Leu	Ala	Ile	Ser	Asp	Leu	Leu	Phe	Leu	Ile	Thr
1				5					10					15	
Leu	Pro	Leu	Trp	Ala	Asp	Asp									
			20												

<210> 12

<211> 24

<212> PRT

<213> Artificial Sequence

<220>

<223> CCR2-TM-2-2 GPCR CCR2

<400> 12

Leu	Leu	Phe	Leu	Ile	Thr	Leu	Pro	Leu	Trp	Ala	His	Ser	Ala	Ala	Asn
1				5					10					15	
Glu	Trp	Val	Phe	Gly	Asn	Asp	Asp								
			20												

<210> 13

<211> 26

<212> PRT

<213> Artificial Sequence

<220>

<223> CCR2-TM-4-1 GPCR CCR2

<400> 13

Phe	Gly	Val	Val	Thr	Ser	Val	Ile	Thr	Trp	Leu	Val	Ala	Val	Phe	Ala
1				5					10					15	
Ser	Val	Pro	Gly	Ile	Ile	Phe	Thr	Asp	Asp						
			20					25							

<210> 14

<211> 28

<212> PRT

<213> Artificial Sequence

<220>

<223> CCR2-TM-6-1 GPCR CCR2

<400> 14

Val	Ile	Phe	Thr	Ile	Met	Ile	Val	Tyr	Phe	Leu	Phe	Trp	Thr	Pro	Tyr
1				5					10					15	
Asn	Ile	Val	Ile	Leu	Leu	Asn	Thr	Phe	Gln	Glu	Asp				
			20					25							

<210> 15

<211> 25

<212> PRT

<213> Artificial Sequence

<220>

<223> CCR2-TM-7-1 GPCR CCR2

64E

<400> 15
 Asp Asp Ala Thr Gln Val Thr Glu Thr Leu Gly Met Thr His Cys Cys
 1 5 10 15
 Ile Asn Pro Ile Ile Tyr Ala Phe Val
 20 25

<210> 16
 <211> 25
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCR3-TM-2-1 GPCR CCR3

<400> 16
 Leu Leu Phe Leu Val Thr Leu Pro Phe Trp Ile His Tyr Val Arg Gly
 1 5 10 15
 His Asn Trp Val Phe Gly Asp Asp Asp
 20 25

<210> 17
 <211> 28
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCR3-TM-4-1 GPCR CCR3

<400> 17
 Phe Gly Val Ile Thr Ser Ile Val Thr Trp Gly Leu Ala Val Leu Ala
 1 5 10 15
 Ala Leu Pro Glu Phe Ile Phe Tyr Glu Thr Glu Asp
 20 25

<210> 18
 <211> 28
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCR3-TM-6-1 GPCR CCR3

<400> 18
 Ile Phe Val Ile Met Ala Val Phe Phe Ile Phe Trp Thr Pro Tyr Asn
 1 5 10 15
 Val Ala Ile Leu Leu Ser Ser Tyr Gln Ser Asp Asp
 20 25

<210> 19
 <211> 26
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCR3-TM-7-1 GPCR CCR3

64F

<400> 19
 Asp Asp Leu Val Met Leu Val Thr Glu Val Ile Ala Tyr Ser His Cys
 1 5 10 15
 Cys Met Asn Pro Val Ile Tyr Ala Phe Val
 20 25

<210> 20
 <211> 30
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCKAR-TM-1-6 GPCR CCKAR

<400> 20
 Asp Asp Glu Trp Gln Ser Ala Leu Gln Ile Leu Leu Tyr Ser Ile Ile
 1 5 10 15
 Phe Leu Leu Ser Val Leu Gly Asn Thr Leu Val Ile Thr Val
 20 25 30

<210> 21
 <211> 22
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCKAR-TM-2-1 GPCR CCKAR

<400> 21
 Phe Leu Leu Ser Leu Ala Val Ser Asp Leu Met Leu Cys Leu Phe Cys
 1 5 10 15
 Met Pro Phe Asn Leu Pro
 20

<210> 22
 <211> 24
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCKAR-TM-2-2 GPCR CCKAR

<400> 22
 Phe Leu Leu Ser Leu Ala Val Ser Asp Leu Met Leu Cys Leu Phe Cys
 1 5 10 15
 Met Pro Phe Asn Leu Ile Asp Asp
 20

<210> 23
 <211> 26
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCKAR-TM-6-4 GPCR CCKAR

64G

<400> 23
 Ile Val Val Leu Phe Phe Leu Cys Trp Met Pro Ile Phe Ser Ala Asn
 1 5 10 15
 Ala Trp Arg Ala Tyr Asp Thr Val Asp Asp
 20 25

<210> 24
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> GPCRAelegans GPCR TM1

<400> 24
 His Pro Cys Glu Asp Ile Met Gly Tyr Val Trp Leu Thr Val Val Ser
 1 5 10 15
 Phe Met Val Gly Ala Val Ala Leu Val Ala Asn Leu Val Val Ala Leu
 20 25 30
 Val Leu Leu Thr Ser Gln Arg Arg Leu Asn Val
 35 40

<210> 25
 <211> 50
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> GRH GPCR TM1

<400> 25
 Asn Leu Pro Thr Leu Thr Leu Ser Gly Lys Ile Arg Val Thr Val Thr
 1 5 10 15
 Phe Phe Leu Phe Leu Leu Ser Ala Thr Phe Asn Ala Ser Phe Leu Leu
 20 25 30
 Lys Leu Gln Lys Trp Thr Gln Lys Lys Glu Lys Gly Lys Lys Leu Ser
 35 40 45
 Arg Met
 50

<210> 26
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> TRH GPCR TM1

<400> 26
 Arg Ala Val Val Ala Leu Glu Tyr Gln Val Val Thr Ile Leu Leu Val
 1 5 10 15
 Leu Ile Ile Cys Gly Leu Gly Ile Val Gly Asn Ile Met Val Val Leu
 20 25 30
 Val Val Met Arg Thr Lys His Met Arg Thr Pro
 35 40

<210> 27
 <211> 43
 <212> PRT

64H

<213> Artificial Sequence

<220>

<223> FSHprec GPCR TM1

<400> 27

Asn	Pro	Cys	Glu	Asp	Ile	Met	Gly	Tyr	Asn	Ile	Leu	Arg	Val	Leu	Ile
1				5				10						15	
Trp	Phe	Ile	Ser	Ile	Leu	Ala	Ile	Thr	Gly	Asn	Ile	Ile	Val	Leu	Val
			20					25					30		
Ile	Leu	Thr	Thr	Ser	Gln	Tyr	Lys	Leu	Thr	Val					
		35					40								

<210> 28

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> TSHprec GPCR TM1

<400> 28

Asn	Pro	Cys	Glu	Asp	Ile	Met	Gly	Tyr	Lys	Phe	Leu	Arg	Ile	Val	Val
1				5				10						15	
Trp	Phe	Val	Ser	Leu	Leu	Ala	Leu	Leu	Gly	Asn	Val	Phe	Val	Leu	Leu
			20					25					30		
Ile	Leu	Leu	Thr	Ser	His	Tyr	Lys	Leu	Asn	Val					
		35					40								

<210> 29

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> LH_CGprec GPCR TM1

<400> 29

Asn	Pro	Cys	Glu	Asp	Ile	Met	Gly	Tyr	Asp	Phe	Leu	Arg	Val	Leu	Ile
1				5				10						15	
Trp	Leu	Ile	Asn	Ile	Leu	Ala	Ile	Met	Gly	Asn	Met	Thr	Val	Leu	Phe
			20					25					30		
Val	Leu	Leu	Thr	Ser	Arg	Tyr	Lys	Leu	Thr	Val					
		35					40								

<210> 30

<211> 49

<212> PRT

<213> Artificial Sequence

<220>

<223> PGE_EP1 GPCR TM1

<400> 30

Pro	Asn	Thr	Ser	Ala	Val	Pro	Pro	Ser	Gly	Ala	Ser	Pro	Ala	Leu	Pro
1				5				10						15	
Ile	Phe	Ser	Met	Thr	Leu	Gly	Ala	Val	Ser	Asn	Leu	Leu	Ala	Leu	Ala
			20					25					30		

64I

Leu Leu Ala Gln Ala Ala Gly Arg Leu Arg Arg Arg Arg Ser Ala Thr
 35 40 45
 Thr

<210> 31
 <211> 45
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PGE_EP2 GPCR TM1

<400> 31
 Ser Ala Ser Leu Ser Pro Asp Arg Leu Asn Ser Pro Val Thr Ile Pro
 1 5 10 15
 Ala Val Met Phe Ile Phe Gly Val Val Gly Asn Leu Val Ala Ile Val
 20 25 30
 Val Leu Cys Lys Ser Arg Lys Glu Gln Lys Glu Thr Thr
 35 40 45

<210> 32
 <211> 45
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PGE_EP3 GPCR TM1

<400> 32
 Gln Trp Leu Pro Pro Gly Glu Ser Pro Ala Ile Ser Ser Val Met Phe
 1 5 10 15
 Ser Ala Gly Val Leu Gly Asn Leu Ile Ala Leu Ala Leu Leu Ala Arg
 20 25 30
 Arg Trp Arg Ser Ala Gly Arg Arg Ser Ser Leu Ser Leu
 35 40 45

<210> 33
 <211> 48
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PGF GPCR TM1

<400> 33
 Ser Asn Thr Thr Cys Gln Thr Glu Asn Arg Leu Ser Val Phe Phe Ser
 1 5 10 15
 Val Ile Phe Met Thr Val Gly Ile Leu Ser Asn Ser Leu Ala Ile Ala
 20 25 30
 Ile Leu Met Lys Ala Tyr Gln Arg Phe Arg Gln Lys Ser Lys Ala Ser
 35 40 45

<210> 34
 <211> 44
 <212> PRT
 <213> Artificial Sequence

<220>

64J

<223> PGI GPCR TM1

<400> 34

Cys	Arg	Asn	Leu	Thr	Tyr	Val	Arg	Gly	Ser	Val	Gly	Pro	Ala	Thr	Ser
1				5					10					15	
Thr	Leu	Met	Phe	Val	Ala	Gly	Val	Val	Gly	Asn	Gly	Leu	Ala	Leu	Gly
			20					25					30		
Ile	Leu	Ser	Ala	Arg	Arg	Pro	Ala	Arg	Pro	Ser	Ala				
		35					40								

<210> 35

<211> 47

<212> PRT

<213> Artificial Sequence

<220>

<223> TXA2 GPCR TM1

<400> 35

Asn	Ile	Thr	Leu	Glu	Glu	Arg	Arg	Leu	Ile	Ala	Ser	Pro	Trp	Phe	Ala
1				5					10					15	
Ala	Ser	Phe	Cys	Val	Val	Gly	Leu	Ala	Ser	Asn	Leu	Leu	Ala	Leu	Ser
			20					25					30		
Val	Leu	Ala	Gly	Ala	Arg	Gln	Gly	Gly	Ser	His	Thr	Arg	Ser	Ser	
		35					40					45			

<210> 36

<211> 46

<212> PRT

<213> Artificial Sequence

<220>

<223> PAF GPCR TM1

<400> 36

His	Met	Asp	Ser	Glu	Phe	Arg	Tyr	Thr	Leu	Phe	Pro	Ile	Val	Tyr	Ser
1				5					10					15	
Ile	Ile	Phe	Val	Leu	Gly	Val	Ile	Ala	Asn	Gly	Tyr	Val	Leu	Trp	Val
			20					25					30		
Phe	Ala	Arg	Leu	Tyr	Pro	Cys	Lys	Lys	Phe	Asn	Glu	Ile	Lys		
		35					40					45			

<210> 37

<211> 40

<212> PRT

<213> Artificial Sequence

<220>

<223> M2 GPCR TM1

<400> 37

Tyr	Lys	Thr	Phe	Glu	Val	Val	Phe	Ile	Val	Leu	Val	Ala	Gly	Ser	Leu
1				5					10					15	
Ser	Leu	Val	Thr	Ile	Ile	Gly	Asn	Ile	Leu	Val	Met	Val	Ser	Ile	Lys
			20					25					30		
Val	Asn	Arg	His	Leu	Gln	Thr	Val								
		35					40								

<210> 38

64K

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> M4 GPCR TM1

<400> 38
 His Asn Arg Tyr Glu Thr Val Glu Met Val Phe Ile Ala Thr Val Thr
 1 5 10 15
 Gly Ser Leu Ser Leu Val Thr Val Val Gly Asn Ile Leu Val Met Leu
 20 25 30
 Ser Ile Lys Val Asn Arg Gln Leu Gln Thr Val
 35 40

<210> 39
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> M1 GPCR TM1

<400> 39
 Gly Lys Gly Pro Trp Gln Val Ala Phe Ile Gly Ile Thr Thr Gly Leu
 1 5 10 15
 Leu Ser Leu Ala Thr Val Thr Gly Asn Leu Leu Val Leu Ile Ser Phe
 20 25 30
 Lys Val Asn Thr Glu Leu Lys Thr Val
 35 40

<210> 40
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> M3 GPCR TM1

<400> 40
 Leu Gly Gly His Thr Val Trp Gln Val Val Phe Ile Ala Phe Leu Thr
 1 5 10 15
 Gly Ile Leu Ala Leu Val Thr Ile Ile Gly Asn Ile Leu Val Ile Val
 20 25 30
 Ser Phe Lys Val Asn Lys Gln Leu Lys Thr Val
 35 40

<210> 41
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> M5 GPCR TM1

<400> 41
 Leu Glu Arg His Arg Leu Trp Glu Val Ile Thr Ile Ala Ala Val Thr
 1 5 10 15

64L

Ala Val Val Ser Leu Ile Thr Ile Val Gly Asn Val Leu Val Met Ile
 20 25 30
 Ser Phe Lys Val Asn Ser Gln Leu Lys Thr Val
 35 40

<210> 42
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> H1 GPCR TM1

<400> 42
 Lys Thr Thr Met Ala Ser Pro Gln Leu Met Pro Leu Val Val Val Leu
 1 5 10 15
 Ser Thr Ile Cys Leu Val Thr Val Gly Leu Asn Leu Leu Val Leu Tyr
 20 25 30
 Ala Val Arg Ser Glu Arg Lys Leu His Thr Val
 35 40

<210> 43
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> H2 GPCR TM1

<400> 43
 Phe Cys Leu Asp Ser Thr Ala Cys Lys Ile Thr Ile Thr Val Val Leu
 1 5 10 15
 Ala Val Leu Ile Leu Ile Thr Val Ala Gly Asn Val Val Val Cys Leu
 20 25 30
 Ala Val Gly Leu Asn Arg Arg Leu Arg Asn Leu
 35 40

<210> 44
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT1A GPCR TM1

<400> 44
 Ile Ser Asp Val Thr Val Ser Tyr Gln Val Ile Thr Ser Leu Leu Leu
 1 5 10 15
 Gly Thr Leu Ile Phe Cys Ala Val Leu Gly Asn Ala Cys Val Val Ala
 20 25 30
 Ala Ile Ala Leu Glu Arg Ser Leu Gln Asn Val
 35 40

<210> 45
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64M

<223> 5HT1B GPCR TM1

<400> 45

Gln	Asp	Ser	Ile	Ser	Leu	Pro	Trp	Lys	Val	Leu	Leu	Val	Met	Leu	Leu
1				5					10					15	
Ala	Leu	Ile	Thr	Leu	Ala	Thr	Thr	Leu	Ser	Asn	Ala	Phe	Val	Ile	Ala
			20					25					30		
Thr	Val	Tyr	Arg	Thr	Arg	Lys	Leu	His	Thr	Pro					
		35					40								

<210> 46

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> 5HT1D GPCR TM1

<400> 46

Asp	Pro	Arg	Thr	Leu	Gln	Ala	Leu	Lys	Ile	Ser	Leu	Ala	Val	Val	Leu
1				5					10					15	
Ser	Val	Ile	Thr	Leu	Ala	Thr	Val	Leu	Ser	Asn	Ala	Phe	Val	Leu	Thr
			20					25					30		
Thr	Ile	Leu	Leu	Thr	Arg	Lys	Leu	His	Thr	Pro					
		35					40								

<210> 47

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> 5HT1E GPCR TM1

<400> 47

Ile	Arg	Pro	Lys	Thr	Ile	Thr	Glu	Lys	Met	Leu	Ile	Cys	Met	Thr	Leu
1				5					10					15	
Val	Val	Ile	Thr	Thr	Leu	Thr	Thr	Leu	Leu	Asn	Leu	Ala	Val	Ile	Met
			20					25					30		
Ala	Ile	Gly	Thr	Thr	Lys	Lys	Leu	His	Gln	Pro					
		35					40								

<210> 48

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> 5HT1F GPCR TM1

<400> 48

Glu	Leu	Leu	Asn	Arg	Met	Pro	Ser	Lys	Ile	Leu	Val	Ser	Leu	Thr	Leu
1				5					10					15	
Ser	Gly	Leu	Ala	Leu	Met	Thr	Thr	Thr	Ile	Asn	Ser	Leu	Val	Ile	Ala
			20					25					30		
Ala	Ile	Ile	Val	Thr	Arg	Lys	Leu	His	His	Pro					
		35					40								

<210> 49

64N

<211> 37
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT2A GPCR TM1

<400> 49
 Gln Glu Lys Asn Trp Ser Ala Leu Leu Thr Ala Val Val Ile Ile Leu
 1 5 10 15
 Thr Ile Ala Gly Asn Ile Leu Val Ile Met Ala Val Ser Leu Glu Lys
 20 25 30
 Lys Leu Gln Asn Ala
 35

<210> 50
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT2B GPCR TM1

<400> 50
 Ile Val Glu Glu Gln Gly Asn Lys Leu His Trp Ala Ala Leu Leu Ile
 1 5 10 15
 Leu Met Val Ile Ile Pro Thr Ile Gly Gly Asn Thr Leu Val Ile Leu
 20 25 30
 Ala Val Ser Leu Glu Lys Lys Leu Gln Tyr Ala
 35 40

<210> 51
 <211> 35
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT2C GPCR TM1

<400> 51
 Gln Asn Trp Pro Ala Leu Ser Ile Val Ile Ile Ile Ile Met Thr Ile
 1 5 10 15
 Gly Gly Asn Ile Leu Val Ile Met Ala Val Ser Met Glu Lys Lys Leu
 20 25 30
 His Asn Ala
 35

<210> 52
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT5A GPCR TM1

<400> 52
 Ser Ser Pro Leu Leu Ser Val Phe Gly Val Leu Ile Leu Thr Leu Leu
 1 5 10 15

640

Gly Phe Leu Val Ala Ala Thr Phe Ala Trp Asn Leu Leu Val Leu Ala
 20 25 30

Thr Ile Leu Arg Val Arg Thr Phe His Arg Val
 35 40

<210> 53
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT5Brat GPCR TM1

 <400> 53
 Arg Glu Pro Pro Phe Ser Ala Phe Thr Val Leu Val Val Thr Leu Leu
 1 5 10 15
 Val Leu Leu Ile Ala Ala Thr Phe Leu Trp Asn Leu Leu Val Leu Val
 20 25 30
 Thr Ile Leu Arg Val Arg Ala Phe His Arg Val
 35 40

<210> 54
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT6rat GPCR TM1

 <400> 54
 Gly Pro Pro Pro Ala Pro Gly Gly Ser Gly Trp Val Ala Ala Ala Leu
 1 5 10 15
 Cys Val Val Ile Val Leu Thr Ala Ala Ala Asn Ser Leu Leu Ile Val
 20 25 30
 Leu Ile Cys Thr Gln Pro Ala Val Arg Asn Thr
 35 40

<210> 55
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT7 GPCR TM1

 <400> 55
 Gln Ile Asn Tyr Gly Arg Val Glu Lys Val Val Ile Gly Ser Ile Leu
 1 5 10 15
 Thr Leu Ile Thr Leu Leu Thr Ile Ala Gly Asn Cys Leu Val Val Ile
 20 25 30
 Ser Val Cys Phe Val Lys Lys Leu Arg Gln Pro
 35 40

<210> 56
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64P

<223> alpha1A GPCR TM1

<400> 56

Gly	Gly	Leu	Val	Val	Ser	Ala	Gln	Gly	Val	Gly	Val	Gly	Val	Phe	Leu
1				5				10						15	
Ala	Ala	Phe	Ile	Leu	Met	Ala	Val	Ala	Gly	Asn	Leu	Leu	Val	Ile	Leu
			20					25					30		
Ser	Val	Ala	Cys	Asn	Arg	His	Leu	Gln	Thr	Val					
		35					40								

<210> 57

<211> 41

<212> PRT

<213> Artificial Sequence

<220>

<223> alpha1B GPCR TM1

<400> 57

Gln	Leu	Asp	Ile	Thr	Arg	Ala	Ile	Ser	Val	Gly	Leu	Val	Leu	Gly	Ala
1				5				10						15	
Phe	Ile	Leu	Phe	Ala	Ile	Val	Gly	Asn	Ile	Leu	Val	Ile	Leu	Ser	Val
			20					25					30		
Ala	Cys	Asn	Arg	His	Leu	Arg	Thr	Pro							
		35					40								

<210> 58

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> alpha1C GPCR TM1

<400> 58

Pro	Ala	Pro	Val	Asn	Ile	Ser	Lys	Ala	Ile	Leu	Leu	Gly	Val	Ile	Leu
1				5				10						15	
Gly	Gly	Leu	Ile	Leu	Phe	Gly	Val	Leu	Cys	Asn	Ile	Leu	Val	Ile	Leu
			20					25					30		
Ser	Val	Ala	Cys	His	Arg	His	Leu	His	Ser	Val					
		35					40								

<210> 59

<211> 40

<212> PRT

<213> Artificial Sequence

<220>

<223> alpha2A GPCR TM1

<400> 59

Tyr	Ser	Leu	Gln	Val	Thr	Leu	Thr	Leu	Val	Cys	Leu	Ala	Gly	Leu	Leu
1				5				10						15	
Met	Leu	Leu	Thr	Val	Phe	Gly	Asn	Val	Leu	Val	Ile	Ile	Ala	Val	Phe
			20					25					30		
Thr	Ser	Arg	Ala	Leu	Lys	Ala	Pro								
		35					40								

<210> 60

64Q

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> alpha2B GPCR TM1

<400> 60
 Gln Asp Pro Tyr Ser Val Gln Ala Thr Ala Ala Ile Ala Ala Ala Ile
 1 5 10 15
 Thr Phe Leu Ile Leu Phe Thr Ile Phe Gly Asn Ala Leu Val Ile Leu
 20 25 30
 Ala Val Leu Thr Ser Arg Ser Leu Arg Ala Pro
 35 40

<210> 61
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> alpha2C1 and alpha2C2 GPCR TM1

<400> 61
 Arg Gly Gln Tyr Ser Ala Gly Ala Val Ala Gly Leu Ala Ala Val Val
 1 5 10 15
 Gly Phe Leu Ile Val Phe Thr Val Val Gly Asn Val Leu Val Val Ile
 20 25 30
 Ala Val Leu Thr Ser Arg Ala Leu Arg Ala Pro
 35 40

<210> 62
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> beta1 GPCR TM1

<400> 62
 Glu Pro Leu Ser Gln Gln Trp Thr Ala Gly Met Gly Leu Leu Met Ala
 1 5 10 15
 Leu Ile Val Leu Leu Ile Val Ala Gly Asn Val Leu Val Ile Val Ala
 20 25 30
 Ile Ala Lys Thr Pro Arg Leu Gln Thr Leu
 35 40

<210> 63
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> beta2 GPCR TM1

<400> 63
 Gln Gln Arg Asp Glu Val Trp Val Val Gly Met Gly Ile Val Met Ser
 1 5 10 15

64R

Leu Ile Val Leu Ala Ile Val Phe Gly Asn Val Leu Val Ile Thr Ala
 20 25 30
 Ile Ala Lys Phe Glu Arg Leu Gln Thr Val
 35 40

<210> 64
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> beta3 GPCR TM1

<400> 64
 Gly Leu Pro Gly Val Pro Trp Glu Ala Ala Leu Ala Gly Ala Leu Leu
 1 5 10 15
 Ala Leu Ala Val Leu Ala Thr Val Gly Gly Asn Leu Leu Val Ile Val
 20 25 30
 Ala Ile Ala Trp Thr Pro Arg Leu Gln Thr Met
 35 40

<210> 65
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> beta4turkey GPCR TM1

<400> 65
 Ser Trp Ala Ala Val Leu Ser Arg Gln Trp Ala Val Gly Ala Ala Leu
 1 5 10 15
 Ser Ile Thr Ile Leu Val Ile Val Ala Gly Asn Leu Leu Val Ile Val
 20 25 30
 Ala Ile Ala Lys Thr Pro Arg Leu Gln Thr Met
 35 40

<210> 66
 <211> 44
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> D1A GPCR TM1

<400> 66
 Val Val Glu Arg Asp Phe Ser Val Arg Ile Leu Thr Ala Cys Phe Leu
 1 5 10 15
 Ser Leu Leu Ile Leu Ser Thr Leu Leu Gly Asn Thr Leu Val Cys Ala
 20 25 30
 Ala Val Ile Arg Phe Arg His Leu Arg Ser Lys Val
 35 40

<210> 67
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64S

<223> D2 GPCR TM1

<400> 67

Asp	Gly	Lys	Ala	Asp	Arg	Pro	His	Tyr	Asn	Tyr	Tyr	Ala	Thr	Leu	Leu
1				5					10					15	
Thr	Leu	Leu	Ile	Ala	Val	Ile	Val	Phe	Gly	Asn	Val	Leu	Val	Cys	Met
			20					25					30		
Ala	Val	Ser	Arg	Glu	Lys	Ala	Leu	Gln	Thr	Thr					
		35					40								

<210> 68

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> D3 GPCR TM1

<400> 68

Thr	Gly	Ala	Ser	Gln	Ala	Arg	Pro	His	Ala	Tyr	Tyr	Ala	Leu	Ser	Tyr
1				5					10					15	
Cys	Ala	Leu	Ile	Leu	Ala	Ile	Val	Phe	Gly	Asn	Gly	Leu	Val	Cys	Met
			20					25					30		
Ala	Val	Leu	Lys	Glu	Arg	Ala	Leu	Gln	Thr	Thr					
		35					40								

<210> 69

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> D4 GPCR TM1

<400> 69

Ala	Ser	Ala	Gly	Leu	Ala	Gly	Gln	Gly	Ala	Ala	Ala	Leu	Val	Gly	Gly
1				5					10					15	
Val	Leu	Leu	Ile	Gly	Ala	Val	Leu	Ala	Gly	Asn	Ser	Leu	Val	Cys	Val
			20					25					30		
Ser	Val	Ala	Thr	Glu	Arg	Ala	Leu	Gln	Thr	Pro					
		35					40								

<210> 70

<211> 44

<212> PRT

<213> Artificial Sequence

<220>

<223> D5 GPCR TM1

<400> 70

Gly	Ala	Pro	Pro	Leu	Gly	Pro	Ser	Gln	Val	Val	Thr	Ala	Cys	Leu	Leu
1				5					10					15	
Thr	Leu	Leu	Ile	Ile	Trp	Thr	Leu	Leu	Gly	Asn	Val	Leu	Val	Cys	Ala
			20					25					30		
Ala	Ile	Val	Arg	Ser	Arg	His	Leu	Arg	Ala	Asn	Met				
		35					40								

<210> 71

64T

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> A1 GPCR TM1

<400> 71
 Met Pro Pro Ser Ile Ser Ala Phe Gln Ala Ala Tyr Ile Gly Ile Glu
 1 5 10 15
 Val Leu Ile Ala Leu Val Ser Val Pro Gly Asn Val Leu Val Ile Trp
 20 25 30
 Ala Val Lys Val Asn Gln Ala Leu Arg Asp Ala
 35 40

<210> 72
 <211> 40
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> A2a GPCR TM1

<400> 72
 Met Pro Ile Met Gly Ser Ser Val Tyr Ile Thr Val Glu Leu Ala Ile
 1 5 10 15
 Ala Val Leu Ala Ile Leu Gly Asn Val Leu Val Cys Trp Ala Val Trp
 20 25 30
 Leu Asn Ser Asn Leu Gln Asn Val
 35 40

<210> 73
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> A2b GPCR TM1

<400> 73
 Met Leu Leu Glu Thr Gln Asp Ala Leu Tyr Val Ala Leu Glu Leu Val
 1 5 10 15
 Ile Ala Ala Leu Ser Val Ala Gly Asn Val Leu Val Cys Ala Ala Val
 20 25 30
 Gly Thr Ala Asn Thr Leu Gln Thr Pro
 35 40

<210> 74
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> A3 GPCR TM1

<400> 74
 Asn Ser Thr Thr Leu Ser Leu Ala Asn Val Thr Tyr Ile Thr Met Glu
 1 5 10 15

64U

Ile Phe Ile Gly Leu Cys Ala Ile Val Gly Asn Val Leu Val Ile Cys
 20 25 30
 Val Val Lys Leu Asn Pro Ser Leu Gln Thr Thr
 35 40

<210> 75
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OCdrome GPCR TM1

<400> 75
 Leu Ala Val Pro Glu Trp Glu Ala Leu Leu Thr Ala Leu Val Leu Ser
 1 5 10 15
 Val Ile Ile Val Leu Thr Ile Ile Gly Asn Ile Leu Val Ile Leu Ser
 20 25 30
 Val Phe Thr Tyr Lys Pro Leu Arg Ile Val
 35 40

<210> 76
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> ACTH GPCR TM1

<400> 76
 Arg Asn Asn Ser Asp Cys Pro Arg Val Val Leu Pro Glu Glu Ile Phe
 1 5 10 15
 Phe Thr Ile Ser Ile Val Gly Val Leu Glu Asn Leu Ile Val Leu Leu
 20 25 30
 Ala Val Phe Lys Asn Lys Asn Leu Gln Ala Pro
 35 40

<210> 77
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MSH GPCR TM1

<400> 77
 Gln Thr Gly Ala Arg Cys Leu Glu Val Ser Ile Ser Asp Gly Leu Phe
 1 5 10 15
 Leu Ser Leu Gly Leu Val Ser Leu Val Glu Asn Ala Leu Val Val Ala
 20 25 30
 Thr Ile Ala Lys Asn Arg Asn Leu His Ser Pro
 35 40

<210> 78
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64V

<223> MC3 GPCR TM1

<400> 78

Ser	Ser	Ser	Ala	Phe	Cys	Glu	Gln	Val	Phe	Ile	Lys	Pro	Glu	Ile	Phe
1				5					10					15	
Leu	Ser	Leu	Gly	Ile	Val	Ser	Leu	Leu	Glu	Asn	Ile	Leu	Val	Ile	Leu
			20					25					30		
Ala	Val	Val	Arg	Asn	Gly	Asn	Leu	His	Ser	Pro					
		35					40								

<210> 79

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> MC4 GPCR TM1

<400> 79

Ser	Asp	Gly	Gly	Cys	Tyr	Glu	Gln	Leu	Phe	Val	Ser	Pro	Glu	Val	Phe
1				5					10					15	
Val	Thr	Leu	Gly	Val	Ile	Ser	Leu	Leu	Glu	Asn	Ile	Leu	Val	Ile	Val
			20					25					30		
Ala	Ile	Ala	Lys	Asn	Lys	Asn	Leu	His	Ser	Pro					
		35					40								

<210> 80

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> MC5 GPCR TM1

<400> 80

Asn	Lys	Ser	Ser	Pro	Cys	Glu	Asp	Met	Gly	Ile	Ala	Val	Glu	Val	Phe
1				5					10					15	
Leu	Thr	Leu	Gly	Val	Ile	Ser	Leu	Leu	Glu	Asn	Ile	Leu	Val	Ile	Gly
			20					25					30		
Ala	Ile	Val	Lys	Asn	Lys	Asn	Leu	His	Ser	Pro					
		35					40								

<210> 81

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> melatonin GPCR TM1

<400> 81

Asp	Gly	Ala	Arg	Pro	Ser	Trp	Leu	Ala	Ser	Ala	Leu	Ala	Cys	Val	Leu
1				5					10					15	
Ile	Phe	Thr	Ile	Val	Val	Asp	Ile	Leu	Gly	Asn	Leu	Leu	Val	Ile	Leu
			20					25					30		
Ser	Val	Tyr	Arg	Asn	Lys	Lys	Leu	Arg	Asn	Ala					
		35					40								

<210> 82

64W

<211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> oxytocin GPCR TM1

<400> 82
 Arg Arg Asn Glu Ala Leu Ala Arg Val Glu Val Ala Val Leu Cys Leu
 1 5 10 15
 Ile Leu Leu Leu Ala Leu Ser Gly Asn Ala Cys Val Leu Leu Ala Leu
 20 25 30
 Arg Thr Thr Arg Gln Lys His Ser Arg
 35 40

<210> 83
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> conopressinLs GPCR TM1

<400> 83
 Phe His Gly Val Asp Glu Asp Leu Leu Lys Ile Glu Ile Ala Val Gln
 1 5 10 15
 Ala Thr Ile Leu Tyr Met Thr Leu Phe Gly Asn Gly Ile Val Leu Leu
 20 25 30
 Val Leu Arg Leu Arg Arg Gln Lys Leu Thr Arg
 35 40

<210> 84
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> V1A GPCR TM1

<400> 84
 Arg Asp Val Arg Asn Glu Glu Leu Ala Lys Leu Glu Ile Ala Val Leu
 1 5 10 15
 Ala Val Thr Phe Ala Val Ala Val Leu Gly Asn Ser Ser Val Leu Leu
 20 25 30
 Ala Leu His Arg Thr Pro Arg Lys Thr Ser Arg
 35 40

<210> 85
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> V1B GPCR TM1

<400> 85
 Trp Leu Gly Arg Asp Glu Glu Leu Ala Lys Val Glu Ile Gly Val Leu
 1 5 10 15

64X

Ala Thr Val Leu Val Leu Ala Thr Gly Gly Asn Leu Ala Val Leu Leu
 20 25 30
 Thr Leu Gly Gln Leu Gly Arg Lys Arg Ser Arg
 35 40

<210> 86
 <211> 45
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> V2 GPCR TM1

<400> 86
 Leu Asp Thr Arg Asp Pro Leu Leu Ala Arg Ala Glu Leu Ala Leu Leu
 1 5 10 15
 Ser Ile Val Phe Val Ala Val Ala Leu Ser Asn Gly Leu Val Leu Ala
 20 25 30
 Ala Leu Ala Arg Arg Gly Arg Arg Gly His Trp Ala Pro
 35 40 45

<210> 87
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCK_A GPCR TM1

<400> 87
 Pro Arg Pro Ser Lys Glu Trp Gln Pro Ala Val Gln Ile Leu Leu Tyr
 1 5 10 15
 Ser Leu Ile Phe Leu Leu Ser Val Leu Gly Asn Thr Leu Val Ile Thr
 20 25 30
 Val Leu Ile Arg Asn Lys Arg Met Arg Thr Val
 35 40

<210> 88
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CCK_B GPCR TM1

<400> 88
 Gly Ala Gly Thr Arg Glu Leu Glu Leu Ala Ile Arg Ile Thr Leu Tyr
 1 5 10 15
 Ala Val Ile Phe Leu Met Ser Val Gly Gly Asn Met Leu Ile Ile Val
 20 25 30
 Val Leu Gly Leu Ser Arg Arg Leu Arg Thr Val
 35 40

<210> 89
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64Y

<223> NPY1 GPCR TM1

<400> 89

Asp	Cys	His	Leu	Pro	Leu	Ala	Met	Ile	Phe	Thr	Leu	Ala	Leu	Ala	Tyr
1				5					10					15	
Gly	Ala	Val	Ile	Ile	Leu	Gly	Val	Ser	Gly	Asn	Leu	Ala	Leu	Ile	Ile
			20					25					30		
Ile	Ile	Leu	Lys	Gln	Lys	Glu	Met	Arg	Asn	Val					
		35					40								

<210> 90

<211> 46

<212> PRT

<213> Artificial Sequence

<220>

<223> NTR GPCR TM1

<400> 90

Asp	Val	Asn	Thr	Asp	Ile	Tyr	Ser	Lys	Val	Leu	Val	Thr	Ala	Val	Tyr
1				5					10					15	
Leu	Ala	Leu	Phe	Val	Val	Gly	Thr	Val	Gly	Asn	Thr	Val	Thr	Ala	Phe
			20					25					30		
Thr	Leu	Ala	Arg	Lys	Lys	Ser	Leu	Gln	Ser	Leu	Gln	Ser	Thr		
		35					40					45			

<210> 91

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> NK1 GPCR TM1

<400> 91

Gln	Phe	Val	Gln	Pro	Ala	Trp	Gln	Ile	Val	Leu	Trp	Ala	Ala	Ala	Tyr
1				5					10					15	
Thr	Val	Ile	Val	Val	Thr	Ser	Val	Val	Gly	Asn	Val	Val	Val	Met	Trp
			20					25					30		
Ile	Ile	Leu	Ala	His	Lys	Arg	Met	Arg	Thr	Val					
		35					40								

<210> 92

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> NK2 GPCR TM1

<400> 92

Ala	Phe	Ser	Met	Pro	Ser	Trp	Gln	Leu	Ala	Leu	Trp	Ala	Pro	Ala	Tyr
1				5					10					15	
Leu	Ala	Leu	Val	Leu	Val	Ala	Val	Thr	Gly	Asn	Ala	Ile	Val	Ile	Trp
			20					25					30		
Ile	Ile	Leu	Ala	His	Arg	Arg	Met	Arg	Thr	Val					
		35					40								

<210> 93

64Z

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> NK3 GPCR TM1

<400> 93
 Gln Phe Val Gln Pro Ser Trp Arg Ile Ala Leu Trp Ser Leu Ala Tyr
 1 5 10 15
 Gly Val Val Val Ala Val Ala Val Leu Gly Asn Leu Ile Val Ile Trp
 20 25 30
 Ile Ile Leu Ala His Lys Arg Met Arg Thr Val
 35 40

<210> 94
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> blueops GPCR TM1

<400> 94
 Tyr His Ile Ala Pro Val Trp Ala Phe Tyr Leu Gln Ala Ala Phe Met
 1 5 10 15
 Gly Thr Val Phe Leu Ile Gly Phe Pro Leu Asn Ala Met Val Leu Val
 20 25 30
 Ala Thr Leu Arg Tyr Lys Lys Leu Arg Gln Pro
 35 40

<210> 95
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> greenops GPCR TM1

<400> 95
 Tyr His Ile Ala Pro Arg Trp Val Tyr His Leu Thr Ser Val Trp Met
 1 5 10 15
 Ile Phe Val Val Ile Ala Ser Val Phe Thr Asn Gly Leu Val Leu Ala
 20 25 30
 Ala Thr Met Lys Phe Lys Lys Leu Arg His Pro
 35 40

<210> 96
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> redops GPCR TM1

<400> 96
 Tyr His Ile Ala Pro Arg Trp Val Tyr His Leu Thr Ser Val Trp Met
 1 5 10 15

64AA

Ile Phe Val Val Thr Ala Ser Val Phe Thr Asn Gly Leu Val Leu Ala
 20 25 30
 Ala Thr Met Lys Phe Lys Lys Leu Arg His Pro
 35 40

<210> 97
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> rhodopsin GPCR TM1

<400> 97
 Tyr Tyr Leu Ala Glu Pro Trp Gln Phe Ser Met Leu Ala Ala Tyr Met
 1 5 10 15
 Phe Leu Leu Ile Val Leu Gly Phe Pro Ile Asn Phe Leu Thr Leu Tyr
 20 25 30
 Val Thr Val Gln His Lys Lys Leu Arg Thr Pro
 35 40

<210> 98
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> violetopsGg GPCR TM1

<400> 98
 Tyr His Ile Ala Pro Pro Trp Ala Phe Tyr Leu Gln Thr Ala Phe Met
 1 5 10 15
 Gly Ile Val Phe Ala Val Gly Thr Pro Leu Asn Ala Val Val Leu Trp
 20 25 30
 Val Thr Val Arg Tyr Lys Arg Leu Arg Gln Pro
 35 40

<210> 99
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> opsin_crab GPCR TM1

<400> 99
 Phe Pro Pro Met Asn Pro Leu Trp Tyr Ser Ile Leu Gly Val Ala Met
 1 5 10 15
 Ile Ile Leu Gly Ile Ile Cys Val Leu Gly Asn Gly Met Val Ile Tyr
 20 25 30
 Leu Met Met Thr Thr Lys Ser Leu Arg Thr Pro
 35 40

<210> 100
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64BB

<223> ET_Aprec GPCR TM1

<400> 100

Gln	Thr	Lys	Ile	Thr	Ser	Ala	Phe	Lys	Tyr	Ile	Asn	Thr	Val	Ile	Ser
1				5					10					15	
Cys	Thr	Ile	Phe	Ile	Val	Gly	Met	Val	Gly	Asn	Ala	Thr	Leu	Leu	Arg
			20					25					30		
Ile	Ile	Tyr	Gln	Asn	Lys	Cys	Met	Arg	Asn	Gly					
		35					40								

<210> 101

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> ET_Bprec GPCR TM1

<400> 101

Pro	Ile	Glu	Ile	Lys	Glu	Thr	Phe	Lys	Tyr	Ile	Asn	Thr	Val	Val	Ser
1				5					10					15	
Cys	Leu	Val	Phe	Val	Leu	Gly	Ile	Ile	Gly	Asn	Ser	Thr	Leu	Leu	Arg
			20					25					30		
Ile	Ile	Tyr	Lys	Asn	Lys	Cys	Met	Arg	Asn	Gly					
		35					40								

<210> 102

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> ET_Cfrog GPCR TM1

<400> 102

Arg	Ala	Lys	Ile	Arg	His	Ala	Phe	Lys	Tyr	Val	Thr	Thr	Ile	Leu	Ser
1				5					10					15	
Cys	Val	Ile	Phe	Leu	Val	Gly	Ile	Val	Gly	Asn	Ser	Thr	Leu	Leu	Arg
			20					25					30		
Ile	Ile	Tyr	Lys	Asn	Lys	Cys	Met	Arg	Asn	Gly					
		35					40								

<210> 103

<211> 45

<212> PRT

<213> Artificial Sequence

<220>

<223> galanin GPCR TM1

<400> 103

Pro	Leu	Phe	Gly	Ile	Gly	Val	Glu	Asn	Phe	Val	Thr	Leu	Val	Val	Phe
1				5					10					15	
Gly	Leu	Ile	Phe	Ala	Leu	Gly	Val	Leu	Gly	Asn	Ser	Leu	Val	Ile	Thr
			20					25					30		
Val	Leu	Ala	Arg	Ser	Lys	Pro	Gly	Lys	Pro	Arg	Ser	Thr			
		35					40					45			

<210> 104

64CC

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> NMB GPCR TM1

<400> 104
 Gly Thr Thr Thr Glu Leu Val Ile Arg Cys Val Ile Pro Ser Leu Tyr
 1 5 10 15
 Leu Leu Ile Ile Thr Val Gly Leu Leu Gly Asn Ile Met Leu Val Lys
 20 25 30
 Ile Phe Ile Thr Asn Ser Ala Met Arg Ser Val
 35 40

<210> 105
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> GRP GPCR TM1

<400> 105
 Asp Asp Trp Ser His Pro Gly Ile Leu Tyr Val Ile Pro Ala Val Tyr
 1 5 10 15
 Gly Val Ile Ile Leu Ile Gly Leu Ile Gly Asn Ile Thr Leu Ile Lys
 20 25 30
 Ile Phe Cys Thr Val Lys Ser Met Arg Asn Val
 35 40

<210> 106
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> BRS3 GPCR TM1

<400> 106
 Asp Asn Ser Pro Gly Ile Glu Ala Leu Cys Ala Ile Tyr Ile Thr Tyr
 1 5 10 15
 Ala Val Ile Ile Ser Val Gly Ile Leu Gly Asn Ala Ile Leu Ile Lys
 20 25 30
 Val Phe Phe Lys Thr Lys Ser Met Gln Thr Val
 35 40

<210> 107
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> deltaOP GPCR TM1

<400> 107
 Gly Ser Ala Ser Ser Leu Ala Leu Ala Ile Ala Ile Thr Ala Leu Tyr
 1 5 10 15

64DD

Ser Ala Val Cys Ala Val Gly Leu Leu Gly Asn Val Leu Val Met Phe
 20 25 30
 Gly Ile Val Arg Tyr Thr Lys Met Lys Thr Ala
 35 40

<210> 108
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> kappaOP GPCR TM1

<400> 108
 Pro Ala His Ile Ser Pro Ala Ile Pro Val Ile Ile Thr Ala Val Tyr
 1 5 10 15
 Ser Val Val Phe Val Val Gly Leu Val Gly Asn Ser Leu Val Met Phe
 20 25 30
 Val Ile Ile Arg Tyr Thr Lys Met Lys Thr Ala
 35 40

<210> 109
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> muOP GPCR TM1

<400> 109
 Gly Ser Pro Ser Met Ile Thr Ala Ile Thr Ile Met Ala Leu Tyr Ser
 1 5 10 15
 Ile Val Cys Val Val Gly Leu Phe Gly Asn Phe Leu Val Met Tyr Val
 20 25 30
 Ile Val Arg Tyr Thr Lys Met Lys Thr Ala
 35 40

<210> 110
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OPRX GPCR TM1

<400> 110
 Gly Ala Phe Leu Pro Leu Gly Leu Lys Val Thr Ile Val Gly Leu Tyr
 1 5 10 15
 Leu Ala Val Cys Val Gly Gly Leu Leu Gly Asn Cys Leu Val Met Tyr
 20 25 30
 Val Ile Leu Arg His Thr Lys Met Lys Thr Ala
 35 40

<210> 111
 <211> 44
 <212> PRT
 <213> Artificial Sequence

<220>

64EE

<223> CB1 GPCR TM1

<400> 111

Phe	Met	Val	Leu	Asn	Pro	Ser	Gln	Gln	Leu	Ala	Ile	Ala	Val	Leu	Ser
1				5					10					15	
Leu	Thr	Leu	Gly	Thr	Phe	Thr	Val	Leu	Glu	Asn	Leu	Leu	Val	Leu	Cys
			20					25					30		
Val	Ile	Leu	His	Ser	Arg	Ser	Leu	Arg	Cys	Arg	Pro				
		35					40								

<210> 112

<211> 44

<212> PRT

<213> Artificial Sequence

<220>

<223> CB2 GPCR TM1

<400> 112

Tyr	Met	Ile	Leu	Ser	Gly	Pro	Gln	Lys	Thr	Ala	Val	Ala	Val	Leu	Cys
1				5				10						15	
Thr	Leu	Leu	Gly	Leu	Leu	Ser	Ala	Leu	Glu	Asn	Val	Ala	Val	Leu	Tyr
			20					25					30		
Leu	Ile	Leu	Ser	Ser	His	Gln	Leu	Arg	Arg	Lys	Pro				
		35					40								

<210> 113

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> SSTR1 GPCR TM1

<400> 113

Thr	Leu	Ser	Glu	Gly	Gln	Gly	Ser	Ala	Ile	Leu	Ile	Ser	Phe	Ile	Tyr
1				5				10						15	
Ser	Val	Val	Cys	Leu	Val	Gly	Leu	Cys	Gly	Asn	Ser	Met	Val	Ile	Tyr
			20					25					30		
Val	Ile	Leu	Arg	Tyr	Ala	Lys	Met	Lys	Thr	Ala					
		35					40								

<210> 114

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> SSTR2 GPCR TM1

<400> 114

Glu	Pro	Tyr	Tyr	Asp	Leu	Thr	Ser	Asn	Ala	Val	Leu	Thr	Phe	Ile	Tyr
1				5				10						15	
Phe	Val	Val	Cys	Ile	Ile	Gly	Leu	Cys	Gly	Asn	Thr	Leu	Val	Ile	Tyr
			20					25					30		
Val	Ile	Leu	Arg	Tyr	Ala	Lys	Met	Lys	Thr	Ile					
		35					40								

<210> 115

64FF

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> SSTR3 GPCR TM1

<400> 115
 Ser Pro Ala Gly Leu Ala Val Ser Gly Val Leu Ile Pro Leu Val Tyr
 1 5 10 15
 Leu Val Val Cys Val Val Gly Leu Leu Gly Asn Ser Leu Val Ile Tyr
 20 25 30
 Val Val Leu Arg His Thr Ala Ser Pro Ser Val
 35 40

<210> 116
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> SSTR4 GPCR TM1

<400> 116
 Gly Asp Ala Arg Ala Ala Gly Met Val Ala Ile Gln Cys Ile Tyr Ala
 1 5 10 15
 Leu Val Cys Leu Val Gly Leu Val Gly Asn Ala Leu Val Ile Phe Val
 20 25 30
 Ile Leu Arg Tyr Ala Lys Met Lys Thr Ala
 35 40

<210> 117
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> SSTR5 GPCR TM1

<400> 117
 Pro Ala Pro Ser Ala Gly Ala Arg Ala Val Leu Val Pro Val Leu Tyr
 1 5 10 15
 Leu Leu Val Cys Ala Ala Gly Leu Gly Gly Asn Thr Leu Val Ile Tyr
 20 25 30
 Val Val Leu Arg Phe Ala Lys Met Lys Thr Val
 35 40

<210> 118
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> IL8A GPCR TM1

<400> 118
 Met Leu Glu Thr Glu Thr Leu Asn Lys Tyr Val Val Ile Ile Ala Tyr
 1 5 10 15

64GG

Ala Leu Val Phe Leu Leu Ser Leu Leu Gly Asn Ser Leu Val Met Leu
 20 25 30
 Val Ile Leu Tyr Ser Arg Val Gly Arg Ser Val
 35 40

<210> 119
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> IL8B GPCR TM1

<400> 119
 Glu Pro Glu Ser Leu Glu Ile Asn Lys Tyr Phe Val Val Ile Ile Tyr
 1 5 10 15
 Ala Leu Val Phe Leu Leu Ser Leu Leu Gly Asn Ser Leu Val Met Leu
 20 25 30
 Val Ile Leu Tyr Ser Arg Val Gly Arg Ser Val
 35 40

<210> 120
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> AT1 AND AT1brat GPCR TM1

<400> 120
 Lys Ala Gly Arg His Asn Tyr Ile Phe Val Met Ile Pro Thr Leu Tyr
 1 5 10 15
 Ser Ile Ile Phe Val Val Gly Ile Phe Gly Asn Ser Leu Val Val Ile
 20 25 30
 Val Ile Tyr Phe Tyr Met Lys Leu Lys Thr Val
 35 40

<210> 121
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> AT2 GPCR TM1

<400> 121
 Gln Lys Pro Ser Asp Lys His Leu Asp Ala Ile Pro Ile Leu Tyr Tyr
 1 5 10 15
 Ile Ile Phe Val Ile Gly Phe Leu Val Asn Ile Val Val Val Thr Leu
 20 25 30
 Phe Cys Cys Gln Lys Gly Pro Lys Lys Val
 35 40

<210> 122
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64HH

<223> BK1 GPCR TM1

<400> 122

Ala	Pro	Glu	Ala	Trp	Asp	Leu	Leu	His	Arg	Val	Leu	Pro	Thr	Phe	Ile
1				5					10					15	
Ile	Ser	Ile	Cys	Phe	Phe	Gly	Leu	Leu	Gly	Asn	Leu	Phe	Val	Leu	Leu
			20					25					30		
Val	Phe	Leu	Leu	Pro	Arg	Arg	Gln	Leu	Asn	Val					
		35					40								

<210> 123

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> BK2 GPCR TM1

<400> 123

Gln	Val	Glu	Trp	Leu	Gly	Trp	Leu	Asn	Thr	Ile	Gln	Pro	Pro	Phe	Leu
1				5					10					15	
Trp	Val	Leu	Phe	Val	Leu	Ala	Thr	Leu	Glu	Asn	Ile	Phe	Val	Leu	Ser
			20					25					30		
Val	Phe	Cys	Leu	His	Lys	Ser	Ser	Cys	Thr	Val					
		35					40								

<210> 124

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> P2Y7 GPCR TM1

<400> 124

Pro	Ser	Leu	Gly	Val	Glu	Phe	Ile	Ser	Leu	Leu	Ala	Ile	Ile	Leu	Leu
1				5					10					15	
Ser	Val	Ala	Leu	Ala	Val	Gly	Leu	Pro	Gly	Asn	Ser	Phe	Val	Val	Trp
			20					25					30		
Ser	Ile	Leu	Lys	Arg	Met	Gln	Lys	Arg	Ser	Val					
		35					40								

<210> 125

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> P2Y6 GPCR TM1

<400> 125

Cys	Val	Tyr	Arg	Glu	Asp	Phe	Lys	Arg	Leu	Leu	Leu	Pro	Pro	Val	Tyr
1				5					10					15	
Ser	Val	Val	Leu	Val	Val	Gly	Leu	Pro	Leu	Asn	Val	Cys	Val	Ile	Ala
			20					25					30		
Gln	Ile	Cys	Ala	Ser	Arg	Arg	Thr	Leu	Thr	Arg					
		35					40								

<210> 126

64II

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P2Y5 GPCR TM1

<400> 126
 Cys Ser Thr Glu Asp Ser Phe Lys Tyr Thr Leu Tyr Gly Cys Val Phe
 1 5 10 15
 Ser Met Val Phe Val Leu Gly Leu Ile Ala Asn Cys Val Ala Ile Tyr
 20 25 30
 Ile Phe Thr Phe Thr Leu Lys Val Arg Asn Glu
 35 40

<210> 127
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P2Y4 GPCR TM1

<400> 127
 Cys Trp Phe Asp Glu Asp Phe Lys Phe Ile Leu Leu Pro Val Ser Tyr
 1 5 10 15
 Ala Val Val Phe Val Leu Gly Leu Gly Leu Asn Ala Pro Thr Leu Trp
 20 25 30
 Leu Phe Ile Phe Arg Leu Arg Pro Trp Asp Ala
 35 40

<210> 128
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P2Y3chick GPCR TM1

<400> 128
 Cys Thr Phe His Glu Glu Phe Lys Gln Val Leu Leu Pro Leu Val Tyr
 1 5 10 15
 Ser Val Val Phe Leu Leu Gly Leu Pro Leu Asn Ala Val Val Ile Gly
 20 25 30
 Gln Ile Trp Leu Ala Arg Lys Ala Leu Thr Arg
 35 40

<210> 129
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P2Y2 GPCR TM1

<400> 129
 Cys Arg Phe Asn Glu Asp Phe Lys Tyr Val Leu Leu Pro Val Ser Tyr
 1 5 10 15

64JJ

Gly Val Val Cys Val Leu Gly Leu Cys Leu Asn Ala Val Gly Leu Tyr
 20 25 30
 Ile Phe Leu Cys Arg Leu Lys Thr Trp Asn Ala
 35 40

<210> 130
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P2Y1 GPCR TM1

<400> 130
 Ala Leu Thr Lys Thr Gly Phe Gln Phe Tyr Tyr Leu Pro Ala Val Tyr
 1 5 10 15
 Ile Leu Val Phe Ile Ile Gly Phe Leu Gly Asn Ser Val Ala Ile Trp
 20 25 30
 Met Phe Val Phe His Met Lys Pro Trp Ser Gly
 35 40

<210> 131
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> THRprec GPCR TM1

<400> 131
 Gly Tyr Leu Thr Ser Ser Trp Leu Thr Leu Phe Val Pro Ser Val Tyr
 1 5 10 15
 Thr Gly Val Phe Val Val Ser Leu Pro Leu Asn Ile Met Ala Ile Val
 20 25 30
 Val Phe Ile Leu Lys Met Lys Val Lys Lys Pro
 35 40

<210> 132
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> C5a GPCR TM1

<400> 132
 Thr Ser Asn Thr Leu Arg Val Pro Asp Ile Leu Ala Leu Val Ile Phe
 1 5 10 15
 Ala Val Val Phe Leu Val Gly Val Leu Gly Asn Ala Leu Val Val Trp
 20 25 30
 Val Thr Ala Phe Glu Ala Lys Arg Thr Ile
 35 40

<210> 133
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64KK

<223> GP01mouse GPCR TM1

<400> 133

Ala	Glu	Ser	Glu	Pro	Glu	Leu	Val	Val	Asn	Pro	Trp	Asp	Ile	Val	Leu
1				5					10					15	
Cys	Ser	Ser	Gly	Thr	Leu	Ile	Cys	Cys	Glu	Asn	Ala	Val	Val	Val	Leu
			20					25					30		
Ile	Ile	Phe	His	Ser	Pro	Ser	Leu	Arg	Ala	Pro					
		35					40								

<210> 134

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> R334rat GPCR TM1

<400> 134

Val	Glu	Ser	Glu	Pro	Glu	Leu	Val	Val	Asn	Pro	Trp	Asp	Ile	Val	Leu
1				5					10					15	
Cys	Ser	Ser	Gly	Thr	Leu	Ile	Cys	Cys	Glu	Asn	Ala	Val	Val	Val	Leu
			20					25					30		
Ile	Ile	Phe	His	Ser	Pro	Ser	Leu	Arg	Ala	Pro					
		35					40								

<210> 135

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> GP21mouse GPCR TM1

<400> 135

Gly	Pro	Ala	Thr	Leu	Leu	Pro	Ser	Pro	Arg	Ala	Trp	Asp	Val	Val	Leu
1				5					10					15	
Cys	Ile	Ser	Gly	Thr	Leu	Val	Ser	Cys	Glu	Asn	Ala	Leu	Val	Val	Ala
			20					25					30		
Ile	Ile	Val	Gly	Thr	Pro	Ala	Phe	Arg	Ala	Pro					
		35					40								

<210> 136

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> GCRCmouse GPCR TM1

<400> 136

Ala	Glu	Ser	Gln	Asn	Pro	Thr	Val	Lys	Ala	Leu	Leu	Ile	Val	Ala	Tyr
1				5					10					15	
Ser	Phe	Thr	Ile	Val	Phe	Ser	Leu	Phe	Gly	Asn	Val	Leu	Val	Cys	His
			20					25					30		
Val	Ile	Phe	Lys	Asn	Gln	Arg	Met	His	Ser	Ala					
		35					40								

<210> 137

64LL

<211> 40
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> TXKR GPCR TM1

<400> 137
 Gln Pro Pro Trp Ala Val Ala Leu Trp Ser Leu Ala Tyr Gly Ala Val
 1 5 10 15
 Val Ala Val Ala Val Leu Gly Asn Leu Val Val Ile Trp Ile Val Leu
 20 25 30
 Ala His Lys Arg Met Arg Thr Val
 35 40

<210> 138
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> G10Drat GPCR TM1

<400> 138
 Met Glu Leu Asn Glu Asn Thr Lys Gln Val Val Leu Phe Val Phe Tyr
 1 5 10 15
 Leu Ala Ile Phe Val Val Gly Leu Val Glu Asn Val Leu Val Ile Cys
 20 25 30
 Val Asn Cys Arg Arg Ser Gly Arg Val Gly Met
 35 40

<210> 139
 <211> 44
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> RDC1 GPCR TM1

<400> 139
 Asn Met Pro Asn Lys Ser Val Leu Leu Tyr Thr Leu Ser Phe Ile Tyr
 1 5 10 15
 Ile Phe Ile Phe Val Ile Gly Met Ile Ala Asn Ser Val Val Val Trp
 20 25 30
 Val Asn Ile Gln Ala Lys Thr Thr Gly Tyr Asp Thr
 35 40

<210> 140
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> BLR1 GPCR TM1

<400> 140
 Met Ala Ser Phe Lys Ala Val Phe Val Pro Val Ala Tyr Ser Leu Ile
 1 5 10 15

64MM

Phe Leu Leu Gly Val Ile Gly Asn Val Leu Val Leu Val Ile Leu Glu
 20 25 30
 Arg His Arg Gln Thr Arg Ser Ser Thr Glu
 35 40

<210> 141
 <211> 48
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CL5 AND LCR1 GPCR TM1

<400> 141
 Arg Glu Glu Asn Ala Asn Phe Asn Lys Ile Phe Leu Pro Thr Ile Tyr
 1 5 10 15
 Ser Ile Ile Phe Leu Thr Gly Ile Val Gly Asn Gly Leu Val Ile Leu
 20 25 30
 Val Met Gly Tyr Gln Lys Lys Leu Arg Ser Met Thr Asp Lys Tyr Arg
 35 40 45

<210> 142
 <211> 47
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> EBI1 GPCR TM1

<400> 142
 Lys Lys Asp Val Arg Asn Phe Lys Ala Trp Phe Leu Pro Ile Met Tyr
 1 5 10 15
 Ser Ile Ile Cys Phe Val Gly Leu Leu Gly Asn Gly Leu Val Val Leu
 20 25 30
 Thr Tyr Ile Tyr Phe Lys Arg Leu Lys Thr Met Thr Asp Thr Tyr
 35 40 45

<210> 143
 <211> 47
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> RBS1rat GPCR TM1

<400> 143
 Leu Gly Asp Ile Val Ala Phe Gly Thr Ile Phe Leu Ser Ile Phe Tyr
 1 5 10 15
 Ser Leu Val Phe Thr Phe Gly Leu Val Gly Asn Leu Leu Val Val Leu
 20 25 30
 Ala Leu Thr Asn Ser Arg Lys Ser Lys Ser Ile Thr Asp Ile Tyr
 35 40 45

<210> 144
 <211> 47
 <212> PRT
 <213> Artificial Sequence

<220>

64NN

<223> EBI2 GPCR TM1

<400> 144

Leu	Tyr	Ala	His	His	Ser	Thr	Ala	Arg	Ile	Val	Met	Pro	Leu	His	Tyr
1				5					10					15	
Ser	Leu	Val	Phe	Ile	Ile	Gly	Leu	Val	Gly	Asn	Leu	Leu	Ala	Leu	Val
			20					25					30		
Val	Ile	Val	Gln	Asn	Arg	Lys	Lys	Ile	Asn	Ser	Thr	Thr	Leu	Tyr	
		35					40					45			

<210> 145

<211> 47

<212> PRT

<213> Artificial Sequence

<220>

<223> GCRTchick GPCR TM1

<400> 145

Cys	Ser	Thr	Glu	Asp	Ser	Phe	Lys	Tyr	Thr	Leu	Tyr	Gly	Cys	Val	Phe
1				5					10					15	
Ser	Met	Val	Phe	Val	Leu	Gly	Leu	Ile	Ala	Asn	Cys	Val	Ala	Ile	Tyr
			20					25					30		
Ile	Phe	Thr	Phe	Thr	Leu	Lys	Val	Arg	Asn	Glu	Thr	Thr	Thr	Tyr	
		35					40					45			

<210> 146

<211> 46

<212> PRT

<213> Artificial Sequence

<220>

<223> APJ GPCR TM1

<400> 146

Glu	Tyr	Thr	Asp	Trp	Lys	Ser	Ser	Gly	Ala	Leu	Ile	Pro	Ala	Ile	Tyr
1				5					10					15	
Met	Leu	Val	Phe	Leu	Leu	Gly	Thr	Thr	Gly	Asn	Gly	Leu	Val	Leu	Trp
			20					25					30		
Thr	Val	Phe	Arg	Ser	Ser	Arg	Glu	Lys	Arg	Arg	Ser	Ala	Asp		
		35					40					45			

<210> 147

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> RTArat GPCR TM1

<400> 147

Glu	Gln	Ile	Ala	Thr	Leu	Pro	Pro	Pro	Ala	Val	Thr	Asn	Tyr	Ile	Phe
1				5					10					15	
Leu	Leu	Leu	Cys	Leu	Cys	Gly	Leu	Val	Gly	Asn	Gly	Leu	Val	Leu	Trp
			20					25					30		
Phe	Phe	Gly	Phe	Ser	Ile	Lys	Arg	Thr	Pro						
		35					40								

<210> 148

6400

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> UHRrat GPCR TM1

<400> 148
 Ser Leu Gln Leu Val His Gln Leu Lys Gly Leu Ile Val Met Leu Tyr
 1 5 10 15
 Ser Ile Val Val Val Val Gly Leu Val Gly Asn Cys Leu Leu Val Leu
 20 25 30
 Val Ile Ala Arg Val Arg Arg Leu His Asn Val
 35 40

<210> 149
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> FMRL1N GPCR TM1

<400> 149
 Glu Pro Ala Gly His Thr Val Leu Trp Ile Phe Ser Leu Leu Val His
 1 5 10 15
 Gly Val Thr Phe Val Phe Gly Val Leu Gly Asn Gly Leu Val Ile Trp
 20 25 30
 Val Ala Gly Phe Arg Met Thr Arg Thr Val
 35 40

<210> 150
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> FMRL2 GPCR TM1

<400> 150
 Glu Ser Ala Gly Tyr Thr Val Leu Arg Ile Leu Pro Leu Val Val Leu
 1 5 10 15
 Gly Val Thr Phe Val Leu Gly Val Leu Gly Asn Gly Leu Val Ile Trp
 20 25 30
 Val Ala Gly Phe Arg Met Thr Arg Thr Val
 35 40

<210> 151
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> fMLP GPCR TM1

<400> 151
 Val Ser Ala Gly Tyr Leu Phe Leu Asp Ile Ile Thr Tyr Leu Val Phe
 1 5 10 15

<220>

64QQ

<223> OLF32Acatfish GPCR TM1

<400> 155

Ser	Gly	Phe	Ser	Gly	Ile	Pro	Phe	Ser	Gln	Tyr	Tyr	Phe	Ala	Phe	Leu
1				5					10					15	
Ile	Phe	Ile	Tyr	Ile	Ile	Ser	Leu	Cys	Gly	Asn	Ser	Ile	Val	Leu	Phe
			20					25					30		
Met	Ile	Leu	Val	Asp	Arg	Thr	Leu	His	Ile	Pro					
		35					40								

<210> 156

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF32Bcatfish, OLF32Ccatfish AND OLF32Dcatfish
GPCR TM1

<400> 156

Ser	Gly	Phe	Ser	Gly	Ile	Pro	Phe	Ser	Gln	Tyr	Tyr	Phe	Val	Phe	Leu
1				5					10					15	
Ile	Phe	Ile	Tyr	Ile	Ile	Ser	Leu	Cys	Gly	Asn	Ser	Ile	Val	Leu	Phe
			20					25					30		
Met	Ile	Leu	Val	Asp	Arg	Thr	Leu	His	Ile	Pro					
		35					40								

<210> 157

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF47catfish GPCR TM1

<400> 157

Ile	Ala	Tyr	Asn	Ser	Leu	Gly	Asn	Lys	Asn	Tyr	Leu	Ile	Leu	Ala	Leu
1				5					10					15	
Gly	Ile	Ile	Tyr	Leu	Ile	Thr	Leu	Leu	Cys	Asn	Phe	Thr	Leu	Leu	Ala
			20					25					30		
Ile	Ile	Leu	Met	Asn	Ser	Ser	Leu	Gln	Asn	Pro					
		35					40								

<210> 158

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF202catfish GPCR TM1

<400> 158

Phe	Pro	Gly	Leu	Pro	Pro	Asn	Tyr	Tyr	Gly	Leu	Val	Ser	Val	Val	Met
1				5					10					15	
Phe	Phe	Val	Tyr	Val	Cys	Thr	Leu	Ile	Gly	Asn	Cys	Thr	Phe	Phe	Thr
			20					25					30		
Leu	Phe	Leu	Arg	Glu	Lys	Ser	Leu	Gln	Lys	Pro					
		35					40								

64RR

<210> 159
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFCOR1chicken GPCR TM1

<400> 159
 Leu Thr Asp Asn Pro Gly Leu Gln Met Pro Leu Phe Met Val Phe Leu
 1 5 10 15
 Ala Ile Tyr Thr Ile Thr Leu Leu Thr Asn Leu Gly Leu Ile Ala Leu
 20 25 30
 Ile Ser Val Asp Leu His Leu Gln Thr Pro
 35 40

<210> 160
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFCOR2chicken GPCR TM1

<400> 160
 Leu Thr Asp Asn Pro Arg Leu Gln Met Pro Leu Phe Met Val Phe Leu
 1 5 10 15
 Val Ile Tyr Thr Thr Thr Leu Leu Thr Asn Leu Gly Leu Ile Ala Leu
 20 25 30
 Ile Gly Met Asp Leu His Leu Gln Thr Pro
 35 40

<210> 161
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFCOR3chicken AND OLFCOR4chicken GPCR TM1

<400> 161
 Leu Thr Asp Asn Pro Gly Leu Gln Met Pro Leu Phe Met Val Phe Leu
 1 5 10 15
 Ala Ile Tyr Thr Ile Thr Leu Leu Thr Asn Leu Gly Leu Ile Arg Leu
 20 25 30
 Ile Ser Val Asp Leu His Leu Gln Thr Pro
 35 40

<210> 162
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFCOR5chicken GPCR TM1

<400> 162
 Leu Thr Asp Asn Pro Arg Leu Gln Met Pro Leu Phe Met Val Phe Leu
 1 5 10 15

64SS

Ala Ile Tyr Thr Ile Thr Leu Leu Ala Asn Leu Gly Leu Ile Ala Leu
 20 25 30
 Ile Ser Val Asp Phe His Leu Gln Thr Pro
 35 40

<210> 163
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFCOR6chicken GPCR TM1

<400> 163
 Leu Thr Asp Asn Pro Gly Leu Gln Met Pro Leu Phe Met Val Phe Leu
 1 5 10 15
 Ala Ile Tyr Thr Ile Thr Leu Leu Thr Asn Leu Gly Leu Ile Ala Leu
 20 25 30
 Ile Arg Ile Asp Leu Gln Leu Gln Thr Pro
 35 40

<210> 164
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFdog GPCR TM1

<400> 164
 Leu Pro Ile Asp Pro Asp Gln Arg Asp Leu Phe Tyr Ala Leu Phe Leu
 1 5 10 15
 Ala Met Tyr Val Thr Thr Ile Leu Gly Asn Leu Leu Ile Ile Val Leu
 20 25 30
 Ile Gln Leu Asp Ser His Leu His Thr Pro
 35 40

<210> 165
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLF07E GPCR TM1

<400> 165
 Met Ser Glu Ser Pro Glu Gln Gln Gln Ile Leu Phe Trp Met Phe Leu
 1 5 10 15
 Ser Met Tyr Leu Val Thr Val Val Gly Asn Val Leu Ile Ile Leu Ala
 20 25 30
 Ile Ser Ser Asp Ser Arg Leu His Thr Pro
 35 40

<210> 166
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>

64TT

<223> OLF07I GPCR TM1

<400> 166

Leu	Pro	Ile	Gln	Pro	Glu	Gln	Gln	Asn	Leu	Cys	Tyr	Ala	Leu	Phe	Leu
1				5					10					15	
Ala	Met	Tyr	Leu	Thr	Thr	Leu	Leu	Gly	Asn	Leu	Leu	Ile	Ile	Val	Leu
			20					25					30		
Ile	Arg	Leu	Asp	Ser	His	Leu	His	Thr	Pro						
		35					40								

<210> 167

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF07J GPCR TM1

<400> 167

Phe	Ser	Ser	Phe	His	Glu	Gln	Gln	Ile	Thr	Leu	Phe	Gly	Val	Phe	Leu
1				5					10					15	
Ala	Leu	Tyr	Ile	Leu	Thr	Leu	Ala	Gly	Asn	Ile	Ile	Ile	Val	Thr	Ile
			20					25					30		
Ile	Arg	Ile	Asp	Leu	His	Leu	His	Thr	Pro						
		35					40								

<210> 168

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF0R3mouse GPCR TM1

<400> 168

Val	Ser	Asp	His	Pro	His	Leu	Glu	Ile	Ile	Phe	Phe	Ala	Val	Ile	Leu
1				5					10					15	
Ala	Ser	Tyr	Leu	Leu	Thr	Leu	Val	Gly	Asn	Leu	Thr	Ile	Ile	Leu	Leu
			20					25					30		
Ser	Arg	Leu	Asp	Ala	Arg	Leu	His	Thr	Pro						
		35					40								

<210> 169

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> OLFrat GPCR TM1

<400> 169

Leu	Thr	Lys	Gln	Pro	Glu	Leu	Leu	Leu	Pro	Leu	Phe	Phe	Leu	Phe	Leu
1				5					10					15	
Val	Ile	Tyr	Val	Leu	Thr	Val	Val	Gly	Asn	Leu	Gly	Met	Ile	Leu	Leu
			20					25					30		
Ile	Ile	Val	Ser	Pro	Leu	Leu	His	Thr	Pro						
		35					40								

<210> 170

64UU

<211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFF3rat GPCR TM1

<400> 170
 Phe Val Glu Asn Lys Asp Leu Gln Pro Leu Ile Tyr Gly Leu Phe Leu
 1 5 10 15
 Ser Met Tyr Leu Val Thr Val Ile Gly Asn Ile Ser Ile Ile Val Ala
 20 25 30
 Ile Ile Ser Asp Pro Cys Leu His Thr Pro
 35 40

<210> 171
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFF5rat GPCR TM1

<400> 171
 Leu Ser Arg Gln Pro Gln Gln Gln Gln Leu Leu Phe Leu Leu Phe Leu
 1 5 10 15
 Ile Met Tyr Leu Ala Thr Val Leu Gly Asn Leu Leu Ile Ile Leu Ala
 20 25 30
 Ile Gly Thr Asp Ser Arg Leu His Thr Pro
 35 40

<210> 172
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFF6rat GPCR TM1

<400> 172
 Phe Pro Gly Pro Arg Ser Met Arg Ile Gly Leu Phe Leu Leu Phe Leu
 1 5 10 15
 Val Met Tyr Leu Leu Thr Val Val Gly Asn Leu Ala Ile Ile Ser Leu
 20 25 30
 Val Gly Ala His Arg Cys Leu Gln Thr Pro
 35 40

<210> 173
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFF12rat GPCR TM1

<400> 173
 Phe Thr Glu Asn Pro Gln Leu His Phe Leu Ile Phe Ala Leu Phe Leu
 1 5 10 15

64VV

Ser Met Tyr Leu Val Thr Val Leu Gly Asn Leu Leu Ile Ile Met Ala
 20 25 30
 Ile Ile Thr Gln Ser His Leu His Thr Pro
 35 40

<210> 174
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFI3rat GPCR TM1

<400> 174
 Leu Pro Ile Pro Glu Glu His Gln His Leu Phe Tyr Ala Leu Phe Leu
 1 5 10 15
 Val Met Tyr Leu Thr Thr Ile Leu Gly Asn Leu Leu Ile Ile Val Leu
 20 25 30
 Val Gln Leu Asp Ser Gln Leu His Thr Pro
 35 40

<210> 175
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFI7rat GPCR TM1

<221> MOD_RES
 <222> (18)...(18)
 <223> Xaa = unknown amino acid

<400> 175
 Phe Pro Ala Pro Ala Pro Leu Arg Val Leu Leu Phe Phe Leu Ser Leu
 1 5 10 15
 Leu Xaa Tyr Val Leu Val Leu Thr Glu Asn Met Leu Ile Ile Ile Ala
 20 25 30
 Ile Arg Asn His Pro Thr Leu His Lys Pro
 35 40

<210> 176
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFI8rat GPCR TM1

<400> 176
 Leu Pro Ile Pro Pro Glu His Gln Gln Leu Phe Phe Ala Leu Phe Leu
 1 5 10 15
 Ile Met Tyr Leu Thr Thr Phe Leu Gly Asn Leu Leu Ile Val Val Leu
 20 25 30
 Val Gln Leu Asp Ser His Leu His Thr Pro
 35 40

<210> 177
 <211> 42

64WW

<212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFI9rat GPCR TM1

<400> 177
 Leu Pro Phe Pro Pro Glu Tyr Gln His Leu Phe Tyr Ala Leu Phe Leu
 1 5 10 15
 Ala Met Tyr Leu Thr Thr Leu Leu Gly Asn Leu Ile Ile Ile Ile Leu
 20 25 30
 Ile Leu Leu Asp Ser His Leu His Thr Pro
 35 40

<210> 178
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFI14rat GPCR TM1

<400> 178
 Leu Pro Ile Pro Ser Glu Tyr His Leu Leu Phe Tyr Ala Leu Phe Leu
 1 5 10 15
 Ala Met Tyr Leu Thr Ile Ile Leu Gly Asn Leu Leu Ile Ile Val Leu
 20 25 30
 Val Arg Leu Asp Ser His Leu His Met Pro
 35 40

<210> 179
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFI15rat GPCR TM1

<400> 179
 Leu Pro Ile Pro Ser Glu His Gln His Val Phe Tyr Ala Leu Phe Leu
 1 5 10 15
 Ser Met Tyr Leu Thr Thr Val Leu Gly Asn Leu Ile Ile Ile Ile Leu
 20 25 30
 Ile His Leu Asp Ser His Leu His Thr Pro
 35 40

<210> 180
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFOR17_40 GPCR TM1

<400> 180
 Leu Leu Glu Ala Pro Gly Leu Gln Pro Val Val Phe Val Leu Phe Leu
 1 5 10 15
 Phe Ala Tyr Leu Val Thr Val Arg Gly Asn Leu Ser Ile Leu Ala Ala
 20 25 30

64XX

Val Leu Val Glu Pro Lys Leu His Thr Pro
 35 40

<210> 181
 <211> 39
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> GUST27rat GPCR TM1

<400> 181
 Met Ile Leu Asn Cys Asn Pro Phe Ser Gly Leu Phe Leu Ser Met Tyr
 1 5 10 15
 Leu Val Thr Val Leu Gly Asn Leu Leu Ile Ile Leu Ala Val Ser Ser
 20 25 30
 Asn Ser His Leu His Asn Leu
 35

<210> 182
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> RPE GPCR TM1

<400> 182
 Pro Thr Gly Phe Gly Glu Leu Glu Val Leu Ala Val Gly Met Val Leu
 1 5 10 15
 Leu Val Glu Ala Leu Ser Gly Leu Ser Leu Asn Thr Leu Thr Ile Phe
 20 25 30
 Ser Phe Cys Lys Thr Pro Glu Leu Arg Thr Pro
 35 40

<210> 183
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> HHRF1 GPCR TM1

<400> 183
 Phe Thr Asp Val Leu Asn Gln Ser Lys Pro Val Thr Leu Phe Leu Tyr
 1 5 10 15
 Gly Val Val Phe Leu Phe Gly Ser Ile Gly Asn Phe Leu Val Ile Phe
 20 25 30
 Thr Ile Thr Trp Arg Arg Arg Ile Gln Cys Ser
 35 40

<210> 184
 <211> 44
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> HHRF2 GPCR TM1

64YY

<400> 184
 Asn Ser Thr Glu Ile Tyr Gln Leu Phe Glu Tyr Thr Arg Leu Gly Val
 1 5 10 15
 Trp Leu Met Cys Ile Val Gly Thr Phe Leu Asn Val Leu Val Ile Thr
 20 25 30
 Thr Ile Leu Tyr Tyr Arg Arg Lys Lys Lys Ser Pro
 35 40

<210> 185
 <211> 45
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> HHRF3 GPCR TM1

<400> 185
 Met Thr Gly Pro Leu Phe Ala Ile Arg Thr Thr Glu Ala Val Leu Asn
 1 5 10 15
 Thr Phe Ile Ile Phe Val Gly Gly Pro Leu Asn Ala Ile Val Leu Ile
 20 25 30
 Thr Gln Leu Leu Thr Asn Arg Val Leu Gly Tyr Ser Thr
 35 40 45

<210> 186
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCP-1A AND MCP-1B GPCR TM1

<400> 186
 Asp Val Lys Gln Ile Gly Ala Gln Leu Leu Pro Pro Leu Tyr Ser Leu
 1 5 10 15
 Val Phe Ile Phe Gly Phe Val Gly Asn Met Leu Val Val Leu Ile Leu
 20 25 30
 Ile Asn Cys Lys Lys Leu Lys Cys Leu
 35 40

<210> 187
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PPR1bovine GPCR TM1

<400> 187
 Glu Val Arg Lys Phe Ala Lys Val Phe Leu Pro Ala Phe Phe Thr Ile
 1 5 10 15
 Ala Phe Ile Ile Gly Leu Ala Gly Asn Ser Thr Val Val Ala Ile Tyr
 20 25 30
 Ala Tyr Tyr Lys Lys Arg Arg Thr Lys
 35 40

<210> 188
 <211> 50
 <212> PRT

64ZZ

<213> Artificial Sequence

<220>

<223> GPCRAelegans GPCR TM2

<400> 188

Thr	Arg	Phe	Leu	Met	Cys	Asn	Leu	Ala	Phe	Ala	Asp	Phe	Ile	Leu	Gly
1				5				10						15	
Leu	Tyr	Ile	Phe	Ile	Leu	Thr	Ser	Val	Ser	Ala	Val	Thr	Arg	Gly	Asp
			20					25					30		
Tyr	His	Asn	Tyr	Val	Gln	Gln	Trp	Gln	Asn	Gly	Ala	Gly	Cys	Lys	Ile
		35					40					45			
Leu	Gly														
50															

<210> 189

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> GRH GPCR TM2

<400> 189

Lys	Leu	Leu	Leu	Lys	His	Leu	Thr	Leu	Ala	Asn	Leu	Leu	Glu	Thr	Leu
1				5				10					15		
Ile	Val	Met	Pro	Leu	Asp	Gly	Met	Trp	Asn	Ile	Thr	Val	Gln	Trp	Tyr
			20					25					30		
Ala	Gly	Glu	Leu	Leu	Cys	Lys	Val	Leu	Ser						
		35					40								

<210> 190

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> TRH GPCR TM2

<400> 190

Thr	Asn	Cys	Tyr	Leu	Val	Ser	Leu	Ala	Val	Ala	Asp	Leu	Met	Val	Leu
1				5				10					15		
Val	Ala	Ala	Gly	Leu	Pro	Asn	Ile	Thr	Asp	Ser	Ile	Tyr	Gly	Ser	Trp
			20					25					30		
Val	Tyr	Gly	Tyr	Val	Gly	Cys	Leu	Cys	Ile	Thr					
		35					40								

<210> 191

<211> 50

<212> PRT

<213> Artificial Sequence

<220>

<223> FSHprec GPCR TM2

<400> 191

Pro	Arg	Phe	Leu	Met	Cys	Asn	Leu	Ala	Phe	Ala	Asp	Leu	Cys	Ile	Gly
1				5				10					15		

64AAA

Ile Tyr Leu Leu Leu Ile Ala Ser Val Asp Ile His Thr Lys Ser Gln
 20 25 30
 Tyr His Asn Tyr Ala Ile Asp Trp Gln Thr Gly Ala Gly Cys Asp Ala
 35 40 45
 Ala Gly
 50

<210> 192
 <211> 50
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> TSHprec GPCR TM2

<400> 192
 Pro Arg Phe Leu Met Cys Asn Leu Ala Phe Ala Asp Phe Cys Met Gly
 1 5 10 15
 Met Tyr Leu Leu Leu Ile Ala Ser Val Asp Leu Tyr Thr His Ser Glu
 20 25 30
 Tyr Tyr Asn His Ala Ile Asp Trp Gln Thr Gly Pro Gly Cys Asn Thr
 35 40 45
 Ala Gly
 50

<210> 193
 <211> 50
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> LH_CGprec GPCR TM2

<400> 193
 Pro Arg Phe Leu Met Cys Asn Leu Ser Phe Ala Asp Phe Cys Met Gly
 1 5 10 15
 Leu Tyr Leu Leu Leu Ile Ala Ser Val Asp Ser Gln Thr Lys Gly Gln
 20 25 30
 Tyr Tyr Asn His Ala Ile Asp Trp Gln Thr Gly Ser Gly Cys Ser Thr
 35 40 45
 Ala Gly
 50

<210> 194
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PGE_EP1 GPCR TM2

<400> 194
 Phe Leu Leu Phe Val Ala Ser Leu Leu Ala Thr Asp Leu Ala Gly His
 1 5 10 15
 Val Ile Pro Gly Ala Leu Val Leu Arg Leu Tyr Thr Ala Gly Arg Ala
 20 25 30
 Pro Ala Gly Gly Ala Cys His Phe Leu Gly
 35 40

64BBB

<210> 195
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PGE_EP2 GPCR TM2

<400> 195
 Phe Tyr Thr Leu Val Cys Gly Leu Ala Val Thr Asp Leu Leu Gly Thr
 1 5 10 15
 Leu Leu Val Ser Pro Val Thr Ile Ala Thr Tyr Met Lys Gly Gln Trp
 20 25 30
 Pro Gly Gly Gln Pro Leu Cys Glu Tyr Ser Thr
 35 40

<210> 196
 <211> 47
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PGE_EP3 GPCR TM2

<400> 196
 Phe His Val Leu Val Thr Glu Leu Val Phe Thr Asp Leu Leu Gly Thr
 1 5 10 15
 Cys Leu Ile Ser Pro Val Val Leu Ala Ser Tyr Ala Arg Asn Gln Thr
 20 25 30
 Leu Val Ala Leu Ala Pro Glu Ser Arg Ala Cys Thr Tyr Phe Ala
 35 40 45

<210> 197
 <211> 47
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PGF GPCR TM2

<400> 197
 Phe Leu Leu Leu Ala Ser Gly Leu Val Ile Thr Asp Phe Phe Gly His
 1 5 10 15
 Leu Ile Asn Gly Ala Ile Ala Val Phe Val Tyr Ala Ser Asp Lys Glu
 20 25 30
 Trp Ile Arg Phe Asp Gln Ser Asn Val Leu Cys Ser Ile Phe Gly
 35 40 45

<210> 198
 <211> 48
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PGI GPCR TM2

<400> 198
 Phe Ala Val Leu Val Thr Gly Leu Ala Ala Thr Asp Leu Leu Gly Thr
 1 5 10 15

64CCC

Ser Phe Leu Ser Pro Ala Val Phe Val Ala Tyr Ala Arg Asn Ser Ser
 20 25 30
 Leu Leu Gly Leu Ala Arg Gly Gly Pro Ala Leu Cys Asp Ala Phe Ala
 35 40 45

<210> 199
 <211> 47
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> TXA2 GPCR TM2

<400> 199
 Phe Leu Thr Phe Leu Cys Gly Leu Val Leu Thr Asp Phe Leu Gly Leu
 1 5 10 15
 Leu Val Thr Gly Thr Ile Val Val Ser Gln His Ala Ala Leu Phe Glu
 20 25 30
 Trp His Ala Val Asp Pro Gly Cys Arg Leu Cys Arg Phe Met Gly
 35 40 45

<210> 200
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> PAF GPCR TM2

<400> 200
 Ile Phe Met Val Asn Leu Thr Met Ala Asp Met Leu Phe Leu Ile Thr
 1 5 10 15
 Leu Pro Leu Trp Ile Val Tyr Tyr Gln Asn Gln Gly Asn Trp Ile Leu
 20 25 30
 Pro Lys Phe Leu Cys Asn Val Ala Gly
 35 40

<210> 201
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> M2 GPCR TM2

<400> 201
 Asn Asn Tyr Phe Leu Phe Ser Leu Ala Cys Ala Asp Leu Ile Ile Gly
 1 5 10 15
 Val Phe Ser Met Asn Leu Tyr Thr Leu Tyr Thr Val Ile Gly Tyr Trp
 20 25 30
 Pro Leu Gly Pro Val Val Cys Asp Leu Trp Leu
 35 40

<210> 202
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64DDD

<223> M4 GPCR TM2

<400> 202

Asn	Asn	Tyr	Phe	Leu	Phe	Ser	Leu	Ala	Cys	Ala	Asp	Leu	Ile	Ile	Gly
1				5					10					15	
Ala	Phe	Ser	Met	Asn	Leu	Tyr	Thr	Val	Tyr	Ile	Ile	Lys	Gly	Tyr	Trp
			20					25					30		
Pro	Leu	Gly	Ala	Val	Val	Cys	Asp	Leu	Trp	Leu					
		35					40								

<210> 203

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> M1 GPCR TM2

<400> 203

Asn	Asn	Tyr	Phe	Leu	Leu	Ser	Leu	Ala	Cys	Ala	Asp	Leu	Ile	Ile	Gly
1				5					10					15	
Thr	Phe	Ser	Met	Asn	Leu	Tyr	Thr	Thr	Tyr	Leu	Leu	Met	Gly	His	Trp
			20					25					30		
Ala	Leu	Gly	Thr	Leu	Ala	Cys	Asp	Leu	Trp	Leu					
		35					40								

<210> 204

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> M3 GPCR TM2

<400> 204

Asn	Asn	Tyr	Phe	Leu	Leu	Ser	Leu	Ala	Cys	Ala	Asp	Leu	Ile	Ile	Gly
1				5					10					15	
Val	Ile	Ser	Met	Asn	Leu	Phe	Thr	Thr	Tyr	Ile	Ile	Met	Asn	Arg	Trp
			20					25					30		
Ala	Leu	Gly	Asn	Leu	Ala	Cys	Asp	Leu	Trp	Leu					
		35					40								

<210> 205

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> M5 GPCR TM2

<400> 205

Asn	Asn	Tyr	Tyr	Leu	Leu	Ser	Leu	Ala	Cys	Ala	Asp	Leu	Ile	Ile	Gly
1				5					10					15	
Ile	Phe	Ser	Met	Asn	Leu	Tyr	Thr	Thr	Tyr	Ile	Leu	Met	Gly	Arg	Trp
			20					25					30		
Ala	Leu	Gly	Ser	Leu	Ala	Cys	Asp	Leu	Trp	Leu					
		35					40								

<210> 206

64EEE

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> H1 GPCR TM2

<400> 206
 Gly Asn Leu Tyr Ile Val Ser Leu Ser Val Ala Asp Leu Ile Val Gly
 1 5 10 15
 Ala Val Val Met Pro Met Asn Ile Leu Tyr Leu Leu Met Ser Lys Trp
 20 25 30
 Ser Leu Gly Arg Pro Leu Cys Leu Phe Trp Leu
 35 40

<210> 207
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> H2 GPCR TM2

<400> 207
 Thr Asn Cys Phe Ile Val Ser Leu Ala Ile Thr Asp Leu Leu Leu Gly
 1 5 10 15
 Leu Leu Val Leu Pro Phe Ser Ala Ile Tyr Gln Leu Ser Cys Lys Trp
 20 25 30
 Ser Phe Gly Lys Val Phe Cys Asn Ile Tyr Thr
 35 40

<210> 208
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT1A GPCR TM2

<400> 208
 Ala Asn Tyr Leu Ile Gly Ser Leu Ala Val Thr Asp Leu Met Val Ser
 1 5 10 15
 Val Leu Val Leu Pro Met Ala Ala Leu Tyr Gln Val Leu Asn Lys Trp
 20 25 30
 Thr Leu Gly Gln Val Thr Cys Asp Leu Phe Ile
 35 40

<210> 209
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT1B GPCR TM2

<400> 209
 Ala Asn Tyr Leu Ile Ala Ser Leu Ala Val Thr Asp Leu Leu Val Ser
 1 5 10 15

64FFF

Ile Leu Val Met Pro Ile Ser Thr Met Tyr Thr Val Thr Gly Arg Trp
 20 25 30
 Thr Leu Gly Gln Val Val Cys Asp Phe Trp Leu
 35 40

<210> 210
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT1D GPCR TM2

<400> 210
 Ala Asn Tyr Leu Ile Gly Ser Leu Ala Thr Thr Asp Leu Leu Val Ser
 1 5 10 15
 Ile Leu Val Met Pro Ile Ser Ile Ala Tyr Thr Ile Thr His Thr Trp
 20 25 30
 Asn Phe Gly Gln Ile Leu Cys Asp Ile Trp Leu
 35 40

<210> 211
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT1E GPCR TM2

<400> 211
 Ala Asn Tyr Leu Ile Cys Ser Leu Ala Val Thr Asp Leu Leu Val Ala
 1 5 10 15
 Val Leu Val Met Pro Leu Ser Ile Ile Tyr Ile Val Met Asp Arg Trp
 20 25 30
 Lys Leu Gly Tyr Phe Leu Cys Glu Val Trp Leu
 35 40

<210> 212
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT1F GPCR TM2

<400> 212
 Ala Asn Tyr Leu Ile Cys Ser Leu Ala Val Thr Asp Phe Leu Val Ala
 1 5 10 15
 Val Leu Val Met Pro Phe Ser Ile Val Tyr Ile Val Arg Glu Ser Trp
 20 25 30
 Ile Met Gly Gln Val Val Cys Asp Ile Trp Leu
 35 40

<210> 213
 <211> 44
 <212> PRT
 <213> Artificial Sequence

<220>

64GGG

<223> 5HT2A GPCR TM2

<400> 213

Thr	Asn	Tyr	Phe	Leu	Met	Ser	Leu	Ala	Ile	Ala	Asp	Met	Leu	Leu	Gly
1				5					10					15	
Phe	Leu	Val	Met	Pro	Val	Ser	Met	Leu	Thr	Ile	Leu	Tyr	Gly	Tyr	Arg
			20					25					30		
Trp	Pro	Leu	Pro	Ser	Lys	Leu	Cys	Ala	Val	Trp	Ile				
		35					40								

<210> 214

<211> 44

<212> PRT

<213> Artificial Sequence

<220>

<223> 5HT2B GPCR TM2

<400> 214

Thr	Asn	Tyr	Phe	Leu	Met	Ser	Leu	Ala	Val	Ala	Asp	Leu	Leu	Val	Gly
1				5					10					15	
Leu	Phe	Val	Met	Pro	Ile	Ala	Leu	Leu	Thr	Ile	Met	Phe	Glu	Ala	Met
			20					25					30		
Trp	Pro	Leu	Pro	Leu	Val	Leu	Cys	Pro	Ala	Trp	Leu				
		35					40								

<210> 215

<211> 44

<212> PRT

<213> Artificial Sequence

<220>

<223> 5HT2C GPCR TM2

<400> 215

Thr	Asn	Tyr	Phe	Leu	Met	Ser	Leu	Ala	Ile	Ala	Asp	Met	Leu	Val	Gly
1				5					10					15	
Leu	Leu	Val	Met	Pro	Leu	Ser	Leu	Leu	Ala	Ile	Leu	Tyr	Asp	Tyr	Val
			20					25					30		
Trp	Pro	Leu	Pro	Arg	Tyr	Leu	Cys	Pro	Val	Trp	Ile				
		35					40								

<210> 216

<211> 44

<212> PRT

<213> Artificial Sequence

<220>

<223> 5HT5A GPCR TM2

<400> 216

Pro	His	Asn	Leu	Val	Ala	Ser	Met	Ala	Val	Ser	Asp	Val	Leu	Val	Ala
1				5					10					15	
Ala	Leu	Val	Met	Pro	Leu	Ser	Leu	Val	His	Glu	Leu	Ser	Gly	Arg	Arg
			20					25					30		
Trp	Gln	Leu	Gly	Arg	Arg	Leu	Cys	Gln	Leu	Trp	Ile				
		35					40								

<210> 217

64HHH

<211> 45
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT5Brat GPCR TM2

<400> 217
 Pro His Asn Leu Val Ala Ser Thr Ala Val Ser Asp Val Leu Val Ala
 1 5 10 15
 Ala Leu Val Met Pro Leu Ser Leu Val Ser Glu Leu Ser Ala Gly Arg
 20 25 30
 Arg Trp Gln Leu Gly Arg Ser Leu Cys His Val Trp Ile
 35 40 45

<210> 218
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT6rat GPCR TM2

<400> 218
 Ser Asn Phe Phe Leu Val Ser Leu Phe Thr Ser Asp Leu Met Val Gly
 1 5 10 15
 Leu Val Val Met Pro Pro Ala Met Leu Asn Ala Leu Tyr Gly Arg Trp
 20 25 30
 Val Leu Ala Arg Gly Leu Cys Leu Leu Trp Thr
 35 40

<210> 219
 <211> 44
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 5HT7 GPCR TM2

<400> 219
 Ser Asn Tyr Leu Ile Val Ser Leu Ala Leu Ala Asp Leu Ser Val Ala
 1 5 10 15
 Val Ala Val Met Pro Phe Val Ser Val Thr Asp Leu Ile Gly Gly Lys
 20 25 30
 Trp Ile Phe Gly His Phe Phe Cys Asn Val Phe Ile
 35 40

<210> 220
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> alpha1A GPCR TM2

<400> 220
 Thr Asn Tyr Phe Ile Val Asn Leu Ala Val Ala Asp Leu Leu Leu Ser
 1 5 10 15

64III

Ala Thr Val Leu Pro Phe Ser Ala Thr Met Glu Val Leu Gly Phe Trp
 20 25 30
 Ala Phe Gly Arg Ala Phe Cys Asp Val Trp Ala
 35 40

<210> 221
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> alpha1B GPCR TM2

<400> 221
 Thr Asn Tyr Phe Ile Val Asn Leu Ala Met Ala Asp Leu Leu Leu Ser
 1 5 10 15
 Phe Thr Val Leu Pro Phe Ser Ala Ala Leu Glu Val Leu Gly Tyr Trp
 20 25 30
 Val Leu Gly Arg Ile Phe Cys Asp Ile Trp Ala
 35 40

<210> 222
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> alpha1C GPCR TM2

<400> 222
 Thr His Tyr Tyr Ile Val Asn Leu Ala Val Ala Asp Leu Leu Leu Thr
 1 5 10 15
 Ser Thr Val Leu Pro Phe Ser Ala Ile Phe Glu Val Leu Gly Tyr Trp
 20 25 30
 Ala Phe Gly Arg Val Phe Cys Asn Ile Trp Ala
 35 40

<210> 223
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> alpha2A GPCR TM2

<400> 223
 Gln Asn Leu Phe Leu Val Ser Leu Ala Ser Ala Asp Ile Leu Val Ala
 1 5 10 15
 Thr Leu Val Ile Pro Phe Ser Leu Ala Asn Glu Val Met Gly Tyr Trp
 20 25 30
 Tyr Phe Gly Lys Ala Trp Cys Glu Ile Tyr Leu
 35 40

<210> 224
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>

64JJJ

<223> alpha2B GPCR TM2

<400> 224

Gln	Asn	Leu	Phe	Leu	Val	Ser	Leu	Ala	Ala	Ala	Asp	Ile	Leu	Val	Ala
1				5				10						15	
Thr	Leu	Ile	Ile	Pro	Phe	Ser	Leu	Ala	Asn	Glu	Leu	Leu	Gly	Tyr	Trp
			20					25					30		
Tyr	Phe	Arg	Arg	Thr	Trp	Cys	Glu	Val	Tyr	Leu					
		35					40								

<210> 225

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> alpha2C1 AND alpha2C2 GPCR TM2

<400> 225

Gln	Asn	Leu	Phe	Leu	Val	Ser	Leu	Ala	Ser	Ala	Asp	Ile	Leu	Val	Ala
1				5				10						15	
Thr	Leu	Val	Met	Pro	Phe	Ser	Leu	Ala	Asn	Glu	Leu	Met	Ala	Tyr	Trp
			20					25					30		
Tyr	Phe	Gly	Gln	Val	Trp	Cys	Gly	Val	Tyr	Leu					
		35					40								

<210> 226

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> beta1 GPCR TM2

<400> 226

Thr	Asn	Leu	Phe	Ile	Met	Ser	Leu	Ala	Ser	Ala	Asp	Leu	Val	Met	Gly
1				5				10						15	
Leu	Leu	Val	Val	Pro	Phe	Gly	Ala	Thr	Ile	Val	Val	Trp	Gly	Arg	Trp
			20					25					30		
Glu	Tyr	Gly	Ser	Phe	Phe	Cys	Glu	Leu	Trp	Thr					
		35					40								

<210> 227

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> beta2 GPCR TM2

<400> 227

Thr	Asn	Tyr	Phe	Ile	Thr	Ser	Leu	Ala	Cys	Ala	Asp	Leu	Val	Met	Gly
1				5				10						15	
Leu	Ala	Val	Val	Pro	Phe	Gly	Ala	Ala	His	Ile	Leu	Met	Lys	Met	Trp
			20					25					30		
Thr	Phe	Gly	Asn	Phe	Trp	Cys	Glu	Phe	Trp	Thr					
		35					40								

<210> 228

64KKK

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> beta3 GPCR TM2

<400> 228
 Thr Asn Val Phe Val Thr Ser Leu Ala Ala Ala Asp Leu Val Met Gly
 1 5 10 15
 Leu Leu Val Val Pro Pro Ala Ala Thr Leu Ala Leu Thr Gly His Trp
 20 25 30
 Pro Leu Gly Ala Thr Gly Cys Glu Leu Trp Thr
 35 40

<210> 229
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> beta4turkey GPCR TM2

<400> 229
 Thr Asn Val Phe Val Thr Ser Leu Ala Cys Ala Asp Leu Val Met Gly
 1 5 10 15
 Leu Leu Val Val Pro Pro Gly Ala Thr Ile Leu Leu Ser Gly His Trp
 20 25 30
 Pro Tyr Gly Thr Val Val Cys Glu Leu Trp Thr
 35 40

<210> 230
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> D1A GPCR TM2

<400> 230
 Thr Asn Phe Phe Val Ile Ser Leu Ala Val Ser Asp Leu Leu Val Ala
 1 5 10 15
 Val Leu Val Met Pro Trp Lys Ala Val Ala Glu Ile Ala Gly Phe Trp
 20 25 30
 Pro Phe Gly Ser Phe Cys Asn Ile Trp Val
 35 40

<210> 231
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> D2 GPCR TM2

<400> 231
 Thr Asn Tyr Leu Ile Val Ser Leu Ala Val Ala Asp Leu Leu Val Ala
 1 5 10 15

64LLL

Thr Leu Val Met Pro Trp Val Val Tyr Leu Glu Val Val Gly Glu Trp
 20 25 30
 Lys Phe Ser Arg Ile His Cys Asp Ile Phe Val
 35 40

<210> 232
 <211> 44
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> D3 GPCR TM2

<400> 232
 Thr Asn Tyr Leu Val Val Ser Leu Ala Val Ala Asp Leu Leu Val Ala
 1 5 10 15
 Thr Leu Val Met Pro Trp Val Val Tyr Leu Glu Val Thr Gly Gly Val
 20 25 30
 Trp Asn Phe Ser Arg Ile Cys Cys Asp Val Phe Val
 35 40

<210> 233
 <211> 44
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> D4 GPCR TM2

<400> 233
 Thr Asn Ser Phe Ile Val Ser Leu Ala Ala Ala Asp Leu Leu Leu Ala
 1 5 10 15
 Leu Leu Val Leu Pro Leu Phe Val Tyr Ser Glu Val Gln Gly Gly Ala
 20 25 30
 Trp Leu Leu Ser Pro Arg Leu Cys Asp Ala Leu Met
 35 40

<210> 234
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> D5 GPCR TM2

<400> 234
 Thr Asn Val Phe Ile Val Ser Leu Ala Val Ser Asp Leu Phe Val Ala
 1 5 10 15
 Leu Leu Val Met Pro Trp Lys Ala Val Ala Glu Val Ala Gly Tyr Trp
 20 25 30
 Pro Phe Gly Ala Phe Cys Asp Val Trp Val
 35 40

<210> 235
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>

64MMM

<223> A1 GPCR TM2

<400> 235

Thr	Phe	Cys	Phe	Ile	Val	Ser	Leu	Ala	Val	Ala	Asp	Val	Ala	Val	Gly
1				5					10					15	
Ala	Leu	Val	Ile	Pro	Leu	Ala	Ile	Leu	Ile	Asn	Ile	Gly	Pro	Gln	Thr
			20					25					30		
Tyr	Phe	His	Thr	Cys	Leu	Met	Val	Ala							
		35					40								

<210> 236

<211> 41

<212> PRT

<213> Artificial Sequence

<220>

<223> A2a GPCR TM2

<400> 236

Thr	Asn	Tyr	Phe	Val	Val	Ser	Leu	Ala	Ala	Ala	Asp	Ile	Ala	Val	Gly
1				5					10					15	
Val	Leu	Ala	Ile	Pro	Phe	Ala	Ile	Thr	Ile	Ser	Thr	Gly	Phe	Cys	Ala
			20					25					30		
Ala	Cys	His	Gly	Cys	Leu	Phe	Ile	Ala							
		35					40								

<210> 237

<211> 41

<212> PRT

<213> Artificial Sequence

<220>

<223> A2b GPCR TM2

<400> 237

Thr	Asn	Tyr	Phe	Leu	Val	Ser	Leu	Ala	Ala	Ala	Asp	Val	Ala	Val	Gly
1				5					10					15	
Leu	Phe	Ala	Ile	Pro	Phe	Ala	Ile	Thr	Ile	Ser	Leu	Gly	Phe	Cys	Thr
			20					25					30		
Asp	Phe	Tyr	Gly	Cys	Leu	Phe	Leu	Ala							
		35					40								

<210> 238

<211> 41

<212> PRT

<213> Artificial Sequence

<220>

<223> A3 GPCR TM2

<400> 238

Thr	Phe	Tyr	Phe	Ile	Val	Ser	Leu	Ala	Leu	Ala	Asp	Ile	Ala	Val	Gly
1				5					10					15	
Val	Leu	Val	Met	Pro	Leu	Ala	Ile	Val	Val	Ser	Leu	Gly	Ile	Thr	Ile
			20					25					30		
His	Phe	Tyr	Ser	Cys	Leu	Phe	Met	Thr							
		35					40								

<210> 239

64NNN

<211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OCdrome GPCR TM2

<400> 239
 Gln Asn Phe Phe Ile Val Ser Leu Ala Val Ala Asp Leu Thr Val Ala
 1 5 10 15
 Leu Leu Val Leu Pro Phe Asn Val Ala Tyr Ser Ile Leu Gly Arg Trp
 20 25 30
 Glu Phe Gly Ile His Leu Cys Lys Leu Trp Leu
 35 40

<210> 240
 <211> 49
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> ACTH GPCR TM2

<400> 240
 Met Tyr Phe Phe Ile Cys Ser Leu Ala Ile Ser Asp Met Leu Gly Ser
 1 5 10 15
 Leu Tyr Lys Ile Leu Glu Asn Ile Leu Ile Ile Leu Arg Asn Met Gly
 20 25 30
 Tyr Leu Lys Pro Arg Gly Ser Phe Glu Thr Thr Ala Asp Asp Ile Ile
 35 40 45
 Asp

<210> 241
 <211> 49
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MSH GPCR TM2

<400> 241
 Met Tyr Cys Phe Ile Cys Cys Leu Ala Leu Ser Asp Leu Leu Val Ser
 1 5 10 15
 Gly Thr Asn Val Leu Glu Thr Ala Val Ile Leu Leu Leu Glu Ala Gly
 20 25 30
 Ala Leu Val Ala Arg Ala Ala Val Leu Gln Gln Leu Asp Asn Val Ile
 35 40 45
 Asp

<210> 242
 <211> 49
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MC3 GPCR TM2

64000

<400> 242

Met Tyr Phe Phe Leu Cys Ser Leu Ala Val Ala Asp Met Leu Val Ser
 1 5 10 15
 Val Ser Asn Ala Leu Glu Thr Ile Met Ile Ala Ile Val His Ser Asp
 20 25 30
 Asp Tyr Thr Phe Glu Asp Gln Phe Ile Gln His Met Asp Asn Ile Phe
 35 40 45
 Asp

<210> 243

<211> 48

<212> PRT

<213> Artificial Sequence

<220>

<223> MC4 GPCR TM2

<400> 243

Met Tyr Phe Phe Ile Cys Ser Leu Ala Val Ala Asp Met Leu Val Ser
 1 5 10 15
 Val Ser Asn Gly Ser Glu Thr Ile Ile Ile Thr Leu Leu Asn Ser Thr
 20 25 30
 Asp Thr Asp Ala Gln Ser Phe Thr Val Asn Ile Asp Asn Val Ile Asp
 35 40 45

<210> 244

<211> 49

<212> PRT

<213> Artificial Sequence

<220>

<223> MC5 GPCR TM2

<400> 244

Met Tyr Phe Phe Val Cys Ser Leu Ala Val Ala Asp Met Leu Val Ser
 1 5 10 15
 Met Ser Ser Ala Trp Glu Thr Ile Thr Ile Tyr Leu Leu Asn Asn Lys
 20 25 30
 His Leu Val Ile Ala Asp Ala Phe Val Arg His Ile Asp Asn Val Phe
 35 40 45
 Asp

<210> 245

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> melatonin GPCR TM2

<400> 245

Gly Asn Ile Phe Val Val Ser Leu Ala Val Ala Asp Leu Val Val Ala
 1 5 10 15
 Ile Tyr Pro Tyr Pro Leu Val Leu Met Ser Ile Phe Asn Asn Gly Trp
 20 25 30
 Asn Leu Gly Tyr Leu His Cys Gln Val Ser Gly
 35 40

64PPP

<210> 246
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> oxytocin GPCR TM2

<400> 246
 Leu Phe Phe Phe Met Lys His Leu Ser Ile Ala Asp Leu Val Val Ala
 1 5 10 15
 Val Phe Gln Val Leu Pro Gln Leu Leu Trp Asp Ile Thr Phe Arg Phe
 20 25 30
 Tyr Gly Pro Asp Leu Leu Cys Arg Leu Val Lys
 35 40

<210> 247
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> conopressinLs GPCR TM2

<400> 247
 Met Gln Trp Phe Ile Ala His Leu Ala Phe Ala Asp Ile Phe Val Gly
 1 5 10 15
 Phe Phe Asn Ile Leu Pro Gln Leu Ile Ser Asp Val Thr Ile Val Phe
 20 25 30
 His Gly Asp Asp Phe Thr Cys Arg Phe Ile Lys
 35 40

<210> 248
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> V1A GPCR TM2

<400> 248
 Met His Leu Phe Ile Arg His Leu Ser Leu Ala Asp Leu Ala Val Ala
 1 5 10 15
 Phe Phe Gln Val Leu Pro Gln Met Cys Trp Asp Ile Thr Tyr Arg Phe
 20 25 30
 Arg Gly Pro Asp Trp Leu Cys Arg Val Val Lys
 35 40

<210> 249
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> V1B GPCR TM2

64QQQ

<400> 249

Met	His	Leu	Phe	Val	Leu	His	Leu	Ala	Leu	Thr	Asp	Leu	Ala	Val	Ala
1				5					10					15	
Leu	Phe	Gln	Val	Leu	Pro	Gln	Leu	Leu	Trp	Asp	Ile	Thr	Tyr	Arg	Phe
			20					25					30		
Gln	Gly	Pro	Asp	Leu	Leu	Cys	Arg	Ala	Val	Lys					
		35					40								

<210> 250

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> V2 GPCR TM2

<400> 250

Ile	His	Val	Phe	Ile	Gly	His	Leu	Cys	Leu	Ala	Asp	Leu	Ala	Val	Ala
1				5					10					15	
Leu	Phe	Gln	Val	Leu	Pro	Gln	Leu	Ala	Trp	Lys	Ala	Thr	Asp	Arg	Phe
			20					25					30		
Arg	Gly	Pro	Asp	Ala	Leu	Cys	Arg	Ala	Val	Lys					
		35					40								

<210> 251

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> CCK_A GPCR TM2

<400> 251

Thr	Asn	Ile	Phe	Leu	Leu	Ser	Leu	Ala	Val	Ser	Asp	Leu	Met	Leu	Cys
1				5					10					15	
Leu	Phe	Cys	Met	Pro	Phe	Asn	Leu	Ile	Pro	Asn	Leu	Leu	Lys	Asp	Phe
			20					25					30		
Ile	Phe	Gly	Ser	Ala	Val	Cys	Lys	Thr	Thr	Thr					
		35					40								

<210> 252

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> CCK_B GPCR TM2

<400> 252

Thr	Asn	Ala	Phe	Leu	Leu	Ser	Leu	Ala	Val	Ser	Asp	Leu	Leu	Leu	Ala
1				5					10					15	
Val	Ala	Cys	Met	Pro	Phe	Thr	Leu	Leu	Pro	Asn	Leu	Met	Gly	Thr	Phe
			20					25					30		
Ile	Phe	Gly	Thr	Val	Ile	Cys	Lys	Ala	Val	Ser					
		35					40								

<210> 253

<211> 43

<212> PRT

64RRR

<213> Artificial Sequence

<220>

<223> NPY1 GPCR TM2

<400> 253

Thr	Asn	Ile	Leu	Ile	Val	Asn	Leu	Ser	Phe	Ser	Asp	Leu	Leu	Val	Ala
1				5					10					15	
Ile	Met	Cys	Leu	Pro	Phe	Thr	Phe	Val	Tyr	Thr	Leu	Met	Asp	His	Trp
			20					25					30		
Val	Phe	Gly	Glu	Ala	Met	Cys	Lys	Leu	Asn	Pro					
		35					40								

<210> 254

<211> 45

<212> PRT

<213> Artificial Sequence

<220>

<223> NTR GPCR TM2

<400> 254

Val	His	Tyr	His	Leu	Gly	Ser	Leu	Ala	Leu	Ser	Asp	Leu	Leu	Thr	Leu
1				5					10					15	
Leu	Leu	Ala	Met	Pro	Val	Glu	Leu	Tyr	Asn	Phe	Ile	Trp	Val	His	His
			20					25					30		
Pro	Trp	Ala	Phe	Gly	Asp	Ala	Gly	Cys	Arg	Gly	Tyr	Tyr			
		35					40					45			

<210> 255

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> NK1 GPCR TM2

<400> 255

Thr	Asn	Tyr	Phe	Leu	Val	Asn	Leu	Ala	Phe	Ala	Glu	Ala	Ser	Met	Ala
1				5					10					15	
Ala	Phe	Asn	Thr	Val	Val	Asn	Phe	Thr	Tyr	Ala	Val	His	Asn	Glu	Trp
			20					25					30		
Tyr	Tyr	Gly	Leu	Phe	Tyr	Cys	Lys	Phe	His	Asn					
		35					40								

<210> 256

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> NK2 GPCR TM2

<400> 256

Thr	Asn	Tyr	Phe	Ile	Val	Asn	Leu	Ala	Leu	Ala	Asp	Leu	Cys	Met	Ala
1				5					10					15	
Ala	Phe	Asn	Ala	Ala	Phe	Asn	Phe	Val	Tyr	Ala	Ser	His	Asn	Ile	Trp
			20					25					30		

64SSS

Tyr Phe Gly Arg Ala Phe Cys Tyr Phe Gln Asn
 35 40

<210> 257
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> NK3 GPCR TM2

<400> 257
 Thr Asn Tyr Phe Leu Val Asn Leu Ala Phe Ser Asp Ala Ser Met Ala
 1 5 10 15
 Ala Phe Asn Thr Leu Val Asn Phe Ile Tyr Ala Leu His Ser Glu Trp
 20 25 30
 Tyr Phe Gly Ala Asn Tyr Cys Arg Phe Gln Asn
 35 40

<210> 258
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> blueops GPCR TM2

<400> 258
 Leu Asn Tyr Ile Leu Val Asn Val Ser Phe Gly Gly Phe Leu Leu Cys
 1 5 10 15
 Ile Phe Ser Val Phe Pro Val Phe Val Ala Ser Cys Asn Gly Tyr Phe
 20 25 30
 Val Phe Gly Arg His Val Cys Ala Leu Glu Gly
 35 40

<210> 259
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> greenops GPCR TM2

<400> 259
 Leu Asn Trp Ile Leu Val Asn Leu Ala Val Ala Asp Leu Ala Glu Thr
 1 5 10 15
 Val Ile Ala Ser Thr Ile Ser Val Val Asn Gln Val Tyr Gly Tyr Phe
 20 25 30
 Val Leu Gly His Pro Met Cys Val Leu Glu Gly
 35 40

<210> 260
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> redops GPCR TM2

64TTT

<400> 260

Leu	Asn	Trp	Ile	Leu	Val	Asn	Leu	Ala	Val	Ala	Asp	Leu	Ala	Glu	Thr
1				5					10					15	
Val	Ile	Ala	Ser	Thr	Ile	Ser	Ile	Val	Asn	Gln	Val	Ser	Gly	Tyr	Phe
			20					25					30		
Val	Leu	Gly	His	Pro	Met	Cys	Val	Leu	Glu	Gly					
		35					40								

<210> 261

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> rhodopsin GPCR TM2

<400> 261

Leu	Asn	Tyr	Ile	Leu	Leu	Asn	Leu	Ala	Val	Ala	Asp	Leu	Phe	Met	Val
1				5					10					15	
Leu	Gly	Gly	Phe	Thr	Ser	Thr	Leu	Tyr	Thr	Ser	Leu	His	Gly	Tyr	Phe
			20					25					30		
Val	Phe	Gly	Pro	Thr	Gly	Cys	Asn	Leu	Glu	Gly					
		35					40								

<210> 262

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> violetopsGg GPCR TM2

<400> 262

Leu	Asn	Tyr	Ile	Leu	Val	Asn	Ile	Ser	Ala	Ser	Gly	Phe	Val	Ser	Cys
1				5					10					15	
Val	Leu	Ser	Val	Phe	Val	Val	Phe	Val	Ala	Ser	Ala	Arg	Gly	Tyr	Phe
			20					25					30		
Val	Phe	Gly	Lys	Arg	Val	Cys	Glu	Leu	Glu	Ala					
		35					40								

<210> 263

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> opsin_crab GPCR TM2

<400> 263

Thr	Asn	Leu	Leu	Val	Val	Asn	Leu	Ala	Phe	Ser	Asp	Phe	Cys	Met	Met
1				5					10					15	
Ala	Phe	Met	Met	Pro	Thr	Met	Thr	Ser	Asn	Cys	Phe	Ala	Glu	Thr	Trp
			20					25					30		
Ile	Leu	Gly	Pro	Phe	Met	Cys	Glu	Val	Tyr	Gly					
		35					40								

<210> 264

<211> 48

<212> PRT

64UUU

<213> Artificial Sequence

<220>

<223> ET_Aprec GPCR TM2

<400> 264

Pro	Asn	Ala	Leu	Ile	Ala	Ser	Leu	Ala	Leu	Gly	Asp	Leu	Ile	Tyr	Val
1				5					10					15	
Val	Ile	Asp	Leu	Pro	Ile	Asn	Val	Phe	Lys	Leu	Leu	Ala	Gly	Arg	Trp
			20					25					30		
Pro	Phe	Asp	His	Asn	Asp	Phe	Gly	Val	Phe	Leu	Cys	Lys	Leu	Phe	Pro
		35					40					45			

<210> 265

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> ET_Bprec GPCR TM2

<400> 265

Pro	Asn	Ile	Leu	Ile	Ala	Ser	Leu	Ala	Leu	Gly	Asp	Leu	Leu	His	Ile
1				5					10					15	
Val	Ile	Asp	Ile	Pro	Ile	Asn	Val	Tyr	Lys	Leu	Leu	Ala	Glu	Asp	Trp
			20					25					30		
Pro	Phe	Gly	Ala	Glu	Met	Cys	Lys	Leu	Val	Pro					
		35					40								

<210> 266

<211> 39

<212> PRT

<213> Artificial Sequence

<220>

<223> ET_Cfrog GPCR TM2

<400> 266

Pro	Asn	Val	Leu	Ile	Ala	Ser	Leu	Ala	Leu	Gly	Asp	Leu	Phe	Tyr	Ile
1				5					10					15	
Leu	Ile	Ala	Ile	Pro	Ile	Ile	Ser	Ile	Ser	Phe	Trp	Leu	Ser	Thr	Gly
			20					25					30		
His	Ser	Glu	Tyr	Ile	Tyr	Gln									
		35													

<210> 267

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> galatin GPCR TM2

<400> 267

Thr	Asn	Leu	Phe	Ile	Leu	Asn	Leu	Ser	Ile	Ala	Asp	Leu	Ala	Tyr	Leu
1				5					10					15	
Leu	Phe	Cys	Ile	Pro	Phe	Gln	Ala	Thr	Val	Tyr	Ala	Leu	Pro	Thr	Trp
			20					25					30		

64VVV

Val Leu Gly Ala Phe Ile Cys Lys Phe Ile His
 35 40

<210> 268
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> NMBGPCR TM2

<400> 268
 Pro Asn Ile Phe Ile Ser Asn Leu Ala Ala Gly Asp Leu Leu Leu Leu
 1 5 10 15
 Leu Thr Cys Val Pro Val Asp Ala Ser Arg Tyr Phe Phe Asp Glu Trp
 20 25 30
 Met Phe Gly Lys Val Gly Cys Lys Leu Ile Pro
 35 40

<210> 269
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> GRP GPCR TM2

<400> 269
 Pro Asn Leu Phe Ile Ser Ser Leu Ala Leu Gly Asp Leu Leu Leu Leu
 1 5 10 15
 Ile Thr Cys Ala Pro Val Asp Ala Ser Arg Tyr Leu Ala Asp Arg Trp
 20 25 30
 Leu Phe Gly Arg Ile Gly Cys Lys Leu Ile Pro
 35 40

<210> 270
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> BRS3 GPCR TM2

<400> 270
 Pro Asn Ile Phe Ile Thr Ser Leu Ala Phe Gly Asp Leu Leu Leu Leu
 1 5 10 15
 Leu Thr Cys Val Pro Val Asp Ala Thr His Tyr Leu Ala Glu Gly Trp
 20 25 30
 Leu Phe Gly Arg Ile Gly Cys Lys Val Leu Ser
 35 40

<210> 271
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> deltaOP GPCR TM2

64WWW

<400> 271

Thr	Asn	Ile	Tyr	Ile	Phe	Asn	Leu	Ala	Leu	Ala	Asp	Ala	Leu	Ala	Thr
1				5					10					15	
Ser	Thr	Leu	Pro	Phe	Gln	Ser	Ala	Lys	Tyr	Leu	Met	Glu	Thr	Trp	Pro
			20					25					30		
Phe	Gly	Glu	Leu	Leu	Cys	Lys	Ala	Val	Leu						
		35					40								

<210> 272

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> kappaOP GPCR TM2

<400> 272

Thr	Asn	Ile	Tyr	Ile	Phe	Asn	Leu	Ala	Leu	Ala	Asp	Ala	Leu	Val	Thr
1				5					10					15	
Thr	Thr	Met	Pro	Phe	Gln	Ser	Thr	Val	Tyr	Leu	Met	Asn	Ser	Trp	Pro
			20					25					30		
Phe	Gly	Asp	Val	Leu	Cys	Lys	Ile	Val	Ile						
		35					40								

<210> 273

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> muOP GPCR TM2

<400> 273

Thr	Asn	Ile	Tyr	Ile	Phe	Asn	Leu	Ala	Leu	Ala	Asp	Ala	Leu	Ala	Thr
1				5					10					15	
Ser	Thr	Leu	Pro	Phe	Gln	Ser	Val	Asn	Tyr	Leu	Met	Gly	Thr	Trp	Pro
			20					25					30		
Phe	Gly	Thr	Ile	Leu	Cys	Lys	Ile	Val	Ile						
		35					40								

<210> 274

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> OPRX GPCR TM2

<400> 274

Thr	Asn	Ile	Tyr	Ile	Phe	Asn	Leu	Ala	Leu	Ala	Asp	Thr	Leu	Val	Leu
1				5					10					15	
Leu	Thr	Leu	Pro	Phe	Gln	Gly	Thr	Asp	Ile	Leu	Leu	Gly	Phe	Trp	Pro
			20					25					30		
Phe	Gly	Asn	Ala	Leu	Cys	Lys	Thr	Val	Ile						
		35					40								

<210> 275

<211> 42

<212> PRT

64XXX

<213> Artificial Sequence

<220>

<223> CB1 GPCR TM2

<400> 275

Ser	Tyr	His	Phe	Ile	Gly	Ser	Leu	Ala	Val	Ala	Asp	Leu	Leu	Gly	Ser
1				5					10					15	
Val	Ile	Phe	Val	Tyr	Ser	Phe	Ile	Asp	Phe	His	Val	Phe	His	Arg	Lys
			20					25					30		
Asp	Ser	Arg	Asn	Val	Phe	Leu	Phe	Lys	Leu						
		35					40								

<210> 276

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> CB2 GPCR TM2

<400> 276

Ser	Tyr	Leu	Phe	Ile	Gly	Ser	Leu	Ala	Gly	Ala	Asp	Phe	Leu	Ala	Ser
1				5					10					15	
Val	Val	Phe	Ala	Cys	Ser	Phe	Val	Asn	Phe	His	Val	Phe	His	Gly	Val
			20					25					30		
Asp	Ser	Lys	Ala	Val	Phe	Leu	Leu	Lys	Ile						
		35					40								

<210> 277

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> SSTR1 GPCR TM2

<400> 277

Thr	Asn	Ile	Tyr	Ile	Leu	Asn	Leu	Ala	Ile	Ala	Asp	Glu	Leu	Leu	Met
1				5					10					15	
Leu	Ser	Val	Pro	Phe	Leu	Val	Thr	Ser	Thr	Leu	Leu	Arg	His	Trp	Pro
			20					25					30		
Phe	Gly	Ala	Leu	Leu	Cys	Arg	Leu	Val	Leu						
		35					40								

<210> 278

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> SSTR2 GPCR TM2

<400> 278

Thr	Asn	Ile	Tyr	Ile	Leu	Asn	Leu	Ala	Ile	Ala	Asp	Glu	Leu	Phe	Met
1				5					10					15	
Leu	Gly	Leu	Pro	Phe	Leu	Ala	Met	Gln	Val	Ala	Leu	Val	His	Trp	Pro
			20					25					30		

64YYY

Phe Gly Lys Ala Ile Cys Arg Val Val Met
 35 40

<210> 279
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> SSTR3 GPCR TM2

<400> 279
 Thr Asn Val Tyr Ile Leu Asn Leu Ala Leu Ala Asp Glu Leu Phe Met
 1 5 10 15
 Leu Gly Leu Pro Phe Leu Ala Ala Gln Asn Ala Leu Ser Tyr Trp Pro
 20 25 30
 Phe Gly Ser Leu Met Cys Arg Leu Val Met
 35 40

<210> 280
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> SSTR4 GPCR TM2

<400> 280
 Thr Asn Ile Tyr Leu Leu Asn Leu Ala Val Ala Asp Glu Leu Phe Met
 1 5 10 15
 Leu Ser Val Pro Phe Val Ala Ser Ser Ala Ala Leu Arg His Trp Pro
 20 25 30
 Phe Gly Ser Val Leu Cys Arg Ala Val Leu
 35 40

<210> 281
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> SSTR5 GPCR TM2

<400> 281
 Thr Asn Ile Tyr Ile Leu Asn Leu Ala Val Ala Asp Val Leu Tyr Met
 1 5 10 15
 Leu Gly Leu Pro Phe Leu Ala Thr Gln Asn Ala Ala Ser Phe Trp Pro
 20 25 30
 Phe Gly Pro Val Leu Cys Arg Leu Val Met
 35 40

<210> 282
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> IL8A and IL8B GPCR TM2

64ZZZ

<400> 282

Thr	Asp	Val	Tyr	Leu	Leu	Asn	Leu	Ala	Leu	Ala	Asp	Leu	Leu	Phe	Ala
1				5					10					15	
Leu	Thr	Leu	Pro	Ile	Trp	Ala	Ala	Ser	Lys	Val	Asn	Gly	Trp	Ile	Phe
			20					25					30		
Gly	Thr	Phe	Leu	Cys	Lys	Val	Val	Ser							
		35					40								

<210> 283

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> AT1a GPCR TM2

<400> 283

Ala	Ser	Val	Phe	Leu	Leu	Asn	Leu	Ala	Leu	Ala	Asp	Leu	Cys	Phe	Leu
1				5					10					15	
Leu	Thr	Leu	Pro	Leu	Trp	Ala	Val	Tyr	Thr	Ala	Met	Glu	Tyr	Arg	Trp
			20					25					30		
Pro	Phe	Gly	Asn	Tyr	Leu	Cys	Lys	Ile	Ala	Ser					
		35					40								

<210> 284

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> AT1brat GPCR TM2

<400> 284

Ala	Ser	Val	Phe	Leu	Leu	Asn	Leu	Ala	Leu	Ala	Asp	Leu	Cys	Phe	Leu
1				5					10					15	
Leu	Thr	Leu	Pro	Leu	Trp	Ala	Val	Tyr	Thr	Ala	Met	Glu	Tyr	Arg	Trp
			20					25					30		
Pro	Phe	Gly	Asn	His	Leu	Cys	Lys	Ile	Ala	Ser					
		35					40								

<210> 285

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> AT2 GPCR TM2

<400> 285

Ser	Ser	Ile	Tyr	Ile	Phe	Asn	Leu	Ala	Val	Ala	Asp	Leu	Leu	Leu	Leu
1				5					10					15	
Ala	Thr	Leu	Pro	Leu	Trp	Ala	Thr	Tyr	Tyr	Ser	Tyr	Arg	Tyr	Asp	Trp
			20					25					30		
Leu	Phe	Gly	Pro	Val	Met	Cys	Lys	Val	Phe	Gly					
		35					40								

<210> 286

<211> 43

<212> PRT

64AAAA

<213> Artificial Sequence

<220>

<223> BK1 GPCR TM2

<400> 286

Ala	Glu	Ile	Tyr	Leu	Ala	Asn	Leu	Ala	Ala	Ser	Asp	Leu	Val	Phe	Val
1				5				10						15	
Leu	Gly	Leu	Pro	Phe	Trp	Ala	Glu	Asn	Ile	Trp	Asn	Gln	Phe	Asn	Trp
			20				25						30		
Pro	Phe	Gly	Ala	Leu	Leu	Cys	Arg	Val	Ile	Asn					
		35					40								

<210> 287

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> BK2 GPCR TM2

<400> 287

Ala	Glu	Ile	Tyr	Leu	Gly	Asn	Leu	Ala	Ala	Ala	Asp	Leu	Ile	Leu	Ala
1				5				10						15	
Cys	Gly	Leu	Pro	Phe	Trp	Ala	Ile	Thr	Ile	Ser	Asn	Asn	Phe	Asp	Trp
			20				25						30		
Leu	Phe	Gly	Glu	Thr	Leu	Cys	Arg	Val	Val	Asn					
		35					40								

<210> 288

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> P2Y7 GPCR TM2

<400> 288

Thr	Ala	Leu	Met	Val	Leu	Asn	Leu	Ala	Leu	Ala	Asp	Leu	Ala	Val	Leu
1				5				10						15	
Leu	Thr	Ala	Pro	Phe	Phe	Leu	His	Phe	Leu	Ala	Gln	Gly	Thr	Trp	Ser
			20				25						30		
Phe	Gly	Leu	Ala	Gly	Cys	Arg	Leu	Cys	His						
		35					40								

<210> 289

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> P2Y6 GPCR TM2

<400> 289

Ser	Ala	Val	Tyr	Thr	Leu	Asn	Leu	Ala	Leu	Ala	Asp	Leu	Leu	Tyr	Ala
1				5				10						15	
Cys	Ser	Leu	Pro	Leu	Leu	Ile	Tyr	Asn	Tyr	Ala	Arg	Gly	Asp	His	Trp
			20				25						30		

64BBBB

Pro Phe Gly Asp Leu Ala Cys Arg Leu Val Arg
 35 40

<210> 290
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P2Y5 GPCR TM2

<400> 290
 Thr Thr Thr Tyr Met Leu Asn Leu Ala Ile Ser Asp Leu Leu Phe Val
 1 5 10 15
 Phe Thr Leu Pro Phe Arg Ile Tyr Tyr Phe Val Val Arg Asn Trp Pro
 20 25 30
 Phe Gly Asp Val Leu Cys Lys Ile Ser Val
 35 40

<210> 291
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P2Y4 GPCR TM2

<400> 291
 Thr Ala Thr Tyr Met Phe His Leu Ala Leu Ser Asp Thr Leu Tyr Val
 1 5 10 15
 Val Ser Leu Pro Thr Leu Ile Tyr Tyr Tyr Ala Ala His Asn His Trp
 20 25 30
 Pro Phe Gly Thr Glu Ile Cys Lys Phe Val Arg
 35 40

<210> 292
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P2Y3chick GPCR TM2

<400> 292
 Thr Thr Ile Tyr Met Leu Asn Leu Ala Met Ala Asp Leu Leu Tyr Val
 1 5 10 15
 Cys Ser Leu Pro Leu Leu Ile Tyr Asn Tyr Thr Gln Lys Asp Tyr Trp
 20 25 30
 Pro Phe Gly Asp Phe Thr Cys Lys Phe Val Arg
 35 40

<210> 293
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P2Y2 GPCR TM2

64CCCC

<400> 293

Ser	Thr	Thr	Tyr	Met	Phe	His	Leu	Ala	Val	Ser	Asp	Ala	Leu	Tyr	Ala
1				5					10					15	
Ala	Ser	Leu	Pro	Leu	Leu	Val	Tyr	Tyr	Tyr	Ala	Arg	Gly	Asp	His	Trp
			20					25					30		
Pro	Phe	Ser	Thr	Val	Leu	Cys	Lys	Leu	Val	Arg					
		35					40								

<210> 294

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> P2Y1 GPCR TM2

<400> 294

Ile	Ser	Val	Tyr	Met	Phe	Asn	Leu	Ala	Leu	Ala	Asp	Phe	Leu	Tyr	Val
1				5					10					15	
Leu	Thr	Leu	Pro	Ala	Leu	Ile	Phe	Tyr	Tyr	Phe	Asn	Lys	Thr	Asp	Trp
			20					25					30		
Ile	Phe	Gly	Asp	Ala	Met	Cys	Lys	Leu	Gln	Arg					
		35					40								

<210> 295

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> THRprec GPCR TM2

<400> 295

Ala	Val	Val	Tyr	Met	Leu	His	Leu	Ala	Thr	Ala	Asp	Val	Leu	Phe	Val
1				5					10					15	
Ser	Val	Leu	Pro	Phe	Lys	Ile	Ser	Tyr	Tyr	Phe	Ser	Gly	Ser	Asp	Trp
			20					25					30		
Gln	Phe	Gly	Ser	Glu	Leu	Cys	Arg	Phe	Val	Thr					
		35					40								

<210> 296

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> C5a GPCR TM2

<400> 296

Asn	Ala	Ile	Trp	Phe	Leu	Asn	Leu	Ala	Val	Ala	Asp	Phe	Leu	Ser	Cys
1				5					10					15	
Leu	Ala	Leu	Pro	Ile	Leu	Phe	Thr	Ser	Ile	Val	Gln	His	His	Trp	Pro
			20					25					30		
Phe	Gly	Gly	Ala	Ala	Cys	Ser	Ile	Leu	Pro						
		35					40								

<210> 297

<211> 39

<212> PRT

64DDDD

<213> Artificial Sequence

<220>

<223> GP01mouse AND R334rat GPCR TM2

<400> 297

Met	Phe	Leu	Leu	Ile	Gly	Ser	Leu	Ala	Leu	Ala	Asp	Leu	Leu	Ala	Gly
1				5					10					15	
Leu	Gly	Leu	Ile	Ile	Asn	Phe	Val	Phe	Ala	Tyr	Leu	Leu	Gln	Ser	Glu
			20					25					30		
Ala	Thr	Lys	Leu	Val	Thr	Ile									
			35												

<210> 298

<211> 39

<212> PRT

<213> Artificial Sequence

<220>

<223> GP21mouse GPCR TM2

<400> 298

Met	Phe	Leu	Leu	Val	Gly	Ser	Leu	Ala	Val	Ala	Asp	Leu	Leu	Ala	Gly
1				5					10					15	
Leu	Gly	Leu	Val	Leu	His	Phe	Ala	Ala	Asp	Phe	Cys	Ile	Gly	Ser	Pro
			20					25					30		
Glu	Met	Ser	Leu	Met	Leu	Val									
			35												

<210> 299

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> GCRCmouse GPCR TM2

<400> 299

Thr	Ser	Leu	Phe	Ile	Val	Asn	Leu	Ala	Val	Ala	Asp	Ile	Met	Ile	Thr
1				5					10					15	
Leu	Leu	Asn	Thr	Pro	Phe	Thr	Leu	Val	Arg	Phe	Val	Asn	Ser	Thr	Trp
			20					25					30		
Val	Phe	Gly	Lys	Gly	Met	Cys	His	Val	Ser	Arg					
		35					40								

<210> 300

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> TXKR GPCR TM2

<400> 300

Thr	Asn	Ser	Phe	Leu	Val	Asn	Leu	Ala	Phe	Ala	Asp	Ala	Ala	Met	Ala
1				5					10					15	
Ala	Leu	Asn	Ala	Leu	Val	Asn	Phe	Ile	Tyr	Ala	Leu	His	Gly	Glu	Trp
			20					25					30		

64EEEE

Tyr Phe Gly Ala Asn Tyr Cys Arg Phe Gln Asn
 35 40

<210> 301
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> G10Drat GPCR TM2

<400> 301
 Leu Asn Leu Tyr Ile Leu Asn Met Ala Val Ala Asp Leu Gly Ile Ile
 1 5 10 15
 Leu Ser Leu Pro Val Trp Met Leu Glu Val Met Leu Glu Tyr Thr Trp
 20 25 30
 Leu Trp Gly Ser Phe Ser Cys Arg Phe Ile His
 35 40

<210> 302
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> RDC1 GPCR TM2

<400> 302
 His Cys Tyr Ile Leu Asn Leu Ala Ile Ala Asp Leu Trp Val Val Leu
 1 5 10 15
 Thr Ile Pro Val Trp Val Val Ser Leu Val Gln His Asn Gln Trp Pro
 20 25 30
 Met Gly Glu Leu Thr Cys Lys Val Thr His
 35 40

<210> 303
 <211> 39
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> BLR1 GPCR TM2

<400> 303
 Thr Phe Leu Phe His Leu Ala Val Ala Asp Leu Leu Leu Val Phe Ile
 1 5 10 15
 Leu Pro Phe Ala Val Ala Glu Gly Ser Val Gly Trp Val Leu Gly Thr
 20 25 30
 Phe Leu Cys Lys Thr Val Ile
 35

<210> 304
 <211> 36
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CL5 and LCR1GPCR TM2

64FFFF

<400> 304

Leu	His	Leu	Ser	Val	Ala	Asp	Leu	Leu	Phe	Val	Ile	Thr	Leu	Pro	Phe
1				5					10					15	
Trp	Ala	Val	Asp	Ala	Val	Ala	Asn	Trp	Tyr	Phe	Gly	Asn	Phe	Leu	Cys
			20					25					30		
Lys	Ala	Val	His												
			35												

<210> 305

<211> 37

<212> PRT

<213> Artificial Sequence

<220>

<223> EBI1 GPCR TM2

<400> 305

Leu	Leu	Asn	Leu	Ala	Val	Ala	Asp	Ile	Leu	Phe	Leu	Leu	Thr	Leu	Pro
1				5					10					15	
Phe	Trp	Ala	Tyr	Ser	Ala	Ala	Lys	Ser	Trp	Val	Phe	Gly	Val	His	Phe
			20					25					30		
Cys	Lys	Leu	Ile	Phe											
			35												

<210> 306

<211> 37

<212> PRT

<213> Artificial Sequence

<220>

<223> RBS1rat GPCR TM2

<400> 306

Leu	Leu	Asn	Leu	Ala	Leu	Ser	Asp	Leu	Leu	Phe	Val	Ala	Thr	Leu	Pro
1				5					10					15	
Phe	Trp	Thr	His	Tyr	Leu	Ile	Ser	His	Glu	Gly	Leu	His	Asn	Ala	Met
			20					25					30		
Cys	Lys	Leu	Thr	Thr											
			35												

<210> 307

<211> 39

<212> PRT

<213> Artificial Sequence

<220>

<223> EBI2 GPCR TM2

<400> 307

Ser	Thr	Asn	Leu	Val	Ile	Ser	Asp	Ile	Leu	Phe	Thr	Thr	Ala	Leu	Pro
1				5					10					15	
Thr	Arg	Ile	Ala	Tyr	Tyr	Ala	Met	Gly	Phe	Asp	Trp	Arg	Ile	Gly	Asp
			20					25					30		
Ala	Leu	Cys	Arg	Ile	Thr	Ala									
			35												

<210> 308

<211> 38

<212> PRT

64GGGG

<213> Artificial Sequence

<220>

<223> GCRTchick GPCR TM2

<400> 308

Met	Leu	Asn	Leu	Ala	Ile	Ser	Asp	Leu	Leu	Phe	Val	Phe	Thr	Leu	Pro
1				5				10						15	
Phe	Arg	Ile	Tyr	Tyr	Phe	Val	Val	Arg	Asn	Trp	Pro	Phe	Gly	Asp	Val
			20					25					30		
Leu	Cys	Lys	Ile	Ser	Val										
			35												

<210> 309

<211> 41

<212> PRT

<213> Artificial Sequence

<220>

<223> APJ GPCR TM2

<400> 309

Ile	Phe	Ile	Ala	Ser	Leu	Ala	Val	Ala	Asp	Leu	Thr	Phe	Val	Val	Thr
1				5				10						15	
Leu	Pro	Leu	Trp	Ala	Thr	Tyr	Thr	Tyr	Arg	Asp	Tyr	Asp	Trp	Pro	Phe
			20					25					30		
Gly	Thr	Phe	Phe	Cys	Lys	Leu	Ser	Ser							
			35				40								

<210> 310

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> RTArat GPCR TM2

<400> 310

Phe	Ser	Ile	Tyr	Phe	Leu	His	Leu	Ala	Ser	Ala	Asp	Gly	Ile	Tyr	Leu
1				5				10						15	
Phe	Ser	Lys	Ala	Val	Ile	Ala	Leu	Leu	Asn	Met	Gly	Thr	Phe	Leu	Gly
			20					25					30		
Ser	Phe	Pro	Asp	Tyr	Val	Arg	Arg	Val	Ser	Arg					
			35				40								

<210> 311

<211> 44

<212> PRT

<213> Artificial Sequence

<220>

<223> UHRrat GPCR TM2

<400> 311

Thr	Asn	Phe	Leu	Ile	Gly	Asn	Leu	Ala	Leu	Ser	Asp	Val	Leu	Met	Cys
1				5				10						15	
Ala	Ala	Cys	Val	Pro	Leu	Thr	Leu	Ala	Tyr	Ala	Phe	Glu	Pro	Arg	Gly
			20					25					30		

64HHHH

Trp Val Phe Gly Gly Gly Leu Cys His Leu Val Phe
 35 40

<210> 312
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> FMRL1 GPCR TM2

<400> 312
 Asn Thr Ile Cys Tyr Leu Asn Leu Ala Leu Ala Asp Phe Ser Phe Ser
 1 5 10 15
 Ala Ile Leu Pro Phe Arg Met Val Ser Val Ala Met Arg Glu Lys Trp
 20 25 30
 Pro Phe Ala Ser Phe Leu Cys Lys Leu Val His
 35 40

<210> 313
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> FMRL2 GPCR TM2

<400> 313
 Thr Thr Ile Cys Tyr Leu Asn Leu Ala Leu Ala Asp Phe Ser Phe Thr
 1 5 10 15
 Ala Thr Leu Pro Phe Leu Ile Val Ser Met Ala Met Gly Glu Lys Trp
 20 25 30
 Pro Phe Gly Trp Phe Leu Cys Lys Leu Ile His
 35 40

<210> 314
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> fMLP GPCR TM2

<400> 314
 Thr Thr Ile Ser Tyr Leu Asn Leu Ala Val Ala Asp Phe Cys Phe Thr
 1 5 10 15
 Ser Thr Leu Pro Phe Phe Met Val Arg Lys Ala Met Gly Gly His Trp
 20 25 30
 Pro Phe Gly Trp Phe Leu Cys Lys Phe Leu Phe
 35 40

<210> 315
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLF1catfish GPCR TM2

64IIII

<400> 315

Lys	Tyr	Ile	Thr	Val	Phe	Asn	Leu	Ala	Leu	Ser	Asp	Leu	Gly	Gly	Ser
1				5					10					15	
Ser	Ala	Leu	Ile	Pro	Lys	Leu	Ile	Asp	Thr	Phe	Leu	Phe	Glu	Asn	Gln
			20					25					30		
Val	Ile	Ser	Tyr	Glu	Ala	Cys	Leu	Ala	Asn	Met					
		35					40								

<210> 316

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF3catfish GPCR TM2

<400> 316

Lys	Tyr	Ile	Ala	Val	Phe	Asn	Leu	Ala	Leu	Ser	Asp	Leu	Cys	Gly	Ser
1				5					10					15	
Ser	Ala	Leu	Ile	Pro	Lys	Leu	Leu	Asp	Met	Leu	Leu	Phe	Glu	Asn	Gln
			20					25					30		
Ser	Ile	Ser	Tyr	Glu	Ala	Cys	Leu	Ser	Asn	Met					
		35					40								

<210> 317

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF8catfish GPCR TM2

<400> 317

Met	Cys	Ile	Leu	Ile	Gly	Leu	Met	Ala	Val	Val	Asp	Leu	Ser	Met	Pro
1				5					10					15	
Ile	Phe	Cys	Val	Pro	Asn	Met	Leu	Leu	Ser	Phe	Leu	Phe	Asn	Trp	Lys
			20					25					30		
Gly	Ile	Ser	Leu	Val	Gly	Cys	Leu	Val	Gln	Met					
		35					40								

<210> 318

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF32Acatfish GPCR TM2

<400> 318

Lys	Tyr	Met	Gly	Ile	Phe	Asn	Leu	Ala	Leu	Ser	Asp	Phe	Gly	Glu	Thr
1				5					10					15	
Asn	Val	Leu	Ile	Pro	Ser	Leu	Val	Lys	Thr	Leu	Phe	Phe	Asp	Ser	Gln
			20					25					30		
Tyr	Ile	Ser	Tyr	Asp	Ala	Cys	Leu	Ala	Asn	Met					
		35					40								

<210> 319

<211> 43

<212> PRT

64JJJJ

<213> Artificial Sequence

<220>

<223> OLF32Bcatfish and OLF32Dcatfish GPCR TM2

<400> 319

Lys	Tyr	Met	Gly	Ile	Phe	Asn	Leu	Ala	Leu	Ser	Asp	Phe	Gly	Glu	Thr
1				5					10					15	
Asn	Ala	Leu	Ile	Pro	Ser	Leu	Val	Lys	Thr	Leu	Phe	Phe	Asp	Ser	Gln
			20					25					30		
Tyr	Ile	Ser	Tyr	Asp	Ala	Cys	Leu	Ala	Asn	Met					
		35					40								

<210> 320

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF32Ccatfish GPCR TM2

<400> 320

Lys	Tyr	Met	Gly	Ile	Phe	Asn	Leu	Ala	Leu	Ser	Asp	Ile	Gly	Glu	Thr
1				5					10					15	
Asn	Ala	Leu	Ile	Pro	Ser	Leu	Val	Lys	Thr	Leu	Phe	Phe	Asp	Ser	Gln
			20					25					30		
Tyr	Ile	Ser	Tyr	Asp	Ala	Cys	Leu	Thr	Asn	Met					
		35					40								

<210> 321

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF47catfish GPCR TM2

<400> 321

Lys	Phe	Leu	Ala	Val	Phe	Asn	Leu	Ala	Val	Val	Asp	Ile	Ser	Ile	Asn
1				5					10					15	
Ser	Val	Ile	Ile	Pro	Gln	Met	Val	Pro	Val	Phe	Val	Phe	Asn	Leu	Asn
			20					25					30		
His	Ile	Ser	Phe	Glu	Ser	Cys	Phe	Ser	Gln	Met					
		35					40								

<210> 322

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF202catfish GPCR TM2

<400> 322

Met	Tyr	Tyr	Ile	Met	Leu	Asn	Leu	Ala	Ala	Ser	Asp	Val	Leu	Phe	Ser
1				5					10					15	
Thr	Thr	Thr	Leu	Pro	Lys	Ile	Ile	Ala	Arg	Tyr	Trp	Phe	Gly	Asp	Gly
			20					25					30		

64KKKK

Ser Ile Ser Phe Val Gly Cys Phe Ile Gln Met
 35 40

<210> 323
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFCOR1chicken, OLFCOR3chicken AND OLFCOR4chicken
 GPCR TM2

<400> 323
 Met Tyr Ile Phe Leu Gln Asn Leu Ser Phe Thr Asp Ala Ala Tyr Ser
 1 5 10 15
 Thr Val Ile Thr Pro Lys Met Leu Ala Thr Phe Leu Glu Glu Arg Lys
 20 25 30
 Thr Ile Ser Tyr Val Gly Cys Ile Leu Gln Tyr
 35 40

<210> 324
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFCOR2chicken and OLFCOR5chicken GPCR TM2

<400> 324
 Met Tyr Ile Phe Leu Gln Asn Leu Ser Phe Thr Asp Ala Ala Tyr Ser
 1 5 10 15
 Thr Val Ile Thr Pro Lys Met Leu Ala Thr Phe Leu Glu Glu Arg Arg
 20 25 30
 Thr Ile Ser Tyr Val Gly Cys Ile Leu Gln Tyr
 35 40

<210> 325
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFCOR6chicken GPCR TM2

<400> 325
 Met Tyr Ile Phe Leu Gln Asn Leu Ser Phe Thr Asp Ala Val Tyr Ser
 1 5 10 15
 Thr Val Ile Thr Pro Lys Met Leu Ala Thr Phe Leu Glu Glu Thr Lys
 20 25 30
 Thr Ile Ser Tyr Val Gly Cys Ile Leu Gln Tyr
 35 40

<210> 326
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFdog GPCR TM2

64LLLL

<400> 326

Met	Tyr	Leu	Phe	Leu	Ser	Asn	Leu	Ser	Phe	Ser	Asp	Leu	Cys	Phe	Ser
1				5					10					15	
Ser	Val	Thr	Met	Pro	Lys	Leu	Leu	Gln	Asn	Met	Gln	Ser	Gln	Val	Pro
			20					25					30		
Ser	Ile	Pro	Tyr	Ala	Gly	Cys	Leu	Thr	Gln	Met					
		35					40								

<210> 327

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF07E GPCR TM2

<400> 327

Val	Tyr	Phe	Phe	Leu	Ala	Asn	Leu	Ser	Phe	Thr	Asp	Leu	Phe	Phe	Val
1				5					10					15	
Thr	Asn	Thr	Ile	Pro	Lys	Met	Leu	Val	Asn	Leu	Gln	Ser	His	Asn	Lys
			20					25					30		
Ala	Ile	Ser	Tyr	Ala	Gly	Cys	Leu	Thr	Gln	Leu					
		35					40								

<210> 328

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF07I GPCR TM2

<400> 328

Met	Tyr	Leu	Phe	Leu	Ser	Asn	Leu	Ser	Phe	Ser	Asp	Leu	Cys	Phe	Ser
1				5					10					15	
Ser	Val	Thr	Ile	Pro	Lys	Leu	Leu	Gln	Asn	Met	Gln	Asn	Gln	Asp	Pro
			20					25					30		
Ser	Ile	Pro	Tyr	Ala	Asp	Cys	Leu	Thr	Gln	Met					
		35					40								

<210> 329

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLF07J GPCR TM2

<400> 329

Met	Tyr	Phe	Phe	Leu	Ser	Met	Leu	Ser	Thr	Ser	Glu	Thr	Val	Tyr	Thr
1				5					10					15	
Leu	Val	Ile	Leu	Pro	Arg	Met	Leu	Ser	Ser	Leu	Val	Gly	Met	Ser	Gln
			20					25					30		
Pro	Met	Ser	Leu	Ala	Gly	Cys	Ala	Thr	Gln	Met					
		35					40								

<210> 330

<211> 43

64MMMM

<212> PRT

<213> Artificial Sequence

<220>

<223> OLFOR3mouse GPCR TM2

<400> 330

Met	Tyr	Phe	Phe	Leu	Ser	Asn	Leu	Ser	Ser	Leu	Asp	Leu	Ala	Phe	Thr
1				5				10						15	
Thr	Ser	Ser	Val	Pro	Gln	Met	Leu	Lys	Asn	Leu	Trp	Gly	Pro	Asp	Lys
			20					25					30		
Thr	Ile	Ser	Tyr	Gly	Gly	Cys	Val	Thr	Gln	Leu					
		35					40								

<210> 331

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLFrat GPCR TM2

<400> 331

Met	Tyr	Tyr	Phe	Leu	Ser	Ser	Leu	Ser	Phe	Val	Asp	Leu	Cys	Tyr	Ser
1				5				10						15	
Thr	Val	Ile	Thr	Pro	Lys	Met	Leu	Val	Asn	Phe	Leu	Gly	Lys	Lys	Asn
			20					25					30		
Phe	Ile	Thr	Tyr	Ser	Glu	Cys	Met	Ala	Gln	Phe					
		35					40								

<210> 332

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLFF3rat GPCR TM2

<400> 332

Met	Tyr	Phe	Phe	Leu	Ser	Asn	Leu	Ser	Phe	Val	Asp	Ile	Cys	Phe	Ile
1				5				10						15	
Ser	Thr	Thr	Val	Pro	Lys	Met	Leu	Val	Asn	Ile	Gln	Thr	Gln	Asn	Asn
			20					25					30		
Val	Ile	Thr	Tyr	Ala	Gly	Cys	Ile	Thr	Gln	Ile					
		35					40								

<210> 333

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLFF5rat GPCR TM2

<400> 333

Met	Tyr	Phe	Phe	Leu	Ser	Asn	Leu	Ser	Phe	Val	Asp	Val	Cys	Phe	Ser
1				5				10						15	
Ser	Thr	Thr	Val	Pro	Lys	Val	Leu	Ala	Asn	His	Ile	Leu	Gly	Ser	Gln
			20					25					30		

64NNNN

Ala Ile Ser Phe Ser Gly Cys Leu Thr Gln Leu
 35 40

<210> 334
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFF6rat GPCR TM2

<400> 334
 Met Tyr Phe Phe Leu Cys Asn Leu Ser Phe Leu Glu Ile Trp Phe Thr
 1 5 10 15
 Thr Ala Cys Val Pro Lys Thr Leu Ala Thr Phe Ala Pro Arg Gly Gly
 20 25 30
 Val Ile Ser Leu Ala Gly Cys Ala Thr Gln Met
 35 40

<210> 335
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFF12rat GPCR TM2

<400> 335
 Met Tyr Phe Phe Leu Ala Asn Leu Ser Phe Val Asp Ile Cys Phe Thr
 1 5 10 15
 Ser Thr Thr Ile Pro Lys Met Leu Val Asn Ile Tyr Thr Gln Ser Lys
 20 25 30
 Ser Ile Thr Tyr Glu Asp Cys Ile Ser Gln Met
 35 40

<210> 336
 <211> 43
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFI3rat GPCR TM2

<400> 336
 Met Tyr Leu Phe Leu Ser Asn Leu Ser Phe Ser Asp Leu Cys Phe Ser
 1 5 10 15
 Ser Val Thr Met Pro Lys Leu Leu Gln Asn Met Arg Ser Gln Asp Thr
 20 25 30
 Ser Ile Pro Tyr Gly Gly Cys Leu Ala Gln Thr
 35 40

<210> 337
 <211> 47
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> OLFI7rat GPCR TM2

640000

<400> 337

Met	Tyr	Phe	Phe	Leu	Ala	Asn	Met	Ser	Phe	Leu	Glu	Ile	Trp	Tyr	Val
1				5					10					15	
Thr	Val	Thr	Ile	Pro	Lys	Met	Leu	Ala	Gly	Phe	Ile	Gly	Ser	Lys	Glu
			20					25					30		
Asn	His	Gly	Gln	Leu	Ile	Ser	Phe	Glu	Ala	Cys	Met	Thr	Gln	Leu	
		35					40					45			

<210> 338

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLFI8rat GPCR TM2

<400> 338

Met	Tyr	Leu	Phe	Leu	Ser	Asn	Leu	Ser	Phe	Ser	Asp	Leu	Cys	Phe	Ser
1				5					10					15	
Ser	Val	Thr	Met	Leu	Lys	Leu	Leu	Gln	Asn	Ile	Gln	Ser	Gln	Val	Pro
			20					25					30		
Ser	Ile	Ser	Tyr	Ala	Gly	Cys	Leu	Thr	Gln	Ile					
		35					40								

<210> 339

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLFI9rat GPCR TM2

<400> 339

Met	Tyr	Leu	Phe	Leu	Ser	Asn	Leu	Ser	Phe	Ala	Asp	Leu	Cys	Phe	Ser
1				5					10					15	
Ser	Val	Thr	Met	Pro	Lys	Leu	Leu	Gln	Asn	Met	Gln	Ser	Gln	Val	Pro
			20					25					30		
Ser	Ile	Pro	Tyr	Ala	Gly	Cys	Leu	Ala	Gln	Ile					
		35					40								

<210> 340

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLFI14rat GPCR TM2

<400> 340

Met	Tyr	Leu	Phe	Leu	Ser	Asn	Leu	Ser	Phe	Ser	Asp	Leu	Cys	Phe	Ser
1				5					10					15	
Ser	Val	Thr	Met	Pro	Lys	Leu	Leu	Gln	Asn	Met	Gln	Ser	Gln	Val	Pro
			20					25					30		
Ser	Ile	Ser	Tyr	Thr	Gly	Cys	Leu	Thr	Gln	Leu					
		35					40								

<210> 341

<211> 43

<212> PRT

64PPPP

<213> Artificial Sequence

<220>

<223> OLF115rat GPCR TM2

<400> 341

Met	Tyr	Leu	Phe	Leu	Ser	Asn	Leu	Ser	Phe	Ser	Asp	Leu	Cys	Phe	Ser
1				5					10					15	
Ser	Val	Thr	Met	Pro	Lys	Leu	Leu	Gln	Asn	Met	Gln	Ser	Gln	Val	Pro
			20					25					30		
Ser	Ile	Pro	Phe	Ala	Gly	Cys	Leu	Thr	Gln	Leu					
		35					40								

<210> 342

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> OLFOR17_40 GPCR TM2

<400> 342

Met	Tyr	Phe	Phe	Leu	Gly	Asn	Leu	Ser	Val	Leu	Asp	Val	Gly	Cys	Ile
1				5					10					15	
Ser	Val	Thr	Val	Pro	Ser	Met	Leu	Ser	Arg	Leu	Leu	Ser	Arg	Lys	Arg
			20					25					30		
Ala	Val	Pro	Cys	Gly	Ala	Cys	Leu	Thr	Gln	Leu					
		35					40								

<210> 343

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> GUST27rat GPCR TM2

<400> 343

Met	Tyr	Phe	Phe	Leu	Ser	Asn	Leu	Ser	Phe	Val	Asp	Ile	Cys	Phe	Ile
1				5					10					15	
Ser	Thr	Thr	Ile	Pro	Lys	Met	Leu	Val	Asn	Ile	His	Ser	Gln	Thr	Lys
			20					25					30		
Asp	Ile	Ser	Tyr	Ile	Glu	Cys	Leu	Ser	Gln	Val					
		35					40								

<210> 344

<211> 42

<212> PRT

<213> Artificial Sequence

<220>

<223> RPE GPCR TM2

<400> 344

Cys	His	Leu	Leu	Val	Leu	Ser	Leu	Ala	Leu	Ala	Asp	Ser	Gly	Ile	Ser
1				5					10					15	
Leu	Asn	Ala	Leu	Val	Ala	Ala	Thr	Ser	Ser	Leu	Leu	Arg	Arg	Trp	Pro
		20						25					30		

64QQQQ

Tyr Gly Ser Asp Gly Cys Gln Ala His Gly
 35 40

<210> 345
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> HHRF1 GPCR TM2

<400> 345
 Gly Asp Val Tyr Phe Ile Asn Leu Ala Ala Ala Asp Leu Leu Phe Val
 1 5 10 15
 Cys Thr Leu Pro Leu Trp Met Gln Tyr Leu Leu Asp His Asn Ser Leu
 20 25 30
 Ala Ser Val Pro Cys Thr Leu Leu Thr
 35 40

<210> 346
 <211> 42
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> HHRF2 GPCR TM2

<400> 346
 Ser Asp Thr Tyr Ile Cys Asn Leu Ala Val Ala Asp Leu Leu Ile Val
 1 5 10 15
 Val Gly Leu Pro Phe Phe Leu Glu Tyr Ala Lys His His Pro Lys Leu
 20 25 30
 Ser Arg Glu Val Val Cys Ser Gly Leu Asn
 35 40

<210> 347
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> HHRF3 GPCR TM2

<400> 347
 Pro Thr Ile Tyr Met Thr Asn Leu Tyr Ser Thr Asn Phe Leu Thr Leu
 1 5 10 15
 Thr Val Leu Pro Phe Ile Val Leu Ser Asn Gln Trp Leu Leu Pro Ala
 20 25 30
 Gly Val Ala Ser Cys Lys Phe Leu Ser
 35 40

<210> 348
 <211> 41
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> MCP-1A and MCP-1B GPCR TM2

64RRRR

<400> 348

Thr	Asp	Ile	Tyr	Leu	Leu	Asn	Leu	Ala	Ile	Ser	Asp	Leu	Leu	Phe	Leu
1				5					10					15	
Ile	Thr	Leu	Pro	Leu	Trp	Ala	His	Ser	Ala	Ala	Asn	Glu	Trp	Val	Phe
			20					25					30		
Gly	Asn	Ala	Met	Cys	Lys	Leu	Phe	Thr							
		35					40								

<210> 349

<211> 41

<212> PRT

<213> Artificial Sequence

<220>

<223> PPR1bovine GPCR TM2

<400> 349

Thr	Asp	Val	Tyr	Ile	Leu	Asn	Leu	Ala	Val	Ala	Asp	Leu	Phe	Leu	Leu
1				5					10					15	
Phe	Thr	Leu	Pro	Phe	Trp	Ala	Val	Asn	Ala	Val	His	Gly	Trp	Val	Leu
			20					25					30		
Gly	Lys	Ile	Met	Cys	Lys	Val	Thr	Ser							
		35					40								

<210> 350

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> F-1-5 GPCR CXCR4

<221> MOD_RES

<222> (18)...(18)

<223> Xaa = valinamide

<400> 350

Asp	Asp	Ile	Phe	Leu	Pro	Thr	Ile	Tyr	Ser	Ile	Ile	Phe	Leu	Thr	Gly
1				5					10					15	
Ile	Xaa														

<210> 351

<211> 22

<212> PRT

<213> Artificial Sequence

<220>

<223> F-2-1 GPCR CXCR4

<400> 351

Leu	Leu	Phe	Val	Ile	Thr	Leu	Pro	Phe	Trp	Ala	Val	Asp	Ala	Val	Ala
1				5					10					15	
Asn	Trp	Tyr	Phe	Gly	Asn										
			20												

<210> 352

<211> 23

<212> PRT

64SSSS

<213> Artificial Sequence

<220>

<223> F-3-1 GPCR CXCR4

<221> MOD_RES

<222> (23)...(23)

<223> Xaa = leucinamide

<400> 352

Lys	Ala	Val	His	Val	Ile	Tyr	Thr	Val	Asn	Leu	Tyr	Ser	Ser	Val	Leu
1				5					10					15	
Ile	Leu	Ala	Phe	Ile	Ser	Xaa									
			20												

<210> 353

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> F-4-1 GPCR CXCR4

<400> 353

Lys	Val	Tyr	Val	Gly	Val	Trp	Ile	Pro	Ala	Leu	Leu	Leu	Thr	Ile	Pro
1				5					10					15	
Asp	Phe	Ile	Phe												
			20												

<210> 354

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> F-5-1 GPCR CXCR4

<221> MOD_RES

<222> (21)...(21)

<223> Xaa = isoleucinamide

<400> 354

His	Ile	Met	Val	Gly	Leu	Ile	Leu	Pro	Gly	Ile	Val	Ile	Leu	Ser	Cys
1				5					10					15	
Tyr	Cys	Ile	Ile	Xaa											
			20												

<210> 355

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> F-7-1 GPCR CXCR4

<221> MOD_RES

<222> (20)...(20)

<223> Xaa = lysinamide

64TTTT

<400> 355
 Ala Leu Ala Phe Phe His Cys Cys Leu Asn Pro Ile Leu Tyr Ala Phe
 1 5 10 15
 Leu Gly Ala Xaa
 20

<210> 356
 <211> 32
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> second transmembrane domain of CXCR4

<400> 356
 His Leu Ser Val Ala Asp Leu Leu Phe Val Ile Thr Leu Pro Phe Trp
 1 5 10 15
 Ala Val Asp Ala Val Ala Asn Trp Tyr Phe Gly Asn Phe Leu Cys Lys
 20 25 30

<210> 357
 <211> 24
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> F-2-8 GPCR CXCR4

<400> 357
 Leu Leu Phe Val Ile Thr Leu Pro Phe Trp Ala Val Asp Ala Val Ala
 1 5 10 15
 Asn Trp Tyr Phe Gly Asn Lys Lys
 20

<210> 358
 <211> 21
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> F-2-4 GPCR CXCR4

<400> 358
 Val Ile Thr Leu Pro Phe Trp Ala Val Asp Ala Val Ala Asn Trp Tyr
 1 5 10 15
 Phe Gly Asn Lys Lys
 20

<210> 359
 <211> 19
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> AcF-2-5 GPCR CXCR4

<221> MOD_RES
 <222> (1)...(1)
 <223> Xaa = acetylated Leu

64UUUU

<400> 359

Xaa	Leu	Phe	Val	Ile	Thr	Leu	Pro	Phe	Trp	Ala	Val	Asp	Ala	Val	Ala
1				5					10					15	
Asn	Asp	Asp													

<210> 360

<211> 24

<212> PRT

<213> Artificial Sequence

<220>

<223> F-2-6 GPCR CXCR4

<400> 360

Leu	Ser	Val	Ala	Asp	Leu	Leu	Phe	Val	Ile	Thr	Leu	Pro	Phe	Trp	Ala
1				5					10					15	
Val	Asp	Ala	Val	Ala	Asn	Asp	Asp								
				20											

<210> 361

<211> 27

<212> PRT

<213> Artificial Sequence

<220>

<223> Rhod-AcF-2-2 GPCR CXCR4

<221> MOD_RES

<222> (1)...(1)

<223> Xaa = acetylated Leu

<221> MOD_RES

<222> (26)...(26)

<223> Xaa = rhodamine linked to Lys

<400> 361

Xaa	Leu	Leu	Phe	Val	Ile	Thr	Leu	Pro	Phe	Trp	Ala	Val	Asp	Ala	Val
1				5					10					15	
Ala	Asn	Trp	Tyr	Phe	Gly	Asn	Asp	Asp	Xaa	Asp					
			20					25							

<210> 362

<211> 28

<212> PRT

<213> Artificial Sequence

<220>

<223> CCKAR-TM-4-2 (#71) GPCR CCKAR

<400> 362

Val	Ile	Ala	Ala	Thr	Trp	Cys	Leu	Ser	Phe	Thr	Ile	Met	Thr	Pro	Tyr
1				5					10					15	
Pro	Ile	Tyr	Ser	Asn	Leu	Val	Pro	Phe	Thr	Asp	Asp				
			20					25							

<210> 363

<211> 24

Asp Asp Gln Thr Phe Leu Leu Leu Ile Leu Phe Leu Leu Pro Gly Ile
1 5 10 15
Val Met Val Val Ala Tyr Gly Leu
20

CLAIMS:

1. An isolated G protein-coupled receptor (GPCR)-modulating molecule comprising a peptide or peptidomimetic having 5 to 50 amino acids that is a structural analog of a portion of a transmembrane domain of a GPCR, wherein

said molecule has an N-terminus and a C-terminus and said molecule has an extracellular end that is negatively charged and an intracellular end that has a neutral charge under physiological conditions;

said molecule has an amidated C-terminal carboxyl group when said C-terminus has a neutral charge, and when said N -terminus is the intracellular end, said peptide or peptidomimetic includes at least four of the first five N -terminal amino acids having neutrally-charged amino acid side chains under physiological conditions;

said molecule spontaneously inserts into a membrane in the same orientation as the transmembrane GPCR domain from which it is derived; and
said molecule modulates a biological activity of said GPCR by disrupting the structure or assembly of said GPCR.

2. The molecule of claim 1, wherein the negative charge of the extracellular end of the molecule is provided by one to three amino acid residues that are acidic under physiological conditions.

3. The molecule of claim 1 or 2, wherein the negative charge of the extracellular end of the molecule is provided by a carboxyl, phosphate, borate, sulfonate, or sulfate group.

4. The molecule of claim 1, wherein the GPCR is CXCR4, said molecule comprising a peptide or peptidomimetic that is a structural analog of a portion of a transmembrane domain of CXCR4, wherein said portion of said transmembrane domain consists of a sequence selected from the group of sequences consisting of:

LLFVITLPFWAVDAVANWYFG N DD (SEQ ID NO: 1),

LLFVITLPFWAVDAVANDD-OH (SEQ ID NO:2),

VYVGWIPALLLTIPDFIFANDD-OH (SEQ ID NO:3),

VILILAFFACWLPYYIGISID-OH (SEQ ID NO:4),

DDEALAFFHCCLNPILYAFL-NH₂ (SEQ ID NO:5),

and

DDSITEALAFFHCCLNPILYAFL-NH₂ (SEQ ID NO:6),

wherein said molecule modulates a biological activity of said CXCR4.

5. The molecule of claim 4, wherein the CXCR4 activity modulated is inhibition of CXCR4-mediated intracellular Ca²⁺ release.

6. The molecule of claim 4, wherein the CXCR4 activity modulated is CXCR4-mediated HIV infection.

7. The molecule of anyone of claims 1-6 for use in modulating the biological property or activity of a target GPCR in a cell that expresses said GPCR.

8. The molecule of claim 1, wherein the molecule comprises a peptide or peptidomimetic that is a structural analog of a portion of a transmembrane domain of the GPCR, selected from the group consisting of:

From the GPCR CXCR4

F-2-2: LLFVITLPFWAVDAVANWYFG N DD (SEQ ID NO: 1),

F-2-5: LLFVITLPFWAVDAVANDD-OH (SEQ ID NO:2)

F-4-2: VYVGWIPALLLTIPDFIFANDD-OH (SEQ ID NO:3)

F-6-1: VILILAFFACWLPPYYIGISID-OH (SEQ ID NO:4)

F-7-3: DDEALAFFHCCLNPILYAFL-NH₂ (SEQ ID NO:5) and

F-7-4: DDSITEALAFFHCCLNPILYAFL-NH₂ (SEQ ID NO:6)

From the GPCR CCR5

CCR5-TM-2-2: LFFL LTVPFWAHYAAAQWDFGDD (SEQ ID NO:7)

CCR5-TM-4-1: FGVVTSVITWVVAVFASLPGIIFTSSDD (SEQ ID NO:8) and

CCR5-TM-6-1: LIFTIMIVYFLFWAPYNNLLNTFQED (SEQ ID NO:9)

From the GPCR CCR2

CCR2-TM-2-1: IYLLNLAISDLLFLITLPLWADD-OH (SEQ ID NO:11)

CCR2-TM-2-2: LLFLITLPLWAHSAANEWVFGNDD-OH (SEQ ID NO:12)

CCR2-TM-4-1: FGVVTSVITWLVAVF ASVPGIIFTDD (SEQ ID NO:13) and
CCR2-TM-6-1: VIFTIMIVYFLFWTPYN NILLNTFQED (SEQ ID NO:14)

From the GPCR CCR3

CCR3-TM-2-1: LLFLVTLPFW IHYVRGHNWVFGDDD (SEQ ID NO:16)
CCR3-TM-4-1: FGV1TSNTWGLAVLAALPEFI FYETED (SEQ ID NO: 17) and
CCR3-TM-6-1: IFVIMAVFFI FWTPYNVAILLSSYQSDD (SEQ ID NO:18)
and

From the GPCR CCKAR

CCKAR-TM-1-6: DDEWQSALQILLYSIIFLLSVLGNTLVITV (SEQ ID NO:20)
CCKAR-TM-2-2: FLLSLAVSDLMLCLFCMPFNLIDD (SEQ ID NO:22) and
CCKAR-TM-6-4: IVVLFFLCWMPIFSANAWRAYDTVDD (SEQ ID NO:23).

9. Use of the molecule of anyone of claims 1-3 for modulating the biological activity of a target GPCR in a cell that expresses said GPCR.
10. Use of the molecule of anyone of claims 1-3 for formulating a medicament for modulating the biological property or activity of a target GPCR in a cell that expresses said GPCR.
11. The use of the molecule according to claim 9 or 10, wherein the concentration of the molecule is about 0.01 to about 100 micromolar.
12. The use of the molecule according to claim 9, 10 or 11, wherein the molecule inhibits the biological activity of the GPCR.
13. The use of the molecule according to anyone of claims 9-12, wherein the inhibited biological activity is selected from the group consisting of ion flux or translocation, phosphorylation, protein synthesis or degradation, cellular morphology, secretion, production of soluble inositol phosphates, binding of a virus and subsequent infection, tumor growth, chemotaxis, mitogenic response, cell growth activation, muscle contraction, vasopressing and vasodepressing activity, synaptic transmission, and release of intracellular

calcium.

14. The use of the molecule according to anyone of claims 9-13, wherein the concentration of the molecule is about 0.2 to about 10 micromolar.
15. The use of the molecule according to anyone of claims 9-14, wherein the target GPCR is CXCR4 and the modulated biological activity is inhibition of CXCR4-mediated HIV infection.
16. The use of the molecule according to anyone of claims 9-14, wherein the target GPCR is CCR5 and the modulated biological activity is inhibition of CCR5-mediated HIV infection.
17. The use of the molecule according to anyone of claims 9-14 or 16, wherein the target GPCR is CCR5 and the modulated biological activity is inhibition of CCR5-mediated intracellular Ca^{2+} release.
18. The use of the molecule according to anyone of claims 9-14, wherein the target GPCR is CCKAR and the modulated biological activity is inhibition of CCKAR-mediated intracellular Ca^{2+} release.
19. The use of the molecule according to anyone of claims 9-14 or 18, wherein the target GPCR is CCKAR and the modulated biological activity is inhibition of tumor growth or progression.
20. The molecule of anyone of claims 1-6, for use in inhibiting HIV -1 infection in a cell that expresses the GPCR, and wherein the GPCR binds HIV-1.
21. Use of the molecule of anyone of claims 1-3, for inhibiting HIV-l infection in a cell that expresses the GPCR, and wherein the GPCR binds HIV-1.

22. Use of the molecule of anyone of claims 1-3, to formulate a medicament for inhibiting HIV-1 infection in a cell that expresses the GPCR, and wherein the GPCR binds HIV-1.

23. The use of the molecule according to claim 21 or 22, wherein the GPCR is CXCR4, CCR2, CCR3, or CCR5.

24. Use of the molecule of anyone of claims 4-6, for inhibiting HIV -1 infection in a cell that expresses the GPCR, and wherein the GPCR binds HIV-1.

25. Use of the molecule of anyone of claims 4-6, to formulate a medicament for inhibiting HIV-1 infection in a cell that expresses the GPCR, and wherein the GPCR binds HIV-1.

26. The use of the molecule according to anyone of claims 21 to 25, wherein the molecule comprises a CXCR 4 transmembrane peptide consisting of a sequence selected from the group of sequences consisting of:

LLFVITLPFWAVDAVANWYFG N DD (SEQ ID NO: 1),

LLFVITLPFWAVDAVANDD-OH (SEQ ID NO:2),

VYVGWIPALLLTIPDFIFANDD-OH (SEQ ID NO:3),

VILILAFFACWLPYYIGISID-OH (SEQ ID NO:4),

DDEALAFFHCCLNPILYAFL-NH₂ (SEQ ID NO:5),

and

DDSITEALAFFHCCLNPILYAFL-NH₂ (SEQ ID NO:6),

and wherein said molecule inhibits CXCR4-mediated infection of CXCR4-expressing cells with said HIV virus.

27. Use of the molecule of anyone of claims 4-6, for modulating the biological activity of a target GPCR in a cell that expresses said GPCR.

28. Use of the molecule of any one of claims 4-6, for formulating a medicament for

modulating the biological property or activity of a target GPCR in a cell that expresses said GPCR.

29. A pharmaceutical composition comprising the molecule of anyone of claims 1-8 and a pharmaceutically acceptable excipient.

30. The pharmaceutical composition of claim 29, wherein the GPCR is CXCR4 and the portion of a trans membrane domain of CXCR4 is LLFVITLPFW A VDA V ANWYFGNDD (SEQ ID NO:1).

31. Use of the molecule of anyone of claims 1-8 for inhibiting cell growth in a cell that expresses the GPCR.

32. Use of the molecule of anyone of claims 1-8 to formulate a medicament for inhibiting cell growth in a cell that expresses the GPCR.

33. The use of claim 31, wherein the cell is a cancer cell.

34. The use of claim 32, wherein the cell is a cancer cell.

35. The use of claim 33, wherein the cancer cell is in a human patient.

36. The use of claim 34, wherein the cancer cell is in a human patient.

37. The use of claim 35, wherein the molecule is in a formulation for oral, parenteral, enterical, gastrical, topical, subcutaneous, rectal, local, or systemic use.

38. The use of claim 36, wherein the medicament is formulated for oral, parenteral, enterical, gastrical, topical, subcutaneous, rectal, local, or systemic use.

39. The use of claim 37, wherein the molecule is in a formulation for use by injection into the

bloodstream of the patient.

40. The use of claim 38, wherein the medicament is formulated for use by injection into the bloodstream of the patient.

41. The use of claim 35, wherein the molecule is in a formulation for daily use in the amount of 0.1 μ g to 100 mg.

42. The use of claim 41, wherein the molecule is in a formulation for daily use in the amount of 0.1 mg to 10 mg.

43. The use of claim 36, wherein the medicament is formulated in a daily dose format comprising the molecule in the amount of 0.1 μ g to 100 mg.

44. The use of claim 43, wherein the molecule is in the amount of 0.1 mg to 10 mg.

45. The use of anyone of claims 31 to 44, wherein the GPCR is CXCR4 and the portion of a trans membrane domain of CXCR4 is LLFVITLPFWAVDAVANWYFGNDD (SEQ ID NO:1).

1/3




Figure 1.

2/3




Figure 2.




Figure 3.

