


**(12) PATENT ABRIDGMENT (11) Document No. AU-B-49226/90**  
**(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 630912**

- (54) Title  
**INSULIN ANALOGS**
- International Patent Classification(s)  
 (51)<sup>s</sup> **C07K 007/40 A61K 037/26**
- (21) Application No. : **49226/90** (22) Application Date : **07.02.90**
- (30) Priority Data
- (31) Number (32) Date (33) Country  
**308352 09.02.89 US UNITED STATES OF AMERICA**  
**388201 04.08.89 US UNITED STATES OF AMERICA**
- (43) Publication Date : **16.08.90**
- (44) Publication Date of Accepted Application : **12.11.92**
- (71) Applicant(s)  
**ELI LILLY AND COMPANY**
- (72) Inventor(s)  
**RONALD EUGENE CHANCE; RICHARD DENNIS DIMARCHI; BRUCE HILL FRANK; JAMES EDWIN SHIELDS**
- (74) Attorney or Agent  
**SPRUSON & FERGUSON , GPO Box 3898, SYDNEY NSW 2001**
- (57) Claim

1. An insulin analog of the formula



or a pharmaceutically acceptable salt thereof, wherein A21 is alanine, asparagine, aspartic acid, glutamine,

(11) AU-B-49226/90  
(10) 630912

-2-

glutamic acid, glycine, threonine, or serine; B1 is phenylalanine, aspartic acid, or is absent; B2 is valine or may be absent when B1 is absent; B3 is asparagine or aspartic acid; B9 is serine or aspartic acid; B10 is histidine or aspartic acid; B27 is threonine or is absent; B28 is any amino acid, B29 is L-proline, D-proline, D-hydroxyproline, L-hydroxyproline, L-(N-methyllysine), D-lysine, L-(N-methylarginine) or D-arginine; B30 is alanine, threonine or is absent; Z is -OH, -NH<sub>2</sub>, -OCH<sub>3</sub>, or -OCH<sub>2</sub>CH<sub>3</sub>; X is Arg, Arg-Arg, Lys, Lys-Lys, Arg-Lys, Lys-Arg, or is absent; and Y may be present only when X is present and, if present, is Glu or an amino acid sequence which comprises all or a portion of the sequence -Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-Leu-Gly-Gly-Gly-Pro-Gly-Ala-Gly-Ser-Leu-Gln-Pro-Leu-Ala-Leu-Glu-Gly-Ser-Leu-Gln-Lys-Arg- and which begins at the N-terminus Glu of such sequence.

25. A process for preparing a compound of Formula (I) and as defined in any one of Claims 1 to 24, which comprises:

- A) combining the A-chain of a compound of Formula (I) in which A21 is as defined in Claim 1, with the B-chain of a compound of Formula (I) in which B1, B2, B3, B9, B10, B27, B28, B29, B30, X, Y, and Z are as defined in Claim 1 to form the appropriate disulfide bonds; or
- B) reacting a modified compound of Formula (I) lacking the peptide sequence beginning at B23 and in which A21, B1, B2, B3, B9 and B10 are as defined in Claim 1 with a peptide having the sequence Gly-Phe-Phe-Tyr-B27-B28-B29-B30-X-Y-Z in which B27, B28, B29, B30, X, Y, and Z are as defined in Claim 1; or

(11) AU-B-49226/90  
(10) 630912

-3-

- C) cleaving a modified proinsulin molecule comprising the amino acid sequence of a compound of Formula (I) in which A21, B1, B2, B3, B9, B10, B27, B28, B29, B30, X, Y, and Z are as defined in Claim 1 to remove all or a portion of the C-peptide from such modified proinsulin.

34. A pharmaceutical composition for the treatment of diabetes mellitus in a mammal, which composition comprises an amount effective for the treatment of diabetes mellitus in said mammal of an insulin analog according to Claim 29 together with a pharmaceutically acceptable carrier, diluent and/or adjuvant.

FORM 10

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952

COMPLETE SPECIFICATION

630912

(ORIGINAL)

FOR OFFICE USE:

Class Int Class

Complete Specification Lodged:  
Accepted:  
Published:

Priority:

Related Art:

Name and Address  
of Applicant:

Eli Lilly and Company  
Lilly Corporate Center  
City of Indianapolis State of Indiana  
UNITED STATES OF AMERICA

Address for Service:

Spruson & Ferguson, Patent Attorneys  
Level 33 St Martins Tower, 31 Market Street  
Sydney, New South Wales, 2001, Australia

Complete Specification for the invention entitled:

Insulin Analogs


The following statement is a full description of this invention, including the best method of performing it known to me/us

INSULIN ANALOGS

The present invention relates to analogs of insulin modified at amino acid position 29 of the native human insulin B chain and optionally at other positions.

Said insulin analogs are less prone to dimerization or self-association to higher molecular weight forms thereby possessing a comparatively more rapid onset of activity while retaining the biological activity of native human insulin.

*According to a first embodiment, the*  
~~The~~ present invention provides insulin analogs of the formula



wherein A21 is alanine, asparagine, aspartic acid, glutamine, glutamic acid, glycine, threonine, or serine; B1 is phenylalanine, aspartic acid, or is absent; B2 is valine or may be absent when B1 is absent; B3 is asparagine or aspartic acid; B9 is serine or aspartic acid; B10 is histidine or aspartic acid; <sup>B27 is threonine or is absent;</sup> B28 is any amino acid, B29 is L-proline, D-proline, D-hydroxyproline, L-hydroxyproline, L-(N-methyllysine), D-lysine, L-(N-methylarginine) or D-arginine; B30 is alanine, threonine or is absent; Z is -OH, -NH<sub>2</sub>, -OCH<sub>3</sub>, or -OCH<sub>2</sub>CH<sub>3</sub>; X is Arg, Arg-Arg, Lys, Lys-Lys, Arg-Lys, Lys-Arg, or is absent; and Y may be present only when X is present and, if present, is Glu or an amino acid sequence which comprises all or a portion of the sequence -Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-Leu-Gly-Gly-Gly-Pro-Gly-Ala-Gly-Ser-Leu-Gln-Pro-Leu-Ala-Leu-Glu-Gly-Ser-Leu-Gln-Lys-Arg- and which begins at the N-terminus Glu of such sequence.

Also disclosed and claimed is a method of treating hyperglycemia by administering to a patient in need thereof an effective amount of an insulin analog of formula I. Further, pharmaceutical compositions containing an effective amount of an insulin analog of formula I in combination with one or more pharmaceutically acceptable excipients are disclosed and claimed.

~~The figures provided herein are not drawn exactly to scale.~~



Thus, according to a second embodiment of this invention, there is provided a process for preparing a compound of Formula (I) and according to the first embodiment, which comprises:

- 5 A) combining the A-chain of a compound of Formula (I) in which A21 is as defined in the first embodiment, with the B-chain of a compound of Formula (I) in which B1, B2, B3, B9, B10, B27, B28, B29, B30, X, Y, and Z are as defined in the first embodiment to form the appropriate disulfide bonds; or
- 10 B) reacting a modified compound of Formula (I) lacking the peptide sequence beginning at B23 and in which A21, B1, B2, B3, B9 and B10 are as defined in the first embodiment with a peptide having the sequence Gly-Phe-Phe-Tyr-B27-B28-B29-B30-X-Y-Z in which B27, B28, B29, B30, X, Y, and Z are as defined in the first embodiment; or
- 15 C) cleaving a modified proinsulin molecule comprising the amino acid sequence of a compound of Formula (I) in which A21, B1, B2, B3, B9, B10, B27, B28, B29, B30, X, Y, and Z are as defined in the first embodiment to remove all or a portion of the C-peptide from such modified proinsulin.

20 According to a third embodiment of this invention, there is provided a pharmaceutical formulation comprising, as an active ingredient, a compound of Formula (I) according to the first embodiment, associated with one or more pharmaceutically-acceptable carriers therefor.

25 According to a fourth embodiment of this invention, there is provided a method for treating diabetes mellitus in mammals afflicted therewith, which comprises administering to such mammal a compound of Formula (I) according to the first embodiment.

30 According to a fifth embodiment of this invention, there is provided a compound of Formula (I) according to the first embodiment, whenever produced by the process of the third embodiment.

The figures provided herein are not drawn exactly to scale.



Figure 1 - A restriction site and function map of plasmid pKC283.

Figure 2 - A restriction site and function map of plasmid pKC283PX.

5 Figure 3 - A restriction site and function map of plasmid pKC283-L.

Figure 4 - A restriction site and function map of plasmid pKC283-LB.

10 Figure 5 - A restriction site and function map of plasmid pKC283-PRS.

Figure 6 - A restriction site and function map of plasmid pL32.

Figure 7 - A restriction site and function map of plasmid pNM789.

15 Figure 8 - A schematic outline of the construction of plasmid 120.

Figure 9 - A restriction site and function map of plasmid pL47.

20 Figure 10 - A restriction site and function map of plasmid pPR12.

Figure 11 - A restriction site and function map of plasmid pPR12AR1.

Figure 12 - A restriction site and function map of plasmid pL110.

25 Figure 13 - A schematic outline of the construction of plasmid pL110C.



Figure 14 - A restriction site and function map of plasmid pCZR126S.

Figure 15 - The nucleotide sequence of the synthetic human proinsulin gene.

5           Figure 16 - A restriction site and function map of plasmid pRB145.

Figure 17 - A restriction site and function map of plasmid pRB164A.

10           Figure 18 - A restriction site and function map of plasmid pRB172.

Figure 19 - A restriction site and function map of plasmid pRB173.

Figure 20 - A restriction site and function map of plasmid pRB175.

15           Formula I is a short-hand representation of the amino acid sequence of the insulin analogs of the present invention. The amino acid abbreviations have the following conventional meanings:

AbbreviationAmino Acid

Aba

 $\alpha$ -Aminobutyric acid

Ala

Alanine

Arg

Arginine

5

Asn

Asparagine

Asp

Aspartic acid

Cya

Cysteic acid

Cys

Cysteine

Gln

Glutamine

10

Glu

Glutamic acid

Gly

Glycine

His

Histidine

Ile

Isoleucine

Leu

Leucine

15

Lys

Lysine

Met

Methionine

Nle

Norleucine

Nva

Norvaline

Orn

Ornithine

20

Phe

Phenylalanine

Pro

Proline

Ser

Serine

|     |            |
|-----|------------|
| Thr | Threonine  |
| Trp | Tryptophan |
| Tyr | Tyrosine   |
| Val | Valine     |

5

B28 may be any natural or non-naturally occurring amino acid. Preferably, said amino acid is aspartic acid, valine, leucine, isoleucine, norleucine, proline, arginine, histidine, citrulline, ornithine, lysine, phenylalanine, alanine or glycine. Of the above amino acids, a particularly preferred subgroup thereof is aspartic acid, valine, leucine, isoleucine, norleucine, proline, arginine, histidine, ornithine or lysine. Lysine is the most particularly preferred amino acid at B28. B29 preferably is L-proline, D-proline, D-hydroxyproline or L-hydroxyproline. A particularly preferred insulin analog of the present invention is one wherein B28 is lysine and B29 is proline, i.e., an inversion of the native human insulin amino acid sequence at positions 28 and 29 of the B-chain.

In a further embodiment, as noted above, when B28 is L-proline, then position B29 preferably is L-(N-methyllysine), D-lysine, L-(N-methylarginine) or D-arginine. A preferred analog of this embodiment is that wherein B28 is L-proline and B29 is L-(N-methyllysine) or D-lysine.

Further modifications of the insulin analogs of the present invention are also contemplated hereunder, i.e., modifications to the insulin analogs at positions other than at positions B28 and B29. Specifically, it may be desirable to optionally substitute the Asn residue from position 21 of the A chain (i.e., the carboxy terminus) with Ala, Asp, Gln, Glu, Gly, Thr or Ser, and, if so substituted, preferably with Ala. Similarly, it may be desirable to substitute the Asn residue at position 3 of the B-chain with aspartic acid (Asp). Such optional substitutions have the effect of increasing the stability of the analogs at pH extremes since Asn is particularly sensitive to deamidation and rearrangement reactions at both low and high pH. The skilled artisan will also readily appreciate that the glutamine residues of the insulin analogs may similarly be sensitive to deamidation and rearrangement. Accordingly, the substitution therefor by glutamic acid is also within the scope of the present invention. Additional, optional modifications to the insulin analogs of the present invention include (in any combination) replacement of the histidine residue at position B10 with the amino acid aspartic acid; replacement of the phenylalanine residue at position B1 with aspartic acid; replacement of the threonine residue at position B30 with alanine; replacement of the serine residue at position B9 with aspartic acid; deletion of amino acids at position B1 (des-B1) alone or in combination with a deletion at position B2 (des-B2); and deletion of threonine from position B-30 (des-B30).

The insulin analogs of this invention may also be modified at the B30 terminus by addition of any of

the following amino acids or dipeptides: Arg, Arg-Arg, Lys, Lys-Lys, Arg-Lys, or Lys-Arg. These, when present, are designated in Formula I by the group X. Preferably, when such an extension is present, it is Arg-Arg.

5                   In addition, should the foregoing extension at B30 be present, the analog may be further modified by addition at the resulting B30-lengthened terminus of glutamic acid (Glu) or of an amino acid sequence comprising all or a portion of the sequence -Glu-Ala-Glu-  
 10 Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-Leu-Gly-Gly-Gly-Pro-Gly-Ala-Gly-Ser-Leu-Gln-Pro-Leu-Ala-Leu-Glu-Gly-Ser-Leu-Gln-Lys-Arg-, which sequence begins at its N-terminus Glu. This amino acid or sequence, when present, is designated in Formula I by the group Y. When the sequence represents  
 15 only a portion of the foregoing, it will be any of those portions which begin at the N-terminus of the sequence, i.e., at the glutamic acid (Glu) residue.

                  In the foregoing, X or the combination of X and Y represents all or a portion of the connecting peptide  
 20 found in human proinsulin, a molecule which is the biological precursor in the formation of native human insulin.

                  Preferred sequences for Y are:

-Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-  
 Leu-Gly-Gly-Gly-Pro-Gly-Ala-Gly-Ser-Leu-Gln-Pro-Leu-Ala-  
 25 Leu-Glu-Gly-Ser-Leu-Gln-Lys-Arg-;  
 -Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-  
 Leu-Gly-Gly-Gly-Pro-Gly-Ala-Gly-Ser-Leu-Gln-Pro-Leu-Ala-  
 Leu-Glu-Gly-Ser-Leu-Gln-Lys-;  
 -Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-  
 30 Leu-Gly-Gly-Gly-Pro-Gly-Ala-Gly-Ser-Leu-Gln-Pro-Leu-Ala-  
 Leu-Glu-Gly-Ser-Leu-Gln-;  
 -Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-

Leu-Gly-Gly-Gly-Pro-Gly-Ala-Gly-Ser-Leu-Gln-Pro-Leu-Ala-  
Leu-Glu-;

-Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-

Leu-Gly-Gly-Gly-Pro-Gly-Ala-Gly-Ser-Leu-Gln-Pro-Leu-Ala-;

5 and

-Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-.

In addition, whether X alone or X and Y together  
are present or absent, the terminus group Z may be any of  
10 the following: -OH, -NH<sub>2</sub>, -OCH<sub>3</sub>, or -OCH<sub>2</sub>CH<sub>3</sub>. Preferably,  
Z is -OH.

As mentioned hereinabove, the invention includes  
pharmaceutically acceptable salts of the insulin analogs.  
Preferred such salts are those of zinc, sodium, potassium,  
15 magnesium, or calcium.

The insulin analogs of this invention can  
be prepared by any of a variety of recognized peptide  
synthesis techniques including classical (solution)  
methods, solid-phase methods, semisynthetic methods and  
20 the more recently available recombinant DNA methods.

In the solid-phase technique, the amino acid  
sequence is constructed sequentially from an initial,  
insoluble, resin-supported C-terminal amino acid.  
Techniques for the solid phase method are described  
25 by J. Stewart *et al.*, Solid-Phase Peptide Synthesis,  
Freeman and Co., San Francisco, 1969.

In general, in the solid-phase method, the  
amino acid corresponding to the C-terminal amino acid  
residue of the desired peptide is anchored to an  
30 insoluble resin support, and the peptide chain then  
is formed beginning at the resin-supported C-terminal

amino acid. Individual amino acids are introduced sequentially until the desired amino acid sequence is obtained. Alternatively, small peptide fragments can be prepared and introduced into the peptide chain in the desired order. The peptide chain remains attached to the resin throughout synthesis, and, upon completion of the chain, the peptide is cleaved from the resin.

The peptide chain is attached to the polystyrene resin by means of an ester linkage formed between the carboxyl group of the C-terminal moiety and a specific methylene group present on the resin matrix as a site for such attachment.

The amino acids are coupled using techniques well known in the art for the formation of peptide bonds. One method involves converting the amino acid to a derivative that will render the carboxyl group more susceptible to reaction with the free N-terminal amino group of the peptide fragment. For example, the amino acid can be converted to a mixed anhydride by reaction of a protected amino acid with ethyl chloroformate, phenyl chloroformate, sec-butyl chloroformate, or isobutyl chloroformate. Alternatively, the amino acid can be converted to an active ester such as a 2,4,5-trichlorophenyl ester, a pentachlorophenyl ester, a p-nitrophenyl ester, a N-hydroxysuccinimide ester, or an ester formed from 1-hydroxybenzotriazole.

Another coupling method involves use of a suitable coupling agent, such as N,N'-dicyclohexylcarbodiimide (DCC) or N,N'-diisopropylcarbodiimide (DIC). Other appropriate coupling agents will be apparent to those skilled in the art. See Schroder and Lubke,

The Peptides, Academic Press, 1965, Chapter III which is incorporated herein by reference.

5 It should be recognized that the  $\alpha$ -amino group of each amino acid employed in the peptide synthesis must be protected during the coupling reaction to prevent side reactions involving the reactive  $\alpha$ -amino function. It should also be recognized that certain amino acids contain reactive side-chain functional groups (e.g., sulfhydryl,  $\epsilon$ -amino,  $\beta$ - and  $\gamma$ -carboxyl, 10 imidazole, guanido and hydroxyl), and that such functional groups must also be protected both during the initial and subsequent coupling steps. Suitable protecting groups are known in the art. See for example, Protective Groups In Organic Chemistry, M. McOmie, 15 Editor, Plenum Press, N.Y., 1973 and U.S. Patent 4,617,119 which is incorporated herein by reference.

20 In selecting a particular protecting group, certain conditions must be observed. An  $\alpha$ -amino protecting group (1) must render the  $\alpha$ -amino function inert under the conditions employed in the coupling reaction, (2) must be readily removable after the coupling reaction under conditions that will not remove side chain protecting groups and will not alter the structure of the peptide fragment, and (3) must eliminate the possibility of racemization upon activation 25 immediately prior to coupling. A side chain protecting group (1) must render the side chain functional group inert under the conditions employed in the coupling reaction, (2) must be stable under the conditions 30 employed in removing the  $\alpha$ -amino protecting group, and (3) must be readily removable upon completion of the



desired amino acid sequence under reaction conditions that will not alter the structure of the peptide chain.

It will be apparent to those skilled in the art that the protecting groups known to be useful for peptide synthesis will vary in reactivity to the agents employed for their removal. For example, certain protecting groups, such as triphenyl methyl and 2-(p-biphenyl)isopropylloxycarbonyl are very labile and can be cleaved under mild acid conditions. Other protecting groups, such as t-butylloxycarbonyl, t-amylloxycarbonyl, adamantylloxycarbonyl, and p-methoxybenzylloxycarbonyl, are less labile and require moderately strong acids, such as trifluoroacetic, hydrochloric, or boron trifluoride in acetic acid, for their removal. Still other protecting groups, such as benzylloxycarbonyl, halobenzylloxycarbonyl, p-nitrobenzylloxycarbonyl, cycloalkylloxycarbonyl, and isopropylloxycarbonyl, are even less labile and require stronger acids, such as hydrogen fluoride, hydrogen bromide, or boron trifluoroacetate in trifluoroacetic acid, for their removal.

Upon completion of the desired peptide sequence, the protected peptide must be cleaved from the resin support, and all protecting groups must be removed. The cleavage reaction and removal of the protecting groups may be accomplished simultaneously or stepwise. When the resin support is a chloromethylated polystyrene resin, the bond anchoring the peptide to the resin is an ester linkage formed between the free carboxyl group of the C-terminal moiety and one of the many chloromethyl groups present on the resin matrix.

It will be recognized that the anchoring bond can be cleaved by reagents which are known to be capable of breaking an ester linkage and of penetrating the resin matrix. One especially convenient method is by treatment with liquid anhydrous hydrogen fluoride. This reagent not only will cleave the peptide from the resin but will also remove all protecting groups. Hence, use of this reagent will directly afford the fully deprotected peptide. When it is desired to cleave the peptide without removing protecting groups, the protected peptide-resin can undergo methanolysis to give the protected peptide in which the C-terminal carboxyl group is methylated. The methyl ester can then be hydrolyzed under mild, alkaline conditions to give the free C-terminal carboxyl. The protecting groups on the peptide chain then can be removed by treatment with a strong acid, such as liquid hydrogen fluoride. A particularly useful technique for methanolysis is that of G. Moore et al., Peptides, Proc. 5th Amer. Pept. Symp., M. Goodman and J. Meienhofer, Eds., John Wiley, N.Y., 1977, pp. 518-521, in which the protected peptide-resin is treated with methanol and potassium cyanide in the presence of crown ether.

Another method for cleaving the protected peptide from the resin is by ammonolysis or by treatment with hydrazine. If desired, the resulting C-terminal amide or hydrazide can be hydrolyzed to the free C-terminal carboxyl, and the protecting groups can be removed conventionally.

It will also be recognized that the protecting group present on the N-terminal  $\alpha$ -amino group may be

removed preferentially either before or simultaneous with the cleavage of the protected peptide from the resin support.

5 The A and B chains of the insulin analogs of the present invention can also be prepared via recombinant DNA methodology. In their preparation, a nucleotide sequence coding for the desired peptide of the A or B chain is prepared using now-routine techniques for such synthesis. These methods generally  
10 involve preparation of oligonucleotides coding both for fragments of the desired coding sequence and for the complementary sequence thereof. The oligonucleotides are designed to provide overlap of one fragment of the coding sequence with two fragments of the complementary  
15 sequence and vice versa. The oligonucleotides are paired and joined, ultimately producing the desired gene sequence.

The sequence is inserted into a cloning vector at a location which permits the peptide product for  
20 which it codes to be expressed. A suitable cloning vector contains at least a portion of a gene's expression control sequence.

The A and B chains of the insulin analogs of the present invention may also be prepared via a pro-  
25 insulin-like precursor molecule using recombinant DNA techniques. See Frank et al., Peptides: Synthesis-Structure-Function, Proc. Seventh Am. Pept. Symp., Eds. D. Rich and E. Gross (1981) which is incorporated herein by reference.

30 The step of combining the individual A and B chains, however produced, may be achieved by the method

of Chance et al., Peptides: Synthesis, Structure and Function: Proc. of Seventh American Peptide Symposium (1981) which is incorporated herein by reference.

5 The following examples are provided as a means of illustrating the present invention and are not to be construed as a limitation thereon.

Example 1

Lys(B28), Pro(B29) Human Insulin

10

A. Preparation of Recombinant-Derived A Chain

Human insulin A-chain was prepared via recombinant DNA technology from the chemical synthesis of the  
15 gene encoding the A-chain and the expression thereof in E. coli. Briefly, the gene for the A-chain was synthesized from various trinucleotides into synthetic fragments, deca- to pentadecanucleotides in length, by a block phosphotriester method. The gene had single-  
20 stranded cohesive termini for the Eco RI and Bam HI restriction endonucleases. Insertion thereof into an appropriate expression vector containing the  $\beta$ -galactosidase gene ( $\beta$ -gal) rendered a chimeric plasmid having the A-chain linked to the  $\beta$ -gal gene via a  
25 methionine codon. Preferably, the tryptophan synthetase gene (trp LE') is used as a promoter rather than the  $\beta$ -gal gene to achieve higher levels of expression. The chimeric plasmid was transformed into E. coli resulting in the expression of a precursor protein,  
30 i.e.,  $\beta$ -gal-met-A-chain (or trp LE'-met-A-chain when the trp LE' promoter system is used). Treatment of the

precursor protein with cyanogen bromide cleaved the methionine bond to render, after purification, the human insulin A-chain. Oxidative sulfitolysis rendered the S-sulfonated A-chain which was utilized for combination  
5 with the B-chain (S-sulfonate) as described, infra.

For complete details on the chemical synthesis of the genes for the human insulin A-chain see Crea et al., Proc. Natl. Acad. Sci. USA, 75, 5765-5769, 1978 and references cited therein which are incorporated herein  
10 by reference. For complete details of the expression in E. coli of the chemically synthesized gene for the human insulin A-chain see Goedel et al., Proc. Natl. Acad. Sci. USA, 76, 101-110, 1979 and references cited therein which are incorporated herein by reference.

15

#### B. Preparation of B Chain Analog [Lys(B28), Pro(B29)]

An Applied Biosystems 430A peptide synthesizer (including software revision 1.4) was used to prepare  
20 a crude peptidyl resin. 0.5 millimoles (mMol) of the starting solid phase resin (t-BOC-Thr(Bzl)OCH<sub>2</sub> Pam resin) was used (.76 mMol/g x .658 g). All amino acids used were BOC protected and, except for glutamic acid and histidine, all were used directly as received (i.e.,  
25 in cartridges from Applied Biosystems, Inc., each cartridge contained approximately 2 mmol of protected amino acid). Glutamic acid and histidine were obtained from Peptides International Corporation and transferred to cartridges such that each cartridge contained approx-  
30 imately 2 mmol of the desired protected amino acid. After drying of the crude peptidyl resin (under vacuum

at room temperature overnight) its weight was determined and compared to the starting weight to assure reasonable weight gain. A small portion of sample was submitted for amino acid analysis to ensure that the desired amino acids were added in the correct amounts.

The peptide was cleaved from the peptidyl resin and side-chain deprotected by stirring for approximately 1 hour at 0°C in a solution of 10 parts (v/w) HF (containing 5% v/v ethyl mercaptan and 5% v/v m-cresol) to 1 part peptidyl resin. After removal of most of the HF by vacuum the peptide was precipitated in ethyl ether. After several rinses with ethyl ether followed by vacuum filtration, the peptide was dissolved in approximately 200 ml of 7M deionized urea containing 0.1M TRIS, 0.1M Na<sub>2</sub>SO<sub>3</sub> and 0.01M Na<sub>2</sub>S<sub>4</sub>O<sub>6</sub>. The solution was adjusted to pH 8.5 with 5N NaOH and allowed to stir vigorously overnight at 4°C.

The resulting S-sulfonated peptide solution was loaded onto a 5 x 215 cm column of Sephadex G-25 (Fine) at room temperature. The sample was eluted at 20 ml/min at room temperature using 50 mM ammonium bicarbonate. The effluent was monitored at 276 nm. 20 ml fractions were collected and a pool of the desired fractions was made and further purified by high performance liquid chromatography (HPLC) as follows.

The pool of the desired fractions was pumped onto a 2.5 x 30 cm DuPont C8, 9-12μ HPLC column and eluted using a linear gradient of increasing acetonitrile in 100 mM ammonium bicarbonate at room temperature (2.6 ml/min). The effluent was monitored at 280 nm. 25 ml fractions were collected. Analytical HPLC

analyses were conducted on selected fractions to determine which fractions to retain. The desired fractions were pooled and lyophilized and used in the following combination with the A chain prepared as described above.

### C. Preparation of Lys(B28), Pro(B29) Human Insulin

The combination of the A and B chain was accomplished by the procedure of Chance et al., supra. 700 mg of the recombinant DNA-derived A-chain S-sulfonate and 140 mg of the synthetic Lys(B28), Pro(B29) B-chain S-sulfonate (both obtained as described above) were each dissolved in 70 ml and 14 ml, respectively, of 0.1M glycine buffer at ambient temperature, each adjusted to pH 10.5 with 5N NaOH and then cooled to 5°C. 6 ml of a dithiothreitol (DTT) solution at 10.5 mg/ml was prepared in 0.1M glycine buffer at ambient temperature, adjusted to pH 10.5 with 5N NaOH and then cooled to 5°C.

The A- and B-chain solutions were combined, then 5.21 ml of the DTT solution was quickly added ( $\text{SH}/\text{SSO}_3^- = 0.90$ ). The reaction solution was stirred at 5°C in an open 200 ml glass centrifuge bottle for 2.5 hours at 5°C. 45 ml of glacial acetic acid was added and the solution was allowed to stand at 5°C overnight.

The resultant precipitated mixture was centrifuged for 20 minutes at 2000 rpm at 5°C. The supernatant was combined with a 1M acetic acid wash of the pellet and placed on a 5 x 200 cm Sephadex G-50 (superfine) column in molar acetic acid at 5°C and eluted by

gravity. Twenty-minute fractions were collected for three days. The fractions were examined at 276 nm and some by analytical HPLC. The fractions containing the Lys(B28), Pro(B29) sequence of the insulin analog were pooled and lyophilized giving a 125 mg sample. This sample was further purified by reverse-phase HPLC (using a 2.12 x 25 cm DuPont C8 column eluted at room temperature, at 2.6 ml/min, using a linear gradient of increasing acetonitrile in 0.1M NaH<sub>2</sub>PO<sub>4</sub>, pH 2.2). The effluent was monitored at 276 nm. Selected fractions were assayed by analytical HPLC. The desired fractions were pooled and further purified using pH 7 HPLC as follows.

The pool from the low pH HPLC preparation run was diluted approximately 2 times in an ice bath with 0.1M (NH<sub>4</sub>)<sub>2</sub>HPO<sub>4</sub>. The pH was adjusted to 7 with cold 2N NaOH in an ice bath. The sample was loaded onto and eluted from the same HPLC column using the same conditions as the low pH preparation run except the eluting buffer was 0.1M, pH 7 (NH<sub>4</sub>)<sub>2</sub>HPO<sub>4</sub>/acetonitrile.

The pool from the pH 7 HPLC preparative run was chilled in an ice bath and diluted two times with 0.1% aqueous trifluoroacetic acid (TFA). 1N HCl was added (cold, sample in ice bath) to lower the pH to 3. The sample was loaded onto a Vydac C4 or, alternatively a DuPont C8 HPLC column (2.12 x 25 cm) and eluted with a linear gradient of increasing acetonitrile in 0.1% aqueous TFA. The effluent was monitored at 214 nm or 276 nm. The desired fractions were pooled and lyophilized giving a sample yield of 41 mg of the desired analog of greater than 97 percent purity by reverse-phase HPLC.



Example 2Lys(B28), Pro(B29) Human Insulin

5                   A second method for preparing Lys(B28),  
Pro(B29) human insulin employed enzymatic semisynthesis  
(reverse proteolysis) to combine des-octapeptide insulin  
(A<sub>1-21</sub> - B<sub>1-22</sub>) with a synthetic octapeptide. The  
des-octapeptide insulin was obtained from a tryptic  
10 digest of natural pork or human insulin as described by  
Bromer and Chance, "Preparation and Characterization of  
Des-octapeptide-Insulin, Biochim. Biophys. Acta, 133:219-  
223 (1967) which is incorporated herein by reference.  
The synthetic octapeptide, Gly-Phe-Phe-Tyr-Thr-Lys-  
15 Pro-Thr, was prepared by automatic solid-phase synthesis  
in the manner described above.

Des-octapeptide insulin (435 mg.) and 465 mg.  
of synthetic Gly-Phe-Phe-Tyr-Thr-Lys-Pro-Thr were  
combined in 15 ml of a solution containing 1 part  
20 dimethyl sulfoxide, 2 parts 1,4-butanediol and 1 part  
0.25M tris acetate buffer pH 7.3. The peptides were  
completely dissolved by warming the solution on a hot  
plate. The solution was then incubated at 37°C and  
90 mg of pork trypsin was added. The solution was  
25 stirred occasionally for 90 minutes at 37°C. The  
reaction was stopped by combining the solution with  
135 ml of 0.05N HCl.

The titled insulin analog was purified by  
loading the acidified solution containing the analog  
30 onto a 2.5 x 25 cm C-8 Zorbax HPLC column and eluting  
it with a linear gradient of increasing acetonitrile

in 0.1M sodium monobasic phosphate pH 2.2 buffer. The effluent was monitored at 276 nm. The desired fractions were pooled, diluted two-fold with water and loaded onto a 1 x 25 cm C-8 Ultrasphere HPLC column. The analog  
5 was eluted with a linear gradient of increasing acetonitrile in 0.5% aqueous TFA. The effluent was monitored at 276 nm. The desired fractions were again pooled and lyophilized to give 125 mg of the purified analog. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

MS: 5809.2 (Theory: 5808.7).

Fast Atom Bombardment - Mass Spectroscopy analyses provided herein were determined using a  
15 VG-ZAB-25E double focusing mass spectrometer at a resolution of approximately 1500. The human insulin analogs were dissolved in a mixture of glycerol and thioglycerol containing oxalic acid. Cesium iodide was used to calibrate the instrument which was scanned  
20 magnetically from m/z 5300 to m/z 6500. Resulting data are presented as average mass + 1.

### Example 3

25 Aba(B28), Pro(B29) Human Insulin

Pork des-octapeptide insulin (384 mg.) and synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Aba-Pro-Thr  
30 (362 mg.) were combined in 13 ml. of a solution containing one part dimethyl sulfoxide, two parts 1,4-butanediol, and one part 0.25M tris pH 7.3 buffer at 37°C. Pork

trypsin (75 mg.) was added. The solution was mixed well and stirred occasionally for 120 min. at 37°C.

The reaction was stopped at this time by adding the mixture to 137 ml. of 0.05N HCl. The entire solution  
5 was pumped onto a 21 x 250 mm C-8 Zorbax column, and the products were eluted in a shallow acetonitrile gradient in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by analytical HPLC, were pooled, diluted two-fold with  
10 water, and pumped onto a 25 x 300 mm C-18 Vydac column. The desalted protein was eluted from the column with an acetonitrile gradient in 0.5% trifluoroacetic acid. The fractions containing the purified insulin analog were pooled and lyophilized to yield 59 mg. The structure  
15 was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

MS: 5765.7 (Theory: 5765.6).

#### Example 4

20

#### Ala(B28), Pro(B29) Human Insulin

Pork des-octapeptide insulin (290 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Ala-Pro-Thr  
25 (310 mg.) were combined in 10 ml. of a solution containing one part dimethyl sulfoxide, two parts 1,4-butanediol, and one part 0.25M tris pH 7.3 buffer at 37°C. Pork trypsin (60 mg.) was added. The solution was mixed well and stirred occasionally for 60 min. at 37°C.

The reaction was stopped at this time by adding the mixture to 90 ml. of 0.05N HCl. The entire solution was pumped onto a 21 x 250 mm C-8 Zorbax column, and the products were eluted in a shallow acetonitrile gradient in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by analytical HPLC, were pooled, diluted two-fold with water, and pumped onto a 10 x 250 mm C-8 Ultrasphere column. The desalted protein was eluted from the column with an acetonitrile gradient in 0.5% trifluoroacetic acid. The fractions containing the purified insulin analog were pooled and lyophilized to yield 43 mg. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

MS: 5752.3 (Theory: 5751.6).

#### Example 5

##### Arg(B28), Pro(B29) Human Insulin

Pork des-octapeptide insulin (290 mg.) and synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Arg-Pro-Thr (310 mg.) were combined in 10 ml. of a solution containing one part dimethyl sulfoxide, two parts 1,4-butanediol, and one part 0.25M tris pH 7.3 buffer at 37°C. Pork trypsin (60 mg.) was added. The solution was mixed well and stirred occasionally for 60 min. at 37°C.

The reaction was stopped at this time by adding the mixture to 90 ml. of 0.05N HCl. The entire solution was pumped onto a 21 x 250 mm C-8 Zorbax column, and the products were eluted in a shallow acetonitrile gradient in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by analytical HPLC, were pooled, diluted two-fold with water, and pumped onto a 10 x 250 mm C-8 Ultrasphere column. The desalted protein was eluted from the column with an acetonitrile gradient in 0.5% trifluoroacetic acid. The fractions containing the purified insulin analog were pooled and lyophilized to yield 103 mg. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

MS: 5836.1 (Theory: 5836.7).

#### Example 6

#### Asn(B28), Pro(B29) Human Insulin

15

Pork des-octapeptide insulin (409 mg.) and synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Asn-Pro-Thr (398 mg.) were combined in 14 ml. of a solution containing one part dimethyl sulfoxide, two parts 1,4-butanediol, and one part 0.25M tris pH 7.3 buffer at 37°C. Pork trypsin (81 mg.) was added. The solution was mixed well and stirred occasionally for 120 min. at 37°C.

The reaction was stopped at this time by adding the mixture to 136 ml. of 0.05N HCl. The entire solution was pumped onto a 21 x 250 mm C-8 Zorbax column, and the products were eluted in a shallow acetonitrile gradient in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by analytical HPLC, were pooled, diluted two-fold with water, and pumped onto a 25 x 300 mm C-18 Vydac column. The desalted protein was eluted from the column with an

acetonitrile gradient in 0.5% trifluoroacetic acid. The fractions containing the purified insulin analog were pooled and lyophilized to yield 56 mg. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

MS: 5794.7 (Theory: 5794.6).

#### Example 7

#### 10                   Asp(B28), Pro(B29) Human Insulin

Pork des-octapeptide insulin (400 mg.) and synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Asp-Pro-Thr (388 mg.) were combined in 13 ml. of a solution containing one part dimethyl sulfoxide, two parts 1,4-butanediol, and one part 0.25M tris pH 7.3 buffer at 37°C. Pork trypsin (78 mg.) was added. The solution was mixed well and stirred occasionally for 120 min. at 37°C.

The reaction was stopped at this time by adding the mixture to 137 ml. of 0.05N HCl. The entire solution was pumped onto a 21 x 250 mm C-8 Zorbax column, and the products were eluted in a shallow acetonitrile gradient in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by analytical HPLC, were pooled, diluted four-fold with water, and pumped onto a 25 x 300 mm C-18 Vydac column. The desalted protein was eluted from the column with an acetonitrile gradient in 0.1% trifluoroacetic acid. The fractions containing the purified insulin analog were pooled and lyophilized to yield 85 mg. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

MS: 5795.7 (Theory: 5795.6).

Example 8Asp(B10), Lys(B28), Pro(B29) Human Insulin

## 5 A. Preparation of Asp(B10), Lys(B28), Pro(B29) Human Insulin B-Chain.

An Applied Biosystems 430A peptide synthesizer (including software revision 1.4) was used to prepare a crude peptidyl resin. 0.5 Millimoles (mMol) of the starting solid phase resin (t-BOC-Thr(Bzl)OCH<sub>2</sub> Pam resin) was used (.72 mMol/g. x .705 g.). All amino acids used were BOC protected and, except for glutamic acid, aspartic acid, and histidine, all were used directly as received (i.e., in cartridges from Applied Biosystems, Inc.; each cartridge contained approximately 2 mmol of protected amino acid). Glutamic acid, aspartic acid, and histidine were obtained from commercial sources and transferred to cartridges such that each cartridge contained approximately 2 mmol of the desired protected amino acid. The crude peptidyl resin was dried under vacuum at room temperature overnight and its weight compared to the starting weight to assure reasonable weight gain. A small portion of sample was submitted for amino acid analysis to ensure that the desired amino acids were added in the correct proportions.

The peptide was cleaved from the peptidyl resin and side-chain deprotected by stirring for approximately 1 hour at 0°C in a solution of 10 parts (v/w) HF (containing 5% v/v p-thiocresol and 5% v/v m-cresol) to 1 part peptidyl resin. After removal of most of the HF

by vacuum the peptide was precipitated in ethyl ether. After several rinses with ethyl ether followed by vacuum filtration, the peptide was dissolved in approximately 120 ml. of 8M guanidine HCl, pH 11, containing 0.1M TRIS, 5 35 mg/ml. Na<sub>2</sub>SO<sub>3</sub> and 25 mg/ml. Na<sub>2</sub>S<sub>4</sub>O<sub>6</sub>. The solution was adjusted to pH 8.8 with 5N NaOH and allowed to stir vigorously for 3 hours at room temperature.

The resulting S-sulfonated peptide solution was loaded onto a 5 x 215 cm column of Sephadex G-25 10 (medium) at room temperature. The sample was eluted at 21 ml./min. at room temperature using 50 mM ammonium bicarbonate. The effluent was monitored at 276 nm. Fractions of 25 ml. each were collected, and a pool of the desired fractions was made and further purified by 15 high performance liquid chromatography (HPLC) as follows.

The pool of the desired fractions was pumped onto a 2.5 x 30 cm DuPont C8, 9-12μ HPLC column and eluted using a linear gradient of increasing acetonitrile in 100 mM ammonium bicarbonate at room temperature (2.6 20 ml./min.). The effluent was monitored at 280 nm. Fractions (25 ml. each) were collected. Analytical HPLC analyses were conducted on selected fractions to determine which fractions to retain. The desired fractions were pooled and lyophilized and used in the following 25 combination with the A chain.

B. Combination of Asp(B10), Lys(B28), Pro(B29) Human Insulin B-Chain with Human Insulin A-Chain.

30 The combination of the A and B Chains was accomplished by the procedure of Chance et al., supra.



Two g. of the recombinant DNA-derived A-chain S-sulfonate and 400 mg. of the synthetic Asp(B10), Lys(B28), Pro(B29) B-chain S-sulfonate were each dissolved in 200 ml. and 40 ml., respectively, of 0.1M glycine buffer at ambient temperature, each adjusted to pH 10.5 with 5N NaOH and then cooled to 5°C. A dithiothreitol (DTT) solution at 15.5 mg./ml. was prepared in 0.1M glycine buffer at ambient temperature, adjusted to pH 10.5 with 5N NaOH and then cooled to 5°C.

10 The A- and B-chain solutions were combined, then 15.9 ml. of the DTT solution were quickly added (SH/SSO<sub>3</sub><sup>-</sup> = 1.0). The reaction solution was stirred at 4°C in an open 200 ml. glass centrifuge bottle for 19.6 hours at 4°C. Glacial acetic acid (129 ml.) was added, and the solution was allowed to stand at 4°C for one hour.

The resultant precipitated mixture was centrifuged for 30 minutes at 2000 rpm at 4°C. The supernatant was combined with 292 ml. milli-Q water and 73 ml. acetonitrile and further purified by reversed-phase HPLC (using a 2.5 x 30 cm Vydac C18 column eluted at room temperature, at 2.6 ml./min., using a linear gradient of increasing acetonitrile in 0.1M NaH<sub>2</sub>PO<sub>4</sub>, pH 2.1). The effluent was monitored at 280 nm. Selected fractions were assayed by analytical HPLC and the desired fractions pooled and diluted two-fold with 0.1% aqueous trifluoroacetic acid (TFA), then loaded onto an Ultrasphere octyl HPLC column (1.0 x 25 cm) and eluted with a linear gradient of increasing acetonitrile in 0.1% aqueous TFA. The effluent was monitored at 280 nm. Selected fractions were assayed by analytical HPLC and the desired fractions

pooled and repurified using the same column and conditions as above with a slightly different gradient of acetonitrile. The appropriate fractions were pooled and lyophilized giving a yield of 65 mg. of the insulin analog of greater than 93% purity by reversed-phase HPLC. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

MS: 5786.1 (Theory: 5786.7).

10

Example 9Cya(B28), Pro(B29) Human Insulin

15

Performic acid oxidation was performed to convert the cysteine of the octapeptide to the cysteic acid form. The synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Cys-Pro-Thr (363 mg.) was dissolved in 36 ml. freshly prepared performic acid in an ice-cooled flask, and allowed to stir gently for one hour. The oxidized material was diluted ten-fold with water and lyophilized. The lyophilizate was used in the semisynthesis.

20

25

Pork des-octapeptide insulin (222 mg.) and synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Cya-Pro-Thr (225 mg.) were combined in 18 ml. of a solution containing one part dimethyl sulfoxide, two parts 1,4-butanediol, and one part 0.25M tris pH 7.3 buffer at 37°C. Pork trypsin (45 mg.) was added. The solution was mixed well and stirred occasionally for 120 min. at 37°C.

The reaction was stopped at this time by adding the mixture to 242 ml. of 0.05N HCl. The entire solution was pumped onto a 21 x 250 mm C-8 Zorbax column, and the products were eluted in a shallow acetonitrile gradient in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by analytical HPLC, were pooled, diluted two-fold with water, and pumped onto a 25 x 300 mm C-18 Vydac column. The desalted protein was eluted from the column with an acetonitrile gradient in 0.5% trifluoroacetic acid. The fractions containing the purified insulin analog were pooled and lyophilized to yield 16 mg. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

MS: 5831.5 (Theory: 5831.7).

#### Example 10

##### Gln(B28), Pro(B29) Human Insulin

Pork des-octapeptide insulin (290 mg.) and synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Gln-Pro-Thr (310 mg.) were combined in 10 ml. of a solution containing one part dimethyl sulfoxide, two parts 1,4-butanediol, and one part 0.25M tris pH 7.3 buffer at 37°C. Pork trypsin (60 mg.) was added. The solution was mixed well and stirred occasionally for 60 min. at 37°C.

The reaction was stopped at this time by adding the mixture to 90 ml. of 0.05N HCl. The entire solution was pumped onto a 21 x 250 mm C-8 Zorbax column, and the products were eluted in a shallow acetonitrile gradient in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by analytical HPLC, were pooled, diluted two-fold with water, and pumped onto a 10 x 250 mm C-8 Ultrasphere column. The desalted protein was eluted from the column with an acetonitrile gradient in 0.5% trifluoroacetic acid. The fractions containing the purified insulin analog were pooled and lyophilized to yield 87 mg. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

10 MS: 5809.4 (Theory: 5808.6).

Example 11

Glu(B28), Pro(B29) Human Insulin

15

Pork des-octapeptide insulin (402 mg.) and synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Glu-Pro-Thr (398 mg.) were combined in 14 ml. of a solution containing one part dimethyl sulfoxide, two parts 1,4-butanediol, and one part 0.25M tris pH 7.3 buffer at 37°C. Pork trypsin (80 mg.) was added. The solution was mixed well and stirred occasionally for 120 min. at 37°C.

20

The reaction was stopped at this time by adding the mixture to 136 ml. of 0.05N HCl. The entire solution was pumped onto a 21 x 250 mm C-8 Zorbax column, and the products were eluted in a shallow acetonitrile gradient in 0.1 M sodium monobasic phosphate, pH 2 buffer.

25

The appropriate fractions, as determined by analytical HPLC, were pooled, diluted four-fold with water, and pumped onto a 25 x 300 mm C-18 Vydac column. The desalted protein was eluted from the column with an

30

acetonitrile gradient in 0.5% trifluoroacetic acid. The fractions containing the purified insulin analog were pooled and lyophilized to yield 59 mg. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

MS: 5809.6 (Theory: 5809.6).

#### Example 12

#### 10 Gly(B28), Pro(B29) Human Insulin

Pork des-octapeptide insulin (412 mg.) and synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Gly-Pro-Thr (376 mg.) were combined in 13 ml. of a solution containing one part dimethyl sulfoxide, two parts 1,4-butanediol, and one part 0.25M tris pH 7.3 buffer at 37°C. Pork trypsin (79 mg.) was added. The solution was mixed well and stirred occasionally for 180 min. at 37°C.

15 The reaction was stopped at this time by adding the mixture to 147 ml. of 0.05N HCl. The entire solution was pumped onto a 21 x 250 mm C-8 Zorbax column, and the products were eluted in a shallow acetonitrile gradient in 0.1 M sodium monobasic phosphate, pH 2 buffer.

20 The appropriate fractions, as determined by analytical HPLC, were pooled, diluted four-fold with water, and pumped onto a 25 x 300 mm C-18 Vydac column. The desalted protein was eluted from the column with an acetonitrile gradient in 0.1% trifluoroacetic acid. The fractions containing the purified insulin analog were pooled and lyophilized to yield 11 mg. The structure was verified by amino acid analysis (Table I) and mass spectroscopy (MS).

25 30 MS: 5737.2 (Theory: 5737.6).

Example 13His(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (400 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-His-Pro-Thr  
(398 mg.) were combined in 13 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10 trypsin (79 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

The reaction was stopped at this time by adding  
the mixture to 237 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15 products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted four-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20 The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.1% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 79 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25 spectroscopy (MS).

MS: 5816.9 (Theory: 5817.7).

Example 14Ile(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (409 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Ile-Pro-Thr  
(398 mg.) were combined in 13 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
10                   and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
trypsin (81 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 136 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20                   The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.5% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 57 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25                   spectroscopy (MS).

MS: 5793.7 (Theory: 5793.7).

Example 15Leu(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (418 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Leu-Pro-Thr  
(410 mg.) were combined in 14 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10   trypsin (83 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

The reaction was stopped at this time by adding  
the mixture to 136 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20   The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.5% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 74 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25   spectroscopy (MS).

MS: 5793.8 (Theory: 5793.7).



Example 16Nle(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (290 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Nle-Pro-Thr  
(310 mg.) were combined in 10 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10                   trypsin (60 mg.) was added. The solution was mixed well  
and stirred occasionally for 60 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 90 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
20                   water, and pumped onto a 10 x 250 mm C-8 Ultrasphere  
column. The desalted protein was eluted from the column  
with an acetonitrile gradient in 0.5% trifluoroacetic  
acid. The fractions containing the purified insulin  
analog were pooled and lyophilized to yield 54 mg. The  
structure was verified by amino acid analysis (Table I)  
25                   and mass spectroscopy (MS).

                  MS: 5794.6 (Theory: 5793.7).

Example 17D-Lys(B29) Human Insulin

5                   Pork des-octapeptide insulin (392 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Pro-D-Lys-Thr  
(387 mg.) were combined in 13.5 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10                   trypsin (78 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 136.5 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted four-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20                   The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.5% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 94 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25                   spectroscopy (MS).

                  MS: 5809.0 (Theory: 5808.7).

Example 18Met(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (350 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Met-Pro-Thr  
(366 mg.) were combined in 12 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10                   trypsin (71 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 118 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted four-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20                   The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.1% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 72 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25                   spectroscopy (MS).

MS: 5811.8 (Theory: 5811.7).

Example 19Orn(B28), Pro(B29) Human Insulin

5           Pork des-octapeptide insulin (290 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Orn-Pro-Thr  
(310 mg.) were combined in 10 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10    trypsin (60 mg.) was added. The solution was mixed well  
and stirred occasionally for 90 min. at 37°C.

          The reaction was stopped at this time by adding  
the mixture to 90 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15    products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

          The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
water, and pumped onto a 10 x 250 mm C-8 Ultrasphere  
20    column. The desalted protein was eluted from the column  
with an acetonitrile gradient in 0.5% trifluoroacetic  
acid. The fractions containing the purified insulin  
analog were pooled and lyophilized to yield 89 mg. The  
structure was verified by amino acid analysis (Table I)  
25    and mass spectroscopy (MS).

MS: 5795.2 (Theory: 5794.7).

Example 20Phe(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (290 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Phe-Pro-Thr  
(310 mg.) were combined in 10 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10                   trypsin (60 mg.) was added. The solution was mixed well  
and stirred occasionally for 80 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 90 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted four-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20                   The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.1% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 17 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25                   spectroscopy (MS).

                  MS: 5827.9 (Theory: 5827.7).

Example 21Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (339 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Pro-Pro-Thr  
(363 mg.) were combined in 9 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10                   trypsin (70 mg.) was added. The solution was mixed well  
and stirred occasionally for 80 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 108 ml. of 0.05N HCl. The entire solution  
was pumped onto a 10 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
water, and pumped onto a 10 x 250 mm C-8 Ultrasphere  
20                   column. The desalted protein was eluted from the column  
with an acetonitrile gradient in 0.5% trifluoroacetic  
acid. The fractions containing the purified insulin  
analog were pooled and lyophilized to yield 97 mg. The  
structure was verified by amino acid analysis (Table I)  
25                   and mass spectroscopy (MS).

                  MS: 5778.6 (Theory: 5777.6).

Example 22Ser(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (412 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Ser-Pro-Thr  
(390 mg.) were combined in 13 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10                   trypsin (80 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 137 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20                   The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.5% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 37 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25                   spectroscopy (MS).

                  MS: 5768.1 (Theory: 5767.6).

Example 23Thr(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (437 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Thr-Pro-Thr  
(420 mg.) were combined in 14.5 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10                   trypsin (86 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 135.5 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20                   The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.5% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 78 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25                   spectroscopy (MS).

                  MS: 5781.9 (Theory: 5781.6).



Example 24Trp(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (310 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Trp-Pro-Thr  
(325 mg.) were combined in 10.5 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10                   trypsin (64 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 140 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20                   The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.5% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 47 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25                   spectroscopy (MS).

                  MS: 5866.2 (Theory: 5866.7).

Example 25Tyr(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (391 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Tyr-Pro-Thr  
(400 mg.) were combined in 13 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10                   trypsin (79 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

                  The reaction was stopped at this time by adding  
the mixture to 137 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15                   products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

                  The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20                   The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.5% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 30 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25                   spectroscopy (MS).

                  MS: 5843.7 (Theory: 5843.7).

Example 26Val(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (400 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Val-Pro-Thr  
(383 mg.) were combined in 12 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10    trypsin (78 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

The reaction was stopped at this time by adding  
the mixture to 238 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15    products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted four-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20    The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.1% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 74 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25    spectroscopy (MS).

MS: 5780.0 (Theory: 5779.6).

Example 27Nva(B28), Pro(B29) Human Insulin

5                   Pork des-octapeptide insulin (292 mg.) and  
synthetic octapeptide Gly-Phe-Phe-Tyr-Thr-Nva-Pro-Thr  
(279 mg.) were combined in 10 ml. of a solution contain-  
ing one part dimethyl sulfoxide, two parts 1,4-butanediol,  
and one part 0.25M tris pH 7.3 buffer at 37°C. Pork  
10 trypsin (57 mg.) was added. The solution was mixed well  
and stirred occasionally for 120 min. at 37°C.

The reaction was stopped at this time by adding  
the mixture to 240 ml. of 0.05N HCl. The entire solution  
was pumped onto a 21 x 250 mm C-8 Zorbax column, and the  
15 products were eluted in a shallow acetonitrile gradient  
in 0.1 M sodium monobasic phosphate, pH 2 buffer.

The appropriate fractions, as determined by  
analytical HPLC, were pooled, diluted two-fold with  
water, and pumped onto a 25 x 300 mm C-18 Vydac column.  
20 The desalted protein was eluted from the column with an  
acetonitrile gradient in 0.5% trifluoroacetic acid. The  
fractions containing the purified insulin analog were  
pooled and lyophilized to yield 51 mg. The structure  
was verified by amino acid analysis (Table I) and mass  
25 spectroscopy (MS).

MS: 5780.0 (Theory: 5779.6).

Example 28

Utilizing the procedures set forth herein, the following additional insulin analogs are prepared:

5

(a) Asp(B1),Lys(B28),Pro(B29) Human Insulin

(b) des(Phe-B1),Lys(B28),Pro(B29) Human  
Insulin

10

(c) des(Phe-B1),Asp(B10),Lys(B28),Pro(B29)  
Human Insulin

(d) des(Phe-B1,Val-B2),Lys(B28),Pro(B29)  
Human Insulin

(e) des(Phe-B1,Val-B2),Asp(B10),Lys(B28),  
Pro(B29) Human Insulin

15

(f) Gly(A21),Asp(B10),Lys(B28),Pro(B29) Human  
Insulin

(g) Ala(A21),Asp(B10),Lys(B28),Pro(B29) Human  
Insulin

(h) des(Thr-B30),Lys(B28),Pro(B29) Human  
Insulin

20

(i) Asp(B10),Arg(B28),Pro(B29) Human Insulin

(j) Ala(A21),Arg(B28),Pro(B29) Human Insulin

(k) Asp(B1),Arg(B28),Pro(B29) Human Insulin

Example 29Lys(B28), Pro(B29) Human Insulin5           Construction of Recombinant Vectors and HostsA.   Construction of Plasmid pCZR126S1.   Isolation of Plasmid pKC283

10

Lyophils of E. coli K12 BE1201/pKC283 are obtained from the Northern Regional Research Laboratory, Peoria, Illinois 61604, under the accession number NRRL B-15830. The lyophils are decanted into tubes containing 10 ml LB medium (10 g Bacto-tryptone, 5 g Bacto-yeast extract, and 10 g NaCl per liter; pH is adjusted to 7.5) and incubated two hours at 32°C, at which time the cultures are made 50 µg/ml in ampicillin and then incubated at 32°C overnight. The E. coli K12 BE1201/pKC283 cells were cultured at 32°C, because the cells comprise a temperature-sensitive cI repressor gene integrated into the cellular DNA. When cells that comprise a wild-type lambda pL repressor gene or do not comprise a lambda pL promoter are utilized in this plasmid isolation procedure, as described in subsequent Examples herein, the temperature of incubation is 37°C.

25

A small portion of the overnight culture is placed on LB-agar (LB medium with 15 g/l Bacto-agar) plates containing 50 µg/ml ampicillin in a manner so as to obtain a single colony isolate of E. coli K12 BE1201/pKC283. The single colony obtained was inocu-

30

lated into 10 ml of LB medium containing 50 µg/ml ampicillin and incubated overnight at 32°C with vigorous shaking. The 10 ml overnight culture was inoculated into 500 ml LB medium containing 50 µg/ml ampicillin and  
5 incubated at 32°C with vigorous shaking until the culture reached stationary phase.

The following procedure is adapted from Maniatis *et al.*, 1982, Molecular Cloning (Cold Spring Harbor Laboratory).

10 The cells were harvested by centrifugation at 4000 g for 10 minutes at 4°C, and the supernatant was discarded. The cell pellet was washed in 100 ml of ice-cold STE buffer (0.1 M NaCl; 10 mM Tris-HCl, pH 7.8; and 1 mM EDTA). After washing, the cell pellet was  
15 resuspended in 10 ml of Solution 1 (50 mM glucose; 25 mM Tris-HCl, pH 8.0; and 10 mM EDTA) containing 5 mg/ml lysozyme and left at room temperature for 10 minutes. Twenty ml of Solution 2 (0.2 N NaOH and 1% SDS) were then added to the lysozyme-treated cells, and the  
20 solution was gently mixed by inversion. The mixture was incubated on ice for 10 minutes.

Fifteen ml of ice-cold 5 M potassium acetate, pH 4.8, were added to the lysed-cell mixture and the solution mixed by inversion. The solution was incubated  
25 on ice for 10 minutes. The 5 M potassium acetate solution was prepared by adding 11.5 ml of glacial acetic acid to 28.5 ml of water and 60 ml of 5 M potassium acetate; the resulting solution is 3 M with respect to potassium and 5 M with respect to acetate.

30 The lysed cell mixture was centrifuged in a Beckman SW27 (or its equivalent) at 20,000 rpm for 20

minutes at 4°C. The cell DNA and debris formed a pellet on the bottom of the tube. About 36 ml of supernatant were recovered, and 0.6 volumes of isopropanol were added, mixed, and the resulting solution left at room temperature for 15 minutes. The plasmid DNA was collected by centrifugation at 12,000 g for 30 minutes at room temperature. The supernatant was discarded, and the DNA pellet was washed with 70% ethanol at room temperature. The ethanol wash was decanted, and the pellet was dried in a vacuum desiccator. The pellet was then resuspended in 8 ml of TE buffer (10 mM Tris-HCl, pH 8.0, and 1 mM EDTA).

Eight grams of CsCl were added to the DNA solution. About 0.8 ml of a 10 mg/ml solution of ethidium bromide in water were added for each 10 ml of CsCl-DNA solution. The final density of the solution was about 1.55 g/ml, and the ethidium bromide concentration was about 600 µg/ml. The solution was transferred to a Beckman Type 50 centrifuge tube, filled to the top with paraffin oil, sealed, and centrifuged at 45,000 rpm for 24 hours at 20°C. After centrifugation, two bands of DNA were visible in ordinary light. After removing the cap from the tube, the lower DNA band was removed by using a syringe with a #21 hypodermic needle inserted through the side of the centrifuge tube.

The ethidium bromide was removed by several extractions with water-saturated 1-butanol. The CsCl was removed by dialysis against TE buffer. After extractions with buffered phenol and then chloroform, the DNA was precipitated, washed with 70% ethanol, and dried. About 1 mg of plasmid pKC283 was obtained and



stored at 4°C in TE buffer at a concentration of about 1 µg/µl. A restriction site and function map of plasmid pKC283 is presented in Figure 1 of the accompanying drawings.

5

## 2. Construction of Plasmid pKC283PX

About 10 µl of the plasmid pKC283 DNA prepared in Example 1 were mixed with 20 µl 10 X medium-salt  
10 restriction buffer (500 mM NaCl; 100 mM Tris-HCl, pH 7.5; 100 mM MgCl<sub>2</sub>; and 10 mM DTT), 20 µl 1 mg/ml BSA, 5 µl restriction enzyme PvuII (~50 Units, as defined by Bethesda Research Laboratories (BRL), from which all restriction enzymes used herein were obtained), and 145  
15 µl of water, and the resulting reaction was incubated at 37°C for 2 hours. Restriction enzyme reactions described herein were routinely terminated by phenol and then chloroform extractions, which were followed by precipitation of the DNA, an ethanol wash, and resus-  
20 pension of the DNA in TE buffer. After terminating the PvuII digestion as described above, the PvuII-digested plasmid pKC283 DNA was precipitated and then resuspended in 5 µl of TE buffer.

About 600 picomoles (pM) of XhoI linkers  
25 (5'-CCTCGAGG-3') were kinased in a mixture containing 10 µl 5 X Kinase Buffer (300 mM Tris-HCl, pH 7.8; 50 mM MgCl<sub>2</sub>; and 25 mM DTT), 5 µl 5 mM ATP, 24 µl H<sub>2</sub>O, 0.5 µl of T4 polynucleotide kinase (about 2.5 units as defined by P-L Biochemicals), 5 µl 1 mg/ml BSA, and 5 µl of  
30 10 mM spermidine by incubating the mixture at 37°C for 30 minutes.

About 12.5  $\mu$ l of the kinased XhoI linkers were added to the 5  $\mu$ l of PvuII-digested plasmid pKC283 DNA, and then 2.5  $\mu$ l of 10 X ligase buffer (300 mM Tris-HCl, pH 7.6; 100 mM MgCl<sub>2</sub>; and 50 mM DTT), 2.5  $\mu$ l of 1 mg/ml BSA, 7  $\mu$ l of 5 mM ATP, 2.5  $\mu$ l (about 2.5 units as defined by P-L Biochemicals) of T4 DNA ligase, 2.5  $\mu$ l of 10 mM spermidine, and 3  $\mu$ l of water were added to the DNA. The resulting ligation reaction was incubated at 4°C overnight. After the ligation reaction, the reaction mixture was adjusted to have the composition of high-salt buffer (0.1 M NaCl; 0.05 M Tris-HCl, pH 7.5; 10.0 mM MgCl<sub>2</sub>; and 1 mM DTT). About 10  $\mu$ l (100 units) of restriction enzyme XhoI were added to the mixture, and the resulting reaction was incubated at 37°C for 2 hours.

The reaction was terminated, and the XhoI-digested DNA was precipitated, resuspended, and ligated as described above, except that no XhoI linkers were added to the ligation mixture. The ligated DNA constituted the desired plasmid pKC283PX. A restriction site and function map of plasmid pKC283PX is presented in Figure 2 of the accompanying drawings.

### 3. Construction of *E. coli* K12 MO( $\lambda^+$ )/pKC283PX

*E. coli* K12 MO( $\lambda^+$ ) can be obtained from the Northern Regional Research Laboratories in lyophilized form under the accession number NRRL B-15993. *E. coli* K12 MO( $\lambda^+$ ) comprises the wild-type lambda pL cI repressor gene, so that transcription from the hybrid pL-lpp

promoter of the present invention does not occur in E. coli K12 MO( $\lambda^+$ ) cells. The lyophils are reconstituted, single colonies of MO( $\lambda^+$ ) are isolated, and a 10 ml overnight culture of the MO( $\lambda^+$ ) cells is prepared in  
5 substantial accordance with the procedure of Example 29A1, except that the temperature of incubation is 37°C and no ampicillin is used in the growth media.

Fifty  $\mu$ l of the overnight culture were used to inoculate 5 ml of LB media which also contained 10 mM  
10 MgSO<sub>4</sub> and 10 mM MgCl<sub>2</sub>. The culture was incubated at 37°C overnight with vigorous shaking. The following morning, the culture was diluted to 200 ml with LB media containing 10 mM MgSO<sub>4</sub> and 10 mM MgCl<sub>2</sub>. The diluted culture was incubated at 37°C with vigorous  
15 shaking until the absorbance at 550 nm ( $A_{550}$ ) was about 0.5, which indicated a cell density of about  $1 \times 10^8$  cells/ml. The culture was cooled for ten minutes in an ice-water bath, and the cells were then collected by centrifugation at 4000 g for 10 minutes at 4°C. The  
20 cell pellet was resuspended in 100 ml of cold 10 mM MgSO<sub>4</sub> and then immediately re-pelleted by centrifugation. The cell pellet was resuspended in 100 ml of 30 mM CaCl<sub>2</sub> and incubated on ice for 20 minutes.

The cells were again collected by centri-  
25 fugation and resuspended in 10 ml of 30 mM CaCl<sub>2</sub>. A one-half ml aliquot of the cells was added to the ligated DNA prepared in Example 29A2; the DNA had been made 30 mM in CaCl<sub>2</sub>. The cell-DNA mixture was incubated on ice for one hour, heat-shocked at 42°C for 90  
30 seconds, and then chilled on ice for about two minutes. The cell-DNA mixture was diluted into 10 ml of LB media

in 125 ml flasks and incubated at 37°C for one hour. One hundred  $\mu$ l aliquots were plated on LB-agar plates containing ampicillin and incubated at 37°C until colonies appeared.

5                   The colonies were individually cultured, and the plasmid DNA of the individual colonies was examined by restriction enzyme analysis and gel electrophoresis. Plasmid DNA isolation was performed on a smaller scale in accordance with the procedure of Example 29A1, but  
10 the CsCl gradient step was omitted until the desired E. coli K12 MO( $\lambda^+$ )/pKC283PX transformants were identified. A restriction site and function map of plasmid pKC283PX is presented in Figure 2 of the accompanying drawings.

15                   4. Construction of E. coli K12 MO( $\lambda^+$ )/pKC283-L

                  Ten  $\mu$ g of plasmid pKC283PX DNA prepared in accordance with the procedure of Example 29A1 were dis-  
20 solved in 20  $\mu$ l of 10X high-salt buffer, 20  $\mu$ l 1 mg/ml BSA, 5  $\mu$ l ( $\sim$ 50 units) restriction enzyme BglII, 5 $\mu$ l ( $\sim$ 50 units) restriction enzyme XhoI, and 150  $\mu$ l of water, and the resulting reaction was incubated at 37°C for two hours. The reaction was stopped, and after precip-  
25 itating the BglII-XhoI digested DNA, the DNA was resuspended in 5  $\mu$ l of TE buffer.

                  A DNA linker with single-stranded DNA ends characteristic of BglII and XhoI restriction enzyme cleavage was synthesized and kinased. The linker was  
30 kinased in substantial accordance with the procedure of Example 29A2. The DNA linker had the following structure:

5'-GATCTATTA ACTCAATCTAGAC-3'  
|||||||  
3'-ATAATTGAGTTAGATCTGAGCT-5'

5

The linker depicted above was synthesized from single-stranded deoxyoligonucleotides by procedures well known in the art. The single-stranded deoxyoligonucleotides can be synthesized with commercially available instruments, such as the 380A DNA Synthesizer marketed by Applied Biosystems (850 Lincoln Centre Drive, Foster City, CA 94404), which utilizes phosphoramidite chemistry. Other procedures for synthesizing DNA are also known in the art. The conventional modified phosphotriester method of synthesizing single stranded DNA is described in Itakura *et al.*, 1977, Science 198:1056 and in Crea *et al.*, 1978, Proc. Nat. Acad. Sci. USA 75:5765. In addition, an especially preferred method of synthesizing DNA is disclosed in Hsiung *et al.*, 1983, Nucleic Acid Research 11:3227 and Narang *et al.*, 1980, Methods in Enzymology 68:90.

The linker and *Bgl*II-*Xho*I-digested plasmid pKC283PX were ligated in substantial accordance with the procedure of Example 29A2. The ligated DNA constituted the desired plasmid pKC283-L. A restriction site and function map of plasmid pKC283-L is presented in Figure 3 of the accompanying drawings. The plasmid pKC283-L DNA was used to transform *E. coli* K12 MO( $\lambda^+$ ) and the resulting *E. coli* K12 MO( $\lambda^+$ )/pKC283-L transformants were identified in substantial accordance with the procedure of Example 29A3.

5. Construction of *E. coli* K12 MO( $\lambda^+$ )/pKC283-LB

About 10  $\mu$ g of plasmid pKC283-L DNA, prepared  
5 in substantial accordance with the procedures of Example  
29A1, were dissolved in 20  $\mu$ l 10X high-salt buffer,  
20  $\mu$ l 1 mg/ml BSA, 5  $\mu$ l ( $\sim$ 50 units) restriction enzyme  
XhoI, and 155  $\mu$ l of H<sub>2</sub>O, and the resulting reaction was  
incubated at 37°C for two hours. The XhoI-digested  
10 plasmid pKC283-L DNA was then precipitated from the  
reaction mixture by the addition of three volumes of  
95% ethanol and one-tenth volume of 3 M sodium acetate,  
incubation in a dry ice-ethanol bath for five minutes,  
and centrifugation. The resulting DNA pellet was washed  
15 with 70% ethanol, dried, and resuspended in 2  $\mu$ l 10X  
nick-translation buffer (0.5 M Tris-HCl, pH 7.2; 0.1  
M MgSO<sub>4</sub>; and 1 mM DTT), 1  $\mu$ l of a solution 2 mM in each  
of the deoxynucleotide triphosphates, 15  $\mu$ l of H<sub>2</sub>O, 1  $\mu$ l  
( $\sim$ 6 units as defined by P-L Biochemicals) of Klenow,  
20 which is the large fragment of *E. coli* DNA polymerase I,  
and 1  $\mu$ l of 1 mg/ml BSA. The resulting reaction was  
incubated at 25°C for 30 minutes; the reaction was  
stopped by incubating the solution at 70°C for five  
minutes.  
25 BamHI linkers (5'-CGGGATCCCG-3') were kinased  
and ligated to the XhoI-digested, Klenow-treated plasmid  
pKC283-L DNA in substantial accordance with the pro-  
cedure of Example 29A2. After the ligation reaction,  
the DNA was digested with about 100 units of BamHI for  
30 about 2 hours at 37°C in high-salt buffer. After the  
BamHI digestion, the DNA was prepared for ligation in  
substantial accordance with the procedure of Example 29A2.

The ~5.9 kb BamHI restriction fragment was circularized by ligation and transformed into E. coli K12 MO( $\lambda^+$ ) in substantial accordance with the procedures of Examples 29A2 and 29A3. The E. coli K12 MO( $\lambda^+$ )/pKC283-LB transformants were identified, and then plasmid pKC283-LB DNA was prepared in substantial accordance with the procedure of Example 29A1. A restriction site and function map of plasmid pKC283-LB is presented in Figure 4 of the accompanying drawings.

#### 6. Construction of E. coli K12 MO( $\lambda^+$ )/pL32

About 10  $\mu$ g of plasmid pKC283PX were digested with restriction enzyme SalI in high-salt buffer, treated with Klenow, and ligated to EcoRI linkers (5'-GAGGAATTCCTC-3') in substantial accordance with the procedure of Example 29A5, with the exception of the starting plasmid, restriction enzymes, and linkers used. After digestion with restriction enzyme EcoRI, which results in the excision of ~2.1 kb of DNA, the ~4.0 kb EcoRI restriction fragment was circularized by ligation to yield plasmid pKC283PRS. The ligated DNA was used to transform E. coli K12 MO( $\lambda^+$ ) in substantial accordance with the procedure of Example 29A3. After the E. coli K12 MO( $\lambda^+$ )/pKC283PRS transformants were identified, plasmid pKC283PRS DNA was prepared in substantial accordance with the procedure of Example 29A1. A restriction site and function map of plasmid pKC283PRS is presented in Figure 5 of the accompanying drawings.

About 10  $\mu\text{g}$  of plasmid pKC283PRS were digested in 200  $\mu\text{l}$  of high-salt buffer with about 50 units each of restriction enzymes PstI and SphI. After incubating the reaction at 37°C for about 2 hours, the reaction mixture was electrophoresed on a 0.6% low-gelling-temperature agarose (FMC Corporation, Marine Colloids Division, Rockland, Maine 04841) gel for 2-3 hours at  $\sim 130$  V and  $\sim 75$  mA in Tris-Acetate buffer.

The gel was stained in a dilute solution of ethidium bromide, and the band of DNA constituting the  $\sim 0.85$  kb PstI-SphI restriction fragment, which was visualized with long-wave UV light, was cut from the gel in a small segment. The volume of the segment was determined by weight and density of the segment, and an equal volume of 10 mM Tris-HCl, pH 7.6, was added to the tube containing the segment. The segment was then melted by incubation at 72°C. About 1  $\mu\text{g}$  of the  $\sim 0.85$  kb PstI-SphI restriction fragment of plasmid pKC283PRS was obtained in a volume of about 100  $\mu\text{l}$ . In an analogous manner, plasmid pKC283-LB was digested with restriction enzymes PstI and SphI, and the resulting  $\sim 3.0$  kb restriction fragment was isolated by agarose gel electrophoresis and prepared for ligation.

The  $\sim 0.85$  kb PstI-SphI restriction fragment of plasmid pKC283PRS was ligated to the  $\sim 3.0$  kb PstI-SphI restriction fragment of plasmid pKC283-LB in substantial accordance with the procedure of Example 29A2. The ligated DNA constituted the desired plasmid pL32. A restriction site and function map of plasmid pL32 is presented in Figure 6 of the accompanying drawings. Plasmid pL32 was transformed into E. coli K12 MO( $\lambda^+$ ) cells in substantial



accordance with the procedure of Example 29A3. Plasmid pL32 DNA was prepared from the E. coli K12 MO( $\lambda^+$ )/pL32 transformants in substantial accordance with the procedure of Example 29A1. Analysis of the plasmid pL32 DNA demonstrated that more than one EcoRI linker attached to the Klenow-treated, SalI ends of plasmid pKC283PX. The presence of more than one EcoRI linker does not affect the utility of plasmid pL32 or derivatives of plasmid pL32 and can be detected by the presence of an XhoI restriction site, which is generated whenever two of the EcoRI linkers are ligated together. Alternatively, plasmid pL32 may be constructed by carrying out the SalI-EcoRI excision and ligation of the first paragraph of this Example upon plasmid pKC283-LB.

15

7. Construction of E. coli K12 MO( $\lambda^+$ )/pL47

E. coli K12 RV308/pNM789 can be obtained from the Northern Regional Research Laboratories in lyophilized form under the accession number NRRL B-18216. A restriction site and function map of pNM789 is presented in Figure 7 of the accompanying drawings. Plasmid DNA is extracted from the culture in substantial accordance with the teaching of Example 1, except that the temperature of incubation is 37°C. Ten micrograms of pNM789 are suspended in 200  $\mu$ l PvuII buffer (50 mM Tris-HCl (pH 7.5), 60 mM NaCl and 6mM MgCl<sub>2</sub>). One unit of PvuII is added and the reaction mix is incubated for 5 minutes at 37°C. The enzyme is inactivated by heating 10 minutes at 65°C. 30  $\mu$ l of 10X BamHI buffer (200 mM

20  
25  
30

Tris-HCl (pH 8.0), 1M NaCl and 70 mM MgCl<sub>2</sub>), 70 µl H<sub>2</sub>O and 10 units of BamHI are next added and the reaction is incubated for 1 hour at 37°C. This is followed by the addition of 5 units of alkaline phosphatase and  
5 incubation for 1 hour at 65°C. The DNA fragments are separated on a 1 percent agarose gel, and a DNA fragment (Figure 8) the size of a single cut fragment is purified.

A DNA linker with a blunt end and a BamHI end is synthesized in substantial accordance with the teaching of Example 29A4. This linker (shown at 118 in Figure 8) has the following structure:

15                   5'-CTGTGCCTTCTAG-3'  
                      |||||||  
                      3'-GACACGGAAGATCCTAG-5'

The linker is kinased and ligated into the BamHI-PvuII digested plasmid pNM789 in substantial accordance with the teaching of Example 29A2. This ligation mixture is  
20 used to transform E. coli K12 RV308 cells and plasmid isolation is performed upon these transformants in substantial accordance with the teaching of Example 29A3. Several plasmids are selected which contain the appropriate size PvuII fragment (494bp) and XbaI-BamHI  
25 fragment (628bp). The sequence of at least two of these is determined by sequencing from the BamHI site toward the unique SmaI site and one clone is selected with the desired sequence. This intermediate plasmid is designated plasmid 120. A schematic outline of this  
30 procedure and a restriction site and function map of plasmid 120 is presented in Figure 8 of the accompanying drawings.

To isolate the EK-BGH-encoding DNA, about 10  $\mu$ g of plasmid 120 were digested in 200  $\mu$ l of high-salt buffer containing about 50 units each of restriction enzymes XbaI and BamHI. The digestion products were separated by agarose gel electrophoresis, and the  $\sim$ 0.6 kb XbaI-BamHI restriction fragment which encodes EK-BGH was isolated and prepared for ligation in substantial accordance with the procedure of Example 29A6.

Plasmid pL32 was also digested with restriction enzymes XbaI and BamHI, and the  $\sim$ 3.9 kb restriction fragment was isolated and prepared for ligation. The  $\sim$ 3.9 kb XbaI-BamHI restriction fragment of plasmid pL32 was ligated to the  $\sim$ 0.6 kb XbaI-BamHI restriction fragment of plasmid 120 in substantial accordance with the procedure of Example 29A2 to yield plasmid pL47. A restriction site and function map of plasmid pL47 is presented in Figure 9 of the accompanying drawings. Plasmid pL47 was transformed into E. coli K12 MO( $\lambda^+$ ) in substantial accordance with the procedure of Example 29A3, and the E. coli K12 MO( $\lambda^+$ )/pL47 transformants were identified. Plasmid pL47 DNA was prepared from the transformants in substantial accordance with the procedures of Example 29A1.

25

#### 8. Construction of E. coli K12 RV308/pPR12AR1

Plasmid pPR12 comprises the temperature-sensitive pL repressor gene cI857 and the plasmid pBR322 tetracycline resistance-conferring gene. Plasmid pPR12

30

is disclosed and claimed in U.S. Patent #4,436,815, issued 13 March 1984. A restriction site and function map of plasmid pPR12 is presented in Figure 10 of the accompanying drawings.

5           About 10  $\mu$ g of plasmid pPR12 were digested with about 50 units of restriction enzyme EcoRI in 200  $\mu$ l of high-salt buffer at 37°C for two hours. The EcoRI-digested plasmid pPR12 DNA was precipitated and treated with Klenow in substantial accordance with the  
10           procedure of Example 29A5. After the Klenow reaction, the EcoRI-digested, Klenow-treated plasmid pPR12 DNA was recircularized by ligation in substantial accordance with the procedure of Example 29A2. The ligated DNA, which constituted the desired plasmid pPR12 $\Delta$ R1, was used  
15           to transform E. coli K12 RV308 in substantial accordance with the procedure of Example 29A3, except that selection was based on tetracycline (5  $\mu$ g/ml) resistance, not ampicillin resistance. E. coli K12 RV308 is available from the NRRL under the  
20           accession number NRRL B-15624. After the E. coli K12 RV308/pPR12 $\Delta$ R1 transformants were identified, plasmid pPR12 $\Delta$ R1 DNA was prepared from the transformants in substantial accordance with the procedure of Example 29A11.  
25           About 10  $\mu$ g of plasmid pPR12 $\Delta$ R1 were digested with about 50 units of restriction enzyme AvaI in 200  $\mu$ l of medium-salt buffer at 37°C for 2 hours. The AvaI-digested plasmid pPR12 $\Delta$ R1 DNA was precipitated and treated with Klenow in substantial accordance with the  
30           procedure of Example 29A5. After the Klenow reaction, the AvaI-digested, Klenow-treated plasmid pPR12 $\Delta$ R1 DNA was

ligated to EcoRI linkers (5'-GAGGAATTCCTC-3') in substantial accordance with the procedure of Example 29A2. After the linker ligation, the DNA was precipitated and then resuspended in about 200  $\mu$ l of high-salt buffer  
5 containing about 50 units of restriction enzyme EcoRI. The resulting reaction was incubated at 37°C for about 2 hours. After the EcoRI digestion, the reaction mixture was loaded onto an agarose gel, and the  $\sim$ 5.1 kb  
10 EcoRI restriction fragment was purified in substantial accordance with the procedure of Example 29A6. The  $\sim$ 5.1 kb EcoRI restriction fragment was recircularized by ligation in substantial accordance with the procedure of Example 29A2. The ligated DNA constituted the desired plasmid pPR12AR1. The plasmid pPR12AR1 DNA was transformed into  
15 E. coli K12 RV308 in substantial accordance with the procedure of Example 29A3, except that selection was based on tetracycline resistance, not ampicillin resistance. After identifying the E. coli K12 RV308/pPR12AR1 transformants, plasmid pPR12AR1 DNA was prepared in substantial accordance with the procedure of Example 29A1.  
20 A restriction site and function map of plasmid pPR12AR1 is presented in Figure 11 of the accompanying drawings.

25 9. Construction of E. coli K12 RV308/pL110

About 10  $\mu$ g of plasmid pPR12AR1 DNA were suspended in about 200 ml of high-salt buffer containing about 50 units each of restriction enzymes PstI and  
30 EcoRI, and the digestion reaction was incubated at 37°C for about 2 hours. The reaction mixture was then loaded

onto an agarose gel, and the ~2.9 kb PstI-EcoRI restriction fragment of plasmid pPR12AR1 was isolated and prepared for ligation in substantial accordance with the procedure of Example 29A6.

5           About 10 ug of plasmid pL47 were digested with restriction enzymes PstI and BamHI in 200 ul of high-salt buffer at 37°C for two hours. The PstI-BamHI-digested DNA was loaded onto an agarose gel, and the ~2.7 kb PstI-BamHI restriction fragment that comprised the  
10           origin of replication and a portion of the ampicillin resistance-conferring gene was isolated and prepared for ligation in substantial accordance with the procedure of Example 29A6. In a separate reaction, about 10 ug of plasmid pL47 DNA were digested with restriction enzymes  
15           EcoRI and BamHI in 200 ul of high-salt buffer at 37°C for two hours, and the ~1.03 kb EcoRI-BamHI restriction fragment that comprised the novel transcriptional and translational activating sequence and the EK-BGH-encoding DNA was isolated and prepared for ligation in  
20           substantial accordance with the procedure of Example 29A6. The ~2 ug of the ~1.03 kb EcoRI-BamHI restriction fragment obtained were used in the construction of plasmid pL110.

25           The ~2.7 kb PstI-BamHI and ~1.03 kb EcoRI-BamHI restriction fragments of plasmid pL47 were ligated to the ~2.9 kb PstI-EcoRI restriction fragment of plasmid pPR12AR1 to construct plasmid pL110, and the  
30           ligated DNA was used to transform E. coli K12 RV308 in substantial accordance with the procedure of Examples 29A2 and 29A3, except that tetracycline resistance, not ampicillin resistance, was used as the basis for selecting transformants.

Two PstI restriction enzyme recognition sites are present in the EK-BGH coding region that are not depicted in the restriction site and function maps presented in the accompanying drawings. A restriction site and function map of plasmid pL110 is presented in Figure 12 of the accompanying drawings.

10. Construction of *E. coli* K12 RV308/pL110C

10 a. Construction of *E. coli* K12 RV308/pL110A

About 1  $\mu$ g of plasmid pL110 DNA was digested with restriction enzyme NdeI in 20  $\mu$ l total volume containing 2  $\mu$ l of 10X high-salt buffer (1.0 M NaCl; 0.50 M Tris-HCl, pH=7.5; 0.10 M MgCl<sub>2</sub>; and 10 mM dithiothreitol) and 3 units of NdeI enzyme for 1 hour at 37°C. The reaction mixture was extracted with phenol/chloroform and the DNA precipitated with ethanol. The NdeI-digested plasmid pL110 DNA was dissolved in 50  $\mu$ l of 1X Klenow buffer (40 mM KPO<sub>4</sub>, pH=7.5; 6.6 mM MgCl<sub>2</sub>; 1.0 mM 2-mercaptoethanol; 33  $\mu$ M dATP; 33  $\mu$ M dCTP; 33  $\mu$ M dGTP; and 33  $\mu$ M TTP). Two  $\mu$ l (~10 units, New England Biolabs) of the large fragment of *E. coli* DNA polymerase I, known as Klenow, were added to and mixed with the DNA, and the resulting reaction was incubated at 16°C for 1 hour. The reaction was terminated by phenol extraction and the DNA conventionally purified. The NdeI-digested, Klenow-treated DNA was then ligated with T4 DNA ligase at 4°C for 16 hours. The resulting DNA was used to conventionally transform *E. coli* K12 strain RV308 (NRRL B-15624). Transformants were selected on

L-agar plates containing 100 µg/ml ampicillin and plasmids isolated from resistant colonies by the rapid alkaline extraction procedure described by Birnboim and Doly. A plasmid (pL110A in Figure 13) lacking an NdeI site was selected.

b. Construction of Phage pL110B by Site-Specific Mutagenesis

The protocol for eliminating the BamHI site in the tetracycline resistance-conferring gene by site-specific mutagenesis is shown on the right hand side of Figure 13 of the accompanying drawings.

b(i) Construction of Phage M13Tc3

15

Plasmid pL110 served as the source of the tetracycline resistance-conferring gene. About 50 µg of plasmid pL110 in 50 µl of TE buffer were added to 25 µl of 10X HindIII buffer and 170 µl of H<sub>2</sub>O. About 5 µl (~50 units) of restriction enzyme HindIII were added to the solution of plasmid pL110 DNA, and the resulting reaction was incubated at 37°C for 2 hours. About 13 µl of 2 M Tris·HCl, pH=7.4, and 5 µl (~50 units) of restriction enzyme EcoRI were added to the HindIII-digested plasmid pL110 DNA, and the reaction was incubated for 2 more hours at 37°C. The reaction was stopped by extracting the reaction mixture with TE-saturated phenol; the phenol was removed by chloroform extractions. The EcoRI-HindIII-digested plasmid pL110 DNA was then collected by precipitation and centrifugation, loaded into a 1% agarose gel, and the large



~4.3 kb EcoRI-HindIII restriction fragment was isolated and purified.

5 About 5  $\mu$ g of phage m13mp18 (New England Biolabs) were dissolved in 50  $\mu$ l of TE buffer and then digested with HindIII and EcoRI as described above. The HindIII-EcoRI-cut phage M13mp18 DNA was purified as described for pL110 except that an ~7.25 kb restriction fragment was isolated and purified.

10 About 100 nanograms of the ~4.3 kb HindIII-EcoRI fragment of plasmid pL110 were mixed with about 100 nanograms of the ~7.25 kb HindIII-EcoRI fragment of phage M13mp18, 2  $\mu$ l of 10X ligase buffer, 1  $\mu$ l (~100 units) of T4 DNA ligase, and 14  $\mu$ l of H<sub>2</sub>O. The ligation reaction was incubated at 15°C for 1.5 hours; the  
15 ligated DNA constituted the desired phage m13Tc3 DNA. A restriction site and function map of phage m13Tc3 is presented in Figure 13 of the accompanying drawings.

One ml of an overnight culture of E. coli K12 JM109 (E. coli K12 JM101, available from New England  
20 Biolabs, can be used instead of E. coli K12 JM109) was used to inoculate 50 ml of L broth, and the resulting culture was incubated at 37°C with aeration until the O.D.<sub>660</sub> was between 0.3 and 0.4. The cells were resuspended in 25 ml of 10 mM NaCl, incubated on ice for  
25 10 minutes, and collected by centrifugation. The cells were resuspended in 1.25 ml of 75 mM CaCl<sub>2</sub>; a 200  $\mu$ l aliquot of the cells was removed, added to 10  $\mu$ l of the ligated DNA prepared above, and incubated on ice for  
30 about 40 minutes. The cell-DNA mixture was then incubated at 42°C for 2 minutes, and varying aliquots (1, 10, and 100  $\mu$ l) were removed and added to 3 ml of top

agar (L broth with 0.5% agar kept molten at 45°C) that also contained 50 µl of 2% X-Gal, 50 µl of 100 mM IPTG, and 200 µl of E. coli K12 JM109 in logarithmic growth phase. The cell-top agar mixture was then plated on  
5 L-agar plates containing 40 µg/ml X-Gal (5-bromo-4-chloro-3-indolyl-β-D-thiogalactoside) and 0.1 mM IPTG (isopropyl-β-D-thiogalactoside), and the plates were incubated at 37°C overnight.

The following morning, several clear, as  
10 opposed to blue, plaques were individually used to inoculate 2 ml of L broth, and the resulting cultures were incubated at 37°C with aeration for 2 hours. The absence of blue color indicates the desired DNA insertion occurred. Then, the cultures were centrifuged, and  
15 200 µl of the resulting supernatant were added to 10 ml cultures (O.D.<sub>550</sub> = 0.5) of E. coli K12 JM109 growing at 37°C with aeration. These cultures were incubated for another 30 minutes at 37°C; then, the cells were pelleted by centrifugation and used to prepare the replicative-  
20 form of the recombinant phage they contained. Double-stranded, replicative form phage DNA was isolated from the cells using a scaled-down version of the procedure described in Example 1. Transformants containing phage m13Tc3 DNA were identified by restriction enzyme  
25 analysis of their phage DNA.

b(ii) Preparation of Single-Stranded Phage m13Tc3 DNA

One and one-half ml of an overnight culture  
30 of E. coli K12 JM109/m13Tc3 were centrifuged, and 100 µl of the phage m13Tc3-containing supernatant were

used to inoculate a 25 ml culture of E. coli JM109 at an O.D.<sub>680</sub> of about 0.4-0.5. The culture was incubated for 6 hours at 37°C with aeration, at which time the culture was centrifuged and the resulting supernatant, about 20 ml, transferred to a new tube. About 2 ml of a solution containing 20% polyethylene glycol (PEG) 6000 and 14.6% NaCl were added to the supernatant, which was then incubated on ice for 20 minutes.

The supernatant was centrifuged for 25 minutes at 7000 rpm, and the resulting pellet, which contained single-stranded phage ml3Tc3 DNA, was resuspended in 500 µl of TE buffer. The DNA solution was extracted twice with TE-saturated phenol and twice with chloroform. The single-stranded DNA was then precipitated using NaOAc and ethanol and centrifuged. The resulting pellet was washed with 70% ethanol, dried, and then dissolved in 60 µl of H<sub>2</sub>O.

b(iii) Mutagenesis

The single-stranded DNA fragment used in the mutagenesis was synthesized on an automated DNA synthesizer. The fragment has the sequence, 5'-CCCGTCCTGTGGATACTCTACGCCGA-3', and is homologous to the region surrounding the BamHI site (5'-GGATCC-3') in the tetracycline resistance-conferring gene from plasmid pBR322, except that the A residue second from the 5' end (or third from the 3' end) is a C in plasmid pBR322. This change does not alter the amino acid composition of the tetracycline resistance-conferring protein but eliminates the BamHI site.

About 10 picomoles of the mutagenic primer and the M13 universal primer (Bethesda Research Laboratories (BRL), P.O. Box 6009, Gaithersburg, MD 20760) were individually treated with 10 units (BRL) of T4 polynucleotide kinase in 20  $\mu$ l of 1X kinase buffer (60 mM Tris-HCl, pH = 7.8; 15 mM 2-mercaptoethanol; 10 mM MgCl<sub>2</sub>; and 0.41  $\mu$ M ATP) for 30 minutes at 37°C. The kinase-treated DNAs were used in the mutagenesis procedure described below.

10 The annealing reaction was carried out mixing together 300 nanograms (1.2  $\mu$ l) of single-stranded phage m13Tc3, 1 picomole (2  $\mu$ l) of the universal primer, 1 picomole (2  $\mu$ l) of the mutagenic primer, 2  $\mu$ l of 10X annealing buffer (100 mM Tris-HCl, pH=7.5; 1 mM EDTA; 15 and 500 mM NaCl), and 12.8  $\mu$ l of H<sub>2</sub>O. The reaction was incubated at 80°C for 2 minutes, at 50°C for 5 minutes, and then allowed to cool to room temperature.

The extension reaction was carried out by adding 5  $\mu$ l of 10X extension buffer (500 mM Tris-HCl, pH=8; 1 mM EDTA; and 120 mM MgCl<sub>2</sub>); 5  $\mu$ l of 2 mM dATP; 1  $\mu$ l of a solution 6 mM in each of dGTP, TTP, and dCTP; 1  $\mu$ l (~2 units, Pharmacia P-L Biochemicals, 800 Centennial Avenue, Piscataway, NJ 08854) of Klenow enzyme; 1  $\mu$ l (100 units) of T4 DNA ligase; and 17  $\mu$ l of 25 H<sub>2</sub>O to the mixture of annealed DNA. The extension reaction was incubated at room temperature for 1 hour, then at 37°C for 2.5 hours, and then overnight at 4°C.

The reaction was stopped by two extractions with TE-saturated phenol, which were followed by two 30 extractions with CHCl<sub>3</sub>. The DNA was precipitated with ethanol and NaOAc. The DNA was collected by centri-

fugation and resuspended in 50  $\mu$ l of H<sub>2</sub>O, and 6  $\mu$ l of 10X S1 buffer were then added to the solution of DNA.

The solution of DNA was split equally into three tubes. About 200 units (Miles Laboratories) of S1  
5 nuclease were added to two of the tubes. One S1 reaction was incubated at room temperature for 5 minutes, the other for 10 minutes. The reactions were stopped by extracting the reaction mixture twice with TE-saturated phenol. The phenol extractions were followed by two  
10 extractions with chloroform; then, the DNA was precipitated from the reaction mixture with NaOAc and ethanol. The untreated sample of DNA served as a negative control. The S1-treated samples were kept separate from each other throughout the remainder of the procedure but  
15 gave similar results.

The DNA pellets were resuspended in 20  $\mu$ l of H<sub>2</sub>O, and 10  $\mu$ l of the resulting solution were used to transform E. coli K12 JM109 (E. coli K12 JM101 could also be used) in accordance with the procedure used  
20 during the construction of phage m13Tc3, except that no IPTG or X-Gal was added to the plates.

Double-stranded replicative form DNA from about 48 plaques was isolated as described above and screened for the presence of a BamHI restriction site.  
25 Isolates without a BamHI site were further screened by preparing single-stranded DNA as described above. The single-stranded DNA was sequenced using the dideoxy sequencing method (J.H. Smith, 1980, Methods in Enzymology 65: 560-580). The desired isolate was  
30 designated pL110B (Figure 13).

c. Construction of Plasmid pL110C

About 50  $\mu$ g of the replicative form of phage pL110B DNA were digested in 250  $\mu$ l of 1X NheI buffer (50 mM NaCl; 6 mM Tris·HCl, pH=7.5; 6 mM MgCl<sub>2</sub>; and 6 mM  $\beta$ -mercaptoethanol) containing ~50 units of NheI restriction enzyme at 37°C for 2 hours. Five  $\mu$ l of 5 M NaCl were then added to the NheI-digested phage pL110B DNA, followed by 5  $\mu$ l (~50 units) of SalI restriction enzyme. Digestion was continued for 2 hours at 37°C. The desired ~422 bp NheI-SalI fragment containing the mutated region of the tetracycline resistance-conferring gene was then isolated from an acrylamide gel, according to well known standard procedures.

Plasmid pL110A DNA was digested with NheI and SalI under identical conditions, except that plasmid pL110A was substituted for phage pL110B. The ~6.1 kb NheI-SalI restriction fragment of plasmid pL110A was purified from agarose.

The desired plasmid pL110C was constructed by ligating together 100 nanograms each of the NheI-SalI fragments of pL110A (~6.1 kb) and pL110B (~422 bp) using conventional procedures. A restriction site and function map of plasmid pL110C is presented in Figure 13 of the accompanying drawings. The desired plasmid pL110C confers tetracycline resistance to 10  $\mu$ g/ml tetracycline in E. coli but lacks a BamHI site in the tetracycline resistance-conferring gene.

11. Construction of Plasmid pCZR111

Plasmid pL110C contains a single ClaI restriction site which was removed by running the following reactions. About 1  $\mu$ g of plasmid pL110C was digested with ClaI in substantial accordance with the teaching of Example 29A2, except restriction enzyme ClaI and 10X ClaI Buffer (500 mM NaCl, 100 mM Tris-HCl (pH 7.9) and 100 mM MgCl<sub>2</sub>) were used. The ClaI-digested DNA was then treated with Klenow in substantial accordance with the teaching of Example 29A5, except only dCTP, rather than all four dNTPs, was added.

The DNA was then precipitated and resuspended in 50  $\mu$ l of Mung Bean Nuclease Buffer (50 mM Sodium Acetate (pH 5.0), 30 mM NaCl and 1 mM ZnSO<sub>4</sub>). One unit of Mung Bean Nuclease (commercially available from New England Biolabs) was added and the reaction was incubated at 30°C for 30 minutes. The tube was then placed in ice and NaCl was added to 0.2 M, then the mixture was phenol/chloroform extracted, ethanol precipitated and resuspended in 10 mM Tris-HCl (pH 8.0). The DNA was then self-ligated and transformed into E. coli cells in substantial accordance with the teaching of Examples 29A3 and 29A4. The resultant plasmid was designated plasmid pCZR111.

12. Construction of Plasmid pCZR126S

About 26  $\mu$ g of plasmid pCZR111 was digested with XbaI as follows. 10X XbaI buffer consists of 600mM Tris-HCl, 100mM MgCl<sub>2</sub>, 1 M NaCl, and 10mM

2-mercaptoethanol, pH 7.5 (at 37°C). 50 ul of 10X XbaI buffer, 15 ul of Xba I (10U/ul), and 185 ul of H<sub>2</sub>O were added to the 250 ul of water containing about 25 ug of plasmid pL110. The digestion proceeded at 37°C for 1  
5 hour. XbaI digested pL110 was then extracted in phenol, a 1/10 volume 3M CH<sub>3</sub>COO-Na was added, 3 volumes of ethanol were added; the mixture was incubated in a dry ice-ethanol bath for 5 minutes, and then centrifuged. The precipitated DNA was resuspended in 50 ul H<sub>2</sub>O.

10 The XbaI digested plasmid pCZR111 was digested with BamHI as follows. 0.2 ul of BamHI (10 U/ul), 10 ul of BamHI buffer (100mM Tris-HCl, 50mM MgCl<sub>2</sub>, 1 M NaCl, and 10mM 2-Mercaptoethanol, pH 8.0 [at 37°C]), and 90 ul of H<sub>2</sub>O was added to the 50 ul of XbaI digested  
15 pL110 obtained hereinabove. The digest proceeded for 5 minutes at 37°C. The digested pCZR111 was extracted in phenol, a 1/10 volumes of CH<sub>3</sub>COONa+ was added, followed by addition of 3 volumes of ethanol. Precipitated DNA was resuspended in 50 ul of 10mM Tris, 1mM EDTA, pH 8.0  
20 buffer.

The XbaI and BamHI digested pCZR111 was then loaded onto an agarose gel and the DNA band at about 5.8 kb was isolated. Plasmid pCZR126S was produced by ligating the ~5.8 kb fragment of pCZR111 to an XbaI to  
25 NdeI linker and a synthetic gene encoding EK-bovine growth hormone, which contains an NdeI site on its 5' end and a BamHI site on its 3' end. The XbaI to NdeI sequence was produced using standard oligonucleotide sequence methodology and consists of the following  
30 sequence:



5' CTAGAGGGTATTAATAATGTATATTG:TTTTAATAAGGAGGAATAATCA 3'  
 |||  
 TCCATAATTATTACATATAACTAAAATTATTCCTCCTTATTAGTAT 5'

5

The above sequence was constructed by chemical synthesis of both strands, followed by mixing to allow hybridization. The gene encoding EK bGH was constructed from 16 chemically synthesized pieces of single-stranded DNA, ranging from 71 to 83 nucleotides long, which together comprise both complementary strands of the entire gene. The synthesis was done using an Applied Biosystems (ABS) machine and consists of the following sequence:

15

5' TATGTTCCCATTTGGATGATGATGATAAGTTCAGCCATGTCCTT  
 |||  
 ACAAGGGTAACCTACTACTACTATTCAAGGGTCGGTACAGGAA

20

GTCCGGCCTGTTTGCCAACGCTGTGCTCCGGGCTCAGCACCTGCATCAGCTGGCTGCTGA  
 |||  
 CAGGCCGGACAAACGGTTGCGACACGAGGCCGAGTCGTGGACGTAGTCGACCGACGACT

25

CACCTTCAAAGAGTTTGAGCGCACCTACATCCCGGAGGGACAGAGATACTCCATCCAGAA  
 |||  
 GTGGAAGTTTCTCAAACCTCGCGTGGATGTAGGGCCTCCCTGTCTCTATGAGGTAGGTCTT

30

CACCCAGGTTGCCTTCTGCTTCTCTGAAACCATCCCGGCCCCACGGGCAAGAATGAGGC  
 |||  
 GTGGGTCCAACGGAAGACGAAGAGACTTTGGTAGGGCCGGGGTGCCCGTTCTTACTCCG

35

CCAGCAGAAATCAGACTTGGAGCTGCTTCGCATCTCACTGCTCCTCATCCAGTCGTGGCT  
 |||  
 GGTCTCTTTAGTCTGAACCTCGACGAAGCGTAGAGTGACGAGGAGTAGGTGAGCACCGA

40

TGGGCCCTGCAGTTCCTCAGCAGAGTCTTACCAACAGCTTGGTGTGTTGGCACCTCGGA  
 |||  
 ACCCGGGACGTC AAGGAGTCTCTCAGAAGTGTTGTGAACCACAAACCGTGGAGCCT

CCGTGTCTATGAGAAGCTGAAGGACCTGGAGGAAGGCATCCTGGCCCTGATGCGGGAGCT  
 |||  
 GGCACAGATACTCTTCGACTTCTTGACCTCCTTCCGTAGGACCGGGACTACGCCCTCGA

```

GGAAGATGGCACCCCCGGGCTGGGCAGATCCTCAAGCAGACCTATGACAAATTTGACAC
|||||
CCTTCTACCGTGGGGGGCCCGACCCGCTTAGGAGTTCGTCTGGATACTGTTAAACTGTG

5  AAACATGCGCAGTGACGACGCGCTGCTCAAGAACTACGGTCTGCTCTCCTGCTTCCGGAA
   |||||||
   TTTGTACGCGTCACTGCTGCGCGACGAGTTCTTGATGCCAGACGAGAGGACGAAGGCCTT

GGACCTGCATAAGACGGAGACGTACCTGAGGGTCATGAAGTGCCGCCGCTTCGGGGAGGC
10 |||||||
   CCTGGACGTATTCTGCCTCTGCATGGACTCCCAGTACTTCACGGCGGCGAAGCCCCTCCG

CAGCTGTGCCTTCTAG 3'
   |||||||
15 GTCGACACGGAAGATCCTAG 5'

```

Construction of plasmid pCZR126S was accomplished by ligation of the following site components:

~0.28 ug of the 5.8 kb fragment obtained from plasmid

20 pL110 after complete digestion with XbaI and partial digestion with BamHI in a total volume of 2 ul, ~0.18 ug of the synthetic gene encoding a bovine growth factor derivative which has a 5' termini corresponding to a XbaI site and a 3' termini corresponding to a BamHI site

25 in a total volume of 2.5 ul, 8.75 picomoles of the chemically synthesized XbaI to NdeI linker in 1 ul. The plasmid components were added to 6 ul of 5x ligation buffer: 250 mM Tris-HCl, 50 mM MgCl<sub>2</sub>, 5 mM ATP, 5 mM DTT, 25% v/v polyethylene glycol 8,000, pH 7.6, 2 ul of

30 ligase, and 16.5 ul of H<sub>2</sub>O. The ligation mixture was incubated overnight at 16°C. The circularized plasmid pCZR126S was then used to transform E. coli RV308 cells in substantial accord with the method of Example 29A3. A restriction site and function map of plasmid pCZR126S

35 is presented in Figure 14 of the accompanying drawings.

B. Construction of Human Proinsulin Expression Plasmid pRB172

1. Construction of plasmid pRB145

5

The human proinsulin gene was first custom synthesized and cloned into PUC18 plasmid (commercially available from BRL). The gene was synthesized using standard techniques and comprises the sequence:

10

```

HindIII  NdeI                               DraIII
5' |AGCTTCAT|ATGATTTTGTAAACCAACACCTGTGCGGCTCCACCTG|GTGGAAGCTCT
    AGTA TACATAAAACAATTGGTTGTGGACACGCCGAGGGTGGAC CACCTTCGAGA
  
```

15

```

GTACCTGGTGTGCGGTGAACGTGGCTTCTTCTACACCCGAAGACCCGCCGTGAGGCA
CATGGACCACACGCCACTTGCACCGAAGAAGATGTGGGGCTTCTGGGGCCACTCCGT
  
```

AvaII

XmaI

```

GAG|GACCTGCAGGTGGGTGAGGTGGAGCTGGGCGGTGGC|CCGGGTGCAGGCAGCCTGC
CTC CTGGACGTCCACCCAGTCCACCTCGACCCGCCACCG GCGCCACGTCCGTGCGACG
  
```

20

```

AGCCGCTGGCCCTGGAGGGTCCCTGCAGAAGCGTGGCATTGTGGAACAATGCTGTAC
TCGGCGACCGGGACCTCCAAGGGACGTCTTCGCACCGTAACACCTTGTTACGACATG
  
```

BamHI

```

CAGCATCTGCTCCCTGTACCAGCTGGAGAACTACTGCAACTAG|GATCCG      3'
  
```

25

```

GTCGTAGACGAGGGACATGGTCGACCTCTTGATGACGTTGATC CTAGGCTTAA|  5'
  
```

EcoRI

One of the clones having the correct sequence was selected for the production of cesium chloride purified DNA. The plasmid was isolated by the standard procedure

30

[see, for example, Molecular Cloning, A Laboratory Manual, (1982) ed. by Maniatis, T; Fritsch, E.F. and Sambrook, J., Cold Spring Harbour Laboratory Publications, New York the entire teaching of which is herein  
5 incorporated by reference]. About 6  $\mu$ l (20  $\mu$ g) of this plasmid DNA was added to 20  $\mu$ l of 10X NdeI buffer (150 mM NaCl, 10 mM Tris-HCl (pH 7.8), 6 mM MgCl<sub>2</sub>, 6 mM 2-mercaptoethanol, 100  $\mu$ g/ml BSA), 5  $\mu$ l NdeI restriction enzyme (~40 units) and 169  $\mu$ l H<sub>2</sub>O. After mixing, the  
10 reaction was incubated at 37°C for 2 hours. The DNA was precipitated by making the mix 0.3 M NaOAc, adding three volumes of ethanol, mixing and chilling to -70°C and centrifuging. The DNA pellet was washed with 70% ethanol (1 ml), dried and dissolved in 20  $\mu$ l of 10X BamHI  
15 buffer (150 mM NaCl, 6 mM Tris-HCl (pH 7.9), 6 mM MgCl<sub>2</sub>, 100  $\mu$ g/ml BSA), 2  $\mu$ l of BamHI restriction enzyme (40 units) and 178  $\mu$ l H<sub>2</sub>O. After gentle mixing, the reaction was incubated at 37°C for 2 hours. The DNA was again precipitated with three volumes of ethanol as above  
20 and electrophoresed on a 1% low melting agarose gel (FMC, sea plaque agarose). The desired DNA fragment corresponding to about 270 bp was sliced from the gel and then DNA was recovered by melting the agarose and passing through an Elutip-d column (Schleicher & Schuell,  
25 Keene, N.H.) according to the procedure recommended by the vendor. After precipitation and drying, the DNA was stored in 30  $\mu$ l of 10 mM Tris-HCl pH 8.0.

About 15  $\mu$ g of plasmid pCZR126S (from Example 29A12) was suspended in 20  $\mu$ l of 10X NdeI buffer, 5  $\mu$ l  
30 of NdeI restriction enzyme (40 units) and H<sub>2</sub>O (175  $\mu$ l), gently mixed and incubated at 37°C for 2 hours. After

the incubation, the DNA was precipitated with three volumes of ethanol as above, dried and then resuspended in 20  $\mu$ l of 10X BamHI buffer, 2  $\mu$ l of BamHI restriction enzyme (40 units) and 178  $\mu$ l water. After gentle  
5 mixing, the reaction was incubated at 37°C for 2 hours. The DNA was again precipitated with three volumes of ethanol and electrophoreses on a 1% low-melting agarose gel. The larger fragment corresponding to the vector  
10 DNA was sliced from this gel and the DNA was recovered by the Elutip-d column procedure as described above. After precipitation and drying the vector DNA was stored in 35  $\mu$ l of 10 mM Tris-HCl pH 8.0.

About 2.5  $\mu$ l of the vector DNA was mixed with 12  $\mu$ l of the purified Human insulin gene fragment from  
15 above, 4  $\mu$ l of 10 mM ATP, 0.5  $\mu$ l of 1M dithiothreitol, 5  $\mu$ l of 10X Ligase buffer (500 mM Tris-HCl pH 7.6, and 100 mM  $MgCl_2$ ), 26  $\mu$ l of water and 0.5  $\mu$ l of  $T_4$ -DNA Ligase (Pharmacia, 3.5 units). The reaction was  
20 incubated at 4°C for 16 hours. The ligated mixture was diluted with 50  $\mu$ l of 10 mM Tris-HCl (pH 7.6) and 3  $\mu$ l of 1 M  $CaCl_2$  and then subsequently transformed into E. coli K12 RV308 in substantial accordance with the  
25 teaching of Example 29A3. The cells were plated on TY plates supplemented with 5  $\mu$ g/ml tetracycline then incubated overnight at 32°C.

Plasmids from 3 mL cultures were isolated from tetracycline resistant colonies by the rapid alkaline extraction procedure described in Molecular Cloning, A Laboratory Manual (1982) ed. by Maniatis, T.,  
30 Fritsch, E.F., and Sambrook, J. Cold Spring Harbour publications, New York, (page 368-369). The presence

of the correct human proinsulin gene fragment was found by miniscreen procedure using polyacrylamide gel electrophoresis to analyze the XbaI/BamHI digested fragment. Those plasmids with the correct size (about 5 315 bp) inserts were selected for amplification and purification. The plasmid containing the human proinsulin gene was pRB145. A restriction site and function map of plasmid pRB145 is presented in Figure 16 of the accompanying drawings.

10

## 2. Construction of plasmid pRB164A

About 30  $\mu$ g of plasmid pRB145 was suspended in 20  $\mu$ l of 10X NdeI buffer, 5  $\mu$ l of NdeI restriction enzyme (New England Biolabs, 40 units), and 175  $\mu$ l H<sub>2</sub>O gently mixed and incubated at 37°C for 1 hour. Two  $\mu$ l of BamHI restriction enzyme (New England Biolabs, 40 units) was then added to the reaction mixture and the incubation at 37°C was continued for another 2 hours. 15 The DNA was precipitated with three volumes of ethanol and 0.3M NaOAc and electrophoresed on a 1% low melting agarose gel. The smaller (about 270 bp) 20 NdeI/BamHI restriction fragment encoding the human proinsulin gene was sliced from the gel and the DNA was recovered by passing through an Elutip-d column as described previously. After precipitation and drying, 25 the DNA was stored in 30  $\mu$ l of 10 mM Tris pH 8.0.

To this DNA (30  $\mu$ l) was then added 20  $\mu$ l of 10X AvaII buffer (50 mM NaCl, 6 mM Tris-HCl, pH 8.0, 30 10 mM MgCl<sub>2</sub>, 6 mM 2-mercaptoethanol, 100  $\mu$ g/ml BSA), 5  $\mu$ l of AvaII restriction enzyme (New England Biolabs,

20 units), and 175  $\mu$ l of H<sub>2</sub>O. After gently mixing, this reaction was incubated at 37°C for 2 hours. The DNA was precipitated with three volumes of ethanol and 3M NaOAc (20  $\mu$ l) and then electrophoresed on a 1.2% low melting agarose gel. The larger AvaII/BamHI restriction fragment (about 156 bp) was sliced from the gel and then DNA was recovered by passing through an Elutip-d column as described above. After precipitation and drying, the DNA was stored in 30  $\mu$ l of 10 mM tris pH 8.0.

10 The DNA ( $\sim$ 115 bp) corresponding to the NdeI/AvaII restriction fragment of human proinsulin gene was synthetically prepared. The first step consisted of synthesis of four single stranded deoxyribooligonucleotides by the DNA synthesizer (Applied Biosystems, model 380B). The nucleotide sequences of these four oligonucleotides are

1. TATGCGTATGTTTGTAAACCAACACCTGTGCGGCTCCACCTG  
GTGGAAGCTCTGTACCT (60 mer)
- 20 2. GGTGTGCGGTGAACGTGGCTTCTTCTACACCAAGCCGACC  
CGCCGTGAGGCAGAG (55 mer)
3. CACCAGGTACAGAGCTTCCACCAGGTGGGAGCC  
25 GCACAGGTGTTGGTTAACAAACATACGCA (62 mer)
4. GTCCTCTGCCTCACGGCGGGTCGGCTTGGTGTAGAA  
GAAGCCACGTTACCGCA (54 mer)

After purifying each oligonucleotide by polyacrylamide gel electrophoresis, oligonucleotides two and three were phosphorylated according to the teachings of Brown, E.L., Belagaje, R., Ryan, M.J., and Khorana, H.G. (1979) *Methods in Enzymology*, Ed. by Wu, R., Academic Press, N.Y. 68, 109-151 the entire teaching of which is herein incorporated by reference. The phosphorylated oligonucleotides two and three (~715 pmoles of each) were then mixed with oligonucleotides one and two (~860 pmoles), annealed and ligated in a buffer (200  $\mu$ l) containing 50 mM Tris-HCl (pH 7.6), 10 mM MgCl<sub>2</sub>, 10 mM DTT, 50  $\mu$ M ATP and 20 units of T<sub>4</sub> DNA ligase (Pharmacia) for 16 hours at 4°C. The ligation product was purified on a 15% polyacrylamide gel. The DNA was recovered from the gel slice electrophoretically followed by desalting on a Sephadex G-50 column. The yield of the desired ligated product was 485 pmoles.

About 100 pmoles of this DNA were phosphorylated in a buffer (50  $\mu$ l) containing 50 mM Tris (pH 7.6), 10 mM MgCl<sub>2</sub>, 10 mM DTT and ATP, as described in Brown, E.L. et al., (1979), *Methods in Enzymology* 68, 109-151 the teaching of which is herein incorporated by reference. After filtration through a column of Sephadex G-50, the DNA was stored in 50  $\mu$ l of 10 mM Tris, pH 8.0.

About 2.5  $\mu$ l of the vector DNA (NdeI-BamHI digested pCZR126S) was mixed with 18  $\mu$ l of AvaII/BamHI restriction fragment from plasmid pRB145 and 10  $\mu$ l (10 pmoles) NdeI/AvaII synthetic linker just constructed in a buffer (50  $\mu$ l) containing 50 mM Tris (pH 7.6),



10 mM MgCl<sub>2</sub>, 10 mM DTT, 800 µl ATP and 3.5 units of T<sub>4</sub> DNA ligase. The reaction was incubated at 4°C overnight and then transformed into E. coli K12 RV308 in accordance with the procedure of Example 29A3. The  
5 desired transformant E. coli K12 RV308/pRB164A was identified by analysis of its plasmid DNA. The proper transformant was grown at 30°C in TY media containing 5 µg/ml tetracycline to an O.D.<sub>550</sub> of about  
10 0.2 (early log phase) and then shifted to 42°C for 3 to 3.5 hours to induce the synthesis of the human proinsulin. The cells were pelleted and lysed by the addition of sample buffer (0.125 M Tris-HCl (pH 6.8), 2% SDS, 30% glycerol, 1M 2-mercaptoethanol, 6M urea). Samples were heated to 97°C for 2 minutes prior to  
15 loading on an acrylamide gel. Bands were easily visualized by staining with Coomassie Brilliant Blue dye. A scanning gel densitometer was used to quantitate the percentages of cell protein. A restriction site and  
20 function map of plasmid pRB164A is presented in Figure 17 of the accompanying drawings.

### 3. Construction of plasmid pRB172

25 About 25 µg of plasmid pRB145 was suspended in 15 µl of 10X DraIII buffer (200 mM NaCl, 50 mM Tris-HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 1 mM DTT, 10 µg/ml BSA) 6 µl of DraIII restriction enzyme (Boehringer Mannheim, 27 units) and 129 µl of H<sub>2</sub>O, gently mixed and incubated at  
30 37°C for 6 hours. After the incubation, the DNA was precipitated, dried and resuspended in 10 µl of 10X XbaI buffer (150 mM NaCl, 6 mM Tris-HCl pH 7.9, 6 mM

MgCl<sub>2</sub>, 7 mM 2 mercaptoethanol, 100 µg/ml BSA), 3 µl of XbaI restriction enzyme (Boehringer Mannheim, 36 units) and 77 µl of H<sub>2</sub>O. The reaction was gently mixed and incubated at 37°C for 4 hours. The DNA was again  
5 precipitated with three volumes of ethanol and NaOAc and electrophoresed on a 1% low melting agarose gel. The lower band corresponding to the about 85 bp XbaI/DraIII restriction fragment of human proinsulin gene was sliced from the gel and the DNA was recovered by the Elutip-d  
10 column procedure. After precipitation and drying, the DNA was stored in 30 µl of 10 mM Tris pH 8.0.

About 15 µg of plasmid pRB164A was cut with the restriction enzyme DraIII and XbaI in accordance with the procedure described above. The upper band  
15 corresponding to the XbaI/DraIII vector fragment was isolated from the agarose gel by the Elutip-d column procedure. After precipitation and drying, the DNA was stored in 30 µl of 10 mM Tris pH 8.0.

About 5 µl of the XbaI/DraIII digested pRB164A  
20 vector DNA was mixed with 5 µl of the XbaI/DraIII restriction fragment from plasmid pBR145 in a buffer (50 µl) containing 50 mM Tris-HCl (pH 7.6), 10 mM MgCl<sub>2</sub>, 10 mM DTT, 800 µM ATP and 6 units of T<sub>4</sub> DNA ligase. The reaction was incubated at 4°C overnight and used to  
25 transform E. coli K12/RV308 cells made competent by a standard CaCl<sub>2</sub> treatment.

Transformants were selected on TY agar plates containing 5 µg/ml tetracycline. Plasmids were isolated from tetracycline resistant colonies by the rapid  
30 alkaline extraction procedure and analyzed by digestion with XbaI and BamHI restriction endonucleases. DNA

from positive clones was sequenced using the sequenase system (U.S. Biochemicals). Those plasmids with the correct desired sequence were selected for amplification and purification. In this manner, the E. coli K12  
5 RV308/pRB172 transformant was isolated. Expression and accumulation of human proinsulin harboring this plasmid was analyzed by visualization of the total cellular protein following electrophoretic separation in a 15% polyacrylamide matrix. A restriction site and function  
10 map of plasmid pRB172 is presented in Figure 18 of the accompanying drawings.

C. Construction of Escherichia coli RV308/pRB173 and pRB174

15 About 20 µg of plasmid pRB172 or pRB145 was suspended in 20 µl of 10X XbaI buffer (150 mM NaCl, 6 mM Tris-HCl, pH 7.9, 6 mM MgCl<sub>2</sub>, 7 mM 2-mercaptoethanol, 100 µg/ml BSA) 2 µl of XbaI restriction enzyme (Boehringer Mannheim, 24 units) was added, gently mixed  
20 and incubated at 37°C for 4 hours. After the incubation, the DNA was precipitated, dried and resuspended in 10 µl of 10X BamHI buffer (100 mM NaCl, 10 mM Tris-HCl (pH 8.0), 5 mM MgCl<sub>2</sub>, 1 mM 2-mercaptoethanol,  
25 100 µg/ml BSA), 5 µl of BamHI restriction enzyme  
Boehringer Mannheim (40 units) and 75 µl of H<sub>2</sub>O. The reaction was gently mixed and incubated at 37°C for 4 hours. The DNA was again precipitated with ethanol and NaOAc and electrophoresed on a 1% low melting agarose gel. The about 320 bp XbaI/BamHI restriction  
30 fragment corresponding to the human proinsulin gene was isolated from the gel by the Elutip-d column procedure.

After precipitation and drying, the DNA was resuspended in 20  $\mu$ l of 10X XmaI buffer (25 mM NaCl, 10 mM Tris-HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 10 mM 2-mercaptoethanol, 100  $\mu$ g/ml BSA), 10  $\mu$ l of XmaI restriction enzyme (New England Biolabs 10 units) and 170  $\mu$ l of water. The reaction was gently mixed and incubated at 37°C for 4 hours. After the incubation, the DNA was precipitated with ethanol and NaOAc and electrophoresed on a 1% low melting agarose gel. The larger DNA corresponding to the about 200 bp XbaI/XmaI restriction fragment was sliced from the gel and the DNA was recovered by the Elutip-d column as described before. The DNA was stored in 30  $\mu$ l of 10 mM Tris pH 8.0.

The 51 bp DNA corresponding to an XmaI/BamHI restriction fragment of human proinsulin gene was constructed synthetically as follows. Two single stranded deoxyribooligonucleotides:

CCGGGTGCAGGCAGCCTGCAGCCGCTGGCCCTGGAGGGTTCCTGCAGTAG (51 mer) and

GATCCTACTGCAGGGAACCCCTCCAGGGCCAGCGGCTGCAGGCTGCCTGCAC (51 mer)

were synthesized on an applied Biosystems DNA Synthesizer (model 380B) and purified by polyacrylamide gel electrophoresis in presence of 7 M urea. After isolation, the two oligonucleotides were phosphorylated according to the teachings from Brown, E.L., Belagaje, R., Ryan, M.J., and Khorana, H.G. (1979) Methods in Enzymology, Ed. by Wu, R., Academic Press, N.Y. 68, 109-151, and annealed to form the desired linker.

About 10  $\mu$ g of plasmid pCZR126S (from Example 29A12) was suspended in 20  $\mu$ l of 10X XbaI buffer (150 mM NaCl, 6 mM Tris-HCl (pH 7.9), 6 mM MgCl<sub>2</sub>, 7 mM 2-

mercaptoethanol, 100 µg/ml BSA) 2 µl of XbaI restriction enzyme (Boehringer Mannheim, 24 units) and 178 µl of water. The reaction was gently mixed and incubated at 37°C for 2 hours. After this incubation, 4 µl of 5M NaCl and 2 µl of BamHI restriction enzyme (New England Biolabs, 20 units) were added, and the incubation was continued for another 2 hours at 37°C. The DNA was then precipitated with ethanol and NaOAc by the standard procedure and electrophoresed on a 1% low melting agarose gel. The upper band corresponding to XbaI/BamHI vector DNA was isolated with a Elutip-d column. After precipitation and drying, the DNA was stored in 30 µl of 10 mM Tris, pH 8.0.

About 5 µl of the XbaI/BamHI-digested pCZR126S vector DNA was mixed with 10 µl of the XbaI/XmaI restriction fragment from pRB145 or pRB172 and 4 µl of the Kinased XmaI/BamHI linker prepared above in a buffer (50 µl) containing 50 µM Tris (pH 7.6), 10 mM MgCl<sub>2</sub>, 10 mM DTT, 800 µM ATP and 6 units of T<sub>4</sub> DNA ligase. The reaction was incubated at 4°C overnight and used to transform E. coli K12 RV308 strain in accordance with the procedure of Example 29A3. The desired transformants, E. coli K12 RV308/pRB173 (if made from the pRB172 vector fragment) and E. coli K12 RV308/pRB174 (if made from the pRB145 vector fragment) were identified by analysis of their plasmid DNA and protein production before and after induction at 42°C. A restriction site and function map of plasmid pRB173 is presented in Figure 19 of the accompanying drawings.

About 200 ng of plasmid DNA from the mini-prep were also transformed into E. coli L201 cells and the

transformants were selected on 2 x TY agar plates containing 15 µg/ml tetracycline. The proteins expressed by the transformants were analyzed and the desired transformants were confirmed by their production of  
5 human proinsulin. In this matter, E. coli L201/pRB173 and E. coli L201/pRB174 were isolated.

D. Construction of Escherichia coli K12 RV308/pRB175, pRB176, pRB177 and pRB178

10

About 12 µg of plasmid pRB172 was suspended in 15 µl of 10X NdeI buffer (100 mM NaCl, 50 mM Tris-HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 1 mM DTT, 100 µg/ml BSA), 2.5 µl of NdeI restriction enzyme (20 units) and 133 µl of  
15 water. The reaction was gently mixed and incubated at 37°C overnight. After the incubation, the DNA was precipitated with ethanol and NaOAc by the standard procedure, dried and resuspended in 15 µl of 10X DraIII buffer (50 mM Tris-HCl (pH 7.5), 200 mM NaCl, 10 mM  
20 MgCl<sub>2</sub>, 1 mM DTT) 5 µl of DraIII restriction enzyme (20 units) and 130 µl of H<sub>2</sub>O. After gently mixing, the reaction was incubated at 37°C overnight. The DNA was again precipitated with ethanol and NaOAc as above and electrophoresed on a 1% low melting agarose. The  
25 desired larger restriction fragment was sliced from the gel and the DNA was recovered by the Elutip-d column procedure. The DNA was stored in 30 µl of 10 mM Tris HCl pH 8.0.

The following DNA linkers were chemically synthesized by the Applied Biosystems DNA Synthesizer (model 380B):

- 5 (i) 5' TATGTACGACCAACACCTGTGCGGCTCCCATCTG 3'  
3' ACATGCTGGTTGTGGACACGCCGAGGGTA 5' for pRB175
- (ii) 5' TATGTACGACGTTAACCAACACCTGTGCGGCTCCCATCTG 3'  
3' ACATGCTGCAATTGGTTGTGGACACGCCGAGGGTA 5' for pRB176
- 10 (iii) 5' TATGTATAACCAACACCTGTGCGGCTCCCATCTG 3'  
3' ACATATTGGTTGTGGACACGCCGAGGGTA 5' for pRB177
- (iv) 5' TATGTACGTTAACCAACACCTGTGCGGCTCCCATCTG 3'  
15 3' ACATGCAATTGGTTGTGGACACGCCGAGGGTA 5' for pRB178

After purifying each oligonucleotide by polyacrylamide gel electrophoresis, they were phosphorylated according to the teachings mentioned previously and annealed in order to facilitate the ligation and construction of the human proinsulin gene analog-encoding DNA fragments. The linkers (i) to (iv) were used to construct the plasmids pRB175, pRB176, pRB177 and pRB178 respectively.

About 3  $\mu$ l of the NdeI/DraIII-digested pRB172 vector DNA fragment was mixed with 4  $\mu$ l of each of the separate Kinased linkers (i-iv) in a buffer (50  $\mu$ l) containing 50  $\mu$ M Tris-HCl (pH 7.6), 10 mM MgCl<sub>2</sub>, 10 mM DTT, 800  $\mu$ M ATP and 6 units of T<sub>4</sub> DNA ligase. The

reaction was incubated at 4°C overnight and this mixture was used to transform E. coli K12 RV308 cells in accordance with the procedure of Example 29A3. The desired transformants, E. coli K12 RV308/pRB175, E. coli K12 RV308/pRB176, E. coli RV308/pRB177 and E. coli K12 RV308/pRB178 were identified by analysis of their plasmid DNA and protein production before and after induction of the cells at 42°C. A restriction site and function map of plasmid pRB175 is presented in Figure 20 of the accompanying drawings.

Table I: Human Proinsulin Analogs

|    | <u>Analog</u>   | <u>Plasmid</u> | <u>% Protein</u> |
|----|---|----------------|------------------|
| 15 | Met-Tyr-Human Proinsulin (HPI)  | pRB145         | 11.3             |
|    | Met-Arg-Met-[Lys(B28),Pro(B29)HPI]  | pRB164A        | 10.5             |
|    | Met-Tyr-[Lys(B28),Pro(B29)HPI]  | pRB172         | 7.3              |
|    | Met-Tyr[Lys(B28),Pro(B29)B-chain]-Q*  | pRB173         | ND               |
| 20 | Met-Tyr-[B-chain]-Q*  | pRB174         | ND               |
|    | Met-Tyr-[Des(B1),Des(B2),Asp(B3),<br>Lys(B28),Pro(B29)HPI]  | pRB175         | 8.5              |
|    | Met-Tyr-[Asp(B1),Lys(B28),<br>Pro(B29)HPI]  | pRB176         | 9.3              |
| 25 | Met-Tyr-[Des(B1),Des(B2),Lys(B28),<br>Pro(B29)HPI]  | pRB177         | 9.4              |
|    | Met-Tyr-[Des(B1),Lys(B28),<br>Pro(B29)HPI]  | pRB178         | 8.5              |
| 30 | *Q = -Arg-Arg-Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-<br>Leu-Gly-Gly-Gly-Pro-Gly-Ala-Gly-Ser-Leu-Gln-Pro-Leu-<br>Ala-Leu-Glu-Gly-Ser-Leu-Gln-; |                |                  |



E. A stock culture of E. coli K12 RV308/pRB172 was formed into a permanent stock culture in the following manner.

Isolated colonies are spotted on the following  
5 phenotype plates: L-agar + 5 mcg./ml. Tc (tetracycline),  
L-agar + Sm (streptomycin sulfate), M-9 medium agar (a  
salt based medium containing MgSO<sub>4</sub>, thiamine, HYCASE,  
and glucose), and M-9 medium agar without glucose and  
with lactose. One L-agar/Tc plate is incubated at 42°C,  
10 and the remaining plates are incubated at 30°C for 24  
hours. A phenotypically correct colony selected from  
the 30°C L-agar/Tc plate is used to inoculate a flask  
containing 50 ml. L-broth (contains tryptone, yeast  
extract, NaCl, and glucose) plus 5 mcg./ml. Tc medium.  
15 The flask is incubated at 30°C with shaking for 7 hours.  
The material in the flask (1.2 ml.) is dispensed into  
bio-freeze vials and frozen in vapor-phase liquid nitro-  
gen. The material then is thawed, and 1 ml. is used to  
inoculate a flask containing 50 ml. of L-broth plus  
20 5 mcg./ml. Tc. The flask is incubated at 30°C with  
shaking for 4 hours. Two flasks containing 500 ml. of  
BG-7 medium (contains MgSO<sub>4</sub>, thiamine, glucose, Tc, and  
trace salts) are inoculated with 1 ml. each from the  
foregoing, and the flasks are incubated with shaking  
25 for 9 hours at 30°C.

A 150-liter fermenter containing 80 liters of  
medium (contains citric acid, ferrous ammonium sulfate,  
K<sub>2</sub>HPO<sub>4</sub>, NH<sub>4</sub>Cl, NaH<sub>2</sub>PO<sub>4</sub>, CaCl<sub>2</sub>, MgSO<sub>4</sub>, glucose, and trace  
inorganic salts) is inoculated with 1 liter of the BG-7  
30 medium broth from above. The fermenter is operated at  
30°C until the carbon dioxide evolution rate reaches

1.0 mM/L/min. At that point, 50 liters of broth are transferred to the production fermenter to provide the cell inoculum.

5 A 2500-liter fermenter containing 1500 liters of the same medium used in the 150-liter fermenter is operated at 30°C until the carbon dioxide evolution rate reaches 1.0 mM/L/min, and then the temperature is raised to 40C. Operation is continued for an additional 8  
10 hours. The broth is then heat inactivated at 61°C for 7 minutes. The following yield data were obtained from the heat inactivated broth: Cell Yield, 13.1 grams/L dry weight; Product Potency, 273 mcg. Des-formylate/ml. of broth; Specific Activity (e.g. grams product/gram of cell), 2.1 percent; Percent Formylated Product, 13  
15 percent.

Fermentation broth is transferred into a stainless steel holding tank and the temperature maintained between 2° and 8°C. The whole broth is then centrifuged or filtered to collect the E. coli cells such that a  
20 solids dry weight of approximately 15 to 20 percent is obtained. The collected cell solids are reslurried in process water. The cells are then disrupted using high pressure homogenization or other suitable methods to achieve lysis of 90 to 99 percent of the cells. The  
25 homogenized slurry is diluted to 5 percent dry weight with process water. The pH of the diluted disrupted cell slurry is adjusted to between pH 7 and 10 by adding sodium hydroxide. The inclusion bodies (granules) are separated from the cell debris by differential centrifuga-  
30 tion. The resulting concentrate is from 15 to 20 percent solids.

The inclusion bodies are solubilized at alkaline pH in the presence of cysteine and urea. These solubilized granules are subjected to cation exchange chromatography on SP Sepharose-superflow in order to  
5 resolve the mixed disulfide form of the proinsulin analog from the bulk of granular material.

The methionyl-tyrosine extension of the amino terminus of the proinsulin analog is removed by incubation with cathepsin C, a dipeptidylaminopeptidase. The  
10 cathepsin cleavage reaction is terminated by sulfitolysis with potassium tetrathionate and cysteine at pH 8.8. Following solvent-exchange across Sephadex G-25, the resulting S-sulphonate of the proinsulin analog is allowed to fold by incubation at pH 10.6 in the presence  
15 of cysteine. The folding process is quenched by adjusting the pH to 2.4. Properly folded proinsulin analog is resolved from improperly folded material as well as contaminants remaining from the cathepsin C cleavage step by hydrophobic interaction chromatography on SP20SS  
20 resin. This separation occurs across an acetone gradient so the organic solvent must be removed from the mainstream pool by evaporation before further processing.

Following evaporation of the organic solvent, properly folded proinsulin analog is incubated with  
25 trypsin and carboxypeptidase B. These enzymes excise the C peptide of the proinsulin analog and generate Lys(B28), Pro(B29) human insulin. The insulin analog is then further purified by cation exchange chromatography on S-Sepharose followed by high performance reverse  
30 phase chromatography on a 10  $\mu$ m C-8 resin. Acetonitrile is removed from the reverse phase mainstream by solvent

exchange across Sephadex G-25 and the resulting material is concentrated using a spiral wound ultrafiltration system. Concentrated material is subjected to size exclusion chromatography on Sephadex G-50 resin and  
5 lyophilized to dryness.

In Table II following are provided the amino acid analyses of compounds of the foregoing examples.

Table II

| Example No.        | Amino Acid Analysis on Human Insulin Analogs <sup>1,2</sup> |                   |         |         |         |         |         |         |         |
|--------------------|---|-------------------|---------|---------|---------|---------|---------|---------|---------|
|                    | 2   | 3                 | 4       | 5       | 6       | 7       | 8       | 9       | 10      |
| Amino Acid Residue |   |                   |         |         |         |         |         |         |         |
| Asp                | 3.00(3)   | 3.00(3)           | 3.00(3) | 3.00(3) | 4.00(4) | 4.00(4) | 4.00(4) | 3.00(3) | 3.00(3) |
| Thr                | 2.82(3)   | 2.81(3)           | 2.64(3) | 2.80(3) | 2.80(3) | 2.94(3) | 2.81(3) | 2.71(3) | 2.73(3) |
| Ser                | 2.76(3)   | 2.73(3)           | 2.70(3) | 2.75(3) | 2.71(3) | 2.46(3) | 2.71(3) | 2.74(3) | 2.72(3) |
| Glu                | 7.16(7)   | 7.05(7)           | 6.93(7) | 7.08(7) | 7.01(7) | 6.69(7) | 7.18(7) | 6.99(7) | 7.94(8) |
| Pro                | 1.02(1)   | 1.11(1)           | 0.92(1) | 0.98(1) | 1.01(1) | 0.98(1) | 1.02(1) | 1.09(1) | 0.98(1) |
| Gly                | 3.93(4)   | 4.00(4)           | 3.85(4) | 3.99(4) | 4.02(4) | 3.97(4) | 4.01(4) | 3.95(4) | 3.90(4) |
| Ala                | 1.01(1)   | 1.04(1)           | 1.93(2) | 1.02(1) | 1.05(1) | 1.05(1) | 1.03(1) | 1.03(1) | 1.02(1) |
| Cys(1/2)           | 5.13(6)   | -(6) <sup>3</sup> | 5.13(6) | 5.17(6) | 5.32(6) | 4.91(6) | 5.53(6) | 5.23(6) | 5.33(6) |
| Val                | 3.45(4)   | 3.44(4)           | 3.49(4) | 3.51(4) | 3.41(4) | 3.67(4) | 3.54(4) | 3.45(4) | 3.48(4) |
| Ile                | 1.47(2)   | 1.47(2)           | 1.49(2) | 1.58(2) | 1.44(2) | 1.65(2) | 1.68(2) | 1.48(2) | 1.47(2) |
| Leu                | 6.04(6)   | 6.01(6)           | 5.86(6) | 6.19(6) | 6.06(6) | 5.67(6) | 5.99(6) | 6.02(6) | 5.98(6) |
| Tyr                | 3.83(4)   | 3.74(4)           | 3.52(4) | 4.19(4) | 3.90(4) | 3.55(4) | 3.82(4) | 3.73(4) | 3.58(4) |
| Phe                | 2.84(3)   | 2.85(3)           | 2.64(3) | 2.85(3) | 2.84(3) | 3.02(3) | 2.86(3) | 2.72(3) | 2.73(3) |
| His                | 2.35(2)   | 2.10(2)           | 2.02(2) | 2.14(2) | 1.60(2) | 2.09(2) | 1.07(1) | 2.02(2) | 2.10(2) |
| Lys                | 0.97(1)   |                   |         |         |         |         | 0.97(1) |         |         |
| Arg                | 0.99(1)   | 0.90(1)           | 0.88(1) | 1.96(2) | 0.91(1) | 0.81(1) | 0.91(1) | 0.89(1) | 0.98(1) |
| Aba                |   | -(1) <sup>3</sup> |         |         |         |         |         |         |         |
| Cya                |   |                   |         |         |         |         |         | 0.90(1) |         |
| Nle                |   |                   |         |         |         |         |         |         |         |
| Orn                |   |                   |         |         |         |         |         |         |         |
| Nva                |   |                   |         |         |         |         |         |         |         |
| Met                |   |                   |         |         |         |         |         |         |         |
| Trp                |   |                   |         |         |         |         |         |         |         |

<sup>1</sup> Molar unity based on aspartic acid.

<sup>2</sup> Number in parenthesis is theoretical amino acid content.

<sup>3</sup> Half-cystine plus aminobutyric acid totaled 6.49.

X-7512A

-96-

Table II, cont'd.

| Example No.               | Amino Acid Analysis on Human Insulin Analogs <sup>1,2</sup> |         |         |         |         |         |         |         |         |
|---------------------------|---|---------|---------|---------|---------|---------|---------|---------|---------|
|                           | 11  | 12      | 13      | 14      | 15      | 16      | 17      | 18      | 19      |
| <u>Amino Acid Residue</u> |   |         |         |         |         |         |         |         |         |
| Asp                       | 3.00(3)   | 3.00(3) | 3.00(3) | 3.00(3) | 3.00(3) | 3.00(3) | 3.00(3) | 3.00(3) | 3.00(3) |
| Thr                       | 2.76(3)   | 3.59(3) | 2.80(3) | 2.80(3) | 2.70(3) | 2.83(3) | 2.85(3) | 3.03(3) | 2.80(3) |
| Ser                       | 2.69(3)   | 2.64(3) | 2.43(3) | 2.78(3) | 2.87(3) | 2.78(3) | 2.97(3) | 2.68(3) | 2.72(3) |
| Glu                       | 8.04(8)   | 6.84(7) | 6.89(7) | 7.09(7) | 7.04(7) | 7.08(7) | 6.94(7) | 6.99(7) | 7.01(7) |
| Pro                       | 1.11(1)   | 1.48(1) | 1.09(1) | 1.12(1) | 0.93(1) | 1.16(1) | 1.10(1) | 1.14(1) | 1.02(1) |
| Gly                       | 4.04(4)   | 5.70(5) | 3.89(4) | 4.03(4) | 3.98(4) | 4.00(4) | 4.12(4) | 4.08(4) | 3.92(4) |
| Ala                       | 1.05(1)   | 1.14(1) | 1.04(1) | 1.05(1) | 1.08(1) | 1.04(1) | 1.09(1) | 1.07(1) | 1.03(1) |
| Cys(1/2)                  | 5.26(6)   | 4.54(6) | 4.57(6) | 5.19(6) | 5.00(6) | 4.75(6) | 5.20(6) | 4.82(6) | 5.30(6) |
| Val                       | 3.60(4)   | 3.52(4) | 3.58(4) | 3.43(4) | 3.41(4) | 3.51(4) | 3.49(4) | 3.63(4) | 3.51(4) |
| Ile                       | 1.62(2)   | 1.82(2) | 1.70(2) | 2.36(3) | 1.44(2) | 1.51(2) | 1.52(2) | 1.75(2) | 1.55(2) |
| Leu                       | 5.99(6)   | 5.79(6) | 5.95(6) | 5.98(6) | 6.89(7) | 6.10(6) | 5.95(6) | 5.98(6) | 6.11(6) |
| Tyr                       | 3.73(4)   | 3.89(4) | 3.75(4) | 3.74(4) | 3.64(4) | 3.86(4) | 3.74(4) | 3.80(4) | 3.84(4) |
| Phe                       | 2.88(3)   | 3.47(3) | 3.01(3) | 2.85(3) | 2.71(3) | 2.86(3) | 2.87(3) | 3.13(3) | 2.89(3) |
| His                       | 2.08(2)   | 2.24(2) | 3.39(3) | 2.07(2) | 2.30(2) | 2.48(2) | 2.20(2) | 2.29(2) | 2.10(2) |
| Lys                       |   |         |         |         |         |         | 0.99(1) |         |         |
| Arg                       | 0.91(1)   | 0.80(1) | 0.95(1) | 0.89(1) | 0.86(1) | 1.00(1) | 0.87(1) | 0.80(1) | 1.03(1) |
| Aba                       |   |         |         |         |         |         |         |         |         |
| Cya                       |   |         |         |         |         |         |         |         |         |
| Nle                       |   |         |         |         |         | 0.93(1) |         |         |         |
| Orn                       |   |         |         |         |         |         |         |         | 0.96(1) |
| Nva                       |   |         |         |         |         |         |         |         |         |
| Met                       |   |         |         |         |         |         |         | 0.98(1) |         |
| Trp                       |   |         |         |         |         |         |         |         |         |

X-7512A

Table II, cont'd.

| Example No.        | Amino Acid Analysis on Human Insulin Analogs <sup>1,2</sup> |         |         |         |         |         |         |         |
|--------------------|---|---------|---------|---------|---------|---------|---------|---------|
|                    | 20  | 21      | 22      | 23      | 24      | 25      | 26      | 27      |
| Amino Acid Residue |   |         |         |         |         |         |         |         |
| Asp                | 3.00(3)   | 3.00(3) | 3.00(3) | 3.00(3) | 3.00(3) | 3.00(3) | 3.00(3) | 3.00(3) |
| Thr                | 2.51(3)   | 3.04(3) | 2.76(3) | 3.73(4) | 2.70(3) | 2.80(3) | 3.16(3) | 2.75(3) |
| Ser                | 2.55(3)   | 2.69(3) | 3.60(4) | 2.74(3) | 2.76(3) | 2.78(3) | 2.75(3) | 2.63(3) |
| Glu                | 7.25(7)   | 7.06(7) | 6.99(7) | 7.07(7) | 7.01(7) | 7.04(7) | 6.98(7) | 7.02(7) |
| Pro                | 0.93(1)   | 2.20(2) | 0.95(1) | 1.00(1) | 0.95(1) | 1.11(1) | 1.22(1) | 1.16(1) |
| Gly                | 3.92(4)   | 3.93(4) | 3.91(4) | 3.97(4) | 3.94(4) | 3.97(4) | 4.26(4) | 3.98(4) |
| Ala                | 1.09(1)   | 1.02(1) | 1.07(1) | 1.03(1) | 1.08(1) | 1.04(1) | 1.14(1) | 1.03(1) |
| Cys(1/2)           | 4.53(6)   | 5.57(6) | 4.31(6) | 5.01(6) | 4.05(6) | 5.17(6) | 4.81(6) | 5.16(6) |
| Val                | 3.72(4)   | 3.61(4) | 3.37(4) | 3.42(4) | 3.41(4) | 3.43(4) | 4.80(5) | 3.59(4) |
| Ile                | 1.72(2)   | 1.56(2) | 1.42(2) | 1.48(2) | 1.44(2) | 1.48(2) | 1.67(2) | 1.59(2) |
| Leu                | 6.01(6)   | 6.09(6) | 5.95(6) | 6.02(6) | 6.01(6) | 5.98(6) | 5.96(6) | 5.99(6) |
| Tyr                | 3.55(4)   | 3.93(4) | 3.60(4) | 3.72(4) | 3.78(4) | 4.67(5) | 3.77(4) | 3.75(4) |
| Phe                | 3.59(4)   | 3.13(3) | 2.76(3) | 2.85(3) | 2.74(3) | 2.83(3) | 3.28(3) | 2.87(3) |
| His                | 2.14(2)   | 2.10(2) | 2.55(2) | 2.32(2) | 2.82(2) | 2.08(2) | 2.25(2) | 2.02(2) |
| Lys                |   |         |         |         |         |         |         |         |
| Arg                | 0.87(1)   | 1.02(1) | 0.85(1) | 0.92(1) | 1.05(1) | 0.86(1) | 1.02(1) | 0.87(1) |
| Aba                |   |         |         |         |         |         |         |         |
| Cya                |   |         |         |         |         |         |         |         |
| Nle                |   |         |         |         |         |         |         |         |
| Orn                |   |         |         |         |         |         |         |         |
| Nva                |   |         |         |         |         |         |         | 0.98(1) |
| Met                |   |         |         |         |         |         |         |         |
| Trp                |   |         |         |         | 0.89(1) |         |         |         |

X-7512A

-98-

The physiological effects of the insulin analogs of the present invention were shown in the following in vivo assay system.

Normal male Sprague Dawley rats from the  
5 Charles River Laboratories (Portage, MI) were used as  
test animals. They were obtained at a weight range  
of 160-180 gms and maintained for one week in animal  
rooms at 75°F with a controlled light cycle (lights  
on 7:00 a.m. - 7:00 p.m., lights off 7:00 p.m. - 7:00  
10 a.m.) The animals were fed Purina rat chow 5001  
ad libitum. Rats used for each assay were fasted for  
16 hours before being used. They weighed about 200 gms  
when first used. On reaching a fasted weight of about  
275 gm (over a period of three weeks), the animals were  
15 no longer used. One group of ten male rats was used  
each day for each compound tested (i.e., biosynthetic  
human insulin, porcine insulin and human insulin analog).  
Each group was used only once during the week. The ten  
rats were divided into two groups of five rats each.  
20 One group served as control and was given vehicle alone  
subcutaneously. The other group of five rats was given  
the compound to be tested. The proteins were dissolved  
in 0.05N HCl (pH 1.6) to provide a stock solution of 100  
µgm per ml. From this, a number of dilutions were made  
25 in normal saline which was injected subcutaneously into  
the rats. A 100 µl sample of blood was taken from the  
tail vein of each rat at zero time and 30 minutes, 1  
hour, 2 hours, 3 hours and 4 hours after administration.  
Glucose was determined colorimetrically by a glucose  
30 oxidase method (Sigma Chemical Co.). The percent change  
in blood glucose from the zero time value was calculated



for each rat and the final results were expressed as the mean percent change  $\pm$  SEM in the experimental group corrected for the mean change in the control group for that day.

- 5                    A dose-response curve was drawn from tests with 4 to 7 different concentrations of the compound tested using the maximal response at a given time (usually one or two hours after administration of the protein). From this curve an ED<sub>50</sub> value was determined
- 10 as that subcutaneous dose ( $\mu\text{g}/\text{kg}$ ) of the protein which gave half the maximal hypoglycemic response. The results are shown in Table III following.

Table III<sup>a</sup>

|    | <u>Compound</u>                  | <u>Maximum<br/>Effect (%)</u> | <u>ED<sub>50</sub><sup>b</sup></u> | <u>Biological<br/>Activity<sup>c</sup></u> |
|----|----------------------------------|-------------------------------|------------------------------------|--|
| 5  | Human Insulin (HI) <sup>d</sup>  | 66                            | 7.95                               | 100  |
|    | Porcine Insulin                  | 60                            | 7.70                               | 103  |
|    | Lys(B28),Pro(B29)HI              | 66                            | 6.8                                | 117  |
|    | Ala(B28),Pro(B29)HI              | 71                            | 15.0                               | 53   |
|    | Asp(B28),Pro(B29)HI              | 64                            | 10.2                               | 78   |
| 10 | Asp(B10),Lys(B28),<br>Pro(B29)HI | 53                            | 10.7                               | 74   |
|    | Gln(B28),Pro(B29)HI              | 69                            | 6.6                                | 120  |
|    | Gly(B28),Pro(B29)HI              | 58                            | 10.5                               | 76   |
|    | Met(B28),Pro(B29)HI              | 51                            | 14.6                               | 54   |
| 15 | Nle(B28),Pro(B29)HI              | 54                            | 7.0                                | 114  |
|    | Orn(B28),Pro(B29)HI              | 60                            | 12.7                               | 63   |
|    | Pro(B29)HI                       | 63                            | 8.7                                | 91   |
|    | Phe(B28),Pro(B29)HI              | 57                            | 9.7                                | 82   |
|    | Val(B28),Pro(B29)HI              | 53                            | 8.7                                | 91   |

20 <sup>a</sup> All values shown refer to blood samples taken one hour after administration of the compound shown

<sup>b</sup> Expressed in  $\mu\text{gm}/\text{kg}$ , subcutaneous

25 <sup>c</sup> Relative to human insulin

<sup>d</sup> Biosynthetic human insulin

As noted previously, the insulin analogs of the present invention have a reduced propensity to dimerize or otherwise self-associate to higher molecular weight forms. Thus, upon administration of one or more of said analogs, a rapid onset of activity is achieved. The insulin analogs of the present invention are effective in treating hyperglycemia by administering to a patient in need thereof an effective amount of an insulin analog of formula I. As used herein the term "effective amount" refers to that amount of one or more insulin analogs of the present invention needed to lower or maintain blood sugar levels either therapeutically or prophylactically. This amount typically may range from about 10 units up to about 60 units or more per day (or about 0.3 to about 2 mg assuming approximately 29 units per mg). However, it is to be understood that the amount of the insulin analog(s) actually administered will be determined by a physician in light of the relevant circumstances including the condition being treated (i.e., the cause of the hyperglycemia) the particular analog to be administered, the chosen parenteral route of administration, the age, weight and response of the individual patient and the severity of the patient's symptoms. Therefore, the above dosage ranges are not intended to limit the scope of the invention in any manner.

The insulin analogs of the invention are administered to a patient in need thereof (i.e., a patient suffering from hyperglycemia) by means of pharmaceutical compositions containing an effective amount of at least one insulin analog of formula I in

combination with one or more pharmaceutically acceptable excipients or carriers. For these purposes, the pharmaceutical compositions may typically be formulated so as to contain about 100 units per ml or similar concentrations containing an effective amount of the insulin analog(s). These compositions are typically, though not necessarily, parenteral in nature and may be prepared by any of a variety of techniques using conventional excipients or carriers for parenteral products which are well known in the art. See, for example, Remington's Pharmaceutical Sciences, 17th Edition, Mack Publishing Company, Easton, PA, USA (1985) which is incorporated herein by reference. For example, dosage forms for parenteral administration may be prepared by suspending or dissolving the desired amount of at least one insulin analog of formula I in a non-toxic liquid vehicle suitable for injection such as an aqueous medium and sterilizing the suspension or solution. Alternatively, a measured amount of the compound may be placed in a vial and the vial and its contents sterilized and sealed. An accompanying vial or vehicle can be provided for purposes of mixing prior to administration. Pharmaceutical compositions adapted for parenteral administration employ diluents, excipients and carriers such as water and water-miscible organic solvents such as glycerin, sesame oil, groundnut oil, aqueous propylene glycol, N,N'-dimethylformamide and the like. Examples of such pharmaceutical compositions include sterile, isotonic, aqueous saline solutions of the insulin analogs of formula I which can be buffered with a pharmaceutically acceptable buffer and which are

pyrogen free. Additionally, the parenteral pharmaceutical formulation may contain preservatives such as meta-cresol or other agents to adjust pH of the final product such as sodium hydroxide or hydrochloric acid.

5           The insulin analogs of the present invention may also be formulated into pharmaceutical compositions suitable for intranasal administration. Such compositions are disclosed in detail in European Patent Application 0200383 A3 which is incorporated herein by refer-  
10           ence. Briefly, such compositions are formulated with one or more pharmaceutically acceptable diluents, a pharmaceutically acceptable amount of an alkali metal salt, the ammonium salt, or the free acid of a substantially zinc-free insulin, and optionally, an absorption  
15           enhancing amount of at least one absorption enhancing agent selected from the group consisting of (1) oleic acid or an ester or salt thereof, (2) a liquid form sorbitan fatty acid ester, (3) a liquid form polyoxy-  
            ethylene derivative of a sorbitan fatty acid ester,  
20           and (4) a liquid form hydroxypolyoxyethylene-polyoxypropylene-polyoxyethylene copolymer.

            To demonstrate the efficacy of intranasal  
            administration of the insulin analog Lys(B28), Pro(B29)  
            human insulin versus human sodium insulin the following  
25           study was conducted. Male beagle dogs weighing 10 to 12 kg were maintained in excellent physical condition prior to and during the course of the study. The dogs were fasted for 16 hours but had access to water to  
            ensure against unexpected fluctuations in insulin  
30           levels. During the entire course of the study the dogs were anesthetized with sodium pentobarbitol and

their body temperature maintained with heating pads to minimize glucose fluctuation. The insulin test samples (i.e., Lys(B28), Pro(B29) human insulin and human sodium insulin) were dissolved in distilled water and the pH of the solution was adjusted to 7.5 with sodium hydroxide. The final insulin concentration was 70 units per ml. An insulin dose of 0.8 units per kg of Lys(B28), Pro(B29) human insulin was administered to each of eight dogs and similarly, a dose of 0.8 units per kg of human sodium insulin was administered to each of four dogs. All doses were administered to the nasal cavity by delivering one spray per nostril with a metered-dose nebulizer and a modified nasal applicator. Blood samples were drawn from the jugular vein at 30, 15 and 0 minutes before administration and at 10, 20, 30, 45, 60, 90, 120, 180 and 240 minutes after administration for the measurement of blood glucose reduction. The results of this study are shown in Table IV.

Table IV<sup>a</sup>

| Time (min) | Lys(B28), Pro(B29) Human Insulin (% of Initial) | Human Sodium Insulin (% of Initial) |            |
|------------|---|-------------------------------------|------------|
| -30        | 95.4 ± 2.5                                      | 101.6 ± 2.4                         |            |
| 25         | 98.2 ± 2.8                                      | 102.4 ± 2.9                         |            |
| 0          | 100   | 100                                 |            |
| 10         | 95.4 ± 2.1                                      | 100.8 ± 1.9                         |            |
| 20         | 83.5 ± 3.3                                      | 96.0 ± 2.1                          |            |
| 30         | 72.2 ± 4.5                                      | 91.6 ± 2.8                          |            |
| 30         | 61.6 ± 6.9                                      | 89.1 ± 2.3                          |            |
| 60         | 60.0 ± 6.0                                      | 93.6 ± 4.5                          |            |
| 90         | 71.6 ± 4.6                                      | 92.2 ± 1.6                          |            |
| 120        | 76.1 ± 3.7                                      | 91.3 ± 2.8                          |            |
| 180        | 91.2 ± 1.8                                      | 93.8 ± 1.8                          |            |
| 35         | 240   | 85.4 ± 2.7                          | 90.7 ± 2.8 |

<sup>a</sup> Blood Glucose levels: mean ± S.E.M.

Utilizing the protocol described above, a comparison of the blood glucose reduction profile of Lys(B28), Pro(B29) human insulin via intranasal administration was compared to both intravenous and subcutaneous administration of the analog as follows.

5 For intravenous administration, the insulin solution containing the Lys(B28), Pro(B29) human insulin analog was administered by bolus injection (0.1 units per kg) into the saphenous vein of each of four dogs. Blood

10 samples were drawn at 30, 15 and 0 minutes before administration and at 5, 10, 15, 20, 30, 45, 60, 90, 120, 180 and 240 minutes after administration. For subcutaneous administration, the insulin solution

15 containing the Lys(B28), Pro(B29) human insulin analog was administered beneath the epidermis (0.2 units per kg) of the outer side of the flank of each of six dogs. Blood samples were drawn at the same protocol as described after nasal administration as described

20 above. The results of this comparative study are shown in Table IV.

TABLE V<sup>a</sup>

|    | <u>Time</u><br><u>(min)</u> | <u>I.V.</u><br><u>(% of initial)</u> | <u>S.C.</u><br><u>(% of initial)</u> | <u>Nasal</u><br><u>(% of initial)</u> |
|----|-----------------------------|--------------------------------------|--------------------------------------|---------------------------------------|
| 5  | -30                         | 96.6 ± 6.5                           | 95.6 ± 2.1                           | 95.4 ± 2.5                            |
|    | -15                         | 101.3 ± 1.5                          | 98.7 ± 2.7                           | 98.2 ± 2.8                            |
|    | 0                           | 100                                  | 100                                  | 100                                   |
| 10 | 5                           | 99.9 ± 2.5                           | -                                    | -                                     |
|    | 10                          | 84.8 ± 4.0                           | 99.2 ± 4.8                           | 95.4 ± 2.1                            |
|    | 15                          | 68.1 ± 2.7                           | -                                    | -                                     |
|    | 20                          | 54.5 ± 3.3                           | 91.9 ± 2.4                           | 83.5 ± 3.3                            |
|    | 30                          | 43.3 ± 2.8                           | 81.6 ± 3.0                           | 72.2 ± 4.5                            |
|    | 45                          | -                                    | 55.8 ± 3.5                           | 61.6 ± 6.9                            |
| 15 | 60                          | 55.3 ± 3.1                           | 35.3 ± 3.1                           | 60.6 ± 6.0                            |
|    | 90                          | 77.2 ± 6.4                           | 39.5 ± 3.4                           | 71.6 ± 4.6                            |
|    | 120                         | 85.2 ± 4.7                           | 36.1 ± 2.6                           | 76.1 ± 3.7                            |
| 20 | 180                         | 97.1 ± 6.4                           | 64.5 ± 7.0                           | 91.2 ± 1.8                            |
|    | 240                         | 100.0 ± 9.6                          | 81.3 ± 2.4                           | 85.4 ± 2.7                            |

<sup>a</sup> Blood glucose levels: mean ± S.E.M.



Microorganisms were deposited with the Agricultural Research Culture Collection, of 1815 North University Street, Peoria, Illinois 61064, United States of America as set out below:

Microorganism      Escherichia coli MCB 13 K12/pKC283  
Deposit Date:      3 August 1984  
Accession No.:      NRRL B-15830

Microorganism      Escherichia coli MCB2605  
Deposit Date:      28 September 1983  
Accession No.:      NRRL B-15624


Microorganism      Escherichia coli MCB 3605 K12 (MO) ( $\lambda$ +)  
Deposit Date:      14 August 1985  
Accession No.:      NRRL B-15993

Microorganism      Escherichia coli MCB 2619 K12 RV308/pNM789  
Deposit Date:      4 May 1987  
Accession No.:      NRRL B-18216.

~~What we claim is:~~

**The claims defining the invention are as follows:**

1. An insulin analog of the formula



or a pharmaceutically acceptable salt thereof, wherein

35 A21 is alanine, asparagine, aspartic acid, glutamine, glutamic acid, glycine, threonine, or serine; B1 is phenylalanine, aspartic acid, or is absent; B2 is valine or may be absent when B1 is absent; B3 is asparagine or aspartic acid; B9 is serine or aspartic acid; B10 is

40 histidine or aspartic acid; B27 is threonine or is absent; B28 is any amino acid, B29 is L-proline, D-proline,

D-hydroxyproline, L-hydroxyproline, L-(N-methyllysine),  
D-lysine, L-(N-methylarginine) or D-arginine; B30 is  
alanine, threonine or is absent; Z is -OH, -NH<sub>2</sub>, -OCH<sub>3</sub>,  
or -OCH<sub>2</sub>CH<sub>3</sub>; X is Arg, Arg-Arg, Lys, Lys-Lys, Arg-Lys,  
5 Lys-Arg, or is absent; and Y may be present only when  
X is present and, if present, is Glu or an amino acid  
sequence which comprises all or a portion of the sequence  
-Glu-Ala-Glu-Asp-Leu-Gln-Val-Gly-Gln-Val-Glu-Leu-Gly-  
10 Glu-Gly-Ser-Leu-Gln-Lys-Arg- and which begins at the  
N-terminus Glu of such sequence.

2. The insulin analog as claimed in Claim 1,  
in which X and Y are absent and Z is OH.

3. The insulin analog as claimed in Claim 1  
15 or 2, wherein B29 is L-proline, D-proline, D-hydroxy-  
proline or L-hydroxyproline; and B30 is threonine.

4. The insulin analog as claimed in any one  
of Claims 1 to 3 wherein B28 is aspartic acid, valine,  
leucine, isoleucine, norleucine, proline, arginine,  
20 histidine, citrulline, ornithine, lysine, phenylalanine,  
alanine or glycine.

5. The insulin analog as claimed in any one  
of Claims 1 to 4 wherein B28 is aspartic acid, valine,  
leucine, isoleucine, norleucine, proline, arginine,  
25 histidine, ornithine or lysine.

6. The insulin analog as claimed in any one  
of Claims 1 to 5 wherein B28 is lysine.

7. The insulin analog as claimed in any one  
of Claims 1 to 6 wherein B28 is lysine; B29 is  
30 L-proline; and B30 is threonine.

8. The insulin analog as claimed in any one of Claims 1 to 7 wherein A21 is alanine.

9. The insulin analog as claimed in any one of Claims 1 to 8 wherein B1 is absent.

5 10. The insulin analog as claimed in any one of Claims 1 to 9 wherein B10 is aspartic acid.

11. The insulin analog as claimed in any one of Claims 1 to 10 wherein B2 is absent.

10 12. The insulin analog as claimed in Claim 11 wherein B10 is aspartic acid.

13. The insulin analog as claimed in any one of Claims 1 to 12 wherein B30 is absent.

14. The insulin analog as claimed in any one of Claims 1 to 13 wherein B30 is alanine.

15 15. The insulin analog as claimed in Claim 1 or 2, wherein B28 is L-proline, and B29 is L-(N-methyllysine), D-lysine, L-(N-methylarginine) or D-arginine.

20 16. The insulin analog as claimed in any one of Claims 1, 2 and 15, wherein B28 is L-proline and B29 is L-(N-methyllysine) or D-lysine.

17. The insulin analog as claimed in any one of Claims 1, 2, 15 and 16 wherein A21 is alanine.

25 18. The insulin analog as claimed in any one of Claims 1, 2, and 15 to 17, wherein B1 is absent.

19. The insulin analog as claimed in any one of Claims 1, 2 and 15 to 18, wherein B10 is <sup>histidine</sup>~~aspartic acid~~.

5 20. The insulin analog as claimed in any one of Claims 1, 2, and 15 to 19, wherein B2 is absent.

21. The insulin analog as claimed in any one of Claims 1, 2, and 15 to <sup>18</sup>~~20~~, wherein B10 is aspartic acid.

10 22. The insulin analog as claimed in any one of Claims 1, 2, and 15 to 21, wherein B30 is absent.

23. The insulin analog as claimed in any one of Claims 1, 2, and 15 to <sup>21</sup>~~22~~, wherein B30 is alanine.

15 24. The insulin analog as claimed in Claim 1 wherein A21 is asparagine, B1 is phenylalanine, B2 is valine, B3 is asparagine, B9 is serine, B10 is histidine, B27 is threonine, B28 is lysine, B29 is L-proline, B30 is threonine, Z is OH, X is absent, and Y is absent.

20 25. A process for preparing a compound of Formula (I) and as defined in any one of Claims 1 to 24, which comprises:

25 A) combining the A-chain of a compound of Formula (I) in which A21 is as defined in Claim 1, with the B-chain of a compound of Formula (I) in which B1, B2, B3, B9, B10, B27, B28, B29, B30, X, Y, and Z are as defined in Claim 1 to form the appropriate disulfide bonds; or



5 B) reacting a modified compound of Formula (I) lacking the peptide sequence beginning at B23 and in which A21, B1, B2, B3, B9 and B10 are as defined in Claim 1 with a peptide having the sequence Gly-Phe-Phe-Tyr-B27-B28-B29-B30-X-Y-Z in which B27, B28, B29, B30, X, Y, and Z are as defined in Claim 1; or

10 C) cleaving a modified proinsulin molecule comprising the amino acid sequence of a compound of Formula (I) in which A21, B1, B2, B3, B9, B10, B27, B28, B29, B30, X, Y, and Z are as defined in Claim 1 to  
15 remove all or a portion of the C-peptide from such modified proinsulin.

20 26. A pharmaceutical formulation comprising, as an active ingredient, a compound of Formula (I) as claimed in any one of Claims 1 to 24, associated with one or more pharmaceutically-acceptable carriers therefor.

25 27. A method for treating diabetes mellitus in mammals afflicted therewith, which comprises administering to such mammal a compound of Formula (I) as claimed in any one of Claims 1 to 24.

28. A compound of Formula (I) as claimed in any one of Claims 1 to 24, whenever produced by the process of Claim 25.



29. An insulin analog as defined in Claim 1 and substantially as hereinbefore described with reference to any one of the Examples.

30. A process for preparing an insulin analog as defined in Claim 1, which process is substantially as hereinbefore described with reference to any  
5 one of Examples 1 to 28.

31. A plasmid for the expression of proinsulin from which an insulin analog as defined in Claim 1 may be derived, which plasmid is substantially as hereinbefore described with reference to Example 29.

32. A plasmid for the expression of proinsulin from which an insulin  
10 analog as defined in Claim 1 may be derived, which plasmid is substantially as hereinbefore described with reference to any one of Figures 17 to 20.

33. A process for preparing a plasmid for the expression of proinsulin from which an insulin analog as defined in Claim 1 may be derived, which process is substantially as hereinbefore described with reference to Example  
15 29.

34. A pharmaceutical composition for the treatment of diabetes mellitus in a mammal, which composition comprises an amount effective for the treatment of diabetes mellitus in said mammal of an insulin analog according to Claim 29 together with a pharmaceutically acceptable carrier, diluent and/or  
20 adjuvant.

35. A method of treating diabetes mellitus in a mammal, which method comprises administering to a mammal in need of treatment for diabetes mellitus, an amount effective for the treatment of diabetes mellitus in said mammal of an insulin analog according to Claim 29, or a composition  
25 according to Claim 34.

DATED this SECOND day of JULY 1992

Eli Lilly and Company

Patent Attorneys for the Applicant

SPRUSON & FERGUSON

