

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 200124463 B2
(10) Patent No. 780136

(54) Title
Method for preparing solid delivery system for encapsulated and non-encapsulated pharmaceuticals

(51)⁷ International Patent Classification(s)
A61K 009/50 A61K 009/10

(21) Application No: **200124463**

(22) Application Date: **2000.12.20**

(87) WIPO No: **WO01/49272**

(30) Priority Data

(31) Number	(32) Date	(33) Country
09/476483	1999.12.30	US
09/656297	2000.09.06	US

(43) Publication Date : **2001.07.16**

(43) Publication Journal Date : **2001.09.27**

(44) Accepted Journal Date : **2005.03.03**

(71) Applicant(s)
Shear/Kershman Laboratories, Inc.

(72) Inventor(s)
Jeff Shear; Al Kershman

(74) Agent/Attorney
SPRUSON and FERGUSON,GPO Box 3898,SYDNEY NSW 2001

(56) Related Art
WO 1999/147122
US 4581381
US 4749575

AU 200124463

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 July 2001 (12.07.2001)

PCT

(10) International Publication Number
WO 01/49272 A3

- (51) International Patent Classification⁷: A61K 9/50, 9/10
- (21) International Application Number: PCT/US00/34776
- (22) International Filing Date:
20 December 2000 (20.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/476,483 30 December 1999 (30.12.1999) US
09/656,297 6 September 2000 (06.09.2000) US
- (71) Applicant (for all designated States except US):
SHEAR/KERSHMAN LABORATORIES, INC.
[US/US]; Suite F, 701 Crown Industrial Court, Chesterfield, MO 63005 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): SHEAR, Jeff [US/US]; 1421 Wildhorse Parkway Drive, Chesterfield, MO 63005 (US). KERSHMAN, Al [US/US]; 1220 Woodbridge Crossing Drive, Chesterfield, MO 63005 (US).
- (74) Agent: ROBBINS, Glenn, K., II: Greensfelder, Hemker & Gale, P.C., 2000 Equitable Building, 10 South Broadway, St. Louis, MO 63102-1774 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- (88) Date of publication of the international search report:
31 January 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/49272 A3

(54) Title: METHOD FOR PREPARING SOLID DELIVERY SYSTEM FOR ENCAPSULATED AND NON-ENCAPSULATED PHARMACEUTICALS

(57) Abstract: The present invention is an oral drug delivery system for delivering unpalatable pharmaceuticals, wherein the pharmaceutical delivery system comprises a lipid, dry particles including at least one pharmaceutical and at least one filler, and a surfactant, wherein the dry particles are continuously coated by the lipid and form a suspension with the lipid, making the pharmaceutical more palatable.

METHOD FOR PREPARING SOLID DELIVERY SYSTEM FOR ENCAPSULATED AND NON-ENCAPSULATED PHARMACEUTICALS

Field of the Invention

5 The present invention relates to oral delivery systems suitable for human and
veterinary applications. More specifically, the present invention relates to an oral
delivery system for cats, dogs and horses, wherein the normally unpalatable
pharmaceutical, such as a drug, vitamin, mineral or food supplement, is made palatable,
and therefore easily administered. The pharmaceutical is administered in a solid dosage
10 form, wherein the pharmaceutical is carried in a lipid suspension.

Background of the Invention

It is known by those skilled in the art that orally administering pharmaceuticals to
animals, specifically cats, dogs and horses, can be exceedingly difficult. The animal may
reject the entire dose, or parts of the dose, leaving the vet or the owner frustrated in trying
15 to administer the proper amount.

The true test of a delivery system for animals is whether the pharmaceutical is
successfully administered repeatedly. As is often the case, when treating an illness or
even a permanent condition of an animal, doses are repeated daily or even several times a
day. Once an animal has tasted the unpalatable pharmaceutical that the vet or owner has
20 attempted to hide or disguise, it becomes uncooperative when the administration is
repeated. Applicants have discovered a delivery system that presents the pharmaceutical

in a palatable form such that the drug can be successfully and easily administered repeatedly.

Heretofore, attempts to alleviate the problem of administering pharmaceuticals have been made a number of ways, including hiding the drug in food. Usually the keen smell of the animal will allow it to detect and reject the pharmaceutical hidden therein. The taste and smell of pharmaceuticals can be masked through the use of flavorings and/or masking agents. Typically, this is only partially successful, as the taste and smell are not completely eliminated.

The pharmaceutical may be microencapsulated, and administered in a gel capsule, but the animal will usually chew the capsule, rupture it, taste the unpalatable pharmaceutical and reject it. Future administrations of the encapsulated pharmaceuticals are thwarted. The use of gel capsules has the additional disadvantages of being costly for delivering large doses of pharmaceuticals. The microencapsulated pharmaceutical may be tableted using a conventional tablet press. However, such a process invariably damages or ruptures the capsules, defeating the benefits of encapsulation.

The present invention is an oral delivery system, wherein the system delivers microencapsulated pharmaceuticals, wherein the microencapsulation is intact and the pharmaceutical is delivered in a palatable form. Further, the present invention provides the pharmaceutical in a low cost, concentrated form that is easily administered. The delivery system comprises at least one lipid, dry particles including at least one pharmaceutical and at least one filler, and a surfactant, wherein the dry particles are continuously coated by the lipid and form a suspension with the lipid. The suspension exhibits thixotropic or pseudoplastic flow properties. The fillers include cellulose, starch and whey, and comprise from about 60 to 80 % of the system (all percentages stated herein are weight percent, unless otherwise indicated). Optionally, flavorings can be added to the dry particles and include cheese, butter, cream, and egg flavorings. Optionally, the pharmaceutical can comprise all of the dry particles, and serve as filler as well as pharmaceutical.

The process for preparing the present delivery system comprises melting the lipid and mixing with the surfactant, blending the dry particles

which include the pharmaceutical, filler and, optionally, flavorings with the melted lipid, and pouring or molding the suspension to provide the dose.

U.S. patent 4,327,076 discloses a compressed chewable antacid tablet which contains from about 10 to 50 % active ingredient, from about 2 to 45 %
5 fat, from about 25 to 75% fat-sorbing materials, such as starch, and from about 20 to 60% tablet bonders, such as sugars, and surfactants and flavorings. The process disclosed by 4,327,076 involves a pretreatment step wherein the fat is melted and is mixed with fat-sorbing material, tablet bonders and flavorings, forming a fatty powder. The second step involves mixing the active ingredient
10 and surfactant with the pretreated fat and forming a free flowing powder wherein each active ingredient particle is coated with the pretreated fat. The powder thus produced is compressed into chewable pellets.

The above patent fails to disclose the claimed delivery system, wherein the pellets are not compressed into tablets but molded or poured.

U.S. patent 4,581,381 discloses a chewable antacid pill or pellet
15 containing solid antacid particles having a particle size under 100 millimicrons coated with a fatty material, a surfactant, and a flavoring. The pill or pellet is formed by molding and provides a non-chalky, non-gritty pellet. The antacid particles are coated by melting the fatty material and mixing in the surfactant,
20 the antacid and the flavorings.

The above patent requires very small particles (100 millimicrons), which are not required in the present process.

Neither of the above-referenced patents disclosed the present invention as disclosed and claimed herein.

Summary of the Invention

The present invention relates to an oral delivery system comprising at least one lipid, at least one surfactant, dry particles including at least one pharmaceutical, and at least one filler, wherein the dry particles are continuously coated by the lipid and form a homogeneous suspension with the lipid. The suspension exhibits thixotropic and/or pseudoplastic properties. The suspension is formed into the desired dose by molding or pouring the suspension when in a liquid or semi-liquid state.

According to a first aspect of the invention there is provided an oral pharmaceutical delivery system comprising:

- A. at least one lipid;
 - B. at least one surfactant; and
 - C. dry particles;
- wherein, the dry particles contain at least one pharmaceutical;
- wherein, the dry particles are continuously coated with the lipid and form a homogeneous suspension with the lipid;
- wherein the suspension exhibits pseudoplastic and/or thixotropic properties,
- wherein the suspension is formed or shaped into the appropriate solid dosage form by molding or pouring the suspension when in a liquid or semi-liquid state; and
- wherein said pharmaceutical is selected from the group consisting of analgesics, antibiotics, anti-inflammatory agents, cardiovascular preparations, gastrointestinal medications, hormone products, and laxatives.

According to a second aspect of the invention there is provided an oral pharmaceutical delivery system comprising:

- A. at least one lipid;
 - B. at least one surfactant; and
 - C. dry particles;
- wherein, the dry particles contain at least one pharmaceutical;
- wherein, the dry particles are continuously coated with the lipid and form a homogeneous suspension with the lipid;
- wherein the suspension exhibits pseudoplastic and/or thixotropic properties;
- and
- wherein the suspension is formed or shaped into the appropriate solid dosage form by molding or pouring the suspension when in a liquid or semi-liquid state; and

wherein said pharmaceutical particles are present in said suspension as a microencapsulated particle.

According to a third aspect of the invention there is provided an oral pharmaceutical delivery system comprising:

- 5 A. at least one lipid;
 B. at least one surfactant; and
 C. dry particles;
 wherein, the dry particles contain at least one pharmaceutical;
 wherein, the dry particles are continuously coated with the lipid and form a
10 homogenous suspension with the lipid;
 wherein the suspension exhibits pseudoplastic and/or thixotropic properties;
and

 wherein the suspension is formed or shaped into the appropriate solid dosage
form by molding or pouring the suspension when in a liquid or semi-liquid state; and
15 wherein said lipid suspension contains a rupturing agent.

According to a fourth aspect of the invention there is provided an oral pharmaceutical delivery system comprising:

- 20 A. at least one lipid;
 B. at least one surfactant; and
 C. dry particles;
 wherein, the dry particles contain at least one pharmaceutical;
 wherein, the dry particles are continuously coated with the lipid and form a
homogenous suspension with the lipid;
 wherein the suspension exhibits pseudoplastic and/or thixotropic properties;
25 and
 wherein the suspension is formed or shaped into the appropriate solid dosage
form by molding or pouring the suspension when in a liquid or semi-liquid state; and
 wherein said filler comprises whey.

According to a fifth aspect of the invention there is provided a method for preparing
30 an oral pharmaceutical delivery system comprising:

- melting at least one lipid;
 mixing in at least one surfactant to said melted lipid;
 mixing in dry particles to said melted lipid and surfactant,
 wherein said dry particles contain at least one pharmaceutical, and at least one
35 filler;

wherein the dry particles are continuously coated with the lipid and form a homogenous suspension with the lipid;

wherein the suspension exhibits pseudoplastic and/or thixotropic properties,

wherein the suspension is formed or shaped into the appropriate dose by molding or pouring the suspension when in a liquid or semi-liquid state; and

wherein said pharmaceutical is selected from the group consisting of analgesics, antibiotics, anti-inflammatory agents, cardiovascular preparations, gastrointestinal medications, hormone products, and laxatives.

According to a sixth aspect of the invention there is provided a method for preparing an oral pharmaceutical delivery system comprising:

melting at least one lipid;

mixing in at least one surfactant to said melted lipid;

mixing in dry particles to said melted lipid and surfactant, wherein said dry particles contain at least one pharmaceutical, and at least one filler;

wherein the dry particles are continuously coated with the lipid and form a homogenous suspension with the lipid;

wherein the suspension exhibits pseudoplastic and/or thixotropic properties;

wherein the suspension is formed or shaped into the appropriate dose by molding or pouring the suspension when in a liquid or semi-liquid state; and

wherein said lipid suspension contains a rupturing agent.

According to a seventh aspect of the invention there is provided a method for preparing an oral pharmaceutical delivery system comprising:

melting at least one lipid;

mixing in at least one surfactant to said melted lipid;

mixing in dry particles to said melted lipid and surfactant, wherein said dry particles contain at least one pharmaceutical, and at least one filler;

wherein the dry particles are continuously coated with the lipid and form a homogenous suspension with the lipid;

wherein the suspension exhibits pseudoplastic and/or thixotropic properties,

wherein the suspension is formed or shaped into the appropriate dose by molding or pouring the suspension when in a liquid or semi-liquid state; and

wherein said dry particles include artificial flavourings.

According to an eighth aspect of the invention there is provided a method for preparing an oral pharmaceutical delivery system comprising the steps of:

microencapsulating pharmaceutical particles;

melting at least one lipid;


mixing in a surfactant to said melted lipid;

5 dry-mixing dry particles comprising at least one filler and said
microencapsulated pharmaceutical;

mixing the dry particles with said melted lipid to form a suspension such that
said dry particles are continuously coated by said lipid such that said suspension exhibits
pseudoplastic and/or thixotropic properties; and

pouring or molding said suspension into a dosage form.

10 According to a ninth aspect of the invention there is provided an oral
pharmaceutical delivery system prepared according to the method of any one of the fifth
to eighth aspects of the invention.



Detailed Description of the Invention

The lipids of the present invention may be of animal, vegetable or mineral origin, which are substantially water-insoluble, inert, non-toxic hydrocarbon fats and oils and derivatives thereof, and may comprise any of the commonly commercially available fats or oils approved by the Food & Drug Administration, having melting points in the range of about 90 to 160°F. The lipid may comprise a vegetable oil base commonly known as hard butter. Hard butters are hydrogenated, press fractionated or other processed oils that are processed or recombined to have a solid fat index (percent solid fat vs. temperature) similar to that of cocoa butter. However, other lipids may be used that are relatively hard or solid at room temperature, but melt rapidly in the mouth at a temperature of about 98°F (mouth temperature). The lipid is employed in the amounts within the range of from about 20 to 40%. Above about 40%, the suspension flows too readily and does not exhibit thixotropic or pseudoplastic flow properties. When present below about 20%, the amount of lipid is not sufficient to completely coat the dry particles.

Examples of suitable lipids include tallow, hydrogenated tallow, hydrogenated vegetable oil, almond oil, coconut oil, corn oil, cottonseed oil, light liquid petrolatum, heavy liquid petrolatum, olein, olive oil, palm oil, peanut oil, persic oil, sesame oil, soybean oil or safflower oil. Additionally, stearines can be used as a lipid in the present invention. The addition of stearines to the product provides the favorable property of mold-release. Further, the addition of stearines raises the melting point of the composition as high as about 100°F, which is particularly beneficial when the product is shipped or stored in unrefrigerated compartments.

The fillers of the present invention are pharmacologically inert and optionally nutritionally beneficial to humans and animals. Such fillers include cellulose such as microcrystalline cellulose, grain starches such as cornstarch, tapioca, dextrin, sugars and sugar alcohols such as sucrose sorbitol, xylitol, mannitol and the like. Preferred fillers include non-fat milk powder, whey, grain brans such as oat bran, and fruit and vegetable pulps. Preferred fillers are finely divided and have a preferred average particle size in the range of

about 10 to 500 microns. The fillers are present in the drug delivery device in a concentration of about 60 to 80%. Optionally, the pharmaceutical particles can also serve as filler in the delivery system.

5 Any emulsifier or surfactant approved for use in foods by the Food and Drug Administration and having a relatively low HLB value, in the range of about 1 to 3, is suitable for use in the present invention. The appropriate surfactant minimizes the surface tension of the lipid, allowing it to oil wet and encapsulate the non-oil solid particles. Typically, the surfactant is present in the delivery system in the concentration of about 0.1 to 1.0%. Suitable
10 surfactants include alkyl aryl sulfonate, or alkyl sulfonates, or sulfonated amides or amines, or sulfated or sulfonated esters or ethers, or alkyl sulfonates, of dioctyl sulfonosuccinate and the like, or a hydrated aluminum silicate such as bentonite or kaolin, triglycerol monostearate, triglycerol monoshortening, octaglycerol monooleate, octaglyceron monostearate, and
15 decaglycerol decaloeate. The preferred surfactant is lecithin.

In the preferred embodiment, the pharmaceutical is microencapsulated. Such microencapsulation includes sustained release encapsulation. Any known method of encapsulation is suitable in the present invention. A preferred method involves slowly blending the drug with a filming agent
20 solution to form granulated particles. The granulated particles are allowed to dry on a tray and are sieved to the desired size, typically in the range of from about 200 to 500 microns. In another embodiment of the present invention, the pharmaceutical is not microencapsulated, but suspended in the lipid as dry particles. Typically the pharmaceutical is present in the delivery device in a
25 concentration of 10% or less. However, the pharmaceutical can comprise all of the dried particles, to provide the necessary dose.

The pharmaceuticals contemplated in the present invention are administered orally. The pharmaceuticals include, but are not limited to, drugs such as analgesics, anti-inflammatory agents, gastrointestinal medications,
30 hormone products, cardiovascular preparations, laxatives and antibiotics. Specific drugs include aspirin, acetaminophen, carprofen, enalapril maleate, furosemide, levothyroxine sodium and prednisolone. Pharmaceuticals further includes vitamins and minerals, as are well known in the art. Pharmaceuticals

also includes synthetic and natural food supplements, such as glucosamine, chondroitin, bee pollin, St. John's wort, echninaesia, etc. Additional pharmaceuticals are contemplated for the present invention, and are disclosed in U.S. patent 4,369,172, which is hereby incorporated by reference.

5 Optionally, the dry particles include flavorings that make the device taste and smell appealing to humans or animals. The flavorings can be natural or synthetic, and can include butter, milk, cream, egg or cheese. The flavorings are typically present in the device in the range of about 0.05 to 50.0%.

10 The delivery device may also include other pharmaceutically acceptable agents, such as sweetening agents, including hydrogenated starch hydrolysates, synthetic sweeteners such as sorbitol, xylitol, saccharin salts, L-aspartyl-L-phenylalanine methyl ester, as well as coloring agents, other binding agents, lubricants, such as calcium stearate, stearic acid, magnesium
15 stearate, antioxidants such as butylated hydroxy toluene, antifatulants such as simethicone and the like.

 Optionally, rupturing agents are used to rapidly deliver the pharmaceutical into the recipient's system. A typical rupturing agent is a starch that swells in the presence of water. A preferred rupturing agent is
20 sodium starch glycolate. When ingested, the capsule or pellet swells in the presence of gastric juices and ruptures. In one embodiment of the present invention, the rupturing agent is present inside the microcapsule. As water penetrates the microcapsule, it swells the starch and ruptures the capsule, rapidly delivering the pharmaceutical to the system.

25 In another embodiment, the rupturing agent is present in the lipid suspension, which ruptures the pellet, but leaves the microcapsules intact. This allows the delayed delivery of the drug farther along in the digestive system, or in the intestines. The present invention is particularly effective in this embodiment, in that the ingested pellet is chewable, yet the pellet cleaves
30 in the lipid suspension when chewed, leaving the microcapsules intact. Tablets or gel capsules, when chewed, typically result in damage to or rupturing of the microcapsules defeating the effectiveness of the microcapsules.

In yet another embodiment, multiple drugs have multiple encapsulations, each containing an rupturing agent. The filming agents used for encapsulation are selected to disintegrate at selected pH conditions, which rupture and release each drug at desired locations in the digestive system.

5 The process for preparing the above delivery system comprises melting the lipid and mixing with the surfactant. The dry particles are mixed with the melted lipid mixture to form a suspension exhibiting pseudoplastic and/or thixotropic flow properties, and poured or molded to provide solid dosage forms.

10 The dry particles, which include the pharmaceutical, filler and optional flavorings and additives, are pre-blended and typically have a particle size in the range of from about 50 to 150 microns. The pre-blended particles are gradually added to the heated lipid base until a high solid suspension is obtained, typically in the range of about 60 to 80% particles and from about 40
15 to 20 % lipid.

 Slow addition of the dry particles is critical in the production of the device, to insure that the particles are suspended in their micronized state and not as agglomerated clumps. Moreover, rapid addition can cause the mixing process to fail in that the melted suspension will not have the desired flow
20 properties, but instead will be a granular oily mass (a sign of product failure). The mixing step is accomplished in a heated mixing device that insures thorough mixing of all materials with minimal shear, such as a planetary mixer or a scrape surface mixer. After the suspension is formed, the product is poured into molds and allowed to cool. De-molding and packaging are then
25 performed. Alternatively, the suspension can be super-cooled and sheeted in a semi-soft format. The sheet is processed through forming rolls containing a design or configuration that embosses and forms the final shape.

 The following examples are to illustrate the claimed invention and are not intended to limit the claims in any way. All of the percentages are by
30 weight unless otherwise indicated.

Examples

Examples I-V were prepared according to the following procedure.

5 Microencapsulation of the Drug

The pharmaceutical particles were dry-blended with the rupturing agent, (sodium starch glycolate sold under the trademark Explotab®) in a Hobart 5 Quart planetary mixer jacketed with a heating mantle. A 4% mixture of the filming agent, (ethylcellulose, sold commercially under the trademark Ethocel®) and ethanol was slowly added to the dry mix thereby forming wet granules. The wet granules were tray dried and screened to a particle of between about 200 and 500 microns.

Forming the Suspension

The first lipid (vegetable stearines sold under the trademark Duratex®) was heated in a Hobart 5 Quart planetary mixer jacketed with a heating mantle in the range of about 140 to 150°F and melted. The second lipid (98°F vegetable hard butter sold under the trademark Kalomel®) was added to the jacketed mixer and melted with mixing. The surfactant, lecithin, was added to the lipids with mixing, and the mixture was allowed to cool to about 135°F.

The dry particles, including the microencapsulated pharmaceutical, the flavorings, and the filler (whey) were screened to a particle size in the range of about 200 and 500 microns and dry-blended. The dry particles were slowly added incrementally to the lipid/surfactant mixture with mixing over a period of about 1 hour, to provide a smooth suspension with no lumps or agglomerations. The suspension was molded and cooled to about 70°F. The suspension shrank as it cooled, and easily released from the mold when inverted.

EXAMPLE I

TABLE A
Microencapsulation of Carprofen

BATCH FORMULA		
Ingredient	Weight (grams)	%
Carprofen (Pharmaceutical)	7.580	18.95%
Ethycellulose (30gm of 4% ethanol solution)	1.200	3.00%
Sodium starch gluconate	31.220	78.05%
<i>Totals</i>	40	100.00%

TABLE B
Forming a Suspension of
Carprofen in a 100mg Dose

5

BATCH FORMULA		
Ingredient	Weight (grams)	%
Vegetable stearine	0.4536	25.20%
Vegetable hard butter	0.1944	10.80%
Lecithin	0.0108	0.60%
Chedlong #1 (flavoring)	0.2430	13.50%
Cheese flavor 2517	0.0090	0.50%
Cheese flavor 1200s	0.0270	1.50%
Salt	0.0180	1.00%
Whey	0.2642	14.68%
Microencapsulated carprofen (drug from table A)	0.5800	32.22%
Totals	1.8	100.00%

EXAMPLE II

TABLE A
Microencapsulation of Enalapril Maleate

5

BATCH FORMULA		
Ingredient	Weight (grams)	%
Enalapril maleate (Pharmaceutical)	1.460	3.65%
Ethylcellulose (20gm of 4% ethanol solution)	0.800	2.00%
Sodium starch glycolate	37.740	94.35%
Totals	40	100.00%

10

TABLE B
Forming a Suspension of
Enapril in a 5mg Dose

BATCH FORMULA		
Ingredient	Weight (grams)	%
Vegetable stearines	0.3780	25.20%
Vegetable hard butter	0.1620	10.80%
Sodium starch glycolate	0.2535	16.90%
Lccithin	0.0090	0.60%
Chedlong #1 (flavoring)	0.2025	13.50%
Cheese flavor 2517	0.0075	0.50%
Cheese flavor 1200s	0.0225	1.50%
Salt	0.0150	1.00%
Whey	0.3000	20.00%
Microencapsulated enapril (From Table A)	0.1500	10.00%
Totals	1.5	100.00%

EXAMPLE III

TABLE A
Microencapsulation of Furosemide

5

BATCH FORMULA		
Ingredient	Weight (grams)	%
Furosemide (Pharmaceutical)	3.66	9.15%
Ethylcellulose (20gm of 4% ethanol solution)	0.80	2.00%
Sodium starch glycolate	35.54	88.85%
<i>Totals</i>	40	100.00%

10

TABLE B
Forming a Suspension of Furosemide in a 12.5mg Dose

BATCH FORMULA		
Ingredient	Weight (grams)	%
Vegetable stearines	0.3780	25.20%
Vegetable hard butter	0.1620	10.80%
Sodium starch glycolate	0.2535	16.90%
Lecithin	0.0090	0.60%
Chedlong #1 (flavoring)	0.2025	13.50%
Cheese flavor 2517	0.0075	0.50%
Cheesc flavor 1200s	0.0225	1.50%
Salt	0.0150	1.00%
Whey	0.3000	20.00%
Microencapsulated furosemide (From Table A)	0.1500	10.00%
<i>Totals</i>	1.5	100.00%

EXAMPLE IV

TABLE A
Microencapsulation of Levothyroxine Sodium

5

BATCH FORMULA		
Ingredient	Weight (grams)	%
Levothyroxine Sodium (Pharmaceutical)	0.120	0.30%
Ethylcellulose (20gm of 4% ethanol solution)	0.800	2.00%
Sodium starch glycolate	39.080	97.70%
<i>Totals</i>	40	100.00%

10

TABLE B
Forming a Suspension of Levothyroxine in a 0.4mg Dose

BATCH FORMULA		
Ingredient	Weight (grams)	%
Vegetable stearines	0.3780	25.20%
Vegetable hard butter	0.1620	10.80%
Sodium starch glycolate	0.2535	16.90%
Lecithin	0.0090	0.60%
Chedlong #1 (flavoring)	0.2025	13.50%
Cheese flavor 2517	0.0075	0.50%
Cheese flavor 1200s	0.0225	1.50%
Salt	0.0150	1.00%
Whey	0.3000	20.00%
Microencapsulated Levothyroxine (From Table A)	0.1500	10.00%
<i>Totals</i>	1.5	100.00%

EXAMPLE V

TABLE A
Microencapsulation of Prednisolone

5

BATCH FORMULA		
Ingredient	Weight (grams)	%
Prednisolone (Pharmaceutical)	1.460	3.65%
Ethylcellulose (20gm of 4% ethanol solution)	0.800	2.00%
Sodium starch glycolate	37.740	94.35%
Totals	40	100.00%

10

TABLE B
Forming a Suspension of
Prednisolone in a 5mg Dose

BATCH FORMULA		
Ingredient	Weight (grams)	%
Vegetable stearines	0.3780	25.20%
Vegetable hard butter	0.1620	10.80%
Sodium starch glycolate	0.2535	16.90%
Lecithin	0.0090	0.60%
Chedlong #1 (flavoring)	0.2025	13.50%
Cheese flavor 2517	0.0075	0.50%
Cheese flavor 1200s	0.0225	1.50%
Salt	0.0150	1.00%
Whey	0.3000	20.00%
Microencapsulated prednisolone (From Table A)	0.1500	10.00%
Totals	1.5	100.00%

The claims defining the invention are as follows:

1. An oral pharmaceutical delivery system comprising:
- A. at least one lipid;
 - B. at least one surfactant; and
 - C. dry particles;
- wherein, the dry particles contain at least one pharmaceutical;
- wherein, the dry particles are continuously coated with the lipid and form a homogenous suspension with the lipid;
- wherein the suspension exhibits pseudoplastic and/or thixotropic properties,
- wherein the suspension is formed or shaped into the appropriate solid dosage form by molding or pouring the suspension when in a liquid or semi-liquid state; and
- wherein said pharmaceutical is selected from the group consisting of analgesics, antibiotics, anti-inflammatory agents, cardiovascular preparations, gastrointestinal medications, hormone products, and laxatives.
2. The pharmaceutical delivery system of claim 1 in which said lipid forms 20% to 40% by weight of said suspension, and said dry particles form 60% to 80% by weight of said suspension.
3. The pharmaceutical delivery system of claim 1 or 2 in which said surfactant is lecithin.
4. The pharmaceutical delivery system of claim 1 or 2 in which said surfactant has a hydrophilic to lipophilic balance ratio in a range of between about 1 to about 3.
5. The pharmaceutical delivery system of any one of claims 1 to 4 in which said lipid comprises a hard butter
6. The pharmaceutical delivery system of any one of claims 1 to 4 in which said lipid comprises a combination of hard butters and stearines.
7. The pharmaceutical delivery system of any one of claims 1 to 6 in which said pharmaceutical is present in the suspension in a concentration range of about 0.1 to 50 %.
8. An oral pharmaceutical delivery system comprising:
- A. at least one lipid;
 - B. at least one surfactant; and
 - C. dry particles;
- wherein, the dry particles contain at least one pharmaceutical;
- wherein, the dry particles are continuously coated with the lipid and form a homogenous suspension with the lipid;

wherein the suspension exhibits pseudoplastic and/or thixotropic properties;
and

wherein the suspension is formed or shaped into the appropriate solid dosage form by molding or pouring the suspension when in a liquid or semi-liquid state; and

5 wherein said pharmaceutical particles are present in said suspension as a microencapsulated particle.

9. The pharmaceutical delivery system of claim 8 in which said microencapsulated particles have an encapsulating film comprised of ethylcellulose.

10 The pharmaceutical delivery system of claim 9 in which said microencapsulated pharmaceutical particles contain therein a rupturing agent.

11. The pharmaceutical delivery system of claim 10 in which said rupturing agent comprises sodium starch glycolate.

12. An oral pharmaceutical delivery system comprising:

- 15 A. at least one lipid;
B. at least one surfactant; and
C. dry particles;

wherein, the dry particles contain at least one pharmaceutical;

wherein, the dry particles are continuously coated with the lipid and form a homogenous suspension with the lipid;

20 wherein the suspension exhibits pseudoplastic and/or thixotropic properties;
and

wherein the suspension is formed or shaped into the appropriate solid dosage form by molding or pouring the suspension when in a liquid or semi-liquid state; and

wherein said lipid suspension contains a rupturing agent.

25 13. An oral pharmaceutical delivery system comprising:

- A. at least one lipid;
B. at least one surfactant; and
C. dry particles;

wherein, the dry particles contain at least one pharmaceutical;

30 wherein, the dry particles are continuously coated with the lipid and form a homogenous suspension with the lipid;

wherein the suspension exhibits pseudoplastic and/or thixotropic properties;
and

wherein the suspension is formed or shaped into the appropriate solid dosage form by molding or pouring the suspension when in a liquid or semi-liquid state; and wherein said filler comprises whey.

5 14. A method for preparing an oral pharmaceutical delivery system comprising:
melting at least one lipid;
mixing in at least one surfactant to said melted lipid;
mixing in dry particles to said melted lipid and surfactant,
wherein said dry particles contain at least one pharmaceutical, and at least one
filler;

10 wherein the dry particles are continuously coated with the lipid and form a
homogenous suspension with the lipid;

wherein the suspension exhibits pseudoplastic and/or thixotropic properties,
wherein the suspension is formed or shaped into the appropriate dose by
molding or pouring the suspension when in a liquid or semi-liquid state; and

15 wherein said pharmaceutical is selected from the group consisting of
analgesics, antibiotics, anti-inflammatory agents, cardiovascular preparations,
gastrointestinal medications, hormone products, and laxatives.

15. The method of claim 14 in which said lipid is selected from the group
consisting of a hard butter, petroleum wax, vegetable fat or vegetable stearines.

20 16. The method of claim 14 or 15 in which said surfactant is lecithin.

17. The method of claim 14 or 15 in which said surfactant has a hydrophilic to
lipophilic balance ratio in the range of about 1 to about 3.

25 18. The method of any one of claims 14 to 17 in which said lipid forms from
about 20% to 40% by weight of said suspension and said dry particles form from about
60% to 80% by weight of said suspension.

30 19. A method for preparing an oral pharmaceutical delivery system comprising:
melting at least one lipid;
mixing in at least one surfactant to said melted lipid;
mixing in dry particles to said melted lipid and surfactant, wherein said dry
particles contain at least one pharmaceutical, and at least one filler;

wherein the dry particles are continuously coated with the lipid and form a
homogenous suspension with the lipid;

35 wherein the suspension exhibits pseudoplastic and/or thixotropic properties;
wherein the suspension is formed or shaped into the appropriate dose by
molding or pouring the suspension when in a liquid or semi-liquid state; and

wherein said lipid suspension contains a rupturing agent.

20. The method of claim 19 in which said rupturing agent is sodium starch glycolate.

21. A method for preparing an oral pharmaceutical delivery system comprising:
5 melting at least one lipid;
mixing in at least one surfactant to said melted lipid;
mixing in dry particles to said melted lipid and surfactant, wherein said dry particles contain at least one pharmaceutical, and at least one filler;
wherein the dry particles are continuously coated with the lipid and form a
10 homogenous suspension with the lipid;
wherein the suspension exhibits pseudoplastic and/or thixotropic properties,
wherein the suspension is formed or shaped into the appropriate dose by molding or pouring the suspension when in a liquid or semi-liquid state; and
wherein said dry particles include artificial flavourings.

15 22. The method of claim 21 in which said lipid is selected from the group consisting of a hard butter, petroleum wax, vegetable fat or animal stearines.

23. The method of claim 21 or 22 in which said lipid suspension contains a rupturing agent.

24. The method of claim 23 in which said rupturing agent is sodium starch
20 glycolate.

25. The method of any one of claims 21 to 24 in which the dry particles include artificial flavourings.

26. A method for preparing an oral pharmaceutical delivery system comprising the steps of:

25 microencapsulating pharmaceutical particles;
melting at least one lipid;
mixing in a surfactant to said melted lipid;
dry-mixing dry particles comprising at least one filler and said microencapsulated pharmaceutical;
30 mixing the dry particles with said melted lipid to form a suspension such that said dry particles are continuously coated by said lipid such that said suspension exhibits pseudoplastic and/or thixotropic properties; and
pouring or molding said suspension into a dosage form.

27. The method of claim 26 in which said pharmaceutical particles are
35 microencapsulated with a filming agent, said filming agent comprising ethylcellulose.

28. The method of claim 26 in which said pharmaceutical particles are microencapsulated with a rupturing agent.

29. The method of claim 28 in which said rupturing agent is sodium starch glycolate.

5 30. The method of claim 29 in which said lipid source forms 20% to 40% by weight of said suspension, and said dry particles form 60% to 80% by weight of said suspension.

31. The method of claim 30 in which said fillers have a size ranging from 10 to 500 microns in diameter and comprise whey.

10 32. An oral pharmaceutical delivery system, substantially as hereinbefore described with reference to any one of the examples.

33. A method for preparing an oral pharmaceutical delivery system, substantially as hereinbefore described with reference to any one of the examples.

15 34. An oral pharmaceutical delivery system prepared according to the method of any one of claims 14 to 31 or 33.

Dated 25 October, 2004

Shear/Kershman Laboratories, Inc.

**Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON**

20