

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 2937418 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 12 N 15/11 (2006.01)** **A 61 K 31/713 (2006.01)** **C 12 N 15/113 (2010.01)**
- (45) Oversættelsen bekendtgjort den: **2018-01-22**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2017-10-18**
- (86) Europæisk ansøgning nr.: **15151452.8**
- (86) Europæisk indleveringsdag: **2009-10-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-10-28**
- (30) Prioritet: **2008-10-20 US 106956 P** **2008-11-18 US 115738 P**
2009-03-02 US 156670 P **2009-06-09 US 185545 P**
2009-09-15 US 242783 P **2009-09-22 US 244794 P**
- (62) Stamansøgningsnr: **09810834.3**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR**
- (73) Patenthaver: **Alnylam Pharmaceuticals, Inc., 300 Third Street, 3rd Floor, Cambridge, MA 02142, USA**
- (72) Opfinder: **Sah, Dinah Wen-yee, c/o Alnylam Pharmaceuticals Inc., 300 Third Street, Third Floor, Cambridge, MA 02142, USA**
HINKLE, Gregory, C/o Alnylam Pharmaceuticals Inc., 300 Third Street Third Floor, Cambridge, MA 02142, USA
ALVAREZ, Rene, C/o Alnylam Pharmaceuticals Inc., 300 Third Street, Third Floor, Cambridge, MA 02142, USA
MILSTEIN, Stuart, C/o Alnylam Pharmaceuticals Inc., 300 Third Street, Third Floor, Cambridge, MA 02142, USA
CHEN, Qingmin, C/o Alnylam Pharmaceuticals Inc., 300 Third Street, Third Floor, Cambridge, MA 02142, USA
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **Sammensætninger og fremgangsmåder til inhibering af ekspresion af transthyretin**
- (56) Fremdragne publikationer:
STEIN THOR D ET AL: "Neutralization of transthyretin reverses the neuroprotective effects of secreted amyloid precursor protein (APP) in APPSW mice resulting in tau phosphorylation and loss of hippocampal neurons: support for the amyloid hypothesis", JOURNAL OF NEUROSCIENCE, THE SOCIETY, WASHINGTON, DC, US, vol. 24, no. 35, 1 September 2004 (2004-09-01), pages 7707-7717, XP002577897, ISSN: 1529-2401, DOI: 10.1523/JNEUROSCI.2211-04.2004
KUROSAWA T ET AL: "Selective silencing of a mutant transthyretin allele by small interfering RNAs", BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 337, no. 3, 25 November 2005 (2005-11-25), pages 1012-1018, XP027218514, ISSN: 0006-291X, DOI: 10.1016/J.BBRC.2005.09.142 [retrieved on 2005-10-18]
ZIMMERMANN TRACY S ET AL: "RNAi-mediated gene silencing in non-human primates", NATURE, NATURE PUBLISHING GROUP, UNITED KINGDOM, vol. 441, no. 7089, 4 May 2006 (2006-05-04), pages 111-114, XP002577898, ISSN: 0028-0836, DOI: 10.1038/NATURE04688 [retrieved on 2006-03-26]
AKINC AKIN ET AL: "A combinatorial library of lipid-like materials for delivery of RNAi therapeutics", NATURE

Fortsættes ...

BIOTECHNOLOGY, NATURE PUBLISHING GROUP, US, vol. 26, no. 5, 1 May 2008 (2008-05-01), pages 561-569,
XP002577899, ISSN: 1087-0156, DOI: 10.1038/NBT1402 [retrieved on 2008-04-27]

DESCRIPTION

[0001] The disclosure relates to a double-stranded ribonucleic acid (dsRNA) targeting a transthyretin (TTR) gene, and methods of using the dsRNA to inhibit expression of TTR.

[0002] Transthyretin (TTR) is a secreted thyroid hormone-binding protein. TTR binds and transports retinol binding protein (RBP)/ Vitamin A, and serum thyroxine (T4) in plasma and cerebrospinal fluid.

[0003] Both normal-sequence TTR and variant-sequence TTR cause amyloidosis. Normal-sequence TTR causes cardiac amyloidosis in people who are elderly and is termed senile systemic amyloidosis (SSA) (also called senile cardiac amyloidosis (SCA)). SSA often is accompanied by microscopic deposits in many other organs. TTR mutations accelerate the process of TTR amyloid formation and are the most important risk factor for the development of clinically significant TTR amyloidosis (also called ATTR (amyloidosis-transthyretin type)). More than 85 amyloidogenic TTR variants are known to cause systemic familial amyloidosis. The liver is the major site of TTR expression. Other significant sites of expression include the choroid plexus, retina and pancreas.

[0004] TTR amyloidosis manifests in various forms. When the peripheral nervous system is affected more prominently, the disease is termed familial amyloidotic polyneuropathy (FAP). When the heart is primarily involved but the nervous system is not, the disease is called familial amyloidotic cardiomyopathy (FAC). A third major type of TTR amyloidosis is called leptomeningeal/CNS (Central Nervous System) amyloidosis.

[0005] Double-stranded RNA molecules (dsRNA) have been shown to block gene expression in a highly conserved regulatory mechanism known as RNA interference (RNAi). WO 99/32619 (Fire et al.) disclosed the use of a dsRNA of at least 25 nucleotides in length to inhibit the expression of genes in *C. elegans*. dsRNA has also been shown to degrade target RNA in other organisms, including plants (see, e.g., WO 99/53050, Waterhouse et al.; and WO 99/61631, Heifetz et al.), *Drosophila* (see, e.g., Yang, D., et al., Curr. Biol. (2000) 10:1191-1200), and mammals (see WO 00/44895, Limmer; and DE 101 00 586.5, Kreutzer et al.).

[0006] U.S. 20070207974 discloses functional and hyperfunctional siRNAs. U.S. 20090082300 discloses antisense molecules directed against TTR. U.S. Pat. No. 7,250,496 discloses microRNAs directed against TTR. Stein et al. (The Journal of Neuroscience, September 1, 2004, 24(35):7707-7717) describes that neutralization of transthyretin reverses the neuroprotective effects of secreted Amyloid Precursor Protein (APP) in APP_{Sw} Mice.

[0007] In one embodiment, the invention provides a double-stranded ribonucleic acid (dsRNA) for inhibiting expression of transthyretin (TTR), wherein (a) said dsRNA comprises a sense strand and an antisense strand, the antisense strand comprising a region complementary to a part of a mRNA encoding transthyretin (TTR), wherein said region of complementarity is 19 nucleotides in length, the antisense strand comprises SEQ ID NO:170, and each strand of the dsRNA is 19, 20, 21, 22, 23, or 24 nucleotides in length; or (b) said dsRNA comprises an antisense strand comprising a region complementary to 19 nucleotides of nucleotides 618-648 of SEQ ID NO: 1331 and wherein said antisense strand base pairs with the guanine at position 628 of SEQ ID NO:1331; and wherein the dsRNA is formulated in a lipid formulation comprising a cationic lipid, a non-cationic lipid, and a lipid that prevents aggregation of the particle.

[0008] Disclosed is a double-stranded ribonucleic acid (dsRNA) for inhibiting expression of transthyretin (TTR), wherein said dsRNA comprises a sense strand and an antisense strand, the antisense strand comprising a region complementary to a part of a mRNA encoding transthyretin (TTR), wherein said region of complementarity is less than 30 nucleotides in length and the antisense strand comprises 15 or more contiguous nucleotides of SEQ ID NO:170, SEQ ID NO:450, SEQ ID NO:730, or SEQ ID NO:1010. In a related implementation, the sense strand comprises 15 or more contiguous nucleotides of SEQ ID NO:169, SEQ ID NO:449, SEQ ID NO:729, or SEQ ID NO:1009. In yet another related embodiment, the sense strand consists of SEQ ID NO:449 and the antisense strand consists of SEQ ID NO:450. In yet another related embodiment, the sense strand consists of SEQ ID NO:729 and the antisense strand consists of SEQ ID NO:730. In still another related embodiment, the sense strand consists of SEQ ID NO:1009 and the antisense strand consists of SEQ ID NO:1010. In yet another related implementation, the dsRNA comprises a sense strand selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16, and an antisense strand selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16.

[0009] In certain implementations, the region of complementarity between the antisense strand of the dsRNA and the mRNA encoding transthyretin is 19 nucleotides in length. In another implementation, the region of complementarity consists of SEQ ID NO:169. In other implementations, each strand of the dsRNA is 19, 20, 21, 22, 23, or 24 nucleotides in length. In still another

implementation, each strand is 21 nucleotides in length.

[0010] In certain implementations, the dsRNA for inhibiting expression of transthyretin does not cleave a TTR mRNA between the adenine nucleotide at position 637 of SEQ ID NO:1331 and the guanine nucleotide at position 638 of SEQ ID NO:1331. In other implementations, the dsRNA cleaves a TTR mRNA between the guanine nucleotide at position 636 of SEQ ID NO:1331 and the adenine nucleotide at position 637 of SEQ ID NO:1331. In certain implementations, the dsRNA anneals to a TTR mRNA between the guanine nucleotide at position 628 of SEQ ID NO:1331 and the uracil nucleotide at position 646 of SEQ ID NO: 1331.

[0011] In still other related embodiments, the invention provides dsRNA as claimed for inhibiting expression of transthyretin wherein the dsRNA comprises one or more modified nucleotides. In related embodiments, at least one modified nucleotide (or nucleotides) is chosen from the group consisting of: a 2'-O-methyl modified nucleotide, a nucleotide comprising a 5'-phosphorothioate group, and a terminal nucleotide linked to a cholesteryl derivative or dodecanoic acid bisdecylamide group. In another related embodiment, the modified nucleotide is chosen from the group of: a 2'-deoxy-2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a locked nucleotide, an abasic nucleotide, 2'-amino-modified nucleotide, 2'-alkyl-modified nucleotide, morpholino nucleotide, a phosphoramidate, and a non-natural base comprising nucleotide. In certain embodiments, the dsRNA comprises at least one 2'-O-methyl modified nucleotide.

[0012] In other implementations, a dsRNA as described above for inhibiting expression of transthyretin is conjugated to a ligand, or formulated in a lipid formulation. In certain implementations, the lipid formulation may be a LNP formulation, a LNP01 formulation, a XTC-SNALP formulation, or a SNALP formulation. In related implementations, the XTC-SNALP formulation is as follows: using 2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC) with XTC/DPPC/Cholesterol/PEG-cDMA in a ratio of 57.1/7.1/34.4/1.4 and a lipid:siRNA ratio of about 7. In still other related implementations, the sense strand of the dsRNA consists of SEQ ID NO:1009 and the antisense strand consists of SEQ ID NO:1010, and the dsRNA is formulated in a XTC-SNALP formulation as follows: using 2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC) with a XTC/DPPC/Cholesterol/PEG-cDMA in a ratio of 57.1/7.1/34.4/1.4 and a lipid:siRNA ratio of about 7. Alternatively, a dsRNA such as those described above can be formulated in a LNP09 formulation as follows: using XTC/DSPC/Chol/PEG₂₀₀₀-C14 in a ratio of 50/10/38.5/1.5 mol% and a lipid:siRNA ratio of about 11:1. In another variation, the dsRNA is formulated in a LNP 11 formulation as follows: using MC3/DSPC/Chol/PEG₂₀₀₀-C14 in a ratio of 50/10/38.5/1.5 mol% and a lipid:siRNA ratio of about 11:1. In still another implementation, the dsRNA is formulated in a LNP09 formulation or a LNP11 formulation and reduces TTR mRNA levels by about 85 to 90% at a dose of 0.3mg/kg, relative to a PBS control group. In yet another implementation, the dsRNA is formulated in a LNP09 formulation or a LNP 11 formulation and reduces TTR mRNA levels by about 50% at a dose of 0.1 mg/kg, relative to a PBS control group. In yet another implementation, the dsRNA is formulated in a LNP09 formulation or a LNP 11 formulation and reduces TTR protein levels in a dose-dependent manner relative to a PBS control group as measured by a western blot. In yet another implementation, the dsRNA is formulated in a SNALP formulation as follows: using DlinDMA with a DlinDMA/DPPC/Cholesterol/PEG2000-cDMA in a ratio of 57.1/7.1/34.4/1.4 and a lipid:siRNA ratio of about 7.

[0013] In certain implementations, the disclosure provides a dsRNA such as those described above for inhibiting expression of transthyretin, wherein administration of the dsRNA to a cell results in about 95% inhibition of TTR mRNA expression as measured by a real time PCR assay, wherein the cell is a HepG2 cell or a Hep3B cell, and wherein the concentration of the dsRNA is 10nM. In related implementations, administration of the dsRNA to a cell results in about 74% inhibition of TTR mRNA expression as measured by a branched DNA assay, wherein the cell is a HepG2 cell or a Hep3B cell, and wherein the concentration of the dsRNA is 10nM. In other related implementations, the dsRNA has an IC₅₀ of less than 10 pM in a HepG2 cell, wherein the concentration of the dsRNA is 10nM. In still other related implementations, the dsRNA has an ED₅₀ of about 1 mg/kg. In still other related implementations, administration of the dsRNA reduces TTR mRNA by about 80% in cynomolgus monkey liver, wherein the concentration of the dsRNA is 3 mg/kg. In still other related implementations, administration of the dsRNA does not result in immunostimulatory activity in human peripheral blood mononuclear cells (PBMCs) as measured by IFN- α and TNF- α ELISA assays. In still other related implementations, administration of the dsRNA reduces liver TTR mRNA levels by about 97% or serum TTR protein levels by about 90%, wherein the concentration of the dsRNA is 6 mg/kg. In still other related implementations, administration of the dsRNA reduces liver TTR mRNA levels and/or serum TTR protein levels up to 22 days, wherein the concentration of the dsRNA is 6 mg/kg or 3 mg/kg. In still other related implementations, the dsRNA suppresses serum TTR protein levels up to day 14 post-treatment when administered to a subject in need thereof at 1 mg/kg or 3 mg/kg. In still other related implementations, the dsRNA reduces TTR expression by 98.9% in a Hep3B cell at a concentration of 0.1nM as measured by real-time PCR. In still other related implementations, the dsRNA reduces TTR expression by 99.4% in a Hep3B cell at a concentration of 10nM as measured by real-time PCR.

[0014] In other implementations, the disclosure provides a double-stranded ribonucleic acid (dsRNA) for inhibiting expression

of transthyretin (TTR), wherein said dsRNA comprises a sense strand and an antisense strand, the antisense strand comprising a region complementary to a part of a mRNA encoding transthyretin (TTR), wherein said region of complementarity is less than 30 nucleotides in length and wherein the dsRNA comprises a sense strand selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16, and an antisense strand selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16.

[0015] In another implementation, the disclosure provides a double-stranded ribonucleic acid (dsRNA) for inhibiting expression of transthyretin (TTR), wherein said dsRNA comprises an antisense strand comprising a region complementary to 15-30 nucleotides of nucleotides 618-648 of SEQ ID NO: 1331 and wherein said antisense strand base pairs with the guanine at position 628 of SEQ ID NO:1331.

[0016] Described is a cell containing any of the dsRNAs described in the Summary, above. Also described is a vector comprising a nucleotide sequence that encodes at least one strand of any of the dsRNAs described in the Summary, above. In certain implementations, the vector is in a cell.

[0017] In other embodiments, the invention provides a pharmaceutical composition for inhibiting expression of a TTR gene comprising any of the dsRNAs as claimed, above, and a pharmaceutically acceptable carrier.

[0018] Described is a pharmaceutical composition for inhibiting expression of a TTR gene comprising a dsRNA and a SNALP formulation, wherein the dsRNA comprises an antisense strand which is less than 30 nucleotides in length and comprises 15 or more contiguous nucleotides of SEQ ID NO:170, SEQ ID NO:450, SEQ ID NO:730, or SEQ ID NO:1010, and wherein the SNALP formulation comprises DLinDMA, DPPC, Cholesterol and PEG2000-cDMA in a ratio of 57.1/7.1/34.4/1.4 respectively.

[0019] Described is a method of inhibiting TTR expression in a cell, the method comprising: (a) contacting the cell with any of dsRNAs described in the Summary, above; and (b) maintaining the cell produced in step (a) for a time sufficient to obtain degradation of the mRNA transcript of a TTR gene, thereby inhibiting expression of the TTR gene in the cell.

[0020] Described is a method of treating a disorder mediated by TTR expression comprising administering to a human in need of such treatment a therapeutically effective amount of any of the dsRNAs describe in the Summary, above. In related embodiments, the dsRNA is administered to the human at about 0.01, 0.1, 0.5, 1.0, 2.5, or 5.0 mg/kg. In yet another related implementation, the dsRNA is administered to the human at about 1.0 mg/kg. In yet another related implementation, the human being treated has transthyretin amyloidosis, and/or a liver disorder. In a related implementation, the human is further provided a liver transplant. In yet another implementation, administration of the dsRNA reduces TTR mRNA by about 80% in human liver, wherein the concentration of the dsRNA is 3 mg/kg. In yet another related implementation, administration of the dsRNA does not result in immunostimulatory activity in the human as measured by IFN-alpha and TNF-alpha ELISA assays. In yet another related implementation, administration of the dsRNA reduces liver TTR mRNA levels by about 97% or serum TTR protein levels by about 90%, wherein the concentration of the dsRNA is 6 mg/kg. In yet another related implementation, administration of the dsRNA reduces liver TTR mRNA levels and/or serum TTR protein levels up to 22 days, wherein the concentration of the dsRNA is 6 mg/kg or 3 mg/kg. In yet another related implementation, the dsRNA is formulated in a LNP09 formulation as follows: using XTC/DSPC/Chol/PEG₂₀₀₀-C14 in a ratio of 50/10/38.5/1.5 mol% and a lipid:siRNA ratio of about 11:1. In yet another related implementation, the dsRNA is formulated in a LNP11 formulation as follows: using MC3/DSPC/Chol/PEG₂₀₀₀-C14 in a ratio of 50/10/38.5/1.5 mol% and a lipid:siRNA ratio of about 11:1. In yet another related implementation, the dsRNA is formulated in a LNP09 formulation or a LNP11 formulation and reduces TTR mRNA levels by about 85 to 90% at a dose of 0.3mg/kg, relative to a PBC control group. In yet another related implementation, the dsRNA is formulated in a LNP09 formulation or a LNP11 formulation and reduces TTR mRNA levels by about 50% at a dose of 0.1 mg/kg, relative to a PBC control group. In still another related implementation, the dsRNA is formulated in a LNP09 formulation or a LNP 11 formulation and reduces TTR protein levels in a dose-dependent manner relative to a PBC control group as measured by a western blot. In still another related implementation, administration of the dsRNA suppresses serum TTR protein levels up to day 14 post-treatment when administered to human at 1 mg/kg or 3 mg/kg. In still another related implementation, the dsRNA is formulated in a SNALP formulation as follows: using DLinDMA with a DLinDMA/DPPC/Cholesterol/PEG2000-cDMA in a ratio of 57.1/7.1/34.4/1.4 and a lipid:siRNA ratio of about 7.

[0021] Described is the use of a dsRNA for treating a disorder mediated by TTR expression comprising administering to a human in need of such treatment a therapeutically effective amount of any of the dsRNAs described in the Summary, above. In related implementations, the dsRNA is administered to the human at about 0.01, 0.1, 0.5, 1.0, 2.5, or 5.0 mg/kg. In a particular related implementation, the dsRNA is administered to the human at about 1.0 mg/kg. In another related implementation, the human has transthyretin amyloidosis, and/or a liver disorder. In yet another implementation of the use provided by the invention, the treated human is further provided a liver transplant.

[0022] Described is the use of a dsRNA in a method for inhibiting TTR expression in a cell, wherein the method comprises (a) contacting the cell with a dsRNA described in the Summary, above; and (b) maintaining the cell produced in step (a) for a time sufficient to obtain degradation of the mRNA transcript of a TTR gene, thereby inhibiting expression of the TTR gene in the cell.

FIG. 1 is a graph of TNFalpha and IFNalpha levels in cultured human PBMCs following transfection with TTR siRNAs.

FIG. 2A and 2B are dose response curves for AD-18324 and AD-18328, respectively, in HepG2 cells.

FIG. 3 is a dose response curve for AD-18246 in HepG2 cells.

FIG. 4A and FIG. 4B show inhibition of liver mRNA and plasma protein levels, respectively, in transgenic H129-mTTR-KO/iNOS-KO/hTTR mice by an intravenous bolus administration of TTR-dsRNA (AD-18324, AD-18328 and AD-18246) formulated in LNP01.

FIG. 5 is a graph summarizing the measurements of TTR mRNA levels in livers of non-human primates following 15-minute intravenous infusion of TTR-dsRNA (AD-18324 and AD-18328) formulated in SNALP.

FIG. 6A and FIG. 6B show inhibition of human V30M TTR liver mRNA and serum protein levels, respectively, in transgenic mice by an intravenous bolus administration of SNALP-18328. Group means were determined, normalized to the PBS control group, and then plotted. Error bars represent standard deviations. The percentage reduction of the group mean, relative to PBS, is indicated for the SNALP-1955 and SNALP-18328 groups. (** $p < 0.001$, One-way ANOVA, with Dunn's post-hoc test).

FIG. 7A and FIG. 7B show the durability of reduction of human V30M TTR liver mRNA and serum protein levels, respectively, in transgenic mice over 22 days following a single intravenous bolus administration of SNALP-18328. Group means were determined. TTR/GAPDH mRNA levels were normalized to day 0 levels and plotted. The percent reduction of normalized TTR mRNA levels relative to SNALP-1955 for each time point were calculated and are indicated for the SNALP-18328 groups. (** $p < 0.001$, One-way ANOVA, with Dunn's post-hoc test).

FIG. 8 shows the timecourse of TTR serum protein levels in non-human primates over 14 days following a single 15-minute intravenous infusion of SNALP-18328.

FIG. 9 shows reduction of TTR-immunoreactivity in various tissues of human V30M TTR/HSF-1 knock-out mice following intravenous bolus administration of SNALP-18328. E, esophagus; S, stomach; I1, intestine/duodenum; I4, intestine/colon; N, nerve; D, dorsal root ganglia.

FIG. 10 shows the measurements of TTR mRNA levels in livers of non-human primates following 15-minute intravenous infusion of XTC-SNALP-18328.

FIGs. 11A and 11B show the measurements of TTR mRNA and serum protein levels, respectively, in livers of non-human primates following 15-minute intravenous infusion of LNP09-18328 or LNP11-18328. FIG. 11C shows the timecourse of TTR serum protein levels over 28 days following a 15-minute intravenous infusion of 0.3mg/kg LNP09-18328, as compared to the PBS control group.

FIG. 12 shows the sequence of human TTR mRNA (Ref. Seq. NM_000371.3, SEQ ID NO:1331).

FIGs. 13A and 13B are the sequences of human and rat TTR mRNA, respectively. FIG. 13A is the sequence of human TTR mRNA (Ref. Seq. NM_000371.2, SEQ ID NO:1329). FIG. 13B is the sequence of rat TTR mRNA (Ref. Seq. NM_012681.1, SEQ ID NO:1330).

FIG. 14 shows the nucleotide alignment of NM_000371.3, NM_000371.2, and AD-18328.

FIG. 15 illustrates symptoms and mutations in TTR associated with familial amyloidotic neuropathy, familial amyloidotic cardiomyopathy and CNS amyloidosis.

FIG. 16 shows reduction of TTR mRNA levels in the liver with SNALP-18534 with different infusion durations. Groups of animals ($n=4$ /group) were administered 1 mg/kg SNALP-18534 via a 15-minute, or 1, 2, or 3 hour infusion. Forty-eight hours later, rats were euthanized and livers harvested. TTR and GAPDH mRNA levels were measured from liver lysates using the Quantigene bDNA assay. The ratio of TTR to GAPDH mRNA levels was calculated for each animal. Group means were determined and normalized to a PBS control group, and then plotted. Error bars represent standard deviations. (** $p < 0.001$, One-way ANOVA with Bonferroni post-hoc test, relative to PBS).

FIG. 17 shows the measurements of TTR mRNA levels in livers of rats following 15-minute intravenous infusion of LNP07-18534 or LNP08-18534.

FIG. 18 shows *in vivo* inhibition of endogenous TTR mRNA levels in livers of Sprague-Dawley Rats following a 15-min IV infusion of LNP09-18534 or LNP11-18534. Groups of animals (n=4/group) were intravenously administered 0.01, 0.03, 0.1, or 0.3 mg/kg LNP09-18534, LNP-11-18534; or PBS via a 15-minute infusion. Forty-eight hours later, animals were euthanized and livers harvested. TTR and GAPDH mRNA levels were measured from liver biopsy lysates using the Quantigene bDNA assay. The ratio of TTR to GAPDH mRNA levels was calculated for each animal. Group means were determined, normalized to the PBS control group, and then plotted. Error bars represent standard deviations.

[0023] The invention is defined by the claims.

[0024] Described are dsRNAs and methods of using the dsRNAs for inhibiting the expression of a TTR gene in a cell or a mammal where the dsRNA targets a TTR gene. The disclosure also provides compositions and methods for treating pathological conditions and diseases, such as a TTR amyloidosis, in a mammal caused by the expression of a TTR gene. dsRNA directs the sequence-specific degradation of mRNA through a process known as RNA interference (RNAi).

[0025] The dsRNAs of the compositions featured herein include an RNA strand (the antisense strand) having a region which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an mRNA transcript of a TTR gene. The use of these dsRNAs enables the targeted degradation of mRNAs of genes that are implicated in pathologies associated with TTR expression in mammals. Very low dosages of TTR dsRNAs in particular can specifically and efficiently mediate RNAi, resulting in significant inhibition of expression of a TTR gene. Using cell-based assays, the present inventors have demonstrated that dsRNAs targeting TTR can specifically and efficiently mediate RNAi, resulting in significant inhibition of expression of a TTR gene. Thus, methods and compositions including these dsRNAs are useful for treating pathological processes that can be mediated by down regulating TTR, such as in the treatment of a liver disorder or a TTR amyloidosis, e.g., FAP.

[0026] The methods and compositions containing a TTR dsRNA are useful for treating pathological processes mediated by TTR expression, such as a TTR amyloidosis. In an implementation, a method of treating a disorder mediated by TTR expression includes administering to a human in need of such treatment a therapeutically effective amount of a dsRNA targeted to TTR. In an embodiment, a dsRNA is administered to the human at about 0.01, 0.1, 0.5, 1.0, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 mg/kg.

[0027] The following detailed description discloses how to make and use the compositions containing dsRNAs to inhibit the expression of a TTR gene, as well as compositions and methods for treating diseases and disorders caused by the expression of this gene. The pharmaceutical compositions include a dsRNA having an antisense strand comprising a region of complementarity which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an RNA transcript of a TTR gene, together with a pharmaceutically acceptable carrier. The compositions also include a dsRNA having an antisense strand having a region of complementarity which is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and is substantially complementary to at least part of an RNA transcript of a TTR gene.

[0028] The sense strand of a dsRNA can include 15, 16, 17, 18, 19, 20, 21, or more contiguous nucleotides of SEQ ID NO:169, SEQ ID NO:449, SEQ ID NO:729, or SEQ ID NO:1009. The antisense strand of a dsRNA can include 15, 16, 17, 18, 19, 20, 21, or more contiguous nucleotides of SEQ ID NO:170, SEQ ID NO:450, SEQ ID NO:730, or SEQ ID NO:1010. In an embodiment, the sense strand of a dsRNA can consist of SEQ ID NO:449 and the antisense strand can consist of SEQ ID NO:450. In an embodiment, the sense strand of a dsRNA can consist of SEQ ID NO:729 and the antisense strand can consist of SEQ ID NO:730. In an embodiment, the sense strand of a dsRNA can consist of SEQ ID NO:1009 and the antisense strand can consist of SEQ ID NO:1010.

[0029] In an implementation, a dsRNA can include at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more modified nucleotides. In an embodiment, a modified nucleotide can include a 2'-O-methyl modified nucleotide, a nucleotide comprising a 5'-phosphorothioate group, and/or a terminal nucleotide linked to a cholesteryl derivative or dodecanoic acid bisdecylamide group. In an embodiment, a modified nucleotide can include a 2'-deoxy-2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a locked nucleotide, an abasic nucleotide, 2'-amino-modified nucleotide, 2'-alkyl-modified nucleotide, morpholino nucleotide, a phosphoramidate, and/or a non-natural base comprising nucleotide.

[0030] In an implementation, the region of complementarity of a dsRNA is at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, or more nucleotides in length. In an implementation, the region of complementarity includes 10, 11, 12, 13, 14, 15, 16, 17,

18, 19, 20, 21, or more contiguous nucleotides of SEQ ID NO:169.

[0031] In an implementation, each strand of a dsRNA is 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more nucleotides in length. In an implementation, the dsRNA includes a sense strand, or 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 21 nucleotide fragment thereof, selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16, and an antisense strand, or 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 21 nucleotide fragment thereof, selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16.

[0032] In an implementation, administration of a dsRNA to a cell results in about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 95% or more inhibition of TTR mRNA expression as measured by a real time PCR assay. In an implementation, administration of a dsRNA to a cell results in about 40% to 45%, 45% to 50%, 50% to 55%, 55% to 60%, 60% to 65%, 65% to 70%, 70% to 75%, 75% to 80%, 80% to 85%, 85% to 90%, 90% to 95% or more inhibition of TTR mRNA expression as measured by a real time PCR assay. In an implementation, administration of a dsRNA to a cell results in about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 95% or more inhibition of TTR mRNA expression as measured by a branched DNA assay. In an implementation, administration of a dsRNA to a cell results in about 40% to 45%, 45% to 50%, 50% to 55%, 55% to 60%, 60% to 65%, 65% to 70%, 70% to 75%, 75% to 80%, 80% to 85%, 85% to 90%, 90% to 95% or more inhibition of TTR mRNA expression as measured by a branched DNA assay.

[0033] In an implementation, a dsRNA has an IC₅₀ of less than 0.01pM, 0.1pM, 1pM, 5pM, 10 pM, 100pM, or 1000pM. In an implementation, a dsRNA has an ED₅₀ of about 0.01, 0.1, 1, 5, or 10 mg/kg.

[0034] In an implementation, administration of a dsRNA can reduce TTR mRNA by about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 95% or more in cynomolgus monkeys. In an implementation, administration of a dsRNA reduces liver TTR mRNA levels by about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 95% or more or serum TTR protein levels by about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 95% or more. In an implementation, administration of a dsRNA reduces liver TTR mRNA levels and/or serum TTR protein levels up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more days.

[0035] In an implementation, a dsRNA is formulated in a LNP formulation and reduces TTR mRNA levels by about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 95% or more at a dose of 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, or 1 mg/kg, relative to a PBC control group. In an implementation, a dsRNA is formulated in a LNP formulation and reduces TTR protein levels about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 95% or more relative to a PBC control group as measured by a western blot. In an implementation, a dsRNA suppresses serum TTR protein levels up to day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 post-treatment when administered to a subject in need thereof at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 mg/kg.

[0036] Accordingly, in some aspects, pharmaceutical compositions containing a TTR dsRNA and a pharmaceutically acceptable carrier, methods of using the compositions to inhibit expression of a TTR gene, and methods of using the pharmaceutical compositions to treat diseases caused by expression of a TTR gene are disclosed.

I. Definitions

[0037] For convenience, the meaning of certain terms and phrases used in the specification, examples, and appended claims, are provided below. If there is an apparent discrepancy between the usage of a term in other parts of this specification and its definition provided in this section, the definition in this section shall prevail.

[0038] "G," "C," "A" and "U" each generally stand for a nucleotide that contains guanine, cytosine, adenine, and uracil as a base, respectively. "T" and "dT" are used interchangeably herein and refer to a deoxyribonucleotide wherein the nucleobase is thymine, e.g., deoxyribothymine. However, it will be understood that the term "ribonucleotide" or "nucleotide" or "deoxyribonucleotide" can also refer to a modified nucleotide, as further detailed below, or a surrogate replacement moiety. The skilled person is well aware that guanine, cytosine, adenine, and uracil may be replaced by other moieties without substantially altering the base pairing properties of an oligonucleotide comprising a nucleotide bearing such replacement moiety. For example, without limitation, a nucleotide comprising inosine as its base may base pair with nucleotides containing adenine, cytosine, or uracil. Hence, nucleotides containing uracil, guanine, or adenine may be replaced in the nucleotide sequences of the invention by a nucleotide containing, for example, inosine. Sequences comprising such replacement moieties are embodiments of the invention.

[0039] As used herein, "transthyretin" ("TTR") refers to a gene in a cell. TTR is also known as ATTR, HsT2651, PALB,

prealbumin, TBPA, and transthyretin (prealbumin, amyloidosis type I). The sequence of a human TTR mRNA transcript can be found at NM_000371. The sequence of mouse TTR mRNA can be found at NM_013697.2, and the sequence of rat TTR mRNA can be found at NM_012681.1.

[0040] As used herein, "target sequence" refers to a contiguous portion of the nucleotide sequence of an mRNA molecule formed during the transcription of a TTR gene, including mRNA that is a product of RNA processing of a primary transcription product.

[0041] As used herein, the term "strand comprising a sequence" refers to an oligonucleotide comprising a chain of nucleotides that is described by the sequence referred to using the standard nucleotide nomenclature.

[0042] As used herein, and unless otherwise indicated, the term "complementary," when used to describe a first nucleotide sequence in relation to a second nucleotide sequence, refers to the ability of an oligonucleotide or polynucleotide comprising the first nucleotide sequence to hybridize and form a duplex structure under certain conditions with an oligonucleotide or polynucleotide comprising the second nucleotide sequence, as will be understood by the skilled person. Such conditions can, for example, be stringent conditions, where stringent conditions may include: 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50°C or 70°C for 12-16 hours followed by washing. Other conditions, such as physiologically relevant conditions as may be encountered inside an organism, can apply. The skilled person will be able to determine the set of conditions most appropriate for a test of complementarity of two sequences in accordance with the ultimate application of the hybridized nucleotides.

[0043] This includes base-pairing of the oligonucleotide or polynucleotide comprising the first nucleotide sequence to the oligonucleotide or polynucleotide comprising the second nucleotide sequence over the entire length of the first and second nucleotide sequence. Such sequences can be referred to as "fully complementary" with respect to each other herein. However, where a first sequence is referred to as "substantially complementary" with respect to a second sequence herein, the two sequences can be fully complementary, or they may form one or more, but generally not more than 4, 3 or 2 mismatched base pairs upon hybridization, while retaining the ability to hybridize under the conditions most relevant to their ultimate application. However, where two oligonucleotides are designed to form, upon hybridization, one or more single stranded overhangs, such overhangs shall not be regarded as mismatches with regard to the determination of complementarity. For example, a dsRNA comprising one oligonucleotide 21 nucleotides in length and another oligonucleotide 23 nucleotides in length, wherein the longer oligonucleotide comprises a sequence of 21 nucleotides that is fully complementary to the shorter oligonucleotide, may yet be referred to as "fully complementary" for the purposes described herein.

[0044] "Complementary" sequences, as used herein, may also include, or be formed entirely from, non-Watson-Crick base pairs and/or base pairs formed from non-natural and modified nucleotides, in as far as the above requirements with respect to their ability to hybridize are fulfilled. Such non-Watson-Crick base pairs includes, but not limited to, G:U Wobble or Hoogsteen base pairing.

[0045] The terms "complementary," "fully complementary" and "substantially complementary" herein may be used with respect to the base matching between the sense strand and the antisense strand of a dsRNA, or between the antisense strand of a dsRNA and a target sequence, as will be understood from the context of their use.

[0046] As used herein, a polynucleotide that is "substantially complementary to at least part of" a messenger RNA (mRNA) refers to a polynucleotide that is substantially complementary to a contiguous portion of the mRNA of interest (*e.g.*, an mRNA encoding TTR) including a 5' UTR, an open reading frame (ORF), or a 3' UTR. For example, a polynucleotide is complementary to at least a part of a TTR mRNA if the sequence is substantially complementary to a non-interrupted portion of an mRNA encoding TTR.

[0047] The term "double-stranded RNA" or "dsRNA," as used herein, refers to a complex of ribonucleic acid molecules, having a duplex structure comprising two anti-parallel and substantially complementary, as defined above, nucleic acid strands. In general, the majority of nucleotides of each strand are ribonucleotides, but as described in detail herein, each or both strands can also include at least one non-ribonucleotide, *e.g.*, a deoxyribonucleotide and/or a modified nucleotide. In addition, as used in this specification, "dsRNA" may include chemical modifications to ribonucleotides, including substantial modifications at multiple nucleotides and including all types of modifications disclosed herein or known in the art. Any such modifications, as used in an siRNA type molecule, are encompassed by "dsRNA" for the purposes of this specification and claims.

[0048] The two strands forming the duplex structure may be different portions of one larger RNA molecule, or they may be separate RNA molecules. Where the two strands are part of one larger molecule, and therefore are connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming the duplex structure, the connecting RNA chain is referred to as a "hairpin loop." Where the two strands are connected covalently by means other than an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming the duplex structure, the connecting structure is referred to as a "linker." The RNA strands may have the same or a different number of nucleotides. The maximum number of base pairs is the number of nucleotides in the shortest strand of the dsRNA minus any overhangs that are present in the duplex. In addition to the duplex structure, a dsRNA may comprise one or more nucleotide overhangs. The term "siRNA" is also used herein to refer to a dsRNA as described above.

[0049] As used herein, a "nucleotide overhang" refers to the unpaired nucleotide or nucleotides that protrude from the duplex structure of a dsRNA when a 3'-end of one strand of the dsRNA extends beyond the 5'-end of the other strand, or vice versa. "Blunt" or "blunt end" means that there are no unpaired nucleotides at that end of the dsRNA, *i.e.*, no nucleotide overhang. A "blunt ended" dsRNA is a dsRNA that is double-stranded over its entire length, *i.e.*, no nucleotide overhang at either end of the molecule.

[0050] The term "antisense strand" refers to the strand of a dsRNA which includes a region that is substantially complementary to a target sequence. As used herein, the term "region of complementarity" refers to the region on the antisense strand that is substantially complementary to a sequence, for example a target sequence, as defined herein. Where the region of complementarity is not fully complementary to the target sequence, the mismatches are most tolerated in the terminal regions and, if present, are generally in a terminal region or regions, *e.g.*, within 6, 5, 4, 3, or 2 nucleotides of the 5' and/or 3' terminus.

[0051] The term "sense strand," as used herein, refers to the strand of a dsRNA that includes a region that is substantially complementary to a region of the antisense strand.

[0052] As used herein, the term "SNALP" refers to a stable nucleic acid-lipid particle. A SNALP represents a vesicle of lipids coating a reduced aqueous interior comprising a nucleic acid such as a dsRNA or a plasmid from which a dsRNA is transcribed. SNALP are described, *e.g.*, in U.S. Patent Application Publication Nos. 20060240093, 20070135372, and USSN 61/045,228 filed on April 15, 2008.

[0053] "Introducing into a cell," when referring to a dsRNA, means facilitating uptake or absorption into the cell, as is understood by those skilled in the art. Absorption or uptake of dsRNA can occur through unaided diffusive or active cellular processes, or by auxiliary agents or devices. The meaning of this term is not limited to cells *in vitro*; a dsRNA may also be "introduced into a cell," wherein the cell is part of a living organism. In such instance, introduction into the cell will include the delivery to the organism. For example, for *in vivo* delivery, dsRNA can be injected into a tissue site or administered systemically. *In vitro* introduction into a cell includes methods known in the art such as electroporation and lipofection. Further approaches are described herein or known in the art.

[0054] The terms "silence," "inhibit the expression of," "down-regulate the expression of," "suppress the expression of" and the like in as far as they refer to a TTR gene, herein refer to the at least partial suppression of the expression of a TTR gene, as manifested by a reduction of the amount of mRNA which may be isolated from a first cell or group of cells in which a TTR gene is transcribed and which has or have been treated such that the expression of a TTR gene is inhibited, as compared to a second cell or group of cells substantially identical to the first cell or group of cells but which has or have not been so treated (control cells). The degree of inhibition is usually expressed in terms of

$$\frac{(\text{mRNA in control cells}) - (\text{mRNA in treated cells})}{(\text{mRNA in control cells})} \bullet 100\%$$

[0055] Alternatively, the degree of inhibition may be given in terms of a reduction of a parameter that is functionally linked to TTR gene expression, *e.g.*, the amount of protein encoded by a TTR gene which is secreted by a cell, or the number of cells displaying a certain phenotype, *e.g.*, apoptosis. In principle, TTR gene silencing may be determined in any cell expressing the target, either constitutively or by genomic engineering, and by any appropriate assay. However, when a reference is needed in order to determine whether a given dsRNA inhibits the expression of a TTR gene by a certain degree and therefore is encompassed by the instant invention, the assays provided in the Examples below shall serve as such reference.

[0056] For example, in certain instances, expression of a TTR gene is suppressed by at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, or 50% by administration of the double-stranded oligonucleotide featured in the invention. In

some embodiments, a TTR gene is suppressed by at least about 60%, 70%, or 80% by administration of the double-stranded oligonucleotide featured in the invention. In some embodiments, a TTR gene is suppressed by at least about 85%, 90%, or 95% by administration of the double-stranded oligonucleotide featured in the invention.

[0057] As used herein in the context of TTR expression, the terms "treat," "treatment," and the like, refer to relief from or alleviation of pathological processes mediated by TTR expression. In the context of the present invention insofar as it relates to any of the other conditions recited herein below (other than pathological processes mediated by TTR expression), the terms "treat," "treatment," and the like mean to relieve or alleviate at least one symptom associated with such condition, or to slow or reverse the progression of such condition, such as the slowing the progression of a TTR amyloidosis, such as FAP. Symptoms of TTR amyloidosis include sensory neuropathy (e.g. paresthesia, hypesthesia in distal limbs), autonomic neuropathy (e.g., gastrointestinal dysfunction, such as gastric ulcer, or orthostatic hypotension), motor neuropathy, seizures, dementia, myelopathy, polyneuropathy, carpal tunnel syndrome, autonomic insufficiency, cardiomyopathy, vitreous opacities, renal insufficiency, nephropathy, substantially reduced mBMI (modified Body Mass Index), cranial nerve dysfunction, and corneal lattice dystrophy.

[0058] As used herein, the phrases "therapeutically effective amount" and "prophylactically effective amount" refer to an amount that provides a therapeutic benefit in the treatment, prevention, or management of pathological processes mediated by TTR expression or an overt symptom of pathological processes mediated by TTR expression. The specific amount that is therapeutically effective can be readily determined by an ordinary medical practitioner, and may vary depending on factors known in the art, such as, for example, the type of pathological processes mediated by TTR expression, the patient's history and age, the stage of pathological processes mediated by TTR expression, and the administration of other anti-pathological processes mediated by TTR expression agents.

[0059] As used herein, a "pharmaceutical composition" comprises a pharmacologically effective amount of a dsRNA and a pharmaceutically acceptable carrier. As used herein, "pharmacologically effective amount," "therapeutically effective amount" or simply "effective amount" refers to that amount of an RNA effective to produce the intended pharmacological, therapeutic or preventive result. For example, if a given clinical treatment is considered effective when there is at least a 25% reduction in a measurable parameter associated with a disease or disorder, a therapeutically effective amount of a drug for the treatment of that disease or disorder is the amount necessary to effect at least a 25% reduction in that parameter. For example, a therapeutically effective amount of a dsRNA targeting TTR can reduce TTR serum levels by at least 25%. In another example, a therapeutically effective amount of a dsRNA targeting TTR can improve liver function or renal function by at least 25%.

[0060] The term "pharmaceutically acceptable carrier" refers to a carrier for administration of a therapeutic agent. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The term specifically excludes cell culture medium. For drugs administered orally, pharmaceutically acceptable carriers include, but are not limited to pharmaceutically acceptable excipients such as inert diluents, disintegrating agents, binding agents, lubricating agents, sweetening agents, flavoring agents, coloring agents and preservatives. Suitable inert diluents include sodium and calcium carbonate, sodium and calcium phosphate, and lactose, while corn starch and alginic acid are suitable disintegrating agents. Binding agents may include starch and gelatin, while the lubricating agent, if present, will generally be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or glyceryl distearate, to delay absorption in the gastrointestinal tract.

[0061] As used herein, a "transformed cell" is a cell into which a vector has been introduced from which a dsRNA molecule may be expressed.

II. Double-stranded ribonucleic acid (dsRNA)

[0062] As described in more detail herein, the disclosure provides double-stranded ribonucleic acid (dsRNA) molecules for inhibiting the expression of a TTR gene in a cell or mammal, e.g., in a human having a TTR amyloidosis, where the dsRNA includes an antisense strand having a region of complementarity which is complementary to at least a part of an mRNA formed in the expression of a TTR gene, and where the region of complementarity is less than 30 nucleotides in length, generally 19-24 nucleotides in length, and where said dsRNA, upon contact with a cell expressing said TTR gene, inhibits the expression of said TTR gene by at least 30% as assayed by, for example, a PCR or branched DNA (bDNA)-based method, or by a protein-based method, such as by Western blot. Expression of a TTR gene can be reduced by at least 30% when measured by an assay as described in the Examples below. For example, expression of a TTR gene in cell culture, such as in Hep3B cells, can be assayed by measuring TTR mRNA levels, such as by bDNA or TaqMan assay, or by measuring protein levels, such as by ELISA assay. The dsRNA of the invention can further include one or more single-stranded nucleotide

overhangs.

[0063] The dsRNA can be synthesized by standard methods known in the art as further discussed below, *e.g.*, by use of an automated DNA synthesizer, such as are commercially available from, for example, Biosearch, Applied Biosystems, Inc. The dsRNA includes two RNA strands that are sufficiently complementary to hybridize to form a duplex structure. One strand of the dsRNA (the antisense strand) includes a region of complementarity that is substantially complementary, and generally fully complementary, to a target sequence, derived from the sequence of an mRNA formed during the expression of a TTR gene, the other strand (the sense strand) includes a region that is complementary to the antisense strand, such that the two strands hybridize and form a duplex structure when combined under suitable conditions. Generally, the duplex structure is between 15 and 30 or between 25 and 30, or between 18 and 25, or between 19 and 24, or between 19 and 21, or 19, 20, or 21 base pairs in length. In one embodiment the duplex is 19 base pairs in length. In another embodiment the duplex is 21 base pairs in length. When two different siRNAs are used in combination, the duplex lengths can be identical or can differ.

[0064] Each strand of the dsRNA is generally between 15 and 30, or between 18 and 25, or 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides in length. In other embodiments, each strand is 25-30 nucleotides in length. Each strand of the duplex can be the same length or of different lengths. When two different siRNAs are used in combination, the lengths of each strand of each siRNA can be identical or can differ.

[0065] The dsRNA can include one or more single-stranded overhang(s) of one or more nucleotides. In one implementation, at least one end of the dsRNA has a single-stranded nucleotide overhang of 1 to 4, generally 1 or 2 nucleotides. In another implementation, the antisense strand of the dsRNA has 1-10 nucleotides overhangs each at the 3' end and the 5' end over the sense strand. In further implementations, the sense strand of the dsRNA has 1-10 nucleotides overhangs each at the 3' end and the 5' end over the antisense strand.

[0066] A dsRNAs having at least one nucleotide overhang can have unexpectedly superior inhibitory properties than the blunt-ended counterpart. In some implementations the presence of only one nucleotide overhang strengthens the interference activity of the dsRNA, without affecting its overall stability. A dsRNA having only one overhang has proven particularly stable and effective *in vivo*, as well as in a variety of cells, cell culture mediums, blood, and serum. Generally, the single-stranded overhang is located at the 3'-terminal end of the antisense strand or, alternatively, at the 3'-terminal end of the sense strand. The dsRNA can also have a blunt end, generally located at the 5'-end of the antisense strand. Such dsRNAs can have improved stability and inhibitory activity, thus allowing administration at low dosages, *i.e.*, less than 5 mg/kg body weight of the recipient per day. Generally, the antisense strand of the dsRNA has a nucleotide overhang at the 3'-end, and the 5'-end is blunt. In another implementation, one or more of the nucleotides in the overhang is replaced with a nucleoside thiophosphate.

[0067] In one implementation, a TTR gene is a human TTR gene. In specific implementations, the sense strand of the dsRNA is one of the sense sequences from Tables 3A, 3B, 4, 6A, 6B, or 7, and the antisense strand is one of the antisense sequences of Tables 3A, 3B, 4, 6A, 6B, or 7. Alternative antisense agents that target elsewhere in the target sequence provided in Tables 3A, 3B, 4, 6A, 6B, or 7 can readily be determined using the target sequence and the flanking TTR sequence.

[0068] The skilled person is well aware that dsRNAs having a duplex structure of between 20 and 23, but specifically 21, base pairs have been hailed as particularly effective in inducing RNA interference (Elbashir et al., EMBO 2001, 20:6877-6888). However, others have found that shorter or longer dsRNAs can be effective as well. In the embodiments described above, by virtue of the nature of the oligonucleotide sequences provided in Tables 3A, 3B, 4, 6A, 6B, and 7, the dsRNAs featured in the invention can include at least one strand of a length described herein. It can be reasonably expected that shorter dsRNAs having one of the sequences of Tables 3A, 3B, 4, 6A, 6B, or 7 minus only a few nucleotides on one or both ends may be similarly effective as compared to the dsRNAs described above. Hence, dsRNAs having a partial sequence of at least 15, 16, 17, 18, 19, 20, or more contiguous nucleotides from one of the sequences of Tables 3, 4, 6 or 7, and differing in their ability to inhibit the expression of a TTR gene in an assay as described herein below by not more than 5, 10, 15, 20, 25, or 30 % inhibition from a dsRNA comprising the full sequence, are contemplated. Further, dsRNAs that cleave within a desired TTR target sequence can readily be made using the corresponding TTR antisense sequence and a complementary sense sequence.

[0069] In addition, the dsRNAs provided in Tables 3A, 3B, 4, 6A, 6B, or 7 identify a site in a TTR that is susceptible to RNAi based cleavage. As such, the present disclosure further features dsRNAs that target within the sequence targeted by one of the agents of the present disclosure. As used herein, a second dsRNA is said to target within the sequence of a first dsRNA if the second dsRNA cleaves the message anywhere within the mRNA that is complementary to the antisense strand of the first

dsRNA. Such a second dsRNA will generally consist of at least 15 contiguous nucleotides from one of the sequences provided in Tables 3A, 3B, 4, 6A, 6B, or 7 coupled to additional nucleotide sequences taken from the region contiguous to the selected sequence in a TTR gene.

[0070] The dsRNA featured herein can contain one or more mismatches to the target sequence. In one implementation, the dsRNA featured in the disclosure contains no more than 3 mismatches. If the antisense strand of the dsRNA contains mismatches to a target sequence, it is preferable that the area of mismatch not be located in the center of the region of complementarity. If the antisense strand of the dsRNA contains mismatches to the target sequence, it is preferable that the mismatch be restricted to 5 nucleotides from either end, for example 5, 4, 3, 2, or 1 nucleotide from either the 5' or 3' end of the region of complementarity. For example, for a 23 nucleotide dsRNA strand which is complementary to a region of a TTR gene, the dsRNA generally does not contain any mismatch within the central 13 nucleotides. The methods described within the disclosure can be used to determine whether a dsRNA containing a mismatch to a target sequence is effective in inhibiting the expression of a TTR gene. Consideration of the efficacy of dsRNAs with mismatches in inhibiting expression of a TTR gene is important, especially if the particular region of complementarity in a TTR gene is known to have polymorphic sequence variation within the population.

Modifications

[0071] In yet another embodiment, the dsRNA is chemically modified to enhance stability. The nucleic acids featured in the invention may be synthesized and/or modified by methods well established in the art, such as those described in "Current protocols in nucleic acid chemistry," Beaucage, S.L. et al. (Eds.), John Wiley & Sons, Inc., New York, NY, USA, which is hereby incorporated herein by reference. Specific examples of dsRNA compounds useful in this invention include dsRNAs containing modified backbones or no natural internucleoside linkages. As defined in this specification, dsRNAs having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified dsRNAs that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

[0072] Modified dsRNA backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5'linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

[0073] Representative U.S. patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,195; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,316; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050.

[0074] Modified dsRNA backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatoms and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

[0075] Representative U.S. patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,64,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and, 5,677,439, each of which is herein incorporated by reference.

[0076] In other suitable dsRNA mimetics, both the sugar and the internucleoside linkage, *i.e.*, the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, a dsRNA mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar backbone of a dsRNA is replaced with an amide

containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative U.S. patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

[0077] Other embodiments of the invention are dsRNAs with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular --CH₂--NH--CH₂--, --CH₂--N(CH₃)--O--CH₂--[known as a methylene (methylimino) or MMI backbone], --CH₂--O--N(CH₃)--CH₂--, --CH₂--N(CH₃)--N(CH₃)--CH₂-- and --N(CH₃)--CH₂--CH₂--[wherein the native phosphodiester backbone is represented as --O--P--O--CH₂--] of the above-referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above-referenced U.S. Pat. No. 5,602,240. Also preferred are dsRNAs having morpholino backbone structures of the above-referenced U.S. Pat. No. 5,034,506.

[0078] Modified dsRNAs may also contain one or more substituted sugar moieties. Preferred dsRNAs comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other preferred dsRNAs comprise one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an dsRNA, or a group for improving the pharmacodynamic properties of an dsRNA, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O--CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) *i.e.*, an alkoxy-alkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, *i.e.*, a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples herein below, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), *i.e.*, 2'-O--CH₂--O--CH₂--N(CH₂)₂, also described in examples herein below.

[0079] Other preferred modifications include 2'-methoxy (2'-OCH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the dsRNA, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked dsRNAs and the 5' position of 5' terminal nucleotide. DsRNAs may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative U.S. patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920.

[0080] dsRNAs may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo, particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-daazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J. L., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie, International Edition*, 1991, 30, 613, and those disclosed by Sanghvi, Y. S., Chapter 15, *DsRNA Research and Applications*, pages 289-302, Crooke, S. T. and Lebleu, B., Eds., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds featured in the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C. (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., Eds., *DsRNA Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are exemplary base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

[0081] Representative U.S. patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Pat. No. 3,687,808, as well as U.S. Pat. Nos.

4,845,205; 5,130,30; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; and 5,681,941, and U.S. Pat. No. 5,750,692.

Conjugates

[0082] Another modification of the dsRNAs involves chemically linking to the dsRNA one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the dsRNA. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., Proc. Natl. Acad. Sci. USA, 1989, 86: 6553-6556), cholic acid (Manoharan et al., Biorg. Med. Chem. Lett., 1994, 4:1053-1060), a thioether, e.g., beryl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660:306-309; Manoharan et al., Biorg. Med. Chem. Lett., 1993, 3:2765-2770), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20:533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., EMBO J, 1991, 10:1111-1118; Kabanov et al., FEBS Lett., 1990, 259:327-330; Svinarchuk et al., Biochimie, 1993, 75:49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-Hphosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36:3651-3654; Shea et al., Nucl. Acids Res., 1990, 18:3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14:969-973), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36:3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264:229-237), or an octadecylamine or hexylamino-carbonyloxycholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277:923-937).

[0083] Representative U.S. patents that teach the preparation of such dsRNA conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941.

[0084] It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within a dsRNA. The present disclosure also includes dsRNA compounds which are chimeric compounds. "Chimeric" dsRNA compounds or "chimeras," in the context of this disclosure, are dsRNA compounds, particularly dsRNAs, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of a dsRNA compound. These dsRNAs typically contain at least one region wherein the dsRNA is modified so as to confer upon the dsRNA increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the dsRNA may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of dsRNA inhibition of gene expression. Consequently, comparable results can often be obtained with shorter dsRNAs when chimeric dsRNAs are used, compared to phosphorothioate deoxydsRNAs hybridizing to the same target region.

[0085] Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art. The cleavage site on the target mRNA of a dsRNA can be determined using methods generally known to one of ordinary skill in the art, e.g., the 5'-RACE method described in Soutschek et al., Nature; 2004, Vol. 432, pp. 173-178 (which is herein incorporated by reference for all purposes). In an embodiment, using the 5'-RACE method described by Soutschek et al., ALN-18328 was determined to cleave a TTR mRNA between the guanine nucleotide at position 636 of SEQ ID NO:1331 (N_000371.3) and the adenine nucleotide at position 637 of SEQ ID NO:1331. In an embodiment, it was determined that ALN-18328 does not cleave a TTR mRNA between the adenine nucleotide at position 637 of SEQ ID NO:1331 and the guanine nucleotide at position 638 of SEQ ID NO:1331.

[0086] In certain instances, the dsRNA may be modified by a non-ligand group. A number of non-ligand molecules have been conjugated to dsRNAs in order to enhance the activity, cellular distribution or cellular uptake of the dsRNA, and procedures for performing such conjugations are available in the scientific literature. Such non-ligand moieties have included lipid moieties, such as cholesterol (Letsinger et al., Proc. Natl. Acad. Sci. USA, 1989, 86:6553), cholic acid (Manoharan et al., Bioorg. Med. Chem. Lett., 1994, 4:1053), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660:306; Manoharan et al., Bioorg. Med. Chem. Lett., 1993, 3:2765), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20:533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., EMBO J., 1991, 10:111; Kabanov et al., FEBS Lett., 1990, 259:327; Svinarchuk et al., Biochimie, 1993, 75:49), a phospholipid, e.g., di-hexadecyl-rac-

glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36:3651; Shea et al., Nucl. Acids Res., 1990, 18:3777), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14:969), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36:3651), a palmitoyl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264:229), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277:923). Representative United States patents that teach the preparation of such dsRNA conjugates have been listed above. Typical conjugation protocols involve the synthesis of dsRNAs bearing an aminolinker at one or more positions of the sequence. The amino group is then reacted with the molecule being conjugated using appropriate coupling or activating reagents. The conjugation reaction may be performed either with the dsRNA still bound to the solid support or following cleavage of the dsRNA in solution phase. Purification of the dsRNA conjugate by HPLC typically affords the pure conjugate.

Vector encoded dsRNAs

[0087] In another aspect, TTR dsRNA molecules are expressed from transcription units inserted into DNA or RNA vectors (see, e.g., Couture, A, et al., TIG. (1996), 12:5-10; Skillern, A., et al., International PCT Publication No. WO 00/22113, Conrad, International PCT Publication No. WO 00/22114, and Conrad, U.S. Pat. No. 6,054,299). These transgenes can be introduced as a linear construct, a circular plasmid, or a viral vector, which can be incorporated and inherited as a transgene integrated into the host genome. The transgene can also be constructed to permit it to be inherited as an extrachromosomal plasmid (Gassmann, et al., Proc. Natl. Acad. Sci. USA (1995) 92:1292).

[0088] The individual strands of a dsRNA can be transcribed by promoters on two separate expression vectors and co-transfected into a target cell. Alternatively each individual strand of the dsRNA can be transcribed by promoters both of which are located on the same expression plasmid. In one embodiment, a dsRNA is expressed as an inverted repeat joined by a linker polynucleotide sequence such that the dsRNA has a stem and loop structure.

[0089] The recombinant dsRNA expression vectors are generally DNA plasmids or viral vectors. dsRNA expressing viral vectors can be constructed based on, but not limited to, adeno-associated virus (for a review, see Muzyczka, et al., Curr. Topics Micro. Immunol. (1992) 158:97-129); adenovirus (see, for example, Berkner, et al., BioTechniques (1998) 6:616), Rosenfeld et al. (1991, Science 252:431-434), and Rosenfeld et al. (1992), Cell 68:143-155); or alphavirus as well as others known in the art. Retroviruses have been used to introduce a variety of genes into many different cell types, including epithelial cells, *in vitro* and/or *in vivo* (see, e.g., Eglitis, et al., Science (1985) 230:1395-1398; Danos and Mulligan, Proc. Natl. Acad. Sci. USA (1998) 85:6460-6464; Wilson et al., 1988, Proc. Natl. Acad. Sci. USA 85:3014-3018; Armentano et al., 1990, Proc. Natl. Acad. Sci. USA 87:61416145; Huber et al., 1991, Proc. Natl. Acad. Sci. USA 88:8039-8043; Ferry et al., 1991, Proc. Natl. Acad. Sci. USA 88:8377-8381; Chowdhury et al., 1991, Science 254:1802-1805; van Beusechem. et al., 1992, Proc. Natl. Acad. Sci. USA 89:7640-19; Kay et al., 1992, Human Gene Therapy 3:641-647; Dai et al., 1992, Proc. Natl. Acad. Sci. USA 89:10892-10895; Hwu et al., 1993, J. Immunol. 150:4104-4115; U.S. Patent No. 4,868,116; U.S. Patent No. 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT Application WO 89/05345; and PCT Application WO 92/07573). Recombinant retroviral vectors capable of transducing and expressing genes inserted into the genome of a cell can be produced by transfecting the recombinant retroviral genome into suitable packaging cell lines such as PA317 and Psi-CRIP (Comette et al., 1991, Human Gene Therapy 2:5-10; Cone et al., 1984, Proc. Natl. Acad. Sci. USA 81:6349). Recombinant adenoviral vectors can be used to infect a wide variety of cells and tissues in susceptible hosts (e.g., rat, hamster, dog, and chimpanzee) (Hsu et al., 1992, J. Infectious Disease, 166:769), and also have the advantage of not requiring mitotically active cells for infection.

[0090] Any viral vector capable of accepting the coding sequences for the dsRNA molecule(s) to be expressed can be used, for example vectors derived from adenovirus (AV); adeno-associated virus (AAV); retroviruses (e.g. lentiviruses (LV), Rhabdoviruses, murine leukemia virus); herpes virus, and the like. The tropism of viral vectors can be modified by pseudotyping the vectors with envelope proteins or other surface antigens from other viruses, or by substituting different viral capsid proteins, as appropriate.

[0091] For example, lentiviral vectors featured in the invention can be pseudotyped with surface proteins from vesicular stomatitis virus (VSV), rabies, Ebola, Mokola, and the like. AAV vectors featured in the invention can be made to target different cells by engineering the vectors to express different capsid protein serotypes. For example, an AAV vector expressing a serotype 2 capsid on a serotype 2 genome is called AAV 2/2. This serotype 2 capsid gene in the AAV 2/2 vector can be replaced by a serotype 5 capsid gene to produce an AAV 2/5 vector. Techniques for constructing AAV vectors which express different capsid protein serotypes are within the skill in the art; see, e.g., Rabinowitz J E et al. (2002), J Virol 76:791-801.

[0092] Selection of recombinant viral vectors, methods for inserting nucleic acid sequences for expressing the dsRNA into the vector, and methods of delivering the viral vector to the cells of interest are within the skill in the art. See, for example, Dornburg R (1995), *Gene Therap.* 2: 301-310; Eglitis M A (1988), *Biotechniques* 6: 608-614; Miller A D (1990), *Hum Gene Therap.* 1: 5-14; Anderson W F (1998), *Nature* 392: 25-30; and Robinson D A et al., *Nat. Genet.* 33: 401-406.

[0093] Viral vectors can be derived from AV and AAV. In one implementation, the dsRNA featured in the invention is expressed as two separate, complementary single-stranded RNA molecules from a recombinant AAV vector having, for example, either the U6 or H1 RNA promoters, or the cytomegalovirus (CMV) promoter.

[0094] A suitable AV vector for expressing the dsRNA featured in the invention, a method for constructing the recombinant AV vector, and a method for delivering the vector into target cells, are described in Xia H et al. (2002), *Nat. Biotech.* 20: 1006-1010.

[0095] Suitable AAV vectors for expressing the dsRNA featured in the invention, methods for constructing the recombinant AV vector, and methods for delivering the vectors into target cells are described in Samulski R et al. (1987), *J. Virol.* 61: 3096-3101; Fisher K J et al. (1996), *J. Virol.* 70: 520-532; Samulski R et al. (1989), *J. Virol.* 63: 3822-3826; U.S. Pat. No. 5,252,479; U.S. Pat. No. 5,139,941; International Patent Application No. WO 94/13788; and International Patent Application No. WO 93/24641.

[0096] The promoter driving dsRNA expression in either a DNA plasmid or viral vector featured in the invention may be a eukaryotic RNA polymerase I (e.g., ribosomal RNA promoter), RNA polymerase II (e.g., CMV early promoter or actin promoter or U1 snRNA promoter) or generally RNA polymerase III promoter (e.g., U6 snRNA or 7SK RNA promoter) or a prokaryotic promoter, for example the T7 promoter, provided the expression plasmid also encodes T7 RNA polymerase required for transcription from a T7 promoter. The promoter can also direct transgene expression to the pancreas (see, e.g., the insulin regulatory sequence for pancreas (Bucchini et al., 1986, *Proc. Natl. Acad. Sci. USA* 83:2511-2515)).

[0097] In addition, expression of the transgene can be precisely regulated, for example, by using an inducible regulatory sequence and expression systems such as a regulatory sequence that is sensitive to certain physiological regulators, e.g., circulating glucose levels, or hormones (Docherty et al., 1994, *FASEB J.* 8:20-24). Such inducible expression systems, suitable for the control of transgene expression in cells or in mammals include regulation by ecdysone, by estrogen, progesterone, tetracycline, chemical inducers of dimerization, and isopropyl-beta-D1-thiogalactopyranoside (IPTG). A person skilled in the art would be able to choose the appropriate regulatory/promoter sequence based on the intended use of the dsRNA transgene.

[0098] Generally, recombinant vectors capable of expressing dsRNA molecules are delivered as described below, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of dsRNA molecules. Such vectors can be repeatedly administered as necessary. Once expressed, the dsRNAs bind to target RNA and modulate its function or expression. Delivery of dsRNA expressing vectors can be systemic, such as by intravenous or intramuscular administration, by administration to target cells ex-planted from the patient followed by reintroduction into the patient, or by any other means that allows for introduction into a desired target cell.

[0099] dsRNA expression DNA plasmids are typically transfected into target cells as a complex with cationic lipid carriers (e.g., Oligofectamine) or non-cationic lipid-based carriers (e.g., Transit-TKO™). Multiple lipid transfections for dsRNA-mediated knockdowns targeting different regions of a single TTR gene or multiple TTR genes over a period of a week or more are also contemplated by the invention. Successful introduction of vectors into host cells can be monitored using various known methods. For example, transient transfection can be signaled with a reporter, such as a fluorescent marker, such as Green Fluorescent Protein (GFP). Stable transfection of cells *ex vivo* can be ensured using markers that provide the transfected cell with resistance to specific environmental factors (e.g., antibiotics and drugs), such as hygromycin B resistance.

[0100] TTR specific dsRNA molecules can also be inserted into vectors and used as gene therapy vectors for human patients. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Patent 5,328,470) or by stereotactic injection (see e.g., Chen et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can include a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g., retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

III. Pharmaceutical compositions containing dsRNA

[0101] In one embodiment, the invention provides pharmaceutical compositions containing a dsRNA as claimed, and a pharmaceutically acceptable carrier. The pharmaceutical composition containing the dsRNA is useful for treating a disease or disorder associated with the expression or activity of a TTR gene, such as pathological processes mediated by TTR expression. Such pharmaceutical compositions are formulated based on the mode of delivery. One example is compositions that are formulated for systemic administration via parenteral delivery, e.g., by intravenous (IV) delivery. Another example is compositions that are formulated for direct delivery into the brain parenchyma, e.g., by infusion into the brain, such as by continuous pump infusion.

[0102] The pharmaceutical compositions featured herein are administered in dosages sufficient to inhibit expression of TTR genes.

[0103] In general, a suitable dose of dsRNA will be in the range of 0.01 to 200.0 milligrams per kilogram body weight of the recipient per day, generally in the range of 1 to 50 mg per kilogram body weight per day. For example, the dsRNA can be administered at 0.0059 mg/kg, 0.01 mg/kg, 0.0295 mg/kg, 0.05 mg/kg, 0.0590 mg/kg, 0.163 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 0.543 mg/kg, 0.5900 mg/kg, 0.6 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg, 1 mg/kg, 1.1 mg/kg, 1.2 mg/kg, 1.3 mg/kg, 1.4 mg/kg, 1.5 mg/kg, 1.628 mg/kg, 2 mg/kg, 3 mg/kg, 5.0 mg/kg, 10 mg/kg, 20 mg/kg, 30 mg/kg, 40 mg/kg, or 50 mg/kg per single dose.

[0104] In one implementation, the dosage is between 0.01 and 0.2 mg/kg. For example, the dsRNA can be administered at a dose of 0.01 mg/kg, 0.02 mg/kg, 0.3 mg/kg, 0.04 mg/kg, 0.05 mg/kg, 0.06 mg/kg, 0.07 mg/kg, 0.08 mg/kg, 0.09 mg/kg, 0.10 mg/kg, 0.11 mg/kg, 0.12 mg/kg, 0.13 mg/kg, 0.14 mg/kg, 0.15 mg/kg, 0.16 mg/kg, 0.17 mg/kg, 0.18 mg/kg, 0.19 mg/kg, or 0.20 mg/kg.

[0105] In one implementation, the dosage is between 0.005 mg/kg and 1.628 mg/kg. For example, the dsRNA can be administered at a dose of 0.0059 mg/kg, 0.0295 mg/kg, 0.0590 mg/kg, 0.163 mg/kg, 0.543 mg/kg, 0.5900 mg/kg, or 1.628 mg/kg.

[0106] In one implementation, the dosage is between 0.2 mg/kg and 1.5 mg/kg. For example, the dsRNA can be administered at a dose of 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 0.6 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg, 1 mg/kg, 1.1 mg/kg, 1.2 mg/kg, 1.3 mg/kg, 1.4 mg/kg, or 1.5 mg/kg.

[0107] The pharmaceutical composition may be administered once daily, or the dsRNA may be administered as two, three, or more sub-doses at appropriate intervals throughout the day or even using continuous infusion or delivery through a controlled release formulation. In that case, the dsRNA contained in each sub-dose must be correspondingly smaller in order to achieve the total daily dosage. The dosage unit can also be compounded for delivery over several days, e.g., using a conventional sustained release formulation which provides sustained release of the dsRNA over a several day period. Sustained release formulations are well known in the art and are particularly useful for delivery of agents at a particular site, such as could be used with the agents of the present invention. In this embodiment, the dosage unit contains a corresponding multiple of the daily dose.

[0108] The effect of a single dose on TTR levels is long lasting, such that subsequent doses are administered at not more than 3, 4, or 5 day intervals, or at not more than 1, 2, 3, or 4 week intervals, or at not more than 5, 6, 7, 8, 9, or 10 week intervals.

[0109] The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a composition can include a single treatment or a series of treatments. Estimates of effective dosages and *in vivo* half-lives for the individual dsRNAs encompassed by the invention can be made using conventional methodologies or on the basis of *in vivo* testing using an appropriate animal model, as described elsewhere herein.

[0110] Advances in mouse genetics have generated a number of mouse models for the study of various human diseases, such as pathological processes mediated by TTR expression. Such models are used for *in vivo* testing of dsRNA, as well as for determining a therapeutically effective dose. A suitable mouse model is, for example, a mouse containing a plasmid expressing human TTR. Another suitable mouse model is a transgenic mouse carrying a transgene that expresses human

TTR.

[0111] The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of compositions featured in the invention lies generally within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the methods featured in the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range of the compound or, when appropriate, of the polypeptide product of a target sequence (*e.g.*, achieving a decreased concentration of the polypeptide) that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0112] The dsRNAs featured in the invention can be administered in combination with other known agents effective in treatment of pathological processes mediated by target gene expression. In any event, the administering physician can adjust the amount and timing of dsRNA administration on the basis of results observed using standard measures of efficacy known in the art or described herein.

Administration

[0113] The present invention also includes pharmaceutical compositions and formulations which include the dsRNA compounds featured in the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical, pulmonary, *e.g.*, by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal, oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, *e.g.*, intraparenchymal, intrathecal or intraventricular, administration.

[0114] The dsRNA can be delivered in a manner to target a particular tissue, such as the liver (*e.g.*, the hepatocytes of the liver).

[0115] The present disclosure includes pharmaceutical compositions that can be delivered by injection directly into the brain. The injection can be by stereotactic injection into a particular region of the brain (*e.g.*, the substantia nigra, cortex, hippocampus, striatum, or globus pallidus), or the dsRNA can be delivered into multiple regions of the central nervous system (*e.g.*, into multiple regions of the brain, and/or into the spinal cord). The dsRNA can also be delivered into diffuse regions of the brain (*e.g.*, diffuse delivery to the cortex of the brain).

[0116] In one implementation, a dsRNA targeting TTR can be delivered by way of a cannula or other delivery device having one end implanted in a tissue, *e.g.*, the brain, *e.g.*, the substantia nigra, cortex, hippocampus, striatum, corpus callosum or globus pallidus of the brain. The cannula can be connected to a reservoir of the dsRNA composition. The flow or delivery can be mediated by a pump, *e.g.*, an osmotic pump or minipump, such as an Alzet pump (Durect, Cupertino, CA). In one embodiment, a pump and reservoir are implanted in an area distant from the tissue, *e.g.*, in the abdomen, and delivery is effected by a conduit leading from the pump or reservoir to the site of release. Infusion of the dsRNA composition into the brain can be over several hours or for several days, *e.g.*, for 1, 2, 3, 5, or 7 days or more. Devices for delivery to the brain are described, for example, in U.S. Patent Nos. 6,093,180, and 5,814,014.

[0117] Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Suitable topical formulations include those in which the dsRNAs featured in the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Suitable lipids and liposomes include neutral (*e.g.*, dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (*e.g.*, dimyristoylphosphatidyl glycerol DMPG) and cationic (*e.g.*, dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). DsRNAs featured in the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, dsRNAs may be complexed to lipids, in particular to cationic lipids. Suitable fatty acids and esters include but are not limited to arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid,

linoleic acid, linolenic acid, dicaprates, tricaprates, monoolein, dilaurin, glyceryl 1-monocaprates, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C₁₋₁₀ alkyl ester (e.g., isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in U.S. Patent No. 6,747,014, which is incorporated herein by reference.

Liposomal formulations

[0118] There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

[0119] Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

[0120] In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

[0121] Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

[0122] Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes and as the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

[0123] Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

[0124] Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

[0125] Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

[0126] Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).

[0127] One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl

phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

[0128] Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g., as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

[0129] Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

[0130] Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G_{M1}, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al., *Cancer Research*, 1993, 53, 3765).

[0131] Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside G_{M1}, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Pat. No. 4,837,028 and WO 88/04924, both to Allen *et al.*, disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G_{M1} or a galactocerebroside sulfate ester. U.S. Pat. No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

[0132] Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C₁₂15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Pat. Nos. 4,426,330 and 4,534,899). Klibanov et al. (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle *et al.* (U.S. Pat. Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Pat. No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Pat. No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Pat. No. 5,540,935 (Miyazaki et al.) and U.S. Pat. No. 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

[0133] A number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Pat. No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include a dsRNA. U.S. Pat. No.

5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes comprising dsRNAs targeted to the raf gene.

[0134] Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g., they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

[0135] Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

[0136] If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

[0137] If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

[0138] If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

[0139] If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

[0140] The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., 1988, p. 285).

Nucleic acid lipid particles

[0141] In one implementation, a TTR dsRNA featured in the invention is fully encapsulated in the lipid formulation, e.g., to form a SPLP, pSPLP, SNALP, or other nucleic acid-lipid particle. As used herein, the term "SNALP" refers to a stable nucleic acid-lipid particle, including SPLP. As used herein, the term "SPLP" refers to a nucleic acid-lipid particle comprising plasmid DNA encapsulated within a lipid vesicle. SNALPs and SPLPs typically contain a cationic lipid, a non-cationic lipid, and a lipid that prevents aggregation of the particle (e.g., a PEG-lipid conjugate). SNALPs and SPLPs are extremely useful for systemic applications, as they exhibit extended circulation lifetimes following intravenous (i.v.) injection and accumulate at distal sites (e.g., sites physically separated from the administration site). SPLPs include "pSPLP," which include an encapsulated condensing agent-nucleic acid complex as set forth in PCT Publication No. WO 00/03683. The particles of the present invention typically have a mean diameter of about 50 nm to about 150 nm, more typically about 60 nm to about 130 nm, more typically about 70 nm to about 110 nm, most typically about 70 nm to about 90 nm, and are substantially nontoxic. In addition, the nucleic acids when present in the nucleic acid-lipid particles of the present invention are resistant in aqueous solution to degradation with a nuclease. Nucleic acid-lipid particles and their method of preparation are disclosed in, e.g., U.S. Patent

Nos. 5,976,567; 5,981,501; 6,534,484; 6,586,410; 6,815,432; and PCT Publication No. WO 96/40964.

[0142] In one implementation, the lipid to drug ratio (mass/mass ratio) (e.g., lipid to dsRNA ratio) will be in the range of from about 1:1 to about 50:1, from about 1:1 to about 25:1, from about 3:1 to about 15:1, from about 4:1 to about 10:1, from about 5:1 to about 9:1, or about 6:1 to about 9:1.

[0143] The cationic lipid may be, for example, N,N-dioleoyl-N,N-dimethylammonium chloride (DODAC), N,N-distearyl-N,N-dimethylammonium bromide (DDAB), N-(1-(2,3-dioleoyloxy)propyl)-N,N,N-trimethylammonium chloride (DOTAP), N-(1-(2,3-dioleoyloxy)propyl)-N,N,N-trimethylammonium chloride (DOTMA), N,N-dimethyl-2,3-dioleoyloxypropylamine (DODMA), 1,2-DiLinoleoyloxy-N,N-dimethylaminopropane (DLinDMA), 1,2-Dilinolenyloxy-N,N-dimethylaminopropane (DLenDMA), 1,2-Dilinoleylcarbamoxyloxy-3-dimethylaminopropane (DLin-C-DAP), 1,2-Dilinoleoyloxy-3-(dimethylamino)acetoxyp propane (DLin-DAC), 1,2-Dilinoleoyloxy-3-morpholinopropane (DLin-MA), 1,2-Dilinoleoyl-3-dimethylaminopropane (DLinDAP), 1,2-Dilinoleylthio-3-dimethylaminopropane (DLin-S-DMA), 1-Linoleoyl-2-linoleoyloxy-3-dimethylaminopropane (DLin-2-DMA), 1,2-Dilinoleoyloxy-3-trimethylaminopropane chloride salt (DLin-TMA.Cl), 1,2-Dilinoleoyl-3-trimethylaminopropane chloride salt (DLin-TAP.Cl), 1,2-Dilinoleoyloxy-3-(N-methylpiperazino)propane (DLin-MPZ), or 3-(N,N-Dilinoleylamino)-1,2-propanediol (DLinAP), 3-(N,N-Dioleylamino)-1,2-propanediol (DOAP), 1,2-Dilinoleoyloxy-3-(2-N,N-dimethylamino)ethoxypropane (DLin-EG-DMA), 1,2-Dilinolenyloxy-N,N-dimethylaminopropane (DLinDMA), 2,2-Dilinoleyl-4-dimethylaminomethyl-[1,3]-dioxolane (DLin-K-DMA) or analogs thereof, (3aR,5s,6aS)-N,N-dimethyl-2,2-di((9Z,12Z)-octadeca-9,12-dienyl)tetrahydro-3aH-cyclopenta[d][1,3]dioxol-5-amine (ALN100), (6Z,9Z,28Z,31Z)-heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (MC3), 1,1'-(2-(4-(2-((bis(2-hydroxydodecyl)amino)ethyl)(2-hydroxydodecyl)amino)ethyl)piperazin-1-yl)ethylazanediyldidodecan-2-ol (Tech G1), or a mixture thereof. The cationic lipid may comprise from about 20 mol % to about 50 mol % or about 40 mol % of the total lipid present in the particle.

[0144] In another implementation, the compound 2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane can be used to prepare lipid-siRNA nanoparticles. Synthesis of 2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane is described in United States provisional patent application number 61/107,998 filed on October 23, 2008, which is herein incorporated by reference.

[0145] In one implementation, the lipid-siRNA particle includes 40% 2, 2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane: 10% DSPC: 40% Cholesterol: 10% PEG-C-DOMG (mole percent) with a particle size of 63.0 ± 20 nm and a 0.027 siRNA/Lipid Ratio.

[0146] The non-cationic lipid may be an anionic lipid or a neutral lipid including, but not limited to, distearoylphosphatidylcholine (DSPC), dioleoylphosphatidylcholine (DOPC), dipalmitoylphosphatidylcholine (DPPC), dioleoylphosphatidylglycerol (DOPG), dipalmitoylphosphatidylglycerol (DPPG), dioleoyl-phosphatidylethanolamine (DOPE), palmitoyloleoylphosphatidylcholine (POPC), palmitoyloleoylphosphatidylethanolamine (POPE), dioleoyl-phosphatidylethanolamine 4-(N-maleimidomethyl)-cyclohexane-1-carboxylate (DOPE-mal), dipalmitoyl phosphatidyl ethanolamine (DPPE), dimyristoylphosphoethanolamine (DMPE), distearoyl-phosphatidyl-ethanolamine (DSPE), 16-O-monomethyl PE, 16-O-dimethyl PE, 18-1 -trans PE, 1 -stearoyl-2-oleoyl- phosphatidylethanolamine (SOPE), cholesterol, or a mixture thereof. The non-cationic lipid may be from about 5 mol % to about 90 mol %, about 10 mol %, or about 58 mol % if cholesterol is included, of the total lipid present in the particle.


[0147] The conjugated lipid that inhibits aggregation of particles may be, for example, a polyethyleneglycol (PEG)-lipid including, without limitation, a PEG-diacylglycerol (DAG), a PEG-dialkyloxypropyl (DAA), a PEG-phospholipid, a PEG-ceramide (Cer), or a mixture thereof. The PEG-DAA conjugate may be, for example, a PEG-dilauryloxypropyl (C_{12}), a PEG-dimyristyloxypropyl (C_{14}), a PEG-dipalmitoyloxypropyl (C_{16}), or a PEG- distearyloxypropyl (C_{18}). The conjugated lipid that prevents aggregation of particles may be from 0 mol % to about 20 mol % or about 2 mol % of the total lipid present in the particle.

[0148] In some implementations, the nucleic acid-lipid particle further includes cholesterol at, e.g., about 10 mol % to about 60 mol % or about 48 mol % of the total lipid present in the particle.

LNP01

[0149] In one implementation, the lipidoid ND98-4HCl (MW 1487) (Formula 1), Cholesterol (Sigma-Aldrich), and PEG-Ceramide C16 (Avanti Polar Lipids) can be used to prepare lipid-siRNA nanoparticles (*i.e.*, LNP01 particles). Stock solutions of each in ethanol can be prepared as follows: ND98, 133 mg/ml; Cholesterol, 25 mg/ml, PEG-Ceramide C16, 100 mg/ml. The ND98, Cholesterol, and PEG-Ceramide C16 stock solutions can then be combined in a, *e.g.*, 42:48:10 molar ratio. The

combined lipid solution can be mixed with aqueous siRNA (e.g., in sodium acetate pH 5) such that the final ethanol concentration is about 35-45% and the final sodium acetate concentration is about 100-300 mM. Lipid-siRNA nanoparticles typically form spontaneously upon mixing. Depending on the desired particle size distribution, the resultant nanoparticle mixture can be extruded through a polycarbonate membrane (e.g., 100 nm cut-off) using, for example, a thermobarrel extruder, such as Lipex Extruder (Northern Lipids, Inc). In some cases, the extrusion step can be omitted. Ethanol removal and simultaneous buffer exchange can be accomplished by, for example, dialysis or tangential flow filtration. Buffer can be exchanged with, for example, phosphate buffered saline (PBS) at about pH 7, e.g., about pH 6.9, about pH 7.0, about pH 7.1, about pH 7.2, about pH 7.3, or about pH 7.4.



Formula 1

[0150] LNP01 formulations are described, e.g., in International Application Publication No. WO 2008/042973.

[0151] Additional exemplary lipid-siRNA formulations are as follows:

	Cationic Lipid	cationic lipid/non-cationic lipid/cholesterol/PEG-lipid conjugate	Process
		Lipid:siRNA ratio	
SNALP	1,2-Dilinolenyloxy-N,N-dimethylaminopropane (DLinDMA)	DLinDMA/DPPC/Cholesterol/PEG-cDMA	
		(57.1/7.1/34.4/1.4)	
		lipid:siRNA ~ 7:1	
SNALP-XTC	2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC)	XTC/DPPC/Cholesterol/PEG-cDMA	
		57.1/7.1/34.4/1.4	
		lipid:siRNA ~ 7:1	
LNP05	2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC)	XTC/DSPC/Cholesterol/PEG-DMG	Extrusion
		57.5/7.5/31.5/3.5	
		lipid:siRNA ~ 6:1	
LNP06	2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC)	XTC/DSPC/Cholesterol/PEG-DMG	Extrusion
		57.5/7.5/31.5/3.5	
		lipid:siRNA ~ 11:1	
LNP07	2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC)	XTC/DSPC/Cholesterol/PEG-DMG	In-line mixing
		60/7.5/31/1.5,	
		lipid:siRNA ~ 6:1	
LNP08	2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC)	XTC/DSPC/Cholesterol/PEG-DMG	In-line mixing
		60/7.5/31/1.5,	
		lipid:siRNA ~ 11:1	
LNP09	2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC)	XTC/DSPC/Cholesterol/PEG-DMG	In-line mixing
		50/10/38.5/1.5	
		Lipid:siRNA 10:1	
		ALN100/DSPC/Cholesterol/PEG-	

	Cationic Lipid	cationic lipid/non-cationic lipid/cholesterol/PEG-lipid conjugate	Process
		Lipid:siRNA ratio	
LNP10	(3aR,5s,6aS)-N,N-dimethyl-2,2-di((9Z,12Z)-octadeca-9,12-dienyl)tetrahydro-3aH-cyclopenta[d][1,3]dioxol-5-amine (ALN100)	DMG	In-line mixing
		50/10/38.5/1.5	
		Lipid:siRNA 10:1	
LNP11	(6Z,9Z,28Z,31Z)-heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (MC3)	MC-3/DSPC/Cholesterol/PEG-DMG	In-line mixing
		50/10/38.5/1.5	
		Lipid:siRNA 10:1	
LNP12	1,1'-(2-(4-(2-((2-(bis(2-hydroxydodecyl)amino)ethyl)(2-hydroxydodecyl)amino)ethyl)piperazin-1-yl)ethylazanediyldidodecan-2-ol (Tech G1)	Tech G1/DSPC/Cholesterol/PEG-DMG	In-line mixing
		50/10/38.5/1.5	
		Lipid:siRNA 10:1	

[0152] LNP09 formulations and XTC comprising formulations are described, e.g., in U.S. Provisional Serial No. 61/239,686, filed September 3, 2009, which is hereby incorporated by reference. LNP11 formulations and MC3 comprising formulations are described, e.g., in U.S. Provisional Serial No. 61/244,834, filed September 22, 2009.

[0153] Formulations prepared by either the standard or extrusion-free method can be characterized in similar manners. For example, formulations are typically characterized by visual inspection. They should be whitish translucent solutions free from aggregates or sediment. Particle size and particle size distribution of lipid-nanoparticles can be measured by light scattering using, for example, a Malvern Zetasizer Nano ZS (Malvern, USA). Particles should be about 20-300 nm, such as 40-100 nm in size. The particle size distribution should be unimodal. The total siRNA concentration in the formulation, as well as the entrapped fraction, is estimated using a dye exclusion assay. A sample of the formulated siRNA can be incubated with an RNA-binding dye, such as Ribogreen (Molecular Probes) in the presence or absence of a formulation disrupting surfactant, e.g., 0.5% Triton-X100. The total siRNA in the formulation can be determined by the signal from the sample containing the surfactant, relative to a standard curve. The entrapped fraction is determined by subtracting the "free" siRNA content (as measured by the signal in the absence of surfactant) from the total siRNA content. Percent entrapped siRNA is typically >85%. For SNALP formulation, the particle size is at least 30 nm, at least 40 nm, at least 50 nm, at least 60 nm, at least 70 nm, at least 80 nm, at least 90 nm, at least 100 nm, at least 110 nm, and at least 120 nm. The suitable range is typically about at least 50 nm to about at least 110 nm, about at least 60 nm to about at least 100 nm, or about at least 80 nm to about at least 90 nm.

[0154] Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. In some embodiments, oral formulations are those in which dsRNAs featured in the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Suitable surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Suitable bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxychenodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucolic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate and sodium glycodihydrofusidate. Suitable fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein, dilaurin, glyceryl 1-monocaprinate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g., sodium). In some embodiments, combinations of penetration enhancers are used, for example, fatty acids/salts in combination with bile acids/salts. One exemplary combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. DsRNAs featured in the invention may be delivered orally, in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. DsRNA complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and starches; polyalkylcyanoacrylates; DEAE-derivatized polyimines, pullulans, celluloses and starches. Suitable complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines, protamine, polyvinylpyridine, polythiodiethylaminomethylethylene P(TDAE), polyaminostyrene (e.g., p-amino),

poly(methylcyanoacrylate), poly(ethylcyanoacrylate), poly(butylcyanoacrylate), poly(isobutylcyanoacrylate), poly(isohexylcyanoacrylate), DEAE-methacrylate, DEAE-hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid), poly(DL-lactic-co-glycolic acid (PLGA), alginate, and polyethyleneglycol (PEG). Oral formulations for dsRNAs and their preparation are described in detail in U.S. Patent 6,887,906, US Publn. No. 20030027780, and U.S. Patent No. 6,747,014, each of which is incorporated herein by reference.

[0155] Compositions and formulations for parenteral, intraparenchymal (into the brain), intrathecal, intraventricular or intrahepatic administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

[0156] Pharmaceutical compositions of the present disclosure include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids. Particularly preferred are formulations that target the liver when treating hepatic disorders such as hepatic carcinoma.

[0157] The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

[0158] The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Emulsions

[0159] The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogeneous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μm in diameter (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa., 1985, p. 301). Emulsions are often biphasic systems comprising two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be of either the water-in-oil (w/o) or the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase, the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase, the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases, and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous phase provides an o/w/o emulsion.

[0160] Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

[0161] Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

[0162] Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgate, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

[0163] A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

[0164] Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

[0165] Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

[0166] The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of ease of formulation, as well as efficacy from an absorption and bioavailability standpoint (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

[0167] In one implementation, the compositions of dsRNAs and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and

geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa., 1985, p. 271).

[0168] The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

[0169] Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

[0170] Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., Pharmaceutical Research, 1994, 11, 1385-1390; Ritschel, Meth. Find. Exp. Clin. Pharmacol., 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., Pharmaceutical Research, 1994, 11, 1385; Ho et al., J. Pharm. Sci., 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or dsRNAs. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of dsRNAs and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of dsRNAs and nucleic acids.

[0171] Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the dsRNAs and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories--surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p. 92). Each of these classes has been discussed above.

Penetration Enhancers

[0172] In one implementation, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly dsRNAs, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

[0173] Penetration enhancers may be classified as belonging to one of five broad categories, *i.e.*, surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

[0174] Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of dsRNAs through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

[0175] Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprinate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C.sub.1-10 alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (*i.e.*, oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, *etc.*) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

[0176] Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. Suitable bile salts include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: Remington's *Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, Pa., 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

[0177] Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of dsRNAs through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Suitable chelating agents include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-diketones (enamines)(Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

[0178] Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of dsRNAs through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Carriers

[0179] Certain compositions of the present disclosure also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (*i.e.*, does not possess biological activity per se) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the

recovery of a partially phosphorothioate dsRNA in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., DsRNA Res. Dev., 1995, 5, 115-121; Takakura et al., DsRNA & Nucl. Acid Drug Dev., 1996, 6, 177-183).

Excipients

[0180] In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, *etc.*, when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (*e.g.*, pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, *etc.*); fillers (*e.g.*, lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, *etc.*); lubricants (*e.g.*, magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, *etc.*); disintegrants (*e.g.*, starch, sodium starch glycolate, *etc.*); and wetting agents (*e.g.*, sodium lauryl sulphate, *etc.*).

[0181] Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

[0182] Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

[0183] Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Other Components

[0184] The compositions of the present disclosure may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, *e.g.*, lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

[0185] Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

[0186] In some implementations, pharmaceutical compositions featured in the invention include (a) one or more dsRNA compounds and (b) one or more anti-cytokine biologic agents which function by a non-RNAi mechanism. Examples of such biologies include, biologies that target IL1 β (*e.g.*, anakinra), IL6 (tocilizumab), or TNF (etanercept, infliximab, adlimumab, or certolizumab).

[0187] Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the

therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds that exhibit high therapeutic indices are preferred.

[0188] The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of compositions featured in the invention lies generally within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the methods featured in the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range of the compound or, when appropriate, of the polypeptide product of a target sequence (*e.g.*, achieving a decreased concentration of the polypeptide) that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0189] In addition to their administration, as discussed above, the dsRNAs featured in the invention can be administered in combination with other known agents effective in treatment of pathological processes mediated by TTR expression. In any event, the administering physician can adjust the amount and timing of dsRNA administration on the basis of results observed using standard measures of efficacy known in the art or described herein.

Methods for treating diseases caused by expression of a TTR gene

[0190] The disclosure relates in particular to the use of a dsRNA targeting TTR and compositions containing at least one such dsRNA for the treatment of a TTR-mediated disorder or disease. For example, a dsRNA targeting a TTR gene can be useful for the treatment of a TTR amyloidosis, such as familial amyloidotic polyneuropathy (FAP), familial amyloidotic cardiomyopathy (FAC), leptomeningeal/CNS amyloidosis, amyloidosis VII form (also known as leptomeningeal or meningocerebrovascular amyloidosis), hyperthyroxinemia, and cardiac amyloidosis (also called senile systemic amyloidosis (SSA) and senile cardiac amyloidosis (SCA)).

[0191] FIG. 15 illustrates symptoms and mutations in TTR associated with familial amyloidotic neuropathy, familial amyloidotic cardiomyopathy and CNS amyloidosis. The invention includes compositions and methods for treatment of these diseases and symptoms, and directed to these mutant versions of TTR,

[0192] A dsRNA targeting a TTR gene is also used for treatment of symptoms and disorders, such as TTR amyloidosis. Symptoms associated with such amyloidosis include, *e.g.*, seizures, dementia, myelopathy, polyneuropathy, carpal tunnel syndrome, autonomic insufficiency, cardiomyopathy, gastrointestinal dysfunction (*e.g.*, gastric ulcers, diarrhea, constipation, malabsorption), weight loss, hepatomegaly, lymphadenopathy, goiter, vitreous opacities, renal insufficiency (including proteinuria and kidney failure), nephropathy, cranial nerve dysfunction, corneal lattice dystrophy, and congestive heart failure with generalized weakness and difficulties breathing from fluid retention.

[0193] Owing to the inhibitory effects on TTR expression, a composition according to the invention or a pharmaceutical composition prepared therefrom can enhance the quality of life.

[0194] The disclosure further relates to the use of a dsRNA or a pharmaceutical composition thereof, *e.g.*, for treating a TTR amyloidosis, in combination with other pharmaceuticals and/or other therapeutic methods, *e.g.*, with known pharmaceuticals and/or known therapeutic methods, such as, for example, those which are currently employed for treating these disorders. In one example, a dsRNA targeting TTR can be administered in combination with a liver transplant. In other examples, a dsRNA targeting TTR can be administered in combination with a pharmaceutical or therapeutic method for treating a symptom of a TTR disease, such as diuretics, ACE (angiotensin converting enzyme) inhibitors, angiotensin receptor blockers (ARBs), or dialysis, *e.g.*, for management of renal function.

[0195] The dsRNA and an additional therapeutic agent can be administered in the same combination, *e.g.*, parenterally, or the additional therapeutic agent can be administered as part of a separate composition or by another method described herein.

[0196] The disclosure features a method of administering a dsRNA targeting TTR to a patient having a disease or disorder mediated by TTR expression, such as a TTR amyloidosis, *e.g.*, FAP. Administration of the dsRNA can stabilize and improve peripheral neurological function, for example, in a patient with FAP. Patients can be administered a therapeutic amount of

dsRNA, such as 0.1 mg/kg, 0.2 mg/kg, 0.5 mg/kg, 1.0 mg/kg, 1.5 mg/kg, 2.0 mg/kg, or 2.5 mg/kg dsRNA. The dsRNA can be administered by intravenous infusion over a period of time, such as over a 5 minute, 10 minute, 15 minute, 20 minute, 25 minute, 60 minute, 120 minute or 180 minute period. The administration is repeated, for example, on a regular basis, such as biweekly (*i. e.*, every two weeks) for one month, two months, three months, four months or longer. After an initial treatment regimen, the treatments can be administered on a less frequent basis. For example, after administration biweekly for three months, administration can be repeated once per month, for six months or a year or longer. Administration of the dsRNA can reduce TTR levels in the blood or urine of the patient by at least 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80 % or 90% or more.

[0197] Before administration of a full dose of the dsRNA, patients can be administered a smaller dose, such as a dose that is 5% of the full dose, and monitored for adverse effects, such as an allergic reaction or a change in liver function. For example, in patients monitored for changes in liver function, a low incidence of LFT (Liver Function Test) change (*e.g.*, a 10-20% incidence of LFT) is acceptable (*e.g.*, a reversible, 3-fold increase in ALT (alanine aminotransferase) and/or AST (aspartate aminotransferase) levels).

[0198] Many TTR-associated diseases and disorders are hereditary. Therefore, a patient in need of a TTR dsRNA can be identified by taking a family history. A healthcare provider, such as a doctor, nurse, or family member, can take a family history before prescribing or administering a TTR dsRNA. A DNA test may also be performed on the patient to identify a mutation in the TTR gene, before a TTR dsRNA is administered to the patient.

[0199] The patient may have a biopsy performed before receiving a TTR dsRNA. The biopsy can be, for example, on a tissue, such as the gastric mucosa, peripheral nerve, skin, abdominal fat, liver, or kidney, and the biopsy may reveal amyloid plaques, which are indicative of a TTR-mediated disorder. Upon the identification of amyloid plaques, the patient is administered a TTR dsRNA.

Methods for inhibiting expression of a TTR gene

[0200] In yet another aspect, the disclosure provides a method for inhibiting the expression of a TTR gene in a mammal. The method includes administering a composition featured in the invention to the mammal such that expression of the target TTR gene is silenced.

[0201] When the organism to be treated is a mammal such as a human, the composition may be administered by any means known in the art including, but not limited to oral or parenteral routes, including intracranial (*e.g.*, intraventricular, intraparenchymal and intrathecal), intravenous, intramuscular, subcutaneous, transdermal, airway (aerosol), nasal, rectal, and topical (including buccal and sublingual) administration. In certain embodiments, the compositions are administered by intravenous infusion or injection.

[0202] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

EXAMPLES

Example 1, dsRNA synthesis

Source of reagents

[0203] Where the source of a reagent is not specifically given herein, such reagent may be obtained from any supplier of reagents for molecular biology at a quality/purity standard for application in molecular biology.

siRNA synthesis


[0204] Single-stranded RNAs were produced by solid phase synthesis on a scale of 1 μ mole using an Expedite 8909 synthesizer (Applied Biosystems, Applera Deutschland GmbH, Darmstadt, Germany) and controlled pore glass (CPG, 500Å, Proligo Biochemie GmbH, Hamburg, Germany) as solid support. RNA and RNA containing 2'-O-methyl nucleotides were generated by solid phase synthesis employing the corresponding phosphoramidites and 2'-O-methyl phosphoramidites, respectively (Proligo Biochemie GmbH, Hamburg, Germany). These building blocks were incorporated at selected sites within the sequence of the oligoribonucleotide chain using standard nucleoside phosphoramidite chemistry such as described in Current protocols in nucleic acid chemistry, Beaucage, S.L. et al. (Edrs.), John Wiley & Sons, Inc., New York, NY, USA. Phosphorothioate linkages were introduced by replacement of the iodine oxidizer solution with a solution of the Beaucage reagent (Chruachem Ltd, Glasgow, UK) in acetonitrile (1%). Further ancillary reagents were obtained from Mallinckrodt Baker (Griesheim, Germany).

[0205] Deprotection and purification of the crude oligoribonucleotides by anion exchange HPLC were carried out according to established procedures. Yields and concentrations were determined by UV absorption of a solution of the respective RNA at a wavelength of 260 nm using a spectral photometer (DU 640B, Beckman Coulter GmbH, Unterschleißheim, Germany). Double stranded RNA was generated by mixing an equimolar solution of complementary strands in annealing buffer (20 mM sodium phosphate, pH 6.8; 100 mM sodium chloride), heated in a water bath at 85 - 90°C for 3 minutes and cooled to room temperature over a period of 3 - 4 hours. The annealed RNA solution was stored at -20°C until use.

[0206] For the synthesis of 3'-cholesterol-conjugated siRNAs (herein referred to as -Chol-3'), an appropriately modified solid support was used for RNA synthesis. The modified solid support was prepared as follows:

Diethyl-2-azabutane-1,4-dicarboxylate AA


[0207]



[0208] A 4.7 M aqueous solution of sodium hydroxide (50 mL) was added into a stirred, ice-cooled solution of ethyl glycinate hydrochloride (32.19 g, 0.23 mole) in water (50 mL). Then, ethyl acrylate (23.1 g, 0.23 mole) was added and the mixture was stirred at room temperature until completion of the reaction was ascertained by TLC. After 19 h the solution was partitioned with dichloromethane (3 x 100 mL). The organic layer was dried with anhydrous sodium sulfate, filtered and evaporated. The residue was distilled to afford AA (28.8 g, 61 %).

3-{Ethoxycarbonylmethyl-[6-(9H-fluoren-9-ylmethoxycarbonyl-amino)-hexanoyl]-amino}-propionic acid ethyl ester AB


[0209]



[0210] Fmoc-6-amino-hexanoic acid (9.12 g, 25.83 mmol) was dissolved in dichloromethane (50 mL) and cooled with ice. Diisopropylcarbodiimide (3.25 g, 3.99 mL, 25.83 mmol) was added to the solution at 0°C. It was then followed by the addition of Diethyl-azabutane-1,4-dicarboxylate (5 g, 24.6 mmol) and dimethylamino pyridine (0.305 g, 2.5 mmol). The solution was brought to room temperature and stirred further for 6 h. Completion of the reaction was ascertained by TLC. The reaction mixture was concentrated under vacuum and ethyl acetate was added to precipitate diisopropyl urea. The suspension was filtered. The filtrate was washed with 5% aqueous hydrochloric acid, 5% sodium bicarbonate and water. The combined organic layer was dried over sodium sulfate and concentrated to give the crude product which was purified by column chromatography (50 % EtOAc/Hexanes) to yield 11.87 g (88%) of AB.

3-[(6-Amino-hexanoyl)-ethoxycarbonylmethyl-amino]-propionic acid ethyl ester AC


[0211]



[0212] 3-{Ethoxycarbonylmethyl-[6-(9H-fluoren-9-ylmethoxycarbonylamino)-hexanoyl]-amino}-propionic acid ethyl ester AB (11.5 g, 21.3 mmol) was dissolved in 20% piperidine in dimethylformamide at 0°C. The solution was continued stirring for 1 h. The reaction mixture was concentrated under vacuum, water was added to the residue, and the product was extracted with ethyl acetate. The crude product was purified by conversion into its hydrochloride salt.

3-{6-[17-(1,5-Dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yloxy-carbonylamino]-hexanoyl}ethoxycarbonylmethyl-amino}-propionic acid ethyl ester AD


[0213]



[0214] The hydrochloride salt of 3-[(6-Amino-hexanoyl)-ethoxycarbonylmethyl-amino]-propionic acid ethyl ester AC (4.7 g, 14.8 mmol) was taken up in dichloromethane. The suspension was cooled to 0°C on ice. To the suspension diisopropylethylamine (3.87 g, 5.2 mL, 30 mmol) was added. To the resulting solution cholesteryl chloroformate (6.675 g, 14.8 mmol) was added. The reaction mixture was stirred overnight. The reaction mixture was diluted with dichloromethane and washed with 10% hydrochloric acid. The product was purified by flash chromatography (10.3 g, 92%).

1-{6-[17-(1,5-Dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yloxy-carbonylamino]-hexanoyl}-4-oxo-pyrrolidine-3-carboxylic acid ethyl ester AE

[0215]




[0216] Potassium t-butoxide (1.1 g, 9.8 mmol) was slurried in 30 mL of dry toluene. The mixture was cooled to 0°C on ice and 5 g (6.6 mmol) of diester AD was added slowly with stirring within 20 mins. The temperature was kept below 5°C during the addition. The stirring was continued for 30 mins at 0°C and 1 mL of glacial acetic acid was added, immediately followed by 4 g of NaH₂PO₄·H₂O in 40 mL of water. The resultant mixture was extracted twice with 100 mL of dichloromethane each and

the combined organic extracts were washed twice with 10 mL of phosphate buffer each, dried, and evaporated to dryness. The residue was dissolved in 60 mL of toluene, cooled to 0°C and extracted with three 50 mL portions of cold pH 9.5 carbonate buffer. The aqueous extracts were adjusted to pH 3 with phosphoric acid, and extracted with five 40 mL portions of chloroform which were combined, dried and evaporated to dryness. The residue was purified by column chromatography using 25% ethylacetate/hexane to afford 1.9 g ofb-ketoester (39%).

[6-(3-Hydroxy-4-hydroxymethyl-pyrrolidin-1-yl)-6-oxo-hexyl]-carbamic acid 17-(1,5-dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yl ester AF


[0217]



[0218] Methanol (2 mL) was added dropwise over a period of 1 h to a refluxing mixture ofb-ketoester AE (1.5 g, 2.2 mmol) and sodium borohydride (0.226 g, 6 mmol) in tetrahydrofuran (10 mL). Stirring was continued at reflux temperature for 1 h. After cooling to room temperature, 1 N HCl (12.5 mL) was added, the mixture was extracted with ethylacetate (3 x 40 mL). The combined ethylacetate layer was dried over anhydrous sodium sulfate and concentrated under vacuum to yield the product which was purified by column chromatography (10% MeOH/CHCl₃) (89%).

(6-[3-[Bis-(4-methoxy-phenyl)-phenyl-methoxymethyl]-4-hydroxy-pyrrolidin-1-yl]-6-oxo-hexyl)-carbamic acid 17-(1,5-dimethyl-hexyl)-10,13-dimethyl-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H-cyclopenta[a]phenanthren-3-yl ester AG


[0219]



[0220] Diol AF (1.25 gm 1.994 mmol) was dried by evaporating with pyridine (2 x 5 mL) *in vacuo*. Anhydrous pyridine (10 mL) and 4,4'-dimethoxytritylchloride (0.724 g, 2.13 mmol) were added with stirring. The reaction was carried out at room temperature overnight. The reaction was quenched by the addition of methanol. The reaction mixture was concentrated under vacuum and to the residue dichloromethane (50 mL) was added. The organic layer was washed with 1M aqueous sodium bicarbonate. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The residual pyridine was removed by evaporating with toluene. The crude product was purified by column chromatography (2% MeOH/Chloroform, R_f = 0.5 in 5% MeOH/CHCl₃) (1.75 g, 95%).

Succinic acid mono-(4-[bis-(4-methoxy-phenyl)-phenyl-methoxymethyl]-1-{6-[17-(1,5-dimethyl-hexyl)-10,13-dimethyl 2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1H cyclopenta[a]phenanthren-3-yloxycarbonylamino]-hexanoyl)-pyrrolidin-3-yl) ester AH


[0221]

**AH**

[0222] Compound AG (1.0 g, 1.05 mmol) was mixed with succinic anhydride (0.150 g, 1.5 mmol) and DMAP (0.073 g, 0.6 mmol) and dried in a vacuum at 40°C overnight. The mixture was dissolved in anhydrous dichloroethane (3 mL), triethylamine (0.318 g, 0.440 mL, 3.15 mmol) was added and the solution was stirred at room temperature under argon atmosphere for 16 h. It was then diluted with dichloromethane (40 mL) and washed with ice cold aqueous citric acid (5 wt%, 30 mL) and water (2 X 20 mL). The organic phase was dried over anhydrous sodium sulfate and concentrated to dryness. The residue was used as such for the next step.

Cholesterol derivatised CPG AI

[0223]

**AI**

[0224] Succinate AH (0.254 g, 0.242 mmol) was dissolved in a mixture of dichloromethane/acetonitrile (3:2, 3 mL). To that solution DMAP (0.0296 g, 0.242 mmol) in acetonitrile (1.25 mL), 2,2'-Dithio-bis(5-nitropyridine) (0.075 g, 0.242 mmol) in acetonitrile/dichloroethane (3:1, 1.25 mL) were added successively. To the resulting solution triphenylphosphine (0.064 g, 0.242 mmol) in acetonitrile (0.6 mL) was added. The reaction mixture turned bright orange in color. The solution was agitated briefly using a wrist-action shaker (5 mins). Long chain alkyl amine-CPG (LCAA-CPG) (1.5 g, 61 mM) was added. The suspension was agitated for 2 h. The CPG was filtered through a sintered funnel and washed with acetonitrile, dichloromethane and ether successively. Unreacted amino groups were masked using acetic anhydride/pyridine. The achieved loading of the CPG was measured by taking UV measurement (37 mM/g).

[0225] The synthesis of siRNAs bearing a 5'-12-dodecanoic acid bisdecylamide group (herein referred to as "5'-C32-") or a 5'-cholesteryl derivative group (herein referred to as "5'-Chol-") was performed as described in WO 2004/065601, except that, for the cholesteryl derivative, the oxidation step was performed using the Beaucage reagent in order to introduce a phosphorothioate linkage at the 5'-end of the nucleic acid oligomer.

[0226] Nucleic acid sequences are represented below using standard nomenclature, and specifically the abbreviations of Table 1.

Table 1: Abbreviations of nucleotide monomers used in nucleic acid sequence representation. It will be understood that these monomers, when present in an oligonucleotide, are mutually linked by 5'-3'-phosphodiester bonds.

Abbreviation	Nucleotide(s)
A	adenosine-3'-phosphate
C	cytidine-3'-phosphate

Abbreviation	Nucleotide(s)
G	guanosine-3'-phosphate
T	5-methyluridine-3'-phosphate
U	uridine-3'-phosphate
N	any nucleotide (G, A, C, or T)
a	2'-O-methyladenosine-3'-phosphate
C	2'-O-methylcytidine-3'-phosphate
g	2'-O-methylguanosine-3'-phosphate
u	2'-O-methyluridine-3'-phosphate
dT	2'-deoxythymidine-3'-phosphate
sT; sdT	2'-deoxy-thymidine-5'phosphate-phosphorothioate

Example 2A. TTR siRNA Design

Transcripts

[0227] siRNA design was carried out to identify siRNAs targeting the gene transthyretin from human (symbol TTR) and rat (symbol Ttr). The design used the TTR transcripts NM_000371.2 (SEQ ID NO:1329) (human) and NM_012681.1 (SEQ ID NO:1330) (rat) from the NCBI Refseq collection. The siRNA duplexes were designed with 100% identity to their respective TTR genes.

siRNA Design and Specificity Prediction

[0228] The predicted specificity of all possible 19mers was determined for each sequence. The TTR siRNAs were used in a comprehensive search against the human and rat transcriptomes (defined as the set of NM_ and XM_ records within the NCBI Refseq set) using the FASTA algorithm. The Python script 'offtargetFasta.py' was then used to parse the alignments and generate a score based on the position and number of mismatches between the siRNA and any potential 'off-target' transcript. The off-target score is weighted to emphasize differences in the 'seed' region of siRNAs, in positions 2-9 from the 5' end of the molecule. The off-target score is calculated as follows: mismatches between the oligo and the transcript are given penalties. A mismatch in the seed region in positions 2-9 of the oligo is given a penalty of 2.8; mismatches in the putative cleavage sites 10 and 11 are given a penalty of 1.2, and mismatches in positions 12-19 a penalty of 1. Mismatches in position 1 are not considered. The off-target score for each oligo-transcript pair is then calculated by summing the mismatch penalties. The lowest off-target score from all the oligo-transcript pairs is then determined and used in subsequent sorting of oligos. Both siRNA strands were assigned to a category of specificity according to the calculated scores: a score above 3 qualifies as highly specific, equal to 3 as specific, and between 2.2 and 2.8 as moderately specific. In picking which oligos to synthesize, off-target scores of the antisense strand were sorted from high to low, and the 144 best (lowest off-target score) oligo pairs from human, and the best 26 pairs from rat were selected.

siRNA sequence selection

[0229] A total of 140 sense and 140 antisense human TTR derived siRNA oligos were synthesized and formed into duplexes. A total of 26 sense and 26 antisense rat TTR derived siRNA oligos were synthesized and formed into duplexes. Duplexes included The oligos are presented in Tables 2-4 (human TTR) and Tables 5-7 (rat TTR).

Table 2. Identification numbers for human TTR dsRNAs

See Table 4 for sequences and modifications of oligos.		
Duplex #	Sense Oligo #	Antisense Oligo #
AD-18243	A-32153	A-32154
AD-18244	A-32155	A-32156

See Table 4 for sequences and modifications of oligos.		
Duplex #	Sense Oligo #	Antisense Oligo #
AD-18245	A-32157	A-32158
AD-18246	A-32159	A-32160
AD-18247	A-32163	A-32164
AD-18248	A-32165	A-32166
AD-18249	A-32167	A-32168
AD-18250	A-32169	A-32170
AD-18251	A-32171	A-32172
AD-18252	A-32175	A-32176
AD-18253	A-32177	A-32178
AD-18254	A-32179	A-32180
AD-18255	A-32181	A-32182
AD-18256	A-32183	A-32184
AD-18257	A-32187	A-32188
AD-18258	A-32189	A-32190
AD-18259	A-32191	A-32192
AD-18260	A-32193	A-32194
AD-18261	A-32195	A-32196
AD-18262	A-32199	A-32200
AD-18263	A-32201	A-32202
AD-18264	A-32203	A-32204
AD-18265	A-32205	A-32206
AD-18266	A-32207	A-32208
AD-18267	A-32211	A-32212
AD-18268	A-32213	A-32214
AD-18269	A-32215	A-32216
AD-18270	A-32217	A-32218
AD-18271	A-32219	A-32220
AD-18272	A-32221	A-32222
AD-18273	A-32223	A-32224
AD-18274	A-32225	A-32226
AD-18275	A-32227	A-32228
AD-18276	A-32229	A-32230
AD-18277	A-32231	A-32232
AD-18278	A-32233	A-32234
AD-18279	A-32235	A-32236
AD-18280	A-32237	A-32238
AD-18281	A-32239	A-32240
AD-18282	A-32241	A-32242
AD-18283	A-32243	A-32244
AD-18284	A-32247	A-32248
AD-18285	A-32249	A-32250
AD-18286	A-32251	A-32252
AD-18287	A-32253	A-32254
AD-18288	A-32255	A-32256
AD-18289	A-32259	A-32260

See Table 4 for sequences and modifications of oligos.		
Duplex #	Sense Oligo #	Antisense Oligo #
AD-18290	A-32261	A-32262
AD-18291	A-32263	A-32264
AD-18292	A-32265	A-32266
AD-18293	A-32267	A-32268
AD-18294	A-32269	A-32270
AD-18295	A-32271	A-32272
AD-18296	A-32273	A-32274
AD-18297	A-32275	A-32276
AD-18298	A-32277	A-32278
AD-18299	A-32279	A-32280
AD-18300	A-32281	A-32282
AD-18301	A-32283	A-32284
AD-18302	A-32285	A-32286
AD-18303	A-32287	A-32288
AD-18304	A-32289	A-32290
AD-18305	A-32291	A-32292
AD-18306	A-32295	A-32296
AD-18307	A-32297	A-32298
AD-18308	A-32299	A-32300
AD-18309	A-32301	A-32302
AD-18310	A-32303	A-32304
AD-18311	A-32307	A-32308
AD-18312	A-32309	A-32310
AD-18313	A-32311	A-32312
AD-18314	A-32313	A-32314
AD-18315	A-32315	A-32316
AD-18316	A-32319	A-32320
AD-18317	A-32321	A-32322
AD-18318	A-32323	A-32324
AD-18319	A-32325	A-32326
AD-18320	A-32327	A-32328
AD-18321	A-32331	A-32332
AD-18322	A-32333	A-32334
AD-18323	A-32335	A-32336
AD-18324	A-32337	A-32338
AD-18325	A-32339	A-32340
AD-18326	A-32341	A-32342
AD-18327	A-32343	A-32344
AD-18328	A-32345	A-32346
AD-18329	A-32347	A-32348
AD-18330	A-32349	A-32350
AD-18331	A-32351	A-32352
AD-18332	A-32353	A-32354
AD-18333	A-32355	A-32356
AD-18334	A-32357	A-32358

See Table 4 for sequences and modifications of oligos.

Duplex #	Sense Oligo #	Antisense Oligo #
AD-18335	A-32359	A-32360
AD-18336	A-32363	A-32364
AD-18337	A-32367	A-32368
AD-18338	A-32369	A-32370
AD-18339	A-32371	A-32372
AD-18340	A-32373	A-32374
AD-18341	A-32375	A-32376
AD-18342	A-32379	A-32380
AD-18343	A-32381	A-32382
AD-18344	A-32383	A-32384
AD-18345	A-32385	A-32386
AD-18346	A-32387	A-32388
AD-18347	A-32391	A-32392
AD-18348	A-32393	A-32394
AD-18349	A-32395	A-32396
AD-18350	A-32397	A-32398
AD-18351	A-32399	A-32400
AD-18352	A-32401	A-32402
AD-18353	A-32403	A-32404
AD-18354	A-32405	A-32406
AD-18355	A-32407	A-32408
AD-18356	A-32409	A-32410
AD-18357	A-32411	A-32412
AD-18358	A-32415	A-32416
AD-18359	A-32417	A-32418
AD-18360	A-32419	A-32420
AD-18361	A-32421	A-32422
AD-18362	A-32423	A-32424
AD-18363	A-32427	A-32428
AD-18364	A-32429	A-32430
AD-18446	A-32161	A-32162
AD-18447	A-32173	A-32174
AD-18448	A-32185	A-32186
AD-18449	A-32197	A-32198
AD-18450	A-32209	A-32210
AD-18451	A-32245	A-32246
AD-18452	A-32257	A-32258
AD-18453	A-32293	A-32294
AD-18454	A-32305	A-32306
AD-18455	A-32317	A-32318
AD-18456	A-32329	A-32330
AD-18457	A-32361	A-32362
AD-18458	A-32365	A-32366
AD-18459	A-32377	A-32378
AD-18460	A-32389	A-32390

See Table 4 for sequences and modifications of oligos.		
Duplex #	Sense Oligo #	Antisense Oligo #
AD-18461	A-32401	A-32402
AD-18462	A-32413	A-32414
AD-18463	A-32425	A-32426

Table 3A. Sense and antisense strand sequences of human TTR dsRNAs

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)					
Strand	Position	Sequence (5' to 3')	SEQ ID NO:	Sequence with 3' dinucleotide overhang (5' to 3')	SEQ ID NO:
S	100	CCGGUGAAUCCAAGUGUCC	1	CCGGUGAAUCCAAGUGUCCNN	281
as	118	GGACACUUGGAUUCACCGG	2	GGACACUUGGAUUCACCGGNN	282
S	11	ACUCAUUCUUGGCAGGAUG	3	ACUCAUUCUUGGCAGGAUGNN	283
as	29	CAUCCUGCCAAGAAUGAGU	4	CAUCCUGCCAAGAAUGAGUNN	284
S	111	AAGUGUCCUCUGAUGGUCA	5	AAGUGUCCUCUGAUGGUCCANN	285
as	129	UGACCAUCAGAGGACACUU	6	UGACCAUCAGAGGACACUUNN	286
S	13	UCAUUCUUGGCAGGAUGGC	7	UCAUUCUUGGCAGGAUGGCNN	287
as	31	GCCAUCCUGCCAAGAAUGA	8	GCCAUCCUGCCAAGAAUGANN	288
s	130	AAGUUCUAGAUGCUGUCCG	9	AAGUUCUAGAUGCUGUCCGNN	289
as	148	CGGACAGCAUCUAGAACUU	10	CGGACAGCAUCUAGAACUUNN	290
s	132	GUUCUAGAUGCUGUCCGAG	11	GUUCUAGAUGCUGUCCGAGNN	291
as	150	CUCGGACAGCAUCUAGAAC	12	CUCGGACAGCAUCUAGAACNN	292
s	135	CUAGAUGCUGUCCGAGGCA	13	CUAGAUGCUGUCCGAGGCANN	293
as	153	UGCCUCGGACAGCAUCUAG	14	UGCCUCGGACAGCAUCUAGNN	294
s	138	GAUGCUGUCCGAGGCAGUC	15	GAUGCUGUCCGAGGCAGUCNN	295
as	156	GACUGCCUCGGACAGCAUC	16	GACUGCCUCGGACAGCAUCNN	296
s	14	CAUUCUUGGCAGGAUGGCU	17	CAUUCUUGGCAGGAUGGCUNN	297
as	32	AGCCAUCCUGCCAAGAAUG	18	AGCCAUCCUGCCAAGAAUGNN	298
s	140	UGCUGUCCGAGGCAGUCCU	19	UGCUGUCCGAGGCAGUCCUNN	299
as	158	AGGACUGCCUCGGACAGCA	20	AGGACUGCCUCGGACAGCANN	300
s	146	CCGAGGCAGUCCUGCCAUC	21	CCGAGGCAGUCCUGCCAUCNN	301
as	164	GAUGGCAGGACUGCCUCGG	22	GAUGGCAGGACUGCCUCGGNN	302
s	152	CAGUCCUGCCAUCAUGUG	23	CAGUCCUGCCAUCAUGUGNN	303
as	170	CACAUUGAUGGCAGGACUG	24	CACAUUGAUGGCAGGACUGNN	304
s	164	CAAUGUGGCCGUGCAUGUG	25	CAAUGUGGCCGUGCAUGUGNN	305
as	182	CACAUGCACGGCCACAUUG	26	CACAUGCACGGCCACAUUGNN	306
s	178	AUGUGUUCAGAAAGGCUGC	27	AUGUGUUCAGAAAGGCUGCNN	307
as	196	GCAGCCUUUCUGAACACAU	28	GCAGCCUUUCUGAACACAUNN	308
s	2	CAGAAGUCCACUCAUUCUU	29	CAGAAGUCCACUCAUUCUUNN	309
as	20	AAGAAUGAGUGGACUUCUG	30	AAGAAUGAGUGGACUUCUGNN	310
s	21	GGCAGGAUGGCUUCUCAUC	31	GGCAGGAUGGCUUCUCAUCNN	311
as	39	GAUGAGAAGCCAUCCUGCC	32	GAUGAGAAGCCAUCCUGCCNN	312
s	210	GAGCCAUUUGCCUCUGGGA	33	GAGCCAUUUGCCUCUGGGANN	313
as	228	UCCCAGAGGCAAUUGGCUC	34	UCCCAGAGGCAAUUGGCUCNN	314
s	23	CAGGAUGGCUUCUCAUCGU	35	CAGGAUGGCUUCUCAUCGUNN	315
as	41	ACGAUGAGAAGCCAUCCUG	36	ACGAUGAGAAGCCAUCCUGNN	316
s	24	AGGAUGGCUUCUCAUCGUC	37	AGGAUGGCUUCUCAUCGUCNN	317
as	42	GACGAUGAGAAGCCAUCCU	38	GACGAUGAGAAGCCAUCCUNN	318
s	245	AGAGCUGCAUGGGCUCACA	39	AGAGCUGCAUGGGCUCACANN	319

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)					
Strand	Position	Sequence (5' to 3')	SEQ ID NO:	Sequence with 3' dinucleotide overhang (5' to 3')	SEQ ID NO:
as	263	UGUGAGCCCAUGCAGCUCU	40	UGUGAGCCCAUGCAGCUCUNN	320
s	248	GCUGCAUGGGCUCACAACU	41	GCUGCAUGGGCUCACAACUNN	321
as	266	AGUUGUGAGCCCAUGCAGC	42	AGUUGUGAGCCCAUGCAGCNN	322
s	25	GGAUGGCUUCUCAUCGUCU	43	GGAUGGCUUCUCAUCGUCUNN	323
as	43	AGACGAUGAGAAGCCAUCC	44	AGACGAUGAGAAGCCAUCCNN	324
s	251	GCAUGGGCUCACAACUGAG	45	GCAUGGGCUCACAACUGAGNN	325
as	269	CUCAGUUGUGAGCCCAUGC	46	CUCAGUUGUGAGCCCAUGCNN	326
s	253	AUGGGCUCACAACUGAGGA	47	AUGGGCUCACAACUGAGGANN	327
as	271	UCCUCAGUUGUGAGCCCAU	48	UCCUCAGUUGUGAGCCCAUNN	328
s	254	UGGGCUCACAACUGAGGAG	49	UGGGCUCACAACUGAGGANN	329
as	272	CUCCUCAGUUGUGAGCCCA	50	CUCCUCAGUUGUGAGCCCANN	330
s	270	GAGGAAUUUGUAGAAGGGA	51	GAGGAAUUUGUAGAAGGGANN	331
as	288	UCCCUUCUACAAAUCCUC	52	UCCCUUCUACAAAUCCUCNN	332
s	276	UUUGUAGAAGGGAUUAUACA	53	UUUGUAGAAGGGAUUAUACANN	333
as	294	UGUAUAUCCCUUCUACAAA	54	UGUAUAUCCCUUCUACAAANN	334
s	277	UUGUAGAAGGGAUUAUACAA	55	UUGUAGAAGGGAUUAUACAANN	335
as	295	UUGUAUAUCCCUUCUACAA	56	UUGUAUAUCCCUUCUACAANN	336
s	278	UGUAGAAGGGAUUAUACAAA	57	UGUAGAAGGGAUUAUACAAANN	337
as	296	UUUGUAUAUCCCUUCUACA	58	UUUGUAUAUCCCUUCUACANN	338
s	281	AGAAGGGAUUAUACAAAGUG	59	AGAAGGGAUUAUACAAAGUGNN	339
as	299	CACUUUGUAUAUCCCUUCU	60	CACUUUGUAUAUCCCUUCUNN	340
s	295	AAGUGGAAAUAGACACCAA	61	AAGUGGAAAUAGACACCAANN	341
as	313	UUGGUGUCUAUUUCCACUU	62	UUGGUGUCUAUUUCCACUUNN	342
s	299	GGAAAUAGACACCAAUUCU	63	GGAAAUAGACACCAAUUCUNN	343
as	317	AGAUUUGGUGUCUAUUUCC	64	AGAUUUGGUGUCUAUUUCCNN	344
s	300	GAAAUAGACACCAAUUCUU	65	GAAAUAGACACCAAUUCUUNN	345
as	318	AAGAUUUGGUGUCUAUUUC	66	AAGAUUUGGUGUCUAUUUCNN	346
s	303	AUAGACACCAAUUCUACU	67	AUAGACACCAAUUCUACUNN	347
as	321	AGUAAGAUUUGGUGUCUAU	68	AGUAAGAUUUGGUGUCUAUNN	348
s	304	UAGACACCAAUUCUACUG	69	UAGACACCAAUUCUACUGNN	349
as	322	CAGUAAGAUUUGGUGUCUA	70	CAGUAAGAUUUGGUGUCUANN	350
s	305	AGACACCAAUUCUACUGG	71	AGACACCAAUUCUACUGGNN	351
as	323	CCAGUAAGAUUUGGUGUCU	72	CCAGUAAGAUUUGGUGUCUNN	352
s	317	UUACUGGAAGGCACUUGGC	73	UUACUGGAAGGCACUUGGCNN	353
as	335	GCCAAGUGCCUUCAGUAA	74	GCCAAGUGCCUUCAGUAANN	354
s	32	UUCUCAUCGUCUGCUCCUC	75	UUCUCAUCGUCUGCUCCUCNN	355
as	50	GAGGAGCAGACGAUGAGAA	76	GAGGAGCAGACGAUGAGAANN	356
s	322	GGAAGGCACUUGGCAUCUC	77	GGAAGGCACUUGGCAUCUCNN	357
as	340	GAGAUGCCAAGUGCCUUC	78	GAGAUGCCAAGUGCCUUCNN	358
s	326	GGCACUUGGCAUCUCCCCA	79	GGCACUUGGCAUCUCCCCANN	359
as	344	UGGGGAGAUGCCAAGUGCC	80	UGGGGAGAUGCCAAGUGCCNN	360
s	333	GGCAUCUCCCCAUUCCAUG	81	GGCAUCUCCCCAUUCCAUGNN	361
as	351	AUGGAAUGGGGAGAUGCCTT	82	AUGGAAUGGGGAGAUGCCTTNN	362
s	334	GCAUCUCCCCAUUCCAUGA	83	GCAUCUCCCCAUUCCAUGANN	363

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)					
Strand	Position	Sequence (5' to 3')	SEQ ID NO:	Sequence with 3' dinucleotide overhang (5' to 3')	SEQ ID NO:
as	352	UCAUGGAAUGGGGAGAUGC	84	UCAUGGAAUGGGGAGAUGCNN	364
s	335	CAUCUCCCCAUUCCAUGAG	85	CAUCUCCCCAUUCCAUGAGNN	365
as	353	CUCAUGGAAUGGGGAGAUG	86	CUCAUGGAAUGGGGAGAUGNN	366
s	336	AUCUCCCCAUUCCAUGAGC	87	AUCUCCCCAUUCCAUGAGCNN	367
as	354	GCUCAUGGAAUGGGGAGAU	88	GCUCAUGGAAUGGGGAGAUNN	368
s	338	CUCCCCAUUCCAUGAGCAU	89	CUCCCCAUUCCAUGAGCAUNN	369
as	356	AUGCUC AUGGAAUGGGGAG	90	AUGCUC AUGGAAUGGGGAGNN	370
s	341	CCCAUCCAUGAGCAUGCA	91	CCCAUCCAUGAGCAUGCANN	371
as	359	UGCAUGCUC AUGGAAUGGG	92	UGCAUGCUC AUGGAAUGGGNN	372
s	347	CCAUGAGCAUGCAGAGGUG	93	CCAUGAGCAUGCAGAGGUGNN	373
as	365	CACCUCUGCAUGCUC AUGG	94	CACCUCUGCAUGCUC AUGGNN	374
s	352	AGCAUGCAGAGGUGGUAUU	95	AGCAUGCAGAGGUGGUAUUNN	375
as	370	AAUACCACCUCUGCAUGCU	96	AAUACCACCUCUGCAUGCUNN	376
s	354	CAUGCAGAGGUGGUAUUCA	97	CAUGCAGAGGUGGUAUUCANN	377
as	372	UGAAUACCACCUCUGCAUG	98	UGAAUACCACCUCUGCAUGNN	378
s	355	AUGCAGAGGUGGUAUUCAC	99	AUGCAGAGGUGGUAUUCACNN	379
as	373	GUGAAUACCACCUCUGCAU	100	GUGAAUACCACCUCUGCAUNN	380
s	362	GGUGGUAUUCACAGCCAAC	101	GGUGGUAUUCACAGCCAACNN	381
as	380	GUUGGCUGUGAAUACCACC	102	GUUGGCUGUGAAUACCACNN	382
s	363	GUGGUAUUCACAGCCAACG	103	GUGGUAUUCACAGCCAACGNN	383
as	381	CGUUGGCUGUGAAUACCAC	104	CGUUGGCUGUGAAUACCACNN	384
s	364	UGGUAUUCACAGCCAACGA	105	UGGUAUUCACAGCCAACGANN	385
as	382	UCGUUGGCUGUGAAUACCA	106	UCGUUGGCUGUGAAUACCANN	386
s	365	GGUAUUCACAGCCAACGAC	107	GGUAUUCACAGCCAACGACNN	387
as	383	GUCGUUGGCUGUGAAUACC	108	GUCGUUGGCUGUGAAUACNN	388
s	366	GUAUUCACAGCCAACGACU	109	GUAUUCACAGCCAACGACUNN	389
as	384	AGUCGUUGGCUGUGAAUAC	110	AGUCGUUGGCUGUGAAUACNN	390
s	367	UAUUCACAGCCAACGACUC	111	UAUUCACAGCCAACGACUCNN	391
as	385	GAGUCGUUGGCUGUGAAUA	112	GAGUCGUUGGCUGUGAAUANN	392
s	370	UCACAGCCAACGACUCCGG	113	UCACAGCCAACGACUCCGNN	393
as	388	CCGGAGUCGUUGGCUGUGA	114	CCGGAGUCGUUGGCUGUGANN	394
s	390	CCCCGCCGCUACACCAUUG	115	CCCCGCCGCUACACCAUUGNN	395
as	408	CAAUGGUGUAGCGGCGGGG	116	CAAUGGUGUAGCGGCGGGNN	396
s	4	GAAGUCCACUCAUUCUUGG	117	GAAGUCCACUCAUUCUUGGNN	397
as	22	CCAAGAAUGAGUGGACUUC	118	CCAAGAAUGAGUGGACUUCNN	398
s	412	CCCUGCUGAGCCCCUACUC	119	CCCUGCUGAGCCCCUACUCNN	399
as	430	GAGUAGGGGCUCAGCAGGG	120	GAGUAGGGGCUCAGCAGGNN	400
s	417	CUGAGCCCCUACUCCUAUU	121	CUGAGCCCCUACUCCUAUUNN	401
as	435	AAUAGGAGUAGGGGCUCAG	122	AAUAGGAGUAGGGGCUCAGNN	402
s	418	UGAGCCCCUACUCCUAUUC	123	UGAGCCCCUACUCCUAUUCNN	403
as	436	GAAUAGGAGUAGGGGCUCA	124	GAAUAGGAGUAGGGGCUCANN	404
s	422	CCCCUACUCCUAUUCACC	125	CCCCUACUCCUAUUCACCNN	405
as	440	GGUGGAAUAGGAGUAGGGG	126	GGUGGAAUAGGAGUAGGGGNN	406
s	425	CUACUCCUAUUCACCACG	127	CUACUCCUAUUCACCACGNN	407
as	443	CGUGGUGGAAUAGGAGUAG	128	CGUGGUGGAAUAGGAGUAGNN	408

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)					
Strand	Position	Sequence (5' to 3')	SEQ ID NO:	Sequence with 3' dinucleotide overhang (5' to 3')	SEQ ID NO:
s	426	UACUCCUAUUCCACCACGG	129	UACUCCUAUUCCACCACGGNN	409
as	444	CCGUGGUGGAAUAGGAGUA	130	CCGUGGUGGAAUAGGAGUANN	410
s	427	ACUCCUAUUCCACCACGGC	131	ACUCCUAUUCCACCACGGCNN	411
as	445	GCCGUGGUGGAAUAGGAGU	132	GCCGUGGUGGAAUAGGAGUNN	412
s	429	UCCUAUUCCACCACGGCUG	133	UCCUAUUCCACCACGGCUGNN	413
as	447	CAGCCGUGGUGGAAUAGGA	134	CAGCCGUGGUGGAAUAGGANN	414
s	432	UAUUCCACCACGGCUGUCG	135	UAUUCCACCACGGCUGUCGNN	415
as	450	CGACAGCCGUGGUGGAAUA	136	CGACAGCCGUGGUGGAAUANN	416
s	433	AUUCCACCACGGCUGUCGU	137	AUUCCACCACGGCUGUCGUNN	417
as	451	ACGACAGCCGUGGUGGAAU	138	ACGACAGCCGUGGUGGAAUNN	418
s	437	CACCACGGCUGUCGUCACC	139	CACCACGGCUGUCGUCACCNN	419
as	455	GGUGACGACAGCCGUGGUG	140	GGUGACGACAGCCGUGGUGNN	420
s	438	ACCACGGCUGUCGUCACCA	141	ACCACGGCUGUCGUCACCANN	421
as	456	UGGUGACGACAGCCGUGGU	142	UGGUGACGACAGCCGUGGUNN	422
s	439	CCACGGCUGUCGUCACCAA	143	CCACGGCUGUCGUCACCAANN	423
as	457	UUGGUGACGACAGCCGUGG	144	UUGGUGACGACAGCCGUGGNN	424
s	441	ACGGCUGUCGUCACCAAUC	145	ACGGCUGUCGUCACCAAUCNN	425
as	459	GAUUGGUGACGACAGCCGU	146	GAUUGGUGACGACAGCCGUNN	426
s	442	CGGCUGUCGUCACCAAUCC	147	CGGCUGUCGUCACCAAUCCNN	427
as	460	GGAUUGGUGACGACAGCCG	148	GGAUUGGUGACGACAGCCGNN	428
s	449	CGUCACCAAUCCCAAGGAA	149	CGUCACCAAUCCCAAGGAANN	429
as	467	UUCCUUGGGAUUGGUGACG	150	UUCCUUGGGAUUGGUGACGNN	430
s	455	CAAUCCCAAGGAAUGAGGG	151	CAAUCCCAAGGAAUGAGGGNN	431
as	473	CCCUCAUUCUUGGGAUUG	152	CCCUCAUUCUUGGGAUUGNN	432
s	491	CCUGAAGGACGAGGGAUGG	153	CCUGAAGGACGAGGGAUGGNN	433
as	509	CCAUCCCUUGUCCUUCAGG	154	CCAUCCCUUGUCCUUCAGGNN	434
s	497	GGACGAGGGAUGGGAUUUC	155	GGACGAGGGAUGGGAUUUCNN	435
as	515	GAAAUCCCAUCCUUCGUCC	156	GAAAUCCCAUCCUUCGUCCNN	436
s	5	AAGUCCACUCAUUCUUGGC	157	AAGUCCACUCAUUCUUGGCNN	437
as	23	GCCAAGAAUGAGUGGACU	158	GCCAAGAAUGAGUGGACUUNN	438
s	508	GGGAUUUCAUGUAACCAAG	159	GGGAUUUCAUGUAACCAAGNN	439
as	526	CUUGGUUACAUGAAAUCCC	160	CUUGGUUACAUGAAAUCCCNN	440
s	509	GGAUUUCAUGUAACCAAGA	161	GGAUUUCAUGUAACCAAGANN	441
as	527	UCUUGGUUACAUGAAAUCC	162	UCUUGGUUACAUGAAAUCCNN	442
s	514	UCAUGUAACCAAGAGUAUU	163	UCAUGUAACCAAGAGUAUUNN	443
as	532	AAUACUCUUGGUUACAUGA	164	AAUACUCUUGGUUACAUGANN	444
s	516	AUGUAACCAAGAGUAUUC	165	AUGUAACCAAGAGUAUUCNN	445
as	534	GGAAUACUCUUGGUUACA	166	GGAAUACUCUUGGUUACAUNN	446
s	517	UGUAACCAAGAGUAUUC	167	UGUAACCAAGAGUAUUCANN	447
as	535	UGGAUACUCUUGGUUACA	168	UGGAUACUCUUGGUUACANN	448
s	518	GUAACCAAGAGUAUUC	169	GUAACCAAGAGUAUUCANN	449
as	536	AUGGAUACUCUUGGUUAC	170	AUGGAUACUCUUGGUUACNN	450
s	54	UGCCUUGCUGGACUGGUAU	171	UGCCUUGCUGGACUGGUAUNN	451
as	72	AUACCAGUCCAGCAAGGCA	172	AUACCAGUCCAGCAAGGCANN	452

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)					
Strand	Position	Sequence (5' to 3')	SEQ ID NO:	Sequence with 3' dinucleotide overhang (5' to 3')	SEQ ID NO:
s	543	UAAAGCAGUGUUUUUACCU	173	UAAAGCAGUGUUUUUACCUUNN	453
as	561	AGGUGAAAACACUGCUUUA	174	AGGUGAAAACACUGCUUUANN	454
s	55	GCCUUGCUGGACUGGUAU	175	GCCUUGCUGGACUGGUAUUNN	455
as	73	AAUACCAGUCCAGCAAGGC	176	AAUACCAGUCCAGCAAGGCNN	456
s	551	UGUUUUCACCUCAUAUGCU	177	UGUUUUCACCUCAUAUGCUNN	457
as	569	AGCAUAUGAGGUGAAAACA	178	AGCAUAUGAGGUGAAAACANN	458
s	552	GUUUUCACCUCAUAUGCUA	179	GUUUUCACCUCAUAUGCUANN	459
as	570	UAGCAUAUGAGGUGAAAAC	180	UAGCAUAUGAGGUGAAAACNN	460
s	553	UUUUCACCUCAUAUGCUAU	181	UUUUCACCUCAUAUGCUAUNN	461
as	571	AUAGCAUAUGAGGUGAAAA	182	AUAGCAUAUGAGGUGAAAAANN	462
s	555	UUCACCUCAUAUGCUAUGU	183	UUCACCUCAUAUGCUAUGUNN	463
as	573	ACAUAGCAUAUGAGGUGAA	184	ACAUAGCAUAUGAGGUGAANN	464
s	557	CACCUCAUAUGCUAUGUUA	185	CACCUCAUAUGCUAUGUUANN	465
as	575	UAACAUAGCAUAUGAGGUG	186	UAACAUAGCAUAUGAGGUGNN	466
s	56	CCUUGCUGGACUGGUAUUU	187	CCUUGCUGGACUGGUAUUUNN	467
as	74	AAUACCAGUCCAGCAAGG	188	AAUACCAGUCCAGCAAGGNN	468
s	563	AUAUGCUAUGUUAGAAGUC	189	AUAUGCUAUGUUAGAAGUCNN	469
as	581	GACUUCUAACAUAGCAUAU	190	GACUUCUAACAUAGCAUAUNN	470
s	564	UAUGCUAUGUUAGAAGUCC	191	UAUGCUAUGUUAGAAGUCCNN	471
as	582	GGACUUCUAACAUAGCAUA	192	GGACUUCUAACAUAGCAUANN	472
s	566	UGCUAUGUUAGAAGUCCAG	193	UGCUAUGUUAGAAGUCCAGNN	473
as	584	CUGGACUUCUAACAUAGCA	194	CUGGACUUCUAACAUAGCANN	474
s	57	CUUGCUGGACUGGUAUUUG	195	CUUGCUGGACUGGUAUUUGNN	475
as	75	CAAUACCAGUCCAGCAAG	196	CAAUACCAGUCCAGCAAGNN	476
s	578	AGUCCAGGCAGAGACAAUA	197	AGUCCAGGCAGAGACAAUANN	477
as	596	AUUGUCUCUGCCUGGACUTT	198	AUUGUCUCUGCCUGGACUTTN	478
s	580	UCCAGGCAGAGACAAUAAA	199	UCCAGGCAGAGACAAUAAANN	479
as	598	UUUAAUUGUCUCUGCCUGGA	200	UUUAAUUGUCUCUGCCUGGANN	480
s	607	GUGAAAGGCACUUUUCAUU	201	GUGAAAGGCACUUUUCAUUNN	481
as	625	AAUGAAAAGUGCCUUUCAC	202	AAUGAAAAGUGCCUUUCACNN	482
s	62	UGGACUGGUAUUUGUGUCU	203	UGGACUGGUAUUUGUGUCUNN	483
as	80	AGACACAAAUACCAGUCCA	204	AGACACAAAUACCAGUCCANN	484
s	77	GUCUGAGGCUGGCCUACG	205	GUCUGAGGCUGGCCUACGNN	485
as	95	CGUAGGGCCAGCCUCAGAC	206	CGUAGGGCCAGCCUCAGACNN	486
s	79	CUGAGGCUGGCCUACGGG	207	CUGAGGCUGGCCUACGGGNN	487
as	97	CCCGUAGGGCCAGCCUCAG	208	CCCGUAGGGCCAGCCUCAGNN	488
s	81	GAGGCUGGCCUACGGGCA	209	GAGGCUGGCCUACGGGCANN	489
as	99	UGCCCGUAGGGCCAGCCUC	210	UGCCCGUAGGGCCAGCCUCNN	490
s	82	AGGCUGGCCUACGGGCAC	211	AGGCUGGCCUACGGGCACNN	491
as	100	GUGCCCGUAGGGCCAGCCU	212	GUGCCCGUAGGGCCAGCCUNN	492
s	84	GCUGGCCCUACGGGCACCG	213	GCUGGCCCUACGGGCACCGNN	493
as	102	CGGUGCCCGUAGGGCCAGC	214	CGGUGCCCGUAGGGCCAGCNN	494
s	85	CUGGCCCUACGGGCACCGG	215	CUGGCCCUACGGGCACCGGNN	495
as	103	CCGGUGCCCGUAGGGCCAG	216	CCGGUGCCCGUAGGGCCAGNN	496
s	87	GGCCCUACGGGCACCGGUG	217	GGCCCUACGGGCACCGGUGNN	497

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)					
Strand	Position	Sequence (5' to 3')	SEQ ID NO:	Sequence with 3' dinucleotide overhang (5' to 3')	SEQ ID NO:
as	105	CACCGGUGCCCGUAGGGCC	218	CACCGGUGCCCGUAGGGCCNN	498
s	9	CCACUCAUUCUUGGCAGGA	219	CCACUCAUUCUUGGCAGGANN	499
as	27	UCCUGCCAAGAAUGAGUGG	220	UCCUGCCAAGAAUGAGUGGNN	500
s	90	CCUACGGGCACCGGUGAAU	221	CCUACGGGCACCGGUGAAUNN	501
as	108	AUUCACCGGUGCCCGUAGG	222	AUUCACCGGUGCCCGUAGGNN	502
s	91	CUACGGGCACCGGUGAAUC	223	CUACGGGCACCGGUGAAUCNN	503
as	109	GAUUCACCGGUGCCCGUAG	224	GAUUCACCGGUGCCCGUAGNN	504
s	92	UACGGGCACCGGUGAAUCC	225	UACGGGCACCGGUGAAUCCNN	505
as	110	GGAUUCACCGGUGCCCGUA	226	GGAUUCACCGGUGCCCGUANN	506
s	93	ACGGGCACCGGUGAAUCCA	227	ACGGGCACCGGUGAAUCCANN	507
as	111	UGGAUUCACCGGUGCCCGU	228	UGGAUUCACCGGUGCCCGUNN	508
s	97	GCACCGGUGAAUCCAAGUG	229	GCACCGGUGAAUCCAAGUGNN	509
as	115	CACUUGGAUUCACCGGUGC	230	CACUUGGAUUCACCGGUGCNN	510
s	98	CACCGGUGAAUCCAAGUGU	231	CACCGGUGAAUCCAAGUGUNN	511
as	116	ACACUUGGAUUCACCGGUG	232	ACACUUGGAUUCACCGGUGNN	512
s	167	UGUGGCCAUGCAUGUGUUC	233	UGUGGCCAUGCAUGUGUUCNN	513
as	185	GAACACAUGCAUGGCCACA	234	GAACACAUGCAUGGCCACANN	514
s	168	GUGGCCAUGCAUGUGUUCA	235	GUGGCCAUGCAUGUGUUCANN	515
as	186	UGAACACAUGCAUGGCCAC	236	UGAACACAUGCAUGGCCACNN	516
s	171	GCCAUGCAUGUGUUCAGAA	237	GCCAUGCAUGUGUUCAGAANN	517
as	189	UUCUGAACACAUGCAUGGC	238	UUCUGAACACAUGCAUGGCNN	518
s	432	UAUUCCACCACGGCUGUCA	239	UAUUCCACCACGGCUGUCANN	519
as	449	UGACAGCCGUGGUGGAAUA	240	UGACAGCCGUGGUGGAAUANN	520
s	447	GUCAUCACCAAUCCCAAGG	241	GUCAUCACCAAUCCCAAGGNN	521
as	465	CCUUGGGAUUGGUGAUGAC	242	CCUUGGGAUUGGUGAUGACNN	522
s	115	GUCCUCUGAUGGUCAAAGU	243	GUCCUCUGAUGGUCAAAGUNN	523
as	133	ACUUUGACCAUCAGAGGAC	244	ACUUUGACCAUCAGAGGACNN	524
s	122	GAUGGUCAAAGUUCUAGAU	245	GAUGGUCAAAGUUCUAGAUNN	525
as	140	AUCUAGAACUUUGACCAUC	246	AUCUAGAACUUUGACCAUCNN	526
s	139	AUGCUGUCCGAGGCAGUCC	247	AUGCUGUCCGAGGCAGUCCNN	527
as	157	GGACUGCCUCGGACAGCAU	248	GGACUGCCUCGGACAGCAUNN	528
s	172	CCGUGCAUGUGUUCAGAAA	249	CCGUGCAUGUGUUCAGAAANN	529
as	190	UUUCUGAACACAUGCACGG	250	UUUCUGAACACAUGCACGGNN	530
s	238	AGUCUGGAGAGCUGCAUGG	251	AGUCUGGAGAGCUGCAUGGNN	531
as	256	CCAUGCAGCUCUCCAGACU	252	CCAUGCAGCUCUCCAGACUNN	532
s	252	CAUGGGCUCACAACUGAGG	253	CAUGGGCUCACAACUGAGGNN	533
as	270	CCUCAGUUGUGAGCCCAUG	254	CCUCAGUUGUGAGCCCAUGNN	534
s	33	UCUCAUCGUCUGCUCCUCC	255	UCUCAUCGUCUGCUCCUCCNN	535
as	51	GGAGGAGCAGACGAUGAGA	256	GGAGGAGCAGACGAUGAGANN	536
s	340	CCCCAUUCCAUGAGCAUGC	257	CCCCAUUCCAUGAGCAUGCNN	537
as	358	GCAUGCUC AUGGAAUGGGG	258	GCAUGCUC AUGGAAUGGGGNN	538
s	421	GCCCCUACUCCUAUUCAC	259	GCCCCUACUCCUAUUCACNN	539
as	439	GUGGAAUAGGAGUAGGGGC	260	GUGGAAUAGGAGUAGGGGCNN	540
s	431	CUAUUCCACCACGGCUGUC	261	CUAUUCCACCACGGCUGUCNN	541

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)

Strand	Position	Sequence (5' to 3')	SEQ ID NO:	Sequence with 3' dinucleotide overhang (5' to 3')	SEQ ID NO:
as	449	GACAGCCGUGGUGGAAUAG	262	GACAGCCGUGGUGGAAUAGNN	542
s	440	CACGGCUGUCGUCACCAAU	263	CACGGCUGUCGUCACCAAUNN	543
as	458	AUUGGUGACGACAGCCGUG	264	AUUGGUGACGACAGCCGUGNN	544
s	496	AGGACGAGGGAUGGGAUUU	265	AGGACGAGGGAUGGGAUUUNN	545
as	514	AAAUCCCAUCCUCGUCCU	266	AAAUCCCAUCCUCGUCCUNN	546
s	556	UCACCUCAUAUGCUAUGUU	267	UCACCUCAUAUGCUAUGUUNN	547
as	574	AACAUAGCAUAUGAGGUGA	268	AACAUAGCAUAUGAGGUGANN	548
s	559	CCUCAUAUGCUAUGUUAGA	269	CCUCAUAUGCUAUGUUAGANN	549
as	577	UCUAACAUAGCAUAUGAGG	270	UCUAACAUAGCAUAUGAGGNN	550
s	570	AUGUUAGAAGUCCAGGCAG	271	AUGUUAGAAGUCCAGGCAGNN	551
as	588	CUGCCUGGACUUCUAACAU	272	CUGCCUGGACUUCUAACAUNN	552
s	78	UCUGAGGCUGGCCCUACGG	273	UCUGAGGCUGGCCCUACGGNN	553
as	96	CCGUAGGGCCAGCCUCAGA	274	CCGUAGGGCCAGCCUCAGANN	554
s	87	GGCCCUACGGGCACCGGUG	275	GGCCCUACGGGCACCGGUGNN	555
as	105	CACCGGUGCCCGUAGGGCC	276	CACCGGUGCCCGUAGGGCCNN	556
s	95	GGGCACCGGUGAAUCCAAG	277	GGGCACCGGUGAAUCCAAGNN	557
as	113	CUUGGAUUCACCGGUGCCC	278	CUUGGAUUCACCGGUGCCCN	558
s	167	CCAUGCAUGUGUUCAGAAA	279	CCAUGCAUGUGUUCAGAAANN	559
as	185	UUUCUGAACACAUGCAUGG	280	UUUCUGAACACAUGCAUGGNN	560

Table 3B. Sense and antisense strand sequences of human TTR dsRNAs

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)

Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
s	100	CCGGUGAAUCCAAGUGUCCdTdT	561
as	118	GGACACUUGGAUUCACCGGdTdT	562
s	11	ACUCAUUCUUGGCAGGAUGdTdT	563
as	29	CAUCCUGCCAAGAAUGAGUdTdT	564
s	111	AAGUGUCCUCUGAUGGUCAdTdT	565
as	129	UGACCAUCAGAGGACACUdTdT	566
s	13	UCAUUCUUGGCAGGAUGGCdTdT	567
as	31	GCCAUCCUGCCAAGAAUGAdTdT	568
s	130	AAGUUCUAGAUGCUGUCCGdTdT	569
as	148	CGGACAGCAUCUAGAACUdTdT	570
s	132	GUUCUAGAUGCUGUCCGAGdTdT	571
as	150	CUCGGACAGCAUCUAGAACdTdT	572
s	135	CUAGAUGCUGUCCGAGGCAdTdT	573
as	153	UGCCUCGGACAGCAUCUAGdTdT	574
s	138	GAUGCUGUCCGAGGCAGUCdTdT	575
as	156	GACUGCCUCGGACAGCAUCdTdT	576
s	14	CAUUCUUGGCAGGAUGGCUdTdT	577
as	32	AGCCAUCCUGCCAAGAAUGdTdT	578
s	140	UGCUGUCCGAGGCAGUCCUdTdT	579
as	158	AGGACUGCCUCGGACAGCAdTdT	580
s	146	CCGAGGCAGUCCUGCCAUCdTdT	581
as	164	GAUGGCAGGACUGCCUCGGdTdT	582

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)			
Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
s	152	CAGUCCUGCCAUCAUGUGdTdT	583
as	170	CACAUUGAUGGCAGGACUGdTdT	584
s	164	CAAUGUGGCCGUGCAUGUGdTdT	585
as	182	CACAUGCACGGCCACAUUGdTdT	586
s	178	AUGUGUUCAGAAAGGCUGCdTdT	587
as	196	GCAGCCUUUCUGAACACAUDdTdT	588
s	2	CAGAAGUCCACUCAUUCUUDdTdT	589
as	20	AAGAAUGAGUGGACUUCUGdTdT	590
s	21	GGCAGGAUGGCUUCUCAUCdTdT	591
as	39	GAUGAGAAGCCAUCCUGCCdTdT	592
s	210	GAGCCAUUUGCCUCUGGGAdTdT	593
as	228	UCCCAGAGGCCAAUUGGCUCdTdT	594
s	23	CAGGAUGGCUUCUCAUCGUDdTdT	595
as	41	ACGAUGAGAAGCCAUCCUGdTdT	596
s	24	AGGAUGGCUUCUCAUCGUCdTdT	597
as	42	GACGAUGAGAAGCCAUCCUDdTdT	598
s	245	AGAGCUGCAUGGGCUCACAdTdT	599
as	263	UGUGAGCCCAUGCAGCUCUDdTdT	600
s	248	GCUGCAUGGGCUCACAACUDdTdT	601
as	266	AGUUGUGAGCCCAUGCAGCdTdT	602
s	25	GGAUGGCUUCUCAUCGUCUDdTdT	603
as	43	AGACGAUGAGAAGCCAUCCdTdT	604
s	251	GCAUGGGCUCACAACUGAGdTdT	605
as	269	CUCAGUUGUGAGCCCAUGCdTdT	606
s	253	AUGGGCUCACAACUGAGGAdTdT	607
as	271	UCCUCAGUUGUGAGCCCAUDdTdT	608
s	254	UGGGCUCACAACUGAGGAGdTdT	609
as	272	CUCCUCAGUUGUGAGCCCAdTdT	610
s	270	GAGGAAUUUGUAGAAGGGAdTdT	611
as	288	UCCCUUCUACAAAUUCCUCdTdT	612
s	276	UUUGUAGAAGGGAUAUACAdTdT	613
as	294	UGUAUAUCCCUUCUACAAAdTdT	614
s	277	UUGUAGAAGGGAUAUACAAAdTdT	615
as	295	UUGUAUAUCCCUUCUACAAAdTdT	616
s	278	UGUAGAAGGGAUAUACAAAdTdT	617
as	296	UUUGUAUAUCCCUUCUACAdTdT	618
s	281	AGAAGGGAUAUACAAAGUGdTdT	619
as	299	CACUUUGUAUAUCCCUUCUDdTdT	620
s	295	AAGUGGAAAUAGACACCAAdTdT	621
as	313	UUGGUGUCUAUUUCCACUUDdTdT	622
s	299	GGAAAUAGACACCAAUCUDdTdT	623
as	317	AGAUUUGGUGUCUAUUUCCdTdT	624
s	300	GAAAUAGACACCAAUCUUDdTdT	625
as	318	AAGAUUUGGUGUCUAUUUCdTdT	626
s	303	AUAGACACCAAUCUUAUCUDdTdT	627

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)			
Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
as	321	AGUAAGAUUUGGUGUCUAUdTdT	628
s	304	UAGACACCAAAUCUUACUGdTdT	629
as	322	CAGUAAGAUUUGGUGUCUAdTdT	630
s	305	AGACACCAAAUCUUACUGGdTdT	631
as	323	CCAGUAAGAUUUGGUGUCUdTdT	632
s	317	UUACUGGAAGGCACUUGGCdTdT	633
as	335	GCCAAGUGCCUCCAGUAAdTdT	634
s	32	UUCUCAUCGUCUGCUCCUCdTdT	635
as	50	GAGGAGCAGACGAUGAGAAdTdT	636
s	322	GGAAGGCACUUGGCAUCUCdTdT	637
as	340	GAGAUGCCAAGUGCCUCCdTdT	638
s	326	GGCACUUGGCAUCUCCCCAdTdT	639
as	344	UGGGGAGAUGCCAAGUGCCdTdT	640
s	333	GGCAUCUCCCCAUUCCAUGdTdT	641
as	351	AUGGAAUGGGGAGAUGCCTTdTdT	642
s	334	GCAUCUCCCCAUUCCAUGAdTdT	643
as	352	UCAUGGAAUGGGGAGAUGCdTdT	644
s	335	CAUCUCCCCAUUCCAUGAGdTdT	645
as	353	CUCAUGGAAUGGGGAGAUGdTdT	646
s	336	AUCUCCCCAUUCCAUGAGCdTdT	647
as	354	GCUCAUGGAAUGGGGAGAUdTdT	648
s	338	CUCCCCAUUCCAUGAGCAUdTdT	649
as	356	AUGCUC AUGGAAUGGGGAGdTdT	650
s	341	CCCAUCCAUGAGCAUGCAdTdT	651
as	359	UGCAUGCUC AUGGAAUGGGdTdT	652
s	347	CCAUGAGCAUGCAGAGGUGdTdT	653
as	365	CACCUCUGCAUGCUCAUGGdTdT	654
s	352	AGCAUGCAGAGGUGGUUUdTdT	655
as	370	AAUACCACCUCUGCAUGCAdTdT	656
s	354	CAUGCAGAGGUGGUUUCAdTdT	657
as	372	UGAAUACCACCUCUGCAUGdTdT	658
s	355	AUGCAGAGGUGGUUUACAdTdT	659
as	373	GUGAAUACCACCUCUGCAUdTdT	660
s	362	GGUGGUUUUACAGCCAACdTdT	661
as	380	GUUGGCUGUGAAUACCACCdTdT	662
s	363	GUGGUUUUACAGCCAACGdTdT	663
as	381	CGUUGGCUGUGAAUACCAdTdT	664
s	364	UGGUUUUACAGCCAACGAdTdT	665
as	382	UCGUUGGCUGUGAAUACCAdTdT	666
s	365	GGUUUUACAGCCAACGAdTdT	667
as	383	GUCGUUGGCUGUGAAUACCdTdT	668
s	366	GUUUUACAGCCAACGACUdTdT	669
as	384	AGUCGUUGGCUGUGAAUACdTdT	670
s	367	UAUUUACAGCCAACGACUCdTdT	671

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)			
Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
as	385	GAGUCGUUGGCUGUGAAUAdTdT	672
s	370	UCACAGCCAACGACUCCGGdTdT	673
as	388	CCGGAGUCGUUGGCUGUGAdTdT	674
s	390	CCCCGCCGCUACACCAUUGdTdT	675
as	408	CAAUGGUGUAGCGGGCGGGdTdT	676
s	4	GAAGUCCACUCAUUCUUGdTdT	677
as	22	CCAAGAAUGAGUGGACUUCdTdT	678
s	412	CCCUGCUGAGCCCCUACUCdTdT	679
as	430	GAGUAGGGGCUCAGCAGGGdTdT	680
s	417	CUGAGCCCCUACUCCUAUUDTdT	681
as	435	AAUAGGAGUAGGGGCUCAGdTdT	682
s	418	UGAGCCCCUACUCCUAUUCdTdT	683
as	436	GAAUAGGAGUAGGGGCUCAdTdT	684
s	422	CCCCUACUCCUAUUCCACCDdTdT	685
as	440	GGUGGAAUAGGAGUAGGGGdTdT	686
s	425	CUACUCCUAUUCCACCACGdTdT	687
as	443	CGUGGUGGAAUAGGAGUAGdTdT	688
s	426	UACUCCUAUUCCACCACGGdTdT	689
as	444	CCGUGGUGGAAUAGGAGUAdTdT	690
s	427	ACUCCUAUUCCACCACGGCdTdT	691
as	445	GCCGUGGUGGAAUAGGAGUdTdT	692
s	429	UCCUAUUCCACCACGGCUGdTdT	693
as	447	CAGCCGUGGUGGAAUAGGAdTdT	694
s	432	UAUUCCACCACGGCUGUCGdTdT	695
as	450	CGACAGCCGUGGUGGAAUAdTdT	696
s	433	AUUCCACCACGGCUGUCGUdTdT	697
as	451	ACGACAGCCGUGGUGGAAUdTdT	698
s	437	CACCACGGCUGUCGUCACCDdTdT	699
as	455	GGUGACGACAGCCGUGGUGdTdT	700
s	438	ACCACGGCUGUCGUCACCAdTdT	701
as	456	UGGUGACGACAGCCGUGGUdTdT	702
s	439	CCACGGCUGUCGUCACCAAdTdT	703
as	457	UUGGUGACGACAGCCGUGGdTdT	704
s	441	ACGGCUGUCGUCACCAAUCDdTdT	705
as	459	GAUUGGUGACGACAGCCGUdTdT	706
s	442	CGGCUGUCGUCACCAAUCCdTdT	707
as	460	GGAUUGGUGACGACAGCCGdTdT	708
s	449	CGUCACCAAUCCCAAGGAAdTdT	709
as	467	UUCCUUGGGAUUGGUGACGdTdT	710
s	455	CAAUCCCAAGGAUAGGGdTdT	711
as	473	CCCUCAUUCCUUGGGAUUGdTdT	712
s	491	CCUGAAGGACGAGGGAUGGdTdT	713
as	509	CCAUCCCUUGUCCUUCAGGdTdT	714
s	497	GGACGAGGGAUGGGAUUUCdTdT	715
as	515	GAAAUCCCAUCCCUUGUCCdTdT	716

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)			
Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
s	5	AAGUCCACUCAUUCUUGGCdTdT	717
as	23	GCCAAGAAUGAGUGGACUdTdT	718
s	508	GGGAUUUCAUGUAACCAAGdTdT	719
as	526	CUUGGUUACAUGAAAUCCdTdT	720
s	509	GGAUUUCAUGUAACCAAGdTdT	721
as	527	UCUUGGUUACAUGAAAUCCdTdT	722
s	514	UCAUGUAACCAAGAGUAUdTdT	723
as	532	AAUACUCUUGGUUACAUGdTdT	724
s	516	AUGUAACCAAGAGUAUCCdTdT	725
as	534	GGAAUACUCUUGGUUACAUdTdT	726
s	517	UGUAACCAAGAGUAUCCAdTdT	727
as	535	UGGAUACUCUUGGUUACAdTdT	728
s	518	GUAACCAAGAGUAUCCAUdTdT	729
as	536	AUGGAUACUCUUGGUUACdTdT	730
s	54	UGCCUUGCUGGACUGGUAdTdT	731
as	72	AUACCAGUCCAGCAAGGCAdTdT	732
s	543	UAAAGCAGUGUUUUCACCUdTdT	733
as	561	AGGUGAAAACACUGCUUUAdTdT	734
s	55	GCCUUGCUGGACUGGUAdTdT	735
as	73	AAUACCAGUCCAGCAAGGCdTdT	736
s	551	UGUUUUCACCUCAUAUGCUdTdT	737
as	569	AGCAUAUGAGGUGAAAACAdTdT	738
s	552	GUUUUCACCUCAUAUGCUAdTdT	739
as	570	UAGCAUAUGAGGUGAAAACdTdT	740
s	553	UUUUCACCUCAUAUGCUAdTdT	741
as	571	AUAGCAUAUGAGGUGAAAAdTdT	742
s	555	UUCACCUCAUAUGCUAUGUdTdT	743
as	573	ACAUAGCAUAUGAGGUGAAAdTdT	744
s	557	CACCUCAUAUGCUAUGUAdTdT	745
as	575	UAACAUAGCAUAUGAGGUGdTdT	746
s	56	CCUUGCUGGACUGGUUUUdTdT	747
as	74	AAAUACCAGUCCAGCAAGGdTdT	748
s	563	AUAUGCUAUGUUAGAAGUCdTdT	749
as	581	GACUUCUAACAUAGCAUAdTdT	750
s	564	UAUGCUAUGUUAGAAGUCCdTdT	751
as	582	GGACUUCUAACAUAGCAUAdTdT	752
s	566	UGCUAUGUUAGAAGUCCAGdTdT	753
as	584	CUGGACUUCUAACAUAGCAAdTdT	754
s	57	CUUGCUGGACUGGUUUUGdTdT	755
as	75	CAAAUACCAGUCCAGCAAGdTdT	756
s	578	AGUCCAGGCAGAGACAAUAdTdT	757
as	596	AUUGUCUCUGCCUGGACUdTdT	758
s	580	UCCAGGCAGAGACAAUAAAdTdT	759
as	598	UUUAUUGUCUCUGCCUGGAdTdT	760

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)			
Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
s	607	GUGAAAGGCACUUUUUCAUdTdT	761
as	625	AAUGAAAAGUGCCUUUCACdTdT	762
s	62	UGGACUGGUUUUGUGUCdTdT	763
as	80	AGACACAAAUACCAGUCCAdTdT	764
s	77	GUCUGAGGCUGGCCCCUACGdTdT	765
as	95	CGUAGGGCCAGCCUCAGAdTdT	766
s	79	CUGAGGCUGGCCCCUACGGGdTdT	767
as	97	CCCGUAGGGCCAGCCUCAGdTdT	768
s	81	GAGGCUGGCCCCUACGGGCAdTdT	769
as	99	UGCCCGUAGGGCCAGCCUCdTdT	770
s	82	AGGCUGGCCCCUACGGGCAdTdT	771
as	100	GUGCCCGUAGGGCCAGCCUdTdT	772
s	84	GCUGGCCCCUACGGGCACCGdTdT	773
as	102	CGGUGCCCGUAGGGCCAGCdTdT	774
s	85	CUGGCCCCUACGGGCACCGdTdT	775
as	103	CCGGUGCCCGUAGGGCCAGdTdT	776
s	87	GGCCCUACGGGCACCGGUGdTdT	777
as	105	CACCGGUGCCCGUAGGGCCdTdT	778
s	9	CCACUCAUUCUUGGCAGGAdTdT	779
as	27	UCCUGCCAAGAAUGAGUGGdTdT	780
s	90	CCUACGGGCACCGGUGAAUdTdT	781
as	108	AUUCACCGGUGCCCGUAGGdTdT	782
s	91	CUACGGGCACCGGUGAAUCdTdT	783
as	109	GAUUCACCGGUGCCCGUAGdTdT	784
s	92	UACGGGCACCGGUGAAUCCdTdT	785
as	110	GGAUUCACCGGUGCCCGUAdTdT	786
s	93	ACGGGCACCGGUGAAUCCAdTdT	787
as	111	UGGAUUCACCGGUGCCCGUdTdT	788
s	97	GCACCGGUGAAUCCAAGUGdTdT	789
as	115	CACUUGGAUUCACCGGUGCdTdT	790
s	98	CACCGGUGAAUCCAAGUGUdTdT	791
as	116	ACACUUGGAUUCACCGGUGdTdT	792
s	167	UGUGGCAUGCAUGUGUUCdTdT	793
as	185	GAACACAUGCAUGGCCACAdTdT	794
s	168	GUGGCAUGCAUGUGUUCAdTdT	795
as	186	UGAACACAUGCAUGGCCACdTdT	796
s	171	GCCAUGCAUGUGUUCAGAAdTdT	797
as	189	UUCUGAACACAUGCAUGGCdTdT	798
s	432	UAUUCCACCACGGCUGUCAdTdT	799
as	449	UGACAGCCGUGGUGGAAUAdTdT	800
s	447	GUCAUCACCAAUCCCAAGGdTdT	801
as	465	CCUUGGGAUUGGUGAUGACdTdT	802
s	115	GUCCUCUGAUGGUCAAAGUdTdT	803
as	133	ACUUUGACCAUCAGAGGACdTdT	804
s	122	GAUGGUCAAAGUUCUAGAUdTdT	805

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)			
Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
as	140	AUCUAGAACUUUGACCAUCdTdT	806
s	139	AUGCUGUCCGAGGCAGUCCdTdT	807
as	157	GGACUGCCUCGGACAGCAUdTdT	808
s	172	CCGUGCAUGUGUUCAGAAAdTdT	809
as	190	UUUCUGAACACAUGCACGGdTdT	810
s	238	AGUCUGGAGAGCUGCAUGGdTdT	811
as	256	CCAUGCAGCUCUCCAGACUdTdT	812
s	252	CAUGGGCUCACAACUGAGGdTdT	813
as	270	CCUCAGUUGUGAGCCCAUGdTdT	814
s	33	UCUCAUCGUCUGCUCCUCCdTdT	815
as	51	GGAGGAGCAGACGAUGAGAdTdT	816
s	340	CCCCAUUCCAUGAGCAUGCdTdT	817
as	358	GCAUGCUC AUGGAAUGGGGdTdT	818
s	421	GCCCCUACUCCUAUUCCACdTdT	819
as	439	GUGGAAUAGGAGUAGGGGCdTdT	820
s	431	CUAUUCCACCACGGCUGUCdTdT	821
as	449	GACAGCCGUGGUGGAAUAGdTdT	822
s	440	CACGGCUGUCGUCACCAAUdTdT	823
as	458	AUUGGUGACGACAGCCGUGdTdT	824
s	496	AGGACGAGGGAUGGGAUUUdTdT	825
as	514	AAAUCCCAUCCUCGUCCUdTdT	826
s	556	UCACCUCAUAUGCUAUGUUdTdT	827
as	574	AACAUAGCAUAUGAGGUGAdTdT	828
s	559	CCUCAUAUGCUAUGUUAGAdTdT	829
as	577	UCUAACAUAGCAUAUGAGGdTdT	830
s	570	AUGUUAGAAGUCCAGGCAGdTdT	831
as	588	CUGCCUGGACUUCUAACAAdTdT	832
s	78	UCUGAGGCUGGCCCCUACGGdTdT	833
as	96	CCGUAGGGCCAGCCUCAGAdTdT	834
s	87	GGCCCUACGGGCACCGGUGdTdT	835
as	105	CACCGGUGCCCGUAGGGCCdTdT	836
s	95	GGGCACCGGUGAAUCCAAGdTdT	837
as	113	CUUGGAUUCACCGGUGCCCDdTdT	838
s	167	CCAUGCAUGUGUUCAGAAAAdTdT	839
as	185	UUUCUGAACACAUGCAUGGdTdT	840

Table 4. Chemically modified sense and antisense strand sequences of human TTR

dsRNAs

See Table 2 for duplex #. Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.2, SEQ ID NO:1329)

Strand	Oligo #	Position	Sequence (5' to 3')	SEQ ID NO:
s	A-32153	100	ACCGUGAUAUCCAAUUGGdTdT	841
as	A-32154	118	GGACACUUGGAUUCACCGdTdT	842
s	A-32155	11	ACUUAUUCUUUCACGAGdTdT	843
as	A-32156	29	CAUCCUGCAAGAAUGAGdTdT	844
s	A-32157	111	AAGUUCUUCUGAUGGUCdTdT	845
as	A-32158	129	UGACUAUACAGGACACUdTdT	846
s	A-32163	13	UCAUUCCUUGCAAGGUGGdTdT	847
as	A-32164	31	GCACUCCUGCCAAAGAdTdT	848

s	A-32165	150	AAATGCTGAGAGGAGGAGGCTGCTG	849
as	A-32166	148	CGGAGAGCAGCAGGAGAGCTGdT	850
s	A-32167	132	GATTCACAGCGGCGCCAGdTdT	851
as	A-32168	150	CUCGGAAGAGAGGAGAGCTdT	852
s	A-32169	135	CGCAGGCGGCGGCGGAGdTdT	853
as	A-32170	153	UCCGCGGAGAGAGAGAGdTdT	854
s	A-32171	138	GAGGCGGCGGAGGAGAGdTdT	855
as	A-32172	156	GACGCGCGCGGAGAGAGdTdT	856
s	A-32175	14	CAAGGCGGCGGAGAGAGdTdT	857
as	A-32176	32	AGGCGGCGGCGGAGAGdTdT	858
s	A-32177	140	AGGCGGCGGCGGAGAGdTdT	859
as	A-32178	158	AGGCGGCGGCGGAGAGdTdT	860
s	A-32179	146	AGGCGGCGGCGGAGAGdTdT	861
as	A-32180	164	GAGGCGGCGGCGGAGAGdTdT	862
s	A-32181	152	GAGGCGGCGGCGGAGAGdTdT	863
as	A-32182	170	CAGGCGGCGGCGGAGAGdTdT	864
s	A-32183	164	CAGGCGGCGGCGGAGAGdTdT	865
as	A-32184	182	CAGGCGGCGGCGGAGAGdTdT	866
s	A-32187	178	AGGCGGCGGCGGAGAGdTdT	867
as	A-32188	196	AGGCGGCGGCGGAGAGdTdT	868
s	A-32189	2	CAGGCGGCGGCGGAGAGdTdT	869
as	A-32190	20	AGGCGGCGGCGGAGAGdTdT	870
s	A-32191	21	AGGCGGCGGCGGAGAGdTdT	871
as	A-32192	39	GAGGCGGAGGCGGAGAGdTdT	872
s	A-32193	210	GAGGCGGAGGCGGAGAGdTdT	873
as	A-32194	228	GAGGCGGAGGCGGAGAGdTdT	874
s	A-32195	23	GAGGCGGAGGCGGAGAGdTdT	875
as	A-32196	41	AGGCGGAGGCGGAGAGdTdT	876

Strand	Oligo #	Position	Sequence (5' to 3')	SEQ ID NO:
s	A-32199	24	AGGAuGGcuucucAuGGuCdTdT	877
as	A-32200	42	GACGAUGGAGGcAcAUCUdTdT	878
s	A-32201	245	AGAGcuucGauGGGcuAcAdTdT	879
s	A-32202	263	UGUGAGCCcAuGcAGCUCUdTdT	880
s	A-32203	248	GcGcAuGGGcuAcAAdTdTdT	881
as	A-32204	266	AGUUGUGAGCCcAuGcAGCcTdT	882
s	A-32205	25	GGUuGGcuucucAuGGuCdTdI	883
as	A-32206	43	AGACGAUGAGAAGCcAUCCdTdT	884
s	A-32207	251	GcAuGGGcuAcAcAcuGAGdTdT	885
as	A-32208	269	CUcAGUUGUGAGCCcAuGcAdTdT	886
s	A-32211	253	uAGGGcuAcAcAcuGAGGAdTdT	887
as	A-32212	271	UCCUcAGUUGUGAGCCcAUdTdT	888
s	A-32213	254	uGGGcuAcAcAcuGAGGAGdTdT	889
s	A-32214	272	CUCCUcAGUUGUGAGCCcAdTdT	890
as	A-32215	270	GAGGAuAuAGAGAGGGAdTdT	891
as	A-32216	288	UCCCUUCuAcAAAUUCCUCdTdT	892
s	A-32217	276	uuuGuAGAGGGGAuAuAcAdTdT	893
as	A-32218	294	UGcAuAUCCCUUUCuAcAAAdTdT	894
s	A-32219	277	uuGuAGAGGGGAuAuAcAdTdT	895
s	A-32220	295	UUUGuAuUCCCUUCuAcAdTdT	896
s	A-32221	278	uGuAGAGGGGAuAuAcAAAdTdT	897
as	A-32222	296	UUUGuAuUCCCUUCuAcAdTdT	898
s	A-32223	281	AGAGGGAuAuAcAAAGGdTdT	899
s	A-32224	299	cACUUUGuAuUCCCUUCdTdT	900
as	A-32225	295	AAGuGGAAuAGAcAcAdTdTdT	901
as	A-32226	313	UUGCGUcUuAUUUUcACAUdTdT	902
s	A-32227	299	GGAAuAGAcAccAAAcuCdTdT	903
as	A-32228	317	AGAUUUUGUGUcUuAUUCCdTdT	904
s	A-32229	300	GAAuAuAGAcAcceAAuCuCdTdT	905
as	A-32230	318	AAGAuuUGUGUcUuAUUCCdTT	906
s	A-32231	303	AuAGAcAccAAAUcuuAcuCdT	907
as	A-32232	321	AGuAAGAUUUUGUGUcUAdTdT	908
s	A-32233	304	uAGAcAcAAuucuuAcuGdTdT	909
s	A-32234	322	cAGuAAGAUUUUGUGUcAdTdT	910
s	A-32235	305	AGAcAccAAuucuuAcuGGdTT	911
as	A-32236	323	CcAGuAAGAUUUUGUGUcUdTdT	912
s	A-32237	317	uuAcuGGAGGcAcuuGGcdTdT	913
as	A-32238	335	GACcAUUGCCUUCcAuGdTTdT	914
s	A-32239	32	uuucucAucGucugGcuuccdTdT	915
s	A-32240	50	GAGGAGcAGAGcAGUGAGAdTdT	916
s	A-32241	322	CGAAGGcAcuuCCcAuucdTdT	917
as	A-32242	340	GAGAGGcAGUGGCCUCCdTTdT	918
s	A-32243	326	GGcAcuuGGcAuucucccAcAdTdT	919
as	A-32244	344	UGGGGAGAUUGcAAUGGcCdTdT	920
s	A-32247	333	GGcAuucucccAuuccAcuGdTdT	921
as	A-32248	351	cAUGGAUUGGGGAGAUGCCdTdT	922
s	A-32249	334	GcAuucucccAuuccAcuGAdTdT	923
as	A-32250	352	UcAUUGAAUUGGGGAGAUcAdTdT	924
s	A-32251	335	cAuucuccccAuuccAcuGAGdTT	925
as	A-32252	353	CUcAUUGGAUUGGGGAGUGdTT	926
s	A-32253	336	AuucuccccAuuccAcuAGcdTdT	927
as	A-32254	354	GCUcAUUGGAUUGGGGAGAUdTdT	928
s	A-32255	338	cuuccccAuuccAcuGAGAuTdT	929
as	A-32256	356	AUGCUcAUUGGAUUGGGGAGdTdT	930
s	A-32259	341	cccAuucccAuGAGcAuGcdTdT	931
as	A-32260	359	UGcAUGCUcAUUGGAGUGGGdTT	932
s	A-32261	347	ccAuGAGcAuGcAAGGUGcdTdT	933
as	A-32262	365	cACCCUcUGcAUGCUcAUUGgdTdT	934

Strand	Oligo #	Position	Sequence (5' to 3')	SEQ ID NO:
s	A-32263	352	AGcAuGcAGAGGgUGGuAuNdTdT	935
s	A-32264	370	AuAuAcCACCUGcAuGcUdTdT	936
as	A-32265	354	cAuGcAGAGGUGAuGcAdTdT	937

as	A-32266	372	UGAAuACcACCUCUGcAUGdTdT	938
s	A-32267	355	AuGcAGAGGuGGuAuucAcdTdT	939
as	A-32268	373	GUuGAuACcACCUCUGcAUGdTdT	940
s	A-32269	362	GSuGGuAuucAcAGccAcdTdT	941
as	A-32270	380	GUUGGUGUGAAuACcACCdTdT	942
s	A-32271	363	GuGGuAuucAcAGccAcGdTdT	943
as	A-32272	381	CGUUGGUGUGAAuACcAcdTdT	944
s	A-32273	364	uGGuAuucAcAGccAcGdTdT	945
as	A-32274	382	UCGUUGGUGUGAAuACcAcdTdT	946
s	A-32275	365	GSuAuucAcAGccAcGdTdT	947
as	A-32276	383	GUCGUUGGUGUGAAuACcAcdTdT	948
s	A-32277	366	GuAuucAcAGccAcGdTdT	949
as	A-32278	384	AGUCGUUGGUGUGAAuACdTdT	950
s	A-32279	367	uAuucAcAGccAcGdTdT	951
as	A-32280	385	GAGUCGUUGGUGUGAAuAdTdT	952
s	A-32281	370	uAcAGccAcAGcuccGdTdT	953
as	A-32282	388	CCGGAGUCGUUGGUGUGAdTdT	954
s	A-32283	390	ccccGccGcuAcAccAuugdTdT	955
as	A-32284	408	cAAUGGUGuAGCGGCGGGdTdT	956
s	A-32285	4	GAAGuccAcucAuucuuGGdTdT	957
as	A-32286	22	CcAAGAAUGAGUGGACUUCdTdT	958
s	A-32287	412	cccuGcuAGGcccuAcucdTdT	959
as	A-32288	430	GAGuAGGGGCUcAGcAGGGdTdT	960
s	A-32289	417	cUGAGcccuAcuccuAuudTdT	961
as	A-32290	435	AAuAGGAGuAGGGGCUcAgdTdT	962
s	A-32291	418	uGAGcccuAcuccuAuudTdT	963
as	A-32292	436	GAuAGGAGuAGGGGCUcAdTdT	964
s	A-32295	422	cccuAcuccuAuuccAcdTdT	965
as	A-32296	440	GGUGAAuAGGAGuAGGGdTdT	966
s	A-32297	425	cuAcuccuAuuccAcAcGdTdT	967
as	A-32298	443	CGUGGUGAAuAGGAGuAGdTdT	968
s	A-32299	426	uAcuccuAuuccAcAcGdTdT	969
as	A-32300	444	CCGUGGUGAAuAGGAGuAdTdT	970
s	A-32301	427	AcuccuAuuccAcAcGdTdT	971
as	A-32302	445	GCCGUGGUGAAuAGGAGuAdTdT	972
s	A-32303	429	uccuAuuccAcAcGcGdTdT	973
as	A-32304	447	cAGCCUGGUGAAuAGGAdTdT	974
s	A-32307	432	uAuuccAcAcGcGcGdTdT	975
as	A-32308	450	CGAcAGCCGUGGUGAAuAdTdT	976
s	A-32309	433	AuuccAcAcGcGcGcGdTdT	977
as	A-32310	451	ACGAcAGCCGUGGUGAAuAdTdT	978
s	A-32311	437	cAcAcGcGcGcGcGdTdT	979
as	A-32312	455	GGUGACGAcAGCCGUGGdTdT	980
s	A-32313	438	AcAcGcGcGcGcAcAdTdT	981
as	A-32314	456	UGGUGACGAcAGCCGUGGdTdT	982
s	A-32315	439	ccAcGcGcGcGcAcAdTdT	983
as	A-32316	457	UUGGUGACGAcAGCCGUGGdTdT	984
s	A-32319	441	AcGGGcGcGcGcAcAdTdT	985
as	A-32320	459	GAUUGGUGACGAcAGCCGdTdT	986
s	A-32321	442	cGGGcGcGcGcAcAcAdTdT	987
as	A-32322	460	GGAUUGGUGACGAcAGCCGdTdT	988
s	A-32323	449	cGcAcAcAuuccAAAGAdTdT	989
as	A-32324	467	UUCGUUGGGAUUGGUGACGdTdT	990
s	A-32325	455	cAAuccAAAGAAuAGGGdTdT	991
as	A-32326	473	CCCUcAUUCGUUGGAUUGdTdT	992

Strand	Oligo #	Position	Sequence (5' to 3')	SEQ ID NO:
s	A-32327	491	ccuGAGGAcAGGGAuGGdTdT	993
as	A-32328	509	CcAUCCUCUGUCCUucAGGdTdT	994
s	A-32331	497	GGAcAGGGAuGGGAuucdTdT	995
as	A-32332	515	GAAAUCCcAUCCUCUGUCCdTdT	996
s	A-32333	5	AAGuccAcucAuucuuGGdTdT	997
as	A-32334	23	GCCAAAGAUAGUGGACUdTdT	998
s	A-32335	508	GGGAuucAuGuAAccAAcAdTdT	999
as	A-32336	526	CUUGGUuAcAUGAAUCCdTdT	1000
s	A-32337	509	GGAuucAuGuAAccAAcAdTdT	1001
as	A-32338	527	UCUUGGUuAcAUGAAUCCdTdT	1002
s	A-32339	514	ucAuGuAAccAAcAGuAuudTdT	1003
as	A-32340	532	AAuACUCUUGGUuAcAUGAdTdT	1004
s	A-32341	516	AuGuAAccAAcAGuAuucdTdT	1005
as	A-32342	534	GGAuACUCUUGGUuAcAUGdTdT	1006
s	A-32343	517	uGuAAccAAcAGuAuucAdTdT	1007
as	A-32344	535	UGGAAuACUCUUGGUuAcAdTdT	1008
s	A-32345	518	GuAAccAAcAGuAuucAdTdT	1009
as	A-32346	536	AUGGAuACUCUUGGUuAcAdTdT	1010
s	A-32347	54	uGcuuGcuGGAcuGGuAdTdT	1011
as	A-32348	72	AuACcAGUCcAGcAAGGcAdTdT	1012
s	A-32349	543	uAAAGcAGuGuuuucAccdTdT	1013
as	A-32350	561	AGGUGAAAcACUGCUuAdTdT	1014
s	A-32351	55	GccuuGcuGGAcuGGuAuudTdT	1015
as	A-32352	73	AAuACcAGUCcAGcAAGGcAdTdT	1016
s	A-32353	551	uGuuuucAccucAuAuGcAdTdT	1017
as	A-32354	569	AGcAuAUGAGGUGAAAcAdTdT	1018
s	A-32355	552	GuuuucAccucAuAuGcAdTdT	1019
as	A-32356	570	uAGcAuAUGAGGUGAAAcAdTdT	1020
s	A-32357	553	uuuuAccucAuAuGcAdTdT	1021
as	A-32358	571	AuAGcAuAUGAGGUGAAAcAdTdT	1022
s	A-32359	555	uucAccucAuAuGcAuGdTdT	1023
as	A-32360	573	AcAuAGcAuAUGAGGUGAdTdT	1024
s	A-32363	557	cAccucAuAuGcAuGuuAdTdT	1025
as	A-32364	575	uAAcAuAGcAuAUGAGGUGdTdT	1026
s	A-32367	56	ccuuGcuGGAcuGGuAuudTdT	1027
as	A-32368	74	AAuACcAGUCcAGcAAGGdTdT	1028
s	A-32369	563	AuAuGcuAuGuuAGAAgucdTdT	1029
as	A-32370	581	GACUUCuAAcAuAGcAuAdTdT	1030
s	A-32371	564	uAuGcuAuGuuAGAAgucdTdT	1031

as	A-32372	582	GGACUUCuAAcAuAGcAuAdTdT	1032
s	A-32373	566	uGcuAuGsuAGAAGuccAGdTdT	1033
as	A-32374	584	CUGGACUUCuAAcAuAGcAdTdT	1034
s	A-32375	57	cuuGcuGGAcuGGuAuuuGdTdT	1035
as	A-32376	75	cAAuAuACcAGUCcAGcAGdTdT	1036
s	A-32379	578	AGuccAGGcAGAGAcAAuAdTdT	1037
as	A-32380	596	uAUUCUCUCUGCCUGGACuAdTdT	1038
s	A-32381	580	uccAGGcAGAGAcAAuAAAdTdT	1039
as	A-32382	598	UUuAUUGUCUCUGCCUGGAdTdT	1040
s	A-32383	607	GuGAAAGGcAcuuuucAuudTdT	1041
as	A-32384	625	AAUGAAAGUGCCUUUCACdTdT	1042
s	A-32385	62	uGGAcuGGuAuuuGuGucudTdT	1043
as	A-32386	80	AGAcAcAAuAuACcAGUCcAdTdT	1044
s	A-32387	77	GucuAGGcGuGGccuAcGdTdT	1045
as	A-32388	95	CGuAGGGCcAGCCUcAGAcdTdT	1046
s	A-32391	79	cuGAGGcuGGccuAcGGGdTdT	1047
as	A-32392	97	CCCGuAGGGCcAGCCUcAGdTdT	1048
s	A-32393	81	GAGGcuGGccuAcGGGcAdTdT	1049
as	A-32394	99	UGCCCGuAGGGCcAGCCUCdTdT	1050

Strand	Oligo #	Position	Sequence (5' to 3')	SEQ ID NO:
s	A-32395	82	AGGcuGGccuAcGGGcAdTdT	1051
as	A-32396	100	GUGCCCGuAGGGCcAGCCUdTdT	1052
s	A-32397	84	GcuGGccuAcGGGcAccGdTdT	1053
as	A-32398	102	CGGUGCCCGuAGGGCcAGCdTdT	1054
s	A-32399	85	cuGGccuAcGGGcAccGGdTdT	1055
as	A-32400	103	CCGGUGCCCGuAGGGCcAGdTdT	1056
s	A-32401	87	GGccuAcGGGcAccGGuGdTdT	1057
as	A-32402	105	cACGGUGCCCGuAGGGCcAdTdT	1058
s	A-32403	9	ccAcuAuucuuGGcAGGAdTdT	1059
as	A-32404	27	UCCUGCcAAGAAUGACUGGdTdT	1060
s	A-32405	90	ccuAcGGGcAccGGuAAuAdTdT	1061
as	A-32406	108	AUUCACCGUGGCCCGuAGGdTdT	1062
s	A-32407	91	cuAcGGGcAccGGuGAuAdTdT	1063
as	A-32408	109	GAUUCACCGUGGCCCGuAGdTdT	1064
s	A-32409	92	uAcGGGcAccGGuGAuuccdTdT	1065
as	A-32410	110	GGAUUCACCGUGGCCCGuAdTdT	1066
s	A-32411	93	AcGGGcAccGGuGAuuccAdTdT	1067
as	A-32412	111	UGGAUUCACCGUGGCCCGuAdTdT	1068
s	A-32415	97	GcAcGGGcAAuuccAAGuGdTdT	1069
as	A-32416	115	cACUUGGAUUCACCGUGGcAdTdT	1070
s	A-32417	98	cAccGGuGAuuccAAGuGdTdT	1071
as	A-32418	116	AcACUUGGAUUCACCGUGGdTdT	1072
s	A-32419	167	uGuGGCcAuGcAuGuGuuAdTdT	1073
as	A-32420	185	GAACAcAuGcAuGGCcAcAdTdT	1074
s	A-32421	168	GuGGCcAuGcAuGuGuuAdTdT	1075
as	A-32422	186	UGAAcAcAuGcAuGGCcAcAdTdT	1076
s	A-32423	171	GccAuGcAuGuGuuAcAGAdTdT	1077
as	A-32424	189	UUCUGAAcAcAuGcAuGGCcAdTdT	1078
s	A-32427	432	uAuuccAccAcGGGcGuAdTdT	1079
as	A-32428	449	UGAcAGCCGUGGUGGAuAdTdT	1080
s	A-32429	447	GucAuAcAAuuccAAGGdTdT	1081
as	A-32430	465	CCUUGGGAUUGUGGAUAGAdTdT	1082
s	A-32159	115	GuccuGuAuGGGcAAAGuAdTdT	1083
as	A-32160	133	ACUUUGACcAuGcAGAGGAcAdTdT	1084
s	A-32161	122	GAuGGGcAAAGuucAuAGAdTdT	1085
as	A-32162	140	AUCuAGAAcUUUGAcAuCAdTdT	1086
s	A-32173	139	AuGcuGuuGAGGcAGuccdTdT	1087
as	A-32174	157	GGACUGCCUGGAcAGcAuAdTdT	1088
s	A-32185	172	ccGuScAuGuGuuAcAGAAAdTdT	1089
as	A-32186	190	UUUCUGAAcAcAuGcACGGdTdT	1090
s	A-32197	238	AGucuGGAGAGcuGcAuGGdTdT	1091
as	A-32198	256	CcAuGcAGCUCUCcAGAcAdTdT	1092
s	A-32209	252	cAuGGGcAcAAcAGGdTdT	1093
as	A-32210	270	CCUcAGUUGUGAGCCcAUGdTdT	1094
s	A-32245	33	ucucAuGcuGcuGcuuccdTdT	1095
as	A-32246	51	GGAGGAcAGAcGAUGAGAdTdT	1096
s	A-32257	340	ccccAuuccAuGAGcAuGdTdT	1097
as	A-32258	358	GcAUGCUcAUGGAUUGGGdTdT	1098
s	A-32293	421	GcccuAcuccuAuuccAdTdT	1099
as	A-32294	439	GUGGAuAGGAGuAGGGCAdTdT	1100
s	A-32305	431	cuAuuccAccAcGGGcGuGdTdT	1101
as	A-32306	449	GAcAGCCGUGGUGGAuAGdTdT	1102
s	A-32317	440	cAcGGcGuGucGucAccAAuAdTdT	1103
as	A-32318	458	AUUGGUGACGAcAGCCGUGdTdT	1104
s	A-32329	496	AGGAcGAGGGAUGGGAuudTdT	1105
as	A-32330	514	AAAUCCcAUCCUCGUCCdTdT	1106
s	A-32361	556	ucAccuAuAuGcuAuGuuAdTdT	1107
as	A-32362	574	AAcAuAGcAuAUGAGGUGAdTdT	1108

Strand	Oligo #	Position	Sequence (5' to 3')	SEQ ID NO:
s	A-32365	559	cuuAuAuGcuAuGuuAGAdTdT	1109
as	A-32366	577	UCuAAcAuAGcAuUGAGGdTdT	1110
s	A-32377	570	AuGuuAGAGGcAcAGGcAdTdT	1111
as	A-32378	588	CUGCCUGGAcUUCuAAcAdTdT	1112
s	A-32389	78	uGGAGGcUGGccuAcGGdTdT	1113
as	A-32390	96	CCGAGGGCcAGCCUcAGAdTdT	1114
s	A-32401	87	GcCCuAcGGGcAcGGGdTdT	1115
as	A-32402	105	cACGGGcCCCGuAGGccAdTdT	1116
s	A-32413	95	GGGcAcGGGcAAuuccAGdTdT	1117
as	A-32414	113	CUUGGAUUCACCGGcCCGdTdT	1118
s	A-32425	167	ccAuGcAuGGGcAGAAAdTdT	1119
as	A-32426	185	GUUCUGAAcAcAuGcAUGGdTdT	1120

Table 5: Identification numbers for rat TTR dsRNAs

See Table 7 for sequences.

Duplex #	Sense Oligo #	Antisense Oligo #
AD-18529	A-32745	A-32746
AD-18530	A-32747	A-32748
AD-18531	A-32749	A-32750
AD-18532	A-32751	A-32752
AD-18533	A-32753	A-32754
AD-18534	A-32755	A-32756
AD-18535	A-32757	A-32758
AD-18536	A-32759	A-32760
AD-18537	A-32761	A-32762
AD-18538	A-32763	A-32764
AD-18539	A-32159	A-32160
AD-18540	A-32765	A-32766
AD-18541	A-32767	A-32768
AD-18542	A-32769	A-32770
AD-18543	A-32771	A-32772
AD-18544	A-32773	A-32774
AD-18545	A-32775	A-32776
AD-18546	A-32777	A-32778
AD-18547	A-32779	A-32780
AD-18548	A-32781	A-32782
AD-18549	A-32783	A-32784
AD-18550	A-32785	A-32786
AD-18551	A-32787	A-32788
AD-18552	A-32791	A-32792
AD-18553	A-32793	A-32794
AD-18554	A-32795	A-32796

Table 6A. Sense and antisense strand sequences for rat TTR dsRNAs

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_012681.1, SEQ ID NO:1330)

Strand	Position	Sequence (5' to 3')	SEQ ID NO:	Sequence with 3' dinucleotide overhang (5' to 3')	SEQ ID NO:
s	115	GUCCUCUGAUGGUCAAAGU	1121	GUCCUCUGAUGGUCAAAGUNN	1173
as	133	ACUUUGACCAUCAGAGGAC	1122	ACUUUGACCAUCAGAGGACNN	1174
s	537	UUCUUGCUCUAUAAACCGU	1123	UUCUUGCUCUAUAAACCGUNN	1175
as	555	ACGGUUUAUAGAGCAAGAA	1124	ACGGUUUAUAGAGCAAGAANN	1176
s	543	CUCUAUAAACCGUGUUAGC	1125	CUCUAUAAACCGUGUUAGCNN	1177
as	561	GCUAACACGGUUUAUAGAG	1126	GCUAACACGGUUUAUAGAGNN	1178
s	392	UCGCCACUACACCAUCGCA	1127	UCGCCACUACACCAUCGCANN	1179
as	410	UGCGAUGGUGUAGUGGCGA	1128	UGCGAUGGUGUAGUGGCGANN	1180
s	538	UCUUGCUCUAUAAACCGUG	1129	UCUUGCUCUAUAAACCGUGNN	1181
as	556	CACGGUUUAUAGAGCAAGA	1130	CACGGUUUAUAGAGCAAGANN	1182
s	541	UGCUCUAUAAACCGUGUUA	1131	UGCUCUAUAAACCGUGUUAANN	1183
as	559	UAACACGGUUUAUAGAGCA	1132	UAACACGGUUUAUAGAGCANN	1184
s	532	CAGUGUUCUUGCUCUAUAA	1133	CAGUGUUCUUGCUCUAUAANN	1185
as	550	UUAUAGAGCAAGAACACUG	1134	UUAUAGAGCAAGAACACUGNN	1186
s	542	GCUCUAUAAACCGUGUUAG	1135	GCUCUAUAAACCGUGUUAGNN	1187

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_012681.1, SEQ ID NO:1330)					
Strand	Position	Sequence (5' to 3')	SEQ ID NO:	Sequence with 3' dinucleotide overhang (5' to 3')	SEQ ID NO:
as	560	CUAACACGGUUUAUAGAGC	1136	CUAACACGGUUUAUAGAGCNN	1188
s	134	CCUGGAUGCUGUCCGAGGC	1137	CCUGGAUGCUGUCCGAGGCNN	1189
as	152	GCCUCGGACAGCAUCCAGG	1138	GCCUCGGACAGCAUCCAGGNN	1190
s	119	UCUGAUGGUCAAAGUCCUG	1139	UCUGAUGGUCAAAGUCCUGNN	1191
as	137	CAGGACUUUGACCAUCAGA	1140	CAGGACUUUGACCAUCAGANN	1192
s	241	CUGGAGAGCUGCACGGGCU	1141	CUGGAGAGCUGCACGGGCUNN	1193
as	259	AGCCCGUGCAGCUCUCCAG	1142	AGCCCGUGCAGCUCUCCAGNN	1194
s	544	UCUAUAAACCGUGUUAGCA	1143	UCUAUAAACCGUGUUAGCANN	1195
as	562	UGCUAACACGGUUUAUAGA	1144	UGCUAACACGGUUUAUAGANN	1196
s	530	AACAGUGUUCUUGCUCUAU	1145	AACAGUGUUCUUGCUCUAUNN	1197
as	548	AUAGAGCAAGAACACUGUU	1146	AUAGAGCAAGAACACUGUUNN	1198
s	118	CUCUGAUGGUCAAAGUCCU	1147	CUCUGAUGGUCAAAGUCCUNN	1199
as	136	AGGACUUUGACCAUCAGAG	1148	AGGACUUUGACCAUCAGAGNN	1200
s	140	UGCUGUCCGAGGCAGCCCU	1149	UGCUGUCCGAGGCAGCCUNN	1201
as	158	AGGGCUGCCUCGGACAGCA	1150	AGGGCUGCCUCGGACAGCANN	1202
s	239	GUCUGGAGAGCUGCACGGG	1151	GUCUGGAGAGCUGCACGGGNN	1203
as	257	CCCGUGCAGCUCUCCAGAC	1152	CCCGUGCAGCUCUCCAGACNN	1204
s	531	ACAGUGUUCUUGCUCUAUA	1153	ACAGUGUUCUUGCUCUAUANN	1205
as	549	UAUAGAGCAAGAACACUGU	1154	UAUAGAGCAAGAACACUGUNN	1206
s	117	CCUCUGAUGGUCAAAGUCC	1155	CCUCUGAUGGUCAAAGUCCNN	1207
as	135	GGACUUUGACCAUCAGAGG	1156	GGACUUUGACCAUCAGAGGNN	1208
s	131	AGUCCUGGAUGCUGUCCGA	1157	AGUCCUGGAUGCUGUCCGANN	1209
as	149	UCGGACAGCAUCCAGGACU	1158	UCGGACAGCAUCCAGGACUNN	1210
s	217	UUGCCUCUGGGAAGACCGC	1159	UUGCCUCUGGGAAGACCGCNN	1211
as	235	GCGGUCUUCCAGAGGCAA	1160	GCGGUCUUCCAGAGGCAANN	1212
s	242	UGGAGAGCUGCACGGGCUC	1161	UGGAGAGCUGCACGGGCUCNN	1213
as	260	GAGCCCGUGCAGCUCUCCA	1162	GAGCCCGUGCAGCUCUCCANN	1214
s	244	GAGAGCUGCACGGGCUCAC	1163	GAGAGCUGCACGGGCUCACNN	1215
as	262	GUGAGCCCGUGCAGCUCUC	1164	GUGAGCCCGUGCAGCUCUCNN	1216
s	246	GAGCUGCACGGGCUCACCA	1165	GAGCUGCACGGGCUCACCANN	1217
as	264	UGGUGAGCCCGUGCAGCUC	1166	UGGUGAGCCCGUGCAGCUCNN	1218
s	399	UACACCAUCGCAGCCUGC	1167	UACACCAUCGCAGCCUGCNN	1219
as	417	GCAGGGCUGCGAUGGUGUA	1168	GCAGGGCUGCGAUGGUGUANN	1220
s	132	GUCCUGGAUGCUGUCCGAG	1169	GUCCUGGAUGCUGUCCGAGNN	1221
as	150	CUCGGACAGCAUCCAGGAC	1170	CUCGGACAGCAUCCAGGACNN	1222
s	245	AGAGCUGCACGGGCUCACC	1171	AGAGCUGCACGGGCUCACCNN	1223
as	263	GGUGAGCCCGUGCAGCUCU	1172	GGUGAGCCCGUGCAGCUCUNN	1224

Table 6B. Sense and antisense strand sequences for rat TTR dsRNAs

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_012681.1, SEQ ID NO:1330)			
Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
s	115	GUCCUCUGAUGGUCAAAGUdTdT	1225
as	133	ACUUUGACCAUCAGAGGACdTdT	1226
s	537	UUCUUGCUCUAUAAACCGUdTdT	1227
as	555	ACGGUUUAUAGAGCAAGAAdTdT	1228

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_012681.1, SEQ ID NO:1330)			
Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
s	543	CUCUAUAAACCGUGUUAGCdTdT	1229
as	561	GCUAACACGGUUUUAUAGAGdTdT	1230
s	392	UCGCCACUACACCAUCGCAdTdT	1231
as	410	UGCGAUGGUGUAGUGGCGAdTdT	1232
s	538	UCUUGCUCUAUAAACCGUGdTdT	1233
as	556	CACGGUUUUAUAGAGCAAGAdTdT	1234
s	541	UGCUCUAUAAACCGUGUUAdTdT	1235
as	559	U AACACGGUUUUAUAGAGCAdTdT	1236
s	532	CAGUGUUCUUGCUCUAUAAdTdT	1237
as	550	UUUAUAGAGCAAGAACACUGdTdT	1238
s	542	GCUCUAUAAACCGUGUUAGdTdT	1239
as	560	CUAACACGGUUUUAUAGAGCdTdT	1240
s	134	CCUGGAUGCUGUCCGAGGCdTdT	1241
as	152	GCCUCGGACAGCAUCCAGGdTdT	1242
s	119	UCUGAUGGUCAAAGUCCUGdTdT	1243
as	137	CAGGACUUUGACCAUCAGAdTdT	1244
s	241	CUGGAGAGCUGCACGGGCUdTdT	1245
as	259	AGCCCGUGCAGCUCUCCAGdTdT	1246
s	544	UCUAUAAACCGUGUUAGCAdTdT	1247
as	562	UGCUAACACGGUUUUAUAGAdTdT	1248
s	530	AACAGUGUUCUUGCUCUAUdTdT	1249
as	548	AUAGAGCAAGAACACUGUdTdT	1250
s	118	CUCUGAUGGUCAAAGUCCUdTdT	1251
as	136	AGGACUUUGACCAUCAGAGdTdT	1252
s	140	UGCUGUCCGAGGCAGCCCUdTdT	1253
as	158	AGGGCUGCCUCGGACAGCAdTdT	1254
s	239	GUCUGGAGAGCUGCACGGGdTdT	1255
as	257	CCCGUGCAGCUCUCCAGACdTdT	1256
s	531	ACAGUGUUCUUGCUCUAUAdTdT	1257
as	549	UAUAGAGCAAGAACACUGUdTdT	1258
s	117	CCUCUGAUGGUCAAAGUCCdTdT	1259
as	135	GGACUUUGACCAUCAGAGGdTdT	1260
s	131	AGUCCUGGAUGCUGUCCGAdTdT	1261
as	149	UCGGACAGCAUCCAGGACUdTdT	1262
s	217	UUGCCUCUGGGAAGACCGCdTdT	1263
as	235	GCGGUCUCCAGAGGCAAdTdT	1264
s	242	UGGAGAGCUGCACGGGCUCdTdT	1265
as	260	GAGCCCGUGCAGCUCUCCAdTdT	1266
s	244	GAGAGCUGCACGGGCUCACdTdT	1267
as	262	GUGAGCCCGUGCAGCUCUCdTdT	1268
s	246	GAGCUGCACGGGCUCACCAAdTdT	1269
as	264	UGGUGAGCCCGUGCAGCUCdTdT	1270
s	399	UACACCAUCGCAGCCCUdCdTdT	1271
as	417	GCAGGGCUGCGAUGGUGUAdTdT	1272

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_012681.1, SEQ ID NO:1330)			
Strand	Position	Sequence with 3'deoxythimidine overhang (5' to 3')	SEQ ID NO:
s	132	GUCCUGGAUGCUGUCCGAGdTdT	1273
as	150	CUCGGACAGCAUCCAGGACdTdT	1274
s	245	AGAGCUGCACGGGCUCACCDdTdT	1275
as	263	GGUGAGCCCGUGCAGCUCUdTdT	1276

Table 7. Chemically modified sense and antisense strand sequences for rat TTR dsRNAs

See Table 5 for duplex # (dsRNA name). Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_012681.1, SEQ ID NO:1330)

Strand	Oligo #	Position	Sequence (5' to 3')	SEQ ID NO:
s	A-32159	115	GuccuGUAuGGGucAAAGdTdT	1277
as	A-32160	133	ACUUUGACcAUcAGAGGAcdTdT	1278
s	A-32745	537	uucuuGcucuAuAAAccGdTdT	1279
as	A-32746	555	ACGGUUuAuAGAGcAAGAdTdT	1280
s	A-32747	543	cucuAuAAAccGGuuAGdTdT	1281
as	A-32748	561	GCuAAcACGGUUuAuAGAdTdT	1282
s	A-32749	392	ucGccAcuAcAccAucGcAdTdT	1283
as	A-32750	410	UGCGAUGGUGuACUGGCGAdTdT	1284
s	A-32751	538	ucuuGcucuAuAAAccGugdTdT	1285
as	A-32752	556	cACGGUUuAuAGAGcAAGAdTdT	1286
s	A-32753	541	uGcucuAuAAAccGGuuAdTdT	1287
as	A-32754	559	uAAcACGGUUuAuAGAGcAdTdT	1288
s	A-32755	532	cAGUGuucuuGcucuAuAAdTdT	1289
as	A-32756	550	UuAuAGAGcAAGAAcACUGdTdT	1290
s	A-32757	542	GcucuAuAAAccGGuuAGdTdT	1291
as	A-32758	560	CuAAcACGGUUuAuAGAGcTdT	1292
s	A-32759	134	ccuGGAuGcuGuccGAGGdTdT	1293
as	A-32760	152	GCCUCGGAcAGcAUcAGGdTdT	1294
s	A-32761	119	ucUGAuGGucAAAGuccgdTdT	1295
as	A-32762	137	cAGCACUUUGAcAUcAGAdTdT	1296
s	A-32763	241	cuGCGAGGcGcAcGGGcudTdT	1297
as	A-32764	259	AGCCCGUGcAGCUCUCcAGdTdT	1298
s	A-32765	544	ucuuAuAAAccGGuuAGcAdTdT	1299
as	A-32766	562	UGCuAAcACGGUUuAuAGAdTdT	1300
s	A-32767	530	AAcAGuGuucuuGcucuAuAdTdT	1301
as	A-32768	548	AuAGAGcAAGAAcACUGUdTdT	1302
s	A-32769	118	cucUGAAcGucAAAGuccdTdT	1303
as	A-32770	136	AGGACUUUGAcAUcAGAGdTdT	1304
s	A-32771	140	uGcuGuccGAGGcAGccudTdT	1305
as	A-32772	158	AGGGCUGCCUCGGAcAGcAdTdT	1306
s	A-32773	239	GcucGGAGAcGcGcAcGGGdTdT	1307
as	A-32774	257	CCCGUGcAGCUCUCcAGAcTdT	1308
s	A-32775	531	AcAGuGuucuuGcucuAuAdTdT	1309
as	A-32776	549	GAuAGAGcAAGAAcACUGUdTdT	1310
s	A-32777	117	ccuGcAGGcGucAAAGuccdTdT	1311
as	A-32778	135	GGACUUUGAcAUcAGAGdTdT	1312
s	A-32779	131	AcuGcGGAuGcuGuccGAdTdT	1313
as	A-32780	149	UCGCAcGcAcAGcAGGcUdTdT	1314
s	A-32781	217	AGGcucUGGGAAGAcGdTdT	1315
as	A-32782	235	CCGUCUCCcAGAGGcAdTdT	1316
s	A-32783	242	uGGAGAGcGcAuAGGcGcAdTdT	1317
as	A-32784	260	GAGCCCGUGcAGCUCUCAdTdT	1318

Strand	Oligo #	Position	Sequence (5' to 3')	SEQ ID NO:
s	A-32785	244	GAGAGcUGcAcGGGcucAcdTdT	1319
as	A-32786	262	GUGAGCCCGUGcAGCUCUCdTdT	1320
s	A-32787	246	GAGcUGcAcGGGcucAccAdTdT	1321
as	A-32788	264	UGGUGAGCCCGUGcAGCUCdTdT	1322
s	A-32791	399	uAcAccAucGcAGccuGcdTdT	1323
as	A-32792	417	GcAGGGCUGCGAUGGUGuAdTdT	1324
s	A-32793	132	GuccuGGAuGcuGuccGAdTdT	1325
as	A-32794	150	CUCGGAcAGcAUcAGGAcdTdT	1326
s	A-32795	245	AGAGcUGcAcGGGcucAcAdTdT	1327
as	A-32796	263	GGUGAGCCCGUGcAGCUCUdTdT	1328

Synthesis of TTR Sequences

[0230] TTR sequences were synthesized on MerMade 192 synthesizer at 1μmol scale. For all the sequences in the list, 'endolight' chemistry was applied as detailed below.

- All pyrimidines (cytosine and uridine) in the sense strand were replaced with corresponding 2'-O-Methyl bases (2'-O-Methyl C and 2'-O-Methyl U)

- In the antisense strand, pyrimidines adjacent to (towards 5' position) ribo A nucleoside were replaced with their corresponding 2-O-Methyl nucleosides
- A two base dTdT extension at 3' end of both sense and antisense sequences was introduced
- The sequence file was converted to a text file to make it compatible for loading in the MerMade 192 synthesis software

[0231] The synthesis of TTR sequences used solid supported oligonucleotide synthesis using phosphoramidite chemistry. The synthesis of the above sequences was performed at 1 μ m scale in 96 well plates. The amidite solutions were prepared at 0.1M concentration and ethyl thio tetrazole (0.6M in Acetonitrile) was used as activator.

[0232] The synthesized sequences were cleaved and deprotected in 96 well plates, using methylamine in the first step and triethylamine.3HF in the second step. The crude sequences thus obtained were precipitated using acetone: ethanol mix and the pellet were re-suspended in 0.5M sodium acetate buffer. Samples from each sequence were analyzed by LC-MS and the resulting mass data confirmed the identity of the sequences. A selected set of samples were also analyzed by IEX chromatography.

[0233] The next step in the process was purification. All sequences were purified on an AKTA explorer purification system using Source 15Q column. A single peak corresponding to the full length sequence was collected in the eluent and was subsequently analyzed for purity by ion exchange chromatography.

[0234] The purified sequences were desalted on a Sephadex G25 column using AKTA purifier. The desalted TTR sequences were analyzed for concentration and purity. The single strands were then annealed to form TTR-dsRNA.

Example 2B: *In vitro* screening of TTR siRNAs for mRNA suppression

[0235] Human TTR targeting dsRNAs (Table 2) were assayed for inhibition of endogenous TTR expression in HepG2 and Hep3B cells, using qPCR (real time PCR) and bDNA (branched DNA) assays to quantify TTR mRNA. Rodent TTR targeting dsRNA (Table 5) were synthesized and assayed for inhibition of endogenous TTR expression using bDNA assays in H.4.II.E cells. Results from single dose assays were used to select a subset of TTR dsRNA duplexes for dose response experiments to calculate IC₅₀'s. IC₅₀ results were used to select TTR dsRNAs for further testing.

Cell culture and transfections:

[0236] The hepatocyte cell lines HepG2, Hep3B and H.4.II.E cells (ATCC, Manassas, VA) were grown to near confluence at 37°C in an atmosphere of 5% CO₂ in Dulbecco's modified Eagle's medium (ATCC) supplemented with 10% FBS, streptomycin, and glutamine (ATCC) before being released from the plate by trypsinization. H.4.II.E cells were also grown in Earle's minimal essential medium. Reverse transfection was carried out by adding 5 μ l of Opti-MEM to 5 μ l of siRNA duplexes per well into a 96-well plate along with 10 μ l of Opti-MEM plus 0.2 μ l of Lipofectamine RNAiMax per well (Invitrogen, Carlsbad CA. cat # 13778-150) and incubated at room temperature for 15 minutes. 80 μ l of complete growth media without antibiotics containing 4x10⁴ (HepG2), 2x10⁴ (Hep3B) or 2x10⁴ (H.4.II.E) cells were then added. Cells were incubated for 24 hours prior to RNA purification. Single dose experiments were performed at 10 nM final duplex concentration and dose response experiments were done with 10, 1, 0.5, 0.1, 0.05, 0.01, 0.005, 0.001, 0.0005, 0.0001, 0.00005, 0.00001 nM.

Total RNA isolation using MagMAX-96 Total RNA Isolation Kit (Applied Biosystems, Foster City CA, part #: AM1830):

[0237] Cells were harvested and lysed in 140 μ l of Lysis/Binding Solution then mixed for 1 minute at 850rpm using and Eppendorf Thermomixer (the mixing speed was the same throughout the process). Twenty micro liters of magnetic beads were added into cell-lysate and mixed for 5 minutes. Magnetic beads were captured using magnetic stand and the supernatant was removed without disturbing the beads. After removing supernatant, magnetic beads were washed with Wash Solution 1 (isopropanol added) and mixed for 1 minute. Beads were captured again and supernatant removed. Beads were then washed with 150 μ l Wash Solution 2 (Ethanol added), captured and supernatant was removed. 50 μ l of DNase mixture (MagMax turbo DNase Buffer and Turbo DNase) was then added to the beads and they were mixed for 10 to 15 minutes. After mixing, 100 μ l of RNA Rebinding Solution was added and mixed for 3 minutes. Supernatant was removed and magnetic

beads were washed again with 150µl Wash Solution 2 and mixed for 1 minute and supernatant was removed completely. The magnetic beads were mixed for 2 minutes to dry before RNA it was eluted with 50µl of water.

cDNA synthesis using ABI High capacity cDNA reverse transcription kit (Applied Biosystems, Foster City, CA, Cat #4368813):

[0238] A master mix of 2µl 10X Buffer, 0.8µl 25X dNTPs, 2µl Random primers, 1µl Reverse Transcriptase, 1µl RNase inhibitor and 3.2µl of H₂O per reaction were added into 10µl total RNA. cDNA was generated using a Bio-Rad C-1000 or S-1000 thermal cycler (Hercules, CA) through the following steps: 25°C 10 min, 37°C 120 min, 85°C 5 sec, 4°C hold.

Real time PCR:

[0239] 2µl of cDNA was added to a master mix of 1µl 18S TaqMan Probe (Applied Biosystems Cat # 4319413E), 1µl TTR TaqMan probe (Applied Biosystems cat # HS00174914 M1) and 101µl TaqMan Universal PCR Master Mix (Applied Biosystems Cat #4324018) per well in a MicroAmp Optical 96 well plate (Applied Biosystems cat # 4326659). Real time PCR was done in an ABI 7000 Prism or an ABI 7900HT Real Time PCR system (Applied Biosystems) using the $\Delta\Delta$ Ct(RQ) assay. All reactions were done in triplicate.

[0240] Real time data were analyzed using the $\Delta\Delta$ Ct method and normalized to assays performed from cells transfected with 10nM BlockIT fluorescent Oligo (Invitrogen Cat # 2013) or 10nM AD-1955 (a control duplex that targets the non-mammalian luciferase gene) to calculate fold change.

Branched DNA assays- QuantiGene 1.0 (Panomics, Fremont, CA. cat #: QG0004)- Used to screen rodent specific duplexes

[0241] H.4.ILE cells (ATCC) were transfected with 10 nM siRNA. After removing media, H.4.ILE were lysed in 100ul of Diluted Lysis Mixture (a mixture of 1 volume of Lysis mixture, 2 volume of nuclease-free water and 10ul of Proteinase-K per ml for the final concentration of 20mg/ml) then incubated at 65 °C for 35 minutes. Then, 80µl of Working Probe Set (a mixture of TTR or GAPDH probe) and 20µl of cell-lysate were added into the Capture Plate. Capture Plates were incubated at 53 °C $\pm$ 1 °C overnight (approximately 16-20hrs). Capture Plates were washed 3 times with 1X Wash Buffer (a mixture of nuclease-free water, Buffer Component 1 and Wash Buffer Component 2), then dried by centrifuging for 1 minute at 1000rpm. 100µl of Amplifier Working Reagent was added into the Capture Plate, which was then sealed and incubated for 1 hour at 46°C $\pm$ 1°C. Wash and dry steps were repeated after 1 hour of incubation and 100µl of Label Solution Reagent was added. The plate was then washed, dried and 100µl Substrate (a mixture of Lithium Lauryl Sulfate and Substrate solution) was added. Capture Plates were placed in the incubator for 30 minutes at 46 °C $\pm$ 1 °C. Capture Plates were then removed from the incubator and incubated at room temperature for 30 minutes. Finally, the Capture Plates were read using the Victor Luminometer (Perkin Elmer, Waltham, MA).

Branched DNA assays- QuantiGene 2.0 (Panomics cat #: QS0011): Used to screen all other duplexes

[0242] After a 24 hour incubation at the dose or doses stated, media was removed and cells were lysed in 100ul Lysis Mixture (1 volume lysis mixture, 2 volumes nuclease-free water and 10µl of Proteinase-K/ml for a final concentration of 20mg/ml) then incubated at 65°C for 35 minutes. 20µl Working Probe Set (TTR probe for gene target and GAPDH for endogenous control) and 80µl of cell-lysate were then added to the Capture Plates. Capture Plates were incubated at 55°C $\pm$ 1°C (approx. 16-20hrs). The next day, the Capture Plates were washed 3 times with 1X Wash Buffer (nuclease-free water, Buffer Component 1 and Wash Buffer Component 2), then dried by centrifuging for 1 minute at 240g. 100µl of pre-Amplifier Working Reagent was added to the Capture Plates, which were sealed with aluminum foil and incubated for 1 hour at 55°C $\pm$ 1°C. Following a 1 hour incubation, the wash step was repeated, then 100µl Amplifier Working Reagent was added. After 1 hour, the wash and dry steps were repeated, and 100µl Label Probe was added. Capture plates were incubated 50°C $\pm$ 1°C for 1 hour. The plates were then washed with 1X Wash Buffer and dried, and then 100µl Substrate was added to the Capture Plates. Capture Plates were read using the SpectraMax Luminometer (Molecular Devices, Sunnyvale, CA) following 5 to 15 minutes incubation.

bDNA data analysis:

[0243] bDNA data were analyzed by (i) subtracting the average background from each triplicate sample, (ii) averaging the resultant triplicate GAPDH (control probe) and TTR (experimental probe) values, and then (iii) taking the ratio: (experimental probe-background)/(control probe-background).

Results

[0244] A summary of the single dose and IC50 results for TTR-dsRNAs (TTR siRNAs) are presented below in Table 8. Single dose results are expressed as % TTR mRNA relative to control, assayed in HepG2 cells. IC50s were determined in HepG2 and/or Hep3B cells, as indicated.

Table 8. Single dose and IC50 results of *in vitro* screens of TTR siRNAs

ND: no data; * indicates result that represents average of two experiments.						
Duplex #	Single Dose at 10nM % relative to control		IC50 (nM)			
	HepG2		HepG2		Hep3B	
	qPCR	bDNA	qPCR	bDNA	qPCR	bDNA
AD-18243	50.35	141.53	ND	ND	ND	ND
AD-18244	64.26	158.55	ND	ND	ND	ND
AD-18245	56.89	107.22	ND	ND	ND	ND
AD-18246	10.53	32.51*	0.265	0.086	ND	ND
AD-18247	125.56	69.57	ND	ND	ND	ND
AD-18248	127.78	66.97	ND	ND	ND	ND
AD-18249	48.77	48.76	ND	ND	ND	ND
AD-18250	96.94	86.42	ND	ND	ND	ND
AD-18251	170.41	129.15	ND	ND	ND	ND
AD-18252	73.52	81.90	ND	ND	ND	ND
AD-18253	25.25	61.25	ND	ND	ND	ND
AD-18254	95.13	103.96	ND	ND	ND	ND
And-18255	119.46	ND	ND	ND	ND	ND
AD-18256	42.64	95.67	ND	ND	ND	ND
And-18257	146.25	141.75	ND	ND	ND	ND
AD-18258	10.20	13.41*	0.007	0.005	0.004	0.005
AD-18259	9.30	20.91*	0.102	0.005	ND	ND
AD-18260	125.37	81.36	ND	ND	ND	ND
AD-18261	14.27	19.40*	0.210	ND	ND	ND
AD-18262	84.95	104.05	ND	ND	ND	ND
AD-18263	16.32	23.25*	0.110	ND	ND	ND
AD-18264	104.18	83.69	ND	ND	ND	ND
AD-18265	41.62	64.87	ND	ND	ND	ND
AD-18266	39.98	110.53	ND	ND	ND	ND
AD-18267	149.64	ND	ND	ND	ND	ND
AD-18268	152.93	174.04	ND	ND	ND	ND
AD-18269	37.27	92.28	ND	ND	ND	ND
AD-18270	99.44	164.75	ND	ND	ND	ND
AD-18271	18.89	28.33*	0.503	0.004	ND	ND
AD-18272	128.32	132.58	ND	ND	ND	ND
AD-18273	115.78	201.95	ND	ND	ND	ND

ND: no data; * indicates result that represents average of two experiments.

	Single Dose at 10nM % relative to control		IC50 (nM)			
	HepG2		HepG2		Hep3B	
Duplex #	qPCR	bDNA	qPCR	bDNA	qPCR	bDNA
AD-18274	8.97	20.04*	0.009	0.176	0.036	0.012
AD-18275	4.09	22.25*	0.026	0.118	ND	ND
AD-18276	19.73	45.22*	0.198	0.677	ND	ND
AD-18277	10.55	26.31*	0.121	0.426	ND	ND
AD-18278	108.86	116.26	ND	ND	ND	ND
AD-18279	66.59	ND	ND	ND	ND	ND
AD-18280	103.26	170.52	ND	ND	ND	ND
AD-18281	87.98	123.88	ND	ND	ND	ND
AD-18282	82.47	140.32	ND	ND	ND	ND
AD-18283	106.54	182.78	ND	ND	ND	ND
AD-18284	106.93	151.78	ND	ND	ND	ND
AD-18285	26.58	60.05*	ND	0.089	ND	ND
AD-18286	109.95	173.66	ND	ND	ND	ND
AD-18287	54.23	155.45	ND	ND	ND	ND
AD-18288	73.52	174.09	ND	ND	ND	ND
AD-18289	103.36	174.76	ND	ND	ND	ND
AD-18290	17.06	52.04*	1.253	0.181	ND	ND
AD-18291	7.71	169.29*	1.304	0.019	ND	ND
AD-18292	7.51	210.03*	0.604	0.005	ND	ND
AD-18293	3.61	62.53*	0.078	0.003	ND	ND
AD-18294	111.53	107.56	ND	ND	ND	ND
AD-18295	115.88	105.37	ND	ND	ND	ND
AD-18296	57.03	38.03	ND	ND	ND	ND
AD-18297	87.69	73.87	ND	ND	ND	ND
AD-18298	10.39	7.25*	0.455	0.008	ND	ND
AD-18299	18.79	18.06*	0.895	0.014	ND	ND
AD-18300	108.70	ND	ND	ND	ND	ND
AD-18301	114.22	70.50	ND	ND	ND	ND
AD-18302	116.19	122.40	ND	ND	ND	ND
AD-18303	124.89	ND	ND	ND	ND	ND
AD-18304	132.99	89.54	ND	ND	ND	ND
AD-18305	153.10	ND	ND	ND	ND	ND
AD-18306	159.22	ND	ND	ND	ND	ND
AD-18307	116.83	84.57	ND	ND	ND	ND
AD-18308	156.72	87.80	ND	ND	ND	ND
AD-18309	113.22	101.97	ND	ND	ND	ND
AD-18310	132.33	ND	ND	ND	ND	ND
AD-18311	161.68	92.92	ND	ND	ND	ND
AD-18312	103.01	71.17	ND	ND	ND	ND
AD-18313	120.65	53.26	ND	ND	ND	ND
AD-18314	116.33	ND	ND	ND	ND	ND
AD-18315	115.13	ND	ND	ND	ND	ND
AD-18316	118.73	122.34	ND	ND	ND	ND

ND: no data; * indicates result that represents average of two experiments.						
	Single Dose at 10nM % relative to control		IC50 (nM)			
	HepG2		HepG2		Hep3B	
Duplex #	qPCR	bDNA	qPCR	bDNA	qPCR	bDNA
AD-18317	114.03	121.10	ND	ND	ND	ND
AD-18318	80.85	122.57	ND	ND	ND	ND
AD-18319	119.14	148.87	ND	ND	ND	ND
AD-18320	22.86	55.43*	ND	0.023	0.403	ND
AD-18321	6.44	31.56*	0.001	0.033	ND	ND
AD-18322	54.21	100.46	ND	ND	ND	ND
AD-18323	6.37	28.71*	0.005	0.023	ND	ND
AD-18324	2.53	15.98*	0.002	0.006	0.005	0.014
AD-18325	2.52	11.96*	0.001	0.016	ND	ND
AD-18326	18.34	43.16*	0.025	0.186	ND	ND
AD-18327	18.28	13.90*	0.044	0.215	ND	ND
AD-18328	4.53	26.04*	0.003	0.004	0.006	0.006
AD-18329	96.93	131.54	ND	ND	ND	ND
AD-18330	11.80	45.18*	0.0004	0.010	0.020	ND
AD-18331	117.77	163.07	ND	ND	ND	ND
AD-18332	11.53	35.09*	0.001	0.076	0.065	ND
AD-18333	12.24	46.94*	0.001	0.115	0.075	ND
AD-18334	16.27	55.28*	0.0004	0.181	1.071	ND
AD-18335	53.52	112.80	ND	ND	ND	ND
AD-18336	6.39	33.00*	0.001	0.112	0.081	ND
AD-18337	51.77	105.33	ND	ND	ND	ND
AD-18338	48.21	102.86	ND	ND	ND	ND
AD-18339	6.48	26.56*	0.004	0.002	0.018	0.029
AD-18340	4.53	30.76*	0.002	0.002	ND	ND
AD-18341	31.27	100.41	ND	ND	ND	ND
AD-18342	7.60	42.89*	ND	0.016	0.076	ND
AD-18343	3.42	17.45*	ND	0.001	ND	ND
AD-18344	75.08	134.31	ND	ND	ND	ND
AD-18345	13.62	42.75*	0.002	0.013	ND	ND
AD-18346	59.25	121.10	ND	ND	ND	ND
AD-18347	91.23	139.54	ND	ND	ND	ND
AD-18348	89.95	159.29	ND	ND	ND	ND
AD-18349	108.01	144.96	ND	ND	ND	ND
AD-18350	123.65	125.87	ND	ND	ND	ND
AD-18351	108.36	104.02	ND	ND	ND	ND
AD-18352	87.82	128.72	ND	ND	ND	ND
AD-18353	14.40	65.77	0.012	0.027	ND	ND
AD-18354	99.27	123.53	ND	ND	ND	ND
AD-18355	135.04	150.88	ND	ND	ND	ND
AD-18356	100.76	178.96	ND	ND	ND	ND
AD-18357	125.30	162.85	ND	ND	ND	ND
AD-18358	103.15	136.01	ND	ND	ND	ND
AD-18359	34.74	140.48	ND	ND	ND	ND

ND: no data; * indicates result that represents average of two experiments.						
	Single Dose at 10nM % relative to control		IC50 (nM)			
	HepG2		HepG2		Hep3B	
Duplex #	qPCR	bDNA	qPCR	bDNA	qPCR	bDNA
AD-18360	103.86	146.86	ND	ND	ND	ND
AD-18361	105.74	152.74	ND	ND	ND	ND
AD-18362	106.96	188.22	ND	ND	ND	ND
AD-18363	124.22	58.46	ND	ND	ND	ND
AD-18364	113.75	66.87	ND	ND	ND	ND
AD-18446	29.73	13.30	ND	ND	ND	ND
AD-18447	109.74	53.63	ND	ND	ND	ND
AD-18448	22.96	8.81	ND	ND	ND	ND
AD-18449	112.59	50.11	ND	ND	ND	ND
AD-18450	89.41	34.89	ND	ND	ND	ND
AD-18451	74.35	23.88	ND	ND	ND	ND
AD-18452	125.25	54.86	ND	ND	ND	ND
AD-18453	126.98	56.31	ND	ND	ND	ND
AD-18454	113.88	52.48	ND	ND	ND	ND
AD-18455	163.00	48.89	ND	ND	ND	ND
AD-18456	15.70	10.52	ND	ND	ND	ND
AD-18457	12.86	8.22	ND	ND	ND	ND
AD-18458	13.00	7.00	ND	ND	ND	ND
AD-18459	14.41	10.72	ND	ND	ND	ND
AD-18460	121.16	74.87	ND	ND	ND	ND
AD-18461	100.53	71.87	ND	ND	ND	ND
AD-18462	47.75	29.35	ND	ND	ND	ND
AD-18463	58.98	44.79	ND	ND	ND	ND

[0245] The dose response data used to identify the IC50 for 5 TTR-dsRNAs (AD-18258, AD-18274, AD-18324, AD-18328, and AD-18339), are presented in detail below in Table 9. All 5 siRNAs were determined to have pM IC50s. The IC50 data for dsRNAs in Table 8 is a summary of the data presented in Table 9 below.

Table 9. Dose response data for 5 TTR-dsRNAs

		% inhibition relative to control AD-1955												
Duplex AD-18258		Dose of duplex (nM)												
Cell type	Detection method	10	1	0.5	0.1	0.05	0.01	0.005	0.001	0.0005	0.0001	0.00005	0.00001	IC50 (nM)
HepG2	qPCR	14.4	14.1	16.2	23.9	27.26	40.19	68.46	78.1	74.48	104.37	98.28	113.68	0.007
HepG2	bDNA	14.3	14.5	11.1	12.8	18.82	19.77	51.21	56.03	63.63	58.35	43.64	51.05	0.005
Hep3B	qPCR	11.9	8.62	12.4	16.4	28.35	30.49	58.36	54.57	81.26	89.43	81.85	101.87	0.004
Hep3B	bDNA	7.65	7.5	11.3	12.6	28.85	27.89	64.57	73.48	72.03	91.44	86.71	89.31	0.005
		% inhibition relative to control AD-1955												
Duplex AD-18274		Dose of duplex (nM)												
Cell type	Detection method	10	1	0.5	0.1	0.05	0.01	0.005	0.001	0.0005	0.0001	0.00005	0.00001	IC50 (nM)
HepG2	qPCR	6.68	8.45	11.7	24.2	42.08	49.89	56.95	62.99	64.47	54.92	67.39	72.67	0.009
HepG2	bDNA	27.5	69	25.2	34.2	73.03	103.4	121.57	97.31	154.93	156.7	Nd	152.25	0.176

		% inhibition relative to control AD-1955												
Duplex AD-18274		Dose of duplex (nM)												
Cell type	Detection method	10	1	0.5	0.1	0.05	0.01	0.005	0.001	0.0005	0.0001	0.00005	0.00001	IC50 (nM)
Hep3B	qPCR	7.58	17	15.6	43.9	42.22	60.55	78.8	77.81	79.97	85.84	86.13	83.99	0.036
Hep3B	bDNA	3.77	4.92	7.51	15	35.21	51.66	72.45	70.12	78.31	77.52	90.72	83.01	0.012
Duplex AD-18324		Dose of duplex (nM)												
Cell type	Detection method	10	1	0.5	0.1	0.05	0.01	0.005	0.001	0.0005	0.0001	0.00005	0.00001	IC50 (nM)
HepG2	qPCR	2.07	2.27	2.74	6.36	8.18	15.23	28.82	52.79	90.86	94.72	116.07	98.97	0.002
HepG2	bDNA	14.5	7.88	11.8	15.9	17.2	46.44	40.4	91.86	0	95.57	0	52.15	0.006
Hep3B	qPCR	2.07	3.48	5.76	16.2	18.73	44.54	49.77	68.88	63.48	76.61	74.7	77.83	0.005
Hep3B	bDNA	3.48	3.8	5.15	15.2	30.84	55.36	74.75	99.39	88.89	110.83	96.55	110.26	0.014
		% inhibition relative to control AD-1955												
Duplex AD-18328		Dose of duplex (nM)												
Cell type	Detection method	10	1	0.5	0.1	0.05	0.01	0.005	0.001	0.0005	0.0001	0.00005	0.00001	IC50 (nM)
HepG2	qPCR	5.85	3.97	3.32	5.62	8	16.75	55.01	39.76	122.41	102.37	114.02	124.09	0.003
HepG2	bDNA	12.3	10.7	10.7	11.9	20.06	25	69.52	57.29	112.28	98.14	142.26	148.92	0.004
Hep3B	qPCR	3.17	5.52	11.7	13.8	27.68	39.58	61.21	61.87	90.51	87.56	106.03	108.72	0.006
Hep3B	bDNA	3.08	3.66	4.19	7.25	21.05	22.1	73.74	63.19	105.55	96.27	105.97	96.46	0.006
		% inhibition relative to control AD-1955												
Duplex AD-18339		Dose of duplex (nM)												
Cell type	Detection method	10	1	0.5	0.1	0.05	0.01	0.005	0.001	0.0005	0.0001	0.00005	0.00001	IC50 (nM)
HepG2	qPCR	6.27	7.28	Nd	11	15.25	38.69	38.78	71.7	84.09	62.2	75.61	85.46	0.004
HepG2	bDNA	15.1	8.14	5.13	6.89	12.17	32.14	42.98	64.01	60.76	79.95	81.97	95.43	0.002
Hep3B	qPCR	8.3	9.47	13.2	34.5	44.54	77.38	81.04	81.41	93.95	81.04	75.61	78.28	0.018
Hep3B	bDNA	10.5	9.43	11.7	27.1	44.88	72.32	79.88	79.6	87.46	96.53	95.13	89.88	0.029

[0246] A summary of the single dose results for rodent specific TTR-dsRNAs (TTR siRNAs) are presented below in Table 10. Single dose results are expressed as % TTR mRNA relative to control, assayed in rat H.4.II.E cells, after transfection of rodent specific TTR siRNAs at 10 nM. These results show that some rodent specific TTR siRNAs are effective in suppressing endogenous rat TTR mRNA *in vitro*.

Table 10. Single dose results of *in vitro* screen of rodent specific TTR-dsRNAs (TTR siRNAs)

Duplex #	% Relative to control at 10 nM	Duplex #	% Relative to control at 10 nM
AD-18529	19.83	AD-18542	6.3
AD-18530	44.49	AD-18543	16.46
AD-18531	6.01	AD-18544	17.55
AD-18532	24.06	AD-18545	3.53
AD-18533	37.78	AD-18546	2.75
AD-18534	8.19	AD-18547	7.01
AD-18535	10.18	AD-18548	5.02
AD-18536	16.13	AD-18549	1.61

Duplex #	% Relative to control at 10 nM	Duplex #	% Relative to control at 10 nM
AD-18537	15.88	AD-18550	9.58
AD-18538	19.93	AD-18551	7.74
AD-18539	49.24	AD-18552	3.74
AD-18540	2.99	AD-18553	50.39
AD-18541	1.32	AD-18554	111.06

Example 3. *In vitro* assay of TTR siRNAs for induction of TNF- α and IFN- α secretion

[0247] To evaluate potential for immunostimulation, TTR siRNAs were assayed *in vitro* for induction of TNF- α and IFN- α secretion.

[0248] Human PBMC were isolated from freshly collected buffy coats obtained from healthy donors (Research Blood Components, Inc., Boston, MA) by a standard Ficoll-Hypaque density centrifugation. Freshly isolated cells (1×10^5 /well/100 μ l) were seeded in 96-well plates and cultured in RPMI 1640 GlutaMax medium (Invitrogen) supplemented with 10% heat-inactivated fetal bovine serum and 1% antibiotic/antimycotic (Invitrogen).

[0249] siRNAs were transfected into PBMC using DOTAP transfection reagent (Roche Applied Science). The DOTAP was first diluted in Opti-MEM (Invitrogen) for 5 minutes before mixing with an equal volume of Opti-MEM containing the siRNA. siRNA/DOTAP complexes were incubated as specified by the manufacturer's instructions and subsequently added to PBMC (50 μ l/well) which were then cultured for 24 hours. Positive and negative control siRNAs were included in all assays. AD-5048 was used as a positive control siRNA. AD-5048 corresponds to a sequence that targets human Apolipoprotein B (Soutschek *et al.*, 2004) and elicits secretion of both IFN- α and TNF- α in this assay. AD-1955, which does not elicit IFN- α and TNF- α secretion in this assay, was used as a negative control siRNA. All siRNAs were used at a final concentration of 133 nM. The ratio of RNA to transfection reagent was 16.5 pmoles per μ g of DOTAP.

[0250] Cytokines were detected and quantified in culture supernatants with a commercially available ELISA kit for IFN- α (BMS216INST) and TNF- α (BMS223INST), both from Bender MedSystems (Vienna, Austria). TTR siRNA cytokine induction is expressed as percent IFN- α or TNF- α produced relative to the positive control siRNA AD-5048.

[0251] IFN- α and TNF- α stimulation results for a number of TTR siRNAs are presented in FIG. 1 (mean of quadruplicate wells $\pm$ SD) and below in Table 11 (percentage compared with AD-5048). None of the TTR siRNAs evaluated induced significant TNF- α or IFN- α secretion by cultured human PBMCs.

Table 11. IFN- α and TNF- α stimulation results for TTR siRNAs

Duplex #	IFN- α (% of AD-5048)	TNF- α (% of AD-5048)
AD-18246	0	4
AD-18258	0	0
AD-18259	0	0
AD-18261	0	0
AD-18263	0	0
AD-18271	0	0
AD-18274	2	1
AD-18275	0	0
AD-18276	0	0
AD-18277	0	0
AD-18285	0	0
AD-18290	0	0
AD-18291	0	0
AD-18292	0	0
AD-18293	0	0

Duplex #	IFN- α (% of AD-5048)	TNF- α (% of AD-5048)
AD-18298	0	0
AD-18299	0	0
AD-18320	0	0
AD-18321	0	0
AD-18323	0	0
AD-18324	0	0
AD-18325	0	0
AD-18326	0	0
AD-18327	0	0
AD-18328	0	0
AD-18330	0	0
AD-18332	1	0
AD-18333	0	1
AD-18334	0	1
AD-18336	1	0
AD-18339	0	0
AD-18340	0	0
AD-18342	0	0
AD-18343	0	0
AD-18345	0	0
AD-18353	0	0
AD-18448	0	0
AD-18456	0	0
AD-18457	0	0
AD-18458	0	0
AD-18459	0	0

[0252] The five lead TTR targeting dsRNAs (TTR siRNAs) were selected based on IC₅₀s in the pM range in the human hepatocyte cell lines HepG2 and Hep3B, and the absence of immunostimulatory activity. Duplexes without any mismatches are more likely to achieve significant knockdown of the target transcript than duplexes with mismatches between the oligo and the mRNA. To better enable interpretation of cross-species toxicology data and to have the broadest applicability to human patients, duplexes that have 100% identity in orthologous genes from rat, cynomolgus monkey and human, and that do not target regions with known polymorphisms are generally preferred. The five lead compounds were selected based on IC₅₀ in hepatocyte cell lines in the pM range, the absence of immunostimulatory activity, specificity to the human TTR transcripts, and absence of known polymorphisms (mutations) in the region of the mRNA targeted by the duplex. In the case of TTR, no 19 base oligos were found with complete identity in human, rat and cynomolgus monkey. A summary of these data are presented in Table 12, which also includes information on known TTR mutations in the region targeted by the duplex and cross-species reactivity.

Table 12. Summary of data for five most potent TTR dsRNAs.

Duplex #	IC ₅₀ (qPCR): nM HepG2	IC ₅₀ (bDNA): nM HepG2	IFN α /TNF α	Mutations not covered	Cross-species reactivity
AD-18258	0.007	0.005	Negative	None (non-coding region)	Cyno: 1 mismatch @ position 14 A to G Rat: no homology at any position
AD-18274	0.009	0.176	Negative	Lys70Asn; Val71Ala; Ile73Val; Asp74His	Cyno: no mismatch Rat: no homology at any position
AD-18324	0.002	0.006	Negative	None (non-coding)	Cyno: no mismatch

Duplex #	IC50 (qPCR): nM HepG2	IC50 (bDNA): nM HepG2	IFNa/TNFa	Mutations not covered	Cross-species reactivity
				region)	Rat: no homology at any position
AD-18328	0.003	0.004	Negative	None (non-coding region)	Cyno: no mismatch Rat: 7 mismatches
AD-18339	0.004	0.002	Negative	None (non-coding region)	None

Example 4. *In vivo* reduction of liver TTR mRNA and plasma TTR protein by LNP01-18324, LNP01-18328 and LNP01-18246 in transgenic mice

[0253] Two TTR siRNAs, AD-18324 and AD-18328, were chosen for *in vivo* evaluation. These duplexes exhibited potent dose-dependent *silencing in vitro* in hepatocyte cell lines (e.g. HepG2). FIG. 2A and FIG. 2B show the dose responses in HepG2 cells after transfection with AD-18324 (FIG. 2A) or AD-18328 (FIG. 2B) where the doses are expressed in nM on the x-axis and the responses are expressed as fraction TTR mRNA remaining relative to control, on the y-axis. In HepG2 cells, the IC50s of AD-18324 and AD-18328 were determined to be 2 pM and 3 pM, respectively. The TTR target sites for both lead dsRNA candidates are in the 3' untranslated region of the TTR mRNA, in a region where there are no reported mutations in the literature.

[0254] The sequences of each strand of the two lead candidates are reproduced below from the Tables. Strand: s= sense; as= antisense; Position: position of 5' base on transcript NM_000371.2.

Duplex #	Strand	Oligo #	Position*	Sequence 5' to 3'	SEQ ID NO:
AD-18324	s	A-32337	509	GGAuuuAuGuAAccAAGAdTdT	1001
AD-18324	as	A-32338	527	UCUUGGUuAcAUGAAAUCCdTdT	1002
AD-18328	s	A-32345	518	GuAAccAAGAGuAuuccAudTdT	1009
AD-18328	as	A-32346	536	AUGGAuACUCUUGGUuACdTdT	1010

[0255] In addition, a rodent cross-reactive TTR dsRNA, AD-18246, was chosen for further evaluation *in vivo*. AD-18246 targets a sequence beginning at position 88 of the open reading frame, where there are three mutations reported in the literature. A dose response curve for AD-18246 in HepG2 cells is shown in FIG. 3. AD-18246 is substantially less potent than AD-18324 and AD-18328; the IC50 of AD-18246 was determined to be 265 pM.

[0256] AD-18324, AD-18328, and AD-18246 were administered to transgenic mice after formulation in LNP01. 3-5 month old H129-mTTR-KO/iNOS-KO/hTTR transgenic mice (mouse transthyretin knock-out/ inducible nitric oxide synthase knock-out/human transthyretin transgenic) were intravenously (IV) administered 200 µl of LNP01-formulated transthyretin-specific siRNA (AD-18324, AD-18328, or AD-18246), LNP01-formulated control siRNA targeting the non-mammalian luciferase gene (AD-1955) or PBS via the tail vein at concentrations of 1.0 mg/kg, 3.0 mg/kg, or 6.0 mg/kg for siRNAs AD-18324 and AD-18328, 3.0 mg/kg for siRNA AD-18246, and 6.0 mg/kg for siRNA AD-1955. LNP01 is a lipidoid formulation comprised of ND98, Cholesterol, and PEG-Ceramide C16.

[0257] After approximately forty-hours, mice were anesthetized with 200 µl of ketamine, and then exsanguinated by severing the right caudal artery. Whole blood was isolated and plasma was isolated and stored at -80°C until assaying. Liver tissue was collected, flash-frozen and stored at -80°C until processing.

[0258] Efficacy of treatment was evaluated by (i) measurement of TTR mRNA in liver at 48 hours post-dose, and (ii) measurement of TTR protein in plasma at prebleed and at 48 hours post-dose. TTR liver mRNA levels were assayed utilizing the Branched DNA assays-QuantiGene 2.0 (Panomics cat #: QS0011). Briefly, mouse liver samples were ground and tissue lysates were prepared. Liver lysis mixture (a mixture of 1 volume of lysis mixture, 2 volume of nuclease-free water and 10ul of Proteinase-K/ml for a final concentration of 20mg/ml) was incubated at 65 °C for 35 minutes. 20µl of Working Probe Set (TTR probe for gene target and GAPDH for endogenous control) and 80ul of tissue-lysate were then added into the Capture Plate. Capture Plates were incubated at 55 °C ±1 °C (aprx. 16-20hrs). The next day, the Capture Plate were washed 3 times with 1X Wash Buffer (nuclease-free water, Buffer Component 1 and Wash Buffer Component 2), then dried by centrifuging for 1

minute at 240g. 100µl of pre-Amplifier Working Reagent was added into the Capture Plate, which was sealed with aluminum foil and incubated for 1 hour at 55°C ±1°C. Following 1 hour incubation, the wash step was repeated, then 100µl of Amplifier Working Reagent was added. After 1 hour, the wash and dry steps were repeated, and 100µl of Label Probe was added. Capture plates were incubated 50 °C ±1 °C for 1 hour. The plate was then washed with 1X Wash Buffer, dried and 100µl Substrate was added into the Capture Plate. Capture Plates were read using the SpectraMax Luminometer following a 5 to 15 minute incubation. bDNA data were analyzed by subtracting the average background from each triplicate sample, averaging the resultant triplicate GAPDH (control probe) and TTR (experimental probe) values, and then computing the ratio: (experimental probe-background)/(control probe-background).

[0259] TTR plasma levels were assayed utilizing the commercially available kit "AssayMax Human Prealbumin ELISA Kit" (AssayPro, St. Charles, MO, Catalog # EP3010-1) according to manufacturer's guidelines. Briefly, mouse plasma was diluted 1:10,000 in 1X mix diluents and added to pre-coated plates along with kit standards, and incubated for 2 hours at room temperature followed by 5X washes with kit wash buffer. Fifty microliters of biotinylated prealbumin antibody was added to each well and incubated for 1 hr at room temperature, followed by 5X washes with wash buffer. Fifty microliters of streptavidin-peroxidase conjugate was added to each well and plates were incubated for 30 minutes at room temperature followed by washing as previously described. The reaction was developed by the addition of 50 µl/well of chromogen substrate and incubation for 10 minutes at room temperature with stopping of reaction by the addition of 50 µl/well of stop solution. Absorbance at 450 nm was read on a Versamax microplate reader (Molecular Devices, Sunnyvale, CA) and data were analyzed utilizing the Softmax 4.6 software package (Molecular Devices).

[0260] LNP01-18324 and LNP01-18328 were found to reduce liver TTR mRNA (FIG. 4A) and plasma TTR protein (FIG. 4B) levels in a dose-dependent manner with IV bolus administration. The mRNA ED50 of LNP01-18328 was determined to be ~1 mg/kg whereas the ED50 of LNP01-18324 was determined to be ~2 mg/kg. The effects of LNP01-18324 and LNP01-18328 were specific, because the control, LNP01-1955 at 6 mg/kg, did not significantly affect liver TTR mRNA levels, as compared with the PBS group. LNP01-18324 and LNP01-18328 reduced plasma TTR protein levels relative to the PBS group, with potencies that were similar to those on TTR mRNA levels. At 3 mg/kg, LNP01-18246 reduced liver TTR mRNA levels to a lesser extent than 3 mg/kg LNP01-18324 or LNP01-18328.

[0261] These results demonstrate that LNP01-18324 and LNP01-18328, administered by IV bolus, substantially reduce human TTR mRNA expressed by the transgenic mouse liver, which results in reduction of human TTR protein in the circulation.

Example 5. *In vivo* reduction of wild-type TTR mRNA in the non-human primate liver by SNALP-18324 and SNALP-18328

[0262] To evaluate the efficacy of TTR siRNAs AD-18324 and AD-18328 in non-human primates on liver TTR mRNA levels, the siRNAs were formulated in SNALP and administered by 15-minute IV infusion. Cynomolgus monkeys (*Macaca fascicularis*) (2 to 5 kg, 3 animals per group) were administered 15-minute IV infusions of SNALP-18324 (0.3, 1.0 or 3.0 mg/kg), SNALP-18328 (0.3, 1 or 3 mg/kg), or SNALP-1955 (3 mg/kg, with negative control siRNA AD-1955 which targets the non-mammalian gene luciferase). At forty-eight hours post-dosing, monkeys were anesthetized with sodium pentobarbital and exsanguinated. Liver tissue for TTR mRNA determination was collected, flash-frozen, and stored at -80°C until processing.

[0263] TTR mRNA levels in the liver were assayed utilizing a custom designed Branched DNA assay, utilizing the QuantiGene1.0 technology. Briefly, monkey liver samples were ground and tissue lysates were prepared. Liver lysis mixture (1 volume lysis mixture, 2 volume nuclease-free water, and 10µl of Proteinase-K/ml for a final concentration of 20mg/ml) was incubated at 65°C for 35 minutes. 20µl Working Probe Set (TTR probe for gene target and GAPDH for endogenous control) and 80µl tissue-lysate were then added into the Capture Plate. Capture Plates were incubated at 55°C ±1°C (approx. 16-20hrs). The next day, the Capture Plates were washed three times with 1X Wash Buffer (nuclease-free water, Buffer Component 1 and Wash Buffer Component 2), then dried by centrifuging for 1 minute at 240g. 100µl of pre-Amplifier Working Reagent was added into the Capture Plate, which was sealed with aluminum foil and incubated for 1 hour at 55°C ±1°C. Following a 1-hour incubation, the wash step was repeated, and then 100µl Amplifier Working Reagent was added. After 1 hour, the wash and dry steps were repeated, and 100µl Label Probe was added. Capture plates were incubated 50°C ±1°C for 1 hour. The plates were then washed with 1X Wash Buffer and dried, and then 100µl Substrate was added into the Capture Plate. Capture Plates were read using the SpectraMax Luminometer following a 5 to 15 minute incubation. bDNA data were analyzed by (i) subtracting the average background from each triplicate sample, (ii) averaging the resultant GAPDH (control probe) and TTR (experimental probe) values, and then (iii) taking the ratio: (experimental probe-background)/(control probe-background).

[0264] The results are shown in FIG. 5. SNALP -18324 and SNALP -18328 reduced TTR mRNA levels in the liver in a dose-dependent manner, compared to the negative control SNALP-1955. The mRNA ED50s of SNALP-18328 and SNALP-18324 were determined to be ~0.3 and ~ 1 mg/kg, respectively.

[0265] These results demonstrate that SNALP-18324 and SNALP-18328 are effective in suppressing wild-type TTR mRNA in non-human primate liver when administered by IV infusion.

Example 6. *In vivo* reduction of mutant (V30M) TTR mRNA and protein by SNALP-18328 in the transgenic mouse

[0266] To evaluate the efficacy of TTR siRNA AD-18328 on mutant (V30M) TTR mRNA in the liver and mutant (V30M) TTR protein in the serum, AD-18328 was formulated in SNALP and administered by IV bolus to V30M hTTR transgenic mice. 8 to 12-week old V30M hTTR transgenic mice (5 animals/ group) were intravenously (IV) administered 200 μ l SNALP-18328 (0.03, 0.3 or 3 mg/kg), SNALP-1955 (3 mg/kg, with negative control siRNA AD-1955 which targets the non-mammalian gene luciferase), or PBS. Mice used were the Mus musculus strain H129-hTTR KO from Institute of Molecular and Cellular Biology, Porto, Portugal. Briefly, hTTR H129 transgenic mice were crossed with a H129 endogenous TTR KO mice (null mice to generate the H129-hTTR transgenic mice, in a null mouse TTR background (Maeda, S., (2003), Use of genetically altered mice to study the role of serum amyloid P component in amyloid deposition. Amyloid Suppl. 1, 17-20.).

[0267] At 48 hrs post-injection, animals in all five treatment groups were given a lethal dose of ketamine/xylazine. Serum samples were collected and stored at -80°C until analysis. Liver tissue was collected, flash-frozen and stored at -80°C until processing.

[0268] For TTR mRNA quantitation, frozen liver tissue was ground into powder, and lysates were prepared. TTR mRNA levels relative to those of GAPDH mRNA were determined in the lysates by using a branched DNA assay (QuantiGene Reagent System, Panomics, Fremont, CA). Briefly, the QuantiGene assay (Genospectra) was used to quantify mRNA levels in tissue sample lysates according to the manufacturer's instructions. The mean level of TTR mRNA was normalized to the mean level of GAPDH mRNA for each sample. Group means of the normalized values were then further normalized to the mean value for the PBS treated group, to obtain the relative level of TTR mRNA expression.

[0269] For TTR protein quantitation, serum was assayed using the AssayPro (St. Charles, MO) Assaymax PreAlbumin ELISA Kit according to the manufacturer's protocol.

[0270] The results are shown in FIG. 6A and FIG. 6B for liver mRNA and serum protein, respectively. SNALP-18328 treated V30M hTTR transgenic mice had a dose-dependent and significant decrease in liver TTR mRNA levels relative to the PBS control group, reaching a maximum reduction of 97% ($p < 0.001$) at 3 mg/kg SNALP-18328, and a 50% reduction (ED50) at ~ 0.15 mg/kg SNALP-18328. Serum TTR protein was also suppressed in a dose-dependent manner, with a maximum reduction of serum TTR protein of 99% ($p < 0.01$) (relative to pre-dose levels) at 3 mg/kg SNALP-18328, consistent with the reduction in TTR mRNA levels. SNALP-1955 at 3 mg/kg did not have a statistically significant effect on either TTR mRNA or protein levels, compared to PBS.

[0271] These results demonstrate that SNALP-18328, when administered IV, is active in suppressing mutant V30M TTR mRNA in the transgenic mouse liver, which results in reduction of mutant V30M TTR protein in the circulation.

Example 7. Durability of TTR mRNA and protein suppression by SNALP-18328 in the transgenic mouse

[0272] To evaluate the durability of TTR mRNA and protein suppression by SNALP-18328, AD-18328 was formulated in SNALP and administered by IV bolus to V30M hTTR transgenic mice. At various timepoints post-dose, liver TTR mRNA levels and serum TTR protein levels were quantified. 8- to 12-week old V30M hTTR transgenic mice (4 animals/ group) were intravenously (IV) administered 200 μ l SNALP-18328 (1 mg/kg) or SNALP-1955 (1 mg/kg, with negative control siRNA AD-1955 which targets the non-mammalian gene luciferase). Mice used were Mus musculus strain H129-hTTR KO from Institute of Molecular and Cellular Biology, Porto, Portugal. Briefly, hTTR H129 transgenic mice were crossed with a H129 endogenous TTR KO mice (null mice to generate the H129-hTTR transgenic mice, in a null mouse TTR background (Maeda, S., (2003), Use of genetically altered mice to study the role of serum amyloid P component in amyloid deposition. Amyloid Suppl. 1, 17-20). Days 3, 8, 15, or 22 post-dose, animals in both treatment groups were given a lethal dose of ketamine/xylazine. Serum samples were collected and stored at -80°C until analysis. Liver tissue was collected, flash-frozen and stored at -80°C until

processing.

[0273] For TTR mRNA quantitation, frozen liver tissue was ground into powder, and lysates were prepared. TTR mRNA levels relative to those of GAPDH mRNA were determined in the lysates by using a branched DNA assay (QuantiGene Reagent System, Panomics, Fremont, CA). Briefly, the QuantiGene assay (Genospectra) was used to quantify mRNA levels in tissue sample lysates according to the manufacturer's instructions. The mean level of TTR mRNA was normalized to the mean level of GAPDH mRNA for each sample. Group means of the normalized values were then further normalized to the mean value for the PBS treated group, to obtain the relative level of TTR mRNA expression.

[0274] For TTR protein quantitation, serum was assayed using the AssayPro (St. Charles, MO) Assaymax PreAlbumin ELISA Kit according to the manufacturer's protocol.

[0275] The results are shown in FIG. 7A and FIG. 7B for liver mRNA and serum protein, respectively. A single IV bolus administration of SNALP-18328 in the hTTR V30M transgenic mouse resulted in durable inhibition of TTR mRNA levels in the liver and TTR protein levels in the serum. Compared to the control group (1 mg/ml SNALP-1955), a single IV administration of SNALP-18328 at 1 mg/kg significantly reduced relative TTR mRNA levels on Days 3, 8, 15 and 22 post-dose by 96% ($p < 0.001$), 90% ($p < 0.001$), 82% ($p < 0.001$) and 73% ($p < 0.001$), respectively, and did not return to baseline levels at termination of the study (Day 22 post-dose). Protein levels also decreased with a maximum reduction of serum TTR of 97% ($p < 0.001$) (relative to SNALP-1955) at Day 3 post-dose. At Days 8, 15, and 22 post-dose, TTR protein levels were suppressed by 72% ($p < 0.05$), 32% ($p < 0.05$), and 40% ($p < 0.001$), respectively, relative to SNALP-1955.

[0276] These results demonstrate that a single IV administration of SNALP-18328 produces durable suppression of target liver mRNA and serum protein levels in the V30M hTTR transgenic mouse, with significant reductions of both liver TTR mRNA and serum TTR protein at 22 days post-dose.

Example 8. Durability of serum TTR protein suppression by SNALP-18328 in the non-human primate

[0277] To evaluate the durability of serum TTR protein suppression by SNALP-18328, AD-18328 was formulated in SNALP and administered by IV infusion to non-human primates. At various timepoints post-dose, serum TTR protein levels were quantified.

[0278] Cynomolgus monkeys (*Macaca fascicularis*) ($n = 5$ animals/group for SNALP-18328 groups and $n = 3$ animals/group for SNALP-1955 and PBS groups) were administered a 15-minute IV infusion of SNALP-18328 (0.3, 1 or 3 mg/kg), SNALP-1955 (3 mg/kg) with negative control siRNA AD-1955 which targets the non-mammalian gene luciferase), or PBS. At Days 0, 1, 2, 3, 4, 5, 7, 10, and 14 of the dosing phase, serum samples were collected and stored at -80°C until analysis.

[0279] Western blot analysis was used to evaluate TTR protein levels in serum samples. Serum samples from each group were pooled and diluted 1:1 with Laemmli sample buffer (β -mercaptoethanol was added at a 1:20 dilution). The samples were heated at 95°C for 10 minutes. 12.5 μl of each sample was loaded in each lane of a 10-20% Criterion (Biorad, Hercules, CA) prep gel and separated by SDS-PAGE at 120V for 1.5 hrs, then transferred to a nitrocellulose membrane using a semi-dry system at 15V for 1 hour. The membrane was blocked overnight at 4°C in LiCOR (Lincoln, NE) blocking buffer diluted 1:1 with 1X PBS. The blot was probed first with primary antibodies (goat anti-TTR from Santa Cruz (Santa Cruz, CA) at a dilution of 1:1000 diluted in LiCOR blocking buffer/PBS on a rocker for 1 hr at room temperature. Blots were washed 4X with PBS + 0.2% Tween 20 (10 minutes per wash). The fluorescent labeled secondary antibodies (anti-goat 680nm from Invitrogen (Carlsbad, CA) were added at a dilution of 1:10,000 in LiCOR blocking buffer/PBS and the blot was incubated for 1 hour at room temperature. After incubation, blots were washed 4X with PBS + 0.2% Tween 20 followed by one wash with 1X PBS. The LiCOR's Odyssey Infrared Imaging System was used to detect the protein bands. TTR monomer migrates at 15 kDa.

[0280] The results are shown in FIG. 8. Serum TTR protein levels showed a dose-dependent reduction with 1 or 3 mg/kg SNALP-18328, as compared to pre-dose (Day 0) levels. The duration of suppression, following a single IV administration of SNALP-18328 is at least 14 days after 1 or 3 mg/kg SNALP-18328 treatment.

[0281] These results demonstrate that a single IV administration of SNALP-18328 produces durable suppression of TTR protein in the circulation in the non-human primate (*Macaca fascicularis*), with significant reduction of TTR protein at 14 days post-dose.

Example 9: *In vivo* reduction of mutant (V30M) TTR in peripheral tissues by SNALP-18328 in the transgenic mouse

[0282] To evaluate the efficacy of SNALP-18328 in reducing TTR in peripheral tissues, hTTR V30M/HSF-1 knock-out mice were evaluated with immunohistochemical staining for TTR. Two-month old hTTR V30M/HSF-1 knock-out mice (Maeda, S., (2003), Use of genetically altered mice to study the role of serum amyloid P component in amyloid deposition. *Amyloid Suppl.* 1, 17-20) were administered an IV bolus of 3 mg/kg SNALP-18328 (12 animals), 3 mg/kg SNALP-1955 (with negative control siRNA AD-1955 which targets the non-mammalian gene luciferase, 4 animals), or PBS (4 animals) once every two weeks for a total of four doses on days 0, 14, 28, and 42. TTR liver mRNA levels and TTR-immunoreactivity in multiple peripheral tissues were evaluated at 8 weeks post-first dose on day 56.

[0283] Mice were anesthetised with 1 mg/kg medetomidine, and given a lethal dose of ketamine. Tissues and organs of interest were collected. For immunohistochemistry, esophagus (E), stomach (S), intestine (duodenum (I1) and colon (I4)), nerve (N) and dorsal root ganglia (D) were fixed in neutral buffered formalin and embedded in paraffin. For TTR detection, rabbit anti-human TTR primary antibody (1:1000, DAKO, Denmark), and anti-rabbit biotin-conjugated secondary antibody (1:20 Sigma, USA) were followed by extravidin labelling (1:20, Sigma, USA) in order to stain for the TTR protein. The reaction was developed with 3-amino-9-ethyl carbazole, AEC (Sigma, USA). Semi-quantitative analysis of immunohistochemical slides was performed using Scion image quant program that measures the area occupied by the substrate reaction color and normalizes this value to the total image area. Mean values of % occupied area are displayed with the corresponding standard deviation. Each animal tissue was evaluated in four different areas. The presence of human TTR in parasympathetic ganglia of the stomach and intestine was studied by double immunofluorescent staining with rabbit anti-human TTR (1:1000, DAKO, Denmark) and mouse anti-PGP9.5 (1:40, Serotec, USA) as the primary antibodies; secondary antibodies were, respectively: anti-rabbit Alexa Fluor 488 (Molecular probes, UK) and goat anti-mouse Alexa Fluor 568 (Molecular probes, UK). Slides were mounted with vectashield (Vector) and visualized in a Zeiss Cell Observer System microscope (Carl Zeiss, Germany) equipped with filters for FITC and rhodamine.

[0284] The results are graphed in FIG. 9. In contrast with PBS and SNALP-1955 treated animals, SNALP-18328 treated animals had a significant reduction of TTR-immunoreactivity in all tissues examined (esophagus (E), stomach (S), intestine (duodenum (I1) and colon (I4)), nerve (N) and dorsal root ganglia (D)).

[0285] These results demonstrate that SNALP-18328 administration to hTTR V30M/HSF-1 knock-out mice causes a significant reduction of TTR protein in peripheral tissues and organs, including esophagus, stomach, intestine (duodenum and colon), nerve, and dorsal root ganglion.

Example 10. *In vivo* reduction of wild-type TTR mRNA in the non-human primate liver by XTC-SNALP-18328

[0286] To evaluate the efficacy of the novel lipid nanoparticle formulation XTC-SNALP for delivery of siRNA in non-human primate, TTR siRNA AD-18328 was formulated in XTC-SNALP (XTC-SNALP-18328) and administered by 15-minute IV infusion, and liver TTR mRNA was quantified. Cynomolgus monkeys (*Macaca fascicularis*) were administered 15-minute IV infusions of XTC-SNALP-18328 (0.03, 0.1, 0.3 or 1 mg/kg) or XTC-SNALP-1955 (1 mg/kg, with negative control siRNA AD-1955 which targets the non-mammalian gene luciferase). At forty-eight hours post-dosing, monkeys were anesthetized with sodium pentobarbital and exsanguinated. Liver tissue for TTR mRNA determination was collected, flash-frozen, and stored at -80°C until processing. Methods used for TTR mRNA quantitation in liver tissue were similar to those described in Example 5 above.

[0287] The results are shown in FIG. 10. XTC-SNALP -18328 reduced TTR mRNA levels in the liver in a dose-dependent manner, compared to the negative control XTC-SNALP -1955. The mRNA ED50 was determined to be ~ 0.1 mg/kg XTC-SNALP -18328.

[0288] These results demonstrate that XTC-SNALP-18328 is effective in suppressing wild-type TTR mRNA in non-human primate liver when administered by IV infusion.

Example 11: *In vivo* reduction of wild-type TTR mRNA in the non-human primate liver by LNP09-18328 and LNP11-18328

[0289] To evaluate the efficacy of two novel lipid nanoparticle formulations, LNP09 and LNP11, for delivery of siRNA in non-human primate, TTR siRNA AD-18328 was formulated in LNP09 (LNP09-18328) or LNP11 (LNP11-18328), and administered

by 15-minute IV infusion, and liver TTR mRNA and serum TTR protein levels were assayed. Cynomolgus monkeys (*Macaca fascicularis*) were administered 15-minute IV infusions of LNP09-18328 (0.03, 0.1, or 0.3 mg/kg), LNP11-18328 (0.03, 0.1, or 0.3 mg/kg), or PBS. Liver biopsy samples were collected at 48 hrs post-dosing, flash-frozen, and stored at -80°C until processing. Serum was collected before dosing (pre-bleed), and on Days 1, 2, 4, 7, 14, 21 and 28 post-dosing and stored at -80°C until processing. Methods used for TTR mRNA quantitation in liver tissue and serum TTR protein evaluation were similar to those described in Examples 5 and 8 above.

[0290] The results are shown in FIG. 11A for mRNA, and in FIG. 11B and FIG. 11C for protein. LNP09-18328 and LNP11-18328 treated animals showed a dose-dependent decrease in TTR mRNA levels in the liver, reaching a maximum reduction at 0.3 mg/kg of ~ 85% (LNP09-18328) and ~ 90% (LNP11-18328) mRNA relative to the PBS control. The mRNA ED50 was determined to be ~ 0.02 mg/kg for both LNP09-18328 and LNP11-18328. At Day 7 post-dosing, serum samples also exhibited a dose-dependent reduction of TTR protein for 0.1 and 0.3 mg/kg LNP09-18328 and LNP11-18328, compared to PBS control levels. FIG 11C shows a decrease in TTR protein levels with a 0.3mg/kg dose of LNP09-18328 that persisted over at least 28 days post-dosing, as compared to the PBS control group and as compared with the pre-bleed samples.

[0291] These results demonstrate that LNP09-18328 and LNP11-18328 are effective in suppressing wild-type TTR mRNA in non-human primate liver and wild-type TTR protein in the circulation, when administered by IV infusion. Furthermore, the suppression with LNP09-18328 is durable, persisting for at least 28 days following the IV infusion.

Example 12. Synthesis of TTR Tiled Sequences

[0292] A set of TTR duplexes ("tiled duplexes") were designed that targeted the TTR gene near the target region of AD-18328, which targets the human TTR gene starting at nucleotide 628 of NM_000371.3.

[0293] In the examples below, the numbering representing the position of the 5' base of an siRNA on the transcript is based on NM_000371.3 (FIG. 12; SEQ ID NO:1331). In the examples shown above, the numbering for siRNA targeting human siRNA was based on NM_000371.2 (FIG. 13A). NM_000371.3 extends the sequence of the 5' UTR by 110 bases compared to NM_000371.2, as shown in FIG. 14. Thus, as an example, the starting position of AD-18328 is 628 on NM_000371.3 and 518 on NM_000371.2 (FIG. 14).

[0294] TTR tiled sequences were synthesized on MerMade 192 synthesizer at 1umol scale. For all the sequences in the list, 'endolight' chemistry was applied as detailed below.

- All pyrimidines (cytosine and uridine) in the sense strand contained 2'-O-Methyl bases (2' O-Methyl C and 2'-O-Methyl U)
- In the antisense strand, pyrimidines adjacent to (towards 5' position) ribo A nucleoside were replaced with their corresponding 2-O-Methyl nucleosides
- A two base dTdT extension at 3' end of both sense and anti sense sequences was introduced
- The sequence file was converted to a text file to make it compatible for loading in the MerMade 192 synthesis software

Synthesis, Cleavage and deprotection:

[0295] The synthesis of TTR sequences used solid supported oligonucleotide synthesis using phosphoramidite chemistry. The synthesis of the sequences was performed at 1 um scale in 96 well plates. The amidite solutions were prepared at 0.1 M concentration and ethyl thio tetrazole (0.6M in Acetonitrile) was used as activator. The synthesized sequences were cleaved and deprotected in 96 well plates, using methylamine in the first step and fluoride reagent in the second step. The crude sequences were precipitated using acetone: ethanol (80:20) mix and the pellet were re-suspended in 0.2M sodium acetate buffer. Samples from each sequence were analyzed by LC-MS to confirm the identity, UV for quantification and a selected set of samples by IEX chromatography to determine purity.

Purification and desalting:

[0296] TTR tiled sequences were purified on AKTA explorer purification system using Source 15Q column. A column

temperature of 65C was maintained during purification. Sample injection and collection was performed in 96 well (1.8mL - deep well) plates. A single peak corresponding to the full length sequence was collected in the eluent. The purified sequences were desalted on a Sephadex G25 column using AKTA purifier. The desalted TTR sequences were analyzed for concentration (by UV measurement at A260) and purity (by ion exchange HPLC). The single strands were then submitted for annealing.

TTR Single strands and duplexes:

[0297] A detailed list of TTR tiled duplexes and corresponding single strands (sense and antisense) are shown in the table below (Table 13).

Table 13: TTR tiled duplexes and corresponding single strands

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.3, SEQ ID NO:1331).					
Duplex #	Position	Oligo #	Strand	Sequence (5' to 3')	SEQ ID NO:
AD-18323	618	A-32335	S	GGGAuuucAuGuAAccAAGdTdT	1332
		A-32336	AS	CUUGGUuAcAUGAAAUCCdTdT	1333
AD-18324	619	A-32337	S	GGAuuucAuGuAAccAAGAdTdT	1334
		A-32338	AS	UCUUGGUuAcAUGAAAUCCdTdT	1335
AD-23000	620	A-42927	S	GAuuucAuGuAAccAAGAGdTdT	1336
		A-42928	AS	CUCUUGGUuAcAUGAAAUCdTdT	1337
AD-23001	621	A-42929	S	AuuucAuGuAAccAAGAGudTdT	1338
		A-42930	AS	ACUCUUGGUuAcAUGAAAUdTdT	1339
AD-23002	622	A-42931	S	uuucAuGuAAccAAGAGuAdTdT	1340
		A-42932	AS	uACUCUUGGUuAcAUGAAAdTdT	1341
AD-23003	623	A-42933	S	uucAuGuAAccAAGAGuAdTdT	1342
		A-42934	AS	AuACUCUUGGUuAcAUGAAAdTdT	1343
AD-18325	624	A-32339	S	ucAuGuAAccAAGAGuAuudTdT	1344
		A-32340	AS	AAuACUCUUGGUuAcAUGAdTdT	1345
AD-23004	625	A-42935	S	cAuGuAAccAAGAGuAuucdTdT	1346
		A-42936	AS	GAAuACUCUUGGUuAcAUGdTdT	1347
AD-18326	626	A-32341	S	AuGuAAccAAGAGuAuuccdTdT	1348
		A-32342	AS	GGAAuACUCUUGGUuAcAUdTdT	1349
AD-18327	627	A-32343	S	uGuAAccAAGAGuAuuccAdTdT	1350
		A-32344	AS	UGGAAuACUCUUGGUuAcAdTdT	1351
AD-23005	628	A-42937	S	uAAccAAGAGuAuuccAuudTdT	1352
		A-42938	AS	AAUGGAAuACUCUUGGUuAdTdT	1353
AD-23006	629	A-42939	S	AAccAAGAGuAuuccAuudTdT	1354
		A-42940	AS	AAAUGGAAuACUCUUGGUUdTdT	1355
AD-23007	631	A-42941	S	AccAAGAGuAuuccAuuuudTdT	1356
		A-42942	AS	AAAAUGGAAuACUCUUGGUdTdT	1357
AD-23008	632	A-42943	S	ccAAGAGuAuuccAuuuuudTdT	1358
		A-42944	AS	AAAAAUGGAAuACUCUUGGdTdT	1359
AD-23009	633	A-42945	S	cAAGAGuAuuccAuuuuuAdTdT	1360
		A-42946	AS	uAAAAAUGGAAuACUCUUGdTdT	1361
AD-23010	634	A-42947	S	AAGAGuAuuccAuuuuuAcdTdT	1362
		A-42948	AS	GuAAAAAUGGAAuACUCUUDdTdT	1363
AD-23011	635	A-42949	S	AGAGuAuuccAuuuuuAcudTdT	1364
		A-42950	AS	AGuAAAAAUGGAAuACUCUdTdT	1365
AD-23012	636	A-42951	S	GAGuAuuccAuuuuuAcuAdTdT	1366

Strand: s= sense; as= antisense; Position: position of 5' base on transcript (NM_000371.3, SEQ ID NO:1331).					
Duplex #	Position	Oligo #	Strand	Sequence (5' to 3')	SEQ ID NO:
		A-42952	AS	uAGuAAAAAUGGAAuACUCdTdT	1367
AD-23013	637	A-42953	S	AGuAuuccAuuuuuAcuAAdTdT	1368
		A-42954	AS	UuAGuAAAAAUGGAAuACUdTdT	1369
AD-23014	638	A-42955	S	GuAuuccAuuuuuAcuAAAdTdT	1370
		A-42956	AS	UUuAGuAAAAAUGGAAuACdTdT	1371
AD-23015	639	A-42957	S	uAuuccAuuuuuAcuAAAGdTdT	1372
		A-42958	AS	CUUuAGuAAAAAUGGAAuAdTdT	1373
AD-23016	640	A-42959	S	AuuccAuuuuuAcuAAAGcdTdT	1374
		A-42960	AS	GCUUuAGuAAAAAUGGAAUdTdT	1375
AD-23017	641	A-42961	S	uuccAuuuuuAcuAAAGcAdTdT	1376
		A-42962	AS	UGC UUuAGuAAAAAUGGAAdTdT	1377
AD-23018	642	A-42963	S	uccAuuuuuAcuAAAGcAGdTdT	1378
		A-42964	AS	CUGCUUuAGuAAAAAUGGAdTdT	1379
AD-23019	643	A-42965	S	ccAuuuuuAcuAAAGcAGudTdT	1380
		A-42966	AS	ACUGCUUuAGuAAAAAUGGdTdT	1381
AD-23020	644	A-42967	S	cAuuuuuAcuAAAGcAGuGdTdT	1382
		A-42968	AS	cACUGCUUuAGuAAAAAUGdTdT	1383
AD-23021	645	A-42969	S	AuuuuuAcuAAAGcAGuGudTdT	1384
		A-42970	AS	AcACUGCUUuAGuAAAAAUdTdT	1385
AD-23022	646	A-42971	S	uuuuuAcuAAAGcAGuGuudTdT	1386
		A-42972	AS	AAcACUGCUUuAGuAAAAAdTdT	1387
AD-23023	647	A-42973	S	uuuuuAcuAAAGcAGuGuuudTdT	1388
		A-42974	AS	AAAcACUGCUUuAGuAAAAAdTdT	1389
AD-23024	648	A-42975	S	uuuAcuAAAGcAGuGuuuudTdT	1390
		A-42976	AS	AAAACACUGCUUuAGuAAAAdTdT	1391
AD-23025	649	A-42977	S	uuAcuAAAGcAGuGuuuucdTdT	1392
		A-42978	AS	GAAAACACUGCUUuAGuAAAdTdT	1393
AD-23026	650	A-42979	S	uAcuAAAGcAGuGuuuucAdTdT	1394
		A-42980	AS	UGAAAACACUGCUUuAGuAdTdT	1395
AD-23027	651	A-42981	S	AcuAAAGcAGuGuuuucAcdTdT	1396
		A-42982	AS	GUGAAAACACUGCUUuAGUdTdT	1397
AD-23028	652	A-42983	S	cuAAAGcAGuGuuuucAccdTdT	1398
		A-42984	AS	GGUGAAAACACUGCUUuAGdTdT	1399
AD-18330	653	A-32349	S	uAAAGcAGuGuuuucAccudTdT	1400
		A-32350	AS	AGGUGAAAACACUGCUUuAdTdT	1401
AD-23029	654	A-42985	S	AAAGcAGuGuuuucAccucdTdT	1402
		A-42986	AS	GAGGUGAAAACACUGCUUuUdTdT	1403
AD-23030	655	A-42987	S	AAGcAGuGuuuucAccucAdTdT	1404
		A-42988	AS	UGAGGUGAAAACACUGCUUdTdT	1405
AD-23031	656	A-42989	S	AGcAGuGuuuucAccucAudTdT	1406
		A-42990	AS	AUGAGGUGAAAACACUGCUdTdT	1407
AD-18328	628	A-32345	S	GuAAccAAGAGuAuuccAudTdT	1408
		A-32346	AS	AUGGAAuACUCUUGGUuACdTdT	1409

Example 13. *In vitro* Screening of TTR Tiled siRNAs

[0298] Tiled TTR duplexes were assayed in Hep3B cells for inhibition of endogenous TTR expression using real time PCR assays.

[0299] Cell culture and transfection: Hep3B cells (ATCC, Manassas, VA) were grown to near confluence at 37°C in an atmosphere of 5% CO₂ in Eagle's Minimum Essential Medium (EMEM, ATCC) supplemented with 10% FBS, streptomycin, and glutamine (ATCC) before being released from the plate by trypsinization. Reverse transfection was carried out by adding 5µl of Opti-MEM to 5µl of each siRNA in individual wells of a 96-well plate. To this 10µl of Opti-MEM plus 0.2µl of Lipofectamine RNAiMax was added per well (Invitrogen, Carlsbad CA. cat # 13778-150) and the mixture was incubated at room temperature for 15 minutes. 80µl of complete growth media described above, but without antibiotic containing 2.0 x10⁴ Hep3B cells were then added. Cells were incubated for 24 hours prior to RNA purification. Experiments were performed at 0.1 or 10nM final duplex concentration.

[0300] Total RNA isolation using MagMAX-96 Total RNA Isolation Kit (Applied Biosystems, Foster City CA, part #: AM1830): Cells were harvested and lysed in 140µl of Lysis/Binding Solution then mixed for 1 minute at 850rpm using an Eppendorf Thermomixer (the mixing speed was the same throughout the process). Twenty micro liters of magnetic beads and Lysis/Binding Enhancer mixture were added into cell-lysate and mixed for 5 minutes. Magnetic beads were captured using magnetic stand and the supernatant was removed without disturbing the beads. After removing supernatant, magnetic beads were washed with Wash Solution 1 (isopropanol added) and mixed for 1 minute. Beads were capture again and supernatant removed. Beads were then washed with 150µl Wash Solution 2 (Ethanol added), captured and supernatant was removed. 50µl of DNase mixture (MagMax turbo DNase Buffer and Turbo DNase) was then added to the beads and they were mixed for 10 to 15 minutes. After mixing, 100µl of RNA Rebinding Solution was added and mixed for 3 minutes. Supernatant was removed and magnetic beads were washed again with 150µl Wash Solution 2 and mixed for 1 minute and supernatant was removed completely. The magnetic beads were mixed for 2 minutes to dry before RNA was eluted with 50µl of water.

[0301] cDNA synthesis using ABI High capacity cDNA reverse transcription kit (Applied Biosystems, Foster City, CA, Cat #4368813): A master mix of 2µl 10X Buffer, 0.8µl 25X dNTPs, 2µl Random primers, 1µl Reverse Transcriptase, 1µl RNase inhibitor and 3.2µl of H₂O per reaction were added into 10µl total RNA. cDNA was generated using a Bio-Rad C-1000 or S-1000 thermal cycler (Hercules, CA) through the following steps: 25°C 10 min, 37°C 120 min, 85°C 5 sec, 4°C hold.

[0302] Real time PCR: 2 µl of cDNA were added to a master mix containing 0.5µl GAPDH TaqMan Probe (Applied Biosystems Cat # 4326317E), 0.5µl TTR TaqMan probe (Applied Biosystems cat #HS00174914 M1) and 10µl Roche Probes Master Mix (Roche Cat # 04887301001) per well in a LightCycler 480 384 well plate (Roche cat # 0472974001). Real time PCR was done in a LightCycler 480 Real Time PCR machine (Roche). Each duplex was tested in two independent transfections and each transfection was assayed in duplicate.

[0303] Real time data were analyzed using the $\Delta\Delta CT$ method. Each sample was normalized to GAPDH expression and knockdown was assessed relative to cells transfected with the non-targeting duplex AD-1955. Table 14 shows the knockdown of TTR using the siRNAs. Data are expressed as the percent of message remaining relative to cells targeted with AD-1955.

[0304] Many but not all tiled TTR-dsRNAs, targeting TTR near the target of AD-18328, reduced TTR mRNA by at least 70% when transfected into Hep3B cells at 0.1 nM.

Table 14: Inhibition of TTR by tiled dsRNA targeting TTR near target of AD-18328.

Duplex #	% message remaining 0.1nM	% SD 0.1nM	% message remaining 10nM	% SD 10nM
AD-18323	6.7	1.90	1.7	0.02
AD-18324	1.8	0.58	0.9	0.10
AD-23000	5.5	0.93	2.1	0.87
AD-23001	15.2	4.89	4.9	1.74
AD-23002	3.1	1.12	1.4	0.55
AD-23003	17.3	3.13	1.7	0.06
AD-18325	1.5	0.27	1.4	0.66
AD-23004	9.0	0.15	10.5	0.96
AD-18326	22.0	1.85	7.6	0.78
AD-18327	11.6	2.64	9.6	1.67

Duplex #	% message remaining 0.1nM	% SD 0.1nM	% message remaining 10nM	% SD 10nM
AD-18328	1.1	0.70	0.6	0.16
AD-23005	0.8	0.31	0.6	0.21
AD-23006	1.5	0.46	1.2	0.43
AD-23007	2.4	0.91	1.9	0.46
AD-23008	0.6	0.10	0.8	0.26
AD-23009	1.0	0.13	0.9	0.22
AD-23010	60.1	15.66	66.2	22.71
AD-23011	56.5	16.99	53.6	4.70
AD-23012	7.7	2.36	7.7	3.25
AD-23013	7.0	0.64	8.0	1.06
AD-23014	0.7	0.01	0.6	0.10
AD-23015	15.4	0.25	16.5	7.07
AD-23016	27.1	0.37	6.7	1.80
AD-23017	4.5	1.26	1.4	0.40
AD-23018	44.6	9.45	7.5	1.09
AD-23019	2.2	0.68	0.8	0.10
AD-23020	52.7	6.45	29.7	1.17
AD-23021	95.4	16.16	45.0	3.00
AD-23022	70.1	3.01	60.8	12.11
AD-23023	2.7	1.12	1.8	0.07
AD-23024	1.7	0.30	1.8	0.33
AD-23025	64.2	13.21	10.5	1.34
AD-23026	1.9	0.15	1.9	0.78
AD-23027	2.5	0.21	1.6	0.49
AD-23028	6.7	4.41	1.2	0.50
AD-18330	6.0	0.56	5.7	1.15
AD-23029	4.5	0.47	1.6	0.10
AD-23030	3.9	0.25	3.3	0.84
AD-23031	3.4	0.78	1.7	0.02

Example 14. Evaluation of Infusion Duration on Efficacy of a Single Intravenous Administration of SNALP-18534 in Sprague-Dawley Rats

Objectives

[0305] To determine the effect of infusion duration on efficacy of a single IV infusion of SNALP-18534 on liver TTR mRNA levels in Sprague-Dawley rats.

Table 15: Abbreviations and definitions used

SNALP-18534	Rodent transthyretin specific siRNA formulated in SNALP
SNALP-1955	Non-mammalian luciferase specific siRNA formulated in SNALP

[0306] The sequences of the sense and antisense strands of AD-18534 are reproduced below from the tables above:

Strand	Oligo #	Position	Sequence 5' to 3'	SEQ ID NO:
s	A-32755	532	cAGuGuucuuGcucuAuAAdTdT	1289

Strand	Oligo #	Position	Sequence 5' to 3'	SEQ ID NO:
as	A-32756	550	UuAuAGAGcAAGAAcACUGdTdT	1290

Study Materials

Test Article(s)

[0307] SNALP-18534 is comprised of an siRNA targeting rodent TTR mRNA (AD-18534), formulated in stable nucleic acid lipid particles (SNALP) for delivery to target tissues. The SNALP formulation (lipid particle) consists of a novel aminolipid (DLinDMA), a PEGylated lipid (mPEG2000-C-DMA), a neutral lipid (DPPC) and cholesterol. The ratio of lipid:nucleic acid in the SNALP formulation is approximately 5.8:1 (w:w). SNALP-1955 contains an siRNA targeting the non-mammalian luciferase mRNA, is formulated with the identical lipid particle as SNALP-18534, and serves as a non-pharmacologically active control. Dose levels are expressed as mg/kg based on the weight of siRNA content.

Study Design & Procedures

Animals and test article administration:

[0308] The study was comprised of 9 groups of Sprague-Dawley rats (4 males/ group). The animals were allowed to have at least a 2 day acclimation period before the study and all animals were 7 weeks old at the initiation of dosing. The dose administered was calculated based upon body weight data collected prior to dosing on Day 1. The test and control articles were administered as a single 15-minute, 1-hour, 2-hour, or 3-hour IV infusion via the tail vein using a 24G ¾" cannula sealed with a Baxter Injection Site septum connected via 27G Terumo butterfly needle to a Baxter AS40A Syringe Pump. The dose volume was 3 ml/kg, the infusion rate was 12 ml/kg/hr, and animals were freely moving in the cages during dosing. Rats were divided into nine treatment groups and administered a single IV infusion of SNALP-18534, SNALP-1955, or PBS as shown in Table 16:

Table 16: Test Animal Dosage Groups

Group	N	Test Article	Infusion Duration	Dose
A	4	PBS	15 minute	---
B	4	PBS	3 hour	---
C	4	SNALP -1955	1 hour	1 mg/kg
D	4	SNALP -1955	2 hour	1 mg/kg
E	4	SNALP -1955	3 hour	1 mg/kg
F	4	SNALP-18534	15 minute	1 mg/kg
G	4	SNALP-18534	1 hour	1 mg/kg
H	4	SNALP-18534	2 hour	1 mg/kg
I	4	SNALP-18534	3 hour	1 mg/kg

Tissue collection and RNA isolation:

[0309] On Day 0, animals were anesthetized by isoflurane inhalation and pre-dosing blood samples were collected into serum separator tubes by retro-orbital bleed. The blood samples were allowed to clot at room temperature for approximately 30 minutes prior to centrifugation at 4°C. Serum samples were then stored at -80°C until analysis was performed. On Day 3, animals in all nine treatment groups were given a lethal dose of ketamine/xylazine. Blood was collected via caudal vena cava into serum separation tubes, and then allowed to clot at room temperature for approximately 30 minutes prior to centrifugation at 4°C. Serum samples were stored at - 80°C until analysis was performed. Liver tissue was harvested and snap frozen on dry ice. Frozen liver tissue was ground and tissue lysates were prepared for liver mRNA quantitation.

TTR mRNA Quantitation:

[0310] TTR mRNA levels relative to those of GAPDH mRNA were determined in the lysates by using a branched DNA assay (QuantiGene Reagent System, Panomics, Fremont, CA). Briefly, the QuantiGene assay (Genospectra) was used to quantify mRNA levels in tissue sample lysates according to the manufacturer's instructions. The mean level of TTR mRNA was normalized to the mean level of GAPDH mRNA for each sample.

[0311] To obtain the relative level of TTR mRNA expression, group mean values for SNALP-1955 and SNALP-18534 treated groups with 15-minute, 1 hour and 2 hour infusion durations were then normalized to the mean value for the PBS treated group with 15-minute infusion whereas group mean values for SNALP-1955 and SNALP-18534 treated groups with 3 hour infusion duration were then normalized to the mean value for the PBS treated group with 3 hour infusion duration.

Results

[0312] As shown in FIG. 16, a single IV infusion of 1 mg/kg SNALP-18534 with different infusion durations of 15 minutes to 3 hours results in comparable inhibition of liver TTR mRNA levels measured two days after dosing. A single IV infusion of 1 mg/kg SNALP-18534 also showed durable TTR downregulation over 29 days following a single 15 minute IV infusion, as compared to SNALP-1955 control (data not shown). Compared to the PBS-treated group, a single 15-minute, 1-hour, 2-hour, or 3-hour IV infusion of SNALP-18534 at 1 mg/kg significantly reduced relative TTR mRNA expression levels by 94% ($p < 0.001$), 94% ($p < 0.001$), 92% ($p < 0.001$) and 93% ($p < 0.001$), respectively. Specificity of SNALP-18534 activity is demonstrated by lack of significant target inhibition by SNALP-1955 administration via 1-hour, 2-hour, or 3-hour IV infusion at the same dose level.

Conclusions

[0313] This study demonstrates that varying the infusion duration from 15 minutes to up to 3 hours does not affect the efficacy of a single IV administration of 1 mg/kg SNALP-18534 in rats, as assessed by reduction of TTR mRNA levels in the liver.

Example 15. *In vivo* reduction of wild-type TTR mRNA in the rat liver by LNP07-18534 and LNP08-18534

[0314] To evaluate the efficacy of 2 novel lipid nanoparticle formulations, LNP07 and LNP08, for delivery of siRNA in the rat, the rodent-specific TTR siRNA, AD-18534, was formulated in LNP07 (LNP07-18534) or LNP08 (LNP08-18534), and administered by 15-minute IV infusion, and liver TTR mRNA was quantified. Sprague-Dawley rats (4 animals per group) were administered 15-minute IV infusions of LNP07-18534 (0.03, 0.1, 0.3 or 1 mg/kg), LNP08-18534 (0.01, 0.03 or 0.1 mg/kg), or LNP07-1955 (1 mg/kg) or LNP08-1955 (0.1 mg/kg) containing the negative control siRNA AD-1955 which targets the non-mammalian gene luciferase. Forty-eight hours later, animals were euthanized and liver tissue was collected, flash-frozen and stored at -80°C until processing.

[0315] For TTR mRNA quantitation, frozen liver tissue was ground into powder, and lysates were prepared. TTR mRNA levels relative to those of GAPDH mRNA were determined in the lysates by using a branched DNA assay (QuantiGene Reagent System, Panomics, Fremont, CA). Briefly, the QuantiGene assay (Genospectra) was used to quantify mRNA levels in tissue sample lysates according to the manufacturer's instructions. The mean level of TTR mRNA was normalized to the mean level of GAPDH mRNA for each sample. Group means of the normalized values were then further normalized to the mean value for the PBS treated group, to obtain the relative level of TTR mRNA expression.

[0316] The results are shown in FIG. 17. LNP07-18534 reduced TTR mRNA levels in the liver in a dose-dependent manner, with 94% suppression of TTR mRNA at 1 mg/kg. The effect was specific, since the negative control LNP07-1955 at 1 mg/kg did not significantly affect TTR mRNA levels compared to the PBS control. The mRNA ED50 was determined to be ~ 0.05 mg/kg LNP07-18534. LNP08-18534 reduced TTR mRNA levels in the liver in a dose-dependent manner, with 86% suppression of TTR mRNA at 0.1 mg/kg. The effect was specific, since the negative control LNP08-1955 at 0.1 mg/kg did not significantly affect TTR mRNA levels compared to the PBS control. The mRNA ED50 was determined to be ~ 0.02 mg/kg

LNP08-18534.

[0317] These results demonstrate that LNP07-18534 and LNP08-18534 are effective in suppressing wild-type TTR mRNA in the rat liver when administered by IV infusion, and that LNP07 and LNP08 are effective formulations for delivering siRNA to the liver.

Example 16: Reduction of TTR liver mRNA by a single intravenous administration of LNP09-18534 or LNP11-18534 in Sprague-Dawley Rats

Objective:

[0318] To evaluate the efficacy of two novel lipid nanoparticle (LNP) formulations for delivery of the rodent TTR-specific siRNA, AD-18534 in the Sprague-Dawley rat for reducing endogenous (wild type) liver TTR mRNA levels. Rats were intravenously dosed via a 15 minute infusion with either 0.01, 0.03, 0.1, or 0.3 mg/kg LNP09-18534, LNP11-18534, or phosphate buffered saline (PBS) and TTR liver mRNA levels were assayed at 48 hrs post-treatment.

Material and Methods:

[0319] LNP09 formulation: (XTC/DSPC/Chol/PEG₂₀₀₀-C14) = 50/10/38.5/1.5 mol%; Lipid:siRNA ~ 11:1. LNP11 formulation: (MC3/DSPC/Chol/PEG₂₀₀₀-C14) = 50/10/38.5/1.5 mol%; Lipid:siRNA ~ 11:1

[0320] Tissue collection and RNA isolation: On Day 3, animals in all treatment groups were given a lethal dose of ketamine/xylazine. Blood was collected via caudal vena cava into serum separation tubes, and then allowed to clot at room temperature for approximately 30 minutes prior to centrifugation at 4°C. Serum samples were stored at -80°C until for future analysis. Liver tissues were harvested and snap frozen on dry ice. Frozen liver tissue was ground and tissue lysates were prepared for liver mRNA quantitation.

[0321] TTR mRNA Quantitation: TTR mRNA levels relative to those of GAPDH mRNA were determined in the lysates by using a branched DNA assay (QuantiGene Reagent System, Panomics, Fremont, CA). Briefly, the QuantiGene assay (Genospectra) was used to quantify mRNA levels in tissue sample lysates according to the manufacturer's instructions. The mean level of TTR mRNA was normalized to the mean level of GAPDH mRNA for each sample. Group mean values were then normalized to the mean value for the PBS treated group, to obtain the relative level of TTR mRNA expression.

Results:

[0322] As shown in FIG. 18, in contrast with PBS treated animals, LNP09-18534 and LNP11-18534 treated animals had a significant dose-dependent decrease in TTR mRNA levels in the liver, reaching maximum reduction of ~ 90% mRNA reduction for both LNP09 and LNP11 formulated groups, relative to PBC control group at 0.3 mg/kg, and a dose achieving 50% reduction (ED₅₀) of < 0.03 mg/kg for LNP11-18534 and < 0.1 mg/kg for LNP09-18534.

Conclusions

[0323] This study demonstrates that a single 15 minute IV infusion of LNP09-18534 or LNP11-18534 in Sprague-Dawley rats results in a dose-dependent reduction of liver TTR mRNA. These data demonstrate the efficacy of LNP09-18328 and LNP11-18328 in reducing endogenously expressed (wild type) TTR mRNA with ED₅₀ levels of <0.03 and <0.1 mg/kg for LNP11-18534 and LNP09-18534, respectively.

Example 17: Inhibition of TTR in humans

[0324] A human subject is treated with a dsRNA targeted to a TTR gene to inhibit expression of the TTR gene to treat a

condition.

[0325] A subject in need of treatment is selected or identified. The subject can have a liver disorder, transthyretin amyloidosis, and/or a transplanted liver.

[0326] The identification of the subject can occur in a clinical setting, or elsewhere, e.g., in the subject's home through the subject's own use of a self-testing kit.

[0327] At time zero, a suitable first dose of an anti-TTR siRNA is administered to the subject. The dsRNA is formulated as described herein. After a period of time following the first dose, e.g., 7 days, 14 days, and 21 days, the subject's condition is evaluated, e.g., by measuring liver function. This measurement can be accompanied by a measurement of TTR expression in said subject, and/or the products of the successful siRNA-targeting of TTR mRNA. Other relevant criteria can also be measured. The number and strength of doses are adjusted according to the subject's needs.

[0328] After treatment, the subject's tumor growth rate is lowered relative to the rate existing prior to the treatment, or relative to the rate measured in a similarly afflicted but untreated subject.

[0329] The present disclosure furthermore relates to the following items:

1. 1. A double-stranded ribonucleic acid (dsRNA) for inhibiting expression of transthyretin (TTR), wherein said dsRNA comprises a sense strand and an antisense strand, the antisense strand comprising a region complementary to a part of a mRNA encoding transthyretin (TTR), wherein said region of complementarity is less than 30 nucleotides in length and the antisense strand comprises 15 or more contiguous nucleotides of SEQ ID NO:170, SEQ ID NO:450, SEQ ID NO:730, or SEQ ID NO:1010.
2. 2. The dsRNA of item 1, wherein the sense strand comprises 15 or more contiguous nucleotides of SEQ ID NO:169, SEQ ID NO:449, SEQ ID NO:729, or SEQ ID NO:1009.
3. 3. The dsRNA of item 1, wherein the sense strand consists of SEQ ID NO:449 and the antisense strand consists of SEQ ID NO:450.
4. 4. The dsRNA of item 1, wherein the sense strand consists of SEQ ID NO:729 and the antisense strand consists of SEQ ID NO:730.
5. 5. The dsRNA of item 1, wherein the sense strand consists of SEQ ID NO:1009 and the antisense strand consists of SEQ ID NO:1010.
6. 6. The dsRNA of item 1, wherein the dsRNA comprises a sense strand selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16, and an antisense strand selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16.
7. 7. The dsRNA of item 1 or 2 or 6, wherein the region of complementarity is 19 nucleotides in length.
8. 8. The dsRNA of item 1 or 2 or 6, wherein the region of complementarity consists of SEQ ID NO:169.
9. 9. The dsRNA of item 1 or 2 or 6, wherein each strand of the dsRNA is 19, 20, 21, 22, 23, or 24 nucleotides in length.
10. 10. The dsRNA of item 1 or 2 or 6, wherein each strand is 21 nucleotides in length.
11. 11. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein the dsRNA does not cleave a TTR mRNA between the adenine nucleotide at position 637 of SEQ ID NO:1331 and the guanine nucleotide at position 638 of SEQ ID NO:1331.
12. 12. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein the dsRNA cleaves a TTR mRNA between the guanine nucleotide at position 636 of SEQ ID NO:1331 and the adenine nucleotide at position 637 of SEQ ID NO:1331.
13. 13. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein the dsRNA anneals to a TTR mRNA between the guanine nucleotide at position 628 of SEQ ID NO:1331 and the uracil nucleotide at position 646 of SEQ ID NO: 1331.
14. 14. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein said dsRNA comprises at least one modified nucleotide.
15. 15. The dsRNA of item 14, wherein at least one of said modified nucleotides is chosen from the group of: a 2'-O-methyl modified nucleotide, a nucleotide comprising a 5'-phosphorothioate group, and a terminal nucleotide linked to a cholesteryl derivative or dodecanoic acid bisdecylamide group.
16. 16. The dsRNA of items 14, wherein said modified nucleotide is chosen from the group of: a 2'-deoxy-2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a locked nucleotide, an abasic nucleotide, 2'-amino-modified nucleotide, 2'-alkyl-modified nucleotide, morpholino nucleotide, a phosphoramidate, and a non-natural base comprising nucleotide.
17. 17. The dsRNA of items 14, comprising at least one 2'-O-methyl modified nucleotide.
18. 18. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein the dsRNA is conjugated to a ligand.
19. 19. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein the dsRNA is formulated in a lipid formulation.
20. 20. The dsRNA of item 19, wherein the dsRNA is formulated in a LNP formulation, a LNP01 formulation, a XTC-SNALP formulation, or a SNALP formulation.
21. 21. The dsRNA of item 19, wherein the dsRNA is formulated in a XTC-SNALP formulation as follows: using 2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC) with a XTC/DPPC/Cholesterol/PEG-cDMA in a ratio of 57.1/7.1/34.4/1.4 and a lipid:siRNA ratio of about 7.

22. 22. The dsRNA of item 19, wherein the sense strand consists of SEQ ID NO:1009 and the antisense strand consists of SEQ ID NO:1010, and wherein the dsRNA is formulated in a XTC-SNALP formulation as follows: using 2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (XTC) with a XTC/DPPC/Cholesterol/PEG-cDMA in a ratio of 57.1/7.1/34.4/1.4 and a lipid:siRNA ratio of about 7.
23. 23. The dsRNA of item 19, wherein the dsRNA is formulated in a LNP09 formulation as follows: using XTC/DSPC/Chol/PEG₂₀₀₀-C14 in a ratio of 50/10/38.5/1.5 mol% and a lipid:siRNA ratio of about 11:1.
24. 24. The dsRNA of item 19, wherein the dsRNA is formulated in a LNP11 formulation as follows: using MC3/DSPC/Chol/PEG₂₀₀₀-C14 in a ratio of 50/10/38.5/1.5 mol% and a lipid:siRNA ratio of about 11:1.
25. 25. The dsRNA of item 19, wherein the dsRNA is formulated in a LNP09 formulation or a LNP11 formulation and reduces TTR mRNA levels by about 85 to 90% at a dose of 0.3mg/kg, relative to a PBS control group.
26. 26. The dsRNA of item 19, wherein the dsRNA is formulated in a LNP09 formulation or a LNP 11 formulation and reduces TTR mRNA levels by about 50% at a dose of 0.1 mg/kg, relative to a PBS control group.
27. 27. The dsRNA of item 19, wherein the dsRNA is formulated in a LNP09 formulation or a LNP11 formulation and reduces TTR protein levels in a dose-dependent manner relative to a PBS control group as measured by a western blot.
28. 28. The dsRNA of item 19, wherein the dsRNA is formulated in a SNALP formulation as follows: using DlinDMA with a DlinDMA/DPPC/Cholesterol/PEG2000-cDMA in a ratio of 57.1/7.1/34.4/1.4 and a lipid:siRNA ratio of about 7.
29. 29. The dsRNA of items 1, 2, 3, 4, 5, or 6,, wherein administration of the dsRNA to a cell results in about 95% inhibition of TTR mRNA expression as measured by a real time PCR assay, wherein the cell is a HepG2 cell or a Hep3B cell, and wherein the concentration of the dsRNA is 10nM.
30. 30. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein administration of the dsRNA to a cell results in about 74% inhibition of TTR mRNA expression as measured by a branched DNA assay, wherein the cell is a HepG2 cell or a Hep3B cell, and wherein the concentration of the dsRNA is 10nM.
31. 31. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein the dsRNA has an IC₅₀ of less than 10 pM in a HepG2 cell, and wherein the concentration of the dsRNA is 10nM.
32. 32. The dsRNA of items 1,2, 3, 4, 5, or 6, wherein the dsRNA has an ED₅₀ of about 1 mg/kg.
33. 33. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein administration of the dsRNA reduced TTR mRNA by about 80% in cynomolgus monkey liver, wherein the concentration of the dsRNA is 3 mg/kg.
34. 34. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein administration of the dsRNA does not result in immunostimulatory activity in human peripheral blood mononuclear cells (PBMCs) as measured by IFN-alpha and TNF-alpha ELISA assays.
35. 35. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein administration of the dsRNA reduces liver TTR mRNA levels by about 97% or serum TTR protein levels by about 90%, wherein the concentration of the dsRNA is 6 mg/kg.
36. 36. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein administration of the dsRNA reduces liver TTR mRNA levels and/or serum TTR protein levels up to 22 days, wherein the concentration of the dsRNA is 6 mg/kg or 3 mg/kg.
37. 37. The dsRNA of items 1, 2, 3, 4, 5, or 6, where the dsRNA suppresses serum TTR protein levels up to day 14 post-treatment when administered to a subject in need thereof at 1 mg/kg or 3 mg/kg.
38. 38. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein the dsRNA reduces TTR expression by 98.9% in a Hep3B cell at a concentration of 0.1nM as measured by real-time PCR.
39. 39. The dsRNA of items 1, 2, 3, 4, 5, or 6, wherein the dsRNA reduces TTR expression by 99.4% in a Hep3B cell at a concentration of 10nM as measured by real-time PCR.
40. 40. A double-stranded ribonucleic acid (dsRNA) for inhibiting expression of transthyretin (TTR), wherein said dsRNA comprises a sense strand and an antisense strand, the antisense strand comprising a region complementary to a part of a mRNA encoding transthyretin (TTR), wherein said region of complementarity is less than 30 nucleotides in length and wherein the dsRNA comprises a sense strand selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16, and an antisense strand selected from Tables 3A, 3B, 4, 6A, 6B, 7, and 16.
41. 41. A double-stranded ribonucleic acid (dsRNA) for inhibiting expression of transthyretin (TTR), wherein said dsRNA comprises an antisense strand comprising a region complementary to 15-30 nucleotides of nucleotides 618-648 of SEQ ID NO: 1331 and wherein said antisense strand base pairs with the guanine at position 628 of SEQ ID NO:1331.
42. 42. A cell containing the dsRNA of items 1-41.
43. 43. A vector comprising a nucleotide sequence that encodes at least one strand of the dsRNA of items 1-41.
44. 44. A cell comprising the vector of item 43.
45. 45. A pharmaceutical composition for inhibiting expression of a TTR gene comprising the dsRNA of items 1-41 and a pharmaceutically acceptable carrier.
46. 46. A pharmaceutical composition for inhibiting expression of a TTR gene comprising a dsRNA and a SNALP formulation, wherein the dsRNA comprises an antisense strand which is less than 30 nucleotides in length and comprises 15 or more contiguous nucleotides of SEQ ID NO:170, SEQ ID NO:450, SEQ ID NO:730, or SEQ ID NO:1010, and wherein the SNALP formulation comprises DlinDMA, DPPC, Cholesterol and PEG2000-cDMA in a ratio of

57.1/7.1/34.4/1.4 respectively.

47. 47. A method of inhibiting TTR expression in a cell, the method comprising:
 1. (a) contacting the cell with the dsRNA of items 1-41; and
 2. (b) maintaining the cell produced in step (a) for a time sufficient to obtain degradation of the mRNA transcript of a TTR gene, thereby inhibiting expression of the TTR gene in the cell.
48. 48. A method of treating a disorder mediated by TTR expression comprising administering to a human in need of such treatment a therapeutically effective amount of the dsRNA of items 1-41.
49. 49. The method of item 48, wherein the dsRNA is administered to the human at about 0.01, 0.1, 0.5, 1.0, 2.5, or 5.0 mg/kg.
50. 50. The method of items 48-49, wherein the dsRNA is administered to the human at about 1.0 mg/kg.
51. 51. The method of items 48-50, wherein the human has transthyretin amyloidosis.
52. 52. The method of items 48-51, wherein the human has a liver disorder.
53. 53. The method of items 48-52, wherein the human is further provided a liver transplant.
54. 54. The method of items 48-53, wherein administration of the dsRNA reduces TTR mRNA by about 80% in human liver, wherein the concentration of the dsRNA is 3 mg/kg.
55. 55. The method of items 48-54, wherein administration of the dsRNA does not result in immunostimulatory activity in the human as measured by IFN-alpha and TNF-alpha ELISA assays.
56. 56. The method of items 48-55, wherein administration of the dsRNA reduces liver TTR mRNA levels by about 97% or serum TTR protein levels by about 90%, wherein the concentration of the dsRNA is 6 mg/kg.
57. 57. The method of items 48-56, wherein administration of the dsRNA reduces liver TTR mRNA levels and/or serum TTR protein levels up to 22 days, wherein the concentration of the dsRNA is 6 mg/kg or 3 mg/kg.
58. 58. The method of items 48-57, wherein the dsRNA is formulated in a LNP09 formulation as follows: using XTC/DSPC/Chol/PEG₂₀₀₀-C14 in a ratio of 50/10/38.5/1.5 mol% and a lipid:siRNA ratio of about 11:1.
59. 59. The method of items 48-58, wherein the dsRNA is formulated in a LNP11 formulation as follows: using MC3/DSPC/Chol/PEG₂₀₀₀-C14 in a ratio of 50/10/38.5/1.5 mol% and a lipid:siRNA ratio of about 11:1.
60. 60. The method of items 48-59, wherein the dsRNA is formulated in a LNP09 formulation or a LNP 11 formulation and reduces TTR mRNA levels by about 85 to 90% at a dose of 0.3mg/kg, relative to a PBC control group.
61. 61. The method of items 48-60, wherein the dsRNA is formulated in a LNP09 formulation or a LNP 11 formulation and reduces TTR mRNA levels by about 50% at a dose of 0.1 mg/kg, relative to a PBC control group.
62. 62. The method of items 48-61, wherein the dsRNA is formulated in a LNP09 formulation or a LNP 11 formulation and reduces TTR protein levels in a dose-dependent manner relative to a PBC control group as measured by a western blot.
63. 63. The method of items 48-62, where the dsRNA suppresses serum TTR protein levels up to day 14 post-treatment when administered to human at 1 mg/kg or 3 mg/kg.
64. 64. The method of items 48-63, wherein the dsRNA is formulated in a SNALP formulation as follows: using DlinDMA, with a DlinDMA/DPPC/Cholesterol/PEG2000-cDMA in a ratio of 57.1/7.1/34.4/1.4 and a lipid:siRNA ratio of about 7.
65. 65. Use of a dsRNA for treating a disorder mediated by TTR expression comprising administering to a human in need of such treatment a therapeutically effective amount of the dsRNA of items 1-41.
66. 66. The use of item 65, wherein the dsRNA is administered to the human at about 0.01, 0.1, 0.5, 1.0, 2.5, or 5.0 mg/kg.
67. 67. The use of items 65-66, wherein the dsRNA is administered to the human at about 1.0 mg/kg.
68. 68. The use of items 65-67, wherein the human has transthyretin amyloidosis.
69. 69. The use of items 65-68, wherein the human has a liver disorder.
70. 70. The use of items 65-69, wherein the human is further provided a liver transplant.
71. 71. Use of a dsRNA for inhibiting TTR expression in a cell, the use comprising:
 1. (a) contacting the cell with the dsRNA of items 1-41; and
 2. (b) maintaining the cell produced in step (a) for a time sufficient to obtain degradation of the mRNA transcript of a TTR gene, thereby inhibiting expression of the TTR gene in the cell.

SEQUENCE LISTING

[0330]

<110> ALNYLAM PHARMACEUTICALS, INC.

<120> COMPOSITIONS AND METHODS FOR INHIBITING EXPRESSION OF TRANSTHYRETIN

<130> T1668 EP/1

<150> US 61/244,794

<151> 2009-09-22

<150> US 61/242,783

<151> 2009-09-15

<150> US 61/185,545

<151> 2009-06-09

<150> US 61/156,670

<151> 2009-03-02

<150> US 61/115,738

<151> 2008-11-18

<150> US 61/106,956

<151> 2008-10-20

<160> 1410

<170> PatentIn version 3.5

<210> 1

<211> 19

<212> RNA

<213> Homo sapiens

<400> 1

ccggugaauc caagugucc 19

<210> 2

<211> 19

<212> RNA

<213> Homo sapiens

<400> 2

ggacacuugg auucaccgg 19

<210> 3

<211> 19

<212> RNA

<213> Homo sapiens

<400> 3

acucauucuu ggcaggaug 19

<210> 4

<211> 19

<212> RNA

<213> Homo sapiens

<400> 4

cauccugcca agaaugagu 19

<210> 5

<211> 19

<212> RNA

<213> Homo sapiens

<400> 5

aaguguccuc ugaugguca 19

<210> 6

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 6
 ugaccaucag aggacacuu 19

 <210> 7
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 7
 ucauucuugg caggauggc 19

 <210> 8
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 8
 gccauccugc caagaauga 19

 <210> 9
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 9
 aaguucuaga ugcuguccg 19

 <210> 10
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 10
 cggacagcau cuagaacuu 19

 <210> 11
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 11
 guucuagaug cuguccgag 19

 <210> 12
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 12
 cucggacagc aucuagaac 19

 <210> 13
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 13
 cuagaugcug uccgaggca 19

 <210> 14

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 14
 ugccucggac agcaucuag 19

 <210> 15
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 15
 gaugcugucc gaggcaguc 19

 <210> 16
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 16
 gacugccucg gacagcauc 19

 <210> 17
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 17
 cauucuuggc aggauggcu 19

 <210> 18
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 18
 agccauccug ccaagaaug 19

 <210> 19
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 19
 ugcuguccga ggcaguccu 19

 <210> 20
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 20
 aggacugccu cggacagca 19

 <210> 21
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 21
 ccgaggcagu ccugccauc 19

 <210> 22

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 22
 gauggcagga cugccucgg 19

 <210> 23
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 23
 caguccugcc aucaaugug 19

 <210> 24
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 24
 cacauugaug gcaggacug 19

 <210> 25
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 25
 caauguggcc gugcaugug 19

 <210> 26
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 26
 cacaugcacg gccacauug 19

 <210> 27
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 27
 auguguucag aaaggcugc 19

 <210> 28
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 28
 gcagccuuuc ugaacacau 19

 <210> 29
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 29
 cagaagucca cucauucuu 19

 <210> 30

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 30
 aagaaugagu ggacuucug 19

 <210> 31
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 31
 ggcaggaugg cuucucauc 19

 <210> 32
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 32
 gaugagaagc cauccugcc 19

 <210> 33
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 33
 gagccauuug ccucuggga 19

 <210> 34
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 34
 ucccagaggc aaauggcuc 19

 <210> 35
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 35
 caggauggcu ucucaucgu 19

 <210> 36
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 36
 acgaugagaa gccauccug 19

 <210> 37
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 37
 aggauggcuu cucaucguc 19

 <210> 38

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 38
 gacgaugaga agccauccu 19

 <210> 39
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 39
 agagcugcau gggcucaca 19

 <210> 40
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 40
 ugugagccca ugcagcucu 19

 <210> 41
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 41
 gcugcauggg cucacaacu 19

 <210> 42
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 42
 aguugugagc ccaugcagc 19

 <210> 43
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 43
 ggauggcuuc ucaucgucu 19

 <210> 44
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 44
 agacgaugag aagccaucc 19

 <210> 45
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 45
 gcaugggcuc acaacugag 19

 <210> 46

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 46
 cucaguugug agcccaugc 19

 <210> 47
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 47
 augggcucac aacugagga 19

 <210> 48
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 48
 uccucaguug ugagcccau 19

 <210> 49
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 49
 ugggcucaca acugaggag 19

 <210> 50
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 50
 cuccucaguu gugagccca 19

 <210> 51
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 51
 gaggaauuug uagaaggga 19

 <210> 52
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 52
 ucccuucuc aaauuccuc 19

 <210> 53
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 53
 uuuguagaag ggauauaca 19

 <210> 54

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 54
 uguauauccc uucuacaaa 19

 <210> 55
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 55
 uuguagaagg gauauacaa 19

 <210> 56
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 56
 uuguauaucc cuucuacaa 19

 <210> 57
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 57
 uguagaaggg auauacaaa 19

 <210> 58
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 58
 uuuguauauc ccuucuaca 19

 <210> 59
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 59
 agaagggaug uacaaagug 19

 <210> 60
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 60
 cacuuuguau aucccuucu 19

 <210> 61
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 61
 aaguggaaau agacaccaa 19

 <210> 62

<211> 19
<212> RNA
<213> Homo sapiens

<400> 62
uuggugucua uuuccacuu 19

<210> 63
<211> 19
<212> RNA
<213> Homo sapiens

<400> 63
ggaaauagac accaaaucu 19

<210> 64
<211> 19
<212> RNA
<213> Homo sapiens

<400> 64
agauuuggug ucuuuuucc 19

<210> 65
<211> 19
<212> RNA
<213> Homo sapiens

<400> 65
gaaauagaca ccaaaucuu 19

<210> 66
<211> 19
<212> RNA
<213> Homo sapiens

<400> 66
aagauuuggu gucuuuuuc 19

<210> 67
<211> 19
<212> RNA
<213> Homo sapiens

<400> 67
auagacacca aaucuuacu 19

<210> 68
<211> 19
<212> RNA
<213> Homo sapiens

<400> 68
aguaagauuu ggugucua 19

<210> 69
<211> 19
<212> RNA
<213> Homo sapiens

<400> 69
uagacaccaa aucuuacug 19

<210> 70

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 70
 caguaagauu uggugucua 19

 <210> 71
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 71
 agacaccaa ucuuacugg 19

 <210> 72
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 72
 ccaguaagau uggugucu 19

 <210> 73
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 73
 uuacuggaag gcacuuggc 19

 <210> 74
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 74
 gccaaugucc uuccaguaa 19

 <210> 75
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 75
 uucucaucgu cugcuccuc 19

 <210> 76
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 76
 gaggagcaga cgaugagaa 19

 <210> 77
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 77
 ggaaggcacu uggcaucuc 19

 <210> 78

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 78
 gagaugccaa gugccuucc 19

 <210> 79
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 79
 ggcacuuggc aucucccca 19

 <210> 80
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 80
 uggggagaug ccaagugcc 19

 <210> 81
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 81
 ggcaucuccc cauuccaug 19

 <210> 82
 <211> 20
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <400> 82
 auggauggg gagaugcctt 20

 <210> 83
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 83
 gcaucucccc auuccauga 19

 <210> 84
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 84
 ucauggaaug gggagaugc 19

 <210> 85
 <211> 19
 <212> RNA
 <213> Homo sapiens

<400> 85	
caucucccca uuccaugag	19
<210> 86	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 86	
cucauggaau ggggagaug	19
<210> 87	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 87	
aucuccccau uccaugagc	19
<210> 88	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 88	
gcucauggaa uggggagau	19
<210> 89	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 89	
cuccccauuc caugagcau	19
<210> 90	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 90	
augcucaugg aauggggag	19
<210> 91	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 91	
cccauuccau gagcaugca	19
<210> 92	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 92	
ugcaugcuca uggaauagg	19
<210> 93	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 93	
ccaugagcau gcagaggug	19
<210> 94	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 94	
caccucugca ugcucaugg	19
<210> 95	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 95	
agcaugcaga ggugguauu	19
<210> 96	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 96	
aauaccaccu cugcaugcu	19
<210> 97	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 97	
caugcagagg uggauuua	19
<210> 98	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 98	
ugaauaccac cucugcaug	19
<210> 99	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 99	
augcagaggu gguauucac	19
<210> 100	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 100	
gugaauacca ccucugcau	19
<210> 101	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 101
ggugguauuc acagccaac 19

<210> 102
<211> 19
<212> RNA
<213> Homo sapiens

<400> 102
guuggcugug aauaccacc 19

<210> 103
<211> 19
<212> RNA
<213> Homo sapiens

<400> 103
gugguauuca cagccaacg 19

<210> 104
<211> 19
<212> RNA
<213> Homo sapiens

<400> 104
cguuggcugu gaauaccac 19

<210> 105
<211> 19
<212> RNA
<213> Homo sapiens

<400> 105
ugguauucac agccaacga 19

<210> 106
<211> 19
<212> RNA
<213> Homo sapiens

<400> 106
ucguuggcug ugaauacca 19

<210> 107
<211> 19
<212> RNA
<213> Homo sapiens

<400> 107
ggauuucaca gccaacgac 19

<210> 108
<211> 19
<212> RNA
<213> Homo sapiens

<400> 108
gucguuggcu gugauuacc 19

<210> 109
<211> 19
<212> RNA
<213> Homo sapiens

<400> 109	
guauucacag ccaacgacu	19
<210> 110	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 110	
agucguuggc ugugaauac	19
<210> 111	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 111	
uauucacagc caacgacuc	19
<210> 112	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 112	
gagucguugg cugugaaua	19
<210> 113	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 113	
ucacagccaa cgacuccgg	19
<210> 114	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 114	
ccggagucgu uggcuguga	19
<210> 115	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 115	
ccccgccgcu acaccauug	19
<210> 116	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 116	
caauggugua gcggcgggg	19
<210> 117	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 117	
gaaguccacu cauucuugg	19
<210> 118	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 118	
ccaagaauga guggacuuc	19
<210> 119	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 119	
cccugcugag ccccuacuc	19
<210> 120	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 120	
gaguaggggc ucagcaggg	19
<210> 121	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 121	
cugagcccu acuccauu	19
<210> 122	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 122	
aauaggagua ggggcucag	19
<210> 123	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 123	
ugagcccu cuccuauuc	19
<210> 124	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 124	
gaauaggagu aggggcuca	19
<210> 125	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 125	
ccccuacucc uauuccacc	19
<210> 126	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 126	
gguggaauag gaguagggg	19
<210> 127	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 127	
cuacuccuau uccaccacg	19
<210> 128	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 128	
cgugguggaa uaggaguag	19
<210> 129	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 129	
uacuccuauu ccaccacgg	19
<210> 130	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 130	
ccguggugga auaggagua	19
<210> 131	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 131	
acuccuauuc caccacggc	19
<210> 132	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 132	
gccguggugg aauaggagu	19
<210> 133	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 133	
uccuauucca ccacggcug	19
<210> 134	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 134	
cagccguggu ggaauagga	19
<210> 135	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 135	
uauuccacca cggcugucg	19
<210> 136	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 136	
cgacagccgu gguggaaua	19
<210> 137	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 137	
auuccaccac ggcugucgu	19
<210> 138	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 138	
acgacagccg ugguggaau	19
<210> 139	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 139	
caccacggcu gucgucacc	19
<210> 140	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 140	
ggugacgaca gccguggug	19
<210> 141	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 141	
accacggcug ucgucacca	19
<210> 142	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 142	
uggugacgac agccguggu	19
<210> 143	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 143	
ccacggcugu cgucaccaa	19
<210> 144	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 144	
uuggugacga cagccgugg	19
<210> 145	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 145	
acggcugucg ucaccaauc	19
<210> 146	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 146	
gauuggugac gacagccgu	19
<210> 147	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 147	
cggcugucgu caccaaucc	19
<210> 148	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 148	
ggauugguga cgacagccg	19
<210> 149	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 149	
cgucaccaau cccaaggaa	19
<210> 150	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 150	
uuccuuggga uuggugacg	19
<210> 151	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 151	
caauccaag gaaugaggg	19
<210> 152	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 152	
ccucauucc uugggauug	19
<210> 153	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 153	
ccugaaggac gagggauug	19
<210> 154	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 154	
ccaucccucg uccuucagg	19
<210> 155	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 155	
ggacgaggga ugggauuuc	19
<210> 156	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 156	
gaaaucccau ccucgucc	19
<210> 157	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 157
aaguccacuc auucuuggc 19

<210> 158
<211> 19
<212> RNA
<213> Homo sapiens

<400> 158
gccaaagaug aguggacuu 19

<210> 159
<211> 19
<212> RNA
<213> Homo sapiens

<400> 159
gggaauuau gaaaccaag 19

<210> 160
<211> 19
<212> RNA
<213> Homo sapiens

<400> 160
cuugguuaca ugaaauccc 19

<210> 161
<211> 19
<212> RNA
<213> Homo sapiens

<400> 161
ggauuucaug uaaccaaga 19

<210> 162
<211> 19
<212> RNA
<213> Homo sapiens

<400> 162
ucuugguuac augaaaucc 19

<210> 163
<211> 19
<212> RNA
<213> Homo sapiens

<400> 163
ucaguaacc aagaguauu 19

<210> 164
<211> 19
<212> RNA
<213> Homo sapiens

<400> 164
aaucucuug guuacauga 19

<210> 165
<211> 19
<212> RNA
<213> Homo sapiens

<400> 165	
auguaaccaa gagauuucc	19
<210> 166	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 166	
ggaauacucu ugguuacau	19
<210> 167	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 167	
uguaaccaag aguauucca	19
<210> 168	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 168	
uggaauacuc uugguuaca	19
<210> 169	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 169	
guaaccaaga guauuccau	19
<210> 170	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 170	
auggaauacu cuugguuac	19
<210> 171	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 171	
ugccuugcug gacugguau	19
<210> 172	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 172	
auaccagucc agcaaggca	19
<210> 173	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 173	
uaaagcagug uuuucaccu	19
<210> 174	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 174	
aggugaaaac acugcuuua	19
<210> 175	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 175	
gccuugcugg acugguauu	19
<210> 176	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 176	
aauaccaguc cagcaaggc	19
<210> 177	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 177	
uguuuucacc ucuaugcu	19
<210> 178	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 178	
agcauagag gugaaaaca	19
<210> 179	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 179	
guuuucaccu cauauugcua	19
<210> 180	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 180	
uagcauaua ggugaaaac	19
<210> 181	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 181	
uuuucaccuc auaugcuau	19
<210> 182	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 182	
auagcauauag aggugaaaa	19
<210> 183	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 183	
uucaccucau augcuau	19
<210> 184	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 184	
acauagcaua ugaggugaa	19
<210> 185	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 185	
caccucauau gcuauuu	19
<210> 186	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 186	
uaacauagca uauaggug	19
<210> 187	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 187	
ccuugcugga cugguuuu	19
<210> 188	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 188	
aaauaccagu ccagcaagg	19
<210> 189	
<211> 19	
<212> RNA	
<213> Homo sapiens	

<400> 189	
auaugcuaug uuagaaguc	19
<210> 190	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 190	
gacuucuaac auagcauau	19
<210> 191	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 191	
uaugcuaugu uagaagucc	19
<210> 192	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 192	
ggacuucuaa cauagcaua	19
<210> 193	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 193	
ugcuauguua gaaguccag	19
<210> 194	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 194	
cuggacuucu aacauagca	19
<210> 195	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 195	
cuugcuggac ugguauuug	19
<210> 196	
<211> 19	
<212> RNA	
<213> Homo sapiens	
<400> 196	
caaauaccag uccagcaag	19
<210> 197	
<211> 19	
<212> RNA	
<213> Homo sapiens	

```

<400> 197
aguccaggca gagacaaua      19

<210> 198
<211> 20
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 198
auugucucug ccuggacutt     20

<210> 199
<211> 19
<212> RNA
<213> Homo sapiens

<400> 199
uccaggcaga gacaauaaa      19

<210> 200
<211> 19
<212> RNA
<213> Homo sapiens

<400> 200
uuuauugucu cugccugga      19

<210> 201
<211> 19
<212> RNA
<213> Homo sapiens

<400> 201
gugaaaggca cuuuucauu      19

<210> 202
<211> 19
<212> RNA
<213> Homo sapiens

<400> 202
aaugaagaagu gccuuucac      19

<210> 203
<211> 19
<212> RNA
<213> Homo sapiens

<400> 203
uggacuggua uuugugucu      19

<210> 204
<211> 19
<212> RNA
<213> Homo sapiens

<400> 204
agacacaaau accagucca      19

<210> 205

```

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 205
 gucugaggcu ggccuacg 19

 <210> 206
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 206
 cguaggcca gccucagac 19

 <210> 207
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 207
 cugaggcugg ccuacggg 19

 <210> 208
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 208
 cccguagggc cagccucag 19

 <210> 209
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 209
 gaggcuggcc cuacgggca 19

 <210> 210
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 210
 ugcccguagg gccagccuc 19

 <210> 211
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 211
 aggcuggccc uacgggcac 19

 <210> 212
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 212
 gugcccguag ggccagccu 19

 <210> 213

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 213
 gcuggcccua cgggcaccg 19

 <210> 214
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 214
 cggugcccgu agggccagc 19

 <210> 215
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 215
 cuggcccuac gggcaccgg 19

 <210> 216
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 216
 ccggugcccg uagggccag 19

 <210> 217
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 217
 ggcccuacgg gcaccggug 19

 <210> 218
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 218
 caccggugcc cguagggcc 19

 <210> 219
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 219
 ccacucauuc uuggcagga 19

 <210> 220
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 220
 uccugccaag aaugagugg 19

 <210> 221

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 221
 ccuacgggca ccggugaau 19

 <210> 222
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 222
 auucaccggu gcccguaagg 19

 <210> 223
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 223
 cuacgggcac cggugaau 19

 <210> 224
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 224
 gauucaccgg ugcccguag 19

 <210> 225
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 225
 uacgggcacc ggugaaucc 19

 <210> 226
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 226
 ggauucaccg gugcccgua 19

 <210> 227
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 227
 acgggcaccg gugaaucca 19

 <210> 228
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 228
 uggauucacc ggugcccgu 19

 <210> 229

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 229
 gcaccgguga auccaagug 19

 <210> 230
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 230
 cacuuggauu caccggugc 19

 <210> 231
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 231
 caccggugaa uccaagugu 19

 <210> 232
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 232
 acacuuggau ucaccggug 19

 <210> 233
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 233
 uguggccaug cauguguuc 19

 <210> 234
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 234
 gaacacaugc auggccaca 19

 <210> 235
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 235
 guggccaugc auguguuca 19

 <210> 236
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 236
 ugaacacaug cauggccac 19

 <210> 237

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 237
 gccaugcaug uguucagaa 19

 <210> 238
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 238
 uucugaacac augcauggc 19

 <210> 239
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 239
 uauuccacca cggcuguca 19

 <210> 240
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 240
 ugacagccgu gguggaaua 19

 <210> 241
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 241
 gucaucacca auccaagg 19

 <210> 242
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 242
 ccuugggauu ggugaugac 19

 <210> 243
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 243
 guccucugau ggucaaagu 19

 <210> 244
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 244
 acuuugacca ucagaggac 19

 <210> 245

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 245
 gauggucaa guucuagau 19

 <210> 246
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 246
 aucuagaacu uugaccauc 19

 <210> 247
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 247
 augcuguccg aggcagucc 19

 <210> 248
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 248
 ggacugccuc ggacagcau 19

 <210> 249
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 249
 ccgugcaugu guucagaaa 19

 <210> 250
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 250
 uuucugaaca caugcacgg 19

 <210> 251
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 251
 agucuggaga gcugcaugg 19

 <210> 252
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 252
 ccaugcagcu cuccagacu 19

 <210> 253

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 253
 caugggcuca caacugagg 19

 <210> 254
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 254
 ccucaguugu gagcccaug 19

 <210> 255
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 255
 ucucaucguc ugcuccucc 19

 <210> 256
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 256
 ggaggagcag acgaugaga 19

 <210> 257
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 257
 ccccauucca ugagcaugc 19

 <210> 258
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 258
 gcaugcucau ggaauagggg 19

 <210> 259
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 259
 gccccuacuc cuauuccac 19

 <210> 260
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 260
 guggaauagg aguaggggc 19

 <210> 261

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 261
 cuauuccacc acggcuguc 19

 <210> 262
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 262
 gacagccgug guggaaauag 19

 <210> 263
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 263
 cacggcuguc gucaccaau 19

 <210> 264
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 264
 auuggugacg acagccgug 19

 <210> 265
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 265
 aggacgaggg augggauuu 19

 <210> 266
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 266
 aaaucccauc ccucguccu 19

 <210> 267
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 267
 ucaccucaua ugcuauguu 19

 <210> 268
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 268
 aacauagcau augagguga 19

 <210> 269

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 269
 ccucauauagc uauguuaga 19

 <210> 270
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 270
 ucuaacauag cauauagg 19

 <210> 271
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 271
 auguuagaag uccaggcag 19

 <210> 272
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 272
 cugccuggac uucuaacau 19

 <210> 273
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 273
 ucugaggcug gcccuacgg 19

 <210> 274
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 274
 ccguagggcc agccucaga 19

 <210> 275
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 275
 ggcccuacgg gcaccggug 19

 <210> 276
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 276
 caccggugcc cguagggcc 19

 <210> 277

<211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 277
 gggcaccggu gaauccaag 19

 <210> 278
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 278
 cuuggauuca ccggugccc 19

 <210> 279
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 279
 ccaugcaugu guucagaaa 19

 <210> 280
 <211> 19
 <212> RNA
 <213> Homo sapiens

 <400> 280
 uuucugaaca caugcaugg 19

 <210> 281
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

 <400> 281
 ccggugaauca caaguguccn n 21

 <210> 282
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

 <400> 282

ggacacuugg auucacggn n 21

<210> 283

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 283

acucauucuu ggcaggaugn n 21

<210> 284

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 284

cauccugcca agaaugagun n 21

<210> 285

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 285

aaguguccuc ugauggucan n 21

<210> 286

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 286
ugaccaucag aggacacuun n 21

<210> 287
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 287
ucauucuugg caggauggcn n 21

<210> 288
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 288
gccauccugc caagaugan n 21

<210> 289
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 289
aaguucuaga ugcuguccgn n 21

<210> 290
<211> 21
<212> DNA
<213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 290
 cggacagcau cuagaacuun n 21

<210> 291
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 291
 guucuagaug cuguccgagn n 21

<210> 292
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 292
 cucggacagc aucuagaacn n 21

<210> 293
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 293
 cuagaugcug uccgaggcan n 21

<210> 294
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 294
ugccucggac agcaucuagn n 21

<210> 295
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 295
gaugcugucc gaggcagucn n 21

<210> 296
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 296
gacugccucg gacagcaucn n 21

<210> 297
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 297

cauucuuggc aggauggcun n 21

<210> 298

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 298

agccauccug ccaagaaugn n 21

<210> 299

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 299

ugcuguccga ggcaguccun n 21

<210> 300

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 300

aggacugccu cggacagcan n 21

<210> 301

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 301

ccgaggcagu ccugccaucn n 21

<210> 302

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 302

gauggcagga cugccucggn n 21

<210> 303

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 303

caguccugcc aucaaugugn n 21

<210> 304

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 304

cacauugaug gcaggacugn n 21

<210> 305

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 305
caauguggcc gugcaugugn n 21

<210> 306
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 306
cacaugcacg gccacauugn n 21

<210> 307
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 307
auguguucag aaaggcugcn n 21

<210> 308
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 308
gcagccuuuc ugaacacaun n 21

<210> 309
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 309
cagaagucca cucauucuun n 21

<210> 310
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 310
aagaagagu ggacuucugn n 21

<210> 311
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 311
ggcaggaugg cuucucauch n 21

<210> 312
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 312
gaugagaagc cauccugccn n 21

<210> 313
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 313
gagccauuug ccucugggan n 21

<210> 314
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 314
ucccagaggc aaauggcucn n 21

<210> 315
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 315
caggauggcu ucucaucgun n 21

<210> 316
<211> 21
<212> DNA
<213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 316
 acgaugagaa gccauccugn n 21

<210> 317
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 317
 aggauggcuu cucaucguch n 21

<210> 318
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 318
 gacgaugaga agccauccun n 21

<210> 319
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 319
 agagcugcau gggcucacan n 21

<210> 320
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 320
ugugagccca ugcagcucun n 21

<210> 321
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 321
gcugcauggg cucacaacun n 21

<210> 322
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 322
aguugugagc ccaugcagcn n 21

<210> 323
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 323

ggauggcuuc ucaucgucun n 21

<210> 324

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 324

agacgaugag aagccaucn n 21

<210> 325

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 325

gcaugggcuc acaacugagn n 21

<210> 326

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 326

cucaguugug agcccaugcn n 21

<210> 327

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 327

augggcucac aacugaggan n 21

<210> 328

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 328

uccucaguug ugagccaun n 21

<210> 329

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 329

ugggcucaca acugaggagn n 21

<210> 330

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 330

cuccucaguu gugagccan n 21

<210> 331

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 331
gaggaauuug uagaaggan n 21

<210> 332
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 332
ucccuucucac aaauuccucn n 21

<210> 333
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 333
uuuguagaag ggauauacan n 21

<210> 334
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 334
uguauauccc uucuacaaan n 21

<210> 335
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20) .. (21)
<223> a, c, t, g, unknown or other

<400> 335
uuguagaagg gauauacaan n 21

<210> 336
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 336
uuguauaucc cuucuacaan n 21

<210> 337
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 337
uguagaaggg auauacaaan n 21

<210> 338
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 338
uuuguauauc ccuucuacan n 21

<210> 339
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 339
agaagggaau uacaaagugn n 21

<210> 340
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 340
cacuuuguau aucccuucun n 21

<210> 341
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 341
aaguggaaau agacaccaan n 21

<210> 342
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 342
uuggugucua uuuccacuun n 21

<210> 343
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 343
ggaaauagac accaaaucun n 21

<210> 344
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 344
agauuuggug ucuauuuccn n 21

<210> 345
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 345
gaaauagaca ccaaaucun n 21

<210> 346
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 346
aagauuuggu gucuauuucn n 21

<210> 347
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 347
auagacacca aaucuuacn n 21

<210> 348
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 348
aguaagauuu ggugucuaun n 21

<210> 349
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 349

uagacaccaa aucuuacugn n 21

<210> 350

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 350

caguaagauu uggugucuan n 21

<210> 351

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 351

agacaccaaa ucuuacugn n 21

<210> 352

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 352

ccaguaagau uggugucun n 21

<210> 353

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 353

uuacuggaag gcacuuggcn n 21

<210> 354

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 354

gccaaugucc uaccaguaan n 21

<210> 355

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 355

uucucaucgu cugcuccucn n 21

<210> 356

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 356

gaggagcaga cgaugagaan n 21

<210> 357

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 357
ggaaggcacu uggcaucucn n 21

<210> 358
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 358
gagaugccaa gugccuuccn n 21

<210> 359
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 359
ggcacuuggc aucuccccan n 21

<210> 360
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 360
ugggggagaug ccaagugccn n 21

<210> 361
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 361
ggcaucuccc cauuccaugn n 21

<210> 362
<211> 22
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (21) .. (22)
<223> a, c, t, g, unknown or other

<400> 362
auggaauggg gagaucctt nn 22

<210> 363
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 363
gcaucucccc auuccaugan n 21

<210> 364
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 364
ucauggaag gggagaugcn n 21

<210> 365
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 365
caucucccca uccaugagn n 21

<210> 366
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 366
cucauggaau ggggagaugn n 21

<210> 367
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 367
aucuccccau uccaugagcn n 21

<210> 368
<211> 21
<212> DNA
<213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 368
 gcucauggaa uggggagaun n 21

<210> 369
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 369
 cucccauuc caugagcaun n 21

<210> 370
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 370
 augcucaugg aauggggagn n 21

<210> 371
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 371
 ccgauuccau gagcaugcan n 21

<210> 372
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 372
ugcaugcuca uggaugggn n 21

<210> 373
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 373
ccaugagcau gcagaggugn n 21

<210> 374
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 374
caccucugca ugcucauggn n 21

<210> 375
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 375

agcaugcaga ggugguauun n 21

<210> 376

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 376

aaauaccaccu cugcaugcun n 21

<210> 377

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 377

caugcagagg ugguauucan n 21

<210> 378

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 378

ugaauaccac cucugcaugn n 21

<210> 379

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 379

augcagaggu gguauucacn n 21

<210> 380

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20) .. (21)

<223> a, c, t, g, unknown or other

<400> 380

gugaauacca ccucugcaun n 21

<210> 381

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 381

ggugguauuc acagccaacn n 21

<210> 382

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 382

guuggcugug aaauaccacn n 21

<210> 383

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 383
gugguauuca cagccaacgn n 21

<210> 384
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 384
cguuggcugu gaauaccacn n 21

<210> 385
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 385
ugguauucac agccaacgan n 21

<210> 386
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 386
ucguuggcug ugaauaccan n 21

<210> 387
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 387
gguaauucaca gccaacgacn n 21

<210> 388
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 388
gucguuggcu gugaauaccn n 21

<210> 389
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 389
guaauucacag ccaacgacun n 21

<210> 390
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 390
agucguuggc ugugaauacn n 21

<210> 391
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 391
uauucacagc caacgacucn n 21

<210> 392
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 392
gagucguugg cugugaaun n 21

<210> 393
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 393
ucacagccaa cgacuccggn n 21

<210> 394
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 394
ccggagucgu uggcugugan n 21

<210> 395
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 395
ccccgccgu acaccuugn n 21

<210> 396
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 396
caauggugua gcggcggggn n 21

<210> 397
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 397
gaaguccacu cauucuuggn n 21

<210> 398
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20) .. (21)
<223> a, c, t, g, unknown or other

<400> 398
ccaagaauga guggacuucn n 21

<210> 399
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 399
cccugcugag ccccuacucn n 21

<210> 400
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 400
gaguaggggc ucagcagggn n 21

<210> 401
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 401

cugagccccc acuccuauun n 21

<210> 402

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20) .. (21)

<223> a, c, t, g, unknown or other

<400> 402

aauaggagua ggggcucagn n 21

<210> 403

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 403

ugagcccua cuccuauucn n 21

<210> 404

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 404

gaauaggagu aggggcucan n 21

<210> 405

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 405

ccccuacucc uauuccacn n 21

<210> 406

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 406

gguggaauag gaguaggggn n 21

<210> 407

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 407

cuacuccuau uccaccacgn n 21

<210> 408

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 408

cgugguggaa uaggaguagn n 21

<210> 409

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 409
uacuccuauu ccaccacggn n 21

<210> 410
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 410
ccguggugga auaggaguan n 21

<210> 411
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 411
acuccuauuc caccacgcn n 21

<210> 412
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 412
gccguggugg aauaggagun n 21

<210> 413
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 413
uccuauucca ccacggcugn n 21

<210> 414
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 414
cagccguggu ggaauaggan n 21

<210> 415
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 415
uauuccacca cggcugucgn n 21

<210> 416
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 416
cgacagccgu gguggaauan n 21

<210> 417
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 417
auuccaccac ggcugucgun n 21

<210> 418
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 418
acgacagccg ugguggaun n 21

<210> 419
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 419
caccacggcu gucgucacn n 21

<210> 420
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 420
ggugacgaca gccguggugn n 21

<210> 421
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 421
accacggcug ucgucaccan n 21

<210> 422
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 422
uggugacgac agccguggun n 21

<210> 423
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 423
ccacggcugu cgucaccaan n 21

<210> 424
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 424
uuggugacga cagccguggn n 21

<210> 425
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 425
acggcugucg ucaccaaucn n 21

<210> 426
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 426
gauuggugac gacagccgun n 21

<210> 427
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 427

cggcugucgu caccaauccn n 21

<210> 428

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 428

ggauugguga cgacagccgn n 21

<210> 429

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 429

cgucaccaau cccaaggaan n 21

<210> 430

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 430

uuccuuggga uuggugacgn n 21

<210> 431

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 431

caaucccaag gaaugagggn n 21

<210> 432

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 432

ccucauucc uugggauugn n 21

<210> 433

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 433

ccugaaggac gagggauugn n 21

<210> 434

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 434

ccaucuccug uccuucaggn n 21

<210> 435

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 435
ggacgaggga ugggauuucn n 21

<210> 436
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 436
gaaaucccau ccucgucn n 21

<210> 437
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 437
aaguccacuc auucuuggcn n 21

<210> 438
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 438
gccagaag aguggacuun n 21

<210> 439
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 439
gggaauuau gaaaccaagn n 21

<210> 440
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 440
cuugguuaca ugaaucccn n 21

<210> 441
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 441
ggauuucaug uaaccaagan n 21

<210> 442
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 442
ucuugguuac augaaaucn n 21

<210> 443
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20) .. (21)
<223> a, c, t, g, unknown or other

<400> 443
ucauguaacc aagaguauun n 21

<210> 444
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 444
aauacucuug guuacaugan n 21

<210> 445
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 445
auguaaccaa gaguauuccn n 21

<210> 446
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 446
ggaaauacucu ugguuacaun n 21

<210> 447
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 447
uguaaccaag aguauuccan n 21

<210> 448
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 448
uggaauacuc uugguuacan n 21

<210> 449
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 449
guaaccaaga guauuccaun n 21

<210> 450
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 450
auggaaauacu cuugguuacn n 21

<210> 451
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 451
ugccuugcug gacugguaun n 21

<210> 452
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 452
auaccagucc agcaaggcan n 21

<210> 453
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 453

uaaagcagug uuuucaccun n 21

<210> 454

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 454

aggugaaaac acugcuuan n 21

<210> 455

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 455

gccuugcugg acugguauun n 21

<210> 456

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 456

aauaccaguc cagcaaggcn n 21

<210> 457

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 457

uguuuucacc ucauauaucun n 21

<210> 458

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 458

agcauauagag gugaaaacan n 21

<210> 459

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 459

guuuuacaccu cauauaucuan n 21

<210> 460

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 460

uagcauauaga ggugaaaacn n 21

<210> 461

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20) .. (21)
<223> a, c, t, g, unknown or other

<400> 461
uuuucaccuc auaugcuaun n 21

<210> 462
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 462
auagcauau aggugaaaa n 21

<210> 463
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 463
uucaccucau augcuaugun n 21

<210> 464
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 464
 acauagcaua ugaggugaan n 21

 <210> 465
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20) .. (21)
 <223> a, c, t, g, unknown or other

 <400> 465
 caccucauau gcuauuguau n 21

 <210> 466
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

 <400> 466
 uaacauagca uaugaggugn n 21

 <210> 467
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

 <400> 467
 ccuugcugga cugguauuun n 21

 <210> 468
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 468
aaaauaccagu ccagcaaggn n 21

<210> 469
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 469
auaugcuaug uuagaaguch n 21

<210> 470
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 470
gacuucuaac auagcauun n 21

<210> 471
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 471
uaugcuaugu uagaaguccn n 21

<210> 472
<211> 21
<212> DNA
<213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 472
 ggacuucuaa cauagcauan n 21

<210> 473
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 473
 ugcuauguua gaaguccagn n 21

<210> 474
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 474
 cuggacuucu aacauagcan n 21

<210> 475
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 475
 cuugcuggac ugguauuugn n 21

<210> 476
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

 <400> 476
 caaauaccag uccagcaagn n 21

 <210> 477
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

 <400> 477
 aguccaggca gagacaauan n 21

 <210> 478
 <211> 22
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (21) .. (22)
 <223> a, c, t, g, unknown or other

 <400> 478
 auugucucug ccuggacutt nn 22

 <210> 479
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 479

uccaggcaga gacaauaaan n 21

<210> 480

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 480

uuuauugucu cugccuggan n 21

<210> 481

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 481

gugaaaggca cuuuucauun n 21

<210> 482

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 482

aaugaaaagu gccuuucacn n 21

<210> 483

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 483

uggacuggua uuugugucun n 21

<210> 484

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 484

agacacaaau accaguccan n 21

<210> 485

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 485

gucugaggcu ggccuacgn n 21

<210> 486

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 486

cguagggcca gccucagacn n 21

<210> 487

<211> 21

<212> DNA
 <213> Homo sapiens
 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 487
 cugaggcugg ccuacgggn n 21

<210> 488
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 488
 cccguagggc cagccucagn n 21

<210> 489
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 489
 gaggcuggcc cuacgggcan n 21

<210> 490
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 490
ugcccgguagg gccagccucn n 21

<210> 491
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 491
aggcuggccc uacgggcacn n 21

<210> 492
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 492
gugcccgua ggcagccun n 21

<210> 493
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 493
gcuggccua cgggcaccgn n 21

<210> 494
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 494
cggugcccg agggccagcn n 21

<210> 495
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 495
cuggcccuac gggcaccggn n 21

<210> 496
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 496
ccggugcccg uaggccagn n 21

<210> 497
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 497
ggcccuacgg gcaccggugn n 21

<210> 498
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 498
caccggugcc cguagggccn n 21

<210> 499
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 499
ccacucauuc uuggcaggan n 21

<210> 500
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 500
uccugccaag aaugaguggn n 21

<210> 501
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 501
ccuacgggca ccggugaaun n 21

<210> 502
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 502
auucaccggu gcccguaagn n 21

<210> 503
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 503
cuacgggcac cggugaau n 21

<210> 504
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 504
gauucaccgg ugcccguagn n 21

<210> 505
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 505

uacgggcacc ggugaauccn n 21

<210> 506

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20) .. (21)

<223> a, c, t, g, unknown or other

<400> 506

ggauucaccg gugcccguan n 21

<210> 507

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 507

acgggcaccg gugaauccan n 21

<210> 508

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 508

uggauucacc ggugcccgun n 21

<210> 509

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 509

gcaccgguga auccaagugn n 21

<210> 510

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 510

cacuuggauu caccggugcn n 21

<210> 511

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 511

caccggugaa uccaagugun n 21

<210> 512

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 512

acacuuggau ucaccggugn n 21

<210> 513

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 513
uguggccaug cauguguuch n 21

<210> 514
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 514
gaacacaugc auggccacan n 21

<210> 515
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 515
guggccaugc auguguucan n 21

<210> 516
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 516
ugaacacaug cauggccacn n 21

<210> 517
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 517
gccaugcaug uguucagaan n 21

<210> 518
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 518
uucugaacac augcauggcn n 21

<210> 519
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 519
uauuccacca cggcugucan n 21

<210> 520
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 520
ugacagccgu gguggaauan n 21

<210> 521
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 521
gucaucacca aucccaaggn n 21

<210> 522
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 522
ccuugggauu ggugaugacn n 21

<210> 523
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 523
guccucugau ggucaaagun n 21

<210> 524
<211> 21
<212> DNA
<213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20) .. (21)
 <223> a, c, t, g, unknown or other

<400> 524
 acuuugacca ucagaggacn n 21

<210> 525
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 525
 gauggucaaaa guucuagaun n 21

<210> 526
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 526
 aucuagaacu uugaccaucn n 21

<210> 527
 <211> 21
 <212> DNA
 <213> Homo sapiens

<220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

<400> 527
 augcuguccg aggaguccn n 21

<210> 528
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20) .. (21)
<223> a, c, t, g, unknown or other

<400> 528
ggacugccuc ggacagcaun n 21

<210> 529
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 529
ccgugcaugu guucagaaan n 21

<210> 530
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 530
uuucugaaca caugcacggn n 21

<210> 531
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 531

agucuggaga gcugcauggn n 21

<210> 532

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 532

ccaugcagcu caccagacun n 21

<210> 533

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 533

caugggcuca caacugaggn n 21

<210> 534

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 534

ccucaguugu gagcccaugn n 21

<210> 535

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 535

ucucaucguc ugcuccuccn n 21

<210> 536

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 536

ggaggagcag acgaugagan n 21

<210> 537

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 537

ccccauucca ugagcaugcn n 21

<210> 538

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 538

gcaugcucau ggaugggggn n 21

<210> 539

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 539
gccccuacuc cuauuccacn n 21

<210> 540
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 540
guggaauagg aguaggggcn n 21

<210> 541
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 541
cuauuccacc acggcugucn n 21

<210> 542
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 542
 gacagccgug guggaauagn n 21

 <210> 543
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

 <400> 543
 cacggcuguc gucaccaun n 21

 <210> 544
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

 <400> 544
 auuggugacg acagccgugn n 21

 <210> 545
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

 <220>
 <221> modified_base
 <222> (20)..(21)
 <223> a, c, t, g, unknown or other

 <400> 545
 aggacgaggg augggauun n 21

 <210> 546
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 546
aaaucccauc ccucguccun n 21

<210> 547
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 547
ucaccucaua ugcuauguun n 21

<210> 548
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 548
aacauagcau augaggugan n 21

<210> 549
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 549
ccucauagc uauguuagan n 21

<210> 550
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 550
ucuaacauag cauugaggn n 21

<210> 551
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 551
auguuagaag uccaggcagn n 21

<210> 552
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 552
cugccuggac uucuaacaun n 21

<210> 553
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 553
ucugaggcug gcccuacggn n 21

<210> 554
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 554
ccguagggcc agccucagan n 21

<210> 555
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 555
ggcccuacgg gcaccggugn n 21

<210> 556
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 556
caccggugcc cguagggccn n 21

<210> 557
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 557

gggcaccggu gaauccaagn n 21

<210> 558

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 558

cuuggauuca ccggugcccn n 21

<210> 559

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 559

ccaugcaugu guucagaaan n 21

<210> 560

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 560

uuucugaaca caugcauggn n 21

<210> 561

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 561

ccggugaauc caagugucct t 21

<210> 562

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 562

ggacacuugg auucaccggt t 21

<210> 563

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 563

acucauucuu ggcaggaugt t 21

<210> 564

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 564

cauccugcca agaaugagut t 21

<210> 565

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 565

aaguguccuc ugauggucat t 21

<210> 566

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 566

ugaccaucag aggacacuat t 21

<210> 567
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 567
ucauucuugg caggauggct t 21

<210> 568
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 568
gccauccugc caagaaugat t 21

<210> 569
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 569
aaguucuaga ugcuguccgt t 21

<210> 570
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 570
cggacagcau cuagaacuut t 21

<210> 571
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 571
guucuagaug cuguccgagt t 21

<210> 572
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 572

cucggacagc aucuagaact t 21

<210> 573

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 573

cuagaugcug uccgagcat t 21

<210> 574

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 574

ugccucggac agcaucuagt t 21

<210> 575

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 575

gaugcugucc gaggcaguct t 21

<210> 576

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 576

gacugccucg gacagcauct t 21

<210> 577

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 577

cauucuuggc aggauggcut t 21

<210> 578

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 578

agccauccug ccaagaugt t 21

<210> 579

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 579

ugcuguccga ggcaguccut t 21

<210> 580

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 580

aggacugccu cggacagcat t 21

<210> 581

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 581

ccgaggcagu ccugccauct t 21

<210> 582

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 582

gauggcagga cugccucgt t 21

<210> 583
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 583
caguccugcc aucaaugugt t 21

<210> 584
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 584
cacauugaug gcaggacugt t 21

<210> 585
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 585
caauguggcc gugcaugugt t 21

<210> 586
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 586
cacaugcacg gccacauugt t 21

<210> 587
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 587
auguguucag aaaggcugct t 21

<210> 588
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 588

gcagccuuuc ugaacacaut t 21

<210> 589

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 589

cagaagucca cucauuuut t 21

<210> 590

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 590

aagaaugagu ggacuucgt t 21

<210> 591

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 591

ggcaggaugg cuucauct t 21

<210> 592

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 592

gaugagaagc cauccugcct t 21

<210> 593

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 593

gagccauuug ccucugggat t 21

<210> 594

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 594

ucccagaggc aaauuggcuct t 21

<210> 595

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 595

caggauggcu ucucaucgut t 21

<210> 596

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 596

acgaugagaa gccauccugt t 21

<210> 597

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 597

aggauggcuu cucaucguct t 21

<210> 598

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 598

gacgaugaga agccaucut t 21

<210> 599
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 599
agagcugcau gggcucacat t 21

<210> 600
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 600
ugugagccca ugcagcucut t 21

<210> 601
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 601
gcugcauggg cucacaacut t 21

<210> 602
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 602
aguugugagc ccaugcagct t 21

<210> 603
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 603
ggauggcuuc ucaucgucut t 21

<210> 604
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 604
agacgaugag aagccaucct t 21

<210> 605
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 605
gcaugggcuc acaacugagt t 21

<210> 606
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 606
cucaguugug agcccaugct t 21

<210> 607
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 607
augggcucac aacugaggat t 21

<210> 608
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 608
uccucaguug ugagcccaut t 21

<210> 609
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 609

ugggcucaca acugaggagt t 21

<210> 610

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 610

cuccucaguu gugagcccat t 21

<210> 611

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 611

gaggaauuug uagaaggat t 21

<210> 612

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 612

ucccuucucac aaauuccuct t 21

<210> 613

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 613

uuuguagaag ggauauacat t 21

<210> 614

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 614

uguauauccc uucuacaaat t 21

<210> 615
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 615
uuguagaagg gauauacaat t 21

<210> 616
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 616
uuguauaucc cuucuacaat t 21

<210> 617
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 617
uguagaaggg auauacaaat t 21

<210> 618
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 618
uuuguauauc ccuucuacat t 21

<210> 619
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 619
agaagggaaua uacaaagugt t 21

<210> 620
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 620
cacuuuguau aucccuucut t 21

<210> 621
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 621
aaguggaaau agacaccaat t 21

<210> 622
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 622
uuggugucua uuuccacuut t 21

<210> 623
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 623
ggaaauagac accaaauucut t 21

<210> 624
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 624
agauuuggug ucuauuucct t 21

<210> 625
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 625

gaaauagaca ccaaaucuut t 21

<210> 626

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 626

aagauuuggu gucuauuuct t 21

<210> 627

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 627

auagacacca aaucuuacut t 21

<210> 628

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 628

aguaagauuu ggugucuat t 21

<210> 629

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 629

uagacaccaa aucuuacugt t 21

<210> 630

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 630

caguaagauu uggugucuat t 21

<210> 631
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 631
agacaccaaa ucuuacuggt t 21

<210> 632
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 632
ccaguaagau uuggugucut t 21

<210> 633
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 633
uuacuggaag gcacuuggct t 21

<210> 634
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 634
gccaagugcc uuccaguaat t 21

<210> 635
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 635
uucucaucgu cugcuccuct t 21

<210> 636
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 636

gaggagcaga cgaugagaat t 21

<210> 637

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 637

ggaaggcacu uggaucuct t 21

<210> 638

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 638

gagaugccaa gugccuuct t 21

<210> 639

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 639

ggcacuuggc aucucccat t 21

<210> 640

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 640

uggggagaug ccaagugcct t 21

<210> 641

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 641

ggcaucuccc cauuccaagt t 21

<210> 642

<211> 22

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 642

auggaauagg gagaugcctt tt 22

<210> 643

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 643

gcaucucccc auuccaagt t 21

<210> 644

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 644

ucauggaau gggagaugt t 21

<210> 645

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 645

caucucccca uuccaagagt t 21

<210> 646

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 646

cucauggaau gggagaugt t 21

<210> 647
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 647
aucuccccau uccaugagct t 21

<210> 648
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 648
gcucauggaa uggggagaut t 21

<210> 649
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 649
cuccccauuc caugagcaut t 21

<210> 650
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 650
augcucaugg aauggggagt t 21

<210> 651
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 651
cccauuccau gagcaugcat t 21

<210> 652
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 652

ugcaugcuca uggaauagggt t 21

<210> 653

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 653

ccaugagcau gcagaggugt t 21

<210> 654

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 654

caccucugca ugcucauggt t 21

<210> 655

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 655

agcaugcaga ggugguauut t 21

<210> 656

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 656

aauaccaccu cugcaugcut t 21

<210> 657

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 657

caugcagagg uggauuuc t 21

<210> 658

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 658

ugaauaccac cucugcaug t 21

<210> 659

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 659

augcagaggu ggauuuc t 21

<210> 660

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 660

gugaauacca ccucugcaut t 21

<210> 661

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 661

ggugguauuc acagccaact t 21

<210> 662

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 662

guuggcugug aaauccacct t 21

<210> 663
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 663
gugguauuca cagccaacgt t 21

<210> 664
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 664
cguuggcugu gaauaccact t 21

<210> 665
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 665
ugguauucac agccaacgat t 21

<210> 666
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 666
ucguuggcug ugaauaccat t 21

<210> 667
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 667
ggauuucaca gccaacgact t 21

<210> 668
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 668

gucguuggcu gugaaucct t 21

<210> 669

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 669

guauucacag ccaacgacut t 21

<210> 670

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 670

agucguuggc ugugaauact t 21

<210> 671

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 671

uauucacagc caacgacut t 21

<210> 672

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 672

gagucguugg cugugaaat t 21

<210> 673

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 673

ucacagccaa cgacuccggt t 21

<210> 674

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 674

ccggagucgu uggcugugat t 21

<210> 675

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 675

ccccgccgc acaccuugt t 21

<210> 676

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 676

caauggugua gcggcggggt t 21

<210> 677

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 677

gaaguccacu cauucuugt t 21

<210> 678

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 678

ccaagaauga guggacuuct t 21

<210> 679
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 679
cccugcugag ccccuacuct t 21

<210> 680
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 680
gaguaggggc ucagcaggg t 21

<210> 681
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 681
cugagcccu acuccuau t 21

<210> 682
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 682
aauaggagua ggggcucagt t 21

<210> 683
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 683
ugagcccu cuccuau t 21

<210> 684
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 684

gaauaggagu aggggcucac t 21

<210> 685

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 685

ccccuacucc uauuccacac t 21

<210> 686

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 686

gguggaauag gaguaggggt t 21

<210> 687

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 687

cuacuccuau uccaccacgt t 21

<210> 688

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 688

cgugguggaa uaggaguagt t 21

<210> 689

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 689

uacuccuauu ccaccacggt t 21

<210> 690

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 690

ccguggugga auaggagat t 21

<210> 691

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 691

acuccuauuc caccacggt t 21

<210> 692

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 692

gccguggugg aauaggagut t 21

<210> 693

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 693

uccuauucca ccacgcgct t 21

<210> 694

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 694

cagccguggu ggaauaggat t 21

<210> 695
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 695
uauuccacca cggcugucgt t 21

<210> 696
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 696
cgacagccgu gguggaauat t 21

<210> 697
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 697
auuccaccac ggcugucgut t 21

<210> 698
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 698
acgacagccg ugguggaaut t 21

<210> 699
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 699
caccacggcu gucgucacct t 21

<210> 700
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 700
ggugacgaca gccguggugt t 21

<210> 701
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 701
accacggcug ucgucaccat t 21

<210> 702
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 702
uggugacgac agccguggut t 21

<210> 703
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 703
ccacggcugu cgucaccaat t 21

<210> 704
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 704
uuggugacga cagccguggt t 21

<210> 705
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 705

acggcugucg ucaccauact t 21

<210> 706

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 706

gauuggugac gacagccgut t 21

<210> 707

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 707

cggcugucgu caccaaucct t 21

<210> 708

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 708

ggauugguga cgacagccgt t 21

<210> 709

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 709

cgucaccaau cccaaggaat t 21

<210> 710

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 710

uuccuuggga uuggugacgt t 21

<210> 711
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 711
caauccaag gaugagggt t 21

<210> 712
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 712
ccucauucc ugggauugt t 21

<210> 713
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 713
ccugaaggac gaggauggt t 21

<210> 714
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 714
ccaucccug uccuucaggt t 21

<210> 715
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 715
ggacgaggga ugggauuuct t 21

<210> 716
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 716
gaaaucccau ccucgucct t 21

<210> 717
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 717
aaguccacuc auucuuggct t 21

<210> 718
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 718
gccagaug aguggacuut t 21

<210> 719
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 719
gggauucau gaaaccaagt t 21

<210> 720
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 720
cuugguaca ugaaaucct t 21

<210> 721
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 721

ggauuucaug uaaccaagat t 21

<210> 722

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 722

ucuugguuac augaaucct t 21

<210> 723

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 723

ucauguaacc aagaguauut t 21

<210> 724

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 724

aaucucuug guuacaugat t 21

<210> 725

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 725

auguaaccaa gagauucct t 21

<210> 726

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 726

ggauuacucu ugguacaut t 21

<210> 727
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 727
uguaaccaag aguauccat t 21

<210> 728
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 728
uggaauacuc uugguacat t 21

<210> 729
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 729
guaaccaaga guauuccat t 21

<210> 730
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 730
auggaauacu cuugguact t 21

<210> 731
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 731
ugccuugcug gacugguaut t 21

<210> 732
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 732
auaccagucc agcaaggcat t 21

<210> 733
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 733
uaaagcagug uuuucaccut t 21

<210> 734
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 734
aggugaaaac acugcuuat t 21

<210> 735
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 735
gccuugcugg acugguauut t 21

<210> 736
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 736
aauaccaguc cagcaaggct t 21

<210> 737
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 737

uguuuucacc ucauauaucut t 21

<210> 738

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 738

agcauauagag gugaaaacat t 21

<210> 739

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 739

guuuucaccu cauauaucuat t 21

<210> 740

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 740

uagcauauaga ggugaaaact t 21

<210> 741

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 741

uuuucaccuc auauaucuat t 21

<210> 742

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 742

auagcauauag aggugaaaat t 21

<210> 743
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 743
uucaccucau augcuaugut t 21

<210> 744
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 744
acauagcaua ugaggugaat t 21

<210> 745
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 745
caccucauau gcuauguuat t 21

<210> 746
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 746
uaacauagca uaugaggugt t 21

<210> 747
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 747
ccuugcugga cugguauuut t 21

<210> 748
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 748
aaauaccagu ccagcaaggt t 21

<210> 749
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 749
auaugcuaug uuagaaguct t 21

<210> 750
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 750
gacuucuaac auagcauat t 21

<210> 751
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 751
uaugcuaugu uagaagucct t 21

<210> 752
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 752
ggacuucuaa cauagcauat t 21

<210> 753
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 753

ugcuauguua gaaguccagt t 21

<210> 754

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 754

cuggacuucu aacauagcat t 21

<210> 755

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 755

cuugcuggac ugguauuugt t 21

<210> 756

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 756

caaauaccag uccagcaagt t 21

<210> 757

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 757

aguccaggca gagacaauat t 21

<210> 758

<211> 22

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 758

auugucucug ccuggacutt tt 22

<210> 759
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 759
uccaggcaga gacaauaaat t 21

<210> 760
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 760
uuuauugucu cugccuggat t 21

<210> 761
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 761
gugaaaggca cuuuucaut t 21

<210> 762
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 762
aaugaaaagu gccuuucact t 21

<210> 763
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 763
uggacuggua uuugugucut t 21

<210> 764
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 764

agacacaaau accaguccat t 21

<210> 765

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 765

gucugaggcu ggccuacgt t 21

<210> 766

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 766

cguaggcca gccucagact t 21

<210> 767

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 767

cugaggcugg ccuacgggt t 21

<210> 768

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 768

ccguagggc cagccucagt t 21

<210> 769

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 769

gaggcugggcc cuacgggcat t 21

<210> 770

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 770

ugcccguagg gccagccuct t 21

<210> 771

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 771

aggcuggccc uacgggcact t 21

<210> 772

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 772

gugcccguag ggcagccut t 21

<210> 773

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 773

gcuggcccua cgggcaccgt t 21

<210> 774

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 774

cggugcccgu agggccagct t 21

<210> 775
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 775
cuggcccuac gggcacggt t 21

<210> 776
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 776
ccggugcccg uaggccagt t 21

<210> 777
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 777
ggcccuacgg gcaccggugt t 21

<210> 778
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 778
cacggugcc cguaggcct t 21

<210> 779
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 779
ccacucauuc uuggcaggat t 21

<210> 780
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 780
uccugccaag aaugaguggt t 21

<210> 781
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 781
ccuacgggca ccggugaaut t 21

<210> 782
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 782
auucaccggu gcccguaagt t 21

<210> 783
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 783
cuacgggcac cggugaauct t 21

<210> 784
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 784
gauucaccgg ugcccguagt t 21

<210> 785
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 785

uacgggcacc ggugaaucct t 21

<210> 786

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 786

ggauucaccg gugcccgua t 21

<210> 787

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 787

acgggcaccg gugaauccat t 21

<210> 788

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 788

uggauucacc ggugcccgut t 21

<210> 789

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 789

gcaccgguga auccaagut t 21

<210> 790

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 790

cacuuggauu caccggugct t 21

<210> 791
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 791
caccggugaa uccaagugut t 21

<210> 792
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 792
acacuuggau ucaccggugt t 21

<210> 793
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 793
uguggccaug cauguguuct t 21

<210> 794
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 794
gaacacaugc auggccacat t 21

<210> 795
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 795
guggccaugc auguguucat t 21

<210> 796
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 796

ugaacacaug cauggccact t 21

<210> 797

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 797

gccaugcaug uguucagaat t 21

<210> 798

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 798

uucugaacac augcauggct t 21

<210> 799

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 799

uauuccacca cggcugucat t 21

<210> 800

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 800

ugacagccgu gguggaauat t 21

<210> 801

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 801

gucaucacca aucccaaggt t 21

<210> 802

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 802

ccuugggauu ggugaugact t 21

<210> 803

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 803

guccucugau ggucaaagut t 21

<210> 804

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 804

acuuugacca ucagaggact t 21

<210> 805

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 805

gauggucaaa guucuagaut t 21

<210> 806

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 806

aucuagaacu uugaccaut t 21

<210> 807
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 807
augcuguccg aggagucct t 21

<210> 808
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 808
ggacugccuc ggacagcaut t 21

<210> 809
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 809
ccgugcaugu guucagaaat t 21

<210> 810
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 810
uuucugaaca caugcacggt t 21

<210> 811
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 811
agucuggaga gcugcauggt t 21

<210> 812
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 812

ccaugcagcu cuccagacut t 21

<210> 813

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 813

caugggcuca caacugaggt t 21

<210> 814

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 814

ccucaguugu gagcccaugt t 21

<210> 815

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 815

ucucaucguc ugcuccucct t 21

<210> 816

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 816

ggaggagcag acgaugagat t 21

<210> 817

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 817

ccccauucca ugagcaugct t 21

<210> 818

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 818

gcaugcucau ggauggggt t 21

<210> 819

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 819

gccccuacuc cuauuccact t 21

<210> 820

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 820

guggaauagg aguaggggt t 21

<210> 821

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 821

cuauuccacc acggcuguct t 21

<210> 822

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 822

gacagccgug guggaauagt t 21

<210> 823
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 823
cacggcuguc gucaccaaut t 21

<210> 824
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 824
auuggugacg acagccgugt t 21

<210> 825
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 825
aggacgaggg augggauuut t 21

<210> 826
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 826
aaaucccauc ccucguccut t 21

<210> 827
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 827
ucaccucaua ugcuauguut t 21

<210> 828
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 828
aacauagcau augaggugat t 21

<210> 829
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 829
ccucauaugc uauguuagat t 21

<210> 830
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 830
ucuaacauag cauagaggt t 21

<210> 831
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 831
auguuagaag uccaggcagt t 21

<210> 832
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 832
cugccuggac uucuaacaut t 21

<210> 833
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 833

ucugaggcug gcccuacggt t 21

<210> 834

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 834

ccguagggcc agccucagat t 21

<210> 835

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 835

ggcccuacgg gcaccggugt t 21

<210> 836

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 836

caccggugcc cguagggcct t 21

<210> 837

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 837

gggcaccggu gaauccaagt t 21

<210> 838

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 838

cuuggauuca ccggugccct t 21

<210> 839
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 839
ccaugcaugu guucagaaat t 21

<210> 840
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 840
uuucugaaca caugcauggt t 21

<210> 841
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 841
ccggugaauc caagugucct t 21

<210> 842
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 842
ggacacuugg auucaccggt t 21

<210> 843
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 843
acucauucuu ggcaggaugt t 21

<210> 844
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 844
cauccugcca agaaugagut t 21

<210> 845
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 845
aaguguccuc ugauggucat t 21

<210> 846
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 846
ugaccaucag aggacacuut t 21

<210> 847
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 847
ucauucuugg caggauggct t 21

<210> 848
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 848
gccauccugc caagaaugat t 21

<210> 849
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 849

aaguucuaga ugcuguccgt t 21

<210> 850

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 850

cggacagcau cuagaacuut t 21

<210> 851

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 851

guucuagaug cuguccgagt t 21

<210> 852

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 852

cucggacagc aucuagaact t 21

<210> 853

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 853

cuagaugcug uccgaggcat t 21

<210> 854

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 854

ugccucggac agcaucuagt t 21

<210> 855
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 855
gaugcugucc gaggcaguct t 21

<210> 856
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 856
gacugccucg gacagcauct t 21

<210> 857
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 857
cauucuuggc aggauggcut t 21

<210> 858
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 858
agccauccug ccaagaauht t 21

<210> 859
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 859
ugcuguccga ggcaguccut t 21

<210> 860
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 860
aggacugccu cggacagcat t 21

<210> 861
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 861
ccgaggcagu ccugccauct t 21

<210> 862
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 862
gauggcagga cugccucggt t 21

<210> 863
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 863
caguccugcc aucaaugugt t 21

<210> 864
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 864
cacauaugaug gcaggacugt t 21

<210> 865
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 865

caauguggcc gugcaugugt t 21

<210> 866

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 866

cacaugcacg gccacauugt t 21

<210> 867

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 867

auguguucag aaaggcugct t 21

<210> 868

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 868

gcagccuuuc ugaacacaut t 21

<210> 869

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 869

cagaagucca cucauucuut t 21

<210> 870

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 870

aagaaugagu ggacuucugt t 21

<210> 871
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 871
ggcaggaugg cuucauct t 21

<210> 872
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 872
gaugagaagc cauccugct t 21

<210> 873
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 873
gagccauug ccucuggat t 21

<210> 874
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 874
ucccagaggc aaauugcuct t 21

<210> 875
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 875
caggauggcu ucucaugct t 21

<210> 876
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 876

acgaugagaa gccauccugt t 21

<210> 877

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 877

aggauggcuu cucaucguct t 21

<210> 878

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 878

gacgaugaga agccaucct t 21

<210> 879

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 879

agagcugcau gggcucacat t 21

<210> 880

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 880

ugugagccca ugcagcucut t 21

<210> 881

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 881

gcugcauggg cucacaacut t 21

<210> 882

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 882

aguugugagc ccaugcagct t 21

<210> 883

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 883

ggauggcuuc ucaucgucut t 21

<210> 884

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 884

agacgaugag aagccaucct t 21

<210> 885

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 885

gcaugggcuc acaacugagt t 21

<210> 886

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 886

cucaguugug agccaugct t 21

<210> 887
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 887
augggcucac aacugaggat t 21

<210> 888
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 888
uccucaguug ugagccaut t 21

<210> 889
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 889
ugggcucaca acugaggagt t 21

<210> 890
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 890
cuccucaguu gugagccat t 21

<210> 891
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 891
gaggaauuug uagaaggat t 21

<210> 892
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 892

ucccuucucac aaauuccuct t 21

<210> 893

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 893

uuuguagaag ggauuacat t 21

<210> 894

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 894

uguauauccc uucuacaaat t 21

<210> 895

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 895

uuguagaagg gauuacaat t 21

<210> 896

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 896

uuguauaucc cuucuacaat t 21

<210> 897

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 897

uguagaaggg auauacaaat t 21

<210> 898

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 898

uuuguauauc ccuucuat t 21

<210> 899

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 899

agaagggaau uacaaagug t 21

<210> 900

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 900

cacuuuguau aucccuuc t 21

<210> 901

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 901

aaguggaaau agacaccaat t 21

<210> 902

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 902

uuggugucua uuuccacuat t 21

<210> 903
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 903
ggaaauagac accaaaucut t 21

<210> 904
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 904
agauuuggug ucuauuucct t 21

<210> 905
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 905
gaaauagaca ccaaaucut t 21

<210> 906
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 906
aagauuuggu gucuauuuct t 21

<210> 907
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 907
auagacacca aaucuuacut t 21

<210> 908
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 908
aguaagauuu ggugucuat t 21

<210> 909
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 909
uagacaccaa aucuuacugt t 21

<210> 910
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 910
caguaagauu uggugucuat t 21

<210> 911
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 911
agacaccaaa ucuuacuggt t 21

<210> 912
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 912
ccaguaagau uggugucut t 21

<210> 913
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 913

uuacuggaag gcacuuggct t 21

<210> 914

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 914

gccaaugucc uuccagaaat t 21

<210> 915

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 915

uucucaucgu cugcuccuct t 21

<210> 916

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 916

gaggagcaga cgaugagaat t 21

<210> 917

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 917

ggaaggcacu uggaucuct t 21

<210> 918

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 918

gagaugccaa gugccuuct t 21

<210> 919
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 919
ggcacuuggc aucucccat t 21

<210> 920
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 920
uggggagaug ccaagugcct t 21

<210> 921
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 921
ggcaucuccc cauuccaagt t 21

<210> 922
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 922
cauggaaugg ggagaucct t 21

<210> 923
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 923
gcaucucccc auuccaagt t 21

<210> 924
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 924

ucauggaaug gggagagct t 21

<210> 925

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 925

caucucccca uuccaugagt t 21

<210> 926

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 926

cucauggaau ggggagagt t 21

<210> 927

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 927

aucuccccau uccaugagct t 21

<210> 928

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 928

gcucauggaa uggggagaut t 21

<210> 929

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 929

cuccccauuc caugagcaut t 21

<210> 930

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 930

augcucaugg aauggggagt t 21

<210> 931

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 931

cccauuccau gagcaugcat t 21

<210> 932

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 932

ugcaugcuca uggaaugggt t 21

<210> 933

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 933

ccaugagcau gcagaggugt t 21

<210> 934

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 934

caccucugca ugcucauggt t 21

<210> 935
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 935
agcaugcaga ggugguauut t 21

<210> 936
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 936
aauaccaccu cugcaugcut t 21

<210> 937
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 937
caugcagagg uggauuuc t 21

<210> 938
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 938
ugaauaccac cucugcaugt t 21

<210> 939
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 939
augcagaggu gguauucact t 21

<210> 940
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 940
gugaauacca ccucugcaut t 21

<210> 941
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 941
ggugguauuc acagccaact t 21

<210> 942
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 942
guuggcugug aauaccact t 21

<210> 943
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 943
gugguauuca cagccaacgt t 21

<210> 944
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 944
cguuggcugu gaauaccact t 21

<210> 945
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 945

ugguauucac agccaacgat t 21

<210> 946

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 946

ucguuggcug ugaauaccat t 21

<210> 947

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 947

ggauuucaca gccaacgact t 21

<210> 948

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 948

gucguuggcu gugauuacct t 21

<210> 949

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 949

guauucacag ccaacgacut t 21

<210> 950

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 950

agucguuggc ugugaauact t 21

<210> 951
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 951
uauucacagc caacgacuct t 21

<210> 952
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 952
gagucguugg cugugaauat t 21

<210> 953
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 953
ucacagccaa cgacuccggt t 21

<210> 954
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 954
ccggagucgu uggcugugat t 21

<210> 955
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 955
ccccgccgc acaccauugt t 21

<210> 956
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 956

caauggugua gcggcggggt t 21

<210> 957

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 957

gaaguccacu cauucuuggt t 21

<210> 958

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 958

ccaagaauga guggacuuct t 21

<210> 959

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 959

cccugcugag ccccuacuct t 21

<210> 960

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 960

gaguaggggc ucagcagggt t 21

<210> 961

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 961

cugagccccc acuccauut t 21

<210> 962

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 962

aaaggagua ggggcucagt t 21

<210> 963

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 963

ugagcccua cuccauuuct t 21

<210> 964

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 964

gaauaggagu aggggcucat t 21

<210> 965

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 965

cccuacucc uauuccacct t 21

<210> 966

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 966

gguggaauag gaguaggggt t 21

<210> 967
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 967
cuacuccuau uccaccacgt t 21

<210> 968
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 968
cgugguggaa uaggaguagt t 21

<210> 969
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 969
uacuccuauu ccaccacggt t 21

<210> 970
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 970
ccguggugga auaggaguat t 21

<210> 971
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 971
acuccuauuc caccacggct t 21

<210> 972
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 972

gccguggugg aauaggagut t 21

<210> 973

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 973

uccuauucca ccacggcugt t 21

<210> 974

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 974

cagccguggu ggaauaggat t 21

<210> 975

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 975

uauuccacca cggcugucgt t 21

<210> 976

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 976

cgacagccgu gguggaauat t 21

<210> 977

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 977

auuccaccac ggcugucgut t 21

<210> 978

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 978

acgacagccg ugguggaaut t 21

<210> 979

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 979

caccacggcu gucgucacct t 21

<210> 980

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 980

ggugacgaca gccguggugt t 21

<210> 981

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 981

accacggcug ucgucaccat t 21

<210> 982

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 982

uggugacgac agccguggut t 21

<210> 983
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 983
ccacggcugu cgucaccaat t 21

<210> 984
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 984
uuggugacga cagccguggt t 21

<210> 985
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 985
acggcugucg ucaccaauct t 21

<210> 986
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 986
gauuggugac gacagccgut t 21

<210> 987
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 987
cggcugucgu caccaaucct t 21

<210> 988
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 988

ggauugguga cgacagccgt t 21

<210> 989

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 989

cgucaccaau cccaaggaat t 21

<210> 990

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 990

uuccuuggga uuggugacgt t 21

<210> 991

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 991

caauccaag gaaugagggt t 21

<210> 992

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 992

cccucauucc uugggauugt t 21

<210> 993

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 993

ccugaaggac gagggauugt t 21

<210> 994

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 994

ccaucccucg uccuucaggt t 21

<210> 995

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 995

ggacgaggga ugggauuuct t 21

<210> 996

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 996

gaaaucccau ccucgucct t 21

<210> 997

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 997

aaguccacuc auucuuggct t 21

<210> 998

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 998

gccaagaug aguggacuut t 21

<210> 999
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 999
gggauuucau gaaaccaagt t 21

<210> 1000
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1000
cuugguuaca ugaaauccct t 21

<210> 1001
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1001
ggauuucaug uaaccaagat t 21

<210> 1002
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1002
ucuugguuac augaaucct t 21

<210> 1003
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1003
ucaguaaacc aagaguauut t 21

<210> 1004
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1004
aauacucuug guuacauat t 21

<210> 1005
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1005
auguaaccaa gaguauucct t 21

<210> 1006
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1006
ggauuacucu ugguuacat t 21

<210> 1007
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1007
uguaaccaag aguauuccat t 21

<210> 1008
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1008
uggauuacuc uggguacat t 21

<210> 1009
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1009

guaaccaaga guauuccaut t 21

<210> 1010

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1010

auggaaauacu cuugguuact t 21

<210> 1011

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1011

ugccuugcug gacugguaut t 21

<210> 1012

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1012

auaccagucc agcaaggcat t 21

<210> 1013

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1013

uaaagcagug uuuucacut t 21

<210> 1014

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1014

aggugaaaac acugcuuat t 21

<210> 1015
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1015
gccuugcugg acugguauut t 21

<210> 1016
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1016
aauaccaguc cagcaaggct t 21

<210> 1017
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1017
uguuuucacc ucauauugcut t 21

<210> 1018
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1018
agcauauugag gugaaaacat t 21

<210> 1019
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1019
guuuuaccu cauauugcuat t 21

<210> 1020
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1020
uagcauauga ggugaaaact t 21

<210> 1021
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1021
uuuucaccuc auaugcuaut t 21

<210> 1022
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1022
auagcauag aggugaaaat t 21

<210> 1023
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1023
uucaccucau augcuaut t 21

<210> 1024
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1024
acauagcaua ugaggugaat t 21

<210> 1025
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1025

caccucauau gcuauguuat t 21

<210> 1026

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1026

uaacauagca uaugaggugt t 21

<210> 1027

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1027

ccuugcugga cugguauuut t 21

<210> 1028

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1028

aaauaccagu ccagcaaggt t 21

<210> 1029

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1029

auaugcuaug uuagaaguct t 21

<210> 1030

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1030

gacuucuaac auagcauaut t 21

<210> 1031
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1031
uaugcuaugu uagaagucct t 21

<210> 1032
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1032
ggacuucuaa cauagcauat t 21

<210> 1033
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1033
ugcuauuuua gaaguccagt t 21

<210> 1034
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1034
cuggacuucu aacauagcat t 21

<210> 1035
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1035
cuugcuggac uggauuuugt t 21

<210> 1036
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1036

caaauaccag uccagcaagt t 21

<210> 1037

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1037

aguccaggca gagacaauat t 21

<210> 1038

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1038

uauugucucu gccuggacut t 21

<210> 1039

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1039

uccaggcaga gacaauaaat t 21

<210> 1040

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1040

uuuauugucu cugccuggat t 21

<210> 1041

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1041
gugaaaggca cuuuucauut t 21

<210> 1042
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1042
aaugaaaagu gccuuucact t 21

<210> 1043
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1043
uggacuggua uuugugucut t 21

<210> 1044
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1044
agacacaaau accaguccat t 21

<210> 1045
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1045
gucugaggcu ggcccuacgt t 21

<210> 1046
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1046
cguagggcca gccucagact t 21

<210> 1047
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1047
cugaggcugg ccuacgggt t 21

<210> 1048
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1048
cccguagggc cagccucagt t 21

<210> 1049
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1049
gaggcuggcc cuacgggcat t 21

<210> 1050
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1050
ugcccguagg gccagccuct t 21

<210> 1051
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1051
aggcuggccc uacgggcact t 21

<210> 1052
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1052
gugcccguag gccagccut t 21

<210> 1053
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1053
gcuggcccua cgggcaccgt t 21

<210> 1054
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1054
cggugcccgu agggccagct t 21

<210> 1055
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1055
cuggcccuac gggcaccggt t 21

<210> 1056
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1056
ccggugcccg uaggccagt t 21

<210> 1057
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1057

ggcccuacgg gcaccggugt t 21

<210> 1058

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1058

caccggugcc cguagggcct t 21

<210> 1059

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1059

ccacucauuc uuggcaggat t 21

<210> 1060

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1060

uccugccaag aaugaguggt t 21

<210> 1061

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1061

ccuacgggca ccggugaaut t 21

<210> 1062

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1062

auucaccggu gcccgagggt t 21

<210> 1063
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1063
cuacgggcac cggugaauct t 21

<210> 1064
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1064
gauuacccgg ugcccguagt t 21

<210> 1065
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1065
uacgggcacc ggugaauoct t 21

<210> 1066
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1066
ggauuacccg gugcccguat t 21

<210> 1067
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1067
acgggcaccg gugaauccat t 21

<210> 1068
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1068
uggauucacc ggugcccgut t 21

<210> 1069
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1069
gcaccgguga auccaagugt t 21

<210> 1070
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1070
cacuuggauu caccggugct t 21

<210> 1071
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1071
caccggugaa uccaagugut t 21

<210> 1072
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1072
acacuuggau ucaccggugt t 21

<210> 1073
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1073

uguggccaug cauguguuct t 21

<210> 1074

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1074

gaacacaugc auggccacat t 21

<210> 1075

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1075

guggccaugc auguguucat t 21

<210> 1076

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1076

ugaacacaug cauggccact t 21

<210> 1077

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1077

gccaugcaug uguucagaat t 21

<210> 1078

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1078

uucugaacac augcauggct t 21

<210> 1079
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1079
uauuccacca cggcugucat t 21

<210> 1080
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1080
ugacagccgu gguggaauat t 21

<210> 1081
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1081
gucaucacca auccaaggt t 21

<210> 1082
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1082
ccuugggauu ggugaugact t 21

<210> 1083
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1083
guccucugau ggucaaagut t 21

<210> 1084
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1084

acuuugacca ucagaggact t 21

<210> 1085

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1085

gauggucaaa guucuagaut t 21

<210> 1086

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1086

aucuagaacu uugaccaut t 21

<210> 1087

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1087

augcuguccg aggcagucct t 21

<210> 1088

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1088

ggacugccuc ggacagcaut t 21

<210> 1089

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1089

ccgugcaugu guucagaaat t 21

<210> 1090

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1090

uuucugaaca caugcacggt t 21

<210> 1091

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1091

agucuggaga gcugcauggt t 21

<210> 1092

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1092

ccaugcagcu caccagacut t 21

<210> 1093

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1093

caugggcuca caacugaggt t 21

<210> 1094

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1094

ccucaguugu gagcccaugt t 21

<210> 1095
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1095
ucucaucguc ugcuccucct t 21

<210> 1096
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1096
ggaggagcag acgaugagat t 21

<210> 1097
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1097
ccccauucca ugagcaugct t 21

<210> 1098
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1098
gcaugcuc au ggauggggt t 21

<210> 1099
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1099
gccccuacuc cuauuccact t 21

<210> 1100
<211> 21
<212> DNA

<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1100
guggaauagg aguaggggt t 21

<210> 1101
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1101
cuauuccacc acggcuguct t 21

<210> 1102
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1102
gacagccgug guggaauagt t 21

<210> 1103
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1103
cacggcuguc gucaccaut t 21

<210> 1104
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1104
auuggugacg acagccgugt t 21

<210> 1105
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1105
aggacgaggg augggauuut t 21

<210> 1106
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1106
aaaucccauc ccucgucut t 21

<210> 1107
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1107
ucaccucaua ugcuauguut t 21

<210> 1108
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1108
aacauagcau augaggugat t 21

<210> 1109
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1109
ccucauaugc uauguuagat t 21

<210> 1110
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1110
ucuaacauag cauagaggt t 21

<210> 1111
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1111
auguuagaag uccaggcagt t 21

<210> 1112
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1112
cugccuggac uucuaacaut t 21

<210> 1113
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1113
ucugaggcug gcccuacgt t 21

<210> 1114
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1114
ccguagggcc agccucagat t 21

<210> 1115
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1115
ggcccuacgg gcaccggugt t 21

<210> 1116
<211> 21
<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1116

caccggugcc cguagggcct t 21

<210> 1117

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1117

gggcaccggu gaauccaagt t 21

<210> 1118

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1118

cuuggauuca ccggugccct t 21

<210> 1119

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1119

ccaugcaugu guucagaaat t 21

<210> 1120

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1120

uuucugaaca caugcauggt t 21

<210> 1121

<211> 19

<212> RNA

<213> Rattus norvegicus

<400> 1121

guccucugau ggucaaagu 19

<210> 1122
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1122
 acuuugacca ucagaggac 19

 <210> 1123
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1123
 uucuugcucu auaaacgu 19

 <210> 1124
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1124
 acgguuuaua gagcaagaa 19

 <210> 1125
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1125
 cucuaaaaac cguguuagc 19

 <210> 1126
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1126
 gcuaacacgg uuuaugag 19

 <210> 1127
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1127
 ucgccacuac accaucgca 19

 <210> 1128
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1128
 ugcgauggug uaguggcga 19

 <210> 1129
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1129
 ucuugcucua uaaaccgug 19

<210> 1130
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1130
 cacgguuuau agagcaaga 19

 <210> 1131
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1131
 ugcucuauaa accguguua 19

 <210> 1132
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1132
 uaacacgguu uauagagca 19

 <210> 1133
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1133
 caguguucuu gcucuauaa 19

 <210> 1134
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1134
 uuauagagca agaacacug 19

 <210> 1135
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1135
 gcucuauaaa ccguguuag 19

 <210> 1136
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1136
 cuaacacggu uuauagagc 19

 <210> 1137
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1137
 ccuggaugcu guccgaggc 19

<210> 1138
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1138
 gccucggaca gcauccagg 19

 <210> 1139
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1139
 ucugaugguc aaaguccug 19

 <210> 1140
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1140
 caggacuuug accaucaga 19

 <210> 1141
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1141
 cuggagagcu gcacgggcu 19

 <210> 1142
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1142
 agcccuguca gcucuccag 19

 <210> 1143
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1143
 ucuauaaacc guguuagca 19

 <210> 1144
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1144
 ugcuaacacg guuuauaga 19

 <210> 1145
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1145
 aacaguguuc uugcucuai 19

<210> 1146
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1146
 auagagcaag aacacuguu 19

 <210> 1147
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1147
 cucugauggu caaaguccu 19

 <210> 1148
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1148
 aggacuuuga ccaucagag 19

 <210> 1149
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1149
 ugcuguccga ggcagccu 19

 <210> 1150
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1150
 agggcugccu cggacagca 19

 <210> 1151
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1151
 gucuggagag cugcacggg 19

 <210> 1152
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1152
 cccgugcagc ucuccagac 19

 <210> 1153
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1153
 acaguguucu ugcucuaua 19

<210> 1154
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1154
 uauagagcaa gaacacugu 19

 <210> 1155
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1155
 ccucugaugg ucaaagucc 19

 <210> 1156
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1156
 ggacuuugac caucagagg 19

 <210> 1157
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1157
 aguccggau gcuguccga 19

 <210> 1158
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1158
 ucggacagca uccaggacu 19

 <210> 1159
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1159
 uugccucugg gaagaccgc 19

 <210> 1160
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1160
 gcggucuucc cagaggcaa 19

 <210> 1161
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1161
 uggagagcug cacgggcuc 19

<210> 1162
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1162
 gagcccgugc agcucucca 19

 <210> 1163
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1163
 gagagcugca cgggcucac 19

 <210> 1164
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1164
 gugagcccgu gcagcucuc 19

 <210> 1165
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1165
 gacugcacg ggcucacca 19

 <210> 1166
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1166
 uggugagccc gugcagcuc 19

 <210> 1167
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1167
 uacaccaucg cagcccugc 19

 <210> 1168
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1168
 gcagggcugc gauggugua 19

 <210> 1169
 <211> 19
 <212> RNA
 <213> Rattus norvegicus

 <400> 1169
 guccuggaug cuguccgag 19

<210> 1170
<211> 19
<212> RNA
<213> Rattus norvegicus

<400> 1170
cucggacagc auccaggac 19

<210> 1171
<211> 19
<212> RNA
<213> Rattus norvegicus

<400> 1171
agagcugcac gggcucacc 19

<210> 1172
<211> 19
<212> RNA
<213> Rattus norvegicus

<400> 1172
ggugagcccg ugcagcucu 19

<210> 1173
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1173
guccucugau ggucaaagun n 21

<210> 1174
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1174
acuuugacca ucagaggacn n 21

<210> 1175
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1175
uucuugcucu auaaacgun n 21

<210> 1176
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1176
acgguuuua gagcaagaan n 21

<210> 1177
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1177
cucuauaac cguguuagcn n 21

<210> 1178
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1178
gcuaacacgg uuuaugagn n 21

<210> 1179
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1179
ucgccacuac accaucgan n 21

<210> 1180
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1180
ugcgauaggug uaguggcgan n 21

<210> 1181
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1181
ucuugcucua uaaaccgugn n 21

<210> 1182
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1182

cacgguuuau agagcaagan n 21

<210> 1183

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1183

ugcucuaauaa accguguuan n 21

<210> 1184

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1184

uaacacgguu uauagagcan n 21

<210> 1185

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1185

caguguucuu gcucuaaau n 21

<210> 1186

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1186

uuauagagca agaacacugn n 21

<210> 1187

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1187

gcucuauaaa ccguguuagn n 21

<210> 1188

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1188

cuaacacggu uuauagagcn n 21

<210> 1189

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1189

ccuggaugcu guccgaggcn n 21

<210> 1190

<211> 21

<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1190
gccucggaca gcauccagn n 21

<210> 1191
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1191
ucugaugguc aaaguccugn n 21

<210> 1192
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1192
caggacuuug accaucagan n 21

<210> 1193
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1193
cuggagagcu gcacgggcun n 21

<210> 1194
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1194
agcccgugca gcucuccagn n 21

<210> 1195
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1195
ucuauaaacc guguuagcan n 21

<210> 1196
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1196
ugcuaaacgc guuuauagan n 21

<210> 1197
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1197

aacaguguuc uugcucuaun n 21

<210> 1198

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1198

auagagcaag aacacuguun n 21

<210> 1199

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1199

cucugauggu caaaguccun n 21

<210> 1200

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1200

aggacuuuga ccaucagagn n 21

<210> 1201

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1201
ugcuguccga ggcagcccun n 21

<210> 1202
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1202
agggcugccu cggacagcan n 21

<210> 1203
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1203
gucuggagag cugcacgggn n 21

<210> 1204
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1204
cccgucagc ucuccagacn n 21

<210> 1205
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1205
acaguguucu ugcucuaauan n 21

<210> 1206
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1206
uauagagcaa gaacacugun n 21

<210> 1207
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1207
ccucugaugg ucaaaguccn n 21

<210> 1208
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1208

ggacuuugac caucagaggn n 21

<210> 1209

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1209

aguccggau gcuguccgan n 21

<210> 1210

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1210

ucggacagca uccaggacun n 21

<210> 1211

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1211

uugccucugg gaagaccgcn n 21

<210> 1212

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20) .. (21)

<223> a, c, t, g, unknown or other

<400> 1212

gcggucuucc cagaggcaan n 21

<210> 1213

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1213

uggagagcug cacgggcucn n 21

<210> 1214

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1214

gagcccuguc agcucuccan n 21

<210> 1215

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>

<221> modified_base

<222> (20)..(21)

<223> a, c, t, g, unknown or other

<400> 1215

gagagcugca cgggcucacn n 21

<210> 1216

<211> 21

<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1216
gugagcccg gcagcucucn n 21

<210> 1217
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1217
gagcugcacg ggcucaccan n 21

<210> 1218
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1218
uggugagccc ggcagcucn n 21

<210> 1219
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1219
uacaccaucg cagcccugcn n 21

<210> 1220
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1220
gcagggcugc gaugguguan n 21

<210> 1221
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1221
guccuggaug cuguccgagn n 21

<210> 1222
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1222
cucggacagc auccaggacn n 21

<210> 1223
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1223
agagcugcac gggcucaccn n 21

<210> 1224
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<220>
<221> modified_base
<222> (20)..(21)
<223> a, c, t, g, unknown or other

<400> 1224
ggugagcccg ugcagcucun n 21

<210> 1225
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1225
guccucugau ggucaaagut t 21

<210> 1226
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1226
acuuugacca ucagaggact t 21

<210> 1227
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1227
uucuugcucu auaaaccgut t 21

<210> 1228
<211> 21

<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1228
acgguuuuaa gagcaagaat t 21

<210> 1229
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1229
cucuauaaac cguguuagct t 21

<210> 1230
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1230
gcuaacacgg uuuauagat t 21

<210> 1231
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1231
ucgccacuac accaucgat t 21

<210> 1232
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1232
ugcgauggug uaguggcgaat t 21

<210> 1233
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1233
ucuugcucua uaaaccgugt t 21

<210> 1234
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1234
cacgguuuau agagcaagat t 21

<210> 1235
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1235
ugcucuauaa accguguuat t 21

<210> 1236
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1236
uaacacgguu uauagagcat t 21

<210> 1237
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1237
caguguucuu gcucuauaat t 21

<210> 1238
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1238

uuauagagca agaacacugt t 21

<210> 1239

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1239

gcucuauaaa ccguguuagt t 21

<210> 1240

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1240

cuaacacggu uuauagagct t 21

<210> 1241

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1241

ccuggaugcu guccgaggct t 21

<210> 1242

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1242

gccucggaca gcauccaggt t 21

<210> 1243

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1243

ucugaugguc aaaguccugt t 21

<210> 1244

<211> 21

<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1244
caggacuuug accaucagat t 21

<210> 1245
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1245
cuggagagcu gcacgggcut t 21

<210> 1246
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1246
agcccgugca gcucuccagt t 21

<210> 1247
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1247
ucuauaaacc guguuagcat t 21

<210> 1248
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1248
ugcuaacacg guuuauagat t 21

<210> 1249
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1249
aacaguguuc ugcucuaut t 21

<210> 1250
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1250
auagagcaag aacacuguut t 21

<210> 1251
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1251
cucugauggu caaaguccut t 21

<210> 1252
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1252
aggacuuuga ccaucagagt t 21

<210> 1253
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1253
ugcuguccga ggcagcccut t 21

<210> 1254
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1254

agggcugccu cggacagcat t 21

<210> 1255

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1255

gucuggagag cugcacgggt t 21

<210> 1256

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1256

cccugcagc ucuccagact t 21

<210> 1257

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1257

acaguguucu ugcucuauat t 21

<210> 1258

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1258

uauagagcaa gaacacugut t 21

<210> 1259

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1259

ccucugaugg ucaaagucct t 21

<210> 1260

<211> 21

<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1260
ggacuuugac caucagaggt t 21

<210> 1261
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1261
aguccuggau gcuguccgat t 21

<210> 1262
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1262
ucggacagca uccaggacut t 21

<210> 1263
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1263
uugccucugg gaagaccgct t 21

<210> 1264
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1264
gcgguuucc cagaggcaat t 21

<210> 1265
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1265
uggagagcug cacgggcuct t 21

<210> 1266
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1266
gagcccgugc agcucucct t 21

<210> 1267
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1267
gagagcugca cgggcucact t 21

<210> 1268
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1268
gugagcccgu gcagcucuct t 21

<210> 1269
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1269
gagcugcacg ggcucacat t 21

<210> 1270
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1270

uggugagccc gugcagcuct t 21

<210> 1271

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1271

uacaccaucg cagcccugct t 21

<210> 1272

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1272

gcagggcugc gaugguguat t 21

<210> 1273

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1273

guccuggaug cuguccgagt t 21

<210> 1274

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1274

cucggacagc auccaggact t 21

<210> 1275

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1275

agagcugcac gggcucacct t 21

<210> 1276

<211> 21

<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1276
ggugagcccg ugcagcucut t 21

<210> 1277
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1277
guccucugau ggucaaagut t 21

<210> 1278
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1278
acuuugacca ucagaggact t 21

<210> 1279
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1279
uucuugcucu auaaaccgut t 21

<210> 1280
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1280
acgguuuuaa gagcaagaat t 21

<210> 1281
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1281
cucuauaaac cguguuagct t 21

<210> 1282
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1282
gcuaacacgg uuuauagagt t 21

<210> 1283
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1283
ucgccacuac accaucgcat t 21

<210> 1284
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1284
ugcgauggug uaguggcgat t 21

<210> 1285
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1285
ucuugcucua uaaaccgugt t 21

<210> 1286
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1286

cacgguuuau agagcaagat t 21

<210> 1287

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1287

ugcucuauaa accguguuat t 21

<210> 1288

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1288

uaacacgguu uauagagcat t 21

<210> 1289

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1289

caguguucuu gcucuauaat t 21

<210> 1290

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1290

uuauagagca agaacacugt t 21

<210> 1291

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1291

gcucuauaaa ccguguuagt t 21

<210> 1292

<211> 21

<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1292
cuaacacggu uuauagagct t 21

<210> 1293
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1293
ccuggaugcu guccgaggct t 21

<210> 1294
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1294
gccucggaca gcauccaggt t 21

<210> 1295
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1295
ucugaugguc aaaguccugt t 21

<210> 1296
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1296
caggacuuug accaucagat t 21

<210> 1297
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1297
cuggagagcu gcacgggcut t 21

<210> 1298
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1298
agcccgugca gcucuccagt t 21

<210> 1299
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1299
ucuauaaacc guguuagcat t 21

<210> 1300
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1300
ugcuaacacg guuuauagat t 21

<210> 1301
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1301
aacaguguuc ugcucuaut t 21

<210> 1302
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1302

auagagcaag aacacuguut t 21

<210> 1303

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1303

cucugauggu caaaguccut t 21

<210> 1304

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1304

aggacuuuga ccaucagagt t 21

<210> 1305

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1305

ugcuguccga ggcagcccut t 21

<210> 1306

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1306

agggcugccu cggacagcat t 21

<210> 1307

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1307

gucuggagag cugcacgggt t 21

<210> 1308

<211> 21

<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1308
cccgugcagc ucuccagact t 21

<210> 1309
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1309
acaguguucu ugcucuauat t 21

<210> 1310
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1310
uauagagcaa gaacacugut t 21

<210> 1311
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1311
ccucugaugg ucaaagucct t 21

<210> 1312
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1312
ggacuuugac caucagaggt t 21

<210> 1313
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1313
aguccuggau gcuguccgat t 21

<210> 1314
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1314
ucggacagca uccaggacut t 21

<210> 1315
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1315
uugccucugg gaagaccgct t 21

<210> 1316
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1316
gcggucuucc cagaggcaat t 21

<210> 1317
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1317
uggagagcug cacgggcuct t 21

<210> 1318
<211> 21
<212> DNA
<213> Rattus norvegicus

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1318

gagcccgugc agcucucctat t 21

<210> 1319

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1319

gagagcugca cgggcucact t 21

<210> 1320

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1320

gugagcccgu gcagcucuct t 21

<210> 1321

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1321

gagcugcacg ggcucacat t 21

<210> 1322

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1322

uggugagccc ggcagcuct t 21

<210> 1323

<211> 21

<212> DNA

<213> Rattus norvegicus

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

<400> 1323

uacaccaucg cagcccugct t 21

<210> 1324

<211> 21

<212> DNA
 <213> Rattus norvegicus

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

 <400> 1324
 gcagggcugc gaugguguat t 21

 <210> 1325
 <211> 21
 <212> DNA
 <213> Rattus norvegicus

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

 <400> 1325
 guccuggaug cuguccgagt t 21

 <210> 1326
 <211> 21
 <212> DNA
 <213> Rattus norvegicus

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

 <400> 1326
 cucggacagc auccaggact t 21

 <210> 1327
 <211> 21
 <212> DNA
 <213> Rattus norvegicus

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

 <400> 1327
 agagcugcac gggcucacct t 21

 <210> 1328
 <211> 21
 <212> DNA
 <213> Rattus norvegicus

 <220>
 <221> source
 <223> /note="Description of Combined DNA/RNA Molecule: Rattus norvegicus"

 <400> 1328
 ggugagcccg ugcagcucut t 21

 <210> 1329
 <211> 650
 <212> DNA
 <213> Homo sapiens

 <400> 1329

```

acagaagtcc actcattctt ggcaggatgg atttctcatg totgtctctc ctctgccttg      60
ctggactggt atttgtgtct gaggtgggcc ctacgggcac cggatgaatcc aagtgtcctc      120
tgatggtcaa agttcttagat gctgtccgag gcagtcctgc catcaatgtg gccgtgcatg      180
tgttcagaaa ggotgtgat gacacctggg agccatttgc ctctgggaaa accagtga      240
ctggagagct gcatgggctc acaactgagg aggaatttgt agaagggata tacaaagtgg      300
aaatagacac caaatcttac tggaaggcac ttggcatctc cccattccat gagcatgcag      360
aggtggtatt cacagccaac gactccggcc cccgccgcta caccattgcc gccctgctga      420
gccctactc ctattccacc acggctgtcg tcaccaatcc caaggaatga gggacttctc      480
ctccagtga cctgaaggac gagggatggg atttcatgta accaagagta ttccattttt      540
actaaagcag tgttttcacc tcatatgcta tgtagaagt ccaggcagag acaataaaac      600
attcctgtga aaggcacttt tcattccaaa aaaaaaaaaa aaaaaaaaaa      650

```

<210> 1330

<211> 595

<212> DNA

<213> Rattus norvegicus

<400> 1330

```

cctgacagga tggcttccct tcgcctgttc ctccctctgcc tcgctggact gatatttgcg      60
tctgaagctg gccctggggg tgctggagaa tccaagtgtc ctctgatggt caaagtcctg      120
gatgctgtcc gaggcagccc tgctgtcgat gtggccgtga aagtgttcaa aaggactgca      180
gacggaagct gggagccgtt tgctctggg aagaccgccc agtctggaga gctgcacggg      240
ctcaccacag atgagaagtt cacggaaggg gtgtacaggg tagaactgga caccaaatca      300
tactggaagg ctcttggcat ttccccattc catgaatacg cagaggtggt tttcacagcc      360
aatgactctg gtcatcgcca ctacaccatc gcagcccctg tcagccccta ctccacagc      420
accactgctg tcgtcagtaa ccccagaaac tgagggaccc agcccacgag gaccaagatc      480
ttgccaaagc agtagctccc atttgaactg aaacagtgtt cttgtctat aaacctgtt      540
agcaactcgg gaagatgccg tgaaacgttc ttattaaacc acctttattt cattc      595

```

<210> 1331

<211> 938

<212> DNA

<213> Homo sapiens

<400> 1331

```

gttgactaag tcaataatca gaatcagcag gtttgcagtc agattggcag ggataagcag      60
cctagctcag gagaagtga gataaaagcc ccaggctggg agcagccatc acagaagtcc      120
actcattctt ggcaggatgg atttctcatg totgtctctc ctctgccttg ctggactggt      180
atttgtgtct gaggtgggcc ctacgggcac cggatgaatcc aagtgtcctc tgatggtcaa      240
agttcttagat gctgtccgag gcagtcctgc catcaatgtg gccgtgcatg tgttcagaaa      300
ggctgtgat gacacctggg agccatttgc ctctgggaaa accagtga      360
gcctgggctc acaactgagg aggaatttgt agaagggata tacaaagtgg aaatagacac      420
caaatcttac tggaaggcac ttggcatctc cccattccat gagcatgcag aggtggtatt      480
cacagccaac gactccggcc cccgccgcta caccattgcc gccctgctga gccctactc      540
ctattccacc acggctgtcg tcaccaatcc caaggaatga gggacttctc ctccagtgga      600
cctgaaggac gagggatggg atttcatgta accaagagta ttccattttt actaaagcag      660
tgttttcacc tcatatgcta tgtagaagt ccaggcagag acaataaaac attcctgtga      720
aaggcacttt tcattccact ttaacttgat tttttaaat cccttattgt cccttcctaaa      780
aaaaagagaa tcaaaatctt acaagaatc aaaggaattc tagaaagtat ctgggcagaa      840
cgctaggaga gatccaaatt tcattgtct tgcaagcaaa gcacgtatta aatatgatct      900
gcagccatta aaaagacaca ttctgtaaaa aaaaaaaaaa      938

```

<210> 1332

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1332
gggauuucau gaaaccaagt t 21

<210> 1333
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1333
cuugguuaca ugaaauccct t 21

<210> 1334
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1334
ggauuucaug uaaccaagat t 21

<210> 1335
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1335
ucuugguuac augaaauccct t 21

<210> 1336
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1336
gauuucaugu aaccaagagt t 21

<210> 1337
<211> 21
<212> DNA
<213> Homo sapiens

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1337
cucuugguua caugaaauct t 21

<210> 1338
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1338
auuucaugua accaagagut t 21

<210> 1339
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1339
acucuugguu acaugaaa t 21

<210> 1340
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1340
uuucauguaa ccaagaguat t 21

<210> 1341
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1341
uacucuuggu uacaugaaat t 21

<210> 1342
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1342

uucauguaac caagaguaut t 21

<210> 1343

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1343

auacucuugg uuacaugaat t 21

<210> 1344

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1344

ucauguaacc aagaguauut t 21

<210> 1345

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1345

aauacucuug guuacaugat t 21

<210> 1346

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1346

cauguaacca agaguauuct t 21

<210> 1347

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1347

gaauacucuu gguuacaugt t 21

<210> 1348

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1348
auguaaccaa gaguauucct t 21

<210> 1349
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1349
ggaauacucu ugguuacaut t 21

<210> 1350
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1350
uguaaccaag aguauuccat t 21

<210> 1351
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1351
uggaauacuc uugguuacat t 21

<210> 1352
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1352
uaaccaagag uauuccauut t 21

<210> 1353
<211> 21
<212> DNA
<213> Homo sapiens

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1353
aauggaauac ucuugguat t 21

<210> 1354
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1354
aaccaagagu auuccauuut t 21

<210> 1355
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1355
aaauggaaua cucuugguut t 21

<210> 1356
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1356
accaagagua uuccauuuut t 21

<210> 1357
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1357
aaaauggaau acucuuggut t 21

<210> 1358
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1358

ccaagaguau uccauuuuut t 21

<210> 1359
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1359
aaaaauggaa uacucuugt t 21

<210> 1360
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1360
caagaguauu ccuuuuuat t 21

<210> 1361
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1361
uaaaaaugga auacucuugt t 21

<210> 1362
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1362
aagaguauuc cauuuuuact t 21

<210> 1363
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1363
guaaaaaugg aaucucuut t 21

<210> 1364
<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1364
agaguauucc auuuuuacut t 21

<210> 1365
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1365
aguaaaaaug gaauacucut t 21

<210> 1366
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1366
gaguauucca uuuuuacuat t 21

<210> 1367
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1367
uaguaaaaau ggaauacuct t 21

<210> 1368
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1368
aguauuccau uuuuacuaat t 21

<210> 1369
<211> 21
<212> DNA
<213> Homo sapiens

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1369
uuaguaaaaa uggaauacut t 21

<210> 1370
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1370
guauuccauu uuuacuaaat t 21

<210> 1371
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1371
uuuaguaaaa auggaauact t 21

<210> 1372
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1372
uauuccauuu uuacuaaagt t 21

<210> 1373
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1373
cuuuaguaaa aauggaauat t 21

<210> 1374
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1374

auuccauuuu uacuaagct t 21

<210> 1375
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1375
gcuuuaguaa aaauggaaut t 21

<210> 1376
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1376
uuccauuuuu acuaaagcat t 21

<210> 1377
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1377
ugcuuuagua aaaauggaat t 21

<210> 1378
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1378
uccauuuuuu cuaaagcagt t 21

<210> 1379
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1379
cugcuuuagu aaaaauggat t 21

<210> 1380
<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1380
ccauuuuuac uaaagcagut t 21

<210> 1381
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1381
acugcuuuag uaaaaauggt t 21

<210> 1382
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1382
cauuuuuacu aaagcagugt t 21

<210> 1383
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1383
cacugcuuua guaaaaaugt t 21

<210> 1384
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1384
auuuuuacua aagcagugut t 21

<210> 1385
<211> 21
<212> DNA
<213> Homo sapiens

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1385
acacugcuu aguaaaaat t 21

<210> 1386
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1386
uuuuuacuaa agcaguguut t 21

<210> 1387
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1387
aacacugcuu uaguaaaaat t 21

<210> 1388
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1388
uuuuacuaaa gcaguguuut t 21

<210> 1389
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1389
aaacacugcu uuaguaaaaat t 21

<210> 1390
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1390

uuuacuaaag caguguuuut t 21

<210> 1391

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1391

aaaacacugc uuuguuaat t 21

<210> 1392

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1392

uuacuaaagc aguguuuuct t 21

<210> 1393

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1393

gaaaacacug cuuuaguaat t 21

<210> 1394

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1394

uacuaaagca guguuuucat t 21

<210> 1395

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1395

ugaaaacacu gcuuuaguat t 21

<210> 1396

<211> 21

<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1396
acuaaagcag uguuuucact t 21

<210> 1397
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1397
gugaaaacac ugcuuuagut t 21

<210> 1398
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1398
cuaaagcagu guuuucacct t 21

<210> 1399
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1399
ggugaaaaca cugcuuuagt t 21

<210> 1400
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1400
uaaagcagug uuuucacut t 21

<210> 1401
<211> 21
<212> DNA
<213> Homo sapiens

<220>

<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1401
aggugaaaac acugcuuat t 21

<210> 1402
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1402
aaagcagugu uuucaccuct t 21

<210> 1403
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1403
gaggugaaaa cacugcuut t 21

<210> 1404
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1404
aagcaguguu uucaccucat t 21

<210> 1405
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1405
ugaggugaaa acacugcuut t 21

<210> 1406
<211> 21
<212> DNA
<213> Homo sapiens

<220>
<221> source
<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1406

agcaguguuu ucaccuaut t 21

<210> 1407

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1407

augaggugaa aacacugcut t 21

<210> 1408

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1408

guaaccaaga guauuccaut t 21

<210> 1409

<211> 21

<212> DNA

<213> Homo sapiens

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Homo sapiens"

<400> 1409

auggaaauacu cuugguuact t 21

<210> 1410

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<221> source

<223> /note="Description of Artificial Sequence: Synthetic oligonucleotide"

<220>

<221> source

<223> /note="Description of Combined DNA/RNA Molecule: Synthetic oligonucleotide"

<400> 1410

gtaaccaa gagtattccat 19

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO9932619A \[0005\]](#)
- [WO9953050A \[0005\]](#)
- [WO9961631A \[0005\]](#)
- [WO0044895A \[0005\]](#)
- [DE10100586 \[0005\]](#)
- [US20070207974A \[0006\]](#)
- [US20090082300A \[0006\]](#)
- [US7250496B \[0006\]](#)
- [US20060240093A \[0052\]](#)
- [US20070135372A \[0052\]](#)
- [US61045228B \[0052\]](#)
- [US3687808A \[0073\] \[0080\] \[0081\]](#)
- [US4469863A \[0073\]](#)
- [US4476301A \[0073\]](#)
- [US5023243A \[0073\]](#)
- [US5177195A \[0073\]](#)
- [US5188897A \[0073\]](#)
- [US5264423A \[0073\]](#)
- [US5276019A \[0073\]](#)
- [US5278302A \[0073\]](#)
- [US5286717A \[0073\]](#)
- [US5321131A \[0073\]](#)
- [US5399676A \[0073\]](#)
- [US5405939A \[0073\]](#)
- [US5453496A \[0073\]](#)
- [US5455233A \[0073\]](#)
- [US5466677A \[0073\] \[0075\]](#)
- [US5476925A \[0073\]](#)
- [US5519126A \[0073\]](#)
- [US5536821A \[0073\]](#)
- [US5541316A \[0073\]](#)
- [US5550111A \[0073\]](#)
- [US5563253A \[0073\]](#)
- [US5571799A \[0073\]](#)
- [US5587361A \[0073\]](#)
- [US5625050A \[0073\]](#)
- [US5034506A \[0075\] \[0077\]](#)
- [US5166315A \[0075\]](#)
- [US5185444A \[0075\]](#)
- [US5214134A \[0075\]](#)
- [US5216141A \[0075\]](#)
- [US5235033A \[0075\]](#)
- [US564562A \[0075\]](#)
- [US5264564A \[0075\]](#)
- [US5405938A \[0075\]](#)
- [US5434257A \[0075\]](#)
- [US5470967A \[0075\]](#)
- [US5489677A \[0075\] \[0077\]](#)
- [US5541307A \[0075\]](#)
- [US5561225A \[0075\]](#)
- [US5596086A \[0075\]](#)
- [US5602240A \[0075\] \[0077\]](#)
- [US5608046A \[0075\] \[0083\]](#)
- [US5610289A \[0075\]](#)

- [US5618704A \[0075\]](#)
- [US5623070A \[0075\]](#)
- [US5663312A \[0075\]](#)
- [US5633360A \[0075\]](#)
- [US5677437A \[0075\]](#)
- [US5677439A \[0075\]](#)
- [US5539082A \[0076\]](#)
- [US5714331A \[0076\]](#)
- [US5719262A \[0076\]](#)
- [US4981957A \[0079\]](#)
- [US5118800A \[0079\]](#)
- [US5319080A \[0079\]](#)
- [US5359044A \[0079\]](#)
- [US5393878A \[0079\]](#)
- [US5446137A \[0079\]](#)
- [US5466786A \[0079\]](#)
- [US5514785A \[0079\] \[0083\]](#)
- [US5519134A \[0079\]](#)
- [US5567611A \[0079\]](#)
- [US5576427A \[0079\]](#)
- [US5591722A \[0079\]](#)
- [US5597909A \[0079\]](#)
- [US5610300A \[0079\]](#)
- [US5627053A \[0079\]](#)
- [US5639873A \[0079\]](#)
- [US5646265A \[0079\]](#)
- [US5658873A \[0079\]](#)
- [US5670633A \[0079\]](#)
- [US5700920A \[0079\]](#)
- [US4845205A \[0081\]](#)
- [US513030A \[0081\]](#)
- [US5134066A \[0081\]](#)
- [US5175273A \[0081\]](#)
- [US5367066A \[0081\]](#)
- [US5432272A \[0081\]](#)
- [US5457187A \[0081\]](#)
- [US5459255A \[0081\]](#)
- [US5484908A \[0081\]](#)
- [US5502177A \[0081\]](#)
- [US5525711A \[0081\]](#)
- [US5552540A \[0081\]](#)
- [US5567469A \[0081\]](#)
- [US5594121A \[0081\]](#)
- [US5596091A \[0081\]](#)
- [US5614617A \[0081\]](#)
- [US5681941A \[0081\]](#)
- [US5750692A \[0081\]](#)
- [US4828979A \[0083\]](#)
- [US4948882A \[0083\]](#)
- [US5218105A \[0083\]](#)
- [US5525465A \[0083\]](#)
- [US5541313A \[0083\]](#)
- [US5545730A \[0083\]](#)
- [US5552538A \[0083\]](#)
- [US5578717A \[0083\]](#)
- [US5580731A \[0083\]](#)
- [US5591584A \[0083\]](#)
- [US5109124A \[0083\]](#)

- [US5116802A \[0083\]](#)
- [US5138045A \[0083\]](#)
- [US5414077A \[0083\]](#)
- [US5486603A \[0083\]](#)
- [US5512439A \[0083\]](#)
- [US5578718A \[0083\]](#)
- [US4587044A \[0083\]](#)
- [US4605735A \[0083\]](#)
- [US4667025A \[0083\]](#)
- [US4762779A \[0083\]](#)
- [US4789737A \[0083\]](#)
- [US4824941A \[0083\]](#)
- [US4835263A \[0083\]](#)
- [US4876335A \[0083\]](#)
- [US4904582A \[0083\]](#)
- [US4958013A \[0083\]](#)
- [US5082830A \[0083\] \[0083\]](#)
- [US5112963A \[0083\] \[0083\]](#)
- [US5214136A \[0083\] \[0083\]](#)
- [US5245022A \[0083\]](#)
- [US5254469A \[0083\]](#)
- [US5258506A \[0083\]](#)
- [US5262536A \[0083\]](#)
- [US5272250A \[0083\]](#)
- [US5292873A \[0083\]](#)
- [US5317098A \[0083\]](#)
- [US5371241A \[0083\]](#)
- [US5391723A \[0083\]](#)
- [US5416203A \[0083\]](#)
- [US5451463A \[0083\]](#)
- [US5510475A \[0083\]](#)
- [US5512667A \[0083\]](#)
- [US5565552A \[0083\]](#)
- [US5567810A \[0083\]](#)
- [US5574142A \[0083\]](#)
- [US5585481A \[0083\]](#)
- [US5587371A \[0083\]](#)
- [US5595726A \[0083\]](#)
- [US5597696A \[0083\]](#)
- [US5599923A \[0083\]](#)
- [US5599928A \[0083\]](#)
- [US5688941A \[0083\]](#)
- [WO0022113A \[0087\]](#)
- [WO0022114A \[0087\]](#)
- [US6054299A \[0087\]](#)
- [US4868116A \[0089\]](#)
- [US4980286A \[0089\]](#)
- [WO8907136A \[0089\]](#)
- [WO8902468A \[0089\]](#)
- [WO8905345A \[0089\]](#)
- [WO9207573A \[0089\]](#)
- [US5252479A \[0095\]](#)
- [US5139941A \[0095\]](#)
- [WO9413788A \[0095\]](#)
- [WO9324641A \[0095\]](#)
- [US5328470A \[0100\]](#)
- [US6093180A \[0116\]](#)
- [US5814014A \[0116\]](#)

- [US6747014B \[0117\] \[0154\]](#)
- [US4837028A \[0131\]](#)
- [WO8804924A \[0131\]](#)
- [US5543152A \[0131\]](#)
- [WO9713499A \[0131\]](#)
- [US4426330A \[0132\]](#)
- [US4534899A \[0132\]](#)
- [EP0445131B1 \[0132\]](#)
- [WO9004384A \[0132\]](#)
- [US5013556A \[0132\]](#)
- [US5356633A \[0132\]](#)
- [US5213804A \[0132\]](#)
- [EP0496813B1 \[0132\]](#)
- [WO9105545A \[0132\]](#)
- [US5225212A \[0132\]](#)
- [WO9420073A \[0132\]](#)
- [WO9610391A \[0132\]](#)
- [US5540935A \[0132\]](#)
- [US5556948A \[0132\]](#)
- [WO9640062A \[0133\]](#)
- [US5264221A \[0133\]](#)
- [US5665710A \[0133\]](#)
- [WO9704787A \[0133\]](#)
- [WO0003683A \[0141\]](#)
- [US5976567A \[0141\]](#)
- [US5981501A \[0141\]](#)
- [US6534464B \[0141\]](#)
- [US6586410B \[0141\]](#)
- [US6815432B \[0141\]](#)
- [WO9640964A \[0141\]](#)
- [US6110799B \[0144\]](#)
- [WO2008042973A \[0150\]](#)
- [US6123968B \[0152\]](#)
- [US61244834B \[0152\]](#)
- [US6887906B \[0154\]](#)
- [US20030027780A \[0154\]](#)
- [WO2004065601A \[0225\]](#)
- [US61244794B \[0330\]](#)
- [US61242783B \[0330\]](#)
- [US61185545B \[0330\]](#)
- [US61156670B \[0330\]](#)
- [US61115738B \[0330\]](#)
- [US61108956B \[0330\]](#)

Non-patent literature cited in the description

- **YANG, D. et al.** Curr. Biol., 2000, vol. 10, 1191-1200 [0005]
- **STEIN et al.** The Journal of Neuroscience, 2004, vol. 24, 357707-7717 [0006]
- **ELBASHIR et al.** EMBO, 2001, vol. 20, 6877-6888 [0068]
- Current protocols in nucleic acid chemistry John Wiley & Sons, Inc. [0071] [0204]
- **NIELSEN et al.** Science, 1991, vol. 254, 1497-1500 [0076]
- The Concise Encyclopedia Of Polymer Science And Engineering John Wiley & Sons 19900000858-859 [0080]
- **ENGLISH et al.** Angewandte Chemie International Edition 19910000vol. 30, 613- [0090]
- **SANGHVI, Y S.** DsRNA Research and Applications CRC Press 19930000289-302 [0080]
- DsRNA Research and Applications CRC Press 19930000276-278 [0080]

- LETSINGER et al. Proc. Natl. Acad. Sci. USA, 1989, vol. 86, 6553-6556 [0082]
- MANOHARAN et al. Biorg. Med. Chem. Lett., 1994, vol. 4, 1053-1060 [0082]
- MANOHARAN et al. Ann. N.Y. Acad. Sci., 1992, vol. 660, 306-309 [0082]
- MANOHARAN et al. Biorg. Med. Chem. Lett., 1993, vol. 3, 2765-2770 [0082]
- OBERHAUSER et al. Nucl. Acids Res., 1992, vol. 20, 533-538 [0082]
- SAISON-BEHMOARAS et al. EMBO J, 1991, vol. 10, 1111-1118 [0082]
- KABANOV et al. FEBS Lett., 1990, vol. 259, 327-330 [0082]
- SVINARCHUK et al. Biochimie, 1993, vol. 75, 49-54 [0082]
- MANOHARAN et al. Tetrahedron Lett., 1995, vol. 36, 3651-3654 [0082] [0082]
- SHEA et al. Nucl. Acids Res., 1990, vol. 18, 3777-3783 [0082]
- MANOHARAN et al. Nucleosides & Nucleotides, 1995, vol. 14, 969-973 [0082]
- MISHRA et al. Biochim. Biophys. Acta, 1995, vol. 1264, 229-237 [0082]
- CROOKE et al. J. Pharmacol. Exp. Ther., 1996, vol. 277, 923-937 [0082]
- SOUTSCHEK et al. Nature, 2004, vol. 432, 173-178 [0085]
- LETSINGER et al. Proc. Natl. Acad. Sci. USA, 1989, vol. 86, 6553- [0086]
- MANOHARAN et al. Bioorg. Med. Chem. Lett., 1994, vol. 4, 1053- [0086]
- MANOHARAN et al. Ann. N.Y. Acad. Sci., 1992, vol. 660, 306- [0086]
- MANOHARAN et al. Bioorg. Med. Chem. Lett., 1993, vol. 3, 2765- [0086]
- OBERHAUSER et al. Nucl. Acids Res., 1992, vol. 20, 533- [0086]
- SAISON-BEHMOARAS et al. EMBO J., 1991, vol. 10, 111- [0086]
- KABANOV et al. FEBS Lett., 1990, vol. 259, 327- [0086]
- SVINARCHUK et al. Biochimie, 1993, vol. 75, 49- [0086]
- MANOHARAN et al. Tetrahedron Lett., 1995, vol. 36, 3651- [0086] [0086]
- SHEA et al. Nucl. Acids Res., 1990, vol. 18, 3777- [0086]
- MANOHARAN et al. Nucleosides & Nucleotides, 1995, vol. 14, 969- [0086]
- MISHRA et al. Biochim. Biophys. Acta, 1995, vol. 1264, 229- [0086]
- CROOKE et al. J. Pharmacol. Exp. Ther., 1996, vol. 277, 923- [0086]
- COUTURE, A et al. TIG, 1996, vol. 12, 5-10 [0087]
- GASSMANN et al. Proc. Natl. Acad. Sci. USA, 1995, vol. 92, 1292- [0087]
- MUZYCZKA et al. Curr. Topics Micro. Immunol., 1992, vol. 158, 97-129 [0089]
- BERKNER et al. BioTechniques, 1998, vol. 6, 616- [0089]
- ROSENFELD et al. Science, 1991, vol. 252, 431-434 [0089]
- ROSENFELD et al. Cell, 1992, vol. 68, 143-155 [0089]
- EGLITIS et al. Science, 1985, vol. 230, 1395-1398 [0089]
- DANOSMULLIGAN Proc. Natl. Acad. Sci. USA, 1998, vol. 85, 6460-6464 [0089]
- WILSON et al. Proc. Natl. Acad. Sci. USA, 1988, vol. 85, 3014-3018 [0089]
- ARMENTANO et al. Proc. Natl. Acad. Sci. USA, 1990, vol. 87, 6141-6145- [0089]
- HUBER et al. Proc. Natl. Acad. Sci. USA, 1991, vol. 88, 8039-8043 [0089]
- FERRY et al. Proc. Natl. Acad. Sci. USA, 1991, vol. 88, 8377-8381 [0089]
- CHOWDHURY et al. Science, 1991, vol. 254, 1802-1805 [0089]
- VAN BEUSECHEM. et al. Proc. Natl. Acad. Sci. USA, 1992, vol. 89, 7640-19 [0089]
- KAY et al. Human Gene Therapy, 1992, vol. 3, 641-647 [0089]
- DAI et al. Proc. Natl. Acad. Sci. USA, 1992, vol. 89, 10892-10895 [0089]
- HWU et al. J. Immunol., 1993, vol. 150, 4104-4115 [0089]
- COMETTE et al. Human Gene Therapy, 1991, vol. 2, 5-10 [0089]
- CONE et al. Proc. Natl. Acad. Sci. USA, 1984, vol. 81, 6349- [0089]
- HSU et al. J. Infectious Disease, 1992, vol. 166, 769- [0089]
- RABINOWITZ J E et al. J Virol, 2002, vol. 76, 791-801 [0091]
- DORNBURG R Gene Therap., 1995, vol. 2, 301-310 [0092]
- EGLITIS M A Biotechniques, 1988, vol. 6, 608-614 [0092]
- MILLER A D Hum Gene Therap., 1990, vol. 1, 5-14 [0092]
- ANDERSON W F Nature, 1998, vol. 392, 25-30 [0092]
- RUBINSON D A et al. Nat. Genet., vol. 33, 401-406 [0092]
- XIA H et al. Nat. Biotech., 2002, vol. 20, 1006-1010 [0094]
- SAMULSKI R et al. J. Virol., 1987, vol. 61, 3096-3101 [0095]
- FISHER K J et al. J. Virol., 1996, vol. 70, 520-532 [0095]
- SAMULSKI R et al. J. Virol., 1989, vol. 63, 3822-3826 [0095]
- BUCCHINI et al. Proc. Natl. Acad. Sci. USA, 1986, vol. 83, 2511-2515 [0096]

- **DOCHERTY et al.**FASEB J., 1994, vol. 8, 20-24 [\[0097\]](#)
- **CHEN et al.**Proc. Natl. Acad. Sci. USA, 1994, vol. 91, 3054-3057 [\[0100\]](#)
- **ROSOFF**Pharmaceutical Dosage FormsMarcel Dekker, Inc.19880000vol. 1, 245- [\[0121\]](#) [\[0159\]](#) [\[0166\]](#) [\[0167\]](#) [\[0168\]](#)
- **WANG et al.**Biochem. Biophys. Res. Commun., 1987, vol. 147, 980-985 [\[0125\]](#)
- **ZHOU et al.**Journal of Controlled Release, 1992, vol. 19, 269-274 [\[0126\]](#)
- **WEINER et al.**Journal of Drug Targeting, 1992, vol. 2, 405-410 [\[0128\]](#)
- **DU PLESSIS et al.**Antiviral Research, 1992, vol. 18, 259-265 [\[0128\]](#)
- **HU et al.**S.T.Pharma. Sci., 1994, vol. 4, 6466- [\[0129\]](#)
- **ALLEN et al.**FEBS Letters, 1987, vol. 223, 42- [\[0130\]](#)
- **WU et al.**Cancer Research, 1993, vol. 53, 3765- [\[0130\]](#)
- **PAPAHADJOPOULOS et al.**Ann. N.Y. Acad. Sci., 1987, vol. 507, 64- [\[0131\]](#)
- **GABIZON et al.**Proc. Natl. Acad. Sci. U.S.A., 1988, vol. 85, 6949- [\[0131\]](#)
- **SUNAMOTO et al.**Bull. Chem. Soc. Jpn., 1980, vol. 53, 2778- [\[0132\]](#)
- **ILLUM et al.**FEBS Lett., 1984, vol. 167, 79- [\[0132\]](#)
- **KLIBANOV et al.**FEBS Lett., 1990, vol. 268, 235- [\[0132\]](#)
- **BLUME et al.**Biochimica et Biophysica Acta, 1990, vol. 1029, 91- [\[0132\]](#)
- **RIEGER**Pharmaceutical Dosage FormsMarcel Dekker, Inc.19880000285- [\[0135\]](#) [\[0140\]](#)
- **IDSON**Pharmaceutical Dosage FormsMarcel Dekker, Inc.19880000vol. 1, 199- [\[0159\]](#) [\[0160\]](#) [\[0161\]](#) [\[0163\]](#) [\[0166\]](#) [\[0166\]](#)
- Block in Pharmaceutical Dosage FormsMarcel Dekker, Inc.19880000vol. 2, 335- [\[0159\]](#)
- **HIGUCHI et al.**Remington's Pharmaceutical SciencesMack Publishing Co.19850000301- [\[0159\]](#)
- **RIEGER**Pharmaceutical Dosage FormsMarcel Dekker, Inc.19880000vol. 1, 285- [\[0161\]](#) [\[0161\]](#)
- **BLOCK**Pharmaceutical Dosage FormsMarcel Dekker, Inc.19880000vol. 1, 335- [\[0163\]](#) [\[0168\]](#)
- **LEUNGSHAH**Controlled Release of Drugs: Polymers and Aggregate SystemsVCH Publishers19890000185-215 [\[0167\]](#)
- **SCHOTT**Remington's Pharmaceutical SciencesMack Publishing Co.19850000271- [\[0167\]](#)
- **CONSTANTINIDES et al.**Pharmaceutical Research, 1994, vol. 11, 1385-1390 [\[0170\]](#)
- **RITSCHEL**Meth. Find. Exp. Clin. Pharmacol., 1993, vol. 13, 205- [\[0170\]](#)
- **CONSTANTINIDES et al.**Pharmaceutical Research, 1994, vol. 11, 1385- [\[0170\]](#)
- **HO et al.**J. Pharm. Sci., 1996, vol. 85, 138-143 [\[0170\]](#)
- **LEE et al.**Critical Reviews in Therapeutic Drug Carrier Systems, 1991, 92- [\[0171\]](#) [\[0173\]](#) [\[0174\]](#) [\[0175\]](#) [\[0176\]](#) [\[0177\]](#) [\[0178\]](#)
- **TAKAHASHI et al.**J. Pharm. Pharmacol., 1988, vol. 40, 252- [\[0174\]](#)
- **MURANISHI**Critical Reviews in Therapeutic Drug Carrier Systems, 1990, vol. 7, 1-33 [\[0175\]](#) [\[0176\]](#) [\[0177\]](#) [\[0178\]](#)
- **EL HARIRI et al.**J. Pharm. Pharmacol., 1992, vol. 44, 651-654 [\[0175\]](#)
- **BRUNTON et al.**Goodman & Gilman's The Pharmacological Basis of TherapeuticsMcGraw-Hill19960000934-935 [\[0176\]](#)
- **SWINYARD**Remington's Pharmaceutical SciencesMack Publishing Co.19900000782-783 [\[0176\]](#)
- **YAMAMOTO et al.**J. Pharm. Exp. Ther., 1992, vol. 263, 25- [\[0176\]](#)
- **YAMASHITA et al.**J. Pharm. Sci., 1990, vol. 79, 579-583 [\[0176\]](#)
- **JARRETT**J. Chromatogr., 1993, vol. 618, 315-339 [\[0177\]](#)
- **BUUR et al.**J. Control Rel., 1990, vol. 14, 43-51 [\[0177\]](#)
- **YAMASHITA et al.**J. Pharm. Pharmacol., 1987, vol. 39, 621-626 [\[0178\]](#)
- **MIYAO et al.**DsRNA Res. Dev., 1995, vol. 5, 115-121 [\[0179\]](#)
- **TAKAKURA et al.**DsRNA & Nucl. Acid Drug Dev., 1996, vol. 6, 177-183 [\[0179\]](#)
- Amyloid, vol. 1, 17-20 [\[0266\]](#) [\[0272\]](#) [\[0262\]](#)

Patentkrav

1. Dobbeltstrenget ribonukleinsyre (dsRNA) til inhibering af
ekspression af transthyretin (TTR), hvor

5 (a) dsRNA'et omfatter en sense-streng og en antisense-streng,
idet antisense-strengen omfatter en region, der er
komplementær til en del af et mRNA, der koder for
transthyretin (TTR), hvor regionen med komplementaritet er
19 nukleotider lang, antisense-strengen omfatter SEQ ID
10 NO: 170, og hver streng af dsRNA'et er 19, 20, 21, 22, 23
eller 24 nukleotider lang; eller

(b) dsRNA'et omfatter en antisense-streng, der omfatter en
region, der er komplementær til 19 nukleotider af
nukleotiderne 618-648 af SEQ ID NO: 1331, og hvor antisense-
15 strengen baseparrer med guanin i position 628 af SEQ ID
NO: 1331;

og hvor dsRNA'et er formuleret i en nukleinsyre-lipidpartikel,
der omfatter et kationisk lipid, et ikke-kationisk lipid og et
lipid, der forhindrer aggregering af partiklen.

20

2. dsRNA ifølge krav 1, hvor det kationiske lipid er
(6Z,9Z,28Z,31Z)-heptatriaconta-6,9,28,31-tetraen-19-yl-4-
(dimethylamino)-butanoat (MC3).

25 3. dsRNA ifølge krav 1, hvor det ikke-kationiske lipid er
distearoylphosphatidylcholin (DSPC).

4. dsRNA ifølge krav 1, hvor lipidet, der forhindrer
aggregering af partiklen, er et polyethylenglycol (PEG)-lipid.

30

5. dsRNA ifølge et hvilket som helst af kravene 1 til 4,
hvor formuleringen yderligere omfatter kolesterol.

6. dsRNA ifølge krav 1, hvor formuleringen omfatter MC3,
35 DSPC, kolesterol og PEG₂₀₀₀-C14.

7. dsRNA ifølge krav 1, hvor formuleringen omfatter
MC3/DSPC/kolesterol/PEG₂₀₀₀-C14 i et forhold på

50/10/38,5/1,5 mol-%.

8. dsRNA ifølge et hvilket som helst af kravene 1 til 7, hvor

- 5 (a) sense-strengen består af SEQ ID NO: 449, og antisense-strengen består af SEQ ID NO: 450;
(b) sense-strengen består af SEQ ID NO: 729, og antisense-strengen består af SEQ ID NO: 730; eller
(c) sense-strengen består af SEQ ID NO: 1009, og antisense-
10 strengen består af SEQ ID NO: 1010.

9. dsRNA ifølge et hvilket som helst af kravene 1 til 7, hvor sense-strengen omfatter SEQ ID NO: 169.

- 15 10. dsRNA ifølge et hvilket som helst af kravene 1 til 7, hvor dsRNA'et omfatter mindst ét modificeret nukleotid, hvor fortrinsvis mindst ét af de modificerede nukleotider er valgt fra gruppen, der består af: et 2'-O-methylmodificeret nukleotid, et nukleotid, der omfatter en 5'-
20 phosphorthioatgruppe, et terminalt nukleotid, der er koblet til et cholesterylderivat eller en dodecansyrebisdecylamidgruppe, et 2'-deoxy-2'-fluormodificeret nukleotid, et 2'-deoxymodificeret nukleotid, et låst nukleotid, et abasisk nukleotid, et 2'-aminomodificeret
25 nukleotid, et 2'-alkylmodificeret nukleotid, et morpholinonukleotid, et phosphoramidat og et nukleotid, der omfatter en ikke-naturlig base.

11. dsRNA ifølge et hvilket som helst af kravene 1 til 10,
30 hvor dsRNA'et er konjugeret til en ligand.

12. Farmaceutisk sammensætning til inhibering af ekspres-
af et TTR-gen, der omfatter dsRNA'et ifølge et hvilket som
helst af kravene 1 til 11 og et farmaceutisk acceptabelt
35 bæremiddel.

13. Fremgangsmåde til inhibering af TTR-ekspres-
sion i en celle, hvilken fremgangsmåde omfatter:

(a) etablering af kontakt mellem cellen og dsRNA'et ifølge et hvilket som helst af kravene 1 til 11; og

(b) opretholdelse af cellen dannet i trin (a) i et tidsrum, der er tilstrækkeligt til opnåelse af degradering af mRNA-transkriptet af et TTR-gen til derved inhibering af

ekspression af TTR-genet i cellen;
forudsat at enhver metode til behandling af menneske- eller dyrekroppen ved hjælp af terapi er eksklusive.

14. dsRNA ifølge et hvilket som helst af kravene 1 til 11 til anvendelse til behandling hos et menneske af en forstyrrelse, der medieres af TTR-ekspression, hvor

(a) mennesket har transthyretin-amyloidose; og/eller

(b) mennesket har en leverforstyrrelse,

hvor transthyretin-amyloidosen fortrinsvis er valgt blandt familiær amyloidotisk polyneuropati (FAP), familiær amyloidotisk kardiomyopati (FAC), leptomeningeal/CNS-amyloidose, senil systemisk amyloidose (SSA) og senil kardiel amyloidose (SCA).

15. dsRNA til anvendelse ifølge krav 14, hvor dsRNA'et er til administration til mennesket ved ca. 0,01, 0,1, 0,2, 0,3, 0,4, 0,5, 0,543, 0,59, 0,6, 0,7, 0,8, 0,9, 1,0, 1,1, 1,2, 1,3, 1,4, 2,5 eller 5,0 mg/kg.


16. dsRNA til anvendelse ifølge krav 14 eller 15, hvor mennesket yderligere modtager en terapeutisk metode til behandling af en TTR-amyloidose, hvilken terapeutiske metode er valgt fra gruppen, der består af diuretika, inhibitorer af angiotensinkonverterende enzym, angiotensinreceptorblokkere, dialyse og et levertransplantat.

17. dsRNA til anvendelse ifølge et hvilket som helst af kravene 14 til 16, hvor dsRNA'et er til administration til mennesket med intervaller på højst 1, 2, 3, eller 4 uger.

18. dsRNA til anvendelse ifølge et hvilket som helst af kravene 14 til 17, hvor dsRNA'et er til intravenøs



administration eller intravenøs infusion over 15 minutter.


DRAWINGS



candidate siRNAs

FIG. 1

**FIG. 2A****FIG. 2B**

**FIG. 3**




FIG. 4A




FIG. 4B





FIG. 5






FIG. 6B




FIG. 7A




FIG. 7B

SNALP-18328




FIG. 9




FIG. 9




FIG. 10




FIG. 11A





FIG. 11B



Lanes 1, 3, 5, 7, 9, 11, 13, and 15: PBS animals
Lanes 2, 4, 6, 8, 10, 12, 14, and 16: 0.3 mg/kg LNP09-18328

FIG. 11C

```

gttgactaag tcaataatca gaatcagcag
gtttgcagtc agattggcag ggataagcag
cctagctcag gagaagtgag tataaaagcc
ccaggctggg agcagccatc acagaagtcc
actcattctt ggcaggatgg cttctcatcg
tctgctcctc ctctgccttg ctggactggt
at ttgtgtct gaggctggcc ctacgggcac
cggtgaatcc aagtgtcctc tgatggtcaa
agttctagat gctgtccgag gcagtcctgc
catcaatgtg gccgtgcatg tgttcagaaa
ggctgctgat gacacctggg agccatttgc
ctctgggaaa accagtgagt ctggagagct
gcatgggctc acaactgagg aggaatttgt
agaagggata tacaaagtgg aaatagacac
caaatcttac tggaaggcac ttggcatctc
cccattccat gagcatgcag aggtggtatt
cacagccaac gactccggcc cccgccgcta
caccattgcc gccctgctga gcccctactc
ctattccacc acggctgtcg tcaccaatcc
caaggaatga gggacttctc ctccagtgga
cctgaaggac gagggatggg atttcatgta
accaagagta ttccattttt actaaagcag
tgttttcacc tcatatgcta tgttagaagt
ccaggcagag acaataaaaac attcctgtga
aaggcacttt tcattccact ttaacttgat
tttttaaatt cccttattgt cccttccaaa
aaaaagagaa tcaaaatttt acaaagaatc
aaaggaattc tagaaagtat ctgggcagaa
cgctaggaga gatccaaatt tccattgtct
tgcaagcaaa gcacgtatta aatatgatct
gcagccatta aaaagacaca ttctgtaaaa
aaaaaaaa (SEQ ID NO: 1331)

```

FIG. 12

FIG. 13A

ACAGAAGTCCACTCATTCTTGGCAGGATGGCTTCTCATCGTCTGCTCCTCCT
CTGCCTTGCTGGACTGGTATTTGTGTCTGAGGCTGGCCCTACGGGCACCGGT
GAATCCAAGTGTCTCTGATGGTCAAAGTTCTAGATGCTGTCCGAGGCAGTC
CTGCCATCAATGTGGCCGTGCATGTGTTTCAGAAAGGCTGCTGATGACACCTG
GGAGCCATTTGCCTCTGGGAAAACCAAGTGAGTCTGGAGAGCTGCATGGGCTC
ACAACTGAGGAGGAATTTGTAGAAAGGATATACAAAGTGGAAATAGACACCA
AATCTTACTGGAAGGCACTTGGCATCTCCCCATTCCATGAGCATGCAGAGGT
GGTATTCACAGCCAACGACTCCGGCCCCCGCCGCTACACCATTGCCGCCCTG
CTGAGCCCCTACTCCTATTCCACCACGGCTGTCGTCACCAATCCCAAGGAAT
GAGGGACTTCTCCTCCAGTGGACCTGAAGGACGAGGGATGGGATTTTCATGTA
ACCAAGAGTATTCATTTTTACTAAAGCAGTGTTCACCTCATATGCTATG
TTAGAAGTCCAGGCAGAGACAATAAAACATTCTGTGAAAGGCACTTTTCAT
TCCAAAAAAAAAAAAAAAAAAAAA (SEQ ID NO:1329)

FIG. 13B

CCTGACAGGATGGCTTCCCTTCGCCTGTTCTCCTCTGCCTCGCTGGACTGA
TATTTGCGTCTGAAGCTGGCCCTGGGGGTGCTGGAGAATCCAAGTGTCTCT
GATGGTCAAAGTCTTGGATGCTGTCCGAGGCAGCCCTGCTGTGATGTGGCC
GTGAAAGTGTTCAAAAGGACTGCAGACGGAAGCTGGGAGCCGTTTGCCTCTG
GGAAGACCGCCGAGTCTGGAGAGCTGCACGGGCTCACCACAGATGAGAAGTT
CACGGAAGGGGTGTACAGGGTAGAACTGGACACCAAATCATACTGGAAGGCT
CTTGGCATTTCCCCATTCCATGAATACGCAGAGGTGGTTTTTCACAGCCAATG
ACTCTGGTCATCGCCACTACACCATCGCAGCCCTGCTCAGCCCGTACTCCTA
CAGCACCCTGCTGTGTCAGTAACCCCCAGAACTGAGGGACCCAGCCACG
AGGACCAAGATCTTGCCAAAGCAGTAGCTCCCATTGTACTGAAACAGTGT
CTTGCTCTATAAACCGTGTTAGCAACTCGGGAAGATGCCGTGAAACGTTCTT
ATTAAACCACCTTTATTTCATTC
(SEQ ID NO:1330)

NM_000371.3 GTTGACTAAGTCAATAATCAGAATCAGCAGGTTTCGACTCAGATTGGCAGGGATAAGCAGCCTAGCTC 68
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 AGGAGAAGTCAGCTATAAAAGCCCCAGCTGGGAGCAGCCATCACAGAAAGTCCACTEATTCTTGGCAGG 136
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 ATGGCTTTCATCGTCTGCTCTCTCTCTGCTTCTCGACTGCTATTCTGTCTGAGGCTGGCCCTAC 204
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 GGGCACCGGTGAATCCAAAGTCTCTCTGATGCTCAAAGTTCTAGATGCTGTCGAGGCAGTCTTCCCA 272
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 TCAATGTGGCGCTCATGTCTTCAGAAAGCTGCTGTATGACACCTGGGAGCCATTTCCTCTGGGAAA 340
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 ACCAGTGAGTCTGGAGAGCTGCATGGCTCACAACGAGGAGGAATTTGAGAAAGGATATACAAAGT 408
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 GGAAATAGACACCAAACTCTTACTGGAAGGCATTTGGCATCTCCCATTCATGAGATGAGAGGTGG 476
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 TATTCAAGCCAAAGCACTCCGGCCCCCGCCGTACACATTGCCGCCCTGCTGAGCCCTACTCTCTAT 544
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 TCCACCACGGCTGTGCTCAACATCCCAAGGAATCAGGAGCTTCTCTCAAGTGACCTGAAGGACGA 612
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 GGGATGGGATTTCACTAACCAGAGTATTCATTTTACTAAGCAGCTTTTCACTCATATGCTA 680
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 TGTTAGAAGTCCAGGCAGAGACAATAAAACATTCTGTGAAAGCACTTTTCACTTCACTTAACTTG 748
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 ATTTTTAAATTCCTTATTGTCCTTCCAAAAAAGAGAAATCAAATTTTACAAGAATCAAAGCA 816
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 ATTCTAGAAAGTATCTGGGCAAGACCTAGGAGAGATCCAAATTTCACTGTCTTGCAGGCAAAAGCAC 884
 NM_000371.2 -----
 AD-18328_sense -----

NM_000371.3 GTATTAAATATGATCTCGAGCCATTAAAAAGACACATTCTGTAAAAAATAAATA 938
 NM_000371.2 -----
 AD-18328_sense -----

FIG. 14

Phenotype	Features	Genotypes (associated mutation in TTR)
Familial amyloidotic neuropathy (FAP)	Early: Impotence Sensorimotor polyneuropathy of the legs Carpal tunnel syndrome Autonomic dysfunction Constipation/diarrhea Late: Cardiomyopathy Vitreous opacities Nephropathy	V28M L58H V30M L58R K70N Y78F I84S Y114H V30A K35N G47V S50R T60A Y114C
Familial amyloidotic cardiomyopathy (FAC)	Cardiomegaly Congestive heart failure Conduction abnormalities, arrhythmias Angina Sudden death	D18N D18E V20I P24S E42D A45T T49P S50I H56R I68L A81T Q92K R103S L111M V122I T60A
CNS amyloidosis (CNSA)	Dementia, ataxia, spasticity, seizures, hemorrhage (Intracerebellar and/or subarachnoid), psychosis, hydrocephalus	L12P D18G A25T V30G A36P G53E F64S Y69H Y114C

FIG. 15




FIG. 16




FIG. 17

