




(86) Date de dépôt PCT/PCT Filing Date: 2012/11/15
(87) Date publication PCT/PCT Publication Date: 2013/05/23
(85) Entrée phase nationale/National Entry: 2014/03/19
(86) N° demande PCT/PCT Application No.: EP 2012/072699
(87) N° publication PCT/PCT Publication No.: 2013/072406
(30) Priorités/Priorities: 2011/11/15 (US61/560,144);
2011/11/15 (US61/560,183); 2011/11/15 (US61/560,178);
2011/11/15 (US61/560,162); 2011/11/15 (US61/560,149);
2012/05/24 (US61/651,474); 2012/05/24 (US61/651,486)

(51) Cl.Int./Int.Cl. *C07K 16/28* (2006.01),
A61K 39/395 (2006.01), *A61P 35/02* (2006.01),
A61P 37/00 (2006.01)
(71) Demandeurs/Applicants:
AMGEN RESEARCH (MUNICH) GMBH, DE;
BOEHRINGER INGELHEIM INTERNATIONAL GMBH,
DE
(72) Inventeurs/Inventors:
KUFER, PETER, DE;
RAUM, TOBIAS, DE;
HOFFMANN, PATRICK, DE;
KISCHEL, ROMAN, DE;
...

(54) Titre : MOLECULES DE LIAISON POUR BCMA ET CD3
(54) Title: BINDING MOLECULES FOR BCMA AND CD3

Figure 16



(57) Abrégé/Abstract:

The present invention relates to a binding molecule which is at least bispecific comprising a first and a second binding domain, wherein the first binding domain is capable of binding to epitope cluster3 of BCMA, and the second binding domain is

(72) **Inventeurs(suite)/Inventors(continued)**: LUTTERBUESE, RALF, DE; RAU, DORIS, DE; ADAM, PAUL, DE; BORGES, ERIC, DE; HEBEIS, BARBARA, DE; HIPPEL, SUSANNE, DE

(74) **Agent**: GOWLING LAFLEUR HENDERSON LLP

(57) **Abrégé(suite)/Abstract(continued)**:

capable of binding to the Tcell CD3 receptor complex. Moreover, the invention provides a nucleic acid sequence encoding the binding molecule, a vector comprising said nucleic acid sequence and a host cell transformed or transfected with said vector. Furthermore, the invention provides a process for the production of the binding molecule of the invention, a medical use of said binding molecule and a kit comprising said binding molecule.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

23 May 2013 (23.05.2013)



(10) International Publication Number

WO 2013/072406 A1

(51) International Patent Classification:

C07K 16/28 (2006.01) A61P 35/02 (2006.01)

A61K 39/395 (2006.01) A61P 37/00 (2006.01)

(21) International Application Number:

PCT/EP2012/072699

(22) International Filing Date:

15 November 2012 (15.11.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/560,144 15 November 2011 (15.11.2011) US

61/560,149 15 November 2011 (15.11.2011) US

61/560,162 15 November 2011 (15.11.2011) US

61/560,178 15 November 2011 (15.11.2011) US

61/560,183 15 November 2011 (15.11.2011) US

61/651,486 24 May 2012 (24.05.2012) US

61/651,474 24 May 2012 (24.05.2012) US

(71) Applicants: AMGEN RESEARCH (MUNICH) GMBH

[DE/DE]; Staffelsestr. 2, 81477 Munich (DE).

BOEHRINGER INGELHEIM INTERNATIONAL

GMBH [DE/DE]; Binger Strasse 173, 55216 Ingelheim

am Rhein (DE).

(72) Inventors: KUFER, Peter; c/o Amgen Research (Munich)

GmbH, Staffelsestr. 2, 81477 Munich (DE). RAUM, To-

bias; c/o Amgen Research (Munich) GmbH, Staffelsestr.

2, 81477 Munich (DE). HOFFMANN, Patrick; c/o Am-

gen Research (Munich) GmbH, Staffelsestr. 2, 81477 Munich (DE). KISCHEL, Roman; c/o Amgen Research (Munich) GmbH, Staffelsestr. 2, 81477 Munich (DE). LUTTERBUESE, Ralf; c/o Amgen Research (Munich) GmbH, Staffelsestr. 2, 81477 Munich (DE). RAU, Doris; c/o Amgen Research (Munich) GmbH, Staffelsestr. 2, 81477 Munich (DE). ADAM, Paul; c/o Boehringer Ingelheim GmbH, Corporate Patents, Binger Strasse 173, 55216 Ingelheim am Rhein (DE). BORGES, Eric; c/o Boehringer Ingelheim GmbH, Corporate Patents, Binger Strasse 173, 55216 Ingelheim am Rhein (DE). HEBEIS, Barbara; c/o Boehringer Ingelheim GmbH, Corporate Patents, Binger Strasse 173, 55216 Ingelheim am Rhein (DE). HIPPE, Susanne; c/o Boehringer Ingelheim GmbH, Corporate Patents, Binger Strasse 173, 55216 Ingelheim am Rhein (DE).


(74) Agents: WEINZIERL, Gerhard et al.; Schiweck Weinzierl Koch, European Patent Attorneys, Landsberger Str. 98, 80339 Munich (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,

[Continued on next page]

(54) Title: BINDING MOLECULES FOR BCMA AND CD3

Figure 16



(57) Abstract: The present invention relates to a binding molecule which is at least bispecific comprising a first and a second binding domain, wherein the first binding domain is capable of binding to epitope cluster3 of BCMA, and the second binding domain is capable of binding to the Tcell CD3 receptor complex. Moreover, the invention provides a nucleic acid sequence encoding the binding molecule, a vector comprising said nucleic acid sequence and a host cell transformed or transfected with said vector. Furthermore, the invention provides a process for the production of the binding molecule of the invention, a medical use of said binding molecule and a kit comprising said binding molecule.

WO 2013/072406 A1

WO 2013/072406 A1

TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.	SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
(84) Designated States <i>(unless otherwise indicated, for every kind of regional protection available):</i> ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,	Published: — <i>with international search report (Art. 21(3))</i> — <i>before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))</i>

5

Binding molecules for BCMA and CD3

The present invention relates to a binding molecule which is at least bispecific comprising a first and a second binding domain, wherein the first binding domain is capable of binding to epitope
10 cluster 3 of BCMA, and the second binding domain is capable of binding to the T cell CD3 receptor complex. Moreover, the invention provides a nucleic acid sequence encoding the binding molecule, a vector comprising said nucleic acid sequence and a host cell transformed or transfected with said vector. Furthermore, the invention provides a process for the production of the binding molecule of the invention, a medical use of said binding molecule and a kit
15 comprising said binding molecule.

BCMA (B-cell maturation antigen, TNFRSF17, CD269) is a transmembrane protein belonging to the TNF receptor super family. BCMA is originally reported as an integral membrane protein in the Golgi apparatus of human mature B lymphocytes, i.e., as an intracellular protein (Gras et al.,
20 (1995) International Immunol 7(7):1093-1105) showing that BCMA seems to have an important role during B-cell development and homeostasis. The finding of Gras et al. might be associated with the fact that the BCMA protein that was described in Gras et al. is, because of a chromosomal translocation, a fusion protein between BCMA and IL-2. Meanwhile BCMA is, however, established to be a B-cell marker that is essential for B-cell development and
25 homeostasis (Schliemann et al., (2001) Science 293 (5537):2111-2114) due to its presumably essential interaction with its ligands BAFF (B cell activating factor), also designated as TALL-1 or TNFSF13B, and APRIL (A proliferation-inducing ligand).

BCMA expression is restricted to the B-cell lineage and mainly present on plasma cells and plasmablasts and to some extent on memory B-cells, but virtually absent on peripheral and
30 naive B-cells. BCMA is also expressed on multiple myeloma (MM) cells. Together with its family members transmembrane activator and cyclophilin ligand interactor (TACI) and B cell activation factor of TNF family receptor (BAFF-R), BCMA regulates different aspects of humoral immunity, B-cell development and homeostasis. Expression of BCMA appears rather late in B-cell differentiation and contributes to the long term survival of plasmablasts and plasma cells in the
35 bone marrow. Targeted deletion of the BCMA gene in mice does not affect the generation of mature B-cells, the quality and magnitude of humoral immune responses, formation of germinal

center and the generation of short-lived plasma cells. However, such mice have significantly reduced numbers of long-lived plasma cells in the bone marrow, indicating the importance of BCMA for their survival (O'Connor et al., 2004).

In line with this finding, BCMA also supports growth and survival of multiple myeloma (MM) cells. Novak et al found that MM cell lines and freshly isolated MM cells express BCMA and TACI protein on their cell surfaces and have variable expression of BAFF-R protein on their cell surface (Novak et al., (2004) Blood 103(2):689-694).

Multiple myeloma (MM) is the second most common hematological malignancy and constitutes 2% of all cancer deaths. MM is a heterogenous disease and caused by mostly by chromosome translocations *inter alia* t(11;14),t(4;14),t(8;14),del(13),del(17) (Drach et al., (1998) Blood 92(3):802-809; Gertz et al., (2005) Blood 106(8):2837-2840; Facon et al., (2001) Blood 97(6):1566-1571). MM-affected patients may experience a variety of disease-related symptoms due to, bone marrow infiltration, bone destruction, renal failure, immunodeficiency, and the psychosocial burden of a cancer diagnosis. As of 2006, the 5-year relative survival rate for MM was approximately 34% highlighting that MM is a difficult-to-treat disease where there are currently no curative options.

Exciting new therapies such as chemotherapy and stem cell transplantation approaches are becoming available and have improved survival rates but often bring unwanted side effects, and thus MM remains still incurable (Lee et al., (2004) J Natl Compr Canc Netw 8 (4): 379-383). To date, the two most frequently used treatment options for patients with multiple myeloma are combinations of steroids, thalidomide, lenalidomide, bortezomib or various cytotoxic agents, and for younger patients high dose chemotherapy concepts with autologous stem cell transplantation.

Most transplants are of the autologous type, i.e. using the patient's own cells. Such transplants, although not curative, have been shown to prolong life in selected patients. They can be performed as initial therapy in newly diagnosed patients or at the time of relapse. Sometimes, in selected patients, more than one transplant may be recommended to adequately control the disease.

Chemotherapeutic agents used for treating the disease are Cyclophosphamid, Doxorubicin, Vincristin and Melphalan, combination therapies with immunomodulating agents such as thalidomide (Thalomid®), lenalidomide (Revlimid®), bortezomib (Velcade®) and corticosteroids (e.g. Dexamethasone) have emerged as important options for the treatment of myeloma, both in

newly diagnosed patients and in patients with advanced disease in whom chemotherapy or transplantation have failed.

The currently used therapies are usually not curative. Stem cell transplantation may not be an option for many patients because of advanced age, presence of other serious illness, or other physical limitations. Chemotherapy only partially controls multiple myeloma, it rarely leads to complete remission. Thus, there is an urgent need for new, innovative treatments.

Bellucci et al. (Blood, 2005; 105(10) identified BCMA-specific antibodies in multiple myeloma patients after they had received donor lymphocyte infusions (DLI). Serum of these patients was capable of mediating BCMA-specific cell lysis by ADCC and CDC and was solely detected in patients with anti-tumor responses (4/9), but not in non-responding patients (0/6). The authors speculate that induction of BCMA-specific antibodies contributes to elimination of myeloma cells and long-term remission of patients.

Ryan et al. (Mol. Cancer Ther. 2007; 6(11) reported the generation of an antagonistic BCMA-specific antibody that prevents NF- κ B activation which is associated with a potent pro-survival signaling pathway in normal and malignant B-cells. In addition, the antibody conferred potent antibody-dependent cell-mediated cytotoxicity (ADCC) to multiple myeloma cell lines in vitro which was significantly enhanced by Fc-engineering.

Other approaches in fighting blood-borne tumors or autoimmune disorders focus on the interaction between BAFF and APRIL, i.e., ligands of the TNF ligand super family, and their receptors TACI, BAFF-R and BCMA which are activated by BAFF and/or APRIL. For example, by fusing the Fc-domain of human immunoglobulin to TACI, Zymogenetics, Inc. has generated Atacicept (TACI-Ig) to neutralize both these ligands and prevent receptor activation. Atacicept is currently in clinical trials for the treatment of Systemic Lupus Erythematosus (SLE, phase III), multiple sclerosis (MS, phase II) and rheumatoid arthritis (RA, phase II), as well as in phase I clinical trials for the treatment of the B-cell malignancies chronic lymphocytic leukaemia (CLL), non-Hodgkins lymphoma (NHL) and MM. In preclinical studies atacicept reduces growth and survival of primary MM cells and MM cell lines in vitro (Moreaux et al, Blood, 2004, 103) and in vivo (Yaccoby et al, Leukemia, 2008, 22, 406-13), demonstrating the relevance of TACI ligands for MM cells. Since most MM cells and derived cell lines express BCMA and TACI, both receptors might contribute to ligand-mediated growth and survival. These data suggest that antagonizing both BCMA and TACI might be beneficial in the treatment of plasma cell disorders.

In addition, BCMA-specific antibodies that cross react with TACI have been described (WO 02/066516).

Human Genome Sciences and GlaxoSmithKline have developed an antibody targeting BAFF which is called Belimumab. Belimumab blocks the binding of soluble BAFF to its receptors BAFF-R, BCMA and TACI on B cells. Belimumab does not bind B cells directly, but by binding BAFF, belimumab inhibits the survival of B cells, including autoreactive B cells, and reduces the differentiation of B cells into immunoglobulin-producing plasma cells.

Nevertheless, despite the fact that BCMA; BAFF-R and TACI, i.e., B cell receptors belonging to the TNF receptor super family, and their ligands BAFF and APRIL are subject to therapies in fighting against cancer and/or autoimmune disorders, there is still a need for having available further options for the treatment of such medical conditions.

Accordingly, there is provided herewith means and methods for the solution of this problem in the form of a binding molecule which is at least bispecific with one binding domain to cytotoxic cells, i.e., cytotoxic T cells, and with a second binding domain to BCMA.

Thus, in a first aspect the present invention provides a binding molecule which is at least bispecific comprising a first and a second binding domain, wherein

(a) the first binding domain is capable of binding to epitope cluster 3 of BCMA (CQLRCSSNTPPLTCQRYC) (SEQ ID NO:1016); and

(b) the second binding domain is capable of binding to the T cell CD3 receptor complex; and wherein epitope cluster 3 of BCMA corresponds to amino acid residues 24 to 41 of the sequence as depicted in SEQ ID NO: 1002.

It must be noted that as used herein, the singular forms "a", "an", and "the", include plural references unless the context clearly indicates otherwise. Thus, for example, reference to "a reagent" includes one or more of such different reagents and reference to "the method" includes reference to equivalent steps and methods known to those of ordinary skill in the art that could be modified or substituted for the methods described herein.

Unless otherwise indicated, the term "at least" preceding a series of elements is to be understood to refer to every element in the series. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific

embodiments of the invention described herein. Such equivalents are intended to be encompassed by the present invention.

5 The term "and/or" wherever used herein includes the meaning of "and", "or" and "all or any other combination of the elements connected by said term".

The term "about" or "approximately" as used herein means within $\pm 20\%$, preferably within $\pm 15\%$, more preferably within $\pm 10\%$, and most preferably within $\pm 5\%$ of a given value or range.

10 Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integer or step. When used herein the term "comprising" can be substituted with the term "containing" or "including" or sometimes when used herein with
15 the term "having".

When used herein "consisting of" excludes any element, step, or ingredient not specified in the claim element. When used herein, "consisting essentially of" does not exclude materials or steps that do not materially affect the basic and novel characteristics of the claim.

20 In each instance herein any of the terms "comprising", "consisting essentially of" and "consisting of" may be replaced with either of the other two terms.

Epitope cluster 3 is comprised by the extracellular domain of BCMA. The "BCMA extracellular
25 domain" or "BCMA ECD" refers to a form of BCMA which is essentially free of transmembrane and cytoplasmic domains of BCMA. It will be understood by the skilled artisan that the transmembrane domain identified for the BCMA polypeptide of the present invention is identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than
30 about 5 amino acids at either end of the domain specifically mentioned herein. A preferred BCMA ECD is shown in SEQ ID NO: 1007.

The T cell CD3 receptor complex is a protein complex and is composed of four distinct chains. In mammals, the complex contains a CD3 γ chain, a CD3 δ chain, and two CD3 ϵ (epsilon)

chains. These chains associate with a molecule known as the T cell receptor (TCR) and the ζ chain to generate an activation signal in T lymphocytes.

The redirected lysis of target cells via the recruitment of T cells by bispecific molecules involves
5 cytolytic synapse formation and delivery of perforin and granzymes. The engaged T cells are capable of serial target cell lysis, and are not affected by immune escape mechanisms interfering with peptide antigen processing and presentation, or clonal T cell differentiation; see, for example, WO 2007/042261.

10 The term "binding molecule" in the sense of the present disclosure indicates any molecule capable of (specifically) binding to, interacting with or recognizing the target molecules BCMA and CD3. According to the present invention, binding molecules are preferably polypeptides. Such polypeptides may include proteinaceous parts and non-proteinaceous parts (e.g. chemical linkers or chemical cross-linking agents such as glutaraldehyde).

15 A binding molecule, so to say, provides the scaffold for said one or more binding domains so that said binding domains can bind/interact with the target molecules BCMA and CD3. For example, such a scaffold could be provided by protein A, in particular, the Z-domain thereof (affibodies), ImmE7 (immunity proteins), BPTI/APPI (Kunitz domains), Ras-binding protein AF-6 (PDZ-domains), charybdotoxin (Scorpion toxin), CTLA-4, Min-23 (knottins), lipocalins
20 (anticalins), neokarzinostatin, a fibronectin domain, an ankyrin consensus repeat domain or thioredoxin (Skerra, Curr. Opin. Biotechnol. 18, 295-304 (2005); Hosse et al., Protein Sci. 15, 14-27 (2006); Nicaise et al., Protein Sci. 13, 1882-1891 (2004) ; Nygren and Uhlen, Curr. Opin. Struc. Biol. 7, 463-469 (1997)). A preferred binding molecule is an antibody.

It is envisaged that the binding molecule is produced by (or obtainable by) phage-display or
25 library screening methods rather than by grafting CDR sequences from a pre-existing (monoclonal) antibody into a scaffold, for example, a scaffold as disclosed herein.

The term "bispecific" as used herein refers to a binding molecule which comprises at least a first and a second binding domain, wherein the first binding domain is capable of binding to one
30 antigen or target, and the second binding domain is capable of binding to another antigen or target. The "binding molecule" of the invention also comprises multispecific binding molecules such as e.g. trispecific binding molecules, the latter ones including three binding domains.

It is also envisaged that the binding molecule of the invention has, in addition to its function to
35 bind to the target molecules BCMA and CD3, a further function. In this format, the binding

molecule is a tri-or multifunctional binding molecule by targeting plasma cells through binding to BCMA, mediating cytotoxic T cell activity through CD3 binding and providing a further function such as a fully functional Fc constant domain mediating antibody-dependent cellular cytotoxicity through recruitment of effector cells like NK cells, a label (fluorescent etc.), a therapeutic agent
5 such as, e.g. a toxin or radionuclide, and/or means to enhance serum half-life, etc.

The term "binding domain" characterizes in connection with the present invention a domain which is capable of specifically binding to / interacting with a given target epitope or a given target site on the target molecules BCMA and CD3.

10 Binding domains can be derived from a binding domain donor such as for example an antibody, protein A, ImmE7 (immunity proteins), BPTI/APPI (Kunitz domains), Ras-binding protein AF-6 (PDZ-domains), charybdotoxin (Scorpion toxin), CTLA-4, Min-23 (knottins), lipocalins (anticalins), neokarzinostatin, a fibronectin domain, an ankyrin consensus repeat domain or thioredoxin (Skerra, Curr. Opin. Biotechnol. 18, 295-304 (2005); Hosse et al., Protein Sci. 15,
15 14-27 (2006); Nicaise et al., Protein Sci. 13, 1882-1891 (2004) ; Nygren and Uhlen, Curr. Opin. Struc. Biol. 7, 463-469 (1997)). A preferred binding domain is derived from an antibody. It is envisaged that a binding domain of the present invention comprises at least said part of any of the aforementioned binding domains that is required for binding to/interacting with a given target epitope or a given target site on the target molecules BCMA and CD3.

20 It is envisaged that the binding domain of the aforementioned binding domain donors is characterized by that part of these donors that is responsible for binding the respective target, i.e. when that part is removed from the binding domain donor, said donor loses its binding capability. "Loses" means a reduction of at least 50% of the binding capability when compared
25 with the binding donor. Methods to map these binding sites are well known in the art – it is therefore within the standard knowledge of the skilled person to locate/map the binding site of a binding domain donor and, thereby, to "derive" said binding domain from the respective binding domain donors.

30 The term "epitope" refers to a site on an antigen to which a binding domain, such as an antibody or immunoglobulin or derivative or fragment of an antibody or of an immunoglobulin, specifically binds. An "epitope" is antigenic and thus the term epitope is sometimes also referred to herein as "antigenic structure" or "antigenic determinant". Thus, the binding domain is an "antigen-
35 interaction-site". Said binding/interaction is also understood to define a "specific recognition". In one example, said binding domain which (specifically) binds to / interacts with a given target

epitope or a given target site on the target molecules BCMA and CD3 is an antibody or immunoglobulin, and said binding domain is a VH and/or VL region of an antibody or of an immunoglobulin.

“Epitopes” can be formed both by contiguous amino acids or non-contiguous amino acids juxtaposed by tertiary folding of a protein. A "linear epitope" is an epitope where an amino acid primary sequence comprises the recognized epitope. A linear epitope typically includes at least 3 or at least 4, and more usually, at least 5 or at least 6 or at least 7, for example, about 8 to about 10 amino acids in a unique sequence.

A "conformational epitope", in contrast to a linear epitope, is an epitope wherein the primary sequence of the amino acids comprising the epitope is not the sole defining component of the epitope recognized (e.g., an epitope wherein the primary sequence of amino acids is not necessarily recognized by the binding domain). Typically a conformational epitope comprises an increased number of amino acids relative to a linear epitope. With regard to recognition of conformational epitopes, the binding domain recognizes a three-dimensional structure of the antigen, preferably a peptide or protein or fragment thereof (in the context of the present invention, the antigen for one of the binding domains is comprised within the BCMA protein). For example, when a protein molecule folds to form a three-dimensional structure, certain amino acids and/or the polypeptide backbone forming the conformational epitope become juxtaposed enabling the antibody to recognize the epitope. Methods of determining the conformation of epitopes include, but are not limited to, x-ray crystallography, two-dimensional nuclear magnetic resonance (2D-NMR) spectroscopy and site-directed spin labelling and electron paramagnetic resonance (EPR) spectroscopy. Moreover, the provided examples describe a further method to test whether a given binding domain binds to one or more epitope cluster(s) of a given protein, in particular BCMA.

In one aspect, the first binding domain of the present invention is capable of binding to epitope cluster 3 of human BCMA, preferably human BCMA ECD. Accordingly, when the respective epitope cluster in the human BCMA protein is exchanged with the respective epitope cluster of a murine BCMA antigen (resulting in a construct comprising human BCMA, wherein human epitope cluster 3 is replaced with murine epitope cluster 3; see SEQ ID NO: 1011), a decrease in the binding of the binding domain will occur. Said decrease is preferably at least 10%, 20%, 30%, 40%, 50%; more preferably at least 60%, 70%, 80%, 90%, 95% or even 100% in comparison to the respective epitope cluster in the human BCMA protein, whereby binding to the respective epitope cluster in the human BCMA protein is set to be 100%. It is envisaged that the aforementioned human BCMA/murine BCMA chimeras are expressed in CHO cells. It is

also envisaged that the human BCMA/murine BCMA chimeras are fused with a transmembrane domain and/or cytoplasmic domain of a different membrane-bound protein such as EpCAM; see Figure 2a.

A method to test this loss of binding due to exchange with the respective epitope cluster of a non-human (e.g. murine) BCMA antigen is described in the appended Examples, in particular in Examples 1-3. A further method to determine the contribution of a specific residue of a target antigen to the recognition by a given binding molecule or binding domain is alanine scanning (see e.g. Morrison KL & Weiss GA. Cur Opin Chem Biol. 2001 Jun;5(3):302-7), where each residue to be analyzed is replaced by alanine, e.g. via site-directed mutagenesis. Alanine is used because of its non-bulky, chemically inert, methyl functional group that nevertheless mimics the secondary structure references that many of the other amino acids possess. Sometimes bulky amino acids such as valine or leucine can be used in cases where conservation of the size of mutated residues is desired. Alanine scanning is a mature technology which has been used for a long period of time.

As used herein, the term "epitope cluster" denotes the entirety of epitopes lying in a defined contiguous stretch of an antigen. An epitope cluster can comprise one, two or more epitopes. The epitope clusters that were defined – in the context of the present invention – in the extracellular domain of BCMA are described above and depicted in Figure 1.

The terms "(capable of) binding to", "specifically recognizing", "directed to" and "reacting with" mean in accordance with this invention that a binding domain is capable of specifically interacting with one or more, preferably at least two, more preferably at least three and most preferably at least four amino acids of an epitope.

As used herein, the terms "specifically interacting", "specifically binding" or "specifically bind(s)" mean that a binding domain exhibits appreciable affinity for a particular protein or antigen and, generally, does not exhibit significant reactivity with proteins or antigens other than BCMA or CD3. "Appreciable affinity" includes binding with an affinity of about 10^{-6} M (KD) or stronger. Preferably, binding is considered specific when binding affinity is about 10^{-12} to 10^{-8} M, 10^{-12} to 10^{-9} M, 10^{-12} to 10^{-10} M, 10^{-11} to 10^{-8} M, preferably of about 10^{-11} to 10^{-9} M. Whether a binding domain specifically reacts with or binds to a target can be tested readily by, *inter alia*, comparing the reaction of said binding domain with a target protein or antigen with the reaction of said binding domain with proteins or antigens other than BCMA or CD3. Preferably, a binding domain of the invention does not essentially bind or is not capable of binding to proteins or antigens other than BCMA or CD3 (i.e. the first binding domain is not capable of binding to

proteins other than BCMA and the second binding domain is not capable of binding to proteins other than CD3).

The term “does not essentially bind”, or “is not capable of binding” means that a binding domain of the present invention does not bind another protein or antigen other than BCMA or CD3, i.e.,
5 does not show reactivity of more than 30%, preferably not more than 20%, more preferably not more than 10%, particularly preferably not more than 9%, 8%, 7%, 6% or 5% with proteins or antigens other than BCMA or CD3, whereby binding to BCMA or CD3, respectively, is set to be 100%.

10 Specific binding is believed to be effected by specific motifs in the amino acid sequence of the binding domain and the antigen. Thus, binding is achieved as a result of their primary, secondary and/or tertiary structure as well as the result of secondary modifications of said structures. The specific interaction of the antigen-interaction-site with its specific antigen may result in a simple binding of said site to the antigen. Moreover, the specific interaction of the
15 antigen-interaction-site with its specific antigen may alternatively or additionally result in the initiation of a signal, e.g. due to the induction of a change of the conformation of the antigen, an oligomerization of the antigen, etc.

In one aspect, the first binding domain of the present invention binds to epitope cluster 3 of
20 human BCMA and is further capable of binding to epitope cluster 3 of macaque BCMA such as BCMA from *Macaca mulatta* (SEQ ID NO:1017) or *Macaca fascicularis* (SEQ ID NO:1017). It is envisaged that the first binding domain does or does not bind to murine BCMA.

Accordingly, in one embodiment, a binding domain which binds to human BCMA, in particular to
25 epitope cluster 3 of the extracellular protein domain of BCMA formed by amino acid residues 24 to 41 of the human sequence as depicted in SEQ ID NO: 1002, also binds to macaque BCMA, in particular to epitope cluster 3 of the extracellular protein domain of BCMA formed by amino acid residues 24 to 41 of the macaque BCMA sequence as depicted in SEQ ID NO: 1006.

30 In one embodiment, a first binding domain of a binding molecule is capable of binding to epitope cluster 3 of BCMA, wherein epitope cluster 3 of BCMA corresponds to amino acid residues 24 to 41 of the sequence as depicted in SEQ ID NO: 1002 (human BCMA full-length polypeptide) or SEQ ID NO: 1007 (human BCMA extracellular domain: amino acids 1-54 of SEQ ID NO: 1002).

35

In one aspect of the present invention, the first binding domain of the binding molecule is additionally or alternatively capable of binding to epitope cluster 3 of *Callithrix jacchus*, *Saguinus oedipus* and/or *Saimiri sciureus* BCMA.

5 Proteins (including fragments thereof, preferably biologically active fragments, and peptides, usually having less than 30 amino acids) comprise one or more amino acids coupled to each other via a covalent peptide bond (resulting in a chain of amino acids). The term "polypeptide" as used herein describes a group of molecules, which consist of more than 30 amino acids. Polypeptides may further form multimers such as dimers, trimers and higher oligomers, i.e.
10 consisting of more than one polypeptide molecule. Polypeptide molecules forming such dimers, trimers etc. may be identical or non-identical. The corresponding higher order structures of such multimers are, consequently, termed homo- or heterodimers, homo- or heterotrimers etc. An example for a heteromultimer is an antibody molecule, which, in its naturally occurring form, consists of two identical light polypeptide chains and two identical heavy polypeptide chains.
15 The terms "polypeptide" and "protein" also refer to naturally modified polypeptides/proteins wherein the modification is effected e.g. by post-translational modifications like glycosylation, acetylation, phosphorylation and the like. A "polypeptide" when referred to herein may also be chemically modified such as pegylated. Such modifications are well known in the art.

In another aspect of the invention, the second binding domain is capable of binding to CD3
20 epsilon. In still another aspect of the invention, the second binding domain is capable of binding to human CD3 and to macaque CD3, preferably to human CD3 epsilon and to macaque CD3 epsilon. Additionally or alternatively, the second binding domain is capable of binding to *Callithrix jacchus*, *Saguinus oedipus* and/or *Saimiri sciureus* CD3 epsilon. According to these embodiments, one or both binding domains of the binding molecule of the invention are
25 preferably cross-species specific for members of the mammalian order of primates. Cross-species specific CD3 binding domains are, for example, described in WO 2008/119567.

It is particularly preferred for the binding molecule of the present invention that the second binding domain capable of binding to the T cell CD3 receptor complex comprises a VL region
30 comprising CDR-L1, CDR-L2 and CDR-L3 selected from:

(a) CDR-L1 as depicted in SEQ ID NO: 27 of WO 2008/119567, CDR-L2 as depicted in SEQ ID NO: 28 of WO 2008/119567 and CDR-L3 as depicted in SEQ ID NO: 29 of WO 2008/119567;

(b) CDR-L1 as depicted in SEQ ID NO: 117 of WO 2008/119567, CDR-L2 as depicted in SEQ ID NO: 118 of WO 2008/119567 and CDR-L3 as depicted in SEQ ID NO: 119 of WO 2008/119567; and

(c) CDR-L1 as depicted in SEQ ID NO: 153 of WO 2008/119567, CDR-L2 as depicted in SEQ ID NO: 154 of WO 2008/119567 and CDR-L3 as depicted in SEQ ID NO: 155 of WO 2008/119567.

In an alternatively preferred embodiment of the binding molecule of the present invention, the second binding domain capable of binding to the T cell CD3 receptor complex comprises a VH region comprising CDR-H 1, CDR-H2 and CDR-H3 selected from:

(a) CDR-H1 as depicted in SEQ ID NO: 12 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 13 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 14 of WO 2008/119567;

(b) CDR-H1 as depicted in SEQ ID NO: 30 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 31 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 32 of WO 2008/119567;

(c) CDR-H1 as depicted in SEQ ID NO: 48 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 49 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 50 of WO 2008/119567;

(d) CDR-H1 as depicted in SEQ ID NO: 66 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 67 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 68 of WO 2008/119567;

(e) CDR-H1 as depicted in SEQ ID NO: 84 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 85 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 86 of WO 2008/119567;

(f) CDR-H1 as depicted in SEQ ID NO: 102 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 103 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 104 of WO 2008/119567;

(g) CDR-H1 as depicted in SEQ ID NO: 120 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 121 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 122 of WO 2008/119567;

(h) CDR-H1 as depicted in SEQ ID NO: 138 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 139 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 140 of WO 2008/119567;

(i) CDR-H1 as depicted in SEQ ID NO: 156 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 157 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 158 of WO 2008/119567; and

(j) CDR-H1 as depicted in SEQ ID NO: 174 of WO 2008/119567, CDR-H2 as depicted in SEQ ID NO: 175 of WO 2008/119567 and CDR-H3 as depicted in SEQ ID NO: 176 of WO 2008/119567.

It is further preferred for the binding molecule of the present invention that the second binding domain capable of binding to the T cell CD3 receptor complex comprises a VL region selected from the group consisting of a VL region as depicted in SEQ ID NO: 35, 39, 125, 129, 161 or 165 of WO 2008/119567.

It is alternatively preferred that the second binding domain capable of binding to the T cell CD3 receptor complex comprises a VH region selected from the group consisting of a VH region as depicted in SEQ ID NO: 15, 19, 33, 37, 51, 55, 69, 73, 87, 91, 105, 109, 123, 127, 141, 145, 159, 163, 177 or 181 of WO 2008/119567.

More preferably, the binding molecule of the present invention is characterized by the second binding domain capable of binding to the T cell CD3 receptor complex comprising a VL region and a VH region selected from the group consisting of:

(a) a VL region as depicted in SEQ ID NO: 17 or 21 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 15 or 19 of WO 2008/119567;

(b) a VL region as depicted in SEQ ID NO: 35 or 39 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 33 or 37 of WO 2008/119567;

(c) a VL region as depicted in SEQ ID NO: 53 or 57 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 51 or 55 of WO 2008/119567;

(d) a VL region as depicted in SEQ ID NO: 71 or 75 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 69 or 73 of WO 2008/119567;

(e) a VL region as depicted in SEQ ID NO: 89 or 93 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 87 or 91 of WO 2008/119567;

(f) a VL region as depicted in SEQ ID NO: 107 or 111 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 105 or 109 of WO 2008/119567;

(g) a VL region as depicted in SEQ ID NO: 125 or 129 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 123 or 127 of WO 2008/119567;

(h) a VL region as depicted in SEQ ID NO: 143 or 147 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 141 or 145 of WO 2008/119567;

(i) a VL region as depicted in SEQ ID NO: 161 or 165 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 159 or 163 of WO 2008/119567; and

5 (j) a VL region as depicted in SEQ ID NO: 179 or 183 of WO 2008/119567 and a VH region as depicted in SEQ ID NO: 177 or 181 of WO 2008/119567.

According to a preferred embodiment of the binding molecule of the present invention, in particular the second binding domain capable of binding to the T cell CD3 receptor complex, the
10 pairs of VH-regions and VL-regions are in the format of a single chain antibody (scFv). The VH and VL regions are arranged in the order VH-VL or VL-VH. It is preferred that the VH-region is positioned N-terminally to a linker sequence. The VL-region is positioned C-terminally of the linker sequence.

15 A preferred embodiment of the above described binding molecule of the present invention is characterized by the second binding domain capable of binding to the T cell CD3 receptor complex comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 23, 25, 41, 43, 59, 61, 77, 79, 95, 97, 113, 115, 131, 133, 149, 151, 167, 169, 185 or 187 of WO 2008/119567.

20 The affinity of the first binding domain for human BCMA is preferably ≤ 15 nM, more preferably ≤ 10 nM, even more preferably ≤ 5 nM, even more preferably ≤ 1 nM, even more preferably ≤ 0.5 nM, even more preferably ≤ 0.1 nM, and most preferably ≤ 0.05 nM. The affinity of the first binding domain for macaque BCMA is preferably ≤ 15 nM, more preferably ≤ 10 nM, even more
25 preferably ≤ 5 nM, even more preferably ≤ 1 nM, even more preferably ≤ 0.5 nM, even more preferably ≤ 0.1 nM, and most preferably ≤ 0.05 nM or even ≤ 0.01 nM. The affinity can be measured for example in a Biacore assay or in a Scatchard assay, e.g. as described in the Examples. The affinity gap for binding to macaque BCMA versus human BCMA is preferably [1:10-1:5] or [5:1-10:1], more preferably [1:5-5:1], and most preferably [1:2-3:1] or even [1:1-
30 3:1]. Other methods of determining the affinity are well-known to the skilled person.

Cytotoxicity mediated by BCMA/CD3 bispecific binding molecules can be measured in various ways. Effector cells can be e.g. stimulated enriched (human) CD8 positive T cells or unstimulated (human) peripheral blood mononuclear cells (PBMC). If the target cells are of
35 macaque origin or express or are transfected with macaque BCMA, the effector cells should

also be of macaque origin such as a macaque T cell line, e.g. 4119LnPx. The target cells should express (at least the extracellular domain of) BCMA, e.g. human or macaque BCMA. Target cells can be a cell line (such as CHO) which is stably or transiently transfected with BCMA, e.g. human or macaque BCMA. Alternatively, the target cells can be a BCMA positive
5 natural expresser cell line, such as the human multiple myeloma cell line L363 or NCI-H929. Usually EC₅₀-values are expected to be lower with target cell lines expressing higher levels of BCMA on the cell surface. The effector to target cell (E:T) ratio is usually about 10:1, but can also vary. Cytotoxic activity of BCMA/CD3 bispecific binding molecules can be measured in an
10 51-chromium release assay (incubation time of about 18 hours) or in a in a FACS-based cytotoxicity assay (incubation time of about 48 hours). Modifications of the assay incubation time (cytotoxic reaction) are also possible. Other methods of measuring cytotoxicity are well-known to the skilled person and comprise MTT or MTS assays, ATP-based assays including bioluminescent assays, the sulforhodamine B (SRB) assay, WST assay, clonogenic assay and the ECIS technology.

15 The cytotoxic activity mediated by BCMA/CD3 bispecific binding molecules of the present invention is preferably measured in a cell-based cytotoxicity assay. It is represented by the EC₅₀ value, which corresponds to the half maximal effective concentration (concentration of the binding molecule which induces a cytotoxic response halfway between the baseline and
20 maximum). Preferably, the EC₅₀ value of the BCMA/CD3 bispecific binding molecules is ≤20.000 pg/ml, more preferably ≤5000 pg/ml, even more preferably ≤1000 pg/ml, even more preferably ≤500 pg/ml, even more preferably ≤350 pg/ml, even more preferably ≤320 pg/ml, even more preferably ≤250 pg/ml, even more preferably ≤100 pg/ml, even more preferably ≤50 pg/ml, even more preferably ≤10 pg/ml, and most preferably ≤5 pg/ml.

25 Any of the above given EC₅₀ values can be combined with any one of the indicated scenarios of a cell-based cytotoxicity assay. For example, when (human) CD8 positive T cells or a macaque T cell line are used as effector cells, the EC₅₀ value of the BCMA/CD3 bispecific binding molecule is preferably ≤1000 pg/ml, more preferably ≤500 pg/ml, even more preferably
30 ≤250 pg/ml, even more preferably ≤100 pg/ml, even more preferably ≤50 pg/ml, even more preferably ≤10 pg/ml, and most preferably ≤5 pg/ml. If in this assay the target cells are (human or macaque) BCMA transfected cells such as CHO cells, the EC₅₀ value of the BCMA/CD3 bispecific binding molecule is preferably ≤150 pg/ml, more preferably ≤100 pg/ml, even more preferably ≤50 pg/ml, even more preferably ≤30 pg/ml, even more preferably ≤10 pg/ml, and
35 most preferably ≤5 pg/ml.

If the target cells are a BCMA positive natural expresser cell line, then the EC₅₀ value is preferably ≤ 350 pg/ml, more preferably ≤ 320 pg/ml, even more preferably ≤ 250 pg/ml, even more preferably ≤ 200 pg/ml, even more preferably ≤ 100 pg/ml, even more preferably ≤ 150 pg/ml, even more preferably ≤ 100 pg/ml, and most preferably ≤ 50 pg/ml, or lower.

- 5 When (human) PBMCs are used as effector cells, the EC₅₀ value of the BCMA/CD3 bispecific binding molecule is preferably ≤ 1000 pg/ml, more preferably ≤ 750 pg/ml, more preferably ≤ 500 pg/ml, even more preferably ≤ 350 pg/ml, even more preferably ≤ 320 pg/ml, even more preferably ≤ 250 pg/ml, even more preferably ≤ 100 pg/ml, and most preferably ≤ 50 pg/ml, or lower.
- 10 In a particularly preferred embodiment, the BCMA/CD3 bispecific binding molecules of the present invention are characterized by an EC₅₀ of ≤ 350 pg/ml or less, more preferably ≤ 320 pg/ml or less. In that embodiment the target cells are L363 cells and the effector cells are unstimulated human PBMCs. The skilled person knows how to measure the EC₅₀ value without further ado. Moreover, the specification teaches a specific instruction how to measure the EC₅₀
- 15 value; see, for example, Example 8.3, below. A suitable protocol is as follows:
 - a) Prepare human peripheral blood mononuclear cells (PBMC) by Ficoll density gradient centrifugation from enriched lymphocyte preparations (buffy coats)
 - b) Optionally wash with Dulbecco's PBS (Gibco)
 - c) Remove remaining erythrocytes from PBMC via incubation with erythrocyte lysis buffer
 - 20 (155 mM NH₄Cl, 10 mM KHCO₃, 100 μ M EDTA)
 - c) Remove platelets via the supernatant upon centrifugation of PBMC at 100 x g
 - d) Deplete CD14⁺ cells and NK cells
 - e) Isolate CD14/CD56 negative cells using, e.g. LS Columns (Miltenyi Biotec, #130-042-401)
 - 25 f) Culture PBMC w/o CD14⁺/CD56⁺ cells, e.g. in RPMI complete medium i.e. RPMI1640 (Biochrom AG, #FG1215) supplemented with 10% FBS (Biochrom AG, #S0115), 1x non-essential amino acids (Biochrom AG, #K0293), 10 mM Hepes buffer (Biochrom AG, #L1613), 1 mM sodium pyruvate (Biochrom AG, #L0473) and 100 U/mL penicillin/streptomycin (Biochrom AG, #A2213) at 37°C in an incubator until needed.
 - 30 g) Label target cells
 - h) Mix effector and target cells, preferably at equal volumes, so as to have an E:T cell ratio of 10:1
 - i) Add the binding molecule, preferably in a serial dilution
 - j) Proceed for 48 hours in a 7% CO₂ humidified incubator

- k) Monitor target cell membrane integrity, e.g., by adding propidium iodide (PI) at a final concentration of 1 µg/mL, for example, by flow cytometry
- l) Calculate EC50, e.g., according to the following formula:

$$\text{Cytotoxicity [\%]} = \frac{n_{\text{dead target cells}}}{n_{\text{target cells}}} \times 100$$

5 n = number of events

Using GraphPad Prism 5 software (Graph Pad Software, San Diego), the percentage of cytotoxicity was plotted against the corresponding bispecific antibody concentrations. Dose response curves can be analyzed with the four parametric logistic regression models for evaluation of sigmoid dose response curves with fixed hill slope and EC50 values were
10 calculated.

In view of the above, it is preferred that the binding molecule of the present invention is characterized by an EC50 (pg/ml) of 350 or less, preferably 320 or less.

15 The present invention also relates to binding molecules described herein which are characterized by an EC50 (pg/ml) which equates to the EC50 (pg/ml) of any one of the BCMA/CD3 bispecific binding molecules BCMA-83 x CD3, BCMA-62 x CD3, BCMA-5 x CD3, BCMA-98 x CD3, BCMA-71 x CD3, BCMA-34 x CD3, BCMA-74 x CD3, BCMA-20 x CD3. In order to determine as to whether the EC50 of a binding molecule as described herein equates
20 to the EC50 of any one of BCMA-83 x CD3, BCMA-62 x CD3, BCMA-5 x CD3, BCMA-98 x CD3, BCMA-71 x CD3, BCMA-34 x CD3, BCMA-74 x CD3, BCMA-20 x CD3, it is envisaged that for the determination of the EC50 value the same assay is applied. The term “equates to” includes thereby a deviation of +/- 10%, preferably +/- 7.5%, more preferably +/- 5%, even more preferably +/- 2.5% of the respective EC50 value.

25 The BCMA/CD3 bispecific binding molecules BCMA-83 x CD3, BCMA-62 x CD3, BCMA-5 x CD3, BCMA-98 x CD3, BCMA-71 x CD3, BCMA-34 x CD3, BCMA-74 x CD3, BCMA-20 x CD3 that serve as “reference” binding molecules in the above described assay are preferably produced in CHO cells.

30 The difference in cytotoxic activity between the monomeric and the dimeric isoform of individual BCMA/CD3 bispecific binding molecules (such as antibodies) is referred to as “potency gap”. This potency gap can e.g. be calculated as ratio between EC₅₀ values of the molecule’s monomeric and dimeric form. Potency gaps of the BCMA/CD3 bispecific binding molecules of

the present invention are preferably ≤ 5 , more preferably ≤ 4 , even more preferably ≤ 3 , even more preferably ≤ 2 and most preferably ≤ 1 .

Preferably, the BCMA/CD3 bispecific binding molecules of the present invention do not bind to, interact with, recognize or cross-react with human BAFF-R and/or human TACI. Methods to
5 detect cross-reactivity with human BAFF-R and/or human TACI are disclosed in Example 9.

It is also preferred that the BCMA/CD3 bispecific binding molecules of the present invention present with very low dimer conversion after a number of freeze/thaw cycles. Preferably the
10 dimer percentages are $\leq 5\%$, more preferably $\leq 4\%$, even more preferably $\leq 3\%$, even more preferably $\leq 2.5\%$, even more preferably $\leq 2\%$, even more preferably $\leq 1.5\%$, and most preferably $\leq 1\%$, for example after three freeze/thaw cycles. A freeze-thaw cycle and the determination of the dimer percentage can be carried out in accordance with Example 16.

15 The BCMA/CD3 bispecific binding molecules (such as antibodies) of the present invention preferably show a favorable thermostability with melting temperatures above 60°C .

To determine potential interaction of BCMA/CD3 bispecific binding molecules (such as antibodies) with human plasma proteins, a plasma interference test can be carried out (see e.g.
20 Example 18). In a preferred embodiment, there is no significant reduction of target binding of the BCMA/CD3 bispecific binding molecules mediated by plasma proteins. The relative plasma interference value is preferably ≤ 2 .

It is furthermore envisaged that the BCMA/CD3 bispecific binding molecules of the present
25 invention are capable of exhibiting therapeutic efficacy or anti-tumor activity. This can be assessed e.g. in a study as disclosed in Example 19 (advanced stage human tumor xenograft model). The skilled person knows how to modify or adapt certain parameters of this study, such as the number of injected tumor cells, the site of injection, the number of transplanted human T cells, the amount of BCMA/CD3 bispecific binding molecules to be administered, and the
30 timelines, while still arriving at a meaningful and reproducible result. Preferably, the tumor growth inhibition T/C [%] is 70 or 60 or lower, more preferably 50 or 40 or lower, even more preferably at least 30 or 20 or lower and most preferably 10 or lower, 5 or lower or even 2.5 or lower.

Preferably, the BCMA/CD3 bispecific binding molecules of the present invention do not induce / mediate lysis or do not essentially induce / mediate lysis of BCMA negative cells such as HL60, MES-SA, and SNU-16. The term “do not induce lysis”, “do not essentially induce lysis”, “do not mediate lysis” or “do not essentially mediate lysis” means that a binding molecule of the present invention does not induce or mediate lysis of more than 30%, preferably not more than 20%, more preferably not more than 10%, particularly preferably not more than 9%, 8%, 7%, 6% or 5% of BCMA negative cells, whereby lysis of a BCMA positive cell line such as NCI-H929, L-363 or OPM-2 is set to be 100%. This applies for concentrations of the binding molecule of at least up to 500 nM. The skilled person knows how to measure cell lysis without further ado. Moreover, the specification teaches a specific instruction how to measure cell lysis; see e.g. Example 20 below.

In one embodiment, the first or the second binding domain is or is derived from an antibody. In another embodiment, both binding domains are or are derived from an antibody.

The definition of the term “antibody” includes embodiments such as monoclonal, chimeric, single chain, humanized and human antibodies. In addition to full-length antibodies, the definition also includes antibody derivatives and antibody fragments, like, *inter alia*, Fab fragments. Antibody fragments or derivatives further comprise F(ab')₂, Fv, scFv fragments or single domain antibodies such as domain antibodies or nanobodies, single variable domain antibodies or immunoglobulin single variable domain comprising merely one variable domain, which might be VHH, VH or VL, that specifically bind an antigen or epitope independently of other V regions or domains; see, for example, Harlow and Lane (1988) and (1999), loc. cit.; Kontermann and Dübel, Antibody Engineering, Springer, 2nd ed. 2010 and Little, Recombinant Antibodies for Immunotherapy, Cambridge University Press 2009. Said term also includes diabodies or Dual-Affinity Re-Targeting (DART) antibodies. Further envisaged are (bispecific) single chain diabodies, tandem diabodies (Tandab's), „minibodies“ exemplified by a structure which is as follows: (VH-VL-CH3)₂, (scFv-CH3)₂ or (scFv-CH3-scFv)₂, „Fc DART“ antibodies and „IgG DART“ antibodies, and multibodies such as triabodies. Immunoglobulin single variable domains encompass not only an isolated antibody single variable domain polypeptide, but also larger polypeptides that comprise one or more monomers of an antibody single variable domain polypeptide sequence.

Various procedures are known in the art and may be used for the production of such antibodies and/or fragments. Thus, (antibody) derivatives can be produced by peptidomimetics. Further,

techniques described for the production of single chain antibodies (see, *inter alia*, US Patent 4,946,778, Kontermann and Dübel (2010), loc. cit. and Little (2009), loc. cit.) can be adapted to produce single chain antibodies specific for elected polypeptide(s). Also, transgenic animals may be used to express humanized antibodies specific for polypeptides and fusion proteins of this invention. For the preparation of monoclonal antibodies, any technique, providing antibodies produced by continuous cell line cultures can be used. Examples for such techniques include the hybridoma technique (Köhler and Milstein Nature 256 (1975), 495-497), the trioma technique, the human B cell hybridoma technique (Kozbor, Immunology Today 4 (1983), 72) and the EBV hybridoma technique to produce human monoclonal antibodies (Cole *et al.*, Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc. (1985), 77-96). Surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of a target polypeptide, such as CD3 epsilon (Schier, Human Antibodies Hybridomas 7 (1996), 97-105; Malmberg, J. Immunol. Methods 183 (1995), 7-13). It is also envisaged in the context of this invention that the term "antibody" comprises antibody constructs, which may be expressed in a host as described herein below, e.g. antibody constructs which may be transfected and/or transduced via, *inter alia*, viruses or plasmid vectors.

Furthermore, the term "antibody" as employed herein also relates to derivatives or variants of the antibodies described herein which display the same specificity as the described antibodies. Examples of "antibody variants" include humanized variants of non-human antibodies, "affinity matured" antibodies (see, e.g. Hawkins *et al.* J. Mol. Biol. 254, 889-896 (1992) and Lowman *et al.*, Biochemistry 30, 10832- 10837 (1991)) and antibody mutants with altered effector function(s) (see, e.g., US Patent 5, 648, 260, Kontermann and Dübel (2010), loc. cit. and Little(2009), loc. cit.).

The terms "antigen-binding domain", "antigen-binding fragment" and "antibody binding region" when used herein refer to a part of an antibody molecule that comprises amino acids responsible for the specific binding between antibody and antigen. The part of the antigen that is specifically recognized and bound by the antibody is referred to as the "epitope" as described herein above. As mentioned above, an antigen-binding domain may typically comprise an antibody light chain variable region (VL) and an antibody heavy chain variable region (VH); however, it does not have to comprise both. Fd fragments, for example, have two VH regions and often retain some antigen-binding function of the intact antigen-binding domain. Examples of antigen-binding fragments of an antibody include (1) a Fab fragment, a monovalent fragment

having the VL, VH, CL and CH1 domains; (2) a F(ab')₂ fragment, a bivalent fragment having two Fab fragments linked by a disulfide bridge at the hinge region; (3) an Fd fragment having the two VH and CH1 domains; (4) an Fv fragment having the VL and VH domains of a single arm of an antibody, (5) a dAb fragment (Ward *et al.*, (1989) Nature 341 :544-546), which has a VH domain; (6) an isolated complementarity determining region (CDR), and (7) a single chain Fv (scFv), the latter being preferred (for example, derived from a scFV-library). Although the two domains of the Fv fragment, VL and VH are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see e.g., Huston *et al.* (1988) Proc. Natl. Acad. Sci USA 85:5879-5883). These antibody fragments are obtained using conventional techniques known to those with skill in the art, and the fragments are evaluated for function in the same manner as are intact antibodies.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translation modifications (e.g., isomerizations, amidations) that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, Nature, 256: 495 (1975), or may be made by recombinant DNA methods (see, e.g., U. S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, Nature, 352: 624-628 (1991) and Marks *et al.*, J. Mol. Biol., 222: 581-597 (1991), for example.

The monoclonal antibodies of the present invention specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or

homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain (s) is (are) identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U. S. Patent No. 4,816, 567; Morrison et al., Proc. Natl. Acad. Sci. USA, 81: 6851-6855 (1984)). Chimeric antibodies of interest herein include "primitized" antibodies comprising variable domain antigen-binding sequences derived from a non-human primate (e.g., Old World Monkey, Ape etc.) and human constant region sequences.

"Humanized" forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) of mostly human sequences, which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region (also CDR) of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, "humanized antibodies" as used herein may also comprise residues which are found neither in the recipient antibody nor the donor antibody. These modifications are made to further refine and optimize antibody performance. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, Nature, 321: 522-525 (1986); Reichmann *et al.*, Nature, 332: 323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2: 593-596 (1992).

The term "human antibody" includes antibodies having variable and constant regions corresponding substantially to human germline immunoglobulin sequences known in the art, including, for example, those described by Kabat *et al.* (See Kabat *et al.* (1991) loc. cit.). The human antibodies of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs, and in particular, CDR3. The human antibody can have at least one, two, three, four, five, or more positions replaced with an amino acid residue that is not encoded by the human germline immunoglobulin sequence.

As used herein, "*in vitro* generated antibody" refers to an antibody where all or part of the variable region (e.g., at least one CDR) is generated in a non-immune cell selection (e.g., an *in vitro* phage display, protein chip or any other method in which candidate sequences can be tested for their ability to bind to an antigen). This term thus preferably excludes sequences generated by genomic rearrangement in an immune cell.

A "bispecific" or "bifunctional" antibody or immunoglobulin is an artificial hybrid antibody or immunoglobulin having two different heavy/light chain pairs and two different binding sites. Bispecific antibodies can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai & Lachmann, Clin. Exp. Immunol. 79:315-321 (1990). Numerous methods known to those skilled in the art are available for obtaining antibodies or antigen-binding fragments thereof. For example, antibodies can be produced using recombinant DNA methods (U.S. Patent No. 4,816,567). Monoclonal antibodies may also be produced by generation of hybridomas (see e.g., Kohler and Milstein (1975) Nature, 256: 495-499) in accordance with known methods. Hybridomas formed in this manner are then screened using standard methods, such as enzyme-linked immunosorbent assay (ELISA) and surface plasmon resonance (BIAcore™) analysis, to identify one or more hybridomas that produce an antibody that specifically binds with a specified antigen. Any form of the specified antigen may be used as the immunogen, e.g., recombinant antigen, naturally occurring forms, any variants or fragments thereof, as well as antigenic peptide thereof."

One exemplary method of making antibodies includes screening protein expression libraries, e.g., phage or ribosome display libraries. Phage display is described, for example, in Ladner *et al.*, U.S. Patent No. 5,223,409; Smith (1985) Science 228:1315-1317; Clackson *et al.* (1991) Nature, 352: 624-628.

In addition to the use of display libraries, the specified antigen can be used to immunize a non-human animal, e.g., a rodent, e.g., a mouse, hamster, or rat. In one embodiment, the non-human animal includes at least a part of a human immunoglobulin gene. For example, it is possible to engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci. Using the hybridoma technology, antigen-specific monoclonal antibodies derived from the genes with the desired specificity may be produced and selected. See, e.g., XENOMOUSE™, Green *et al.* (1994) Nature Genetics 7:13-21, US 2003- 0070185, WO 96/34096, and WO96/33735.

A monoclonal antibody can be obtained from a non-human animal, and then modified, e.g., humanized, deimmunized, chimeric, may be produced using recombinant DNA techniques known in the art. A variety of approaches for making chimeric antibodies have been described. See e.g., Morrison *et al.*, Proc. Natl. Acad. Sci. U.S.A. 81:6851, 1985; Takeda *et al.*, Nature 314:452, 1985, Cabilly *et al.*, U.S. Patent No. 4,816,567; Boss *et al.*, U.S. Patent No. 4,816,397; Tanaguchi *et al.*, EP 0171496; EP 0173494, GB 2177096. Humanized antibodies may also be produced, for example, using transgenic mice that express human heavy and light chain genes, but are incapable of expressing the endogenous mouse immunoglobulin heavy and light chain genes. Winter describes an exemplary CDR-grafting method that may be used to prepare the humanized antibodies described herein (U.S. Patent No. 5,225,539). All of the CDRs of a particular human antibody may be replaced with at least a portion of a non-human CDR, or only some of the CDRs may be replaced with non-human CDRs. It is only necessary to replace the number of CDRs required for binding of the humanized antibody to a predetermined antigen.

Humanized antibodies or fragments thereof can be generated by replacing sequences of the Fv variable domain that are not directly involved in antigen binding with equivalent sequences from human Fv variable domains. Exemplary methods for generating humanized antibodies or fragments thereof are provided by Morrison (1985) Science 229:1202-1207; by Oi *et al.* (1986) BioTechniques 4:214; and by US 5,585,089; US 5,693,761; US 5,693,762; US 5,859,205; and US 6,407,213. Those methods include isolating, manipulating, and expressing the nucleic acid sequences that encode all or part of immunoglobulin Fv variable domains from at least one of a heavy or light chain. Such nucleic acids may be obtained from a hybridoma producing an antibody against a predetermined target, as described above, as well as from other sources. The recombinant DNA encoding the humanized antibody molecule can then be cloned into an appropriate expression vector.

A humanized antibody can be optimized by the introduction of conservative substitutions, consensus sequence substitutions, germline substitutions and/or back mutations. Such altered immunoglobulin molecules can be made by any of several techniques known in the art, (e.g., Teng *et al.*, Proc. Natl. Acad. Sci. U.S.A., 80: 7308-7312, 1983; Kozbor *et al.*, Immunology Today, 4: 7279, 1983; Olsson *et al.*, Meth. Enzymol., 92: 3-16, 1982), and may be made according to the teachings of EP 239 400.

An antibody or fragment thereof may also be modified by specific deletion of human T cell epitopes or "deimmunization" by the methods disclosed in WO 98/52976 and WO 00/34317.

Briefly, the heavy and light chain variable domains of an antibody can be analyzed for peptides

that bind to MHC class II; these peptides represent potential T cell epitopes (as defined in WO 98/52976 and WO 00/34317). For detection of potential T cell epitopes, a computer modeling approach termed "peptide threading" can be applied, and in addition a database of human MHC class II binding peptides can be searched for motifs present in the VH and VL sequences, as described in WO 98/52976 and WO 00/34317. These motifs bind to any of the 18 major MHC class II DR allotypes, and thus constitute potential T cell epitopes. Potential T-cell epitopes detected can be eliminated by substituting small numbers of amino acid residues in the variable domains, or preferably, by single amino acid substitutions. Typically, conservative substitutions are made. Often, but not exclusively, an amino acid common to a position in human germline antibody sequences may be used. Human germline sequences, e.g., are disclosed in Tomlinson, *et al.* (1992) J. Mol. Biol. 227:776-798; Cook, G.P. *et al.* (1995) Immunol. Today Vol. 16 (5): 237-242; and Tomlinson *et al.* (1995) EMBO J. 14: 14:4628-4638. The V BASE directory provides a comprehensive directory of human immunoglobulin variable region sequences (compiled by Tomlinson, LA. *et al.* MRC Centre for Protein Engineering, Cambridge, UK). These sequences can be used as a source of human sequence, e.g., for framework regions and CDRs. Consensus human framework regions can also be used, e.g., as described in US Patent No. 6,300,064.

The pairing of a VH and VL together forms a single antigen-binding site. The CH domain most proximal to VH is designated as CH1. Each L chain is linked to an H chain by one covalent disulfide bond, while the two H chains are linked to each other by one or more disulfide bonds depending on the H chain isotype. The VH and VL domains consist of four regions of relatively conserved sequences called framework regions (FR1, FR2, FR3, and FR4), which form a scaffold for three regions of hypervariable sequences (complementarity determining regions, CDRs). The CDRs contain most of the residues responsible for specific interactions of the antibody with the antigen. CDRs are referred to as CDR 1, CDR2, and CDR3. Accordingly, CDR constituents on the heavy chain are referred to as H1, H2, and H3, while CDR constituents on the light chain are referred to as L1, L2, and L3.

The term "variable" refers to the portions of the immunoglobulin domains that exhibit variability in their sequence and that are involved in determining the specificity and binding affinity of a particular antibody (i.e., the "variable domain(s)"). Variability is not evenly distributed throughout the variable domains of antibodies; it is concentrated in sub-domains of each of the heavy and light chain variable regions. These sub-domains are called "hypervariable" regions or "complementarity determining regions" (CDRs). The more conserved (i.e., non-hypervariable) portions of the variable domains are called the "framework" regions (FRM). The variable

domains of naturally occurring heavy and light chains each comprise four FRM regions, largely adopting a β -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the β -sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRM and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site (see Kabat *et al.*, loc. cit.). The constant domains are not directly involved in antigen binding, but exhibit various effector functions, such as, for example, antibody-dependent, cell-mediated cytotoxicity and complement activation.

It is also preferred for the binding molecule of the invention that first and the second domain form a molecule that is selected from the group of (scFv)₂, (single domain mAb)₂, scFv-single domain mAb, diabody or oligomeres thereof.

The terms "CDR", and its plural "CDRs", refer to a complementarity determining region (CDR) of which three make up the binding character of a light chain variable region (CDRL1, CDRL2 and CDRL3) and three make up the binding character of a heavy chain variable region (CDRH1, CDRH2 and CDRH3). CDRs contribute to the functional activity of an antibody molecule and are separated by amino acid sequences that comprise scaffolding or framework regions. The exact definitional CDR boundaries and lengths are subject to different classification and numbering systems. CDRs may therefore be referred to by Kabat, Chothia, contact or any other boundary definitions, including the numbering system described herein. Despite differing boundaries, each of these systems has some degree of overlap in what constitutes the so called "hypervariable regions" within the variable sequences. CDR definitions according to these systems may therefore differ in length and boundary areas with respect to the adjacent framework region. See for example Kabat, Chothia, and/or MacCallum (Kabat *et al.*, loc. cit.; Chothia *et al.*, J. Mol. Biol, 1987, 196: 901; and MacCallum *et al.*, J. Mol. Biol, 1996, 262: 732). However, the numbering in accordance with the so-called Kabat system is preferred.

The term "amino acid" or "amino acid residue" typically refers to an amino acid having its art recognized definition such as an amino acid selected from the group consisting of: alanine (Ala or A); arginine (Arg or R); asparagine (Asn or N); aspartic acid (Asp or D); cysteine (Cys or C); glutamine (Gln or Q); glutamic acid (Glu or E); glycine (Gly or G); histidine (His or H); isoleucine (Ile or I); leucine (Leu or L); lysine (Lys or K); methionine (Met or M); phenylalanine (Phe or F); proline (Pro or P); serine (Ser or S); threonine (Thr or T); tryptophan (Trp or W); tyrosine (Tyr or Y); and valine (Val or V), although modified, synthetic, or rare amino acids may be used as

desired. Generally, amino acids can be grouped as having a nonpolar side chain (e.g., Ala, Cys, He, Leu, Met, Phe, Pro, Val); a negatively charged side chain (e.g., Asp, Glu); a positively charged sidechain (e.g., Arg, His, Lys); or an uncharged polar side chain (e.g., Asn, Cys, Gln, Gly, His, Met, Phe, Ser, Thr, Trp, and Tyr).

5

The term "hypervariable region" (also known as "complementarity determining regions" or CDRs) when used herein refers to the amino acid residues of an antibody which are (usually three or four short regions of extreme sequence variability) within the V-region domain of an immunoglobulin which form the antigen-binding site and are the main determinants of antigen specificity. There are at least two methods for identifying the CDR residues: (1) An approach based on cross-species sequence variability (i. e., Kabat *et al.*, loc. cit.); and (2) An approach based on crystallographic studies of antigen-antibody complexes (Chothia, C. *et al.*, J. Mol. Biol. 196: 901-917 (1987)). However, to the extent that two residue identification techniques define regions of overlapping, but not identical regions, they can be combined to define a hybrid CDR. However, in general, the CDR residues are preferably identified in accordance with the so-called Kabat (numbering) system.

The term "framework region" refers to the art-recognized portions of an antibody variable region that exist between the more divergent (i.e., hypervariable) CDRs. Such framework regions are typically referred to as frameworks 1 through 4 (FR1, FR2, FR3, and FR4) and provide a scaffold for the presentation of the six CDRs (three from the heavy chain and three from the light chain) in three dimensional space, to form an antigen-binding surface.

Typically, CDRs form a loop structure that can be classified as a canonical structure. The term "canonical structure" refers to the main chain conformation that is adopted by the antigen binding (CDR) loops. From comparative structural studies, it has been found that five of the six antigen binding loops have only a limited repertoire of available conformations. Each canonical structure can be characterized by the torsion angles of the polypeptide backbone. Correspondent loops between antibodies may, therefore, have very similar three dimensional structures, despite high amino acid sequence variability in most parts of the loops (Chothia and Lesk, J. Mol. Biol., 1987, 196: 901; Chothia *et al.*, Nature, 1989, 342: 877; Martin and Thornton, J. Mol. Biol, 1996, 263: 800, each of which is incorporated by reference in its entirety). Furthermore, there is a relationship between the adopted loop structure and the amino acid sequences surrounding it. The conformation of a particular canonical class is determined by the length of the loop and the amino acid residues residing at key positions within the loop, as well

as within the conserved framework (i.e., outside of the loop). Assignment to a particular canonical class can therefore be made based on the presence of these key amino acid residues. The term "canonical structure" may also include considerations as to the linear sequence of the antibody, for example, as catalogued by Kabat (Kabat *et al.*, loc. cit.). The
5 Kabat numbering scheme (system) is a widely adopted standard for numbering the amino acid residues of an antibody variable domain in a consistent manner and is the preferred scheme applied in the present invention as also mentioned elsewhere herein. Additional structural considerations can also be used to determine the canonical structure of an antibody. For example, those differences not fully reflected by Kabat numbering can be described by the
10 numbering system of Chothia *et al* and/or revealed by other techniques, for example, crystallography and two or three-dimensional computational modeling. Accordingly, a given antibody sequence may be placed into a canonical class which allows for, among other things, identifying appropriate chassis sequences (e.g., based on a desire to include a variety of canonical structures in a library). Kabat numbering of antibody amino acid sequences and
15 structural considerations as described by Chothia *et al.*, loc. cit. and their implications for construing canonical aspects of antibody structure, are described in the literature.

CDR3 is typically the greatest source of molecular diversity within the antibody-binding site. H3, for example, can be as short as two amino acid residues or greater than 26 amino acids. The
20 subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known in the art. For a review of the antibody structure, see Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, eds. Harlow *et al.*, 1988. One of skill in the art will recognize that each subunit structure, e.g., a CH, VH, CL, VL, CDR, FR structure, comprises active fragments, e.g., the portion of the VH, VL, or CDR subunit the binds to the antigen, i.e.,
25 the antigen-binding fragment, or, e.g., the portion of the CH subunit that binds to and/or activates, e.g., an Fc receptor and/or complement. The CDRs typically refer to the Kabat CDRs, as described in Sequences of Proteins of immunological Interest, US Department of Health and Human Services (1991), eds. Kabat *et al.* Another standard for characterizing the antigen binding site is to refer to the hypervariable loops as described by Chothia. See, e.g., Chothia, *et al.* (1987; J. Mol. Biol. 227:799-817); and Tomlinson *et al.* (1995) EMBO J. 14: 4628-4638. Still
30 another standard is the AbM definition used by Oxford Molecular's AbM antibody modeling software. See, generally, e.g., Protein Sequence and Structure Analysis of Antibody Variable Domains. In: Antibody Engineering Lab Manual (Ed.: Duebel, S. and Kontermann, R., Springer-Verlag, Heidelberg). Embodiments described with respect to Kabat CDRs can alternatively be

implemented using similar described relationships with respect to Chothia hypervariable loops or to the AbM-defined loops.

The sequence of antibody genes after assembly and somatic mutation is highly varied, and these varied genes are estimated to encode 10^{10} different antibody molecules (Immunoglobulin Genes, 2nd ed., eds. Jonio et al., Academic Press, San Diego, CA, 1995). Accordingly, the immune system provides a repertoire of immunoglobulins. The term "repertoire" refers to at least one nucleotide sequence derived wholly or partially from at least one sequence encoding at least one immunoglobulin. The sequence(s) may be generated by rearrangement in vivo of the V, D, and J segments of heavy chains, and the V and J segments of light chains. Alternatively, the sequence(s) can be generated from a cell in response to which rearrangement occurs, e.g., in vitro stimulation. Alternatively, part or all of the sequence(s) may be obtained by DNA splicing, nucleotide synthesis, mutagenesis, and other methods, see, e.g., U.S. Patent 5,565,332. A repertoire may include only one sequence or may include a plurality of sequences, including ones in a genetically diverse collection.

In one embodiment, the first binding domain of the binding molecule of the invention comprises a VH region comprising CDR-H1, CDR-H2 and CDR-H3 and a VL region comprising CDR-L1, CDR-L2 and CDR-L3 selected from the group consisting of:

- (1) CDR-H1 as depicted in SEQ ID NO: 1, CDR-H2 as depicted in SEQ ID NO: 2, CDR-H3 as depicted in SEQ ID NO: 3, CDR-L1 as depicted in SEQ ID NO: 4, CDR-L2 as depicted in SEQ ID NO: 5 and CDR-L3 as depicted in SEQ ID NO: 6;
- (2) CDR-H1 as depicted in SEQ ID NO: 11, CDR-H2 as depicted in SEQ ID NO: 12, CDR-H3 as depicted in SEQ ID NO: 13, CDR-L1 as depicted in SEQ ID NO: 14, CDR-L2 as depicted in SEQ ID NO: 15 and CDR-L3 as depicted in SEQ ID NO: 16;
- (3) CDR-H1 as depicted in SEQ ID NO: 21, CDR-H2 as depicted in SEQ ID NO: 22, CDR-H3 as depicted in SEQ ID NO: 23, CDR-L1 as depicted in SEQ ID NO: 24, CDR-L2 as depicted in SEQ ID NO: 25 and CDR-L3 as depicted in SEQ ID NO: 26;
- (4) CDR-H1 as depicted in SEQ ID NO: 31, CDR-H2 as depicted in SEQ ID NO: 32, CDR-H3 as depicted in SEQ ID NO: 33, CDR-L1 as depicted in SEQ ID NO: 34, CDR-L2 as depicted in SEQ ID NO: 35 and CDR-L3 as depicted in SEQ ID NO: 36;
- (5) CDR-H1 as depicted in SEQ ID NO: 41, CDR-H2 as depicted in SEQ ID NO: 42, CDR-H3 as depicted in SEQ ID NO: 43, CDR-L1 as depicted in SEQ ID NO: 44, CDR-L2 as depicted in SEQ ID NO: 45 and CDR-L3 as depicted in SEQ ID NO: 46;

- (6) CDR-H1 as depicted in SEQ ID NO: 51, CDR-H2 as depicted in SEQ ID NO: 52, CDR-H3 as depicted in SEQ ID NO: 53, CDR-L1 as depicted in SEQ ID NO: 54, CDR-L2 as depicted in SEQ ID NO: 55 and CDR-L3 as depicted in SEQ ID NO: 56;
- 5 (7) CDR-H1 as depicted in SEQ ID NO: 61, CDR-H2 as depicted in SEQ ID NO: 62, CDR-H3 as depicted in SEQ ID NO: 63, CDR-L1 as depicted in SEQ ID NO: 64, CDR-L2 as depicted in SEQ ID NO: 65 and CDR-L3 as depicted in SEQ ID NO: 66;
- (8) CDR-H1 as depicted in SEQ ID NO: 71, CDR-H2 as depicted in SEQ ID NO: 72, CDR-H3 as depicted in SEQ ID NO: 73, CDR-L1 as depicted in SEQ ID NO: 74, CDR-L2 as depicted in SEQ ID NO: 75 and CDR-L3 as depicted in SEQ ID NO: 76;
- 10 (9) CDR-H1 as depicted in SEQ ID NO: 161, CDR-H2 as depicted in SEQ ID NO: 162, CDR-H3 as depicted in SEQ ID NO: 163, CDR-L1 as depicted in SEQ ID NO: 164, CDR-L2 as depicted in SEQ ID NO: 165 and CDR-L3 as depicted in SEQ ID NO: 166;
- (10) CDR-H1 as depicted in SEQ ID NO: 171, CDR-H2 as depicted in SEQ ID NO: 172, CDR-H3 as depicted in SEQ ID NO: 173, CDR-L1 as depicted in SEQ ID NO: 174, CDR-L2 as depicted in SEQ ID NO: 175 and CDR-L3 as depicted in SEQ ID NO: 176;
- 15 (11) CDR-H1 as depicted in SEQ ID NO: 181, CDR-H2 as depicted in SEQ ID NO: 182, CDR-H3 as depicted in SEQ ID NO: 183, CDR-L1 as depicted in SEQ ID NO: 184, CDR-L2 as depicted in SEQ ID NO: 185 and CDR-L3 as depicted in SEQ ID NO: 186;
- (12) CDR-H1 as depicted in SEQ ID NO: 191, CDR-H2 as depicted in SEQ ID NO: 192, CDR-H3 as depicted in SEQ ID NO: 193, CDR-L1 as depicted in SEQ ID NO: 194, CDR-L2 as depicted in SEQ ID NO: 195 and CDR-L3 as depicted in SEQ ID NO: 196;
- 20 (13) CDR-H1 as depicted in SEQ ID NO: 201, CDR-H2 as depicted in SEQ ID NO: 202, CDR-H3 as depicted in SEQ ID NO: 203, CDR-L1 as depicted in SEQ ID NO: 204, CDR-L2 as depicted in SEQ ID NO: 205 and CDR-L3 as depicted in SEQ ID NO: 206;
- 25 (14) CDR-H1 as depicted in SEQ ID NO: 211, CDR-H2 as depicted in SEQ ID NO: 212, CDR-H3 as depicted in SEQ ID NO: 213, CDR-L1 as depicted in SEQ ID NO: 214, CDR-L2 as depicted in SEQ ID NO: 215 and CDR-L3 as depicted in SEQ ID NO: 216;
- (15) CDR-H1 as depicted in SEQ ID NO: 221, CDR-H2 as depicted in SEQ ID NO: 222, CDR-H3 as depicted in SEQ ID NO: 223, CDR-L1 as depicted in SEQ ID NO: 224, CDR-L2 as depicted in SEQ ID NO: 225 and CDR-L3 as depicted in SEQ ID NO: 226;
- 30 (16) CDR-H1 as depicted in SEQ ID NO: 311, CDR-H2 as depicted in SEQ ID NO: 312, CDR-H3 as depicted in SEQ ID NO: 313, CDR-L1 as depicted in SEQ ID NO: 314, CDR-L2 as depicted in SEQ ID NO: 315 and CDR-L3 as depicted in SEQ ID NO: 316;

- (17) CDR-H1 as depicted in SEQ ID NO: 321, CDR-H2 as depicted in SEQ ID NO: 322, CDR-H3 as depicted in SEQ ID NO: 323, CDR-L1 as depicted in SEQ ID NO: 324, CDR-L2 as depicted in SEQ ID NO: 325 and CDR-L3 as depicted in SEQ ID NO: 326;
- 5 (18) CDR-H1 as depicted in SEQ ID NO: 331, CDR-H2 as depicted in SEQ ID NO: 332, CDR-H3 as depicted in SEQ ID NO: 333, CDR-L1 as depicted in SEQ ID NO: 334, CDR-L2 as depicted in SEQ ID NO: 335 and CDR-L3 as depicted in SEQ ID NO: 336;
- (19) CDR-H1 as depicted in SEQ ID NO: 341, CDR-H2 as depicted in SEQ ID NO: 342, CDR-H3 as depicted in SEQ ID NO: 343, CDR-L1 as depicted in SEQ ID NO: 344, CDR-L2 as depicted in SEQ ID NO: 345 and CDR-L3 as depicted in SEQ ID NO: 346;
- 10 (20) CDR-H1 as depicted in SEQ ID NO: 351, CDR-H2 as depicted in SEQ ID NO: 352, CDR-H3 as depicted in SEQ ID NO: 353, CDR-L1 as depicted in SEQ ID NO: 354, CDR-L2 as depicted in SEQ ID NO: 355 and CDR-L3 as depicted in SEQ ID NO: 356;
- (21) CDR-H1 as depicted in SEQ ID NO: 361, CDR-H2 as depicted in SEQ ID NO: 362, CDR-H3 as depicted in SEQ ID NO: 363, CDR-L1 as depicted in SEQ ID NO: 364, CDR-L2 as depicted in SEQ ID NO: 365 and CDR-L3 as depicted in SEQ ID NO: 366;
- 15 (22) CDR-H1 as depicted in SEQ ID NO: 371, CDR-H2 as depicted in SEQ ID NO: 372, CDR-H3 as depicted in SEQ ID NO: 373, CDR-L1 as depicted in SEQ ID NO: 374, CDR-L2 as depicted in SEQ ID NO: 375 and CDR-L3 as depicted in SEQ ID NO: 376;
- (23) CDR-H1 as depicted in SEQ ID NO: 381, CDR-H2 as depicted in SEQ ID NO: 382, CDR-H3 as depicted in SEQ ID NO: 383, CDR-L1 as depicted in SEQ ID NO: 384, CDR-L2 as depicted in SEQ ID NO: 385 and CDR-L3 as depicted in SEQ ID NO: 386;
- 20 (24) CDR-H1 as depicted in SEQ ID NO: 581, CDR-H2 as depicted in SEQ ID NO: 582, CDR-H3 as depicted in SEQ ID NO: 583, CDR-L1 as depicted in SEQ ID NO: 584, CDR-L2 as depicted in SEQ ID NO: 585 and CDR-L3 as depicted in SEQ ID NO: 586;
- 25 (25) CDR-H1 as depicted in SEQ ID NO: 591, CDR-H2 as depicted in SEQ ID NO: 592, CDR-H3 as depicted in SEQ ID NO: 593, CDR-L1 as depicted in SEQ ID NO: 594, CDR-L2 as depicted in SEQ ID NO: 595 and CDR-L3 as depicted in SEQ ID NO: 596;
- (26) CDR-H1 as depicted in SEQ ID NO: 601, CDR-H2 as depicted in SEQ ID NO: 602, CDR-H3 as depicted in SEQ ID NO: 603, CDR-L1 as depicted in SEQ ID NO: 604, CDR-L2 as depicted in SEQ ID NO: 605 and CDR-L3 as depicted in SEQ ID NO: 606;
- 30 (27) CDR-H1 as depicted in SEQ ID NO: 611, CDR-H2 as depicted in SEQ ID NO: 612, CDR-H3 as depicted in SEQ ID NO: 613, CDR-L1 as depicted in SEQ ID NO: 614, CDR-L2 as depicted in SEQ ID NO: 615 and CDR-L3 as depicted in SEQ ID NO: 616;

- (28) CDR-H1 as depicted in SEQ ID NO: 621, CDR-H2 as depicted in SEQ ID NO: 622, CDR-H3 as depicted in SEQ ID NO: 623, CDR-L1 as depicted in SEQ ID NO: 624, CDR-L2 as depicted in SEQ ID NO: 625 and CDR-L3 as depicted in SEQ ID NO: 626;
- 5 (29) CDR-H1 as depicted in SEQ ID NO: 631, CDR-H2 as depicted in SEQ ID NO: 632, CDR-H3 as depicted in SEQ ID NO: 633, CDR-L1 as depicted in SEQ ID NO: 634, CDR-L2 as depicted in SEQ ID NO: 635 and CDR-L3 as depicted in SEQ ID NO: 636;
- (30) CDR-H1 as depicted in SEQ ID NO: 641, CDR-H2 as depicted in SEQ ID NO: 642, CDR-H3 as depicted in SEQ ID NO: 643, CDR-L1 as depicted in SEQ ID NO: 644, CDR-L2 as depicted in SEQ ID NO: 645 and CDR-L3 as depicted in SEQ ID NO: 646;
- 10 (31) CDR-H1 as depicted in SEQ ID NO: 651, CDR-H2 as depicted in SEQ ID NO: 652, CDR-H3 as depicted in SEQ ID NO: 653, CDR-L1 as depicted in SEQ ID NO: 654, CDR-L2 as depicted in SEQ ID NO: 655 and CDR-L3 as depicted in SEQ ID NO: 656;
- (32) CDR-H1 as depicted in SEQ ID NO: 661, CDR-H2 as depicted in SEQ ID NO: 662, CDR-H3 as depicted in SEQ ID NO: 663, CDR-L1 as depicted in SEQ ID NO: 664, CDR-L2 as depicted in SEQ ID NO: 665 and CDR-L3 as depicted in SEQ ID NO: 666;
- 15 (33) CDR-H1 as depicted in SEQ ID NO: 671, CDR-H2 as depicted in SEQ ID NO: 672, CDR-H3 as depicted in SEQ ID NO: 673, CDR-L1 as depicted in SEQ ID NO: 674, CDR-L2 as depicted in SEQ ID NO: 675 and CDR-L3 as depicted in SEQ ID NO: 676;
- (34) CDR-H1 as depicted in SEQ ID NO: 681, CDR-H2 as depicted in SEQ ID NO: 682, CDR-H3 as depicted in SEQ ID NO: 683, CDR-L1 as depicted in SEQ ID NO: 684, CDR-L2 as depicted in SEQ ID NO: 685 and CDR-L3 as depicted in SEQ ID NO: 686;
- 20 (35) CDR-H1 as depicted in SEQ ID NO: 691, CDR-H2 as depicted in SEQ ID NO: 692, CDR-H3 as depicted in SEQ ID NO: 693, CDR-L1 as depicted in SEQ ID NO: 694, CDR-L2 as depicted in SEQ ID NO: 695 and CDR-L3 as depicted in SEQ ID NO: 696;
- 25 (36) CDR-H1 as depicted in SEQ ID NO: 701, CDR-H2 as depicted in SEQ ID NO: 702, CDR-H3 as depicted in SEQ ID NO: 703, CDR-L1 as depicted in SEQ ID NO: 704, CDR-L2 as depicted in SEQ ID NO: 705 and CDR-L3 as depicted in SEQ ID NO: 706;
- (37) CDR-H1 as depicted in SEQ ID NO: 711, CDR-H2 as depicted in SEQ ID NO: 712, CDR-H3 as depicted in SEQ ID NO: 713, CDR-L1 as depicted in SEQ ID NO: 714, CDR-L2 as depicted in SEQ ID NO: 715 and CDR-L3 as depicted in SEQ ID NO: 716;
- 30 (38) CDR-H1 as depicted in SEQ ID NO: 721, CDR-H2 as depicted in SEQ ID NO: 722, CDR-H3 as depicted in SEQ ID NO: 723, CDR-L1 as depicted in SEQ ID NO: 724, CDR-L2 as depicted in SEQ ID NO: 725 and CDR-L3 as depicted in SEQ ID NO: 726;

- (39) CDR-H1 as depicted in SEQ ID NO: 731, CDR-H2 as depicted in SEQ ID NO: 732, CDR-H3 as depicted in SEQ ID NO: 733, CDR-L1 as depicted in SEQ ID NO: 734, CDR-L2 as depicted in SEQ ID NO: 735 and CDR-L3 as depicted in SEQ ID NO: 736;
- 5 (40) CDR-H1 as depicted in SEQ ID NO: 741, CDR-H2 as depicted in SEQ ID NO: 742, CDR-H3 as depicted in SEQ ID NO: 743, CDR-L1 as depicted in SEQ ID NO: 744, CDR-L2 as depicted in SEQ ID NO: 745 and CDR-L3 as depicted in SEQ ID NO: 746;
- (41) CDR-H1 as depicted in SEQ ID NO: 751, CDR-H2 as depicted in SEQ ID NO: 752, CDR-H3 as depicted in SEQ ID NO: 753, CDR-L1 as depicted in SEQ ID NO: 754, CDR-L2 as depicted in SEQ ID NO: 755 and CDR-L3 as depicted in SEQ ID NO: 756;
- 10 (42) CDR-H1 as depicted in SEQ ID NO: 761, CDR-H2 as depicted in SEQ ID NO: 762, CDR-H3 as depicted in SEQ ID NO: 763, CDR-L1 as depicted in SEQ ID NO: 764, CDR-L2 as depicted in SEQ ID NO: 765 and CDR-L3 as depicted in SEQ ID NO: 766;
- (43) CDR-H1 as depicted in SEQ ID NO: 771, CDR-H2 as depicted in SEQ ID NO: 772, CDR-H3 as depicted in SEQ ID NO: 773, CDR-L1 as depicted in SEQ ID NO: 774, CDR-L2 as depicted in SEQ ID NO: 775 and CDR-L3 as depicted in SEQ ID NO: 776;
- 15 (44) CDR-H1 as depicted in SEQ ID NO: 781, CDR-H2 as depicted in SEQ ID NO: 782, CDR-H3 as depicted in SEQ ID NO: 783, CDR-L1 as depicted in SEQ ID NO: 784, CDR-L2 as depicted in SEQ ID NO: 785 and CDR-L3 as depicted in SEQ ID NO: 786;
- (45) CDR-H1 as depicted in SEQ ID NO: 791, CDR-H2 as depicted in SEQ ID NO: 792, CDR-H3 as depicted in SEQ ID NO: 793, CDR-L1 as depicted in SEQ ID NO: 794, CDR-L2 as depicted in SEQ ID NO: 795 and CDR-L3 as depicted in SEQ ID NO: 796;
- 20 (46) CDR-H1 as depicted in SEQ ID NO: 801, CDR-H2 as depicted in SEQ ID NO: 802, CDR-H3 as depicted in SEQ ID NO: 803, CDR-L1 as depicted in SEQ ID NO: 804, CDR-L2 as depicted in SEQ ID NO: 805 and CDR-L3 as depicted in SEQ ID NO: 806;
- 25 (47) CDR-H1 as depicted in SEQ ID NO: 811, CDR-H2 as depicted in SEQ ID NO: 812, CDR-H3 as depicted in SEQ ID NO: 813, CDR-L1 as depicted in SEQ ID NO: 814, CDR-L2 as depicted in SEQ ID NO: 815 and CDR-L3 as depicted in SEQ ID NO: 816;
- (48) CDR-H1 as depicted in SEQ ID NO: 821, CDR-H2 as depicted in SEQ ID NO: 822, CDR-H3 as depicted in SEQ ID NO: 823, CDR-L1 as depicted in SEQ ID NO: 824, CDR-L2 as depicted in SEQ ID NO: 825 and CDR-L3 as depicted in SEQ ID NO: 826;
- 30 (49) CDR-H1 as depicted in SEQ ID NO: 831, CDR-H2 as depicted in SEQ ID NO: 832, CDR-H3 as depicted in SEQ ID NO: 833, CDR-L1 as depicted in SEQ ID NO: 834, CDR-L2 as depicted in SEQ ID NO: 835 and CDR-L3 as depicted in SEQ ID NO: 836;

- (50) CDR-H1 as depicted in SEQ ID NO: 961, CDR-H2 as depicted in SEQ ID NO: 962, CDR-H3 as depicted in SEQ ID NO: 963, CDR-L1 as depicted in SEQ ID NO: 964, CDR-L2 as depicted in SEQ ID NO: 965 and CDR-L3 as depicted in SEQ ID NO: 966;
- (51) CDR-H1 as depicted in SEQ ID NO: 971, CDR-H2 as depicted in SEQ ID NO: 972, CDR-H3 as depicted in SEQ ID NO: 973, CDR-L1 as depicted in SEQ ID NO: 974, CDR-L2 as depicted in SEQ ID NO: 975 and CDR-L3 as depicted in SEQ ID NO: 976;
- (52) CDR-H1 as depicted in SEQ ID NO: 981, CDR-H2 as depicted in SEQ ID NO: 982, CDR-H3 as depicted in SEQ ID NO: 983, CDR-L1 as depicted in SEQ ID NO: 984, CDR-L2 as depicted in SEQ ID NO: 985 and CDR-L3 as depicted in SEQ ID NO: 986; and
- (53) CDR-H1 as depicted in SEQ ID NO: 991, CDR-H2 as depicted in SEQ ID NO: 992, CDR-H3 as depicted in SEQ ID NO: 993, CDR-L1 as depicted in SEQ ID NO: 994, CDR-L2 as depicted in SEQ ID NO: 995 and CDR-L3 as depicted in SEQ ID NO: 996.

In yet another embodiment, the first binding domain of the binding molecule comprises a VH region selected from the group consisting of a VH region as depicted in SEQ ID NO: 7, SEQ ID NO: 17, SEQ ID NO: 27, SEQ ID NO: 37, SEQ ID NO: 47, SEQ ID NO: 57, SEQ ID NO: 67, SEQ ID NO: 77, SEQ ID NO: 167, SEQ ID NO: 177, SEQ ID NO: 187, SEQ ID NO: 197, SEQ ID NO: 207, SEQ ID NO: 217, SEQ ID NO: 227, SEQ ID NO: 317, SEQ ID NO: 327, SEQ ID NO: 337, SEQ ID NO: 347, SEQ ID NO: 357, SEQ ID NO: 367, SEQ ID NO: 377, SEQ ID NO: 387, SEQ ID NO: 587, SEQ ID NO: 597, SEQ ID NO: 607, SEQ ID NO: 617, SEQ ID NO: 627, SEQ ID NO: 637, SEQ ID NO: 647, SEQ ID NO: 657, SEQ ID NO: 667, SEQ ID NO: 677, SEQ ID NO: 687, SEQ ID NO: 697, SEQ ID NO: 707, SEQ ID NO: 717, SEQ ID NO: 727, SEQ ID NO: 737, SEQ ID NO: 747, SEQ ID NO: 757, SEQ ID NO: 767, SEQ ID NO: 777, SEQ ID NO: 787, SEQ ID NO: 797, SEQ ID NO: 807, SEQ ID NO: 817, SEQ ID NO: 827, SEQ ID NO: 837, SEQ ID NO: 967, SEQ ID NO: 977, SEQ ID NO: 987, and SEQ ID NO: 997.

In another embodiment, the first binding domain of the binding molecule comprises a VL region selected from the group consisting of a VL region as depicted in SEQ ID in SEQ ID NO: 8, SEQ ID NO: 18, SEQ ID NO: 28, SEQ ID NO: 38, SEQ ID NO: 48, SEQ ID NO: 58, SEQ ID NO: 68, SEQ ID NO: 78, SEQ ID NO: 168, SEQ ID NO: 178, SEQ ID NO: 188, SEQ ID NO: 198, SEQ ID NO: 208, SEQ ID NO: 218, SEQ ID NO: 228, SEQ ID NO: 318, SEQ ID NO: 328, SEQ ID NO: 338, SEQ ID NO: 348, SEQ ID NO: 358, SEQ ID NO: 368, SEQ ID NO: 378, SEQ ID NO: 388, SEQ ID NO: 588, SEQ ID NO: 598, SEQ ID NO: 608, SEQ ID NO: 618, SEQ ID NO: 628, SEQ ID NO: 638, SEQ ID NO: 648, SEQ ID NO: 658, SEQ ID

NO: 668, SEQ ID NO: 678, SEQ ID NO: 688, SEQ ID NO: 698, SEQ ID NO: 708, SEQ ID NO: 718, SEQ ID NO: 728, SEQ ID NO: 738, SEQ ID NO: 748, SEQ ID NO: 758, SEQ ID NO: 768, SEQ ID NO: 778, SEQ ID NO: 788, SEQ ID NO: 798, SEQ ID NO: 808, SEQ ID NO: 818, SEQ ID NO: 828, SEQ ID NO: 838, SEQ ID NO: 968, SEQ ID NO: 978, SEQ ID NO: 988, and SEQ ID NO: 998.

In one embodiment, the first binding domain of the binding molecule comprises a VH region and a VL region selected from the group consisting of:

- (1) a VH region as depicted in SEQ ID NO: 7, and a VL region as depicted in SEQ ID NO: 8;
- (2) a VH region as depicted in SEQ ID NO: 17, and a VL region as depicted in SEQ ID NO: 18;
- (3) a VH region as depicted in SEQ ID NO: 27, and a VL region as depicted in SEQ ID NO: 28;
- (4) a VH region as depicted in SEQ ID NO: 37, and a VL region as depicted in SEQ ID NO: 38;
- (5) a VH region as depicted in SEQ ID NO: 47, and a VL region as depicted in SEQ ID NO: 48;
- (6) a VH region as depicted in SEQ ID NO: 57, and a VL region as depicted in SEQ ID NO: 58;
- (7) a VH region as depicted in SEQ ID NO: 67, and a VL region as depicted in SEQ ID NO: 68;
- (8) a VH region as depicted in SEQ ID NO: 77, and a VL region as depicted in SEQ ID NO: 78;
- (9) a VH region as depicted in SEQ ID NO: 167, and a VL region as depicted in SEQ ID NO: 168;
- (10) a VH region as depicted in SEQ ID NO: 177, and a VL region as depicted in SEQ ID NO: 178;
- (11) a VH region as depicted in SEQ ID NO: 187, and a VL region as depicted in SEQ ID NO: 188;
- (12) a VH region as depicted in SEQ ID NO: 197, and a VL region as depicted in SEQ ID NO: 198;
- (13) a VH region as depicted in SEQ ID NO: 207, and a VL region as depicted in SEQ ID NO: 208;

- (14) a VH region as depicted in SEQ ID NO: 217, and a VL region as depicted in SEQ ID NO: 218;
- (15) a VH region as depicted in SEQ ID NO: 227, and a VL region as depicted in SEQ ID NO: 228;
- 5 (16) a VH region as depicted in SEQ ID NO: 317, and a VL region as depicted in SEQ ID NO: 318;
- (17) a VH region as depicted in SEQ ID NO: 327, and a VL region as depicted in SEQ ID NO: 328;
- 10 (18) a VH region as depicted in SEQ ID NO: 337, and a VL region as depicted in SEQ ID NO: 338;
- (19) a VH region as depicted in SEQ ID NO: 347, and a VL region as depicted in SEQ ID NO: 348;
- (20) a VH region as depicted in SEQ ID NO: 357, and a VL region as depicted in SEQ ID NO: 358;
- 15 (21) a VH region as depicted in SEQ ID NO: 367, and a VL region as depicted in SEQ ID NO: 368;
- (22) a VH region as depicted in SEQ ID NO: 377, and a VL region as depicted in SEQ ID NO: 378;
- 20 (23) a VH region as depicted in SEQ ID NO: 387, and a VL region as depicted in SEQ ID NO: 388;
- (24) a VH region as depicted in SEQ ID NO: 587, and a VL region as depicted in SEQ ID NO: 588;
- (25) a VH region as depicted in SEQ ID NO: 597, and a VL region as depicted in SEQ ID NO: 598;
- 25 (26) a VH region as depicted in SEQ ID NO: 607, and a VL region as depicted in SEQ ID NO: 608;
- (27) a VH region as depicted in SEQ ID NO: 617, and a VL region as depicted in SEQ ID NO: 618;
- 30 (28) a VH region as depicted in SEQ ID NO: 627, and a VL region as depicted in SEQ ID NO: 628;
- (29) a VH region as depicted in SEQ ID NO: 637, and a VL region as depicted in SEQ ID NO: 638;
- (30) a VH region as depicted in SEQ ID NO: 647, and a VL region as depicted in SEQ ID NO: 648;

- (31) a VH region as depicted in SEQ ID NO: 657, and a VL region as depicted in SEQ ID NO: 658;
- (32) a VH region as depicted in SEQ ID NO: 667, and a VL region as depicted in SEQ ID NO: 668;
- 5 (33) a VH region as depicted in SEQ ID NO: 677, and a VL region as depicted in SEQ ID NO: 678;
- (34) a VH region as depicted in SEQ ID NO: 687, and a VL region as depicted in SEQ ID NO: 688;
- (35) a VH region as depicted in SEQ ID NO: 697, and a VL region as depicted in SEQ ID NO: 698;
- 10 (36) a VH region as depicted in SEQ ID NO: 707, and a VL region as depicted in SEQ ID NO: 708;
- (37) a VH region as depicted in SEQ ID NO: 717, and a VL region as depicted in SEQ ID NO: 718;
- 15 (38) a VH region as depicted in SEQ ID NO: 727, and a VL region as depicted in SEQ ID NO: 728;
- (39) a VH region as depicted in SEQ ID NO: 737, and a VL region as depicted in SEQ ID NO: 738;
- (40) a VH region as depicted in SEQ ID NO: 747, and a VL region as depicted in SEQ ID NO: 748;
- 20 (41) a VH region as depicted in SEQ ID NO: 757, and a VL region as depicted in SEQ ID NO: 758;
- (42) a VH region as depicted in SEQ ID NO: 767, and a VL region as depicted in SEQ ID NO: 768;
- 25 (43) a VH region as depicted in SEQ ID NO: 777, and a VL region as depicted in SEQ ID NO: 778;
- (44) a VH region as depicted in SEQ ID NO: 787, and a VL region as depicted in SEQ ID NO: 788;
- (45) a VH region as depicted in SEQ ID NO: 797, and a VL region as depicted in SEQ ID NO: 798;
- 30 (46) a VH region as depicted in SEQ ID NO: 807, and a VL region as depicted in SEQ ID NO: 808;
- (47) a VH region as depicted in SEQ ID NO: 817, and a VL region as depicted in SEQ ID NO: 818;

- (48) a VH region as depicted in SEQ ID NO: 827, and a VL region as depicted in SEQ ID NO: 828;
- (49) a VH region as depicted in SEQ ID NO: 837, and a VL region as depicted in SEQ ID NO: 838;
- 5 (50) a VH region as depicted in SEQ ID NO: 967, and a VL region as depicted in SEQ ID NO: 968;
- (51) a VH region as depicted in SEQ ID NO: 977, and a VL region as depicted in SEQ ID NO: 978;
- (52) a VH region as depicted in SEQ ID NO: 987, and a VL region as depicted in SEQ ID NO: 988; and
- 10 (53) a VH region as depicted in SEQ ID NO: 997, and a VL region as depicted in SEQ ID NO: 998.

In one example, the first binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 9, SEQ ID NO: 19, SEQ ID NO: 29, SEQ ID NO: 39, SEQ ID NO: 49, SEQ ID NO: 59, SEQ ID NO: 69, SEQ ID NO: 79, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, SEQ ID NO: 229, SEQ ID NO: 319, SEQ ID NO: 329, SEQ ID NO: 339, SEQ ID NO: 349, SEQ ID NO: 359, SEQ ID NO: 369, SEQ ID NO: 379, SEQ ID NO: 389, SEQ ID NO: 589, SEQ ID NO: 599, SEQ ID NO: 609, SEQ ID NO: 619, SEQ ID NO: 629, SEQ ID NO: 639, SEQ ID NO: 649, SEQ ID NO: 659, SEQ ID NO: 669, SEQ ID NO: 679, SEQ ID NO: 689, SEQ ID NO: 699, SEQ ID NO: 709, SEQ ID NO: 719, SEQ ID NO: 729, SEQ ID NO: 739, SEQ ID NO: 749, SEQ ID NO: 759, SEQ ID NO: 769, SEQ ID NO: 779, SEQ ID NO: 789, SEQ ID NO: 799, SEQ ID NO: 809, SEQ ID NO: 819, SEQ ID NO: 829, SEQ ID NO: 839, SEQ ID NO: 969, SEQ ID NO: 979, SEQ ID NO: 989, and SEQ ID NO: 999.

It is preferred that a binding molecule of the present invention has a CDR-H3 region of 12 amino acids in length, wherein a tyrosine (Y) residue is present at position 3, 4 and 12. A preferred CDR-H3 is shown in SEQ ID NOs: 43, 193, 333, 613, 703, 733, 823, or 973. Accordingly, a binding molecule of the present invention has in a preferred embodiment a CDR-H3 shown in of SEQ ID NOs: 43, 193, 333, 613, 703, 733, 823, or 973.

Preferred is a binding molecule having the amino acid sequence shown in SEQ ID NO: 340. Also preferred is a binding molecule having the amino acid sequence shown in or SEQ ID NO: 980.

The binding molecule of the present invention is preferably an "isolated" binding molecule. "Isolated" when used to describe the binding molecule disclosed herein, means a binding molecule that has been identified, separated and/or recovered from a component of its production environment. Preferably, the isolated binding molecule is free of association with all other components from its production environment. Contaminant components of its production environment, such as that resulting from recombinant transfected cells, are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the binding molecule will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Ordinarily, however, an isolated antibody will be prepared by at least one purification step.

Amino acid sequence modifications of the binding molecules described herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of the binding molecules are prepared by introducing appropriate nucleotide changes into the binding molecules nucleic acid, or by peptide synthesis.

Such modifications include, for example, deletions from, and/or insertions into, and/or substitutions of, residues within the amino acid sequences of the binding molecules. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the binding molecules, such as changing the number or position of glycosylation sites. Preferably, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids may be substituted in a CDR, while 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 25 amino acids may be substituted in the framework regions (FRs). The substitutions are preferably conservative substitutions as described herein. Additionally or alternatively, 1, 2, 3, 4, 5, or 6 amino acids may be inserted or deleted in each of the CDRs (of course, dependent on their length), while 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or 25 amino acids may be inserted or deleted in each of the FRs.

A useful method for identification of certain residues or regions of the binding molecules that are preferred locations for mutagenesis is called "alanine scanning mutagenesis" as described by

Cunningham and Wells in Science, 244: 1081-1085 (1989). Here, a residue or group of target residues within the binding molecule is/are identified (e.g. charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with the epitope.

5

Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* needs not to be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at a target codon or region and the expressed binding molecule variants are screened for the desired activity.

10

Preferably, amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 residues to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. An insertional variant of the binding molecule includes the fusion to the N-or C-terminus of the antibody to an enzyme or a fusion to a polypeptide which increases the serum half-life of the antibody.

15

Another type of variant is an amino acid substitution variant. These variants have preferably at least 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 amino acid residues in the binding molecule replaced by a different residue. The sites of greatest interest for substitutional mutagenesis include the CDRs of the heavy and/or light chain, in particular the hypervariable regions, but FR alterations in the heavy and/or light chain are also contemplated.

20

25

For example, if a CDR sequence encompasses 6 amino acids, it is envisaged that one, two or three of these amino acids are substituted. Similarly, if a CDR sequence encompasses 15 amino acids it is envisaged that one, two, three, four, five or six of these amino acids are substituted.

30

Generally, if amino acids are substituted in one or more or all of the CDRs of the heavy and/or light chain, it is preferred that the then-obtained "substituted" sequence is at least 60%, more preferably 65%, even more preferably 70%, particularly preferably 75%, more particularly preferably 80% identical to the "original" CDR sequence. This means that it is dependent of the length of the CDR to which degree it is identical to the "substituted" sequence. For example, a

35

CDR having 5 amino acids is preferably 80% identical to its substituted sequence in order to have at least one amino acid substituted. Accordingly, the CDRs of the binding molecule may have different degrees of identity to their substituted sequences, e.g., CDRL1 may have 80%, while CDRL3 may have 90%.

5

Preferred substitutions (or replacements) are conservative substitutions. However, any substitution (including non-conservative substitution or one or more from the "exemplary substitutions" listed in Table 1, below) is envisaged as long as the binding molecule retains its capability to bind to BCMA via the first binding domain and to CD3 epsilon via the second binding domain and/or its CDRs have an identity to the then substituted sequence (at least 60%, more preferably 65%, even more preferably 70%, particularly preferably 75%, more particularly preferably 80% identical to the "original" CDR sequence).

10

Conservative substitutions are shown in Table 1 under the heading of "preferred substitutions". If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions" in Table 1, or as further described below in reference to amino acid classes, may be introduced and the products screened for a desired characteristic.

15

Table 1: Amino Acid Substitutions

Original	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val, leu, ile	val
Arg (R)	lys, gln, asn	lys
Asn (N)	gln, his, asp, lys, arg	gln
Asp (D)	glu, asn	glu
Cys (C)	ser, ala	ser
Gln (Q)	asn, glu	asn
Glu (E)	asp, gln	asp
Gly (G)	ala	ala
His (H)	asn, gln, lys, arg	arg
Ile (I)	leu, val, met, ala, phe	leu
Leu (L)	norleucine, ile, val, met, ala	ile
Lys (K)	arg, gln, asn	arg
Met (M)	leu, phe, ile	leu
Phe (F)	leu, val, ile, ala, tyr	tyr

Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr, phe	tyr
Tyr (Y)	trp, phe, thr, ser	phe
Val (V)	ile, leu, met, phe, ala	leu

Substantial modifications in the biological properties of the binding molecule of the present invention are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties: (1) hydrophobic: norleucine, met, ala, val, leu, ile; (2) neutral hydrophilic: cys, ser, thr; (3) acidic: asp, glu; (4) basic: asn, gin, his, lys, arg; (5) residues that influence chain orientation: gly, pro; and (6) aromatic : trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Any cysteine residue not involved in maintaining the proper conformation of the binding molecule may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

A particularly preferred type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (e. g. a humanized or human antibody). Generally, the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sites (e. g. 6-7 sites) are mutated to generate all possible amino acid substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (e. g. binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to

identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the binding domain and, e.g., human BCMA. Such contact residues and neighbouring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

Other modifications of the binding molecule are contemplated herein. For example, the binding molecule may be linked to one of a variety of non-proteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol. The binding molecule may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatine-microcapsules and poly (methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nanoparticles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Oslo, A., Ed., (1980).

The binding molecules disclosed herein may also be formulated as immuno-liposomes. A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang *et al.*, Proc. Natl Acad. Sci. USA, 77: 4030 (1980); US Pat. Nos. 4,485,045 and 4,544,545; and W0 97/38731 published October 23, 1997. Liposomes with enhanced circulation time are disclosed in US Patent No. 5,013, 556. Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.* J. Biol. Chem. 257: 286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent is optionally contained within the liposome. See Gabizon *et al.* J. National Cancer Inst. 81 (19) 1484 (1989).

When using recombinant techniques, the binding molecule can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the binding molecule is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, are removed, for example, by centrifugation or ultrafiltration. Carter *et al.*, Bio/Technology 10: 163-167 (1992) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*.

The binding molecule composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique.

In a further aspect, the present invention relates to a nucleic acid sequence encoding a binding molecule of the invention. The term "nucleic acid" is well known to the skilled person and encompasses DNA (such as cDNA) and RNA (such as mRNA). The nucleic acid can be double stranded and single stranded, linear and circular. Said nucleic acid molecule is preferably comprised in a vector which is preferably comprised in a host cell. Said host cell is, e.g. after transformation or transfection with the nucleic acid sequence of the invention, capable of expressing the binding molecule. For that purpose the nucleic acid molecule is operatively linked with control sequences.

A vector is a nucleic acid molecule used as a vehicle to transfer (foreign) genetic material into a cell. The term "vector" encompasses – but is not restricted to – plasmids, viruses, cosmids and artificial chromosomes. In general, engineered vectors comprise an origin of replication, a multicloning site and a selectable marker. The vector itself is generally a nucleotide sequence, commonly a DNA sequence, that comprises an insert (transgene) and a larger sequence that serves as the "backbone" of the vector. Modern vectors may encompass additional features besides the transgene insert and a backbone: promoter, genetic marker, antibiotic resistance, reporter gene, targeting sequence, protein purification tag. Vectors called expression vectors (expression constructs) specifically are for the expression of the transgene in the target cell, and generally have control sequences such as a promoter sequence that drives expression of the transgene. Insertion of a vector into the target cell is usually called "transformation" for bacterial cells, "transfection" for eukaryotic cells, although insertion of a viral vector is also called "transduction".

As used herein, the term "host cell" is intended to refer to a cell into which a nucleic acid encoding the binding molecule of the invention is introduced by way of transformation, transfection and the like. It should be understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications
5 may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

As used herein, the term "expression" includes any step involved in the production of a binding
10 molecule of the invention including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are
15 suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

A nucleic acid is "operably linked" when it is placed into a functional relationship with another
20 nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked"
25 means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The terms "host cell," "target cell" or "recipient cell" are intended to include any individual cell or cell culture that can be or has/have been recipients for vectors or the incorporation of exogenous nucleic acid molecules, polynucleotides and/or proteins. It also is intended to include progeny of a single cell, and the progeny may not necessarily be completely identical (in morphology or in genomic or total DNA complement) to the original parent cell due to natural,
30 accidental, or deliberate mutation. The cells may be prokaryotic or eukaryotic, and include but

are not limited to bacterial cells, yeast cells, animal cells, and mammalian cells, e.g., murine, rat, macaque or human.

5 Suitable host cells include prokaryotes and eukaryotic host cells including yeasts, fungi, insect cells and mammalian cells.

The binding molecule of the invention can be produced in bacteria. After expression, the binding molecule of the invention, preferably the binding molecule is isolated from the *E. coli* cell paste in a soluble fraction and can be purified through, e.g., affinity chromatography and/or size
10 exclusion. Final purification can be carried out similar to the process for purifying antibody expressed e. g, in CHO cells.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for the binding molecule of the invention. *Saccharomyces*
15 *cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe*, Kluyveromyces hosts such as, e.g., *K. lactis*, *K. fragilis* (ATCC 12424), *K. bulgaricus* (ATCC 16045), *K. wickerhamii* (ATCC 24178), *K. waltii* (ATCC 56500), *K. drosophilarum* (ATCC 36906), *K. thermotolerans*, and
20 *K. marxianus*; yarrowia (EP 402 226); *Pichia pastoris* (EP 183 070); *Candida*; *Trichoderma reesia* (EP 244 234); *Neurospora crassa*; Schwanniomyces such as *Schwanniomyces occidentalis*; and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

25 Suitable host cells for the expression of glycosylated binding molecule of the invention, preferably antibody derived binding molecules are derived from multicellular organisms. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruit fly), and *Bombyx mori* have been identified. A variety of viral strains for
30 transfection are publicly available, e. g. , the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, Arabidopsis and tobacco can also be utilized as hosts. Cloning and expression vectors useful in the production of proteins in plant cell culture are known to those of skill in the art. See e.g. Hiatt *et al.*, Nature (1989) 342: 76-78, Owen *et al.* (1992) Bio/Technology 10: 790-794, Artsaenko *et al.* (1995) The Plant J 8: 745-750, and Fecker *et al.* (1996) Plant Mol Biol 32: 979-986.

However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen Virol. 36 : 59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub *et al.*, Proc. Natl. Acad. Sci. USA 77: 4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23: 243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL1587) ; human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, 1413 8065); mouse mammary tumor (MMT 060562, ATCC CCL5 1); TRI cells (Mather *et al.*, Annals N. Y Acad. Sci. 383 : 44-68 (1982)) ; MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

When using recombinant techniques, the binding molecule of the invention can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the binding molecule is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, are removed, for example, by centrifugation or ultrafiltration. Carter *et al.*, Bio/Technology 10: 163-167 (1992) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 min. Cell debris can be removed by centrifugation. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

The binding molecule of the invention prepared from the host cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique.

5 The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly (styrenedivinyl) benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the binding molecule of the invention comprises a CH3 domain, the Bakerbond ABXMresin (J. T. Baker, Phillipsburg, NJ) is useful for purification. Other techniques for protein
10 purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™ chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromato-focusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

15 In another aspect, processes are provided for producing binding molecules of the invention, said processes comprising culturing a host cell defined herein under conditions allowing the expression of the binding molecule and recovering the produced binding molecule from the culture.

20 The term "culturing" refers to the in vitro maintenance, differentiation, growth, proliferation and/or propagation of cells under suitable conditions in a medium.

In an alternative embodiment, compositions are provided comprising a binding molecule of the
25 invention, or produced according to the process of the invention. Preferably, said composition is a pharmaceutical composition.

As used herein, the term "pharmaceutical composition" relates to a composition for administration to a patient, preferably a human patient. The particular preferred pharmaceutical
30 composition of this invention comprises the binding molecule of the invention. Preferably, the pharmaceutical composition comprises suitable formulations of carriers, stabilizers and/or excipients. In a preferred embodiment, the pharmaceutical composition comprises a composition for parenteral, transdermal, intraluminal, intraarterial, intrathecal and/or intranasal administration or by direct injection into tissue. It is in particular envisaged that said composition
35 is administered to a patient via infusion or injection. Administration of the suitable compositions

may be effected by different ways, e.g., by intravenous, intraperitoneal, subcutaneous, intramuscular, topical or intradermal administration. In particular, the present invention provides for an uninterrupted administration of the suitable composition. As a non-limiting example, uninterrupted, i.e. continuous administration may be realized by a small pump system worn by the patient for metering the influx of therapeutic agent into the body of the patient. The pharmaceutical composition comprising the binding molecule of the invention can be administered by using said pump systems. Such pump systems are generally known in the art, and commonly rely on periodic exchange of cartridges containing the therapeutic agent to be infused. When exchanging the cartridge in such a pump system, a temporary interruption of the otherwise uninterrupted flow of therapeutic agent into the body of the patient may ensue. In such a case, the phase of administration prior to cartridge replacement and the phase of administration following cartridge replacement would still be considered within the meaning of the pharmaceutical means and methods of the invention together make up one "uninterrupted administration" of such therapeutic agent.

The continuous or uninterrupted administration of these binding molecules of the invention may be intravenous or subcutaneous by way of a fluid delivery device or small pump system including a fluid driving mechanism for driving fluid out of a reservoir and an actuating mechanism for actuating the driving mechanism. Pump systems for subcutaneous administration may include a needle or a cannula for penetrating the skin of a patient and delivering the suitable composition into the patient's body. Said pump systems may be directly fixed or attached to the skin of the patient independently of a vein, artery or blood vessel, thereby allowing a direct contact between the pump system and the skin of the patient. The pump system can be attached to the skin of the patient for 24 hours up to several days. The pump system may be of small size with a reservoir for small volumes. As a non-limiting example, the volume of the reservoir for the suitable pharmaceutical composition to be administered can be between 0.1 and 50 ml.

The continuous administration may be transdermal by way of a patch worn on the skin and replaced at intervals. One of skill in the art is aware of patch systems for drug delivery suitable for this purpose. It is of note that transdermal administration is especially amenable to uninterrupted administration, as exchange of a first exhausted patch can advantageously be accomplished simultaneously with the placement of a new, second patch, for example on the surface of the skin immediately adjacent to the first exhausted patch and immediately prior to removal of the first exhausted patch. Issues of flow interruption or power cell failure do not arise.

The inventive compositions may further comprise a pharmaceutically acceptable carrier. Examples of suitable pharmaceutical carriers are well known in the art and include solutions, e.g. phosphate buffered saline solutions, water, emulsions, such as oil/water emulsions, various
5 types of wetting agents, sterile solutions, liposomes, etc. Compositions comprising such carriers can be formulated by well known conventional methods. Formulations can comprise carbohydrates, buffer solutions, amino acids and/or surfactants. Carbohydrates may be non-reducing sugars, preferably trehalose, sucrose, octasulfate, sorbitol or xylitol. In general, as used herein, "pharmaceutically acceptable carrier" means any and all solvents, dispersion
10 media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed and include: additional buffering agents; preservatives; co-solvents; antioxidants, including ascorbic acid and
15 methionine; chelating agents such as EDTA; metal complexes (e.g., Zn-protein complexes); biodegradable polymers, such as polyesters; salt-forming counter-ions, such as sodium, polyhydric sugar alcohols; amino acids, such as alanine, glycine, asparagine, 2-phenylalanine, and threonine; sugars or sugar alcohols, such as trehalose, sucrose, octasulfate, sorbitol or xylitol stachyose, mannose, sorbose, xylose, ribose, myoinisitol, galactose, lactitol, ribitol,
20 myoinisitol, galactitol, glycerol, cyclitols (e.g., inositol), polyethylene glycol; sulfur containing reducing agents, such as glutathione, thiocetic acid, sodium thioglycolate, thioglycerol, [alpha]-monothioglycerol, and sodium thio sulfate; low molecular weight proteins, such as human serum albumin, bovine serum albumin, gelatin, or other immunoglobulins; and hydrophilic polymers, such as polyvinylpyrrolidone. Such formulations may be used for continuous administrations
25 which may be intravenous or subcutaneous with and/or without pump systems. Amino acids may be charged amino acids, preferably lysine, lysine acetate, arginine, glutamate and/or histidine. Surfactants may be detergents, preferably with a molecular weight of >1.2 KD and/or a polyether, preferably with a molecular weight of >3 KD. Non-limiting examples for preferred detergents are Tween 20, Tween 40, Tween 60, Tween 80 or Tween 85. Non-limiting examples
30 for preferred polyethers are PEG 3000, PEG 3350, PEG 4000 or PEG 5000. Buffer systems used in the present invention can have a preferred pH of 5-9 and may comprise citrate, succinate, phosphate, histidine and acetate.

The compositions of the present invention can be administered to the subject at a suitable dose
35 which can be determined e.g. by dose escalating studies by administration of increasing doses

of the polypeptide of the invention exhibiting cross-species specificity described herein to non-chimpanzee primates, for instance macaques. As set forth above, the binding molecule of the invention exhibiting cross-species specificity described herein can be advantageously used in identical form in preclinical testing in non-chimpanzee primates and as drug in humans. These compositions can also be administered in combination with other proteinaceous and non-proteinaceous drugs. These drugs may be administered simultaneously with the composition comprising the polypeptide of the invention as defined herein or separately before or after administration of said polypeptide in timely defined intervals and doses. The dosage regimen will be determined by the attending physician and clinical factors. As is well known in the medical arts, dosages for any one patient depend upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently.

Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, inert gases and the like. In addition, the composition of the present invention might comprise proteinaceous carriers, like, e.g., serum albumin or immunoglobulin, preferably of human origin. It is envisaged that the composition of the invention might comprise, in addition to the polypeptide of the invention defined herein, further biologically active agents, depending on the intended use of the composition. Such agents might be drugs acting on the gastro-intestinal system, drugs acting as cytostatica, drugs preventing hyperurikemia, drugs inhibiting immunoreactions (e.g. corticosteroids), drugs modulating the inflammatory response, drugs acting on the circulatory system and/or agents such as cytokines known in the art. It is also envisaged that the binding molecule of the present invention is applied in a co-therapy, i.e., in combination with another anti-cancer medicament.

The biological activity of the pharmaceutical composition defined herein can be determined for instance by cytotoxicity assays, as described in the following examples, in WO 99/54440 or by Schlereth *et al.* (Cancer Immunol. Immunother. 20 (2005), 1-12). "Efficacy" or "*in vivo* efficacy"

as used herein refers to the response to therapy by the pharmaceutical composition of the invention, using e.g. standardized NCI response criteria. The success or *in vivo* efficacy of the therapy using a pharmaceutical composition of the invention refers to the effectiveness of the composition for its intended purpose, i.e. the ability of the composition to cause its desired effect, i.e. depletion of pathologic cells, e.g. tumor cells. The *in vivo* efficacy may be monitored by established standard methods for the respective disease entities including, but not limited to white blood cell counts, differentials, Fluorescence Activated Cell Sorting, bone marrow aspiration. In addition, various disease specific clinical chemistry parameters and other established standard methods may be used. Furthermore, computer-aided tomography, X-ray, nuclear magnetic resonance tomography (e.g. for National Cancer Institute-criteria based response assessment [Cheson BD, Horning SJ, Coiffier B, Shipp MA, Fisher RI, Connors JM, Lister TA, Vose J, Grillo-Lopez A, Hagenbeek A, Cabanillas F, Klippensten D, Hiddemann W, Castellino R, Harris NL, Armitage JO, Carter W, Hoppe R, Canellos GP. Report of an international workshop to standardize response criteria for non-Hodgkin's lymphomas. NCI Sponsored International Working Group. J Clin Oncol. 1999 Apr;17(4):1244]), positron-emission tomography scanning, white blood cell counts, differentials, Fluorescence Activated Cell Sorting, bone marrow aspiration, lymph node biopsies/histologies, and various lymphoma specific clinical chemistry parameters (e.g. lactate dehydrogenase) and other established standard methods may be used.

Another major challenge in the development of drugs such as the pharmaceutical composition of the invention is the predictable modulation of pharmacokinetic properties. To this end, a pharmacokinetic profile of the drug candidate, i.e. a profile of the pharmacokinetic parameters that affect the ability of a particular drug to treat a given condition, can be established. Pharmacokinetic parameters of the drug influencing the ability of a drug for treating a certain disease entity include, but are not limited to: half-life, volume of distribution, hepatic first-pass metabolism and the degree of blood serum binding. The efficacy of a given drug agent can be influenced by each of the parameters mentioned above.

"Half-life" means the time where 50% of an administered drug are eliminated through biological processes, e.g. metabolism, excretion, etc.

By "hepatic first-pass metabolism" is meant the propensity of a drug to be metabolized upon first contact with the liver, i.e. during its first pass through the liver.

"Volume of distribution" means the degree of retention of a drug throughout the various compartments of the body, like e.g. intracellular and extracellular spaces, tissues and organs, etc. and the distribution of the drug within these compartments.

- 5 "Degree of blood serum binding" means the propensity of a drug to interact with and bind to blood serum proteins, such as albumin, leading to a reduction or loss of biological activity of the drug.

10 Pharmacokinetic parameters also include bioavailability, lag time (Tlag), Tmax, absorption rates, more onset and/or Cmax for a given amount of drug administered. "Bioavailability" means the amount of a drug in the blood compartment. "Lag time" means the time delay between the administration of the drug and its detection and measurability in blood or plasma.

15 "Tmax" is the time after which maximal blood concentration of the drug is reached, and "Cmax" is the blood concentration maximally obtained with a given drug. The time to reach a blood or tissue concentration of the drug which is required for its biological effect is influenced by all parameters. Pharmacokinetic parameters of bispecific single chain antibodies exhibiting cross-species specificity, which may be determined in preclinical animal testing in non-chimpanzee primates as outlined above, are also set forth e.g. in the publication by Schlereth *et al.* (Cancer Immunol. Immunother. 20 (2005), 1-12).

25 The term "toxicity" as used herein refers to the toxic effects of a drug manifested in adverse events or severe adverse events. These side events might refer to a lack of tolerability of the drug in general and/or a lack of local tolerance after administration. Toxicity could also include teratogenic or carcinogenic effects caused by the drug.

30 The term "safety", "*in vivo* safety" or "tolerability" as used herein defines the administration of a drug without inducing severe adverse events directly after administration (local tolerance) and during a longer period of application of the drug. "Safety", "*in vivo* safety" or "tolerability" can be evaluated e.g. at regular intervals during the treatment and follow-up period. Measurements include clinical evaluation, e.g. organ manifestations, and screening of laboratory abnormalities. Clinical evaluation may be carried out and deviations to normal findings recorded/coded according to NCI-CTC and/or MedDRA standards. Organ manifestations may include criteria such as allergy/immunology, blood/bone marrow, cardiac arrhythmia, coagulation and the like, 35 as set forth e.g. in the Common Terminology Criteria for adverse events v3.0 (CTCAE).

Laboratory parameters which may be tested include for instance hematology, clinical chemistry, coagulation profile and urine analysis and examination of other body fluids such as serum, plasma, lymphoid or spinal fluid, liquor and the like. Safety can thus be assessed e.g. by physical examination, imaging techniques (i.e. ultrasound, x-ray, CT scans, Magnetic Resonance Imaging (MRI), other measures with technical devices (i.e. electrocardiogram), vital signs, by measuring laboratory parameters and recording adverse events. For example, adverse events in non-chimpanzee primates in the uses and methods according to the invention may be examined by histopathological and/or histochemical methods.

10 The term "effective dose" or "effective dosage" is defined as an amount sufficient to achieve or at least partially achieve the desired effect. The term "therapeutically effective dose" is defined as an amount sufficient to cure or at least partially arrest the disease and its complications in a patient already suffering from the disease. Amounts effective for this use will depend upon the severity of the infection and the general state of the subject's own immune system. The term
15 "patient" includes human and other mammalian subjects that receive either prophylactic or therapeutic treatment.

The term "effective and non-toxic dose" as used herein refers to a tolerable dose of an inventive binding molecule which is high enough to cause depletion of pathologic cells, tumor elimination, tumor shrinkage or stabilization of disease without or essentially without major toxic effects.
20 Such effective and non-toxic doses may be determined e.g. by dose escalation studies described in the art and should be below the dose inducing severe adverse side events (dose limiting toxicity, DLT).

25 The above terms are also referred to e.g. in the Preclinical safety evaluation of biotechnology-derived pharmaceuticals S6; ICH Harmonised Tripartite Guideline; ICH Steering Committee meeting on July 16, 1997.

The appropriate dosage, or therapeutically effective amount, of the binding molecule of the invention will depend on the condition to be treated, the severity of the condition, prior therapy, and the patient's clinical history and response to the therapeutic agent. The proper dose can be adjusted according to the judgment of the attending physician such that it can be administered to the patient one time or over a series of administrations. The pharmaceutical composition can be administered as a sole therapeutic or in combination with additional therapies such as anti-
35 cancer therapies as needed.

The pharmaceutical compositions of this invention are particularly useful for parenteral administration, i.e., subcutaneously, intramuscularly, intravenously, intra-articular and/or intra-synovial. Parenteral administration can be by bolus injection or continuous infusion.

5

If the pharmaceutical composition has been lyophilized, the lyophilized material is first reconstituted in an appropriate liquid prior to administration. The lyophilized material may be reconstituted in, e.g., bacteriostatic water for injection (BWFI), physiological saline, phosphate buffered saline (PBS), or the same formulation the protein had been in prior to lyophilization.

10 Preferably, the binding molecule of the invention or produced by a process of the invention is used in the prevention, treatment or amelioration of a disease selected from a proliferative disease, a tumorous disease, or an immunological disorder.

15 An alternative embodiment of the invention provides a method for the prevention, treatment or amelioration of a disease selected from a proliferative disease, a tumorous disease, or an immunological disorder comprising the step of administering to a patient in the need thereof the binding molecule of the invention or produced by a process of the invention.

20 The formulations described herein are useful as pharmaceutical compositions in the treatment, amelioration and/or prevention of the pathological medical condition as described herein in a patient in need thereof. The term "treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Treatment includes the application or administration of the formulation to the body, an isolated tissue, or cell from a patient who has a disease/disorder, a symptom of a disease/disorder, or a predisposition toward a disease/disorder, with the
25 purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disease, the symptom of the disease, or the predisposition toward the disease.

Those "in need of treatment" include those already with the disorder, as well as those in which the disorder is to be prevented. The term "disease" is any condition that would benefit from
30 treatment with the protein formulation described herein. This includes chronic and acute disorders or diseases including those pathological conditions that predispose the mammal to the disease in question. Non-limiting examples of diseases/disorders to be treated herein include proliferative disease, a tumorous disease, or an immunological disorder.

Preferably, the binding molecule of the invention is for use in the prevention, treatment or amelioration of B cell disorders that correlate with BCMA (over)expression such as plasma cell disorders, and/or autoimmune diseases. The autoimmune disease is, for example, systemic lupus erythematoses or rheumatoid arthritis.

5

Also provided by the present invention is a method for the treatment or amelioration of B cell disorders that correlate with BCMA (over)expression such as plasma cell disorders, and/or autoimmune diseases, comprising the step of administering to a subject in need thereof the binding molecule of the invention. The autoimmune disease is, for example, systemic lupus erythematoses or rheumatoid arthritis.

10

In plasma cell disorders, one clone of plasma cells multiplies uncontrollably. As a result, this clone produces vast amounts of a single (monoclonal) antibody known as the M-protein. In some cases, such as with monoclonal gammopathies, the antibody produced is incomplete, consisting of only light chains or heavy chains. These abnormal plasma cells and the antibodies they produce are usually limited to one type. Preferably, the plasma cell disorder is selected from the group consisting of multiple myeloma, plasmacytoma, plasma cell leukemia, macroglobulinemia, amyloidosis, Waldenstrom's macroglobulinemia, solitary bone plasmacytoma, extramedullary plasmacytoma, osteosclerotic myeloma, heavy chain diseases, monoclonal gammopathy of undetermined significance, and smoldering multiple myeloma.

15

20

In another aspect, kits are provided comprising a binding molecule of the invention, a nucleic acid molecule of the invention, a vector of the invention, or a host cell of the invention. The kit may comprise one or more vials containing the binding molecule and instructions for use. The kit may also contain means for administering the binding molecule of the present invention such as a syringe, pump, infuser or the like, means for reconstituting the binding molecule of the invention and/or means for diluting the binding molecule of the invention.

25

Furthermore, the present invention relates to the use of epitope cluster 3 of BCMA, preferably human BCMA, for the generation of a binding molecule, preferably an antibody, which is capable of binding to BCMA, preferably human BCMA. The epitope cluster 3 of BCMA preferably corresponds to amino acid residues 24 to 41 of the sequence as depicted in SEQ ID NO: 1002.

30

In addition, the present invention provides a method for the generation of an antibody, preferably a bispecific binding molecule, which is capable of binding to BCMA, preferably human BCMA, comprising

- (a) immunizing an animal with a polypeptide comprising epitope cluster 3 of BCMA, preferably human BCMA, wherein epitope cluster 3 of BCMA corresponds to amino acid residues 24 to 41 of the sequence as depicted in SEQ ID NO: 1002,
- (b) obtaining said antibody, and
- (c) optionally converting said antibody into a bispecific binding molecule which is capable of binding to human BCMA and preferably to the T cell CD3 receptor complex.

Preferably, step (b) includes that the obtained antibody is tested as follows:

when the respective epitope cluster in the human BCMA protein is exchanged with the respective epitope cluster of a murine BCMA antigen (resulting in a construct comprising human BCMA, wherein human epitope cluster 3 is replaced with murine epitope cluster 3; see SEQ ID NO: 1011), a decrease in the binding of the antibody will occur. Said decrease is preferably at least 10%, 20%, 30%, 40%, 50%; more preferably at least 60%, 70%, 80%, 90%, 95% or even 100% in comparison to the respective epitope cluster in the human BCMA protein, whereby binding to the respective epitope cluster in the human BCMA protein is set to be 100%. It is envisaged that the aforementioned human BCMA/murine BCMA chimeras are expressed in CHO cells. It is also envisaged that the human BCMA/murine BCMA chimeras are fused with a transmembrane domain and/or cytoplasmic domain of a different membrane-bound protein such as EpCAM; see Figure 2a.

A method to test this loss of binding due to exchange with the respective epitope cluster of a non-human (e.g. murine) BCMA antigen is described in the appended Examples, in particular in Examples 1-3.

The method may further include testing as to whether the antibody binds to epitope cluster 3 of human BCMA and is further capable of binding to epitope cluster 3 of macaque BCMA such as BCMA from *Macaca mulatta* (SEQ ID NO:1017) or *Macaca fascicularis* (SEQ ID NO:1017).

The present invention also provides binding molecules comprising any one of the amino acid sequences shown in SEQ ID NOs: 1-1000 and 1022-1093.

Preferably, a binding molecule comprises three VH CDR sequences (named "VH CDR1", "VH CDR2", "VH CDR3", see 4th column of the appended Sequence Table) from a binding molecule termed "BCMA-(X)", wherein X is 1-100 (see 2nd column of the appended Sequence Table) and/or three VL CDR sequences (named "VL CDR1", "VH CDR2", "VH CDR3", see 4th column

of the appended Sequence Table) from a binding molecule term BCMA-X, wherein X is 1-100 (see 2nd column of the appended Sequence Table).

Preferably, a binding molecule comprises a VH and/or VL sequence as is given in the appended Sequence Table (see 4th column of the appended Sequence Table: "VH" and "VL").

5 Preferably, a binding molecule comprises a scFv sequence as is given in the appended Sequence Table (see 4th column of the appended Sequence Table: "scFv").

Preferably, a binding molecule comprises a bispecific molecule sequence as is given in the appended Sequence Table (see 4th column of the appended Sequence Table: "bispecific molecule").

10

The present invention also relates to a bispecific binding agent comprising at least two binding domains, comprising a first binding domain and a second binding domain, wherein said first binding domain binds to the B cell maturation antigen BCMA and wherein said second binding domain binds to CD3 (item 1) also including the following items:

15 Item 2. The bispecific binding agent of item 1, wherein said first binding domain binds to the extracellular domain of BCMA and said second binding domain binds to the ϵ chain of CD3.

20 Item 3. A bispecific binding agent of item 1 or 2 which is in the format of a full-length antibody or an antibody fragment.

Item 4. A bispecific binding agent of item 3 in the format of a full-length antibody, wherein said first BCMA-binding domain is derived from mouse said and wherein said second CD3-binding domain is derived from rat.

25

Item 5. A bispecific binding agent of item 3, which is in the format of an antibody fragment in the form of a diabody that comprises a heavy chain variable domain connected to a light chain variable domain on the same polypeptide chain such that the two domains do not pair.

30

Item 6. A bispecific binding agent of item 1 or 2 which is in the format of a bispecific single chain antibody that consists of two scFv molecules connected via a linker peptide or by a human serum albumin molecule.

Item 7. The bispecific binding agent of item 6, heavy chain regions (VH) and the corresponding variable light chain regions (VL) are arranged, from N-terminus to C-terminus, in the order

VH(BCMA)-VL(BCMA) -VH(CD3)-VL(CD3),
VH(CD3)-VL(CD3) -VH(BCMA)-VL(BCMA) or
VH CD3)-VL(CD3)-VL(BCMA)-VH(BCMA).

Item 8. A bispecific binding agent of item 1 or 2, which is in the format of a single domain immunoglobulin domain selected from VHHs or VHs.

Item 9. The bispecific binding agent of item 1 or 2, which is in the format of an Fv molecule that has four antibody variable domains with at least two binding domains, wherein at least one binding domain is specific to human BCMA and at least one binding domain is specific to human CD3.

Item 10. A bispecific binding agent of item 1 or 2, which is in the format of a single-chain binding molecule consisting of a first binding domain specific for BCMA, a constant sub-region that is located C-terminal to said first binding domain, a scorpion linker located C-terminal to the constant sub-region, and a second binding domain specific for CD3, which is located C-terminal to said constant sub-region.

Item 11. The bispecific binding agent of item 1 or 2, which is in the format of an antibody-like molecule that binds to BCMA via the two heavy chain/light chain Fv of an antibody or an antibody fragment and which binds to CD3 via a binding domain that has been engineered into non-CDR loops of the heavy chain or the light chain of said antibody or antibody fragment.

Item 12. A bispecific binding agent of item 1 which is in the format of a bispecific ankyrin repeat molecule.

Item 13. A bispecific binding agent of item 1, wherein said first binding domain has a format selected from the formats defined in any one of items 3 to 12 and wherein said second binding domain has a different format selected from the formats defined in any one of items 3 to 12.

Item 14. A bispecific binding agent of item 1 which is a bicyclic peptide.

Item 15. A pharmaceutical composition containing at least one bispecific binding agent of any one of items 1 to 14.

5

Item 16. A bispecific binding agent of any one of items 1 to 14 or a pharmaceutical composition of item 14 for the treatment of plasma cell disorders or other B cell disorders that correlate with BCMA expression and for the treatment of autoimmune diseases.

10 Item 17. A bispecific binding agent of any one of items 1 to 14 or a pharmaceutical composition of item 15 for the treatment of plasma cell disorders selected from plasmacytoma, plasma cell leukemia, multiple myeloma, macroglobulinemia, amyloidosis, Waldenstrom's macroglobulinemia, solitary bone plasmacytoma, extramedullary plasmacytoma, osteosclerotic myeloma, heavy chain diseases,
15 monoclonal gammopathy of undetermined significance, smoldering multiple myeloma.

Variations of the above items are derivable from EP 10 191 418.2 which are also included herein.

It should be understood that the inventions herein are not limited to particular methodology,
20 protocols, or reagents, as such can vary. The discussion and examples provided herein are presented for the purpose of describing particular embodiments only and are not intended to limit the scope of the present invention, which is defined solely by the claims.

All publications and patents cited throughout the text of this specification (including all patents, patent applications, scientific publications, manufacturer's specifications, instructions, etc.),
25 whether supra or infra, are hereby incorporated by reference in their entirety. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention. To the extent the material incorporated by reference contradicts or is inconsistent with this specification, the specification will supersede any such material.

30

The Figures show:

Figure 1:

Sequence alignment of the extracellular domain (ECD) of human BCMA (amino acid residues 1-54 of the full-length protein) and murine BCMA (amino acid residues 1-49 of the full-length protein). Highlighted are the regions (domains or amino acid residues) which were exchanged in the chimeric constructs, as designated for the epitope clustering. Cysteines are depicted by black boxes. Disulfide bonds are indicated.

Figure 2:

Epitope mapping of the BCMA constructs. Human and murine BCMA (figure 2a) as well as seven chimeric human-murine BCMA constructs (figure 2b) expressed on the surface of CHO cells as shown by flow cytometry. The expression of human BCMA on CHO was detected with a monoclonal anti-human BCMA antibody. Murine BCMA expression was detected with a monoclonal anti-murine BCMA-antibody. Bound monoclonal antibody was detected with an anti-rat IgG-Fc-gamma-specific antibody conjugated to phycoerythrin.

Figure 3:

Examples of binding molecules specific for epitope cluster E3, as detected by epitope mapping of the chimeric BCMA constructs (see example 3).

Figure 4:

Determination of binding constants of bispecific binding molecules (anti BCMA x anti CD3) on human and macaque BCMA using the Biacore system. Antigen was immobilized in low to intermediate density (100 RU) on CM5 chip. Dilutions of binders were floated over the chip surface and binding determined using BiaEval Software. Respective off-rates and the binding constant (KD) of the respective binders are depicted below every graph.

Figure 5:

Cytotoxic activity of BCMA bispecific antibodies as measured in an 18-hour ⁵¹chromium release assay. Effector cells: stimulated enriched human CD8 T cells. Target cells: Human BCMA transfected CHO cells (left figure) and macaque BCMA transfected CHO cells (right figure). Effector to target cell (E:T) ratio: 10:1.

Figure 6:

Determination of binding constants of BCMA/CD3 bispecific antibodies of epitope cluster E3 on human and macaque BCMA and on human and macaque CD3 using the Biacore system. Antigen was immobilized in low to intermediate density (100-200 RU) on CM5 chip. Dilutions of bispecific antibodies were floated over the chip surface and binding determined using BiaEval
5 Software. Respective on- and off-rates and the resulting binding constant (KD) of the respective bispecific antibodies are depicted below every graph.

Figure 7:

FACS analysis of BCMA/CD3 bispecific antibodies of epitope cluster E3 on indicated cell lines:
10 1) human BCMA transfected CHO cells, 2) human CD3 positive human T cell line HBP-ALL, 3) macaque BCMA transfected CHO cells, 4) macaque T cell line 4119 LnPx , 5) BCMA-positive human multiple myeloma cell line NCI-H929 and 6) untransfected CHO cells. Negative controls [1) to 6)]: detection antibodies without prior BCMA/CD3 bispecific antibody.

Figure 8:

Scatchard analysis of BCMA/CD3 bispecific antibodies on BCMA-expressing cells. Cells were incubated with increasing concentrations of monomeric antibody until saturation. Antibodies were detected by flow cytometry. Values of triplicate measurements were plotted as hyperbolic curves and as sigmoid curves to demonstrate a valid concentration range used. Maximal
20 binding was determined using Scatchard evaluation, and the respective KD values were calculated.

Figure 9:

Cytotoxic activity of BCMA/CD3 bispecific antibodies of epitope cluster E3, as measured in an
25 18-hour 51-chromium release assay against CHO cells transfected with human BCMA. Effector cells: stimulated enriched human CD8 T cells. Effector to target cell (E:T) ratio: 10:1.

Figure 10:

Cytotoxic activity of BCMA/CD3 bispecific antibodies of epitope cluster E3 as measured in a 48-
30 hour FACS-based cytotoxicity assay. Effector cells: unstimulated human PBMC. Target cells: CHO cells transfected with human BCMA. Effector to target cell (E:T)-ratio: 10:1.

Figure 11:

FACS analysis of BCMA/CD3 bispecific antibodies of epitope cluster E3 on BAFF-R and TACI transfected CHO cells. Cell lines: 1) human BAFF-R transfected CHO cells, 2) human TACI transfected CHO cells 3) multiple myeloma cell line L363; negative controls: detection antibodies without prior BCMA/CD3 bispecific antibody. Positive controls: BAFF-R detection: 5 goat anti hu BAFF-R (R&D AF1162; 1:20) detected by anti-goat antibody-PE (Jackson 705-116-147; 1:50) TACI-detection: rabbit anti TACI antibody (abcam AB 79023; 1:100) detected by goat anti rabbit-antibody PE (Sigma P9757; 1:20).

Figure 12:

10 Cytotoxic activity of BCMA/CD3 bispecific antibodies as measured in an 18-hour ⁵¹-chromium release assay. Effector cells: stimulated enriched human CD8 T cells. Target cells: BCMA-positive human multiple myeloma cell line L363 (i.e. natural expresser). Effector to target cell (E:T) ratio: 10:1.

15 **Figure 13:**

Cytotoxic activity of BCMA/CD3 bispecific antibodies as measured in a 48-hour FACS-based cytotoxicity assay. Effector cells: unstimulated human PBMC. Target cells: human multiple myeloma cell line L363 (natural BCMA expresser). Effector to target cell (E:T)-ratio: 10:1.

20 **Figure 14:**

Cytotoxic activity of BCMA/CD3 bispecific antibodies as measured in a 48-hour FACS-based cytotoxicity assay. Effector cells: unstimulated human PBMC. Target cells: BCMA-positive human multiple myeloma cell line NCI-H929. Effector to target cell (E:T)-ratio: 10:1.

25 **Figure 15:**

Cytotoxic activity of BCMA/CD3 bispecific antibodies as measured in a 48-hour FACS-based cytotoxicity assay. Effector cells: macaque T cell line 4119LnPx. Target cells: CHO cells transfected with macaque BCMA. Effector to target cell (E:T) ratio: 10:1.

30 **Figure 16:**

Anti-tumor activity of BCMA/CD3 bispecific antibodies of epitope cluster E3 in an advanced-stage NCI-H929 xenograft model (see Example 16).

Figure 17:

FACS-based cytotoxicity assay using human multiple myeloma cell lines NCI-H929, L-363 and OPM-2 as target cells and human PBMC as effector cells (48h; E:T = 10:1). The figure depicts the cytokine levels [pg/ml] which were determined for IL-2, IL-6, IL-10, TNF and IFN-gamma at increasing concentrations of the BCMA/CD3 bispecific antibodies of epitope cluster E3 (see
5 Example 22).

Examples:

The following examples illustrate the invention. These examples should not be construed as to limit the scope of this invention. The examples are included for purposes of illustration, and the present invention is limited only by the claims.

5

Example 1**Generation of CHO cells expressing chimeric BCMA**

For the construction of the chimeric epitope mapping molecules, the amino acid sequence of the respective epitope domains or the single amino acid residue of human BCMA was changed to the murine sequence. The following molecules were constructed:

10

- Human BCMA ECD / E1 murine (SEQ ID NO: 1009)

15

Chimeric extracellular BCMA domain: Human extracellular BCMA domain wherein epitope cluster 1 (amino acid residues 1-7 of SEQ ID NO: 1002 or 1007) is replaced by the respective murine cluster (amino acid residues 1-4 of SEQ ID NO: 1004 or 1008)
→ deletion of amino acid residues 1-3 and G6Q mutation in SEQ ID NO: 1002 or 1007

- Human BCMA ECD / E2 murine (SEQ ID NO: 1010)

20

Chimeric extracellular BCMA domain: Human extracellular BCMA domain wherein epitope cluster 2 (amino acid residues 8-21 of SEQ ID NO: 1002 or 1007) is replaced by the respective murine cluster (amino acid residues 5-18 of SEQ ID NO: 1004 or 1008)
→ S9F, Q10H, and N11S mutations in SEQ ID NO: 1002 or 1007

- Human BCMA ECD / E3 murine (SEQ ID NO: 1011)

25

Chimeric extracellular BCMA domain: Human extracellular BCMA domain wherein epitope cluster 3 (amino acid residues 24-41 of SEQ ID NO: 1002 or 1007) is replaced by the respective murine cluster (amino acid residues 21-36 of SEQ ID NO: 1004 or 1008)
→ deletion of amino acid residues 31 and 32 and Q25H, S30N, L35A, and R39P mutation in SEQ ID NO: 1002 or 1007

30

- Human BCMA ECD / E4 murine (SEQ ID NO: 1012)

Chimeric extracellular BCMA domain: Human extracellular BCMA domain wherein epitope cluster 4 (amino acid residues 42-54 of SEQ ID NO: 1002 or 1007) is replaced by the respective murine cluster (amino acid residues 37-49 of SEQ ID NO: 1004 or 1008)
→ N42D, A43P, N47S, N53Y and A54T mutations in SEQ ID NO: 1002 or 1007

35

- Human BCMA ECD / E5 murine (SEQ ID NO: 1013)

Chimeric extracellular BCMA domain: Human extracellular BCMA domain wherein the amino acid residue at position 22 of SEQ ID NO: 1002 or 1007 (isoleucine) is replaced by its
 5 respective murine amino acid residue of SEQ ID NO: 1004 or 1008 (lysine, position 19)
 → I22K mutation in SEQ ID NO: 1002 or 1007

- Human BCMA ECD / E6 murine (SEQ ID NO: 1014)

Chimeric extracellular BCMA domain: Human extracellular BCMA domain wherein the amino
 10 acid residue at position 25 of SEQ ID NO: 1002 or 1007 (glutamine) is replaced by its respective
 murine amino acid residue of SEQ ID NO: 1004 or 1008 (histidine, position 22)
 → Q25H mutation in SEQ ID NO: 1002 or 1007

- Human BCMA ECD / E7 murine (SEQ ID NO: 1015)

Chimeric extracellular BCMA domain: Human extracellular BCMA domain wherein the amino
 15 acid residue at position 39 of SEQ ID NO: 1002 or 1007 (arginine) is replaced by its respective
 murine amino acid residue of SEQ ID NO: 1004 or 1008 (proline, position 34)
 → R39P mutation in SEQ ID NO: 1002 or 1007

20 A) The cDNA constructs were cloned into the mammalian expression vector pEF-DHFR and
 stably transfected into CHO cells. The expression of human BCMA on CHO cells was verified in
 a FACS assay using a monoclonal anti-human BCMA antibody. Murine BCMA expression was
 demonstrated with a monoclonal anti-mouse BCMA-antibody. The used concentration of the
 BCMA antibodies was 10 µg/ml in PBS/2%FCS. Bound monoclonal antibodies were detected
 25 with an anti-rat-IgG-Fcy-PE (1:100 in PBS/2%FCS; Jackson-Immuno-Research #112-116-071).
 As negative control, cells were incubated with PBS/2% FCS instead of the first antibody. The
 samples were measured by flow cytometry on a FACSCanto II instrument (Becton Dickinson)
 and analyzed by FlowJo software (Version 7.6). The surface expression of human-murine
 BCMA chimeras, transfected CHO cells were analyzed and confirmed in a flow cytometry assay
 30 with different anti-BCMA antibodies (Figure 2).

B) For the generation of CHO cells expressing human, macaque, mouse and human/mouse
 chimeric transmembrane BCMA, the coding sequences of human, macaque, mouse BCMA and
 the human-mouse BCMA chimeras (BCMA sequences as published in GenBank, accession
 35 numbers NM_001192 [human]; NM_011608 [mouse] and XM_001106892 [macaque]) were

obtained by gene synthesis according to standard protocols. The gene synthesis fragments were designed as to contain first a Kozak site for eukaryotic expression of the constructs and the coding sequence of a 19 amino acid immunoglobulin leader peptide, followed in frame by the coding sequence of the BCMA proteins respectively in case of the chimeras with the
5 respective epitope domains of the human sequence exchanged for the murine sequence.

Except for the human BCMA ECD / E4 murine and human BCMA constructs the coding sequence of the extracellular domain of the BCMA proteins was followed in frame by the coding sequence of an artificial Ser1-Gly4-Ser1-linker followed by the intracellular domain of human
10 EpCAM (amino acids 226-314; sequence as published in GenBank accession number NM_002354).

All coding sequences were followed by a stop codon. The gene synthesis fragments were also designed as to introduce suitable restriction sites. The gene synthesis fragments were cloned
15 into a plasmid designated pEF-DHFR (pEF-DHFR is described in Raum *et al.* Cancer Immunol Immunother 50 (2001) 141-150). All aforementioned procedures were carried out according to standard protocols (Sambrook, Molecular Cloning; A Laboratory Manual, 3rd edition, Cold Spring Harbour Laboratory Press, Cold Spring Harbour, New York (2001)). For each antigen a clone with sequence-verified nucleotide sequence was transfected into DHFR deficient CHO
20 cells for eukaryotic expression of the constructs. Eukaryotic protein expression in DHFR deficient CHO cells was performed as described by Kaufman R.J. (1990) Methods Enzymol. 185, 537-566. Gene amplification of the constructs was induced by increasing concentrations of methotrexate (MTX) to a final concentration of up to 20 nM MTX.

Example 2

2.1 Transient expression in HEK 293 cells

Clones of the expression plasmids with sequence-verified nucleotide sequences were used for transfection and protein expression in the FreeStyle 293 Expression System (Invitrogen GmbH,
30 Karlsruhe, Germany) according to the manufacturer's protocol. Supernatants containing the expressed proteins were obtained, cells were removed by centrifugation and the supernatants were stored at -20 C.

2.2 Stable expression in CHO cells

Clones of the expression plasmids with sequence-verified nucleotide sequences were transfected into DHFR deficient CHO cells for eukaryotic expression of the constructs. Eukaryotic protein expression in DHFR deficient CHO cells was performed as described by Kaufman R.J. (1990) Methods Enzymol. 185, 537-566. Gene amplification of the constructs was induced by increasing concentrations of methotrexate (MTX) to a final concentration of 20 nM MTX. After two passages of stationary culture the cells were grown in roller bottles with nucleoside-free HyQ PF CHO liquid soy medium (with 4.0 mM L-Glutamine with 0.1 % Pluronic F – 68; HyClone) for 7 days before harvest. The cells were removed by centrifugation and the supernatant containing the expressed protein was stored at -20 C.

2.3 Protein purification

Purification of soluble BCMA proteins was performed as follows: Äkta® Explorer System (GE Healthcare) and Unicorn® Software were used for chromatography. Immobilized metal affinity chromatography ("IMAC") was performed using a Fractogel EMD chelate® (Merck) which was loaded with ZnCl₂ according to the protocol provided by the manufacturer. The column was equilibrated with buffer A (20 mM sodium phosphate buffer pH 7.2, 0.1 M NaCl) and the filtrated (0.2 µm) cell culture supernatant was applied to the column (10 ml) at a flow rate of 3 ml/min. The column was washed with buffer A to remove unbound sample. Bound protein was eluted using a two-step gradient of buffer B (20 mM sodium phosphate buffer pH 7.2, 0.1 M NaCl, 0.5 M imidazole) according to the following procedure:

Step 1: 10 % buffer B in 6 column volumes

Step 2: 100 % buffer B in 6 column volumes

Eluted protein fractions from step 2 were pooled for further purification. All chemicals were of research grade and purchased from Sigma (Deisenhofen) or Merck (Darmstadt).

Gel filtration chromatography was performed on a HiLoad 16/60 Superdex 200 prep grade column (GE/Amersham) equilibrated with Equi-buffer (10 mM citrate, 25 mM lysine-HCl, pH 7.2 for proteins expressed in HEK cells and PBS pH 7.4 for proteins expressed in CHO cells). Eluted protein samples (flow rate 1 ml/min) were subjected to standard SDS-PAGE and Western Blot for detection. Protein concentrations were determined using OD₂₈₀ nm.

Proteins obtained via transient expression in HEK 293 cells were used for immunizations. Proteins obtained via stable expression in CHO cells were used for selection of binders and for measurement of binding.

Example 3

Epitope clustering of murine scFv-fragments

Cells transfected with human or murine BCMA, or with chimeric BCMA molecules were stained with crude, undiluted periplasmic extract containing scFv binding to human/maaque BCMA. Bound scFv were detected with 1 µg/ml of an anti-FLAG antibody (Sigma F1804) and a R-PE-labeled anti-mouse Fc gamma-specific antibody (1:100; Dianova #115-116-071). All antibodies
5 were diluted in PBS with 2% FCS. As negative control, cells were incubated with PBS/2% FCS instead of the periplasmic extract. The samples were measured by flow cytometry on a FACSCanto II instrument (Becton Dickinson) and analyzed by FlowJo software (Version 7.6); see Figure 3.

10 **Example 4**

Procurement of different recombinant forms of soluble human and macaque BCMA

A) The coding sequences of human and rhesus BCMA (as published in GenBank, accession numbers NM_001192 [human], XM_001106892 [rhesus]) coding sequences of human albumin, human Fcγ1 and murine albumin were used for the construction of artificial cDNA sequences
15 encoding soluble fusion proteins of human and macaque BCMA respectively and human albumin, human IgG1 Fc and murine albumin respectively as well as soluble proteins comprising only the extracellular domains of BCMA. To generate the constructs for expression of the soluble human and macaque BCMA proteins, cDNA fragments were obtained by PCR mutagenesis of the full-length BCMA cDNAs described above and molecular cloning according
20 to standard protocols.

For the fusions with human albumin, the modified cDNA fragments were designed as to contain first a Kozak site for eukaryotic expression of the constructs followed by the coding sequence of the human and rhesus (or *Macaca mulatta*) BCMA proteins respectively, comprising amino acids 1 to 54 and 1 to 53 corresponding to the extracellular domain of human and rhesus
25 BCMA, respectively, followed in frame by the coding sequence of an artificial Ser1-Gly4-Ser1-linker, followed in frame by the coding sequence of human serum albumin, followed in frame by the coding sequence of a Flag tag, followed in frame by the coding sequence of a modified histidine tag (SGHHGGHHGGHH) and a stop codon.

For the fusions with murine IgG1, the modified cDNA fragments were designed as to contain first a Kozak site for eukaryotic expression of the constructs followed by the coding sequence of the human and macaque BCMA proteins respectively, comprising amino acids 1 to 54 and 1 to
30 53 corresponding to the extracellular domain of human and rhesus BCMA, respectively, followed in frame by the coding sequence of an artificial Ser1-Gly4-Ser1-linker, followed in frame by the coding sequence of the hinge and Fc gamma portion of human IgG1, followed in
35 frame by the coding sequence of a hexahistidine tag and a stop codon.

For the fusions with murine albumin, the modified cDNA fragments were designed as to contain first a Kozak site for eukaryotic expression of the constructs followed by the coding sequence of the human and macaque BCMA proteins respectively, comprising amino acids 1 to 54 and 1 to 53 corresponding to the extracellular domain of human and rhesus BCMA, respectively, followed in frame by the coding sequence of an artificial Ser1-Gly4-Ser1-linker, followed in frame by the coding sequence of murine serum albumin, followed in frame by the coding sequence of a Flag tag, followed in frame by the coding sequence of a modified histidine tag (SGHHGGHHGGHH) and a stop codon.

For the soluble extracellular domain constructs, the modified cDNA fragments were designed as to contain first a Kozak site for eukaryotic expression of the constructs followed by the coding sequence of the human and macaque BCMA proteins respectively, comprising amino acids 1 to 54 and 1 to 53 corresponding to the extracellular domain of human and rhesus BCMA, respectively, followed in frame by the coding sequence of an artificial Ser1-Gly1-linker, followed in frame by the coding sequence of a Flag tag, followed in frame by the coding sequence of a modified histidine tag (SGHHGGHHGGHH) and a stop codon.

The cDNA fragments were also designed to introduce restriction sites at the beginning and at the end of the fragments. The introduced restriction sites, EcoRI at the 5' end and Sall at the 3' end, were utilized in the following cloning procedures. The cDNA fragments were cloned via EcoRI and Sall into a plasmid designated pEF-DHFR (pEF-DHFR is described in Raum et al. Cancer Immunol Immunother 50 (2001) 141-150). The aforementioned procedures were all carried out according to standard protocols (Sambrook, Molecular Cloning; A Laboratory Manual, 3rd edition, Cold Spring Harbour Laboratory Press, Cold Spring Harbour, New York (2001)).

B) The coding sequences of human and macaque BCMA as described above and coding sequences of human albumin, human Fcγ1, murine Fcγ1, murine Fcγ2a, murine albumin, rat albumin, rat Fcγ1 and rat Fcγ2b were used for the construction of artificial cDNA sequences encoding soluble fusion proteins of human and macaque BCMA respectively and human albumin, human IgG1 Fc, murine IgG1 Fc, murine IgG2a Fc, murine albumin, rat IgG1 Fc, rat IgG2b and rat albumin respectively as well as soluble proteins comprising only the extracellular domains of BCMA. To generate the constructs for expression of the soluble human and macaque BCMA proteins cDNA fragments were obtained by PCR mutagenesis of the full-length BCMA cDNAs described above and molecular cloning according to standard protocols.

For the fusions with albumins the modified cDNA fragments were designed as to contain first a Kozak site for eukaryotic expression of the constructs and the coding sequence of a 19 amino acid immunoglobulin leader peptide, followed in frame by the coding sequence of the

extracellular domain of the respective BCMA protein followed in frame by the coding sequence of an artificial Ser1-Gly4-Ser1-linker, followed in frame by the coding sequence of the respective serum albumin, followed in frame by the coding sequence of a Flag tag, followed in frame by the coding sequence of a modified histidine tag (SGHHGGHHGGHH) and a stop codon.

5 For the fusions with IgG Fcs the modified cDNA fragments were designed as to contain first a Kozak site for eukaryotic expression of the constructs and the coding sequence of a 19 amino acid immunoglobulin leader peptide, followed in frame by the coding sequence of the extracellular domain of the respective BCMA protein followed in frame by the coding sequence of an artificial Ser1-Gly4-Ser1-linker, except for human IgG1 Fc where an artificial Ser1-Gly1-
10 linker was used, followed in frame by the coding sequence of the hinge and Fc gamma portion of the respective IgG, followed in frame by the coding sequence of a Flag tag, followed in frame by the coding sequence of a modified histidine tag (SGHHGGHHGGHH) and a stop codon.

For the soluble extracellular domain constructs the modified cDNA fragments were designed as to contain first a Kozak site for eukaryotic expression of the constructs and the coding sequence
15 of a 19 amino acid immunoglobulin leader peptide, followed in frame by the coding sequence of the extracellular domain of the respective BCMA protein followed in frame by the coding sequence of an artificial Ser1-Gly1-linker, followed in frame by the coding sequence of a Flag tag, followed in frame by the coding sequence of a modified histidine tag (SGHHGGHHGGHH) and a stop codon.

20 For cloning of the constructs suitable restriction sites were introduced. The cDNA fragments were all cloned into a plasmid designated pEF-DHFR (pEF-DHFR is described in Raum *et al.* 2001). The aforementioned procedures were all carried out according to standard protocols (Sambrook, 2001).

The following constructs were designed to enable directed panning on distinct epitopes. The
25 coding sequence of murine-human BCMA chimeras and murine-macaque BCMA chimeras (mouse, human and macaque BCMA sequences as described above) and coding sequences of murine albumin and murine Fcγ1 were used for the construction of artificial cDNA sequences encoding soluble fusion proteins of murine-human and murine-macaque BCMA chimeras respectively and murine IgG1 Fc and murine albumin, respectively. To generate the constructs
30 for expression of the soluble murine-human and murine-macaque BCMA chimeras cDNA fragments of murine BCMA (amino acid 1-49) with the respective epitope domains mutated to the human and macaque sequence respectively were obtained by gene synthesis according to standard protocols. Cloning of constructs was carried out as described above and according to standard protocols (Sambrook, 2001).

35 The following molecules were constructed:

- amino acid 1-4 human, murine IgG1 Fc
- amino acid 1-4 human, murine albumin
- amino acid 1-4 rhesus, murine IgG1 Fc
- amino acid 1-4 rhesus, murine albumin
- 5 • amino acid 5-18 human, murine IgG1 Fc
- amino acid 5-18 human, murine albumin
- amino acid 5-18 rhesus, murine IgG1 Fc
- amino acid 5-18 rhesus, murine albumin
- amino acid 37-49 human, murine IgG1 Fc
- 10 • amino acid 37-49 human, murine albumin
- amino acid 37-49 rhesus, murine IgG1 Fc
- amino acid 37-49 rhesus, murine albumin

15 **Example 5**

5.1 Biacore-based determination of bispecific antibody affinity to human and macaque BCMA and CD3

Biacore analysis experiments were performed using recombinant BCMA fusion proteins with human serum albumin (ALB) to determine BCMA target binding. For CD3 affinity
 20 measurements, recombinant fusion proteins having the N-terminal 27 amino acids of the CD3 epsilon (CD3e) fused to human antibody Fc portion were used. This recombinant protein exists in a human CD3e1-27 version and in a cynomolgous CD3e version, both bearing the epitope of the CD3 binder in the bispecific antibodies.

In detail, CM5 Sensor Chips (GE Healthcare) were immobilized with approximately 100 to
 25 150 RU of the respective recombinant antigen using acetate buffer pH4.5 according to the manufacturer's manual. The bispecific antibody samples were loaded in five concentrations: 50 nM, 25 nM, 12.5 nM, 6.25 nM and 3.13 nM diluted in HBS-EP running buffer (GE Healthcare). Flow rate was 30 to 35 µl/min for 3 min, then HBS-EP running buffer was applied for 8 min again at a flow rate of 30 to 35 µl/ml. Regeneration of the chip was performed using
 30 10 mM glycine 0.5 M NaCl pH 2.45. Data sets were analyzed using BiaEval Software (see Figure 4). In general two independent experiments were performed.

5.2 Binding affinity to human and macaque BCMA

Binding affinities of BCMA/CD3 bispecific antibodies to human and macaque BCMA were determined by Biacore analysis using recombinant BCMA fusion proteins with mouse albumin (ALB).

- 5 In detail, CM5 Sensor Chips (GE Healthcare) were immobilized with approximately 150 to 200 RU of the respective recombinant antigen using acetate buffer pH4.5 according to the manufacturer's manual. The bispecific antibody samples were loaded in five concentrations: 50 nM, 25 nM, 12.5 nM, 6.25 nM and 3.13 nM diluted in HBS-EP running buffer (GE Healthcare). For BCMA affinity determinations the flow rate was 35 μ l/min for 3 min, then HBS-EP running
10 buffer was applied for 10, 30 or 60 min again at a flow rate of 35 μ l/ml. Regeneration of the chip was performed using a buffer consisting of a 1:1 mixture of 10 mM glycine 0.5 M NaCl pH 1.5 and 6 M guanidine chloride solution. Data sets were analyzed using BiaEval Software (see Figure 6). In general two independent experiments were performed.
- 15 Confirmative human and macaque CD3 epsilon binding was performed in single experiments using the same concentrations as applied for BCMA binding; off-rate determination was done for 10 min dissociation time.

All BCMA/CD3 bispecific antibodies of epitope cluster E3 showed high affinities to human
20 BCMA in the sub-nanomolar range down to 1-digit picomolar range. Binding to macaque BCMA was balanced, also showing affinities in the 1-digit nanomolar down to subnanomolar range. Affinities and affinity gaps of BCMA/CD3 bispecific antibodies are shown in Table 2.

Table 2: Affinities of BCMA/CD3 bispecific antibodies of the epitope cluster E3 to human and
25 macaque BCMA as determined by Biacore analysis, and calculated affinity gaps (ma BCMA : hu BCMA).

<i>BCMA/CD3 antibody</i>	<i>bispecific</i>	<i>hu BCMA [nM]</i>	<i>ma BCMA [nM]</i>	<i>Affinity gap ma BCMA : hu BCMA</i>
BCMA-83		0.031	0.077	2.5
BCMA-98		0.025	0.087	3.5
BCMA-71		0.60	2.2	3.7
BCMA-34		0.051	0.047	1 : 1.1
BCMA-74		0.088	0.12	1.4

BCMA-20	0.0085	0.016	1.9
---------	--------	-------	-----

5.3 Biacore-based determination of the bispecific antibody affinity to human and macaque BCMA

5 The affinities of BCMA/CD3 bispecific antibodies to recombinant soluble BCMA on CM5 chips in Biacore measurements were repeated to reconfirm KDs and especially off-rates using longer dissociation periods (60 min instead of 10 min as used in the previous experiment). All of the tested BCMA/CD3 bispecific antibodies underwent two independent affinity measurements with five different concentrations each.

10 The affinities of the BCMA/CD3 bispecific antibodies of the epitope cluster E3 were clearly subnanomolar down to 1-digit picomolar, see examples in Table 3.

15 **Table 3:** Affinities (KD) of BCMA/CD3 bispecific antibodies of the epitope cluster E3 from Biacore experiments using extended dissociation times (two independent experiments each).

<i>BCMA/CD3 bispecific antibody</i>	<i>KD [nM] human BCMA</i>	<i>KD [nM] macaque BCMA</i>
BCMA-83	0.053 ± 0.017	0.062 ± 0.011
BCMA-98	0.025 ± 0.003	0.060 ± 0.001
BCMA-71	0.242 ± 0.007	0.720 ± 0.028
BCMA-34	0.089 ± 0.019	0.056 ± 0.003
BCMA-74	0.076 ± 0.002	0.134 ± 0.010
BCMA-20	0.0095 ± 0.0050	0.0060 ± 0.0038

Example 6

20 Bispecific binding and interspecies cross-reactivity

For confirmation of binding to human and macaque BCMA and CD3, bispecific antibodies were tested by flow cytometry using CHO cells transfected with human and macaque BCMA, respectively, the human multiple myeloma cell line NCI-H929 expressing native human BCMA,

CD3-expressing human T cell leukemia cell line HPB-ALL (DSMZ, Braunschweig, ACC483) and the CD3-expressing macaque T cell line 4119LnPx (Knappe A, et al., Blood, 2000, 95, 3256-3261). Moreover, untransfected CHO cells were used as negative control.

- 5 For flow cytometry 200,000 cells of the respective cell lines were incubated for 30 min on ice with 50 µl of purified bispecific antibody at a concentration of 5 µg/ml. The cells were washed twice in PBS/2% FCS and binding of the constructs was detected with a murine PentaHis antibody (Qiagen; diluted 1:20 in 50 µl PBS/2% FCS). After washing, bound PentaHis antibodies were detected with an Fc gamma-specific antibody (Dianova) conjugated to
10 phycoerythrin, diluted 1:100 in PBS/2% FCS. Samples were measured by flow cytometry on a FACSCanto II instrument and analyzed by FACSDiva software (both from Becton Dickinson).

The BCMA/CD3 bispecific antibodies of epitope cluster E3 stained CHO cells transfected with human and macaque BCMA, the human BCMA-expressing multiple myeloma cell line NCI-
15 H929 as well as human and macaque T cells. Moreover, there was no staining of untransfected CHO cells (see Figure 7).

Example 7

Scatchard-based determination of bispecific-antibody affinity to human and macaque 20 BCMA

For Scatchard analysis, saturation binding experiments are performed using a monovalent detection system developed at Micromet (anti-His Fab/Alexa 488) to precisely determine monovalent binding of the bispecific antibodies to the respective cell line.

2 x 10⁴ cells of the respective cell line (recombinantly human BCMA-expressing CHO cell line,
25 recombinantly macaque BCMA-expressing CHO cell line) are incubated with each 50 µl of a triplet dilution series (eight dilutions at 1:2) of the respective BCMA bispecific antibody starting at 100 nM followed by 16 h incubation at 4°C under agitation and one residual washing step. Then, the cells are incubated for further 30 min with 30 µl of an anti-His Fab/Alexa488 solution (Micromet; 30 µg/ml). After one washing step, the cells are resuspended in 150 µl FACS buffer
30 containing 3.5 % formaldehyde, incubated for further 15 min, centrifuged, resuspended in FACS buffer and analyzed using a FACS Cantoll machine and FACS Diva software. Data are generated from two independent sets of experiments. Values are plotted as hyperbole binding curves. Respective Scatchard analysis is calculated to extrapolate maximal binding (Bmax). The concentrations of bispecific antibodies at half-maximal binding are determined reflecting the

respective KDs. Values of triplicate measurements are plotted as hyperbolic curves. Maximal binding is determined using Scatchard evaluation and the respective KDs are calculated.

The affinities of BCMA/CD3 bispecific antibodies to CHO cells transfected with human or macaque BCMA were determined by Scatchard analysis as the most reliable method for measuring potential affinity gaps between human and macaque BCMA.

Cells expressing the BCMA antigen were incubated with increasing concentrations of the respective monomeric BCMA/CD3 bispecific antibody until saturation was reached (16 h). Bound bispecific antibody was detected by flow cytometry. The concentrations of BCMA/CD3 bispecific antibodies at half-maximal binding were determined reflecting the respective KDs.

Values of triplicate measurements were plotted as hyperbolic curves and as S-shaped curves to demonstrate proper concentration ranges from minimal to optimal binding. Maximal binding (Bmax) was determined (Figure 8) using Scatchard evaluation and the respective KDs were calculated. Values depicted in Table 4 were derived from two independent experiments per BCMA/CD3 bispecific antibody.

Cell based Scatchard analysis confirmed that the BCMA/CD3 bispecific antibodies of the epitope cluster E3 are subnanomolar in affinity to human BCMA and present with a small interspecies BCMA affinity gap of below five.

Table 4: Affinities (KD) of BCMA/CD3 bispecific antibodies of the epitope cluster E3 from cell based Scatchard analysis (two independent experiments each) with the calculated affinity gap KD macaque BCMA / KD human BCMA.

<i>BCMA/CD3 bispecific antibody</i>	<i>KD [nM] human BCMA</i>	<i>KD [nM] macaque BCMA</i>	<i>x-fold KD difference KD ma vs. KD hu BCMA</i>
BCMA-83	0.40 ± 0.13	1.22 ± 0.25	3.1
BCMA-98	0.74 ± 0.02	1.15 ± 0.64	1.6
BCMA-71	0.78 ± 0.07	3.12 ± 0.26	4.0
BCMA-34	0.77 ± 0.11	0.97 ± 0.33	1.3
BCMA-74	0.67 ± 0.03	0.95 ± 0.06	1.4
BCMA-20	0.78 ± 0.10	0.85 ± 0.01	1.1

Example 8**Cytotoxic activity****8.1 Chromium release assay with stimulated human T cells**

5 Stimulated T cells enriched for CD8⁺ T cells were obtained as described below.

A petri dish (145 mm diameter, Greiner bio-one GmbH, Kremsmünster) was coated with a commercially available anti-CD3 specific antibody (OKT3, Orthoclone) in a final concentration of 1 µg/ml for 1 hour at 37°C. Unbound protein was removed by one washing step with PBS. 3 – 5 x 10⁷ human PBMC were added to the precoated petri dish in 120 ml of RPMI 1640 with stabilized glutamine / 10% FCS / IL-2 20 U/ml (Proleukin®, Chiron) and stimulated for 2 days. On the third day, the cells were collected and washed once with RPMI 1640. IL-2 was added to a final concentration of 20 U/ml and the cells were cultured again for one day in the same cell culture medium as above.

15 CD8⁺ cytotoxic T lymphocytes (CTLs) were enriched by depletion of CD4⁺ T cells and CD56⁺ NK cells using Dynal-Beads according to the manufacturer's protocol.

Macaque or human BCMA-transfected CHO target cells were washed twice with PBS and labeled with 11.1 MBq ⁵¹Cr in a final volume of 100 µl RPMI with 50% FCS for 60 minutes at 37°C. Subsequently, the labeled target cells were washed 3 times with 5 ml RPMI and then used in the cytotoxicity assay. The assay was performed in a 96-well plate in a total volume of 200 µl supplemented RPMI with an E:T ratio of 10:1. A starting concentration of 0.01 – 1 µg/ml of purified bispecific antibody and threefold dilutions thereof were used. Incubation time for the assay was 18 hours. Cytotoxicity was determined as relative values of released chromium in the supernatant relative to the difference of maximum lysis (addition of Triton-X) and spontaneous lysis (without effector cells). All measurements were carried out in quadruplicates. Measurement of chromium activity in the supernatants was performed in a Wizard 3" gamma counter (Perkin Elmer Life Sciences GmbH, Köln, Germany). Analysis of the results was carried out with Prism 5 for Windows (version 5.0, GraphPad Software Inc., San Diego, California, USA). EC50 values calculated by the analysis program from the sigmoidal dose response curves were used for comparison of cytotoxic activity (see Figure 5).

30 **8.2 Potency of redirecting stimulated human effector T cells against human BCMA-transfected CHO cells**

The cytotoxic activity of BCMA/CD3 bispecific antibodies was analyzed in a 51-chromium (⁵¹Cr) release cytotoxicity assay using CHO cells transfected with human BCMA as target cells, and

stimulated enriched human CD8 T cells as effector cells. The experiment was carried out as described in Example 8.1.

All BCMA/CD3 bispecific antibodies of epitope cluster E3 showed very potent cytotoxic activity against human BCMA transfected CHO cells with EC50-values in the 1-digit pg/ml range or even below (Figure 9 and Table 5). So the epitope cluster E3 presents with a very favorable epitope-activity relationship supporting very potent bispecific antibody mediated cytotoxic activity.

Table 5: EC50 values [pg/ml] of BCMA/CD3 bispecific antibodies of the epitope cluster E3 analyzed in a 51-chromium (⁵¹Cr) release cytotoxicity assay using CHO cells transfected with human BCMA as target cells, and stimulated enriched human CD8 T cells as effector cells.

<i>BCMA/CD3 bispecific antibody</i>	<i>EC50 [pg/ml]</i>	<i>R square value</i>
BCMA-83	0.38	0.79
BCMA-98	0.27	0,85
BCMA-71	3.2	0.85
BCMA-34	3.4	0.81
BCMA-74	0.73	0.80
BCMA-20	0.83	0.82

8.3 FACS-based cytotoxicity assay with unstimulated human PBMC

Isolation of effector cells

Human peripheral blood mononuclear cells (PBMC) were prepared by Ficoll density gradient centrifugation from enriched lymphocyte preparations (buffy coats), a side product of blood banks collecting blood for transfusions. Buffy coats were supplied by a local blood bank and PBMC were prepared on the same day of blood collection. After Ficoll density centrifugation and extensive washes with Dulbecco's PBS (Gibco), remaining erythrocytes were removed from PBMC via incubation with erythrocyte lysis buffer (155 mM NH₄Cl, 10 mM KHCO₃, 100 μM EDTA). Platelets were removed via the supernatant upon centrifugation of PBMC at 100 x g. Remaining lymphocytes mainly encompass B and T lymphocytes, NK cells and monocytes. PBMC were kept in culture at 37°C/5% CO₂ in RPMI medium (Gibco) with 10% FCS (Gibco).

Depletion of CD14⁺ and CD56⁺ cells

For depletion of CD14⁺ cells, human CD14 MicroBeads (Milty Biotec, MACS, #130-050-201) were used, for depletion of NK cells human CD56 MicroBeads (MACS, #130-050-401). PBMC were counted and centrifuged for 10 min at room temperature with 300 x g. The supernatant was discarded and the cell pellet resuspended in MACS isolation buffer [80 µL/ 10⁷ cells; PBS (Invitrogen, #20012-043), 0.5% (v/v) FBS (Gibco, #10270-106), 2 mM EDTA (Sigma-Aldrich, #E-6511)]. CD14 MicroBeads and CD56 MicroBeads (20 µL/10⁷ cells) were added and incubated for 15 min at 4 - 8°C. The cells were washed with MACS isolation buffer (1 - 2 mL/10⁷ cells). After centrifugation (see above), supernatant was discarded and cells resuspended in MACS isolation buffer (500 µL/10⁸ cells). CD14/CD56 negative cells were then isolated using LS Columns (Milty Biotec, #130-042-401). PBMC w/o CD14⁺/CD56⁺ cells were cultured in RPMI complete medium i.e. RPMI1640 (Biochrom AG, #FG1215) supplemented with 10% FBS (Biochrom AG, #S0115), 1x non-essential amino acids (Biochrom AG, #K0293), 10 mM Hepes buffer (Biochrom AG, #L1613), 1 mM sodium pyruvate (Biochrom AG, #L0473) and 100 U/mL penicillin/streptomycin (Biochrom AG, #A2213) at 37°C in an incubator until needed.

Target cell labeling

For the analysis of cell lysis in flow cytometry assays, the fluorescent membrane dye DiOC₁₈ (DiO) (Molecular Probes, #V22886) was used to label human BCMA- or macaque BCMA-transfected CHO cells as target cells and distinguish them from effector cells. Briefly, cells were harvested, washed once with PBS and adjusted to 10⁶ cell/mL in PBS containing 2 % (v/v) FBS and the membrane dye DiO (5 µL/10⁶ cells). After incubation for 3 min at 37°C, cells were washed twice in complete RPMI medium and the cell number adjusted to 1.25 x 10⁵ cells/mL. The vitality of cells was determined using 0.5 % (v/v) isotonic EosinG solution (Roth, #45380).

Flow cytometry based analysis

This assay was designed to quantify the lysis of macaque or human BCMA-transfected CHO cells in the presence of serial dilutions of BCMA bispecific antibodies.

Equal volumes of DiO-labeled target cells and effector cells (i.e., PBMC w/o CD14⁺ cells) were mixed, resulting in an E:T cell ratio of 10:1. 160 µL of this suspension were transferred to each well of a 96-well plate. 40 µL of serial dilutions of the BCMA bispecific antibodies and a negative control bispecific (an CD3-based bispecific antibody recognizing an irrelevant target antigen) or RPMI complete medium as an additional negative control were added. The bispecific antibody-mediated cytotoxic reaction proceeded for 48 hours in a 7% CO₂ humidified incubator. Then cells were transferred to a new 96-well plate and loss of target cell membrane integrity was monitored by adding propidium iodide (PI) at a final concentration of 1 µg/mL. PI is a membrane impermeable dye that normally is excluded from viable cells, whereas dead cells take it up and become identifiable by fluorescent emission.

Samples were measured by flow cytometry on a FACSCanto II instrument and analyzed by FACSDiva software (both from Becton Dickinson).

Target cells were identified as DiO-positive cells. PI-negative target cells were classified as living target cells. Percentage of cytotoxicity was calculated according to the following formula:

$$\text{Cytotoxicity [\%]} = \frac{n_{\text{dead target cells}}}{n_{\text{target cells}}} \times 100$$

n = number of events

Using GraphPad Prism 5 software (Graph Pad Software, San Diego), the percentage of cytotoxicity was plotted against the corresponding bispecific antibody concentrations. Dose response curves were analyzed with the four parametric logistic regression models for evaluation of sigmoid dose response curves with fixed hill slope and EC50 values were calculated.

8.4 Unstimulated human PBMC against human BCMA-transfected target cells

The cytotoxic activity of BCMA/CD3 bispecific antibodies was analyzed in a FACS-based cytotoxicity assay using CHO cells transfected with human BCMA as target cells, and unstimulated human PBMC as effector cells. The assay was carried out as described above (Example 8.3).

The results of the FACS-based cytotoxicity assays with unstimulated human PBMC as effector cells and human BCMA-transfected CHO cells as targets are shown in Figure 10 and Table 6.

Table 6: EC50 values [pg/ml] of BCMA/CD3 bispecific antibodies of epitope cluster E3 as measured in a 48-hour FACS-based cytotoxicity assay with unstimulated human PBMC as effector cells and CHO cells transfected with human BCMA as target cells.

<i>BCMA/CD3 bispecific antibody</i>	<i>EC50 [pg/ml]</i>	<i>R square value</i>
BCMA-83	212	0.97
BCMA-7	102	0.97
BCMA-5	58.4	0.94
BCMA-98	53.4	0.95
BCMA-71	208	0.94

BCMA-34	149	0.94
BCMA-74	125	0.97
BCMA-20	176	0.98

Example 9

9.1 Exclusion of cross-reactivity with BAFF-receptor

5 For flow cytometry, 200,000 cells of the respective cell lines were incubated for 30 min on ice with 50 μ l of purified bispecific molecules at a concentration of 5 μ g/ml. The cells were washed twice in PBS with 2% FCS and binding of the constructs was detected with a murine PentaHis antibody (Qiagen; diluted 1:20 in 50 μ l PBS with 2% FCS). After washing, bound PentaHis antibodies were detected with an Fc gamma-specific antibody (Dianova) conjugated to
10 phycoerythrin, diluted 1:100 in PBS with 2% FCS. Samples were measured by flow cytometry on a FACSCanto II instrument and analyzed by FACSDiva software (both from Becton Dickinson). The bispecific binders were shown to not be cross-reactive with BAFF receptor.

9.2 Exclusion of BCMA/CD3 bispecific antibody cross-reactivity with human BAFF-receptor (BAFF-R) and TACI

For exclusion of binding to human BAFF-R and TACI, BCMA/CD3 bispecific antibodies were tested by flow cytometry using CHO cells transfected with human BAFF-R and TACI, respectively. Moreover, L363 multiple myeloma cells were used as positive control for binding to
20 human BCMA. Expression of BAFF-R and TACI antigen on CHO cells was confirmed by two positive control antibodies. Flow cytometry was performed as described in the previous example.

Flow cytometric analysis confirmed that none of the BCMA/CD3 bispecific antibodies of the
25 epitope cluster E3 cross-reacts with human BAFF-R or human TACI (see Figure 11).

Example 10

Cytotoxic activity

The potency of human-like BCMA bispecific antibodies in redirecting effector T cells against
30 BCMA-expressing target cells is analyzed in five additional in vitro cytotoxicity assays:

1. The potency of BCMA bispecific antibodies in redirecting stimulated human effector T cells against a BCMA-positive (human) tumor cell line is measured in a 51-chromium release assay.

2. The potency of BCMA bispecific antibodies in redirecting the T cells in unstimulated human PBMC against human BCMA-transfected CHO cells is measured in a FACS-based cytotoxicity assay.

3. The potency of BCMA bispecific antibodies in redirecting the T cells in unstimulated human PBMC against a BCMA-positive (human) tumor cell line is measured in a FACS-based cytotoxicity assay.

4. For confirmation that the cross-reactive BCMA bispecific antibodies are capable of redirecting macaque T cells against macaque BCMA-transfected CHO cells, a FACS-based cytotoxicity assay is performed with a macaque T cell line as effector T cells.

5. The potency gap between monomeric and dimeric forms of BCMA bispecific antibodies is determined in a 51-chromium release assay using human BCMA-transfected CHO cells as target cells and stimulated human T cells as effector cells.

Example 11

Stimulated human T cells against the BCMA-positive human multiple myeloma cell line L363

The cytotoxic activity of BCMA/CD3 bispecific antibodies was analyzed in a 51-chromium (⁵¹Cr) release cytotoxicity assay using the BCMA-positive human multiple myeloma cell line L363 (DSMZ No. ACC49) as source of target cells, and stimulated enriched human CD8 T cells as effector cells. The assay was carried out as described in Example 8.1.

In accordance with the results of the 51-chromium release assays with stimulated enriched human CD8 T lymphocytes as effector cells and human BCMA-transfected CHO cells as targets, BCMA/CD3 bispecific antibodies of epitope cluster E3 are very potent in cytotoxic activity (Figure 12 and Table 7).

Another group of antibodies was identified during epitope clustering (see Examples 1 and 3), which is capable of binding to epitope clusters 1 and 4 of BCMA ("E1/E4"). Unexpectedly, BCMA/CD3 bispecific antibodies of epitope cluster E1/E4 – although potent in cytotoxic activity against CHO cell transfected with human BCMA – proved to be rather weakly cytotoxic against the human multiple myeloma cell line L363 expressing native BCMA at low density on the cell

surface (Figure 12 and Table 7). Without wishing to be bound by theory, the inventors believe that the E1/E4 epitope of human BCMA might be less well accessible on natural BCMA expressers than on BCMA-transfected cells.

- 5 **Table 7:** EC50 values [pg/ml] of BCMA/CD3 bispecific antibodies of epitope clusters E1/E4 (rows 1 and 2) and E3 (rows 3 to 8) analyzed in an 18-hour 51-chromium (⁵¹Cr) release cytotoxicity assay with the BCMA-positive human multiple myeloma cell line L363 as source of target cells, and stimulated enriched human CD8 T cells as effector cells.

	<i>BCMA/CD3 bispecific antibody</i>	<i>EC50 [pg/ml]</i>	<i>R square value</i>
1	BCMA-54	685	0.84
2	BCMA-53	1107	0.82
3	BCMA-83	28	0.83
4	BCMA-98	10	0.81
5	BCMA-71	125	0.86
6	BCMA-34	42	0.81
7	BCMA-74	73	0.79
8	BCMA-20	21	0.85

10

Example 12

Unstimulated human PBMC against the BCMA-positive human multiple myeloma cell line L363

15

The cytotoxic activity of BCMA/CD3 bispecific antibodies was furthermore analyzed in a FACS-based cytotoxicity assay using the BCMA-positive human multiple myeloma cell line L363 (DSMZ, ACC49) - showing the weakest surface expression of native BCMA of all tested target T cell lines - as source of target cells and unstimulated human PBMC as effector cells. The assay was carried out as described above (Example 8.3).

20

As observed in the 51-chromium release assay with stimulated enriched human CD8 T lymphocytes against the human multiple myeloma cell line L363, the BCMA/CD3 bispecific antibodies of epitope cluster E1/E4 – in contrast to their potent cytotoxic activity against CHO cell transfected with human BCMA – proved to be again less potent in redirecting the cytotoxic activity of unstimulated PBMC against the human multiple myeloma cell line L363 expressing

25

native BCMA at low density on the cell surface. This is in line with the theory provided hereinabove, i.e., the E1/E4 epitope of human BCMA may be less well accessible on natural BCMA expressers than on BCMA-transfected cells. BCMA/CD3 bispecific antibodies of the epitope cluster E3 presented with 3-digit pg/ml EC50-values in this assay (see Figure 13 and Table 8).

Table 8: EC50 values [pg/ml] of BCMA/CD3 bispecific antibodies of epitope clusters E1/E4 (rows 1 and 2) and E3 (rows 3 to 8) as measured in a 48-hour FACS-based cytotoxicity assay with unstimulated human PBMC as effector cells and the human multiple myeloma cell line L363 as source of target cells.

	<i>BCMA/CD3 bispecific antibody</i>	<i>EC50 [pg/ml]</i>	<i>R square value</i>
1	BCMA-54	3162	0.99
2	BCMA-53	2284	0.98
3	BCMA-83	241	0.99
4	BCMA-98	311	0.99
5	BCMA-71	284	0.99
6	BCMA-34	194	0.99
7	BCMA-74	185	0.99
8	BCMA-20	191	0.99

Expectedly, EC50-values were higher in cytotoxicity assays with unstimulated PBMC as effector cells than in cytotoxicity assays using enriched stimulated human CD8 T cells.

Example 13

Unstimulated human PBMC against the BCMA-positive human multiple myeloma cell line NCI-H929

The cytotoxic activity of BCMA/CD3 bispecific antibodies was analyzed in a FACS-based cytotoxicity assay using the BCMA-positive human multiple myeloma cell line NCI-H929 (ATCC CRL-9068) as source of target cells and unstimulated human PBMC as effector cells. The assay was carried out as described above (Example 8.3).

The results of this assay with another human multiple myeloma cell line (i.e. NCI-H929) expressing native BCMA on the cell surface confirm those obtained with human multiple myeloma cell line L363. Again, BCMA/CD3 bispecific antibodies of epitope cluster E1/E4 – in contrast to their potent cytotoxic activity against CHO cell transfected with human BCMA – proved to be less potent in redirecting the cytotoxic activity of unstimulated PBMC against human multiple myeloma cells confirming the theory that the E1/E4 epitope of human BCMA may be less well accessible on natural BCMA expressers than on BCMA-transfected cells. Such an activity gap between BCMA-transfected target cells and natural expressers as seen for the E1/E4 binders was not found for the E3. BCMA/CD3 bispecific antibodies of the epitope cluster E3 presented with 2- to 3-digit pg/ml EC50-values and hence redirected unstimulated PBMC against NCI-H929 target cells with very good EC50-values (see Figure 14 and Table 9).

Table 9: EC50 values [pg/ml] of BCMA/CD3 bispecific antibodies of epitope clusters E1/E4 (rows 1 and 2) and E3 (rows 3 to 8) as measured in a 48-hour FACS-based cytotoxicity assay with unstimulated human PBMC as effector cells and the human multiple myeloma cell line NCI-H929 as source of target cells.

	<i>BCMA/CD3 bispecific antibody</i>	<i>EC50 [pg/ml]</i>	<i>R square value</i>
1	BCMA-54	2604	0.99
2	BCMA-53	2474	0.99
3	BCMA-83	154	0.93
4	BCMA-98	67.6	0.87
5	BCMA-71	50.7	0.96
6	BCMA-34	227	0.99
7	BCMA-74	103	0.97
8	BCMA-20	123	0.97

As expected, EC50-values were lower with the human multiple myeloma cell line NCI-H929, which expresses higher levels of BCMA on the cell surface compared to L363.

Example 14

Macaque T cells against macaque BCMA-expressing target cells

Finally, the cytotoxic activity of BCMA/CD3 bispecific antibodies was analyzed in a FACS-based cytotoxicity assay using CHO cells transfected with macaque BCMA as target cells, and a macaque T cell line as source of effector cells.

- 5 The macaque T cell line 4119LnPx (Knappe et al. Blood 95:3256-61 (2000)) was used as source of effector cells. Target cell labeling of macaque BCMA-transfected CHO cells and flow cytometry based analysis of cytotoxic activity was performed as described above.

10 Macaque T cells from cell line 4119LnPx were induced to efficiently kill macaque BCMA-transfected CHO cells by BCMA/CD3 bispecific antibodies of the E3 epitope cluster. The antibodies presented very potently with 1-digit to low 2-digit pg/ml EC50-values in this assay, confirming that these antibodies are very active in the macaque system. On the other hand, BCMA/CD3 bispecific antibodies of the epitope cluster E1/E4 showed a significantly weaker potency with EC50-values in the 2-digit to 3-digit pg/ml range (see Figure 15 and Table 10). The
15 E3 specific antibodies are hence about 3 to almost 100 times more potent in the macaque system.

Table 10: EC50 values [pg/ml] of BCMA/CD3 bispecific antibodies of epitope clusters E1/E4 (rows 1 and 2) and E3 (rows 3 to 8) as measured in a 48-hour FACS-based cytotoxicity assay
20 with macaque T cell line 4119LnPx as effector cells and CHO cells transfected with macaque BCMA as target cells.

	<i>BCMA/CD3 bispecific antibody</i>	<i>EC50 [pg/ml]</i>	<i>R square value</i>
1	BCMA-54	78.5	0.98
2	BCMA-53	183	0.96
3	BCMA-83	10.9	0.97
4	BCMA-98	2.5	0.89
5	BCMA-71	3.2	0.97
6	BCMA-34	2.1	0.95
7	BCMA-74	2.0	0.95
8	BCMA-20	26	0.98

25 **Example 15**

Potency gap between BCMA/CD3 bispecific antibody monomer and dimer

In order to determine the difference in cytotoxic activity between the monomeric and the dimeric isoform of individual BCMA/CD3 bispecific antibodies (referred to as potency gap), a 51-chromium release cytotoxicity assay as described hereinabove (Example 8.1) was carried out with purified BCMA/CD3 bispecific antibody monomer and dimer. The potency gap was calculated as ratio between EC50 values of the bispecific antibody's monomer and dimer. Potency gaps of the tested BCMA/CD3 bispecific antibodies of the epitope cluster E3 were between 0.03 and 1.2. There is hence no substantially more active dimer compared to its respective monomer.

Example 16

Monomer to dimer conversion after three freeze/thaw cycles

Bispecific BCMA/CD3 antibody monomer were subjected to three freeze/thaw cycles followed by high performance SEC to determine the percentage of initially monomeric antibody, which had been converted into antibody dimer.

15 µg of monomeric antibody were adjusted to a concentration of 250 µg/ml with generic buffer and then frozen at -80°C for 30 min followed by thawing for 30 min at room temperature. After three freeze/thaw cycles the dimer content was determined by HP-SEC. To this end, 15 µg aliquots of the monomeric isoforms of the antibodies were thawed and equalized to a concentration of 250 µg/ml in the original SEC buffer (10 mM citric acid – 75 mM lysine HCl – 4% trehalose - pH 7.2) followed by incubation at 37°C for 7 days. A high resolution SEC Column TSK Gel G3000 SWXL (Tosoh, Tokyo-Japan) was connected to an Äkta Purifier 10 FPLC (GE Lifesciences) equipped with an A905 Autosampler. Column equilibration and running buffer consisted of 100 mM KH₂PO₄ – 200 mM Na₂SO₄ adjusted to pH 6.6. After 7 days of incubation, the antibody solution (15 µg protein) was applied to the equilibrated column and elution was carried out at a flow rate of 0.75 ml/min at a maximum pressure of 7 MPa. The whole run was monitored at 280, 254 and 210 nm optical absorbance. Analysis was done by peak integration of the 210 nm signal recorded in the Äkta Unicorn software run evaluation sheet. Dimer content was calculated by dividing the area of the dimer peak by the total area of monomer plus dimer peak.

The BCMA/CD3 bispecific antibodies of the epitope cluster E3 presented with dimer percentages of 0.7 to 1.1% after three freeze/thaw cycles, which is considered good. However,

the dimer conversion rates of BCMA/CD3 bispecific antibodies of the epitope cluster E1/E4 reached unfavorably high values, exceeding the threshold to disadvantageous dimer values of $\geq 2.5\%$ (4.7% and 3.8%, respectively), see Table 11.

- 5 **Table 11:** Percentage of monomeric versus dimeric BCMA/CD3 bispecific antibodies of epitope clusters E1/E4 (rows 1 and 2) and E3 (rows 3 to 8) after three freeze/thaw cycles as determined by High Performance Size Exclusion Chromatography (HP-SEC).

	<i>BCMA/CD3 bispecific antibody</i>	<i>Monomer [%]</i>	<i>Dimer [%]</i>
1	BCMA-54	95.3	4.7
2	BCMA-53	96.2	3.8
3	BCMA-83	99.1	0.9
4	BCMA-98	99.1	0.9
5	BCMA-71	99.1	0.9
6	BCMA-34	98.9	1.1
7	BCMA-74	99.3	0.7
8	BCMA-20	99.2	0.8

10

Example 17

Thermostability

15 Temperature melting curves were determined by Differential Scanning Calorimetry (DSC) to determine intrinsic biophysical protein stabilities of the BCMA/CD3 bispecific antibodies. These experiments were performed using a MicroCal LLC (Northampton, MA, U.S.A) VP-DSC device. The energy uptake of a sample containing BCMA/CD3 bispecific antibody was recorded from 20 to 90°C compared to a sample which just contained the antibody's formulation buffer.

20 In detail, BCMA/CD3 bispecific antibodies were adjusted to a final concentration of 250 µg/ml in storage buffer. 300 µl of the prepared protein solutions were transferred into a deep well plate and placed into the cooled autosampler rack position of the DSC device. Additional wells were filled with the SEC running buffer as reference material for the measurement. For the measurement process the protein solution was transferred by the autosampler into a capillary.

25 An additional capillary was filled with the SEC running buffer as reference. Heating and

recording of required heating energy to heat up both capillaries at equal temperature ranging from 20 to 90°C was done for all samples.

For recording of the respective melting curve, the overall sample temperature was increased stepwise. At each temperature T energy uptake of the sample and the formulation buffer reference was recorded. The difference in energy uptake C_p (kcal/mole/°C) of the sample minus the reference was plotted against the respective temperature. The melting temperature is defined as the temperature at the first maximum of energy uptake.

All tested BCMA/CD3 bispecific antibodies of the epitope cluster E3 showed favorable thermostability with melting temperatures above 60°C, more precisely between 61.62°C and 63.05°C.

Example 18

Exclusion of plasma interference by flow cytometry

To determine potential interaction of BCMA/CD3 bispecific antibodies with human plasma proteins, a plasma interference test was established. To this end, 10 µg/ml of the respective BCMA/CD3 bispecific antibodies were incubated for one hour at 37°C in 90 % human plasma. Subsequently, the binding to human BCMA expressing CHO cells was determined by flow cytometry.

For flow cytometry, 200,000 cells of the respective cell lines were incubated for 30 min on ice with 50 µl of purified antibody at a concentration of 5 µg/ml. The cells were washed twice in PBS/2% FCS and binding of the constructs was detected with a murine PentaHis antibody (Qiagen; diluted 1:20 in 50 µl PBS/2% FCS). After washing, bound PentaHis antibodies were detected with an Fc gamma-specific antibody (Dianova) conjugated to phycoerythrin, diluted 1:100 in PBS/2% FCS. Samples were measured by flow cytometry on a FACSCanto II instrument and analyzed by FACSDiva software (both from Becton Dickinson).

The obtained data were compared with a control assay using PBS instead of human plasma. Relative binding was calculated as follows:
(signal PBS sample / signal w/o detection agent) / (signal plasma sample / signal w/o detection agent).

In this experiment it became obvious that there was no significant reduction of target binding of the respective BCMA/CD3 bispecific antibodies of the epitope cluster E3 mediated by plasma proteins. The relative plasma interference value was below a value of 2 in all cases, more precisely between 1.29 ± 0.25 and 1.70 ± 0.26 (with a value of "2" being considered as lower threshold for interference signals).

Example 19

Therapeutic efficacy of BCMA/CD3 bispecific antibodies in human tumor xenograft models

On day 1 of the study, 5×10^6 cells of the human cancer cell line NCI-H929 were subcutaneously injected in the right dorsal flank of female NOD/SCID mice.

On day 9, when the mean tumor volume had reached about 100 mm^3 , *in vitro* expanded human CD3^+ T cells were transplanted into the mice by injection of about 2×10^7 cells into the peritoneal cavity of the animals. Mice of vehicle control group 1 (n=5) did not receive effector cells and were used as an untransplanted control for comparison with vehicle control group 2 (n=10, receiving effector cells) to monitor the impact of T cells alone on tumor growth.

The antibody treatment started on day 13, when the mean tumor volume had reached about 200 mm^3 . The mean tumor size of each treatment group on the day of treatment start was not statistically different from any other group (analysis of variance). Mice were treated with 0.5 mg/kg/day of the BCMA/CD3 bispecific antibodies BCMA-98 x CD3 (group 3, n=7) or BCMA-34 x CD3 (group 4, n=6) by intravenous bolus injection for 17 days.

Tumors were measured by caliper during the study and progress evaluated by intergroup comparison of tumor volumes (TV). The tumor growth inhibition T/C [%] was determined by calculating TV as $\text{T/C\%} = 100 \times (\text{median TV of analyzed group}) / (\text{median TV of control group 2})$. The results are shown in Table 12 and Figure 16.

Table 12: Median tumor volume (TV) and tumor growth inhibition (T/C) at days 13 to 30.

Dose group	Data	d13	d14	d15	d16	d18	d19	d21	d23	d26	d28	d30
1 Vehi. control	med.TV [mm³]	238	288	395	425	543	632	863	1067	1116	1396	2023

w/o T cells	T/C [%]	120	123	127	118	104	114	122	113	87	85	110
<u>2</u> Vehicle control	med.TV [mm³]	198	235	310	361	525	553	706	942	1290	1636	1839
	T/C [%]	100	100	100	100	100	100	100	100	100	100	100
<u>3</u> BCMA- 98	med.TV [mm³]	207	243	248	235	164	137	93.5	46.2	21.2	0.0	0.0
	T/C [%]	105	104	79.7	65.0	31.2	24.7	13.2	4.9	1.6	0.0	0.0
<u>4</u> BCMA- 34	med.TV [mm³]	206	233	212	189	154	119	56.5	17.4	0.0	0.0	0.0
	T/C [%]	104	99.2	68.2	52.3	29.4	21.5	8.0	1.8	0.0	0.0	0.0

Example 20**Exclusion of lysis of target negative cells**

5

An *in vitro* lysis assay was carried out using the BCMA-positive human multiple myeloma cell line NCI-H929 and purified T cells at an effector to target cell ratio of 5:1 and with an incubation time of 24 hours. BCMA/CD3 bispecific antibodies of epitope cluster E3 (BCMA-34 and BCMA-98) showed high potency and efficacy in the lysis of NCI-H929. However, no lysis was detected in the BCMA negative cell lines HL60 (AML / myeloblast morphology), MES-SA (uterus sarcoma, fibroblast morphology), and SNU-16 (stomach carcinoma, epithelial morphology) for up to 500 nM of the respective antibody.

10

Example 21**15 Induction of T cell activation of different PBMC subsets**

A FACS-based cytotoxicity assay (48h; E:T = 10:1) was carried out using human multiple myeloma cell lines NCI-H929, L-363 and OPM-2 as target cells and different subsets of human PBMC (CD4⁺ / CD8⁺ / CD25⁺ / CD69⁺) as effector cells. The results (see Table 13) show that the degree of activation, as measured by the EC₅₀ value, is essentially in the same range for the different analyzed PBMC subsets.

20

Table 13: EC₅₀ values [ng/ml] of BCMA/CD3 bispecific antibodies of epitope cluster E3 as measured in a 48-hour FACS-based cytotoxicity assay with different subsets of human PBMC as effector cells and different human multiple myeloma cell lines as target cells.

5

		EC ₅₀ [ng/ml]	
Cell line	PBMC	BCMA-98 x CD3	BCMA-34 x CD3
NCI-H929	CD4 ⁺ / CD25 ⁺	1.46	1.20
	CD8 ⁺ / CD25 ⁺	0.53	0.49
	CD4 ⁺ / CD69 ⁺	0.59	0.47
	CD8 ⁺ / CD69 ⁺	0.21	0.21
OPM-2	CD4 ⁺ / CD25 ⁺	2.52	4.88
	CD8 ⁺ / CD25 ⁺	1.00	1.20
	CD4 ⁺ / CD69 ⁺	1.65	2.27
	CD8 ⁺ / CD69 ⁺	0.48	0.42
L-363	CD4 ⁺ / CD25 ⁺	0.54	0.62
	CD8 ⁺ / CD25 ⁺	0.24	0.28
	CD4 ⁺ / CD69 ⁺	0.35	0.34
	CD8 ⁺ / CD69 ⁺	0.12	0.11

Example 22

Induction of cytokine release

10

A FACS-based cytotoxicity assay (48h; E:T = 10:1) was carried out using human multiple myeloma cell lines NCI-H929, L-363 and OPM-2 as target cells and human PBMC as effector cells. The levels of cytokine release [pg/ml] were determined at increasing concentrations of BCMA/CD3 bispecific antibodies of epitope cluster E3. The following cytokines were analyzed:

15 IL-2, IL-6, IL-10, TNF and IFN-gamma. The results are shown in Table 14 and Figure 17.

Table 14: Release of IL-2, IL-6, IL-10, TNF and IFN-gamma [pg/ml] induced by 2.5 µg/ml of BCMA/CD3 bispecific antibodies of epitope cluster E3 (BCMA-98 and BCMA-34) in a 48-hour FACS-based cytotoxicity assay with human PBMC as effector cells and different human multiple myeloma cell lines as target cells (E:T = 10:1).

5

	Cytokine levels [pg/ml]				
	NCI-H929				
	IL-2	IL-6	IL-10	TNF	IFN-gamma
BCMA-98	1357	699	2798	10828	73910
BCMA-34	1327	631	3439	6675	77042
	OPM-2				
	IL-2	IL-6	IL-10	TNF	IFN-gamma
	IL-2	IL-6	IL-10	TNF	IFN-gamma
BCMA-98	41	118	990	5793	33302
BCMA-34	28	109	801	4913	23214
	L-363				
	IL-2	IL-6	IL-10	TNF	IFN-gamma
	IL-2	IL-6	IL-10	TNF	IFN-gamma
BCMA-98	97	314	2433	5397	64981
BCMA-34	168	347	2080	5930	75681

SEQ ID NO	Designation	Designation	Format / source	Type	Sequence
1	BCMA-1	BC 5G9 91-C7-B10	VH CDR1	aa	NYDMA
2	BCMA-1	BC 5G9 91-C7-B10	VH CDR2	aa	SIITSGDATYYRDSVKG
3	BCMA-1	BC 5G9 91-C7-B10	VH CDR3	aa	HDYYDGSYGFA Y
4	BCMA-1	BC 5G9 91-C7-B10	VL CDR1	aa	KASQSVGINVD
5	BCMA-1	BC 5G9 91-C7-B10	VL CDR2	aa	GASNRHT
6	BCMA-1	BC 5G9 91-C7-B10	VL CDR3	aa	LQYGSIPFT
7	BCMA-1	BC 5G9 91-C7-B10	VH	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPKGLEWVASIITSGDATYYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDY YDGSYGFAYWGQGT LVTVSS
8	BCMA-1	BC 5G9 91-C7-B10	VL	aa	EIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGREFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
9	BCMA-1	BC 5G9 91-C7-B10	scFv	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPKGLEWVASIITSGDATYYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDY YDGSYGFAYWGQGT LVTVSSGGSGSGSGGGGSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGGREFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
10	BCMA-1 HL x CD3 HL	BC 5G9 91-C7-B10 HL x CD3 HL	bispecific molecule	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPKGLEWVASIITSGDATYYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDY YDGSYGFAYWGQGT LVTVSSGGSGSGSGGGGGSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGGREFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWNVRQAPKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVTVSSGGSGSGSGGGGSGT VVTQEPSLT VSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYCYVLWYSNRWVFGGGTKLTVL
11	BCMA-2	BC 5G9 91-C7-D8	VH CDR1	aa	NYDMA

12	BCMA-2	BC 5G9 91- C7-D8	VH CDR2	aa	SIITSGDMTYYRDSVKG
13	BCMA-2	BC 5G9 91- C7-D8	VH CDR3	aa	HDYYDGSYGFAY
14	BCMA-2	BC 5G9 91- C7-D8	VL CDR1	aa	KASQSVGINVD
15	BCMA-2	BC 5G9 91- C7-D8	VL CDR2	aa	GASNRHT
16	BCMA-2	BC 5G9 91- C7-D8	VL CDR3	aa	LQYGSIPFT
17	BCMA-2	BC 5G9 91- C7-D8	VH	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAYWGQGLVTVSSSSGGGGSGGG FTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAYWGQGLVTVSS
18	BCMA-2	BC 5G9 91- C7-D8	VL	aa	EIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGS GREFTLTISLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
19	BCMA-2	BC 5G9 91- C7-D8	scFv	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAYWGQGLVTVSSSSGGGGSGGGGG GSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGREFTLTISLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
20	BCMA-2 HL x CD3 HL	BC 5G9 91- C7-D8 HL x CD3 HL	bispecific molecule	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAYWGQGLVTVSSSSGGGGSGGGGG GSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGREFTLTISLQSEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSSSGGGGSGGGGQT VVTQEP S LTVS PGGT VTLTCGSS TGAVTSGNYPNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWY SNRWVFGGGTKLTVL
21	BCMA-3	BC 5G9 91- E4-B10	VH CDR1	aa	NYDMA
22	BCMA-3	BC 5G9 91- E4-B10	VH CDR2	aa	SIITSGDATYYRDSVKG
23	BCMA-3	BC 5G9 91- E4-B10	VH CDR3	aa	HDYYDGSYGFAY
24	BCMA-3	BC 5G9 91- E4-B10	VL CDR1	aa	KASQSVGINVD

25	BCMA-3	BC 5G9 91-E4-B10	VL CDR2	aa	GASNRHT	
26	BCMA-3	BC 5G9 91-E4-B10	VL CDR3	aa	LQYGSIPFT	
27	BCMA-3	BC 5G9 91-E4-B10	VH	aa	QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDATYYRDSVKGRFTISRDN SKNTLYLQMNSLRSEDTAVYYCVRHDYYDGSYGFA YWGQGTLLVTVSS	
28	BCMA-3	BC 5G9 91-E4-B10	VL	aa	EIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK	
29	BCMA-3	BC 5G9 91-E4-B10	scFv	aa	QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDATYYRDSVKGRFTISRDN SKNTLYLQMNSLRSEDTAVYYCVRHDYYDGSYGFA YWGQGTLLVTVSSGGGGSGGGGGG GSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGGTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK	
30	BCMA-3 HL x CD3 HL	BC 5G9 91-E4-B10 HL x CD3 HL	bispecific molecule	aa	QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDATYYRDSVKGRFTISRDN SKNTLYLQMNSLRSEDTAVYYCVRHDYYDGSYGFA YWGQGTLLVTVSSGGGGSGGGGGG GSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGGTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIKS GGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGGGGQT VVTQEP SLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYVCVLWYSNRWVFGGGTKLTVL	
31	BCMA-4	BC 5G9 91-E4-D8	VH CDR1	aa	NYDMA	
32	BCMA-4	BC 5G9 91-E4-D8	VH CDR2	aa	SIITSGDMTYRDSVKG	
33	BCMA-4	BC 5G9 91-E4-D8	VH CDR3	aa	HDYYDGSYGFA Y	
34	BCMA-4	BC 5G9 91-E4-D8	VL CDR1	aa	KASQSVGINVD	
35	BCMA-4	BC 5G9 91-E4-D8	VL CDR2	aa	GASNRHT	
36	BCMA-4	BC 5G9 91-E4-D8	VL CDR3	aa	LQYGSIPFT	
37	BCMA-4	BC 5G9 91-E4-D8	VH	aa	QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYRDSVKGRFTISRDN SKNTLYLQMNSLRSEDTAVYYCVRHDYYDGSYGFA YWGQGTLLVTVSS	

38	BCMA-4	BC 5G9 91-E4-D8	VL	aa	EIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLTITSSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
39	BCMA-4	BC 5G9 91-E4-D8	scFv	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYYRDSVKGRFTISRDN SKNTLYLQMNSLRSEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVTVSSGGGGSGGGGGGSGEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLTITSSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
40	BCMA-4 HL x CD3 HL	BC 5G9 91-E4-D8 HL x CD3 HL	bispecific molecule	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYYRDSVKGRFTISRDN SKNTLYLQMNSLRSEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVTVSSGGGGSGGGGGGSGEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLTITSSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIKLSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVTVSSGGGGSGGGGQT VVTQEPSLTVSPGGTVTLTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
41	BCMA-5	BC 5G9 91-D2-B10	VH CDR1	aa	NYDMA
42	BCMA-5	BC 5G9 91-D2-B10	VH CDR2	aa	SIITSGDATYYRDSVKG
43	BCMA-5	BC 5G9 91-D2-B10	VH CDR3	aa	HDYYDGSYGFA Y
44	BCMA-5	BC 5G9 91-D2-B10	VL CDR1	aa	KASQSVGINVD
45	BCMA-5	BC 5G9 91-D2-B10	VL CDR2	aa	GASNRHT
46	BCMA-5	BC 5G9 91-D2-B10	VL CDR3	aa	LQYGSIPFT
47	BCMA-5	BC 5G9 91-D2-B10	VH	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDATYYRDSVKGRFTISRDN AKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVTVSS
48	BCMA-5	BC 5G9 91-D2-B10	VL	aa	EIVMTQSPASMSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLTITSSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
49	BCMA-5	BC 5G9 91-D2-B10	scFv	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDATYYRDSVKGRFTISRDN AKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVTVSSGGGGSGGGGGGSGEIVMTQSPASMSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLTITSSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK

50	BCMA-5 HL x CD3 HL	BC 5G9 91-D2-B10 HL x CD3 HL	bispecific molecule	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDATYYRDSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSSGGGGSGGGSGGSGEIVMTQSPASMSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGGTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGSGGQTVVTQEPSLTVS PGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
51	BCMA-6	BC 5G9 91-D2-D8	VH CDR1	aa	NYDMA
52	BCMA-6	BC 5G9 91-D2-D8	VH CDR2	aa	SIITSGDMTYRDSVKG
53	BCMA-6	BC 5G9 91-D2-D8	VH CDR3	aa	HDYYDGSYGFAF
54	BCMA-6	BC 5G9 91-D2-D8	VL CDR1	aa	KASQSVGINVD
55	BCMA-6	BC 5G9 91-D2-D8	VL CDR2	aa	GASNRHT
56	BCMA-6	BC 5G9 91-D2-D8	VL CDR3	aa	LQYGSIPFT
57	BCMA-6	BC 5G9 91-D2-D8	VH	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYRDSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSS
58	BCMA-6	BC 5G9 91-D2-D8	VL	aa	EIVMTQSPASMSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
59	BCMA-6	BC 5G9 91-D2-D8	scFv	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYRDSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSSGGGGSGGGSGGSGEIVMTQSPASMSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGGTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
60	BCMA-6 HL x CD3 HL	BC 5G9 91-D2-D8 HL x CD3 HL	bispecific molecule	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYRDSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSSGGGGSGGGSGGSGEIVMTQSPASMSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGGTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN

						NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGGGGGSGQT VVTQEPSLTVS PGGTVTLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYCVLWYSNRWVFGGGTKLTVL
61	BCMA-7	BC 5G9 92- E10-B10	VH CDR1	aa		NYDMA
62	BCMA-7	BC 5G9 92- E10-B10	VH CDR2	aa		SIITSGDATYYRDSVKG
63	BCMA-7	BC 5G9 92- E10-B10	VH CDR3	aa		HDYYDGSYGFA Y
64	BCMA-7	BC 5G9 92- E10-B10	VL CDR1	aa		KASQSVGINVD
65	BCMA-7	BC 5G9 92- E10-B10	VL CDR2	aa		GASNRHT
66	BCMA-7	BC 5G9 92- E10-B10	VL CDR3	aa		LQYGSIPFT
67	BCMA-7	BC 5G9 92- E10-B10	VH	aa		QVQLVESGGGVQPGRSRLSCAASGFTFSNYDMAWVRQAPKGLEWVASIITSGDATYYRDSVKGR FTVSRDNSKNTLYLQMNSLRAEDTAVYYCVRHDY YDGSYGFA YWGQGTLLVTVSS
68	BCMA-7	BC 5G9 92- E10-B10	VL	aa		EIVMTQSPATLSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGS GTEFTLTISSLQAEDFAVYYCLQYGSIPFTFGPGTKVDIK
69	BCMA-7	BC 5G9 92- E10-B10	scFv	aa		QVQLVESGGGVQPGRSRLSCAASGFTFSNYDMAWVRQAPKGLEWVASIITSGDATYYRDSVKGR FTVSRDNSKNTLYLQMNSLRAEDTAVYYCVRHDY YDGSYGFA YWGQGTLLVTVSSGGGGSGGGGG GSEIVMTQSPATLSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGTEFTLTISSLQAEDFAVYYCLQYGSIPFTFGPGTKVDIK
70	BCMA-7 HL x CD3 HL	BC 5G9 92- E10B10 HL x CD3 HL	bispecific molecule	aa		QVQLVESGGGVQPGRSRLSCAASGFTFSNYDMAWVRQAPKGLEWVASIITSGDATYYRDSVKGR FTVSRDNSKNTLYLQMNSLRAEDTAVYYCVRHDY YDGSYGFA YWGQGTLLVTVSSGGGGSGGGGG GSEIVMTQSPATLSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGTEFTLTISSLQAEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWNVRQAPKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGGGGGSGQT VVTQEPSLTVS PGGTVTLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYCVLWYSNRWVFGGGTKLTVL
71	BCMA-8	BC 5G9 92- E10-D8	VH CDR1	aa		NYDMA

72	BCMA-8	BC 5G9 92- E10-D8	VH CDR2	aa	SIITSGDMTYYRDSVKG
73	BCMA-8	BC 5G9 92- E10-D8	VH CDR3	aa	HDYYDGSYGFAY
74	BCMA-8	BC 5G9 92- E10-D8	VL CDR1	aa	KASQSVGINVD
75	BCMA-8	BC 5G9 92- E10-D8	VL CDR2	aa	GASNRHT
76	BCMA-8	BC 5G9 92- E10-D8	VL CDR3	aa	LQYGSIPFT
77	BCMA-8	BC 5G9 92- E10-D8	VH	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYYRDSVKGR FTVSRDNSKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSS
78	BCMA-8	BC 5G9 92- E10-D8	VL	aa	EIVMTQSPATLSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGS GTEFTLTISSLQAEDFAVYYCLQYGSIPFTFGPGTKVDIK
79	BCMA-8	BC 5G9 92- E10-D8	scFv	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYYRDSVKGR FTVSRDNSKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSSGGGGSGGGG GSEIVMTQSPATLSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGTEFTLTISSLQAEDFAVYYCLQYGSIPFTFGPGTKVDIK
80	BCMA-8 HL x CD3 HL	BC 5G9 92- E10-D8 HL x CD3 HL	bispecific molecule	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGDMTYYRDSVKGR FTVSRDNSKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSSGGGGSGGGG GSEIVMTQSPATLSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGTEFTLTISSLQAEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGGQT VVTQEPSLTVS PGGT VTLTCGSS TGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
81	BCMA-9	BC H1 38- D2-A4	VH CDR1	aa	NYWIIH
82	BCMA-9	BC H1 38- D2-A4	VH CDR2	aa	AIYPGNSDTHYNQKFQG
83	BCMA-9	BC H1 38- D2-A4	VH CDR3	aa	SSYYDGSLEFAS
84	BCMA-9	BC H1 38- D2-A4	VL CDR1	aa	RSSQSIVHSNGNTYLY

85	BCMA-9	BC H1 38- D2-A4	VL CDR2	aa	RVSNRFS	
86	BCMA-9	BC H1 38- D2-A4	VL CDR3	aa	FQGSTLLPFT	
87	BCMA-9	BC H1 38- D2-A4	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQ GK VTITRDTASASTAYMELSSLTSEDTA VYYCTRSSYYDGS LFA SWGQGT LVT VSS	
88	BCMA-9	BC H1 38- D2-A4	VL	aa	DI VMTQTPLSLSVSPGQPASISCRSSQSI VHSNGNTYLYWYLQKPGQPPQ LLIYRVSNRFS GVPDRF SGSGSGTDFTLKISRVEAEDVGVYYCFQGSTL PFTFGQGTKLEIK	
89	BCMA-9	BC H1 38- D2-A4	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQ GK VTITRDTASASTAYMELSSLTSEDTA VYYCTRSSYYDGS LFA SWGQGT LVT VSSGGSGGGSGGG GSDI VMTQTPLSLSVSPGQPASISCRSSQSI VHSNGNTYLYWYLQKPGQPPQ LLIYRVSNRFS GVPD RFSGSGSGTDFTLKISRVEAEDVGVYYCFQGSTL PFTFGQGTKLEIK	
90	BCMA-9 HL x CD3 HL	BC H1 38- D2-A4 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQ GK VTITRDTASASTAYMELSSLTSEDTA VYYCTRSSYYDGS LFA SWGQGT LVT VSSGGSGGGSGGG GSDI VMTQTPLSLSVSPGQPASISCRSSQSI VHSNGNTYLYWYLQKPGQPPQ LLIYRVSNRFS GVPD RFSGSGSGTDFTLKISRVEAEDVGVYYCFQGSTL PFTFGQGTKLEIKSGGGSEVQLVESGGLVQP GGSLKLSCAASGFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTA Y LQMNNLKTEDTA VYYCVRHGNFGNSYI SYWAYWGQGT LVT VSSGGSGGGSGGGSTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQ QKPGQAPRGLIGGTKFLAPGTPARFSGSL LGGKAAL TL SGVQPEDEAEYYCVLWYSNRWVFGGTKLTVL	
91	BCMA-10	BC H1 38- D2-F12	VH CDR1	aa	NYWIH	
92	BCMA-10	BC H1 38- D2-F12	VH CDR2	aa	AIYPGNSDTHYNQKFQG	
93	BCMA-10	BC H1 38- D2-F12	VH CDR3	aa	SSYYYDGS LFAS	
94	BCMA-10	BC H1 38- D2-F12	VL CDR1	aa	RSSQSI VHSNGNTYLY	
95	BCMA-10	BC H1 38- D2-F12	VL CDR2	aa	RVSNRFS	
96	BCMA-10	BC H1 38- D2-F12	VL CDR3	aa	FQGS HLPFT	
97	BCMA-10	BC H1 38- D2-F12	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFITNYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQ GK	

					VTITRDTSASTAYMELSSLTSEDTA VYYCTRSSYYDGS L FASWGQGLVTVSS
98	BCMA-10	BC H1 38- D2-F12	VL	aa	DI VMTQTPLSLSVSPGQPASISCRSSQSI VHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPDRF SGSGSGTDFTLKISRVEAEDVGVYYCFQGSHLPFTFGQGTKLEIK
99	BCMA-10	BC H1 38- D2-F12	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQ GK VTITRDTASTAYMELSSLTSEDTA VYYCTRSSYYDGS L FASWGQGLVTVSSGGGGSGGGSGGG GSDIVMTQTPLSLSVSPGQPASISCRSSQSI VHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPD RFSGSGSGTDFTLKISRVEAEDVGVYYCFQGSHLPFTFGQGTKLEIK
100	BCMA-10 HL x CD3 HL	BC H1 38- D2-F12 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQ GK VTITRDTASTAYMELSSLTSEDTA VYYCTRSSYYDGS L FASWGQGLVTVSSGGGGSGGGSGGG GSDIVMTQTPLSLSVSPGQPASISCRSSQSI VHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPD RFSGSGSGTDFTLKISRVEAEDVGVYYCFQGSHLPFTFGQGTKLEIKSGGGSEVQLVESGGGLVQ P GGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTA Y LQMNNLKTEDTA VYYCVRHGNFGNSYI SYWAYWGQGLVTVSSGGGGSGGGSGGT VVTQEP SLTVSPGGTVTLTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAAL TL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
101	BCMA-11	BC H1 38- C1-A4	VH CDR1	aa	NYWIH
102	BCMA-11	BC H1 38- C1-A4	VH CDR2	aa	AIYPGNSDTHYNQKFQG
103	BCMA-11	BC H1 38- C1-A4	VH CDR3	aa	SSYYYDGS LFAS
104	BCMA-11	BC H1 38- C1-A4	VL CDR1	aa	KSSQSI VHSNGNTYLY
105	BCMA-11	BC H1 38- C1-A4	VL CDR2	aa	RVS NRFS
106	BCMA-11	BC H1 38- C1-A4	VL CDR3	aa	FQGSTLPFT
107	BCMA-11	BC H1 38- C1-A4	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQ GK VTITRDTSASTAYMELSSLTSEDTA VYYCTRSSYYDGS L FASWGQGLVTVSS
108	BCMA-11	BC H1 38- C1-A4	VL	aa	DI VMTQTPLSLSVTPGQQASISCKSSQSI VHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPDRF SGSGSGTDFTLKISRVEAEDVGVYYCFQGSHLPFTFGQGTKLEIK
109	BCMA-11	BC H1 38- C1-A4	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQ GK

					VTITRDTSASTAYMELSSLTSED ~~TAVYYCT~~ RSSYYD ~~GS~~ L ~~FAS~~ WGQ ~~GT~~ LVTVSSGGGGGGGGGGGG GSDI ~~VMTQTPLSLSV~~ TPGQ ~~QASISCKSSQ~~ SI ~~VHS~~ NGNTYLYWYLQ ~~KPGQ~~ PPQ ~~LLIYRV~~ SNRFS ~~GV~~ PD R ~~FS~~ GS ~~GS~~ GT ~~DTL~~ KISR ~~VEA~~ EDVG ~~VY~~ YCFQ ~~GS~~ TL ~~PFT~~ FGQ ~~GT~~ KL ~~EIK~~
110	BCMA-11 HL x CD3 HL	BC H1 38- C1-A4 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTF ~~TNY~~ WI ~~HWVK~~ QAPGQ ~~R~~ LEWIGAIYPGNSD ~~THYNQ~~ KFQ ~~GK~~ VTITRDT ~~SASTAYMELSSLT~~ SED ~~TAVYYCT~~ RSSYYD ~~GS~~ L ~~FAS~~ WGQ ~~GT~~ LVTVSSGGGGGGGGGGGG GSDI ~~VMTQTPLSLSV~~ TPGQ ~~QASISCKSSQ~~ SI ~~VHS~~ NGNTYLYWYLQ ~~KPGQ~~ PPQ ~~LLIYRV~~ SNRFS ~~GV~~ PD R ~~FS~~ GS ~~GS~~ GT ~~DTL~~ KISR ~~VEA~~ EDVG ~~VY~~ YCFQ ~~GS~~ TL ~~PFT~~ FGQ ~~GT~~ KL ~~EIK~~ KS ~~GGSE~~ VQ ~~LVE~~ SG ~~GLV~~ Q ~~P~~ G ~~SL~~ KLSCAASGFTFNKYAMN ~~WVRQ~~ APGK ~~GLEW~~ VARIRSKYNNYATYYADSVKDRFTISR ~~DD~~ SKNTA YLQMN ~~N~~ LKTEDTAVYYCVRHGNFGNSYISYWAYWGQ ~~GT~~ LVTVSSGGGGGGGGGGGG ~~Q~~ TVVTQEP SLTVSPGGTVTLTCGSS ~~TG~~ AVTSGNYPNWVQ ~~QKPGQ~~ APRGLIGG ~~TK~~ FLAPGTPARFSGSLLGGKAAAL T ~~LSGVQ~~ PEDEAEYYCVLWYSNRWVFGGGTKLT ~~VL~~
111	BCMA-12	BC H1 38- C1-F12	VH CDR1	aa	NYW ~~I~~ H
112	BCMA-12	BC H1 38- C1-F12	VH CDR2	aa	AIYPGNSD ~~THYNQ~~ KFQ ~~G~~
113	BCMA-12	BC H1 38- C1-F12	VH CDR3	aa	SSYYD ~~GS~~ L ~~FAS~~
114	BCMA-12	BC H1 38- C1-F12	VL CDR1	aa	KSSQSI ~~VHS~~ NGNTYLY
115	BCMA-12	BC H1 38- C1-F12	VL CDR2	aa	R ~~V~~ SNRFS
116	BCMA-12	BC H1 38- C1-F12	VL CDR3	aa	FQ ~~GS~~ HL ~~PFT~~
117	BCMA-12	BC H1 38- C1-F12	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTF ~~TNY~~ WI ~~HWVK~~ QAPGQ ~~R~~ LEWIGAIYPGNSD ~~THYNQ~~ KFQ ~~GK~~ VTITRDT ~~SASTAYMELSSLT~~ SED ~~TAVYYCT~~ RSSYYD ~~GS~~ L ~~FAS~~ WGQ ~~GT~~ LVTVSS
118	BCMA-12	BC H1 38- C1-F12	VL	aa	DI ~~VMTQTPLSLSV~~ TPGQ ~~QASISCKSSQ~~ SI ~~VHS~~ NGNTYLYWYLQ ~~KPGQ~~ PPQ ~~LLIYRV~~ SNRFS ~~GV~~ PD ~~R~~ F SGSGGT ~~DTL~~ KISR ~~VEA~~ EDVG ~~VY~~ YCFQ ~~GS~~ HL ~~PFT~~ FGQ ~~GT~~ KL ~~EIK~~
119	BCMA-12	BC H1 38- C1-F12	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTF ~~TNY~~ WI ~~HWVK~~ QAPGQ ~~R~~ LEWIGAIYPGNSD ~~THYNQ~~ KFQ ~~GK~~ VTITRDT ~~SASTAYMELSSLT~~ SED ~~TAVYYCT~~ RSSYYD ~~GS~~ L ~~FAS~~ WGQ ~~GT~~ LVTVSSGGGGGGGGGGGG GSDI ~~VMTQTPLSLSV~~ TPGQ ~~QASISCKSSQ~~ SI ~~VHS~~ NGNTYLYWYLQ ~~KPGQ~~ PPQ ~~LLIYRV~~ SNRFS ~~GV~~ PD R ~~FS~~ GS ~~GS~~ GT ~~DTL~~ KISR ~~VEA~~ EDVG ~~VY~~ YCFQ ~~GS~~ HL ~~PFT~~ FGQ ~~GT~~ KL ~~EIK~~
120	BCMA-12 HL x CD3 HL	BC H1 38- C1-F12 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTF ~~TNY~~ WI ~~HWVK~~ QAPGQ ~~R~~ LEWIGAIYPGNSD ~~THYNQ~~ KFQ ~~GK~~ VTITRDT ~~SASTAYMELSSLT~~ SED ~~TAVYYCT~~ RSSYYD ~~GS~~ L ~~FAS~~ WGQ ~~GT~~ LVTVSSGGGGGGGGGGGG

					GSDIVMTQTPLSLSVTPGQQASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPDRFSGSGSGTDFTLKISRVEAEDVGYVYCFQGSHLPFTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAAL TLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
121	BCMA-13	BC H1 39- B2-A4	VH CDR1	aa	NYWIIH
122	BCMA-13	BC H1 39- B2-A4	VH CDR2	aa	AIYPGNSDTHYNQKFQG
123	BCMA-13	BC H1 39- B2-A4	VH CDR3	aa	SSYYDGS LFAS
124	BCMA-13	BC H1 39- B2-A4	VL CDR1	aa	KSSQSIVHSNGNTYLY
125	BCMA-13	BC H1 39- B2-A4	VL CDR2	aa	RVS NRFS
126	BCMA-13	BC H1 39- B2-A4	VL CDR3	aa	FQGSTLPFT
127	BCMA-13	BC H1 39- B2-A4	VH	aa	QVQLVQSGAVVAKPGASVKVSKKASGYTFITNYWIHWVKQAPGQRLIEWMGA IYPGNSDTHYNQKFQGRVTLTDTDSASTAYMELSSLRNEDTAVYYCTRSSYYDGS LFASWGQGTLLVTVSS
128	BCMA-13	BC H1 39- B2-A4	VL	aa	DI VMTQTPLSLSVTPGQQASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPDRFSGSGSGTDFTLKISRVEAEDVGYVYCFQGSHLPFTFGQGTKLEIK
129	BCMA-13	BC H1 39- B2-A4	scFv	aa	QVQLVQSGAVVAKPGASVKVSKKASGYTFITNYWIHWVKQAPGQRLIEWMGA IYPGNSDTHYNQKFQGRVTLTDTDSASTAYMELSSLRNEDTAVYYCTRSSYYDGS LFASWGQGTLLVTVSSGGGGSGGGGGGSDI VMTQTPLSLSVTPGQQASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPDRFSGSGSGTDFTLKISRVEAEDVGYVYCFQGSHLPFTFGQGTKLEIK
130	BCMA-13 HL x CD3 HL	BC H1 39- B2-A4 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAVVAKPGASVKVSKKASGYTFITNYWIHWVKQAPGQRLIEWMGA IYPGNSDTHYNQKFQGRVTLTDTDSASTAYMELSSLRNEDTAVYYCTRSSYYDGS LFASWGQGTLLVTVSSGGGGSGGGGGGSDI VMTQTPLSLSVTPGQQASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPDRFSGSGSGTDFTLKISRVEAEDVGYVYCFQGSHLPFTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAAL

						TLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
131	BCMA-14	BC H1 39- B2-F12	VH CDR1	aa		NYWIIH
132	BCMA-14	BC H1 39- B2-F12	VH CDR2	aa		AIYPGNSDTHYNQKFQG
133	BCMA-14	BC H1 39- B2-F12	VH CDR3	aa		SSYYDGSLFAS
134	BCMA-14	BC H1 39- B2-F12	VL CDR1	aa		KSSQSIVHSNGNTYLY
135	BCMA-14	BC H1 39- B2-F12	VL CDR2	aa		RVSNRFS
136	BCMA-14	BC H1 39- B2-F12	VL CDR3	aa		FQGSHLPFT
137	BCMA-14	BC H1 39- B2-F12	VH	aa		QVQLVQSGAVVAKPGASVKVSKKASGYTFFTNYWIHWVKQAPGQRLLEWMGAIYPGNSDTHYNQKFQGR VTLTDTDSASTAYMELSSLRNEDTAVYYCTRSSYYDGSLFASWGQGLTVTVSSSGGGGGTGLTVTVSS
138	BCMA-14	BC H1 39- B2-F12	VL	aa		DIVMTQTPLSLSVTPGQQASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQQLLIYRVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVYYCFQGSHLPFTFGQGTKLEIK
139	BCMA-14	BC H1 39- B2-F12	scFv	aa		QVQLVQSGAVVAKPGASVKVSKKASGYTFFTNYWIHWVKQAPGQRLLEWMGAIYPGNSDTHYNQKFQGR VTLTDTDSASTAYMELSSLRNEDTAVYYCTRSSYYDGSLFASWGQGLTVTVSSSGGGGGTGLTVTVSSSGGGGG GSDIVMTQTPLSLSVTPGQQASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQQLLIYRVSNRFSGVPPD RFSGSGSGTDFTLKISRVEAEDVGVYYCFQGSHLPFTFGQGTKLEIK
140	BCMA-14 HL x CD3 HL	BC H1 39- B2-F12 HL x CD3 HL	bispecific molecule	aa		QVQLVQSGAVVAKPGASVKVSKKASGYTFFTNYWIHWVKQAPGQRLLEWMGAIYPGNSDTHYNQKFQGR VTLTDTDSASTAYMELSSLRNEDTAVYYCTRSSYYDGSLFASWGQGLTVTVSSSGGGGGTGLTVTVSSSGGGGG GSDIVMTQTPLSLSVTPGQQASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQQLLIYRVSNRFSGVPPD RFSGSGSGTDFTLKISRVEAEDVGVYYCFQGSHLPFTFGQGTKLEIKSFGGGSEVQLVESGGGLVQRP GGSLKLSCAAAGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTA YLMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSSGGGGGTGLTVTVSSSGGGGGTGLTVTVSS SLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLGGKAAL TLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
141	BCMA-15	BC H1 39- C9-A4	VH CDR1	aa		SYWIIH
142	BCMA-15	BC H1 39- C9-A4	VH CDR2	aa		AIYPGNSDTHYNQKFQG

143	BCMA-15	BC H1 39- C9-A4	VH CDR3	aa	SSYYDGSLEFAD	
144	BCMA-15	BC H1 39- C9-A4	VL CDR1	aa	KSSQSIVHSNGNTYLY	
145	BCMA-15	BC H1 39- C9-A4	VL CDR2	aa	RVSNRFS	
146	BCMA-15	BC H1 39- C9-A4	VL CDR3	aa	FQGSTLPFT	
147	BCMA-15	BC H1 39- C9-A4	VH	aa	QVQLVQSGAEVKKPGTSVKVSKKASGYTFTSYWIHWVKQAPGQRLEWIGAIYPGNSDTHYNQKFQGR VTLTRDTSASTAYMELSSLRSEDSAVYYCTRSSYYDGSLEFADWGQGTLVTVSS	
148	BCMA-15	BC H1 39- C9-A4	VL	aa	DI VMTQTPLSLSVTPGQPASISCKSSQSI VHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVYYCFQGSTLPFTFGQGTKLEIK	
149	BCMA-15	BC H1 39- C9-A4	scFv	aa	QVQLVQSGAEVKKPGTSVKVSKKASGYTFTSYWIHWVKQAPGQRLEWIGAIYPGNSDTHYNQKFQGR VTLTRDTSASTAYMELSSLRSEDSAVYYCTRSSYYDGSLEFADWGQGTLVTVSSGGSGGGSGGG GSDI VMTQTPLSLSVTPGQPASISCKSSQSI VHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPPD RFSGSGSGTDFTLKISRVEAEDVGVYYCFQGSTLPFTFGQGTKLEIK	
150	BCMA-15 HL x CD3 HL	BC H1 39- C9-A4 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGTSVKVSKKASGYTFTSYWIHWVKQAPGQRLEWIGAIYPGNSDTHYNQKFQGR VTLTRDTSASTAYMELSSLRSEDSAVYYCTRSSYYDGSLEFADWGQGTLVTVSSGGSGGGSGGG GSDI VMTQTPLSLSVTPGQPASISCKSSQSI VHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVPPD RFSGSGSGTDFTLKISRVEAEDVGVYYCFQGSTLPFTFGQGTKLEIKSGGGSEVQLVESGGGLVQP GGSLKLSCAAAGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTA YLMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLVTVSSGGSGGGSGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAAL TLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL	
151	BCMA-16	BC H1 39- C9-F12	VH CDR1	aa	SYWIH	
152	BCMA-16	BC H1 39- C9-F12	VH CDR2	aa	AIYPGNSDTHYNQKFQG	
153	BCMA-16	BC H1 39- C9-F12	VH CDR3	aa	SSYYDGSLEFAD	
154	BCMA-16	BC H1 39- C9-F12	VL CDR1	aa	KSSQSIVHSNGNTYLY	
155	BCMA-16	BC H1 39- C9-F12	VL CDR2	aa	RVSNRFS	

156	BCMA-16	BC H1 39- C9-F12	VL CDR3	aa	FQGSHLPPFT
157	BCMA-16	BC H1 39- C9-F12	VH	aa	QVQLVQSGAEVKKPGTSVKVSCKASGYTFTSYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQGR VTLTRDTSASTAYMELSSLRSEDSAVYYCTRSSYYDGS LFADWGQGLVTVSS
158	BCMA-16	BC H1 39- C9-F12	VL	aa	DIVMTQTPLSLSVTPGQPASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVDPDRF SGSGSGTDFTLKISRVEAEDVGVYCFQGS HLPFTFGQGTKLEIK
159	BCMA-16	BC H1 39- C9-F12	scFv	aa	QVQLVQSGAEVKKPGTSVKVSCKASGYTFTSYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQGR VTLTRDTSASTAYMELSSLRSEDSAVYYCTRSSYYDGS LFADWGQGLVTVSSGGGGSGGGSGGG GSDIVMTQTPLSLSVTPGQPASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVDP RFSGSGSGTDFTLKISRVEAEDVGVYCFQGS HLPFTFGQGTKLEIK
160	BCMA-16 HL x CD3 HL	BC H1 39- C9-F12 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGTSVKVSCKASGYTFTSYWIHWVKQAPGQRLIEWIGAIYPGNSDTHYNQKFQGR VTLTRDTSASTAYMELSSLRSEDSAVYYCTRSSYYDGS LFADWGQGLVTVSSGGGGSGGGSGGG GSDIVMTQTPLSLSVTPGQPASISCKSSQSIVHSNGNTYLYWYLQKPGQPPQLLIYRVSNRFSGVDP RFSGSGSGTDFTLKISRVEAEDVGVYCFQGS HLPFTFGQGTKLEIKSGGGSEVQLVESGGGLVQP GGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTA YLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGSGGGSQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGTKFLAPGTPARFSGSLLGKKAAL TLSGVQPEDEAEY YCVLWY SNRWVFGGGTKLTVL
161	BCMA-17	BC C3 33- D7-E6	VH CDR1	aa	NFDMA
162	BCMA-17	BC C3 33- D7-E6	VH CDR2	aa	SITTGADHAIYADSVKG
163	BCMA-17	BC C3 33- D7-E6	VH CDR3	aa	HGYDGYHLFDY
164	BCMA-17	BC C3 33- D7-E6	VL CDR1	aa	RASQGISNYLN
165	BCMA-17	BC C3 33- D7-E6	VL CDR2	aa	YTSNLQS
166	BCMA-17	BC C3 33- D7-E6	VL CDR3	aa	QQYDISSYT
167	BCMA-17	BC C3 33- D7-E6	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGR FTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSS
168	BCMA-17	BC C3 33- D7-E6	VL	aa	DIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKGAPKPLIYYTSLNLSQGVPSRFSGSGS

		D7-E6				GTDYTLTISSLQPEDEFATYYCQQYDISSYTFGQGTKLEIK
169	BCMA-17	BC C3 33-D7-E6	scFv	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSSGGGGGGGGGGGSDIQMTQSPSSLSASVGDRTITCRASQGISNYLNWYQQKPGKAPKPLIYYTSNQLQSGVPSRFSGSGSGTDYTLTISSLQPEDEFATYYCQQYDISSYTFGQGTKLEIK
170	BCMA-17 HL x CD3 HL	BC C3 33-D7-E6 HL x CD3 HL	bispecific molecule	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSSGGGGGGGGGGGSDIQMTQSPSSLSASVGDRTITCRASQGISNYLNWYQQKPGKAPKPLIYYTSNQLQSGVPSRFSGSGSGTDYTLTISSLQPEDEFATYYCQQYDISSYTFGQGTKLEIK
171	BCMA-18	BC C3 33-D7-E6B1	VH CDR1	aa		NFDMA
172	BCMA-18	BC C3 33-D7-E6B1	VH CDR2	aa		SITTGADHAIYADSVKG
173	BCMA-18	BC C3 33-D7-E6B1	VH CDR3	aa		HGY YDGYHLFDY
174	BCMA-18	BC C3 33-D7-E6B1	VL CDR1	aa		RASQGISNYLN
175	BCMA-18	BC C3 33-D7-E6B1	VL CDR2	aa		YTSNLQS
176	BCMA-18	BC C3 33-D7-E6B1	VL CDR3	aa		MGQTISSYT
177	BCMA-18	BC C3 33-D7-E6B1	VH	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSS
178	BCMA-18	BC C3 33-D7-E6B1	VL	aa		DIQMTQSPSSLSASVGDRTITCRASQGISNYLNWYQQKPGKAPKPLIYYTSNQLQSGVPSRFSGSGSGTDYTLTISSLQPEDEFATYYCMGQTISSYTFGQGTKLEIK
179	BCMA-18	BC C3 33-D7-E6B1	scFv	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSSGGGGGGGGGGGSDIQMTQSPSSLSASVGDRTITCRASQGISNYLNWYQQKPGKAPKPLIYYTSNQLQSGVPSRFSGSGSGTDYTLTISSLQPEDEFATYYCMGQTISSYTFGQGTKLEIK

180	BCMA-18 HL x CD3 HL	BC C3 33- D7-E6B1 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLLFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDRTVITCRASQGISNVLNWIYQKPGKAPKPLIYYTSNQLQSGVPSRFSGSGSGTDYTLTISSLPEDFATYYCMGQTISSYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGSGGTVVVTQEPSLTVS PGGTVTLTCGSSTGAVTSGNYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
181	BCMA-19	BC C3 33- F8-E6	VH CDR1	aa	NFDMA
182	BCMA-19	BC C3 33- F8-E6	VH CDR2	aa	SITTGADHAIYADSVKG
183	BCMA-19	BC C3 33- F8-E6	VH CDR3	aa	HGYDGYHLLFDY
184	BCMA-19	BC C3 33- F8-E6	VL CDR1	aa	RASQGISNYLN
185	BCMA-19	BC C3 33- F8-E6	VL CDR2	aa	YTSNLQS
186	BCMA-19	BC C3 33- F8-E6	VL CDR3	aa	QQYDISSYT
187	BCMA-19	BC C3 33- F8-E6	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLLFDYWGQGLVTVSS
188	BCMA-19	BC C3 33- F8-E6	VL	aa	DIQMTQSPSSLSASVGDRTVITCRASQGISNVLNWIYQKPGKAPKPLIYYTSNQLQSGVPSRFSGSGSGTDYTLTISSLPEDFATYYCQYDISSYTFGQGTKLEIK
189	BCMA-19	BC C3 33- F8-E6	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLLFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDRTVITCRASQGISNVLNWIYQKPGKAPKPLIYYTSNQLQSGVPSRFSGSGSGTDYTLTISSLPEDFATYYCQYDISSYTFGQGTKLEIK
190	BCMA-19 HL x CD3 HL	BC C3 33- F8-E6 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLLFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDRTVITCRASQGISNVLNWIYQKPGKAPKPLIYYTSNQLQSGVPSRFSGSGSGTDYTLTISSLPEDFATYYCQYDISSYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN

						NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGGGGGSGGSGQTVVVTQEPSLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
191	BCMA-20	BC C3 33-F8-E6B1	VH CDR1	aa		NFDMA
192	BCMA-20	BC C3 33-F8-E6B1	VH CDR2	aa		SITTGADHAIYADSVKG
193	BCMA-20	BC C3 33-F8-E6B1	VH CDR3	aa		HGYDGYHLEFDY
194	BCMA-20	BC C3 33-F8-E6B1	VL CDR1	aa		RASQGISNYLN
195	BCMA-20	BC C3 33-F8-E6B1	VL CDR2	aa		YTSNLQS
196	BCMA-20	BC C3 33-F8-E6B1	VL CDR3	aa		MGQTISSYT
197	BCMA-20	BC C3 33-F8-E6B1	VH	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYVDGYHLEFDYWGQGTLLTVVSS
198	BCMA-20	BC C3 33-F8-E6B1	VL	aa		DIQMTQSPSSLSASVGDRTVITCRASQGISNYLNWYQQKPKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK
199	BCMA-20	BC C3 33-F8-E6B1	scFv	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYVDGYHLEFDYWGQGTLLTVVSSGGGGGGSGGGGGSDIQMTQSPSSLSASVGDRTVITCRASQGISNYLNWYQQKPKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK
200	BCMA-20 HL x CD3 HL	BC C3 33-F8-E6B1 HL x CD3 HL	bispecific molecule	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYVDGYHLEFDYWGQGTLLTVVSSGGGGGGSGGGGGSDIQMTQSPSSLSASVGDRTVITCRASQGISNYLNWYQQKPKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN
201	BCMA-21	BC C3 33-F9-E6	VH CDR1	aa		NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGGGGGSGGSGQTVVVTQEPSLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL

202	BCMA-21	BC C3 33-F9-E6	VH CDR2	aa	SITTGADHAIYADSVKG
203	BCMA-21	BC C3 33-F9-E6	VH CDR3	aa	HGYDGYHLFDY
204	BCMA-21	BC C3 33-F9-E6	VL CDR1	aa	RASQGISNYLN
205	BCMA-21	BC C3 33-F9-E6	VL CDR2	aa	YTSNLQS
206	BCMA-21	BC C3 33-F9-E6	VL CDR3	aa	QQYDISSYT
207	BCMA-21	BC C3 33-F9-E6	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMDSLRSIEDTAVYYCVRHGYYDGYHLFDYWGQGLVTVSS
208	BCMA-21	BC C3 33-F9-E6	VL	aa	DIQMTQSPSSLSASVGDRTVITSCRASQGISNYLNWYQQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCQQYDISSYTFGQGTKLEIK
209	BCMA-21	BC C3 33-F9-E6	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMDSLRSIEDTAVYYCVRHGYYDGYHLFDYWGQGLVTVSSGGGGSGGGGGGSDIQMTQSPSSLSASVGDRTVITSCRASQGISNYLNWYQQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCQQYDISSYTFGQGTKLEIK
210	BCMA-21 HL x CD3 HL	BC C3 33-F9-E6 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYADSVKGRFTISRDNAKNTLYLQMDSLRSIEDTAVYYCVRHGYYDGYHLFDYWGQGLVTVSSGGGGSGGGGGGGSDIQMTQSPSSLSASVGDRTVITSCRASQGISNYLNWYQQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCQQYDISSYTFGQGTKLEIK
211	BCMA-22	BC C3 33-F9-E6B1-E	VH CDR1	aa	NFDMA
212	BCMA-22	BC C3 33-F9-E6B1-E	VH CDR2	aa	SITTGADHAIYAESVKG
213	BCMA-22	BC C3 33-F9-E6B1-E	VH CDR3	aa	HGYDGYHLFDY
214	BCMA-22	BC C3 33-F9-E6B1-E	VL CDR1	aa	RASQGISNYLN

215	BCMA-22	BC C3 33-F9-E6B1-E	VL CDR2	aa	YTSNLQS	
216	BCMA-22	BC C3 33-F9-E6B1-E	VL CDR3	aa	MGQTISSYT	
217	BCMA-22	BC C3 33-F9-E6B1-E	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYAESVKGR FTISRDNAKNTLYLQMDSLRSEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSS	
218	BCMA-22	BC C3 33-F9-E6B1-E	VL	aa	DIQMTQSPSSLSASVGDRVTISCRASQGISNYLNWYQQKPKGAPKPLIYYTSTNLQSGVPSRFSGSGS GTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK	
219	BCMA-22	BC C3 33-F9-E6B1-E	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYAESVKGR FTISRDNAKNTLYLQMDSLRSEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSSGGGGGGGGGG GSDIQMTQSPSSLSASVGDRVTISCRASQGISNYLNWYQQKPKGAPKPLIYYTSTNLQSGVPSRFSGSGS GGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK	
220	BCMA-22 HL x CD3 HL	BC C3 33-F9-E6B1-E HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGADHAIYAESVKGR FTISRDNAKNTLYLQMDSLRSEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSSGGGGGGGGGG GSDIQMTQSPSSLSASVGDRVTISCRASQGISNYLNWYQQKPKGAPKPLIYYTSTNLQSGVPSRFSGSGS GGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGGGGGGGQTVVTQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYCVLWYSNRWVFGGGTKLTVL	
221	BCMA-23	BC C3 33-F10-E6B1	VH CDR1	aa	NFDMA	
222	BCMA-23	BC C3 33-F10-E6B1	VH CDR2	aa	SITTGADHAIYADSVKG	
223	BCMA-23	BC C3 33-F10-E6B1	VH CDR3	aa	HGYDGYHLFDY	
224	BCMA-23	BC C3 33-F10-E6B1	VL CDR1	aa	RASQGISNYLN	
225	BCMA-23	BC C3 33-F10-E6B1	VL CDR2	aa	YTSNLQS	
226	BCMA-23	BC C3 33-F10-E6B1	VL CDR3	aa	MGQTISSYT	
227	BCMA-23	BC C3 33-F10-E6B1	VH	aa	EVQLVESGGGLVQPGRSLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTGADHAIYADSVKGR	

					FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLFDYWGQGLVTVSS
228	BCMA-23	BC C3 33- F10-E6B1	VL	aa	DIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKAPKPLIYYTNSNLQSGVPSRFSGSGS GTDFTLLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK
229	BCMA-23	BC C3 33- F10-E6B1	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTGADHAIYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLFDYWGQGLVTVSSGGGGGGGGGGGG GSDIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKAPKPLIYYTNSNLQSGVPSRFSGSGS GSGTDFTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK
230	BCMA-23 HL x CD3 HL	BC C3 33- F10-E6B1 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTGADHAIYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLFDYWGQGLVTVSSGGGGGGGGGGGG GSDIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKAPKPLIYYTNSNLQSGVPSRFSGSGS GSGTDFTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGGGGGGGGGQTVVTTQEPSLTVS PGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYCVLWYSNRWVFGGGTKLTVL
231	BCMA-24	BC B6 64- H5-A4	VH CDR1	aa	DYYIN
232	BCMA-24	BC B6 64- H5-A4	VH CDR2	aa	WIYFASGNSEYNQKFTG
233	BCMA-24	BC B6 64- H5-A4	VH CDR3	aa	LYDYDWYFDV
234	BCMA-24	BC B6 64- H5-A4	VL CDR1	aa	KSSQSLVHSNGNTYLH
235	BCMA-24	BC B6 64- H5-A4	VL CDR2	aa	KVSNRFS
236	BCMA-24	BC B6 64- H5-A4	VL CDR3	aa	AETSHVPWT
237	BCMA-24	BC B6 64- H5-A4	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSEFPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
238	BCMA-24	BC B6 64- H5-A4	VL	aa	DIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLVQKPGQSPQLLIYKVSNRFSGVPDFR SGSGGTDFTLKINRVEAEDVGVIYCAETSHVPWTFGQGTKLEIK
239	BCMA-24	BC B6 64- H5-A4	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSEFPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR

					VTMTRDTSSSTAYMELSSLRSED TAVYFCASLYDYDWYFDVWGQGT MTVTVSSGGGGSGGGSGGGGS DIVMTQTPLSLSVTPGQPASIS CKSSQSLVHSNGNTYLHWY LQKPGQSPQLLIYKVS NRFSGVPDRFSGSGGTDFTL KINRVEAEDVGVIYCAETSH VPWTFGQGTKLEIK
240	BCMA-24 HL x CD3 HL	BC B6 64- H5-A4 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVK VCKASGYSPFDYINWVRQAP GQGLEWMGWIFYFASGNSE YNQKFTGRVTMTRDTSSST AYMELSSLRSED TAVYFCASLYDYDWYFDV WGQGTMTVTVSSGGGGSG GGSGGGGS DIVMTQTPLSLSVTPGQPAS ISCKSSQSLVHSNGNTYLH WYILQKPGQSPQLLIYKVS NRFSGVPDRFSGSGGTDFT LKINRVEAEDVGVIYCAET SHVPWTFGQGTKLEIKS GGGSEVQLVESGGGLVQ PFGSLKLSCAASGFTFNKY AMNWVRQAPGKGLEWVAR IRSKYNNYATYYADSVKDR FTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNF GNSYISYWAYWGQGT LTVTVSSGGGGSGGGSG GGSTVVTQE FSLTVSPGGTVTLTCGSS TGAVTSGNYPNWVQQK PGQAPRGLIGG TKFLAPGTPARFSGS LLGGKAALTLSGVQPE DEAEYYCVLWYSNRW VFGGTKLTVL
241	BCMA-25	BC B6 64- H5-H9	VH CDR1	aa	DYIIN
242	BCMA-25	BC B6 64- H5-H9	VH CDR2	aa	WIYFASGNSEYNQKFTG
243	BCMA-25	BC B6 64- H5-H9	VH CDR3	aa	LYDYDWYFDV
244	BCMA-25	BC B6 64- H5-H9	VL CDR1	aa	KSSQSLVHSNGNTYLH
245	BCMA-25	BC B6 64- H5-H9	VL CDR2	aa	KVSNRFS
246	BCMA-25	BC B6 64- H5-H9	VL CDR3	aa	LTTS HVPWT
247	BCMA-25	BC B6 64- H5-H9	VH	aa	QVQLVQSGAEVKKPGASVK VCKASGYSPFDYINWVRQAP GQGLEWMGWIFYFASGNSE YNQKFTGRVTMTRDTSSST AYMELSSLRSED TAVYFCASLYDYDWYFDV WGQGTMTVTVSS
248	BCMA-25	BC B6 64- H5-H9	VL	aa	DIVMTQTPLSLSVTPGQPAS ISCKSSQSLVHSNGNTYLH WYILQKPGQSPQLLIYKVS NRFSGVPDRFSGSGGTDFT LKINRVEAEDVGVIYCLT TSHVPWTFGQGTKLEIK
249	BCMA-25	BC B6 64- H5-H9	scFv	aa	QVQLVQSGAEVKKPGASVK VCKASGYSPFDYINWVRQAP GQGLEWMGWIFYFASGNSE YNQKFTGRVTMTRDTSSST AYMELSSLRSED TAVYFCASLYDYDWYFDV WGQGTMTVTVSSGGGGSG GGSGGGGS DIVMTQTPLSLSVTPGQPAS ISCKSSQSLVHSNGNTYLH WYILQKPGQSPQLLIYKVS NRFSGVPDRFSGSGGTDFT LKINRVEAEDVGVIYCLT TSHVPWTFGQGTKLEIK
250	BCMA-25 HL x CD3 HL	BC B6 64- H5-H9 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVK VCKASGYSPFDYINWVRQAP GQGLEWMGWIFYFASGNSE YNQKFTGRVTMTRDTSSST AYMELSSLRSED TAVYFCASLYDYDWYFDV WGQGTMTVTVSSGGGGSG GGSGGGGS

					DI VMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPDRF SGSGSGTDFTLLKINRVEAEDVGVIYCLTTTSHVPWTFGQGTKLEIKSGGGSEVQLVESGGLVQPGG SLKLSCAAAGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDS KNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQTLVTVSSGGSGGGSGGGSQTVVTQEPSL TVSPGGT VTLTTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
251	BCMA-26	BC B6 65- B5-A4	VH CDR1	aa	DYIIN
252	BCMA-26	BC B6 65- B5-A4	VH CDR2	aa	WIYFASGNSEYNQKFTG
253	BCMA-26	BC B6 65- B5-A4	VH CDR3	aa	LYDYDWYFDV
254	BCMA-26	BC B6 65- B5-A4	VL CDR1	aa	KSSQSLVHSNGNTYLH
255	BCMA-26	BC B6 65- B5-A4	VL CDR2	aa	KVSNRFS
256	BCMA-26	BC B6 65- B5-A4	VL CDR3	aa	AETSHVPWT
257	BCMA-26	BC B6 65- B5-A4	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
258	BCMA-26	BC B6 65- B5-A4	VL	aa	DI VMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPDRF SGSGSGTDFTLLKISRVEAEDVGVIYCAETSHVPWTFGQGTKLEIK
259	BCMA-26	BC B6 65- B5-A4	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGGS DI VMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPDRF SGSGSGTDFTLLKISRVEAEDVGVIYCAETSHVPWTFGQGTKLEIK
260	BCMA-26 HL x CD3 HL	BC B6 65- B5-A4 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGGS DI VMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPDRF SGSGSGTDFTLLKISRVEAEDVGVIYCAETSHVPWTFGQGTKLEIKSGGGSEVQLVESGGLVQPGG SLKLSCAAAGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDS KNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQTLVTVSSGGSGGGSGGGSQTVVTQEPSL TVSPGGT VTLTTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT

						SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
261	BCMA-27	BC B6 65- B5-H9	VH CDR1	aa		DYYIN
262	BCMA-27	BC B6 65- B5-H9	VH CDR2	aa		WIYFASGNSEYNQKFTG
263	BCMA-27	BC B6 65- B5-H9	VH CDR3	aa		LYDYDWYFDV
264	BCMA-27	BC B6 65- B5-H9	VL CDR1	aa		KSSQSLVHSNGNTYLH
265	BCMA-27	BC B6 65- B5-H9	VL CDR2	aa		KVSNRFS
266	BCMA-27	BC B6 65- B5-H9	VL CDR3	aa		LTTSHPWT
267	BCMA-27	BC B6 65- B5-H9	VH	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
268	BCMA-27	BC B6 65- B5-H9	VL	aa		DIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVYYCLTTSHPWTFGQGTKLEIK
269	BCMA-27	BC B6 65- B5-H9	scFv	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGGS DIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVYYCLTTSHPWTFGQGTKLEIK
270	BCMA-27 HL x CD3 HL	BC B6 65- B5-H9 HL x CD3 HL	bispecific molecule	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGGS DIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVYYCLTTSHPWTFGQGTKLEIKS GGGGSEVQLVESGGGLVQPGG SLKLSCAAAGFTFNKYAMNNWVRQAPGKGLEWVARIRSKYNNYAITYYADSVKDRFTISRDDSKNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLVTVSSGGSGGGSGGGSGQTVVVTQEPSL TVSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKKAALLTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
271	BCMA-28	BC B6 65- H7-A4	VH CDR1	aa		DYYIN
272	BCMA-28	BC B6 65- H7-A4	VH CDR2	aa		WIYFASGNSEYNQKFTG

273	BCMA-28	BC B6 65- H7-A4	VH CDR3	aa	LYDYDWYFDV
274	BCMA-28	BC B6 65- H7-A4	VL CDR1	aa	KSSQSLVHSNGNTYLH
275	BCMA-28	BC B6 65- H7-A4	VL CDR2	aa	KVSNRFS
276	BCMA-28	BC B6 65- H7-A4	VL CDR3	aa	AETSHVPWT
277	BCMA-28	BC B6 65- H7-A4	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDY _Y INWVRQAPGQGLEWMGW _{IY} FASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
278	BCMA-28	BC B6 65- H7-A4	VL	aa	DI VMTQTPLSLSPGQPASISCKSSQSLVHSNGNTY _L HWYLQKPGQSPQ _{LL} IYKVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVY _Y CAETSHVPWTFGQGTKLEIK
279	BCMA-28	BC B6 65- H7-A4	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDY _Y INWVRQAPGQGLEWMGW _{IY} FASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGGS DI VMTQTPLSLSPGQPASISCKSSQSLVHSNGNTY _L HWYLQKPGQSPQ _{LL} IYKVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVY _Y CAETSHVPWTFGQGTKLEIK
280	BCMA-28 HL x CD3 HL	BC B6 65- H7-A4 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDY _Y INWVRQAPGQGLEWMGW _{IY} FASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGGS DI VMTQTPLSLSPGQPASISCKSSQSLVHSNGNTY _L HWYLQKPGQSPQ _{LL} IYKVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVY _Y CAETSHVPWTFGQGTKLEIKS _{GGG} SEVQLVESGGGLVQPGG SLKLSCAAAGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATY _Y ADSVKDRFTISRDDSKNTAYL QMNNLKTEDTAVY _Y CVRHGNFGNSYISYWAYWGQGT _L TVTVSSGGSGGGSGGGGSQ _T VVTVQEPSL TVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGKAAALTL SGVQPEDEAEY _Y CVLWYSNRWVFGGGTKLTVL
281	BCMA-29	BC B6 65- H7-H9	VH CDR1	aa	DY _Y IN
282	BCMA-29	BC B6 65- H7-H9	VH CDR2	aa	WIYFASGNSEYNQKFTG
283	BCMA-29	BC B6 65- H7-H9	VH CDR3	aa	LYDYDWYFDV
284	BCMA-29	BC B6 65- H7-H9	VL CDR1	aa	KSSQSLVHSNGNTYLH
285	BCMA-29	BC B6 65- H7-H9	VL CDR2	aa	KVSNRFS

286	BCMA-29	BC B6 65- H7-H9	VL CDR3	aa	LTTSHPVWT	
287	BCMA-29	BC B6 65- H7-H9	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFPPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS	
288	BCMA-29	BC B6 65- H7-H9	VL	aa	DIVMTQTPLSLSPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF SGSGGTDFTLKISRVEAEDVGVYYCLTTSHPVPTFGQGTKLEIK	
289	BCMA-29	BC B6 65- H7-H9	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFPPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGGS DIVMTQTPLSLSPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF SGSGGTDFTLKISRVEAEDVGVYYCLTTSHPVPTFGQGTKLEIK	
290	BCMA-29 HL x CD3 HL	BC B6 65- H7-H9 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFPPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGGS DIVMTQTPLSLSPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF SGSGGTDFTLKISRVEAEDVGVYYCLTTSHPVPTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGG SLKLSCAAAGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGGGGSGGGSGGGGSQTVVTTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGTKFLAPGTPARFSGSLLGGKAALL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL	
291	BCMA-30	BC B6 65- H8-A4	VH CDR1	aa	DYIN	
292	BCMA-30	BC B6 65- H8-A4	VH CDR2	aa	WIYFASGNSEYNQKFTG	
293	BCMA-30	BC B6 65- H8-A4	VH CDR3	aa	LYDYDWYFDV	
294	BCMA-30	BC B6 65- H8-A4	VL CDR1	aa	KSSQSLVHSNGNTYLH	
295	BCMA-30	BC B6 65- H8-A4	VL CDR2	aa	KVSNRFS	
296	BCMA-30	BC B6 65- H8-A4	VL CDR3	aa	AETSHVPWT	
297	BCMA-30	BC B6 65- H8-A4	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFPPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS	
298	BCMA-30	BC B6 65-	VL	aa	DIVMTQTPLSLVTPGEPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF	

	H8-A4				SGSGSGADFTLLKISRVEAEDVGVIYCAETSHVPWTFGQGTKLEIK
299	BCMA-30	BC B6 65-H8-A4	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYINWVRQAPGQGLEWMGWIFYFASGNSEYNQKFTGRVTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSDIVMTQTPLSLSVTPGEPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGADFTLLKISRVEAEDVGVIYCAETSHVPWTFGQGTKLEIK
300	BCMA-30 HL x CD3 HL	BC B6 65-H8-A4 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYINWVRQAPGQGLEWMGWIFYFASGNSEYNQKFTGRVTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSDIVMTQTPLSLSVTPGEPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGADFTLLKISRVEAEDVGVIYCAETSHVPWTFGQGTKLEIK
301	BCMA-31	BC B6 65-H8-H9	VH CDR1	aa	DYYIN
302	BCMA-31	BC B6 65-H8-H9	VH CDR2	aa	WIYFASGNSEYNQKFTG
303	BCMA-31	BC B6 65-H8-H9	VH CDR3	aa	LYDYDWYFDV
304	BCMA-31	BC B6 65-H8-H9	VL CDR1	aa	KSSQSLVHSNGNTYLH
305	BCMA-31	BC B6 65-H8-H9	VL CDR2	aa	KVSNRFS
306	BCMA-31	BC B6 65-H8-H9	VL CDR3	aa	LTTSHPWT
307	BCMA-31	BC B6 65-H8-H9	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYINWVRQAPGQGLEWMGWIFYFASGNSEYNQKFTGRVTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
308	BCMA-31	BC B6 65-H8-H9	VL	aa	DIVMTQTPLSLSVTPGEPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGADFTLLKISRVEAEDVGVIYCLTTSHPWTFGQGTKLEIK
309	BCMA-31	BC B6 65-H8-H9	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYINWVRQAPGQGLEWMGWIFYFASGNSEYNQKFTGRVTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSDIVMTQTPLSLSVTPGEPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGADFTLLKISRVEAEDVGVIYCLTTSHPWTFGQGTKLEIK

310	BCMA-31 HL x CD3 HL	BC B6 65- H8-H9 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSPFDY ^Y INWVRQAPGQGLEWMGW ^I YFASGNSEYNQKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGGS DIVMTQTPLSLSVTPGEPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYK ^V SNRFSGVPDRF SGSGGADFTLKISRVEAEDVGVYYCLT ^T SHVPWTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGG SLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT ^L TVSSGGGGSGGGGSQTVVTQEPSL TVSPGGTVT ^L TCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKKAALT ^L SGVQPEDEAEYYCVLWYSNRWVFGGGTK ^L TVL
311	BCMA-32	BC A7 27- A6-G7	VH CDR1	aa	NHIIH
312	BCMA-32	BC A7 27- A6-G7	VH CDR2	aa	YINPYPGYHAYNEKFQG
313	BCMA-32	BC A7 27- A6-G7	VH CDR3	aa	DGY ^R RD ^T DVLDY
314	BCMA-32	BC A7 27- A6-G7	VL CDR1	aa	QASQDISNYLN
315	BCMA-32	BC A7 27- A6-G7	VL CDR2	aa	YTSRLHT
316	BCMA-32	BC A7 27- A6-G7	VL CDR3	aa	QQGNTLPWT
317	BCMA-32	BC A7 27- A6-G7	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHI ^I HWVRQAPGQGLEWMGYINPYPGYHAYNEKFQGR ATMTSDTSTSTVYMELSSLRSEDTAVYYCARDGY ^R YRDTDVLDYWGQGT ^L TVTVSS
318	BCMA-32	BC A7 27- A6-G7	VL	aa	DIQMTQSPSSLSASLGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSRFSGSGS GTDFTFTISSLQQEDIA ^T Y ^C QQGNTLPWTFGQGTKVEIK
319	BCMA-32	BC A7 27- A6-G7	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHI ^I HWVRQAPGQGLEWMGYINPYPGYHAYNEKFQGR ATMTSDTSTSTVYMELSSLRSEDTAVYYCARDGY ^R YRDTDVLDYWGQGT ^L TVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASLGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSRFSGS GSGTDFFTISSLQQEDIA ^T Y ^C QQGNTLPWTFGQGTKVEIK
320	BCMA-32 HL x CD3 HL	BC A7 27- A6-G7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHI ^I HWVRQAPGQGLEWMGYINPYPGYHAYNEKFQGR ATMTSDTSTSTVYMELSSLRSEDTAVYYCARDGY ^R YRDTDVLDYWGQGT ^L TVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASLGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSRFSGS GSGTDFFTISSLQQEDIA ^T Y ^C QQGNTLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN

					NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGGGGGSGQT VVTQEPSLTVS PGGTVTLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
321	BCMA-33	BC A7 27- A6-H11	VH CDR1	aa	NHIIH
322	BCMA-33	BC A7 27- A6-H11	VH CDR2	aa	YINPYDGGWDYNEKFQG
323	BCMA-33	BC A7 27- A6-H11	VH CDR3	aa	DGYIRDADVLDY
324	BCMA-33	BC A7 27- A6-H11	VL CDR1	aa	QASQDISNYLN
325	BCMA-33	BC A7 27- A6-H11	VL CDR2	aa	YTSRLHT
326	BCMA-33	BC A7 27- A6-H11	VL CDR3	aa	QQGNTLPWT
327	BCMA-33	BC A7 27- A6-H11	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHIIHWVRQAPGQGLEWMGYINPYDGGWDYNEKFQGR ATMTSDTSTVYMELSSLRSEDTAVYYCARDGY YRDADVLDYWGQGLTVTVSS
328	BCMA-33	BC A7 27- A6-H11	VL	aa	DIQMTQSPSSLSASLGDRVITITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSPRFSGSGS GTDFTFTISSLQQEDIAITYYCQQGNTLPWTFGQGTKVEIK
329	BCMA-33	BC A7 27- A6-H11	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHIIHWVRQAPGQGLEWMGYINPYDGGWDYNEKFQGR ATMTSDTSTVYMELSSLRSEDTAVYYCARDGY YRDADVLDYWGQGLTVTVSSGGGGGGSGGGG GSDIQMTQSPSSLSASLGDRVITITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSPRFSGSGS GGTDFTFTISSLQQEDIAITYYCQQGNTLPWTFGQGTKVEIK
330	BCMA-33 HL x CD3 HL	BC A7 27- A6-H11 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHIIHWVRQAPGQGLEWMGYINPYDGGWDYNEKFQGR ATMTSDTSTVYMELSSLRSEDTAVYYCARDGY YRDADVLDYWGQGLTVTVSSGGGGGGSGGGG GSDIQMTQSPSSLSASLGDRVITITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSPRFSGSGS GGTDFTFTISSLQQEDIAITYYCQQGNTLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGGGGGSGQT VVTQEPSLTVS PGGTVTLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
331	BCMA-34	BC A7 27- C4-G7	VH CDR1	aa	NHIIH

332	BCMA-34	BC A7 27- C4-G7	VH CDR2	aa	YINPYPGYHAYNEKFQG
333	BCMA-34	BC A7 27- C4-G7	VH CDR3	aa	DGYRDTDVLDY
334	BCMA-34	BC A7 27- C4-G7	VL CDR1	aa	QASQDISNYLN
335	BCMA-34	BC A7 27- C4-G7	VL CDR2	aa	YTSRLHT
336	BCMA-34	BC A7 27- C4-G7	VL CDR3	aa	QQGNTLPWT
337	BCMA-34	BC A7 27- C4-G7	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHIIHWVRQAPGQGLEWMGYINPYPGYHAYNEKFQGR ATMTSDTSTVYMELSSLRSEDTAVYYCARDGYRDTDVLDYWGQGLVTVSSGGGGSGGG ATMTSDTSTVYMELSSLRSEDTAVYYCARDGYRDTDVLDYWGQGLVTVSS
338	BCMA-34	BC A7 27- C4-G7	VL	aa	DIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSRFSGSGS GTDFTFTISSLEPEDIAITYYCQQGNTLPWTFGQGTKVEIK
339	BCMA-34	BC A7 27- C4-G7	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHIIHWVRQAPGQGLEWMGYINPYPGYHAYNEKFQGR ATMTSDTSTVYMELSSLRSEDTAVYYCARDGYRDTDVLDYWGQGLVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSRFSGSGS GSGTDFTTISSLEPEDIAITYYCQQGNTLPWTFGQGTKVEIK
340	BCMA-34 HL x CD3 HL	BC A7 27- C4-G7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHIIHWVRQAPGQGLEWMGYINPYPGYHAYNEKFQGR ATMTSDTSTVYMELSSLRSEDTAVYYCARDGYRDTDVLDYWGQGLVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSRFSGSGS GSGTDFTTISSLEPEDIAITYYCQQGNTLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGGSGQT VVTQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
341	BCMA-35	BC A7 27- C4-H11	VH CDR1	aa	NHIIH
342	BCMA-35	BC A7 27- C4-H11	VH CDR2	aa	YINPYDGWGDYNEKFQG
343	BCMA-35	BC A7 27- C4-H11	VH CDR3	aa	DGYRDADVLDY
344	BCMA-35	BC A7 27- C4-H11	VL CDR1	aa	QASQDISNYLN

345	BCMA-35	BC A7 27- C4-H11	VL CDR2	aa	YTSRLHT	
346	BCMA-35	BC A7 27- C4-H11	VL CDR3	aa	QQGNLTLPWT	
347	BCMA-35	BC A7 27- C4-H11	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHIIHWVRQAPGQGLEWMGYINPYDGWGDYNEKFQGR ATMTSDTSTSTVYMELSSLRSEDTAVYYCARDGYIRDADVLDYWGQGLTVTVSS	
348	BCMA-35	BC A7 27- C4-H11	VL	aa	DIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPKGKAPKLLIYYTSTRLLHTGVPSRFSGSGS GTDFTFTISSLEPEDIAITYYCQQGNLTLPWTFGQGTKVEIK	
349	BCMA-35	BC A7 27- C4-H11	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHIIHWVRQAPGQGLEWMGYINPYDGWGDYNEKFQGR ATMTSDTSTSTVYMELSSLRSEDTAVYYCARDGYIRDADVLDYWGQGLTVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPKGKAPKLLIYYTSTRLLHTGVPSRFSGSGS GSGTDFTFTISSLEPEDIAITYYCQQGNLTLPWTFGQGTKVEIK	
350	BCMA-35 HL x CD3 HL	BC A7 27- C4-H11 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFTNHIIHWVRQAPGQGLEWMGYINPYDGWGDYNEKFQGR ATMTSDTSTSTVYMELSSLRSEDTAVYYCARDGYIRDADVLDYWGQGLTVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPKGKAPKLLIYYTSTRLLHTGVPSRFSGSGS GSGTDFTFTISSLEPEDIAITYYCQQGNLTLPWTFGQGTKVEIKSGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGSGGGQT VVTQEPSLTVS PGGT VTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL	
351	BCMA-36	BC A7 15- H2-G7	VH CDR1	aa	NHIIH	
352	BCMA-36	BC A7 15- H2-G7	VH CDR2	aa	YINPYPGYHAYNQKFQG	
353	BCMA-36	BC A7 15- H2-G7	VH CDR3	aa	DGY YRDTDVLDY	
354	BCMA-36	BC A7 15- H2-G7	VL CDR1	aa	QASQDISNYLN	
355	BCMA-36	BC A7 15- H2-G7	VL CDR2	aa	YTSRLHT	
356	BCMA-36	BC A7 15- H2-G7	VL CDR3	aa	QQGNLTLPWT	
357	BCMA-36	BC A7 15- H2-G7	VH	aa	QVQLVQSGAKVIKPGASVKVSCKASGYTFTNHIIHWVRQKPGQGLEWMGYINPYPGYHAYNQKFQGR	

						VTMTRDKSTSTVYMELSSLTSED	TAVYYCARDGY	YRDTDVLDYWGQ	TLVTVSS															
358	BCMA-36	BC A7 15- H2-G7	VL	aa		DIQMTQSPSSLSASVGDRVTITCQASQDI	SNYLNWYQQKPGRAPKLLIYYTSRLHTGVP	SRFSGSGS	GTDYSFTISSLQPED	IATYYCQQGNTLPWTFGQGTKVEIK														
359	BCMA-36	BC A7 15- H2-G7	scFv	aa		QVQLVQSGAKVIKPGASVKV	SCKASGYTFTNHI	IHWVRQKPGQGLEWMGYINPY	PGYHAYNQKFQGR	VTMTRDKSTSTVYMELSSLTSED	TAVYYCARDGY	YRDTDVLDYWGQ	TLVTVSSGGSGGGSGGG	GSDIQMTQSPSSLSASVGDRVTITCQASQDI	SNYLNWYQQKPGRAPKLLIYYTSRLHTGVP	SRFSGSGS	GGTDYSFTISSLQPED	IATYYCQQGNTLPWTFGQGTKVEIK						
360	BCMA-36 HL x CD3 HL	BC A7 15- H2-G7 HL x CD3 HL	bispecific molecule	aa		QVQLVQSGAKVIKPGASVKV	SCKASGYTFTNHI	IHWVRQKPGQGLEWMGYINPY	PGYHAYNQKFQGR	VTMTRDKSTSTVYMELSSLTSED	TAVYYCARDGY	YRDTDVLDYWGQ	TLVTVSSGGSGGGSGGG	GSDIQMTQSPSSLSASVGDRVTITCQASQDI	SNYLNWYQQKPGRAPKLLIYYTSRLHTGVP	SRFSGSGS	GGTDYSFTISSLQPED	IATYYCQQGNTLPWTFGQGTKVEIK						
						LSCAASGFTFNKYAMN	WVRQAPGKGLEW	VARIRSKYNNYATYYADSVKDRFTISR	DDSKNTAYLQMN	NLKTEDTAVYYCVRHGN	FGNSYISYWAYWGQ	TLVTVSSGGSGGGSGG	GSQT	VVTQEP	SLTVS	PGGT	VTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGG	TKFLAPGTPARFSGS	LLGGKAALTLSGV	QPEDEAEY	YCVLWYSNRWVFGG	TKLTVL		
361	BCMA-37	BC A7 15- H2-H11	VH CDR1	aa																				
362	BCMA-37	BC A7 15- H2-H11	VH CDR2	aa																				
363	BCMA-37	BC A7 15- H2-H11	VH CDR3	aa																				
364	BCMA-37	BC A7 15- H2-H11	VL CDR1	aa																				
365	BCMA-37	BC A7 15- H2-H11	VL CDR2	aa																				
366	BCMA-37	BC A7 15- H2-H11	VL CDR3	aa																				
367	BCMA-37	BC A7 15- H2-H11	VH	aa																				
368	BCMA-37	BC A7 15- H2-H11	VL	aa																				
369	BCMA-37	BC A7 15- H2-H11	scFv	aa																				

					VTMTRDKSTSTVYMELSSLTSED TAVYYCARDGYIRDADVLDYWGQGT LVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGD RVTITCQASQDISNYLNWYQQK PGRAPKLLIYYTSRLHTGVP SRFSGSGGTDYSFTISSLQPE DIATYYCQQGNTLPWTFGQGT KVEIK
370	BCMA-37 HL x CD3 HL	BC A7 15- H2-H11 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVIKPGASVKV SCKASGYTFTNHIIHWVRQK PGQGLEWMGYINPYDGGDYNQ KFQGRVTMTRDKSTSTVYMEL SSLTSED TAVYYCARDGYIRDADVLDYWG QGT LVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGD RVTITCQASQDISNYLNWYQQK PGRAPKLLIYYTSRLHTGVP SRFSGSGGTDYSFTISSLQPE DIATYYCQQGNTLPWTFGQGT KVEIKLSCAASGFTFNKYAMN WVRQAPGKGLEWVARIRSKYNN YATYYADSVKDRFTISRDDSK NTAYLQMN NLKTEDTAVYYCVRHGNFGNS YISYWAYWGQGT LVTVSSGGGGSGGGSGGGQ T VVTQEPSLT VSPGGTVTLTCGSSSTGAVT SGNYPNWVQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKA ALTLSGVQPEDEAEYCYVLWY SNRWVFGGGTKLTVL
371	BCMA-38	BC A7 15- H8-G7	VH CDR1	aa	NHIIH
372	BCMA-38	BC A7 15- H8-G7	VH CDR2	aa	YINPYPGYHAYNQKFQG
373	BCMA-38	BC A7 15- H8-G7	VH CDR3	aa	DGY YRDTDVLDY
374	BCMA-38	BC A7 15- H8-G7	VL CDR1	aa	QASQDISNYLN
375	BCMA-38	BC A7 15- H8-G7	VL CDR2	aa	YTSRLHT
376	BCMA-38	BC A7 15- H8-G7	VL CDR3	aa	QQGNTLPWT
377	BCMA-38	BC A7 15- H8-G7	VH	aa	QVQLVQSGAEVIKPGASVKV SCKASGYTFTNHIIHWVRQK PGQGLEWIGYINPYGYHAYN QKFQGV VTMTRD TSTSTVYMELSSLTSED TAVYYCARDGY YRDTDVLDY WGQGT LVTVSS
378	BCMA-38	BC A7 15- H8-G7	VL	aa	DIQMTQSPSSLSASLGDRVT ITCQASQDISNYLNWYQQK PGKAPKLLIYYTSRLHTGVP SRFSGSGSDFTFTISSLQQE DIATYYCQQGNTLPWTFGQGT KVEIK
379	BCMA-38	BC A7 15- H8-G7	scFv	aa	QVQLVQSGAEVIKPGASVKV SCKASGYTFTNHIIHWVRQK PGQGLEWIGYINPYGYHAYN QKFQGV VTMTRD TSTSTVYMELSSLTSED TAVYYCARDGY YRDTDVLDY WGQGT LVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASLGDRVT ITCQASQDISNYLNWYQQK PGKAPKLLIYYTSRLHTGVP SRFSGSGGTDFTFTISSLQQE DIATYYCQQGNTLPWTFGQGT KVEIK
380	BCMA-38 HL x CD3 HL	BC A7 15- H8-G7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVIKPGASVKV SCKASGYTFTNHIIHWVRQK PGQGLEWIGYINPYGYHAYN QKFQGV VTMTRD TSTSTVYMELSSLTSED TAVYYCARDGY YRDTDVLDY WGQGT LVTVSSGGGGSGGGSGGG

					GSDIQMTQSPSSLSASLGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSPRFSGS GSGTDFTTISSLQQEDIAYYCQQGNTLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGQTVVTVTQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWWVQOKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
381	BCMA-39	BC A7 15- H8-H11	VH CDR1	aa	NHIIH
382	BCMA-39	BC A7 15- H8-H11	VH CDR2	aa	YINPYDGGDYNQKFQG
383	BCMA-39	BC A7 15- H8-H11	VH CDR3	aa	DGYRRDADVLDY
384	BCMA-39	BC A7 15- H8-H11	VL CDR1	aa	QASQDISNYLN
385	BCMA-39	BC A7 15- H8-H11	VL CDR2	aa	YTSRLHT
386	BCMA-39	BC A7 15- H8-H11	VL CDR3	aa	QQGNTLPWT
387	BCMA-39	BC A7 15- H8-H11	VH	aa	QVQLVQSGAEVIKPGASVKVSKKASGYTFTNHIIHWVRQKPGQGLEWIGYINPYDGGDYNQKFQ GK VTMTRDTSTVYMELSSLTSEDTAVYYCARDGYIRDADVLDYWGQGLTVTVSSGGGGSGGGG VTMTRDTSTVYMELSSLTSEDTAVYYCARDGYIRDADVLDYWGQGLTVTVSSGGGGSGGGG
388	BCMA-39	BC A7 15- H8-H11	VL	aa	DIQMTQSPSSLSASLGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSPRFSGS GTDFTFTISSLQQEDIAYYCQQGNTLPWTFGQGTKVEIK
389	BCMA-39	BC A7 15- H8-H11	scFv	aa	QVQLVQSGAEVIKPGASVKVSKKASGYTFTNHIIHWVRQKPGQGLEWIGYINPYDGGDYNQKFQ GK VTMTRDTSTVYMELSSLTSEDTAVYYCARDGYIRDADVLDYWGQGLTVTVSSGGGGSGGGG VTMTRDTSTVYMELSSLTSEDTAVYYCARDGYIRDADVLDYWGQGLTVTVSSGGGGSGGGG
390	BCMA-39 HL x CD3 HL	BC A7 15- H8-H11 HL x CD3 HL	bispecific molecule	aa	GSDIQMTQSPSSLSASLGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYYTSRLHTGVPSPRFSGS GSGTDFTTISSLQQEDIAYYCQQGNTLPWTFGQGTKVEIK LSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGQTVVTVTQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWWVQOKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV

						QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
391	BCMA-40	BC 7A4 96-D4-A12	VH CDR1	aa		DYYIN
392	BCMA-40	BC 7A4 96-D4-A12	VH CDR2	aa		WIYFASGNSEYNQKFTG
393	BCMA-40	BC 7A4 96-D4-A12	VH CDR3	aa		LYDYDWYFDV
394	BCMA-40	BC 7A4 96-D4-A12	VL CDR1	aa		KSSQSLVHSNGNTYLH
395	BCMA-40	BC 7A4 96-D4-A12	VL CDR2	aa		KVSNRFS
396	BCMA-40	BC 7A4 96-D4-A12	VL CDR3	aa		SQSSTAPWT
397	BCMA-40	BC 7A4 96-D4-A12	VH	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
398	BCMA-40	BC 7A4 96-D4-A12	VL	aa		DI VMTQTPLSLPVTLGQPASISCKSSQSLVHSNGNTYLHWY LQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSSTAPWTFGQGTKLEIK
399	BCMA-40	BC 7A4 96-D4-A12	scFv	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGSGSDI VMTQTPLSLPVTLGQPASISCKSSQSLVHSNGNTYLHWY LQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSSTAPWTFGQGTKLEIK
400	BCMA-40 HL x CD3 HL	BC 7A4 96-D4-A12 HL x CD3 HL	bispecific molecule	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGSGSDI VMTQTPLSLPVTLGQPASISCKSSQSLVHSNGNTYLHWY LQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSSTAPWTFGQGTKLEIK
401	BCMA-41	BC 7A4 96-D4-D7	VH CDR1	aa		DYYIN
402	BCMA-41	BC 7A4 96-D4-D7	VH CDR2	aa		WIYFASGNSEYNQKFTG

403	BCMA-41	BC 7A4 96-D4-D7	VH CDR3	aa	LYDYDWYFDV
404	BCMA-41	BC 7A4 96-D4-D7	VL CDR1	aa	KSSQSLVHSNGNTYLH
405	BCMA-41	BC 7A4 96-D4-D7	VL CDR2	aa	KVSNRFS
406	BCMA-41	BC 7A4 96-D4-D7	VL CDR3	aa	SQSSIIYPWT
407	BCMA-41	BC 7A4 96-D4-D7	VH	aa	QVQLVQSGAEVKKPGASVKVCKASGYSPFDYYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTSS
408	BCMA-41	BC 7A4 96-D4-D7	VL	aa	DIVMTQTPLSLPVTLGQPASISCKSSQSLVHSNGNTYLVHWYLQKPGQSPQLLIYKVSNRFSGVDPDRFSGSGSGTDFTLKISRVEAEDVGYVYCSQSSIIYPWTFGQGTKLEIK
409	BCMA-41	BC 7A4 96-D4-D7	scFv	aa	QVQLVQSGAEVKKPGASVKVCKASGYSPFDYYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTSSGGGGSGGGSGGGSGS DIVMTQTPLSLPVTLGQPASISCKSSQSLVHSNGNTYLVHWYLQKPGQSPQLLIYKVSNRFSGVDPDRFSGSGSGTDFTLKISRVEAEDVGYVYCSQSSIIYPWTFGQGTKLEIK
410	BCMA-41 HL x CD3 HL	BC 7A4 96-D4-D7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVCKASGYSPFDYYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTSSGGGGSGGGSGGGSGS DIVMTQTPLSLPVTLGQPASISCKSSQSLVHSNGNTYLVHWYLQKPGQSPQLLIYKVSNRFSGVDPDRFSGSGSGTDFTLKISRVEAEDVGYVYCSQSSIIYPWTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLVQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSGGGSQTVVTQEPSLTVSPGGTFTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
411	BCMA-42	BC 7A4 96-D4-E7	VH CDR1	aa	DYIIN
412	BCMA-42	BC 7A4 96-D4-E7	VH CDR2	aa	WIYFASGNSEYNQKFTG
413	BCMA-42	BC 7A4 96-D4-E7	VH CDR3	aa	LYDYDWYFDV
414	BCMA-42	BC 7A4 96-D4-E7	VL CDR1	aa	KSSQSLVHSNGNTYLH
415	BCMA-42	BC 7A4 96-D4-E7	VL CDR2	aa	KVSNRFS

416	BCMA-42	BC 7A4 96-D4-E7	VL CDR3	aa	SQSTYPEFT	
417	BCMA-42	BC 7A4 96-D4-E7	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFPPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS	
418	BCMA-42	BC 7A4 96-D4-E7	VL	aa	DIVMTQTPLSLPVTLLGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQQLLIYKVSNRFSGVPPDRFSGSGGTDFTLKISRVEAEDVGVYYCSQSTYPEFTFGQGTKLEIK	
419	BCMA-42	BC 7A4 96-D4-E7	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFPPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSGGGSDIVMTQTPLSLPVTLLGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQQLLIYKVSNRFSGVPPDRFSGSGGTDFTLKISRVEAEDVGVYYCSQSTYPEFTFGQGTKLEIK	
420	BCMA-42 HL x CD3 HL	BC 7A4 96-D4-E7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFPPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSGGGSDIVMTQTPLSLPVTLLGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQQLLIYKVSNRFSGVPPDRFSGSGGTDFTLKISRVEAEDVGVYYCSQSTYPEFTFGQGTKLEIKSLEKSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAWWGQGTLLTVSSGGGGSGGGSGGGSGQTVVTTQEPSSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALLTSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL	
421	BCMA-43	BC 7A4 96-F4-A12	VH CDR1	aa	DYIN	
422	BCMA-43	BC 7A4 96-F4-A12	VH CDR2	aa	WIYFASGNSEYNQKFTG	
423	BCMA-43	BC 7A4 96-F4-A12	VH CDR3	aa	LYDYDWYFDV	
424	BCMA-43	BC 7A4 96-F4-A12	VL CDR1	aa	KSSQSLVHSNGNTYLH	
425	BCMA-43	BC 7A4 96-F4-A12	VL CDR2	aa	KVSNRFS	
426	BCMA-43	BC 7A4 96-F4-A12	VL CDR3	aa	SQSSTAPWT	
427	BCMA-43	BC 7A4 96-F4-A12	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFPPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS	
428	BCMA-43	BC 7A4 96-	VL	aa	DIVMTQTPLSLVTPGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQQLLIYKVSNRFSGVPPDRF	

		F4-A12				SGSGSGTDFTLKISRVEAEDVGVYYCSQSSTAPWTFGQGTKLEIK
429	BCMA-43	BC 7A4 96-F4-A12	scFv	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSDIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYILHWYLQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSSTAPWTFGQGTKLEIK
430	BCMA-43 HL x CD3 HL	BC 7A4 96-F4-A12 HL x CD3 HL	bispecific molecule	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSDIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYILHWYLQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSSTAPWTFGQGTKLEIK
431	BCMA-44	BC 7A4 96-F4-D7	VH CDR1	aa		DYYIN
432	BCMA-44	BC 7A4 96-F4-D7	VH CDR2	aa		WIIYFASGNSEYNQKFTG
433	BCMA-44	BC 7A4 96-F4-D7	VH CDR3	aa		LYDYYDWYFDV
434	BCMA-44	BC 7A4 96-F4-D7	VL CDR1	aa		KSSQSLVHSNGNTYLH
435	BCMA-44	BC 7A4 96-F4-D7	VL CDR2	aa		KVSNRFS
436	BCMA-44	BC 7A4 96-F4-D7	VL CDR3	aa		SQSSIYPWT
437	BCMA-44	BC 7A4 96-F4-D7	VH	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
438	BCMA-44	BC 7A4 96-F4-D7	VL	aa		DIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYILHWYLQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSIIYPWTFGQGTKLEIK
439	BCMA-44	BC 7A4 96-F4-D7	scFv	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSPFDYYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSDIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYILHWYLQKPGQSPQLLIYKVSNRFSGVPPDRFSGSGSGTDFTLKISRVEAEDVGVYYCSQSIIYPWTFGQGTKLEIK

440	BCMA-44 HL x CD3 HL	BC 7A4 96-F4-D7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSDIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLVQKPGQSPQQLLIYKVSNRFSGVPPDRFSGSGGTDFTLKISRVEAEDVGVYCSQSSIYPWTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSGQTVVTTQEPSSLTVSPGGTVTLLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKKAALLTSGVQPEDEAEYICVLWYSNRWVFGGGTKLTVL
441	BCMA-45	BC 7A4 96-F4-E7	VH CDR1	aa	DYIIN
442	BCMA-45	BC 7A4 96-F4-E7	VH CDR2	aa	WIYFASGNSEYNQKFTG
443	BCMA-45	BC 7A4 96-F4-E7	VH CDR3	aa	LYDYDWYFDV
444	BCMA-45	BC 7A4 96-F4-E7	VL CDR1	aa	KSSQSLVHSNGNTYLH
445	BCMA-45	BC 7A4 96-F4-E7	VL CDR2	aa	KVSNRFS
446	BCMA-45	BC 7A4 96-F4-E7	VL CDR3	aa	SQSTYPEFT
447	BCMA-45	BC 7A4 96-F4-E7	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
448	BCMA-45	BC 7A4 96-F4-E7	VL	aa	DIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLVQKPGQSPQQLLIYKVSNRFSGVPPDRFSGSGGTDFTLKISRVEAEDVGVYCSQSTYPEFTFGQGTKLEIK
449	BCMA-45	BC 7A4 96-F4-E7	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSDIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLVQKPGQSPQQLLIYKVSNRFSGVPPDRFSGSGGTDFTLKISRVEAEDVGVYCSQSTYPEFTFGQGTKLEIK
450	BCMA-45 HL x CD3 HL	BC 7A4 96-F4-E7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNQKFTGRVTMTRDTSISTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGSGGGSDIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWYLVQKPGQSPQQLLIYKVSNRFSGVPPDRFSGSGGTDFTLKISRVEAEDVGVYCSQSTYPEFTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYL

						QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGGGSGQTVVVTQEP SLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLG GKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
451	BCMA-46	BC 7A4 96- G2-A12	VH CDR1	aa		DYYIN
452	BCMA-46	BC 7A4 96- G2-A12	VH CDR2	aa		WIYFASGNSEYNEKFTG
453	BCMA-46	BC 7A4 96- G2-A12	VH CDR3	aa		LYDYDWYFDV
454	BCMA-46	BC 7A4 96- G2-A12	VL CDR1	aa		KSSQSLVHSNGNTYLH
455	BCMA-46	BC 7A4 96- G2-A12	VL CDR2	aa		KVSNRFS
456	BCMA-46	BC 7A4 96- G2-A12	VL CDR3	aa		SQSSSTAPWT
457	BCMA-46	BC 7A4 96- G2-A12	VH	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSEFPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNEKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
458	BCMA-46	BC 7A4 96- G2-A12	VL	aa		DIVMTQTPLSLSVSLGQPASISCKSSQSLVHSNGNTYLHWYILQKPGQSPQLLIYKVS NRFSGVPDFRSGSGSGTDFTTLKISRVEAEDVGVYYCSQSSSTAPWTFGQGTKLEIK
459	BCMA-46	BC 7A4 96- G2-A12	scFv	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSEFPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNEKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGGGGS DIVMTQTPLSLSVSLGQPASISCKSSQSLVHSNGNTYLHWYILQKPGQSPQLLIYKVS NRFSGVPDFRSGSGSGTDFTTLKISRVEAEDVGVYYCSQSSSTAPWTFGQGTKLEIK
460	BCMA-46 HL x CD3 HL	BC 7A4 96- G2-A12 HL x CD3 HL	bispecific molecule	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSEFPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNEKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGGSGGGGGGS DIVMTQTPLSLSVSLGQPASISCKSSQSLVHSNGNTYLHWYILQKPGQSPQLLIYKVS NRFSGVPDFRSGSGSGTDFTTLKISRVEAEDVGVYYCSQSSSTAPWTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGG SLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGGGSGQTVVVTQEP SLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLG GKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
461	BCMA-47	BC 7A4 96- G2-D7	VH CDR1	aa		DYYIN

462	BCMA-47	BC 7A4 96- G2-D7	VH CDR2	aa	WIYFASGNSEYNEKFTG
463	BCMA-47	BC 7A4 96- G2-D7	VH CDR3	aa	LYDYDWYFDV
464	BCMA-47	BC 7A4 96- G2-D7	VL CDR1	aa	KSSQSLVHSNGNTYLH
465	BCMA-47	BC 7A4 96- G2-D7	VL CDR2	aa	KVSNRFS
466	BCMA-47	BC 7A4 96- G2-D7	VL CDR3	aa	SQSSIYPWT
467	BCMA-47	BC 7A4 96- G2-D7	VH	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNEKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS
468	BCMA-47	BC 7A4 96- G2-D7	VL	aa	DIVMTQTPLSLSVSLGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVYYCSQSSIYPWTFGQGTKLEIK
469	BCMA-47	BC 7A4 96- G2-D7	scFv	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNEKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGGS DIVMTQTPLSLSVSLGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVYYCSQSSIYPWTFGQGTKLEIK
470	BCMA-47 HL x CD3 HL	BC 7A4 96- G2-D7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSCKASGYSFDPDYINWVRQAPGQGLEWMGWIIYFASGNSEYNEKFTGR VTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGSGGGSGGGGS DIVMTQTPLSLSVSLGQPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNRFSGVPPDRF SGSGSGTDFTLKISRVEAEDVGVYYCSQSSIYPWTFGQGTKLEIKSFGGGSEVQLVESGGGLVQPGG SLKLSCAAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLVTVSSGGSGGGSGGGGSQTVVVTQEPSL TVSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKKAALLTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
471	BCMA-48	BC 7A4 96- G2-E7	VH CDR1	aa	DYYIN
472	BCMA-48	BC 7A4 96- G2-E7	VH CDR2	aa	WIYFASGNSEYNEKFTG
473	BCMA-48	BC 7A4 96- G2-E7	VH CDR3	aa	LYDYDWYFDV
474	BCMA-48	BC 7A4 96- G2-E7	VL CDR1	aa	KSSQSLVHSNGNTYLH

475	BCMA-48	BC 7A4 96- G2-E7	VL CDR2	aa	KVSNRFS	
476	BCMA-48	BC 7A4 96- G2-E7	VL CDR3	aa	SQSTYPEFT	
477	BCMA-48	BC 7A4 96- G2-E7	VH	aa	QVQLVQSGAEVKKPGASVKVCKASGYSPFDY _Y INWVRQAPGQGLEWMGW _{IY} FASGNSEYNEKFTGR VTMTRDTSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSS	
478	BCMA-48	BC 7A4 96- G2-E7	VL	aa	DIVMTQTPLSLSVSLGQPASISCKSSQSLVHSNGNTY _L LHWY _L LQKPGQSPQ _{LL} IYKVSNRFSGVPDRF SGSGGTDFTLKISRVEAEDVGVY _Y CSQSTYPEFTFGQGTKLEIK	
479	BCMA-48	BC 7A4 96- G2-E7	scFv	aa	QVQLVQSGAEVKKPGASVKVCKASGYSPFDY _Y INWVRQAPGQGLEWMGW _{IY} FASGNSEYNEKFTGR VTMTRDTSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGSGGGSGGGGS DIVMTQTPLSLSVSLGQPASISCKSSQSLVHSNGNTY _L LHWY _L LQKPGQSPQ _{LL} IYKVSNRFSGVPDRF SGSGGTDFTLKISRVEAEDVGVY _Y CSQSTYPEFTFGQGTKLEIK	
480	BCMA-48 HL x CD3 HL	BC 7A4 96- G2-E7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVCKASGYSPFDY _Y INWVRQAPGQGLEWMGW _{IY} FASGNSEYNEKFTGR VTMTRDTSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTVSSGGGSGGGSGGGGS DIVMTQTPLSLSVSLGQPASISCKSSQSLVHSNGNTY _L LHWY _L LQKPGQSPQ _{LL} IYKVSNRFSGVPDRF SGSGGTDFTLKISRVEAEDVGVY _Y CSQSTYPEFTFGQGTKLEIKSGGGSEVQLVESGGLVQPGG SLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATY _Y ADSVKDRFTISRDDSKNTAY _L QMNNLKTEDTAVY _Y CVRHGNFGNSY _I SYWAYWGQGT _L TVSSGGGSGGGSGGGSQTVVTTQEP _{SL} TVSPGGTV _{TL} TCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKAA _{LTL} SGVQPEDEAEY _Y CVLWY _Y SNRWVFGGTK _{LTVL}	
481	BCMA-49	BC 7A4 97- A3-A12	VH CDR1	aa	DY _Y IN	
482	BCMA-49	BC 7A4 97- A3-A12	VH CDR2	aa	WIYFASGNSEYNQKFTG	
483	BCMA-49	BC 7A4 97- A3-A12	VH CDR3	aa	LYDYDWYFDV	
484	BCMA-49	BC 7A4 97- A3-A12	VL CDR1	aa	KSSQSLVHSNGNTY _L H	
485	BCMA-49	BC 7A4 97- A3-A12	VL CDR2	aa	KVSNRFS	
486	BCMA-49	BC 7A4 97- A3-A12	VL CDR3	aa	SQSSAPWT	
487	BCMA-49	BC 7A4 97- A3-A12	VH	aa	QVQLVQSGAEVKKPGASVKVCKASGYSPFDY _Y INWVRQAPGQGLEWMGW _{IY} FASGNSEYNQKFTGR	

					VTMTRDTSINTAYMELSSLTSED	TAVYFCASLYDYDWYFDVWGQGTMTVTVSS			VTMTRDTSINTAYMELSSLTSED	TAVYFCASLYDYDWYFDVWGQGTMTVTVSS	
488	BCMA-49	BC 7A4 97-A3-A12	VL	aa		DIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWY LQKPGQSPQLLIYKVS	NRFS	GV	PD	RF	
489	BCMA-49	BC 7A4 97-A3-A12	scFv	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSF	FPDYYINWVRQAPGQGLEWMGWIYFAS	GN	SE	YNQKFTGR	
490	BCMA-49 HL x CD3 HL	BC 7A4 97-A3-A12 HL x CD3 HL	bispecific molecule	aa		VTMTRDTSINTAYMELSSLTSED	TAVYFCASLYDYDWYFDVWGQGTMTVTVSS	GGSGGGSGGGSGGGGS			
491	BCMA-50	BC 7A4 97-A3-D7	VH CDR1	aa		DY	YIN				
492	BCMA-50	BC 7A4 97-A3-D7	VH CDR2	aa		WIYFASGNSEYNQKFTG					
493	BCMA-50	BC 7A4 97-A3-D7	VH CDR3	aa		LYDYDWYFDV					
494	BCMA-50	BC 7A4 97-A3-D7	VL CDR1	aa		KSSQSLVHSNGNTYLH					
495	BCMA-50	BC 7A4 97-A3-D7	VL CDR2	aa		KV	SNRFS				
496	BCMA-50	BC 7A4 97-A3-D7	VL CDR3	aa		SQSSIYPWT					
497	BCMA-50	BC 7A4 97-A3-D7	VH	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSF	FPDYYINWVRQAPGQGLEWMGWIYFAS	GN	SE	YNQKFTGR	
498	BCMA-50	BC 7A4 97-A3-D7	VL	aa		DIVMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWY LQKPGQSPQLLIYKVS	NRFS	GV	PD	RF	
499	BCMA-50	BC 7A4 97-A3-D7	scFv	aa		QVQLVQSGAEVKKPGASVKVSCKASGYSF	FPDYYINWVRQAPGQGLEWMGWIYFAS	GN	SE	YNQKFTGR	

					VTMTRDTSINTAYMELSSLTSED TAVYFCASLYDYDWYFDVWGQGT MTVTVSSGGGGSGGGSGGGGS DIVMTQTPLSLVTPGQPASISCKSSQSLVHSNGNTY LHWYLQKPGQSPQLLIYKVS NRFSGVVPDRFSGSGGTDFTLKISRVEAEDVGIY YCSQSSIYPWTFGQGTKLEIK
500	BCMA-50 HL x CD3 HL	BC 7A4 97- A3-D7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSC KASGYSFPDYYINWVRQAPGQ GLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSINTAYMELSSLTSED TAVYFCASLYDYDWYFDVWGQGT MTVTVSSGGGGSGGGSGGGGS DIVMTQTPLSLVTPGQPASISCKSSQSLVHSNGNTY LHWYLQKPGQSPQLLIYKVS NRFSGVVPDRFSGSGGTDFTLKISRVEAEDVGIY YCSQSSIYPWTFGQGTKLEIKS GGGSEVQLVESGGGLVQPPGG SLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIR SKYNNYATYYADSVKDRFTISRDDSKNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVTVSSGGGGSGGGSGGGSTVVTQE PSLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQ QKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAA LTL SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
501	BCMA-51	BC 7A4 97- A3-E7	VH CDR1	aa	DYYIN
502	BCMA-51	BC 7A4 97- A3-E7	VH CDR2	aa	WIYFASGNSEYNQKFTG
503	BCMA-51	BC 7A4 97- A3-E7	VH CDR3	aa	LYDYDWYFDV
504	BCMA-51	BC 7A4 97- A3-E7	VL CDR1	aa	KSSQSLVHSNGNTYLH
505	BCMA-51	BC 7A4 97- A3-E7	VL CDR2	aa	KVSNRFS
506	BCMA-51	BC 7A4 97- A3-E7	VL CDR3	aa	SQSTYPEFT
507	BCMA-51	BC 7A4 97- A3-E7	VH	aa	QVQLVQSGAEVKKPGASVKVSC KASGYSFPDYYINWVRQAPGQ GLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSINTAYMELSSLTSED TAVYFCASLYDYDWYFDVWGQGT MTVTVSS
508	BCMA-51	BC 7A4 97- A3-E7	VL	aa	DIVMTQTPLSLVTPGQPASISCKSSQSLVHSNGNTY LHWYLQKPGQSPQLLIYKVS NRFSGVVPDRFSGSGGTDFTLKISRVEAEDVGIY YCSQSTYPEFTFGQGTKLEIK
509	BCMA-51	BC 7A4 97- A3-E7	scFv	aa	QVQLVQSGAEVKKPGASVKVSC KASGYSFPDYYINWVRQAPGQ GLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSINTAYMELSSLTSED TAVYFCASLYDYDWYFDVWGQGT MTVTVSSGGGGSGGGSGGGGS DIVMTQTPLSLVTPGQPASISCKSSQSLVHSNGNTY LHWYLQKPGQSPQLLIYKVS NRFSGVVPDRFSGSGGTDFTLKISRVEAEDVGIY YCSQSTYPEFTFGQGTKLEIK
510	BCMA-51 HL x CD3 HL	BC 7A4 97- A3-E7 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGAEVKKPGASVKVSC KASGYSFPDYYINWVRQAPGQ GLEWMGWIIYFASGNSEYNQKFTGR VTMTRDTSINTAYMELSSLTSED TAVYFCASLYDYDWYFDVWGQGT MTVTVSSGGGGSGGGSGGGGS

					DI VMTQTPLSLSVTPGQPASISCKSSQSLVHSNGNTYLHWY LQKPGQSPQLLIYKVS NRFSGVPDRFSGSGGTDFTLKISRVEAEDVGIYYCSQSSTYPEFTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDS KNTAYLQMN LKTEDTAVYYCVRHGNFGNSYISYWAYWGQTLVTVSSGGSGGGSGGSGQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
511	BCMA-52	BC E11 19-F11-F8	VH CDR1	aa	NAWMD
512	BCMA-52	BC E11 19-F11-F8	VH CDR2	aa	QITAKSNNNYATYYAEPVKG
513	BCMA-52	BC E11 19-F11-F8	VH CDR3	aa	DGYH
514	BCMA-52	BC E11 19-F11-F8	VL CDR1	aa	RASEDIRNGLA
515	BCMA-52	BC E11 19-F11-F8	VL CDR2	aa	NANSLHT
516	BCMA-52	BC E11 19-F11-F8	VL CDR3	aa	EDTSKYPYT
517	BCMA-52	BC E11 19-F11-F8	VH	aa	EVQLVESGGGLVKPGE SLRLSCAASGFTFSNAWMDWVRQAPGKRLEWVAQITAKSNNNYATYYAEPVKGRFTISRDDS KNTLYLQMN SLKTEDTAVYYCTDDGYHWGQGTLLVTVSS
518	BCMA-52	BC E11 19-F11-F8	VL	aa	AIQMTQSPSSLSASVGETVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVP SRFSGSGSGTEFTLKISSLPQPEDEATYYCEDTSKYPYTFGQGTKLEIK
519	BCMA-52	BC E11 19-F11-F8	scFv	aa	EVQLVESGGGLVKPGE SLRLSCAASGFTFSNAWMDWVRQAPGKRLEWVAQITAKSNNNYATYYAEPVKGRFTISRDDS KNTLYLQMN SLKTEDTAVYYCTDDGYHWGQGTLLVTVSSGGSGGGSGGSGAIQM TQSPSSLSASVGETVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVP SRFSGSGSGTEFTLKISSLPQPEDEATYYCEDTSKYPYTFGQGTKLEIK
520	BCMA-52 HL x CD3 HL	BC E11 19-F11-F8 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVKPGE SLRLSCAASGFTFSNAWMDWVRQAPGKRLEWVAQITAKSNNNYATYYAEPVKGRFTISRDDS KNTLYLQMN SLKTEDTAVYYCTDDGYHWGQGTLLVTVSSGGSGGGSGGSGAIQM TQSPSSLSASVGETVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVP SRFSGSGSGTEFTLKISSLPQPEDEATYYCEDTSKYPYTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDS KNTAYLQMN LKTEDTAVYYCVRHGNFGNSYISYWAYWGQTLVTVSSGGSGGGSGGSGQTVVTQEP SLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEA

						EYYCVLWYSNRWVFGGGTKLTVL
521	BCMA-53	BC E11 19- G3-F8	VH CDR1	aa		NAWMD
522	BCMA-53	BC E11 19- G3-F8	VH CDR2	aa		QITAKSNNNYATYYAAPVKG
523	BCMA-53	BC E11 19- G3-F8	VH CDR3	aa		DGYH
524	BCMA-53	BC E11 19- G3-F8	VL CDR1	aa		RASEDIRNGLA
525	BCMA-53	BC E11 19- G3-F8	VL CDR2	aa		NANSLHS
526	BCMA-53	BC E11 19- G3-F8	VL CDR3	aa		EDTSKYPYT
527	BCMA-53	BC E11 19- G3-F8	VH	aa		EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNNYATYYAAPVK GRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGTLVTVSS
528	BCMA-53	BC E11 19- G3-F8	VL	aa		AIQMTQSPSSLSASVGDRVTIKCRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHSGVPSRFSGSGS GTDFTLTISSMQPEDEGTYYCEDTSKYPYTFGQGTKLEIK
529	BCMA-53	BC E11 19- G3-F8	scFv	aa		EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNNYATYYAAPVK GRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGTLVTVSSGGSGGGSGGSAIQM TQSPSSLSASVGDRVTIKCRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHSGVPSRFSGSGGTD TLTISSMQPEDEGTYYCEDTSKYPYTFGQGTKLEIK
530	BCMA-53 HL x CD3 HL	BC E11 19- G3-F8 HL x CD3 HL	bispecific molecule	aa		EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNNYATYYAAPVK GRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGTLVTVSSGGSGGGSGGSAIQM TQSPSSLSASVGDRVTIKCRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHSGVPSRFSGSGGTD TLTISSMQPEDEGTYYCEDTSKYPYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAAS GFTFNKYAMNNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTED TAVYYCVRHGNFNSYISYWAYWGQGLVTVSSGGSGGGSGGGSQT VVTQEPSSLTVSPGGT VT LTCGSSSTGAVTSGNYPNWWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEA EYYCVLWYSNRWVFGGGTKLTVL
531	BCMA-54	BC E11 19- B2-F8	VH CDR1	aa		NAWMD
532	BCMA-54	BC E11 19- B2-F8	VH CDR2	aa		QITAKSNNNYATYYAAPVKG

533	BCMA-54	BC E11 19- B2-F8	VH CDR3	aa	DGYH	
534	BCMA-54	BC E11 19- B2-F8	VL CDR1	aa	RASEDIRNGLA	
535	BCMA-54	BC E11 19- B2-F8	VL CDR2	aa	NANSLHT	
536	BCMA-54	BC E11 19- B2-F8	VL CDR3	aa	EDTSKYPYT	
537	BCMA-54	BC E11 19- B2-F8	VH	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNNYATYYAAPVK GRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGTLVTVSS	
538	BCMA-54	BC E11 19- B2-F8	VL	aa	AIQMTQSPSSLSASVGDRTVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVPSPRFSGSGS GTDFTLTITSSLPQPEDEAIYYCEDTSKYPYTFGQGTKLEIK	
539	BCMA-54	BC E11 19- B2-F8	scFv	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNNYATYYAAPVK GRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGTLVTVSSGGSGGGSGGSAIQM TQSPSSLSASVGDRTVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVPSPRFSGSGGTDF TLTITSSLPQPEDEAIYYCEDTSKYPYTFGQGTKLEIK	
540	BCMA-54 HL x CD3 HL	BC E11 19- B2-F8 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNNYATYYAAPVK GRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGTLVTVSSGGSGGGSGGSAIQM TQSPSSLSASVGDRTVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVPSPRFSGSGGTDF TLTITSSLPQPEDEAIYYCEDTSKYPYTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAAS GFTFNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTED TAVYYCVRHGNFNSYISYWAYWGQTLVTVSSGGSGGGSGGSGQTVVTQEPSTVSPGGTVT LTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKKAALTLSGVQPEDEA EYCVLWYSNRWVFGGGTKLTVL	
541	BCMA-55	BC E11-20- H9-E9	VH CDR1	aa	NAWMD	
542	BCMA-55	BC E11-20- H9-E9	VH CDR2	aa	QITAKSNNYATYYAAPVKG	
543	BCMA-55	BC E11-20- H9-E9	VH CDR3	aa	DGYH	
544	BCMA-55	BC E11-20- H9-E9	VL CDR1	aa	RASEDIRNGLA	
545	BCMA-55	BC E11-20- H9-E9	VL CDR2	aa	NANSLHT	
546	BCMA-55	BC E11-20- H9-E9	VL CDR3	aa	EETLKYPYT	

547	BCMA-55	BC E11-20-H9-E9	VH	aa	EVQLVESGGSLVKPGGSLRLSCAASGFTFSNAWMDWVRQAPGKRLEWVAQITAKSNNYATYYAAPVKGRFTISRDDSKNTLYLQMNSLKEEDTAVYYCTDDGYHWGQGTLVTVSS
548	BCMA-55	BC E11-20-H9-E9	VL	aa	AIQMTQSPSSLSASVGDRVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVPSRFSGSGSGTDFTLTISNLPQPEDEATYYCEETLKYPYTFGQGTKLEIK
549	BCMA-55	BC E11-20-H9-E9	scFv	aa	EVQLVESGGSLVKPGGSLRLSCAASGFTFSNAWMDWVRQAPGKRLEWVAQITAKSNNYATYYAAPVKGRFTISRDDSKNTLYLQMNSLKEEDTAVYYCTDDGYHWGQGTLVTVSSGGSGGGSGGSAIQMTQSPSSLSASVGDRVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVPSRFSGSGGTDFTLTISNLPQPEDEATYYCEETLKYPYTFGQGTKLEIK
550	BCMA-55 HL x CD3 HL	BC E11-20-H9-E9 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGSLVKPGGSLRLSCAASGFTFSNAWMDWVRQAPGKRLEWVAQITAKSNNYATYYAAPVKGRFTISRDDSKNTLYLQMNSLKEEDTAVYYCTDDGYHWGQGTLVTVSSGGSGGGSGGSAIQMTQSPSSLSASVGDRVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVPSRFSGSGGTDFTLTISNLPQPEDEATYYCEETLKYPYTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFNSYISYWAYWGQTLVTVSSGGSGGGSGGSGQTVVTQEPSSLTVSPGGTVT LTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYFCVLWYSNRWVFGGGTKLTVL
551	BCMA-56	BC E11-19-F11-E9	VH CDR1	aa	NAWMD
552	BCMA-56	BC E11-19-F11-E9	VH CDR2	aa	QITAKSNNYATYYAEPVKG
553	BCMA-56	BC E11-19-F11-E9	VH CDR3	aa	DGYH
554	BCMA-56	BC E11-19-F11-E9	VL CDR1	aa	RASEDIRNGLA
555	BCMA-56	BC E11-19-F11-E9	VL CDR2	aa	NANSLHT
556	BCMA-56	BC E11-19-F11-E9	VL CDR3	aa	EETLKYPYT
557	BCMA-56	BC E11-19-F11-E9	VH	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWVAQITAKSNNYATYYAEPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTDDGYHWGQGTLVTVSS
558	BCMA-56	BC E11-19-F11-E9	VL	aa	AIQMTQSPSSLSASVGETVTIACRASEDIRNGLAWYQQKPGKAPKLLIYNANSLHTGVPSRFSGSGSGTEFTLKISSLPQPEDEATYYCEETLKYPYTFGQGTKLEIK

559	BCMA-56	BC E11-19-F11-E9	scFv	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWVAQITAKSNNYATYYAEFPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTDDGYHWGQGLVTVSSGGSGGGSGGSAIQMTQSPSSLSASVGETVTIACRASEDIRNGLAWYQQKPKGAPKLLIYNANSLHTGVPSRFSGSGGTEFTLKISLQPEDEATYYCEEETLKYPYTFGQGTKLEIK
560	BCMA-56 HL x CD3 HL	BC E11-19-F11-E9 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWVAQITAKSNNYATYYAEFPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTDDGYHWGQGLVTVSSGGSGGGSGGSAIQMTQSPSSLSASVGETVTIACRASEDIRNGLAWYQQKPKGAPKLLIYNANSLHTGVPSRFSGSGGTEFTLKISLQPEDEATYYCEEETLKYPYTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFNSYISYWAYWGQGLVTVSSGGSGGGSGGSGQTVVVTQEPSSLTVSPGGTVELTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKKAALTLSGVQPEDEAEYCYCLWYSNRWVFGGGTKLTVL
561	BCMA-57	BC E11-19-B2-E9	VH CDR1	aa	NAWMD
562	BCMA-57	BC E11-19-B2-E9	VH CDR2	aa	QITAKSNNYATYYAAPVKG
563	BCMA-57	BC E11-19-B2-E9	VH CDR3	aa	DGYH
564	BCMA-57	BC E11-19-B2-E9	VL CDR1	aa	RASEDIRNGLA
565	BCMA-57	BC E11-19-B2-E9	VL CDR2	aa	NANSLHT
566	BCMA-57	BC E11-19-B2-E9	VL CDR3	aa	EETLKYPYT
567	BCMA-57	BC E11-19-B2-E9	VH	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNYATYYAAPVKGGRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGLVTVSS
568	BCMA-57	BC E11-19-B2-E9	VL	aa	AIQMTQSPSSLSASVGDVRTIACRASEDIRNGLAWYQQKPKGAPKLLIYNANSLHTGVPSRFSGSGSGTDFTLTISLQPEDEAIYYCEEETLKYPYTFGQGTKLEIK
569	BCMA-57	BC E11-19-B2-E9	scFv	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNYATYYAAPVKGGRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGLVTVSSGGSGGGSGGSAIQMTQSPSSLSASVGDVRTIACRASEDIRNGLAWYQQKPKGAPKLLIYNANSLHTGVPSRFSGSGGTDFTLTISLQPEDEAIYYCEEETLKYPYTFGQGTKLEIK

570	BCMA-57 HL x CD3 HL	BC E11-19-B2-E9 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNYATYYAAPVKGRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGLVTVSSGGGGSGGGSGGSAIQMTQSPSSLSASVGDRTIACRASEDIRNGLAWYQQKPKAPKLLIYNANSLHTGVPSRFSGSGGTDFTLTISSLQPEDEAIYYCEEETLKYPYTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFNSYISYWAYWGQGLVTVSSGGGGSGGGSGGTVVVTQEPSTVSPGGTVTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKKAALTLSGVQPEDEAEYCYCLWYSNRWVFGGGTKLTVL
571	BCMA-58	BC E11-19-G3-E9	VH CDR1	aa	NAWMD
572	BCMA-58	BC E11-19-G3-E9	VH CDR2	aa	QITAKSNNYATYYAAPVKG
573	BCMA-58	BC E11-19-G3-E9	VH CDR3	aa	DGYH
574	BCMA-58	BC E11-19-G3-E9	VL CDR1	aa	RASEDIRNGLA
575	BCMA-58	BC E11-19-G3-E9	VL CDR2	aa	NANSLHS
576	BCMA-58	BC E11-19-G3-E9	VL CDR3	aa	EETLKYPYT
577	BCMA-58	BC E11-19-G3-E9	VH	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNYATYYAAPVKGRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGLVTVSS
578	BCMA-58	BC E11-19-G3-E9	VL	aa	AIQMTQSPSSLSASVGDRTVIKCRASEDIRNGLAWYQQKPKAPKLLIYNANSLHSGVPSRFSGSGSGTDFTLTISSTMQPEDEGTYYCEEETLKYPYTFGQGTKLEIK
579	BCMA-58	BC E11-19-G3-E9	scFv	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNYATYYAAPVKGRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGLVTVSSGGGGSGGGSGGSAIQMTQSPSSLSASVGDRTVIKCRASEDIRNGLAWYQQKPKAPKLLIYNANSLHSGVPSRFSGSGGTDFTLTISSMQPEDEGTYYCEEETLKYPYTFGQGTKLEIK
580	BCMA-58 HL x CD3 HL	BC E11-19-G3-E9 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVKPGESLRLSCAASGFTFSNAWMDWVRQAPGKRLEWIAQITAKSNNYATYYAAPVKGRFTISRDDSKNTLYLQMNSLKKEDTAVYYCTDDGYHWGQGLVTVSSGGGGSGGGSGGSAIQMTQSPSSLSASVGDRTVIKCRASEDIRNGLAWYQQKPKAPKLLIYNANSLHSGVPSRFSGSGGTDFTLTISSMQPEDEGTYYCEEETLKYPYTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTED

						TAVYYCVRHGNFNSYISYWAYWGQGTLLVTVSSGGGGGGGGSGQTVVVTQEPSLTVSPGGTVT LTCGSSSTGAVTSGNYPNWWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEA EYCVLWYSNRWVFGGGTKLTVL
581	BCMA-59	BC 5G9- 91-D2	VH CDR1	aa		NYDMA
582	BCMA-59	BC 5G9- 91-D2	VH CDR2	aa		SIITSGGDNYRDSVKG
583	BCMA-59	BC 5G9- 91-D2	VH CDR3	aa		HDYYDGSYGFAI
584	BCMA-59	BC 5G9- 91-D2	VL CDR1	aa		KASQSVGINVD
585	BCMA-59	BC 5G9- 91-D2	VL CDR2	aa		GASNRHT
586	BCMA-59	BC 5G9- 91-D2	VL CDR3	aa		LQYGSIPFT
587	BCMA-59	BC 5G9- 91-D2	VH	aa		QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPKGLEWVASIITSGGDNYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAIYWGQGTLLVTVSS
588	BCMA-59	BC 5G9- 91-D2	VL	aa		EIVMTQSPASMSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGS GTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
589	BCMA-59	BC 5G9- 91-D2	scFv	aa		QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPKGLEWVASIITSGGDNYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAIYWGQGTLLVTVSSGGGGGGSGGG GSEIVMTQSPASMSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
590	BCMA-59 HL x CD3 HL	BC 5G9- 91-D2 HL x CD3 HL	bispecific molecule	aa		QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPKGLEWVASIITSGGDNYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAIYWGQGTLLVTVSSGGGGGGSGGG GSEIVMTQSPASMSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGTEFTLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWNVRQAPKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFNSYISYWAYWGQGTLLVTVSSGGGGGGGGSGQTVVVTQEPSLTVS PGGTVTLTCGSSSTGAVTSGNYPNWWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYCVLWYSNRWVFGGGTKLTVL
591	BCMA-60	BC 5G9- 91-C7	VH CDR1	aa		NYDMA

592	BCMA-60	BC 5G9-91-C7	VH CDR2	aa	SIITSGGDNYRDSVKG
593	BCMA-60	BC 5G9-91-C7	VH CDR3	aa	HDYYDGSYGfAY
594	BCMA-60	BC 5G9-91-C7	VL CDR1	aa	KASQSVGINVD
595	BCMA-60	BC 5G9-91-C7	VL CDR2	aa	GASNRHT
596	BCMA-60	BC 5G9-91-C7	VL CDR3	aa	LQYGSIPFT
597	BCMA-60	BC 5G9-91-C7	VH	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIIITSGGDNYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGfAYWGQGLVTVSS
598	BCMA-60	BC 5G9-91-C7	VL	aa	EIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGS GREFTLTISLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
599	BCMA-60	BC 5G9-91-C7	scFv	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIIITSGGDNYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGfAYWGQGLVTVSSGGGSGGGSGGG GSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGREFTLTISLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK
600	BCMA-60 HL x CD3 HL	BC 5G9-91-C7 HL x CD3 HL	bispecific molecule	aa	QVQLVESGGGVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIIITSGGDNYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGfAYWGQGLVTVSSGGGSGGGSGGG GSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGREFTLTISLQSEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGSGGGSGGGSQTVVTQEP SLTVS PGGTVTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGKAAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
601	BCMA-61	BC 5G9-91-E4	VH CDR1	aa	NYDMA
602	BCMA-61	BC 5G9-91-E4	VH CDR2	aa	SIITSGGDNYRDSVKG
603	BCMA-61	BC 5G9-91-E4	VH CDR3	aa	HDYYDGSYGfAY
604	BCMA-61	BC 5G9-91-E4	VL CDR1	aa	KASQSVGINVD

605	BCMA-61	BC 5G9-91-E4	VL CDR2	aa	GASNRHT	
606	BCMA-61	BC 5G9-91-E4	VL CDR3	aa	LQYGSIPFT	
607	BCMA-61	BC 5G9-91-E4	VH	aa	QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGGDNYYRDSVKGR FTISRDN SKNTLYLQMNSLRSEDTAVYYCVRHDYYDGSYGFA YWGQGTLLVTVSS	
608	BCMA-61	BC 5G9-91-E4	VL	aa	EIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGS GTEFTLLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK	
609	BCMA-61	BC 5G9-91-E4	scFv	aa	QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGGDNYYRDSVKGR FTISRDN SKNTLYLQMNSLRSEDTAVYYCVRHDYYDGSYGFA YWGQGTLLVTVSSGGGGSGGGSGGG GSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGTEFTLLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIK	
610	BCMA-61 HL x CD3 HL	BC 5G9-91-E4 HL x CD3 HL	bispecific molecule	aa	QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGGDNYYRDSVKGR FTISRDN SKNTLYLQMNSLRSEDTAVYYCVRHDYYDGSYGFA YWGQGTLLVTVSSGGGGSGGGSGGG GSEIVMTQSPATLSVSPGERVTLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGS GSGTEFTLLTISSLQSEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSGGGQT VVTQEPSLTVS PGGT VTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL	
611	BCMA-62	BC 5G9-92-E10	VH CDR1	aa	NYDMA	
612	BCMA-62	BC 5G9-92-E10	VH CDR2	aa	SIITSGGDNYYRDSVKG	
613	BCMA-62	BC 5G9-92-E10	VH CDR3	aa	HDYYDGSYGFA Y	
614	BCMA-62	BC 5G9-92-E10	VL CDR1	aa	KASQSVGINVD	
615	BCMA-62	BC 5G9-92-E10	VL CDR2	aa	GASNRHT	
616	BCMA-62	BC 5G9-92-E10	VL CDR3	aa	LQYGSIPFT	
617	BCMA-62	BC 5G9-92-E10	VH	aa	QVQLVESGGGVQPGRSRLRSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGGDNYYRDSVKGR	

						FTVSRDNSKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSS
618	BCMA-62	BC 5G9-92-E10	VL	aa		EIVMTQSPATLSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLLTISSLQAEDFAVYYCLQYGSIPFTFGPGTKVDIK
619	BCMA-62	BC 5G9-92-E10	scFv	aa		QVQLVESGGGVVQPGRSRLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGGDNYYRDSVKGRFTVSRDNSKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSSGGGGSGGGSGGGGSEIVMTQSPATLSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLLTISSLQAEDFAVYYCLQYGSIPFTFGPGTKVDIK
620	BCMA-62 HL x CD3 HL	BC 5G9-92-E10 HL x CD3 HL	bispecific molecule	aa		QVQLVESGGGVVQPGRSRLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVASIITSGGDNYYRDSVKGRFTVSRDNSKNTLYLQMNSLRAEDTAVYYCVRHDYYDGSYGFAFWGQGLVTVSSGGGGSGGGSGGGGSEIVMTQSPATLSVSPGERATLSCKASQSVGINVDWYQQKPGQAPRLLIYGASNRHTGIPARFSGSGSGTEFTLLTISSLQAEDFAVYYCLQYGSIPFTFGPGTKVDIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLIKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGSGGGGQTVVTTQEPSLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYCVLWYSNRWVFGGGTKLTVL
621	BCMA-63	BC 3A4-37-C8	VH CDR1	aa		NYDMA
622	BCMA-63	BC 3A4-37-C8	VH CDR2	aa		SISTRGDITSYRDSVKG
623	BCMA-63	BC 3A4-37-C8	VH CDR3	aa		QDYITDYMGFAY
624	BCMA-63	BC 3A4-37-C8	VL CDR1	aa		RASEDIYNGLA
625	BCMA-63	BC 3A4-37-C8	VL CDR2	aa		GASSLQD
626	BCMA-63	BC 3A4-37-C8	VL CDR3	aa		QQSYKYPLT
627	BCMA-63	BC 3A4-37-C8	VH	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLVTVSS
628	BCMA-63	BC 3A4-37-C8	VL	aa		AIQMTQSPSSLSASVGDVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSFGSGSGTDYTLTISSLQPEDEATYYCQQSYKYPLTFGGGTKVEIK
629	BCMA-63	BC 3A4-37-C8	scFv	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR

630	BCMA-63 HL x CD3 HL	BC 3A4-37-C8 HL x CD3 HL	bispecific molecule	aa	FTISRDNKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLVTVSSGGGGGGGGGGGGG GSAIQMTQSPSSLSASVGDVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GGTDYTLTISSLQPEDEATYYCQQSYKYPLTFGGGTKVEIK
631	BCMA-64	BC 3A4-37-C9	VH CDR1	aa	NYDMA
632	BCMA-64	BC 3A4-37-C9	VH CDR2	aa	SISTRGDITSYRDSVKG
633	BCMA-64	BC 3A4-37-C9	VH CDR3	aa	QDY YTDYMGFAY
634	BCMA-64	BC 3A4-37-C9	VL CDR1	aa	RASEDIYNGLA
635	BCMA-64	BC 3A4-37-C9	VL CDR2	aa	GASSLQD
636	BCMA-64	BC 3A4-37-C9	VL CDR3	aa	QQSYKYPLT
637	BCMA-64	BC 3A4-37-C9	VH	aa	EVQLLESGGGLVQPGRSRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLVTVSS
638	BCMA-64	BC 3A4-37-C9	VL	aa	AIQMTQSPSSLSASVGDVRTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GTDFTLTISSMQPEDEATYYCQQSYKYPLTFGGGTKVEIK
639	BCMA-64	BC 3A4-37-C9	scFv	aa	EVQLLESGGGLVQPGRSRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLVTVSSGGGGGGGGGGGGG GSAIQMTQSPSSLSASVGDVRTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GGTDFTLTISSMQPEDEATYYCQQSYKYPLTFGGGTKVEIK
640	BCMA-64 HL x CD3 HL	BC 3A4-37-C9 HL x CD3 HL	bispecific molecule	aa	EVQLLESGGGLVQPGRSRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLVTVSSGGGGGGGGGGGGG

					GSAIQMTQSPSSLSASVGDRVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTDFTLTIISSMQPEDEATYYCQQSYKYPLTFGGGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGGGGSGGGGQTVVTVTQEPSLTVS PGGTVTLTCCSSTGAVTSGNYPNWNVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYCVLWYSNRWVFGGGTKLTVL
641	BCMA-65	BC 3A4-37- E11	VH CDR1	aa	NYDMA
642	BCMA-65	BC 3A4-37- E11	VH CDR2	aa	SISTRGDITSYRDSVKG
643	BCMA-65	BC 3A4-37- E11	VH CDR3	aa	QDYITDYMGFAY
644	BCMA-65	BC 3A4-37- E11	VL CDR1	aa	RASEDIYNGLA
645	BCMA-65	BC 3A4-37- E11	VL CDR2	aa	GASSLQD
646	BCMA-65	BC 3A4-37- E11	VL CDR3	aa	QQSYKYPLT
647	BCMA-65	BC 3A4-37- E11	VH	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGTLLTVTVSS
648	BCMA-65	BC 3A4-37- E11	VL	aa	AIQMTQSPSSLSASVGDRVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGS GTHYTLTISSLQPEDEATYYCQQSYKYPLTFGGGTKVEIK
649	BCMA-65	BC 3A4-37- E11	scFv	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGTLLTVTVSSGGGGSGGGGGG GSAIQMTQSPSSLSASVGDRVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTHYTLTISSLQPEDEATYYCQQSYKYPLTFGGGTKVEIK
650	BCMA-65 HL x CD3 HL	BC 3A4-37- E11 HL x CD3 HL	bispecific molecule	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGTLLTVTVSSGGGGSGGGGGG GSAIQMTQSPSSLSASVGDRVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTHYTLTISSLQPEDEATYYCQQSYKYPLTFGGGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGGGGSGGGGQTVVTVTQEPSLTVS PGGTVTLTCCSSTGAVTSGNYPNWNVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV

						QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
651	BCMA-66	BC 3A4-37- C8-G1	VH CDR1	aa		NYDMA
652	BCMA-66	BC 3A4-37- C8-G1	VH CDR2	aa		SISTRGDITSYRDSVKG
653	BCMA-66	BC 3A4-37- C8-G1	VH CDR3	aa		QDYITDYMGFAY
654	BCMA-66	BC 3A4-37- C8-G1	VL CDR1	aa		RASEDIYNGLA
655	BCMA-66	BC 3A4-37- C8-G1	VL CDR2	aa		GASSLQD
656	BCMA-66	BC 3A4-37- C8-G1	VL CDR3	aa		AGPHKYPLT
657	BCMA-66	BC 3A4-37- C8-G1	VH	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLTVTVSS
658	BCMA-66	BC 3A4-37- C8-G1	VL	aa		AIQMTQSPSSLSASVGDVTITTCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGS GTDYTLTISSLQPEDEATYYCAGPHKYPLTFGGGTKVEIK
659	BCMA-66	BC 3A4-37- C8-G1	scFv	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLTVTVSSGGGGSGGGSGG GSAIQMTQSPSSLSASVGDVTITTCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTDYTLTISSLQPEDEATYYCAGPHKYPLTFGGGTKVEIK
660	BCMA-66 HL x CD3 HL	BC 3A4-37- C8-G1 HL x CD3 HL	bispecific molecule	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLTVTVSSGGGGSGGGSGG GSAIQMTQSPSSLSASVGDVTITTCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTDYTLTISSLQPEDEATYYCAGPHKYPLTFGGGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGQT VVTQEPSLTVS PGGT VTLTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
661	BCMA-67	BC 3A4-37- E11-G1	VH CDR1	aa		NYDMA
662	BCMA-67	BC 3A4-37- E11-G1	VH CDR2	aa		SISTRGDITSYRDSVKG

663	BCMA-67	BC 3A4-37-E11-G1	VH CDR3	aa	QDY YTDYMGFAY
664	BCMA-67	BC 3A4-37-E11-G1	VL CDR1	aa	RASEDIYNGLA
665	BCMA-67	BC 3A4-37-E11-G1	VL CDR2	aa	GASSLQD
666	BCMA-67	BC 3A4-37-E11-G1	VL CDR3	aa	AGPHKYPLT
667	BCMA-67	BC 3A4-37-E11-G1	VH	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGT LVTVSS
668	BCMA-67	BC 3A4-37-E11-G1	VL	aa	AIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGS GTHYTLTISSLQPEDEATYYCAGPHKYPLTFGGG TKVEIK
669	BCMA-67	BC 3A4-37-E11-G1	scFv	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGT LVTVSSGGGSGGGSGGG GSAIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTHYTLTISSLQPEDEATYYCAGPHKYPLTFGGG TKVEIK
670	BCMA-67 HL x CD3 HL	BC 3A4-37-E11-G1 HL x CD3 HL	bispecific molecule	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGT LVTVSSGGGSGGGSGGG GSAIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTHYTLTISSLQPEDEATYYCAGPHKYPLTFGGG TKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGT LVTVSSGGGSGGGSGGGQT VVTQEPSLTVS PGGT VTLTCGSSSTGAVTSGNYPNWWVQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
671	BCMA-68	BC 3A4-37-C8-G8	VH CDR1	aa	NYDMA
672	BCMA-68	BC 3A4-37-C8-G8	VH CDR2	aa	SISTRGDITSYRDSVKG
673	BCMA-68	BC 3A4-37-C8-G8	VH CDR3	aa	QDY YTDYMGFAY
674	BCMA-68	BC 3A4-37-C8-G8	VL CDR1	aa	RASEDIYNGLA
675	BCMA-68	BC 3A4-37-C8-G8	VL CDR2	aa	GASSLQD

676	BCMA-68	BC 3A4-37- C8-G8	VL CDR3	aa	QQSRNYQQT	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLVTVSS
677	BCMA-68	BC 3A4-37- C8-G8	VH	aa		AIQMTQSPSSLSASVGDVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGS GTDYTLTISSLQPEDEATYYCQQSRNYQQTFGGGTKVEIK
678	BCMA-68	BC 3A4-37- C8-G8	VL	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLVTVSSGGGGSGGGGGG GSAIQMTQSPSSLSASVGDVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTDYTLTISSLQPEDEATYYCQQSRNYQQTFGGGTKVEIK
679	BCMA-68	BC 3A4-37- C8-G8	scFv	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLVTVSSGGGGSGGGGGG GSAIQMTQSPSSLSASVGDVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTDYTLTISSLQPEDEATYYCQQSRNYQQTFGGGTKVEIK
680	BCMA-68 HL x CD3 HL	BC 3A4-37- C8-G8 HL x CD3 HL	bispecific molecule	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLVTVSSGGGGSGGGGGG GSAIQMTQSPSSLSASVGDVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTDYTLTISSLQPEDEATYYCQQSRNYQQTFGGGTKVEIKSGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGGSGQTVVVTQEPSSLTVS PGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKKAALTLSGV QPEDEAEYCVLWYSNRWVFGGGTKLTVL
681	BCMA-69	BC 3A4-37- E11-G8	VH CDR1	aa	NYDMA	
682	BCMA-69	BC 3A4-37- E11-G8	VH CDR2	aa	SISTRGDITSYRDSVKG	
683	BCMA-69	BC 3A4-37- E11-G8	VH CDR3	aa	QDYITDYMGFAY	
684	BCMA-69	BC 3A4-37- E11-G8	VL CDR1	aa	RASEDIYNGLA	
685	BCMA-69	BC 3A4-37- E11-G8	VL CDR2	aa	GASSLQD	
686	BCMA-69	BC 3A4-37- E11-G8	VL CDR3	aa	QQSRNYQQT	
687	BCMA-69	BC 3A4-37- E11-G8	VH	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDNKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLVTVSS	
688	BCMA-69	BC 3A4-37-	VL	aa	AIQMTQSPSSLSASVGDVRTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGS	

		E11-G8				GTHYTLTISSLQPEDEATYYCQQSRNYQQTFGGGTKVEIK
689	BCMA-69	BC 3A4-37- E11-G8	scFv	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLTVTVSSGGGGSGGGSGG GSAIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTHYTLTISSLQPEDEATYYCQQSRNYQQTFGGGTKVEIK
690	BCMA-69 HL x CD3 HL	BC 3A4-37- E11-G8 HL x CD3 HL	bispecific molecule	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLTVTVSSGGGGSGGGSGG GSAIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTHYTLTISSLQPEDEATYYCQQSRNYQQTFGGGTKVEIK
691	BCMA-70	BC 3A4-37- A11-G8	VH CDR1	aa		NYDMA
692	BCMA-70	BC 3A4-37- A11-G8	VH CDR2	aa		SISTRGDITSYRDSVKG
693	BCMA-70	BC 3A4-37- A11-G8	VH CDR3	aa		QDY YTDYMGFAY
694	BCMA-70	BC 3A4-37- A11-G8	VL CDR1	aa		RASEDIYNGLA
695	BCMA-70	BC 3A4-37- A11-G8	VL CDR2	aa		GASSLQD
696	BCMA-70	BC 3A4-37- A11-G8	VL CDR3	aa		QQSRNYQQT
697	BCMA-70	BC 3A4-37- A11-G8	VH	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLTVTVSS
698	BCMA-70	BC 3A4-37- A11-G8	VL	aa		AIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GTEFTLTISSLQPEDEATYYCQQSRNYQQTFGGGTKVEIK
699	BCMA-70	BC 3A4-37- A11-G8	scFv	aa		EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLTVTVSSGGGGSGGGSGG GSAIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTEFTLTISSLQPEDEATYYCQQSRNYQQTFGGGTKVEIK

700	BCMA-70 HL x CD3 HL	BC 3A4-37-A11-G8 HL x CD3 HL	bispecific molecule	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSI STRGDITSYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLVTVSSGGGGSGGGSGG GSAIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTEFTLTISSLQPEDEATY CQQSRNYQQTFGGG TKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFNSYISYWAYWGQGLVTVSSGGGGSGGGSQTVVTQEPSLTVS PGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEY YCVLWYSNRWVFGGGTKLTVL
701	BCMA-71	BC 3A4-37-A11-G1	VH CDR1	aa	NYDMA
702	BCMA-71	BC 3A4-37-A11-G1	VH CDR2	aa	SISTRGDITSYRDSVKG
703	BCMA-71	BC 3A4-37-A11-G1	VH CDR3	aa	QDY YTDYMGFAY
704	BCMA-71	BC 3A4-37-A11-G1	VL CDR1	aa	RASEDIYNGLA
705	BCMA-71	BC 3A4-37-A11-G1	VL CDR2	aa	GASSLQD
706	BCMA-71	BC 3A4-37-A11-G1	VL CDR3	aa	AGPHKYPLT
707	BCMA-71	BC 3A4-37-A11-G1	VH	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSI STRGDITSYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLVTVSS
708	BCMA-71	BC 3A4-37-A11-G1	VL	aa	AIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTEFTLTISSLQPEDEATY CAGPHKYPLTFGGG TKVEIK
709	BCMA-71	BC 3A4-37-A11-G1	scFv	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSI STRGDITSYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLVTVSSGGGGSGGGSGG GSAIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTEFTLTISSLQPEDEATY CAGPHKYPLTFGGG TKVEIK
710	BCMA-71 HL x CD3 HL	BC 3A4-37-A11-G1 HL x CD3 HL	bispecific molecule	aa	EVQLLESGGGLVQPGGSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSI STRGDITSYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGLVTVSSGGGGSGGGSGG GSAIQMTQSPSSLSASVGD RVTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGS GSGTEFTLTISSLQPEDEATY CAGPHKYPLTFGGG TKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN

						NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGGGGGSGGSGQTVVVTQEPSLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYCYVLWYSNRWVFGGGTKLTVL
711	BCMA-72	BC 3A4-37-C9-G1	VH CDR1	aa		NYDMA
712	BCMA-72	BC 3A4-37-C9-G1	VH CDR2	aa		SISTRGDITSYRDSVKG
713	BCMA-72	BC 3A4-37-C9-G1	VH CDR3	aa		QDYYYTDYMGFAY
714	BCMA-72	BC 3A4-37-C9-G1	VL CDR1	aa		RASEDIYNGLA
715	BCMA-72	BC 3A4-37-C9-G1	VL CDR2	aa		GASSLQD
716	BCMA-72	BC 3A4-37-C9-G1	VL CDR3	aa		AGPHKYPLT
717	BCMA-72	BC 3A4-37-C9-G1	VH	aa		EVQLLESGGGLVQPGRSRLRSCAASGFTFSNYDMAWVRQAPKGLEWVSSISTRGDITSYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGTLLVTVSS
718	BCMA-72	BC 3A4-37-C9-G1	VL	aa		AIQMTQSPSSLSASVGDRTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGSGTDFTLTISSMQPEDEATYYCAGPHKYPLTFGGGTKVEIK
719	BCMA-72	BC 3A4-37-C9-G1	scFv	aa		EVQLLESGGGLVQPGRSRLRSCAASGFTFSNYDMAWVRQAPKGLEWVSSISTRGDITSYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGTLLVTVSSGGGGGGSGGGGSAIQMTQSPSSLSASVGDRTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGSGTDFTLTISSMQPEDEATYYCAGPHKYPLTFGGGTKVEIK
720	BCMA-72 HL x CD3 HL	BC 3A4-37-C9-G1 HL x CD3 HL	bispecific molecule	aa		EVQLLESGGGLVQPGRSRLRSCAASGFTFSNYDMAWVRQAPKGLEWVSSISTRGDITSYRDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDY YTDYMGFAYWGQGTLLVTVSSGGGGGGSGGGGSAIQMTQSPSSLSASVGDRTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGSGTDFTLTISSMQPEDEATYYCAGPHKYPLTFGGGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGGGGGSGGSGQTVVVTQEPSLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYCYVLWYSNRWVFGGGTKLTVL
721	BCMA-73	BC 3A4-37-C9-G8	VH CDR1	aa		NYDMA

722	BCMA-73	BC 3A4-37- C9-G8	VH CDR2	aa	SISTRGDITSYRDSVKG
723	BCMA-73	BC 3A4-37- C9-G8	VH CDR3	aa	QDYITDYMGFAY
724	BCMA-73	BC 3A4-37- C9-G8	VL CDR1	aa	RASEDIYNGLA
725	BCMA-73	BC 3A4-37- C9-G8	VL CDR2	aa	GASSLQD
726	BCMA-73	BC 3A4-37- C9-G8	VL CDR3	aa	QQSRNYQQT
727	BCMA-73	BC 3A4-37- C9-G8	VH	aa	EVQLLESGGGLVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLVTVSS
728	BCMA-73	BC 3A4-37- C9-G8	VL	aa	AIQMTQSPSSLSASVGDRTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGS GTDFTLTISMQPEDEATYYCQQSRNYQQTFGGGTKVEIK
729	BCMA-73	BC 3A4-37- C9-G8	scFv	aa	EVQLLESGGGLVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLVTVSSGGGGSGGGGG GSAIQMTQSPSSLSASVGDRTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGS GGTDFTLTISMQPEDEATYYCQQSRNYQQTFGGGTKVEIK
730	BCMA-73 HL x CD3 HL	BC 3A4-37- C9-G8 HL x CD3 HL	bispecific molecule	aa	EVQLLESGGGLVQPGRSLRLSCAASGFTFSNYDMAWVRQAPGKGLEWVSSISTRGDITSYRDSVKGR FTISRDN SKNTLYLQMNSLRAEDTAVYYCARQDYITDYMGFAYWGQGLVTVSSGGGGSGGGGGGG GSAIQMTQSPSSLSASVGDRTITCRASEDIYNGLAWYQQKPGKAPKLLIYGASSLQDGVPSRFSGSGS GGTDFTLTISMQPEDEATYYCQQSRNYQQTFGGGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGGSGTAVTQEPSTVS PGGTVTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYFCVLWYSNRWVFGGGTKLTVL
731	BCMA-74	BC C3-33- D7-B1	VH CDR1	aa	NFDMA
732	BCMA-74	BC C3-33- D7-B1	VH CDR2	aa	SITTTGGGDTYYADSVKG
733	BCMA-74	BC C3-33- D7-B1	VH CDR3	aa	HGYDGYHLFDY
734	BCMA-74	BC C3-33- D7-B1	VL CDR1	aa	RASQGISNYLN

735	BCMA-74	BC C3-33-D7-B1	VL CDR2	aa	YTSNLQS
736	BCMA-74	BC C3-33-D7-B1	VL CDR3	aa	MGQTISSYT
737	BCMA-74	BC C3-33-D7-B1	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSS
738	BCMA-74	BC C3-33-D7-B1	VL	aa	DIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKGKAPKPLIYYTSNLQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK
739	BCMA-74	BC C3-33-D7-B1	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKGKAPKPLIYYTSNLQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK
740	BCMA-74 HL x CD3 HL	BC C3-33-D7-B1 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKGKAPKPLIYYTSNLQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSGGGGQT VVTQEPSLT VSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYCVLWYSNRWVFGGGTKLTVL
741	BCMA-75	BC C3-33-F8-B1	VH CDR1	aa	NFDMA
742	BCMA-75	BC C3-33-F8-B1	VH CDR2	aa	SITTTGGGDTYYADSVKKG
743	BCMA-75	BC C3-33-F8-B1	VH CDR3	aa	HGYDGYHLFDY
744	BCMA-75	BC C3-33-F8-B1	VL CDR1	aa	RASQGISNYLN
745	BCMA-75	BC C3-33-F8-B1	VL CDR2	aa	YTSNLQS
746	BCMA-75	BC C3-33-F8-B1	VL CDR3	aa	MGQTISSYT
747	BCMA-75	BC C3-33-F8-B1	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR

					FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSS
748	BCMA-75	BC C3-33-F8-B1	VL	aa	DIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK
749	BCMA-75	BC C3-33-F8-B1	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSSGGGGGGGGGGGGGSDIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK
750	BCMA-75 HL x CD3 HL	BC C3-33-F8-B1 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSSGGGGGGGGGGGGGSDIQMTQSPSSLSASVGDRVTITCRASQGISNYLNWYQQKPKGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNLTKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGGGGGGGGGSGTQVVTQEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEY YCVLWYSNRWVFGGGTKLTVL
751	BCMA-76	BC C3-33-F9-B1	VH CDR1	aa	NFDMA
752	BCMA-76	BC C3-33-F9-B1	VH CDR2	aa	SITTTGGGDTYYADSVKKG
753	BCMA-76	BC C3-33-F9-B1	VH CDR3	aa	HGY YDGYHLFDY
754	BCMA-76	BC C3-33-F9-B1	VL CDR1	aa	RASQGISNYLN
755	BCMA-76	BC C3-33-F9-B1	VL CDR2	aa	YTSNLQS
756	BCMA-76	BC C3-33-F9-B1	VL CDR3	aa	MGQTISSYT
757	BCMA-76	BC C3-33-F9-B1	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMDSLRS EDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSS
758	BCMA-76	BC C3-33-F9-B1	VL	aa	DIQMTQSPSSLSASVGDRVTISCRASQGISNYLNWYQQKPKGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIK
759	BCMA-76	BC C3-33-F9-B1	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR

					FTISRDNAKNTLYLQMDSLRSED TAVYYCVRHGY YDGYHLFDYWGQ GTLVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGD RVTISCRASQGISNYLNWYQ QKPGKAPKPLIYYTSNLT QSGVPSRFSGS GGTDYTLTIISSLQPEDFA TY YCMGQTISSYTFGQTK LEIK
760	BCMA-76 HL x CD3 HL	BC C3-33- F9-B1 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRL SCAASGFTFSNFDMAWVRQ APGKGLVWVSSITTGGGDT YYADSVKGR FTISRDNAKNTLYLQMDSL RSED TAVYYCVRHGY YDGYHLFD YWGQ GTLVTVSSGGGGSGGGSGG GSDIQMTQSPSSLSASVGD RVTISCRASQGISNYLNWYQ QKPGKAPKPLIYYTSNLT QSGVPSRFSGS GGTDYTLTIISSLQPEDFA TY YCMGQTISSYTFGQTK LEIKSGGGSEVQLVESGG GLVQPGGSLK LSCAASGFTFNKYAMNWRQ APGKGLEWVARIRSKYNNY ATYYADSVKDRFTISRDDS KNTAYLQMN NLKTEDTAVYYCVRHGNF GNSYISYWAYWGQ GTLVTVSSGGGGSGGGSQ TVVTQEPSLT VSPGGTVTLTCGSS TGAVTSGNYPNWVQKPGQ APRGLIGG TKFLAPGTPARFSGSLLG GKAALTLSGV QPEDEAEYCYVLWYSNRW VFGGGTKLT VL
761	BCMA-77	BC C3-33- F10B1	VH CDR1	aa	NFDMA
762	BCMA-77	BC C3-33- F10B1	VH CDR2	aa	SITTTGGGDTYYADSVK G
763	BCMA-77	BC C3-33- F10B1	VH CDR3	aa	HGY YDGYHLFDY
764	BCMA-77	BC C3-33- F10B1	VL CDR1	aa	RASQGISNYLN
765	BCMA-77	BC C3-33- F10B1	VL CDR2	aa	YTSNLQS
766	BCMA-77	BC C3-33- F10B1	VL CDR3	aa	MGQTISSYT
767	BCMA-77	BC C3-33- F10B1	VH	aa	EVQLVESGGGLVQPGRSL RLS CAASGFTFSNFDMAWVRQ APAKGLEWVSSITTGGGDT YYADSVKGR FTISRDNAKNTLYLQMN SLRAEDTAVYYCVRHGY YDGYHLFDYWGQ GTLVTVSS
768	BCMA-77	BC C3-33- F10B1	VL	aa	DIQMTQSPSSLSASVGD RVTITCRASQGISNYLNWY QKPGKAPKPLIYYTSNLT QSGVPSRFSGS GTDFTLTIISSLQPEDFA TY YCMGQTISSYTFGQTK LEIK
769	BCMA-77	BC C3-33- F10B1	scFv	aa	EVQLVESGGGLVQPGRSL RLS CAASGFTFSNFDMAWVRQ APAKGLEWVSSITTGGGDT YYADSVKGR FTISRDNAKNTLYLQMN SLRAEDTAVYYCVRHGY YDGYHLFDYWGQ GTLVTVSSGGGGSGGGSG GSDIQMTQSPSSLSASVGD RVTITCRASQGISNYLNWY QKPGKAPKPLIYYTSNLT QSGVPSRFSGS GGTDFTLTIISSLQPEDFA TY YCMGQTISSYTFGQTK LEIK
770	BCMA-77 HL x CD3 HL	BC C3-33- F10B1 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGRSL RLS CAASGFTFSNFDMAWVRQ APAKGLEWVSSITTGGGDT YYADSVKGR FTISRDNAKNTLYLQMN SLRAEDTAVYYCVRHGY YDGYHLFDYWGQ GTLVTVSSGGGGSGGGSG

					GSDIQMTQSPSSLSASVGD RVTITCRASQGISNYLNWYQQKPGKAPKPLIYYTSNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCMGQTISSYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGGGGSGGGGSGQT VVTQEPSLTVS PGGTVTLT CGSSTGAVTSGNYPNWVQOKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYCVLWYSNRWVFGGGTKLTVL
771	BCMA-78	BC E5-33- A11-A10	VH CDR1	aa	NFDMA
772	BCMA-78	BC E5-33- A11-A10	VH CDR2	aa	SITTTGGGDTYYADSVKG
773	BCMA-78	BC E5-33- A11-A10	VH CDR3	aa	HGYDGYHLFDY
774	BCMA-78	BC E5-33- A11-A10	VL CDR1	aa	RASQGISNHLN
775	BCMA-78	BC E5-33- A11-A10	VL CDR2	aa	YTSNLQS
776	BCMA-78	BC E5-33- A11-A10	VL CDR3	aa	QQYFDRPYT
777	BCMA-78	BC E5-33- A11-A10	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLFDYWGQGTLLTVTVSS
778	BCMA-78	BC E5-33- A11-A10	VL	aa	DIQMTQSPSSLSASVGD RVTISCRASQGISNHLNWFQQKPGRAPKPLIYYTSNLQSGVPSRFSGSGS GTDFTLTITISLQPEDFATYYCQQYFDRPYTFGGGTKVEIK
779	BCMA-78	BC E5-33- A11-A10	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLFDYWGQGTLLTVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGD RVTISCRASQGISNHLNWFQQKPGRAPKPLIYYTSNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQQYFDRPYTFGGGTKVEIK
780	BCMA-78 HL x CD3 HL	BC E5-33- A11-A10 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLFDYWGQGTLLTVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGD RVTISCRASQGISNHLNWFQQKPGRAPKPLIYYTSNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQQYFDRPYTFGGGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLTVSSGGGGSGGGGSGQT VVTQEPSLTVS PGGTVTLT CGSSTGAVTSGNYPNWVQOKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV

						QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
781	BCMA-79	BC E5-33-B11-A10	VH CDR1	aa		NFDMA
782	BCMA-79	BC E5-33-B11-A10	VH CDR2	aa		SITTTGGGDTYYADSVKG
783	BCMA-79	BC E5-33-B11-A10	VH CDR3	aa		HGYDGYHLFDY
784	BCMA-79	BC E5-33-B11-A10	VL CDR1	aa		RASQGISNHLN
785	BCMA-79	BC E5-33-B11-A10	VL CDR2	aa		YTSNLQS
786	BCMA-79	BC E5-33-B11-A10	VL CDR3	aa		QQYFDRPYT
787	BCMA-79	BC E5-33-B11-A10	VH	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLFDYWGQGLVTVSS
788	BCMA-79	BC E5-33-B11-A10	VL	aa		DIQMTQSPSSLSASVGDRVTISCRASQGISNHLNWYQQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGS GTDYTLTISSLQPEDFATYYCQQYFDRPYTFGGGTKVEIK
789	BCMA-79	BC E5-33-B11-A10	scFv	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLFDYWGQGLVTVSSGGGGSGGGSGG GSDIQMTQSPSSLSASVGDRVTISCRASQGISNHLNWYQQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGS GGTDYTLTISSLQPEDFATYYCQQYFDRPYTFGGGTKVEIK
790	BCMA-79 HL x CD3 HL	BC E5-33-B11-A10 HL x CD3 HL	bispecific molecule	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLFDYWGQGLVTVSSGGGGSGGGSGG GSDIQMTQSPSSLSASVGDRVTISCRASQGISNHLNWYQQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGS GGTDYTLTISSLQPEDFATYYCQQYFDRPYTFGGGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGGSGQT VVTQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGKAAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
791	BCMA-80	BC E5-33-G11-A10	VH CDR1	aa		NFDMA
792	BCMA-80	BC E5-33-G11-A10	VH CDR2	aa		SITTTGGGDTYYADSVKG

793	BCMA-80	BC E5-33- G11-A10	VH CDR3	aa	HGYDGYHLFDY
794	BCMA-80	BC E5-33- G11-A10	VL CDR1	aa	RASQGISNHLN
795	BCMA-80	BC E5-33- G11-A10	VL CDR2	aa	YTSNLQS
796	BCMA-80	BC E5-33- G11-A10	VL CDR3	aa	QQYFDRPYT
797	BCMA-80	BC E5-33- G11-A10	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSS
798	BCMA-80	BC E5-33- G11-A10	VL	aa	DIQMTQSPSSLSASVGDRVTITCRASQGISNHLNWFQQKPGKAPKPLIYYTSNLSQSGVPSRFSGSGS GTDFTLTITSSLPEDFATYYCQQYFDRPYTFGGGTKVEIK
799	BCMA-80	BC E5-33- G11-A10	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSSGGGGSGGGGG GSDIQMTQSPSSLSASVGDRVTITCRASQGISNHLNWFQQKPGKAPKPLIYYTSNLSQSGVPSRFSGSGS GGTDFTLTITSSLPEDFATYYCQQYFDRPYTFGGGTKVEIK
800	BCMA-80 HL x CD3 HL	BC E5-33- G11-A10 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSSGGGGSGGGGG GSDIQMTQSPSSLSASVGDRVTITCRASQGISNHLNWFQQKPGKAPKPLIYYTSNLSQSGVPSRFSGSGS GGTDFTLTITSSLPEDFATYYCQQYFDRPYTFGGGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGGSGTAVVTVQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLGKGKAAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
801	BCMA-81	BC E5-33- G12-A10	VH CDR1	aa	NFDMA
802	BCMA-81	BC E5-33- G12-A10	VH CDR2	aa	SITTTGGGDTYYADSVKG
803	BCMA-81	BC E5-33- G12-A10	VH CDR3	aa	HGYDGYHLFDY
804	BCMA-81	BC E5-33- G12-A10	VL CDR1	aa	RASQGISNHLN
805	BCMA-81	BC E5-33- G12-A10	VL CDR2	aa	YTSNLQS

806	BCMA-81	BC E5-33- G12-A10	VL CDR3	aa	QQYFDRPYT	EVQLVESGGGLVQPGRSRLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHFLFDYWGQGTLVTVSS
807	BCMA-81	BC E5-33- G12-A10	VH	aa		DIQMTQSPSSLSASVGERVTITCRASQGISNHLNWIYQKPGKAPKSLIYYTSNLTQSGVPSRFSGSGS GTDFTLTITSSLPEDFATYYCQQYFDRPYTFGGGTKVEIK
808	BCMA-81	BC E5-33- G12-A10	VL	aa		EVQLVESGGGLVQPGRSRLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHFLFDYWGQGTLVTVSSGGGSGGGGGG GSDIQMTQSPSSLSASVGERVTITCRASQGISNHLNWIYQKPGKAPKSLIYYTSNLTQSGVPSRFSGSGS GGTDFTLTITSSLPEDFATYYCQQYFDRPYTFGGGTKVEIK
809	BCMA-81	BC E5-33- G12-A10	scFv	aa		EVQLVESGGGLVQPGRSRLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHFLFDYWGQGTLVTVSSGGGSGGGGGG GSDIQMTQSPSSLSASVGERVTITCRASQGISNHLNWIYQKPGKAPKSLIYYTSNLTQSGVPSRFSGSGS GGTDFTLTITSSLPEDFATYYCQQYFDRPYTFGGGTKVEIK
810	BCMA-81 HL x CD3 HL	BC E5-33- G12-A10 HL x CD3 HL	bispecific molecule	aa		EVQLVESGGGLVQPGRSRLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHFLFDYWGQGTLVTVSSGGGSGGGGGG GSDIQMTQSPSSLSASVGERVTITCRASQGISNHLNWIYQKPGKAPKSLIYYTSNLTQSGVPSRFSGSGS GGTDFTLTITSSLPEDFATYYCQQYFDRPYTFGGGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQTLVTVSSGGGSGGGGSGTQVVTQEPSTLTVS PGGTVTLTGSGSTGAVTSGNYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGKAAALTLSGV QPEDEAEYCVLWYSNRWVFGGGTKLTVL
811	BCMA-82	BC E5-33- A11-B8	VH CDR1	aa	NFDMA	
812	BCMA-82	BC E5-33- A11-B8	VH CDR2	aa	SITTTGGGDTYYADSVKG	
813	BCMA-82	BC E5-33- A11-B8	VH CDR3	aa	HGYDGYHFLDY	
814	BCMA-82	BC E5-33- A11-B8	VL CDR1	aa	RASQGISNHLN	
815	BCMA-82	BC E5-33- A11-B8	VL CDR2	aa	YTSNLQS	
816	BCMA-82	BC E5-33- A11-B8	VL CDR3	aa	QQYSNLPYT	
817	BCMA-82	BC E5-33- A11-B8	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHFLFDYWGQGTLVTVSS	
818	BCMA-82	BC E5-33- A11-B8	VL	aa	DIQMTQSPSSLSASVGDRTVITISCRASQGISNHLNWFQQKPGRAPKPLIYYTSNLTQSGVPSRFSGSGS	

		A11-B8				GTDFTLTISSLQPEDFATYYCQQYSNLPYTFGGGTKVEIK
819	BCMA-82	BC E5-33- A11-B8	scFv	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLTVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGDRTVISCRAEQGISNHLNWFQQKPGRAPKPLIYYTSNQLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQQYSNLPYTFGGGTKVEIK
820	BCMA-82 HL x CD3 HL	BC E5-33- A11-B8 HL x CD3 HL	bispecific molecule	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLTVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGDRTVISCRAEQGISNHLNWFQQKPGRAPKPLIYYTSNQLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQQYSNLPYTFGGGTKVEIK
821	BCMA-83	BC E5-33- B11-B8	VH CDR1	aa		NFDMA
822	BCMA-83	BC E5-33- B11-B8	VH CDR2	aa		SITTTGGGDTYYADSVKG
823	BCMA-83	BC E5-33- B11-B8	VH CDR3	aa		HGY YDGYHLFDY
824	BCMA-83	BC E5-33- B11-B8	VL CDR1	aa		RASQGISNHLN
825	BCMA-83	BC E5-33- B11-B8	VL CDR2	aa		YTSNLQS
826	BCMA-83	BC E5-33- B11-B8	VL CDR3	aa		QQYSNLPYT
827	BCMA-83	BC E5-33- B11-B8	VH	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLTVTVSS
828	BCMA-83	BC E5-33- B11-B8	VL	aa		DIQMTQSPSSLSASVGDRTVISCRAEQGISNHLNWFQQKPGKAPKPLIYYTSNQLQSGVPSRFSGS GTDYTLTISSLQPEDFATYYCQQYSNLPYTFGGGTKVEIK
829	BCMA-83	BC E5-33- B11-B8	scFv	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGY YDGYHLFDYWGQGLTVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGDRTVISCRAEQGISNHLNWFQQKPGKAPKPLIYYTSNQLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQQYSNLPYTFGGGTKVEIK

830	BCMA-83 HL x CD3 HL	BC E5-33-B11-B8 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMDSLRAEDTAVYYCVRHGYDGYHLLFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDRTVITCRASQGISNHLNWIYQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLPEDFATYYCQQYSNLPYTFGGGTKEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGSGGTVVTTQEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGKAAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
831	BCMA-84	BC E5-33-G12-B8	VH CDR1	aa	NFDMA
832	BCMA-84	BC E5-33-G12-B8	VH CDR2	aa	SITTTGGGDTYYADSVKGG
833	BCMA-84	BC E5-33-G12-B8	VH CDR3	aa	HGYDGYHLLFDY
834	BCMA-84	BC E5-33-G12-B8	VL CDR1	aa	RASQGISNHLN
835	BCMA-84	BC E5-33-G12-B8	VL CDR2	aa	YTSNLQS
836	BCMA-84	BC E5-33-G12-B8	VL CDR3	aa	QQYSNLPYT
837	BCMA-84	BC E5-33-G12-B8	VH	aa	EVQLVESGGGLVQPGRSLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLLFDYWGQGLVTVSS
838	BCMA-84	BC E5-33-G12-B8	VL	aa	DIQMTQSPSSLSASVGERVTITCRASQGISNHLNWIYQKPGKAPKSLIYYTSNLTQSGVPSRFSGSGSGTDFTLTITSSLPEDFATYYCQQYSNLPYTFGGGTKEIK
839	BCMA-84	BC E5-33-G12-B8	scFv	aa	EVQLVESGGGLVQPGRSLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLLFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGERVTITCRASQGISNHLNWIYQKPGKAPKSLIYYTSNLTQSGVPSRFSGSGSGTDFTLTITSSLPEDFATYYCQQYSNLPYTFGGGTKEIK
840	BCMA-84 HL x CD3 HL	BC E5-33-G12-B8 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGRSLRLSCAASGFTFSNFDMAWVRQAPAKGLEWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYDGYHLLFDYWGQGLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGERVTITCRASQGISNHLNWIYQKPGKAPKSLIYYTSNLTQSGVPSRFSGSGSGTDFTLTITSSLPEDFATYYCQQYSNLPYTFGGGTKEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN

					NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGGGGGSGQT VVTQEPSLTVS PGGTVTLLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
841	BCMA-85	BC C6-97- G5	VH CDR1	aa	NFGMN
842	BCMA-85	BC C6-97- G5	VH CDR2	aa	WINTYTGESIYADDFKG
843	BCMA-85	BC C6-97- G5	VH CDR3	aa	GGVYGGYDAMDY
844	BCMA-85	BC C6-97- G5	VL CDR1	aa	RASQDISNYLN
845	BCMA-85	BC C6-97- G5	VL CDR2	aa	YTSRLHS
846	BCMA-85	BC C6-97- G5	VL CDR3	aa	QQGNTLPWT
847	BCMA-85	BC C6-97- G5	VH	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLTVTVSS
848	BCMA-85	BC C6-97- G5	VL	aa	DIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPKDKAPKLLIYYTSRLLHSGVPSRFSGSGS GTDYTLTISSELEPEDIAITYYCQQGNTLPWTFGQGTKVEIK
849	BCMA-85	BC C6-97- G5	scFv	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLTVTVSSGGGGGGSGGGG GSDIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPKDKAPKLLIYYTSRLLHSGVPSRFSGSGS GSGTDYTLTISSELEPEDIAITYYCQQGNTLPWTFGQGTKVEIK
850	BCMA-85 HL x CD3 HL	BC C6-97- G5 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLTVTVSSGGGGGGSGGGG GSDIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPKDKAPKLLIYYTSRLLHSGVPSRFSGSGS GSGTDYTLTISSELEPEDIAITYYCQQGNTLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGGGGGSGQT VVTQEPSLTVS PGGTVTLLTCGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
851	BCMA-86	BC C6-98- C8	VH CDR1	aa	NFGMN

852	BCMA-86	BC C6-98-C8	VH CDR2	aa	WINTYTGESIIYADDFKG
853	BCMA-86	BC C6-98-C8	VH CDR3	aa	GGVYGGYDAMDY
854	BCMA-86	BC C6-98-C8	VL CDR1	aa	RASQDISNYLN
855	BCMA-86	BC C6-98-C8	VL CDR2	aa	YTSRLHS
856	BCMA-86	BC C6-98-C8	VL CDR3	aa	QQGNTLPWT
857	BCMA-86	BC C6-98-C8	VH	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMMGWINTYTGESIIYADDFKGR FVFSSDTSVSTAYLQINSLKAEDTAVYFCARGGVYGGYDAMDYWGQGTLLVTVSSSSGGGGSGGGGG
858	BCMA-86	BC C6-98-C8	VL	aa	DIQMTQTPSSLSASVGDRTVITCRASQDISNYLNWYQQKPGKALKLLIYYTSRLLHSGVPSRFSGSGS GTDYSLTISNLQPEDIIATYYCQQGNTLPWTFGQGTKEIK
859	BCMA-86	BC C6-98-C8	scFv	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMMGWINTYTGESIIYADDFKGR FVFSSDTSVSTAYLQINSLKAEDTAVYFCARGGVYGGYDAMDYWGQGTLLVTVSSSSGGGGSGGGGG GSDIQMTQTPSSLSASVGDRTVITCRASQDISNYLNWYQQKPGKALKLLIYYTSRLLHSGVPSRFSGSGS GSGTDYSLTISNLQPEDIIATYYCQQGNTLPWTFGQGTKEIK
860	BCMA-86 HL x CD3 HL	BC C6-98-C8 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMMGWINTYTGESIIYADDFKGR FVFSSDTSVSTAYLQINSLKAEDTAVYFCARGGVYGGYDAMDYWGQGTLLVTVSSSSGGGGSGGGGG GSDIQMTQTPSSLSASVGDRTVITCRASQDISNYLNWYQQKPGKALKLLIYYTSRLLHSGVPSRFSGSGS GSGTDYSLTISNLQPEDIIATYYCQQGNTLPWTFGQGTKEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSSSGGGGSGGGGQTAVTQEPSTVS PGGTVTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
861	BCMA-87	BC C6-97-A6	VH CDR1	aa	NFGMN
862	BCMA-87	BC C6-97-A6	VH CDR2	aa	WINTYTGESIIYADDFKG
863	BCMA-87	BC C6-97-A6	VH CDR3	aa	GGVYGGYDAMDY
864	BCMA-87	BC C6-97-A6	VL CDR1	aa	RASQDISNYLN

865	BCMA-87	BC C6-97-A6	VL CDR2	aa	YTSRLHS	
866	BCMA-87	BC C6-97-A6	VL CDR3	aa	Q̄QGNLTLPWT	
867	BCMA-87	BC C6-97-A6	VH	aa	QVQLVQSGSELKKGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVGGYDAMDYWGQGTLLVTVSS	
868	BCMA-87	BC C6-97-A6	VL	aa	DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPKGAPKLLIYYTSRLLHSGVPSRFSGSGS GTDYTLTISSELEQEDIATYFCQ̄Q̄GNLTLPWTFGQ̄GTKVEIK	
869	BCMA-87	BC C6-97-A6	scFv	aa	QVQLVQSGSELKKGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVGGYDAMDYWGQGTLLVTVSSGGGGGGGGGG GSDIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPKGAPKLLIYYTSRLLHSGVPSRFSGSGS GSGTDYTLTISSELEQEDIATYFCQ̄Q̄GNLTLPWTFGQ̄GTKVEIK	
870	BCMA-87 HL x CD3 HL	BC C6-97-A6 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGSELKKGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVGGYDAMDYWGQGTLLVTVSSGGGGGGGGGG GSDIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPKGAPKLLIYYTSRLLHSGVPSRFSGSGS GSGTDYTLTISSELEQEDIATYFCQ̄Q̄GNLTLPWTFGQ̄GTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGGGGGGGGQT VVTQEP S LTVS PGGT VTLTCGSS TGA VTS GNY PNVVQ̄Q̄KPGQ̄APRGLIGG TKFLAPGT PARFSGSLLGGKAALTLSGV Q̄PEDEAEYYCVLWYSNRWVFGGGTKLTVL	
871	BCMA-88	BC C6-98-C8-E3	VH CDR1	aa	NFGMN	
872	BCMA-88	BC C6-98-C8-E3	VH CDR2	aa	WINTYTGESIYADDFKG	
873	BCMA-88	BC C6-98-C8-E3	VH CDR3	aa	GGVYGGYDAMDY	
874	BCMA-88	BC C6-98-C8-E3	VL CDR1	aa	RASQDISNYLN	
875	BCMA-88	BC C6-98-C8-E3	VL CDR2	aa	YTSRLHS	
876	BCMA-88	BC C6-98-C8-E3	VL CDR3	aa	Q̄SFATLTPWT	
877	BCMA-88	BC C6-98-C8-E3	VH	aa	QVQLVQSGSELKKGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR	

					FVFSSDTSVSTAYLQINSLKAEDTAVYFCARGGVGGYDAMDYWGQGLVTVSS
878	BCMA-88	BC C6-98- C8-E3	VL	aa	DIQMTQTPSSLSASVGDVRTITCRASQDISNYLNWYQQKPGKALKLLIYYTSRLHSGVPSRFSGSGS GTDYSLTISNLQPEDIAIYYCQSFATLPWTFGQGTKVEIK
879	BCMA-88	BC C6-98- C8-E3	scFv	aa	QVQLVQSGSELKKPGASVKVSCKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSSDTSVSTAYLQINSLKAEDTAVYFCARGGVGGYDAMDYWGQGLVTVSSGGGGSGGGSGGG GSDIQMTQTPSSLSASVGDVRTITCRASQDISNYLNWYQQKPGKALKLLIYYTSRLHSGVPSRFSGSGS GSGTDYSLTISNLQPEDIAIYYCQSFATLPWTFGQGTKVEIK
880	BCMA-88 HL x CD3 HL	BC C6-98- C8-E3 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGSELKKPGASVKVSCKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSSDTSVSTAYLQINSLKAEDTAVYFCARGGVGGYDAMDYWGQGLVTVSSGGGGSGGGSGGG GSDIQMTQTPSSLSASVGDVRTITCRASQDISNYLNWYQQKPGKALKLLIYYTSRLHSGVPSRFSGSGS GSGTDYSLTISNLQPEDIAIYYCQSFATLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGGSGGTVVVTQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEIYCVLWYSNRWVFGGGTKLTVL
881	BCMA-89	BC C6-98- A1-E3	VH CDR1	aa	NFGMN
882	BCMA-89	BC C6-98- A1-E3	VH CDR2	aa	WINTYTGESIYADDFKG
883	BCMA-89	BC C6-98- A1-E3	VH CDR3	aa	GGVYGGYDAMDY
884	BCMA-89	BC C6-98- A1-E3	VL CDR1	aa	RASQDISNYLN
885	BCMA-89	BC C6-98- A1-E3	VL CDR2	aa	YTSRLHS
886	BCMA-89	BC C6-98- A1-E3	VL CDR3	aa	QSFATLPWT
887	BCMA-89	BC C6-98- A1-E3	VH	aa	QVQLVQSGSELKKPGASVKISCKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSSDTSVSTAYLQINNLLKAEDTAVYYCARGGVGGYDAMDYWGQGLVTVSS
888	BCMA-89	BC C6-98- A1-E3	VL	aa	DIQMTQSPSSLSASVGDVRTISCRASQDISNYLNWYQQKPGKAPKLLIYYTSRLHSGVPSRFSGSGS GTDYTFITISNLQPEDIAIYYCQSFATLPWTFGQGTKVEIK
889	BCMA-89	BC C6-98- A1-E3	scFv	aa	QVQLVQSGSELKKPGASVKISCKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR

890	BCMA-89 HL x CD3 HL	BC C6-98-A1-E3 HL x CD3 HL	bispecific molecule	aa	EVFSSDTSVSTAYLQINNLLKAEDTAVYYCARGGVYGGYDAMDYWGQGLVTVSSGGGGGGGGGGGGGSDIQMTQSPSSLSASVGDRTVITCRASQDISNYLNWYQQKPKAPKLLIYYTSRLHSGVPSRFSGSGSGTDYTFITISNLQPEDIAIYYCQSFATLPWTFGQGTKVEIK
891	BCMA-90	BC C6-97-G5-E3	VH CDR1	aa	QVQLVQSGSELKKPGASVKISCKASGYTFITNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR
892	BCMA-90	BC C6-97-G5-E3	VH CDR2	aa	FVFSSDTSVSTAYLQINNLLKAEDTAVYYCARGGVYGGYDAMDYWGQGLVTVSSGGGGGGGGGGGGGSDIQMTQSPSSLSASVGDRTVITCRASQDISNYLNWYQQKPKAPKLLIYYTSRLHSGVPSRFSGSGSGTDYTFITISNLQPEDIAIYYCQSFATLPWTFGQGTKVEIK
893	BCMA-90	BC C6-97-G5-E3	VH CDR3	aa	QVQLVQSGSELKKPGASVKISCKASGYTFITNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR
894	BCMA-90	BC C6-97-G5-E3	VL CDR1	aa	LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN
895	BCMA-90	BC C6-97-G5-E3	VL CDR2	aa	NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGGGGGGGGGQTVVTQEPSLTVSPGGTVTLTCGSSGTGAVTSGNYPNWVQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV
896	BCMA-90	BC C6-97-G5-E3	VL CDR3	aa	QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
897	BCMA-90	BC C6-97-G5-E3	VH	aa	NFGMN
898	BCMA-90	BC C6-97-G5-E3	VL	aa	WINTYTGESIYADDFKG
899	BCMA-90	BC C6-97-G5-E3	scFv	aa	GGVYGGYDAMDY
900	BCMA-90 HL x CD3 HL	BC C6-97-G5-E3 HL x CD3 HL	bispecific molecule	aa	RASQDISNYLN
				aa	YTSRLHS
				aa	QSFATLPWT
				aa	QVQLVQSGSELKKPGASVKVSKKASGYTFITNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR
				aa	FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLVTVSS
				aa	DIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPDKAPKLLIYYTSRLHSGVPSRFSGSGSGTDYTLTISLLEPEDIAIYYCQSFATLPWTFGQGTKVEIK
				aa	QVQLVQSGSELKKPGASVKVSKKASGYTFITNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR
				aa	FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLVTVSSGGGGGGGGGGGGGSDIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPDKAPKLLIYYTSRLHSGVPSRFSGSGSGTDYTLTISLLEPEDIAIYYCQSFATLPWTFGQGTKVEIK
				aa	QVQLVQSGSELKKPGASVKVSKKASGYTFITNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR
				aa	FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLVTVSSGGGGGGGGGGGGGSDIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPDKAPKLLIYYTSRLHSGVPSRFSGSGSGTDYTLTISLLEPEDIAIYYCQSFATLPWTFGQGTKVEIK
				aa	QVQLVQSGSELKKPGASVKVSKKASGYTFITNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR
				aa	FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLVTVSSGGGGGGGGGGGGGSDIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPDKAPKLLIYYTSRLHSGVPSRFSGSGSGTDYTLTISLLEPEDIAIYYCQSFATLPWTFGQGTKVEIK

					GSDIQMTQSPSSLSASLGD RVTITCRASQDISNYLNWYQQKPDKAPKLLIYYTSRLHSGVPSRFSGSGGTDYTLTISSELPEDIAYYCQSFATLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGQTVVTVTQEPSLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQOKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYCYVLWYSNRWVFGGGTKLTVL
901	BCMA-91	BC C6-97-A6-E3	VH CDR1	aa	NFGMN
902	BCMA-91	BC C6-97-A6-E3	VH CDR2	aa	WINTYTGESIYADDFKG
903	BCMA-91	BC C6-97-A6-E3	VH CDR3	aa	GGVYGGYDAMDY
904	BCMA-91	BC C6-97-A6-E3	VL CDR1	aa	RASQDISNYLN
905	BCMA-91	BC C6-97-A6-E3	VL CDR2	aa	YTSRLHS
906	BCMA-91	BC C6-97-A6-E3	VL CDR3	aa	QSFATLPWT
907	BCMA-91	BC C6-97-A6-E3	VH	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGRFVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLTVTVSS
908	BCMA-91	BC C6-97-A6-E3	VL	aa	DIQMTQSPSSLSASVGD RVTITCRASQDISNYLNWYQQKPGKAPKLLIYYTSRLHSGVPSRFSGSGSGTDYTLTISSELEQEDIAYFCQSFATLPWTFGQGTKVEIK
909	BCMA-91	BC C6-97-A6-E3	scFv	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGRFVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLTVTVSSGGGGSGGGSGGGSDIQMTQSPSSLSASVGD RVTITCRASQDISNYLNWYQQKPGKAPKLLIYYTSRLHSGVPSRFSGSGSGTDYTLTISSELEQEDIAYFCQSFATLPWTFGQGTKVEIK
910	BCMA-91 HL x CD3 HL	BC C6-97-A6-E3 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGRFVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLTVTVSSGGGGSGGGSGGGSDIQMTQSPSSLSASVGD RVTITCRASQDISNYLNWYQQKPGKAPKLLIYYTSRLHSGVPSRFSGSGSGTDYTLTISSELEQEDIAYFCQSFATLPWTFGQGTKVEIKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGQTVVTVTQEPSLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQOKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV

						QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
911	BCMA-92	BC C6-97- G5-G9	VH CDR1	aa		NFGMN
912	BCMA-92	BC C6-97- G5-G9	VH CDR2	aa		WINTYTGESIYADDFKG
913	BCMA-92	BC C6-97- G5-G9	VH CDR3	aa		GGVYGGYDAMDY
914	BCMA-92	BC C6-97- G5-G9	VL CDR1	aa		RASQDISNYLN
915	BCMA-92	BC C6-97- G5-G9	VL CDR2	aa		YTSRLHS
916	BCMA-92	BC C6-97- G5-G9	VL CDR3	aa		QHFRTLPWTT
917	BCMA-92	BC C6-97- G5-G9	VH	aa		QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLTGLTVSS
918	BCMA-92	BC C6-97- G5-G9	VL	aa		DIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPDKAPKLLIYYTSRLLHSGVPSRFSGSGS GTDYTLTISSLEPEDIAITYYCQHFRTLPWTFGQGTKVEIK
919	BCMA-92	BC C6-97- G5-G9	scFv	aa		QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLTGLTVSSGGGGSGGGGG GSDIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPDKAPKLLIYYTSRLLHSGVPSRFSGSGS GGTDYTLTISSLEPEDIAITYYCQHFRTLPWTFGQGTKVEIK
920	BCMA-92 HL x CD3 HL	BC C6-97- G5-G9 HL x CD3 HL	bispecific molecule	aa		QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVYGGYDAMDYWGQGLTGLTVSSGGGGSGGGGG GSDIQMTQSPSSLSASLGDRVTITCRASQDISNYLNWYQQKPDKAPKLLIYYTSRLLHSGVPSRFSGSGS GGTDYTLTISSLEPEDIAITYYCQHFRTLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGGGGSGQTVVTQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
921	BCMA-93	BC C6-98- C8-G9	VH CDR1	aa		NFGMN
922	BCMA-93	BC C6-98- C8-G9	VH CDR2	aa		WINTYTGESIYADDFKG

923	BCMA-93	BC C6-98- C8-G9	VH CDR3	aa	GGVYGGYDAMDY
924	BCMA-93	BC C6-98- C8-G9	VL CDR1	aa	RASQDISNYLN
925	BCMA-93	BC C6-98- C8-G9	VL CDR2	aa	YTSRLHS
926	BCMA-93	BC C6-98- C8-G9	VL CDR3	aa	QHFRTLPWT
927	BCMA-93	BC C6-98- C8-G9	VH	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSSDTSVSTAYLQINSLKAEDTAVYFCARGGVYGGYDAMDYWGQGTLLVTVSS
928	BCMA-93	BC C6-98- C8-G9	VL	aa	DIQMTQTPSSLSASVGDRTITTCRASQDISNYLNWYQQKPGKALKLLIYYTSRLHSGVPSRFSGSGS GTDYSLTISNLQPEDIATYYCQHFRTLPWTFGQGTKVEIK
929	BCMA-93	BC C6-98- C8-G9	scFv	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSSDTSVSTAYLQINSLKAEDTAVYFCARGGVYGGYDAMDYWGQGTLLVTVSSGGGGSGGGGG GSDIQMTQTPSSLSASVGDRTITTCRASQDISNYLNWYQQKPGKALKLLIYYTSRLHSGVPSRFSGSGS GSGTDYSLTISNLQPEDIATYYCQHFRTLPWTFGQGTKVEIK
930	BCMA-93 HL x CD3 HL	BC C6-98- C8-G9 HL x CD3 HL	bispecific molecule	aa	QVQLVQSGSELKKPGASVKVSKKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR FVFSSDTSVSTAYLQINSLKAEDTAVYFCARGGVYGGYDAMDYWGQGTLLVTVSSGGGGSGGGGG GSDIQMTQTPSSLSASVGDRTITTCRASQDISNYLNWYQQKPGKALKLLIYYTSRLHSGVPSRFSGSGS GSGTDYSLTISNLQPEDIATYYCQHFRTLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGGSGQTVVTVQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
931	BCMA-94	BC C6-97- A6-G9	VH CDR1	aa	NFGMN
932	BCMA-94	BC C6-97- A6-G9	VH CDR2	aa	WINTYTGESIYADDFKG
933	BCMA-94	BC C6-97- A6-G9	VH CDR3	aa	GGVYGGYDAMDY
934	BCMA-94	BC C6-97- A6-G9	VL CDR1	aa	RASQDISNYLN
935	BCMA-94	BC C6-97- A6-G9	VL CDR2	aa	YTSRLHS

936	BCMA-94	BC C6-97- A6-G9	VL CDR3	aa	QHFRTLPWTT	QVQLVQSGSELEKPKGASVKVSCKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVGGYDAMDYWGQGTLLVTVSS
937	BCMA-94	BC C6-97- A6-G9	VH	aa		DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPKGAPKLLIYYTSRRLHSGVPSRFSGSGS GTDYTLTISSLEQEDIATYFCQHFRTLPWTFGQGTKVEIK
938	BCMA-94	BC C6-97- A6-G9	VL	aa		QVQLVQSGSELEKPKGASVKVSCKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVGGYDAMDYWGQGTLLVTVSSGGGGSGGGGG GSDIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPKGAPKLLIYYTSRRLHSGVPSRFSGSGS GGTDYTLTISSLEQEDIATYFCQHFRTLPWTFGQGTKVEIK
939	BCMA-94	BC C6-97- A6-G9	scFv	aa		QVQLVQSGSELEKPKGASVKVSCKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVGGYDAMDYWGQGTLLVTVSSGGGGSGGGGG GSDIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPKGAPKLLIYYTSRRLHSGVPSRFSGSGS GGTDYTLTISSLEQEDIATYFCQHFRTLPWTFGQGTKVEIK
940	BCMA-94 HL x CD3 HL	BC C6-97- A6-G9 HL x CD3 HL	bispecific molecule	aa		QVQLVQSGSELEKPKGASVKVSCKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIIYADDFKGR FVFSLDTSVTTAYLQINSLKDEDTAVYYCARGGVGGYDAMDYWGQGTLLVTVSSGGGGSGGGGG GSDIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPKGAPKLLIYYTSRRLHSGVPSRFSGSGS GGTDYTLTISSLEQEDIATYFCQHFRTLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGGSGQTVVTQEPSTLTVS PGGTVTLTCSGSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGKAAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
941	BCMA-95	BC C6-98- A1-G9	VH CDR1	aa	NFGMN	
942	BCMA-95	BC C6-98- A1-G9	VH CDR2	aa	WINTYTGESIIYADDFKG	
943	BCMA-95	BC C6-98- A1-G9	VH CDR3	aa	GGVYGGYDAMDY	
944	BCMA-95	BC C6-98- A1-G9	VL CDR1	aa	RASQDISNYLN	
945	BCMA-95	BC C6-98- A1-G9	VL CDR2	aa	YTSRLLHS	
946	BCMA-95	BC C6-98- A1-G9	VL CDR3	aa	QHFRTLPWTT	
947	BCMA-95	BC C6-98- A1-G9	VH	aa		QVQLVQSGSELEKPKGASVKISCKASGYTFTNFGMNWVRQAPGQGLEWMGWINTYTGESIIYADDFKGR FVFSDDTSVSTAYLQINNLLKAEDTAVYYCARGGVGGYDAMDYWGQGTLLVTVSS
948	BCMA-95	BC C6-98- A1-G9	VL	aa		DIQMTQSPSSLSASVGDRVTISCRASQDISNYLNWYQQKPKGAPKLLIYYTSRRLHSGVPSRFSGSGS

		A1-G9				GTDYFTTISNLQPEDIAITYFCQHFRTLPWTFGQGTKVEIK
949	BCMA-95	BC C6-98-A1-G9	scFv	aa		QVQLVQSGSELKKPGASVKISCKASGYFTTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGRFVFSSDTSVSTAYLQINNLKAEDTAVYYCARGGVYGGYDAMDYWGQGTLLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDVRTISCRASQDISNYLNWYQQKPGKAPKLLIYYTSRLLHSGVPSRFSGSGSGTDYFTTISNLQPEDIAITYFCQHFRTLPWTFGQGTKVEIK
950	BCMA-95 HL x CD3 HL	BC C6-98-A1-G9 HL x CD3 HL	bispecific molecule	aa		QVQLVQSGSELKKPGASVKISCKASGYFTTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGRFVFSSDTSVSTAYLQINNLKAEDTAVYYCARGGVYGGYDAMDYWGQGTLLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDVRTISCRASQDISNYLNWYQQKPGKAPKLLIYYTSRLLHSGVPSRFSGSGSGTDYFTTISNLQPEDIAITYFCQHFRTLPWTFGQGTKVEIKSGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGSGGGSQTVVTQEPSLTVSPGGTVTLTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYCYVLWYSNRWVFGGGTKLTVL
951	BCMA-96	BC C6 98-A1	VH CDR1	aa		NFGMN
952	BCMA-96	BC C6 98-A1	VH CDR2	aa		WINTYTGESIYADDFKG
953	BCMA-96	BC C6 98-A1	VH CDR3	aa		GGVYGGYDAMDY
954	BCMA-96	BC C6 98-A1	VL CDR1	aa		RASQDISNYLN
955	BCMA-96	BC C6 98-A1	VL CDR2	aa		YTSRLHS
956	BCMA-96	BC C6 98-A1	VL CDR3	aa		QQGNTLTPWT
957	BCMA-96	BC C6 98-A1	VH	aa		QVQLVQSGSELKKPGASVKISCKASGYFTTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGRFVFSSDTSVSTAYLQINNLKAEDTAVYYCARGGVYGGYDAMDYWGQGTLLVTVSS
958	BCMA-96	BC C6 98-A1	VL	aa		DIQMTQSPSSLSASVGDVRTISCRASQDISNYLNWYQQKPGKAPKLLIYYTSRLLHSGVPSRFSGSGSGTDYFTTISNLQPEDIAITYYCQQGNTLPWTFGQGTKVEIK
959	BCMA-96	BC C6 98-A1	scFv	aa		QVQLVQSGSELKKPGASVKISCKASGYFTTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGRFVFSSDTSVSTAYLQINNLKAEDTAVYYCARGGVYGGYDAMDYWGQGTLLVTVSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDVRTISCRASQDISNYLNWYQQKPGKAPKLLIYYTSRLLHSGVPSRFSGSGSGTDYFTTISNLQPEDIAITYYCQQGNTLPWTFGQGTKVEIK
960	BCMA-96 HL x	BC C6 98-	bispecific	aa		QVQLVQSGSELKKPGASVKISCKASGYFTTNFGMNWVRQAPGQGLEWMGWINTYTGESIYADDFKGR

	CD3 HL	A1 HL x CD3 HL	molecule		EVFSSDTSVSTAYLQINNLLKAEDTAVYYCARGGVYGGYDAMDYWGQGTLLVTVSSGGGGSGGGSGGG GSDIQMTQSPSSLSASVGDRTVITSCRASQDISNVLNWIYQQKPGKAPKLLIYYTSRLHSGVPSRFSGS GSGTDYTFITISNLQPEDIAIYCCQQGNTLPWTFGQGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGGSGQTIVVTQEPSLTVS PGGTVTLTTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
961	BCMA-97	BC B12-33- G2-B2	VH CDR1	aa	NFDMA
962	BCMA-97	BC B12-33- G2-B2	VH CDR2	aa	SITTTGGGDTYYADSVKKG
963	BCMA-97	BC B12-33- G2-B2	VH CDR3	aa	HGYDDGYHLFDY
964	BCMA-97	BC B12-33- G2-B2	VL CDR1	aa	RASQGISNNLN
965	BCMA-97	BC B12-33- G2-B2	VL CDR2	aa	YTSNLQS
966	BCMA-97	BC B12-33- G2-B2	VL CDR3	aa	QQFTSLPYT
967	BCMA-97	BC B12-33- G2-B2	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYVDGYHLFDYWGQGTLLVTVSS
968	BCMA-97	BC B12-33- G2-B2	VL	aa	DIQMTQSPSSMSASVGDRTVITCRASQGISNNLNWIYQQKPGKAPKSLIYYTSNLQSGVPSRFSGS GTDYTLTITSSLQPEDFATYYCQQFTSLPYTFGQGTKLEIK
969	BCMA-97	BC B12-33- G2-B2	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYVDGYHLFDYWGQGTLLVTVSSGGGGSGGGSGGG GSDIQMTQSPSSMSASVGDRTVITCRASQGISNNLNWIYQQKPGKAPKSLIYYTSNLQSGVPSRFSGS GSGTDYTLTITSSLQPEDFATYYCQQFTSLPYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGGSGQTIVVTQEPSLTVS PGGTVTLTTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV
970	BCMA-97 HL x CD3 HL	BC B12-33- G2-B2 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGR FTISRDNAKNTLYLQMNSLRAEDTAVYYCVRHGYVDGYHLFDYWGQGTLLVTVSSGGGGSGGGSGGG GSDIQMTQSPSSMSASVGDRTVITCRASQGISNNLNWIYQQKPGKAPKSLIYYTSNLQSGVPSRFSGS GSGTDYTLTITSSLQPEDFATYYCQQFTSLPYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGSGGGGSGQTIVVTQEPSLTVS PGGTVTLTTCGSSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGV

						QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
971	BCMA-98	BC B12-33-A4-B2	VH CDR1	aa		NFDMA
972	BCMA-98	BC B12-33-A4-B2	VH CDR2	aa		SITTTGGGDTYYADSVKG
973	BCMA-98	BC B12-33-A4-B2	VH CDR3	aa		HGYDGYHLFDY
974	BCMA-98	BC B12-33-A4-B2	VL CDR1	aa		RANQGISNNLN
975	BCMA-98	BC B12-33-A4-B2	VL CDR2	aa		YTSNLQS
976	BCMA-98	BC B12-33-A4-B2	VL CDR3	aa		QQFTSLPYT
977	BCMA-98	BC B12-33-A4-B2	VH	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKSTLYLQMDSLRSEDTAVYYCVRHGYDGYHLFDYWGQGLVTVSS
978	BCMA-98	BC B12-33-A4-B2	VL	aa		DIQMTQSPSSLSASVGDRVTITCRANQGISNNLNWYQQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLPQPEDFATYYCQQFTSLPYTFGQGTKLEIK
979	BCMA-98	BC B12-33-A4-B2	scFv	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKSTLYLQMDSLRSEDTAVYYCVRHGYDGYHLFDYWGQGLVTVSSGGGGSGGGGGGGSDIQMTQSPSSLSASVGDRVTITCRANQGISNNLNWYQQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLPQPEDFATYYCQQFTSLPYTFGQGTKLEIK
980	BCMA-98 HL x CD3 HL	BC B12-33-A4-B2 HL x CD3 HL	bispecific molecule	aa		EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKSTLYLQMDSLRSEDTAVYYCVRHGYDGYHLFDYWGQGLVTVSSGGGGSGGGGGGGSDIQMTQSPSSLSASVGDRVTITCRANQGISNNLNWYQQKPGKAPKPLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLPQPEDFATYYCQQFTSLPYTFGQGTKLEIKKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGGSQT VVTQEPSLT VSPGGTVTLTCGSSGTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
981	BCMA-99	BC B12-33-A5-B2	VH CDR1	aa		NFDMA
982	BCMA-99	BC B12-33-A5-B2	VH CDR2	aa		SITTTGGGDTYYADSVKG

983	BCMA-99	BC B12-33-A5-B2	VH CDR3	aa	HGYDGYHLFDY
984	BCMA-99	BC B12-33-A5-B2	VL CDR1	aa	RASQGISNNLN
985	BCMA-99	BC B12-33-A5-B2	VL CDR2	aa	YTSNLQS
986	BCMA-99	BC B12-33-A5-B2	VL CDR3	aa	QQFTSLPYT
987	BCMA-99	BC B12-33-A5-B2	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRSEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSS
988	BCMA-99	BC B12-33-A5-B2	VL	aa	DIQMTQSPSSMSASVGDRTVITCRASQGISNNLNWYQQKPGKAPKSLIYYTSNLSQGVPSRFSGSGS GTDYTLTISSLQPEDFATYYCQQFTSLPYTFGQGTKLEIK
989	BCMA-99	BC B12-33-A5-B2	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRSEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSSGGGGGGGGGG GSDIQMTQSPSSMSASVGDRTVITCRASQGISNNLNWYQQKPGKAPKSLIYYTSNLSQGVPSRFSGSGS GGTDYTLTISSLQPEDFATYYCQQFTSLPYTFGQGTKLEIK
990	BCMA-99 HL x CD3 HL	BC B12-33-A5-B2 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTGGGDTYYADSVKGR FTISRDNAKNTLYLQMDSLRSEDTAVYYCVRHGYDGYHLFDYWGQGTLLVTVSSGGGGGGGGGG GSDIQMTQSPSSMSASVGDRTVITCRASQGISNNLNWYQQKPGKAPKSLIYYTSNLSQGVPSRFSGSGS GGTDYTLTISSLQPEDFATYYCQQFTSLPYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWNVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLLVTVSSGGGGGGGGGGQTIVVTQEPSLTVS PGGTVTLTGSSSTGAVTSGNYPNWWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLGKGKAAALTLSGV QPEDEAEYYCVLWYSNRWVFGGGTKLTVL
991	BCMA-100	BC B12-33-A5-C10	VH CDR1	aa	NFDMA
992	BCMA-100	BC B12-33-A5-C10	VH CDR2	aa	SITTTGGGDTYYADSVKG
993	BCMA-100	BC B12-33-A5-C10	VH CDR3	aa	HGYDGYHLFDY
994	BCMA-100	BC B12-33-A5-C10	VL CDR1	aa	RASQGISNNLN
995	BCMA-100	BC B12-33-A5-C10	VL CDR2	aa	YTSNLQS

996	BCMA-100	BC B12-33-A5-C10	VL CDR3	aa	QQFAHLPYT
997	BCMA-100	BC B12-33-A5-C10	VH	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMDSLRS EDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSS
998	BCMA-100	BC B12-33-A5-C10	VL	aa	DIQMTQSPSSMSASVGDRVTITCRASQGISNNLNWYQQKPKGKAPKSLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCQQFAHLPYTFGQGTKLEIK
999	BCMA-100	BC B12-33-A5-C10	scFv	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMDSLRS EDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSSGGGGSGGGSGGGGGS DIQMTQSPSSMSASVGDRVTITCRASQGISNNLNWYQQKPKGKAPKSLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCQQFAHLPYTFGQGTKLEIK
1000	BCMA-100 HL x CD3 HL	BC B12-33-A5-C10 HL x CD3 HL	bispecific molecule	aa	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNFDMAWVRQAPGKGLVWVSSITTTGGGDTYYADSVKGRFTISRDNAKNTLYLQMDSLRS EDTAVYYCVRHGY YDGYHLFDYWGQGLVTVSSGGGGSGGGSGGGGGS DIQMTQSPSSMSASVGDRVTITCRASQGISNNLNWYQQKPKGKAPKSLIYYTSNLTQSGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCQQFAHLPYTFGQGTKLEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMN NLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVTVSSGGGGSGGGSGGGSGQT VVTQEPSLT VSPGGTVTLTCGSSTGAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGKKAALTLSGVQPEDEAEY YCVLWY SNRWVFGGGTKLTVL
1001	human BCMA		human	na	atgttgcatggctgggcagtgctcccaaaatgaataatttgacagtttgttgcatgttgcttgcatatcctgtgtaacttcgatgttcttctaatactcctcctcctaacatgtcagcgttatgtgaatgcaagtgtgaccaattcagtgaaaggaaacgaatgcgattctctggacctgttgggactgagcttaataatttctttggcagtttctgtgctaattgttttgctaaggaaataaactctgaaccattaaaggacgagtttaaaaacacaggatcaggctctcctgggcattggctaacattgacctggaaaagacgaggaactggtgatgaattattcttccgagaggcctcgagtacacggtggaagaatgcacctgtgaagactgcatcaagagcaaccgaaaggtcgactctgaccttgcttccactccagctatggaggaaaggcgcaaccattcttgtcaccacgaaaacgaatgactattgcaagagcctgccagctgcttggagtgcctacggagatagagaaatcaatttctgctaggtaa
1002	human BCMA		human	aa	MLQMAGQCSQNEYFDSL LHACIPQLRCSSNTPPLTCQRYCNASVTNSVKGTNAILLWTC LGLSLIISLAVFVLMFLLRKINSEPLKDEFKNTGSGLLGMANIDLEKSRTGDEIILPRGLE YTVVEECTCEDCIKSKPKVDS DHCFPLPAMEEGATILVTTKTNDYCKSLPAALSATEIEKSISAR
1003	mouse BCMA		murine	na	atggcgcaacagtggttccacagtgaaataatttgacagtcctgctgcatgcttgcaaacccgtgtcacttgcgatgttccaaccctcctgcaacctgtcagccttactgtgatccaagcgtgaccagttcagtgaa

				agggacgtacacgggtgctctggatcttcttggggctgaccttggtcctctcttggcacttttcaca atctcattcttgctgaggaagatgaaccccgagccctgaaggacgacctcaagcccaggtcagc ttgacggatcggctcagctggacaagccgacaccgagctgactaggatcagggctggtgacgacag gatcttccccgaagcctggagtatacagtggaagagtgcacctgtgaggactgtgcaagagcaaa ccaagggggtctgaccatttcttccgcttcagccatggaggagggggaaccattcttgtca ccacaaaaacgggtgactacgggcaagtcaagtgtgccaaactgcttggcaagtgtcatggggatgga gaagccaaactcacactagataa
1004	mouse BCMA	murine	aa	MAQQCFHSEYFDSLHACKPCHLRCSNPPATCQPYCDPSVTSSVKGTYTVLWIFLGLTLVLSLALFT ISFLLRKMNPEALKDEPQSPGQLDGSAQLDKADTELTRIRAGDDRIFFPRSLEYTVEECTCEDCVKSK PKGDSDDHFFPLPAMEEGATILVTTKTGDYGKSSVPTALQSVMGMEKPTHTR
1005	macaque BCMA	rhesus	na	atgttgcatggctcggcagtgctcccaaaatgaataatttgacagtttgtgcatgattgcaaac cttgtcaacttcgatgttctagtactcctcctctaacatgtcagcgttattgcaatgcaagtatgac caattcagtgaaaggaatgaatcgattctctggacctgttgggactgagcttgataatttctttg gcagtttctgctaaacgtttttgctaaggaaagatgagctctgaaaccattaaaggatgagtttaaaa acacaggatcaggtctcctgggcattggttaacattgacctggaaaaaggcaggactggtgatgaaat tgttcttccaagagccctggagtacacgggtggaagaatgcacctgtgaagactgcatcaagaataaa ccaaggttgattctgaccattgcttccactccagcatggaggaaaggcgaaccattctcgtca ccacgaaaaacgaatgactattgcaatagcctgtcagctgcttgcaggtgttacggagatagagaaatc aatttctgctaggtaa
1006	macaque BCMA	rhesus	aa	MLQMARQCSQNEYFDSLHDKPCQLRCSSTPPLTCQRYCNASMTNSVKGMAIILWCLGLSLIISL AVFVLTFLLRKMSSEPLKDEFKNTGSGLLGMANIDLEKRTGDEIVLPRGLEYYTVEECTCEDCIKNK PKVDSDDHCFPLPAMEEGATILVTTKTNDYCNLSAALSVEIEKISAR
1007	hu BCMA ECD = positions 1-54 of SEQ ID NO: 1002	human	aa	MLQMAGQCSQNEYFDSLHACIPCQLRCSSTPPLTCQRYCNASVTNSVKGTNA
1008	mu BCMA ECD = positions 1-49 of SEQ ID NO: 1004	murine	aa	MAQQCFHSEYFDSLHACKPCHLRCSNPPATCQPYCDPSVTSSVKGTYT
1009	hu BCMA ECD / E1 murine	chimeric hu / mu	aa	MAQQCSQNEYFDSLHACIPCQLRCSSTPPLTCQRYCNASVTNSVKGTNA
1010	hu BCMA ECD / E2 murine	chimeric hu / mu	aa	MLQMAGQCFHSEYFDSLHACIPCQLRCSSTPPLTCQRYCNASVTNSVKGTNA
1011	hu BCMA ECD / E3 murine	chimeric hu / mu	aa	MLQMAGQCSQNEYFDSLHACIPCCHLRCSNPPATCQPYCNASVTNSVKGTNA
1012	hu BCMA ECD / E4 murine	chimeric hu / mu	aa	MLQMAGQCSQNEYFDSLHACIPCQLRCSSTPPLTCQRYCDPSVTSSVKGTYT

1013	hu BCMA ECD / E5 murine	chimeric hu / mu	aa	MLQMAGQCSQNEYFDLSLLHACKPCQLRCSSNTPPLTCQRYCNASVTNSVKGTNA
1014	hu BCMA ECD / E6 murine	chimeric hu / mu	aa	MLQMAGQCSQNEYFDLSLLHACIPCHLRCSSNTPPLTCQRYCNASVTNSVKGTNA
1015	hu BCMA ECD / E7 murine	chimeric hu / mu	aa	MLQMAGQCSQNEYFDLSLLHACIPCQLRCSSNTPPLTCQPYCNASVTNSVKGTNA
1016	hu BCMA epitope cluster 3	human	aa	CQLRCSSNTPPLTCQRYC
1017	mac BCMA epitope cluster 3	macaque	aa	CQLRCSSTPPLTCQRYC
1018	hu BCMA epitope cluster 1	human	aa	MLQMAGQ
1019	hu BCMA epitope cluster 4	human	aa	NASVTNSVKGTNA
1020	mac BCMA epitope cluster 1	macaque	aa	MLQMARQ
1021	mac BCMA epitope cluster 4	macaque	aa	NASMTNSVKGMNA
1022	BCMA-101	VH CDR1	aa	GFTFSNYDMA
1023	BCMA-101	VH CDR2	aa	SIITSGGDNYRDSVKG
1024	BCMA-101	VH CDR3	aa	HDYYDGSYGfAY
1025	BCMA-101	VL CDR1	aa	KASQSVGINVD
1026	BCMA-101	VL CDR2	aa	GASNRHT
1027	BCMA-101	VL CDR3	aa	LQYGSIPFT
1028	BCMA-101	VH	aa	EVQLVESGGGLVQPGRSLKLSCAASGFTFSNYDMAWVRQAPTkgLEWVASIIITSGGDNYRDSVKGR FTVSRDnAKSTLYLQMDSLRSEDtATYYCVRHDYYDGSYGfAYWGQGTlLVTVSS
1029	BCMA-101	VL	aa	ETVMTQSPtSMSTsIGERVtLNCKASQSVGINVDWYQQTPGQSPKLLIYGASNRHTGVPDRFTGSgF GRDFtLLtISNVEAE DLAVYYCLQYGSIPFTFGSGTKLELk
1030	BCMA-101	scFv	aa	EVQLVESGGGLVQPGRSLKLSCAASGFTFSNYDMAWVRQAPTkgLEWVASIIITSGGDNYRDSVKGR FTVSRDnAKSTLYLQMDSLRSEDtATYYCVRHDYYDGSYGfAYWGQGTlLVTVSSGGGGGGGGGGGG GGSEtVMTQSPtSMSTsIGERVtLNCKASQSVGINVDWYQQTPGQSPKLLIYGASNRHTGVPDRFTG SGfGRDFtLLtISNVEAE DLAVYYCLQYGSIPFTFGSGTKLELk

1031	BCMA-102	BC 244-A7	VH CDR1	aa	GYTFTNHIIH
1032	BCMA-102	BC 244-A7	VH CDR2	aa	YINPYNDDTEYNEKFKG
1033	BCMA-102	BC 244-A7	VH CDR3	aa	DGY YRDMDVMDY
1034	BCMA-102	BC 244-A7	VL CDR1	aa	RASQDISNYLN
1035	BCMA-102	BC 244-A7	VL CDR2	aa	YTSRLHS
1036	BCMA-102	BC 244-A7	VL CDR3	aa	QQGNTLLPWT
1037	BCMA-102	BC 244-A7	VH	aa	EVQLVEQSGPELVKPGASVKMSCKASGYTFTNHI IHWVKQKPGQGLEWIGYINPYNDDTEYNEKFKG KATLTSDKSSTTAYMELSSLTSEDSAVYYCARDGY YRDMDVMDYWGQGT'TVTVSS
1038	BCMA-102	BC 244-A7	VL	aa	ELVMTQTTPSSLSASLGDRVTTISCRASQDISNYLNWYQQKPDGTVKLLIYYTSRLHSGVPSRFSGSGS GTDYSLTISNLEQEDIATYFCQQQGNLTPWTFGGGTKLEIK
1039	BCMA-102	BC 244-A7	scFv	aa	EVQLVEQSGPELVKPGASVKMSCKASGYTFTNHI IHWVKQKPGQGLEWIGYINPYNDDTEYNEKFKG KATLTSDKSSTTAYMELSSLTSEDSAVYYCARDGY YRDMDVMDYWGQGT'TVTVSSGGSGGGSGG GGSELVMTQTTPSSLSASLGDRVTTISCRASQDISNYLNWYQQKPDGTVKLLIYYTSRLHSGVPSRFSG SGSGTDYSLTISNLEQEDIATYFCQQQGNLTPWTFGGGTKLEIK
1040	BCMA-103	BC 263-A4	VH CDR1	aa	GFTFSNYDMA
1041	BCMA-103	BC 263-A4	VH CDR2	aa	SISTRGDITSYRDSVKG
1042	BCMA-103	BC 263-A4	VH CDR3	aa	QDY YTDYMGFAY
1043	BCMA-103	BC 263-A4	VL CDR1	aa	RASEDIYNGLA
1044	BCMA-103	BC 263-A4	VL CDR2	aa	GASSLQD
1045	BCMA-103	BC 263-A4	VL CDR3	aa	QQSYKYPLT
1046	BCMA-103	BC 263-A4	VH	aa	EVQLVEESGGGLLQPGRSLKLSCAASGFTFSNYDMAWVRQAPT KGLEWVASISTRGDITSYRDSVKG RFTISRDNAKSTLYLQMDSLRSEDTATYYCARQDY YTDYMGFAYWGQGT'LVTVSS
1047	BCMA-103	BC 263-A4	VL	aa	ELVMTQSPASLSASLGTEVTTIECRASEDIYNGLAWYQQKPGKSPQLLIYGASSLQDGVPSRFSGSGS GTQYSLKISGMQPEDEANYFCQQQSYKYPLTFGSGTKLELK

1048	BCMA-103	BC 263-A4	scFv	aa	EVQLVEESGGGLLPGRSLKLSCAASGFTFSNYDMAWVRQAPTRGLEWVASISTRGDITSYRDSVKG RFTISRDNAKSTLYLQMDSLRSEDATATYYCARQDYITDYMGFAYWGQGTLLTVSSGGGGSGGGSGG GELVMTQSPASLSASLGSETVTIECRASEDIYNGLAWYQQKPKSPQLLIYGASSLQDGVPSRFSGSG SGTQYSLKISGMQPEDEANYFCQQQSYKYPLTFGSGTKLELKG
1049	BCMA-104	BC 271-C3	VH CDR1	aa	GFTFSNFDMA
1050	BCMA-104	BC 271-C3	VH CDR2	aa	SITTTGGGDTYYRDSVKG
1051	BCMA-104	BC 271-C3	VH CDR3	aa	HGYDGYHLEFDY
1052	BCMA-104	BC 271-C3	VL CDR1	aa	RASQGISNYL
1053	BCMA-104	BC 271-C3	VL CDR2	aa	YTSNLQS
1054	BCMA-104	BC 271-C3	VL CDR3	aa	QQYDISSYT
1055	BCMA-104	BC 271-C3	VH	aa	EVQLVEESGGGLVQPGRSLKLSCAASGFTFSNFDMAWVRQAPTRGLEWVASITTTGGGDTYYRDSVKG RFTISRDNAKSTLYLQMDSLRSEDATATYYCVRHGYDGYHLEFDYWGQGASVTVSS
1056	BCMA-104	BC 271-C3	VL	aa	ELVMTQTPSSMPASLGERVTISCRASQGISNYLNWYQQKPDGTIKPLIYYTSNLQSGVPSRFSGSGS GTDYSLTINSLEPEDFAVYYCQQYDISSYTFGAGTKLEIK
1057	BCMA-104	BC 271-C3	scFv	aa	EVQLVEESGGGLVQPGRSLKLSCAASGFTFSNFDMAWVRQAPTRGLEWVASITTTGGGDTYYRDSVKG RFTISRDNAKSTLYLQMDSLRSEDATATYYCVRHGYDGYHLEFDYWGQGASVTVSSGGGGSGGGSGG GGSELVMTQTPSSMPASLGERVTISCRASQGISNYLNWYQQKPDGTIKPLIYYTSNLQSGVPSRFSG SGSGTDYSLTINSLEPEDFAVYYCQQYDISSYTFGAGTKLEIK
1058	BCMA-105	BC 265-E5	VH CDR1	aa	GFTFSNFDMA
1059	BCMA-105	BC 265-E5	VH CDR2	aa	SITTTGGGDTYYRDSVKG
1060	BCMA-105	BC 265-E5	VH CDR3	aa	HGYDGYHLEFDY
1061	BCMA-105	BC 265-E5	VL CDR1	aa	RASQGISNHLN
1062	BCMA-105	BC 265-E5	VL CDR2	aa	YTSNLQS
1063	BCMA-105	BC 265-E5	VL CDR3	aa	QQYDSFPPLT

1064	BCMA-105	BC 265-E5	VH	aa	EVQLVEESGGGLVQPGRSLKLSCAASGFTFSNFDMAWVRQAPTRGLEWVASITTTGGDYYRDSVKG RFTISRDNAKSTLYLQMDSLRSEDATATYYCVRHGYDGYHLLFDYWGGQGLVTVSS
1065	BCMA-105	BC 265-E5	VL	aa	ELVMTQTSPSSMPASLGERVTISCRASQGISNHLNWYQQKPDGTIKPLIYYTSNLTQSGVPSRFSGSGS GTDYSLTISLLEPEDFAMYYCQQYDSFPPLTFGSGTKLEIK
1066	BCMA-105	BC 265-E5	scFv	aa	EVQLVEESGGGLVQPGRSLKLSCAASGFTFSNFDMAWVRQAPTRGLEWVASITTTGGDYYRDSVKG RFTISRDNAKSTLYLQMDSLRSEDATATYYCVRHGYDGYHLLFDYWGGQGLVTVSSGGSGGGSGG GGSELVMTQTSPSSMPASLGERVTISCRASQGISNHLNWYQQKPDGTIKPLIYYTSNLTQSGVPSRFSG SGSGTDYSLTISLLEPEDFAMYYCQQYDSFPPLTFGSGTKLEIK
1067	BCMA-106	BC271-B12	VH CDR1	aa	GFTFSNFDMA
1068	BCMA-106	BC271-B12	VH CDR2	aa	SITTTGGDYYRDSVKG
1069	BCMA-106	BC271-B12	VH CDR3	aa	HGYDGYHLLFDY
1070	BCMA-106	BC271-B12	VL CDR1	aa	RASQGISNNLN
1071	BCMA-106	BC271-B12	VL CDR2	aa	YTSNLQS
1072	BCMA-106	BC271-B12	VL CDR3	aa	QQFDTSPYT
1073	BCMA-106	BC271-B12	VH	aa	EVQLVEESGGGLVQPGRSLKLSCAASGFTFSNFDMAWVRQAPTRGLEWVASITTTGGDYYRDSVKG RFTISRDNAKSTLYLQMDSLRSEDATATYYCVRHGYDGYHLLFDYWGGQGMVTVSS
1074	BCMA-106	BC271-B12	VL	aa	ELVMTQTSPSSMPASLGERVTISCRASQGISNHLNWYQQKPDGTIKPLIYYTSNLTQSGVPSRFSGSGS GTDYSLTISLLEPEDFAMYYCQQFDTSPTYTFGAGTKLEIK
1075	BCMA-106	BC271-B12	scFv	aa	EVQLVEESGGGLVQPGRSLKLSCAASGFTFSNFDMAWVRQAPTRGLEWVASITTTGGDYYRDSVKG RFTISRDNAKSTLYLQMDSLRSEDATATYYCVRHGYDGYHLLFDYWGGQGMVTVSSGGSGGGSGG GGSELVMTQTSPSSMPASLGERVTISCRASQGISNHLNWYQQKPDGTIKPLIYYTSNLTQSGVPSRFSG SGSGTDYSLTISLLEPEDFAMYYCQQFDTSPTYTFGAGTKLEIK
1076	BCMA-107	BC 247-A4	VH CDR1	aa	GYSFPDYIN
1077	BCMA-107	BC 247-A4	VH CDR2	aa	WIYFASGNSEYNE
1078	BCMA-107	BC 247-A4	VH CDR3	aa	LYDYDWYFDV

1079	BCMA-107	BC 247-A4	VL CDR1	aa	RSSQSLVHSNGNTYLH
1080	BCMA-107	BC 247-A4	VL CDR2	aa	KVSNRFS
1081	BCMA-107	BC 247-A4	VL CDR3	aa	SQSTHVPYT
1082	BCMA-107	BC 247-A4	VH	aa	EVQLVEQSGPELVKPGASVKISCKVSGYSFPDYINWVKQRPQGQGLEWIGWIYFASGNSEYNERFTG KATLTVDTSSENTAYMQLSLTSSEDTAVYFCASLYDYDWYFDVWVGQGT'TVTVSS
1083	BCMA-107	BC 247-A4	VL	aa	ELVMTQTPLSLPVSLGDQASISCRSSQSLVHSNGNTYLHWY LQKPGQSPKLLIYKVSNRFSGVPDRF SGSGSGADF TTKISRVEAEDLGVYFCQSQSTHVPYTFGGG TKLEIK
1084	BCMA-107	BC 247-A4	scFv	aa	EVQLVEQSGPELVKPGASVKISCKVSGYSFPDYINWVKQRPQGQGLEWIGWIYFASGNSEYNERFTG KATLTVDTSSENTAYMQLSLTSSEDTAVYFCASLYDYDWYFDVWVGQGT'TVTVSSGGSGGGGG SELVMTQTPLSLPVSLGDQASISCRSSQSLVHSNGNTYLHWY LQKPGQSPKLLIYKVSNRFSGVPDR FSGSGSGADF TTKISRVEAEDLGVYFCQSQSTHVPYTFGGG TKLEIK
1085	BCMA-108	BC 246-B6	VH CDR1	aa	GYSFDPYIIN
1086	BCMA-108	BC 246-B6	VH CDR2	aa	WIYFASGNSEYNE
1087	BCMA-108	BC 246-B6	VH CDR3	aa	LYD YDWYFDV
1088	BCMA-108	BC 246-B6	VL CDR1	aa	RSSQSLVHSNGNTYLH
1089	BCMA-108	BC 246-B6	VL CDR2	aa	KVSNRFS
1090	BCMA-108	BC 246-B6	VL CDR3	aa	FQGSHPVPT
1091	BCMA-108	BC 246-B6	VH	aa	EVQLVEQSGPQLVKPGASVKISCKVSGYSFPDYINWVKQRPQGQGLEWIGWIYFASGNSEYNERFTG KATLTVDTSSENTAYMQLSLTSSEDTAVYFCASLYDYDWYFDVWVGQGT'TVTVSS
1092	BCMA-108	BC 246-B6	VL	aa	ELVMTQTPLSLPVSLGDQASISCRSSQSLVHSNGNTYLHWY LQKPGQSPKLLIYKVSNRFSGVPGRF SGSGSGTDF TTKINRVEAEDLGVYCFQGSHPVPTFGGG TKLEIK
1093	BCMA-108	BC 246-B6	scFv	aa	EVQLVEQSGPQLVKPGASVKISCKVSGYSFPDYINWVKQRPQGQGLEWIGWIYFASGNSEYNERFTG KATLTVDTSSENTAYMQLSLTSSEDTAVYFCASLYDYDWYFDVWVGQGT'TVTVSSGGSGGGGG SELVMTQTPLSLPVSLGDQASISCRSSQSLVHSNGNTYLHWY LQKPGQSPKLLIYKVSNRFSGVPGR FSGSGSGTDF TTKINRVEAEDLGVYCFQGSHPVPTFGGG TKLEIK

Claims

1. A binding molecule which is at least bispecific comprising a first and a second binding domain, wherein
 - (a) the first binding domain is capable of binding to epitope cluster 3 of BCMA (CQLRCSSNTPPLTCQRYC); and
 - (b) the second binding domain is capable of binding to the T cell CD3 receptor complex; andwherein epitope cluster 3 of BCMA corresponds to amino acid residues 24 to 41 of the sequence as depicted in SEQ ID NO: 1002.
2. The binding molecule according to claim 1, wherein the first binding domain is further capable of binding to epitope cluster 3 of macaque BCMA (CQLRCSSTPPLTCQRYC).
3. The binding molecule according to claim 1 or 2, wherein the second binding domain is capable of binding to CD3 epsilon.
4. The binding molecule according to any one of the preceding claims, wherein the second binding domain is capable of binding to human CD3 and to macaque CD3.
5. The binding molecule according to any one of the preceding claims, wherein the first and/or the second binding domain are derived from an antibody.
6. The binding molecule according to claim 5, which is selected from the group consisting of (scFv)₂, (single domain mAb)₂, scFv-single domain mAb, diabodies and oligomers thereof.
7. The binding molecule according to any one of the preceding claims, wherein the first binding domain comprises a VH region comprising CDR-H1, CDR-H2 and CDR-H3 and a VL region comprising CDR-L1, CDR-L2 and CDR-L3 selected from the group consisting of:
 - (1) CDR-H1 as depicted in SEQ ID NO: 1, CDR-H2 as depicted in SEQ ID NO: 2, CDR-H3 as depicted in SEQ ID NO: 3, CDR-L1 as depicted in SEQ ID NO: 4, CDR-L2 as depicted in SEQ ID NO: 5 and CDR-L3 as depicted in SEQ ID NO: 6;

- (2) CDR-H1 as depicted in SEQ ID NO: 11, CDR-H2 as depicted in SEQ ID NO: 12, CDR-H3 as depicted in SEQ ID NO: 13, CDR-L1 as depicted in SEQ ID NO: 14, CDR-L2 as depicted in SEQ ID NO: 15 and CDR-L3 as depicted in SEQ ID NO: 16;
- (3) CDR-H1 as depicted in SEQ ID NO: 21, CDR-H2 as depicted in SEQ ID NO: 22, CDR-H3 as depicted in SEQ ID NO: 23, CDR-L1 as depicted in SEQ ID NO: 24, CDR-L2 as depicted in SEQ ID NO: 25 and CDR-L3 as depicted in SEQ ID NO: 26;
- (4) CDR-H1 as depicted in SEQ ID NO: 31, CDR-H2 as depicted in SEQ ID NO: 32, CDR-H3 as depicted in SEQ ID NO: 33, CDR-L1 as depicted in SEQ ID NO: 34, CDR-L2 as depicted in SEQ ID NO: 35 and CDR-L3 as depicted in SEQ ID NO: 36;
- (5) CDR-H1 as depicted in SEQ ID NO: 41, CDR-H2 as depicted in SEQ ID NO: 42, CDR-H3 as depicted in SEQ ID NO: 43, CDR-L1 as depicted in SEQ ID NO: 44, CDR-L2 as depicted in SEQ ID NO: 45 and CDR-L3 as depicted in SEQ ID NO: 46;
- (6) CDR-H1 as depicted in SEQ ID NO: 51, CDR-H2 as depicted in SEQ ID NO: 52, CDR-H3 as depicted in SEQ ID NO: 53, CDR-L1 as depicted in SEQ ID NO: 54, CDR-L2 as depicted in SEQ ID NO: 55 and CDR-L3 as depicted in SEQ ID NO: 56;
- (7) CDR-H1 as depicted in SEQ ID NO: 61, CDR-H2 as depicted in SEQ ID NO: 62, CDR-H3 as depicted in SEQ ID NO: 63, CDR-L1 as depicted in SEQ ID NO: 64, CDR-L2 as depicted in SEQ ID NO: 65 and CDR-L3 as depicted in SEQ ID NO: 66;
- (8) CDR-H1 as depicted in SEQ ID NO: 71, CDR-H2 as depicted in SEQ ID NO: 72, CDR-H3 as depicted in SEQ ID NO: 73, CDR-L1 as depicted in SEQ ID NO: 74, CDR-L2 as depicted in SEQ ID NO: 75 and CDR-L3 as depicted in SEQ ID NO: 76;
- (9) CDR-H1 as depicted in SEQ ID NO: 161, CDR-H2 as depicted in SEQ ID NO: 162, CDR-H3 as depicted in SEQ ID NO: 163, CDR-L1 as depicted in SEQ ID NO: 164, CDR-L2 as depicted in SEQ ID NO: 165 and CDR-L3 as depicted in SEQ ID NO: 166;
- (10) CDR-H1 as depicted in SEQ ID NO: 171, CDR-H2 as depicted in SEQ ID NO: 172, CDR-H3 as depicted in SEQ ID NO: 173, CDR-L1 as depicted in SEQ ID NO: 174,

- CDR-L2 as depicted in SEQ ID NO: 175 and CDR-L3 as depicted in SEQ ID NO: 176;
- (11) CDR-H1 as depicted in SEQ ID NO: 181, CDR-H2 as depicted in SEQ ID NO: 182, CDR-H3 as depicted in SEQ ID NO: 183, CDR-L1 as depicted in SEQ ID NO: 184, CDR-L2 as depicted in SEQ ID NO: 185 and CDR-L3 as depicted in SEQ ID NO: 186;
- (12) CDR-H1 as depicted in SEQ ID NO: 191, CDR-H2 as depicted in SEQ ID NO: 192, CDR-H3 as depicted in SEQ ID NO: 193, CDR-L1 as depicted in SEQ ID NO: 194, CDR-L2 as depicted in SEQ ID NO: 195 and CDR-L3 as depicted in SEQ ID NO: 196;
- (13) CDR-H1 as depicted in SEQ ID NO: 201, CDR-H2 as depicted in SEQ ID NO: 202, CDR-H3 as depicted in SEQ ID NO: 203, CDR-L1 as depicted in SEQ ID NO: 204, CDR-L2 as depicted in SEQ ID NO: 205 and CDR-L3 as depicted in SEQ ID NO: 206;
- (14) CDR-H1 as depicted in SEQ ID NO: 211, CDR-H2 as depicted in SEQ ID NO: 212, CDR-H3 as depicted in SEQ ID NO: 213, CDR-L1 as depicted in SEQ ID NO: 214, CDR-L2 as depicted in SEQ ID NO: 215 and CDR-L3 as depicted in SEQ ID NO: 216;
- (15) CDR-H1 as depicted in SEQ ID NO: 221, CDR-H2 as depicted in SEQ ID NO: 222, CDR-H3 as depicted in SEQ ID NO: 223, CDR-L1 as depicted in SEQ ID NO: 224, CDR-L2 as depicted in SEQ ID NO: 225 and CDR-L3 as depicted in SEQ ID NO: 226;
- (16) CDR-H1 as depicted in SEQ ID NO: 311, CDR-H2 as depicted in SEQ ID NO: 312, CDR-H3 as depicted in SEQ ID NO: 313, CDR-L1 as depicted in SEQ ID NO: 314, CDR-L2 as depicted in SEQ ID NO: 315 and CDR-L3 as depicted in SEQ ID NO: 316;
- (17) CDR-H1 as depicted in SEQ ID NO: 321, CDR-H2 as depicted in SEQ ID NO: 322, CDR-H3 as depicted in SEQ ID NO: 323, CDR-L1 as depicted in SEQ ID NO: 324, CDR-L2 as depicted in SEQ ID NO: 325 and CDR-L3 as depicted in SEQ ID NO: 326;
- (18) CDR-H1 as depicted in SEQ ID NO: 331, CDR-H2 as depicted in SEQ ID NO: 332, CDR-H3 as depicted in SEQ ID NO: 333, CDR-L1 as depicted in SEQ ID NO: 334, CDR-L2 as depicted in SEQ ID NO: 335 and CDR-L3 as depicted in SEQ ID NO: 336;

- (19) CDR-H1 as depicted in SEQ ID NO: 341, CDR-H2 as depicted in SEQ ID NO: 342, CDR-H3 as depicted in SEQ ID NO: 343, CDR-L1 as depicted in SEQ ID NO: 344, CDR-L2 as depicted in SEQ ID NO: 345 and CDR-L3 as depicted in SEQ ID NO: 346;
- (20) CDR-H1 as depicted in SEQ ID NO: 351, CDR-H2 as depicted in SEQ ID NO: 352, CDR-H3 as depicted in SEQ ID NO: 353, CDR-L1 as depicted in SEQ ID NO: 354, CDR-L2 as depicted in SEQ ID NO: 355 and CDR-L3 as depicted in SEQ ID NO: 356;
- (21) CDR-H1 as depicted in SEQ ID NO: 361, CDR-H2 as depicted in SEQ ID NO: 362, CDR-H3 as depicted in SEQ ID NO: 363, CDR-L1 as depicted in SEQ ID NO: 364, CDR-L2 as depicted in SEQ ID NO: 365 and CDR-L3 as depicted in SEQ ID NO: 366;
- (22) CDR-H1 as depicted in SEQ ID NO: 371, CDR-H2 as depicted in SEQ ID NO: 372, CDR-H3 as depicted in SEQ ID NO: 373, CDR-L1 as depicted in SEQ ID NO: 374, CDR-L2 as depicted in SEQ ID NO: 375 and CDR-L3 as depicted in SEQ ID NO: 376;
- (23) CDR-H1 as depicted in SEQ ID NO: 381, CDR-H2 as depicted in SEQ ID NO: 382, CDR-H3 as depicted in SEQ ID NO: 383, CDR-L1 as depicted in SEQ ID NO: 384, CDR-L2 as depicted in SEQ ID NO: 385 and CDR-L3 as depicted in SEQ ID NO: 386;
- (24) CDR-H1 as depicted in SEQ ID NO: 581, CDR-H2 as depicted in SEQ ID NO: 582, CDR-H3 as depicted in SEQ ID NO: 583, CDR-L1 as depicted in SEQ ID NO: 584, CDR-L2 as depicted in SEQ ID NO: 585 and CDR-L3 as depicted in SEQ ID NO: 586;
- (25) CDR-H1 as depicted in SEQ ID NO: 591, CDR-H2 as depicted in SEQ ID NO: 592, CDR-H3 as depicted in SEQ ID NO: 593, CDR-L1 as depicted in SEQ ID NO: 594, CDR-L2 as depicted in SEQ ID NO: 595 and CDR-L3 as depicted in SEQ ID NO: 596;
- (26) CDR-H1 as depicted in SEQ ID NO: 601, CDR-H2 as depicted in SEQ ID NO: 602, CDR-H3 as depicted in SEQ ID NO: 603, CDR-L1 as depicted in SEQ ID NO: 604, CDR-L2 as depicted in SEQ ID NO: 605 and CDR-L3 as depicted in SEQ ID NO: 606;
- (27) CDR-H1 as depicted in SEQ ID NO: 611, CDR-H2 as depicted in SEQ ID NO: 612, CDR-H3 as depicted in SEQ ID NO: 613, CDR-L1 as depicted in SEQ ID NO: 614,

- CDR-L2 as depicted in SEQ ID NO: 615 and CDR-L3 as depicted in SEQ ID NO: 616;
- (28) CDR-H1 as depicted in SEQ ID NO: 621, CDR-H2 as depicted in SEQ ID NO: 622, CDR-H3 as depicted in SEQ ID NO: 623, CDR-L1 as depicted in SEQ ID NO: 624, CDR-L2 as depicted in SEQ ID NO: 625 and CDR-L3 as depicted in SEQ ID NO: 626;
- (29) CDR-H1 as depicted in SEQ ID NO: 631, CDR-H2 as depicted in SEQ ID NO: 632, CDR-H3 as depicted in SEQ ID NO: 633, CDR-L1 as depicted in SEQ ID NO: 634, CDR-L2 as depicted in SEQ ID NO: 635 and CDR-L3 as depicted in SEQ ID NO: 636;
- (30) CDR-H1 as depicted in SEQ ID NO: 641, CDR-H2 as depicted in SEQ ID NO: 642, CDR-H3 as depicted in SEQ ID NO: 643, CDR-L1 as depicted in SEQ ID NO: 644, CDR-L2 as depicted in SEQ ID NO: 645 and CDR-L3 as depicted in SEQ ID NO: 646;
- (31) CDR-H1 as depicted in SEQ ID NO: 651, CDR-H2 as depicted in SEQ ID NO: 652, CDR-H3 as depicted in SEQ ID NO: 653, CDR-L1 as depicted in SEQ ID NO: 654, CDR-L2 as depicted in SEQ ID NO: 655 and CDR-L3 as depicted in SEQ ID NO: 656;
- (32) CDR-H1 as depicted in SEQ ID NO: 661, CDR-H2 as depicted in SEQ ID NO: 662, CDR-H3 as depicted in SEQ ID NO: 663, CDR-L1 as depicted in SEQ ID NO: 664, CDR-L2 as depicted in SEQ ID NO: 665 and CDR-L3 as depicted in SEQ ID NO: 666;
- (33) CDR-H1 as depicted in SEQ ID NO: 671, CDR-H2 as depicted in SEQ ID NO: 672, CDR-H3 as depicted in SEQ ID NO: 673, CDR-L1 as depicted in SEQ ID NO: 674, CDR-L2 as depicted in SEQ ID NO: 675 and CDR-L3 as depicted in SEQ ID NO: 676;
- (34) CDR-H1 as depicted in SEQ ID NO: 681, CDR-H2 as depicted in SEQ ID NO: 682, CDR-H3 as depicted in SEQ ID NO: 683, CDR-L1 as depicted in SEQ ID NO: 684, CDR-L2 as depicted in SEQ ID NO: 685 and CDR-L3 as depicted in SEQ ID NO: 686;
- (35) CDR-H1 as depicted in SEQ ID NO: 691, CDR-H2 as depicted in SEQ ID NO: 692, CDR-H3 as depicted in SEQ ID NO: 693, CDR-L1 as depicted in SEQ ID NO: 694, CDR-L2 as depicted in SEQ ID NO: 695 and CDR-L3 as depicted in SEQ ID NO: 696;

- (36) CDR-H1 as depicted in SEQ ID NO: 701, CDR-H2 as depicted in SEQ ID NO: 702, CDR-H3 as depicted in SEQ ID NO: 703, CDR-L1 as depicted in SEQ ID NO: 704, CDR-L2 as depicted in SEQ ID NO: 705 and CDR-L3 as depicted in SEQ ID NO: 706;
- (37) CDR-H1 as depicted in SEQ ID NO: 711, CDR-H2 as depicted in SEQ ID NO: 712, CDR-H3 as depicted in SEQ ID NO: 713, CDR-L1 as depicted in SEQ ID NO: 714, CDR-L2 as depicted in SEQ ID NO: 715 and CDR-L3 as depicted in SEQ ID NO: 716;
- (38) CDR-H1 as depicted in SEQ ID NO: 721, CDR-H2 as depicted in SEQ ID NO: 722, CDR-H3 as depicted in SEQ ID NO: 723, CDR-L1 as depicted in SEQ ID NO: 724, CDR-L2 as depicted in SEQ ID NO: 725 and CDR-L3 as depicted in SEQ ID NO: 726;
- (39) CDR-H1 as depicted in SEQ ID NO: 731, CDR-H2 as depicted in SEQ ID NO: 732, CDR-H3 as depicted in SEQ ID NO: 733, CDR-L1 as depicted in SEQ ID NO: 734, CDR-L2 as depicted in SEQ ID NO: 735 and CDR-L3 as depicted in SEQ ID NO: 736;
- (40) CDR-H1 as depicted in SEQ ID NO: 741, CDR-H2 as depicted in SEQ ID NO: 742, CDR-H3 as depicted in SEQ ID NO: 743, CDR-L1 as depicted in SEQ ID NO: 744, CDR-L2 as depicted in SEQ ID NO: 745 and CDR-L3 as depicted in SEQ ID NO: 746;
- (41) CDR-H1 as depicted in SEQ ID NO: 751, CDR-H2 as depicted in SEQ ID NO: 752, CDR-H3 as depicted in SEQ ID NO: 753, CDR-L1 as depicted in SEQ ID NO: 754, CDR-L2 as depicted in SEQ ID NO: 755 and CDR-L3 as depicted in SEQ ID NO: 756;
- (42) CDR-H1 as depicted in SEQ ID NO: 761, CDR-H2 as depicted in SEQ ID NO: 762, CDR-H3 as depicted in SEQ ID NO: 763, CDR-L1 as depicted in SEQ ID NO: 764, CDR-L2 as depicted in SEQ ID NO: 765 and CDR-L3 as depicted in SEQ ID NO: 766;
- (43) CDR-H1 as depicted in SEQ ID NO: 771, CDR-H2 as depicted in SEQ ID NO: 772, CDR-H3 as depicted in SEQ ID NO: 773, CDR-L1 as depicted in SEQ ID NO: 774, CDR-L2 as depicted in SEQ ID NO: 775 and CDR-L3 as depicted in SEQ ID NO: 776;
- (44) CDR-H1 as depicted in SEQ ID NO: 781, CDR-H2 as depicted in SEQ ID NO: 782, CDR-H3 as depicted in SEQ ID NO: 783, CDR-L1 as depicted in SEQ ID NO: 784,

- CDR-L2 as depicted in SEQ ID NO: 785 and CDR-L3 as depicted in SEQ ID NO: 786;
- (45) CDR-H1 as depicted in SEQ ID NO: 791, CDR-H2 as depicted in SEQ ID NO: 792, CDR-H3 as depicted in SEQ ID NO: 793, CDR-L1 as depicted in SEQ ID NO: 794, CDR-L2 as depicted in SEQ ID NO: 795 and CDR-L3 as depicted in SEQ ID NO: 796;
- (46) CDR-H1 as depicted in SEQ ID NO: 801, CDR-H2 as depicted in SEQ ID NO: 802, CDR-H3 as depicted in SEQ ID NO: 803, CDR-L1 as depicted in SEQ ID NO: 804, CDR-L2 as depicted in SEQ ID NO: 805 and CDR-L3 as depicted in SEQ ID NO: 806;
- (47) CDR-H1 as depicted in SEQ ID NO: 811, CDR-H2 as depicted in SEQ ID NO: 812, CDR-H3 as depicted in SEQ ID NO: 813, CDR-L1 as depicted in SEQ ID NO: 814, CDR-L2 as depicted in SEQ ID NO: 815 and CDR-L3 as depicted in SEQ ID NO: 816;
- (48) CDR-H1 as depicted in SEQ ID NO: 821, CDR-H2 as depicted in SEQ ID NO: 822, CDR-H3 as depicted in SEQ ID NO: 823, CDR-L1 as depicted in SEQ ID NO: 824, CDR-L2 as depicted in SEQ ID NO: 825 and CDR-L3 as depicted in SEQ ID NO: 826;
- (49) CDR-H1 as depicted in SEQ ID NO: 831, CDR-H2 as depicted in SEQ ID NO: 832, CDR-H3 as depicted in SEQ ID NO: 833, CDR-L1 as depicted in SEQ ID NO: 834, CDR-L2 as depicted in SEQ ID NO: 835 and CDR-L3 as depicted in SEQ ID NO: 836;
- (50) CDR-H1 as depicted in SEQ ID NO: 961, CDR-H2 as depicted in SEQ ID NO: 962, CDR-H3 as depicted in SEQ ID NO: 963, CDR-L1 as depicted in SEQ ID NO: 964, CDR-L2 as depicted in SEQ ID NO: 965 and CDR-L3 as depicted in SEQ ID NO: 966;
- (51) CDR-H1 as depicted in SEQ ID NO: 971, CDR-H2 as depicted in SEQ ID NO: 972, CDR-H3 as depicted in SEQ ID NO: 973, CDR-L1 as depicted in SEQ ID NO: 974, CDR-L2 as depicted in SEQ ID NO: 975 and CDR-L3 as depicted in SEQ ID NO: 976;
- (52) CDR-H1 as depicted in SEQ ID NO: 981, CDR-H2 as depicted in SEQ ID NO: 982, CDR-H3 as depicted in SEQ ID NO: 983, CDR-L1 as depicted in SEQ ID NO: 984, CDR-L2 as depicted in SEQ ID NO: 985 and CDR-L3 as depicted in SEQ ID NO: 986; and

- (53) CDR-H1 as depicted in SEQ ID NO: 991, CDR-H2 as depicted in SEQ ID NO: 992, CDR-H3 as depicted in SEQ ID NO: 993, CDR-L1 as depicted in SEQ ID NO: 994, CDR-L2 as depicted in SEQ ID NO: 995 and CDR-L3 as depicted in SEQ ID NO: 996.
8. The binding molecule according to any one of the preceding claims, wherein the first binding domain comprises a VH region selected from the group consisting of VH regions as depicted in SEQ ID NO: 7, SEQ ID NO: 17, SEQ ID NO: 27, SEQ ID NO: 37, SEQ ID NO: 47, SEQ ID NO: 57, SEQ ID NO: 67, SEQ ID NO: 77, SEQ ID NO: 167, SEQ ID NO: 177, SEQ ID NO: 187, SEQ ID NO: 197, SEQ ID NO: 207, SEQ ID NO: 217, SEQ ID NO: 227, SEQ ID NO: 317, SEQ ID NO: 327, SEQ ID NO: 337, SEQ ID NO: 347, SEQ ID NO: 357, SEQ ID NO: 367, SEQ ID NO: 377, SEQ ID NO: 387, SEQ ID NO: 587, SEQ ID NO: 597, SEQ ID NO: 607, SEQ ID NO: 617, SEQ ID NO: 627, SEQ ID NO: 637, SEQ ID NO: 647, SEQ ID NO: 657, SEQ ID NO: 667, SEQ ID NO: 677, SEQ ID NO: 687, SEQ ID NO: 697, SEQ ID NO: 707, SEQ ID NO: 717, SEQ ID NO: 727, SEQ ID NO: 737, SEQ ID NO: 747, SEQ ID NO: 757, SEQ ID NO: 767, SEQ ID NO: 777, SEQ ID NO: 787, SEQ ID NO: 797, SEQ ID NO: 807, SEQ ID NO: 817, SEQ ID NO: 827, SEQ ID NO: 837, SEQ ID NO: 967, SEQ ID NO: 977, SEQ ID NO: 987, and SEQ ID NO: 997.
9. The binding molecule according to any one of the preceding claims, wherein the first binding domain comprises a VL region selected from the group consisting of VL regions as depicted in SEQ ID NO: 8, SEQ ID NO: 18, SEQ ID NO: 28, SEQ ID NO: 38, SEQ ID NO: 48, SEQ ID NO: 58, SEQ ID NO: 68, SEQ ID NO: 78, SEQ ID NO: 168, SEQ ID NO: 178, SEQ ID NO: 188, SEQ ID NO: 198, SEQ ID NO: 208, SEQ ID NO: 218, SEQ ID NO: 228, SEQ ID NO: 318, SEQ ID NO: 328, SEQ ID NO: 338, SEQ ID NO: 348, SEQ ID NO: 358, SEQ ID NO: 368, SEQ ID NO: 378, SEQ ID NO: 388, SEQ ID NO: 588, SEQ ID NO: 598, SEQ ID NO: 608, SEQ ID NO: 618, SEQ ID NO: 628, SEQ ID NO: 638, SEQ ID NO: 648, SEQ ID NO: 658, SEQ ID NO: 668, SEQ ID NO: 678, SEQ ID NO: 688, SEQ ID NO: 698, SEQ ID NO: 708, SEQ ID NO: 718, SEQ ID NO: 728, SEQ ID NO: 738, SEQ ID NO: 748, SEQ ID NO: 758, SEQ ID NO: 768, SEQ ID NO: 778, SEQ ID NO: 788, SEQ ID NO: 798, SEQ ID NO: 808, SEQ ID NO: 818, SEQ ID NO: 828, SEQ ID NO: 838, SEQ ID NO: 968, SEQ ID NO: 978, SEQ ID NO: 988, and SEQ ID NO: 998.

10. The binding molecule according to any one of the preceding claims, wherein the first binding domain comprises a VH region and a VL region selected from the group consisting of:
- (1) a VH region as depicted in SEQ ID NO: 7, and a VL region as depicted in SEQ ID NO: 8;
 - (2) a VH region as depicted in SEQ ID NO: 17, and a VL region as depicted in SEQ ID NO: 18;
 - (3) a VH region as depicted in SEQ ID NO: 27, and a VL region as depicted in SEQ ID NO: 28;
 - (4) a VH region as depicted in SEQ ID NO: 37, and a VL region as depicted in SEQ ID NO: 38;
 - (5) a VH region as depicted in SEQ ID NO: 47, and a VL region as depicted in SEQ ID NO: 48;
 - (6) a VH region as depicted in SEQ ID NO: 57, and a VL region as depicted in SEQ ID NO: 58;
 - (7) a VH region as depicted in SEQ ID NO: 67, and a VL region as depicted in SEQ ID NO: 68;
 - (8) a VH region as depicted in SEQ ID NO: 77, and a VL region as depicted in SEQ ID NO: 78;
 - (9) a VH region as depicted in SEQ ID NO: 167, and a VL region as depicted in SEQ ID NO: 168;
 - (10) a VH region as depicted in SEQ ID NO: 177, and a VL region as depicted in SEQ ID NO: 178;
 - (11) a VH region as depicted in SEQ ID NO: 187, and a VL region as depicted in SEQ ID NO: 188;
 - (12) a VH region as depicted in SEQ ID NO: 197, and a VL region as depicted in SEQ ID NO: 198;
 - (13) a VH region as depicted in SEQ ID NO: 207, and a VL region as depicted in SEQ ID NO: 208;
 - (14) a VH region as depicted in SEQ ID NO: 217, and a VL region as depicted in SEQ ID NO: 218;
 - (15) a VH region as depicted in SEQ ID NO: 227, and a VL region as depicted in SEQ ID NO: 228;
 - (16) a VH region as depicted in SEQ ID NO: 317, and a VL region as depicted in SEQ ID NO: 318;

- (17) a VH region as depicted in SEQ ID NO: 327, and a VL region as depicted in SEQ ID NO: 328;
- (18) a VH region as depicted in SEQ ID NO: 337, and a VL region as depicted in SEQ ID NO: 338;
- (19) a VH region as depicted in SEQ ID NO: 347, and a VL region as depicted in SEQ ID NO: 348;
- (20) a VH region as depicted in SEQ ID NO: 357, and a VL region as depicted in SEQ ID NO: 358;
- (21) a VH region as depicted in SEQ ID NO: 367, and a VL region as depicted in SEQ ID NO: 368;
- (22) a VH region as depicted in SEQ ID NO: 377, and a VL region as depicted in SEQ ID NO: 378;
- (23) a VH region as depicted in SEQ ID NO: 387, and a VL region as depicted in SEQ ID NO: 388;
- (24) a VH region as depicted in SEQ ID NO: 587, and a VL region as depicted in SEQ ID NO: 588;
- (25) a VH region as depicted in SEQ ID NO: 597, and a VL region as depicted in SEQ ID NO: 598;
- (26) a VH region as depicted in SEQ ID NO: 607, and a VL region as depicted in SEQ ID NO: 608;
- (27) a VH region as depicted in SEQ ID NO: 617, and a VL region as depicted in SEQ ID NO: 618;
- (28) a VH region as depicted in SEQ ID NO: 627, and a VL region as depicted in SEQ ID NO: 628;
- (29) a VH region as depicted in SEQ ID NO: 637, and a VL region as depicted in SEQ ID NO: 638;
- (30) a VH region as depicted in SEQ ID NO: 647, and a VL region as depicted in SEQ ID NO: 648;
- (31) a VH region as depicted in SEQ ID NO: 657, and a VL region as depicted in SEQ ID NO: 658;
- (32) a VH region as depicted in SEQ ID NO: 667, and a VL region as depicted in SEQ ID NO: 668;
- (33) a VH region as depicted in SEQ ID NO: 677, and a VL region as depicted in SEQ ID NO: 678;


- (34) a VH region as depicted in SEQ ID NO: 687, and a VL region as depicted in SEQ ID NO: 688;
- (35) a VH region as depicted in SEQ ID NO: 697, and a VL region as depicted in SEQ ID NO: 698;
- (36) a VH region as depicted in SEQ ID NO: 707, and a VL region as depicted in SEQ ID NO: 708;
- (37) a VH region as depicted in SEQ ID NO: 717, and a VL region as depicted in SEQ ID NO: 718;
- (38) a VH region as depicted in SEQ ID NO: 727, and a VL region as depicted in SEQ ID NO: 728;
- (39) a VH region as depicted in SEQ ID NO: 737, and a VL region as depicted in SEQ ID NO: 738;
- (40) a VH region as depicted in SEQ ID NO: 747, and a VL region as depicted in SEQ ID NO: 748;
- (41) a VH region as depicted in SEQ ID NO: 757, and a VL region as depicted in SEQ ID NO: 758;
- (42) a VH region as depicted in SEQ ID NO: 767, and a VL region as depicted in SEQ ID NO: 768;
- (43) a VH region as depicted in SEQ ID NO: 777, and a VL region as depicted in SEQ ID NO: 778;
- (44) a VH region as depicted in SEQ ID NO: 787, and a VL region as depicted in SEQ ID NO: 788;
- (45) a VH region as depicted in SEQ ID NO: 797, and a VL region as depicted in SEQ ID NO: 798;
- (46) a VH region as depicted in SEQ ID NO: 807, and a VL region as depicted in SEQ ID NO: 808;
- (47) a VH region as depicted in SEQ ID NO: 817, and a VL region as depicted in SEQ ID NO: 818;
- (48) a VH region as depicted in SEQ ID NO: 827, and a VL region as depicted in SEQ ID NO: 828;
- (49) a VH region as depicted in SEQ ID NO: 837, and a VL region as depicted in SEQ ID NO: 838;
- (50) a VH region as depicted in SEQ ID NO: 967, and a VL region as depicted in SEQ ID NO: 968;

- (51) a VH region as depicted in SEQ ID NO: 977, and a VL region as depicted in SEQ ID NO: 978;
 - (52) a VH region as depicted in SEQ ID NO: 987, and a VL region as depicted in SEQ ID NO: 988; and
 - (53) a VH region as depicted in SEQ ID NO: 997, and a VL region as depicted in SEQ ID NO: 998.
11. The binding molecule according to claim 10, wherein the first binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 9, SEQ ID NO: 19, SEQ ID NO: 29, SEQ ID NO: 39, SEQ ID NO: 49, SEQ ID NO: 59, SEQ ID NO: 69, SEQ ID NO: 79, SEQ ID NO: 169, SEQ ID NO: 179, SEQ ID NO: 189, SEQ ID NO: 199, SEQ ID NO: 209, SEQ ID NO: 219, SEQ ID NO: 229, SEQ ID NO: 319, SEQ ID NO: 329, SEQ ID NO: 339, SEQ ID NO: 349, SEQ ID NO: 359, SEQ ID NO: 369, SEQ ID NO: 379, SEQ ID NO: 389, SEQ ID NO: 589, SEQ ID NO: 599, SEQ ID NO: 609, SEQ ID NO: 619, SEQ ID NO: 629, SEQ ID NO: 639, SEQ ID NO: 649, SEQ ID NO: 659, SEQ ID NO: 669, SEQ ID NO: 679, SEQ ID NO: 689, SEQ ID NO: 699, SEQ ID NO: 709, SEQ ID NO: 719, SEQ ID NO: 729, SEQ ID NO: 739, SEQ ID NO: 749, SEQ ID NO: 759, SEQ ID NO: 769, SEQ ID NO: 779, SEQ ID NO: 789, SEQ ID NO: 799, SEQ ID NO: 809, SEQ ID NO: 819, SEQ ID NO: 829, SEQ ID NO: 839, SEQ ID NO: 969, SEQ ID NO: 979, SEQ ID NO: 989, and SEQ ID NO: 999.
 12. The binding molecule according to any one of claims 1-6 having the amino acid sequence shown in SEQ ID NO:340 or SEQ ID NO: 980.
 13. The binding molecule according to any one of the preceding claims, characterized by an EC50 (pg/ml) of 350 or less, preferably 320 or less.
 14. The binding molecule according to any one of the preceding claims, characterized by an EC50 (pg/ml) which equates to the EC50 (pg/ml) of any one of BC E5 33-B11-B8, BC 5G9 92-E10, BC 5G9 91-D2-B10, BC B12 33-A4-B2, BC 3A4 37-A11-G1, BC A7-27 C4-G7, BC C3 33-D7-B1, BC C3 33-F8-E6 B1.
 15. A nucleic acid sequence encoding a binding molecule as defined in any one of claims 1 to 14

16. A vector comprising a nucleic acid sequence as defined in claim 15.
17. A host cell transformed or transfected with the nucleic acid sequence as defined in claim 15 or with the vector as defined in claim 16.
18. A process for the production of a binding molecule according to any one of claims 1 to 14, said process comprising culturing a host cell as defined in claim 17 under conditions allowing the expression of the binding molecule as defined in any one of claims 1 to 14 and recovering the produced binding molecule from the culture.
19. A pharmaceutical composition comprising a binding molecule according to any one of claims 1 to 14, or produced according to the process of claim 18.
20. The binding molecule according to any one of claims 1 to 14, or produced according to the process of claim 18 for use in the prevention, treatment or amelioration of a disease selected from the group consisting of plasma cell disorders, other B cell disorders that correlate with BCMA expression and autoimmune diseases.
21. A method for the treatment or amelioration of a disease selected from the group consisting of plasma cell disorders, other B cell disorders that correlate with BCMA expression and autoimmune diseases, comprising the step of administering to a subject in need thereof the binding molecule according to any one of claims 1 to 14, or produced according to the process of claim 18.
22. The method according to claim 21, wherein the plasma cell disorder is selected from the group consisting of multiple myeloma, plasmacytoma, plasma cell leukemia, macroglobulinemia, amyloidosis, Waldenstrom's macroglobulinemia, solitary bone plasmacytoma, extramedullary plasmacytoma, osteosclerotic myeloma, heavy chain diseases, monoclonal gammopathy of undetermined significance, and smoldering multiple myeloma.
23. The method according to claim 21, wherein the autoimmune disease is systemic lupus erythematoses.

24. A kit comprising a binding molecule as defined in any one of claims 1 to 14, a nucleic acid molecule as defined in claim 15, a vector as defined in claim 16, and/or a host cell as defined in claim 17.
25. Use of epitope cluster 3 of BCMA for the generation of a binding molecule, preferably an antibody, which is capable of binding to BCMA, wherein epitope cluster 3 of BCMA corresponds to amino acid residues 24 to 41 of the sequence as depicted in SEQ ID NO: 1002.
26. A method for the generation of an antibody, preferably a bispecific binding molecule, which is capable of binding to BCMA, comprising
 - (a) immunizing an animal with a polypeptide comprising epitope cluster 3 of BCMA, wherein epitope cluster 3 of BCMA corresponds to amino acid residues 24 to 41 of the sequence as depicted in SEQ ID NO: 1002,
 - (b) obtaining said antibody, and
 - (c) optionally converting said antibody into a bispecific binding molecule which is capable of binding to human BCMA and preferably to the T cell CD3 receptor complex.

Figure 1



- E1: N-terminal domain
- E2: First disulfide bound defined domain
- E3: Second (double) disulfide bound defined domain
- E4: C-terminal extracellular domain
- E5: Inter-E2/E3 point mutation
- E6+E7: E3 point mutations

Figure 2a




Figure 2b




Figure 3




Figure 3 (continued)






Figure 4

Human BCMA

Macaque BCMA



BCMA 101 x CD3



KD = 0.5 nM

k off = $1.1 \times 10^{-4} \text{ s}^{-1}$

KD = 0.8 nM

k off = $1.5 \times 10^{-4} \text{ s}^{-1}$

5/18



BCMA 102 x CD3

KD = 0.6 nM

k off = $1.1 \times 10^{-4} \text{ s}^{-1}$

KD = 0.3 nM

k off = $0.54 \times 10^{-4} \text{ s}^{-1}$

Figure 5




Figure 6






Figure 7



BCMA-98 x CD3


Figure 8

Human BCMA / CHO Macaque BCMA / CHO



BCMA-20 x CD3

Figure 9



11/18

Figure 10






Figure 11


13/18

Figure 12




14/18

Figure 13




15/18

Figure 14



16/18

Figure 15



17/18

Figure 16




Figure 17




Figure 16

