

Related U.S. Application Data

- (60) Provisional application No. 60/946,287, filed on Jun. 26, 2007, provisional application No. 61/049,826, filed on May 2, 2008.

(51) Int. Cl.

C07K 16/00 (2006.01)

C07K 16/32 (2006.01)

(52) U.S. Cl.

CPC **C07K 2317/52** (2013.01); **C07K 2317/569** (2013.01); **C07K 2318/10** (2013.01); **C40B 30/04** (2013.01); **G01N 2333/70535** (2013.01)

(56) References Cited

U.S. PATENT DOCUMENTS

5,475,100	A	12/1995	Hashino et al.	536/23.53
5,536,814	A	7/1996	Ruoslakhti et al.	530/329
5,723,323	A	3/1998	Kaurrman et al.	435/172.3
5,759,817	A	6/1998	Barbas	435/69.7
5,763,192	A	6/1998	Kauffman et al.	435/7.1
5,814,476	A	9/1998	Kauffman et al.	435/69.1
5,817,483	A	10/1998	Kauffman et al.	435/69.1
5,824,514	A	10/1998	Kauffman et al.	435/91.1
5,844,094	A	12/1998	Hudson et al.	530/387.3
5,892,019	A	4/1999	Schlom et al.	536/23.53
6,057,098	A	5/2000	Buechler et al.	435/6
6,294,654	B1	9/2001	Bogen et al.	530/387.3
6,352,842	B1	3/2002	Short et al.	435/69.1
6,358,709	B1	3/2002	Short et al.	435/69.1
6,361,974	B1	3/2002	Short et al.	435/69.1
6,365,377	B1	4/2002	Patten et al.	435/91.1
6,376,246	B1	4/2002	Cramer et al.	435/440
6,562,617	B1	5/2003	Anderson et al.	435/325
6,602,684	B1	8/2003	Umaña et al.	435/69.1
7,442,778	B2	10/2008	Gegg et al.	530/391.7
7,632,497	B2	12/2009	Stavenhagen	424/133.1
7,645,861	B2	1/2010	Gegg et al.	530/391.7
7,655,764	B2	2/2010	Gegg et al.	530/391.7
7,655,765	B2	2/2010	Gegg et al.	530/391.7
7,662,931	B2	2/2010	Gegg et al.	530/391.7
7,750,127	B2	7/2010	Gegg et al.	530/391.7
7,750,128	B2	7/2010	Gegg et al.	530/391.7
7,858,090	B2	12/2010	Koide	424/145.1
8,008,453	B2	8/2011	Gegg et al.	530/391.7
8,580,927	B2	11/2013	Dimitrov	530/387.3
8,859,738	B2	10/2014	Himmeler et al.	
8,921,279	B2	12/2014	Himmeler et al.	506/9
9,045,528	B2	6/2015	Rüker et al.	
2002/0103345	A1	8/2002	Zhu	530/388.15
2002/0106370	A1	8/2002	Cardy et al.	424/133.1
2003/0027213	A1	2/2003	Zhu et al.	435/7.1
2003/0129188	A1	7/2003	Barbas et al.	424/144.1
2003/0148372	A1	8/2003	Tomlinson et al.	435/7.1
2003/0157091	A1	8/2003	Hoogenboom	424/130.1
2004/0018508	A1	1/2004	Friedman	435/6
2004/0043424	A1	3/2004	Baughn et al.	435/7.1
2004/0063924	A1	4/2004	Tang et al.	536/23.5
2004/0071690	A1	4/2004	Hudon et al.	424/130.1
2004/0082508	A1	4/2004	Yue et al.	514/12
2004/0097711	A1	5/2004	Yue et al.	530/387.1
2004/0101905	A1	5/2004	Brekke et al.	435/7.1
2004/0132101	A1	7/2004	Lazar et al.	435/7.1
2004/0146976	A1	7/2004	Wittrup et al.	435/69.1
2005/0009025	A1	1/2005	Jakobsen et al.	435/6
2005/0054832	A1	3/2005	Lazar et al.	530/387.3
2005/0069549	A1	3/2005	Herman	424/178.1
2005/0158829	A1	7/2005	Fandl et al.	435/69.1
2005/0244403	A1	11/2005	Lazar et al.	424/130.1
2005/0255548	A1	11/2005	Lipovsck et al.	435/69.1
2005/0266000	A1	12/2005	Bond et al.	424/143.1
2006/0140934	A1	6/2006	Gegg et al.	424/133.1
2008/0227958	A1	9/2008	Thompson et al.	530/387.3
2009/0298195	A1	12/2009	Rüker et al.	436/501
2010/0048877	A1	2/2010	Rüker et al.	530/387.3

2011/0251375	A1	10/2011	Rüker et al.	530/387.3
2012/0010388	A1	1/2012	Himmeler	530/387.3
2012/0028839	A1	2/2012	Rüker et al.	506/14
2012/0094874	A1	4/2012	Rüker et al.	506/18
2012/0276104	A1	11/2012	Woissetschlager	424/136.1
2015/0153359	A1	6/2015	Himmeler et al.	

FOREIGN PATENT DOCUMENTS

CN	1606566	A	4/2005	C07K 5/08
EP	0 640 130	B1	7/1993	C12N 15/12
EP	1 752 471	A1	2/2007	C07K 16/00
EP	1 772 465	A1	4/2007	C07K 16/00
EP	1 797 127	A2	6/2007	A61K 47/48
EP	2 028 193	A1	2/2009	C07K 16/00
EP	1 699 826	B1	3/2009	C07K 16/00
EP	2 158 220	A1	3/2010	C07K 16/00
EP	2 407 487	A1	1/2012	C07K 16/32
EP	2 451 838		5/2012	C12N 15/09
JP	2002-58479	A	2/2002	C12N 15/09
JP	2003-518377		6/2003	C12N 15/09
JP	2003-518377	A	6/2003	C12N 15/09
WO	WO 90/07861		7/1990	C12P 21/00
WO	WO 92/09690		6/1992	C12N 15/00
WO	WO 93/08278		4/1993	C12N 15/10
WO	WO 93/23537		11/1993	C12N 15/12
WO	WO 96/22377		7/1996	C12N 15/62
WO	WO 97/20858		6/1997	C07K 16/32
WO	WO 97/34631		9/1997	A61K 39/395
WO	WO 98/39482		9/1998	C12Q 1/68
WO	WO 00/42561		7/2000	G06F 19/00
WO	WO 00/71694		11/2000	C12N 15/06
WO	WO 01/01748		1/2001	
WO	WO 01/48145	A2	7/2001	
WO	WO 01/55366	A1	8/2001	C12N 12/10
WO	WO 01/57211	A1	8/2001	C07K 14/725
WO	WO 01/62908	A2	8/2001	C12N 15/10
WO	WO 01/70947		9/2001	C12N 15/10
WO	WO 01/83525		11/2001	C07K 14/00
WO	WO 01/88159		11/2001	C12N 15/62
WO	WO 02/06469		1/2002	C12N 15/10
WO	WO 02/32925		4/2002	
WO	WO 02/44215		6/2002	C07K 16/00
WO	WO 02/059263	A2	8/2002	
WO	WO 02/060919		8/2002	
WO	WO 02/066636	A2	8/2002	C12N 15/10
WO	WO 02/088171		11/2002	
WO	WO 03/012100		2/2003	C12N 15/10
WO	WO 03/029456		4/2003	C12N 15/00
WO	WO 03/075840		9/2003	
WO	WO 2004/018674		3/2004	C12N 15/10
WO	WO 2004/033685	A1	4/2004	C12N 15/12
WO	WO 2004/041862	A2	5/2004	C07K 16/00
WO	WO 2004/044004		5/2004	C07K 14/705
WO	WO 2004/044011		5/2004	
WO	WO 2004/050705		6/2004	C07K 14/705
WO	WO 2004/074322		9/2004	C07K 14/725
WO	WO 2005/021595		3/2005	C07K 19/00
WO	WO 2005/113595	A2	12/2005	C07K 14/705
WO	WO 2005/114215		12/2005	G01N 33/68
WO	WO 2005/116646	A1	12/2005	G01N 33/50
WO	WO 2006/033700		3/2006	
WO	WO 2006/036834		4/2006	C07K 19/00
WO	WO 2006/037960	A2	4/2006	C07K 14/705
WO	WO 2006/054096	A2	5/2006	C12N 15/62
WO	WO 2006/056733	A1	6/2006	C12N 15/12
WO	WO 2006/072620		7/2006	C07K 16/00
WO	WO 2006/087637		8/2006	C07K 16/32
WO	WO 2008/003103		1/2008	C07K 16/00
WO	WO 2008/003116		1/2008	C12N 15/09
WO	WO 2008/119096		10/2008	C07K 16/00
WO	WO 2009/000006		12/2008	C07K 16/00
WO	WO 2009/099961		8/2009	C07K 16/00
WO	WO 2009/132876	A1	11/2009	A61K 39/395

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO 2011/003811	1/2011	C07K 16/00
WO	WO 2012/007167	1/2012	C07K 16/32
WO	WO 2015/049537	4/2015	C12Q 1/68

OTHER PUBLICATIONS

Short et al. (Dec. 1, 1995) *Journal of Biological Chemistry* vol. 270 pp. 28541 to 28550.*

Bird, et al., "Single-Chain Antigen-Binding Proteins," *Science*, vol. 242, No. 4877, p. 423, Oct. 21, 1988.

Brawley, et al., "Complementarity-Determining Region 1 Sequence Requirements Drive Limited V α Usage in Response to Influenza Hemagglutinin in 307-319 Peptide¹," *The Journal of Immunology*, vol. 168, No. 8, pp. 3894-3901, Apr. 15, 2002.

Brekke, et al., "Therapeutic antibodies for human diseases at the dawn of the twenty-first century," *nature Reviews Drug Discovery*, vol. 2, No. 1, pp. 52-62, Jan. 2003.

Bunn, Jr., et al., "Expression of Her-2/ *neu* in Human Lung Cancer Cell Lines by Immunohistochemistry and Fluorescence in Situ Hybridization," *Clinical Cancer Research*, vol. 7, pp. 3239-3250, Oct. 2001.

Chlewicki, et al., "High-Affinity, Peptide-Specific T Cell Receptors Can be Generated by Mutations in CDR1, CDR2 or CDR3," *Journal of Molecular Biology*, vol. 346, No. 1, pp. 223-239, Feb. 11, 2005.

DiGiusto, et al., "An Analysis of Sequence Variation in the β Chain Framework and Complementarity Determining Regions of an Allo-Reactive T Cell Receptor," *Molecular Immunology*, vol. 31, No. 9, pp. 693-699, Jan. 1994.

Dunn, et al., "Directed Evolution of Human T Cell Receptor CDR2 Residues by Phage Display Dramatically Enhances Affinity for Cognate Peptide-MHC Without Increasing Apparent Cross-Reactivity," *Protein Science*, vol. 15, pp. 710-721, Published by Cold Spring Harbor Laboratory Press, 2006.

Esteve, et al., "Molecular Predictors of Response to Trastuzumab and Lapatinib in Breast Cancer," *Nature Reviews, Clinical Oncology*, vol. 7, No. 2, Feb. 1, 2010, pp. 98-107.

Fountzilas, et al., "A randomized phase III study comparing three anthracycline-free taxane-based regimens, as first line chemotherapy, in metastatic breast cancer," *Breast Cancer Research and Treatment*, vol. 115, pp. 87-99, 2009.

Hoogenboom, H.R., "Selecting and screening recombinant antibody libraries," *Nature in Biotechnology*, vol. 23, No. 9, pp. 1105-1116, Sep. 2005.

Iyengar, et al., "A Pilot Study of Dose-Dense Paclitaxel With Trastuzumab and Lapatinib for Node-negative HER2-Overexpressed Breast Cancer," *Clinical Breast Cancer*, vol. 16, No. 2, pp. 87-94, Apr. 2016.

Krebber et al., "Selectively-infective Phage (SIP): A Mechanistic Dissection of a Novel in vivo Selection for Protein-ligand Interactions," *Journal of Molecular Biology*, vol. 268, No. 9, pp. 607-618, May 9, 1997.

Laugel, et al., "Design of Soluble Recombinant T Cell Receptors for Antigen Targeting and T Cell Inhibition," *Theo Journal of Biological Chemistry*, vol. 280, No. 3, pp. 1882-1892, Jan. 2005.

Leung, et al., "A HER2-specific Modified Fc Fragment (Fcab) Induces Antitumor Effects Through Degradation of HER2 and Apoptosis," *Molecular Therapy*, vol. 23, No. 11, pp. 1722-1733, Nov. 2015.

Liang, et al., "Cross-species Vascular Endothelial Growth Factor (VEGF)-blocking Antibodies Completely Inhibit the Growth of Human Tumor Xenografts and Measure the Contribution of Stromal VEGF," *The Journal of Biological Chemistry*, vol. 28, No. 2, pp. 951-961, Jan. 13, 2006.

Molloy, et al., "Soluble T Cell Receptors: Novel Immunotherapies," *Current Opinion in Pharmacology*, vol. 5, Issue 4, pp. 438-443, Aug. 2005.

Nakauchi, et al., "Molecular cloning of Lyt-2, a Membrane Glycoprotein Marking a Subset of Mouse T Lymphocytes: Molecu-

lar Homology to its Human Counterpart, Leu-2/T8, and to Immunoglobulin Variable Regions," *Proceeding of the National Academy of Science of the United States of America*, vol. 82, No. 15, pp. 5126-5130, Aug. 1, 1985.

Philippidis, "Companion Diagnostics: 52 Pick-Up," *Genetic Engineering & Biotechnology News, Insight & Intelligence*, 12 pages, May 13, 2013.

Richman, et al., "Development of a Novel Strategy for Engineering High-Affinity Proteins by Yeast Display," *Protein Engineering, Design & Selection*, vol. 19, No. 6, pp. 255-264, 2006.

Richman, et al., "Structural features of T cell receptor variable regions that enhance domain stability and enable expression as single-chain V α V β fragments," *Molecular Immunology*, vol. 46, Issue 5, pp. 902-916, Feb. 2009 (Abstract).

Stagg, et al., "Anti-ErbB-2 mAb therapy requires type I and II interferons and synergizes with anti-PD-1 or anti-CD137 mAb therapy," *Proceedings of the National Academy of Sciences, USA*, vol. 108, No. 17, pp. 7142-7147, Apr. 26, 2011.

Tolaney, et al., "Adjuvant Paclitaxel and Trastuzumab for Node-Negative, HER2-Positive Breast Cancer," *The New England journal of Medicine*, vol. 372, pp. 134-141, Jan. 8, 2015.

Traxlmayr et al., "Directed Evolution of Her2/neu-binding IgG1-Fc for Improved Stability and Resistance to Aggregation by Using Yeast Surface Display," *Protein Engineering, Design & Selection*, vol. 26, No. 4, pp. 255-265, 2013.

Willcox, et al., "Production of Soluble $\alpha\beta$ T-cell Receptor Heterodimers Suitable for Biophysical Analysis of Ligand Binding," *Protein Science*, vol. 8, No. 11, pp. 2418-2423, Nov. 1999.

Wülfing, et al., "Correctly Folded T-cell Receptor Fragments in the Periplasm of *Escherichia coli*: Influence of Folding Catalysts," *Journal of Molecular Biology*, vol. 242, Issue 5, pp. 655-669, Oct. 6, 1994.

International Searching Authority, Written Opinion of International Searching Authority, Application No. PCT/GB2014/052994, 6 pages, Jan. 9, 2015.

European Patent Office, International Search Report, Application No. PCT/GB2014/052994, 5 pages, Jan. 9, 2015.

European Patent Office, International Search Report and Written Opinion, Application No. PCT/EP2016/057800, 18 pages, Jun. 24, 2016.

The International Bureau of WIPO, Switzerland, Authorized Officer Athina Nickitas-Etienne, International Preliminary Report on Patentability with Written Opinion of the International Searching Authority, International Application PCT/GB2014/052994, 7 pages, Apr. 5, 2016.

U.S. Appl. No. 11/722,517, filed Jun. 21, 2007.

U.S. Appl. No. 12/307,582, filed Jan. 5, 2009.

U.S. Appl. No. 12/307,578, filed Jun. 26, 2007.

U.S. Appl. No. 12/307,569, filed Sep. 21, 2009.

U.S. Appl. No. 13/086,897, filed Apr. 14, 2011.

U.S. Appl. No. 13/149,871, filed May 31, 2011.

U.S. Appl. No. 13/151,207, filed Jun. 1, 2011.

U.S. Appl. No. 13/228,559, filed Sep. 9, 2011.

U.S. Appl. No. 13/377,817, filed Dec. 12, 2011.

U.S. Appl. No. 13/434,765, filed Mar. 29, 2012.

U.S. Appl. No. 13/482,926, filed May 29, 2012.

U.S. Appl. No. 14/470,425, filed Aug. 27, 2014.

U.S. Appl. No. 14/559,662, filed Dec. 1, 2014.

U.S. Appl. No. 14/629,760, filed Feb. 24, 2015.

U.S. Appl. No. 14/853,919, filed Sep. 14, 2015.

U.S. Appl. No. 15/004,692, filed Jan. 22, 2016.

U.S. Appl. No. 15/087,272, filed Mar. 31, 2016.

Adachi et al., "Interaction Between the Antigen and Antibody Is Controlled by the Constant Domains: Normal Mode Dynamics of the HEL-HyHEL-10 Complex," *Protein Science*, vol. 12, No. 10, pp. 2125-2131, Oct. 2003.

Adib-Conquy et al., "Effect of Amino Acid Substitutions in the Heavy Chain CDR3 of an Autoantibody On Its Reactivity," *International Immunology*, vol. 10, No. 3, pp. 341-346, Mar. 1998.

Altschul et al., "Local Alignment Statistics," *Methods in Enzymology* vol. 266, pp. 460-480, 1996.

(56)

References Cited

OTHER PUBLICATIONS

- Amstutz et al., "In vitro Display Technologies: Novel Developments and Applications," *Current Opinion Biotechnology*, vol. 12, No. 4, pp. 400-405, Aug. 2001.
- Asano et al., "Humanization of the Bispecific Epidermal Growth Factor Receptor \times CD3 Diabody and Its Efficacy as a Potential Clinical Reagent," *Clinical Cancer Research*, vol. 12, No. 13, p. 4036-4042, Jul. 1, 2006.
- Auf der Maur et al., "Antigen-Independent Selection of intracellular Stable Antibody Frameworks," *Methods*, vol. 34, No. 2, pp. 215-224, Oct. 2004.
- Barbas III, et al., "Semisynthetic Combinatorial Antibody Libraries: A Chemical Solution to the Diversity Problem," *Proceeding of the National Academy of Science, USA*, vol. 89, No. 10, pp. 4457-4461, May 15, 1992.
- Barclay, "Membrane Proteins With Immunoglobulin-Like Domains—A Master Superfamily of interaction Molecules," *Seminars in Immunology*, vol. 15, No. 4, pp. 215-223, Aug. 2003.
- Batey et al., "Abstract B123: Preclinical Evaluation of FS102: A HER2-Specific Fcab With a Novel Mechanism of Action," *Molecular Cancer Therapeutics*, vol. 12, Supplement 11, B123, Nov. 2013.
- Batey et al., Poster: "Pre-Clinical Evaluation of FS102: A HER2 Specific Fcab With a Novel Mechanism of Action," AACR-NCI-EORTC International Conference on Molecular Targets and Cancer Therapeutics, Boston, 1 page, Oct. 21, 2013.
- Benhar et al., "Highly Efficient Selection of Phage Antibodies Mediated by Display of Antigen as Lpp-OmpA' Fusions on Live Bacteria," *Journal of Molecular Biology*, vol. 301, No. 4, pp. 893-904, Aug. 25, 2000.
- Berntzen et al., "Prolonged and Increased Expression of Soluble Fc Receptors, IgG and a TCR-Ig Fusion Protein by Transiently Transfected Adherent 293E Cells," *Journal Immunological Methods*, vol. 298, No. 1-2, pp. 93-104, Mar. 2005.
- Berntzen et al., "Characterization of an Fc γ /RI-Binding Peptide Selected by Phage Display," *Protein Engineering, Design & Selection*, vol. 19, No. 3, pp. 121-128, Jan. 19, 2006.
- Berry et al., "Development of Functional Human Monoclonal Single-Chain Variable Fragment Antibody Against HIV-1 from Human Cervical B Cells," *Hybrid. Hybridoma*, vol. 22, No. 2, pp. 97-108, Apr. 2003.
- Binz et al., "High-Affinity Binders Selected From Designed Ankyrin Repeat Protein Libraries," *Nature Biotechnology*, vol. 22, No. 5, pp. 575-582, May 2004.
- Binz et al., "Engineering Novel Binding Proteins From Nonimmunoglobulin Domains," *Nature Biotechnology*, vol. 23, pp. 1257-1268, Oct. 6, 2005.
- Boder et al., "Yeast Surface Display for Screening Combinatorial Polypeptide Libraries," *Nature Biotechnology*, vol. 15, No. 6, pp. 553-557, Jun. 1997.
- Boder et al., "Yeast Surface Display for Directed Evolution of Protein Expression, Affinity, and Stability," *Methods in Enzymology*, vol. 328, pp. 430-444, 2000.
- Boder et al., "Directed Evolution of Antibody Fragments With Monovalent Femtomolar Antigen-Binding Affinity," *Proceedings of the National Academy of Science*, vol. 97, No. 20, pp. 10701-10705, Sep. 26, 2000.
- Bork et al., "The Immunoglobulin Fold. Structural Classification, Sequence Patterns and Common Core," *Journal of Molecular Biology*, vol. 242, pp. 309-320, 1994.
- Boulter et al., "Stable, Soluble, High-Affinity, Engineered T Cell Receptors: Novel Antibody-Like Proteins for Specific Targeting of Peptide Antigens," *Clinical and Experimental Immunology*, pp. 454-460, 2005.
- Bowie et al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions" *Science*, vol. 247, No. 4948, pp. 1306-1310, Mar. 16, 1990.
- Burgess et al., "Possible Dissociation of the Heparin-binding and Mitogenic Activities of Heparin-binding (Acidic Fibroblast) Growth Factor-1 from Its Receptor-binding Activities by Site Directed Mutagenesis of a Single Lysine Residue," *Journal of Cell Biology*, vol. 111, pp. 2129-2138, Nov. 1990.
- Cabilly et al., "Generation of Antibody Activity From Immunoglobulin Polypeptide Chains Produced in *Escherichia coli*," *Proceedings of National Academy of Science USA*, vol. 81, pp. 3273-3277, Jun. 1984.
- Caldas et al., "Humanization of the Anti-CD18 Antibody 6.7: An Unexpected Effect of a Framework Residue in Binding to Antigen," *Molecular Immunology*, vol. 39, No. 15, pp. 941-952, May 2003.
- Carter, "Bispecific Human IgG by Design," *Journal of Immunological Methods*, vol. 248, pp. 7-15, 2001.
- Carter et al., "High Level *Escherichia coli* Expression and Production of a Bivalent Humanized Antibody Fragment," *Biotechnology*, vol. 10, No. 2, pp. 163-167, Feb. 1992.
- Chen et al., "Isolation of High-Affinity Ligand-Binding Proteins by Periplasmic Expression With Cytometric Screen (PECS)," *Nature Biotechnology* vol. 19, pp. 537-542, Jun. 2001.
- Chien et al., "Significant Structural and Functional Change of an Antigen-Binding Site by a Distant Amino Acid Substitution: Proposal of a Structural Mechanism," *Proceeding of the National Academy of Science USA*, vol. 86, No. 14, pp. 5532-5536, Jul. 1989.
- Chirino et al., "Minimizing the Immunogenicity of Protein Therapeutics," *Drug Discovery Today*, vol. 9, No. 2, pp. 82-90, Jan. 2004.
- Cho et al., "Structure of the Extracellular Region of HER2 Alone and in Complex with the Herceptin Fab," *Nature*, vol. 421, pp. 756-760, Feb. 13, 2003.
- Coco et al., "DNA Shuffling Method for Generating Highly Recombined Genes and Evolved Enzymes," *Nature Publishing Group*, vol. 19, pp. 354-359, Apr. 2001.
- Conrath et al., "Antigen Binding and Solubility Effects upon the Veneering of a Camel VHH in Framework-2 to Mimic a VH," *Journal of Molecular Biology*, vol. 350, pp. 112-125, 2005.
- Cornish-Bowden, "Nomenclature for Incompletely Specified Bases in Nucleic Acid Sequences: Recommendations 1984," *Nucleic Acids Research*, vol. 3, No. 9, pp. 3021-3030, May 10, 1985.
- Cortez-Retamozo et al., "Efficient Tumor Targeting by Single-Domain Antibody Fragments of Camels," *International Journal of Cancer*, vol. 98, pp. 456-462, 2002.
- Cramer et al., "DNA Shuffling of a Family of Genes From Diverse Species Accelerates Directed Evolution," *Nature* vol. 391, pp. 288-291, Jan. 15, 1998.
- Dall'Acqua et al., "Increasing the Affinity of a Human IgG1 for the Neonatal Fc Receptor: Biological Consequences," *The Journal of Immunology*, vol. 169, pp. 5171-5180, 2002.
- Dall'Acqua et al., "Modulation of the Effector Functions of a Human IgG1 through Engineering of Its Hinge Region," *The Journal of Immunology*, vol. 177, pp. 1129-1138, 2006.
- de Jager et al., "Simultaneous Detection of 15 Human Cytokines in a Single Sample of Stimulated Peripheral Blood Mononuclear Cells," *Clinical Diagnostic Laboratory Immunology*, vol. 10, No. 1, pp. 133-139, 2003.
- Doi et al., "Screening of Conformationally Constrained Random Polypeptide Libraries Displayed on a Protein Scaffold," *Cellular and Molecular Life Sciences*, vol. 54, No. 5, pp. 394-404, May 1998.
- Dottorini et al., "Crystal Structure of a Human VH: Requirements for Maintaining a Monomeric Fragment," *Biochemistry*, vol. 43, No. 3, pp. 622-628, Jan. 27, 2004.
- Ewert et al., "Stability Improvement of Antibodies for Extracellular and Intracellular Applications: CDR Grafting to Stable Frameworks and Structure-Based Framework Engineering," *Methods*, vol. 34, pp. 184-199, 2004.
- F-star, "F-star Alpha: A New Asset Centric Company," 15 pages, Feb. 11, 2014.
- Felgenhauer et al., "Nucleotide Sequences of the Cdnas Encoding the V-Regions of H- and L-Chains of a Human Monoclonal Antibody Specific to HIV-1-gp41," *Nucleic Acids Research*, vol. 18, No. 16, pp. 4927, 1990.
- Fellouse et al., "Synthetic Antibodies From a Four-Amino-Acid Code: A Dominant Role for Tyrosine in Antigen Recognition," *Proceeding of the National Academy of Science*, vol. 101, No. 34, pp. 12467-12472, Aug. 24, 2004.

(56)

References Cited

OTHER PUBLICATIONS

- Fellouse et al., "Molecular Recognition by a Binary Code," *The Journal of Molecular Biology*, vol. 348, No. 5, pp. 1153-1162, May 20, 2005.
- Fellouse et al., "Tyrosine Plays a Dominant Functional Role in the Paratope of a Synthetic Antibody Derived from a Four Amino Acid Code," *The Journal of Molecular Biology*, vol. 357, pp. 100-114, 2006.
- Fields et al., "A Novel Genetic System to Detect Protein-Protein Interactions," *Nature* vol. 340, pp. 245-246, Jul. 20, 1989.
- Fitzgerald, "In vitro Display Technologies—New Tools for Drug Discovery," vol. 5, No. 6, pp. 253-258, Jun. 2000.
- Foote, "Antibody Framework Residues Affecting the Conformation of the Hypervariable Loops," *Journal of Molecular Biology*, vol. , No. 2, pp. 487-499, Mar. 20, 1992.
- Gao et al., "Making Artificial Antibodies: A Format for Phage Display of Combinatorial Heterodimeric Arrays," *Proceedings of the National Academy of Sciences of USA*, vol. 96, No. 11 pp. 6025-6030, May 25, 1999.
- Georgiou et al., "Practical Applications of Engineering Ggram-Negative Bacterial Cell Surfaces," *Trends in Biotechnology*, vol. 11, No. 1, pp. 6-10, Jan. 1993.
- Georgiou et al., "Display of Heterologous Proteins on the Surface of Microorganisms: From the Screening of Combinatorial Libraries to Live Recombinant Vaccines," *Nature Biotechnology* vol. 15, No. 1, pp. 29-34, Jan. 1997.
- Ghahroudi et al., "Selection and Identification of Single Domain Antibody Fragments from Camel Heavy-Chain Antibodies," *FEBS Letters*, vol. 414, No. 3, pp. 521-526, Sep. 15, 1997.
- Giusti et al., "Somatic Diversification of S107 from an Antiphosphocholine to an Anti-DNA Autoantibody is Due to a Single Base Change in its Heavy Chain Variable Region," *Proceedings of the National Academy of Science, USA*, vol. 84, No. 9, pp. 2926-2930, May 1987.
- Goncalves, "Fluorescent Labeling of Biomolecules with Organic Probes," *Chemical Review*, vol. 109, No. 1, pp. 190-212, 2009.
- Gram et al., "In vitro Selection and Affinity Maturation of Antibodies from a Naive Combinatorial Immunoglobulin Library," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 89, No. 8, pp. 3576-3580, Apr. 15, 1992.
- Halaby et al., "The Immunoglobulin Fold Family: Sequence Analysis and 3D Structure Comparisons," *Protein Engineering*, vol. 12, No. 7, pp. 563-571, Jul. 1999.
- Hanes et al., "In vitro Selection and Evolution of Functional Proteins by Using Ribosome Display," *Proceedings of the National Academy of Science, USA*, vol. 94, No. 19, pp. 4937-4942, May 1997.
- Harriman et al., "Multiplexed Elispot Assay," *Journal Immunology Methods*, vol. 341, No. 1-2, pp. 127-134, Feb. 28, 2009.
- Hasenhindl et al., "Stability Assessment on a Library Scale: A Rapid Method for the Evaluation of the Commutability and Insertion of Residues in C-Terminal Loops of the CH3 domains of IgG1-Fc," *Protein Engineering, Design and Selection*, vol. 26, issue 10, pp. 675-682, Oct. 2013.
- Hasenhindl et al., "Creating Stable Stem Regions for Loop Elongation in Fcabs—Insights from Combining Yeast Surface Display, in Sihco Loop Reconstruction and Molecular Dynamics Simulations," *Biochimica et Biophysica Acta*, vol. 1844, No. 9, pp. 1530-1540, Sep. 2014.
- Haurum, "How to Leverage Oncogene Addiction: Targeted Biological Therapy Inducing Growth Factor Receptor Internalization and Degradation," *PEPtalk: The Protein Science Week* Jan. 19, 2015.
- Hayhurst et al., "High-throughput Antibody Isolation," *Current Opinion in Chemical Biology*, vol. 5, No. 6, pp. 683-689, Dec. 2001.
- He et al., "Structure of a Human Monoclonal Antibody Fab Fragment Against gp41 of Human Immunodeficiency Virus Type 1," *Proceedings of the National Academy of Science, USA* vol. 89, No. 15, pp. 7154-7158, Aug. 1, 1992.
- Hermeling et al., "Structure-Immunogenicity Relationships of Therapeutic Proteins," *Pharmaceutical Research*, vol. 21, No. 6, pp. 897-903, Jun. 2004.
- Holler et al., "In vitro Evolution of a T Cell Receptor with High Affinity for Peptide/MHC," *Proceedings of the National Academy of Science, USA*, vol. 97, No. 10, pp. 5387-5392, May 9, 2000.
- Holliger et al., "Engineered Antibody Fragments and the Rise of Single Domains," *Nature Biotechnology*, vol. 23, No. 9, pp. 1126-1136, Sep. 2005.
- Hoogenboom et al., "Multi-Subunit Proteins on the Surface of Filamentous Phage: Methodologies for Displaying Antibody (Fab) Heavy and Light Chains," *Nucleic Acids Research*, vol. 19, No. 15, pp. 4133-4137, Aug. 11, 1991.
- Hoover et al., "DNAWorks: An Automated Method for Designing Oligonucleotides for PCR-based Gene Synthesis," *Nucleic Acids Research*, vol. 30, No. 10, May 15, 2002.
- Hosse et al., "A New Generation of Protein Display Scaffolds for Molecular Recognition," *Protein Science*, vol. 15, No. 1, pp. 14-27, Jan. 2006.
- Hufton et al., "Development and Application of Cytotoxic T Lymphocyte-Associated Antigen 4 As a Protein Scaffold for the Generation of Novel Binding Ligand," *FEBS Letters*, vol. 465, No. 3, pp. 225-231, Jun. 23, 2000.
- Huston et al., "Protein Engineering of Antibody Binding Sites: Recovery of Specific Activity in an Anti-Digoxin Single-Chain Fv Analogue Produced in *Escherichia coli*," *Proceedings of the National Academy of Science, USA*, vol. 85, No. 16, pp. 5879-5883, Aug. 1988.
- Isaac et al., Poster: "Pre-Clinical Evaluation of FS102: A HER2-Specific Fcab with a Novel Mechanism of Action," *AstraZeneca-MedImmune—Cambridge Cancer Centre Symposium*, 1 page, Mar. 25, 2014.
- Janeway et al., "Immunobiology, the Immune System in Health and Disease," 6th Edition, Garland Science, 2005.
- Jez et al., "Significant Impact of Single N-Glycan Residues on the Biological Activity of Fc-Based Antibody-Like Fragments," *The Journal of Biological Chemistry*, vol. 287, No. 9, pp. 24313-24319, Jul. 13, 2012.
- Johnsson et al., "Split Ubiquitin As a Sensor of Protein interactions in vivo," *Proceedings of the National Academy of Science, USA* vol. 91, No. 22, pp. 10340-10344, Oct. 1994.
- Jones et al., "Replacing the Complementarity-Determining Regions in a Human Antibody With Those From a Mouse," *Nature*, vol. 321, No. 6069, pp. 522-525, May 29, 1986.
- Jung et al., "Surface Display of Zymomonas Mobilis Levansucrase by Using the Ice-Nucleation Protein of Pseudomonas Syringae," *Nature Biotechnology* vol. 16, No. 6, 576-580, Jun. 1998.
- Kainer et al., "Correlation Between CD16a Binding and Immuno Effector Functionality of an Antigen Specific Immunoglobulin Fc Fragment (Fcab)," *Archives of Biochemistry and Biophysics*, vol. 526, No. 2, pp. 154-158, Oct. 23, 2012.
- Kang et al., "Linkage of Recognition and Replication Functions by Assembling Combinatorial Antibody Fab Libraries Along Phage Surfaces," *Proceedings of the National Academy of Science, USA*, vol. 88, No. 10, pp. 4363-4366, May 15, 1991.
- Kang et al., "Human Neutralizing Fab Molecules against Severe Acute Respiratory Syndrome Coronavirus Generated by Phage Display," *Clinical and Vaccine Immunology*, vol. 13, No. 8, pp. 953-957, Aug. 2006.
- Kashmiri et al., "SDR Grafting—A New Approach to Antibody Humanization," *Methods*, vol. 36, No. 1, pp. 25-34, May 2005.
- Kay et al., "Phage Display of Peptides and Proteins: A Laboratory Manual," Academic Press, 1996.
- Kettleborough et al., "Humanization of a Mouse Monoclonal Antibody by CDR-Grafting: the importance of Framework Residues on Loop Conformation," *Protein Engineering*, vol. 4, No. 7, pp. 773-783, Oct. 1991.
- Kieke et al., "Selection of Functional T Cell Receptor Mutants From a Yeast Surface-Display Library," *Proceedings of the National Academy of Science, USA*, vol. 96, No. 10, pp. 5651-5656, May 1999.

(56)

References Cited

OTHER PUBLICATIONS

- Kikuchi et al., "Novel Family Shuffling Methods for the in vitro Evolution of Enzymes," *Gene*, vol. 236, No. 1, pp. 159-167, Aug. 5, 1999.
- Kikuchi et al., "An Effective Family Shuffling Method Using Single-Stranded DNA," *Gene*, vol. 243, No. 1-2, pp. 133-137, Feb. 8, 2000.
- Kohl et al., "Cloning and Expression of an HIV-1 Specific Single-Chain Fv Region Fused to *Escherichia coli* Alkaline Phosphatase," *Annals of the New York Academy of Science*, vol. 646, pp. 106-114, Dec. 27, 1991.
- Koide et al., "High-Affinity Single-Domain Binding Proteins with a Binary-Code Interface," *Proceedings of the National Academy of Science, USA*, vol. 104, No. 16, pp. 6632-6637, Apr. 17, 2007.
- Koivunen et al., "Selection of Peptides Binding to the $\alpha 5 \beta 1$ Integrin from Phage Display Library," *The Journal of Biological Chemistry*, vol. 268, No. 27, pp. 20205-20210, Sep. 25, 1993.
- Kolkman et al., "Directed Evolution of Proteins by Exon Shuffling," *Nature Biotechnology*, vol. 19, pp. 423-428, May 2001.
- Konig, "Interactions Between MHC Molecules and Co-Receptors of the TCR," *Current Opinion in Immunology*, pp. 75-83, Mar. 2002.
- Kontermann, "Dual Targeting Strategies With Bispecific Antibodies," *MAbs*, vol. 4, No. 2, pp. 182-197, Mar. 2012.
- Koren et al., "Immune Responses to Therapeutic Proteins in Humans—Clinical Significance, Assessment and Prediction," *Current Pharmaceutical Biotechnology*, vol. 3, pp. 349-360, 2002.
- Kufer et al., "A Revival of Bispecific Antibodies," *Trends in Biotechnology*, vol. 22, No. 5, pp. 238-244, May 2004.
- Kunkel et al., "Rapid and Efficient Site-Specific Mutagenesis Without Phenotype Selection," *Proceedings of National Academy of Sciences*, vol. 82, pp. 488-492, Jan. 1985.
- Laffly et al., "Monoclonal and Recombinant Antibodies, 30 years after . . .," *Human Antibiotics*, vol. 14, pp. 33-55, 2005.
- Lauvrak et al., "Identification and Characterisation of C1q-Binding Phage Displayed Peptides," *Biology Chemistry*, vol. 378, No. 12, pp. 1509-1519, Dec. 1997.
- Lazar et al., "Transforming Growth Factor α : Mutation of Aspartic Acid 47 and Leucine 48 Results in Different Biological Activities," *Molecular and Cellular Biology*, pp. 1247-1252, Mar. 1988.
- Le Gall et al., "Effect of Linker Sequences Between the Antibody Variable Domains on the Formation, Stability and Biological Activity of a Bispecific Tandem Diabody," *Protein Engineering, Design & Selection*, vol. 17, No. 44, pp. 357-366, 2004.
- Lea et al., "Analysis of Antigenic Surfaces of Proteins," *Federation of American Societies for Experimental Biology*, vol. 9, No. 1, pp. 87-93, Jan. 1995.
- Lederman et al., "A Single Amino Acid Substitution in a Common African Allele of the CD4 Molecule Abolishes Binding of the Monoclonal Antibody, OKT4," *Molecular Immunology* vol. 28, No. 11, pp. 1171-1181, Nov. 1991.
- Lee et al., "Surface-Displayed Viral Antigens on *Salmonella* Carrier Vaccine," *Nature Biotechnology*, vol. 18, No. 6, pp. 645-648, Jun. 2000.
- Lefranc et al., "IMGT, the International ImMunoGeneTics Database," *Nucleic Acids Research*, vol. 27, No. 1, pp. 209-212, Jan. 1, 1999.
- Lefranc, "IMGT, the International ImMunoGeneTics database," *Nucleic Acids Research*, vol. 29, No. 1, pp. 207-209, 2001.
- Lefranc, "IMGT, the International ImMunoGeneTics database," *Nucleic Acids Research*, vol. 31, No. 1, pp. 307-310, 2003.
- Lefranc et al., "IMGT Unique Numbering for Immunoglobulin and T Cell Receptor Constant Domains and Ig Superfamily C-like Domains," *Developmental and Comparative Immunology* vol. 27, pp. 55-77, 2003.
- Lefranc et al., "IMGT Unique Numbering for immunoglobulin and T Cell Receptor Constant Domains and Ig Superfamily C-like Domains," *Developmental and Comparative Immunology*, vol. 29, pp. 185-203, 2005.
- Lefranc et al., "IMGT, the International ImMunoGeneTics Information System," *Nucleic Acids Research*, vol. 33, Database issue, pp. D593-D597, 2005.
- Li et al., "Beta-Endorphin Omission Analogs: Dissociation of Immunoreactivity From Other Biological Activities," *Proceedings of the National Academy of Science, USA*, vol. 77, No. 6, pp. 3211-3214, Jun. 1980.
- Li et al., "Directed Evolution of Human T-Cell Receptors With Picomolar Affinities by Phage Display," *Nature Biotechnology*, vol. 23, No. 3, pp. 349-354, Mar. 2005.
- Lo Conte et al., "The Atomic Structure of Protein-Protein Recognition Sites," *Journal of Molecular Biology*, vol. 285, pp. 2177-2198, 1999.
- Lowman et al., "Selecting High-Affinity Binding Proteins by Monovalent Phage Display," *Biochemistry*, vol. 30, pp. 10832-10838, 1991.
- Lutz et al., "Creating Multiple Crossover DNA Libraries Independent of Sequence Identity," *Proceedings of the National Academy of Science, USA*, vol. 98, No. 20, pp. 11248-11253, Sep. 25, 2001.
- Malborg et al., "Selective Phage Infection Mediated by Epitope Expression on F Pilus," *Journal of Molecular Biology*, vol. 273, pp. 544-551, 1997.
- Marvin et al., "Recombinant Approaches to IgG-like Bispecific Antibodies," *Acta Pharmacologica Sinica*, vol. 6, pp. 649-658, Jun. 2005.
- Masuda et al., "The Role of Interface Framework Residues in Determining Antibody VH/VL, Interaction Strength and Antigen-Binding Affinity," *The FEBS Journal*, vol. 273, pp. 2184-2194, 2006.
- Mattheakis et al., "An in vitro Polysome Display System for Identifying Ligands From Very Large Peptide Libraries," *Proceedings of the National Academy of Science, USA*, vol. 91, pp. 9022-9026, Sep. 1994.
- Maynard et al., "Antibody Engineering," *Annual Review of Biomedical Engineering*, vol. 2, pp. 339-376, 2000.
- McCall et al., "Isolation and Characterization of an Anti-CD16 Single-chain Fv Fragment and Construction of an Anti-HER2/neu/anti-CD16 Bispecific scFv that Triggers CD16-dependent Tumor Cytolysis," *Molecular Immunology*, vol. 36, pp. 433-446, 1999.
- McCall et al., "Increasing the Affinity for Tumor Antigen Enhances Bispecific Antibody Cytotoxicity," *The Journal of immunology*, vol. 166, pp. 6112-6117, 2001.
- Merz et al., "The Protein Folding Problem and Tertiary Structure Prediction," Chapter 1, Authors Adrian Roitberg and Ron Elber, "Modeling Side Chains in Peptides and Proteins with Locally Enhanced Sampling/Simulated Annealing Method", Birkhauser, 584 pages, 1994.
- Merz et al., "The Protein Folding Problem and Tertiary Structure Prediction," Chapter 14, Authors J. Thomas Ngo, Joe Marks and Martin Karplus, "Computational Complexity. Protein Structure Prediction, and the Levinthal Paradox", Birkhauser, 9 pages, 1994.
- Miyazaki et al., "Changes in the Specificity of Antibodies by Site-Specific Mutagenesis Followed by Random Mutagenesis," *Protein Engineering*, vol. 12, No. 5, pp. 407-415, 1999.
- Moza et al., "Long-Range Cooperative Binding Effects in a T Cell Receptor Variable Domain," *Proceedings of the National Academy of Science, USA*, vol. 103, No. 26, pp. 9867-9872, Jun. 27, 2006.
- Munoz-Olaya, "Advancing Novel Modular Antibody Technology for Generating Bispecific Antibodies," *PEGS Lisbon*, Nov. 3-4, 2014.
- Nemoto et al., "In vitro virus: Bonding of mRNA Bearing Puromycin at the 3'-Terminal End to the C-terminal End of its Encoded Protein on the Ribosome in vitro," *FEBS Letters*, vol. 414, pp. 405-408, 1997.
- Nygren et al., "Scaffold for Engineering Novel Binding Sites in Proteins," *Current Biology, Engineering and Design*, vol. 7, pp. 463-469, 1997.
- Park et al., "Rationally Designed Anti-HER2/neu Peptide Mimetic Disables P185 HER2/neu Tyrosine Kinases in vitro and in vivo," *Nature Biotechnology*, vol. 18, pp. 194-198, Feb. 2000.
- Paul, *Fundamental Immunology*, Chapter 9, "Structure and Function of Immunoglobulins", Third Edition, Raven Press, pp. 292-295, 1993.

(56)

References Cited

OTHER PUBLICATIONS

- Pelletier et al., "Oligomerization Domain-Directed Reassembly of Active Dihydrofolate Reductase From Rationally Designed Fragments," *Proceedings of the National Academy of Science, USA*, vol. 95, pp. 12141-12146, Oct. 1998.
- Perosa et al., "CD20 Mimicry by a MAbs Rituximab-Specific Linear Peptide, A Potential Tool for Active Immunotherapy of Autoimmune Diseases," *Annals New York Academy of Sciences*, pp. 672-683, 2005.
- Presta et al., "Engineering Therapeutic Antibodies for Improved Function," *Biochemical Society*, vol. 30, No. 4, pp. 487-490, Mar. 2002.
- Privezentzev, Poster: "F-star: Advancing Novel Bispecific Antibody Biologics," *Gordon Research Conference; Antibody Biology & Engineering*, 1 page, Mar. 2014.
- Riechmann et al., "Single Domain Antibodies: Comparison of Camel VH and Camelized Human VH Domains," *Journal of Immunological Methods*, vol. 231, pp. 25-38, 1999.
- Roberts et al., "RNA-Peptide Fusions for the in vitro Selection of Peptides and Proteins," *Proceedings of the National Academy of Sciences, USA*, vol. 94, pp. 12297-12302, Nov. 1997.
- Rondot et al., "A Helper Phage to Improve Single-Chain Antibody Presentation in Phage Display," *Nature Biotechnology*, vol. 19, pp. 75-78, Jan. 2001.
- Roovers et al., "Efficient Inhibition of EGFR Signalling and of Tumour Growth by Antagonistic Anti-EGFR Nanobodies," *Cancer Immunology Immunotherapy*, vol. 56, pp. 303-317, 2007.
- Rudikoff et al., "Single Amino Acid Substitution Altering Antigen-Binding Specificity," *Proceedings of the National Academy of Sciences*, vol. 79, pp. 1979-1983, Mar. 1982.
- Ruiz et al., "IMGT, the International ImmunoGeneTics database," *Nucleic Acids Research*, vol. 28, No. 1, pp. 219-221, 2000.
- Rüker et al., "Expression of a Human Monoclonal Anti-HIV-1 Antibody in CHO Cells" *Annals of New York Academy of Sciences*, vol. 646, pp. 212-219, Dec. 27, 1991.
- Saerens et al., "Identification of a Universal VHH Framework to Graft Non-canonical Antigen-binding Loops of Camel Single-domain Antibodies," *Journal of Molecular Biology*, vol. 352, pp. 597-607, 2005.
- Salfield, "Isotype Selection in Antibody Engineering," *Nature Biotechnology*, vol. 25, No. 12, pp. 1369-1372, 2007.
- Schaffitzel et al., "Ribosome Display: an in vitro Method for Selection and Evolution of Antibodies From Libraries," *Journal of Immunological Methods*, vol. 231, pp. 119-135, 1999.
- Schmittl et al., "Application of the IFN- γ ELISPOT Assay to Quantify T Cell Responses Against Proteins," *Journal of Immunological Methods*, vol. 247, pp. 17-24, 2001.
- Shao et al., "Random-priming in vitro Recombination: An Effective Tool for Directed Evolution," *Nucleic Acids Research*, vol. 26, No. 2, pp. 681-683, 1998.
- Shields et al., "High Resolution Mapping of the Binding Site on Human IgG1 for Fc γ RI, Fc γ RII, Fc γ RIII, and Fc γ Rn and Design of IgG1 Variants with Improved Binding to the Fc γ RI," *The Journal of Biological Chemistry*, vol. 276, No. 9, pp. 6591-6604, 2001.
- Shusta et al., "Directed Evolution of a Stable Scaffold for T-cell Receptor Engineering," *Nature Biotechnology*, vol. 19, 6 pages, Jul. 2000.
- Sidhu et al., "Synthetic Therapeutic Antibodies," *Nature Chemical Biology*, vol. 2, No. 12, pp. 682-688, Dec. 2006.
- Simon et al., "A Functional Antibody Mutant With an Insertion in the Framework Region 3 Loop of the VH Domain: Implications for Antibody Engineering," *Protein Engineering*, vol. 5, No. 3, pp. 229-234, 1992.
- Skolnick et al., "From Genes to Protein Structure and Function: Novel Applications of Computational Approaches in the Genomic Era," *Trends in Biotechnology*, vol. 18, pp. 34-39, Jan. 2000.
- Smith, "Filamentous Fusion Phage: Novel Expression Vectors That Display Cloned Antigens on the Virion Surface," *Science*, vol. 228, No. 4705, pp. 1315-1317, Jun. 14, 1985.
- Spiridon et al., "Targeting Multiple Her-2 Epitopes with Monoclonal Antibodies Results in Improved Antigrowth Activity of a Human Breast Cancer Cell Line in Vviro and in Vivo," *Clinical Cancer Research*, vol. 8, pp. 1720-1730, Jun. 2002.
- Tangri et al., "Rationally Engineered Therapeutic Proteins with Reduced Immunogenicity," *The Journal of Immunology*, pp. 3187-3196, 2005.
- Traxlmayr et al., "Construction of pH-Sensitive Her2-binding IgG1-Fc by Directed Evolution," *Biotechnology*, vol. 9, pp. 1013-1022, 2014.
- Uhlenbroich, "F-star: Advancing Novel Bispecific Antibody Biologics," *Empowered Antibodies Congress*, 1 page, Jun. 18-19, 2014—Abstract.
- Vajdos et al., "Comprehensive Functional Maps of the Antigen-binding Site of an Anti-ErbB2 Antibody Obtained with Shotgun Scanning Mutagenesis," *Journal of Molecular Biology*, vol. 320 pp. 416-428, 2002.
- Virnekas et al., "Trinucleotide Phosphoramidites: Ideal Reagents for the Synthesis of Mixed Oligonucleotides for Random Mutagenesis," *Nucleic Acids Research*, vol. 22, No. 25, pp. 5600-5607, 1994.
- Visintin et al., Selection of Antibodies for Intracellular Function Using a Two-hybrid in vivo System, *Proceedings of the National Academy of Science, USA*, vol. 96, No. 21, pp. 11723-11728, Oct. 12, 1999.
- Vogt et al., "Construction of an Artificial Receptor Protein ("Anticalin") Based on the Human Apolipoprotein D," *ChemBioChem*, vol. 5, No. 2, pp. 191-199, Feb. 6, 2004.
- Wang et al., "Expression Patterns and Transcript Processing of ftt-1 and ftt-2, two C. elegans 14-3-3 Homologues," *Journal of Molecular Biology*, vol. 268, pp. 619-630, 1997.
- Wang et al., "Retargeting T Cells for HER2-Positive Tumor Killing by a Bispecific Fv-Fc Antibody," *PLOS One*, vol. 8, No. 9, pp. e75589-1-e75589-11, Sep. 23, 2013.
- Weaver-Feldhaus et al., "Yeast Mating for Combinatorial Fab Library Generation and Surface Display," *FEBS Letters*, vol. 564, No. 1-2, pp. 24-34, Apr. 23, 2004.
- Weber et al., "Class II-Restricted T Cell Receptor Engineered In Vitro for Higher Affinity Retains Peptide Specificity and Function," *Proceedings of the National Academy of Sciences*, vol. 102, No. 52, pp. 19033-19038, Dec. 27, 2005.
- Weiner et al., "Site-Directed Mutagenesis of Double-Stranded DNA by the Polymerase Chain Reaction," *Gene*, vol. 151, No. 1-2, pp. 119-124, Dec. 30, 1994.
- Whitehorn et al., "A Generic Method for Expression and Use of "Tagged" Soluble Versions of Cell Surface Receptors," *BioTechnology*, vol. 13, No. 11, pp. 1215-1219, Nov. 1995.
- Winkler et al., "Changing the Antigen Binding Specificity by Single Point Mutations of an Anti-p24 (HIV-1) Antibody," *The Journal of Immunology*, vol. 165, No. 8, pp. 4505-4514, Oct. 15, 2000.
- Winter et al., "Humanized Antibodies," *Immunology Today*, vol. 14, No. 6, pp. 243-246, 1993.
- Wittrup, "Protein Engineering by Cell-Surface Display," *Current Opinion in Biotechnology*, vol. 12, No. 4, pp. 395-399, Aug. 2001.
- Woisetschlager et al., "In vivo and in vitro Activity of an Immunoglobulin Fc Fragment (Fcab) with Engineered Her-2/neu Binding Sites," *Biotechnology Journal*, vol. 9, No. 6, pp. 844-851, Jun. 2014.
- Wozniak-Knopp et al., "Introducing Antigen-Binding Sites in Structural Loops of Immunoglobulin Constant Domains: Fc Fragments with Engineered HER2/neu-binding Sites and Antibody Properties," *Protein Engineering, Design & Selection*, vol. 23, No. 4, pp. 289-297, Apr. 2010.
- Wu et al., "Humanization of a Murine Monoclonal Antibody by Simultaneous Optimization of Framework and CDR Residues," *Journal of Molecular Biology*, vol. 294, No. 1, pp. 151-162, Nov. 19, 1999.
- Xiao et al., "A Large Library Based on a Novel (CH2) Scaffold: identification of HIV-1 Inhibitors," *Biochemical and Biophysical Research Communications*, vol. 387, No. 2, pp. 387-392, Sep. 18, 2009.
- Yanez et al., "Combinatorial Condon-Based Amino Acid Substitutions," *Nucleic Acids Research*, vol. 32, No. 20, pp. 1-10, 2004.

(56)

References Cited

OTHER PUBLICATIONS

Yau et al., "Affinity Maturation of a VhH by Mutational Hotspot Randomization," *Journal of Immunological Methods*, vol. 297, No. 1-2, pp. 213-224, Feb. 2005.

Zemlin et al., "Expressed Murine and Human CDR-H3 Intervals of Equal Length Exhibit Distinct Repertoires that Differ in their Amino Acid Composition and Predicted Range of Structures," *Journal of Molecular Biology*, vol. 334, No. 4, pp. 733-749, Dec. 5, 2003.

Zhao et al., "A Novel Strategy by the Action of Ricin that Connects Phenotype and Genotype without Loss of the Diversity of Libraries," *Journal of American Chemical Society* vol. 124, No. 4, pp. 538-543, Jan. 30, 2002.

Zhou et al., "A Novel Strategy by the Action of Ricin that Connects Phenotype and Genotype without Loss of the Diversity of Libraries," *Journal of American Chemical Society* vol. 124, No. 4, pp. 538-543, Jan. 30, 2002.

European Patent Office—Munich, Extended European Search Report, Application No. EP 14191631.2-1405, 12 pages, dated Jun. 16, 2015.

European Patent Office, Authorized Officer, Isabel Perez-Mato, International Searching Report, Application No. PCT/AT2008/000232, 4 pages, dated Oct. 13, 2008, 4 pages.

International Bureau of WIPO, Authorized officer Yolaine Cussac, International Preliminary Report on Patentability, International Application PCT/AT2008/000232, 10 pages, dated Jan. 5, 2010.

International Searching Authority, Written Opinion pertaining to International Application PCT/AT2008/000232, dated Oct. 12, 2008.

European Patent Office—Rijswijk Authorized Officer Rebecca Hix, International Search Report, dated Feb. 1, 2008, Application No. PCT/AT2007/000313, 3 pages.

International Searching Authority, Authorized Officer, Yolaine Cussac, Written Opinion pertaining to Application No. PCT/EP2009/052506, 8 pages, dated Nov. 2, 2010.

International Searching Authority, Authorized Officer Marie-Paul Toussaint, International Search Report and Written Opinion, Application No. PCT/EP2009/052509, 17 pages, dated Jun. 3, 2009.

European Patent Office, Communication pursuant to Article 94(3) EPC, European Patent Application No. 08 756 842.4, dated Jun. 14, 2010.

Rüker, F. "Modular Antibody Technology", F-Star Fact Sheet, online, Feb. 2008, www.boku.ac.at/fileadmin/BOKU-Topstories/20080702_Rueker_Factsheet.pdf.

* cited by examiner

Figure 1.





Figure 2



DISPLAY OF BINDING AGENTS

CROSS REFERENCE

This application is a continuation of U.S. application Ser. No. 12/666,618 filed Dec. 23, 2009, which is a 371 U.S. national phase application of International Application No. PCT/AT08/00232 filed Jun. 26, 2008, which claims the benefit of U.S. Provisional Application No. 60/946,287 filed Jun. 26, 2007 and U.S. Provisional Application No. 61/049,826 filed May 2, 2008. The contents of all of the above applications are hereby incorporated by reference in their entirety.

SEQUENCE LISTING

The attached sequence listing "3906_1029_revised_seq_listing_2_23" created Feb. 24, 2016, size 128 kb) is hereby incorporated by reference in its entirety.

The invention relates to a method of preparing a genetic package displaying oligomers of modular antibody domains binding to a target and to a scaffold ligand and vectors and libraries of bivalent genetic packages produced by these methods. The invention further relates to methods of selecting suitable linker sequences for use in such oligomer display.

Monoclonal antibodies have been widely used as a scaffold for binding agents. The basic antibody structure will be explained here using as example an intact IgG1 immunoglobulin.

Two identical heavy (H) and two identical light (L) chains combine to form the Y-shaped antibody molecule. The heavy chains each have four domains. The amino terminal variable domains (VH) are at the tips of the Y. These are followed by three constant domains: CH1, CH2, and the carboxy-terminal CH3, at the base of the Y's stem. A short stretch, the switch, connects the heavy chain variable and constant regions. The hinge connects CH2 and CH3 (the Fc fragment) to the remainder of the antibody (the Fab fragments). One Fc and two identical Fab fragments can be produced by proteolytic cleavage of the hinge in an intact antibody molecule. The light chains are constructed of two domains, variable (VL) and constant (CL), separated by a switch.

Disulfide bonds in the hinge region connect the two heavy chains. The light chains are coupled to the heavy chains by additional disulfide bonds. Asn-linked carbohydrate moieties are attached at different positions in constant domains depending on the class of immunoglobulin. For IgG1 two disulfide bonds in the hinge region, between Cys235 and Cys238 pairs, unite the two heavy chains. The light chains are coupled to the heavy chains by two additional disulfide bonds, between Cys229s in the CH1 domains and Cys214s in the CL domains. Carbohydrate moieties are attached to Asn306 of each CH2, generating a pronounced bulge in the stem of the Y.

These features have profound functional consequences. The variable regions of both the heavy and light chains (VH) and (VL) lie at the "tips" of the Y, where they are positioned to react with antigen. This tip of the molecule is the side on which the N-terminus of the amino acid sequence is located. The stem of the Y projects in a way to efficiently mediate effector functions such as the activation of complement and interaction with Fc receptors, or ADCC and ADCP. Its CH2 and CH3 domains bulge to facilitate interaction with effector

proteins. The C-terminus of the amino acid sequence is located on the opposite side of the tip, which can be termed "bottom" of the Y.

Two types of light chain, termed lambda (λ) and kappa (κ), are found in antibodies. A given immunoglobulin either has K chains or λ chains, never one of each. No functional difference has been found between antibodies having λ or κ light chains.

Each domain in an antibody molecule has a similar structure of two beta sheets packed tightly against each other in a compressed antiparallel beta barrel. This conserved structure is termed the immunoglobulin fold. The immunoglobulin fold of constant domains contains a 3-stranded sheet packed against a 4-stranded sheet. The fold is stabilized by hydrogen bonding between the beta strands of each sheet, by hydrophobic bonding between residues of opposite sheets in the interior, and by a disulfide bond between the sheets. The 3-stranded sheet comprises strands C, F, and G, and the 4-stranded sheet has strands A, B, E, and D. The letters A through G denote the sequential positions of the beta strands along the amino acid sequence of the immunoglobulin fold.

The fold of variable domains has 9 beta strands arranged in two sheets of 4 and 5 strands. The 5-stranded sheet is structurally homologous to the 3-stranded sheet of constant domains, but contains the extra strands C' and C". The remainder of the strands (A, B, C, D, E, F, G) have the same topology and similar structure as their counterparts in constant domain immunoglobulin folds. A disulfide bond links strands B and F in opposite sheets, as in constant domains.

The variable domains of both light and heavy immunoglobulin chains contain three hypervariable loops, or complementarity-determining regions (CDRs). The three CDRs of a V domain (CDR1, CDR2, CDR3) cluster at one end of the beta barrel. The CDRs are loops that connect beta strands B-C, C'-C", and F-G of the immunoglobulin fold. The residues in the CDRs vary from one immunoglobulin molecule to the next, imparting antigen specificity to each antibody.

The VL and VH domains at the tips of antibody molecules are closely packed such that the 6 CDRs (3 on each domain) cooperate in constructing a surface (or cavity) for antigen-specific binding. The natural antigen binding site of an antibody thus is composed of the loops which connect strands B-C, C'-C", and F-G of the light chain variable domain and strands B-C, C'-C", and F-G of the heavy chain variable domain.

The loops which are not CDR-loops in a native immunoglobulin, or not part of the antigen-binding pocket as determined by the CDR loops and optionally adjacent loops within the CDR loop region, do not have antigen binding or epitope binding specificity, but contribute to the correct folding of the entire immunoglobulin molecule and/or its effector or other functions and are therefore called structural loops for the purpose of this invention.

Prior art documents show that the immunoglobulin-like scaffold has been employed so far for the purpose of manipulating the existing antigen binding site, thereby introducing novel binding properties. In most cases the CDR regions have been engineered for antigen binding, in other words, in the case of the immunoglobulin fold, only the natural antigen binding site has been modified in order to change its binding affinity or specificity. A vast body of literature exists which describes different formats of such manipulated immunoglobulins, frequently expressed in the form of single-chain Fv fragments (scFv) or Fab fragments,

either displayed on the surface of phage particles or solubly expressed in various prokaryotic or eukaryotic expression systems.

W006072620A1 describes a method of engineering an immunoglobulin which comprises a modification in a structural loop region to obtain new antigen binding sites. This method is broadly applicable to immunoglobulins and may be used to produce a series of immunoglobulins targeting a variety of antigens. A CH3 library has been shown to be useful for selecting specific binders to an antigen.

Although multivalent display of proteins on genetic packages has been described (such as direct phage cloning and display, bacterial display, yeast display), prior art refers to monomeric monovalent display of binding domains, in general. W09209690 describes phagemid particles displaying a single copy of a fusion protein on the surface of the particle. Thereby it was described to obtain high affinity binders from a library of phagemid particles, also called bacteriophages. Replicable expression vectors comprising genes encoding a binding polypeptide and a phage coat protein are provided so to form a gene fusion encoding a fusion protein, which is a chimeric protein of a phagemid particle, the phage coat protein and the binding polypeptide.

U.S. Pat. No. 5,223,409 generally describes the method of fusing a gene encoding a protein of interest to the N-terminal domain of the gene III coat protein of the filamentous phage M13. The gene fusion is mutated to form a library of structurally related fusion proteins that are expressed in low quantity on the surface of a phagemid particle. Biological selection and screening is employed to identify novel ligands useful as drug candidates.

However, there are some limitations in using such "fusion phage" or monovalent phage display and respective single fusion proteins. Many biologicals naturally occur in oligomeric form. For the purpose of the present invention oligomeric means dimeric, trimeric or even higher polymeric forms, up to 24 monomers.

The fusion phages according to the prior art are described to display monomeric fusion proteins, mainly because it was believed that binders of highest affinity could only be selected from a library if single fusion proteins are displayed by the phagemid particles. Native proteins are however often assembled as a dimer or even at a higher degree of oligomerization. To obtain dimeric display with a single fusion protein, some techniques have been developed that involve conditional stop codons located between the coat protein and the binding polypeptide (Dall'Acqua et al *The Journal of Immunology*, 2002, 169: 5171-5180). Thereby soluble monomers of the polypeptides in addition to those fused to the phage are expressed, thus enabling the formation of a dimer. However, such stop codons requires propagation in specific suppressor host cells that may translate a stop codon in an amino acid, to provide an appropriate amount of fusion proteins in addition to the soluble binding polypeptides. WO 03/029456 describes the use of multi-chain eukaryotic display vectors for the selection of immunoglobulin Fab fragments on the surface of yeast cells.

Prior art fusion proteins involve in some cases linker sequences to display larger binding polypeptides. Linker sequences of up to 24 amino acids are usually employed for standard purposes of displaying variable domains of an antibody. See for example, the display vector pCOMB3x (Hybrid. Hybridomics. 2003 April; 22(2):97-108. Development of functional human monoclonal single-chain variable fragment antibody against HIV-1 from human cervical B cells. Berry J D, Rutherford J, Silverman G J, Kaul R, Elia M, Gobuty S, Fuller R, Plummer F A, Barbas C F.)

It is an object of this invention to provide an effective method for the preparation of oligomers of modular antibody domains and to prepare such oligomers displayed on the surface of a replicable genetic package.

BRIEF DESCRIPTION OF THE INVENTION

The objects are solved by the subject matter of the present invention.

According to the invention a method of preparing a genetic package displaying oligomers of modular antibody domains binding to a target and to a scaffold ligand comprising

providing a genetic package, and
displaying at least two of the antibody modular domains by fusing to the outer surface of the package is covered.

The genetic package can be displayed in a mobilized or cellular system, wherein according to the invention the mobilized system can be selected from viruses, phages, phagemids, in-vitro display systems, mRNA systems and ribosomal display systems. Alternatively, a cellular system can be selected using yeast, mammalian cells, bacterial cells, bacterial spores or insect cells.

Oligomers are possibly formed by oligomerization motifs associated with the structure of said agents, such as leucine zipper, disulfide bonds, electrostatic or hydrophobic motifs.

According to one embodiment of the invention the oligomers can be dimers, trimers or tetramers, involving the same type of monomers (homomers) or different types (heteromers). The preferred method according to the invention is specifically useful for providing homomers, in particular homodimers of oligonucleotides on a genetic package, such as a phagemid particle or a phage or yeast.

The method according to the invention can be applied to oligomers which are polypeptides with a target binding site that directs towards the surface of the genetic package and is close to said surface that are biologicals, such as polypeptides.

Alternatively, the appropriate design of such a polypeptide is employed with a binding site that is closer to the surface of the genetic package than to the surrounding environment of the genetic package. This may be advantageous for a binding polypeptide with a potential binding site that is closer to the C-terminus than to the N-terminus of the polypeptide, in particular when the binding site is engineered in a C-terminal loop position. When the potential binding site is engineered at a position that is adjacent to the site where the genetic package particle is bound, for instance to a surface structure of a cell or a virus, it is advantageous to choose a stable construct with defined accessibility of the binding partner of said binding agent. C-terminal loop positions are, for instance, less accessible than N-terminal loop positions, when they are fused to the N-terminus of the protein 3 of a filamentous phage because they are exposed and in sterical proximity to the genetic package.

According to a preferred embodiment of the invention, however, a defined structure, such as a true oligomeric or dimeric fusion protein is provided to enable the efficient engineering of the potential binding site at an N-terminal loop position. One embodiment of the invention refers to polypeptides with at least two target binding sites, possibly engineered at the monomeric or the oligomeric target binding agent. In some cases the interaction between the monomeric structures enables additional variations of structures and thus additional potential binding sites.

5

According to a preferred embodiment of the invention the modular antibody is an antibody, Fc fragment, an antibody fragment with a CDR region and combinations thereof, possibly also comprising a fragment with a binding site at a structural loop position.

It can also be an antibody fragment with a CDR region like for example Fab, dAb, scFv, diabody, unibody, SMIPs, TANDABS, Fc fusion proteins and combinations thereof.

In case the genetic package is a filamentous bacteriophage, the preferred fusion structure employed with a bacteriophage involves at least part of an outer surface protein, such as p3, p6, or p8, however, p9 or p10 may also be used for the purpose of the invention.

In case the genetic package is yeast it is preferred that the oligomer is a fusion protein comprising one of the proteins of yeast cell surface receptors selected from the group consisting of alpha-agglutinin, a-agglutinin, Aga1p, Aga2p or FL01.

The appropriate genetic package is preferably provided in a particular form, and containing a vector encoding at least one of said fusion proteins. According to the invention there is, for example, provided a cassette vector, containing sequences encoding one or more than one fusion protein operatively linked to the genetic package. Thus, at least two of the chimeric fusion proteins are bound at the surface of the vector particle. The vector can be, for example, a phagemid.

An expression system for expressing oligomers of modular antibody domains bound at the surface of a genetic package produced according to the inventive method is also covered wherein the oligomers are encoded by a single gene and the fusion protein is displayed with at least two copies on the surface of the genetic package.

By using the method and means according to the invention it is possible to display oligomers of modular antibodies without the need of controlled expression of a soluble form of one of the oligomerization partners. The technique can be utilized for molecules such as antibody fragments, even those containing more than two immunoglobulin domains, e.g. at least four immunoglobulin domains. Thus difficult constructs involving stop codons, such as amber stop codons, can be avoided. There will be no need to get a mixture of fusion proteins and soluble monomers for the dimer display. Thus, a preferred technique of managing mixtures of a variety of binding agents can easily be employed, while the risk of insufficient matching of the monomers is reduced. It is also possible to avoid the suppressor strains as a host cell. Conventional host cells, with non-suppressor function, can serve in a standard way to propagate the oligomeric fusion proteins.

In another embodiment according to the invention, the fusion between the binding partner and the surface protein of the genetic package is such that no conditional stop codon (i.e. no amber, ochre, opal or other similar conditional stop codon) is present in between. In such a situation, upon infection with helper phage, the binding partner is present only as a fusion protein and not in soluble form. In order for the dimer (trimer or higher) to form on the surface of the genetic package, the linker which connects the binding partner to the genetic package needs to be of sufficient length and sufficient flexibility. Linkers that fulfil this requirement can be selected using the method described above.

In another embodiment according to the invention, a helper phage can be used that has specific properties that favour the display of more than one copy of the fusion protein on the surface of the genetic package. An example for such a helper phage is the so-called hyperphage (Rondot

6

S, Koch J, Breitling F, Dubel S. A helper phage to improve single-chain antibody presentation in phage display. *Nat Biotechnol.* 2001 January; 19(1):75-78), which itself is devoid of p3, and therefore depends totally on the p3 fusion protein provided by the phagemid in order to be infective. Using such helper phage leads preferentially to phage particles that carry more than one copies of the fusion protein on their surface, and thus favors the formation of dimers of the binding protein on the surface of the genetic package.

According to a preferred embodiment of the invention each of the fusion protein monomers are prepared in the context of an oligomer, so that the portion of soluble binding agents is less than 20%, more preferably less than 10%, most preferably less than 1%.

According to the inventive method a genetic package can be prepared preferably displaying at least two fusion proteins. In a specific embodiment each monomer of the inventive oligomer is bound to the outer surface of the genetic package. The invention also provides bivalent phage displaying two fusion proteins from oligomers each containing an oligomer that dimerizes upon expression.

A library of genetic packages is also claimed, which can exemplarily comprise at least 10 variant genetic packages, wherein said variant genetic packages can display heteromers of modular antibody domains. The heteromer can be based on a scaffold comprising a target binding site. According to a specific embodiment of the invention the target binding site and the scaffold binding site can be similar or identical. A scaffold ligand according to the invention can also be a CDR target. For example, the scaffold can be a parent Fab and at least 20%, preferably at least 30%, more preferred at least 40% of the parent Fab variants are binding to the CDR-target of said parent Fab.

The inventive library can also contain variants of the oligomer produced according to the invention having differences in the amino acid sequence.

This can be provided by modifying the amino acid sequences by at least one insertion or substitution to introduce at least one foreign amino acid, and by a deletion. Foreign amino acids can be introduced by a randomization technique. The foreign amino acids can also be selected from a specific group of amino acids to obtain a library enriched with specific amino acids at the randomized positions. When the foreign amino acid is selected from a specific group of amino acids, such as amino acids with specific polarity, or hydrophobicity, a library enriched in the specific group of amino acids at the randomized positions can be obtained according to the invention. Such libraries are also called "focused" libraries.

According to the invention there is further provided a method of selecting a linker for binding a polypeptide to the outer surface of a genetic package, comprising

- providing a library of genetic packages containing a variety of linkers to connect a first polypeptide to the genetic package,
- determining the member of the library containing a linker that does not significantly interfere with the function of said first polypeptide, and
- selecting said linker for connecting a second polypeptide to said genetic package.

Thereby suitable linkers to be used for fusion proteins of the same type can be obtained. By the same type of fusion proteins those are meant that link the same formats of genetic packages and binding agent to each other.

Such a method according to the invention can further be used to prepare enriched libraries containing a preselected group of linker variants that fulfil at least one selection

criterion, such as flexibility and sterical accessibility of the binding site to the binding partner. This can be determined by measuring the binding properties of a well-known binding agent in the presence of a variety of linker sequences. The enriched library may thus be further selected for another criterion, such as protease resistance, which is e.g. important for the stability in a medium containing bacterial proteases.

The selected specific linker sequence can then be used to prepare libraries of fusion proteins of the same format, however, with variants of the binding agents, thus enabling the identification, selection and preparation of those agents with the best binding properties in predetermined test systems.

In a preferred embodiment according to the invention such a linker is at least 20 amino acids long, preferably at least 25 amino acid residues, more preferably at least 30 amino acid residues, up to 50 amino acid residues. Especially when amino acids such as Gly, Ser or Ala are involved, which are responsible for the flexibility of a linker, such a linker is advantageously used for the display of a potential binding site, which is close to the surface of the genetic package and which potential binding site may not be able to bind to its partner for sterical reasons.

The linker between the protein to be displayed and the anchor protein of the genetic package (in case of filamentous phage e.g. p3, p8, pX, pIX, pVII) is especially important if the potential binding site of the displayed molecule is in spatial vicinity of the phage particle. In antibody libraries utilizing variable domains and antigen binding sites formed by CDR-loops and display of the library members as amino-terminal fusion to p3 the potential antigen binding site is directed away from the phage particle. Therefore, the linker structure between library members and the phage coat protein is not important. Engineering the bottom loops of immunoglobulin domains and performing phage display may however be an inefficient process and decrease yields of antigen binding clones or even preclude it. Varying the linker between a library member protein and its fusion partner on the surface can solve or may at least reduce this problem.

In order to select for optimal linker sequences (in terms of length and flexibility as well as stability) a library of linkers can be prepared in which the anchor protein at the surface of the genetic replicable package is fused to a known binding protein which is for sterical reasons notoriously difficult to select for.

This library of sequences can be varied in length and amino acid content.

Selection methods of the linker library for optimal linkers depend on the application but basically it should be for selecting all properties one wishes to have in a certain methodology. Enrichment against an antigen that is difficult to select may yield linker sequences which allow library members a good access to the antigen. Incubation in protease solutions or under other harsh conditions or frequent passaging through host cells under proteolytic conditions (e.g. old microbial cultures) may be an appropriate selection for stable display linkers.

A library of linkers may be produced by any well known library technology. Synthetic linker sequence lengths may vary between 10-500 amino acids. Alternatively, linker can be complete proteins known to be of flexible nature.

The invention also provides a method of producing an oligomer of modular antibody domains binding to a target comprising the steps of:

providing a library of oligomers of modular antibody domains produced according to the inventive method as described

contacting said library with said target in the presence of a scaffold ligand,

selecting a library member binding to said target in the presence of a scaffold ligand, and

manufacturing a preparation of the functional oligomer.

The scaffold ligand can be selected from the group consisting of an effector molecule, FcRn, serum albumin, Protein A, Protein G, Protein L or a CDR target. As an example, the effector molecule can be selected from the group consisting of CD64, CD16, CD32, Fc receptors.

The oligomers can be dimers selected from the group of VH/VL, CH1/CL, CH2/CH2, CH3/CH3, Fc and Fab, or single chains thereof.

The method according to the invention can provide a library containing at least 10^2 independent clones expressing functional oligomers of modular antibody domains or variants thereof. The library member can then be selected according to the requested binding affinity, preferably it has a target binding affinity of $K_d < 10^{-8} M$. According to the invention it is also provided a pool of preselected independent clones, which is e.g. affinity matured, which pool comprises preferably at least 10, more preferably at least 100, more preferably at least 1000, more preferably at least 10000, even more than 100000 independent clones. Those libraries, which contain the preselected pools, are preferred sources to select the high affinity modular antibodies according to the invention.

Preferably the library is a yeast library and the yeast host cell exhibits at the surface of the cell the oligomers with the biological activity. The yeast host cell is preferably selected from the genera *Saccharomyces*, *Pichia*, *Hansenula*, *Schizosaccharomyces*, *Kluyveromyces*, *Yarrowia* and *Candida*. Most preferred, the host cell is *Saccharomyces cerevisiae*.

According to a specific embodiment of the invention, the target is a receptor of the erbB class. In this case, by the method of the invention an immunoglobulin can be obtained that binds to a receptor of the erbB class.

The invention further provides a high quality library containing at least 10^6 independent clones of functional dimers of modular antibody domains or variants thereof, or the pools of optimized or preselected clones, e.g. the affinity matured clones, which pools are containing at least 10 independent clones that are binding to a target and to a scaffold ligand. The target can be a ligand binding to a parent molecule subject to amino acid variation. The parent molecule can be a functional Fc or a functional Fab, or part thereof.

According to a specific embodiment of the invention the parent molecules can be varied by random or site specific mutagenesis.

The library can contain functional dimers of modular antibody domains that are binding to a target and to a scaffold ligand, and at least 20%, preferably at least 30%, more preferred at least 40% of the functional dimers are binding to CD64. This is particularly preferred with a modular antibody that contains CH2 domains, such as an Fc scaffold.

Alternatively, the library can contain functional dimers of modular antibody domains that are binding to a target and to a scaffold ligand, and at least 20%, preferably at least 30%, more preferred at least 40% of the functional dimers are binding to protein A. This is particularly preferred with a modular antibody that contains CH2 and CH3 domains, such as an Fc scaffold.

Alternatively, the library can contain functional dimers of modular antibody domains that are binding to a target and to a scaffold ligand, and at least 20%, preferably at least 30%, more preferred at least 40% of the functional dimers are binding to the same CDR target. This is particularly preferred with modular antibodies containing a variable region, such as an Fab scaffold with specificity to a single CDR target.

FIGURES

FIG. 1:

Schematic presentation of the PCRs used for production of the fragments used for assembly of the library FcabO1. PCR primers are indicated by arrows with their respective 5'-3' orientation, and vertical lines indicate the approximate positions of the introduced restriction sites which were used for assembly of the mutated gene. The restriction sites are contained on the primers for ligations of the PCR fragments.

FIG. 2:

Amino acid sequence and secondary structure of a CH3 domain (IMGT numbering). The randomization scheme is provided for the libraries FcabO1 to FcabO6.

Randomized positions in the AB and EF loop are marked with a circle. X stands for all 20 amino acids (encoded by NNB), z only for Ala, Asp, Ser, Tyr (encoded by KMT; focused library).

DETAILED DESCRIPTION OF THE INVENTION

The oligomers of the modular antibody domains according to the present invention will be useful as stand-alone molecules, as well as fusion proteins or derivatives, most typically fused before or after modification in such a way as to be part of larger structures, e.g. of complete antibody molecules, or parts thereof. Immunoglobulins or fusion proteins as produced according to the invention thus also comprise Fc fragments, Fab fragments, Fv fragments, single domain antibodies, single chain antibodies, in particular single-chain Fv fragments, bior multispecific scFv, diabodies, unibodies, multibodies, multivalent or multimers of immunoglobulin domains and others. It will be possible to use the engineered proteins to produce molecules which are monospecific, bispecific, trispecific, and may even carry more specificities. By the invention it is possible to control and preselect the valency of binding at the same time according to the requirements of the planned use of such molecules.

Specific terms as used throughout the specification have the following meaning.

The term "immunoglobulin" as used according to the present invention is defined as polypeptides or proteins that may exhibit mono- or bi- or multi-specific, or mono-, bi- or multivalent binding properties, preferably at least two, more preferred at least three specific binding sites for epitopes of e.g. antigens, effector molecules or proteins either of pathogen origin or of human structure, like self-antigens including cell-associated or serum proteins. The term immunoglobulin as used according to the invention also includes functional fragments of an antibody, such as Fc, Fab, scFv, single chain dimers of CH1/CL domains, Fv, dimers like VH/VL, CH1/CL, CH2/CH2, CH3/CH3, or other derivatives or combinations of the immunoglobulins, like single chains of pairs of immunoglobulin domains. The definition further includes domains of the heavy and light chains of the variable region (such as dAb, Fd, V1, Vk, Vh, VHH) and the constant region

or individual domains of an intact antibody such as CH1, CH2, CH3, CH4, C1 and Ck, as well as mini-domains consisting of at least two beta-strands of an immunoglobulin domain connected by a structural loop.

"Modular antibodies" as used according to the invention are defined as antigen-binding molecules, like human antibodies, composed of at least one polypeptide module or protein domain, preferably in the natural form. The term "modular antibodies" includes antigen-binding molecules that are either immunoglobulins, immunoglobulin-like proteins, or other proteins exhibiting modular formats and antigen-binding properties similar to immunoglobulins or antibodies, which can be used as antigen-binding scaffolds, preferably based on human proteins.

The term "immunoglobulin-like molecule" as used according to the invention refers to any antigen-binding protein, in particular to a human protein, which has a domain structure that can be built in a modular way. Immunoglobulin-like molecules as preferably used for the present invention are T-cell receptors (TCR), fibronectin, transferrin, CTLA-4, single-chain antigen receptors, e.g. those related to T-cell receptors and antibodies, antibody mimetics, adnectins, anticalins, phylomers, repeat proteins such as ankyrin repeats, avimers, Versabodies™, scorpion toxin based molecules, and other non-antibody protein scaffolds with antigen binding properties.

Ankyrin repeat (AR), armadillo repeat (ARM), leucine-rich repeat (LRR) and tetratricopeptide repeat (TPR) proteins are the most prominent members of the protein class of repeat proteins. Repeat proteins are composed of homologous structural units (repeats) that stack to form elongated domains. The binding interaction is usually mediated by several adjacent repeats, leading to large target interaction surfaces.

Avimers contain A-domains as strings of multiple domains in several cell-surface receptors. Domains of this family bind naturally over 100 different known targets, including small molecules, proteins and viruses. Truncation analysis has shown that a target is typically contacted by multiple A-domains with each domain binding independently to a unique epitope. The avidity generated by combining multiple binding domains is a powerful approach to increase affinity and specificity, which these receptors have exploited during evolution.

Anticalins are engineered human proteins derived from the lipocalin scaffold with defined binding properties typical for humanized antibodies. Lipocalins comprise 160-180 amino acids and form conical beta-barrel proteins with a ligand-binding pocket surrounded by four loops. Small hydrophobic compounds are the natural ligands of lipocalins, and different lipocalin variants with new compound specificities (also termed 'anticalins') could be isolated after randomizing residues in this binding pocket.

Single chain or single domain antigen receptors contain a single variable domain and are 20% smaller than camelid single domain antibodies.

Phylomers are peptides derived from biodiverse natural protein fragments.

It is understood that the term "modular antibody", "immunoglobulin", "immunoglobulin-like proteins" includes a derivative thereof as well. A derivative is any combination of one or more modular antibodies of the invention and or a fusion protein in which any domain or minidomain of the modular antibody of the invention may be fused at any position of one or more other proteins (such as other modular antibodies, immunoglobulins, ligands, scaffold proteins, enzymes, toxins and the like). A derivative of the

modular antibody of the invention may also be obtained by association or binding to other substances by various chemical techniques such as covalent coupling, electrostatic interaction, di-sulphide bonding etc. The other substances bound to the immunoglobulins may be lipids, carbohydrates, nucleic acids, organic and inorganic molecules or any combination thereof (e.g. PEG, prodrugs or drugs). A derivative would also comprise an antibody with the same amino acid sequence but made completely or partly from non-natural or chemically modified amino acids.

A “structural loop” or “non-CDR-loop” according to the present invention is to be understood in the following manner: modular antibodies, immunoglobulins or immunoglobulin-like substances are made of domains with a so called immunoglobulin fold. In essence, strands of antiparallel beta sheets are connected by loops to form a compressed antiparallel beta barrel. In the variable region, some of the loops of the domains contribute essentially to the specificity of the antibody, i.e. the binding to an antigen by the natural binding site of an antibody. These loops are called CDR-loops. The CDR loops are located within the CDR loop region, which may in some cases include also the variable framework region (called “VFR”) that is adjacent to the CDR loops. It is known that VFRs may contribute to the antigen binding pocket of an antibody, which generally is mainly determined by the CDR loops. Thus, those VFRs are considered as part of the CDR loop region, and would not be appropriately used for engineering new antigen binding sites. Contrary to those VFRs within the CDR loop region or located proximal to the CDR loops, other VFRs of variable domains would be particularly suitable for use according to the invention. Those are the structural loops of the VFRs located opposite to the CDR loop region, or at the C-terminal side of a variable immunoglobulin domain.

The term “antigen” or “target” as used according to the present invention shall in particular include all antigens and target molecules capable of being recognised by a binding site of a modular antibody. Specifically preferred antigens as targeted by the receptor molecule according to the invention are those antigens or molecules, which have already been proven to be or are capable of being immunologically or therapeutically relevant, especially those, for which a clinical efficacy has been tested.

The term “target” or “antigen” as used herein shall comprise molecules selected from the group consisting of allergens, tumor associated antigens, self antigens including cell surface receptors, enzymes, Fc-receptors, FcRn, RSA, IgG, interleukins or cytokines, proteins of the complement system, transport proteins, serum molecules, bacterial antigens, fungal antigens, protozoan antigens and viral antigens, also molecules responsible for transmissible spongiform encephalitis (TSE), such as prions, infective or not, and markers or molecules that relate to inflammatory conditions, such as pro-inflammatory factors, multiple sclerosis or alzheimer disease, or else haptens.

The term “cell surface antigens” shall include all antigens capable of being recognised by an antibody structure on the surface of a cell, and fragments of such molecules. Preferred cell surface antigens are those antigens, which have already been proven to be or which are capable of being immunologically or therapeutically relevant, especially those, for which a preclinical or clinical efficacy has been tested. Those cell surface molecules are specifically relevant for the purpose of the present invention, which mediate cell killing activity. Upon binding of the immunoglobulin according to the invention to preferably at least two of those cell surface

molecules the immune system provides for cytolysis or cell death, thus a potent means for attacking human cells may be provided.

The antigen is either recognized as a whole target molecule or as a fragment of such molecule, especially substructures of targets, generally referred to as epitopes.

Substructures of antigens are generally referred to as “epitopes” (e.g. B-cell epitopes, T-cell epitopes), as long as they are immunologically relevant, i.e. are also recognisable by natural or monoclonal antibodies. The term “epitope” as used herein according to the present invention shall mean a molecular structure which may completely make up a specific binding partner or be part of a specific binding partner to a binding site of a modular antibody or an immunoglobulin of the present invention. The term epitope may also refer to haptens. Chemically, an epitope may either be composed of a carbohydrate, a peptide, a fatty acid, an organic, biochemical or inorganic substance or derivatives thereof and any combinations thereof. If an epitope is a polypeptide, it will usually include at least 3 amino acids, preferably 8 to 50 amino acids, and more preferably between about 10-20 amino acids in the peptide. There is no critical upper limit to the length of the peptide, which could comprise nearly the full length of a polypeptide sequence of a protein. Epitopes can be either linear or conformational epitopes. A linear epitope is comprised of a single segment of a primary sequence of a polypeptide chain. Linear epitopes can be contiguous or overlapping. Conformational epitopes are comprised of amino acids brought together by folding of the polypeptide to form a tertiary structure and the amino acids are not necessarily adjacent to one another in the linear sequence. Specifically, epitopes are at least part of diagnostically relevant molecules, i.e. the absence or presence of an epitope in a sample is qualitatively or quantitatively correlated to either a disease or to the health status of a patient or to a process status in manufacturing or to environmental and food status. Epitopes may also be at least part of therapeutically relevant molecules, i.e. molecules which can be targeted by the specific binding domain which changes the course of the disease.

As used herein, the term “specifically binds” or “specific binding” refers to a binding reaction which is determinative of the cognate ligand of interest in a heterogeneous population of molecules. Thus, under designated conditions (e.g. immunoassay conditions), the modular antibody binds to its particular target and does not bind in a significant amount to other molecules present in a sample. The specific binding means that binding is selective in terms of target identity, high, medium or low binding affinity or avidity, as selected. Selective binding is usually achieved if the binding constant or binding dynamics is at least 10 fold different, preferably the difference is at least 100 fold, and more preferred a least 1000 fold.

The term “expression system” refers to nucleic acid molecules containing a desired coding sequence and control sequences in operable linkage, so that hosts transformed or transfected with these sequences are capable of producing the encoded proteins. In order to effect transformation, the expression system may be included on a vector; however, the relevant DNA may also be integrated into the host chromosome.

Alternatively, an expression system can be used for in vitro transcription/translation. The expression system preferably is employing a host cell that is either a eukaryotic or prokaryotic host cell, preferably a mammalian or yeast host cell, as well as a bacterial host cell.

The preferred expression system for the fusion proteins is a non-suppressor host cell, which would be sensitive to a stop codon, such as an amber stop codon, and would thus stop translation thereafter. In the absence of such a stop codon such non-suppressor host cells, preferably *E. coli*, are preferably used. In the presence of such a stop codon suppressor host cells would be used.

All numbering of the amino acid sequences of the immunoglobulins is according to the IMGT numbering scheme (IMGT, the international ImmunoGeneTics, Lefranc et al., 1999, Nucleic Acids Res. 27: 209-212).

For the purposes of this invention, the term "binding agent" or "ligand" refers to a member of a binding pair, in particular binding polypeptides having the potential of serving as a binding domain for a binding partner. Examples of binding partners include pairs of binding agents with functional interactions, such as receptor binding to ligands, antibody binding to antigen or receptors, a drug binding to a target, and enzyme binding to a substrate.

The term "fusion protein" or "chimeric fusion protein" as used for the purpose of the invention shall mean the molecule composed of a genetic package, at least part of an outer surface structure, such as a coat protein or part thereof, optionally a linker sequence, and a binding agent. The fusion protein is encoded by a vector with the gene of the binding agent and information to display a copy of the binding agent at the surface of the genetic package.

The term "cytotoxic activity" as used for the purpose of the invention shall mean the activity on effector cells resulting in activation of cytotoxic T-cells or on cells, which mediate antibody-dependent cell cytotoxicity (ADCC), complement dependent cytotoxicity (CDC) and/or antibody-dependent cellular phagocytosis (ADCP). Modular antibodies according to the invention thus kill antibody-coated target cells, which they optionally bind with their Fc receptors.

"Scaffold" shall mean a temporary framework either natural or artificial used to support the molecular structure of a polypeptide in the construction of variants or a repertoire of the polypeptide. It is usually a modular system of polypeptide domains that maintains the tertiary structure or the function of the parent molecule. Exemplary scaffolds are modular antibodies, which may be mutagenized to produce variants within said scaffold, to obtain a library.

The term "scaffold ligand" as used for the purpose of the invention shall mean a ligand that binds to a scaffold or the backbone of modular antibodies, thus determining the molecular structure or primary function and specificity of said modular antibody. In preferred cases the scaffold ligand is a functional ligand, mediating a biological function upon binding, like an effector ligand. In an alternative embodiment the scaffold ligand is a functional ligand, which is a specific target bound by the CDR region, non-structural loop region or structural loop region. The same scaffold ligand can bind many variants of a modular antibody regardless of their target specificities. In general, the presence of scaffold ligand binding site indicates that the variant is expressed and folded correctly. Thus, binding of the scaffold ligand to its binding site provides a method for preselecting functional polypeptides from a repertoire of polypeptides. Designing variants of modular antibodies that keep the binding property to a scaffold ligand avoids the preparation of variants that are non-functional, for example as a result of the introduction of mutations, folding mutants or expression mutants which would be or are incapable of binding to substantially any target or effector ligand. Such non-functional mutants sometimes are generated by the normal

randomisation and variation procedures employed in the construction of polypeptide repertoires. Providing functional mutants that bind to a scaffold ligand permits the person skilled in the art to prepare a library of modular antibodies which is enriched in functional, well folded and highly expressed library members.

The term "effector ligand" as used for the purpose of the invention shall mean a ligand mediating effector functions, like an effector molecule. Exemplary effector ligands are Pc receptors or Pc receptor-like molecules interfering with immunoglobulins. An Pc receptor is a protein found on the surface of certain cells—including natural killer cells, macrophages, neutrophils, and mast cells—that contribute to the protective functions of the immune system. Its name is derived from its binding specificity for a part of an antibody known as the Fc (Fragment-crystallizable) region. Pc receptors bind to antibodies that are attached to infected cells or invading pathogens. Their activity stimulates phagocytic or cytotoxic cells to destroy microbes, or infected cells by antibody-mediated cellular phagocytosis (ADCP) or antibody-dependent cell-mediated cytotoxicity (ADCC). There are several different types of Pc receptors, which are classified based on the type of antibody that they recognize; those that bind the most common class of antibody, IgG, are called Pc-gamma receptors (FcγR), those that bind IgA are called Pc-alpha receptors (FcαR) and those that bind IgE are called Pc-epsilon receptors (FcεR). Equivalent to an effector ligand and thus incorporated into the definition is any surrogate ligand that recognizes the same or similar binding site within the modular antibody, such as Protein A.

All FcγRs belong to the immunoglobulin superfamily and are the most important Fc receptors for inducing phagocytosis of opsonized (coated) microbes. This family includes several members that differ in their antibody affinities due to their different molecular structure: FcγRI (CD64), FcγRIIA (CD32a), FcγRIIB (CD32b), FcγRIIIA (CD16a), FcγRIIIB (CD16b). For instance, FcγRI binds to IgG more strongly than FcγRII and FcγRIII, and has an extracellular portion composed of three immunoglobulin (Ig)-like domains, one more domain than FcγRII and FcγRIII. These properties allow activation of FcγRI by a sole IgG molecule (or monomer), while the latter two Fcγ receptors must bind multiple IgG molecules within an immune complex to be activated.

Another FcR is expressed on multiple cell types and is similar in structure to MHC class I. This receptor also binds IgG and is involved in preservation of this antibody in order to increase its biological half-life in vivo. However, since this Fc receptor is also involved in transferring IgG from a mother either via the placenta to her fetus or in milk to her suckling infant, it is called the neonatal Fc receptor (FcRn). Recently this receptor has been implicated in being involved in homeostasis of IgG serum levels.

Antibody-Dependent Cell-Mediated Cytotoxicity (ADCC) is a mechanism of cell-mediated immunity whereby an effector cell of the immune system actively lyses a target cell that has been bound by specific antibodies. It is one of the mechanisms through which antibodies, as part of the humoral immune response, can act to limit and contain infection. Classical ADCC is mediated by natural killer (NK) cells; monocytes and eosinophils can also mediate ADCC. For example Eosinophils can kill certain parasitic worms known as helminths through ADCC. ADCC is part of the adaptive immune response due to its dependence on a prior antibody response.

The term "foreign" in the context of amino acids shall mean the newly introduced amino acids being naturally

occurring, but foreign to the site of modification, or substitutes of naturally occurring amino acids. "Foreign" with reference to an antigen binding sites means that the antigen binding site is not naturally formed by the specific binding region of the agent, and a foreign binding partner, but not the natural binding partner of the agent, is bound by the newly engineered binding site.

The term "variable binding region" sometimes called "CDR region" as used herein refers to molecules with varying structures capable of binding interactions with antigens. Those molecules can be used as such or integrated within a larger protein, thus forming a specific region of such protein with binding function. The varying structures can be derived from natural repertoires of binding proteins such as immunoglobulins or phylomers or synthetic diversity, including repeat-proteins, avimers and anticalins. The varying structures can as well be produced by randomization techniques, in particular those described herein. These include mutagenized CDR or non-CDR regions, loop regions of immunoglobulin variable domains or constant domains.

Modified binding agents with different modifications at specific sites are referred to as "variants". Variants of a scaffold are preferably grouped to form libraries of binding agents, which can be used for selecting members of the library with predetermined functions. In accordance therewith, a loop region of a binding agent comprising positions within one or more loops potentially contributing to a binding site, is preferably mutated or modified to produce libraries, preferably by random, semi-random or, in particular, by site-directed random mutagenesis methods, in particular to delete, exchange or introduce randomly generated inserts into loops, preferably into structural loops. Alternatively preferred is the use of combinatorial approaches. Any of the known mutagenesis methods may be employed, among them cassette mutagenesis. These methods may be used to make amino acid modifications at desired positions of the immunoglobulin of the present invention. In some cases positions are chosen randomly, e.g. with either any of the possible amino acids or a selection of preferred amino acids to randomize loop sequences, or amino acid changes are made using simplistic rules. For example all residues may be mutated preferably to specific amino acids, such as alanine, referred to as amino acid or alanine scanning. Such methods may be coupled with more sophisticated engineering approaches that employ selection methods to screen higher levels of sequence diversity.

The preferred cytotoxic modular antibody according to the invention with a molecular weight of less than 60 kD or up to 60 kD has a small size as compared to full length antibodies. The preferred size is up to 55 kD. Modular antibody single domains usually have a molecular size of 10-15 kD, thus a molecule based on 4 modular antibody domains would have a molecular size of 40-60 kD, depending on the glycosylation or any additional conjugation of pharmacologically active substances, like toxins or peptides.

The preferred format is an oligomer, composed of modular antibody domains, preferably up to 4 domains, more preferred 3 domains, and even more preferred made up of 2 domains. Formats based on the combination of 5 modular antibody domains or more are commonly thought not to exert the specific advantages of small sized antibody fragments, which are e.g. ease of expression in various expression systems and tissue penetration.

It is feasible to provide the preferred modular antibody of the invention as a single domain antibody. However, antibody domains tend to dimerize upon expression, either as a

homodimer, like an Fc, or a heterodimer, like an Fab. The dimeric structure is thus considered as a basis for the preferred stable molecule. The preferred dimers of immunoglobulin domains are selected from the group consisting of single domain dimers, like VH/VL, CH1/CL (kappa or lambda), CH2/CH2 and CH3/CH3. Dimers or oligomers of modular antibody domains can also be provided as single chain or two chain molecules, in particular those linking the C-terminus of one domain to the N-terminus of another.

Binding partners are agents that specifically bind to one another, usually through non-covalent interactions. Examples of binding partners include pairs of binding agents with functional interactions, such as receptor binding to ligands, antibody binding to antigen, a drug binding to a target, and enzyme binding to a substrate. Binding partners have found use in many therapeutic, diagnostic, analytical and industrial applications. Most prominent binding partners, also called binding pairs, are antibodies or immunoglobulins, fragments or derivatives thereof. In most cases the binding of such binding agents is required to mediate a biological effect or a function, a "functional interaction".

According to a specific embodiment of the present invention the binding agent is an immunoglobulin of human or murine origin, and may be employed for various purposes, in particular in pharmaceutical compositions. Of course, the modified immunoglobulin may also be a humanized or chimeric immunoglobulin.

The binding agent which is a human immunoglobulin is preferably selected or derived from the group consisting of IgA1, IgA2, IgD, IgE, IgG1, IgG2, IgG3, IgG4 and IgM. The murine immunoglobulin binding agent is preferably selected or derived from the group consisting of IgA, IgD, IgE, IgG1, IgG2A, IgG2B, IgG2C, IgG3 and IgM.

Such a binding agent comprises preferably a heavy and/or light chain or a part thereof. A modified immunoglobulin according to the invention may comprise a heavy and/or light chain, at least one variable and/or constant domain, or a part thereof including a minidomain.

A constant domain is an immunoglobulin fold unit of the constant part of an immunoglobulin molecule, also referred to as a domain of the constant region (e.g. CH1, CH2, CH3, CH4, Ck, Cl).

A variable domain is an immunoglobulin fold unit of the variable part of an immunoglobulin, also referred to as a domain of the variable region (e.g. Vh, Vk, V1, Vd).

An exemplary modular antibody according to the invention consists of a constant domain selected from the group consisting of CH1, CH2, CH3, CH4, Igk-C, Igl-C, combinations, derivatives or a part thereof including a minidomain, with at least one loop region, and is characterised in that said at least one loop region comprises at least one amino acid modification forming at least one modified loop region, wherein said at least one modified loop region binds specifically to at least one epitope of an antigen.

Another modular antibody according to the invention can consist of a variable domain of a heavy or light chain, combinations, derivatives or a part thereof including a minidomain, with at least one loop region, and is characterised in that said at least one loop region comprises at least one amino acid modification forming at least one modified loop region, wherein said at least one modified loop region binds specifically to at least one epitope of an antigen.

The modular antibody according to the present invention may comprise one or more domains (e.g. at least two, three, four, five, six, ten domains). If more than one domain is present in the modular antibody these domains may be of the same type or of varying types (e.g. CH1-CH1-CH2, CH3-

CH3, (CH2)2-(CH3)2, with or without the hinge region). Of course also the order of the single domains may be of any kind (e.g. CH1-CH3-CH2, CH4-CH1-CH3-CH2).

The invention preferably refers to parts of antibodies, such as IgG, IgA, IgM, IgD, IgE and the like. The modular antibodies of the invention may also be a functional antibody fragment such as Fab, Fab2, scFv, Fv, Fc, FcabTM, an antigen-binding Fc, or parts thereof, or other derivatives or combinations of the immunoglobulins such as minibodies, domains of the heavy and light chains of the variable region (such as dAb, Fd (binding site made up of one or more single domains), VL, including Vlambda (V1) and Vkappa (Vk), VH, VHH) as well as mini-domains consisting of two beta-strands of an immunoglobulin domain connected by at least two structural loops, as isolated domains or in the context of naturally associated molecules. A particular embodiment of the present invention refers to the Fc fragment of an antibody molecule, either as antigen-binding Fc fragment (FcabTM) through modifications of the amino acid sequence or as conjugates or fusions to receptors, peptides or other antigen-binding modules, such as scFv.

The modular antibodies can be used as isolated polypeptides or as combination molecules, e.g. through recombination, fusion or conjugation techniques, with other peptides or polypeptides. The peptides are preferably homologous to immunoglobulin domain sequences, and are preferably at least 5 amino acids long, more preferably at least 10 or even at least 50 or 100 amino acids long, and constitute at least partially the loop region of the immunoglobulin domain. The preferred binding characteristics relate to predefined epitope binding, affinity and avidity.

The modular antibody according to the invention is possibly further combined with one or more modified modular antibodies or with unmodified modular antibodies, or parts thereof, to obtain a combination modular antibody. Combinations are preferably obtained by recombination techniques, but also by binding through adsorption, electrostatic interactions or the like, or else through conjugation or chemical binding with or without a linker. The preferred linker sequence is either a natural linker sequence or a functionally suitable artificial sequence.

In general the modular antibody according to the invention may be used as a building block to molecularly combine other modular antibodies or biologically active substances or molecules. It is preferred to molecularly combine at least one antibody binding to the specific partner via the variable or non-variable sequences, like structural loops, with at least one other binding molecule which can be an antibody, antibody fragment, a soluble receptor, a ligand or another antibody domain, or a binding moiety thereof. Other combinations refer to proteinaceous molecules, nucleic acids, lipids, organic molecules and carbohydrates.

The engineered molecules according to the present invention will be useful as stand-alone proteins as well as fusion proteins or derivatives, most typically fused in such a way as to be part of larger antibody structures or complete antibody molecules, or parts or fragments thereof, such as Fab fragments, Fc fragments, Fv fragments and others. It will be possible to use the engineered proteins to produce molecules which are monospecific, bispecific, trispecific, and maybe even carry more specificities at the same time, and it will be possible at the same time to control and preselect the valency of binding at the same time according to the requirements of the planned use of such molecules.

According to the present invention, the modular antibody optionally exerts one or more binding regions to antigens, including the binding site binding specifically to the cell

surface target and the binding sites mediating effector function. Antigen binding sites to one or more antigens may be presented by the CDR-region or any other natural receptor binding structure, or be introduced into a structural loop region of an antibody domain, either of a variable or constant domain structure. The antigens as used for testing the binding properties of the binding sites may be naturally occurring molecules or chemically synthesized molecules or recombinant molecules, either in solution or in suspension, e.g. located on or in particles such as solid phases, on or in cells or on viral surfaces. It is preferred that the binding of an immunoglobulin to an antigen is determined when the antigen is still adhered or bound to molecules and structures in the natural context. Thereby it is possible to identify and obtain those modified immunoglobulins that are best suitable for the purpose of diagnostic or therapeutic use.

Modular antibody or immunoglobulin domains may be modified according to the present invention (as used herein the terms immunoglobulin and antibody are interchangeable) which modifications are preferably effected in immunoglobulin domains or parts thereof that contain a loop, either a CDR-loop or a non-CDR loop, structural loops being the preferred sites of modifications or mutagenesis. In some cases it is preferable to use a defined modified structural loop or a structural loop region, or parts thereof, as isolated molecules for binding or combination purposes.

It is particularly preferred that the modular antibody according to the invention is binding to said cell surface target through at least part of a structural loop and/or CDR loop.

In an alternate embodiment it is preferred that the modular antibody according to the invention is binding to said effector ligand, or a surrogate ligand for such an effector ligand, like protein A, through at least part of a structural loop and/or CDR loop, thus mediating the effector function.

In a preferred embodiment the binding agent is binding with its native or modified binding structure or newly formed binding site, specifically to at least two such epitopes that are identical or differ from each other, either of the same antigen or of different antigens.

In a preferred domain structure of a binding agent it is preferred to modify at least one loop region resulting in a substitution, deletion and/or insertion of one or more nucleotides or amino acids, preferably a point mutation, or even the exchange of whole loops, more preferred the change of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15, up to 30 amino acids. Thereby the modified sequence comprises amino acids not included in the conserved regions of the loops, the newly introduced amino acids being naturally occurring, but foreign to the site of modification, or substitutes of naturally occurring amino acids.

However, the maximum number of amino acids inserted into a loop region of a binding agent preferably may not exceed the number of 30, preferably 25, more preferably 20 amino acids at a maximum. The substitution and the insertion of the amino acids occurs preferably randomly or semi-randomly using all possible amino acids or a selection of preferred amino acids for randomization purposes, by methods known in the art and as disclosed in the present patent application.

The site of modification may be at a specific single loop or a loop region, in particular a structural loop or a structural loop region. A loop region usually is composed of at least one, preferably at least two, preferably at least 3 or at least 4 loops that are at the tip or the bottom of a domain, in proximity or adjacent to each other, and which may contribute to the binding of an antigen through forming an

antigen binding site or antigen binding pocket. It is preferred that the one or more sites of modification are located within the area of 10 amino acids, more preferably within 20, 30, 40, 50, 60, 70, 80, 90 up to 100 amino acids, in particular within a structural region to form a surface or pocket where the antigen can sterically access the loop regions.

In this regard the preferred modifications are engineered in the loop regions of CH1, CH2, CH3 and CH4, in particular in the range of amino acids 7 to 21, amino acids 25 to 39, amino acids 41 to 81, amino acids 83 to 85, amino acids 89 to 103 and amino acids 106 to 117.

In another preferred embodiment a modification in the structural loop region comprising amino acids 92 to 98 is combined with a modification in the structural loop region comprising amino acids 8 to 20.

The above identified amino acid regions of the respective immunoglobulins comprise loop regions to be modified. Preferably, a modification in the structural loop region comprising amino acids 92 to 98 is combined with a modification in one or more of the other structural loops.

In a preferred embodiment a modification in the structural loop region comprising amino acids 92 to 98 is combined with a modification in the structural loop region comprising amino acids 41 to 45.2.

Most preferably each of the structural loops comprising amino acids 92 to 98, amino acids 41 to 45.2 and amino acids 8 to 20 contain at least one amino acid modification.

In another preferred embodiment each of the structural loops comprising amino acids 92 to 98, amino acids 41 to 45.2, and amino acids 8 to 20 contain at least one amino acid modification.

According to another preferred embodiment the amino acid residues in the area of positions 15 to 17, 29 to 34, 41 to 45.2, 84 to 85, 92 to 100, and/or 108 to 115 of CH3 are modified.

The preferred modifications of Igk-C and Igl-C of human origin are engineered in the loop regions in the area of amino acids 8 to 20, amino acids 26 to 36, amino acids 41 to 82, amino acids 83 to 88, amino acids 92 to 100, amino acids 107 to 124 and amino acids 123 to 126.

The preferred modifications of loop regions of Igk-C and Igl-C of murine origin are engineered at sites in the area of amino acids 8 to 20, amino acids 26 to 36, amino acids 43 to 79, amino acids 83 to 85, amino acids 90 to 101, amino acids 108 to 116 and amino acids 122 to 126.

Another preferred immunoglobulin preferably used as a therapeutic according to the invention consists of a variable domain of a heavy or light chain, or a part thereof including a minidomain, with at least one loop region, preferably a structural loop region, and is characterised in that said at least one loop region comprises at least one amino acid modification forming at least one modified loop region, wherein said at least one modified loop region forms a relevant binding site as described above.

According to a specific embodiment the immunoglobulin preferably used according to the invention may contain a modification within the variable domain, which is selected from the group of VH, Vkappa, Vlambd, VHH and combinations thereof. More specifically, they comprise at least one modification within amino acids 7 to 22, amino acids 39 to 55, amino acids 66 to 79, amino acids 77 to 89 or amino acids 89 to 104, where the numbering of the amino acid position of the domains is that of the IMGT.

In a specific embodiment, the immunoglobulin preferably used according to the invention is characterised in that the loop regions of VH or Vkappa or Vlambd of human origin comprise at least one modification within amino acids 7 to

22, amino acids 43 to 51, amino acids 67 to 77, amino acids 77 to 88, and amino acids 89 to 104, most preferably amino acid positions 12 to 17, amino acid positions 45 to 50, amino acid positions 68 to 77, amino acids 79 to 88, and amino acid positions 92 to 99, where the numbering of the amino acid position of the domains is that of the IMGT.

The structural loop regions of the variable domain of the immunoglobulin of human origin, as possible selected for modification purposes are preferably located in the area of amino acids 8 to 20, amino acids 44 to 50, amino acids 67 to 76, amino acids 78 to 87, and amino acids 89 to 101.

According to a preferred embodiment the structural loop regions of the variable domain of the immunoglobulin of murine origin as possible selected for modification purposes are preferably located in the area of amino acids 6 to 20, amino acids 43 to 52, amino acids 67 to 79, amino acids 79 to 87, and amino acids 91 to 100.

The immunoglobulin preferably used as a therapeutic according to the invention may also be of camelid origin. Camel antibodies comprise only one heavy chain and have the same antigen affinity as normal antibodies consisting of light and heavy chains. Consequently camel antibodies are much smaller than, e.g., human antibodies, which allows them to penetrate dense tissues to reach the antigen, where larger proteins cannot. Moreover, the comparative simplicity, high affinity and specificity and the potential to reach and interact with active sites, camel's heavy chain antibodies present advantages over common antibodies in the design, production and application of clinically valuable compounds.

According to another preferred embodiment of the present invention the structural loop regions of a modular antibody or an immunoglobulins of camelid origin are modified, e.g. within a VHH, in the region of amino acids 7 to 19, amino acids 43 to 55, amino acids 68 to 76, amino acids 80 to 87 and amino acids 91 to 101.

The preferred method of producing the modular antibody according to the invention refers to engineering a modular antibody that is binding specifically to at least one first epitope and comprising modifications in each of at least two structural loop regions, and determining the specific binding of said at least two loop regions to at least one second epitope, wherein the unmodified structural loop region (non-CDR region) does not specifically bind to said at least one second epitope. Thus, an antibody or antigen-binding structure specific for a first antigen may be improved by adding another valency or specificity against a second antigen, which specificity may be identical, either targeting different epitopes or the same epitope, to increase valency or to obtain bi-, oligo- or multispecific molecules.

On the other hand it is preferred to make use of those modular antibodies that contain native structures interacting with effector molecules or immune cells. Those native structures either remain unchanged or are modulated for an increased effector function. Binding sites for e.g. Fc receptors are described to be located in a CH2 and/or CH3 domain region, and may be mutagenized by well known techniques.

ADCC, antibody-dependent cell-mediated cytotoxicity is the killing of antibody-coated target cells by cells with Fc receptors that recognize the constant region of the bound antibody. Most ADCC is mediated by NK cells that have the Fc receptor FcγRIII or CD16 on their surface. Typical assays employ target cells, like Ramos cells, incubated with serially diluted antibody prior to the addition of freshly isolated effector cells. The ADCC assay is then further incubated for several hours and % cytotoxicity detected.

Usually the Target: Effector ratio is about 1:16, but may be 1:1 up to 1:50.

Complement-dependent cytotoxicity (CDC) is a mechanism of killing cells in which antibody bound to the target cell surface fixes complement, which results in assembly of the membrane attack complex that punches holes in the target cell membrane resulting in subsequent cell lysis. The commonly used CDC assay follows the same procedure as for ADCC determination, however, with complement containing serum instead of effector cells.

The cytotoxic activity as determined by either of ADCC and CDC assay is proven for a modular antibody according to the invention, if there is a significant increase in the percentage of cytolysis as compared to a control. The absolute percentage increase preferably is higher than 5%, more preferably higher than 10%, even more preferred higher than 20%.

The antibody-dependent cellular phagocytosis, ADCP, sometimes called ADPC, is usually investigated side by side with cytolysis of cultured human cells. Phagocytosis by phagocytes, usually human monocytes or monocyte-derived macrophages, as mediated by an antibody can be determined as follows. Purified monocytes may be cultured with cytokines to enhance expression of FcγRs or to induce differentiation into macrophages. ADCP and ADCC assays are then performed with target cells. Phagocytosis is determined as the percentage of positive cells measured by flow cytometry. The positive ADCP activity is proven with a significant uptake of the antibody-antigen complex by the phagocytes. The absolute percentage preferably is higher than 5%, more preferably higher than 10%, even more preferred higher than 20%.

In a typical assay PBMC or monocytes or monocyte derived macrophages are resuspended in RPMI 1640 supplemented with 2% FCS in 96-well plates at a concentration of 1×10^5 viable cells in 100 μl/well. Appropriate target cells, expressing the target antigen, e.g. Her2/neu antigen and SKBR3 cells, are stained with PKH2 green fluorescence dye. Subsequently 1×10^4 PKH2-labeled target cells and an Her 2 specific (IgG1) antibody (or modular antibody) or mouse IgG1 isotype control (or modular antibody control) are added to the well of PBMC's in different concentrations (e.g. 1-100 pg/ml) and incubated in a final volume of 200 μl at 37° C. for 24 h. Following the incubation, PBMCs or monocytes or monocyte derived macrophages and target cells are harvested with EDTA-PBS and transferred to 96-well V-bottomed plates. The plates are centrifuged and the supernatant is aspirated. Cells are counterstained with a 100-μl mixture of RPE-conjugated anti-CD11b, anti-CD14, and human IgG, mixed and incubated for 60 min on ice. The cells are washed and fixed with 2% formaldehyde-PBS. Two-color flow cytometric analysis is performed with e.g. a FACS Calibur under optimal gating. PKH2-labeled target cells (green) are detected in the FL-1 channel (emission wavelength, 530 nm) and RPE-labeled PBMC or monocytes or monocyte derived macrophages (red) are detected in the FL-2 channel (emission wavelength, 575 nm). Residual target cells are defined as cells that are PKH2⁺/RPE⁻. Dual-labeled cells (PKH2⁺/RPE⁺) are considered to represent phagocytosis of targets by PBMC or monocytes or monocyte derived macrophages. Phagocytosis of target cells is calculated with the following equation: percent phagocytosis = $100 \times [(\text{percent dual positive}) / (\text{percent dual positive} + \text{percent residual targets})]$. All tests are usually performed in duplicate or triplicate and the results are expressed as mean ± 6 SD.

The effector function of the modular antibody according to the invention usually differs from any synthetic cytotoxic activity, e.g. through a toxin that may be conjugated to an immunoglobulin structure. Toxins usually do not activate effector molecules and the biological defence mechanism. Thus, the preferred cytotoxic activity of the modular antibodies according to the invention is a biological cytotoxic activity, which usually is immunostimulatory, leading to effective cytolysis.

The modular antibody according to the invention may specifically bind to any kind of binding molecules or structures, in particular to antigens, proteinaceous molecules, proteins, peptides, polypeptides, nucleic acids, glycans, carbohydrates, lipids, organic molecules, in particular small organic molecules, anorganic molecules, or combinations or fusions thereof, including PEG, prodrugs or drugs. The preferred modular antibody according to the invention may comprise at least two loops or loop regions whereby each of the loops or loop regions may specifically bind to different molecules or epitopes.

Preferably the target antigen is selected from cell surface antigens, including receptors, in particular from the group consisting of erbB receptor tyrosine kinases (such as EGFR, HER2, HER3 and HER4, in particular those epitopes of the extracellular domains of such receptors, e.g. the 4D5 epitope), molecules of the TNF-receptor superfamily, such as Apo-1 receptor, TNFR1, TNFR2, nerve growth factor receptor NGFR, CD40, T-cell surface molecules, T-cell receptors, T-cell antigen 0x40, TACI-receptor, BCMA, Apo-3, DR4, DR5, DR6, decoy receptors, such as DcR1, DcR2, CAR1, HVEM, GITR, ZTNFR-5, NTR1, TNFL1 but not limited to these molecules, B-cell surface antigens, such as CD10, CD19, CD20, CD21, CD22, antigens or markers of solid tumors or hematologic cancer cells, cells of lymphoma or leukaemia, other blood cells including blood platelets, but not limited to these molecules.

According to a further preferred embodiment the target antigen is selected from those antigens presented by cells, like epithelial cells, cells of solid tumors, infected cells, blood cells, antigen-presenting cells and mononuclear cells. Those target antigens expressed or overexpressed by cells are preferably targeted, which are selected from the group consisting of tumor associated antigens, in particular EpCAM, tumor-associated glycoprotein-72 (TAG-72), tumor-associated antigen CA 125, Prostate specific membrane antigen (PSMA), High molecular weight melanoma-associated antigen (HMW-MAA), tumor-associated antigen expressing Lewis Y related carbohydrate, Carcinoembryonic antigen (CEA), CEACAM5, HMFG PEM, mucin MUC1, MUC18 and cytokeratin tumor-associated antigen, bacterial antigens, viral antigens, allergens, allergy related molecules IgE, cKIT and Fc-epsilon-receptor1, IRp60, IL-5 receptor, CCR3, red blood cell receptor (CR1), human serum albumin, mouse serum albumin, rat serum albumin, Fc receptors, like neonatal Fc-gamma-receptor FcRn, Pc-gamma-receptors Pc-gamma RI, Pc-gamma-Rh, Pc-gamma RiII, Pc-alpha-receptors, Pc-epsilon-receptors, fluorescein, lysozyme, toll-like receptor 9, erythropoietin, CD2, CD3, CD3E, CD4, CD11, CD11a, CD14, CD16, CD18, CD19, CD20, CD22, CD23, CD25, CD28, CD29, CD30, CD32, CD33 (p67 protein), CD38, CD40, CD40L, CD52, CD54, CD56, CD64, CD80, CD147, GD3, IL-1, IL-1R, IL-2, IL-2R, IL-4, IL-5, IL-6, IL-6R, IL-8, IL-12, IL-15, IL-17, IL-18, IL-23, IL-28, IL-30, IL-31, IL-32, IL-33, IL-34, IL-35, IL-36, IL-37, IL-38, IL-39, IL-40, IL-41, IL-42, IL-43, IL-44, IL-45, IL-46, IL-47, IL-48, IL-49, IL-50, IL-51, IL-52, IL-53, IL-54, IL-55, IL-56, IL-57, IL-58, IL-59, IL-60, IL-61, IL-62, IL-63, IL-64, IL-65, IL-66, IL-67, IL-68, IL-69, IL-70, IL-71, IL-72, IL-73, IL-74, IL-75, IL-76, IL-77, IL-78, IL-79, IL-80, IL-81, IL-82, IL-83, IL-84, IL-85, IL-86, IL-87, IL-88, IL-89, IL-90, IL-91, IL-92, IL-93, IL-94, IL-95, IL-96, IL-97, IL-98, IL-99, IL-100, IL-101, IL-102, IL-103, IL-104, IL-105, IL-106, IL-107, IL-108, IL-109, IL-110, IL-111, IL-112, IL-113, IL-114, IL-115, IL-116, IL-117, IL-118, IL-119, IL-120, IL-121, IL-122, IL-123, IL-124, IL-125, IL-126, IL-127, IL-128, IL-129, IL-130, IL-131, IL-132, IL-133, IL-134, IL-135, IL-136, IL-137, IL-138, IL-139, IL-140, IL-141, IL-142, IL-143, IL-144, IL-145, IL-146, IL-147, IL-148, IL-149, IL-150, IL-151, IL-152, IL-153, IL-154, IL-155, IL-156, IL-157, IL-158, IL-159, IL-160, IL-161, IL-162, IL-163, IL-164, IL-165, IL-166, IL-167, IL-168, IL-169, IL-170, IL-171, IL-172, IL-173, IL-174, IL-175, IL-176, IL-177, IL-178, IL-179, IL-180, IL-181, IL-182, IL-183, IL-184, IL-185, IL-186, IL-187, IL-188, IL-189, IL-190, IL-191, IL-192, IL-193, IL-194, IL-195, IL-196, IL-197, IL-198, IL-199, IL-200, IL-201, IL-202, IL-203, IL-204, IL-205, IL-206, IL-207, IL-208, IL-209, IL-210, IL-211, IL-212, IL-213, IL-214, IL-215, IL-216, IL-217, IL-218, IL-219, IL-220, IL-221, IL-222, IL-223, IL-224, IL-225, IL-226, IL-227, IL-228, IL-229, IL-230, IL-231, IL-232, IL-233, IL-234, IL-235, IL-236, IL-237, IL-238, IL-239, IL-240, IL-241, IL-242, IL-243, IL-244, IL-245, IL-246, IL-247, IL-248, IL-249, IL-250, IL-251, IL-252, IL-253, IL-254, IL-255, IL-256, IL-257, IL-258, IL-259, IL-260, IL-261, IL-262, IL-263, IL-264, IL-265, IL-266, IL-267, IL-268, IL-269, IL-270, IL-271, IL-272, IL-273, IL-274, IL-275, IL-276, IL-277, IL-278, IL-279, IL-280, IL-281, IL-282, IL-283, IL-284, IL-285, IL-286, IL-287, IL-288, IL-289, IL-290, IL-291, IL-292, IL-293, IL-294, IL-295, IL-296, IL-297, IL-298, IL-299, IL-300, IL-301, IL-302, IL-303, IL-304, IL-305, IL-306, IL-307, IL-308, IL-309, IL-310, IL-311, IL-312, IL-313, IL-314, IL-315, IL-316, IL-317, IL-318, IL-319, IL-320, IL-321, IL-322, IL-323, IL-324, IL-325, IL-326, IL-327, IL-328, IL-329, IL-330, IL-331, IL-332, IL-333, IL-334, IL-335, IL-336, IL-337, IL-338, IL-339, IL-340, IL-341, IL-342, IL-343, IL-344, IL-345, IL-346, IL-347, IL-348, IL-349, IL-350, IL-351, IL-352, IL-353, IL-354, IL-355, IL-356, IL-357, IL-358, IL-359, IL-360, IL-361, IL-362, IL-363, IL-364, IL-365, IL-366, IL-367, IL-368, IL-369, IL-370, IL-371, IL-372, IL-373, IL-374, IL-375, IL-376, IL-377, IL-378, IL-379, IL-380, IL-381, IL-382, IL-383, IL-384, IL-385, IL-386, IL-387, IL-388, IL-389, IL-390, IL-391, IL-392, IL-393, IL-394, IL-395, IL-396, IL-397, IL-398, IL-399, IL-400, IL-401, IL-402, IL-403, IL-404, IL-405, IL-406, IL-407, IL-408, IL-409, IL-410, IL-411, IL-412, IL-413, IL-414, IL-415, IL-416, IL-417, IL-418, IL-419, IL-420, IL-421, IL-422, IL-423, IL-424, IL-425, IL-426, IL-427, IL-428, IL-429, IL-430, IL-431, IL-432, IL-433, IL-434, IL-435, IL-436, IL-437, IL-438, IL-439, IL-440, IL-441, IL-442, IL-443, IL-444, IL-445, IL-446, IL-447, IL-448, IL-449, IL-450, IL-451, IL-452, IL-453, IL-454, IL-455, IL-456, IL-457, IL-458, IL-459, IL-460, IL-461, IL-462, IL-463, IL-464, IL-465, IL-466, IL-467, IL-468, IL-469, IL-470, IL-471, IL-472, IL-473, IL-474, IL-475, IL-476, IL-477, IL-478, IL-479, IL-480, IL-481, IL-482, IL-483, IL-484, IL-485, IL-486, IL-487, IL-488, IL-489, IL-490, IL-491, IL-492, IL-493, IL-494, IL-495, IL-496, IL-497, IL-498, IL-499, IL-500, IL-501, IL-502, IL-503, IL-504, IL-505, IL-506, IL-507, IL-508, IL-509, IL-510, IL-511, IL-512, IL-513, IL-514, IL-515, IL-516, IL-517, IL-518, IL-519, IL-520, IL-521, IL-522, IL-523, IL-524, IL-525, IL-526, IL-527, IL-528, IL-529, IL-530, IL-531, IL-532, IL-533, IL-534, IL-535, IL-536, IL-537, IL-538, IL-539, IL-540, IL-541, IL-542, IL-543, IL-544, IL-545, IL-546, IL-547, IL-548, IL-549, IL-550, IL-551, IL-552, IL-553, IL-554, IL-555, IL-556, IL-557, IL-558, IL-559, IL-560, IL-561, IL-562, IL-563, IL-564, IL-565, IL-566, IL-567, IL-568, IL-569, IL-570, IL-571, IL-572, IL-573, IL-574, IL-575, IL-576, IL-577, IL-578, IL-579, IL-580, IL-581, IL-582, IL-583, IL-584, IL-585, IL-586, IL-587, IL-588, IL-589, IL-590, IL-591, IL-592, IL-593, IL-594, IL-595, IL-596, IL-597, IL-598, IL-599, IL-600, IL-601, IL-602, IL-603, IL-604, IL-605, IL-606, IL-607, IL-608, IL-609, IL-610, IL-611, IL-612, IL-613, IL-614, IL-615, IL-616, IL-617, IL-618, IL-619, IL-620, IL-621, IL-622, IL-623, IL-624, IL-625, IL-626, IL-627, IL-628, IL-629, IL-630, IL-631, IL-632, IL-633, IL-634, IL-635, IL-636, IL-637, IL-638, IL-639, IL-640, IL-641, IL-642, IL-643, IL-644, IL-645, IL-646, IL-647, IL-648, IL-649, IL-650, IL-651, IL-652, IL-653, IL-654, IL-655, IL-656, IL-657, IL-658, IL-659, IL-660, IL-661, IL-662, IL-663, IL-664, IL-665, IL-666, IL-667, IL-668, IL-669, IL-670, IL-671, IL-672, IL-673, IL-674, IL-675, IL-676, IL-677, IL-678, IL-679, IL-680, IL-681, IL-682, IL-683, IL-684, IL-685, IL-686, IL-687, IL-688, IL-689, IL-690, IL-691, IL-692, IL-693, IL-694, IL-695, IL-696, IL-697, IL-698, IL-699, IL-700, IL-701, IL-702, IL-703, IL-704, IL-705, IL-706, IL-707, IL-708, IL-709, IL-710, IL-711, IL-712, IL-713, IL-714, IL-715, IL-716, IL-717, IL-718, IL-719, IL-720, IL-721, IL-722, IL-723, IL-724, IL-725, IL-726, IL-727, IL-728, IL-729, IL-730, IL-731, IL-732, IL-733, IL-734, IL-735, IL-736, IL-737, IL-738, IL-739, IL-740, IL-741, IL-742, IL-743, IL-744, IL-745, IL-746, IL-747, IL-748, IL-749, IL-750, IL-751, IL-752, IL-753, IL-754, IL-755, IL-756, IL-757, IL-758, IL-759, IL-760, IL-761, IL-762, IL-763, IL-764, IL-765, IL-766, IL-767, IL-768, IL-769, IL-770, IL-771, IL-772, IL-773, IL-774, IL-775, IL-776, IL-777, IL-778, IL-779, IL-780, IL-781, IL-782, IL-783, IL-784, IL-785, IL-786, IL-787, IL-788, IL-789, IL-790, IL-791, IL-792, IL-793, IL-794, IL-795, IL-796, IL-797, IL-798, IL-799, IL-800, IL-801, IL-802, IL-803, IL-804, IL-805, IL-806, IL-807, IL-808, IL-809, IL-810, IL-811, IL-812, IL-813, IL-814, IL-815, IL-816, IL-817, IL-818, IL-819, IL-820, IL-821, IL-822, IL-823, IL-824, IL-825, IL-826, IL-827, IL-828, IL-829, IL-830, IL-831, IL-832, IL-833, IL-834, IL-835, IL-836, IL-837, IL-838, IL-839, IL-840, IL-841, IL-842, IL-843, IL-844, IL-845, IL-846, IL-847, IL-848, IL-849, IL-850, IL-851, IL-852, IL-853, IL-854, IL-855, IL-856, IL-857, IL-858, IL-859, IL-860, IL-861, IL-862, IL-863, IL-864, IL-865, IL-866, IL-867, IL-868, IL-869, IL-870, IL-871, IL-872, IL-873, IL-874, IL-875, IL-876, IL-877, IL-878, IL-879, IL-880, IL-881, IL-882, IL-883, IL-884, IL-885, IL-886, IL-887, IL-888, IL-889, IL-890, IL-891, IL-892, IL-893, IL-894, IL-895, IL-896, IL-897, IL-898, IL-899, IL-900, IL-901, IL-902, IL-903, IL-904, IL-905, IL-906, IL-907, IL-908, IL-909, IL-910, IL-911, IL-912, IL-913, IL-914, IL-915, IL-916, IL-917, IL-918, IL-919, IL-920, IL-921, IL-922, IL-923, IL-924, IL-925, IL-926, IL-927, IL-928, IL-929, IL-930, IL-931, IL-932, IL-933, IL-934, IL-935, IL-936, IL-937, IL-938, IL-939, IL-940, IL-941, IL-942, IL-943, IL-944, IL-945, IL-946, IL-947, IL-948, IL-949, IL-950, IL-951, IL-952, IL-953, IL-954, IL-955, IL-956, IL-957, IL-958, IL-959, IL-960, IL-961, IL-962, IL-963, IL-964, IL-965, IL-966, IL-967, IL-968, IL-969, IL-970, IL-971, IL-972, IL-973, IL-974, IL-975, IL-976, IL-977, IL-978, IL-979, IL-980, IL-981, IL-982, IL-983, IL-984, IL-985, IL-986, IL-987, IL-988, IL-989, IL-990, IL-991, IL-992, IL-993, IL-994, IL-995, IL-996, IL-997, IL-998, IL-999, IL-1000, IL-1001, IL-1002, IL-1003, IL-1004, IL-1005, IL-1006, IL-1007, IL-1008, IL-1009, IL-1010, IL-1011, IL-1012, IL-1013, IL-1014, IL-1015, IL-1016, IL-1017, IL-1018, IL-1019, IL-1020, IL-1021, IL-1022, IL-1023, IL-1024, IL-1025, IL-1026, IL-1027, IL-1028, IL-1029, IL-1030, IL-1031, IL-1032, IL-1033, IL-1034, IL-1035, IL-1036, IL-1037, IL-1038, IL-1039, IL-1040, IL-1041, IL-1042, IL-1043, IL-1044, IL-1045, IL-1046, IL-1047, IL-1048, IL-1049, IL-1050, IL-1051, IL-1052, IL-1053, IL-1054, IL-1055, IL-1056, IL-1057, IL-1058, IL-1059, IL-1060, IL-1061, IL-1062, IL-1063, IL-1064, IL-1065, IL-1066, IL-1067, IL-1068, IL-1069, IL-1070, IL-1071, IL-1072, IL-1073, IL-1074, IL-1075, IL-1076, IL-1077, IL-1078, IL-1079, IL-1080, IL-1081, IL-1082, IL-1083, IL-1084, IL-1085, IL-1086, IL-1087, IL-1088, IL-1089, IL-1090, IL-1091, IL-1092, IL-1093, IL-1094, IL-1095, IL-1096, IL-1097, IL-1098, IL-1099, IL-1100, IL-1101, IL-1102, IL-1103, IL-1104, IL-1105, IL-1106, IL-1107, IL-1108, IL-1109, IL-1110, IL-1111, IL-1112, IL-1113, IL-1114, IL-1115, IL-1116, IL-1117, IL-1118, IL-1119, IL-1120, IL-1121, IL-1122, IL-1123, IL-1124, IL-1125, IL-1126, IL-1127, IL-1128, IL-1129, IL-1130, IL-1131, IL-1132, IL-1133, IL-1134, IL-1135, IL-1136, IL-1137, IL-1138, IL-1139, IL-1140, IL-1141, IL-1142, IL-1143, IL-1144, IL-1145, IL-1146, IL-1147, IL-1148, IL-1149, IL-1150, IL-1151, IL-1152, IL-1153, IL-1154, IL-1155, IL-1156, IL-1157, IL-1158, IL-1159, IL-1160, IL-1161, IL-1162, IL-1163, IL-1164, IL-1165, IL-1166, IL-1167, IL-1168, IL-1169, IL-1170, IL-1171, IL-1172, IL-1173, IL-1174, IL-1175, IL-1176, IL-1177, IL-1178, IL-1179, IL-1180, IL-1181, IL-1182, IL-1183, IL-1184, IL-1185, IL-1186, IL-1187, IL-1188, IL-1189, IL-1190, IL-1191, IL-1192, IL-1193, IL-1194, IL-1195, IL-1196, IL-1197, IL-1198, IL-1199, IL-1200, IL-1201, IL-1202, IL-1203, IL-1204, IL-1205, IL-1206, IL-1207, IL-1208, IL-1209, IL-1210, IL-1211, IL-1212, IL-1213, IL-1214, IL-1215, IL-1216, IL-1217, IL-1218, IL-1219, IL-1220, IL-1221, IL-1222, IL-1223, IL-1224, IL-1225, IL-1226, IL-1227, IL-1228, IL-1229, IL-1230, IL-1231, IL-1232, IL-1233, IL-1234, IL-1235, IL-1236, IL-1237, IL-1238, IL-1239, IL-1240, IL-1241, IL-1242, IL-1243, IL-1244, IL-1245, IL-1246, IL-1247, IL-1248, IL-1249, IL-1250, IL-1251, IL-1252, IL-1253, IL-1254, IL-1255, IL-1256, IL-1257, IL-1258, IL-1259, IL-1260, IL-1261, IL-1262, IL-1263, IL-1264, IL-1265, IL-1266, IL-1267, IL-1268, IL-1269, IL-1270, IL-1271, IL-1272, IL-1273, IL-1274, IL-1275, IL-1276, IL-1277, IL-1278, IL-1279, IL-1280, IL-1281, IL-1282, IL-1283, IL-1284, IL-1285, IL-1286, IL-1287, IL-1288, IL-1289, IL-1290, IL-1291, IL-1292, IL-1293, IL-1294, IL-1295, IL-1296, IL-1297, IL-1298, IL-1299, IL-1300, IL-1301, IL-1302, IL-1303, IL-1304, IL-1305, IL-1306, IL-1307, IL-1308, IL-1309, IL-1310, IL-1311, IL-1312, IL-1313, IL-1314, IL-1315, IL-1316, IL-1317, IL-1318, IL-1319, IL-1320, IL-1321, IL-1322, IL-1323, IL-1324, IL-1325, IL-1326, IL-1327, IL-1328, IL-1329, IL-1330, IL-1331, IL-1332, IL-1333, IL-1334, IL-1335, IL-1336, IL-1337, IL-1338, IL-1339, IL-1340, IL-1341, IL-1342, IL-1343, IL-1344, IL-1345, IL-1346, IL-1347, IL-1348, IL-1349, IL-1350, IL-1351, IL-1352, IL-1353, IL-1354, IL-1355, IL-1356, IL-1357, IL-1358, IL-1359, IL-1360, IL-1361, IL-1362, IL-1363, IL-1364, IL-1365, IL-1366, IL-1367, IL-1368, IL-1369, IL-1370, IL-1371, IL-1372, IL-1373, IL-1374, IL-1375, IL-1376, IL-1377, IL-1378, IL-1379, IL-1380, IL-1381, IL-1382, IL-1383, IL-1384, IL-1385, IL-1386, IL-1387, IL-1388, IL-1389, IL-1390, IL-1391, IL-1392, IL-1393, IL-1394, IL-1395, IL-1396, IL-1397, IL-1398, IL-1399, IL-1400, IL-1401, IL-1402, IL-1403, IL-1404, IL-1405, IL-1406, IL-1407, IL-1408, IL-1409, IL-1410, IL-1411, IL-1412, IL-1413, IL-1414, IL-1415, IL-1416, IL-1417, IL-1418, IL-1419, IL-1420, IL-1421, IL-1422, IL-1423, IL-1424, IL-1425, IL-1426, IL-1427, IL-1428, IL-1429, IL-1430, IL-1431, IL-1432, IL-1433, IL-1434, IL-1435, IL-1436, IL-1437, IL-1438, IL-1439, IL-1440, IL-1441, IL-1442, IL-1443, IL-1444, IL-1445, IL-1446, IL-1447, IL-1448, IL-1449, IL-1450, IL-1451, IL-1452, IL-1453, IL-1454, IL-1455, IL-1456, IL-1457, IL-1458, IL-1459, IL-1460, IL-1461, IL-1462, IL-1463, IL-1464, IL-1465, IL-1466, IL-1467, IL-1468, IL-1469, IL-1470, IL-1471, IL-1472, IL-1473, IL-1474, IL-1475, IL-1476, IL-1477, IL-1478, IL-1479, IL-1480, IL-1481, IL-1482, IL-1483, IL-1484, IL-1485, IL-1486, IL-1487, IL-1488, IL-1489, IL-1490, IL-1491, IL-1492, IL-1493, IL-1494, IL-1495, IL-1496, IL-1497, IL-1498, IL-1499, IL-1500, IL-1501, IL-1502, IL-1503, IL-1504, IL-1505, IL-1506, IL-1507, IL-1508, IL-1509, IL-1510, IL-1511, IL-1512, IL-1513, IL-1514, IL-1515, IL-1516, IL-1517, IL-1518, IL-1519, IL-1520, IL-1521, IL-1522, IL-1523, IL-1524, IL-1525, IL-1526, IL-1527, IL-1528, IL-1529, IL-1530, IL-1531, IL-1532, IL-1533, IL-1534, IL-1535, IL-1536, IL-1537, IL-1538, IL-1539, IL-1540, IL-1541, IL-1542, IL-1543, IL-1544, IL-1545, IL-1546, IL-1547, IL-1548, IL-1549, IL-1550, IL-1551, IL-1552, IL-1553, IL-1554, IL-1555, IL-1556, IL-1557, IL-1558, IL-1559, IL-1560, IL-1561, IL-1562, IL-1563, IL-1564, IL-1565, IL-1566, IL-1567, IL-1568, IL-1569, IL-1570, IL-1571, IL-1572, IL-1573, IL-1574, IL-1575, IL-1576, IL-1577, IL-1578, IL-1579, IL-1580, IL-1581, IL-1582, IL-1583, IL-1584, IL-1585, IL-1586, IL-1587, IL-1588, IL-1589, IL-1590, IL-1591, IL-1592, IL-1593, IL-1594, IL-1595, IL-1596, IL-1597, IL-1598, IL-1599, IL-1600, IL-1601, IL-1602, IL-1603, IL-1604, IL-1605, IL-1606, IL-1607, IL-1608, IL-1609, IL-1610, IL-1611, IL-1612, IL-1613, IL-1614, IL-1615, IL-1616, IL

LIGHT, VEG1, OX4OL, TRAIL Receptor-1, AI Adenosine Receptor, Lymphotoxin Beta Receptor, TACI, BAFF-R, EPO; LFA-3, ICAM-1, ICAM-3, integrin beta1, integrin beta2, integrin alpha4/beta7, integrin alpha2, integrin alpha3, integrin alpha4, integrin alpha5, integrin alpha6, integrin alphav, alphaVbeta3 integrin, FGFR-3, Keratinocyte Growth Factor, GM-CSF, M-CSF, RANKL, VLA-1, VLA-4, L-selectin, anti-Id, E-selectin, HLA, HLA-DR, CTLA-4, T cell receptor, B7-1, 87-2, VNRintegrin, TGE-beta1, TGFbeta2, eotaxin1, BLyS (B-lymphocyte Stimulator), complement C5, IgE, IgA, IgD, IgM, IgG, factor VII, CBL, NCA 90, EGFR (ErbB-1), Her2/neu (ErbB2), Her3 (ErbB-3), Her4 (ErbB4), Tissue Factor, VEGF, VEGFR, endothelin receptor, VLA-4, carbohydrates such as blood group antigens and related carbohydrates, Galili-Glycosylation, Gastrin, Gastrin receptors, tumor associated carbohydrates, Hapten NP-cap or NIP-cap, T cell receptor alpha/beta, E-selectin, P-glycoprotein, MRP3, MRP5, glutathione-S-transferase pi (multi drug resistance proteins), alpha-granule membrane protein (GMP) 140, digoxin, placental alkaline phosphatase (PLAP) and testicular PLAP-like alkaline phosphatase, transferrin receptor, Heparanase I, human cardiac myosin, Glycoprotein IIb/IIIa (GPIIb/IIIa), human cytomegalovirus (HCMV) gH envelope glycoprotein, HIV gp120, HCMV, respiratory syncytial virus RSV F, RSVF Fgp, VNRintegrin, Hep B gp120, CMV, gp120, HIV IIIB gp120 V3 loop, respiratory syncytial virus (RSV) Fgp, Herpes simplex virus (HSV) gD glycoprotein, HSV gB glycoprotein, HCMV gB envelope glycoprotein, *Clostridium perfringens* toxin and fragments thereof.

Preferred modular antibodies according to the invention are binding said target antigen with a high affinity, in particular with a high on and/or a low off rate, or a high avidity of binding. Usually a binder is considered a high affinity binder with a K_d of $<10^{-9}$ M. Medium affinity binders with a K_d of less than 10^{-6} up to 10^{-9} M may be provided according to the invention as well, preferably in conjunction with an affinity maturation process.

Affinity maturation is the process by which antibodies with increased affinity for antigen are produced. With structural changes of an antibody, including amino acid mutagenesis or as a consequence of somatic mutation in immunoglobulin gene segments, variants of a binding site to an antigen are produced and selected for greater affinities. Affinity matured modular antibodies may exhibit a several logfold greater affinity than a parent antibody. Single parent antibodies may be subject to affinity maturation. Alternatively pools of modular antibodies with similar binding affinity to the target antigen may be considered as parent structures that are varied to obtain affinity matured single antibodies or affinity matured pools of such antibodies.

The preferred affinity matured variant of a modular antibody according to the invention exhibits at least a 10 fold increase in affinity of binding, preferably at least a 100 fold increase. The affinity maturation may be employed in the course of the selection campaigns employing respective libraries of parent molecules, either with modular antibodies having medium binding affinity to obtain the modular antibody of the invention having the specific target binding property of a $K_d < 10^{-8}$ M and/or a potency of $IC_{50} < 10^{-8}$ M. Alternatively, the binding potency or affinity may be even more increased by affinity maturation of the modular antibody according to the invention to obtain the high values corresponding to a K_d or IC_{50} of less than 10^{-9} M, preferably less than 10^{-10} M or even less than 10^{-11} M, most preferred in the picomolar range.

The IC_{50} , also called 50% saturation concentration, is a measure for the binding potency of a modular antibody. It is the molar concentration of a binder, which produces 50% of the maximum possible binding at equilibrium or under saturation. The potency of an antagonist is usually defined by its IC_{50} value. This can be calculated for a given antagonist by determining the concentration of antagonist needed to elicit half saturation of the maximum binding of an agonist. Elucidating an IC_{50} value is useful for comparing the potency of antibodies or antibody variants with similar efficacies; however the dose-response curves produced by both drug antagonists must be similar. The lower the IC_{50} , the greater the potency of the antagonist, and the lower the concentration of drug that is required to inhibit the maximum biological response, like effector function or cytotoxic activity. Lower concentrations of drugs may also be associated with fewer side effects.

Usually the affinity of an antibody correlates well with the IC_{50} . The affinity of an antagonist for its binding site (K_i), is understood as its ability to bind to a receptor, which determines the duration of binding and respective agonist activity. Measures to increase the affinity by affinity maturation usually also increase the potency of binding, resulting in the respective reduction of IC_{50} values in the same range of the K_d values.

The IC_{50} and K_d values may be determined using the saturation binding assays well-known in the art.

The modular antibody according to the invention is preferably conjugated to a label or reporter molecule, selected from the group consisting of organic molecules, enzyme labels, radioactive labels, colored labels, fluorescent labels, chromogenic labels, luminescent labels, haptens, digoxigenin, biotin, metal complexes, metals, colloidal gold and mixtures thereof. Modified immunoglobulins conjugated to labels or reporter molecules may be used, for instance, in assay systems or diagnostic methods.

The modular antibody according to the invention may be conjugated to other molecules which allow the simple detection of said conjugate in, for instance, binding assays (e.g. ELISA) and binding studies.

In a preferred embodiment, antibody variants are screened using one or more cell-based or in vivo assays. For such assays, purified or unpurified modified immunoglobulins are typically added exogenously such that cells are exposed to individual immunoglobulins or pools of immunoglobulins belonging to a library. These assays are typically, but not always, based on the function of the immunoglobulin; that is, the ability of the antibody to bind to its target and mediate some biochemical event, for example effector function, ligand/receptor binding inhibition, apoptosis, and the like. Such assays often involve monitoring the response of cells to the antibody, for example cell survival, cell death, change in cellular morphology, or transcriptional activation such as cellular expression of a natural gene or reporter gene. For example, such assays may measure the ability of antibody variants to elicit ADCC, ADCP, or CDC. For some assays additional cells or components, that is in addition to the target cells, may need to be added, for example serum complement, or effector cells such as peripheral blood monocytes (PBMCs), NK cells, macrophages, and the like. Such additional cells may be from any organism, preferably humans, mice, rat, rabbit, and monkey. Modular antibodies may cause apoptosis of certain cell lines expressing the target, or they may mediate attack on target cells by immune cells which have been added to the assay. Methods for monitoring cell death or viability are known in the art, and include the use of dyes, immunochemical, cytochemical,

25

and radioactive reagents. For example, caspase staining assays may enable apoptosis to be measured, and uptake or release of radioactive substrates or fluorescent dyes such as alamar blue may enable cell growth or activation to be monitored.

In a preferred embodiment, the DELFIART EuTDA-based cytotoxicity assay (Perkin Elmer, MA) may be used.

Alternatively, dead or damaged target cells may be monitored by measuring the release of one or more natural intracellular components, for example lactate dehydrogenase.

Transcriptional activation may also serve as a method for assaying function in cell-based assays. In this case, response may be monitored by assaying for natural genes or immunoglobulins which may be upregulated, for example the release of certain interleukins may be measured, or alternatively readout may be via a reporter construct. Cell-based assays may also involve the measure of morphological changes of cells as a response to the presence of modular antibodies. Cell types for such assays may be prokaryotic or eukaryotic, and a variety of cell lines that are known in the art may be employed. Alternatively, cell-based screens are performed using cells that have been transformed or transfected with nucleic acids encoding the variants. That is, antibody variants are not added exogenously to the cells. For example, in one embodiment, the cell-based screen utilizes cell surface display. A fusion partner can be employed that enables display of modified immunoglobulins on the surface of cells (Wittrup, 2001, *Curr Opin Biotechnol*, 12:395-399).

In a preferred embodiment, the immunogenicity of the modular antibodies may be determined experimentally using one or more cell-based assays. In a preferred embodiment, ex vivo T-cell activation assays are used to experimentally quantitate immunogenicity. In this method, antigen presenting cells and naive T cells from matched donors are challenged with a peptide or whole antibody of interest one or more times. Then, T cell activation can be detected using a number of methods, for example by monitoring production of cytokines or measuring uptake of tritiated thymidine. In the most preferred embodiment, interferon gamma production is monitored using Elispot assays.

The biological properties of the modular antibody according to the invention may be characterized ex vivo in cell, tissue, and whole organism experiments. As is known in the art, drugs are often tested in vivo in animals, including but not limited to mice, rats, rabbits, dogs, cats, pigs, and monkeys, in order to measure a drug's efficacy for treatment against a disease or disease model, or to measure a drug's pharmacokinetics, pharmacodynamics, toxicity, and other properties. The animals may be referred to as disease models. Therapeutics are often tested in mice, including but not limited to nude mice, SCID mice, xenograft mice, and transgenic mice (including knockins and knockouts). Such experimentation may provide meaningful data for determination of the potential of the antibody to be used as a therapeutic with the appropriate half-life, effector function, apoptotic activity, cytotoxic or cytolytic activity. Any organism, preferably mammals, may be used for testing. For example because of their genetic similarity to humans, primates, monkeys can be suitable therapeutic models, and thus may be used to test the efficacy, toxicity, pharmacokinetics, pharmacodynamics, half-life, or other property of the modular antibody according to the invention. Tests of the substances in humans are ultimately required for approval as drugs, and thus of course these experiments are contemplated. Thus the modular antibodies of the present invention may be tested in humans to determine their therapeutic

26

efficacy, toxicity, immunogenicity, pharmacokinetics, and/or other clinical properties. Especially those modular antibodies according to the invention that bind to single cell or a cellular complex through at least two binding motifs, preferably binding of at least three structures cross-linking target cells, would be considered effective in effector activity or preapoptotic or apoptotic activity upon cell targeting and cross-linking. Multivalent binding provides a relatively large association of binding partners, also called cross-linking, which is a prerequisite for apoptosis and cell death.

The modular antibody of the present invention may find use in a wide range of antibody products. In one embodiment the modular antibody of the present invention is used for therapy or prophylaxis, e.g. as an active or passive immunotherapy, for preparative, industrial or analytic use, as a diagnostic, an industrial compound or a research reagent, preferably a therapeutic. The modular antibody may find use in an antibody composition that is monoclonal or polyclonal. In a preferred embodiment, the modular antibodies of the present invention are used to capture or kill target cells that bear the target antigen, for example cancer cells. In an alternate embodiment, the modular antibodies of the present invention are used to block, antagonize, or agonize the target antigen, for example by antagonizing a cytokine or cytokine receptor.

In an alternately preferred embodiment, the modular antibodies of the present invention are used to block, antagonize, or agonize growth factors or growth factor receptors and thereby mediate killing the target cells that bear or need the target antigen.

In an alternately preferred embodiment, the modular antibodies of the present invention are used to block, antagonize, or agonize enzymes and substrate of enzymes.

In a preferred embodiment, a modular antibody is administered to a patient to treat a specific disorder. A "patient" for the purposes of the present invention includes both humans and other animals, preferably mammals and most preferably humans. By "specific disorder" herein is meant a disorder that may be ameliorated by the administration of a pharmaceutical composition comprising a modified immunoglobulin of the present invention.

In one embodiment, a modular antibody according to the present invention is the only therapeutically active agent administered to a patient. Alternatively, the modular antibody according to the present invention is administered in combination with one or more other therapeutic agents, including but not limited to cytotoxic agents, chemotherapeutic agents, cytokines, growth inhibitory agents, anti-hormonal agents, kinase inhibitors, anti-angiogenic agents, cardioprotectants, or other therapeutic agents. The modular antibody may be administered concomitantly with one or more other therapeutic regimens. For example, a modular antibody of the present invention may be administered to the patient along with chemotherapy, radiation therapy, or both chemotherapy and radiation therapy. In one embodiment, the modular antibody of the present invention may be administered in conjunction with one or more antibodies, which may or may not comprise a modular antibody of the present invention. In accordance with another embodiment of the invention, the modular antibody of the present invention and one or more other anti-cancer therapies is employed to treat cancer cells ex vivo. It is contemplated that such ex vivo treatment may be useful in bone marrow transplantation and particularly, autologous bone marrow transplantation. It is of course contemplated that the antibodies of the invention can be employed in combination with still other therapeutic techniques such as surgery.

A variety of other therapeutic agents may find use for administration with the modular antibody of the present invention. In one embodiment, the modular antibody is administered with an anti-angiogenic agent, which is a compound that blocks, or interferes to some degree, the development of blood vessels. The anti-angiogenic factor may, for instance, be a small molecule or a protein, for example an antibody, Fc fusion molecule, or cytokine, that binds to a growth factor or growth factor receptor involved in promoting angiogenesis. The preferred anti-angiogenic factor herein is an antibody that binds to Vascular Endothelial Growth Factor (VEGF). In an alternate embodiment, the modular antibody is administered with a therapeutic agent that induces or enhances adaptive immune response, for example an antibody that targets CTLA-4. In an alternate embodiment, the modified immunoglobulin is administered with a tyrosine kinase inhibitor, which is a molecule that inhibits to some extent tyrosine kinase activity of a tyrosine kinase. In an alternate embodiment, the modular antibody of the present invention is administered with a cytokine. By "cytokine" as used herein is meant a generic term for proteins released by one cell population that act on another cell as intercellular mediators including chemokines.

Pharmaceutical compositions are contemplated wherein modular antibodies of the present invention and one or more therapeutically active agents are formulated. Stable formulations of the modular antibodies of the present invention are prepared for storage by mixing said immunoglobulin having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers, in the form of lyophilized formulations or aqueous solutions. The formulations to be used for in vivo administration are preferably sterile. This is readily accomplished by filtration through sterile filtration membranes or other methods. The modular antibody and other therapeutically active agents disclosed herein may also be formulated as immunoliposomes, and/or entrapped in microcapsules.

Administration of the pharmaceutical composition comprising a modular antibody of the present invention, preferably in the form of a sterile aqueous solution, may be done in a variety of ways, including, but not limited to, orally, subcutaneously, intravenously, intranasally, intraotically, transdermally, mucosal, topically (e.g., gels, salves, lotions, creams, etc.), intraperitoneally, intramuscularly, intrapulmonary (e.g., *AERX*TM inhalable technology commercially available from Aradigm, or *I_{hance}*TM pulmonary delivery system commercially available from Inhale Therapeutics), vaginally, parenterally, rectally, or intraocularly.

A preferred method according to the invention refers to a randomly modified nucleic acid molecule coding for an immunoglobulin, immunoglobulin domain or a part thereof which comprises at least one nucleotide repeating unit within a structural loop coding region having the sequence 5T-NNS-3', 5T-NNN-3', 5'-NNB-3' or 5'-NNK-3'. In some embodiments the modified nucleic acid comprises nucleotide codons selected from the group of TNT, WMT, BMT, RMC, RMG, MRT, SRC, KMT, RST, YMT, MKC, RSA, RRC, NNK, NNN, NNS or any combination thereof (the coding is according to IUPAC).

The modification of the nucleic acid molecule may be performed by introducing synthetic oligonucleotides into a larger segment of nucleic acid or by de novo synthesis of a complete nucleic acid molecule. Synthesis of nucleic acid may be performed with tri-nucleotide building blocks which would reduce the number of nonsense sequence combina-

tions if a subset of amino acids is to be encoded (e.g. Yanez et al. *Nucleic Acids Res.* (2004) 32:e158; Virnekas et al. *Nucleic Acids Res.* (1994) 22:5600-5607)

The randomly modified nucleic acid molecule may comprise the above identified repeating units, which code for all known naturally occurring amino acids or a subset thereof. Those libraries that contain modified sequences wherein a specific subset of amino acids are used for modification purposes are called "focused" libraries. The members of such libraries have an increased probability of an amino acid of such a subset at the modified position, which is at least two times higher than usual, preferably at least 3 times or even at least 4 times higher. Such libraries have also a limited or lower number of library members, so that the number of actual library members reaches the number of theoretical library members. In some cases the number of library members of a focused library is not less than 10³ times the theoretical number, preferably not less than 10² times, most preferably not less than 10 times.

Usually libraries according to the invention comprise at least 10 fusion proteins or potential binding agents or variants of scaffold proteins, preferably at least 100, more preferred at least 1000, more preferred at least 10⁴, more preferred at least 10⁵, more preferred at least 10⁶, more preferred at least 10⁷, more preferred at least 10⁸, more preferred at least 10% more preferred at least 10¹⁰, more preferred at least 10¹¹, up to 10¹², in cases of in vitro display methods, such as ribosomal display, even higher number are feasible.

Various alternatives are available for the manufacture of the gene encoding the randomized library. It is possible to produce the DNA by a completely synthetic approach, in which the sequence is divided into overlapping fragments which are subsequently prepared as synthetic oligonucleotides. These oligonucleotides are mixed together, and annealed to each other by first heating to ca. 100° C. and then slowly cooling down to ambient temperature. After this annealing step, the synthetically assembled gene can be either cloned directly, or it can be amplified by PCR prior to cloning.

Alternatively, other methods for site directed mutagenesis can be employed for generation of the library insert, such as the Kunkel method (Kunkel T.A. *Rapid and efficient site-specific mutagenesis without phenotypic selection.* *Proc Natl Acad Sci USA.* 1985 January; 82(2):488-92) or the DpnI method (Weiner M P, Costa G L, Schoettlin W, Cline J, Mathur E, Bauer J C. *Site-directed mutagenesis of double-stranded DNA by the polymerase chain reaction.* *Gene.* 1994 Dec. 30; 151(1-2):119-23.).

For various purposes, it may be advantageous to introduce silent mutations into the sequence encoding the library insert. For example, restriction sites can be introduced which facilitate cloning or modular exchange of parts of the sequence. Another example for the introduction of silent mutations is the ability to "mark" libraries, that means to give them a specific codon at a selected position, allowing them (or selected clones derived from them) e.g. to be recognized during subsequent steps, in which for example different libraries with different characteristics can be mixed together and used as a mixture in the selection or panning procedure.

The invention also provides a method of producing an oligomer of modular antibody domains binding to a target comprising the steps of:

providing a library of oligomers of modular antibody domains produced according to the inventive method as described

contacting said library with said target in the presence of a scaffold ligand,
selecting a library member binding to said target in the presence of a scaffold ligand, and
manufacturing a preparation of the functional oligomer.

The scaffold ligand can be selected from the group consisting of an effector molecule, FcRn, Protein A and CDR target. As an example, the effector molecule can be selected from the group consisting of CD64, CD32, CD16, Fc receptors.

The oligomers can be dimers selected from the group of VH/VL, CH1/CL, CH2/CH2, CH3/CH3, Fc and Fab, or single chains thereof

The method according to the invention can provide a library containing at least 10^2 independent clones expressing functional oligomers of modular antibody domains or variants thereof.

Libraries as used according to the invention preferably comprise at least 10^2 library members, more preferred at least 10^3 , more preferred at least 10^4 , more preferred at least 10^5 , more preferred at least 10^6 library members, more preferred at least 10^7 , more preferred at least 10^8 , more preferred at least 10% more preferred at least 10^{10} , more preferred at least 10^{11} , up to 10^{12} members of a library, preferably derived from a parent molecule, which is a functional modular antibody as a scaffold containing at least one specific function or binding moiety, and derivatives thereof to engineer a new binding site apart from the original functional binding region of said parent moiety.

Usually the libraries according to the invention further contain variants of the modular antibody, resulting from mutagenesis or randomization techniques. These variants include inactive or non-functional antibodies. Thus, it is preferred that any such libraries be screened with the appropriate assay for determining the functional effect. Preferred libraries, according to the invention, comprise at least 10^2 variants of such modular antibodies, more preferred at least 10^3 , more preferred at least 10^4 , more preferred at least 10^5 , more preferred at least 10^6 , more preferred at least 10^7 , more preferred at least 10^8 , more preferred at least 10% more preferred at least 10^{10} , more preferred at least 10^{11} , up to 10^{12} variants or higher to provide a highly diverse repertoire of antibodies for selecting the best suitable binders. Any such synthetic libraries may be generated using mutagenesis methods as disclosed herein.

Preferably the library is a yeast library and the yeast host cell exhibits at the surface of the cell the oligomers, or monomers that form oligomers, with the biological activity. The yeast host cell is preferably selected from the genera *Saccharomyces*, *Pichia*, *Hansenula*, *Schizosaccharomyces*, *Kluyveromyces*, *Yarrowia* and *Candida*. Most preferred, the host cell is *Pichia* or *Saccharomyces cerevisiae*.

The invention further provides a high quality library containing at least 10^2 independent clones of functional dimers of modular antibody domains or variants thereof that are binding to a target and to a scaffold ligand. The target can be a ligand binding to a parent molecule subject to amino acid variation. The parent molecule can be a functional oligomer, in particular a functional Pc or a functional Fab, or part thereof.

As is well-known in the art, there is a variety of display and selection technologies that may be used for the identification and isolation of proteins with certain binding characteristics and affinities, including, for example, display technologies such as cellular and non-cellular, in particular mobilized display systems. Among the cellular systems the phage display, virus display, yeast or other eukaryotic cell

display, such as mammalian or insect cell display, may be used. Mobilized systems are relating to display systems in the soluble form, such as in vitro display systems, among them ribosome display, mRNA display or nucleic acid display.

Methods for production and screening of antibody variants are well-known in the art. General methods for antibody molecular biology, expression, purification, and screening are described in Antibody Engineering, edited by Duebel & Kontermann, Springer-Verlag, Heidelberg, 2001; and Hayhurst & Georgiou, 2001, Curr Opin Chem Biol 5:683-689; Maynard & Georgiou, 2000, Annu Rev Biomed Eng 2:339-76.

A library according to the invention may be designed as a dedicated library that contains at least 50% specific formats, preferably at least 60%, more preferred at least 70%, more preferred at least 80%, more preferred at least 90%, or those that mainly consist of specific antibody formats. Such a preferred library mainly contains the same kind of library members having similar structural features. Specific antibody formats are preferred, such that the preferred library according to the invention is selected from the group consisting of a VH library, VHH library, Vkappa library, Vlambda library, Fab library, a CH1/CL library, an Fe library and a CH3 library. Libraries characterized by the content of composite molecules containing more than one antibody domains, such as an IgG library or Fe library are specially preferred. Other preferred libraries are those containing T-cell receptors, forming T-cell receptor libraries. Further preferred libraries are epitope or peptide libraries, wherein the fusion protein comprises a molecule with a variant of an epitope, also enabling the selection of competitive molecules having similar binding function, but different functionality. Exemplary is a TNFalpha library, wherein trimers of the TNFalpha fusion protein are displayed by a single genetic package.

Another important aspect of the invention is that each potential binding domain remains physically associated with the particular DNA or RNA molecule which encodes it, and in addition, the fusion proteins oligomerize at the surface of a genetic package to present the binding polypeptide in the native and functional oligomeric structure. Once successful binding domains are identified, one may readily obtain the gene for expression, recombination or further engineering purposes. The form that this association takes is a "replicable genetic package", such as a virus, cell or spore which replicates and expresses the binding domain-encoding gene, and transports the binding domain to its outer surface. Another form is an in-vitro replicable genetic package such as ribosomes that link coding RNA with the translated protein. In ribosome display the genetic material is replicated by enzymatic amplification with polymerases.

Those cells or viruses or nucleic acid bearing the binding agents which recognize the target molecule are isolated and, if necessary, amplified. The genetic package preferably is M13 phage, and the protein includes the outer surface transport signal of the M13 gene III protein.

Preferably in the method of this invention the vector or plasmid of the genetic package is under tight control of the transcription regulatory element, and the culturing conditions are adjusted so that the amount or number of vector or phagemid particles displaying less than two copies of the fusion protein on the surface of the particle is less than about 20%. More preferably, the amount of vector or phagemid particles displaying less than two copies of the fusion

31

protein is less than 10% the amount of particles displaying one or more copies of the fusion protein. Most preferably the amount is less than 1%.

The expression vector preferably used according to the invention is capable of expressing a binding polypeptide, and may be produced as follows: First a binding polypeptide gene library is synthesized by introducing a plurality of polynucleotides encoding different binding sequences. The plurality of polynucleotides may be synthesized in an appropriate amount to be joined in operable combination into a vector that can be propagated to express a fusion protein of said binding polypeptide. Alternatively the plurality of polynucleotides can also be amplified by polymerase chain reaction to obtain enough material for expression. However, this would only be advantageous if the binding polypeptide would be encoded by a large polynucleotide sequence, e.g. longer than 200 base pairs or sometimes longer than 300 base pairs. Thus, a diverse synthetic library is preferably formed, ready for selecting from said diverse library at least one expression vector capable of producing binding polypeptides having the desired preselected function and binding property, such as specificity.

The foregoing description will be more fully understood with reference to the following examples. Such examples are, however, merely representative of methods of practicing one or more embodiments of the present invention and should not be read as limiting the scope of invention.

EXAMPLES

Example 1

Construction of the Non-Focussed Fcab Library (Fcab01) and Phage Surface Display

The crystal structure of an IgG1 Fc fragment, which is published in the Brookhaven Database as entry 10Q0.pdb was used to aid in the design of the Fcab library.

The sequence which was used as the basis for construction of the Fcab library is given in SEQ ID No. 1. In this sequence, the first amino acid corresponds to Glu 216 of human IgG1 (EU numbering; according to the IMGT database (http://imgt.cines.fr/textes/IMGTrepertoire/Proteins/protein/human/IGH/IGHC/Hu_IGHC.allgenes.html; lookup 2007 06 25), it is the first residue of the human IgG1 hinge region, which is given as: (E) PKSCDKTHTCPPCP) of the heavy constant chain hinge region of human IgG1.) The second-last residue of SEQ ID No. 1 corresponds to Gly 446 of human IgG1 (EU numbering; IMGT: residue number 129 of the CH3 domain of human IgG1).

After detailed analysis of the structure of logo.pdb and by visual inspection of the residues forming the loops which connect the beta strands, it was decided to randomize residues 144, 145 and 146, which are part of the loop connecting beta strand A-B as well as 198, 199, 200, 203 and 204, which are part of the loop connecting beta strand E-F of SEQ ID No.1. In addition to the mutated residues, 5 residues were inserted at residue number 198 of SEQ ID No.1. In SEQ ID No. 2, the sequence of the library insert of library FcabO1 is given in which all randomized residue positions as well as the 5 inserted residues are designated with the letter X.

The engineered gene was produced by a series of PCR reactions using degenerate primers followed by ligation of the resulting PCR products. To facilitate ligation, some of the codons of the nucleotide sequence coding for SEQ ID No. 1 were modified to produce restriction sites without

32

changing the amino acid sequences (silent mutations). For insertion into the cloning vector pHEN1 (Nucleic Acids Res. 1991 Aug. 11; 19(15):4133-7. Multi-subunit proteins on the surface of filamentous phage: methodologies for displaying antibody (Fab) heavy and light chains. Hoogenboom H R, Griffiths A D, Johnson K S, Chiswell D J, Hudson P, Winter G.) in frame with the pelB secretion signal, the NcoI restriction site close to the 3' end of the pelB secretion signal was used. For the randomized residues, the codon NNS (IUPAC code, where S means nucleotides C and G) was chosen which encodes all 20 naturally occurring amino acids, but avoids 2 out of 3 stop codons. Other codons such as for example the NNB (B meaning nucleotides T, C and G) can also be used. The engineered sequence is given as a nucleotide sequence in SEQ ID No. 3. This sequence also includes the restriction sites used for cloning into the phagemid display vector pHEN1, namely an NcoI site at the 5' end and a NotI site at the 3' end.

The sequences of the PCR primers used for assembly of the mutated CH3 domain are given in SEQ ID No. 4 through SEQ ID No. 9.

SEQ ID No. 4 (PCR primer EPKSNCO) ccatggccgagc-
ccaaatcttgtgacaaaactc

SEQ ID No. 5 (PCR primer CH3LSAC) agtcgagctcgt-
cacgggatgggggcaggg

SEQ ID No. 6 (PCR primer CH3CSAC) gtac-
gagctennsnnsnnscaagtcagcctgacctgacctgg

SEQ ID No. 7 (PCR primer CH3CHIN) tgccaagcttgctga-
gaggaagaaggagccg

SEQ ID No. 8 (PCR primer CH3RHIN) tgccaagettaccgt-
gnnnsnnsnnsaggtggnnsnnsgggaacgtcttctcatgctccg

SEQ ID No. 9 (PCR primer CH3RNOT) agttggggccgctttac-
coggagacagggagag

FIG. 1 shows a schematic presentation of the PCR fragments generated for assembly of the mutated gene, and the primers used therefore.

cDNA of the heavy chain of the human monoclonal antibody 3D6 (Felgenhauer M, Kohl J, RUKer F. Nucleotide sequences of the cDNAs encoding the V-regions of H- and L-chains of a human mono-clonal antibody specific to HIV-1-gp41. Nucleic Acids Res. 1990 Aug. 25; 18(16): 4927.) was used as template for the PCR reactions. The 3 PCR products were digested with Sac^I and/or HindIII respectively and ligated together. The ligation product was further digested with NcoI and NotI and ligated into the surface display phagmid vector pHEN1, which had previously been digested with NcoI and NotI. The ligation product was then transformed into *E. coli* by electroporation. A number of selected clones were controlled by restriction analysis and by DNA sequencing and were found to contain the insert as planned, including the correctly inserted randomized sequences. For the following steps of phage preparation, standard protocols were followed. Briefly, the ligation mixture was transformed into *E. coli* TG1 cells by electroporation. Subsequently, phage particles were rescued from *E. coli* TG1 cells with helper phage M13-K07. Phage particles were then precipitated from culture supernatant with PEG/NaCl in 2 steps, dissolved in water and used for selection by panning or, alternatively, they were stored at minus 80° C.

Example 2

Construction of the Focussed Fcab Library (Fcab02) and Phage Surface Display

As described in example 1, an Fcab library was prepared in which the randomized library positions are fully random-

ized, i.e. they are encoded by a codon such as NNS, NNB, NNN, NNN or others are used.

For clarity, the meaning of the letters such as N, B, S or K is defined by the IUPAC nucleotide ambiguity code, which is given in the following table:

TABLE 1

IUPAC nucleotide ambiguity code		
Symbol	Meaning	Nucleic Acid
A	A	Adenine
C	C	Cytosine
G	G	Guanine
T		Thymine
U		Uracil
M	A or C	
R	A or G	
W	A or T	
S	C or G	
Y	C or T	
K	G or T	
V	A or C or G	
H	A or C or T	
D	A or G or T	
B	C or G or T	
X	G or A or T or C	
N	G or A or T or C	

Source: Nomenclature for incompletely specified bases in nucleic acid sequences: recommendations 1984. A Cornish-Bowden, *Nucleic Acids Res.* 1985 May 10; 13(9): 3021-3030.

These codons given above are designed such that all 20 amino acids are encoded by them. It may be preferable to choose subsets out of the possible amino acids. Examples can be found in the literature (Fellouse F A, Li B, Compaan D M, Peden A A, Hymowitz S G, Sidhu S S. Molecular recognition by a binary code. *J Mol Biol.* 2005 May 20; 348(5):1153-62. Epub 2005 Apr. 1.; Fellouse F A, Wiesmann C, Sidhu S S. Synthetic antibodies from a four-amino-acid code: a dominant role for tyrosine in antigen recognition. *Proc Natl Acad Sci USA.* 2004 Aug. 24; 101(34):12467-72. Epub 2004 Aug. 11.). Focused libraries which for example allow for only 4 different amino acid types can be constructed e.g. by employing the codon KMT, which codes for the amino acids Ser, Tyr, Ala and Asp.

A focused Fcab library, designated Fcab02, has been constructed in the same way as described in example 1, except that the NNS codons were replaced by KMT codons.

Therefore, the letter "X" in SEQ ID No. 2 now means "S, Y, A and D" (Ser, Tyr, Ala and Asp) in order to describe the focused library Fcab02

Example 3

Construction of a Phage Surface Display Library with Additional Amino Acid Residues Between the Library Insert (Binding Partner) and p3

In order to investigate accessibility of the potential binding site of the displayed protein a binding assay is performed: the phage suspension is reacted with anti-myc mAb 9E10-coated microplates (or immunotubes). After washing, the bound phages are detected with anti-M13-enzyme conjugate. As a control, helper phage—which does not display the protein fusion and the myc-tag is reacted with the plates. Other controls are reaction of phages with non-coated plates and reaction of phages with antiserum recognizing the p3-fusion partner of the phages.

Ideally, the anti-myc-reactivity of phages displaying the p3-fusion protein should give very clear ELISA readouts

whereas helper phage reactions to anti-myc-mAb should not be above background (non-coated plates). The structure of a CH3 dimer displayed at the surface of an M13 phage through binding to protein III as an anchor is such, that each CH3 is anchored to protein III using various linker length and compositions. Thus, the CH3 dimer is preferably displayed by two anchors.

Linker Optimization:

The linker between the protein to be displayed and the anchor protein of the genetic package (in case of filamentous phage e.g. p3, p8, pX, pIX, pVII) is especially important if the potential binding site of the displayed molecule is in spatial vicinity of the phage particle. In antibody libraries utilizing variable domains and antigen binding sites formed by CDR-loops and display of the library members as amino-terminal fusion to p3 the potential antigen binding site is directed away from the phage particle. Therefore, the linker structure between library members and the phage coat protein is less important. Engineering the bottom loops of immunoglobulin domains and performing phage display may however be an inefficient process and decrease yields of antigen binding clones or even preclude it. Varying the linker between a library member protein and its fusion partner on the surface can solve or may at least reduce this problem.

In order to select for optimal linker sequences (in terms of length and flexibility as well as stability) a library of linkers can be prepared in which the anchor protein at the surface of the genetic replicable package is fused to a known binding protein which is for sterical reasons notoriously difficult to select for.

This library of sequences can be varied in length and amino acid content.

Selection methods of the linker library for optimal linkers depend on the application but basically it should be for selecting all properties one wishes to have in a certain methodology. Enrichment against a difficult to select for antigen may yield linker sequences which allow library members a good access to the antigen. Incubation in protease solutions or under other harsh conditions or frequent passaging through host cells under proteolytic conditions (e.g. old microbial cultures) may be an appropriate selection for stable display linkers.

A library of linkers may be produced by any well known library technology. Synthetic linker sequence lengths may vary between 10-500 amino acids. Alternatively, linker can be complete proteins known to be of flexible nature.

Linker Optimization Fcab01:

As an example, library Fcab01 (as described in example 1) can be used. Originally, this library is cloned in the phagmid display vector pHEN1, using NcoI and NotI restriction sites. When cloned in this manner, 18 amino acid residues are in between the C-terminal amino acid residue of the Fcab01 library insert and the N-terminal amino acid residue of phage M13 p3. The sequence of this junction region is given in SEQ ID No. 10 SPGKAAAEQKLISEEDLNGAATVES—and is explained as follows: the first 4 residues, SPGK (SEQ ID No. 83), are the 4 C-terminal residues of the Fcab01 library insert, followed by the amino acid sequence AAA, which is the amino acid residues encoded by the NotI restriction site, followed by the sequence EQKLISEEDL (SEQ ID No. 84), which is the myc epitope, followed by NGAA (SEQ ID No. 85), after which there is an amber stop codon, which is translated to Glutamine (Q) in amber suppressor strains of *E. coli* such as TG1.

The C-terminal 4 residues of SEQ ID No. 10, TVES (SEQ ID No. 86), are the N-terminal 4 residues of phage M13 p3 as present in the vector pHEN1.

In order to construct a phage which displays an Fcab insert with an increased distance between the Fcab (the binding partner) and the body of the phage (the genetic package), 5 additional residues were inserted at the C-terminus of the Fcab insert FcabRGD4, directly upstream of the NotI cloning site, resulting in the clone FcabRGD4L. FcabRGD4 is an Fcab that has an integrin-binding RGD motif inserted in the EF-loop of the CH3 domain and which binds to $\alpha\beta$ -integrin in ELISA. As an increased-length linker sequence, the amino acid sequence EGGGS (SEQ ID No.89), which appears 8 times in the phage M13 p3 sequence was used. The resulting amino acid sequence of FcabRGD4L as expressed after cloning in pHEN1 is given in SEQ ID No.11. In SEQ ID No.11, amino acid residues 198-204 represent the RGD motif, amino acid residue 237 is the C-terminal residue of the Fcab insert, residues 238-242 represent the inserted linker sequence (which is the difference to unmodified pHEN1), which is followed by myc tag, amber stop codon and the p3 sequence.

For cloning of the construct, the FcabRGD4 sequence was amplified from pHENFcabRGD4 (SEQ ID No. 12) using PCR primers EPKSNCO (SEQ ID No. 4) and CH3rlink actagcggccgcagagccaccacctctaccggagacagggagag (SEQ ID No. 13) and cloned via NcoI and NotI restriction sites into the vector pHEN1. The resulting vector, pHENFcabRGD4L (SEQ ID. No. 14) has the additional linker sequence at nucleotide positions 3057-3071.

The two phagemid vectors, pHENFcabRGD4 and pHENFcabRGD4L were transformed into *E. coli* TG1. Subsequently, phage particles were rescued from *E. coli* TG1 cells with helper phage M13-1K07. Phage particles were then precipitated from culture supernatant with PEG/NaCl in 2 steps, dissolved in water and used for ELISA.

Phage ELISA was performed as follows: The phage suspension is reacted with α V133-integrin-coated microplates (or immunotubes). After washing, the bound phages are detected with anti-M13-enzyme conjugate. As controls, helper phage—which does not display the protein fusion and the myc-tag is reacted with the plates as well as phage particles carrying wtFcab on their surface. Other controls are reaction of phages with non-coated plates and reaction of phages with antiserum recognizing the Fcab-fusion partner of the phages. Phage particles with the increased-length linker resulting from pHENFcabRGD4L react more readily with α v3 integrin than phage particles with the original linker as contained in pHENFcabRGD4, and therefore give a stronger signal in ELISA.

Phage selections can be performed in which phage particles with wtFcab are mixed with small amounts of phage particles carrying either FcabRGD4 or FcabRGD4L. After several (typically 3-5) rounds of panning, preferentially phages displaying FcabRGD4L are selected.

Example 4

Fcab Library Design

("Fcab" is a registered trademark of f-star Biotechnologische Forschungs- und Entwicklungsges.m.b.H.)

Design of Fcab Libraries (illustrated in FIG. 2): amino acid positions in non CDR-loops of CH3 constant domains of antibodies are considered for randomization. Especially loops A-B, C-D and E-F are considered as they are on one

side of the domain. Some of the design criteria for randomization at a certain position are described herein.

Amino acids frequently involved in antigen antibody interactions are described herein to be included in a focused library. Here the amino acids Ala, Asp, Ser and Tyr are used to design the focused library.

Libraries with restricted amino acid utilization have been shown to be sufficient to generate binders against virtually any antigen (Sidhu & Fellhouse, NATURE CHEMICAL BIOLOGY VOLUME 2 page 682ff; Koide et al PNAS, volume 104 p6632-6637). The advantage of such restricted (or focused) libraries is that they can be covered completely by current technologies. Ideally, the amino acid utilization reflects a natural amino acid utilization of ligand receptor binding. However, even libraries utilizing only 2 amino acids (Tyrosine and Serine) have been reported to yield good selection results (in terms of frequency of binders against different binders and in terms of affinity).

Loop Flexibility:

Certain loop structures may be required by the scaffold protein in order to keep the overall natural structure. Randomizing many amino acid positions in loops and even elongation of loops may be facilitated by building certain sequences either on one or on both sides of the randomized positions. These sequences may be flexible sequences in order to allow to compensate for any tensions with certain library sequences in such a position.

TABLE 2

Exemplary Fcabm libraries, focused and non-focused			
	# of randomized positions	Theoretical diversity on amino acid level	Number of independent bacterial clones
Fcab01	13	8.2×10^{16}	0.6×10^9
Fcab02	13, focused	6.7×10^7	0.6×10^9
Fcab03	13	8.2×10^{16}	1.0×10^9
Fcab04	13, focused	6.7×10^7	0.8×10^9
Fcab05	15	1.3×10^{18}	0.8×10^9
Fcab06	15, focused	1.3×10^9	1.0×10^9

Fcab01 library is described in the examples above. The sequence space of the focused library designs Fcab02, Fcab04 and Fcab06 are covered by the actual bacterial library sizes of approximately 10^9 . In contrast, the completely randomized libraries Fcab01, Fcab03 and Fcab05 are actually grossly underrepresented.

Example 5

Cloning of Yeast Display Libraries by Homologous Recombination

Vector

pYD1 (Invitrogen) is used as the basic vector. The vector is modified as follows, in order to remove an XhoI site: pYD1 is cleaved with XhoI, treated with Klenow fragment of DNA polymerase and religated. The resulting sequence is given in pYD1dX (SEQ ID No. 15).

pYD1dX contains a unique BamEI restriction site at position 921/925 and a unique NotI restriction site at position 963/967. It is opened with these two restriction enzymes.

An insert encoding CH1-hinge-CH2-CH3 from human IgG1 is prepared by PCR from cDNA encoding the heavy chain of a human IgG1 monoclonal antibody. In this insert, a point mutation is introduced using standard procedures to

mutate the C-terminal Cystein residue of the CH1 domain to a Serine. The insert is amplified using PCR primers that attached a BamHI and a Not restriction site to both ends respectively. These restriction sites are then used for cloning the insert into pYD1dX to yield the display vector pYD1dXFc (SEQ ID No. 16). The mutated codon at the C-terminus of the CH1 domain (Cys to Ser) is at positions 1233-1235 in the sequence pYD1dXFc. The stop codon of the insert is at position 1917/1919.

This vector is used as a positive control for the display of human CH1-hinge-CH2-CH3 on the surface of yeast and as a starting point for the construction of the vector pYD1CH12 (see below).

Cloning of Libraries

Cloning of libraries in which mutations are introduced into structural loops of CH3 domains is performed in yeast by homologous recombination (gap repair). For this purpose, a recipient vector is prepared that lacks the CH3 domain: pYD1dXFc is cleaved with XhoI (position 1603/1607) and NotI (position 1921/1925), the large fragment is prepared by preparative gel electrophoresis, treated with Klenow fragment of DNA polymerase and re-ligated. This procedure reconstitutes a unique XhoI site (position 1603/1607) and yielded vector pYD1CH12 (SEQ ID No. 17). pYD1CH12 is subsequently cleaved with XhoI and is used as recipient vector for gap repair in yeast.

As a source of insert, Fcab libraries Fcab01 (SEQ ID No. 18), Fcab02 (SEQ ID No. 19), Fcab03 (SEQ ID No. 20), Fcab04 (SEQ ID No. 21), Fcab05 (SEQ ID No. 22) and Fcab06 (SEQ ID No. 23) are used. These libraries are prepared by standard DNA synthesis, and contain randomized residues as well as inserted residues in the AB loop (between residues 359 and 361 (EU numbering)) as well as in the EF loop (between residues 413 and 419 (EU numbering)) of the CH3 domain of human IgG1. From this synthetic DNA, the insert for gap repair in yeast is amplified by PCR using PCR primer pair gapch35 caacaaggccctgcctgcccccat cgagaagaccatctccaaggccaaggccagcctcgagaac-cacaggtgtacacctgccc (SEQ ID No. 24) and gapfcs3 gagac-cgaggagagggttagggataggcttacct tcgaaggccctctagactcgatcgagcggccgctcatttaccggagacagggagagctc ttc (SEQ ID No. 25). 100 pg of XhoI cleaved vector pYD1CH12 and 100 pg of insert are mixed and transformed in *Saccharomyces* strain EBY100 (Invitrogen) using the Lithium acetate procedure according to the following protocol, which is upscaled by a factor 100 to transform the required amount of cells and of DNA. Briefly, for a single transformation of 1 pg vector DNA and 1 pg insert DNA, 10 ml of YPD (2% peptone, 2% dextrose (D-glucose)) are inoculated with a yeast colony and shaken overnight at 30° C. The OD600 of the overnight culture is determined and the culture diluted to an OD600 of 0.4 in 50 ml of YPD and grown for an additional 6 hours. Cells are pelleted at 2500 rpm and resuspended in 40 ml distilled water. Cells are pelleted again at 2500 rpm and resuspended in 100 mM LiAc, followed by incubation at 30° C. for 30 minutes. 1 pg vector DNA, 1 pg insert and 50 pl denatured sheared salmon sperm DNA (2 mg/ml) are mixed with 300 pl of the yeast suspension. 700 pl of a solution of 200 mM Li-acetate and 40% PEG-3350 are added and mixed with the yeast/DNA suspension, followed by incubation at 30° C. for 30 minutes. 88 pl DMSO are added, mixed and the mixture is incubated at 42° C. for 40 minutes, followed by centrifugation in a microcentrifuge for 10 seconds. The supernatant is then removed, the cell pellet is resuspended in 1 ml distilled water. The pellet is then resuspended in 50-100 pl TE and plated on minimal dextrose plates containing leucine

(10 g/l yeast nitrogen base, 20 g/l dextrose, 0.1 g/l leucine, 15 g/l agar). After incubation of the plates at 30° C. for 2 to 4 days single colonies appeared that are subsequently harvested.

Cultivation—Induction

The harvested yeast libraries (yFcab libraries) are inoculated in 10 ml SD-CAA medium (10 g/l yeast nitrogen base, 10 g/l casamino acids, and 20 g/l dextrose, 0.1 g/l leucine, 9.67 g/l NaH₂PO₄-2H₂O and 10.19 g/l Na₂HPO₄-7H₂O) and grown on a shaker at 250 rpm at 28° C. for 6-8 hours. The OD600 of the culture is determined, and the culture is diluted to an OD600 of 0.2, and grown under the same conditions until an OD600 of 1-2 is reached. Cells are harvested by centrifugation (3000 rpm 5 min/4° C.) and resuspended in induction medium SG/R-CAA (10 g/l yeast nitrogen base, 10 g/l casamino acids, and 20 g/l galactose, 10 g/l raffinose, 0.1 g/l leucine, 9.67 g/l NaH₂PO₄-2H₂O and 10.19 g/l Na₂HPO₄-7H₂O). Cultures are induced by incubation for 2 days on a shaker at 250 rpm at 20° C. and subsequently analysed and sorted.

Quality Control of yFcab Libraries.

yFcab libraries are tested for their expression level and quality of expressed Fcab's two days after induction with SD-CAA medium. The expression level is tested using a polyclonal anti human IgG-Fc antiserum (Sigma). For this purpose 0.5×10⁶ library cells are diluted in 1 ml staining buffer (SB), which comprises PBS with 2% BSA. Cells are pelleted and stained with 100 pl SB containing 1/2000 diluted anti human IgG-Pc-PE antiserum (Sigma) for 30 min on ice, washed twice with SB and subsequently analyzed in the FACS. In general 70%-80% of all cells in each library express Fcabs on their cell surface. To test correct folding of Fcabs, staining with Protein A is performed. Again 0.5×10⁶ library cells are diluted in 1 ml staining buffer SB, cells are pelleted and stained with 100 pl SB containing 1 pg/ml Prot-A-FITC (Fluka) for 30' on ice, washed twice with SB and subsequently analyzed in the FACS. In general, the yFcab libraries as described above show >40% Prot A positive cells. In order to test whether the Fcabs are expressed as dimers on the surface of the cells a staining with human CD64 is performed. The affinity of CD64 is too low for efficient monomeric binding therefore CD64 complexes with dimers are used. For this purpose e.g. lug recombinant CD64 from R&D Systems (containing a HIS-tag) is mixed with 1 ug anti Penta HIS-alexafluor 488 (from Qiagen) in 1 ml SB (total volume). yFcab libraries are tested for binding to CD64 by incubating the 5×10⁵ cells with 100 pl of the complex-mixture for 30 minutes on ice, as control the cells are incubated with equivalent of the anti HIS-alexafluor 488 alone (1/200 dilution in SB). After incubation the cells are washed twice with ice cold SB and analysed in the FACS. In general >50% of all cells in each library express dimeric Fcabs on their cell surface.

Biotinylation of Antigen (her2)

Recombinant antigen e.g. Her2 (Bendermedsystems) was done with the EZ link system of Pierce according to the manufacturers instruction. In short, the antigen is dialyzed against PBS, diluted to 1 mg/ml in PBS and mixed with 10 ml sulfo-LC-LC⁻ biotin (EZ link, Pierce), which was pre-dissolved in water. The final ratio between antigen and biotin is 1:3 and the mixture is incubated at room temperature from 30'. Afterwards the mixture is "dialyzed" against PBS using Vivaspin MWC03000 (Sartorius) columns (5×8', 4000 rpm). Finally the concentration of the biotinylated antigen (Her2) is tested by HPLC and aliquots are stored at -20° C.

The quality of the biotinylated antigen is tested by ELISA. First the plates are coated with an anti-Her2 antibody (e.g. Herceptin) at 10 pg/ml in PBS, 100 pl/well overnight at 4° C., after this the plate is washed 3× with washing buffer (WB)(PBS+0.05% Tween20) and blocked by blocking buffer (BB) (PBS+2% BSA) 1 h at room temperature. After 3× washing with WB, different concentrations of Her2-biotin are added in 100 pl/well BB for 1 h at room temperature, followed by 3× washing with WB. Finally the plate is incubated with 1:25000 streptavidin-HRP (GE healthcare) in BB for 1 h at room temperature and washed 3× with WB. Colour is developed by adding 100 pl/well of the substrate TMB (Sigma) after -10 minutes the reaction is stopped by adding 100 pl/well of 30% H2504. The results is analysed with an ELISA reader at 450-630 nm.

Example 6

Production of Antigen Specific (her2) Fcabs

Selection of antigen specific (Her2) Fcabs using FACS First Selection Round:

Two days before FACS sorting a yeast library containing 2.5×10^8 individual Fcab clones is induced with SG/R-CAA medium to express the Fcabs on their cell surface as described above. After two days, the amount of cells covering e.g. 10 times the library ($=2.5 \times 10^9$) is incubated for 30 minutes on ice with 500 nM biotinylated antigen (Her2) in 201 SB. Then the cells are washed once with cold SB and subsequently incubated for 30' on ice with streptavidin-PE (from R&D systems) diluted 1:100 in SB. The cells are washed twice with ice cold SB and diluted to an end concentration of 1×10^9 cells/ml. Control stainings with 5×10^6 cell/ml in 100 pl are made with streptavidin-PE only, in the absence of antigen. Both the complete library and the control stainings are analysed in e.g. a FACS ARIA from BD. To set the gates for sorting the control cells are used. First a FSC/SSC gate (G1) is set to identify healthy yeast cells, from G1 a FSC-width versus FSC-area plot is made and only non-aggregating cells are selected in a new gate (G2). Cells in G2 are subsequently analysed for reactivity with streptavidin-PE using FSC versus FL-2 (PE channel). G3 is set to include 0.1% of (false) positive cells. Subsequently, at least 5×10^8 stained cells (twice the library size ideally more) are analysed with the settings as indicated above and the cells in G3 are sorted into a tube containing 2-3 ml SD-CAA medium. Roughly 5×10^5 cells (Pooh) are harvested in the first round of selection and propagated for 1 to 2 days, after which the cells can be stored at -80° C. and aliquots can be induced to express the Fcabs as described above. After two more days the next selection round can take place.

Second Selection Round:

Pooh 1 selected in round 1 are induced to express the Fcab on their surface as described above. At least 5×10^6 cells (comprising multiple copies of Pooh) are incubated for 30' on ice with 500 nM biotinylated antigen (Her2) in 1 ml SB. Then the cells are washed once with cold SB and subsequently incubated for 30' on ice with streptavidin-PE (from R&D systems) diluted 1 in 100 in SB together with 2 g/ml Protein A-FITC (Fluka). Next the cells are washed twice with ice cold SB and diluted to an end concentration of 2×10^6 cells/ml. In addition, control stainings are made in which 5×10^6 cells/ml of Pooh 1 in 100 pl cells are incubated with a mixture of Prot A and streptavidin-PE as indicated above, but without the incubation with the antigen (Her2). In addition, 5×10^5 cell in 100 pl of a yeast clone expressing

Fcab wt non randomized Pc fragment) is stained with Prot A-FITC as described above in the absence of streptavidin-PE. Fcab-wt expressing cells are analysed in e.g. a FACS ARIA from BD to set gates for sorting. First a FSC/SSC gate (G1) is set to identify healthy yeast cells, from G1 a FSC-width versus FSC-area plot is made and only non aggregating cells are selected in new gate (G2).

Cells in G2 are subsequently analysed for Protein A expression using FSC versus FL-1 (FITC). G3 is set to cover strong Prot A positive cells (50-60% of parent gate) and G4 is set to cover weak Prot A positive cells (20-30% of parent cells). G3+G4 will include roughly 70-80% of all cells in G2. Now the Pool cells stained for streptavidin-PE in the presence of Prot A-FITC are used to set the rest of the sorting gates. First G1 and G2 are checked with the Pool cells and if necessary adjusted. Pool cells will have lesser events in G3 and maybe also in G4 indicating that not all cells in Pooh 1 express Fcabs that are folded as the Fcab-wt. Using the control stained Pool cells a new gate is prepared both for G3 and G4. The new gates are set in a plot FSC and FL-2 (PE). Gate (G5) is prepared that includes 0.1% (false) streptavidin positive cells in G3 and the same is done for cells in G4 resulting in G6. In the next step at least 5×10^6 cells stained for Her2-biotin+streptavidin-PE and Prot A-FITC are sorted by the FACS-ARIA. Cells are collected from G5 (Poo12.1 and G6 (Poo12.2) in separate tubes containing 2-3 ml yeast culture medium. Between 10 and 1000 clones can be expected from both gates. Both new pools are propagated for 1 or 2 days and stored at -80° C. Cells from 2.1 and 2.2 may be either used for direct further sorting in a third round or they may be subjected, (preferably after mixing the two clone together again) to a round of additional randomization of the AB loop (affinity maturation) before they are further sorted in FACS.

Affinity Maturation for Selected Clones/Pools

For affinity maturation, diversity is introduced in selected clones or in pools of selected clones in the AB loop. For this purpose, a PCR is made with a primer that contained degenerate codons at positions 359, 360 and 361 (EU numbering) (primer Abmut, gaaccacaggtgtacacctgocccatccoggatgagctgnbnnbnnbca ggtcagctgacctgcc tggtaaaag, SEQ ID No. 26) or alternatively with a primer that contained degenerate codons at positions 358, 359, 360, 361 and 362 (EU numbering) (primer Abmut2LR, gaaccacaggtgtacacctgccccatcccggtgaggnbnnbnnbnnbnnbgtcagcctgacctgctgtcaaaag, SEQ ID No. 27). The second primer used in these PCRs is gapfcs3 in both cases. In order to create flanking sequences for efficient gap repair in yeast, the resulting PCR products are further amplified with the primer pair gapch35 and gapfcs3 and subsequently transformed in *Saccharomyces cerevisiae* strain EBY100 by Lithiumacetate transformation together with XhoI cleaved pYD1CH12 as described above. As alternative primers for randomization of the described residues in the AB loop, primers such as AbmutIL (gaaccacaggtgtacacctgccccatcccggtgaggnbnnbnnbnnbcaggtcagc ctgacctgctgtgtcaaaag, SEQ ID No. 28) or AbmutILR (gaaccacaggtgtgta caccctgccccatcccggtgagctgnbnnbnnbnnbnnbgtcagcctgacctgctgtgtaaaag, SEQ ID No. 29) can also be used. In an analogous manner, residues in the EF loop can be randomized e.g. by total randomization or by randomization using spiked oligonucleotides as primers or by similar mutagenesis techniques. The Abmut primer will result in 8000 new variants (Poo12.3) of each clone and the Abmut2LR primer will lead to 3×10^6 new variants (Poo12.4). Therefore Pools 2.3. and 2.4 will both results in

41

new libraries of approximately 10e8 individual since the starting material (P0012.1+2.2) already contains approximately 10-1000 clones.

Third Selection Round

Affinity matured pools 2.3 and 2.4 and if necessary 5 P0012.1 (only the Prot A positive cells are preferred) are induced to express Fcabs on their cell surface as described above and subsequently sorted as described for "Second selection round", with exception that the Pools 2.3 and 2.4 are much bigger and therefore staining volumes for the pools are equal to those of the library staining described in "First selection round". In the third selection round, only Her2 positive/Prot A positive cells are sorted. Pools derived from these selections contain typically >20% Her2/Prot A positive cells. If not then a fourth and fifth (or even more) round(s) of selection combining Prot A with Her2 can be performed. Clone Analyses:

Individual clones from pools containing Her2/Prot A cells (>20% is preferred) are prepared either by plating the pools on agar plates with SD-CAA medium or by sorting the singles cells (=clones) directly from the FACS ARIA onto the plates without generating a pool. Clones are allowed to grow and are transferred to liquid cultures and stored in -80° C. Aliquots of the clones are subsequently induced to express Fcabs on their cell surface as described above and screened for a number of parameters in the FACS. These parameters may be: a dose response range of the antigen used for selection (Her2) with and without the presence of Prot A-FITC, CD64 staining as described above. In addition using similar staining protocols a number of irrelevant biotinylated antigen can be screened to identify non-cross reacting Fcabs.

It is to be expected that, after several rounds of selecting antigen (Her2)+Prot A positive cells, a large percentage of clones show >25% antigen (Her2) positivity when stained with 500 nM antigen (Her2) and >70% Prot A positivity when stained with 2 pg/ml Prot A-FITC. In most of the cases these clones will also show >50% CD64 binding. Thus mimicking the Prot A and CD64 levels of non-randomized Pc fragments (Fcab wt) expressed on yeast.

Clones selected as described above with characteristics as described above are now ready to be produced as soluble molecules. This can be done by transient transfection or by stable transfection of the Fcab DNA into new host cells. For this purpose the DNA from individual yeast clones is isolated using standard procedures. The relevant DNA coding for the complete CH3 domain or only the part of the CH3 domain that is randomized in the library is amplified by PCR and transferred into a new expression vector containing the missing part of the Fcab+a suitable promoter and one of more selection markers such as G418, that allows selection of transfected cells out of a pool of non transfected cells. The new vector is then e.g. transiently transfected into a new host cell such as HEK293 or CHO. The host cells are allowed to recover and are subsequently cultured for a number of days. The supernatant of the cultures with contain the soluble Fcab which can be used for further testing with or without purification over e.g. Prot A. Stable cell lines can also be made by standard procedures.

42

TABLE 2

Sequences of selected Her2 clones: with reference to numbering of Seq. ID No. 1		
Clone name	AB loop AA143ff	EF Loop AA198ff
Fcab wt	LTKNQ (SEQ ID No. 87)	-----DKSRWQQ (SEQ ID No. 88)
y ⁻ Her.C2- P3.1-1	LDNSQ (SEQ ID No. 30)	IRSSVGSRRWWS (SEQ ID No. 51)
y ⁻ Her.C2- P3.1-3	YEGSS (SEQ ID No. 31)	ARYSPRMLRWAH (SEQ ID No. 52)
y ⁻ Her.C2- P3.1-5	YMSAD (SEQ ID No. 32)	SRDSSLLRWAH (SEQ ID No. 53)
y ⁻ Her.C2- P3.1-6	YRRGD (SEQ ID No. 33)	APGSKGYRRWAL (SEQ ID No. 54)
y ⁻ Her.C2- P3.1-8	LMSRQ (SEQ ID No. 34)	DKPFWGTSRWSR (SEQ ID No. 55)
y ⁻ Her.C2- P3.1-16	LHLAQ (SEQ ID No. 35)	SINDLINHRWPY (SEQ ID No. 56)
y ⁻ Her.C2- P3.1-18	YLSKD (SEQ ID No. 36)	MWGSRDYWRWSH (SEQ ID No. 57)
y ⁻ Her.C2- P3.2-3	YRSGS (SEQ ID No. 37)	NSGSAMMVRWAH (SEQ ID No. 58)
y ⁻ Her.C2- P3.2-9	LRDGQ (SEQ ID No. 38)	QRSRLSRQRWWR (SEQ ID No. 59)
y ⁻ Her.C2. P4.2-1	YSANT (SEQ ID No. 39)	ARYSPRMLRWAH (SEQ ID No. 60)
y ⁻ Her.C2. P4.2-3	YASNT (SEQ ID No. 40)	ARYSPRMLRWAH (SEQ ID No. 61)
y ⁻ Her.C2. P4.2-4	YSDGD (SEQ ID No. 41)	ARYSPRMLRWAH (SEQ ID No. 62)
y ⁻ Her.C2. P4.2-5	YSGGS (SEQ ID No. 42)	ARYSPRMLRWAH (SEQ ID No. 63)
y ⁻ Her.C2. P4.2-6	YGRDS (SEQ ID No. 43)	ARYSPRMLRWAH (SEQ ID No. 64)
y ⁻ Her.C2. P4.2-8	YAGGT (SEQ ID No. 44)	ARYSPRMLRWAH (SEQ ID No. 65)
y ⁻ Her.C2. P4.2-10	YSSDS (SEQ ID No. 45)	ARYSPRMLRWAH (SEQ ID No. 66)
y ⁻ Her.C2. P4.2-12	YHSGS (SEQ ID No. 46)	ARYSPRMLRWAH (SEQ ID No. 67)
y ⁻ Her.C2. P4.2-15	YLTNS (SEQ ID No. 47)	ARYSPRMLRWAH (SEQ ID No. 68)
y ⁻ Her.C2. P4.2-18	YGSEE (SEQ ID No. 48)	ARYSPRMLRWAH (SEQ ID No. 69)
y ⁻ Her.C2. P4.2-19	YRSGE (SEQ ID No. 49)	ARYSPRMLRWAH (SEQ ID No. 70)
y ⁻ Her.C2. P4.2-20	YGTDD (SEQ ID No. 50)	ARYSPRMLRWAH (SEQ ID No. 71)

43

Example 7

Yeast Display of 4D5 Fab

For the display of a Fab fragment on yeast, the yeast display vector pYD1 (Invitrogen) (SEQ ID No. 72) is modified as follows:

A NheI restriction site is introduced by site directed mutagenesis at position 581/586 to yield the modified vector pYD1Nhe (SEQ ID No. 73). This vector is restricted with NheI and PmeI, to yield 3 fragments. The largest fragment is the remaining vector backbone, in which a synthetic oligonucleotide linker is inserted to yield the vector pYD1lnk (SEQ ID No. 74). A cassette which includes the MATa transcription termination region is then amplified by PCR from the vector pYD1 and is cloned into pYD1lnk via BamHI and PstI restriction and ligation. The resulting vector is pYD1mata (SEQ ID No. 75). A cassette that contains the GAL1 promoter, the gene coding for Aga2 and a synthetic linker with NotI and SfiI cloning sites is amplified by PCR from pYD1 and cloned in pYD1mata via EcoRI and Pad restriction to yield the vector pYD1gal (SEQ. ID No. 76).

As an example for a Fab to be displayed on yeast, the genes coding for VH-CH1 and VL-CL respectively of the antibody 4D5 (Herceptin) are made synthetically (sequences 4D5H (SEQ ID No. 77) and 4D5L (SEQ ID No. 78)).

4D5H is flanked by SfiI and NotI restriction sites, and cloned into the vector pYD1gal to yield the vector pYD4D5hc (SEQ ID No. 79). In this vector, the N-terminus of 4D5H is fused to the C-terminus of Aga2, and at the C-terminus of 4D5H, a hexahistidine tag is attached, followed by the stop codon. The amino acid sequence of VH-CH1 of 4D5 is given in 4D5 hp (SEQ ID No. 80).

4D5L is flanked by NcoI and Ascl restriction sites, and cloned into the vector pYD4D5hc to yield the vector pYD4D5h1 (SEQ ID No. 81). 4D5L is preceded by an Aga2 secretion signal, and carries a stop codon after the C-terminal Cysteine residue of the CL domain. The amino acid sequence of VL-CL of 4D5 is given in 4D51p (SEQ ID No. 82).

44

For display of the 4D5 Fab, the vector pYD4D5h1 is transformed into the yeast strain EBY100 (Invitrogen), transformants are selected on minimal medium without tryptophan, and expression of the recombinant protein is induced by growth on galactose containing medium according to standard protocols (Invitrogen).

Example 8

Construction of a Library with Randomized Residues in Structural Loops of the CL Domain of 4D5 Fab

As first step in the yeast display library construction, the wildtype CL (C kappa) domain is cut out from the display vector pYD4D5h1 with restriction enzymes BsiWI and Ascl. A synthetic gene encoding human C kappa domain flanked by BsiWI and Ascl sites (in the context according to pYD4D5h1) is prepared in which random mutations and insertions respectively are introduced in the AB and EF loops. In this particular example, insertions of 3, 4 or 5 NNB codons are made between amino acid positions 16 and 17 of the human C kappa domain, and residue positions 92, 93, 94, 95, 97, 98 and 99 are replaced by NNB codons. (IMGT numbering, see FIG. 2). An NNB codon contains all 4 nucleotides at positions 1 and 2, and C, G and T at position 3. NNB therefore encodes all 20 naturally encoded amino acids.

The library is prepared and selected following standard procedures.

As a scaffold ligand the CDR target Her2neu and 4D5 epitope is used. Those members of the library are selected for production of a cytotoxic modular antibody according to the invention, that have a binding site engineered into the CL domain, which is specifically binding to an effector molecule, such as an Fcgamma receptor, or a half-life prolonging protein, such as serum albumin. The resulting Fab is tested for (i) Her2neu binding with a $K_d < 10^{-8}$ M and an $1050 < 10^{-8}$ M, and (ii) effector function using a CDC and/or ADCC assay, and alternatively to determine albumin binding.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 96

<210> SEQ ID NO 1

<211> LENGTH: 232

<212> TYPE: PRT

<213> ORGANISM: human

<400> SEQUENCE: 1

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
1 5 10 15

Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
20 25 30

Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
35 40 45

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
50 55 60

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
65 70 75 80

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
85 90 95

-continued

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
100 105 110

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
115 120 125

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr
130 135 140

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
145 150 155 160

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
165 170 175

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
180 185 190

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
195 200 205

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
210 215 220

Ser Leu Ser Leu Ser Pro Gly Lys
225 230

<210> SEQ ID NO 2

<211> LENGTH: 237

<212> TYPE: PRT

<213> ORGANISM: artificial

<220> FEATURE:

<223> OTHER INFORMATION: human IgG including randomized amino acid
modifications

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (144)..(146)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (198)..(205)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (208)..(209)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 2

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
1 5 10 15

Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
20 25 30

Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
35 40 45

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
50 55 60

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
65 70 75 80

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
85 90 95

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
100 105 110

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
115 120 125

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Xaa
130 135 140

Xaa Xaa Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser

-continued

145	150	155	160
Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr			
	165	170	175
Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr			
	180	185	190
Ser Lys Leu Thr Val Xaa Xaa Xaa Xaa Xaa Xaa Xaa Arg Trp Xaa			
	195	200	205
Xaa Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn			
	210	215	220
His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys			
225	230	235	

```

<210> SEQ ID NO 3
<211> LENGTH: 728
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: DNA Sequence for Cloning of modified IgG
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (438)..(439)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (441)..(442)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (444)..(445)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (600)..(601)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (603)..(604)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (606)..(607)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (609)..(610)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (612)..(613)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (615)..(616)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (618)..(619)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (621)..(622)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (630)..(631)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (633)..(634)
<223> OTHER INFORMATION: n is a, c, g, or t
<400> SEQUENCE: 3

```

-continued

```

ccatggccga gcccaaatct tgtgacaaaa ctcacacatg cccaccgtgc ccagcacctg      60
aactcctggg gggaccgtca gtcttctctt tcccccaaa acccaaggac accctcatga      120
tctcccgga ccttgaggtc acatgcgtgg tggtaggacgt gagccacgaa gacctgagg      180
tcaagttcaa ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca aagccgcggg      240
aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg caccaggact      300
ggctgaatgg caaggagtac aagtgaagg tctccaacaa agccctccca gcccctatcg      360
agaaaaccat ctccaaagcc aaagggcagc ctcgagaacc acaggtgtac accctgcccc      420
catcccgta cgagctcnns nnsnnscaag tcagcctgac ctgcctggtc aaaggcttct      480
atcccagcga catgcgcgtg gaggtaggaga gcaatgggca gccggagaa aactacaaga      540
ccacgcctcc cgtgctggac tccgacggct ctttcttct ctacagcaag cttaccgtgn      600
nnsnnsnnn snnsnnsnns nnsaggtggn nsnsgggaa cgtcttctca tgctccgtga      660
tgcatgaggc tctgcacaac cactacacac agaagagcct ctccctgtct ccgggtaaaag      720
cggccgca                                           728

```

```

<210> SEQ ID NO 4
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer EPKSNCO

```

```

<400> SEQUENCE: 4

```

```

ccatggccga gcccaaatct tgtgacaaaa ctc                                           33

```

```

<210> SEQ ID NO 5
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer CH3LSAC

```

```

<400> SEQUENCE: 5

```

```

agtcgagctc gtcacgggat gggggcaggg                                           30

```

```

<210> SEQ ID NO 6
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer CH3CSAC
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (11)..(12)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (14)..(15)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (17)..(18)
<223> OTHER INFORMATION: n is a, c, g, or t

```

```

<400> SEQUENCE: 6

```

```

gtacgagctc nnsnnsnsc aagtcagcct gacctgcctg g                                           41

```

```

<210> SEQ ID NO 7
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: artificial

```

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: PCR primer CH3CHIN

<400> SEQUENCE: 7

tgccaagctt gctgtagagg aagaaggagc cg                               32

<210> SEQ ID NO 8
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer CH3RHIN
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (17)..(18)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (20)..(21)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (23)..(24)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (32)..(33)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (35)..(36)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 8

tgccaagctt accgtgnnsn nsnnnsaggtg gnnsnnsggg aacgtcttct catgctccg       59

<210> SEQ ID NO 9
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer CH3RNOT

<400> SEQUENCE: 9

agttgcggcc gctttaccgc gagacaggga gag                               33

<210> SEQ ID NO 10
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: junction region

<400> SEQUENCE: 10

Ser Pro Gly Lys Ala Ala Ala Glu Gln Lys Leu Ile Ser Glu Glu Asp
1          5          10          15

Leu Asn Gly Ala Ala Thr Val Glu Ser
          20          25

<210> SEQ ID NO 11
<211> LENGTH: 662
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: amino acid sequence of FcabRGD4L

<400> SEQUENCE: 11

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
1          5          10          15

```

Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Lys	Pro
Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val
Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr
Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu
Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His
Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys
Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln
Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu
Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro
Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn
Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu
Ser	Lys	Leu	Thr	Val	Gly	Cys	Arg	Gly	Asp	Cys	Leu	Ser	Arg	Trp
Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His
His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys	Glu	Gly
Gly	Ser	Ala	Ala	Ala	Glu	Gln	Lys	Leu	Ile	Ser	Glu	Glu	Asp	Leu
Gly	Ala	Ala	Thr	Val	Glu	Ser	Cys	Leu	Ala	Lys	Pro	His	Thr	Glu
Ser	Phe	Thr	Asn	Val	Trp	Lys	Asp	Asp	Lys	Thr	Leu	Asp	Arg	Tyr
Asn	Tyr	Glu	Gly	Cys	Leu	Trp	Asn	Ala	Thr	Gly	Val	Val	Val	Cys
Gly	Asp	Glu	Thr	Gln	Cys	Tyr	Gly	Thr	Trp	Val	Pro	Ile	Gly	Leu
Ile	Pro	Glu	Asn	Glu	Gly	Gly	Gly	Ser	Glu	Gly	Gly	Gly	Ser	Glu
Gly	Gly	Ser	Glu	Gly	Gly	Gly	Thr	Lys	Pro	Pro	Glu	Tyr	Gly	Asp
Pro	Ile	Pro	Gly	Tyr	Thr	Tyr	Ile	Asn	Pro	Leu	Asp	Gly	Thr	Tyr
Pro	Gly	Thr	Glu	Gln	Asn	Pro	Ala	Asn	Pro	Asn	Pro	Ser	Leu	Glu
Ser	Gln	Pro	Leu	Asn	Thr	Phe	Met	Phe	Gln	Asn	Asn	Arg	Phe	Arg
Arg	Gln	Gly	Ala	Leu	Thr	Val	Tyr	Thr	Gly	Thr	Val	Thr	Gln	Gly
Asp	Pro	Val	Lys	Thr	Tyr	Tyr	Gln	Tyr	Thr	Pro	Val	Ser	Ser	Lys

-continued

Met Tyr Asp Ala Tyr Trp Asn Gly Lys Phe Arg Asp Cys Ala Phe His
435 440 445

Ser Gly Phe Asn Glu Asp Pro Phe Val Cys Glu Tyr Gln Gly Gln Ser
450 455 460

Ser Asp Leu Pro Gln Pro Pro Val Asn Ala Gly Gly Gly Ser Gly Gly
465 470 475 480

Gly Ser Gly Gly Gly Ser Glu Gly Gly Gly Ser Glu Gly Gly Gly Ser
485 490 495

Glu Gly Gly Gly Ser Glu Gly Gly Gly Ser Gly Gly Gly Ser Gly Ser
500 505 510

Gly Asp Phe Asp Tyr Glu Lys Met Ala Asn Ala Asn Lys Gly Ala Met
515 520 525

Thr Glu Asn Ala Asp Glu Asn Ala Leu Gln Ser Asp Ala Lys Gly Lys
530 535 540

Leu Asp Ser Val Ala Thr Asp Tyr Gly Ala Ala Ile Asp Gly Phe Ile
545 550 555 560

Gly Asp Val Ser Gly Leu Ala Asn Gly Asn Gly Ala Thr Gly Asp Phe
565 570 575

Ala Gly Ser Asn Ser Gln Met Ala Gln Val Gly Asp Gly Asp Asn Ser
580 585 590

Pro Leu Met Asn Asn Phe Arg Gln Tyr Leu Pro Ser Leu Pro Gln Ser
595 600 605

Val Glu Cys Arg Pro Tyr Val Phe Gly Ala Gly Lys Pro Tyr Glu Phe
610 615 620

Ser Ile Asp Cys Asp Lys Ile Asn Leu Phe Arg Gly Val Phe Ala Phe
625 630 635 640

Leu Leu Tyr Val Ala Thr Phe Met Tyr Val Phe Ser Thr Phe Ala Asn
645 650 655

Ile Leu His Lys Glu Ser
660

<210> SEQ ID NO 12

<211> LENGTH: 5200

<212> TYPE: DNA

<213> ORGANISM: artificial

<220> FEATURE:

<223> OTHER INFORMATION: vector PHENFcabRGD4

<400> SEQUENCE: 12

```

gacgaaaggg cctcgtgata cgccatattt tatagggttaa tgtcatgata ataatgggtt      60
cttagacgtc aggtggcact ttccggggaa atgtgcgcgg aaccctatt tgtttatatt      120
tctaaataca ttcaaatatg tatccgctca tgagacaata accctgataa atgcttcaat      180
aatattgaaa aaggaagagt atgagtattc aacatttccg tgcgcgcctt attccctttt      240
ttgcggcatt ttgccttcct gtttttgctc acccagaaac gctggtgaaa gtaaaagatg      300
ctgaagatca gttgggtgca cgagtgggtt acatcgaact ggatctcaac agcggtaaga      360
tccttgagag ttttcgcccc gaagaacgtt ttccaatgat gagcactttt aaagtctctg      420
tatgtggcgc ggtattatcc cgtattgacg ccgggcaaga gcaactcggg cgccgcatac      480
actattctca gaatgacttg gttgagtact caccagtcac agaaaagcat cttacggatg      540
gcatgacagt aagagaatta tgcagtgtg ccataacatc gagtataaac actgcggcca      600
acttacttct gacaacgacg ggaggaccga aggagctaac cgcttttttg cacaacatgg      660
gggatcatgt aactcgcctt gatcgttggg aaccggagct gaatgaagcc ataccaaacy      720

```

-continued

acgagcgtga caccacgatg cctgtagcaa tggcaacaac gttgcgcaaa ctattaactg	780
gcgaactact tactctagct tcccggaac aattaataga ctggatggag gcggataaag	840
ttgcaggacc acttctgcgc tcggcccttc cggtggctg gtttattgct gataaatctg	900
gagccggtga gcgtgggtct cgcggtatca ttgcagcact ggggccagat ggtaagccct	960
cccgtatcgt agttatctac acgacgggga gtcaggcaac tatggatgaa cgaaatagac	1020
agatcgctga gatagggtgc tcaactgatta agcattggta actgtcagac caagtttact	1080
catatatact ttagattgat ttaaaacttc atttttaatt taaaaggatc taggtgaaga	1140
tcctttttga taatctcatg accaaaatcc cttaacgtga gttttcgttc cactgagcgt	1200
cagaccccg agaaaagatc aaaggatctt cttagatcc tttttttctg cgcgtaatct	1260
gctgcttgca aacaaaaaaa ccaccgctac cagcgggtgt ttgtttgccg gatcaagagc	1320
taccaactct ttttcgaag gtaactggct tcagcagagc gcagatacca aatactgtcc	1380
ttctagtgtg gccgtagtta ggccaccact tcaagaactc tgtagcaccg cctacatacc	1440
tcgctctgct aatcctgtta ccagtggctg ctgccagtgg cgataagtcg tgtcttaccg	1500
ggttggactc aagacgatag ttaccggata aggcgcagcg gtcgggctga acgggggggt	1560
cgtgcacaca gccacgttg gagcgaacga cctacaccga actgagatac ctacagcgtg	1620
agcattgaga aagcgccacg cttccgaag ggagaaaggc ggacaggtat ccggtaaagc	1680
gcagggtcgg aacaggagag cgcacgagg agcttccagg gggaaacgcc tggatatctt	1740
atagtcctgt cgggtttcgc cacctctgac ttgagcgtcg atttttgtga tgcctctcag	1800
gggggcgag cctatggaaa aacgccagca acgcggcctt tttaacgttc ctggcctttt	1860
gctggccttt tgctcacatg ttctttcctg cgttatcccc tgattctgtg gataaccgta	1920
ttaccgcctt tgagttagct gataccgtc gccgcagccg aacgaccgag cgcagcgagt	1980
cagtgagcga ggaagcggaa gagcgcccaa tacgcaaacc gcctctcccc gcgcgttggc	2040
cgattcatta atgcagctgg cagcagaggt ttcccgactg gaaagcgggc agtgagcgca	2100
acgcaattaa tgtgagttag ctcaactcatt aggcacccca ggctttacac tttatgcttc	2160
cggctcgat gttgtgtgga attgtgagcg gataacaatt tcacacagga aacagctatg	2220
accatgatta cgccaagctt aagcttgcac gcaaattcta tttcaaggag acagtcataa	2280
tgaataacct attgcctacg gcagccgctg gattgttatt actcgcggcc cagccggcca	2340
tggccgagcc caaatcttgt gacaaaactc acacatgccc accgtgcccc gcacctgaac	2400
tcctgggggg accgtcagtc ttctctctcc ccccaaaacc caaggacacc ctcatgatct	2460
cccggacccc tgaggtcaca tgcgtgggtg tggacgtgag ccacgaagac cctgagggtca	2520
agttcaactg gtacgtggac ggcgtggagg tgcataatgc caagacaaag ccgcgaggag	2580
agcagtacaa cagcagctac cgtgtggtca gcgtcctcac cgtcctgcac caggactggc	2640
tgaatggcaa ggagtacaag tgcaaggctt ccaacaaagc cctcccagcc cccatcgaga	2700
aaaccatctc caaagccaaa gggcagcccc gagaaccaca ggtgtacacc ctgcccccat	2760
cccgggatga gctgaccaag aaccagggtc gcctgacctg cctgggtcaaa ggcttctatc	2820
ccagcgacat cgccgtggag tgggagagca atgggcagcc ggagaacaac tacaagacca	2880
cgctcccggt gctggactcc gacggctcct tcttctctca cagcaagctt accgtgggtt	2940
gccgcggtga ttgtctgagc aggtggcagc aggggaacgt cttctcatgc tccgtgatgc	3000
atgaggctct gcacaaccac tacacgcaga agagcctctc cctgtctccg ggtaaagcgg	3060
ccgcagaaca aaaactcatc tcagaagagg atctgaatgg ggccgcatag actgttgaaa	3120

-continued

```

gttgttttagc aaaacctcat acagaaaatt catttactaa cgtctggaaa gacgacaaaa 3180
ctttagatcg ttacgctaac tatgagggtc gtctgtggaa tgctacaggc gttgtgggtt 3240
gtactggtga cgaaactcag tgttacggta catgggttcc tattgggctt gctatccctg 3300
aaaatgaggg tgggtgctct gaggggtggc gttctgaggg tggcggttct gaggggtggc 3360
gtactaaacc tcctgagtac ggtgatacac ctattccggg ctatacttat atcaaccctc 3420
tcgacggcac ttatccgctt ggtactgagc aaaacccgc taatccta atcctctcttg 3480
aggagtctca gcctcttaat actttcatgt ttcagaataa taggttcga aataggcagg 3540
gtgcattaac tgtttatacg ggcactgtta ctcaaggcac tgaccccggt aaaacttatt 3600
accagtacac tcctgtatca tcaaaagcca tgtatgacgc ttactggaac ggtaaattca 3660
gagactgcgc ttccattctt ggtcttaatg aggatccatt cgtttgtgaa tatcaaggcc 3720
aatcgtctga cctgcctcaa cctcctgtca atgctggcgg cggctctggt ggtgggtctg 3780
gtggcggctc tgagggtggc ggccttgagg gtggcgggtc tgagggtggc ggctctgagg 3840
gtggcgggtc cggtggcggc tccggttccg gtgattttga ttatgaaaa atggcaaacg 3900
ctaataaggg ggctatgacc gaaaatgccc atgaaaacgc gctacagtct gacgctaaag 3960
gcaaacttga ttctgtcgct actgattacg gtgctgctat cgatgggttc attggtgacg 4020
tttcggcctt tgctaagtgt aatggtgcta ctggtgattt tgetggctct aattcccaaa 4080
tggctcaagt cggtgacggt gataattcac ctttaatgaa taatttccgt caatatttac 4140
cttctttgcc tcagtcggtt gaatgtccc cttatgtctt tggcgctggt aaaccatatg 4200
aattttctat tgattgtgac aaaataaact tattccgtgg tgtctttgcg tttcttttat 4260
atgttgccac ctttatgtat gtattttcga cgtttgctaa catactgcat aaggagtctt 4320
aataagaatt cactggcctt cgttttacaa cgtcgtgact gggaaaaccc tggcggttac 4380
caacttaatc gccttgacgc acatccccct ttcgccagct ggcgtaatag cgaagaggcc 4440
cgcaccgatc gcccttccca acagttgcgc agcctgaatg gcgaatggcg cctgatgcgg 4500
tattttctcc ttacgcatct gtgcggtatt tcacaccgca cgtcaaagca accatagtac 4560
gcgccttgta gcggcgcatc aagcgcggcg ggtgtggtgg ttacgcgcag cgtgaccgct 4620
acacttgcca gcgccttagc gcccgctcct ttcgctttct tcccttcctt tctgccacg 4680
ttcgccggct tccccgta agctctaaat cgggggctcc ctttagggtt ccgatttagt 4740
gctttacggc acctgcaccc caaaaaactt gatttgggtg atggttcacg tagtgggcca 4800
tcgcctgat agacggtttt tcgccctttg acgttgaggt ccacgttctt taatagtga 4860
ctcttgttcc aaactggaac aacactcaac cctatctcgg gctattcttt tgatttataa 4920
gggattttgc cgatttcggc ctattgggtt aaaaatgagc tgatttaaca aaaatttaac 4980
gcgaatttta acaaaatatt aacgtttaca attttatggt gcaactctcag tacaactctg 5040
tctgatgcgc catagttaag ccagcccgca caccgcgcaa caccgctga cgcgcctga 5100
cgggcttgtc tgctccggc atccgcttac agacaagctg tgaccgtctc cgggagctgc 5160
atgtgtcaga ggttttcacc gtcacaccg aaacgcgcga 5200

```

<210> SEQ ID NO 13

<211> LENGTH: 48

<212> TYPE: DNA

<213> ORGANISM: artificial

<220> FEATURE:

<223> OTHER INFORMATION: CH3rlink

-continued

<400> SEQUENCE: 13

actagcggcc gcagagccac caccctcctt acccgagac agggagag 48

<210> SEQ ID NO 14

<211> LENGTH: 5215

<212> TYPE: DNA

<213> ORGANISM: artificial

<220> FEATURE:

<223> OTHER INFORMATION: vector pHENFcabRGD4L

<400> SEQUENCE: 14

gacgaaaggg cctcgtgata cgcctatatt tataggtaa tgtcatgata ataattggttt 60

cttagacgtc aggtggcact ttctggggaa atgtgcgcgg aacctctatt tgtttatatt 120

tctaaataca ttcaaatatg tatccgctca tgagacaata accctgataa atgcttcaat 180

aatattgaaa aaggaagagt atgagtattc aacatttccg tgctgcctct attccctttt 240

ttgcggcatt ttgccttctt gtttttctc acccagaaac gctggtgaaa gtaaaagatg 300

ctgaagatca gttgggtgca cgagtgggtt acatcgaact ggatctcaac agcggtaaga 360

tccttgagag ttttcgcccc gaagaacgtt ttccaatgat gagcactttt aaagtctctg 420

tatgtggcgc ggtattatcc cgtattgacg ccgggcaaga gcaactcggc cgcgcatac 480

actattctca gaatgacttg gttgagtact caccagtcac agaaaagcat cttacggatg 540

gcatgacagt aagagaatta tgcagtgtg ccataaccat gagtataac actgcggcca 600

acttacttct gacaacgacg ggaggaccga aggagctaac cgtttttttg cacaacatgg 660

gggatcatgt aactcgcctt gatcgttggg aaccggagct gaatgaagcc ataccaaacg 720

acgagcgtga caccacgatg cctgtagcaa tggcaacaac gttgcgcaaa ctattaactg 780

gogaactact tactctagct tcccggcaac aattaataga ctggatggag gcggataaag 840

ttgcaggacc acttctgcgc tcggcccttc cggctggctg gtttattgct gataaatctg 900

gagccggtga gcgtgggtct cgcggtatca ttgcagcact ggggccagat ggtaagccct 960

cccgatcgt agttatctac acgacgggga gtcaggcaac tatggatgaa cgaaatagac 1020

agatcgtga gatagggtgc tcaactgatta agcattggta actgtcagac caagtctact 1080

catatatact ttgatattgat ttaaaacttc atttttaatt taaaaggatc taggtgaaga 1140

tcctttttga taatctcatg accaaaatcc cttaacgtga gttttcgttc cactgagcgt 1200

cagaccccg agaaaagatc aaaggatctt cttgagatcc tttttttctg cgcgtaatct 1260

gctgcttgca aacaaaaaaa ccaccgctac cagcgggtgg ttgtttgccg gatcaagagc 1320

taccaactct ttttcgaag gtaactggct tcagcagagc gcagatacca aatactgtcc 1380

ttctagtgt gccgtagtta ggccaccact tcaagaactc tgtagcaccg cctacatacc 1440

tcgctctgct aatcctgtta ccagtggctg ctgccagtgg cgataagtcg tgtcttaccg 1500

ggttggactc aagacgatag ttaccggata aggcgcagcg gtcgggctga acgggggggt 1560

cgtgcacaca gccagcttg gagcgaacga cctacaccga actgagatac ctacagcgtg 1620

agcattgaga aagcgccacg cttcccgaag ggagaaaggc ggacaggtat ccggtaagcg 1680

gcagggtcgg aacaggagag cgcacgaggg agcttcaggc gggaaacgcc tggatatctt 1740

atagtcctgt cgggtttctc cacctctgac ttgagcgtcg atttttgtga tgctcgtcag 1800

gggggaggag cctatggaaa aacgccagca acgcggcctt tttacggttc ctggcctttt 1860

gctggccttt tgetcacatg ttctttctc cgttatcccc tgattctgtg gataaccgta 1920

ttaccgcctt tgagttagct gataccgctc gccgcagccg aacgaccgag cgcagcgagt 1980

-continued

cagtgcgcga ggaagcggaa gagcgcccaa tacgcaaacc gcctctcccc gcgcgttggc	2040
cgattcatta atgcagctgg cacgacaggt ttcccgaactg gaaagcgggc agtgagcgca	2100
acgcaattaa tgtgagtttag ctcaactcatt aggcacccca ggctttacac tttatgcttc	2160
cggctcgtat gttgtgtgga attgtgagcg gataacaatt tcacacagga aacagctatg	2220
accatgatta cgccaagctt aagcttgcac gcaaattcta tttcaaggag acagtcataa	2280
tgaataacct attgcctacg gcagccgctg gattgttatt actcgcggcc cagccggcca	2340
tggccgagcc caaatcttgt gacaaaactc acacatgcc accgtgcccc gcacctgaac	2400
tcttgggggg accgtcagtc ttctcttcc ccccaaaacc caaggacacc ctcatgatct	2460
cccgaccccc tgaggtcaca tgcgtggtgg tggacgtgag ccacgaagac cctgaggcca	2520
agttcaactg gtacgtggac ggcgtggagg tgcataatgc caagacaaag ccgcgggagg	2580
agcagtacaa cagcacgtac cgtgtggtca gcgtcctcac cgtcctgcac caggactggc	2640
tgaatggcaa ggagtacaag tgcaaggctc ccaacaaagc cctcccagcc cccatcgaga	2700
aaaccatctc caaagccaaa ggcgagcccc gagaaccaca ggtgtacacc ctgcccccat	2760
cccggtatga gctgaccaag aaccaggcca gctgacctg cctggtcaaa ggcttctatc	2820
ccagcgacat cgcctggag tgggagagca atgggcagcc ggagaacaac tacaagacca	2880
cgcctcccg tctggactcc gacggctcct tcttctctca cagcaagctt accgtgggtt	2940
gccgcggtga ttgtctgagc aggtggcagc aggggaacgt cttctcatgc tccgtgatgc	3000
atgaggctct gcacaaccac tacacgcaga agagcctctc cctgtctccg ggtaaggagg	3060
gtggtggctc tgcggcgcca gaacaaaaac tcatctcaga agaggatctg aatggggccg	3120
catagactgt tgaaagtgtg ttagcaaaac ctcatacaga aaattcattt actaacgtct	3180
ggaaagacga caaaacttta gatcgcttac ctaactatga gggctgtctg tggaatgcta	3240
caggcgttgt ggtttgtact ggtgacgaaa ctcaagtgtta cggtagatgg gttcctattg	3300
ggcttgctat ccctgaaaat gaggggtgtg gctctgaggg tggcggttct gaggggtggc	3360
gttctgaggg tggcggtact aaacctcctg agtacgggtg tacacctatt ccgggctata	3420
cttatatcaa cctctctgac ggcacttata cgcctggtac tgagcaaaac ccgctaatac	3480
ctaatacttc tcttgaggag tctcagctc ttaatacttt catgtttcag aataataggt	3540
tccgaaatag gcagggtgca ttaactgttt atacgggcac tgttactcaa ggcactgacc	3600
ccgttaaaac ttattaccag tacactcctg tatcatcaaa agccatgtat gacgcttact	3660
ggaacggtaa attcagagac tgcgctttcc attctggctt taatgaggat ccattcgttt	3720
gtgaatatca aggccaatcg tctgacctgc ctcaacctcc tgtcaatgct ggcgcggt	3780
ctggtggtgg ttctggtggc ggcctcgagg gtggcggtcc tgaggggtggc ggttctgagg	3840
gtggcggtcc tgaggggtggc ggttccggtg gcggctccgg ttccggtgat tttgattatg	3900
aaaaaatggc aaacgctaata aaggggggcta tgaccgaaaa tgccgatgaa aacgcgctac	3960
agtctgacgc taaaggcaaa cttgattctg tcgctactga ttacgggtgct gctatcgatg	4020
gtttcattgg tgacgtttcc ggccttgcta atggtaatgg tgctactggt gatthtgctg	4080
gctctaattc ccaaatggct caagtcggtg acggtgataa ttcaccttta atgaataatt	4140
tccgtcaata ttaccttct ttgcctcagt cgggtgaatg tcgcccctat gtctttggcg	4200
ctggtaaacc atatgaattt tctattgatt gtgacaaaaa aaacttatcc cgtggtgtct	4260
ttgcgtttct tttatatgtt gccaccttta tgtatgtatt ttgcagttt gctaacatac	4320

-continued

tgcataagga gtcttaataa gaattcactg gccgtcggtt tacaacgtcg tgactgggaa	4380
aaccttggcg ttaccaact taatcgctt gcagcacatc cccctttcgc cagctggcgt	4440
aatagcgaag aggcccgac cgategccct tccaacagt tgcgcagcct gaatggcgaa	4500
tggcgctga tgcggtatct tctccttacg catctgtgcg gtatttcaca ccgcacgtca	4560
aagcaacat agtacgcgc ctgtagcggc gcattaagcg cggcggtgt ggtggttacg	4620
cgcagcgtga ccgtacact tgcagcgcc ctagecgccg ctcccttcgc tttcttcct	4680
tccctttctg ccacgttcgc cggctttccc cgtcaagctc taaatcgagg gctcccttta	4740
gggttcgat ttagtgcttt acggcacctc gaccccaaaa aacttgattt ggggtgatgt	4800
tcacgtagtg ggccatcgcc ctgatagacg gtttttcgcc ctttgacgtt ggagtccacg	4860
ttctttaata gtggactctt gttccaaact ggaacaacac tcaaccctat ctgggctat	4920
tcttttgatt tataagggat tttgccgatt tcggcctatt ggttaaaaaa tgagctgatt	4980
taacaaaaat ttaacgcgaa ttttaacaaa atattaacgt ttacaatttt atgggtgact	5040
ctcagtacaa tctgctctga tgcgcgatag ttaagccagc cccgacacc gccaacacc	5100
gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca agctgtgacc	5160
gtctccggga gctgcatgtg tcagagggtt tcaccgctat caccgaaacg cgca	5215

<210> SEQ ID NO 15
 <211> LENGTH: 5013
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: vector pYD1dX

<400> SEQUENCE: 15

acggattaga agccgcgcag cgggtgacag cctccgaag gaagactctc ctccgtgcgt	60
cctcgtcttc accggtcgcg ttctgaaac gcagatgtgc ctgcgcgcg actgctccga	120
acaataaaga ttctacaata ctgactttta tgggtatgaa gaggaataa tggcagtaac	180
ctggccccc aaaccttcaa atgaacgaat caaattaaca accataggat gataatgcga	240
ttagtttttt agccttattt ctggggtaat taatcagcga agcagtgatt tttgatctat	300
taacagatat ataaatgcaa aaactgcata accactttaa ctaatacttt caacattttc	360
ggtttgattt acttcttatt caaatgtaat aaaagtatca acaaaaaatt gttaatatat	420
ctctatactt taacgtcaag gagaaaaaac cccggatcgg actactagca gctgtaatac	480
gactcactat agggaaatatt aagctaattc tacttcatac attttcaatt aagatgcagt	540
tacttcgctg tttttcaata ttttctgtta ttgcttcagt tttagcacag gaactgacaa	600
ctatatgcga gcaaatcccc tcaccaactt tagaatcgac gccgtactct ttgtcaacga	660
ctactatttt ggccaacggg aaggcaatgc aaggagtttt tgaatattac aaatcagtaa	720
cgtttgtcag taattgcggt tctcaccct caacaactag caaaggcagc cccataaaca	780
cacagtatgt ttttaagctt ctgcaggcta gtggtggtgg tggttctggt ggtggtggtt	840
ctggtggtgg tggttctgct agcatgactg gtggacagca aatgggtcgg gatctgtacg	900
acgatgacga taaggtaacca ggatccagtg tgggtgaatt ctgcagatat ccagcacagt	960
ggcgccgcgt cgatecagtc tagagggcc ttcgaagga agcctatccc taacctctc	1020
ctcggtctcg attctacgcg taccggtcat catcaccatc accattgagt ttaaacccgc	1080
tgatctgata acaacagtgt agatgtaaca aaatcgactt tgttcccact gtacttttag	1140
ctcgtacaaa atacaatata cttttcattt ctccgtaaac aacatgtttt ccatgtaat	1200

-continued

atccttttct	atctttcggt	ccgtttacaa	ctttacacat	actttatata	gtatttcact	1260
tctatacact	aaaaaactaa	gacaatttta	atcttgctgc	ctgccatatt	tcaatttggt	1320
ataaattcct	ataatttatc	ctattagtag	ctaaaaaaag	atgaatgtga	atcgaatcct	1380
aagagaattg	ggcaagtgc	caaacaatac	ttaaataaat	actactcagt	aataacctat	1440
ttcttagcat	tttgacgaa	atctgctatt	ttgtagagt	ctttacacc	atttgtctcc	1500
acacctccgc	ttacatcaac	accaataacg	ccatttaato	taagcgcac	accaacattt	1560
tctggcgta	gtccaccagc	taacataaaa	tgtaagctct	cggggctctc	ttgccttcca	1620
accagtcag	aatcgaggt	ccaatccaaa	agttcacctg	tcccacctgc	ttctgaatca	1680
aacaaggga	taaacgaatg	aggtttctgt	gaagctgcac	tgagtagtat	gttgagctct	1740
tttgaaata	cgagtctttt	aataactggc	aaaccgagga	actcttggtg	ttcttgccac	1800
gactcatctc	cgtgcagttg	gacgatatac	atgccgtaat	cattgaccag	agccaaaaca	1860
tcctccttag	gttgattacg	aaacacgcca	accaagtatt	tcggagtgc	tgaactattt	1920
ttatatgctt	ttacaagact	tgaatttttc	cttgcaataa	cggggtcaat	tggtctcttt	1980
ctattgggca	cacatataat	accagcaag	tcagcatcgg	aatctagagc	acattctgcg	2040
gcctctgtgc	tctgcaagcc	gcaaaccttc	accaatggac	cagaactacc	tgtgaaatta	2100
ataacagaca	tactccaagc	tgctttgtg	tgcttaatac	cgtataactc	cgtgctcaat	2160
agtcaccaat	gccctccctc	ttggccctct	ccttttcttt	tttcgaccga	atttcttgaa	2220
gacgaaagg	cctcgtgata	cgctattttt	tatagggtta	tgctcatgata	ataatgggtt	2280
cttaggacgg	atcgcttgcc	tgtaacttac	acgcgcctcg	tatcttttaa	tgatggaata	2340
atttgggaat	ttactctgtg	tttatttatt	tttatgtttt	gtatttggat	tttagaaagt	2400
aaataaagaa	ggtagaagag	ttacggaatg	aagaaaaaaa	aataaacaaa	ggtttaaaaa	2460
atttcaacaa	aaagcgtact	ttacatatat	atttattaga	caagaaaagc	agattaaata	2520
gatatacatt	cgattaacga	taagtaaaat	gtaaaatcac	aggattttcg	tgtgtggtct	2580
tctacacaga	caagatgaaa	caattcggca	ttaataacctg	agagcaggaa	gagcaagata	2640
aaagtagta	ttgtttggcg	atccccctag	agtcttttac	atcttcggaa	aacaaaaact	2700
attttttctt	taattttctt	ttttactttc	tatttttaat	ttatatattt	atattaaaaa	2760
atttaaat	taattatttt	tatagcacgt	gatgaaaagg	accaggtgg	cacttttcgg	2820
ggaaatgtgc	gcggaacccc	tatttggtta	tttttctaaa	tacattcaaa	tatgtatccg	2880
ctcatgagac	aataacctcg	ataaatgctt	caataatatt	gaaaaaggaa	gagtatgagt	2940
attcaacatt	tccgtgtcgc	ccttattccc	ttttttgcgg	cattttgcct	tcctgttttt	3000
gctcaccag	aaacgctggt	gaaagtataa	gatgctgaag	atcagttggg	tgacagagtg	3060
ggttacatcg	aactggatct	caacagcggg	aagatccttg	agagttttcg	ccccgaagaa	3120
cgttttccaa	tgatgagcac	ttttaagtt	ctgctatgtg	gcgcggtatt	atcccggtgt	3180
gacgcccggc	aagagcaact	cggtcgccgc	atacactatt	ctcagaatga	cttggttgag	3240
tactcaccag	tcacagaaaa	gcattcttac	gatggcatga	cagtaagaga	attatgcagt	3300
gctgccataa	ccatgagtga	taacactgcg	gccaaactac	ttctgacaac	gatcggagga	3360
ccgaaggagc	taaccgcttt	tttgacaaac	atgggggatc	atgtaactcg	ccttgatcgt	3420
tgggaaccgg	agctgaatga	agccatacca	aacgacgagc	gtgacaccac	gatgcctgta	3480
gcaatggcaa	caacgttgcg	caaactatta	actggcgaac	tacttaactc	agcttcccgg	3540

-continued

caacaattaa tagactggat ggaggcggat aaagttgcag gaccacttct gcgctcggcc	3600
cttcgggctg gctgggttat tgcgtataaa tctggagccg gtgagcgtgg gtctcgcggt	3660
atcattgcag cactggggcc agatggtaag cctcccgta tcgtagttat ctacacgacg	3720
ggcagtcagg caactatgga tgaacgaaat agacagatcg ctgagatagg tgcctcactg	3780
attaagcatt ggtaactgtc agaccaagtt tactcatata tacttttagat tgatttaaaa	3840
cttcattttt aatttaaaag gatctagggt aagatccttt ttgataatct catgacaaaa	3900
atcccttaac gtgagttttc gttccactga gcgtcagacc ccgtagaaaa gatcaaagga	3960
tcttcttgag atcctttttt tctgcgcgta atctgctgct tgcaaaaaaaa aaaaccaccg	4020
ctaccagcgg tggtttgttt gccggatcaa gagctaccaa ctctttttcc gaaggtaact	4080
ggcttcagca gagcgcagat accaaatact gtccttctag tgtagccgta gttaggccac	4140
cacttcaaga actctgtagc accgcctaca tacctcgctc tgctaatect gttaccagt	4200
gctgctgcca gtggcgataa gtcgtgtctt accgggttgg actcaagacg atagttaccg	4260
gataaggcgc agcggctcggg ctgaacgggg gggtcgtgca cacagcccag cttggagcga	4320
acgacctaca ccgaactgag atacctacag cgtgagcatt gagaagcgc cacgcttccc	4380
gaaggggagaa aggcggacag gtatccggtg agcggcaggg tcggaacagg agagcgcacg	4440
agggagcttc caggggggaa cgcctgggtat ctttatagtc ctgtcgggtt tcgccacctc	4500
tgacttgagc gtcgattttt gtgatgctcg tcaggggggc cgagcctatg gaaaaacgcc	4560
agcaacgcgg cctttttacg gttcctggcc ttttctgggc cttttgctca catgttcttt	4620
cctgcgttat cccctgattc tgtggataac cgtattaccg cctttgagtg agctgatacc	4680
gctcgccgca gccgaacgac cgagcgcagc gagtcagtga gcgaggaagc ggaagagcgc	4740
ccaatacgca aaccgcctct ccccgcgctg tggccgattc attaatgcag ctggcacgac	4800
aggtttcccg actggaaagc gggcagtgag cgcaacgcaa ttaatgtgag ttacctcact	4860
cattaggcac ccagggtttt acactttatg cttccggctc ctatgttggtg tggaattgtg	4920
agcggataac aatttcacac aggaacacgc tatgaacctg attacgcaa gctcggaatt	4980
aaccctcact aaagggaaca aaagctggct agt	5013

<210> SEQ ID NO 16

<211> LENGTH: 5971

<212> TYPE: DNA

<213> ORGANISM: artificial

<220> FEATURE:

<223> OTHER INFORMATION: vector pYD1dXFc

<400> SEQUENCE: 16

acggattaga agccgccgag cgggtgacag cctccgaag gaagactctc ctccgtgcgt	60
cctcgtcttc accggtcgcg ttccctgaaac gcagatgtgc ctgcgcgcgc actgctccga	120
acaataaaga ttctacaata ctagctttta tggttatgaa gaggaaaaaat tggcagtaac	180
ctggccccac aaaccttcaa atgaacgaat caaattaaca accataggat gataatgcga	240
ttagtttttt agccttattt ctggggtaat taatcagcga agcgatgatt tttgatctat	300
taacagatat ataaatgcaa aaactgcata accactttta ctaatacttt caacattttc	360
ggtttgtatt acttcttatt caaatgtaat aaaagtatca acaaaaaatt gttaatatac	420
ctctatactt taacgtcaag gagaaaaaac cccggatcgg actactagca gctgtaatac	480
gactcactat agggaaatatt aagctaattc tacttcatac attttcaatt aagatgcagt	540
tacttcgctg tttttcaata ttttctgtta ttgcttcagt tttagcacag gaactgacaa	600

-continued

ctatatgcga gcaaatcccc tcaccaactt tagaatcgac gccgtactct ttgtcaacga	660
ctactatttt ggccaacggg aaggcaatgc aaggagtttt tgaatattac aaatcagtaa	720
cgtttgtcag taattgCGgt tctaccccct caacaactag caaaggcagc cccataaaca	780
cacagtatgt ttttaagctt ctgcaggcta gtgggtgggtg tggttctggt ggtgggtggt	840
ctgggtgggtg tggttctgct agcatgactg gtggacagca aatgggtcgg gatctgtacg	900
acgatgacga taaggtagca ggatccgcta gcaccaaggg cccagcgtg ttccctctgg	960
ccccagctc caagagcacc tccggcggca ccgcgcctt gggctgcctg gtgaaggatt	1020
acttcccaga gcccgtagc gtgagctgga acagcggcgc cctgaccagc ggcgtgcaca	1080
cctttcccgc cgtgctcag tccagcggcc tgtactccct gagcagcgtg gtgaccgtgc	1140
ccagcagcag cctgggcacc cagacctaca tctgcaatgt gaaccacaag cccagcaata	1200
ccaaggtgga taagaagggtg gagccaaga gcagcgacaa gacacacacg tgtccccat	1260
gtcccgcctc tgagctcgtg ggcggccctt ccgtgttctt gttccctccc aagccaaagg	1320
acaccctgat gatctcccg acccctgagg tgacctgtgt ggtgggtggac gtgagccacg	1380
aggaccaga ggtgaagtgc aactggtacg tggacggcgt ggaggtgcac aacgccaaga	1440
ccaagcctag agaggagcag tacaacagca cctaccgctg ggtgagcgtg ctgaccgtgc	1500
tgcaccagga ttggctgaat ggcaaggagt acaagtgcaa ggtgagcaac aaggccctgc	1560
ctgcccccat cgagaagacc atctccaagg ccaagggcca gcctcgagaa ccacaggtgt	1620
acaccctgcc cccatcccg gatgagctga ccaagaacca ggtcagcctg acctgcctgg	1680
tcaaaggctt ctatcccagc gacatcgccg tggagtggga gagcaatggg cagccggaga	1740
acaactacaa gaccacgct cccgtgctgg actccgacgg ctcttcttc ctctacagca	1800
agctcacctg ggacaagagc aggtggcagc aggggaacgt cttctcatgc tccgtgatgc	1860
atgaggctct gcacaaccac tacacacaga agagcctctc cctgtctccg ggtaaatgag	1920
cggccgctcg atcgagtcta gagggccctt cgaaggtaag cctatcccta accctctcct	1980
cggctcctgat tctacgcta ccgggtcatca tcaccatcac cattgagttt aaaccgctg	2040
atctgataac aacagtgtag atgtaacaaa atcgactttg ttcccactgt acttttagct	2100
cgtacaaaat acaatatact ttctatttct ccgtaaacaa catgttttcc catgtaatat	2160
ccttttctat ttttcttcc gttaccaact ttacacatac tttatatagc tattcacttc	2220
tatacactaa aaaactaaga caattttaat tttgtgcct gccatatttc aatttgttat	2280
aaattcctat aatttctct attagtagct aaaaaagat gaatgtgaat cgaatcctaa	2340
gagaattggg caagtgcaca aacaatactt aaataaatac tactcagtaa taacctattt	2400
cttagcattt ttgacgaat ttgctatttt gttagagtct tttacaccat ttgtctccac	2460
acctccgctt acatcaacac caataacgcc atttaacta agcgcacac caacattttc	2520
tggcgtcagt ccaccagcta acataaaatg taagctctcg gggctctctt gccttccaac	2580
ccagtcagaa atcgagtcc aatccaaaag ttcacctgct ccacctgctt ctgaatcaaa	2640
caagggaata aacgaatgag gtttctgtga agctgcactg agtagtatgt tgcagtcttt	2700
tggaaatacg agtcttttaa taactggcaa accgaggaac tcttggtatt cttgccacga	2760
ctcatctccg tgcagtggga cgatatcaat gccgtaatca ttgaccagag ccaaaacatc	2820
ctccttaggt tgattacgaa acacgccaac caagtatttc ggagtgcctg aactattttt	2880
atatgctttt acaagacttg aaattttcct tgcaataacc gggtoaattg ttctctttct	2940

-continued

attgggcaca	catataatac	ccagcaagtc	agcatcggaa	tctagagcac	attctgcggc	3000
ctctgtgctc	tgcaagcgcg	aaactttcac	caatggacca	gaactacctg	tgaaattaat	3060
aacagacata	ctccaagctg	cctttgtgtg	cttaatcacg	tatactcacg	tgctcaatag	3120
tcaccaatgc	cctccctctt	ggccctctcc	ttttcttttt	tcgaccgaat	ttcttgaaga	3180
cgaaagggcc	tcgtgatacg	cctattttta	taggttaatg	tcagtataat	aatggtttct	3240
taggacggat	cgcttgctg	taacttacac	gcgcctcgta	tcctttaatg	atggaataat	3300
ttgggaattt	actctgtgtt	tatttatatt	tatgttttgt	atttgatttt	tagaaagtaa	3360
ataaagaagg	tagaagagtt	acggaatgaa	gaaaaaaaaa	taaacaaagg	tttaaaaaat	3420
ttcaacaaaa	agcgactctt	acatatatat	ttattagaca	agaaaagcag	attaaataga	3480
tatacattcg	attaacgata	agtaaaatgt	aaaatcacag	gattttcgtg	tgtggtcttc	3540
tacacagaca	agatgaaaca	attcggcatt	aatacctgag	agcaggaaga	gcaagataaa	3600
aggtagtatt	tgttggcgat	ccccctagag	tcctttacat	cttcggaaaa	caaaaactat	3660
ttttctttta	atttcttttt	ttactttcta	tttttaattt	atatatttat	attaaaaaat	3720
ttaaattata	attattttta	tagcacgtga	tgaaaaggac	ccagggtggca	cttttcgggg	3780
aatgtgctgc	ggaaccccta	ttgttttatt	tttctaaata	cattcaaata	tgtatccgct	3840
catgagacaa	taaccctgat	aaatgcttca	ataatattga	aaaaggaaga	gtatgagtat	3900
tcaacatttc	cgtgtcgccc	ttattccctt	ttttgcggca	ttttgccttc	ctgtttttgc	3960
tcacccgaaa	acgtcgttga	aagtaaaaga	tgctgaagat	cagttgggtg	cacgagtggg	4020
ttacatcgaa	ctggatctca	acagcggtaa	gatccttgag	agttttcgcc	ccgaagaacg	4080
ttttccaatg	atgagcactt	ttaaagttct	gctatgtggc	gcggtattat	cccggttgta	4140
cgccggggcaa	gagcaactcg	gtcgccgcat	acactattct	cagaatgact	tggttgagta	4200
ctcaccagtc	acagaaaagc	atcttacgga	tgccatgaca	gtaagagaat	tatgcagtgc	4260
tgccataacc	atgagtatac	acactgcggc	caacttactt	ctgacaacga	tcggaggacc	4320
gaaggagcta	accgcttttt	tgacacaacat	gggggatcat	gtaactcgcc	ttgatcgttg	4380
ggaaccggag	ctgaatgaag	ccataccaaa	cgacgagcgt	gacaccacga	tgccgttagc	4440
aatggcaaca	acgttgcgca	aactattaac	tgccgaacta	cttactctag	cttcccgcca	4500
acaattaata	gactggatgg	aggcggataa	agttgcagga	ccacttctgc	gctcggccct	4560
tccggctggc	tggtttattg	ctgataaaat	tgagccgggt	gagcgtgggt	ctcgcggtat	4620
cattgcagca	ctggggccag	atggtaagcc	ctcccgatc	gtagtattct	acacgacggg	4680
cagtcaggca	actatggatg	aacgaaatag	acagatcgct	gagatagggt	cctcactgat	4740
taagcattgg	taactgtcag	accaagttta	ctcatatata	cttttagattg	atttaaaact	4800
tcatttttaa	tttaaaagga	tctaggtgaa	gatccttttt	gataatctca	tgacaaaaat	4860
cccttaacgt	gagttttctg	tccactgagc	gtcagacccc	gtagaaaaga	tcaaaggatc	4920
ttcttgagat	cctttttttc	tgccgcta	ctgctgcttg	caaacaaaaa	aaccaccgct	4980
accagcgggt	gtttgtttgc	cggatcaaga	gctaccaact	ctttttccga	aggttaactgg	5040
cttcagcaga	gcgcagatac	caaatactgt	ccttctagtg	tagccgtagt	taggccacca	5100
cttcaagaac	tctgtagcac	gcctacata	cctcgctctg	ctaactcctgt	taccagtggc	5160
tgctgccagt	ggcgataagt	cgtgtcttac	cgggttgagc	tcaagacgat	agttaccgga	5220
taaggcgcag	cggtcgggct	gaacgggggg	ttcgtgcaca	cagcccagct	tgagcgaac	5280
gacctacacc	gaactgagat	acctacagcg	tgagcattga	gaaagcgcca	cgcttcccca	5340

-continued

```

aggagaaaag gcgacaggt atccggttaag cggcagggtc ggaacaggag agcgacagag 5400
ggagcttcca ggggggaacg cctggtatct ttatagtcct gtcgggttcc gccacctctg 5460
acttgagcgt cgatttttgt gatgctcgtc aggggggccc agcctatgga aaaacgccag 5520
caacgcggcc tttttacggt tctggcctt ttgtggcct tttgctcaca tgttctttcc 5580
tgcgttatcc cctgattctg tggataaccg tattaccgcc tttagtgag ctgataccgc 5640
tcgccgcagc cgaacgaccg agcgacagca gtcagtgcgc gaggaagcgg aagagcgccc 5700
aatacgcaaa ccgctctcc ccgcgcgttg gccgattcat taatgcagct ggcacgacag 5760
gtttcccgac tggaaagcgg gcagtgcgc caacgcaatt aatgtgagtt acctcactca 5820
ttaggcaccc caggctttac actttatgct tccggctcct atgttgtgtg gaattgtgag 5880
cggataacaa tttcacacag gaaacagcta tgacctgat tacgccaagc tcggaattaa 5940
ccctcactaa agggaacaaa agctggctag t 5971

```

```

<210> SEQ ID NO 17
<211> LENGTH: 5657
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: pYD1CH12

```

```

<400> SEQUENCE: 17

```

```

acggattaga agccgccgag cgggtgacag ccctccgaag gaagactctc ctccgtgcgt 60
cctcgtcttc accggtgcgc ttctgaaac gcagatgtgc ctgcgcgcgc actgctccga 120
acaataaaga ttctacaata ctagctttta tggttatgaa gaggaaaaaat tggcagtaac 180
ctggccccac aaaccttcaa atgaacgaat caaattaaca accataggat gataatgcga 240
ttagtttttt agccttattt ctggggtaat taatcagcga agcgatgatt ttgatctat 300
taacagatat ataaatgcaa aaactgcata accactttaa ctaatacttt caacattttc 360
ggtttgattt acttcttatt caaatgtaat aaaagtatca acaaaaaatt gttaatatat 420
ctctatactt taacgtcaag gagaaaaaac ccggatcgg actactagca gctgtaatac 480
gactcactat agggaaatatt aagctaattc tacttcatac attttcaatt aagatgcagt 540
tacttcgctg tttttcaata ttttctgtta ttgcttcagt tttagcacag gaactgacaa 600
ctatatgcga gcaaatcccc tcaccaactt tagaatcgac gccgtactct ttgtcaacga 660
ctactatttt ggccaacggg aaggcaatgc aaggagtttt tgaatattac aaatcagtaa 720
cgtttgcag taattgcggt tctcaccct caacaactag caaaggcagc ccataaaca 780
cacagtatgt ttttaagctt ctgcaggcta gtggtgggtg tggttctggt ggtgggtggt 840
ctggtgggtg tggttctgct agcatgactg gtggacagca aatgggtcgg gatctgtacg 900
acgatgacga taaggtacca ggtccgcta gcaccaaggg cccagcgtg ttccctctgg 960
ccccagctc caagagcacc tccggcggca ccgcgcctc gggctgcctg gtgaaggatt 1020
acttcccaga gcccgtagcc gtgagctgga acagcggcgc cctgaccagc ggcgtgcaca 1080
cctttccgcg cgtgctgcag tccagcggcc tgtactccct gagcagcgtg gtgacggtgc 1140
ccagcagcag cctgggcacc cagacctaca tctgcaatgt gaaccacaag cccagcaata 1200
ccaagggtga taagaagggt gagccaaga gcagcgacaa gacacacacg tgtccccat 1260
gtcccgcccc tgagctgctg ggcggccctt ccgtgttcc tttccctccc aagccaaagg 1320
acaccctgat gatctcccg acccctgagg tgacctgtgt ggtgggtggac gtgagccacg 1380

```


-continued

aggacccaga ggtgaagtac aactggtacg tggacggcgt ggaggtgcac aacgccaaga	1440
ccaagcctag agaggagcag tacaacagca cctaccgctg ggtgagcgtg ctgaccgtgc	1500
tgcaccagga ttggtgaat ggcaaggagt acaagtgcac ggtgagcaac aaggccctgc	1560
ctgccccat cgagaagacc atctccaagg ccaagggcca gcctcgagggc cgctcgatcg	1620
agtctagagg gcccttcgaa ggtaagccta tccctaacc tctcctcggg ctcgattcta	1680
cgcgtagcgg tcatcatcac catcaccatt gagtttaaac ccgctgatct gataacaaca	1740
gtgtagatgt aacaaaatcg actttgttcc cactgtactt ttagctcgta caaaatacaa	1800
tatacttttc atttctcctg aaacaacatg ttttcccatg taatatcctt ttctattttt	1860
cgttccggtta ccaactttac acatacttta tatagctatt cacttctata cactaaaaaa	1920
ctaagacaat tttaatTTtg ctgcctgcca tatttcaatt tgttataaat tcctataatt	1980
tatcctatta gtacttaaaa aaagatgaat gtgaatcgaa tcctaagaga attgggcaag	2040
tgcacaaaaca atactttaat aaatactact cagtaataac ctatttctta gcatttttga	2100
cgaaatttgc tattttgtta gactctttta caccatttgt ctccacacct ccgcttacat	2160
caacaccaat aacgccattt aatctaagcg catcaccaac attttctggc gtcagtcac	2220
cagctaacat aaaatgtaag ctctcggggc tctcttgctt tccaaccag tcagaaatcg	2280
agttccaatc caaaagtta cctgtccac ctgcttctga atcaacaag ggaataaacg	2340
aatgaggttt ctgtgaagct gcactgagta gtatgttgca gtcttttga aatacgagtc	2400
ttttaataac tggcaaacg aggaactctt ggtattcttg ccacgactca tctcgtgca	2460
gttgagcgt atcaatgccc taatcattga ccagagccaa aacatcctcc ttaggttgat	2520
tacgaaacac gccaccaag tatttcggag tgcctgaact atttttatat gcttttaca	2580
gacttgaat tttccttgca ataaccgggt caattgttct ctttctattg ggcacacata	2640
taataccag caagtacga tcggaatcta gagcacattc tgcggcctct gtgctctgca	2700
agccgcaaac tttaccaat ggaccagaac tacctgtgaa attaataaca gacatactcc	2760
aagctgcctt tgtgtgctta atcacgtata ctcacgtgct caatagtcac caatgcctc	2820
cctcttgccc ctctcctttt cttttttcga ccgaatttct tgaagacgaa agggcctcgt	2880
gatacgcta tttttatagg ttaatgtcat gataataatg gtttcttagg acggatcgct	2940
tgcctgtaac ttacacgcgc ctcgatctt ttaatgatgg aataatttgg gaatttactc	3000
tgtgtttatt tatttttatg ttttgattt ggattttaga aagtaataa agaaggtaga	3060
agagttacgg aatgaagaaa aaaaaataaa caaagggtta aaaaatttca acaaaaagcg	3120
tactttacat atatatttat tagacaagaa aagcagatta aatagatata cattcgatta	3180
acgataagta aaatgtaaaa tcacaggatt ttcgtgtgtg gtcttctaca cagacaagat	3240
gaaacaattc ggcathtaata cctgagagca ggaagagcaa gataaaagg agtatttgtt	3300
ggcgatcccc ctagagtctt ttacatcttc ggaaaaaaa aactattttt tctttaattt	3360
ctttttttac tttctatttt taatttatat atttatatta aaaaatttaa attataatta	3420
tttttatagc acgtgatgaa aaggacccag gtggcacttt tcggggaaat gtgcgcggaa	3480
cccctatttg tttatttttc taaatacatt caaatatgta tccgctcatg agacaataac	3540
cctgataaat gcttcaataa tattgaaaaa ggaagagtat gagtattcaa ctttccgtg	3600
tcgccttat tccctttttt gcggcatttt gccttctgt ttttgcac ccagaaacgc	3660
tggtgaaagt aaaagatgct gaagatcagt tgggtgcacg agtgggttac atcgaactgg	3720
atctcaacag cggtaagatc cttgagagtt ttcgccccga agaacgtttt ccaatgatga	3780

-continued

```

gcacttttaa agttctgcta tgtggcgcggt tattatcccg tgttgacgcc gggcaagagc 3840
aactcggctcg ccgcatacac tattctcaga atgacttggt tgagtactca ccagtcacag 3900
aaaagcatct tacggatggc atgacagtaa gagaattatg cagtgtctgcc ataaccatga 3960
gtgataaacac tgcggccaac ttacttctga caacgatcgg aggaccgaag gagctaaccg 4020
cttttttgca caacatgggg gatcatgtaa ctgcgcttga tcgttgggaa ccggagctga 4080
atgaagccat accaaacgac gagcgtgaca ccacgatgcc tgtagcaatg gcaacaacgt 4140
tgcgcaaaact attaaactggc gaactactta ctctagcttc ccggcaacaa ttaatagact 4200
ggatggaggc ggataaagtt gcaggaccac ttctgcgctc ggcccttcgg gctggctggt 4260
ttattgctga taaatctgga gccggtgagc gtgggtctcg cgggtatcatt gcagcactgg 4320
ggccagatgg taagccctcc cgtatcgtag ttatctacac gacgggcagt caggcaacta 4380
tggatgaacg aaatagacag atcgtgaga taggtgcctc actgattaag catttgtaac 4440
tgtcagacca agtttactca tatatacttt agattgattt aaaacttcat ttttaattta 4500
aaaggatcta ggtgaagatc ctttttgata atctcatgac caaaatccct taacgtgagt 4560
tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa aggatcttct tgagatcctt 4620
tttttctgcg cgtaatctgc tgcttgcaaa caaaaaaacc accgctacca gcggtgggtt 4680
gtttgccgga tcaagagcta ccaactcttt ttccgaaggt aactggcttc agcagagcgc 4740
agataccaaa tactgtcctt ctagtgtagc cgtagttagg ccaccacttc aagaactctg 4800
tagcacccgc tacatactc gctctgctaa tctgttacc agtggctgct gccagtggcg 4860
ataagtcgtg tcttaccggg ttggactcaa gacgatagtt accggataag gcgcagcggg 4920
cgggctgaac gggggggtcg tgcacacagc ccagcttggg gcgaacgacc tacaccgaac 4980
tgagatacct acagcgtgag cattgagaaa gcgccacgct tcccgaaggg agaaaggcgg 5040
acaggtatcc ggtaagcggc aggggtcggaa caggagagcg cacgaggagg cttccagggg 5100
ggaacgcctg gtatctttat agtctgtcg ggtttcgcca cctctgactt gagcgtcgat 5160
ttttgtgatg ctgctcaggg gggccgagcc tatggaaaa cgccagcaac gcggcctttt 5220
tacggttctt ggccttttgc tggccttttg ctccatggt ctttctgctg ttatccctg 5280
attctgtgga taaccgtatt accgccttg agtgagctga taccgctcgc cgcagccgaa 5340
cgaccgagcg cagcagagta gtgagcgagg aagcgggaag gcgcccaata cgcaaaccgc 5400
ctctccccgc gcgttgccg attcattaat gcagctggca cgacagggtt cccgactgga 5460
aagcgggcag tgagcgcaac gcaattaatg tgagttacct cactcattag gcaccccagg 5520
ctttacactt tatgcttcgg gtcctatgt tgtgtggaat tgtgagcgga taacaatttc 5580
acacaggaaa cagctatgac catgattacg ccaagctcgg aattaaccct cactaaaggg 5640
aacaaaagct ggctagt 5657

```

```

<210> SEQ ID NO 18
<211> LENGTH: 738
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: Fcab01
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (449)..(450)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (452)..(453)

```

-continued

```

<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (455)..(456)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (611)..(612)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (614)..(615)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (617)..(618)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (620)..(621)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (623)..(624)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (626)..(627)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (629)..(630)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (632)..(633)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (641)..(642)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (644)..(645)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 18

ggcccgccg gccatggccg agcccaaacc ttgtgacaaa actcacacat gccaccgtg      60
cccagcacct gaactcctgg ggggaccgtc agtcttcttc ttcccccaa aacccaagga      120
caccctcatg atctcccgga cccctgaggt cacatgctgt gtggtggacg tgagccacga      180
agaccctgag gtcaagttca actggtacgt ggacggcgtg gaggtgcata atgccaagac      240
aaagccgcgg gaggagcagt acaacagcac gtaccgtgtg gtcagcgtcc tcaccgtcct      300
gcaccaggac tggtgtaagt gcaaggagta caagtgaag gtctccaaca aagccctccc      360
agcccccacg gagaaaacca tctccaaagc caaagggcag cctcgagaac cacaggtgta      420
caccctgccc ccattccggg atgaactggn bnnbnnbcag gtcagcctga cctgcctggt      480
caaaggcttc tatcccgagc acatcgccgt ggagtgggag agcaatgggc agccggagaa      540
caactacaag accacgcctc ccgtgctgga ctccgacggc tccttcttcc tctacagcaa      600
gtcaccgtgt nnnnnnnnnn nnnnnnnnnn bnnbaggtgg nnnnnbaggga acgtcttctc      660
atgctccgtg atgcatgagg ctctgcacaa ccaactacag cagaagagcc tctccctgtc      720
tccgggtaaa gcggccgc                                738

<210> SEQ ID NO 19
<211> LENGTH: 738
<212> TYPE: DNA
<213> ORGANISM: artificial

```

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Fcab02

<400> SEQUENCE: 19

ggcccagccg gccatggccg agcccaaata ttgtgacaaa actcacacat gccacccgtg      60
cccagcacct gaactectgg ggggaccgtc agtcttcttc tccccccaa aaccaagga      120
caccctcatg atctcccgga cccctgaggt cacatgcgtg gtggtggacg tgagccacga      180
agaccctgag gtcaagtcca actggtacgt ggacggcggt gaggtgcata atgccaagac      240
aaagccgcgg gaggagcagt acaacagcac gtaccgtgtg gtcagcgccc tcaccgtcct      300
gcaccaggac tggctgaatg gcaaggagta caagtgaag gtctccaaca aagccctccc      360
agcccccatc gagaaaacca tctccaaagc caaagggcag cctcgagaac cacagggtga      420
caccctgccc ccattccggg atgagctgkm tkmtkmtcag gtgagcctga cctgcctggt      480
caaaggcttc tatcccagcg acatcgccgt ggagtgggag agcaatgggc agccggagaa      540
caactacaag accacgcctc ccgtgtctga ctccgacggc tccttcttcc tctacagcaa      600
gtcacccgtg kmtkmtkmtk mtkmtkmtkm tkmtagtggt kmtkmtggga acgtcttctc      660
atgctccgtg atgcatgagg ctctgcacaa ccactacacg cagaagagcc tctccctgtc      720
tccgggtaaa gcggccgc                                          738

```

```

<210> SEQ ID NO 20
<211> LENGTH: 750
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: Fcab03
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (449)..(450)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (452)..(453)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (455)..(456)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (617)..(618)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (620)..(621)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (623)..(624)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (626)..(627)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (629)..(630)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (632)..(633)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (635)..(636)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:

```

-continued

```

<221> NAME/KEY: misc_feature
<222> LOCATION: (638)..(639)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (653)..(654)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (656)..(657)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 20

ggcccagccg gccatggccg agcccaaata ttgtgacaaa actcacacat gccaccgctg      60
cccagcacct gaactcctgg ggggaccgtc agtcttcttc tccccccaa aaccaagga      120
caccctcatg atctcccgga ccctgaggt caccatgcgtg gtggtggacg tgagccacga      180
agaccctgag gtcaagtcca actggtacgt ggacggcgctg gaggtgcata atgccaagac      240
aaagccgcgg gaggagcagt acaacagcac gtaccgtgtg gtcagcgtcc tcaccgtcct      300
gcaccaggac tggctgaatg gcaaggagta caagtgaag gtctccaaca aagccctccc      360
agcccccatc gagaaaacca tctccaaagc caaagggcag cctcgagaac cacaggtgta      420
caccctgccc ccttccggg atgagctggn bnnbnnbcag gtcagcctga cctgcctggt      480
caaaggcttc tatccagcg acatcgccgt ggagtgggag agcaatgggc agccggagaa      540
caactacaag accacgcctc ccgtgctgga ctccgacggc tccttcttcc tctacagcaa      600
gtcacccgtg ggttctnnbn nbnbnnnbn bnnbnnnnb agcggcaggt ggnnbnnbgb      660
gaacgtcttc tcattgctcg tgatgcatga ggctctgcac aaccactaca cgcagaagag      720
cctctccctg tctccgggta aagcggccgc      750

```

```

<210> SEQ ID NO 21
<211> LENGTH: 750
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: Fcab04

```

```

<400> SEQUENCE: 21

ggcccagccg gccatggccg agcccaaata ttgtgacaaa actcacacat gccaccgctg      60
cccagcacct gaactcctgg ggggaccgtc agtcttcttc tccccccaa aaccaagga      120
caccctcatg atctcccgga ccctgaggt caccatgcgtg gtggtggacg tgagccacga      180
agaccctgag gtcaagtcca actggtacgt ggacggcgctg gaggtgcata atgccaagac      240
aaagccgcgg gaggagcagt acaacagcac gtaccgtgtg gtcagcgtcc tcaccgtcct      300
gcaccaggac tggctgaatg gcaaggagta caagtgaag gtctccaaca aagccctccc      360
agcccccatc gagaaaacca tctccaaagc caaagggcag cctcgagaac cacaggtgta      420
caccctgccc ccatctcggg atgagctgkm tkmtkmtcag gtcagcctga cctgcctggt      480
caaaggcttc tatccagcg acatcgccgt ggagtgggag agcaatgggc agccggagaa      540
caactacaag accacgcctc ccgtgctgga ctccgacggc tccttcttcc tctacagcaa      600
gtcacccgtg ggttctkmtk mtkmtkmtkm tkmtkmtkmt agcggcaggt ggkmtkmtgg      660
gaacgtcttc tcattgctcg tgatgcatga ggctctgcac aaccactaca cgcagaagag      720
cctctccctg tctccgggta aagcggccgc      750

```

```

<210> SEQ ID NO 22
<211> LENGTH: 738

```

-continued

```

<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: Fcab05
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (449)..(450)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (452)..(453)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (455)..(456)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (611)..(612)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (614)..(615)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (617)..(618)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (620)..(621)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (623)..(624)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (626)..(627)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (629)..(630)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (632)..(633)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (641)..(642)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (644)..(645)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 22

ggccccagcgc gccatggcgc agcccaaatc ttgtgacaaa actcacacat gccaccgctg      60
cccagcacct gaactcctgg ggggaccgtc agtcttcttc tccccccaa aaccaagga      120
caccctcatg atctcccgga ccctgaggt ccatgcgtg gtggtggacg tgagccacga      180
agaccctgag gtcaagtcca actggtacgt ggacggcgtg gaggtgcata atgccaagac      240
aaagccgcgc gaggagcagt acaacagcac gtaccgtgtg gtcagcgtcc tcaccgtcct      300
gcaccaggac tggtggaatg gcaaggagta caagtgaag gtctccaaca aagccctccc      360
agcccccatc gagaaaacca tctccaaagc caaagggcag cctcgagaac cacaggtgta      420
caccctgccc ccattccgtg atgagkmtnn bnnbnnbkmt gtcagcctga cctgcctggt      480
caaaggcttc tatccagcg acatcgccgt ggagtgggag agcaatgggc agccggagaa      540
caactacaag accacgcctc ccgtgctgga ctccgacggc tccttcttcc tctacagcaa      600

```

-continued

gctcaccgtg nnnnnnnnnn nnnnnnnnnn bnnbaggtgg nnnnnbggga acgtcttctc	660
atgctccgtg atgcatgagg ctctgcacaa ccactacacg cagaagagcc tctccctgtc	720
tccgggtaaa gcggccgc	738

<210> SEQ ID NO 23
 <211> LENGTH: 738
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Fcab06

<400> SEQUENCE: 23

ggcccagccg gccatggccg agcccaaatc ttgtgacaaa actcacacat gccaccgtg	60
cccagcacct gaactcctgg ggggaccgtc agtcttctc tccccccaa aaccaagga	120
cacctcatg atctcccgga ccctgaggt cacatgcgtg gtggtggacg tgagccacga	180
agacctgag gtcaagtcca actggtacgt ggacggcgtg gaggtgcata atgccaagac	240
aaagccgcgg gaggagcagt acaacagcac gtaccgtgtg gtcagcgtcc tcaccgtcct	300
gcaccaggac tggctgaatg gcaaggagta caagtgaag gtctccaaca aagccctccc	360
agcccccatc gagaaaacca tctccaaagc caaagggcag cctcgagaac cacaggtgta	420
cacctgtccc ccctccggg acgagkmtkm tkmtkmtkmt gtcagcctga cctgcctggt	480
caaaggcttc tatccagcg acatcgccgt ggagtgggag agcaatgggc agccggagaa	540
caactacaag accacgcctc ccgtgctgga ctccgacggc tccttcttcc tctacagcaa	600
gctcaccgtg kmtkmtkmtk mtktkmtkmt tkmtagggtg kmtkmtggga acgtcttctc	660
atgctccgtg atgcatgagg ctctgcacaa ccactacacg cagaagagcc tctccctgtc	720
tccgggtaaa gcggccgc	738

<210> SEQ ID NO 24
 <211> LENGTH: 85
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer gapch35

<400> SEQUENCE: 24

caacaaggcc ctgctgtccc ccatcgagaa gaccatctcc aaggccaagg gccagcctcg	60
agaaccacag gtgtacaccc tgccc	85

<210> SEQ ID NO 25
 <211> LENGTH: 99
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer gapfcs3

<400> SEQUENCE: 25

gagaccgagg agagggttag ggataggctt accttcgaag ggcctctag actcgatcga	60
gcggccgctc atttaccgg agacaggag aggcctctc	99

<210> SEQ ID NO 26
 <211> LENGTH: 79
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: primer Abmut
 <220> FEATURE:
 <221> NAME/KEY: misc_feature

-continued

```

<222> LOCATION: (43)..(44)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (46)..(47)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (49)..(50)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 26

gaaccacagg tgtacaccct gcccccatcc cgggatgagc tgnnbnnbnn bcaggtcagc      60
ctgacctgcc tgggtcaaag                                                    79

<210> SEQ ID NO 27
<211> LENGTH: 79
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: primer Abmut2LR
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (40)..(41)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (43)..(44)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (46)..(47)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (49)..(50)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (52)..(53)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 27

gaaccacagg tgtacaccct gcccccatcc cgggatgagn nbnnbnnbnn bnnbgtcagc      60
ctgacctgcc tgggtcaaag                                                    79

<210> SEQ ID NO 28
<211> LENGTH: 79
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: primer Abmut1L
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (40)..(41)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (43)..(44)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (46)..(47)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (49)..(50)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 28

gaaccacagg tgtacaccct gcccccatcc cgggatgagn nbnnbnnbnn bcaggtcagc      60

```


-continued

ctgacctgcc tggtaaag

79

<210> SEQ ID NO 29
 <211> LENGTH: 79
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: primer Abmut1R
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (43)..(44)
 <223> OTHER INFORMATION: n is a, c, g, or t
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (46)..(47)
 <223> OTHER INFORMATION: n is a, c, g, or t
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (49)..(50)
 <223> OTHER INFORMATION: n is a, c, g, or t
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (52)..(53)
 <223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 29

gaaccacagg tgtacacct gcccccatcc cgggatgagc tgnnbnnbnn bnnbgtcagc

60

ctgacctgcc tggtaaag

79

<210> SEQ ID NO 30
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 30

Leu Asp Asn Ser Gln
 1 5

<210> SEQ ID NO 31
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 31

Tyr Glu Gly Ser Ser
 1 5

<210> SEQ ID NO 32
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 32

Tyr Met Ser Ala Asp
 1 5

<210> SEQ ID NO 33
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: AB-loop sequence AA143ff

-continued

<400> SEQUENCE: 33

Tyr Arg Arg Gly Asp
1 5

<210> SEQ ID NO 34
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 34

Leu Met Ser Arg Gln
1 5

<210> SEQ ID NO 35
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 35

Leu His Leu Ala Gln
1 5

<210> SEQ ID NO 36
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 36

Tyr Leu Ser Lys Asp
1 5

<210> SEQ ID NO 37
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 37

Tyr Arg Ser Gly Ser
1 5

<210> SEQ ID NO 38
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 38

Leu Arg Asp Gly Gln
1 5

<210> SEQ ID NO 39
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 39

-continued

Tyr Ser Ala Asn Thr
1 5

<210> SEQ ID NO 40
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 40

Tyr Ala Ser Asn Thr
1 5

<210> SEQ ID NO 41
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 41

Tyr Ser Asp Gly Asp
1 5

<210> SEQ ID NO 42
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 42

Tyr Ser Gly Gly Ser
1 5

<210> SEQ ID NO 43
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 43

Tyr Gly Arg Asp Ser
1 5

<210> SEQ ID NO 44
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 44

Tyr Ala Gly Gly Thr
1 5

<210> SEQ ID NO 45
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 45

-continued

Tyr Ser Ser Asp Ser
1 5

<210> SEQ ID NO 46
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 46

Tyr His Ser Gly Ser
1 5

<210> SEQ ID NO 47
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 47

Tyr Leu Thr Asn Ser
1 5

<210> SEQ ID NO 48
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 48

Tyr Gly Ser Glu Glu
1 5

<210> SEQ ID NO 49
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 49

Tyr Arg Ser Gly Glu
1 5

<210> SEQ ID NO 50
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 50

Tyr Gly Thr Asp Asp
1 5

<210> SEQ ID NO 51
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 51

Ile Arg Ser Ser Val Gly Ser Arg Arg Trp Trp Ser

-continued

1 5 10

<210> SEQ ID NO 52
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 52

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
 1 5 10

<210> SEQ ID NO 53
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 53

Ser Arg Arg Asp Ser Ser Leu Leu Arg Trp Ala His
 1 5 10

<210> SEQ ID NO 54
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 54

Ala Pro Gly Ser Lys Gly Tyr Arg Arg Trp Ala Leu
 1 5 10

<210> SEQ ID NO 55
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 55

Asp Lys Pro Phe Trp Gly Thr Ser Arg Trp Ser Arg
 1 5 10

<210> SEQ ID NO 56
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 56

Ser Ile Asn Asp Leu Ile Asn His Arg Trp Pro Tyr
 1 5 10

<210> SEQ ID NO 57
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 57

Met Trp Gly Ser Arg Asp Tyr Trp Arg Trp Ser His
 1 5 10

-continued

<210> SEQ ID NO 58
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 58

Asn Ser Gly Ser Ala Met Met Val Arg Trp Ala His
1 5 10

<210> SEQ ID NO 59
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 59

Gln Arg Ser Arg Leu Ser Arg Gln Arg Trp Trp Arg
1 5 10

<210> SEQ ID NO 60
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 60

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

<210> SEQ ID NO 61
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 61

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

<210> SEQ ID NO 62
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 62

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

<210> SEQ ID NO 63
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 63

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

-continued

<210> SEQ ID NO 64
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 64

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

<210> SEQ ID NO 65
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 65

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

<210> SEQ ID NO 66
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 66

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

<210> SEQ ID NO 67
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 67

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

<210> SEQ ID NO 68
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 68

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

<210> SEQ ID NO 69
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

<400> SEQUENCE: 69

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1 5 10

-continued

```

<210> SEQ ID NO 70
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

```

```

<400> SEQUENCE: 70

```

```

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1           5           10

```

```

<210> SEQ ID NO 71
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF-loop sequence AA198ff

```

```

<400> SEQUENCE: 71

```

```

Ala Arg Tyr Ser Pro Arg Met Leu Arg Trp Ala His
1           5           10

```

```

<210> SEQ ID NO 72
<211> LENGTH: 5009
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: vector pYD1

```

```

<400> SEQUENCE: 72

```

```

acggattaga agccgccgag cgggtgacag cctccgaag gaagactctc ctccgtgcgt      60
cctcgctctc accggtcgcg ttccctgaaac gcagatgtgc ctccgccgcg actgctccga    120
acaataaaga ttctacaata ctgactttta tggttatgaa gaggaataat tggcagtaac    180
ctggccccac aaacctcaa atgaacgaat caaatataca accataggat gataatgcga    240
ttagtttttt agccttattt ctggggtaat taatcagcga agcgatgatt ttgtatctat    300
taacagatat ataaatgcaa aaactgcata accactttta ctaatacttt caacattttc    360
ggtttgtatt acttcttatt caaatgtaat aaaagtatca acaaaaaatt gttaatatatac  420
ctctatactt taacgtcaag gagaaaaaac cccggatcgg actactagca gctgtaatac    480
gactcactat agggaatatt aagctaattc tacttcatat attttcaatt aagatgcagt    540
tacttcgctg tttttcaata ttttctgtta ttgcttcagt tttagcacag gaactgacaa    600
ctatatgcga gcaaatcccc tcaccaactt tagaatcgac gccgtactct ttgtcaacga    660
ctactatttt ggccaacggg aaggcaatgc aaggagtttt tgaatattac aaatcagtaa    720
cgtttgtcag taattgcggg tctcaccctt caacaactag caaaggcagc ccataaaca    780
cacagtatgt ttttaagctt ctgcaggcta gtggtggtgg tggttctggg ggtggtggtt    840
ctggtggtgg tggttctgct agcatgactg gtggacagca aatgggtcgg gatctgtacg    900
acgatgacga taaggtaacca ggatccagtg tggtggaatt ctgcagatat ccagcacagt    960
ggcggccgct cgagtctaga gggcccttcg aaggtaagcc tatccctaac cctctcctcg  1020
gtctcgattc tacgcgtacc ggtcatcatc accatcacca ttgagttaa acccgctgat    1080
ctgataacaa cagtgtagat gtaacaaaat cgactttgtt cccactgtac ttttagctcg    1140
tacaaaatac aatatacttt tcattttctc gtaaacacaa tgttttccca tgtaatatcc    1200
ttttctattt ttctgtccgt taccaacttt acacatactt tatatageta ttcacttcta    1260
tacactaaaa aactaagaca attttaattt tgctgcctgc catatttcaa tttgttataa    1320

```


-continued

attcctataa tttatcctat tagtagctaa aaaaagatga atgtgaatcg aatcctaaga	1380
gaattgggca agtgcacaaa caatacttaa ataaatacta ctacagtaata acctatttct	1440
tagcattttt gacgaaatgt gctattttgt tagagtcctt tacaccattt gtctccacac	1500
ctccgcttac atcaacacca ataacgccat ttaatctaag cgcacacca acattttctg	1560
gcgtcagtc accagctaac ataaaatgta agctctcggg gctctcttgc ctccaaccc	1620
agtcagaaat cgagttccaa tccaaaagt cactgtccc acctgcttct gaatacaaca	1680
agggataaaa cgaatgaggt ttctgtgaag ctgcactgag tagtatgttg cagtcttttg	1740
gaaatacgag tcttttaata actggcaaac cgaggaactc ttggtattct tgccacgact	1800
catctccgtg cagttggacg atataaatgc cgtaatcatt gaccagagcc aaaacatcct	1860
ccttaggttg attacgaaac acgccaacca agtatttcgg agtgccctgaa ctatttttat	1920
atgcttttac aagacttgaa attttccttg caataaccgg gtcaattgtt ctctttctat	1980
tgggcacaca tataataccc agcaagtcag catcggaatc tagagcacat tctgcggcct	2040
ctgtgctctg caagccgcaa actttcacca atggaccaga actacctgtg aaattaataa	2100
cagacatact ccaagctgcc ttgtgtgtct taatcacgta tactcacgtg ctcaatagtc	2160
accaatgccc tccctcttgg cctctcctt ttcttttttc gaccgaattt cttgaagacg	2220
aaagggcctc gtgatacgcc tatttttata ggtaaatgtc atgataataa tggtttctta	2280
ggacggatcg cttgcctgta acttacacgc gcctcgatc ttttaatgat ggaataattt	2340
gggaatttac tctgtgttta ttatttttta tgttttgtat ttggatttta gaaagtaaat	2400
aaagaaggta gaagagttac ggaatgaaga aaaaaaata aacaaagggt taaaaattt	2460
caacaaaaag cgtactttac atatatattt attagacaag aaaagcagat taaatagata	2520
tacattcgat taacgataag taaaatgtaa aatcacagga ttttcgtgtg tggctctcta	2580
cacagacaag atgaaacaat tcggcattaa tacctgagag caggaagagc aagataaaag	2640
gtagtatttg ttggcgatcc ccttagagtc ttttacatct tcggaaaaca aaaactattt	2700
ttcttttaat ttcttttttt actttctatt tttaatttat atattttatat taaaaattt	2760
aaattataat tatttttata gcacgtgatg aaaaggaccc aggtggcact ttccggggaa	2820
atgtgcgcgg aacccttatt tgtttatttt tctaaataca ttcaaatatg tatccgctca	2880
tgagacaata acctgataa atgcttcaat aatattgaaa aaggaagagt atgagtattc	2940
aacatttcog tgcgcctctt attccctttt ttgcggcatt ttgccttctt gtttttgctc	3000
accagaaaac gctggtgaaa gtaaaagatg ctgaagatca gttgggtgca cgagtgggtt	3060
acatcgaaact ggtatctcaac agcggtaaga tccctgagag ttttcgcccc gaagaacgtt	3120
ttccaatgat gagcactttt aaagtctctg tatgtggcgc ggtattatcc cgtgttgacg	3180
cgggcaaga gcaactcggg cgcgcatac actatttcca gaatgacttg gttgagtact	3240
caccagtcac agaaaagcat cttacggatg gcatgacagt aagagaatta tgcagtgtg	3300
ccataaccat gagtgataac actgcggcca acttacttct gacaacgacg ggaggaccga	3360
aggagctaac cgcttttttg cacaacatgg gggatcatgt aactgcctt gatcgttggg	3420
aaccggagct gaatgaagcc ataccaaagc acgagcgtga caccacgatg cctgtagcaa	3480
tggcaacaac gttgcgcaaa ctattaactg gcgaactact tactctagct tcccggcaac	3540
aattaataga ctggatggag gcggataaag ttgcaggacc acttctgcgc tcggcccttc	3600
cggctggctg gtttatgtct gataaatctg gagccggtga gcgtgggtct cgcggtatca	3660
ttgcagcact ggggccagat ggtaagccct cccgtatcgt agttatctac acgacgggca	3720

-continued

```

gtcaggcaac tatggatgaa cgaaatagac agatcgctga gatagggtgcc tcaactgatta 3780
agcattggta actgtcagac caagtttact catatatact ttagattgat ttaaaacttc 3840
atttttaatt taaaaggatc taggtgaaga tcctttttga taatctcatg accaaaaatcc 3900
cttaacgtga gttttcgctc cactgagcgt cagaccccggt agaaaagatc aaaggatcctt 3960
cttgagatcc tttttttctg cgcgtaactc gctgcttgca aacaaaaaaa ccaccgctac 4020
cagcggtgggt ttgtttgcgc gatcaagagc taccaactct ttttcggaag gtaactggct 4080
tcagcagagc gcagatacca aatactgtcc ttctagtgtg gccgtagtta ggccaccact 4140
tcaagaactc tgtagcaccg cctacatacc tcgctctgct aatcctgtta ccagtggctg 4200
ctgccagtgg cgataagtcg tgtcttaccg ggttggaactc aagacgatag ttaccggata 4260
aggcgagcgc gtcgggctga acgggggggt cgtgcacaca gccagccttg gagcgaaacga 4320
cctacaccga actgagatac ctacagcgtg agcattgaga aagcgccacg cttcccgaag 4380
ggagaaaaggc ggacaggtat ccggtgaagc gcagggtcgg aacaggagag cgcacgaggg 4440
agcttccagg ggggaacgcc tggtatcttt atagtctgtg cgggtttcgc cactctgac 4500
ttgagcgtcg atttttgtga tgctcgtcag gggggccgag cctatggaaa aacgccagca 4560
acgcggcctt tttacggttc ctggcctttt gctggccttt tgctcactg ttctttcctg 4620
cgttatcccc tgattctgtg gataaccgta ttaccgcctt tgagtgaact gataccgctc 4680
gccgcagccg aacgaccgag cgcagcaggt cagtgaagca ggaagcggaa gagcgcccaa 4740
tacgcaaacc gcctctcccc gcgcgttggt cgattcatta atgcagctgg cagcagaggt 4800
ttcccgactg gaaagcgggc agtgagcga acgcaattaa tgtgagttac ctcaactatt 4860
aggcacccca ggcttttacac tttatgcttc cggctcctat gttgtgtgga attgtgagcg 4920
gataacaatt tcacacagga aacagctatg accatgatta cgccaagctc ggaattaacc 4980
ctcactaaag ggaacaaaag ctggctagt 5009

```

```

<210> SEQ ID NO 73
<211> LENGTH: 5009
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: modified vector pYD1Nhe

```

```

<400> SEQUENCE: 73

```

```

acggattaga agccgcgcgag cgggtgacag cctccgaag gaagactctc ctccgtgcgt 60
cctcgtcttc accggtcgcg ttctgaaac gcagatgtgc ctccgcgcgc actgctccga 120
acaataaaga ttctacaata ctgactttta tggttatgaa gaggaataat tggcagtaac 180
ctggcccccac aaaccttcaa atgaacgaat caaattaaca accataggat gataatgcga 240
ttagtttttt agccttattt ctggggtaat taatcagcga agcagtgatt ttgatctat 300
taacagatat ataaatgcaa aaactgcata accactttaa ctaatacttt caacattttc 360
ggtttgtatt acttcttatt caaatgtaat aaaagtatca acaaaaaatt gttaatatat 420
ctctatactt taacgtcaag gagaaaaaac cccggatcgg actactagca gctgtaatac 480
gactcactat agggaaatatt aagctaattc tacttcatac attttcaatt aagatgcagt 540
tacttcgctg tttttcaata ttttctgtta ttgcttcagt gctagcacag gaactgacaa 600
ctatatgcga gcaaatcccc tcaccaactt tagaatcgac gccgtactct ttgtcaacga 660
ctactatttt ggccaacggg aaggcaatgc aaggagtttt tgaatattac aaatcagtaa 720

```

-continued

cgtttgtag	taattgcggt	tctcaccct	caacaactag	caaaggcagc	cccataaaca	780
cacagtatgt	ttttaagctt	ctgcaggcta	gtggtgggtg	tggttctggt	ggtgggtggt	840
ctggtgggtg	tggttctgct	agcatgactg	gtggacagca	aatgggtcgg	gatctgtacg	900
acgatgacga	taaggtagca	ggatccagtg	tggtggaatt	ctgcagatat	ccagcacagt	960
ggcgcccgct	cgagtctaga	gggcccttcg	aaggtaagcc	tatccctaac	cctctcctcg	1020
gtctcgattc	tacgcgtacc	ggatcatc	accatcacca	ttgagtttaa	accgctgat	1080
ctgataacaa	cagtgtagat	gtaacaaaat	cgactttggt	cccactgtac	tttagctcg	1140
tacaaaat	aataactttt	tctttctctc	gtaaacaaca	tggtttccca	tgtaatatcc	1200
ttttctat	ttcggttcgt	taccaacttt	acacatactt	tatatagcta	ttcacttcta	1260
tacactaaaa	aactaagaca	attttaattt	tgctgctg	catatttcaa	ttgttataa	1320
attctataa	tttatctat	tagtagctaa	aaaaagatga	atgtgaatcg	aatcctaaga	1380
gaattgggca	agtgacacaa	caatacttaa	ataaatacta	ctcagtaata	acctatttct	1440
tagcattttt	gacgaaat	gctattttgt	tagagtcttt	tacaccattt	gtctccacac	1500
ctccgcttac	atcaacacca	ataacgccat	ttaatcctaag	cgcatcacca	acattttctg	1560
gogtcagtc	accagctaac	ataaaatgta	agctctcggt	gctctcttgc	cttccaaacc	1620
agtcagaaat	cgagttccaa	tccaaaagtt	cacctgtccc	acctgcttct	gaatcaaaca	1680
aggggaataa	cgaatgaggt	tctgtgaag	ctgcactgag	tagtatgttg	cagtcttttg	1740
gaaatacgag	tcttttaata	actggcaaac	cgaggaactc	ttggtattct	tgccacgact	1800
catctccgtg	cagttggacg	atatcaatgc	cgtaatcatt	gaccagagcc	aaaacatcct	1860
ccttaggttg	attacgaac	acgccaacca	agtatttcgg	agtgccctgaa	ctatttttat	1920
atgcttttac	aagacttgaa	attttccttg	caataaccgg	gtcaattggt	ctctttctat	1980
tgggcacaca	tataataccc	agcaagtcag	catcggaatc	tagagcacat	tctgcggcct	2040
ctgtgctctg	caagccgcaa	actttcacca	atggaccaga	actacctgtg	aaattaataa	2100
cagacatact	ccaagctgcc	tttgtgtgct	taatcacgta	tactcacgtg	ctcaatagtc	2160
accaatgccc	tccctcttgg	ccctctcctt	ttcttttttc	gaccgaattt	cttgaagacg	2220
aaagggcctc	gtgatacgcc	tatttttata	ggttaatgtc	atgataataa	tggtttctta	2280
ggacggatcg	cttgccgtga	acttacacgc	gcctcgatc	ttttaatgat	ggaataattt	2340
gggaatttac	tctgtgttta	tttattttta	tggtttgtat	ttggatttta	gaaagtaaat	2400
aaagaaggtg	gaagagttac	ggaatgaaga	aaaaaaaata	aacaaagggt	taaaaaattt	2460
caacaaaaag	cgtagctttac	atatatat	attagacaag	aaaagcagat	taaatagata	2520
tacattcgat	taacgataag	taaaatgtaa	aatcacagga	tttctgtgtg	tggtcttcta	2580
cacagacaag	atgaacaat	tcggcattaa	tacctgagag	caggaagagc	aagataaaaag	2640
gtagtatttg	ttggcgatcc	ccctagagtc	ttttacatct	tcggaaaaaca	aaaactattt	2700
tttctttaat	ttcttttttt	actttctatt	tttaatttat	atatttatat	taaaaaattt	2760
aaattataat	tatttttata	gcacgtgatg	aaaaggaccc	aggtggcact	tttcggggaa	2820
atgtgcgcgg	aaccctattt	tggtttattt	tctaaatata	ttcaaatatg	tatccgctca	2880
tgagacaata	acctgataa	atgcttcaat	aatattgaaa	aaggaagagt	atgagtattc	2940
aacatttcgg	tgtagccctt	attccctttt	ttgcggcatt	ttgccttctc	gtttttgctc	3000
accagaaaac	gctggtgaaa	gtaaaagatg	ctgaagatca	gttgggtgca	cgagtgggtt	3060
acatcgaaat	ggatctcaac	agcggtaaga	tccttgagag	ttttcgcccc	gaagaacggt	3120

-continued

```

ttccaatgat gagcactttt aaagttctgc tatgtggcgc ggtattatcc cgtgttgacg 3180
ccgggcaaga gcaactcggc cgccgcatac actattctca gaatgacttg gttgagtact 3240
caccagtcac agaaaagcat cttacggatg gcatgacagt aagagaatta tgcagtgctg 3300
ccataaccat gagtgataac actgcggcca acttacttct gacaacgacg ggaggaccga 3360
aggagctaac cgcttttttg cacaacatgg gggatcatgt aactcgctt gatcgttggg 3420
aaccggagct gaatgaagcc ataccaaacg acgagcgtga caccacgacg cctgtagcaa 3480
tggcaacaac gttgcgcaaa ctattaactg gcgaactact tactctagct tcccgcaac 3540
aattaataga ctggatggag gcggataaag ttgcaggacc acttctgcgc tcggcccttc 3600
cggctggctg gttttattgct gataaatctg gagccgggtg gcgtgggtct cgcggtatca 3660
ttgcagcact ggggccagat ggtaagccct ccgctatcgt agttatctac acgacgggca 3720
gtcaggcaac tatggatgaa cgaaatagac agatcgtga gatagggtgcc tcaactgatta 3780
agcattggta actgtcagac caagtttact catatatact ttagattgat ttaaaacttc 3840
atthtttaatt taaaaggatc taggtgaaga tcctttttga taatctcatg accaaaatcc 3900
cttaacgtga gttttcgttc cactgagcgt cagaccccg agaaaagatc aaaggatctt 3960
cttgagatcc tttttttctg cgcgtaatct gctgcttgca aacaaaaaaa ccaccgctac 4020
cagcgggtggt ttgtttgcg gatcaagagc taccaactct ttttcgaag gtaactggct 4080
tcagcagagc gcagatacca aatactgtcc ttctagtgtg gccgtagtta ggccaccact 4140
tcaagaactc tgtagcacgc cctacatacc tcgctctgct aatcctgtta ccagtggctg 4200
ctgccagtgg cgataagtgc tgtcttaccg ggttggactc aagacgatag ttaccggata 4260
aggcgcagcg gtcgggctga acgggggggt cgtgcacaca gccagcttg gagcgaaacga 4320
cctacaccga actgagatac ctacagcgtg agcattgaga aagcgccacg cttcccgaag 4380
ggagaaaggc ggacaggtat ccggtaagcg gcagggtcgg aacaggagag cgcacgaggg 4440
agcttccagg ggggaacgcc tggatatctt atagtctgt cggttttcgc cactctgac 4500
ttgagcgtcg attttttgta tgctcgtcag gggggccgag cctatggaaa aacgccagca 4560
acggcgctt tttacggttc ctggcctttt gctggccttt tgctcacatg ttctttctg 4620
cgttatcccc tgattctgtg gataaccgta ttaccgcctt tgagttagct gataccgctc 4680
gccgcagcgc aacgaccgag cgcagcagat cagttagcga ggaagcggaa gagcgcccaa 4740
tacgcaaaac gcctctcccc gcgcgttggc cgattcatta atgcagctgg cagcacaggt 4800
ttcccgaactg gaaagcgggc agtgagcgca acgcaattaa tgtgagttac ctactcatt 4860
aggcacccca ggctttacac tttatgcttc cggctcctat gttgtgtgga attgtgagcg 4920
gataacaatt tcacacagga aacagctatg accatgatta cgccaagctc ggaattaacc 4980
ctcactaaag ggaacaaaag ctggctagt 5009

```

```

<210> SEQ ID NO 74
<211> LENGTH: 4605
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: vector pYD1lnk

```

```

<400> SEQUENCE: 74

```

```

acggattaga agccgcgag cgggtgacag cctccgaag gaagactctc ctccgtgcgt 60
ctcgtcttc accggtcgcg ttctgaaac gcagatgtgc ctgcgcgcg actgctccga 120

```

-continued

acaataaaga	ttctacaata	ctagctttta	tggttatgaa	gaggaaaaat	tggcagtaac	180
ctggcccccac	aaaccttcaa	atgaacgaat	caaattaaca	accataggat	gataatgcga	240
ttagttttttt	agccttattt	ctggggtaat	taatcagcga	agcgatgatt	tttgatctat	300
taacagatat	ataaatgcaa	aaactgcata	accactttaa	ctaatacttt	caacattttc	360
ggtttgtatt	acttcttatt	caaagtgaat	aaaagtatca	acaaaaaatt	gttaatatatac	420
ctctatactt	taacgtcaag	gagaaaaaac	cccgatcgg	actactagca	gctgtaatac	480
gactcactat	agggaaatatt	aagctaattc	tacttcatac	atthttcaatt	aagatgcagt	540
tacttcgctg	tttttcaata	ttttctgtta	ttgcttcagt	gctagccgct	ggggccatgg	600
ttactgattg	gcgcgcggga	tccttcagac	tctgcagaat	tcggccgcgt	acttaattaa	660
gtttaaaccc	gctgatctga	taacaacagt	gtagatgtaa	caaaatcgac	tttgttccca	720
ctgtactttt	agctcgtaca	aaatacaata	tacttttcat	ttctccgtaa	acaacatggt	780
ttcccatgta	atatcctttt	ctatttttctg	ttccgttacc	aactttacac	atactttata	840
tagctattca	cttctatata	ctaaaaaact	aagacaattt	taattttgct	gcctgccata	900
tttcaatttg	ttataaattc	ctataattta	tcctattagt	agctaaaaaa	agatgaatgt	960
gaatcgaatc	ctaagagaat	tgggcaagt	cacaaacaat	acttaataaa	atactactca	1020
gtaataacct	atttcttagc	atttttgacg	aaatttgcta	ttttgttaga	gtcttttaca	1080
ccatttgtct	ccacacctcc	gcttacatca	acaccaataa	cgccatttaa	tctaagcgca	1140
tcaccaacat	tttctggcgt	cagtcaccca	gctaacataa	aatgtaagct	ctcggggctc	1200
tcttgccctc	caaccagtc	agaaatcgag	ttccaatcca	aaagttcacc	tgtcccacct	1260
gcttctgaat	caaacaagg	aataaacgaa	tgaggtttct	gtgaagctgc	actgagtagt	1320
atgttgtagt	cttttggaat	tacgagtcct	ttaataactg	gcaaacggag	gaactcttgg	1380
tattcttgcc	acgactcatc	tcctgtagt	tggaagatata	caatgccgta	atcattgacc	1440
agagccaaaa	catcctcctt	agggttgatta	cgaaacacgc	caaccaagta	tttcggagtg	1500
cctgaactat	ttttatatgc	ttttacaaga	cttgaaattt	tccttgcaat	aacgggtgca	1560
attgttctct	ttctattggg	cacacatata	ataccagca	agtcagcatc	ggaatctaga	1620
gcacattctg	cggcctctgt	gctctgcaag	cgcgaactt	tcaccaatgg	accagaacta	1680
cctgtgaaat	taataacaga	catactccaa	gctgcctttg	tgtgcttaat	cacgtatact	1740
cacgtgctca	atagtcacca	atgccctccc	tcttgccct	ctccttttct	tttttcgacc	1800
gaatttcttg	aagacgaag	ggcctcgtga	tacgcctatt	tttatagggt	aatgtcatga	1860
taataatggg	ttcttaggac	ggatcgcttg	cctgtaactt	acacgcgcct	cgtatctttt	1920
aatgatggaa	taatttgga	atttactctg	tgtttattta	tttttatggt	ttgtatttgg	1980
atttttagaaa	gtaataaag	aaggtagaag	agttacggaa	tgaagaaaaa	aaaataaaca	2040
aagggtttaa	aaatttcaac	aaaaagcgta	ctttacatat	atatttatta	gacaagaaaa	2100
gcagattaaa	tagatatata	ttcgattaac	gataagtaaa	atgtaaaatc	acaggattht	2160
cgtgtgtggg	cttctacaca	gacaagatga	aacaattcgg	cattaatacc	tgagagcagg	2220
aagagcaaga	taaaaggtag	tatttggttg	cgatccccct	agagtctttt	acatcttcgg	2280
aaaacaaaaa	ctattttttc	tttaatttct	ttttttactt	tctattttta	atttatatat	2340
ttatatataa	aaattttaa	tataattatt	tttatagcac	gtgatgaaaa	ggaccaggt	2400
ggcacttttc	ggggaaatgt	gcgcggaacc	cctatttggt	tatttttcta	aatacattca	2460
aatatgtatc	cgctcatgag	acaataaccc	tgataaatgc	ttcaataata	ttgaaaaagg	2520

-continued

```

aagagtatga gtattcaaca ttccgtgtc gcccttattc ccttttttgc ggcattttgc 2580
cttcctgttt ttgctcacc cagaacgctg gtgaaagtaa aagatgctga agatcagttg 2640
gggtgcacgag tgggttacat cgaactggat ctcaacacgc gtaagatcct tgagagtttt 2700
cgccccgaag aacgttttcc aatgatgagc actttttaag ttctgctatg tggcgcggtg 2760
ttatcccgtg ttgacgcgg gcaagagcaa ctcggtcgcc gcatacacta ttctcagaat 2820
gacttggttg agtactcacc agtcacagaa aagcatctta cggatggcat gacagtaaga 2880
gaattatgca gtgctgccat aacctgagt gataaactg cggccaactt acttctgaca 2940
acgatcggag gaccgaagga gctaaccgct tttttgcaca acatggggga tcatgtaact 3000
cgccttgatc gttgggaacc ggagctgaat gaagccatc caaacgacga gcgtgacacc 3060
acgatgcctg tagcaatggc aacaacgttg cgcaactat taactggcga actacttact 3120
ctagcttccc ggcaacaatt aatagactgg atggaggcgg ataaagttgc aggaccactt 3180
ctgcgctcgg cccttcgggc tggttggttt attgctgata aatctggagc cggtgagcgt 3240
gggtctcgcg gtatcattgc agcactgggg ccagatggta agccctcccg tatcgtagtt 3300
atctacacga cgggcagtcg ggcaactatg gatgaacgaa atagacagat cgctgagata 3360
gggtgcctcac tgattaagca ttggtaactg tcagaccaag tttactcata tatactttag 3420
attgatttaa aacttcattt ttaatttaaa aggatctagg tgaagatcct ttttgataat 3480
ctcatgacca aaatccctta acgtgagttt tcgttcact gagcgtcaga ccccgtagaa 3540
aagatcaaag gatcttcttg agatcctttt tttctgcgcg taatctgctg cttgcaaaca 3600
aaaaaaccac cgctaccagc ggtggtttgt ttgcccgcgc aagagctacc aactcttttt 3660
ccgaaggtaa ctggtctcag cagagcgcag ataccaaata ctgtccttct agtgtagccg 3720
tagttaggcc accacttcaa gaactctgta gcaccgccta catacctcgc tctgctaata 3780
ctgttaccag tggtctgctg cagtggcgat aagtcgtgtc ttaccgggtt ggactcaaga 3840
cgatagttag cggataaggc gcagcggctg ggctgaacgg ggggttcgtg cacacagccc 3900
agcttgagagc gaacgaccta caccgaactg agatacctac agcgtgagca ttgagaaagc 3960
gccacgcttc ccgaaggagc aaaggcggac aggtatccgg taagcggcag ggtcggaaca 4020
ggagagcgca cgaggagct tccagggggg aacgcctggg atctttatag tcctgtcggg 4080
tttcgccacc tctgacttga gcgtcgattt ttgtgatgct cgtcaggggg gccgagccta 4140
tggaaaaaag ccagcaacgc ggccctttta cggttcctgg ccttttctg gctttttgct 4200
cacatgttct tcctgcggtt atcccctgat tctgtggata accgtattac cgcccttgag 4260
tgagctgata ccgctcgcgc cagccgaacg accgagcgca gcgagtcagt gagcgaggaa 4320
gcggaagagc gcccaatacg caaacgcct ctcccgcgcg gttggccgat tcattaatgc 4380
agctggcacg acaggtttcc cgactggaaa gcgggcagtg agcgcaacgc aattaatgtg 4440
agttacctca ctcattaggc accccaggct ttacacttta tgcttcgggc tcctatgttg 4500
tgtggaattg tgagcggata acaatttcac acaggaaaca gctatgacca tgattacgcc 4560
aagctcggaa ttaaccctca ctaaaggga caaaagctgg ctagt 4605

```

<210> SEQ ID NO 75

<211> LENGTH: 4886

<212> TYPE: DNA

<213> ORGANISM: artificial

<220> FEATURE:

<223> OTHER INFORMATION: vector pYD1mata

-continued

<400> SEQUENCE: 75

acggattaga agccgcgcgag cgggtgacag cctccgaag gaagactctc ctccgtgcgt	60
cctcgtcttc accggtcgcg ttctgaaac gcagatgtgc ctgcgcgcgc actgctccga	120
acaataaaga ttctacaata ctgactttta tggttatgaa gaggaataat tggcagtaac	180
ctggcccccac aaaccttcaa atgaacgaat caaatataca accataggat gataatgcga	240
ttagtttttt agccttattt ctggggtaat taatcagcga agcgatgatt ttgatctat	300
taacagatat ataaatgcaa aaactgcata accactttaa ctaatacttt caacattttc	360
ggtttgtatt acttcttatt caaatgtaat aaaagtatca acaaaaaatt gttaataatac	420
ctctataact taacgtcaag gagaaaaaac cccggatcgg actactagca gctgtaatac	480
gactcactat agggaaatatt aagctaattc tacttcatac attttcaatt aagatgcagt	540
tacttcgctg tttttcaata ttttctgtta ttgcttcagt gctagccgct ggggccatgg	600
ttactgattg gcgcgcgcga tccgatgtaa caaaatcgac tttgttccca ctgtactttt	660
agctcgtaca aaatacaata tacttttcat ttctccgtaa acaacatgtt ttcccatgta	720
atatcctttt ctatttttctg ttccgttacc aactttacac atactttata tagctattca	780
cttctataca ctaaaaaact aagacaattt taattttgct gcctgccata tttcaatttg	840
ttataaaatc ctataattta tcctattagt agctaaaaaa agatgaatgt gaatcgaatc	900
ctaagagaat tgctgcagaa ttccggccgcg tacttaatta agtttaaacc cgctgatctg	960
ataacaacag tgtagatgta acaaaatcga ctttgttccc actgtacttt tagctcgta	1020
aaaatacaat atacttttca tttctccgta aacaacatgt tttcccatgt aatatccttt	1080
tctatttttc gttccgttac caactttaca catactttat atagctattc acttctatac	1140
actaaaaaac taagacaatt ttaattttgc tgccctgcat atttcaattt gttataaatt	1200
cctataattt atcctattag tagctaaaaa aagatgaatg tgaatcgaat cctaagagaa	1260
ttgggcaagt gcacaaaaca tacttaataa aatactactc agtaataacc tatttcttag	1320
catttttgac gaaatttgct attttgttag agtcttttac accatttgtc tccacacctc	1380
cgcttacatc aacaccaata acgccattta atctaagcgc atcaccaaca ttttctggcg	1440
tcagtcacac agctaacata aaatgtaagc tctcggggct ctcttgccct ccaacccagt	1500
cagaaatcga gttccaatcc aaaagttcac ctgtcccacc tgcttctgaa tcaacaag	1560
gaataaacga atgaggtttc tgtgaagctg cactgagtag tatgttgacg tcttttgaa	1620
atacagagtct ttaataaact ggcaaacga ggaactcttg gtattcttgc cagcactcat	1680
ctccgtgcag ttggacgata tcaatgccgt aatcattgac cagagccaaa acatcctcct	1740
taggttgatt acgaaacacg ccaaccaagt atttcggagt gcctgaacta tttttatatg	1800
cttttacaag acttgaaatt ttccctgcaa taaccgggtc aattgttctc tttctatttg	1860
gcacacatat aatacccagc aagtcagcat cggaatctag agcacattct gcggcctctg	1920
tgctctgcaa gccgcaaaact ttcaccaatg gaccagaact acctgtgaaa ttaataacag	1980
acatactcca agctgccttt gtgtgcttaa tcacgtatac tcacgtgctc aatagtcacc	2040
aatgccctcc ctcttgggcc tctccttttc ttttttcgac cgaatttctt gaagacgaaa	2100
gggcctcgtg atacgcctat ttttataggt taatgtcatg ataataatgg tttcttagga	2160
cggatcgctt gcctgtaact tacacgcgc tcgtatcttt taatgatgga ataatttggg	2220
aatttactct gtgtttattt atttttatgt tttgtatttg gattttagaa agtaataaaa	2280
gaaggtagaa gagttacgga atgaagaaaa aaaaataaac aaaggtttta aaaatttcaa	2340

-continued

caaaaagcgt	actttacata	tatatattatt	agacaagaaa	agcagattaa	atagatatatac	2400
attcgattaa	cgataagtaa	aatgtaaaat	cacaggattt	tcgtgtgtgg	tcttctacac	2460
agacaagatg	aaacaattcg	gcattaatac	ctgagagcag	gaagagcaag	ataaaaggta	2520
gtatttggtg	gcgatccccc	tagagtcttt	tacatcttcg	gaaaacaaaa	actatTTTTT	2580
ctttaatttc	TTTTTTTact	tTctatTTTT	aatttatata	tttatattaa	aaaatttaaa	2640
ttataattat	TTTTatagca	cgtgatgaaa	aggacccagg	tggcactttt	cggggaaatg	2700
tgcgcgggaa	ccctatttgt	ttatTTTTct	aaatacattc	aaatatgtat	ccgctcatga	2760
gacaataacc	ctgataaatg	cttcaataat	attgaaaaag	gaagagtatg	agtattcaac	2820
atttccgtgt	cgccttattt	ccctTTTTtg	cggcattttg	ccttccctgt	tttgcctcac	2880
cagaaacgct	ggtgaaagta	aaagatgctg	aagatcagtt	gggtgcacga	gtgggttaca	2940
tcgaactgga	tctcaacagc	ggtaagatcc	ttgagagttt	tcgccccgaa	gaacgttttc	3000
caatgatgag	cacttttaaa	gttctgctat	gtggcgcggt	attatcccg	gttgacgccc	3060
ggcaagagca	actcggctgc	cgcatacact	attctcagaa	tgacttggtt	gagtactcac	3120
cagtcacaga	aaagcatctt	acggatggca	tgacagtaag	agaattatgc	agtgcctcca	3180
taaccatgag	tgataaacact	gcggccaact	tacttctgac	aacgatcggg	ggaccgaagg	3240
agctaaccgc	TTTTTTgcac	aacatggggg	atcatgtaac	tcgccttgat	cgttgggaa	3300
cggagctgaa	tgaagccata	ccaaacgacg	agcgtgacac	cacgatgcct	gtagcaatgg	3360
caacaacggt	gcgcaaaact	ttaactggcg	aactacttac	tctagcttcc	cggaacaat	3420
taatagactg	gatggaggcg	gataaagttg	caggaccact	tctgcgctcg	gcccttcggt	3480
ctggctgggt	tattgctgat	aaatctggag	ccggtgagcg	tgggtctcgc	ggtatcattg	3540
cagcactggg	gccagatggg	aagccctccc	gtatcgtagt	tatctacacg	acgggcagtc	3600
aggcaactat	ggatgaacga	aatagacaga	tcgctgagat	aggtgcctca	ctgattaagc	3660
attggtaact	gtcagaccaa	gtttactcat	atatacttta	gattgattta	aaacttcatt	3720
tttaatttaa	aaggatctag	gtgaagatcc	TTTTtgataa	tctcatgacc	aaaatccctt	3780
aacgtgagtt	ttcgtttccac	tgagcgtcag	accccgtaga	aaagatcaaa	ggatcttctt	3840
gagatccttt	TTTTctgcgc	gtaatctgct	gcttgcaaac	aaaaaaacca	ccgctaccag	3900
cgggtgggtt	tttgccggat	caagagctac	caactctttt	tccgaaggta	actggcttca	3960
gcagagcgca	gataccaaat	actgtccttc	tagtgtagcc	gtagttaggc	caccacttca	4020
agaactctgt	agcaccgcct	acatacctcg	ctctgctaat	cctgttacca	gtggctgctg	4080
ccagtggcga	taagtctgtg	cttaccgggt	tggactcaag	acgatagtta	ccggataagg	4140
cgcagcggtc	gggctgaacg	gggggttcgt	gcacacagcc	cagcttgagg	cgaacgacct	4200
acaccgaact	gagataccta	cagcgtgagc	attgagaaag	cgccacgctt	cccgaaggga	4260
gaaaggcgga	caggatatccg	gtaagcggca	gggtcgggaa	aggagagcgc	acgaggggag	4320
ttccaggggg	gaacgcctgg	tatctttata	gtcctgtcgg	gtttgcgca	ctctgacttg	4380
agcgtcgatt	tttgtgatgc	tcgtcagggg	ggccgagcct	atggaaaaac	gccagcaacg	4440
cggccttttt	acggtttctg	gcctttttgt	ggccttttgc	tcacatgttc	tttctgcgt	4500
tatccccga	ttctgtggat	aaccgtatta	ccgcctttga	gtgagctgat	accgctcgcc	4560
gcagccgaac	gaccgagcgc	agcgagtcag	tgagcgagga	agcgggaagag	cgcccaatac	4620
gcaaaccgcc	tctccccgcg	cgttggccga	ttcattaatg	cagctggcac	gacagggttc	4680

-continued

```

ccgactggaa agcgggcagc gagcgcaacg caattaatgt gagttacctc actcattagg 4740
caccgccaggc ttctacacttt atgcttcocgg ctccatgtgt gtgtggaatt gtgagcggat 4800
aacaatttca cacaggaaac agctatgacc atgattacgc caagctcggg attaacccctc 4860
actaaaggga acaaaagctg gctagt 4886

```

```

<210> SEQ ID NO 76
<211> LENGTH: 5801
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: vector pYD1gal

```

```

<400> SEQUENCE: 76

```

```

acggattaga agccgccgag cgggtgacag ccctccgaag gaagactctc ctccgtgcgt 60
cctcgtcttc accggtcgcg ttctgaaac gcagatgtgc ctgcgccgc actgctccga 120
acaataaaga ttctacaata ctacttttta tggttatgaa gaggaataat tggcagtaac 180
ctggccccac aaaccttcaa atgaacgaat caaattaaca accataggat gataatgcga 240
ttagtttttt agccttattt ctggggtaat taatcagcga agcgatgatt ttgatctat 300
taacagatat ataatgcaa aaactgcata accactttaa ctaatacttt caacattttc 360
ggtttgattt acttcttatt caaatgtaat aaaagtatca acaaaaaatt gttaatatat 420
ctctatactt taacgtcaag gagaaaaaac ccggatcgg actactagca gctgtaatac 480
gactcactat agggaaatatt aagctaattc tacttcatac attttcaatt aagatgcagt 540
tacttcgctg tttttcaata ttttctgtta ttgcttcagt gctagccgct ggggccatgg 600
ttactgattg gcgcgccgga tccgatgtaa caaatcgac tttgttccca ctgtactttt 660
agctcgtaca aaatacaata tacttttcat ttctccgtaa acaacatgtt tcccatgta 720
atatcctttt ctatttttctg ttccgttacc aactttacac atactttata tagctattca 780
cttctataca ctaaaaaact aagacaattt taattttgct gcctgccata tttcaatttg 840
ttataaatc ctataattta tctatttagt agctaaaaaa agatgaatgt gaatcgaatc 900
ctaagagaat tgctgcagaa ttcacggatt agaagccgcc gagcgggtga cagccctccg 960
aaggaaagact ctctccgtg cgtccctcgtc ttcaccggtc gcgttcctga aacgcagatg 1020
tgctcgcgc cgcactgctc cgaacaataa agattctaca atactagctt ttatggttat 1080
gaagaggaaa aattggcagt aacctggccc cacaacactt caaatgaacg aatcaaatca 1140
acaaccatag gatgataatg cgattagttt tttagcctta tttctggggt aattaatcag 1200
cgaagcagatg atttttgatc tattaacaga tatataaatg caaaaactgc ataaccactt 1260
taactaatat tttcaacatt ttcggtttgt attacttctt attcaaatgt aataaaagta 1320
tcaacaaaaa attgttaata tacctctata cttaacgctc aaggagaaaa aaccgccgat 1380
cggactacta gcagctgtaa tacgactcac tatagggaat attaaagtaa ttctacttca 1440
tacattttca attaatgac agttacttcg ctgtttttca atattttctg ttattgcttc 1500
agtttttagc caggaaactg caactatatg cgagcaaact ccctcaccia ctttagaatc 1560
gacgccgtac tctttgtcaa cgactactat tttggccaac gggaaggcaa tgcaaggagt 1620
ttttgaatat tacaaatcag taacgtttgt cagtaattgc gggtctcacc cctcaacaac 1680
tagcaaaaggc agcccccata acacacagta tgtttttaag cttctgcagg ctagtgggtg 1740
tggtggttct ggtggtggtg gttctggtgg tgggtggtct gctagcatga ctggtggcca 1800
gcaaggccta attctgatgc ggccgcacat catcaccatc accattgatt aattaagttt 1860

```

-continued

aaacccgctg atctgataac aacagtgtag atgtaacaaa atcgactttg ttcccactgt	1920
acttttagct cgtacaaaat acaatatact tttcatttct cegtaacaaa catgttttcc	1980
catgtaatat ccttttctat ttttcgttcc gttaccaact ttacacatac tttatatagc	2040
tattcacttc tatacactaa aaaactaaga caattttaat tttgctgect gccatatttc	2100
aatttggtat aaattcctat aatttatcct attagtagct aaaaaagat gaatgtgaat	2160
cgaatcctaa gagaattggg caagtgcaca aacaatactt aaataaatac tactcagtaa	2220
taacctatct cttagcattt ttgacgaaat ttgctatttt gttagagtct tttacaccat	2280
ttgtctccac acctccgett acatcaacac caataacgcc atttaatcta agcgcacac	2340
caacattttc tggcgctcagt ccaccagcta acataaaatg taagctctcg gggctctctt	2400
gccttccaac ccagtcagaa atcgagttcc aatccaaaag ttcacctgtc ccacctgett	2460
ctgaatcaaa caagggaata aacgaatgag gtttctgtga agctgcactg agtagtatgt	2520
tgcagtcttt tggaaatcac agtcttttaa taactggcaa accgaggaac tcttggtatt	2580
cttgccacga ctcactctcg tgcagttgga cgatatcaat gccgtaatca ttgaccagag	2640
ccaaaacatc ctccttaggt tgattacgaa acacgccaac caagtatttc ggagtgcctg	2700
aactattttt atatgctttt acaagacttg aaattttcct tgcaataacc gggtcatttg	2760
ttctctttct attgggcaca catataatac ccagcaagtc agcatcgga tctagagcac	2820
attctgcggc ctctgtgtct tgcaagccgc aaactttcac caatggacca gaactacctg	2880
tgaaattaat aacagacata ctccaagctg cctttgtgtg cttaatcacg tatactcacg	2940
tgctcaatag tcaccaatgc cctccctctt ggccctctcc ttttctttt tcgaccgaat	3000
ttcttgaaga cgaaagggcc tcgtgatacg cctattttta taggttaatg tcatgataat	3060
aatggtttct taggacggat cgcttgccctg taacttacac gcgcctcgta tcttttaattg	3120
atggaataat ttgggaattt actctgtgtt tatttatctt tatgttttgt atttgattt	3180
tagaaagtaa ataaagaagg tagaagagtt acggaatgaa gaaaaaaaa taacaaaagg	3240
tttaaaaaat ttcaacaaaa agcgtacttt acatatatat ttattagaca agaaaagcag	3300
attaaataga tatacattcg attaacgata agtaaaatgt aaatcacag gattttcgtg	3360
tgtggtcttc tacacagaca agatgaaaca attcggcatt aatacctgag agcaggaaga	3420
gcaagataaa aggtagtatt tgttgccgat cccctagag tcttttacat cttcggaata	3480
caaaaactat tttttcttta atttctttt ttactttcta tttttaattt atatatttat	3540
attaaaaaat ttaaattata attattttta tagcacgtga tgaaaaggac ccagggtggca	3600
cttttcgggg aaatgtgcgc ggaaccccta tttgtttatt tttctaaata cattcaaata	3660
tgtatccgct catgagacaa taacctgat aaatgcttca ataatttga aaaaggaaga	3720
gtatgagtat tcaacatttc cgtgtcgccc ttattccctt ttttgcggca ttttgccttc	3780
ctgtttttgc tcaaccagaa acgctgggtga aagtaaaaga tgctgaagat cagttgggtg	3840
cacgagtggg ttacatcgaa ctggatctca acageggtaa gatccttgag agttttcgcc	3900
ccgaagaacg ttttccaatg atgagcactt ttaaagttct gctatgtggc gcggtattat	3960
cccgtgttga cgccgggcaa gagcaactcg gtcgccgat acactattct cagaatgact	4020
tgggttagta ctcaccagtc acagaaaagc atcttacgga tggcatgaca gtaagagaat	4080
tatgcagtgc tgccataacc atgagtata acaactgcgc caacttactt ctgacaacga	4140
tcggaggacc gaaggagcta accgcttttt tgcacaacat gggggatcat gtaactcgcc	4200

-continued

ttgatcgttg ggaaccggag ctgaatgaag ccataccaaa cgacgagcgt gacaccacga	4260
tgcctgtagc aatggcaaca acgttgcgca aactattaac tggcgaacta cttactctag	4320
cttcccggca acaattaata gactggatgg aggcggataa agttgcagga ccacttctgc	4380
gctcggccct tccggctggc tggtttattg ctgataaatc tggagccggg gagcgtgggt	4440
ctcgcggtat cattgcagca ctggggccag atggtaagcc ctcccgtatc gtagttatct	4500
acacgacggg cagtcaggca actatggatg aacgaaatag acagatcgct gagatagggtg	4560
cctcactgat taagcattgg taactgtcag accaagttta ctcatatata ctttagattg	4620
atttaaaact tcatttttaa tttaaaagga tctaggtgaa gatccttttt gataatctca	4680
tgacaaaat cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga	4740
tcaaaggatc ttcttgagat cctttttttc tgcgcgtaat ctgctgcttg caaacaaaaa	4800
aaccacgct accagcgggt gtttgtttgc cggatcaaga gctaccaact ctttttccga	4860
aggtaactgg cttcagcaga gcgcagatac caaatactgt ccttctagt tagccgtagt	4920
taggccacca cttcaagaac tctgtagcac cgcctacata cctcgtcttg ctaatcctgt	4980
taccagtggc tgctgccagt ggcgataagt cgtgtcttac cgggttgga tcaagacgat	5040
agttaccgga taaggcgcag cggtcggggt gaacgggggg ttcgtgcaca cagcccagct	5100
tggagcgaac gacctacacc gaactgagat acctacagcg tgagcattga gaaagcgcca	5160
cgcttcccgaggagaaaag gcggacaggt atccggtaag cggcagggtc ggaacaggag	5220
agcgcacgag ggagcttcca ggggggaacg cctggtatct ttatagtct gtcgggtttc	5280
gccacctctg acttgagcgt cgatttttgt gatgctcgtc agggggggcc agcctatgga	5340
aaaacgccag caacgcggcc tttttacggt tcttggcctt ttgctggcct tttgctcaca	5400
tgttctttcc tgcgttatcc cctgattctg tggataaccg tattaccgcc tttgagttag	5460
ctgataccgc tcgccgcagc cgaacgaccg agcgcagcga gtcagtgagc gaggaagcgg	5520
aagagcgcgc aatacgcaaa ccgcctctcc ccgcgcgttg gccgattcat taatgcagct	5580
ggcagcagag gtttcccgac tggaaagcgg gcagtgagcg caacgcaatt aatgtgagtt	5640
acctcactca ttaggcaccc caggctttac actttatgct tccggctcct atgttgtgtg	5700
gaattgtgag cggataacaa tttcacacag gaaacagcta tgaccatgat tacgccaagc	5760
tcggaattaa ccctcactaa agggaacaaa agctggctag t	5801

<210> SEQ ID NO 77

<211> LENGTH: 692

<212> TYPE: DNA

<213> ORGANISM: artificial

<220> FEATURE:

<223> OTHER INFORMATION: 4D5H

<400> SEQUENCE: 77

ggccagcaag gccaaagagt tcaactagt gagtctggcg gtggcctggg gcagccaggg	60
ggctcactcc gtttgtcctg tgcagcttct ggcttcaaca ttaaagacac ctatatacac	120
tgggtgcgtc agggcccggt taaggcctg gaatgggttg caaggattta tcctacgaat	180
ggttatacta gatatgcoga tagcgtcaag ggccgtttca ctataagcg agacacatcc	240
aaaaacacag cctacctgca gatgaacagc ctgcgtgctg aggacactgc cgtctattat	300
tgttctagat ggggagggga cggcttctat gctatggact actgggggtca aggaacctg	360
gtcaccgtct cctcggctag caccaagggc ccagcgtgt tccctctggc cccagctcc	420
aagagcacct ccggcggcac cgcgcgcctg ggctgcctgg tgaaggatta cttcccagag	480

-continued

```

cccgtgaccg  tgagctggaa  cagcggcgcc  ctgaccagcg  gcgtgcacac  ctttcccgcc  540
gtgctgcagt  ccagcggcct  gtactccctg  agcagcgtgg  tgaccgtgcc  cagcagcagc  600
ctgggcaccc  agacctacat  ctgcaatgtg  aaccacaagc  ccagcaatac  caagggtgat  660
aagaagggtg  agcccaagag  ctgcgcggcc  gc  692

```

```

<210> SEQ ID NO 78
<211> LENGTH: 661
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: 4D5L

```

```

<400> SEQUENCE: 78

```

```

ccatggcgga  tateccagatg  acccagtcct  cgagctccct  gtccgcctct  gtgggcgata  60
gggtcaccat  cacctgcctg  gccagtcagg  atgtgaatac  tgctgtagcc  tggatatcaac  120
agaaaccagg  aaaagctccg  aaactactga  ttactctcgc  atccttcctc  tactctggag  180
tcccttctcg  cttctctgga  tccagatctg  ggacggattt  cactctgacc  atcagcagtc  240
tgcagccgga  agacttcgca  acttattact  gtcagcaaca  ttatactact  cctcccacgt  300
tcggacaggg  taccaagggt  gagatcaaac  gtacggtgcc  ggcgccatct  gtcttcatct  360
tcccgccatc  tgatgagcag  cttaagtctg  gaactgcctc  tgttggtgtg  ctgctgaata  420
acttctatcc  cagagaggcc  aaagtacagt  ggaagggtga  taacgccctc  caatcgggta  480
actcccagga  gagtgtcaca  gacgaggaca  gcaaggacag  cacctacagc  ctcagcagca  540
ccttgacgct  gagcaaagca  gactacgaga  aacacaaagt  ctacgcctgc  gaagtcaccc  600
atcagggcct  gagctcgccc  gtcacaaaga  gcttcaacag  gggagagtgt  tgaggcgcgc  660
c  661

```

```

<210> SEQ ID NO 79
<211> LENGTH: 6468
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: vector pYD4D5hc

```

```

<400> SEQUENCE: 79

```

```

acggattaga  agccgcgcag  cgggtgacag  cctccgaag  gaagactctc  ctccgtgcgt  60
cctcgtcttc  accggtcgcg  ttccgtgaaac  gcagatgtgc  ctccgcgcgc  actgctccga  120
acaataaaga  ttctacaata  ctgactttta  tggttatgaa  gaggaaaaat  tggcagtaac  180
ctggccccac  aaaccttcaa  atgaacgaat  caaatcaaca  accataggat  gataatgcga  240
ttagtttttt  agccttattt  ctggggtaat  taatcagcga  agcagtgatt  tttgatctat  300
taacagatat  ataaatgcaa  aaactgcata  accactttaa  ctaatacttt  caacattttc  360
ggttttgtatt  acttcttatt  caaatgtaat  aaaagtatca  acaaaaaatt  gttaatatatac  420
ctctatactt  taacgtcaag  gagaaaaaac  cccggatcgg  actactagca  gctgtaatac  480
gactcactat  agggaatatt  aagctaattc  tacttcatac  attttcaatt  aagatgcagt  540
tacttcgctg  tttttcaata  ttttctgtta  ttgcttcagt  gctagccgct  gggggccatgg  600
ttactgattg  gcgcgcgcga  tccgatgtaa  caaaatcgac  tttgttccca  ctgtactttt  660
agctcgtaca  aaatacaata  tacttttcat  ttctcgttaa  acaacatggt  ttcccatgta  720
atatcctttt  ctatttttct  ttccgttacc  aactttacac  atactttata  tagctattca  780

```

-continued

cttctataca	ctaaaaaact	aagacaattt	taatttttgc	gcctgccata	tttcaatttg	840
ttataaatc	ctataattta	tctatttagt	agctaaaaa	agatgaatgt	gaatcgaatc	900
ctaagagaat	tgctgcagaa	ttcacggatt	agaagccgcc	gagcgggtga	cagccctccg	960
aaggaagact	ctcctccgtg	cgctcctgtc	ttcacccgtc	gcgttctga	aacgcagatg	1020
tgccctcgcg	cgcactgctc	cgaacaataa	agattctaca	atactagctt	ttatggttat	1080
gaagaggaaa	aattggcagt	aacctggccc	cacaaacctt	caaatgaacg	aatcaaatca	1140
acaaccatag	gatgataatg	cgattagttt	tttagcctta	tttctggggg	aattaatcag	1200
cgaagcgatg	atttttgatc	tattaacaga	tataataatg	caaaaactgc	ataaccactt	1260
taactaatac	tttcaacatt	ttcggtttgt	attacttctt	attcaaatgt	aataaaagta	1320
tcaacaaaaa	attgttaata	tacctctata	ctttaacgtc	aaggagaaaa	aaccccgcat	1380
cggactacta	gcagctgtaa	tacgactcac	tatagggaat	attaagctaa	ttctacttca	1440
tacattttca	attaagatgc	agttaactcg	ctgtttttca	atattttctg	ttattgcttc	1500
agtttttagc	caggaactga	caactatatg	cgagcaaatc	ccctcaccaa	ctttagaatc	1560
gacgccgtac	tctttgtcaa	cgactactat	tttgcccaac	gggaaggcaa	tgcaaggagt	1620
ttttgaatat	tacaaatcag	taacgtttgt	cagtaattgc	ggttctcacc	cctcaacaac	1680
tagcaaaggc	agccccataa	acacacagta	tgtttttaag	cttctgcagg	ctagtgggtg	1740
tggtgggtct	gggtgggtg	gttctgggtg	tggtgggtct	gctagcatga	ctggtggcca	1800
gcaaggccaa	ggttctgagg	ttcaactagt	ggagtctggc	gggtggcctg	tgagccagg	1860
gggctcactc	cgtttgcctc	gtgcagcttc	tggttcaac	attaaagaca	cctatatata	1920
ctgggtgcgt	caggcccccg	gtaagggcct	ggaatgggtt	gcaaggattt	atcctacgaa	1980
tggttatact	agatatgcg	atagcgtcaa	gggcggttcc	actataagcg	cagacacatc	2040
caaaaacaca	gcctacctgc	agatgaacag	cctgcgtgct	gaggacactg	ccgtctatta	2100
ttgttctaga	tggggagggg	acggcttcta	tgctatggac	tactgggggc	aaggaaccct	2160
ggtcaccgtc	tctcgggcta	gcaccaaggg	ccccagcgtg	ttccctctgg	ccccagctc	2220
caagagcacc	tccggcggca	ccgccgcctc	gggctgcctg	gtgaaggatt	acttcccaga	2280
gcccgtgacc	gtgagctgga	acagcggcgc	cctgaccagc	ggcgtgcaca	cctttcccgc	2340
cgtgctgcag	tccagcggcc	tgtactccct	gagcagcgtg	gtgaccgtgc	ccagcagcag	2400
cctgggcacc	cagacctaca	tctgcaatgt	gaaccacaag	cccagcaata	ccaaggtgga	2460
taagaagggtg	gagcccaaga	gctgcgcggc	cgcacatcat	caccatcacc	attgattaat	2520
taagtttaaa	cccgtgatc	tgataacaac	agtgtagatg	taacaaaatc	gactttgttc	2580
ccactgtact	tttagctcgt	acaaaatata	atatactttt	catttctccg	taacaacat	2640
gttttcccat	gtaatatcct	tttctatttt	tcgttccgtt	accaacttta	cacatacttt	2700
atatagctat	tcacttctat	acactaaaaa	actaagacaa	ttttaatttt	gctgcctgcc	2760
atatttcaat	ttgttataaa	ttcctataat	ttatcctatt	agtagctaaa	aaaagatgaa	2820
tgtgaatcga	atcctaagag	aattggggcaa	gtgcacaaac	aatacttaaa	taaatactac	2880
tcagtaataa	cctatttctt	agcatttttg	acgaaaattg	ctattttgtt	agagtctttt	2940
acaccatttg	tctccacacc	tccgcttaca	tcaacaccaa	taacgccatt	taatctaagc	3000
gcacacacaa	cattttcttg	cgctagtcga	ccagetaaca	taaaatgtaa	gctctcgggg	3060
ctctcttgcc	ttccaaccca	gtcagaaatc	gagttccaat	ccaaaagtgc	acctgtccca	3120
cctgcttctg	aatcaaacaa	gggaataaac	gaatgaggtt	tctgtgaagc	tgactgagtg	3180

-continued

agtatgttgc agtcttttgg aaatacagagt cttttaataa ctggcaaacc gaggaactct	3240
tggtattctt gccacgactc atctccgtgc agttggacga tatcaatgcc gtaatcattg	3300
accagagcca aaacatcctc cttagggtga ttacgaaaca cgccaaccaa gtatttcgga	3360
gtgcctgaac tatttttata tgcttttaca agacttgaaa ttttccttgc aataaccggg	3420
tcaattgttc tctttctatt gggcacacat ataataccca gcaagtcagc atcggaatct	3480
agagcacatt ctgcggcctc tgtgctctgc aagccgcaaa ctttcaccaa tggaccagaa	3540
ctacctgtga aattaataac agacatactc caagctgcct ttgtgtgctt aatcacgtat	3600
actcacgtgc tcaatagtca ccaatgcctt cctctttggc cctctccttt tcttttttcg	3660
accgaatttc ttgaagacga aagggcctcg tgatacgctt atttttatag gttaatgtca	3720
tgataataat ggtttcttag gacggatgc ttgectgtaa cttacacgcg cctcgatatc	3780
tttaatgatg gaataatttg ggaatttact ctgtgtttat ttatttttat gttttgtatt	3840
tggatttttag aaagtaataa aagaaggtag aagagttacg gaatgaagaa aaaaaataa	3900
acaaagggtt aaaaaatttc aacaaaaagc gtactttaca tatatattta ttagacaaga	3960
aaagcagatt aaatagatat acattcgatt aacgataagt aaaatgtaa atcacaggat	4020
tttcgtgtgt ggtcttctac acagacaaga tgaaacaatt cggcattaat acctgagagc	4080
aggaagagca agataaaagg tagtatttgt tggcgatccc cctagagtct ttacatctt	4140
cggaaaaaca aaactatttt ttctttaatt tcttttttta ctttctattt ttaatttata	4200
tatttatatt aaaaaattta aattataatt atttttatag cacgtgatga aaaggaccca	4260
ggtggcactt ttcggggaaa tgtgcgcgga acccctattt gtttattttt ctaaatacat	4320
tcaaatatgt atccgctcat gagacaataa ccttgataaa tgcttcaata atattgaaaa	4380
aggaagagta tgagtattca acatttcctg tgcgccctta ttcccttttt tgcggcattt	4440
tgcttctctg tttttgtcga ccagaaaacg ctggtgaaag taaaagatgc tgaagatcag	4500
ttgggtgcac gagtgggtta catcgaaact gatctcaaca gcggaagat ccttgagagt	4560
tttcgccccg aagaacgttt tccaatgatg agcactttta aagttctgct atgtggcgcg	4620
gtattatccc gtgttgacgc cgggcaagag caactcggtc gccgcataca ctattctcag	4680
aatgacttgg ttgagtactc accagtcaca gaaaagcatc ttacggatgg catgacagta	4740
agagaattat gcagtgtctc cataaccatg agtgataaca ctgcggccaa cttacttctg	4800
acaacgatcg gaggaccgaa ggagctaacg gcttttttgc acaacatggg ggatcatgta	4860
actcgccttg atcgttggga accggagctg aatgaagcca taccaaacga cgagcgtgac	4920
accacgatgc ctgtagcaat ggcaacaacg ttgcgcaaac tattaactgg cgaactactt	4980
actctagctt cccggcaaca attaatagac tggatggagg cggataaagt tgcaggacca	5040
cttctgcgct cggcccttcc ggctggctgg tttattgctg ataaatctgg agcgggtgag	5100
cgtgggtctc gcggtatcat tgcagactg gggccagatg gtaagccctc ccgtatcgta	5160
gttatctaca cgacgggcag tcaggcaact atggatgaac gaaatagaca gatcgctgag	5220
ataggtgcct cactgattaa gcatttgtaa ctgtcagacc aagtttactc atatatactt	5280
tagattgatt taaaacttca tttttaattt aaaaggatct aggtgaagat cttttttgat	5340
aatctcatga ccaaaatccc ttaacgtgag ttttcgttcc actgagcgtc agaccccgta	5400
gaaaagatca aaggatcttc ttgagatcct ttttttctgc gcgtaatctg ctgcttgcaa	5460
acaaaaaac caccgctacc agcggtggtt tgtttgcggg atcaagagct accaactctt	5520

-continued

```

tttccgaagg taactggctt cagcagagcg cagataccaa atactgtcct tctagtgtag 5580
ccgtagtttag gccaccactt caagaactct gtagcaccgc ctacatacct cgctctgcta 5640
atcctgtttac cagtggctgc tgccagtggc gataagtcgt gtcttacggg gttggactca 5700
agacgatagt taccggataa ggcgcagcgg tcgggctgaa cgggggggttc gtgcacacag 5760
cccagcttgg agcgaacgac ctacaccgaa ctgagatacc tacagcgtga gcattgagaa 5820
agcgccacgc ttcccgaagg gagaaaggcg gacaggtatc cggtaaagcgg cagggtcgga 5880
acaggagagc gcacgagggg gcttccaggg gggaacgcct ggtatcttta tagtcctgtc 5940
gggttttcgcc acctctgact tgagcgtcga tttttgtgat gctcgtcagg gggggccgagc 6000
ctatggaaaa acgccagcaa cgcggccttt ttacgggttc tggccttttg ctggcctttt 6060
gctcacatgt tctttcctgc gttatccctt gattctgtgg ataaccgtat taccgccttt 6120
gagtgagctg ataccgctcg ccgcagccga acgaccgagc gcagcgagtc agtgagcgag 6180
gaagcggaag agcgcccaat acgcaaaccg cctctccccg cgcgttggcc gattcattaa 6240
tgcagctggc acgacaggtt tcccgaactg aaagcgggca gtgagcgcaa cgcaattaat 6300
gtgagttacc tcactcatta ggcaccccag gctttacact ttatgcttcc ggctcctatg 6360
ttgtgtggaa ttgtgagcgg ataacaattt cacacaggaa acagctatga ccatgattac 6420
gccaaagctg gaattaacct tcactaaagg gaacaaaagc tggctagt 6468

```

```

<210> SEQ ID NO 80
<211> LENGTH: 223
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: 4D5hp

```

```

<400> SEQUENCE: 80

```

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1           5           10          15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Asn Ile Lys Asp Thr
20          25          30
Tyr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35          40          45
Ala Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr Ala Asp Ser Val
50          55          60
Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr
65          70          75          80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85          90          95
Ser Arg Trp Gly Gly Asp Gly Phe Tyr Ala Met Asp Tyr Trp Gly Gln
100         105         110
Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
115         120         125
Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
130         135         140
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
145         150         155         160
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
165         170         175
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
180         185         190
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys

```

-continued

195	200	205	
Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys			
210	215	220	
<210> SEQ ID NO 81			
<211> LENGTH: 7100			
<212> TYPE: DNA			
<213> ORGANISM: artificial			
<220> FEATURE:			
<223> OTHER INFORMATION: vector pYD4D5h1			
<400> SEQUENCE: 81			
acggattaga agccgccgag cgggtgacag ccctccgaag gaagactctc ctccgtgcgt			60
cctcgtcttc accggtcgag ttctgaaac gcagatgtgc ctgcgccgc actgctccga			120
acaataaaga ttctacaata ctgactttta tggttatgaa gaggaataat tggcagtaac			180
ctggccccac aaaccttcaa atgaacgaat caaattaaca accataggat gataatgcga			240
ttagtttttt agccttattt ctggggtaat taatcagcga agcgatgatt tttgatctat			300
taacagatat ataaatgcaa aaactgcata accactttaa ctaatacttt caacattttc			360
ggtttgattt acttcttatt caaatgtaat aaaagtatca acaaaaaatt gttaatatat			420
ctctatactt taacgtcaag gagaaaaaac cccggatcgg actactagca gctgtaatac			480
gactcactat agggaaatatt aagctaattc tacttcatac attttcaatt aagatgcagt			540
tacttcgctg tttttcaata ttttctgtta ttgcttcagt gctagccgct ggggccatgg			600
cggatatcca gatgaccag tccccgagct cctgtccgc ctctgtgggc gataggggtca			660
ccatcacctg cgtgccagt caggatgtga atactgctgt agcctggat caacagaaac			720
caggaaaagc tccgaaacta ctgatttact cggcatcctt cctctactct ggagtccctt			780
ctcgtctctc tggatccaga tctgggacgg atttcaactc gaccatcagc agtctgcagc			840
cggaagactt cgcaacttat tactgtcagc aacattatac tactcctccc acgttcggac			900
aggggtacca ggtggagatc aaacgtacgg tggcggcgcc atctgtcttc atcttccgc			960
catctgatga gcagcttaag tctggaactg cctctgttgt gtgcctgctg aataacttct			1020
atcccagaga ggccaaagta cagtggagg tggataacgc cctccaatcg ggtaactccc			1080
aggagagtgt cacagagcag gacagcaagg acagcaccta cagcctcagc agcaccctga			1140
cgtgagcaa agcagactac gagaaacaca aagtctacgc ctgcgaagtc accatcagg			1200
gctgagctc gcccgtcaca aagagcttca acaggggaga gtgttgaggc gcgccggatc			1260
cgatgtaaca aaatcgactt tgttcccact gtacttttag ctctacaaa atacaatata			1320
cttttcattt ctccgtaaac aacatgtttt cccatgtaat atccttttct atttttcgtt			1380
ccgttaccaa ctttacacat actttatata gctattcact tctatacact aaaaaactaa			1440
gacaatttta attttgctgc ctgccatatt tcaatttgtt ataaattcct ataatttatc			1500
ctattagtag ctaaaaaaag atgaatgtga atcgaatcct aagagaattg ctgcagaatt			1560
cacggattag aagccgcga gcgggtgaca gccctccgaa ggaagactct cctccgtgcg			1620
tccctgtctt caccggtcgc gttcctgaaa cgcagatgtg cctcgcgccg cactgctccg			1680
aacaataaag attctacaat actagctttt atggttatga agaggaaaaa tggcagtaa			1740
cctggcccca caaaccttca aatgaacgaa tcaaatatc aaccatagga tgataatgcg			1800
attagttttt tagccttatt tctggggtaa ttaatcagcg aagcgatgat ttttgatcta			1860
ttaacagata tataaatgca aaaactgcat aaccacttta actaatactt tcaacatttt			1920

-continued

cggtttgtat	tacttcttat	tcaaatgtaa	taaaagtatc	aacaaaaaat	tgtaatatata	1980
cctctatact	ttaacgtcaa	ggagaaaaaa	ccccggatcg	gactactagc	agctgtaata	2040
cgactcacta	tagggaatat	taagctaatt	ctacttcata	cattttcaat	taagatgcag	2100
ttacttcgct	gtttttcaat	attttctgtt	attgcttcag	ttttagcaca	ggaactgaca	2160
actatatgcg	agcaaatccc	ctcaccaact	ttagaatcga	cgccgtactc	tttgtcaacg	2220
actactatth	tggccaacgg	gaaggcaatg	caaggagttt	tgaaatatta	caaatcagta	2280
acgtttgtca	gtaattgogg	ttctcaccoc	tcaacaacta	gcaaaggcag	ccccataaac	2340
acacagtatg	tttttaagct	tctgcaggct	agtgggtggg	gtgggtctcg	tggtgggtgg	2400
tctgtgggtg	gtggttctgc	tagcatgact	ggtggccagc	aaggccaaga	ggttcaacta	2460
gtggagtctg	gcggtggcct	ggtgcagcca	gggggctcac	tccgtttgtc	ctgtgcagct	2520
tctggcttca	acattaaaga	cacctatata	cactgggtgc	gtcaggcccc	gggtaagggc	2580
ctggaatggg	ttgcaaggat	ttatctctac	aatgggtata	ctagatatgc	cgatagcgct	2640
aagggccggt	tcactataag	cgacagacac	tccaaaaaca	cagcctacct	gcagatgaac	2700
agcctgcgtg	ctgaggacac	tgccgtctat	tattgttcta	gatggggagg	ggacggcttc	2760
tatgctatgg	actactgggg	tcaaggaacc	ctggtcacog	tctcctcggc	tagcaccaag	2820
ggccccagcg	tggtccctct	ggccccagc	tccaagagca	cctccggcgg	caccgcccgc	2880
ctgggctgcc	tggtgaagga	ttacttccca	gagcccgtag	ccgtgagctg	gaacagcggc	2940
gccctgacca	gcggcgtgca	cacctttccc	gccgtgctgc	agtcacggcg	cctgtactcc	3000
ctgagcagcg	tggtgaccgt	gcccagcagc	agcctgggca	cccagacctc	catctgcaat	3060
gtgaaccaca	agcccagcaa	taccaagggt	gataagaagg	tggaagccaa	gagctgcgcg	3120
gccgcacatc	atcaccatca	ccattgatta	attaagttta	aaccgcgtga	tctgataaca	3180
acagtgtaga	tgtaacaaaa	tcgactttgt	tcccactgta	cttttagctc	gtacaaaata	3240
caataatact	ttcattttct	cgtaaacac	atgttttccc	atgtaatatc	cttttctatt	3300
tttcgttcog	ttaccaactt	tacacatact	ttatatagct	attcacttct	atacactaaa	3360
aaactaagac	aattttaatt	ttgctgctg	ccatatttca	atttggtata	aattcctata	3420
atttatccca	ttagtagcta	aaaaaagatg	aatgtgaatc	gaatcctaag	agaattgggc	3480
aagtcacaaa	acaatactta	aataaatact	actcagtaat	aacctatttc	ttagcattht	3540
tgacgaaatt	tgctattttg	ttagagtctt	ttacaccatt	tgtctccaca	cctccgctta	3600
catcaacacc	aataacgcca	tttaattctaa	gcgcacacc	aacattttct	ggcgtcagtc	3660
caccagctaa	cataaaatgt	aagctctcgg	ggctctcttg	ccttccaacc	cagtcagaaa	3720
tcgagttcca	atccaaaagt	tcacctgtcc	cacctgcttc	tgaatcaaac	aagggaataa	3780
acgaatgagg	tttctgtgaa	gctgcactga	gtagtatggt	gcagtctttt	ggaaatacga	3840
gtcttttaat	aactggcaaa	ccgaggaact	cttggtatct	ttgccacgac	tcatctccgt	3900
gcagttggac	gatatcaatg	ccgtaatcat	tgaccagagc	caaaacatcc	tccttagggt	3960
gattacgaaa	cacgccaacc	aagtatttct	gagtgcctga	actattttta	tatgctttta	4020
caagacttga	aattttcctt	gcaataaccg	ggtcaattgt	tctctttcta	ttgggcacac	4080
atataatacc	cagcaagtca	gcacgggaat	ctagagcaca	ttctgcgggc	tctgtgctct	4140
gcaagccgca	aactttcacc	aatggaccag	aactacctgt	gaaattaata	acagacatac	4200
tccaagctgc	ctttgtgtgc	ttaatcacgt	atactcacgt	gctcaatagt	caccaatgcc	4260
ctccctcttg	gccctctcct	tttctttttt	cgaccgaatt	tcttgaagac	gaaagggcct	4320

-continued

cgtgatacgc	ctatTTTTat	aggTTaatgt	catgataata	atggtttctt	aggacggatc	4380
gcttgccctgt	aacttacacg	cgcctcgtat	cttttaataga	tggaataaatt	tggaattta	4440
ctctgtgttt	atTTatTTTT	atgttttgta	tttggtttt	agaaagtaa	taaagaaggt	4500
agaagagtta	cggaatgaag	aaaaaaaaat	aaacaaaggt	ttaaaaaatt	tcaacaaaa	4560
gcgtacttta	catatatatt	tattagacaa	gaaaagcaga	ttaaatagat	atacattcga	4620
ttaacgataa	gtaaaatgta	aaatcacagg	attttcgtgt	gtggtcttct	acacagacaa	4680
gatgaaacaa	ttcggcatta	atacctgaga	gcaggaagag	caagataaaa	ggtagtattt	4740
gttggcgatc	cccctagagt	cttttacatc	ttcggaaaac	aaaaactatt	tttctttaa	4800
tttctTTTT	tactttctat	ttttaattta	tatatTTata	ttaaaaaatt	taaattataa	4860
ttatTTTTat	agcacgtgat	gaaaaggacc	caggtggcac	tttctgggga	aatgtgcgcg	4920
gaacccctat	ttgtttatTT	ttctaaatac	attcaaatat	gtatccgctc	atgagacaat	4980
aaccttgata	aatgcttcaa	taatatTgaa	aaaggaagag	tatgagtatt	caacatttcc	5040
gtgtgcgcc	tattcccttt	tttgcggcat	tttgccctcc	tgTTTTgtct	caccagaaa	5100
cgtcggtgaa	agtaaaagat	gctgaagatc	agttgggtgc	acgagtgggt	tacatcgaa	5160
tggatctcaa	cagcggtaag	atccttgaga	gttttcgccc	cgaagaacgt	tttccaatga	5220
tgagcacttt	taaagtctct	ctatgtggcg	cggatttacc	ccgtgttgac	gccgggcaag	5280
agcaactcgg	tcgccgcata	cactattctc	agaatgactt	ggttgagtac	tcaccagtca	5340
cagaaaagca	tcttacggat	ggcatgacag	taagagaatt	atgcagtgct	gccataacca	5400
tgagtgataa	cactgcggcc	aacttacttc	tgacaacgat	cggaggaccg	aaggagctaa	5460
ccgctTTTT	gcacaacatg	ggggatcatg	taactcgcct	tgatcgttgg	gaaccggagc	5520
tgaatgaagc	cataccaaac	gacgagcgtg	acaccacgat	gcctgtagca	atggcaacaa	5580
cgttcgcgaa	actattaact	ggcgaaactac	ttactctagc	ttcccgcaa	caattaatag	5640
actggatgga	ggcggataaa	gttgacggac	cacttctgcg	ctcggccctt	ccggctggct	5700
ggtttatTgc	tgataaatct	ggagccgggtg	agcgtgggtc	tcgcggtatc	attgcagcac	5760
tggggccaga	tggttaagccc	tcccgatcgc	tagttatcta	cacgacgggc	agtcaggcaa	5820
ctatggatga	acgaaataga	cagatcgcgtg	agatagggtgc	ctcactgatt	aagcattggt	5880
aactgtcaga	ccaagtTtac	tcatatatac	tttagattga	tttaaaactt	catttttaat	5940
ttaaaaggat	ctaggtgaag	atcctTTTTg	ataatctcat	gacaaaaatc	ccttaacgtg	6000
agttttcgtt	ccactgagcg	tcagaccccg	tagaaaagat	caaaggatct	tcttgagatc	6060
ctttttttct	gcgcgtaatc	tgtctgtTgc	aaacaaaaaa	accaccgcta	ccagcggTgg	6120
tttgtttgcc	ggatcaagag	ctaccaactc	tttttcgaa	ggtaactggc	ttcagcagag	6180
cgcagatacc	aaatactgtc	cttctagtgt	agccgtagtt	aggccaccac	ttcaagaact	6240
ctgtagcacc	gcctacatac	ctcgcctctgc	taatcctgtt	accagtggct	gctgccagtgc	6300
gcgataagtc	gtgtcttacc	gggttggact	caagacgata	gttaccggat	aaggcgcagc	6360
ggtcgggctg	aacgggggggt	tcgtgcacac	agcccagctt	ggagcgaacg	acctacaccg	6420
aactgagata	cctacagcgt	gagcattgag	aaagcggcac	gcttcccgaa	gggagaaagg	6480
cggacaggta	tccggtaagc	ggcagggctg	gaacaggaga	gcgcacgagg	gagcttccag	6540
gggggaacgc	ctggtatctt	tatagtctctg	tcgggtttcg	ccacctctga	cttgagcgtc	6600
gatttttTgtg	atgctcgtca	ggggggccga	gcctatggaa	aaacgccagc	aacgcggcct	6660

-continued

```

ttttacgggt cctggccttt tgcggcctt ttgctcacat gttctttcct gcgttatccc 6720
ctgattctgt ggataaccgt attaccgctt ttgagtgagc tgataccgct cgccgcagcc 6780
gaacgaccga gcgcagcgag tcagtgcgag aggaagcgga agagcgccca atacgcaaac 6840
cgctctctcc cgcgcggttg ccgattcatt aatgcagctg gcacgacagg ttccccgact 6900
ggaaagcggg cagtgcgagc aacgcaatta atgtgagtta cctcactcat taggcacccc 6960
aggttttaca ctttatgctt ccggctccta tgttgtgtgg aattgtgagc ggataacaat 7020
ttcacacagg aaacagctat gaccatgatt acgccaagct cggaattaac cctcactaaa 7080
gggaacaaaa gctggctagt 7100

```

```

<210> SEQ ID NO 82
<211> LENGTH: 214
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: 4D5lp

```

```

<400> SEQUENCE: 82

```

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1      5      10      15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Asn Thr Ala
20     25     30
Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35     40     45
Tyr Ser Ala Ser Phe Leu Tyr Ser Gly Val Pro Ser Arg Phe Ser Gly
50     55     60
Ser Arg Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65     70     75     80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln His Tyr Thr Thr Pro Pro
85     90     95
Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
100    105    110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
115    120    125
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
130    135    140
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145    150    155    160
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
165    170    175
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
180    185    190
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
195    200    205
Phe Asn Arg Gly Glu Cys
210

```

```

<210> SEQ ID NO 83
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: C-terminal residues of junction region (SEQ ID
      No. 10)

```

```

<400> SEQUENCE: 83

```

-continued

Ser Pro Gly Lys
1

<210> SEQ ID NO 84
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: Myc epitope of junction region (SEQ ID No. 10)

<400> SEQUENCE: 84

Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu
1 5 10

<210> SEQ ID NO 85
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: Residues 18-21 of junction region (SEQ ID No. 10)

<400> SEQUENCE: 85

Asn Gly Ala Ala
1

<210> SEQ ID NO 86
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: N-terminal residues of junction region (SEQ ID No. 10)

<400> SEQUENCE: 86

Thr Val Glu Ser
1

<210> SEQ ID NO 87
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: AB-loop sequence AA143ff

<400> SEQUENCE: 87

Leu Thr Lys Asn Gln
1 5

<210> SEQ ID NO 88
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: EF loop sequence AA198ff

<400> SEQUENCE: 88

Asp Lys Ser Arg Trp Gln Gln
1 5

<210> SEQ ID NO 89
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: linker

<400> SEQUENCE: 89

-continued

Glu Gly Gly Gly Ser
1 5

<210> SEQ ID NO 90
<211> LENGTH: 105
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: engineered CH3 domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (19)..(21)
<223> OTHER INFORMATION: Motif
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (72)..(80)
<223> OTHER INFORMATION: Motif

<400> SEQUENCE: 90

Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp
1 5 10 15
Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe
20 25 30
Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu
35 40 45
Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
50 55 60
Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly
65 70 75 80
Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr
85 90 95
Thr Gln Lys Ser Leu Ser Leu Ser Pro
100 105

<210> SEQ ID NO 91
<211> LENGTH: 110
<212> TYPE: PRT
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: engineered CH3 domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (19)..(21)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (73)..(80)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (83)..(84)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 91

Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp
1 5 10 15
Glu Leu Xaa Xaa Xaa Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe
20 25 30
Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu
35 40 45
Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
50 55 60
Phe Leu Tyr Ser Lys Leu Thr Val Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
65 70 75 80

-continued

Arg Trp Xaa Xaa Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 85 90 95

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 100 105 110

<210> SEQ ID NO 92
 <211> LENGTH: 110
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: engineered CH3 domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (19)..(21)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (73)..(80)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (83)..(84)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr

<400> SEQUENCE: 92

Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp
 1 5 10 15

Glu Leu Xaa Xaa Xaa Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe
 20 25 30

Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu
 35 40 45

Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
 50 55 60

Phe Leu Tyr Ser Lys Leu Thr Val Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 65 70 75 80

Arg Trp Xaa Xaa Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 85 90 95

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 100 105 110

<210> SEQ ID NO 93
 <211> LENGTH: 114
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: engineered CH3 domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (19)..(21)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (75)..(82)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (87)..(88)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 93

Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp
 1 5 10 15

Glu Leu Xaa Xaa Xaa Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe
 20 25 30

Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu

-continued

35	40	45
----	----	----

Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
 50 55 60

Phe Leu Tyr Ser Lys Leu Thr Val Gly Ser Xaa Xaa Xaa Xaa Xaa Xaa
 65 70 75 80

Xaa Xaa Ser Gly Arg Trp Xaa Xaa Gly Asn Val Phe Ser Cys Ser Val
 85 90 95

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 100 105 110

Ser Pro

<210> SEQ ID NO 94
 <211> LENGTH: 114
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: engineered CH3 domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (19)..(21)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (75)..(82)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (87)..(88)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr

<400> SEQUENCE: 94

Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp
1 5 10 15
Glu Leu Xaa Xaa Xaa Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe
20 25 30
Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu
35 40 45
Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
50 55 60
Phe Leu Tyr Ser Lys Leu Thr Val Gly Ser Xaa Xaa Xaa Xaa Xaa Xaa
65 70 75 80
Xaa Xaa Ser Gly Arg Trp Xaa Xaa Gly Asn Val Phe Ser Cys Ser Val
85 90 95
Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
100 105 110

Ser Pro

<210> SEQ ID NO 95
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: engineered CH3 domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (19)..(23)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (20)..(22)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (75)..(82)

-continued

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (85)..(86)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 95

Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp
 1 5 10 15

Glu Leu Xaa Xaa Xaa Xaa Gln Val Ser Leu Thr Cys Leu Val Lys
 20 25 30

Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
 35 40 45

Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly
 50 55 60

Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Xaa Xaa Xaa Xaa Xaa
 65 70 75 80

Xaa Xaa Arg Trp Xaa Xaa Gly Asn Val Phe Ser Cys Ser Val Met His
 85 90 95

Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 100 105 110

<210> SEQ ID NO 96
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: engineered CH3 domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (19)..(23)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (75)..(82)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (85)..(86)
 <223> OTHER INFORMATION: Xaa can be any one of Ala, Asp, Ser or Tyr

<400> SEQUENCE: 96

Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp
 1 5 10 15

Glu Leu Xaa Xaa Xaa Xaa Gln Val Ser Leu Thr Cys Leu Val Lys
 20 25 30

Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
 35 40 45

Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly
 50 55 60

Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Xaa Xaa Xaa Xaa Xaa Xaa
 65 70 75 80

Xaa Xaa Arg Trp Xaa Xaa Gly Asn Val Phe Ser Cys Ser Val Met His
 85 90 95

Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
 100 105 110

157

We claim:

1. A method of producing a functional binding agent, wherein the functional binding agent binds to both a target antigen and a scaffold ligand, wherein the scaffold ligand binds to the backbone of the oligomer of modular antibody domains regardless of the target antigen specificity of the functional binding agent, wherein the functional binding agent is an antibody Fc fragment comprising a CH2 domain and a CH3 domain, with at least one structural loop region, characterized in that said at least one loop region comprises at least one amino acid modification forming at least one modified loop region, wherein said at least one modified loop region binds specifically to said target antigen,

the method comprising the steps of:

(a) providing a library of the functional binding agents by a method comprising

(i) providing a genetic package, wherein the genetic package is yeast, and

(ii) displaying at least two of the functional binding agents by fusing them to the outer surface of the genetic package, so that a genetic package displaying functional binding agents capable of binding both to a target antigen and to a scaffold ligand is prepared,

(b) contacting said library with said target antigen in the presence of a scaffold ligand, wherein said target antigen and said scaffold ligand are different molecules, wherein said scaffold ligand is selected from the group consisting of CD64, CD16, CD32, FcRn and Protein A,

(c) selecting a library member binding to said target antigen in the presence of said scaffold ligand to obtain a functional binding agent which binds to both the target antigen and the scaffold ligand, and

158

(d) producing a preparation of the selected functional binding agent which binds to both the target antigen and the scaffold ligand.

2. The method of claim 1, wherein said library of the functional binding agents contains at least 10^2 independent clones expressing oligomers of modular antibody domains.

3. The method of claim 1, wherein said library of the functional binding agents contains at least 10^6 independent clones expressing oligomers of modular antibody domains.

4. The method of claim 1, wherein said target antigen is a receptor of the erbB class.

5. The method of claim 1, wherein said target antigen is human Her2.

6. The method of claim 1, wherein the selected functional binding agent has a molecular weight of less than 60 kD.

7. The method of claim 1, wherein the selected functional binding agent has a target antigen binding affinity of $K_d < 10^{-8}$ M.

8. The method of claim 1, further comprising the steps of:
(e) affinity maturing the selected functional binding agent by amino acid variation to obtain an affinity matured pool of binding agents, and

(f) selecting a member of said pool which binds to the target antigen in the presence of the scaffold ligand to obtain an affinity matured functional binding agent.

9. The method of claim 8, wherein the selected affinity matured functional binding agent from step 8(f) exhibits at least a 10 fold increase in affinity of binding to the target antigen compared to the selected functional binding agent from step 1(c).

* * * * *