




(86) Date de dépôt PCT/PCT Filing Date: 1999/06/10
(87) Date publication PCT/PCT Publication Date: 1999/12/16
(45) Date de délivrance/Issue Date: 2012/10/23
(85) Entrée phase nationale/National Entry: 2000/12/08
(86) N° demande PCT/PCT Application No.: DK 1999/000314
(87) N° publication PCT/PCT Publication No.: 1999/064619
(30) Priorités/Priorities: 1998/06/10 (US09/111,256);
1998/10/20 (DK PA 1998 01341);
1998/10/20 (DK PA 1998 01340);
1998/10/28 (US60/105,970); 1998/10/28 (US60/106,054);
1998/12/23 (DK PA 1998 01725);
1999/03/05 (DK PA 1999 00309);

(51) Cl.Int./Int.Cl. *C12N 15/56* (2006.01),
A23F 5/24 (2006.01), *C11D 3/386* (2006.01),
C12N 15/10 (2006.01), *C12N 9/24* (2006.01),
C12N 9/42 (2006.01)
(72) Inventeurs/Inventors:
KAUPPINEN, MARKUS SAKARI, DK;
SCHULEIN, MARTIN, DK;
SCHNORR, KIRK, DK;
ANDERSEN, LENE NONBOE, DK;
BJORNVAD, MADSEN, ESKEJUNG, DK
(73) Propriétaire/Owner:
NOVOZYMES A/S, DK
(74) Agent: NORTON ROSE CANADA
S.E.N.C.R.L./S.R.L./LLP

(54) Titre : MANNANASES ISOLEES POUR LE TRAITEMENT DE FIBRES CELLULOSIQUES OU SYNTHETIQUES
(54) Title: ISOLATED MANNANASES FOR USE IN TREATING CELLULOSIC OR SYNTHETIC FIBERS



0.10

(57) Abrégé/Abstract:

Novel mannanases comprising e.g. an amino acid sequence as shown in positions 31-330 of SEQ ID NO: 2 or their homologues may be derived from e.g. Bacillus sp. 1633, or may be encoded by polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO: 1 from nucleotide 91 to nucleotide 990, polynucleotide molecules that encode a polypeptide that is at least 65 % identical to the amino acid sequence of SEQ ID NO: 2 from amino acid residue 31 to amino acid residue 330, or degenerate nucleotide sequences thereof. The mannanases are alkaline and are useful e.g. in cleaning compositions, in a fracturing fluid useful to fracture a subterranean formation, for modifying plant material, and for treatment of cellulosic fibres.

(30) **Priorités(suite)/Priorities(continued):** 1999/03/05 (DKPA 1999 00306); 1999/03/05 (DKPA 1999 00307);
1999/03/05 (DKPA 1999 00308); 1999/03/09 (US60/123,543); 1999/03/10 (US60/123,641); 1999/03/10 (US60/123,623);
1999/03/11 (US60/123,642)

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q	A2	(11) International Publication Number: WO 99/64619 (43) International Publication Date: 16 December 1999 (16.12.99)																																										
(21) International Application Number: PCT/DK99/00314 (22) International Filing Date: 10 June 1999 (10.06.99) (30) Priority Data: <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;">09/111,256</td> <td style="width: 40%;">10 June 1998 (10.06.98)</td> <td style="width: 30%;">US</td> </tr> <tr> <td>PA 1998 01341</td> <td>20 October 1998 (20.10.98)</td> <td>DK</td> </tr> <tr> <td>PA 1998 01340</td> <td>20 October 1998 (20.10.98)</td> <td>DK</td> </tr> <tr> <td>60/105,970</td> <td>28 October 1998 (28.10.98)</td> <td>US</td> </tr> <tr> <td>60/106,054</td> <td>28 October 1998 (28.10.98)</td> <td>US</td> </tr> <tr> <td>PA 1998 01725</td> <td>23 December 1998 (23.12.98)</td> <td>DK</td> </tr> <tr> <td>PA 1999 00308</td> <td>5 March 1999 (05.03.99)</td> <td>DK</td> </tr> <tr> <td>PA 1999 00307</td> <td>5 March 1999 (05.03.99)</td> <td>DK</td> </tr> <tr> <td>PA 1999 00306</td> <td>5 March 1999 (05.03.99)</td> <td>DK</td> </tr> <tr> <td>PA 1999 00309</td> <td>5 March 1999 (05.03.99)</td> <td>DK</td> </tr> <tr> <td>60/123,543</td> <td>9 March 1999 (09.03.99)</td> <td>US</td> </tr> <tr> <td>60/123,641</td> <td>10 March 1999 (10.03.99)</td> <td>US</td> </tr> <tr> <td>60/123,623</td> <td>10 March 1999 (10.03.99)</td> <td>US</td> </tr> <tr> <td>60/123,642</td> <td>11 March 1999 (11.03.99)</td> <td>US</td> </tr> </table> (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).		09/111,256	10 June 1998 (10.06.98)	US	PA 1998 01341	20 October 1998 (20.10.98)	DK	PA 1998 01340	20 October 1998 (20.10.98)	DK	60/105,970	28 October 1998 (28.10.98)	US	60/106,054	28 October 1998 (28.10.98)	US	PA 1998 01725	23 December 1998 (23.12.98)	DK	PA 1999 00308	5 March 1999 (05.03.99)	DK	PA 1999 00307	5 March 1999 (05.03.99)	DK	PA 1999 00306	5 March 1999 (05.03.99)	DK	PA 1999 00309	5 March 1999 (05.03.99)	DK	60/123,543	9 March 1999 (09.03.99)	US	60/123,641	10 March 1999 (10.03.99)	US	60/123,623	10 March 1999 (10.03.99)	US	60/123,642	11 March 1999 (11.03.99)	US	(72) Inventors; and (75) Inventors/Applicants (for US only): KAUPPINEN, Markus, Sakari [FI/DK]; Egegade 10,5, DK-2200 Copenhagen N. (DK). SCHÜLEIN, Martin [DK/DK]; Wiedeweltsgade 51, DK-2100 Copenhagen Ø (DK). SCHNORR, Kirk [US/DK]; Nørrebrogade 44A, 1. tv., DK-2200 Copenhagen N. (DK). ANDERSEN, Lene, Nonboe [DK/DK]; Laksevej 11, DK-3450 Allerød (DK). BJØRNVAD, Mads, Eskelund [DK/DK]; Dr. Abildgaards Allé 8, 3th, DK-1955 Frederiksberg (DK). (74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i> <i>With an indication in relation to deposited biological material furnished under Rule 13bis separately from the description.</i>
09/111,256	10 June 1998 (10.06.98)	US																																										
PA 1998 01341	20 October 1998 (20.10.98)	DK																																										
PA 1998 01340	20 October 1998 (20.10.98)	DK																																										
60/105,970	28 October 1998 (28.10.98)	US																																										
60/106,054	28 October 1998 (28.10.98)	US																																										
PA 1998 01725	23 December 1998 (23.12.98)	DK																																										
PA 1999 00308	5 March 1999 (05.03.99)	DK																																										
PA 1999 00307	5 March 1999 (05.03.99)	DK																																										
PA 1999 00306	5 March 1999 (05.03.99)	DK																																										
PA 1999 00309	5 March 1999 (05.03.99)	DK																																										
60/123,543	9 March 1999 (09.03.99)	US																																										
60/123,641	10 March 1999 (10.03.99)	US																																										
60/123,623	10 March 1999 (10.03.99)	US																																										
60/123,642	11 March 1999 (11.03.99)	US																																										
(54) Title: NOVEL MANNANASES																																												
0.10																																												
(57) Abstract Novel mannanases comprising e.g. an amino acid sequence as shown in positions 31-330 of SEQ ID NO: 2 or their homologues may be derived from e.g. <i>Bacillus</i> sp. 1633, or may be encoded by polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO: 1 from nucleotide 91 to nucleotide 990, polynucleotide molecules that encode a polypeptide that is at least 65 % identical to the amino acid sequence of SEQ ID NO: 2 from amino acid residue 31 to amino acid residue 330, or degenerate nucleotide sequences thereof. The mannanases are alkaline and are useful e.g. in cleaning compositions, in a fracturing fluid useful to fracture a subterranean formation, for modifying plant material, and for treatment of cellulosic fibres.																																												

ISOLATED MANNANASES FOR USE IN TREATING CELLULOSIC OR SYNTHETIC FIBERS

The present invention relates to microbial mannanases, more specifically to microbial enzymes exhibiting mannanase activity as their major enzymatic activity in the neutral and alkaline pH ranges; to a method of producing such enzymes; and to methods for using such enzymes in the paper and pulp, textile, oil drilling, cleaning, laundering, detergent and cellulose fiber processing industries.

10

BACKGROUND OF THE INVENTION

Mannan containing polysaccharides are a major component of the hemicellulose fraction in woods and endosperm in many leguminous seeds and in some mature seeds of non-leguminous plants. Essentially unsubstituted linear beta-1,4-mannan is found in some non-leguminous plants. Unsubstituted beta-1,4-mannan which is present e.g. in ivory nuts resembles cellulose in the conformation of the individual polysaccharide chains, and is water-insoluble. In leguminous seeds, water-soluble galactomannan is the main storage carbohydrate comprising up to 20% of the total dry weight. Galactomannans have a linear beta-1,4-mannan backbone substituted with single alpha-1,6-galactose, optionally substituted with acetyl groups. Mannans are also found in several monocotyledonous plants and are the most abundant polysaccharides in the cell wall material in palm kernel meal. Glucomannans are linear polysaccharides with a backbone of beta-1,4-linked mannose and glucose alternating in a more or less regular manner, the backbone optionally being substituted with galactose and/or acetyl groups. Mannans, galactomannans, glucomannans and galactoglucomannans (i.e. glucomannan backbones with branched galactose) contribute to more than 50% of the softwood hemicellulose. Moreover, the

30

cellulose of many red algae contains a significant amount of mannose.

Mannanases have been identified in several *Bacillus* organisms. For example, Talbot et al., Appl. Environ.

5 Microbiol., Vol.56, No. 11, pp. 3505-3510 (1990) describes a beta-mannanase derived from *Bacillus stearothermophilus* in dimer form having molecular weight of 162 kDa and an optimum pH of 5.5-7.5. Mendoza et al., World J. Microbiol. Biotech., Vol. 10, No. 5, pp. 551-555 (1994) describes a beta-mannanase derived
10 from *Bacillus subtilis* having a molecular weight of 38 kDa, an optimum activity at pH 5.0 and 55°C and a pI of 4.8. JP-A-03047076 discloses a beta-mannanase derived from *Bacillus* sp., having a molecular weight of 37 ± 3 kDa measured by gel filtration, an optimum pH of 8-10 and a pI of 5.3-5.4. JP-A-
15 63056289 describes the production of an alkaline, thermostable beta-mannanase which hydrolyses beta-1,4-D-mannopyranoside bonds of e.g. mannans and produces manno-oligosaccharides. JP-A-63036775 relates to the *Bacillus* microorganism FERM P-8856 which produces beta-mannanase and beta-mannosidase at an alkaline pH.
20 JP-A-08051975 discloses alkaline beta-mannanases from alkalophilic *Bacillus* sp. AM-001 having molecular weights of 43 ± 3 kDa and 57 ± 3 kDa and optimum pH of 8-10. A purified mannanase from *Bacillus amyloliquefaciens* useful in the bleaching of pulp and paper and a method of preparation thereof
25 is disclosed in WO 97/11164. WO 91/18974 describes a hemicellulase such as a glucanase, xylanase or mannanase active at an extreme pH and temperature. WO 94/25576 discloses an enzyme from *Aspergillus aculeatus*, CBS 101.43, exhibiting mannanase activity which may be useful for degradation or
30 modification of plant or algae cell wall material. WO 93/24622 discloses a mannanase isolated from *Trichoderma reesei* useful for bleaching lignocellulosic pulps.

WO 95/35362 discloses cleaning compositions containing plant cell wall degrading enzymes having pectinase and/or hemicellulase and optionally cellulase activity for the removal of stains of vegetable origin and further discloses an alkaline
5 mannanase from the strain C11SB.G17.

It is an object of the present invention to provide a novel and efficient enzyme exhibiting mannanase activity also in the alkaline pH range, e.g. when applied in cleaning compositions or different industrial processes.

10

SUMMARY OF THE INVENTION

The inventors have now found novel enzymes having substantial mannanase activity, i.e. enzymes exhibiting
15 mannanase activity which may be obtained from a bacterial strain of the genus *Bacillus* and have succeeded in identifying DNA sequences encoding such enzymes. The DNA sequences are listed in the sequence listing as SEQ ID No. 1, 5, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29 and 31; and the deduced amino acid sequences
20 are listed in the sequence listing as SEQ ID No. 2, 6, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30 and 32, respectively. It is believed that the novel enzymes will be classified according to the Enzyme Nomenclature in the Enzyme Class EC 3.2.1.78.

In a first aspect, the present invention relates to a
25 mannanase which is i) a polypeptide produced by *Bacillus* sp. I633, ii) a polypeptide comprising an amino acid sequence as shown in positions 31-330 of SEQ ID NO:2, or iii) an analogue of the polypeptide defined in i) or ii) which is at least 65% homologous with said polypeptide, is derived from said
30 polypeptide by substitution, deletion or addition of one or several amino acids, or is immunologically reactive with a polyclonal antibody raised against said polypeptide in purified

form.

Within one aspect, the present invention provides an isolated polynucleotide molecule selected from the group consisting of (a) polynucleotide molecules encoding a polypeptide having
5 mannanase activity and comprising a sequence of nucleotides as shown in SEQ ID NO: 1 from nucleotide 91 to nucleotide 990; (b) species homologs of (a); (c) polynucleotide molecules that encode a polypeptide having mannanase activity that is at least 65% identical to the amino acid sequence of SEQ ID NO: 2 from
10 amino acid residue 31 to amino acid residue 330; (d) molecules complementary to (a), (b) or (c); and (e) degenerate nucleotide sequences of (a), (b), (c) or (d).

The plasmid pBXM3 comprising the polynucleotide molecule (the DNA sequence) encoding a mannanase of the present invention
15 has been transformed into a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH,
20 Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 29 May 1998 under the deposition number DSM 12197.

Within another aspect of the invention there is provided an expression vector comprising the following operably linked elements: a transcription promoter; a DNA segment selected from
25 the group consisting of (a) polynucleotide molecules encoding a polypeptide having mannanase activity and comprising a sequence of nucleotides as shown in SEQ ID NO: 1 from nucleotide 91 to nucleotide 990; (b) species homologs of (a); (c) polynucleotide molecules that encode a polypeptide having mannanase activity
30 that is at least 65% identical to the amino acid sequence of SEQ ID NO: 2 from amino acid residue 31 to amino acid residue 330;

and (d) degenerate nucleotide sequences of (a), (b), or (c); and a transcription terminator.

Within yet another aspect of the present invention there is provided a cultured cell into which has been introduced an expression vector as disclosed above, wherein said cell expresses the polypeptide encoded by the DNA segment.

Further aspects of the present invention provide an isolated polypeptide having mannanase activity selected from the group consisting of (a) polypeptide molecules comprising a sequence of amino acid residues as shown in SEQ ID NO:2 from amino acid residue 31 to amino acid residue 330; (b) species homologs of (a); and a fusion protein having mannanase activity comprising a first polypeptide part exhibiting mannanase activity and a second polypeptide part exhibiting cellulose binding function, the second polypeptide preferably being a cellulose binding domain (CBD), such as a fusion protein represented by SEQ ID NO:4.

Within another aspect of the present invention there is provided a composition comprising a purified polypeptide according to the invention in combination with other polypeptides.

Within another aspect of the present invention there are provided methods for producing a polypeptide according to the invention comprising culturing a cell into which has been introduced an expression vector as disclosed above, whereby said cell expresses a polypeptide encoded by the DNA segment and recovering the polypeptide.

The novel enzyme of the present invention is useful for the treatment of cellulosic material, especially cellulose-containing fiber, yarn, woven or non-woven fabric, treatment of mechanical paper-making pulps, kraft pulps or recycled waste paper, and for retting of fibres. The treatment can be carried out during the processing of cellulosic material into a material

ready for manufacture of paper or of garment or fabric, the latter e.g. in the desizing or scouring step; or during industrial or household laundering of such fabric or garment.

Accordingly, in further aspects the present invention
5 relates to a cleaning or detergent composition comprising the enzyme of the invention; and to use of the enzyme of the invention for the treatment, eg cleaning, of cellulose-containing fibers, yarn, woven or non-woven fabric, as well as synthetic or partly synthetic fabric.

10 It is contemplated that the enzyme of the invention is useful in an enzymatic scouring process and/or desizing (removal of mannan size) in the preparation of cellulosic material e.g. for proper response in subsequent dyeing operations. The enzyme is also useful for removal of mannan containing print paste.
15 Further, detergent compositions comprising the novel enzyme are capable of removing or bleaching certain soils or stains present on laundry, especially soils and spots resulting from mannan containing food, plants, and the like. Further, treatment with cleaning or detergent compositions comprising the novel enzyme
20 can improve whiteness as well as prevent binding of certain soils to the cellulosic material.

Accordingly, the present invention also relates to cleaning compositions, including laundry, dishwashing, hard surface cleaner, personal cleansing and oral/dental compositions, com-
25 prising the mannanase of the invention. Further, the present invention relates to such cleaning compositions comprising a mannanase and an enzyme selected from cellulases, proteases, lipases, amylases, pectin degrading enzymes and xyloglucanases, such compositions providing superior cleaning performance, i.e.
30 superior stain removal, dingy cleaning or whiteness maintenance.

DEFINITIONS

Prior to discussing this invention in further detail, the following terms will first be defined.

- 5 The term "ortholog" (or "species homolog") denotes a polypeptide or protein obtained from one species that has homology to an analogous polypeptide or protein from a different species.

The term "paralog" denotes a polypeptide or protein obtained
10 from a given species that has homology to a distinct polypeptide or protein from that same species.

The term "expression vector" denotes a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for
15 its transcription. Such additional segments may include promoter and terminator sequences, and may optionally include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, and the like. Expression vectors are generally derived from plasmid or viral DNA, or may
20 contain elements of both. The expression vector of the invention may be any expression vector that is conveniently subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which the vector is to be introduced. Thus, the vector may be an autonomously replicating
25 vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the
30 chromosome(s) into which it has been integrated.

The term "recombinant expressed" or "recombinantly expressed" used herein in connection with expression of a

polypeptide or protein is defined according to the standard definition in the art. Recombinantly expression of a protein is generally performed by using an expression vector as described immediately above.

5 The term "isolated", when applied to a polynucleotide molecule, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such
10 isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as
15 promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, Nature 316:774-78, 1985). The term "an isolated polynucleotide" may alternatively be termed "a cloned polynucleotide".

20 When applied to a protein/polypeptide, the term "isolated" indicates that the protein is found in a condition other than its native environment. In a preferred form, the isolated protein is substantially free of other proteins, particularly other homologous proteins (*i.e.* "homologous impurities" (see below)).
25 It is preferred to provide the protein in a greater than 40% pure form, more preferably greater than 60% pure form.

Even more preferably it is preferred to provide the protein in a highly purified form, *i.e.*, greater than 80% pure, more preferably greater than 95% pure, and even more preferably
30 greater than 99% pure, as determined by SDS-PAGE.

The term "isolated protein/polypeptide may alternatively be termed "purified protein/polypeptide".

The term "homologous impurities" means any impurity (e.g. another polypeptide than the polypeptide of the invention) which originate from the homologous cell where the polypeptide of the invention is originally obtained from.

5 The term "obtained from" as used herein in connection with a specific microbial source, means that the polynucleotide and/or polypeptide is produced by the specific source (homologous expression), or by a cell in which a gene from the source have been inserted (heterologous expression).

10 The term "operably linked", when referring to DNA segments, denotes that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription initiates in the promoter and proceeds through the coding segment to the terminator

15 The term "polynucleotide" denotes a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic
20 molecules.

The term "complements of polynucleotide molecules" denotes polynucleotide molecules having a complementary base sequence and reverse orientation as compared to a reference sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.
25

The term "degenerate nucleotide sequence" denotes a sequence of nucleotides that includes one or more degenerate codons (as compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of
30 nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "promoter" denotes a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

5 The term "secretory signal sequence" denotes a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger peptide is commonly cleaved to remove the
10 secretory peptide during transit through the secretory pathway.

The term "enzyme core" denotes a single domain enzyme which may or may not have been modified or altered, but which has retained its original activity; the catalytic domain as known in the art has remained intact and functional.

15 By the term "linker" or "spacer" is meant a polypeptide comprising at least two amino acids which may be present between the domains of a multidomain protein, for example, an enzyme comprising an enzyme core and a binding domain such as a cellulose binding domain (CBD) or any other enzyme hybrid, or
20 between two proteins or polypeptides expressed as a fusion polypeptide, for example a fusion protein comprising two core enzymes. For example, the fusion protein of an enzyme core with a CBD is provided by fusing a DNA sequence encoding the enzyme core, a DNA sequence encoding the linker and a DNA sequence
25 encoding the CBD sequentially into one open reading frame and expressing this construct.

The term "mannanase" or "galactomannanase" denotes a mannanase enzyme defined according to the art as officially being named mannan endo-1,4-beta-mannosidase and having the alternative names beta-mannanase and endo-1,4-mannanase and catalysing
30 hydrolyses of 1,4-beta-D-mannosidic linkages in mannans, galactomannans, glucomannans, and galactoglucomannans which enzyme

is classified according to the Enzyme Nomenclature as EC 3.2.1.78.

DETAILED DESCRIPTION OF THE INVENTION

5 HOW TO USE A SEQUENCE OF THE INVENTION TO GET OTHER RELATED SEQUENCES: The disclosed sequence information herein relating to a polynucleotide sequence encoding a mannanase of the invention can be used as a tool to identify other homologous mannanases. For instance, polymerase chain reaction (PCR) can be used to
10 amplify sequences encoding other homologous mannanases from a variety of microbial sources, in particular of different *Bacillus* species.

ASSAY FOR ACTIVITY TEST

15 A polypeptide of the invention having mannanase activity may be tested for mannanase activity according to standard test procedures known in the art, such as by applying a solution to be tested to 4 mm diameter holes punched out in agar plates containing 0.2% AZCL galactomannan (carob), i.e. substrate for
20 the assay of endo-1,4-beta-D-mannanase available as CatNo.I-AZGMA from the company Megazyme.

POLYNUCLEOTIDES

25 Within preferred embodiments of the invention an isolated polynucleotide of the invention will hybridize to similar sized regions of SEQ ID NO: 1, or a sequence complementary thereto, under at least medium stringency conditions.

 In particular polynucleotides of the invention will
30 hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:1 or a partial sequence comprising the segment shown in positions 91-990 of SEQ

ID NO:1 which segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in positions 91-990 of SEQ ID NO:1 which subsequence has a length of at least about 100
5 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail below. Suitable experimental conditions for determining hybridization at medium, or high stringency between a nucleotide probe and a homologous DNA or RNA sequence involves presoaking
10 of the filter containing the DNA fragments or RNA to hybridize in 5 x SSC (Sodium chloride/Sodium citrate, Sambrook et al. 1989) for 10 min, and prehybridization of the filter in a solution of 5 x SSC, 5 x Denhardt's solution (Sambrook et al. 1989), 0.5 % SDS and 100 µg/ml of denatured sonicated salmon
15 sperm DNA (Sambrook et al. 1989), followed by hybridization in the same solution containing a concentration of 10ng/ml of a random-primed (Feinberg, A. P. and Vogelstein, B. (1983) Anal. Biochem. 132:6-13), 32P-dCTP-labeled (specific activity higher than 1×10^9 cpm/µg) probe for 12 hours at ca. 45°C. The filter
20 is then washed twice for 30 minutes in 2 x SSC, 0.5 % SDS at least 60°C (medium stringency), still more preferably at least 65°C (medium/high stringency), even more preferably at least 70°C (high stringency), and even more preferably at least 75°C (very high stringency).

25 Molecules to which the oligonucleotide probe hybridizes under these conditions are detected using a x-ray film.

Other useful isolated polynucleotides are those which will hybridize to similar sized regions of SEQ ID NO: 5, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17,
30 SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29 or SEQ ID NO: 31, respectively, or a sequence complementary thereto, under at least medium stringency

conditions.

Particularly useful are polynucleotides which will hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:5 or a partial sequence comprising the segment shown in positions 94-1032 of SEQ ID NO:5 which segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in positions 94-1032 of SEQ ID NO:5 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above; as well as polynucleotides which will hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:9 or a partial sequence comprising the segment shown in positions 94-1086 of SEQ ID NO:9 which segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in positions 94-1086 of SEQ ID NO:9 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above; as well as polynucleotides which will hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:11 or a partial sequence comprising the segment shown in positions 97-993 of SEQ ID NO:11 which segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in positions 97-993 of SEQ ID NO:11 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above; as well as polynucleotides which will hybridize to a denatured double-

stranded DNA probe comprising either the full sequence shown in SEQ ID NO:13 or a partial sequence comprising the segment shown in positions 498-1464 of SEQ ID NO:13 which segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in positions 498-1464 of SEQ ID NO:13 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above; as well as

10 polynucleotides which will hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:15 or a partial sequence comprising the segment shown in positions 204-1107 of SEQ ID NO:15 which segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in positions 204-1107 of SEQ ID NO:15 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above; as well as

20 polynucleotides which will hybridize to a denatured double-stranded DNA probe comprising either the sequence shown in SEQ ID NO:17 or any probe comprising a subsequence of SEQ ID NO:17 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at

25 high stringency conditions as described in detail above; as well as polynucleotides which will hybridize to a denatured double-stranded DNA probe comprising either the sequence shown in SEQ ID NO:19 or any probe comprising a subsequence of SEQ ID NO:19 which subsequence has a length of at least about 100 base pairs

30 under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above; as well as polynucleotides which will hybridize to a denatured double-

stranded DNA probe comprising either the full sequence shown in SEQ ID NO:21 or a partial sequence comprising the segment shown in positions 88-960 of SEQ ID NO:21 which segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in positions 88-960 of SEQ ID NO:21 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above; as well as

10 polynucleotides which will hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:23 or any probe comprising a subsequence of SEQ ID NO:23 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but

15 preferably at high stringency conditions as described in detail above; as well as polynucleotides which will hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:25 or a partial sequence comprising the segment shown in positions 904-1874 of SEQ ID NO:25 which

20 segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in positions 904-1874 of SEQ ID NO:25 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high

25 stringency conditions as described in detail above; as well as polynucleotides which will hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:27 or a partial sequence comprising the segment shown in positions 498-1488 of SEQ ID NO:27 which segment encodes for

30 the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in positions 498-1488 of SEQ ID NO:27 which subsequence has a

length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above; as well as polynucleotides which will hybridize to a denatured double-
5 stranded DNA probe comprising either the full sequence shown in SEQ ID NO:29 or a partial sequence comprising the segment shown in positions 79-1083 of SEQ ID NO:29 which segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in
10 positions 79-1083 of SEQ ID NO:29 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above; as well as polynucleotides which will hybridize to a denatured double-
15 stranded DNA probe comprising either the full sequence shown in SEQ ID NO:31 or a partial sequence comprising the segment shown in positions 1779-2709 of SEQ ID NO:31 which segment encodes for the catalytically active domain or enzyme core of the mannanase of the invention or any probe comprising a subsequence shown in
20 positions 1779-2709 of SEQ ID NO:31 which subsequence has a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail above.

As previously noted, the isolated polynucleotides of the
25 present invention include DNA and RNA. Methods for isolating DNA and RNA are well known in the art. DNA and RNA encoding genes of interest can be cloned in Gene Banks or DNA libraries by means of methods known in the art.

Polynucleotides encoding polypeptides having mannanase
30 activity of the invention are then identified and isolated by, for example, hybridization or PCR.

The present invention further provides counterpart polypeptides and polynucleotides from different bacterial strains (orthologs or paralogs). Of particular interest are mannanase polypeptides from gram-positive alkalophilic strains, including species of *Bacillus* such as *Bacillus* sp., *Bacillus agaradhaerens*, *Bacillus halodurans*, *Bacillus clausii* and *Bacillus licheniformis*; and mannanase polypeptides from *Thermoanaerobacter* group, including species of *Caldicellulosiruptor*. Also mannanase polypeptides from the fungus *Humicola* or *Scytalidium*, in particular the species *Humicola insolens* or *Scytalidium thermophilum*, are of interest.

Species homologues of a polypeptide with mannanase activity of the invention can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a DNA sequence of the present invention can be cloned using chromosomal DNA obtained from a cell type that expresses the protein. Suitable sources of DNA can be identified by probing Northern or Southern blots with probes designed from the sequences disclosed herein. A library is then prepared from chromosomal DNA of a positive cell line. A DNA sequence of the invention encoding an polypeptide having mannanase activity can then be isolated by a variety of methods, such as by probing with probes designed from the sequences disclosed in the present specification and claims or with one or more sets of degenerate probes based on the disclosed sequences. A DNA sequence of the invention can also be cloned using the polymerase chain reaction, or PCR (Mullis, U.S. Patent 4,683,202), using primers designed from the sequences disclosed herein. Within an additional method, the DNA library can be used to transform or transfect host cells, and expression of the DNA of interest can be detected with an antibody (mono-clonal or polyclonal) raised

against the mannanase cloned from *B.sp.*, expressed and purified as described in Materials and Methods and Example 1, or by an activity test relating to a polypeptide having mannanase activity.

5 The mannanase encoding part of the DNA sequence (SEQ ID NO:1) cloned into plasmid pBXM3 present in *Escherichia coli* DSM 12197 and/or an analogue DNA sequence of the invention may be cloned from a strain of the bacterial species *Bacillus sp.* I633, or another or related organism as described herein.

10 The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:5) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of
15 Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 18 May 1998 under the deposition number DSM 12180; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:5) and/or
20 an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Bacillus agaradhaerens*, for example from the type strain DSM 8721, or another or related organism as described herein.

The mannanase encoding part of the polynucleotide molecule
25 (the DNA sequence of SEQ ID NO:9) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen
30 und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 7 October 1998 under the deposition number DSM 12433; this mannanase encoding part of the

polynucleotide molecule (the DNA sequence of SEQ ID NO:9) and/or an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Bacillus* sp. AAI12 or another or related organism as described herein.

5 The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:11) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of
10 Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 9 October 1998 under the deposition number DSM 12441; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:11)
15 and/or an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Bacillus halodurans* or another or related organism as described herein.

 The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:13) was transformed a strain of
20 the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig,
25 Federal Republic of Germany, on 11 May 1995 under the deposition number DSM 9984; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:13) and/or an analogue DNA sequence thereof may be cloned from a strain of the fungal species *Humicola insolens* or another or
30 related organism as described herein.

 The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:15) was transformed a strain of

the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 5 October 1998 under the deposition number DSM 12432; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:15) and/or an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Bacillus sp.* AA349 or another or related organism as described herein.

The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:17) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 4 June 1999 under the deposition number DSM 12847; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:17) and/or an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Bacillus sp.* or another or related organism as described herein.

The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:19) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 4 June 1999 under the deposition

number DSM 12848; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:19) and/or an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Bacillus* sp. or another or
5 related organism as described herein.

The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:21) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International
10 Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 4 June 1999 under the deposition number DSM 12849; this mannanase encoding part of the
15 polynucleotide molecule (the DNA sequence of SEQ ID NO:21) and/or an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Bacillus clausii* or another or related organism as described herein.

The mannanase encoding part of the polynucleotide molecule
20 (the DNA sequence of SEQ ID NO:23) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen
25 und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 4 June 1999 under the deposition number DSM 12850; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:23) and/or an analogue DNA sequence thereof may be cloned from a
30 strain of the bacterial species *Bacillus* sp. or another or related organism as described herein.

The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:25) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International
5 Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 4 June 1999 under the deposition number DSM 12846; this mannanase encoding part of the
10 polynucleotide molecule (the DNA sequence of SEQ ID NO:25) and/or an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Bacillus* sp. or another or related organism as described herein.

The mannanase encoding part of the polynucleotide molecule
15 (the DNA sequence of SEQ ID NO:27) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen
20 und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 4 June 1999 under the deposition number DSM 12851; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:27) and/or an analogue DNA sequence thereof may be cloned from a
25 strain of the bacterial species *Bacillus* sp. or another or related organism as described herein.

The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:29) was transformed a strain of the *Escherichia coli* which was deposited by the inventors
30 according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen

und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 4 June 1999 under the deposition number DSM 12852; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:29) and/or an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Bacillus licheniformis* or another or related organism as described herein.

The mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:31) was transformed a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 5 October 1998 under the deposition number DSM 12436; this mannanase encoding part of the polynucleotide molecule (the DNA sequence of SEQ ID NO:31) and/or an analogue DNA sequence thereof may be cloned from a strain of the bacterial species *Caldicellulosiruptor* sp. or another or related organism as described herein.

Alternatively, the analogous sequence may be constructed on the basis of the DNA sequence obtainable from the plasmid present in *Escherichia coli* DSM 12197 (which is believed to be identical to the attached SEQ ID NO:1), the plasmid present in *Escherichia coli* DSM 12180 (which is believed to be identical to the attached SEQ ID NO:5), the plasmid present in *Escherichia coli* DSM 12433 (which is believed to be identical to the attached SEQ ID NO:9), the plasmid present in *Escherichia coli* DSM 12441 (which is believed to be identical to the attached SEQ ID NO:11), the plasmid present in *Escherichia coli* DSM 9984 (which is believed to be identical to the attached SEQ ID NO:13), the plasmid present in *Escherichia coli* DSM 12432 (which

is believed to be identical to the attached SEQ ID NO:15), the plasmid present in *Escherichia coli* DSM 12847 (which is believed to be identical to the attached SEQ ID NO:17), the plasmid present in *Escherichia coli* DSM 12848 (which is believed to be identical to the attached SEQ ID NO:19), the plasmid present in *Escherichia coli* DSM 12849 (which is believed to be identical to the attached SEQ ID NO:21), the plasmid present in *Escherichia coli* DSM 12850 (which is believed to be identical to the attached SEQ ID NO:23), the plasmid present in *Escherichia coli* DSM 12846 (which is believed to be identical to the attached SEQ ID NO:25), the plasmid present in *Escherichia coli* DSM 12851 (which is believed to be identical to the attached SEQ ID NO:27), the plasmid present in *Escherichia coli* DSM 12852 (which is believed to be identical to the attached SEQ ID NO:29) or the plasmid present in *Escherichia coli* DSM 12436 (which is believed to be identical to the attached SEQ ID NO:31), e.g be a sub-sequence thereof, and/or by introduction of nucleotide substitutions which do not give rise to another amino acid sequence of the mannanase encoded by the DNA sequence, but which corresponds to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions which may give rise to a different amino acid sequence (i.e. a variant of the mannan degrading enzyme of the invention).

25

POLYPEPTIDES

The sequence of amino acids in positions 31-490 of SEQ ID NO: 2 is a mature mannanase sequence. The sequence of amino acids nos. 1-30 of SEQ ID NO: 2 is the signal peptide. It is believed that the subsequence of amino acids in positions 31-330 of SEQ ID NO: 2 is the catalytic domain of the mannanase enzyme and that the mature enzyme additionally comprises a linker in

positions 331-342 and at least one C-terminal domain of unknown function in positions 343-490. Since the object of the present invention is to obtain a polypeptide which exhibits mannanase activity, the present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 31-330 of SEQ ID NO: 2, ie a catalytical domain, optionally operably linked, either N-terminally or C-terminally, to one or two or more than two other domains of a different functionality. The domain having the subsequence of amino acids nos. 343-490 of SEQ ID NO: 2 is a domain of the mannanase enzyme of unknown function, this domain being highly homologous with similar domains in known mannanases, cf. example 1.

The sequence of amino acids in positions 32-494 of SEQ ID NO:6 is a mature mannanase sequence. The sequence of amino acids nos. 1-31 of SEQ ID NO:6 is the signal peptide. It is believed that the subsequence of amino acids in positions 32-344 of SEQ ID NO:6 is the catalytic domain of the mannanase enzyme and that the mature enzyme additionally comprises at least one C-terminal domain of unknown function in positions 345-494. Since the object of the present invention is to obtain a polypeptide which exhibits mannanase activity, the present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 32-344 of SEQ ID NO: 6, ie a catalytical domain, optionally operably linked, either N-terminally or C-terminally, to one or two or more than two other domains of a different functionality.

The sequence of amino acids in positions 32-586 of SEQ ID NO:10 is a mature mannanase sequence. The sequence of amino acids nos. 1-31 of SEQ ID NO:10 is the signal peptide. It is believed that the subsequence of amino acids in positions 32-362 of SEQ ID NO:10 is the catalytic domain of the mannanase enzyme and that the mature enzyme additionally comprises at least one C-terminal domain of unknown function in positions 363-586.

Since the object of the present invention is to obtain a polypeptide which exhibits mannanase activity, the present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 32-362 of SEQ ID NO: 10, ie a catalytical domain, optionally operably linked, either N-terminally or C-terminally, to one or two or more than two other domains of a different functionality.

The sequence of amino acids in positions 33-331 of SEQ ID NO:12 is a mature mannanase sequence. The sequence of amino acids nos. 1-32 of SEQ ID NO:12 is the signal peptide. It is believed that the subsequence of amino acids in positions 33-331 of SEQ ID NO:12 is the catalytic domain of the mannanase enzyme. This mannanase enzyme core comprising the sequence of amino acids nos. 33-331 of SEQ ID NO: 12, ie a catalytical domain, may or may not be operably linked, either N-terminally or C-terminally, to one or two or more than two other domains of a different functionality, ie being part of a fusion protein.

The sequence of amino acids in positions 22-488 of SEQ ID NO:14 is a mature mannanase sequence. The sequence of amino acids nos. 1-21 of SEQ ID NO:14 is the signal peptide. It is believed that the subsequence of amino acids in positions 166-488 of SEQ ID NO:14 is the catalytic domain of the mannanase enzyme and that the mature enzyme additionally comprises at least one N-terminal domain of unknown function in positions 22-164. Since the object of the present invention is to obtain a polypeptide which exhibits mannanase activity, the present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 166-488 of SEQ ID NO: 14, ie a catalytical domain, optionally operably linked, either N-terminally or C-terminally, to one or two or more than two other domains of a different functionality.

The sequence of amino acids in positions 26-369 of SEQ ID NO:16 is a mature mannanase sequence. The sequence of amino acids nos. 1-25 of SEQ ID NO:16 is the signal peptide. It is believed that the subsequence of amino acids in positions 68-369 of SEQ ID NO:16 is the catalytic domain of the mannanase enzyme and that the mature enzyme additionally comprises at least one N-terminal domain of unknown function in positions 26-67. Since the object of the present invention is to obtain a polypeptide which exhibits mannanase activity, the present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 68-369 of SEQ ID NO:16, ie a catalytical domain, optionally operably linked, either N-terminally or C-terminally, to one or two or more than two other domains of a different functionality.

The sequence of amino acids of SEQ ID NO:18 is a partial sequence forming part of a mature mannanase sequence. The present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 1-305 of SEQ ID NO: 18.

The sequence of amino acids of SEQ ID NO:20 is a partial sequence forming part of a mature mannanase sequence. The present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 1-132 of SEQ ID NO:20.

The sequence of amino acids in positions 29-320 of SEQ ID NO:22 is a mature mannanase sequence. The sequence of amino acids nos. 1-28 of SEQ ID NO:22 is the signal peptide. It is believed that the subsequence of amino acids in positions 29-320 of SEQ ID NO:22 is the catalytic domain of the mannanase enzyme. This mannanase enzyme core comprising the sequence of amino acids nos. 29-320 of SEQ ID NO:22, ie a catalytical domain, may or may not be operably linked, either N-terminally or C-terminally, to one or two or more than two other domains of a different functionality, ie being part of a fusion protein.

The sequence of amino acids of SEQ ID NO:24 is a partial sequence forming part of a mature mannanase sequence. The present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 29-188 of SEQ ID NO:24.

5 The sequence of amino acids in positions 30-815 of SEQ ID NO:26 is a mature mannanase sequence. The sequence of amino acids nos. 1-29 of SEQ ID NO:26 is the signal peptide. It is believed that the subsequence of amino acids in positions 301-625 of SEQ ID NO:26 is the catalytic domain of the mannanase
10 enzyme and that the mature enzyme additionally comprises at least two N-terminal domain of unknown function in positions 44-166 and 195-300, respectively, and a C-terminal domain of unknown function in positions 626-815. Since the object of the present invention is to obtain a polypeptide which exhibits
15 mannanase activity, the present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 301-625 of SEQ ID NO:26, ie a catalytical domain, optionally operably linked, either N-terminally or C-terminally, to one or two or more than two other domains of a different functionality.

20 The sequence of amino acids in positions 38-496 of SEQ ID NO:28 is a mature mannanase sequence. The sequence of amino acids nos. 1-37 of SEQ ID NO:28 is the signal peptide. It is believed that the subsequence of amino acids in positions 166-496 of SEQ ID NO:28 is the catalytic domain of the mannanase
25 enzyme and that the mature enzyme additionally comprises at least one N-terminal domain of unknown function in positions 38-165. Since the object of the present invention is to obtain a polypeptide which exhibits mannanase activity, the present invention relates to any mannanase enzyme comprising the
30 sequence of amino acids nos. 166-496 of SEQ ID NO:28, ie a catalytical domain, optionally operably linked, either N-terminally or C-terminally, to one or two or more than two other

domains of a different functionality.

The sequence of amino acids in positions 26-361 of SEQ ID NO:30 is a mature mannanase sequence. The sequence of amino acids nos. 1-25 of SEQ ID NO:30 is the signal peptide. It is
5 believed that the subsequence of amino acids in positions 26-361 of SEQ ID NO:30 is the catalytic domain of the mannanase enzyme. This mannanase enzyme core comprising the sequence of amino acids nos. 26-361 of SEQ ID NO:30, ie a catalytical domain, may or may not be optionally operably linked, either N-terminally or
10 C-terminally, to one or two or more than two other domains of a different functionality.

The sequence of amino acids in positions 23-903 of SEQ ID NO:32 is a mature mannanase sequence. The sequence of amino acids nos. 1-22 of SEQ ID NO:32 is the signal peptide. It is
15 believed that the subsequence of amino acids in positions 593-903 of SEQ ID NO:32 is the catalytic domain of the mannanase enzyme and that the mature enzyme additionally comprises at least three N-terminal domains of unknown function in positions 23-214, 224-424 and 434-592, respectively. Since the object of
20 the present invention is to obtain a polypeptide which exhibits mannanase activity, the present invention relates to any mannanase enzyme comprising the sequence of amino acids nos. 593-903 of SEQ ID NO:32, ie a catalytical domain, optionally operably linked, either N-terminally or C-terminally, to one or
25 two or more than two other domains of a different functionality.

The present invention also provides mannanase polypeptides that are substantially homologous to the polypeptides of SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID
30 NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30 and SEQ ID NO:32, respectively, and species homologs (paralogs or orthologs) thereof. The term "substantially homologous" is used

herein to denote polypeptides having 65%, preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, and even more preferably at least 90%, sequence identity to the sequence shown in amino acids nos. 33-340 or nos. 33-490 of SEQ ID NO:2 or their orthologs or
5 paralog; or to the sequence shown in amino acids nos. 32-344 or nos. 32-494 of SEQ ID NO:6 or their orthologs or paralog; or to the sequence shown in amino acids nos. 32-362 or nos. 32-586 of SEQ ID NO:10 or their orthologs or paralog; or to the sequence
10 shown in amino acids nos. 33-331 of SEQ ID NO:12 or its orthologs or paralog; or to the sequence shown in amino acids nos. 166-488 or nos. 22-488 of SEQ ID NO:14 or their orthologs or paralog; or to the sequence shown in amino acids nos. 68-369 or nos. 32-369 of SEQ ID NO:16 or their orthologs or paralog;
15 or to the sequence shown in amino acids nos. 1-305 of SEQ ID NO:18 or its orthologs or paralog; or to the sequence shown in amino acids nos. 1-132 of SEQ ID NO:20 or its orthologs or paralog; or to the sequence shown in amino acids nos. 29-320 of SEQ ID NO:22 or its orthologs or paralog; or to the sequence
20 shown in amino acids nos. 29-188 of SEQ ID NO:24 or its orthologs or paralog; or to the sequence shown in amino acids nos. 301-625 or nos. 30-625 of SEQ ID NO:26 or their orthologs or paralog; or to the sequence shown in amino acids nos. 166-496 or nos. 38-496 of SEQ ID NO:28 or their orthologs or
25 paralog; or to the sequence shown in amino acids nos. 26-361 of SEQ ID NO:30 or its orthologs or paralog; or to the sequence shown in amino acids nos. 593-903 or nos. 23-903 of SEQ ID NO:32 or their orthologs or paralog.

Such polypeptides will more preferably be at least 95%
30 identical, and most preferably 98% or more identical to the sequence shown in amino acids nos. 31-330 or nos. 31-490 of SEQ ID NO:2 or its orthologs or paralog; or to the sequence shown

in amino acids nos. 32-344 or nos. 32-494 of SEQ ID NO:6 or its orthologs or paralog; or to the sequence shown in amino acids nos. 32-362 or nos. 32-586 of SEQ ID NO:10 or its orthologs or paralog; or to the sequence shown in amino acids nos. 33-331 of
5 SEQ ID NO:12 or its orthologs or paralog; or to the sequence shown in amino acids nos. 166-488 or nos. 22-488 of SEQ ID NO:14 or its orthologs or paralog; or to the sequence shown in amino acids nos. 68-369 or nos. 32-369 of SEQ ID NO:16 or its orthologs or paralog; or to the sequence shown in amino acids
10 nos. 1-305 of SEQ ID NO:18 or its orthologs or paralog; or to the sequence shown in amino acids nos. 1-132 of SEQ ID NO:20 or its orthologs or paralog; or to the sequence shown in amino acids nos. 29-320 of SEQ ID NO:22 or its orthologs or paralog; or to the sequence shown in amino acids nos. 29-188 of SEQ ID
15 NO:24 or its orthologs or paralog; or to the sequence shown in amino acids nos. 301-625 or nos. 30-625 of SEQ ID NO:26 or its orthologs or paralog; or to the sequence shown in amino acids nos. 166-496 or nos. 38-496 of SEQ ID NO:28 or its orthologs or paralog; or to the sequence shown in amino acids nos. 26-361 of
20 SEQ ID NO:30 or its orthologs or paralog; or to the sequence shown in amino acids nos. 593-903 or nos. 23-903 of SEQ ID NO:32 or its orthologs or paralog.

Percent sequence identity is determined by conventional methods, by means of computer programs known in the art such as
25 GAP provided in the GCG program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711) as disclosed in Needleman, S.B. and Wunsch, C.D., (1970), Journal of Molecular Biology, 48, 443-453.

30 GAP is used with the following settings for polypeptide sequence comparison: GAP creation penalty of 3.0 and GAP extension penalty of 0.1.

Sequence identity of polynucleotide molecules is determined by similar methods using GAP with the following settings for DNA sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3.

The enzyme preparation of the invention is preferably derived from a microorganism, preferably from a bacterium, an archaea or a fungus, especially from a bacterium such as a bacterium belonging to *Bacillus*, preferably to a *Bacillus* strain which may be selected from the group consisting of the species *Bacillus* sp. and highly related *Bacillus* species in which all species preferably are at least 95%, even more preferably at least 98%, homologous to *Bacillus* sp. I633, *Bacillus halodurans* or *Bacillus* sp. AAI12 based on aligned 16S rDNA sequences.

5 These species are claimed based on phylogenetic relationships identified from aligned 16S rDNA sequences from RDP (Ribosomal Database Project) (Bonne L. Maidak, Neils Larson, Michael J. McCaughey, Ross Overbeek, Gary J. Olsen, Karl Fogel, James Blandy, and Carl R. Woese, Nucleic Acids Research, 1994, 10 Vol. 22, No17, p. 3485-3487, The Ribosomal Database Project). The alignment was based on secondary structure. Calculation of sequence similarities were established using the "Full matrix calculation" with default settings of the neighbor joining method integrated in the ARB program package (Oliver Strunk and 15 Wolfgang Ludwig, Technical University of Munich, Germany).

Information derived from table II are the basis for the claim for all family 5 mannanases from the highly related *Bacillus* species in which all species over 93% homologous to *Bacillus* sp. I633 are claimed. These include: *Bacillus sporother-* 20 *modurans*, *Bacillus acalophilus*, *Bacillus pseudoalcalophilus* and *Bacillus clausii*. See Figure 1: Phylogenetic tree generated from ARP program relating closest species to *Bacillus* sp. I633. The 16S RNA is shown in SEQ ID NO:33.

Table II: 16S ribosomal RNA homology index for select *Bacillus* species

	BaiSpor2	BaiAlcal	BaiSpec3	BaiSpec5	B.sp.I633
BaiSpor2		92.75%	92.98%	92.41%	93.43%
5 BaiAlcal			98.11%	94.69%	97.03%
BaiSpec3				94.49%	96.39%
BaiSpec5					93.67%

BaiSpor2 = *B. sporothermodurans*, u49079

BaiAlcal = *B. B. alcalophilus*, x76436

10 BaiSpec3 = *B. pseudoalcalophilus*, x76449

BaiSpec5 = *B. clausii*, x76440

Other useful family 5 mannanases are those derived from the highly related *Bacillus* species in which all species show more than 93% homology to *Bacillus halodurans* based on aligned
 15 16S sequences. These *Bacillus* species include: *Sporolactobacillus laevis*, *Bacillus agaradhaerens* and *Marinococcus halophilus*. See Figure 2: Phylogenic tree generated from ARP program relating closest species to *Bacillus halodurans*.

Table III: 16S ribosomal RNA homology index for selected
 20 *Bacillus* species

	SplLaev3	BaiSpec6	BaiSpe11	MaoHalo2	NN
SplLaev3		90.98%	87.96%	85.94%	91.32%
BaiSpec6			91.63%	87.96%	99.46%
BaiSpe11				89.04%	92.04%
25 MaoHalo2					88.17%

NN

SplLaev3 = *Sporolactobacillus laevis*, D16287

BaiSpec6 = *B. halodurans*, X76442

BaiSpe11 = *B. agaradhaerens*, X76445

30 MaoHalo2 = *Marinococcus halophilus*, X62171

NN = donor organism of the invention (*B. halodurans*)

Other useful family 5 mannanases are those derived from a strain selected from the group consisting of the species *Bacillus agaradhaerens* and highly related *Bacillus* species in which all species preferably are at least 95%, even more preferably at least 98%, homologous to *Bacillus agaradhaerens*, DSM 8721, based on aligned 16S rDNA sequences.

Useful family 26 mannanases are for example those derived from the highly related *Bacillus* species in which all species over 93% homologous to *Bacillus* sp. AAI12 are claimed. These include: *Bacillus sporothermodurans*, *Bacillus alcalophilus*, *Bacillus pseudoalcalophilus* and *Bacillus clausii*. See Figure 3: Phylogenetic tree generated from ARP program relating closest species to *Bacillus* sp. AAI 12. The 16S RNA is shown in SEQ ID NO:34.

Table IV: 16S ribosomal RNA homology index for selected *Bacillus* species

	BaiSpor2	BaiAlcal	BaiSpec3	BaiSpec5	B.sp.AAI12
BaiSpor2		92.75%	92.98%	92.41%	92.24%
BaiAlcal			98.11%	94.69%	97.28%
BaiSpec3				94.49%	96.10%
BaiSpec5					93.83%

BaiSpor2 = *B. sporothermodurans*, u49079

BaiAlcal = *B. B. alcalophilus*, x76436

BaiSpec3 = *B. pseudoalcalophilus*, x76449

BaiSpec5 = *B. clausii*, x76440

Other useful family 26 mannanases are those derived from a strain selected from the group consisting of the species *Bacillus licheniformis* and highly related *Bacillus* species in which all species preferably are at least 95%, even more preferably at least 98%, homologous to *Bacillus licheniformis* based on aligned 16S rDNA sequences.

Substantially homologous proteins and polypeptides are characterized as having one or more amino acid substitutions, deletions or additions. These changes are preferably of a minor nature, that is conservative amino acid substitutions (see Table 5 2) and other substitutions that do not significantly affect the folding or activity of the protein or polypeptide; small deletions, typically of one to about 30 amino acids; and small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue, a small linker peptide of up to 10 about 20-25 residues, or a small extension that facilitates purification (an affinity tag), such as a poly-histidine tract, protein A (Nilsson et al., EMBO J. 4:1075, 1985; Nilsson et al., Methods Enzymol. 198:3, 1991. See, in general Ford et al., Protein Expression and Purification 2: 95-107, 1991.

15 DNAs encoding affinity tags are available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, NJ; New England Biolabs, Beverly, MA).

However, even though the changes described above preferably are of a minor nature, such changes may also be of a larger 20 nature such as fusion of larger polypeptides of up to 300 amino acids or more both as amino- or carboxyl-terminal extensions to a Mannanase polypeptide of the invention.

Table 1

25 Conservative amino acid substitutions

	Basic:	arginine
		lysine
		histidine
30	Acidic:	glutamic acid
		aspartic acid
	Polar:	glutamine

		asparagine
	Hydrophobic:	leucine
		isoleucine
		valine
5	Aromatic:	phenylalanine
		tryptophan
		tyrosine
	Small:	glycine
		alanine
10		serine
		threonine
		methionine

In addition to the 20 standard amino acids, non-standard amino acids (such as 4-hydroxyproline, 6-N-methyl lysine, 2-aminoisobutyric acid, isovaline and α -methyl serine) may be substituted for amino acid residues of a polypeptide according to the invention. A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic code, and unnatural amino acids may be substituted for amino acid residues. "Unnatural amino acids" have been modified after protein synthesis, and/or have a chemical structure in their side chain(s) different from that of the standard amino acids. Unnatural amino acids can be chemically synthesized, or preferably, are commercially available, and include pipecolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, and 3,3-dimethylproline.

Essential amino acids in the mannanase polypeptides of the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, Science 244: 1081-1085, 1989). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resul-

tant mutant molecules are tested for biological activity (i.e. mannanase activity) to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., J. Biol. Chem. 271:4699-4708, 1996. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., Science 255:306-312, 1992; Smith et al., J. Mol. Biol. 224:899-904, 1992; Wlodaver et al., FEBS Lett. 309:59-64, 1992.

The identities of essential amino acids can also be inferred from analysis of homologies with polypeptides which are related to a polypeptide according to the invention.

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis, recombination and/or shuffling followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer (Science 241:53-57, 1988), Bowie and Sauer (Proc. Natl. Acad. Sci. USA 86:2152-2156, 1989), WO95/17413, or WO 95/22625. Briefly, these authors disclose methods for simultaneously randomizing two or more positions in a polypeptide, or recombination/shuffling of different mutations (WO95/17413, WO95/22625), followed by selecting for functional a polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman et al., Biochem. 30:10832-10837, 1991; Ladner et al., U.S. Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204) and region-directed mutagenesis (Derbyshire et al., Gene 46:145, 1986; Ner et al., DNA 7:127, 1988).

Mutagenesis/shuffling methods as disclosed above can be combined with high-throughput, automated screening methods to

detect activity of cloned, mutagenized polypeptides in host cells. Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using modern equipment. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

Using the methods discussed above, one of ordinary skill in the art can identify and/or prepare a variety of polypeptides that are substantially homologous to residues 33-340 or 33-490 of SEQ ID NO:2; or to residues 32-344 or 32-494 of SEQ ID NO:6; or to residues 32-362 or 32-586 of SEQ ID NO:10; or to residues 33-331 of SEQ ID NO:12; or to residues 166-488 or 22-488 of SEQ ID NO:14; or to residues 68-369 or 32-369 of SEQ ID NO:16; or to residues 1-305 of SEQ ID NO:18; or to residues 1-132 of SEQ ID NO:20; or to residues 29-320 of SEQ ID NO:22; or to residues 29-188 of SEQ ID NO:24; or to residues 301-625 or 30-625 of SEQ ID NO:26; or to residues 166-496 or 38-496 of SEQ ID NO:28; or to residues 26-361 of SEQ ID NO:30; or to residues 593-903 or 23-903 of SEQ ID NO:32 and retain the mannanase activity of the wild-type protein.

The mannanase enzyme of the invention may, in addition to the enzyme core comprising the catalytically domain, also comprise a cellulose binding domain (CBD), the cellulose binding domain and enzyme core (the catalytically active domain) of the enzyme being operably linked. The cellulose binding domain (CBD) may exist as an integral part of the encoded enzyme, or a CBD from another origin may be introduced into the mannan degrading enzyme thus creating an enzyme hybrid. In this context, the term "cellulose-binding domain" is intended to be understood as defined by Peter Tomme et al. "Cellulose-Binding Domains: Classification and Properties" in "Enzymatic Degradation

of Insoluble Carbohydrates", John N. Saddler and Michael H. Penner (Eds.), ACS Symposium Series, No. 618, 1996. This definition classifies more than 120 cellulose-binding domains into 10 families (I-X), and demonstrates that CBDs are found in various enzymes such as cellulases, xylanases, mannanases, arabinofuranosidases, acetyl esterases and chitinases. CBDs have also been found in algae, e.g. the red alga *Porphyra purpurea* as a non-hydrolytic polysaccharide-binding protein, see Tomme et al., *op.cit.* However, most of the CBDs are from cellulases and xylanases, CBDs are found at the N and C termini of proteins or are internal. Enzyme hybrids are known in the art, see e.g. WO 90/00609 and WO 95/16782, and may be prepared by transforming into a host cell a DNA construct comprising at least a fragment of DNA encoding the cellulose-binding domain ligated, with or without a linker, to a DNA sequence encoding the mannan degrading enzyme and growing the host cell to express the fused gene. Enzyme hybrids may be described by the following formula:



wherein CBD is the N-terminal or the C-terminal region of an amino acid sequence corresponding to at least the cellulose-binding domain; MR is the middle region (the linker), and may be
5 a bond, or a short linking group preferably of from about 2 to about 100 carbon atoms, more preferably of from 2 to 40 carbon atoms; or is preferably from about 2 to to about 100 amino acids, more preferably of from 2 to 40 amino acids; and X is an N-terminal or C-terminal region of the mannanase of the
10 invention. SEQ ID NO:4 discloses the amino acid sequence of an enzyme hybrid of a mannanase enzyme core and a CBD.

Preferably, the mannanase enzyme of the present invention has its maximum catalytic activity at a pH of at least 7, more preferably of at least 8, more preferably of at least 8.5, more

preferably of at least 9, more preferably of at least 9.5, more preferably of at least 10, even more preferably of at least 10.5, especially of at least 11; and preferably the maximum activity of the enzyme is obtained at a temperature of at least 40°C, more preferably of at least 50°C, even more preferably of at least 55°C.

Preferably, the cleaning composition of the present invention provides, eg when used for treating fabric during a washing cycle of a machine washing process, a washing solution having a pH typically between about 8 and about 10.5. Typically, such a washing solution is used at temperatures between about 20°C and about 95°C, preferably between about 20°C and about 60°C, preferably between about 20°C and about 50°C.

15 PROTEIN PRODUCTION:

The proteins and polypeptides of the present invention, including full-length proteins, fragments thereof and fusion proteins, can be produced in genetically engineered host cells according to conventional techniques. Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA and grown in culture, and include bacteria, fungal cells, and cultured higher eukaryotic cells. Bacterial cells, particularly cultured cells of gram-positive organisms, are preferred. Gram-positive cells from the genus of *Bacillus* are especially preferred, such as from the group consisting of *Bacillus subtilis*, *Bacillus lentus*, *Bacillus clausii*, *Bacillus agaradhaerens*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus*, *Bacillus thuringiensis*, *Bacillus licheniformis*, and *Bacillus sp.*, in particular *Bacillus sp.* I633, *Bacillus sp.* AAI12, *Bacillus*

clausii, *Bacillus agaradhaerens* and *Bacillus licheniformis*.

In another preferred embodiment, the host cell is a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota (as defined by
5 Hawksworth et al., In, Ainsworth and Bisby's Dictionary of The Fungi, 8th edition, 1995, CAB International, University Press, Cambridge, UK) as well as the Oomycota (as cited in Hawksworth et al., 1995, supra, page 171) and all mitosporic fungi (Hawksworth et al., 1995, supra). Representative groups of Ascomycota
10 include, e.g., *Neurospora*, *Eupenicillium* (= *Penicillium*), *Emmericella* (= *Aspergillus*), *Eurotium* (= *Aspergillus*), and the true yeasts listed above. Examples of Basidiomycota include mushrooms, rusts, and smuts. Representative groups of Chytridiomycota include, e.g., *Allomyces*, *Blastocladiella*,
15 *Coelomomyces*, and aquatic fungi. Representative groups of Oomycota include, e.g., Saprolegniomycetous aquatic fungi (water molds) such as *Achlya*. Examples of mitosporic fungi include *Aspergillus*, *Penicillium*, *Candida*, and *Alternaria*. Representative groups of Zygomycota include, e.g., *Rhizopus* and
20 *Mucor*.

In yet another preferred embodiment, the fungal host cell is a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth et al., 1995, supra). In a more preferred
25 embodiment, the filamentous fungal host cell is a cell of a species of, but not limited to, *Acremonium*, *Aspergillus*, *Fusarium*, *Humicola*, *Mucor*, *Myceliophthora*, *Neurospora*, *Penicillium*, *Thielavia*, *Tolypocladium*, and *Trichoderma* or a teleomorph or synonym thereof.

30 In particular, the cell may belong to a species of *Trichoderma*, preferably *Trichoderma harzianum* or *Trichoderma reesei*, or a species of *Aspergillus*, most preferably *Aspergillus*

oryzae or *Aspergillus niger*, or a species of *Fusarium*, most preferably a *Fusarium* sp. having the identifying characteristic of *Fusarium* ATCC 20334, as further described in PCT/US/95/07743.

Fungal cells may be transformed by a process involving
5 protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* host cells are described in EP 238 023 and Yelton et al., 1984, *Proceedings of the National Academy of Sciences USA* 81:1470-1474. A suitable
10 method of transforming *Fusarium* species is described by Malardier et al., 1989, *Gene* 78:147-156 or in copending U.S. Patent No. 5,837,847. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology*,
15 *Methods in Enzymology*, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito et al., 1983, *Journal of Bacteriology* 153:163; and Hinnen et al., 1978, *Proceedings of the National Academy of Sciences USA* 75:1920. Mammalian cells may be transformed by direct uptake using the calcium phosphate
20 precipitation method of Graham and Van der Eb (1978, *Virology* 52:546).

Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd
25 ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989; Ausubel et al. (eds.), Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY, 1987; and "Bacillus subtilis and Other Gram-Positive Bacteria", Sonensheim et al., 1993, American Society for Microbiology, Washington D.C.

30 In general, a DNA sequence encoding a mannanase of the present invention is operably linked to other genetic elements

required for its expression, generally including a transcription promoter and terminator within an expression vector. The vector will also commonly contain one or more selectable markers and one or more origins of replication, although those skilled in the art will recognize that within certain systems selectable markers may be provided on separate vectors, and replication of the exogenous DNA may be provided by integration into the host cell genome. Selection of promoters, terminators; selectable markers, vectors and other elements is a matter of routine design within the level of ordinary skill in the art. Many such elements are described in the literature and are available through commercial suppliers.

To direct a polypeptide into the secretory pathway of a host cell, a secretory signal sequence (also known as a leader sequence, prepro sequence or pre sequence) is provided in the expression vector. The secretory signal sequence may be that of the polypeptide, or may be derived from another secreted protein or synthesized *de novo*. Numerous suitable secretory signal sequences are known in the art and reference is made to "Bacillus subtilis and Other Gram-Positive Bacteria", Sonensheim et al., 1993, American Society for Microbiology, Washington D.C.; and Cutting, S. M. (eds.) "Molecular Biological Methods for Bacillus", John Wiley and Sons, 1990, for further description of suitable secretory signal sequences especially for secretion in a Bacillus host cell. The secretory signal sequence is joined to the DNA sequence in the correct reading frame. Secretory signal sequences are commonly positioned 5' to the DNA sequence encoding the polypeptide of interest, although certain signal sequences may be positioned elsewhere in the DNA sequence of interest (see, e.g., Welch et al., U.S. Patent No. 5,037,743; Holland et al., U.S. Patent No. 5,143,830).

The expression vector of the invention may be any expression vector that is conveniently subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which the vector it is to be introduced.

5 Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell
10 genome and replicated together with the chromosome(s) into which it has been integrated.

Examples of suitable promoters for use in filamentous fungus host cells are, e.g. the ADH3 promoter (McKnight et al., The EMBO J. 4 (1985), 2093 - 2099) or the tpiA promoter.

15 Examples of other useful promoters are those derived from the gene encoding *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral α -amylase, *Aspergillus niger* acid stable α -amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (gluA), *Rhizomucor miehei*
20 lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase or *Aspergillus nidulans* acetamidase.

Transformed or transfected host cells are cultured according to conventional procedures in a culture medium containing nutrients and other components required for the growth of the chosen
25 host cells. A variety of suitable media, including defined media and complex media, are known in the art and generally include a carbon source, a nitrogen source, essential amino acids, vitamins and minerals. Media may also contain such components as growth factors or serum, as required. The growth medium will
30 generally select for cells containing the exogenously added DNA by, for example, drug selection or deficiency in an essential

nutrient which is complemented by the selectable marker carried on the expression vector or co-transfected into the host cell.

PROTEIN ISOLATION

5 When the expressed recombinant polypeptide is secreted the polypeptide may be purified from the growth media. Preferably the expression host cells are removed from the media before purification of the polypeptide (e.g. by centrifugation).

 When the expressed recombinant polypeptide is not secreted
10 from the host cell, the host cell are preferably disrupted and the polypeptide released into an aqueous "extract" which is the first stage of such purification techniques. Preferably the expression host cells are collected from the media before the cell disruption (e.g. by centrifugation).

15 The cell disruption may be performed by conventional techniques such as by lysozyme digestion or by forcing the cells through high pressure. See (Robert K. Scobes, Protein Purification, Second edition, Springer-Verlag) for further description of such cell disruption techniques.

20 Whether or not the expressed recombinant polypeptides (or chimeric polypeptides) is secreted or not it can be purified using fractionation and/or conventional purification methods and media.

 Ammonium sulfate precipitation and acid or chaotrope extrac-
25 tion may be used for fractionation of samples. Exemplary purification steps may include hydroxyapatite, size exclusion, FPLC and reverse-phase high performance liquid chromatography. Suitable anion exchange media include derivatized dextrans, agarose, cellulose, polyacrylamide, specialty silicas, and the like. PEI,
30 DEAE, QAE and Q derivatives are preferred, with DEAE Fast-Flow Sepharose* (Pharmacia, Piscataway, NJ) being particularly preferred. Exemplary chromatographic media include those media

*Trade-mark

derivatized with phenyl, butyl, or octyl groups, such as Phenyl-Sepharose FF (Pharmacia), Toyopearl* butyl 650 (Toso Haas, Montgomeryville, PA), Octyl-Sepharose (Pharmacia) and the like; or polyacrylic resins, such as Amberchrom CG 71 (Toso Haas) and the like. Suitable solid supports include glass beads, silica-based resins, cellulosic resins, agarose beads, cross-linked agarose beads, polystyrene beads, cross-linked polyacrylamide resins and the like that are insoluble under the conditions in which they are to be used. These supports may be modified with reactive groups that allow attachment of proteins by amino groups, carboxyl groups, sulfhydryl groups, hydroxyl groups and/or carbohydrate moieties. Examples of coupling chemistries include cyanogen bromide activation, N-hydroxysuccinimide activation, epoxide activation, sulfhydryl activation, hydrazide activation, and carboxyl and amino derivatives for carbodiimide coupling chemistries. These and other solid media are well known and widely used in the art, and are available from commercial suppliers.

Selection of a particular method is a matter of routine design and is determined in part by the properties of the chosen support. See, for example, Affinity Chromatography: Principles & Methods, Pharmacia LKB Biotechnology, Uppsala, Sweden, 1988.

Polypeptides of the invention or fragments thereof may also be prepared through chemical synthesis. Polypeptides of the invention may be monomers or multimers; glycosylated or non-glycosylated; pegylated or non-pegylated; and may or may not include an initial methionine amino acid residue.

Based on the sequence information disclosed herein a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 1, at least the DNA sequence from position 94 to position 990, or, alternatively, the DNA sequence from position 94 to position 1470, may

be cloned. Likewise may be cloned a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 5, at least the DNA sequence from position 94 to position 1032, or, alternatively, the DNA sequence from position 94 to position 1482; and a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 9, at least the DNA sequence from position 94 to position 1086, or, alternatively, the DNA sequence from position 94 to position 1761; and a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 11, at least the DNA sequence from position 97 to position 993; and a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 13, at least the DNA sequence from position 498 to position 1464, or, alternatively, the DNA sequence from position 64 to position 1464; and a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 15, at least the DNA sequence from position 204 to position 1107, or, alternatively, the DNA sequence from position 76 to position 1107; and a DNA sequence partially encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 17; and a DNA sequence partially encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 19; and a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 21, at least the DNA sequence from position 88 to position 960; and a DNA sequence partially encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 23; and a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 25, at least the DNA sequence from position 904 to position 1875, or,

alternatively, the DNA sequence from position 88 to position 2445; and a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 27, at least the DNA sequence from position 498 to position 1488, or, alternatively, the DNA sequence from position 112 to position 1488; and a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 29, at least the DNA sequence from position 79 to position 1083; and a full length DNA sequence encoding a mannanase of the invention and comprising the DNA sequence shown in SEQ ID No 31, at least the DNA sequence from position 1779 to position 2709, or, alternatively, the DNA sequence from position 67 to position 2709.

Cloning is performed by standard procedures known in the art such as by,

- preparing a genomic library from a *Bacillus* strain, especially a strain selected from *B. sp. I633*, *B. sp. AAI12*, *B. sp. AA349*, *Bacillus agaradhaerens*, *Bacillus halodurans*, *Bacillus clausii* and *Bacillus licheniformis*, or from a fungal strain, especially the strain *Humicola insolens*;
- plating such a library on suitable substrate plates;
- identifying a clone comprising a polynucleotide sequence of the invention by standard hybridization techniques using a probe based on any of the sequences SEQ ID Nos. 1, 5, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31; or by
- identifying a clone from said genomic library by an Inverse PCR strategy using primers based on sequence information from SEQ ID No 1, 5, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31. Reference is made to M.J. MCPerson et al. ("PCR A practical approach" Information Press Ltd, Oxford England) for further details relating to Inverse PCR.

Based on the sequence information disclosed herein (SEQ ID Nos 1, 2, 5, 6, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32) is it routine work for a person skilled in the art to isolate homologous polynucleotide sequences encoding homologous mannanase of the invention by a similar strategy using genomic libraries from related microbial organisms, in particular from genomic libraries from other strains of the genus *Bacillus* such as alkalophilic species of *Bacillus* sp., or from fungal strains such as species of *Humicola*.

Alternatively, the DNA encoding the mannan or galactomannan-degrading enzyme of the invention may, in accordance with well-known procedures, conveniently be cloned from a suitable source, such as any of the above mentioned organisms, by use of synthetic oligonucleotide probes prepared on the basis of the DNA sequence obtainable from the plasmid present any of the strains *Escherichia coli* DSM 12197, DSM 12180, DSM 12433, DSM 12441, DSM 9984, DSM 12432, DSM 12436, DSM 12846, DSM 12847, DSM 12848, DSM 12849, DSM 12850, DSM 12851 and DSM 12852.

Accordingly, the polynucleotide molecule of the invention may be isolated from any of *Escherichia coli*, DSM 12197, DSM 12180, DSM 12433, DSM 12441, DSM 9984, DSM 12432, DSM 12436, DSM 12846, DSM 12847, DSM 12848, DSM 12849, DSM 12850, DSM 12851 and DSM 12852, in which the plasmid obtained by cloning such as described above is deposited. Also, the present invention relates to an isolated substantially pure biological culture of any of the strains *Escherichia coli*, DSM 12197, DSM 12180, DSM 12433, DSM 12441, DSM 9984, DSM 12432, DSM 12436, DSM 12846, DSM 12847, DSM 12848, DSM 12849, DSM 12850, DSM 12851 and DSM 12852.

In the present context, the term "enzyme preparation" is intended to mean either a conventional enzymatic fermentation

product, possibly isolated and purified, from a single species of a microorganism, such preparation usually comprising a number of different enzymatic activities; or a mixture of monocomponent enzymes, preferably enzymes derived from bacterial or fungal species by using conventional recombinant techniques, which enzymes have been fermented and possibly isolated and purified separately and which may originate from different species, preferably fungal or bacterial species; or the fermentation product of a microorganism which acts as a host cell for expression of a recombinant mannanase, but which microorganism simultaneously produces other enzymes, e.g. pectin degrading enzymes, proteases, or cellulases, being naturally occurring fermentation products of the microorganism, i.e. the enzyme complex conventionally produced by the corresponding naturally occurring microorganism.

The mannanase preparation of the invention may further comprise one or more enzymes selected from the group consisting of proteases, cellulases (endo- β -1,4-glucanases), β -glucanases (endo- β -1,3(4)-glucanases), lipases, cutinases, peroxidases, laccases, amylases, glucoamylases, pectinases, reductases, oxidases, phenoloxidases, ligninases, pullulanases, hemicellulases, pectate lyases, xyloglucanases, xylanases, pectin acetyl esterases, rhamnogalacturonan acetyl esterases, polygalacturonases, rhamnogalacturonases, pectin lyases, pectin methylesterases, cellobiohydrolases, transglutaminases; or mixtures thereof. In a preferred embodiment, one or more or all enzymes in the preparation is produced by using recombinant techniques, i.e. the enzyme(s) is/are mono-component enzyme(s) which is/are mixed with the other enzyme(s) to form an enzyme preparation with the desired enzyme blend.

In another aspect, the present invention also relates to a method of producing the enzyme preparation of the invention, the

method comprising culturing a microorganism, eg a wild-type strain, capable of producing the mannanase under conditions permitting the production of the enzyme, and recovering the enzyme from the culture. Culturing may be carried out using
5 conventional fermentation techniques, e.g. culturing in shake flasks or fermentors with agitation to ensure sufficient aeration on a growth medium inducing production of the mannanase enzyme. The growth medium may contain a conventional N-source such as peptone, yeast extract or casamino acids, a reduced
10 amount of a conventional C-source such as dextrose or sucrose, and an inducer such as guar gum or locust bean gum. The recovery may be carried out using conventional techniques, e.g. separation of bio-mass and supernatant by centrifugation or filtration, recovery of the supernatant or disruption of cells
15 if the enzyme of interest is intracellular, perhaps followed by further purification as described in EP 0 406 314 or by crystallization as described in WO 97/15660.

Examples of useful bacteria producing the enzyme or the enzyme preparation of the invention are Gram positive bacteria,
20 preferably from the *Bacillus/Lactobacillus* subdivision, preferably a strain from the genus *Bacillus*, more preferably a strain of *Bacillus sp.*

In yet another aspect, the present invention relates to an isolated mannanase having the properties described above and
25 which is free from homologous impurities, and is produced using conventional recombinant techniques.

IMMUNOLOGICAL CROSS-REACTIVITY

30 Polyclonal antibodies to be used in determining immunological cross-reactivity may be prepared by use of a purified mannanase enzyme. More specifically, antiserum against

the mannanase of the invention may be raised by immunizing rabbits (or other rodents) according to the procedure described by N. Axelsen et al. in: A Manual of Quantitative Immuno-electrophoresis, Blackwell Scientific Publications, 1973, Chapter 23, or A. Johnstone and R. Thorpe, Immunochemistry in Practice, Blackwell Scientific Publications, 1982 (more specifically p. 27-31). Purified immunoglobulins may be obtained from the antisera, for example by salt precipitation $((\text{NH}_4)_2 \text{SO}_4)$, followed by dialysis and ion exchange chromatography, e.g. on DEAE-Sephadex*. Immunochemical characterization of proteins may be done either by Ouchterlony double-diffusion analysis (O. Ouchterlony in: Handbook of Experimental Immunology (D.M. Weir, Ed.), Blackwell Scientific Publications, 1967, pp. 655-706), by crossed immuno-electrophoresis (N. Axelsen et al., supra, Chapters 3 and 4), or by rocket immuno-electrophoresis (N. Axelsen et al., Chapter 2).

Use in the detergent industry

In further aspects, the present invention relates to a detergent composition comprising the mannanase or mannanase preparation of the invention, to a process for machine treatment of fabrics comprising treating fabric during a washing cycle of a machine washing process with a washing solution containing the mannanase or mannanase preparation of the invention, and to cleaning compositions, including laundry, dishwashing, hard surface cleaner, personal cleansing and oral/dental compositions, comprising a mannanase and optionally another enzyme selected among cellulases, amylases, pectin degrading enzymes and xyloglucanases and providing superior cleaning performance, i.e. superior stain removal, dingy cleaning and whiteness maintenance.

*Trade-mark

Without being bound to this theory, it is believed that the mannanase of the present invention is capable of effectively degrading or hydrolysing any soiling or spots containing galactomannans and, accordingly, of cleaning laundry comprising
5 such soilings or spots.

The cleaning compositions of the invention must contain at least one additional detergent component. The precise nature of these additional components, and levels of incorporation thereof will depend on the physical form of the composition, and the
10 nature of the cleaning operation for which it is to be used.

The cleaning compositions of the present invention preferably further comprise a detergent ingredient selected from a selected surfactant, another enzyme, a builder and/or a bleach system.

15 The cleaning compositions according to the invention can be liquid, paste, gels, bars, tablets, spray, foam, powder or granular. Granular compositions can also be in "compact" form and the liquid compositions can also be in a "concentrated" form.

20 The compositions of the invention may for example, be formulated as hand and machine dishwashing compositions, hand and machine laundry detergent compositions including laundry additive compositions and compositions suitable for use in the soaking and/or pretreatment of stained fabrics, rinse added
25 fabric softener compositions, and compositions for use in general household hard surface cleaning operations. Compositions containing such carbohydrases can also be formulated as sanitization products, contact lens cleansers and health and beauty care products such as oral/dental care and personal cleaning
30 compositions.

When formulated as compositions for use in manual dishwashing methods the compositions of the invention preferably contain

a surfactant and preferably other detergent compounds selected from organic polymeric compounds, suds enhancing agents, group II metal ions, solvents, hydrotropes and additional enzymes.

When formulated as compositions suitable for use in a laundry machine washing method, the compositions of the invention preferably contain both a surfactant and a builder compound and additionally one or more detergent components preferably selected from organic polymeric compounds, bleaching agents, additional enzymes, suds suppressors, dispersants, lime-soap dispersants, soil suspension and anti-redeposition agents and corrosion inhibitors. Laundry compositions can also contain softening agents, as additional detergent components. Such compositions containing carbohydrase can provide fabric cleaning, stain removal, whiteness maintenance, softening, colour appearance, dye transfer inhibition and sanitization when formulated as laundry detergent compositions.

The compositions of the invention can also be used as detergent additive products in solid or liquid form. Such additive products are intended to supplement or boost the performance of conventional detergent compositions and can be added at any stage of the cleaning process.

If needed the density of the laundry detergent compositions herein ranges from 400 to 1200 g/litre, preferably 500 to 950 g/litre of composition measured at 20°C.

The "compact" form of the compositions herein is best reflected by density and, in terms of composition, by the amount of inorganic filler salt; inorganic filler salts are conventional ingredients of detergent compositions in powder form; in conventional detergent compositions, the filler salts are present in substantial amounts, typically 17-35% by weight of the total composition. In the compact compositions, the filler salt is present in amounts not exceeding 15% of the total composi-

tion, preferably not exceeding 10%, most preferably not exceeding 5% by weight of the composition. The inorganic filler salts, such as meant in the present compositions are selected from the alkali and alkaline-earth-metal salts of sulphates and chlorides. A preferred filler salt is sodium sulphate.

Liquid detergent compositions according to the present invention can also be in a "concentrated form", in such case, the liquid detergent compositions according the present invention will contain a lower amount of water, compared to conventional liquid detergents. Typically the water content of the concentrated liquid detergent is preferably less than 40%, more preferably less than 30%, most preferably less than 20% by weight of the detergent composition.

15 Cleaning compositions

Surfactant system

The cleaning or detergent compositions according to the present invention comprise a surfactant system, wherein the surfactant can be selected from nonionic and/or anionic and/or cationic and/or ampholytic and/or zwitterionic and/or semi-polar surfactants.

The surfactant is typically present at a level from 0.1% to 60% by weight. The surfactant is preferably formulated to be compatible with enzyme hybrid and enzyme components present in the composition. In liquid or gel compositions the surfactant is most preferably formulated in such a way that it promotes, or at least does not degrade, the stability of any enzyme hybrid or enzyme in these compositions.

Suitable systems for use according to the present invention comprise as a surfactant one or more of the nonionic and/or anionic surfactants described herein.

Polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols are suitable for use as the nonionic surfactant of the surfactant systems of the present invention, with the polyethylene oxide condensates being preferred. These compounds include the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 14 carbon atoms, preferably from about 8 to about 14 carbon atoms, in either a straight chain or branched-chain configuration with the alkylene oxide. In a preferred embodiment, the ethylene oxide is present in an amount equal to from about 2 to about 25 moles, more preferably from about 3 to about 15 moles, of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include IgepalTM CO-630, marketed by the GAF Corporation; and TritonTM X-45, X-114, X-100 and X-102, all marketed by the Rohm & Haas Company. These surfactants are commonly referred to as alkylphenol alkoxylates (e.g., alkyl phenol ethoxylates).


The condensation products of primary and secondary aliphatic alcohols with about 1 to about 25 moles of ethylene oxide are suitable for use as the nonionic surfactant of the nonionic surfactant systems of the present invention. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Preferred are the condensation products of alcohols having an alkyl group containing from about 8 to about 20 carbon atoms, more preferably from about 10 to about 18 carbon atoms, with from about 2 to about 10 moles of ethylene oxide per mole of alcohol. About 2 to about 7 moles of ethylene oxide and most preferably from 2 to 5 moles of ethylene oxide per mole of alcohol are present in said condensation products. Examples of commercially available nonionic surfactants of this type include TergitolTM 15-S-9 (The condensation product

of C₁₁-C₁₅ linear alcohol with 9 moles ethylene oxide), TergitolTM 24-L-6 NMW (the condensation product of C₁₂-C₁₄ primary alcohol with 6 moles ethylene oxide with a narrow molecular weight distribution), both marketed by Union Carbide Corporation;
5 NeodolTM 45-9 (the condensation product of C₁₄-C₁₅ linear alcohol with 9 moles of ethylene oxide), NeodolTM 23-3 (the condensation product of C₁₂-C₁₃ linear alcohol with 3.0 moles of ethylene oxide), NeodolTM 45-7 (the condensation product of C₁₄-C₁₅ linear alcohol with 7 moles of ethylene oxide), NeodolTM 45-5 (the
10 condensation product of C₁₄-C₁₅ linear alcohol with 5 moles of ethylene oxide) marketed by Shell Chemical Company, KyroTM EOB (the condensation product of C₁₃-C₁₅ alcohol with 9 moles ethylene oxide), marketed by The Procter & Gamble Company, and Genapol^{*} LA 050 (the condensation product of C₁₂-C₁₄ alcohol with
15 5 moles of ethylene oxide) marketed by Hoechst. Preferred range of HLB in these products is from 8-11 and most preferred from 8-10.

Also useful as the nonionic surfactant of the surfactant systems of the present invention are alkylpolysaccharides
20 disclosed in US 4,565,647, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g. a polyglycoside, hydrophilic group containing from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably
25 from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties (optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose
30 or galactose as opposed to a glucoside or galactoside). The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6-

positions on the preceding saccharide units.

The preferred alkylpolyglycosides have the formula




wherein R^2 is selected from the group consisting of alkyl,
5 alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures
thereof in which the alkyl groups contain from about 10 to about
18, preferably from about 12 to about 14, carbon atoms; n is 2
or 3, preferably 2; t is from 0 to about 10, preferably 0; and
10 x is from about 1.3 to about 10, preferably from about 1.3 to
about 3, most preferably from about 1.3 to about 2.7. The
glycosyl is preferably derived from glucose. To prepare these
compounds, the alcohol or alkylpolyethoxy alcohol is formed
first and then reacted with glucose, or a source of glucose, to
form the glucoside (attachment at the 1-position). The
15 additional glycosyl units can then be attached between their 1-
position and the preceding glycosyl units 2-, 3-, 4-, and/or 6-
position, preferably predominantly the 2-position.

The condensation products of ethylene oxide with a
hydrophobic base formed by the condensation of propylene oxide
20 with propylene glycol are also suitable for use as the
additional nonionic surfactant systems of the present invention.
The hydrophobic portion of these compounds will preferably have
a molecular weight from about 1500 to about 1800 and will
exhibit water insolubility. The addition of polyoxyethylene
25 moieties to this hydrophobic portion tends to increase the water
solubility of the molecule as a whole, and the liquid character
of the product is retained up to the point where the
polyoxyethylene content is about 50% of the total weight of the
condensation product, which corresponds to condensation with up
30 to about 40 moles of ethylene oxide. Examples of compounds of
this type include certain of the commercially available
PluronicTM surfactants, marketed by BASF.

Also suitable for use as the nonionic surfactant of the nonionic surfactant system of the present invention, are the condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and
 5 ethylenediamine. The hydrophobic moiety of these products consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from about 2500 to about 3000. This hydrophobic moiety is condensed with ethylene oxide to the extent that the condensation product
 10 contains from about 40% to about 80% by weight of polyoxyethylene and has a molecular weight of from about 5,000 to about 11,000. Examples of this type of nonionic surfactant include certain of the commercially available Tetronic™ compounds, marketed by BASF.

15 Preferred for use as the nonionic surfactant of the surfactant systems of the present invention are polyethylene oxide condensates of alkyl phenols, condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethyleneoxide, alkylpolysaccharides, and
 20 mixtures hereof. Most preferred are C₈-C₁₄ alkyl phenol ethoxylates having from 3 to 15 ethoxy groups and C₈-C₁₈ alcohol ethoxylates (preferably C₁₀ avg.) having from 2 to 10 ethoxy groups, and mixtures thereof.

Highly preferred nonionic surfactants are polyhydroxy
 25 fatty acid amide surfactants of the formula



30 wherein R¹ is H, or R¹ is C₁₋₄ hydrocarbyl, 2-hydroxyethyl, 2-hydroxypropyl or a mixture thereof, R² is C₅₋₃₁ hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain


with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative thereof. Preferably, R^1 is methyl, R^2 is straight C_{11-15} alkyl or C_{16-18} alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z is derived from a
5 reducing sugar such as glucose, fructose, maltose or lactose, in a reductive amination reaction.

Highly preferred anionic surfactants include alkyl alkoxyated sulfate surfactants. Examples hereof are water soluble salts or acids of the formula $RO(A)_mSO_3M$ wherein R is an
10 unsubstituted $C_{10}-C_{24}$ alkyl or hydroxyalkyl group having a $C_{10}-C_{24}$ alkyl component, preferably a $C_{12}-C_{20}$ alkyl or hydro-xyalkyl, more preferably $C_{12}-C_{18}$ alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3,
15 and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxyated sulfates as well as alkyl propoxyated sulfates are contemplated herein. Specific examples of substituted ammonium cations
20 include methyl-, dimethyl, trimethyl-ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like. Exemplary surfactants are $C_{12}-C_{18}$ alkyl
25 polyethoxylate (1.0) sulfate ($C_{12}-C_{18}E(1.0)M$), $C_{12}-C_{18}$ alkyl polyethoxylate (2.25) sulfate ($C_{12}-C_{18}(2.25)M$), and $C_{12}-C_{18}$ alkyl polyethoxylate (3.0) sulfate ($C_{12}-C_{18}E(3.0)M$), and $C_{12}-C_{18}$ alkyl polyethoxylate (4.0) sulfate ($C_{12}-C_{18}E(4.0)M$), wherein M is conveniently selected from sodium and potassium.

30 Suitable anionic surfactants to be used are alkyl ester sulfonate surfactants including linear esters of C_8-C_{20} carboxylic acids (i.e., fatty acids) which are sulfonated with

gaseous SO₃ according to "The Journal of the American Oil Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm oil, etc.

5 The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester sulfonate surfactants of the structural formula:



wherein R³ is a C₈-C₂₀ hydrocarbyl, preferably an alkyl, or
 15 combination thereof, R⁴ is a C₁-C₆ hydrocarbyl, preferably an alkyl, or combination thereof, and M is a cation which forms a water soluble salt with the alkyl ester sulfonate. Suitable salt-forming cations include metals such as sodium, potassium, and lithium, and substituted or unsubstituted ammonium cations,
 20 such as monoethanolamine, diethanolamine, and triethanolamine. Preferably, R³ is C₁₀-C₁₆ alkyl, and R⁴ is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates wherein R³ is C₁₀-C₁₆ alkyl.

Other suitable anionic surfactants include the alkyl
 25 sulfate surfactants which are water soluble salts or acids of the formula ROSO₃M wherein R preferably is a C₁₀-C₂₄ hydrocarbyl, preferably an alkyl or hydroxyalkyl having a C₁₀-C₂₀ alkyl component, more preferably a C₁₂-C₁₈ alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium,
 30 potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and

dimethyl piperdinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like). Typically, alkyl chains of C_{12} - C_{16} are preferred for lower wash temperatures (e.g. below about 50°C) and C_{16} - C_{18} alkyl chains are preferred for higher wash temperatures (e.g. above about 50°C).

Other anionic surfactants useful for deterative purposes can also be included in the laundry detergent compositions of the present invention. These can include salts (including, for example, sodium, potassium, ammonium, and substituted ammonium salts such as mono- di- and triethanolamine salts) of soap, C_8 - C_{22} primary or secondary alkanesulfonates, C_8 - C_{24} olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, C_8 - C_{24} alkylpolyglycoethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated C_{12} - C_{18} monoesters) and diesters of sulfosuccinates (especially saturated and unsaturated C_6 - C_{12} diesters), acyl sarcosinates, sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), branched primary alkyl sulfates, and alkyl polyethoxy carboxylates such as those of the formula $\text{RO}(\text{CH}_2\text{CH}_2\text{O})_k\text{-CH}_2\text{COO-M}^+$ wherein R is a C_8 - C_{22} alkyl, k is an integer from 1 to 10, and M is a soluble salt forming cation. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated

resin acids present in or derived from tall oil.


Alkylbenzene sulfonates are highly preferred. Especially preferred are linear (straight-chain) alkyl benzene sulfonates (LAS) wherein the alkyl group preferably contains from 10 to 18 carbon atoms.

Further examples are described in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in US 3,929,678, (Column 23, line 58 through Column 29, line 23).

When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to about 40%, preferably from about 3% to about 20% by weight of such anionic surfactants.

The cleaning or laundry detergent compositions of the present invention may also contain cationic, ampholytic, zwitterionic, and semi-polar surfactants, as well as the nonionic and/or anionic surfactants other than those already described herein.

Cationic deterative surfactants suitable for use in the laundry detergent compositions of the present invention are those having one long-chain hydrocarbyl group. Examples of such cationic surfactants include the ammonium surfactants such as alkyltrimethylammonium halogenides, and those surfactants having the formula:



wherein R^2 is an alkyl or alkyl benzyl group having from about 8 to about 18 carbon atoms in the alkyl chain, each R^3 is selected from the group consisting of $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}(\text{CH}_3)-$, $-\text{CH}_2\text{CH}(\text{CH}_2\text{OH})-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, and mixtures thereof; each R^4 is
 5 selected from the group consisting of C_1 - C_4 alkyl, C_1 - C_4 hydroxyalkyl, benzyl ring structures formed by joining the two R^4 groups, $-\text{CH}_2\text{CHOHCHOH}\text{COR}^6\text{CHOHCH}_2\text{OH}$, wherein R^6 is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when y is not 0; R^5 is the same as R^4 or is an alkyl
 10 chain, wherein the total number of carbon atoms or R^2 plus R^5 is not more than about 18; each y is from 0 to about 10, and the sum of the y values is from 0 to about 15; and X is any compatible anion.

Highly preferred cationic surfactants are the water
 15 soluble quaternary ammonium compounds useful in the present composition having the formula:




wherein R_1 is C_8 - C_{16} alkyl, each of R_2 , R_3 and R_4 is independently
 20 C_1 - C_4 alkyl, C_1 - C_4 hydroxy alkyl, benzyl, and $-(\text{C}_2\text{H}_4)_x\text{H}$ where x has a value from 2 to 5, and X is an anion. Not more than one of R_2 , R_3 or R_4 should be benzyl.

The preferred alkyl chain length for R_1 is C_{12} - C_{15} , particularly where the alkyl group is a mixture of chain lengths
 25 derived from coconut or palm kernel fat or is derived synthetically by olefin build up or OXO alcohols synthesis.

Preferred groups for R_2 , R_3 and R_4 are methyl and hydroxyethyl groups and the anion X may be selected from halide, methosulphate, acetate and phosphate ions.

30 Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are:

- coconut trimethyl ammonium chloride or bromide;
 coconut methyl dihydroxyethyl ammonium chloride or bromide;
 decyl triethyl ammonium chloride;
 decyl dimethyl hydroxyethyl ammonium chloride or bromide;
 5 C₁₂₋₁₅ dimethyl hydroxyethyl ammonium chloride or bromide;
 coconut dimethyl hydroxyethyl ammonium chloride or bromide;
 myristyl trimethyl ammonium methyl sulphate;
 lauryl dimethyl benzyl ammonium chloride or bromide;
 lauryl dimethyl (ethenoxy)₄ ammonium chloride or bromide;
 10 choline esters (compounds of formula (i) wherein R₁ is



- 15 di-alkyl imidazolines [compounds of formula (i)].

Other cationic surfactants useful herein are also described in US 4,228,044 and in EP 000 224.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about
 20 25%, preferably from about 1% to about 8% by weight of such cationic surfactants.

Ampholytic surfactants are also suitable for use in the laundry detergent compositions of the present invention. These surfactants can be broadly described as aliphatic derivatives of
 25 secondary or tertiary amines, or aliphatic derivatives of heterocyclic secondary and tertiary amines in which the aliphatic radical can be straight- or branched-chain. One of the aliphatic substituents contains at least about 8 carbon atoms, typically from about 8 to about 18 carbon atoms, and at least
 30 one contains an anionic water-solubilizing group, e.g. carboxy, sulfonate, sulfate. See US 3,929,678 (column 19, lines 18-35) for examples of ampholytic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such ampholytic surfactants.

5 Zwitterionic surfactants are also suitable for use in laundry detergent compositions. These surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or
10 tertiary sulfonium compounds. See US 3,929,678 (column 19, line 38 through column 22, line 48) for examples of zwitterionic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about
15 15%, preferably from about 1% to about 10% by weight of such zwitterionic surfactants.

Semi-polar nonionic surfactants are a special category of nonionic surfactants which include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon
20 atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; watersoluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and
25 hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety from about 10 to about 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms.

30 Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula:

O

↑



5 wherein R^3 is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms; R^4 is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof; x is from 0 to about 3; and each R^5 is an alkyl or hydroxyalkyl group
 10 containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The R^5 groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.

15 These amine oxide surfactants in particular include C_{10} - C_{18} alkyl dimethyl amine oxides and C_8 - C_{12} alkoxy ethyl dihydroxy ethyl amine oxides.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about
 20 15%, preferably from about 1% to about 10% by weight of such semi-polar nonionic surfactants.

Builder system

The compositions according to the present invention may
 25 further comprise a builder system. Any conventional builder system is suitable for use herein including aluminosilicate materials, silicates, polycarboxylates and fatty acids, materials such as ethylenediamine tetraacetate, metal ion sequestrants such as aminopolyphosphonates, particularly
 30 ethylenediamine tetramethylene phosphonic acid and diethylene triamine pentamethylenephosphonic acid. Though less preferred for obvious environmental reasons, phosphate builders can also

be used herein.

Suitable builders can be an inorganic ion exchange material, commonly an inorganic hydrated aluminosilicate material, more particularly a hydrated synthetic zeolite such as
5 hydrated zeolite A, X, B, HS or MAP.

Another suitable inorganic builder material is layered silicate, e.g. SKS-6* (Hoechst). SKS-6 is a crystalline layered silicate consisting of sodium silicate ($\text{Na}_2\text{Si}_2\text{O}_5$).

Suitable polycarboxylates containing one carboxy group
10 include lactic acid, glycolic acid and ether derivatives thereof as disclosed in Belgian Patent Nos. 831,368, 821,369 and 821,370. Polycarboxylates containing two carboxy groups include the water-soluble salts of succinic acid, malonic acid, (ethylenedioxy) diacetic acid, maleic acid, diglycollic acid,
15 tartaric acid, tartronic acid and fumaric acid, as well as the ether carboxylates described in German Offenle-enschrift 2,446,686, and 2,446,487, US 3,935,257 and the sulfinyl carboxylates described in Belgian Patent No. 840,623. Polycarboxylates containing three carboxy groups include, in
20 particular, water-soluble citrates, aconitrates and citraconates as well as succinate derivatives such as the carboxymethyloxysuccinates described in British Patent No. 1,379,241, lactoxysuccinates described in Netherlands Application 7205873, and the oxypolycarboxylate materials such
25 as 2-oxa-1,1,3-propane tricarboxylates described in British Patent No. 1,387,447.

Polycarboxylates containing four carboxy groups include oxydisuccinates disclosed in British Patent No. 1,261,829, 1,1,2,2,-ethane tetracarboxylates, 1,1,3,3-propane
30 tetracarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in British Patent Nos. 1,398,421 and 1,398,422 and in US 3,936,448, and the sulfonated

pyrolysed citrates described in British Patent No. 1,082,179, while polycarboxylates containing phosphone substituents are disclosed in British Patent No. 1,439,000.

Alicyclic and heterocyclic polycarboxylates include
5 cyclopentane-cis,cis-cis-tetracarboxylates, cyclopentadienide pentacarboxylates, 2,3,4,5-tetrahydro-furan - cis, cis, cis-tetracarboxylates, 2,5-tetrahydro-furan-cis, discarboxylates, 2,2,5,5,-tetrahydrofuran - tetracarboxylates, 1,2,3,4,5,6-hexane - hexacarboxylates and carboxymethyl derivatives of polyhydric
10 alcohols such as sorbitol, mannitol and xylitol. Aromatic polycarboxylates include mellitic acid, pyromellitic acid and the phthalic acid derivatives disclosed in British Patent No. 1,425,343.

Of the above, the preferred polycarboxylates are hydroxy-
15 carboxylates containing up to three carboxy groups per molecule, more particularly citrates.

Preferred builder systems for use in the present compositions include a mixture of a water-insoluble aluminosilicate builder such as zeolite A or of a layered
20 silicate (SKS-6), and a water-soluble carboxylate chelating agent such as citric acid.

A suitable chelant for inclusion in the detergent compositions in accordance with the invention is ethylenediamine-N,N'-disuccinic acid (EDDS) or the alkali metal, alkaline earth
25 metal, ammonium, or substituted ammonium salts thereof, or mixtures thereof. Preferred EDDS compounds are the free acid form and the sodium or magnesium salt thereof. Examples of such preferred sodium salts of EDDS include Na_2EDDS and Na_4EDDS . Examples of such preferred magnesium salts of EDDS include
30 MgEDDS and Mg_2EDDS . The magnesium salts are the most preferred for inclusion in compositions in accordance with the invention.

Preferred builder systems include a mixture of a water-insoluble aluminosilicate builder such as zeolite A, and a water soluble carboxylate chelating agent such as citric acid.

Other builder materials that can form part of the builder system for use in granular compositions include inorganic materials such as alkali metal carbonates, bicarbonates, silicates, and organic materials such as the organic phosphonates, amino polyalkylene phosphonates and amino polycarboxylates.

Other suitable water-soluble organic salts are the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms.

Polymers of this type are disclosed in GB-A-1,596,756. Examples of such salts are polyacrylates of MW 2000-5000 and their copolymers with maleic anhydride, such copolymers having a molecular weight of from 20,000 to 70,000, especially about 40,000.

Detergency builder salts are normally included in amounts of from 5% to 80% by weight of the composition. Preferred levels of builder for liquid detergents are from 5% to 30%.

Enzymes:

Mannanase is incorporated into the cleaning or detergent compositions in accordance with the invention preferably at a level of from 0.0001% to 2%, more preferably from 0.0005% to 0.5%, most preferred from 0.001% to 0.1% pure enzyme by weight of the composition.

The cleaning compositions of the present invention may further comprise as an essential element a carbohydrase selected from the group consisting of cellulases, amylases, pectin degrading enzymes and xyloglucanases. Preferably, the cleaning

compositions of the present invention will comprise a mannanase, an amylase and another bioscouring-type of enzyme selected from the group consisting of cellulases, pectin degrading enzymes and xyloglucanases.

5 The cellulases usable in the present invention include both bacterial or fungal cellulases. Preferably, they will have a pH optimum of between 5 and 12 and a specific activity above 50 CEVU/mg (Cellulose Viscosity Unit). Suitable cellulases are disclosed in U.S. Patent 4,435,307, J61078384 and WO96/02653
10 which discloses fungal cellulase produced from *Humicola insolens*, *Trichoderma*, *Thielavia* and *Sporotrichum*, respectively. EP 739 982 describes cellulases isolated from novel *Bacillus* species. Suitable cellulases are also disclosed in GB-A-2075028; GB-A-2095275; DE-OS-22 47 832 and WO95/26398.

15 Examples of such cellulases are cellulases produced by a strain of *Humicola insolens* (*Humicola grisea* var. *thermoidea*), particularly the strain *Humicola insolens*, DSM 1800. Other suitable cellulases are cellulases originated from *Humicola insolens* having a molecular weight of about 50kD, an isoelectric
20 point of 5.5 and containing 415 amino acids; and a ~43kD endo-beta-1,4-glucanase derived from *Humicola insolens*, DSM 1800; a preferred cellulase has the amino acid sequence disclosed in PCT Patent Application No. WO 91/17243. Also suitable cellulases are the EGIII cellulases from *Trichoderma longibrachiatum* described
25 in WO94/21801. Especially suitable cellulases are the cellulases having color care benefits. Examples of such cellulases are the cellulases described in WO96/29397, EP-A-0495257, WO 91/17243, WO91/17244 and WO91/21801. Other suitable cellulases for fabric care and/or cleaning properties are described in WO96/34092,
30 WO96/17994 and WO95/24471.

Said cellulases are normally incorporated in the detergent composition at levels from 0.0001% to 2% of pure enzyme by

weight of the detergent composition.

Preferred cellulases for the purpose of the present invention are alkaline cellulases, i.e. enzyme having at least 25%, more preferably at least 40% of their maximum activity at a pH ranging from 7 to 12. More preferred cellulases are enzymes having their maximum activity at a pH ranging from 7 to 12. A preferred alkaline cellulase is the cellulase sold under the tradename Carezyme® by Novo Nordisk A/S.

Amylases (α and/or β) can be included for removal of carbohydrate-based stains. WO94/02597, Novo Nordisk A/S published February 03, 1994, describes cleaning compositions which incorporate mutant amylases. See also WO95/10603, Novo Nordisk A/S, published April 20, 1995. Other amylases known for use in cleaning compositions include both α - and β -amylases. α -Amylases are known in the art and include those disclosed in US Pat. no. 5,003,257; EP 252,666; WO/91/00353; FR 2,676,456; EP 285,123; EP 525,610; EP 368,341; and British Patent specification no. 1,296,839 (Novo). Other suitable amylases are stability-enhanced amylases described in WO94/18314, published August 18, 1994 and WO96/05295, Genencor, published February 22, 1996 and amylase variants having additional modification in the immediate parent available from Novo Nordisk A/S, disclosed in WO 95/10603, published April 95. Also suitable are amylases described in EP 277 216, WO95/26397 and WO96/23873 (all by Novo Nordisk).

Examples of commercial α -amylases products are Purafect Ox Am® from Genencor and Termamyl®, Ban®, Fungamyl® and Duramyl®, all available from Novo Nordisk A/S Denmark. WO95/26397 describes other suitable amylases : α -amylases characterised by having a specific activity at least 25% higher than the specific activity of Termamyl® at a temperature range of 25°C to 55°C and at a pH value in the range of 8 to 10, measured by the Phadebas

® α -amylase activity assay. Suitable are variants of the above enzymes, described in WO96/23873 (Novo Nordisk). Other amylolytic enzymes with improved properties with respect to the activity level and the combination of thermostability and a higher activity level are described in WO95/35382.

Preferred amylases for the purpose of the present invention are the amylases sold under the tradename Termamyl, Duramyl and Maxamyl and or the α -amylase variant demonstrating increased thermostability disclosed as SEQ ID No. 2 in WO96/23873.

Preferred amylases for specific applications are alkaline amylases, ie enzymes having an enzymatic activity of at least 10%, preferably at least 25%, more preferably at least 40% of their maximum activity at a pH ranging from 7 to 12. More preferred amylases are enzymes having their maximum activity at a pH ranging from 7 to 12.

The amylolytic enzymes are incorporated in the detergent compositions of the present invention a level of from 0.0001% to 2%, preferably from 0.00018% to 0.06%, more preferably from 0.00024% to 0.048% pure enzyme by weight of the composition.

The term "pectin degrading enzyme" is intended to encompass arabinanase (EC 3.2.1.99), galactanases (EC 3.2.1.89), polygalacturonase (EC 3.2.1.15) exo-polygalacturonase (EC 3.2.1.67), exo-poly-alpha-galacturonidase (EC 3.2.1.82), pectin lyase (EC 4.2.2.10), pectin esterase (EC 3.2.1.11), pectate lyase (EC 4.2.2.2), exo-polygalacturonate lyase (EC 4.2.2.9) and hemicellulases such as endo-1,3- β -xylosidase (EC 3.2.1.32), xylan-1,4- β -xylosidase (EC 3.2.1.37) and α -L-arabinofuranosidase (EC 3.2.1.55). The pectin degrading enzymes are natural mixtures of the above mentioned enzymatic activities. Pectin enzymes therefore include the pectin methylesterases which hydrolyse the pectin methyl ester linkages, polygalacturonases which cleave the glycosidic bonds between galacturonic acid molecules, and

the pectin transeliminases or lyases which act on the pectic acids to bring about non-hydrolytic cleavage of α -1 $\rightarrow$ 4 glycosidic linkages to form unsaturated derivatives of galacturonic acid.

5 Pectin degrading enzymes are incorporated into the compositions in accordance with the invention preferably at a level of from 0.0001 % to 2 %, more preferably from 0.0005% to 0.5%, most preferred from 0.001 % to 0.1 % pure enzyme by weight of the total composition.

10 Preferred pectin degrading enzymes for specific applications are alkaline pectin degrading enzymes, ie enzymes having an enzymatic activity of at least 10%, preferably at least 25%, more preferably at least 40% of their maximum activity at a pH ranging from 7 to 12. More preferred pectin degrading enzymes
15 are enzymes having their maximum activity at a pH ranging from 7 to 12. Alkaline pectin degrading enzymes are produced by alkalophilic microorganisms e.g. bacterial, fungal and yeast microorganisms such as *Bacillus* species. Preferred microorganisms are *Bacillus firmus*, *Bacillus circulans* and *Bacillus subtilis* as
20 described in JP 56131376 and JP 56068393. Alkaline pectin decomposing enzymes include galacturan-1,4- α -galacturonase (EC 3.2.1.67), poly-galacturonase activities (EC 3.2.1.15, pectin esterase (EC 3.1.1.11), pectate lyase (EC 4.2.2.2) and their iso enzymes and they can be produced by the *Erwinia* species. Preferred
25 are *E. chrysanthemi*, *E. carotovora*, *E. amylovora*, *E. herbicola*, *E. dissolvens* as described in JP 59066588, JP 63042988 and in World J. Microbiol. Microbiotechnol. (8, 2, 115-120) 1992. Said alkaline pectin enzymes can also be produced by *Bacillus* species as disclosed in JP 73006557 and Agr. Biol. Chem. (1972), 36(2) 285-93.
30

The term xyloglucanase encompasses the family of enzymes described by Vincken and Voragen at Wageningen University

[Vincken et al (1994) Plant Physiol., **104**, 99-107] and are able to degrade xyloglucans as described in Hayashi et al (1989) Plant. Physiol. Plant Mol. Biol., **40**, 139-168. Vincken et al demonstrated the removal of xyloglucan coating from cellulose of the isolated apple cell wall by a xyloglucanase purified from *Trichoderma viride* (endo-IV-glucanase). This enzyme enhances the enzymatic degradation of cell wall-embedded cellulose and work in synergy with pectic enzymes. Rapidase LIQ+ from Gist-Brocades contains an xyloglucanase activity.

10 This xyloglucanase is incorporated into the cleaning compositions in accordance with the invention preferably at a level of from 0.0001% to 2%, more preferably from 0.0005% to 0.5%, most preferred from 0.001% to 0.1 % pure enzyme by weight of the composition.

15 Preferred xyloglucanases for specific applications are alkaline xyloglucanases, ie enzymes having an enzymatic activity of at least 10%, preferably at least 25%, more preferably at least 40% of their maximum activity at a pH ranging from 7 to 12. More preferred xyloglucanases are enzymes having their
20 maximum activity at a pH ranging from 7 to 12.

The above-mentioned enzymes may be of any suitable origin, such as vegetable, animal, bacterial, fungal and yeast origin. Origin can further be mesophilic or extremophilic (psychrophilic, psychrotrophic, thermophilic, barophilic, alkalophilic, acidophilic, halophilic, etc.). Purified or non-
25 purified forms of these enzymes may be used. Nowadays, it is common practice to modify wild-type enzymes via protein / genetic engineering techniques in order to optimise their performance efficiency in the cleaning compositions of the invention.
30 For example, the variants may be designed such that the compatibility of the enzyme to commonly encountered ingredients of such compositions is increased. Alternatively, the variant may be

designed such that the optimal pH, bleach or chelant stability, catalytic activity and the like, of the enzyme variant is tailored to suit the particular cleaning application.

In particular, attention should be focused on amino acids sensitive to oxidation in the case of bleach stability and on surface charges for the surfactant compatibility. The isoelectric point of such enzymes may be modified by the substitution of some charged amino acids, e.g. an increase in isoelectric point may help to improve compatibility with anionic surfactants. The stability of the enzymes may be further enhanced by the creation of e.g. additional salt bridges and enforcing metal binding sites to increase chelant stability.

Bleaching agents:

Additional optional detergent ingredients that can be included in the detergent compositions of the present invention include bleaching agents such as PB1, PB4 and percarbonate with a particle size of 400-800 microns. These bleaching agent components can include one or more oxygen bleaching agents and, depending upon the bleaching agent chosen, one or more bleach activators. When present oxygen bleaching compounds will typically be present at levels of from about 1% to about 25%. In general, bleaching compounds are optional added components in non-liquid formulations, e.g. granular detergents.

A bleaching agent component for use herein can be any of the bleaching agents useful for detergent compositions including oxygen bleaches, as well as others known in the art.

A bleaching agent suitable for the present invention can be an activated or non-activated bleaching agent.

One category of oxygen bleaching agent that can be used encompasses percarboxylic acid bleaching agents and salts thereof. Suitable examples of this class of agents include

magnesium monoperoxyphthalate hexahydrate, the magnesium salt of meta-chloro perbenzoic acid, 4-nonylamino-4-oxoperoxybutyric acid and diperoxydodecanedioic acid. Such bleaching agents are disclosed in US 4,483,781, US 740,446, EP 0 133 354 and US 5 4,412,934. Highly preferred bleaching agents also include 6-nonylamino-6-oxoperoxy-caproic acid as described in US 4,634,551.

Another category of bleaching agents that can be used encompasses the halogen bleaching agents. Examples of hypohalite bleaching agents, for example, include trichloro isocyanuric 10 acid and the sodium and potassium dichloroisocyanurates and N-chloro and N-bromo alkane sulphonamides. Such materials are normally added at 0.5-10% by weight of the finished product, preferably 1-5% by weight.

The hydrogen peroxide releasing agents can be used in 15 combination with bleach activators such as tetra-acetylenethylenediamine (TAED), nonanoyloxybenzenesulfonate (NOBS, described in US 4,412,934), 3,5-trimethyl-hexsanoyloxybenzenesulfonate (ISONOBS, described in EP 120 591) or pentaacetylglucose (PAG), which are perhydrolyzed to form a 20 peracid as the active bleaching species, leading to improved bleaching effect. In addition, very suitable are the bleach activators C8(6-octanamido-caproyl) oxybenzene-sulfonate, C9(6-nonanamido caproyl) oxybenzenesulfonate and C10 (6-decanamido caproyl) oxybenzenesulfonate or mixtures thereof. Also suitable 25 activators are acylated citrate esters such as disclosed in European Patent Application No. 91870207.7.

Useful bleaching agents, including peroxyacids and bleaching systems comprising bleach activators and peroxygen bleaching compounds for use in cleaning compositions according 30 to the invention are described in U.S. Patent No. 5,677,272.

The hydrogen peroxide may also be present by adding an enzymatic system (i.e. an enzyme and a substrate therefore)

which is capable of generation of hydrogen peroxide at the beginning or during the washing and/or rinsing process. Such enzymatic systems are disclosed in European Patent Application EP 0 537 381.

5 Bleaching agents other than oxygen bleaching agents are also known in the art and can be utilized herein. One type of non-oxygen bleaching agent of particular interest includes photoactivated bleaching agents such as the sulfonated zinc and/or aluminium phthalocyanines. These materials can be deposited
10 upon the substrate during the washing process. Upon irradiation with light, in the presence of oxygen, such as by hanging clothes out to dry in the daylight, the sulfonated zinc phthalocyanine is activated and, consequently, the substrate is bleached. Preferred zinc phthalocyanine and a photoactivated
15 bleaching process are described in US 4,033,718. Typically, detergent composition will contain about 0.025% to about 1.25%, by weight, of sulfonated zinc phthalocyanine.

Bleaching agents may also comprise a manganese catalyst. The manganese catalyst may, e.g., be one of the compounds
20 described in "Efficient manganese catalysts for low-temperature bleaching", Nature 369, 1994, pp. 637-639.

Suds suppressors:

Another optional ingredient is a suds suppressor,
25 exemplified by silicones, and silica-silicone mixtures. Silicones can generally be represented by alkylated polysiloxane materials, while silica is normally used in finely divided forms exemplified by silica aerogels and xerogels and hydrophobic silicas of various types. These materials can be incorporated
30 as particulates, in which the suds suppressor is advantageously releasably incorporated in a water-soluble or water-dispersible, substantially non surface-active detergent impermeable carrier.

Alternatively the suds suppressor can be dissolved or dispersed in a liquid carrier and applied by spraying on to one or more of the other components.

A preferred silicone suds controlling agent is disclosed in US 3,933,672. Other particularly useful suds suppressors are the self-emulsifying silicone suds suppressors, described in German Patent Application DTOS 2,646,126. An example of such a compound is DC-544, commercially available from Dow Corning, which is a siloxane-glycol copolymer. Especially preferred suds controlling agent are the suds suppressor system comprising a mixture of silicone oils and 2-alkyl-alkanols. Suitable 2-alkyl-alkanols are 2-butyl-octanol which are commercially available under the trade name Isofol^{*} 12 R.

Such suds suppressor system are described in European Patent Application EP 0 593 841.

Especially preferred silicone suds controlling agents are described in European Patent Application No. 92201649.8. Said compositions can comprise a silicone/ silica mixture in combination with fumed nonporous silica such as Aerosil^R.

The suds suppressors described above are normally employed at levels of from 0.001% to 2% by weight of the composition, preferably from 0.01% to 1% by weight.

Other components:

Other components used in detergent compositions may be employed, such as soil-suspending agents, soil-releasing agents, optical brighteners, abrasives, bactericides, tarnish inhibitors, coloring agents, and/or encapsulated or nonencapsulated perfumes.

Especially suitable encapsulating materials are water soluble capsules which consist of a matrix of polysaccharide and polyhydroxy compounds such as described in GB 1,464,616.

^{*}Trade-mark

Other suitable water soluble encapsulating materials comprise dextrans derived from ungelatinized starch acid esters of substituted dicarboxylic acids such as described in US 3,455,838. These acid-ester dextrans are, preferably, prepared from such starches as waxy maize, waxy sorghum, sago, tapioca and potato. Suitable examples of said encapsulation materials include N-Lok manufactured by National Starch. The N-Lok encapsulating material consists of a modified maize starch and glucose. The starch is modified by adding monofunctional substituted groups such as octenyl succinic acid anhydride.


Antiredeposition and soil suspension agents suitable herein include cellulose derivatives such as methylcellulose, carboxymethylcellulose and hydroxyethylcellulose, and homo- or co-polymeric polycarboxylic acids or their salts..Polymers of this type include the polyacrylates and maleic anhydride-acrylic acid copolymers previously mentioned as builders, as well as copolymers of maleic anhydride with ethylene, methylvinyl ether or methacrylic acid, the maleic anhydride constituting at least 20 mole percent of the copolymer. These materials are normally used at levels of from 0.5% to 10% by weight, more preferably from 0.75% to 8%, most preferably from 1% to 6% by weight of the composition.

Preferred optical brighteners are anionic in character, examples of which are disodium 4,4'-bis-(2-diethanolamino-4-anilino -s- triazin-6-ylamino)stilbene-2:2' disulphonate, disodium 4, - 4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino-stilbene-2:2' - disulphonate, disodium 4,4' - bis-(2,4-dianilino-s-triazin-6-ylamino)stilbene-2:2' - disulphonate, monosodium 4',4'' - bis-(2,4-dianilino-s-tri-azin-6 ylamino)stilbene-2-sulphonate, disodium 4,4' -bis-(2-anilino-4-(N-methyl-N-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2' - disulphonate, di-sodium 4,4' -bis-(4-phenyl-2,1,3-

triazol-2-yl)-stilbene-2,2' disulphonate, di-so-dium 4,4'-bis(2-anilino-4-(1-methyl-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2'disulphonate, sodium 2(stilbyl-4''-(naphtho-1',2':4,5)-1,2,3, - triazole-2''-sulphonate and 4,4'-bis(2-sulphostyryl)biphenyl.

Other useful polymeric materials are the polyethylene glycols, particularly those of molecular weight 1000-10000, more particularly 2000 to 8000 and most preferably about 4000. These are used at levels of from 0.20% to 5% more preferably from 0.25% to 2.5% by weight. These polymers and the previously mentioned homo- or co-polymeric poly-carboxylate salts are valuable for improving whiteness maintenance, fabric ash deposition, and cleaning performance on clay, proteinaceous and oxidizable soils in the presence of transition metal impurities.

Soil release agents useful in compositions of the present invention are conventionally copolymers or terpolymers of terephthalic acid with ethylene glycol and/or propylene glycol units in various arrangements. Examples of such polymers are disclosed in US 4,116,885 and 4,711,730 and EP 0 272 033. A particular preferred polymer in accordance with EP 0 272 033 has the formula:



where PEG is $-(\text{OC}_2\text{H}_4)_n-$, PO is $(\text{OC}_3\text{H}_6\text{O})$ and T is $(\text{pOOC}_6\text{H}_4\text{CO})$.

Also very useful are modified polyesters as random copolymers of dimethyl terephthalate, dimethyl sulfoisophthalate, ethylene glycol and 1,2-propanediol, the end groups consisting primarily of sulphobenzoate and secondarily of mono esters of ethylene glycol and/or 1,2-propanediol. The target is to obtain a polymer capped at both end by sulphobenzoate groups, "primarily", in the present context most

of said copolymers herein will be endcapped by sulphobenzoate groups. However, some copolymers will be less than fully capped, and therefore their end groups may consist of monoester of ethylene glycol and/or 1,2-propanediol, thereof consist "secondarily" of such species.

The selected polyesters herein contain about 46% by weight of dimethyl terephthalic acid, about 16% by weight of 1,2-propanediol, about 10% by weight ethylene glycol, about 13% by weight of dimethyl sulfobenzoic acid and about 15% by weight of sulfoisophthalic acid, and have a molecular weight of about 3.000. The polyesters and their method of preparation are described in detail in EP 311 342.

Softening agents:

Fabric softening agents can also be incorporated into laundry detergent compositions in accordance with the present invention. These agents may be inorganic or organic in type. Inorganic softening agents are exemplified by the smectite clays disclosed in GB-A-1 400898 and in US 5,019,292. Organic fabric softening agents include the water insoluble tertiary amines as disclosed in GB-A1 514 276 and EP 0 011 340 and their combination with mono C_{12} - C_{14} quaternary ammonium salts are disclosed in EP-B-0 026 528 and di-long-chain amides as disclosed in EP 0 242 919. Other useful organic ingredients of fabric softening systems include high molecular weight polyethylene oxide materials as disclosed in EP 0 299 575 and 0 313 146.

Levels of smectite clay are normally in the range from 5% to 15%, more preferably from 8% to 12% by weight, with the material being added as a dry mixed component to the remainder of the formulation. Organic fabric softening agents such as the water-insoluble tertiary amines or dilong chain amide materials

are incorporated at levels of from 0.5% to 5% by weight, normally from 1% to 3% by weight whilst the high molecular weight polyethylene oxide materials and the water soluble cationic materials are added at levels of from 0.1% to 2%, normally from 0.15% to 1.5% by weight. These materials are normally added to the spray dried portion of the composition, although in some instances it may be more convenient to add them as a dry mixed particulate, or spray them as molten liquid on to other solid components of the composition.

10

Polymeric dye-transfer inhibiting agents:

The detergent compositions according to the present invention may also comprise from 0.001% to 10%, preferably from 0.01% to 2%, more preferably from 0.05% to 1% by weight of polymeric dye-transfer inhibiting agents. Said polymeric dye-transfer inhibiting agents are normally incorporated into detergent compositions in order to inhibit the transfer of dyes from colored fabrics onto fabrics washed therewith. These polymers have the ability of complexing or adsorbing the fugitive dyes washed out of dyed fabrics before the dyes have the opportunity to become attached to other articles in the wash.

Especially suitable polymeric dye-transfer inhibiting agents are polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylpyrrolidone polymers, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof.

Addition of such polymers also enhances the performance of the enzymes according the invention.

30

Use in the paper pulp industry

Further, it is contemplated that the mannanase of the present invention is useful in chlorine-free bleaching processes for paper pulp (chemical pulps, semichemical pulps, mechanical pulps or kraft pulps) in order to increase the brightness thereof, thus decreasing or eliminating the need for hydrogen peroxide in the bleaching process.

Use in the textile and cellulosic fiber processing industries

The mannanase of the present invention can be used in combination with other carbohydrate degrading enzymes (for instance xyloglucanase, xylanase, various pectinases) for preparation of fibers or for cleaning of fibers in combination with detergents.

In the present context, the term "cellulosic material" is intended to mean fibers, sewn and unsewn fabrics, including knits, wovens, denims, yarns, and toweling, made from cotton, cotton blends or natural or manmade cellulosics (e.g. originating from xylan-containing cellulose fibers such as from wood pulp) or blends thereof. Examples of blends are blends of cotton or rayon/viscose with one or more companion material such as wool, synthetic fibers (e.g. polyamide fibers, acrylic fibers, polyester fibers, polyvinyl alcohol fibers, polyvinyl chloride fibers, polyvinylidene chloride fibers, polyurethane fibers, polyurea fibers, aramid fibers), and cellulose-containing fibers (e.g. rayon/viscose, ramie, hemp, flax/linen, jutè, cellulose acetate fibers, lyocell).

The processing of cellulosic material for the textile industry, as for example cotton fiber, into a material ready for garment manufacture involves several steps: spinning of the fiber into a yarn; construction of woven or knit fabric from the yarn and subsequent preparation, dyeing and finishing operations. Woven goods are constructed by weaving a filling yarn between a series of warp yarns; the yarns could be two

different types.

Desizing: polymeric size like e.g. mannan, starch, CMC or PVA is added before weaving in order to increase the warp speed; This material must be removed before further processing. The
5 enzyme of the invention is useful for removal of mannan containing size.

Degradation of thickeners

Galactomannans such as guar gum and locust bean gum are
10 widely used as thickening agents e.g. in food and print paste for textile printing such as prints on T-shirts. The enzyme or enzyme preparation according to the invention can be used for reducing the viscosity of eg residual food in processing equipment and thereby facilitate cleaning after processing. Further,
15 it is contemplated that the enzyme or enzyme preparation is useful for reducing viscosity of print paste, thereby facilitating wash out of surplus print paste after textile printins.

Degradation or modification of plant material

20 The enzyme or enzyme preparation according to the invention is preferably used as an agent for degradation or modification of mannan, galactomannan, glucomannan or galactoglucomannan containing material originating from plants. Examples of such material is guar gum and locust bean gum.

25 The mannanase of the invention may be used in modifying the physical-chemical properties of plant derived material such as the viscosity. For instance, the mannanase may be used to reduce the viscosity of feed or food which contain mannan and to promote processing of viscous mannan containing material.

30

Coffee extraction

The enzyme or enzyme preparation of the invention may also be used for hydrolysing galactomannans present in a liquid coffee extract, preferably in order to inhibit gel formation during freeze drying of the (instant) coffee. Preferably, the mannanase
5 of the invention is immobilized in order to reduce enzyme consumption and avoid contamination of the coffee. This use is further disclosed in EP-A-676 145.

Use in the fracturing of a subterranean formation (oil drilling)

10 Further, it is contemplated that the enzyme of the present invention is useful as an enzyme breaker as disclosed in US patent nos. 5,806,597, 5,562,160, 5,201,370 and 5,067,566 to BJ Services Company (Houston, TX, U.S.A.).

15 Accordingly, the mannanase of the present invention is useful in a method of fracturing a subterranean formation in a well bore in which a gellable fracturing fluid is first formed by blending together an aqueous fluid, a hydratable polymer, a suitable cross-linking agent for cross-linking the hydratable
20 polymer to form a polymer gel and an enzyme breaker, ie the enzyme of the invention. The cross-linked polymer gel is pumped into the well bore under sufficient pressure to fracture the surrounding formation. The enzyme breaker is allowed to degrade the cross-linked polymer with time to reduce the viscosity of
25 the fluid so that the fluid can be pumped from the formation back to the well surface.

The enzyme breaker may be an ingredient of a fracturing fluid or a breaker-crosslinker-polymer complex which further comprises a hydratable polymer and a crosslinking agent. The
30 fracturing fluid or complex may be a gel or may be gellable. The complex is useful in a method for using the complex in a fracturing fluid to fracture a subterranean formation that surrounds

a well bore by pumping the fluid to a desired location within the well bore under sufficient pressure to fracture the surrounding subterranean formation. The complex may be maintained in a substantially non-reactive state by maintaining specific
5 conditions of pH and temperature, until a time at which the fluid is in place in the well bore and the desired fracture is completed. Once the fracture is completed, the specific conditions at which the complex is inactive are no longer maintained. When the conditions change sufficiently, the complex becomes
10 active and the breaker begins to catalyze polymer degradation causing the fracturing fluid to become sufficiently fluid to be pumped from the subterranean formation to the well surface.

MATERIALS AND METHODS

15 Assay for activity test

A polypeptide of the invention having mannanase activity may be tested for mannanase activity according to standard test procedures known in the art, such as by applying a solution to be tested to 4 mm diameter holes punched out in agar plates
20 containing 0.2% AZCL galactomannan (carob), i.e. substrate for the assay of endo-1,4-beta-D-mannanase available as CatNo.I-AZGMA from the company Megazyme.

25 Determination of catalytic activity (ManU) of mannanase **Colorimetric Assay**

Substrate: 0.2% AZCL-Galactomannan (Megazyme, Australia) from carob in 0.1 M Glycin buffer, pH 10.0.

The assay is carried out in an Eppendorf Micro tube 1.5 ml
30 on a thermomixer with stirring and temperature control of 40°C. Incubation of 0.750 ml substrate with 0.05 ml enzyme for 20 min, stop by centrifugation for 4 minutes at 15000 rpm. The

colour of the supernatant is measured at 600 nm in a 1 cm cuvette.

One ManU (Mannanase units) gives 0.24 abs in 1 cm.

5 Strains and donor organism

The *Bacillus* sp. I633 mentioned above comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:1.

E.coli DSM 12197 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:1).

10 The *Bacillus agaradhaerens* NCIMB 40482 mentioned above comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:5.

E.coli DSM 12180 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:5).

15 The *Bacillus* sp. AAI12 mentioned above comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:9.

E.coli DSM 12433 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:9).

20 The *Bacillus halodurans* mentioned above comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:11.

E.coli DSM 12441 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:11).

The *Humicola insolens* mentioned above comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:13.

25 *E.coli* DSM 9984 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:13).

The *Bacillus* sp. AA349 mentioned above comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:15.

30 *E.coli* DSM 12432 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:15).

E.coli DSM 12847 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:17).

E.coli DSM 12848 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:19).

The *Bacillus clausii* mentioned above comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:21.

5 *E.coli* DSM 12849 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:21).

E.coli DSM 12850 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:23).

10 *Bacillus sp.* comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:25.

E.coli DSM 12846 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:25).

Bacillus sp. comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:27.

15 *E.coli* DSM 12851 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:27).

The *Bacillus licheniformis* mentioned above comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:29.

20 *E.coli* DSM 12852 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:29).

Bacillus sp. comprises the beta-1,4-mannanase encoding DNA sequence shown in SEQ.ID.NO:31.

E.coli DSM 12436 comprises the plasmid containing the DNA encoding the beta-1,4-mannanase of the invention (SEQ.ID.NO:31).

25 *E. coli* strain: Cells of *E. coli* SJ2 (Diderichsen, B., Wedsted, U., Hedegaard, L., Jensen, B. R., Sjøholm, C. (1990) Cloning of aldB, which encodes alpha-acetolactate decarboxylase, an exoenzyme from *Bacillus brevis*. J. Bacteriol., 172, 4315-4321), were prepared for and transformed by electroporation
30 using a Gene Pulser™ electroporator from BIO-RAD as described by the supplier.

B. subtilis PL2306. This strain is the *B. subtilis* DN1885 with disrupted *apr* and *npr* genes (Diderichsen, B., Wedsted, U., Hedegaard, L., Jensen, B. R., Sjøholm, C. (1990) Cloning of *aldB*, which encodes alpha-acetolactate decarboxylase, an exoenzyme from *Bacillus brevis*. J. Bacteriol., 172, 4315-4321) disrupted in the transcriptional unit of the known *Bacillus subtilis* cellulase gene, resulting in cellulase negative cells. The disruption was performed essentially as described in (Eds. A.L. Sonenshein, J.A. Hoch and Richard Losick (1993) *Bacillus subtilis* and other Gram-Positive Bacteria, American Society for microbiology, p.618).

Competent cells were prepared and transformed as described by Yasbin, R.E., Wilson, G.A. and Young, F.E. (1975) Transformation and transfection in lysogenic strains of *Bacillus subtilis*: evidence for selective induction of prophage in competent cells. J. Bacteriol, 121:296-304.

General molecular biology methods:

Unless otherwise stated all the DNA manipulations and transformations were performed using standard methods of molecular biology (Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for *Bacillus*". John Wiley and Sons, 1990).

Enzymes for DNA manipulations were used according to the manufacturer's instructions (e.g. restriction endonucleases, ligases etc. are obtainable from New England Biolabs, Inc.).

Plasmids

pSJ1678: (see International Patent Application published as WO 94/19454).

pBK-CMV (Stratagene inc., La Jolla Ca.)

pMOL944. This plasmid is a pUB110 derivative essentially
 5 containing elements making the plasmid propagatable in *Bacillus subtilis*, kanamycin resistance gene and having a strong promoter and signal peptide cloned from the *amyL* gene of *B.licheniformis* ATCC14580. The signal peptide contains a *SacII* site making it
 10 convenient to clone the DNA encoding the mature part of a protein in-fusion with the signal peptide. This results in the expression of a Pre-protein which is directed towards the exterior of the cell.

The plasmid was constructed by means of ordinary genetic engineering and is briefly described in the following.

15 Construction of pMOL944:

The pUB110 plasmid (McKenzie, T. et al., 1986, Plasmid 15:93-103) was digested with the unique restriction enzyme *NciI*. A PCR fragment amplified from the *amyL* promoter encoded on the plasmid pDN1981 (P.L. Jørgensen et al., 1990, Gene, 96, p37-41.)
 20 was digested with *NciI* and inserted in the *NciI* digested pUB110 to give the plasmid pSJ2624.

The two PCR primers used have the following sequences:

LWN5494 5'-GTCGCCGGGGCGGCCGCTATCAATTGGTAACTGTATCTCAGC -3'

LWN5495 5'-GTCGCCCGGGAGCTCTGATCAGGTACCAAGCTTGTCGACCTGCAGAA

25 TGAGGCAGCAAGAAGAT -3'

The primer #LWN5494 inserts a *NotI* site in the plasmid.

The plasmid pSJ2624 was then digested with *SacI* and *NotI* and a new PCR fragment amplified on *amyL* promoter encoded on the pDN1981 was digested with *SacI* and *NotI* and this DNA fragment
 30 was inserted in the *SacI*-*NotI* digested pSJ2624 to give the plasmid pSJ2670.

This cloning replaces the first amyL promoter cloning with the same promoter but in the opposite direction. The two primers used for PCR amplification have the following sequences:

5 #LWN5938 5'-GTCGGCGGCCGCTGATCACGTACCAAGCTTGTCGACCTGCAGAATG
AGGCAGCAAGAAGAT -3'
#LWN5939 5'-GTCGGAGCTCTATCAATTGGTAACTGTATCTCAGC -3'

The plasmid pSJ2670 was digested with the restriction en-
10 zymes PstI and BclI and a PCR fragment amplified from a cloned DNA sequence encoding the alkaline amylase SP722 (Patent # WO9526397-A1) was digested with PstI and BclI and inserted to give the plasmid pMOL944. The two primers used for PCR amplification have the following sequence:

15 #LWN7864 5' -AACAGCTGATCACGACTGATCTTTTAGCTTGGCAC-3'
#LWN7901 5' -AACTGCAGCCGCGGCACATCATAATGGGACAAATGGG -3'

The primer #LWN7901 inserts a SacII site in the plasmid.

Cultivation of donor strains and isolation of genomic DNA

20 The relevant strain of *Bacillus*, eg *Bacillus* sp. I633, was grown in TY with pH adjusted to approximately pH 9.7 by the addition of 50 ml of 1M Sodium-Sesquicarbonat per 500 ml TY. After 24 hours incubation at 30°C and 300 rpm, the cells were harvested, and genomic DNA was isolated by the method described
25 by Pitcher et al. [Pitcher, D. G., Saunders, N. A., Owen, R. J; Rapid extraction of bacterial genomic DNA with guanidium thiocyanate; Lett Appl Microbiol 1989 **8** 151-156].

Media

30 **TY** (as described in Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995).

LB agar (as described in Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995).

LBPG is LB agar supplemented with 0.5% Glucose and 0.05 M potassium phosphate, pH 7.0

AZCL-galactomannan is added to LBPG-agar to 0.5 % AZCL-galactomannan is from Megazyme, Australia.

BPX media is described in EP 0 506 780 (WO 91/09129).

NZY agar (per liter) 5 g of NaCl, 2 g of MgSO₄, 5 g of yeast extract, 10 g of NZ amine (casein hydrolysate), 15 g of agar; add deionized water to 1 liter, adjust pH with NaOH to pH 7.5 and autoclave

NZY broth (per liter) 5 g of NaCl, 2 g of MgSO₄, 5 g of yeast extract, 10 g of NZ amine (casein hydrolysate); add deionized water to 1 liter, adjust pH with NaOH to pH 7.5 and autoclave

NZY Top Agar (per liter) 5 g of NaCl, 2 g of MgSO₄, 5 g of yeast extract, 10 g of NZ amine (casein hydrolysate), 0.7 % (w/v) agarose; add deionized water to 1 liter, adjust pH with NaOH to pH 7.5 and autoclave.

The following non-limiting examples illustrate the invention.

25 **EXAMPLE 1**

Mannanase derived from *Bacillus sp* (I633)

Construction of a genomic library from *Bacillus sp*. I633 in the lambdaZAPExpress vector

30 Genomic DNA of *Bacillus sp*. I633 was partially digested with restriction enzyme *Sau3A*, and size-fractionated by electrophoresis on a 0.7 % agarose gel (SeaKem* agarose, FMC, USA).

*Trade-mark

Fragments between 1.5 and 10 kb in size were isolated and concentrated to a DNA band by running the DNA fragments backwards on a 1.5 % agarose gel followed by extraction of the fragments from the agarose gel slice using the Qiaquick gel extraction kit according to the manufacturer's instructions (Qiagen Inc., USA). To construct a genomic library, ca. 100ng of purified, fractionated DNA from above was ligated with 1 ug of BamHI-cleaved, dephosphorylated lambdaZAPexpress vector arms (Stratagene, La Jolla CA, USA) for 24 hours at + 4 °C according to the manufacturer's instructions. A 3-ul aliquot of the ligation mixture was packaged directly using the GigaPackIII Gold packaging extract (Stratagene, USA) according to the manufacturers instructions (Stratagene). The genomic lambdaZAPExpress phage library was titered using the E. coli XL1-Blue MRF- strain from Stratagene (La Jolla, USA). The unamplified genomic library comprised of 3×10^7 plaque-forming units (pfu) with a vector background of less than 1 %.

Screening for beta-mannanase clones by functional expression in lambdaZAPExpress

Approximately 5000 plaque-forming units (pfu) from the genomic library were plated on NZY-agar plates containing 0.1 % AZCL-galactomannan (MegaZyme, Australia, cat. no. I-AZGMA), using E. coli XL1-Blue MRF' (Stratagene, USA) as a host, followed by incubation of the plates at 37 °C for 24 hours. Mannanase-positive lambda clones were identified by the formation of blue hydrolysis halos around the positive phage clones. These were recovered from the screening plates by coring the TOP-agar slices containing the plaques of interest into 500 ul of SM buffer and 20 ul of chloroform. The mannanase-positive lambdaZAPExpress clones were plaque-purified by plating an aliquot of the cored phage stock on NZY plates containing 0.1 % AZCL-

galactomannan as above. Single, mannanase-positive lambda clones were cored into 500 ul of SM buffer and 20 ul of chloroform, and purified by one more plating round as described above.

5 Single-clone in vivo excision of the phagemids from the mannanase-positive lambdaZAPExpress clones

E. coli XL1-Blue' cells (Stratagene, La Jolla Ca.) were prepared and resuspended in 10mM MgSO₄ as recommended by Stratagene (La Jolla, USA). 250-ul aliquots of the pure phage
10 stocks from the mannanase-positive clones were combined in Falcon 2059 tubes with 200uls of XL1-Blue MRF' cells (OD₆₀₀=1.0) and > 10⁶ pfus/ml of the ExAssist M13 helper phage (Stratagene), and the mixtures were incubated at 37°C for 15 minutes. Three mls of NZY broth was added to each tube and the tubes were incubated at
15 37 C for 2.5 hours. The tubes were heated at 65°C for 20 minutes to kill the E. coli cells and bacteriophage lambda; the phagemids being resistant to heating. The tubes were spun at 3000 rpm for 15 minutes to remove cellular debris and the supernatants were decanted into clean Falcon 2059 tubes. Aliquots of
20 the supernatants containing the excised single-stranded phagemids were used to infect 200uls of E. coli XL0LR cells (Stratagene, OD₆₀₀=1.0 in 10mM MgSO₄) by incubation at 37°C for 15 minutes. 350uls of NZY broth was added to the cells and the tubes were incubated for 45 min at 37°C. Aliquots of the cells
25 were plated onto LB kanamycin agar plates and incubated for 24 hours at 37°C. Five excised single colonies were re-streaked onto LB kanamycin agar plates containing 0.1 % AZCL-galactomannan (MegaZyme, Australia). The mannanase-positive phagemid clones were characterized by the formation of blue
30 hydrolysis halos around the positive colonies. These were further analysed by restriction enzyme digests of the isolated plagemid DNA (QiaSpin kit, Qiagen, USA) with EcoRI, PstI, EcoRI-

PstI, and HindIII followed by agarose gel electrophoresis.

Nucleotide sequence analysis

The nucleotide sequence of the genomic beta-1,4-mannanase
5 clone pBXM3 was determined from both strands by the dideoxy
chain-termination method (Sanger, F., Nicklen, S., and Coulson,
A. R. (1977) Proc. Natl. Acad. Sci. U. S. A. 74, 5463-5467)
using 500 ng of Qiagen-purified template (Qiagen, USA), the Taq
deoxy-terminal cycle sequencing kit (Perkin-Elmer, USA), fluo-
10 rescent labeled terminators and 5 pmol of either pBK-CMV
polylinker primers (Stratagene, USA) or synthetic oligonucleo-
tide primers. Analysis of the sequence data was performed ac-
cording to Devereux et al., 1984 (Devereux, J., Haeberli, P.,
and Smithies, O. (1984) Nucleic Acids Res. 12, 387-395).

15

Sequence alignment

A multiple sequence alignment of the glycohydrolase family
5 beta-1,4-mannanase from *Bacillus* sp. I633 of the present
invention (ie SEQ ID NO:2), *Bacillus circulans* (GenBank/EMBL
20 accession no. 066185), *Vibrio* sp. (acc. no. 069347),
Streptomyces lividans (acc. no. P51529), and
Caldicellulosiruptor saccharolyticus (acc. no. P22533). The
multiple sequence alignment was created using the PileUp program
of the GCG Wisconsin software package, version 8.1.; with gap
25 creation penalty 3.00 and gap extension penalty 0.10.

Sequence Similarities

The deduced amino acid sequence of the family 5 beta-1,4-
mannanase of the present invention cloned from *Bacillus* sp. I633
30 shows 75 % similarity and 60.1 % sequence identity to the beta-
1,4-mannanase of *Bacillus circulans* (GenBank/EMBL accession no.
066185), 64.4 % similarity and 44.6 % identity to the beta-1,4-

mannanase from *Vibrio* sp. (acc. no. 069347), 63 % similarity and 43.2 % identity to the beta-1,4-mannanase from *Streptomyces lividans* (acc. no. P51529), 52.5 % similarity and 34.4 % sequence identity to the beta-1,4-mannanase from

5 *Caldicellulosiruptor saccharolyticus* (acc. no. P2253). The sequences were aligned using the GAP program of the GCG Wisconsin software package, version 8.1.; with gap creation penalty 3.00 and gap extension penalty 0.10.

10 **Cloning of *Bacillus* sp (I633) mannanase gene**

A. Subcloning and expression of a catalytic core mannanase enzyme in *B.subtilis*:

The mannanase encoding DNA sequence of the invention was PCR amplified using the PCR primer set consisting of the following two oligo nucleotides:

BXM2.upper.SacII

5'-GTT GAG AAA GCG GCC GCC TTT TTT CTA TTC TAC AAT CAC ATT ATC-3'

BXM2.core.lower.NotI

20 5'-GAC GAC GTA CAA GCG GCC GCT CAC TAC GGA GAA GTT CCT CCA TCA G-3'

Restriction sites SacII and NotI are underlined.

Chromosomal DNA isolated from *Bacillus* sp. I633 as described above was used as template in a PCR reaction using
25 Amplitaq DNA Polymerase (Perkin Elmer) according to manufacturers instructions. The PCR reaction was set up in PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.01 % (w/v) gelatin) containing 200 µM of each dNTP, 2.5 units of AmpliTaq polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of
30 each primer.

The PCR reactions was performed using a DNA thermal

cycler (Landgraf, Germany). One incubation at 94°C for 1 min followed by thirty cycles of PCR performed using a cycle profile of denaturation at 94°C for 30 sec, annealing at 60°C for 1 min, and extension at 72 °C for 2 min. Five-µl aliquots of the amplification product was analysed by electrophoresis in 0.7 % agarose gels (NuSieve, FMC). The appearance of a DNA fragment size 1.0 kb indicated proper amplification of the gene segment.

Subcloning of PCR fragment:

Fortyfive-µl aliquots of the PCR products generated as described above were purified using QIAquick PCR purification kit (Qiagen, USA) according to the manufacturer's instructions. The purified DNA was eluted in 50 µl of 10mM Tris-HCl, pH 8.5. 5 µg of pMOL944 and twentyfive-µl of the purified PCR fragment was digested with SacII and NotI, electrophoresed in 0.8 % low gelling temperature agarose (SeaPlaque* GTG, FMC) gels, the relevant fragments were excised from the gels, and purified using QIAquick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the SacII-NotI digested and purified pMOL944. The ligation was performed overnight at 16°C using 0.5 µg of each DNA fragment, 1 U of T4 DNA ligase and T4 ligase buffer (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent B.subtilis PL2306. The transformed cells were plated onto LBPG-10 µg/ml of Kanamycin-agar plates. After 18 hours incubation at 37°C colonies were seen on plates. Several clones were analyzed by isolating plasmid DNA from overnight culture broth.

One such positive clone was restreaked several times on agar plates as used above, this clone was called MB748. The clone MB748 was grown overnight in TY-10µg/ml Kanamycin at 37°C, and next day 1 ml of cells were used to isolate plasmid from the

cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for *B.subtilis* plasmid preparations. This DNA was DNA sequenced and revealed the DNA sequence corresponding to the mature part of the mannanase
 5 (corresponding to positions 91-990 in the appended DNA sequence SEQ ID NO:1 and positions 31-330 in the appended protein sequence SEQ ID NO:2) with introduced stop codon replacing the amino acid residue no 331 corresponding to the base pair positions 1201-1203 in SEQ ID NO:1.

10

B. Subcloning and expression of mature full length mannanase in *B.subtilis*.

The mannanase encoding DNA sequence of the invention was PCR amplified using the PCR primer set consisting of these two
 15 oligo nucleotides:

BXM2.upper.SacII

5'-CAT TCT GCA GCC GCG GCA AAT TCC GGA TTT TAT GTA AGC GG-3'

BXM2.lower.NotI

5'-GTT GAG AAA GCG GCC GCC TTT TTT CTA TTC TAC AAT CAC ATT ATC -
 20 3'

Restriction sites SacII and NotI are underlined

Chromosomal DNA isolated from *Bacillus sp.* (I633) as described above was used as template in a PCR reaction using Amplitaq DNA Polymerase (Perkin Elmer) according to
 25 manufacturers instructions. The PCR reaction was set up in PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.01 % (w/v) gelatin) containing 200 μM of each dNTP, 2.5 units of AmpliTaQ polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of each primer

30

The PCR reactions was performed using a DNA thermal cycler (Landgraf, Germany). One incubation at 94°C for 1 min

followed by thirty cycles of PCR performed using a cycle profile of denaturation at 94°C for 30 sec, annealing at 60°C for 1 min, and extension at 72 °C for 2 min. Five- μ l aliquots of the amplification product was analysed by electrophoresis in 0.7 %

5 agarose gels (NuSieve, FMC) . The appearance of a DNA fragment size 1.5 kb indicated proper amplification of the gene segment.

Subcloning of PCR fragment:

Fortyfive- μ l aliquots of the PCR products generated as described above were purified using QIAquick PCR purification
10 kit (Qiagen, USA) according to the manufacturer's instructions. The purified DNA was eluted in 50 μ l of 10mM Tris-HCl, pH 8.5. 5 μ g of pMOL944 and twentyfive- μ l of the purified PCR fragment was digested with SacII and NotI, electrophoresed in 0.8 % low gelling temperature agarose (SeaPlaque GTG, FMC) gels, the
15 relevant fragments were excised from the gels, and purified using QIAquick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the SacII-NotI digested and purified pMOL944. The ligation was performed overnight at 16°C using 0.5 μ g of
20 each DNA fragment, 1 U of T4 DNA ligase and T4 ligase buffer (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent B.subtilis PL2306. The transformed cells were plated onto LBPG- 10 μ g/ml of Kanamycin-agar plates. After 18 hours incubation at
25 37°C colonies were seen on plates. Several clones were analyzed by isolating plasmid DNA from overnight culture broth.

One such positive clone was restreaked several times on agar plates as used above, this clone was called MB643. The clone MB643 was grown overnight in TY-10 μ g/ml Kanamycin at 37°C,
30 and next day 1 ml of cells were used to isolate plasmid from the cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for B.subtilis plasmid

preparations. This DNA was DNA sequenced and revealed the DNA sequence corresponding to the mature part of the mannanase position 317-1693 in SEQ ID NO. 1 and 33-490 in the SEQ ID NO. 2.

5 The clone MB643 was grown in 25 x 200 ml BPX media with 10 µg/ml of Kanamycin in 500 ml two baffled shakeflasks for 5 days at 37°C at 300 rpm.

The DNA sequence encoding the C-terminal domain of unknown function from amino acid residue no. 341 to amino acid residue
10 no. 490 shows high homology to a domain denoted X18 from a known mannanase. This X18 is found in EMBL entry AB007123 from: Yoshida S., Sako Y., Uchida A.: "Cloning, sequence analysis, and expression in *Escherichia coli* of a gene coding for an enzyme from *Bacillus circulans* K-1 that degrades guar gum" in Biosci.
15 Biotechnol. Biochem. 62:514-520 (1998). This gene codes for the signal peptide (aa 1-34), the catalytic core of a family 5 mannanase (aa 35-335), a linker (aa 336-362) and finally the X18 domain of unknown function (aa 363-516).

This X18 domain is also found in *Bacillus subtilis* beta-
20 mannanase Swiss protein database entry P55278 which discloses a gene coding for a signal peptide (aa 1-26), a catalytic core family 26 mannanase (aa 27-360) and this X18 protein domain of unknown function (aa 361-513); (Cloning and sequencing of beta-mannanase gene from *Bacillus subtilis* NM-39, Mendoza NS ; Arai
25 M ; Sugimoto K ; Ueda M ; Kawaguchi T ; Joson LM , Phillippines. In Biochimica Et Biophysica Acta Vol. 1243, No. 3 pp. 552-554 (1995)).

EXAMPLE 2

30 **Expression, purification and characterisation of mannanase from *Bacillus* sp. I633**

The clone MB748 obtained as described in Example 1 and under Materials and Methods was grown in 25 x 200ml BPX media with 10 µg/ml of Kanamycin in 500ml two baffled shakeflasks for 5 days at 37°C at 300 rpm.

5 4500 ml of the shake flask culture fluid of the clone MB748 was collected and pH was adjusted to 5.6. 100 ml of cationic agent (10% C521) and 180 ml of anionic agent (A130) was added during agitation for flocculation. The flocculated material was separated by centrifugation using a Sorval RC 3B centrifuge at 10 9000 rpm for 20 min at 6°C. The supernatant was clarified using Whatman glass filters GF/D and C and finally concentrated on a filtron with a cut off of 10 kDa.

700 ml of this concentrate was adjusted to pH 7.5 using sodium hydroxide. The clear solution was applied to anion-exchange 15 chromatography using a 1000 ml Q-Sepharose column equilibrated with 50 mmol Tris pH 7.5. The mannanase activity bound was eluted in 1100ml using a sodium chloride gradient. This was concentrated to 440 ml using a Filtron membrane. For obtaining highly pure mannanase the concentrate was passed over a Superdex 20 200column equilibrated with 0.1M sodium acetate, pH 6.0.

The pure enzyme gave a single band in SDS-PAGE with a molecular weight of 34 kDa.

Steady state kinetic using locust bean gum:

25 The assay was carried out using different amounts of the substrate locust bean gum, incubating for 20 min at 40°C at pH 10 in 0.1 M Glycine buffer, followed by the determination of formation of reducing sugars. Glucose was used as standard for calculation of micromole formation of reducing sugar during 30 steady state.

The following data was obtained for the highly purified mannanase of the invention:

KCat of 467 per sec with a standard deviation of 13;
kM of 0.7 with a standard deviation of 0.07.

The computer program grafit by Leatherbarrow from Erithacus Software U.K. was used for calculations. Reducing sugar was
5 determined using the PHBAH method (Lever, M. (1972), A new reaction for colormetric determination of carbohydrates. Anal. Biochem. 47, 273-279.)

The following N-terminal sequence of the purified protein was determined: ANSGFYVSGTTLYDANG.

10 Stability: The mannanase was fully stable between pH 6.0 and 11 after incubation for 2 days at room temperature. The enzyme precipitated at low pH.

The pH activity profile shows that the enzyme is more than 60% active between pH 7.5 and pH 10.

15 Temperature optimum was found to be 50°C at pH 10.

DSC differential scanning calometry gave 66°C as melting point at pH 6.0 in sodium acetate buffer indicating that this mannanase enzyme is thermostable.

Immunological properties: Rabbit polyclonal monospecific
20 serum was raised against the highly purified cloned mannanase using conventional techniques at the Danish company DAKO. The serum formed a nice single precipitate in agarose gels with the crude non purified mannanase of the invention.

25 **EXAMPLE 3**

Use of the enzyme of example 2 in detergents

Using commercial detergents instead of buffer and incubation for 20 minutes at 40°C with 0.2% AZCL-Galactomannan (Megazyme, Australia) from carob degree as described above
30 followed by determination of the formation of blue color, the enzyme obtained as described in example 2 was active in European powder detergent Ariel Futur with 60% relative activity, Euro-

pean liquid detergent Ariel Futur with 80% relative activity, in US Tide powder with 45% relative activity and in US Tide liquid detergent with 37% relative activity to the activity measured in Glycine buffer. In these tests, the detergent concentration was
5 as recommended on the commercial detergent packages and the wash water was tap water having 18 degrees German hardness under European (Ariel Futur) conditions and 9 degree under US conditions (US Tide).

10 **EXAMPLE 4**

Construction and expression of fusion protein between the mannanase of *Bacillus* sp. I633 (example 1 and 2) and a cellulose binding domain (CBD)

The CBD encoding DNA sequence of the CipB gene from
15 *Clostridium thermocellum* strain YS (Poole D M; Morag E; Lamed R; Bayer EA; Hazlewood GP; Gilbert HJ (1992) Identification of the cellulose-binding domain of the cellulosome subunit S1 from *Clostridium thermocellum* YS, Fems Microbiology Letters Vol. 78 , No. 2-3 pp. 181-186 had previously been introduced to a vector
20 pMOL1578. Chromosomal DNA encoding the CBD can be obtained as described in Poole DM; Morag E; Lamed R; Bayer EA; Hazlewood GP Gilbert HJ (1992) Identification of the cellulose-binding domain of the cellulosome subunit S1 from *Clostridium thermocellum* YS, Fems Microbiology Letters Vol. 78 , No. 2-3 pp.
25 181-186. A DNA sample encoding the CBD was used as template in a PCR and the CBD was cloned in an appropriate plasmid pMB993 based on the pMOL944 vector.

The pMB993 vector contains the CipB CBD with a peptide linker preceeding the CBD. The linker consists of the following
30 peptide sequence ASPEPTPEPT and is directly followed by the CipB CBD. The AS aminoacids are derived from the DNA sequence that

constitute the Restriction Endonuclease site *NheI*, which in the following is used to clone the mannase of the invention.

Mannanase.Upper.SacII

5 5'-CAT TCT GCA GCC GCG GCA AAT TCC GGA TTT TAT GTA AGC GG -3'

Mannanase.Lower.NheI

5'-CAT CAT GCT AGC TGT AAA AAC GGT GCT TAA TCT CG -3'

10 Restriction sites *NheI* and *SacII* are underlined.

Chromosomal DNA isolated from *Bacillus* sp. I633 as described above was used as template in a PCR reaction using Amplitaq DNA Polymerase (Perkin Elmer) according to manufacturers instructions. The PCR reaction was set up in PCR buffer (10 mM
15 Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.01 % (w/v) gelatin) containing 200 µM of each dNTP, 2.5 units of AmpliTag polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of each primer.

The PCR reactions was performed using a DNA thermal
cycler (Landgraf, Germany). One incubation at 94°C for 1 min
20 followed by thirty cycles of PCR performed using a cycle profile of denaturation at 94°C for 30 sec, annealing at 60°C for 1 min, and extension at 72 °C for 2 min. Five-µl aliquots of the amplification product was analysed by electrophoresis in 0.7 % agarose gels (NuSieve, FMC). The appearance of a DNA fragment
25 size 0.9 kb indicated proper amplification of the gene segment.

Subcloning of PCR fragment:

Fortyfive-µl aliquots of the PCR products generated as described above were purified using QIAquick PCR purification
30 kit (Qiagen, USA) according to the manufacturer's instructions. The purified DNA was eluted in 50 µl of 10mM Tris-HCl, pH 8.5.

5 µg of pMB993 and twentyfive-µl of the purified PCR fragment was digested with SacII and NheI, electrophoresed in 0.7 % low gelling temperature agarose (SeaPlaque GTG, FMC) gels, the relevant fragments were excised from the gels, and purified using QIAquick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the SacII-NheI digested and purified pMB993. The ligation was performed overnight at 16°C using 0.5 µg of each DNA fragment, 1 U of T4 DNA ligase and T4 ligase buffer (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent *B.subtilis* PL2306. The transformed cells were plated onto LBPG-10 µg/ml of Kanamycin-agar plates. After 18 hours incubation at 37°C colonies were seen on plates. Several clones were analyzed by isolating plasmid DNA from overnight culture broth.

One such positive clone was restreaked several times on agar plates as used above, this clone was called MB1014. The clone MB1014 was grown overnight in TY-10µg/ml Kanamycin at 37°C, and next day 1 ml of cells were used to isolate plasmid from the cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for *B.subtilis* plasmid preparations. This DNA was DNA sequenced and revealed the DNA sequence corresponding to the mature part of the Mannanase-linker-cbd as represented in SEQ ID NO:3 and in the appended protein sequence SEQ ID NO: 4.

Thus the final construction contains the following expression relevant elements: (amyL-promoter)-(amyL-signalpeptide)-mannanase-linker-CBD.

30 Expression and detection of mannanase-CBD fusion protein

MB1014 was incubated for 20 hours in TY-medium at 37°C and 250 rpm. 1 ml of cell-free supernatant was mixed with 200 µl of

10% Avicel* (Merck, Darmstadt, Germany) in Millipore H₂O. The mixture was left for ½ hour incubation at 0°C. After this binding of BXM2-Linker-CBD fusion protein to Avicel the Avicel with bound protein was spun 5 min at 5000g. The pellet was resuspended in 100 µl of SDS-page buffer, boiled at 95°C for 5 min, spun at 5000g for 5 min and 25 µl was loaded on a 4-20% Laemmli Tris-Glycine, SDS-PAGE NOVEX gel (Novex, USA). The samples were electrophoresed in a Xcell™ Mini-Cell (NOVEX, USA) as recommended by the manufacturer, all subsequent handling of gels including staining with comassie, destaining and drying were performed as described by the manufacturer.

The appearance of a protein band of approx. 53 kDa, verified the expression in *B.subtilis* of the full length Mannanase-Linker-CBD fusion encoded on the plasmid pMB1014.

15

EXAMPLE 5

Mannanase derived from *Bacillus agaradhaerens*

Cloning of the mannanase gene from *Bacillus agaradherens*

Genomic DNA preparation

20 Strain *Bacillus agaradherens* NCIMB 40482 was propagated in liquid medium as described in W094/01532. After 16 hours incubation at 30°C and 300 rpm, the cells were harvested, and genomic DNA isolated by the method described by Pitcher et al. (Pitcher, D. G., Saunders, N. A., Owen, R. J. (1989). Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. Lett. Appl. Microbiol., 8, 151-156).

Genomic library construction

Genomic DNA was partially digested with restriction enzyme Sau3A, and size-fractionated by electrophoresis on a 0.7 % agarose gel. Fragments between 2 and 7 kb in size was isolated by electrophoresis onto DEAE-cellulose paper (Dretzen, G.,

*Trade-mark

Bellard, M., Sassone-Corsi, P., Chambon, P. (1981) A reliable method for the recovery of DNA fragments from agarose and acrylamide gels. Anal. Biochem., 112, 295-298).

Isolated DNA fragments were ligated to BamHI digested pSJ1678 plasmid DNA, and the ligation mixture was used to transform *E. coli* SJ2.

Identification of positive clones

A DNA library in *E. coli*, constructed as described above, was screened on LB agar plates containing 0.2% AZCL-galactomannan (Megazyme) and 9 µg/ml Chloramphenicol and incubated overnight at 37°C. Clones expressing mannanase activity appeared with blue diffusion halos. Plasmid DNA from one of these clone was isolated by Qiagen plasmid spin preps on 1 ml of overnight culture broth (cells incubated at 37°C in TY with 9 µg/ml Chloramphenicol and shaking at 250 rpm).

This clone (MB525) was further characterized by DNA sequencing of the cloned Sau3A DNA fragment. DNA sequencing was carried out by primerwalking, using the Taq deoxy-terminal cycle sequencing kit (Perkin-Elmer, USA), fluorescent labelled terminators and appropriate oligonucleotides as primers.

Analysis of the sequence data was performed according to Devereux et al. (1984) Nucleic Acids Res. 12, 387-395. The sequence encoding the mannanase is shown in SEQ ID No 5. The derived protein sequence is shown in SEQ ID No.6.

Subcloning and expression of *B. agaradhaerens* mannanase in *B. subtilis*

The mannanase encoding DNA sequence of the invention was PCR amplified using the PCR primer set consisting of these two oligo nucleotides:

Mannanase.upper.SacII

5'-CAT TCT GCA GCC GCG GCA GCA AGT ACA GGC TTT TAT GTT GAT GG-3'

Mannanase.lower.NotI

5 5'-GAC GAC GTA CAA GCG GCC GCG CTA TTT CCC TAA CAT GAT GAT ATT
TTC G -3'

Restriction sites SacII and NotII are underlined.

Chromosomal DNA isolated from *B.agaradherens* NCIMB 40482 as described above was used as template in a PCR reaction using
10 Amplitaq DNA Polymerase (Perkin Elmer) according to manufacturers instructions. The PCR reaction was set up in PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.01 % (w/v) gelatin) containing 200 µM of each dNTP, 2.5 units of AmpliTaq polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of
15 each primer.

The PCR reaction was performed using a DNA thermal cycler (Landgraf, Germany). One incubation at 94°C for 1 min followed by thirty cycles of PCR performed using a cycle profile of denaturation at 94°C for 30 sec, annealing at 60°C for 1 min,
20 and extension at 72°C for 2 min. Five-µl aliquots of the amplification product was analysed by electrophoresis in 0.7 % agarose gels (NuSieve, FMC). The appearance of a DNA fragment size 1.4 kb indicated proper amplification of the gene segment.

25 Subcloning of PCR fragment

Fortyfive-µl aliquots of the PCR products generated as described above were purified using QIAquick PCR purification kit (Qiagen, USA) according to the manufacturer's instructions. The purified DNA was eluted in 50 µl of 10mM Tris-HCl, pH 8.5.
30 5 µg of pMOL944 and twentyfive-µl of the purified PCR fragment was digested with SacII and NotI, electrophoresed in 0.8% low gelling temperature agarose (SeaPlaque GTG, FMC) gels, the

relevant fragments were excised from the gels, and purified using QIAquick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the SacII-NotI digested and purified pMOL944.

5 The ligation was performed overnight at 16°C using 0.5µg of each DNA fragment, 1 U of T4 DNA ligase and T4 ligase buffer (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent *B.subtilis* PL2306. The transformed cells were plated onto LBPG-
10 10 µg/ml of Kanamycin plates. After 18 hours incubation at 37°C colonies were seen on plates. Several clones were analysed by isolating plasmid DNA from overnight culture broth.

One such positive clone was restreaked several times on agar plates as used above, this clone was called MB594. The
15 clone MB594 was grown overnight in TY-10 µg/ml kanamycin at 37°C, and next day 1 ml of cells were used to isolate plasmid from the cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for *B.subtilis* plasmid preparations. This DNA was DNA sequenced and
20 revealed the DNA sequence corresponding to the mature part of the mannanase, i.e. positions 94-1404 of the appended SEQ ID NO:7. The derived mature protein is shown in SEQ ID NO:8. It will appear that the 3' end of the mannanase encoded by the sequence of SEQ ID NO:5 was changed to the one shown in SEQ ID
25 NO:7 due to the design of the lower primer used in the PCR. The resulting amino acid sequence is shown in SEQ ID NO:8 and it is apparent that the C terminus of the SEQ ID NO:6 (SHHVREIGVQFSAADNSSGQTALYVDNVTLR) is changed to the C terminus of SEQ ID NO:8 (IIMLGK).

30

EXAMPLE 6

Expression, purification and characterisation of mannanase from *Bacillus agaradhaerens*

The clone MB 594 obtained as described in example 5 was grown in 25 x 200ml BPX media with 10 µg/ml of Kanamycin in 5 500ml two baffled shakeflasks for 5 days at 37°C at 300 rpm.

6500 ml of the shake flask culture fluid of the clone MB 594 (batch #9813) was collected and pH adjusted to 5.5. 146 ml of cationic agent (C521) and 292 ml of anionic agent (A130) was added during agitation for flocculation. The flocculated material was separated by centrifugation using a Sorval RC 3B centrifuge at 9000 rpm for 20 min at 6°C. The supernatant was clarified using Whatman glass filters GF/D and C and finally concentrated on a filtron with a cut off of 10 kDa.

750 ml of this concentrate was adjusted to pH 7.5 using sodium hydroxide. The clear solution was applied to anion-exchange chromatography using a 900 ml Q-Sepharose column equilibrated with 50 mmol Tris pH 7.5. The mannanase activity bound was eluted using a sodium chloride gradient.

The pure enzyme gave a single band in SDS-PAGE with a 20 molecular weight of 38 kDa.

The amino acid sequence of the mannanase enzyme, i.e. the translated DNA sequence, is shown in SEQ ID No.6.

Determination of kinetic constants:

Substrate: Locust bean gum (carob) and reducing sugar analysis (PHBAH). Locust bean gum from Sigma (G-0753).

Kinetic determination using different concentrations of locust bean gum and incubation for 20 min at 40°C at pH 10 gave

K_{cat}: 467 per sec.

K_m: 0.08 gram per l

30 MW: 38kDa

pI (isoelectric point): 4.2

The temperature optimum of the mannanase was found to be 60°C.

The pH activity profile showed maximum activity between pH 8 and 10.

5 DSC differential scanning calorimetry gives 77°C as melting point at pH 7.5 in Tris buffer indicating that this enzyme is very thermostable.

Detergent compatibility using 0.2% AZCL-Galactomannan from carob as substrate and incubation as described above at 40°C.
10 shows excellent compability with conventional liquid detergents and good compability with conventional powder detergents.

EXAMPLE 7

Use of the enzyme of the invention in detergents

15 The purified enzyme obtained as described in example 6 (batch #9813) showed improved cleaning performance when tested at a level of 1 ppm in a miniwash test using a conventional commercial liquid detergent. The test was carried out under conventional North American wash conditions.

20

EXAMPLE 8

Mannanase derived from *Bacillus* sp. AAI12

Construction of a genomic library from *Bacillus* sp. AAI12

25 Genomic DNA of *Bacillus* sp. was partially digested with restriction enzyme Sau3A, and size-fractionated by electrophoresis on a 0.7 % agarose gel (SeaKem agarose, FMC, USA). Fragments between 1.5 and 10 kb in size were isolated and concentrated to a DNA band by running the DNA fragments backwards
30 on a 1.5 % agarose gel followed by extraction of the fragments from the agarose gel slice using the Qiaquick gel extraction kit according to the manufacturer's instructions (Qiagen Inc., USA).

To construct a genomic library, ca. 100ng of purified, fractionated DNA from above was ligated with 1 ug of BamHI-cleaved, dephosphorylated lambdaZAPexpress vector arms (Stratagene, La Jolla CA, USA) for 24 hours at + 4 °C according to the manufacturer's instructions. A 3-ul aliquot of the ligation mixture was packaged directly using the GigaPackIII Gold packaging extract (Stratagene, USA) according to the manufacturers instructions (Stratagene). The genomic lambdaZAPExpress phage library was titered using the E. coli XL1-Blue MRF- strain from Stratagene (La Jolla, USA). The unamplified genomic library comprised of 7.8×10^7 plaque-forming units (pfu) with a vector background of less than 1 %.

Screening for beta-mannanase clones by functional expression in lambdaZAPExpress

Approximately 5000 plaque-forming units (pfu) from the genomic library were plated on NZY-agar plates containing 0.1 % AZCL-galactomannan (MegaZyme, Australia, cat. no. I-AZGMA), using E. coli XL1-Blue MRF' (Stratagene, USA) as a host, followed by incubation of the plates at 37 °C for 24 hours. Mannanase-positive lambda clones were identified by the formation of blue hydrolysis halos around the positive phage clones. These were recovered from the screening plates by coring the TOP-agar slices containing the plaques of interest into 500 ul of SM buffer and 20 ul of chloroform. The mannanase-positive lambdaZAPExpress clones were plaque-purified by plating an aliquot of the cored phage stock on NZY plates containing 0.1 % AZCL-galactomannan as above. Single, mannanase-positive lambda clones were cored into 500 ul of SM buffer and 20 ul of chloroform, and purified by one more plating round as described above.

Single-clone in vivo excision of the phagemids from the mannase-positive lambdaZAPExpress clones

E. coli XL1-Blue cells (Stratagene, La Jolla Ca.) were prepared and resuspended in 10mM MgSO₄ as recommended by
5 Stratagene (La Jolla, USA). 250-ul aliquots of the pure phage stocks from the mannase-positive clones were combined in Falcon 2059 tubes with 200uls of XL1-Blue MRF' cells (OD₆₀₀=1.0) and > 106 pfus/ml of the ExAssist M13 helper phage (Stratagene), and the mixtures were incubated at 37 C for 15 minutes. Three mls of
10 NZY broth was added to each tube and the tubes were incubated at 37 C for 2.5 hours. The tubes were heated at 65 C for 20 minutes to kill the E. coli cells and bacteriophage lambda; the phagemids being resistant to heating. The tubes were spun at 3000 rpm for 15 minutes to remove cellular debris and the super-
15 natants were decanted into clean Falcon 2059 tubes. Aliquots of the supernatants containing the excised single-stranded phagemids were used to infect 200uls of E. coli XL0LR cells (Stratagene, OD₆₀₀=1.0 in 10mM MgSO₄) by incubation at 37°C for 15 minutes. 350uls of NZY broth was added to the cells and the
20 tubes were incubated for 45 min at 37°C. Aliquots of the cells were plated onto LB kanamycin agar plates and incubated for 24 hours at 37°C. Five excised single colonies were re-streaked onto LB kanamycin agar plates containing 0.1 % AZCL-galactomannan (MegaZyme, Australia). The mannase-positive
25 phagemid clones were characterized by the formation of blue hydrolysis halos around the positive colonies. These were further analysed by restriction enzyme digests of the isolated phagemid DNA (QiaSpin kit, Qiagen, USA) with EcoRI, PstI, EcoRI-PstI, and HindIII followed by agarose gel electrophoresis.

30

Nucleotide sequence analysis

The nucleotide sequence of the genomic beta-1,4-mannanase clone pBXM1 was determined from both strands by the dideoxy chain-termination method (Sanger, F., Nicklen, S., and Coulson, A. R. (1977) Proc. Natl. Acad. Sci. U. S. A. 74, 5463-5467) using 500 ng of Qiagen-purified template (Qiagen, USA), the Taq deoxy-terminal cycle sequencing kit (Perkin-Elmer, USA), fluorescent labeled terminators and 5 pmol of either pBK-CMV polylinker primers (Stratagene, USA) or synthetic oligonucleotide primers. Analysis of the sequence data was performed according to Devereux et al., 1984 (Devereux, J., Haeberli, P., and Smithies, O. (1984) Nucleic Acids Res. 12, 387-395).

Sequence alignment

A multiple sequence alignment of the glycohydrolase family 26 beta-1,4-mannanases from *Bacillus* sp. AAI 12 of the present invention (ie SEQ ID NO: 10), *Caldicellulosiruptor saccharolyticus* (GenBank/EMBL accession no. P77847), *Dictyoglomus thermophilum* (acc. no. 030654), *Rhodothermus marinus* (acc. no. P49425), *Piromyces* sp. encoded by ManA (acc. no. P55296), *Bacillus* sp. (acc. no. P91007), *Bacillus subtilis* (acc. no. 005512) and *Pseudomonas fluorescens* (acc. no P49424). was created using the PileUp program of the GCG Wisconsin software package, version 8.1. (see above); with gap creation penalty 3.00 and gap extension penalty 0.10.

Sequence Similarities

The deduced amino acid sequence of the family 26 beta-1,4-mannanase of the invention cloned from *Bacillus* sp. AAI 12 shows 45 % sequence similarity and 19.8 % sequence identity to the beta-1,4-mannanase from *Caldicellulosiruptor saccharolyticus* (GenBank/EMBL accession no. P77847), 49 % similarity and 25.1. %

identity to the beta-1,4-mannanase from *Dictyoglomus thermophilum* (acc. no. 030654), 48.2 % similarity and 26.8 % identity to the beta-1,4-mannanase from *Rhodothermus marinus* (acc. no. P49425), 46 % similarity and 19.5 % sequence identity to the ManA-encoded beta-1,4-mannanase from *Piromyces* sp. (acc. no. P55296), 47.2 % similarity and 22 % identity to the beta-1,4-mannanase from *Bacillus* sp. (acc. no. P91007), 52.4 % similarity and 27.5 % sequence identity to the beta-1,4-mannanase from *Bacillus subtilis* (acc. no. 005512) and 60.6 % similarity and 37.4 % identity to the beta-1,4-mannanase from *Pseudomonas fluorescens* (acc. no P49424. The sequences were aligned using the GAP program of the GCG Wisconsin software package, version 8.1.; with gap creation penalty 3.00 and gap extension penalty 0.10.

15

Cloning of the *Bacillus* sp (AAI 12) mannanase gene

Subcloning and expression of mannanase in *B.subtilis*

The mannanase encoding DNA sequence of the invention was PCR amplified using the PCR primer set consisting of these two oligo nucleotides:

BXM1.upper.SacII

5' - CAT TCT GCA GCC GCG GCA TTT TCT GGA AGC GTT TCA GC-3'

BXM1.lower.NotI

5'-CAG CAG TAG CGG CCG CCA CTT CCT GCT GGT ACA TAT GC -3'

25 Restriction sites SacII and NotI are underlined.

Chromosomal DNA isolated from *Bacillus* sp. AAI 12 as described above was used as template in a PCR reaction using Amplitaq DNA Polymerase (Perkin Elmer) according to manufacturers instructions. The PCR reaction was set up in PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.01 % (w/v) gelatin) containing 200 µM of each dNTP, 2.5 units of

30

AmpliTaq polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of each primer.

The PCR reactions was performed using a DNA thermal cyclor (Landgraf, Germany). One incubation at 94°C for 1 min
5 followed by thirty cycles of PCR performed using a cycle profile of denaturation at 94°C for 30 sec, annealing at 60°C for 1 min, and extension at 72 °C for 2 min. Five-µl aliquots of the amplification product was analysed by electrophoresis in 0.7 % agarose gels (NuSieve, FMC). The appearance of a DNA fragment
10 size 1.0 kb indicated proper amplification of the gene segment.

Subcloning of PCR fragment

Fortyfive-µl aliquots of the PCR products generated as described above were purified using QIAquick PCR purification
15 kit (Qiagen, USA) according to the manufacturer's instructions. The purified DNA was eluted in 50 µl of 10mM Tris-HCl, pH 8.5. 5 µg of pMOL944 and twentyfive-µl of the purified PCR fragment was digested with SacII and NotI, electrophoresed in 0.8 % low gelling temperature agarose (SeaPlaque GTG, FMC) gels, the
20 relevant fragments were excised from the gels, and purified using QIAquick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the SacII-NotI digested and purified pMOL944. The ligation was performed overnight at 16°C using 0.5 µg of
25 each DNA fragment, 1 U of T4 DNA ligase and T4 ligase buffer (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent B.subtilis PL2306. The transformed cells were plated onto LBPG-
10 µg/ml of Kanamycin-agar plates. After 18 hours incubation at
30 37°C colonies were seen on plates. Several clones were analyzed by isolating plasmid DNA from overnight culture broth.

One such positive clone was restreaked several times on agar plates as used above, this clone was called MB747. The clone MB747 was grown overnight in TY-10µg/ml Kanamycin at 37°C, and next day 1 ml of cells were used to isolate plasmid from the 5 cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for B.subtilis plasmid preparations. This DNA was DNA sequenced and revealed the DNA sequence corresponding to the mature part of the mannanase in the SEQ ID NO. 9.

10

Expression, purification and characterisation of mannanase from *Bacillus sp.* AAI 12

The clone MB747 obtained as described above was grown in 25 x 200ml BPX media with 10 µg/ml of Kanamycin in 500ml two baffled shakeflasks for 5 days at 37°C at 300 rpm. 15

4100 ml of the shake flask culture fluid of the clone MB747 was collected, pH was adjusted to 7.0, and EDTA was added to a final concentration of 2mM. 185 ml of cationic agent (10% C521) and 370 ml of anionic agent (A130) was added during agitation 20 for flocculation. The flocculated material was separated by centrifugation using a Sorval RC 3B centrifuge at 9000 rpm for 20 min at 6°C. The supernatant was clarified using Whatman glass filters GF/D and C and finally concentrated on a filtron with a cut off of 10 kDa.

25 1500 ml of this concentrate was adjusted to pH 7.5 using sodium hydroxide. The clear solution was applied to anion-exchange chromatography using a 1000 ml Q-Sepharose column equilibrated with 25 mmol Tris pH 7.5. The mannanase activity bound was eluted in 1100ml using a sodium chloride gradient. 30 This was concentrated to 440 ml using a Filtron membrane. For obtaining highly pure mannanase the concentrate was passed over a Superdex column equilibrated with 0.1M sodium acetate, pH 6.0.

The pure enzyme gave a single band in SDS-PAGE with a molecular weight of 62 kDa.

The amino acid sequence of the mannanase enzyme, i.e. the translated DNA sequence, is shown in SEQ ID No.10.

5 The following N-terminal sequence was determined:
FSGSVSASGQELKMTDQN.

pI (isoelectric point): 4.5

DSC differential scanning calorimetry gave 64°C as melting point at pH 6.0 in sodium acetate buffer indicating that this
10 mannanase enzyme is thermostable.

It was found that the catalytic activity increases with ionic strength indicating that the specific activity of the enzyme may be increased by using salt of phosphate buffer with high ionic strength.

15 The mannanase activity of the polypeptide of the invention is inhibited by calcium ions.

Immunological properties: Rabbit polyclonal monospecific serum was raised against the highly purified mannanase of the invention using conventional techniques at the Danish company DAKO.
20 The serum formed a nice single precipitate in agarose gels with the crude mannanase of the invention.

EXAMPLE 9

25 Use of the enzyme of example 8 in detergents

Using commercial detergents instead of buffer and incubation for 20 minutes at 40°C with 0.2% AZCL-Galactomannan (Megazyme, Australia) from carob degree as described above followed by determination of the formation of blue color, the
30 enzyme obtained as described in example 8 was active in European powder detergent Ariel Futur with 132% relative activity, in US Tide powder with 108% relative activity and in US Tide liquid

detergent with 86% relative activity to the activity measured in Glycine buffer. In these tests, the detergent concentration was as recommended on the commercial detergent packages and the wash water was tap water having 18 degrees German hardness under European (Ariel Futur) conditions and 9 degree under US conditions (US Tide).

EXAMPLE 10

Mannanase derived from *Bacillus halodurans*

10

Construction of a genomic library from *Bacillus halodurans* in the pSJ1678 vector

Genomic DNA of *Bacillus halodurans* was partially digested with restriction enzyme *Sau3A*, and size-fractionated by electrophoresis on a 0.7 % agarose gel (SeaKem agarose, FMC, USA). DNA fragments between 2 and 10 kb in size was isolated by electrophoresis onto DEAE-cellulose paper (Dretzen, G., Bellard, M., Sassone-Corsi, P., Chambon, P. (1981) A reliable method for the recovery of DNA fragments from agarose and acrylamide gels. Anal. Biochem., 112, 295-298). Isolated DNA fragments were ligated to BamHI-digested pSJ1678 plasmid DNA, and the ligation mixture was used to transform *E. coli* SJ2.

Screening for beta-mannanase clones by functional expression in *Escherichia coli*

Approximately 10.000 colony-forming units (cfu) from the genomic library were plated on LB-agar plates containing containing 9 µg/ml chloramphenicol and 0.1 % AZCL-galactomannan (MegaZyme, Australia, cat. no. I-AZGMA), using *E. coli* SJ2 as a host, followed by incubation of the plates at 37°C for 24 hours. Mannanase-positive *E. coli* colonies were identified by the formation of blue hydrolysis halos around the positive plasmid

clones. The mannanase-positive clones in pSJ1678 were colony-purified by re-streaking the isolated colonies on LB plates containing 9 μ g/ml Chloramphenicol and 0.1 % AZCL-galactomannan as above. Single, mannanase-positive plasmid clones were inoculated into 5 ml of LB medium containing 9 μ g/ml Chloramphenicol, for purification of the plasmid DNA.

Nucleotide sequence analysis

The nucleotide sequence of the genomic beta-1,4-mannanase clone pBXM5 was determined from both strands by the dideoxy chain-termination method (Sanger, F., Nicklen, S., and Coulson, A. R. (1977) Proc. Natl. Acad. Sci. U. S. A. 74, 5463-5467) using 500 ng of Qiagen-purified template (Qiagen, USA), the Taq deoxy-terminal cycle sequencing kit (Perkin-Elmer, USA), fluorescent labeled terminators and 5 pmol of either pBK-CMV polylinker primers (Stratagene, USA) or synthetic oligonucleotide primers. Analysis of the sequence data was performed according to Devereux et al., 1984 (Devereux, J., Haeberli, P., and Smithies, O. (1984) Nucleic Acids Res. 12, 387-395).

Sequence alignment

A multiple sequence alignment of the glycohydrolase family beta-1,4-mannanase from *Bacillus halodurans* of the present invention (ie SEQ ID NO:12), *Bacillus circulans* (GenBank/EMBL accession no. 066185), *Vibrio sp.* (acc. no. 069347), *Streptomyces lividans* (acc. no. P51529), and *Caldicellulosiruptor saccharolyticus* (acc. no. P22533). The multiple sequence alignment was created using the PileUp program of the GCG Wisconsin software package, version 8.1.; with gap creation penalty 3.00 and gap extension penalty 0.10.

Sequence Similarities

The deduced amino acid sequence of the family 5 beta-1,4-mannanase of the present invention cloned from *Bacillus halodurans* shows 77% similarity and 60% sequence identity to the 5 beta-1,4-mannanase of *Bacillus circulans* (GenBank/EMBL accession no. O66185), 64.2% similarity and 46% identity to the beta-1,4-mannanase from *Vibrio* sp. (acc. no. O69347), 63% similarity and 41.8% identity to the beta-1,4-mannanase from *Streptomyces lividans* (acc. no. P51529), 60.3% similarity and 42% sequence 10 identity to the beta-1,4-mannanase from *Caldicellulosiruptor saccharolyticus* (acc. no. P2253). The sequences were aligned using the GAP program of the GCG Wisconsin software package, version 8.1.; with gap creation penalty 3.00 and gap extension penalty 0.10.

15

Cloning of *Bacillus halodurans* mannanase gene

Subcloning and expression of mature full length mannanase in *B. subtilis*

The mannanase encoding DNA sequence of the invention was 20 PCR amplified using the PCR primer set consisting of these two oligo nucleotides:

BXM5.upper.SacII

5'-CAT TCT GCA GCC GCG GCA CAT CAC AGT GGG TTC CAT G-3'

25

BXM5.lower.NotI

5'-GCG TTG AGA CGC GCG GCC GCT TAT TGA AAC ACA CTG CTT CTT TTA G-3'

Restriction sites SacII and NotI are underlined

30

Chromosomal DNA isolated from *Bacillus halodurans* as described above was used as template in a PCR reaction using

Amplitaq DNA Polymerase (Perkin Elmer) according to manufacturers instructions. The PCR reaction was set up in PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.01 % (w/v) gelatin) containing 200 μ M of each dNTP, 2.5 units of 5 AmpliTag polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of each primer.

The PCR reactions was performed using a DNA thermal cycler (Landgraf, Germany). One incubation at 94°C for 1 min followed by thirty cycles of PCR performed using a cycle profile 10 of denaturation at 94°C for 30 sec, an-nea-ling at 60°C for 1 min, and extension at 72°C for 2 min. Five- μ l aliquots of the ampli-fication product was analysed by electrophoresis in 0.7 % agarose gels (NuSieve, FMC). The appearance of a DNA fragment size 0.9 kb indicated proper amplification of the gene segment.

15

Subcloning of PCR fragment:

Fortyfive- μ l aliquots of the PCR products generated as described above were purified using QIA-quick PCR purification kit (Qiagen, USA) according to the manufacturer's instructions. The 20 purified D-NA was eluted in 50 μ l of 10mM Tris-HCl, pH 8.5.

5 μ g of pMOL944 and twentyfive- μ l of the purified PCR fragment was digested with SacII and NotI, electrophoresed in 0.8 % low gelling temperature agarose (SeaPlaque GTG, FMC) gels, the relevant fragments were excised from the gels, and purified 25 using QIA-quick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the SacII-NotI digested and purified pMOL944. The ligation was performed overnight at 16°C using 0.5 μ g of each DNA fragment, 1 U of T4 DNA ligase and T4 ligase 30 buffer (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent B.subtilis PL2306. The transformed cells were plated onto LBPG-

10 $\mu\text{g/ml}$ of Kanamycin-agar plates. After 18 hours incubation at 37°C colonies were seen on plates. Several clones were analyzed by isolating plasmid DNA from overnight culture broth.

One such positive clone was restreaked several times on
5 agar plates as used above, this clone was called MB878. The clone MB878 was grown overnight in TY- $10\mu\text{g/ml}$ Kanamycin at 37°C , and next day 1 ml of cells were used to isolate plasmid from the cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for *B.subtilis* plasmid
10 preparations. This DNA was DNA sequenced and revealed the DNA sequence corresponding to the mature part of the mannanase position 97-993 in SEQ ID NO. 11 and 33-331 in the SEQ ID NO. 12.

15 **Expression, purification and characterisation of mannanase from *Bacillus halodurans***

The clone MB878 obtained as described above was grown in 25 x 200ml BPX media with $10\mu\text{g/ml}$ of Kanamycin in 500ml two baffled shakeflasks for 5 days at 37°C at 300 rpm.

20 5000 ml of the shake flask culture fluid of the clone MB878 was collected and pH was adjusted to 6.0. 125 ml of cationic agent (10% C521) and 250 ml of anionic agent (A130) was added during agitation for flocculation. The flocculated material was separated by centrifugation using a Sorval RC 3B centrifuge at
25 9000 rpm for 20 min at 6°C . The supernatant was adjusted to pH 8.0 using NaOH and clarified using Whatman glass filters GF/D and C. Then 50 g of DEAE A-50 Sephadex was equilibrated with 0.1M Sodium acetate, pH 6.0, and added to the filtrate, the enzyme was bound and left overnight at room temperature. The
30 bound enzyme was eluted with 0.5 M NaCl in the acetate buffer. Then the pH was adjusted to pH 8.0 using sodium hydroxide and then concentrated on a Filtron with a 10 kDa cut off to 450 ml

and then stabilized with 20% glycerol, 20% MPG and 2% Berol. The product was used for application trials.

2 ml of this concentrate was adjusted to pH 8.5 using sodium hydroxide. For obtaining highly pure mannanase the concentrate was passed over a Superdex column equilibrated with 0.1 M sodium phosphate, pH 8.5.

The pure enzyme gave a single band in SDS-PAGE with a molecular weight of 34 kDa.

The amino acid sequence of the mannanase enzyme, i.e. the translated DNA sequence, is shown in SEQ ID NO:12.

The following N-terminal sequence of the purified protein was determined: AHHS GFHVNGTTLYDA.

The pH activity profile using the ManU assay (incubation for 20 minutes at 40°C) shows that the enzyme has a relative activity higher than 50% between pH 7.5 and pH 10.

Temperature optimum was found (using the ManU assay; glycine buffer) to be between 60°C and 70°C at pH 10.

Immunological properties: Rabbit polyclonal monospecific serum was raised against the highly purified cloned mannanase using conventional techniques at the Danish company DAKO. The serum formed a nice single precipitate in agarose gels with the crude non purified mannanase of the invention.

EXAMPLE 11

25 Use of the mannanase enzyme of example 10 in detergents

Using commercial detergents instead of buffer and incubation for 20 minutes at 40°C with 0.2% AZCL-Galactomannan (Megazyme, Australia) from carob degree as described above followed by determination of the formation of blue color, the mannanase enzyme obtained as described in example 10 was active with an activity higher than 40% relative to the activity in buffer in European liquid detergent Ariel Futur, in US Tide

powder and in US Tide liquid detergent. In these tests, the detergent concentration was as recommended on the commercial detergent packages and the wash water was tap water having 18 degrees German hardness under European (Ariel Futur) conditions 5 and 9 degree under US conditions (US Tide).

EXAMPLE 12

Mannanase derived from *Bacillus sp.* AA349

10 Cloning of *Bacillus sp.* (AA349) mannanase gene

Subcloning and expression of a catalytic core mannanase enzyme in *B. subtilis*:

The mannanase encoding DNA sequence of the invention was PCR amplified using the PCR primer set consisting of the follow-
15 ing two oligo nucleotides:

BXM7.upper.SacII

5'-CAT TCT GCA GCC GCG GCA AGT GGA CAT GGG CAA ATG C-3'

BXM7.lower.NotI

5'-GCG TTG AGA CGC GCG GCC GCT TAT TTT TTG TAT ACA CTA ACG ATT
20 TC-3'

Restriction sites SacII and NotI are underlined.

Chromosomal DNA isolated from *Bacillus sp.* AA349 as described above was used as template in a PCR reaction using Amplitaq DNA Polymerase (Perkin Elmer) according to
25 manufacturers instructions. The PCR reaction was set up in PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.01 % (w/v) gelatin) containing 200 µM of each dNTP, 2.5 units of AmpliTaq polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of each primer.

30 The PCR reactions was performed using a DNA thermal cycler (Landgraf, Germany). One incubation at 94°C for 1 min

followed by thirty cycles of PCR performed using a cycle profile of denaturation at 94°C for 30 sec, annealing at 60°C for 1 min, and extension at 72 °C for 2 min. Five-µl aliquots of the amplification product was analysed by electrophoresis in 0.7 % agarose gels (NuSieve, FMC). The appearance of a DNA fragment approximate size of 1.0 kb indicated proper amplification of the gene segment.

Subcloning of PCR fragment:

Fortyfive-µl aliquots of the PCR products generated as described above were purified using QIAquick PCR purification kit (Qiagen, USA) according to the manufacturer's instructions. The purified DNA was eluted in 50 µl of 10mM Tris-HCl, pH 8.5. 5 µg of pMOL944 and twentyfive-µl of the purified PCR fragment was digested with SacII and NotI, electrophoresed in 0.8 % low gelling temperature agarose (SeaPlaque GTG, FMC) gels, the relevant fragments were excised from the gels, and purified using QIAquick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the SacII-NotI digested and purified pMOL944. The ligation was performed overnight at 16°C using 0.5 µg of each DNA fragment, 1 U of T4 DNA ligase and T4 ligase buffer (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent B.subtilis PL2306. The transformed cells were plated onto LBPG-10 µg/ml of Kanamycin-agar plates. After 18 hours incubation at 37°C colonies were seen on plates. Several clones were analyzed by isolating plasmid DNA from overnight culture broth.

One such positive clone was restreaked several times on agar plates as used above, this clone was called MB879. The clone MB879 was grown overnight in TY-10µg/ml Kanamycin at 37°C, and next day 1 ml of cells were used to isolate plasmid from the

cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for B.subtilis plasmid preparations. This DNA was DNA sequenced and revealed the DNA sequence corresponding to the mature part of the mannanase
5 (corresponding to positions 204-1107 in the appended DNA sequence SEQ ID NO:15 and positions 26-369 in the appended protein sequence SEQ ID NO:16.

Expression, purification and characterisation of mannanase from
10 ***Bacillus* sp. AA349**

The clone MB879 obtained as described above was grown in 25 x 200ml BPX media with 10 µg/ml of Kanamycin in 500ml two baffled shakeflasks for 5 days at 37°C at 300 rpm.

400 ml of the shake flask culture fluid of the clone MB879
15 was collected and pH was 6.5. 19 ml of cationic agent (10% C521) and 38 ml of anionic agent (A130) was added during agitation for flocculation. The flocculated material was separated by centrifugation using a Sorval RC 3B centrifuge at 5000 rpm for 25 min at 6°C. The then concentrated and washed with water to
20 reduce the conductivity on a Filtron with a 10 kDa cut off to 150 ml. then the pH was adjusted to 4.0 and the liquid applied to S-Sepharose column chromatography in a 50 mM Sodium acetate buffer pH 4.0. The column was first eluted with a NaCl gradient to 0.5 M then the mannanase eluted using 0.1 M glycine buffer pH
25 10. The mannanase active fraction was pooled and they gave a single band in SDS-PAGE with a molecular weight of 38 kDa.

The amino acid sequence of the mannanase enzyme, i.e. the translated DNA sequence, is shown in SEQ ID NO:16.

The pH activity profile using the ManU assay (incubation
30 for 20 minutes at 40°C) shows that the enzyme has a relative activity higher than 30% between pH 5 and pH 10.

Temperature optimum was found (using the ManU assay; gly-

cine buffer) to be between 60°C and 70°C at pH 10.

Immunological properties: Rabbit polyclonal monospecific serum was raised against the highly purified cloned mannanase using conventional techniques at the Danish company DAKO. The
5 serum formed a nice single precipitate in agarose gels with the crude non purified mannanase of the invention.

EXAMPLE 13

Use of the mannanase enzyme of example 12 in detergents

10 Using commercial detergents instead of buffer and incubation for 20 minutes at 40°C with 0.2% AZCL-Galactomannan (Megazyme, Australia) from carob degree as described above followed by determination of the formation of blue color, the mannanase enzyme obtained as described in example 12 was active
15 with an activity higher than 65% relative to the activity in buffer in European liquid detergent Ariel Futur and in US Tide liquid detergent. The mannanase was more than 35% active in powder detergents from Europe, Ariel Futur and in US tide powder. In these tests, the detergent concentration was as recommended on the commercial detergent packages and the wash water
20 was tap water having 18 degrees German hardness under European (Ariel Futur) conditions and 9 degree under US conditions (US Tide).

25 EXAMPLE 14

Mannanase derived from the fungal strain *Humicola insolens* DSM 1800

Expression cloning of a family 26 beta-1,4-mannanase from *Humicola insolens*
30

Fungal strain and cultivation conditions

Humicola insolens strain DSM 1800 was fermented as described in WO 97/32014, the mycelium was harvested after 5 days growth at 26 °C, immediately frozen in liquid N₂, and stored at - 80 °C..

5

Preparation of RNase-free glassware, tips and solutions

All glassware used in RNA isolations were baked at + 220 °C for at least 12 h. Eppendorf tubes, pipet tips and plastic columns were treated in 0.1 % diethylpyrocarbonate (DEPC) in EtOH for 12 h, and autoclaved. All buffers and water (except Tris-containing buffers) were treated with 0.1 % DEPC for 12 h at 37 °C, and autoclaved.

Extraction of total RNA

15 The total RNA was prepared by extraction with guanidinium thiocyanate followed by ultracentrifugation through a 5.7 M CsCl cushion (Chirgwin et al., 1979) using the following modifications. The frozen mycelia was ground in liquid N₂ to fine powder with a mortar and a pestle, followed by grinding in a precooled
20 coffee mill, and immediately suspended in 5 vols of RNA extraction buffer (4 M GuSCN, 0.5 % Na-laurylsarcosine, 25 mM Na-citrate, pH 7.0, 0.1 M β-mercaptoethanol). The mixture was stirred for 30 min. at RT° and centrifuged (30 min., 5000 rpm, RT°, Heraeus Megafuge 1.0 R) to pellet the cell debris. The
25 supernatant was collected, carefully layered onto a 5.7 M CsCl cushion (5.7 M CsCl, 0.1 M EDTA, pH 7.5, 0.1 % DEPC; autoclaved prior to use) using 26.5 ml supernatant per 12.0 ml CsCl cushion, and centrifuged to obtain the total RNA (Beckman, SW 28 rotor, 25 000 rpm, RT°, 24h). After centrifugation the supernatant
30 was carefully removed and the bottom of the tube containing the RNA pellet was cut off and rinsed with 70 % EtOH. The total RNA pellet was transferred into an Eppendorf tube, sus-

pended in 500 ml TE, pH 7.6 (if difficult, heat occasionally for 5 min at 65 °C), phenol extracted and precipitated with ethanol for 12 h at - 20 °C (2.5 vols EtOH, 0.1 vol 3M NaAc, pH 5.2). The RNA was collected by centrifugation, washed in 70 % EtOH, and resuspended in a minimum volume of DEPC-DIW. The RNA concentration was determined by measuring OD_{260/280}.

Isolation of poly(A)⁺RNA

The poly(A)⁺RNAs were isolated by oligo(dT)-cellulose affinity chromatography (Aviv & Leder, 1972). Typically, 0.2 g of oligo(dT) cellulose (Boehringer Mannheim, check for binding capacity) was preswollen in 10 ml of 1 x column loading buffer (20 mM Tris-Cl, pH 7.6, 0.5 M NaCl, 1 mM EDTA, 0.1 % SDS), loaded onto a DEPC-treated, plugged plastic column (Poly Prep Chromatography Column, Bio Rad), and equilibrated with 20 ml 1 x loading buffer. The total RNA was heated at 65 °C for 8 min., quenched on ice for 5 min, and after addition of 1 vol 2 x column loading buffer to the RNA sample loaded onto the column. The eluate was collected and reloaded 2-3 times by heating the sample as above and quenching on ice prior to each loading. The oligo(dT) column was washed with 10 vols of 1 x loading buffer, then with 3 vols of medium salt buffer (20 mM Tris-Cl, pH 7.6, 0.1 M NaCl, 1 mM EDTA, 0.1 % SDS), followed by elution of the poly(A)⁺ RNA with 3 vols of elution buffer (10 mM Tris-Cl, pH 7.6, 1 mM EDTA, 0.05 % SDS) preheated to + 65 °C, by collecting 500 ml fractions. The OD₂₆₀ was read for each collected fraction, and the mRNA containing fractions were pooled and ethanol precipitated at - 20 °C for 12 h. The poly(A)⁺ RNA was collected by centrifugation, resuspended in DEPC-DIW and stored in 5-10 mg aliquots at - 80 °C.

cDNA synthesis**First strand synthesis**

Double-stranded cDNA was synthesized from 5 mg of *Humicola insolens* poly(A)⁺ RNA by the RNase H method (Gubler & Hoffman 1983, Sambrook et al., 1989) using the hair-pin modification developed by F. S. Hagen. The poly(A)⁺RNA (5 mg in 5 ml of DEPC-treated water) was heated at 70°C for 8 min., quenched on ice, and combined in a final volume of 50 ml with reverse transcriptase buffer (50 mM Tris-Cl, pH 8.3, 75 mM KCl, 3 mM MgCl₂, 10 mM DTT, Bethesda Research Laboratories) containing 1 mM each dNTP (Pharmacia), 40 units of human placental ribonuclease inhibitor (RNasin, Promega), 10 mg of oligo(dT)₁₂₋₁₈ primer (Pharmacia) and 1000 units of SuperScript II RNase H⁻ reverse transcriptase (Bethesda Research Laboratories). First-strand cDNA was synthesized by incubating the reaction mixture at 45 °C for 1 h.

Second strand synthesis

After synthesis 30 ml of 10 mM Tris-Cl, pH 7.5, 1 mM EDTA was added, and the mRNA:cDNA hybrids were ethanol precipitated for 12 h at - 20 °C by addition of 40 mg glycogen carrier (Boehringer Mannheim) 0.2 vols 10 M NH₄Ac and 2.5 vols 96 % EtOH. The hybrids were recovered by centrifugation, washed in 70 % EtOH, air dried and resuspended in 250 ml of second strand buffer (20 mM Tris-Cl, pH 7.4, 90 mM KCl, 4.6 mM MgCl₂, 10 mM (NH₄)₂SO₄, 16 mM βNAD⁺) containing 100 mM each dNTP, 44 units of E. coli DNA polymerase I (Amersham), 6.25 units of RNase H (Bethesda Research Laboratories) and 10.5 units of E. coli DNA ligase (New England Biolabs). Second strand cDNA synthesis was performed by incubating the reaction tube at 16 °C for 3 h, and the reaction was stopped by addition of EDTA to 20 mM final concentration followed by phenol extraction.

Mung bean nuclease treatment

The double-stranded (ds) cDNA was ethanol precipitated at -20°C for 12 h by addition of 2 vols of 96 % EtOH, 0.1 vol 3 M NaAc, pH 5.2, recovered by centrifugation, washed in 70 % EtOH, dried (SpeedVac), and resuspended in 30 ml of Mung bean nuclease buffer (30 mM NaAc, pH 4.6, 300 mM NaCl, 1 mM ZnSO₄, 0.35 mM DTT, 2 % glycerol) containing 36 units of Mung bean nuclease (Bethesda Research Laboratories). The single-stranded hair-pin DNA was clipped by incubating the reaction at 30 °C for 30 min, followed by addition of 70 ml 10 mM Tris-Cl, pH 7.5, 1 mM EDTA, phenol extraction, and ethanol precipitation with 2 vols of 96 % EtOH and 0.1 vol 3M NaAc, pH 5.2 at -20 °C for 12 h.

15 Blunt-ending with T4 DNA polymerase

The ds cDNA was blunt-ended with T4 DNA polymerase in 50 ml of T4 DNA polymerase buffer (20 mM Tris-acetate, pH 7.9, 10 mM MgAc, 50 mM KAc, 1 mM DTT) containing 0.5 mM each dNTP and 7.5 units of T4 DNA polymerase (Invitrogen) by incubating the reaction mixture at + 37 °C for 15 min. The reaction was stopped by addition of EDTA to 20 mM final concentration, followed by phenol extraction and ethanol precipitation.

Adaptor ligation and size selection

25 After the fill-in reaction the cDNA was ligated to non-palindromic BstX I adaptors (1 mg/ml, Invitrogen) in 30 ml of ligation buffer (50 mM Tris-Cl, pH 7.8, 10 mM MgCl₂, 10 mM DTT, 1 mM ATP, 25 mg/ml bovine serum albumin) containing 600 pmol BstX I adaptors and 5 units of T4 ligase (Invitrogen) by incubating the reaction mix at + 16 °C for 12 h. The reaction was stopped by heating at + 70 °C for 5 min, and the adapted cDNA was size-fractionated by agarose gel electrophoresis (0.8 % HSB-

agarose, FMC) to separate unligated adaptors and small cDNAs. The cDNA was size-selected with a cut-off at 0.7 kb, and the cDNA was electroeluted from the agarose gel in 10 mM Tris-Cl, pH 7.5, 1 mM EDTA for 1 h at 100 volts, phenol extracted and ethanol precipitated at - 20 °C for 12 h as above.

Construction of the *Humicola insolens* cDNA library

The adapted, ds cDNAs were recovered by centrifugation, washed in 70 % EtOH and resuspended in 25 ml DIW. Prior to large-scale library ligation, four test ligations were carried out in 10 ml of ligation buffer (same as above) each containing 1 ml ds cDNA (reaction tubes #1 - #3), 2 units of T4 ligase (Invitrogen) and 50 ng (tube #1), 100 ng (tube #2) and 200 ng (tubes #3 and #4) Bst XI cleaved pYES 2.0 vector (Invitrogen). The ligation reactions were performed by incubation at + 16 °C for 12 h, heated at 70 °C for 5 min, and 1 ml of each ligation electroporated (200 W, 2.5 kV, 25 mF) to 40 ml competent *E. coli* 1061 cells (OD600 = 0.9 in 1 liter LB-broth, washed twice in cold DIW, once in 20 ml of 10 % glycerol, resuspended in 2 ml 10 % glycerol). After addition of 1 ml SOC to each transformation mix, the cells were grown at + 37 °C for 1 h, 50 ml plated on LB + ampicillin plates (100 mg/ml) and grown at + 37 °C for 12h.

Using the optimal conditions a large-scale ligation was set up in 40 ml of ligation buffer containing 9 units of T4 ligase, and the reaction was incubated at + 16°C for 12 h. The ligation reaction was stopped by heating at 70°C for 5 min, ethanol precipitated at - 20°C for 12 h, recovered by centrifugation and resuspended in 10 ml DIW. One ml aliquots were transformed into electrocompetent *E. coli* 1061 cells using the same electroporation conditions as above, and the transformed cells were titered and the library plated on LB + ampicillin plates with 5000-7000 c.f.u./plate. The cDNA library, comprising of 1×10^6 recombi-

nant clones, was stored as 1) individual pools (5000-7000 c.f.u./pool) in 20 % glycerol at - 80°C, 2) cell pellets of the same pools at - 20°C, and 3) Qiagen purified plasmid DNA from individual pools at - 20°C (Qiagen Tip 100, Qiagen).

5

Expression cloning in *Saccharomyces cerevisiae* of beta-1,4 mannanase cDNAs from *Humicola insolens*

One ml aliquots of purified plasmid DNA (100 ng/ml) from individual pools were electroporated (200 W, 1.5 kV, 25 mF) into 40 ml of electrocompetent *S. cerevisiae* W3124 (MATa; ura 3-52; leu 2-3, 112; his 3-D200; pep 4-1137; prc1::HIS3, prb1::LEU2; cir+) cells (OD600 = 1.5 in 500 ml YPD, washed twice in cold DIW, once in cold 1 M sorbitol, resuspended in 0.5 ml 1 M sorbitol, Becker & Guarante, 1991). After addition of 1 ml 1M cold sorbitol, 80 ml aliquots were plated on SC + glucose - uracil to give 250-400 colony forming units per plate and incubated at 30°C for 3 - 5 days. The plates were replicated on SC + galactose - uracil plates, containing AZCl-galactomannan (MegaZyme, Australia) incorporated in the agar plates. In total, ca. 50 000 yeast colonies from the *H. insolens* library were screened for mannanase-positive clones.

The positive clones were identified by the formation of blue hydrolysis halos around the corresponding yeast colonies. The clones were obtained as single colonies, the cDNA inserts were amplified directly from yeast cell lysates using biotinylated pYES 2.0 polylinker primers, purified by magnetic beads (Dynabead M-280, Dynal) system and characterized individually by sequencing the 5'-end of each cDNA clone using the chain-termination method (Sanger et al., 1977) and the Sequenase system (United States Biochemical).

The mannanase-positive yeast colonies were inoculated into 20 ml YPD broth in a 50 ml tubes. The tubes were shaken for 2

days at 30°C, and the cells were harvested by centrifugation for 10 min. at 3000 rpm. Total yeast DNA was isolated according to WO 94/14953, dissolved in 50 ml of autoclaved water, and transformed into *E. coli* by electroporation as above. The insert-
5 containing pYES 2.0 cDNA clones were rescued by plating on LB + ampicillin agar plates, the plasmid DNA was isolated from *E. coli* using standard procedures, and analyzed by digesting with restriction enzymes.

10 Nucleotide sequence analysis

The nucleotide sequence of the full-length *H. insolens* beta-1,4-mannanase cDNA clone pC1M59 was determined from both strands by the dideoxy chain-termination method (Sanger et al. 1977), using 500 ng of Qiagen-purified template (Qiagen, USA)
15 template, the Taq deoxy-terminal cycle sequencing kit (Perkin-Elmer, USA), fluorescent labeled terminators and 5 pmol of the pYES 2.0 polylinker primers (Invitrogen, USA). Analysis of the sequence data were performed according to Devereux et al. (1984).

20

Heterologous expression in *Aspergillus oryzae*

Transformation of *Aspergillus oryzae*

Transformation of *Aspergillus oryzae* was carried out as described by Christensen et al., (1988), Biotechnology 6, 1419-
25 1422.

Construction of the beta-1,4-mannanase expression cassette for *Aspergillus* expression

Plasmid DNA was isolated from the mannanase clone pC1M59
30 using standard procedures and analyzed by restriction enzyme analysis. The cDNA insert was excised using appropriate restriction enzymes and ligated into the *Aspergillus* expression vector

pHD414, which is a derivative of the plasmid p775 (described in EP 238023). The construction of pHD414 is further described in WO 93/11249.

5 **Transformation of *Aspergillus oryzae* or *Aspergillus niger***

General procedure: 100 ml of YPD (Sherman et al., Methods in Yeast Genetics, Cold Spring Harbor Laboratory, 1981) is inoculated with spores of *A. oryzae* or *A. niger* and incubated with shaking at 37°C for about 2 days. The mycelium is harvested
10 by filtration through miracloth and washed with 200 ml of 0.6 M MgSO₄. The mycelium is suspended in 15 ml of 1.2 M MgSO₄. 10 mM NaH₂PO₄, pH = 5.8. The suspension is cooled on ice and 1 ml of buffer containing 120 mg of Novozym® 234 is added. After 5 minutes 1 ml of 12 mg/ml BSA is added and incubation with gentle
15 agitation continued for 1.5-2.5 hours at 37°C until a large number of protoplasts is visible in a sample inspected under the microscope. The suspension is filtered through miracloth, the filtrate transferred to a sterile tube and overlaid with 5 ml of 0.6 M sorbitol, 100 mM Tris-HCl, pH = 7.0. Centrifugation is
20 performed for 15 minutes at 100 g and the protoplasts are collected from the top of the MgSO₄ cushion. 2 volumes of STC are added to the protoplast suspension and the mixture is centrifuged for 5 minutes at 1000 g. The protoplast pellet is resuspended in 3 ml of STC and repelleted. This is repeated. Finally
25 the protoplasts are resuspended in 0.2-1 ml of STC. 100 µl of protoplast suspension is mixed with 5-25 µg of the appropriate DNA in 10 µl of STC. Protoplasts are mixed with p3SR2 (an *A. nidulans* amdS gene carrying plasmid). The mixture is left at room temperature for 25 minutes. 0.2 ml of 60% PEG 4000. 10 mM
30 CaCl₂ and 10 mM Tris-HCl, pH 7.5 is added and carefully mixed (twice) and finally 0.85 ml of the same solution is added and carefully mixed. The mixture is left at room temperature for 25

minutes, spun at 2500 g for 15 minutes and the pellet is resuspended in 2 ml of 1.2 M sorbitol. After one more sedimentation the protoplasts are spread on the appropriate plates. Protoplasts are spread on minimal plates to inhibit background growth. After incubation for 4-7 days at 37°C spores are picked and spread for single colonies. This procedure is repeated and spores of a single colony after the second re-isolation is stored as a defined transformant.

10 **Purification of the *Aspergillus oryzae* transformants**

Aspergillus oryzae colonies are purified through conidial spores on AmdS+-plates (+ 0,01% Triton X-100) and growth in YPM for 3 days at 30°C.

15 **Identification of mannanase-positive *Aspergillus oryzae* transformants**

The supernatants from the *Aspergillus oryzae* transformants were assayed for beta-1,4-mannanase activity on agar plates containing 0.2 % AZCl-galactomannan (MegaZyme, Australia) as substrate. Positive transformants were identified by analyzing the plates for blue hydrolysis halos after 24 hours of incubation at 30°C.

SDS-PAGE analysis

SDS-PAGE analysis of supernatants from beta-1,4-mannanase producing *Aspergillus oryzae* transformants. The transformants were grown in 5 ml YPM for three days. 10 µl of supernatant was applied to 12% SDS-polyacrylamide gel which was subsequently stained with Coomassie Brilliant Blue.

Purification and characterisation of the *Humicola insolens* mannanase

The gene was transformed into *A. oryzae* ads described above and the transformed strain was grown in a fermentor using standard medium of Maltose syrup, sucrose, MgSO_4 , K_2PO_4 and K_2SO_4 and citric acid yeast extract and trace metals. Incubation for 6 days at 34°C with air.

The fermentation broth (5000 ml) was harvested and the mycelium separated from the liquid by filtration. The clear liquid was concentrated on a filtron to 275 ml.

The mannanase was purified using Cationic chromatography. A S-Spharose column was equilibrated with 25 mM citric acid pH 4.0 and the mannanase bound to the column and was eluted using a sodium chloride gradient (0-0.5 M). The mannanase active fractions was pooled and the pH adjusted to 7.3. The 100 ml pooled mannanase was then concentrated to 5 ml with around 13 mg protein per ml and used for applications trials. For further purification 2 ml was applied to size chromatography on Superdex 200 in sodium acetate buffer pH 6.1. The mannanase active fraction showed to equal stained bands in SDS-PAGE with a MW of 45 kDa and 38 kDa, indicating proteolytic degradation of the N-terminal non-catalytic domain.

The amino acid sequence of the mannanase enzyme, i.e. the translated DNA sequence, is shown in SEQ ID NO:14.

The DNA sequence of SEQ ID NO:13 codes for a signal peptide in positions 1 to 21. A domain of unknown function also found in other mannanases is represented in the amino acid sequence SEQ ID NO: 14 in positions 22 to 159 and the catalytic active domain is found in positions 160 to 488 of SEQ ID NO:14.

Highest sequence homology was found to *DICTYOGLOMUS THERMOPHILUM* (49% identity); Mannanase sequence EMBL; AF013989 submitted by REEVES R.A., GIBBS M.D., BERGQUIST P.L. submitted in July 1997.

Molecular Weight: 38 kDa.

DSC in sodium acetate buffer pH 6.0 was 65°.

The pH activity profile using the ManU assay (incubation for 20 minutes at 40°C) shows that the enzyme has optimum activity at pH 8.

Temperature optimum was found (using the ManU assay; Megazyme AZCL locust bean gum as substrate) to be 70°C at pH 10.

Immunological properties: Rabbit polyclonal monospecific serum was raised against the highly purified cloned mannanase using conventional techniques at the Danish company DAKO. The serum formed a nice single precipitate in agarose gels with the crude non purified mannanase of the invention.

EXAMPLE 15

15 Wash evaluation of Humicola Insolens family 26 mannanase

Wash performance was evaluated by washing locust bean gum coated swatches in a detergent solution with the mannanase of the invention. After wash the effect were visualised by soiling the swatches with iron oxide.

20 Preparation of locust bean gum swatches: Clean cotton swatches were soaked in a solution of 2 g/l locust bean gum and dried overnight at room temperature. The swatches were prewashed in water and dried again.

Wash: Small circular locust bean gum swatches were placed in a beaker with 6,7 g/l Ariel Futur liquid in 15°dH water and incubated for 30 min at 40°C with magnetic stirring. The swatches were rinsed in tap water and dried.

Soiling: The swatches were placed in a beaker with 0.25 g/l Fe₂O₃ and stirred for 3 min. The swatches were rinsed in tap water and dried.

Evaluation: Remission of the swatches was measured at 440 nm using a MacBeth ColorEye 7000 remission spectrophotometer.

The results are expressed as

$$\text{delta remission} = (R_{\text{after wash}} - R_{\text{before wash}})_{\text{enzyme}} - (R_{\text{after wash}} - R_{\text{before wash}})_{\text{control}}$$

where R is the remission at 440 nm.

5 The mannanase of this invention is clearly effective on locust bean gum swatches with a wash performance slightly better than the control mannanase from *Bacillus sp.* I633.

Wash performance of *Humicola insolens* family 26 mannanase compared to the mannanase from *Bacillus sp.* I633 (examples 1-3) given as delta remission values:

Enzyme dose in mg/l	<i>Humicola insolens</i> mannanase	<i>Bacillus sp.</i> I633mannanase
0	0	0
0.01	6.6	5.0
0.1	9.3	8.6
1.0	10.2	7.7
10.0	10.5	9.7

EXAMPLES 16-40

The following examples are meant to exemplify compositions of the present invention, but are not necessarily meant to limit or otherwise define the scope of the invention.

In the detergent compositions, the enzymes levels are expressed by pure enzyme by weight of the total composition and unless otherwise specified, the detergent ingredients are expressed by weight of the total compositions. The abbreviated component identifications therein have the following meanings:

LAS : Sodium linear C₁₁₋₁₃ alkyl benzene sulphate.

TAS : Sodium tallow alkyl sulphate.

- CxyAS : Sodium C_{1x} - C_{1y} alkyl sulfate.
- CxySAS : Sodium C_{1x} - C_{1y} secondary (2,3) alkyl sulfate.
- CxyEz : C_{1x} - C_{1y} predominantly linear primary alcohol condensed with an average of z moles of ethylene oxide.
- CxyEzS : C_{1x} - C_{1y} sodium alkyl sulfate condensed with an average of z moles of ethylene oxide.
- QAS : R₂.N+(CH₃)₂(C₂H₄OH) with R₂ = C₁₂-C₁₄.
- QAS 1 : R₂.N+(CH₃)₂(C₂H₄OH) with R₂ = C₈-C₁₁.
- APA : C₈-10 amido propyl dimethyl amine.
- Soap : Sodium linear alkyl carboxylate derived from a 80/20 mixture of tallow and coconut fatty acids.
- Nonionic : C₁₃-C₁₅ mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5.
- Neodol 45-13 : C₁₄-C₁₅ linear primary alcohol ethoxylate, sold by Shell Chemical CO.
- STS : Sodium toluene sulphonate.
- CFAA : C₁₂-C₁₄ alkyl N-methyl glucamide.
- TFAA : C₁₆-C₁₈ alkyl N-methyl glucamide.
- TPKFA : C₁₂-C₁₄ topped whole cut fatty acids.
- Silicate : Amorphous Sodium Silicate (SiO₂:Na₂O ratio = 1.6-3.2).
- Metasilicate : Sodium metasilicate (SiO₂:Na₂O ratio = 1.0).

- Zeolite A : Hydrated Sodium Aluminosilicate of formula $\text{Na}_{12}(\text{AlO}_2\text{SiO}_2)_{12} \cdot 27\text{H}_2\text{O}$ having a primary particle size in the range from 0.1 to 10 micrometers (Weight expressed on an anhydrous basis).
- Na-SKS-6 : Crystalline layered silicate of formula $\delta\text{-Na}_2\text{Si}_2\text{O}_5$.
- Citrate : Tri-sodium citrate dihydrate of activity 86.4% with a particle size distribution between 425 and 850 micrometres.
- Citric : Anhydrous citric acid.
- Borate : Sodium borate
- Carbonate : Anhydrous sodium carbonate with a particle size between 200 and 900 micrometres.
- Bicarbonate : Anhydrous sodium hydrogen carbonate with a particle size distribution between 400 and 1200 micrometres.
- Sulphate : Anhydrous sodium sulphate.
- Mg Sulphate : Anhydrous magnesium sulfate.
- STPP : Sodium tripolyphosphate.
- TSPP : Tetrasodium pyrophosphate.
- MA/AA : Random copolymer of 4:1 acrylate/maleate, average molecular weight about 70,000-80,000.
- MA/AA 1 : Random copolymer of 6:4 acrylate/maleate, average molecular weight about 10,000.
- AA : Sodium polyacrylate polymer of average molecular weight 4,500.
- PA30 : Polyacrylic acid of average molecular weight of between about 4,500 - 8,000.

- 480N : Random copolymer of 7:3 acrylate/methacrylate, average molecular weight about 3,500.
- Polygel/carbopo 1 : High molecular weight crosslinked polyacrylates.
- PB1 : Anhydrous sodium perborate monohydrate of nominal formula $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$.
- PB4 : Sodium perborate tetrahydrate of nominal formula $\text{NaBO}_2 \cdot 3\text{H}_2\text{O} \cdot \text{H}_2\text{O}_2$.
- Percarbonate : Anhydrous sodium percarbonate of nominal formula $2\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}_2$.
- NaDCC : Sodium dichloroisocyanurate.
- TAED : Tetraacetylenediamine.
- NOBS : Nonanoyloxybenzene sulfonate in the form of the sodium salt.
- NACA-OBS : (6-nonamidocaproyl) oxybenzene sulfonate.
- DTPA : Diethylene triamine pentaacetic acid.
- HEDP : 1,1-hydroxyethane diphosphonic acid.
- DETPMP : Diethyltriamine penta (methylene) phosphonate, marketed by Monsanto under the Trade name Dequest 2060.
- EDDS : Ethylenediamine-N,N'-disuccinic acid, (S,S) isomer in the form of its sodium salt
- MnTACN : Manganese 1,4,7-trimethyl-1,4,7-triazacyclononane.
- Photoactivated Bleach : Sulfonated zinc phthalocyanine encapsulated in dextrin soluble polymer.
- Photoactivated Bleach 1 : Sulfonated alumino phthalocyanine encapsulated in dextrin soluble polymer.
- PAAC : Pentaamine acetate cobalt(III) salt.

- Paraffin : Paraffin oil sold under the tradename Winog 70 by Wintershall.
- NaBz : Sodium benzoate.
- BzP : Benzoyl Peroxide.
- Mannanase : As described herein
- Protease : Proteolytic enzyme sold under the trade-name Savinase, Alcalase, Durazym by Novo Nordisk A/S, Maxacal, Maxapem sold by Gist-Brocades and proteases described in patents WO91/06637 and/or WO95/10591 and/or EP 251 446.
- Amylase : Amylolytic enzyme sold under the tradename Purafact Ox Am^R described in WO 94/18314, WO96/05295 sold by Genencor; Termamyl[®], Fungamyl[®] and Duramyl[®], all available from Novo Nordisk A/S and those described in WO95/26397.
- Lipase : Lipolytic enzyme sold under the tradename Lipolase, Lipolase Ultra by Novo Nordisk A/S and Lipomax by Gist-Brocades.
- Cellulase : Cellulytic enzyme sold under the tradename Carezyme, Celluzyme and/or Endolase by Novo Nordisk A/S.
- CMC : Sodium carboxymethyl cellulose.
- PVP : Polyvinyl polymer, with an average molecular weight of 60,000.
- PVNO : Polyvinylpyridine-N-Oxide, with an average molecular weight of 50,000.
- PVPVI : Copolymer of vinylimidazole and vinylpyrrolidone, with an average molecular weight of 20,000.

- Brightener 1 : Disodium 4,4'-bis(2-sulphostyryl)biphenyl.
- Brightener 2 : Disodium 4,4'-bis(4-anilino-6-morpholino-1.3.5-triazin-2-yl) stilbene-2:2'-disulfonate.
- Silicone anti-foam : Polydimethylsiloxane foam controller with siloxane-oxyalkylene copolymer as dispersing agent with a ratio of said foam controller to said dispersing agent of 10:1 to 100:1.
- Suds Suppressor : 12% Silicone/silica, 18% stearyl alcohol, 70% starch in granular form.
- Opacifier : Water based monostyrene latex mixture, sold by BASF Aktiengesellschaft under the tradename Lytron 621.
- SRP 1 : Anionically end capped poly esters.
- SRP 2 : Diethoxylated poly (1,2 propylene terephthalate) short block polymer.
- QEA : $\text{bis}((\text{C}_2\text{H}_5\text{O})(\text{C}_2\text{H}_4\text{O})_n)(\text{CH}_3) - \text{N}^+ - \text{C}_6\text{H}_{12} - \text{N}^+ - (\text{CH}_3) \text{bis}((\text{C}_2\text{H}_5\text{O}) - (\text{C}_2\text{H}_4\text{O}))_n$, wherein $n =$ from 20 to 30.
- PEI : Polyethyleneimine with an average molecular weight of 1800 and an average ethoxylation degree of 7 ethyleneoxy residues per nitrogen.
- SCS : Sodium cumene sulphonate.
- HMWPEO : High molecular weight polyethylene oxide.
- PEGx : Polyethylene glycol, of a molecular weight of x .
- PEO : Polyethylene oxide, with an average molecular weight of 5,000.
- TEPAE : Tetraethylenepentaamine ethoxylate.

WO 99/64619

147

PCT/DK99/00314

BTA : Benzotriazole.

PH : Measured as a 1% solution in distilled
water at 20°C.

WO 99/64619

148

PCT/DK99/00314

Example 16

The following high density laundry detergent compositions were prepared according to the present invention :

	I	II	III	IV	V	VI
LAS	8.0	8.0	8.0	2.0	6.0	6.0
TAS	-	0.5	-	0.5	1.0	0.1
C46(S)AS	2.0	2.5	-	-	-	-
C25AS	-	-	-	7.0	4.5	5.5
C68AS	2.0	5.0	7.0	-	-	-
C25E5	-	-	3.4	10.0	4.6	4.6
C25E7	3.4	3.4	1.0	-	-	-
C25E3S	-	-	-	2.0	5.0	4.5
QAS	-	0.8	-	-	-	-
QAS 1	-	-	-	0.8	0.5	1.0
Zeolite A	18.1	18.0	14.1	18.1	20.0	18.1
Citric	-	-	-	2.5	-	2.5
Carbonate	13.0	13.0	27.0	10.0	10.0	13.0
Na-SKS-6	-	-	-	10.0	-	10.0
Silicate	1.4	1.4	3.0	0.3	0.5	0.3
Citrate	-	1.0	-	3.0	-	-
Sulfate	26.1	26.1	26.1	6.0	-	-
Mg sulfate	0.3	-	-	0.2	-	0.2
MA/AA	0.3	0.3	0.3	4.0	1.0	1.0
CMC	0.2	0.2	0.2	0.2	0.4	0.4
PB4	9.0	9.0	5.0	-	-	-
Percarbonate	-	-	-	-	18.0	18.0
TAED	1.5	0.4	1.5	-	3.9	4.2
NACA-OBS	-	2.0	1.0	-	-	-
DETPMP	0.25	0.25	0.25	0.25	-	-
SRP 1	-	-	-	0.2	-	0.2

WO 99/64619

149

PCT/DK99/00314

	I	II	III	IV	V	VI
EDDS	-	0.25	0.4	-	0.5	0.5
CFAA	-	1.0	-	2.0	-	-
HEDP	0.3	0.3	0.3	0.3	0.4	0.4
QEA	-	-	-	0.2	-	0.5
Protease	0.009	0.009	0.01	0.04	0.05	0.03
Mannanase	0.05	0.009	0.03	0.009	0.03	0.009
Amylase	0.002	0.002	0.002	0.006	0.008	0.008
Cellulase	0.0007	-	-	0.0007	0.0007	0.0007
Lipase	0.006	-	-	0.01	0.01	0.01
Photoactivated bleach (ppm)	15	15	15	-	20	20
PVNO/PVPVI	-	-	-	0.1	-	-
Brightener 1	0.09	0.09	0.09	-	0.09	0.09
Perfume	0.3	0.3	0.3	0.4	0.4	0.4
Silicone anti- foam	0.5	0.5	0.5	-	0.3	0.3
Density in g/litre	850	850	850	850	850	850
Miscellaneous and minors	Up to 100%					

Example 17

The following granular laundry detergent compositions of particular utility under European machine wash conditions were prepared according to the present invention :

5

	I	II	III	IV	V	VI
LAS	5.5	7.5	5.0	5.0	6.0	7.0
TAS	1.25	1.9	-	0.8	0.4	0.3
C24AS/C25AS	-	2.2	5.0	5.0	5.0	2.2
C25E3S	-	0.8	1.0	1.5	3.0	1.0
C45E7	3.25	-	-	-	-	3.0
TFAA	-	-	2.0	-	-	-
C25E5	-	5.5	-	-	-	-
QAS	0.8	-	-	-	-	-
QAS 1	-	0.7	1.0	0.5	1.0	0.7
STPP	19.7	-	-	-	-	-
Zeolite A	-	19.5	25.0	19.5	20.0	17.0
NaSKS-6/citric acid (79:21)	-	10.6	-	10.6	-	-
Na-SKS-6	-	-	9.0	-	10.0	10.0
Carbonate	6.1	21.4	9.0	10.0	10.0	18.0
Bicarbonate	-	2.0	7.0	5.0	-	2.0
Silicate	6.8	-	-	0.3	0.5	-
Citrate	-	-	4.0	4.0	-	-
Sulfate	39.8	-	-	5.0	-	12.0
Mg sulfate	-	-	0.1	0.2	0.2	-
MA/AA	0.5	1.6	3.0	4.0	1.0	1.0
CMC	0.2	0.4	1.0	1.0	0.4	0.4
PB4	5.0	12.7	-	-	-	-
Percarbonate	-	-	-	-	18.0	15.0
TAED	0.5	3.1	-	-	5.0	-

WO 99/64619

151

PCT/DK99/00314

	I	II	III	IV	V	VI
NACA-OBS	1.0	3.5	-	-	-	2.5
DETPMP	0.25	0.2	0.3	0.4	-	0.2
HEDP	-	0.3	-	0.3	0.3	0.3
QEA	-	-	1.0	1.0	1.0	-
Protease	0.009	0.03	0.03	0.05	0.05	0.02
Mannanase	0.03	0.03	0.001	0.03	0.005	0.009
Lipase	0.003	0.003	0.006	0.006	0.006	0.004
Cellulase	0.000	0.000	0.000	0.000	0.000	0.000
	6	6	5	5	7	7
Amylase	0.002	0.002	0.006	0.006	0.01	0.003
PVNO/PVPVI	-	-	0.2	0.2	-	-
PVP	0.9	1.3	-	-	-	0.9
SRP 1	-	-	0.2	0.2	0.2	-
Photoactivated bleach (ppm)	15	27	-	-	20	20
Photoactivated bleach 1 (ppm)	15	-	-	-	-	-
Brightener 1	0.08	0.2	-	-	0.09	0.15
Brightener 2	-	0.04	-	-	-	-
Perfume	0.3	0.5	0.4	0.3	0.4	0.3
Silicone anti- foam	0.5	2.4	0.3	0.5	0.3	2.0
Density in g/litre	750	750	750	750	750	750
Miscellaneous and minors	Up to 100%					

WO 99/64619

152

PCT/DK99/00314

Example 18

The following detergent compositions of particular utility under European machine wash conditions were prepared according to the present invention :

5

	I	II	III	IV
Blown Powder				
LAS	6.0	5.0	11.0	6.0
TAS	2.0	-	-	2.0
Zeolite A	24.0	-	-	20.0
STPP	-	27.0	24.0	-
Sulfate	4.0	6.0	13.0	-
MA/AA	1.0	4.0	6.0	2.0
Silicate	1.0	7.0	3.0	3.0
CMC	1.0	1.0	0.5	0.6
Brightener 1	0.2	0.2	0.2	0.2
Silicone antifoam	1.0	1.0	1.0	0.3
DETPMP	0.4	0.4	0.2	0.4
Spray On				
Brightener	0.02	-	-	0.02
C45E7	-	-	-	5.0
C45E2	2.5	2.5	2.0	-
C45E3	2.6	2.5	2.0	-
Perfume	0.5	0.3	0.5	0.2
Silicone antifoam	0.3	0.3	0.3	-
Dry additives				
QEA	-	-	-	1.0
EDDS	0.3	-	-	-
Sulfate	2.0	3.0	5.0	10.0
Carbonate	6.0	13.0	15.0	14.0
Citric	2.5	-	-	2.0

WO 99/64619

153

PCT/DK99/00314

	I	II	III	IV
QAS 1	0.5	-	-	0.5
Na-SKS-6	10.0	-	-	-
Percarbonate	18.5	-	-	-
PB4	-	18.0	10.0	21.5
TAED	2.0	2.0	-	2.0
NACA-OBS	3.0	2.0	4.0	-
Protease	0.03	0.03	0.03	0.03
Mannanase	0.009	0.01	0.03	0.001
Lipase	0.008	0.008	0.008	0.004
Amylase	0.003	0.003	0.003	0.006
Brightener 1	0.05	-	-	0.05
Miscellaneous and minors			Up to 100%	

WO 99/64619

154

PCT/DK99/00314

Example 19

The following granular detergent compositions were prepared according to the present invention :

	I	II	III	IV	V	VI
Blown Powder						
LAS	23.0	8.0	7.0	9.0	7.0	7.0
TAS	-	-	-	-	1.0	-
C45AS	6.0	6.0	5.0	8.0	-	-
C45AES	-	1.0	1.0	1.0	-	-
C45E35	-	-	-	-	2.0	4.0
Zeolite A	10.0	18.0	14.0	12.0	10.0	10.0
MA/AA	-	0.5	-	-	-	2.0
MA/AA 1	7.0	-	-	-	-	-
AA	-	3.0	3.0	2.0	3.0	3.0
Sulfate	5.0	6.3	14.3	11.0	15.0	19.3
Silicate	10.0	1.0	1.0	1.0	1.0	1.0
Carbonate	15.0	20.0	10.0	20.7	8.0	6.0
PEG 4000	0.4	1.5	1.5	1.0	1.0	1.0
DTPA	-	0.9	0.5	-	-	0.5
Brightener 2	0.3	0.2	0.3	-	0.1	0.3
Spray On						
C45E7	-	2.0	-	-	2.0	2.0
C25E9	3.0	-	-	-	-	-
C23E9	-	-	1.5	2.0	-	2.0
Perfume	0.3	0.3	0.3	2.0	0.3	0.3
Agglomerates						
C45AS	-	5.0	5.0	2.0	-	5.0
LAS	-	2.0	2.0	-	-	2.0
Zeolite A	-	7.5	7.5	8.0	-	7.5

WO 99/64619		155					PCT/DK99/00314
	I	II	III	IV	V	VI	
Carbonate	-	4.0	4.0	5.0	-	4.0	
PEG 4000	-	0.5	0.5	-	-	0.5	
Misc (Water etc.)	-	2.0	2.0	2.0	-	2.0	
Dry additives							
QAS	-	-	-	-	1.0	-	
Citric	-	-	-	-	2.0	-	
PB4	-	-	-	-	12.0	1.0	
PB1	4.0	1.0	3.0	2.0	-	-	
Percarbonate	-	-	-	-	2.0	10.0	
Carbonate	-	5.3	1.8	-	4.0	4.0	
NOBS	4.0	-	6.0	-	-	0.6	
Methyl cellu- lose	0.2	-	-	-	-	-	
Na-SKS-6	8.0	-	-	-	-	-	
STS	-	-	2.0	-	1.0	-	
Culmene sulfo- nic acid	-	1.0	-	-	-	2.0	
Protease	0.02	0.02	0.02	0.01	0.02	0.02	
Mannanase	0.009	0.01	0.03	0.009	0.01	0.001	
Lipase	0.004	-	0.004	-	0.004	0.008	
Amylase	0.003	-	0.002	-	0.003	-	
Cellulase	0.0005	0.0005	0.000	0.000	0.000	0.000	
			5	7	5	5	
PVPVI	-	-	-	-	0.5	0.1	
PVP	-	-	-	-	0.5	-	
PVNO	-	-	0.5	0.3	-	-	
QEA	-	-	-	-	1.0	-	
SRP 1	0.2	0.5	0.3	-	0.2	-	

WO 99/64619

156

PCT/DK99/00314

	I	II	III	IV	V	VI
Silicone anti-foam	0.2	0.4	0.2	0.4	0.1	-
Mg sulfate	-	-	0.2	-	0.2	-
Miscellaneous and minors				Up to 100%		

Example 20

The following nil bleach-containing detergent compositions of particular use in the washing of colored clothing were prepared according to the present invention :

5

	I	II	III
Blown Powder			
Zeolite A	15.0	15.0	-
Sulfate	-	5.0	-
LAS	3.0	3.0	-
DETPMP	0.4	0.5	-
CMC	0.4	0.4	-
MA/AA	4.0	4.0	-
Agglomerates			
C45AS	-	-	11.0
LAS	6.0	5.0	-
TAS	3.0	2.0	-
Silicate	4.0	4.0	-
Zeolite A	10.0	15.0	13.0
CMC	-	-	0.5
MA/AA	-	-	2.0
Carbonate	9.0	7.0	7.0
Spray-on			
Perfume	0.3	0.3	0.5
C45E7	4.0	4.0	4.0
C25E3	2.0	2.0	2.0
Dry additives			
MA/AA	-	-	3.0
Na-SKS-6	-	-	12.0
Citrate	10.0	-	8.0
Bicarbonate	7.0	3.0	5.0

WO 99/64619

158

PCT/DK99/00314

	I	II	III
Carbonate	8.0	5.0	7.0
PVPVI/PVNO	0.5	0.5	0.5
Protease	0.03	0.02	0.05
Mannanase	0.001	0.004	0.03
Lipase	0.008	0.008	0.008
Amylase	0.01	0.01	0.01
Cellulase	0.001	0.001	0.001
Silicone antifoam	5.0	5.0	5.0
Sulfate	-	9.0	-
Density (g/litre)	700	700	700
Miscellaneous and minors			Up to 100%

Example 21

The following detergent compositions were prepared according to the present invention :

	I	II	III	IV
Base granule				
Zeolite A	30.0	22.0	24.0	10.0
Sulfate	10.0	5.0	10.0	7.0
MA/AA	3.0	-	-	-
AA	-	1.6	2.0	-
MA/AA 1	-	12.0	-	6.0
LAS	14.0	10.0	9.0	20.0
C45AS	8.0	7.0	9.0	7.0
C45AES	-	1.0	1.0	-
Silicate	-	1.0	0.5	10.0
Soap	-	2.0	-	-
Brightener 1	0.2	0.2	0.2	0.2
Carbonate	6.0	9.0	10.0	10.0
PEG 4000	-	1.0	1.5	-
DTPA	-	0.4	-	-
Spray On				
C25E9	-	-	-	5.0
C45E7	1.0	1.0	-	-
C23E9	-	1.0	2.5	-
Perfume	0.2	0.3	0.3	-
Dry additives				
Carbonate	5.0	10.0	18.0	8.0
PVPVI/PVNO	0.5	-	0.3	-
Protease	0.03	0.03	0.03	0.02
Mannanase	0.002	0.009	0.015	0.03
Lipase	0.008	-	-	0.008

WO 99/64619

160

PCT/DK99/00314

	I	II	III	IV
Amylase	0.002	-	-	0.002
Cellulase	0.0002	0.0005	0.0005	0.0002
NOBS	-	4.0	-	4.5
PB1	1.0	5.0	1.5	6.0
Sulfate	4.0	5.0	-	5.0
SRP 1	-	0.4	-	-
Suds suppressor	-	0.5	0.5	-
Miscellaneous and minors			Up to 100%	

Example 22

The following granular detergent compositions were prepared according to the present invention :

	I	II	III
Blown Powder			
Zeolite A	20.0	-	15.0
STPP	-	20.0	-
Sulfate	-	-	5.0
Carbonate	-	-	5.0
TAS	-	-	1.0
LAS	6.0	6.0	6.0
C68AS	2.0	2.0	-
Silicate	3.0	8.0	-
MA/AA	4.0	2.0	2.0
CMC	0.6	0.6	0.2
Brightener 1	0.2	0.2	0.1
DETPMP	0.4	0.4	0.1
STS	-	-	1.0
Spray On			
C45E7	5.0	5.0	4.0
Silicone antifoam	0.3	0.3	0.1
Perfume	0.2	0.2	0.3
Dry additives			
QEA	-	-	1.0
Carbonate	14.0	9.0	10.0
PB1	1.5	2.0	-
PB4	18.5	13.0	13.0
TAED	2.0	2.0	2.0
QAS	-	-	1.0

WO 99/64619	162			PCT/DK99/00314
	I	II	III	
Photoactivated bleach	15 ppm	15 ppm	15 ppm	
Na-SKS-6	-	-	3.0	
Protease	0.03	0.03	0.007	
Mannanase	0.001	0.005	0.02	
Lipase	0.004	0.004	0.004	
Amylase	0.006	0.006	0.003	
Cellulase	0.0002	0.0002	0.0005	
Sulfate	10.0	20.0	5.0	
Density (g/litre)	700	700	700	
Miscellaneous and minors			Up to 100%	

Example 23

The following detergent compositions were prepared according to the present invention :

	I	II	III
Blown Powder			
Zeolite A	15.0	15.0	15.0
Sulfate	-	5.0	-
LAS	3.0	3.0	3.0
QAS	-	1.5	1.5
DETPMP	0.4	0.2	0.4
EDDS	-	0.4	0.2
CMC	0.4	0.4	0.4
MA/AA	4.0	2.0	2.0
Agglomerate			
LAS	5.0	5.0	5.0
TAS	2.0	2.0	1.0
Silicate	3.0	3.0	4.0
Zeolite A	8.0	8.0	8.0
Carbonate	8.0	8.0	4.0
Spray On			
Perfume	0.3	0.3	0.3
C45E7	2.0	2.0	2.0
C25E3	2.0	-	-
Dry Additives			
Citrate	5.0	-	2.0
Bicarbonate	-	3.0	-
Carbonate	8.0	15.0	10.0
TAED	6.0	2.0	5.0
PB1	14.0	7.0	10.0
PEO	-	-	0.2

WO 99/64619

164

PCT/DK99/00314

	I	II	III
Bentonite clay	-	-	10.0
Protease	0.03	0.03	0.03
Mannanase	0.001	0.005	0.01
Lipase	0.008	0.008	0.008
Cellulase	0.001	0.001	0.001
Amylase	0.01	0.01	0.01
Silicone antifoam	5.0	5.0	5.0
Sulfate	-	3.0	-
Density (g/litre)	850	850	850
Miscellaneous and minors			Up to 100%

Example 24

The following detergent compositions were prepared according to the present invention :

	I	II	III	IV
LAS	18.0	14.0	24.0	20.0
QAS	0.7	1.0	-	0.7
TFAA	-	1.0	-	-
C23E56.5	-	-	1.0	-
C45E7	-	1.0	-	-
C45E3S	1.0	2.5	1.0	-
STPP	32.0	18.0	30.0	22.0
Silicate	9.0	5.0	9.0	.8.0
Carbonate	11.0	7.5	10.0	5.0
Bicarbonate	-	7.5	-	-
PB1	3.0	1.0	-	-
PB4	-	1.0	-	-
NOBS	2.0	1.0	-	-
DETPMP	-	1.0	-	-
DTPA	0.5	-	0.2	0.3
SRP 1	0.3	0.2	-	0.1
MA/AA	1.0	1.5	2.0	0.5
CMC	0.8	0.4	0.4	.0.2
PEI	-	-	0.4	-
Sulfate	20.0	10.0	20.0	30.0
Mg sulfate	0.2	-	0.4	0.9
Mannanase	0.001	0.005	0.01	0.015
Protease	0.03	0.03	0.02	0.02
Amylase	0.008	0.007	-	0.004
Lipase	0.004	-	0.002	-
Cellulase	0.0003	-	-	0.0001

WO 99/64619

166

PCT/DK99/00314

	I	II	III	IV
Photoactivated bleach	30 ppm	20 ppm	-	10 ppm
Perfume	0.3	0.3	0.1	0.2
Brightener 1/2	0.05	0.02	0.08	0.1
Miscellaneous and minors			up to 100%	

Example 25

The following liquid detergent formulations were prepared according to the present invention (Levels are given in parts per weight, enzyme are expressed in pure enzyme) :

5

	I	II	III	IV	V
LAS	11.5	8.8	-	3.9	-
C25E2.5S	-	3.0	18.0	-	16.0
C45E2.25S	11.5	3.0	-	15.7	-
C23E9	-	2.7	1.8	2.0	1.0
C23E7	3.2	-	-	-	-
CFAA	-	-	5.2	-	3.1
TPKFA	1.6	-	2.0	0.5	2.0
Citric (50%)	6.5	1.2	2.5	4.4	2.5
Ca formate	0.1	0.06	0.1	-	-
Na formate	0.5	0.06	0.1	0.05	0.05
SCS	4.0	1.0	3.0	1.2	-
Borate	0.6	-	3.0	2.0	2.9
Na hydroxide	5.8	2.0	3.5	3.7	2.7
Ethanol	1.75	1.0	3.6	4.2	2.9
1,2 Propanediol	3.3	2.0	8.0	7.9	5.3
Monoethanolamine	3.0	1.5	1.3	2.5	0.8
TEPAE	1.6	-	1.3	1.2	1.2
Mannanase	0.001	0.01	0.015	0.015	0.001
Protease	0.03	0.01	0.03	0.02	0.02
Lipase	-	-	0.002	-	-
Amylase	-	-	-	0.002	-
Cellulase	-	-	0.0002	0.0005	0.0001
SRP 1	0.2	-	0.1	-	-
DTPA	-	-	0.3	-	-
PVNO	-	-	0.3	-	0.2

WO 99/64619

168

PCT/DK99/00314

	I	II	III	IV	V
Brightener 1	0.2	0.07	0.1	-	-
Silicone antifoam	0.04	0.02	0.1	0.1	0.1
Miscellaneous and water					

Example 26

The following liquid detergent formulations were prepared according to the present invention (Levels are given in parts per weight, enzyme are expressed in pure enzyme) :

5

	I	II	III	IV
LAS	10.0	13.0	9.0	-
C25AS	4.0	1.0	2.0	10.0
C25E3S	1.0	-	-	3.0
C25E7	6.0	8.0	13.0	2.5
TFAA	-	-	-	4.5
APA	-	1.4	-	-
TPKFA	2.0	-	13.0	7.0
Citric	2.0	3.0	1.0	1.5
Dodecenyl / tetradecenyl succinic acid	12.0	10.0	-	-
Rapeseed fatty acid	4.0	2.0	1.0	-
Ethanol	4.0	4.0	7.0	2.0
1,2 Propanediol	4.0	4.0	2.0	7.0
Monoethanolamine	-	-	-	5.0
Triethanolamine	-	-	8.0	-
TEPAE	0.5	-	0.5	0.2
DETPMP	1.0	1.0	0.5	1.0
Mannanase	0.001	0.015	0.01	0.03
Protease	0.02	0.02	0.01	0.008
Lipase	-	0.002	-	0.002
Amylase	0.004	0.004	0.01	0.008
Cellulase	-	-	-	0.002
SRP 2	0.3	-	0.3	0.1
Boric acid	0.1	0.2	1.0	2.0
Ca chloride	-	0.02	-	0.01

WO 99/64619

PCT/DK99/00314

170

	I	II	III	IV
Brightener 1	-	0.4	-	-
Suds suppressor	0.1	0.3	-	0.1
Opacifier	0.5	0.4	-	0.3
NaOH up to pH	8.0	8.0	7.6	7.7
Miscellaneous and water				

Example 27

The following liquid detergent compositions were prepared according to the present invention (Levels are given in parts per weight, enzyme are expressed in pure enzyme) :

5

	I	II	III	IV
LAS	25.0	-	-	-
C25AS	-	13.0	18.0	15.0
C25E3S	-	2.0	2.0	4.0
C25E7	-	-	4.0	4.0
TFAA	-	6.0	8.0	8.0
APA	3.0	1.0	2.0	-
TPKFA	-	15.0	11.0	11.0
Citric	1.0	1.0	1.0	1.0
Dodecenyl / tetradecenyl succinic acid	15.0	-	-	-
Rapeseed fatty acid	1.0	-	3.5	-
Ethanol	7.0	2.0	3.0	2.0
1,2 Propanediol	6.0	8.0	10.0	13.0
Monoethanolamine	-	-	9.0	9.0
TEPAE	-	-	0.4	0.3
DETPMP	2.0	1.2	1.0	-
Mannanase	0.001	0.0015	0.01	0.01
Protease	0.05	0.02	0.01	0.02
Lipase	-	-	0.003	0.003
Amylase	0.004	0.01	0.01	0.01
Cellulase	-	-	0.004	0.003
SRP 2	-	-	0.2	0.1
Boric acid	1.0	1.5	2.5	2.5
Bentonite clay	4.0	4.0	-	-
Brightener 1	0.1	0.2	0.3	-

WO 99/64619

PCT/DK99/00314

172

	I	II	III	IV
Suds suppressor	0.4	-	-	-
Opacifier	0.8	0.7	-	-
NaOH up to pH	8.0	7.5	8.0	8.2
Miscellaneous and water				

Example 28

The following liquid detergent compositions were prepared according to the present invention (Levels are given in parts by weight, enzyme are expressed in pure enzyme) :

5

	I	II
LAS	27.6	18.9
C45AS	13.8	5.9
C13E8	3.0	3.1
Oleic acid	3.4	2.5
Citric	5.4	5.4
Na hydroxide	0.4	3.6
Ca Formate	0.2	0.1
Na Formate	-	0.5
Ethanol	7.0	-
Monoethanolamine	16.5	8.0
1,2 propanediol	5.9	5.5
Xylene sulfonic acid	-	2.4
TEPAE	1.5	0.8
Protease	0.05	0.02
Mannanase	0.001	0.01
PEG	-	0.7
Brightener 2	0.4	0.1
Perfume	0.5	0.3
Water and Minors		

Example 29

The following granular fabric detergent compositions which provide "softening through the wash" capability were prepared according to the present invention :

5

	I	II
C45AS	-	10.0
LAS	7.6	-
C68AS	1.3	-
C45E7	4.0	-
C25E3	-	5.0
Coco-alkyl-dimethyl hydroxy-ethyl ammonium chloride	1.4	1.0
Citrate	5.0	3.0
Na-SKS-6	-	11.0
Zeolite A	15.0	15.0
MA/AA	4.0	4.0
DETPMP	0.4	0.4
PB1	15.0	-
Percarbonate	-	15.0
TAED	5.0	5.0
Smectite clay	10.0	10.0
HMWPEO	-	0.1
Mannanase	0.001	0.01
Protease	0.02	0.01
Lipase	0.02	0.01
Amylase	0.03	0.005
Cellulase	0.001	-
Silicate	3.0	5.0
Carbonate	10.0	10.0
Suds suppressor	1.0	4.0

WO 99/64619

175

PCT/DK99/00314

	I	II
CMC	0.2	0.1
Miscellaneous and minors		Up to 100%

Example 30

The following rinse added fabric softener composition was prepared according to the present invention :

DEQA (2)	20.0
Mannanase	0.0008
Cellulase	0.001
HCL	0.03
Antifoam agent	0.01
Blue dye	25ppm
CaCl ₂	0.20
Perfume	0.90
5 Miscellaneous and water	Up to 100%

Example 31

The following fabric softener and dryer added fabric conditioner compositions were prepared according to the present invention :

5

	I	II	III	IV	V
DEQA	2.6	19.0	-	-	-
DEQA (2)	-	-	-	-	51.8
DTMAMS	-	-	-	26.0	-
SDASA	-	-	70.0	42.0	40.2
Stearic acid of IV=0	0.3	-	-	-	-
Neodol 45-13	-	-	13.0	-	-
Hydrochloride acid	0.02	0.02	-	-	-
Ethanol	-	-	1.0	-	-
Mannanase	0.0008	0.0002	0.0005	0.005	0.0002
Perfume	1.0	1.0	0.75	1.0	1.5
Glycoperse S-20	-	-	-	-	15.4
Glycerol	-	-	-	26.0	-
monostearate					
Digeranyl Succinate	-	-	0.38	-	-
Silicone antifoam	0.01	0.01	-	-	-
Electrolyte	-	0.1	-	-	-
Clay	-	-	-	3.0	-
Dye	10ppm	25ppm	0.01	-	-
Water and minors	100%	100%	-	-	-

Example 32

The following laundry bar detergent compositions were prepared according to the present invention (Levels are given in parts per weight, enzyme are expressed in pure enzyme) :

5

	I	II	III	VI	V	III	VI	V
LAS	-	-	19.0	15.0	21.0	6.75	8.8	-
C28AS	30.0	13.5	-	-	-	15.7	11.2	22.5
						5		
Na Laurate	2.5	9.0	-	-	-	-	-	-
Zeolite A	2.0	1.25	-	-	-	1.25	1.25	1.25
Carbonate	20.0	3.0	13.0	8.0	10.0	15.0	15.0	10.0
Ca Carbon- ate	27.5	39.0	35.0	-	-	40.0	-	40.0
Sulfate	5.0	5.0	3.0	5.0	3.0	-	-	5.0
TSP	5.0	-	-	-	-	5.0	2.5	-
STP	5.0	15.0	10.0	-	-	7.0	8.0	10.0
Bentonite clay	-	10.0	-	-	5.0	-	-	-
DETPMP	-	0.7	0.6	-	0.6	0.7	0.7	0.7
CMC	-	1.0	1.0	1.0	1.0	-	-	1.0
Talc	-	-	10.0	15.0	10.0	-	-	-
Silicate	-	-	4.0	5.0	3.0	-	-	-
PVNO	0.02	0.03	-	0.01	-	0.02	-	-
MA/AA	0.4	1.0	-	-	0.2	0.4	0.5	0.4
SRP 1	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Mannanase	0.00	0.00	0.01	0.01	0.01	0.00	0.05	0.01
	1	1			5	1		
Amylase	-	-	0.01	-	-	-	0.00	-

WO 99/64619

179

PCT/DK99/00314

	I	II	III	VI	V	III	VI	V
Protease	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	1	4	1	3	3	1	1	3
Lipase	-	0.00	-	0.00	-	-	-	-
		2		2				
Cellulase	-	.000	-	-	.000	.000	-	-
		3			3	2		
PEO	-	0.2	-	0.2	0.3	-	-	0.3
Perfume	1.0	0.5	0.3	0.2	0.4	-	-	0.4
Mg sulfate	-	-	3.0	3.0	3.0	-	-	-
Brightener	0.15	0.1	0.15	-	-	-	-	0.1
Photoactiva	-	15.0	15.0	15.0	15.0	-	-	15.0
ted bleach								
(ppm)								

Example 33

The following detergent additive compositions were prepared according to the present invention :

	I	II	III
LAS	-	5.0	5.0
STPP	30.0	-	20.0
Zeolite A	-	35.0	20.0
PB1	20.0	15.0	-
TAED	10.0	8.0	-
Mannanase	0.001	0.01	0.01
Protease	0.3	0.3	0.3
Amylase	-	0.06	0.06
5 Minors, water and miscellaneous			Up to 100%

Example 34

The following compact high density (0.96Kg/l) dishwashing detergent compositions were prepared according to the present invention :

5

	I	II	III	IV	V	VI	VII	VIII
STPP	-	-	54.3	51.4	51.4	-	-	50.9
Citrate	35.0	17.0	-	-	-	46.1	40.2	-
Carbonate	-	17.5	14.0	14.0	14.0	-	8.0	32.1
Bicarbonat e	-	-	-	-	-	25.4	-	-
Silicate	32.0	14.8	14.8	10.0	10.0	1.0	25.0	3.1
Metasilica te	-	2.5	-	9.0	9.0	-	-	-
PB1	1.9	9.7	7.8	7.8	7.8	-	-	-
PB4	8.6	-	-	-	-	-	-	-
Percarbona te	-	-	-	-	-	6.7	11.8	4.8
Nonionic	1.5	2.0	1.5	1.7	1.5	2.6	1.9	5.3
TAED	5.2	2.4	-	-	-	2.2	-	1.4
HEDP	-	1.0	-	-	-	-	-	-
DETPMP	-	0.6	-	-	-	-	-	-
MnTACN	-	-	-	-	-	-	0.00	-
							8	
PAAC	-	-	0.00	0.01	0.00	-	-	-
			8		7			
BzP	-	-	-	-	1.4	-	-	-
Paraffin	0.5	0.5	0.5	0.5	0.5	0.6	-	-
Mannanase	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
	1	1	2	2	1	3	2	2

[illegible]

Example 35

The following granular dishwashing detergent compositions of bulk density 1.02Kg/L were prepared according to the present invention :

5

	I	II	III	IV	V	VI	VII	VII
STPP	30.0	30.0	33.0	34.2	29.6	31.1	26.6	17.6
Carbonate	30.5	30.5	31.0	30.0	23.0	39.4	4.2	45.0
Silicate	7.4	7.4	7.5	7.2	13.3	3.4	43.7	12.4
Metasilicat e	-	-	4.5	5.1	-	-	-	-
Percarbonat e	-	-	-	-	-	4.0	-	-
PB1	4.4	4.2	4.5	4.5	-	-	-	-
NADCC	-	-	-	-	2.0	-	1.6	1.0
Nonionic	1.2	1.0	0.7	0.8	1.9	0.7	0.6	0.3
TAED	1.0	-	-	-	-	0.8	-	-
PAAC	-	0.00	0.004	0.00	-	-	-	-
		4		4				
BzP	-	-	-	1.4	-	-	-	-
Paraffin	0.25	0.25	0.25	0.25	-	-	-	-
Mannanase	0.00	0.00	0.001	0.00	0.00	0.00	0.00	0.00
	1	1		1	1	1	1	1
Protease	0.03	0.01	0.03	0.02	-	0.03	-	-
	6	5		8				
Amylase	0.00	0.00	0.01	0.00	-	0.01	-	-
	3	3		6				
Lipase	0.00	-	0.001	-	-	-	-	-
	5							
BTA	0.15	0.15	0.15	0.15	-	-	-	-
Perfume	0.2	0.2	0.2	0.2	0.1	0.2	0.2	-

Example 36

The following tablet detergent compositions were prepared according to the present invention by compression of a granular dishwashing detergent composition at a pressure of 13KN/cm² using a standard 12 head rotary press:

	I	II	III	IV	V	VI
STPP	-	48.8	49.2	38.0	-	46.8
Citrate	26.4	-	-	-	31.1	-
Carbonate	-	5.0	14.0	15.4	14.4	23.0
Silicate	26.4	14.8	15.0	12.6	17.7	2.4
Mannanase	0.001	0.001	0.001	0.001	0.001	0.02
Protease	0.058	0.072	0.041	0.033	0.052	0.013
Amylase	0.01	0.03	0.012	0.007	0.016	0.002
Lipase	0.005	-	-	-	-	-
PB1	1.6	7.7	12.2	10.6	15.7	-
PB4	6.9	-	-	-	-	14.4
Nonionic	1.5	2.0	1.5	1.65	0.8	6.3
PAAC	-	-	0.02	0.009	-	-
MnTACN	-	-	-	-	0.007	-
TAED	4.3	2.5	-	-	1.3	1.8
HEDP	0.7	-	-	0.7	-	0.4
DETPMP	0.65	-	-	-	-	-
Paraffin	0.4	0.5	0.5	0.55	-	-
BTA	0.2	0.3	0.3	0.3	-	-
PA30	3.2	-	-	-	-	-
MA/AA	-	-	-	-	4.5	0.55
Perfume	-	-	0.05	0.05	0.2	0.2
Sulphate	24.0	13.0	2.3	-	10.7	3.4
Weight of tablet	25g	25g	20g	30g	18g	20g

WO 99/64619

186

PCT/DK99/00314

	I	II	III	IV	V	VI
pH	10.6	10.6	10.7	10.7	10.9	11.2
Miscellaneous and water					Up to 100%	

Example 37

The following liquid dishwashing detergent compositions of density 1.40Kg/L were prepared according to the present invention :

5

	I	II	III	IV
STPP	17.5	17.5	17.2	16.0
Carbonate	2.0	-	2.4	-
Silicate	5.3	6.1	14.6	15.7
NaOCl	1.15	1.15	1.15	1.25
Polygen/carbopol	1.1	1.0	1.1	1.25
Nonionic	-	-	0.1	-
NaBz	0.75	0.75	-	-
Mannanase	0.001	0.005	0.01	0.001
NaOH	-	1.9	-	3.5
KOH	2.8	3.5	3.0	-
pH	11.0	11.7	10.9	11.0
Sulphate, miscellaneous and water	up to 100%			

Example 38

The following liquid dishwashing compositions were prepared according to the present invention :

	I	II	III	IV	V
C17ES	28.5	27.4	19.2	34.1	34.1
Amine oxide	2.6	5.0	2.0	3.0	3.0
C12 glucose amide	-	-	6.0	-	-
Betaine	0.9	-	-	2.0	2.0
Xylene sulfonate	2.0	4.0	-	2.0	-
Neodol C11E9	-	-	5.0	-	-
Polyhydroxy fatty acid amide	-	-	-	6.5	6.5
Sodium diethylene penta acetate (40%)	-	-	0.03	-	-
TAED	-	-	-	0.06	0.06
Sucrose	-	-	-	1.5	1.5
Ethanol	4.0	5.5	5.5	9.1	9.1
Alkyl diphenyl oxide disulfonate	-	-	-	-	2.3
Ca formate	-	-	-	0.5	1.1
Ammonium citrate	0.06	0.1	-	-	-
Na chloride	-	1.0	-	-	-
Mg chloride	3.3	-	0.7	-	-
Ca chloride	-	-	0.4	-	-
Na sulfate	-	-	0.06	-	-
Mg sulfate	0.08	-	-	-	-
Mg hydroxide	-	-	-	2.2	2.2
Na hydroxide	-	-	-	1.1	1.1
Hydrogen peroxide	200ppm	0.16	0.006	-	-

WO 99/64619

PCT/DK99/00314

189

	I	II	III	IV	V
Mannanase	0.001	0.05	0.001	0.00	0.01
				1	5
Protease	0.017	0.005	0.035	0.00	0.00
				3	2
Perfume	0.18	0.09	0.09	0.2	0.2
Water and minors				Up to 100%	

Example 39

The following liquid hard surface cleaning compositions were prepared according to the present invention :

	I	II	III	IV	V
Mannanase	0.001	0.0015	0.0015	0.05	0.01
Amylase	0.01	0.002	0.005	-	-
Protease	0.05	0.01	0.02	-	-
Hydrogen peroxide	-	-	-	6.0	6.8
Acetyl triethyl citrate	-	-	-	2.5	-
DTPA	-	-	-	0.2	-
Butyl hydroxy toluene	-	-	-	0.05	-
EDTA*	0.05	0.05	0.05	-	-
Citric / Citrate	2.9	2.9	2.9	1.0	-
LAS	0.5	0.5	0.5	-	-
C12 AS	0.5	0.5	0.5	-	-
C10AS	-	-	-	-	1.7
C12(E)S	0.5	0.5	0.5	-	-
C12,13 E6.5 non- ionic	7.0	7.0	7.0	-	-
Neodol 23-6.5	-	-	-	12.0	-
Dobanol 23-3	-	-	-	-	1.5
Dobanol 91-10	-	-	-	-	1.6
C25AE1.8S	-	-	-	6.0	
Na paraffin sul- phonate	-	-	-	6.0	
Perfume	1.0	1.0	1.0	0.5	0.2
Propanediol	-	-	-	1.5	

WO 99/64619

191

PCT/DK99/00314

	I	II	III	IV	V
Ethoxylated tetra- ethylene pen- tamine	-	-	-	1.0	-
2, Butyl octanol	-	-	-	-	0.5
Hexyl carbitol**	1.0	1.0	1.0	-	-
SCS	1.3	1.3	1.3	-	-
pH adjusted to	7-12	7-12	7-12	4	-
Miscellaneous and water				Up to 100%	

*Na4 ethylenediamine diacetic acid

**Diethylene glycol monohexyl ether

Example 40

The following spray composition for cleaning of hard surfaces and removing household mildew was prepared according to the present invention :

5

Mannanase	0.01
Amylase	0.01
Protease	0.01
Na octyl sulfate	2.0
Na dodecyl sulfate	4.0
Na hydroxide	0.8
Silicate	0.04
Butyl carbitol*	4.0
Perfume	0.35
Water/minors	up to 100%

*Diethylene glycol monobutyl ether

LITERATURE

Aviv, H. & Leder, P. 1972. Proc. Natl. Acad. Sci. U. S. A. 69: 1408-1412.

5

Becker, D. M. & Guarante, L. 1991. Methods Enzymol. 194: 182-187.

Chirgwin, J. M., Przybyla, A. E., MacDonald, R. J. & Rutter, W. J. 1979. Biochemistry 18: 5294-5299.

Gubler, U. & Hoffman, B. J. 1983. Gene 25: 263-269.

15 Sambrook, J., Fritsch, E. F. & Maniatis, T. 1989. Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Lab., Cold Spring Harbor, NY.

Sanger, F., Nicklen, S. & Coulson, A. R. 1977. Proc. Natl. Acad. Sci. U. S. A. 74: 5463-5467.

Lever, M. (1972) A new reaction for colormetric determination of carbohydrates. Anal. Biochem. 47, 273-279.

25 N. C. Carpita and D. M. Gibeaut (1993) The Plant Journal 3:1-30.

Diderichsen, B., Wedsted, U., Hedegaard, L., Jensen, B. R., Sjøholm, C. (1990) Cloning of aldB, which encodes alpha-acetolactate decarboxylase, an exoenzyme from *Bacillus brevis*. J. Bacteriol. 172:4315-4321.

30

193a

SEQUENCE LISTING

<110> NOVOZYMES A/S

<120> NOVEL MANNANASES

<130> 15194-1 FC/gc

<140> 2,331,199

<141> 1999-06-10

<150> PCT/DK99/00314

<151> 1999-06-10

<150> 09/111,256

<151> 1999-06-10

<150> DK PA 1998 01341

<151> 1998-10-20

<150> DK PA 1998 01340

<151> 1998-10-20

<150> 60/105,970

<151> 1998-10-28

<150> 60/106,054

<151> 1998-10-28

<150> DK PA 1998 01725

<151> 1998-12-23

<150> DK PA 1999 00308

<151> 1999-03-05

<150> DK PA 1999 00307

<151> 1999-03-05

<150> DK PA 1999 00306

<151> 1999-03-05

<150> DK PA 1999 00309

<151> 1999-03-05

<150> 60/123,543

<151> 1999-03-09

<150> 60/123,641

<151> 1999-03-10

<150> 60-123,623

<151> 1999-03-10

<150> 60-123,642

<151> 1999-03-11

193b

<160> 34

<170> PatentIn Ver. 2.1

<210> 1

<211> 1470

<212> DNA

<213> Bacillus sp. I633

<400> 1

```

ttgaataatg gttttaaaaa aatttttttct ataacattat cattactatt agctagctct 60
attctgttcg tttcaggaac ttctacagct aatgcaaatt ccggatttta tgtaagcggg 120
accactctat acgatgccaa tggaaaccca tttgtaatga gagggattaa ccatgggcac 180
gcatgggtata aagaccaggc aactactgca attgaaggga ttgcaaatac cggtgctaat 240
acggtccegga ttgtgttatc tgatggggga caatggacaa aagatgacat ccatacagta 300
agaaacctta tctcttttagc ggaagataat catttggttg ctgttcttga agttcatgat 360
gctaccgggtt atgattccat tgcttcgctc aatcgtgctg ttgattattg gattgaaatg 420
agaagtgcct taattggaaa ggaagatacc gtcattatta atattgcgaa tgaatgggtt 480
ggttcgtggg aaggggatgc ttgggctgac ggggtataaac aagcaatccc gcgattgcgt 540
aacgccggtc taaaccatac cttgatggta gatgctgcgg ggtggggaca atttccacaa 600
tcgattcatg attatggaag agaagttttt aatgctgacc ctcaacgaaa tacaatgttt 660
tcgattcata tgtatgaata tgcagggtgt aatgcacgc aagttcgta taatattgac 720
cgagttctta atcaagacct cgcattagtc attgggtgaat ttggacaccg tcatacaaat 780
ggtgacgtcg atgaagcaac gattatgagc tattctgaac aaagaggagt tgggtgggtg 840
gcgtgggtcat ggaaaggga cggcccagaa tgggagtatt tagaccttc gaatgattgg 900
gctggaaata accttacagc ttgggggaaat acaatagtga atggtccata tggtttaaga 960
gaaacttcga gattaagcac cgtttttaca ggtggaggat ctgatggagg aacttctccg 1020
acaactcttt atgattttga aggtagtatg caaggatgga ctggaagtag cttgagcgga 1080
ggtccttggg ctgtgacaga gtggtcttct aaaggaagtc attctttaaa agcggatatt 1140
caattgtcgt caaattcaca acattactta catgttatc aaaatacgtc tttacagcag 1200
aatagtagga tacaagctac tgttaaacad gcaaattggg gaagtgttgg taatggaatg 1260
actgcgcgtc tttatgtgaa aacaggacat gggtatacat ggtactctgg aagctttgtg 1320
ccgattaacg gttcatctgg aacaacgcta tctctagatt tatcaaattg ccaaaatctt 1380
tctcaagtaa gggaaattgg agttcagttc caatcagcga gtgatagtag tggacaaaca 1440
tcgatttata ttgataatgt gattgtagaa 1470

```

<210> 2

<211> 490

<212> PRT

<213> Bacillus sp. I633

<400> 2

```

Leu Asn Asn Gly Phe Lys Lys Ile Phe Ser Ile Thr Leu Ser Leu Leu
 1              5              10              15

Leu Ala Ser Ser Ile Leu Phe Val Ser Gly Thr Ser Thr Ala Asn Ala
      20              25              30

Asn Ser Gly Phe Tyr Val Ser Gly Thr Thr Leu Tyr Asp Ala Asn Gly
      35              40              45

Asn Pro Phe Val Met Arg Gly Ile Asn His Gly His Ala Trp Tyr Lys
      50              55              60

```

193c

Asp Gln Ala Thr Thr Ala Ile Glu Gly Ile Ala Asn Thr Gly Ala Asn
 65 70 75 80
 Thr Val Arg Ile Val Leu Ser Asp Gly Gly Gln Trp Thr Lys Asp Asp
 85 90 95
 Ile His Thr Val Arg Asn Leu Ile Ser Leu Ala Glu Asp Asn His Leu
 100 105 110
 Val Ala Val Pro Glu Val His Asp Ala Thr Gly Tyr Asp Ser Ile Ala
 115 120 125
 Ser Leu Asn Arg Ala Val Asp Tyr Trp Ile Glu Met Arg Ser Ala Leu
 130 135 140
 Ile Gly Lys Glu Asp Thr Val Ile Ile Asn Ile Ala Asn Glu Trp Phe
 145 150 155 160
 Gly Ser Trp Glu Gly Asp Ala Trp Ala Asp Gly Tyr Lys Gln Ala Ile
 165 170 175
 Pro Arg Leu Arg Asn Ala Gly Leu Asn His Thr Leu Met Val Asp Ala
 180 185 190
 Ala Gly Trp Gly Gln Phe Pro Gln Ser Ile His Asp Tyr Gly Arg Glu
 195 200 205
 Val Phe Asn Ala Asp Pro Gln Arg Asn Thr Met Phe Ser Ile His Met
 210 215 220
 Tyr Glu Tyr Ala Gly Gly Asn Ala Ser Gln Val Arg Thr Asn Ile Asp
 225 230 235 240
 Arg Val Leu Asn Gln Asp Leu Ala Leu Val Ile Gly Glu Phe Gly His
 245 250 255
 Arg His Thr Asn Gly Asp Val Asp Glu Ala Thr Ile Met Ser Tyr Ser
 260 265 270
 Glu Gln Arg Gly Val Gly Trp Leu Ala Trp Ser Trp Lys Gly Asn Gly
 275 280 285
 Pro Glu Trp Glu Tyr Leu Asp Leu Ser Asn Asp Trp Ala Gly Asn Asn
 290 295 300
 Leu Thr Ala Trp Gly Asn Thr Ile Val Asn Gly Pro Tyr Gly Leu Arg
 305 310 315 320
 Glu Thr Ser Arg Leu Ser Thr Val Phe Thr Gly Gly Gly Ser Asp Gly
 325 330 335
 Gly Thr Ser Pro Thr Thr Leu Tyr Asp Phe Glu Gly Ser Met Gln Gly
 340 345 350
 Trp Thr Gly Ser Ser Leu Ser Gly Gly Pro Trp Ala Val Thr Glu Trp
 355 360 365

193d

Ser Ser Lys Gly Ser His Ser Leu Lys Ala Asp Ile Gln Leu Ser Ser
 370 375 380

Asn Ser Gln His Tyr Leu His Val Ile Gln Asn Thr Ser Leu Gln Gln
 385 390 395 400

Asn Ser Arg Ile Gln Ala Thr Val Lys His Ala Asn Trp Gly Ser Val
 405 410 415

Gly Asn Gly Met Thr Ala Arg Leu Tyr Val Lys Thr Gly His Gly Tyr
 420 425 430

Thr Trp Tyr Ser Gly Ser Phe Val Pro Ile Asn Gly Ser Ser Gly Thr
 435 440 445

Thr Leu Ser Leu Asp Leu Ser Asn Val Gln Asn Leu Ser Gln Val Arg
 450 455 460

Glu Ile Gly Val Gln Phe Gln Ser Ala Ser Asp Ser Ser Gly Gln Thr
 465 470 475 480

Ser Ile Tyr Ile Asp Asn Val Ile Val Glu
 485 490

<210> 3

<211> 1438

<212> DNA

<213> Bacillus sp. I633

<400> 3

gcaaattccg gatttttatgt aagcgggtacc actctatacgt atgccaatgg aaacccattt 60
 gtaatgagag ggattaacca tgggcacgca tgggtataaag accaggcaac tactgcaatt 120
 gaagggattg caaataaccgg tgctaatacgt gtccggattg tgttatctga tgggggacaa 180
 tggacaaaag atgacatcca tacagtaaga aacctttatct ctttagcgga agataatcat 240
 ttggttgctg ttcttgaagt tcatgatgct accggttatg attccattgc ttcgctcaat 300
 cgtgctggtg attattggat tgaaatgaga agtgctttaa ttggaaagga agataccgtc 360
 attattaata ttgcgaatga atggtttggg tcgtgggaag gggatgcttg ggctgacggg 420
 tataaacaag caatcccgcg attgcgtaac gccgggtctaa accatacctt gatggtagat 480
 gctgcggggg ggggacaatt tccacaatcg attcatgatt atggaagaga agtttttaat 540
 gctgaccctc aacgaaatac aatgttttcg attcatatgt atgaatatgc aggtgggtaat 600
 gcatcgcaag ttctgtactaa tattgaccga gttcttaatc aagacctcgc attagtcatt 660
 ggtgaatttg gacaccgtca taaaaatggg gacgtcgatg aagcaacgat tatgagctat 720
 tctgaacaaa gaggagttgg gtgggttggcg tgggtcatgga aaggggaacgg cccagaatgg 780
 gagtathtag acctttcgaa tgattgggct ggaaataacc ttacagcttg gggaaataca 840
 atagtgaatg gtccatatgg tttaagagaa acttcgagat taagcaccgt ttttacagct 900
 agcccgggaa caacaccaga gccgaccgca aatacaccgg tatcaggcaa tttgaagggt 960
 gaattctaca acagcaatcc ttcagatact actaactcaa tcaatcctca gttcaagggt 1020
 actaataccg gaagcagtgc aattgatttg tccaaactca cattgagata ttattataca 1080
 gtagacggac agaaagatca gaccttctgg tgtgaccatg ctgcaataat cggcagtaac 1140
 ggcagctaca acggaattac ttcaaatgta aaaggaacat ttgtaaaaat gagttcctca 1200
 acaaataacg cagacaccta ccttgaaata agctttacag gcggaactct tgaaccgggt 1260
 gcacatgttc agatacaagg tagatttgca aagaatgact ggagtaacta tacacagtca 1320
 aatgactact cattcaagtc tcgttcacag tttgttgaat gggatcaggt aacagcatac 1380
 ttgaacgggtg ttcttgtatg gggtaaagaa cccgggtggca gtgtagtata gcggccgc 1438

193e

<210> 4
 <211> 476
 <212> PRT
 <213> Bacillus sp.

<400> 4
 Ala Asn Ser Gly Phe Tyr Val Ser Gly Thr Thr Leu Tyr Asp Ala Asn
 1 5 10 15
 Gly Asn Pro Phe Val Met Arg Gly Ile Asn His Gly His Ala Trp Tyr
 20 25 30
 Lys Asp Gln Ala Thr Thr Ala Ile Glu Gly Ile Ala Asn Thr Gly Ala
 35 40 45
 Asn Thr Val Arg Ile Val Leu Ser Asp Gly Gly Gln Trp Thr Lys Asp
 50 55 60
 Asp Ile His Thr Val Arg Asn Leu Ile Ser Leu Ala Glu Asp Asn His
 65 70 75 80
 Leu Val Ala Val Pro Glu Val His Asp Ala Thr Gly Tyr Asp Ser Ile
 85 90 95
 Ala Ser Leu Asn Arg Ala Val Asp Tyr Trp Ile Glu Met Arg Ser Ala
 100 105 110
 Leu Ile Gly Lys Glu Asp Thr Val Ile Ile Asn Ile Ala Asn Glu Trp
 115 120 125
 Phe Gly Ser Trp Glu Gly Asp Ala Trp Ala Asp Gly Tyr Lys Gln Ala
 130 135 140
 Ile Pro Arg Leu Arg Asn Ala Gly Leu Asn His Thr Leu Met Val Asp
 145 150 155 160
 Ala Ala Gly Trp Gly Gln Phe Pro Gln Ser Ile His Asp Tyr Gly Arg
 165 170 175
 Glu Val Phe Asn Ala Asp Pro Gln Arg Asn Thr Met Phe Ser Ile His
 180 185 190
 Met Tyr Glu Tyr Ala Gly Gly Asn Ala Ser Gln Val Arg Thr Asn Ile
 195 200 205
 Asp Arg Val Leu Asn Gln Asp Leu Ala Leu Val Ile Gly Glu Phe Gly
 210 215 220
 His Arg His Thr Asn Gly Asp Val Asp Glu Ala Thr Ile Met Ser Tyr
 225 230 235 240
 Ser Glu Gln Arg Gly Val Gly Trp Leu Ala Trp Ser Trp Lys Gly Asn
 245 250 255

193f

Gly Pro Glu Trp Glu Tyr Leu Asp Leu Ser Asn Asp Trp Ala Gly Asn
 260 265 270
 Asn Leu Thr Ala Trp Gly Asn Thr Ile Val Asn Gly Pro Tyr Gly Leu
 275 280 285
 Arg Glu Thr Ser Arg Leu Ser Thr Val Phe Thr Ala Ser Pro Glu Pro
 290 295 300
 Thr Pro Glu Pro Thr Ala Asn Thr Pro Val Ser Gly Asn Leu Lys Val
 305 310 315 320
 Glu Phe Tyr Asn Ser Asn Pro Ser Asp Thr Thr Asn Ser Ile Asn Pro
 325 330 335
 Gln Phe Lys Val Thr Asn Thr Gly Ser Ser Ala Ile Asp Leu Ser Lys
 340 345 350
 Leu Thr Leu Arg Tyr Tyr Tyr Thr Val Asp Gly Gln Lys Asp Gln Thr
 355 360 365
 Phe Trp Cys Asp His Ala Ala Ile Ile Gly Ser Asn Gly Ser Tyr Asn
 370 375 380
 Gly Ile Thr Ser Asn Val Lys Gly Thr Phe Val Lys Met Ser Ser Ser
 385 390 395 400
 Thr Asn Asn Ala Asp Thr Tyr Leu Glu Ile Ser Phe Thr Gly Gly Thr
 405 410 415
 Leu Glu Pro Gly Ala His Val Gln Ile Gln Gly Arg Phe Ala Lys Asn
 420 425 430
 Asp Trp Ser Asn Tyr Thr Gln Ser Asn Asp Tyr Ser Phe Lys Ser Arg
 435 440 445
 Ser Gln Phe Val Glu Trp Asp Gln Val Thr Ala Tyr Leu Asn Gly Val
 450 455 460
 Leu Val Trp Gly Lys Glu Pro Gly Gly Ser Val Val
 465 470 475

<210> 5

<211> 1482

<212> DNA

<213> Bacillus agaradhaerens

<400> 5

atgaaaaaaa agttatcaca gattttatcat ttaattattt gcacacttat aataagtgtg 60
 ggaataatgg ggattacaac gtcccatca gcagcaagta caggctttta tgttgatggc 120
 aatacgttat atgacgcaaa tgggcagcca tttgtcatga gaggtattaa ccatggacat 180
 gcttggtata aagacaccgc ttcaacagct attcctgccca ttgcagagca aggcgccaac 240
 acgattcgta ttgttttatac agatggcggt caatgggaaa aagacgacat tgacaccatt 300
 cgtgaagtca ttgagcttgc ggagcaaaat aaaatggtgg ctgtcgttga agttcatgat 360

193g

```

gccacggggtc gcgattcgcg cagtgattta aatcgagccg ttgattattg gatagaaatg 420
aaagatgcgc ttatcggtaa agaagatacg gttattatta acattgcaaa cgagtgggtat 480
gggagttggg atggctcagc ttggggccgat ggctatatattg atgtcattcc gaagcttcgc 540
gatgccggct taacacacac cttaatgggt gatgcagcag gatgggggca atatccgcaa 600
tctattcatg attacggaca agatgtgttt aatgcagatc cgttaaaaaa tacgatgttc 660
tccatccata tgtatgagta tgctgggtgg gatgctaaca ctgttagatc aaatattgat 720
agagtcatag atcaagacct tgctctcgta atagggtgaat tcggtcatag acatactgat 780
ggtgatgttg atgaagatac aatccttagt tattctgaag aaactggcac aggggtggctc 840
gcttgggtcct ggaaaggcaa cagtaccgaa tgggactatt tagaccttc agaagactgg 900
gctgggtcaac atttaactga ttgggggaat agaattgtcc acggggccga tggcttacag 960
gaaacctcca aaccatccac cgtatttaca gatgataacg gtggtcaccc tgaaccgcca 1020
actgctacta ccttgatga ctttgaagga agcacacaag ggtggcatgg aagcaacgtg 1080
accggtggcc cttgggtccgt aacagaatgg ggtgcttcag gtaactactc tttaaaagcc 1140
gatgtaaatt taacctcaaa ttcttcacat gaactgtata gtgaacaaag tcgtaatcta 1200
cacggatact ctcagctcaa cgcaaccgtt cgccatgcc attggggaaa tcccggtaat 1260
ggcatgaatg caagacttta cgtgaaaacg ggctctgatt atacatggca tagcggtcct 1320
tttacacgta tcaatagctc caactcagga acaacgttat cttttgattt aaacaacatc 1380
gaaaatagtc atcatgttag ggaaataggc gtgcaatttt cagcggcaga taatagcagt 1440
ggtcaaactg ctctatacgt tgataacggt actttaagat ag 1482

```

<210> 6

<211> 493

<212> PRT

<213> Bacillus agaradhaerens

<400> 6

```

Met Lys Lys Lys Leu Ser Gln Ile Tyr His Leu Ile Ile Cys Thr Leu
  1              5              10              15

Ile Ile Ser Val Gly Ile Met Gly Ile Thr Thr Ser Pro Ser Ala Ala
      20              25              30

Ser Thr Gly Phe Tyr Val Asp Gly Asn Thr Leu Tyr Asp Ala Asn Gly
      35              40              45

Gln Pro Phe Val Met Arg Gly Ile Asn His Gly His Ala Trp Tyr Lys
      50              55              60

Asp Thr Ala Ser Thr Ala Ile Pro Ala Ile Ala Glu Gln Gly Ala Asn
      65              70              75              80

Thr Ile Arg Ile Val Leu Ser Asp Gly Gly Gln Trp Glu Lys Asp Asp
      85              90              95

Ile Asp Thr Ile Arg Glu Val Ile Glu Leu Ala Glu Gln Asn Lys Met
      100             105             110

Val Ala Val Val Glu Val His Asp Ala Thr Gly Arg Asp Ser Arg Ser
      115             120             125

Asp Leu Asn Arg Ala Val Asp Tyr Trp Ile Glu Met Lys Asp Ala Leu
      130             135             140

```

193h

Ile	Gly	Lys	Glu	Asp	Thr	Val	Ile	Ile	Asn	Ile	Ala	Asn	Glu	Trp	Tyr	145	150	155	160
Gly	Ser	Trp	Asp	Gly	Ser	Ala	Trp	Ala	Asp	Gly	Tyr	Ile	Asp	Val	Ile	165	170	175	
Pro	Lys	Leu	Arg	Asp	Ala	Gly	Leu	Thr	His	Thr	Leu	Met	Val	Asp	Ala	180	185	190	
Ala	Gly	Trp	Gly	Gln	Tyr	Pro	Gln	Ser	Ile	His	Asp	Tyr	Gly	Gln	Asp	195	200	205	
Val	Phe	Asn	Ala	Asp	Pro	Leu	Lys	Asn	Thr	Met	Phe	Ser	Ile	His	Met	210	215	220	
Tyr	Glu	Tyr	Ala	Gly	Gly	Asp	Ala	Asn	Thr	Val	Arg	Ser	Asn	Ile	Asp	225	230	235	240
Arg	Val	Ile	Asp	Gln	Asp	Leu	Ala	Leu	Val	Ile	Gly	Glu	Phe	Gly	His	245	250	255	
Arg	His	Thr	Asp	Gly	Asp	Val	Asp	Glu	Asp	Thr	Ile	Leu	Ser	Tyr	Ser	260	265	270	
Glu	Glu	Thr	Gly	Thr	Gly	Trp	Leu	Ala	Trp	Ser	Trp	Lys	Gly	Asn	Ser	275	280	285	
Thr	Glu	Trp	Asp	Tyr	Leu	Asp	Leu	Ser	Glu	Asp	Trp	Ala	Gly	Gln	His	290	295	300	
Leu	Thr	Asp	Trp	Gly	Asn	Arg	Ile	Val	His	Gly	Ala	Asp	Gly	Leu	Gln	305	310	315	320
Glu	Thr	Ser	Lys	Pro	Ser	Thr	Val	Phe	Thr	Asp	Asp	Asn	Gly	Gly	His	325	330	335	
Pro	Glu	Pro	Pro	Thr	Ala	Thr	Thr	Leu	Tyr	Asp	Phe	Glu	Gly	Ser	Thr	340	345	350	
Gln	Gly	Trp	His	Gly	Ser	Asn	Val	Thr	Gly	Gly	Pro	Trp	Ser	Val	Thr	355	360	365	
Glu	Trp	Gly	Ala	Ser	Gly	Asn	Tyr	Ser	Leu	Lys	Ala	Asp	Val	Asn	Leu	370	375	380	
Thr	Ser	Asn	Ser	Ser	His	Glu	Leu	Tyr	Ser	Glu	Gln	Ser	Arg	Asn	Leu	385	390	395	400
His	Gly	Tyr	Ser	Gln	Leu	Asn	Ala	Thr	Val	Arg	His	Ala	Asn	Trp	Gly	405	410	415	
Asn	Pro	Gly	Asn	Gly	Met	Asn	Ala	Arg	Leu	Tyr	Val	Lys	Thr	Gly	Ser	420	425	430	
Asp	Tyr	Thr	Trp	His	Ser	Gly	Pro	Phe	Thr	Arg	Ile	Asn	Ser	Ser	Asn	435	440	445	

193i

Ser Gly Thr Thr Leu Ser Phe Asp Leu Asn Asn Ile Glu Asn Ser His
 450 455 460

His Val Arg Glu Ile Gly Val Gln Phe Ser Ala Ala Asp Asn Ser Ser
 465 470 475 480

Gly Gln Thr Ala Leu Tyr Val Asp Asn Val Thr Leu Arg
 485 490

<210> 7

<211> 1407

<212> DNA

<213> Bacillus agaradhaerens

<400> 7

```

atgaaaaaaaa agttatcaca gatttatcat ttaattatatt gcacacttat aataagtgtg 60
ggaataatgg ggattacaac gtcccatca gcagcaagta caggctttta tgttgatggc 120
aatacgttat atgacgcaaa tgggcagcca tttgtcatga gaggtattaa ccatggacat 180
gcttggtata aagacaccgc ttcaacagct attcctgcca ttgcagagca aggcgccaac 240
acgattcgta ttgttttatc agatggcggc caatgggaaa aagacgacat tgacaccatt 300
cgtgaagtca ttgagcttgc ggagcaaaat aaaatggtgg ctgtcgttga agttcatgat 360
gccacgggtc gcgattcgcg cagtgattta aatcgagccg ttgattattg gatagaaatg 420
aaagatgcgc ttatcggtaa agaagatacg gttattatta acattgcaaa cgagtgggat 480
gggagttggg atggctcagc ttgggccgat ggctatattg atgtcattcc gaagcttcgc 540
gatgccggct taacacacac cttaatggtt gatgcagcag gatgggggca atatccgcaa 600
tctattcatg attacggaca agatgtgttt aatgcagatc cgttaaaaaa tacgatgttc 660
tccatccata tgtatgagta tgctgggtgg gatgctaaca ctgttagatc aaatattgat 720
agagtcatag atcaagacct tgctctcgta ataggtgaat tcggtcatag acatactgat 780
ggtgatgttg atgaagatac aatccttagt tattctgaag aaactggcac aggggtggctc 840
gcttggtcct ggaaaggcaa cagtaccgaa tgggactatt tagaccttc agaagactgg 900
gctgggtcaac atttaactga ttgggggaat agaattgtcc acggggccga tggcttacag 960
gaaacctcca aaccatccac cgtatttaca gatgataacg gtggtcacc tgaaccgcca 1020
actgctacta ccttgtatga ctttgaagga agcacacaag ggtggcatgg aagcaacgtg 1080
accggtggcc cttggtccgt aacagaatgg ggtgcttcag gtaactactc tttaaaagcc 1140
gatgtaaatt taacctcaaa ttcttcacat gaactgtata gtgaacaaag tcgtaatcta 1200
cacggatact ctcaagctcaa cgcaaccggt cgccatgcca attggggaaa tcccggtaat 1260
ggcatgaatg caagacttta cgtgaaaacg ggctctgatt atacatggca tagcggtcct 1320
tttacacgta tcaatagctc caactcagga acaacgttat cttttgattt aaacaacatc 1380
gaaaatatca tcatgttagg gaaatag 1407

```

<210> 8

<211> 468

<212> PRT

<213> Bacillus agaradhaerens

<400> 8

Met Lys Lys Lys Leu Ser Gln Ile Tyr His Leu Ile Ile Cys Thr Leu
 1 5 10 15

Ile Ile Ser Val Gly Ile Met Gly Ile Thr Thr Ser Pro Ser Ala Ala
 20 25 30

193j

Ser Thr Gly Phe Tyr Val Asp Gly Asn Thr Leu Tyr Asp Ala Asn Gly
 35 40 45
 Gln Pro Phe Val Met Arg Gly Ile Asn His Gly His Ala Trp Tyr Lys
 50 55 60
 Asp Thr Ala Ser Thr Ala Ile Pro Ala Ile Ala Glu Gln Gly Ala Asn
 65 70 75 80
 Thr Ile Arg Ile Val Leu Ser Asp Gly Gly Gln Trp Glu Lys Asp Asp
 85 90 95
 Ile Asp Thr Ile Arg Glu Val Ile Glu Leu Ala Glu Gln Asn Lys Met
 100 105 110
 Val Ala Val Val Glu Val His Asp Ala Thr Gly Arg Asp Ser Arg Ser
 115 120 125
 Asp Leu Asn Arg Ala Val Asp Tyr Trp Ile Glu Met Lys Asp Ala Leu
 130 135 140
 Ile Gly Lys Glu Asp Thr Val Ile Ile Asn Ile Ala Asn Glu Trp Tyr
 145 150 155 160
 Gly Ser Trp Asp Gly Ser Ala Trp Ala Asp Gly Tyr Ile Asp Val Ile
 165 170 175
 Pro Lys Leu Arg Asp Ala Gly Leu Thr His Thr Leu Met Val Asp Ala
 180 185 190
 Ala Gly Trp Gly Gln Tyr Pro Gln Ser Ile His Asp Tyr Gly Gln Asp
 195 200 205
 Val Phe Asn Ala Asp Pro Leu Lys Asn Thr Met Phe Ser Ile His Met
 210 215 220
 Tyr Glu Tyr Ala Gly Gly Asp Ala Asn Thr Val Arg Ser Asn Ile Asp
 225 230 235 240
 Arg Val Ile Asp Gln Asp Leu Ala Leu Val Ile Gly Glu Phe Gly His
 245 250 255
 Arg His Thr Asp Gly Asp Val Asp Glu Asp Thr Ile Leu Ser Tyr Ser
 260 265 270
 Glu Glu Thr Gly Thr Gly Trp Leu Ala Trp Ser Trp Lys Gly Asn Ser
 275 280 285
 Thr Glu Trp Asp Tyr Leu Asp Leu Ser Glu Asp Trp Ala Gly Gln His
 290 295 300
 Leu Thr Asp Trp Gly Asn Arg Ile Val His Gly Ala Asp Gly Leu Gln
 305 310 315 320
 Glu Thr Ser Lys Pro Ser Thr Val Phe Thr Asp Asp Asn Gly Gly His
 325 330 335

193k

Pro Glu Pro Pro Thr Ala Thr Thr Leu Tyr Asp Phe Glu Gly Ser Thr
 340 345 350

Gln Gly Trp His Gly Ser Asn Val Thr Gly Gly Pro Trp Ser Val Thr
 355 360 365

Glu Trp Gly Ala Ser Gly Asn Tyr Ser Leu Lys Ala Asp Val Asn Leu
 370 375 380

Thr Ser Asn Ser Ser His Glu Leu Tyr Ser Glu Gln Ser Arg Asn Leu
 385 390 395 400

His Gly Tyr Ser Gln Leu Asn Ala Thr Val Arg His Ala Asn Trp Gly
 405 410 415

Asn Pro Gly Asn Gly Met Asn Ala Arg Leu Tyr Val Lys Thr Gly Ser
 420 425 430

Asp Tyr Thr Trp His Ser Gly Pro Phe Thr Arg Ile Asn Ser Ser Asn
 435 440 445

Ser Gly Thr Thr Leu Ser Phe Asp Leu Asn Asn Ile Glu Asn Ile Ile
 450 455 460

Met Leu Gly Lys
 465

<210> 9

<211> 1761

<212> DNA

<213> Bacillus halodurans

<400> 9

```

atgaaaagta taaagaaatt ggtagtcggt tgcattggcat ttctattaat ttttccatcg 60
acgtcatttg ctttttctgg aagcgtttca gcttcaggtc aagagcttaa aatgacagat 120
caaaacgcat ctcaatatac aaaagagttg tttgcctttt tacgtgatgt aagtggtaaa 180
caagttttat ttggtcaaca acacgcaact gatgagggat taacacttag aggaacaggat 240
aaccgaattg gttcaacaga atcagaagtg aaaaatgctg ttggtgatta tcctgctggt 300
tttggttggg atacaaacag tctagatggt agagaaaagc ccggtaatga tgaaccgagt 360
caagaacaaa gaatcttaaa tacagcagct tcaatgaagg cagctcacga cttaggtggg 420
attatcacac taagtatgca tcctgataac tttgtaacag gaggggctta tggcgataca 480
actggaaatg ttgtacaaga aattcttcct ggtggatcaa agcatgaaga attcaatgca 540
tggttggata acctagcggc tttagctcac gaattaaagg atgacaacgg gaaacacatt 600
ccaattattt tccgtccttt ccatgagcaa acagggttctt ggttctggtg gggagcaagc 660
acaacaactc cagaacagta taaagctatt tacagatata cggttgaata cttacgtgac 720
gtaaaaggag caaacaactt cttatacggg ttttctcctg gtgcaggtec agctggcgat 780
ttaaatcggt atatggaaac ttaccctggt gatgattatg tcgatatctt tggattatgat 840
aactatgaca ataaatcaaa tgctggatca gaagcttgga tacaagggtg tgtaaccgat 900
ttagctatgc ttgttgattt agctgaagaa aaaggaaaga ttgctgcgtt taccgagtat 960
ggttacagtg caacaggat gaatcgtact ggtaacacat tggattggta tactcgttta 1020
cttaatgcaa taaaagaaga tccaaaagca agtaagattt cttacatgct tacatgggca 1080
aactttgggt tccctaacaa tatgtatggt ccttacaag acattcacgg tgatttaggt 1140
ggagatcatg aactccttcc agatttcac aaattttttg aagatgatta ctcagctttc 1200
acaggagata tcaagggaaa tgtgtatgat acaggaattg aatatactgt agcaccacat 1260

```

1931

```

gaacgttttaa tgtatgtgct ttcgcctatt actggaacaa cgataacaga tactgtttaca 1320
ttacgagcta aagtattaaa cgatgataac gcagttgtta cgtacagggt tgaagggttct 1380
gacgttgaac atgaaatgac gttagctgac tcgggatact acacagctaa gtattctccg 1440
acggcagaag taaatgggtg atcagttgat ttaacagtta cgtactgggc tggagaagaa 1500
aaagtacaag atgaagtgat tagactttat gtaaaggctt cagaaatctc actttacaag 1560
cttacgtttg atgaggatat taatggaatt aagtcgaatg gcacttggcc tgaagatggg 1620
attacatctg acgttttctca tgtcagtttt gacggaaatg ggaaattgaa gtttgcagtt 1680
aatggaatgt catccgaaga gtggtggcaa gaacttaa atagaattaac agatctttct 1740
gatgtgaatt tagccaagta a 1761

```

<210> 10

<211> 586

<212> PRT

<213> Bacillus halodurans

<400> 10

```

Met Lys Ser Ile Lys Lys Leu Val Val Val Cys Met Ala Phe Leu Leu
  1              5              10              15

```

```

Ile Phe Pro Ser Thr Ser Phe Ala Phe Ser Gly Ser Val Ser Ala Ser
              20              25              30

```

```

Gly Gln Glu Leu Lys Met Thr Asp Gln Asn Ala Ser Gln Tyr Thr Lys
              35              40              45

```

```

Glu Leu Phe Ala Phe Leu Arg Asp Val Ser Gly Lys Gln Val Leu Phe
  50              55              60

```

```

Gly Gln Gln His Ala Thr Asp Glu Gly Leu Thr Leu Arg Gly Thr Gly
  65              70              75              80

```

```

Asn Arg Ile Gly Ser Thr Glu Ser Glu Val Lys Asn Ala Val Gly Asp
              85              90              95

```

```

Tyr Pro Ala Val Phe Gly Trp Asp Thr Asn Ser Leu Asp Gly Arg Glu
              100              105              110

```

```

Lys Pro Gly Asn Asp Glu Pro Ser Gln Glu Gln Arg Ile Leu Asn Thr
              115              120              125

```

```

Ala Ala Ser Met Lys Ala Ala His Asp Leu Gly Gly Ile Ile Thr Leu
              130              135              140

```

```

Ser Met His Pro Asp Asn Phe Val Thr Gly Gly Ala Tyr Gly Asp Thr
  145              150              155              160

```

```

Thr Gly Asn Val Val Gln Glu Ile Leu Pro Gly Gly Ser Lys His Glu
              165              170              175

```

```

Glu Phe Asn Ala Trp Leu Asp Asn Leu Ala Ala Leu Ala His Glu Leu
              180              185              190

```

```

Lys Asp Asp Asn Gly Lys His Ile Pro Ile Ile Phe Arg Pro Phe His
              195              200              205

```

193m

Glu Gln Thr Gly Ser Trp Phe Trp Trp Gly Ala Ser Thr Thr Thr Pro
 210 215 220

Glu Gln Tyr Lys Ala Ile Tyr Arg Tyr Thr Val Glu Tyr Leu Arg Asp
 225 230 235 240

Val Lys Gly Ala Asn Asn Phe Leu Tyr Gly Phe Ser Pro Gly Ala Gly
 245 250 255

Pro Ala Gly Asp Leu Asn Arg Tyr Met Glu Thr Tyr Pro Gly Asp Asp
 260 265 270

Tyr Val Asp Ile Phe Gly Ile Asp Asn Tyr Asp Asn Lys Ser Asn Ala
 275 280 285

Gly Ser Glu Ala Trp Ile Gln Gly Val Val Thr Asp Leu Ala Met Leu
 290 295 300

Val Asp Leu Ala Glu Glu Lys Gly Lys Ile Ala Ala Phe Thr Glu Tyr
 305 310 315 320

Gly Tyr Ser Ala Thr Gly Met Asn Arg Thr Gly Asn Thr Leu Asp Trp
 325 330 335

Tyr Thr Arg Leu Leu Asn Ala Ile Lys Glu Asp Pro Lys Ala Ser Lys
 340 345 350

Ile Ser Tyr Met Leu Thr Trp Ala Asn Phe Gly Phe Pro Asn Asn Met
 355 360 365

Tyr Val Pro Tyr Lys Asp Ile His Gly Asp Leu Gly Gly Asp His Glu
 370 375 380

Leu Leu Pro Asp Phe Ile Lys Phe Phe Glu Asp Asp Tyr Ser Ala Phe
 385 390 395 400

Thr Gly Asp Ile Lys Gly Asn Val Tyr Asp Thr Gly Ile Glu Tyr Thr
 405 410 415

Val Ala Pro His Glu Arg Leu Met Tyr Val Leu Ser Pro Ile Thr Gly
 420 425 430

Thr Thr Ile Thr Asp Thr Val Thr Leu Arg Ala Lys Val Leu Asn Asp
 435 440 445

Asp Asn Ala Val Val Thr Tyr Arg Val Glu Gly Ser Asp Val Glu His
 450 455 460

Glu Met Thr Leu Ala Asp Ser Gly Tyr Tyr Thr Ala Lys Tyr Ser Pro
 465 470 475 480

Thr Ala Glu Val Asn Gly Gly Ser Val Asp Leu Thr Val Thr Tyr Trp
 485 490 495

Ser Gly Glu Glu Lys Val Gln Asp Glu Val Ile Arg Leu Tyr Val Lys
 500 505 510

193n

Ala Ser Glu Ile Ser Leu Tyr Lys Leu Thr Phe Asp Glu Asp Ile Asn
515 520 525

Gly Ile Lys Ser Asn Gly Thr Trp Pro Glu Asp Gly Ile Thr Ser Asp
530 535 540

Val Ser His Val Ser Phe Asp Gly Asn Gly Lys Leu Lys Phe Ala Val
545 550 555 560

Asn Gly Met Ser Ser Glu Glu Trp Trp Gln Glu Leu Lys Leu Glu Leu
565 570 575

Thr Asp Leu Ser Asp Val Asn Leu Ala Lys
580 585

<210> 11

<211> 995

<212> DNA

<213> Bacillus sp. AAI12

<400> 11

```

gtgtataagc ttaccatac gtattttggt gcgttaattt gttctatatt gatctttgct 60
ggggtttttaa atacttcttc ttcacaagca gaagcccatc acagtgggtt ccatgttaat 120
ggtacaacat tatatgatgc aaatggaaac ccttttggtta tgagagggat taatcatgga 180
catgcttggt ttaaacaaga actagaaaca tccatgagag ggattagtca aacaggggca 240
aatacgattc gtgtcgtttt gtctaattggg caaagatggc aaaaagatga tcgaaacatg 300
gtagcttcgg ttatttcttt ggcagagcag catcaaata ga ttgccgtttt agaagttcat 360
gatgctactg gtagcaataa tttctccgat ctgcaagctg ctgtggacta ttggattgag 420
atgaaggatg ttttgcaggg gaaagaggac atagtgatca ttaatatcgc caatgaatgg 480
tacgggtgctt gggacggagg cgcattgggca cgagggtatc agaatgcgat acgtcagctt 540
cgaaatgcag gcttgtcaca tacatttatg gttgacgctg ccggttatgg ccagtacct 600
caatcggtag ttgattatgg tcaagaagta ttaaattgctg acccacagag aaacacaatg 660
ttttctgttc atatgtatga atatgcaggc ggagatgcta atacagtaag acgaaacatt 720
gactcgatct taagccagaa cttagctctt gtcattggtg aattcgggca ttggcattat 780
gacggtgatg ttgatgagga caccatttta agctattcac agcaaagaaa tgtgggatgg 840
ttggcgtgga gctggcatgg caatagtga ggggtcgaat atcttgattt atcgaatgac 900
tttgctggta atcgactgac atgggtgggt gatcgaatag taaacgggtc gaatgggatt 960
cgtcaaacct ctaaaagaag cagtgtgttt caata 995

```

<210> 12

<211> 331

<212> PRT

<213> Bacillus sp. AAI12

<400> 12

Val Tyr Lys Leu Thr His Thr Tyr Phe Val Ala Leu Ile Cys Ser Ile
1 5 10 15

Leu Ile Phe Ala Gly Val Leu Asn Thr Ser Ser Ser Gln Ala Glu Ala
20 25 30

His	His	Ser	Gly	Phe	His	Val	Asn	Gly	Thr	Thr	Leu	Tyr	Asp	Ala	Asn
		35					40					45			
Gly	Asn	Pro	Phe	Val	Met	Arg	Gly	Ile	Asn	His	Gly	His	Ala	Trp	Phe
	50					55					60				
Lys	Gln	Glu	Leu	Glu	Thr	Ser	Met	Arg	Gly	Ile	Ser	Gln	Thr	Gly	Ala
65					70					75					80
Asn	Thr	Ile	Arg	Val	Val	Leu	Ser	Asn	Gly	Gln	Arg	Trp	Gln	Lys	Asp
				85					90					95	
Asp	Arg	Asn	Met	Val	Ala	Ser	Val	Ile	Ser	Leu	Ala	Glu	Gln	His	Gln
			100					105					110		
Met	Ile	Ala	Val	Leu	Glu	Val	His	Asp	Ala	Thr	Gly	Ser	Asn	Asn	Phe
		115					120					125			
Ser	Asp	Leu	Gln	Ala	Ala	Val	Asp	Tyr	Trp	Ile	Glu	Met	Lys	Asp	Val
	130					135					140				
Leu	Gln	Gly	Lys	Glu	Asp	Ile	Val	Ile	Ile	Asn	Ile	Ala	Asn	Glu	Trp
145					150					155					160
Tyr	Gly	Ala	Trp	Asp	Gly	Gly	Ala	Trp	Ala	Arg	Gly	Tyr	Gln	Asn	Ala
				165					170					175	
Ile	Arg	Gln	Leu	Arg	Asn	Ala	Gly	Leu	Ser	His	Thr	Phe	Met	Val	Asp
			180					185					190		
Ala	Ala	Gly	Tyr	Gly	Gln	Tyr	Pro	Gln	Ser	Val	Val	Asp	Tyr	Gly	Gln
		195					200					205			
Glu	Val	Leu	Asn	Ala	Asp	Pro	Gln	Arg	Asn	Thr	Met	Phe	Ser	Val	His
	210					215					220				
Met	Tyr	Glu	Tyr	Ala	Gly	Gly	Asp	Ala	Asn	Thr	Val	Arg	Arg	Asn	Ile
225					230					235					240
Asp	Ser	Ile	Leu	Ser	Gln	Asn	Leu	Ala	Leu	Val	Ile	Gly	Glu	Phe	Gly
				245					250					255	
His	Trp	His	Tyr	Asp	Gly	Asp	Val	Asp	Glu	Asp	Thr	Ile	Leu	Ser	Tyr
			260					265					270		
Ser	Gln	Gln	Arg	Asn	Val	Gly	Trp	Leu	Ala	Trp	Ser	Trp	His	Gly	Asn
		275					280					285			
Ser	Glu	Gly	Val	Glu	Tyr	Leu	Asp	Leu	Ser	Asn	Asp	Phe	Ala	Gly	Asn
	290					295					300				
Arg	Leu	Thr	Trp	Trp	Gly	Asp	Arg	Ile	Val	Asn	Gly	Pro	Asn	Gly	Ile
305					310					315					320
Arg	Gln	Thr	Ser	Lys	Arg	Ser	Ser	Val	Phe	Gln					
				325					330						

193p

<210> 13
 <211> 1464
 <212> DNA
 <213> Humicola insolens

<400> 13
 atggcaaagg ctctgaagta ctttgccctgg ggccttgctg ccctggcctc gggcgctggt 60
 gccgctcctt actgtgctcc ccagccgctc acaacctctc aggagcctac gagcactccg 120
 tcgcctgtgc ccggtccgcg gaccttcgaa gcggaggatg ccatcctcac gggcacgagg 180
 gttgagtcga gcctcgccgg ctactctggt accggatatg tagcgggctt cgacgagccc 240
 agtgacaaga tcacgttcca cgtggacagc gagaccacac ggctgtacga cctcaccatc 300
 cgcggtggccg ccatctatgg cgagaagcgc accaccgtcg tgctcaataa cggcgcgcca 360
 agtgaggtct acttcccggc aggcgattcg ttcgtcgaca tcgctgccgg ccaggctcctg 420
 ctgaaccagg gcgacaacac catcgacatt gtcaacaact ggggatggta cctgatcgac 480
 tccatcacca tcaccccctc cgccccgcga cccctcacc aaatcaacce ttcccccgctc 540
 aaccctgccg ccgacgacaa cgcgcgggcg ttgtacgcat acctccgctc catctacggc 600
 aagaaaatcc tttccggcca gcaggagctt tcctgggcca actggatcgc ccaacagacg 660
 ggcaaaactc ccgcgctggt gtcgctcgat atgatggatt attcccctag tcgggtggaa 720
 agaggcactg tcgggtctgc cgtcgaggag gccatcgagc atcacccggc cggcgccatt 780
 gtctcggtgt tgtggcactg gaacgcgcc acggggctgt acgacacgcc cgagcgccgg 840
 tgggtggagcg ggttctacac ggacgcgacc gactttgacg tcgcgcgggc gctggcggat 900
 acgacgaatg ccaactacac gctgctgatc cgggatatcg acgcgatcgc ggtgcagctc 960
 aagaggttgc gggacgcggg cgtgccgggtg ctttggcgcc cgctgcacga ggccgagggc 1020
 ggttggtttt ggtggggagc gaagggcccg gaggcataca agaagctgtg ggggattctg 1080
 tatgaccgac tcacgaacta ccatgggctg aataacctgc tgtgggtgtg gaactcgatc 1140
 ctacccgagt ggtatcccgg agacgaaaca gtagacattg tcagcgcgga cgtgtacgcg 1200
 cagggtaatg ggcccatgtc gacgcagtat aaccagctca tcgagctggg caaggacaag 1260
 aagatgatcg cggcgactga ggtcggggcc gcgccgctgc cggacctgtt gcaggcctat 1320
 gaggctcact ggttgtgggt cgctgttttg ggagacacgt tcatacaaa ccctcagtgg 1380
 aactcgatcg agaccttgaa gacgatctac aatagcgact atgttctcac tctcgatgag 1440
 attcaggggt ggaggaacgc gcaa 1464

<210> 14
 <211> 488
 <212> PRT
 <213> Humicola insolens

<400> 14
 Met Ala Lys Ala Leu Lys Tyr Phe Ala Trp Gly Leu Ala Ala Leu Ala
 1 5 10 15
 Ser Gly Ala Val Ala Ala Pro Tyr Cys Ala Pro Gln Pro Ser Thr Thr
 20 25 30
 Ser Gln Glu Pro Thr Ser Thr Pro Ser Pro Val Pro Gly Pro Arg Thr
 35 40 45
 Phe Glu Ala Glu Asp Ala Ile Leu Thr Gly Thr Arg Val Glu Ser Ser
 50 55 60
 Leu Ala Gly Tyr Ser Gly Thr Gly Tyr Val Ala Gly Phe Asp Glu Pro
 65 70 75 80

193q

Ser	Asp	Lys	Ile	Thr	Phe	His	Val	Asp	Ser	Glu	Thr	Thr	Arg	Leu	Tyr	85	90	95	
Asp	Leu	Thr	Ile	Arg	Val	Ala	Ala	Ile	Tyr	Gly	Glu	Lys	Arg	Thr	Thr	100	105	110	
Val	Val	Leu	Asn	Asn	Gly	Ala	Ala	Ser	Glu	Val	Tyr	Phe	Pro	Ala	Gly	115	120	125	
Asp	Ser	Phe	Val	Asp	Ile	Ala	Ala	Gly	Gln	Val	Leu	Leu	Asn	Gln	Gly	130	135	140	
Asp	Asn	Thr	Ile	Asp	Ile	Val	Asn	Asn	Trp	Gly	Trp	Tyr	Leu	Ile	Asp	145	150	155	160
Ser	Ile	Thr	Ile	Thr	Pro	Ser	Ala	Pro	Arg	Pro	Pro	His	Gln	Ile	Asn	165	170	175	
Pro	Ser	Pro	Val	Asn	Pro	Ala	Ala	Asp	Asp	Asn	Ala	Arg	Ala	Leu	Tyr	180	185	190	
Ala	Tyr	Leu	Arg	Ser	Ile	Tyr	Gly	Lys	Lys	Ile	Leu	Ser	Gly	Gln	Gln	195	200	205	
Glu	Leu	Ser	Trp	Ala	Asn	Trp	Ile	Ala	Gln	Gln	Thr	Gly	Lys	Thr	Pro	210	215	220	
Ala	Leu	Val	Ser	Val	Asp	Met	Met	Asp	Tyr	Ser	Pro	Ser	Arg	Val	Glu	225	230	235	240
Arg	Gly	Thr	Val	Gly	Ser	Ala	Val	Glu	Glu	Ala	Ile	Glu	His	His	Arg	245	250	255	
Arg	Gly	Gly	Ile	Val	Ser	Val	Leu	Trp	His	Trp	Asn	Ala	Pro	Thr	Gly	260	265	270	
Leu	Tyr	Asp	Thr	Pro	Glu	Arg	Arg	Trp	Trp	Ser	Gly	Phe	Tyr	Thr	Asp	275	280	285	
Ala	Thr	Asp	Phe	Asp	Val	Ala	Arg	Ala	Leu	Ala	Asp	Thr	Thr	Asn	Ala	290	295	300	
Asn	Tyr	Thr	Leu	Leu	Ile	Arg	Asp	Ile	Asp	Ala	Ile	Ala	Val	Gln	Leu	305	310	315	320
Lys	Arg	Leu	Arg	Asp	Ala	Gly	Val	Pro	Val	Leu	Trp	Arg	Pro	Leu	His	325	330	335	
Glu	Ala	Glu	Gly	Gly	Trp	Phe	Trp	Trp	Gly	Ala	Lys	Gly	Pro	Glu	Ala	340	345	350	
Tyr	Lys	Lys	Leu	Trp	Gly	Ile	Leu	Tyr	Asp	Arg	Leu	Thr	Asn	Tyr	His	355	360	365	
Gly	Leu	Asn	Asn	Leu	Leu	Trp	Val	Trp	Asn	Ser	Ile	Leu	Pro	Glu	Trp	370	375	380	

193r

Tyr Pro Gly Asp Glu Thr Val Asp Ile Val Ser Ala Asp Val Tyr Ala
385 390 395 400

Gln Gly Asn Gly Pro Met Ser Thr Gln Tyr Asn Gln Leu Ile Glu Leu
405 410 415

Gly Lys Asp Lys Lys Met Ile Ala Ala Thr Glu Val Gly Ala Ala Pro
420 425 430

Leu Pro Asp Leu Leu Gln Ala Tyr Glu Ala His Trp Leu Trp Phe Ala
435 440 445

Val Trp Gly Asp Thr Phe Ile Asn Asn Pro Gln Trp Asn Ser Ile Glu
450 455 460

Thr Leu Lys Thr Ile Tyr Asn Ser Asp Tyr Val Leu Thr Leu Asp Glu
465 470 475 480

Ile Gln Gly Trp Arg Asn Ala Gln
485

<210> 15

<211> 1107

<212> DNA

<213> Bacillus sp. AA349

<400> 15

```

atgagaagta tgaagctttt atttgctatg tttatttttag ttttttcttc ttttactttt 60
aacttagtag ttgcgcaagc tagtggacat gggcaaagtc ataaagtacc ttgggcacca 120
caagctgaag cacctggaaa aacggctgaa aatggagtct gggataaagt tcgaaataat 180
cctggaaaag ccaatcctcc agcaggaaaa gtcaatgggt tttatataga tggaacaacc 240
ttatatgatg caaatggtaa gccatttgtg atgctgtgaa ttaaccacgg tcattcatgg 300
tacaagcctc acatagaaac cgcgatggag gcaattgctg atactggagc aaactccatt 360
cgtgtagttc tctcagatgg acaacagtgg accaaagatg atgttgacga agtagcaaaa 420
attatatctt tagcagaaaa acattcttta gttgctgctc ttgaggtaca tgatgcactc 480
ggaacagatg atattgaacc attacttaaa acagttgatt actggattga gatcaaagat 540
gctttaatcg gaaaagagga caaagtaatt attaacattt ctaatgaatg gtttggttct 600
tggagcagtg aagggtgggc agatggatat aaaaaagcaa ttcctttact aagagaggcg 660
ggtctaaaac atacacttat ggttgacgca gctgggtggg gacaatttcc tagatctatt 720
catgaaaaag gattagaagt ttttaactca gaccatttaa agaatacaat gttttccatt 780
catatgtatg aatgggcagc gggtaatcct caacaagtaa aagacaatat tgacgggtgtt 840
cttgaaaaga atttagctgt agtaattggg gagttcggtc atcatcacta cggaagagat 900
gttgctgttg atacaatctt aagtcattct gagaagtatg atgtaggttg gcttgcttgg 960
tcttggcacg gaaatagtgg tgggtgtagag tatcttgact tagcaacaga tttctcaggg 1020
acacaactaa ctgaatgggg agaaagaatt gtacacggtc cgaatgggtt aaaagaaact 1080
tctgaaatcg ttagtgtata caaaaaa 1107

```

<210> 16

<211> 369

<212> PRT

<213> Bacillus sp. AA349

193s

<400> 16

Met	Arg	Ser	Met	Lys	Leu	Leu	Phe	Ala	Met	Phe	Ile	Leu	Val	Phe	Ser
1				5					10					15	
Ser	Phe	Thr	Phe	Asn	Leu	Val	Val	Ala	Gln	Ala	Ser	Gly	His	Gly	Gln
			20					25					30		
Met	His	Lys	Val	Pro	Trp	Ala	Pro	Gln	Ala	Glu	Ala	Pro	Gly	Lys	Thr
		35					40					45			
Ala	Glu	Asn	Gly	Val	Trp	Asp	Lys	Val	Arg	Asn	Asn	Pro	Gly	Lys	Ala
	50					55					60				
Asn	Pro	Pro	Ala	Gly	Lys	Val	Asn	Gly	Phe	Tyr	Ile	Asp	Gly	Thr	Thr
65					70					75					80
Leu	Tyr	Asp	Ala	Asn	Gly	Lys	Pro	Phe	Val	Met	Arg	Gly	Ile	Asn	His
				85					90					95	
Gly	His	Ser	Trp	Tyr	Lys	Pro	His	Ile	Glu	Thr	Ala	Met	Glu	Ala	Ile
			100					105					110		
Ala	Asp	Thr	Gly	Ala	Asn	Ser	Ile	Arg	Val	Val	Leu	Ser	Asp	Gly	Gln
		115					120					125			
Gln	Trp	Thr	Lys	Asp	Asp	Val	Asp	Glu	Val	Ala	Lys	Ile	Ile	Ser	Leu
	130					135					140				
Ala	Glu	Lys	His	Ser	Leu	Val	Ala	Ala	Leu	Glu	Val	His	Asp	Ala	Leu
145					150					155					160
Gly	Thr	Asp	Asp	Ile	Glu	Pro	Leu	Leu	Lys	Thr	Val	Asp	Tyr	Trp	Ile
				165					170					175	
Glu	Ile	Lys	Asp	Ala	Leu	Ile	Gly	Lys	Glu	Asp	Lys	Val	Ile	Ile	Asn
			180					185					190		
Ile	Ser	Asn	Glu	Trp	Phe	Gly	Ser	Trp	Ser	Ser	Glu	Gly	Trp	Ala	Asp
		195					200					205			
Gly	Tyr	Lys	Lys	Ala	Ile	Pro	Leu	Leu	Arg	Glu	Ala	Gly	Leu	Lys	His
	210					215					220				
Thr	Leu	Met	Val	Asp	Ala	Ala	Gly	Trp	Gly	Gln	Phe	Pro	Arg	Ser	Ile
225					230					235					240
His	Glu	Lys	Gly	Leu	Glu	Val	Phe	Asn	Ser	Asp	Pro	Leu	Lys	Asn	Thr
				245					250					255	
Met	Phe	Ser	Ile	His	Met	Tyr	Glu	Trp	Ala	Ala	Gly	Asn	Pro	Gln	Gln
			260					265					270		
Val	Lys	Asp	Asn	Ile	Asp	Gly	Val	Leu	Glu	Lys	Asn	Leu	Ala	Val	Val
		275					280					285			

193t

Ile Gly Glu Phe Gly His His His Tyr Gly Arg Asp Val Ala Val Asp
 290 295 300

Thr Ile Leu Ser His Ser Glu Lys Tyr Asp Val Gly Trp Leu Ala Trp
 305 310 315 320

Ser Trp His Gly Asn Ser Gly Gly Val Glu Tyr Leu Asp Leu Ala Thr
 325 330 335

Asp Phe Ser Gly Thr Gln Leu Thr Glu Trp Gly Glu Arg Ile Val His
 340 345 350

Gly Pro Asn Gly Leu Lys Glu Thr Ser Glu Ile Val Ser Val Tyr Lys
 355 360 365

Lys

<210> 17

<211> 915

<212> DNA

<213> Bacillus sp.

<400> 17

```

atctcaacac tcagaaatgc cggatttcgc aatacaatcg ttgtggatgc atcgggggtgg 60
ggacaaaatt catcgccaat taaagcttat ggcaacgaag tgttaaacca tgatccgcag 120
cgcaatgtta tgttctccat acacatgtac ggttcctgga ataatcagtc gcgaatcggc 180
agcgaattgc aggccatcaa agaccttggg cttgctgtca tgattgggtga attcgggatac 240
aactacaaca acggcaataa caacttgggg agtcagggtta acgcccagga aatcatgaat 300
caggcgcaag caaaaggaat cggctacatg ccgtggtcgt ggactggcaa tgacgcggct 360
aactcttggg tggatatgac aacaaacgat tggcaaacac ttacatcatg ggggaatcta 420
gttgtaaatg gaaccaacgg cattcgagct acgtctgtcc cagcaactgt atttaataca 480
caaacaacaa tttatgattt tgaaggcggc aatgcccagg gctggtcagg ttccggtttg 540
agcggggggc cttgggtctgt taatgaatgg gcggcgagcg gtagttattc tctcaaagcg 600
aatatatctc taggcgccac tcaaaaagct ttgcaaacca cagcgtccca taatttcagc 660
ggccgggtcta cattatccgt aagagtaaag catgcagcat ggggaaatca cggcagcggt 720
atgcaagcca agttatatgt gaaaacaggg gccggttacg cctgggtatga tggcggcact 780
gtaaacatca acagctcggg caacacattg acgctaaacc tggcaggcat tcctaattctg 840
aacgacgtca gagaactcgg aattgaattt ataacacctg caaattcgag tggttctttc 900
gcaatttatg ttgac 915

```

<210> 18

<211> 305

<212> PRT

<213> Bacillus sp.

<400> 18

Ile Ser Thr Leu Arg Asn Ala Gly Ile Arg Asn Thr Ile Val Val Asp
 1 5 10 15

Ala Ser Gly Trp Gly Gln Asn Ser Ser Pro Ile Lys Ala Tyr Gly Asn
 20 25 30

[illegible]

193v

<210> 19
 <211> 397
 <212> DNA
 <213> *Bacillus clausii*

<400> 19
 atctctcagg gcttggtagg agtcattatt ctcttataca tggcatttag tcaagagaga 60
 ggattggcgc aaactggatt tcaagtaaca gggacccagt tgcttgatgg agagggcaat 120
 ccgtatgtga tgcgtggagt caatcacgga cattcatggt tcaaacaaga ccttgatata 180
 gcaataccag ctattgcagc gactggcgct aatacgggtga gaatcgtttt atcgaatggc 240
 caacaatggg agcgagatac cgtagcggaa gttgaaagag tgcttgcagt taccgaagag 300
 gaaggcttga cggctgtact tgaagttcat gatgcgacgg gaagtgatga tccaaacgat 360
 ttgtttactg cagtggagta ttggtcagag agaggat 397

<210> 20
 <211> 132
 <212> PRT
 <213> *Bacillus clausii*

<400> 20
 Ile Ser Gln Gly Leu Val Gly Val Ile Ile Leu Leu Tyr Met Ala Phe
 1 5 10 15
 Ser Gln Glu Arg Gly Leu Ala Gln Thr Gly Phe Gln Val Thr Gly Thr
 20 25 30
 Gln Leu Leu Asp Gly Glu Gly Asn Pro Tyr Val Met Arg Gly Val Asn
 35 40 45
 His Gly His Ser Trp Phe Lys Gln Asp Leu Asp Thr Ala Ile Pro Ala
 50 55 60
 Ile Ala Ala Thr Gly Ala Asn Thr Val Arg Ile Val Leu Ser Asn Gly
 65 70 75 80
 Gln Gln Trp Glu Arg Asp Thr Val Ala Glu Val Glu Arg Val Leu Ala
 85 90 95
 Val Thr Glu Glu Glu Gly Leu Thr Ala Val Leu Glu Val His Asp Ala
 100 105 110
 Thr Gly Ser Asp Asp Pro Asn Asp Leu Phe Thr Ala Val Glu Tyr Trp
 115 120 125
 Ser Glu Arg Gly
 130

<210> 21
 <211> 960
 <212> DNA
 <213> *Bacillus* sp.

193w

<400> 21

```

atgaatcgta agcgggttaca atggggttgga gcactagtgg tgggtggtggg tttgttttgta 60
tacagtagcg gtttagcatc tgcacaaagc ggctttcacg taaaagggtac agagttggtg 120
gacaaaaatg gcgatcctta cgttatgcgt ggcgtcaacc atggacattc ttggttttaa 180
caagatttag aggaggcaat ccctgccata gcagaaacag gggcgaacac agtgagaatc 240
gtcttatcca atggacagca atgggaaaaa gatgatgcct ctgagcttgc ccgtgtgctt 300
gctgccacag aaacatatgg gttgacaacc gtgctggaag tccacgatgc tacaggaagt 360
gataatcccg atgatttaga taaagcagtc gattactgga tcgaaatggc tgatgttcta 420
aaggggacag aagaccgggt aatcattaac attgccaatg aatggtatgg ggcgtggagg 480
agtgacgttt gggcagaggc atacgcacaa gcgatcccg cgttgcgag tgctggcctc 540
gcccatacgt taatagttga tgcggcaggt tggggacagt accctgcctc tatccatgag 600
cggggagccg acgtatttgc ctccgatcca ttaaaaaaca caatgttttc catccatatg 660
tacgaatatg caggagcgga tagggcgaca gtttctgaaa acatcgacgg tgtacttgct 720
gaaaatcttg ctgtggtaat cgggtgaattt ggccataggc atcatgatgg cgatgtcgat 780
gaagatgcga ttttggccta tacagcagag cggcaagtgg gctggcttgc ctggatcatg 840
tatggcaata gcgggggtgt tgaatacttg gatttaactg aaggcccatc aggtccatta 900
acgagttggg gcgaacggat tgtctatggg gaaatgggct taaaagtaat tgatcacttg 960

```

<210> 22

<211> 320

<212> PRT

<213> Bacillus sp.

<400> 22

```

Met Asn Arg Lys Arg Leu Gln Trp Val Gly Ala Leu Val Val Val Leu
  1              5              10              15

Val Leu Phe Val Tyr Ser Ser Gly Leu Ala Ser Ala Gln Ser Gly Phe
      20              25              30

His Val Lys Gly Thr Glu Leu Leu Asp Lys Asn Gly Asp Pro Tyr Val
      35              40              45

Met Arg Gly Val Asn His Gly His Ser Trp Phe Lys Gln Asp Leu Glu
      50              55              60

Glu Ala Ile Pro Ala Ile Ala Glu Thr Gly Ala Asn Thr Val Arg Ile
      65              70              75              80

Val Leu Ser Asn Gly Gln Gln Trp Glu Lys Asp Asp Ala Ser Glu Leu
      85              90              95

Ala Arg Val Leu Ala Ala Thr Glu Thr Tyr Gly Leu Thr Thr Val Leu
      100              105              110

Glu Val His Asp Ala Thr Gly Ser Asp Asn Pro Asp Asp Leu Asp Lys
      115              120              125

Ala Val Asp Tyr Trp Ile Glu Met Ala Asp Val Leu Lys Gly Thr Glu
      130              135              140

Asp Arg Val Ile Ile Asn Ile Ala Asn Glu Trp Tyr Gly Ala Trp Arg
      145              150              155              160

```

193x

Ser Asp Val Trp Ala Glu Ala Tyr Ala Gln Ala Ile Pro Arg Leu Arg
165 170 175

Ser Ala Gly Leu Ala His Thr Leu Ile Val Asp Ala Ala Gly Trp Gly
180 185 190

Gln Tyr Pro Ala Ser Ile His Glu Arg Gly Ala Asp Val Phe Ala Ser
195 200 205

Asp Pro Leu Lys Asn Thr Met Phe Ser Ile His Met Tyr Glu Tyr Ala
210 215 220

Gly Ala Asp Arg Ala Thr Val Ser Glu Asn Ile Asp Gly Val Leu Ala
225 230 235 240

Glu Asn Leu Ala Val Val Ile Gly Glu Phe Gly His Arg His His Asp
245 250 255

Gly Asp Val Asp Glu Asp Ala Ile Leu Ala Tyr Thr Ala Glu Arg Gln
260 265 270

Val Gly Trp Leu Ala Trp Ser Trp Tyr Gly Asn Ser Gly Gly Val Glu
275 280 285

Tyr Leu Asp Leu Thr Glu Gly Pro Ser Gly Pro Leu Thr Ser Trp Gly
290 295 300

Glu Arg Ile Val Tyr Gly Glu Met Gly Leu Lys Val Ile Asp His Leu
305 310 315 320

<210> 23
<211> 564
<212> DNA
<213> Bacillus sp.

<400> 23
atgaatcgta agcgggttaca atggggttgga gcactagtgg cgggtgttggt tttgtttgta 60
tacagtagcg gtttagcatc tgcacaaagc ggcttttcacg taaaagggtac agagttgttg 120
gacaaaaatg gcgatcctta cgttatgcgt ggcgtcaacc atggacattc ttggttttaa 180
caagatttag aggaggcaat ccctgccata gcagaaacag gggcgaacac agtgagaatc 240
gtcttatcca atggacagca atgggaaaaa gatgatgcct ctgagcttgc ccgtgtgctt 300
gctgccacag aaacatatgg gttgacaacc gtgctggaag tccacgatgc tacaggaagt 360
gataatcccg atgatttaga taaagcagtc gattactgga tcgaaatggc tgatgttcta 420
aaggggacag aagaccgggt aatcattaac attgccaatg aatgggtatgg ggcgtggagg 480
agtgaccttt gggcaaaagc atacgcacaa gcgatcccg cgttgccgag tgctggcctc 540
gcccatacgt taataattga tgcc 564

<210> 24
<211> 188
<212> PRT
<213> Bacillus sp.

193y

<400> 24

Met Asn Arg Lys Arg Leu Gln Trp Val Gly Ala Leu Val Ala Val Leu
 1 5 10 15

Val Leu Phe Val Tyr Ser Ser Gly Leu Ala Ser Ala Gln Ser Gly Phe
 20 25 30

His Val Lys Gly Thr Glu Leu Leu Asp Lys Asn Gly Asp Pro Tyr Val
 35 40 45

Met Arg Gly Val Asn His Gly His Ser Trp Phe Lys Gln Asp Leu Glu
 50 55 60

Glu Ala Ile Pro Ala Ile Ala Glu Thr Gly Ala Asn Thr Val Arg Ile
 65 70 75 80

Val Leu Ser Asn Gly Gln Gln Trp Glu Lys Asp Asp Ala Ser Glu Leu
 85 90 95

Ala Arg Val Leu Ala Ala Thr Glu Thr Tyr Gly Leu Thr Thr Val Leu
 100 105 110

Glu Val His Asp Ala Thr Gly Ser Asp Asn Pro Asp Asp Leu Asp Lys
 115 120 125

Ala Val Asp Tyr Trp Ile Glu Met Ala Asp Val Leu Lys Gly Thr Glu
 130 135 140

Asp Arg Val Ile Ile Asn Ile Ala Asn Glu Trp Tyr Gly Ala Trp Arg
 145 150 155 160

Ser Asp Leu Trp Ala Lys Ala Tyr Ala Gln Ala Ile Pro Arg Leu Arg
 165 170 175

Ser Ala Gly Leu Ala His Thr Leu Ile Ile Asp Ala
 180 185

<210> 25

<211> 2445

<212> DNA

<213> Bacillus sp.

<400> 25

atgaacaaac aaccgttaaa gactgcattt attatgttgt tatgtagcgt gtttatgttt 60
 caaagcctac cttactatgt gaacgctatc aatgaaggcg agagagaagc ttttgcaccc 120
 gcagggagat atgatgctga acaggcgact acgacaggaa atgccgtatt cacgaccgag 180
 cctgttgagg acggcgagta cgccgggtccg ggctacattt ccttccttttc tgaagattcc 240
 tcgccacctt cttcatcgac aacctttcac attcaggccg ataaaacgga gctctatcat 300
 ttatctatcg gatactatgc tccatacggg aacaaggga ccacaattct ggtgaacggt 360
 gcaggtaacg gagagtttat gttgccagcg cccgaggacg gggcagtctc cgccgaagtg 420
 gaaattagca aaatcctgct cgaagaagga aataatacga ttacattcac aagaggctgg 480
 ggttattacg gcattgaata tattcgggtc gagccggtta atccaacgtt accgactata 540
 ttatttgaag cagaagaaga ttacgaagcg actggaaatg ttagcggtac caatgaaatc 600
 gaaggttatt ccggagcagg ctatttggtc aaccaagagg ggacaattca ttggaatgta 660

193z

```

acctcaccgg aaacctctat atatgaagta atcggtgcct atgcagctcc ttatggcgac 720
aaacaaacaa atctgacagt gaatggacag ggtaccgtca atcttgactt gaaagagaca 780
gaagtcttcg tggagttgaa tgtcggcatc gtaagtctca atgaaggcga aaacacacta 840
acactccata gtggttgggg atggtacaat atcgattata tcaagcttgt acctgtggtc 900
agttcggatc ccgaaccgca tcaggtcgaa aaaacactgg tgaatccgga cgcctcacct 960
gaggcaagag cgctaattaa ttatctcgta gaccagtacg ggaacaaaat tctatcaggt 1020
caaaccgagt tgaaagacgc caggtggatc catgaacagg tgggcaaata tcctgcggtt 1080
atggcagttg attttatgga ctacagcccg tcccgcgtag tgcattggcg aactggaact 1140
gcggttgagg aagcgattga gtgggcagag atgggtggga tcattacctt ccactggcat 1200
tggaacgcgc caaaggacct gcttaatgta cccggcaatg agtgggtggc cgggtttttat 1260
acctgtgcca caacgtttga tgtggagtac gcttttagaga accgggaatc tgaggatttc 1320
caattgttga ttagcgacat ggatgtgatc gccgagcaat tgaagcggct gcaggcagag 1380
aacatccctg tgttatggag accgcttcat gaggcggaag gcggctggtt ctggtggggc 1440
gccaaaggtc cagaggcggc aatagagctc tacaggctga tgtacgatcg ttacaccaat 1500
caccataaac taaacaattt gatatggatg tggaattcgg aagcgggaaga atggtatccg 1560
ggcgatgatg tcgtggacat gatcagtagc gatatttata atcctgtcgg agatttcagt 1620
cccagcatca acaagtatga gcatctaaag gaattggtac aggataagaa gctggttgcc 1680
ttgcctgaaa ccggcattat tccggatccc gatcagcttc agctgttcaa tgcgaactgg 1740
agttgggttc ccacctggac tggagactat atcagggacg gcactctcaa ccctatagaa 1800
cacctgcaaa aggtgtttca tcatgactac gtcatcacc tggatgaatt gccggagaac 1860
ctgtcccgtt acggtattat tgaaggagtc tggaagagcg acgcgatct atccgtaaaa 1920
acgaggacga cctccgaaat tacagtgaac tgggtcaaat ccattcaata tgattccgtt 1980
aatggctata aattaattaa agatggtgta gagaccgttt cagttgaagg cggcgtgcaa 2040
gagtatacct tcacaaattt attgccgggc acgcagtata cgataaaaag agaggcactg 2100
gaccaggatg accgatggac cgccgacgga ccggtcgccg ttgtatctac attatccaac 2160
gctccgatat cctatcctcc ggctgtcact cctgatgagc cgaatgaaga actgtcggag 2220
ggagagtata cgctcttggc agatgactta tccagccagg atggtgttct ggaagtaagt 2280
cttgagccga cagttacgaa gctcattatt ccttctgcac tagccggcac attagacgga 2340
gacttgagaa tcggttatgg ggacgtctgg atcgtcatcc cacacgaaca gcttgggggt 2400
gacgagcagc aatccggcag cgcgtatgag ttagtgctgg agatc 2445

```

<210> 26

<211> 815

<212> PRT

<213> Bacillus sp.

<400> 26

```

Met Asn Lys Gln Pro Leu Lys Thr Ala Phe Ile Met Leu Leu Cys Ser
  1                      5                      10                      15

```

```

Val Phe Met Phe Gln Ser Leu Pro Tyr Tyr Val Asn Ala Ile Asn Glu
      20                      25                      30

```

```

Gly Glu Arg Glu Ala Phe Ala Ser Ala Gly Arg Tyr Asp Ala Glu Gln
      35                      40                      45

```

```

Ala Thr Thr Thr Gly Asn Ala Val Phe Thr Thr Glu Pro Val Glu Asp
      50                      55                      60

```

```

Gly Glu Tyr Ala Gly Pro Gly Tyr Ile Ser Phe Phe Ser Glu Asp Ser
      65                      70                      75                      80

```

```

Ser Pro Pro Ser Ser Ser Thr Thr Phe His Ile Gln Ala Asp Lys Thr
      85                      90                      95

```

193aa

Glu	Leu	Tyr	His	Leu	Ser	Ile	Gly	Tyr	Tyr	Ala	Pro	Tyr	Gly	Asn	Lys	100	105	110
Gly	Thr	Thr	Ile	Leu	Val	Asn	Gly	Ala	Gly	Asn	Gly	Glu	Phe	Met	Leu	115	120	125
Pro	Ala	Pro	Glu	Asp	Gly	Ala	Val	Ser	Ala	Glu	Val	Glu	Ile	Ser	Lys	130	135	140
Ile	Leu	Leu	Glu	Glu	Gly	Asn	Asn	Thr	Ile	Thr	Phe	Thr	Arg	Gly	Trp	145	150	155
Gly	Tyr	Tyr	Gly	Ile	Glu	Tyr	Ile	Arg	Val	Glu	Pro	Val	Asn	Pro	Thr	165	170	175
Leu	Pro	Thr	Ile	Phe	Ile	Glu	Ala	Glu	Glu	Asp	Tyr	Glu	Ala	Thr	Gly	180	185	190
Asn	Val	Ser	Val	Thr	Asn	Glu	Ile	Glu	Gly	Tyr	Ser	Gly	Ala	Gly	Tyr	195	200	205
Leu	Phe	Asn	Gln	Glu	Gly	Thr	Ile	His	Trp	Asn	Val	Thr	Ser	Pro	Glu	210	215	220
Thr	Ser	Ile	Tyr	Glu	Val	Ile	Val	Ala	Tyr	Ala	Ala	Pro	Tyr	Gly	Asp	225	230	235
Lys	Gln	Thr	Asn	Leu	Thr	Val	Asn	Gly	Gln	Gly	Thr	Val	Asn	Leu	Asp	245	250	255
Leu	Lys	Glu	Thr	Glu	Val	Phe	Val	Glu	Leu	Asn	Val	Gly	Ile	Val	Ser	260	265	270
Leu	Asn	Glu	Gly	Glu	Asn	Thr	Leu	Thr	Leu	His	Ser	Gly	Trp	Gly	Trp	275	280	285
Tyr	Asn	Ile	Asp	Tyr	Ile	Lys	Leu	Val	Pro	Val	Val	Ser	Ser	Asp	Pro	290	295	300
Glu	Pro	His	Gln	Val	Glu	Lys	Thr	Leu	Val	Asn	Pro	Asp	Ala	Ser	Pro	305	310	315
Glu	Ala	Arg	Ala	Leu	Ile	Asn	Tyr	Leu	Val	Asp	Gln	Tyr	Gly	Asn	Lys	325	330	335
Ile	Leu	Ser	Gly	Gln	Thr	Glu	Leu	Lys	Asp	Ala	Arg	Trp	Ile	His	Glu	340	345	350
Gln	Val	Gly	Lys	Tyr	Pro	Ala	Val	Met	Ala	Val	Asp	Phe	Met	Asp	Tyr	355	360	365
Ser	Pro	Ser	Arg	Val	Val	His	Gly	Ala	Thr	Gly	Thr	Ala	Val	Glu	Glu	370	375	380
Ala	Ile	Glu	Trp	Ala	Glu	Met	Gly	Gly	Ile	Ile	Thr	Phe	His	Trp	His	385	390	395

193bb

Trp	Asn	Ala	Pro	Lys	Asp	Leu	Leu	Asn	Val	Pro	Gly	Asn	Glu	Trp	Trp	405	410	415	
Ser	Gly	Phe	Tyr	Thr	Arg	Ala	Thr	Thr	Phe	Asp	Val	Glu	Tyr	Ala	Leu	420	425	430	
Glu	Asn	Arg	Glu	Ser	Glu	Asp	Phe	Gln	Leu	Leu	Ile	Ser	Asp	Met	Asp	435	440	445	
Val	Ile	Ala	Glu	Gln	Leu	Lys	Arg	Leu	Gln	Ala	Glu	Asn	Ile	Pro	Val	450	455	460	
Leu	Trp	Arg	Pro	Leu	His	Glu	Ala	Glu	Gly	Gly	Trp	Phe	Trp	Trp	Gly	465	470	475	480
Ala	Lys	Gly	Pro	Glu	Ala	Ala	Ile	Glu	Leu	Tyr	Arg	Leu	Met	Tyr	Asp	485	490	495	
Arg	Tyr	Thr	Asn	His	His	Lys	Leu	Asn	Asn	Leu	Ile	Trp	Met	Trp	Asn	500	505	510	
Ser	Glu	Ala	Glu	Glu	Trp	Tyr	Pro	Gly	Asp	Asp	Val	Val	Asp	Met	Ile	515	520	525	
Ser	Thr	Asp	Ile	Tyr	Asn	Pro	Val	Gly	Asp	Phe	Ser	Pro	Ser	Ile	Asn	530	535	540	
Lys	Tyr	Glu	His	Leu	Lys	Glu	Leu	Val	Gln	Asp	Lys	Lys	Leu	Val	Ala	545	550	555	560
Leu	Pro	Glu	Thr	Gly	Ile	Ile	Pro	Asp	Pro	Asp	Gln	Leu	Gln	Leu	Phe	565	570	575	
Asn	Ala	Asn	Trp	Ser	Trp	Phe	Ala	Thr	Trp	Thr	Gly	Asp	Tyr	Ile	Arg	580	585	590	
Asp	Gly	Ile	Ser	Asn	Pro	Ile	Glu	His	Leu	Gln	Lys	Val	Phe	His	His	595	600	605	
Asp	Tyr	Val	Ile	Thr	Leu	Asp	Glu	Leu	Pro	Glu	Asn	Leu	Ser	Arg	Tyr	610	615	620	
Gly	Leu	Ser	Glu	Gly	Val	Trp	Lys	Ser	Asp	Ala	Asp	Leu	Ser	Val	Lys	625	630	635	640
Thr	Arg	Thr	Thr	Ser	Glu	Ile	Thr	Val	Asn	Trp	Ser	Asn	Ala	Ile	Gln	645	650	655	
Tyr	Asp	Ser	Val	Asn	Gly	Tyr	Lys	Leu	Ile	Lys	Asp	Gly	Val	Glu	Thr	660	665	670	
Val	Ser	Val	Glu	Gly	Gly	Val	Gln	Glu	Tyr	Thr	Phe	Thr	Asn	Leu	Leu	675	680	685	
Pro	Gly	Thr	Gln	Tyr	Thr	Ile	Lys	Val	Glu	Ala	Leu	Asp	Gln	Asp	Asp	690	695	700	

193cc

Arg	Trp	Thr	Ala	Asp	Gly	Pro	Val	Ala	Val	Val	Ser	Thr	Leu	Ser	Asn
705					710					715					720
Ala	Pro	Ile	Ser	Tyr	Pro	Pro	Ala	Val	Thr	Pro	Asp	Glu	Pro	Asn	Glu
				725					730					735	
Glu	Leu	Ser	Glu	Gly	Glu	Tyr	Thr	Leu	Leu	Ala	Asp	Asp	Leu	Ser	Ser
			740					745					750		
Gln	Asp	Gly	Val	Leu	Glu	Val	Ser	Leu	Glu	Pro	Thr	Val	Thr	Lys	Leu
		755					760					765			
Ile	Ile	Pro	Ser	Ala	Leu	Ala	Gly	Thr	Leu	Asp	Gly	Asp	Leu	Arg	Ile
	770						775				780				
Gly	Tyr	Gly	Asp	Val	Trp	Ile	Val	Ile	Pro	His	Glu	Gln	Leu	Gly	Gly
785					790					795					800
Asp	Glu	Gln	Gln	Ser	Gly	Ser	Ala	Tyr	Glu	Leu	Val	Leu	Glu	Ile	
			805						810					815	

<210> 27

<211> 1488

<212> DNA

<213> Bacillus sp.

<400> 27

atgaggaatg	aaaaaatcag	gccatttact	aaaataaagg	caagtgttgt	tactagtgtt	60
ttactattaa	ctatttcctt	aatttttact	ataggaaata	tagcaaatgc	tgaatctgag	120
gtaagaatat	ttgaagctga	agatgctatt	ttaaatgggc	tgactattaa	aaattctgaa	180
ccagggtttt	ctggtaccgg	atatgtaggt	gactttgaaa	atagctctca	gagtgtgacg	240
tttcaaattg	aggctcctaa	agccggttta	tacaacttaa	atattggata	tggcgcgatt	300
tatggaagtg	gaaaagtagc	taatgttatt	gtaaatggag	agaagctaag	tacttttaca	360
atgggaagtg	gcttttggtta	agcgtcagca	ggaaagggtat	tacttaattc	aggcttaaat	420
actatctcga	ttactcctaa	ttggacatgg	tttaccattg	attatattga	agttatacat	480
gcaccggaac	cggaaaacca	taatgtagaa	aagacgttaa	ttaacccaaa	tgcaacggat	540
gaagccaaag	ctttaataag	ctatctagtt	gataactttg	gtgagaaaat	tcttgcaggg	600
caacatgatt	atccaaatac	acgaccacga	gatttagaat	atatttatga	aactactggg	660
aagtatcctg	ctgtttttagg	tttagacttt	attgataaca	gtccttctag	agttgagcgc	720
ggagcctctg	ctgatgaaac	accagtagct	attgactggg	ggaataaagg	gggaattggt	780
actttcacct	ggcattggaa	tgctcccaaa	gatttattag	atgaaccagg	aaatgaatgg	840
tggagtgggt	tttatacgag	agcaacaact	tttgacgtag	aatatgcttt	aaaacatccg	900
aagtcggagg	actacatgct	tctaatacgt	gatattgatg	taatagctgg	tgaactaaag	960
aaattgcagg	aagcaaatgt	tcctgtttta	tggaggccac	ttcatgaggc	tgaaggcggg	1020
tggttctggt	ggggggcaaa	aggtcctgaa	tcaaccaagg	agctatggag	attaatgtat	1080
gatagaatga	cgaactacca	taacttaaat	aatttaatat	gggtatggaa	ttccattgaa	1140
gaggattggt	atcctggaga	tgagtatgtc	gatattgtaa	gcttcgattc	atatccaggt	1200
gaatataact	atagtccaat	gagccgtgag	tatgaagcac	ttaaagagtt	gtctagtaac	1260
aagaaactta	tagcaatagc	agaaaatgga	ccaataccag	atcctgattt	actacaactt	1320
taccatgcta	actatagttg	gtttgctaca	tggaatggag	atatattaag	aaatcaaaat	1380
agcgaagagc	acctaagaaa	agtatataat	catgattatg	tgattaccct	aaataaatta	1440
cctaacctta	aaacatatag	gggaagatgc	acttatacac	acactatc		1488

```

<400> 28
Met Arg Asn Glu Lys Ile Arg Pro Phe Thr Lys Ile Lys Ala Ser Val
  1          5          10          15
Val Thr Ser Val Leu Leu Leu Thr Ile Ser Leu Ile Phe Thr Ile Gly
      20          25          30
Asn Ile Ala Asn Ala Glu Ser Glu Val Arg Ile Phe Glu Ala Glu Asp
      35          40          45
Ala Ile Leu Asn Gly Leu Thr Ile Lys Asn Ser Glu Pro Gly Phe Ser
      50          55          60
Gly Thr Gly Tyr Val Gly Asp Phe Glu Asn Ser Ser Gln Ser Val Thr
  65          70          75          80
Phe Gln Ile Glu Ala Pro Lys Ala Gly Leu Tyr Asn Leu Asn Ile Gly
      85          90          95
Tyr Gly Ala Ile Tyr Gly Ser Gly Lys Val Ala Asn Val Ile Val Asn
      100          105          110
Gly Glu Lys Leu Ser Thr Phe Thr Met Gly Ser Gly Phe Gly Lys Ala
      115          120          125
Ser Ala Gly Lys Val Leu Leu Asn Ser Gly Leu Asn Thr Ile Ser Ile
      130          135          140
Thr Pro Asn Trp Thr Trp Phe Thr Ile Asp Tyr Ile Glu Val Ile His
  145          150          155          160
Ala Pro Glu Pro Glu Asn His Asn Val Glu Lys Thr Leu Ile Asn Pro
      165          170          175
Asn Ala Thr Asp Glu Ala Lys Ala Leu Ile Ser Tyr Leu Val Asp Asn
      180          185          190
Phe Gly Glu Lys Ile Leu Ala Gly Gln His Asp Tyr Pro Asn Thr Arg
      195          200          205
Pro Arg Asp Leu Glu Tyr Ile Tyr Glu Thr Thr Gly Lys Tyr Pro Ala
      210          215          220
Val Leu Gly Leu Asp Phe Ile Asp Asn Ser Pro Ser Arg Val Glu Arg
  225          230          235          240
Gly Ala Ser Ala Asp Glu Thr Pro Val Ala Ile Asp Trp Trp Asn Lys
      245          250          255

```

```
<210> 29
<211> 1086
<212> DNA
<213> Bacillus licheniformis
```

```
<400> 29
atgtacaaaa aatttggaat ctcttttattg cttgctttat taatcgtttc agctttctcg 60
cagacggcat ctgctcatac agtgaatccg gtgaaccaa atgccagtc gacaacgaag 120
gaqctqatga attggcttgc tcattctgccg aaccgatcgg aaaatcgcg actgtcaggt 180
```

193ff

```

gcattcggcg gatattctaa tgcgacgttt tctatgaaag aagccaatcg aatcaaagat 240
gctacagggc agtcacctgt cgtgtatgct tgtgattatt cgagaggatg gctggagaca 300
gctcatattg ctgatgcgat cgattatagc tgtaacagcg atctaattct tcattggaag 360
agcggaggca tacctcagat cagcatgcat cttcctaacc ctgcgtttca atccggcaat 420
tacaaaacaa agatctcaaa cagtcagtat gaaaaaatct tagactcatc aaccacagaa 480
ggcaaacgat tggatgctgt actgagcaag gttgcagatg gccttcagca gttaaaaaat 540
gaaggcgttc cagttctttt cagacctctt cacgaaatga acggagaatg gttctggtgg 600
gggcttaccg gctataacca aaaggatagc gagcgaatat cactatacaa acagctttac 660
caaaaaatct atcattatat gaccgataca agaggattgg acaacttgat ttgggtttat 720
gcaccagacg ccaaccgcga ctttaagaca gacttttata ctggggattc atatgttgat 780
attgtcggat tagacgcgta tttctcagat gcttattcga tcaaaggata tgacgagtta 840
acggcgctta ataagccatt tgcctttaca gaagtcggtc cgcaaacaac aaacggcagc 900
ctggattatt ctcaatttat caatgcagtt aaacaaaaat atccgaaaac catttatattc 960
ttagcttggg atgagggttg gagccctgcg gctaatacagg gtgcctttaa tctctataat 1020
gacagttgga cgctgaataa gggagagcta tgggaaggca gctcacttac accggcagcc 1080
gaataa                                     1086

```

<210> 30
 <211> 361
 <212> PRT
 <213> *Bacillus licheniformis*

<400> 30
 Met Tyr Lys Lys Phe Gly Ile Ser Leu Leu Leu Ala Leu Leu Ile Val
 1 5 10 15
 Ser Ala Phe Ser Gln Thr Ala Ser Ala His Thr Val Asn Pro Val Asn
 20 25 30
 Gln Asn Ala Gln Ser Thr Thr Lys Glu Leu Met Asn Trp Leu Ala His
 35 40 45
 Leu Pro Asn Arg Ser Glu Asn Arg Val Leu Ser Gly Ala Phe Gly Gly
 50 55 60
 Tyr Ser Asn Ala Thr Phe Ser Met Lys Glu Ala Asn Arg Ile Lys Asp
 65 70 75 80
 Ala Thr Gly Gln Ser Pro Val Val Tyr Ala Cys Asp Tyr Ser Arg Gly
 85 90 95
 Trp Leu Glu Thr Ala His Ile Ala Asp Ala Ile Asp Tyr Ser Cys Asn
 100 105 110
 Ser Asp Leu Ile Ser His Trp Lys Ser Gly Gly Ile Pro Gln Ile Ser
 115 120 125
 Met His Leu Pro Asn Pro Ala Phe Gln Ser Gly Asn Tyr Lys Thr Lys
 130 135 140
 Ile Ser Asn Ser Gln Tyr Glu Lys Ile Leu Asp Ser Ser Thr Thr Glu
 145 150 155 160

193gg

Gly Lys Arg Leu Asp Ala Val Leu Ser Lys Val Ala Asp Gly Leu Gln
 165 170 175
 Gln Leu Lys Asn Glu Gly Val Pro Val Leu Phe Arg Pro Leu His Glu
 180 185 190
 Met Asn Gly Glu Trp Phe Trp Trp Gly Leu Thr Gly Tyr Asn Gln Lys
 195 200 205
 Asp Ser Glu Arg Ile Ser Leu Tyr Lys Gln Leu Tyr Gln Lys Ile Tyr
 210 215 220
 His Tyr Met Thr Asp Thr Arg Gly Leu Asp Asn Leu Ile Trp Val Tyr
 225 230 235 240
 Ala Pro Asp Ala Asn Arg Asp Phe Lys Thr Asp Phe Tyr Pro Gly Asp
 245 250 255
 Ser Tyr Val Asp Ile Val Gly Leu Asp Ala Tyr Phe Ser Asp Ala Tyr
 260 265 270
 Ser Ile Lys Gly Tyr Asp Glu Leu Thr Ala Leu Asn Lys Pro Phe Ala
 275 280 285
 Phe Thr Glu Val Gly Pro Gln Thr Thr Asn Gly Ser Leu Asp Tyr Ser
 290 295 300
 Gln Phe Ile Asn Ala Val Lys Gln Lys Tyr Pro Lys Thr Ile Tyr Phe
 305 310 315 320
 Leu Ala Trp Asp Glu Gly Trp Ser Pro Ala Ala Asn Gln Gly Ala Phe
 325 330 335
 Asn Leu Tyr Asn Asp Ser Trp Thr Leu Asn Lys Gly Glu Leu Trp Glu
 340 345 350
 Gly Ser Ser Leu Thr Pro Ala Ala Glu
 355 360

<210> 31

<211> 3041

<212> DNA

<213> Caldocellulosiruptor sp.

<400> 31

caatgggctt gaagattggt attcactggg gtgctgattt tgtaatagcc aatatcaagg 60
 ttgaagaggt aactcagtaa aagaggcttt ttgctggtga gcacaccgct gaagagaaaa 120
 gtaaggttat gttaaagaag cgggtgtgcc accggcttta aaaaaataaa aaagggggaga 180
 gtgccaggat tatgagaaag ggcttaaaga ttacatcttt aatagtgagc cttgtatttt 240
 tacttgggct tttgccgaca ggaatttttg gtgctggtga gacatctgtt caaagctatg 300
 ttttcgactt tgaagatggc accacaatga cattcggtga ggcttgggga gactcattaa 360
 aatgtatcaa aaaggtgtca gtttctactg atttgcagcg acctggtaac aagtatgcgc 420
 tcaggcttga tgttgagttc aacgagaaca atggatggga ccagggcgac cttggtgcat 480
 ggataggtgg tgttgtcgaa gggcagtttg actttacaaa ctacaagtct gttgagtttg 540

193hh

```

aatgtttgt tccatacgac gagtttgcaa aagcaaaagg tggctttgct tacaaggttg 600
tattgaatga tggatggaaa gaacttggaa gcgaatttag cattacagta aatgctggca 660
aaaagggtgaa gataaacggc aaggactata tggtcattca caaggcgttt gcaattccag 720
atgatttttag aaccaaaaaag cgtgcacagc ttgtgttcca atttgcaggt caaaactgca 780
actacaaagg acctatctac cttgacaata taagagtaag acctgaggat gcgtcaaacc 840
tctcaaaaga agactatgga agtagcgaag aagaggaaat ttctgaggac tttttcacag 900
gggttaccct tgtgtatcca caggaaggca aaaactttgt gtacaatttt gaaaaagaca 960
caatgggatt ttataaatac tcgggtgatg gatttgcaaa gaaaacaaaag tcaatggaat 1020
tttcacagga cttgaagaca tcaacaaatg caggcagcct caaactcaat gctaattttc 1080
agggtactgc gtttgaagaa atgaacattg ctgtaaagct cacagacaaa gaaggaaaac 1140
tttttgacct tggcaaatac tccgcacttg agtatacaat ctacattcca aatccagaca 1200
aagttgcggg gaaaatcatg tctgcaagtg ctgtggacag tccatggaag ataatcaaag 1260
actttacact tcttaactac aaagataaga caacatggaa agagataaac ggaaagactt 1320
atgcggtcat aaagtgaag gataatcttt acaatgtaaa agaaaaagca ggtgtatttg 1380
ttttgaggat tgcgggggtct tatgtaaagt atacaggccc catctacatt gataacgtaa 1440
cattaattgc tggaaagaag gttgcaccaa aggtggagag aatatcactt ccaaatccaa 1500
agacatacta taaagttaag attgaagctg agagtgaag tgatggctgg gcttacagcg 1560
ttgagaaaga aaatgcaaag ttttctggga aaggctatgt acttttgttt gggaacaaca 1620
tgggcaatac cctttataac atcaagggtc cgaagacagg acattacatc ttcactcttg 1680
caatctcaac ccttgggctt gtaaaggatg gtagcattga tatctggata gacggtgatt 1740
tgaaagggtg ggcaaagggt ccaaacgtaa agggcaagtt ccaggaagtt gttgtcagaa 1800
aaaagattta tttaacagcg ggtgagcaca caatatcact gcaaaaatct ggcggatata 1860
caattgcagt tgactatttt gtgatagaag agcttggtgc ggcaaataaa tcaaagcttt 1920
cggtttcttc aaagttagtg accccaaatc cacaccccaa tgcccaaagg ctcataaatt 1980
atgtgtcaag catttacggt gaaaagattt tgtctggtca gcagagcagc ggtgaaggca 2040
aagaggttca gatgattttt gatgtcacia agagatatcc agctgttaga agctttgatt 2100
tcatggacta ctcaccaagc agagtgcagc atggtacaaa aggtacagat gttgatgagg 2160
caataaagtg gtggaagagc ggcggcatag ttgcattttg ctggcactgg aacgcaccaa 2220
caggtcttat tgaccagccg ggcaaagagt ggtggagagg tttttacaca gaggctacaa 2280
catttgacct caagaaagcc atggacaatc caaattctga agaataataa ctcattttga 2340
gagatataga cgctattgct gagcagctca aaaaattgca ggctgaagggt gtgccagttc 2400
ttttcagacc gcttcacgag gcctctggcg gctgggtctg gtgggggtgca aaagggtccag 2460
agccgtatat aaagcttttg aagctcatgt ttgacaggct tgtaaactat cacaaaatca 2520
acaaccta atgggttatgg aacggtcagg atgctgcctg gtatccgggt gaccagtatg 2580
ttgatataat tgcagaagat atatatgagg aaaaagctca gtactacca tatacagaga 2640
ggttcgtgaa agctctcaag tacacaaatg caaacaagat gatagcactt tctgagtgcg 2700
gaactattcc tgaccgggt gtgctaaaac aagaagggtg ttcgtggctg tggttttctg 2760
tatgggcagg aagctatgtc atgacaggca gcaagtacaa cgatgaatgg aacgacaatc 2820
acatgctaag aaagatttac aacaatgact atgtaataac aaaagatgaa ctacctgata 2880
taaagagcat tccactcaaa tagaatgaga tatattttgg aatatccaaa atcaactgtc 2940
agcctgtgag aggagagaag ttcaaaaaag acctctccc tttttgggtc ttgcaaaata 3000
atcaattttt ggttttgaca tctcaaacat gttaattaaa a 3041

```

<210> 32

<211> 903

<212> PRT

<213> Caldocellulosiruptor sp.

<400> 32

Met Arg Lys Gly Leu Lys Ile Thr Ser Leu Ile Val Ser Leu Val Phe
1 5 10 15

Leu Leu Gly Leu Leu Pro Thr Gly Ile Phe Gly Ala Val Glu Thr Ser
20 25 30

Val	Gln	Ser	Tyr	Val	Phe	Asp	Phe	Glu	Asp	Gly	Thr	Thr	Met	Thr	Phe	
		35					40					45				
Gly	Glu	Ala	Trp	Gly	Asp	Ser	Leu	Lys	Cys	Ile	Lys	Lys	Val	Ser	Val	
	50					55					60					
Ser	Thr	Asp	Leu	Gln	Arg	Pro	Gly	Asn	Lys	Tyr	Ala	Leu	Arg	Leu	Asp	
65					70					75					80	
Val	Glu	Phe	Asn	Glu	Asn	Asn	Gly	Trp	Asp	Gln	Gly	Asp	Leu	Gly	Ala	
				85					90					95		
Trp	Ile	Gly	Gly	Val	Val	Glu	Gly	Gln	Phe	Asp	Phe	Thr	Asn	Tyr	Lys	
			100					105					110			
Ser	Val	Glu	Phe	Glu	Met	Phe	Val	Pro	Tyr	Asp	Glu	Phe	Ala	Lys	Ala	
		115					120					125				
Lys	Gly	Gly	Phe	Ala	Tyr	Lys	Val	Val	Leu	Asn	Asp	Gly	Trp	Lys	Glu	
	130					135					140					
Leu	Gly	Ser	Glu	Phe	Ser	Ile	Thr	Val	Asn	Ala	Gly	Lys	Lys	Val	Lys	
145					150					155					160	
Ile	Asn	Gly	Lys	Asp	Tyr	Met	Val	Ile	His	Lys	Ala	Phe	Ala	Ile	Pro	
				165					170					175		
Asp	Asp	Phe	Arg	Thr	Lys	Lys	Arg	Ala	Gln	Leu	Val	Phe	Gln	Phe	Ala	
			180					185					190			
Gly	Gln	Asn	Cys	Asn	Tyr	Lys	Gly	Pro	Ile	Tyr	Leu	Asp	Asn	Ile	Arg	
		195					200					205				
Val	Arg	Pro	Glu	Asp	Ala	Ser	Asn	Leu	Ser	Lys	Glu	Asp	Tyr	Gly	Ser	
	210					215					220					
Ser	Glu	Glu	Glu	Glu	Ile	Ser	Glu	Asp	Phe	Phe	Thr	Gly	Val	Thr	Leu	
225					230					235					240	
Val	Tyr	Pro	Gln	Glu	Gly	Lys	Asn	Phe	Val	Tyr	Asn	Phe	Glu	Lys	Asp	
				245					250					255		
Thr	Met	Gly	Phe	Tyr	Lys	Tyr	Ser	Gly	Asp	Gly	Phe	Ala	Lys	Lys	Thr	
			260					265					270			
Lys	Ser	Met	Glu	Phe	Ser	Gln	Asp	Leu	Lys	Thr	Ser	Thr	Asn	Ala	Gly	
		275					280					285				
Ser	Leu	Lys	Leu	Asn	Ala	Asn	Phe	Gln	Gly	Thr	Ala	Phe	Glu	Glu	Met	
	290					295					300					
Asn	Ile	Ala	Val	Lys	Leu	Thr	Asp	Lys	Glu	Gly	Lys	Leu	Phe	Asp	Leu	
305					310					315					320	
Gly	Lys	Tyr	Ser	Ala	Leu	Glu	Tyr	Thr	Ile	Tyr	Ile	Pro	Asn	Pro	Asp	
				325					330					335		

193jj

Lys Val Ala Gly Lys Ile Met Ser Ala Ser Ala Val Asp Ser Pro Trp
 340 345 350

Lys Ile Ile Lys Asp Phe Thr Leu Leu Asn Tyr Lys Asp Lys Thr Thr
 355 360 365

Trp Lys Glu Ile Asn Gly Lys Thr Tyr Ala Val Ile Lys Cys Lys Asp
 370 375 380

Asn Leu Tyr Asn Val Lys Glu Lys Ala Gly Val Leu Val Leu Arg Ile
 385 390 395 400

Ala Gly Ser Tyr Val Lys Tyr Thr Gly Pro Ile Tyr Ile Asp Asn Val
 405 410 415

Thr Leu Ile Ala Gly Lys Lys Val Ala Pro Lys Val Glu Arg Ile Ser
 420 425 430

Leu Pro Asn Pro Lys Thr Tyr Tyr Lys Val Lys Ile Glu Ala Glu Ser
 435 440 445

Ala Ser Asp Gly Trp Ala Tyr Ser Val Glu Lys Glu Asn Ala Lys Phe
 450 455 460

Ser Gly Lys Gly Tyr Val Leu Leu Phe Gly Asn Asn Met Gly Asn Thr
 465 470 475 480

Leu Tyr Asn Ile Lys Val Pro Lys Thr Gly His Tyr Ile Phe Thr Leu
 485 490 495

Ala Ile Ser Thr Leu Gly Leu Val Lys Asp Gly Ser Ile Asp Ile Trp
 500 505 510

Ile Asp Gly Asp Leu Lys Gly Gly Ala Lys Val Pro Asn Val Lys Gly
 515 520 525

Lys Phe Gln Glu Val Val Val Arg Lys Lys Ile Tyr Leu Thr Ala Gly
 530 535 540

Glu His Thr Ile Ser Leu Gln Lys Ser Gly Gly Tyr Thr Ile Ala Val
 545 550 555 560

Asp Tyr Phe Val Ile Glu Glu Leu Val Ala Ala Asn Lys Ser Lys Leu
 565 570 575

Ser Val Ser Ser Lys Leu Val Thr Pro Asn Pro His Pro Asn Ala Gln
 580 585 590

Arg Leu Ile Asn Tyr Leu Ser Ser Ile Tyr Gly Glu Lys Ile Leu Ser
 595 600 605

Gly Gln Gln Ser Ser Gly Glu Gly Lys Glu Val Gln Met Ile Phe Asp
 610 615 620

Val Thr Lys Arg Tyr Pro Ala Val Arg Ser Phe Asp Phe Met Asp Tyr
 625 630 635 640

193kk

Ser Pro Ser Arg Val Gln His Gly Thr Lys Gly Thr Asp Val Asp Glu
 645 650 655
 Ala Ile Lys Trp Trp Lys Ser Gly Gly Ile Val Ala Phe Cys Trp His
 660 665 670
 Trp Asn Ala Pro Thr Gly Leu Ile Asp Gln Pro Gly Lys Glu Trp Trp
 675 680 685
 Arg Gly Phe Tyr Thr Glu Ala Thr Thr Phe Asp Leu Lys Lys Ala Met
 690 695 700
 Asp Asn Pro Asn Ser Glu Glu Tyr Lys Leu Ile Leu Arg Asp Ile Asp
 705 710 715 720
 Ala Ile Ala Glu Gln Leu Lys Lys Leu Gln Ala Glu Gly Val Pro Val
 725 730 735
 Leu Phe Arg Pro Leu His Glu Ala Ser Gly Gly Trp Phe Trp Trp Gly
 740 745 750
 Ala Lys Gly Pro Glu Pro Tyr Ile Lys Leu Trp Lys Leu Met Phe Asp
 755 760 765
 Arg Leu Val Asn Tyr His Lys Ile Asn Asn Leu Ile Trp Val Trp Asn
 770 775 780
 Gly Gln Asp Ala Ala Trp Tyr Pro Gly Asp Gln Tyr Val Asp Ile Ile
 785 790 795 800
 Ala Glu Asp Ile Tyr Glu Glu Lys Ala Gln Tyr Ser Pro Tyr Thr Glu
 805 810 815
 Arg Phe Val Lys Ala Leu Lys Tyr Thr Asn Ala Asn Lys Met Ile Ala
 820 825 830
 Leu Ser Glu Cys Gly Thr Ile Pro Asp Pro Ala Val Leu Lys Gln Glu
 835 840 845
 Gly Val Ser Trp Leu Trp Phe Ser Val Trp Ala Gly Ser Tyr Val Met
 850 855 860
 Thr Gly Ser Lys Tyr Asn Asp Glu Trp Asn Asp Asn His Met Leu Arg
 865 870 875 880
 Lys Ile Tyr Asn Asn Asp Tyr Val Ile Thr Lys Asp Glu Leu Pro Asp
 885 890 895
 Ile Lys Ser Ile Pro Leu Lys
 900

<210> 33

<211> 1450

<212> RNA

<213> Bacillus sp. I633

19311

<400> 33

```

gcucccugau guuagcggcg gacgggugag uaacacgugg gcaaccugcc cuguagacug 60
ggauaacauc gagaaaucgg ugcuaauacc ggauaauaga uggaauugca uauuucuaau 120
uuaaaagaug gcuccggcua ucacuacagg augggcccgcc ggcgcauuag cuaguuggua 180
agguaacggc uuaccaaggc gacgaugcgu agccgaccug agagggugau cggccacacu 240
gggacugaga cacggcccag acuccuacgg gaggcagcag uagggaauuc uccgcaaugg 300
acgaaagucu gacggagcaa cgccgcguga gcgaugaagg ccuucggguu guaaagcucu 360
guuguuaggg aagaacaagu gccauucaa uaggguggca ccuugacggg accuaaccag 420
aaagccacgg cuaacuacgu gccagcagcc gcgguauuac guagguggca agcguugucc 480
ggaauuauug ggcguaaagc gcgcgcaggg gguuucuaaa gucugaugug aaagcccccg 540
gcucaaccgg ggagggucau uggaaacugg gagacuugag uacagaagag gagaguggaa 600
uuccacgugu agcggugaaa ugcguagaua uguggaggaa caccaguggc gaaggcgacu 660
cucuggucug uaacugacgc ugaggcgcg aagcgugggg agcaaacagg auuagauacc 720
cugguagucc acgccguaaa cgaugagugc uagguguuag gguuucgau gcccuuagug 780
ccgaaguuaa cacaguaagc acuccgccug gggaguacgg ccgcaaggcu gaaacucaaa 840
ggaauugacg ggggcccgcg caagcggugg agcauguggu uuaauucgaa gcaacgcgaa 900
gaaccuuacc aggucuugac auccuuugac aaccuagag auaggggcuu ccccuucggg 960
ggacaaagug acagguggug caugguuguc gucagcucgu gucgugagau guuggguuaa 1020
gucccgcaac gagcgcaacc cuugaucuaa guugccagca uuuaguuggg cacucuaagg 1080
ugacugcccg ugacaaaccg gaggaaggug gggauagcgu caaaucauca ugccccuau 1140
gaccugggcu acacacgugc uacaauaggau gguacaaagg gcagcaaac cgcgaggucg 1200
agccaauccc auaaaaccu ucucaguucg gauuguaggc ugcaacucgc cuacaugaag 1260
ccggaaucgc uaguaaucgc ggaucagcau gccgcgguga auacguuccc gggccuugua 1320
cacaccgcc gucacaccac gagaguugu aacacccgaa gucggugggg uaaccuuuug 1380
gagccagccg ccuaaggugg gacagaugau uggggugaag ucguaacaag guagccguau 1440
cggaaggugc                                     1450

```

<210> 34

<211> 1508

<212> RNA

<213> Bacillus sp. AAI12

<400> 34

```

gacgaacgcu ggcggcgugc cuaauacaug caagucgagc ggacauuuag gagcuugcuc 60
cuaaauguua gcgggcgagc ggugaguaac acgugggcaa ccugcccugu agacugggau 120
aauaucgaga aaucggugcu aaauaccgau aaucuuagag auugcauaau ccucuuguaa 180
aagauggcuc cggcuauac uacgggaugg gcccgcggcg cauuagcuag uugguaaggu 240
aacggcuuac caaggcgagc augcguagcc gaccugagag ggugaucggc cacacuggga 300
cugagacacg gccagacuc cuacgggagg cagcaguagg gaaucuuucc caauggacga 360
aagucugacg gagcaacgcc gcgugaguga ugaaggguuu cggcucguua agcucuguug 420
uuagggaaga acaagugccg uucaauuagg gcggcaccuu gacgguaacc aaccagaaag 480
ccacggcuua cuacgugcca gcagccgcgg uaauacguag guggcaagcg uuguccggaa 540
uuauuggggc uaaagcgcg gcaggcgguu uuuuaagucu gaugugaaau cucggggcuc 600
aaccgccgag ggucuuugga aacugggaga cuugaguaca gaagaggaga guggaaaucc 660
acguguagcg gugaaaugcg uagauaugug gaggaacacc aguggcgaag gcgacucucu 720
ggucuguaac ugacgcugag gcgcgaaagc guggggagca aacaggauua gauaccucgg 780
uaguccacgc cguaaacgau gagugcuagg uguuaggggu uucgaugccc uuagugccga 840
aguuaacaca uuaagcacuc cgccugggga guacgaccgc aagguugaaa cucaaaggaa 900
uugacggggg cccgcacaag caguggagca ugugguuuua uucgaagcaa cgcgaagaac 960
cuuaccaggu cuugacaucc uuaugaccuc ccuagagaua gggauuuccc uucggggaca 1020
uaagugacag guggugcaug guugucguca gcucgugucg ugagauguug gguuaagucc 1080
cgcaacgagc gcaaccuuug aucuuaguug ccagcauuua guugggcacu cuaaggugac 1140
ugccggugau aaaccggagg aaggugggga ugacgucaaa ucaucaugcc ccuuaugacc 1200
ugggcuaacac acgugcuaca auggauggua caaagagcag caaaaccgcg agguccagcc 1260

```

193mm

aaucucauaa	agccauucuc	aguucggauu	guaggcugca	acucgccuac	augaagccgg	1320
aaugcuagu	aaucgcggau	cagcaugccg	cggugaauac	guucccgggc	cuuguacaca	1380
ccgcccguca	caccacgaga	guuuguaaca	cccgaagucg	guggaguaac	ccuuacggga	1440
gcuagccgcc	uaagguggga	cagaugauug	gggugaaguc	guaacaaggu	agccguaucg	1500
gaaggugc						1508

CLAIMS:

1. An isolated mannanase which is
 - (a) a polypeptide comprising an amino acid sequence as shown in positions 31-330 of SEQ ID NO: 2,
 - (b) a polypeptide encoded by the DNA sequence as shown in positions 91-990 or positions 91-1470 of SEQ ID NO: 1, or
 - (c) a polypeptide which is at least 80 % identical with a polypeptide defined in (a) or (b) and wherein said polypeptide has mannanase activity.
2. The mannanase according to claim 1, said mannanase being from a strain of *Bacillus* sp.
3. The mannanase according to claim 2 which has
 - (i) a relative mannanase activity of at least 60 % in the pH range 7.5-10, measured at 40°C;
 - (ii) a molecular weight of 34 ± 10 kDa, as determined by SDS-PAGE; or
 - (iii) the N-terminal sequence ANSGFYVSGTTLYDANG.
4. The mannanase according to claim 2 which has
 - (i) a relative mannanase activity of at least 60 % in the pH range 7.5-10, measured at 40°C;
 - (ii) a molecular weight of 34 ± 10 kDa, as determined by SDS-PAGE; and
 - (iii) the N-terminal sequence ANSGFYVSGTTLYDANG.
5. An isolated polynucleotide molecule comprising a DNA sequence encoding an enzyme exhibiting mannanase activity, which DNA sequence comprises:

- (a) the DNA sequence shown in positions 91-1470 in SEQ ID NO: 1;
- (b) a DNA sequence which is at least 80 % identical with the DNA sequence defined in (a);
- (c) a DNA sequence which hybridizes with a double-stranded DNA probe comprising the sequence shown in positions 91-990 in SEQ ID NO: 1 under the following conditions: a filter containing DNA fragments or RNA to hybridize is presoaked in 5 x SSC for 10 min, and prehybridization of the filter in a solution of 5 x SSC, 5 x Denhardt's solution, 0.5 % SDS and 100 µg/ml of denatured sonicated salmon sperm DNA is performed, followed by hybridization in the same solution containing a concentration of 10ng/ml of a random-primed, 32P-dCTP-labeled probe for 12 hours at 45°C, the filter is then washed twice for 30 minutes in 2 x SSC, 0.5 % SDS at 70°C; or
- (d) a DNA sequence which, because of the degeneracy of the genetic code, does not hybridize with the sequences of (a), (b) or (c) under the conditions set out in (c), but which codes for a poly-peptide having exactly the same amino acid sequence as the polypeptide encoded by the DNA sequence of (a), (b) or (c).

6. The polynucleotide molecule of claim 5, wherein the DNA sequence encoding the enzyme exhibiting mannanase activity comprises the DNA sequence shown in positions 91-990 in SEQ ID NO: 1.

7. The cloned DNA sequence according to claim 5 or 6, in which the DNA sequence encoding an enzyme exhibiting mannanase activity is obtained from *Bacillus* sp I633, *Bacillus* sp AAI12, *Bacillus* sp AA349 or *Humicola insolens*.

8. An isolated polynucleotide molecule encoding a polypeptide having mannanase activity which polynucleotide molecule hybridizes to a denatured double-stranded DNA probe under the following conditions: a filter containing DNA fragments or RNA to hybridize is presoaked in 5 x SSC for 10 min, and prehybridization of the filter in a solution of 5 x SSC, 5 x Denhardt's solution, 0.5 % SDS and 100 µg/ml of denatured sonicated salmon sperm DNA is performed, followed by hybridization in the same solution containing a concentration of 10ng/ml of a random-primed, 32P-dCTP-labeled probe for 12 hours at 45°C, the filter is then washed twice for 30 minutes in 2 x SSC, 0.5 % SDS at 70°C, wherein the probe is selected from the group consisting of DNA probes comprising the sequence shown in positions 91-990 of SEQ ID NO: 1, the sequence shown in positions 91-1470 of SEQ ID NO: 1 and DNA probes comprising a subsequence of positions 91-990 of SEQ ID NO: 1 having a length of at least about 100 base pairs.

9. An expression vector comprising the following operably linked elements: a transcription promoter; a DNA segment selected from the group consisting of (a) polynucleotide molecules encoding a polypeptide having mannanase activity comprising a nucleotide sequence as shown in SEQ ID NO: 1 from nucleotide 91 to nucleotide

990, (b) polynucleotide molecules encoding a polypeptide having mannanase activity that is at least 80 % identical to the amino acid sequence of SEQ ID NO: 2 from amino acid residue 31 to amino acid residue 330, and (c) degenerate nucleotide sequences of (a) or (b); and a transcription terminator.

10. A cultured cell into which has been introduced an expression vector according to claim 9, wherein said cell expresses the polypeptide encoded by the DNA segment.

11. An isolated polypeptide having mannanase activity selected from the group consisting of:

- (a) polypeptide molecules comprising an amino acid sequence as shown in SEQ ID NO: 2 from residue 31 to residue 330; and
- (b) polypeptide molecules that are at least 80 % identical to the amino acids of SEQ ID NO: 2 from amino acid residue 31 to amino acid residue 330, and wherein said polypeptide has mannanase activity.

12. The polypeptide according to claim 11 which is produced by *Bacillus* sp I633.

13. An enzyme preparation comprising a purified polypeptide according to claim 11 or 12.

14. A method of producing a polypeptide having mannanase activity comprising culturing a cell into which has been introduced an expression vector according to claim 9, whereby said cell expresses a polypeptide

encoded by the DNA segment; and recovering the polypeptide.

15. The preparation according to claim 13 which further comprises at least one enzyme selected from the group consisting of proteases, cellulases, β -glucanases, hemicellulases, lipases, peroxidases, laccases, α -amylases, glucoamylases, cutinases, pectinases, reductases, oxidases, phenoloxidases, ligninases, pullulanases, pectate lyases, xyloglucanases, xylanases, pectin acetyl esterases, polygalacturonases, rhamnogalacturonases, pectin lyases, other mannanases, pectin methylesterases, cellobiohydrolases, and transglutaminases.

16. The preparation according to claim 15, where the cellulase is an endoglucanase.

17. An isolated enzyme having mannanase activity, in which the enzyme is (i) free from homologous impurities, and (ii) produced by the method according to claim 14.

18. A method for improving the properties of cellulosic or synthetic fibres, yarn, woven or non-woven fabric in which method the fibres, yarn or fabric is treated with the preparation according to claim 13, 15 or 16, an effective amount of the mannanase according to any one of claims 1-3 or an effective amount of the enzyme according to claim 17.

19. The method according to claim 18, wherein the enzyme preparation or the enzyme is used in a desizing process step.

20. A method for degradation or modification of plant material in which method the plant material is treated with the preparation according to claim 13, 15 or 16, the mannanase according to any one of claims 1-3 or the enzyme according to claim 17.

21. The method according to claim 20 wherein the plant material is recycled waste paper; mechanical, chemical, semichemical, kraft or other paper-making pulps; fibres subjected to a retting process; or guar gum or locust bean gum containing material.

22. A method for processing liquid coffee extract, in which method the coffee extract is treated with the preparation according to claim 13, 15 or 16 or an effective amount of the mannanase according to any one of claims 1-3 or an effective amount of the enzyme according to claim 17.

23. A cleaning composition comprising the enzyme preparation according to claim 13, 15 or 16, the mannanase according to any one of claims 1-3 or the enzyme according to claim 17, and a detergent component.

24. The cleaning composition according to claim 23 which further comprises an enzyme selected from the group consisting of cellulases, proteases, lipases, amylases, pectin degrading enzymes and xyloglucanases; and a conventional detergent ingredient.

25. The cleaning composition according to claim 23 wherein said enzyme or enzyme preparation is present at a level of from 0.0001 % to 2 % pure enzyme by weight of total composition.

26. The cleaning composition according to claim 23 wherein said enzyme or enzyme preparation is present at a level of from 0.0005 % to 0.5 % pure enzyme by weight of total composition.

27. The cleaning composition according to claim 23 wherein said enzyme or enzyme preparation is present at a level of from 0.001 % to 0.1 % pure enzyme by weight of total composition.

28. The cleaning composition according to claim 24 wherein the enzyme is present at a level of from 0.0001 % to 2 % pure enzyme by weight of total composition.

29. The cleaning composition according to claim 24 wherein said enzyme or enzyme preparation is present at a level of from 0.0005 % to 0.5 % pure enzyme by weight of total composition.

30. The cleaning composition according to claim 24 wherein said enzyme or enzyme preparation is present at a level of from 0.0001 % to 0.1 % pure enzyme by weight of total composition.

31. The cleaning composition according to claim 24 wherein the enzyme is an amylase.

32. The cleaning composition according to claim 31 which further comprises yet another enzyme selected from

the group consisting of cellulases, proteases, lipases, pectin degrading enzymes and xyloglucanases.

33. The cleaning composition according to claim 23 which comprises at least one surfactant selected from anionic, nonionic and cationic surfactants.

34. The cleaning composition according to claim 23 which comprises a bleaching agent.

35. The cleaning composition according to claim 23 which comprises a builder.

36. A cleaning composition according to claim 23 which comprises a cationic surfactant comprising two long carbon chains.

37. A process for machine treatment of fabrics which process comprises treating fabric during a washing cycle of a machine washing process with a washing solution containing the enzyme preparation according to claim 13, 15 or 16, the mannanase according to any one of claims 1-3 or the enzyme according to claim 17.

38. Use of the enzyme preparation according to claim 13, 15 or 16, the mannanase according to any one of claims 1-3 or the enzyme according to claim 17 together with an enzyme selected from the group consisting of cellulases, proteases, lipases, amylases, pectin degrading enzymes and xyloglucanases in a cleaning composition for fabric cleaning or fabric stain removal.

39. Use of the enzyme preparation according to claim 13, 15 or 16, the mannanase according to any one of


claims 1-3 or the enzyme according to claim 17 together with an enzyme selected from the group consisting of cellulases, amylases, proteases, lipases, pectin degrading enzymes and xyloglucanases in a cleaning composition for cleaning hard surfaces.

40. The use of claim 39, wherein the hard surfaces are selected from the group consisting of floors, walls, and bathroom tiles.

41. Use of the enzyme preparation according to claim 13, 15 or 16, the mannanase according to any one of claims 1-3 or the enzyme according to claim 17 together with an enzyme selected from the group consisting of cellulases, amylases, proteases, lipases, pectin degrading enzymes and xyloglucanases in a cleaning composition for hand and machine dishwashing.

42. Use of the enzyme preparation according to claim 13, 15 or 16, the mannanase according to any one of claims 1-3 or the enzyme according to claim 17 together with an enzyme selected from the group consisting of cellulases, amylases, proteases, lipases, pectin degrading enzymes and xyloglucanases in a cleaning composition for oral, dental, contact lenses and personal cleaning applications.

1/3



0.10

Fig. 1




Fig. 2

3/3




Fig. 3

BaiSpec5, *Bacillus clausii*

NN017546

BaiAlcal, *Bacillus alcalophilus*

BaiSpec3, *Bacillus pseudoalcalophilus*

BaiSpor2, *Bacillus sporothermodurans*

0.10