



(51) International Patent Classification:

C07H 19/24 (2006.01) C12N 15/10 (2006.01)

C07H 1/00 (2006.01) C12Q 1/68 (2018.01)

C07H 21/02 (2006.01) G01N 33/58 (2006.01)

(21) International Application Number:

PCT/EP2021/067342

(22) International Filing Date:

24 June 2021 (24.06.2021)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2009705.1 25 June 2020 (25.06.2020) GB

(71) Applicant: **STEALTH LABELS BIOTECH AB** [SE/SE]; c/o Chalmers Ventures AB, Vera Sandbergs Allé 8, 412 96 Gothenburg, Västra Götaland County (SE).

(72) Inventors: **WILHELMSSON, Marcus**; c/o Chalmers University of Technology, SE-412 96 Gothenburg (SE). **ESBJÖRNER, Elin**; c/o Chalmers University of Technology, SE-412 96 Gothenburg (SE). **BALADI, Tom**; c/o Chalmers University of Technology, SE-412 96 Gothenburg (SE).



(74) Agent: **JONES, Teyrnon**; Alembia Intellectual Property Ltd, Biohub at Alderley Park, Alderley Park, Macclesfield Cheshire SK10 4TG (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: FLUORESCENT CYTOSINE ANALOGUES AND THEIR APPLICATION IN TRANSCRIPTION AND TRANSLATION

Figure 1



(57) Abstract: This specification discloses a novel methodology for labelling RNA via enzymatic incorporation of a minimally perturbing fluorescent tricyclic cytosine analogue. This analogue is shown to be 100% incorporated in example transcripts and is fully compatible with both in vitro and in cell transcription. Spectroscopic characterization shows that the incorporation rate of the cytosine analogue is on par with its natural counterpart. Using live cell imaging and flow cytometry, labelled mRNAs are efficiently and correctly translated upon transfection into living cells and cell-free systems. The spectral properties of the modified transcripts and their correct translation product allow for their straightforward and simultaneous visualization. This technology therefore offers a general route to understanding the biological behaviour of RNA of interest, including RNA based drugs. The fluorescent tricyclic cytosine analogue has formula (I):



Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))
- in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE

Fluorescent Cytosine Analogues and Their Application in Transcription and Translation

FIELD

5 [001] This specification relates to a modified fluorescent nucleobase triphosphate that can be used to biosynthetically produce labelled RNA, which is in turn amenable to in cell translation. This enables live cell imaging in which the labelled messenger RNA and its corresponding translation product (when a fluorescent fusion protein) can be visualised simultaneously.

BACKGROUND

10 [002] RNA plays a fundamental role in human biology. It is the main player of the central dogma of biochemistry and a crucial regulator of gene expression *via* for instance micro and small interfering RNA, as well as through its intrinsic catalytic activity. It has, for these reasons, also emerged as a highly promising and versatile new drug modality: since RNA therapeutics have the potential to modify cellular function at the translational level, they may open up new opportunities to address previously undruggable targets.

[003] An increased molecular and mechanistic knowledge of the biological processes involving RNA is therefore vital to understanding diseases and treat them. For example, there is a growing body of evidence suggesting that the key to unleashing the full potential of RNA-based drugs lies in understanding the processes of cell uptake and endosomal release (Dowdy, S. F., *Nat. Biotechnol.* 35, 222-229, [2017]). Regardless of the endocytosis mechanism, the delivery of a nucleic acid cargo to the cytoplasm always relies on endosomal escape, the understanding of which, despite extensive investigations, remains elusive (Crooke, S. T. *et al.*, *Nat. Biotechnol.* 35, 230 [2017]; Pei, D. & Buyanova, M., *Bioconjugate Chem.* [2018]). In this context, tracking of endogenous and exogenous (therapeutic) RNAs inside cells, including their translocation, localization, splicing and degradation, is of great importance.

20 [004] Recent advances have resulted in the development of a broad spectrum of tools and probes by which RNA can be analysed and quantified, but they generally involve heavily modified oligonucleotides with properties significantly different from natural ones, potentially resulting in loss of ability to be recognized and processed by the enzymatic machinery of cells. For example, a drawback of existing fluorescence-based technologies for studying cellular localization of RNA is that they primarily rely on highly amphiphilic and/or bulky external fluorescent constructs which could

impair motility and perturb localization of the RNA and its molecular interactions with (for example) membrane constituents. In addition, a majority of these technologies are incompatible with live cell imaging (Li, Y., Ke, K. & Spitale, R. C., *Biochemistry* 58, 379-386 [2019]).

[005] To overcome these issues and provide an improved method of investigating RNA mechanisms, this specification discloses a modified nucleobase triphosphate (compound (I), "tC⁰TP") which can be used to incorporate minimally perturbing and internal labels ("tC⁰") into functional messenger RNA ("mRNA"), giving it utility in live cell imaging and for drug delivery studies. Once incorporated into RNA, the structure of tC⁰ enables it to retain base-pairing and stacking, so that it minimally perturbs natural biological processes (Fig. 1, where the top schematic structure is tC⁰ labelled RNA compared to a conventional externally labelled RNA below). Therefore, this fluorophore constitutes a native-like alternative label, opening new possibilities not only to track the nucleic acid of interest but also to use fluorescent read-outs to obtain detailed information regarding nucleic acid structure and behaviour.

[006] To exemplify the possible applications of tC⁰, the specification describes successful in vitro transcription and also effective in cell translation of a full-length mRNA internally labelled with this fluorescent nucleobase analogue. In addition, by using a transcript encoding for the histone protein H2B fused to a Green Fluorescent Protein ("H2B:GFP") which localizes to the nucleus, the specification shows that it is possible to visualize the labelled mRNA transcripts inside cells while concomitantly recording the fluorescence emanating from the expressed H2B:GFP protein. This approach should be generally applicable to any fluorescent protein.

[007] Wilhelmsson, M. *et al. Sci. Rep.* 7, 2393 [2017] reports the preparation of RNA molecules labelled with tC⁰, but only discloses very short fluorescent RNA oligomers prepared by non-enzymatic solid-phase oligonucleotide synthesis, as opposed to full length RNAs accessible by biosynthesis. By virtue of their method of preparation the labelled RNA molecules are not amenable to in cell "live" analysis of transcription, translation or delivery of long therapeutic RNAs (which are mRNA-based) and therefore do not allow the same level of mechanistic insight.

[008] WO2011/034895 concerns methods for labelling DNA and RNA. It mentions a structurally different fluorescent ribonucleotide analogue 1,3-diaza-2-oxophenothiazine-ribose-5'-triphosphate ("tCTP") which is used during in vitro transcription reactions to prepare labelled RNA. However, unlike the present specification only non-coding labelled RNA was prepared and there is no disclosure of any in cell biosynthesis or visualisation.

[009] The differently labelled, full-length coding RNA polymers accessible using the technology disclosed in the present specification also have advantageous properties over the labelled RNAs in WO2011/034895, for example 1) improved fluorescence levels and label photostability; 2) improved

in vitro transcription fidelity; and 3) native-like levels of in cell translation of tC⁰-labelled mRNA resulting in the correct protein product and localization.

[010] In summary therefore, this specification discloses a labelling technique that not only allows localisation and tracking of the tagged RNA, but also facilitates analysis of the biological functionality and delivery efficacy of mRNA, an important future drug modality. Since the internal tC⁰ label is compatible with biological processes that RNA participates in it holds a great potential to be used as a powerful imaging tool in live cell microscopy, for example in detailed investigations of cellular uptake, endosomal release, exosomal loading and trafficking. These have significant potential to elucidate how these vital delivery pathways work and can be controlled.

SUMMARY

[011] A primary objective of the present specification is to provide a modified nucleobase triphosphate that can be used to make labelled RNA especially suitable for in vitro and in vivo mechanistic investigations.

[012] Accordingly, this specification describes, in part, a compound of formula (I) or a salt thereof as claimed in claim 1.

[013] This specification also describes, in part, a process for preparing a compound of formula (I) or a salt thereof as claimed in claim 5.

[014] This specification also describes, in part, a composition for preparing a tC⁰ labelled RNA molecule comprising a compound of formula (I) as claimed in claim 16.

[015] This specification also describes, in part, the use of a compound of formula (I) or a salt thereof to enzymatically prepare a tC⁰ labelled RNA molecule as claimed in claim 17.

[016] This specification also describes, in part, a process for preparing a tC⁰ labelled RNA molecule as claimed in claim 19.

[017] This specification also describes, in part, the use of a tC⁰ labelled mRNA molecule to prepare a protein encoded by the mRNA by translation as claimed in claim 20.

ILLUSTRATIVE EMBODIMENTS

[018] The invention detailed in this specification should not be interpreted as being limited to any of the recited embodiments or examples. Other embodiments will be readily apparent to a reader skilled in the art.

General Definitions

[019] “A” or “an” mean “at least one”. In any embodiment where “a” or “an” are used to denote a given material or element, “a” or “an” may mean one. In any embodiment where “a” or “an” are used to denote a given material or element, “a” or “an” may mean 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 100, 1000, 10000, 100000 or 1000000 (1 million).

[020] When an embodiment includes “a” or “an” feature X, subsequent referrals to “the” feature X do not imply only one of the feature is present. Instead the above interpretation of “a” or “an” continues to apply so that “the” also means “at least one”. In other words, embodiments comprising “a feature X, where the feature X is...” should be construed as “at least one feature X, where the at least one feature X is...”.

[021] “Comprising” means that a given embodiment may contain other features. For example, in any embodiment where a material “comprising” certain materials or elements is mentioned, the given material may be formed of at least 10% w/w, at least 20% w/w, at least 30% w/w, or at least 40% w/w of the materials or elements (or combination of materials or elements).

[022] In any embodiment where “comprising” is mentioned, “comprising” may also mean “consisting of” (or “consists of”) or “consisting essentially of” (or “consists essentially of”).

[023] With respect to embodiments of a material, “consisting of” or “consists of” means the material or element is formed entirely of the material or element (or combination of materials or elements). In any embodiment where “consisting of” or “consists of” is mentioned the given material or element may be formed of 100% w/w of the material or element.

[024] With respect to embodiments of a material, “consisting essentially of” or “consists essentially of” means that a given material or element consists almost entirely of that material or element (or combination of materials or elements). In any embodiment where “consisting essentially of” or “consists essentially of” is mentioned the given material or element may be formed of at least 50% w/w, at least 60% w/w, at least 70% w/w, at least 80% w/w, at least 90% w/w, at least 95% w/w or at least 99% w/w of the material or element.

[025] In any embodiment where “is” or “may be” is used to define a material or element, “is” or “may be” may mean the material or element “consists of” or “consists essentially of” the material or element.


[026] When it is mentioned that “in some embodiments...” a certain element may be present, the element may be present in a suitable embodiment in any part of the specification, not just a suitable embodiment in the same section or textual region of the specification.

[027] When a feature is “selected from” a list, the feature is selected from a list consisting of the specified alternatives (i.e. a list of the alternatives specified and no others).

[028] Claims are embodiments.

5 *Modified Nucleobase Triphosphates*

[029] In one embodiment there is provided a compound of formula (I) or a salt thereof:



(I)

10 [030] Compounds and salts described in this specification may exist as a mixture of tautomers (structural isomers resulting from the migration of a hydrogen atom that exist in equilibrium). Relevant embodiments include all tautomers of compounds of formula (I) or salts thereof.

[031] Atoms of the compounds and salts described in this specification may exist as their isotopes. Embodiments include all compounds of formula (I) where an atom is replaced by one or more of its isotopes (for example a compound of formula (I) where one or more carbon atom is an ^{11}C or ^{13}C carbon isotope, or where one or more hydrogen atom is a ^2H or ^3H isotope).

[032] A suitable salt of a compound of formula (I) is for example a base-addition salt. A base-addition salt is formed by bringing the compound of formula (I) into contact with a suitable organic or inorganic base. A base addition salt may be formed using a suitable organic base like a nitrogen base, for example ammonia or a trialkylamine like triethylamine. A base addition salt may also for example be formed using a suitable inorganic base like an alkali metal or rare earth hydroxide, for example potassium hydroxide, sodium hydroxide, magnesium hydroxide or manganese hydroxide.

[033] In one embodiment there is provided a compound of formula (I) which is a free acid.

[034] In one embodiment there is provided a compound of formula (I) which is a salt.

25 [035] In one embodiment there is provided a compound of formula (I) which is a sodium, potassium, magnesium, or ammonium salt.

[036] In one embodiment there is provided a compound of formula (I) which is a sodium, potassium, or ammonium salt.

[037] In one embodiment there is provided a compound of formula (I) which is a sodium or ammonium salt.

5 [038] In one embodiment there is provided a compound of formula (I) which is a monosodium, disodium, trisodium, tetrasodium, monoammonium, diammonium, triammonium or tetraammonium salt.

[039] In one embodiment there is provided a compound of formula (I) which is a monosodium, disodium, trisodium or tetrasodium salt.

10 [040] In one embodiment there is provided a compound of formula (I) which is a monosodium salt.

[041] In one embodiment there is provided a compound of formula (I) which is a disodium salt.

[042] In one embodiment there is provided a compound of formula (I) which is a trisodium salt.

[043] In one embodiment there is provided a compound of formula (I) which is a monoammonium, diammonium, triammonium or tetraammonium salt.

15 [044] In one embodiment there is provided a compound of formula (I) which is a monoammonium salt.


[045] In one embodiment there is provided a compound of formula (I) which is a diammonium salt.

[046] In one embodiment there is provided a compound of formula (I) which is a triammonium salt.

20 *Synthetic Processes*


[047] In one embodiment there is provided a process for preparing a compound of formula (I) or a salt thereof comprising:

- i. Providing a compound of formula (II) or a salt thereof:



Where PG¹ is a suitable protecting group;

- ii. Immobilising the compound of formula (II) or a salt thereof by linking one of its secondary alcohol groups to a suitable support;
- iii. Capping any remaining secondary alcohol groups with a suitable protecting group PG²;
- iv. Removing the protecting group PG¹;
- v. Reacting the exposed primary alcohol group with a compound of formula (III):



Where R¹ is selected from a hydro group and a C₁₋₃alkyl group;

- vi. Oxidising the resultant phosphorus (III) compound to a phosphorus (V) compound;
- vii. Reacting the phosphorus (V) compound with a tetraalkylammonium pyrophosphate to generate a triphosphate;
- viii. Removing the protecting group PG²; and
- ix. Cleaving the resultant triphosphate from the support to generate a compound of formula (I) or salt thereof.

[048] A protecting group ("PG", for example PG¹ and PG²) is any group suitable for temporarily protecting a reactive centre, for example a hydroxyl group. Suitable protecting groups for the reactive centres disclosed herein may be found for example in "Greene's Protective Groups in Organic Synthesis, Fourth Edition", Greene T. W., Wuts P. G. M.; John Wiley & Sons, Inc. **2007**, doi: 10.1002/0470053488), the contents of all of which are herein incorporated by reference.

[049] A "hydro" group is equivalent to a hydrogen atom. Atoms with a hydro group attached to them can be regarded as unsubstituted.

[050] A "C₁₋₃alkyl group" is a straight chain or branched saturated alkyl group with the indicated number of carbons. Example C₁₋₃alkyl groups include methyl, ethyl, propyl and isopropyl.

[051] In step iii) above, the secondary alcohols to be capped may be those on the ribose part of the molecule.

[052] This overall process is an advantageous preparation of the compound of formula (I) for several reasons:


- There is no need for prior protection of the secondary alcohol groups on the starting protected nucleoside;
- After step 1, unreacted nucleoside can be recovered, which minimizes loss of material;
- The solid-supported nucleosides can be stored for up to 3 months without degradation;
- The route is compatible with automated synthesis;
- The process is robust and has good overall reproducibility;
- Clean phosphorylation crude products are generated that are easy to purify; and
- Gives high phosphorylation yields (typically *ca.* 60 % starting from the nucleoside loaded resin).

[053] In some embodiments R^1 may be a hydro group.

[054] In some embodiments R^1 may be a C_{1-3} alkyl group. It has been observed that when R^1 is a C_{1-3} alkyl group, the phosphoramidite reagent preparation is easier and higher yielding, but performs at least as well in step v above as when R^1 is a hydro group.


[055] In some embodiments R^1 may be methyl.

[056] In one embodiment there is provided a compound of formula (III):



Where R^1 is a C_{1-3} alkyl group.

[057] In one embodiment there is provided a compound of formula (IIIa):



[058] In some embodiments the support may be a solid polymer.

[059] In some embodiments the support may be a solid polymer selected from controlled-porosity glass and polystyrene.

[060] In some embodiments the support may be polystyrene.

[061] In some embodiments the support may be controlled-porosity glass.

[062] In some embodiments the support may be functionalised with a primary amino group. This may form the reactive point of attachment to the support.

[063] In some embodiments the support may be controlled-porosity glass functionalised with a primary amino group (for example Amino-SynBase™).

[064] In some embodiments PG¹ may be selected from trityl, dimethoxytrityl and trimethoxytrityl.

[065] In some embodiments PG² may be selected from acetyl, benzoyl, 2,2,2-trichloroethylcarbonyl, paramethoxybenzyl, methyl, tetrahydropyranyl, triethylsilyl, triisopropylsilyl, trimethylsilyl, *tert*-butyldimethylsilyl and methoxyethyl.

[066] In some embodiments PG² may be acetyl. Where an immobilised molecule is base labile, this allows for an efficient synthesis in which removal of the PG² group and cleavage from the resin may be accomplished in a single step.

[067] In some embodiments PG¹ may be dimethoxytrityl and PG² may be acetyl.


[068] In some embodiments immobilisation of the compound of formula (II) in step i) may occur mainly at the 2'-hydroxy position.

[069] When immobilisation occurs mainly at the 2'-hydroxy position, this may be >50%, >60%, >70%, >80%, >90% or 100% of the total immobilisation (i.e. the total covalent binding of both secondary hydroxyl groups to the support).

[070] In some embodiments the tetraalkylammonium pyrophosphate may be tetrabutylammonium pyrophosphate.


[071] In one embodiment there is provided a process for preparing a compound of formula (I) or a salt thereof comprising:

- i. Providing a compound of formula (II) or a salt thereof:



Where PG¹ is selected from trityl, dimethoxytrityl and trimethoxytrityl;

- ii. Immobilising the compound of formula (II) or a salt thereof by linking one of its secondary alcohol groups to a controlled-porosity glass support;
- iii. Capping any remaining secondary alcohol groups with a protecting group PG² selected from acetyl, benzoyl, 2,2,2-trichloroethylcarbonyl, paramethoxybenzyl, methyl, tetrahydropyranyl, triethylsilyl, triisopropylsilyl, trimethylsilyl, *tert*-butyldimethylsilyl and methoxyethyl;
- iv. Removing the protecting group PG¹;
- v. Reacting the exposed primary alcohol group with a compound of formula (III):




Where R¹ is a C₁₋₃alkyl group;

- vi. Oxidising the resultant phosphorus (III) compound to a phosphorus (V) compound;
- vii. Reacting the phosphorus (V) compound with tetrabutylammonium pyrophosphate to generate a triphosphate;
- viii. Removing the protecting group PG²; and
- ix. Cleaving the resultant triphosphate from the support to generate a compound of formula (I) or salt thereof.


[072] In one embodiment there is provided a process for preparing a compound of formula (I) or a salt thereof comprising:

- i. Providing a compound of formula (II) or a salt thereof:



Where PG¹ is dimethoxytrityl;

- ii. Immobilising the compound of formula (II) or a salt thereof by linking one of its secondary alcohol groups to a controlled-porosity glass support;
- iii. Capping any remaining secondary alcohol groups with a protecting group PG² which is acetyl;
- iv. Removing the protecting group PG¹;
- v. Reacting the exposed primary alcohol group with a compound of formula (III):



Where R¹ is a methyl group;

- vi. Oxidising the resultant phosphorus (III) compound to a phosphorus (V) compound;
- vii. Reacting the phosphorus (V) compound with tetrabutylammonium pyrophosphate to generate a triphosphate;
- viii. Removing the protecting group PG²; and
- ix. Cleaving the resultant triphosphate from the support to generate a compound of formula (I) or salt thereof.

[073] Suitable conditions and reagents to effect each of steps i) to ix) above are known to the skilled person or can be found in the Detailed Description.

[074] In some embodiments immobilising the compound of formula (II) or salt thereof in step ii) may be accomplished by a coupling reagent (for example succinic anhydride catalysed by dimethylaminopyridine when the support is functionalised with a primary amino group).

[075] In some embodiments reaction of the exposed primary alcohol group with a compound of formula (III) may be accomplished using an activator (for example BTT activator or Activator 42®).

[076] In some embodiments the phosphorus (III) compound in step vi) may be oxidised to a phosphorus (V) compound using aqueous pyridine and iodine.

[077] In some embodiments cleaving the triphosphate from the support may be accomplished using basic conditions (for example by treating with AMA). When there is a base-labile support and a base-labile protecting group is chosen for PG², using these conditions allows simultaneous deprotection and cleavage.

Labelled RNA Synthesis

[078] In one embodiment there is provided a composition for preparing a tC^O labelled RNA molecule comprising a compound of formula (I) and a natural ribonucleotide triphosphate.

[079] A “natural ribonucleotide triphosphate” comprises the appropriate natural ribonucleoside

5 with a triphosphate group bonded to the 5' hydroxy position. It is equivalent to a natural ribonucleoside triphosphate. In some embodiments a natural ribonucleotide triphosphate may be selected from cytidine 5'-triphosphate, uridine 5'-triphosphate, adenosine 5'-triphosphate and guanine 5'-triphosphate. A composition of natural ribonucleotide triphosphates (i.e. one comprising a ribonucleotide triphosphate as defined herein) may comprise combinations of varying
10 amounts of these building blocks, in amounts sufficient to construct the target RNA molecule (for example as provided in NTP mix).

[080] In one embodiment there is provided the use of a compound of formula (I) or a salt thereof to enzymatically prepare a tC^O labelled RNA molecule.

[081] A tC^O labelled RNA molecule comprises at least one tC^O residue but is otherwise similar to the
15 natural RNA molecule (i.e. one with an unmodified cytosine residue at the same location as the tC^O residue).

[082] In some embodiments a tC^O labelled RNA molecule may comprise >10%, >20%, >30%, >40%, >50%, >60%, >70%, >80%, >90% or 100% of tC^O residues in place of unmodified cytosine residues.

[083] In some embodiments a tC^O labelled RNA molecule may comprise 10%-20%, 10%-30%, 10%-
20 40%, 20%-50%, 30%-60%, 40%-70%, 50%-80% or 50%-90% of tC^O residues in place of unmodified cytosine residues.

[084] In one embodiment there is provided a process for preparing a tC^O labelled RNA molecule comprising providing a DNA template to composition comprising a compound of formula (I) and a natural ribonucleotide triphosphate (for example a combination of varying amounts of cytidine 5'-
25 triphosphate, uridine 5'-triphosphate, adenosine 5'-triphosphate and/or guanine 5'-triphosphate in amounts sufficient to construct the target RNA molecule, for example as provided in NTP mix), then treating the resultant mixture with an RNA polymerase.

[085] In some embodiments the tC^O labelled RNA molecule may be a tC^O labelled mRNA.

[086] In some embodiments the tC^O labelled RNA molecule may encode for a protein fused to a
30 fluorescent protein. Example fusable fluorescent proteins include Green Fluorescent Protein (GFP) and mFruit family proteins. When the target protein is fluorescent (either inherently or due to a tag), it is possible to simultaneously visualise both the labelled RNA molecule and the protein it is being used to synthesise, giving a greater degree of mechanistic insight.

[087] In some embodiments the tC⁰ labelled RNA molecule may encode for a protein selected from H2B, calmodulin, H2B:GFP and calmodulin-3:GFP.

[088] The terminology “:GFP” means that the protein target preceding the colon is fused to a Green Fluorescent Protein (GFP) family protein.

[089] In some embodiments the tC⁰ labelled RNA molecule may encode for H2B:GFP.

[090] In some embodiments the tC⁰ labelled RNA molecule may encode for calmodulin-3.

[091] In some embodiments the RNA polymerase may be selected from T7 polymerase and SP6 polymerase.

[092] In some embodiments a process for preparing a tC⁰ labelled RNA molecule may be carried out in the presence of transcription buffer (e.g. 5X transcription buffer), magnesium salt (e.g. magnesium(II) chloride) and/or an RNase inhibitor (e.g. Ribolock).

[093] In some embodiments a process for preparing a tC⁰ labelled RNA molecule may be carried out in the presence of transcription buffer (e.g. 5X transcription buffer), magnesium salt (e.g. magnesium(II) chloride) and an RNase inhibitor (e.g. Ribolock). In some embodiments a process for preparing a process for preparing a tC⁰ labelled RNA molecule may be carried out substantially as described in the experimental section (e.g. as detailed in the section headed “H2B:GFP RNA transcription and purification”).

[094] In one embodiment there is provided a kit for preparing a tC⁰ labelled RNA molecule comprising:

- i. A compound of formula (I);
- ii. A composition of natural ribonucleotide triphosphates;
- iii. An RNA polymerase; optionally
- iv. A DNA template; and optionally
- v. Instructions for use.

Labelled RNA Translation

[095] In one embodiment there is provided the use of a tC⁰ labelled mRNA molecule to prepare a protein encoded by the mRNA by translation.

[096] “Translation” refers to the central biological process whereby mRNA is decoded in a ribosome to produce a specific polypeptide, which may fold into an active protein before performing its functions in a cell.

[097] In one embodiment there is provided the use of a tC⁰ labelled mRNA molecule to prepare a protein encoded by the mRNA by in vitro translation (for example, substantially as described in the part of the experimental section (e.g. as detailed under the heading “cell-free translation”).

[098] In some embodiments, in vitro translation may be performed using E.coli bacterial lysates and/or the Expressway[®] mini cell-free expression system.

[099] In one embodiment there is provided the use of a tC⁰ labelled mRNA molecule to prepare a protein encoded by the mRNA by in cell translation (for example, substantially as described in the parts of the experimental section (e.g. as detailed under the headings “cell culture” and “electroporation or chemical transfection”).

[100] In some embodiments, in cell translation may be performed in human neuroblastoma cells (e.g. SH-SY5Y cells).

[101] In one embodiment there is provided the translation of a tC⁰ labelled RNA molecule into a protein.

[102] In one embodiment there is provided the in vitro translation of a tC⁰ labelled RNA molecule into a protein.

[103] In one embodiment there is provided the in cell translation of a tC⁰ labelled RNA molecule into a protein.

[104] In some embodiments the encoded protein may be fused to a fluorescent protein (for example a GFP or mFruit family protein). When this is the case, it is possible to simultaneously visualise both the labelled RNA molecule and the protein it is being used to synthesise, giving a greater degree of mechanistic insight.

[105] In some embodiments the tC⁰ labelled mRNA and the encoded protein may be simultaneously analysed spatiotemporally using confocal microscopy (for example fluorescence confocal microscopy). This is a convenient way to simultaneously monitor a labelled RNA and protein, and RNA containing a fluorescent base analogue has never before been used in such live cell visualisation.

FIGURES

[106] Figure 1: Schematic showing minimally perturbing tC⁰ labelled RNA compared to a common externally labelled RNA

[107] Figure 2: Basic synthetic scheme for the preparation of compound (I).

[108] Figure 3: Incorporation of tC⁰ into full length mRNA by T7 RNA polymerase assisted in vitro transcription. Denaturing agarose bleach gels showing RNA transcripts formed at five different

tC⁰TP/CTP ratios (0-100%). Direct visualization of tC⁰ fluorescence (a) and after ethidium bromide staining (b). RNA samples were heat-denatured (65 °C for 5 min, 1.5 % bleach in the gel) prior to loading. (c) Same RNA transcripts upon harsher denaturation (70 °C for 10 min., 2 % bleach in the gel). The RiboRuler High Range RNA ladder was used.

[109] Figure 4: Incorporation of tC⁰ into full length mRNA by SP6 and T7 RNA polymerase assisted in vitro transcription. (a) Denaturing agarose bleach gels showing RNA transcripts formed at five different tC⁰TP/CTP ratios (0-100%). The produced RNA was visualized directly by tC⁰ fluorescence (left image) and after ethidium bromide staining (right image). The RNA samples were heat-denatured (65 °C for 5 min, 1.5 % bleach in the gel) prior to loading on the gel. (b) Denaturing bleach gels of the same RNA transcripts as in (A) but at stronger denaturing conditions (70 °C for 10 min., 2 % bleach in the gel). The RiboRuler High Range RNA ladder was used.

[110] Figure 5: Spectroscopic characterization of in vitro synthesized tC⁰-modified RNA transcripts. Four reactions charged with different molar fractions of tC⁰TP in the total cytosine triphosphate pool (tC⁰TP + CTP) were performed. The product transcripts were purified to wash out unreacted triphosphates prior to characterization. All reactions were performed as independent duplicates and the results are presented as mean ± standard deviation. a) UV-vis absorption spectra normalized to A = 1 at the RNA band, ca. 260 nm with increased tC⁰-absorption centred at 360 nm growing in with an increasing tC⁰ to C ratio. Inset: tC⁰TP absorption normalized to A = 1 at the tC⁰-band λ_{max} (360 nm). b) Plain bars: Fraction of incorporated tC⁰ (relative to the total amount of incorporated cytosines, i.e. tC⁰ + C) in the transcripts. Checkered bars: Ratio of first-order reaction rate constants for CTP vs. tC⁰TP consumption. c) Solid lines: UV-vis absorption spectra (normalized to A = 1 at the RNA band, ca. 260 nm) showing the tC⁰-band centred at 368–369 nm. Dashed lines: Emission spectra normalized to I = 1 at λ_{max} (457 nm and 459 nm for the 25% and 100% transcript, respectively). For clarity, the emission spectra for the 50% and 75% reactions were omitted. d) Plain bars: Fluorescence quantum yields. Striped bars: Fluorescence lifetime.

[111] Figure 6: Cell-free translation of calmodulin-3. a) Coomassie staining and b) Western Blot (WB) of the in-vitro translation reactions. NTC: no template control; +: kit template DNA control. The PageRuler Prestained Protein Ladder was used. c) Quantification by WB and densitometry analysis (mean of 4 replicates).

[112] Figure 7: Translation efficiency of modified RNA constructs in human cells and validation of tC⁰ as an intracellular tracking probe. The H2B:GFP encoded protein was observed by confocal microscopy and quantified by flow cytometry for each tC⁰-incorporated RNA constructs. Representative images (3x zoomed-in, scale bars: 10 μm), scatter plots and histograms, show the

signal distribution in single living cells at (a, b) 24 h post-electroporation or (c) 48 h post-chemical transfection. The boxplots display the GFP mean fluorescence intensities (MFI GFP) up to 72 h from 3 independent experiments performed in triplicate. (d) Cells overexpressing mRFP-Rab5 (early endosome biomarker) were transfected with 75 % tC⁰ mRNA and followed overtime to validate tC⁰ as an intracellular tracking probe (white arrows) not altering the translation, scale bars: 10 μ m. (e) Cells were analysed 24 h post-electroporation or post-transfection with non-labelled (NL) or Cyanine5-labelled (Cy5) eGFP encoding mRNAs (TriLink®), scale bars: 10 μ m. (f) The impact of tC⁰ or Cy5 incorporation on RNA translation was expressed as the ratio of MFI GFP relative to the non-labelled RNA for all constructs.

[113] Figure 8: Translation efficiency of the modified RNA constructs in human cells and cytotoxicity assessment. Representative confocal images (large view, scale bar: 10 μ m) of RNA-tC⁰ constructs and mRNAs from TriLink® transfected by (a, e) electroporation or (b, f) chemical transfection. (c) Percentages of positive cells for H2B:GFP at 24 h, 48 h and 72 h post-transfection with RNA-tC⁰ constructs. (d) Representative histogram of the GFP signal distribution in single living cells at 48 h post-chemical transfection. Cytotoxicity assessment performed 24 h (g) post-electroporation or (h) post-chemical transfection using the LDH cell membrane integrity assay.

DETAILED DESCRIPTION

Example 1: Synthesis of Modified Nucleobase Triphosphates

[114] Compound (I) may be prepared according to the scheme shown in Fig. 2. Unless otherwise noted reagents were commercially available and used without further purification. The following reagents used for the triphosphorylation were bought from Sigma-Aldrich: DCA deblock for ÄKTA, CAP A for ÄKTA, CAP B1 and B2 for ÄKTA, BTT Activator. ¹H (500 MHz) and ¹³C (126 MHz) NMR spectra were recorded at 300 K on a Bruker 500 MHz system equipped with a CryoProbe. ³¹P (202 MHz) NMR spectra were recorded at 300 K on a Bruker 500 MHz system. All shifts are recorded in ppm relative to the deuterated solvent (DMSO-d₆, CDCl₃ or D₂O).

3-((2R,3R,4S,5R)-5-((bis(4-methoxyphenyl)(phenyl)methoxy)methyl)-3,4-dihydroxytetrahydrofuran-2-yl)-3H-benzo[b]pyrimido[4,5-e][1,4]oxazin-2(10H)-one 1

[115] Compound 1 was prepared according to the literature (Füchtbauer, A. F. *et al.*, *Sci . Rep.* 7, 2393 [2017])

[116] MS (ESI-) [M-H]⁻ = 634.5. ¹H NMR (500 MHz, DMSO-d₆) δ 10.61 (bs, 1H), 7.42 (d, J = 7.7 Hz, 2H), 7.27 – 7.35 (m, 7H), 7.22 (t, J = 7.1 Hz, 1H), 6.90 (dd, J = 8.6, 4.2 Hz, 4H), 6.75 – 6.87 (m, 3H), 6.46 (d, J = 7.8 Hz, 1H), 5.71 (d, J = 3.6 Hz, 1H), 5.49 (bs, 1H), 5.18 (bs, 1H), 4.08 (d, J = 5.3 Hz, 1H), 4.04 (s, 1H), 3.94 (s, 1H), 3.71 (s, 3H), 3.70 (s, 3H), 3.29 (d, J = 4.8 Hz, 1H), 3.16 (d, J = 9.1 Hz, 1H).

5

CPG solid support 3

[117] Amino-SynBase™ CPG 500/110 (LCAA) from LinkTech (Nu. 1397-C025, 1 g, 0.08 mmol) was activated by shaking in trichloroacetic acid 3% in DCE (8 mL, 0.08 mmol) for 18 h. The activated support was then filtered off and washed with 9:1 triethylamine:diisopropylethylamine (20 mL), dichloromethane (20 mL) and diethyl ether (20 mL). The activated support was dried under vacuum for 2 days before use. Subsequently, the support (1 g, 0.08 mmol), succinic anhydride (0.345 g, 3.44 mmol) and N,N-dimethylpyridin-4-amine (0.070 g, 0.57 mmol) were suspended in dry Pyridine (3 mL) under N₂. The reaction mixture was then gently shaken at RT for 4 h. After 4 h, solvent was filtered off and the support washed successively with pyridine (20 mL), dichloromethane (20 mL), diethyl ether (20 mL) and air-dried. Negative ninhydrin test on a small portion of support proved full succinylation. Succinylated CPG could thereafter be kept at room temperature for several months.

15

CPG-supported (2R,3R,4R,5R)-5-((bis(4-methoxyphenyl)(phenyl)methoxy)methyl)-4-hydroxy-2-(2-oxo-2,10-dihydro-3H-benzo[b]pyrimido[4,5-e][1,4]oxazin-3-yl)tetrahydrofuran-3-yl acetate 4

[118] In a 10 mL syringe with PTFE filter, succinylated support 3 (1.420 g, 82 μmol/g, 0.12 mmol), DMAP (0.028 g, 0.23 mmol), DIC (719 μL, 4.64 mmol), 1 (0.076 g, 0.12 mmol) and triethylamine (49 μL, 0.35 mmol) were suspended in pyridine (5 mL). The mixture was gently shaken for 18 h at RT. After 18 h, the syringe was purged and the support washed with pyridine (5 mL), dichloromethane (5 mL) and diethyl ether. Subsequently, in the same syringe, DMAP (0.028 g, 0.23 mmol), diisopropylcarbodiimide (719 μL, 4.64 mmol), triethylamine (49 μL, 0.35 mmol) and 2,3,4,5,6-pentachlorophenol (0.309 g, 1.16 mmol) were added to the support and suspended in pyridine (4 mL). The mixture was gently shaken for 4 h at RT before a solution of piperidine (2 mL, 20% in DMF – for capping of the unreacted carboxylic acids on the support) was added for 1 min (longer exposure time will reduce loading as piperidine cleaves the ester bonds with the nucleoside), then quickly washed away with DMF (3x5 mL), dichloromethane (5 mL) and diethyl ether (5 mL). Finally, the resin was shaken in a CAP A + CAP B mix (50/50 v/v) for 2 hours under argon atmosphere, then washed with DMF (5 mL), dichloromethane (5 mL), diethyl ether (5 mL) and argon-dried (final loading: 13 μmol/g – determined by reading optical density of a DMT solution cleaved from a weighed amount of support - ε=70000 M⁻¹.cm⁻¹ at 498 nm).

30

Final loading can be increased by performing a second coupling with 1 in the same conditions before capping (typical loading after second coupling 20-25 $\mu\text{mol/g}$). Concentrating the reaction mixture and washing the residue multiple times with water and diethyl ether allows recovery of nearly 85% of unreacted nucleoside 1.

5

6-chloro-*N,N*-diisopropyl-4*H*-benzo[d][1,3,2]dioxaphosphinin-2-amine 5

[119] Compound 5 was prepared according to the literature (Ducho, C. *et al.*, *J. Med. Chem.* 50, 1335-1346 [2007]). Briefly, 5-chlorosalicylic acid was reduced with LAH (0.5 equiv.) at -20 °C and the resulting 5-chlorosalicylic alcohol was cyclized into 2,6-dichloro-4*H*-benzo[d][1,3,2]dioxaphosphinine using PCl_3 (1.2 equiv.) and triethylamine (2.3 equiv.) at -20 °C under argon. Low temperature and use of triethylamine as the base were decisive in avoiding rapid and quantitative Arbuzov rearrangement of the desired product into the more stable 2,5-dichloro-3*H*-benzo[d][1,2]oxaphosphole 2-oxide. The crude 2,6-dichloro-4*H*-benzo[d][1,3,2]dioxaphosphinine was subsequently treated with diisopropylamine (3 equiv.) for 2 h at room temperature. The mixture was then filtered under argon, concentrated to dryness and taken in 20 % diisopropylamine in heptane. Quick filtration on a small silica gel plug allowed desired compound 5 as a colourless oil, crystallizing over time at -20 °C. Any attempt of more thorough column chromatography on compound 5 would lead to quantitative Arbuzov rearrangement.

[120] ^1H NMR (500 MHz, DMSO-d_6) δ = 7.23 (dd, J = 8.6, 2.6 Hz, 1H), 7.20 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 8.6 Hz, 1H), 5.06 (dd, J = 14.7, 5.2 Hz, 1H), 4.89 (dd, J = 19.6, 14.8 Hz, 1H), 3.53 – 3.63 (m, 2H), 1.15 – 1.19 (dd, J = 8.0, 7.0 Hz, 12H). ^{31}P NMR (202 MHz, DMSO-d_6) δ = 136.00 (s, 1P).

***Bis*(tetrabutylammonium) dihydrogen diphosphate 6**

[121] Compound 6 was prepared according to the literature (Warnecke, S. & Meier, C., *J. Org. Chem.* 74, 3024-3030 [2009]).

[122] ^1H NMR (500 MHz, D_2O) δ 3.04 – 3.13 (m, 16H), 1.53 (bs, 16H), 1.24 (h, J =7.3, 16H), 0.83 (t, J =7.4, 24H). ^{31}P NMR (202 MHz, D_2O) δ = -10.78 (s, 2P).

((2*R*,3*S*,4*R*,5*R*)-3,4-dihydroxy-5-(2-oxo-2,10-dihydro-3*H*-benzo[*b*]pyrimido[4,5-*e*][1,4]oxazin-3-yl)tetrahydrofuran-2-yl)methyl triphosphate 7

[123] Reactions were performed in a 5 mL syringe with PTFE filter loaded with 4 (800 mg, 0.016 mmol) under an argon atmosphere and with shaking.

[124] Steps were performed as following:

- a. *5'-DMT removal*: the support was washed with a flow of DCA deblock until the filtrate was colourless, then washed with ACN (5 x 5 mL).
- b. *Coupling*: N,N-diisopropyl-4H-benzo[d][1,3,2]dioxaphosphinin-2-amine 5 (345 mg, 1.36 mmol) was dissolved in 4.8 mL ACN and reacted portion wise with the support (3 equal couplings with reaction times 60 s – 60 s – 90 s respectively). To each coupling, an activator (e.g. BTT activator (2.4 mL) or Activator 42) was also added. The support was subsequently washed with ACN (3x5 mL).
- c. *Oxidation*: Pyridine/Water/Iodine (9/1/12.7 v/v/w, 5mL) for 45 s, followed by ACN wash (3x5 mL) and drying of the support in an argon flow.
- d. *Triphosphorylation*: Two injections of bis(tetrabutylammonium) dihydrogen diphosphate 6 (0.5 M, 5 ml) for 15 min and 18 hours, respectively. The support was subsequently rinsed with DMF (5 mL), water (3x5 mL), ACN (5 mL) and then dried in an argon flow.
- e. *Cleavage and Purification*: Cleavage of the triphosphate was done in 2 h at room temperature with AMA (50/50 v/v mix of 23 % aq. NH₄OH and 40 % aq. methylamine, 5 mL). After 2 hours, the AMA filtrate was purged in a round-bottom flask and the support was rinsed 3 times with 23% aq. NH₄OH solution. After freeze-drying of the mixture, purification by HPLC (Waters Acquity HSS T3 column, 2.1x50 mm, 0.4 mL/min, 2 to 99 % 50 mM NH₄OAc in water 80:20 EtOH) was performed to allow compound 7 (5.6 mg, 62.0 % determined from UV absorbance) as a light-yellow solid (ammonium salt). The same level of purity could be achieved with ion-exchange HPLC using a semi-preparative Dionex DNAPac PA100 column (9 x 250 mm) on an ÄKTA pure 25 HPLC system using a gradient from water to 20 % 1M NH₄HCO₄ (pH 7.8) in 30 min at a flow rate of 4 mL/min.

[125] HRMS (ESI-TOF) m/z calc. for C₁₅H₁₈N₃O₁₅P₃ [M+H]⁺: 574.0029, found : 574.0013; m/z calc. for C₁₅H₁₈N₃O₁₅P₃ [M-H]⁻: 571.9878, found: 571.9872. ¹H NMR (500 MHz, D₂O) δ 7.44 (s, 1H), 6.84 – 6.94 (m, 3H), 6.79 (dd, J = 7.5, 1.7 Hz, 1H), 5.91 (d, J = 4.9 Hz, 1H), 4.36 (t, J = 4.8 Hz, 1H), 4.29 (t, J = 5.1 Hz, 1H), 4.25 (d, J = 4.1 Hz, 3H). ¹³C NMR (126 MHz, D₂O) δ 155.8, 154.8, 142.4, 129.4, 124.9, 124.3, 122.3, 116.6, 88.8, 82.8, 73.4, 69.7, 64.5. ³¹P NMR (202 MHz, D₂O) δ -10.89 (d, J = 18.5 Hz, 1P), -11.46 (d, J = 19.7 Hz, 1P), -23.21 (t, J = 19 Hz, 1P).

[126] Compound (I) can also be made by a slightly modified route wherein the coupling step (b above) is carried out with a modified phosphoramidite such as 6-chloro-N,N-diisopropyl-4-methyl-4H-benzo[d][1,3,2]dioxaphosphinin-2-amine 8 (compound (IIIa) above). This reagent has been found to be more easily prepared, and compound 8 is obtainable in a yield of 60% compared to around 3-10% for the preparation of compound 5 under the conditions in this specification.

6-chloro-*N,N*-diisopropyl-4-methyl-4*H*-benzo[d][1,3,2]dioxaphosphinin-2-amine 8

[127] 5-chloro-2-hydroxybenzaldehyde was reacted with methylmagnesium bromide (2.5 equiv.) at -20 °C and the resulting 4-chloro-2-(1-hydroxyethyl)phenol was cyclized into 2,6-dichloro-4-methyl-4*H*-benzo[d][1,3,2]dioxaphosphinine using PCl₃ (1.2 equiv.) and triethylamine (2.3 equiv.) at -20 °C under argon. The crude 2,6-dichloro-4-methyl-4*H*-benzo[d][1,3,2]dioxaphosphinine was subsequently treated with diisopropylamine (3 equiv.) for 2 h at room temperature. The mixture was then filtered under argon, concentrated to dryness and taken in 20 % diisopropylamine in heptane. Quick filtration on a small silica gel plug furnished desired compound 8 as a colourless oil.

[128] ¹H NMR (500 MHz, DMSO-*d*₆) δ = 6.96 (d, *J* = 8.5 Hz, 1H), 6.87 (d, *J* = 8.5 Hz, 1H), 6.74 (d, *J* = 8.4 Hz, 1H), 5.19 – 5.26 (m, 1H), 5.16 (dq, *J* = 10.4, 6.6 Hz, 1H), 3.57 (tdt, *J* = 13.6, 10.6, 6.8 Hz, 2H), 1.63 (d, *J* = 6.6 Hz, 3H), 1.55 (d, *J* = 6.4 Hz, 2H), 1.16 – 1.19 (m, 24H). ³¹P NMR (202 MHz, DMSO-*d*₆) δ = 137.63 (s, 1P), 127.90 (s, 1P).

Example 2: Cell-free in vitro transcription

[129] The utility of compound (I) in RNA labelling was demonstrated by its cell-free in vitro transcription to produce fluorescent full-length messenger RNA (mRNA), from a DNA template encoding for H2B histone protein fused to GFP (H2B:GFP).

[130] The template was codon optimized to limit the number of C repeats, preventing self-quenching and improving brightness. Efficient transcription and tC⁰ incorporation was observed using two different bacteriophage RNA polymerases, T7 and SP6 at tC⁰TP/canonical CTP ratios ranging from 0 to 100% (full replacement), as demonstrated by agarose bleach gel electrophoresis (Fig. 3a for T7 and Fig. 4a for SP6).

[131] All RNA transcripts run as one single band on the gels, with a size corresponding to the expected 1247 nt mRNA product (H2B:GFP), demonstrating that the full-length mRNA is formed. The tC⁰-containing mRNA bands could be directly visualized upon 302 nm excitation (Fig. 4a); the increasing band intensities with increasing tC⁰TP/CTP reaction ratio supported successful concentration-dependent incorporation of tC⁰. Re-visualization of the gel after ethidium bromide staining (Fig. 3b) provided a further qualitative indication that tC⁰ incorporation does not reduce the reaction yield.

[132] Furthermore, no shorter transcripts were observed, suggesting that T7 processes tC⁰TP correctly and without premature abortion. Higher order bands are apparent in all lanes of the gel (Fig. 3b and 4b) but were removed upon heat denaturation, suggesting the presence of RNA secondary

structures. Notably, this feature appears independently of the CTP/tC^OTP-ratio, indicating that the effect is not specific to the modified cytosine base.

[133] Therefore, these results demonstrate that tC^O can be successfully incorporated into full-length RNA transcripts even under conditions where all canonical CTP is replaced with tC^OTP (0% CTP; i.e. 100% C-labelling efficiency).

Example 3: Spectroscopic characterization of in vitro synthesized tC^O-modified RNA transcripts

[134] A spectroscopic approach was used to quantify the incorporation efficiency of tC^OTP, compared to the canonical CTP. To enable this, all RNA transcripts were purified using a Monarch RNA Cleanup kit, ensuring complete removal of unreacted tC^OTP. Absorption spectra (Fig. 5a) showed the appearance of a band centred at *ca.* 370 nm in samples with the incorporated cytosine analogue tC^O, consistent with the spectral profile of this fluorescent base analogue (Fig. 5a).

[135] By relating the absorption of the purified RNA transcripts at 260 nm, which reflects their total concentration, to the absorption at 370 nm (emanating exclusively from tC^O), the relative rate constants for the incorporation of CTP and tC^OTP (*k*C and *k*tC^O, respectively, see later for details). The calculated quotients *k*C/*k*tC^O (Fig. 5b) have values close to or slightly above unity (0.96–1.4, Fig. 5b), demonstrating that the T7 polymerase displays no substantial preference for the canonical CTP over tC^OTP in the *in vitro* reactions. This supports that the tricyclic chemical modification of cytosine is indeed minimally perturbing in the transcription process.

[136] The emissive behaviour of tC^O was also investigated in the mRNA transcripts exploring the relation to the tC^OTP fraction added to the initial reaction mixture. A substantial decrease in fluorescence quantum yield (from 0.18 to 0.09, Fig. 5d) was observed with increasing tC^O incorporation. This was accompanied by a decrease in fluorescence lifetime (from 4.3 ns to 3.2 ns) and a slight redshift of the emission spectrum (*ca.* 4 nm, Fig. 5c).

[137] This may be ascribed to electronic interaction (coupling) of molecular states of the tC^O fluorophore and self-quenching effect caused by the expected increasing concentration of vicinal tC^Os. Importantly, this quenching effect at high tC^O fractions is balanced by the large overall number of incorporations and does not prevent visualization of the mRNA, even for transcripts where all Cs are replaced by tC^Os.

Example 4: Translation of tC^O-labelled mRNA in bacterial lysates

[138] In order to verify the functionality of the tC^O-labelled mRNA transcripts the translation of a tC^O-labelled Calmodulin-3 mRNA in cell-free conditions using bacterial lysates was investigated. The

labelled mRNAs encoding for the 17 kDa protein were transcribed from a commercial Calmodulin-3 DNA template plasmid using the same tC^O TP/CTP ratios as for the H2B:GFP encoding mRNA (0 to 100% of tC^O TP). After RNA purification and cell-free translation, the presence of Calmodulin-3 was confirmed by Coomassie staining (Fig. 6a) as well as Western Blot (Fig. 6b). Satisfactorily we observed stable expression levels when increasing the tC^O content of the transcripts, ranging from 80%–137% of that obtained using a commercially available, unlabelled DNA template control (Figure 6c).

Example 5: Translation efficiency of tC^O -labelled mRNA in human cells

[139] Electroporation was used to introduce in vitro-transcribed tC^O -labelled mRNA transcripts into human neuroblastoma SH-SY5Y cells. Taking advantage of them encoding for a fluorescent fusion protein with nuclear localization (H2B:GFP), the translation was detected by fluorescence (Figs. 7 and 8). To improve stability and reduce cytosolic degradation, the mRNAs were capped with a 5'-Cap 0 analogue and 3'-protected by poly-adenylation (by *ca.* 300 nt).

[140] Live-cell confocal microscopy and flow cytometry showed that GFP fluorescence in the cell nuclei could be detected in 32, 25, 18, and 12% of the cells 24 hours post-electroporation for mRNA's containing 25, 50, 75 and 100% of tC^O , respectively (Fig. 7a and Fig. 8a). In comparison, the transfection efficiency with unmodified mRNA was 46%. This provides the first observation that a fluorescent base analogue-modified RNA transcript can be accurately and efficiently translated by human ribosomal machineries, resulting in a correctly localized and folded protein product.

[141] Using flow cytometry, the levels of H2B:GFP fluorescence in the cells was quantified (Fig. 7b), showing a decrease in mean cellular H2B:GFP fluorescence intensity upon increasing the percentage of tC^O in the transcript (approximately one order of magnitude difference between 0% and 100% of tC^O (Fig. 7a and 7f). This suggests that under these conditions, translation, as opposed to transcription, is somewhat impeded by the tC^O modification, especially at the highest incorporation fraction.

[142] Importantly, no evidence was found of mRNA-induced cell toxicity at 24 hours post-electroporation (Fig. 8g). Of significant note is that the mean fluorescence intensity of translated protein in cell cultures electroporated with a commercial enhanced GFP-encoding mRNA tagged with Cy5 *via* conjugation to UTPs (*ca.* 25% of all U positions of this mRNA are labelled) was only 17% of that in cell cultures electroporated with the corresponding non-labelled sequence (Fig. 7f). This is comparable to the effect of a 50%–75% labelled tC^O transcript, which suggests that Cy5 modifications affect the ribosome translation capability more than tC^O does.

[143] It is evident from the images in Fig. 7a and 7e that neither the fluorescent tC^O -labelled mRNA nor the Cy5-labelled mRNA can be detected inside cells after electroporation due to their low/diffused

cytoplasmic concentration. The delivery of the H2B:GFP transcripts was therefore probed using a chemical transfection reagent (lipofectamine), which is also more relevant from a drug delivery perspective. This resulted in the successful production of H2B:GFP (Figs. 7c, 8b and 8d) irrespective of the tC⁰ content, but with albeit lower delivery efficiencies (3.8–4.6 % of H2B:GFP-positive cells for tC⁰-labelled transcripts vs. 12-32% post-electroporation). This reflects the relatively poor transfectability of SH-SY5Y cells compared to many other cell lines rather than the transfectability of tC⁰-labelled constructs, as further supported by the finding that cultures transfected with non-modified mRNA displayed a virtually identical response (4.5 % of cells expressing H2B:GFP).

[144] H2B:GFP fluorescence was found to increase gradually with time between 24 h and 72 h (Fig. 7c), which is a contrasting behaviour compared to electroporated cells (Fig. 7a), suggesting that the lipofectamine-mRNA complexes are continuously internalized and, potentially, that endocytosed complexes progressively release more transcripts with time, counteracting the degradative effect in the cytosol.

[145] When delivered using lipofectamine, the tC⁰-labelled mRNAs were found to promote very similar H2B:GFP translation compared to the corresponding non-labelled mRNA, as indicated by the fluorescence levels in Fig. 7f. The Cy5-tagged mRNA, on the other hand, results in an average fluorescence level that is 80 % lower than that of its corresponding non-labelled transcript. This suggests that tC⁰ does not impair the ability of mRNA to be processed by ribosomes upon chemical transfection, possibly because of an absence of charge and reduced steric hindrance, whereas Cy5, currently the most common commercial fluorophore for mRNA labelling, quite considerably impacts the translation process and/or interferes with a native-like uptake process of mRNA since it introduces significant amphiphilicity to the mRNA and, hence, possibly non-native interactions between the CY5-mRNA and the lipophilic membrane constituents.

[146] Importantly, the complexation of the tC⁰-labelled mRNA with lipofectamine enabled its direct visualization inside cells using live cell confocal microscopy (Fig. 7c). This represents the first observation of fluorescent base analogue-labelled nucleic acids inside live cells. It was also found to be possible to simultaneously visualize, in real time, the uptake and subsequent translation of a fluorescent base analogue-modified mRNA by time-lapse recordings. We observed co-localization of the tC⁰ signal with an mRFP-labelled Rab5 protein, thus highlighting that the mRNA transits through the early endosome (Fig. 7d). Consequently, this technology allows tracking of both the intrinsically labelled mRNA transcripts and their translation products live, to gather spatiotemporal information on the translation product, even with as low as 50 % tC⁰ content. This demonstrates the flexibility and

versatility of this new labelling approach where fine-tuning of tC⁰ content can be utilized to optimize the mRNA for specific drug delivery applications.

Example 6: Biochemical Methods

Generation of H2B:GFP DNA template

[147] The original coding sequence for H2B:GFP was taken from pCS2-H2B:GFP plasmid (Addgene, Plasmid #53744, manually codon-optimized to minimize the occurrence of poly-Cn stretches (n<3), in silico-assembled with an additional T7 promoter and other desired features (Shine-Dalgarno/Kozak consensus sequences for enhancement of translation and a 3xStop, respectively at the 5' and 3' of the coding sequence itself, plus the needed HindIII/SnaBI restriction sites, to generate the ligation-prone sticky ends) and ordered from Twist Bioscience as a synthetic gene block. The obtained sequence was then PCR-amplified, using a Phusion Hot Start High-Fidelity Taq (Thermo Scientific), and subcloned into a HindIII/SnaBI-digested (Fast Digest enzymes, Thermo Scientific) empty pCS2 backbone. After ligation with T4 ligase for 1 h at room temperature (Roche), DH5 α E. coli competent cells (Invitrogen) were transformed following the recommended protocol, and obtained colonies were screened by colony-PCR. The selected colony was then inoculated into a midiprep-scale volume of liquid Luria-Bertani growth medium (VWR) and plasmid DNA isolated using a PureLink Fast Low-Endotoxin Midi Plasmid Purification Kit (Thermo Scientific). The purified plasmid was finally digested again with HindIII/SnaBI and gel-purified, to generate the transcription template with the desired size.

[148] Primers (Eurofins Genomics):

Twist-H2B.F: GAAGTGCCATTCCGCCTGAC

Twist-H2B.R: CACTGAGCCTCCACCTAGCC

H2B:GFP RNA transcription and purification

[149] *In-vitro* transcription reactions, for T7 and SP6 polymerases (Thermo Scientific), were assembled as recommended by the corresponding protocols, with a few modifications that resulted in a consistently increased yield in all conditions:

1. 5X Transcription buffer - 10 μ l
2. NTP Mix, 10 mM each (2 mM final concentration) – volume depending on batch for tC⁰TP
3. Linearized template DNA 1 μ g – volume depending on concentration
4. RiboLock RNase Inhibitor - 1.25 μ l (50 U)
5. T7/T3/SP6 RNA Polymerase - 3 μ l (60 U, double compared to recommendations)

6. MgCl₂ 4 mM final concentration (increased as recommended by Thomen, P. *et al. Biophys. J.* 95, 2423-2433 [2008])

7. DEPC-treated Water qsp 50 µl

[150] *In-vitro* transcriptions were always performed at 20 °C for 14 h, then RNAs were purified using a Monarch RNA Cleanup kit (NEB), or homemade equivalent buffers and regenerated columns following the same rationale. It was possible to partially recover unreacted tC⁰TP from the transcription mixtures by HPLC to re-use for further assays. For cellular studies, each batch of RNA was then enzymatically added with a polyA tail (with a Poly(A) Polymerase, NEB protocol #M0276 with incubation extended to 1 hour) and a Cap 0 analogue (using a Vaccinia capping system, NEB protocol #M2080), following the recommended procedures.

Denaturing bleach-agarose gels

[151] For a qualitative check of all *in vitro* synthesized RNAs, a denaturing agarose gel was run, in presence of 1,5 % bleach (Sigma Aldrich), as recommended in Aranda, P. S., LaJoie, D. M. & Jorcyk, C. L., *Electrophoresis* 33, 366-369 [2012]. RNAs were first mixed with a 6x DNA loading dye (Invitrogen) and then heat-denatured at 70°C for 10 min in a heating block, then immediately transferred and kept on ice. The RiboRuler High Range RNA Ladder (Thermo Scientific) underwent the same treatment; 2 µl of RNA ladder were loaded along the samples and the gel was run at constant voltage (70 V) for 1 h and then imaged, under UV transillumination (302 nm) using a ChemiDoc Touch (BioRad). To counterstain the whole gel, and especially the lanes without tC⁰TP-containing samples, a standard ethidium bromide staining was finally performed at room temperature for 10 min and gentle rocking, followed by two washes in TAE and then a final wash in distilled water (10 min each).

Cell culture

[152] Human neuroblastoma SH-SY5Y cells (Sigma-Aldrich) were grown in a 1:1 mixture of minimal essential medium (HyClone) and nutrient mixture F-12 Ham (Sigma-Aldrich) supplemented with 10% fetal bovine serum (FBS), 1% non-essential amino acids (Lonza) and 2 mM L-glutamine. For the tracking experiments, an in-house generated model of human hepatic Huh-7 cells stably overexpressing mRFP-Rab5 were cultured in DMEM/GlutaMax/High glucose (Gibco) supplemented with 10 % FBS. The cells are detached with trypsin-EDTA 0.05 % (Gibco) and passaged twice a week.

Electroporation or chemical transfection

[153] Cells were electroporated either with 9.7 µg of tC⁰TP (for *in vitro* incorporation experiments) or 100 ng of tC⁰-labelled mRNA per 10⁵ cells (for *in vitro* translation, cytotoxicity assessment, flow

cytometry analysis and confocal microscopy), using a Neon Transfection System (Invitrogen, Carlsbad, CA, US) and following the protocol for 10 μ L Neon Tip provided by the manufacturer, with a triple pulse of 1200 V and a pulse width of 20 ms. For chemical transfection, SH-SY5Y cells were seeded one day prior transfection at a density of 0.8×10^6 cells/mL, in 48-well plate or glass-bottomed culture dishes for flow cytometry or confocal microscopy analysis, respectively. Lipofectamine MessengerMAX was used as chemical reagent for transfection according to the manufacturer's instructions. Briefly, the reagent was diluted and incubated for 10 min at room temperature in Opti-MEM medium. The tC^O-mRNA constructs were added to the reagent to reach a 1:1 final ratio reagent-mRNA (v/w), followed by a 5 min incubation at room temperature allowing the complex mRNA-lipid to form. Cells were incubated with this complex up to 72 h. To address the impact of the dye incorporation on RNA translation, SH-SY5Y cells were electroporated or chemical transfected with commercially available non-labelled (NL) or Cyanine5-labelled (Cy5) eGFP encoding mRNAs (Trilink®) has described here.

Cytotoxicity assessment

[154] Cell membrane integrity was determined using the Pierce™ LDH Cytotoxicity Assay Kit (Invitrogen) according to the manufacturer's instructions. Briefly, LDH released in the supernatants of cells 24 h post-electroporated or post-transfected with tC^O-labelled mRNA, or Cy5-mRNA, was measured with a coupled enzymatic assay which results in the conversion of a tetrazolium salt into a red formazan product. The absorbance was recorded at 490 nm and 680 nm. The toxicity was expressed as the percentage of LDH release in supernatant compared to maximum LDH release (supernatant + cell lysate). Data are means $\pm$ SD from three experiments performed in triplicate.

Flow cytometry

[155] Following electroporation of tC^O-labelled mRNA, cells were seeded in 48-well plate (2.105×10^5 cells/well) and the expression of H2B:GFP in cells was quantified by flow cytometry. Briefly, 24 h, 48 h or 72 h post-electroporation or post-transfection with tC^O-mRNA, non-labelled mRNA or Cy5-mRNA, cells were harvested and analysed on a Guava EasyCyte 8HT flow cytometer (Millipore). Data are mean fluorescence intensities $\pm$ SD of gated single living cells from three experiments performed in triplicate. The average fluorescence intensities were baseline corrected by subtracting the signal for RNase-free water electroporated or transfected cells. All flow cytometry data were analysed in Flowing software (version 2.5.1) and displayed using R (<http://www.R-project.org/>). H2B:GFP: Excitation 488 nm; Emission 525-530 nm.

Confocal microscopy

[156] After electroporation, cells were seeded in glass-bottomed culture dishes (MatTek glass-bottomed or in 4-sectors subdivided CELLview dishes; 2.105 cells/chamber). For tracking experiment, the Huh-7 cells stably overexpressing mRFP-Rab5 were incubated with lipofectamine/tC⁰-mRNA complex and time-lapse was recorded up to 20 h post-chemical transfection. Confocal images were acquired on a Nikon C2+ confocal microscope equipped with a C2-DUVB GaAsP Detector Unit and using an oil-immersion 60 × 1.4 Nikon APO objective (Nikon Instruments, Amsterdam, Netherlands). Data were processed with the Fiji software. H2B:GFP: Excitation 488 nm; Emission 495-558 nm. tC⁰-labelled mRNA: Exc. 405 nm; Em. 447-486 nm. Cy5-labelled mRNA: Exc. 640 nm; Em. 652-700 nm. mRFP-Rab5: Exc. 561 nm; Em. 565-720 nm.

Cell-free translation

[157] Cell-free translation reactions were performed using E. coli bacterial lysates and an Expressway™ Mini Cell-Free Expression System (Thermo Scientific). Calmodulin-like 3 protein is provided as a positive control plasmid (pEXP5-NT/CALML3) in the kit itself; this DNA vector contains a T7 polymerase promoter and a 6xHis tag, hence it was first in-vitro-transcribed in presence of the desired concentrations of tC⁰TP (vide supra). The obtained RNAs, once purified, were used as templates for the cell-free translation reaction according to the manufacturer's recommendations: E. coli slyD- Extract - 20 µl; 2.5X IVPS E. coli Reaction Buffer (-A.A.) - 20 µl; 50 mM Amino Acids (-Met) - 1.25 µl; 75 mM Methionine* - 1 µl; T7 Enzyme Mix - 1 µl (omitted when using tC⁰-labelled RNAs); DNA Template - 1 µg (when testing the tC⁰-labelled RNAs, added the same amount of RNA instead); DNase/RNase-free distilled water qsp 50 µl.

Coomassie staining and Western Blots

[158] Protein samples from in vitro translation experiments were quantified with a Qubit Protein Assay kit (Thermo Scientific), mixed with 6x SDS Laemmli reducing buffer (Alfa Aesar), then heat-denatured at 85 °C for 10 min and kept at room temperature until needed. Samples were generally run in 1 mm polyacrylamide 4-20 % Novex MES/SDS gels (Thermo Scientific) and using a Mini Gel Tank, with the PSU set at constant voltage (200 V). For Coomassie staining, the gel was then washed three times in boiling water, to remove excess of SDS, on a benchtop shaker; a 1x Coomassie non-toxic staining solution was added to the gel and microwaved until initial boiling.

[159] Gel was finally washed after the appropriate incubation time, to remove excess of background noise, in distilled water and imaged using a ChemiDoc Touch. For Western Blot, the gels were blotted

onto PVDF LF ethanol-activated membranes (BioRad) with a TransBlot semi-dry system (BioRad), according to manufacturer's recommendations (settings for 1mm-thick gels and mixed weight proteins). PVDF membranes were then washed 5 min in TBS-T (TBS and 0,1 % Tween-20, Sigma Aldrich), blocked in 5 % milk in TBS for 1 h at room temperature and incubated with the appropriate primary antibody dilutions.

[160] After 3x5 min washes in TBST and an incubation of 1 h with the corresponding HRP-conjugated secondary antibodies, the membrane was washed again three times in TBS-T, once in TBS and once more in distilled water. Finally, membranes were incubated with a minimal volume of SuperSignal West Pico PLUS (Thermo Scientific) and imaged with a ChemiDoc Touch. Primary antibodies: mouse monoclonal anti-6xHistidine tag (Invitrogen) and mouse monoclonal anti-GAPDH (ref. 437000, Invitrogen), both diluted 1:1000 in 3 % BSA/TBS-T. Secondary antibodies: HRP-conjugated polyclonal goat anti-Ms and anti-Rb Cross-Adsorbed IgG (H+L) (ref. A16072 and A16104, Invitrogen), used at 1:10000 dilution in TBS-T.

Example 8: Spectroscopic Methods

[161] The tC⁰-RNA products from the cell-free transcription reactions (prior to polyadenylation and capping, see Methods: Bio for details) were measured as received, i.e. in RNase free Milli Q water. All measurements were carried out at room temperature (ca. 22°C) in a 3.0 mm path length quartz cuvette, with a sample volume of ca. 60 µL.

Steady state absorption

[162] Absorption spectra were recorded on a Cary 5000 (Varian Technologies) spectrophotometer with a wavelength interval of 1.0 nm, integration time of 0.1 s, and a spectral band width (SBW) of 1 nm. All spectra were baseline corrected by subtracting the corresponding absorption from the solvent only. A second-order polynomial Savitzky-Golay (five points) smoothing filter was applied to all spectra. For samples exhibiting significant scattering, as evidenced by characteristic absorption in the long wavelength region (here for $\lambda > 475$ nm), an additional correction was applied. The scattering contribution ($A_{scatter}$) to the absorption was in such cases fitted (using absorption at 550–475 nm as input) to the Rayleigh scattering function (equation S1), where c is a proportionality constant and A_0 a constant, and then subtracted for all wavelengths.

$$A_{scatter}(\lambda) = \log \left(\frac{1}{1 - c \times \lambda^{-4}} \right) + A_0 \quad (S1)$$

Steady state emission

[163] Emission spectra were recorded on a SPEX Fluorolog (Jobin Yvon Horiba) fluorimeter with excitation at 356 nm. Emission was collected at a right angle with an integration time of 0.1 s and wavelength interval of 1 nm. Monochromator slits were adjusted to achieve optimal signal output, leading to SBWs in the interval 1.5–2.5 nm on both the excitation and emission side. Emission spectra were corrected for Raman scattering by subtracting the corresponding emission from a sample containing only solvent. A second-order polynomial Savitzky-Golay (five points) smoothing filter was applied to all spectra.

Fluorescence quantum yield determination

[164] Sample fluorescence quantum yields (Φ_F) were determined relative to a solution of quinine sulphate (Sigma) in 0.5 M H₂SO₄ ($\Phi_{F,REF} = 0.546$) and calculated according to equation S2.

$$\Phi_F = \Phi_{F,REF} \times \frac{\int_{\lambda_i}^{\lambda_f} I_S(\lambda) d\lambda}{\int_{\lambda_i}^{\lambda_f} I_{REF}(\lambda) d\lambda} \times \frac{A_{REF}}{A_S} \times \frac{\eta_S^2}{\eta_{REF}^2} \quad (S2)$$

[165] Emission spectra for the sample, $I_S(\lambda)$ and reference, $I_{REF}(\lambda)$, were integrated between $\lambda_i = 365$ nm and $\lambda_f = 700$ nm. Absorption at the excitation wavelength (356 nm) for the sample (A_S) and reference (A_{REF}) were in the interval 0.05–0.11 for all samples. Adopted solvent refractive indices for the samples (water) and reference (0.5 M H₂SO₄) were $\eta_S = 1.333$ and $\eta_{REF} = 1.339$, respectively. All quantum yields are presented as mean $\pm$ standard deviation of two independent cell-free transcription reactions.

Time-resolved emission

[166] Fluorescence lifetimes were determined using time-correlated single photon counting (TCSPC). Samples were excited using an LDH-P-C-375 (PicoQuant) pulsed laser diode with emission centred at 377 nm (FWHM pulse width was 1 nm and 70 ps with respect to wavelength and time, respectively), operated with a PDL 800-B (PicoQuant) laser driver at a repetition frequency of 10 MHz. Sample emission (458 nm, SBW = 10 nm) was collected at a right angle, through an emission polarizer set at 54.9° (magic angle detection). Photon counts were recorded on a R3809U 50 microchannel plate PMT (Hamamatsu) and fed into a LifeSpec multichannel analyser (Edinburgh Analytical Instruments) with 2048 active channels (24.4 ps/channel), until the stop condition of 104 counts in the top channel was met. The instrument response function (IRF) was determined using a frosted glass (scattering) modular insert while observing the emission at 377 nm (SBW = 10 nm).

Fitting of fluorescence lifetimes

[167] The intensity decays were fitted with IRF re-convolution to the multiexponential model shown in equation S3.

$$I(t) = \int_0^t IRF(t') \sum_{i=1}^n \alpha_i e^{-\frac{t-t'}{\tau_i}} dt' \quad (S3)$$

[168] The least-square re-convolution fitting procedure was carried out using the DecayFit software (<http://www.fluortools.com/software/decayfit>). All decays were fitted to a tri-exponential ($n = 3$) model. The presented lifetimes are amplitude-weighted average lifetimes ($\bar{\tau}$), calculated using the pre-exponential factors α_i and lifetimes (τ_i) according to equation S4. The fitting parameters for the decays are shown in Table 2.

$$\bar{\tau} = \sum_{i=1}^n \alpha_i \tau_i \quad (S4)$$

Table 2: Fitted lifetime parameters for the TCSPC experiments. The χ^2 -value (Chi-Square) was evaluated to indicate goodness of fit.

	transcript	α_1	τ_1 (ns)	α_2	τ_2 (ns)	α_3	τ_3 (ns)	$\bar{\tau}$ (ns)	χ^2
Set 1	25%	0.19	0.67	0.49	3.2	0.31	5.6	3.5	1.09
	50%	0.28	0.71	0.43	2.8	0.30	5.2	2.9	1.06
	75%	0.33	0.68	0.45	2.6	0.23	5.2	2.5	1.12
	100%	0.33	0.48	0.44	2.1	0.23	4.8	2.2	1.00
Set 2	25%	0.21	0.63	0.40	2.8	0.34	5.3	3.3	0.99
	50%	0.26	0.64	0.41	2.7	0.30	5.2	3.0	1.03
	75%	0.29	0.50	0.42	2.1	0.22	4.8	2.4	1.02
	100%	0.37	0.47	0.40	1.9	0.39	4.5	2.0	1.01

Cell-free transcription reaction kinetics

[169] The ratio of the rate constants for cytosine vs. tC^O incorporation (k_C/k_{tC^O}) was calculated using the absorption spectra of the tC^O -RNA transcripts (A_{260} and A_{369}), and triphosphate initial concentrations ($[CTP]_0$ and $[tC^OTP]_0$) as input. Equations S5 and S6 follows upon assuming first order reaction kinetics with respect to the triphosphate species [CTP] and $[tC^OTP]$.

$$\frac{d[CTP]}{dt} = -k_C \times [CTP] \quad (S5)$$

$$\frac{d[tC^OTP]}{dt} = -k_{tC^O} \times [tC^OTP] \quad (S6)$$

[170] Solving S5 and S6 for the respective rate constants renders equation S7, in which [C] and $[tC^O]$ denote the concentration of incorporated C and tC^O , respectively.

$$\frac{k_C}{k_{tC^O}} = \frac{\ln\left(\frac{[CTP]_0}{[CTP]_0 - [C]}\right)}{\ln\left(\frac{[tC^O TP]_0}{[tC^O TP]_0 - [tC^O]}\right)} \quad (S7)$$

[171] Using the Lambert-Beer law, absorption is related to nucleobase concentration according to equations S8 and S9. The following molar absorptivities (unit: $M^{-1}cm^{-1}$) were adopted:

$$\varepsilon_{260}^{tC^O} = 12200, \varepsilon_{260}^C = 7400, \varepsilon_{260}^G = 11800, \varepsilon_{260}^U = 9300, \varepsilon_{260}^A = 15300, \varepsilon_{369}^{tC^O} = 9370.$$

$$A_{260} = 0.9 \times l \times (\varepsilon_{260}^{tC^O} \times [tC^O] + \varepsilon_{260}^C \times [C] + \varepsilon_{260}^G \times [G] + \varepsilon_{260}^U \times [U] + \varepsilon_{260}^A \times [A]) \quad (S8)$$

$$A_{369} = l \times \varepsilon_{369}^{tC^O} \times [tC^O] \quad (S9)$$

[172] Assuming that the product RNA is uniform in size (1247 nucleotides), its base composition (A: 408, U: 272, G: 307, C: 260) allows for equations S10 - S13.

$$[tC^O] + [C] = [RNA] \times 260 \quad (S10)$$

$$[A] = [RNA] \times 408 \quad (S11)$$

$$[U] = [RNA] \times 272 \quad (S12)$$

$$[G] = [RNA] \times 307 \quad (S13)$$

[173] Solving the equation system composed of S7 through S13 allows for quantification of k_C/k_{tC^O} , $[tC^O]$, $[RNA]$, $[C]$, $[A]$, $[U]$, and $[G]$. The average-strand tC^O incorporation degree (θ_{tC^O}) can then be calculated according to equation S14.

$$\theta_{tC^O} = \frac{[tC^O]}{[C] + [tC^O]} \quad (S14)$$

[174] Using the volume of the cell-free reaction (50 μL) and resulting product solution (100 μL), equation S15 was applied to calculate the tC^O incorporation yield (η_{tC^O}).

$$\eta_{tC^O} = \frac{[tC^O] \times 100 \mu L}{[tC^O TP]_0 \times 50 \mu L} \quad (S15)$$

[175] Consequently, the RNA yield (η_{RNA}) was calculated according to equation S16.

$$\eta_{RNA} = \frac{[RNA] \times 100 \mu L}{[tC^O TP]_0 \times 50 \mu L} \quad (S16)$$

Conclusions

[176] This specification demonstrates that an artificial, size-expanded analogue of cytosine takes the role of natural cytosine and is correctly recognized by several enzymatic machineries, including


the ribosome. This fluorescent base analogue, tC^O , is demonstrated to be a suitable intrinsic imaging label of different size RNAs which minimally perturbs native properties and is compatible with enzymatic labelling processes.

[177] Modified transcripts are non-toxic and translationally active both in bacterial lysate and in
5 eukaryotic systems, regardless of their degree of tC^O incorporation. This conveniently allows for simultaneous monitoring of mRNA uptake and translation into H2B:GFP in live-cell confocal microscopy using selective excitation, an approach that should be applicable to the translation of any protein similarly tagged with a GFP family protein.


[178] The intrinsic fluorescence RNA-labelling methodologies disclosed herein are therefore
10 excellent non-invasive ways to, in real time, elucidate cellular trafficking mechanisms such as endosomal escape or exosomes formation, both of which are of fundamental importance for pharmaceutical applications. As such the technology for live cell imaging should enable new and improved delivery strategies for next-generation nucleic acid-based drugs.

Claims

1. A compound of formula (I) or a salt thereof:




2. A compound of formula (I) as claimed in claim 1.
3. A compound of formula (I) as claimed in claim 1 which is a sodium, potassium, or ammonium salt.
4. A compound of formula (I) as claimed in claim 3 which is a monosodium, disodium, trisodium, monoammonium, diammonium or triammonium salt.
5. A process for preparing a compound of formula (I) or a salt thereof as claimed in any of the preceding claims comprising:
- i. Providing a compound of formula (II) or a salt thereof:



Where PG¹ is a suitable protecting group;

- ii. Immobilising the compound of formula (II) or a salt thereof by linking one of its secondary alcohol groups to a suitable support;
- iii. Capping any free secondary alcohol groups with a suitable protecting group PG²;
- iv. Removing the protecting group PG¹;
- v. Reacting the exposed primary alcohol group with a compound of formula (III):



Where R¹ is selected from a hydro group and a C₁₋₃alkyl group;

- vi. Oxidising the resultant phosphorus (III) compound to a phosphorus (V) compound;
 - vii. Reacting the phosphorus (V) compound with a tetraalkylammonium pyrophosphate to generate a triphosphate;
 - viii. Removing the protecting group PG²; and
 - ix. Cleaving the resultant triphosphate from the support to generate a compound of formula (I) or salt thereof.
6. A process for preparing a compound of formula (I) or a salt thereof as claimed in claim 5, where R¹ is a methyl group.
 7. A process for preparing a compound of formula (I) or a salt thereof as claimed in claim 5 or claim 6, where the support is a solid polymer.
 8. A process for preparing a compound of formula (I) or a salt thereof as claimed in claim 7, where the support is selected from controlled-porosity glass and polystyrene.
 9. A process for preparing a compound of formula (I) or a salt thereof as claimed in claim 8, where the support is controlled-porosity glass.

10. A process for preparing a compound of formula (I) or a salt thereof as claimed in any of claims 5 to 9, where PG¹ is selected from trityl, dimethoxytrityl and trimethoxytrityl.
11. A process for preparing a compound of formula (I) or a salt thereof as claimed in any of claims 5 to 9, where PG² is selected from acetyl, benzoyl, 2,2,2-trichloroethylcarbonyl, paramethoxybenzyl, methyl, tetrahydropyranyl, triethylsilyl, triisopropylsilyl, trimethylsilyl, *tert*-butyldimethylsilyl and methoxyethyl.
12. A process for preparing a compound of formula (I) or a salt thereof as claimed in any of claims 5 to 9, where PG¹ is dimethoxytrityl and PG² is acetyl.
13. A process for preparing a compound of formula (I) or a salt thereof as claimed in any of claims 5 to 12, where immobilisation of the compound of formula (II) in step i) occurs mainly at the 2'-hydroxy position.
14. A process for preparing a compound of formula (I) or a salt thereof as claimed in any of claims 5 to 13, where the tetraalkylammonium pyrophosphate is tetrabutylammonium pyrophosphate.
15. A process for preparing a compound of formula (I) or a salt thereof as claimed in any of claims 5 to 14, where the phosphorus (III) compound in step vi) is oxidised to a phosphorus (V) compound using aqueous pyridine and iodine.
16. A composition for preparing a tC⁰ labelled RNA molecule comprising a compound of formula (I) as claimed in any of claims 1 to 4 and a natural ribonucleotide triphosphate.
17. The use of a compound of formula (I) or a salt thereof as claimed in any of claims 1 to 4 to enzymatically prepare a tC⁰ labelled RNA molecule.
18. The use of a compound of formula (I) or a salt thereof as claimed in claim 17 where the RNA molecule is mRNA.

19. A process for preparing a tC^O labelled RNA molecule comprising providing a DNA template to composition comprising a compound of formula (I) and a natural ribonucleotide triphosphate, then treating the resultant mixture with an RNA polymerase.
20. The use of a tC^O labelled mRNA molecule to prepare a protein encoded by the mRNA by translation.
21. The use of a tC^O labelled mRNA molecule as claimed in claim 20, where the encoded protein is fused to a fluorescent protein.
22. The use of a tC^O labelled mRNA molecule as claimed in claim 21, where the tC^O labelled mRNA and the encoded protein are simultaneously analysed spatiotemporally using confocal microscopy.

Figure 1




Figure 2




Figure 3a




Figure 3b




Figure 3c




Figure 4a




Figure 4b





Figure 5a




Figure 5b




Figure 5c




Figure 5d




Figure 6a




Figure 6b




Figure 6c




Figure 7a




Figure 7b




Figure 7c




Figure 7d




Figure 7e





Figure 7f




Figure 8a




Figure 8b




Figure 8c




Figure 8d




Figure 8e




Figure 8f




Figure 8g





Figure 8h



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2021/067342

A. CLASSIFICATION OF SUBJECT MATTER INV. C07H19/24 C07H1/00 C07H21/02 C12N15/10 C12Q1/68 G01N33/58 ADD. According to International Patent Classification (IPC) or to both national classification and IPC														
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07H C40B C12N C12Q G01N Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data, WPI Data														
C. DOCUMENTS CONSIDERED TO BE RELEVANT <table border="1"> <thead> <tr> <th>Category*</th> <th>Citation of document, with indication, where appropriate, of the relevant passages</th> <th>Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td>X</td> <td>WO 2011/034895 A1 (UNIV COLORADO [US]; KUCHTA ROBERT D [US] ET AL.) 24 March 2011 (2011-03-24) cited in the application</td> <td>1-22</td> </tr> <tr> <td>Y</td> <td>paragraph [00023] claims 1-17</td> <td>1-22</td> </tr> <tr> <td>Y</td> <td>----- WO 2018/089582 A1 (SAN DIEGO STATE UNIV RESEARCH FOUNDATION [US] ET AL.) 17 May 2018 (2018-05-17) claim 1 -----</td> <td>1-22</td> </tr> </tbody> </table>			Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	X	WO 2011/034895 A1 (UNIV COLORADO [US]; KUCHTA ROBERT D [US] ET AL.) 24 March 2011 (2011-03-24) cited in the application	1-22	Y	paragraph [00023] claims 1-17	1-22	Y	----- WO 2018/089582 A1 (SAN DIEGO STATE UNIV RESEARCH FOUNDATION [US] ET AL.) 17 May 2018 (2018-05-17) claim 1 -----	1-22
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
X	WO 2011/034895 A1 (UNIV COLORADO [US]; KUCHTA ROBERT D [US] ET AL.) 24 March 2011 (2011-03-24) cited in the application	1-22												
Y	paragraph [00023] claims 1-17	1-22												
Y	----- WO 2018/089582 A1 (SAN DIEGO STATE UNIV RESEARCH FOUNDATION [US] ET AL.) 17 May 2018 (2018-05-17) claim 1 -----	1-22												
☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.														
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family														
Date of the actual completion of the international search 5 October 2021		Date of mailing of the international search report 15/10/2021												
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Mezzato, Stefano												

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP2021/067342

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. ☒ forming part of the international application as filed:
 - ☒ in the form of an Annex C/ST.25 text file.
 - ☐ on paper or in the form of an image file.
 - b. ☐ furnished together with the international application under PCT Rule 13~~ter~~.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
 - c. ☐ furnished subsequent to the international filing date for the purposes of international search only:
 - ☐ in the form of an Annex C/ST.25 text file (Rule 13~~ter~~.1(a)).
 - ☐ on paper or in the form of an image file (Rule 13~~ter~~.1(b) and Administrative Instructions, Section 713).
2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2021/067342

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2011034895	A1	24-03-2011	NONE

WO 2018089582	A1	17-05-2018	US 2020262862 A1 20-08-2020
			WO 2018089582 A1 17-05-2018
