



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2473144 C 2013/05/28

(11)(21) **2 473 144**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2003/02/03
(87) Date publication PCT/PCT Publication Date: 2003/08/14
(45) Date de délivrance/Issue Date: 2013/05/28
(85) Entrée phase nationale/National Entry: 2004/07/12
(86) N° demande PCT/PCT Application No.: US 2003/003097
(87) N° publication PCT/PCT Publication No.: 2003/066662
(30) Priorité/Priority: 2002/02/05 (US60/354,579)

(51) Cl.Int./Int.Cl. *C07K 1/22* (2006.01),
C07K 16/06 (2006.01)
(72) Inventeurs/Inventors:
FAHRNER, ROBERT L., US;
GORRELL, JEFFREY R., US;
LAZZARESCHI, KATHLYN PHAM, US;
LESTER, PHILIP M., US;
PENG, DAVID, US;
BREECE, TIMOTHY N., US
(73) Propriétaire/Owner:
GENENTECH, INC., US
(74) Agent: SMART & BIGGAR

(54) Titre : PURIFICATION DE PROTEINES
(54) Title: PROTEIN PURIFICATION

(57) **Abrégé/Abstract:**

A method for purifying proteins by Protein A chromatography is described which comprises removing contaminants by washing the solid phase with various intermediate wash buffers.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
14 August 2003 (14.08.2003)

PCT

(10) International Publication Number
WO 2003/066662 A3

(51) International Patent Classification⁷: **C07K 1/22**

(74) Agent: **LEE, Wendy, M.**; Genentech, Inc., MS 49, 1 DNA Way, South San Francisco, CA 94080-4990 (US).

(21) International Application Number:
PCT/US2003/003097

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 3 February 2003 (03.02.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/354,579 5 February 2002 (05.02.2002) US

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: **GENENTECH, INC.** [US/US]; 1 DNA Way, South San Francisco, CA 94080 (US).

(72) Inventors: **FAHRNER, Robert, L.**; 3816 Pasadena Drive, San Mateo, CA 94403 (US). **GORRELL, Jeffrey, R.**; 2351 Bennington Drive, San Bruno, CA 94066 (US). **LAZZARESCHI, Kathlyn, Pham**; 304 Catalpa Avenue, San Mateo, CA 94401 (US). **LESTER, Philip, M.**; 15766 Via Colusa, San Lorenzo, CA 94580 (US). **PENG, David**; 55 Parkside Avenue, Daly City, CA 94015 (US). **BREECE, Timothy, N.**; 3330 Scott Street, San Francisco, CA 94123 (US).

Published:

— with international search report

(88) Date of publication of the international search report:
31 December 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROTEIN PURIFICATION

(57) Abstract: A method for purifying proteins by Protein A chromatography is described which comprises removing contaminants by washing the solid phase with various intermediate wash buffers.



WO 2003/066662 A3

5

PROTEIN PURIFICATION**BACKGROUND OF THE INVENTION****Field of the Invention**

10 This invention relates generally to protein purification. In particular, the invention relates to a method for purifying C_H2/C_H3 region-containing proteins, such as antibodies and immunoadhesins, by Protein A affinity chromatography.

Description of Related Art

15 The large-scale, economic purification of proteins is increasingly an important problem for the biotechnology industry. Generally, proteins are produced by cell culture, using either mammalian or bacterial cell lines engineered to produce the protein of interest by insertion of a recombinant plasmid containing the gene for that protein. Since the cell lines used are living organisms, they must be fed with a complex growth medium, containing sugars, amino acids, and growth factors, usually supplied from preparations of animal serum. Separation of the
20 desired protein from the mixture of compounds fed to the cells and from the by-products of the cells themselves to a purity sufficient for use as a human therapeutic poses a formidable challenge.

Procedures for purification of proteins from cell debris initially depend on the site of expression of the protein. Some proteins can be caused to be secreted directly from the cell
25 into the surrounding growth media; others are made intracellularly. For the latter proteins, the first step of a purification process involves lysis of the cell, which can be done by a variety of methods, including mechanical shear, osmotic shock, or enzymatic treatments. Such disruption releases the entire contents of the cell into the homogenate, and in addition produces subcellular fragments that are difficult to remove due to their small size. These are generally
30 removed by differential centrifugation or by filtration. The same problem arises, although on a smaller scale, with directly secreted proteins due to the natural death of cells and release of intracellular host cell proteins in the course of the protein production run.

Once a clarified solution containing the protein of interest has been obtained, its separation from the other proteins produced by the cell is usually attempted using a combination
35 of different chromatography techniques. These techniques separate mixtures of proteins on the basis of their charge, degree of hydrophobicity, or size. Several different chromatography resins are available for each of these techniques, allowing accurate tailoring of the purification scheme to the particular protein involved. The essence of each of these separation methods

is that proteins can be caused either to move at different rates down a long column, achieving a physical separation that increases as they pass further down the column, or to adhere selectively to the separation medium, being then differentially eluted by different solvents. In some cases, the desired protein is separated from impurities when the impurities specifically
5 adhere to the column, and the protein of interest does not, that is, the protein of interest is present in the "flow-through."

Affinity chromatography, which exploits a specific interaction between the protein to be purified and an immobilized capture agent, may also be an option for some proteins. Protein A is a useful adsorbent for affinity chromatography of proteins, such as antibodies, which contain
10 an Fc region. Protein A is a 41kD cell wall protein from *Staphylococcus aureus* which binds with a high affinity (about 10^{-8} M to human IgG) to the Fc region of antibodies.

Proteins may be purified using controlled pore glass (Sulkowski, E. *Protein Purification: Micro to Macro*, pgs 177-195 (1987); Chadha *et al. Preparative Biochemistry* 11(4):467-482 (1981)) or underivatized silica (Reifsnnyder *et al. J. Chromatography* 753:73-80 (1996)).

US Patent Nos. 6,127,526 and 6,333,398 (Blank, G.) describe an intermediate wash
15 step during Protein A chromatography using hydrophobic electrolytes, *e.g.*, tetramethylammonium chloride (TMAC) and tetraethylammonium chloride (TEAC), to remove the contaminants, but not the immobilized Protein A or the protein of interest, bound to the Protein A column.

20

SUMMARY OF THE INVENTION

The present invention provides various intermediate wash buffers, other than TMAC or TEAC, for use in Protein A chromatography.

In one embodiment, the invention provides a method for purifying a protein which
25 comprises a C_{H2}/C_{H3} region, from a contaminated solution thereof by Protein A chromatography comprising: (a) adsorbing the protein to Protein A immobilized on a solid phase; (b) removing contaminants by washing the solid phase with a composition comprising detergent and salt; and (c) recovering the protein from the solid phase.

In another embodiment, the invention provides a method for purifying a protein, which
30 comprises a C_{H2}/C_{H3} region, from a contaminated solution thereof by Protein A chromatography comprising: (a) adsorbing the protein to Protein A immobilized on a solid phase; (b) removing contaminants by washing the solid phase with a composition comprising a buffer at a concentration of greater than about 0.8M; and (c) recovering the protein from the solid phase.

35 The invention also pertains, in another embodiment, to a method for purifying a protein, which comprises a C_{H2}/C_{H3} region, from a contaminated solution thereof by Protein A chromatography comprising: (a) adsorbing the protein to Protein A immobilized on a solid

phase; (b) removing contaminants by washing the solid phase with a composition comprising salt and solvent; and (c) recovering the protein from the solid phase.

Moreover, the invention provides a method for purifying a protein, which comprises a C_H2/C_H3 region, from a contaminated solution thereof by Protein A chromatography comprising:

- 5 (a) adsorbing the protein to Protein A immobilized on a solid phase; (b) removing contaminants by washing the solid phase with a composition comprising salt and polymer; and (c) recovering the protein from the solid phase.

10 In preferred embodiments, the protein is an antibody (*e.g.* one which binds HER2, vascular endothelial growth factor (VEGF), IgE, CD20, CD40, CD11a, tissue factor (TF), prostate stem cell antigen (PSCA), interleukin-8 (IL-8), epidermal growth factor receptor (EGFR), HER3, HER4, $\alpha4\beta7$ or $\alpha5\beta3$) or an immunoadhesin (*e.g.* a TNF receptor immunoadhesin).

Brief description of the drawings

15 Figures 1A-B show the light chain amino acid sequence (SEQ ID NO:1) and heavy chain amino acid sequence (SEQ ID NO:2) of Trastuzumab (HERCEPTIN®).

Figure 2 depicts screening of various intermediate wash buffers with regard to the anti-IgE antibody E26. The amount of Chinese Hamster Ovary Protein (CHOP) contamination in the elution pool (ppm) is depicted.

20 Figure 3 shows screening of intermediate wash buffers containing polyethylene glycol (PEG). The antibody is E26.

Figure 4 illustrates screening of urea intermediate washes, where the antibody is E26.

Figure 5 depicts screening of various intermediate wash buffers with regard to the anti-HER2 antibody Trastuzumab. The amount of CHOP in the elution pool (ppm) is depicted.

25 Figure 6 depicts screening of various intermediate wash buffers with regard to Trastuzumab and a humanized anti-CD11a antibody.

Figure 7 shows CHOP in the Protein A pool for Trastuzumab and an intermediate wash buffer including sodium citrate or sodium acetate salt; or E26 and an intermediate wash buffer including sodium sulfate or sodium citrate salt.

30 Figure 8 represents alteration of polysorbate 20 in the intermediate wash buffer for E26.

Figure 9 illustrates the effect of pH on CHOP in the elution pool for Trastuzumab and E26.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTSDefinitions:

When used herein, the term "Protein A" encompasses Protein A recovered from a native source thereof, Protein A produced synthetically (*e.g.* by peptide synthesis or by recombinant techniques), and variants thereof which retain the ability to bind proteins which have a C_H2/C_H3 region. Protein A can be purchased commercially from Repligen, Pharmacia and Fermatech.

The Protein A is immobilized on a solid phase. By "solid phase" is meant a non-aqueous matrix to which the Protein A can adhere. The solid phase of interest herein is generally one which comprises a glass, silica, agarose or polystyrene surface. The solid phase may be a purification column or a discontinuous phase of discrete particles. In preferred embodiments, the solid phase is a controlled pore glass column or a silicic acid column. In certain embodiments, the solid phase is coated with a reagent (such as glycerol) which is intended to prevent nonspecific adherence of contaminants to the solid phase.

The protein of interest herein is one which comprises a C_H2/C_H3 region and therefore is amenable to purification by Protein A chromatography. The term "C_H2/C_H3 region" when used herein refers to those amino acid residues in the Fc region of an immunoglobulin molecule which interact with Protein A. In preferred embodiments, the C_H2/C_H3 region comprises an intact C_H2 region followed by an intact C_H3 region, and most preferably comprises a Fc region of an immunoglobulin. Examples of C_H2/C_H3 region-containing proteins include antibodies, immunoadhesins and fusion proteins comprising a protein of interest fused to, or conjugated with, a C_H2/C_H3 region.

The "intermediate wash step" is a step performed after the protein of interest is loaded on the solid phase and adsorbed to the Protein A, but before the protein is recovered from the column. The intermediate wash step serves to remove contaminants nonspecifically bound to the solid phase, antibody and/or Protein A, without significantly eluting the protein of interest or Protein A from the solid phase. In the intermediate wash step, the solid phase is washed with the desired "intermediate wash buffer" (*e.g.* the intermediate wash buffer is passed through the Protein A column, where the solid phase is a column).

A "buffer" is a buffered solution that resists changes in pH by the action of its acid-base conjugate components.

An "equilibration buffer" herein is that used to prepare the solid phase (with immobilized Protein A) for loading the protein of interest. The equilibration buffer is preferably isotonic and commonly has a pH in the range from about 6 to about 8. The equilibration buffer of the example was 25mM Tris, 25mM NaCl, 5mM EDTA, pH 7.1.

The "loading buffer" is that which is used to load the mixture of the C_H2/C_H3 region-containing protein and contaminant(s) onto the solid phase to which the Protein A is immobilized. Often, the equilibration and loading buffers are the same.

The "elution buffer" is used to elute the C_H2/C_H3 region-containing protein from the immobilized Protein A. Preferably the elution buffer has a low pH and thereby disrupts interactions between Protein A and the protein of interest. Preferably, the low pH elution buffer has a pH in the range from about 2 to about 5, most preferably in the range from about 3 to about 4. Examples of buffers that will control the pH within this range include phosphate, acetate, citrate and ammonium buffers, as well as combinations of these. The preferred such buffers are citrate and acetate buffers, most preferably sodium citrate or sodium acetate buffers. Other elution buffers are contemplated including high pH buffers (e.g. those having a pH of 9 or more) or buffers comprising a compound or composition such as MgCl₂ (2mM) for eluting the protein of interest.

The "intermediate wash buffer" is the buffer used to remove contaminant(s), such as CHOP, from the immobilized Protein A without removing significant amounts of the protein of interest bound to the Protein A. The intermediate wash buffer preferably comprises (a) salt and detergent (e.g. polysorbate); (b) salt and solvent (e.g. hexylene glycol); (c) high concentration salt (e.g. high molarity Tris buffer); or (d) salt and polymer (e.g. PEG).

A "salt" is a compound formed by the interaction of an acid and a base. The preferred salt herein is acetate (e.g. sodium acetate), citrate (e.g. sodium citrate), chloride (e.g. sodium chloride), sulphate (e.g. sodium sulphate), or a potassium salt.

As used herein, "solvent" refers to a liquid substance capable of dissolving or dispersing one or more other substances to provide a solution. The preferred solvent is an organic, non-polar solvent, such as ethanol, methanol, isopropanol, acetonitrile, hexylene glycol, propylene glycol, and 2,2-thiodiglycol.

The term "detergent" refers to nonionic surfactants such as polysorbates (e.g. polysorbates 20 or 80); poloxamers (e.g. poloxamer 188); TritonTM; sodium dodecyl sulfate (SDS); sodium laurel sulfate; sodium octyl glycoside; lauryl-, myristyl-, linoleyl-, or stearyl-sulfobetaine; lauryl-, myristyl-, linoleyl- or stearyl-sarcosine; linoleyl-, myristyl-, or cetyl-betaine; lauroamidopropyl-, cocamidopropyl-, linoleamidopropyl-, myristamidopropyl-, palmidopropyl-, or isostearamidopropyl-betaine (e.g. lauroamidopropyl); myristamidopropyl-, palmidopropyl-, or isostearamidopropyl-dimethylamine; sodium methyl cocoyl-, or disodium methyl oleyl-taurate; and the MONAQUATTM series (Mona Industries, Inc., Paterson, New Jersey). The preferred detergent is a polysorbate, such as polysorbate 20 (TWEEN 20®) or polysorbate 80 (TWEEN 80®).

A "polymer" herein is a molecule formed by covalent linkage of two or more

monomers, where the monomers are not amino acid residues. Examples of polymers include polyethyl glycol, polypropyl glycol, and copolymers of ethylene and propylene glycol (e.g. Pluronic, PF68 etc). The preferred polymer is polyethylene glycol (PEG), e.g. PEG 400 and PEG 8000.

5 The term "antibody" is used in the broadest sense and specifically covers monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments so long as they retain, or are modified to comprise, a C_H2/C_H3 region as herein defined.

10 "Antibody fragments" comprise a portion of a full length antibody, generally the antigen binding or variable region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; single-chain antibody molecules; diabodies; linear antibodies; and multispecific antibodies formed from antibody fragments.

15 The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant
20 on the antigen. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, *Nature* 256:495 (1975), or may be
25 made by recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature* 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.* 222:581-597 (1991), for example.

30 The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such
35 antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison *et al.*, *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984)).

The term "hypervariable region" when used herein refers to the amino acid residues of an antibody which are responsible for antigen-binding. The hypervariable region comprises amino acid residues from a "complementarity determining region" or "CDR" (*i.e.* residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues from a "hypervariable loop" (*i.e.* residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). "Framework" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined.

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric antibodies which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which hypervariable region residues of the recipient are replaced by hypervariable region residues from a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues which are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992).

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the "binding domain" of a heterologous "adhesin" protein (*e.g.* a receptor, ligand or enzyme) with the effector functions of an immunoglobulin constant domain. Structurally, the immunoadhesins comprise a fusion of the adhesin amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site (antigen combining site) of an antibody (*i.e.* is "heterologous") and an immunoglobulin constant domain sequence. The immunoglobulin constant domain sequence in the immunoadhesin is preferably derived from $\gamma 1$, $\gamma 2$, or $\gamma 4$ heavy chains since immunoadhesins comprising

these regions can be purified by Protein A chromatography (Lindmark *et al.*, *J. Immunol. Meth.* 62:1-13 (1983)).

The term "ligand binding domain" as used herein refers to any native cell-surface receptor or any region or derivative thereof retaining at least a qualitative ligand binding of a corresponding native receptor. In a specific embodiment, the receptor is from a cell-surface polypeptide having an extracellular domain which is homologous to a member of the immunoglobulin supergene family. Other receptors, which are not members of the immunoglobulin supergene family but are nonetheless specifically covered by this definition, are receptors for cytokines, and in particular receptors with tyrosine kinase activity (receptor tyrosine kinases), members of the hematopoietin and nerve growth factor receptor superfamilies, and cell adhesion molecules, *e.g.* (E-, L- and P-) selectins.

The term "receptor binding domain" is used to designate any native ligand for a receptor, including cell adhesion molecules, or any region or derivative of such native ligand retaining at least a qualitative receptor binding ability of a corresponding native ligand. This definition, among others, specifically includes binding sequences from ligands for the above-mentioned receptors.

An "antibody-immunoadhesin chimera" comprises a molecule which combines at least one binding domain of an antibody (as herein defined) with at least one immunoadhesin (as defined in this application). Exemplary antibody-immunoadhesin chimeras are the bispecific CD4-IgG chimeras described in Berg *et al.*, *PNAS (USA)* 88:4723-4727 (1991) and Chamow *et al.*, *J. Immunol.* 153:4268 (1994).

"Trastuzumab" or "HERCEPTIN®" is a humanized anti-HER2 antibody comprising the light chain amino acid sequence of SEQ ID NO:1 and the heavy chain amino acid sequence of SEQ ID NO:2 or amino acid sequence variants thereof which retain the ability to bind HER2 and inhibit growth of tumor cells which overexpress HER2 (see US Patent 5,677,171).

Modes for Carrying Out the Invention

The process herein involves purifying a C_H2/C_H3 region-containing protein from contaminants by Protein A chromatography. In preferred embodiments, the protein to be purified using Protein A chromatography is an antibody, an immunoadhesin or a protein fused to, or conjugated with, a C_H2/C_H3 region. Techniques for generating such molecules will be discussed below.

1. Antibodies

The preferred protein to be purified according to the present invention is an antibody. Antibodies within the scope of the present invention include, but are not limited to: anti-HER2 antibodies including Trastuzumab (HERCEPTIN®) (Carter *et al.*, *Proc. Natl. Acad.*

Sci. USA, 89:4285-4289 (1992), U.S. Patent No. 5,725,856); anti-CD20 antibodies such as chimeric anti-CD20 "C2B8" as in US Patent No. 5,736,137 (RITUXAN®), a chimeric or humanized variant of the 2H7 antibody as in US Patent No. 5,721,108, B1, or Tositumomab (BEXXAR®); anti-IL-8 (St John *et al.*, *Chest*, 103:932 (1993), and International Publication No. WO 95/23865); anti-VEGF antibodies including humanized and/or affinity matured anti-VEGF antibodies such as the humanized anti-VEGF antibody huA4.6.1 AVASTIN® (Kim *et al.*, *Growth Factors*, 7:53-64 (1992), International Publication No. WO 96/30046, and WO 98/45331, published October 15, 1998); anti-PSCA antibodies (WO01/40309); anti-CD40 antibodies, including S2C6 and humanized variants thereof (WO00/75348); anti-CD11a (US Patent No. 5,622,700, WO 98/23761, Steppe *et al.*, *Transplant Intl.* 4:3-7 (1991), and Hourmant *et al.*, *Transplantation* 58:377-380 (1994)); anti-IgE (Presta *et al.*, *J. Immunol.* 151:2623-2632 (1993), and International Publication No. WO 95/19181); anti-CD18 (US Patent No. 5,622,700, issued April 22, 1997, or as in WO 97/26912, published July 31, 1997); anti-IgE (including E25, E26 and E27; US Patent No. 5,714,338, issued February 3, 1998 or US Patent No. 5,091,313, issued February 25, 1992, WO 93/04173 published March 4, 1993, or International Publication No. WO/1999/001556, US Patent No. 5,714,338); anti-Apo-2 receptor antibody (WO 98/51793 published November 19, 1998); anti-TNF- α antibodies including cA2 (REMICADE®), CDP571 and MAK-195 (See, US Patent No. 5,672,347 issued September 30, 1997, Lorenz *et al.* *J. Immunol.* 156(4):1646-1653 (1996), and Dhainaut *et al.* *Crit. Care Med.* 23(9):1461-1469 (1995)); anti-Tissue Factor (TF) (European Patent No. 0 420 937 B1 granted November 9, 1994); anti-human $\alpha_4\beta_7$ integrin (WO 98/06248 published February 19, 1998); anti-EGFR (chimerized or humanized 225 antibody as in WO 96/40210 published December 19, 1996); anti-CD3 antibodies such as OKT3 (US Patent No. 4,515,893 issued May 7, 1985); anti-CD25 or anti-tac antibodies such as CHI-621 (SIMULECT®) and (ZENAPAX®) (See US Patent No. 5,693,762 issued December 2, 1997); anti-CD4 antibodies such as the cM-7412 antibody (Choy *et al.* *Arthritis Rheum* 39(1):52-56 (1996)); anti-CD52 antibodies such as CAMPATH-1H (Riechmann *et al.* *Nature* 332:323-337 (1988)); anti-Fc receptor antibodies such as the M22 antibody directed against Fc γ RI as in Graziano *et al.* *J. Immunol.* 155(10):4996-5002 (1995); anti-carcinoembryonic antigen (CEA) antibodies such as hMN-14 (Sharkey *et al.* *Cancer Res.* 55(23Suppl): 5935s-5945s (1995); antibodies directed against breast epithelial cells including huBrE-3, hu-Mc 3 and CHL6 (Ceriani *et al.* *Cancer Res.* 55(23): 5852s-5856s (1995); and Richman *et al.* *Cancer Res.* 55(23 Supp): 5916s-5920s (1995)); antibodies that bind to colon carcinoma cells such as C242 (Litton *et al.* *Eur J. Immunol.* 26(1):1-9 (1996)); anti-CD38 antibodies, *e.g.* AT 13/5 (Ellis *et al.* *J. Immunol.* 155(2):925-937 (1995)); anti-CD33 antibodies such as Hu M195 (Jurcic *et al.* *Cancer Res* 55(23 Suppl):5908s-5910s (1995) and CMA-676 or CDP771; anti-CD22 antibodies such as LL2 or LymphoCide™ (Juweid

et al. Cancer Res 55(23 Suppl):5899s-5907s (1995)); anti-EpCAM antibodies such as 1/-1A (PANOREX®); anti-GpIIb/IIIa antibodies such as abciximab or c7E3 Fab (REOPRO®); anti-RSV antibodies such as MEDI-493 (SYNAGIS®); anti-CMV antibodies such as PROTOVIR®; anti-HIV antibodies such as PRO542; anti-hepatitis antibodies such as the
 5 anti-Hep B antibody OSTAVIR®; anti-CA 125 antibody OvaRex™; anti-idiotypic GD3 epitope antibody BEC2; anti- $\alpha v \beta 3$ antibody VITAXIN®; anti-human renal cell carcinoma antibody such as ch-G250; ING-1; anti-human 17-1A antibody (3622W94); anti-human colorectal tumor antibody (A33); anti-human melanoma antibody R24 directed against GD3 ganglioside; anti-human squamous-cell carcinoma (SF-25); and anti-human leukocyte
 10 antigen (HLA) antibodies such as Smart™ ID10 and the anti-HLA DR antibody Oncolym™ (Lym-1).

The preferred target antigens for the antibody herein are: HER2 receptor, VEGF, IgE, CD20, CD11a, and CD40.

Aside from the antibodies specifically identified above, the skilled practitioner could generate antibodies directed against an antigen of interest, *e.g.*, using the techniques
 15 described below.

(i) *Antigen selection and preparation*

The antibody herein is directed against an antigen of interest. Preferably, the antigen is a biologically important polypeptide and administration of the antibody to a mammal suffering from a disease or disorder can result in a therapeutic benefit in that
 20 mammal. However, antibodies directed against nonpolypeptide antigens (such as tumor-associated glycolipid antigens; see US Patent 5,091,178) are also contemplated. Where the antigen is a polypeptide, it may be a transmembrane molecule (*e.g.* receptor) or ligand such as a growth factor. Exemplary antigens include those proteins described in section (3) below. Exemplary molecular targets for antibodies encompassed by the present invention
 25 include CD proteins such as CD3, CD4, CD8, CD19, CD20, CD22 and CD34; members of the ErbB receptor family such as the EGF receptor, HER2, HER3 or HER4 receptor; cell adhesion molecules such as LFA-1, Mac1, p150,95, VLA-4, ICAM-1, VCAM and $\alpha v \beta 3$ integrin including either α or β subunits thereof (*e.g.* anti-CD11a, anti-CD18 or anti-CD11b antibodies); growth factors such as VEGF; IgE; blood group antigens; flk2/flt3 receptor;
 30 obesity (OB) receptor; *mpl* receptor; CTLA-4; protein C, or any of the other antigens mentioned herein.

Soluble antigens or fragments thereof, optionally conjugated to other molecules, can be used as immunogens for generating antibodies. For transmembrane molecules, such as
 35 receptors, fragments of these (*e.g.* the extracellular domain of a receptor) can be used as the immunogen. Alternatively, cells expressing the transmembrane molecule can be used as the immunogen. Such cells can be derived from a natural source (*e.g.* cancer cell lines) or may be cells which have been transformed by recombinant techniques to express the

transmembrane molecule.

Other antigens and forms thereof useful for preparing antibodies will be apparent to those in the art.

(ii) *Polyclonal antibodies*

5 Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the antigen to a protein that is immunogenic in the species to be immunized, *e.g.*, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl
10 sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl groups.

Animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, *e.g.*, 100 μg or 5 μg of the protein or conjugate (for rabbits or mice,
15 respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of antigen or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Preferably, the
20 animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitably used to enhance the immune response.

(iii) *Monoclonal antibodies*

25 Monoclonal antibodies may be made using the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster or macaque monkey, is immunized as hereinabove described to elicit lymphocytes
30 that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)).

35 The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the

enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, Protein A-SepharoseTM, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography. Preferably the Protein A chromatography procedure described herein is used.

DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the monoclonal antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells.

The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, *et al.*, *Proc. Natl Acad. Sci. USA*, 81:6851 (1984)), or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody, or they are substituted for the variable domains of one antigen-combining site of an antibody to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an antigen and another antigen-combining site having specificity for a different antigen.

In a further embodiment, monoclonal antibodies can be isolated from antibody phage libraries generated using the techniques described in McCafferty *et al.*, *Nature*, 348:552-554 (1990). Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks *et al.*, *Bio/Technology*, 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse *et al.*, *Nuc. Acids. Res.*, 21:2265-2266 (1993)). Thus, these techniques are viable alternatives to traditional hybridoma techniques for isolation of monoclonal antibodies.

(iv) *Humanized and human antibodies*

A humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, *Nature*, 321:522-525 (1986); Riechmann *et al.*, *Nature*, 332:323-327 (1988); Verhoeyen *et al.*, *Science*, 239:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The

human sequence which is closest to that of the rodent is then accepted as the human FR for the humanized antibody (Sims *et al.*, *J. Immunol.*, 151:2296 (1993)). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta *et al.*, *J. Immunol.*, 151:2623 (1993)).

It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

Alternatively, it is now possible to produce transgenic animals (*e.g.*, mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, *e.g.*, Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993); Jakobovits *et al.*, *Nature*, 362:255-258 (1993); Bruggermann *et al.*, *Year in Immuno.*, 7:33 (1993); and Duchosal *et al.*, *Nature* 355:258 (1992). Human antibodies can also be derived from phage-display libraries (Hoogenboom *et al.*, *J. Mol. Biol.*, 227:381 (1991); Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991); Vaughan *et al.*, *Nature Biotech* 14:309 (1996)).

(v) Antibody fragments

Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, *e.g.*, Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992) and Brennan *et al.*, *Science*, 229:81 (1985)). However, these fragments can now be

produced directly by recombinant host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, Bio/Technology 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). See WO 93/16185.

(vi) *Multispecific antibodies*

10 Multispecific antibodies have binding specificities for at least two different antigens. While such molecules normally will only bind two antigens (*i.e.* bispecific antibodies, BsAbs), antibodies with additional specificities such as trispecific antibodies are encompassed by this expression when used herein.

15 Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein *et al.*, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure.

20 Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker *et al.*, *EMBO J.*, 10:3655-3659 (1991).

25 According to another approach described in WO96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (*e.g.* tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (*e.g.* alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

35 Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using

any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science*, 229: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby *et al.*, *J. Exp. Med.*, 175: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, *J. Immunol.*, 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et*

al., J. Immunol., 152:5368 (1994). Alternatively, the antibodies can be "linear antibodies" as described in Zapata *et al. Protein Eng.* 8(10):1057-1062 (1995). Briefly, these antibodies comprise a pair of tandem Fd segments (V_H - C_H1 - V_H - C_H1) which form a pair of antigen binding regions. Linear antibodies can be bispecific or monospecific.

5 Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al. J. Immunol.* 147: 60 (1991).

2. Immunoadhesins

10 The simplest and most straightforward immunoadhesin design combines the binding domain(s) of the adhesin (*e.g.* the extracellular domain (ECD) of a receptor) with the hinge and Fc regions of an immunoglobulin heavy chain. Ordinarily, when preparing the immunoadhesins of the present invention, nucleic acid encoding the binding domain of the adhesin will be fused C-terminally to nucleic acid encoding the N-terminus of an immunoglobulin constant domain sequence, however N-terminal fusions are also possible.

15 Typically, in such fusions the encoded chimeric polypeptide will retain at least functionally active hinge, C_H2 and C_H3 domains of the constant region of an immunoglobulin heavy chain. Fusions are also made to the C-terminus of the Fc portion of a constant domain, or immediately N-terminal to the C_H1 of the heavy chain or the corresponding region of the light chain. The precise site at which the fusion is made is not critical; 20 particular sites are well known and may be selected in order to optimize the biological activity, secretion, or binding characteristics of the immunoadhesin.

 In a preferred embodiment, the adhesin sequence is fused to the N-terminus of the Fc domain of immunoglobulin G_1 (IgG₁). It is possible to fuse the entire heavy chain constant region to the adhesin sequence. However, more preferably, a sequence beginning 25 in the hinge region just upstream of the papain cleavage site which defines IgG Fc chemically (*i.e.* residue 216, taking the first residue of heavy chain constant region to be 114), or analogous sites of other immunoglobulins is used in the fusion. In a particularly preferred embodiment, the adhesin amino acid sequence is fused to (a) the hinge region and C_H2 and C_H3 or (b) the C_H1 , hinge, C_H2 and C_H3 domains, of an IgG heavy chain.

30 For bispecific immunoadhesins, the immunoadhesins are assembled as multimers, and particularly as heterodimers or heterotetramers. Generally, these assembled immunoglobulins will have known unit structures. A basic four chain structural unit is the form in which IgG, IgD, and IgE exist. A four chain unit is repeated in the higher molecular weight immunoglobulins; IgM generally exists as a pentamer of four basic units held 35 together by disulfide bonds. IgA globulin, and occasionally IgG globulin, may also exist in multimeric form in serum. In the case of multimer, each of the four units may be the same or different.

Various exemplary assembled immunoadhesins within the scope herein are schematically diagrammed below:

- (a) AC_L-AC_L ;
- (b) $AC_H-(AC_H, AC_L-AC_H, AC_L-V_HC_H, \text{ or } V_LC_L-AC_H)$;
- 5 (c) $AC_L-AC_H-(AC_L-AC_H, AC_L-V_HC_H, V_LC_L-AC_H, \text{ or } V_LC_L-V_HC_H)$
- (d) $AC_L-V_HC_H-(AC_H, \text{ or } AC_L-V_HC_H, \text{ or } V_LC_L-AC_H)$;
- (e) $V_LC_L-AC_H-(AC_L-V_HC_H, \text{ or } V_LC_L-AC_H)$; and
- (f) $(A-Y)_n-(V_LC_L-V_HC_H)_2$,

wherein each A represents identical or different adhesin amino acid sequences;

10 V_L is an immunoglobulin light chain variable domain;

V_H is an immunoglobulin heavy chain variable domain;

C_L is an immunoglobulin light chain constant domain;

C_H is an immunoglobulin heavy chain constant domain;

n is an integer greater than 1;

15 Y designates the residue of a covalent cross-linking agent.

In the interests of brevity, the foregoing structures only show key features; they do not indicate joining (J) or other domains of the immunoglobulins, nor are disulfide bonds shown. However, where such domains are required for binding activity, they shall be constructed to be present in the ordinary locations which they occupy in the immunoglobulin

20 molecules.

Alternatively, the adhesin sequences can be inserted between immunoglobulin heavy chain and light chain sequences, such that an immunoglobulin comprising a chimeric heavy chain is obtained. In this embodiment, the adhesin sequences are fused to the 3' end of an immunoglobulin heavy chain in each arm of an immunoglobulin, either between the hinge

25 and the C_H2 domain, or between the C_H2 and C_H3 domains. Similar constructs have been reported by Hoogenboom, *et al.*, *Mol. Immunol.* 28:1027-1037 (1991).

Although the presence of an immunoglobulin light chain is not required in the immunoadhesins of the present invention, an immunoglobulin light chain might be present either covalently associated to an adhesin-immunoglobulin heavy chain fusion polypeptide,

30 or directly fused to the adhesin. In the former case, DNA encoding an immunoglobulin light chain is typically coexpressed with the DNA encoding the adhesin-immunoglobulin heavy chain fusion protein. Upon secretion, the hybrid heavy chain and the light chain will be covalently associated to provide an immunoglobulin-like structure comprising two disulfide-linked immunoglobulin heavy chain-light chain pairs. Methods suitable for the preparation of

35 such structures are, for example, disclosed in U.S. Patent No. 4,816,567, issued 28 March 1989.

Immunoadhesins are most conveniently constructed by fusing the cDNA sequence encoding the adhesin portion in-frame to an immunoglobulin cDNA sequence. However, fusion to genomic immunoglobulin fragments can also be used (see, *e.g.* Aruffo *et al.*, *Cell* 61:1303-1313 (1990); and Stamenkovic *et al.*, *Cell* 66:1133-1144 (1991)). The latter type of fusion requires the presence of Ig regulatory sequences for expression. cDNAs encoding IgG heavy-chain constant regions can be isolated based on published sequences from cDNA libraries derived from spleen or peripheral blood lymphocytes, by hybridization or by polymerase chain reaction (PCR) techniques. The cDNAs encoding the "adhesin" and the immunoglobulin parts of the immunoadhesin are inserted in tandem into a plasmid vector that directs efficient expression in the chosen host cells.

3. Other C_H2/C_H3 region-containing proteins

In other embodiments, the protein to be purified is one which is fused to, or conjugated with, a C_H2/C_H3 region. Such fusion proteins may be produced so as to increase the serum half-life of the protein and/or to facilitate purification of the protein by Protein A chromatography.

Examples of biologically important proteins which can be conjugated this way include renin; a growth hormone, including human growth hormone and bovine growth hormone; growth hormone releasing factor; parathyroid hormone; thyroid stimulating hormone; lipoproteins; alpha-1-antitrypsin; insulin A-chain; insulin B-chain; proinsulin; follicle stimulating hormone; calcitonin; luteinizing hormone; glucagon; clotting factors such as factor VIIIc, factor IX, tissue factor, and von Willebrands factor; anti-clotting factors such as Protein C; atrial natriuretic factor; lung surfactant; a plasminogen activator, such as urokinase or human urine or tissue-type plasminogen activator (t-PA); bombesin; thrombin; hemopoietic growth factor; tumor necrosis factor-alpha and -beta; enkephalinase; RANTES (regulated on activation normally T-cell expressed and secreted); human macrophage inflammatory protein (MIP-1-alpha); a serum albumin such as human serum albumin; Muellerian-inhibiting substance; relaxin A-chain; relaxin B-chain; prorelaxin; mouse gonadotropin-associated peptide; a microbial protein, such as beta-lactamase; DNase; IgE; a cytotoxic T-lymphocyte associated antigen (CTLA), such as CTLA-4; inhibin; activin; vascular endothelial growth factor (VEGF); receptors for hormones or growth factors; Protein A or D; rheumatoid factors; a neurotrophic factor such as bone-derived neurotrophic factor (BDNF), neurotrophin-3, -4, -5, or -6 (NT-3, NT-4, NT-5, or NT-6), or a nerve growth factor such as NGF-β; platelet-derived growth factor (PDGF); fibroblast growth factor such as aFGF and bFGF; epidermal growth factor (EGF); transforming growth factor (TGF) such as TGF-alpha and TGF-beta, including TGF-β1, TGF-β2, TGF-β3, TGF-β4, or TGF-β5; insulin-like growth factor-I and -II (IGF-I and IGF-II); des(1-3)-IGF-I (brain IGF-I), insulin-like growth

factor binding proteins; CD proteins such as CD3, CD4, CD8, CD19 and CD20; erythropoietin; osteoinductive factors; immunotoxins; a bone morphogenetic protein (BMP); an interferon such as interferon-alpha, -beta, and -gamma; colony stimulating factors (CSFs), *e.g.*, M-CSF, GM-CSF, and G-CSF; interleukins (ILs), *e.g.*, IL-1 to IL-10; superoxide
5 dismutase; T-cell receptors; surface membrane proteins; decay accelerating factor; viral antigen such as, for example, a portion of the AIDS envelope; transport proteins; homing receptors; addressins; regulatory proteins; integrins such as CD11a, CD11b, CD11c, CD18, an ICAM, VLA-4 and VCAM; a tumor associated antigen such as HER2, HER3 or HER4 receptor; and fragments of any of the above-listed polypeptides.

10

4. Protein Purification

The protein to be purified using the method described herein is generally produced using recombinant techniques. Methods for producing recombinant proteins are described, *e.g.*, in US Pat No's 5,534,615 and 4,816,567, specifically incorporated herein by reference.

15

In preferred embodiments, the protein of interest is produced in a CHO cell (see, *e.g.* WO 94/11026). Examples of proteins which can be purified using the process described herein have been described above.

When using recombinant techniques, the protein can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the protein is produced
20 intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, is removed, for example, by centrifugation or ultrafiltration. Where the protein is secreted into the medium, the recombinant host cells may be separated from the cell culture medium by tangential flow filtration, for example.

Protein A immobilized on a solid phase is used to purify the C_H2/C_H3 region-
25 containing protein. The solid phase is preferably a column comprising a glass, silica, agarose or polystyrene surface for immobilizing the Protein A. Preferably, the solid phase is a controlled pore glass column or a silicic acid column. Sometimes, the column has been coated with a reagent, such as glycerol, in an attempt to prevent nonspecific adherence to the column. The PROSEP A™ column, commercially available from Bioprocessing Limited,
30 is an example of a Protein A controlled pore glass column which is coated with glycerol. Other examples of columns contemplated herein include the POROS 50 A™ (polystyrene) column or rProtein A SEPHAROSE FAST FLOW™ (agarose) column.

The solid phase for the Protein A chromatography is equilibrated with a suitable buffer. For example, the equilibration buffer may be 25mM Tris, 25mM NaCl, 5mM EDTA,
35 pH 7.1.

The contaminated preparation derived from the recombinant host cells is loaded on the equilibrated solid phase using a loading buffer which may be the same as the

equilibration buffer. As the contaminated preparation flows through the solid phase, the protein is adsorbed to the immobilized Protein A and other contaminants (such as Chinese Hamster Ovary Proteins, CHOP, where the protein is produced in a CHO cell) may bind nonspecifically to the solid phase.

5 The next step performed sequentially entails removing the contaminants bound to the solid phase, antibody and/or Protein A, by washing the solid phase in an intermediate wash step. After loading, the solid phase may be equilibrated with equilibration buffer before beginning the intermediate wash step.

10 The intermediate wash buffer may comprise salt and a further compound, where the further compound is (a) detergent (preferably polysorbate, *e.g.* polysorbate 20 or polysorbate 80); (b) solvent (preferably hexylene glycol); and (c) polymer (preferably PEG).

15 The salt employed may be selected based on the protein of interest, but preferably is acetate (*e.g.* sodium acetate), especially where the antibody is an anti-HER2 antibody such as Trastuzumab; or citrate (*e.g.* sodium citrate), especially where the antibody is an anti-IgE antibody such as E26.

20 The amounts of the salt and further compound in the composition are such that the combined amount elutes the contaminant(s), without substantially removing the protein of interest. Preferred salt concentrations in such wash buffers are from about 0.1 to about 2M, and more preferably from about 0.2M to about 0.6M. Useful detergent concentrations are from about 0.01 to about 5%, more preferably from about 0.1 to 1%, and most preferably about 0.5%, *e.g.* where the detergent is polysorbate. Exemplary solvent concentrations are from about 1% to 40%, preferably from about 5 to about 25%. For instance, in the examples herein, the preferred concentration of the solvent (hexylene glycol) for E26 was about 20%, whereas for Trastuzumab the preferred concentration of the solvent (again hexylene glycol) was about 10%. Where the further compound is a polymer (*e.g.* PEG 400 or PEG 8000), the concentration thereof may, for example, be from about 1% to about 20%, preferably from about 5% to about 15%.

30 In another embodiment, the intermediate wash step involves the use of a highly concentrated buffer solution, *e.g.* a buffer at a concentration of greater than about 0.8M, *e.g.* up to about 2M, and preferably in the range from about 0.8M to about 1.5M, most preferably about 1M. In this embodiment, the buffer is preferably a Tris buffer, such as Tris acetate.

 The pH of the intermediate wash buffer is preferably from about 4 to about 8, more preferably from about 4.5 to about 5.5, and most preferably about 5.0. In another preferred embodiment, the pH is about 7.0.

35 Following the intermediate wash step of the preceding paragraph, the protein of interest is recovered from the column. This is normally achieved using a suitable elution buffer. The protein may, for example, be eluted from the column using an elution buffer

having a low pH, *e.g.* in the range from about 2 to about 5, and preferably in the range from about 2.5 to about 3.5. Examples of elution buffers for this purpose include citrate or acetate buffers.

The eluted protein preparation may be subjected to additional purification steps either prior to, or after, the Protein A chromatography step. Exemplary further purification steps include hydroxylapatite chromatography; dialysis; affinity chromatography using an antibody to capture the protein; hydrophobic interaction chromatography (HIC); ammonium sulphate precipitation; anion or cation exchange chromatography; ethanol precipitation; reverse phase HPLC; chromatography on silica; chromatofocusing; and gel filtration. In the examples herein, the Protein A chromatography step is followed by downstream cation exchange (SP-Sepharose) and anion exchange (Q-Sepharose) purification steps.

The protein thus recovered may be formulated in a pharmaceutically acceptable carrier and is used for various diagnostic, therapeutic or other uses known for such molecules.

The following examples are offered by way of illustration and not by way of limitation.

EXAMPLE 1

Intermediate Wash Solutions

The anti-HER2 antibody Trastuzumab (Figs. 1A-B; HERCEPTIN®), anti-IgE antibody (E26; US Patent No. 5,994,511, Lowman *et al.*) and humanized anti-CD11a antibody (XANELIM®; US Patent No. 6,037,454) were recombinantly produced in Chinese Hamster Ovary (CHO) cells and purified by Protein A chromatography as a first chromatographic step to remove contaminating CHO proteins (CHOP). However, CHOP tends to binding non-specifically to ProSepA, the resin used for this step. ProSep A has the antibody-binding Protein A immobilized on glycerol-coated controlled-pore glass. Although the glycerol coat reduces non-specific binding, some CHOP still adheres to the resin's glass backbone. During the elution phase of the Protein A operation, any non-specifically bound CHOP will co-elute with the antibody, compromising the purity of the product pool. To remove this CHOP before the elution phase, US Patent Nos. 6,127,526 and 6,333,398 (Blank, G.) exemplify an intermediate wash step using tetramethylammonium chloride (TMAC) to remove CHOP. Although TMAC is effective at removing non-specifically bound CHOP, it is difficult to handle and dispense, is toxic, requires costly disposal as a hazardous waste, and is corrosive at high concentration and low pH. The following study shows that alternative wash compositions, without the drawbacks of TMAC, can be used in an intermediate wash step. The other buffers (equilibration, load, elution and regeneration buffers) were as in the Example in US Patent Nos. 6,127,526 and 6,333,398.

A wide variety of "intermediate wash buffers" were screened with regard to the anti-IgE antibody E26, the anti-HER2 antibody Trastuzumab, and the anti-CD11a antibody XANELIM™. The classes of wash buffers were: (a) detergent and salt; (b) solvent and salt; (c) polymer and salt; (d) high concentration buffer; and (e) urea.

5 The protein yield, CHOP removal, and protein aggregation in the Protein A pools were determined for each of the intermediate wash buffers. For all runs, yield was greater than 94%, except the intermediate wash solution with 20% hexylene glycol where the protein was Trastuzumab. CHOP removal achieved with the various intermediate wash buffers is depicted in Figs. 2-6. Protein aggregation as determined by size exclusion chromatography
10 was less than 1.5% for all runs.

The preferred intermediate wash buffers, taking into consideration CHOP clearance, final protein yield and ease of use, were: (a) polysorbate/salt; (b) hexylene glycol/salt; and (c) high concentration Tris buffer. The polysorbate/salt intermediate wash buffer was selected for further studies.

15

EXAMPLE 2

Alteration of the Intermediate Wash Buffer

The effect of (a) salt type and concentration, (b) polysorbate concentration, and (c) pH of the intermediate wash buffer on CHOP removal was evaluated. The antibodies were the anti-IgE antibody E26, and the anti-HER2 antibody Trastuzumab.

20 Fig. 7 depicts the effect of salt type and concentration on CHOP removal. For E26, CHOP level in the elution pool was not significantly affected by concentration of citrate or sulphate in the wash solution. For Trastuzumab, CHOP level in the elution pool was affected by the concentration of salt in the intermediate wash buffer. Preferred concentrations of the salt were from about 0.3 to about 0.6M.

25 The effect of polysorbate concentration on CHOP level in the elution pool was also evaluated. As shown in Fig. 8, as the concentration of polysorbate in the intermediate wash buffer increased, the amount of CHOP contamination decreased. The preferred concentration of polysorbate is about 0.5% to about 1%.

The effect of pH on CHOP was also assessed and the results of these experiments
30 are shown in Fig. 9. A lower pH resulted in less CHOP contamination in the eluted protein. The preferred pH is about 5.

EXAMPLE 3

Downstream performance

The downstream performance in terms of CHOP removal, yield, etc (SDS-PAGE,
35 HPLC-IEC, size exclusion chromatography (SEC), and protein A leaching), were determined for E26 and Trastuzumab. The downstream purification steps were cation exchange chromatography (SP-Sepharose) and anion exchange chromatography (Q-Sepharose).

For E26, the intermediate wash buffers were: (a) 0.5% polysorbate 20/0.2M sodium citrate, (b) 20% hexylene glycol/0.2M sodium citrate, and (c) 1M Tris acetate. The results are shown in the following Table.

Table 1
E26 Downstream

	CHOP (ppm)				
Step	TMAC	Polys/Salt	Hex Gly/Salt	TrisOAc	Equil
Pro A	320	260	230	310	600
SP-Seph	50	40	30	40	90
Q-Seph	<1	<1	<1	<1	4
Overall % Yield:	74	72	67	73	72

For Trastuzumab, the intermediate wash buffers were: (a) 0.5% polysorbate 20/0.5M sodium acetate, (b) 10% hexylene glycol/0.5M sodium acetate, and (c) 1M Tris acetate. The results are summarized in the Tables below.

Table 2
Trastuzumab Downstream

	CHOP (ppm)				
Step	TMAC	Polys/Salt	Hex Gly/Salt	TrisOAc	Equil
Pro A	390	490	460	560	1540
SP-Seph	45	40	40	50	70
Q-Seph*	<2	<2	<1	<1	6
Overall % yield:	66	66	66	65	69

*Concentrated Q-Seph pool >25 mg/ml

Table 3
Trastuzumab Downstream - Contribution of polysorbate 20 and salt

	CHOP (ppm)				
Step	TMAC	Polys/NaOAc	Polys	NaOAc	Equil
Pro A	449	386	1328	628	2130
SP-Seph	12	13	13	22	24
Q-Seph	<1	<1	<3	4	4
Overall % yield:	84	84	69*	83	84

* Low yield (83%) at Protein A step

The yields and CHOP removal were similar for the two antibodies tested.

Polysorbate, hexylene glycol and Tris showed good CHOP clearance and yield after Protein A chromatography and subsequent ion exchange chromatography steps.

Size Exclusion Chromatography (SEC) for measuring percentage aggregate and

SDS-PAGE were performed on Q-Sepharose pools. Ion Exchange analysis (IEX) for measuring percent main peak, and SEC for evaluating aggregation were performed on Protein A and SP-Sepharose pools. For both Trastuzumab and E26, the assays showed similar results between the positive control (TMAC) and the alternative intermediate wash
5 buffers.

EXAMPLE 4

Column Re-Use

The purpose of this experiment was to determine whether polysorbate 20 in the intermediate wash buffer affects resin lifetime. Buffers (equilibration, polysorbate 20
10 intermediate wash, elution and regeneration) were recycled on a 6.84ml column (0.66 cm x 20cm) over 400 cycles. A HER2 breakthrough curve is performed every 50 cycles to determine resin binding capacity. The current specification for maximum binding capacity is 20 grams of antibody per liter of resin. A drop in binding capacity with increasing cycles would indicate that polysorbate 20 reduces resin lifetime. This experiment demonstrated
15 that polysorbate 20 does not reduce resin lifetime. Actual re-use has been completed for 140 cycles and no change in yield has been seen.

Sequence Listing

<110> Genentech, Inc.

<120> PROTEIN PURIFICATION

<130> 81014-91

<140> PCT/US2003/003097

<141> 2003-02-03

<150> US 60/354,579

<151> 2002-02-05

<160> 2

<210> 1

<211> 214

<212> PRT

<213> Artificial Sequence

<220>

<223> Sequence is synthesized.

<400> 1

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val
1				5				10						15

Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Asp	Val	Asn
				20				25						30

Thr	Ala	Val	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys
				35				40						45

Leu	Leu	Ile	Tyr	Ser	Ala	Ser	Phe	Leu	Tyr	Ser	Gly	Val	Pro	Ser
				50				55						60

Arg	Phe	Ser	Gly	Ser	Arg	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile
				65				70						75

Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln
				80				85						90

His	Tyr	Thr	Thr	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu
				95				100						105

Ile	Lys	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro
				110				115						120

Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu
				125				130						135

Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val
				140				145						150

Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu
				155				160						165

Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr
				170				175						180

Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu
 185 190 195

Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn
 200 205 210

Arg Gly Glu Cys

<210> 2

<211> 449

<212> PRT

<213> Artificial Sequence

<220>

<223> Sequence is synthesized.

<400> 2

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15

Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Asn Ile Lys
 20 25 30

Asp Thr Tyr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 35 40 45

Glu Trp Val Ala Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr
 50 55 60

Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser
 65 70 75

Lys Asn Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 80 85 90

Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr
 95 100 105

Ala Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 110 115 120

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser
 125 130 135

Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys
 140 145 150

Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala
 155 160 165

Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser
 170 175 180

Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
 185 190 195

Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser
 200 205 210

Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	215	220	225
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	230	235	240
Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	245	250	255
Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	260	265	270
His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	275	280	285
Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	290	295	300
Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	305	310	315
Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	320	325	330
Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	335	340	345
Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	350	355	360
Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	365	370	375
Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	380	385	390
Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	395	400	405
Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	410	415	420
Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	425	430	435
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly		440	445	

Claims:

1. A method for purifying a protein, which comprises a C_H2/C_H3 region, from a contaminated solution thereof by Protein A chromatography comprising:
 - (a) adsorbing the protein from said contaminated solution to Protein A immobilized on a solid phase;
 - (b) removing contaminants by washing the solid phase with a composition comprising detergent at a concentration between 0.01% and 5% and salt at a concentration of between 0.1M and 2M, the composition having a pH of pH 4.5 to 5.5; and
 - (c) recovering the protein from the solid phase with an elution buffer having a pH in the range from 2 to 5.
2. The method of claim 1 wherein the solid phase comprises silica, glass, agarose, or polystyrene.
3. The method of claim 1 or 2 wherein the solid phase comprises silica or glass.
4. The method of claim 1, 2, or 3 wherein the protein is an antibody or an immunoadhesin.
5. The method of claim 1, 2, or 3 wherein the protein is an antibody.
6. The method of claim 5 wherein the antibody binds an antigen selected from the group consisting of HER2, vascular endothelial growth factor (VEGF), IgE, CD20, CD40, CD11a, tissue factor (TF), prostate stem cell antigen (PSCA), interleukin-8 (IL-8), epidermal growth factor receptor (EGFR), HER3, HER4, α 4 β 7 and α 5 β 3.
7. The method of claim 5 wherein the antibody is an anti-HER2 antibody.
8. The method of claim 5 wherein the antibody is an anti-IgE antibody.
9. The method of any one of claims 1-8 wherein the detergent is polysorbate.

10. The method of claim 9 wherein the concentration of the polysorbate in the composition is from 0.1 to 1%.

11. The method of any one of claims 1-10 wherein the salt is acetate or citrate.

12. The method of any one of claims 1-11 wherein the concentration of the salt in the composition is from 0.2 to 0.6 M.

LIGHT CHAIN

1 D I Q M T Q S P S S L S A S V G D R V T I T C R A S Q D V N T A V A W Y Q Q K P G K A P K 45

46 L L I Y S A S F L Y S G V P S R F S G S R S G T D F T L T I S S L Q P E D F A T Y Y C Q Q 90

91 H Y T T P P T F G Q G T K V E I K R T V A A P S V F I F P P S D E Q L K S G T A S V V C L 135

136 L N N F Y P R E A K V Q W K V D N A L Q S G N S Q E S V T E Q D S K D S T Y S L S S T L T 180

181 L S K A D Y E K H K V Y A C E V T H Q G L S S P V T K S F N R G E C 214

SEQ ID NO: 1

FIG.-1A

HEAVY CHAIN

153045

467590

91120135

136165180

181210225

226255270

271300315

316345360


361390405

406435449


EVQLVESGGGLVQPGGSLRLSCAASGFNIKDTYIHWVRQAPGKGL
EWVARIYPTNGYTTRYADSVKGRFTISADTSKNTAYLQMNSLRAED
TAVYYCSRWGGDGFYAMDYWGQGTLVTVSSASTKGPSVFPLAPSS
KSTSGGTAAALGCLVKDYFPEPVTVSWNSGALLTSGVHTFPAVLQSS
GLYSSLSSVTVPPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDK
THTCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVS
HEDPEVKFNWYVDGVEVHNAAKTKPREEQYNSTVRVVSVLTVLHQD
WLVNGKEVKCKVSNKALPAPIEKTKAKGQPREPQVYTLPPSREE
MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPPVLDSDG
SFFLYSKLTVDKSRWQQGNVFFSCSVMH⁴EALHNHYTQKSLSPG

FIG.-1B


3 / 9

Screening: E26**FIG. 2**


4 / 9

**FIG. 3**


5 / 9

**FIG. 4**



6 / 9

**FIG._5**


7 / 9

Screening: Anti-HER2 & anti-CD11a/Tris**FIG._6**

8 / 9

**FIG. 7****FIG. 8**

9 / 9

**FIG. 9**