



(51) International Patent Classification:

C07D 419/12 (2006.01) *A61K 31/553* (2006.01)
C07D 413/12 (2006.01) *A61P 35/00* (2006.01)
A61K 31/554 (2006.01)

(21) International Application Number:

PCT/US2012/031245

(22) International Filing Date:

29 March 2012 (29.03.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/469,054 29 March 2011 (29.03.2011) US

(71) Applicants (for all designated States except US): **THE BROAD INSTITUTE, INC.** [US/US]; 7 Cambridge Center, Cambridge, MA 02142 (US). **PRESIDENT & FELLOWS OF HARVARD COLLEGE** [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HUSSAIN, Mahmud** [BD/BD]; 44/6 Maddhay Paikpara, Mirpur, Dhaka, 1216 (BD). **ITO, Daisuke** [JP/JP]; 1-30-20- Kamikashiwada, Ushiku, Ibaraki, 300-1232 (JP). **LAW, Jason** [US/US]; 812 Memorial Dr., Apt 1309, Cambridge, MA 02139 (US). **LEIENDECKER, Matthias** [DE/DE]; Ahornweg 58, 54296 Trier (DE). **LIU, Ke** [CN/US]; 144 Spring Street, Apt. 1, Cambridge, MA 02141 (US). **MUNOZ, Benito** [CA/US]; 377 Lowell Ave., Newtonville, MA 02460 (US). **SCHREIBER, Stuart** [US/US]; 2 Avery Street, Boston, MA 02111 (US). **SHAMJI, Alykhan** [CA/US]; 21 Burn-

side Avenue, Apt. 2, Somerville, MA 02144 (US). **STERN, Andrew** [US/US]; 255 Commonwealth Ave., #5, Boston, MA 02115 (US).

(74) Agents: **ISSAC, Boy, P.** et al.; Elmore Patent Law Group, P.C., 484 Groton Road, Westford, MA 01886 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: COMPOUNDS AND METHODS FOR THE TREATMENT OF ISOCITRATE DEHYDROGNASE RELATED DISEASES



(57) Abstract: The invention relates to compounds of Formula I pharmaceutically acceptable salt, ester or prodrug thereof:

Inventors: Mahmud Hussain, Daisuke Ito, Jason Law, Matthias Leiendecker, Ke Liu, Benito Munoz, Stuart Schreiber, Alykhan Shamji and Andrew Stern

COMPOUNDS AND METHODS FOR THE TREATMENT OF ISOCITRATE DEHYDROGENASE RELATED DISEASES

RELATED APPLICATION

5 This application claims the benefit of U.S. Provisional Application No. 61/469,054, filed on March 29, 2011. The entire teachings of the above application are incorporated herein by reference.

BACKGROUND

Cancer genomics is revealing comprehensively somatic mutations that may
10 constitute the root cause of disease. These findings suggest novel mechanisms for cancer initiation and progression, and new ways we might treat cancer in the future. Recently, a genome-wide sequencing study identified mutations in the metabolic enzyme IDH1 in samples from patients with glioblastoma multiformes, which are among the most lethal cancers with survival of only months after their diagnosis. (Parsons D.W. et al., Science
15 321, 1807-1812, 2008). Subsequent analyses revealed that mutations in IDH1 are common (70%-80%) in grade II, III gliomas and secondary glioblastomas and that patients lacking IDH1 mutations often harbor mutations in IDH2, which shares 70% identity with IDH1. (Yan H. et al., N Engl J Med 360, 765-773, 2009; Bleeker F.E. et al., Hum Mutat 30, 7-11, 2009; Balss J. et al., Acta Neuropathol 116, 597-602, 2008). More recently,
20 IDH1/2 mutations have been observed in acute myeloid leukemia (AML) and rare cases have been reported in other cancers. (Ward P.S. et al., Cancer Cell 17, 225-234, 2010; Exp. Med. 207, 339-344, 2010; Mardis E. R. et al., N Engl J Med 361, 1058-1066, 2009). Notably, all mutations in IDH1/2 are heterozygous and the majority of them affect a particular codon (R132 in IDH1 and the analogous codon, R172, in IDH2) suggesting the
25 gene may contribute to carcinogenesis as an oncogene rather than a tumor suppressor through a gain of function.

IDH1/2 are NADP⁺-dependent isocitrate dehydrogenases that normally mediate oxidative decarboxylation of isocitrate to α -ketoglutarate (α -KG) via the conversion of NADP⁺ to NADPH. Mutations in IDH1/2 appear to have two functional consequences

for the enzymes' activities: i) a decreased ability to convert isocitrate to α -KG; and ii) a new ability to reduce α -KG to (R)-2-hydroxyglutarate (2-HG) using NADPH (Ward et al., *supra*; Dang, L. et al., *Nature* 462, 739-744, 2009). Indeed 2-HG levels are elevated >50-fold in samples from patients with IDH1/2-mutations. This observation has motivated
5 study of 2-HG as a disease biomarker, as well as deeper study into the molecular mechanism by which this putative 'oncometabolite' might contribute to disease. Indeed, 2-HG could interfere with a wide range of processes, such as those regulated by α -KG-dependent, iron-dependent dioxygenases; these processes include the response to hypoxic stress (mediated by EglN prolyl hydroxylases), DNA modification (mediated by TET2, a
10 5-methylcytosine hydroxylase), and histone methylation (mediated by JmjC-containing demethylases), among others (Figuroa M.E. et al., *Cancer Cell*, 18, 553-567, 2010; Christensen, B.C. et al., *J. Natl. Cancer Inst.* 103, 2, 143-53, 2011; Zhao, S. et al., *Science*, 324, 261-265, 2009; Xu, W. et al., *Cancer Cell*, 19, 17-30, 2011). Thus, the development of small molecules that inhibit the 2-HG-generating activity of IDH1/2 mutants in cells is
15 important in cancer cell biology and drug development.

Point mutations IDH1 and IDH2 occur early in the pathogenesis of gliomas. Reitman reports that the study of 200 metabolites in human oligodendroglioma (HOG) cells to determine the effects of expression of IDH1 and IDH2 mutants showed that the levels of amino acids, glutathione metabolites, choline derivatives, and tricarboxylic acid
20 (TCA) cycle intermediates were altered in mutant IDH1- and IDH2-expressing cells. (Reitman Z. J. et al., *Proc. Natl. Acad. Sci.* 2011, 108(8) 3270-3275). Furthermore, N-acetyl-aspartyl-glutamate (NAAG), a common dipeptide in brain, was 50-fold reduced in cells expressing IDH1 mutants and 8.3-fold reduced in cells expressing IDH2 mutants.


Hartmann et al., (US 20100291590) discloses a method for the diagnosis of a brain
25 tumor using the presence/absence of a particular IDH1 mutation as a marker. Vogelstein et al. (WO 2010/028099) discloses that mutations in IDH1 and IDH2 are related to astrocytomas, oligodendrogliomas and glioblastomas. Dang et al., (WO 2010/105243) discloses methods for the treatment of isocitrate dehydrogenase related proliferative disorders.

30 Glioblastoma is the most frequent and most malignant human brain tumor. The prognosis remains very poor, with most patients dying within 1 year after diagnosis. (Ohgaki et al. *American Journal of Pathology*, 170(5), 2007, 1445-1453). There exists a need to develop effective treatments against proliferative disorders including glioblastoma

and acute myeloid leukemia. Thus, there exists a need to focus on developing compounds that inhibit mutated IDH1/2 with selectivity over wild-type IDH1/2 with the goal of targeting cancer cells selectively over normal cell. That said, there is also a need for advancing compounds that target both mutant and wild-type IDH1/2 since cancer cells which harbor mutant IDH alleles are dependent on the wild-type allele for proliferation, suggesting inhibition of wild-type IDH may also prove valuable for treating cancer (Ward P.S. et al., Cancer Cell 17, 225-234, 2010).

SUMMARY OF THE INVENTION

The invention relates to compounds of Formula I or a pharmaceutically acceptable salt, ester or prodrug thereof and their use in the manufacture of a medicament, in particular, for the treatment of a cell proliferative disease:



Formula I

each n and m is independently 0, 1, 2 or 3;

each p and q is independently 0, 1, 2, 3, 4, 5, 6 or 7;

X_1 is $-C(O)N(R_A)-$, $-C(S)N(R_A)-$, or $-S(O)_2N(R_A)-$;

wherein R_A is hydrogen, aliphatic, substituted aliphatic, heteroaryl, substituted heteroaryl, aryl or substituted aryl;

X_2 is $-S-$, $-O-$, $-S(O)_2-$, $-C(R_{20})(R_{21})-$ or $-N(R_B)-$;

wherein R_B is hydrogen, aliphatic, substituted aliphatic, heteroaryl, substituted heteroaryl, aryl or substituted aryl;

each R_1 and R_2 is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl;

each R₁₀ is independently absent, hydrogen, halogen, -OR₂₀, -SR₂₀, -NR₂₀R₂₁, -CF₃, -CN, -NO₂, -N₃, -C(O)OR₂₀, -C(O)R₂₀, -C(O)N R₂₀R₂₁, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two of R₁₀ groups together with the atoms to which they are attached and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring;

wherein each R₂₀ and R₂₁ is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl;

each R₁₁ is independently absent, hydrogen, halogen, -OR₂₀, -S R₂₀, -NR₂₀R₂₁, -CF₃, -CN, -NO₂, -N₃, -C(O)OR₂₀, -C(O)R₂₀, -C(O)N R₂₀R₂₁, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two of R₁₁ groups together with the atoms to which they are attached and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring; and,

Cyl is an optionally substituted aryl or optionally substituted heteroaryl.

The invention further relates to the use of a compound of Formula I in the manufacture of a medicament. The invention further relates to the use of a compound of Formula I for the treatment of a cell proliferative disease and the use of a compound of Formula I in the manufacture of a medicament for the treatment of a cell proliferative disease.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1(A): Diagram of the assay principle for the IDH1-R132H enzymatic assay.

Figure 1(B): Optimization of resazurin concentration to give the highest signal-to-background ratio.

Figure 1(C): Reaction time course of the IDH1-R132H enzyme assay in buffers with magnesium or manganese.

Figure 1 (D): K_m of α-Ketoglutarate with 20 μM or 5 μM of NADPH.

Figure 1 (E) The effect of DMSO on the IDH1-R132H assay.


Figure 1 (F) Reaction time course of the IDH1-R132H assay.

Figure 1 (G) Concentration-response of the IDH1-R132H enzyme. The assay was linear at 8 µg/ml.

Figure 2(A-B): Both (A) consumption of NADPH by IDH1-R132H, and (B) production of NADPH by wild-type IDH1 can be monitored by measuring absorbance at
5 340 nm.

DETAILED DESCRIPTION OF THE INVENTION

The invention relates to compounds of Formula I or a pharmaceutically acceptable salt, ester or prodrug thereof, and their use in the manufacture of a medicament, in
10 particular, for the treatment of a cell proliferative disease:



Formula I

each n and m is independently 0, 1, 2 or 3;

15 each p and q is independently 0, 1, 2, 3, 4, 5, 6 or 7;

X_1 is $-C(O)N(R_A)-$, $-C(S)N(R_A)-$, or $-S(O)_2N(R_A)-$;

wherein R_A is hydrogen, aliphatic, substituted aliphatic, heteroaryl, substituted heteroaryl, aryl or substituted aryl;

X_2 is $-S-$, $-O-$, $-S(O)_2-$, $-C(R_{20})(R_{21})-$ or $-N(R_B)-$;

20 wherein R_B is hydrogen, aliphatic, substituted aliphatic, heteroaryl, substituted heteroaryl, aryl or substituted aryl;

each R_1 and R_2 is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl;

25 each R_{10} is independently absent, hydrogen, halogen, $-OR_{20}$, $-SR_{20}$, $-NR_{20}R_{21}$, $-CF_3$, $-CN$, $-NO_2$, $-N_3$, $-C(O)OR_{20}$, $-C(O)R_{20}$, $-C(O)N R_{20}R_{21}$, acyl, alkoxy, substituted


alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two of R_{10} groups together with the atoms to which they are attached and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring;

wherein each R_{20} and R_{21} is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl;


each R_{11} is independently absent, hydrogen, halogen, $-OR_{20}$, $-SR_{20}$, $-NR_{20}R_{21}$, $-CF_3$, $-CN$, $-NO_2$, $-N_3$, $-C(O)OR_{20}$, $-C(O)R_{20}$, $-C(O)NR_{20}R_{21}$, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two of R_{11} groups together with the atoms to which they are attached and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring; and,

Cy1 is an optionally substituted aryl or optionally substituted heteroaryl.

In one embodiment, the invention relates to a compound of Formula II-III or a pharmaceutically acceptable salt, ester or prodrug thereof:




Formula II



Formula III

wherein, n , p , q , X_2 , Cy1, R_1 , R_2 , R_{10} , and R_{11} are as defined above; and R_3 is hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl.


In a preferred embodiment, the invention relates to a compound of Formula IIIA:




Formula IIIA

- wherein R_{3a} is selected from alkyl, aryl, alkyl substituted with aryl, straight chain or
 5 branched C₁-C₁₀ alkyl, C₁-C₁₀ alkenyl, C₁-C₁₀ alkynyl, C₁-C₁₀ alkoxy, alkoxyC₁-
 C₁₀alkoxy, C₁-C₁₀alkylamino, alkoxyC₁-C₁₀alkylamino, C₁-C₁₀ alkylcarbonylamino, C₁-
 C₁₀ alkylaminocarbonyl, aryloxyC₁-C₁₀ alkoxy, aryloxyC₁-C₁₀alkylamino, aryloxyC₁-C₁₀
 alkylamino carbonyl, C₁-C₁₀-alkylaminoalkylaminocarbonyl, C₁-C₁₀alkyl(N-
 alkyl)aminoalkyl-aminocarbonyl, alkylaminoalkylamino, alkylcarbonylaminoalkylamino,
 10 alkyl(N-alkyl)aminoalkylamino, (N-alkyl)alkylcarbonylaminoalkylamino,
 alkylaminoalkyl, alkylaminoalkylaminoalkyl, alkylpiperazinoalkyl, piperazinoalkyl,
 alkylpiperazino, alkenylaryloxyC₁-C₁₀ alkoxy, alkenylarylaminoc₁-C₁₀ alkoxy,
 alkenylarylalkylamino C₁-C₁₀ alkoxy, alkenylaryloxyC₁-C₁₀ alkylamino,
 alkenylaryloxyC₁-C₁₀alkylaminocarbonyl, piperazinoalkylaryl, heteroarylC₁-C₁₀ alkyl,
 15 heteroarylC₁-C₁₀alkenyl, heteroarylC₁-C₁₀ alkynyl, heteroarylC₁-C₁₀ alkylamino,
 heteroarylC₁-C₁₀alkoxy, heteroaryloxyC₁-C₁₀alkyl, heteroaryloxyC₁-C₁₀ alkenyl,
 heteroaryloxyC₂-C₁₀ alkynyl, heteroaryloxyC₁-C₁₀alkylamino, heteroaryloxyC₁-C₁₀
 alkoxy.

- In one embodiment, the invention relates to a compound of Formula IV-V or a
 20 pharmaceutically acceptable salt, ester or prodrug thereof:



Formula IV



Formula V
















wherein p, q, X₂, R₁, R₂, R₃, R₁₀, and R₁₁ are as defined above.


In a preferred embodiment, the invention relates to a compound of Formula V wherein R₃ is R_{3a}, and R_{3a} is as defined above.

- In a preferred embodiment, the invention relates to a compound of Formula I, II, III, IV or V wherein, R₁ is an optionally substituted alkyl. In a preferred embodiment, R₁ is an optionally substituted C₁-C₆ alkyl, more preferably selected from methyl, ethyl, propyl, cyclopropyl, isopropyl, *n*-butyl, *tert*-butyl, cyclobutyl, *n*-pentyl, neopentyl, cyclopentyl, *n*-hexyl and cyclohexyl.

- In one embodiment, the invention relates to a compound of Formula I-III wherein
10 Cyl is selected from Table A:

TABLE A



wherein  represents a single or double bond;


each R_{12} is independently absent, hydrogen, $-C(O)R_{20}$, $-C(O)OR_{20}$, $-C(O)NR_{20}R_{21}$, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl, substituted aryl, heteroaryl or substituted heteroaryl;


d is 0, 1 or 2;

alternatively, two R_{11} and R_{12} groups may form an optionally substituted 3, 4, 5, 6, or 7 membered ring.

In one embodiment the invention relates to a compound of Formula I-V and R_{11} is selected from Table B:

TABLE B





wherein, t is 0, 1, 2, 3, 4, 5, 6 or 7;

Cy2 is an optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocyclyl, optionally substituted aryl or optionally substituted heteroaryl;

each R₁₃, R₁₄ and R₁₅ is independently absent, hydrogen, halogen, -OR₂₀, -SR₂₀, -NR₂₀R₂₁,

- 5 -CF₃, -CN, -NO₂, -N₃, -C(O)R₂₀, -C(O)OR₂₀, -C(O)N R₂₀R₂₁, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino,

substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two R_{13} groups together with the atoms to which they are attached and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring; and,

- 5 each R_{16} , R_{17} and R_{18} is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl substituted aryl, heteroaryl or substituted heteroaryl;

alternatively two of R_{13} , R_{14} , R_{15} , R_{16} , R_{17} and R_{18} groups together with the atoms to which they are attached, and any intervening atoms may form an optionally substituted 3, 4, 5, 6 or 7 membered ring.

- 10 In one embodiment the invention relates to a compound of Formula I-V wherein R_2 is selected from Table C:

TABLE C

wherein u and w is independently 0, 1, 2, 3, 4, 5 or 6; and,

Cy3 is optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocyclyl, optionally substituted aryl or optionally substituted heteroaryl.

In a preferred embodiment, u is 1. In a preferred embodiment, w is 1. In a preferred embodiment, Cy3 is an optionally substituted aryl, more preferably an optionally substituted heteroaryl.


In a preferred embodiment, the invention relates to a compound of Formula I-III wherein Cy1 is selected from Table A, q is 1 and R₁₁ is selected from Table B, and R₂ is selected from Table C.

In one embodiment, the invention relates to a compound of Formula II-III wherein n is 0, X₂ is O, and Cy1 is an optionally substituted aryl. In a preferred embodiment, Cy1 is an optionally substituted phenyl ring. In a preferred embodiment Cy1 is a phenyl group and R₁₁ is selected from Table B, and R₂ is selected from Table C.

In one embodiment the invention relates to a compound of Formula I-V wherein R₃ is selected from Table D:

15
















TABLE D





In a preferred embodiment, the invention relates to a compound of Formula I-V wherein Cy1 is selected from Table A, q is 1 and R₁₁ is selected from Table B, R₂ is selected from Table C, and R₃ is selected from Table D. In a more preferred embodiment, R₄ is an optionally substituted alkyl group, preferably a C1-C8 alkyl.

- 5 In one embodiment the invention relates to a compound of Formula III wherein R₃ is selected from Table E:

TABLE E





wherein, t, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇ and R₁₈ are as defined above;

r is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15;

R₃₀ is alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl;

each R₃₁, and R₃₂ is independently hydrogen, alkyl, substituted alkyl, aryl, substituted aryl,

5 heteroaryl or substituted heteroaryl; and,

R₃₃ is alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, -OR₂₀, -SR₂₀, -NR₂₀R₂₁, -CF₃, -CN, -NO₂, -N₃, -C(O)OR₂₀, -C(O)R₂₀, -C(O)N R₂₀R₂₁, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted




10 alkylsulfonyl wherein R₂₀ and R₂₁ are as defined above.





In a preferred embodiment, R₃₀ is C₁-C₁₀ alkyl, C₁-C₁₀alkylaryl, or C₁-C₁₀alkylarylalkoxy.





In a preferred embodiment, R₁₆ is aryl, substituted aryl, heteroaryl or substituted heteroaryl.





In a preferred embodiment, a compound of Formula I is selected from Table 1:





TABLE 1





1	 (4R,5R)-8-(cyclopent-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
2	 1-(4-fluorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
3	 (4R,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-8-phenyl-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide





4	 N-(((4R,5R)-8-(cyclohexylethynyl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-4-yl)acetamide
5	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
6	 N-(((4R,5R)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-8-(4-methylpent-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide
7	 N-((2S,3S)-5-((R)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-3,4,5,6-tetrahydro-2H-benzo[b][1,5]oxazin-10-yl)benzo[d]thiazole-2-carboxamide





8	 (4R,5R)-8-(cyclopropylethynyl)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
9	 (4R,5S)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-4-ylmethyl)amino)methyl)-8-(p-tolyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
10	 (4S,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-8-(pent-1-yn-1-yl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
11	 N-(((4R,5S)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-8-(4-methylpent-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide





12	 (4R,5S)-8-(cyclohex-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
13	 (4S,5S)-8-(cyclopent-1-en-1-yl)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-4-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
14	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)cyclohexanecarboxamide
15	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-2-(((4-methoxybenzyl)(methyl)amino)methyl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)cyclohexanecarboxamide




16	 (4S,5S)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-4-ylmethyl)amino)methyl)-8-(o-tolyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
17	 (2S,3S)-5-((R)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyrimidin-5-ylmethyl)amino)methyl)-8-((E)-styryl)-4,5-dihydro-2H-pyrido[2,3-b][1,5]oxazocin-6(3H)-one
18	 4-((((4R,5R)-8-(3-cyclopentylprop-1-yn-1-yl)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)(methyl)amino)methyl)benzoic acid
19	 N-(((4R,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-8-(3-methylbut-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methylnicotinamide




20	 N-(((4R,5R)-8-(cyclohex-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide
21	 N-(((4S,5S)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-8-(4-methylpent-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide
22	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-isopropylurea
23	 1-(4-bromophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea




24	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(4-methoxyphenyl)urea
25	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(pyridin-3-yl)urea
26	 1-cyclopentyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
27	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(2-morpholinopyridin-4-yl)urea





28	 1-ethyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
29	 1-(4-chlorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
30	 1-(cyclohexylmethyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
31	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)-3-phenylurea




32	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)-3-(naphthalen-1-yl)urea
33	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)-3-(4-(trifluoromethyl)phenyl)urea
34	 1-(benzo[d][1,3]dioxol-5-yl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
35	 1-(3,5-dimethylisoxazol-4-yl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea



36	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)cyclopropanecarboxamide
37	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)isonicotinamide
38	 1-(2-fluorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea

39	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(1,2,3,4-tetrahydronaphthalen-1-yl)urea
40	 1-(4-cyanophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
41	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(3-(pyrimidin-2-yl)phenyl)urea


42	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(2-(piperidin-1-yl)phenyl)urea
43	 1-(benzofuran-5-yl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
44	 1-cyclohexyl-3-((2R,3R)-2-(((3-fluoropyridin-4-yl)methyl)(methyl)amino)methyl)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea

45	 1-((2R,3R)-2-((((3-chloropyridin-4-yl)methyl)(methyl)amino)methyl)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)-3-cyclohexylurea
46	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyrimidin-4-yl)methyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
47	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(quinolin-4-yl)methyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
48	 1-((2R,3R)-2-((((1H-imidazol-2-yl)methyl)(methyl)amino)methyl)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)-3-cyclohexylurea

49	 1-cyclohexyl-3-((2R,3R)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-5-((S)-1-(2-(pyrrolidin-1-yl)ethoxy)propan-2-yl)-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
50	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-((4-methoxybenzyl)oxy)propan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
51	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-(4-(4-methoxyphenyl)-1H-1,2,3-triazol-1-yl)propan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea

52	 1-cyclohexyl-3-((2R,3R)-5-isopropyl-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
53	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-((4-methoxybenzyl)oxy)propan-2-yl)-3-methyl-2-((methyl((3-methylpyridin-4-yl)methyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea


The invention further relates to a process for preparing compounds of Formula I-V. Scheme 1 shows a general methodology for the synthesis of medium-sized ring scaffolds from a common linear intermediate used for the synthesis of final compounds.



Scheme 1: General protocol for synthesis of compounds of Formula I

R_C and R_D are each independently hydrogen, halogen, $-OR_{20}$, $-SR_{20}$, $-NR_{20}R_{21}$, $-CF_3$, $-CN$, $-NO_2$, $-N_3$, $-C(O)R_{20}$, $-C(O)OR_{20}$, $-C(O)NR_{20}R_{21}$, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl.

For this synthetic route (Scheme 1), intermediates and medium-sized ring scaffolds were synthesized using the procedures similar to Marcaurelle, L. A. et al., *J. Am. Chem. Soc.* **2010**, *132*, 16962-16976.



Scheme 2. Synthesis of intermediates and final compounds.

The compounds shown in the synthetic route (Scheme 2), intermediates and final compounds were synthesized using a procedure similar to that reported in Marcaurelle, L.A. et al., *J. Am. Chem. Soc.* **2010**, *132*, 16962-16976.

ABBREVIATIONS

Abbreviations which may appear in the synthetic schemes and examples are:

- Ac for acetyl;
- Alloc for allyloxycarbonyl;
- Boc for *tert*-butoxycarbonyl;
- DCM for dichloromethane;
- DMF for dimethyl formamide;
- DMSO for dimethyl sulfoxide;
- EtOAc for ethyl acetate;

iPr for isopropyl;

IPA for isopropyl alcohol;

MeOH for methanol;

TEA for triethylamine; and

5 TFA for trifluoroacetic acid.

The invention further provides methods for the prevention or treatment of diseases or conditions involving aberrant proliferation, differentiation or survival of cells. The invention relates to the treatment of cell proliferative disease, such as cancer, by the administration of a compound of Formula I-V to a patient in need thereof.

10 In a preferred embodiment, the cancer is selected from glioma, acute myeloid leukemia (AML), Burkitt's leukemia/lymphoma (B-ALL), melanoma and prostate carcinoma. In a preferred embodiment, the disease is selected from Grade I, II, III or IV glioma. In a preferred embodiment, the disease is selected from astrocytomas, oligodendrogliomas, ependymomas and glioblastoma multiforme (GBM).

15 The invention further relates to a method of treating a disease related to a defect in isocitrate dehydrogenase comprising the step of administering a compound according to any of the above claims to a patient in need thereof, in particular wherein said defect in isocitrate dehydrogenase is a somatic mutation at codon 132 isocitrate dehydrogenase (IDH1) or at codon 172 in isocitrate dehydrogenase 2 (IDH2). The invention further
20 relates to the treatment of Grade I, II, III and IV glioma. The glioma can be selected from astrocytomas, oligodendrogliomas, ependymomas and glioblastoma multiforme (GBM).

The invention further provides methods for the prevention or treatment of diseases related to defects in isocitrate dehydrogenase (IDH1 or IDH2) by administration of a compound of Formula I-III. The invention relates to diseases associated with somatic
25 mutations at codon 132 isocitrate dehydrogenase (IDH1), and at codon 172 and 140 in isocitrate dehydrogenase 2 (IDH2). The invention further relates to the treatment of patients with one more mutations at codons 132 of IDH1 and 172 or 140 of IDH2. The invention further relates to the treatment of diseases associated with mutations to IDH1 selected from R132H, R132C, R132S, R132L, and R132G. The invention further relates
30 to treatment of diseases associated with mutations to IDH2 selected from R172M, R172G, R172K and R140Q. In one embodiment, the disease associated with the above mentioned mutations is a cell proliferative disease, in particular cancer. In a preferred embodiment,

the cell proliferative disease is selected from glioma, acute myeloid leukemia (AML), Burkitt's leukemia/lymphoma (B-ALL), melanoma and prostate carcinoma. In a preferred embodiment, the disease is selected from Grade I, II, III or IV glioma. In a preferred embodiment, the disease is selected from astrocytomas, oligodendrogliomas, endependymomas and glioblastoma multiforme (GBM).

In one embodiment, the invention relates to the inhibition of IDH1/2 by administration of a compound of Formula I-V. In one embodiment, the compound of Formula I-V inhibits the wild type IDH1. In one embodiment, the compound of Formula I-V inhibits wild type IDH2.

In one embodiment, the invention relates to the administration of a compound of Formula I-V in patient that exhibit abnormal 2-hydroxyglutarate (2-HG) production. In one embodiment, the patient exhibit an increase in 2-HG production of more than about 20 fold, or more than about 40 fold or more than about 50 fold or more than about 100 fold or more than about 200 fold compared to normal tissue. The increased 2-HG production can be localized to tumor tissue. In one embodiment, the abnormal 2-HG production is due to a mutation in IDH1 or IDH2.

In one embodiment, the invention relates to a method of treating a disease related to a defect in isocitrate dehydrogenase comprising the step of administering a compound according to a compound of Formula I-V to a patient in need thereof, wherein the compound selectively inhibit mutant of IDH1 or a mutant of IDH2 over the respective wild type IDH. In one embodiment, a compound of Formula I-V selectively inhibits a mutant of IDH1 over the wild type IDH1. In one embodiment, the ratio of inhibitory activity against an IDH1 mutant over the wild type IDH1 is about 2 to about 1000, preferably about 5 to about 500, preferably about 10 to about 100, preferably about 25 to about 100. In one embodiment, the mutation to IDH1 is selected from R132H, R132C, R132S, R132L, and R132G. In one embodiment, a compound of Formula I-V selectively inhibits a mutant of IDH2 over the wild type IDH2. In one embodiment, the ratio of inhibitory activity against an IDH2 mutant over the wild type IDH2 is about 2 to about 1000, preferably about 5 to about 500, preferably about 10 to about 100, preferably about 25 to about 100. In one embodiment, the mutation to IDH2 is selected from R172M, R172G, R172K and R140Q.

The invention further relates to the use of a compound of Formula I in the manufacture of a medicament. The invention further relates to the use of a compound of Formula I for the treatment of a cell proliferative disease and the use of a compound of Formula I in the manufacture of a medicament for the treatment of a cell proliferative disease.

In one embodiment, the invention relates to the administration of a compound of Formula I-V in combination with a second pharmacological agent for the treatment of a cell proliferative disease. In one embodiment, the second pharmacological agent is selected from vatalanib (PTK-787/ZK222584), SU-5416, SU-6668, SU-11248, SU-14813, AZD-6474, AZD-2171, CP-547632, CEP-7055, AG-013736, IM-842 or GW-786034, gefitinib, erlotinib, HKI-272, CI-1033 or GW-2016, iressa (ZD-1839), tarceva (OSI-774), PKI-166, EKB-569, herceptin, BAY-43-9006, BAY-57-9006, atrasentan, rituximab, cetuximab, bevacizumab, bivatuzumab mertansine, IMC-1C11, erbitux (C-225), DC-101, EMD-72000, vitaxin, imatinib or dasatinib, VEGFtrap, melphalan, cyclophosphamide, an oxazaphosphorine, cisplatin, carboplatin, oxaliplatin, satraplatin, tetraplatin, iproplatin, mitomycin, streptozocin, carmustine (BCNU), lomustine (CCNU), busulfan, ifosfamide, streptozocin, thiotepa, chlorambucil, mechlorethamine, an ethyleneimine compound, an alkylsulphonate, daunorubicin, doxorubicin (adriamycin), liposomal doxorubicin (doxil), epirubicin, idarubicin, mitoxantrone, amsacrine, dactinomycin, distamycin or a derivative thereof, netropsin, pibenzimol, mitomycin, CC-1065, a duocarmycin, mithramycin, chromomycin, olivomycin, propamidine or stilbamidine, an anthramycin, an aziridine, a nitrosourea or a derivative thereof, cytarabine, 5-fluorouracil (5-FU), pemetrexed, tegafur/uracil, uracil mustard, fludarabine, gemcitabine, capecitabine, mercaptopurine, cladribine, thioguanine, methotrexate, pentostatin, hydroxyurea, or folic acid, a phleomycin, a bleomycin or a derivative or salt thereof, CHPP, BZPP, MTPP, BAPP, liblomycin, an acridine or a derivative thereof, a rifamycin, an actinomycin, adramycin, a camptothecin such as irinotecan (camptosar) or topotecan, an amsacrine or analogue thereof, a tricyclic carboxamide, a histone deacetylase inhibitor such as SAHA, MD-275, trichostatin A, CBHA, LAQ824, or valproic acid, an anti-cancer drug from plants such as paclitaxel (taxol), docetaxel or taxotere, navelbine, vinblastin, vincristin, vindesine, vinorelbine, colchicine or a derivative thereof, maytansine, an ansamitocin or rhizoxin, phomopsin, dolastatin, an epipodophyllotoxin or a derivative of podophyllotoxin, etoposide, teniposide, a steganacin, combretastatin, amphetinile, procarbazine,

bortezomib, asparaginase, pegylated asparaginase (pegaspargase), a thymidine-phosphorylase inhibitor, a gestagen, an estrogen, estramustine (T-66), megestrol, an anti-androgen, flutamide, casodex, anandron or cyproterone acetate, aminoglutethimide, anastrozole, formestan, exemestane, letrozole, leuporelin, buserelin, goserelin, triptorelin,

5 an anti-estrogen, tamoxifen or its citrate salt, droloxifene, trioxifene, raloxifene, zindoxifene, an estrogen receptor antagonist such as fulvestrant, a derivative of 17.β-estradiol, ICI 164,384, ICI 182,780, aminoglutethimide, formestane, fadrozole, finasteride, ketoconazole, a LH-RH antagonist, leuprolide, a steroid, prednisone, prednisolone, methylprednisolone, dexamethasone, budenoside, fluocortolone, triamcinolone, interferon

10 .β., IL-10, IL-12, an anti-TNF.α. antibody, etanercept, TNF-α. (tasonermin), thalidomide and its R- and S-enantiomers and its derivatives, revimid (CC-5013), a leukotrien antagonist, mitomycin C, BMY-42355, AZQ or EO-9, a 2-nitroimidazole misonidazole, NLP-1 or NLA-1, a nitroacridine, a nitroquinoline, a nitropyrazoloacridine, RSU-1069, RB-6145, CB-1954, nitromin, an anti-CD3 or anti-CD25 antibody, a tolerance

15 induction agent, minodronic acid and its derivatives (YM-529, Ono-5920, YH-529), zoledronic acid monohydrate, ibandronate sodium hydrate, clodronate disodium, metronidazole, misonidazole, benznidazole, nimorazole, RSU-1069, SR-4233, bromodeoxyuridine, iododeoxyuridine, WR-2721, porfimer, photofrin, a benzoporphyrin derivative, a pheophorbide derivative, merocyanin 540 (MC-540), tin etioporphyrin, an ant-

20 template, an anti-sense RNA or DNA, oblimersen, a non-steroidal inflammatory drug, acetylsalicylic acid, mesalazin, ibuprofen, naproxen, flurbiprofen, fenoprofen, fenbufen, ketoprofen, indoprofen, piroprofen, carprofen, oxaprozin, pranoprofen, miroprofen, tioxaprofen, suprofen, alminoprofen, tiaprofenic acid, fluprofen, indomethacin, sulindac, tolmetin, zomepirac, nabumetone, diclofenac, fenclofenac, alclofenac, bromfenac,

25 ibufenac, aceclofenac, acemetacin, fentiazac, clidanac, etodolac, oxpinac, mefenamic acid, meclofenamic acid, flufenamic acid, niflumonic acid, tolfenamic acid, diflunisal, flufenisal, piroxicam, tenoxicam, lornoxicam, nimesulide, meloxicam, celecoxib, rofecoxib, a pharmaceutically acceptable salt of a non-steroidal inflammatory drug, a cytotoxic antibiotic, an antibody targeting the surface molecules of cancer cells, apolizumab,

30 1D09C3, TIMP-1, TIMP-2, Zinc, an inhibitor of oncogenes, P53, R.sup.b, heterocyclic complexes of lanthanides, PUVA, an inhibitor of the transcription factor complex ESX/DRIP130/Sur-2, an inhibitor of HER-2 expression, the heat shock protein HSP90 modulator geldanamycin and its derivative 17-allylaminogeldanamycin or 17-AAG, therapeutic agent selected from IM-842, tetrathiomolybdate, squalamine, combrestatin A4,

TNP-470, marimastat, neovastat, bicalutamide, abarelix, oregovomab, mitumomab, TLK-286, alemtuzumab, ibritumomab, temozolomide, denileukin diftitox, aldesleukin, dacarbazine, floxuridine, plicamycin, mitotane, pipobroman, plicamycin, tamoxifen and testolactone.

5 In one embodiment, the invention further provides for the use of one or more compounds of the invention in the manufacture of a medicament for halting or decreasing diseases involving aberrant proliferation, differentiation, or survival of cells. In one embodiment, the invention relates to a method of treating cancer in a subject in need of treatment comprising administering to said subject a therapeutically effective amount of a
10 compound of the invention.

Definitions

 Listed below are definitions of various terms used to describe this invention. These definitions apply to the terms as they are used throughout this specification and claims, unless otherwise limited in specific instances, either individually or as part of a larger
15 group.

 The term "aliphatic group" or "aliphatic" refers to a non-aromatic moiety that may be saturated (e.g. single bond) or contain one or more units of unsaturation, e.g., double and/or triple bonds. An aliphatic group may be straight chained, branched or cyclic, contain carbon, hydrogen or, optionally, one or more heteroatoms and may be substituted
20 or unsubstituted. In addition to aliphatic hydrocarbon groups, aliphatic groups include, for example, polyalkoxyalkyls, such as polyalkylene glycols, polyamines, and polyimines, for example. Such aliphatic groups may be further substituted. It is understood that aliphatic groups may include alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, and substituted or unsubstituted cycloalkyl groups as described herein.

25 The term "acyl" refers to a carbonyl substituted with hydrogen, alkyl, partially saturated or fully saturated cycloalkyl, partially saturated or fully saturated heterocycle, aryl, or heteroaryl. For example, acyl includes groups such as (C₁-C₆) alkanoyl (e.g., formyl, acetyl, propionyl, butyryl, valeryl, caproyl, t-butylacetyl, etc.), (C₃-C₆)cycloalkylcarbonyl (e.g., cyclopropylcarbonyl, cyclobutylcarbonyl, cyclopentylcarbonyl, cyclohexylcarbonyl, etc.), heterocyclic carbonyl (e.g.,
30 pyrrolidinylcarbonyl, pyrrolid-2-one-5-carbonyl, piperidinylcarbonyl, piperazinylcarbonyl, tetrahydrofuranylcarbonyl, etc.), aroyl (e.g., benzoyl) and heteroaroyl (e.g., thiophenyl-2-

carbonyl, thiophenyl-3-carbonyl, furanyl-2-carbonyl, furanyl-3-carbonyl, 1H-pyrrolyl-2-carbonyl, 1H-pyrrolyl-3-carbonyl, benzo[b]thiophenyl-2-carbonyl, etc.). In addition, the alkyl, cycloalkyl, heterocycle, aryl and heteroaryl portion of the acyl group may be any one of the groups described in the respective definitions. When indicated as being

5 "optionally substituted", the acyl group may be unsubstituted or optionally substituted with one or more substituents (typically, one to three substituents) independently selected from the group of substituents listed below in the definition for "substituted" or the alkyl, cycloalkyl, heterocycle, aryl and heteroaryl portion of the acyl group may be substituted as described above in the preferred and more preferred list of substituents, respectively.

10 The term "alkyl" is intended to include both branched and straight chain, substituted or unsubstituted saturated aliphatic hydrocarbon radicals/groups having the specified number of carbons. Preferred alkyl groups comprise about 1 to about 24 carbon atoms ("C₁-C₂₄"). Other preferred alkyl groups comprise at about 1 to about 8 carbon atoms ("C₁-C₈") such as about 1 to about 6 carbon atoms ("C₁-C₆"), or such as about 1 to
15 about 3 carbon atoms ("C₁-C₃"). Examples of C₁-C₆ alkyl radicals include, but are not limited to, methyl, ethyl, propyl, isopropyl, *n*-butyl, *tert*-butyl, *n*-pentyl, neopentyl and *n*-hexyl radicals.

The term "alkenyl" refers to linear or branched radicals having at least one carbon-carbon double bond. Such radicals preferably contain from about two to about twenty-
20 four carbon atoms ("C₂-C₂₄"). Other preferred alkenyl radicals are "lower alkenyl" radicals having two to about ten carbon atoms ("C₂-C₁₀") such as ethenyl, allyl, propenyl, butenyl and 4-methylbutenyl. Preferred lower alkenyl radicals include 2 to about 6 carbon atoms ("C₂-C₆"). The terms "alkenyl", and "lower alkenyl", embrace radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations.

25 The term "alkynyl" refers to linear or branched radicals having at least one carbon-carbon triple bond. Such radicals preferably contain from about two to about twenty-four carbon atoms ("C₂-C₂₄"). Other preferred alkynyl radicals are "lower alkynyl" radicals having two to about ten carbon atoms such as propargyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne and 1-pentyne. Preferred lower alkynyl radicals include 2 to about 6
30 carbon atoms ("C₂-C₆").

The term "cycloalkyl" refers to saturated carbocyclic radicals having three to about twelve carbon atoms ("C₃-C₁₂"). The term "cycloalkyl" embraces saturated carbocyclic

radicals having three to about twelve carbon atoms. Examples of such radicals include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term "cycloalkenyl" refers to partially unsaturated carbocyclic radicals having three to twelve carbon atoms. Cycloalkenyl radicals that are partially unsaturated carbocyclic radicals that contain two double bonds (that may or may not be conjugated) can be called "cycloalkyldienyl". More preferred cycloalkenyl radicals are "lower cycloalkenyl" radicals having four to about eight carbon atoms. Examples of such radicals include cyclobutenyl, cyclopentenyl and cyclohexenyl.

The term "alkylene," as used herein, refers to a divalent group derived from a straight chain or branched saturated hydrocarbon chain having the specified number of carbons atoms. Examples of alkylene groups include, but are not limited to, ethylene, propylene, butylene, 3-methyl-pentylene, and 5-ethyl-hexylene.

The term "alkenylene," as used herein, denotes a divalent group derived from a straight chain or branched hydrocarbon moiety containing the specified number of carbon atoms having at least one carbon-carbon double bond. Alkenylene groups include, but are not limited to, for example, ethenylene, 2-propenylene, 2-butenylene, 1-methyl-2-buten-1-yne, and the like.

The term "alkynylene," as used herein, denotes a divalent group derived from a straight chain or branched hydrocarbon moiety containing the specified number of carbon atoms having at least one carbon-carbon triple bond. Representative alkynylene groups include, but are not limited to, for example, propynylene, 1-butyne, 2-methyl-3-hexynylene, and the like.

The term "alkoxy" refers to linear or branched oxy-containing radicals each having alkyl portions of one to about twenty-four carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkoxy radicals are "lower alkoxy" radicals having one to about ten carbon atoms and more preferably having one to about eight carbon atoms. Examples of such radicals include methoxy, ethoxy, propoxy, butoxy and tert-butoxy.

The term "alkoxyalkyl" refers to alkyl radicals having one or more alkoxy radicals attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl radicals.

The term "aryl", alone or in combination, means an aromatic system containing one, two or three rings wherein such rings may be attached together in a pendent manner

or may be fused. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indane furanyl, quinazolinyl, pyridyl and biphenyl.

The terms "heterocyclyl", "heterocycle" "heterocyclic" or "heterocyclo" refer to saturated, partially unsaturated and unsaturated heteroatom-containing ring-shaped radicals, which can also be called "heterocyclyl", "heterocycloalkenyl" and "heteroaryl" correspondingly, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. Examples of saturated heterocyclyl radicals include saturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms (e.g. pyrrolidinyl, imidazolidinyl, piperidino, piperazinyl, etc.); saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. morpholinyl, etc.); saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., thiazolidinyl, etc.). Examples of partially unsaturated heterocyclyl radicals include dihydrothiophene, dihydropyran, dihydrofuran and dihydrothiazole. Heterocyclyl radicals may include a pentavalent nitrogen, such as in tetrazolium and pyridinium radicals. The term "heterocycle" also embraces radicals where heterocyclyl radicals are fused with aryl or cycloalkyl radicals. Examples of such fused bicyclic radicals include benzofuran, benzothiophene, and the like.

The term "heteroaryl" refers to unsaturated aromatic heterocyclyl radicals. Examples of heteroaryl radicals include unsaturated 3 to 6 membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl (e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.), tetrazolyl (e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.), etc.; unsaturated condensed heterocyclyl group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indolizinyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl (e.g., tetrazolo[1,5-b]pyridazinyl, etc.), etc.; unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl, etc.; unsaturated 3 to 6-membered heteromonocyclic group containing a sulfur atom, for example, thienyl, etc.; unsaturated 3- to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.) etc.; unsaturated condensed heterocyclyl group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. benzoxazolyl, benzoxadiazolyl, etc.); unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1

to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl (e.g., 1,2,4- thiadiazolyl, 1,3,4- thiadiazolyl, 1,2,5-thiadiazolyl, etc.) etc.; unsaturated condensed heterocyclyl group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., benzothiazolyl, benzothiadiazolyl, etc.) and the like.

5 The term "heterocycloalkyl" refers to heterocyclo-substituted alkyl radicals. More preferred heterocycloalkyl radicals are "lower heterocycloalkyl" radicals having one to six carbon atoms in the heterocyclo radical.

 The term "alkylthio" refers to radicals containing a linear or branched alkyl radical, of one to about ten carbon atoms attached to a divalent sulfur atom. Preferred alkylthio
10 radicals have alkyl radicals of one to about twenty-four carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkylthio radicals have alkyl radicals which are "lower alkylthio" radicals having one to about ten carbon atoms. Most preferred are alkylthio radicals having lower alkyl radicals of one to about eight carbon atoms. Examples of such lower alkylthio radicals include methylthio, ethylthio, propylthio,
15 butylthio and hexylthio.

 The terms "aralkyl" or "arylalkyl" refer to aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenylethyl, and diphenylethyl.

 The term "aryloxy" refers to aryl radicals attached through an oxygen atom to other radicals.

20 The terms "aralkoxy" or "arylalkoxy" refer to aralkyl radicals attached through an oxygen atom to other radicals.

 The term "aminoalkyl" refers to alkyl radicals substituted with amino radicals. Preferred aminoalkyl radicals have alkyl radicals having about one to about twenty-four carbon atoms or, preferably, one to about twelve carbon atoms. More preferred
25 aminoalkyl radicals are "lower aminoalkyl" that have alkyl radicals having one to about ten carbon atoms. Most preferred are aminoalkyl radicals having lower alkyl radicals having one to eight carbon atoms. Examples of such radicals include aminomethyl, aminoethyl, and the like.

 The term "alkylamino" denotes amino groups which are substituted with one or
30 two alkyl radicals. Preferred alkylamino radicals have alkyl radicals having about one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkylamino radicals are "lower alkylamino" that have alkyl radicals having one

to about ten carbon atoms. Most preferred are alkylamino radicals having lower alkyl radicals having one to about eight carbon atoms. Suitable lower alkylamino may be monosubstituted N-alkylamino or disubstituted N,N-alkylamino, such as N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino or the like.

5 The term "substituted" refers to the replacement of one or more hydrogen radicals in a given structure with the radical of a specified substituent including, but not limited to: halo, alkyl, alkenyl, alkynyl, aryl, heterocyclyl, thiol, alkylthio, arylthio, alkylthioalkyl, arylthioalkyl, alkylsulfonyl, alkylsulfonylalkyl, arylsulfonylalkyl, alkoxy, aryloxy, aralkoxy, aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, alkoxycarbonyl,
10 aryloxycarbonyl, haloalkyl, amino, trifluoromethyl, cyano, nitro, alkylamino, arylamino, alkylaminoalkyl, arylaminoalkyl, aminoalkylamino, hydroxy, alkoxyalkyl, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, acyl, aralkoxycarbonyl, carboxylic acid, sulfonic acid, sulfonyl, phosphonic acid, aryl, heteroaryl, heterocyclic, and aliphatic. It is understood that the substituent may be further substituted.

15 For simplicity, chemical moieties that are defined and referred to throughout can be univalent chemical moieties (e.g., alkyl, aryl, etc.) or multivalent moieties under the appropriate structural circumstances clear to those skilled in the art. For example, an "alkyl" moiety can be referred to a monovalent radical (e.g. $\text{CH}_3\text{-CH}_2\text{-}$), or in other instances, a bivalent linking moiety can be "alkyl," in which case those skilled in the art
20 will understand the alkyl to be a divalent radical (e.g., $\text{-CH}_2\text{-CH}_2\text{-}$), which is equivalent to the term "alkylene." Similarly, in circumstances in which divalent moieties are required and are stated as being "alkoxy", "alkylamino", "aryloxy", "alkylthio", "aryl", "heteroaryl", "heterocyclic", "alkyl", "alkenyl", "alkynyl", "aliphatic", or "cycloalkyl", those skilled in the art will understand that the terms alkoxy", "alkylamino", "aryloxy",
25 "alkylthio", "aryl", "heteroaryl", "heterocyclic", "alkyl", "alkenyl", "alkynyl", "aliphatic", or "cycloalkyl" refer to the corresponding divalent moiety.

 The terms "halogen" or "halo" as used herein, refers to an atom selected from fluorine, chlorine, bromine and iodine.

 The terms "compound", "drug", and "prodrug" as used herein all include
30 pharmaceutically acceptable salts, co-crystals, solvates, hydrates, polymorphs, enantiomers, diastereoisomers, racemates and the like of the compounds, drugs and prodrugs having the formulas as set forth herein.

Substituents indicated as attached through variable points of attachments can be attached to any available position on the ring structure.

As used herein, the term "effective amount of the subject compounds," with respect to the subject method of treatment, refers to an amount of the subject compound which, when delivered as part of desired dose regimen, brings about management of the disease or disorder to clinically acceptable standards.

"Treatment" or "treating" refers to an approach for obtaining beneficial or desired clinical results in a patient. For purposes of this invention, beneficial or desired clinical results include, but are not limited to, one or more of the following: alleviation of symptoms, diminishment of extent of a disease, stabilization (i.e., not worsening) of a state of disease, preventing spread (i.e., metastasis) of disease, preventing occurrence or recurrence of disease, delay or slowing of disease progression, amelioration of the disease state, and remission (whether partial or total).

The term "cancer" refers to any cancer caused by the proliferation of malignant neoplastic cells, such as tumors, neoplasms, carcinomas, sarcomas, leukemias, lymphomas and the like. For example, cancers include, but are not limited to, mesothelioma, leukemias and lymphomas such as cutaneous T-cell lymphomas (CTCL), noncutaneous peripheral T-cell lymphomas, lymphomas associated with human T-cell lymphotropic virus (HTLV) such as adult T-cell leukemia/lymphoma (ATLL), B-cell lymphoma, acute nonlymphocytic leukemias, chronic lymphocytic leukemia, chronic myelogenous leukemia, acute myelogenous leukemia, lymphomas, and multiple myeloma, non-Hodgkin lymphoma, acute lymphatic leukemia (ALL), chronic lymphatic leukemia (CLL), Hodgkin's lymphoma, Burkitt lymphoma, adult T-cell leukemia lymphoma, acute-myeloid leukemia (AML), chronic myeloid leukemia (CML), or hepatocellular carcinoma. Further examples include myelodysplastic syndrome, childhood solid tumors such as brain tumors, neuroblastoma, retinoblastoma, Wilms' tumor, bone tumors, and soft-tissue sarcomas, common solid tumors of adults such as head and neck cancers (*e.g.*, oral, laryngeal, nasopharyngeal and esophageal), genito urinary cancers (*e.g.*, prostate, bladder, renal, uterine, ovarian, testicular), lung cancer (*e.g.*, small-cell and non-small cell), breast cancer, pancreatic cancer, melanoma and other skin cancers, stomach cancer, brain tumors, tumors related to Gorlin's syndrome (*e.g.*, medulloblastoma, meningioma, etc.), and liver cancer. Additional exemplary forms of cancer which may be treated by the subject compounds include, but are not limited to, cancer of skeletal or smooth muscle, stomach cancer,

cancer of the small intestine, rectum carcinoma, cancer of the salivary gland, endometrial cancer, adrenal cancer, anal cancer, rectal cancer, parathyroid cancer, and pituitary cancer.

Additional cancers that the compounds described herein may be useful in preventing, treating and studying are, for example, colon carcinoma, familial
5 adenomatous polyposis carcinoma and hereditary non-polyposis colorectal cancer, or melanoma. Further, cancers include, but are not limited to, labial carcinoma, larynx carcinoma, hypopharynx carcinoma, tongue carcinoma, salivary gland carcinoma, gastric carcinoma, adenocarcinoma, thyroid cancer (medullary and papillary thyroid carcinoma), renal carcinoma, kidney parenchyma carcinoma, cervix carcinoma, uterine corpus
10 carcinoma, endometrium carcinoma, chorion carcinoma, testis carcinoma, urinary carcinoma, melanoma, brain tumors such as glioblastoma, astrocytoma, meningioma, medulloblastoma and peripheral neuroectodermal tumors, gall bladder carcinoma, bronchial carcinoma, multiple myeloma, basalioma, teratoma, retinoblastoma, choroidea melanoma, seminoma, rhabdomyosarcoma, craniopharyngeoma, osteosarcoma,
15 chondrosarcoma, myosarcoma, liposarcoma, fibrosarcoma, Ewing sarcoma, and plasmocytoma. In one aspect of the invention, the present invention provides for the use of one or more compounds of the invention in the manufacture of a medicament for the treatment of cancer.

In one embodiment, the present invention includes the use of one or more
20 compounds of the invention in the manufacture of a medicament that prevents further aberrant proliferation, differentiation, or survival of cells. For example, compounds of the invention may be useful in preventing tumors from increasing in size or from reaching a metastatic state. The subject compounds may be administered to halt the progression or advancement of cancer. In addition, the instant invention includes use of the subject
25 compounds to prevent a recurrence of cancer.

This invention further embraces the treatment or prevention of cell proliferative disorders such as hyperplasias, dysplasias and pre-cancerous lesions. Dysplasia is the earliest form of pre-cancerous lesion recognizable in a biopsy by a pathologist. The subject compounds may be administered for the purpose of preventing said hyperplasias,
30 dysplasias or pre-cancerous lesions from continuing to expand or from becoming cancerous. Examples of pre-cancerous lesions may occur in skin, esophageal tissue, breast and cervical intra-epithelial tissue.

"Combination therapy" includes the administration of the subject compounds in further combination with other biologically active ingredients (such as, but not limited to, a second and different antineoplastic agent) and non-drug therapies (such as, but not limited to, surgery or radiation treatment). For instance, the compounds of the invention can be used in combination with other pharmaceutically active compounds, preferably compounds that are able to enhance the effect of the compounds of the invention. The compounds of the invention can be administered simultaneously (as a single preparation or separate preparation) or sequentially to the other drug therapy. In general, a combination therapy envisions administration of two or more drugs during a single cycle or course of therapy.

Pharmaceutical Compositions

The pharmaceutical compositions of the present invention comprise a therapeutically effective amount of a compound of the present invention formulated together with one or more pharmaceutically acceptable carriers or excipients.

As used herein, the term "pharmaceutically acceptable carrier or excipient" means a non-toxic, inert solid, semi-solid, gel or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Some examples of materials which can serve as pharmaceutically acceptable carriers are sugars such as lactose, glucose and sucrose; cyclodextrins such as alpha- (α), beta- (β) and gamma- (γ) cyclodextrins; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols such as propylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

The pharmaceutical compositions of this invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an

implanted reservoir. In a preferred embodiment, administration is parenteral administration by injection.

The pharmaceutical compositions of this invention may contain any conventional non-toxic pharmaceutically-acceptable carriers, adjuvants or vehicles. In some cases, the pH of the formulation may be adjusted with pharmaceutically acceptable acids, bases or buffers to enhance the stability of the formulated compound or its delivery form. The term parenteral as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, intrathecal, intralesional and intracranial injection or infusion techniques.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions, may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable suspension or emulsion, such as INTRALIPID®, LIPOSYN® or OMEGAVEN®, or solution, in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. INTRALIPID® is an intravenous fat emulsion containing 10-30% soybean oil, 1-10% egg yolk phospholipids, 1-10% glycerin and water. LIPOSYN® is also an intravenous fat emulsion containing 2-15% safflower oil, 2-15% soybean oil, 0.5-5% egg phosphatides 1-10% glycerin and water. OMEGAVEN® is an emulsion for infusion containing about 5-25% fish oil, 0.5-10% egg phosphatides, 1-10% glycerin and water. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, USP and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For

this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

5 The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

10 Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

15 Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or: a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents
20 such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and
25 mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

30 The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and

can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

5 Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, eye ointments, powders and solutions are
10 also contemplated as being within the scope of this invention.

The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.


15 Powders and sprays can contain, in addition to the compounds of this invention, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder; or mixtures of these substances. Sprays can additionally contain customary propellants such as chlorofluorohydrocarbons.

Transdermal patches have the added advantage of providing controlled delivery of
20 a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

For pulmonary delivery, a therapeutic composition of the invention is formulated
25 and administered to the patient in solid or liquid particulate form by direct administration e.g., inhalation into the respiratory system. Solid or liquid particulate forms of the active compound prepared for practicing the present invention include particles of respirable size: that is, particles of a size sufficiently small to pass through the mouth and larynx upon inhalation and into the bronchi and alveoli of the lungs. Delivery of aerosolized
30 therapeutics is known in the art (see, for example U.S. Pat. No. 5,767,068 to VanDevanter *et al.*, U.S. Pat. No. 5,508,269 to Smith *et al.*, and WO 98/43650 by Montgomery).

EXAMPLES

Example 1: Synthesis of (4R,5R)-8-(cyclopent-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide



5

Scaffold loading: To a flame-dried flask containing silicon-functionalized Lanterns was added a freshly prepared solution of TfOH in anhydrous DCM (9.0 equiv, 5 g of TfOH/100 mL of DCM) was added. Each flask was shaken at RT for 10 min at which time the Lanterns had turned bright orange. The deep red TfOH solution was removed via

10 cannula and anhydrous 2,6-lutidine (12.0 equiv relative to Si) was added. Once the Lantern color had changed from orange to white, (9H-fluoren-9-yl)methyl (((4R,5R)-8-bromo-2-((S)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)(methyl)carbamate, the scaffold, (1.2

15 equiv. relative to Si) was added as a solution in anhydrous DCM (0.4 mL/Lantern) and the reaction mixture was shaken for 48h overnight. The loading mixture was removed and set aside (to recover any unreacted alcohol) and the Lanterns were washed with the following solvents for 30 min intervals: DCM, THF, 3:1 THF/IPA, 3:1 THF/H₂O, DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. The Lanterns were then dried on a lyophilizer overnight and carried on to the next step.

20 **Fmoc removal:** To a flask containing Lanterns was added a solution of 20% piperidine in DMF (0.8mL/Lantern). After shaking at rt for 30 min, the piperidine solution was removed and the Lanterns were washed with the following solvents for 30 min intervals: DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material. The Lanterns were then dried on a

25 lyophilizer overnight and carried on to the next step.


N-Capping/Aldehydes: To a flask containing Lanterns was added DMF with 2% AcOH (0.800 mL/Lantern) followed by 4-pyridinecarboxaldehyde (20 equiv). The reaction

mixture was shaken at rt for 1 hr then sodium triacetoxymethylborohydride (20 equiv) was added and shaking was continued. After 3 days the reaction mixture was removed and the Lanterns were washed with the following solvents for 30 min intervals: DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material. The Lanterns were then dried on a lyophilizer overnight and carried on to the next step.

Cross-Coupling/Suzuki: To each flask containing Lanterns was added ethanol (0.800 mL/Lantern) followed by cyclopent-1-en-1-ylboronic acid (20 equiv), triethylamine (40 equiv) and Pd(PPh₃)₂Cl₂ (1 equiv). The resulting mixture was degassed with a stream of N₂ before shaking at 60 °C. After 4 days, the reaction mixture was removed and the Lanterns were washed with following solvents for 30 min intervals: DCM, DMF, NaCN solution (0.1M) in 1:1 THF/H₂O, DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material.

Cleavage Protocol: To a 96-well plate containing Lanterns was added a 15% solution of HF/pyridine in stabilized THF (350 µL/Lantern). After 2 h the cleavage solution was quenched with TMSOMe (700 µL/Lantern) and the contents of each well were transferred to a pre-weighed 2-mL vial. The Lanterns were washed with an additional 200 µL of stabilized THF (or THF/MeOH) and the solution was transferred to the 2-mL vial. The samples were concentrated on a GENEVAC® solvent evaporation system overnight without heating to afford the title compound (4R,5R)-8-(cyclopent-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide (7.5 mg, 86% yield). LCMS RT 0.81 min, observed [M+1]⁺ 485.23, calculated [M+1]⁺ 485.245). Loading masses for each alcohol was determined on a FLEXIWEIGH® system.

Example 2: Synthesis of 1-(4-fluorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea



- 5 **Scaffold loading:** To a flame-dried flask containing silicon-functionalized Lanterns was added a freshly prepared solution of TfOH in anhydrous DCM (9.0 equiv, 5 g of TfOH/100 mL of DCM) was added. Each flask was shaken at RT for 10 min at which time the Lanterns had turned bright orange. The deep red TfOH solution was removed via cannula and anhydrous 2,6-lutidine (12.0 equiv relative to Si) was added. Once the
- 10 Lantern color had changed from orange to white, allyl (((2R,3R)-9-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-2-yl)methyl)(methyl)carbamate, the scaffold (1.2 equiv. relative to Si) was added as a solution in anhydrous DCM (0.4 mL/Lantern) and the reaction mixture was shaken for 48 h overnight. The loading mixture was removed and
- 15 set aside (to recover any unreacted alcohol) and the Lanterns were washed with the following solvents for 30 min intervals: DCM, THF, 3:1 THF/IPA, 3:1 THF/H₂O, DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. The Lanterns were then dried on a lyophilizer overnight and carried on to the next step.

- Fmoc removal:** To a flask containing Lanterns was added a solution of 20% piperidine in DMF (0.8mL/Lantern). After shaking at rt for 30 min, the piperidine solution was removed and the Lanterns were washed with the following solvents for 30 min intervals: DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material. The Lanterns were then dried on a lyophilizer overnight and carried on to the next step.

- 25 **N-Capping/Isocyanates:** To a flask containing Lanterns was added DCM (0.8 mL/Lantern) followed by 4-fluorophenyl isocyanate (15 equiv). The Lanterns were shaken at rt overnight and then washed with the following solvents for 30 min intervals:

DCM, DMF, 3:1 THF/ H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material (a stereoisomer is shown). The Lanterns were then dried on a lyophilizer overnight and carried on to the next step.

Alloc removal: To the reaction vessel containing Lanterns, THF (0.8 mL/Lantern) was added, followed by Pd(PPh₃)₄ (1 equiv) and 1,3-dimethylbarbituric acid (30 equiv). The flask was sealed and shaken at rt for 1 day. The reaction mixture was removed and the Lanterns were washed with DMF until the washings were clear (without any yellow color). Subsequently the Lanterns were washed with the following solvents for 30 min intervals: 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material. The Lanterns were then dried on a lyophilizer overnight and carried on to the next step.

N-Capping/Aldehydes: To a flask containing Lanterns was added DMF with 2% AcOH (0.800 mL/Lantern) followed by 4-pyridinecarboxaldehyde (20 equiv). The reaction mixture was shaken at rt for 1 hr then sodium triacetoxymethylborohydride (20 equiv) was added and shaking was continued. After 3 days the reaction mixture was removed and the Lanterns were washed with the following solvents for 30 min intervals: DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material.

Cleavage Protocol: To a 96-well plate containing Lanterns was added a 15% solution of HF/pyridine in stabilized THF (350 µL/Lantern). After 2 h the cleavage solution was quenched with TMSOMe (700 µL/Lantern) and the contents of each well were transferred to a pre-weighed 2-mL vial. The Lanterns were washed with an additional 200 µL of stabilized THF (or THF/MeOH) and the solution was transferred to the 2-mL vial. The samples were concentrated on a GENEVAC® solvent evaporation system overnight without heating to afford the title compound 1-(4-fluorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea (7.5 mg, 84% yield). LCMS RT 0.69 min, observed [M+1]⁺ 549.28, calculated [M+1]⁺ 549.28). Loading masses for each alcohol was determined on a FLEXIWEIGH® system.

30

Example 3: Synthesis of (4R,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-8-phenyl-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide

Scaffold loading: To a flame-dried flask containing silicon-functionalized Lanterns was added a freshly prepared solution of TfOH in anhydrous DCM (9.0 equiv, 5 g of TfOH/100 mL of DCM) was added. Each flask was shaken at RT for 10 min at which time the Lanterns had turned bright orange. The deep red TfOH solution was removed via cannula and anhydrous 2,6-lutidine (12.0 equiv relative to Si) was added. Once the Lantern color had changed from orange to white, (9H-fluoren-9-yl)methyl (((4R,5R)-8-bromo-2-((S)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)(methyl)carbamate, the scaffold (1.2 equiv. relative to Si) was added as a solution in anhydrous DCM (0.4 mL/Lantern) and the reaction mixture was shaken for 48h overnight. The loading mixture was removed and set aside (to recover any unreacted alcohol) and the Lanterns were washed with the following solvents for 30 min intervals: DCM, THF, 3:1 THF/IPA, 3:1 THF/H₂O, DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. The Lanterns were then dried on a lyophilizer overnight and carried on to the next step.

Fmoc removal: To a flask containing Lanterns was added a solution of 20% piperidine in DMF (0.8mL/Lantern). After shaking at rt for 30 min, the piperidine solution was removed and the Lanterns were washed with the following solvents for 30 min intervals: DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material. The Lanterns were then dried on a lyophilizer overnight and carried on to the next step.

N-Capping/Aldehydes: To a flask containing Lanterns was added DMF with 2% AcOH (0.800 mL/Lantern) followed by 4-pyridinecarboxaldehyde (20 equiv). The reaction mixture was shaken at rt for 1 hr then sodium triacetoxyborohydride (20 equiv) was added and shaking was continued. After 3 days the reaction mixture was removed and the Lanterns were washed with the following solvents for 30 min intervals: DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material. The Lanterns were then dried on a lyophilizer overnight and carried on to the next step.

Cross-Coupling/Suzuki: To each flask containing Lanterns was added ethanol (0.800 mL/Lantern) followed by phenylboronic acid (20 equiv), triethylamine (40 equiv) and

Pd(PPh₃)₂Cl₂ (1 equiv). The resulting mixture was degassed with a stream of N₂ before shaking at 60 °C. After 4 days, the reaction mixture was removed and the Lanterns were washed with following solvents for 30 min intervals: DCM, DMF, NaCN solution (0.1M) in 1:1 THF/H₂O, DMF, 3:1 THF/H₂O, 3:1 THF/IPA, THF, DCM. QC analysis of an aliquot cleaved from the lantern shows no remaining starting material.

Cleavage Protocol: To a 96-well plate containing Lanterns was added a 15% solution of HF/pyridine in stabilized THF (350 µL/Lantern). After 2 h the cleavage solution was quenched with TMSOMe (700 µL/Lantern) and the contents of each well were transferred to a pre-weighed 2-mL vial. The Lanterns were washed with an additional 200 µL of stabilized THF (or THF/MeOH) and the solution was transferred to the 2-mL vial. The samples were concentrated on a GENEVAC® solvent evaporation system overnight without heating to afford the title compound (4R,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-8-phenyl-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide (3.1 mg, 31% yield). LCMS RT 0.78 min, observed [M+1]⁺ 495.22, calculated [M+1]⁺ 495.22). Loading masses for each alcohol was determined on a FLEXIWEIGH® system.

Compounds 1-53 were synthesized following similar procedure as above. The synthetic characterization data (LCMS) for Compounds 1-53 is given in Table 2.

TABLE 2

Compound No.	Observed [M+1] ⁺	Expected (M+H) ⁺
1.	485.23	485.235
2.	549.28	549.275
3.	495.22	495.219
4.	553.26	553.261
5.	537.33	537.332
6.	527.25	527.245
7.	559.23	559.225
8.	483.22	483.219
9.	509.23	509.235
10.	485.23	485.235
11.	527.25	527.245
12.	499.25	499.25
13.	485.23	485.235
14.	522.32	522.321
15.	551.34	551.336
16.	509.23	509.235
17.	487.26	487.258
18.	568.26	568.261

19.	499.21	499.214
20.	527.25	527.245
21.	527.25	527.245
22.	498.30	498.308
23.	610.44	610.203
24.	562.51	562.303
25.	533.24	533.288
26.	524.3	524.324
27.	618.33	618.341
28.	484.25	484.293
29.	566.45	566.253
30.	552.72	552.355
31.	532.26	532.292
32.	582.56	582.308
33.	600.50	600.280
34.	576.53	576.282
35.	551.53	551.298
36.	481.52	481.281
37.	518.52	518.277
38.	550.36	550.283
39.	586.73	586.339
40.	557.48	557.288
41.	610.54	610.314
42.	615.59	615.366
43.	572.50	572.287
44.	556.53	556.330
45.	572.50	572.301
46.	539.40	539.335
47.	588.57	588.355
48.	527.31	527.335
49.	635.38	635.429
50.	658.41	658.397
51.	695.49	695.403
52.	522.26	522.344
53.	672.49	672.413

Example 4: A biochemical assay to identify small molecules that inhibit the neomorphic activity of purified IDH1-R132H was used to identify compounds of interest. C-terminally His8-tagged IDH1-R132H protein was expressed in the E. coli Rosetta strain and purified by metal chelate affinity chromatography. The ability of IDH1-R132H to convert α -KG to 2-HG using NADPH is assayed by measuring consumption of NADPH in a diaphorase-coupled assay. Resazurin is a dark blue reagent that has little intrinsic fluorescence. In the presence of NADPH, Resazurin is reduced by diaphorase to resorufin, which is highly fluorescent with an excitation peak at 579 nm and an emission peak at 584 nm (Figures 1A-G).

To validate the above assay, a set of compounds was screened in a multi-well format in duplicate. The average signal-to-background ratio was about 10; the Z' factor ranged from 0.7 to 0.85 and CV was <8% for each of the four plates. The scale-up was facile and robust, with an average Z' of about 0.74 for 101 plates, and yielding over
5 hundred hits with $\geq 30\%$ inhibition in both replicates. A more stringent cutoff of $\geq 40\%$ inhibition in both replicates, corresponding to >4 times the sum of the standard deviations of the negative and positive controls, yielded a smaller set.

To identify assay artifacts, such as compounds that activate diaphorase directly, another assay was developed as an alternative method to measure the activity of IDH1-R132H based on absorbance at 340 nm by NADPH (Figure 2A-B). The assay solution
10 contained 100 mM Tris.HCl pH 7.4, 150 mM NaCl, 5 mM MnCl₂, 0.03% BSA, 20 μ M NADPH, 0.4 mM α -Ketoglutarate and 87 nM of recombinant IDH1-R132H. Absorbance is measured on a SpectraMax spectrophotometer (Molecular Devices) in kinetic mode.

Compound Selectivity Test: To define their selectivity, confirmed inhibitors of IDH1-R132H were tested in dose for their ability to inhibit: (i) wild-type IDH1, (ii) another α -Ketoglutarate-dependent enzyme, the histone demethylase GASC1, and, (iii) another
15 NADP⁺-dependent dehydrogenase, G6PDH. Selectivity over the following criteria was studied:

- (i) wild-type IDH1. The ability of the compound to inhibit IDH1-R132H compared to
20 wild-type IDH1.
- (ii, iii) Selectivity over GASC1 and G6PDH. Ideal compounds should not inhibit either GASC1 or G6PDH (IC₅₀ difference > 20 -fold).

Step Description of assay conditions:

1. Dispense 5 nl of compounds to black 1536-well plates using Echo liquid handling
25 system.
2. Dispense 2.5 μ l of assay buffer to positive control wells of assay plates using Multidrop Combi dispenser.
3. Dispense 2.5 μ l of enzyme mix to assay wells using Multidrop Combi dispenser
4. Incubate at room temperature (about 20-25°C) for 20 min.
- 30 5. Dispense 2.5 μ l of substrate mix to initiate the reaction using Multidrop Combi dispenser.

6. Incubate at room temperature for 50 min. Pre-read to identify fluorescent compounds.
7. Dispense 2.5 μ l of detection mix using Multidrop Combi dispenser.
8. Incubate at room temperature for 30 min.
- 5 9. Read plates on Envision (Ex 535, Em 595).


The activity of Compounds to inhibit IDH1 (wild type) and IDH1-R132H is given in Table 3.

TABLE 3



Example No.	IC ₅₀ IDH1 (R132H)	IC ₅₀ IDH1 (Wild Type)	GASC1	G6PDH
1	I	I	ND	IV
2	I	IV	IV	IV
3	I	IV	IV	IV
4	I	IV	ND	IV
5	I	IV	ND	IV
6	I	IV	ND	IV
7	I	IV	ND	IV
8	I	IV	ND	IV
9	I	IV	IV	IV
10	I	IV	ND	IV
11	III	III	IV	ND
12	II	IV	IV	IV
13	III	II	ND	IV
14	III	IV	IV	IV
15	II	III	II	IV
16	II	III	IV	IV
17	III	IV	I	IV
18	III	III	ND	ND
19	II	III	ND	ND
20	II	IV	IV	IV
21	III	III	IV	IV
22	III	IV	ND	IV

23	II		ND	ND
24	I		ND	ND
25	I		ND	ND
26	I		ND	ND
27	I		ND	ND
28	III		ND	ND
29	II		ND	ND
30	I		ND	ND
31	I		ND	ND
32	III		ND	ND
33	II		ND	ND
34	I		ND	ND
35	III		ND	ND
36	II		ND	ND
37	I		ND	ND
38	I		ND	ND
39	I		ND	ND
40	III		ND	ND
41	II		ND	ND
42	II		ND	ND
43	II		ND	ND
44	II		ND	ND
45	I		ND	ND
46	I		ND	ND
47	III		ND	ND
48	I		ND	ND
49	I		ND	ND
50	I		ND	ND
51	I		ND	ND
52	II		ND	ND
53	I		ND	ND

Protocol for the synthesis of Example-52




Example-52

5 Step 1: Synthesis of **b**

Compound **a** was synthesized according to published procedure (Marcaurelle et al. *J. Am. Chem. Soc.* 2010, 132, 16962-16976). The remaining reagents for the following reaction are commercially available. To an oven-dried round-bottom flask equipped with a magnetic stirrer was added **a** (945 mg, 2.61 mmol, 1.0 equiv) and 10 mL
 5 dichloromethane. PyBOP (1.50 g, 2.88 mmol, 1.1 equiv) and diisopropyl ethylamine (1.01 g, 7.84 mmol, 3.0 equiv) were added. The resulting mixture was cooled in an ice bath before isopropylamine (185 mg, 3.14 mmol, 1.2 equiv) was added as a solution in dichloromethane (3 mL) dropwise over 20 minutes. The mixture was stirred for three days
 10 at room temperature. The reaction was quenched with water and extracted with dichloromethane. The combined organic extracts were dried over magnesium sulfate, filtered, and concentrated to yield a white solid. The soluble portions of the mixture were taken up in diethyl ether and the insoluble phosphoramidate byproducts were removed *via* filtration. The solvent was removed *in vacuo* and the crude product was isolated. Flash
 15 chromatography on silica gel (0-50% Ethyl acetate in hexanes) gave the product (540 mg, 51%).


Step 2: Synthesis of **c**



To an oven-dried 2-necked round-bottom flask equipped with a magnetic stirbar and condenser was added **b** (530 mg, 1.32 mmol, 1 equiv) in 15 mL tetrahydrofuran (final
 20 concentration 0.05 M). Borane-dimethylsulfide complex (500 mg, 5.68 mmol, 5.0 equiv) was added dropwise via syringe and the reaction was heated at 65 °C for 5 hr. The flask was placed on an ice bath and excess hydride was quenched by addition of methanol until bubbling stopped. The mixture was concentrated to an oil and evaporated three times with
 25 methanol to remove excess $\text{B}(\text{OMe})_3$. The oil was redissolved in a mixture of aqueous sodium potassium tartrate (10 mL, 0.5 M) and methanol (10 mL) and the resulting slurry was heated at reflux (85 °C) for 12 hours to disrupt the boron-nitrogen complex. Volatiles were removed under reduced pressure and the resulting aqueous mixture was extracted


with ethyl acetate. The combined organic extracts were washed once with brine, dried over magnesium sulfate, filtered, and concentrated to provide the desired amine **c** as a colorless oil in quantitative crude yield (527.6 mg). The material was taken on without purification.

5 Step 3: Synthesis of **d**



To a flame-dried round-bottom flask equipped with stirbar and purged with nitrogen was added **c** (427 mg, 1.10 mmol, 1.0 equiv) in dichloromethane (final volume 15 mL, concentration 0.04 M). The solution cooled in an ice bath and triethylamine (556 mg, 5.49 mmol, 5.0 equiv) and 2-fluoro-5-nitrophenylacetic acid chloride (598 mg, 2.75 mmol, 2.5 equiv)(prepared as described in Marcuarelle et. al. *J. Am. Chem. Soc.* 2010, 132, 16962-16976) were added via syringe. The reaction mixture took on a dark red-orange color. The vessel was warmed to room temperature and allowed to stir overnight. The reaction was quenched with water and extracted with dichloromethane. The combined
 10 organic extracts were dried over magnesium sulfate and concentrated to yield the crude product. The material was purified by flash chromatography on silica gel (0-50% EtOAc in hexanes) to give the product **d** (526 mg, 84%).


Step 4: Synthesis of **e**



To a flame-dried 50 mL flask that had been purged with nitrogen was added ammonium fluoride (171 mg, 4.61 mmol, 5.0 equiv) followed by **d** (526 mg, 0.922 mmol, 1.0 equiv) in tetrahydrofuran (14 mL, final concentration 0.07 M) and tetrabutylammonium fluoride (1.21 g, 4.61 mmol, 5.0 equiv). The mixture was stirred at room temperature overnight. The reaction was quenched with saturated aqueous ammonium chloride. The organic and aqueous layers were separated and the aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with acetic acid solution (1.0 M in water) and brine and dried over magnesium sulfate. The mixture was filtered and the filtrate was concentrated. Flash chromatography on silica gel (20-60% EtOAc in hexanes) yielded the deprotected alcohol (215 mg). This intermediate was dissolved in THF (15 mL, final concentration 0.03 M) and placed in a flame-dried round-bottom flask with stirbar that had been purged with nitrogen. Sodium hydride (94 mg, 2.36 mmol, 5.0 equiv) was added in one portion and the resulting mixture was allowed to stir overnight at room temperature. The reaction was quenched with not-quite-saturated aqueous ammonium chloride solution. The organic and aqueous layers were separated and the aqueous layer was extracted with ethyl acetate. The combined organic extracts were dried over magnesium sulfate and filtered. The filtrate was concentrated to yield the crude product. Flash chromatography on silica gel (50-70% EtOAc in hexanes) afforded the product as a white crystalline solid (122 mg, 30%).

¹H NMR (300 MHz, CDCl₃, 22 °C) δ 8.18 (d, *J* = 3 Hz, 1H), 8.06 (dd, *J* = 3, 9 Hz, 1H), 7.09 (d, *J* = 9 Hz, 1H), 4.56 (m, 1H), 4.13 (m, 1H), 3.64 (m, 2H), 3.47 (m, 3H), 2.92 (s, 3H), 2.17 (m, 1H), 1.70 (m, 1H), 1.40 (s, 9H), 1.22 (m, 6H), 0.98 (d, *J* = 7 Hz, 3H). ¹³C NMR (300 MHz, CDCl₃, 22 °C) δ 162.8, 144.0, 131.4, 127.7, 123.9, 123.2, 80.2, 52.3, 51.5, 37.4, 36.2, 28.3, 21.2, 19.8, 15.8.


Step 5: Synthesis of **f**



To a flame-dried flask equipped with a stirbar was added 10% palladium on carbon (28.2 mg, 26 μmol, 0.1 equiv), followed by purging of the flask with nitrogen. **e** (115 mg,


256 μmol , 1.0 equiv) was added as a solution in ethanol (11 mL, final concentration 0.025 M). The mixture was heated to 35 $^{\circ}\text{C}$ and a hydrogen atmosphere was applied via balloon. The reaction was allowed to progress for two hours, after which the mixture was removed from heat and filtered through Celite and the solvent was removed *in vacuo* to afford the crude product (95 mg, crude yield 88%). The material was taken on crude.

Step 6: Synthesis of **g**



To a flame-dried 5 mL flask with stirbar that had been purged with nitrogen was added **f** (9.2 mg, 23 μmol , 1.0 equiv) and 500 μL dichloromethane. Cyclohexyl isocyanate (12.8 mg, 102 μmol , 4.5 equiv) was added and the mixture was stirred for four days. Extra dichloromethane was periodically added to replace that which evaporated over the course of the reaction. The reaction was quenched with water and extracted with methylene chloride. The combined organic extracts were dried over sodium sulfate; the solution was decanted and concentrated *in vacuo* to afford crude product. Flash chromatography on silica gel (0-10% MeOH in dichloromethane) provided the product (10.4 mg, 86%).


Step 7: Synthesis of **h**





To a small vial equipped with a stirbar was added **g** (10.4 mg, 20 μmol , 1.0 equiv) in dichloromethane (600 μL , 0.02 M). Trifluoroacetic acid (444 mg, 3.89 mmol, 200 equiv) was added and the mixture was stirred for 30 minutes. The mixture was then concentrated *in vacuo* and mixed with saturated aqueous sodium bicarbonate. The resulting aqueous mixture was extracted with dichloromethane; the combined organic

extracts were dried over sodium sulfate. The solution was decanted and the solvent was removed by evaporation under a stream of dry nitrogen. The material was taken on crude.

Step 8: Synthesis of Example-52



- 5 To a vial equipped with a stirbar was added **9** (9.6 mg, 22 μ mol, 1.0 equiv) dissolved in 2% acetic acid in dimethylformamide (612 μ L, 0.036 M). Isonicotinaldehyde (5.7 mg, 53 μ mol, 2.4 equiv) was added via syringe and the reaction was allowed to proceed for 90 minutes. Sodium triacetoxyborohydride (9.0 mg, 42 μ mol, 1.9 equiv) was added and the mixture was stirred for two days. The solvent was removed *in vacuo* at
- 10 50°C and the residue was taken up in saturated aqueous sodium bicarbonate solution. The aqueous mixture was extracted with dichloromethane; the combined organic extracts were dried over sodium sulfate. The solution was decanted and concentrated *in vacuo*. The residue was purified by high-pressure liquid chromatography to yield the product (4.9 mg, 42%).
- 15 ¹H NMR (300 MHz, CDCl₃, 22 °C) δ 7.43 (m, 2H), 7.04 (m, 2H), 6.94 (m, 1H), 5.45 (m, 1H), 4.78 (m, 1H), 4.17 (m, 1H), 3.75 (m, 1H), 3.51 (m, 2H), 3.34 (m, 1H), 3.11 (m, 1H), 2.79 (m, 1H), 2.59 (m, 1H), 2.41 (m, 1H), 2.22 (s, 3H), 1.88 (m, 5H), 1.56 (m, 6H), 1.23 (m, 6H), 0.94 (m, 3H).

Synthesis of Example -50**Step 1 and 2 - Reduction and N-capping with isocyanates**

- 5 Lactam **i** (1 equiv.) and palladium (10% on activated carbon, 0.1 equiv.) were stirred in EtOH (25 mM) at 35 °C under a hydrogen atmosphere. The reaction was monitored by LCMS for complete consumption of starting material (1-2h). The mixture was cooled, filtered through Celite and concentrated to give the aniline derivative **j**.


- The crude product was dried under vacuum and then dissolved in DCM (150 mM) and transferred into a dried round-bottom flask flushed with nitrogen. Cyclohexyl isocyanate (1.5 equiv.) was added with a syringe. It was stirred overnight at room temperature. The mixture was concentrated and the product was isolated by column chromatography (DCM/MeOH gradient from 0 to 20% MeOH). Yield: 96%.

- $[\alpha]_D^{20} = -2.3^\circ$ (c 1.0, CHCl₃). IR (cm⁻¹) = 3346, 2927, 2853, 2361, 2342, 1692, 1611, 1551, 1497, 1451, 1393, 1365, 1249, 1217, 1149, 1086, 819. ¹H NMR (300 MHz, CDCl₃-d) δ = 7.34 (s, 1H), 7.24-7.16 (m, 3H), 6.86 (d, J = 8 Hz, 2H), 6.67 (m, 1H), 5.52 (s, 1H), 4.44 (s, 2H), 4.15 (m, 1H), 3.81-3.79 (m, 6H), 3.61-3.59 (m, 4H), 3.36-3.31 (m, 2H), 2.99-2.90 (m, 3 H), 2.61 (s, 3H), 2.09 (m, 1H), 1.90 (m, 2H), 1.69-1.55 (m, 4H), 1.43

(s, 9H), 1.35 (d, $J = 7$ Hz, 4H) 1.25 (s, 3H), 0.87-0.83 (m, 3H). ^{13}C NMR (75 MHz, $\text{CDCl}_3\text{-d}$) $\delta = 172.3, 159.5, 155.7, 152.5, 136.4, 130.7, 129.6, 124.4, 123.02, 120.0, 114.1, 80.2, 77.6, 73.2, 72.8, 59.2, 55.6, 48.8, 41.4, 38.0, 36.0, 34.1, 34.0, 30.0, 28.8, 26.1, 25.3, 25.2, 23.0, 15.4$. HRMS (ESI) calcd for $\text{C}_{37}\text{H}_{54}\text{N}_4\text{O}_7$ $[\text{M}+\text{H}]^+$: 667.3993. Found: 667.843.

5

Step 3 - Boc removal




2,6-Lutidine (1,26 mL, 10.8 mmol, 4 equiv.) and TBSOTf (1.86 mL, 8.1 mmol, 3 equiv.) were added to a solution of **j** (1.8 g, 2.7 mmol, 1 equiv.) in DCM (27 mL, 0.1 M) at room temperature and stirred overnight. The reaction was quenched with saturated NH_4Cl solution and extracted with EtOAc. The combined organic extracts were dried over MgSO_4 , filtered, and concentrated to the crude silyl carbamate.


The resulting oil was dissolved in THF (22.5 mL, 1.2 M) and subsequently HF/pyridine (70% HF; 360 μL , 1 equiv. HF) was added. The mixture was stirred for 30 min, quenched with saturated NH_4Cl and extracted with EtOAc. The combined organic extracts were dried over MgSO_4 , filtered, and concentrated. Product **k** was used without further purification. Yield: 98%.

$[\alpha]_{\text{D}}^{20} = -25.7^{\circ}$ (c 1.0, CHCl_3). IR (cm^{-1}) 3342, 2927, 2853, 2361, 1610, 1551, 1513, 1498, 1249, 1218, 1085, 1034, 820. ^1H NMR (300 MHz, $\text{CDCl}_3\text{-d}$) $\delta = 7.55$ (s, 1H), 7.26-7.24 (m, 3H), 6.87 (d, $J = 8$ Hz, 2H) 6.44 (m, 1H), 5.76 (m, 1H), 4.46 (s, 3H), 4.38 (m, 1H), 4.15 (d, $J = 14$ Hz, 1H), 3.86 (m, 5H), 3.59-3.30 (m, 4H), 3.09 (d, $J = 13$ Hz, 1H), 2.81 (d, $J = 11$ Hz, 1H), 2.58-2.35 (m, 6H), 1.91-1.54 (m, 6H), 1.34-0.84 (m, 10H). ^{13}C NMR (75 MHz, $\text{CDCl}_3\text{-d}$) $\delta = 172.7, 159.5, 155.9, 151.4, 136.5, 130.8, 130.0, 129.6, 125.0, 123.2, 119.7, 114.1, 89.0, 73.1, 72.7, 55.6, 54.0, 51.9, 48.6, 37.6, 37.4, 34.1, 33.2, 30.0, 26.1, 25.3, 15.6$. HRMS (ESI) calcd for $\text{C}_{32}\text{H}_{46}\text{N}_4\text{O}_5$ $[\text{M}+\text{H}]^+$: 567.3468. Found: 567.4246.

Step 4 - Reductive amination




Example-50




Example-50

Starting material **k** (0.7 g, 1.24 mmol, 1 equiv.) was diluted in DMF (containing
 5 2% Acetic acid; 88 mM). Isonicotinic aldehyd (0.128 mL, 1.36 mmol, 1.1 equiv.) was
 added and the mixture was stirred for 1h at room temperature. Subsequently sodium
 triacetoxyborohydride (340 mg, 1.61 mmol, 1.3 equiv.) was added and the mixture was
 stirred for 3 days. The solvents were evaporated *in vacuo* and product Example-50 was
 purified by column chromatography (DCM/MeOH gradient from 0 to 20% MeOH). Yield:
 10 93%.

$[\alpha]_D^{20} = -6.2^0$ (c 1.0, CHCl₃). IR (cm⁻¹) 3349, 2929, 2852, 1604, 1551, 1497, 1451,
 1414, 1248, 1217, 1084, 820. ¹H NMR (300 MHz, CDCl₃-d) δ = 8.51 (d, J = 5 Hz, 2H),
 7.53 (s, 1H), 7.22-7.14 (m, 4H), 6.85 (d, J = 8 Hz, 2H), 6.68 (dd, J₁ = 33 Hz, J₂ = 8 Hz,
 2H), 5.69 (d, J = 8 Hz, 1H), 4.43-4.34 (m, 3H), 4.14 (d, J = 13 Hz, 1H), 3.89-3.29 (m,
 15 12H), 3.06 (d, J = 11 Hz, 1H), 2.72 (d, J = 13 Hz, 1H), 2.53 (d, J = 13 Hz, 1H), 2.39 (m,
 1H), 2.19 (s, 3H), 1.90-1.53 (m, 6H), 1.33-0.88 (m, 10H). ¹³C NMR (75 MHz, CDCl₃-d) δ
 = 172.6, 159.5, 155.9, 151.8, 150.1, 149.0, 136.6, 130.7, 130.0, 129.5, 124.6, 124.0, 123.6,
 119.5, 114.2, 89.4, 73.1, 72.8, 62.6, 59.2, 55.6, 53.3, 48.7, 44.2, 37.5, 34.6, 34.1, 26.1,
 25.2, 15.9, 15.6. HRMS (ESI) calcd for C₃₈H₅₁N₅O₅ [M+H]⁺: 658.3890. Found: 658.3772.



To a solution of BRD5667 in DMF (10 mg in 0.372 mL, 50 mM) at room temperature was added sodium hydride (1.3 mg, 0.055 mmol). The mixture was stirred for 10 minutes and 4-(2-chloroethyl)pyrrolidine hydrochloride (4.4 mg, 0.026 mmol) was added to the reaction mixture. The mixture was stirred at 40 °C for 24 h. Yield after HPLC purification: 0.494 mg.



BRD5667 was distilled azeotropically three times with 20% dichloromethane in benzene and lyophilized overnight.

To a solution of BRD5667 in anhydrous tetrahydrofuran (83.3 mg in 3.87 mL, 40 mM) under argon was added DBU (0.093 mL, 0.62 mmol) followed by diphenyl phosphorazidate (0.1 mL, 0.465 mmol). The mixture was stirred for 24 h. The solvent was removed under vacuum and the azide was purified by column chromatography (0-20% MeOH in DCM). Yield after purification: 10 mg, 11%.

The azide (10mg, 0.018 mmol) was diluted in tetrahydrofuran (0.35 mL, 50 mM) in a dried LCMS vial. Triethylamine (10 uL, 0.07 mmol), the alkyne (7 uL, 0.05 mmol), and copper(I) iodide (0.34 mg, 0.0018 mmol) were added. It was stirred overnight at room temperature. After filtration, the solvent was removed and the product was purified by HPLC. Yield after HPLC purification: 0.674 mg.

The patent and scientific literature referred to herein establishes the knowledge that is available to those with skill in the art. All United States patents and published or unpublished United States patent applications cited herein are incorporated by reference.


- 5 All published foreign patents and patent applications cited herein are hereby incorporated by reference. All other published references, documents, manuscripts and scientific literature cited herein are hereby incorporated by reference.

- While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various
10 changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.


CLAIMS

What is claimed is:


1. A compound of Formula IA, IB, IC or ID, or a pharmaceutically acceptable salt, ester or prodrug thereof:




Formula IA



Formula IB



10 Formula IC



Formula ID

wherein each n and m is independently 0, 1, 2 or 3;

each p and q is independently 0, 1, 2, 3, 4, 5, 6 or 7;

5 X_1 is $-C(O)N(R_A)-$ or $-C(S)N(R_A)-$;

wherein R_A is independently hydrogen, aliphatic, substituted aliphatic, heteroaryl, substituted heteroaryl, aryl or substituted aryl;

X_2 is $-S-$, $-O-$, $-S(O)_2-$, $-C(R_{20})(R_{21})-$ or $-N(R_B)-$;

10 wherein R_B is independently hydrogen, aliphatic, substituted aliphatic, heteroaryl, substituted heteroaryl, aryl or substituted aryl;

each R_1 and R_2 is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl;

R_3 is hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl;

15 each R_{10} , and R_{28} is independently absent, hydrogen, halogen, $-OR_{20}$, $-SR_{20}$, $-NR_{20}R_{21}$, $-CF_3$, $-CN$, $-NO_2$, $-N_3$, $-C(O)OR_{20}$, $-C(O)R_{20}$, $-C(O)N R_{20}R_{21}$, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two of R_{10} groups together with the atoms to which they are attached
20 and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring;

wherein each R_{20} and R_{21} is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl;

each R_{11} is independently absent, hydrogen, halogen, $-OR_{20}$, $-SR_{20}$, $-NR_{20}R_{21}$, $-CF_3$, $-CN$, $-NO_2$, $-N_3$, $-C(O)R_{20}$, $-C(O)OR_{20}$, $-C(O)NR_{20}R_{21}$, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two of R_{11} groups together with the atoms to which they are attached and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring;


each G_1 and G_2 is independently, absent, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl or substituted alkynyl, preferably $-CH_2-$;

each R_{27} and R_{29} is independently hydrogen, alkyl, substituted alkyl, aryl or substituted aryl, preferably hydrogen;

wherein R_{3a} is selected from alkyl, aryl, alkyl substituted with aryl, straight chain or branched C_1 - C_{10} alkyl, C_1 - C_{10} alkenyl, C_1 - C_{10} alkynyl, C_1 - C_{10} alkoxy, alkoxy C_1 - C_{10} alkoxy, C_1 - C_{10} alkylamino, alkoxy C_1 - C_{10} alkylamino, C_1 - C_{10} alkylcarbonylamino, C_1 - C_{10} alkylaminocarbonyl, aryloxy C_1 - C_{10} alkoxy, aryloxy C_1 - C_{10} alkylamino, aryloxy C_1 - C_{10} alkylamino carbonyl, C_1 - C_{10} -alkylaminoalkylaminocarbonyl, C_1 - C_{10} alkyl(N-alkyl)aminoalkyl-aminocarbonyl, alkylaminoalkylamino, alkylcarbonylaminoalkylamino, alkyl(N-alkyl)aminoalkylamino, (N-alkyl)alkylcarbonylaminoalkylamino, alkylaminoalkyl, alkylaminoalkylaminoalkyl, alkylpiperazinoalkyl, piperazinoalkyl, alkylpiperazino, alkenylaryloxy C_1 - C_{10} alkoxy, alkenylarylaminoc C_1 - C_{10} alkoxy, alkenylaryllalkylamino C_1 - C_{10} alkoxy, alkenylaryloxy C_1 - C_{10} alkylamino, alkenylaryloxy C_1 - C_{10} alkylaminocarbonyl, piperazinoalkylaryl, heteroaryl C_1 - C_{10} alkyl, heteroaryl C_1 - C_{10} alkenyl, heteroaryl C_1 - C_{10} alkynyl, heteroaryl C_1 - C_{10} alkylamino, heteroaryl C_1 - C_{10} alkoxy, heteroaryloxy C_1 - C_{10} alkyl, heteroaryloxy C_1 - C_{10} alkenyl, heteroaryloxy C_2 - C_{10} alkynyl, heteroaryloxy C_1 - C_{10} alkylamino, heteroaryloxy C_1 - C_{10} alkoxy;

Cy1 is an optionally substituted aryl or optionally substituted heteroaryl; and,


R_{40} is selected from:




- 5 wherein, R_{41} is hydrogen, halogen, aliphatic, substituted aliphatic, aryl, substituted aryl, $-OR_{20}$, $-SR_{20}$, $-NR_{20}R_{21}$, $-CF_3$, $-CN$, $-NO_2$, $-N_3$, $-C(O)R_{20}$, $-C(O)OR_{20}$, $-C(O)NR_{20}R_{21}$, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl.

10

2. A compound of claim 1 having Formula IIA-III A or a pharmaceutically acceptable salt, ester or prodrug thereof:




Formula IIA




Formula IIIA

wherein, n , p , q , X_2 , $Cy1$, R_1 , R_2 , R_3 , R_{3a} , R_{10} , R_{11} , and R_{40} are as defined above.

- 5 3. A compound of claim 1 having Formula IVA or VA a pharmaceutically acceptable salt, ester or prodrug thereof:



Formula IVA



10 Formula VA


wherein p , q , X_2 , R_1 , R_2 , R_3 , R_{3a} , R_{10} , and R_{11} are as defined above.

4. The compound according to any of the above claims, wherein R_1 is an optionally substituted alkyl.
5. The compound according to any of the above claims, wherein R_1 is an optionally substituted C_1 - C_6 alkyl, preferably selected from methyl, ethyl, propyl, cyclopropyl, isopropyl, *n*-butyl, *tert*-butyl, cyclobutyl, *n*-pentyl, neopentyl, cyclopentyl, *n*-hexyl and cyclohexyl.
6. A compound according to any of the above claims, wherein Cy1 is selected from Table

10

A:

TABLE A





wherein  represents a single or double bond;

- each R_{12} is independently absent, hydrogen, $-C(O)R_{20}$, $-C(O)OR_{20}$, $-C(O)NR_{20}R_{21}$, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl substituted aryl, heteroaryl or substituted heteroaryl;
- 5 d is 0, 1 or 2;
- alternatively, two R_{11} and R_{12} groups may form an optionally substituted 3, 4, 5, 6, or 7 membered ring.

- 10 7. A compound according to any of the above claims, wherein R_{11} is selected from Table B:

TABLE B





wherein, t is 0, 1, 2, 3, 4, 5, 6 or 7;

Cy2 is an optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocyclyl, optionally substituted aryl or optionally substituted heteroaryl,

- 5 each R_{13} , R_{14} and R_{15} is independently absent, hydrogen, halogen, $-OR_{20}$, $-SR_{20}$, $-NR_{20}R_{21}$, $-CF_3$, $-CN$, $-NO_2$, $-N_3$, $-C(O)R_{20}$, $-C(O)OR_{20}$, $-C(O)NR_{20}R_{21}$, acyl, alkoxy,

substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two R_{13} groups together with the atoms to which they are attached and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring; and,

each R_{16} , R_{17} and R_{18} is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl, substituted aryl, heteroaryl or substituted heteroaryl; alternatively two R_{13} , R_{14} , R_{15} , R_{16} , R_{17} and R_{18} groups together with the atoms to which they are attached, and any intervening atoms may form an optionally substituted 3, 4, 5, 6 or 7 membered ring; and,

alternatively when two R_{13} , R_{14} , R_{15} , R_{16} , R_{17} and R_{18} groups are attached to a carbon atom, together said two groups and said carbon atom may form a carbonyl or an optionally substituted vinyl group.


15

8. A compound according to any of the above claims, wherein q is 1.

9. A compound according to any of the above claims, wherein R_2 is selected from Table C:

20

TABLE C



wherein R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{17} and R_{18} are as defined above;

alternatively, R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{17} and R_{18} groups together with the atoms to which they are attached, and any intervening atoms may form an optionally substituted 3, 4, 5, 6 or 7 membered ring;

- 5 alternatively when two R_{13} , R_{14} , R_{15} , R_{16} , R_{17} and R_{18} groups are attached to a carbon atom, together said two groups and said carbon atom may form a carbonyl or an optionally substituted vinyl group;

u and w is independently 0, 1, 2, 3, 4, 5 or 6; and,

- 10 Cy3 is optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted heterocyclyl, optionally substituted aryl or optionally substituted heteroaryl.

10. A compound according to any of the above claims, wherein R_3 is selected from Table D, or R_{3a} selected from Table E:

- 15 TABLE D





TABLE E











11. A compound selected from Table 1A or a pharmaceutically acceptable salt, ester or prodrug thereof:





5





TABLE 1A





1	(4R,5R)-8-(cyclopent-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
2	1-(4-fluorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea





4	 N-(((4R,5R)-8-(cyclohexylethynyl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-4-yl)acetamide
6	 N-(((4R,5R)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-8-(4-methylpent-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide
7	 N-((2S,3S)-5-((R)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-3,4,5,6-tetrahydro-2H-benzo[b][1,5]oxazocin-10-yl)benzo[d]thiazole-2-carboxamide
8	 (4R,5R)-8-(cyclopropylethynyl)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide





9	 (4R,5S)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-4-ylmethyl)amino)methyl)-8-(p-tolyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
10	 (4S,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-8-(pent-1-yn-1-yl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
11	 N-(((4R,5S)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-8-(4-methylpent-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide
12	 (4R,5S)-8-(cyclohex-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide




13	 (4S,5S)-8-(cyclopent-1-en-1-yl)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-4-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
14	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)cyclohexanecarboxamide
16	 (4S,5S)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-4-ylmethyl)amino)methyl)-8-(o-tolyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
17	 (2S,3S)-5-((R)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyrimidin-5-ylmethyl)amino)methyl)-8-((E)-styryl)-4,5-dihydro-2H-pyrido[2,3-b][1,5]oxazocin-6(3H)-one




18	 4-((((4R,5R)-8-(3-cyclopentylprop-1-yn-1-yl)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)(methyl)amino)methyl)benzoic acid
19	 N-(((4R,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-8-(3-methylbut-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methylnicotinamide
20	 N-(((4R,5R)-8-(cyclohex-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide
21	 N-(((4S,5S)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-8-(4-methylpent-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide





23	 1-(4-bromophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
24	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(4-methoxyphenyl)urea
25	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(pyridin-3-yl)urea
26	 1-cyclopentyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea




27	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(2-morpholinopyridin-4-yl)urea
28	 1-ethyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
29	 1-(4-chlorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
30	 1-(cyclohexylmethyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea



32	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(naphthalen-1-yl)urea
34	 1-(benzo[d][1,3]dioxol-5-yl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
37	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)isonicotinamide
38	 1-(2-fluorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea

39	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(1,2,3,4-tetrahydronaphthalen-1-yl)urea
40	 1-(4-cyanophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
41	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(3-(pyrimidin-2-yl)phenyl)urea


42	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(2-(piperidin-1-yl)phenyl)urea
43	 1-(benzofuran-5-yl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
44	 1-cyclohexyl-3-((2R,3R)-2-(((3-fluoropyridin-4-yl)methyl)(methyl)amino)methyl)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea

45	 1-((2R,3R)-2-(((3-chloropyridin-4-yl)methyl)(methyl)amino)methyl)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-cyclohexylurea
46	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyrimidin-4-yl)methyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
47	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(quinolin-4-yl)methyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
48	 1-((2R,3R)-2-(((1H-imidazol-2-yl)methyl)(methyl)amino)methyl)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-cyclohexylurea

49	 1-cyclohexyl-3-((2R,3R)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-5-((S)-1-(2-(pyrrolidin-1-yl)ethoxy)propan-2-yl)-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
50	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-((4-methoxybenzyl)oxy)propan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
51	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-(4-(4-methoxyphenyl)-1H-1,2,3-triazol-1-yl)propan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea

52	 1-cyclohexyl-3-((2R,3R)-5-isopropyl-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
53	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-((4-methoxybenzyl)oxy)propan-2-yl)-3-methyl-2-((methyl((3-methylpyridin-4-yl)methyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea

12. A method of treating a disease related to a defect in isocitrate dehydrogenase comprising the step of administering a compound of Formula I to a patient in need thereof:



Formula I

each n and m is independently 0, 1, 2 or 3;

each p and q is independently 0, 1, 2, 3, 4, 5, 6 or 7;

X_1 is $-C(O)N(R_A)-$, $-C(S)N(R_A)-$, or $-S(O)_2N(R_A)-$;

5 wherein R_A is hydrogen, aliphatic, substituted aliphatic, heteroaryl, substituted heteroaryl, aryl or substituted aryl;

X_2 is $-S-$, $-O-$, $-S(O)_2-$, $-C(R_{20})(R_{21})-$ or $-N(R_B)-$;

wherein R_B is hydrogen, aliphatic, substituted aliphatic, heteroaryl, substituted heteroaryl, aryl or substituted aryl;

10 each R_1 and R_2 is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl;

each R_{10} is independently absent, hydrogen, halogen, $-OR_{20}$, $-SR_{20}$, $-NR_{20}R_{21}$, $-CF_3$, $-CN$, $-NO_2$, $-N_3$, $-C(O)OR_{20}$, $-C(O)R_{20}$, $-C(O)N R_{20}R_{21}$, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, 15 aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two of R_{10} groups together with the atoms to which they are attached and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring;

wherein each R_{20} and R_{21} is independently hydrogen, halogen, aliphatic, substituted aliphatic, aryl or substituted aryl;

20 each R_{11} is independently absent, hydrogen, halogen, $-OR_{20}$, $-S R_{20}$, $-NR_{20}R_{21}$, $-CF_3$, $-CN$, $-NO_2$, $-N_3$, $-C(O)OR_{20}$, $-C(O)R_{20}$, $-C(O)N R_{20}R_{21}$, acyl, alkoxy, substituted alkoxy, alkylamino, substituted alkylamino, dialkylamino, substituted dialkylamino, substituted or unsubstituted alkylthio, substituted or unsubstituted alkylsulfonyl, aliphatic, substituted aliphatic, aryl or substituted aryl; alternatively two of R_{11} groups 25 together with the atoms to which they are attached and any intervening atoms may form an additional optionally substituted, 3, 4, 5, 6 or 7 membered ring; and,


$Cy1$ is an optionally substituted aryl or optionally substituted heteroaryl.





13. The method according to claim 12, wherein said defect in isocitrate dehydrogenase is a somatic mutation at codon 132 isocitrate dehydrogenase (IDH1) or at codon 172 in isocitrate dehydrogenase 2 (IDH2) or at codon 140 of IDH2.
- 5 14. The method according to claim 12, wherein said patient has mutations on both IDH1 and IDH2.
15. The method according to claim 12, wherein said mutation is selected from R132H, R132C, R132S, R132L, R132G in IDH1, or R172M, R172G, R172K or R140Q in
10 IDH2.
16. The method according to any of claims 12-15, wherein said disease is a cell proliferative disease.
- 15 17. The method according to any of claims 12-16, wherein said disease is selected from Grade I, II, III or IV glioma.
18. The method according to any of claims 12-17, wherein said disease is selected from astrocytomas, oligodendrogliomas, ependymomas and glioblastoma multiforme
20 (GBM).
19. A method of treating a patient exhibiting abnormal 2-hydroxyglutarate (2-HG) production comprising administering a compound according to any of claims 1-11.
- 25 20. The method according to claim 19, wherein said abnormal 2-HG production is an increase of more than about 20 fold, or more than about 40 fold or more than about 50 fold or more than about 100 fold or more than about 200 fold compared to a corresponding normal cell or tissue or plasma concentration of 2-HG.





21. The method according to claim 20, wherein said increase is due to a mutation in IDH1.
22. The method according to claim 20, wherein said increase is due to a mutation in IDH2.
- 5 23. A method of treating a disease related to a defect in isocitrate dehydrogenase comprising the step of administering a compound according to any of claims 1-11 to a patient in need thereof, wherein said compound has a selectivity for inhibition of a mutant of IDH1 or a mutant of IDH2.
- 10 24. The method according to claim 23, wherein the ratio of inhibitory activity against an IDH1 mutant over the wild type IDH1 is about 2 to about 1000, preferably about 5 to about 500, preferably about 10 to about 100, preferably about 25 to about 100.
- 15 25. The method according to claim 23, wherein the ratio of inhibitory activity against an IDH2 mutant over the wild type IDH2 is about 2 to about 1000, preferably about 5 to about 500, preferably about 10 to about 100, preferably about 25 to about 100.
- 20 26. A method of treating a cell proliferative disease comprising the step of administering a compound according any of claims 1-11 to a patient in need thereof.
27. The method according to claim 26, wherein said cell proliferative disease is cancer.
28. The method according to claim 27, wherein said cancer is selected from glioma, acute myeloid leukemia (AML), Burkitt's leukemia/lymphoma (B-ALL), melanoma and
25 prostate carcinoma.
29. The method according to claim 28, wherein said glioma is selected from astrocytomas, oligodendrogliomas, ependymomas and glioblastoma multiforme.





30. The method according to claim 26, wherein said compound is an inhibitor of a mutant of IDH1 wherein said mutation is selected from R132H, R132C, R132S, R132L, R132G in IDH1.
- 5 31. The method according to claim 26, wherein said compound is an inhibitor of a mutant of IDH2, wherein said mutation is selected from R172M, R172G, R172K or R140Q.
32. The method according to claim 26, wherein said compound is an inhibitor of wild type IDH1.
- 10 33. The method according to claim 26, wherein said compound is an inhibitor of wild type IDH2.
34. The method according to any of claims 12-33, wherein said disease is associated with an increase in the production of 2-hydroxyglutarate in a tissue or plasma.
- 15 35. The method according to any of claims 12-34, wherein said compound of Formula I is a compound according to any of claims 1-11.
- 20 36. The method according to any of claims 12-34, wherein said compound of Formula I is selected from Table 1:





TABLE 1





1	 (4R,5R)-8-(cyclopent-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
---	---





2	 1-(4-fluorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
3	 (4R,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-8-phenyl-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
4	 N-(((4R,5R)-8-(cyclohexylethynyl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-4-yl)acetamide
5	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea





6	 N-(((4R,5R)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-8-(4-methylpent-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide
7	 N-((2S,3S)-5-((R)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-3,4,5,6-tetrahydro-2H-benzo[b][1,5]oxazocin-10-yl)benzo[d]thiazole-2-carboxamide
8	 (4R,5R)-8-(cyclopropylethynyl)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
9	 (4R,5S)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-4-ylmethyl)amino)methyl)-8-(p-tolyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide





10	 (4S,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-8-(pent-1-yn-1-yl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
11	 N-(((4R,5S)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-8-(4-methylpent-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide
12	 (4R,5S)-8-(cyclohex-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-3-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
13	 (4S,5S)-8-(cyclopent-1-en-1-yl)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-4-ylmethyl)amino)methyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide





14	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)cyclohexanecarboxamide
15	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-2-(((4-methoxybenzyl)(methyl)amino)methyl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)cyclohexanecarboxamide
16	 (4S,5S)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-5-((methyl(pyridin-4-ylmethyl)amino)methyl)-8-(o-tolyl)-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocine 1,1-dioxide
17	 (2S,3S)-5-((R)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyrimidin-5-ylmethyl)amino)methyl)-8-((E)-styryl)-4,5-dihydro-2H-pyrido[2,3-b][1,5]oxazocin-6(3H)-one





18	 4-((((4R,5R)-8-(3-cyclopentylprop-1-yn-1-yl)-2-((R)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)(methyl)amino)methyl)benzoic acid
19	 N-(((4R,5R)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-8-(3-methylbut-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methylnicotinamide
20	 N-(((4R,5R)-8-(cyclohex-1-en-1-yl)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide
21	 N-(((4S,5S)-2-((S)-1-hydroxypropan-2-yl)-4-methyl-8-(4-methylpent-1-yn-1-yl)-1,1-dioxido-2,3,4,5-tetrahydrobenzo[b][1,4,5]oxathiazocin-5-yl)methyl)-N-methyl-2-(pyridin-3-yl)acetamide




22	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-isopropylurea
23	 1-(4-bromophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
24	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(4-methoxyphenyl)urea
25	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(pyridin-3-yl)urea





26	 1-cyclopentyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
27	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)-3-(2-morpholinopyridin-4-yl)urea
28	 1-ethyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
29	 1-(4-chlorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea




30	 1-((cyclohexylmethyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
31	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)-3-phenylurea
32	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)-3-(naphthalen-1-yl)urea
33	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)-3-(4-(trifluoromethyl)phenyl)urea



34	 1-(benzo[d][1,3]dioxol-5-yl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
35	 1-(3,5-dimethylisoxazol-4-yl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
36	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)cyclopropanecarboxamide
37	 N-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)isonicotinamide

38	 1-(2-fluorophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
39	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(1,2,3,4-tetrahydronaphthalen-1-yl)urea
40	 1-(4-cyanophenyl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
41	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(3-(pyrimidin-2-yl)phenyl)urea

42	 1-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-(2-(piperidin-1-yl)phenyl)urea
43	 1-(benzofuran-5-yl)-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
44	 1-cyclohexyl-3-((2R,3R)-2-(((3-fluoropyridin-4-yl)methyl)(methyl)amino)methyl)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea

45	 1-((2R,3R)-2-(((3-chloropyridin-4-yl)methyl)(methyl)amino)methyl)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-cyclohexylurea
46	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(pyrimidin-4-yl)methyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
47	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-2-((methyl(quinolin-4-yl)methyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
48	 1-((2R,3R)-2-(((1H-imidazol-2-yl)methyl)(methyl)amino)methyl)-5-((S)-1-hydroxypropan-2-yl)-3-methyl-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)-3-cyclohexylurea

49	 1-cyclohexyl-3-((2R,3R)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-5-((S)-1-(2-(pyrrolidin-1-yl)ethoxy)propan-2-yl)-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
50	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-((4-methoxybenzyl)oxy)propan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea
51	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-(4-(4-methoxyphenyl)-1H-1,2,3-triazol-1-yl)propan-2-yl)-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazin-9-yl)urea

52	 1-cyclohexyl-3-((2R,3R)-5-isopropyl-3-methyl-2-((methyl(pyridin-4-ylmethyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea
53	 1-cyclohexyl-3-((2R,3R)-5-((S)-1-((4-methoxybenzyl)oxy)propan-2-yl)-3-methyl-2-((methyl((3-methylpyridin-4-yl)methyl)amino)methyl)-6-oxo-2,3,4,5,6,7-hexahydrobenzo[h][1,5]oxazonin-9-yl)urea




FIG. 1A

B Optimization of resazurin concentration




FIG. 1B

C Reaction time course in different buffers




FIG. 1C




FIG. 1E

FIG. 1D

Reaction time course




FIG. 1F

Concentration-response of enzyme




FIG. 1G




FIG. 2