




(86) Date de dépôt PCT/PCT Filing Date: 2006/12/14	(51) Cl.Int./Int.Cl. <i>C07K 16/18</i> (2006.01), <i>C12N 15/13</i> (2006.01)
(87) Date publication PCT/PCT Publication Date: 2007/10/25	
(45) Date de délivrance/Issue Date: 2015/12/22	(72) Inventeurs/Inventors: GORDON, NATHANIEL C., US; KELLEY, ROBERT F., US; PHAM, ANH, US; HYMOWITZ, SARAH, US
(85) Entrée phase nationale/National Entry: 2008/06/10	
(86) N° demande PCT/PCT Application No.: US 2006/062115	(73) Propriétaire/Owner: GENENTECH, INC., US
(87) N° publication PCT/PCT Publication No.: 2007/120334	
(30) Priorités/Priorities: 2005/12/15 (US60/751,081); 2006/04/21 (US60/793,980)	(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : METHODES ET COMPOSITIONS DE CIBLAGE DE LA POLYUBIQUITINE
(54) Title: METHODS AND COMPOSITIONS FOR TARGETING POLYUBIQUITIN

1 MET GLN ILE PHE VAL LYS THR LEU THR GLY LYS THR ILE THR
15 LEU GLU VAL GLU PRO SER ASP THR ILE GLU ASN VAL LYS ALA
29 LYS ILE GLN ASP LYS GLU GLY ILE PRO PRO ASP GLN GLN ARG
43 LEU ILE PHE ALA GLY LYS GLN LEU GLU ASP GLY ARG THR LEU
57 SER ASP TYR ASN ILE GLN LYS GLU SER THR LEU HIS LEU VAL
71 LEU ARG LEU ARG GLY GLY

A



(57) Abrégé/Abstract:
Anti-polyubiquitin monoclonal antibodies, and methods for using the antibodies, are provided.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 October 2007 (25.10.2007)

PCT

(10) International Publication Number
WO 2007/120334 A2

(51) International Patent Classification: Not classified

(21) International Application Number:
PCT/US2006/062115

(22) International Filing Date:
14 December 2006 (14.12.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/751,081 15 December 2005 (15.12.2005) US
60/793,980 21 April 2006 (21.04.2006) US

(71) Applicant (for all designated States except US): **GENEN-TECH, INC.** [US/US]; 1 DNA Way, South San Francisco, California 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GORDON, Nathaniel C.** [US/US]; 1415 Sharp Court, Campbell,

California 95008 (US). **KELLEY, Robert F.** [US/US]; 1029 San Felipe Avenue, San Bruno, California 94066 (US). **PHAM, Anh** [US/US]; 400 North Chapel Avenue, #102, Alhambra, California 91801 (US). **HYMOWITZ, Sarah** [US/US]; 1050 Francisco Street, San Francisco, California 94109 (US).

(74) Agents: **ELLIOTT, Jennifer** et al.; c/o GENENTECH, INC., 1 DNA Way MS 49, South San Francisco, California 94080 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

[Continued on next page]


(54) Title: METHODS AND COMPOSITIONS FOR TARGETING POLYUBIQUITIN

1 MET GLN ILE PHE VAL LYS THR LEU THR GLY LYS THR ILE THR
15 LEU GLU VAL GLU PRO SER ASP THR ILE GLU ASN VAL LYS ALA
29 LYS ILE GLN ASP LYS GLU GLY ILE PRO PRO ASP GLN GLN ARG
43 LEU ILE PHE ALA GLY LYS GLN LEU GLU ASP GLY ARG THR LEU
57 SER ASP TYR ASN ILE GLN LYS GLU SER THR LEU HIS LEU VAL
71 LEU ARG LEU ARG GLY GLY

A



B



(57) Abstract: Anti-polyubiquitin monoclonal antibodies, and methods for using the antibodies, are provided.

WO 2007/120334 A2

WO 2007/120334 A2

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *without international search report and to be republished upon receipt of that report*
- *with sequence listing part of description published separately in electronic form and available upon request from the International Bureau*

(48) Date of publication of this corrected version:

3 April 2008

(15) Information about Correction:

see Notice of 3 April 2008

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

ELECTRONICALLY FILED: December 14, 2006
ATTORNEY DOCKET NO: P2260R1

METHODS AND COMPOSITIONS FOR TARGETING POLYUBIQUITIN

FIELD OF THE INVENTION

This invention relates to the field of anti-polyubiquitin antibodies, and more particularly to anti-polyubiquitin antibodies that do not specifically bind to monoubiquitin and that can discriminate between polyubiquitins having different isopeptide linkages.

BACKGROUND

Ubiquitin is a small protein that has important regulatory roles in a wide variety of cellular pathways. The best known of these is ubiquitin's role in protein degradation, where covalent attachment of ubiquitin to a target protein enables that target protein to be recognized and destroyed by the 26S proteasome (*see* Wilkinson, Semin. Cell Devel. Biol. 11(3): 141-148 (2000)). Protein kinase regulation of various signaling pathways has also been correlated with ubiquitination (*see* Sun and Chen, Curr. Opin. Cell Biol. 16: 119-126 (2004)). For example, phosphorylation of I κ B by I κ B kinase permits ubiquitination of I κ B and subsequent degradation by the 26S proteasome; because I κ B is an inhibitor of NF κ B, the degradation of I κ B activates NF κ B (Ghosh and Karin, Cell 109 (Suppl.): S81-S96 (2002); Palombella et al., Cell 78: 773-785 (1994)). Ubiquitination also mediates DNA repair (*see* Sun and Chen, Curr. Opin. Cell Biol. 16:119-126 (2004)). After DNA is damaged, monoubiquitination of proliferating cell nuclear antigen (PCNA) activates damage-tolerant polymerases which are able to synthesize DNA despite any DNA lesions (Stelter and Ulrich, Nature 425: 188-191 (2003)). Other physiological processes in which ubiquitination is known to be involved include cell division, cell growth, cell movement, and apoptosis/cell death (Johnson, Nat. Cell Biol. 4:E295-E298 (2002); Pickart, Mol. Cell. 8: 499-504 (2001)).

The covalent attachment of ubiquitin, a 76 amino acid protein, to a target protein is a three-step enzymatic process (Pickart, Annu. Rev. Biochem. 70: 503-533 (2001)). First, ubiquitin-activating enzyme E1 forms an ubiquitin-E1 thioester in an ATP-dependent reaction. The ubiquitin is transferred from the ubiquitin-E1 thioester to a member of the ubiquitin-conjugating enzyme (E2) family in the second step. In the third step, with the assistance of a ubiquitin-protein ligase (E3), an isopeptide bond is formed between the carboxyl terminus of ubiquitin and the ϵ -amino group of a lysine residue on the target protein. Enzymes termed deubiquitinases remove ubiquitin moieties from target proteins (Guterman and Glickman, Curr. Prot. Pep. Sci. 5: 201-210 (2004)). Highlighting ubiquitin's role as an important regulatory molecule, the human genome contains many different proteins involved in ubiquitination or deubiquitination: at least 40 different E2s, 500 different E3s,

and 80 different deubiquitinases have been identified thus far (Wong et al., *Drug. Discov. Today* 8: 746-754 (2003)).

Ubiquitin contains seven lysine residues (Lys6, Lys22, Lys27, Lys33, Lys29, Lys48, and Lys63), and thus ubiquitin itself may serve as a target protein for ubiquitination (Peng et al., *Nat. Biotechnol.* 21: 921-926 (2003); Pickart and Fushman, *Curr. Opin. Chem. Biol.* 8:610-616 (2004)). The molecule produced upon ubiquitination of a ubiquitin protein is termed a polyubiquitin molecule, and may comprise two or more ubiquitin moieties. Ubiquitination of ubiquitin may theoretically occur at any of the seven lysine residues (Peng et al., *Nat. Biotechnol.* 21: 921-926 (2003)), so that different species of polyubiquitins exist having isopeptide bonds to different lysine residues within ubiquitin. It is possible that a single polyubiquitin molecule with greater than two ubiquitin moieties may have more than one type of lysine linkage. Studies have shown that the E2 enzyme influences the type of lysine linkage created between one ubiquitin molecule and another (Tenno et al., *Genes to Cells* 9: 865-875 (2004); Deng et al. (2000); Hofmann and Pickart (2001)). Polyubiquitin and ubiquitin exist both as free molecules and in covalent attachment with a target protein.

Like ubiquitin, polyubiquitin involvement has been found in many cellular processes, including intracellular trafficking, endocytosis, gene expression/silencing, proteolysis, kinase activation, translation, and DNA repair (Hoege et al., *Nature* 419:135-141 (2002); Spence et al., *Mol. Cell. Biol.* 15:1265-1273 (1995); Hofmann and Pickart, *Cell* 96: 645-653 (1999). Polyubiquitin and polyubiquitination can have strikingly different physiological roles than monoubiquitin and monoubiquitination in the same pathways, however. For example, whereas monoubiquitination of PCNA after DNA damage results in the activation of error-prone DNA polymerases, polyubiquitination of PCNA at the identical residue where monoubiquitination is observed results in activation of error-free DNA repair (Stelter and Ulrich, *Nature* 425: 188-191 (2003); Hoege et al., *Nature* 419:135-141 (2002); Spence et al., *Mol. Cell. Biol.* 15:1265-1273 (1995); and Hofmann and Pickart, *Cell* 96: 645-653 (1999)).

Even polyubiquitins having different lysine linkages appear to play different physiological roles. The two best-studied are the Lys48-linked and Lys63-linked polyubiquitins, and structural studies of the two suggest that different lysine-linked polyubiquitins may adopt markedly different conformations, thus permitting different interactions with selected binding partners (Tenno et al., *Genes to Cells* 9: 865-875 (2004)). Covalent modification by Lys48-linked polyubiquitin typically marks the target protein for proteolytic degradation, though there is some evidence that Lys48-linked polyubiquitin may also regulate certain proteins by non-proteolytic means (Chau et al., *Science* 243: 1576-1583 (1989); Finley et al., *Mol. Cell. Biol.* 14: 5501-5509 (1994); Flick et al., *Nat. Cell. Biol.* 6:634-641 (2004)). Lys63-linked polyubiquitins, in contrast, have been linked to a variety of nonproteolytic intracellular pathways, including DNA repair (yeast cells expressing K63R-ubiquitin are defective in DNA repair), kinase activation, intracellular trafficking, and translation (Pickart and Fushman, *Curr. Opin. Chem. Biol.* 8: 610-616 (2004); Hicke and Dunn, *Annu Rev. Cell Dev. Biol.*

19: 141-172 (2003); Spece et al., Mol. Cell Biol. 15: 1265-1273 (1995); Ulrich, Eukaryot. Cell 1: 1-10 (2002); Spence et al., Cell 102: 67-76 (2000); Seibenhener et al., Mol. Cell. Biol. 24(18): 8055-8068 (2004)). In one specific example, synphilin-1 is normally ubiquitinated with K63-linked polyubiquitin by parkin in a proteasomal-independent manner, but synphilin-1 can alternately be
5 targeted for destruction by ubiquitination with K48-linked polyubiquitin (Lim et al., J. Neurosci. 25(8): 2002-9 (2005)). An analysis of subjects with Parkinson's disease shows that K63-polyubiquitination of synphilin-1 may be involved in the formation of Lewy body inclusions associated with that disease (Lim et al., J. Neurosci. 25(8): 2002-9 (2005)).

Other lysine-linked polyubiquitins have not been studied extensively, largely because of the
10 difficulty in distinguishing between them. Studies have thus far relied on cells expressing mutagenized ubiquitins in which one or more lysines have been removed, on enzymatically synthesized polyubiquitins of particular linkages, or on techniques such as mass spectrometry to distinguish between one type of polyubiquitin and another. Each of those methodologies is ill-suited or cumbersome for analysis of the normal physiological behavior of particular lysine-linked
15 polyubiquitins. While antibodies exist that are specific for polyubiquitin as opposed to monoubiquitin (Fujimoro et al., FEBS Lett. 349: 173-180 (1994)), there are as yet no antibodies that can distinguish between polyubiquitins of different lysine linkages.

Unsurprisingly, given their important roles in a variety of cellular processes, ubiquitin and polyubiquitins have also been implicated in many diseases *see* (Argiles, Ubiquitin and Disease, R. G. Landes (1998)). Ubiquitin dysregulation is observed in muscle wasting (Mitch and Goldberg, New Engl. J. Med. 335: 1897-905 (1996); Bodine et al., Science 294: 1704-1708 (2001)). Several genetic diseases have been linked to aberrant ubiquitin activity, including cystic fibrosis (Ward et al., Cell 83: 121-127 (1995)), Angelman's syndrome (Kishino et al., Nature Genet. 15: 70-73 (1997)), and Liddle syndrome (Staub et al., EMBO J 16: 6325-6336 (1997)). Ubiquitin also plays a role in immune and
25 inflammatory responses; for example, extracellular ubiquitin has been found to act as a sort of cytokine, inhibiting the TNF α response to endotoxin in peripheral blood mononuclear cells and regulating endotoxin hyporesponsiveness (Majetschak et al., Blood 101: 1882-1890 (2003); Ciechanover, EMBO J 17: 7151-7160 (1998)). Also, both ubiquitin and polyubiquitin have been found in human serum, with higher levels of both molecules observed in the serum of patients having
30 parasitic and allergic disease (Takada et al., Clinical Chem. 43: 1188-1195 (1997)).

Dysregulation of several ubiquitin-mediated pathways are also involved in cancer (Spataro et al., Br. J. Cancer 77: 448-55 (1998); Beckmann et al., Hum. Mutat. 25: 507-12 (2005)). For example, mutations in the heterodimeric ubiquitin ligase BRCA1-BARD1 are correlated with breast cancer (Hashizume et al., J. Biol. Chem. 276: 14537-40 (2001)), mutations that disrupt the ability of Myc to
35 be degraded by the ubiquitin pathway activate the oncogenic potential of c-Myc (Salghetti et al., EMBO J. 18: 717-726 (1999)), and transformed v-Jun is unable to be ubiquitinated and degraded as

its non-oncogenic correlate, c-Jun, is, giving rise to uncontrolled growth (Ciechanover, EMBO J. 17: 7151-7160 (1998); Trier et al., Cell 78: 787-798 (1994)).

Ubiquitin and polyubiquitin have particularly been studied in the context of neurological diseases (Chung et al., TINS 24(11 Suppl.) S7-S14 (2001)). The inclusions, bodies, and neurofibrillary tangles that accumulate in Huntington's disease, Spinocerebellar ataxia, prion encephalopathies, Pick's disease, Lewy body disease, Parkinson's disease, and Alzheimer's disease stain immunopositively for mono and/or polyubiquitin (Alves-Rodrigues et al., Trends Neurosci. 21: 516-520 (1998); Cammarata et al., Neurosci Lett. 156: 96-98 (1993); Kalchman et al., J. Biol. Chem. 271: 19385-94 (1996); Holmberg et al., Human Mol. Genet. 7: 913-918 (1998); Yedidia et al., EMBO J. 20: 5383-91 (2001); Mori et al., Science 235: 1641-44 (1987); Leigh et al., Acta Neuropathol. (Berl.) 79: 61-72 (1989); and Kuzuhara et al., Acta Neuropathologica 75: 345-353 (1988)). Several forms of Parkinson's disease have been linked to mutations in the ubiquitin carboxy-terminal hydrolase L1 (UCH-L1) gene, a deubiquitinase (Leroy et al., Nature 395: 451-452 (1998)), while other forms of Parkinson's have been linked to inactivating mutations in Parkin, an E2-dependent ubiquitin-protein ligase known to interact with the ubiquitin-conjugating enzyme UbcH7 and to ubiquitinate synphilin-1 (Shimura et al., Nature Genet. 25: 302-305 (2000), Zhang et al., Proc. Natl. Acad. Sci. 97: 13354-13359 (2000); Lim et al., J. Neurosci. 25(8): 2002-9 (2005)). Both types of mutations result in aberrant proteolytic processing and the inappropriate aggregation of proteins (*see* McNaught et al., Nature Rev. Neurosci. 2: 589-594 (2001)). UCH-L1 mutations have also been found to segregate with Huntington's disease (Naze et al., Neurosci. Lett. 328: 1: 1-4 (2002)). A mutant form of ubiquitin has been identified in the brains of Alzheimer's patients that is very efficiently incorporated into polyubiquitin chains, but is refractory to deubiquitination once formed, potentially leading to dominant inhibition of the normal cellular proteolytic processing system (Lam et al., Proc. Natl. Acad. Sci. 97: 9902-9906 (2000)).

It is clear that it would be beneficial not only to have compositions and methods that can distinguish between polyubiquitins of different lysine linkages, but also to have compositions and methods that are effective in targeting and modulating ubiquitin and polyubiquitin-mediated pathways. The invention provided herein relates to such compositions and methods.

30

DISCLOSURE OF THE INVENTION

The invention provides novel antibodies capable of binding to and/or regulating biological activities associated with polyubiquitin.

35 In one embodiment, an isolated antibody that specifically binds to polyubiquitin is provided, wherein the antibody does not specifically bind to monoubiquitin. In one embodiment, an isolated

antibody that specifically binds a first polyubiquitin comprising a first lysine linkage is provided, wherein the antibody does not specifically bind a second polyubiquitin comprising a second lysine linkage, and wherein the first lysine linkage differs from the second lysine linkage. In one aspect, the antibody further specifically binds lysine 6-linked polyubiquitin, lysine 11-linked polyubiquitin, lysine 27-linked polyubiquitin, lysine 29-linked polyubiquitin, lysine 33-linked polyubiquitin, lysine 48-linked polyubiquitin, or lysine 63-linked polyubiquitin.

In one embodiment, an isolated antibody that specifically binds a first K48-linked polyubiquitin is provided, wherein the antibody does not specifically bind a second polyubiquitin comprising a different lysine-linked form of polyubiquitin (i.e., not K48-linked polyubiquitin). In one embodiment, the second polyubiquitin is K63-linked polyubiquitin.

In one embodiment, an isolated antibody that specifically binds a first K63-linked polyubiquitin is provided, wherein the antibody does not specifically bind a second polyubiquitin comprising a different lysine-linked form of polyubiquitin (i.e., not K63-linked polyubiquitin). In one embodiment, the second polyubiquitin is K48-linked polyubiquitin.

In one embodiment, an isolated antibody that specifically binds both a first polyubiquitin comprising a first lysine linkage and a second polyubiquitin comprising a second lysine linkage is provided, wherein the first lysine linkage differs from the second lysine linkage, wherein the antibody does not specifically bind monoubiquitin, and wherein the antibody binds the second polyubiquitin with a substantially reduced binding affinity as compared to the binding affinity of the antibody for the first polyubiquitin.

In one embodiment, an isolated antibody that specifically binds lysine-48-linked polyubiquitin is provided, wherein the antibody does not specifically bind monoubiquitin. In one embodiment, the antibody further comprises at least one hypervariable (HVR) sequence selected from HVR-H1, HVR-H2, HVR-H3, and HVR-L3 of any of SEQ ID NOs: 1-25, 151-175, 265-279, 392-459, and 695-704; SEQ ID NOs: 27-51, 177-201, 281-295, 461-528, and 706-715; SEQ ID NOs: 53-77, 203-227, 297-311, 530-597, and 717-726; and SEQ ID NOs: 313-327 and 728-737, respectively. In one embodiment, the antibody further comprises at least one sequence selected from HVR-H1, HVR-H2, HVR-H3, wherein HVR-H1 comprises the amino acid sequence a b c d e f g h i j (SEQ ID NO: 825), wherein amino acid a is glycine; amino acid b is phenylalanine; amino acid c is asparagine; amino acid d is selected from valine, phenylalanine, leucine, and isoleucine; amino acid e is selected from serine and tyrosine; amino acid f is tyrosine; amino acid g is selected from serine and tyrosine; amino acid h is selected from serine and tyrosine; amino acid i is selected from isoleucine and methionine; and amino acid j is histidine; wherein HVR-H2 comprises the amino acid sequence k l m n o p q r s t u v w x y z a' (SEQ ID NO: 826), wherein amino acid k is serine; amino acid l is isoleucine; amino acid m is selected from serine and tyrosine; amino acid n is selected from proline and serine; amino acid o is tyrosine; amino acid p is tyrosine; amino acid q is selected from serine and glycine; amino acid r is selected from serine and tyrosine; amino acid s is threonine, amino acid t is

selected from serine and tyrosine; amino acid u is tyrosine; amino acid v is alanine; amino acid w is aspartic acid; amino acid x is serine; amino acid y is valine; amino acid z is lysine; and amino acid a' is glycine; and wherein HVR-H3 comprises the amino acid sequence b'c' d' e' f' g' h' i' j' k' l', wherein amino acid b' is selected from glutamic acid, serine, glycine, and tyrosine; amino acid c' is selected from glycine, tyrosine, serine, and asparagine; amino acid d' is selected from tyrosine, serine, lysine, phenylalanine, and glutamic acid; amino acid e' is selected from serine, tyrosine, glycine, and tryptophan; amino acid f' is selected from glutamine, tyrosine, serine, and glycine; amino acid g' is selected from glycine, serine, tyrosine, methionine, and alanine; amino acid h' is selected from glycine, alanine, proline, and isoleucine; amino acid i' is selected from phenylalanine, isoleucine, methionine, alanine, and leucine, or is not present; amino acid j' is phenylalanine or is not present; amino acid k' is aspartic acid; and amino acid l' is tyrosine. In one embodiment, the antibody further comprises HVR-H1, HVR-H2, and HVR-H3 sequences corresponding to those set forth for clones apu01, apu02, apu03, apu04, apu05, apu06, apu07, apu08, apu09, apu10, apu11, apu12, apu13, apu14, or apu15 in Figures 10A and 10B.

In one embodiment, the antibody comprises at least one sequence selected from HVR-H1, HVR-H2, HVR-H3, wherein HVR-H1 comprises the amino acid sequence a b c d e f g h i j (SEQ ID NO: 827), wherein amino acid a is glycine; amino acid b is phenylalanine; amino acid c is asparagine; amino acid d is isoleucine; amino acid e is selected from serine and phenylalanine; amino acid f is tyrosine; amino acid g is selected from serine and glycine; amino acid h is selected from serine and glycine; amino acid i is selected from isoleucine and methionine; and amino acid j is histidine; wherein HVR-H2 comprises the amino acid sequence k l m n o p q r s t u v w x y z a' (SEQ ID NO: 828), wherein amino acid k is serine; amino acid l is isoleucine; amino acid m is tyrosine; amino acid n is serine; amino acid o is tyrosine; amino acid p is tyrosine; amino acid q is serine; amino acid r is tyrosine; amino acid s is threonine, amino acid t is serine; amino acid u is tyrosine; amino acid v is alanine; amino acid w is aspartic acid; amino acid x is serine; amino acid y is valine; amino acid z' is lysine; and amino acid a' is glycine; and wherein HVR-H3 comprises the amino acid sequence b' c' d' e' f' g' h' i' j' k' (SEQ ID NO: 829), wherein amino acid b' is selected from serine and glycine; amino acid c' is tyrosine; amino acid d' is serine; amino acid e' is selected from tyrosine and tryptophan; amino acid f' is selected from serine, tyrosine, arginine, phenylalanine, and histidine; amino acid g' is selected from glutamic acid, serine, leucine, phenylalanine, methionine, asparagine, and valine; amino acid h' is selected from alanine and glycine; amino acid i' is selected from leucine, methionine, phenylalanine, and isoleucine; amino acid j' is aspartic acid; and amino acid k' is tyrosine. In one embodiment, the antibody further comprises HVR-H1, HVR-H2, and HVR-H3 sequences corresponding to those set forth for clones apu2.01, apu2.02, apu2.03, apu2.04, apu2.05, apu2.06, apu2.07, apu2.08, apu2.09, or apu2.10 in Figure 16A.

In one embodiment, the antibody further comprises a HVR-L3 sequence comprising the amino acid sequence m' n' o' p' q' r' s' t' u' v' w' (SEQ ID NO: 830), wherein amino acid m' is

glutamine; amino acid n' is glutamine; amino acid o' is selected from serine and tyrosine; amino acid p' is selected from serine and tyrosine; amino acid q' is selected from serine and tyrosine; amino acid r' is selected from serine and tyrosine; amino acid s' is selected from serine and tyrosine; amino acid t' is selected from leucine, serine, proline, and tyrosine; amino acid u' is proline or is not present;

5 amino acid v' is selected from phenylalanine, isoleucine, valine, and leucine; and amino acid w' is threonine. In one embodiment, the antibody further comprises an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence corresponding to the HVR-L3 sequence set forth for clones apu01, apu02, apu03, apu04, apu05, apu06, apu07, apu08, apu09, apu10, apu11, apu12, apu13, apu14, or apu15 in Figure 10C. In one embodiment, the antibody further
10 comprises a HVR-L3 sequence comprising the amino acid sequence Q-Q-S-S-Y-S-S-L-I-T (SEQ ID NO: 728). In one embodiment, the antibody further comprises an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence corresponding to the HVR-L3 sequence set forth for clones apu2.01, apu2.02, apu2.03, apu2.04, apu2.05, apu2.06, apu2.07, apu2.08, apu2.09, or apu2.10 in Figure 16B.

15 In one embodiment, an isolated antibody that specifically binds lysine-48-linked polyubiquitin is provided, wherein the antibody does not specifically bind monoubiquitin, and wherein the antibody comprises an HVR-H1 sequence of SEQ ID NO: 269, an HVR-H2 sequence of SEQ ID NO: 285, an HVR-H3 sequence of SEQ ID NO: 301, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 317. In one
20 embodiment, an isolated antibody that specifically binds lysine-48-linked polyubiquitin is provided, wherein the antibody does not specifically bind monoubiquitin, and wherein the antibody comprises an HVR-H1 sequence of SEQ ID NO: 701, an HVR-H2 sequence of SEQ ID NO: 712, an HVR-H3 sequence of SEQ ID NO: 723, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 734. In one embodiment, an isolated
25 antibody that specifically binds lysine-48-linked polyubiquitin is provided, wherein the antibody does not specifically bind monoubiquitin, and wherein the antibody comprises an HVR-H1 sequence of SEQ ID NO: 701, an HVR-H2 sequence of SEQ ID NO: 712, an HVR-H3 sequence of SEQ ID NO: 723, and HVR-L1 sequence of SEQ ID NO: 79, and HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 734.

30 In one embodiment, an isolated antibody that specifically binds to lysine-63-linked polyubiquitin is provided, wherein the antibody does not specifically bind to monoubiquitin. In one embodiment the antibody further comprises at least one hypervariable (HVR) sequence selected from HVR-H1, HVR-H2, HVR-H3, and HVR-L3 of any of SEQ ID NOs: 81-89, 229-239, 329-336, 599-629, and 739-748; SEQ ID NOs: 91-99, 241-251; 338-345, 631-661, and 750-759; SEQ ID NOs: 101-
35 109, 253-263, 347-354, 663-693, and 761-770; and SEQ ID NOs: 356-363 and 772-781, respectively. In one embodiment, the antibody comprises at least one sequence selected from HVR-H1, HVR-H2, HVR-H3, wherein HVR-H1 comprises the amino acid sequence a b c d e f g h i j (SEQ ID NO: 831),

wherein amino acid a is glycine; amino acid b is phenylalanine; amino acid c is asparagine; amino acid d is selected from valine, isoleucine, and phenylalanine; amino acid e is selected from serine and tyrosine; amino acid f is selected from serine and tyrosine; amino acid g is selected from serine and tyrosine; amino acid h is selected from serine and tyrosine; amino acid i is selected from isoleucine and methionine; and amino acid j is histidine; wherein HVR-H2 comprises the amino acid sequence k l m n o p q r s t u v w x y z a' (SEQ ID NO: 832), wherein amino acid k is selected from serine and tyrosine; amino acid l is isoleucine; amino acid m is selected from serine and tyrosine; amino acid n is selected from proline and serine; amino acid o is selected from serine and tyrosine; amino acid p is selected from serine and tyrosine; amino acid q is selected from serine and glycine; amino acid r is selected from serine and tyrosine; amino acid s is threonine, amino acid t is selected from serine and tyrosine; amino acid u is tyrosine; amino acid v is alanine; amino acid w is aspartic acid; amino acid x is serine; amino acid y is valine; amino acid z is lysine; and amino acid a is glycine; and wherein HVR-H3 comprises the amino acid sequence b' c' d' e' f' g' h' i' j' k' l' m' n' o' p' q' r' s' t' u' v', wherein amino acid b' is selected from serine, glutamic acid, glycine, and tryptophan; amino acid c' is selected from glycine, tyrosine, isoleucine, glutamine, and serine; amino acid d' is selected from tyrosine, methionine, glycine, and isoleucine; amino acid e' is selected from tyrosine, arginine, phenylalanine, tryptophan, alanine, and proline; amino acid f' is selected from tyrosine, tryptophan, serine, and glycine; amino acid g' is selected from glutamine, tyrosine, serine, phenylalanine, and valine; amino acid h' is selected from glycine, threonine, tryptophan, lysine, and proline; amino acid i' is selected from tyrosine, alanine, tryptophan, glutamic acid, proline, and serine; amino acid j' is selected from tryptophan, isoleucine, tyrosine, and alanine; amino acid k' is selected from tryptophan, tyrosine, glycine, and aspartic acid, or is not present; amino acid l' is selected from tyrosine, serine, phenylalanine, and tryptophan, or is not present; amino acid m' is selected from tyrosine, aspartic acid, and serine, or is not present; amino acid n' is selected from tyrosine and alanine, or is not present; amino acid o' is selected from threonine, serine, valine, glycine, and tyrosine, or is not present; amino acid p' is selected from glycine, aspartic acid, serine, methionine, and tyrosine, or is not present; amino acid q' is selected from tyrosine, alanine, and glycine, or is not present; amino acid r' is selected from tyrosine, leucine, and glycine, or is not present; amino acid s' is glycine or is not present; amino acid t' is selected from methionine and leucine, or is not present; amino acid u' is aspartic acid; and amino acid v' is tyrosine. In one embodiment, the antibody further comprises HVR-H1, HVR-H2, and HVR-H3 sequences corresponding to those set forth for clones apu17, apu18, apu19, apu20, apu21, apu22, apu23, and apu24 in Figures 11A and 11B.

In one embodiment, the antibody comprises at least one sequence selected from HVR-H1, HVR-H2, HVR-H3, wherein HVR-H1 comprises the amino acid sequence a b c d e f g h i j (SEQ ID NO: 833), wherein amino acid a is glycine; amino acid b is phenylalanine; amino acid c is asparagine; amino acid d is selected from isoleucine, valine, and leucine; amino acid e is selected from serine, lysine, and valine; amino acid f is selected from serine, tryptophan, glycine, and threonine; amino acid

g is selected from serine, asparagine, and glycine; amino acid h is selected from tyrosine, isoleucine, leucine, and phenylalanine; amino acid i is selected from isoleucine and methionine; and amino acid j is histidine; wherein HVR-H2 comprises the amino acid sequence k l m n o p q r s t u v w x y z a' (SEQ ID NO: 834), wherein amino acid k is selected from tyrosine, phenylalanine, aspartic acid, histidine, and alanine; amino acid l is isoleucine; amino acid m is selected from serine, alanine, and glutamine; amino acid n is proline; amino acid o is tyrosine; amino acid p is selected from leucine, tyrosine, and phenylalanine; amino acid q is selected from serine and glycine; amino acid r is selected from serine, threonine, and tryptophan; amino acid s is threonine, amino acid t is selected from serine, asparagine, lysine, and isoleucine; amino acid u is tyrosine; amino acid v is alanine; amino acid w is aspartic acid; amino acid x is serine; amino acid y is valine; amino acid z' is lysine; and amino acid a' is glycine; and wherein HVR-H3 comprises the amino acid sequence b' c' d' e' f' g' h' i' j' k' l' (SEQ ID NO: 908), wherein amino acid b' is glutamic acid; amino acid c' is tyrosine; amino acid d' is tyrosine; amino acid e' is arginine; amino acid f' is tryptophan; amino acid g' is tyrosine; amino acid h' is threonine; amino acid i' is alanine; amino acid j' is isoleucine; amino acid k' is aspartic acid; and amino acid l' is tyrosine. In one embodiment, the antibody comprises HVR-H1, HVR-H2, and HVR-H3 sequences corresponding to those set forth for clones apu2.11, apu2.12, apu2.13, apu2.14, apu2.15, apu2.16, apu2.17, apu2.18, apu2.19, and apu2.20 in Figure 17A.

In one embodiment, the antibody comprises at least one sequence selected from HVR-H1, HVR-H2, HVR-H3, wherein HVR-H1 comprises the amino acid sequence a b c d e f g h i j (SEQ ID NO: 835), wherein amino acid a is glycine; amino acid b is phenylalanine; amino acid c is asparagine; amino acid d is selected from isoleucine, valine, and leucine; amino acid e is selected from lysine and methionine; amino acid f is selected from threonine, methionine, asparagine, arginine, and isoleucine; amino acid g is selected from glycine, valine, and phenylalanine; amino acid h is selected from tyrosine, isoleucine, leucine, and phenylalanine; amino acid i is selected from isoleucine and methionine; and amino acid j is histidine; wherein HVR-H2 comprises the amino acid sequence k l m n o p q r s t u v w x y z a' b' (SEQ ID NO: 836), wherein amino acid k is alanine; amino acid l is tyrosine; amino acid m is isoleucine; amino acid n is selected from serine, isoleucine, and threonine; amino acid o is proline; amino acid p is tyrosine; amino acid q is selected from leucine, tyrosine, aspartic acid, serine, and tryptophan; amino acid r is glycine; amino acid s is selected from tryptophan, valine, serine, asparagine, arginine, and tyrosine; amino acid t is threonine, amino acid u is selected from arginine, asparagine, valine, threonine, serine, and lysine; amino acid v is tyrosine; amino acid w is alanine; amino acid x is aspartic acid; amino acid y is serine; amino acid z is valine; amino acid a' is lysine; and amino acid b' is glycine; and wherein HVR-H3 comprises the amino acid sequence c' d' e' f' g' h' i' j' k' l' m' n' o' (SEQ ID NO: 837), wherein amino acid c' is serine; amino acid d' is arginine; amino acid e' is glutamic acid; amino acid f' is tyrosine; amino acid g' is tyrosine; amino acid h' is arginine; amino acid i' is tryptophan; amino acid j' is tyrosine; amino acid k' is threonine; amino acid l' is alanine; amino acid m' is isoleucine; amino acid n' is aspartic acid; and

amino acid o' is tyrosine. In one embodiment, the antibody comprises HVR-H1, HVR-H2, and HVR-H3 sequences corresponding to those set forth for clones apu3.01, apu3.02, apu3.03, apu3.04, apu3.05, apu3.06, apu3.07, apu3.08, apu3.09, apu3.10, and 3.11 in Figures 23A and 23B.

In one embodiment, the antibody comprises a HVR-L3 sequence comprising the amino acid sequence w' x' y' z' A B C D E F G, wherein amino acid w' is glutamine; amino acid x' is glutamine; amino acid y' is selected from serine and tyrosine; amino acid z' is selected from serine and tyrosine; amino acid A is selected from serine and tyrosine; amino acid B is selected from serine and tyrosine; amino acid C is selected from proline, serine and leucine; amino acid D is selected from serine, proline, and tyrosine, or is not present; amino acid E is selected from leucine and phenylalanine, or is not present; amino acid F is selected from phenylalanine, valine, threonine, and isoleucine; and amino acid G is selected from arginine, threonine, and phenylalanine. In one embodiment, the antibody comprises an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence corresponding to the HVR-L3 sequence set forth for clones apu17, apu18, apu19, apu20, apu21, apu22, apu23, and apu24 in Figure 11C. In one embodiment, the antibody comprises a HVR-L3 sequence comprising the amino acid sequence Q-Q-Y-S-S-Y-S-S-L-F-T (SEQ ID NO: 772). In one embodiment, the antibody comprises an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence corresponding to the HVR-L3 sequence set forth for clones apu2.11, apu2.12, apu2.13, apu2.14, apu2.15, apu2.16, apu2.17, apu2.18, apu2.19, and apu2.20 in Figure 17B. In one embodiment, the antibody comprises an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence corresponding to the HVR-L3 sequence of SEQ ID NO: 777.

In one embodiment, an isolated antibody that specifically binds lysine-63-linked polyubiquitin is provided, wherein the antibody does not specifically bind monoubiquitin, and wherein the antibody comprises an HVR-H1 sequence of SEQ ID NO: 330, an HVR-H2 sequence of SEQ ID NO: 339, an HVR-H3 sequence of SEQ ID NO: 348, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 357. In one embodiment, an isolated antibody that specifically binds lysine-63-linked polyubiquitin is provided, wherein the antibody does not specifically bind monoubiquitin, and wherein the antibody comprises an HVR-H1 sequence of SEQ ID NO: 739, an HVR-H2 sequence of SEQ ID NO: 750, an HVR-H3 sequence of SEQ ID NO: 761, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 772. In one embodiment, an isolated antibody that specifically binds lysine-63-linked polyubiquitin is provided, wherein the antibody does not specifically bind monoubiquitin, and wherein the antibody comprises an HVR-H1 sequence of SEQ ID NO: 740, an HVR-H2 sequence of SEQ ID NO: 751, an HVR-H3 sequence of SEQ ID NO: 762, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 773. In one embodiment, an isolated antibody that specifically binds lysine-63-linked polyubiquitin is provided, wherein the antibody does not specifically bind

monoubiquitin, and wherein the antibody comprises an HVR-H1 sequence of SEQ ID NO: 744, an HVR-H2 sequence of SEQ ID NO: 755, an HVR-H3 sequence of SEQ ID NO: 766, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 777. In one embodiment, an isolated antibody that specifically binds lysine-63-linked polyubiquitin is provided, wherein the antibody does not specifically bind monoubiquitin, and wherein the antibody comprises an HVR-H1 sequence of SEQ ID NO: 795, an HVR-H2 sequence of SEQ ID NO: 807, an HVR-H3 sequence of SEQ ID NO: 819, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 777.

In one aspect, an isolated antibody that binds to the same antigenic determinant on polyubiquitin as the antibody of any of the aforementioned antibodies is provided, wherein the antibody does not specifically bind to monoubiquitin. In one aspect, an isolated antibody that competes with any of the aforementioned antibodies for binding to polyubiquitin is provided, wherein the antibody does not specifically bind to monoubiquitin.

In one aspect, any of the aforementioned antibodies specifically binds to a polyubiquitinated protein. In one aspect, the antibody further inhibits degradation of the polyubiquitinated protein. In one aspect, the antibody further modulates at least one polyubiquitin-mediated signaling pathway. In one aspect, the antibody further inhibits at least one polyubiquitin-mediated signaling pathway. In one aspect, the antibody further stimulates at least one polyubiquitin-mediated signaling pathway.

In one aspect, a nucleic acid molecule encoding an antibody of the invention is provided. In one aspect, a vector that comprises the nucleic acid is provided. In one aspect, a host cell comprising the vector is provided. In one aspect, a cell line capable of producing an antibody of the invention is provided. In one aspect, a method of producing an antibody of the invention is provided, comprising culturing a host cell comprising a nucleic acid molecule encoding the antibody under conditions wherein the antibody is produced. In one aspect, a composition comprising an effective amount of an antibody of the invention and a pharmaceutically acceptable carrier is provided.

In one aspect, a method of identifying the presence of polyubiquitin or a polyubiquitinated protein in a sample is provided, comprising contacting the sample with at least one antibody of the invention. In one aspect, a method for the treatment of a disease or condition associated with dysregulation of polyubiquitin in a patient is provided, the method comprising administering to the patient an effective amount of at least one antibody of the invention. In one aspect, the patient is a mammalian patient. In one aspect, the patient is human. In one aspect, the disease is selected from cancer, a muscular disorder, a ubiquitin-pathway-related genetic disorder, an immune/inflammatory disorder, and a neurological disorder. In one aspect, the disease is selected from carcinoma, lymphoma, blastoma, sarcoma, leukemia, muscular dystrophy, multiple sclerosis, amyotrophic lateral sclerosis, cystic fibrosis, Angelman's syndrome, Liddle syndrome, Alzheimer's disease, Parkinson's disease, Pick's disease, and Paget's disease.

In one aspect, a method of determining the presence of a polyubiquitin or polyubiquitinated protein in a sample suspected of containing a polyubiquitin or polyubiquitinated protein is provided, comprising exposing the sample to at least one antibody of the invention and determining the binding of the at least one antibody to a polyubiquitin or polyubiquitinated protein in the sample.

5 In one aspect, a method of separating polyubiquitinated protein from non-polyubiquitinated protein in a sample is provided, comprising contacting the sample with at least one antibody of the invention.

In one aspect, a method of determining the function and/or activity of polyubiquitin in a cell is provided, comprising contacting the cell with at least one antibody of the invention and assessing
10 the effect of said contacting step on the cell.

In one aspect, a method of determining the function and/or activity of polyubiquitin in a sample is provided, comprising contacting the sample with at least one antibody of the invention and assessing the effect of said contacting step on the sample.

In another embodiment, an isolated antibody that specifically binds to a lysine-63-linked
15 polyubiquitin is provided, wherein the antibody binds to an epitope in the lysine-63-linked polyubiquitin. In one aspect, the epitope includes residues in both a first ubiquitin subunit and a second ubiquitin subunit of the lysine-63-linked polyubiquitin. In another such aspect, the epitope includes at least one residue in a first ubiquitin subunit selected from Glu-18, Pro-19, Ser-20, Asp-21, Thr-55, Leu-56, Ser-57, Asp-58, Asn-60, Ile-61, and Gln-62. In another such aspect, the epitope
20 includes at least one residue in a second ubiquitin subunit selected from Leu-8, Thr-9, Glu-34, Gly-35, Ile-36, Pro-37, Asp-39, Gln-40, Leu-71, Arg-72, Leu-73, Arg-74, and Gly-75. In another such aspect, the epitope includes at least one residue in a first ubiquitin subunit selected from Glu-18, Pro-19, Ser-20, Asp-21, Thr-55, Leu-56, Ser-57, Asp-58, Asn-60, Ile-61, and Gln-62, and at least one residue in a second ubiquitin subunit selected from Leu-8, Thr-9, Glu-34, Gly-35, Ile-36, Pro-37, Asp-39, Gln-40,
25 Leu-71, Arg-72, Leu-73, Arg-74, and Gly-75.

In one embodiment, an isolated antibody that specifically binds to a first polyubiquitin comprising at least one isopeptide bond to a first lysine residue at a first amino acid position of a ubiquitin molecule is provided, wherein the antibody does not specifically bind to a second polyubiquitin comprising at least one isopeptide bond to a second lysine residue at a second amino
30 acid position of a ubiquitin molecule, and wherein the first and the second amino acid positions differ. An antibody of the invention can be in any number of forms. For example, an antibody of the invention can be a chimeric antibody, a humanized antibody or a human antibody. In one embodiment, an antibody of the invention is not a human antibody, for example it is not an antibody produced in a xenomouse™ (e.g., as described in WO/96/33735). An antibody of the invention can be
35 full length or a fragment thereof (e.g., a fragment comprising an antigen binding component). In another embodiment, the invention provides an antigen-binding fragment of any of the above-described antibodies.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the primary structure of ubiquitin and schematic views of certain polyubiquitin isopeptide linkages. Figure 1a shows the amino acid sequence of human ubiquitin (SEQ ID NO: 377), with the lysine residues indicated in bold, underlined text. Figure 1b shows a schematic depiction of the bond formed between the lysine-48 or lysine-63 of a first ubiquitin molecule and the C-terminal glycine residue of a second ubiquitin molecule.

Figures 2A-2C show heavy chain HVR loop sequences of anti-polyubiquitin antibody molecules specifically recognizing K48-linked polyubiquitin, as described in Example 1(A). The designator “48” indicates that the antibody molecule specifically recognized K48-linked polyubiquitin. The designator “both” indicates that the antibody molecule recognized both K48-linked and K63-linked polyubiquitin. The designator “all” indicates that the antibody molecule recognized both K48-linked and K63-linked polyubiquitin as well as monoubiquitin. The designator “n.p.” indicates that certain clones did not have an amino acid at the indicated position. The figures show the heavy chain HVR sequences, H1, H2, and H3. Amino acid positions are numbered according to the Kabat numbering system as described below.

Figures 3A-3B show heavy chain HVR loop sequences of anti-polyubiquitin antibody molecules specifically recognizing K63-linked polyubiquitin, as described in Example 1(A). The designator “63” indicates that the antibody molecule specifically recognized K63-linked polyubiquitin. The designator “both” indicates that the antibody molecule recognized both K63-linked and K48-linked polyubiquitin. The designator “all” indicates that the antibody molecule recognized both K63-linked and K48-linked polyubiquitin as well as monoubiquitin. The designator “n.p.” indicates that certain clones did not have an amino acid at the indicated position. The figures show the heavy chain HVR sequences, H1, H2, and H3. Amino acid positions are numbered according to the Kabat numbering system as described below.

Figures 4A and 4B and Figure 5 depict exemplary acceptor human consensus framework sequences for use in practicing the instant invention with sequence identifiers as follows:

Variable heavy (VH) consensus frameworks (Figures 4A and 4B)

Human VH subgroup I consensus framework minus Kabat CDRs (SEQ ID NOs: 111, 839, 858, and 877)

Human VH subgroup I consensus framework minus extended hypervariable regions (SEQ ID NOs: 112-114, 840-842, 859-861, and 878-880)

Human VH subgroup II consensus framework minus Kabat CDRs (SEQ ID NOs: 115, 843, 862, and 881)

Human VH subgroup II consensus framework minus extended hypervariable regions (SEQ ID NOs: 116-118, 844-846, 863-865, and 882-884)

Human VH subgroup III consensus framework minus Kabat CDRs (SEQ ID NOs: 119, 847, 866, and 885)

Human VH subgroup III consensus framework minus extended hypervariable regions (SEQ ID NOs: 120-122, 848-850, 867-869, and 886-888)

Human VH acceptor framework minus Kabat CDRs (SEQ ID NOs: 123, 851, 870, and 889)

Human VH acceptor framework minus extended hypervariable regions (SEQ ID NOs: 124-
5 125, 852-853, 871-872, and 890-891)

Human VH acceptor 2 framework minus Kabat CDRs (SEQ ID NOs: 126, 854, 873, and 892)

Human VH acceptor 2 framework minus extended hypervariable regions (SEQ ID NOs: 127-
129, 855-857, 874-876, and 893-895)

Variable light (VL) consensus frameworks (Figure 5)

10 Human VL kappa subgroup I consensus framework (SEQ ID NOs: 130, 896, 900, and 904)

Human VL kappa subgroup II consensus framework (SEQ ID NOs: 131, 897, 901, and 905)

Human VL kappa subgroup III consensus framework (SEQ ID NOs: 132, 898, 902, and 906)

Human VL kappa subgroup IV consensus framework (SEQ ID NOs: 133, 899, 903, and 907)

Figure 6 depicts framework region sequences of huMAb4D5-8 light and heavy chains.

15 Numbers in superscript/bold indicate amino acid positions according to Kabat.

Figure 7 depicts modified/variant framework region sequences of huMAb4D5-8 light and heavy chains. Numbers in superscript/bold indicate amino acid positions according to Kabat.

Figures 8A-8C show heavy chain HVR loop sequences of anti-polyubiquitin antibody molecules specifically recognizing K48-linked polyubiquitin, as described in Example 1A. The
20 figures show the heavy chain HVR sequences, H1, H2, and H3. The designator “n.p.” indicates that certain clones did not have an amino acid at the indicated position. Amino acid positions are numbered according to the Kabat numbering system as described below.

Figures 9A-9B show heavy chain HVR loop sequences of anti-polyubiquitin antibody molecules specifically recognizing K63-linked polyubiquitin, as described in Example 1A. The
25 figures show the heavy chain HVR sequences, H1, H2, and H3. The designator “n.p.” indicates that certain clones did not have an amino acid at the indicated position. Amino acid positions are numbered according to the Kabat numbering system as described below.

Figures 10A-10C show heavy chain and light chain HVR loop sequences of anti-polyubiquitin antibody molecules apu01-apu15 that specifically recognize K48-linked polyubiquitin
30 and which were recognized by an antibody specific for pentahistidine, as described in Example 1(B). The figure shows the heavy chain HVR sequences, H1, H2, and H3, and light chain HVR sequence, L3. The designator “n.p.” indicates that certain clones did not have an amino acid at the indicated position. Amino acid positions are numbered according to the Kabat numbering system as described below.

35 Figures 11A-11C show heavy chain and light chain HVR loop sequences of anti-polyubiquitin antibody molecules apu17-apu24 that specifically recognize K63-linked polyubiquitin and which were recognized by an antibody specific for pentahistidine, as described in Example 1(B).

The figures show the heavy chain HVR sequences, H1, H2, and H3, and light chain HVR sequence, L3. The designator “n.p.” indicates that certain clones did not have an amino acid at the indicated position. Amino acid positions are numbered according to the Kabat numbering system as described below.

5 Figure 12 depicts the binding interactions between various concentrations of the anti-polyubiquitin Fab apu09 and K48-linked or K63-linked polyubiquitin observed using BIACORE[®] analysis, as described in Example 1(C).

 Figures 13A – 13B depict the binding interactions between various concentrations of the anti-polyubiquitin Fab apu18 and K48-linked or K63-linked polyubiquitin observed using
10 BIACORE[®] analysis, as described in Example 1(C).

 Figures 14A-14F show heavy chain HVR loop sequences of second generation anti-polyubiquitin antibody molecules based on the sequence of Fab apu05, which specifically recognize K48-linked polyubiquitin, as described in Example 2. The figures show the heavy chain HVR sequences, H1, H2, and H3. The designator “n.p.” indicates that certain clones did not have an
15 amino acid at the indicated position. Amino acid positions are numbered according to the Kabat numbering system as described below. Shaded text indicates that the sequence is identical to the amino acid sequence of the corresponding HVR sequence in the Fab apu05. Bold text indicates that the antibody demonstrated strong binding in the phage ELISA assay described in Example 2.

 Figures 15A-15C show heavy chain HVR loop sequences of second generation anti-polyubiquitin antibody molecules based on the sequence of Fab apu18, which specifically recognize
20 K63-linked polyubiquitin, as described in Example 2. The figures show the heavy chain HVR sequences, H1, H2, and H3. The designator “n.p.” indicates that certain clones did not have an amino acid at the indicated position. Amino acid positions are numbered according to the Kabat numbering system as described below. Shaded text indicates that the sequence is identical to the
25 amino acid sequence of the corresponding HVR sequence in the Fab apu18. Bold text indicates that the antibody demonstrated strong binding in the phage ELISA assay described in Example 2.

 Figures 16A and 16B show the amino acid sequences of the heavy chain hypervariable regions of Fab molecules derived from mutagenized apu05 which specifically recognized K48-linked polyubiquitin (apu2.01-apu2.10) and which were recognized by an antibody specific for
30 pentahistidine, as described in Example 2. The figures show the heavy chain HVR sequences, H1, H2, and H3, and light chain HVR sequence, L3. The designator “n.p.” indicates that certain clones did not have an amino acid at the indicated position. Amino acid positions are numbered according to the Kabat numbering system as described below. Shaded text indicates that the sequence is identical to the amino acid sequence of the corresponding HVR sequence in the Fab apu05.

35 Figures 17A and 17B show the amino acid sequences of the heavy chain hypervariable regions of Fab molecules derived from mutagenized apu18 which specifically recognized K63-linked polyubiquitin (apu2.11-apu2.20) and which were recognized by an antibody specific for

pentahistidine, as described in Example 2. The figures show the heavy chain HVR sequences, H1, H2, and H3, and light chain HVR sequence, L3. The designator “n.p.” indicates that certain clones did not have an amino acid at the indicated position. Amino acid positions are numbered according to the Kabat numbering system as described below. Shaded text indicates that the sequence is identical to the amino acid sequence of the corresponding HVR sequence in the Fab apu18.

Figure 18 depicts the results of the phage ELISA assay described in Example 2 where the binding of each of second-generation Fabs apu2.01-2.20 to K48-linked polyubiquitin, K63-linked polyubiquitin, monoubiquitin, and bovine serum albumin was assessed.

Figures 19A and 19B depict the results of western blotting experiments described in Example 1. Figure 19A shows the binding of the Fabs produced from clones apu01 to apu15 to immobilized K48-linked tetraubiquitin. Figure 19B shows an absence of binding of the Fabs produced from clones apu18 to apu24 to immobilized K63-linked polyubiquitin.

Figures 20A and 20B depict the results of western blotting experiments described in Example 2. Figure 20A shows the binding of apu2.01-apu2.10 to immobilized K48-linked tetraubiquitin and the absence of binding to immobilized K63-linked diubiquitin. Figure 20B shows the binding of apu2.11-apu2.20 to immobilized K63-linked tetraubiquitin and the absence of binding to immobilized K48-linked diubiquitin.

Figures 21A and 21B show Western blots from immunoprecipitation experiments to detect the ubiquitination state of RIP, as described in Example 3. The blot in Figure 21A includes samples that were immunoprecipitated with apu2.16 IgG to capture K63-linked polyubiquitinated proteins. The blot in Figure 21B includes samples that were immunoprecipitated with apu2.07 IgG to capture K48-linked polyubiquitinated proteins. Both blots were stained with an anti-RIP antibody.

Figure 22 depicts the results of the phage ELISA assay described in Example 4 where the binding of each of third-generation clones apu3.01-3.12 to K48-linked polyubiquitin, K63-linked polyubiquitin, monoubiquitin, and bovine serum albumin was assessed.

Figures 23A and 23B show the amino acid sequences of the heavy chain hypervariable regions of clones derived from mutagenized apu2.16 which specifically recognized K63-linked polyubiquitin (apu3.01-apu3.12), as described in Example 4. The figures show the heavy chain HVR sequences, H1, H2, and H3. The designator “ND.” indicates that the sequence was not determined. Amino acid positions are numbered according to the Kabat numbering system as described below. Shaded text indicates that the sequence is identical to the amino acid sequence of the corresponding HVR sequence in apu2.16.

Figures 24A-24D depict the results of western blotting experiments described in Example 4. Figure 24A shows the binding of apu2.07 IgG to immobilized K48-linked tri- to heptaubiquitin and the absence of binding to immobilized K63-linked tri- to heptaubiquitin or monoubiquitin. Figure 24B shows the binding of apu3.07 IgG to immobilized K63-linked tri- to heptaubiquitin and the absence of binding to immobilized K48-linked tri- to heptaubiquitin or monoubiquitin. Figure 24C

shows concentration-dependent binding of apu2.07 IgG to immobilized K48-linked tetraubiquitin and the absence of binding to immobilized K63-linked tetraubiquitin. Figure 24D shows concentration-dependent binding of apu3.07 IgG to immobilized K63-linked tetraubiquitin and the absence of binding to immobilized K48-linked tetraubiquitin.

5 Figure 25 depicts the results of a western blotting experiment described in Example 4. The figure shows the binding of an anti-ubiquitin polyclonal antibody, apu2.07 IgG, and apu3.07 IgG to immobilized lysates from 293T cells treated (+) or untreated (-) with Velcade®.

Figures 26A-26C show the interactions between a K63-linked polyubiquitin-specific fab of the invention and K63-linked polyubiquitin as determined by crystallographic analysis. Figure 26A depicts the complex formed between K63-linked polyubiquitin-specific fab apu2.16 and a K63-linked diubiquitin. Apu2.16 is shown in ribbon diagram at the bottom of the figure while K63-linked diubiquitin is shown in globular form at the top of the figure. Figure 26B depicts the surface of K63-linked diubiquitin, with those residues within 4.5 Å of the fab colored dark grey and residues of interest labeled. Figure 26C depicts the surface of apu2.16, with those residues within 4.5 Å of the K63-linked ubiquitin dimer colored dark grey. The CDRs are labeled.

MODES FOR CARRYING OUT THE INVENTION

General Techniques

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, such as, "Molecular Cloning: A Laboratory Manual", third edition (Sambrook et al., 2001); "Oligonucleotide Synthesis" (M. J. Gait, ed., 1984); "Animal Cell Culture" (R. I. Freshney, ed., 1987); "Methods in Enzymology" (Academic Press, Inc.); "Current Protocols in Molecular Biology" (F. M. Ausubel et al., eds., 1987, and periodic updates); "PCR: The Polymerase Chain Reaction", (Mullis et al., ed., 1994); PCR 2: A Practical Approach (M. J. MacPherson, B. D. Hames and G.R. Taylor eds. (1995)); Harlow and Lane, eds. (1988) Antibodies, A Laboratory Manual; "A Practical Guide to Molecular Cloning" (Perbal Bernard V., 1988); and "Phage Display: A Laboratory Manual" (Barbas et al., 2001).

30

Definitions

As used herein, the terms “ubiquitin” and “monoubiquitin” are used interchangeably, and are defined as all species of native human and synthetic ubiquitin substantially similar to a 76-amino acid protein having at least one lysine residue at amino acid 6, amino acid 22, amino acid 27, amino acid
5 29, amino acid 33, amino acid 48, and/or amino acid 63.

As used herein, the term “polyubiquitin” is defined as all species of native human and synthetic polymeric chains of ubiquitin which fall within human and synthetic classes of different polymeric linkages of ubiquitin, including, but not limited to, K6-linked polyubiquitin, K22-linked polyubiquitin, K27-linked polyubiquitin, K29-linked polyubiquitin, K33-linked polyubiquitin, K48-
10 linked polyubiquitin and K63-linked polyubiquitin. Polyubiquitin may be of any length, and includes at least two ubiquitin moieties. Polyubiquitin is distinguished from tandem repeats of ubiquitin that are originally expressed as a single protein.

As used herein, the terms “K*-linked polyubiquitin” and “Lys*-linked polyubiquitin” are interchangeable, and refer to a polyubiquitin molecule comprising at least one isopeptide bond
15 between the C-terminus of one ubiquitin moiety and a lysine at position * in another ubiquitin moiety. For example, a “K63-linked polyubiquitin” is used interchangeably with a “Lys63-linked polyubiquitin”, and both terms refer to a polyubiquitin molecule comprising an isopeptide bond between the C-terminus of one of the ubiquitin moieties in the molecule and the lysine at position 63 in another ubiquitin moiety in the molecule.

As used herein, a statement that a first lysine linkage “differs” from a second lysine linkage indicates that the first lysine linkage between one ubiquitin moiety and another ubiquitin moiety involves a different lysine residue (e.g., K6, K22, K27, K29, K33, K48, and/or K63) than the second lysine linkage between one ubiquitin moiety and another ubiquitin moiety.

As used herein, the term “ubiquitin pathway” refers to a biochemical pathway in a cell or
25 reconstituted *in vitro* that includes ubiquitin and/or polyubiquitin.

An “isolated” antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with research, diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In one
30 embodiment, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by, for example, the Lowry method, and in some embodiments more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of, for example, a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using, for example, Coomassie blue or silver stain. Isolated
35 antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

As used herein, the terms “anti-ubiquitin antibody” and “anti-monoubiquitin antibody” are used interchangeably, and refer to an antibody that is capable of specifically binding to a ubiquitin molecule.

As used herein, the term “anti-polyubiquitin antibody” refers to an antibody that is capable of specifically binding to a polyubiquitin molecule.

As used herein, the term “anti-K48-linked polyubiquitin antibody” refers to an antibody that is capable of specifically binding to K48-linked polyubiquitin.

As used herein, the term “anti-K63-linked polyubiquitin antibody” refers to an antibody that is capable of binding to K63-linked polyubiquitin.

The phrase “substantially similar,” “substantially the same”, “equivalent”, or “substantially equivalent”, as used herein, denotes a sufficiently high degree of similarity between two numeric values (for example, one associated with a molecule and the other associated with a reference/comparator molecule) such that one of skill in the art would consider the difference between the two values to be of little or no biological and/or statistical significance within the context of the biological characteristic measured by said values (e.g., K_d values, anti-viral effects, etc.). The difference between said two values is, for example, less than about 50%, less than about 40%, less than about 30%, less than about 20%, and/or less than about 10% as a function of the value for the reference/comparator molecule.

The phrase “substantially reduced,” or “substantially different”, as used herein, denotes a sufficiently high degree of difference between two numeric values (generally one associated with a molecule and the other associated with a reference/comparator molecule) such that one of skill in the art would consider the difference between the two values to be of statistical significance within the context of the biological characteristic measured by said values (e.g., K_d values). The difference between said two values is, for example, greater than about 10%, greater than about 20%, greater than about 30%, greater than about 40%, and/or greater than about 50% as a function of the value for the reference/comparator molecule.

“Binding affinity” generally refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (e.g., an antibody) and its binding partner (e.g., an antigen). Unless indicated otherwise, as used herein, “binding affinity” refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (e.g., antibody and antigen). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K_d). Affinity can be measured by common methods known in the art, including those described herein. Low-affinity antibodies generally bind antigen slowly and tend to dissociate readily, whereas high-affinity antibodies generally bind antigen faster and tend to remain bound longer. A variety of methods of measuring binding affinity are known in the art, any of which can be used for purposes of the present invention. Specific illustrative embodiments are described in the following.

In one embodiment, the “Kd” or “Kd value” according to this invention is measured by a radiolabeled antigen binding assay (RIA) performed with the Fab version of an antibody of interest and its antigen as described by the following assay. Solution binding affinity of Fabs for antigen is measured by equilibrating Fab with a minimal concentration of (^{125}I)-labeled antigen in the presence of a titration series of unlabeled antigen, then capturing bound antigen with an anti-Fab antibody-coated plate (Chen, et al., (1999) *J. Mol Biol* 293:865-881). To establish conditions for the assay, microtiter plates (Dynex) are coated overnight with 5 $\mu\text{g}/\text{ml}$ of a capturing anti-Fab antibody (Cappel Labs) in 50 mM sodium carbonate (pH 9.6), and subsequently blocked with 2% (w/v) bovine serum albumin in PBS for two to five hours at room temperature (approximately 23°C). In a non-adsorbent plate (Nunc #269620), 100 pM or 26 pM [^{125}I]-antigen are mixed with serial dilutions of a Fab of interest (e.g., consistent with assessment of an anti-VEGF antibody, Fab-12, in Presta et al., (1997) *Cancer Res.* 57:4593-4599). The Fab of interest is then incubated overnight; however, the incubation may continue for a longer period (e.g., 65 hours) to insure that equilibrium is reached. Thereafter, the mixtures are transferred to the capture plate for incubation at room temperature (e.g., for one hour). The solution is then removed and the plate washed eight times with 0.1% Tween-20 in PBS. When the plates have dried, 150 $\mu\text{l}/\text{well}$ of scintillant (MicroScintTM-20; Packard) is added, and the plates are counted on a TopcountTM gamma counter (Packard) for ten minutes. Concentrations of each FAB that give less than or equal to 20% of maximal binding are chosen for use in competitive binding assays. According to another embodiment the Kd or Kd value is measured by using surface plasmon resonance assays using a BIAcoreTM-2000 or a BIAcoreTM-3000 (BIAcore, Inc., Piscataway, NJ) at 25°C with immobilized antigen CM5 chips at ~10 response units (RU). Briefly, carboxymethylated dextran biosensor chips (CM5, BIAcore Inc.) are activated with *N*-ethyl-*N'*-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and *N*-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 $\mu\text{g}/\text{ml}$ (~0.2 μM) before injection at a flow rate of 5 $\mu\text{l}/\text{minute}$ to achieve approximately 10 response units (RU) of coupled protein. Following the injection of antigen, 1 M ethanolamine is injected to block unreacted groups. For kinetics measurements, two-fold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with 0.05% TweenTM 20 (PBST) at 25°C at a flow rate of approximately 25 $\mu\text{l}/\text{min}$. Association rates (k_{on}) and dissociation rates (k_{off}) are calculated using a simple one-to-one Langmuir binding model (BIAcore Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (Kd) is calculated as the ratio $k_{\text{off}}/k_{\text{on}}$. See, e.g., Chen, Y., et al., (1999) *J. Mol Biol* 293:865-881. If the on-rate exceeds $10^6 \text{ M}^{-1} \text{ s}^{-1}$ by the surface plasmon resonance assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase or decrease in fluorescence emission intensity (excitation = 295 nm; emission = 340 nm, 16 nm band-pass) at 25°C of a 20 nM

anti-antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such as a stop-flow equipped spectrophotometer (Aviv Instruments) or a 8000-series SLM-Aminco spectrophotometer (ThermoSpectronic) with a stirred cuvette.

5 An "on-rate" or "rate of association" or "association rate" or " k_{on} " according to this invention can also be determined with the same surface plasmon resonance technique described above using a BIAcoreTM-2000 or a BIAcoreTM-3000 (BIAcore, Inc., Piscataway, NJ) at 25°C with immobilized antigen CM5 chips at ~10 response units (RU). Briefly, carboxymethylated dextran biosensor chips (CM5, BIAcore Inc.) are activated with *N*-ethyl-*N*'-(3-dimethylaminopropyl)-carbodiimide
10 hydrochloride (EDC) and *N*-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 µg/ml (~0.2 µM) before injection at a flow rate of 5 µl/minute to achieve approximately 10 response units (RU) of coupled protein. Following the injection of antigen, 1M ethanolamine is injected to block unreacted groups. For kinetics measurements, two-fold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with
15 0.05% Tween 20 (PBST) at 25°C at a flow rate of approximately 25 µl/min. Association rates (k_{on}) and dissociation rates (k_{off}) are calculated using a simple one-to-one Langmuir binding model (BIAcore Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgram. The equilibrium dissociation constant (K_d) was calculated as the ratio k_{off}/k_{on} . See, e.g., Chen, Y., et al., (1999) *J. Mol Biol* 293:865-881. However, if the on-rate exceeds $10^6 \text{ M}^{-1} \text{ s}^{-1}$
20 by the surface plasmon resonance assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase or decrease in fluorescence emission intensity (excitation = 295 nm; emission = 340 nm, 16 nm band-pass) at 25°C of a 20 nM anti-antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such as a stop-flow equipped spectrophotometer (Aviv Instruments) or a
25 8000-series SLM-Aminco spectrophotometer (ThermoSpectronic) with a stirred cuvette.

The term "vector," as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a phage vector. Another type of vector is a viral vector, wherein additional DNA
30 segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are
35 operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "recombinant vectors"). In general, expression vectors of utility in recombinant DNA techniques are often

in the form of plasmids. In the present specification, "plasmid" and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector.

"Polynucleotide," or "nucleic acid," as used interchangeably herein, refer to polymers of nucleotides of any length, and include DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a polymer by DNA or RNA polymerase, or by a synthetic reaction. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and their analogs. If present, modification to the nucleotide structure may be imparted before or after assembly of the polymer. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after synthesis, such as by conjugation with a label. Other types of modifications include, for example, "caps," substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, carbamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), those containing pendant moieties, such as, for example, proteins (e.g., nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotides(s). Further, any of the hydroxyl groups ordinarily present in the sugars may be replaced, for example, by phosphonate groups, phosphate groups, protected by standard protecting groups, or activated to prepare additional linkages to additional nucleotides, or may be conjugated to solid or semi-solid supports. The 5' and 3' terminal OH can be phosphorylated or substituted with amines or organic capping group moieties of from 1 to 20 carbon atoms. Other hydroxyls may also be derivatized to standard protecting groups. Polynucleotides can also contain analogous forms of ribose or deoxyribose sugars that are generally known in the art, including, for example, 2'-O-methyl-, 2'-O-allyl, 2'-fluoro- or 2'-azido-ribose, carbocyclic sugar analogs, α -anomeric sugars, epimeric sugars such as arabinose, xyloses or lyxoses, pyranose sugars, furanose sugars, sedoheptuloses, acyclic analogs and basic nucleoside analogs such as methyl riboside. One or more phosphodiester linkages may be replaced by alternative linking groups. These alternative linking groups include, but are not limited to, embodiments wherein phosphate is replaced by P(O)S ("thioate"), P(S)S ("dithioate"), "(O)NR₂" ("amidate"), P(O)R, P(O)OR', CO or CH₂ ("formacetal"), in which each R or R' is independently H or substituted or unsubstituted alkyl (1-20 C) optionally containing an ether (-O-) linkage, aryl, alkenyl, cycloalkyl, cycloalkenyl or araldyl. Not all linkages in a polynucleotide need be identical. The preceding description applies to all polynucleotides referred to herein, including RNA and DNA.

“Oligonucleotide,” as used herein, generally refers to short, generally single-stranded, generally synthetic polynucleotides that are generally, but not necessarily, less than about 200 nucleotides in length. The terms “oligonucleotide” and “polynucleotide” are not mutually exclusive. The description above for polynucleotides is equally and fully applicable to oligonucleotides.

5 "Antibodies" (Abs) and "immunoglobulins" (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific antigen, immunoglobulins include both antibodies and other antibody-like molecules which generally lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas.

10 The terms "antibody" and “immunoglobulin” are used interchangeably in the broadest sense and include monoclonal antibodies (*e.g.*, full length or intact monoclonal antibodies), polyclonal antibodies, monovalent, multivalent antibodies, multispecific antibodies (*e.g.*, bispecific antibodies so long as they exhibit the desired biological activity) and may also include certain antibody fragments (as described in greater detail herein). An antibody can be chimeric, human, humanized and/or
15 affinity matured.

The “variable region” or “variable domain” of an antibody refers to the amino-terminal domains of heavy or light chain of the antibody. These domains are generally the most variable parts of an antibody and contain the antigen-binding sites.

The term “variable” refers to the fact that certain portions of the variable domains differ
20 extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity-determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the
25 framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a beta-sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the beta-sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat et al., Sequences of
30 Proteins of Immunological Interest, Fifth Edition, National Institute of Health, Bethesda, MD (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

Papain digestion of antibodies produces two identical antigen-binding fragments, called “Fab”
35 fragments, each with a single antigen-binding site, and a residual “Fc” fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

“Fv” is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. In a two-chain Fv species, this region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. In a single-chain Fv species, one heavy- and one light-chain variable domain can be covalently linked by a flexible peptide linker such that the light and heavy chains can associate in a “dimeric” structure analogous to that in a two-chain Fv species. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The “light chains” of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

Depending on the amino acid sequences of the constant domains of their heavy chains, antibodies (immunoglobulins) can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known and described generally in, for example, Abbas et al. Cellular and Mol. Immunology, 4th ed. (2000). An antibody may be part of a larger fusion molecule, formed by covalent or non-covalent association of the antibody with one or more other proteins or peptides.

The terms “full length antibody,” “intact antibody” and “whole antibody” are used herein interchangeably, to refer to an antibody in its substantially intact form, not antibody fragments as defined below. The terms particularly refer to an antibody with heavy chains that contain the Fc region.

“Antibody fragments” comprise only a portion of an intact antibody, wherein the portion retains at least one, and as many as most or all, of the functions normally associated with that portion when present in an intact antibody. In one embodiment, an antibody fragment comprises an antigen binding site of the intact antibody and thus retains the ability to bind antigen. In another embodiment,

an antibody fragment, for example one that comprises the Fc region, retains at least one of the biological functions normally associated with the Fc region when present in an intact antibody, such as FcRn binding, antibody half life modulation, ADCC function and complement binding. In one embodiment, an antibody fragment is a monovalent antibody that has an *in vivo* half life substantially similar to an intact antibody. For example, such an antibody fragment may comprise on antigen binding arm linked to an Fc sequence capable of conferring *in vivo* stability to the fragment.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies. Such monoclonal antibody typically includes an antibody comprising a polypeptide sequence that binds a target, wherein the target-binding polypeptide sequence was obtained by a process that includes the selection of a single target binding polypeptide sequence from a plurality of polypeptide sequences. For example, the selection process can be the selection of a unique clone from a plurality of clones, such as a pool of hybridoma clones, phage clones or recombinant DNA clones. It should be understood that the selected target binding sequence can be further altered, for example, to improve affinity for the target, to humanize the target binding sequence, to improve its production in cell culture, to reduce its immunogenicity *in vivo*, to create a multispecific antibody, etc., and that an antibody comprising the altered target binding sequence is also a monoclonal antibody of this invention. In contrast to polyclonal antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody of a monoclonal antibody preparation is directed against a single determinant on an antigen. In addition to their specificity, the monoclonal antibody preparations are advantageous in that they are typically uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of techniques, including, for example, the hybridoma method (e.g., Kohler et al., Nature, 256: 495 (1975); Harlow et al., Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling et al., in: Monoclonal Antibodies and T-Cell hybridomas 563-681 (Elsevier, N.Y., 1981)), recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567), phage display technologies (See, e.g., Clackson et al., Nature, 352: 624-628 (1991); Marks et al., J. Mol. Biol. 222: 581-597 (1992); Sidhu et al., J. Mol. Biol. 338(2): 299-310 (2004); Lee et al., J. Mol. Biol. 340(5): 1073-1093 (2004); Fellouse, Proc. Natl. Acad. Sci. USA 101(34): 12467-12472 (2004); and Lee et al., J. Immunol. Methods 284(1-2): 119-132(2004), and technologies for producing human or human-like antibodies in animals that have parts or all of the human immunoglobulin loci or genes encoding human immunoglobulin sequences (see, e.g., WO98/24893;

WO96/34096; WO96/33735; WO91/10741; Jakobovits et al., Proc. Natl. Acad. Sci. USA 90: 2551 (1993); Jakobovits et al., Nature 362: 255-258 (1993); Bruggemann et al., Year in Immunol. 7:33 (1993); U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016; Marks et al., Bio.Technology 10: 779-783 (1992); Lonberg et al., Nature 368: 856-859 (1994); Morrison, Nature 368: 812-813 (1994); Fishwild et al., Nature Biotechnol. 14: 845-851 (1996); Neuberger, Nature Biotechnol. 14: 826 (1996) and Lonberg and Huszar, Intern. Rev. Immunol. 13: 65-93 (1995).

The monoclonal antibodies herein specifically include "chimeric" antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison *et al.*, Proc. Natl. Acad. Sci. USA 81:6851-6855 (1984)).

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. In one embodiment, a humanized antibody is a human immunoglobulin (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and/or capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally will also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, Nature 321:522-525 (1986); Riechmann *et al.*, Nature 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol. 2:593-596 (1992). See also the following review articles and references cited therein: Vaswani and Hamilton, Ann. Allergy, Asthma & Immunol. 1:105-115 (1998); Harris, Biochem. Soc. Transactions 23:1035-1038 (1995); Hurle and Gross, Curr. Op. Biotech. 5:428-433 (1994).

The term "hypervariable region", "HVR", or "HV", when used herein refers to the regions of an antibody variable domain which are hypervariable in sequence and/or form structurally defined loops. Generally, antibodies comprise six hypervariable regions; three in the VH (H1, H2, H3), and three in the VL (L1, L2, L3). A number of hypervariable region delineations are in use and are encompassed herein. The Kabat Complementarity Determining Regions (CDRs) are based on sequence variability and are the most commonly used (Kabat *et al.*, Sequences of Proteins of

Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). Chothia refers instead to the location of the structural loops (Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). The AbM hypervariable regions represent a compromise between the Kabat CDRs and Chothia structural loops, and are used by Oxford Molecular's AbM antibody modeling software. The “contact” hypervariable regions are based on an analysis of the available complex crystal structures. The residues from each of these hypervariable regions are noted below.

Loop	Kabat	AbM	Chothia	Contact
----	-----	---	-----	-----
L1	L24-L34	L24-L34	L26-L32	L30-L36
L2	L50-L56	L50-L56	L50-L52	L46-L55
L3	L89-L97	L89-L97	L91-L96	L89-L96
H1	H31-H35B	H26-H35B	H26-H32	H30-H35B
	(Kabat Numbering)			
H1	H31-H35	H26-H35	H26-H32	H30-H35
	(Chothia Numbering)			
H2	H50-H65	H50-H58	H53-H55	H47-H58
H3	H95-H102	H95-H102	H96-H101	H93-H101

Hypervariable regions may comprise “extended hypervariable regions” as follows: 24-36 or 24-34 (L1), 46-56 or 50-56 (L2) and 89-97 or 89-96 (L3) in the VL and 26-35 (H1), 50-65 or 49-65 (H2) and 93-102, 94-102, or 95-102 (H3) in the VH. The variable domain residues are numbered according to Kabat et al., *supra*, for each of these definitions.

"Framework" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined.

The term “variable domain residue numbering as in Kabat” or “amino acid position numbering as in Kabat,” and variations thereof, refers to the numbering system used for heavy chain variable domains or light chain variable domains of the compilation of antibodies in Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991). Using this numbering system, the actual linear amino acid sequence may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, a FR or HVR of the variable domain. For example, a heavy chain variable domain may include a single amino acid insert (residue 52a according to Kabat) after residue 52 of H2 and inserted residues (e.g. residues 82a, 82b, and 82c, etc. according to Kabat) after heavy chain FR residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a “standard” Kabat numbered sequence.

“Single-chain Fv” or “scFv” antibody fragments comprise the VH and VL domains of antibody, wherein these domains are present in a single polypeptide chain. Generally, the scFv

polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv see Pluckthun, in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

5 The term “diabodies” refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) in the same polypeptide chain (VH-VL). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are
10 described more fully in, for example, EP 404,097; WO93/1161; and Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90: 6444-6448 (1993).

A “human antibody” is one which possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies as disclosed herein. This definition of a human antibody specifically
15 excludes a humanized antibody comprising non-human antigen-binding residues.

An “affinity matured” antibody is one with one or more alterations in one or more HVRs thereof which result in an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). In one embodiment, an affinity matured antibody has nanomolar or even picomolar affinities for the target antigen. Affinity matured
20 antibodies are produced by procedures known in the art. Marks *et al.* *Bio/Technology* 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas *et al.* *Proc Nat. Acad. Sci. USA* 91:3809-3813 (1994); Schier *et al.* *Gene* 169:147-155 (1995); Yelton *et al.* *J. Immunol.* 155:1994-2004 (1995); Jackson *et al.*, *J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al.*, *J. Mol. Biol.* 226:889-896
25 (1992).

A “blocking” antibody or an “antagonist” antibody is one which inhibits or reduces biological activity of the antigen it binds. Certain blocking antibodies or antagonist antibodies substantially or completely inhibit the biological activity of the antigen.

An “agonist antibody”, as used herein, is an antibody which mimics at least one of the
30 functional activities of a polypeptide of interest.

A “disorder” is any condition that would benefit from treatment with an antibody of the invention. This includes chronic and acute disorders or diseases including those pathological conditions which predispose the mammal to the disorder in question. Non-limiting examples of disorders to be treated herein include cancer, muscular disorders, ubiquitin-pathway-related genetic
35 disorders, immune/inflammatory disorders, neurological disorders, and other ubiquitin pathway-related disorders.

The terms “cell proliferative disorder” and “proliferative disorder” refer to disorders that are associated with some degree of abnormal cell proliferation. In one embodiment, the cell proliferative disorder is cancer.

“Tumor,” as used herein, refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues. The terms “cancer,” “cancerous,” “cell proliferative disorder,” “proliferative disorder” and “tumor” are not mutually exclusive as referred to herein.

The terms “cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth/proliferation. Examples of cancer include, but are not limited to, carcinoma, lymphoma (e.g., Hodgkin’s and non-Hodgkin’s lymphoma), blastoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney cancer, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, leukemia and other lymphoproliferative disorders, and various types of head and neck cancer.

The term “muscular disorder” refers to or describes the physiological condition in muscle-containing animals that is typically characterized by deterioration or weakening of skeletal and/or smooth muscle such that normal muscular function is significantly reduced. Examples of muscular disorders include, but are not limited to, muscular dystrophy, multiple sclerosis, amyotrophic lateral sclerosis, Isaac’s syndrome; stiff-person syndrome; familial periodic paralyses, myopathy, myotonia, rhabdomyolyses, muscle atrophy, and various types of muscle weakness and muscle rigidity.

The term “ubiquitin pathway-related genetic disorder” refers to or describes a genetically-based disorder that is typically characterized by or contributed to by aberrant functioning of the ubiquitin pathway. Examples of ubiquitin pathway-related genetic disorders include, but are not limited to, cystic fibrosis, Angelman’s syndrome, and Liddle syndrome.

The terms “neurological disorder” or “neurological disease” refer to or describe a disease or disorder of the central and/or peripheral nervous system in mammals that is typically characterized by deterioration of nervous tissue or deterioration of communication between cells in nervous tissue. Examples of neurological disorders include, but are not limited to, neurodegenerative diseases (including, but not limited to, Lewy body disease, postpoliomyelitis syndrome, Shy-Draeger syndrome, olivopontocerebellar atrophy, Parkinson’s disease, multiple system atrophy, striatonigral degeneration, tauopathies (including, but not limited to, Alzheimer disease and supranuclear palsy), prion diseases (including, but not limited to, bovine spongiform encephalopathy, scrapie, Creutzfeldt-Jakob syndrome, kuru, Gerstmann-Straussler-Scheinker disease, chronic wasting disease, and fatal

familial insomnia), bulbar palsy, motor neuron disease, and nervous system heterodegenerative disorders (including, but not limited to, Canavan disease, Huntington's disease, neuronal ceroid-lipofuscinosis, Alexander's disease, Tourette's syndrome, Menkes kinky hair syndrome, Cockayne syndrome, Halervorden-Spatz syndrome, lafora disease, Rett syndrome, hepatolenticular
 5 degeneration, Lesch-Nyhan syndrome, and Unverricht-Lundborg syndrome), dementia (including, but not limited to, Pick's disease, and spinocerebellar ataxia.

The terms "inflammatory disorder" and "immune disorder" refer to or describe disorders caused by aberrant immunologic mechanisms and/or aberrant cytokine signaling. Examples of inflammatory and immune disorders include, but are not limited to, autoimmune diseases,
 10 immunologic deficiency syndromes, and hypersensitivity. An "autoimmune disease" herein is a non-malignant disease or disorder arising from and directed against an individual's own tissues. The autoimmune diseases herein specifically exclude malignant or cancerous diseases or conditions, especially excluding B cell lymphoma, acute lymphoblastic leukemia (ALL), chronic lymphocytic leukemia (CLL), Hairy cell leukemia and chronic myeloblastic leukemia. Examples of autoimmune
 15 diseases or disorders include, but are not limited to, inflammatory responses such as inflammatory skin diseases including psoriasis and dermatitis (*e.g.* atopic dermatitis); systemic scleroderma and sclerosis; responses associated with inflammatory bowel disease (such as Crohn's disease and ulcerative colitis); respiratory distress syndrome (including adult respiratory distress syndrome; ARDS); dermatitis; meningitis; encephalitis; uveitis; colitis; glomerulonephritis; allergic conditions
 20 such as eczema and asthma and other conditions involving infiltration of T cells and chronic inflammatory responses; atherosclerosis; leukocyte adhesion deficiency; rheumatoid arthritis; systemic lupus erythematosus (SLE) (including but not limited to lupus nephritis, cutaneous lupus); diabetes mellitus (*e.g.* Type I diabetes mellitus or insulin dependent diabetes mellitus); multiple sclerosis; Reynaud's syndrome; autoimmune thyroiditis; Hashimoto's thyroiditis; allergic
 25 encephalomyelitis; Sjogren's syndrome; juvenile onset diabetes; and immune responses associated with acute and delayed hypersensitivity mediated by cytokines and T-lymphocytes typically found in tuberculosis, sarcoidosis, polymyositis, granulomatosis and vasculitis; pernicious anemia (Addison's disease); diseases involving leukocyte diapedesis; central nervous system (CNS) inflammatory disorder; multiple organ injury syndrome; hemolytic anemia (including, but not limited to
 30 cryoglobulinemia or Coombs positive anemia) ; myasthenia gravis; antigen-antibody complex mediated diseases; anti-glomerular basement membrane disease; antiphospholipid syndrome; allergic neuritis; Graves' disease; Lambert-Eaton myasthenic syndrome; pemphigoid bullous; pemphigus; autoimmune polyendocrinopathies; Reiter's disease; stiff-man syndrome; Behcet disease; giant cell arteritis; immune complex nephritis; IgA nephropathy; IgM polyneuropathies; immune thrombocytopenic
 35 purpura (ITP) or autoimmune thrombocytopenia, etc.

Examples of immunologic deficiency syndromes include, but are not limited to, ataxia telangiectasia, leukocyte-adhesion deficiency syndrome, lymphopenia, dysgammaglobulinemia, HIV

or deltaretrovirus infections, common variable immunodeficiency, severe combined immunodeficiency, phagocyte bactericidal dysfunction, agammaglobulinemia, DiGeorge syndrome, and Wiskott-Aldrich syndrome. Examples of hypersensitivity include, but are not limited to, allergies, asthma, dermatitis, hives, anaphylaxis, Wissler's syndrome, and thrombocytopenic purpura.

As used herein, "treatment" refers to clinical intervention in an attempt to alter the natural course of the individual or cell being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Desirable effects of treatment include preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing or decreasing inflammation and/or tissue/organ damage, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. In some embodiments, antibodies of the invention are used to delay development of a disease or disorder.

An "individual" is a vertebrate. In certain embodiments, the vertebrate is a mammal. Mammals include, but are not limited to, farm animals (such as cows), sport animals, pets (such as cats, dogs, and horses), primates, mice and rats. In certain embodiments, the vertebrate is a human.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, etc. In certain embodiments, the mammal is human.

An "effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result.

A "therapeutically effective amount" of a substance/molecule of the invention may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the substance/molecule, to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the substance/molecule are outweighed by the therapeutically beneficial effects. A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically but not necessarily, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount would be less than the therapeutically effective amount.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g., At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³², Pb²¹² and radioactive isotopes of Lu), chemotherapeutic agents (e.g., methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), doxorubicin, melphalan, mitomycin C, chlorambucil, daunorubicin or other intercalating agents, enzymes and fragments thereof such as nucleolytic enzymes, antibiotics, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof, and the various antitumor or anticancer

agents disclosed below. Other cytotoxic agents are described below. A tumoricidal agent causes destruction of tumor cells.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as 5 benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); delta-9-tetrahydrocannabinol (dronabinol, MARINOL®); beta-lapachone; lapachol; colchicines; betulinic 10 acid; a camptothecin (including the synthetic analogue topotecan (HYCAMTIN®), CPT-11 (irinotecan, CAMPTOSAR®), acetylcamptothecin, scoplectin, and 9-aminocamptothecin); bryostatin; callistatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); podophyllotoxin; podophyllinic acid; teniposide; cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and 15 CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e. g., calicheamicin, 20 especially calicheamicin gammaII and calicheamicin omegaII (see, e.g., Agnew, Chem Intl. Ed. Engl., 33: 183-186 (1994)); dynemicin, including dynemicin A; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabacin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5- 25 oxo-L-norleucine, ADRIAMYCIN® doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as 30 methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitiostanol, mepitiothane, testolactone; anti- adrenals such as 35 aminogluthimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; an epothilone;

etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, OR); razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine (ELDISINE®, FILDESIN®); dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); thiotepa; taxoids, e.g., TAXOL® paclitaxel (Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAXANETM Cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumburg, Illinois), and TAXOTERE® doxorubicin (Rhône-Poulenc Rorer, Antony, France); chloranbucil; gemcitabine (GEMZAR®); 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine (VELBAN®); platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine (ONCOVIN®); oxaliplatin; leucovorin; vinorelbine (NAVELBINE®); novantrone; edatrexate; daunomycin; aminopterin; ibandronate; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine (XELODA®); pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of cyclophosphamide, doxorubicin, vincristine, and prednisolone, and FOLFOX, an abbreviation for a treatment regimen with oxaliplatin (ELOXATINTM) combined with 5-FU and leucovorin.

Also included in this definition are anti-hormonal agents that act to regulate, reduce, block, or inhibit the effects of hormones that can promote the growth of cancer, and are often in the form of systemic, or whole-body treatment. They may be hormones themselves. Examples include anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX® tamoxifen), EVISTA® raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON® toremifene; anti-progesterones; estrogen receptor down-regulators (ERDs); agents that function to suppress or shut down the ovaries, for example, leutinizing hormone-releasing hormone (LHRH) agonists such as LUPRON® and ELIGARD® leuprolide acetate, goserelin acetate, buserelin acetate and triptorelin; other anti-androgens such as flutamide, nilutamide and bicalutamide; and aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, MEGASE® megestrol acetate, AROMASIN® exemestane, formestane, fadrozole, RIVISOR® vorozole, FEMARA® letrozole, and ARIMIDEX® anastrozole. In addition, such definition of chemotherapeutic agents includes bisphosphonates such as clodronate (for example, BONEFOS® or OSTAC®), DIDROCAL® etidronate, NE-58095, ZOMETA® zoledronic acid/zoledronate, FOSAMAX® alendronate, AREDIA® pamidronate, SKELID® tiludronate, or ACTONEL® risedronate; as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine

analog); antisense oligonucleotides, particularly those that inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Raf, H-Ras, and epidermal growth factor receptor (EGF-R); vaccines such as THERATOPE® vaccine and gene therapy vaccines, for example, ALLOVECTIN® vaccine, LEUVECTIN® vaccine, and VAXID® vaccine; LURTOTECAN® topoisomerase 1 inhibitor; ABARELIX® rmRH; lapatinib ditosylate (an ErbB-2 and EGFR dual tyrosine kinase small-molecule inhibitor also known as GW572016); and pharmaceutically acceptable salts, acids or derivatives of any of the above.

Compositions and Methods of Making Same

The present invention provides antibodies that bind specifically to polyubiquitin, but not to monoubiquitin. More specifically, antibodies that are capable of binding specifically to a polyubiquitin comprising a first lysine linkage but not to a polyubiquitin comprising a second, different, lysine linkage are provided.

In one aspect, the invention provides an antibody comprising an HVR-H1 region comprising the sequence of at least one of SEQ ID NOs: 1-25, 81-89, 151-175, 229-239, 265-279, 329-336, 392-459, 599-629, 695-704, 739-748, and 789-799. In one aspect, the invention provides an antibody comprising a HVR-H1 region consensus sequence selected from SEQ ID NOs: 26, 90, 176, 240, 280, 337, 460, 630, 705, 749, and 800. In one aspect, the invention provides an antibody comprising a HVR-H2 region comprising the sequence of at least one of SEQ ID NOs: 27-51, 91-99, 177-201, 241-251, 281-295, 338-345, 461-528, 631-661, 706-715, 750-759, and 801-811. In one aspect, the invention provides an antibody comprising a HVR-H2 region consensus sequence selected from SEQ ID NOs: 52, 100, 202, 252, 296, 346, 529, 662, 716, 760, and 812. In one aspect, the invention provides an antibody comprising a HVR-H3 region comprising the sequence of at least one of SEQ ID NOs: 53-77, 101-109, 203-227, 253-263, 297-311, 347-354, 530-597, 663-693, 717-726, 761-770, and 813-823. In one aspect, the invention provides an antibody comprising a HVR-H3 region consensus sequence selected from SEQ ID NOs: 78, 110, 228, 264, 312, 355, 598, 694, 727, 771, and 824.

In one aspect, the invention provides an antibody comprising a HVR-H1 region comprising the sequence of at least one of SEQ ID NOs: 1-26, 81-90, 151-176, 229-240, 265-280, 329-337, 392-460, 599-630, 695-705, 739-749, and 789-800, and an HVR-H2 region comprising the sequence of at least one of SEQ ID NOs: 27-52, 91-100, 177-202, 241-252, 281-296, 338-346, 461-529, 631-662, 706-716, 750-760, and 801-812. In one aspect, the invention provides an antibody comprising a HVR-H1 region comprising the sequence of at least one of SEQ ID NOs: 1-26, 81-90, 151-176, 229-240, 265-280, 329-337, 392-460, 599-630, 695-705, 739-749, and 789-800, and an HVR-H3 region comprising the sequence of at least one of SEQ ID NOs: 53-78, 101-110, 203-228, 253-264, 297-312, 347-355, 530-598, 663-694, 717-727, 761-771, and 813-824. In one aspect, the invention provides an antibody comprising a HVR-H2 region comprising the sequence of at least one of SEQ ID NO: 27-52,

91-100, 177-202, 241-252, 281-296, 338-346, 461-529, 631-662, 706-716, 750-760, and 801-812, and an HVR-H3 region comprising the sequence of at least one of SEQ ID NOs: 53-78, 101-110, 203-228, 253-264, 297-312, 347-355, 530-598, 663-694, 717-727, 761-771, and 813-824.

In one aspect, the invention provides an antibody comprising a HVR-L3 region comprising the sequence of at least one of SEQ ID NOs: 313-327, 356-363, 728-737, and 772-781. In one aspect, the invention provides an antibody comprising a HVR-L3 region consensus sequence selected from SEQ ID NOs: 328, 364, 738, and 782. In one embodiment, the invention provides an antibody comprising a HVR-L3 region comprising the sequence of at least one of SEQ ID NOs: 313-328, 356-364, 728-738, and 772-782, and further comprising at least one HVR-H1, HVR-H2 or HVR-H3 sequence selected from SEQ ID NOs: 1-26, 81-90, 151-176, 229-240, 265-280, 329-337, 392-460, 599-630, 695-705, 739-749, and 789-800; SEQ ID NOs: 27-52, 91-100, 177-202, 241-252, 281-296, 338-346, 461-529, 631-662, 706-716, 750-760, and 801-812; and SEQ ID NOs: 53-78, 101-110, 203-228, 253-264, 297-312, 347-355, 530-598, 663-694, 717-727, 761-771, and 813-824, respectively.

In one aspect, the invention provides an antibody comprising at least one, at least two, at least three, or all four of the following:

- (i) an HVR-H1 sequence comprising at least one sequence of SEQ ID NOs: 1-26, 81-90, 151-176, 229-240, 265-280, 329-337, 392-460, 599-630, 695-705, 739-749, and 789-800;
- (ii) an HVR-H2 sequence comprising at least one sequence of SEQ ID NOs: 27-52, 91-100, 177-202, 241-252, 281-296, 338-346, 461-529, 631-662, 706-716, 750-760, and 801-812;
- (iii) an HVR-H3 sequence comprising at least one sequence of SEQ ID NOs: 53-78, 101-110, 203-228, 253-264, 297-312, 347-355, 530-598, 663-694, 717-727, 761-771, and 813-824;
- (iv) an HVR-L3 sequence comprising at least one sequence of SEQ ID NOs: 313-328, 356-364, 728-738, and 772-782.

In one aspect, the invention provides an antibody that specifically binds K48-linked polyubiquitin with high affinity but binds K63-linked polyubiquitin with substantially reduced affinity, comprising at least one, at least two, at least three, or all four of the following:

- (i) an HVR-H1 sequence comprising at least one sequence of SEQ ID NOs: 1-26, 151-176, 265-280, 392-460, and 695-705;
- (ii) an HVR-H2 sequence comprising at least one sequence of SEQ ID NOs: 27-52, 177-202, 281-296, 461-529, and 706-716;
- (iii) an HVR-H3 sequence comprising at least one sequence of SEQ ID NOs: 53-78, 203-228, 297-312, 530-598, and 717-727; and
- (iv) an HVR-L3 sequence comprising at least one sequence of SEQ ID NOs: 313-328 and 728-738.

In one aspect, the invention provides an antibody that specifically binds K63-linked polyubiquitin with high affinity but binds K48-linked polyubiquitin with substantially reduced affinity, comprising at least one, at least two, at least three, or all four of the following:

- (i) an HVR-H1 sequence comprising at least one sequence of SEQ ID NOs: 81-90, 229-240, 329-337, 599-630, 739-749, and 789-800;
- (ii) an HVR-H2 sequence comprising at least one sequence of SEQ ID NOs: 91-100, 241-252, 338-346, 631-662, 750-760, and 801-812;
- 5 (iii) an HVR-H3 sequence comprising at least one sequence of SEQ ID NOs: 101-110, 253-264, 347-355, 663-694, 761-771, and 813-824;
- (iv) an HVR-L3 sequence comprising at least one sequence of SEQ ID NOs: 356-364 and 772-782.

The amino acid sequences of SEQ ID NOs: 1-78, 81-106-149, 151-364, 392-782, and 789-824 are numbered with respect to individual HVR (i.e., H1, H2, H3, L3) as indicated in Figures 2, 3, 8, 9, 10, 11, 14, 15, 16, 17, and 22, the numbering being consistent with the Kabat numbering system as described below. In one embodiment, an antibody of the invention comprises one, two, three, or all of the HVR sequences of (i)-(iv) above, and HVR-L1 and/or HVR-L2 comprising a Kabat consensus sequence (e.g., SEQ ID NO: 79 (HVR-L1) and 80 (HVR-L2)).

15 In one aspect, the invention provides antibodies comprising heavy chain HVR sequences as depicted in Figures 2, 3, 8, 9, 10, 11, 14, 15, 16, 17, and 22. In one embodiment, the antibodies further comprise light chain HVR sequences as depicted in Figures 10, 11, 16, and 17.

Some embodiments of antibodies of the invention comprise a light chain variable domain of humanized 4D5 antibody (huMAb4D5-8) (HERCEPTIN®, Genentech, Inc., South San Francisco, CA, USA) (also referred to in U.S. Pat. No. 6,407,213 and Lee et al., J. Mol. Biol. (2004), 20 340(5):1073-93) as depicted in SEQ ID NO: 783 below.

1 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val
 Thr Ile Thr Cys Arg Ala Ser Gln **Asp** Val **Asn** **Thr** Ala Val Ala Trp Tyr Gln Gln Lys
 Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ser Ala Ser **Phe** Leu Tyr Ser Gly Val Pro
 25 Ser Arg Phe Ser Gly Ser **Arg** Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln
 Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln **His** Tyr Thr Thr Pro Pro Thr Phe Gly
 Gln Gly Thr Lys Val Glu Ile Lys 107 (SEQ ID NO: 783) (HVR residues are underlined)

In one embodiment, the huMAb4D5-8 light chain variable domain sequence is modified at one or more of positions 28, 30, 31, 53, 66, and 91 (Asp, Asn, Thr, Phe, Arg, and His as indicated in bold/italics above, respectively). In one embodiment, the modified huMAb4D5-8 sequence comprises Ser in position 28, Ser in position 30, Ser in position 31, Ser in position 53, Gly in position 66, and/or Ser in position 91. Accordingly, in one embodiment, an antibody of the invention comprises a light chain variable domain comprising the sequence depicted in SEQ ID NO: 784 below:

1 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val
 Thr Ile Thr Cys Arg Ala Ser Gln **Ser** Val **Ser** **Ser** Ala Val Ala Trp Tyr Gln Gln Lys

Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ser Ala Ser **Ser** Leu Tyr Ser Gly Val Pro
 Ser Arg Phe Ser Gly Ser **Gly** Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln
 Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln **Ser** Tyr Thr Thr Pro Pro Thr Phe Gly
 Gln Gly Thr Lys Val Glu Ile Lys 107 (SEQ ID NO: 784) (HVR residues are
 5 underlined)

Substituted residues with respect to huMAb4D5-8 are indicated in bold/italics above.

Antibodies of the invention can comprise any suitable framework variable domain sequence, provided binding activity to polyubiquitin including a particular lysine linkage is substantially retained. For example, in some embodiments, antibodies of the invention comprise a human subgroup
 10 III heavy chain framework consensus sequence. In one embodiment of these antibodies, the framework consensus sequence comprises substitution at position 71, 73 and/or 78. In some embodiments of these antibodies, position 71 is A, 73 is T and/or 78 is A. In one embodiment, these antibodies comprise heavy chain variable domain framework sequences of huMAb4D5-8 (HERCEPTIN[®], Genentech, Inc., South San Francisco, CA, USA) (also referred to in U.S. Pat. Nos.
 15 6,407,213 & 5,821,337, and Lee et al., J. Mol. Biol. (2004), 340(5):1073-93). In one embodiment, these antibodies further comprise a human κI light chain framework consensus sequence. In one embodiment, these antibodies comprise at least one, two or all of the light chain HVR sequences of SEQ ID NOs: 79, 80, 313-328, 356-364, 728-738, and 772-78. In one embodiment, these antibodies comprise light chain HVR sequences of huMAb4D5-8 as described in U.S. Pat. Nos. 6,407,213 &
 20 5,821,337.) In one embodiment, these antibodies comprise light chain variable domain sequences of huMAb4D5-8 (SEQ ID NO: 783 and 784) (HERCEPTIN[®], Genentech, Inc., South San Francisco, CA, USA) (also referred to in U.S. Pat. Nos. 6,407,213 & 5,821,337, and Lee et al., J. Mol. Biol. (2004), 340(5):1073-93).

In one embodiment, an antibody of the invention comprises a heavy chain variable domain,
 25 wherein the framework sequence comprises the sequence of at least one of SEQ ID NOs: 111-129, 138-141, 146-149, and 839-895, and HVR H1, H2 and H3 sequences are selected from at least one of SEQ ID NOs: 1-26, 81-90, 151-176, 229-240, 265-280, 329-337, 392-460, 599-630, 695-705, 739-749, and 789-800; 27-52, 91-100, 177-202, 241-252, 281-296, 338-346, 461-529, 631-662, 706-716, 750-760, and 801-812; and 53-78, 101-110, 203-228, 253-264, 297-312, 347-355, 530-598, 663-694,
 30 717-727, 761-771, and 813-824, respectively. In one embodiment, an antibody of the invention comprises a light chain variable domain, wherein the framework sequence comprises the sequence of at least one of SEQ ID NOs: 130-133, 134-137, 142-145, and 896-907, the HVR-L1 sequence is SEQ ID NO: 79, the HVR-L2 sequence is SEQ ID NO: 80, and the HVR-L3 sequence is selected from at least one of SEQ ID NOs: 313-328, 356-364, 728-738, and 772-782.

35 In one embodiment, an antibody of the invention comprises a heavy chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 111-129 and 839-895, and HVR H1, H2 and H3 sequences are SEQ ID NO: 1, 27, and 53, respectively (clone 48-1).

Similarly, in other embodiments, antibodies of each of clones 48-2 through 48-118, clones 63-1 through 63-51, Fabs apu01 through apu24, Fabs apu2.01 through apu2.20, and clones apu3.01 through 3.11 comprise a heavy chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 111-129 and 839-895, and HVR-H1, HVR-H2, and HVR-H3 sequences are those sequences specifically enumerated for each clone or Fab in Figures 2, 3, 8-11, 14-17, and 22. In one embodiment, an antibody of the invention comprises a light chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 130-133 and 896-907, and HVR L1, L2 and L3 sequences are SEQ ID NOs: 79, 80, and 313, respectively (Fab apu01). Similarly, in other embodiments, antibodies of each of Fabs apu01 through apu24 and Fabs apu2.01 through apu2.20 comprise a light chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 130-133 and 896-907, and HVR-L1 is SEQ ID NO: 79, HVR-L2 is SEQ ID NO: 80, and HVR-L3 sequences are those sequences specifically enumerated for each Fab in Figures 10, 11C, 16B, and 17B.

In one embodiment, an antibody of the invention comprises a heavy chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 138-141, and HVR H1, H2 and H3 sequences are SEQ ID NO: 1, 27, and 53, respectively (clone 48-1). Similarly, in other embodiments, antibodies of each of clones 48-2 through 48-118, clones 63-1 through 63-51, Fabs apu01 through apu24, Fabs apu2.01 through apu2.20, and clones apu3.01-3.11 comprise a heavy chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 138-141, and HVR-H1, HVR-H2, and HVR-H3 sequences are those sequences specifically enumerated for each clone or Fab in Figures 2, 3, 8-11, 14-17, and 22. In one embodiment, an antibody of the invention comprises a light chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 134-137, and HVR L1, L2 and L3 sequences are SEQ ID NOs: 79, 80, and 313, respectively (Fab apu01). Similarly, in other embodiments, antibodies of each of Fabs apu01 through apu24 and Fabs apu2.01 through apu2.20 comprise a light chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 134-137, and HVR-L1 is SEQ ID NO: 79, HVR-L2 is SEQ ID NO: 80, and HVR-L3 sequences are those sequences specifically enumerated for each Fab in Figures 10, 11C, 16B, and 17B.

In one embodiment, an antibody of the invention comprises a heavy chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 146-149, and HVR H1, H2 and H3 sequences are SEQ ID NO: 1, 27, and 53, respectively (clone 48-1). Similarly, in other embodiments, antibodies of each of clones 48-2 through 48-118, clones 63-1 through 63-51, Fabs apu01 through apu24, Fabs apu2.01 through apu2.20, and clones apu3.01 through 3.11 comprise a heavy chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 146-149, and HVR-H1, HVR-H2, and HVR-H3 sequences are those sequences specifically enumerated for each clone or Fab in Figures 2, 3, 8-11, 14-17, and 22. In one embodiment, an antibody of the invention comprises a light chain variable domain, wherein the

framework sequence comprises at least one sequence of SEQ ID NOs: 142-145, and HVR L1, L2 and L3 sequences are SEQ ID NOs: 79, 80, and 313, respectively (Fab apu01). Similarly, in other embodiments, antibodies of each of Fabs apu01 through apu24 and Fabs apu2.01 through apu2.20 comprise a light chain variable domain, wherein the framework sequence comprises at least one sequence of SEQ ID NOs: 142-145, and HVR-L1 is SEQ ID NO: 79, HVR-L2 is SEQ ID NO: 80, and HVR-L3 sequences are those sequences specifically enumerated for each Fab in Figures 10, 11C, 16B, and 17B.

In one embodiment, an antibody of the invention is affinity matured to obtain the target binding affinity desired. In one example, an affinity matured antibody of the invention which specifically binds to K48-linked polyubiquitin with high affinity but binds to K63-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H1 amino acid positions 29, 30, 33, and 34. In another example, an affinity matured antibody of the invention which specifically binds to K48-linked polyubiquitin with high affinity but binds to K63-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H2 amino acid positions 52 and 52a. In another example, an affinity matured antibody of the invention which specifically binds to K48-linked polyubiquitin with high affinity but binds to K63-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H3 amino acid positions 99, 100, 100a, and 100b. In another example, an affinity matured antibody of the invention which specifically binds to K48-linked polyubiquitin with high affinity but binds to K63-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H3 amino acid positions 95-100, 100a, and 100b. In another example, an affinity matured antibody of the invention which specifically binds to K48-linked polyubiquitin with high affinity but binds to K63-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-L3 amino acid positions 91 and 96. In another example, an affinity matured antibody of the invention which specifically binds to K63-linked polyubiquitin with high affinity but binds to K48-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H1 amino acid positions 29-34. In another example, an affinity matured antibody of the invention which specifically binds to K63-linked polyubiquitin with high affinity but binds to K48-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H2 amino acid positions 50, 52, 52a, 53-56, and 58. In another example, an affinity matured antibody of the invention which specifically binds to K63-linked polyubiquitin with high affinity but binds to K48-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H3 amino acid positions 95-100, 100a, 100b, and 100c. In another example, an affinity matured antibody of the invention which specifically binds to K63-linked polyubiquitin with high affinity but binds to K48-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-L3 amino acid positions 91-95, 95a, and 95b.

In another example, an affinity matured antibody of the invention which specifically binds to K63-linked polyubiquitin with high affinity but binds to K48-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H1 amino acid positions 29-34. In another example, an affinity matured antibody of the invention which specifically binds to K63-linked polyubiquitin with high affinity but binds to K48-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H2 amino acid positions 50, 52, 54, 56, and 58. In another example, an affinity matured antibody of the invention which specifically binds to K63-linked polyubiquitin with high affinity but binds to K48-linked polyubiquitin with substantially reduced affinity comprises substitution at HVR-H3 amino acid positions 95-100, 100a, 100b, and 100c.

In one embodiment, an antibody of the invention comprises a heavy chain variable domain comprising the sequence of SEQ ID NOs: 265, 281, and 297. In one embodiment, an antibody of the invention comprises a light chain variable domain comprising the sequence of SEQ ID NOs: 79, 80, and 313. In one embodiment, an antibody of the invention comprises a heavy chain variable domain comprising the sequence of SEQ ID NOs: 265, 281, and 297 and also comprises a light chain variable domain comprising the sequence of SEQ ID NOs: 79, 80, and 313. In other embodiments, an antibody of the invention corresponding to a particular clone number comprises a heavy chain variable domain comprising an HVR-H1, HVR-H2, and HVR-H3 sequence as set forth in Figures 2, 3, 8, 9, 10, 11, 14-17, and 22 for that clone number. In other embodiments, an antibody of the invention corresponding to a particular clone number comprises a light chain variable domain comprising an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence as set forth in Figures 10, 11, 16, and 17 for that clone number. In other embodiments, an antibody of the invention corresponding to a particular clone number comprises a heavy chain variable domain comprising an HVR-H1, HVR-H2, and HVR-H3 sequence as set forth in Figures 2, 3, 8, 9, 10, 11, 14-17, and 22 for that clone number and also comprises a light chain variable domain comprising an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence as set forth in Figures 10, 11, 16, and 17 for that clone number.

In one aspect, the invention provides an antibody that competes with any of the above-mentioned antibodies for binding to polyubiquitin. In one aspect, the invention provides an antibody that binds to the same antigenic determinant on polyubiquitin as any of the above-mentioned antibodies.

As shown herein, the antibodies of the invention specifically bind to an isolated polyubiquitin having a specific lysine linkage. As shown herein, the antibodies of the invention also specifically bind to polyubiquitin having a specific lysine linkage when that polyubiquitin is attached to a heterologous protein (see, e.g., Examples 3 and 4).

Compositions comprising at least one anti-polyubiquitin antibody or at least one polynucleotide comprising sequences encoding an anti-polyubiquitin antibody are provided. In certain embodiments, a composition may be a pharmaceutical composition. As used herein,

compositions comprise one or more antibodies that bind to one or more polyubiquitin and/or one or more polynucleotides comprising sequences encoding one or more antibodies that bind to one or more polyubiquitin. These compositions may further comprise suitable carriers, such as pharmaceutically acceptable excipients including buffers, which are well known in the art.

5 Isolated antibodies and polynucleotides are also provided. In certain embodiments, the isolated antibodies and polynucleotides are substantially pure.

In one embodiment, anti-polyubiquitin antibodies are monoclonal. In another embodiment, fragments of the anti-polyubiquitin antibodies (e.g., Fab, Fab'-SH and F(ab')₂ fragments) are provided. These antibody fragments can be created by traditional means, such as enzymatic digestion, 10 or may be generated by recombinant techniques. Such antibody fragments may be chimeric, humanized, or human. These fragments are useful for the diagnostic and therapeutic purposes set forth below.

Generation of anti-polyubiquitin antibodies using a phage display library

15 A variety of methods are known in the art for generating phage display libraries from which an antibody of interest can be obtained. One method of generating antibodies of interest is through the use of a phage antibody library as described in Lee et al., J. Mol. Biol. (2004), 340(5):1073-93.

The anti-polyubiquitin antibodies of the invention can be made by using combinatorial libraries to screen for synthetic antibody clones with the desired activity or activities. In principle, 20 synthetic antibody clones are selected by screening phage libraries containing phage that display various fragments of antibody variable region (Fv) fused to phage coat protein. Such phage libraries are panned by affinity chromatography against the desired antigen. Clones expressing Fv fragments capable of binding to the desired antigen are adsorbed to the antigen and thus separated from the non-binding clones in the library. The binding clones are then eluted from the antigen, and can be further 25 enriched by additional cycles of antigen adsorption/elution. Any of the anti-polyubiquitin antibodies of the invention can be obtained by designing a suitable antigen screening procedure to select for the phage clone of interest followed by construction of a full length anti-polyubiquitin antibody clone using the Fv sequences from the phage clone of interest and suitable constant region (Fc) sequences described in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, Fifth Edition, NIH 30 Publication 91-3242, Bethesda MD (1991), vols. 1-3.

The antigen-binding domain of an antibody is formed from two variable (V) regions of about 110 amino acids, one each from the light (VL) and heavy (VH) chains, that both present three hypervariable loops or complementarity-determining regions (CDRs). Variable domains can be displayed functionally on phage, either as single-chain Fv (scFv) fragments, in which VH and VL are 35 covalently linked through a short, flexible peptide, or as Fab fragments, in which they are each fused to a constant domain and interact non-covalently, as described in Winter *et al.*, *Ann. Rev. Immunol.*,

12: 433-455 (1994). As used herein, scFv encoding phage clones and Fab encoding phage clones are collectively referred to as "Fv phage clones" or "Fv clones".

Repertoires of VH and VL genes can be separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be searched for antigen-binding clones as described in Winter *et al.*, *Ann. Rev. Immunol.*, **12**: 433-455 (1994). Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned to provide a single source of human antibodies to a wide range of non-self and also self antigens without any immunization as described by Griffiths *et al.*, *EMBO J*, **12**: 725-734 (1993). Finally, naive libraries can also be made synthetically by cloning the unrearranged V-gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to accomplish rearrangement in vitro as described by Hoogenboom and Winter, *J. Mol. Biol.*, **227**: 381-388 (1992).

Filamentous phage is used to display antibody fragments by fusion to the minor coat protein pIII. The antibody fragments can be displayed as single chain Fv fragments, in which VH and VL domains are connected on the same polypeptide chain by a flexible polypeptide spacer, e.g. as described by Marks *et al.*, *J. Mol. Biol.*, **222**: 581-597 (1991), or as Fab fragments, in which one chain is fused to pIII and the other is secreted into the bacterial host cell periplasm where assembly of a Fab-coat protein structure which becomes displayed on the phage surface by displacing some of the wild type coat proteins, e.g. as described in Hoogenboom *et al.*, *Nucl. Acids Res.*, **19**: 4133-4137 (1991).

In general, nucleic acids encoding antibody gene fragments are obtained from immune cells harvested from humans or animals. If a library biased in favor of anti-polyubiquitin clones is desired, the subject is immunized with polyubiquitin to generate an antibody response, and spleen cells and/or circulating B cells or other peripheral blood lymphocytes (PBLs) are recovered for library construction. In one embodiment, a human antibody gene fragment library biased in favor of anti-human polyubiquitin clones is obtained by generating an anti-human polyubiquitin antibody response in transgenic mice carrying a functional human immunoglobulin gene array (and lacking a functional endogenous antibody production system) such that polyubiquitin immunization gives rise to B cells producing human antibodies against polyubiquitin. The generation of human antibody-producing transgenic mice is described in Section (III)(b) below.

Additional enrichment for anti- polyubiquitin reactive cell populations can be obtained by using a suitable screening procedure to isolate B cells expressing polyubiquitin-specific membrane bound antibody, e.g., by cell separation with polyubiquitin affinity chromatography or adsorption of cells to fluorochrome-labeled polyubiquitin followed by flow-activated cell sorting (FACS).

Alternatively, the use of spleen cells and/or B cells or other PBLs from an unimmunized donor provides a better representation of the possible antibody repertoire, and also permits the construction of an antibody library using any animal (human or non-human) species in which

polyubiquitin is not antigenic. For libraries incorporating in vitro antibody gene construction, stem cells are harvested from the subject to provide nucleic acids encoding unrearranged antibody gene segments. The immune cells of interest can be obtained from a variety of animal species, such as human, mouse, rat, lagomorpha, luprine, canine, feline, porcine, bovine, equine, and avian species,
5 etc.

Nucleic acid encoding antibody variable gene segments (including VH and VL segments) are recovered from the cells of interest and amplified. In the case of rearranged VH and VL gene libraries, the desired DNA can be obtained by isolating genomic DNA or mRNA from lymphocytes followed by polymerase chain reaction (PCR) with primers matching the 5' and 3' ends of rearranged
10 VH and VL genes as described in Orlandi *et al.*, *Proc. Natl. Acad. Sci. (USA)*, **86**: 3833-3837 (1989), thereby making diverse V gene repertoires for expression. The V genes can be amplified from cDNA and genomic DNA, with back primers at the 5' end of the exon encoding the mature V-domain and forward primers based within the J-segment as described in Orlandi *et al.* (1989) and in Ward *et al.*, *Nature*, **341**: 544-546 (1989). However, for amplifying from cDNA, back primers can also be based
15 in the leader exon as described in Jones *et al.*, *Biotechnol.*, **9**: 88-89 (1991), and forward primers within the constant region as described in Sastry *et al.*, *Proc. Natl. Acad. Sci. (USA)*, **86**: 5728-5732 (1989). To maximize complementarity, degeneracy can be incorporated in the primers as described in Orlandi *et al.* (1989) or Sastry *et al.* (1989). In certain embodiments, the library diversity is maximized by using PCR primers targeted to each V-gene family in order to amplify all available VH
20 and VL arrangements present in the immune cell nucleic acid sample, e.g. as described in the method of Marks *et al.*, *J. Mol. Biol.*, **222**: 581-597 (1991) or as described in the method of Orum *et al.*, *Nucleic Acids Res.*, **21**: 4491-4498 (1993). For cloning of the amplified DNA into expression vectors, rare restriction sites can be introduced within the PCR primer as a tag at one end as described in Orlandi *et al.* (1989), or by further PCR amplification with a tagged primer as described in Clackson
25 *et al.*, *Nature*, **352**: 624-628 (1991).

Repertoires of synthetically rearranged V genes can be derived in vitro from V gene segments. Most of the human VH-gene segments have been cloned and sequenced (reported in Tomlinson *et al.*, *J. Mol. Biol.*, **227**: 776-798 (1992)), and mapped (reported in Matsuda *et al.*, *Nature Genet.*, **3**: 88-94 (1993)); these cloned segments (including all the major conformations of the H1 and
30 H2 loop) can be used to generate diverse VH gene repertoires with PCR primers encoding H3 loops of diverse sequence and length as described in Hoogenboom and Winter, *J. Mol. Biol.*, **227**: 381-388 (1992). VH repertoires can also be made with all the sequence diversity focused in a long H3 loop of a single length as described in Barbas *et al.*, *Proc. Natl. Acad. Sci. USA*, **89**: 4457-4461 (1992). Human V κ and V λ segments have been cloned and sequenced (reported in Williams and Winter, *Eur.*
35 *J. Immunol.*, **23**: 1456-1461 (1993)) and can be used to make synthetic light chain repertoires. Synthetic V gene repertoires, based on a range of VH and VL folds, and L3 and H3 lengths, will encode antibodies of considerable structural diversity. Following amplification of V-gene encoding

DNAs, germline V-gene segments can be rearranged in vitro according to the methods of Hoogenboom and Winter, *J. Mol. Biol.*, **227**: 381-388 (1992).

Repertoires of antibody fragments can be constructed by combining VH and VL gene repertoires together in several ways. Each repertoire can be created in different vectors, and the vectors recombined in vitro, e.g., as described in Hogrefe *et al.*, *Gene*, **128**: 119-126 (1993), or in vivo by combinatorial infection, e.g., the loxP system described in Waterhouse *et al.*, *Nucl. Acids Res.*, **21**: 2265-2266 (1993). The *in vivo* recombination approach exploits the two-chain nature of Fab fragments to overcome the limit on library size imposed by *E. coli* transformation efficiency. Naive VH and VL repertoires are cloned separately, one into a phagemid and the other into a phage vector. The two libraries are then combined by phage infection of phagemid-containing bacteria so that each cell contains a different combination and the library size is limited only by the number of cells present (about 10^{12} clones). Both vectors contain in vivo recombination signals so that the VH and VL genes are recombined onto a single replicon and are co-packaged into phage virions. These huge libraries provide large numbers of diverse antibodies of good affinity (K_d^{-1} of about 10^{-8} M).

Alternatively, the repertoires may be cloned sequentially into the same vector, e.g. as described in Barbas *et al.*, *Proc. Natl. Acad. Sci. USA*, **88**: 7978-7982 (1991), or assembled together by PCR and then cloned, e.g. as described in Clackson *et al.*, *Nature*, **352**: 624-628 (1991). PCR assembly can also be used to join VH and VL DNAs with DNA encoding a flexible peptide spacer to form single chain Fv (scFv) repertoires. In yet another technique, "in cell PCR assembly" is used to combine VH and VL genes within lymphocytes by PCR and then clone repertoires of linked genes as described in Embleton *et al.*, *Nucl. Acids Res.*, **20**: 3831-3837 (1992).

Screening of the libraries can be accomplished by any art-known technique. For example, polyubiquitin can be used to coat the wells of adsorption plates, expressed on host cells affixed to adsorption plates or used in cell sorting, or conjugated to biotin for capture with streptavidin-coated beads, or used in any other art-known method for panning phage display libraries.

The phage library samples are contacted with immobilized polyubiquitin under conditions suitable for binding of at least a portion of the phage particles with the adsorbent. Normally, the conditions, including pH, ionic strength, temperature and the like are selected to mimic physiological conditions. The phages bound to the solid phase are washed and then eluted by acid, e.g. as described in Barbas *et al.*, *Proc. Natl. Acad. Sci USA*, **88**: 7978-7982 (1991), or by alkali, e.g. as described in Marks *et al.*, *J. Mol. Biol.*, **222**: 581-597 (1991), or by polyubiquitin antigen competition, e.g. in a procedure similar to the antigen competition method of Clackson *et al.*, *Nature*, **352**: 624-628 (1991). Phages can be enriched 20-1,000-fold in a single round of selection. Moreover, the enriched phages can be grown in bacterial culture and subjected to further rounds of selection.

The efficiency of selection depends on many factors, including the kinetics of dissociation during washing, and whether multiple antibody fragments on a single phage can simultaneously engage with antigen. Antibodies with fast dissociation kinetics (and weak binding affinities) can be

retained by use of short washes, multivalent phage display and high coating density of antigen in solid phase. The high density not only stabilizes the phage through multivalent interactions, but favors rebinding of phage that has dissociated. The selection of antibodies with slow dissociation kinetics (and good binding affinities) can be promoted by use of long washes and monovalent phage display as described in Bass *et al.*, *Proteins*, **8**: 309-314 (1990) and in WO 92/09690, and a low coating density of antigen as described in Marks *et al.*, *Biotechnol.*, **10**: 779-783 (1992).

It is possible to select between phage antibodies of different affinities, even with affinities that differ slightly, for polyubiquitin. However, random mutation of a selected antibody (e.g. as performed in some of the affinity maturation techniques described above) is likely to give rise to many mutants, most binding to antigen, and a few with higher affinity. With limiting polyubiquitin, rare high affinity phage could be competed out. To retain all the higher affinity mutants, phages can be incubated with excess biotinylated polyubiquitin, but with the biotinylated polyubiquitin at a concentration of lower molarity than the target molar affinity constant for polyubiquitin. The high affinity-binding phages can then be captured by streptavidin-coated paramagnetic beads. Such "equilibrium capture" allows the antibodies to be selected according to their affinities of binding, with sensitivity that permits isolation of mutant clones with as little as two-fold higher affinity from a great excess of phages with lower affinity. Conditions used in washing phages bound to a solid phase can also be manipulated to discriminate on the basis of dissociation kinetics.

Anti-polyubiquitin clones may be activity selected. In one embodiment, the invention provides anti-polyubiquitin antibodies that block the binding between a polyubiquitin ligand and polyubiquitin, but do not block the binding between a polyubiquitin ligand and a second protein. Fv clones corresponding to such anti-polyubiquitin antibodies can be selected by (1) isolating anti-polyubiquitin clones from a phage library as described in Section B(I)(2) above, and optionally amplifying the isolated population of phage clones by growing up the population in a suitable bacterial host; (2) selecting polyubiquitin and a second protein against which blocking and non-blocking activity, respectively, is desired; (3) adsorbing the anti-polyubiquitin phage clones to immobilized polyubiquitin; (4) using an excess of the second protein to elute any undesired clones that recognize polyubiquitin-binding determinants which overlap or are shared with the binding determinants of the second protein; and (5) eluting the clones which remain adsorbed following step (4). Optionally, clones with the desired blocking/non-blocking properties can be further enriched by repeating the selection procedures described herein one or more times.

DNA encoding the Fv clones of the invention is readily isolated and sequenced using conventional procedures (e.g. by using oligonucleotide primers designed to specifically amplify the heavy and light chain coding regions of interest from hybridoma or phage DNA template). Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of the desired monoclonal

antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of antibody-encoding DNA include Skerra *et al.*, *Curr. Opinion in Immunol.*, **5**: 256 (1993) and Pluckthun, *Immunol. Revs.*, **130**: 151 (1992).

DNA encoding the Fv clones of the invention can be combined with known DNA sequences encoding heavy chain and/or light chain constant regions (e.g. the appropriate DNA sequences can be obtained from Kabat *et al.*, *supra*) to form clones encoding full or partial length heavy and/or light chains. It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species. A Fv clone derived from the variable domain DNA of one animal (such as human) species and then fused to constant region DNA of another animal species to form coding sequence(s) for "hybrid", full length heavy chain and/or light chain is included in the definition of "chimeric" and "hybrid" antibody as used herein. In one embodiment, a Fv clone derived from human variable DNA is fused to human constant region DNA to form coding sequence(s) for all human, full or partial length heavy and/or light chains.

The antibodies produced by naive libraries (either natural or synthetic) can be of moderate affinity (K_d^{-1} of about 10^6 to 10^7 M^{-1}), but affinity maturation can also be mimicked *in vitro* by constructing and reselecting from secondary libraries as described in Winter *et al.* (1994), *supra*. For example, mutation can be introduced at random *in vitro* by using error-prone polymerase (reported in Leung *et al.*, *Technique*, **1**: 11-15 (1989)) in the method of Hawkins *et al.*, *J. Mol. Biol.*, **226**: 889-896 (1992) or in the method of Gram *et al.*, *Proc. Natl. Acad. Sci USA*, **89**: 3576-3580 (1992).

Additionally, affinity maturation can be performed by randomly mutating one or more CDRs, e.g. using PCR with primers carrying random sequence spanning the CDR of interest, in selected individual Fv clones and screening for higher affinity clones. WO 9607754 (published 14 March 1996) described a method for inducing mutagenesis in a complementarity determining region of an immunoglobulin light chain to create a library of light chain genes. Another effective approach is to recombine the VH or VL domains selected by phage display with repertoires of naturally occurring V domain variants obtained from unimmunized donors and screen for higher affinity in several rounds of chain reshuffling as described in Marks *et al.*, *Biotechnol.*, **10**: 779-783 (1992). This technique allows the production of antibodies and antibody fragments with affinities in the 10^{-9} M range.

Other methods of generating anti-polyubiquitin antibodies

Other methods of generating and assessing the affinity of antibodies are well known in the art and are described, e.g., in Kohler *et al.*, *Nature* 256: 495 (1975); U.S. Patent No. 4,816,567; Goding, *Monoclonal Antibodies: Principles and Practice*, pp. 59-103 (Academic Press, 1986; Kozbor, *J. Immunol.*, **133**:3001 (1984); Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987; Munson *et al.*, *Anal. Biochem.*, **107**:220 (1980); Engels *et al.*, *Angew. Chem. Int. Ed. Engl.*, **28**: 716-734 (1989); Abrahmsen *et al.*,

EMBO J., **4**: 3901 (1985); *Methods in Enzymology*, vol. 44 (1976); Morrison *et al.*, *Proc. Natl. Acad. Sci. USA*, **81**: 6851-6855 (1984).

General Methods

5 In general, the invention provides anti-polyubiquitin antibodies that are useful for treatment of polyubiquitin-mediated disorders in which a partial or total blockade of one or more polyubiquitin activities is desired. In one embodiment, the anti-polyubiquitin antibodies of the invention are used to treat cancer. In another embodiment, the anti-polyubiquitin antibodies provided herein are used to treat muscular disorders, such as those indicated above. In another embodiment, the anti-
10 polyubiquitin antibodies provided herein are used to treat neurological disorders, such as those indicated above. In another embodiment, the anti-polyubiquitin antibodies provided herein are used to treat genetic disease. In another embodiment, the anti-polyubiquitin antibodies provided herein are used to treat immune/inflammatory disorders.

The unique properties of the anti-polyubiquitin antibodies of the invention make them
15 particularly useful for distinguishing between different lysine-linked forms of polyubiquitin in a cellular system without resorting to cumbersome and expensive genetic manipulation or biophysical methods such as mass spectrometry. The anti-polyubiquitin antibodies of the invention can be used to characterize the function(s) and activities of specific lysine-linked polyubiquitins both *in vitro* and *in vivo*. The anti-polyubiquitin antibodies of the invention can also be used to determine the role of
20 specific lysine-linked polyubiquitins in the development and pathogenesis of disease. The anti-polyubiquitin antibodies of the invention can further be used to treat diseases in which one or more specific lysine-linked polyubiquitins are aberrantly regulated or aberrantly functioning without interfering with the normal activity of polyubiquitins for which the anti-polyubiquitin antibodies are not specific.

25 In another aspect, the anti-polyubiquitin antibodies of the invention find utility as reagents for detection and isolation of polyubiquitin of specific lysine linkages, such as detection of polyubiquitin in various cell types and tissues, including the determination of polyubiquitin density and distribution in cell populations and within a given cell, and cell sorting based on the presence or amount of polyubiquitin.

30 In yet another aspect, the present anti-polyubiquitin antibodies are useful for the development of polyubiquitin antagonists with blocking activity patterns similar to those of the subject antibodies of the invention. For example, anti-K48-linked polyubiquitin antibodies of the invention can be used to determine and identify other antibodies that have the same K48-linked polyubiquitin binding characteristics and/or capabilities of blocking K48-linked polyubiquitin-mediated pathways.

35 Similarly, anti-K63-linked polyubiquitin antibodies of the invention can be used to determine and identify other antibodies that have the same K63-linked polyubiquitin binding characteristics and/or capabilities of blocking K63-linked polyubiquitin-mediated pathways.

As a further example, anti-polyubiquitin antibodies of the invention can be used to identify other anti-polyubiquitin antibodies that bind substantially the same antigenic determinant(s) of polyubiquitin as the antibodies exemplified herein, including linear and conformational epitopes.

The anti-polyubiquitin antibodies of the invention can be used in assays based on the physiological pathways in which polyubiquitin is involved to screen for small molecule antagonists of polyubiquitin having one or more specific lysine linkages which will exhibit similar pharmacological effects in blocking the binding of one or more binding partners to polyubiquitin having those one or more lysine linkages. For example, K48-linked polyubiquitin is known to be involved in targeted proteasomal degradation of certain proteins (*see, e.g.,* Chau et al., Science 243: 1576-1583 (1989); Finley et al., Mol. Cell. Biol. 14: 5501-5509 (1994); Flick et al., Nat. Cell. Biol. 6:634-641 (2004)); thus anti-K48-linked polyubiquitin antibodies may be used in screens to identify small molecule antagonists of K48-linked polyubiquitin-mediated targeted proteasomal degradation by comparing the activity of one or more potential small molecule antagonists to the activity of the anti-K48-linked polyubiquitin antibodies in that pathway. Similarly, in another example, K63-linked polyubiquitin is known to be involved in DNA repair (*see, e.g.,* Pickart and Fushman, Curr. Opin. Chem. Biol. 8: 610-616 (2004)), and thus the activity of anti-K63-linked polyubiquitin antibodies to antagonize a DNA repair pathway may be compared to the activity of one or more potential small molecule antagonists of K63-linked polyubiquitin in that same DNA repair pathway. Similarly, in another example, K63-linked polyubiquitin is known to be involved in formation of Lewy bodies in Parkinson's disease (*see, e.g.,* Lim et al., J. Neurosci. 25(8): 2002-9 (2005)), and thus the activity of anti-K63-linked polyubiquitin antibodies to antagonize the formation of Lewy bodies may be compared to the activity of one or more potential small molecule antagonists of K63-linked polyubiquitin in antagonizing the formation of Lewy bodies.

Generation of candidate antibodies can be achieved using routine skills in the art, including those described herein, such as the hybridoma technique and screening of phage displayed libraries of binder molecules. These methods are well-established in the art.

Briefly, the anti-polyubiquitin antibodies of the invention can be made by using combinatorial libraries to screen for synthetic antibody clones with the desired activity or activities. In principle, synthetic antibody clones are selected by screening phage libraries containing phage that display various fragments of antibody variable region (Fv) fused to phage coat protein. Such phage libraries are panned by affinity chromatography against the desired antigen. Clones expressing Fv fragments capable of binding to the desired antigen are adsorbed to the antigen and thus separated from the non-binding clones in the library. The binding clones are then eluted from the antigen, and can be further enriched by additional cycles of antigen adsorption/elution. Any of the anti-polyubiquitin antibodies of the invention can be obtained by designing a suitable antigen screening procedure to select for the phage clone of interest followed by construction of a full length anti-polyubiquitin antibody clone using the Fv sequences from the phage clone of interest and suitable constant region (Fc) sequences

described in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, Fifth Edition, NIH Publication 91-3242, Bethesda MD (1991), vols. 1-3. See also PCT Pub. WO03/102157, and references cited therein.

In one embodiment, anti-polyubiquitin antibodies of the invention are monoclonal. Also encompassed within the scope of the invention are antibody fragments such as Fab, Fab', Fab'-SH and F(ab')₂ fragments, and variations thereof, of the anti-polyubiquitin antibodies provided herein. These antibody fragments can be created by traditional means, such as enzymatic digestion, or may be generated by recombinant techniques. Such antibody fragments may be chimeric, human or humanized. These fragments are useful for the experimental, diagnostic, and therapeutic purposes set forth herein.

Monoclonal antibodies can be obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies.

The anti-polyubiquitin monoclonal antibodies of the invention can be made using a variety of methods known in the art, including the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or alternatively they may be made by recombinant DNA methods (e.g., U.S. Patent No. 4,816,567).

Vectors, Host Cells and Recombinant Methods

For recombinant production of an antibody of the invention, the nucleic acid encoding it is isolated and inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. DNA encoding the antibody is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody). Many vectors are available. The choice of vector depends in part on the host cell to be used. Host cells include, but are not limited to, cells of either prokaryotic or eukaryotic (generally mammalian) origin. It will be appreciated that constant regions of any isotype can be used for this purpose, including IgG, IgM, IgA, IgD, and IgE constant regions, and that such constant regions can be obtained from any human or animal species.

Generating Antibodies Using Prokaryotic Host Cells

Vector Construction

Polynucleotide sequences encoding polypeptide components of the antibody of the invention can be obtained using standard recombinant techniques. Desired polynucleotide sequences may be isolated and sequenced from antibody producing cells such as hybridoma cells. Alternatively, polynucleotides can be synthesized using nucleotide synthesizer or PCR techniques. Once obtained,

sequences encoding the polypeptides are inserted into a recombinant vector capable of replicating and expressing heterologous polynucleotides in prokaryotic hosts. Many vectors that are available and known in the art can be used for the purpose of the present invention. Selection of an appropriate vector will depend mainly on the size of the nucleic acids to be inserted into the vector and the particular host cell to be transformed with the vector. Each vector contains various components, depending on its function (amplification or expression of heterologous polynucleotide, or both) and its compatibility with the particular host cell in which it resides. The vector components generally include, but are not limited to: an origin of replication, a selection marker gene, a promoter, a ribosome binding site (RBS), a signal sequence, the heterologous nucleic acid insert and a transcription termination sequence.

In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using pBR322, a plasmid derived from an *E. coli* species. pBR322 contains genes encoding ampicillin (Amp) and tetracycline (Tet) resistance and thus provides easy means for identifying transformed cells. pBR322, its derivatives, or other microbial plasmids or bacteriophage may also contain, or be modified to contain, promoters which can be used by the microbial organism for expression of endogenous proteins. Examples of pBR322 derivatives used for expression of particular antibodies are described in detail in Carter et al., U.S. Patent No. 5,648,237.

In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, bacteriophage such as λ GEM.TM.-11 may be utilized in making a recombinant vector which can be used to transform susceptible host cells such as *E. coli* LE392.

The expression vector of the invention may comprise two or more promoter-cistron pairs, encoding each of the polypeptide components. A promoter is an untranslated regulatory sequence located upstream (5') to a cistron that modulates its expression. Prokaryotic promoters typically fall into two classes, inducible and constitutive. Inducible promoter is a promoter that initiates increased levels of transcription of the cistron under its control in response to changes in the culture condition, e.g. the presence or absence of a nutrient or a change in temperature.

A large number of promoters recognized by a variety of potential host cells are well known. The selected promoter can be operably linked to cistron DNA encoding the light or heavy chain by removing the promoter from the source DNA via restriction enzyme digestion and inserting the isolated promoter sequence into the vector of the invention. Both the native promoter sequence and many heterologous promoters may be used to direct amplification and/or expression of the target genes. In some embodiments, heterologous promoters are utilized, as they generally permit greater

transcription and higher yields of expressed target gene as compared to the native target polypeptide promoter.

Promoters suitable for use with prokaryotic hosts include the PhoA promoter, the β -galactamase and lactose promoter systems, a tryptophan (*trp*) promoter system and hybrid promoters such as the *tac* or the *trc* promoter. However, other promoters that are functional in bacteria (such as other known bacterial or phage promoters) are suitable as well. Their nucleotide sequences have been published, thereby enabling a skilled worker operably to ligate them to cistrons encoding the target light and heavy chains (Siebenlist et al. (1980) Cell 20: 269) using linkers or adaptors to supply any required restriction sites.

In one aspect of the invention, each cistron within the recombinant vector comprises a secretion signal sequence component that directs translocation of the expressed polypeptides across a membrane. In general, the signal sequence may be a component of the vector, or it may be a part of the target polypeptide DNA that is inserted into the vector. The signal sequence selected for the purpose of this invention should be one that is recognized and processed (i.e. cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the signal sequences native to the heterologous polypeptides, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group consisting of the alkaline phosphatase, penicillinase, Ipp, or heat-stable enterotoxin II (STII) leaders, LamB, PhoE, PelB, OmpA and MBP. In one embodiment of the invention, the signal sequences used in both cistrons of the expression system are STII signal sequences or variants thereof.

In another aspect, the production of the immunoglobulins according to the invention can occur in the cytoplasm of the host cell, and therefore does not require the presence of secretion signal sequences within each cistron. In that regard, immunoglobulin light and heavy chains are expressed, folded and assembled to form functional immunoglobulins within the cytoplasm. Certain host strains (e.g., the *E. coli trxB* strains) provide cytoplasm conditions that are favorable for disulfide bond formation, thereby permitting proper folding and assembly of expressed protein subunits. Proba and Pluckthun *Gene*, 159:203 (1995).

Antibodies of the invention can also be produced by using an expression system in which the quantitative ratio of expressed polypeptide components can be modulated in order to maximize the yield of secreted and properly assembled antibodies of the invention. Such modulation is accomplished at least in part by simultaneously modulating translational strengths for the polypeptide components.

One technique for modulating translational strength is disclosed in Simmons et al., U.S. Pat. No. 5,840,523. It utilizes variants of the translational initiation region (TIR) within a cistron. For a given TIR, a series of amino acid or nucleic acid sequence variants can be created with a range of

translational strengths, thereby providing a convenient means by which to adjust this factor for the desired expression level of the specific chain. TIR variants can be generated by conventional mutagenesis techniques that result in codon changes which can alter the amino acid sequence. In certain embodiments, changes in the nucleotide sequence are silent. Alterations in the TIR can include, for example, alterations in the number or spacing of Shine-Dalgarno sequences, along with alterations in the signal sequence. One method for generating mutant signal sequences is the generation of a "codon bank" at the beginning of a coding sequence that does not change the amino acid sequence of the signal sequence (i.e., the changes are silent). This can be accomplished by changing the third nucleotide position of each codon; additionally, some amino acids, such as leucine, serine, and arginine, have multiple first and second positions that can add complexity in making the bank. This method of mutagenesis is described in detail in Yansura et al. (1992) *METHODS: A Companion to Methods in Enzymol.* 4:151-158.

In one embodiment, a set of vectors is generated with a range of TIR strengths for each cistron therein. This limited set provides a comparison of expression levels of each chain as well as the yield of the desired antibody products under various TIR strength combinations. TIR strengths can be determined by quantifying the expression level of a reporter gene as described in detail in Simmons et al. U.S. Pat. No. 5, 840,523. Based on the translational strength comparison, the desired individual TIRs are selected to be combined in the expression vector constructs of the invention.

Prokaryotic host cells suitable for expressing antibodies of the invention include Archaeobacteria and Eubacteria, such as Gram-negative or Gram-positive organisms. Examples of useful bacteria include Escherichia (e.g., *E. coli*), Bacilli (e.g., *B. subtilis*), Enterobacteria, Pseudomonas species (e.g., *P. aeruginosa*), Salmonella typhimurium, Serratia marcescans, Klebsiella, Proteus, Shigella, Rhizobia, Vitreoscilla, or Paracoccus. In one embodiment, gram-negative cells are used. In one embodiment, *E. coli* cells are used as hosts for the invention. Examples of *E. coli* strains include strain W3110 (Bachmann, Cellular and Molecular Biology, vol. 2 (Washington, D.C.: American Society for Microbiology, 1987), pp. 1190-1219; ATCC Deposit No. 27,325) and derivatives thereof, including strain 33D3 having genotype W3110 $\Delta fhuA$ ($\Delta tonA$) *ptr3 lac Iq lacL8* $\Delta ompT \Delta (nmpc-fepE)$ *degP41 kan^R* (U.S. Pat. No. 5,639,635). Other strains and derivatives thereof, such as *E. coli* 294 (ATCC 31,446), *E. coli* B, *E. coli* λ 1776 (ATCC 31,537) and *E. coli* RV308 (ATCC 31,608) are also suitable. These examples are illustrative rather than limiting. Methods for constructing derivatives of any of the above-mentioned bacteria having defined genotypes are known in the art and described in, for example, Bass *et al.*, Proteins, 8:309-314 (1990). It is generally necessary to select the appropriate bacteria taking into consideration replicability of the replicon in the cells of a bacterium. For example, *E. coli*, Serratia, or Salmonella species can be suitably used as the host when well known plasmids such as pBR322, pBR325, pACYC177, or pKN410 are used to supply the replicon. Typically the host cell should secrete minimal amounts of

proteolytic enzymes, and additional protease inhibitors may desirably be incorporated in the cell culture.

Antibody Production

Host cells are transformed with the above-described expression vectors and cultured in
5 conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

Transformation means introducing DNA into the prokaryotic host so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integrant. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The
10 calcium treatment employing calcium chloride is generally used for bacterial cells that contain substantial cell-wall barriers. Another method for transformation employs polyethylene glycol/DMSO. Yet another technique used is electroporation.

Prokaryotic cells used to produce the polypeptides of the invention are grown in media known in the art and suitable for culture of the selected host cells. Examples of suitable media include luria
15 broth (LB) plus necessary nutrient supplements. In some embodiments, the media also contains a selection agent, chosen based on the construction of the expression vector, to selectively permit growth of prokaryotic cells containing the expression vector. For example, ampicillin is added to media for growth of cells expressing ampicillin resistant gene.

Any necessary supplements besides carbon, nitrogen, and inorganic phosphate sources may
20 also be included at appropriate concentrations introduced alone or as a mixture with another supplement or medium such as a complex nitrogen source. Optionally the culture medium may contain one or more reducing agents selected from the group consisting of glutathione, cysteine, cystamine, thioglycollate, dithioerythritol and dithiothreitol.

The prokaryotic host cells are cultured at suitable temperatures. For *E. coli* growth, for
25 example, growth occurs at a temperature range including, but not limited to, about 20°C to about 39°C, about 25°C to about 37°C, and at about 30°C. The pH of the medium may be any pH ranging from about 5 to about 9, depending mainly on the host organism. For *E. coli*, the pH can be from about 6.8 to about 7.4, or about 7.0.

If an inducible promoter is used in the expression vector of the invention, protein expression
30 is induced under conditions suitable for the activation of the promoter. In one aspect of the invention, PhoA promoters are used for controlling transcription of the polypeptides. Accordingly, the transformed host cells are cultured in a phosphate-limiting medium for induction. In one embodiment, the phosphate-limiting medium is the C.R.A.P medium (see, e.g., Simmons et al., *J.*

Immunol. Methods (2002), 263:133-147). A variety of other inducers may be used, according to the vector construct employed, as is known in the art.

In one embodiment, the expressed polypeptides of the present invention are secreted into and recovered from the periplasm of the host cells. Protein recovery typically involves disrupting the microorganism, generally by such means as osmotic shock, sonication or lysis. Once cells are disrupted, cell debris or whole cells may be removed by centrifugation or filtration. The proteins may be further purified, for example, by affinity resin chromatography. Alternatively, proteins can be transported into the culture media and isolated therein. Cells may be removed from the culture and the culture supernatant being filtered and concentrated for further purification of the proteins produced. The expressed polypeptides can be further isolated and identified using commonly known methods such as polyacrylamide gel electrophoresis (PAGE) and Western blot assay.

In one aspect of the invention, antibody production is conducted in large quantity by a fermentation process. Various large-scale fed-batch fermentation procedures are available for production of recombinant proteins. Large-scale fermentations have at least 1000 liters of capacity, for example about 1,000 to 100,000 liters of capacity. These fermentors use agitator impellers to distribute oxygen and nutrients, especially glucose (a common carbon/energy source). Small scale fermentation refers generally to fermentation in a fermentor that is no more than approximately 100 liters in volumetric capacity, and can range from about 1 liter to about 100 liters.

In a fermentation process, induction of protein expression is typically initiated after the cells have been grown under suitable conditions to a desired density, e.g., an OD₅₅₀ of about 180-220, at which stage the cells are in the early stationary phase. A variety of inducers may be used, according to the vector construct employed, as is known in the art and described above. Cells may be grown for shorter periods prior to induction. Cells are usually induced for about 12-50 hours, although longer or shorter induction time may be used.

To improve the production yield and quality of the polypeptides of the invention, various fermentation conditions can be modified. For example, to improve the proper assembly and folding of the secreted antibody polypeptides, additional vectors overexpressing chaperone proteins, such as Dsb proteins (DsbA, DsbB, DsbC, DsbD and or DsbG) or FkpA (a peptidylprolyl cis,trans-isomerase with chaperone activity) can be used to co-transform the host prokaryotic cells. The chaperone proteins have been demonstrated to facilitate the proper folding and solubility of heterologous proteins produced in bacterial host cells. Chen et al. (1999) *J Bio Chem* 274:19601-19605; Georgiou et al., U.S. Patent No. 6,083,715; Georgiou et al., U.S. Patent No. 6,027,888; Bothmann and Pluckthun (2000) *J. Biol. Chem.* 275:17100-17105; Ramm and Pluckthun (2000) *J. Biol. Chem.* 275:17106-17113; Arie et al. (2001) *Mol. Microbiol.* 39:199-210.

To minimize proteolysis of expressed heterologous proteins (especially those that are proteolytically sensitive), certain host strains deficient for proteolytic enzymes can be used for the present invention. For example, host cell strains may be modified to effect genetic mutation(s) in the genes encoding known bacterial proteases such as Protease III, OmpT, DegP, Tsp, Protease I, Protease Mi, Protease V, Protease VI and combinations thereof. Some *E. coli* protease-deficient strains are available and described in, for example, Joly et al. (1998), *supra*; Georgiou et al., U.S. Patent No. 5,264,365; Georgiou et al., U.S. Patent No. 5,508,192; Hara et al., *Microbial Drug Resistance*, 2:63-72 (1996).

In one embodiment, *E. coli* strains deficient for proteolytic enzymes and transformed with plasmids overexpressing one or more chaperone proteins are used as host cells in the expression system of the invention.

Antibody Purification

In one embodiment, the antibody protein produced herein is further purified to obtain preparations that are substantially homogeneous for further assays and uses. Standard protein purification methods known in the art can be employed. The following procedures are exemplary of suitable purification procedures: fractionation on immunoaffinity or ion-exchange columns, ethanol precipitation, reverse phase HPLC, chromatography on silica or on a cation-exchange resin such as DEAE, chromatofocusing, SDS-PAGE, ammonium sulfate precipitation, and gel filtration using, for example, Sephadex™ G-75.

In one aspect, Protein A immobilized on a solid phase is used for immunoaffinity purification of the antibody products of the invention. Protein A is a 41kD cell wall protein from *Staphylococcus aureus* which binds with a high affinity to the Fc region of antibodies. Lindmark et al (1983) *J. Immunol. Meth.* 62:1-13. The solid phase to which Protein A is immobilized can be a column comprising a glass or silica surface, or a controlled pore glass column or a silicic acid column. In some applications, the column is coated with a reagent, such as glycerol, to possibly prevent nonspecific adherence of contaminants.

As the first step of purification, the preparation derived from the cell culture as described above can be applied onto a Protein A immobilized solid phase to allow specific binding of the antibody of interest to Protein A. The solid phase would then be washed to remove contaminants non-specifically bound to the solid phase. Finally the antibody of interest is recovered from the solid phase by elution.

Generating antibodies using eukaryotic host cells

The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

5 (i) *Signal sequence component*

A vector for use in a eukaryotic host cell may also contain a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide of interest. The heterologous signal sequence selected generally is one that is recognized and processed (*i.e.*, cleaved by a signal peptidase) by the host cell. In mammalian cell expression, mammalian signal
10 sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, are available.

The DNA for such precursor region is ligated in reading frame to DNA encoding the antibody.

(ii) *Origin of replication*

Generally, an origin of replication component is not needed for mammalian expression
15 vectors. For example, the SV40 origin may typically be used only because it contains the early promoter.

(iii) *Selection gene component*

Expression and cloning vectors may contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*,
20 ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, where relevant, or (c) supply critical nutrients not available from complex media.

One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs
25 neomycin, mycophenolic acid and hygromycin.

Another example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the antibody nucleic acid, such as DHFR, thymidine kinase, metallothionein-I and -II (*e.g.*, primate metallothionein genes), adenosine deaminase, ornithine decarboxylase, *etc.*

30 For example, cells transformed with the DHFR selection gene may first be identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. Appropriate host cells when wild-type DHFR is employed include, for example, the Chinese hamster ovary (CHO) cell line deficient in DHFR activity (*e.g.*, ATCC CRL-9096).

Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding an antibody, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH) can be selected by cell growth in medium containing a selection agent for the selectable marker such as an
5 aminoglycosidic antibiotic, *e.g.*, kanamycin, neomycin, or G418. See U.S. Patent No. 4,965,199.

(iv) Promoter component

Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to nucleic acid encoding a polypeptide of interest (*e.g.*, an antibody). Promoter sequences are known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region
10 located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CNCAAT region where N may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

15 Antibody polypeptide transcription from vectors in mammalian host cells can be controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, *e.g.*, the actin promoter or an immunoglobulin promoter, or from heat-shock
20 promoters, provided such promoters are compatible with the host cell systems.

The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment. A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is
25 disclosed in U.S. Patent No. 4,419,446. A modification of this system is described in U.S. Patent No. 4,601,978. See also Reyes *et al.*, *Nature* 297:598-601 (1982) on expression of human β -interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus. Alternatively, the Rous Sarcoma Virus long terminal repeat can be used as the promoter.

(v) Enhancer element component

30 Transcription of DNA encoding an antibody polypeptide of the invention by higher eukaryotes can often be increased by inserting an enhancer sequence into the vector. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus
35 early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and

adenovirus enhancers. See also Yaniv, *Nature* 297:17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the antibody polypeptide-encoding sequence, but is generally located at a site 5' from the promoter.

(vi) *Transcription termination component*

5 Expression vectors used in eukaryotic host cells will typically also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding an antibody. One useful transcription termination
10 component is the bovine growth hormone polyadenylation region. See WO94/11026 and the expression vector disclosed therein.

(vii) *Selection and transformation of host cells*

Suitable host cells for cloning or expressing the DNA in the vectors herein include higher eukaryote cells described herein, including vertebrate host cells. Propagation of vertebrate cells in
15 culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, *J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub *et al.*, *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); mouse sertoli cells
20 (TM4, Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather *et al.*, *Annals N.Y. Acad. Sci.* 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).
25

Host cells are transformed with the above-described expression or cloning vectors for antibody production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

(viii) *Culturing the host cells*

30 The host cells used to produce an antibody of this invention may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ((MEM), (Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham *et al.*, *Meth. Enz.* 58:44 (1979), Barnes *et al.*, *Anal. Biochem.* 102:255 (1980), U.S. Pat. Nos. 4,767,704; 4,657,866;
35 4,927,762; 4,560,655; or 5,122,469; WO 90/03430; WO 87/00195; or U.S. Patent Re. 30,985 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with

hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as GENTAMYCIN™ drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and
 5 glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

(ix) *Purification of antibody*

10 When using recombinant techniques, the antibody can be produced intracellularly, or directly secreted into the medium. If the antibody is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, are generally removed, for example, by centrifugation or ultrafiltration. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter,
 15 for example, an Amicon™ or Millipore Pellicon™ ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants.

The antibody composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with
 20 affinity chromatography being a generally acceptable purification technique. The suitability of affinity reagents such as protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the antibody. Protein A can be used to purify antibodies that are based on human $\gamma 1$, $\gamma 2$, or $\gamma 4$ heavy chains (Lindmark *et al.*, *J. Immunol. Meth.* 62:1-13 (1983)). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss *et al.*, *EMBO J.*
 25 5:15671575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a C_H3 domain, the Bakerbond ABX™ resin (J. T. Baker, Phillipsburg, NJ) is useful for purification. Other techniques for protein purification such as
 30 fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™ chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

Following any preliminary purification step(s), the mixture comprising the antibody of
 35 interest and contaminants may be subjected to further purification steps, as necessary, for example by

low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, generally performed at low salt concentrations (*e.g.*, from about 0-0.25M salt).

It should be noted that, in general, techniques and methodologies for preparing antibodies for use in research, testing and clinical use are well-established in the art, consistent with the above
5 and/or as deemed appropriate by one skilled in the art for the particular antibody of interest.

Activity Assays

Antibodies of the invention can be characterized for their physical/chemical properties and biological functions by various assays known in the art.

Purified antibodies can be further characterized by a series of assays including, but not limited
10 to, N-terminal sequencing, amino acid analysis, non-denaturing size exclusion high pressure liquid chromatography (HPLC), mass spectrometry, ion exchange chromatography and papain digestion.

Where necessary, antibodies are analyzed for their biological activity. In some embodiments, antibodies of the invention are tested for their antigen binding activity. The antigen binding assays that are known in the art and can be used herein include without limitation any direct or competitive
15 binding assays using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), “sandwich” immunoassays, immunoprecipitation assays, fluorescent immunoassays, and protein A immunoassays.

In one embodiment, the invention contemplates an altered antibody that possesses some but not all effector functions, which make it a desirable candidate for many applications in which the half
20 life of the antibody *in vivo* is important yet certain effector functions (such as complement and ADCC) are unnecessary or deleterious. In certain embodiments, the Fc activities of the antibody are measured to ensure that only the desired properties are maintained. *In vitro* and/or *in vivo* cytotoxicity assays can be conducted to confirm the reduction/depletion of CDC and/or ADCC activities. For example, Fc receptor (FcR) binding assays can be conducted to ensure that the antibody lacks FcγR
25 binding (hence likely lacking ADCC activity), but retains FcRn binding ability. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII and FcγRIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991). An example of an *in vitro* assay to assess ADCC activity of a molecule of interest is described in U.S. Patent No. 5,500,362 or 5,821,337. Useful
30 effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, *e.g.*, in a animal model such as that disclosed in Clynes *et al. PNAS (USA)* 95:652-656 (1998). C1q binding assays may also be carried out to confirm that the antibody is unable to bind C1q and hence lacks CDC activity. To assess complement activation, a CDC assay, *e.g.* as described in

Gazzano-Santoro *et al.*, *J. Immunol. Methods* 202:163 (1996), may be performed. FcRn binding and *in vivo* clearance/half life determinations can also be performed using methods known in the art.

Antibody Fragments

5 The present invention encompasses antibody fragments. In certain circumstances there are advantages of using antibody fragments, rather than whole antibodies. The smaller size of the fragments allows for rapid clearance, and may lead to improved access to solid tumors.

 Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g.,
10 Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992); and Brennan *et al.*, *Science*, 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. Fab, Fv and ScFv antibody fragments can all be expressed in and secreted from *E. coli*, thus allowing the facile production of large amounts of these fragments. Antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be
15 directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, *Bio/Technology* 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Fab and F(ab')₂ fragment with increased *in vivo* half-life comprising salvage receptor binding epitope residues are described in U.S. Pat. No. 5,869,046. Other techniques for the production of antibody fragments will be apparent to the skilled
20 practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). See WO 93/16185; U.S. Pat. Nos. 5,571,894; and 5,587,458. Fv and sFv are the only species with intact combining sites that are devoid of constant regions; thus, they are suitable for reduced nonspecific binding during *in vivo* use. sFv fusion proteins may be constructed to yield fusion of an effector protein at either the amino or the carboxy terminus of an sFv. See *Antibody Engineering*, ed.
25 Borrebaeck, *supra*. The antibody fragment may also be a "linear antibody", e.g., as described in U.S. Pat. No. 5,641,870 for example. Such linear antibody fragments may be monospecific or bispecific.

Humanized Antibodies

 The invention encompasses humanized antibodies. Various methods for humanizing non-
30 human antibodies are known in the art. For example, a humanized antibody can have one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.* (1986) *Nature* 321:522-525; Riechmann *et al.* (1988) *Nature* 332:323-327;
35 Verhoeyen *et al.* (1988) *Science* 239:1534-1536), by substituting hypervariable region sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567) wherein substantially less than an intact human

variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some hypervariable region residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

5 The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies can be important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework for the humanized antibody (Sims *et al.* (1993) *J.*
10 *Immunol.* 151:2296; Chothia *et al.* (1987) *J. Mol. Biol.* 196:901. Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.* (1992) *Proc. Natl. Acad. Sci. USA*, 89:4285; Presta *et al.* (1993) *J. Immunol.*, 151:2623.

 It is further generally desirable that antibodies be humanized with retention of high affinity
15 for the antigen and other favorable biological properties. To achieve this goal, according to one method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display
20 probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic,
25 such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

Human antibodies

 Human anti-polyubiquitin antibodies of the invention can be constructed by combining Fv
30 clone variable domain sequence(s) selected from human-derived phage display libraries with known human constant domain sequences(s) as described above. Alternatively, human monoclonal anti-polyubiquitin antibodies of the invention can be made by the hybridoma method. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described, for example, by Kozbor *J. Immunol.*, **133**: 3001 (1984); Brodeur *et al.*, *Monoclonal*

Antibody Production Techniques and Applications, pp. 51-63 (Marcel Dekker, Inc., New York, 1987); and Boerner *et al.*, *J. Immunol.*, **147**: 86 (1991).

It is now possible to produce transgenic animals (e.g. mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (JH) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, e.g., Jakobovits *et al.*, *Proc. Natl. Acad. Sci USA*, **90**: 2551 (1993); Jakobovits *et al.*, *Nature*, **362**: 255 (1993); Bruggermann *et al.*, *Year in Immunol.*, **7**: 33 (1993).

Gene shuffling can also be used to derive human antibodies from non-human, e.g. rodent, antibodies, where the human antibody has similar affinities and specificities to the starting non-human antibody. According to this method, which is also called "epitope imprinting", either the heavy or light chain variable region of a non-human antibody fragment obtained by phage display techniques as described above is replaced with a repertoire of human V domain genes, creating a population of non-human chain/human chain scFv or Fab chimeras. Selection with antigen results in isolation of a non-human chain/human chain chimeric scFv or Fab wherein the human chain restores the antigen binding site destroyed upon removal of the corresponding non-human chain in the primary phage display clone, i.e. the epitope governs (imprints) the choice of the human chain partner. When the process is repeated in order to replace the remaining non-human chain, a human antibody is obtained (see PCT WO 93/06213 published April 1, 1993). Unlike traditional humanization of non-human antibodies by CDR grafting, this technique provides completely human antibodies, which have no FR or CDR residues of non-human origin.

Bispecific Antibodies

Bispecific antibodies are monoclonal antibodies that have binding specificities for at least two different antigens. In certain embodiments, bispecific antibodies are human or humanized antibodies. In certain embodiments, one of the binding specificities is for polyubiquitin including a specific lysine linkage and the other is for any other antigen. In certain embodiments, bispecific antibodies may bind to two different polyubiquitins having two different lysine linkages. Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies).

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, *Nature*, **305**: 537 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody

molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829 published May 13, 1993, and in Traunecker *et al.*, *EMBO J.*, **10**: 3655 (1991).

5 According to a different embodiment, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion, for example, is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. In certain embodiments, the first heavy-chain constant region (CH1), containing the site necessary for light chain binding, is present in at least
10 one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences
15 for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

In one embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin
20 heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies see, for example,
25 Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986).

According to another approach, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The interface comprises at least a part of the C_H3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody
30 molecule are replaced with larger side chains (*e.g.* tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (*e.g.* alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

35 Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (US Patent No.

4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/00373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in US Patent No. 4,676,980, along with a number of cross-linking techniques.

5 Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science*, **229**: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular
10 disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

15 Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby *et al.*, *J. Exp. Med.*, **175**: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells
20 overexpressing the HER2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

 Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, *J. Immunol.*, **148(5)**:1547-1553 (1992). The leucine
25 zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, **90**:6444-6448 (1993) has provided an alternative mechanism for making
30 bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the VH and VL domains of one fragment are forced to pair with the complementary VL and VH domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of
35 single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, *J. Immunol.*, **152**:5368 (1994).

 Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.* *J. Immunol.* **147**: 60 (1991).

Multivalent Antibodies

A multivalent antibody may be internalized (and/or catabolized) faster than a bivalent antibody by a cell expressing an antigen to which the antibodies bind. The antibodies of the present invention can be multivalent antibodies (which are other than of the IgM class) with three or more antigen binding sites (e.g. tetravalent antibodies), which can be readily produced by recombinant expression of nucleic acid encoding the polypeptide chains of the antibody. The multivalent antibody can comprise a dimerization domain and three or more antigen binding sites. The dimerization domain comprises (or consists of), for example, an Fc region or a hinge region. In this scenario, the antibody will comprise an Fc region and three or more antigen binding sites amino-terminal to the Fc region. In one embodiment, a multivalent antibody comprises (or consists of), for example, three to about eight, or four antigen binding sites. The multivalent antibody comprises at least one polypeptide chain (for example, two polypeptide chains), wherein the polypeptide chain(s) comprise two or more variable domains. For instance, the polypeptide chain(s) may comprise VD1-(X1)_n-VD2-(X2)_n-Fc, wherein VD1 is a first variable domain, VD2 is a second variable domain, Fc is one polypeptide chain of an Fc region, X1 and X2 represent an amino acid or polypeptide, and n is 0 or 1. For instance, the polypeptide chain(s) may comprise: VH-CH1-flexible linker-VH-CH1-Fc region chain; or VH-CH1-VH-CH1-Fc region chain. The multivalent antibody herein may further comprise at least two (for example, four) light chain variable domain polypeptides. The multivalent antibody herein may, for instance, comprise from about two to about eight light chain variable domain polypeptides. The light chain variable domain polypeptides contemplated here comprise a light chain variable domain and, optionally, further comprise a CL domain.

Antibody Variants

In some embodiments, amino acid sequence modification(s) of the antibodies described herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of the antibody are prepared by introducing appropriate nucleotide changes into the antibody nucleic acid, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution can be made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid alterations may be introduced in the subject antibody amino acid sequence at the time that sequence is made.

A useful method for identification of certain residues or regions of the antibody that are preferred locations for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham and Wells (1989) *Science*, 244:1081-1085. Here, a residue or group of target residues

are identified (*e.g.*, charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (*e.g.*, alanine or polyalanine) to affect the interaction of the amino acids with antigen. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* need not be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed immunoglobulins are screened for the desired activity.

Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue or the antibody fused to a cytotoxic polypeptide. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (*e.g.* for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the antibody molecule replaced by a different residue. The sites of greatest interest for substitutional mutagenesis include the hypervariable regions, but FR alterations are also contemplated. Conservative substitutions are shown in Table A under the heading of "preferred substitutions". If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions" in Table A, or as further described below in reference to amino acid classes, may be introduced and the products screened.

TABLE A

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Amino acids may be grouped according to similarities in the properties of their side chains (in A. L. Lehninger, in Biochemistry, second ed., pp. 73-75, Worth Publishers, New York (1975)):

(1) non-polar: Ala (A), Val (V), Leu (L), Ile (I), Pro (P), Phe (F), Trp (W), Met (M)

(2) uncharged polar: Gly (G), Ser (S), Thr (T), Cys (C), Tyr (Y), Asn (N), Gln (Q)

(3) acidic: Asp (D), Glu (E)

(4) basic: Lys (K), Arg (R), His(H)

Alternatively, naturally occurring residues may be divided into groups based on common side-chain properties:

(1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;

(2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;

(3) acidic: Asp, Glu;

(4) basic: His, Lys, Arg;

(5) residues that influence chain orientation: Gly, Pro;

(6) aromatic: Trp, Tyr, Phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, into the remaining (non-conserved) sites.

One type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (*e.g.* a humanized or human antibody). Generally, the resulting variant(s) selected for further development will have modified (*e.g.*, improved) biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sites (*e.g.* 6-7 sites) are mutated to generate all possible amino acid substitutions at each site. The antibodies thus generated are displayed from filamentous phage particles as fusions to at least part of a phage coat protein (*e.g.*, the gene III product of M13) packaged within each particle. The phage-displayed variants are then screened for their biological activity (*e.g.* binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, scanning mutagenesis (*e.g.*, alanine scanning) can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues are candidates for substitution according to techniques known in the art, including those elaborated herein. Once such variants are generated, the panel of variants is subjected to screening using techniques known in the art, including those described herein, and antibodies with superior properties in one or more relevant assays may be selected for further development.

Nucleic acid molecules encoding amino acid sequence variants of the antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the antibody.

It may be desirable to introduce one or more amino acid modifications in an Fc region of antibodies of the invention, thereby generating an Fc region variant. The Fc region variant may comprise a human Fc region sequence (*e.g.*, a human IgG1, IgG2, IgG3 or IgG4 Fc region) comprising an amino acid modification (*e.g.* a substitution) at one or more amino acid positions including that of a hinge cysteine.

In accordance with this description and the teachings of the art, it is contemplated that in some embodiments, an antibody of the invention may comprise one or more alterations as compared to the wild type counterpart antibody, *e.g.* in the Fc region. These antibodies would nonetheless retain substantially the same characteristics required for therapeutic utility as compared to their wild type counterpart. For example, it is thought that certain alterations can be made in the Fc region that

would result in altered (*i.e.*, either improved or diminished) C1q binding and/or Complement Dependent Cytotoxicity (CDC), e.g., as described in WO99/51642. See also Duncan & Winter *Nature* 322:738-40 (1988); U.S. Patent No. 5,648,260; U.S. Patent No. 5,624,821; and WO94/29351 concerning other examples of Fc region variants.

5 In one aspect, the invention provides antibodies comprising modifications in the interface of Fc polypeptides comprising the Fc region, wherein the modifications facilitate and/or promote heterodimerization. These modifications comprise introduction of a protuberance into a first Fc polypeptide and a cavity into a second Fc polypeptide, wherein the protuberance is positionable in the cavity so as to promote complexing of the first and second Fc polypeptides. Methods of generating
10 antibodies with these modifications are known in the art, e.g., as described in U.S. Pat. No. 5,731,168.

Immunoconjugates

In another aspect, the invention provides immunoconjugates, or antibody-drug conjugates (ADC), comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, a drug, a growth inhibitory agent, a toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant,
15 or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate).

The use of antibody-drug conjugates for the local delivery of cytotoxic or cytostatic agents, *i.e.* drugs to kill or inhibit tumor cells in the treatment of cancer (Syrigos and Epenetos (1999) *Anticancer Research* 19:605-614; Niculescu-Duvaz and Springer (1997) *Adv. Drg Del. Rev.* 26:151-172; U.S. patent 4,975,278) allows targeted delivery of the drug moiety to tumors, and intracellular
20 accumulation therein, where systemic administration of these unconjugated drug agents may result in unacceptable levels of toxicity to normal cells as well as the tumor cells sought to be eliminated (Baldwin et al., (1986) *Lancet* pp. (Mar. 15, 1986):603-05; Thorpe, (1985) "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review," in *Monoclonal Antibodies '84: Biological And Clinical Applications*, A. Pinchera et al. (ed.s), pp. 475-506). Maximal efficacy with minimal toxicity
25 is sought thereby. Both polyclonal antibodies and monoclonal antibodies have been reported as useful in these strategies (Rowland et al., (1986) *Cancer Immunol. Immunother.*, 21:183-87). Drugs used in these methods include daunomycin, doxorubicin, methotrexate, and vindesine (Rowland et al., (1986) *supra*). Toxins used in antibody-toxin conjugates include bacterial toxins such as diphtheria toxin, plant toxins such as ricin, small molecule toxins such as geldanamycin (Mandler et al (2000) *Jour. of the Nat. Cancer Inst.* 92(19):1573-1581; Mandler et al (2000) *Bioorganic & Med. Chem. Letters* 10:1025-1028; Mandler et al (2002) *Bioconjugate Chem.* 13:786-791), maytansinoids (EP 1391213; Liu et al., (1996) *Proc. Natl. Acad. Sci. USA* 93:8618-8623), and calicheamicin (Lode et al (1998) *Cancer Res.* 58:2928; Hinman et al (1993) *Cancer Res.* 53:3336-3342). The toxins may effect their cytotoxic and cytostatic effects by mechanisms including tubulin binding, DNA binding, or
30

topoisomerase inhibition. Some cytotoxic drugs tend to be inactive or less active when conjugated to large antibodies or protein receptor ligands.

ZEVALIN® (ibritumomab tiuxetan, Biogen/Idec) is an antibody-radioisotope conjugate composed of a murine IgG1 kappa monoclonal antibody directed against the CD20 antigen found on the surface of normal and malignant B lymphocytes and ^{111}In or ^{90}Y radioisotope bound by a thiourea linker-chelator (Wiseman et al (2000) *Eur. Jour. Nucl. Med.* 27(7):766-77; Wiseman et al (2002) *Blood* 99(12):4336-42; Witzig et al (2002) *J. Clin. Oncol.* 20(10):2453-63; Witzig et al (2002) *J. Clin. Oncol.* 20(15):3262-69). Although ZEVALIN has activity against B-cell non-Hodgkin's Lymphoma (NHL), administration results in severe and prolonged cytopenias in most patients. MYLOTARG™ (gemtuzumab ozogamicin, Wyeth Pharmaceuticals), an antibody drug conjugate composed of a hu CD33 antibody linked to calicheamicin, was approved in 2000 for the treatment of acute myeloid leukemia by injection (*Drugs of the Future* (2000) 25(7):686; U.S. Patent Nos. 4970198; 5079233; 5585089; 5606040; 5693762; 5739116; 5767285; 5773001). Cantuzumab mertansine (Immunogen, Inc.), an antibody drug conjugate composed of the huC242 antibody linked via the disulfide linker SPP to the maytansinoid drug moiety, DM1, is tested for the treatment of cancers that express CanAg, such as colon, pancreatic, gastric, and others. MLN-2704 (Millennium Pharm., BZL Biologics, Immunogen Inc.), an antibody drug conjugate composed of the anti-prostate specific membrane antigen (PSMA) monoclonal antibody linked to the maytansinoid drug moiety, DM1, is tested for the potential treatment of prostate tumors. The auristatin peptides, auristatin E (AE) and monomethylauristatin (MMAE), synthetic analogs of dolastatin, were conjugated to chimeric monoclonal antibodies cBR96 (specific to Lewis Y on carcinomas) and cAC10 (specific to CD30 on hematological malignancies) (Doronina et al (2003) *Nature Biotechnology* 21(7):778-784) and are under therapeutic development.

Chemotherapeutic agents useful in the generation of immunoconjugates are described herein (above). Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. See, e.g., WO 93/21232 published October 28, 1993. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re . Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-

diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary
 5 chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

Conjugates of an antibody and one or more small molecule toxins, such as a calicheamicin, maytansinoids, dolostatins, aurostatins, a trichothecene, and CC1065, and the derivatives of these toxins that have toxin activity, are also contemplated herein.

Maytansine and maytansinoids

10 In some embodiments, the immunoconjugate comprises an antibody of the invention conjugated to one or more maytansinoid molecules.

Maytansinoids are mitotic inhibitors which act by inhibiting tubulin polymerization. Maytansine was first isolated from the east African shrub *Maytenus serrata* (U.S. Patent No. 3,896,111). Subsequently, it was discovered that certain microbes also produce maytansinoids, such
 15 as maytansinol and C-3 maytansinol esters (U.S. Patent No. 4,151,042). Synthetic maytansinol and derivatives and analogues thereof are disclosed, for example, in U.S. Patent Nos. 4,137,230; 4,248,870; 4,256,746; 4,260,608; 4,265,814; 4,294,757; 4,307,016; 4,308,268; 4,308,269; 4,309,428; 4,313,946; 4,315,929; 4,317,821; 4,322,348; 4,331,598; 4,361,650; 4,364,866; 4,424,219; 4,450,254; 4,362,663; and 4,371,533.

20 Maytansinoid drug moieties are attractive drug moieties in antibody drug conjugates because they are: (i) relatively accessible to prepare by fermentation or chemical modification, derivatization of fermentation products, (ii) amenable to derivatization with functional groups suitable for conjugation through the non-disulfide linkers to antibodies, (iii) stable in plasma, and (iv) effective against a variety of tumor cell lines.

25 Exemplary embodiments of maytansinoid drug moieties include: DM1; DM3; and DM4. Immunoconjugates containing maytansinoids, methods of making same, and their therapeutic use are disclosed, for example, in U.S. Patent Nos. 5,208,020, 5,416,064 and European Patent EP 0 425 235 B1. Liu et al., Proc. Natl. Acad. Sci. USA 93:8618-8623 (1996) described immunoconjugates comprising a maytansinoid
 30 designated DM1 linked to the monoclonal antibody C242 directed against human colorectal cancer. The conjugate was found to be highly cytotoxic towards cultured colon cancer cells, and showed antitumor activity in an in vivo tumor growth assay. Chari et al., Cancer Research 52:127-131 (1992) describe immunoconjugates in which a maytansinoid was conjugated via a disulfide linker to the murine antibody A7 binding to an antigen on human colon cancer cell lines, or to another murine
 35 monoclonal antibody TA.1 that binds the HER-2/neu oncogene. The cytotoxicity of the TA.1-maytansinoid conjugate was tested in vitro on the human breast cancer cell line SK-BR-3, which expresses 3×10^5 HER-2 surface antigens per cell. The drug conjugate achieved a degree of

cytotoxicity similar to the free maytansinoid drug, which could be increased by increasing the number of maytansinoid molecules per antibody molecule. The A7-maytansinoid conjugate showed low systemic cytotoxicity in mice.

Antibody-maytansinoid conjugates can be prepared by chemically linking an antibody to a maytansinoid molecule without significantly diminishing the biological activity of either the antibody or the maytansinoid molecule. See, e.g., U.S. Patent No. 5,208,020

. An average of 3-4 maytansinoid molecules conjugated per antibody molecule has shown efficacy in enhancing cytotoxicity of target cells without negatively affecting the function or solubility of the antibody, although even one molecule of toxin/antibody would be expected to enhance cytotoxicity over the use of naked antibody. Maytansinoids are well known in the art and can be synthesized by known techniques or isolated from natural sources. Suitable maytansinoids are disclosed, for example, in U.S. Patent No. 5,208,020 and in the other patents and nonpatent publications referred to hereinabove. Maytansinoids include, but are not limited to, maytansinol and maytansinol analogues modified in the aromatic ring or at other positions of the maytansinol molecule, such as various maytansinol esters.

There are many linking groups known in the art for making antibody-maytansinoid conjugates, including, for example, those disclosed in U.S. Patent No. 5,208,020 or EP Patent 0 425 235 B1, Chari et al., Cancer Research 52:127-131 (1992), and U.S. Patent Application No. 10/960,602, filed Oct. 8, 2004,

reference. Antibody-maytansinoid conjugates comprising the linker component SMCC may be prepared as disclosed in U.S. Patent Application No. 10/960,602, filed Oct. 8, 2004. The linking groups include disulfide groups, thioether groups, acid labile groups, photolabile groups, peptidase labile groups, or esterase labile groups, as disclosed in the above-identified patents. Additional linking groups are described and exemplified herein.

Conjugates of the antibody and maytansinoid may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). Coupling agents include, but are not limited to, N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP) (Carlsson et al., Biochem. J. 173:723-737 (1978)) and N-succinimidyl-4-(2-pyridylthio)pentanoate (SPP) to provide for a disulfide linkage.

The linker may be attached to the maytansinoid molecule at various positions, depending on the type of the link. For example, an ester linkage may be formed by reaction with a hydroxyl group

using conventional coupling techniques. The reaction may occur at the C-3 position having a hydroxyl group, the C-14 position modified with hydroxymethyl, the C-15 position modified with a hydroxyl group, and the C-20 position having a hydroxyl group. In one embodiment, the linkage is formed at the C-3 position of maytansinol or a maytansinol analogue.

5 Auristatins and dolostatins

In some embodiments, the immunoconjugate comprises an antibody of the invention conjugated to dolostatins or dolostatin peptidic analogs and derivatives, the auristatins (U.S. Patent Nos. 5635483; 5780588). Dolostatins and auristatins have been shown to interfere with microtubule dynamics, GTP hydrolysis, and nuclear and cellular division (Woyke et al (2001) Antimicrob. Agents and Chemother. 45(12):3580-3584) and have anticancer (U.S. 5663149) and antifungal activity (Pettit et al (1998) Antimicrob. Agents Chemother. 42:2961-2965). The dolastatin or auristatin drug moiety may be attached to the antibody through the N (amino) terminus or the C (carboxyl) terminus of the peptidic drug moiety (WO 02/088172).

Exemplary auristatin embodiments include the N-terminus linked monomethylauristatin drug moieties DE and DF, disclosed in "Monomethylvaline Compounds Capable of Conjugation to Ligands", U.S. Ser. No. 10/983,340, filed Nov. 5, 2004.

Exemplary auristatin embodiments include MMAE and MMAF. Additional exemplary embodiments comprising MMAE or MMAF and various linker components (described further herein) include Ab-MC-vc-PAB-MMAF, Ab-MC-vc-PAB-MMAE, Ab-MC-MMAE and Ab-MC-MMAF.

Typically, peptide-based drug moieties can be prepared by forming a peptide bond between two or more amino acids and/or peptide fragments. Such peptide bonds can be prepared, for example, according to the liquid phase synthesis method (see E. Schröder and K. Lübke, "The Peptides", volume 1, pp 76-136, 1965, Academic Press) that is well known in the field of peptide chemistry. The auristatin/dolastatin drug moieties may be prepared according to the methods of: U.S. 5,635,483; U.S. 5,780,588; Pettit et al (1989) J. Am. Chem. Soc. 111:5463-5465; Pettit et al (1998) Anti-Cancer Drug Design 13:243-277; Pettit, G.R., et al. Synthesis, 1996, 719-725; and Pettit et al (1996) J. Chem. Soc. Perkin Trans. 1 5:859-863. See also Doronina (2003) Nat Biotechnol 21(7):778-784; "Monomethylvaline Compounds Capable of Conjugation to Ligands", U.S. Ser. No. 10/983,340, filed Nov. 5, 2004, hereby incorporated by reference in its entirety (disclosing, e.g., linkers and methods of preparing monomethylvaline compounds such as MMAE and MMAF conjugated to linkers).

Calicheamicin

In other embodiments, the immunoconjugate comprises an antibody of the invention conjugated to one or more calicheamicin molecules. The calicheamicin family of antibiotics is capable of producing double-stranded DNA breaks at sub-picomolar concentrations. For the preparation of conjugates of the calicheamicin family, see U.S. patents 5,712,374, 5,714,586,

5,739,116, 5,767,285, 5,770,701, 5,770,710, 5,773,001, and 5,877,296 (all to American Cyanamid Company). Structural analogues of calicheamicin which may be used include, but are not limited to, γ_1^I , α_2^I , α_3^I , N-acetyl- γ_1^I , PSAG and θ_1^I (Hinman et al., Cancer Research 53:3336-3342 (1993), Lode et al., Cancer Research 58:2925-2928 (1998) and the aforementioned U.S. patents to American Cyanamid). Another anti-tumor drug to which the antibody can be conjugated is QFA, which is an antifolate. Both calicheamicin and QFA have intracellular sites of action and do not readily cross the plasma membrane. Therefore, cellular uptake of these agents through antibody mediated internalization greatly enhances their cytotoxic effects.

Other cytotoxic agents

Other antitumor agents that can be conjugated to the antibodies of the invention include BCNU, streptozocin, vincristine and 5-fluorouracil, the family of agents known collectively LL-E33288 complex described in U.S. patents 5,053,394, 5,770,710, as well as esperamicins (U.S. patent 5,877,296).

Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. See, for example, WO 93/21232 published October 28, 1993.

The present invention further contemplates an immunoconjugate formed between an antibody and a compound with nucleolytic activity (e.g., a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

For selective destruction of the tumor, the antibody may comprise a highly radioactive atom. A variety of radioactive isotopes are available for the production of radioconjugated antibodies.

Examples include At^{211} , I^{131} , I^{125} , Y^{90} , Re^{186} , Re^{188} , Sm^{153} , Bi^{212} , P^{32} , Pb^{212} and radioactive isotopes of Lu. When the conjugate is used for detection, it may comprise a radioactive atom for scintigraphic studies, for example Tc^{99m} or I^{123} , or a spin label for nuclear magnetic resonance (NMR) imaging (also known as magnetic resonance imaging, MRI), such as iodine-123 again, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

The radio- or other labels may be incorporated in the conjugate in known ways. For example, the peptide may be biosynthesized or may be synthesized by chemical amino acid synthesis using suitable amino acid precursors involving, for example, fluorine-19 in place of hydrogen. Labels such as Tc^{99m} or I^{123} , Re^{186} , Re^{188} and In^{111} can be attached via a cysteine residue in the peptide.

Yttrium-90 can be attached via a lysine residue. The IODOGEN method (Fraker et al (1978)

Biochem. Biophys. Res. Commun. 80: 49-57) can be used to incorporate iodine-123. "Monoclonal Antibodies in Immunoscintigraphy" (Chatal, CRC Press 1989) describes other methods in detail.

Conjugates of the antibody and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science 238:1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026. The linker may be a “cleavable linker” facilitating release of the cytotoxic drug in the cell. For example, an acid-labile linker, peptidase-sensitive linker, photolabile linker, dimethyl linker or disulfide-containing linker (Chari et al., Cancer Research 52:127-131 (1992); U.S. Patent No. 5,208,020) may be used.

The compounds of the invention expressly contemplate, but are not limited to, ADC prepared with cross-linker reagents: BMPS, EMCS, GMBS, HBVS, LC-SMCC, MBS, MPBH, SBAP, SIA, SIAB, SMCC, SMPB, SMPH, sulfo-EMCS, sulfo-GMBS, sulfo-KMUS, sulfo-MBS, sulfo-SIAB, sulfo-SMCC, and sulfo-SMPB, and SVSB (succinimidyl-(4-vinylsulfone)benzoate) which are commercially available (e.g., from Pierce Biotechnology, Inc., Rockford, IL., U.S.A). See pages 467-498, 2003-2004 Applications Handbook and Catalog.

Preparation of antibody drug conjugates

In the antibody drug conjugates (ADC) of the invention, an antibody (Ab) is conjugated to one or more drug moieties (D), e.g. about 1 to about 20 drug moieties per antibody, through a linker (L). The ADC of Formula I may be prepared by several routes, employing organic chemistry reactions, conditions, and reagents known to those skilled in the art, including: (1) reaction of a nucleophilic group of an antibody with a bivalent linker reagent, to form Ab-L, via a covalent bond, followed by reaction with a drug moiety D; and (2) reaction of a nucleophilic group of a drug moiety with a bivalent linker reagent, to form D-L, via a covalent bond, followed by reaction with the nucleophilic group of an antibody. Additional methods for preparing ADC are described herein.






The linker may be composed of one or more linker components. Exemplary linker components include 6-maleimidocaproyl (“MC”), maleimidopropanoyl (“MP”), valine-citrulline (“val-cit”), alanine-phenylalanine (“ala-phe”), p-aminobenzyloxycarbonyl (“PAB”), N-Succinimidyl 4-(2-pyridylthio) pentanoate (“SPP”), N-Succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (“SMCC”), and N-Succinimidyl (4-iodo-acetyl) aminobenzoate (“SIAB”). Additional linker components are known in the art and some are described herein. See also “Monomethylvaline


Compounds Capable of Conjugation to Ligands”, U.S. Ser. No. 10/983,340, filed Nov. 5, 2004.

In some embodiments, the linker may comprise amino acid residues. Exemplary amino acid linker components include a dipeptide, a tripeptide, a tetrapeptide or a pentapeptide. Exemplary dipeptides include: valine-citrulline (vc or val-cit), alanine-phenylalanine (af or ala-phe). Exemplary tripeptides include: glycine-valine-citrulline (gly-val-cit) and glycine-glycine-glycine (gly-gly-gly). Amino acid residues which comprise an amino acid linker component include those occurring naturally, as well as minor amino acids and non-naturally occurring amino acid analogs, such as citrulline. Amino acid linker components can be designed and optimized in their selectivity for enzymatic cleavage by a particular enzyme, for example, a tumor-associated protease, cathepsin B, C and D, or a plasmin protease.


Exemplary linker component structures are shown below (wherein the wavy line indicates sites of covalent attachment to other components of the ADC):




Additional exemplary linker components and abbreviations include (wherein the antibody (Ab) and linker are depicted, and p is 1 to about 8):



Val-cit



MC-val-cit



MC-val-cit-PAB

5

10

15

Nucleophilic groups on antibodies include, but are not limited to: (i) N-terminal amine groups, (ii) side chain amine groups, e.g. lysine, (iii) side chain thiol groups, e.g. cysteine, and (iv) sugar hydroxyl or amino groups where the antibody is glycosylated. Amine, thiol, and hydroxyl groups are nucleophilic and capable of reacting to form covalent bonds with electrophilic groups on linker moieties and linker reagents including: (i) active esters such as NHS esters, HOBt esters, haloformates, and acid halides; (ii) alkyl and benzyl halides such as haloacetamides; (iii) aldehydes, ketones, carboxyl, and maleimide groups. Certain antibodies have reducible interchain disulfides, i.e. cysteine bridges. Antibodies may be made reactive for conjugation with linker reagents by treatment with a reducing agent such as DTT (dithiothreitol). Each cysteine bridge will thus form, theoretically, two reactive thiol nucleophiles. Additional nucleophilic groups can be introduced into antibodies

through the reaction of lysines with 2-iminothiolane (Traut's reagent) resulting in conversion of an amine into a thiol. Reactive thiol groups may be introduced into the antibody (or fragment thereof) by introducing one, two, three, four, or more cysteine residues (e.g., preparing mutant antibodies comprising one or more non-native cysteine amino acid residues).

5 Antibody drug conjugates of the invention may also be produced by modification of the antibody to introduce electrophilic moieties, which can react with nucleophilic substituents on the linker reagent or drug. The sugars of glycosylated antibodies may be oxidized, e.g. with periodate oxidizing reagents, to form aldehyde or ketone groups which may react with the amine group of linker reagents or drug moieties. The resulting imine Schiff base groups may form a stable linkage, or may
10 be reduced, e.g. by borohydride reagents to form stable amine linkages. In one embodiment, reaction of the carbohydrate portion of a glycosylated antibody with either galactose oxidase or sodium meta-periodate may yield carbonyl (aldehyde and ketone) groups in the protein that can react with appropriate groups on the drug (Hermanson, *Bioconjugate Techniques*). In another embodiment, proteins containing N-terminal serine or threonine residues can react with sodium meta-periodate,
15 resulting in production of an aldehyde in place of the first amino acid (Geoghegan & Stroh, (1992) *Bioconjugate Chem.* 3:138-146; U.S. 5362852). Such aldehyde can be reacted with a drug moiety or linker nucleophile.

Likewise, nucleophilic groups on a drug moiety include, but are not limited to: amine, thiol, hydroxyl, hydrazide, oxime, hydrazine, thiosemicarbazone, hydrazine carboxylate, and arylhydrazide
20 groups capable of reacting to form covalent bonds with electrophilic groups on linker moieties and linker reagents including: (i) active esters such as NHS esters, HOBt esters, haloformates, and acid halides; (ii) alkyl and benzyl halides such as haloacetamides; (iii) aldehydes, ketones, carboxyl, and maleimide groups.

Alternatively, a fusion protein comprising the antibody and cytotoxic agent may be made,
25 e.g., by recombinant techniques or peptide synthesis. The length of DNA may comprise respective regions encoding the two portions of the conjugate either adjacent to one another or separated by a region encoding a linker peptide which does not destroy the desired properties of the conjugate.

In yet another embodiment, the antibody may be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pre-targeting wherein the antibody-receptor conjugate is
30 administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) which is conjugated to a cytotoxic agent (e.g., a radionucleotide).

Antibody (Ab)-MC-MMAE may be prepared by conjugation of any of the antibodies provided herein with MC-MMAE as follows. Antibody, dissolved in 500 mM sodium borate and 500
35 mM sodium chloride at pH 8.0 is treated with an excess of 100 mM dithiothreitol (DTT). After incubation at 37 °C for about 30 minutes, the buffer is exchanged by elution over Sephadex G25 resin and eluted with PBS with 1 mM DTPA. The thiol/Ab value is checked by determining the reduced

antibody concentration from the absorbance at 280 nm of the solution and the thiol concentration by reaction with DTNB (Aldrich, Milwaukee, WI) and determination of the absorbance at 412 nm. The reduced antibody dissolved in PBS is chilled on ice. The drug linker reagent, maleimidocaproyl-monomethyl auristatin E (MMAE), i.e. MC-MMAE, dissolved in DMSO, is diluted in acetonitrile and water at known concentration, and added to the chilled reduced antibody 2H9 in PBS. After about one hour, an excess of maleimide is added to quench the reaction and cap any unreacted antibody thiol groups. The reaction mixture is concentrated by centrifugal ultrafiltration and 2H9-MC-MMAE is purified and desalted by elution through G25 resin in PBS, filtered through 0.2 μ m filters under sterile conditions, and frozen for storage.

Antibody-MC-MMAF may be prepared by conjugation of any of the antibodies provided herein with MC-MMAF following the protocol provided for preparation of Ab-MC-MMAE.

Antibody-MC-val-cit-PAB-MMAE is prepared by conjugation of any of the antibodies provided herein with MC-val-cit-PAB-MMAE following the protocol provided for preparation of Ab-MC-MMAE.

Antibody-MC-val-cit-PAB-MMAF is prepared by conjugation of any of the antibodies provided herein with MC-val-cit-PAB-MMAF following the protocol provided for preparation of Ab-MC-MMAE.

Antibody-SMCC-DM1 is prepared by conjugation of any of the antibodies provided herein with SMCC-DM1 as follows. Purified antibody is derivatized with (Succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC, Pierce Biotechnology, Inc) to introduce the SMCC linker. Specifically, antibody is treated at 20 mg/mL in 50 mM potassium phosphate/ 50 mM sodium chloride/ 2 mM EDTA, pH 6.5 with 7.5 molar equivalents of SMCC (20 mM in DMSO, 6.7 mg/mL). After stirring for 2 hours under argon at ambient temperature, the reaction mixture is filtered through a Sephadex G25 column equilibrated with 50mM potassium phosphate/ 50 mM sodium chloride/ 2 mM EDTA, pH 6.5. Antibody-containing fractions are pooled and assayed.

Antibody-SMCC prepared thusly is diluted with 50mM potassium phosphate/50 mM sodium chloride/2 mM EDTA, pH 6.5, to a final concentration of about 10 mg/ml, and reacted with a 10 mM solution of DM1 in dimethylacetamide. The reaction is stirred at ambient temperature under argon for 16.5 hours. The conjugation reaction mixture is filtered through a Sephadex G25 gel filtration column (1.5 x 4.9 cm) with 1 x PBS at pH 6.5. The DM1 drug to antibody ratio (p) may be about 2 to 5, as measured by the absorbance at 252 nm and at 280 nm.

Ab-SPP-DM1 is prepared by conjugation of any of the antibodies provided herein with SPP-DM1 as follows. Purified antibody is derivatized with N-succinimidyl-4-(2-pyridylthio)pentanoate to introduce dithiopyridyl groups. Antibody (376.0 mg, 8 mg/mL) in 44.7 mL of 50 mM potassium phosphate buffer (pH 6.5) containing NaCl (50 mM) and EDTA (1 mM) is treated with SPP (5.3 molar equivalents in 2.3 mL ethanol). After incubation for 90 minutes under argon at ambient temperature, the reaction mixture is gel filtered through a Sephadex G25 column equilibrated with a

35 mM sodium citrate, 154 mM NaCl, 2 mM EDTA buffer. Antibody-containing fractions were pooled and assayed. The degree of modification of the antibody is determined as described above.

Antibody-SPP-Py (about 10 μ moles of releasable 2-thiopyridine groups) is diluted with the above 35 mM sodium citrate buffer, pH 6.5, to a final concentration of about 2.5 mg/mL. DM1 (1.7
5 equivalents, 17 μ moles) in 3.0 mM dimethylacetamide (DMA, 3% v/v in the final reaction mixture) is then added to the antibody solution. The reaction proceeds at ambient temperature under argon for about 20 hours. The reaction is loaded on a Sephacryl™ S3000 gel filtration column (5.0 cm x 90.0 cm, 1.77 L) equilibrated with 35 mM sodium citrate, 154 mM NaCl, pH 6.5. The flow rate may be about 5.0 mL/min and 65 fractions (20.0 mL each) are collected. The number of DM1 drug molecules
10 linked per antibody molecule (p') is determined by measuring the absorbance at 252 nm and 280 nm, and may be about 2 to 4 DM1 drug moieties per 2H9 antibody.

Antibody-BMPEO-DM1 is prepared by conjugation of any of the antibodies provided herein with BMPEO-DM1 as follows. The antibody is modified by the bis-maleimido reagent BM(PEO)₄ (Pierce Chemical), leaving an unreacted maleimido group on the surface of the antibody. This may be
15 accomplished by dissolving BM(PEO)₄ in a 50% ethanol/water mixture to a concentration of 10 mM and adding a tenfold molar excess to a solution containing antibody in phosphate buffered saline at a concentration of approximately 1.6 mg/ml (10 micromolar) and allowing it to react for 1 hour to form an antibody-linker intermediate, 2H9-BMPEO. Excess BM(PEO)₄ is removed by gel filtration (HiTrap column, Pharmacia) in 30 mM citrate, pH 6 with 150 mM NaCl buffer. An approximate 10
20 fold molar excess DM1 is dissolved in dimethyl acetamide (DMA) and added to the 2H9-BMPEO intermediate. Dimethyl formamide (DMF) may also be employed to dissolve the drug moiety reagent. The reaction mixture is allowed to react overnight before gel filtration or dialysis into PBS to remove unreacted DM1. Gel filtration on S200 columns in PBS is used to remove high molecular weight aggregates and to furnish purified 2H9-BMPEO-DM1.

25 Antibody Derivatives

Antibodies of the invention can be further modified to contain additional nonproteinaceous moieties that are known in the art and readily available. In one embodiment, the moieties suitable for derivatization of the antibody are water soluble polymers. Non-limiting examples of water soluble polymers include, but are not limited to, polyethylene glycol (PEG), copolymers of ethylene
30 glycol/propylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), and dextran or poly(n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (e.g., glycerol), polyvinyl alcohol, and mixtures thereof. Polyethylene
35 glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The

polymer may be of any molecular weight, and may be branched or unbranched. The number of polymers attached to the antibody may vary, and if more than one polymer is attached, the polymers can be the same or different molecules. In general, the number and/or type of polymers used for derivatization can be determined based on considerations including, but not limited to, the particular properties or functions of the antibody to be improved, whether the antibody derivative will be used in a therapy under defined conditions, etc.

In another embodiment, conjugates of an antibody and nonproteinaceous moiety that may be selectively heated by exposure to radiation are provided. In one embodiment, the nonproteinaceous moiety is a carbon nanotube (Kam et al., Proc. Natl. Acad. Sci. 102: 11600-11605 (2005)). The radiation may be of any wavelength, and includes, but is not limited to, wavelengths that do not harm ordinary cells, but which heat the nonproteinaceous moiety to a temperature at which cells proximal to the antibody-nonproteinaceous moiety are killed.

Pharmaceutical Formulations

Therapeutic formulations comprising an antibody of the invention are prepared for storage by mixing the antibody having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of aqueous solutions, lyophilized or other dried formulations. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, histidine and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, including, but not limited to those with complementary activities that do not adversely affect each other. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsule prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsule and poly-(methacrylate) microcapsule, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980).

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the immunoglobulin of the invention, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsule. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated immunoglobulins remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Uses

An antibody of the invention may be used in, for example, *in vitro*, *ex vivo* and *in vivo* therapeutic methods. Antibodies of the invention can be used as an antagonist to partially or fully block the specific antigen activity *in vitro*, *ex vivo* and/or *in vivo*. Moreover, at least some of the antibodies of the invention can neutralize antigen activity from other species. Accordingly, antibodies of the invention can be used to inhibit a specific antigen activity, *e.g.*, in a cell culture containing the antigen, in human subjects or in other mammalian subjects having the antigen with which an antibody of the invention cross-reacts (*e.g.* chimpanzee, baboon, marmoset, cynomolgus and rhesus, pig or mouse). In one embodiment, an antibody of the invention can be used for inhibiting antigen activities

by contacting the antibody with the antigen such that antigen activity is inhibited. In one embodiment, the antigen is a human protein molecule.

In one embodiment, an antibody of the invention can be used in a method for inhibiting an antigen in a subject suffering from a disorder in which the antigen activity is detrimental, comprising
5 administering to the subject an antibody of the invention such that the antigen activity in the subject is inhibited. In one embodiment, the antigen is a human protein molecule and the subject is a human subject. Alternatively, the subject can be a mammal expressing the antigen with which an antibody of the invention binds. Still further the subject can be a mammal into which the antigen has been introduced (e.g., by administration of the antigen or by expression of an antigen transgene). An
10 antibody of the invention can be administered to a human subject for therapeutic purposes. Moreover, an antibody of the invention can be administered to a non-human mammal expressing an antigen with which the antibody cross-reacts (e.g., a primate, pig or mouse) for veterinary purposes or as an animal model of human disease. Regarding the latter, such animal models may be useful for evaluating the therapeutic efficacy of antibodies of the invention (e.g., testing of dosages and time courses of
15 administration). Antibodies of the invention can be used to treat, inhibit, delay progression of, prevent/delay recurrence of, ameliorate, or prevent diseases, disorders or conditions associated with abnormal expression and/or activity of polyubiquitins and polyubiquitinated proteins, including but not limited to cancer, muscular disorders, ubiquitin-pathway-related genetic disorders, immune/inflammatory disorders, neurological disorders, and other ubiquitin pathway-related
20 disorders.

In one aspect, a blocking antibody of the invention is specific for a polyubiquitin having a particular lysine linkage, and inhibits normal polyubiquitin activity by blocking or interfering with the interaction between a polyubiquitin having a particular lysine linkage and a protein that interacts with that polyubiquitin, thereby inhibiting the corresponding signal pathway and other associated
25 molecular or cellular events.

In certain embodiments, an immunoconjugate comprising an antibody conjugated with a cytotoxic agent is administered to the patient. In some embodiments, the immunoconjugate and/or antigen to which it is bound is/are internalized by the cell, resulting in increased therapeutic efficacy of the immunoconjugate in killing the target cell to which it binds. In one embodiment, the cytotoxic
30 agent targets or interferes with nucleic acid in the target cell. Examples of such cytotoxic agents include any of the chemotherapeutic agents noted herein (such as a maytansinoid or a calicheamicin), a radioactive isotope, or a ribonuclease or a DNA endonuclease.

Antibodies of the invention can be used either alone or in combination with other compositions in a therapy. For instance, an antibody of the invention may be co-administered with
35 another antibody, and/or adjuvant/therapeutic agents (e.g., steroids). For instance, an antibody of the

invention may be combined with an anti-inflammatory and/or antiseptic in a treatment scheme, e.g. in treating any of the diseases described herein, including cancer, muscular disorders, ubiquitin-pathway-related genetic disorders, immune/inflammatory disorders, neurological disorders, and other ubiquitin pathway-related disorders . Such combined therapies noted above include combined
5 administration (where the two or more agents are included in the same or separate formulations), and separate administration, in which case, administration of the antibody of the invention can occur prior to, and/or following, administration of the adjunct therapy or therapies.

An antibody of the invention (and adjunct therapeutic agent) can be administered by any suitable means, including parenteral, subcutaneous, intraperitoneal, intrapulmonary, and intranasal,
10 and, if desired for local treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. In addition, the antibody is suitably administered by pulse infusion, particularly with declining doses of the antibody. Dosing can be by any suitable route, e.g. by injections, such as intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic.

15 The location of the binding target of an antibody of the invention may be taken into consideration in preparation and administration of the antibody. When the binding target is an intracellular molecule, certain embodiments of the invention provide for the antibody or antigen-binding fragment thereof to be introduced into the cell where the binding target is located. In one embodiment, an antibody of the invention can be expressed intracellularly as an intrabody. The term
20 “intrabody,” as used herein, refers to an antibody or antigen-binding portion thereof that is expressed intracellularly and that is capable of selectively binding to a target molecule, as described in Marasco, Gene Therapy 4: 11-15 (1997); Kontermann, Methods 34: 163-170 (2004); U.S. Patent Nos. 6,004,940 and 6,329,173; U.S. Patent Application Publication No. 2003/0104402, and PCT Publication No. WO2003/077945. Intracellular expression of an intrabody is effected by introducing
25 a nucleic acid encoding the desired antibody or antigen-binding portion thereof (lacking the wild-type leader sequence and secretory signals normally associated with the gene encoding that antibody or antigen-binding fragment) into a target cell. Any standard method of introducing nucleic acids into a cell may be used, including, but not limited to, microinjection, ballistic injection, electroporation, calcium phosphate precipitation, liposomes, and transfection with retroviral, adenoviral, adeno-
30 associated viral and vaccinia vectors carrying the nucleic acid of interest. One or more nucleic acids encoding all or a portion of an anti-polyubiquitin antibody of the invention can be delivered to a target cell, such that one or more intrabodies are expressed which are capable of intracellular binding to a polyubiquitin and modulation of one or more polyubiquitin-mediated cellular pathways.

In another embodiment, internalizing antibodies are provided. Antibodies can possess certain
35 characteristics that enhance delivery of antibodies into cells, or can be modified to possess such characteristics. Techniques for achieving this are known in the art. For example, cationization of an

antibody is known to facilitate its uptake into cells (see, e.g., U.S. Patent No. 6,703,019).

Lipofections or liposomes can also be used to deliver the antibody into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is generally advantageous. For example, based upon the variable-region sequences
5 of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco et al., Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993).

Entry of modulator polypeptides into target cells can be enhanced by methods known in the art. For example, certain sequences, such as those derived from HIV Tat or the Antennapedia
10 homeodomain protein are able to direct efficient uptake of heterologous proteins across cell membranes. See, e.g., Chen et al., Proc. Natl. Acad. Sci. USA (1999), 96:4325-4329.

When the binding target is located in the brain, certain embodiments of the invention provide for the antibody or antigen-binding fragment thereof to traverse the blood-brain barrier. Certain neurodegenerative diseases are associated with an increase in permeability of the blood-brain barrier,
15 such that the antibody or antigen-binding fragment can be readily introduced to the brain. When the blood-brain barrier remains intact, several art-known approaches exist for transporting molecules across it, including, but not limited to, physical methods, lipid-based methods, and receptor and channel-based methods.

Physical methods of transporting the antibody or antigen-binding fragment across the blood-
20 brain barrier include, but are not limited to, circumventing the blood-brain barrier entirely, or by creating openings in the blood-brain barrier. Circumvention methods include, but are not limited to, direct injection into the brain (see, e.g., Papanastassiou et al., Gene Therapy 9: 398-406 (2002)), interstitial infusion/convection-enhanced delivery (see, e.g., Bobo et al., Proc. Natl. Acad. Sci. USA 91: 2076-2080 (1994)), and implanting a delivery device in the brain (see, e.g., Gill et al., Nature
25 Med. 9: 589-595 (2003); and Gliadel Wafers™, Guildford Pharmaceutical). Methods of creating openings in the barrier include, but are not limited to, ultrasound (see, e.g., U.S. Patent Publication No. 2002/0038086), osmotic pressure (e.g., by administration of hypertonic mannitol (Neuwelt, E. A., Implication of the Blood-Brain Barrier and its Manipulation, Vols 1 & 2, Plenum Press, N.Y. (1989))), permeabilization by, e.g., bradykinin or permeabilizer A-7 (see, e.g., U.S. Patent Nos.
30 5,112,596, 5,268,164, 5,506,206, and 5,686,416), and transfection of neurons that straddle the blood-brain barrier with vectors containing genes encoding the antibody or antigen-binding fragment (see, e.g., U.S. Patent Publication No. 2003/0083299).

Lipid-based methods of transporting the antibody or antigen-binding fragment across the blood-brain barrier include, but are not limited to, encapsulating the antibody or antigen-binding
35 fragment in liposomes that are coupled to antibody binding fragments that bind to receptors on the vascular endothelium of the blood-brain barrier (see, e.g., U.S. Patent Application Publication No. 20020025313), and coating the antibody or antigen-binding fragment in low-density lipoprotein

particles (see, e.g., U.S. Patent Application Publication No. 20040204354) or apolipoprotein E (see, e.g., U.S. Patent Application Publication No. 20040131692).

Receptor and channel-based methods of transporting the antibody or antigen-binding fragment across the blood-brain barrier include, but are not limited to, using glucocorticoid blockers to increase permeability of the blood-brain barrier (see, e.g., U.S. Patent Application Publication Nos. 2002/0065259, 2003/0162695, and 2005/0124533); activating potassium channels (see, e.g., U.S. Patent Application Publication No. 2005/0089473), inhibiting ABC drug transporters (see, e.g., U.S. Patent Application Publication No. 2003/0073713); coating antibodies with a transferrin and modulating activity of the one or more transferrin receptors (see, e.g., U.S. Patent Application Publication No. 2003/0129186), and cationizing the antibodies (see, e.g., U.S. Patent No. 5,004,697).

The antibody composition of the invention would be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The antibody need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of antibodies of the invention present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as described herein, or about from 1 to 99% of the dosages described herein, or in any dosage and by any route that is empirically/clinically determined to be appropriate.

For the prevention or treatment of disease, the appropriate dosage of an antibody of the invention (when used alone or in combination with other agents such as chemotherapeutic agents) will depend on the type of disease to be treated, the type of antibody, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments. Depending on the type and severity of the disease, about 1 µg/kg to 15 mg/kg (*e.g.* 0.1mg/kg-10mg/kg) of antibody can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. One typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment would generally be sustained until a desired suppression of disease symptoms occurs. One exemplary dosage of the antibody would be in the range from about 0.05mg/kg to about 10mg/kg. Thus, one or more doses of about 0.5mg/kg, 2.0mg/kg, 4.0mg/kg or 10mg/kg (or any combination thereof) may be administered to the patient. Such doses may be administered

intermittently, *e.g.* every week or every three weeks (*e.g.* such that the patient receives from about two to about twenty, or *e.g.* about six doses of the antibody). An initial higher loading dose, followed by one or more lower doses may be administered. An exemplary dosing regimen comprises administering an initial loading dose of about 4 mg/kg, followed by a weekly maintenance dose of about 2 mg/kg of the antibody. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

Articles of Manufacture

In another aspect of the invention, an article of manufacture containing materials useful for the treatment, prevention and/or diagnosis of the disorders described above is provided. The article of manufacture comprises a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is by itself or when combined with another composition effective for treating, preventing and/or diagnosing the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is an antibody of the invention. The label or package insert indicates that the composition is used for treating the condition of choice. Moreover, the article of manufacture may comprise (a) a first container with a composition contained therein, wherein the composition comprises an antibody of the invention; and (b) a second container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the compositions can be used to treat a particular condition. Alternatively, or additionally, the article of manufacture may further comprise a second (or third) container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

The following are examples of the methods and compositions of the invention. It is understood that various other embodiments may be practiced, given the general description provided above.

EXAMPLES**EXAMPLE 1: ISOLATION AND CHARACTERIZATION OF FIRST GENERATION ANTI-POLYUBIQUITIN ANTIBODIES****A) Library Sorting**

- 5 Phage from naïve antibody libraries were subjected to binding selection against immobilized synthetic peptides including an isopeptide bond mimicking either K48-linked polyubiquitin or K63-linked polyubiquitin. No enrichment was observed after six rounds of selection. The synthetic peptides were lengthened, and the naïve antibody libraries were again screened. Once again, no enrichment was observed after six rounds of binding selection.
- 10 Phage from the naïve YS-B antibody library were subjected to four rounds of binding selection against polyubiquitin chains having different known ubiquitin isopeptide linkages. The YS-B antibody library contains randomized amino acids in all three heavy chain CDRs and light chain CDR3 (see U.S. Published Patent Application No. 2005-0106667), and is based on humanized antibody 4D5.
- 15 Enzymatically synthesized full-length K48-linked or K63-linked polyubiquitin chains of 3 to 7 units in length (Boston Biochem) were immobilized on 96-well Maxisorp™ immunoplates (NUNC). The plates were coated overnight with 5 µg/ml of K48- or K63-linked polyubiquitin in 50 mM carbonate buffer, pH 9.6. The coated plates were washed with phosphate buffered saline (PBS) and blocked with either bovine serum albumin (BSA) or casein, at a concentration of 0.2% in PBS. The
- 20 plates were subsequently washed with PBS containing 0.05% Tween 20 (PBST) at 25°C. Each well was incubated with 100 µl of 10^{12} phage/ml in sorting buffer (BSA or casein, 0.2% in PBST) for 2 hours at room temperature. Each well was washed eight times in PBST to remove unbound phage. Bound phage were eluted by incubation with 0.1 M HCl for 10 minutes, and the eluant was neutralized with 2 M Tris base. The eluted phage were propagated in *Escherichia coli* XL1-blue
- 25 (Stratagene) with the addition of M13-KO7 helper phage (New England Biolabs).
- Amplified phage were used for additional rounds of selection against the same target that was used in the previous round. For selection rounds two to four, 10 µg/ml ubiquitin was included in the sorting buffer as a counterselection. Rounds three and four were also sorted both with and without additional counterselection: either 10 µg/ml K63-linked polyubiquitin in the K48-linked polyubiquitin
- 30 selection or 10 µg/ml K48-linked polyubiquitin in the K63-linked polyubiquitin selection.
- For those clones where selection rounds two to four lacked polyubiquitin counterselection, individual clones were grown in a 96-well format in 400 µl of 2YT broth supplemented with carbenicillin and M13-KO7 helper phage. Supernatants from those cultures were used in high-throughput phage ELISAs to screen clones for binding to K48-linked polyubiquitin, K63-linked
- 35 polyubiquitin, ubiquitin and BSA. All clones were subjected to DNA sequence analysis. A portion of the heavy chain that includes the HVR was sequenced, allowing for analysis of the heavy chain

hypervariable regions. The heavy chain hypervariable regions for the clones recognizing K48-linked polyubiquitin or both K48-linked and K63-linked polyubiquitin are shown in Figures 2A-C. The heavy chain hypervariable regions for the clones recognizing K63-linked polyubiquitin or both K48-linked and K63-linked polyubiquitin are shown in Figures 3A and 3B. The light chain HVR for each clone was not sequenced, but based on the nature of the YS-B library, the sequences of HVR-L1 and HVR-L2 were expected to be invariant, while the HVR-L3 sequence was expected to be clone-specific. The HVR-L1 sequence is RASQSVSSAVA (SEQ ID NO: 79) and the HVR-L2 sequence is SASSLYS (SEQ ID NO: 80), according to the library design. All clones had the same heavy chain and light chain framework sequences (see Figure 6).

A different set of clones included counterselection with 10 µg/ml of polyubiquitin of a different lysine linkage (either K48-linked polyubiquitin or K63-linked polyubiquitin) in selection rounds two to four. Ten µl of the phage eluted from the fourth selection round was used to infect growing *E. coli* CJ236 for 20 minutes, which were then grown overnight on solid agar containing carbenicillin. Fifteen milliliters of 2YT broth supplemented with carbenicillin and chloramphenicol was added to the plate to resuspend the phagemid-containing CJ236 cells. M13-KO7 helper phage was added. After incubation for one hour at 37°C with agitation, 2.5 ml of the suspension was added to 250 ml of 2YT broth supplemented with carbenicillin and kanamycin. The suspension was allowed to grow overnight.

The phage were harvested by polyethylene glycol precipitation, and the Kunkel DNA was isolated using an M13 spin kit (Qiagen). One microgram of Kunkel template was used for Kunkel mutagenesis (see Kunkel, Proc. Natl. Acad. Sci. USA 82:488 (1985)) with the oligonucleotide F1560-2 (TCTTGTGACAAACTCACCATCACCATCACCATCACTA GGGCGGTGGCTCTGGTTCCGGTGATTTT) (SEQ ID NO: 150). The mutagenesis reaction was transformed in *E. coli* XL1-blue and grown on solid agar containing carbenicillin overnight.

Individual clones were then picked and grown as above and screened against K48- and K63-linked polyubiquitin, monoubiquitin, BSA and anti-pentaHis antibody (Qiagen) in a phage ELISA as described above. Clones identified to be specific for K48-linked polyubiquitin or for K63-linked polyubiquitin were subjected to DNA sequence analysis. The amino acid sequences of hypervariable regions HVR-H1, HVR-H2, and HVR-H3 are shown in Figures 8A-C (specific for K48-linked polyubiquitin) and 9A and 9B (specific for K63-linked polyubiquitin). The light chain HVR for each clone was not sequenced, but based on the nature of the YS-B library, the sequences of HVR-L1 and HVR-L2 were expected to be invariant, while the HVR-L3 sequence was expected to be clone-specific. The HVR-L1 sequence is RASQSVSSAVA (SEQ ID NO: 79) and the HVR-L2 sequence is SASSLYS (SEQ ID NO: 80), according to the library design. All clones had the same heavy chain and light chain framework sequences (see Figure 6).

(B) Fab Production

For Fab production, phage supernatants from unique clones positive for the pentahistidine tag were used to infect *E. coli* FF34B8, a cell line in which the F' episome was added to the 34B8 strain by mating with XL1 and culturing in selective solid media. Infected cells were streaked on solid agar containing carbenicillin and grown overnight. Single colonies of FF34B8 containing phagemid were picked from the plates and grown overnight at 37°C in LB containing carbenicillin. Those cultures were then used to inoculate 500 ml complete CRAP media (3.57 g (NH₄)₂SO₄, 0.71 g sodium citrate · 2H₂O, 1.07 g KCl, 5.36 g yeast extract (certified), 5.36 g Hycase™ SF (Sheffield), pH adjusted to 7.3 by addition of KOH and volume adjusted to 872 mL with ultrapure water, autoclaved, cooled to 55 °C, to which was added (per L) 110 mL 1 M MOPS pH 7.3, 11 mL 50% glucose, and 7 mL 1 M MgSO₄) containing carbenicillin, and grown for 24 hours at 30°C with agitation. Cells were harvested by centrifugation and the cell pellets were stored at -20°C. Fabs were purified by resuspending each cell pellet in 35 ml cold wash buffer (PBS + 150 mM NaCl) containing 0.2 mg/ml lysozyme and 0.3 U/mL Dnase I. Resuspended cell pellets were transferred to 50 ml centrifuge tubes and vortexed rapidly at 25°C for 45 minutes. The pellets were centrifuged and the lysate was loaded on 1 ml protein G-sepharose columns preequilibrated with wash buffer at 4°C. The columns were washed with 50 ml cold wash buffer, eluted with 3 ml 0.1 M acetic acid, and neutralized with 150 µl of 2 M Tris base. The eluted Fabs were buffer-exchanged into PBS and concentrated using Centriprep™ 10 centrifuge filters (Millipore). The resulting Fab concentrations were determined spectrophotometrically (1 OD_{280nm} = 1.55mg/mL). The concentrated Fabs were stored at 4°C.

Each Fab was included in an ELISA protein assay, as described above, to determine its relative affinity for K48-linked and K63-linked polyubiquitin, and to confirm that the Fab was not reactive with monoubiquitin or BSA. The Fabs apu01-15 had greater specificity for K48-linked polyubiquitin than for K63-linked polyubiquitin. Fabs apu17-apu24 demonstrated greater specificity for K63-linked polyubiquitin than for K48-linked polyubiquitin in the ELISA. Apu16 was not produced as a Fab.

All Fabs were subjected to DNA sequence analysis. The amino acid sequences of hypervariable regions HVR-H1, HVR-H2, HVR-H3, and HVR-L3 for each Fab that bound specifically to K48-linked polyubiquitin are shown in Figures 10A-C. The amino acid sequences of hypervariable regions HVR-H1, HVR-H2, HVR-H3, and HVR-L3 for each Fab that bound specifically to K63-linked polyubiquitin are shown in Figures 11A-C. The heavy chain and light chain framework sequences for each Fab appear in Figure 6. The first two light chain hypervariable regions, HVR-L1 and HVR-L2, were identical for each clone, according to the library design (see SEQ ID NOs: 79 and 80, above).

(C) Affinity Analysis of Isolated Fabs

The affinities of selected Fabs (see section (B), above) for ubiquitin and its lysine-linked forms were determined by surface plasmon resonance using a BIACORE® 3000 system (Biacore).

Approximately 100 resonance units of ubiquitin, K48- or K63-linked diubiquitin, or K48- or K63-linked polyubiquitin (chain lengths 3 to 7) were immobilized on different flow cells of CM5 chips using the amine coupling protocol supplied by the manufacturer. In each experiment, flow cell 1 was activated and ethanolamine-blocked without immobilizing protein, to be used for reference

5 subtraction. Serial dilutions of Fab proteins (1.6-100 nM) of each of apu01 to apu24 were injected (50 μ l total at a flow rate of 25 μ l/minute) over each flow cell. The signal for each flow cell was recorded and the reference signal was subtracted. Following a dissociation period (5 minutes), the chip surface was regenerated with 13 μ l of 20 mM HCl. Exemplary binding curves for the Fabs apu09 and apu18 are shown in Figures 12 and 13. Apu09 binds to K48-linked polyubiquitin, but not to K63-linked
10 polyubiquitin, as shown in Figure 12. Figure 13A shows the binding curves for apu18 to K48-linked polyubiquitin. While some binding is observed, it is substantially less than the binding observed to K63-linked polyubiquitin (Figure 13B). Similar analyses were performed for each Fab.

Kinetic constants and binding constants were simultaneously calculated by nonlinear regression analysis using software provided by the manufacturer, and are shown in Table B. The
15 language "NB" in Table B indicates that no binding was detected for the indicated interaction. The language "nd" in Table B indicates that no data was measured for the indicated interaction. The results show that the kinetic constants of a particular Fab for binding to diubiquitin are very similar to those for binding to a polyubiquitin. Thus, the Fabs appear to recognize a particular isopeptide linkage between two ubiquitin moieties.

Attorney Docket P2260R1

TABLE B: Kinetic Constants of Anti-Polyubiquitin Fabs as Measured by BIACORE® Analysis

	K48-linked polyubiquitin			K63-linked polyubiquitin			K48-linked diubiquitin			K63-linked diubiquitin		
	k_{on} (1/M·s)	k_{off} (1/s)	K_d (nM)	k_{on} (1/M·s)	k_{off} (1/s)	K_d (nM)	k_{on} (1/M·s)	k_{off} (1/s)	K_d (nM)	k_{on} (1/M·s)	k_{off} (1/s)	K_d (nM)
Apu01	1.92×10^6	3.49×10^{-3}	1.82			NB	3.28×10^6	1.38×10^{-2}	4.20			NB
Apu02	nd	nd	nd	nd	nd	nd	5.46×10^5	6.96×10^{-3}	12.70			NB
Apu03	nd	nd	nd	nd	nd	nd	4.79×10^5	5.33×10^{-3}	11.10	8.27×10^4	0.0112	135.00
Apu04	nd	nd	nd	nd	nd	nd	5.38×10^5	5.89×10^{-3}	10.90	3.70×10^5	0.0232	62.70
Apu05	1.69×10^6	4.16×10^{-3}	2.46			NB	3.70×10^6	1.46×10^{-2}	3.94			NB
Apu06	1.04×10^6	6.58×10^{-3}	6.31			NB	2.60×10^6	2.20×10^{-2}	8.5			NB
Apu07	nd	nd	nd	nd	nd	nd	1.05×10^6	9.93×10^{-3}	9.49			NB
Apu08	9.04×10^5	8.71×10^{-3}	9.64			NB	1.12×10^6	0.0127	11.30			NB
Apu09	1.05×10^6	9.79×10^{-3}	9.36			NB	1.81×10^6	0.0169	9.32			NB
Apu 10	8.45×10^5	5.62×10^{-3}	6.65			NB	1.12×10^6	0.016	14			NB
Apu 11	nd	nd	nd	nd	nd	nd	9.04×10^5	0.0177	19.60			NB
Apu 12	5.73×10^5	7.07×10^{-3}	12.30			NB	3.13×10^5	0.0108	34.50			NB
Apu 13	nd	nd	nd	nd	nd	nd	1.15×10^5	0.016	139.00			NB
Apu 14	1.58×10^6	1.08×10^{-2}	6.85			NB	3.18×10^6	2.66×10^{-2}	8.35			NB
Apu 15	nd	nd	nd	nd	nd	nd	6.91×10^5	7.31×10^{-3}	10.60	1.28	0.0212	1650.00
Apu 16	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd
Apu 17	nd	nd	nd	nd	nd	nd			NB			NB
Apu 18	2.32×10^5	0.0169	72.80	1.09×10^6	8.17×10^{-3}	7.53	1.73×10^6	1.95×10^{-2}	11.30	1.69×10^6	1.75×10^{-2}	10.40
Apu18*							3.16×10^5	1.48×10^{-2}	46.90	1.01×10^6	1.54×10^{-2}	15.30
Apu 19			NB			NB			NB			NB
Apu 20	nd	nd	nd	nd	nd	nd			NB			NB
Apu21	nd	nd	nd	nd	nd	nd			NB			NB
Apu22	nd	nd	nd	1.53×10^5	0.0191	125.00	6.19×10^5	1.28×10^{-2}	20.70	nd	nd	nd
Apu23	nd	nd	nd	nd	nd	nd			NB			NB
Apu24	nd	nd	nd	nd	nd	nd			NB			NB

5 * A second Biacore analysis of Apu18 confirmed the previously obtained kinetic constants for the K63-linked diubiquitin interaction.

(D) Western Blot

Tetraubiquitin (K48-linked or K63-linked as appropriate) and diubiquitin (either K48-linked or K63-linked) (Boston Biochem) were separated in polyacrylamide gels and transferred by electroblotting to nitrocellulose membranes. Non-specific binding sites on the membranes were blocked by incubating the membranes overnight at 4°C in 0.5% Qiagen blocking reagent (Qiagen). The blocked membranes were placed in a miniblottedter apparatus. Fab clones (1 µg/ml) were applied to serial sections of the membrane in 0.5% Qiagen blocking reagent (Qiagen). After a one hour incubation period, the membranes were washed. Anti-ubiquitin antibodies bound to the membrane were revealed using HRP-conjugated anti-penta-histidine antibody (Qiagen) according to the manufacturer's instructions.

The K48-linked polyubiquitin-specific Fabs produced from clones apu01 to apu15 specifically bound to K48-linked tetraubiquitin immobilized to nitrocellulose (see Figure 19B). No binding of the K63-linked polyubiquitin-specific Fabs produced from clones apu17 to apu24 was observed to K63-linked polyubiquitin immobilized on nitrocellulose membranes (see Figure 19A).

EXAMPLE 2: ISOLATION AND CHARACTERIZATION OF SECOND GENERATION ANTI-POLYUBIQUITIN ANTIBODIES

Second generation libraries for Fab display were constructed from the phagemids encoding the previously identified clones apu05 (K48-linked polyubiquitin-selective) and apu18 (K63-linked polyubiquitin-selective) (see Figures 10 and 11). Phage from those clones were used to infect CJ236 cells to prepare Kunkel DNA templates. Those templates were subsequently mutagenized to insert stop codons, and the stop-containing templates were used in library construction as follows.

The Fab apu05 was mutagenized according to two different schemes to create two different apu05-derived libraries. In the first library, only HVR-H3 was mutagenized. HVR-H3 was first modified to include a stop codon in the Kunkel template, followed by a mutagenesis utilizing four mutagenic oligonucleotides. The stop codon-encoding oligonucleotide in all cases was CGTCTATTATTGTGCTCGCTAATAAGACTACTGGGGTCAAGG (SEQ ID NO: 365). The first three mutagenic oligonucleotides were three permutations of the same desired sequence, in which one tyrosine residue was fixed and each remaining tyrosine residue was randomized using the NNS mixed codon set (where N corresponds to G, C, A, or T and S corresponds to G or C); the amino acid at position 100b was selected from phenylalanine, methionine, leucine, and isoleucine; the amino acid at position 100a was selected from glycine and alanine; and the remaining amino acids were soft randomized. Soft randomization in this context indicates that certain nucleotide positions were 70% of the time occupied by the indicated base and 10% of the time occupied by one of the other three bases. For those oligonucleotides that follow, where such soft randomization was included at a particular base, the presence of the soft randomization is indicated by the presence of a number at that

base position. The number “5” indicates that the base adenine is present 70% of the time at that position, while the bases guanine, cytosine, and thymine are each present 10% of the time. Similarly, the number “6” refers to guanine, “7” to cytosine, and “8” to thymine, where in each case, each of the other three bases is present only 10% of the time. The first three mutagenic oligonucleotide sequences were: CGTCTATTATTGTGCTCGC567TAC567NNSNNS567GSTWTSGACTA
 5 CTGGGGTCAAGG (SEQ ID NO: 367), CGTCTATTATTGTGCTCGC567NNS567TACNNS567 GSTWTSGACTACTGGGGTCAAGG (SEQ ID NO: 368), and CGTCTATTATTGTGCTCGC567 NNS567NNSTAC567GSTWTSGACTACTGGGGTCAAGG (SEQ ID NO: 369). The fourth mutagenic oligonucleotide included randomization of the tyrosines at positions 96, 98, and 99 using
 10 the NNS mixed codon set; selection of the amino acid at position 100b from phenylalanine, methionine, leucine, and isoleucine; selection of the amino acid at position 100a from glycine and alanine, and soft randomization at every other position, in keeping with the soft randomization nomenclature described above. The sequence of the fourth oligonucleotide was
 15 CGTCTATTATTGTGCTCGC567NNS567NNSNNS567GSTWTSGACTACTGGGGTCAAGG (SEQ ID NO: 370).

In the second apu05 library, HVR-H1, HVR-H2, HVR-H3, and HVR-L3 were mutagenized. HVR-H1 was modified such that the serines at positions 30 and 33 were randomized using the NNS mixed codon set (where N corresponds to G, C, A, or T and S corresponds to G or C); the amino acid at position 29 was selected from amino acids phenylalanine, leucine, isoleucine, and valine; and the
 20 amino acid at position 34 was selected from isoleucine and methionine. The oligonucleotides used to mutagenize apu05 HVR-H1 were GCAGCTTCTGGCTTCAACTAATAACACTGGGTGCGTCAGG (SEQ ID NO: 371) and GCAGCTTCTGGCTTCAACNTCNNSTACTCTNNSATSCACTGGGTGC GTCAGG (SEQ ID NO: 372). HVR-H2 was modified such that the tyrosine at position 52 was randomized using the NNS mixed codon set and the amino acid at position 52a was selected from
 25 proline and serine. The oligonucleotides used to mutagenize HVR-H2 were GGCCTGGAATGGGTTGCATAATAATATGCCGATAGCGTCAAGG (SEQ ID NO: 373) and GGCCTGGAATGGGTTGCATCTATCNSYCTTACTACTCTTACACCTCTTATGCCGATAGCG TCAAGG (SEQ ID NO: 374). HVR-H3 was modified such that the tyrosine at position 99 and the serine at position 100 were randomized using the NNS mixed codon set; the amino acid at position
 30 100a was selected from glycine and alanine; and the amino acid at position 100b was selected from phenylalanine, methionine, leucine, and isoleucine. The oligonucleotides used to mutagenize HVR-H3 were CGTCTATTATTGTGCTCGCTAATAAGACTACTGGGGTCAAGG (SEQ ID NO: 365) and CGTCTATTATTGTGCTCGCTCTTACTCTTACNNSNNSGSTWTSGACTACTGGGGTCA AGG (SEQ ID NO: 375). HVR-L3 was modified such that position S91 was randomized according
 35 to the NNS mixed codon set and position I96 was selected from phenylalanine, isoleucine, and valine. The oligonucleotides used to mutagenize HVR-L3 were CGCAACTTATTACTGTCAGCAATAATAAACGTTCCGGACAGGGTACC (SEQ ID NO: 376) and

CGCAACTTATTACTGTCAGCAANNSTCTTACTCTTCTCTGDTTACGTTTCGGACAGGGTACC
(SEQ ID NO: 378).

Six different apu18-derived libraries were made by six different mutagenesis schemes of apu18. In the first library, only HVR-H3 was mutagenized. HVR-H3 was first modified to include a stop codon in the Kunkel template, followed by a mutagenesis methodology utilizing seven mutagenic oligonucleotides. The stop-codon-encoding oligonucleotide in all cases was CGTCTATTATTGTGCTCGCTAATAAGACTACTGGGGTCAAGG (SEQ ID NO: 366). The first six mutagenic oligonucleotides were six permutations of the same desired sequence, in which two tyrosine or tryptophan residues were fixed and each remaining tyrosine and tryptophan residue was randomized using the NNS mixed codon set (where N corresponds to G, C, A, or T and S corresponds to G or C); the amino acid at position 100c was selected from phenylalanine, methionine, leucine, and isoleucine; the amino acid at position 100b was selected from glycine and alanine; and the remaining amino acids were soft randomized in keeping with the soft randomization nomenclature described above. The first six mutagenic oligonucleotide sequences were:

CGTCTATTATTGTGCTCGC655TACTAC565NNSNNS577GSTWTSGACTACTGGGGTCAAGG
(SEQ ID NO: 379),

CGTCTATTATTGTGCTCGC655NNSNNS565TGGTAC577GSTWTSGACTACTGGGGTCAAGG
(SEQ ID NO: 380),

CGTCTATTATTGTGCTCGC655TACBBS565NNSTAC577GSTWTSGACTACTGGGGTCAAGG
(SEQ ID NO: 381),

CGTCTATTATTGTGCTCGC655NNSTAC565TGGNNS577GSTWTSGACTACTGGGGTCAAGG
(SEQ ID NO: 382),

CGTCTATTATTGTGCTCGC655TACNNS565TGGNNS577GSTWTSGACTACTGGGGTCAAGG
(SEQ ID NO: 383), and

CGTCTATTATTGTGCTCGC655NNSTAC565NNSTAC577GSTWTSGACTACTGGGGTCAAGG
(SEQ ID NO: 384). The seventh mutagenic oligonucleotide included randomization of the tyrosines

at positions 96, 97, and 100 and the tryptophan at position 99 using the NNS mixed codon set; selection of the amino acid at position 100b from phenylalanine, methionine, leucine, and isoleucine; selection of the amino acid at position 100c from phenylalanine, methionine, leucine, and isoleucine; selection of the amino acid at position 100b from glycine and alanine, and soft randomization at every other position, in keeping with the soft randomization nomenclature described above. The sequence of the seventh oligonucleotide was:

CGTCTATTATTGTGCTCGC655NNSNNS565NNSNNS577GSTWTSGACTACTGGGGTCAAGG
(SEQ ID NO: 385).

In the second apu18 library, only HVR-H2 was mutagenized. HVR-H2 was first modified to include a stop codon in the Kunkel template, followed by a mutagenesis methodology utilizing four mutagenic oligonucleotides. The stop codon-encoding oligonucleotide in all cases was

GGCCTGGAATGGGTTGCATAATAATATGCCGATAGCGTCAAGG (SEQ ID NO: 373). The first three mutagenic oligonucleotides were three permutations of the same desired sequence, in which one tyrosine residue was fixed and each remaining tyrosine and the serine at position 52 were randomized using the NNS mixed codon set (where N corresponds to G, C, A, or T and S corresponds to G or C); the amino acid at position 52a was selected from proline and serine; the amino acid at position 55 was selected from glycine and serine; the isoleucine and position 51 and the threonine at position 57 were fixed; and the remaining amino acids were soft randomized in keeping with the soft randomization nomenclature described above. The first three mutagenic oligonucleotide sequences were:

10 GGCCTGGAATGGGTTGCATACATCNNSYCTNNSNNSRGC567ACC567TATGCCGATAGCGT
CAAGG (SEQ ID NO: 386),
GGCCTGGAATGGGTTGCANNSATCNNSYCTTACNNSRGC567ACC567TATGCCGATAGCGT
CAAGG (SEQ ID NO: 387), and
GGCCTGGAATGGGTTGCANNSATCNNSYCTNNSTACRGC567ACC567TATGCCGATAGCGT
15 CAAGG (SEQ ID NO: 388). The fourth mutagenic oligonucleotide included randomization of the
tyrosines at positions 50, 53, and 54 using the NNS mixed codon set; selection of the amino acid at
position 52a from phenylalanine and serine; selection of the amino acid at position 55 from glycine
and serine; fixing the isoleucine and threonine residues at positions 51 and 57, respectively; and soft
randomization at every other position, in keeping with the soft randomization nomenclature described
20 above. The sequence of the fourth oligonucleotide was:
GGCCTGGAATGGGTTGCANNSATCNNSYCTNNSNNSRGC567ACC567TATGCCGATAGCGT
CAAGG (SEQ ID NO: 389).

In the third apu18 library, HVR-H2 and HVR-H3 were mutagenized. HVR-H2 was modified identically to the modifications made to HVR-H2 in the second apu18 library, using the same four
25 mutagenic oligonucleotides. HVR-H3 was modified identically to the modifications made to HVR-
H3 in the first apu18 library, using the same first six mutagenic oligonucleotides.

In the fourth apu18 library, HVR-H3 and HVR-L3 were mutagenized. HVR-H3 was modified identically to the modifications made to HVR-H3 in the first apu18 library, using the same first six mutagenic oligonucleotides. HVR-L3 was first modified to include a stop codon in the
30 Kunkel template, followed by a mutagenesis using a mutagenic oligonucleotide. Within HVR-L3, the
tyrosines at positions 91 and 94 and the serine at position 95a were randomized using the NNS mixed
codon set; the leucine at position 95b was selected from phenylalanine, isoleucine, and valine; and the
serines at positions 92, 93, and 95 were soft randomized in keeping with the soft randomization
nomenclature described above. The oligonucleotides used for the mutagenesis of HVR-L3 were
35 CGCAACTTATTACTGTCAGCAATAATAACGTTTCGGACAGGGTACC (SEQ ID NO: 376) and
CGCAACTTATTACTGTCAGCAANNS567567NNS567NNSCTGDTTACGTTTCGGACAGGGTA
CC (SEQ ID NO: 390).

In the fifth apu18 library, HVR-H1 and HVR-H2 were mutagenized. HVR-H2 was modified identically to the modifications made to HVR-H2 in the second apu18 library, using the same four mutagenic oligonucleotides. HVR-H1 was modified to include a stop codon; the serine at position 30 and the tyrosine at position 33 were randomized using the NNS mixed codon set; the amino acid at position 29 was selected from phenylalanine, leucine, isoleucine, and valine; the amino acid at position 34 was selected from isoleucine and methionine, and the amino acids at positions 31 and 32 were soft randomized in keeping with the soft randomization nomenclature described above. The oligonucleotides used for the mutagenesis of HVR-H1 were:

GCAGCTTCTGGCTTCAACTAATAACACTGGGTGCGTCAGG (SEQ ID NO: 371) and

GCAGCTTCTGGCTTCAACNTCNNS567567NNSATSCACTGGGTGCGTCAGG (SEQ ID NO: 391).

In the sixth apu18 library, HVR-H1, HVR-H2 and HVR-L3 were mutagenized. HVR-H1 was modified identically to the modifications made to HVR-H1 in the fifth apu18 library, using the same mutagenic oligonucleotide. HVR-H2 was modified identically to the modifications made to HVR-H2 in the second apu18 library, using the same four mutagenic oligonucleotides. HVR-L3 was modified identically to the modifications made to HVR-L3 in the fourth apu18 library, using the same mutagenic oligonucleotide.

Mutagenesis reactions for each of the two apu5-derived libraries and each of the six apu18-derived libraries were transformed into electrocompetent *E. coli* XL-1 by electroporation. Cells were allowed to recover for 30 minutes at 37 °C with agitation in SOC medium. Twenty microliters of the cell-containing SOC medium was reserved to determine the number of transformants, and the remainder was then transferred to 500 ml 2YT containing carbenicillin and 10^{10} M13K07 helper phage per milliliter. After 45 minutes at 37 °C with agitation, the broth was supplemented with kanamycin and grown overnight at 37 °C with agitation. The number of transformants for each library was $>10^9$. Phage were harvested and concentrated from the broth by centrifugation and PEG precipitation, and subsequently used in rounds of selection.

K48-linked polyubiquitin and K63-linked polyubiquitin were immobilized on different Maxisorp plates (NUNC) as described above in Example 1(A). Each library was sorted separately against its respective target (K48-linked polyubiquitin for the two apu05-derived libraries, K63-linked polyubiquitin for the six apu18-derived libraries) for one round with the addition of 3 μ M monoubiquitin in the sorting buffer. The eluted phage were amplified and pooled (two pools, one for each lysine-linked polyubiquitin target) for further rounds of sorting.

Subsequent selection rounds were solution-phase sorted. Phage pools were incubated with biotinylated (Sulfo-NHS-biotin, Pierce) polyubiquitin chains for one to two hours at room temperature in solution-sorting buffer (PBST with 0.5% Superblock (Pierce)). The mixture was diluted five- to ten-fold in solution-sorting buffer and added to neutravidin-coated wells for a brief (5 minute) capture of biotinylated polyubiquitin. A reaction containing unbiotinylated polyubiquitin chains served as a

control to monitor background phage binding. The plates were washed with PBST and eluted with 0.1 M HCl for 10 minutes.

Stringency was modulated in three ways: by the concentration of biotinylated polyubiquitin; by the addition of excess unbiotinylated polyubiquitin to compete for binding before capture on neutravidin-coated wells; and by the duration of the competition. For each round of sorting, monoubiquitin and polyubiquitin of the other linkage was added at a concentration of 30 µg/ml to the sorting buffer during the first incubation step. The first round of solution sorting employed 20 nM biotinylated polyubiquitin incubated with phage for one hour at room temperature. The mixture was then diluted tenfold in solution-sorting buffer, and captured using neutravidin-coated wells for five minutes. For the second round phage was equilibrated with 20 nM biotinylated polyubiquitin, as in round 1, but was diluted tenfold in solution-sorting buffer containing 30 µg/ml of unbiotinylated polyubiquitin (K48-linked for K48 selection, K63-linked for K63 selection) for fifteen minutes of off-rate selection followed by capture on neutravidin coated wells. The third round of solution sorting was as described for the second round, but further included 5 nM biotinylated polyubiquitin and 30 minutes of off-rate selection. After one round of plate sorting and three rounds of solution sorting, individual clones selected from the second generation were grown in a 96 well format as described. Individual clones were screened by phage ELISA and sequenced.

After the first round of solution sorting, up to 40 times enrichment was observed for the libraries based on apu05 and up to 7-fold enrichment for the libraries based on apu18 (see Table C). An additional 11-fold enrichment was obtained for the K48-specific clones and an additional 3-fold enrichment for the K63-specific clones after the second selection (solution sort for slow off rate) (see Table C). The third selection (both affinity sorting and off-rate sorting) resulted in an 18-fold enrichment for the K48-specific clones and a four-fold enrichment for the K63-specific clones (see Table C).

Table C: Results from Second Generation Anti-Polyubiquitin Antibody Library Solution Sorting

First Selection (solution sort for affinity)						
	5 nM		10 nM		20 nM	
	Size	Enrichment	Size	Enrichment	Size	Enrichment
Based on apu05	2.00x10 ⁴	20.00	3.00x10 ⁴	30.00	4.00x10 ⁴	40
Based on apu18	2.20x10 ³	1.83	2.00x10 ³	1.67	8.40x10 ³	7

Second Selection (solution sort for slow off rate)						
	5 min		10 min		15 min	
	Size	Enrichment	Size	Enrichment	Size	Enrichment
Based on apu05	4.00×10^4	17.78	4.00×10^4	17.78	2.50×10^4	11.11
Based on apu18	5.50×10^3	4.07	7.00×10^3	5.19	4.50×10^3	3.33
Third Selection (solution sort for both affinity and slow off rate)						
	5 nM and 30 min					
	Size	Enrichment				
Based on apu05	4.70×10^3	18.60				
Based on apu18	1.00×10^2	4.25				

Sixty-eight unique clones were identified which bound specifically to K48-linked polyubiquitin; those clones were subjected to DNA sequence analysis. The HVR-H1, HVR-H2, and HVR-H3 sequences of those clones are shown in Figures 14A-F. Thirty-one unique clones were identified which bound specifically to K63-linked polyubiquitin; those clones were also subjected to sequence analysis. The HVR-H1, HVR-H2, and HVR-H3 sequences of those clones are shown in Figures 15A-C. The light chain HVR for each of the K48-linked polyubiquitin and K63-linked polyubiquitin-specific clones were not sequenced, but, as with apu05 and apu18, the sequences of HVR-L1 and HVR-L2 were expected to be invariant, while the HVR-L3 sequence was expected to be clone-specific. The HVR-L1 sequence is RASQSVSSAVA (SEQ ID NO: 79) and the HVR-L2 sequence is SASSLYS (SEQ ID NO: 80), according to the library design. All clones had the same heavy chain and light chain framework sequences (see Figure 6).

The twenty clones with the greatest observed binding (ten K48-linked polyubiquitin-specific and ten K63-linked polyubiquitin-specific) were produced as Fabs as described in Example 1. Fabs apu2.01 through apu2.20 were subjected to DNA sequence analysis. The HVR-H1, HVR-H2, HVR-H3, and HVR-L3 sequences of those clones are shown in Figures 16A and B (K48-specific Fabs) and 17A and B (K63-specific Fabs).

Fabs apu2.01 through apu2.20 were included in an ELISA protein assay to determine their relative affinities for K48-linked and K63-linked polyubiquitin, and to confirm that the Fabs was not reactive with monoubiquitin or BSA (see Figure 18). Apu clones 2.11 and 2.12 each demonstrated about 300 times greater affinity for K63-linked polyubiquitin than for K48-linked polyubiquitin in that assay. Apu2.20 and 2.16 had lesser, but still marked differences (about 30 times greater and about 10 times greater, respectively) in affinity for K63-linked polyubiquitin as compared to K48-

linked polyubiquitin. No binding of clones apu2.01-apu2.10 to K63-linked poly- or diubiquitin was detected.

Each Fab was also analyzed by BIAcore as described previously in Example 1(C). The obtained kinetic constants and binding constants are shown in Table D. The language “NB” in Table

5 D indicates that no binding was detected for the indicated interaction.

TABLE D: Kinetic Constants of Anti-Polyubiquitin Fabs as Measured by BIAcore Analysis

	K63-linked polyubiquitin			K48-linked diubiquitin			K63-linked diubiquitin		
	k_{on} (1/M·s)	k_{off} (1/s)	K_d (nM)	k_{on} (1/M·s)	k_{off} (1/s)	K_d (nM)	k_{on} (1/M·s)	k_{off} (1/s)	K_d (nM)
Apu2.01			NB	6.33×10^5	1.49×10^{-2}	23.5			NB
Apu2.02			NB	1.11×10^6	1.01×10^{-2}	9.16			NB
Apu2.03			NB	8.67×10^5	2.66×10^{-3}	3.07			NB
Apu2.04			NB	6.22×10^5	9.39×10^{-3}	15.10			NB
Apu2.05			NB	1.20×10^6	6.09×10^{-3}	5.06			NB
Apu2.06			NB	1.28×10^6	2.40×10^{-3}	1.87			NB
Apu2.07			NB	2.70×10^6	3.18×10^{-3}	1.18			NB
Apu2.08			NB	1.25×10^6	7.21×10^{-3}	5.76			NB
Apu2.09			NB	7.15×10^5	4.78×10^{-3}	6.69			NB
Apu2.10			NB	3.28×10^6	3.99×10^{-3}	1.21			NB
Apu05			NB	2.44×10^6	1.07×10^{-2}	4.37			NB
Apu2.11	2.49×10^5	7.11×10^{-3}	28.6	1.68×10^4	1.24×10^{-2}	738	3.12×10^5	1.02×10^{-2}	32.7
Apu2.12	4.76×10^5	9.68×10^{-3}	20.3	5.66×10^3	7.88×10^{-3}	1390	5.49×10^5	1.51×10^{-2}	27.5
Apu2.13	4.89×10^5	2.41×10^{-3}	4.93	2.08×10^5	2.15×10^{-2}	103	5.83×10^5	4.11×10^{-3}	7.05
Apu2.14	2.09×10^5	7.08×10^{-3}	34.0	4.97×10^3	1.15×10^{-2}	2310	2.14×10^5	1.06×10^{-2}	49.6
Apu2.15	9.07×10^5	9.98×10^{-3}	11.0	1.43×10^4	1.23×10^{-2}	856	1.10×10^6	1.55×10^{-2}	14.0
Apu2.16	1.68×10^5	1.80×10^{-2}	107	4.11×10^1	1.73×10^{-3}	42000	1.38×10^5	1.27×10^{-2}	92.1
Apu2.17	1.19×10^6	1.35×10^{-2}	11.3	6.90×10^3	2.73×10^{-3}	396	1.13×10^6	2.89×10^{-2}	25.6
Apu2.18	1.48×10^5	1.89×10^{-2}	128	4.98×10^3	5.36×10^{-3}	1080	2.66×10^5	2.53×10^{-2}	95.0
Apu2.19	8.97×10^5	8.47×10^{-3}	9.44	2.54×10^5	1.24×10^{-2}	48.9	8.76×10^5	1.41×10^{-2}	16.1
Apu2.20	2.93×10^5	1.77×10^{-2}	60.7	4.02×10^3	3.77×10^{-3}	939	4.19×10^5	2.50×10^{-2}	59.7
Apu18	1.09×10^6	8.17×10^{-3}	7.53	3.16×10^5	1.48×10^{-2}	46.9	1.01×10^6	1.54×10^{-2}	15.3

Several of the Fabs based on apu05 had K_d s lower than that of the Fab corresponding to apu05 for

10 K48-linked diubiquitin, representing tighter binding to polyubiquitin. Each of the apu2.11-2.20 Fabs bound not only to K63-linked polyubiquitin but also, to a lesser extent, to K48-linked diubiquitin.

While only apu2.13 had a lower K_d than apu18, its K_d for K48-linked diubiquitin was larger than that of apu18. Each of apu 2.11, 2.12, 2.16, and 2.20 had better ratios of K_d for K63-linked polyubiquitin to K_d for K48-linked diubiquitin than apu18. The observed kinetic constants for binding of the

apu18-based Fabs to K63-linked polyubiquitin were similar to those observed for binding to K63-linked diubiquitin.

The ability of each Fab to specifically bind to polyubiquitin immobilized to a nitrocellulose membrane was also assessed by Western blot as previously described in Example 1(D).

5 Tetraubiquitin containing either a K48 or K63 linkage, diubiquitin containing the opposite lysine linkage from that of the tetraubiquitin (e.g., K63-linked diubiquitin when K48-linked tetraubiquitin was used, or K48-linked diubiquitin when K63-linked tetraubiquitin was used), and monoubiquitin were immobilized to nitrocellulose membranes, and Fabs apu2.01-2.20 were assessed for their abilities to recognize all three immobilized molecules (Figures 20A and 20B). None of the Fabs specifically
10 recognized monoubiquitin. Apu2.01-apu2.10 each specifically recognized immobilized K48-linked tetraubiquitin but did not recognize immobilized K63-linked diubiquitin (see Figure 20A). Several other bands appeared on the blot which represent contaminant tri-, penta-, and octaubiquitin species in the K48-linked tetraubiquitin preparation. Apu2.11-apu2.20 specifically recognized immobilized K63-linked tetraubiquitin and did not recognize immobilized K48-linked diubiquitin (see Figure
15 20B).

EXAMPLE 3: BINDING OF ANTI-POLYUBIQUITIN ANTIBODIES TO ENDOGENOUSLY POLYUBIQUITINATED PROTEINS

Previous experiments had demonstrated that the activity of Receptor Interacting Protein (RIP), a 140 kD essential mediator of the proximal TNF receptor 1 (TNFR1) signaling complex, is
20 modulated by polyubiquitination (Wertz et al., Nature 430: 694-699 (2004)). When RIP is polyubiquitinated with K63-linked polyubiquitin chains, signaling through TNFR1 is facilitated. Removal of the K63-linked polyubiquitin chains from RIP by the de-ubiquitinating N-terminus of A20 and replacement with K48-linked polyubiquitin chains by the ubiquitin ligase function of the A20 C terminus inactivates RIP and targets it for proteasomal degradation. That mechanism had been
25 elucidated using cell lines expressing mutant ubiquitin capable of forming only K48-linked or K63-linked polyubiquitin.

The ability of two of the anti-K48 and anti-K63-linked polyubiquitin binding proteins of the invention to specifically recognize the differently polyubiquitinated forms of RIP that exist in HeLa S3 cells at different time points after TNF treatment was assessed. Four liters of HeLa S3 cells at
30 approximately 1.5×10^6 cells/mL were treated with 21 μ M MG-132. Immediately after treatment, one liter of cells was removed from the main culture, harvested by centrifugation, washed with 200 mL PBS, and recentrifuged. This sample was used as the time zero time point. The remaining three liters of cell culture were treated with 100 ng/mL TNF. One liter of cells were removed, harvested, washed with 200 mL PBS, and recentrifuged at each of 5, 15, and 25 minutes after treatment with
35 TNF. The cells from each time point were lysed in 30 mL of IP lysis buffer (LB) (20 mM Tris pH 7.5, 150 mM NaCl, 1% Triton™ x-100, 1 mM EDTA, 25 μ M MG-132, 10 mM NEM, 30 μ L of each of phosphatase inhibitor cocktails 1 and 2 (Sigma), and 1 Complete protease inhibitor tablet (Roche))

and incubated with rotation at 4° C for 20-60 minutes. Each lysate was transferred to a 50 mL centrifuge tube and pelleted twice for five minutes at 10,000 x g. The protein concentration of the lysate from each time point was estimated. Each lysate (30 mL for each time point with normalized protein concentrations) was incubated for 1.25-1.5 hours with 1 mL unblocked Protein A/G beads at 4° C. The beads and debris were separated from the lysate by centrifugation at 2000 rpm for 5 minutes. A sample was taken from each lysate for direct Western blot analysis, and the remaining volume was frozen at -80°C.

Four 16 mL samples of each lysate were taken. To each sample was added 25 mM MG-132 and 20 µL NEM. Two samples were each combined with 2.4 µg/mL of an anti-TNFR1 antibody.

Two other samples were each combined with 2.4 µg/mL of a control antibody (anti-myc). All samples were incubated for two hours at 4° C with rotation, followed by the addition of 150 µL of a 50% slurry of unblocked protein A beads. The samples were incubated at 4° C for an additional 5 hours with rotation. The samples were pelleted by centrifugation, and the beads were washed once with 15 mL LB, twice with 10 mL LB containing 1 M NaCl, and twice with 10 mL LB. The washed beads were resuspended in 1.25 mL LB and transferred to microfuge tubes. Each sample was aspirated such that the tube contained a total volume of 950 µL, and 360 mg solid urea was added to each sample for a final concentration in each sample of 6M in urea. The samples were incubated for 15 minutes at room temperature with gentle agitation. The beads in each sample were pelleted by centrifugation. A portion of each supernatant was reserved for Western blot analysis, and the remaining supernatant from each sample (approximately 1 mL per sample) was diluted to 10 mL with dissociation dilution buffer (1% Triton-X100, 0.5% deoxycholate, 120 mM NaCl, 50 mM HEPES pH 7.2, and Complete protease inhibitor cocktail (Roche)).

Each sample was split into two five mL portions. One portion was treated with 2.5 µg apu2.16 that had been reformatted from the fab form to the IgG form, and the other was treated with 2.5 µg apu2.07 that had been reformatted from the fab form to the IgG form. To both samples were added 50 µL protein A beads, and the samples were incubated for 5 hours at 4° C. The protein A beads were pelleted and washed three times in TNFR1 LB, having been transferred to microfuge tubes during the washing process. Sample buffer was added to all samples, and each sample (including the samples previously reserved for Western blot analysis) was reduced and run on 10% tris/gly 1.5 mm 10-well Novex® gels (Invitrogen). After running, the proteins in the gels were transferred to Invitrolon™ PVDF membranes (Invitrogen) according to the manufacturer's instructions. The membranes were blocked with 5% PBS-T and probed with an anti-RIP monoclonal antibody (Becton Dickinson) overnight at room temperature. The blots were then washed, probed with HRP-conjugated goat anti-mouse secondary antibody (Cappel), washed, exposed to the reagent to activate chemiluminescence, and exposed to film. The results are shown in Figures 21A and 21B. Figure 21A depicts the blot containing the samples first immunoprecipitated with TNFR1 or anti-myc and then immunoprecipitated with the apu2.16 IgG (K63-linked polyubiquitin selective). As shown

in Figure 21A, RIP is not visible in either the anti-myc control samples or in the zero time sample. The RIP band was strongest in the 5 minute sample, and then decreased significantly in the 15 and 25 minute samples. Figure 21B depicts the blot containing the samples first immunoprecipitated with TNFR1 or anti-myc and then immunoprecipitated with the apu2.07 IgG (K48-linked polyubiquitin selective). As shown in Figure 21B, no RIP was observed in the anti-myc control lanes. RIP levels in this blot increased over time, and were strongest in the 25 minute time point sample. This data correlates with the earlier findings that upon signaling through the TNFR1, RIP is first K63-linked polyubiquitinated, and then subsequently deubiquitinated and K48-linked polyubiquitinated by A20. Thus, the antibodies of the invention were able to specifically bind to and discriminate between a K63-linked polyubiquitinated polypeptide and a K48-linked polyubiquitinated polypeptide that had been polyubiquitinated in cells.

EXAMPLE 4: ISOLATION AND CHARACTERIZATION OF THIRD GENERATION ANTI-POLYUBIQUITIN ANTIBODIES

Third generation libraries for Fab display were constructed from the phagemid encoding the previously identified clone apu2.16 (K63-linked polyubiquitin-selective) (see Example 2 and Figures 17A and 17B). Phage from that clone was used to infect CJ236 cells to prepare a Kunkel DNA template. That template was subsequently mutagenized to insert stop codons, and the stop-containing templates were used in library construction as follows.

The Fab apu2.16 was mutagenized according to three different schemes to create three different apu2.16-derived libraries. In the first library, HVR-H1 and HVR-H2 were mutagenized. HVR-H1 was mutagenized to include a stop codon in the Kunkel template, followed by a mutagenesis methodology utilizing one mutagenic oligonucleotide. The stop codon-encoding oligonucleotide sequence was: GCAGCTTCTGGCTTCAACTAATAACACTGGGTGCGTCAGG (SEQ ID NO: 371). The mutagenic oligonucleotide permitted the isoleucine at position 29 to be selected from phenylalanine, leucine, isoleucine, and valine; the isoleucine at position 34 to be selected from methionine and isoleucine using the NNS mixed codon set (where N corresponds to G, C, A, or T and S corresponds to G or C); and the soft-randomization of amino acids K30, T31, G32, and L33. Soft randomization in this context indicated that certain nucleotide positions were 70% of the time occupied by the indicated base and 10% of the time occupied by one of the other three bases. For those oligonucleotides that follow, where such soft randomization was included at a particular base, the presence of the soft randomization is indicated by the presence of a number at that base position. The number "5" indicates that the base adenine was present 70% of the time at that position, while the bases guanine, cytosine, and thymine were each present 10% of the time. Similarly, the number "6" refers to guanine, "7" to cytosine, and "8" to thymine, wherein in each case, each of the other three bases was present only 10% of the time. The oligonucleotide sequence used to mutagenize apu3.16

HVR-H1 was: GCAGCTTCTGGCTTCAACNTC556577668788ATSCACTGGGTGCGTCAGG (SEQ ID NO: 785).

HVR-H2 was also modified to include a stop codon in the Kunkel template, followed by a mutagenesis utilizing three mutagenic oligonucleotides. The stop codon-encoding oligonucleotide in all cases was GGCCTGGAATGGGTTGCATAATAATATGCCGATAGCGTCAAGG (SEQ ID NO: 373). The three mutagenic oligonucleotides were two permutations of a first desired sequence, and one second desired sequence. The first desired sequence included: one fixed and one randomized tyrosine residue at either of positions 50 and 54 using the NNS mixed codon set (described above); fixed residues at positions 51 (isoleucine), 52a (proline), 53 (tyrosine), 55 (glycine), and 57 (threonine); and soft randomization of positions S52, S56, and S58 (according to the soft randomization scheme described above). The second desired sequence included: fixed residues at positions 51 (isoleucine), 52a (proline), 53 (tyrosine), 55 (glycine), and 57 (threonine); hard randomization of the tyrosines at positions 50 and 54 using the NNS mixed codon set (described above), and soft randomization of positions S52, S56, and S58 (according to the soft randomization scheme described above.) The oligonucleotides used to mutagenize apu2.16 HVR-H2 were GGCCTGGAATGGGTTGCANNSATC567CCGTACTACGGT567ACC567TATGCCGATAGCGTCAAGG (SEQ ID NO: 786), GGCCTGGAATGGGTTGCATACATC567CCGTACNNSGGT567ACC567TATGCCGATAGCGTCAAGG (SEQ ID NO: 787), and GGCCTGGAATGGGTTGCANNSATC567CCGTACNNSGGT567ACC567TATGCCGATAGCGTCAAGG (SEQ ID NO: 788).

In the second apu2.16 library, HVR-H2 and HVR-H3 were mutagenized. HVR-H2 was mutagenized identically to the modifications made to HVR-H2 in the first apu2.16 library, using the same stop codon-containing oligonucleotide and same three mutagenic oligonucleotides. HVR-H3 was mutagenized identically to the modifications made to HVR-H3 in the first apu18 library (described in Example 2), using the same stop-codon-containing oligonucleotide and same six mutagenic oligonucleotides.

In the third apu2.16 library, HVR-H1 and HVR-H3 were mutagenized. HVR-H1 was mutagenized identically to the modifications made to HVR-H1 in the first apu2.16 library, using the same stop-codon-containing oligonucleotide and mutagenic oligonucleotide. HVR-H3 was mutagenized identically to the modifications made to HVR-H3 in the first apu18 library (described in Example 2), using the same stop-codon-containing oligonucleotide and same six mutagenic oligonucleotides.

Mutagenesis reactions for each of the three apu2.16-derived libraries were transformed into electrocompetent *E. coli* XL-1 by electroporation. Cells were allowed to recover for 30 minutes at 37 °C with agitation in SOC medium. Twenty microliters of the cell-containing SOC medium was reserved to determine the number of transformants, and the remainder was then transferred to 500 ml 2YT containing carbenicillin and 10^{10} M13K07 helper phage per milliliter. After 45 minutes incubation at 37 °C with agitation, the broth was supplemented with kanamycin and grown overnight

at 37 °C with agitation. The number of transformants for each library was $>10^9$. Phage were harvested and concentrated from the broth by centrifugation and PEG precipitation, and used in subsequent rounds of selection.

The three third-generation libraries were each sorted separately against K63-linked polyubiquitin immobilized on Maxisorp plates (NUNC) as described above in Example 1(A). Stringency was modulated in three ways: by the concentration of biotinylated polyubiquitin; by the addition of excess unbiotinylated polyubiquitin to compete for binding before capture on neutravidin-coated wells; and by the duration of the competition. Each round of solution sorting included 3 μ M monoubiquitin and 30 μ g/mL K48-linked polyubiquitin in the sorting buffer during the incubation step. The eluted phage were amplified and pooled for further rounds of sorting. Subsequent selection rounds were solution-phase sorted. The first round of solution sorting included 100 nM biotinylated (Sulfo-NHS-biotin, Pierce) polyubiquitin incubated with the phage pools for one hour at room temperature. The mixture was then diluted tenfold in solution-sorting buffer (PBST with 0.5% Superblock™ (Pierce), and captured using neutravidin-coated wells for five minutes. A reaction containing unbiotinylated polyubiquitin chains served as a control to monitor background phage binding. The plates were washed with PBST and eluted with 0.1 M HCl for 10 minutes. For the second solution-sorted round, phage was equilibrated with 30 nM biotinylated polyubiquitin, as in round one, but was diluted tenfold in solution-sorting buffer containing 30 μ g/mL of unbiotinylated polyubiquitin (K63-linked) for five minutes of off-rate selection followed by capture on neutravidin-coated wells.

After the first round of solution sorting, 6.5 times enrichment was observed for the combined libraries based on apu2.16 (see Table E). An additional 10-fold enrichment was obtained after the second solution sort for slow off rate (see Table E).

Table E: Results from Third Generation Anti-Polyubiquitin Antibody Library Solution Sorting

	Plate Sort		Solution Sort			
	Round 1		Round 2 (100 nM)		Round 3 (30 nM)	
Library	Library size	Enrichment	Library size	Enrichment	Library size	Enrichment
16-1	1.06e+06	ND	1.04e+05	6.5	1.00e+05	10
16-2	5.98e+05	ND				
16-3	1.76e+05	ND				

25

After one round of plate sorting and two rounds of solution sorting, individual clones selected from the third generation were grown in a 96-well format and screened by phage ELISA as described above in Example 2. Seventy-two unique clones were identified by sequencing. Of those clones, twelve demonstrated the greatest degree of specificity for K63-linked polyubiquitin in the phage ELISA assay (Figure 22). Those twelve were designated apu3.01-3.12, and their HVR-H1, HVR-H2,

30

and HVR-H3 sequences appear in Figures 23A and 23B. The light chain HVR for each of the K63-linked polyubiquitin-specific clones were not sequenced, but the sequences of HVR-L1 and HVR-L2 were expected to be invariant, while the HVR-L3 sequence was expected to be identical to that for apu2.16. The HVR-L1 sequence was RASQSVSSAVA (SEQ ID NO: 79) and the HVR-L2 sequence was SASSLYS (SEQ ID NO: 80), according to the library design. All clones had the same heavy chain and light chain framework sequences (see Figure 6).

Apu2.07 (see Example 2 and Figures 16A and 16B)) and apu3.07 (see above and Figures 23A and 23B) were expressed in either CHO or 293 cells as human IgGs. Expression constructs were made by Kunkel mutagenesis of appropriate pRK mammalian expression vectors encoding the heavy and light chains of human IgG (Gorman et al., DNA Prot. Eng. Tech. 2:3-10(1990)). IgGs were purified by affinity chromatography using standard methodologies.

The ability of each IgG to specifically bind to polyubiquitin of the appropriate linkage immobilized to a nitrocellulose membrane was assessed by Western blot. K48- or K63-linked polyubiquitins and monoubiquitin (Boston Biochem) were run 4-20% Tris-glycine polyacrylamide gels (Invitrogen). The contents of the gels were transferred to nitrocellulose by electroblotting. Non-specific binding sites on the nitrocellulose membrane were blocked for 1 hour in 5% non-fat skim milk powder dissolved in Tris-buffered saline containing 0.1% Tween-20 (TBST). The K48-specific or K63-specific antibodies were then added to the blot at a concentration of 2 μ g/mL (apu 2.07 IgG) or 1 μ g/mL (apu3.07 IgG) and incubated for one hour to allow binding to occur. As a positive control, one blot was incubated with rabbit anti-ubiquitin antibodies (Sigma). Blots were washed in TBST and bound antibodies detected by peroxidase-conjugated goat anti-human Ig Fc (ICN) or peroxidase-conjugated anti-rabbit Ig (Amersham) diluted 1:10,000 in TBST containing 5% non-fat milk powder. After one hour, blots were washed in TBST and developed using Super Signal West Dura reagent (Pierce) to reveal peroxidase activity. The results are shown in Figures 24A-24D.

As expected, the apu2.07 IgG specifically recognized immobilized K48-linked tetraubiquitin and K48-linked tri- to heptaubiquitin (Figure 24A), but did not bind to any of the immobilized K63-linked polyubiquitin samples. Similarly, the apu3.07 IgG specifically recognized immobilized K63-linked tetraubiquitin and K63-linked tri- to heptaubiquitin (Figure 24B), but did not bind to any of the immobilized K48-linked polyubiquitin samples. Neither IgG bound to immobilized monoubiquitin. To assess the sensitivity of each IgG, additional western blot analysis was performed with varied concentrations of immobilized K48-linked and K63-linked tetraubiquitin (25-1000 ng/lane) (Figures 24C and 24D). Apu2.07 IgG detected as little as 50 ng immobilized K48-linked tetraubiquitin, and again did not specifically bind to immobilized K63-linked tetraubiquitin (Figure 24C). Apu3.07 IgG detected as little as 50 ng immobilized K63-linked tetraubiquitin, and again did not specifically bind to immobilized K48-linked tetraubiquitin (Figure 24D). In both cases, increased amounts of immobilized tetraubiquitin resulted in increased observed binding.

To determine whether the IgG were able to detect endogenously polyubiquitinated proteins, protein lysates were prepared from the human embryonic kidney cell line 293T treated with or without 20 μ M of the proteasome inhibitor Velcade® (bortezomib) for four hours. Lysates were resolved by SDS-PAGE in a 4-20% Tris-glycine polyacrylamide gel (Invitrogen), and western blotting was performed as described above. The results are shown in Figure 25. A polyclonal anti-ubiquitin antibody (Sigma) detected a large number of high molecular weight proteins (leftmost lanes) both in the presence and absence of Velcade® treatment. The apu2.07 IgG bound to numerous proteins of varied molecular weights (rightmost lanes), and the observed binding was heavier to the immobilized velcade-treated lysates than to the immobilized untreated lysates. Significantly less binding overall was observed with the apu3.07 IgG (center lanes), which was expected to bind to K63-polyubiquitinated proteins, and there was no apparent difference in binding between the velcade-treated and untreated lanes. K48-linked polyubiquitinylation is known to target intracellular proteins for proteolytic degradation (Chau et al., Science 243: 1576-1583 (1989); Finley et al., Mol. Cell. Biol. 14: 5501-5509 (1994); Flick et al., Nat. Cell. Biol. 6:634-641 (2004)). Thus, one explanation for the apu2.07 IgG results was that when proteolytic processing was prevented, an increased amount of K48-polyubiquitinated proteins remained in the lysate, resulting in increased binding by the apu2.07 IgG over the untreated sample. K63-linked polyubiquitinylation is not known to target proteins for degradation (Pickart and Fushman, Curr. Opin. Chem. Biol. 8: 610-616 (2004); Hicke and Dunn, Annu. Rev. Cell Dev. Biol. 19: 141-172 (2003); Spece et al., Mol. Cell Biol. 15: 1265-1273 (1995); Ulrich, Eukaryot. Cell 1: 1-10 (2002); Spence et al., Cell 102: 67-76 (2000); Seibenhener et al., Mol. Cell. Biol. 24(18): 8055-8068 (2004)). Thus, one explanation for the apu3.07 IgG results was that inhibition of proteasomes did not result in an accumulation of K63-polyubiquitinated proteins.

EXAMPLE 5: STRUCTURAL ANALYSIS OF FAB BINDING TO ANTI-K63-LINKED POLYUBIQUITIN

To better understand the interaction of the anti-K63-linked polyubiquitin fab with polyubiquitin, the anti-K63-linked polyubiquitin fab apu2.16 was co-crystallized with K63-linked diubiquitin. Crystals were grown in hanging drops using 1 μ L of an apu2.16 solution (15 mg/mL in 10 mM Tris, 75mM NaCl pH 8.0) and 1 μ L of well solution (0.1M LiCl, 0.1 M Tris pH 8.2, 1 M citrate). To each drop was added 0.5 μ L of 0.1 M cupric chloride, and each drop was streak seeded. Crystal clusters grew over the course of several days and could be manipulated to obtain a single, diffracting crystal. The structure was determined by molecular replacement. Native data were collected at 100K and were processed with HKL2000. Crystals belonged to the C2 space group with cell dimensions $a = 177.7 \text{ \AA}$, $b = 94.9 \text{ \AA}$, $c = 97.9 \text{ \AA}$ and $\beta = 107^\circ$, with two complexes in the asymmetric unit. The structure was solved by molecular replacement using the program Phaser and the coordinates of a variant of the humanized 4d5 fab fragment (PDB for 4d5: PDB code 1FVE). Model building was carried out in program Coot and the structure was refined with Refmac5. The

resolution of the structure is 3.1 Å. The complex has been refined to an R of 24.5% and an R_{free} of 30.4%.

The interaction between apu2.16 and K63-linked diubiquitin is shown in Figures 26A-26C. The structural epitope is a combination of residues which bury at least 25% of their solvent accessible surface area upon binding fab and/or have more than one atom within 4.5 Å of either the heavy or light chain of the fab. The ubiquitin chain which donates K63 is chain A, and the ubiquitin chain which donates the C-terminus is chain B. Fab light chain residues belong to chain L, and fab heavy chain residues belong to chain H. Chain number precedes residue number in the table below, and fab residues are numbered sequentially.

TABLE F: Residues located in the apu2.16-K63-linked diubiquitin binding interface

Ubiquitin Residues	Fab Residues
A 18 Glu	
A 19 Pro	L 31 Ser
A 20 Ser	L 49 Tyr
A 21 Asp	L 50 Ser
A 55 Thr	L 51 Ala
A 56 Leu	L 52 Ser
A 57 Ser	L 53 Ser
A 58 Asp	L 66 Arg
A 60 Asn	L 98 Phe
A 61 Ile	
A 62 Gln	H 30 Lys
	H 31 Thr
B 8 Leu	H 32 Gly
B 9 Thr	H 33 Leu
B 34 Glu	H 50 Tyr
B 35 Gly	H 52 Ser
B 36 Ile	H 54 Tyr
B 37 Pro	H 55 Tyr
B 39 Asp	H 99 Glu
B 40 Gln	H 100 Tyr
B 71 Leu	H 101 Tyr
B 72 Arg	H 102 Arg
B 73 Leu	H 104 Tyr
B 74 Arg	H 105 Thr
B 75 Gly	

As shown in Table F, and as indicated in Figure 26B in dark grey, there were eleven residues in the K63-diubiquitin A chain and thirteen residues in the K63-diubiquitin B chain that were located within 4.5 Å of apu2.16 when bound to apu2.16. As shown in Table F, and as indicated in Figure 26C in dark grey, there were eight residues in the apu2.16 light chain and fourteen residues in the apu2.16 heavy chain that were located within 3.5 Å of K63-linked diubiquitin when bound to that molecule. Based on this data, residues likely to mediate the interaction between the two on K63-linked diubiquitin include Glu-18, Ser-20, Leu-57, and Asp-58 in the A chain and Pro-37, Arg-74, and Gly-

75 in the B chain. Interestingly, the antibody does not interact intimately with the K63-Gly 76 linkage but instead derives specificity via interactions the surface of the di-ubiquitin complex adjoining the linkage.

SEQUENCE LISTING IN ELECTRONIC FORM

This description contains a sequence listing in electronic form in ASCII text format. A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office. The sequences in the sequence listing in electronic form are reproduced in the following Table.

SEQUENCE TABLE

<110> GENENTECH, INC.

<120> METHODS AND COMPOSITIONS FOR TARGETING POLYUBIQUITIN

<130> 81014-257

<140> PCT/US2006/062115

<141> 2006-12-14

<150> 60/751,081

<151> 2005-12-15

<150> 60/793,980

<151> 2006-04-21

<160> 908

<170> PatentIn version 3.3

<210> 1

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 1

Gly	Phe	Asn	Leu	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 2

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 2

Gly	Phe	Asn	Val	Ser	Tyr	Ser	Ser	Ile	His
1				5					10

<210> 3

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 3

Gly	Phe	Asn	Ile	Tyr	Tyr	Ser	Ser	Ile	His
1				5					10

<210> 4

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 4

Gly	Phe	Asn	Ile	Ser	Tyr	Tyr	Tyr	Ile	His
1				5					10

<210> 5

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 5

Gly	Phe	Asn	Val	Ser	Tyr	Tyr	Tyr	Met	His
1				5					10

<210> 6

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 6

Gly	Phe	Asn	Phe	Tyr	Ser	Ser	Tyr	Met	His
1				5					10

<210> 7
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 7
 Gly Phe Asn Leu Tyr Tyr Ser Tyr Met His
 1 5 10

<210> 8
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 8
 Gly Phe Asn Val Tyr Tyr Ser Ser Ile His
 1 5 10

<210> 9
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 9
 Gly Phe Asn Ile Ser Tyr Ser Tyr Met His
 1 5 10

<210> 10
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 10
 Gly Phe Asn Val Tyr Tyr Ser Ser Ile His
 1 5 10

<210> 11
 <211> 10

<212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 11

Gly	Phe	Asn	Val	Ser	Tyr	Ser	Tyr	Met	His
1				5					10

<210> 12

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 12

Gly	Phe	Asn	Leu	Tyr	Tyr	Ser	Tyr	Met	His
1				5					10

<210> 13

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 13

Gly	Phe	Asn	Val	Tyr	Tyr	Ser	Ser	Ile	His
1				5					10

<210> 14

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 14

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 15

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 15

Gly	Phe	Asn	Val	Ser	Tyr	Ser	Ser	Ile	His
1				5					10

<210> 16

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 16

Gly	Phe	Asn	Phe	Tyr	Tyr	Tyr	Tyr	Ile	His
1				5					10

<210> 17

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 17

Gly	Phe	Asn	Val	Ser	Ser	Tyr	Ser	Ile	His
1				5					10

<210> 18

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 18

Gly	Phe	Asn	Val	Ser	Ser	Tyr	Ser	Met	His
1				5					10

<210> 19

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 19

Gly Phe Asn Leu Ser Tyr Tyr Ser Ile His
1 5 10

<210> 20

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 20

Gly Phe Asn Leu Ser Tyr Tyr Ser Ile His
1 5 10

<210> 21

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 21

Gly Phe Asn Val Ser Tyr Ser Tyr Met His
1 5 10

<210> 22

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 22

Gly Phe Asn Val Ser Tyr Tyr Ser Ile His
1 5 10

<210> 23

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 23

Gly Phe Asn Leu Ser Tyr Ser Ser Ile His
 1 5 10

<210> 24
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 24
 Gly Phe Asn Val Ser Tyr Ser Ser Ile His
 1 5 10

<210> 25
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 25
 Gly Phe Asn Leu Ser Tyr Ser Ser Met His
 1 5 10

<210> 26
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<220>
 <221> MOD_RES
 <222> (4)..(4)
 <223> Leu, Val, Ile or Phe

<220>
 <221> MOD_RES
 <222> (5)..(8)
 <223> Ser or Tyr

<220>
 <221> MOD_RES
 <222> (9)..(9)
 <223> Met or Ile

<400> 26

Gly Phe Asn Xaa Xaa Xaa Xaa Xaa Xaa His
 1 5 10

<210> 27
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 27
 Ser Ile Tyr Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 28
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 28
 Ser Ile Ser Pro Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 29
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 29
 Ser Ile Tyr Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 30
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic

peptide

<400> 30

Ser	Ile	Ser	Ser	Tyr	Tyr	Gly	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 31

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 31

Ser	Ile	Ser	Ser	Tyr	Tyr	Gly	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 32

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 32

Ser	Ile	Ser	Pro	Tyr	Tyr	Gly	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 33

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 33

Ser	Ile	Ser	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 34

<211> 17

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 34
 Ser Ile Tyr Pro Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 35
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 35
 Ser Ile Ser Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 36
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 36
 Ser Ile Ser Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 37
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 37
 Ser Ile Ser Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 38
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 38
 Ser Ile Ser Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 39
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 39
 Ser Ile Ser Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 40
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 40
 Ser Ile Tyr Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 41
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 41

Ser	Ile	Tyr	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1			5					10						15	

Gly

<210> 42

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 42

Ser	Ile	Ser	Pro	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1			5					10						15	

Gly

<210> 43

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 43

Ser	Ile	Ser	Ser	Ser	Tyr	Ser	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1			5					10						15	

Gly

<210> 44

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 44

Ser	Ile	Tyr	Ser	Ser	Ser	Gly	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1			5					10						15	

Gly

<210> 45

<211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 45
 Ser Ile Tyr Ser Ser Tyr Ser Ser Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 46
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 46
 Ser Ile Tyr Pro Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 47
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 47
 Ser Ile Ser Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 48
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 48

Ser Ile Tyr Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 49

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 49

Ser Ile Tyr Pro Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 50

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 50

Ser Ile Tyr Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 51

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 51

Ser Ile Tyr Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 52

<211> 17

<212> PRT

<213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<220>
 <221> MOD_RES
 <222> (3)..(3)
 <223> Tyr or Ser

<220>
 <221> MOD_RES
 <222> (4)..(4)
 <223> Pro or Ser

<220>
 <221> MOD_RES
 <222> (5)..(6)
 <223> Tyr or Ser

<220>
 <221> MOD_RES
 <222> (7)..(7)
 <223> Ser or Gly

<220>
 <221> MOD_RES
 <222> (8)..(8)
 <223> Tyr or Ser

<220>
 <221> MOD_RES
 <222> (10)..(10)
 <223> Ser or Tyr

<400> 52
 Ser Ile Xaa Xaa Xaa Xaa Xaa Xaa Thr Xaa Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 53
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 53
 Gly Tyr Glu Gly Gly Met Ala Met Asp Tyr
 1 5 10

<210> 54
 <211> 10

<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 54
Asp Gly Tyr Ala Met Asp Ala Leu Asp Tyr
1 5 10

<210> 55
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 55
Leu Tyr His Asn Thr Leu Gly Met Asp Tyr
1 5 10

<210> 56
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 56
Pro Tyr Ser Tyr Ser Glu Ala Met Asp Tyr
1 5 10

<210> 57
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 57
Glu Tyr Tyr Met Tyr Asp Ala Leu Asp Tyr
1 5 10

<210> 58
<211> 10
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 58

Asp Tyr Tyr Tyr Ile Ser Ala Ile Asp Tyr
1 5 10

<210> 59

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 59

Ser Tyr Ser Tyr Ser Ser Ala Leu Asp Tyr
1 5 10

<210> 60

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 60

Gly Tyr Lys Tyr Trp Ser Ala Phe Asp Tyr
1 5 10

<210> 61

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 61

Ser Tyr Ser Ser Tyr Ser Ala Ile Asp Tyr
1 5 10

<210> 62

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 62

Glu Gly Tyr Ser Gln Gly Gly Phe Asp Tyr
1 5 10

<210> 63

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 63

Ser Tyr Gly Tyr Tyr Val Ala Phe Asp Tyr
1 5 10

<210> 64

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 64

Asp Tyr Lys Phe Gly Tyr Ala Ile Asp Tyr
1 5 10

<210> 65

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 65

Glu Gly Tyr Ser Gln Gly Gly Phe Asp Tyr
1 5 10

<210> 66

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 66

Ser Tyr Ser Tyr Tyr Ser Ala Met Asp Tyr
 1 5 10

<210> 67
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 67
 Gly Tyr Met Trp Tyr Gly Gly Ile Asp Tyr
 1 5 10

<210> 68
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 68
 Glu Tyr Tyr Ser Tyr Leu Gly Ala Ile Asp Tyr
 1 5 10

<210> 69
 <211> 21
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 69
 His Thr Lys Tyr Val Tyr Leu Tyr Thr Tyr Trp Glu Asp Ser Met Asp
 1 5 10 15

Tyr Gly Leu Asp Tyr
 20

<210> 70
 <211> 21
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 70

Ser Ser Ile Ser Glu Trp Tyr Gly Ser Trp Tyr Tyr Phe Trp Glu Ser
 1 5 10 15

Ser Gly Ile Asp Tyr
 20

<210> 71

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 71

Glu Ser Tyr Trp Ser Tyr Ala Met Asp Tyr
 1 5 10

<210> 72

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 72

Ser Tyr Ser Tyr Ser Tyr Gly Ile Asp Tyr
 1 5 10

<210> 73

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 73

Ser Tyr Ser Tyr Tyr Ser Ala Ile Asp Tyr
 1 5 10

<210> 74

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 74

Ser Tyr Ser Tyr Ser Tyr Gly Leu Asp Tyr
 1 5 10

<210> 75
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 75
 Gly Tyr Ile His Trp Glu Ala Leu Asp Tyr
 1 5 10

<210> 76
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 76
 Ser Tyr Ser Tyr Ser Ser Gly Leu Asp Tyr
 1 5 10

<210> 77
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 77
 Ser Tyr Ser Tyr Ser Tyr Gly Met Asp Tyr
 1 5 10

<210> 78
 <211> 21
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<220>
 <221> MOD_RES
 <222> (1)..(1)
 <223> Gly, Asp, Leu, Pro, Glu, Ser or His

<220>
<221> MOD_RES
<222> (2)..(2)
<223> Tyr, Gly, Thr or Ser

<220>
<221> MOD_RES
<222> (3)..(3)
<223> Glu, Tyr, His, Ser, Lys, Gly, Met or Ile

<220>
<221> MOD_RES
<222> (4)..(4)
<223> Gly, Ala, Asn, Tyr, Met, Ser, Phe, Trp or His

<220>
<221> MOD_RES
<222> (5)..(5)
<223> Gly, Met, Thr, Ser, Tyr, Ile, Trp, Gln, Val or Glu

<220>
<221> MOD_RES
<222> (6)..(6)
<223> Met, Asp, Leu, Glu, Ser, Gly, Val, Tyr or Trp

<220>
<221> MOD_RES
<222> (7)..(7)
<223> Ala, Gly, Leu or Tyr

<220>
<221> MOD_RES
<222> (8)..(8)
<223> Met, Leu, Ile, Phe, Ala, Tyr or Gly

<220>
<221> MOD_RES
<222> (9)..(9)
<223> Ile, Thr, Ser or not present

<220>
<221> MOD_RES
<222> (10)..(11)
<223> Tyr, Trp or not present

<220>
<221> MOD_RES
<222> (12)..(12)
<223> Glu, Tyr or not present

<220>
<221> MOD_RES
<222> (13)..(13)
<223> Asp, Phe or not present

<220>
 <221> MOD_RES
 <222> (14)..(14)
 <223> Ser, Trp or not present

<220>
 <221> MOD_RES
 <222> (15)..(15)
 <223> Met, Glu or not present

<220>
 <221> MOD_RES
 <222> (16)..(16)
 <223> Asp, Ser or not present

<220>
 <221> MOD_RES
 <222> (17)..(17)
 <223> Tyr, Ser or not present

<220>
 <221> MOD_RES
 <222> (18)..(18)
 <223> Gly or not present

<220>
 <221> MOD_RES
 <222> (19)..(19)
 <223> Leu, Ile or not present

<400> 78
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10 15

Xaa Xaa Xaa Asp Tyr
 20

<210> 79
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 79
 Arg Ala Ser Gln Ser Val Ser Ser Ala Val Ala
 1 5 10

<210> 80
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 80

Ser Ala Ser Ser Leu Tyr Ser
1 5

<210> 81

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 81

Gly Phe Asn Ile Ser Ser Ser Tyr Ile His
1 5 10

<210> 82

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 82

Gly Phe Asn Phe Ser Tyr Ser Tyr Ile His
1 5 10

<210> 83

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 83

Gly Phe Asn Leu Ser Tyr Tyr Tyr Ile His
1 5 10

<210> 84

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 84

Gly	Phe	Asn	Phe	Tyr	Ser	Ser	Tyr	Ile	His
1			5					10	

<210> 85

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 85

Gly	Phe	Asn	Phe	Tyr	Ser	Ser	Ser	Ile	His
1			5					10	

<210> 86

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 86

Gly	Phe	Asn	Ile	Ser	Ser	Ser	Ser	Ile	His
1			5					10	

<210> 87

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 87

Gly	Phe	Asn	Phe	Tyr	Tyr	Ser	Ser	Ile	His
1			5					10	

<210> 88

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 88

Gly Phe Asn Phe Ser Tyr Ser Ser Ile His
 1 5 10

<210> 89

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<400> 89

Gly Phe Asn Val Ser Ser Tyr Ser Ile His
 1 5 10

<210> 90

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<220>

<221> MOD_RES

<222> (4)..(4)

<223> Ile, Phe, Leu or Val

<220>

<221> MOD_RES

<222> (5)..(8)

<223> Ser or Tyr

<400> 90

Gly Phe Asn Xaa Xaa Xaa Xaa Xaa Ile His
 1 5 10

<210> 91

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<400> 91

Tyr Ile Ser Pro Tyr Tyr Gly Ser Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 92
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 92
Ser Ile Ser Ser Ser Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 93
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 93
Ser Ile Ser Ser Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 94
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 94
Ser Ile Ser Ser Ser Tyr Gly Ser Thr Ser Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 95
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 95

Ser	Ile	Ser	Ser	Tyr	Tyr	Gly	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 96

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 96

Ser	Ile	Ser	Pro	Ser	Tyr	Ser	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 97

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 97

Ser	Ile	Tyr	Ser	Ser	Tyr	Gly	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 98

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 98

Ser	Ile	Tyr	Ser	Ser	Tyr	Ser	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 99

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 99

Ser	Ile	Tyr	Pro	Ser	Ser	Gly	Ser	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 100

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD_RES

<222> (1)..(1)

<223> Tyr or Ser

<220>

<221> MOD_RES

<222> (3)..(3)

<223> Ser or Tyr

<220>

<221> MOD_RES

<222> (4)..(4)

<223> Pro or Ser

<220>

<221> MOD_RES

<222> (5)..(6)

<223> Tyr or Ser

<220>

<221> MOD_RES

<222> (7)..(7)

<223> Gly or Ser

<220>

<221> MOD_RES

<222> (8)..(8)

<223> Ser or Tyr

<220>

<221> MOD_RES

<222> (10)..(10)

<223> Ser or Tyr

<400> 100

Xaa	Ile	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Thr	Xaa	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 101

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 101

Glu	Tyr	Tyr	Arg	Trp	Tyr	Thr	Ala	Ile	Asp	Tyr
1				5					10	

<210> 102

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 102

Glu	Lys	Met	Tyr	Tyr	Ser	Tyr	Gly	Phe	Asp	Tyr
1				5					10	

<210> 103

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 103

Glu	Ser	Tyr	Ser	Ile	His	Phe	Gly	Phe	Asp	Tyr
1				5					10	

<210> 104

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 104

Met Tyr Tyr Ser Tyr Tyr Trp Arg Pro Tyr Gly Asn Ala Ile Asp Tyr
 1 5 10 15

<210> 105

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<400> 105

Gly Ser Ile Pro Ser Tyr Trp Ser Ala Asp Trp Tyr Tyr Tyr Tyr Gly
 1 5 10 15

Leu Asp Tyr

<210> 106

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<400> 106

Tyr Lys Tyr Asn Tyr Tyr Tyr Phe Glu Ser Gly Met Asp Tyr
 1 5 10

<210> 107

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<400> 107

Glu Tyr Tyr Trp Trp Tyr Lys Glu Ala Trp Tyr Ser Ala Gly Met Asp
 1 5 10 15

Tyr

<210> 108

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<400> 108

Gly	Ile	Met	Phe	Ser	Ser	Trp	Trp	Trp	Tyr	Tyr	Asp	Tyr	Ser	Asp	Ala
1				5					10					15	

Leu Asp Tyr

<210> 109

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 109

Ser	Gly	Tyr	Tyr	Tyr	Gln	Gly	Tyr	Trp	Trp	Tyr	Tyr	Tyr	Thr	Gly	Tyr
1				5				10						15	

Tyr Gly Met Asp Tyr
20

<210> 110

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD_RES

<222> (1)..(1)

<223> Glu, Met, Gly, Tyr or Ser

<220>

<221> MOD_RES

<222> (2)..(2)

<223> Tyr, Lys, Ser, Ile or Gly

<220>

<221> MOD_RES

<222> (3)..(3)

<223> Tyr, Met or Ile

<220>

<221> MOD_RES

<222> (4)..(4)

<223> Arg, Tyr, Ser, Pro, Asn, Trp or Phe

<220>

<221> MOD_RES

<222> (5)..(5)

<223> Trp, Tyr, Ile or Ser

<220>

<221> MOD_RES

<222> (6)..(6)

<223> Tyr, Ser, His or Gln

<220>

<221> MOD_RES

<222> (7)..(7)

<223> Thr, Tyr, Phe, Trp, Lys or Gly

<220>

<221> MOD_RES

<222> (8)..(8)

<223> Ala, Gly, Arg, Ser, Phe, Glu, Trp or Tyr

<220>

<221> MOD_RES

<222> (9)..(9)

<223> Ile, Phe, Pro, Ala, Glu or Trp

<220>

<221> MOD_RES

<222> (10)..(10)

<223> Tyr, Asp, Ser, Trp or not present

<220>

<221> MOD_RES

<222> (11)..(11)

<223> Gly, Trp, Tyr or not present

<220>

<221> MOD_RES

<222> (12)..(12)

<223> Asn, Tyr, Met, Ser, Asp or not present

<220>

<221> MOD_RES

<222> (13)..(13)

<223> Ala, Tyr or not present

<220>

<221> MOD_RES

<222> (14)..(14)

<223> Ile, Tyr, Gly, Ser, Thr or not present

<220>

<221> MOD_RES

<222> (15)..(15)

<223> Tyr, Met, Asp, Gly or not present

<220>

<221> MOD_RES

<222> (16)..(16)

<223> Gly, Ala, Tyr or not present

<220>

<221> MOD_RES

<222> (17)..(17)

<223> Leu, Tyr or not present

<220>

<221> MOD_RES

<222> (18)..(18)

<223> Gly or not present

<220>

<221> MOD_RES

<222> (19)..(19)

<223> Met or not present

<400> 110

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1				5					10						15

Xaa	Xaa	Xaa	Asp	Tyr
			20	

<210> 111

<211> 30

<212> PRT

<213> Homo sapiens

<400> 111

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala
1				5					10						15

Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr
			20					25					30

<210> 112

<211> 25

<212> PRT

<213> Homo sapiens

<400> 112

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala
1				5					10						15

Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser
			20					25

<210> 113

<211> 25

<212> PRT

<213> Homo sapiens

<400> 113

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser
 20 25

<210> 114
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 114
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser
 20 25

<210> 115
 <211> 30
 <212> PRT
 <213> Homo sapiens

<400> 115
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Val Ser
 20 25 30

<210> 116
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 116
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser
 20 25

<210> 117
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 117
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser
 20 25

<210> 118

<211> 25
 <212> PRT
 <213> Homo sapiens

<400> 118
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Thr Val Ser
 20 25

<210> 119
 <211> 30
 <212> PRT
 <213> Homo sapiens

<400> 119
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
 20 25 30

<210> 120
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 120
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser
 20 25

<210> 121
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 121
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser
 20 25

<210> 122
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 122
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> 123
<211> 30
<212> PRT
<213> Homo sapiens

<400> 123
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Asn Ile Lys
20 25 30

<210> 124
<211> 25
<212> PRT
<213> Homo sapiens

<400> 124
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> 125
<211> 25
<212> PRT
<213> Homo sapiens

<400> 125
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> 126
<211> 30
<212> PRT
<213> Homo sapiens

<400> 126
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Asn Ile Lys
20 25 30

<210> 127
<211> 25
<212> PRT

<213> Homo sapiens

<400> 127

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> 128

<211> 25

<212> PRT

<213> Homo sapiens

<400> 128

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> 129

<211> 25

<212> PRT

<213> Homo sapiens

<400> 129

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> 130

<211> 23

<212> PRT

<213> Homo sapiens

<400> 130

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys
20

<210> 131

<211> 23

<212> PRT

<213> Homo sapiens

<400> 131

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys
20

<210> 132
<211> 23
<212> PRT
<213> Homo sapiens

<400> 132
Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys
20

<210> 133
<211> 23
<212> PRT
<213> Homo sapiens

<400> 133
Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys
20

<210> 134
<211> 23
<212> PRT
<213> Homo sapiens

<400> 134
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys
20

<210> 135
<211> 15
<212> PRT
<213> Homo sapiens

<400> 135
Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
1 5 10 15

<210> 136
<211> 32
<212> PRT
<213> Homo sapiens

<400> 136

Gly Val Pro Ser Arg Phe Ser Gly Ser Arg Ser Gly Thr Asp Phe Thr
 1 5 10 15

Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys
 20 25 30

<210> 137
 <211> 10
 <212> PRT
 <213> Homo sapiens

<400> 137
 Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 1 5 10

<210> 138
 <211> 25
 <212> PRT
 <213> Homo sapiens

<400> 138
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
 20 25

<210> 139
 <211> 13
 <212> PRT
 <213> Homo sapiens

<400> 139
 Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 1 5 10

<210> 140
 <211> 30
 <212> PRT
 <213> Homo sapiens

<400> 140
 Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr Leu Gln
 1 5 10 15

Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 20 25 30

<210> 141
 <211> 11
 <212> PRT
 <213> Homo sapiens

<400> 141

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
1 5 10

<210> 142
<211> 23
<212> PRT
<213> Homo sapiens

<400> 142
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys
20

<210> 143
<211> 15
<212> PRT
<213> Homo sapiens

<400> 143
Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
1 5 10 15

<210> 144
<211> 32
<212> PRT
<213> Homo sapiens

<400> 144
Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
1 5 10 15

Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys
20 25 30

<210> 145
<211> 10
<212> PRT
<213> Homo sapiens

<400> 145
Phe Arg Gln Gly Thr Lys Val Glu Ile Lys
1 5 10

<210> 146
<211> 25
<212> PRT
<213> Homo sapiens

<400> 146
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser

20

25

<210> 147

<211> 13

<212> PRT

<213> Homo sapiens

<400> 147

Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 1 5 10

<210> 148

<211> 30

<212> PRT

<213> Homo sapiens

<400> 148

Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr Leu Gln
 1 5 10 15

Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 20 25 30

<210> 149

<211> 11

<212> PRT

<213> Homo sapiens

<400> 149

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 1 5 10

<210> 150

<211> 66

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 150

tcttgtgaca aaactcacca tcaccatcac catcactagg gcggtggctc tggttccggt 60

gattttt 66

<210> 151

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 151

Gly	Phe	Asn	Val	Tyr	Tyr	Ser	Ser	Ile	His
1				5					10

<210> 152

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 152

Gly	Phe	Asn	Val	Ser	Tyr	Ser	Tyr	Met	His
1				5					10

<210> 153

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 153

Gly	Phe	Asn	Phe	Ser	Tyr	Tyr	Ser	Met	His
1				5					10

<210> 154

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 154

Gly	Phe	Asn	Leu	Ser	Tyr	Tyr	Ser	Ile	His
1				5					10

<210> 155

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 155

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
 1 5 10

<210> 156
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 156
 Gly Phe Asn Val Tyr Tyr Ser Ser Ile His
 1 5 10

<210> 157
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 157
 Gly Phe Asn Leu Ser Tyr Ser Tyr Ile His
 1 5 10

<210> 158
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 158
 Gly Phe Asn Phe Tyr Tyr Tyr Tyr Ile His
 1 5 10

<210> 159
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 159
 Gly Phe Asn Leu Ser Tyr Ser Ser Ile His
 1 5 10

<210> 160
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 160
Gly Phe Asn Val Ser Tyr Ser Ser Ile His
1 5 10

<210> 161
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 161
Gly Phe Asn Val Ser Tyr Ser Ser Ile His
1 5 10

<210> 162
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 162
Gly Phe Asn Val Tyr Tyr Ser Ser Ile His
1 5 10

<210> 163
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 163
Gly Phe Asn Val Ser Tyr Tyr Tyr Ile His
1 5 10

<210> 164
<211> 10
<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 164

Gly	Phe	Asn	Leu	Ser	Tyr	Ser	Ser	Ile	His
1			5					10	

<210> 165

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 165

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Tyr	Met	His
1			5					10	

<210> 166

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 166

Gly	Phe	Asn	Leu	Tyr	Tyr	Ser	Tyr	Met	His
1			5					10	

<210> 167

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 167

Gly	Phe	Asn	Val	Ser	Tyr	Tyr	Tyr	Met	His
1			5					10	

<210> 168

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 168

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Tyr	Met	His
1				5				10	

<210> 169

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 169

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Ile	His
1				5				10	

<210> 170

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 170

Gly	Phe	Asn	Val	Ser	Tyr	Ser	Ser	Met	His
1				5				10	

<210> 171

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 171

Gly	Phe	Asn	Leu	Ser	Tyr	Tyr	Ser	Ile	His
1				5				10	

<210> 172

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 172

Gly Phe Asn Val Ser Tyr Tyr Ser Ile His
1 5 10

<210> 173

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 173

Gly Phe Asn Ile Ser Tyr Ser Ser Ile His
1 5 10

<210> 174

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 174

Gly Phe Asn Phe Ser Tyr Tyr Ser Ile His
1 5 10

<210> 175

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 175

Gly Phe Asn Leu Ser Tyr Ser Ser Met His
1 5 10

<210> 176

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<220>

<221> MOD_RES

<222> (4)..(4)

<223> Val, Phe, Leu or Ile

<220>

<221> MOD_RES

<222> (5)..(5)

<223> Tyr or Ser

<220>

<221> MOD_RES

<222> (7)..(8)

<223> Ser or Tyr

<220>

<221> MOD_RES

<222> (9)..(9)

<223> Ile or Met

<400> 176

Gly	Phe	Asn	Xaa	Xaa	Tyr	Xaa	Xaa	Xaa	His
1				5					10

<210> 177

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 177

Ser	Ile	Ser	Pro	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 178

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 178

Ser	Ile	Ser	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 179

<211> 17

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 179
 Ser Ile Tyr Pro Tyr Tyr Gly Tyr Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 180
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 180
 Ser Ile Tyr Pro Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 181
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 181
 Ser Ile Tyr Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 182
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 182
 Ser Ile Tyr Pro Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 183
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 183
 Ser Ile Ser Ser Tyr Tyr Gly Ser Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 184
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 184
 Ser Ile Ser Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 185
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 185
 Ser Ile Tyr Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 186
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic

peptide

<400> 186

Ser	Ile	Tyr	Pro	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 187

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 187

Ser	Ile	Tyr	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 188

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 188

Ser	Ile	Tyr	Pro	Tyr	Tyr	Gly	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 189

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 189

Ser	Ile	Ser	Ser	Tyr	Tyr	Gly	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 190

<211> 17

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 190
 Ser Ile Tyr Ser Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 191
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 191
 Ser Ile Ser Pro Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 192
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 192
 Ser Ile Ser Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 193
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 193
 Ser Ile Ser Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 194

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 194

Ser	Ile	Ser	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5				10						15	

Gly

<210> 195

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 195

Ser	Ile	Ser	Ser	Ser	Tyr	Ser	Ser	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5				10						15	

Gly

<210> 196

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 196

Ser	Ile	Ser	Pro	Tyr	Tyr	Gly	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5				10						15	

Gly

<210> 197

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

peptide

<400> 197

Ser	Ile	Tyr	Ser	Ser	Tyr	Ser	Ser	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 198

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 198

Ser	Ile	Tyr	Pro	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 199

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 199

Ser	Ile	Tyr	Pro	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 200

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 200

Ser	Ile	Tyr	Pro	Tyr	Tyr	Ser	Tyr	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 201

<211> 17

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 201
 Ser Ile Tyr Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 202
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<220>
 <221> MOD_RES
 <222> (3)..(3)
 <223> Ser or Tyr

<220>
 <221> MOD_RES
 <222> (4)..(4)
 <223> Pro or Ser

<220>
 <221> MOD_RES
 <222> (5)..(5)
 <223> Tyr or Ser

<220>
 <221> MOD_RES
 <222> (7)..(7)
 <223> Ser or Gly

<220>
 <221> MOD_RES
 <222> (8)..(8)
 <223> Tyr or Ser

<220>
 <221> MOD_RES
 <222> (10)..(10)
 <223> Ser or Tyr

<400> 202
 Ser Ile Xaa Xaa Xaa Tyr Xaa Xaa Thr Xaa Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 203
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 203
 Glu Gly Tyr Ser Gln Gly Gly Phe Asp Tyr
 1 5 10

<210> 204
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 204
 Ser Tyr Ser Tyr Tyr Ser Ala Ile Asp Tyr
 1 5 10

<210> 205
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 205
 Ser Tyr Ser Tyr Ser Tyr Gly Met Asp Tyr
 1 5 10

<210> 206
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 206
 Ser Tyr Ser Tyr Ser Tyr Gly Ile Asp Tyr
 1 5 10

<210> 207
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 207
 Ser Tyr Ser Tyr Tyr Ser Ala Met Asp Tyr
 1 5 10

<210> 208
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 208
 Gly Tyr Lys Tyr Trp Ser Ala Phe Asp Tyr
 1 5 10

<210> 209
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 209
 Glu Ser Phe Tyr Tyr Ser Pro Ala Phe Asp Tyr
 1 5 10

<210> 210
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 210
 Glu Tyr Tyr Ser Tyr Leu Gly Ala Ile Asp Tyr
 1 5 10

<210> 211
 <211> 10
 <212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 211

Gly	Tyr	Glu	Gly	Gly	Met	Ala	Met	Asp	Tyr
1				5					10

<210> 212

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 212

Ser	Tyr	Ser	Tyr	Ser	Ser	Gly	Leu	Asp	Tyr
1				5					10

<210> 213

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 213

Gly	Tyr	Met	Trp	Tyr	Gly	Gly	Ile	Asp	Tyr
1				5					10

<210> 214

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD_RES

<222> (5)..(5)

<223> Variable amino acid

<400> 214

Asp	Cys	Tyr	Tyr	Xaa	Ala	Ala	Phe	Asp	Tyr
1				5					10

<210> 215
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 215
Glu Asn Tyr Trp Trp Ala Ile Asp Tyr
1 5

<210> 216
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 216
Ser Tyr Ser Tyr Tyr Ser Ala Phe Asp Tyr
1 5 10

<210> 217
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 217
Asp Tyr Tyr Phe Phe Ser Ala Ile Asp Tyr
1 5 10

<210> 218
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 218
Ser Tyr Ser Tyr Ser Ser Ala Leu Asp Tyr
1 5 10

<210> 219
<211> 11
<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 219

Glu	Gly	Tyr	Ile	Ser	Gly	Asp	Ala	Ile	Asp	Tyr
1				5					10	

<210> 220

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 220

Ser	Tyr	Ser	Ser	Tyr	Ser	Ala	Ile	Asp	Tyr
1				5					10

<210> 221

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 221

Gly	Tyr	Phe	Glu	Gly	Trp	Tyr	Gly	Leu	Asp	Tyr
1				5					10	

<210> 222

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 222

Glu	Tyr	Ser	Tyr	Tyr	Gly	Gly	Phe	Asp	Tyr
1				5					10

<210> 223

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 223

Glu	Ser	Tyr	Trp	Ser	Tyr	Ala	Met	Asp	Tyr
1				5					10

<210> 224

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 224

Ser	Tyr	Ser	Tyr	Ser	Tyr	Gly	Leu	Asp	Tyr
1				5					10

<210> 225

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 225

Tyr	Tyr	Ser	Tyr	Ser	Ser	Gly	Leu	Asp	Tyr
1				5					10

<210> 226

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 226

Ser	Tyr	Ser	Tyr	Ser	Tyr	Gly	Leu	Asp	Tyr
1				5					10

<210> 227

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 227
 Ser Tyr Ser Tyr Ser Tyr Gly Met Asp Tyr
 1 5 10

<210> 228
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<220>
 <221> MOD_RES
 <222> (1)..(1)
 <223> Glu, Ser, Gly, Asp or Tyr

<220>
 <221> MOD_RES
 <222> (2)..(2)
 <223> Gly, Tyr, Ser, Cys or Asn

<220>
 <221> MOD_RES
 <222> (3)..(3)
 <223> Tyr, Ser, Lys, Phe, Glu or Met

<220>
 <221> MOD_RES
 <222> (4)..(4)
 <223> Ser, Tyr, Gly, Trp, Phe, Ile or Glu

<220>
 <221> MOD_RES
 <222> (5)..(5)
 <223> Variable amino acid

<220>
 <221> MOD_RES
 <222> (6)..(6)
 <223> Gly, Ser, Tyr, Leu, Met, Gly, Ala or Trp

<220>
 <221> MOD_RES
 <222> (7)..(7)
 <223> Gly, Ala, Pro, Ile, Asp or Tyr

<220>
 <221> MOD_RES
 <222> (8)..(8)
 <223> Phe, Ile, Met, Ala, Leu, Gly or not present

<220>
 <221> MOD_RES

<222> (9)..(9)

<223> Phe, Ile, Leu or not present

<400> 228

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Asp	Tyr
1				5					10	

<210> 229

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 229

Gly	Phe	Asn	Ile	Ser	Ser	Ser	Tyr	Ile	His
1				5					10

<210> 230

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 230

Gly	Phe	Asn	Val	Ser	Ser	Tyr	Ser	Ile	His
1				5					10

<210> 231

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 231

Gly	Phe	Asn	Phe	Ser	Tyr	Ser	Ser	Ile	His
1				5					10

<210> 232

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 232

Gly Phe Asn Phe Tyr Ser Ser Ser Ile His
1 5 10

<210> 233

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 233

Gly Phe Asn Phe Tyr Tyr Ser Ser Ile His
1 5 10

<210> 234

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 234

Gly Phe Asn Val Tyr Tyr Ser Ser Ile His
1 5 10

<210> 235

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 235

Gly Phe Asn Phe Tyr Tyr Ser Tyr Ile His
1 5 10

<210> 236

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 236

Gly Phe Asn Val Ser Ser Ser Tyr Ile His
1 5 10

<210> 237
 <211> 10
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic peptide

 <400> 237
 Gly Phe Asn Phe Tyr Tyr Ser Ser Ile His
 1 5 10

 <210> 238
 <211> 10
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic peptide

 <400> 238
 Gly Phe Asn Phe Tyr Ser Ser Tyr Met His
 1 5 10

 <210> 239
 <211> 10
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic peptide

 <400> 239
 Gly Phe Asn Val Ser Ser Tyr Ser Ile His
 1 5 10

 <210> 240
 <211> 10
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic peptide

 <220>
 <221> MOD_RES
 <222> (4)..(4)
 <223> Ile, Val or Phe

 <220>

<221> MOD_RES
 <222> (5)..(8)
 <223> Ser or Tyr

<220>
 <221> MOD_RES
 <222> (9)..(9)
 <223> Ile or Met

<400> 240
 Gly Phe Asn Xaa Xaa Xaa Xaa Xaa Xaa His
 1 5 10

<210> 241
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 241
 Tyr Ile Ser Pro Tyr Tyr Gly Ser Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 242
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 242
 Ser Ile Tyr Pro Ser Ser Gly Ser Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 243
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 243
 Ser Ile Tyr Ser Ser Tyr Ser Ser Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 244

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 244

Ser	Ile	Ser	Ser	Tyr	Tyr	Gly	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 245

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 245

Ser	Ile	Tyr	Ser	Ser	Tyr	Gly	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 246

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 246

Ser	Ile	Ser	Pro	Ser	Tyr	Gly	Tyr	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 247

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

peptide

<400> 247

Tyr	Ile	Ser	Pro	Ser	Tyr	Ser	Ser	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 248

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 248

Ser	Ile	Ser	Ser	Ser	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 249

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 249

Ser	Ile	Tyr	Ser	Ser	Tyr	Ser	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 250

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 250

Tyr	Ile	Ser	Ser	Tyr	Ser	Gly	Tyr	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 251

<211> 17

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 251
 Ser Ile Tyr Pro Ser Ser Gly Ser Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 252
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<220>
 <221> MOD_RES
 <222> (1)..(1)
 <223> Tyr or Ser

<220>
 <221> MOD_RES
 <222> (3)..(3)
 <223> Ser or Tyr

<220>
 <221> MOD_RES
 <222> (4)..(4)
 <223> Pro or Ser

<220>
 <221> MOD_RES
 <222> (5)..(6)
 <223> Tyr or Ser

<220>
 <221> MOD_RES
 <222> (7)..(7)
 <223> Gly or Ser

<220>
 <221> MOD_RES
 <222> (8)..(8)
 <223> Ser or Tyr

<220>
 <221> MOD_RES
 <222> (10)..(10)

<223> Ser or Tyr

<400> 252

Xaa	Ile	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Thr	Xaa	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 253

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 253

Glu	Tyr	Tyr	Arg	Trp	Tyr	Thr	Ala	Ile	Asp	Tyr
1				5					10	

<210> 254

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 254

Ser	Gly	Tyr	Tyr	Tyr	Gln	Gly	Tyr	Trp	Trp	Tyr	Tyr	Tyr	Thr	Gly	Tyr
1					5				10					15	

Tyr	Gly	Met	Asp	Tyr
				20

<210> 255

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 255

Gly	Ile	Met	Phe	Ser	Ser	Trp	Trp	Trp	Tyr	Tyr	Asp	Tyr	Ser	Asp	Ala
1					5				10					15	

Leu Asp Tyr

<210> 256

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 256

Gly	Ser	Ile	Pro	Ser	Tyr	Trp	Ser	Ala	Asp	Trp	Tyr	Tyr	Tyr	Tyr	Gly
1				5					10					15	

Leu Asp Tyr

<210> 257

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 257

Glu	Tyr	Tyr	Trp	Trp	Tyr	Lys	Glu	Ala	Trp	Tyr	Ser	Ala	Gly	Met	Asp
1				5					10					15	

Tyr

<210> 258

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 258

Trp	Gln	Gly	Tyr	Gly	Phe	Lys	Tyr	Tyr	Trp	Ser	Tyr	Tyr	Val	Ser	Tyr
1				5					10					15	

Gly	Gly	Leu	Asp	Tyr
			20	

<210> 259

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 259

Ser	Tyr	Ser	Tyr	Tyr	Tyr	Tyr	Ser	Ser	Tyr	Gly	Phe	Asp	Tyr
1				5					10				

<210> 260
 <211> 13
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 260
 Glu Ser Tyr Ala Gly Val Pro Pro Tyr Gly Phe Asp Tyr
 1 5 10

<210> 261
 <211> 19
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 261
 Gly Ile Met Phe Ser Ser Trp Trp Trp Tyr Tyr Asp Tyr Ser Asp Ala
 1 5 10 15

Leu Asp Tyr

<210> 262
 <211> 19
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 262
 Gly Ile Met Phe Ser Ser Trp Trp Trp Tyr Tyr Asp Tyr Ser Asp Ala
 1 5 10 15

Leu Asp Tyr

<210> 263
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 263
 Glu Tyr Tyr Trp Trp Tyr Lys Glu Ala Trp Tyr Ser Ala Gly Met Asp
 1 5 10 15

Tyr

<210> 264

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD_RES

<222> (1)..(1)

<223> Glu, Ser, Gly or Trp

<220>

<221> MOD_RES

<222> (2)..(2)

<223> Tyr, Gly, Ile, Ser or Gln

<220>

<221> MOD_RES

<222> (3)..(3)

<223> Tyr, Met, Ile, Gly or Ser

<220>

<221> MOD_RES

<222> (4)..(4)

<223> Arg, Tyr, Phe, Pro or Trp

<220>

<221> MOD_RES

<222> (5)..(5)

<223> Trp, Tyr, Ser or Gly

<220>

<221> MOD_RES

<222> (6)..(6)

<223> Tyr, Gln, Ser, Phe or Val

<220>

<221> MOD_RES

<222> (7)..(7)

<223> Thr, Gly, Trp, Lys, Tyr or Pro

<220>

<221> MOD_RES

<222> (8)..(8)

<223> Ala, Tyr, Trp, Ser, Glu or Pro

<220>

<221> MOD_RES

<222> (9)..(9)

<223> Ile, Trp, Ala, Tyr or Ser

<220>

<221> MOD_RES

<222> (10)..(10)

<223> Trp, Tyr, Asp or Gly

<220>

<221> MOD_RES

<222> (11)..(11)

<223> Tyr, Trp, Ser, Gly or Phe

<220>

<221> MOD_RES

<222> (12)..(12)

<223> Tyr, Asp, Ser, Phe or not present

<220>

<221> MOD_RES

<222> (13)..(13)

<223> Tyr, Ala or not present

<220>

<221> MOD_RES

<222> (14)..(14)

<223> Thr, Ser, Tyr, Gly, Val or not present

<220>

<221> MOD_RES

<222> (15)..(15)

<223> Gly, Asp, Tyr, Met, Ser or not present

<220>

<221> MOD_RES

<222> (16)..(16)

<223> Tyr, Ala, Gly or not present

<220>

<221> MOD_RES

<222> (17)..(17)

<223> Tyr, Leu, Gly or not present

<220>

<221> MOD_RES

<222> (18)..(18)

<223> Gly or not present

<220>

<221> MOD_RES

<222> (19)..(19)

<223> Met, Leu or not present

<400> 264

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa

1

5

10

15

Xaa Xaa Xaa Asp Tyr
20

<210> 265
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 265
Gly Phe Asn Val Tyr Tyr Ser Ser Ile His
1 5 10

<210> 266
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 266
Gly Phe Asn Val Ser Tyr Ser Tyr Met His
1 5 10

<210> 267
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 267
Gly Phe Asn Phe Ser Tyr Tyr Ser Met His
1 5 10

<210> 268
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 268
Gly Phe Asn Leu Ser Tyr Tyr Ser Ile His
1 5 10

<210> 269
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 269
 Gly Phe Asn Ile Ser Tyr Ser Ser Met His
 1 5 10

<210> 270
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 270
 Gly Phe Asn Val Tyr Tyr Ser Ser Ile His
 1 5 10

<210> 271
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 271
 Gly Phe Asn Leu Ser Tyr Ser Tyr Ile His
 1 5 10

<210> 272
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 272
 Gly Phe Asn Leu Ser Tyr Ser Ser Met His
 1 5 10

<210> 273
 <211> 10

<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 273

Gly	Phe	Asn	Val	Ser	Tyr	Ser	Ser	Ile	His
1				5					10

<210> 274

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 274

Gly	Phe	Asn	Val	Ser	Tyr	Ser	Ser	Ile	His
1				5					10

<210> 275

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 275

Gly	Phe	Asn	Val	Ser	Tyr	Tyr	Tyr	Ile	His
1				5					10

<210> 276

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 276

Gly	Phe	Asn	Leu	Tyr	Tyr	Ser	Tyr	Met	His
1				5					10

<210> 277

<211> 10

<212> PRT

<213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 277
 Gly Phe Asn Val Ser Tyr Tyr Ser Ile His
 1 5 10

<210> 278
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 278
 Gly Phe Asn Ile Ser Tyr Ser Ser Ile His
 1 5 10

<210> 279
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 279
 Gly Phe Asn Phe Ser Tyr Tyr Ser Ile His
 1 5 10

<210> 280
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<220>
 <221> MOD_RES
 <222> (4)..(4)
 <223> Val, Phe, Leu or Ile

<220>
 <221> MOD_RES
 <222> (5)..(5)
 <223> Tyr or Ser

<220>
 <221> MOD_RES

<222> (7)..(8)
 <223> Ser or Tyr

<220>
 <221> MOD_RES
 <222> (9)..(9)
 <223> Ile or Met

<400> 280
 Gly Phe Asn Xaa Xaa Tyr Xaa Xaa Xaa His
 1 5 10

<210> 281
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 281
 Ser Ile Ser Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 282
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 282
 Ser Ile Ser Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 283
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 283
 Ser Ile Tyr Pro Tyr Tyr Gly Tyr Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 284
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 284
 Ser Ile Tyr Pro Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 285
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 285
 Ser Ile Tyr Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 286
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 286
 Ser Ile Tyr Pro Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 287
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic

peptide

<400> 287

Ser	Ile	Ser	Ser	Tyr	Tyr	Gly	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 288

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 288

Ser	Ile	Tyr	Pro	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 289

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 289

Ser	Ile	Tyr	Pro	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 290

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 290

Ser	Ile	Tyr	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 291

<211> 17

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 291
 Ser Ile Ser Ser Tyr Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 292
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 292
 Ser Ile Ser Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 293
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 293
 Ser Ile Tyr Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 294
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 294
 Ser Ile Tyr Pro Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 295

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 295

Ser	Ile	Tyr	Pro	Tyr	Tyr	Ser	Tyr	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 296

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD_RES

<222> (3)..(3)

<223> Ser or Tyr

<220>

<221> MOD_RES

<222> (4)..(4)

<223> Pro or Ser

<220>

<221> MOD_RES

<222> (7)..(7)

<223> Ser or Gly

<220>

<221> MOD_RES

<222> (8)..(8)

<223> Tyr or Ser

<220>

<221> MOD_RES

<222> (10)..(10)

<223> Ser or Tyr

<400> 296

Ser	Ile	Xaa	Xaa	Tyr	Tyr	Xaa	Xaa	Thr	Xaa	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 297
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 297
 Glu Gly Tyr Ser Gln Gly Gly Phe Asp Tyr
 1 5 10

<210> 298
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 298
 Ser Tyr Ser Tyr Tyr Ser Ala Ile Asp Tyr
 1 5 10

<210> 299
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 299
 Ser Tyr Ser Tyr Ser Tyr Gly Met Asp Tyr
 1 5 10

<210> 300
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 300
 Ser Tyr Ser Tyr Ser Tyr Gly Ile Asp Tyr
 1 5 10

<210> 301
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 301
Ser Tyr Ser Tyr Tyr Ser Ala Met Asp Tyr
1 5 10

<210> 302
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 302
Gly Tyr Lys Tyr Trp Ser Ala Phe Asp Tyr
1 5 10

<210> 303
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 303
Glu Ser Phe Tyr Tyr Ser Pro Ala Phe Asp Tyr
1 5 10

<210> 304
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 304
Gly Tyr Glu Gly Gly Met Ala Met Asp Tyr
1 5 10

<210> 305
<211> 10
<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 305

Ser Tyr Ser Tyr Ser Ser Gly Leu Asp Tyr
1 5 10

<210> 306

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 306

Gly Tyr Met Trp Tyr Gly Gly Ile Asp Tyr
1 5 10

<210> 307

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 307

Glu Asn Tyr Trp Trp Ala Ile Asp Tyr
1 5

<210> 308

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 308

Ser Tyr Ser Tyr Ser Ser Ala Leu Asp Tyr
1 5 10

<210> 309

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 309

Ser	Tyr	Ser	Tyr	Ser	Tyr	Gly	Leu	Asp	Tyr
1				5					10

<210> 310

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 310

Tyr	Tyr	Ser	Tyr	Ser	Ser	Gly	Leu	Asp	Tyr
1				5					10

<210> 311

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 311

Ser	Tyr	Ser	Tyr	Ser	Tyr	Gly	Leu	Asp	Tyr
1				5					10

<210> 312

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD_RES

<222> (1)..(1)

<223> Glu, Ser, Gly or Tyr

<220>

<221> MOD_RES

<222> (2)..(2)

<223> Gly, Tyr, Ser or Asn

<220>

<221> MOD_RES

<222> (3)..(3)

<223> Tyr, Ser, Lys, Phe or Glu

<220>

<221> MOD_RES

<222> (4)..(4)

<223> Ser, Tyr, Gly or Trp

<220>

<221> MOD_RES

<222> (5)..(5)

<223> Gln, Tyr, Ser or Gly

<220>

<221> MOD_RES

<222> (6)..(6)

<223> Gly, Ser, Tyr, Met or Ala

<220>

<221> MOD_RES

<222> (7)..(7)

<223> Gly, Ala, Pro or Ile

<220>

<221> MOD_RES

<222> (8)..(8)

<223> Phe, Ile, Met, Ala, Leu or not present

<220>

<221> MOD_RES

<222> (9)..(9)

<223> Phe or not present

<400> 312

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asp Tyr

1

5

10

<210> 313

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 313

Gln Gln Ser Ser Tyr Ser Ser Leu Phe Thr

1

5

10

<210> 314

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 314

Gln Gln Ser Ser Tyr Ser Ser Leu Phe Thr
1 5 10

<210> 315

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 315

Gln Gln Tyr Ser Ser Tyr Ser Ser Pro Ile Thr
1 5 10

<210> 316

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 316

Gln Gln Tyr Ser Ser Tyr Tyr Ser Pro Val Thr
1 5 10

<210> 317

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 317

Gln Gln Ser Ser Tyr Ser Ser Leu Ile Thr
1 5 10

<210> 318

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 318

Gln Gln Ser Ser Tyr Ser Ser Leu Val Thr
1 5 10

<210> 319

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 319

Gln Gln Ser Tyr Tyr Tyr Ser Leu Phe Thr
1 5 10

<210> 320

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 320

Gln Gln Ser Ser Tyr Ser Ser Leu Val Thr
1 5 10

<210> 321

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 321

Gln Gln Ser Ser Tyr Ser Ser Leu Phe Thr
1 5 10

<210> 322

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 322

Gln Gln Tyr Ser Tyr Ser Ser Leu Phe Thr
1 5 10

<210> 323
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 323
 Gln Gln Ser Tyr Tyr Tyr Tyr Pro Ile Thr
 1 5 10

<210> 324
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 324
 Gln Gln Ser Ser Tyr Ser Ser Leu Val Thr
 1 5 10

<210> 325
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 325
 Gln Gln Tyr Ser Ser Ser Tyr Tyr Pro Phe Thr
 1 5 10

<210> 326
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 326
 Gln Gln Ser Ser Tyr Ser Ser Leu Leu Thr
 1 5 10

<210> 327
 <211> 11

<212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 327
 Gln Gln Tyr Tyr Tyr Tyr Tyr Tyr Pro Ile Thr
 1 5 10

<210> 328
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<220>
 <221> MOD_RES
 <222> (3)..(7)
 <223> Ser or Tyr

<220>
 <221> MOD_RES
 <222> (8)..(8)
 <223> Leu, Ser, Pro or Tyr

<220>
 <221> MOD_RES
 <222> (9)..(9)
 <223> Pro or not present

<220>
 <221> MOD_RES
 <222> (10)..(10)
 <223> Phe, Ile, Val or Leu

<400> 328
 Gln Gln Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Thr
 1 5 10

<210> 329
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 329

Gly Phe Asn Val Ser Ser Tyr Ser Ile His
 1 5 10

<210> 330

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 330

Gly Phe Asn Ile Ser Ser Ser Tyr Ile His
 1 5 10

<210> 331

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 331

Gly Phe Asn Phe Ser Tyr Ser Ser Ile His
 1 5 10

<210> 332

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 332

Gly Phe Asn Val Tyr Tyr Ser Ser Ile His
 1 5 10

<210> 333

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 333

Gly Phe Asn Phe Tyr Tyr Ser Ser Ile His
 1 5 10

<210> 334

<211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 334
 Gly Phe Asn Val Ser Ser Ser Tyr Ile His
 1 5 10

<210> 335
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 335
 Gly Phe Asn Phe Tyr Ser Ser Tyr Met His
 1 5 10

<210> 336
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 336
 Gly Phe Asn Phe Tyr Ser Ser Ser Ile His
 1 5 10

<210> 337
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<220>
 <221> MOD_RES
 <222> (4)..(4)
 <223> Val, Ile or Phe

<220>
 <221> MOD_RES
 <222> (5)..(8)

<223> Ser or Tyr

<220>

<221> MOD_RES

<222> (9)..(9)

<223> Ile or Met

<400> 337

Gly	Phe	Asn	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	His
1			5						10

<210> 338

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 338

Ser	Ile	Tyr	Pro	Ser	Ser	Gly	Ser	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 339

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 339

Tyr	Ile	Ser	Pro	Tyr	Tyr	Gly	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 340

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 340

Ser	Ile	Tyr	Ser	Ser	Tyr	Ser	Ser	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 341
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 341
 Ser Ile Ser Pro Ser Tyr Gly Tyr Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 342
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 342
 Ser Ile Tyr Ser Ser Tyr Gly Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 343
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 343
 Ser Ile Ser Ser Ser Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 344
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 344

Tyr Ile Ser Ser Tyr Ser Gly Tyr Thr Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 345

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 345

Ser Ile Ser Ser Tyr Tyr Gly Ser Thr Ser Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 346

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<220>

<221> MOD_RES

<222> (1)..(1)

<223> Ser or Tyr

<220>

<221> MOD_RES

<222> (3)..(3)

<223> Tyr or Ser

<220>

<221> MOD_RES

<222> (4)..(4)

<223> Pro or Ser

<220>

<221> MOD_RES

<222> (5)..(6)

<223> Ser or Tyr

<220>

<221> MOD_RES

<222> (7)..(7)

<223> Gly or Ser

<220>
 <221> MOD_RES
 <222> (8)..(8)
 <223> Ser or Tyr

<220>
 <221> MOD_RES
 <222> (10)..(10)
 <223> Tyr or Ser

<400> 346
 Xaa Ile Xaa Xaa Xaa Xaa Xaa Xaa Thr Xaa Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 347
 <211> 21
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 347
 Ser Gly Tyr Tyr Tyr Gln Gly Tyr Trp Trp Tyr Tyr Tyr Thr Gly Tyr
 1 5 10 15

Tyr Gly Met Asp Tyr
 20

<210> 348
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 348
 Glu Tyr Tyr Arg Trp Tyr Thr Ala Ile Asp Tyr
 1 5 10

<210> 349
 <211> 19
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 349

Gly Ile Met Phe Ser Ser Trp Trp Trp Tyr Tyr Asp Tyr Ser Asp Ala
 1 5 10 15

Leu Asp Tyr

<210> 350

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<400> 350

Trp Gln Gly Tyr Gly Phe Lys Tyr Tyr Trp Ser Tyr Tyr Val Ser Tyr
 1 5 10 15

Gly Gly Leu Asp Tyr
 20

<210> 351

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<400> 351

Glu Tyr Tyr Trp Trp Tyr Lys Glu Ala Trp Tyr Ser Ala Gly Met Asp
 1 5 10 15

Tyr

<210> 352

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 peptide

<400> 352

Glu Ser Tyr Ala Gly Val Pro Pro Tyr Gly Phe Asp Tyr
 1 5 10

<210> 353

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 353

Gly	Ile	Met	Phe	Ser	Ser	Trp	Trp	Trp	Tyr	Tyr	Asp	Tyr	Ser	Asp	Ala
1				5					10					15	

Leu Asp Tyr

<210> 354

<211> 19

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 354

Gly	Ser	Ile	Pro	Ser	Tyr	Trp	Ser	Ala	Asp	Trp	Tyr	Tyr	Tyr	Tyr	Gly
1				5					10					15	

Leu Asp Tyr

<210> 355

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD_RES

<222> (1)..(1)

<223> Ser, Glu, Gly or Trp

<220>

<221> MOD_RES

<222> (2)..(2)

<223> Gly, Tyr, Ile, Gln or Ser

<220>

<221> MOD_RES

<222> (3)..(3)

<223> Tyr, Met, Gly or Ile

<220>

<221> MOD_RES

<222> (4)..(4)

<223> Tyr, Arg, Phe, Trp, Ala or Pro

<220>

<221> MOD_RES

<222> (5)..(5)
<223> Tyr, Trp, Ser or Gly

<220>
<221> MOD_RES
<222> (6)..(6)
<223> Gln, Tyr, Ser, Phe or Val

<220>
<221> MOD_RES
<222> (7)..(7)
<223> Gly, Thr, Trp, Lys or Pro

<220>
<221> MOD_RES
<222> (8)..(8)
<223> Tyr, Ala, Trp, Glu, Pro or Ser

<220>
<221> MOD_RES
<222> (9)..(9)
<223> Trp, Ile, Tyr or Ala

<220>
<221> MOD_RES
<222> (10)..(10)
<223> Trp, Tyr, Gly, Asp or not present

<220>
<221> MOD_RES
<222> (11)..(11)
<223> Tyr, Ser, Phe, Trp or not present

<220>
<221> MOD_RES
<222> (12)..(12)
<223> Tyr, Asp, Ser or not present

<220>
<221> MOD_RES
<222> (13)..(13)
<223> Tyr, Ala or not present

<220>
<221> MOD_RES
<222> (14)..(14)
<223> Thr, Ser, Val, Gly, Tyr or not present

<220>
<221> MOD_RES
<222> (15)..(15)
<223> Gly, Asp, Ser, Met, Tyr or not present

<220>
<221> MOD_RES

<222> (16)..(16)

<223> Tyr, Ala, Gly or not present

<220>

<221> MOD_RES

<222> (17)..(17)

<223> Tyr, Leu, Gly or not present

<220>

<221> MOD_RES

<222> (18)..(18)

<223> Gly or not present

<220>

<221> MOD_RES

<222> (19)..(19)

<223> Met, Leu or not present

<400> 355

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1			5					10						15	

Xaa	Xaa	Xaa	Asp	Tyr
			20	

<210> 356

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 356

Gln	Gln	Tyr	Ser	Tyr	Tyr	Pro	Phe	Arg
1			5					

<210> 357

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 357

Gln	Gln	Tyr	Ser	Ser	Tyr	Ser	Ser	Leu	Phe	Thr
1			5					10		

<210> 358

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 358

Gln Gln Tyr Ser Ser Ser Leu Val Thr
1 5

<210> 359

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 359

Gln Gln Tyr Ser Ser Ser Ser Pro Phe Thr Phe
1 5 10

<210> 360

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 360

Gln Gln Ser Ser Tyr Ser Pro Ile Thr
1 5

<210> 361

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 361

Gln Gln Tyr Ser Tyr Ser Ser Tyr Leu Ile Thr
1 5 10

<210> 362

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 362
 Gln Gln Ser Tyr Tyr Ser Pro Phe Thr
 1 5

<210> 363
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<400> 363
 Gln Gln Tyr Tyr Ser Ser Leu Val Thr
 1 5

<210> 364
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 peptide

<220>
 <221> MOD_RES
 <222> (3)..(6)
 <223> Tyr or Ser

<220>
 <221> MOD_RES
 <222> (7)..(7)
 <223> Pro, Ser or Leu

<220>
 <221> MOD_RES
 <222> (8)..(8)
 <223> Ser, Pro, Tyr or not present

<220>
 <221> MOD_RES
 <222> (9)..(9)
 <223> Leu, Phe or not present

<220>
 <221> MOD_RES
 <222> (10)..(10)
 <223> Phe, Val, Thr or Ile

<220>
 <221> MOD_RES
 <222> (11)..(11)

<223> Arg, Thr or Phe

<400> 364

Gln Gln Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10

<210> 365

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 365

cgtctattat tgtgctcgct aataagacta ctgggggtcaa gg

42

<210> 366

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 366

cgtctattat tgtgctcgct aataagacta ctgggggtcaa gg

42

<210> 367

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (20)..(22)

<223> a, c, g or t

<220>

<221> modified_base

<222> (26)..(30)

<223> a, c, g or t

<220>

<221> modified_base

<222> (32)..(33)

<223> a, c, g or t

<220>

<221> modified_base

<222> (35)..(37)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 367

cgtctattat tgtgctcgcn nntacnnnnn snnsnnngst wtsgactact ggggtcaagg 60

<210> 368

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (20)..(24)

<223> a, c, g or t

<220>

<221> modified_base

<222> (26)..(28)

<223> a, c, g or t

<220>

<221> modified_base

<222> (32)..(33)

<223> a, c, g or t

<220>

<221> modified_base

<222> (35)..(37)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 368

cgtctattat tgtgctcgcn nnnnsnnnta cnnsnnngst wtsgactact ggggtcaagg 60

<210> 369

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

oligonucleotide

<220>

<221> modified_base

<222> (20)..(24)

<223> a, c, g or t

<220>

<221> modified_base

<222> (26)..(30)

<223> a, c, g or t

<220>

<221> modified_base

<222> (35)..(37)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 369

cgtctattat tgtgctcgcn nnnnsnnnnn stacnnngst wtsgactact ggggtcaagg 60

<210> 370

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (20)..(24)

<223> a, c, g or t

<220>

<221> modified_base

<222> (26)..(30)

<223> a, c, g or t

<220>

<221> modified_base

<222> (32)..(33)

<223> a, c, g or t

<220>

<221> modified_base

<222> (35)..(37)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of

substitutions and preferred embodiments

<400> 370
cgtctattat tgtgctcgcn nnnnsnnnnn snnsnnngst wtsgactact ggggtcaagg 60

<210> 371
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 371
gcagcttctg gcttcaacta ataacactgg gtgcgtcagg 40

<210> 372
<211> 52
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>
<221> modified_base
<222> (19)..(19)
<223> a, c, g or t

<220>
<221> modified_base
<222> (22)..(23)
<223> a, c, g or t

<220>
<221> modified_base
<222> (31)..(32)
<223> a, c, g or t

<400> 372
gcagcttctg gcttcaacnt cnnstactct nnsatscact ggggtgcgtca gg 52

<210> 373
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 373
ggcctggaat gggttgcata ataatatgcc gatagcgtca agg 43

<210> 374
 <211> 67
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (25)..(26)
 <223> a, c, g or t

<400> 374
 ggcttggaat gggttgcatc taccnnsyct tactactctt acacctctta tgccgatagc 60
 gtcaagg 67

<210> 375
 <211> 60
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (32)..(33)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (35)..(36)
 <223> a, c, g or t

<400> 375
 cgtctattat tgtgctcgct cttactctta cnsnnsngst wtsgactact ggggtcaagg 60

<210> 376
 <211> 46
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> 376
 cgcaacttat tactgtcagc aataataaac gttcggacag ggtacc 46

<210> 377
 <211> 76
 <212> PRT
 <213> Homo sapiens

<400> 377
 Met Gln Ile Phe Val Lys Thr Leu Thr Gly Lys Thr Ile Thr Leu Glu
 1 5 10 15
 Val Glu Pro Ser Asp Thr Ile Glu Asn Val Lys Ala Lys Ile Gln Asp
 20 25 30
 Lys Glu Gly Ile Pro Pro Asp Gln Gln Arg Leu Ile Phe Ala Gly Lys
 35 40 45
 Gln Leu Glu Asp Gly Arg Thr Leu Ser Asp Tyr Asn Ile Gln Lys Glu
 50 55 60
 Ser Thr Leu His Leu Val Leu Arg Leu Arg Gly Gly
 65 70 75

<210> 378
 <211> 61
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (23)..(24)
 <223> a, c, g or t

<400> 378
 cgcaacttat tactgtcagc aannstctta ctcttctctg dttacgttcg gacagggtac 60
 c 61

<210> 379
 <211> 63
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (20)..(22)
 <223> a, c, g or t

<220>

<221> modified_base

<222> (29)..(33)

<223> a, c, g or t

<220>

<221> modified_base

<222> (35)..(36)

<223> a, c, g or t

<220>

<221> modified_base

<222> (38)..(40)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 379

cgtctattat tgtgctcgcn nntactacnn nnnsnnnsnnn gstwtsgact actgggggtca 60

agg 63

<210> 380

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (20)..(24)

<223> a, c, g or t

<220>

<221> modified_base

<222> (26)..(27)

<223> a, c, g or t

<220>

<221> modified_base

<222> (29)..(31)

<223> a, c, g or t

<220>

<221> modified_base

<222> (38)..(40)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 380
 cgtctattat tgtgctcgcn nnnnsnnnsnn ntggtacnnn gstwtsgact actgggggtca 60
 agg 63

<210> 381
 <211> 63
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (20)..(22)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (29)..(33)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (38)..(40)
 <223> a, c, g or t

<220>
 <223> see specification as filed for detailed description of
 substitutions and preferred embodiments

<400> 381
 cgtctattat tgtgctcgcn nntacbbsnn nnnstacnnn gstwtsgact actgggggtca 60
 agg 63

<210> 382
 <211> 63
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (20)..(24)
 <223> a, c, g or t

<220>
 <221> modified_base

<222> (29)..(31)

<223> a, c, g or t

<220>

<221> modified_base

<222> (35)..(36)

<223> a, c, g or t

<220>

<221> modified_base

<222> (38)..(40)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 382

cgtctattat tgtgctcgcn nnnnstacnn ntggnnnsnnn gstwtsgact actgggggtca 60

agg 63

<210> 383

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (20)..(22)

<223> a, c, g or t

<220>

<221> modified_base

<222> (26)..(27)

<223> a, c, g or t

<220>

<221> modified_base

<222> (29)..(31)

<223> a, c, g or t

<220>

<221> modified_base

<222> (35)..(36)

<223> a, c, g or t

<220>

<221> modified_base

<222> (38)..(40)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 383

cgtctattat tgtgctcgcn nntacnnsnn ntggnnnsnnn gstwts gact actgggggtca 60

agg 63

<210> 384

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (20)..(24)

<223> a, c, g or t

<220>

<221> modified_base

<222> (29)..(33)

<223> a, c, g or t

<220>

<221> modified_base

<222> (38)..(40)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 384

cgtctattat tgtgctcgcn nnnnstacnn nnnstacnnn gstwts gact actgggggtca 60

agg 63

<210> 385

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (20)..(24)

<223> a, c, g or t

<220>

<221> modified_base

<222> (26)..(27)

<223> a, c, g or t

<220>

<221> modified_base

<222> (29)..(33)

<223> a, c, g or t

<220>

<221> modified_base

<222> (35)..(36)

<223> a, c, g or t

<220>

<221> modified_base

<222> (38)..(40)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 385

cgtctattat tgtgctcgcn nnnnsnnsnn nnnnsnnsnn gstwtsgact actgggggtca 60

agg

63

<210> 386

<211> 67

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (25)..(26)

<223> a, c, g or t

<220>

<221> modified_base

<222> (31)..(32)

<223> a, c, g or t

<220>

<221> modified_base

<222> (34)..(35)

<223> a, c, g or t

<220>
 <221> modified_base
 <222> (40)..(42)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (46)..(48)
 <223> a, c, g or t

<220>
 <223> see specification as filed for detailed description of
 substitutions and preferred embodiments

<400> 386
 ggccctggaat gggttgcata catcnnsyct nnsnnsergn nnaccnnnta tgccgatagc 60
 gtcaagg 67

<210> 387
 <211> 67
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (19)..(20)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (25)..(26)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (34)..(35)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (40)..(42)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (46)..(48)
 <223> a, c, g or t

<220>
 <223> see specification as filed for detailed description of

substitutions and preferred embodiments

<400> 387

ggcctggaat gggttgcann saccnnsyct tacnnsrgcn nnaccnnnta tgccgatagc 60

gtcaagg

67

<210> 388

<211> 67

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (19)..(20)

<223> a, c, g or t

<220>

<221> modified_base

<222> (25)..(26)

<223> a, c, g or t

<220>

<221> modified_base

<222> (31)..(32)

<223> a, c, g or t

<220>

<221> modified_base

<222> (40)..(42)

<223> a, c, g or t

<220>

<221> modified_base

<222> (46)..(48)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 388

ggcctggaat gggttgcann saccnnsyct nnstacrgcn nnaccnnnta tgccgatagc 60

gtcaagg

67

<210> 389

<211> 67

<212> DNA

<213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (19)..(20)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (25)..(26)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (31)..(32)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (34)..(35)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (40)..(42)
 <223> a, c, g or t

<220>
 <221> modified_base
 <222> (46)..(48)
 <223> a, c, g or t

<220>
 <223> see specification as filed for detailed description of
 substitutions and preferred embodiments

<400> 389
 ggcctggaat gggttgcann satcnnsyct nnsnnsrgcn nnaccnnnta tgccgatagc 60
 gtcaagg 67

<210> 390
 <211> 64
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 oligonucleotide

<220>
 <221> modified_base
 <222> (23)..(24)

<223> a, c, g or t

<220>

<221> modified_base

<222> (26)..(33)

<223> a, c, g or t

<220>

<221> modified_base

<222> (35)..(39)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 390

cgcaacttat tactgtcagc aannsnnnnnn nnnsnnnnns ctgdttacgt tcggacaggg 60

tacc 64

<210> 391

<211> 52

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<220>

<221> modified_base

<222> (19)..(19)

<223> a, c, g or t

<220>

<221> modified_base

<222> (22)..(23)

<223> a, c, g or t

<220>

<221> modified_base

<222> (25)..(32)

<223> a, c, g or t

<220>

<223> see specification as filed for detailed description of
substitutions and preferred embodiments

<400> 391

gcagcttctg gcttcaacnt cnnsnnnnnn nnsatscact ggggtgcgtca gg 52

<210> 392

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 392

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 393

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 393

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 394

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 394

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 395

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 395

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 396

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 396

Gly	Phe	Asn	Ile	Gly	Tyr	Ser	Phe	Met	His
1			5						10

<210> 397

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 397

Gly	Phe	Asn	Val	Asp	Tyr	Ser	Tyr	Met	His
1			5						10

<210> 398

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 398

Gly	Phe	Asn	Val	Asp	Tyr	Ser	Tyr	Met	His
1			5						10

<210> 399

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 399

Gly	Phe	Asn	Phe	Ser	Tyr	Ser	Phe	Met	His
1			5						10

<210> 400

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 400

Gly Phe Asn Ile Val Tyr Ser Phe Met His
1 5 10

<210> 401

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 401

Gly Phe Asn Ile Ile Tyr Ser Phe Met His
1 5 10

<210> 402

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 402

Gly Phe Asn Ile Val Tyr Ser Phe Ile His
1 5 10

<210> 403

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 403

Gly Phe Asn Leu Ser Tyr Ser Phe Met His
1 5 10

<210> 404

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 404

Gly Phe Asn Val Asp Tyr Ser Phe Met His
1 5 10

<210> 405
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 405
Gly Phe Asn Val Ile Tyr Ser Phe Met His
1 5 10

<210> 406
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 406
Gly Phe Asn Val Ala Tyr Ser Leu Met His
1 5 10

<210> 407
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 407
Gly Phe Asn Ile Ser Tyr Ser Trp Met His
1 5 10

<210> 408
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 408
Gly Phe Asn Leu Asp Tyr Ser Phe Met His
1 5 10

<210> 409
<211> 10

<212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic peptide

 <400> 409
 Gly Phe Asn Phe Leu Tyr Ser Gly Ile His
 1 5 10

 <210> 410
 <211> 10
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic peptide

 <400> 410
 Gly Phe Asn Ile Ser Tyr Ser Ser Met His
 1 5 10

 <210> 411
 <211> 10
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic peptide

 <400> 411
 Gly Phe Asn Ile Ser Tyr Ser Ser Met His
 1 5 10

 <210> 412
 <211> 10
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic peptide

 <400> 412
 Gly Phe Asn Ile Ser Tyr Ser Ser Met His
 1 5 10

 <210> 413
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 413

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 414

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 414

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 415

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 415

Gly	Phe	Asn	Ile	Leu	Tyr	Ser	Gly	Ile	His
1				5					10

<210> 416

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 416

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 417

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 417

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 418

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 418

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 419

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 419

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 420

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 420

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 421

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 421

Gly Phe Asn Ile Ser Tyr Ser Ser Met His

1 5 10

<210> 422
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 422
 Gly Phe Asn Ile Ser Tyr Ser Ser Met His
 1 5 10

<210> 423
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 423
 Gly Phe Asn Ile Ser Tyr Ser Ser Met His
 1 5 10

<210> 424
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 424
 Gly Phe Asn Ile Ser Tyr Ser Ser Met His
 1 5 10

<210> 425
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 425
 Gly Phe Asn Ile Ser Tyr Ser Ser Met His
 1 5 10

<210> 426

<211> 10
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 426

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1			5					10	

<210> 427

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 427

Gly	Phe	Asn	Ile	Phe	Tyr	Ser	Gly	Ile	His
1			5					10	

<210> 428

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 428

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1			5					10	

<210> 429

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 429

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1			5					10	

<210> 430

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 430

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 431

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 431

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 432

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 432

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 433

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 433

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 434

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

peptide

<400> 434

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 435

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 435

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 436

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 436

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 437

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 437

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 438

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 438

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 439

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 439

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 440

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 440

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 441

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 441

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 442

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 442

Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 443
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 443
Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 444
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 444
Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 445
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 445
Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 446
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 446
Gly Phe Asn Ile Ser Tyr Ser Ser Met His
1 5 10

<210> 447
<211> 10
<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 447

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 448

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 448

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 449

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 449

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 450

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 450

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 451

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 451

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 452

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 452

Gly	Phe	Asn	Ile	Ser	Tyr	Ser	Ser	Met	His
1				5					10

<210> 453

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 453

Gly	Phe	Asn	Leu	Ser	Tyr	Ser	Gly	Met	His
1				5					10

<210> 454

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 454

Gly	Phe	Asn	Leu	Leu	Tyr	Ser	Gly	Met	His
1				5					10

<210> 455

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 455

Gly Phe Asn Val Ala Tyr Ser Gly Ile His
1 5 10

<210> 456

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 456

Gly Phe Asn Val Asp Tyr Ser Gly Met His
1 5 10

<210> 457

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 457

Gly Phe Asn Val Asp Tyr Ser Gly Met His
1 5 10

<210> 458

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 458

Gly Phe Asn Val Ser Tyr Ser Ser Ile His
1 5 10

<210> 459

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 459

Gly Phe Asn Val Val Tyr Ser Gly Ile His
1 5 10

<210> 460
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<220>
 <221> MOD_RES
 <222> (4)..(4)
 <223> Ile, Val, Phe or Leu

<220>
 <221> MOD_RES
 <222> (5)..(5)
 <223> Ser, Gly, Asp, Val, Ile, Leu, Phe or Ala

<220>
 <221> MOD_RES
 <222> (8)..(8)
 <223> Ser, Phe, Tyr, Leu, Trp or Gly

<220>
 <221> MOD_RES
 <222> (9)..(9)
 <223> Met or Ile

<400> 460
 Gly Phe Asn Xaa Xaa Tyr Ser Xaa Xaa His
 1 5 10

<210> 461
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 461
 Ser Ile Tyr Ser Tyr Tyr Ser Tyr Thr Ser Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 462
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 462

Ser	Ile	Tyr	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 463

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 463

Ser	Ile	Tyr	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 464

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 464

Ser	Ile	Tyr	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 465

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 465

Ser	Ile	Ala	Ser	Tyr	Tyr	Ser	Tyr	Thr	Ser	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 466

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

CLAIMS

1. An isolated antibody that specifically binds a first polyubiquitin comprising a first lysine linkage, wherein the antibody does not specifically bind a second polyubiquitin comprising a second lysine linkage or wherein the antibody binds the second polyubiquitin comprising a second lysine linkage with substantially reduced affinity as compared to the binding affinity of the antibody for the first polyubiquitin, and wherein the first lysine linkage differs from the second lysine linkage, wherein the first polyubiquitin is lysine-48-linked and the second polyubiquitin is lysine-63-linked or wherein the first polyubiquitin is lysine-63-linked and the second polyubiquitin is lysine-48-linked.
2. The antibody of claim 1, wherein the first polyubiquitin is lysine-48-linked and the second polyubiquitin is lysine-63-linked.
3. The antibody of claim 1, wherein the first polyubiquitin is lysine-63-linked and the second polyubiquitin is lysine-48-linked.
4. The antibody of claim 2, wherein the antibody does not specifically bind monoubiquitin.
5. The antibody of claim 4, comprising HVR-H1, HVR-H2, HVR-H3, and HVR-L3 sequences corresponding to those set forth for clones apu01, apu02, apu03, apu04, apu05, apu06, apu07, apu08, apu09, apu10, apu11, apu12, apu13, apu14, or apu15 shown as SEQ ID NOs: 265-280; SEQ ID NOs: 281-296; SEQ ID NOs: 297-312; and SEQ ID NOs: 313-328, respectively; and comprising an HVR-L1 sequence of SEQ ID NO: 79 and an HVR-L2 sequence of SEQ ID NO: 80.
6. The antibody of claim 4, comprising HVR-H1, HVR-H2, HVR-H3, and HVR-L3 sequences corresponding to those set forth for clones apu2.01, apu2.02, apu2.03, apu2.04, apu2.05, apu2.06, apu2.07, apu2.08, apu2.09, or apu2.10 shown as SEQ ID NOs: 695-705; SEQ ID NOs: 706-716; SEQ ID NOs: 717-727; and SEQ ID NOs: 728-738, respectively; and comprising an HVR-L1 sequence of SEQ ID NO: 79 and an HVR-L2 sequence of SEQ ID NO: 80.
7. The antibody of claim 4, comprising an HVR-H1 sequence of SEQ ID NO: 269, an HVR-H2 sequence of SEQ ID NO: 285, an HVR-H3 sequence of SEQ ID NO: 301, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 317.

8. The antibody of claim 4, comprising an HVR-H1 sequence of SEQ ID NO: 701, an HVR-H2 sequence of SEQ ID NO: 712, an HVR-H3 sequence of SEQ ID NO: 723, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 734.

9. The antibody of claim 3, wherein the antibody does not specifically bind to monoubiquitin.

10. The antibody of claim 9, comprising HVR-H1, HVR-H2, HVR-H3, and HVR-L3 sequences corresponding to those set forth for clones apu17, apu18, apu19, apu20, apu21, apu22, apu23, and apu24 shown as SEQ ID NOs: 329-337; SEQ ID NOs: 338-346; SEQ ID NOs: 347-355; and SEQ ID NOs: 356-364, respectively; and comprising an HVR-L1 sequence of SEQ ID NO: 79 and an HVR-L2 sequence of SEQ ID NO: 80.

11. The antibody of claim 9, comprising HVR-H1, HVR-H2, HVR-H3, and HVR-L3 sequences corresponding to those set forth for clones apu2.11, apu2.12, apu2.13, apu2.14, apu2.15, apu2.16, apu2.17, apu2.18, apu2.19, and apu2.20 shown as SEQ ID NOs: 739-749; SEQ ID NOs: 750-760; SEQ ID NOs: 761-771; and SEQ ID NOs: 772-782, respectively; and comprising an HVR-L1 sequence of SEQ ID NO: 79 and an HVR-L2 sequence of SEQ ID NO: 80.

12. The antibody of claim 9, comprising an HVR-H1 sequence of SEQ ID NO: 330, an HVR-H2 sequence of SEQ ID NO: 339, an HVR-H3 sequence of SEQ ID NO: 348, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 357.

13. The antibody of claim 9, comprising an HVR-H1 sequence of SEQ ID NO: 739, an HVR-H2 sequence of SEQ ID NO: 750, an HVR-H3 sequence of SEQ ID NO: 761, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 772.

14. The antibody of claim 9, comprising an HVR-H1 sequence of SEQ ID NO: 740, an HVR-H2 sequence of SEQ ID NO: 751, an HVR-H3 sequence of SEQ ID NO: 762, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 773.

15. The antibody of claim 9, comprising HVR-H1, HVR-H2, and HVR-H3 sequences corresponding to those set forth for clones apu3.01, apu3.02, apu3.03, apu3.04, apu3.05,

apu3.06, apu3.07, apu3.08, apu3.09, apu3.10, and 3.11 shown as SEQ ID NOs: 789-800; SEQ ID NOs: 801-812; and SEQ ID NOs: 813-824, respectively; and comprising an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 777.

16. The antibody of claim 9, comprising an HVR-H1 sequence of SEQ ID NO: 744, an HVR-H2 sequence of SEQ ID NO: 755, an HVR-H3 sequence of SEQ ID NO: 766, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 777.

17. The antibody of claim 9, comprising an HVR-H1 sequence of SEQ ID NO: 795, an HVR-H2 sequence of SEQ ID NO: 807, an HVR-H3 sequence of SEQ ID NO: 819, an HVR-L1 sequence of SEQ ID NO: 79, an HVR-L2 sequence of SEQ ID NO: 80, and an HVR-L3 sequence of SEQ ID NO: 777.

18. The antibody of any one of claims 1 to 17, wherein the antibody specifically binds to a polyubiquitinated protein.

19. A nucleic acid molecule encoding the antibody of any one of claims 5 to 8 and 10 to 17.

20. A vector that comprises the nucleic acid of claim 19.

21. A host cell comprising the vector of claim 20.

22. A cell line capable of producing the antibody of any one of claims 1 to 18.

23. A method of producing the antibody of any one of claims 1 to 18, comprising culturing a host cell comprising a nucleic acid molecule encoding the antibody under conditions wherein the antibody is produced.

24. A method of identifying the presence of polyubiquitin or a polyubiquitinated protein in a sample, comprising contacting the sample with at least one antibody of any one of claims 1 to 18.

25. The method of claim 24, further comprising determining the binding of the at least one antibody to a polyubiquitin or polyubiquitinated protein in the sample.

26. A method of separating polyubiquitinated protein from non-polyubiquitinated protein in a sample, comprising contacting the sample with at least one antibody of any one of claims 1 to 18.

27. The isolated antibody of claim 9, wherein the antibody binds to an epitope that includes residues in both a first ubiquitin subunit and a second ubiquitin subunit of the lysine-63-linked polyubiquitin.

28. The isolated antibody of claim 27, wherein the epitope includes at least one residue in a first ubiquitin subunit selected from Glu-18, Pro-19, Ser-20, Asp-21, Thr-55, Leu-56, Ser-57, Asp-58, Asn-60, Ile-61, and Gln-62.

29. The isolated antibody of claim 27, wherein the epitope includes at least one residue in a second ubiquitin subunit selected from Leu-8, Thr-9, Glu-34, Gly-35, Ile-36, Pro-37, Asp-39, Gln-40, Leu-71, Arg-72, Leu-73, Arg-74, and Gly-75.


30. The isolated antibody of claim 27, wherein the epitope includes at least one residue in a first ubiquitin subunit selected from Glu-18, Pro-19, Ser-20, Asp-21, Thr-55, Leu-56, Ser-57, Asp-58, Asn-60, Ile-61, and Gln-62, and at least one residue in a second ubiquitin subunit selected from Leu-8, Thr-9, Glu-34, Gly-35, Ile-36, Pro-37, Asp-39, Gln-40, Leu-71, Arg-72, Leu-73, Arg-74, and Gly-75.

31. An antigen-binding fragment of the antibody of any one of claims 1 to 18 or 27 to 30.

P2260R1

1 / 45

1 MET GLN ILE PHE VAL LYS THR LEU THR GLY LYS THR ILE THR
15 LEU GLU VAL GLU PRO SER ASP THR ILE GLU ASN VAL LYS ALA
29 LYS ILE GLN ASP LYS GLU GLY ILE PRO PRO ASP GLN GLN ARG
43 LEU ILE PHE ALA GLY LYS GLN LEU GLU ASP GLY ARG THR LEU
57 SER ASP TYR ASN ILE GLN LYS GLU SER THR LEU HIS LEU VAL
71 LEU ARG LEU ARG GLY GLY

FIG. 1A**FIG. 1B**

HVR-H1

Clone #	SEQ ID NO:											Bind to
		26	27	28	29	30	31	32	33	34	35	
48-1	1	G	F	N	L	S	Y	S	S	M	H	48
48-2	2	G	F	N	V	S	Y	S	S	I	H	48
48-3	3	G	F	N	I	Y	Y	S	S	I	H	48
48-4	4	G	F	N	I	S	Y	Y	Y	I	H	48
48-5	5	G	F	N	V	S	Y	Y	Y	M	H	48
48-6	6	G	F	N	F	Y	S	S	Y	M	H	48
48-7	7	G	F	N	L	Y	Y	S	Y	M	H	48
48-8	8	G	F	N	V	Y	Y	S	S	I	H	48
48-9	9	G	F	N	I	S	Y	S	Y	M	H	48
48-10	10	G	F	N	V	Y	Y	S	S	I	H	48
48-11	11	G	F	N	V	S	Y	S	Y	M	H	48
48-12	12	G	F	N	L	Y	Y	S	Y	M	H	48
48-13	13	G	F	N	V	Y	Y	S	S	I	H	48
48-14	14	G	F	N	I	S	Y	S	S	M	H	48
48-15	15	G	F	N	V	S	Y	S	S	I	H	48/BOTH
48-16	16	G	F	N	F	Y	Y	Y	Y	I	H	48/BOTH
48-17	17	G	F	N	V	S	S	Y	S	I	H	48
48-18	18	G	F	N	V	S	S	Y	S	M	H	48/BOTH
48-19	19	G	F	N	L	S	Y	Y	S	I	H	48/BOTH
48-20	20	G	F	N	L	S	Y	Y	S	I	H	ALL
48-21	21	G	F	N	V	S	Y	S	Y	M	H	BOTH
48-22	22	G	F	N	V	S	Y	Y	S	I	H	BOTH
48-23	23	G	F	N	L	S	Y	S	S	I	H	BOTH
48-24	24	G	F	N	V	S	Y	S	S	I	H	BOTH
48-25	25	G	F	N	L	S	Y	S	S	M	H	NONE/BOTH
Consensus	26	G	F	N	L/V/ I/F	S/Y	Y/S	S/Y	S/Y	M/I	H	

FIG. 2A

3 / 45

2025

2025

P2260R1

4 / 45

HVR-H3

Clone #	SEQ ID NO:																						Bind to
		95	96	97	98	99	100	100a	100b	100c	100d	100e	100f	100g	100h	100i	100j	100k	100l	100m	101	102	
48-1	53	G	Y	E	G	G	M	A	M											D	Y	48	
48-2	54	D	G	Y	A	M	D	A	L											D	Y	48	
48-3	55	L	Y	H	N	T	L	G	M											D	Y	48	
48-4	56	P	Y	S	Y	S	E	A	M											D	Y	48	
48-5	57	E	Y	Y	M	Y	D	A	L											D	Y	48	
48-6	58	D	Y	Y	Y	I	S	A	I											D	Y	48	
48-7	59	S	Y	S	Y	S	S	A	L											D	Y	48	
48-8	60	G	Y	K	Y	W	S	A	F											D	Y	48	
48-9	61	S	Y	S	S	Y	S	A	I											D	Y	48	
48-10	62	E	G	Y	S	Q	G	G	F											D	Y	48	
48-11	63	S	Y	G	Y	Y	V	A	F											D	Y	48	
48-12	64	D	Y	K	F	G	Y	A	I											D	Y	48	
48-13	65	E	G	Y	S	Q	G	G	F											D	Y	48	
48-14	66	S	Y	S	Y	Y	S	A	M											D	Y	48	
48-15	67	G	Y	M	W	Y	G	G	I											D	Y	48/BOTH	
48-16	68	E	Y	Y	S	Y	L	G	A	I										D	Y	48/BOTH	
48-17	69	H	T	K	Y	V	Y	L	Y	T	Y	W	E	D	S	M	D	Y	G	L	D	Y	48
48-18	70	S	S	I	S	E	W	Y	G	S	W	Y	Y	F	W	E	S	S	G	I	D	Y	48/BOTH
48-19	71	E	S	Y	W	S	Y	A	M											D	Y	48/BOTH	
48-20	72	S	Y	S	Y	S	Y	G	I											D	Y	ALL	
48-21	73	S	Y	S	Y	Y	S	A	I											D	Y	BOTH	
48-22	74	S	Y	S	Y	S	Y	G	L											D	Y	BOTH	
48-23	75	G	Y	I	H	W	E	A	L											D	Y	BOTH	
48-24	76	S	Y	S	Y	S	S	G	L											D	Y	BOTH	
48-25	77	S	Y	S	Y	S	Y	G	M											D	Y	BOTH	
Con-sensus	78	G/D/ L/P/ E/S/ H	Y/ G/ T/S	E/Y/ H/S/ K/G/ M/I	G/A/N/ Y/M/ S/F/ W/H	G/M/T/ S/Y/I/ W/Q/ V/E	M/D/L/ E/S/G/ V/Y/W	A/G/ L/Y	M/L/ I/F/ A/Y/ G	I/T/ S/ n.p.	Y/ W/ n.p.	W/ Y/ n.p.	E/ Y/ n.p.	D/ F/ n.p.	S/ W/ n.p.	M/ E/ n.p.	D/ S/ n.p.	Y/ S/ n.p.	G/ n.p.	L/I/ n.p.	D	Y	

FIG. 2C

5 / 45

HR-12

Clone #	SEQ ID NO:																			Binds to
		50	51	52	52a	53	54	55	56	57	58	59	60	61	62	63	64	65		
63-1	91	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G		
63-2	92	S	I	S	S	S	Y	G	Y	T	S	Y	A	D	S	V	K	G	63/BOTH	
63-3	93	S	I	S	S	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G	ALL	
63-4	94	S	I	S	S	S	Y	G	S	T	S	Y	A	D	S	V	K	G	BOTH	
63-5	95	S	I	S	S	Y	Y	G	S	T	S	Y	A	D	S	V	K	G	63/BOTH	
63-6	96	S	I	S	P	S	Y	S	S	T	S	Y	A	D	S	V	K	G	BOTH	
63-7	97	S	I	Y	S	S	Y	G	Y	T	S	Y	A	D	S	V	K	G	ALL	
63-8	98	S	I	Y	S	S	Y	S	S	T	S	Y	A	D	S	V	K	G	BOTH	
63-9	99	S	I	Y	P	S	S	G	S	T	Y	Y	A	D	S	V	K	G	ALL	
Consensus	100	Y/S	I	S/Y	P/S	Y/S	Y/S	G/S	S/Y	T	S/Y	Y	A	D	S	V	K	G	ALL	

FIG. 3A

6 / 45

BB
GG
FF

P2260R1

7 / 45

I	A	QVQLVQSGAEVKKPGASVKVSCKASGYTFT	-H1-	WVRQAPGQGLEWMG	-H2-	RVTIT
	B	QVQLVQSGAEVKKPGASVKVSCKAS	-H1-	WVRQAPGQGLEWM	-H2-	RVTIT
	C	QVQLVQSGAEVKKPGASVKVSCKAS	-H1-	WVRQAPGQGLEWM	-H2-	RVTIT
	D	QVQLVQSGAEVKKPGASVKVSCKAS	-H1-	WVRQAPGQGLEWM	-H2-	RVTIT
II	A	QVQLQESGPGGLVKKPSQTLSTCTVSGGSVS	-H1-	WIRQPPGKGLEWIG	-H2-	RVTIS
	B	QVQLQESGPGGLVKKPSQTLSTCTVS	-H1-	WIRQPPGKGLEWI	-H2-	RVTIS
	C	QVQLQESGPGGLVKKPSQTLSTCTVS	-H1-	WIRQPPGKGLEWI	-H2-	RVTIS
	D	QVQLQESGPGGLVKKPSQTLSTCTVS	-H1-	WIRQPPGKGLEWI	-H2-	RVTIS
III	A	EVQLVESGGGLVQPGGSLRLSCAASGFTFS	-H1-	WVRQAPGKGLEWVS	-H2-	RFTIS
	B	EVQLVESGGGLVQPGGSLRLSCAAS	-H1-	WVRQAPGKGLEWV	-H2-	RFTIS
	C	EVQLVESGGGLVQPGGSLRLSCAAS	-H1-	WVRQAPGKGLEWV	-H2-	RFTIS
	D	EVQLVESGGGLVQPGGSLRLSCAAS	-H1-	WVRQAPGKGLEWV	-H2-	RFTIS
Acceptor						
A	A	EVQLVESGGGLVQPGGSLRLSCAASGENIK	-H1-	WVRQAPGKGLEWVS	-H2-	RFTIS
	B	EVQLVESGGGLVQPGGSLRLSCAAS	-H1-	WVRQAPGKGLEWV	-H2-	RFTIS
	C	EVQLVESGGGLVQPGGSLRLSCAAS	-H1-	WVRQAPGKGLEWV	-H2-	RFTIS
Second Acceptor						
A	A	EVQLVESGGGLVQPGGSLRLSCAASGENIK	-H1-	WVRQAPGKGLEWVS	-H2-	RFTIS
	B	EVQLVESGGGLVQPGGSLRLSCAAS	-H1-	WVRQAPGKGLEWV	-H2-	RFTIS
	C	EVQLVESGGGLVQPGGSLRLSCAAS	-H1-	WVRQAPGKGLEWV	-H2-	RFTIS
	D	EVQLVESGGGLVQPGGSLRLSCAAS	-H1-	WVRQAPGKGLEWV	-H2-	RFTIS

FIG. 4A

P2260R1

8 / 45

I	A	ADTSTSTAYMELSSLRSED	TA	VYYCAR		-H3-	WGQGT	LV	TVSS	SEQ ID NO.: 111, 839, 858 & 877
	B	ADTSTSTAYMELSSLRSED	TA	VYYCAR		-H3-	WGQGT	LV	TVSS	SEQ ID NO.: 112, 840, 859 & 878
	C	ADTSTSTAYMELSSLRSED	TA	VYYCA		-H3-	WGQGT	LV	TVSS	SEQ ID NO.: 113, 841, 860 & 879
	D	ADTSTSTAYMELSSLRSED	TA	VYYC		-H3-	WGQGT	LV	TVSS	SEQ ID NO.: 114, 842, 861 & 880
II	A	VDTSKNQFSLKLS	SV	TA	ADTA	VYYCAR				SEQ ID NO.: 115, 843, 862 & 881
	B	VDTSKNQFSLKLS	SV	TA	ADTA	VYYCAR				SEQ ID NO.: 116, 844, 863 & 882
	C	VDTSKNQFSLKLS	SV	TA	ADTA	VYYCA				SEQ ID NO.: 117, 845, 864 & 883
	D	VDTSKNQFSLKLS	SV	TA	ADTA	VYYC				SEQ ID NO.: 118, 846, 865 & 884
III	A	RDNSKNTLYLQMN	SL	RA	EDTA	VYYCAR				SEQ ID NO.: 119, 847, 866 & 885
	B	RDNSKNTLYLQMN	SL	RA	EDTA	VYYCAR				SEQ ID NO.: 120, 848, 867 & 886
	C	RDNSKNTLYLQMN	SL	RA	EDTA	VYYCA				SEQ ID NO.: 121, 849, 868 & 887
	D	RDNSKNTLYLQMN	SL	RA	EDTA	VYYC				SEQ ID NO.: 122, 850, 869 & 888
Acceptor										
	A	ADTSKNTAYLQMN	SL	RA	EDTA	VYYC	SR			SEQ ID NO.: 123, 851, 870 & 889
	B	ADTSKNTAYLQMN	SL	RA	EDTA	VYYC	SR			SEQ ID NO.: 124, 852, 871 & 890
	C	ADTSKNTAYLQMN	SL	RA	EDTA	VYYC	SR			SEQ ID NO.: 125, 853, 872 & 891
Second Acceptor										
	A	ADTSKNTAYLQMN	SL	RA	EDTA	VYYCAR				SEQ ID NO.: 126, 854, 873 & 892
	B	ADTSKNTAYLQMN	SL	RA	EDTA	VYYCAR				SEQ ID NO.: 127, 855, 874 & 893
	C	ADTSKNTAYLQMN	SL	RA	EDTA	VYYCA				SEQ ID NO.: 128, 856, 875 & 894
	D	ADTSKNTAYLQMN	SL	RA	EDTA	VYYC				SEQ ID NO.: 129, 857, 876 & 895

FIG. 4B

P2260R1

9 / 45




FIG. 5

P2260R1

10 / 45

Framework Sequences of huMAb4D5-8 Light Chain

- LC-FR1 ¹Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
Asp Arg Val Thr Ile Thr Cys²³ (SEQ ID NO: 134)
- LC-FR2 ³⁵Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr⁴⁹
(SEQ ID NO: 135)
- LC-FR3 ⁵⁷Gly Val Pro Ser Arg Phe Ser Gly Ser Arg Ser Gly Thr Asp Phe Thr
Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys⁸⁸
(SEQ ID NO: 136)
- LC-FR4 ⁹⁸Phe Gly Gln Gly Thr Lys Val Glu Ile Lys¹⁰⁷ (SEQ ID NO: 137)

Framework Sequences of huMAb4D5-8 Heavy Chain

- HC-FR1 ¹Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
Ser Leu Arg Leu Ser Cys Ala Ala Ser²⁵ (SEQ ID NO: 138)
- HC-FR2 ³⁶Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val⁴⁸
(SEQ ID NO: 139)
- HC-FR3 ⁶⁶Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr Leu Gln
Met Asn⁸³ Ser^{83a} Leu^{83b} Arg^{83c} Ala Glu Asp Thr Ala Val Tyr Tyr Cys⁹²
(SEQ ID NO: 140)
- HC-FR4 ¹⁰³Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser¹¹³ (SEQ ID NO: 141)

FIG. 6Framework Sequences of huMAb4D5-8 Light Chain Modified at Position 66 and 99 (Underlined)

- LC-FR1 ¹Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
Asp Arg Val Thr Ile Thr Cys²³ (SEQ ID NO: 142)
- LC-FR2 ³⁵Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr⁴⁹
(SEQ ID NO: 143)
- LC-FR3 ⁵⁷Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys⁸⁸
(SEQ ID NO: 144)
- LC-FR4 ⁹⁸Phe Arg Gln Gly Thr Lys Val Glu Ile Lys¹⁰⁷ (SEQ ID NO: 145)

Framework Sequences of huMAb4D5-8 Heavy Chain Modified at Postions 71, 73 and 78 (Underlined)

- HC-FR1 ¹Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
Ser Leu Arg Leu Ser Cys Ala Ala Ser²⁵ (SEQ ID NO: 146)
- HC-FR2 ³⁶Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val⁴⁸
(SEQ ID NO: 147)
- HC-FR3 ⁶⁶Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr Leu Gln
Met Asn⁸³ Ser^{83a} Leu^{83b} Arg^{83c} Ala Glu Asp Thr Ala Val Tyr Tyr
Cys⁹² (SEQ ID NO: 148)
- HC-FR4 ¹⁰³Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser¹¹³ (SEQ ID NO: 149)

FIG. 7

P2260R1

11 / 45

HVR-H1

Clone #	SEQ ID NO:										
		26	27	28	29	30	31	32	33	34	35
48-26	151	G	F	N	V	Y	Y	S	S	I	H
48-27	152	G	F	N	V	S	Y	S	Y	M	H
48-28	153	G	F	N	F	S	Y	Y	S	M	H
48-29	154	G	F	N	L	S	Y	Y	S	I	H
48-30	155	G	F	N	I	S	Y	S	S	M	H
48-31	156	G	F	N	V	Y	Y	S	S	I	H
48-32	157	G	F	N	L	S	Y	S	Y	I	H
48-33	158	G	F	N	F	Y	Y	Y	Y	I	H
48-34	159	G	F	N	L	S	Y	S	S	I	H
48-35	160	G	F	N	V	S	Y	S	S	I	H
48-36	161	G	F	N	V	S	Y	S	S	I	H
48-37	162	G	F	N	V	Y	Y	S	S	I	H
48-38	163	G	F	N	V	S	Y	Y	Y	I	H
48-39	164	G	F	N	L	S	Y	S	S	I	H
48-40	165	G	F	N	I	S	Y	S	Y	M	H
48-41	166	G	F	N	L	Y	Y	S	Y	M	H
48-42	167	G	F	N	V	S	Y	Y	Y	M	H
48-43	168	G	F	N	I	S	Y	S	Y	M	H
48-44	169	G	F	N	I	S	Y	S	S	I	H
48-45	170	G	F	N	V	S	Y	S	S	M	H
48-46	171	G	F	N	L	S	Y	Y	S	I	H
48-47	172	G	F	N	V	S	Y	Y	S	I	H
48-48	173	G	F	N	I	S	Y	S	S	I	H
48-49	174	G	F	N	F	S	Y	Y	S	I	H
48-50	175	G	F	N	L	S	Y	S	S	M	H
Consensus	176	G	F	N	V/F/ L/I	Y/S	Y	S/Y	S/Y	I/M	H

FIG. 8A

HVR-H2

Clone #	SEQ ID NO:																	
		50	51	52	52a	53	54	55	56	57	58	59	60	61	62	63	64	65
48-26	177	S	I	S	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-27	178	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-28	179	S	I	Y	P	Y	Y	G	Y	T	Y	Y	A	D	S	V	K	G
48-29	180	S	I	Y	P	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
48-30	181	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-31	182	S	I	Y	P	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
48-32	183	S	I	S	S	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
48-33	184	S	I	S	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-34	185	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-35	186	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-36	187	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-37	188	S	I	Y	P	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
48-38	189	S	I	S	S	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
48-39	190	S	I	Y	S	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
48-40	191	S	I	S	P	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
48-41	192	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-42	193	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-43	194	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-44	195	S	I	S	S	S	Y	S	S	T	Y	Y	A	D	S	V	K	G
48-45	196	S	I	S	P	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
48-46	197	S	I	Y	S	S	Y	S	S	T	Y	Y	A	D	S	V	K	G
48-47	198	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-48	199	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-49	200	S	I	Y	P	Y	Y	S	Y	T	Y	Y	A	D	S	V	K	G
48-50	201	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
Consensus	202	S	I	S/Y	P/S	Y/S	Y	S/G	Y/S	T	S/Y	Y	A	D	S	V	K	G

FIG. 8B

P2260R1

13 / 45

HVR-H3

Clone #	SEQ ID NO:											
		95	96	97	98	99	100	100a	100b	100c	101	102
48-26	203	E	G	Y	S	Q	G	G	F		D	Y
48-27	204	S	Y	S	Y	Y	S	A	I		D	Y
48-28	205	S	Y	S	Y	S	Y	G	M		D	Y
48-29	206	S	Y	S	Y	S	Y	G	I		D	Y
48-30	207	S	Y	S	Y	Y	S	A	M		D	Y
48-31	208	G	Y	K	Y	W	S	A	F		D	Y
48-32	209	E	S	F	Y	Y	S	P	A	F	D	Y
48-33	210	E	Y	Y	S	Y	L	G	A	I	D	Y
48-34	211	G	Y	E	G	G	M	A	M		D	Y
48-35	212	S	Y	S	Y	S	S	G	L		D	Y
48-36	213	G	Y	M	W	Y	G	G	I		D	Y
48-37	214	D	C	Y	Y	x	A	A	F		D	Y
48-38	215	E	N	Y	W	W	A	I			D	Y
48-39	216	S	Y	S	Y	Y	S	A	F		D	Y
48-40	217	D	Y	Y	F	F	S	A	I		D	Y
48-41	218	S	Y	S	Y	S	S	A	L		D	Y
48-42	219	E	G	Y	I	S	G	D	A	I	D	Y
48-43	220	S	Y	S	S	Y	S	A	I		D	Y
48-44	221	G	Y	F	E	G	W	Y	G	L	D	Y
48-45	222	E	Y	S	Y	Y	G	G	F		D	Y
48-46	223	E	S	Y	W	S	Y	A	M		D	Y
48-47	224	S	Y	S	Y	S	Y	G	L		D	Y
48-48	225	Y	Y	S	Y	S	S	G	L		D	Y
48-49	226	S	Y	S	Y	S	Y	G	L		D	Y
48-50	227	S	Y	S	Y	S	Y	G	M		D	Y
Con-sensus	228	E/S/ G/D/ Y	G/Y/ S/C/ N	Y/S/ K/F/ E/M	S/Y/ G/W/ F/I/E	Q/Y/ S/W/ G/X/ F	G/S/ Y/L/ M/G/ A/W	G/A P/I/ D/Y	F/I/ M/A/ L/G/ n.p.	F/I/ L/ n.p.	D	Y

FIG. 8C

14 / 45

[illegible]

P2260R1

HVR-H3

Clone #	SEQ ID NO:	95	96	97	98	99	100	100a	100b	100c	100d	100e	100f	100g	100h	100i	100j	100k	100l	100m	101	102
63-10	253	E	Y	Y	R	W	Y	T	A	I											D	Y
63-11	254	S	G	Y	Y	Y	Q	G	Y	W	W	Y	Y	Y	T	G	Y	Y	G	M	D	Y
63-12	255	G	I	M	F	S	S	W	W	W	Y	Y	D	Y	S	D	A	L			D	Y
63-13	256	G	S	I	P	S	Y	W	S	A	D	W	Y	Y	Y	Y	G	L			D	Y
63-14	257	E	Y	Y	W	W	Y	K	E	A	W	Y	S	A	G	M					D	Y
63-15	258	W	Q	G	Y	G	F	K	Y	Y	W	S	Y	Y	V	S	Y	G	G	L	D	Y
63-16	259	S	Y	S	Y	Y	Y	Y	S	S	Y	G	F								D	Y
63-17	260	E	S	Y	A	G	V	P	P	Y	G	F									D	Y
63-18	261	G	I	M	F	S	S	W	W	W	Y	Y	D	Y	S	D	A	L			D	Y
63-19	262	G	I	M	F	S	S	W	W	W	Y	Y	D	Y	S	D	A	L			D	Y
63-20	263	E	Y	Y	W	W	Y	K	E	A	W	Y	S	A	G	M					D	Y
Con-sensus	264	E/S/ G/W	Y/ G/I/ S/G	Y/M/ I/Y/ G/S	R/Y/ F/P/W	W/Y/ S/G	Y/Q/ S/F/V	T/G/ W/K/ Y/P	A/Y/ W/S/ E/P	I/W/ A/Y/ S	W/Y/ D/G	Y/W/ S/G F	Y/D/ S/F/ n.p.	Y/A/ n.p.	T/S/ Y/G/ V/ n.p.	G/D/ Y/M/ S/ n.p.	Y/A/ G/ n.p.	Y/L/ G/ n.p.	G/ n.p.	M/L/ n.p.	D	Y

FIG. 9B

P2260R1

16 / 45

HVR-H1

Clone #	SEQ ID NO:										
		26	27	28	29	30	31	32	33	34	35
apu01	265	G	F	N	V	Y	Y	S	S	I	H
apu02	266	G	F	N	V	S	Y	S	Y	M	H
apu03	267	G	F	N	F	S	Y	Y	S	M	H
apu04	268	G	F	N	L	S	Y	Y	S	I	H
apu05	269	G	F	N	I	S	Y	S	S	M	H
apu06	270	G	F	N	V	Y	Y	S	S	I	H
apu07	271	G	F	N	L	S	Y	S	Y	I	H
apu08	272	G	F	N	L	S	Y	S	S	M	H
apu09	273	G	F	N	V	S	Y	S	S	I	H
apu10	274	G	F	N	V	S	Y	S	S	I	H
apu11	275	G	F	N	V	S	Y	Y	Y	I	H
apu12	276	G	F	N	L	Y	Y	S	Y	M	H
apu13	277	G	F	N	V	S	Y	Y	S	I	H
apu14	278	G	F	N	I	S	Y	S	S	I	H
apu15	279	G	F	N	F	S	Y	Y	S	I	H
Consensus	280	G	F	N	V/F/L/I	Y/S	Y	S/Y	S/Y	I/M	H

HVR-H2

Clone #	SEQ ID NO:																	
		50	51	52	52a	53	54	55	56	57	58	59	60	61	62	63	64	65
apu01	281	S	I	S	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu02	282	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu03	283	S	I	Y	P	Y	Y	G	Y	T	Y	Y	A	D	S	V	K	G
apu04	284	S	I	Y	P	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
apu05	285	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu06	286	S	I	Y	P	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
apu07	287	S	I	S	S	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
apu08	288	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu09	289	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu10	290	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu11	291	S	I	S	S	Y	Y	G	Y	T	S	Y	A	D	S	V	K	G
apu12	292	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu13	293	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu14	294	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu15	295	S	I	Y	P	Y	Y	S	Y	T	Y	Y	A	D	S	V	K	G
Consensus	296	S	I	S/Y	P/S	Y	Y	S/G	Y/S	T	S/Y	Y	A	D	S	V	K	G

FIG. 10A

HVR-H3

Clone #	SEQ ID NO:											
		95	96	97	98	99	100	100a	100b	100c	101	102
apu01	297	E	G	Y	S	Q	G	G	F		D	Y
apu02	298	S	Y	S	Y	Y	S	A	I		D	Y
apu03	299	S	Y	S	Y	S	Y	G	M		D	Y
apu04	300	S	Y	S	Y	S	Y	G	I		D	Y
apu05	301	S	Y	S	Y	Y	S	A	M		D	Y
apu06	302	G	Y	K	Y	W	S	A	F		D	Y
apu07	303	E	S	F	Y	Y	S	P	A	F	D	Y
apu08	304	G	Y	E	G	G	M	A	M		D	Y
apu09	305	S	Y	S	Y	S	S	G	L		D	Y
apu10	306	G	Y	M	W	Y	G	G	I		D	Y
apu11	307	E	N	Y	W	W	A	I			D	Y
apu12	308	S	Y	S	Y	S	S	A	L		D	Y
apu13	309	S	Y	S	Y	S	Y	G	L		D	Y
apu14	310	Y	Y	S	Y	S	S	G	L		D	Y
apu15	311	S	Y	S	Y	S	Y	G	L		D	Y
Con-sensus	312	E/S/ G/Y	G/Y/ S/N	Y/S/ K/F/E	S/Y/ G/W	Q/Y/ S/G	G/S/ Y/M/ A	G/A/ P/I	F/I/ M/A/ L/n.p.	F/ n.p.	D	Y

FIG. 10B

HVR-L3

Clone #	SEQ ID NO:											
		89	90	91	92	93	94	95	95a	95b	96	97
apu01	313	Q	Q	S	S	Y	S	S	L		F	T
apu02	314	Q	Q	S	S	Y	S	S	L		F	T
apu03	315	Q	Q	Y	S	S	Y	S	S	P	I	T
apu04	316	Q	Q	Y	S	S	Y	Y	S	P	V	T
apu05	317	Q	Q	S	S	Y	S	S	L		I	T
apu06	318	Q	Q	S	S	Y	S	S	L		V	T
apu07	319	Q	Q	S	Y	Y	Y	S	L		F	T
apu08	320	Q	Q	S	S	Y	S	S	L		V	T
apu09	321	Q	Q	S	S	Y	S	S	L		F	T
apu10	322	Q	Q	Y	S	Y	S	S	L		F	T
apu11	323	Q	Q	S	Y	Y	Y	Y	P		I	T
apu12	324	Q	Q	S	S	Y	S	S	L		V	T
apu13	325	Q	Q	Y	S	S	S	Y	Y	P	F	T
apu14	326	Q	Q	S	S	Y	S	S	L		L	T
apu15	327	Q	Q	Y	Y	Y	Y	Y	Y	P	I	T
Con-sensus	328	Q	Q	S/Y	S/Y	Y/S	S/Y	S/Y	L/S/ P/Y	P/ n.p.	F/I/ V/L	T

FIG. 10C

P2260R1

18 / 45

THE

[illegible]

2-H-2

[illegible]

FIG. 11A

P2260R1

20 / 45






FIG. 12

P2260R1

21 / 45

**FIG. 13A****FIG. 13B**

P2260R1

22 / 45

HVR-H1

Clone #	SEQ ID NO:										
		26	27	28	29	30	31	32	33	34	35
48-51	392	G	F	N	I	S	Y	S	S	M	H
48-52	393	G	F	N	I	S	Y	S	S	M	H
48-53	394	G	F	N	I	S	Y	S	S	M	H
48-54	395	G	F	N	I	S	Y	S	S	M	H
48-55	396	G	F	N	I	G	Y	S	F	M	H
48-56	397	G	F	N	V	D	Y	S	Y	M	H
48-57	398	G	F	N	V	D	Y	S	Y	M	H
48-58	399	G	F	N	F	S	Y	S	F	M	H
48-59	400	G	F	N	I	V	Y	S	F	M	H
48-60	401	G	F	N	I	I	Y	S	F	M	H
48-61	402	G	F	N	I	V	Y	S	F	I	H
48-62	403	G	F	N	L	S	Y	S	F	M	H
48-63	404	G	F	N	V	D	Y	S	F	M	H
48-64	405	G	F	N	V	I	Y	S	F	M	H
48-65	406	G	F	N	V	A	Y	S	L	M	H
48-66	407	G	F	N	I	S	Y	S	W	M	H
48-67	408	G	F	N	L	D	Y	S	F	M	H
48-68	409	G	F	N	F	L	Y	S	G	I	H
48-69	410	G	F	N	I	S	Y	S	S	M	H
48-70	411	G	F	N	I	S	Y	S	S	M	H
48-71	412	G	F	N	I	S	Y	S	S	M	H
48-72	413	G	F	N	I	S	Y	S	S	M	H
48-73	414	G	F	N	I	S	Y	S	S	M	H
48-74	415	G	F	N	I	L	Y	S	G	I	H
48-75	416	G	F	N	I	S	Y	S	S	M	H
48-76	417	G	F	N	I	S	Y	S	S	M	H
48-77	418	G	F	N	I	S	Y	S	S	M	H
48-78	419	G	F	N	I	S	Y	S	S	M	H
48-79	420	G	F	N	I	S	Y	S	S	M	H
48-80	421	G	F	N	I	S	Y	S	S	M	H
48-81	422	G	F	N	I	S	Y	S	S	M	H
48-82	423	G	F	N	I	S	Y	S	S	M	H
48-83	424	G	F	N	I	S	Y	S	S	M	H
48-84	425	G	F	N	I	S	Y	S	S	M	H
48-85	426	G	F	N	I	S	Y	S	S	M	H
48-86	427	G	F	N	I	F	Y	S	G	I	H
48-87	428	G	F	N	I	S	Y	S	S	M	H
48-88	429	G	F	N	I	S	Y	S	S	M	H
48-89	430	G	F	N	I	S	Y	S	S	M	H
48-90	431	G	F	N	I	S	Y	S	S	M	H
48-91	432	G	F	N	I	S	Y	S	S	M	H
48-92	433	G	F	N	I	S	Y	S	S	M	H
48-93	434	G	F	N	I	S	Y	S	S	M	H
48-94	435	G	F	N	I	S	Y	S	S	M	H
48-95	436	G	F	N	I	S	Y	S	S	M	H
48-96	437	G	F	N	I	S	Y	S	S	M	H

FIG. 14A

P2260R1

23 / 45

HVR-H1 (Con't.)

Clone #	SEQ ID NO:										
		26	27	28	29	30	31	32	33	34	35
48-97	438	G	F	N	I	S	Y	S	S	M	H
48-98	439	G	F	N	I	S	Y	S	S	M	H
48-99	440	G	F	N	I	S	Y	S	S	M	H
48-100	441	G	F	N	I	S	Y	S	S	M	H
48-101	442	G	F	N	I	S	Y	S	S	M	H
48-102	443	G	F	N	I	S	Y	S	S	M	H
48-103	444	G	F	N	I	S	Y	S	S	M	H
48-104	445	G	F	N	I	S	Y	S	S	M	H
48-105	446	G	F	N	I	S	Y	S	S	M	H
48-106	447	G	F	N	I	S	Y	S	S	M	H
48-107	448	G	F	N	I	S	Y	S	S	M	H
48-108	449	G	F	N	I	S	Y	S	S	M	H
48-109	450	G	F	N	I	S	Y	S	S	M	H
48-110	451	G	F	N	I	S	Y	S	S	M	H
48-111	452	G	F	N	I	S	Y	S	S	M	H
48-112	453	G	F	N	L	S	Y	S	G	M	H
48-113	454	G	F	N	L	L	Y	S	G	M	H
48-114	455	G	F	N	V	A	Y	S	G	I	H
48-115	456	G	F	N	V	D	Y	S	G	M	H
48-116	457	G	F	N	V	D	Y	S	G	M	H
48-117	458	G	F	N	V	S	Y	S	S	I	H
48-118	459	G	F	N	V	V	Y	S	G	I	H
Consensus	460	G	F	N	I/V/ F/L	S/G/ D/V/ I/L/ F/A	Y	S	S/F/ Y/L/ W/G	M/I	H

FIG. 14B

HVR-H2

Clone #	SEQ ID NO:																	
		50	51	52	52a	53	54	55	56	57	58	59	60	61	62	63	64	65
48-51	461	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-52	462	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-53	463	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-54	464	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-55	465	S	I	A	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-56	466	S	I	A	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-57	467	S	I	A	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-58	468	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-59	469	S	I	S	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-60	470	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-61	471	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-62	472	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-63	473	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-64	474	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-65	475	S	I	S	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-66	476	S	I	T	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-67	477	S	I	T	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-68	478	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-69	479	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-70	480	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-71	481	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-72	482	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-73	483	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-74	484	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-75	485	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-76	486	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-77	487	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-78	488	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-79	489	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-80	490	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-81	491	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-82	492	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-83	493	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-84	494	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-85	495	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-86	496	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-87	497	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-88	498	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-89	499	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-90	500	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-91	501	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-92	502	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-93	503	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G

FIG. 14C

P2260R1

25 / 45

Clone #	SEQ ID NO:																	
		50	51	52	52a	53	54	55	56	57	58	59	60	61	62	63	64	65
48-94	504	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-95	505	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-96	506	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-97	507	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-98	508	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-99	509	S	I	Y	P	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-100	510	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-101	511	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-102	512	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-103	513	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-104	514	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-105	515	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-106	516	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-107	517	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-108	518	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-109	519	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-110	520	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-111	521	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-112	522	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-113	523	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-114	524	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-115	525	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-116	526	S	I	Y	S	Y	Y	T	Y	T	S	Y	A	D	S	V	K	G
48-117	527	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
48-118	528	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
Con-sensus	529	S	I	Y/A/S/T	S/P	Y	Y	S/T	Y	T	S	Y	A	D	S	V	K	G

FIG. 14D

P2260R1

26 / 45

HVR-H3

Clone #	SEQ ID NO:										
		95	96	97	98	99	100	100a	100b	101	102
48-51	530	S	Y	N	N	T	T	S	I	D	Y
48-52	531	G	Y	S	W	Y	N	A	M	D	Y
48-53	532	G	Y	S	W	F	N	A	I	D	Y
48-54	533	G	Y	Y	W	F	D	A	M	D	Y
48-55	534	S	Y	S	Y	Y	S	A	M	D	Y
48-56	535	S	Y	S	Y	R	E	T	M	D	Y
48-57	536	S	Y	S	Y	R	E	T	M	D	Y
48-58	537	S	Y	S	Y	Y	S	A	M	D	Y
48-59	538	S	Y	S	Y	Y	S	A	M	D	Y
48-60	539	S	Y	S	Y	Y	S	A	M	D	Y
48-61	540	S	Y	S	Y	Y	S	A	M	D	Y
48-62	541	S	Y	S	Y	Y	S	A	M	D	Y
48-63	542	S	Y	S	Y	Y	S	A	M	D	Y
48-64	543	S	Y	S	Y	Y	S	A	M	D	Y
48-65	544	S	Y	S	Y	Y	S	A	M	D	Y
48-66	545	S	Y	S	Y	Y	S	A	M	D	Y
48-67	546	S	Y	S	Y	Y	S	A	M	D	Y
48-68	547	S	Y	S	Y	Y	S	A	M	D	Y
48-69	548	S	Y	S	Y	Y	S	A	M	D	Y
48-70	549	S	Y	S	Y	S	F	G	M	D	Y
48-71	550	S	Y	S	Y	F	M	G	M	D	Y
48-72	551	S	Y	S	Y	H	V	A	F	D	Y
48-73	552	S	Y	S	Y	Y	S	A	M	D	Y
48-74	553	S	Y	S	Y	Y	S	A	M	D	Y
48-75	554	S	Y	S	Y	H	L	A	F	D	Y
48-76	555	S	Y	S	Y	S	L	A	F	D	Y
48-77	556	S	Y	S	Y	Y	I	A	M	D	Y
48-78	557	S	Y	S	Y	Y	M	G	M	D	Y
48-79	558	S	Y	S	Y	S	M	G	M	D	Y
48-80	559	S	Y	S	Y	H	V	A	M	D	Y
48-81	560	S	Y	S	Y	H	M	G	M	D	Y
48-82	561	S	Y	S	Y	H	L	G	M	D	Y
48-83	562	S	Y	S	Y	Y	Q	G	F	D	Y
48-84	563	S	Y	S	Y	S	M	G	M	D	Y
48-85	564	S	Y	S	Y	F	L	A	M	D	Y
48-86	565	S	Y	S	Y	Y	S	A	M	D	Y
48-87	566	S	Y	S	Y	S	E	A	L	D	Y
48-88	567	S	Y	S	Y	S	L	G	M	D	Y
48-89	568	S	Y	S	Y	Y	S	A	M	D	Y
48-90	569	S	Y	S	Y	F	M	G	M	D	Y
48-91	570	S	Y	S	Y	F	L	A	M	D	Y
48-92	571	S	Y	S	Y	F	L	A	M	D	Y
48-93	572	S	Y	S	Y	Y	L	A	M	D	Y
48-94	573	S	Y	S	Y	F	I	G	M	D	Y
48-95	574	S	Y	S	Y	H	L	G	M	D	Y

FIG. 14E

P2260R1

27 / 45

Clone #	SEQ ID NO:										
		95	96	97	98	99	100	100a	100b	101	102
48-96	575	S	Y	S	Y	T	E	A	F	D	Y
48-97	576	S	Y	S	Y	Y	S	A	M	D	Y
48-98	577	S	Y	S	Y	T	Y	G	L	D	Y
48-99	578	S	Y	S	Y	Y	S	A	M	D	Y
48-100	579	S	Y	S	Y	S	L	G	M	D	Y
48-101	580	S	Y	S	Y	W	V	G	M	D	Y
48-102	581	S	Y	S	Y	T	L	A	M	D	Y
48-103	582	S	Y	S	Y	T	M	G	L	D	Y
48-104	583	S	Y	S	Y	F	L	G	M	D	Y
48-105	584	S	Y	S	Y	H	M	G	F	D	Y
48-106	585	S	Y	S	Y	S	L	G	M	D	Y
48-107	586	S	Y	S	Y	Y	E	A	F	D	Y
48-108	587	S	Y	S	Y	R	M	A	F	D	Y
48-109	588	S	Y	S	Y	H	I	A	F	D	Y
48-110	589	S	Y	S	Y	S	V	G	M	D	Y
48-111	590	S	Y	S	Y	T	L	G	M	D	Y
48-112	591	S	Y	S	Y	Y	S	A	M	D	Y
48-113	592	S	Y	S	Y	Y	S	A	M	D	Y
48-114	593	S	Y	S	Y	Y	S	A	M	D	Y
48-115	594	S	Y	S	Y	Y	S	A	M	D	Y
48-116	595	S	Y	S	Y	Y	S	A	M	D	Y
48-117	596	S	Y	S	Y	Y	S	A	M	D	Y
48-118	597	S	Y	S	Y	Y	S	A	M	D	Y
Consensus	598	S/G	Y	S/N/Y	Y/N/ W	T/Y/ F/R/ S/H/ W	T/N/ D/S/ E/F/ M/Y/ L/I/Y	S/A/ T/G	I/M/ F/L	D	Y

FIG. 14F

P2260R1

28 / 45

HVR-H1

Clone #	SEQ ID NO:										
		26	27	28	29	30	31	32	33	34	35
63-21	599	G	F	N	I	S	S	S	Y	I	H
63-22	600	G	F	N	I	S	S	S	Y	I	H
63-23	601	G	F	N	I	S	S	S	Y	I	H
63-24	602	G	F	N	I	S	S	S	Y	I	H
63-25	603	G	F	N	I	S	S	S	Y	I	H
63-26	604	G	F	N	I	S	S	S	Y	I	H
63-27	605	G	F	N	I	S	S	S	Y	I	H
63-28	606	G	F	N	I	S	S	S	Y	I	H
63-29	607	G	F	N	I	S	S	S	Y	I	H
63-30	608	G	F	N	I	S	S	S	Y	I	H
63-31	609	G	F	N	I	S	S	S	Y	I	H
63-32	610	G	F	N	I	S	S	S	Y	I	H
63-33	611	G	F	N	I	S	S	S	Y	I	H
63-34	612	G	F	N	I	S	S	S	Y	I	H
63-35	613	G	F	N	F	S	S	S	Y	I	H
63-36	614	G	F	N	I	K	G	S	L	I	H
63-37	615	G	F	N	I	K	G	S	I	M	H
63-38	616	G	F	N	I	K	S	S	I	M	H
63-39	617	G	F	N	I	S	S	S	Y	I	H
63-40	618	G	F	N	I	S	S	S	Y	I	H
63-41	619	G	F	N	I	S	S	S	Y	I	H
63-42	620	G	F	N	I	S	S	S	Y	I	H
63-43	621	G	F	N	I	S	S	S	Y	I	H
63-44	622	G	F	N	I	S	S	S	Y	I	H
63-45	623	G	F	N	I	S	S	S	Y	I	H
63-46	624	G	F	N	I	S	S	S	Y	I	H
63-47	625	G	F	N	L	A	S	S	F	M	H
63-48	626	G	F	N	L	V	S	S	L	M	H
63-49	627	G	F	N	V	K	T	G	L	I	H
63-50	628	G	F	N	V	K	W	N	Y	I	H
63-51	629	G	F	N	V	V	S	S	F	I	H
Consensus	630	G	F	N	I/F/ L/V	S/K/ A/V	S/G/ T/W	S/G/N	Y/L/ I/F	I/M	H

FIG. 15A

P2260R1

29 / 45

HVR-H2

Clone #	SEQ ID NO:																	
		50	51	52	52a	53	54	55	56	57	58	59	60	61	62	63	64	65
63-21	631	A	I	A	P	Y	L	G	S	T	S	Y	A	D	S	V	K	G
63-22	632	A	I	P	P	F	Y	G	W	T	S	Y	A	D	S	V	K	G
63-23	633	A	I	Q	P	Y	F	G	W	T	I	Y	A	D	S	V	K	G
63-24	634	A	I	S	P	Y	L	G	S	T	S	Y	A	D	S	V	K	G
63-25	635	D	I	A	P	Y	L	G	T	T	K	Y	A	D	S	V	K	G
63-26	636	D	I	S	P	W	Y	G	G	T	S	Y	A	D	S	V	K	G
63-27	637	D	I	S	S	Y	T	G	S	T	D	Y	A	D	S	V	K	G
63-28	638	F	I	Q	P	Y	Y	G	S	T	I	Y	A	D	S	V	K	G
63-29	639	F	I	S	P	Y	L	G	S	T	N	Y	A	D	S	V	K	G
63-30	640	G	I	T	P	Y	L	G	W	T	S	Y	A	D	S	V	K	G
63-31	641	H	I	S	P	Y	L	G	S	T	S	Y	A	D	S	V	K	G
63-32	642	I	I	S	P	Y	L	G	S	T	G	Y	A	D	S	V	K	G
63-33	643	S	I	T	P	Y	Y	G	W	T	R	Y	A	D	S	V	K	G
63-34	644	W	I	S	P	Y	L	G	R	T	S	Y	A	D	S	V	K	G
63-35	645	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-36	646	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-37	647	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-38	648	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-39	649	Y	I	G	P	F	T	G	S	T	N	Y	A	D	S	V	K	G
63-40	650	Y	I	S	P	F	L	S	T	T	S	Y	A	D	S	V	K	G
63-41	651	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-42	652	Y	I	S	P	Y	S	G	S	T	K	Y	A	D	S	V	K	G
63-43	653	Y	I	S	P	Y	Y	S	S	T	S	Y	A	D	S	V	K	G
63-44	654	Y	I	S	P	Y	L	G	S	T	S	Y	A	D	S	V	K	G
63-45	655	Y	I	S	P	Y	L	S	S	T	S	Y	A	D	S	V	K	G
63-46	656	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-47	657	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-48	658	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-49	659	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-50	660	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
63-51	661	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
Con-sensus	662	A/D/ F/G/ H/I/ S/W/ Y	I	A/P/ Q/S/ T/G	P/S	Y/F	L/Y/ F/T	G/S	S/ W/T/ G/R	T	S/I/ K/D/ N/G/ R	Y	A	D	S	V	K	G

FIG. 15B

P2260R1

30 / 45

HVR-H3

Clone #	SEQ ID NO:											
		95	96	97	98	99	100	100a	100b	100c	101	102
63-21	663	E	Y	Y	R	W	Y	T	A	I	D	Y
63-22	664	E	Y	Y	R	W	Y	T	A	I	D	Y
63-23	665	E	Y	Y	R	W	Y	T	A	I	D	Y
63-24	666	E	Y	Y	R	W	Y	T	A	I	D	Y
63-25	667	E	Y	Y	R	W	Y	T	A	I	D	Y
63-26	668	E	Y	Y	R	W	Y	T	A	I	D	Y
63-27	669	E	Y	Y	R	W	Y	T	A	I	D	Y
63-28	670	E	Y	Y	R	W	Y	T	A	I	D	Y
63-29	671	E	Y	Y	R	W	Y	T	A	I	D	Y
63-30	672	E	Y	Y	R	W	Y	T	A	I	D	Y
63-31	673	E	Y	Y	R	W	Y	T	A	I	D	Y
63-32	674	E	Y	Y	R	W	Y	T	A	I	D	Y
63-33	675	E	Y	Y	R	W	Y	T	A	I	D	Y
63-34	676	E	Y	Y	R	W	Y	T	A	I	D	Y
63-35	677	E	Y	Y	R	W	Y	T	A	I	D	Y
63-36	678	E	Y	Y	R	W	Y	T	A	I	D	Y
63-37	679	E	Y	Y	R	W	Y	T	A	I	D	Y
63-38	680	E	Y	Y	R	W	Y	T	A	I	D	Y
63-39	681	E	Y	Y	R	W	Y	T	A	I	D	Y
63-40	682	E	Y	Y	R	W	Y	T	A	I	D	Y
63-41	683	E	Y	Y	R	W	Y	T	A	I	D	Y
63-42	684	E	Y	Y	R	W	Y	T	A	I	D	Y
63-43	685	E	Y	Y	R	W	Y	T	A	I	D	Y
63-44	686	E	Y	Y	R	W	Y	T	A	I	D	Y
63-45	687	E	Y	Y	R	W	Y	T	A	I	D	Y
63-46	688	E	Y	Y	R	W	Y	T	A	I	D	Y
63-47	689	E	Y	Y	R	W	Y	T	A	I	D	Y
63-48	690	E	Y	Y	R	W	Y	T	A	I	D	Y
63-49	691	E	Y	Y	R	W	Y	T	A	I	D	Y
63-50	692	E	Y	Y	R	W	Y	T	A	I	D	Y
63-51	693	E	Y	Y	R	W	Y	T	A	I	D	Y
Consensus	694	E	Y	Y	R	W	Y	T	A	I	D	Y

FIG. 15C

31 / 45

HVR-H1

Clone #	SEQ ID NO:										
		26	27	28	29	30	31	32	33	34	35
apu2.01	695	G	F	N	I	S	Y	S	S	M	H
apu2.02	696	G	F	N	I	F	Y	G	G	I	H
apu2.03	697	G	F	N	I	S	Y	S	S	M	H
apu2.04	698	G	F	N	I	S	Y	S	S	M	H
apu2.05	699	G	F	N	I	S	Y	S	S	M	H
apu2.06	700	G	F	N	I	S	Y	S	S	M	H
apu2.07	701	G	F	N	I	S	Y	S	S	M	H
apu2.08	702	G	F	N	I	S	Y	S	S	M	H
apu2.09	703	G	F	N	I	S	Y	S	S	M	H
apu2.10	704	G	F	N	I	S	Y	S	S	M	H
Consensus	705	G	F	N	I	S/F	Y	S/G	S/G	M/I	H

HVR-H2

Clone #	SEQ ID NO:																	
		50	51	52	52a	53	54	55	56	57	58	59	60	61	62	63	64	65
apu2.01	706	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu2.02	707	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu2.03	708	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu2.04	709	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu2.05	710	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu2.06	711	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu2.07	712	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu2.08	713	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu2.09	714	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
apu2.10	715	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G
Consensus	716	S	I	Y	S	Y	Y	S	Y	T	S	Y	A	D	S	V	K	G

HVR-H3

Clone #	SEQ ID NO:										
		95	96	97	98	99	100	100a	100b	101	102
apu2.01	717	S	Y	S	Y	S	E	A	L	D	Y
apu2.02	718	S	Y	S	Y	Y	S	A	M	D	Y
apu2.03	719	S	Y	S	Y	S	L	A	F	D	Y
apu2.04	720	S	Y	S	Y	S	F	G	M	D	Y
apu2.05	721	S	Y	S	Y	R	M	A	F	D	Y
apu2.06	722	G	Y	S	W	F	N	A	I	D	Y
apu2.07	723	S	Y	S	Y	H	L	G	M	D	Y
apu2.08	724	S	Y	S	Y	S	V	G	M	D	Y
apu2.09	725	S	Y	S	Y	H	V	A	F	D	Y
apu2.10	726	S	Y	S	Y	F	L	A	M	D	Y
Consensus	727	S/G	Y	S	Y/W	S/Y/ R/F/H	E/S/L/ F/M/N/V	A/G	L/M/ F/I	D	Y

FIG. 16A

P2260R1

32 / 45

HVR-L3

Clone #	SEQ ID NO:										
		89	90	91	92	93	94	95	95a	96	97
apu2.01	728	Q	Q	S	S	Y	S	S	L	I	T
apu2.02	729	Q	Q	S	S	Y	S	S	L	I	T
apu2.03	730	Q	Q	S	S	Y	S	S	L	I	T
apu2.04	731	Q	Q	S	S	Y	S	S	L	I	T
apu2.05	732	Q	Q	S	S	Y	S	S	L	I	T
apu2.06	733	Q	Q	S	S	Y	S	S	L	I	T
apu2.07	734	Q	Q	S	S	Y	S	S	L	I	T
apu2.08	735	Q	Q	S	S	Y	S	S	L	I	T
apu2.09	736	Q	Q	S	S	Y	S	S	L	I	T
apu2.10	737	Q	Q	S	S	Y	S	S	L	I	T
Con-sensus	738	Q	Q	S	S	Y	S	S	L	I	T

FIG. 16B

HVR-H1

Clone #	SEQ ID NO:										
		26	27	28	29	30	31	32	33	34	35
apu2.11	739	G	F	N	I	S	S	S	Y	I	H
apu2.12	740	G	F	N	I	S	S	S	Y	I	H
apu2.13	741	G	F	N	V	K	W	N	Y	I	H
apu2.14	742	G	F	N	I	S	S	S	Y	I	H
apu2.15	743	G	F	N	I	K	G	S	I	M	H
apu2.16	744	G	F	N	V	K	T	G	L	I	H
apu2.17	745	G	F	N	I	S	S	S	Y	I	H
apu2.18	746	G	F	N	L	V	S	S	L	M	H
apu2.19	747	G	F	N	V	V	S	S	F	I	H
apu2.20	748	G	F	N	I	S	S	S	Y	I	H
Consensus	749	G	F	N	I/V/L	S/K/V	S/W/G/T	S/N/G	Y/I/L/F	I/M	H

HVR-H2

Clone #	SEQ ID NO:																	
		50	51	52	52a	53	54	55	56	57	58	59	60	61	62	63	64	65
apu2.11	750	Y	I	S	P	Y	L	S	S	T	S	Y	A	D	S	V	K	G
apu2.12	751	F	I	S	P	Y	L	G	S	T	N	Y	A	D	S	V	K	G
apu2.13	752	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
apu2.14	753	D	I	A	P	Y	L	G	T	T	K	Y	A	D	S	V	K	G
apu2.15	754	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
apu2.16	755	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
apu2.17	756	H	I	S	P	Y	L	G	S	T	S	Y	A	D	S	V	K	G
apu2.18	757	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
apu2.19	758	Y	I	S	P	Y	Y	G	S	T	S	Y	A	D	S	V	K	G
apu2.20	759	A	I	Q	P	Y	F	G	W	T	I	Y	A	D	S	V	K	G
Consensus	760	Y/F/ D/H/ A	I	S/A/ Q	P	Y	L/Y/ F	S/ G	S/T/ W	T	S/N/ K/I	Y	A	D	S	V	K	G

HVR-H3

Clone #	SEQ ID NO:											
		95	96	97	98	99	100	100a	100b	100c	101	102
apu2.11	761	E	Y	Y	R	W	Y	T	A	I	D	Y
apu2.12	762	E	Y	Y	R	W	Y	T	A	I	D	Y
apu2.13	763	E	Y	Y	R	W	Y	T	A	I	D	Y
apu2.14	764	E	Y	Y	R	W	Y	T	A	I	D	Y
apu2.15	765	E	Y	Y	R	W	Y	T	A	I	D	Y
apu2.16	766	E	Y	Y	R	W	Y	T	A	I	D	Y
apu2.17	767	E	Y	Y	R	W	Y	T	A	I	D	Y
apu2.18	768	E	Y	Y	R	W	Y	T	A	I	D	Y
apu2.19	769	E	Y	Y	R	W	Y	T	A	I	D	Y
apu2.20	770	E	Y	Y	R	W	Y	T	A	I	D	Y
Consensus	771	E	Y	Y	R	W	Y	T	A	I	D	Y

FIG. 17A

P2260R1

34 / 45

HVR-L3

Clone #	SEQ ID NO:											
		89	90	91	92	93	94	95	95a	95b	96	97
apu2.11	772	Q	Q	Y	S	S	Y	S	S	L	F	T
apu2.12	773	Q	Q	Y	S	S	Y	S	S	L	F	T
apu2.13	774	Q	Q	Y	S	S	Y	S	S	L	F	T
apu2.14	775	Q	Q	Y	S	S	Y	S	S	L	F	T
apu2.15	776	Q	Q	Y	S	S	Y	S	S	L	F	T
apu2.16	777	Q	Q	Y	S	S	Y	S	S	L	F	T
apu2.17	778	Q	Q	Y	S	S	Y	S	S	L	F	T
apu2.18	779	Q	Q	Y	S	S	Y	S	S	L	F	T
apu2.19	780	Q	Q	Y	S	S	Y	S	S	L	F	T
apu2.20	781	Q	Q	Y	S	S	Y	S	S	L	F	T
Con-sensus	782	Q	Q	Y	S	S	Y	S	S	L	F	T

FIG. 17B

P2260R1

35 / 45




FIG. 18

P2260R1

36 / 45






FIG. 19A

FIG. 19B



P2260R1

37 / 45

**FIG. 20A****FIG. 20B**


P2260R1

38 / 45

**FIG. 21A****FIG. 21B**

P2260R1

39 / 45

**FIG. 22**

40 / 45

HVR-H1

Clone #	SEQ ID NO:										
		26	27	28	29	30	31	32	33	34	35
apu3.01	789	G	F	N	V	K	T	G	F	M	H
apu3.02	790	G	F	N	V	K	T	G	F	I	H
apu3.03	791	G	F	N	I	K	M	V	F	M	H
apu3.04	792	G	F	N	V	K	N	F	I	I	H
apu3.05	793	G	F	N	V	K	T	G	F	M	H
apu3.06	794	G	F	N	V	K	R	G	F	M	H
apu3.07	795	G	F	N	L	K	T	G	F	I	H
apu3.08	796	G	F	N	V	K	T	G	Y	M	H
apu3.09	797	G	F	N	V	K	T	G	L	I	H
apu3.10	798	G	F	N	V	M	I	G	I	I	H
apu3.11	799	G	F	N	I	K	T	G	F	M	H
apu3.12	ND										
Consensus	800	G	F	N	V/I/L	K/M	T/M/ N/R/I	G/V/ F	F/I/ Y/L	M/I	H

HVR-H2

Clone #	SEQ ID NO:																	
		50	51	52	52a	53	54	55	56	57	58	59	60	61	62	63	64	65
apu3.01	801	Y	I	S	P	Y	Y	G	W	T	R	Y	A	D	S	V	K	G
apu3.02	802	Y	I	S	P	Y	L	G	V	T	R	Y	A	D	S	V	K	G
apu3.03	803	Y	I	S	P	Y	D	G	S	T	N	Y	A	D	S	V	K	G
apu3.04	804	Y	I	I	P	Y	S	G	N	T	V	Y	A	D	S	V	K	G
apu3.05	805	Y	I	S	P	Y	S	G	R	T	R	Y	A	D	S	V	K	G
apu3.06	806	Y	I	S	P	Y	L	G	S	T	R	Y	A	D	S	V	K	G
apu3.07	807	Y	I	S	P	Y	W	G	S	T	T	Y	A	D	S	V	K	G
apu3.08	808	Y	I	S	P	Y	Y	G	S	T	R	Y	A	D	S	V	K	G
apu3.09	809	Y	I	S	P	Y	F	G	Y	T	S	Y	A	D	S	V	K	G
apu3.10	810	Y	I	I	P	Y	S	G	S	T	K	Y	A	D	S	V	K	G
apu3.11	811	Y	I	T	P	Y	W	G	S	T	K	Y	A	D	S	V	K	G
apu3.12	ND																	
Consensus	812	Y	I	S/I/T	P	Y	Y/L/ D/S/ W	G	W/V/ S/N/ R/Y	T	R/N/ V/T/ S/K	Y	A	D	S	V	K	G

FIG. 23A

P2260R1

41 / 45



HVR-H3

Clone #	SEQ ID NO:											
		95	96	97	98	99	100	100a	100b	100c	101	102
apu3.01	813	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.02	814	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.03	815	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.04	816	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.05	817	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.06	818	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.07	819	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.08	820	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.09	821	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.10	822	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.11	823	E	Y	Y	R	W	Y	T	A	I	D	Y
apu3.12	ND											
Con-sensus	824	E	Y	Y	R	W	Y	T	A	I	D	Y

FIG. 23B



P2260R1

42 / 45


**FIG. 24A****FIG. 24B**

P2260R1

43 / 45


**FIG. 24C****FIG. 24D**

44 / 45

**FIG. 25**



P2260R1

45 / 45



1 MET GLN ILE PHE VAL LYS THR LEU THR GLY LYS THR ILE THR
 15 LEU GLU VAL GLU PRO SER ASP THR ILE GLU ASN VAL LYS ALA
 29 LYS ILE GLN ASP LYS GLU GLY ILE PRO PRO ASP GLN GLN ARG
 43 LEU ILE PHE ALA GLY LYS GLN LEU GLU ASP GLY ARG THR LEU
 57 SER ASP TYR ASN ILE GLN LYS GLU SER THR LEU HIS LEU VAL
 71 LEU ARG LEU ARG GLY GLY

A



B