



US00PP11862P2

(12) **United States Plant Patent**
Roberson

(10) **Patent No.:** **US PP11,862 P2**

(45) **Date of Patent:** **May 8, 2001**

(54) **HEMEROCALLIS HYBRID PLANT NAMED**
'ROBBOBELI'

(76) Inventor: **Robert J. Roberson**, 31706 E. Pink
Hill Rd., Grain Valley, Jackson County,
MO (US) 64029

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/249,101**

(22) Filed: **Feb. 12, 1999**

(51) **Int. Cl.**⁷ **A01H 5/00**

(52) **U.S. Cl.** **Plt./312**

(58) **Field of Search** **Plt./312**

Primary Examiner—Bruce R. Campell

Assistant Examiner—Anne Marie Grünberg

(74) *Attorney, Agent, or Firm*—Rothwell, Figg, Ernst &
Manbeck

(57) **ABSTRACT**

The new and distinct daylily plant as substantially shown
and described, particularly characterized by its ability to
adapt, grow and perform well in a wide range of USDA
zones, it's compact growth habit, it's unique near-white
blooms, it's above average increase (bunch formation) rate
in it's color class, and it's tendency to rebloom late in the
season.

2 Drawing Sheets

1

SUMMARY OF THE INVENTION

This invention relates to a new and distinct daylily
distinguished by it's "near white" (so pale that it is visually
"white" to the eye) colored blooms with diamond dusting,
it's tendency to rebloom late in the season, it's ability to
adapt and perform in a wide range of USDA zones, it's
above average increase rate (bunch formation) in it's color
class, and it's compact growth habit.

BACKGROUND OF THE INVENTION

This new variety of daylily plant originated as a seedling
selected from a group of seedlings planted by me at my
ranch in Grain Valley, Mo., and resulted from my crossing
of the clonal cultivar 'Joan Senior' (not patented) with the
variety 'Sea Gull' (not patented). My objective was to
produce a plant having white blooms and compact growth
habit. A further objective was to produce such a plant that
would be both cold hardy, heat tolerant, have above-average
bunch formation rate and repeat blooming.

This new plant was selected for propagation because it
most closely possessed the advantageous features that I
sought and asexual reproduction of the selected plant was
accomplished by division at the crown or basal plate which
is the area where the roots and foliage meet.

During the course of propagation of this new plant at my
daylily ranch in Grain Valley, Mo., I have observed that the
new plant has the tendency to rebloom late in the season, has
unique "near white" blooms which abcess approximately
30% quicker than the average of other daylilies of this color
category, and has consistently compact growth habit. Fur-
thermore this plant has an above-average increase or bunch
formation rate (7 fans compared to 4 fans for the average
near-white daylily) in its color class, and is cold hardy and
particularly capable of adapting, growing and performing
well across five USDA zones (4-8) according to test data
from the All American Daylily Selection Council (AADSC).
The inventor has determined that the traits described and
illustrated herein are stable and firmly fixed. I have chosen
to identify this new cultivar as Hemerocallis 'Robbobelii'.
This cultivar is being marketed in the United States under
the name LADY ELIZABETH™.

2

BRIEF DESCRIPTION OF THE DRAWING

The accompanying photograph shows the new and dis-
tinct cultivar as nearly true as is reasonably possible to make
the same, in a color illustration of this character.

FIG. 1 illustrates the face view of the flower.

FIG. 2 illustrates the whole plant.

DETAILED DESCRIPTION

The following is a detailed description of my hybrid
daylily cultivar based upon observation of plants grown at
Grain Valley, Mo. during May, 1993 through October, 1997.
All colors being noted as compared with the Pantone Match-
ing System (PMS), except the dominate bloom color (near
white) being noted as compared with The Royal Horticulture
Society Colour Chart (R.H.S.). The measurements and col-
ors were recorded from mature plants grown in the vicinity
of Grain Valley, Mo. and from trial sites for the All American
Daylily Selection Council.

THE PLANT

Origin: It should be noted that the plant was initially selected
from a daylily planting being grown near Grain Valley,
Mo. in a cultivated area and has since been a sexually
reproduced by root division (division at the crown or
basal plate which is the area where the roots and foliage
meet) in the vicinity of Grain Valley, Mo. with the new
and distinct characteristics stated herein, found to be
maintained through successive generations as before
recited.

Parentage:

Seed parent.—Hemerocallis Hybrid 'Joan Senior' (not
patented).

Pollen parent.—Hemerocallis Hybrid 'Seagull' (not
patented).

Classification:

Botanic.—Hemerocallis hybrid.

Commercial.—Daylily cultivar 'Robbobelii'.

Form: Single stem, erect and stately scape from a fan shaped
plant made up of narrow arching grass-like, long keeled
glabrous leaves 2-ranked at the base of the scape.

Foliage:

Quantity.—Abundant, a mature plant having about 12 leaves

Leaf size.—About 1.1" wide and about 19.7" average in length.

Leaf shape.—Linear and long keeled with entire margins.

Texture.—Glabrous.

Color.—Immature leaf, top and under side — Green PMS #370; Mature leaf, top side —Green PMS #364; underside —Green PMS #363.

Clump foliage height (24 month maturity).—16" average in height; 26.6" average in width.

Plant bunch formation rate.—An average increase of 7 plants per 24 month period compared to an average of 4 plants for the same period for the average plant in this color class. This increase or bunch formation data was collected by the AADSC. The clumps should be divided or separated every 3–5 years depending on the USDA zone and the increase rate. This separation is to maintain optimum growth and bloom beauty. The specimens will continue to flower for many years without division although the quality of performance will be diminished.

Scape:

Color.—Green (PMS #364).

Height.—About 17.3".

THE BUD

Form: Modified (Slightly wider at the mid-point) oblanceolate.

Size (day before opening):

Length.—About 5 cm.

Width.—About 2 cm.

Opening rate: About 11 hours.

Length bloom is open: About 16 hours, i.e., an "extended" bloomer per American Hemerocallis Society terminology.

Color: When sepals first divide — pale lime green (PMS #372).

Peduncle:

Character.—Rigid and sturdy.

Color.—Green (PMS #364).

THE FLOWER

Blooming habit: Flowers begin to appear on the established plant in mid-season (or the middle of the hemerocallis blooming season which is referred to as "peak bloom") and then repeat bloom with 1–2 more cycles of scape production, depending on USDA zone. 'Robbobeli' blooms for an average period of 44 days (compared to the average daylily tested by the AADSC, which has an average bloom period of 21 days).

Flower size:

Diameter.—About 5/4".

Depth.—About 2".

Shape: Triangular and recurved

Borne: Singly on the branchlets of a sturdy, erect rachis which is ramulus; each scape has at least four or five peduncles which divides into twelve or fifteen pedicles.

Number of blooms per scape: 10–13 average.

Petalage: The flower consists of six perianth segments (tepals) consisting of three sepals (the outer segments) and three petals (the inner segments) all in an imbricated arrangement.

THE SEPALs

Shape: Oblanceolate with slightly undulated entire margins and an acuminate apex.

Size: About 3" long and 1.5" wide.

Texture: Ribbed, lightly creped, and diamond dusted.

Color: Near-white (RHS 155D) with yellow-green throat (PMS 393) and green heart (PMS 377).

THE PETALS

Shape: Broadly obovate with entire undulated margins and a broadly cuspidate apex.

Color: Near white (RHS 155D) with yellow-green throat (PMS 393) and green (PMS 377) heart.

Texture: Ribbed, lightly creped, and diamond dusted.

Size: About 2.3" wide; 3" long.

Effect of weather: Resistant to sun andrain; petal turgidity holds well.

Fragrance: Slightly sweet floral scent

Disease resistance: This cultivar, as most hemerocallis, has not been noted to be very susceptible to plant disease

Lasting quality: Good lasting quality due to flower segments (tepals) having thick, moderately spongy consistency which aids in fresh and turgid blooms throughout the day.

As is typical for hemerocallis, blooms are only open for one day.

THE REPRODUCTIVE ORGANS

Stamens:

Number.—Six to eight.

Arrangement.—Inserted individually at the summit of the perianth tube.

Anthers.—Arrangement: Introrse; Size: 5 mm; Color: Brown (PMS #438).

Filaments.—Slender and 4 cm in length; Color: White (RHS #155D).

Pollen.—Yellow (PMS #109).

Pistils

Number.—One.

Style.—Length — 7 cm; Color — White (RHS #155D).

Stigma.—Color: White & glistening.

Character of ovaries.—Three celled.

Shape.—Oblong, becoming a loculicidal 3-valved capsule.

THE FRUIT (SEED POD)

Shape: The seed pod is an ovoid capsule

Color: Ranges from pale green (PMS #372) to olive brown (PMS #140) at maturity.

Fertility: Yes, this cultivar is fertile both ways, i.e., the pollen is fertile for application to the pistil of another cultivar and the pistil of this cultivar can receive pollen from another cultivar.

Size (at maturity): About 7/8" wide by 1" in length.

GENERAL OBSERVATIONS

'Robbobeli', with its near-white, diamond dusted blooms, its compact growth habit, its tendency to repeat bloom late in the season, its ability to adapt and perform in a wide range of USDA zones, its dormant foliage, its light fragrance, and its above average bunch formation rate is an improved hemerocallis cultivar in the white color category.

COMPARISON TO KNOWN VARIETIES

'Robbobi' may be compared with known varieties along the following lines:

As compared to the seed parent, 'Joan Senior', 'Robbobi' is a distinct cultivar because of its lighter color of bloom and its improved cold hardiness over 'Joan Senior', its faster bunch formation rate (7 fans in 24 months compared to 5 fans for 'Joan Senior' in the same period), its more compact growth habit (17.3" scape height compared to 25" scape height for 'Joan Senior' and its improved bloom abscission, i.e., drops its spent blooms more quickly than 'Joan Senior' giving a "cleaner" appearance in the landscape.

As compared to the pollen parent 'Seagull', 'Robbobi' is an improved cultivar because it is more compact than 'Seagull' (17.3" in height compared to 26" in height), it is more cold hardy than 'Seagull', has a fuller bloom form than 'Seagull' ('Robbobi' petals are 2.3" in width compared to 'Seagull' petals of 1.7" in width), and has more bloom turgidity which gives superior lasting quality of the bloom.

I claim:

1. The new and distinct daylily plant as shown and described herein.

* * * * *



FIG. 1



FIG. 2