



(86) Date de dépôt PCT/PCT Filing Date: 1992/01/02
(87) Date publication PCT/PCT Publication Date: 1992/10/01
(45) Date de délivrance/Issue Date: 2008/10/07
(85) Entrée phase nationale/National Entry: 1993/09/13
(86) N° demande PCT/PCT Application No.: US 1992/000122
(87) N° publication PCT/PCT Publication No.: 1992/016620
(30) Priorité/Priority: 1991/03/15 (US07/670,085)

(51) Cl.Int./Int.Cl. *C12N 15/16* (2006.01),
A61K 38/24 (2006.01), *A61P 43/00* (2006.01),
C07K 14/72 (2006.01), *C12N 15/63* (2006.01),
C12N 5/10 (2006.01), *C12P 21/02* (2006.01),
G01N 33/566 (2006.01), *G01N 33/74* (2006.01)

(72) Inventeurs/Inventors:
SCHWEICKHARDT, RENE L., US;
NUGENT, NOREEN P., US;
CHENG, SHIRLEY V. Y., US;
KELTON, CHRISTIE A., US

(73) Propriétaire/Owner:
APPLIED RESEARCH SYSTEMS ARS HOLDING N.V.,
AN

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : RECEPTEUR HUMAIN DE LA FOLLICULOSTIMULINE
(54) Title: HUMAN FOLLICLE STIMULATING HORMONE RECEPTOR

(57) **Abrégé/Abstract:**

The present invention embraces essentially pure human FSH receptor, or a fragment or mutant thereof which binds FSH, DNA encoding said receptor, fragment or mutant, expression vectors comprising said DNA, cells transfected with said expression vectors, and methods of producing said receptor, fragment or mutant by culturing said transfected cells. The present invention also includes pharmaceutical compositions comprising said receptor, fragment or mutant, as well as methods of treating patients with such compositions to reduce endogenous FSH bioactivity. An improved assay for human FSH using the receptor, fragment or mutant of the present invention is also disclosed.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C12N 15/12, C07K 13/00 A61K 37/02, 37/43, C07K 15/00	A1	(11) International Publication Number: WO 92/16620 (43) International Publication Date: 1 October 1992 (01.10.92)
(21) International Application Number: PCT/US92/00122 (22) International Filing Date: 2 January 1992 (02.01.92) (30) Priority data: 670,085 15 March 1991 (15.03.91) US (60) Parent Application or Grant (63) Related by Continuation US 670,085 (CIP) Filed on 15 March 1991 (15.03.91) (71) Applicant (for all designated States except US): APPLIED RESEARCH SYSTEMS ARS HOLDING N.V. [NL/NL]; John B. Gorsiraweg, Curacao (AN). (72) Inventors; and (75) Inventors/Applicants (for US only) : KELTON, Christie, Ann [US/US]; 4 Valentine Road, Hopkinton, MA 01748 (US). CHENG, Shirley, Vui, Yen [MY/US]; 1 Longfellow Place, Apt. 2215, Boston, MA 02114 (US). NUGENT, Noreen, Patrice [US/US]; 38 Westgate Road, Framingham, MA 01701 (US). SCHWEICKHARDT, Rene, Lynn [US/US]; 148 Presidents Lane, Apt. 10, Quincy, MA 02169 (US).		(74) Agent: BROWDY, Roger, L.; Browdy & Neimark, 419 Seventh Street, N.W., Washington, DC 20004 (US). (81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC (European patent), MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL (European patent), NO, RO, RU, SD, SE (European patent), SN (OAPI patent), TD (OAPI patent), TG (OAPI patent), US. Published <i>With international search report.</i>
(54) Title: HUMAN FOLLICLE STIMULATING HORMONE RECEPTOR		
(57) Abstract The present invention embraces essentially pure human FSH receptor, or a fragment or mutant thereof which binds FSH, DNA encoding said receptor, fragment or mutant, expression vectors comprising said DNA, cells transfected with said expression vectors, and methods of producing said receptor, fragment or mutant by culturing said transfected cells. The present invention also includes pharmaceutical compositions comprising said receptor, fragment or mutant, as well as methods of treating patients with such compositions to reduce endogenous FSH bioactivity. An improved assay for human FSH using the receptor, fragment or mutant of the present invention is also disclosed.		

HUMAN FOLLICLE STIMULATING HORMONE RECEPTOR

Background of The Invention

This invention relates to human follicle stimulating
10 hormone receptor and its synthesis by recombinant DNA
techniques.

Follicle stimulating hormone (FSH) is a pituitary-
derived heterodimeric glycoprotein hormone which shares
structural similarities with luteinizing hormone (LH) and
15 thyroid stimulating hormone (TSH), both of which are also
produced in the pituitary gland, and chorionic gonadotropin
(CG), which is produced in the placenta. The hormones are
relatively large (28-38 kilodaltons) and are composed of a
common α subunit non-covalently bound to a distinct β
20 subunit that confers receptor binding specificity.

The cellular receptors for these hormones are known
to be members of the G protein-coupled class of membrane-
bound receptors which, when activated, stimulate an
increase in the activity of adenylyl cyclase. This results
25 in an increase in the level of the intracellular second
messenger adenosine 3',5'-monophosphate (cAMP), which in
turn causes increased steroid synthesis and secretion.
Hydropathicity plots of the amino acid sequences of these
receptors reveal three general domains: (1) a hydrophilic
30 amino-terminal region, considered to be the amino-terminal
extracellular domain, (2) seven hydrophobic segments of
membrane-spanning length, considered to be the
transmembrane domain, and (3) a carboxy-terminal region
which contains potential phosphorylation sites (serine,
35 threonine, and tyrosine residues), considered to be the
carboxy-terminal intracellular or cytoplasmic domain. The
glycoprotein hormone receptor family is distinguished from
other G protein-coupled receptors, such as the $\beta 2$ -

adrenergic, rhodopsin, and substance K receptors, by the large size of the hydrophilic amino-terminal domain, which is involved in hormone binding.

The FSH receptor is expressed on testicular Sertoli
5 cells and ovarian granulosa cells. While there has been a recognized need for providing essentially pure human FSH receptor, purification of naturally derived preparations is not practical and would not likely be sufficient to permit determination of the amino acid sequence. Recently, one
10 group has cloned the cDNA encoding the rat FSH receptor, deduced the amino acid sequence, and expressed it in mammalian cells (Sprengel, Mol. Endocrinol. 4:525, 1990). Another group, attempting to clone the TSH receptor, apparently also cloned and identified a portion of the
15 transmembrane region of the human FSH receptor (Parmentier, Science 246:1620, 1989).

Summary of The Invention

20

The present invention embraces essentially pure human FSH receptor, or a fragment or mutant thereof which binds FSH, DNA encoding said receptor, fragment or mutant, expression vectors comprising said DNA, cells transfected
25 with said expression vectors, and methods of producing said receptor, fragment or mutant by culturing said transfected cells. The present invention also includes pharmaceutical compositions comprising said receptor, fragment or mutant, as well as methods of treating patients with such
30 compositions to reduce endogenous FSH bioactivity. An improved assay for human FSH using the receptor, fragment or mutant of the present invention is also disclosed.

2106361

Brief Description of The Drawings

Figure 1 is a map of the human FSH receptor cDNA clones. A partial restriction endonuclease map is shown which was determined by combining DNA sequence data derived from portions of each of the five clones. Size increments of 0.2 kilobase pair (kb) are marked. The regions to which the clones correspond are indicated by the solid lines below the restriction endonuclease map. The clone designation and the approximate size in kb are indicated to the left and right of each solid line, respectively. Approximate locations of the amino-terminal extracellular domain, transmembrane domain, and carboxy-terminal intracellular domain of the encoded protein are indicated by the dashed arrows above the restriction endonuclease map. The position of an insertion (probably an intron or a portion of an intron) in the 5-10 clone is indicated by the open box.

Figures 2A and 2B depict the strategy used for engineering a "full-length" human FSH receptor cDNA for the purpose of expression of the encoded protein in a mammalian cell line.

Figure 3 is a diagram of the expression vector plasmid used to develop CHO-DUKX cell lines stably transfected with the human FSH receptor.

Figure 4 is a diagram of the expression vector plasmid used to develop Y1 cell lines stably transfected with the human FSH receptor.

Figure 5 illustrates intracellular cAMP levels measured in the CHO-DUKX cell line CHOFSHR4Q-13, after treatment with varying doses of either FSH (circles), or LH (triangles). Each point represents the mean value. Standard errors are shown by the vertical bars.

Figure 6 illustrates progesterone levels measured in the culture medium of the Y1 cell line Y1HFSHR4-38, after treatment with varying doses of FSH. Vertical bars depict the standard error in the progesterone measurements at each FSH dose.

Detailed Description of the Invention

As used herein, human FSH receptor, and its FSH-binding fragments or mutants, refers to polypeptides which are capable of recognizing and selectively binding with human FSH and which produce no significant immunological response in humans. Thus, the present invention includes human FSH receptor having the amino acid sequence depicted in Sequence ID No:2, as well as FSH-binding fragments or mutants thereof which retain a very high degree of homology (i.e. at least 95% identity) with the depicted amino acid sequence, or with at least the extracellular portion thereof. Fragments of the human FSH receptor embraced by the present invention include those fragments remaining after the cytoplasmic and/or the transmembrane domains have been deleted from the full polypeptide. An especially preferred fragment embraced by the present invention is the amino-terminal extracellular portion comprising approximately amino acids 1 to 349 of the human FSH receptor amino acid sequence shown in Sequence ID No:2. Since some of the loops which span the transmembrane regions (listed in features section, Sequence ID No:2) are also extracellular, these may be linked (e.g. through appropriate spacer molecules) to a fragment containing the amino-terminal extracellular portion to improve binding. The polypeptide may be glycosylated, as in the natural receptor, or may be partially or completely deglycosylated.

It is further contemplated that only a portion of the above-described amino-terminal extracellular domain may be effectively utilized to bind FSH since it is very likely that the complete extracellular domain is not necessary for this purpose. Thus, a fragment which is somewhat shorter than the 349 amino acids of the complete extracellular domain may be readily produced and tested for effective binding to FSH. So long as the FSH-binding region of the extracellular portion is maintained intact, the length of the overall fragment utilized is not critical. For this reason, it is also expected that non-interfering amino

acids can be added to either end of the extracellular domain, or FSH-binding fragment thereof, without adversely affecting the FSH-binding capacity of the polypeptide. Accordingly, this invention embraces a fragment of the
5 human FSH receptor which comprises a substantial portion of the extracellular domain and which retains substantially the same FSH-binding characteristics as the complete extracellular domain.

Mutant forms of the above-described receptor and
10 fragments are also within the scope of the present invention. Such mutants include conservative substitutions in one to ten amino acid residues, the location and nature of such substitutions being selected so as not to significantly degrade the FSH-binding characteristics of
15 the receptor or fragment thereof which is modified.

Essentially pure human FSH receptor, or a fragment or mutant thereof, is prepared by isolating and cloning the DNA encoding it from a cDNA or genomic library, ligating the DNA into a vector, transfecting host cells with the
20 vector, culturing transfected host cells under conditions which permit expression of the receptor, fragment or mutant, and recovering the receptor, fragment or mutant from the culture.

The DNA which is used to make expression vectors may
25 be genomic DNA or cDNA encoding human FSH receptor, and may contain regions which enhance expression, such as introns, promoters, enhancers, etc. The DNA may be readily modified by substitution, deletion or insertion of nucleotides (e.g. by site-specific mutagenesis) that do not adversely affect
30 the biological or FSH-binding activity of the expressed protein. For example, conservative substitutions (mutations) which alter from one to ten amino acids may be made without adversely affecting the overall structure and activity of the expressed protein (mutant). Likewise,
35 certain portions of the DNA, such as those portions which code for the cytoplasmic and/or transmembrane domains, may be deleted so that only a fragment, such as the soluble extracellular domain, of the protein is expressed. The

2106061

-6-

human FSH receptor or FSH-binding fragment or mutant thereof may also be expressed as a fusion protein. One such fusion protein would include a polypeptide at the carboxy-terminus which would confer characteristics which would facilitate purification of the protein, or immobilization of the purified protein on a solid substrate for use in FSH assays or FSH purification protocols. Another such fusion protein would include a cleavable polypeptide at the amino-terminus which would facilitate expression.

Human FSH receptor produced in accordance with the present invention is essentially pure, meaning that it is substantially free of biological adventitious agents normally associated with FSH receptor extracted from natural sources, such as, for example, bacteria, viruses, and other proteins. The receptor may be formulated into pharmaceutical compositions by mixing with suitable pharmaceutically acceptable carriers in a manner known to those skilled in the art.

Generally, pharmaceutical compositions may be formulated for oral, parenteral (including subcutaneous, intramuscular, and intravenous), vaginal, rectal, buccal (including sublingual), transdermal or intranasal administration. Compositions for parenteral administration are normally in the form of a liquid solution, dispersion, or emulsion, preferably isotonic; for vaginal or rectal administration, as a cream or suppository; for oral or buccal administration, as a tablet or capsule; and for intranasal administration, as a powder, nasal drop or aerosol. Various slow release, depot implant or injectable dosage forms may be utilized. The active component may also be incorporated into polymer matrices, liposomes and microspheres to control delivery.

These compositions may conveniently be administered in unit dosage form and may be prepared by any of the methods well-known in the pharmaceutical art. Formulations for parenteral administration may contain as common excipients sterile water or saline, alkylene glycols such

as propylene glycol, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes and the like. Formulations for vaginal or rectal administration, e.g. suppositories, may contain as
5 excipients, for example, polyalkyleneglycols, vaseline, cocoa butter, and the like. Formulations for nasal administration may be in powder form and contain as excipients, for example, lactose or dextran, or may be aqueous or oily solutions for administration in the form of
10 nasal drops or metered spray. For buccal administration, typical excipients include sugars, calcium stearate, pregelatinated starch and the like. One or more surfactant acids or salts can be added to the solution or powder formulation. Suitable pharmaceutically acceptable
15 surfactant salts will be those which retain the phenomenon of enhanced peptide absorption, as well as the compound's surfactant characteristics and which are not deleterious to the subject or otherwise contraindicated.

The dosage of active ingredient administered, as well
20 as the route and frequency of administration, will obviously depend upon the needs and condition of the patient being treated, the therapeutic effect which is sought, and the judgement of the doctor conducting the treatment.

25 Pharmaceutical compositions comprising human FSH receptor, or FSH-binding fragments or mutants thereof, may be administered to a patient in therapeutically effective doses to bind with endogenous circulating FSH in the patient and thereby control the available level of
30 bioactive FSH. Thus, pharmaceutical compositions of the present invention are effectively utilized to reduce endogenous FSH bioactivity. In a female patient, such treatment may be effectively used to prevent follicle growth and maturation, thereby preventing pregnancy. In a
35 male patient, such treatment may be effectively used to prevent spermatogenesis. A particularly suitable pharmaceutical composition for the above purpose comprises a fragment of the human FSH receptor which comprises the

2106061
amino-terminal extracellular domain or a substantial portion thereof with substantially the same FSH-binding characteristics.

Essentially pure human FSH receptor may also be
5 advantageously utilized in conventional receptor assays for FSH, such as, for example, that disclosed in Reichert, Endocrinology 94:483, 1974. Substitution of the pure receptor of the present invention, or a fragment thereof which binds FSH, will substantially improve the consistency
10 and performance of such assays. A fragment comprising the amino-terminal extracellular domain or an FSH-binding fragment and/or fusion protein thereof can also be linked to an affinity column to purify FSH from fluids, extracts, etc.

15 The receptor can be incorporated into a stable cell line, preferably a eukaryotic cell line, and most preferably a mammalian cell line, which is capable of producing a measurable biological response upon stimulation of the receptor. Measurement of cell response in the
20 presence of FSH under assay (e.g. in test serum, plasma, culture media, tissue homogenates, etc.) would provide an indication of bioactivity, and thereby provide a highly useful diagnostic assay. Such a cell line could also be used to screen chemical libraries for substances that may
25 interact with the FSH receptor or to test peptides or small proteins for their ability to bind to the FSH receptor in a rapid through-put screening system. An example of a rapid through-put screening system is one in which the binding of a ligand to the recombinant FSH receptor results in the
30 production or blockage of production of a measurable product such as cAMP or progesterone. Such a system could be further enhanced by operatively linking an easily measured marker element such as one which produces bioluminescence (e.g. the luciferase gene) to the receptor
35 signal transduction pathway (e.g. by using the cAMP response element).

Essentially pure human FSH receptor or FSH-binding fragments or mutants thereof can also be used in X-ray

crystallographic analysis to develop molecular models. Such models are useful in defining the tertiary structure of the hormone-binding domains of the human FSH receptor. Such information would provide insight into the structure
5 of the actual regions of contact between FSH and its receptor, thus aiding the design of peptides which have FSH agonistic or antagonistic activity.

The recombinant techniques suitable for producing the proteins and DNA of the present invention, including
10 identification of suitable mutation techniques, vectors, host cells, culture conditions, etc., are well-known to those skilled in the art and are adequately described, for example, in US 4,761,371 and WO 88/09818, the disclosures of which are incorporated herein by reference. The
15 experimental protocols, bacterial and bacteriophage culture media and chemical solutions used in the examples below are described in detail in Sambrook, "Molecular Cloning: A Laboratory Manual," 2nd ed., Cold Spring Harbor Laboratory Press, 1989, unless otherwise referenced.

20

Example 1. Isolation and Characterization of Human
FSH Receptor cDNA Clones

25 Library screening.

A rat FSH receptor cDNA clone similar to that described by Sprengel, Mol. Endocrinol. 4:525, 1990 was obtained from Dr. William Moyle, University of Medicine and Dentistry of New Jersey-Robert Wood Johnson Medical School.
30 This cDNA clone was inserted into the SV40 late expression vector pSVL (Pharmacia LKB, product number 27-4509-01) and was designated pSVLFSHR. A 2.1 kb DNA fragment containing the region corresponding to the complete rat FSH receptor protein coding region was excised from the plasmid using
35 the restriction endonuclease sites XbaI and BamHI. The digested DNA was size-fractionated by gel electrophoresis, and the 2.1 kb rat FSH receptor fragment was purified by electroelution from the gel. This purified DNA fragment

was used as a probe for library screening to identify human FSH receptor cDNA clones.

A lambda gt11 cDNA library, constructed from RNA extracted from human testis, was purchased from Clontech, Palo Alto, CA (catalogue number HL1010b) and was amplified prior to use. Twenty aliquots of the amplified library, equivalent to approximately 7.5×10^4 plaque forming units (pfu), were each adsorbed to about 0.5 milliliters (ml) of a plating suspension of E. coli strain Y1088. The suspension had been prepared by growing an overnight culture of Y1088 at 37°C in NZYM or LB supplemented with 0.2% maltose and 10 millimolar (mM) MgSO_4 , pelleting the cells, and then resuspending them in 10 mM MgSO_4 to an O.D.₆₀₀ of 0.5. About 6.5 ml of molten NZYM top agarose (0.7%), at a temperature of 48°C, was added to each phage/cell suspension and the resulting mixture was poured onto one of twenty 150 millimeter (mm) NZYM agar plates (prewarmed to 42°C). The total number of phage plated for the primary screen was about 1.5×10^6 . After a 4 hour incubation at 42°C, followed by chilling at 4°C for several hours, duplicate nitrocellulose filter (Millipore) plaque lifts were generated from each plate according to the procedures of Benton and Davies (Science, 196:180, 1977). The random oligonucleotide priming procedure of Feinberg and Vogelstein (Anal. Biochem. 137:266, 1983) was used to generate, from the template rat FSH receptor DNA fragment described in the preceding paragraph, a ^{32}P -labeled probe of a specific activity of $1-2 \times 10^9$ counts per minute (cpm)/microgram (μg). Prehybridization of the nitrocellulose filter phage lifts was done in a buffer containing 50% formamide, 5X SSC [1X SSC is 0.15 molar (M) sodium chloride, 0.015 M sodium citrate], 20 mM sodium phosphate buffer, pH 7.2, 10X Denhardt's reagent (50X Denhardt's reagent is 1% Ficoll[®], 1% polyvinylpyrrolidone and 1% bovine serum albumin) and 100 $\mu\text{g}/\text{ml}$ tRNA at 37°C for about 6 hours. Hybridization of the filters was done in the same buffer except that the rat FSH receptor DNA probe was added at a concentration of about 3×10^6 cpm/ml

-11-

buffer. After hybridization for 16-24 hours at 37°C, excess probe was washed from the filters in 2X SSC, 0.1% SDS at room temperature for 30 minutes, then 0.2X SSC, 0.1% SDS at 37°C for 60 minutes. The filters were subsequently
5 exposed to XAR film (Kodak) overnight at -70°C. Six duplicate positives were identified from the primary library screen. With the wide end of a pasteur pipette, plaque-containing agar plugs were removed from the 150 mm plates in the regions in which the positive clones were
10 located. The phage were eluted by soaking the plugs in SM. The suspended phage were then replated onto 150 mm NZYM agar plates as described for the primary screen. Plates which contained about 500 pfu per plate were selected for secondary screening. The procedures used for filter lifts
15 and filter hybridizations for the secondary screen were the same as described for the primary screen.

Following the secondary screen, 5 putative human FSH receptor positives were identified and isolated as purified λ gt11 bacteriophage clones. They were assigned the
20 following designations: 1-5, 5-10, 11-11, 13-9, and 15-6.

Determination of the DNA sequences of the putative human FSH receptor cDNA clones.

Bacteriophage DNA was prepared from each of the λ gt11
25 cDNA isolates using the plate lysate method described in Sambrook, "Molecular Cloning: A Laboratory Manual," 2nd ed., Cold Spring Harbor Laboratory Press, 1989, page 2.118. The bacteriophage DNA was digested with EcoRI restriction endonuclease, and then size-fractionated in an agarose gel
30 for purification of the cDNA insert fragments. The purified cDNA inserts were subcloned into the EcoRI site of pUC18 in order to facilitate subsequent cloning and sequencing manipulations. Double-stranded plasmid DNA was purified from 5 ml cultures of the host strain E. coli
35 MC1061 by the small-scale alkaline lysis method (Sambrook, *ibid.* p. 1.25), then further purified by passage over Elutip-d[®] columns (Schleicher & Schuell, Keene, NH) using the manufacturer's protocol. Half of the plasmid DNA

2106061
obtained from each small-scale plasmid preparation was then denatured in 0.2 normal NaOH, 0.2 mM EDTA in a volume of 20 microliters (μ l) at room temperature for 10 minutes. The denatured plasmid DNA was neutralized and ethanol
5 precipitated by adding 7.5 μ l of 7.5 M ammonium acetate and 110 μ l 100% ethanol and chilling the mixture in liquid nitrogen. The DNA precipitate was pelleted by centrifugation in a microfuge for 10 minutes. The DNA pellets were washed in 70% ethanol and dried. Sequencing
10 reactions were done with Sequenase T7 DNA polymerase (United States Biochemical) according to the manufacturer's specifications. Preliminary sequencing reactions were done with forward and reverse sequencing primers (Pharmacia LKB) for the pUC18 polylinker region. Data generated from
15 preliminary sequencing was used to design human FSH receptor-specific sequencing primers. The primers were either synthesized on a model 391 Applied Biosystems DNA synthesizer or ordered from National Biosciences, Inc., Hamel, MN. Some DNA sequence data was obtained by
20 subcloning smaller restriction endonuclease fragments of the original clones into pUC18 and repeating sequencing reactions with the forward and reverse primers.

Preliminary sequence data showed that none of the five cDNA isolates represented the full-length protein
25 coding region of the human FSH receptor, but that the combined sequence data from the clones could be used to infer the complete protein sequence. A schematic diagram of the relative locations of each of the five clones in relation to a map of the complete human FSH receptor cDNA
30 sequence is shown in figure 1. The complete human FSH receptor cDNA sequence, obtained by combining the sequence readings from the overlapping clones using the Genetics Computer Group (GCG) fragment assembly computer program, is depicted in Sequence ID No:1 and the inferred amino acid
35 sequence is depicted in Sequence ID No:2. Analysis of the human FSH receptor DNA sequence resulted in the identification of a long open reading frame of 2085 nucleotides which encoded a protein of 695 amino acids.

-13-

The human FSH receptor is therefore 3 amino acids longer than the rat FSH receptor. The overall percent identities between the rat and human FSH receptor DNA and protein sequences were determined using the GCG Bestfit program and were 87% and 90%, respectively. The extracellular amino-terminal hydrophilic portion of the human FSH receptor is estimated to be 349 amino acids in length, and shares 87% identity with the corresponding region of the rat FSH receptor. The seven membrane spanning regions of the two species, which are bridged by three extracellular and three intracellular loops, share 95% identity, and the carboxy-terminal intracellular regions share only 81% identity. The partial amino acid sequence published by Parmentier, Science 246:1620-1622, 1989, corresponds to amino acids 399 to 525 in Sequence ID No:2.

Clone 5-10 was a variant in that it contained an insertion of 0.25 kb after the T at nucleotide position 448 in Sequence ID No:1. The DNA sequence of the insertion was not similar to any portion of the rat or human FSH receptor DNA sequence and also was not similar to any known sequence in the Genbank[®] or EMBL[®] DNA sequence databases. The corresponding region in clone 11-11 did not contain this insertion. LH receptor cDNA clones isolated from a human thyroid cDNA library contain similar variations (Frazier, Mol. Endocrinol. 4:1264-1276, 1990). These variants are likely derived from incompletely and/or aberrantly spliced mRNA molecules. The presence of a 3' splice consensus sequence (CAG'G) at the 3' junction of the 5-10 insertion is evidence which supports this explanation.

The pUC18 plasmids harboring the 11-11 (called pHFSHR11-11), 15-6 (called pHFSHR15-6), and 5-10 (called pHFSHR5-10) cDNA inserts were deposited with the American Type Culture Collection (ATCC), Rockville, MD, on March 1, 1991 and have received the accession numbers ATCC68538, ATCC68540, and ATCC68539, respectively. These deposits were made in accordance with all of the requirements of the Budapest Treaty.

Example 2. Construction of vectors for the expression of the full-length human FSH receptor in mammalian cells.

2106061

The strategy for engineering a human FSH receptor DNA construct for expression in mammalian cells is shown in Figures 2A and 2B. Referring to Figure 2A, the 5' 691 base pair (bp) NsiI-BamHI fragment which contained the start ATG was purified from pHFSHR11-11 and subcloned into pUC18 digested with PstI and BamHI. The resulting plasmid, pHFSHR11-11nb, was linearized with SphI. The termini were blunted by treatment with the Klenow fragment of DNA polymerase I (New England Biolabs, Beverly, MA). After ligation of XhoI linkers (New England Biolabs, Beverly, MA) to the blunted termini, the mixture was digested with XhoI and BamHI and the approximately 700 bp human FSH receptor 5' fragment was gel purified. A fragment from the middle region of the human FSH receptor cDNA was isolated from pHFSHR11-11 by digestion with BamHI and SphI and gel purification of the appropriate 734 bp piece. The 3' human FSH receptor fragment, which contained the TAA stop codon, was isolated from pHFSHR15-6 by first digesting the plasmid with DraIII, then blunting the termini in a reaction with T4 DNA polymerase. XhoI linkers were ligated to the blunted ends and the resulting mixture was digested with SphI and XhoI and the 3' 690 bp (approximately) SphI-XhoI fragment was isolated and purified.

Referring to Figure 2B, the 5' 700 bp XhoI-BamHI and middle 734 bp BamHI-SphI human FSH receptor cDNA fragments were subcloned into pUC18-XhoI (created in this laboratory by converting the SmaI site in the pUC18 polylinker to an XhoI site with XhoI linkers) digested with XhoI and SphI. The resulting plasmid, pHFSHR1.4xs, was digested with XhoI and SphI. The digested DNA was size-fractionated by electrophoresis in a 5-6% polyacrylamide gel and the approximately 1400 bp fragment containing the 5' and middle regions of the human FSH receptor cDNA was excised and purified by electroelution. The 3' SphI-XhoI fragment, which contains the TAA stop codon, was subcloned into

pUC18-XhoI digested with SphI and XhoI. The resulting plasmid, pHFSHR675sx, was digested with SphI and EcoRI. The digested DNA was fractionated in a 5-6% polyacrylamide gel and the approximately 700 bp piece was excised and
5 purified by electroelution. The complete human FSH receptor coding region was assembled by combining the 5' 1400 bp XhoI-SphI fragment with the 3' 700 bp SphI-EcoRI fragment in a ligation with pUC18-XhoI digested with XhoI and EcoRI. Correct assembly of the construction, called
10 pHFSHRX, was verified by restriction endonuclease digestion and DNA sequencing. The completely engineered human FSH receptor 2.1 kb XhoI fragment, which contains approximately 18 bp of 5' noncoding sequence and 12 bp of 3' noncoding sequence in addition to the entire human FSH receptor
15 protein coding region, was excised from pHFSHRX by XhoI digestion. The digested DNA was size-fractionated by electrophoresis in a 0.7% agarose gel. The 2.1 kb XhoI fragment was excised from the gel, purified by electroelution in a Little Blue TankTM (ISCO) using the
20 procedure recommended by the manufacturer, and inserted into the XhoI cloning sites of the mammalian cell expression vector plasmids. Plasmids containing the insert in the appropriate orientation for human FSH receptor mRNA transcription were selected by restriction endonuclease
25 fragment analysis. Prior to introduction into mammalian cells, expression vector plasmid DNA was purified either by two sequential high-speed centrifugations through cesium chloride density gradients, or by using the plasmid maxi kit (Qiagen, Chatsworth, CA) according to the
30 manufacturer's protocol.

Any suitable mammalian cell expression vector can be used for the expression of the human FSH receptor. The necessary vector plasmid components include the following: a eukaryotic promoter such as the mouse metallothionein I
35 (MMT-I) promoter, Rous Sarcoma Virus promoter, or simian virus 40 early or late promoters for driving transcription of the human FSH receptor mRNA; a marker gene such as the neomycin resistance gene (Neo), dihydrofolate reductase

2206081
(DHFR) gene, or multidrug resistance gene (MDR) for the selection of transfected cells; a polyadenylation signal (poly A), such as the SV40 early polyadenylation region for the 3' processing of the receptor RNA transcript; and a
5 bacterial origin of replication and antibiotic resistance gene such as the pBR322 ori and the ampicillin resistance gene, to enable growth and propagation of the vector plasmid in a suitable E. coli strain. For some cell lines, it may be important to include an intron in the region
10 which is transcribed into human FSH receptor encoding mRNA. It is preferable to place the intron between the promoter and the human FSH receptor cDNA insert, thereby positioning it in the 5' untranslated region of the human FSH receptor RNA transcript. Any suitable intron can be used. The
15 intron used in our experiments was a 2 kb XbaI-PstI portion of intron A in the human glycoprotein α subunit gene (Fiddes, J. Mol. Appl. Genet. 1:13, 1981). In the expression vectors, it was inserted between the the MMT-1 promoter region and the human FSH receptor cDNA fragment.
20 The truncated intron retained the endogenous splice acceptor, but the splice donor was supplied by a synthetic oligonucleotide.

A diagram of the human FSH receptor expression vector used for CHO cell transfections, pD α HFSHRX (human FSH
25 receptor in CLH3AXSV2DHFRh α IVS), is shown in Figure 3 and a diagram of the human FSH receptor expression vector used to stably transfect Y1 cells, pN α HFSHRX (human FSH receptor in CLH3AXSV2NEOh α IVS), is shown in Figure 4. The MMT-I gene sequences (nucleotide numbers 4542-7514 in Figure 3,
30 nucleotide numbers 5192-8164 in Figure 4) which include the promoter region, are the same as those shown upstream of the XhoI site in the construction of CLH3X (Reddy, DNA 6:461, 1987). The human glycoprotein α subunit gene intron A fragment, described in the preceding paragraph, is
35 located between nucleotide numbers 7514 and 9514 in Figure 3, and between nucleotide numbers 8164 and 10164 in Figure 4. Downstream of the XhoI site, the MMT-I introns and polyadenylation signal sequences were removed and replaced

with the SV40 early polyadenylation region (nucleotide numbers 11614-11857 in figure 3, nucleotide numbers 12264-12508 in figure 4). The pML region (Lusky, Nature 293:79, 1981) contains the bacterial origin of replication and
5 ampicillin resistance gene from pBR322 (nucleotide numbers 1925-4542 in Figure 3, nucleotide numbers 2575-5192 in Figure 4). The DHFR gene transcription unit (nucleotide numbers 1 to 1925 in Figure 3), which includes the mouse DHFR cDNA and the SV40 early region promoter, small T
10 intron and early region polyadenylation sequences, corresponds to the PvuII-BamHI fragment in pSV2DHFR (Subramani, Mol. Cell. Biol. 1:854, 1981). The PvuII site was converted to a BamHI site prior to insertion of the fragment into the expression vector plasmid. The neo gene
15 transcription unit (nucleotide numbers 1-2575 in figure 4) was assembled by replacing the DHFR cDNA (nucleotide numbers 342-1077 in Figure 3) with a 1385 bp HindIII-SmaI fragment from pSV2-neo (Southern, J. Mol. Appl. Genet. 1:327, 1982). The SmaI site was converted to a BamHI site
20 prior to insertion of the fragment into the expression vector plasmid.

The plasmid pN α hFSHRX can be used to transfect most continuous mammalian cell lines, and was deposited with the ATCC on November 19, 1991, as accession number ATCC68833.

25 The deposit was made in accordance with all of the requirements of the Budapest Treaty. The plasmid pD α hFSHRX is best suited for the transfection of cell lines which lack dihydrofolate reductase (DHFR) activity.

DNA cloning and engineering techniques can be used by
30 those skilled in the art to modify the human FSH receptor DNA expression construction so that FSH-binding fragments are encoded. These modified DNA fragments can be inserted into expression vectors such as those described herein and used to transfect mammalian cells to develop lines which
35 secrete soluble FSH-binding fragments of the human FSH receptor, such as the amino terminal extracellular domain or an FSH-binding fragment thereof.

Example 3. Development of Mammalian Cell Lines which
Stably Express a Functional Human Recombinant FSH Receptor
or an FSH-binding fragment or mutant thereof.

5 Suitable mammalian cell lines for expression of
recombinant human FSH and its derivatives include Chinese
Hamster Ovary (CHO), mouse adenocarcinoma Y1, rat pituitary
GH₃, human breast carcinoma MCF7, and human embryonic
kidney 293. In this example, the use of Y1 and CHO cells
10 is described.

Y1 cells are a clonal steroid secreting cell strain
initiated from a mouse adrenal cortex tumor (Yasumura,
Cancer Res. 26: 529-536, 1966). These cells were obtained
from the ATCC cell bank (ATCC CCL 79) and were maintained
15 in culture by growth in Ham's F10 medium supplemented with
15% horse serum (HS), 2.5% FBS, and 1% L-glutamine (Y1
growth medium).

CHO-DUKX cells are a clonal mutant of Chinese hamster
ovary cells lacking dihydrofolate reductase activity
20 (Urlaub G. and Chasin, L.A. PNAS 77: 4216-4220, 1980). The
cells were maintained in Minimum Essential Medium Alpha
(MEM- α) supplemented with 10% FBS and 1% L-glutamine (CHO
growth medium).

All cell culturing was done at 37°C in a 5% CO₂,
25 humidified atmosphere.

Calcium phosphate transfections.

The transfection protocol used was a modification of
a published method (Graham, Virology 52: 456, 1973).
30 Approximately 24 hours prior to transfection, cells were
plated in 100 mm diameter dishes at a density of 7×10^5
cells/dish (CHO-DUKX) or 1×10^6 cells/dish (Y1). For
transfections, 10 μ g of the vector plasmid DNA (pD α HFSHRX
for CHO-DUKX cells, pN α HFSHRX for Y1 cells) was added to
35 0.5 ml of transfection buffer. Transfection buffer was
prepared by combining 4 grams (g) NaCl, 0.185 g KCl, 0.05 g
Na₂HPO₄, 0.5 g dextrose, and 2.5 g HEPES in sterile,
distilled water and adjusting the volume to 500 ml, and the

pH to 7.5. To the DNA/transfection buffer mixture was added 31 μ l of 2 M CaCl_2 . The solution was mixed using a vortex mixer and allowed to stand at room temperature for 45 minutes to allow the DNA precipitate to form. The DNA precipitate was layered over the cells after removal of the culture medium. After 20 minutes at room temperature, 5 ml of the appropriate growth medium was added and the cells were cultured for 6 hours. The medium was removed and the cells shocked for 3.5 minutes in 3-4 ml of transfection buffer containing 15% glycerol. The cells were rinsed twice with phosphate buffered saline prior to the addition of 10 ml growth medium. Approximately 48 hours post transfection, the CHO-DUKX cells were subcultured at a 1:10 split ratio and selection medium was added. Y1 cells were allowed to grow for 72 hours post transfection before subculturing at a 1:5 split ratio in selection medium. Selection medium for CHO-DUKX cells consisted of MEM- α without ribonucleosides and deoxyribonucleosides supplemented with 10% dialyzed FBS, 1% L-glutamine, and 0.02 μ M methotrexate (MTX). Selection medium for Y1 cells consisted of Y1 growth medium with 80 μ g/ml G418. MTX-resistant CHO-DUKX cell colonies and G418-resistant Y1 cell colonies formed after approximately 2 to 3 weeks, and were individually picked and cultured until cell numbers were sufficient for cryopreservation and testing for hormone responsiveness.

Demonstration of Human Recombinant FSH Receptor Bioactivity in Stably Transfected CHO Cells.

To determine if the engineered human FSH receptor cDNA construct could produce a protein that was biologically functional in stably transfected CHO cells, the MTX-resistant CHO cell colonies were treated with human recombinant FSH and intracellular cAMP levels were measured. For the cAMP assay, the cells were subcultured in 12 well plates at a density of 2.5×10^4 cells/well in CHO cell selection medium. After 72 hours of culture, each well was washed with 1.5 ml of warm growth medium. To each

2106061

washed well was added 300 μ l of serum-free growth medium containing 0.1 mM 3-isobutyl-1-methylxanthine (Sigma). After 15 minutes of culture, human recombinant FSH was added to the medium in each well so that the final

5 concentration was approximately 335 ng (2413 mIU) per ml. After 30 minutes of culture, the assay was terminated by subjecting the cells to four rapid freeze-thaw cycles. A 50 μ l aliquot of each cell lysate was transferred to a 1.5 ml tube for determination of protein content. Cold ethanol

10 (300 μ l) was added to each remaining lysate sample before transfer to separate 1.5 ml tubes. All lysate samples were centrifuged at 13,000 X g for 15 minutes to remove cell debris. Determination of total soluble protein was done with a protein assay kit purchased from Bio-Rad (catalog

15 number 500-0002) using the protocol supplied by the manufacturer. The ethanol-treated lysate samples were freeze-dried in aliquots of 5-100 μ l, and resuspended in 100 μ l of the assay buffer provided in the Dupont/NEN Medical Products cAMP assay kit (catalog number NEK-033).

20 The cAMP content of the resuspended lysate samples was determined using the acetylated radioimmunoassay protocol supplied with the kit. The samples were not diluted for the assay. In the majority of the human FSH receptor-transfected CHO cell lines, higher intracellular cAMP

25 levels were measured after FSH stimulation than in the controls (untransfected CHO-DUKX cells and CHO-DUKX cells transfected with CLH3AXSV2DHFRh α IVS). The average response was about 16-fold that of the controls.

One nonclonal cell line, CHO_HFSHR4Q-13, was chosen

30 for a hormone dose response analysis. The procedure used was the same as in the preceding paragraph, except that the cells were treated with either human recombinant FSH or LH in concentrations ranging from 0-10,000 mIU/ml. Each dose was analyzed in triplicate. The amount of intracellular

35 cAMP produced was expressed in nanomoles (nmole) per milligram (mg) of soluble protein and plotted versus FSH or LH concentration to generate dose response curves. The FSH and LH dose response curves are shown in Figure 5. The

results show that the CHOFSHR4Q-13 CHO cell line displayed a saturable increase in intracellular cAMP in response to FSH stimulation but not in response to LH stimulation. Half-maximal FSH stimulation (ED₅₀) occurred at a dose of 144 mIU/ml FSH. The human FSH receptor cDNA therefore can be used to produce a biologically functional human FSH receptor protein in CHO cells, enabling the cells to mediate a specific cAMP response to FSH and not LH.

The human FSH receptor content of the CHOFSHR4Q-13 line, or a similar line which produces the human recombinant FSH receptor or FSH-binding fragments or mutants thereof, may be increased by exposing the cells to stepwise increases in MTX concentration. This causes amplification of DHFR cDNA copy number along with linked sequences which include the human FSH receptor cDNA, indirectly increasing synthesis of the encoded proteins. This may be done, if necessary, to increase the sensitivity of the bioassay. Also, mammalian cells which express high levels of human FSH receptor or FSH-binding fragments or mutants thereof, can be used for the manufacture of large amounts of human FSH receptor protein for use as therapeutics or in receptor radioimmunoassays.

The CHOFSHR4Q-13 line, or a similar line, may also be used in an assay for estimating the in vitro bioactivity of pharmaceutical preparations containing FSH, or FSH-like substances. A reporter gene such as luciferase, can be operatively linked to a cAMP response element, and increases in cAMP levels may be indirectly measured by a non-radioactive method, such as bioluminescence, to provide a measurement of FSH bioactivity.

The CHOFSHR4Q-13 line was deposited with the ATCC on November 19, 1991, as accession number ATCC CRL 10921. The deposit was made in accordance with all the requirements of the Budapest Treaty.

Demonstration of Recombinant Human FSH Receptor Bioactivity
in Stably Transfected Mouse Adrenal Y1 Cells.

To determine if the engineered human FSH receptor cDNA construct could produce a protein that was biologically functional in stably transfected Y1 cells, the G418-resistant Y1 cell colonies were treated with human recombinant FSH and the culture media analyzed for progesterone content. For the progesterone assay, the cells were subcultured in 6 well plates (35 mm well diameter) at a density of 4×10^5 cells/well in Y1 cell selection medium. After two days of culture, the Y1 cell selection medium was removed and replaced with 3 ml Ham's F10 medium supplemented with 5% horse serum, 0.8% fetal bovine serum, 1% L-glutamine, and 80 $\mu\text{g/ml}$ G418. After incubation for one day, the medium was removed and the cells were washed twice with 2 ml of Ham's F10 medium supplemented with 1 mg/ml bovine serum albumin (fraction V), 1% L-glutamine, and 80 $\mu\text{g/ml}$ G418 (assay medium). At this time the average cell number per well was estimated for each cell line by removing and counting cells from three test wells. The cells were then cultured for 4 hours with 1 ml assay medium containing 100 mIU/ml human recombinant FSH. After incubation, the culture dishes were placed on ice while the culture medium from each well was transferred into separate glass 12 X 75 test tubes. The tubes containing the culture media were placed in a boiling water bath for 10 minutes, then centrifuged at 1100 X g at 4°C. The supernatants were transferred to clean 12 X 75 glass test tubes and stored at -20°C overnight. Progesterone levels were measured using the Serono Diagnostics Progesterone MAIA, product number 12274, distributed by Ciba Corning, Medfield, MA. The assay was done according to the instructions provided by the manufacturer, except that the culture medium samples and the progesterone for the standard curve were diluted in a solution of phosphate buffered saline containing 0.1% bovine serum albumin and 0.6% sodium azide, instead of the dilution buffer provided in the kit. The progesterone for

the standard curve was obtained from Cal Biochem (catalog number 5341), and was diluted to a concentration of 15 $\mu\text{g/ml}$ in 100% ethanol prior to use. The stock solution was stored at -70°C . The final concentrations of the
5 progesterone standard curve dilutions were the same as those recommended in the assay kit. The diluted progesterone standards were stable at 4°C for one week.

Several FSH-treated Y1 lines, stably transfected with the human FSH receptor, secreted increased levels of
10 progesterone relative to the controls (the same line not treated with FSH, untransfected Y1 cells, Y1 cells transfected with CLH3AXSV2NEOh α IVS). One nonclonal line, Y1HFSHR4-38, was chosen for hormone dose response analyses. The procedure used was the same as that described in the
15 preceding paragraph except that the cells were treated with a range of FSH concentrations, 0-100 mIU/ml. Each dose was done in duplicate. The amount of progesterone produced by the FSH-stimulated Y1 cells was normalized to 1×10^6 cells and plotted versus FSH concentration to generate a dose
20 response curve. The dose response curve is shown in Figure 6. In this particular experiment, a 25-fold increase in progesterone activity was observed at a dose of 20 mIU/ml human recombinant FSH. The linear range was from 2.5 to 20 mIU/ml FSH and the ED_{50} was 6.4 mIU/ml FSH. These results
25 demonstrate that FSH dose dependent increases in progesterone secretion were observed in Y1 cells stably transfected with the human recombinant FSH receptor. The human FSH receptor is therefore functionally active when expressed in mouse adrenal Y1 cells.

30

Cell populations derived from the CHOHFSHR4Q-13 or Y1HFSHR4-38 cell lines, or similar cell lines, can be used to develop sensitive in vitro bioassays for FSH and FSH-like substances. Other FSH in vitro bioassays which are
35 currently used, such as the granulosa cell aromatase bioassay and the Sertoli cell aromatase bioassay, require the establishment of primary cell cultures from rats each time an assay is performed. For the new Y1 cell and CHO

cell bioassays, stably transformed continuous cell lines are used. This enables the practice of in vitro bioassay methods superior to the current methods in simplicity, accuracy, and consistency.

SEQUENCE LISTING

INFORMATION FOR SEQ ID NO: 1

SEQUENCE CHARACTERISTICS

(A) LENGTH: 2179

(B) TYPE: Nucleic acid

(C) STRANDEDNESS: Double

(D) TOPOLOGY: Linear

MOLECULE TYPE: cDNA to mRNA

ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(F) TISSUE TYPE: Testis

IMMEDIATE SOURCE:

(A) LIBRARY: λ gt11 cDNA library, ClonTech #HL1010b

(B) CLONE: pHFSHR11-11, pHFSHR15-6

FEATURE:

(A) NAME/KEY: protein coding region

(B) LOCATION: 75 to 2159

SEQUENCE DESCRIPTION: SEQ ID NO:1:

TGTGGAGCTT CTGAGATCTG TGGAGGTTTT TCTCTGCAAA TGCAGGAAGA AATCAGGTGG	60
ATGGATGCAT AATT ATG GCC CTG CTC CTG GTC TCT TTG CTG GCA TTC CTG	110
Met Ala Leu Leu Leu Val Ser Leu Leu Ala Phe Leu	
-15 -10	
AGC TTG GGC TCA GGA TGT CAT CAT CGG ATC TGT CAC TGC TCT AAC AGG	158
Ser Leu Gly Ser Gly Cys His His Arg Ile Cys His Cys Ser Asn Arg	
-5 1 5 10	
GTT TTT CTC TGC CAA GAG AGC AAG GTG ACA GAG ATT CCT TCT GAC CTC	206
Val Phe Leu Cys Gln Glu Ser Lys Val Thr Glu Ile Pro Ser Asp Leu	
15 20 25	
CCG AGG AAT GCC ATT GAA CTG AGG TTT GTC CTC ACC AAG CTT CGA GTC	254
Pro Arg Asn Ala Ile Glu Leu Arg Phe Val Leu Thr Lys Leu Arg Val	
30 35 40	
ATC CAA AAA GGT GCA TTT TCA GGA TTT GGG GAC CTG GAG AAA ATA GAG	302
Ile Gln Lys Gly Ala Phe Ser Gly Phe Gly Asp Leu Glu Lys Ile Glu	
45 50 55	
ATC TCT CAG AAT GAT GTC TTG GAG GTG ATA GAG GCA GAT GTG TTC TCC	350
Ile Ser Gln Asn Asp Val Leu Glu Val Ile Glu Ala Asp Val Phe Ser	
60 65 70 75	
AAC CTT CCC AAA TTA CAT GAA ATT AGA ATT GAA AAG GCC AAC AAC CTG	398
Asn Leu Pro Lys Leu His Glu Ile Arg Ile Glu Lys Ala Asn Asn Leu	
80 85 90	

CTC	TAC	ATC	AAC	CCT	GAG	GCC	TTC	CAG	AAC	CTT	CCC	AAC	CTT	CAA	TAT	446
Leu	Tyr	Ile	Asn	Pro	Glu	Ala	Phe	Gln	Asn	Leu	Pro	Asn	Leu	Gln	Tyr	
			95					100					105			
CTG	TTA	ATA	TCC	AAC	ACA	GGT	ATT	AAG	CAC	CTT	CCA	GAT	GTT	CAC	AAG	494
Leu	Leu	Ile	Ser	Asn	Thr	Gly	Ile	Lys	His	Leu	Pro	Asp	Val	His	Lys	
		110					115					120				
ATT	CAT	TCT	CTC	CAA	AAA	GTT	TTA	CTT	GAC	ATT	CAA	GAT	AAC	ATA	AAC	542
Ile	His	Ser	Leu	Gln	Lys	Val	Leu	Leu	Asp	Ile	Gln	Asp	Asn	Ile	Asn	
	125					130					135					
ATC	CAC	ACA	ATT	GAA	AGA	AAT	TCT	TTC	GTG	GGG	CTG	AGC	TTT	GAA	AGT	590
Ile	His	Thr	Ile	Glu	Arg	Asn	Ser	Phe	Val	Gly	Leu	Ser	Phe	Glu	Ser	
140					145					150					155	
GTG	ATT	CTA	TGG	CTG	AAT	AAG	AAT	GGG	ATT	CAA	GAA	ATA	CAC	AAC	TGT	638
Val	Ile	Leu	Trp	Leu	Asn	Lys	Asn	Gly	Ile	Gln	Glu	Ile	His	Asn	Cys	
				160				165						170		
GCA	TTC	AAT	GGA	ACC	CAA	CTA	GAT	GAG	CTG	AAT	CTA	AGC	GAT	AAT	AAT	686
Ala	Phe	Asn	Gly	Thr	Gln	Leu	Asp	Glu	Leu	Asn	Leu	Ser	Asp	Asn	Asn	
			175					180					185			
AAT	TTA	GAA	GAA	TTG	CCT	AAT	GAT	GTT	TTC	CAC	GGA	GCC	TCT	GGA	CCA	734
Asn	Leu	Glu	Glu	Leu	Pro	Asn	Asp	Val	Phe	His	Gly	Ala	Ser	Gly	Pro	
		190					195					200				
GTC	ATT	CTA	GAT	ATT	TCA	AGA	ACA	AGG	ATC	CAT	TCC	CTG	CCT	AGC	TAT	782
Val	Ile	Leu	Asp	Ile	Ser	Arg	Thr	Arg	Ile	His	Ser	Leu	Pro	Ser	Tyr	
	205					210					215					
GGC	TTA	GAA	AAT	CTT	AAG	AAG	CTG	AGG	GCC	AGG	TCG	ACT	TAC	AAC	TTA	830
Gly	Leu	Glu	Asn	Leu	Lys	Lys	Leu	Arg	Ala	Arg	Ser	Thr	Tyr	Asn	Leu	
220					225					230					235	
AAA	AAG	CTG	CCT	ACT	CTG	GAA	AAG	CTT	GTC	GCC	CTC	ATG	GAA	GCC	AGC	878
Lys	Lys	Leu	Pro	Thr	Leu	Glu	Lys	Leu	Val	Ala	Leu	Met	Glu	Ala	Ser	
				240					245					250		
CTC	ACC	TAT	CCC	AGC	CAT	TGC	TGT	GCC	TTT	GCA	AAC	TGG	AGA	CGG	CAA	926
Leu	Thr	Tyr	Pro	Ser	His	Cys	Cys	Ala	Phe	Ala	Asn	Trp	Arg	Arg	Gln	
			255					260					265			
ATC	TCT	GAG	CTT	CAT	CCA	ATT	TGC	AAC	AAA	TCT	ATT	TTA	AGG	CAA	GAA	974
Ile	Ser	Glu	Leu	His	Pro	Ile	Cys	Asn	Lys	Ser	Ile	Leu	Arg	Gln	Glu	
		270					275					280				
GTT	GAT	TAT	ATG	ACT	CAG	ACT	AGG	GGT	CAG	AGA	TCC	TCT	CTG	GCA	GAA	1022
Val	Asp	Tyr	Met	Thr	Gln	Thr	Arg	Gly	Gln	Arg	Ser	Ser	Leu	Ala	Glu	
	285					290					295					
GAC	AAT	GAG	TCC	AGC	TAC	AGC	AGA	GGA	TTT	GAC	ATG	ACG	TAC	ACT	GAG	1070
Asp	Asn	Glu	Ser	Ser	Tyr	Ser	Arg	Gly	Phe	Asp	Met	Thr	Tyr	Thr	Glu	
300					305					310					315	

TTT GAC TAT GAC TTA TGC AAT GAA GTG GTT GAC GTG ACC TGC TCC CCT	1118
Phe Asp Tyr Asp Leu Cys Asn Glu Val Val Asp Val Thr Cys Ser Pro	
320 325 330	
AAG CCA GAT GCA TTC AAC CCA TGT GAA GAT ATC ATG GGG TAC AAC ATC	1166
Lys Pro Asp Ala Phe Asn Pro Cys Glu Asp Ile Met Gly Tyr Asn Ile	
335 340 345	
CTC AGA GTC CTG ATA TGG TTT ATC AGC ATC CTG GCC ATC ACT GGG AAC	1214
Leu Arg Val Leu Ile Trp Phe Ile Ser Ile Leu Ala Ile Thr Gly Asn	
350 355 360	
ATC ATA GTG CTA GTG ATC CTA ACT ACC AGC CAA TAT AAA CTC ACA GTC	1262
Ile Ile Val Leu Val Ile Leu Thr Thr Ser Gln Tyr Lys Leu Thr Val	
365 370 375	
CCC AGG TTC CTT ATG TGC AAC CTG GCC TTT GCT GAT CTC TGC ATT GGA	1310
Pro Arg Phe Leu Met Cys Asn Leu Ala Phe Ala Asp Leu Cys Ile Gly	
380 385 390 395	
ATC TAC CTG CTG CTC ATT GCA TCA GTT GAT ATC CAT ACC AAG AGC CAA	1358
Ile Tyr Leu Leu Leu Ile Ala Ser Val Asp Ile His Thr Lys Ser Gln	
400 405 410	
TAT CAC AAC TAT GCC ATT GAC TGG CAA ACT GGG GCA GGC TGT GAT GCT	1406
Tyr His Asn Tyr Ala Ile Asp Trp Gln Thr Gly Ala Gly Cys Asp Ala	
415 420 425	
GCT GGC TTT TTC ACT GTC TTT GCC AGT GAG CTG TCA GTC TAC ACT CTG	1454
Ala Gly Phe Phe Thr Val Phe Ala Ser Glu Leu Ser Val Tyr Thr Leu	
430 435 440	
ACA GCT ATC ACC TTG GAA AGA TGG CAT ACC ATC ACG CAT GCC ATG CAG	1502
Thr Ala Ile Thr Leu Glu Arg Trp His Thr Ile Thr His Ala Met Gln	
445 450 455	
CTG GAC TGC AAG GTG CAG CTC CGC CAT GCT GCC AGT GTC ATG GTG ATG	1550
Leu Asp Cys Lys Val Gln Leu Arg His Ala Ala Ser Val Met Val Met	
460 465 470 475	
GGC TGG ATT TTT GCT TTT GCA GCT GCC CTC TTT CCC ATC TTT GGC ATC	1598
Gly Trp Ile Phe Ala Phe Ala Ala Ala Leu Phe Pro Ile Phe Gly Ile	
480 485 490	
AGC AGC TAC ATG AAG GTG AGC ATC TGC CTG CCC ATG GAT ATT GAC AGC	1646
Ser Ser Tyr Met Lys Val Ser Ile Cys Leu Pro Met Asp Ile Asp Ser	
495 500 505	
CCT TTG TCA CAG CTG TAT GTC ATG TCC CTC CTT GTG CTC AAT GTC CTG	1694
Pro Leu Ser Gln Leu Tyr Val Met Ser Leu Leu Val Leu Asn Val Leu	
510 515 520	
GCC TTT GTG GTC ATC TGT GGC TGC TAT ATC CAC ATC TAC CTC ACA GTG	1742
Ala Phe Val Val Ile Cys Gly Cys Tyr Ile His Ile Tyr Leu Thr Val	
525 530 535	

CGG AAC CCC AAC ATC GTG TCC TCC TCT AGT GAC ACC AGG ATC GCC AAG 1790
Arg Asn Pro Asn Ile Val Ser Ser Ser Ser Asp Thr Arg Ile Ala Lys
540 545 550 555

CGC ATG GCC ATG CTC ATC TTC ACT GAC TTC CTC TGC ATG GCA CCC ATT 1838
Arg Met Ala Met Leu Ile Phe Thr Asp Phe Leu Cys Met Ala Pro Ile
560 565 570

TCT TTC TTT GCC ATT TCT GCC TCC CTC AAG GTG CCC CTC ATC ACT GTG 1886
Ser Phe Phe Ala Ile Ser Ala Ser Leu Lys Val Pro Leu Ile Thr Val
575 580 585

TCC AAA GCA AAG ATT CTG CTG GTT CTG TTT CAC CCC ATC AAC TCC TGT 1934
Ser Lys Ala Lys Ile Leu Leu Val Leu Phe His Pro Ile Asn Ser Cys
590 595 600

GCC AAC CCC TTC CTC TAT GCC ATC TTT ACC AAA AAC TTT CGC AGA GAT 1982
Ala Asn Pro Phe Leu Tyr Ala Ile Phe Thr Lys Asn Phe Arg Arg Asp
605 610 615

TTC TTC ATT CTG CTG AGC AAG TGT GGC TGC TAT GAA ATG CAA GCC CAA 2030
Phe Phe Ile Leu Leu Ser Lys Cys Gly Cys Tyr Glu Met Gln Ala Gln
620 625 630 635

ATT TAT AGG ACA GAA ACT TCA TCC ACT GTC CAC AAC ACC CAT CCA AGG 2078
Ile Tyr Arg Thr Glu Thr Ser Ser Thr Val His Asn Thr His Pro Arg
640 645 650

AAT GGC CAC TGC TCT TCA GCT CCC AGA GTC ACC AAT GGT TCC ACT TAC 2126
Asn Gly His Cys Ser Ser Ala Pro Arg Val Thr Asn Gly Ser Thr Tyr
655 660 665

ATA CTT GTC CCT CTA AGT CAT TTA GCC CAA AAC TAAACACAA TGTGAAAATG 2179
Ile Leu Val Pro Leu Ser His Leu Ala Gln Asn
670 675

INFORMATION FOR SEQ ID NO: 2

SEQUENCE CHARACTERISTICS

(A) LENGTH: 695

(B) TYPE: Amino acid

(D) TOPOLOGY: Linear

MOLECULE TYPE: protein

FEATURE:

(A) NAME/KEY: signal sequence

(B) LOCATION: -17 to -1

(C) IDENTIFICATION METHOD: hydrophobic

FEATURE:

(A) NAME/KEY: putative amino-terminal extracellular domain

(B) LOCATION: 1 to 349

(C) IDENTIFICATION METHOD: similarity with other
dimeric glycoprotein receptor extracellular
domains, hydrophilic

FEATURE:

- (A) NAME/KEY: transmembrane domain
- (B) LOCATION: 350 to 613
- (C) IDENTIFICATION METHOD: similarity to other G
protein-coupled receptor transmembrane domains

FEATURE:

- (A) NAME/KEY: putative transmembrane region I
- (B) LOCATION: 350 to 370
- (C) IDENTIFICATION METHOD: similarity to other G
protein-coupled receptor transmembrane regions,
hydrophobic, about 20-23 amino acids in length

FEATURE:

- (A) NAME/KEY: putative transmembrane region II
- (B) LOCATION: 382 to 404
- (C) IDENTIFICATION METHOD: similarity to other G
protein-coupled receptor transmembrane regions,
hydrophobic, about 20-23 amino acids in length

FEATURE:

- (A) NAME/KEY: putative transmembrane region III
- (B) LOCATION: 427 to 448
- (C) IDENTIFICATION METHOD: similarity to other G
protein-coupled receptor transmembrane regions,
hydrophobic, about 20-23 amino acids in length

FEATURE:

- (A) NAME/KEY: putative transmembrane region IV
- (B) LOCATION: 469 to 491
- (C) IDENTIFICATION METHOD: similarity to other G
protein-coupled receptor transmembrane regions,
hydrophobic, about 20-23 amino acids in length

FEATURE:

- (A) NAME/KEY: putative transmembrane region V
- (B) LOCATION: 512 to 533
- (C) IDENTIFICATION METHOD: similarity to other G
protein-coupled receptor transmembrane regions,
hydrophobic, about 20-23 amino acids in length

FEATURE:

- (A) NAME/KEY: putative transmembrane region VI
- (B) LOCATION: 557 to 580
- (C) IDENTIFICATION METHOD: similarity to other G
protein-coupled receptor transmembrane regions,
hydrophobic, about 20-23 amino acids in length

FEATURE:

- (A) NAME/KEY: putative transmembrane region VII
- (B) LOCATION: 592 to 613
- (C) IDENTIFICATION METHOD: similarity to other G
protein-coupled receptor transmembrane regions,
hydrophobic, about 20-23 amino acids in length

FEATURE:

- (A) NAME/KEY: putative carboxy-terminal intracellular
domain
- (B) LOCATION: 614 to 678

-30-

SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ala Leu Leu Leu Val Ser Leu Leu Ala Phe Leu Ser Leu Gly Ser
-15 -10 -5

Gly Cys His His Arg Ile Cys His Cys Ser Asn Arg Val Phe Leu Cys
1 5 10 15

Gln Glu Ser Lys Val Thr Glu Ile Pro Ser Asp Leu Pro Arg Asn Ala
20 25 30

Ile Glu Leu Arg Phe Val Leu Thr Lys Leu Arg Val Ile Gln Lys Gly
35 40 45

Ala Phe Ser Gly Phe Gly Asp Leu Glu Lys Ile Glu Ile Ser Gln Asn
50 55 60

Asp Val Leu Glu Val Ile Glu Ala Asp Val Phe Ser Asn Leu Pro Lys
65 70 75

Leu His Glu Ile Arg Ile Glu Lys Ala Asn Asn Leu Leu Tyr Ile Asn
80 85 90 95

Pro Glu Ala Phe Gln Asn Leu Pro Asn Leu Gln Tyr Leu Leu Ile Ser
100 105 110

Asn Thr Gly Ile Lys His Leu Pro Asp Val His Lys Ile His Ser Leu
115 120 125

Gln Lys Val Leu Leu Asp Ile Gln Asp Asn Ile Asn Ile His Thr Ile
130 135 140

Glu Arg Asn Ser Phe Val Gly Leu Ser Phe Glu Ser Val Ile Leu Trp
145 150 155

Leu Asn Lys Asn Gly Ile Gln Glu Ile His Asn Cys Ala Phe Asn Gly
160 165 170 175

Thr Gln Leu Asp Glu Leu Asn Leu Ser Asp Asn Asn Asn Leu Glu Glu
180 185 190

Leu Pro Asn Asp Val Phe His Gly Ala Ser Gly Pro Val Ile Leu Asp
195 200 205

Ile Ser Arg Thr Arg Ile His Ser Leu Pro Ser Tyr Gly Leu Glu Asn
210 215 220

Leu Lys Lys Leu Arg Ala Arg Ser Thr Tyr Asn Leu Lys Lys Leu Pro
225 230 235

Thr Leu Glu Lys Leu Val Ala Leu Met Glu Ala Ser Leu Thr Tyr Pro
240 245 250 255

Ser His Cys Cys Ala Phe Ala Asn Trp Arg Arg Gln Ile Ser Glu Leu
260 265 270

His Pro Ile Cys Asn Lys Ser Ile Leu Arg Gln Glu Val Asp Tyr Met
275 280 285

Thr Gln Thr Arg Gly Gln Arg Ser Ser Leu Ala Glu Asp Asn Glu Ser
290 295 300

Ser Tyr Ser Arg Gly Phe Asp Met Thr Tyr Thr Glu Phe Asp Tyr Asp
305 310 315

Leu Cys Asn Glu Val Val Asp Val Thr Cys Ser Pro Lys Pro Asp Ala
320 325 330 335

Phe Asn Pro Cys Glu Asp Ile Met Gly Tyr Asn Ile Leu Arg Val Leu
340 345 350

Ile Trp Phe Ile Ser Ile Leu Ala Ile Thr Gly Asn Ile Ile Val Leu
355 360 365

Val Ile Leu Thr Thr Ser Gln Tyr Lys Leu Thr Val Pro Arg Phe Leu
370 375 380

Met Cys Asn Leu Ala Phe Ala Asp Leu Cys Ile Gly Ile Tyr Leu Leu
385 390 395

Leu Ile Ala Ser Val Asp Ile His Thr Lys Ser Gln Tyr His Asn Tyr
400 405 410 415

Ala Ile Asp Trp Gln Thr Gly Ala Gly Cys Asp Ala Ala Gly Phe Phe
420 425 430

Thr Val Phe Ala Ser Glu Leu Ser Val Tyr Thr Leu Thr Ala Ile Thr
435 440 445

Leu Glu Arg Trp His Thr Ile Thr His Ala Met Gln Leu Asp Cys Lys
450 455 460

Val Gln Leu Arg His Ala Ala Ser Val Met Val Met Gly Trp Ile Phe
465 470 475

Ala Phe Ala Ala Ala Leu Phe Pro Ile Phe Gly Ile Ser Ser Tyr Met
480 485 490 495

Lys Val Ser Ile Cys Leu Pro Met Asp Ile Asp Ser Pro Leu Ser Gln
500 505 510

Leu Tyr Val Met Ser Leu Leu Val Leu Asn Val Leu Ala Phe Val Val
515 520 525

Ile Cys Gly Cys Tyr Ile His Ile Tyr Leu Thr Val Arg Asn Pro Asn
530 535 540

Ile Val Ser Ser Ser Ser Asp Thr Arg Ile Ala Lys Arg Met Ala Met
545 550 555

Leu Ile Phe Thr Asp Phe Leu Cys Met Ala Pro Ile Ser Phe Phe Ala
560 565 570 575

2106061

-32-

Ile Ser Ala Ser Leu Lys Val Pro Leu Ile Thr Val Ser Lys Ala Lys
580 585 590

Ile Leu Leu Val Leu Phe His Pro Ile Asn Ser Cys Ala Asn Pro Phe
595 600 605

Leu Tyr Ala Ile Phe Thr Lys Asn Phe Arg Arg Asp Phe Phe Ile Leu
610 615 620

Leu Ser Lys Cys Gly Cys Tyr Glu Met Gln Ala Gln Ile Tyr Arg Thr
625 630 635

Glu Thr Ser Ser Thr Val His Asn Thr His Pro Arg Asn Gly His Cys
640 645 650 655

Ser Ser Ala Pro Arg Val Thr Asn Gly Ser Thr Tyr Ile Leu Val Pro
660 665 670

Leu Ser His Leu Ala Gln Asn
675

We claim:

1. In a method of assaying human FSH using a molecule which binds human FSH, the improvement wherein said molecule is a polypeptide capable of binding FSH which includes a contiguous portion of the extracellular domain of the human FSH receptor, said domain having the sequence of amino acids 1-349 of SEQ ID NO:2, which portion retains the same FSH-binding characteristics as the complete extracellular domain and has at least 95% identity with SEQ ID NO:2, or a polypeptide which includes a mutein of said portion, the sequence of said mutein differing from the sequence of said portion by substitutions, deletions or insertions affecting a total of no more than ten amino acids, and which mutein retains the same FSH-binding characteristic as said portion.
2. A method in accordance with claim 1, wherein said polypeptide is a polypeptide capable of binding FSH which includes a contiguous portion of said extracellular domain of the human FSH receptor, which portion retains the same FSH-binding characteristics as the complete extracellular domain and has at least 95% identity with SEQ ID NO:2.
3. A method in accordance with claim 1, wherein said polypeptide includes amino acid sequence 1-349 of SEQ ID NO:2 or a mutein differing therefrom by substitutions, deletions or insertions affecting a total of no more than ten amino acids, which mutein retains the same FSH-binding characteristics as said sequence.

4. A method in accordance with claim 3, wherein said polypeptide includes amino acid sequence 1-349 of SEQ ID NO:2.

5. A pure polypeptide capable of binding human follicle stimulating hormone (FSH), having a sequence comprising the sequence of:

(a) the human FSH receptor having the sequence of amino acids 1-678 of SEQ ID NO:2;

(b) a mutein which differs from the sequence of (a) by substitutions, deletions or insertions affecting a total of no more than ten amino acids;

(c) the amino-terminal extracellular portion of the human FSH receptor, having the sequence of amino acids 1-349 of SEQ ID NO:2;

(d) a mutein which differs from the sequence of (c) by substitutions, deletions or insertions affecting a total of no more than ten amino acids;

(e) a fragment of (c) which comprises a substantial portion of said amino-terminal extracellular portion of the human FSH receptor and which retains the same FSH-binding characteristics as the complete extracellular portion and has at least 95% identity with SEQ ID NO:2.

6. A polypeptide capable of binding human FSH, in accordance with claim 5, having a sequence comprising the sequence of the polypeptides of (a) or (b).
7. A pharmaceutical composition comprising a polypeptide in accordance with claim 6 and a pharmaceutically acceptable carrier.
8. A polypeptide capable of binding human FSH, in accordance with claim 5, comprising the amino acid sequence 1-613 of SEQ ID NO:2 or a mutein which differs therefrom by substitutions, deletions or insertions affecting a total of no more than ten amino acids.
9. A pharmaceutical composition comprising a polypeptide in accordance with claim 8 and a pharmaceutically acceptable carrier.
10. A polypeptide capable of binding human FSH, in accordance with claim 5, having a sequence comprising the sequence of the polypeptides of (c) or (d).
11. A pharmaceutical composition comprising a polypeptide in accordance with claim 10 and a pharmaceutically acceptable carrier.
12. A polypeptide capable of binding human FSH, in accordance with claim 5, having a sequence comprising the sequence of the polypeptides of (e).
13. A pharmaceutical composition comprising a polypeptide in accordance with claim 12 and a pharmaceutically acceptable carrier.

14. A pharmaceutical composition comprising a polypeptide in accordance with claim 5 and a pharmaceutically acceptable carrier.

15. In a method of assaying human FSH using a molecule which binds human FSH, the improvement wherein said molecule is a polypeptide in accordance with claim 5.

16. A pure polypeptide capable of binding human follicle stimulating hormone (FSH), consisting of a polypeptide capable of binding FSH which includes a contiguous portion of the extracellular domain of the human FSH receptor, said domain having the sequence of amino acids 1-349 of SEQ ID NO:2, which portion retains the same FSH-binding characteristics as the complete extracellular domain and has at least 95% identity with SEQ ID NO:2, or a polypeptide which includes a mutein of said portion, the sequence of said mutein differing therefrom by substitutions, deletions or insertions affecting a total of no more than ten amino acids, which mutein retains the same FSH-binding characteristics as said portion.

17. A pharmaceutical composition comprising a polypeptide in accordance with claim 16 and a pharmaceutically acceptable carrier.

18. Isolated DNA encoding a polypeptide which binds to human FSH, comprising:

(a) the nucleotide sequence of SEQ ID NO: 1;

(b) a mutant thereof which differs from (a) by substitution, deletion and/or insertion of nucleotides which alter a total of no more than ten amino acids in

the polypeptide encoded by (a), which mutant encodes a polypeptide which binds human FSH; or

(c) a fragment of (a) or (b) encoding a polypeptide which binds human FSH and while polypeptide has at least 95% identity with SEQ ID NO:2.

19. A recombinant expression vector comprising the DNA of claim 18.

20. A cell transfected with the expression vector of claim 19.

21. The cell of claim 20 which is a mammalian cell.

22. A method of producing human FSH receptor, or a fragment or mutant thereof which binds FSH, which comprises culturing the cell of claim 20 in a nutrient medium and recovering the receptor therefrom.

23. The method of claim 22 wherein the cell is a mammalian cell.

24. The isolated DNA of claim 18, comprising the nucleotide sequence of SEQ ID NO:1 or a fragment thereof which encodes a protein that binds human FSH and wherein the protein has at least 95% identity with SEQ ID NO:2.

25. The isolated DNA of claim 24 comprising the nucleotide sequence of Sequence ID No:1.

26. A recombinant expression vector comprising the DNA of claim 25.

27. A cell transfected with the expression vector of claim 26.

28. A method of producing human FSH receptor which comprises culturing the cell of claim 27 in a nutrient medium and recovering the receptor therefrom.

29. The isolated DNA of claim 24 comprising the nucleotide sequence of Sequence ID NO:1 in which the nucleotide sequence encoding the cytoplasmic domain of the FSH receptor is deleted.

30. A recombinant expression vector comprising the DNA of claim 29.

31. A cell transfected with the expression vector of claim 30.

32. A method of producing a human FSH receptor portion which binds FSH, which comprises culturing the cell of claim 31 in a nutrient medium under conditions that result in the expression of said receptor portion and recovering said human FSH receptor portion.

33. The isolated DNA of claim 24 comprising the nucleotide sequence of Sequence ID NO:1 in which the nucleotide sequence encoding the cytoplasmic and transmembrane domains of the FSH receptor are deleted.

34. A recombinant expression vector comprising the DNA of claim 33.

35. A cell transfected with the expression vector of claim 34.

36. A method of producing a human FSH receptor portion which binds FSH, which comprises culturing the cell of claim 35 in a nutrient medium and recovering the receptor therefrom.

37. An isolated DNA encoding a polypeptide which binds to human FSH, comprising a nucleotide sequence encoding:

(a) the human FSH receptor having SEQ ID NO: 2;

(b) a polypeptide which includes the amino-terminal extracellular portion of the human FSH receptor, said portion having the sequence of amino acids 1-349 of SEQ ID NO: 2; or

(c) a polypeptide which includes a fragment of the human FSH receptor which comprises a substantial portion of said amino-terminal extracellular portion of the human FSH receptor and which retains the same FSH-binding characteristics as the complete extracellular portion and has at least 95% identity with SEQ ID NO:2.

38. An isolated DNA in accordance with claim 37, comprising a nucleotide sequence encoding the human FSH receptor of SEQ ID NO:2.

39. A recombinant expression vector comprising the DNA of claim 38.

40. A cell transfected with the expression vector of claim 39.

41. A method of producing a human FSH receptor which comprises culturing the cell of claim 40 in a nutrient medium under conditions that result in the expression of said receptor and recovering said human FSH receptor.

42. An isolated DNA in accordance with claim 37, comprising a nucleotide sequence encoding a polypeptide of (b) or (c).

43. A recombinant expression vector comprising the DNA of claim 42.

44. A cell transfected with the expression vector of claim 43.

45. A method of producing a human FSH receptor portion which binds FSH, which comprises culturing the cell of claim 44 in a nutrient medium under conditions that result in the expression of said receptor portion and recovering said human FSH receptor portion.

46. A recombinant expression vector comprising the DNA of claim 37.

47. A cell transfected with the expression vector of claim 46.

48. A method of producing a human FSH receptor or a portion thereof which binds FSH, which comprises

culturing the cell of claim 47 in a nutrient medium and recovering the receptor therefrom.

49. Use of a polypeptide capable of binding FSH which includes a contiguous portion of the extracellular domain of the human FSH receptor, said domain having the sequence of amino acids 1-349 of SEQ ID NO:2 which portion retains the same FSH-binding characteristics as the complete extracellular domain and has at least 95% identity with SEQ ID NO:2, or a polypeptide which includes a mutein of said portion which mutein retains at least 95% identity with said portion and which mutein retains the same FSH-binding characteristic as said portion to reduce endogenous human follicle stimulating hormone (FSH) bioactivity in a patient.

50. Use of claim 49, wherein said polypeptide is a polypeptide capable of binding FSH which includes a contiguous portion of the extracellular domain of the human FSH receptor, said domain having the sequence of amino acids 1-349 of SEQ ID NO:2 which portion retains the same FSH-binding characteristics as the complete extracellular domain and has at least 95% identity with SEQ ID NO:2.

51. Use of claim 49, wherein said mutein differs from said portion only by conservative substitutions in one to ten amino acid residues, the location and nature of said substitutions being selected so as not to degrade the FSH binding characteristics of said portion.

52. Use of claim 49, wherein said polypeptide includes

amino acid sequence 1-349 of SEQ ID NO:2 or a mutein which retains at least 95% identity with said sequence and retains the same FSH-binding characteristics as said sequence.

53. Use of claim 52, wherein said polypeptide includes amino acid sequence 1-349 of SEQ ID NO:2.

54. Use of a polypeptide capable of binding human follicle stimulating hormone (FSH) which includes a contiguous portion of the extracellular domain of the human FSH receptor, said domain having the sequence of amino acids 1-349 of SEQ ID NO:2 which portion retains the same FSH-binding characteristics as the complete extracellular domain, or a polypeptide which includes a mutein of said portion which mutein retains at least 95% identity with said portion and which mutein retains the same FSH-binding characteristic as said portion to prevent follicle growth and maturation in a female patient.

55. Use of claim 54, wherein said polypeptide is a polypeptide capable of binding FSH which includes a contiguous portion of the extracellular domain of the human FSH receptor, said domain having the sequence of amino acids 1-349 of SEQ ID NO:2 which portion retains the same FSH-binding characteristics as the complete extracellular domain and has at least 95% identity with SEQ ID NO:2.

56. Use of claim 54, wherein said mutein differs from said portion only by conservative substitutions in one to

ten amino acid residues, the location and nature of said substitutions being selected so as not to degrade the FSH binding characteristics of said portion.

57. Use of claim 54, wherein said polypeptide includes amino acid sequence 1-349 of SEQ ID NO:2 or a mutein which retains at least 95% identity with said sequence and retains the same FSH-binding characteristics as said sequence.

58. Use of claim 54, wherein said polypeptide includes amino acid sequence 1-349 of SEQ ID NO:2.

59. Use of a polypeptide capable of binding human follicle stimulating hormone (FSH) which includes a contiguous portion of the extracellular domain of the human FSH receptor, said domain having the sequence of amino acids 1-349 of SEQ ID NO:2 which portion retains the same FSH-binding characteristics as the complete extracellular domain and has at least 95% identity with SEQ ID NO:2, or a polypeptide which includes a mutein of said portion which mutein retains at least 95% identity with said portion and which mutein retains the same FSH-binding characteristic as said portion to preventing spermatogenesis in a male patient.

60. Use of claim 59, wherein said polypeptide is a polypeptide capable of binding FSH which includes a contiguous portion of the extracellular domain of the human FSH receptor, which portion retains the same FSH-binding characteristics as the complete extracellular domain and has at least 95% identity with SEQ ID NO:2.

61. Use of claim 59, wherein said mutein differs from said portion only by conservative substitutions in one to ten amino acid residues, the location and nature of said substitutions being selected so as not to degrade the FSH binding characteristics of said portion.

62. Use of claim 59, wherein said polypeptide includes amino acid sequence 1-349 of SEQ ID NO:2 or a mutein which retains at least 95% identity with said sequence and retains the same FSH-binding characteristics as said sequence.


63. Use of claim 59, wherein said polypeptide includes amino acid sequence 1-349 of SEQ ID NO:2.

2106061

PCT/US
50/US

92 / 00122
29 SEP 1992


FIG. 1



MAP KEY:
(N.N kb) SIZE OF INSERTS AS ESTIMATED BY AGAROSE GEL ELECTROPHORESIS
□ 0.25 kb INSERTION, PROBABLY AN INTRON OR A PORTION OF AN INTRON

2106061

2 / 7




2106061

PCT/US 92 / 00122
POUS 29 SEP 1992

3 / 7

FIG. 2B




SUBSTITUTE SHEET

2106061

4 / 7


FIG. 3



2106061

5 / 7


FIG. 4



2106061

6 / 7

FIG. 5



2106061 7 / 7

FIG. 6

