

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2017248731 B2**

(54) Title
Compositions for treatment of wet age-related macular degeneration

(51) International Patent Classification(s)
C07K 16/22 (2006.01) **C12N 15/86** (2006.01)
A61K 48/00 (2006.01)

(21) Application No: **2017248731** (22) Date of Filing: **2017.04.14**

(87) WIPO No: **WO17/180936**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/331,100	2016.05.03	US
62/442,946	2017.01.05	US
62/323,184	2016.04.15	US
62/460,515	2017.02.17	US
62/466,721	2017.03.03	US

(43) Publication Date: **2017.10.19**

(44) Accepted Journal Date: **2024.08.29**

(71) Applicant(s)
The Trustees of The University of Pennsylvania

(72) Inventor(s)
Tretiakova, Anna P.;Wilson, James M.

(74) Agent / Attorney
WRAYS PTY LTD, L7 863 Hay St, Perth, WA, 6000, AU

(56) Related Art
US 2013/0090375 A1



(51) International Patent Classification:

A61K 48/00 (2006.01) C07K 16/22 (2006.01)
C12N 15/86 (2006.01)

(21) International Application Number:

PCT/US2017/027529

(22) International Filing Date:

14 April 2017 (14.04.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/323,184	15 April 2016 (15.04.2016)	US
62/331,100	3 May 2016 (03.05.2016)	US
62/442,946	5 January 2017 (05.01.2017)	US
62/460,515	17 February 2017 (17.02.2017)	US
62/466,721	3 March 2017 (03.03.2017)	US

(71) Applicant: THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; 3160 Chestnut Street, Suite 200, Philadelphia, PA 19104 (US).

(72) Inventors: TRETIAKOVA, Anna, P.; 2531 Parrish Street, Philadelphia, PA 19103 (US). WILSON, James, M.; 1831 Delancey Street, Philadelphia, PA 19103 (US).

(74) Agents: KODROFF, Cathy, A. et al.; Howson & Howson LLP, 350 Sentry Parkway, Building 620, Suite 210, Blue Bell, PA 19422 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: COMPOSITIONS FOR TREATMENT OF WET AGE-RELATED MACULAR DEGENERATION




FIG 1

(57) Abstract: A recombinant adeno-associated virus (rAAV) having an AAV8 capsid which is suitable for intra-retinal injection is provided herein. The rAAV comprises a vector genome packaged within the capsid which contains, operably linked to regulatory elements which direct expression of anti-human vascular endothelial growth factor (VEGF) antigen binding antibody fragment (aVEGF), a coding sequence for aVEGF, wherein the coding sequence is operably linked to regulatory elements which direct expression of the anti-VEGF Fab in the eye. Also provided herein are liquid suspensions containing these rAAV8.aVEGF and methods of using same for treatment of wet AMD and other ocular conditions.

COMPOSITIONS FOR TREATMENT OF WET AGE-RELATED MACULAR DEGENERATION

INCORPORATION-BY-REFERENCE OF MATERIAL SUBMITTED IN ELECTRONIC 5 FORM

Applicant hereby incorporates by reference the Sequence Listing material filed in electronic form herewith. This file is labeled "UPN-16-7683PCT_ST25.txt".

BACKGROUND OF THE INVENTION

10 Age-related macular degeneration (AMD) is a progressive degenerative macular disease attacking the region of highest visual acuity (VA), the macula, and is the leading cause of blindness in Americans 60 years or older (NIH Medline Plus (2008), Leading cause of blindness, NIH Medline Plus 3(2) 14-15. www.nlm.nih.gov/medlineplus/magazine/issues/summer08/articles/summer08pg14-15.html). The neovascular “wet” form of the
15 disease (nAMD or wet AMD) is characterized by choroidal neovascularization which is marked by proliferation of blood vessels and cells including those of the retinal pigment epithelium (RPE) (Carmeliet (2005) Nature 438: 932–936). Ultimately, photoreceptor death and scar formation result in a severe loss of central vision and the inability to read, write, and recognize faces or drive. Many patients can no longer maintain gainful employment, carry
20 out daily activities and consequently report a diminished quality of life (Mitchell and Bradley (2006), Health Qual Life Outcomes 4: 97). Preventative therapies have demonstrated little effect and therapeutic strategies have focused primarily on treating the neovascular lesion.

Some currently available treatments for wet AMD include laser photocoagulation,
25 photodynamic therapy with verteporfin, and intravitreal (IVT) injections with the vascular endothelial growth factor (VEGF) inhibitors such as pegaptanib, ranibizumab, bevacizumab or aflibercept (Schmidt-Erfurth, (2014) Guidelines for the management of neovascular age-related macular degeneration by the European Society of Retina Specialists (EURETINA) Br J Ophthalmol 98:1144–1167). While these therapies have some effect on best-corrected
30 visual acuity (BCVA), their effects may be limited in restoring visual acuity and in duration

injection. The AAV8 capsid packages a vector genome that provides for the production of a soluble antigen-binding fragment (Fab) of a human monoclonal antibody (MAb) that binds and inhibits human vascular endothelial growth factor (hVEGF) – the expression product is sometimes referred to herein as “anti-hVEGF Fab” or “aVEGF”.

According to a further aspect, the present invention provides a recombinant adeno-associated virus (rAAV) which is suitable for administration to an eye of a human subject wherein the rAAV comprises an expression cassette in an AAV8 capsid, wherein the expression cassette comprises:

a coding sequence for an anti-human vascular endothelial growth factor (hVEGF) antigen binding antibody fragment (Fab) (anti-hVEGF Fab) operably linked to regulatory elements which direct expression thereof, wherein coding sequence comprises:

(i) a nucleotide sequence encoding an anti-hVEGF Fab heavy immunoglobulin chain (HC) having a first interleukin 2 (IL-2) leader sequence operably linked to an anti-hVEGF Fab HC variable region comprising amino acids 1 to 123 of SEQ ID NO: 1,

(ii) a nucleotide sequence encoding an F2A linker,

(iii) a nucleotide sequence encoding anti-hVEGF Fab light chain (LC) having a second IL-2 leader sequence operably linked to nucleotides encoding an anti-hVEGF Fab LC variable region comprising amino acids 1 to 107 of SEQ ID NO: 2; and

wherein the regulatory elements comprise a cytomegalovirus immediate early (CMV IE) enhancer, a chicken beta-actin promoter, and a chicken beta-actin intron.

According to a further aspect, the present invention provides a recombinant adeno-associated virus (rAAV) having an AAV8 capsid which is suitable for administration to a human eye, wherein the rAAV comprises an expression cassette in the AAV8 capsid, said expression cassette comprises nucleotides 198 to 3733 of SEQ ID NO: 14.

According to another aspect, the present invention provides a recombinant adeno-associated virus (rAAV) having an AAV8 capsid, wherein the rAAV comprises a vector genome in the AAV8 capsid, wherein the vector genome comprises a AAV2 5' inverted terminal repeat (ITR), an expression cassette comprising the nucleotides 198 to 3733 of SEQ ID NO: 14, and a AAV2 3' ITR.

According to another aspect, the present invention provides a recombinant adeno-associated virus (rAAV) having an AAV8 capsid which is suitable for administration to an eye,

wherein the rAAV comprises a vector genome packaged within the capsid, said vector genome comprising SEQ ID NO: 14 (ITR-CB7-CI-aVEGFv3-rBG-ITR).

According to a further aspect, the present invention provides a liquid suspension suitable for administration to a human eye, said liquid suspension comprising an aqueous liquid and recombinant adeno-associated virus (rAAV) according to any one of claims 1 to 17 and optionally one or more excipients, preservatives, and/or surfactants.

According to yet a further aspect, the present invention provides use of a rAAV according to the invention or a liquid suspension according to the invention in the preparation of a medicament for subretinal administration to a human subject.

According to another aspect, the present invention provides a recombinant adeno-associated virus (rAAV) according to the invention or the liquid suspension according to the invention when used in subretinal administration to a patient in need thereof.

According to another aspect, the present invention provides a product comprising: (a) a container comprising an rAAV according to the invention and an aqueous liquid, and optionally a second container comprising (b) a diluent, and (c) a needle for administration.

The vector genome packaged within the rAAV8 capsid, comprises: (a) an AAV inverted terminal repeat(s) (ITR(s)) flanking an expression construct for the anti-VEGF Fab; (b) the expression construct having regulatory elements comprising a chicken beta-actin promoter or a ubiquitin C promoter that direct expression in the eye of a transgene encoding anti-hVEGF Fab; and (c) the transgene which encodes the heavy and light chains of the anti-hVEGF Fab, each chain having a heterologous leader sequence added to its amino terminus, and wherein the coding sequences for the heavy and light chains are separated by a coding sequence for a “cleavable” peptide linker or an IRES (internal ribosome entry site) to ensure production of separate heavy and light chain polypeptides, and a polyadenylation signal. The resulting transgene expression products may contain an amino acid residue in addition to those normally found in Fab heavy chains.

In one embodiment, the present invention provides a recombinant adeno-associated virus (rAAV) having an AAV8 capsid which is suitable for subretinal and/or intra-retinal injection, wherein the rAAV comprises a vector genome packaged within the capsid, said vector genome comprising: (a) an AAV inverted terminal repeat (ITR); (b) a coding sequence for an anti-human vascular endothelial growth factor (VEGF) antigen binding antibody fragment (Fab) having an exogenous leader sequence, a heavy immunoglobulin chain, a linker, and a light immunoglobulin chain having an exogenous leader sequence, wherein the coding sequence is operably linked to

regulatory elements which direct expression of the anti-VEGF Fab exogenous leader sequence, heavy immunoglobulin chain, linker, exogenous leader sequence and light immunoglobulin chain in the eye; (c) the regulatory elements of (b), which comprise a promoter selected from a chicken beta-actin promoter or a ubiquitin C promoter; and (d) an AAV ITR.

In particular embodiments, codon sequences for the heavy and light chains optimized for expression in human cells are used. As illustrated by the examples, these can include but are not limited to AAV2/8.CB7.Cl.aVEGFv1.rBG; AAV2/8.CB7.Cl.aVEGFv2.rBG; AAV2/8.CB7.Cl.aVEGFv3.rBG; AAV2/8.CB7.Cl.aVEGFv4.rBG; AAV2/8.CB7.Cl.aVEGFv5.rBG; AAV2/8.CB7.Cl.aVEGFv6.rBG; AAV2/8.CB7.Cl.aVEGFv7.rBG; AAV2/8.CB7.Cl.aVEGFv8.rBG; AAV2/8.CB7.Cl.aVEGFv9.rBG; AAV2/8.CB7.Cl.aVEGFv10.rBG; AAV2/8.CB7.Cl.aVEGFv11.rBG; AAV2/8.CB7.Cl.aVEGFv12.rBG; AAV2/8.CB7.Cl.aVEGFv13.rBG. As used herein, "AAV2/8" and "AAV8" are used interchangeably to refer to a recombinant AAV having an AAV8 capsid and vector genome flanked by AAV2 ITRs.

In yet another aspect, a liquid suspension of any of the foregoing rAAV8.aVEGF for sub-retinal and/or intra-retinal injection is provided. The composition comprises an aqueous liquid and rAAV8.aVEGF as described herein, and optionally one or more excipients, preservatives, and/or surfactants.

In still a further aspect, a method for delivering an anti-hVEGF Fab to a patient having wet age-related related macular degeneration is provided. The method involves subretinally injecting the patient's eye with the liquid suspension comprising the rAAV8 vector carrying the expression construct for the anti-hVEGF Fab.

In certain embodiments, the invention provides a rAAV as described herein, or a liquid suspension, administrable subretinally to a patient. In certain embodiments, use of a rAAV or a liquid suspension, for subretinal administration to a patient is provided. The

patient may have been previously diagnosed with wet age-related macular degeneration, or another ocular condition as defined herein.

In still a further embodiment, a product comprising: (a) a first container comprising an rAAV8.anti-hVEGF Fab and an aqueous liquid, (b) optionally a second container
5 comprising a diluent, and (c) a needle for injection. In certain embodiments, the product is an injection kit.

The invention is illustrated by the examples below which demonstrate that subretinal administration of an rAAV8.aVEGF vector results in gene transfer throughout the retina, and expression of anti-VEGF Fab throughout the retina and in the vitreous and anterior chamber
10 fluids. This result is surprising in view of prior art gene therapy studies that demonstrated that gene transfer spreads laterally outside of the original injection bleb but remains confined to those expanded boundaries and did not achieve gene transfer and transgene expression outside this expanded area of injection (the “bleb” formed in the retina at the injection site); and offers an advantage over standard of care treatment for nAMD in that a single
15 administration of the rAAV8.aVEGF vector should result in (i) continuous delivery of the effective amounts of the VEGF inhibitor throughout the retina which may in turn improve performance as compared to repeated IVT administrations of high dose boluses of the VEGF inhibitor that dissipate over time; and (ii) avoidance of repeated ocular injections which pose additional risks and inconvenience to patients. Each aspect may improve therapeutic
20 outcome.

Still other aspects and advantages of the invention will be apparent from the following detailed description of the invention.

BRIEF DESCRIPTION OF THE FIGURES

25 FIG 1 provides a schematic representation of an AAV8 vector genome containing a gene cassette flanked by the AAV2 inverted terminal repeat (ITRs) and expressing human anti-vascular endothelial growth factor (anti-VEGF) antigen binding antibody fragment (Fab). Control elements include the CB7 promoter consisting of the chicken β -actin promoter and CMV enhancer, chicken β -actin and a rabbit β -globin poly A signal. The nucleic acid
30 sequences coding for the heavy and light chains of anti-VEGF Fab are separated by a self-cleaving furin (F)/F2A linker. A furin recognition site that consists of arginine-lysine-arginine-arginine amino acid sequence was used. In addition, the light and heavy chain each

contain a heterologous leader peptide which directs nascent peptide into appropriate cellular compartment where leader peptide is processed away from the mature protein by the host cellular machinery. These and the other synthetic anti-VEGF constructs are termed herein, AAV.aVEGF.

5 FIG 2 provides the expression levels and kinetics of AAV8.CB7.aVEGFv1 or rAAV8.UbC.aVEGFv1 at various time points post-injection of the left eye (os) or right eye (od). AAV8.CB7.aVEGF-Rv1 is the top line with the closed circle. AAV8.UbC.aVEGF-Rv1 is the bottom line with the closed circle. The middle line with the open circles is AAV8.UbC.aVEGF-Rv1.

10 FIGs 3A - 3D show expression of anti-VEGF Fab in anterior chamber fluid and blood for animals in Groups 2 and 3 as described in Example 3 in which Cynomolgus monkeys were administered a single dose of 1.00×10^{11} GC/eye of AAV2/8 vectors into each eye subretinally. Anterior chamber fluid and blood were collected at prespecified timepoints. Expression of the anti-VEGF Fab was determined using enzyme-linked
15 immunosorbent assay. In FIGs 3A and 3B, the results for animals in Group 2. In FIGS 3C and 3D, the results for Group 3 are presented. In FIGS 3B and 3D, the gray area in the panel presenting the results in serum denotes baseline levels. Circles denote females and squares denote males. Samples were analyzed in duplicate. The results are presented as mean \pm standard deviation. Abbreviations: Fab = fragment antigen-binding; GC = genome
20 copies; OD = right eye; OS = left eye; VEGF = vascular endothelial growth factor.

 FIGs 4A - 4D show expression of anti-VEGF Fab in anterior chamber fluid and blood for animals in Groups 5 and 6 as described in Example 3 in which cynomolgus monkeys were administered a single dose of 1.00×10^{11} GC/eye of AAV2/8 vectors into each eye subretinally. Anterior chamber fluid and blood were collected at prespecified
25 timepoints. Expression of the anti-VEGF Fab was determined using enzyme-linked immunosorbent assay. The results for Group 5 are presented in FIGS 4A and 4B and the results for Group 6 is presented in FIGS 4C and 4D. In FIGS 4B and 4D, the gray area in the panel presenting the results in serum denotes baseline levels. Circles denote females and squares denote males. Samples were analyzed in duplicate. The results are presented as mean
30 \pm standard deviation. Abbreviations: Fab = fragment antigen-binding; GC = genome copies; OD = right eye; OS = left eye; VEGF = vascular endothelial growth factor.

FIGs 5A - 5D provide levels of mRNA for AAV8.aVEGF test vector in retina determined by RT-qPCR. Cynomolgus monkeys were administered a single dose of 1.00×10^{12} GC/eye of a AAV8.aVEGF test vector or FFB-314 into the right eye subretinally. Levels of mRNA for the AAV8.aVEGF test vector were determined in different portions of the dissected retinas by quantitative reverse transcription polymerase chain reaction (RT-qPCR). In left panels, schematics of injection sites are depicted. In middle panels, retinal dissections are presented. In right panels, levels of mRNA for AAV8.aVEGF test vector mRNA (GC per 100 ng of RNA) in 4 sections of retina are depicted. Abbreviations: BV = major blood vessel; F = fovea; GC = genome copies; IB = injection bleb; ID = identification; O = optic disk; UD = undetected.

FIGs 6A - 6D provide results of expression of anti-VEGF Fab in anterior chamber fluid, vitreous, and retina (Group 2, Example 6). Cynomolgus monkeys were administered a single dose of 1.00×10^{11} GC/eye of AAV2/8 vector subretinally. These data represent results from different AAV8.aVEGF vectors than shown in FIGs 5A - 5D. Concentrations of anti-VEGF Fab were determined in anterior chamber fluid, vitreous, and 4 different parts of retina. Eyes were dissected as described in FIGs 5A - 5D. In FIGs 6A and 6C, infrared spectral domain optical coherence tomography images of the retinas with boundaries of injection site are depicted. In FIGs 6B and 6D, graphs of concentrations of anti-VEGF Fab are presented. In this figure, the results for animals in Group 2 in Example 6 are presented. Abbreviations: ACF = anterior chamber fluid; BV = major blood vessel; F = fovea; Fab = fragment antigen-binding; FOV = middle section containing fovea; GC = genome copies; IB = injection bleb; ID = identification; INF = inferior retinal section; O = optic disk; ODI = middle section containing optic disk; SUP = superior retinal section; VEGF = vascular endothelial growth factor; VIT = vitreous.

FIGs 7A - 7D provide results of expression of anti-VEGF Fab in anterior chamber fluid, vitreous, and retina (Group 3, Example 6). Cynomolgus monkeys were administered a single dose of 1.00×10^{11} GC/eye of AAV2/8 vector subretinally. These data represent results from different AAV8.aVEGF vectors than shown in FIGs 5A - 5D. Concentrations of anti-VEGF Fab were determined in anterior chamber fluid, vitreous, and 4 different parts of retina. Eyes were dissected as described in FIGs 5A - 5D. In FIGs 7A and 7C, infrared spectral domain optical coherence tomography images of the retinas with boundaries of injection site are depicted. In the graphs of FIGs 7B and 7D, concentrations of anti-VEGF

Fab are presented. In this figure, the results for animals in Group 3 in Example 6 are presented. Abbreviations: ACF = anterior chamber fluid; BV = major blood vessel; F = fovea; Fab = fragment antigen-binding; FOV = middle section containing fovea; GC = genome copies; IB = injection bleb; ID = identification; INF = inferior retinal section; O = optic disk; ODI = middle section containing optic disk; SUP = superior retinal section; VEGF = vascular endothelial growth factor; VIT = vitreous.

FIGs 8A - 8D provide results of expression of anti-VEGF Fab in anterior chamber fluid, vitreous, and retina (Group 5, Example 6). Cynomolgus monkeys were administered a single dose of 1.00×10^{11} GC/eye of AAV2/8 vector subretinally. These data represent results from different AAV8.aVEGF vectors than shown in FIGs 5A - 5D. Concentrations of anti-VEGF Fab were determined in anterior chamber fluid, vitreous, and 4 different parts of retina. Eyes were dissected as described in FIGs 5A - 5D. In FIGS 8A and 8C, infrared spectral domain optical coherence tomography images of the retinas with boundaries of injection site are depicted. In the graphs of FIGs 8B and 8D, concentrations of anti-VEGF Fab are presented. In this figure, the results for animals in Group 5 in Example 6 are presented. Abbreviations: ACF = anterior chamber fluid; BV = major blood vessel; F = fovea; Fab = fragment antigen-binding; FOV = middle section containing fovea; GC = genome copies; IB = injection bleb; ID = identification; INF = inferior retinal section; O = optic disk; ODI = middle section containing optic disk; SUP = superior retinal section; VEGF = vascular endothelial growth factor; VIT = vitreous.

FIG 9 provides a flow diagram of the manufacturing process.

FIGs 10A - 10D illustrate the results of an rcAAV assay for AAV8. wtAAV8 is spiked into different GC amounts of AAV vector and the cap gene copy number per 1 μ g of 293 cell DNA is determined after three successive passages of the cell lysate onto fresh cells. 3 different spike levels of wtAAV8 [one level per panel: 1×10^2 GC, 1×10^3 GC and 1×10^4 GC] 4 different vector amounts [0 GC (dark square), 1×10^9 GC (gray square) 1×10^{10} GC (triangle) and 1×10^{11} GC (marked with X)] are shown and background levels are indicated (controls).

30 DETAILED DESCRIPTION OF THE INVENTION

Recombinant, replication-defective adeno-associated virus (rAAV) vectors having an AAV8 capsid and compositions containing same which are suitable for subretinal injections

to deliver an anti-VEFG antibody binding fragment (Fab). Also provided are compositions containing same, and in particularly, liquid aqueous suspension. Uses of these compositions are also provided.

The rAAV8 vectors are designed to express an anti-VEGF antibody binding
5 fragment (Fab) in mammalian, and more particularly, human cells. These anti-VEGF Fabs are particularly well suited for treatment of age-related macular degeneration (AMD). For convenience, these vectors are terms rAAV8.AMD. As described herein, a series of novel AAV8.aVEGF constructs have been developed which have demonstrated high yield, expression levels, and/or activity.

10 The invention is illustrated by the examples below which demonstrate that subretinal administration of an rAAV8.aVEGF vector results in gene transfer throughout the retina, and expression of anti-VEGF Fab throughout the retina and in the vitreous and anterior chamber fluids. This result is surprising in view of prior art gene therapy studies that demonstrated
15 that gene transfer spreads laterally outside of the original injection bleb but remains confined to those expanded boundaries and did not achieve gene transfer and transgene expression outside this expanded area of injection (the “bleb” formed in the retina at the injection site); and offers an advantage over standard of care treatment for nAMD in that a single administration of the rAAV8.aVEGF vector should result in (i) continuous delivery of the effective amounts of the VEGF inhibitor throughout the retina which may in turn improve
20 performance as compared to repeated IVT administrations of high dose boluses of the VEGF inhibitor that dissipate over time; and (ii) avoidance of repeated ocular injections which pose additional risks and inconvenience to patients. Each aspect may improve therapeutic outcome.

The present invention provides constructs encoding a novel anti-VEGF Fab having,
25 at a minimum, a heavy chain amino acid sequence of SEQ ID NO: 1 and a light chain amino acid sequence of SEQ ID NO:2, each of which has been engineered to have an exogenous leader sequence for each the heavy chain and light chain. In certain constructs illustrated herein, the leader sequence is derived from a human IL2 leader. Further, in certain constructs illustrated in the working examples, the heavy and light chains are separated by a
30 furin/F2a linker, which may result may result in one or more extra amino acids being added to the heavy chain [SEQ ID NO:1]. In one embodiment, a single arginine [R] is added to the heavy chain. However, in certain embodiments, another linker may be selected and/or a

different system may result in no additional amino acid, or one or more extra amino acids [e.g., R, Lys (K), RK, RKR, RKRR among others]. In prior provisional applications, the resulting constructs were termed herein, aVEGF-R. However, for clarity, these constructs encoding the anti-VEGF Fab transgene product described herein, are referred to as: anti-VEGF Fab, aVEGF, anti-hVEGF, anti-human VEGF, or anti-VEGF Fab transgene product. In the constructs encoding this transgene product, a numerical designation following the term aVEGF, e.g., aVEGFv1, aVEGFv2, aVEGFv3, through aVEGFv13, refers to different nucleic acid coding sequences for the open reading frame of the immunoglobulin heavy chain and light chains.

10 In certain embodiments, the amino acid sequence of the anti-VEGF Fab has 513 amino acids, including anti-VEGF is heavy and light chain separated by extra amino acids as a result of the linker. For example, while each of the following expression cassettes encodes the same anti-VEGF heavy chain and light chain, in one embodiment, there may be one amino acid added to the last position of the heavy chain. In still other embodiments, there
15 may be two, three, four or more extra amino acids attached to the heavy chain. For example, in certain embodiments, the nucleic acid sequences coding for the heavy and light chains of anti-VEGF Fab are separated by a self-cleaving furin (F)/F2A linker. A furin recognition site that consists of arginine-lysine-arginine-arginine amino acid sequence may be used. Due to the mechanism of furin-mediated cleavage, vector-expressed anti-VEGF Fab may contain
20 an additional arginine (R) residue added to the last position of the heavy chain [SEQ ID NO: 1]. In other embodiments, the vector-expressed anti-VEGF Fab may contain the dipeptide arginine-lysine at the end of the heavy chain, the tripeptide arginine-lysine-arginine at the end of the heavy chain, or the polypeptide arginine-lysine-arginine-arginine at the end of the heavy chain. In certain embodiments, the vector expressed anti-VEGF Fab are a
25 heterogeneous mixture of two or more of these Fab products. Other furin cleavage sites can be used (arginine-X-X-arginine, or arginine-X-lysine or arginine-arginine), which can also generate C-terminal heterogeneity. In other words, other vector expressed anti-VEGF Fabs may be a heterogeneous population of the Fab in which the heavy chain has 0, 1, 2, 3, or 4 amino acids at its C-terminus as a result of the linker processing. In addition, the light and
30 heavy chain each contain a heterologous leader peptide which directs nascent peptide into appropriate cellular compartment where leader peptide is processed away from the mature

protein by the host cellular machinery. In certain embodiments, the anti-VEGF Fab contains no HC or LC leader sequences. See, e.g., SEQ ID NO: 33.

In certain embodiments, the anti-VEGF Fab heavy chain has the amino acid sequence of residues 21 - 252 of SEQ ID NO: 33 with a leader sequence. In other
5 embodiments, the anti-VEGF Fab light chain has the amino acid sequence of residues 300 - 513 of SEQ ID NO: 33 with a leader sequence. For example, the leader sequence may be from about 15 to about 25 amino acids, preferably about 20 amino acids. In some
embodiments, the leader has the sequence of amino acids 1-20 of SEQ ID NO: 33.

In one embodiment, the coding sequences for the heavy chain and light chain of anti-
10 VEGFv1 are provided in SEQ ID NO: 24. More particularly, the heavy chain variable region open reading frame (ORF) is provided in nucleotides (nt) 1843 to 2211 and the heavy chain constant region (CH1) ORF is provided in nt 2212-2532, with reference to SEQ ID NO: 24. Thus, the aVEGFv1 heavy chain, without the leader, has the nucleic acid sequence of nt 1843 to 2532. The light chain variable region (VL) ORF is provided in nt 2680 to 3000
15 and the light chain constant region (CL) is provided in nt 3001 to 3321 of SEQ ID NO: 24. Thus, the aVEGFv2 light chain, without the leader, has the nucleic acid sequence of nt 2680 to 3321 of SEQ ID NO: 24.

In another embodiment, the coding sequences for the heavy chain and light chain of anti-VEGFv2 are provided in SEQ ID NO: 3. More particularly, the VH ORF is provided in
20 nt 2059 to 2427 and the CH1 is provided in 2428 to 2748 of SEQ ID NO: 3; the heavy chain without the leader has the nucleic acid sequence of nt 2059 to 2748 of SEQ ID NO: 3. The VL ORF is provided in nt 2896 to 3216 and CL is provided in nt 3217 to 3536 of SEQ ID NO: 3; the light chain without the leader sequence has the nucleic acid sequence of nt 2896 to 3536 of SEQ ID NO: 3.

In yet another embodiment, the coding sequences for the heavy chain and light chain of aVEGFv3 are provided in SEQ ID NO: 19. The VH ORF is provided in nt 1842 to 2210 and the CH1 is provided in nt 2211 to 2531 of SEQ ID NO: 19; the heavy chain without the leader has the nucleic acid sequence of nt 1842 to 2531 of SEQ ID NO: 19. The VL ORF is provided in nt 2679 to 2999 and CL is provided in nt 3000 to 3320 of SEQ ID NO: 19; the
30 light chain without the leader has the nucleic acid sequence of nt 2670 to 3320 of SEQ ID NO: 19.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv4 are provided in SEQ ID NO: 35. The heavy chain leader sequence is encoded by nt 1993 - 2052, the VH ORF is at nt 2053 - 2421 and the CH1 is at nt 2422 - 2742 of SEQ ID NO: 35. As in the other constructs described herein, as a result of the location of the F2A cleavage site, sequences encoding additional amino acids may be retained on the VH chain. The light chain leader sequence is encoded by nt 2830-2889; the VL ORF is provided in nt 2890 - 3210; the CL ORF is located at nt 3211-3531 of SEQ ID NO: 35.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv5 is provided within SEQ ID NO: 36. The heavy chain leader sequence is encoded by nt 1993 - 2052, the VH ORF is encoded by nt 2053 - 2421, and the CH1 is encoded by nt 2422 - 2742 of SEQ ID NO: 36. As in the other constructs described herein, as a result of the location of the F2A cleavage site, sequences encoding additional amino acids may be retained on the VH chain. The light chain leader sequence is encoded by nt 2830-2889; the VL ORF is provided in nt 2890 - 3210; the CL ORF is located at nt 3211-3531 of SEQ ID NO: 36.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv6 is provided within SEQ ID NO: 37. The heavy chain leader sequence is encoded by nt 1993 - 2051, the VH ORF is encoded by nt 2053 - 2421, and the CH1 is encoded by nt 2422 - 2742 of SEQ ID NO: 37. As in the other constructs described herein, as a result of the location of the F2A cleavage site, sequences encoding additional amino acids may be retained on the VH chain. The light chain leader sequence is encoded by nt 2830-2889; the VL ORF is provided in nt 2890 - 3210; the CL ORF is located at nt 3211-3531 of SEQ ID NO: 37.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv7 is provided within SEQ ID NO: 38. The heavy chain leader sequence is encoded by nt 1993 - 2052, the VH ORF is encoded by nt 2053 - 2421, and the CH1 is encoded by nt 2422 - 2742 of SEQ ID NO: 38. As in the other constructs described herein, as a result of the location of the F2A cleavage site, an additional Arg codon is retained on the VH chain. The light chain leader sequence is encoded by nt 2830 - 2889; the VL ORF is provided in nt 2890-3210; the CL ORF is located at nt 3211-3531 of SEQ ID NO: 38.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv8 is provided within SEQ ID NO: 39. The heavy chain leader sequence is encoded

by nt 1993 - 2052, the VH ORF is encoded by nt 205 - 2421, and the CH1 is encoded by nt 2422 - 2742 of SEQ ID NO: 39. As in the other constructs described herein, as a result of the location of the F2A cleavage site, an additional Arg codon is retained on the VH chain. The light chain leader sequence is encoded by nt 2830 - 2889; the VL ORF is provided in
5 2890 - 3210; the CL ORF is located at nt 3211 - 3531 of SEQ ID NO: 39.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv9 is provided within SEQ ID NO: 40. The heavy chain leader sequence is encoded by nt 1999 - 2058, the VH ORF is encoded by nt 2059 - 2427, and the CH1 is encoded by nt 2428 - 2748 of SEQ ID NO: 40. As in the other constructs described herein, as a result of
10 the location of the F2A cleavage site, an additional Arg codon is retained on the VH chain. The light chain leader sequence is encoded by nt 2836 - 2895; the VL ORF is provided in nt 2896 - 3216; the CL ORF is located at nt 3217 - 3637 of SEQ ID NO: 40.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv10 is provided within SEQ ID NO: 41. The heavy chain leader sequence is encoded
15 by nt 1993 - 2052, the VH ORF is encoded by nt 2053 - 2421, and the CH1 is encoded by nt 2422 - 2742 of SEQ ID NO: 41. As in the other constructs described herein, as a result of the location of the F2A cleavage site, an additional Arg codon is retained on the VH chain. The light chain leader sequence is encoded by nt 2830 - 2889; the VL ORF is provided in nt 2890 - 3210; the CL ORF is located at nt 3211 - 3231 of SEQ ID NO: 41.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv11 is provided within SEQ ID NO: 42. The heavy chain leader sequence is encoded
20 by nt 1993 - 2052, the VH ORF is encoded by nt 2053 - 2421, and the CH1 is encoded by nt 2422 - 2742 of SEQ ID NO: 42. As in the other constructs described herein, an F2A cleavage site is located between the end of the heavy chain and the beginning of the light
25 chain. The light chain leader sequence is encoded by nt 2830 - 2889; the VL ORF is provided in nt 2890 - 3210; the CL ORF is located at nt 3211 - 3531 of SEQ ID NO: 42.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv12 is provided within SEQ ID NO: 43. The heavy chain leader sequence is encoded
30 by nt 1993 - 2052, the VH ORF is encoded by nt 2053 - 2421, and the CH1 is encoded by nt 2422 - 2742 of SEQ ID NO: 43. The light chain leader sequence is encoded by nt 2830 - 2889; the VL ORF is provided in nt 2890 - 3210; the CL ORF is located at nt 3211 - 3531 of SEQ ID NO: 43.

In a further embodiment, the coding sequences for the heavy and light chain of aVEGFv13 is provided within SEQ ID NO: 44. The heavy chain leader sequence is encoded by nt 1993 - 2052, the VH ORF is encoded by nt 2053 - 2421, and the CH1 is encoded by nt 2422 - 2742 of SEQ ID NO: 44. The light chain leader sequence is encoded by nt 2830 - 2889; the VL ORF is nt 2890 - 3210; the CL ORF is located at nt 3211 - 3531 of SEQ ID NO: 44.

Ranibizumab is described herein as a positive control and is currently marketed under the brand name Lucentis®. It is described as a Fab moiety of a high affinity version of recombinant humanized monoclonal antibody rhuMAb vascular endothelial growth factor (VEGF). It consists of a 214-residue light chain linked by a disulfide bond at its C-terminus to the 231-residue N-terminal segment of the heavy chain. The expected amino acid sequences of the heavy and light chains are provided in SEQ ID NO: 1 and 2. CAS number 347396-82-1.

As used herein, an “immunoglobulin domain” refers to a domain of an antibody heavy chain or light chain as defined with reference to a conventional, full-length antibody. More particularly, a full-length antibody contains a heavy (H) chain polypeptide which contains four domains: one N-terminal variable (VH) region and three C-terminal constant (CH1, CH2 and CH3) regions and a light (L) chain polypeptide which contains two domains: one N-terminal variable (VL) region and one C-terminal constant (CL) region. An Fc region may contain two domains (CH2 - CH3). A Fab region contains one constant and one variable domain for each the heavy and light chains.

In one embodiment, rAAV.aVEGF vector has an AAV8 capsid and a vector genome packaged therein which comprises at least one element heterologous to AAV8. In one embodiment, the vector genome contains, from 5' to 3': (a) an AAV 5' ITR; (b) an enhancer; (c) a promoter; (d) an intron; (e) a leader sequence and the anti-VEGF heavy chain coding sequence; (f) a furin-F2a linker; (g) a leader sequence and the anti-VEFG light chain coding sequence; (h) a polyA signal; and (i) an AAV3' ITR.

In certain embodiments, the processing of anti-VEGF Fab heavy chain and light chains is directed by leader peptides that are derived from human IL2 protein. In one embodiment the leader sequence is an interleukin (IL) IL-2 leader sequence, which may be the wild-type human IL2, MYRMQLLSICIALSLALVTNS [SEQ ID NO: 29], or a mutated leader, such as MYRMQLLLLIALSLALVTNS [SEQ ID NO: 30] or

MRMQLLLLIALLSLALVTNS [SEQ ID NO: 31]. In another embodiment, a human serpinF1 secretion signal may be used as leader peptides. Other leader sequences can be used, or other leaders exogenous to the heavy and light chain.

As used in the following description of the vector genome unless otherwise specified as the light chain or heavy chain, reference to a coding sequence (e.g., aVEGFv2) encompasses the anti-VEGF heavy chain - furin/F2a linker - anti-VEGF light chain. In one embodiment, a nucleic acid sequence encoding the furin recognition site Arginine-Lysine-Arginine-Arginine is selected. In certain embodiments, nucleic acids encoding a F2A linker which is a 24 amino acid peptide derived from FMDV (GenBank # CAA2436.1) is selected. However, if desired, an IRES sequence, e.g., such as derived from encephalomyocarditis virus (EMCV) : SEQ ID NO: 32:

[TATGCTAGTACGTCTCTCAAGGATAAGTAAGTAATATTAAGGTACGGGAGGTAT
TGGACAGGCCGCAATAAAAATATCTTTATTTTCATTACATCTGTGTGTTGGTTTTTT
GTGTGAATCGATAGTACTAACATACGCTCTCCATCAAAACAAAACGAAACAAAA
15 CAAACTAGCAAAATAGGCTGTCCCCAGTGCAAGTGCAGGTGCCAGAACATTTCT
CTGGCCTAACTGGCCGGTACCTGAGCTCTAGTTTCACTTTCCCTAGTTTCACTTTC
CCTAGTTTCACTTTCCCTAGTTTCACTTTCCCTAGTTTCACTTTCCCTCGAGGAT
ATCAAGATCTGGCCTCGGCCGCCAG], cMyc [Nanbru C, et al (1997). J. Biol. Chem.
272, 32061-32066; Stoneley M, et al., (1998). Oncogene 16, 423-428.], or foot and mouth
20 disease (FMD) may be selected.

Inverted terminal repeats (ITR) from AAV2 may be selected. Vectors having ITRs from a different source than its capsid are termed "pseudotyped". In certain embodiments, ITRs from a source other than AAV2 may be selected for this construct to generate another pseudotyped AAV. Alternatively, ITRs from the same source as the capsid may be selected. In certain embodiments, ITRs may be selected to generate a self-complementary AAV, such as defined *infra*.

In certain embodiments, the promoter is CB7, a hybrid between a cytomegalovirus (CMV) immediate early enhancer (C4) and the chicken beta actin promoter. In other embodiments, the promoter is a ubiquitin C (UbC) promoter. See, e.g., WO 2001/091800. See, e.g., GenBank® accession numbers AF232305 (rat) and D63791 (human), respectively. Still other promoters and/or enhancers may be selected. See, e.g., cytomegalovirus (CMV) immediate early enhancer (260 bp, C4; GenBank # K03104.1). Chicken beta-actin promoter

(281 bp; CB; GenBank # X00182.1). In still other embodiments, multiple enhancers and/or promoters may be included.

In certain embodiments, an intron is included. One suitable intron is a chicken beta-actin intron. In one embodiment, the intron is 875 bp (GenBank # X00182.1). In another
5 embodiment, a chimeric intron available from Promega is used. However, other suitable introns may be selected.

The vector genomes described herein include a polyadenylation signal (polyA). A variety of suitable polyA are known. In one example, the polyA is rabbit beta globin, such as the 127 bp rabbit beta-globin polyadenylation signal (GenBank # V00882.1). In other
10 embodiments, an SV40 polyA signal is selected. Still other suitable polyA sequences may be selected.

Optionally, other suitable vector elements may be selected which may include, e.g., a UTR sequence or a Kozak sequence.

In one embodiment, the vector genome contains, ITR-CB7-CI-aVEGFv2-rBG-ITR,
15 [SEQ ID NO: 3]. In another embodiment, the vector genome contains: ITR-UbC-CI-aVEGFv2-SV40-ITR.[SEQ ID NO: 9]. In one embodiment, the vector genome contains, ITR-CB7-CI-aVEGFv3-rBG-ITR [SEQ ID NO: 14]. In another embodiment, the vector genome contains: ITR-UbC-PI-aVEGFv3-SV40-ITR [SEQ ID NO: 19]. In another embodiment, the vector genome contains: ITR-UbC-PI-aVEGFv1-SV40-ITR [SEQ ID NO:
20 24].In a further embodiment, the vector genome contains AAV2-ITR-CB7.CI.aVEGFv4.rBG-AAV2 ITR [SEQ ID NO: 35]. In a further embodiment, the vector genome contains AAV2-ITR-CB7.CI.aVEGFv5.rBG-AAV2 ITR [SEQ ID NO: 36]. In a further embodiment, the vector genome contains AAV2-ITR-CB7.CI.aVEGFv6.rBG-AAV2 ITR [SEQ ID NO: 37]. In a further embodiment, the vector genome contains AAV2-ITR-
25 CB7.CI.aVEGFv7.rBG-AAV2 ITR [SEQ ID NO: 38]. In a further embodiment, the vector genome contains AAV2-ITR-CB7.CI.aVEGFv8.rBG-AAV2 ITR [SEQ ID NO: 39]. In a further embodiment, the vector genome contains AAV2-ITR-CB7.CI.aVEGFv9.rBG-AAV2 ITR [SEQ ID NO: 40]. In a further embodiment, the vector genome contains AAV2-ITR-CB7.CI.aVEGFv10.rBG-AAV2 ITR [SEQ ID NO: 41]. In a further embodiment, the vector genome contains AAV2-ITR-CB7.CI.aVEGFv11.rBG-AAV2 ITR [SEQ ID NO: 42]. In a
30 further embodiment, the vector genome contains AAV2-ITR-CB7.CI.aVEGFv12.rBG-AAV2 ITR [SEQ ID NO: 43]. In a further embodiment, the vector genome contains AAV2

ITR-CB7.CI.aVEGFv13.rBG-AAV2 ITR [see, SEQ ID NO: 44]. In a further embodiment, the vector genome contains AAV2 ITR-CMV.PI.aVEGFv7.eCMVIres.aVEGF.SV40-AAV2 ITR [SEQ ID NO: 45]. In another embodiment, the vector genome contains AAV2 ITR.CMV.PI.aVEGF.FMDV1IRES.SV40 - ITR [SEQ ID NO: 46]. In still a further
5 embodiment, the vector genome contains AAV2 ITR.CMV.PI.aVEGF.cMycIRES.Fab.SV40 - ITR [SEQ ID NO: 47].

For use in producing an AAV viral vector (*e.g.*, a recombinant (r) AAV), the expression cassettes can be carried on any suitable vector, *e.g.*, a plasmid, which is delivered to a packaging host cell. The plasmids useful in this invention may be engineered such that
10 they are suitable for replication and packaging in prokaryotic cells, mammalian cells, or both. Suitable transfection techniques and packaging host cells are known and/or can be readily designed by one of skill in the art.

Methods for generating and isolating AAVs suitable for use as vectors are known in the art. *See generally, e.g.*, Grieger & Samulski, 2005, "Adeno-associated virus as a gene
15 therapy vector: Vector development, production and clinical applications," *Adv. Biochem. Engin/Biotechnol.* 99: 119-145; Buning *et al.*, 2008, "Recent developments in adeno-associated virus vector technology," *J. Gene Med.* 10:717-733; and the references cited below, each of which is incorporated herein by reference in its entirety. For packaging a
20 transgene into virions, the ITRs are the only AAV components required in *cis* in the same construct as the nucleic acid molecule containing the expression cassettes. The cap and rep genes can be supplied in *trans*.

In one embodiment, the expression cassettes described herein are engineered into a genetic element (*e.g.*, a shuttle plasmid) which transfers the immunoglobulin construct sequences carried thereon into a packaging host cell for production of a viral vector. In one
25 embodiment, the selected genetic element may be delivered to an AAV packaging cell by any suitable method, including transfection, electroporation, liposome delivery, membrane fusion techniques, high velocity DNA-coated pellets, viral infection and protoplast fusion. Stable AAV packaging cells can also be made. Alternatively, the expression cassettes may be used to generate a viral vector other than AAV, or for production of mixtures of
30 antibodies *in vitro*. The methods used to make such constructs are known to those with skill in nucleic acid manipulation and include genetic engineering, recombinant engineering, and

synthetic techniques. *See, e.g.*, Molecular Cloning: A Laboratory Manual, ed. Green and Sambrook, Cold Spring Harbor Press, Cold Spring Harbor, NY (2012).

As used herein, "AAV8 capsid" refers to the AAV8 capsid having the amino acid sequence of GenBank accession: YP_077180 (SEQ ID NO: 48) encoded by nucleic acid
5 sequence of NCBI Reference Sequence: NC_006261.1 (SEQ ID NO: 49), both of which are incorporated by reference herein. Some variation from this encoded sequence is encompassed by the present invention, which may include sequences having about 99% identity to the referenced amino acid sequence in GenBank accession: YP_077180; US Patent 7,282,199, 7,790,449; 8,319,480; 8,962,330; US 8,962,332, (*i.e.*, less than
10 about 1% variation from the referenced sequence). In another embodiment, the AAV8 capsid may have the VP1 sequence of the AAV8 variant described in WO2014/124282, which is incorporated by reference herein. Methods of generating the capsid, coding sequences therefore, and methods for production of rAAV viral vectors have been described. *See, e.g.*, Gao, et al, Proc. Natl. Acad. Sci. U.S.A. 100 (10), 6081-6086 (2003),
15 US 2013/0045186A1, and WO 2014/124282. In certain embodiments, an AAV8 variant which shows tropism for the desired target cell, e.g., photoreceptors, RPE or other ocular cells is selected. For example, an AAV8 capsid may have Y447F, Y733F and T494V mutations (also called "AAV8(C&G+T494V)" and "rep2-cap8(Y447F+733F+T494V)"), as described by Kay et al, Targeting Photoreceptors via Intravitreal Delivery Using Novel,
20 Capsid-Mutated AAV Vectors, PLoS One. 2013; 8(4): e62097. Published online 2013 Apr 26, which is incorporated herein by reference. *See, e.g.*, Mowat et al, Tyrosine capsid-mutant AAV vectors for gene delivery to the canine retina from a subretinal or intravitreal approach, Gene Therapy 21, 96-105 (January 2014), which is incorporated herein by
25 reference. In another embodiment, the AAV capsid is an AAV8 capsid, which preferentially targets bipolar cells. *See*, WO 2014/024282, which is incorporated herein by reference.

As used herein, the term "NAb titer" a measurement of how much neutralizing antibody (e.g., anti-AAV Nab) is produced which neutralizes the physiologic effect of its targeted epitope (e.g., an AAV). Anti-AAV NAb titers may be measured as described in,
30 e.g., Calcedo, R., et al., Worldwide Epidemiology of Neutralizing Antibodies to Adeno-Associated Viruses. Journal of Infectious Diseases, 2009. 199(3): p. 381-390, which is incorporated by reference herein.

The terms “percent (%) identity”, “sequence identity”, “percent sequence identity”, or “percent identical” in the context of amino acid sequences refers to the residues in the two sequences which are the same when aligned for correspondence. Percent identity may be readily determined for amino acid sequences over the full-length of a protein, polypeptide, about 32 amino acids, about 330 amino acids, or a peptide fragment thereof or the corresponding nucleic acid sequence coding sequencers. A suitable amino acid fragment may be at least about 8 amino acids in length, and may be up to about 700 amino acids. Generally, when referring to “identity”, “homology”, or “similarity” between two different sequences, “identity”, “homology” or “similarity” is determined in reference to “aligned” sequences. “Aligned” sequences or “alignments” refer to multiple nucleic acid sequences or protein (amino acids) sequences, often containing corrections for missing or additional bases or amino acids as compared to a reference sequence. Alignments are performed using any of a variety of publicly or commercially available Multiple Sequence Alignment Programs. Sequence alignment programs are available for amino acid sequences, *e.g.*, the “Clustal Omega”, “Clustal X”, “MAP”, “PIMA”, “MSA”, “BLOCKMAKER”, “MEME”, and “Match-Box” programs. Generally, any of these programs are used at default settings, although one of skill in the art can alter these settings as needed. Alternatively, one of skill in the art can utilize another algorithm or computer program which provides at least the level of identity or alignment as that provided by the referenced algorithms and programs. See, *e.g.*, J. D. Thomson et al, Nucl. Acids. Res., “A comprehensive comparison of multiple sequence alignments”, 27(13):2682-2690 (1999). As used herein, the term “operably linked” refers to both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest.

A “replication-defective virus” or “viral vector” refers to a synthetic or artificial viral particle in which an expression cassette containing a gene of interest is packaged in a viral capsid or envelope, where any viral genomic sequences also packaged within the viral capsid or envelope are replication-deficient; *i.e.*, they cannot generate progeny virions but retain the ability to infect target cells. In one embodiment, the genome of the viral vector does not include genes encoding the enzymes required to replicate (the genome can be engineered to be “gutless” - containing only the transgene of interest flanked by the signals required for amplification and packaging of the artificial genome), but these genes may be supplied during production. Therefore, it is deemed safe for use in gene therapy since replication and

infection by progeny virions cannot occur except in the presence of the viral enzyme required for replication.

The abbreviation “sc” refers to self-complementary. “Self-complementary AAV” refers a plasmid or vector having an expression cassette in which a coding region carried by a recombinant AAV nucleic acid sequence has been designed to form an intra-molecular double-stranded DNA template. Upon infection, rather than waiting for cell mediated synthesis of the second strand, the two complementary halves of scAAV will associate to form one double stranded DNA (dsDNA) unit that is ready for immediate replication and transcription. See, e.g., D M McCarty et al, “Self-complementary recombinant adeno-associated virus (scAAV) vectors promote efficient transduction independently of DNA synthesis”, *Gene Therapy*, (August 2001), Vol 8, Number 16, Pages 1248-1254. Self-complementary AAVs are described in, e.g., U.S. Patent Nos. 6,596,535; 7,125,717; and 7,456,683, each of which is incorporated herein by reference in its entirety.

The term “heterologous” when used with reference to a protein or a nucleic acid indicates that the protein or the nucleic acid comprises two or more sequences or subsequences which are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid. For example, in one embodiment, the nucleic acid has a promoter from one gene arranged to direct the expression of a coding sequence from a different gene. Thus, with reference to the coding sequence, the promoter is heterologous.

The term “exogenous” when used with reference to a protein or nucleic acid sequences indicates two or more sequences or subsequences which are from different sources, e.g., an AAV and a human protein.

It is to be noted that the term “a” or “an” refers to one or more. As such, the terms “a” (or “an”), “one or more,” and “at least one” are used interchangeably herein.

The words “comprise”, “comprises”, and “comprising” are to be interpreted inclusively rather than exclusively. The words “consist”, “consisting”, and its variants, are to be interpreted exclusively, rather than inclusively. While various embodiments in the specification are presented using “comprising” language, under other circumstances, a related embodiment is also intended to be interpreted and described using “consisting of” or “consisting essentially of” language.

As used herein, the term "about" means a variability of 10% from the reference given, unless otherwise specified.

Unless defined otherwise in this specification, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art and by reference to published texts, which provide one skilled in the art with a general guide
5 to many of the terms used in the present application.

rAAV8.aVEGF Formulation

The rAAV8.aVEGF formulation is a suspension containing an effective amount of
10 rAAV8.aVEGF vector suspended in an aqueous solution. In certain embodiments, the suspension contains buffered saline, optionally with a surfactant and/or other excipients. A buffered saline typically contains a physiologically compatible salt or mixture of salts, e.g. phosphate buffered saline, sodium chloride, or a mixture thereof.

In one embodiment, the formulation may contain, e.g., about 1×10^8 GC/eye to about
15 7×10^{12} GC/eye, or about 5×10^9 GC/eye to about 1×10^{11} GC/eye, or about 10^{10} GC/eye, or about as measured by oqPCR or digital droplet PCR (ddPCR) as described in, e.g., M. Lock et al, Hu Gene Therapy Methods, Hum Gene Ther Methods. 2014 Apr;25(2):115-25. doi: 10.1089/hgtb.2013.131. Epub 2014 Feb 14, which is incorporated herein by reference.

For example, a suspension as provided herein may contain both NaCl and KCl. The
20 pH may be in the range of 6.5 to 8, or 7.2 to 7.6. pH may be assessed using any suitable method, e.g., USP <791> [US Pharmacopeial Convention, reference standards]. A suitable surfactant, or combination of surfactants, may be selected from among a Poloxamers, i.e., nonionic triblock copolymers composed of a central hydrophobic chain of polyoxypropylene (poly(propylene oxide)) flanked by two hydrophilic chains of polyoxyethylene
25 (poly(ethylene oxide)), SOLUTOL HS 15 (Macrogol-15 Hydroxystearate), LABRASOL (Polyoxy caprylic glyceride), polyoxy 10 oleyl ether, TWEEN (polyoxyethylene sorbitan fatty acid esters), ethanol and polyethylene glycol. In one embodiment, the formulation contains a poloxamer. These copolymers are commonly named with the letter "P" (for poloxamer) followed by three digits: the first two digits x 100 give the approximate
30 molecular mass of the polyoxypropylene core, and the last digit x 10 gives the percentage polyoxyethylene content. In one embodiment Poloxamer 188 is selected. The surfactant may be present in an amount up to about 0.0005 % to about 0.001% of the suspension. In one

embodiment, the rAAV8.aVEGF formulation is a suspension containing at least 1×10^{11} genome copies (GC)/mL, or greater, e.g., about 1×10^{13} GC/mL as measured by oqPCR or digital droplet PCR (ddPCR) as described in, e.g., M. Lock et al, *Hum Gene Therapy Methods*, *Hum Gene Ther Methods*. 2014 Apr;25(2):115-25. doi: 10.1089/hgtb.2013.131. Epub 2014
5 Feb 14, which is incorporated herein by reference. In one embodiment, the vector is suspended in an aqueous solution containing 180 mM sodium chloride, 10 mM sodium phosphate, 0.001% Poloxamer 188, pH 7.3. The formulation is suitable for use in human subjects and is administered subretinally.

In order to ensure that empty capsids are removed from the dose of AAV8.aVEGF
10 that is administered to patients, empty capsids are separated from vector particles during the vector purification process. In one embodiment, the vector particles containing packaged genomes are purified from empty capsids using the process described in International Patent Application No. PCT/US16/65976, filed December 9, 2016 and its priority documents, US Patent Appln Nos. 62/322,098, filed April 13, 2016 and 62/266,341, filed on December 11,
15 2015, and entitled "Scalable Purification Method for AAV8", which is incorporated by reference herein. Briefly, a two-step purification scheme is described which selectively captures and isolates the genome-containing rAAV vector particles from the clarified, concentrated supernatant of a rAAV production cell culture. The process utilizes an affinity capture method performed at a high salt concentration followed by an anion exchange resin
20 method performed at high pH to provide rAAV vector particles which are substantially free of rAAV intermediates.

In one embodiment, the pH used is from 10 to 10.4 (about 10.2) and the rAAV particles are at least about 50% to about 90% purified from AAV8 intermediates, or a pH of 10.2 and about 90% to about 99% purified from AAV8 intermediates. In one embodiment,
25 this is determined by genome copies. A stock or preparation of rAAV8 particles (packaged genomes) is "substantially free" of AAV empty capsids (and other intermediates) when the rAAV8 particles in the stock are at least about 75% to about 100%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least 99% of the rAAV8 in the stock and "empty capsids" are less than about 1%, less than about 5%, less than about 10%,
30 less than about 15% of the rAAV8 in the stock or preparation. In one embodiment, the formulation is be characterized by an rAAV stock having a ratio of "empty" to "full" of 1 or less, preferably less than 0.75, more preferably, 0.5, preferably less than 0.3.

In a further embodiment, the average yield of rAAV particles is at least about 70%. This may be calculated by determining titer (genome copies) in the mixture loaded onto the column and the amount presence in the final elutions. Further, these may be determined based on q-PCR analysis and/or SDS-PAGE techniques such as those described herein or
5 those which have been described in the art.

For example, to calculate empty and full particle content, VP3 band volumes for a selected sample (*e.g.*, an iodixanol gradient-purified preparation where # of GC = # of particles) are plotted against GC particles loaded. The resulting linear equation ($y = mx+c$) is used to calculate the number of particles in the band volumes of the test article peaks. The
10 number of particles (pt) per 20 μ L loaded is then multiplied by 50 to give particles (pt) /mL. Pt/mL divided by GC/mL gives the ratio of particles to genome copies (pt/GC). Pt/mL–GC/mL gives empty pt/mL. Empty pt/mL divided by pt/mL and x 100 gives the percentage of empty particles.

Generally, methods for assaying for empty capsids and AAV vector particles with
15 packaged genomes have been known in the art. See, *e.g.*, Grimm et al., *Gene Therapy* (1999) 6:1322-1330; Sommer et al., *Molec. Ther.* (2003) 7:122-128. To test for denatured capsid, the methods include subjecting the treated AAV stock to SDS-polyacrylamide gel electrophoresis, consisting of any gel capable of separating the three capsid proteins, for example, a gradient gel containing 3-8% Tris-acetate in the buffer, then running the gel until
20 sample material is separated, and blotting the gel onto nylon or nitrocellulose membranes, preferably nylon. Anti-AAV capsid antibodies are then used as the primary antibodies that bind to denatured capsid proteins, preferably an anti-AAV capsid monoclonal antibody, most preferably the B1 anti-AAV-2 monoclonal antibody (Wobus et al., *J. Viral.* (2000) 74:9281-9293). A secondary antibody is then used, one that binds to the primary antibody and
25 contains a means for detecting binding with the primary antibody, more preferably an anti-IgG antibody containing a detection molecule covalently bound to it, most preferably a sheep anti-mouse IgG antibody covalently linked to horseradish peroxidase. A method for detecting binding is used to semi-quantitatively determine binding between the primary and secondary antibodies, preferably a detection method capable of detecting radioactive isotope
30 emissions, electromagnetic radiation, or colorimetric changes, most preferably a chemiluminescence detection kit. For example, for SDS-PAGE, samples from column fractions can be taken and heated in SDS-PAGE loading buffer containing reducing agent

(*e.g.*, DTT), and capsid proteins were resolved on pre-cast gradient polyacrylamide gels (*e.g.*, Novex). Silver staining may be performed using SilverXpress (Invitrogen, CA) according to the manufacturer's instructions. In one embodiment, the concentration of AAV vector genomes (vg) in column fractions can be measured by quantitative real time PCR (Q-PCR).

5 Samples are diluted and digested with DNase I (or another suitable nuclease) to remove exogenous DNA. After inactivation of the nuclease, the samples are further diluted and amplified using primers and a TaqMan™ fluorogenic probe specific for the DNA sequence between the primers. The number of cycles required to reach a defined level of fluorescence (threshold cycle, Ct) is measured for each sample on an Applied Biosystems Prism 7700
10 Sequence Detection System. Plasmid DNA containing identical sequences to that contained in the AAV vector is employed to generate a standard curve in the Q-PCR reaction. The cycle threshold (Ct) values obtained from the samples are used to determine vector genome titer by normalizing it to the Ct value of the plasmid standard curve. End-point assays based on the digital PCR can also be used.

15 In one aspect, an optimized q-PCR method is provided herein which utilizes a broad spectrum serine protease, *e.g.*, proteinase K (such as is commercially available from Qiagen). More particularly, the optimized qPCR genome titer assay is similar to a standard assay, except that after the DNase I digestion, samples are diluted with proteinase K buffer and treated with proteinase K followed by heat inactivation. Suitably samples are diluted with
20 proteinase K buffer in an amount equal to the sample size. The proteinase K buffer may be concentrated to 2 fold or higher. Typically, proteinase K treatment is about 0.2 mg/mL, but may be varied from 0.1 mg/mL to about 1 mg/mL. The treatment step is generally conducted at about 55 °C for about 15 minutes, but may be performed at a lower temperature (*e.g.*, about 37 °C to about 50 °C) over a longer time period (*e.g.*, about 20 minutes to about
25 30 minutes), or a higher temperature (*e.g.*, up to about 60 °C) for a shorter time period (*e.g.*, about 5 to 10 minutes). Similarly, heat inactivation is generally at about 95 °C for about 15 minutes, but the temperature may be lowered (*e.g.*, about 70 to about 90 °C) and the time extended (*e.g.*, about 20 minutes to about 30 minutes). Samples are then diluted (*e.g.*, 1000 fold) and subjected to TaqMan analysis as described in the standard assay.

30 Additionally, or alternatively, droplet digital PCR (ddPCR) may be used. For example, methods for determining single-stranded and self-complementary AAV vector genome titers by ddPCR have been described. See, *e.g.*, M. Lock et al, Hu Gene Therapy

Methods, Hum Gene Ther Methods. 2014 Apr;25(2):115-25. doi: 10.1089/hgtb.2013.131.
Epub 2014 Feb 14.

Manufacturing

The rAAV8.aVEGF vector can be manufactured as shown in the flow diagram
5 shown in FIG 9. Briefly, cells (e.g. HEK 293 cells) are propagated in a suitable cell culture
system and transfected for vector generation. The rAAV8.aVEGF vector can then be
harvested, concentrated and purified to prepare bulk vector which is then filled and finished
in a downstream process. Methods for manufacturing the gene therapy vectors described
herein include methods well known in the art such as generation of plasmid DNA used for
10 production of the gene therapy vectors, generation of the vectors, and purification of the
vectors. In some embodiments, the gene therapy vector is an AAV vector and the plasmids
generated are an AAV cis-plasmid encoding the AAV genome and the gene of interest, an
AAV trans-plasmid containing AAV rep and cap genes, and an adenovirus helper plasmid.
The vector generation process can include method steps such as initiation of cell culture,
15 passage of cells, seeding of cells, transfection of cells with the plasmid DNA, post-
transfection medium exchange to serum free medium, and the harvest of vector-containing
cells and culture media. The harvested vector-containing cells and culture media are referred
to herein as crude cell harvest.

The crude cell harvest may thereafter be subject method steps such as concentration
20 of the vector harvest, diafiltration of the vector harvest, microfluidization of the vector
harvest, nuclease digestion of the vector harvest, filtration of microfluidized intermediate,
purification by chromatography, purification by ultracentrifugation, buffer exchange by
tangential flow filtration, and formulation and filtration to prepare bulk vector.

In a specific embodiment, the methods used for manufacturing the gene therapy
25 vectors are described in the examples herein.

Patient Population

Patients who are candidates for treatment include those with neovascular age-related
macular degeneration, macular edema following retinal vein occlusion (RVO), diabetic
macular edema (DME), diabetic retinopathy (non-proliferative diabetic retinopathy (NPDR),
30 proliferative diabetic retinopathy (PDR) in patients with DME, diabetic retinopathy in
patients with diabetic macular edema. These patients are particularly well suited for
subretinal treatment with an AAV8.aVEGF composition as described herein.

Patients who are candidates for intraocular, including, e.g., subretinal and/or intravitreal administration, with an AAV8.aVEGF as described herein include those with macular degeneration, neovascular/wet/exudative age-related macular degeneration, macular edema following retinal vein occlusion (RVO) (including central retinal vein occlusion (CRVO) and branch retinal vein occlusion (BRVO)) central/hemi/branch retinal vein occlusion, retinal artery occlusion; retinal neovascularization; diabetic macular edema (DME), diabetic retinopathy (non-proliferative diabetic retinopathy (NPDR), proliferative diabetic retinopathy (PDR)) in patients with DME, diabetic retinopathy without macular edema (including pre-treatment of vitrectomy for proliferative diabetic retinopathy); active photocoagulated diabetic retinopathy; choroidal neovascularization, rare causes of choroidal neovascularization (angioid streaks, choroiditis [including choroiditis secondary to ocular histoplasmosis], idiopathic degenerative myopia, retinal dystrophies, rubeosis iridis, and trauma), idiopathic choroidal neovascularization, corneal neovascularization; retinopathy of prematurity, optic nerve head perfusion, retrolental fibroplasia; retinal degeneration; vitreomacular traction syndrome; retinal detachment, diabetic traction retinal detachment, submacular vascularized pigment epithelial detachments, Vogt Koyanagi Harada Disease, pigment epithelial detachment, pigment epithelium rip; vitreoretinopathy proliferative; vitreoretinal surgery in diabetic tractional retinal detachment, polypoidal choroidal vasculopathy; punctate inner choroidopathy (PIC); multifocal choroiditis; central serous chorioretinopathy (CSC), serpiginous choroiditis, vitreous hemorrhage, pars plana vitrectomy for vitreous hemorrhage, diabetic premacular hemorrhage with active fibrovascular proliferation; Choroidal hemorrhage amblyopia; myopia, myopic choroidal neovascularization, choroidal subfoveal/juxtafoveal neovascularization in high myopia; choroidal melanoma; ocular histoplasmosis syndrome, tecalcitrant inflammatory ocular neovascularization (neovascularization, tuberculosis, multifocal serpiginous choroiditis, harada toxoplasmosis); Pseudoxanthoma elasticum; hereditary eye diseases; corneal endothelial cell loss; Vogt Koyanagi Harada Disease, non-arteritic anterior ischemic optic neuropathy; cystoid macular edema; refractory cystoid macular oedema; idiopathic macular telangiectasia; Coat's disease (Coates' disease, also known as exudative retinitis or retinal telangiectasis); glaucoma, neovascular glaucoma, steroid-induced glaucoma, ocular hypertension, Glaucoma surgery; control of wound healing ; uveal melanoma; uveitis; radiation maculopathy, pattern dystrophy, radiation retinopathy, radiation necrosis; Hippel-

Lindau Disease; Von Hippel-Lindau Syndrome; endophthalmitis; neuromyelitis optica spectrum disorder; pterygium, primary pterygium (including as adjunctive therapy for primary pterygium surgery), recurrent pterygium; retinal drusen; eye neoplasms; intraocular melanoma; cataract; corneal graft failure; trabeculectomy; lipid keratopathy, penetrating keratoplasty, herpetic keratopathy, rosacea; retinal angioma; retinovascular disease; vision disorders, vitreoretinopathy proliferative; iris neovascularization (NV); corneal NV, including pannus, pars planitis sarcoid or Eale's disease.

Patients who are candidates for treatment with an AAV8.aVEGF (the anti-VEGF transgene product) in a regimen which involves a combination with, but not limited to 24GyE proton, 16GyE, Xylocaine, Proparacaine Hydrochloride, Tetravisc, Acuvail, Zimura, Triamcinolone acetone, Ranibizumab, or Ozurdex. Examples of suitable indications include those in the preceding paragraph. For example, a combination regimen involving an AAV8.aVEGF with one or more of the drugs listed above, may be used for treatment of exudative age-related macular degeneration, central retinal vein occlusion, idiopathic polypoidal choroidal vasculopathy, and/or diabetic macular edema.

The AAV8.aVEGF composition described herein are also useful in preventing vascularization in a number of cancers, neoplasms and other diseases associated with VEGF. Such compositions may be administered for any suitable route, including, e.g., intravenous, intralesional, direct delivery to a tumor or organ, among others. Such patients may include those with Acute Lymphoblastic Leukemia (ALL), Acute Myeloid Leukemia (AML), Adrenocortical Carcinoma, Adrenocortical Carcinoma, AIDS-Related Cancers, Kaposi Sarcoma, AIDS-Related Lymphoma, Primary CNS Lymphoma, Anal Cancer, Appendix Cancer, Astrocytomas, Atypical Teratoid/Rhabdoid Tumor, Basal Cell Carcinoma of the Skin, Bladder Cancer, Bone Cancer (includes Ewing Sarcoma and Osteosarcoma and Malignant Fibrous Histiocytoma), Brain Tumors, Breast Cancer, Bronchial Tumors, Burkitt Lymphoma, Non-Hodgkin Lymphoma, Carcinoid Tumors, Carcinoma of Unknown Primary, Cardiac Tumors, Central Nervous System Atypical Teratoid/Rhabdoid Tumor, Embryonal Tumors, Germ Cell Tumor, Primary CNS Lymphoma, Cervical Cancer, Unusual Cancers of Childhood, Cholangiocarcinoma, Bile Duct Cancer, Chordoma, Chronic Lymphocytic Leukemia (CLL), Chronic Myelogenous Leukemia (CML), Chronic Myeloproliferative Neoplasms, Colorectal Cancer, Craniopharyngioma, Cutaneous T-Cell Lymphoma, Ductal Carcinoma In Situ (DCIS), Central Nervous System Embryonal Tumors, Endometrial

Cancer, Ependymoma, Esophageal Cancer, Esthesioneuroblastoma, Ewing Sarcoma ,
 Extracranial Germ Cell Tumor, Extragonadal Germ Cell Tumor, Eye Cancer, Intraocular
 Melanoma, Retinoblastoma, Fallopian Tube Cancer, Fibrous Histiocytoma of Bone,
 Malignant, Osteosarcoma, Gallbladder Cancer, Gastric Cancer, Childhood Gastric Cancer,
 5 Gastrointestinal Stromal Tumors (GIST), Germ Cell Tumors, Childhood Central Nervous
 System Germ Cell Tumors, Childhood Extracranial Germ Cell Tumors, Extragonadal Germ
 Cell Tumors, Ovarian Germ Cell Tumors, Testicular Cancer, Gestational Trophoblastic
 Disease, Hairy Cell Leukemia, Head and Neck Cancer, Heart Tumors, Hepatocellular
 Cancer, Histiocytosis, Langerhans Cell, Hodgkin Lymphoma, Hypopharyngeal Cancer,
 10 Intraocular Melanoma, Islet Cell Tumors, Pancreatic Neuroendocrine Tumors, Kaposi
 Sarcoma, Kidney Cancer, Langerhans Cell Histiocytosis, Laryngeal Cancer, Childhood
 Laryngeal Cancer and Papillomatosis, Leukemia, Lip and Oral Cavity Cancer, Liver Cancer,
 Lung Cancer (Non-Small Cell and Small Cell), Childhood Lung Cancer, Lymphoma, Male
 Breast Cancer, Malignant Fibrous Histiocytoma of Bone and Osteosarcoma, Melanoma,
 15 Intraocular Melanoma, Merkel Cell Carcinoma, Malignant Mesothelioma, Metastatic
 Cancer, Metastatic Squamous Neck Cancer with Occult Primary, Midline Tract Carcinoma
 Involving NUT Gene, Mouth Cancer, Multiple Endocrine Neoplasia Syndromes, Multiple
 Myeloma/Plasma Cell Neoplasms, Mycosis Fungoides, Myelodysplastic Syndromes,
 Myelodysplastic, Myeloproliferative Neoplasms, Myelogenous Leukemia, Chronic (CML),
 20 Acute Myeloid Leukemia (AML), Chronic Myeloproliferative Neoplasms, Nasal Cavity and
 Paranasal Sinus Cancer; Nasopharyngeal Cancer, Neuroblastoma, Non-Hodgkin Lymphoma,
 Oral Cancer, Lip and Oral Cavity Cancer and Oropharyngeal Cancer, Osteosarcoma and
 Malignant Fibrous Histiocytoma of Bone, Ovarian Cancer, Pancreatic Cancer, Pancreatic
 Neuroendocrine Tumors, Papillomatosis, Paraganglioma, Paranasal Sinus and Nasal Cavity
 25 Cancer, Parathyroid Cancer, Penile Cancer, Pharyngeal Cancer, Pheochromocytoma, Plasma
 Cell Neoplasm/Multiple Myeloma, Pleuropulmonary Blastoma; Pregnancy and Breast
 Cancer, Primary Central Nervous System (CNS) Lymphoma, Primary Peritoneal Cancer,
 Prostate Cancer, Rectal Cancer, Recurrent Cancer, Renal Cell Cancer, Retinoblastoma,
 Salivary Gland Cancer, Sarcoma, Childhood Rhabdomyosarcoma, Childhood Vascular
 30 Tumors, Ewing Sarcoma, Osteosarcoma, Uterine Sarcoma, Sézary Syndrome, Skin Cancer,
 Small Cell Lung Cancer, Small Intestine Cancer, Soft Tissue Sarcoma, Squamous Cell
 Carcinoma of the Skin, Squamous Neck Cancer with Occult Primary, Metastatic, Stomach

Cancer, Cutaneous T-Cell Lymphoma, Testicular Cancer, Throat Cancer, Nasopharyngeal Cancer, Oropharyngeal Cancer, Hypopharyngeal Cancer, Thymoma and Thymic Carcinoma, Thyroid Cancer, Transitional Cell Cancer of the Renal Pelvis and Ureter (Kidney (Renal Cell) Cancer), Carcinoma of Unknown Primary, Childhood Cancer of Unknown Primary,

5 Unusual Cancers of Childhood, Ureter and Renal Pelvis, Urethral Cancer; Endometrial Uterine Cancer, Uterine Sarcoma, Uterine Leiomyosarcomas, Vaginal Cancer, Vascular Tumors, Vulvar Cancer, Wilms Tumor and Other Childhood Kidney Tumors, abdominal neoplasms (adenocarcinoma; hepatocellular, papillary serous mullerian, Lapatinib, colorectal, ovarian, fallopian tube, peritoneal ncer/neoplasms/carcinoma/tumors);

10 lymphoproliferative disorder; small intestine cancer; acoustic neuroma(e.g. vestibular Schwannoma, Neurofibromatosis Type 2); acute myeloid leukemia, acute respiratory distress syndrome (ARDS), head and neck cancer; squamous cell carcinoma, multiple myeloma, non-hodgkin's lymphoma; B-cell lymphoma, sarcoma, neuroblastoma, advanced cancer, alignant neoplasms of female genital organs; metastatic or unresectable solid tumor,

15 anaplastic astrocytoma, Colon Cancer, Metastatic Melanoma, Malignant Ascites, Renal Cell Carcinoma, Glioblastoma, Gliosarcoma, Colorectal Liver Metastases, Advanced Malignancy, Myeloma, Gestational Trophoblastic Neoplasia, Choriocarcinoma, Placental Site Trophoblastic Tumor, Epithelioid Trophoblastic Tumor, Biliary Tract Cancer, Malignant Glioma, Cervical Cancer, Uterine Cancer, Mesothelioma. Candidates thereof are

20 treated with said composition alone or in combination with anti-cancer treatments, for example but not limited to paclitaxel, carboplatin, oxaliplatin, radiation, capecitabine, irinotecan, fluorouracil, doxorubin hydrochloride liposome, erlotinib hydrochloride, irinotecan hydrochloride, Irinotecan hydrochloride hydrate (CPT-11), gemcitabine hydrochloride, Pazopanib Hydrochloride, topotecan hydrochloride, Trifluridine/tipiracil

25 hydrochloride, Pegylated Liposomal Doxorubicin Hydrochloride, enzastaurin hydrochloride, mitoxantrone hydrochloride; epirubicin hydrochloride, docetaxel, gemcitabine, erlotinib, cisplatin, chemotherapy, cetuximab, FOLFIRI-Cetuximab, 5-Fluorouracil (5-FU), LV5FU2, cyclophosphamide, temozolomide, pemetrexed, levofolinate calcium (l-LV), Leucovorin Calcium, FOLFOX, FOLFOX6, mFOLFOX, FOLFOXIRI, FOLFIRI, doxorubicin,

30 Liposomal Doxorubicin, doxorubicin HCL liposome, Sorafenib Tosylate, sorafenib, triamcinolone, Triamcinolone acetonide, trastuzumab, everolimus, sunitinib, dexamethasone, conventional surgery, xeloda, radiotherapy, temsirolimus, pazopanib, Leucovorin (LV), l-

LV, anitumumab, epirubicin, verteporfin, AMG 655, Amgen 386, AMG 479, AMG 706,
 AMG 951, AMG 102, Folinic Acid, levo-folinic acid, etoposide, BAY 43-9006,
 atezolizumab, Interferon Alfa-2b, Interferon alpha-2a, interferon alfa, Gamma-Interferon-1b,
 Photodynamic Therapy, vinorelbine tartrate, vinorelbine, topotecan, tarceva, pemetrexed
 5 disodium, estramustine phosphate sodium, Imetelstat sodium, XELOX, RAD001,
 pegfilgrastim, paclitaxel albumin-stabilized nanoparticle formulation, ipilimumab,
 Stereotactic Radiosurgery (SRS), Stereotactic Radiation, ozurdex, letrozole, AG-013736
 (axitinib), filgrastim, crizotinib, cediranib maleate, cediranib, bortezomib, abraxane,
 vorinostat, vincristine, TRC105, rituximab, regorafenib, pembrolizumab, methotrexate,
 10 imatinib, Herceptin, tecentriq, oxaliplatin (OXA), lomustine, ixabepilone, CPT-11, CGC-
 11047, vinorelbine tartrat, tartrate, prednisone, nivolumab, fulvestrant, enzastaurin, doxil,
 AZD2014, AZD2281, AZD2171, AZD4547, AZD5363, AZD8931, Vitamin B12, Vitamin
 C, Vitamin D, Valproic acid, mitomycin C, Cediranib Maleate, lenalidomide, lapatinib, HAI
 Abraxane, HAI Irinotecan, GDC-0941, GDC-0449, GDC-0980, bicalutamide, xeliri,
 15 vandetanib, thalidomide, rapamycin, olaparib, NovoTTF100A, Navelbine, metmab, Imatinib
 Mesylate (Gleevec), ifosfamide, hydroxychloroquine, and GM-CSF.

Still other suitable conditions for treatment may include, e.g., Hemophilia, Synovitis,
 Hypertension, keloid, inflammation, Radiation Necrosis, and Neoplastic Meningitis. These
 and the conditions described above may be delivered by any suitable route, except where
 20 subretinal or another type of administration to the eye is specified.

In certain embodiments, patients receive a single dose of rAAV8.aVEGF
 administered subretinally. For example, this is particularly well suited for treatment of
 neovascular age-related macular degeneration, macular edema following retinal vein
 occlusion (RVO), diabetic macular edema (DME), diabetic retinopathy (non-proliferative
 25 diabetic retinopathy (NPDR), proliferative diabetic retinopathy (PDR) in patients with DME,
 diabetic retinopathy in patients with diabetic macular edema.

The dose of rAAV8.aVEGF administered to a patient is at least 1×10^9 GC/eye to 1×10^{13} GC/eye, or at least 1×10^{10} GC/eye to about 7.5×10^{12} GC/eye (as measured by
 oqPCR or ddPCR). However, other doses may be selected. For example, therapeutically
 30 effective subretinal doses of the rAAV8.aVEGF for patients may range from about 6.6×10^9
 GC/eye to about 6.6×10^{11} GC/eye, most preferably, 6.6×10^{10} GC/eye, in an injection
 volume ranging from about 0.1 mL to about 0.5 mL, preferably in 0.1 to 0.15 mL (100 – 150

μl). In still other embodiments, therapeutically effective concentrations may be about 1×10^5 concentration can be 1×10^5 GC/μL to 1×10^9 GC/μL, and the volume of injection for any GC concentration in that range can be from 10 μL to 300 μL.

5 In certain embodiments, patients may receive an rAAV8.aVEGF by subretinal administration by a retinal surgeon under local anesthesia. The procedure may involve standard 3 port pars plana vitrectomy with a core vitrectomy followed by subretinal delivery into the subretinal space by a subretinal cannula (36 to 41 gauge). In certain embodiments, 100 to 150 microliters of rAAV8.aVEGF will be delivered.

10 In some embodiments, rAAV8.aVEGF is administered in combination with one or more therapies for the treatment of wetAMD or another selected disorder. In some embodiments, rAAV.aVEGF is administered in combination with laser coagulation, photodynamic therapy with verteporfin, and intravitreal with anti-VEGF agent, including but not limited to pegaptanib, ranibizumab, aflibercept, or bevacizumab.

15 In certain embodiments, patients for rAAV8.aVEGF therapy may include those which have previously responded to conventional anti-VEGF antibody (Fab) treatment.

The goal of the gene therapy treatment of the invention is to slow or arrest the progression of retinal degeneration, and to slow or prevent loss of vision with minimal intervention/invasive procedures. In certain embodiments, the efficacy of the gene therapy treatment may be indicated by the elimination of or reduction in the number of rescue 20 treatments using standard of care, for example, intravitreal injections with anti-VEGF agents, including but not limited to pegaptanib, ranibizumab, aflibercept, or bevacizumab.

In certain embodiments, efficacy by measured by one or more of the following:
Vision change, visual acuity, including best corrected visual acuity measured by (BCVA) score, Snellen chard or Early Treatment Diabetic Retinopathy (ETDRS) visual acuity score, 25 percentage of subjects losing or gaining measured by ETDRS, distance best corrected visual acuity, reading best corrected visual acuity, change in NEI Visual Functioning Questionnaire-25 (VFQ-25) score , questionnaire of vision-related quality of life, contrast sensitivity measured by Pelli-Robson charts; low-contrast visual acuity on Electronic Visual Acuity Tester; peripheral visual field as measured by Goldmann visual field, mean angle 30 opening distance and trabecular-iris spur area measured by Heidelberg Slit-Lamp Optical Coherence Tomography, Preferential-Hyperacuity-Perimeter (PHP) testing of Age Related Macular Degeneration (AMD) by characterizing central and paracentral metamorphopsia,

retinal sensitivity (mfERG, Nidek MP-1 microperimetry), Visual Analog Scale (VAS), Macular Mapping Test, electrophysiological changes, including electroretinogram (ERG), pattern electroretinography (PERG) and full field (or flash) electroretinography (ffERG), multifocal electroretinography (mfERG), mfERG central ring amplitude density; mean

5 retinal sensitivity (dB) in three concentric rings (4°, 8° & 12°), visual evoked potential (VEP): ECG parameters included PR interval, QRS interval, and corrected QT interval using Fridericia's formula (QTcF). Anatomical changes, including regression of NVE (retinal neovascularization), CNVM (Choroidal Neovascular Membranes), changes measured using optical coherence topography (OCT), including macular volume, macular thickness, central

10 macular subfield thickness, retinal volume (inner retinal volume and outer retinal volume), retinal thickness, central retinal thickness, central subfield retinal thickness (CSRT), subfoveal retinal thickness (SRT), foveal thickness, maximum diameter of foveal avascular zone, integrity of retinal layers, external limiting membrane (ELM) integrity, ellipsoidal line/band integrity, lens status, lens opacity, neovascular membrane regression percentage

15 measured by Optical coherence tomography angiography (OCTA), degree of integrity of the photoreceptors in the inner/outer segments layer in the 1 mm centered in the fovea. Optionally, during trial, AMD lesion size and leakage may be by fluorescein angiography, change in total lesion size and CNV (choroidal neovascularization) size by fluorescein

20 angiography (FA) and Indocyanine green angiography (ICG), active CNV leakage which may include subretinal fluid or hemorrhage, area of leakage, area of macular leakage, change in percentage of lesion hemorrhage, change in drusen size, amount of fluid, intra-retinal cystoid changes (IRCs) volume, vessel density, presence of intra/sub-retinal fluid, sub-retinal fluid (SRF) height and diameter, intraretinal fluid volume, anterior chamber reaction, chorioretinal perfusion (ICG), development of geographic atrophy (GA) as detected by

25 fundus photography (FP) and/or fundus autofluorescence (AF), presence and extension of capillary occlusion, peripheral retinal ischemia, macular sensitivity using microperimetry, neovascularization of the iris, neovascularization of the angle, diabetic retinopathy.

In certain embodiments, subretinal and/or intra-retinal injection of the AAV8.aVEGF results in plasma and serum levels free of the aVEGF.

30 In certain embodiments, efficacy may be monitored by measuring BCVA (Best-Corrected Visual Acuity), intraocular pressure, slit lamp biomicroscopy, indirect ophthalmoscopy, SD-OCT (SD-Optical Coherence Tomography). Signs of vision loss,

infection, inflammation and other safety events, including retinal detachment may also be monitored.

SD-OCT is a useful non-invasive, in vivo cross-sectional retinal microscopy technique. Suitable equipment is commercially available. *See, e.g.*, Spectralis OCT, Heidelberg Engineering, Carlsbad, CA. In brief, this technique may be performed by dilating pupils. En face retinal imaging can be performed with near infrared (NIR) reflectance (REF) and/or with NIR fundus autofluorescence (FAF) using the scanning laser ophthalmoscope of this imaging system. Spectral domain optical coherence tomography scanning can be performed with 9 mm long horizontal and vertical cross-sections through the fovea and overlapping 30×25 mm raster scans extending into the near midperiphery. The parameters may be modified as needed, or other suitable parameters determined comparable.

Retinal function can be evaluated by a full-field electroretinogram (ERG). An ERG is a mass electrical potential generated by the retina in response to light stimulus. Usually, it is recorded by an electrode in contact with the corneal surface. Electroretinograms can be conducted in accordance with the recommendations set by the International Society for Clinical Electrophysiology of Vision (ISCEV; McCulloch, *Doc Ophthalmol.* 2015 Feb;130(1):1-12. 2015). In summary, an electroretinogram (ERG) is usually generated when all retinal cells actively respond to a flash stimulation (a dark-adapted animal, moderate to intense flash). The 2 components are the following: • a-wave: cornea-negative signal, first after the flash. Origin: photoreceptor photocurrent, the most direct signature of photoreceptor function. • b-wave: cornea-positive signal following the a-wave generated mostly by on-bipolar cells (second order neurons downstream from photoreceptors). In the examples described below, the following International Society for Clinical Electrophysiology of Vision (ISCEV) standard and additional protocols were used. However, these parameters may be adjusted as needed or required. Dark-adapted rod ERG: Stimulus intensity: 0.01 to 0.02 cd s m^{-2} . Response: b-wave only, no a-wave. Source: rod “on” bipolar cells (second order neurons driven by input from rods). Meaning: a measure of rod function. Dark-adapted standard flash ERG: Stimulus intensity: 3 cd s m^{-2} . Response: combined rod-cone a- and b-waves; 60% to 70% of the signal being generated by the rod-driven pathway. Source: photoreceptors, both rods and cones (a-wave); higher order neurons driven by both rods and cones. Meaning: a measure of mostly rod function; less sensitive to the state of dark adaptation and less variable than the “dim flash” response. Dark-adapted bright flash ERG:

Stimulus intensity: 10 cd s m^{-2} . Response and meaning: same as for the “standard flash” response, but bright flash response is larger in magnitude and may be less variable. Light-adapted standard flash cone ERG: Stimulus intensity: 3 cd s m^{-2} , delivered in presence of 30 cd m^{-2} background light after 5 minutes of light adaptation. Response: a- and b-waves generated by cone-driven pathways. Meaning: in presence of background light which completely desensitizes rods the ERG is produced exclusively by cones and cone-driven secondary retinal neurons and is a measure of the cone function. Light-adapted bright flash cone ERG (in addition to the ISCEV standard): Stimulus intensity: 10 cd s m^{-2} , delivered in presence of 30 cd m^{-2} background light after 5 minutes of light adaptation. Response and meaning: cone-driven ERG as in case of the “Standard cone ERG”, but of greater magnitude and potentially less variable. ERG measures (a-wave amplitude, a-wave implicit time, b-wave amplitude, b-wave implicit time) were summarized using mean and standard deviation (SD) for treated eyes and control eye.

Another measure of efficacy may include a lack of thickening of the retina.

As illustrated in the examples below, administration of 1×10^{10} GC/eye of an AAV8.aVEFG vector causes no impairment to retinal function. This dose is not a limitation on the therapeutically effective amounts which can be administered.

Measuring Clinical Objectives

Safety of the gene therapy vector after administration can be assessed by the number of adverse events, changes noted on physical examination, and/or clinical laboratory parameters assessed at multiple time points up to about 36 months post vector administration. Although physiological effect may be observed earlier, e.g., in about 1 day to one week, in one embodiment, steady state levels expression levels are reached by about 12 weeks.

Improvement/efficacy resulting from rAAV.aVEGF administration can be assessed as a defined mean change in baseline in visual acuity at about 12 weeks, 12 months, 24 months, 36 months, or at other desired time points. Other improvements/efficacy can be assessed as mean change from baseline in central retinal thickness as measured by spectral domain optical coherence tomography (SD-OCT) at 12, 24 and 36 months.

In some embodiments, treatment with rAAV.aVEGF results in a 5%, 10%, 15%, 20%, 30%, 40%, 50% or more increase in visual acuity from baseline. In some embodiments, treatment with rAAV.aVEGF results in a decrease, e.g., about 5%, about 10%, about 15%, about 20%,

about 30%, about 40%, about 50% or more decrease in central retinal thickness . In other embodiments, the central retinal thickness is stable, i.e., no increase in central retinal thickness. In certain embodiments, a measure of efficacy includes stabilizing retinal thickness, and/or stabilizing/decreasing) exudate and/or drusen.

5 In one embodiment, expression may be observed as early as about 8 hours to about 24 hours post-dosing. One or more of the desired clinical effects described above may be observed within several days to several weeks post-dosing.

 The invention is illustrated by the examples below which demonstrate that subretinal administration of an rAAV8.aVEGF vector results in gene transfer throughout the retina, and
10 expression of anti-VEGF Fab throughout the retina and in the vitreous and anterior chamber fluids. This result is surprising in view of prior art gene therapy studies that demonstrated that gene transfer spreads laterally outside of the original injection bleb but remains confined to those expanded boundaries and did not achieve gene transfer and transgene expression outside this expanded area of injection (the “bleb” formed in the retina at the injection site);
15 and offers an advantage over standard of care treatment for nAMD in that a single administration of the rAAV8.aVEGF vector should result in (i) continuous delivery of the effective amounts of the VEGF inhibitor throughout the retina which may in turn improve performance as compared to repeated IVT administrations of high dose boluses of the VEGF inhibitor that dissipate over time; and (ii) avoidance of repeated ocular injections which pose
20 additional risks and inconvenience to patients. Each aspect may improve therapeutic outcome.

EXAMPLES:

 The following abbreviations are used in the specification: AAV refers to Adeno-Associated Virus. ACF refers to Anterior Chamber Fluid. Ad5 refers to Adenovirus type 5.
25 AE refers to Adverse Event. AMD refers to Age-Related Macular Degeneration. BCA refers to Bicinchoninic Acid. BCVA refers to Best-Corrected Visual Acuity. BH refers to Bulk Harvest. BI refers to Bulk Drug Substance Intermediate. BP refers to Base pairs. CB refers to Chicken Beta Actin Promoter. CB7 refers to a hybrid CMV Enhancer (C4) and Chicken β -Actin Promoter. CBC refers to Complete Blood Count. CI refers to chicken β -Actin Intron.
30 CMC refers to Chemistry, Manufacturing and Control. CMO refers to Contract Manufacturing Organization. CMV refers to Cytomegalovirus. CNV refers to Choroidal Neovascularization. CS-10 refers to Corning 10-layer CellSTACKs® plates. ddPCR refers

to Droplet Digital Polymerase Chain Reaction. DLS refers to Dynamic Light Scattering. DMEM refers to Dulbecco's Modified Eagle Medium. DNA refers to deoxyribonucleic Acid. DP refers to Drug Product. ELISA refers to Enzyme-Linked Immunosorbent Assay. ERG refers to electroretinogram. ELISPOT refers to Enzyme Linked Immunospot. Fab
5 refers to Antigen-Binding Fragment. FBS refers to Fetal Bovine Serum. GC refers to Genome Copies. g refers to gram. GLP refers to Good Laboratory Practices. GMP refers to Good Manufacturing Practices. HEK293 refers to Human Embryonic Kidney Cells. HCP refers to Host Cell Protein. HS-36 refers to Corning 36-layer HYPERStacks®. ICH refers to International Conference on Harmonization. IND refers to Investigational New Drug. IP
10 refers to In-Process. ITR refers to Inverted Terminal Repeat. IU refers to Infectious Unit. IV refers to Intravenous. IVT refers to Intravitreal. KDa refers to KiloDalton. Kg refers to Kilogram. LOQ refers to Limit of Quantification. Lucentis® is a brand name Ranibizumab. MCB refers to Master Cell Bank. MED refers to Minimally Effective Dose. µl refers to microliter. mL refers to milliliter. Mm refers to millimeter. mRNA refers to Messenger
15 RNA. MS refers to Mass Spectrometry. Ng refers to Nanogram. NHP refers to Non-Human Primate. OCT refers to Optical Coherence Tomography. oqPCR refers to Optimized Quantitative Polymerase Chain Reaction. PCR refers to Polymerase Chain Reaction. PD refers to Pharmacodynamics. popPK refers to Population Pharmacokinetics. PEI refers to Polyethylenimine. PK refers to Pharmacokinetics. POC refers to Proof-Of-Concept. PRN
20 refers to pro re nata (as needed). QA refers to Quality Assurance. qPCR refers to Quantitative Polymerase Chain Reaction. rAAV refers to Recombinant Adeno-Associated Virus. RBG refers to Rabbit Beta-Globin. RPE refers to Retinal Pigment Epithelium. S-36 refers to HYPERstack®36-layer. SEND refers to Standards for Exchange of Nonclinical Data. SOC refers to Standard of Care. SOP refers to Standard Operating Procedure. TCID50
25 refers to Tissue Culture Infectious Dose 50%. TFF refers to Tangential Flow Filtration. µL refers to Microliter. VA refers to Visual Acuity. VEGF refers to Vascular Endothelial Growth Factor. WAMD refers to Wet Age-Related Macular Degeneration. YAG refers to Yttrium-aluminum-garnet.

30 EXAMPLE 1: Treating Human Subjects

This Example relates to a gene therapy treatment for patients with neovascular (wet) age-related macular degeneration (nAMD). In this example, the gene therapy vector,

rAAV8.aVEGF, a replication deficient adeno-associated viral vector 8 (AAV8) carrying a coding sequence for a soluble anti-VEGF Fab protein is administered to patients with nAMD. The goal of the gene therapy treatment is to slow or arrest the progression of retinal degeneration and to slow or prevent loss of vision with minimal intervention/invasive procedures.

A. Gene Therapy Vector

The generation of several rAAV8.aVEGF gene therapy vectors is described in Example 2 herein. Moreover, a schematic representation of the rAAV8.aVEGF vector genome is shown in FIG 1. rAAV8.aVEGF is a non-replicating recombinant AAV8 viral vector containing a transgene that leads to the production of a human anti-vascular endothelial growth factor (anti-VEGF) antigen binding antibody fragment (Fab). The gene cassette is flanked by the AAV2 inverted terminal repeats (ITRs). Expression from the cassette is driven by a CB7 promoter, a hybrid of a cytomegalovirus immediate-early enhancer and the chicken β -actin promoter. Transcription from this promoter is enhanced by the presence of the chicken β -actin intron. The polyadenylation signal for the expression cassette is from the rabbit β -globin gene. The nucleic acid sequences coding for the heavy and light chains of anti-VEGF Fab are separated by a self-cleaving furin (F)/F2A linker. The incorporation of the furin-F2A linker ensures expression of about equal amounts of the heavy and the light chain polypeptides.

The final product is supplied as a frozen solution of the AAV vector active ingredient in a formulation buffer in Crystal Zenith[®] vials sealed with latex-free rubber stoppers and aluminum flip-off seals. Vials are stored at $\leq -60^{\circ}\text{C}$.

B. Dosing & Route of Administration

A volume of 250 μL of rAAV8.aVEGF is administered as a single dose via subretinal delivery in the eye of a subject in need of treatment. The subject receives a dose of 3×10^9 GC/eye, 1×10^{10} GC/eye, or 6×10^{10} GC/eye.

rAAV8.aVEGF is administered by a single subretinal delivery by a retinal surgeon with the subject under local anesthesia. The procedure involves a standard 3-port pars plana vitrectomy with a core vitrectomy followed by subretinal delivery of rAAV8.aVEGF into the subretinal space by a subretinal cannula (38 gauge). The delivery is automated via the vitrectomy machine to deliver 250 μL to the subretinal space.

rAAV8.aVEGF can be administered in combination with one or more therapies for the treatment of wet AMD. For example, rAAV8.aVEGF is administered in combination with laser coagulation, photodynamic therapy with verteporfin, and intravitreal with anti-VEGF agent, including but not limited to pegaptanib, ranibizumab, aflibercept, or
 5 bevacizumab.

Starting at about 4 weeks post- rAAV8.aVEGF administration, a patient may receive intravitreal ranibizumab rescue therapy in the affected eye.

C. Patient Subpopulations

Suitable patients may include those:

- 10 Having a diagnosis of nAMD;
 Responsive to anti-VEGF therapy;
 Requiring frequent injections of anti-VEGF therapy;
 Males or females aged 50 years or above;
 Having a BCVA $\leq 20/100$ and $\geq 20/400$ (≤ 65 and ≥ 35 ETDRS letters) in the affected
 15 eye;
 Having a BCVA between $\leq 20/63$ and $\geq 20/400$ (≤ 75 and ≥ 35 ETDRS letters);
 Having a documented diagnosis of subfoveal CNV secondary to AMD in the affected eye;
 Having CNV lesion characteristics as follows: lesion size less than 10 disc areas
 20 (typical disc area is 2.54 mm^2), blood and/or scar $< 50\%$ of the lesion size;
 Having received at least 4 intravitreal injections of an anti-VEGF agent for treatment of nAMD in the affected eye in the 8 months (or less) prior to treatment, with anatomical response documented on SD-OCT; and/or
 Having subretinal or intraretinal fluid present in the affected eye, evidenced on SD-
 25 OCT.
 Prior to treatment, patients are screened and one or more of the following criteria may indicate this therapy is not suitable for the patient:
- CNV or macular edema in the affected eye secondary to any causes other than AMD;
 - 30 • Blood occupying $\geq 50\%$ of the AMD lesion or blood $> 1.0 \text{ mm}^2$ underlying the fovea in the affected eye;

- Any condition preventing VA improvement in the affected eye, e.g., fibrosis, atrophy, or retinal epithelial tear in the center of the fovea;
- Active or history of retinal detachment in the affected eye;
- Advanced glaucoma in the affected eye;
- 5 • Any condition in the affected eye that may increase the risk to the subject, require either medical or surgical intervention to prevent or treat vision loss, or interfere with study procedures or assessments;
- History of intraocular surgery in the affected eye within 12 weeks prior to screening (Yttrium aluminum garnet capsulotomy may be permitted if performed >10 weeks prior to
10 the screening visit.);
- History of intravitreal therapy in the affected eye, such as intravitreal steroid injection or investigational product, other than anti-VEGF therapy, in the 6 months prior to screening;
- Presence of an implant in the affected eye at screening (excluding intraocular lens).
- 15 • History of malignancy requiring chemotherapy and/or radiation in the 5 years prior to screening (Localized basal cell carcinoma may be permitted.);
- History of therapy known to have caused retinal toxicity, or concomitant therapy with any drug that may affect visual acuity or with known retinal toxicity, e.g, chloroquine or hydroxychloroquine;
- 20 • Ocular or periocular infection in the affected eye that may interfere with the surgical procedure;
- Myocardial infarction, cerebrovascular accident, or transient ischemic attacks within the past 6 months of treatment;
- Uncontrolled hypertension (systolic blood pressure [BP] >180 mmHg, diastolic BP
25 >100 mmHg) despite maximal medical treatment;
- Any concomitant treatment that may interfere with ocular surgical procedure or healing process;
- Known hypersensitivity to ranibizumab or any of its components or past hypersensitivity to agents like rAAV8.aVEGF;
- 30 • Any serious or unstable medical or psychological condition that, in the opinion of the Investigator, would compromise the subject's safety or successful participation in the study.

- Aspartate aminotransferase (AST)/alanine aminotransferase (ALT) $>2.5 \times$ upper limit of normal (ULN)
 - Total bilirubin $>1.5 \times$ ULN unless the subject has a previously known history of Gilbert's syndrome and a fractionated bilirubin that shows conjugated bilirubin $<35\%$ of total bilirubin
 - Prothrombin time (PT) $>1.5 \times$ ULN
 - Hemoglobin <10 g/dL for male subjects and <9 g/dL for female subjects
 - Platelets $<100 \times 10^3/\mu\text{L}$
 - Estimated glomerular filtration rate (GFR) <30 mL/min/1.73 m²
- Starting at about 4 weeks post- rAAV8.aVEGF administration, a patient may receive intravitreal ranibizumab rescue therapy in the affected eye for disease activity if 1 or more of the following rescue criteria apply:
- Vision loss of ≥ 5 letters (per Best Corrected Visual Acuity [BCVA]) associated with accumulation of retinal fluid on Spectral Domain Optical Coherence Tomography (SD-OCT)
 - Choroidal neovascularization (CNV)-related increased, new, or persistent subretinal or intraretinal fluid on SD-OCT
 - New ocular hemorrhage
- Further rescue injections may be deferred per the Investigator's discretion if one of the following sets of findings occur:
- Visual acuity is 20/20 or better and central retinal thickness is "normal" as assessed by SD-OCT, or
 - Visual acuity and SD-OCT are stable after 2 consecutive injections.
- If injections are deferred, they are resumed if visual acuity or SD-OCT get worse per the criteria above.
- D. Measuring Clinical Objectives**
- Primary clinical objectives include slowing or arresting the progression of retinal degeneration and slowing or preventing loss of vision. Clinical objectives are indicated by the elimination of or reduction in the number of rescue treatments using standard of care, for example, intravitreal injections with anti-VEGF agents, including but not limited to

pegaptanib, ranibizumab, aflibercept, or bevacizumab. Clinical objectives are also indicated by a decrease or prevention of vision loss and/or a decrease or prevention of retinal detachment.

Clinical objectives are determined by measuring BCVA (Best-Corrected Visual Acuity), intraocular pressure, slit lamp biomicroscopy, indirect ophthalmoscopy, and/or SD-OCT (SD-Optical Coherence Tomography). In particular, clinical objectives are determined by measuring mean change from baseline in BCVA over time, measuring the gain or loss of ≥ 15 letters compared to baseline as per BCVA, measuring mean change from baseline in CRT as measured by SD-OCT over time, measuring mean number of ranibizumab rescue injections over time, measuring time to 1st rescue ranibizumab injection, measuring mean change from baseline in CNV and lesion size and leakage area based on FA over time, measuring mean change from baseline in aqueous aVEGF protein over time, performing vector shedding analysis in serum and urine, and/or measuring immunogenicity to rAAV.aVEGF, i.e., measuring Nabs to AAV, measuring binding antibodies to AAV, measuring antibodies to aVEGF, and/or performing ELISpot.

Clinical objectives are also determined by measuring the mean change from baseline over time in area of geographic atrophy per fundus autofluorescence (FAF), measuring the incidence of new area of geographic atrophy by FAF (in subjects with no geographic atrophy at baseline, measuring the proportion of subjects gaining or losing ≥ 5 and ≥ 10 letters, respectively, compared with baseline as per BCVA, measuring the proportion of subjects who have a reduction of 50% in rescue injections compared with previous year, measuring the proportion of subjects with no fluid on SD-OCT.

Improvement/efficacy resulting from rAAV.aVEGF administration can be assessed as a defined mean change in baseline in visual acuity at about 4 weeks, 12 weeks, 6 months, 12 months, 24 months, 36 months, or at other desired timepoints. Treatment with rAAV.aVEGF can result in a 5%, 10%, 15%, 20%, 30%, 40%, 50% or more increase in visual acuity from baseline. Improvements/efficacy can be assessed as mean change from baseline in central retinal thickness (CRT) as measured by spectral domain optical coherence tomography (SD-OCT) at 4 weeks, 12 weeks, 6 months, 12 months, 24 months and 36 months. Treatment with rAAV.aVEGF can result in a 5%, 10%, 15%, 20%, 30%, 40%, 50% or more increase central retinal thickness from baseline.

EXAMPLE 2: Generation of AAV8.CMV.aVEGF

Each of the aVEGF vectors described herein include an expression cassette including a promoter which drives expression of the anti-VEGF Fab heavy chain and light chain, each of which has an IL2 leader sequence. The Fab coding sequence in the vector genomes carried by the rAAV in the tested composition (suspension) were designed to be identical. The expression cassette is flanked by a 5' AAV2 ITR and a 3' AAV2 ITR. Each of the tested vector genomes contains a coding sequence variant for the same anti-VEGF Fab (previously designated aVEGF-Arg or aVEGF-R). In certain embodiments, the expressed aVEGF Fab is a homogenous population. In certain embodiments, the expressed aVEGF Fab has heterogeneity at the heavy chain carboxy terminus. The open reading frames for the IL2-aVEGF heavy chain and IL2-aVEGF light chain were separated by an encoded furin cleavage site/F2A linker to promote equal molar expression of both, heavy and light chains. This results in expression of an aVEGF heavy chain which optionally further contains 0, 1, 2, 3 or 4 amino acids at its carboxy terminus: an arginine, arginine-lysine, arginine-lysine-arginine, or arginine - lysine - arginine - arginine at its carboxy terminus.

Various coding sequences are designated aVEGFv1, v2, etc. These vector genomes are provided in the Sequence Listing, which is incorporated by reference.

The following elements to be included in the transgene cassette in AAV2/8 vector for expression of anti-VEGF Fab in mice were evaluated.

- 7 different promoters (98 male C57BL/6 mice; Jackson Laboratories) were assessed using a convenient antibody (F16) expressed from AAV2/8. Expression of F16 mAb was measured by ELISA against hemagglutinin (HA) protein;
- 2 different leader peptides (28 male C57BL/6 mice; Jackson Laboratories). Expression of anti-VEGF Fab was measured by ELISA against VEGF;
- 3 different light-heavy chain separators (42 male C57BL/6 mice; Jackson Laboratories) were evaluated using the following vectors.

Group	Treatment	No. of animals		ROA
		Dose (GC/eye)		
		1.00×10 ⁹	5.00×10 ⁹	
1	AAV2/8.CMV.PI.aVEGFv7.EMCVIRES.Fab.SV40; the sequence of the expression cassette is provided in SEQ ID NO: 45	7	7	Subretinal
2	AAV2/8.CMV.PI.aVEGFv7.FMDV1IRES.Fab.SV40; the sequence of the expression cassette is provided in SEQ ID NO: 46	7	7	Subretinal
3	AAV2/8.CMV.PI.aVEGFv7.cMycIRES.Fab.SV40; the sequence of the expression cassette is provided in SEQ ID NO: 47	7	7	Subretinal

Abbreviation: GC = genome copies; No. = number; ROA = route of administration.

- 13 different coding sequences (182 male C57BL/6 mice; Jackson Laboratories).

Expression of anti-VEGF Fab was measured by ELISA against VEGF.

Vectors were delivered into subretinal space of the mouse eye. Expression of reporter genes was determined by enzyme-linked immunosorbent assay (ELISA).

- 5 Seven different promoters were evaluated in another study: 3 viral (cytomegalovirus [CMV], thymidine kinase [TK], simian virus [SV40]), 3 non-viral (phosphoglycerate kinase [PGK], human elongation factor-1 α [EF1 α], ubiquitin C [UbC]), and 1 hybrid (chicken β -actin [CB7]) promoters.

- 10 Two different leader peptides were also evaluated using rAAV8 vectors having identical vector elements and the same coding sequence, i.e., v3, with the exception of the leader sequence (interleukin-2 vs the serpin leader). AAV2/8 = adeno associated virus (AAV) capsid type 8 with AAV2 inverted terminal repeats flanking the transgene; amd201Lead = anti-VEGF Fab with IL2 leader sequence and Furin F2A as light-heavy chain separator; amd201altLead = anti-VEGF Fab with SF1 leader sequence and Furin F2A as
15 light-heavy chain separator; CB7 = chicken β actin promoter; CI = chimeric intron; rBG = rabbit β globin polyadenylation sequence.

Three different internal ribosome entry site (IRES) sequences separating heavy and light chains of anti-VEGF Fab were evaluated in another study. These IRES sequences were derived from encephalomyocarditis virus (EMCV), cMyc, and foot-and-mouth disease virus 1 (FMDV1). In this study, the vectors were identical except for the light and heavy chain
 5 separators (EMC, FMDV1 and cMyc). AAV2/8 = adeno associated virus (AAV) capsid type 8 with AAV2 inverted terminal repeats flanking the transgene; amd201 = codon variant of anti-VEGF Fab with IL2 leader sequence; CMV = cytomegalovirus promoter; EMCV = encephalomyocarditis virus; Fab = fragment antigen-binding region; FMDV1 = foot and mouth disease virus 1; IRES = internal ribosome entry site; PI = Promega intron; SV40 =
 10 simian virus polyadenylation sequence

In another study, thirteen different coding sequences for the anti-VEGF Fab were evaluated. The overall coding sequence variance was between approximately 20% and 30 %. The vectors are described in the following Table.

Vectors with Different Coding Sequences Used. The SEQ ID NO for the expression
 15 cassettes is provided in the following table:

Vector	SEQ ID NO: (vector genome)
AAV2/8.CB7.CI.aVEGFv4.rBG	35
AAV2/8.CB7.CI.aVEGFv5.rBG	36
AAV2/8.CB7.CI.aVEGFv1.rBG	34
AAV2/8.CB7.CI.aVEGFv2.rBG:	3
AAV2/8.CB7.CI.aVEGFv6.rBG	37
AAV2/8.CB7.CI.aVEGFv7.rBG	38
AAV2/8.CB7.CI.aVEGFv8.rBG	39
AAV2/8.CB7.CI.aVEGFv9.rBG	40
AAV2/8.CB7.CI.aVEGFv10.rBG	41
AAV2/8.CB7.CI.aVEGFv11.rBG	42

Vector	SEQ ID NO: (vector genome)
AAV2/8.CB7.CI.aVEGFv12.rBG	43
AAV2/8.CB7.CI.aVEGFv3.rBG	14
AAV2/8.CB7.CI.aVEGFv13.rBG	44

Vectors in all studies were diluted in Dulbecco's phosphate-buffered saline (DPBS).

Animals were assigned into treatment groups and administered 1.00×10^9 or 5.00×10^9 genome copies (GC)/eye of AAV2/8 vectors into the right eye. The left eye was used as an untreated control. Vectors were administered subretinally in a total volume of 1 μ L.

A. Subretinal Injections

Subretinal injections were conducted using aseptic technique and sterile dissecting instruments. Animals were anesthetized with ketamine/xylazine or 3% to 5% isoflurane and administered meloxicam. Animals were then placed under a dissection microscope with the eye to be injected under view (using a 15 \times magnification). The temporal conjunctiva was grasped with jeweler's forceps and carefully cut down to the sclera using the tip of Vannas iridotomy scissors. Conjunctival peritomy was conducted by introducing the lower lip of the scissors through the incision and extending circumferentially both superiorly and inferiorly of the conjunctiva. Any conjunctival debris was carefully removed from the surface of the sclera. The conjunctiva adjacent to the cornea was grasped with the forceps, providing traction to rotate the globe and allow optimal surgical exposure. Using a 30 $\frac{1}{2}$ -gauge needle, a small incision, large enough to allow the blunt-tip needle to pass through, was made.

The tip of a 33-gauge blunt-tip needle mounted on a Hamilton auto-injector syringe was introduced into the incision tangentially to the surface of the globe. The needle was passed along the inner surface of the sclera with the tip entering approximately 1 mm. The 33-gauge needle passed through the sclera and choroid and then terminated in the subretinal space. Up to 1 μ L of vector was delivered. Once the procedure was completed, antibiotic ophthalmic ointment was applied to the eye.

B. Assay Methods

The collected eyes were homogenized by placing entire eyeball into a conical tube with stainless steel beads and 200 μ L of cocktail containing protein lysis and extraction buffer (RIPA) and cOmplete™, Mini Protease Inhibitor Cocktail tablets (1 tablet/10 mL of RIPA buffer). The eyes were homogenized for at least 2 minutes in a TissueLyser (Qiagen, USA) or until fully homogenized. Homogenate was centrifuged for 20 minutes at 12000 RPM at 4°C in a cold room. The supernatants were transferred into fresh tubes and used in analytical assays.

Determination of Protein Concentrations in Eye Homogenates

Protein concentration in eye homogenate was determined using Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific) per the manufacturer's instructions. Equal amounts of protein in all samples were used in ELISA.

Enzyme-Linked Immunosorbent Assay

Ninety-six-well, round-bottom plates were coated with 2 μ g/mL of HA A-Beijing or 1 μ g/mL VEGF overnight at 4°C. After coating, the plates were washed 5 times with 200 μ L of phosphate-buffered saline (PBS) with 0.05% Tween-20 (PBS-T) using a 405 TS Washer (BioTek Instruments, Winooski, VT). Plates were then blocked with 200 μ L/well of 1% bovine serum albumin (BSA) at room temperature (RT) for 1 hour. After washing (as described), 100 μ L/well of sample was loaded into duplicate wells and incubated at 37°C for 1 hour. Following incubation, the plates were washed (as described) and then blocked with 1% BSA at RT for 1 hour. After washing (as described), 100 μ L/well of the primary antibody was added and incubated at RT for 1 hour. Wells were then washed (as described) and incubated with 100 μ L/well of the secondary antibody at RT for 1 hour. Following a final wash (as described), 150 μ L/well of 3,3',5,5'-tetramethylbenzidine, a detection substrate, was added and incubated at RT for 30 minutes protected from light. The reaction was stopped with 50 μ L/well of 2N H₂SO₄. The plates were then read at the excitation/emission of 450 nm/540 nm using spectrophotometer SpectraMax® M3 (Molecular Devices, Sunnyvale, CA).

The following primary antibodies were used: 1.0 mg/mL Goat Anti-Human IgG H&L (Biotin) preadsorbed at a 1:10000 dilution in PBS (Abcam 0.5 mg/mL); 0.5 mg/mL Goat Anti-Human IgG H&L (Biotin) preadsorbed at a 1:5000 dilution in PBS (Abcam, 1

mg/mL). The following secondary antibody was used: 1 mg/mL Streptavidin (HRP) at a 1:30000 dilution in PBS.

Statistical Analyses

Average and standard deviation values for concentration of reporter genes for ELISA were calculated using Microsoft Office Excel 2010.

C. RESULTS

AAV2/8 vectors with 7 different promoters were evaluated for expression of FI6 mAb. Expression of FI6 mAb was not observed in any animal when promoter EF 1- α was used. Expression of FI6 mAb was low when promoters SV40.PI, PGK.PI and TK.PI were used. Promoters CMV.PI, CB7.CI, and UbC.PI demonstrated the highest expression of FI6 mAb. No expression was observed in the untreated left eye in any animal (data on file). AAV2/8 vectors with 2 different leader peptides were evaluated for expression of anti-VEGF Fab. Compared to aVEGFv7 with an IL2 leader, expression of anti-VEGF Fab at low dose was greater when leader peptide aVEGFv7 with SF2 leader was used. At high dose, the expression of anti-VEGF Fab was similar for both leader peptides. No expression was observed in the untreated left eye in any animal (data on file). AAV2/8 vectors with 3 different light-heavy chain separators were evaluated for expression of anti-VEGF Fab. Expression of anti-VEGF Fab was not observed in any animal when cMyc light-heavy chain separator was used. With EMCV and FMDV1 light-heavy chain separators, anti-VEGF Fab was expressed at low levels. No expression was observed in the untreated left eye in any animal (data on file).

AAV2/8 vectors with 13 different coding sequences were evaluated for expression of anti-VEGF Fab transgene product. When coding sequences aVEGFv4, aVEGFv5, aVEGFv6, aVEGFv7, aVEGFv8, and aVEGFv9 were used, expression of anti-VEGF Fab was low. Expression was higher with coding sequences aVEGFv13, aVEGFv10, aVEGFv11, and aVEGFv12. When coding sequences aVEGFv1, aVEGFv2, and aVEGFv3 were used, expression of anti-VEGF Fab was the highest. No expression was observed in the untreated left eye in any animal (data on file).

Each of the vectors encodes the same anti-VEGF transgene product. Based in part on these results, a single replication-defective, recombinant AAV8.aVEGF was selected for further development. This tert vector has an AAV8 capsid and a vector genome in which AAV2 ITRs flank a CB7 promoter, an intron, an anti-VEGF coding sequence selected from

the coding sequence as described earlier and a rBG poly A sequence. This is termed the test vector (alternatively AAV2/8.aVEGF test vector or AAV8.aVEGF test vector) in the following examples, except where specifically specified otherwise.

5 **EXAMPLE 3: Pharmacokinetic (PK) Study in Non-Human Primates**

Macaques were used in this study because they are the closest species to humans for studying retinal diseases. Cynomolgus monkeys and humans have similar eye anatomy, including fovea. The dimensions of the eyes are comparable, which allows determination of the human dose based on relative retinal areas.

10 This study was conducted to select AAV2/8 vector for clinical development and to evaluate toxicity and immunogenicity of AAV2/8 vector and anti-VEGF Fab in cynomolgus monkeys. The study is ongoing. The results presented are based on the data collected at Month 10. Evaluation of toxicity of AAV2/8 vector and anti-VEGF Fab is described. Animals were administered AAV2/8 vectors subretinally. Toxicity was evaluated based on
 15 clinical observations, body weights, indirect ophthalmoscopy, spectral domain optical coherence tomography, hematology, coagulation, clinical chemistry, and gross pathologic findings. The only adverse finding related to AAV2/8 vector or anti-VEGF Fab was some thinning in outer nuclear layer localized to the injection site observed by spectral domain optical coherence tomography in several eyes of animals administered 1.00×10^{11} GC/eye of
 20 AAV2/8 vectors.

Animals were assigned into 4 treatment groups. Animals were administered a single dose of 1.00×10^{11} genome copies (GC)/eye of AAV2/8 vectors into each eye in a total volume of 100 μ L. Vectors were administered subretinally (confirmed visually by appearance of a dome shaped retinal detachment/retinal bleb under microscope) into both
 25 eyes. The following table lists the tested vectors.

Group	Treatment	Dose (GC/eye)	No. of animals	ROA	
				OD	OS
2	AAV2/8.UbC.PI.aVEGFv2.SV40	1.00×10^{11}	2 M, 2F	Subretinal	
3	AAV2/8.UbC.PI.aVEGFv3.SV40	1.00×10^{11}	2 M, 2 F	Subretinal	
5	AAV2/8.CB7.CI.aVEGFv2aVEGFv2.rBG	1.00×10^{11}	2 M, 2 F	Subretinal	

Group	Treatment	Dose (GC/eye)	No. of animals	ROA	
				OD	OS
6	AAV2/8.CB7.CI.aVEGFv3.rBG	1.00×10^{11}	2 M, 2 F	Subretinal	

Abbreviation: F = female; GC = genome copies; M = male; No. = number; OD = right eye; OS = left eye; ROA = route of administration.

Subretinal Injections

For subretinal injections, a needle was inserted through a trocar, introduced by sclerotomy, at the 2 or 10 o'clock position. The needle was advanced through the vitreous to penetrate the retina in the posterior pole. Under the microscopic control, 100 μ L of test article was injected into the subretinal space. This was confirmed by appearance of a dome shaped retinal detachment/retinal bleb. If the first injection attempt did not result in retinal detachment, the cannula was moved to another site in the retina. The injection site may have resulted in a temporary scotoma. The injected solution was reabsorbed within a few hours by the retina. The retinal detachment was made in the peripheral retina and did not result in permanent blindness. The site of sclerotomy was sutured with absorbable suture and the eye dressed with PredG ointment or equivalent. Subconjunctival kenalog or equivalent was administered. Animals were observed daily and administered parenteral analgesics as needed. If vitreal inflammation appeared, animals were treated with topical atropine and PredG ointment or equivalent daily until symptoms resolved.

Collection of Anterior Chamber Fluid

Animals were anesthetized and their head stabilized. Betadine 5% antiseptic solution and Proparacaine or equivalent were applied to each eye. An eye speculum was placed to allow access to the anterior chamber. The procedure was performed with a tuberculin syringe attached to a 27- to 30-gauge hypodermic needle. The eye was held steady with forceps or a cotton tip applicator on the nasal conjunctiva. The needle was inserted bevel up through the paralimbal peripheral clear cornea, anterior to the iris plane. Once the eye was entered, a sampler slowly withdrew the plunger of the syringe to aspirate the aqueous fluid. A maximum of 100 μ L of anterior chamber fluid was collected. Once anterior chamber fluid was drained, the needle was withdrawn from the eye. The anterior chamber fluid was placed on wet ice until use or storage. After the procedure, topical flurbiprofen, PredG ointment,

and antibiotic drops were applied to each eye. Anterior chamber fluid was collected on the following Study days (occasionally adjusted due to weekends, holidays or scheduling issues)

- 0, 15, 29, 43, 57, 71, 85, 120, 149, 183, 212, 247, 274, and 302.

Spectral Domain Optical Coherence Tomography

5 Retinal structure (at a micron-level resolution) was evaluated by *in vivo*, non-invasive, cross-sectional retinal microscopy with SD-OCT (Spectralis OCT, Heidelberg Engineering, Carlsbad, CA). Pupils were dilated with phenylephrine 2.5% and tropicamide 1%. En-face retinal imaging was performed with near infrared (NIR) reflectance (REF) and in a subset of animals with NIR fundus autofluorescence (FAF) using the scanning laser
10 ophthalmoscope of this imaging system. Spectral domain optical coherence tomography scanning was performed with 9 mm long horizontal and vertical cross-sections through the fovea and overlapping 30 × 25 mm raster scans extending into the near midperiphery. See, Aleman, Invest Ophthalmol Vis Sci. 2007 Oct;48(10):4759-65.

Enzyme-Linked Immunosorbent Assay (ELISA)

15 The ELISA was performed essentially as described for the mouse studies above. Ninety-six-well, round-bottom plates were coated with 1 µg/mL of VEGF for expression of anti-VEGF Fab. Plates were coated overnight at 4°C. After coating, the plates were washed 5 times with 200 µL of phosphate-buffered saline (PBS) with 0.05% Tween-20 (PBS-T) using a 405 TS Washer (BioTek Instruments, Winooski, VT). Plates were then blocked with
20 200 µL/well of 1% bovine serum albumin (BSA) at room temperature (RT) for 1 hour. After washing (as described), 100 µL/well of sample was loaded into duplicate wells and incubated at 37°C for 1 hour. Following incubation, the plates were washed (as described) and then blocked with 1% BSA at RT for 1 hour. After washing (as described), 100 µL/well of the primary antibody was added and incubated at RT for 1 hour. Wells were then washed
25 (as described) and incubated with 100 µL/well of the secondary antibody at RT for 1 hour.

Following a final wash (as described), 150 µL/well of 3,3',5,5'-tetramethylbenzidine, a detection substrate, was added and incubated at RT for 30 minutes protected from light. The reaction was stopped with 50 µL/well of 2N H₂SO₄. The plates were then read at the excitation/emission of 450 nm/540 nm using spectrophotometer SpectraMax[®] M3
30 (Molecular Devices, Sunnyvale, CA). The following primary antibodies were used: 1.0 mg/mL Goat Anti-Human IgG H&L (Biotin) preadsorbed at a 1:10000 dilution in PBS; 0.5 mg/mL Goat Anti-Human IgG H&L (Biotin) preadsorbed at a 1:5000 dilution in PBS. The

following secondary antibody was used: 1 mg/mL Streptavidin (HRP) at a 1:30000 dilution in PBS.

Average and standard deviation values for concentrations of the anti-VEGF Fab in anterior chamber fluid and blood were calculated using Microsoft Office Excel 2010.

5 A Pharmacology Results

Four AAV vectors with different promoters and coding sequences were evaluated as described earlier in this example. Vectors were administered subretinally. Expression of anti-VEGF Fab was determined by enzyme-linked immunosorbent assay.

Expression of Anti-VEGF Fab in Anterior Chamber Fluid

10 In anterior chamber fluid of animals in all groups, similar expression kinetics was observed (FIG 3A-3D; FIG 4A-4D). Onset of expression of the anti-VEGF Fab was rapid, generally within 7 days. Steady-state expression levels were achieved within 1 month. All except 2 animals continued to express the anti-VEGF Fab at steady-state levels until the last evaluated timepoint.

15 One animal in Group 2 (FIG 3B) and 1 animal in Group 5 (FIG 4A) lost expression of the anti-VEGF Fab. Loss of expression coincided with appearance of antibodies against the anti-VEGF Fab. No difference in expression of the anti-VEGF Fab between males and females and between the right and the left eye was observed.

20 Generally, vectors controlled by the CB7.CI promoter (FIG 4A-4D) expressed the anti-VEGF Fab at higher levels than vectors controlled by the UbC.PI promoter (FIG 3). Vector AAV2/8.CB7.CI.aVEGFv3aVEGFv3.rBG was selected as a primary vector for clinical development. This selection was based on expression level of transgene, better translatability of relative expression levels from mice to cynomolgus monkeys for aVEGFv3 coding sequence, and greater level of experience with CB7.CI promoter.

25 Expression of Anti-VEGF Fab in Blood

In some patients administered a single IVT injection of Lucentis, ranibizumab was observed in serum (Xu, 2013). To determine if subretinal administration of AAV2/8 vector results in systemic exposure to the anti-VEGF Fab, its concentrations were measured in serum.

30 Expression of the anti-VEGF Fab was around the baseline levels in blood of all animals (FIGs 3A-3D, FIGs 4A-3D).

B. Toxicology

Evaluation of toxicity of AAV2/8 vector and anti-VEGF Fab is described in this subpart B. Animals were administered AAV2/8 vectors subretinally. Toxicity was evaluated based on clinical observations, body weights, indirect ophthalmoscopy, spectral domain optical coherence tomography (SD-OCT), hematology, coagulation, clinical chemistry, and gross pathologic findings.

For each variable in each treatment group, the measurements at each time point were compared to the corresponding baseline values using Wilcoxon rank-sum test. The Wilcoxon rank-sum test is a nonparametric alternative to the two-sample t-test, which is based solely on the order in which the observations from the 2 samples fall. It is a preferable test for dataset with small sample size. Statistical significance was declared at the 0.05 level without the adjustment for multiple testing. The analysis was done using R program (version 3.3.1; cran.r-project.org/) with function “wilcox.test”.

The study is ongoing. The results presented are based on the data collected by Month 10. There were no mortalities in this study. No adverse clinical observations related to AAV2/8 vector or anti-VEGF Fab were noted in any animal. No clinically meaningful changes in body weight during the study were observed for any animal. No adverse observations related to AAV2/8 vector or anti-VEGF Fab were noted during indirect ophthalmoscopy in any animal.

Spectral Domain Optical Coherence Tomography

All 4 animals (8 eyes) in Group 6 were imaged by SD OCT. Injected regions in the eyes that received the intermediate dose level of AAV8.aVEGF test vector (1.00×10^{11} GC/eye) showed intermediate outcomes as compared to 1.00×10^{10} and 1.00×10^{12} dose levels described in Example 7 (esp., subpart B). In two animals (Animal C71896 and Animal C65936), some thinning of ONL was observed (data on file). In addition, in 2 animals (Animal C74422 and Animal C74414), minimal changes were observed (data on file). No clinically significant changes in hematology, coagulation, or clinical chemistry parameters were observed in any animal.

No findings related to AAV2/8 vector or anti-VEGF Fab were observed in 2 animals sacrificed at Month 10. In Animal C65936, a liver mass was observed, which microscopically was focal chronic grade 3 inflammation. In Animal C74414, bilateral grade 3 lymphoid hyperplasia was observed. These findings were not related to AAV2/8 vector or anti-VEGF Fab. At a dose level of 1.00×10^{11} GC/eye of AAV2/8 vector, the only findings

related to AAV2/8 vector or anti-VEGF Fab were minimal vacuolation of the lens of the right eye of Animal C65926. Minimal perivascular mononuclear cell infiltrates around the vasculature of the right optic nerve in Animal C74414 were observed. In the same animal, a minimal mononuclear cell infiltrate in the subconjunctiva of the left eye and a minimal perivascular exocular mononuclear cell infiltrate in right eye were also noted.

The only adverse finding related to AAV2/8 vector or anti-VEGF Fab was some thinning in ONL localized to the injection site observed by SD OCT in several eyes of animals administered 1.00×10^{11} GC/eye of AAV2/8 vectors.

C. Immunology

In this section, evaluation of immunogenicity of AAV2/8 vector and anti-VEGF Fab is described. Vectors were administered subretinally as described earlier in this example. Immunogenicity was assessed by the presence of IgM and IgG antibodies against anti-VEGF Fab, neutralizing antibodies against AAV8 capsid, and cellular immune response against AAV2/8 vector and anti VEGF Fab.

To summarize, one animal each in Groups 2 and 5 had antibodies against anti-VEGF Fab, high levels of NAb to AAV8 capsid, and T-cell responses. Both animals lost expression of anti-VEGF Fab. Animals with pre-existing NAb to AAV8 capsid generally had increased response after administration of AAV2/8 vector when compared to animals without pre-existing NAb. In some animals sacrificed at approximately Study day 300, NAb to AAV8 capsid were observed in vitreous fluid. No antibodies against anti-VEGF Fab and no T-cell responses were observed in animals in Group 6. Only mild fluctuations in the levels of NAb were observed in animals in Group 6 after administration of AAV2/8 vector.

Anti-VEGF Fab was expressed in all animals administered AAV2/8 vectors (Part A of this Example). Two animals (Animal C74440 and Animal C68127) lost expression of the anti-VEGF Fab. Loss of expression coincided with appearance of antibodies against the anti-VEGF Fab.

Overall, levels of IgM and IgG against anti-VEGF Fab were below baseline levels in anterior chamber fluid and serum. Increases above the baseline levels at some timepoints were observed in some animals. In 1 animal each in Groups 2 (Animal C74440) and 5 (Animal C68127), levels of IgG against anti-VEGF Fab in anterior chamber fluid increased above baseline levels approximately at approximately 6 months. The levels generally

increased thereafter. In both animals, IgG against anti-VEGF Fab increased above the baseline levels in serum. These increases in IgG coincided with loss of expression of anti-VEGF Fab. Importantly, IgM and IgG against anti-VEGF Fab in animals in Group 6 were not detected or were below the baseline levels for the duration of the study.

5 In brief, the animal immune system permitted continued localized expression of anti-VEGF transgene product, despite the fact that the transgene product is a human antibody.

Presence of Neutralizing Antibodies Against AAV8 Capsid

Baseline levels of NAbS against AAV8 capsid were determined in serum from blood samples collected on Study day 0. The limit of detection was a 1:5 dilution; titers of < 5 were considered undetectable. In 2 of 16 animals (Animal C63116 and Animal C66122), pre-existing NAbS in serum were not observed. The levels of NAbS in these 2 animals following administration of AAV2/8 vectors remained below the limit detection or were low. In 14 of 16 animals, pre-existing NAbS were observed. In 11 of these animals, levels of NAbS fluctuated throughout the study. In 1 animal each in Groups 2 (Animal C74440) and 5 (Animal C68127), levels of NAbS following administration of AAV2/8 vector increased up to 256 and 128 two-fold dilutions respectively at two months. These increases in NAbS coincided with loss of expression of anti-VEGF Fab. In 6 sacrificed animals from Groups 2, 3 and 5, the presence of NAbS was evaluated in vitreous fluid. In 2 animals (Animal C63116 and Animal C66122), with undetectable NAb in serum time of sacrifice, NAbS were not present in vitreous fluid at sacrifice. In the remaining animals, levels of NAbS at sacrifice did not correlate with levels in serum at time of sacrifice. In all animals in Group 6, pre-existing NAbS were observed. The levels of NAbS in these animals fluctuated mildly throughout the study.

T-Cell Responses to AAV2/8 Vector and Anti-VEGF Fab

25 In 1 animal (Animal C74440) in Group 2, elevated T-cell responses were observed at a single time point. In this animal, antibodies against anti-VEGF Fab and NAbS against AAV8 capsid were also observed. This animal lost expression of anti-VEGF Fab. In 1 animal (Animal C65873) in Group 5, displayed sustained T-cell responses to the pool B peptides of AAV8 capsid were observed including pre-injection baseline sample. The same animal had the highest levels of NAbS after administration of AAV2/8 vector. Another animal (Animal C68127) in the same group developed T-cells to all peptide pools of AAV8 capsid which were not sustained over time. This animal had antibodies against anti-VEGF

Fab and the second highest level of NAb. The animal lost expression of anti-VEGF Fab.

No other sustained T-cell responses to the transgene product were observed. No sustained T-cell responses were observed in animals in Group 6.

5 EXAMPLE 4 - Animal Models Useful For Evaluating AAV2/8.aVEGF and Anti-VEGF Transgene Product

VEGF transgenic mice are used as animal models of Wet AMD. Two such models include the Rho/VEGF mouse model and the Tet/opsin/VEGF model.

A. Rho/VEGF Mouse Model

10 Rho/VEGF mice are transgenic mice in which the rhodopsin promoter drives expression of human vascular endothelial growth factor (VEGF165) in photoreceptors, causing new vessels to sprout from the deep capillary bed of the retina and grow into the subretinal space, starting at postnatal Day 10. The production of VEGF is sustained and therefore the new vessels continue to grow and enlarge and form large nets in the subretinal
15 space similar to those seen in humans with neovascular age-related macular degeneration. See Tobe, Takao, et al. "Evolution of neovascularization in mice with overexpression of vascular endothelial growth factor in photoreceptors." *Investigative ophthalmology & visual science* 39.1 (1998): 180-188.

An enzyme-linked immunosorbent assay (ELISA) can be performed as follows.
20 Briefly, plates are coated with 1µg/mL of VEGF overnight at 4°C. 1% BSA is used as blocking buffer and is allowed to incubate at room temperature for 1 hour at 200µL per well. Samples are loaded in duplicate at 100µL per well and incubated for 1 hour at 37°C, followed by a second blocking buffer incubation. The primary antibody is a goat Anti-Human IgG H&L conjugated with Biotin which is left to incubate for 1 hour at room
25 temperature at 100µL per well. Secondary antibody is a 1:30,000 dilution of Streptavidin, loaded at 100µL per well and incubated at room temperature for 1 hour. TMB solution is used as detection substrate (0.1M NaOAc Citric Buffer (pH 6.0), Hydrogen Peroxide, 100X TMB Stock), loaded at 150 µL per well and incubated at room temperature for 30 minutes without exposure to light. 50µL of stop solution (2N H₂SO₄) is added to each well, and each
30 plate was then read at 450nm-540nm.

In one study performed using this model and a test AAV8.aVEGF as described in the preceding examples, the ELISA results were as follows:

AAV8.aVEGF		Dose (GC/eye)	Eye 1R	Eye 1L	Eye 2R	Eye 2L	Eye 3R	Eye 3L
		1.00E+10	0.00	0.00	0.00	0.00	0.00	0.00
AAV8.aVEGF		1.00E+08	0.00	0.00	0.00	0.00	0.00	313.58
AAV8.aVEGF		3.00E+08	0.00	0.00	0.00	0.00	0.00	0.00
AAV8.aVEGF		1.00E+09	0.00	530.45	0.00	0.00	324.01	0.00
AAV8.aVEGF		3.00E+09	0.00	0.00	0.00	0.00	208.71	0.00
AAV8.aVEGF		1.00E+10	232.23	239.19	139.30	0.00	0.00	0.00

Vector	Dose (GC/eye)	Eye 4R	Eye 4L	Eye 5R	Eye 5L
Empty	1.00E+10	0.00	0.00	0.00	0.00
AAV8.aVEGF	1.00E+08	0.00	251.56	0.00	0.00
AAV8.aVEGF	3.00E+08	0.00	0.00	0.00	0.00
AAV8.aVEGF	1.00E+09	0.00	564.75	0.00	0.00
AAV8.aVEGF	3.00E+09	355.31	207.95		
AAV8.aVEGF	1.00E+10	134.53	0.00	214.05	167.79

Anti-VEGF FAb levels are shown in ng/eye.

5

B. Tet/opsin/VEGF Mouse Model

Tet/opsin/VEGF mice are transgenic mice that are normal until given doxycycline in drinking water. Doxycycline induces very high photoreceptor expression of vascular endothelial growth factor (VEGF), leading to massive vascular leakage, culminating in total exudative retinal detachment in 80-90% of mice within 4 days of induction. See, Ohno-Matsui, Kyoko, et al. "Inducible expression of vascular endothelial growth factor in adult mice causes severe proliferative retinopathy and retinal detachment." The American journal of pathology 160.2 (2002): 711-719.

The ELISA can be performed as described in Part A of this Example. In one study performed using this model and a test AAV8.aVEGF as described in the preceding examples, the ELISA results were as follows. Results are shown in the tables below as the average ± standard deviation (Std).

			Mouse Eye ID's			
			1R	1L	2R	2L
Sample	Vector	Dose (GC/eye)	Avg±Std	Avg±Std	Avg±Std	Avg±Std
Empty	Empty	1.00E+10	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
1.00E+11	AAV8.aVEGF	1.00E+08	0.00±0.00	216.34±14.85	0.00±0.00	0.00±0.00
3.00E+11	AAV8.aVEGF	3.00E+08	88.23±0.10	106.54±1.01	0.00±0.00	0.00±0.00
1.00E+12	AAV8.aVEGF	1.00E+09	424.07±19.26	0.00±0.00	344.51±30.67	0.00±0.00
3.00E+12	AAV8.aVEGF	3.00E+09	581.28±50.45	175.23±20.45	254.13±21.33	477.85±34.54
1.00E+13	AAV8.aVEGF	1.00E+10	366.10±20.76	309.06±2.45	234.42±4.78	173.46±1.86

			Mouse Eye ID's					
			3R	3R	4R	4L	5R	5L
Sample	Vector	Dose (GC/eye)	Avg± Std	Avg± Std	Avg± Std	Avg ± Std	Avg ± Std	Avg ± Std
Empty	Empty	1.00E+10	0.00± 0.00	0.00± 0.00	0.00± 0.00	0.00± 0.00	0.00± 0.00	0.00
1.00E+11	AAV8.aVEGF	1.00E+08	0.00± 0.00	0.00± 0.00	0.00± 0.00	0.00± 0.00	0.00± 0.00	322.39± 23.43
3.00E+11	AAV8.aVEGF	3.00E+08	0.00± 0.00	0.00± 0.00	88.25± 1.96	0.00± 0.00	150.83± 26.63	444.96± 54.45
1.00E+12	AAV8.aVEGF	1.00E+09	0.00± 0.00	0.00± 0.00	537.61± 17.07	0.00± 0.00	0.00± 0.00	0.00± 0.00
3.00E+12	AAV8.aVEGF	3.00E+09	366.10± 23.68	285.85± 27.94	456.01± 27.81	155.71± 12.75	778.20± 143.25	270.15± 10.85
1.00E+13	AAV8.aVEGF	1.00E+10	498.95± 26.64	289.96± 8.95	291.74± 7.96	338.82± 6.03	165.10± 4.85	301.23± 21.10

5

C. Other Animal Models

Other animal models of Wet AMD are utilized. In the laser trauma model, high-powered, focused laser energy is used to induce a break in Bruch's membrane. Subretinal injection of matrigel, VEGF, macrophages, lipid hydroperoxide, and/or polyethylene glycol induces choroidal neovascularization (CNV), a wet AMD pathology. See Pennesi, Mark E.,

Martha Neuringer, and Robert J. Courtney. "Animal models of age related macular degeneration." *Molecular aspects of medicine* 33.4 (2012): 487-509.

Optimized rAAV.aVEGF vectors are generated, diluted and delivered into subretinal space of the transgenic mice eye with dosage described in the previous examples.

- 5 Expressions of reporter genes, VEGF and anti-VEGF antibodies in the eye and/or plasma are determined by PCR, qPCR, ddPCR, oqPCR, Western Blot and ELISA as described in previous Examples. Electron Microscopy and Immunohistochemical analysis are also performed to evaluate the retinal neovascularization. The number of lesions per retina, area per lesion, neovascularization area per retina and traction retinal detachment
- 10 Histopathological Evaluation of Retinas are quantified.

EXAMPLE 5 - Assessment of Expression of Anti-VEGF Fab (Transgene Product) in Cynomolgus Monkeys

- This study was conducted to assess the expression of the anti-VEGF Fab (transgene product) and to evaluate toxicity, immunogenicity, and biodistribution of an AAV8 vector
- 15 expressing the anti-VEGF Fab following its administration in cynomolgus monkeys. In this report, expression of the transgene product and immunogenicity of the vector are described. Animals were administered an AAV2/8.aVEGF vector as described in these Examples or FFB-314 (control article) subretinally. Expression of transgene product in anterior chamber
- 20 fluid and blood was determined by enzyme linked immunosorbent assay (ELISA). Immunogenicity was assessed by the presence of neutralizing antibodies (NAbs) against AAV8 capsid before and after administration. The transgene product is expressed in anterior chamber fluid of all animals administered the vector. The transgene product is not expressed in blood. Increase in levels of NAbs was observed in 1 animal (C73723) administered
- 25 AAV8.aVEGF; this animal had pre-existing NAbs.

Animals in this study were administered a single dose of 1.00×10^{12} genome copies (GC)/eye of AAV8.CB7.CI.aVEGFv3.rBG or formulation buffer, FFB-314.

- AAV8.CB7.CI.aVEGFv3.RBG and FFB-314 were administered subretinally into the right eye (confirmed visually by appearance of a dome shaped retinal detachment/retinal bleb
- 30 under microscope) in a total volume of 100 μ L.

Animals were randomized using www.jamestease.co.uk/team-generator. One of 4 animals was selected using www.randomizer.org/ by random and assigned to Group 2. The

remaining 3 animals were assigned to Group 1. Group designation and dose levels for this study are presented in the following Table.

Group	Treatment ^a	Dose (GC/eye)	No. of animals	Follow-up
1	AAV8.aVEGF	1.00×10^{12}	1 M, 2 F	7 days
2	FFB-314	NA	1 M	

Abbreviation: F = female; GC = genome copies; M = male; NA = not applicable; No. = number. ^a Test and control articles were administered subretinally into the right eye.

Animals were euthanized on Study day 7. Samples of anterior chamber fluid and blood were collected for determination of expression of the anti-VEGF Fab transgene product and/or the presence of NAbS against AAV8 capsid.

Subretinal injections were performed as described in the earlier examples. Collection of anterior chamber fluid was as described in earlier examples. For the ELISA, ninety-six-well, round-bottom plates were coated with 1 µg/mL of VEGF for expression of the anti-VEGF Fab transgene product, or 0.5 µg/mL of a commercial anti-VEGF Fab for expression of IgM and IgG against the Anti-VEGF Fab transgene product. The ELISA methods were as described in the earlier examples.

The following primary antibodies were used: 1.0 mg/mL Goat Anti-Human IgG H&L (Biotin) preadsorbed at a 1:10000 dilution in PBS; 0.5 mg/mL Goat Anti-Human IgG H&L (Biotin) preadsorbed at a 1:5000 dilution in PBS. The following secondary antibody was used: 1 mg/mL Streptavidin (HRP) at a 1:30000 dilution in PBS.

Neutralizing Antibody Assay

Neutralizing antibody responses to AAV8 capsid were analyzed as follows. A Poly D lysine-coated 96-well black-walled/clear-bottom plate was seeded with human embryonic kidney 293 (HEK293) cells at 1×10^5 cells/well (referred to as a cell plate); the plate was incubated at 37°C overnight. The following day, the serum sample was heat-inactivated at 56°C for 35 minutes. The heat-inactivated sample and a recombinant vector (AAV8.CMV.LacZ at 1×10^9 GC/well; provided by the Penn Vector Core at the University of Pennsylvania) were used to formulate a serum-vector plate. The recombinant vector was diluted in serum-free Dulbecco's Modified Eagle Medium (DMEM) and incubated with 2-fold serial dilutions (starting at 1:5) of the heat inactivated samples at 37°C for 1 hour. Prior

to combining the serum vector plate with the cell plate, the HEK293 cells (now at 2×10^5 cells/well) were infected with wild type HAdV5 (90 particles/cell) and incubated at 37°C for 2 hours. After the incubation, the serum–vector plate and the cell plate were combined and incubated at 37°C for 1 hour. Following the incubation, an equal volume of 20% fetal bovine serum (FBS) with DMEM was added to each well and the combined plate was incubated at 37°C for additional 18 to 22 hours. The next day, the combined plate was washed with PBS and the HEK293 cells were lysed, and the lysate was developed using a mammalian β -galactosidase bioluminescence assay kit per the manufacturer’s instructions. As a control, mouse serum was used instead of serum sample. The resulting luminescence was measured using a SpectraMax® M3 microplate luminometer. The resulting NAb titer was reported as the serum dilution that inhibits transduction of vector by at least 50% compared to the mouse serum.

Statistical Analyses

Average and standard deviation values for concentrations of the Anti-VEGF Fab transgene product in anterior chamber fluid and blood were calculated using Microsoft Office Excel 2010.

RESULTS

Expression of Anti-VEGF Fab transgene product in Anterior Chamber Fluid

The Anti-VEGF Fab transgene product was not expressed in anterior chamber fluid of the animal administered FFB-314. The anti-VEGF Fab transgene product was expressed in anterior chamber fluid collected from the right eye of all animals administered AAV8.aVEGF test vector. No expression was observed in the left eye. No difference in expression of the Anti-VEGF Fab transgene product between males and females was observed.

Expression of Anti-VEGF Fab transgene product in Blood

In some patients administered a single IVT injection of Lucentis, ranibizumab was observed in serum (Xu, Invest Ophthalmol Vis Sci, 54: 1616-24(2013)). To determine if subretinal administration of AAV8.aVEGF test vector results in systemic exposure to the Anti-VEGF Fab transgene product, its concentrations were measured in serum.

Expression of the Anti-VEGF Fab transgene product was below nonspecific background levels in blood of the animal administered FFB-314 and all animals administered the AAV8.aVEGF vector as compared to matched pre-injection level.

Presence of Neutralizing Antibodies Against AAV8 capsid

Baseline levels of NABs against AAV8 capsid were determined in serum from blood samples collected on Study day 0. The limit of detection was a 1:5 dilution; titers of < 5 were considered undetectable.

Treatment	Animal Identification	NAb titer	
		Baseline	Study day 7
FFB-314	C64956	5	40
AAV2/8.aVEGF 1.00 × 10 ¹² GC/eye	C73723	10	320
	C74431	< 5	< 5
	C65027	40	10

5 Abbreviations: GC = genome copies; NAb = neutralizing antibody.

Note: the NAb titer values reported are the reciprocal dilutions of serum at which the relative luminescence units (RLUs) were reduced for 50% compared to control wells (without sample). The limit of detection was 1:5 dilution of sample.

10 Animal administered FFB-314 had pre-existing NABs against AAV8 capsid (see preceding Table). One animal (C74431) administered AAV8.aVEGF test vector did not have detectable NABs to AAV8 capsid. In 2 animals (C73723, C65027) administered AAV8.aVEGF test vector, pre-existing NABs against AAV8 capsid were observed, which persisted on Day 7 (see preceding Table).

15 Toxicity was evaluated based on clinical observations, body weights, indirect ophthalmoscopy, hematology, coagulation, clinical chemistry, and gross pathologic findings. There were no mortalities or unscheduled sacrifices in this study. No adverse clinical observations related to AAV8.aVEGF test vector or the Anti-VEGF Fab transgene product were noted for any animal. Several animals exhibited intermittent transient bouts of diarrhea with no impact to the welfare of the animals because body weights remained stable. No
20 clinically meaningful changes in body weight during the study were observed for any animal. No adverse observations related to AAV8.aVEGF Test vector or the Anti-VEGF Fab transgene product were noted during indirect ophthalmoscopy in any animal. No clinically significant changes in hematology, coagulation, or clinical chemistry parameters were observed in any animal. All clinical pathology parameters were within normal ranges in
25 all animals. There were no gross finding in animals C64956 and C74431. The surfaces of

the right and left kidney of C73723 were pale. There was a focal lesion on the liver in C65027. In conclusion, there were no major toxicology findings.

The test vector in Examples 6-11 is rAAV8.CB7.CI.aVEGFrv3.rBG.

5 EXAMPLE 6 - Expression of AAV2/8.aVEGF Vector in Cynomolgus Monkeys

This study was conducted to assess expression of the anti-VEGF transgene product and to evaluate toxicity, immunogenicity, and effect on normal retinal function of AAV2/8.aVEGF and the anti-VEGF transgene product, and shedding of AAV8.aVEGF in cynomolgus monkeys. The study is ongoing.

10 An AAV2/8.aVEGF described earlier in the examples is used in this study. The vector is diluted in Dulbecco's phosphate-buffered saline (DPBS) with 0.001% Pluronic F-68. As a control article, FFB-314 (DPBS with 0.001% Pluronic F-68) was used. The study is ongoing. The results presented are based on the data collected at Month 3.

15 Macaques were used because they are the closest species to humans for studying retinal diseases. These monkeys and humans have similar eye anatomy, including fovea. The dimensions of the eyes are comparable, which allows determination of the human dose based on relative retinal areas.

Animals in this Example were administered a single dose of 1.00×10^{10} genome copies (GC)/eye of AAV8.aVEGF, or 1.00×10^{12} GC/eye of AAV8.aVEGF, or FFB-314. 20 AAV8.aVEGF and FFB-314 were administered subretinally into the right eye (confirmed visually by appearance of a dome-shaped retinal detachment/retinal bleb under microscope) in a total volume of 100 μ L.

Animals were randomly assigned to 6 sets of 4 animals per set using www.jamestease.co.uk/team-generator. After assigning the sets, 1 of 4 animals from each of 25 the 6 sets was selected using www.randomizer.org/ at random and assigned to groups administered FFB-314 for each given administration date (Groups 2, 4, 6, 8, 10, and 12). The remaining 3 animals were assigned to groups administered 1×10^{12} GC/eye or 1×10^{10} GC/eye AAV8.aVEGF (Groups 1, 3, 5, 7, 9, and 11). Group designation and dose levels for Examples 6 and 7 are presented below.

30

Group Designation and Dose Levels

Group	Treatment ^a	Dose (GC/eye)	Number (#) of animals	Follow-up
1	AAV8.aVEGF test vector	1.00×10^{12}	1 M, 2 F	3 months
2	FFB-314	NA	1 M	
3	AAV8.aVEGF test vector	1.00×10^{10}	2 M, 1 F	
4	FFB-314	NA	1 M	
5	AAV8.aVEGF test vector	1.00×10^{12}	2 M, 2 F ^b	1 year
6	FFB-314	NA	1 F	
7	AAV8.aVEGF test vector	1.00×10^{10}	1 M, 2 F	
8	FFB-314	NA	1 F	
9	AAV8. aVEGF test vector	1.00×10^{12}	2 M, 1F	7 days
10	FFB-314	NA	1 M	
11	AAV8.aVEGF test vector	1.00×10^{10}	2 M, 1 F	
12	FFB-314	NA	1 F	

Abbreviation: F = female; GC = genome copies; M = male; NA = not applicable; No. = number.

^a Test and control articles were administered subretinally into the right eye.

^b One female animal was euthanized during the study because of severe eye infection. The animal was replaced.

Samples of anterior chamber fluid and blood were collected for determination of expression of the Anti-VEGF Fab transgene product Subretinal injections were performed as described in earlier examples.

5 . A. Pharmacology

The results presented are based on the data collected at Month 3. In this report, expression of the anti-VEGF Fab transgene product is described.

1. Methods

10 Animals were administered AAV8.aVEGF test vector or FFB-314 (control article) subretinally. Expression of anti-VEGF transgene product in anterior chamber fluid and blood was determined by enzyme linked immunosorbent assay (ELISA) which was performed as described in previous examples.

2. Pharmacology Results

(a) Expression of Transgene Product in Anterior Chamber Fluid:

The transgene product was not expressed in anterior chamber fluid of any animal administered FFB-314. The transgene product was expressed in anterior chamber fluid of all animals administered AAV8.aVEGF test vector. Onset of expression was rapid, generally within 7 days. Steady-state expression levels were achieved within 1 month. All animals continued to express the transgene product at steady-state levels until the last evaluated timepoint. However, overall expression levels of the anti- transgene product were greater in animals administered 1.00×10^{12} GC/eye of AAV8.aVEGF test vector. No difference in expression of the transgene product between males and females was observed.

(b) Expression of Transgene Product in Blood

In some patients administered a single IVT injection of Lucentis, ranibizumab was observed in serum (Xu, Invest Ophthalmol Vis Sci. 2013 Mar 5,54(3):1616-24). To determine if subretinal administration of an AAV2/8.aVEGF test vector described in these examples results in systemic exposure to the anti-VEGF Fab transgene product, its concentrations were measured in serum. Expression of the anti-VEGF Fab transgene product was below nonspecific background levels in blood of all animals administered AAV8.aVEGF test vector compared to matched pre-injection levels.

3. Conclusion:

Anti-VEGF Fab transgene product is expressed in anterior chamber fluid of all animals administered AAV8.aVEGF test vector.

Anti-VEGF Fab transgene product is not expressed in blood of any animal administered AAV8.aVEGF test vector.

B. Toxicology

In this report, evaluation of toxicity of an AAV2/8.aVEGF test vector is described. Animals were administered AAV8.aVEGF test vector or FFB-314 (control article) subretinally. Toxicity was evaluated based on clinical observations, body weights, ocular pressure, indirect ophthalmoscopy, spectral domain optical coherence tomography, hematology, coagulation, clinical chemistry, and gross pathologic findings, and histopathologic findings.

Ocular pressure was evaluated via rebound tonometry (TonoVet). This method is easy to use and does not require topical anesthesia. Rebound tonometry estimates OP by

using an induction coil to magnetize a small, plastic-tipped metal probe that is launched against the cornea. As the probe rebounds back to the instrument, it creates an induction current from which the OP is calculated. Up to 2 readings were taken, from which an average OP was determined, and accuracy of the results was indicated. Application with the device
5 was performed according to the manufacturer's instructions.

Retinal structure (at a micron-level resolution) was evaluated by *in vivo*, non-invasive, cross-sectional retinal microscopy with SD-OCT (Spectralis OCT, Heidelberg Engineering, Carlsbad, CA). Pupils were dilated with phenylephrine 2.5% and tropicamide 1%. En-face retinal imaging was performed with near infrared (NIR) reflectance (REF) and
10 in a subset of animals with NIR fundus autofluorescence (FAF) using the scanning laser ophthalmoscope of this imaging system. Spectral domain optical coherence tomography scanning was performed with 9 mm long horizontal and vertical cross-sections through the fovea and overlapping 30 × 25 mm raster scans extending into the near midperiphery.

The only adverse AAV8.aVEGF test vector - related finding was significant retinal
15 thinning and loss of photoreceptors observed by spectral domain optical coherence tomography in animals administered 1.00×10^{12} GC/eye of test vector.

C. Electroretinogram (ERG)

In this subpart, assessment of effects of AAV8.aVEGF test vector and the anti-VEGF Fab transgene product on normal retinal function is described. Animals were administered
20 AAV8.aVEGF test vector or FFB-314 (control article) subretinally. Retinal function was evaluated by the full-field electroretinogram (ERG). The full-field ERG is a widely used electrophysiologic test of retinal function. Electroretinogram is a mass electrical potential generated by the retina in response to light stimulus. Usually, it is recorded by an electrode in contact with the corneal surface. Electroretinograms in this study were conducted in
25 accord with the recommendations set by the International Society for Clinical Electrophysiology of Vision (ISCEV; McCulloch, Doc Ophthalmol. 2015 Feb;130(1):1-12. 2015). The results presented are based on the data collected at Month 3. In this report, assessment of effects of AAV8.aVEGF test vector and the anti-VEGF Fab transgene product on normal retinal function is described. Animals were administered AAV8.aVEGF test
30 vector or FFB-314 (control article) subretinally. Retinal function was evaluated by the full-field electroretinogram. In summary, administration of 1.00×10^{10} genome copies (GC)/eye

of AAV8.aVEGF test vector do not impair retinal function. In contrast, administration of 1.00×10^{12} GC/eye of AAV8.aVEGF test vector impairs retinal function.

1. Electroretinogram (ERG) Parameters

An electroretinogram (ERG) generated usually when all retinal cells are active respond to a flash stimulation (a dark-adapted animal, moderate to intense flash). The 2 components are the following:

- a-wave: cornea-negative signal, first after the flash. Origin: photoreceptor photocurrent, the most direct signature of photoreceptor function.
- b-wave: cornea-positive signal following the a-wave generated mostly by on-bipolar cells (second order neurons downstream from photoreceptors).

In this study, the following International Society for Clinical Electrophysiology of Vision (ISCEV) standard and additional protocols were used:

- Dark-adapted rod ERG: Stimulus intensity: 0.01 to 0.02 cd s m^{-2} . Response: b-wave only, no a-wave. Source: rod “on” bipolar cells (second order neurons driven by input from rods). Meaning: a measure of rod function. Designation in data sheets: “Dim flash”.
- Dark-adapted standard flash ERG: Stimulus intensity: 3 cd s m^{-2} . Response: combined rod-cone a- and b-waves; 60% to 70% of the signal being generated by the rod-driven pathway. Source: photoreceptors, both rods and cones (a-wave); higher order neurons driven by both rods and cones. Meaning: a measure of mostly rod function; less sensitive to the state of dark adaptation and less variable than the “dim flash” response. Designation in data sheets: “Standard flash”.
- Dark-adapted bright flash ERG: Stimulus intensity: 10 cd s m^{-2} . Response and meaning: same as for the “standard flash” response, but bright flash response is larger in magnitude and may be less variable. Designation in data sheets: “Bright flash”.
- Light-adapted standard flash cone ERG: Stimulus intensity: 3 cd s m^{-2} , delivered in presence of 30 cd m^{-2} background light after 5 minutes of light adaptation. Response: a- and b-waves generated by cone-driven pathways. Meaning: in presence of background light which completely desensitizes rods the ERG is produced exclusively by cones and cone-driven secondary retinal neurons and is a measure of the cone function. Designation in data sheets: “Standard cone ERG”.
- Light-adapted bright flash cone ERG (in addition to the ISCEV standard): Stimulus intensity: 10 cd s m^{-2} , delivered in presence of 30 cd m^{-2} background light after 5 minutes of

light adaptation. Response and meaning: cone-driven ERG as in case of the “Standard cone ERG”, but of greater magnitude and potentially less variable.

ERG measures (a-wave amplitude, a-wave implicit time, b-wave amplitude, b-wave implicit time) were summarized using mean and standard deviation (SD) for treated eyes and control eye, and for each treatment (FFB-314 (vehicle) groups, AAV8.aVEGF test vector 1.00×10^{10} GC/eye groups, AAV8.aVEGF test vector 1.00×10^{12} GC/eye groups). The paired t-test was used for comparing the ERG measures between AAV8.aVEGF test vector (treated) eye and FFB-314 (control eye), and for comparison between post-injection vs. pre-injection. The two-sample t-test was used for comparing the ERG measures between AAV8.aVEGF test vector 1.00×10^{10} GC/eye groups vs. FFB-314 (vehicle) groups, AAV8.aVEGF test vector 1.00×10^{12} GC/eye groups vs. FFB-314 (vehicle) groups, and AAV8.aVEGF test vector 1.00×10^{12} GC/eye groups vs. AAV8.aVEGF test vector 1.00×10^{10} GC/eye groups. The t-test is appropriate even when the sample size is small [Winter JCF. Using the Student’s t-test with extremely small sample sizes. Practical Assessment, Research and Evaluation. 2013;18 (10). Available online: pareonline.net/getvn.asp?v=18&n=10] 1 or the data are not normally distributed. See, Shuster JJ. Diagnostic for assumptions in moderate to large simple clinical trials: do they really help? Statist. Med. 2005;24:2431-2438; Ganju J. D. Comment on “Diagnostic for assumptions in moderate to large simple clinical trials: do they really help?” Statist. Med. 2006;25:1798-1800.] All the statistical analyses were performed in SAS v9.4 (SAS Institute Inc., Cary, NC), and two-sided p-value ≤ 0.05 is considered as statistically significant.

2. Results

Anti-VEGF Fab transgene product was expressed in all animals administered AAV8.aVEGF test vector (see pharmacology results in Part A of this Example). Retinal function 3 months following administration of AAV8.aVEGF test vector or FFB-314 (post-injection) was compared to retinal function before administration (pre-injection) for treated and untreated eyes. An animal in Group 8 was excluded from data analyses due to an unobtainable ERG following administration of FFB-314.

			FFB-314	Low dose	High dose	p-value		
ERG test	Parameter	Stimulus intensity (cd·s·m ⁻²)	Mean (SD)	Mean (SD)	Mean (SD)	Low dose vs. FFB-314	High dose vs. FFB-314	High dose vs. Low dose
Dark-adapted	a-wave amplitude (uv)	3	68.2 (10.0)	58.4 (17.4)	31.3 (12.4)	0.4	0.003	0.01
		10	113.8 (0.9)	109.8 (25.6)	54.1 (21.0)	0.8	0.002	0.002
Light-adapted	a-wave amplitude (uv)	3	19.9 (4.1)	18.7 (3.5)	9.8 (3.7)	0.65	0.007	0.002
		10	36.1 (7.1)	33.3 (7.4)	19.5 (8.6)	0.61	0.02	0.01

Abbreviations: ERG = electroretinogram; GC = genome copies; SD = standard deviation.

Low dose: 1.00×10^{10} GC/eye of AAV8.aVEGF test vector

High dose: 1.00×10^{12} GC/eye of AAV8.aVEGF test vector.

a. Comparison of Retinal Function Between Treatment Groups

For treated eyes, retinal function post-injection was comparable between animals in low-dose group (1.00×10^{10} GC/eye of AAV8.aVEGF test vector) and FFB-314 group (see preceding Table). For treated eyes, retinal function post-injection in animals in high-dose group (1.00×10^{12} GC/eye of AAV8.aVEGF test vector) was significantly reduced compared to animals in FFB-314 group (see preceding Table). For treated eyes, retinal function post-injection in animals in high-dose group was significantly reduced compared to animals in low-dose group (see preceding Table). For untreated eyes, retinal function post-injection was comparable to pre-injection for all groups.

b. Comparison of Retinal Function Within Treatment Groups

For treated eyes, in animals in low-dose and FFB-314 groups, retinal function post-injection was comparable to matched pre-injection baseline. For treated eyes, in animals in high-dose group, retinal function post-injection was significantly reduced

compared to matched pre-injection baseline. For untreated eye, retinal function post-injection was comparable to matched pre-injection baseline.

E. Virus Shedding

Shedding of AAV8.aVEGF test vector was determined by quantitative PCR analysis targeting transgene-specific sequence in samples of tears, nasal secretion, serum, saliva, urine, and feces. Samples were collected before and after administration of AAV8.aVEGF test vector or FFB-314. AAV8.aVEGF test vector DNA was readily detectable in most samples collected from animals administered AAV8.aVEGF test vector. The presence of AAV8.aVEGF DNA was dose-dependent, transient, and decreased over time.

F. Immunogenicity

In this study, immunogenicity of AAV8.aVEGF test vector and the anti-VEGF Fab transgene product is described. Immunogenicity was assessed by the following:

- The presence of IgM and IgG antibodies against the anti-VEGF Fab transgene product using enzyme linked immunosorbent assay (ELISA);
- The presence of neutralizing antibodies (NAbs) against AAV8 capsid using NAb assay;
- T-cell responses to AAV8.aVEGF test vector and the anti-VEGF Fab transgene product using enzyme linked immunospot (ELISPOT) assay.

Animals were administered AAV8.aVEGF test vector or FFB-314 (control article) subretinally as described earlier in this Example. No sustained IgM, IgG, or T-cell responses to the anti-VEGF Fab transgene product were observed in any animal. Animals administered 1.00×10^{12} GC/eye AAV8.aVEGF test vector developed a higher neutralizing antibody (Nab) response to AAV8 capsid than animals administered 1.00×10^{10} GC/eye AAV8.aVEGF test vector. The NAb response was higher in animals with pre-existing NAbs. Slightly increased T-cell responses against AAV8 capsid were observed in 2 of 6 animals administered 1.00×10^{12} GC/eye test vector.

Results

Anti-VEGF Fab transgene product was expressed in all animals administered AAV8.aVEGF test vector (Example 6). There was no significant IgM against the Anti-VEGF Fab transgene product in serum or in anterior chamber fluid of animals administered FFB-314. IgG against the Anti-VEGF Fab transgene product above baseline level was not observed in animals administered FFB-314.

IgM against the anti-VEGF Fab transgene product was elevated above the baseline level in anterior chamber fluid of 1 animal administered 1.00×10^{10} GC/eye of AAV8.aVEGF test vector. However, as there was no corresponding elevation in the serum, therefore this observation was not clinically significant. IgG against the anti-VEGF Fab transgene product above baseline level was not observed in this treatment group.

IgM against the anti-VEGF Fab transgene product above baseline level was observed in anterior chamber fluid of 1 animal administered 1.00×10^{12} GC/eye of AAV8.aVEGF test vector. However, as there was no corresponding elevation in the serum, this observation was not clinically meaningful. IgG against the anti-VEGF Fab transgene product above baseline level was observed in serum and anterior chamber fluid from another animal and in anterior chamber fluid only of a third animal in this treatment group. However, as neither was preceded by any detectable IgM, these observations were not clinically meaningful. The presence of IgG in these animals was not associated with loss of expression of the anti-VEGF Fab transgene product.

Baseline levels of NABs against AAV8 capsid were determined in serum from blood samples collected on Study day 0. The limit of detection was a 1:5 dilution; titers of < 5 were considered undetectable.

In 4 of 6 animals administered FFB-314, pre-existing NABs were not observed. Two animals that were followed by Study day 90 did not develop NABs. In 2 animals administered FFB-314, pre-existing NABs were observed. The levels of NABs in these 2 animals fluctuated no more than 2 two-fold serial dilutions during the study.

In 2 of 9 animals administered 1.00×10^{10} GC/eye of AAV8.aVEGF test vector, pre-existing NABs were not observed. In 1 animal that was followed by Study day 90, NABs were not observed following administration of AAV8.aVEGF test vector. In animals with pre-existing NABs, their levels increased by no more than 4 two-fold serial dilutions following administration of AAV8.aVEGF test vector.

In 4 of 9 animals administered 1.00×10^{12} GC/eye of AAV8.aVEGF test vector, pre-existing NABs were not observed. Regardless of status of pre-existing NABs, in most animals, an increase in NAb response of up to 9 two-fold serial dilutions was observed following administration of AAV8.aVEGF. This response was sustained through the Study day 90.

T-cell responses to AAV8.aVEGF test vector were observed in 1 animal administered FFB-314 at a single timepoint. In 1 animal, non-specific T cell responses were observed at all timepoints.

5 Sustained T-cell responses to AAV8.aVEGF test vector were not observed in animals administered 1.00×10^{10} GC/eye of test vector.

In 4 of 6 animals administered 1.00×10^{12} GC/eye of AAV8.aVEGF test vector, low-level immune response against AAV8.aVEGF test vector was observed. In 2 of 4 animals with low-level immune response, a sustained (more than 2 consecutive time points) response was observed. Sustained T-cell responses to the anti-VEGF Fab transgene product were not
10 observed in any animal.

No sustained IgM, IgG, or T-cell responses to the Anti-VEGF Fab transgene product were observed in any animal.

Animals administered 1.00×10^{12} GC/eye AAV8.aVEGF test vector developed a higher NAb response to AAV8.aVEGF test vector than animals administered 1.00×10^{10}
15 GC/eye of the same test vector. The NAb response was higher in animals with pre-existing NAb. Slightly increased T-cell responses against this AAV8.aVEGF test vector were observed in 2 of 6 animals administered 1.00×10^{12} GC/eye the AAV8.aVEGF test vector.

20 EXAMPLE 7 - Evaluation of Distribution of AAV2/8 Vector mRNA and Anti-VEGF Fragment Antigen-Binding Following Subretinal Administration of AAV2/8 Vectors in Cynomolgus Monkeys

This study was conducted to evaluate retinal distribution of AAV2/8 vector mRNA and distribution of anti-VEGF Fab throughout the eye following subretinal administration of AAV2/8 vector utilizing tissues from Example 3, Example 5 and Example 6. Levels of
25 mRNA in different parts of retina were assessed by quantitative reverse transcription-polymerase chain reaction and by in situ hybridization. Concentrations of anti-VEGF Fab were determined in retinal sections, anterior chamber fluid and vitreous humor by enzyme-linked immunosorbent assay.

mRNA for AAV2/8 vector is distributed throughout the entire retina following
30 subretinal administration. Similarly, anti-VEGF Fab is distributed throughout the entire retina and is detected in both, vitreous and anterior chamber fluid.

Vector
AAV2/8.UbC.PI.aVEGFv2.SV40
AAV2/8.UbC.PI.aVEGFv3.SV40
AAV2/8.CB7.CI.aVEGFv2.rBG
AAV2/8.CB7.CI.aVEGFv3.rBG

Site of subretinal administration is denoted by a retinal bleb, which can be visualized by SD OCT. In all SD OCT images, retinal blebs are visible.

5 Levels of mRNA for AAV2/8.aVEGF Test Vector in Retina Determined by RT-qPCR

mRNA for AAV8.aVEGF test vector was not detected in the retina of the animal administered FFB-314. mRNA for the AAV8.aVEGF test vector was detected in retinas of all animals administered the AAV8.aVEGF test vector. The highest level of mRNA was detected in the retinal sections that incorporated the site of the subretinal injection. However, mRNA for the AAV8.aVEGF test vector was also detected in sections outside of the injection bleb. mRNA levels in these sections were lower than those in the bleb. The levels were up to 4 logs lower in sections most peripheral to the injection blebs. In sections immediately adjacent to the injection bleb, the levels of mRNA were intermediate.

15 Expression of mRNA for AAV2/8 Vectors in Retina Determined by In Situ Hybridization (ISH)

Expression of mRNA for the AAV2/8 vector determined by ISH was high at the injection site. The transduced cells within retinal layers included RPE cells, photoreceptors, and ganglion cells. Expression of mRNA was lower when moving away from the injection site, disappearing almost completely in the areas most distal to the injection site.

20 Concentrations of Anti-VEGF Fab in Anterior Chamber Fluid, Vitreous, and Retina

Anti-VEGF Fab was expressed in retinas, vitreous, and anterior chamber fluid of eyes of all animals administered AAV2/8 vector (FIGs 6-8) . Expression in the vitreous was 3- to 9-fold higher than in the anterior chamber fluid. With the exception of 1 animal (C65873) in Group 5 (FIG 8), maximal expression in the retinal segments was 1.2- to 3.6-fold higher than in the vitreous. This concentration gradient is likely a reflection of the

mechanism of distribution of anti-VEGF Fab. Anti-VEGF Fab is secreted into vitreous by transduced retina and then diffuses from vitreous to the anterior chamber fluid. Of note, expression of anti-VEGF Fab throughout the retina is more uniform than expression of mRNA.

5 Overall, functional AAV2/8 vector is surprisingly distributed throughout the entire retina following subretinal administration as evidenced by the expression of the vector mRNA by the transduced cells, instead of being limited to the injection bleb. Anti-VEGF Fab is also surprisingly distributed throughout the entire retina including retinal segments that are peripheral to the injection bleb, and is detected in both, vitreous and anterior
10 chamber fluid.

EXAMPLE 8 - Determination Of Affinity For Binding Of Anti-VEGF Transgene Product To Recombinant Human VEGF

This study was conducted to determine affinity for binding of the Anti-VEGF Fab
15 heavy and light chains product to recombinant human VEGF. Binding affinity was determined using Biacore 3000 system, based on surface plasmon resonance (SPR) technique. This technique is based on the plane-polarized light hitting a sensor chip under the conditions of total internal reflection. Interaction between immobilized ligands (e.g., VEGF) and interacting molecules (e.g., Anti-VEGF Fab transgene product) on the sensor chip causes
20 a change in angle of reflectivity of plane-polarized light. This change is immediately detected by sensogram in real time as response units (Daghestani, Theory and applications of surface plasmon resonance, resonant mirror, resonant waveguide grating, and dual polarization interferometry biosensors. Sensors (Basel). 2010;10(11): 9630-46.). The equilibrium binding affinity constant for binding of the Anti-VEGF Fab transgene product is
25 consistent with published range for ranibizumab.

EXAMPLE 9: Tissue Cross-Reactivity Study

The objective of this study was to assess using immunohistochemical techniques, the potential cross-reactivity of the Sponsor supplied antibody Fab fragment aVEGF transgene
30 product with histologically prepared cryo-sections from a selected panel of human tissues.

Anti-VEGF Fab transgene product (1 mg/mL) ("Test product") and ranibizumab (0.97 mg/mL) were used for this study. Natural Human IgG Fab Fragment Protein (the

"Control Article") was supplied at a protein concentration of 14.64 mg/mL. To facilitate immunohistochemical detection the Test transgene product, natural human IgG Fab fragment protein and ranibizumab were conjugated with biotin. The respective protein concentrations were 2.79 mg/mL, 2.88 mg/mL and 2.89 mg/mL. Cryo-sections from the control material and the human tissues for examination were prepared. The assessment of tissue viability indicated that the panel of human tissues was viable. Following slide evaluation of the control titration the following three concentrations of Test transgene product-Biotin: 5, 2.5 and 1.25 $\mu\text{g/mL}$, and the following concentration of ranibizumab-Biotin: 2.5 $\mu\text{g/mL}$, were selected for use in the tissue titration. In the tissue titration no specific positive staining was observed with anti-VEGF transgene product-Biotin or ranibizumab-Biotin in any of the tissues examined. All other observed staining was variable and considered to be non-specific.

Under the conditions of this study, antigen-specific binding of Test transgene product-Biotin and ranibizumab-Biotin was demonstrated in the positive control materials (human glioblastoma and VEGF protein spots). No similar staining was observed with Natural human IgG Fab fragment protein-Biotin or the antibody diluent at the concentrations examined in the tissue titration.

EXAMPLE 10 - Clinical Study

A rAAV8.aVEGF vector was selected for further study which provides the advantage of single sub-retinal administration, thereby reducing the burden of repeated injections. Continued expression of anti-VEGF Fab in NHP for over 6 months and reduction in neovascularization in an animal model of WAMD treated with an rAAV8.aVEGF vector have been demonstrated in pre-clinical studies, and safety of sub-retinal injection is evaluated in non-human primates. The initial clinical study evaluates the safety and transgene expression after a single sub-retinal injection of an rAAV8.aVEGF test vector as described above. Once injected sub-retinally, these vectors are expected to continue to release an anti-VEGF Fab transgene product and block the angiogenic signal thereby protecting the retina from further damage.

Each dosing cohort includes 3 subjects. The first 3 subjects enrolled start with the lowest dose and each group escalates. After each first rAAV8.aVEGF dose, there is a 4 week observation period for safety prior to the next patient being dosed. The primary safety endpoint is at 6 weeks post administration of rAAV8.aVEGF.

Primary endpoints

Ocular and non-ocular safety assessment at 6 weeks, 24 weeks, 6 and 12 months post procedure.

5 Secondary endpoints

- Ocular and non-ocular safety over 106 weeks
 - Mean change from baseline in aqueous rAAV8.aVEGF protein over time
 - Mean change from baseline in BCVA over time
 - Proportion of subjects gaining or losing ≥ 15 letters compared to baseline as per
- 10 BCVA at Week 26, Week 54, and Week 106
- Mean change from baseline in CRT as measured by SD-OCT over time
 - Mean number of ranibizumab rescue injections over time
 - Time to 1st rescue ranibizumab injection
 - Mean change from baseline in CNV and lesion size and leakage area based on FA
- 15 over time
- Immunogenicity measurements (NAb to AAV8, binding antibodies to AAV8, antibodies to aVEGF protein, and Enzyme-Linked ImmunoSpot [ELISpot]).
 - Vector shedding analysis in serum and urine.

Exploratory endpoints:

- 20 • Mean change from baseline over time in area geographic atrophy per fundus autofluorescence (FAF) Incidence of new area of geographic atrophy by FAF (in subjects with no geographic atrophy at baseline)
- Proportion of subjects gaining or losing \geq and ≥ 10 letters, respectively, compared with baseline as per BCVA
- 25 • Proportion of subjects who have a reduction of 50% in rescue injections compared with previous year
- Proportion of subjects with no fluid on SD-OCT

For the present study, patients must have a diagnosis of neovascular age-related macular degeneration (wet AMD) and meet the following criteria.

Inclusion Criteria:

In order to be eligible to participate in this study, a subject must meet all of the following criteria. It is understood that one or more of these criteria may not be required for further studies and for treatment of other populations.

- 5 1. Males or females aged 50 years or above.
2. Sentinel subject for each dose cohort must have a BCVA $\leq 20/100$ and $\geq 20/400$ (≤ 65 and ≥ 35 ETDRS letters) in the study eye.
 - a. Following the sentinel subject evaluation, the rest of the subjects in the dose cohort must have a BCVA between $\leq 20/63$ and $\geq 20/400$ (≤ 75 and ≥ 35 ETDRS letters).
- 10 3. In the case both eyes are eligible, study eye must be the subject's worse-seeing eye, as determined by the Investigator.
4. Must have a documented diagnosis of subfoveal CNV secondary to AMD in the study eye.
 - a. CNV lesion characteristics: lesion size needs to be less than 10 disc areas
- 15 5. Must have received at least 4 intravitreal injections of an anti-VEGF agent for treatment of nAMD in the study eye in the 8 months (or less) prior to Visit 1, with anatomical response documented on SD-OCT.
6. Must have subretinal or intraretinal fluid present at Visit 1 in the study eye,
- 20 evidenced on SD-OCT.
7. Must be pseudophakic (status post cataract surgery) in the study eye.
8. Must be willing and able to comply with all study procedures and be available for the duration of the study.
9. Females of childbearing potential must have a negative urine pregnancy test at the
- 25 screening visit, have negative serum results by Day 8, and be willing to have additional pregnancy tests during the study.
10. Sexually active subjects (both female and male) must be willing to use a medically accepted method of barrier contraception (e.g., condom, diaphragm, or abstinence)
- 30 after this point should be discussed with a responsible physician.
11. Must be willing and able to provide written, signed informed consent.

Exclusion Criteria:

Subjects who meet any of the following exclusion criteria are not eligible to participate in the study. It is understood that future studies and treatment of other patient populations may not include any or all of these criteria.

- 5 1. CNV or macular edema in the study eye secondary to any causes other than AMD.
2. Blood occupying $\geq 50\%$ of the AMD lesion or blood $>1.0 \text{ mm}^2$ underlying the fovea in the study eye.
3. Any condition preventing VA improvement in the study eye, eg, fibrosis,
10 atrophy, or retinal epithelial tear in the center of the fovea.
4. Active or history of retinal detachment in the study eye.
5. Advanced glaucoma in the study eye.
6. Any condition in the study eye that, in the opinion of the Investigator, may
15 increase the risk to the subject, require either medical or surgical intervention during the course of the study to prevent or treat vision loss, or interfere with study procedures or assessments.
7. History of intraocular surgery in the study eye within 12 weeks prior to the screening visit. Yttrium aluminum garnet capsulotomy is permitted if performed
 >10 weeks prior to the screening visit.
- 20 8. History of intravitreal therapy in the study eye, such as intravitreal steroid injection or investigational product, other than anti-VEGF therapy, in the 6 months prior to screening.
9. Presence of an implant in the study eye at screening (excluding intraocular lens).
10. History of malignancy requiring chemotherapy and/or radiation in the 5 years
25 prior to screening. Localized basal cell carcinoma is permitted.
11. Receipt of any investigational product within the 30 days of enrollment or 5 half-lives of the investigational product, whichever is longer.
12. Participation in any other gene therapy study.
13. History of therapy known to have caused retinal toxicity, or concomitant
30 therapy with any drug that may affect visual acuity or with known retinal toxicity, eg, chloroquine or hydroxychloroquine.

14. Ocular or periocular infection in the study eye that may interfere with the surgical procedure.
15. Myocardial infarction, cerebrovascular accident, or transient ischemic attacks within the past 6 months.
- 5 16. Uncontrolled hypertension (systolic blood pressure [BP] >180 mmHg, diastolic BP >100 mmHg) despite maximal medical treatment.
17. Any concomitant treatment that, in the opinion of the Investigator, may interfere with ocular surgical procedure or healing process.
18. Known hypersensitivity to ranibizumab or any of its components or past
10 hypersensitivity (in the Investigator's opinion) to agents like rAAV8.aVEFG test vector.
19. Any serious or unstable medical or psychological condition that, in the opinion of the Investigator, would compromise the subject's safety or successful participation in the study.

Criteria for Continuing Study After Receiving Ranibizumab

15 At Visit 2, subjects are assessed for initial anti-VEGF response to ranibizumab. Subjects undergo both SD-OCT and BCVA, which are compared by the Investigator with the Visit 1 values:

1. Responsive (subjects continue in the study): Response is defined as reduction in CRT >50 microns or >30% improvement in fluid by SD-OCT.
- 20 2. Non-responsive (subjects exit the study as early withdrawals): Non-response is defined as not meeting the criteria above. Additional subjects continue to be enrolled until up to 6 subjects in each cohort receive a single dose of rAAV8.aVEFG test vector.

At this visit central lab results are reviewed. Any subjects with the following values are withdrawn:

- 25 3. Aspartate aminotransferase (AST)/alanine aminotransferase (ALT) >2.5 × upper limit of normal (ULN)
4. Total bilirubin >1.5 × ULN unless the subject has a previously known history of Gilbert's syndrome and a fractionated bilirubin that shows conjugated bilirubin <35% of total bilirubin
- 30 5. Prothrombin time (PT) >1.5 × ULN
6. Hemoglobin <10 g/dL for male subjects and <9 g/dL for female subjects
7. Platelets <100 × 10³/μL

8. Estimated glomerular filtration rate (GFR) <30 mL/min/1.73 m²

In the initial study, ranibizumab (LUCENTIS, Genentech) 0.5 mg is administered by intravitreal injection on Visit 1, 14 days prior to rAAV8.aVEGF test vector subretinal delivery. The rAAV8.aVEGF is given by subretinal administration by a retinal surgeon under local anesthesia. The procedure involves standard 3 port pars plana vitrectomy with a core vitrectomy followed by a single subretinal administration into the subretinal space by a subretinal cannula (36 to 41 gauge). 100 – 150 microliters of rAAV8.aVEGF is delivered. Patients receive one of 3, 4 or 5 doses. Three dose levels: 3×10^9 genome copies (GC)/eye, 1×10^{10} GC/eye, and 6×10^{10} GC/eye. Starting at 4 weeks post-rAAV8.aVEGF test vector administration, the subject may receive intravitreal ranibizumab rescue therapy in the study eye for disease activity if 1 or more of the following rescue criteria apply: Vision loss of ≥ 5 letters (per Best Corrected Visual Acuity [BCVA]) associated with accumulation of retinal fluid on Spectral Domain Optical Coherence Tomography (SD-OCT). Choroidal neovascularization (CNV)-related increased, new, or persistent subretinal or intraretinal fluid on SD-OCT. New ocular hemorrhage.

Further rescue injections may be deferred per the clinician's discretion if one of the following sets of findings occur: Visual acuity is 20/20 or better and central retinal thickness is “normal” as assessed by SD-OCT, or Visual acuity and SD-OCT are stable after 2 consecutive injections. If injections are deferred, they are resumed if visual acuity or SD-OCT get worse per the criteria above.

EXAMPLE 11 - Dose Escalation Study

This Phase I, open-label, multiple-cohort, dose-escalation study is designed to evaluate the safety and tolerability of rAAV8.aVEGF gene therapy in subjects with previously treated neovascular AMD (nAMD). Three doses are studied in approximately 18 subjects. Subjects who meet the inclusion/exclusion criteria and have an anatomic response to an initial anti VEGF injection receive a single dose of rAAV8.aVEGF administered by subretinal delivery. rAAV8.aVEGF uses an AAV8 vector that contains a gene that encodes for a monoclonal antibody fragment which binds to and neutralizes VEGF activity. Safety is the primary focus for the initial 24 weeks after rAAV8.aVEGF administration (primary study period). In certain embodiments, the study includes administering an anti-VEGF antibody, e.g., ranibizumab, and response is measured at week 1 (Visit 2) by SD-OCT. For patients

responsive to this treatment, rAAV8.aVEGF may be administered at Visit 3 (week 2), post-anti-VEGF antibody administration and safety is then assessed through week 26 (24 weeks post-rAAV8.aVEGF administration). Following completion of the primary study period, subjects continue to be assessed until 104 weeks following treatment with rAAV8.aVEGF.

5 Subjects who meet the inclusion/exclusion criteria are enrolled and receive a 0.5 mg intravitreal injection of ranibizumab in the study eye (Visit 1). At Visit 2 (7 days after ranibizumab injection), subjects are evaluated by SD-OCT to confirm anatomic response to the initial anti-VEGF activity associated with the ranibizumab injection compared with their baseline assessment. Subjects who do not have an anatomic response are withdrawn from the study. For withdrawn subjects, anyone who has an AE associated with the ranibizumab injections on Visit 1 is followed until the AE resolves (up to 30 days post-injection). At Visit 10 3 (Week 2), subjects receive a single dose of rAAV8.aVEGF Fab administered in an operating room by subretinal delivery. The sentinel subject in each cohort has vision of $\leq 20/100$ and $\geq 20/400$ (≤ 65 and ≥ 35 ETDRS letters). After rAAV8.aVEGF Fab 15 administration to the sentinel subject, there is a 4-week observation period for safety. Up to 5 additional subjects (with expanded vision criteria of $\leq 20/63$ and $\geq 20/400$ [≤ 75 and ≥ 35 ETDRS letters]) may be enrolled in parallel with a minimum of 1 day between each enrollment. If no safety review triggers (SRTs) are observed, then 4 weeks after the last subject is dosed. Subjects have 3 visits within the first 4 weeks after treatment with 20 rAAV8.aVEGF Fab. Starting 4 weeks after rAAV8.aVEGF Fab administration, subjects may receive intravitreal ranibizumab rescue therapy if they meet predefined rescue injection criteria. Immunogenicity to the vector and transgene of rAAV8.aVEGF Fab is assessed throughout the study.

 Safety is the primary focus for the initial 24 weeks after rAAV8.aVEGF 25 administration (primary study period). Following completion of the primary study period, subjects continue to be assessed until 104 weeks following treatment with rAAV8.aVEGF (Week 106). At the end of the study, subjects are invited to participate in a long-term follow-up study. The safety and tolerability of rAAV8.aVEGF are assessed in each dosed subject and are monitored through assessment of ocular and non-ocular AEs and SAEs, chemistry, 30 hematology, coagulation, urinalysis, immunogenicity, ocular examinations and imaging (BCVA, intraocular pressure, slit lamp biomicroscopy, indirect ophthalmoscopy, and SD-OCT), and vital signs.

A. Arms and Interventions

Arms	Assigned Intervention
Dose 1 3 x 10 ⁹ GC of rAAV8.aVEGF	Biological/Vaccine: rAAV8.aVEGF is a recombinant adeno-associated virus (AAV) gene therapy vector carrying a coding sequence for a soluble anti-VEGF protein
Experimental: Dose 2 1 x 10 ¹⁰ GC of rAAV8.aVEGF	Biological/Vaccine: rAAV8.aVEGF is a recombinant adeno-associated virus (AAV) gene therapy vector carrying a coding sequence for a soluble anti-VEGF protein
Experimental: Dose 3 6 x 10 ¹⁰ GC of rAAV8.aVEGF	Biological/Vaccine: rAAV8.aVEGF is a recombinant adeno-associated virus (AAV) gene therapy vector carrying a coding sequence for a soluble anti-VEGF protein

B. Endpoints:

5

Primary outcome measure:

1. Safety: Incidence of ocular adverse events (AE) and non-ocular serious adverse events (SAE) over 26 weeks

Secondary outcome measure:

10 weeks

2. Safety: Incidence of ocular and non-ocular AEs and SAEs over 106

3. Change in best corrected visual acuity (BCVA) over 106 weeks

106 weeks.

5. Rescue injections: mean number of rescue injections over 106 weeks

15

6. Change in choroidal neovascularization and lesion size and leakage area CNV changes as measured by FA over 106 weeks

Criteria: Inclusion Criteria:

1. Patients ≥ 50 years with a diagnosis of subfoveal CNV secondary to AMD in the study eye receiving prior intravitreal anti-VEGF therapy. Selected patient population is not gender based (males and females included).
- 5 2. BCVA between $\leq 20/100$ and $\geq 20/400$ (≤ 65 and ≥ 35 Early Treatment Diabetic Retinopathy Study [ETDRS] letters) for the first patient in each cohort followed by BCVA between $\leq 20/63$ and $\geq 20/400$ (≤ 75 and ≥ 35 ETDRS letters) for the rest of the cohort.
3. History of need for and response to anti-VEGF therapy.
4. Response to anti-VEGF at trial entry (assessed by SD-OCT at week 1 (Visit
10 2)
5. Must be pseudophakic (status post cataract surgery) in the study eye.
6. Aspartate aminotransferase (AST)/ Alanine aminotransferase (ALT) $< 2.5 \times$ upper limit of normal (ULN); Total Bilirubin (TB) $< 1.5 \times$ ULN; Prothrombin time (PT) $< 1.5 \times$ ULN; Hemoglobin (Hb) > 10 g/dL (males) and > 9 g/dL (females); Platelets $> 100 \times$
15 $10^3/\mu\text{L}$; estimated glomerular filtration rate (eGFR) > 30 mL/min/1.73 m²
7. Must be willing and able to provide written, signed informed consent.

Exclusion Criteria:

1. CNV or macular edema in the study eye secondary to any causes other than AMD.
- 20 2. Any condition preventing visual acuity improvement in the study eye, eg, fibrosis, atrophy, or retinal epithelial tear in the center of the fovea.
3. Active or history of retinal detachment in the study eye.
4. Advanced glaucoma in the study eye.
5. History of intravitreal therapy in the study eye, such as intravitreal steroid
25 injection or investigational product, other than anti-VEGF therapy, in the 6 months prior to screening.
6. Presence of an implant in the study eye at screening (excluding intraocular lens).
7. Myocardial infarction, cerebrovascular accident, or transient ischemic attacks
30 within the past 6 months.

8. Uncontrolled hypertension (systolic blood pressure [BP] >180 mmHg, diastolic BP >100 mmHg) despite maximal medical treatment.

EXAMPLE 12 - Vector Production and Manufacturing

5 A. Description of the Manufacturing Process

Cell Seeding: A qualified human embryonic kidney 293 cell line is used for the production process. Cell culture used for vector production is initiated from a single thawed MCB vial, and expanded per a Master Batch Record Document (MBR). Cells are expanded to $5 \times 10^9 - 5 \times 10^{10}$ cells using Corning T-flasks and CS-10, which allow sufficient cell mass to be generated for seeding up to 50 HS-36 for vector production per BDS lot. Cells are cultivated in medium composed of Dulbecco's Modified Eagle Medium (DMEM), supplemented with 10% gamma irradiated, US-sourced, Fetal Bovine Serum (FBS). The cells are anchorage dependent and cell disassociation is accomplished using TrypLE™ Select, an animal product-free cell dissociation reagent. Cell seeding is accomplished using sterile, single-use disposable bioprocess bags and tubing sets. The cells are maintained at 15 37°C ($\pm 2^\circ\text{C}$), in 5% ($\pm 0.5\%$) CO₂ atmosphere.

Transient Transfection: Following approximately 3 days of growth (DMEM media + 10% FBS), HS-36 cell culture media are replaced with fresh, serum free DMEM media and transfected with the 3 production plasmids using an optimized PEI-based transfection method. All plasmids used in the production process are produced in the context of a CMO quality system and infrastructure utilizing controls to ensure traceability, document control, and materials segregation. 20

Sufficient DNA plasmid transfection complex are prepared in the BSC to transfect 50 HS-36 (per BDS batch). Initially a DNA/PEI mixture is prepared containing 7.5 mg of the relevant vector genome plasmid), 150 mg of pAdDeltaF6(Kan), 75 mg of pAAV2/8Kan AAV helper plasmid and GMP grade PEI (PEIPro, PolyPlus Transfection SA). This plasmid ratio is determined to be optimal for AAV production in small scale optimization studies. After mixing well, the solution is allowed to sit at room temperature for 25 min. and then added to serum-free media to quench the reaction and then added to the HS-36's. The transfection mixture is equalized between all 36 layers of the HS-36 and the cells are 30 incubated at 37°C ($\pm 2^\circ\text{C}$) in a 5% ($\pm 0.5\%$) CO₂ atmosphere for 5 days.

Cell Media Harvesting: Transfected cells and media are harvested from each HS-36 using disposable bioprocess bags by aseptically draining the medium out of the units. Following the harvest of media, the ~ 200 liter volume is supplemented with MgCl₂ to a final concentration of 2 mM (co-factor for Benzonase) and Benzonase nuclease (Cat#: 1.016797.0001, Merck Group) is added to a final concentration of 25 units/mL. The product (in a disposable bioprocess bag) is incubated at 37°C for 2 hr in an incubator to provide sufficient time for enzymatic digestion of residual cellular and plasmid DNA present in the harvest as a result of the transfection procedure. This step is performed to minimize the amount of residual DNA in the final vector DP. After the incubation period, NaCl is added to a final concentration of 500 mM to aid in the recovery of the product during filtration and downstream tangential flow filtration.

Clarification: Cells and cellular debris is removed from the product using a depth filter capsule (1.2/0.22 µm) connected in series as a sterile, closed tubing and bag set that is driven by a peristaltic pump. Clarification assures that downstream filters and chromatography columns are protected from fouling and bioburden reduction filtration ensures that at the end of the filter train, any bioburden potentially introduced during the upstream production process is removed before downstream purification. The harvest material is passed through a Sartorius Sartoguard PES capsule filter (1.2/0.22 µm) (Sartorius Stedim Biotech Inc.).

Large-scale Tangential Flow Filtration: Volume reduction (10-fold) of the clarified product is achieved by Tangential Flow Filtration (TFF) using a custom sterile, closed bioprocessing tubing, bag and membrane set. The principle of TFF is to flow a solution under pressure parallel to a membrane of suitable porosity (100 kDa). The pressure differential drives molecules of smaller size through the membrane and effectively into the waste stream while retaining molecules larger than the membrane pores. By recirculating the solution, the parallel flow sweeps the membrane surface preventing membrane pore fouling. By choosing an appropriate membrane pore size and surface area, a liquid sample may be rapidly reduced in volume while retaining and concentrating the desired molecule. Diafiltration in TFF applications involves addition of a fresh buffer to the recirculating sample at the same rate that liquid is passing through the membrane and to the waste stream. With increasing volumes of diafiltration, increasing amounts of the small molecules are removed from the recirculating sample. This results in a modest purification of the clarified

product, but also achieves buffer exchange compatible with the subsequent affinity column chromatography step. Accordingly, a 100 kDa, PES membrane is used for concentration that is then diafiltered with a minimum of 4 diavolumes of a buffer composed of: 20 mM Tris pH 7.5 and 400 mM NaCl. The diafiltered product is stored overnight at 4°C and then further
5 clarified with a 1.2/0.22 µm depth filter capsule to remove any precipitated material.

Affinity Chromatography: The diafiltered product is applied to a Poros™ Capture Select™ AAV8 affinity resin (Life Technologies) that efficiently captures the AAV8 serotype. Under these ionic conditions, a significant percentage of residual cellular DNA and proteins flow through the column, while AAV particles are efficiently captured. Following
10 application, the column is washed to remove additional feed impurities followed by a low pH step elution (400 mM NaCl, 20 mM Sodium Citrate; pH 2.5) that is immediately neutralized by collection into a 1/10th volume of a neutralization buffer (Bis Tris Propane, 200 mM, pH 10.2).

Anion Exchange Chromatography: To achieve further reduction of in-process
15 impurities including empty AAV particles, the Poros-AAV8 elution pool is diluted 50-fold (20 mM Bis Tris Propane, 0.001% Pluronic F68; pH 10.2) to reduce ionic strength to enable binding to a CIMultus™ QA monolith matrix (BIA Separations). Following a low-salt wash, vector product is eluted using a 60 CV NaCl linear salt gradient (10-180 mM NaCl). This shallow salt gradient effectively separates capsid particles without a vector genome (empty
20 particles) from particles containing vector genome (full particles) and results in a preparation enriched for full capsids. Fractions are collected into tubes containing 1/100th volume of 0.1% pluronic F68 and 1/27th volume of Bis Tris pH 6.3 to minimize non-specific binding to tubes and the length of exposure to high pH respectively. The appropriate peak fraction is collected, and the peak area assessed and compared to previous data for determination of the
25 approximate vector yield.

Final Formulation and Bioburden Reduction Filtration to yield the BDS: TFF is used to achieve final formulation on the pooled AEX fractions with a 100 kDa membrane. This is accomplished by diafiltration of formulation buffer (PBS with NaCl and 0.001% Pluronic or PBS with 0.001% Pluronic to be selected following completion of stability studies) and
30 concentrated to yield the BDS Intermediate at a desired target Samples are removed for BDS Intermediate testing (described in the section below). The BDS Intermediate is stored in

sterile polypropylene tubes and frozen at $\leq -60^{\circ}\text{C}$ in a quarantine location until release for Final Fill. Stability studies are underway to assess stability following storage at $< -60^{\circ}\text{C}$.

Final Fill: The frozen BDS is thawed, pooled, adjusted to the target concentration (dilution or concentrating step via TFF) using the final formulation buffer (PBS with NaCl and 0.001% Pluronic or PBS with 0.001% Pluronic to be selected following completion of stability studies). The product is then terminally filtered through a $0.22\ \mu\text{m}$ filter and filled into either West Pharmaceutical's "Ready-to-Use" (pre-sterilized) glass vials or Crystal Zenith (polymer) vials (vial type pending the outcome of comparability studies) and stoppers with crimp seals at a fill volume $\geq 0.1\ \text{mL}$ to $\leq 0.5\ \text{mL}$ per vial. Vials are individually labeled according to the specifications below. Labeled vials are stored at $\leq -60^{\circ}\text{C}$. All doses require dilution in the formulation buffer prior to administration. The dilution is conducted by the pharmacy at the time of dosing.

B. Assay Methods

Sterility and Bacteriostasis/ Fungistasis: This procedure is performed once according to United States Pharmacopeia (USP) $<71>$, to ensure that the sample matrix does not cause inhibition of the assay. Included in the test is the suitability test.

Particle Aggregation: Drug product particle aggregation is assessed using a dynamic light scattering (DLS) assay. DLS measures fluctuations in scattered light intensity due to diffusing particles and is used to characterize the size of various particles in the sample. DLS instrument software typically displays the particle population at different diameters. If the system is monodisperse, only one population is detected and the mean effective diameter of the particles can be determined. In a polydisperse system, such as in the case of aggregation, multiple particle populations are detected and sized using CONTIN analysis.

Residual plasmid DNA: Detection of plasmid DNA sequences is accomplished using qPCR and primer probe sets specific for the kanamycin gene present in the plasmid backbone but not in vector genomes. The assay is performed in both the presence and absence of DNase digestion such that the amount of free plasmid and the amount packaged into vector particles can be determined.

E1 DNA: Adenoviral E1 DNA is a host cell contaminant and is detected by qPCR specific for the gene. The assay is performed in both the presence and absence of DNase digestion such that both free and packaged E1 DNA can be quantified.

Residual Host Cell DNA: Levels of residual host cell DNA (HCDNA) are quantified

using qPCR directed against the human 18s rDNA gene which is a high copy number DNA sequence and thus confers sensitivity. In addition to total residual HCDNA levels, the amount of DNA at various size ranges is also determined.

5 Residual Host Cell Protein: Residual 293 host cell protein (HCP) is detected using commercially available ELISA kits such as that sold by Cygnus Technologies.

Poros-AAV8 Leachable Ligand: An Enzyme-Linked Immunosorbent Assay (ELISA) kit supplied by Life Technologies, the maker of the Poros-AAV8 resin, is used to detect leached camelid antibody in the drug product.

Mycoplasma Detection: Mycoplasma testing is performed according to USP <63>.

10 Bioburden Testing: This test is performed according to USP <61>.

Endotoxin Testing: This assay is performed according to USP <85>.

In vitro Assay for Adventitious Agents: The purpose of the *in vitro* assay for viral contaminants is to detect possible adventitious viruses introduced during AAV8.AMD vector production and is based upon CBER's 1993 Points to Consider and ICH Q5A. The *in vitro* 15 assays use 3 indicator cell lines - human diploid lung (MRC-5) cells, African green monkey kidney (Vero) cells, and human foreskin fibroblast (Hs68) cells. Assay endpoints are observation of cytopathic effects (CPE) over a course of at least 28 days as well as hemadsorption at the end of the assay period, which facilitates the detection of a broad range of viruses.

20 Vector Genome Identity: DNA Sequencing: Viral Vector genomic DNA is isolated and the sequence determined by 2-fold sequencing coverage using primer walking. Sequence alignment is performed and compared to the expected sequence.

Vector Capsid Identity: AAV Capsid Mass spectrometry of VP1: Confirmation of the AAV2/8 serotype of the drug product is achieved by an assay based upon analysis of 25 peptides of the AAV capsid protein.

Genomic Copy (GC) Titer: A droplet digital PCR (ddPCR)-based technique for determining the genome copy (GC) titer for AAV vectors is described in Lock et al. Human Gene Therapy Methods 25:115–125. The assay utilized involves digestion with DNase I, followed by digital PCR analysis to measure encapsulated vector genomic copies. DNA 30 detection is accomplished using sequence specific primers targeting the RBG polyA region in combination with a fluorescently tagged probe hybridizing to this same region. A number

of standards, validation samples and controls (for background and DNA contamination) have been introduced into the assay.

Empty to Full Particle Ratio: The total particle content of the drug product is determined by SDS-PAGE analysis. A reference vector preparation purified on an iodixanol gradient is analyzed by various methods (analytical ultracentrifugation, electron microscopy and absorbance at 260/280 nm) to established percentage of full particles in the preparation. This reference material is serially diluted to known genome copy numbers (and thus by extension, particle numbers) and each dilution is run on an SDS PAGE gel along with a similar dilution series of the drug product. Peak area volumes of both the reference material and drug product VP3 protein bands are determined by densitometry and the reference material volumes are plotted versus particle number. The total particle concentration of the drug product is determined by extrapolation from this curve and the genome copy (GC) titer is then subtracted to obtain the empty particle titer. The empty to full particle ratio is the ratio of the empty particle titer to the GC titer.

Infectious Titer: The infectious unit (IU) assay is used to determine the productive uptake and replication of AAV8.AMD vector in RC32 cells (rep2 expressing HeLa cells). A 96-well end-point format has been employed similar to that previously published. Briefly, RC32 cells are co-infected by serial dilutions of AAV8.AMD BDS and a uniform dilution of Ad5 with 12 replicates at each dilution of rAAV. Seventy-two hours after infection the cells are lysed, and qPCR performed to detect rAAV vector amplification over input. An end-point dilution Tissue Culture Infectious Dose 50% (TCID₅₀) calculation (Spearman-Kärber) is performed to determine a replicative titer expressed as IU/mL. Since “infectivity” values are dependent on particles coming into contact with cells, receptor binding, internalization, transport to the nucleus and genome replication, they are influenced by assay geometry and the presence of appropriate receptors and post-binding pathways in the cell line used. Receptors and post-binding pathways are not usually maintained in immortalized cell lines and thus infectivity assay titers are not an absolute measure of the number of “infectious” particles present. However, the ratio of encapsidated GC to “infectious units” (described as GC/IU ratio) can be used as a measure of product consistency from lot to lot.

Host Cell DNA: A qPCR assay is used to detect residual human 293 DNA. After spiking with a “non-relevant DNA”, total DNA (non-relevant, vector and residual genomic) is extracted from ~1 mL of product. The Host Cell DNA is quantified using qPCR targeting

the 18S rDNA gene. The quantities of DNA detected are normalized based on the recovery of the spiked non-relevant DNA.

Host Cell Protein: An ELISA is performed to measure levels of contaminating host HEK293 cell proteins. The Cygnus Technologies HEK293 Host Cell Proteins 2nd Generation ELISA kit is used according to instructions.

Replication-competent AAV (rcAAV) Assay: A sample is analyzed for the presence of replication competent AAV2/8 (rcAAV) that can potentially arise during the production process.

An example of this type of assay is shown in (FIGs 10A-10D), where wtAAV8 is spiked into different GC amounts of AAV8 vector and the cap gene copy number per 1 µg of 293 cell DNA is determined after 3 successive passages of the cell lysate onto fresh cells. The details of the assay development are included in the CTA submission. These results indicate that the minimum detectable amount of wtAAV8 using this assay is 10⁴ GC. This number is equivalent to approximately 1 TCID₅₀ IU and reflects the lack of infectivity of AAV8 for 293 cells as evidenced by the high GC:IU ratios obtained compared to AAV2. The low sensitivity appears unavoidable with the current assay system but might be overcome in future by engineering a cell line with a yet to be discovered AAV8 cellular receptor or other protein important in post-entry pathways. Spiking the wtAAV8 into AAV8 vector concentrations of up to 10¹¹ GC had little effect on detection and indicates a lack of interference of the vector on wtAAV8 replication at this vector level. While wildtype AAV has been used extensively as a surrogate in the past for rcAAV2 and in our own rcAAV assay development efforts for AAV8, the best surrogate is a AAV8 capsid containing AAV2 ITRs, an AAV2 rep gene and an AAV8 cap gene.

Sample Steps	Test	Analytical Method	Acceptance Criteria ¹
Final Drug Product in vials	Appearance	Visual Inspection	Clear to slightly Opaque, colorless to faint white solution, free of non-product related foreign particulates
	pH	USP<791>	7.4 +/-0.2

Sample Steps	Test	Analytical Method	Acceptance Criteria ¹
	GC Titer	ddPCR	$\geq 1 \times 10^{11}$ GC/mL**
	AAV Vector Genome Identity	Sequencing(Sanger)	Confirm expected sequence
Final Drug Product in vials	Total Protein Content	Micro BCA	Report result
	Osmolality Content	USP<785>	< 400 mOsm
	Empty/Full Particle ratio Purity	PT(by SDS-PAGE)/GC Ratio	Report result
	Empty: Full particle ratio Purity	OD260/280	Report result
	Viral Capsid Purity	SDS-PAGE	Report result
	Aggregation Characterization Purity	Dynamic Light Scattering	Report result
	In vitro potency Potency	HEK293transduction/ ranibizumab ELISA	Conforms to reference standard
	Transgene expression Characterization Identity	In vitro expression and ELISA	Positive for ranibizumab
	Infectious Titer Characterization Potency	TCID50/qPCR	Report results

Sample Steps	Test	Analytical Method	Acceptance Criteria ¹
	rcAAV by triple passage HEK293 + Ad5 Characterization Safety	Cell Culture/qPCR	Report results
	Capsid Identity Characterization	UPLC/Mass Spectrometry	AAV-8 signature peptide detected. Signature peptides for AAV1,2,6,9 hu37 and Rh10 not detected
	Endotoxin Safety	USP<85> Kinetic Chromogenic	<0.80 EU/mg* or (≤ the safety limit based on dose calculations, pending the verification of total protein)
	Sterility Safety	USP<71>	No Growth
	Container Closure Integrity (for stability study only and not for lot release)	Dye Ingress Test	Container is Integral

¹The acceptance criteria is determined upon completion of the first GMP campaign.

*Endotoxin limit calculation is based on dose in Mass. Once the total protein concentration is confirmed from the GMP run, the limit can be recalculated. The current limit is based on the protein concentration from the Tox materials in relation to GC titer. The value is an approximation and not a definitive value. The dual acceptance criteria presented here.

** DP GC Titer criteria may change depending on the final selected dose levels for the study

The clinically suitable surfactant Pluronic F68 is added to the final formulation buffer of AAV8.AMD and is anticipated to minimize this type of loss. The interaction of the drug product with both the storage vial and the clinical delivery device is investigated to determine the amount of vector loss through binding to surfaces. GC titers (oqPCR) of the engineering run drug product are determined before and after vialling and storage at $\leq -60^{\circ}\text{C}$. For the delivery device, the DP is thawed, diluted in the appropriate clinical diluent to the correct dosing concentration and passed through the device. GC titrations are performed on the DP directly after thaw, after dilution and after passage through the device, and the appropriate number of replicates is included to assure statistical significance. Comparison of GC titers in this manner enables an assessment of DP loss during storage and administration to the patient. Parallel studies are also performed in a similar way to assess the activity of the drug product after passing through the delivery device. For this purpose the *in vitro* ranibizumab expression-based potency assay is employed.

15 (Sequence Listing Free Text)

The following information is provided for sequences containing free text under numeric identifier <223>.

SEQ ID NO: (containing free text)	Free text under <223>
1	<223> Humanized anti-VEGF Fab heavy chain <220> <221> MISC_FEATURE <222> (28)..(39) <223> complementarity determining region <220> <221> MISC_FEATURE <222> (54)..(83) <223> complementarity determining region
2	<223> Humanized anti-VEGF Fab <220> <221> MISC_FEATURE

	<p><222> (26)..(37) <223> complementarity determining region <220> <221> MISC_FEATURE <222> (107)..(117) <223> complementarity determining region</p>
<p>3</p>	<p><223> 5'ITR.CB7.CI.aVEGFv2.rBG.3'ITR <220> <221> repeat_region <222> (1)..(130) <223> 5' ITR <220> <221> promoter <222> (198)..(579) <223> CMV IE promoter <220> <221> enhancer <222> (279)..(538) <223> C4 enhancer with 2 mismatches <220> <221> misc_feature <222> (279)..(538) <223> C4 enhancer <220> <221> promoter <222> (582)..(862)</p>

	<p><223> CB promoter</p> <p><220></p> <p><221> TATA_signal</p> <p><222> (836)..(839)</p> <p><220></p> <p><221> misc_feature</p> <p><222> (955)..(1829)</p> <p><223> chicken beta-actin intron</p> <p><220></p> <p><221> Intron</p> <p><222> (956)..(1928)</p> <p><223> chicken beta-actin intron</p> <p><220></p> <p><221> Intron</p> <p><222> (956)..(1928)</p> <p><223> chicken beta-actin promoter</p> <p><220></p> <p><221> 5'UTR</p> <p><222> (1946)..(1993)</p> <p><223> c-myc 5' UTR</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1994)..(1999)</p> <p><223> kozak sequence</p> <p><220></p>
--	---

	<p><221> transit_peptide <222> (1999)..(2058)</p> <p><220> <221> CDS <222> (2059)..(2427) <223> aVEGFv2 VH</p> <p><220> <221> CDS <222> (2428)..(2748) <223> CH1</p> <p><220> <221> misc_feature <222> (2752)..(2763) <223> Furin cleavage site</p> <p><220> <221> CDS <222> (2764)..(2835) <223> F2A linker</p> <p><220> <221> transit_peptide <222> (2836)..(2895)</p> <p><220> <221> CDS <222> (2896)..(3216) <223> aVEGFv2 VL</p>
--	---

	<p><220> <221> CDS <222> (3217)..(3537) <223> CL</p> <p><220> <221> polyA_signal <222> (3613)..(3739) <223> rabbit globin polyA</p> <p><220> <221> repeat_region <222> (3828)..(3957) <223> 3' ITR</p>
4	<223> Synthetic Construct
5	<223> Synthetic Construct
6	<223> Synthetic Construct
7	<223> Synthetic Construct
8	<223> Synthetic Construct
9	<p><213> Artificial Sequence</p> <p><220> <223> AAV25'ITR.UbC.Ci.aVEGFv2.rBG.AAV23'ITR</p> <p><220> <221> repeat_region <222> (17)..(146) <223> 5' ITR</p> <p><220> <221> promoter</p>

	<p><222> (207)..(1435)</p> <p><223> UbC with C insertion at 289 and G insertion at 990</p> <p><220></p> <p><221> Intron</p> <p><222> (1529)..(1661)</p> <p><223> chimeric intron</p> <p><220></p> <p><221> 5'UTR</p> <p><222> (1736)..(1783)</p> <p><223> c-myc 5'UTR</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1784)..(1789)</p> <p><223> kozak</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1789)..(1848)</p> <p><223> kozak</p> <p><220></p> <p><221> CDS</p> <p><222> (1849)..(2217)</p> <p><223> aVEGFv2 VH</p> <p><220></p> <p><221> CDS</p> <p><222> (2218)..(2538)</p>
--	---

	<p><223> CH1</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2542)..(2553)</p> <p><223> furin cleavage site</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2554)..(2625)</p> <p><223> F2a linker</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2626)..(2685)</p> <p><223> Leader</p> <p><220></p> <p><221> CDS</p> <p><222> (2686)..(3006)</p> <p><223> aVEGFv2 VL</p> <p><220></p> <p><221> CDS</p> <p><222> (3007)..(3327)</p> <p><223> CL</p> <p><220></p> <p><221> polyA_signal</p> <p><222> (3345)..(3576)</p> <p><223> SV40 late polyadenylation signal</p>
--	---

	<p><220> <221> repeat_region <222> (3641)..(3770) <223> 3' ITR</p>
10	<223> Synthetic Construct
11	<223> Synthetic Construct
12	<223> Synthetic Construct
13	<223> Synthetic Construct
14	<p><223> ITR.CB7.CI.aVEGRv3.rBG.ITR</p> <p><220> <221> repeat_region <222> (1)..(130) <223> 5'ITR</p> <p><220> <221> promoter <222> (198)..(579) <223> CMV IE promoter</p> <p><220> <221> enhancer <222> (279)..(538) <223> C4 enhancer with 2 mismatches</p> <p><220> <221> promoter <222> (582)..(862) <223> CB promoter</p> <p><220></p>

	<p><221> TATA_signal <222> (836)..(839)</p> <p><220></p> <p><221> Intron <222> (956)..(1928) <223> chicken beta-actin intron</p> <p><220></p> <p><221> 5'UTR <222> (1940)..(1987)</p> <p><220></p> <p><221> misc_feature <222> (1988)..(1993) <223> Kozak</p> <p><220></p> <p><221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220></p> <p><221> CDS <222> (2053)..(2421) <223> aVEGFv3 VH</p> <p><220></p> <p><221> CDS <222> (2422)..(2742) <223> CH1</p>
--	---

	<p><220> <221> misc_feature <222> (2746)..(2757) <223> furin cleavage site</p> <p><220> <221> misc_feature <222> (2758)..(2829) <223> F2a linker</p> <p><220> <221> misc_feature <222> (2830)..(2889) <223> leader</p> <p><220> <221> CDS <222> (2890)..(3210) <223> aVEGFv3 VL</p> <p><220> <221> CDS <222> (3211)..(3531) <223> CL</p> <p><220> <221> polyA_signal <222> (3607)..(3733) <223> rabbit globin polyA</p> <p><220> <221> repeat_region</p>
--	--

	<222> (3822)..(3951) <223> 3' ITR
15	<223> Synthetic Construct
16	<223> Synthetic Construct
17	<223> Synthetic Construct
18	<223> Synthetic Construct
19	<223> ITR.UbC.PI.aVEGFv3.SV40.ITR <220> <221> repeat_region <222> (17)..(146) <223> AAV2 5' ITR <220> <221> promoter <222> (207)..(1434) <223> UbC, with C insert at 289 and G insert at 990 <220> <221> Intron <222> (1528)..(1660) <223> chimeric intron <220> <221> 5'UTR <222> (1729)..(1776) <223> c-myc 5'UTR <220> <221> misc_feature

	<p><222> (1777)..(1782) <223> kozak</p> <p><220> <221> misc_feature <222> (1782)..(1841) <223> leader</p> <p><220> <221> CDS <222> (1842)..(2210) <223> aVEGFv3 VH</p> <p><220> <221> CDS <222> (2211)..(2531) <223> CH1</p> <p><220> <221> misc_feature <222> (2535)..(2546) <223> furin cleavage site</p> <p><220> <221> misc_feature <222> (2547)..(2618) <223> F2a linker</p> <p><220> <221> misc_feature <222> (2619)..(2678) <223> leader</p>
--	---

	<p><220> <221> CDS <222> (2679)..(2999) <223> aVEGFv3 VL</p> <p><220> <221> CDS <222> (3000)..(3320) <223> CL</p> <p><220> <221> polyA_signal <222> (3338)..(3569) <223> SV40 late polyA</p> <p><220> <221> repeat_region <222> (3634)..(3763) <223> AAV2 3'ITR</p>
20	<223> Synthetic Construct
21	<223> Synthetic Construct
22	<223> Synthetic Construct
23	<223> Synthetic Construct
24	<p><223> ITR.UbC.PI.aVEGFv1.SV40.ITR</p> <p><220> <221> repeat_region <222> (17)..(146) <223> ITR</p>

	<p><220> <221> promoter <222> (207)..(1435) <223> UbC</p> <p><220> <221> Intron <222> (1529)..(1661) <223> Promoga chimeric intron</p> <p><220> <221> 5'UTR <222> (1730)..(1777) <223> c-myc 5'UTR</p> <p><220> <221> misc_feature <222> (1778)..(1783) <223> kozak</p> <p><220> <221> misc_feature <222> (1783)..(1842) <223> leader</p> <p><220> <221> CDS <222> (1843)..(2211) <223> aVEGFv1 VH</p> <p><220> <221> CDS</p>
--	---

	<p><222> (2212)..(2532) <223> CH1</p> <p><220> <221> misc_feature <222> (2536)..(2547) <223> furin cleavage site</p> <p><220> <221> misc_feature <222> (2548)..(2619) <223> F2A linker</p> <p><220> <221> misc_feature <222> (2620)..(2679) <223> leader</p> <p><220> <221> CDS <222> (2680)..(3000) <223> aVEGFv1 VL</p> <p><220> <221> CDS <222> (3001)..(3321) <223> CL</p> <p><220> <221> polyA_signal <222> (3339)..(3570) <223> SV40 polyadenylation signal</p>
--	---

	<220> <221> repeat_region <222> (3635)..(3764) <223> ITR
25	<223> Synthetic Construct
26	<223> Synthetic Construct
27	<223> Synthetic Construct
28	<223> Synthetic Construct
30	<223> synthetic leader
31	<223> synthetic leader 2
32	<223> derived from encephalomyocarditts virus
33	<223> aVEGF <220> <221> MISC_FEATURE <222> (1)..(20) <223> leader <220> <221> MISC_FEATURE <222> (21)..(252) <223> aVEGF Heavy Chain <220> <221> MISC_FEATURE <222> (280)..(299) <223> leader <220> <221> MISC_FEATURE

	<p><222> (300)..(513) <223> aVEGF Light Chain</p>
<p>34</p>	<p><223> ITR.CB7.CI.aVEGFv1.rBG.ITR</p> <p><220> <221> repeat_region <222> (1)..(130) <223> 5' ITR</p> <p><220> <221> promoter <222> (204)..(584) <223> CMV IE promoter</p> <p><220> <221> promoter <222> (585)..(862) <223> CB promoter</p> <p><220> <221> TATA_signal <222> (836)..(839)</p> <p><220> <221> Intron <222> (956)..(1928) <223> chicken beta-actin intron</p> <p><220> <221> 5'UTR <222> (1940)..(1987) <223> c-myc 5'UTR</p>

	<p><220> <221> misc_feature <222> (1988)..(1993) <223> kozak</p> <p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220> <221> misc_feature <222> (2053)..(2421) <223> aVEGFv1 VH</p> <p><220> <221> misc_feature <222> (2422)..(2742) <223> CH1</p> <p><220> <221> misc_feature <222> (2746)..(2757) <223> furin cleavage site</p> <p><220> <221> misc_feature <222> (2758)..(2829) <223> F2A linker</p> <p><220></p>
--	---

	<p><221> misc_feature <222> (2830)..(2889) <223> leader</p> <p><220> <221> misc_feature <222> (2890)..(3210) <223> aVEGFv1 VL</p> <p><220> <221> misc_feature <222> (3211)..(3531) <223> CL</p> <p><220> <221> misc_feature <222> (3532)..(3537) <223> stop cassette</p> <p><220> <221> polyA_signal <222> (3607)..(3733) <223> Rabbit globin poly A</p> <p><220> <221> misc_feature <222> (3785)..(3821) <223> part of AAV</p> <p><220> <221> repeat_region <222> (3822)..(3951)</p>
--	---

	<p><223> 3'ITR</p>
<p>35</p>	<p><223> ITR.CB7.CI.aVEGFv4.rBG.ITR</p> <p><220></p> <p><221> repeat_region</p> <p><222> (1)..(130)</p> <p><223> 5'ITR</p> <p><220></p> <p><221> promoter</p> <p><222> (198)..(579)</p> <p><223> CMV IE promoter</p> <p><220></p> <p><221> enhancer</p> <p><222> (279)..(538)</p> <p><223> C4 enhancer</p> <p><220></p> <p><221> promoter</p> <p><222> (582)..(862)</p> <p><223> CB promoter</p> <p><220></p> <p><221> Intron</p> <p><222> (956)..(1928)</p> <p><223> chicken beta-actin intron</p> <p><220></p> <p><221> 5'UTR</p> <p><222> (1940)..(1987)</p> <p><223> c-myc 5'UTR</p>

	<p><220> <221> misc_feature <222> (1988)..(1993) <223> kozak</p> <p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220> <221> misc_feature <222> (2053)..(2421) <223> aVEGFv4 VH</p> <p><220> <221> misc_feature <222> (2422)..(2742) <223> CH1</p> <p><220> <221> misc_feature <222> (2746)..(2757) <223> furin cleavage site</p> <p><220> <221> misc_feature <222> (2758)..(2829) <223> F2A linker</p> <p><220> <221> misc_feature</p>
--	--

	<p><222> (2830)..(2889) <223> leader</p> <p><220> <221> misc_feature <222> (2890)..(3210) <223> aVEGFv4 VL</p> <p><220> <221> misc_feature <222> (3211)..(3531) <223> CL</p> <p><220> <221> misc_feature <222> (3532)..(3537) <223> stop cassette</p> <p><220> <221> polyA_signal <222> (3607)..(3733) <223> Rabbit globin poly A</p> <p><220> <221> repeat_region <222> (3822)..(3951) <223> 3'ITR</p>
36	<p><223> ITR.CB7.CI.aVEGFv5.rBG.ITR</p> <p><220> <221> repeat_region</p>

	<p><222> (1)..(130) <223> 5'ITR</p> <p><220> <221> promoter <222> (198)..(579) <223> CMV IE promoter</p> <p><220> <221> enhancer <222> (279)..(538) <223> C4 enhancer</p> <p><220> <221> promoter <222> (582)..(862) <223> CB promoter</p> <p><220> <221> Intron <222> (956)..(1928) <223> chicken beta-actin intron</p> <p><220> <221> 5'UTR <222> (1940)..(1987) <223> c-myc 5'UTR</p> <p><220> <221> misc_feature <222> (1988)..(1993) <223> kozak</p>
--	---

	<p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p>
	<p><220> <221> misc_feature <222> (2053)..(2421) <223> aVEGFv5 VH</p>
	<p><220> <221> misc_feature <222> (2422)..(2742) <223> CH1</p>
	<p><220> <221> misc_feature <222> (2746)..(2757) <223> furin cleavage site</p>
	<p><220> <221> misc_feature <222> (2758)..(2829) <223> F2A linker</p>
	<p><220> <221> misc_feature <222> (2830)..(2889) <223> leader</p>
	<p><220></p>

	<p><221> misc_feature <222> (2890)..(3210) <223> aVEGFv5 VL</p> <p><220> <221> misc_feature <222> (3211)..(3531) <223> CL</p> <p><220> <221> misc_feature <222> (3532)..(3537) <223> stop cassette</p> <p><220> <221> polyA_signal <222> (3607)..(3733) <223> rabbit globin poly A</p> <p><220> <221> repeat_region <222> (3822)..(3951) <223> 3'ITR</p>
37	<p><223> ITR.CB7.CI.aVEGFv6.rBG.ITR</p> <p><220> <221> repeat_region <222> (1)..(130) <223> 5'ITR</p> <p><220></p>

	<p><221> promoter <222> (198)..(579) <223> CMV IE promoter</p> <p><220> <221> enhancer <222> (279)..(538) <223> C4 enhancer</p> <p><220> <221> promoter <222> (582)..(862) <223> CB promoter</p> <p><220> <221> Intron <222> (956)..(1928) <223> chicken beta-actin intron</p> <p><220> <221> 5'UTR <222> (1940)..(1987) <223> c-myc 5'UTR</p> <p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220> <221> misc_feature <222> (2053)..(2421)</p>
--	---

	<p><223> aVEGFv6 VH</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2422)..(2742)</p> <p><223> CH1</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2746)..(2757)</p> <p><223> furin cleavage site</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2758)..(2829)</p> <p><223> F2A linker</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2830)..(2889)</p> <p><223> leader</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2890)..(3210)</p> <p><223> aVEGFv6 VL</p> <p><220></p> <p><221> misc_feature</p> <p><222> (3211)..(3531)</p> <p><223> CL</p>
--	---

	<p><220></p> <p><221> misc_feature</p> <p><222> (3532)..(3537)</p> <p><223> stop cassette</p> <p><220></p> <p><221> polyA_signal</p> <p><222> (3607)..(3733)</p> <p><223> Rabbit globin poly A</p> <p><220></p> <p><221> repeat_region</p> <p><222> (3822)..(3951)</p> <p><223> 3'ITR</p>
<p>38</p>	<p><223> ITR.CB7.CI.aVEGFv7.rBG.ITR</p> <p><220></p> <p><221> repeat_region</p> <p><222> (1)..(130)</p> <p><223> 5'ITR</p> <p><220></p> <p><221> promoter</p> <p><222> (198)..(579)</p> <p><223> CMV IE promoter</p> <p><220></p> <p><221> enhancer</p> <p><222> (279)..(538)</p> <p><223> C4 enhancer</p>

	<p><220> <221> promoter <222> (582)..(862) <223> CB promoter</p> <p><220> <221> Intron <222> (956)..(1928) <223> chicken beta-actin intron</p> <p><220> <221> 5'UTR <222> (1940)..(1987) <223> c-myc 5'UTR</p> <p><220> <221> misc_feature <222> (1988)..(1993) <223> kozak</p> <p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220> <221> misc_feature <222> (2053)..(2421) <223> aVEGFv7 VH</p> <p><220> <221> misc_feature</p>
--	---

	<p><222> (2422)..(2742) <223> CH1</p> <p><220> <221> misc_feature <222> (2746)..(2757) <223> furin cleavage site</p> <p><220> <221> misc_feature <222> (2758)..(2829) <223> F2A linker</p> <p><220> <221> misc_feature <222> (2830)..(2889) <223> leader</p> <p><220> <221> misc_feature <222> (2890)..(3210) <223> aVEGFv7 VL</p> <p><220> <221> misc_feature <222> (3211)..(3531) <223> CL</p> <p><220> <221> misc_feature <222> (3532)..(3537) <223> stop cassette</p>
--	---

	<p><220></p> <p><221> polyA_signal</p> <p><222> (3607)..(3733)</p> <p><223> rabbit globin poly A</p> <p><220></p> <p><221> repeat_region</p> <p><222> (3822)..(3951)</p> <p><223> 3'ITR</p>
<p>39</p>	<p><223> ITR.CB7.CI.aVEGFv8.rBG.ITR</p> <p><220></p> <p><221> repeat_region</p> <p><222> (1)..(130)</p> <p><223> 5'ITR</p> <p><220></p> <p><221> promoter</p> <p><222> (198)..(579)</p> <p><223> CMV IE promoter</p> <p><220></p> <p><221> enhancer</p> <p><222> (279)..(538)</p> <p><223> C4 enhancer</p> <p><220></p> <p><221> promoter</p> <p><222> (582)..(862)</p> <p><223> CB promoter</p>

	<p><220> <221> Intron <222> (956)..(1928) <223> chicken beta-actin intron</p> <p><220> <221> 5'UTR <222> (1940)..(1987) <223> c-myc 5'UTR</p> <p><220> <221> misc_feature <222> (1988)..(1993) <223> kozak</p> <p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220> <221> misc_feature <222> (2053)..(2421) <223> aVEGFv8 VH</p> <p><220> <221> misc_feature <222> (2422)..(2742) <223> CH1</p> <p><220></p>
--	--

	<p><221> misc_feature <222> (2746)..(2757) <223> furin cleavage site</p> <p><220> <221> misc_feature <222> (2758)..(2829) <223> F2A linker</p> <p><220> <221> misc_feature <222> (2830)..(2889) <223> leader</p> <p><220> <221> misc_feature <222> (2890)..(3210) <223> aVEGFv8 VL</p> <p><220> <221> misc_feature <222> (3211)..(3531) <223> CL</p> <p><220> <221> misc_feature <222> (3532)..(3537) <223> stop cassette</p> <p><220> <221> polyA_signal <222> (3607)..(3733)</p>
--	--

	<p><223> rabbit globin poly A</p> <p><220></p> <p><221> repeat_region</p> <p><222> (3822)..(3951)</p> <p><223> 3'ITR</p>
<p>40</p>	<p><223> ITR.CB7.CI.aVEGFv9.rBG.ITR</p> <p><220></p> <p><221> repeat_region</p> <p><222> (1)..(130)</p> <p><223> 5'ITR</p> <p><220></p> <p><221> promoter</p> <p><222> (198)..(579)</p> <p><223> CMV IE promoter</p> <p><220></p> <p><221> enhancer</p> <p><222> (279)..(538)</p> <p><223> C4 enhancer</p> <p><220></p> <p><221> promoter</p> <p><222> (582)..(862)</p> <p><223> CB promoter</p> <p><220></p> <p><221> Intron</p> <p><222> (956)..(1928)</p>

	<p><223> chicken beta-actin intron</p> <p><220></p> <p><221> 5'UTR</p> <p><222> (1946)..(1993)</p> <p><223> c-myc 5'UTR</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1994)..(1999)</p> <p><223> kozak</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1999)..(2058)</p> <p><223> leader</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2059)..(2427)</p> <p><223> aVEGFv9 VH</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2428)..(2748)</p> <p><223> CH1</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2752)..(2763)</p> <p><223> furin cleavage site</p>
--	---

	<p><220> <221> misc_feature <222> (2764)..(2835) <223> F2A linker</p> <p><220> <221> misc_feature <222> (2836)..(2895) <223> leader</p> <p><220> <221> misc_feature <222> (2896)..(3216) <223> aVEGFv9 VL</p> <p><220> <221> misc_feature <222> (3217)..(3537) <223> CL</p> <p><220> <221> misc_feature <222> (3538)..(3543) <223> Stop Cassette</p> <p><220> <221> polyA_signal <222> (3613)..(3739) <223> rabbit globin poly A</p> <p><220> <221> repeat_region</p>
--	---

	<p><222> (3828)..(3957) <223> 3'ITR</p>
<p>41</p>	<p><223> ITR.CB7.CI.aVEGFv10.rBG.ITR</p> <p><220> <221> repeat_region <222> (1)..(130) <223> 5'ITR</p> <p><220> <221> promoter <222> (198)..(579) <223> CMV IE promoter</p> <p><220> <221> enhancer <222> (279)..(538) <223> C4 enhancer</p> <p><220> <221> promoter <222> (582)..(862) <223> CB promoter</p> <p><220> <221> Intron <222> (956)..(1928) <223> chicken beta-actin intron</p> <p><220> <221> 5'UTR</p>

	<p><222> (1940)..(1987) <223> c-myc 5'UTR</p> <p><220> <221> misc_feature <222> (1988)..(1993) <223> kozak</p> <p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220> <221> misc_feature <222> (2053)..(2421) <223> aVEGFv10 VH</p> <p><220> <221> misc_feature <222> (2422)..(2742) <223> CH1</p> <p><220> <221> misc_feature <222> (2746)..(2757) <223> furin cleavage site</p> <p><220> <221> misc_feature <222> (2758)..(2829) <223> F2A linker</p>
--	---

	<p><220> <221> misc_feature <222> (2830)..(2889) <223> leader</p> <p><220> <221> misc_feature <222> (2890)..(3210) <223> aVEGFv10 VL</p> <p><220> <221> misc_feature <222> (3211)..(3531) <223> CL</p> <p><220> <221> misc_feature <222> (3532)..(3537) <223> stop cassette</p> <p><220> <221> polyA_signal <222> (3607)..(3733) <223> rabbit globin poly A</p> <p><220> <221> repeat_region <222> (3822)..(3951) <223> 3'ITR</p>
42	<223> ITR.CB7.CI.aVEGFv11.rBG.ITR

	<p><220> <221> repeat_region <222> (1)..(130) <223> 5'ITR</p> <p><220> <221> promoter <222> (198)..(579) <223> CMV IE promoter</p> <p><220> <221> enhancer <222> (279)..(538) <223> C4 enhancer</p> <p><220> <221> promoter <222> (582)..(862) <223> CB promoter</p> <p><220> <221> Intron <222> (956)..(1928) <223> chicken beta-actin intron</p> <p><220> <221> 5'UTR <222> (1940)..(1987) <223> c-myc 5' UTR</p> <p><220></p>
--	--

	<p><221> misc_feature <222> (1988)..(1993) <223> kozak</p> <p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220> <221> misc_feature <222> (2053)..(2421) <223> aVEGFv11 VH</p> <p><220> <221> misc_feature <222> (2422)..(2742) <223> CH1</p> <p><220> <221> misc_feature <222> (2746)..(2754) <223> furing cleavage site</p> <p><220> <221> misc_feature <222> (2755)..(2829) <223> F2A linker</p> <p><220> <221> misc_feature <222> (2830)..(2889)</p>
--	---

	<p><223> leader</p> <p><220></p> <p><221> misc_feature</p> <p><222> (2890)..(3210)</p> <p><223> aVEGFv11 VL</p> <p><220></p> <p><221> misc_feature</p> <p><222> (3211)..(3531)</p> <p><223> CL</p> <p><220></p> <p><221> misc_feature</p> <p><222> (3532)..(3537)</p> <p><223> stop cassette</p> <p><220></p> <p><221> polyA_signal</p> <p><222> (3607)..(3733)</p> <p><223> rabbit globin poly A</p> <p><220></p> <p><221> repeat_region</p> <p><222> (3822)..(3951)</p> <p><223> 3' ITR</p>
43	<p><223> ITR.CB7.CI.aVEGFv12.rBG.ITR</p> <p><220></p> <p><221> repeat_region</p> <p><222> (1)..(130)</p>

	<p><223> 5'ITR</p> <p><220></p> <p><221> promoter</p> <p><222> (198)..(579)</p> <p><223> CMV IE promoter</p> <p><220></p> <p><221> enhancer</p> <p><222> (279)..(538)</p> <p><223> C4 enhancer</p> <p><220></p> <p><221> promoter</p> <p><222> (582)..(862)</p> <p><223> CB promoter</p> <p><220></p> <p><221> Intron</p> <p><222> (956)..(1928)</p> <p><223> chicken beta-actin intron</p> <p><220></p> <p><221> 5'UTR</p> <p><222> (1940)..(1987)</p> <p><223> c-myc 5'UTR</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1988)..(1993)</p> <p><223> kozak</p>
--	--

	<p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220> <221> misc_feature <222> (2053)..(2421) <223> aVEGFv12 VH</p> <p><220> <221> misc_feature <222> (2422)..(2742) <223> CH1</p> <p><220> <221> misc_feature <222> (2746)..(2757) <223> furing cleavage site</p> <p><220> <221> misc_feature <222> (2758)..(2829) <223> F2A linker</p> <p><220> <221> misc_feature <222> (2830)..(2889) <223> leader</p> <p><220> <221> misc_feature</p>
--	---

	<p><222> (2890)..(3210) <223> aVEGFv12 VL</p> <p><220> <221> misc_feature <222> (3211)..(3531) <223> CL</p> <p><220> <221> misc_feature <222> (3532)..(3537) <223> stop cassette</p> <p><220> <221> polyA_signal <222> (3607)..(3733) <223> rabbit globin poly A</p> <p><220> <221> repeat_region <222> (3822)..(3951) <223> 3'ITR</p>
44	<p><223> ITR.CB7.CI.aVEGFv13.rBG.ITR</p> <p><220> <221> repeat_region <222> (1)..(130) <223> 5'ITR</p> <p><220> <221> misc_feature</p>

	<p><222> (131)..(167) <223> part of AAV</p> <p><220> <221> promoter <222> (198)..(579) <223> CME IE promoter</p> <p><220> <221> misc_feature <222> (204)..(233) <223> promoter start</p> <p><220> <221> enhancer <222> (279)..(538) <223> C4 enhancer with 2 mismatches</p> <p><220> <221> misc_feature <222> (561)..(584) <223> CMV promoter end</p> <p><220> <221> promoter <222> (582)..(862) <223> CB promoter</p> <p><220> <221> misc_feature <222> (585)..(615) <223> begin promoter</p>
--	--

	<p><220> <221> TATA_signal <222> (836)..(839)</p> <p><220> <221> Intron <222> (956)..(1928) <223> chichen beta-actin intron</p> <p><220> <221> misc_feature <222> (1814)..(1830) <223> end of intron</p> <p><220> <221> 5'UTR <222> (1940)..(1987) <223> c-myc 5'UTR</p> <p><220> <221> misc_feature <222> (1988)..(1993) <223> kozak</p> <p><220> <221> misc_feature <222> (1993)..(2052) <223> leader</p> <p><220> <221> misc_feature</p>
--	---

	<p><222> (2053)..(2421) <223> aVEGFv13 VH</p> <p><220> <221> misc_feature <222> (2422)..(2742) <223> CH1</p> <p><220> <221> misc_feature <222> (2746)..(2757) <223> furin cleavage site</p> <p><220> <221> misc_feature <222> (2758)..(2829) <223> F2A linker</p> <p><220> <221> misc_feature <222> (2830)..(2889) <223> leader</p> <p><220> <221> misc_feature <222> (2890)..(3210) <223> aVEGFv13 VL</p> <p><220> <221> misc_feature <222> (3211)..(3531) <223> CL</p>
--	--

	<p><220></p> <p><221> misc_feature</p> <p><222> (3532)..(3537)</p> <p><223> stop cassette</p> <p><220></p> <p><221> polyA_signal</p> <p><222> (3607)..(3733)</p> <p><223> rabbit globin poly A</p> <p><220></p> <p><221> misc_feature</p> <p><222> (3785)..(3821)</p> <p><223> part of AAV</p> <p><220></p> <p><221> repeat_region</p> <p><222> (3822)..(3951)</p> <p><223> 3'ITR</p>
<p>45</p>	<p><223></p> <p>ITR.CMV.PI.aVEGFv7.eMCV.IRES.SV40.ITR</p> <p><220></p> <p><221> repeat_region</p> <p><222> (1)..(130)</p> <p><223> ITR</p> <p><220></p> <p><221> promoter</p> <p><222> (191)..(932)</p>

	<p><223> human CMV I.E. enhancer & promoter</p> <p><220></p> <p><221> Intron</p> <p><222> (1047)..(1179)</p> <p><223> Promega chimeric intron</p> <p><220></p> <p><221> 5'UTR</p> <p><222> (1248)..(1295)</p> <p><223> c-myc 5'UTR</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1299)..(1307)</p> <p><223> kozak</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1305)..(1364)</p> <p><223> leader</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1365)..(1685)</p> <p><223> aVEGFv7 VL</p> <p><220></p> <p><221> misc_feature</p> <p><222> (1686)..(2006)</p> <p><223> CL</p>
--	---

	<p><220> <221> enhancer <222> (2018)..(2608) <223> IRES</p> <p><220> <221> misc_feature <222> (2606)..(2665) <223> leader</p> <p><220> <221> misc_feature <222> (2666)..(3034) <223> aVEGFv7 VH</p> <p><220> <221> misc_feature <222> (3035)..(3355) <223> CH1</p> <p><220> <221> polyA_signal <222> (3384)..(3615) <223> SV40 late polyadenylation signal</p> <p><220> <221> repeat_region <222> (3680)..(3809) <223> ITR</p>
46	<p><223> ITR.CMV.PI.aVEGFv7.fmdIRES.SV40.ITR</p>

	<p><220> <221> repeat_region <222> (1)..(130) <223> ITR</p> <p><220> <221> promoter <222> (191)..(932) <223> human CMV I.E. enhancer and promoter</p> <p><220> <221> TATA_signal <222> (897)..(901)</p> <p><220> <221> Intron <222> (1047)..(1179) <223> Promega chimeric intron</p> <p><220> <221> 5'UTR <222> (1248)..(1295) <223> c-myc 5'UTR</p> <p><220> <221> misc_feature <222> (1299)..(1307) <223> kozak</p> <p><220> <221> misc_feature</p>
--	---

	<p><222> (1305)..(1313) <223> leader</p> <p><220> <221> misc_feature <222> (1314)..(1364) <223> leader</p> <p><220> <221> misc_feature <222> (1365)..(1685) <223> aVEGFv7 VL</p> <p><220> <221> misc_feature <222> (1686)..(2006) <223> CL</p> <p><220> <221> misc_feature <222> (2021)..(2482) <223> FMDV</p> <p><220> <221> misc_feature <222> (2501)..(2542) <223> leader</p> <p><220> <221> misc_feature <222> (2543)..(2911) <223> aVEGFv7 VH</p>
--	---

	<p><220> <221> misc_feature <222> (2912)..(3232) <223> CH1</p> <p><220> <221> polyA_signal <222> (3261)..(3492) <223> Sv40 late polyadenylation signal</p> <p><220> <221> repeat_region <222> (3557)..(3686) <223> ITR</p>
<p>47</p>	<p><223> ITR.CMV.PI.aVEGFv7.cMycIRES.SV40.ITR</p> <p><220> <221> repeat_region <222> (1)..(130) <223> ITR</p> <p><220> <221> promoter <222> (191)..(932) <223> human CMV I.E. enhancer and promoter</p> <p><220> <221> TATA_signal <222> (897)..(901)</p>

	<p><220> <221> Intron <222> (1047)..(1179) <223> Promega chimeric intron</p> <p><220> <221> 5'UTR <222> (1248)..(1295) <223> c-myc 5'UTR</p> <p><220> <221> misc_feature <222> (1299)..(1307) <223> kozak</p> <p><220> <221> misc_feature <222> (1305)..(1313) <223> leader</p> <p><220> <221> misc_feature <222> (1314)..(1364) <223> leader</p> <p><220> <221> misc_feature <222> (1365)..(1685) <223> aVEGFv7 VL</p> <p><220></p>
--	--

	<p><221> misc_feature <222> (1686)..(2006) <223> CL</p> <p><220> <221> misc_feature <222> (2021)..(2415) <223> IRES c-myc</p> <p><220> <221> misc_feature <222> (2226)..(2273) <223> mini c-myc IRES</p> <p><220> <221> misc_feature <222> (2275)..(2275) <223> this C to T mutaion increases expression</p> <p><220> <221> misc_feature <222> (2434)..(2475) <223> leader</p> <p><220> <221> misc_feature <222> (2476)..(2844) <223> aVEGFv7 VH</p> <p><220> <221> misc_feature <222> (2845)..(3165)</p>
--	---

	<223> CH1 <220> <221> polyA_signal <222> (3194)..(3425) <223> SV40 late polyadenylation signal <220> <221> repeat_region <222> (3490)..(3619) <223> ITR
48	<223> AAV8 capsid
49	<223> Nucleic acid sequence of AAV8 capsid

All publications cited in this specification are incorporated herein by reference in their entireties, as are US Provisional Patent Application No. 62/466,721, filed March 3, 2017, US Provisional Patent Application No. 62/460,515, filed February 17, 2017, US
 5 Provisional Patent Application No. 62/442,946, filed January 5, 2017, US Provisional Patent Application No. 62/331,100, filed May 3, 2016 and US Provisional Patent Application No. 62/323,184, filed April 15, 2016. Similarly, the Sequence Listing filed herewith is hereby incorporated by reference. While the invention has been described with reference to particular embodiments, it will be appreciated that modifications can be made without
 10 departing from the spirit of the invention. Such modifications are intended to fall within the scope of the appended claims.

CLAIMS

1. A recombinant adeno-associated virus (rAAV) which is suitable for administration to an eye of a human subject wherein the rAAV comprises an expression cassette in an AAV8 capsid, wherein the expression cassette comprises:

a coding sequence for an anti-human vascular endothelial growth factor (hVEGF) antigen binding antibody fragment (Fab) (anti-hVEGF Fab) operably linked to regulatory elements which direct expression thereof, wherein coding sequence comprises:

(i) a nucleotide sequence encoding an anti-hVEGF Fab heavy immunoglobulin chain (HC) having a first interleukin 2 (IL-2) leader sequence operably linked to an anti-hVEGF Fab HC variable region comprising amino acids 1 to 123 of SEQ ID NO: 1,

(ii) a nucleotide sequence encoding an F2A linker,

(iii) a nucleotide sequence encoding anti-hVEGF Fab light chain (LC) having a second IL-2 leader sequence operably linked to nucleotides encoding an anti-hVEGF Fab LC variable region comprising amino acids 1 to 107 of SEQ ID NO: 2; and

wherein the regulatory elements comprise a cytomegalovirus immediate early (CMV IE) enhancer, a chicken beta-actin promoter, and a chicken beta-actin intron.

2. The rAAV according to claim 1, wherein the regulatory elements further comprise a polyadenylation (polyA) signal sequence.

3. The rAAV according to claim 2, wherein the polyA signal sequence is a rabbit beta globin polyA signal sequence.

4. The rAAV according to any one of claims 1 to 3, wherein the rAAV comprises a vector genome, wherein said vector genome further comprises, in the following order, an AAV 5' inverted terminal repeat (ITR), the expression cassette, and an AAV 3' ITR.

5. The rAAV according to claim 4, wherein the AAV 5' ITR and AAV 3' ITR are from AAV2.

6. The rAAV according to any one of claims 1 to 5, wherein the coding sequence for the anti-hVEGF Fab HC encodes the amino acid sequence of SEQ ID NO: 1 and the coding sequence for the anti-hVEGF Fab LC encodes the amino acid sequence of SEQ ID NO: 2.

7. The rAAV according to any one of claims 1 to 6, wherein the regulatory elements further comprise a UTR sequence.

8. The rAAV according to any one of claims 1 to 7, wherein the F2A linker further comprises a furin recognition site.

9. The rAAV according to any one of claims 1 to 8, wherein the first IL-2 leader sequence and the second IL-2 leader sequence each comprise the amino acid sequence of SEQ ID NO: 30.

10. The rAAV according to any one of claims 1 to 9, wherein the coding sequence for the anti-hVEGF Fab HC and anti-hVEGF Fab LC comprise nucleotides 1842 to 2210 for SEQ ID NO: 19 for the anti-hVEGF Fab HC variable region and nucleotides 2679 to 2999 of SEQ ID NO: 19 for the anti-hVEGF Fab LC variable chain (aVEGFv3).

11. The rAAV according to any one of claims 1 to 10, wherein the coding sequence for the anti-VEGF Fab HC comprises nucleotides 1842 to 2531 of SEQ ID NO: 19 and the coding sequence for the anti-VEGF Fab LC comprises nucleotides 2679 to 3320 of SEQ ID NO: 19 (aVEGFv3).

12. A recombinant adeno-associated virus (rAAV) having an AAV8 capsid which is suitable for administration to a human eye, wherein the rAAV comprises an expression cassette in the AAV8 capsid, said expression cassette comprises nucleotides 198 to 3733 of SEQ ID NO: 14.

13. A recombinant adeno-associated virus (rAAV) having an AAV8 capsid, wherein the rAAV comprises a vector genome in the AAV8 capsid, wherein the vector genome comprises a AAV2 5' inverted terminal repeat (ITR), an expression cassette comprising the nucleotides 198 to 3733 of SEQ ID NO: 14, and a AAV2 3' ITR.

14. A recombinant adeno-associated virus (rAAV) having an AAV8 capsid which is suitable for administration to an eye, wherein the rAAV comprises a vector genome packaged within the capsid, said vector genome comprising SEQ ID NO: 14 (ITR-CB7-CI-aVEGFv3-rBG-ITR).

15. A liquid suspension suitable for administration to a human eye, said liquid suspension comprising an aqueous liquid and recombinant adeno-associated virus (rAAV) according to any one of claims 1 to 14 and optionally one or more excipients, preservatives, and/or surfactants.

16. A recombinant adeno-associated virus (rAAV) according to any one of claims 1 to 14 or the liquid suspension according to claim 15 when used in subretinal administration to a patient in need thereof.

17. The rAAV according to any one of claims 1 to 14 or the liquid suspension according to claim 15, wherein said human subject has wet age-related macular degeneration.

18. Use of a rAAV according to any one of claims 1 to 14 or a liquid suspension according to claim 15 in the preparation of a medicament for subretinal administration to a human subject.

19. The use according to claim 18, wherein the human subject has wet age-related macular degeneration.

20. The rAAV according to claim 16, or the use according to claim 18, wherein the administration comprises about 1×10^8 genome copies (GC) per eye (GC/eye) to about 1.5×10^{12} GC/eye.

21. A product comprising: (a) a container comprising an rAAV according to any one of claims 1 to 14 and an aqueous liquid, and optionally a second container comprising (b) a diluent, and (c) a needle for administration.

22. The rAAV according to claim 9, wherein the nucleotide sequence encoding the first IL-2 leader sequence comprises nucleotides 1993 to 2052 of SEQ ID NO: 14.

23. The rAAV according to claim 9, wherein the nucleotide sequence encoding the second IL-2 leader sequence comprises nucleotides 2830 to 2889 of SEQ ID NO: 14.

24. The rAAV according to claim 9, wherein the nucleotide sequence encoding the first IL-2 leader sequence comprises nucleotides 1993 to 2052 of SEQ ID NO: 14 and the nucleotide sequence encoding the second IL-2 leader sequence comprises nucleotides 2830 to 2889 of SEQ ID NO: 14.

25. The rAAV according to any one of claims 1 to 10, wherein the CMV IE enhancer, the chicken beta actin promoter, and the chicken beta actin intron together comprise nucleotides 198 through 1928 of SEQ ID NO:14.

26. The rAAV according to claim 3, wherein the rabbit beta globin polyA signal sequence comprises nucleotides 3607 through 3733 of SEQ ID NO:14.

27. The rAAV according to claim 8, wherein the nucleotide sequence encoding the F2A linker further comprising a furin recognition site comprises nucleotides 2746 through 2829 of SEQ ID NO 14.




FIG 1




FIG 2

Replacement Sheet




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D

Replacement Sheet




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D

Replacement Sheet

FIG. 5A

Animal ID: C64956; Group 2; FFB-314




FIG. 5B

Animal ID: C

Group 1; RGX-314; 1.00×10^{12} GC/eye




FIG. 5C

Animal ID: C

Group 1; RGX-314; 1.00×10^{12} GC/eye





FIG. 5D

Animal ID: C

Group 1; RGX-314; 1.00×10^{12} GC/eye



6/10

Replacement Sheet

Animal ID: C62442; Group 2;
 1.00×10^{11} GC/eye; right eye




FIG. 6A




FIG. 6B

Animal ID: C63116; Group 2;
 1.00×10^{11} GC/eye; right eye




FIG. 6C




FIG. 6D

7/10

Replacement Sheet

Animal ID: C66122; Group 3;
1.00 x 10¹¹ GC/eye; right eye




FIG. 7A




FIG. 7B

Animal ID: C71941; Group 3;
1.00 x 10¹¹ GC/eye; right eye




FIG. 7C




FIG. 7D

8/10

Replacement Sheet

Animal ID: C74420; Group 5;
 1.00×10^{11} GC/eye; right eye




FIG. 8A




FIG. 8B

Animal ID: C65873; Group 5;
 1.00×10^{11} GC/eye; right eye




FIG. 8C




FIG. 8D

9/10

Replacement Sheet




FIG. 9

10/10
Replacement Sheet




FIG. 10A

Controls

Spike copies	Vector Copies	Passage 1	Passage 2	Passage 3
blank	blank	7.78E+00	7.47E+00	2.7E+01
0	0	0.00E+00	1.03E+01	5.5E+00
0	1.E+11	1.23E+02	7.91E+00	1.5E+01

Sensitivity : 1×10^4 wt AAV8 GC
(~ 1 TCID50 IU) in 1×10^{11} GC

FIG. 10B




FIG. 10C




FIG. 10D

SEQUENCE LISTING

<110> The Trustees of the University of Pennsylvania

*<120> COMPOSITIONS FOR TREATMENT OF WET AGE-RELATED
MACULAR*

DEGENERATION

<130> UPN-16-7683PCT

<150> US 62/466,721

<151> 2017-03-03

<150> US 62/460,515

<151> 2017-02-17

<150> US 62/442,946

<151> 2017-01-05

<150> US 62/331,100

<151> 2016-05-03

<150> *US 62/323,184*

<151> *2016-04-15*

<160> *49*

<170> *PatentIn version 3.5*

<210> *1*

<211> *231*

<212> *PRT*

<213> *Artificial Sequence*

<220>

<223> *Humanized anti-VEGF Fab heavy chain*

<220>

<221> *MISC_FEATURE*

<222> *(28)..(39)*

<223> *complementarity determining region*

<220>

<221> MISC_FEATURE

<222> (54)..(83)

<223> complementarity determining region

<400> 1

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr

20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe

50 55 60

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr

65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

85 90 95

Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val

100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly

115 120 125

Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly

130 135 140

Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val

145 150 155 160

Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe

165 170 175

Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val

180 185 190

Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val

195 200 205

Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys

210 215 220

Ser Cys Asp Lys Thr His Leu

225

230

<210> 2

<211> 214

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Humanized anti-VEGF Fab*

<220>

<221> MISC_FEATURE

<222> (26)..(37)

<223> *complementarity determining region*

<220>

<221> MISC_FEATURE

<222> (107)..(117)

<223> *complementarity determining region*

<400> 2

Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr

20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile

35 40 45

Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp

85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala

100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly

115 120 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala

130 135 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln

145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser

165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr

180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser

195 200 205

Phe Asn Arg Gly Glu Cys

210

<210> 3

<211> 3957

<212> *DNA*

<213> *Artificial Sequence*

<220>

<223> *5'ITR.CB7.Cl.alphaVEGFv2.rBG.3'ITR*

<220>

<221> *repeat_region*

<222> *(1)..(130)*

<223> *5' ITR*

<220>

<221> *promoter*

<222> *(198)..(579)*

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> *(279)..(538)*

<223> *C4 enhancer with 2 mismatches*

<220>

<221> *misc_feature*

<222> (279)..(538)

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *TATA_signal*

<222> (836)..(839)

<220>

<221> *misc_feature*

<222> (955)..(1829)

<223> *chicken beta-actin intron*

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin promoter*

<220>

<221> *5'UTR*

<222> (1946)..(1993)

<223> *c-myc 5' UTR*

<220>

<221> *misc_feature*

<222> (1994)..(1999)

<223> *kozak sequence*

<220>

<221> *transit_peptide*

<222> (1999)..(2058)

<220>

<221> CDS

<222> (2059)..(2427)

<223> *α*VEGFv2 VH

<220>

<221> CDS

<222> (2428)..(2748)

<223> CH1

<220>

<221> *misc_feature*

<222> (2752)..(2763)

<223> *Furin cleavage site*

<220>

<221> CDS

<222> (2764)..(2835)

<223> *F2A linker*

<220>

<221> *transit_peptide*

<222> (2836)..(2895)

<220>

<221> CDS

<222> (2896)..(3216)

<223> *α*VEGFv2 VL

<220>

<221> CDS

<222> (3217)..(3537)

<223> CL

<220>

<221> *polyA_signal*

<222> (3613)..(3739)

<223> *rabbit globin polyA*

<220>

<221> *repeat_region*

<222> (3828)..(3957)

<223> 3' ITR

<400> 3

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgtcg ggcgaccttt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatgggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgcct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgcca agtacgcccc ctattgacgt
480

caatgacggt aatggcccc cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcat cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctccccccc ctccccacc ccaattttgt atttatttat
660

ttttaatta ttttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720

ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

gagcggcgcg ctccgaaagt ttccttttat ggcgaggcgg cggcggcggc
ggcctataa 840

*aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900*

*cgccgccgcc tcgcgccgcc cgccccggct ctgactgacc gcgttactcc cacagggtgag
960*

*cgggcgggac ggcccttctc ctccgggctg taattagcgc ttggttaat gacggcttg
1020*

*ttcttttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080*

*gagcggctcg gggggtgcgt gcgtgtgtgt gtgcgtgggg agcgcgcgct
gcggctccgc 1140*

*gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200*

*gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260*

*caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320*

*cgggctgcaa cccccctgc acccccctcc ccgagttgct gagcacggcc cggcttcggg
1380*

*tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcgga 1440*

*ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggcg 1500*

*cggcggcccc cggagcgccg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca ccccctctag cgggcgcggg gcgaagcggg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttcttcttt ttctacagc tctgggcaa cgtgctggtt attgtgctgt ctcattcatt
1920*

*tggcaaagaa ttcgctagag ctagcgggca cttgcaactg gaacttaca
cacccgagca 1980*

*aggacgcgac tctccaccat gtatcggatg cagctgctgc tgctgattgc tctgagcctg
2040*

*gctctggtga ccaacagc gaa gtg cag ctg gtg gaa agc ggc ggc ggc ctg
2091*

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu

1 5 10

*gtg cag ccc ggc ggc agc ctg cgg ctg agc tgt gct gct agc ggc tat
2139*

Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr

15 20 25

gac ttt acc cat tat ggc atg aac tgg gtg cgg cag gct ccc ggc aag
2187

Asp Phe Thr His Tyr Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys

30 35 40

ggc ctg gaa tgg gtg ggc tgg att aac acc tat acc ggc gaa ccc acc
2235

Gly Leu Glu Trp Val Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr

45 50 55

tat gct gct gac ttt aag cgg cgg ttt acc ttt agc ctg gac acc agc 2283

Tyr Ala Ala Asp Phe Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser

60 65 70 75

aag agc acc gct tat ctg cag atg aac agc ctg cgg gct gaa gac acc
2331

Lys Ser Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr

80 85 90

gct gtg tat tat tgt gct aag tat ccc tat tat tat ggc acc agc cat 2379

Ala Val Tyr Tyr Cys Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His

95 100 105

tgg tat ttt gac gtg tgg ggc cag ggc acc ctg gtg acc gtg agc agc

2427

Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser

110 115 120

gct agc acc aag ggc ccc agc gtg ttt ccc ctg gct ccc agc agc aag

2475

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys

125 130 135

agc acc agc ggc ggc acc gct gct ctg ggc tgt ctg gtg aag gac tat

2523

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr

140 145 150 155

ttt ccc gaa ccc gtg acc gtg agc tgg aac agc ggc gct ctg acc agc

2571

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser

160

165

170

ggc gtg cat acc ttt ccc gct gtg ctg cag agc agc ggc ctg tat agc
2619

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser

175

180

185

ctg agc agc gtg gtg acc gtg ccc agc agc agc ctg ggc acc cag acc
2667

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr

190

195

200

tat att tgt aac gtg aac cat aag ccc agc aac acc aag gtg gac aag
2715

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

205

210

215

aag gtg gaa ccc aag agc tgt gac aag acc cat ctgcggaagc ggcgg gct
2766

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Ala

220

225

230

ccc gtg aag cag acc ctg aac ttt gac ctg ctg aag ctg gct ggc gac
2814

Pro Val Lys Gln Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp
235 240 245

gtg gaa agc aac ccc ggc ccc atgtatcgga tgcagctgct gctgctgatt
2865

Val Glu Ser Asn Pro Gly Pro
250

gctctgagcc tggctctggt gaccaacagc gac att cag ctg acc cag agc ccc
2919

Asp Ile Gln Leu Thr Gln Ser Pro
255 260

agc agc ctg agc gct agc gtg ggc gac cgg gtg acc att acc tgt agc
2967

Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Ser
265 270 275

gct agc cag gac att agc aac tat ctg aac tgg tat cag cag aag ccc
3015

Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro

280 285 290

ggc aag gct ccc aag gtg ctg att tat ttt acc agc agc ctg cat agc
3063

Gly Lys Ala Pro Lys Val Leu Ile Tyr Phe Thr Ser Ser Leu His Ser

295 300 305 310

ggc gtg ccc agc cgg ttt agc ggc agc ggc agc ggc acc gac ttt acc
3111

Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr

315 320 325

ctg acc att agc agc ctg cag ccc gaa gac ttt gct acc tat tat tgt 3159

Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys

330 335 340

cag cag tat agc acc gtg ccc tgg acc ttt ggc cag ggc acc aag gtg
3207

Gln Gln Tyr Ser Thr Val Pro Trp Thr Phe Gly Gln Gly Thr Lys Val

345 350 355

gaa att aag cgg acc gtg gct gct ccc agc gtg ttt att ttt ccc ccc 3255

Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro

360 365 370

agc gac gaa cag ctg aag agc ggc acc gct agc gtg gtg tgt ctg ctg

3303

Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu

375 380 385 390

aac aac ttt tat ccc cgg gaa gct aag gtg cag tgg aag gtg gac aac

3351

Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn

395 400 405

gct ctg cag agc ggc aac agc cag gaa agc gtg acc gaa cag gac agc

3399

Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser

410 415 420

aag gac agc acc tat agc ctg agc agc acc ctg acc ctg agc aag gct

3447

Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala

425

430

435

gac tat gaa aag cat aag gtg tat gct tgt gaa gtg acc cat cag ggc
3495

Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly

440

445

450

ctg agc agc ccc gtg acc aag agc ttt aac cgg ggc gaa tgt 3537

Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

455

460

465

tgataaagcg gccgcggtac ctctagagtc gacccgggcg gcctcgagga
cggggtgaac 3597

tacgcctgag gatccgatct tttccctct gccaaaaatt atggggacat catgaagccc
3657

cttgagcatc tgacttctgg ctaataaagg aaatttattt tcattgcaat agtgtgttgg
3717

aatTTTTTgt gtctctcact cggaagcaat tcgttgatct gaatttcgac cacccataat
3777

accattacc ctggtagata agtagcatgg cgggtaatc attaactaca aggaaccct
3837

agtgatggag ttggccactc cctctctgcg cgctcgctcg ctactgagg ccgggcgacc
3897

aaaggtcgcc cgacgcccgg gctttgcccg ggcggcctca gtgagcgagc
gagcgcgag 3957

<210> 4

<211> 123

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 4

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr

20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe

50 55 60

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr

65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

85 90 95

Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val

100

105

110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser

115

120

<210> 5

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 5

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys

1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr

20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser

35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser

50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr

65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

85 90 95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His

100

105

<210> 6

<211> 24

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 6

Ala Pro Val Lys Gln Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly

1

5

10

15

Asp Val Glu Ser Asn Pro Gly Pro

20

<210> 7

<211> 107

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 7

Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr

20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile

35 40 45

Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp

85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys

100 105

<210> 8

<211> 107

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 8

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu

1 5 10 15

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe

20 25 30

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln

35 40 45

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser

50 55 60

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu

65 70 75 80

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser

85 90 95

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

100 105

<210> 9

<211> 3770

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *AAV25'ITR.UbC.Ci.aVEGFv2.rBG.AAV23'ITR*

<220>

<221> *repeat_region*

<222> *(17)..(146)*

<223> *5' ITR*

<220>

<221> *promoter*

<222> *(207)..(1435)*

<223> *UbC with C insertion at 289 and G insertion at 990*

<220>

<221> *Intron*

<222> *(1529)..(1661)*

<223> *chimeric intron*

<220>

<221> *5'UTR*

<222> (1736)..(1783)

<223> *c-myc* 5'UTR

<220>

<221> *misc_feature*

<222> (1784)..(1789)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1789)..(1848)

<223> *kozak*

<220>

<221> *CDS*

<222> (1849)..(2217)

<223> *aVEGFv2 VH*

<220>

<221> *CDS*

<222> (2218)..(2538)

<223> CH1

<220>

<221> *misc_feature*

<222> (2542)..(2553)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2554)..(2625)

<223> *F2a linker*

<220>

<221> *misc_feature*

<222> (2626)..(2685)

<223> *Leader*

<220>

<221> CDS

<222> (2686)..(3006)

<223> *aVEGFv2 VL*

<220>

<221> CDS

<222> (3007)..(3327)

<223> CL

<220>

<221> *polyA_signal*

<222> (3345)..(3576)

<223> *SV40 late polyadenylation signal*

<220>

<221> *repeat_region*

<222> (3641)..(3770)

<223> *3' ITR*

<400> 9

taaggcctta attaggctgc gcgctcgctc gctcactgag gccgcccggg caaagcccgg

60

gcgtcgggcg acctttggtc gcccggcctc agtgagcgag cgagcgcgca

gagagggagt 120

ggccaactcc atcactaggg gttccttgta gttaatgatt aacccgccat gctacttatc
180

tacgtagcca tgctctagga agatctggcc tccgcgccgg gttttggcgc ctcccgcggg
240

cgccccctc ctcacggcga gcgctgccac gtcagacgaa gggcgagcg
agcgtcctga 300

tccttccgcc cggacgctca ggacagcggc ccgctgctca taagactcgg ccttagaacc
360

ccagtatcag cagaaggaca ttttaggacg ggacttgggt gactctaggg cactggtttt
420

ctttccagag agcggaacag gcgaggaaaa gtagtccctt ctcggcgatt
ctgcggaggg 480

atctccgtgg ggcggtgaac gccgatgatt atataaggac gcgccgggtg
tggcacagct 540

agttccgtcg cagccgggat ttgggtcgcg gttcttgttt gtggatcgct gtgatcgta
600

cttggtgagt agcgggctgc tgggctggcc ggggcttctg tggccgccgg gccgctcgg
660

gggacggaag cgtgtggaga gaccgccaag ggctgtagtc tgggtccgcg
agcaagggtg 720

ccctgaactg ggggttgggg ggagcgcagc aaaatggcgg ctgtcccga
gtcttgaatg 780

gaagacgctt gtgaggcggg ctgtgaggtc gttgaaacaa ggtggggggc
atggtgggcg 840

gcaagaacct aaggtcttga ggccttcgct aatgcgggaa agctcttatt cgggtgagat
900

gggctggggc accatctggg gaccctgacg tgaagtttgt cactgactgg
agaactcgg 960

ttgtcgtctg ttgcgggggc ggcagttatg gcggtgccgt tgggcagtgc accgtacct
1020

ttgggagcgc gcgccctcgt cgtgtcgtga cgtcacccgt tctgttggct tataatgcag
1080

ggtggggcca cctgccgta ggtgtgcggt aggcttttct ccgtcgcagg acgcagggtt
1140

cgggcctagg gtaggctctc ctgaatcgac aggcgccgga cctctggtga
ggggagggat 1200

aagtgaggcg tcagtttctt tggtcggttt tatgtaccta tcttcttaag tagctgaagc
1260

tccggttttg aactatgcgc tcggggttgg cgagtgtggt ttgtgaagtt ttttaggcac
1320

cttttgaat gtaatcattt gggccaatat gtaattttca gtgtagact agtaaattgt
1380

ccgctaaatt ctggccgttt ttggcttttt tgtagacga agctttattg cggtagtta
1440

tcacagttaa attgctaacg cagtcagtgc ttctgacaca acagtctcga acttaagctg
1500

cagaagttgg tcgtgaggca ctgggcaggt aagtatcaag gttacaagac
aggtttaagg 1560

agaccaatag aaactgggct tgtcgagaca gagaagactc ttgcgtttct
gataggcacc 1620

tattggcttt actgacatcc actttgcctt tctctccaca ggtgtccact cccagttcaa
1680

ttacagctct taaggctaga gtacttaata cgactcacta taggctagag ctagcgggca
1740

ctttgcactg gaacttaca caccgagca aggacgcgac tctccaccat gtatcggatg
1800

cagctgctgc tgctgattgc tctgagcctg gctctggtga ccaacagc gaa gtg cag
1857

Glu Val Gln

1

ctg gtg gaa agc ggc ggc ggc ctg gtg cag ccc ggc ggc agc ctg cgg
1905

Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg

5 10 15

ctg agc tgt gct gct agc ggc tat gac ttt acc cat tat ggc atg aac 1953

Leu Ser Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr Gly Met Asn

20 25 30 35

tgg gtg cgg cag gct ccc ggc aag ggc ctg gaa tgg gtg ggc tgg att
2001

Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly Trp Ile

40 45 50

aac acc tat acc ggc gaa ccc acc tat gct gct gac ttt aag cgg cgg
2049

Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Arg Arg

55 60 65

ttt acc ttt agc ctg gac acc agc aag agc acc gct tat ctg cag atg 2097

Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr Leu Gln Met

70 75 80

aac agc ctg cgg gct gaa gac acc gct gtg tat tat tgt gct aag tat
2145

Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr
85 90 95

ccc tat tat tat ggc acc agc cat tgg tat ttt gac gtg tgg ggc cag 2193

Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val Trp Gly Gln
100 105 110 115

ggc acc ctg gtg acc gtg agc agc gct agc acc aag ggc ccc agc gtg
2241

Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
120 125 130

ttt ccc ctg gct ccc agc agc aag agc acc agc ggc ggc acc gct gct
2289

Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
135 140 145

ctg ggc tgt ctg gtg aag gac tat ttt ccc gaa ccc gtg acc gtg agc
2337

Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser

150 155 160

tgg aac agc ggc gct ctg acc agc ggc gtg cat acc ttt ccc gct gtg
2385

Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val

165 170 175

ctg cag agc agc ggc ctg tat agc ctg agc agc gtg gtg acc gtg ccc
2433

Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro

180 185 190 195

agc agc agc ctg ggc acc cag acc tat att tgt aac gtg aac cat aag
2481

Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys

200 205 210

ccc agc aac acc aag gtg gac aag aag gtg gaa ccc aag agc tgt gac
2529

Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp

215 220 225

aag acc cat ctgcggaagc ggcgggctcc cgtgaagcag accctgaact
2578

Lys Thr His

230

ttgacctgct gaagctggct ggcgacgtgg aaagcaacc cggccccatg
tatcggatgc 2638

agctgctgct gctgattgct ctgagcctgg ctctggtgac caacagc gac att cag
2694

Asp Ile Gln

ctg acc cag agc ccc agc agc ctg agc gct agc gtg ggc gac cgg gtg
2742

Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val

235

240

245

acc att acc tgt agc gct agc cag gac att agc aac tat ctg aac tgg
2790

Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp

250 255 260 265

tat cag cag aag ccc ggc aag gct ccc aag gtg ctg att tat ttt acc
2838

Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Phe Thr

270 275 280

agc agc ctg cat agc ggc gtg ccc agc cgg ttt agc ggc agc ggc agc
2886

Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser

285 290 295

ggc acc gac ttt acc ctg acc att agc agc ctg cag ccc gaa gac ttt
2934

Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe

300 305 310

gct acc tat tat tgt cag cag tat agc acc gtg ccc tgg acc ttt ggc 2982

Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp Thr Phe Gly

315 320 325

cag ggc acc aag gtg gaa att aag cgg acc gtg gct gct ccc agc gtg
3030

Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val

330 335 340 345

ttt att ttt ccc ccc agc gac gaa cag ctg aag agc ggc acc gct agc
3078

Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser

350 355 360

gtg gtg tgt ctg ctg aac aac ttt tat ccc cgg gaa gct aag gtg cag
3126

Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln

365 370 375

tgg aag gtg gac aac gct ctg cag agc ggc aac agc cag gaa agc gtg
3174

Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val

380 385 390

acc gaa cag gac agc aag gac agc acc tat agc ctg agc agc acc ctg
3222

Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu

395 400 405

acc ctg agc aag gct gac tat gaa aag cat aag gtg tat gct tgt gaa
3270

Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu

410 415 420 425

gtg acc cat cag ggc ctg agc agc ccc gtg acc aag agc ttt aac cgg
3318

Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg

430 435 440

ggc gaa tgt tgataaagcg gccgcttcga gcagacatga taagatacat
3367

Gly Glu Cys

tgatgagttt ggacaaacca caactagaat gcagtgaaaa aaatgcttta
tttgtgaaat 3427

ttgtgatgct attgctttat ttgtaacat tataagctgc aataacaag ttaacaaca
3487

caattgcatt cattttatgt ttcaggttca gggggagatg tgggaggttt ttaaagcaa
3547

gtaaacctc tacaatgtg gtaaatcga taaggatctt cctagagcat
ggctacgtag 3607

ataagtagca tggcgggta atcattaact acaaggaacc cctagtgatg
gagttggcca 3667

ctccctctct gcgcgctcgc tcgctcactg aggccgggcg accaaaggtc gcccgacgcc
3727

cgggctttgc ccgggcggcc tcagtgagcg agcgagcgcg cag 3770

<210> 10

<211> 123

<212> PRT

<213> Artificial Sequence

<220>

<223> *Synthetic Construct*

<400> 10

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr

20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe

50 55 60

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr

65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

85 90 95

Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val

100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser

115 120

<210> 11

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> *Synthetic Construct*

<400> 11

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys

1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr

20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser

35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser

50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr

65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

85 90 95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His

100 105

<210> 12

<211> 107

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 12

Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr

20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile

35 40 45

Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp

85

90

95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys

100

105

<210> 13

<211> 107

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 13

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu

1

5

10

15

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe

20

25

30

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln

35

40

45

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser

50

55

60

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu

65

70

75

80

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser

85

90

95

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

100

105

<210> 14

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGRv3.rBG.ITR*

<220>

<221> *repeat_region*

<222> *(1)..(130)*

<223> *5'ITR*

<220>

<221> *promoter*

<222> (198)..(579)

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> (279)..(538)

<223> *C4 enhancer with 2 mismatches*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *TATA_signal*

<222> (836)..(839)

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> 5'UTR

<222> (1940)..(1987)

<220>

<221> *misc_feature*

<222> (1988)..(1993)

<223> *Kozak*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *CDS*

<222> (2053)..(2421)

<223> *aVEGFv3 VH*

<220>

<221> CDS

<222> (2422)..(2742)

<223> CH1

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2a linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> CDS

<222> (2890)..(3210)

<223> *α*VEGFv3 VL

<220>

<221> CDS

<222> (3211)..(3531)

<223> CL

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *rabbit globin polyA*

<220>

<221> *repeat_region*

<222> (3822)..(3951)

<223> *3' ITR*

<400> 14

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgctc ggcgacctt

60

*ggtcgcccgg cctcagtgag cgagcgagcg cgagagagg gagtggccaa
ctccatcact 120*

*aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatgggg
180*

*atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240*

*tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300*

*tggcccgct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360*

*tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420*

*aactgcccac ttggcagtac atcaagtgta tcatatgccca agtacgcccc ctattgacgt
480*

caatgacggt aatggcccg cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcat cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctccccccc ctccccaccc ccaattttgt atttatttat
660

ttttaatta ttttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720

ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

gagcggcggc ctccgaaagt ttccttttat ggcgagggcg cggcggcggc
ggccctataa 840

aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900

cgccgccgcc tcgcgccgcc cgccccggct ctgactgacc gcgttactcc cacaggtgag
960

cgggcgggac ggcccttctc ctccgggctg taattagcgc ttggttaat gacggcttgt
1020

ttcttttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080

gagcggctcg ggggggtgcgt gcgtgtgtgt gtgcgtgggg agcgcgcgct
gcggctccgc 1140

gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200

gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260

caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggc 1320

cgggctgcaa cccccctgc acccccctcc ccgagttgct gagcacggcc cggcttcggg
1380

*tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcggca 1440*

*ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggcg 1500*

*cggcggcccc cggagcggcg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccccttag cgggcgcggg gcgaagcgg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

cttcttcttt ttctacagc tctgggcaa cgtgctggtt attgtgctgt ctcattcattt
1920

tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980

cgactctcca ccatgtaccg tatgcagctg ctgctgctga tcgcaactgtc actggcactg
2040

gttaccaact ca gaa gtt cag ctg gtt gaa tca ggc ggc ggc ctg gtt cag
2091

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln

1 5 10

ccc ggc ggc tca ctg cgt ctg tca tgt gca gca tca ggc tac gat ttc 2139

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Asp Phe

15 20 25

acc cac tac ggc atg aac tgg gtt cgt cag gca ccc ggc aaa ggc ctg
2187

Thr His Tyr Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu

30 35 40 45

gaa tgg gtt ggc tgg atc aac acc tac acc ggc gaa ccc acc tac gca
2235

Glu Trp Val Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala

50 55 60

gca gat ttc aaa cgt cgt ttc acc ttc tca ctg gat acc tca aaa tca 2283

Ala Asp Phe Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser

65 70 75

acc gca tac ctg cag atg aac tca ctg cgt gca gaa gat acc gca gtt
2331

Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val

80 85 90

tac tac tgt gca aaa tac ccc tac tac tac ggc acc tca cac tgg tac 2379

Tyr Tyr Cys Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr

95 100 105

ttc gat gtt tgg ggc cag ggc acc ctg gtt acc gtt tca tca gca tca 2427

Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser

110 115 120 125

acc aaa ggc ccc tca gtt ttc ccc ctg gca ccc tca tca aaa tca acc 2475

Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr

130 135 140

tca ggc ggc acc gca gca ctg ggc tgt ctg gtt aaa gat tac ttc ccc
2523

Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro

145 150 155

gaa ccc gtt acc gtt tca tgg aac tca ggc gca ctg acc tca ggc gtt
2571

Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val

160 165 170

cac acc ttc ccc gca gtt ctg cag tca tca ggc ctg tac tca ctg tca 2619

His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser

175 180 185

tca gtt gtt acc gtt ccc tca tca tca ctg ggc acc cag acc tac atc 2667

Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile

190 195 200 205

tgt aac gtt aac cac aaa ccc tca aac acc aaa gtt gat aaa aaa gtt
2715

Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val

210 215 220

gaa ccc aaa tca tgt gat aaa acc cac ctgctgtaaac gtcgtgcacc
2762

Glu Pro Lys Ser Cys Asp Lys Thr His

225 230

cgtaaacag accctgaact tcgatctgct gaaactggca ggcgatgttg aatcaaacc
2822

cggcccatg taccgatgc agctgctgct gctgatcgca ctgtcactgg cactggttac
2882

caactca gat atc cag ctg acc cag tca ccc tca tca ctg tca gca tca
2931

Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser

235 240

gtt ggc gat cgt gtt acc atc acc tgt tca gca tca cag gat atc tca 2979

Val Gly Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser

245 250 255 260

aac tac ctg aac tgg tac cag cag aaa ccc ggc aaa gca ccc aaa gtt
3027

Asn Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val

265 270 275

ctg atc tac ttc acc tca tca ctg cac tca ggc gtt ccc tca cgt ttc 3075

Leu Ile Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe

280 285 290

tca ggc tca ggc tca ggc acc gat ttc acc ctg acc atc tca tca ctg 3123

Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu

295 300 305

cag ccc gaa gat ttc gca acc tac tac tgt cag cag tac tca acc gtt 3171

Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val

310 315 320

ccc tgg acc ttc ggc cag ggc acc aaa gtt gaa atc aaa cgt acc gtt
3219

Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val

325 330 335 340

gca gca ccc tca gtt ttc atc ttc ccc ccc tca gat gaa cag ctg aaa 3267

Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys

345 350 355

tca ggc acc gca tca gtt gtt tgt ctg ctg aac aac ttc tac ccc cgt 3315

Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg

360 365 370

gaa gca aaa gtt cag tgg aaa gtt gat aac gca ctg cag tca ggc aac
3363

Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn

375 380 385

tca cag gaa tca gtt acc gaa cag gat tca aaa gat tca acc tac tca
3411

Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser

390

395

400

ctg tca tca acc ctg acc ctg tca aaa gca gat tac gaa aaa cac aaa
3459

Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys

405

410

415

420

gtt tac gca tgt gaa gtt acc cac cag ggc ctg tca tca ccc gtt acc 3507

Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr

425

430

435

aaa tca ttc aac cgt ggc gaa tgt tgataaagcg gccgcggtac ctctagagtc
3561

Lys Ser Phe Asn Arg Gly Glu Cys

440

gacccgggcg gcctcgagga cggggtgaac tacgcctgag gatccgatct tttccctct
3621

gccaaaaatt atggggacat catgaagccc cttgagcatc tgacttctgg ctaataaagg
3681

aaatttattt tcattgcaat agtgtgttg aatTTTTgt gtctctcact cggaagcaat
3741

tcgttgatct gaatttcgac cacccataat acccattacc ctggtagata agtagcatgg
3801

cgggttaatc attaactaca aggaaccct agtgatggag ttggccactc cctctctgcg
3861

cgctcgctcg ctactgagg ccgggCGacc aaaggTCGCC cgacgcccgg gctttgcccg
3921

ggcggcctca gtgagcgagc gagcgcgcag 3951

<210> 15

<211> 123

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 15

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr

20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe

50 55 60

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr

65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

85

90

95

Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val

100

105

110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser

115

120

<210> 16

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 16

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys

1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr

20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser

35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser

50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr

65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

85

90

95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His

100

105

<210> 17

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 17

Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr

20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile

35 40 45

Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp

85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys

100

105

<210> 18

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 18

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu

1

5

10

15

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe

20 25 30

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln

35 40 45

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser

50 55 60

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu

65 70 75 80

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser

85 90 95

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

100 105

<210> 19

<211> 3763

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.UbC.PI.alphaVEGFv3.SV40.ITR*

<220>

<221> *repeat_region*

<222> (17)..(146)

<223> *AAV2 5' ITR*

<220>

<221> *promoter*

<222> (207)..(1434)

<223> *UbC, with C insert at 289 and G insert at 990*

<220>

<221> *Intron*

<222> (1528)..(1660)

<223> *chimeric intron*

<220>

<221> *5'UTR*

<222> (1729)..(1776)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1777)..(1782)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1782)..(1841)

<223> *leader*

<220>

<221> CDS

<222> (1842)..(2210)

<223> *aVEGFv3 VH*

<220>

<221> CDS

<222> (2211)..(2531)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2535)..(2546)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2547)..(2618)

<223> *F2a linker*

<220>

<221> *misc_feature*

<222> (2619)..(2678)

<223> *leader*

<220>

<221> *CDS*

<222> (2679)..(2999)

<223> *aVEGFv3 VL*

<220>

<221> *CDS*

<222> (3000)..(3320)

<223> *CL*

<220>

<221> *polyA_signal*

<222> (3338)..(3569)

<223> *SV40 late polyA*

<220>

<221> *repeat_region*

<222> (3634)..(3763)

<223> AAV2 3'ITR

<400> 19

taaggcctta attaggctgc gcgctcgctc gctcactgag gccgcccggg caaagcccgg
60

gcgctcgggcg acctttggtc gcccggcctc agtgagcgag cgagcgcgca
gagagggagt 120

ggccaactcc atcactaggg gttccttgta gttaatgatt aaccgccat gctacttatc
180

tacgtagcca tgctctagga agatctggcc tccgcgccgg gttttggcgc ctcccgggg
240

cgccccctc ctacggcga gcgctgccac gtcagacgaa gggcgcagcg
agcgtcctga 300

tccttcgcc cggacgctca ggacagcggc ccgctgctca taagactcgg ccttagaacc
360

ccagtatcag cagaaggaca ttttaggacg ggacttgggt gactctaggg cactggtttt
420

*ctttccagag agcggaacag gcgaggaaaa gtagtccctt ctcggcgatt
ctgcggaggg 480*

*atctccgtgg ggcggtgaac gccgatgatt atataaggac gcgccgggtg
tggcacagct 540*

*agttccgtcg cagccgggat ttgggtcgcg gttcttgttt gtggatcgct gtgatcgta
600*

*cttggtgagt agcgggctgc tgggctggcc ggggctttcg tggccgccgg gccgctcgg
660*

*gggacggaag cgtgtggaga gaccgccaag ggctgtagtc tgggtccgcg
agcaaggtg 720*

*ccctgaactg ggggttgggg ggagcgcagc aaaatggcgg ctgttcccga
gtcttgaatg 780*

*gaagacgctt gtgaggcggg ctgtgaggtc gttgaaaca ggtggggggc
atggtgggcg 840*

gcaagaaccc aaggtcttga ggccttcgct aatgcgggaa agctcttatt cgggtgagat
900

gggctggggc accatctggg gaccctgacg tgaagtttg cactgactgg
agaactcgg 960

ttgtcgtctg ttgcgggggc ggcagttatg cggtgccggt gggcagtgca cccgtacctt
1020

tgggagcgcg cgccctcgtc gtgtcgtgac gtcacccggt ctggtggctt ataatgcagg
1080

gtggggccac ctgccggtag gtgtgcggtg ggcttttctc cgtcgcagga cgcagggttc
1140

gggcctaggg taggctctcc tgaatcgaca ggcgccggac ctctggtgag
gggagggata 1200

agtgaggcgt cagtttcttt ggtcggtttt atgtacctat cttcttaagt agctgaagct
1260

ccggttttga actatgcgct cggggttggc gagtgtgttt tgtgaagttt ttaggcacc
1320

ttttgaaatg taatcatttg ggtcaatatg taatcttcag tgtagacta gtaaattgct
1380

cgctaaattc tggccgtttt tggctttttt gtagacgaa gctttattgc gtagtttat
1440

cacagttaaa ttgctaacgc agtcagtgct tctgacacaa cagtctcgaa cttagctgc
1500

agaagttggt cgtgaggcac tgggcaggta agtatcaagg ttacaagaca
ggtttaagga 1560

gaccaataga aactgggctt gtcgagacag agaagactct tgcgtttctg ataggcacct
1620

attggtctta ctgacatcca ctttgccttt ctctccacag gtgtccactc ccagttcaat
1680

tacagctctt aaggctagag tacttaatac gactcactat aggctagcgg gcactttgca
1740

ctggaactta caacacccga gcaaggacgc gactctccac catgtaccgt atgcagctgc
1800

tgctgctgat cgcactgtca ctggcactgg ttaccaactc a gaa gtt cag ctg gtt
1856

Glu Val Gln Leu Val

1 5

gaa tca ggc ggc ggc ctg gtt cag ccc ggc ggc tca ctg cgt ctg tca
1904

Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser

10 15 20

tgt gca gca tca ggc tac gat ttc acc cac tac ggc atg aac tgg gtt
1952

Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr Gly Met Asn Trp Val

25 30 35

cgt cag gca ccc ggc aaa ggc ctg gaa tgg gtt ggc tgg atc aac acc
2000

Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly Trp Ile Asn Thr

40 45 50

tac acc ggc gaa ccc acc tac gca gca gat ttc aaa cgt cgt ttc acc
2048

Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Arg Arg Phe Thr
55 60 65

ttc tca ctg gat acc tca aaa tca acc gca tac ctg cag atg aac tca
2096

Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr Leu Gln Met Asn Ser
70 75 80 85

ctg cgt gca gaa gat acc gca gtt tac tac tgt gca aaa tac ccc tac
2144

Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro Tyr
90 95 100

tac tac ggc acc tca cac tgg tac ttc gat gtt tgg ggc cag ggc acc
2192

Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr
105 110 115

ctg gtt acc gtt tca tca gca tca acc aaa ggc ccc tca gtt ttc ccc 2240

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro

120 125 130

ctg gca ccc tca tca aaa tca acc tca ggc ggc acc gca gca ctg ggc
2288

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly

135 140 145

tgt ctg gtt aaa gat tac ttc ccc gaa ccc gtt acc gtt tca tgg aac 2336

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn

150 155 160 165

tca ggc gca ctg acc tca ggc gtt cac acc ttc ccc gca gtt ctg cag 2384

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln

170 175 180

tca tca ggc ctg tac tca ctg tca tca gtt gtt acc gtt ccc tca tca 2432

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser

185 190 195

tca ctg ggc acc cag acc tac atc tgt aac gtt aac cac aaa ccc tca
2480

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser

200 205 210

aac acc aaa gtt gat aaa aaa gtt gaa ccc aaa tca tgt gat aaa acc
2528

Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr

215 220 225

cac ctgctgtaaac gtcgtgcacc cgtaaacag accctgaact tcgatctgct
2581

His

230

gaaactggca ggcgatgttg aatcaaacc cggcccatg taccgtatgc agctgctgct
2641

gctgatcgca ctgtcactgg cactggttac caactca gat atc cag ctg acc cag
2696

Asp Ile Gln Leu Thr Gln

235

tca ccc tca tca ctg tca gca tca gtt ggc gat cgt gtt acc atc acc 2744

Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr

240 245 250

tgt tca gca tca cag gat atc tca aac tac ctg aac tgg tac cag cag
2792

Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln

255 260 265

aaa ccc ggc aaa gca ccc aaa gtt ctg atc tac ttc acc tca tca ctg
2840

Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Phe Thr Ser Ser Leu

270 275 280

cac tca ggc gtt ccc tca cgt ttc tca ggc tca ggc tca ggc acc gat 2888

His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp

285 290 295 300

ttc acc ctg acc atc tca tca ctg cag ccc gaa gat ttc gca acc tac 2936

Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr

305 310 315

tac tgt cag cag tac tca acc gtt ccc tgg acc ttc ggc cag ggc acc 2984

Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp Thr Phe Gly Gln Gly Thr

320 325 330

aaa gtt gaa atc aaa cgt acc gtt gca gca ccc tca gtt ttc atc ttc 3032

Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe

335 340 345

ccc ccc tca gat gaa cag ctg aaa tca ggc acc gca tca gtt gtt tgt
3080

Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys

350 355 360

ctg ctg aac aac ttc tac ccc cgt gaa gca aaa gtt cag tgg aaa gtt
3128

Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val

365 370 375 380

gat aac gca ctg cag tca ggc aac tca cag gaa tca gtt acc gaa cag
3176

Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln

385 390 395

gat tca aaa gat tca acc tac tca ctg tca tca acc ctg acc ctg tca 3224

Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser

400

405

410

aaa gca gat tac gaa aaa cac aaa gtt tac gca tgt gaa gtt acc cac

3272

Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His

415

420

425

cag ggc ctg tca tca ccc gtt acc aaa tca ttc aac cgt ggc gaa tgt

3320

Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

430

435

440

tgataaagcg gccgcttcga gcagacatga taagatacat tgatgagttt

ggacaaacca 3380

caactagaat gcagtgaaaa aaatgcttta tttgtgaaat ttgtgatgct attgctttat

3440

ttgtaacat tataagctgc aataaacaag ttaacaacaa caattgcatt cattttatgt

3500

*ttcaggttca gggggagatg tgggaggttt tttaaagcaa gtaaacctc
tacaaatgtg 3560*

*gtaaaatcga taaggatctt cctagagcat ggctacgtag ataagtagca
tggcgggtta 3620*

*atcattaact acaaggaacc cctagtgatg gagttggcca ctccctctct gcgcgctcgc
3680*

*tcgctcactg aggccgggcg accaaaggtc gcccgacgcc cgggctttgc
ccgggcggcc 3740*

tcagtgagcg agcgagcgcg cag 3763

<210> 20

<211> 123

<212> PRT

<213> Artificial Sequence

<220>

<223> *Synthetic Construct*

<400> 20

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr

20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe

50 55 60

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr

100

65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

85 90 95

Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val

100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser

115 120

<210> 21

<211> 107

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 21

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys

1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr

20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser

35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser

50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr

65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

85 90 95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His

100 105

<210> 22

<211> 107

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 22

Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr

20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile

35 40 45

Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp

85

90

95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys

100

105

<210> 23

<211> 107

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 23

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu

1

5

10

15

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe

20 25 30

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln

35 40 45

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser

50 55 60

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu

65 70 75 80

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser

85 90 95

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

100

105

<210> 24

<211> 3764

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.UbC.PI.alphaVEGFv1.SV40.ITR*

<220>

<221> *repeat_region*

<222> (17)..(146)

<223> *ITR*

<220>

<221> *promoter*

<222> (207)..(1435)

<223> *UbC*

<220>

<221> *Intron*

<222> (1529)..(1661)

<223> *Promoga chimeric intron*

<220>

<221> *5'UTR*

<222> (1730)..(1777)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1778)..(1783)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1783)..(1842)

<223> *leader*

<220>

<221> CDS

<222> (1843)..(2211)

<223> *aVEGFv1 VH*

<220>

<221> CDS

<222> (2212)..(2532)

<223> CH1

<220>

<221> *misc_feature*

<222> (2536)..(2547)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2548)..(2619)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2620)..(2679)

<223> *leader*

<220>

<221> *CDS*

<222> (2680)..(3000)

<223> *aVEGFv1 VL*

<220>

<221> *CDS*

<222> (3001)..(3321)

<223> *CL*

<220>

<221> *polyA_signal*

<222> (3339)..(3570)

<223> *SV40 polyadenylation signal*

<220>

<221> *repeat_region*

<222> (3635)..(3764)

<223> ITR

<400> 24

taaggcctta attaggctgc gcgctcgctc gctcactgag gccgcccggg caaagcccgg
60

gcgctcgggcg accttggtc gcccggcctc agtgagcgag cgagcgcgca
gagagggagt 120

ggccaactcc atcactaggg gttccttgta gttaatgatt aaccgccat gctactatc
180

tacgtagcca tgctctagga agatctggcc tccgcgccgg gttttggcgc ctcccgcggg
240

cgccccctc ctacggcga gcgctgccac gtcagacgaa gggcgcagcg
agcgtcctga 300

tccttccgcc cggacgctca ggacagcggc ccgctgctca taagactcgg ccttagaacc
360

ccagtatcag cagaaggaca ttttaggacg ggacttgggt gactctaggg cactggtttt
420

ctttccagag agcggaacag gcgaggaaaa gtagtccctt ctcggcgatt
ctgcggaggg 480

atctccgtgg ggcggtgaac gccgatgatt atataaggac gcgccgggtg
tggcacagct 540

agttccgtcg cagccgggat ttgggtcgcg gttcttgttt gtggatcgct gtgatcgta
600

cttggtgagt agcgggctgc tgggctggcc ggggctttcg tggccgccgg gccgctcgg
660

gggacggaag cgtgtggaga gaccgccaag ggctgtagtc tgggtccgcg
agcaaggttg 720

ccctgaactg ggggttgggg ggagcgcagc aaaatggcgg ctgttcccga
gtcttgaatg 780

gaagacgctt gtgaggcggg ctgtgaggtc gttgaaaca ggtggggggc
atggtgggcg 840

gcaagaaccc aaggtcttga ggccttcgct aatgcgggaa agctcttatt cgggtgagat
900

gggctggggc accatctggg gaccctgacg tgaagtttgt cactgactgg
agaactcggg 960

ttgtcgtctg ttgcgggggc ggcagttatg gcggtgccgt tgggcagtgc acccgtacct
1020

ttgggagcgc gcgccctcgt cgtgtcgtga cgtcacccgt tctgttggct tataatgcag
1080

ggtggggcca cctgccggta ggtgtgcggt aggcttttct ccgtcgcagg acgcagggtt
1140

cgggcctagg gtaggctctc ctgaatcgac aggcgccgga cctctggtga
ggggagggat 1200

aagtgaggcg tcagtttctt tggtcggttt tatgtacctt tcttcttaag tagctgaagc
1260

tccggttttg aactatgcgc tcggggttgg cgagtgtggt ttgtgaagtt ttttaggcac
1320

cttttgaat gtaatcattt gggccaatat gtaattttca gtgtagact agtaaattgt
1380

ccgctaaatt ctggccggtt ttggcttttt tgtagacga agctttattg cggtagtta
1440

tcacagttaa attgctaacg cagtcagtcg ttctgacaca acagtctcga acttaagctg
1500

cagaagttgg tcgtgaggca ctgggcaggt aagtatcaag gttacaagac
aggtttaagg 1560

agaccaatag aaactgggct tgtcgagaca gagaagactc ttgcgtttct
gataggcacc 1620

tattggtctt actgacatcc actttgcctt tctctccaca ggtgtccact cccagttcaa
1680

ttacagctct taaggctaga gtacttaata cgactcacta taggctagcg ggcactttgc
1740

*actggaactt acaacacccg agcaaggacg cgactctcca ccatgtaccg
aatgcagctg 1800*

*ctgctgctga tcgcactgag cctggcactg gtgaccaaca gc gag gtg cag ctg
1854*

Glu Val Gln Leu

1

*gtg gag agc gga gga gga ctg gtg cag cca gga gga agc ctg cga ctg
1902*

Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu

5 10 15 20

*agc tgt gca gca agc gga tac gac ttc acc cat tac gga atg aac tgg
1950*

Ser Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr Gly Met Asn Trp

25 30 35

*gtg cga cag gca cca gga aag gga ctg gag tgg gtg gga tgg atc aac
1998*

Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly Trp Ile Asn

40 45 50

acc tac acc gga gag cca acc tac gca gca gac ttc aag cga cga ttc
2046

Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe Lys Arg Arg Phe

55 60 65

acc ttc agc ctg gac acc agc aag agc acc gca tac ctg cag atg aac
2094

Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr Leu Gln Met Asn

70 75 80

agc ctg cga gca gag gac acc gca gtg tac tac tgt gca aag tac cca
2142

Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Lys Tyr Pro

85 90 95 100

tac tac tac gga acc agc cat tgg tac ttc gac gtg tgg gga cag gga
2190

Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val Trp Gly Gln Gly

105 110 115

acc ctg gtg acc gtg agc agc gca agc acc aag gga cca agc gtg ttc
2238

Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe

120 125 130

cca ctg gca cca agc agc aag agc acc agc gga gga acc gca gca ctg
2286

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu

135 140 145

gga tgt ctg gtg aag gac tac ttc cca gag cca gtg acc gtg agc tgg
2334

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp

150 155 160

aac agc gga gca ctg acc agc gga gtg cat acc ttc cca gca gtg ctg
2382

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu

165 170 175 180

cag agc agc gga ctg tac agc ctg agc agc gtg gtg acc gtg cca agc
2430

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser

185

190

195

agc agc ctg gga acc cag acc tac atc tgt aac gtg aac cat aag cca
2478

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro

200

205

210

agc aac acc aag gtg gac aag aag gtg gag cca aag agc tgt gac aag
2526

Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys

215

220

225

acc cat ctgcgaaagc gacgagcacc agtgaagcag accctgaact tcgacctgct
2582

Thr His

230

gaagctggca ggagacgtgg agagcaaccc aggaccaatg taccgaatgc
agctgctgct 2642

gctgatcgca ctgagcctgg cactggtgac caacagc gac atc cag ctg acc cag
2697

Asp Ile Gln Leu Thr Gln

235

agc cca agc agc ctg agc gca agc gtg gga gac cga gtg acc atc acc
2745

Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr

240

245

250

tgt agc gca agc cag gac atc agc aac tac ctg aac tgg tac cag cag
2793

Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp Tyr Gln Gln

255

260

265

aag cca gga aag gca cca aag gtg ctg atc tac ttc acc agc agc ctg
2841

Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Phe Thr Ser Ser Leu

270

275

280

cat agc gga gtg cca agc cga ttc agc gga agc gga agc gga acc gac
2889

His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp

285 290 295 300

ttc acc ctg acc atc agc agc ctg cag cca gag gac ttc gca acc tac
2937

Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr

305 310 315

tac tgt cag cag tac agc acc gtg cca tgg acc ttc gga cag gga acc
2985

Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp Thr Phe Gly Gln Gly Thr

320 325 330

aag gtg gag atc aag cga acc gtg gca gca cca agc gtg ttc atc ttc
3033

Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe

335 340 345

cca cca agc gac gag cag ctg aag agc gga acc gca agc gtg gtg tgt
3081

Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys

350 355 360

ctg ctg aac aac ttc tac cca cga gag gca aag gtg cag tgg aag gtg
3129

Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val
365 370 375 380

gac aac gca ctg cag agc gga aac agc cag gag agc gtg acc gag cag
3177

Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln
 385 390 395

gac agc aag gac agc acc tac agc ctg agc agc acc ctg acc ctg agc
3225

Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser
 400 405 410

aag gca gac tac gag aag cat aag gtg tac gca tgt gag gtg acc cat
3273

Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His
 415 420 425

cag gga ctg agc agc cca gtg acc aag agc ttc aac cga gga gag tgt
3321

Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

430 435 440

tgataaagcg gccgcttcga gcagacatga taagatacat tgatgagttt
ggacaaacca 3381

caactagaat gcagtgaaaa aatgcttta tttgtgaaat ttgtgatgct attgctttat
3441

ttgtaacat tataagctgc aataaacaag ttaacaaca caattgcatt cattttatgt
3501

ttcaggttca gggggagatg tgggaggttt tttaaagcaa gtaaacctc
tacaaatgtg 3561

gtaaaatcga taaggatctt cctagagcat ggctacgtag ataagtagca
tggcgggtta 3621

atcattaact acaaggaacc cctagtgatg gagttggcca ctccctctct gcgcgctcgc
3681

tcgctcactg aggccgggcg accaaaggtc gcccgacgcc cgggctttgc
ccgggcggcc 3741

tcagtgagcg agcgagcgcg cag *3764*

<210> 25

<211> 123

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 25

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Asp Phe Thr His Tyr

20 25 30

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35 40 45

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Ala Asp Phe

50 55 60

Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys Ser Thr Ala Tyr

65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

85 90 95

Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp Tyr Phe Asp Val

100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser

115

120

<210> 26

<211> 107

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 26

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys

1

5

10

15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr

20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser

35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser

50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr

65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

85 90 95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His

100 105

<210> 27

<211> 107

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *Synthetic Construct*

<400> 27

Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr

20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile

35 40 45

Tyr Phe Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro

65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp

85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys

100 105

<210> 28

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 28

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu

1 5 10 15

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe

20 25 30

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln

35 40 45

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser

50 55 60

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu

65 70 75 80

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser

85 90 95

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

100 105

<210> 29

<211> 20

<212> PRT

<213> *Homo sapiens*

<400> 29

Met Tyr Arg Met Gln Leu Leu Ser Cys Ile Ala Leu Ser Leu Ala Leu

1 5 10 15

Val Thr Asn Ser

20

<210> 30

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic leader

<400> 30

Met Tyr Arg Met Gln Leu Leu Leu Leu Ile Ala Leu Ser Leu Ala Leu

1 5 10 15

Val Thr Asn Ser

20

<210> 31

<211> 19

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *synthetic leader 2*

<400> 31

Met Arg Met Gln Leu Leu Leu Leu Ile Ala Leu Ser Leu Ala Leu Val

1 5 10 15

Thr Asn Ser

<210> 32

<211> 355

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *derived from encephalomyocarditis virus*

<400> 32

*tatgctagta cgtctctcaa ggataagtaa gtaatattaa ggtacgggag
gtattggaca 60*

*ggccgcaata aaatatcttt attttcatta catctgtgtg ttggtttttt gtgtgaatcg
120*

*atagtactaa catacgctct ccatcaaac aaaacgaaac aaaacaaact
agcaaaatag 180*

gctgtcccca gtgcaagtgc aggtgccaga acatttctct ggcctaactg gccggtacct
240

gagctctagt ttcactttcc ctagtttcac tttccctagt ttcactttcc ctagtttcac
300

tttcctagt ttcactttcc cctcgaggat atcaagatct ggcctcggcg gccag
355

<210> 33

<211> 513

<212> PRT

<213> *Artificial Sequence*

<220>

<223> *aVEGF*

<220>

<221> MISC_FEATURE

<222> (1)..(20)

<223> *leader*

<220>

<221> *MISC_FEATURE*

<222> (21)..(252)

<223> *aVEGF Heavy Chain*

<220>

<221> *MISC_FEATURE*

<222> (280)..(299)

<223> *leader*

<220>

<221> *MISC_FEATURE*

<222> (300)..(513)

<223> *aVEGF Light Chain*

<400> 33

Met Tyr Arg Met Gln Leu Leu Leu Leu Ile Ala Leu Ser Leu Ala Leu

1

5

10

15

Val Thr Asn Ser Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val

20

25

30

Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Tyr Asp

35

40

45

Phe Thr His Tyr Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly

50

55

60

Leu Glu Trp Val Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr

65

70

75

80

Ala Ala Asp Phe Lys Arg Arg Phe Thr Phe Ser Leu Asp Thr Ser Lys

85

90

95

Ser Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala

100

105

110

Val Tyr Tyr Cys Ala Lys Tyr Pro Tyr Tyr Tyr Gly Thr Ser His Trp

115

120

125

Tyr Phe Asp Val Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala

130

135

140

Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser

145

150

155

160

Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe

165

170

175

Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly

180 185 190

Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu

195 200 205

Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr

210 215 220

Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys

225 230 235 240

Val Glu Pro Lys Ser Cys Asp Lys Thr His Leu Arg Lys Arg Arg Ala

245 250 255

Pro Val Lys Gln Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala Gly Asp

260 265 270

Val Glu Ser Asn Pro Gly Pro Met Tyr Arg Met Gln Leu Leu Leu Leu

275 280 285

Ile Ala Leu Ser Leu Ala Leu Val Thr Asn Ser Asp Ile Gln Leu Thr

290 295 300

Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile

305 310 315 320

Thr Cys Ser Ala Ser Gln Asp Ile Ser Asn Tyr Leu Asn Trp Tyr Gln

325 330 335

Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile Tyr Phe Thr Ser Ser

340 345 350

Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr

355 360 365

Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr

370 375 380

Tyr Tyr Cys Gln Gln Tyr Ser Thr Val Pro Trp Thr Phe Gly Gln Gly

385 390 395 400

Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile

405 410 415

Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val

420 425 430

Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys

435

440

445

Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu

450

455

460

Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu

465

470

475

480

Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr

485

490

495

His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu

500

505

510

Cys

<210> 34

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv1.rBG.ITR*

<220>

<221> *repeat_region*

<222> *(1)..(130)*

<223> *5' ITR*

<220>

<221> *promoter*

<222> *(204)..(584)*

<223> *CMV IE promoter*

<220>

<221> *promoter*

<222> (585)..(862)

<223> *CB promoter*

<220>

<221> *TATA_signal*

<222> (836)..(839)

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> *5'UTR*

<222> (1940)..(1987)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1988)..(1993)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2053)..(2421)

<223> *aVEGFv1 VH*

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv1 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *Rabbit globin poly A*

<220>

<221> *misc_feature*

<222> (3785)..(3821)

<223> *part of AAV*

<220>

<221> *repeat_region*

<222> (3822)..(3951)

<223> 3'ITR

<400> 34

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgctc ggcgaccttt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgcagagagg gagtggccaa
ctccatcact 120

aggggttct thtagttaat gattaaccg ccatgctact tatctaccag ggtaatgggg
180

atcctctaga actatagcta gtcgacattg attattgact agtattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgctt ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgcca agtacgcccc ctattgacgt
480

caatgacggt aatggcccc cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcat cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctccccccc ctccccaccc ccaattttgt atttatttat
660

ttttaatta ttttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720

ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

gagcggcggc ctccgaaagt ttccttttat ggcgagggcg cggcggcggc
ggccctataa 840

*aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900*

*cgccgcccgc tcgcgcccgc cgccccggct ctgactgacc gcgttactcc cacaggtgag
960*

*cgggcgggac ggcccttctc ctccgggctg taattagcgc ttggttaat gacggcttgt
1020*

*ttcttttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080*

*gagcggctcg gggggtgcgt gcgtgtgtgt gtgcgtgggg agcgcgcgct
gcggctccgc 1140*

*gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200*

*gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260*

*caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggc 1320*

cgggctgcaa cccccctgc accccctcc ccgagttgct gagcacggcc cggcttcggg
1380

tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcggca 1440

ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggagggggcg 1500

cggcggcccc cggagcggcg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560

atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620

atctgggagg cgccgccgca cccccttag cgggcgcggg gcgaagcgg
gcggcgccgg 1680

caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*ctttttttt ttctacagc tctgggcaa cgtgctggtt attgtgctgt ctcattcattt
1920*

*tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980*

*cgactctcca ccatgtaccg aatgcagctg ctgctgctga tcgactgag cctggcactg
2040*

*gtgaccaaca gcgaggtgca gctggtggag agcggaggag gactggtgca
gccaggagga 2100*

*agcctgacgac tgagctgtgc agcaagcggga tacgacttca cccattacgg
aatgaactgg 2160*

*gtgacgacag caccaggaaa gggactggag tgggtgggat ggatcaaac
ctacaccgga 2220*

*gagccaacct acgcagcaga cttcaagcga cgattcacct tcagcctgga
caccagcaag 2280*

*agcaccgcat acctgcagat gaacagcctg cgagcagagg acaccgcagt
gtactactgt 2340*

*gcaaagtacc catactacta cggaaccagc cattggtact tcgacgtgtg
gggacagggga 2400*

*accctggtga ccgtgagcag cgcaagcacc aagggaccaa gcgtgttccc
actggcacca 2460*

*agcagcaaga gcaccagcgg aggaaccgca gcactgggat gtctggtgaa
ggactacttc 2520*

*ccagagccag tgaccgtgag ctggaacagc ggagcactga ccagcggagt
gcataccttc 2580*

*ccagcagtgc tgcagagcag cggactgtac agcctgagca gcgtggtgac
cgtgccaagc 2640*

*agcagcctgg gaacccagac ctacatctgt aacgtgaacc ataagccaag
caacaccaag 2700*

*gtggacaaga aggtggagcc aaagagctgt gacaagaccc atctgcgaaa
gcgacgagca 2760*

*ccagtgaagc agaccctgaa cttcgacctg ctgaagctgg caggagacgt
ggagagcaac 2820*

*ccaggaccaa tgtaccgaat gcagctgctg ctgctgatcg cactgagcct ggcactggtg
2880*

*accaacagcg acatccagct gacccagagc ccaagcagcc tgagcgcaag
cgtgggagac 2940*

*cgagtgacca tcacctgtag cgcaagccag gacatcagca actacctgaa
ctggtaccag 3000*

*cagaagccag gaaaggcacc aaaggtgctg atctacttca ccagcagcct
gcatagcggg 3060*

*gtgccaagcc gattcagcgg aagcggaagc ggaaccgact tcacctgac
catcagcagc 3120*

ctgcagccag aggacttcgc aacctactac tgtcagcagt acagcaccgt gccatggacc
3180

ttcggacagg gaaccaaggt ggagatcaag cgaaccgtgg cagcaccaag
cgtgttcac 3240

ttcccaccaa gcgacgagca gctgaagagc ggaaccgcaa gcgtggtgtg
tctgctgaac 3300

aacttctacc cacgagaggc aaaggtgcag tggaaggtgg acaacgcact
gcagagcggg 3360

aacagccagg agagcgtgac cgagcaggac agcaaggaca gcacctacag
cctgagcagc 3420

accctgacc tgagcaaggc agactacgag aagcataagg tgtacgcatg
tgaggtgacc 3480

catcagggac tgagcagccc agtgaccaag agcttcaacc gaggagagtg
ttgataaagc 3540

*ggccgcggta cctctagagt cgacccgggc ggcctcgagg acggggtgaa
ctacgcctga 3600*

*ggatccgatc ttttccctc tgccaaaaat tatggggaca tcatgaagcc ccttgagcat
3660*

*ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720*

*tgtctctcac tcggaagcaa ttcgttgatc tgaatttcga ccaccataa taccattac
3780*

*cctggtagat aagtagcatg gcggggttaat cattaactac aaggaacccc
tagtgatgga 3840*

*gttggccact ccctctctgc gcgctcgctc gctcactgag gccgggacgac caaaggctgc
3900*

*ccgacgcccg ggctttgccg gggcggcctc agtgagcgag cgagcgcgca g
3951*

<210> 35

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv4.rBG.ITR*

<220>

<221> *repeat_region*

<222> (1)..(130)

<223> *5'ITR*

<220>

<221> *promoter*

<222> (198)..(579)

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> (279)..(538)

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> *5'UTR*

<222> (1940)..(1987)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1988)..(1993)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2053)..(2421)

<223> *aVEGFv4 VH*

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv4 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *Rabbit globin poly A*

<220>

<221> *repeat_region*

<222> (3822)..(3951)

<223> *3'ITR*

<400> 35

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgctg ggcgacctt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatgggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgccca agtacgcccc ctattgacgt
480

caatgacggg aaatggcccg cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcac cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctcccccc ctccccaccc ccaattttgt atttatttat
660

ttttaatta tttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720

ggcggggcgcg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

gagcggcgcg ctccgaaagt ttcctttat ggcgaggcgg cggcggcggc
ggccctataa 840

aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900

cgccgcccgc tcgcgcccgc cgccccggct ctgactgacc gcgttactcc cacagggtgag
960

cgggcgggac ggcccttctc ctccgggctg taattagcgc ttggttaat gacggcttgt
1020

ttcttttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080

gagcggctcg ggggggtgctg gcgtgtgtgt gtgcgtgggg agcgccgcgt
gcggctccgc 1140

gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200

gcgcgagggg agcgcgcccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260

caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320

cgggctgcaa cccccctgc acccccctcc ccgagttgct gagcacggcc cggcttcggg
1380

tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcggca 1440

ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggagggggcg 1500

*cggcggcccc cggagcgccg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccctctag cgggcgcggg gcgaagcgg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttctctttt ttctacagc tcctgggcaa cgtgctgggtt attgtgctgt ctcattcatt
1920*

*tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980*

*cgactctcca ccatgtacag aatgcagctg ctgctgctga tcgccctgag cctggccctg
2040*

*gtcaccaaca gcgagggtcca gctgggtcgag agcggaggag gactgggtcca
gcccggagga 2100*

*agcctgagac tgagctgctc cgccagcggga tacgacttca cccactacgg
aatgaactgg 2160*

*gtcagacagg cccccggaaa gggactggag tgggtcggat ggatcaacac
ctacaccgga 2220*

*gagcccacct acgccgccga cttcaagaga agattcacct tcagcctgga
caccagcaag 2280*

*agcaccgcct acctgcagat gaacagcctg agagccgagg acaccgccgt
ctactactgc 2340*

*gccaagtacc cctactacta cggaaccagc cactggtact tcgacgtctg
gggacaggga 2400*

accctgggtca ccgtcagcag cgccagcacc aagggaccca gcgtcttccc cctggcccc
2460

agcagcaaga gcaccagcgg aggaaccgcc gccctgggat gcctgggtcaa
ggactacttc 2520

cccgagccccg tcaccgtcag ctggaacagc ggagccctga ccagcggagt
ccacaccttc 2580

cccgccgtcc tgcagagcag cggactgtac agcctgagca gcgtcgtcac cgtccccagc
2640

agcagcctgg gaaccagac ctacatctgc aacgtcaacc acaagcccag
caacaccaag 2700

gtcgacaaga aggtcgagcc caagagctgc gacaagaccc acctgagaaa
gagaagagcc 2760

cccgtaagc agaccctgaa cttcgacctg ctgaagctgg ccggagacgt
cgagagcaac 2820

cccggaccca tgtacagaat gcagctgctg ctgctgatcg ccctgagcct ggccttggtc
2880

accaacagcg acatccagct gacccagagc cccagcagcc tgagcgccag
cgtcggagac 2940

agagtcacca tcacctgcag cgccagccag gacatcagca actacctgaa
ctggtaccag 3000

cagaagcccc gaaaggcccc caaggtcctg atctacttca ccagcagcct
gcacagcgga 3060

gtccccagca gattcagcgg aagcggaagc ggaaccgact tcaccctgac
catcagcagc 3120

ctgcagcccc aggacttcgc cacctactac tgccagcagt acagcaccgt cccctggacc
3180

ttcggacagg gaaccaaggt cgagatcaag agaaccgctg ccgccccag
cgtcttcac 3240

ttcccccca gcgacgagca gctgaagagc ggaaccgcca gcgtcgtctg
cctgctgaac 3300

*aacttctacc ccagagagggc caaggtccag tggaaggtcg acaacgccct
gcagagcgga 3360*

*aacagccagg agagcgtcac cgagcaggac agcaaggaca gcacctacag
cctgagcagc 3420*

*accctgacct tgagcaaggc cgactacgag aagcacaagg tctacgcctg
cgaggtcacc 3480*

*caccagggac tgagcagccc cgtcaccaag agcttcaaca gaggagagtg
ctgataaagc 3540*

*ggccgcggta cctctagagt cgacccgggc ggcctcgagg acggggtgaa
ctacgcctga 3600*

*ggatccgatc ttttccctc tgccaaaaat tatggggaca tcatgaagcc ccttgagcat
3660*

*ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720*

tgtctctcac tcggaagcaa ttcgttgatc tgaatttcga ccaccataa taccattac
3780

cctggtagat aagtagcatg gcgggtaat cattaactac aaggaacccc
tagtgatgga 3840

gttggccact ccctctctgc gcgctcgctc gtcactgag gccgggcgac caaaggtcgc
3900

ccgacgcccg ggctttgccg gggcggcctc agtgagcgag cgagcgcgca g
3951

<210> 36

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv5.rBG.ITR*

<220>

<221> *repeat_region*

<222> (1)..(130)

<223> 5'ITR

<220>

<221> *promoter*

<222> (198)..(579)

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> (279)..(538)

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> *5'UTR*

<222> (1940)..(1987)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1988)..(1993)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2053)..(2421)

<223> *aVEGFv5 VH*

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv5 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *rabbit globin poly A*

<220>

<221> *repeat_region*

<222> *(3822)..(3951)*

<223> *3'ITR*

<400> 36

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgtcg ggcgacctt
60

ggtcgccccg cctcagtgag cgagcgagcg cgcagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgcca agtacgcccc ctattgacgt
480

caatgacggt aatggcccc cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcat cgctattacc atggtcgagg tgagccccac
600

gttctgctt actctcccca tctcccccc ctccccacc ccaatttgt atttattat
660

ttttaatta tttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720

ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

*gagcggcgcg ctccgaaagt ttcctttat ggcgaggcgg cggcggcggc
ggcctataa 840*

*aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttgccccg
tgccccgctc 900*

*cgccgccgcc tcgcgccgcc cgccccggct ctgactgacc gcgttactcc cacagggtgag
960*

*cgggcgggac ggccttctc ctccgggctg taattagcgc ttggttaat gacggcttg
1020*

*ttctttctg tggctgcgtg aaagcctga ggggctccgg gagggccctt tgtgcggggg
1080*

*gagcggctcg gggggtgcgt gcgtgtgtgt gtgcgtgggg agcggcgcgt
gcggctccgc 1140*

*gctgccccgc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200*

*gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260*

*caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320*

*cgggctgcaa cccccctgc accccctcc ccgagttgct gagcacggcc cggcttcggg
1380*

*tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcggca 1440*

*ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggcg 1500*

*cggcggcccc cggagcgccg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccctctag cgggcgcggg gcgaagcggg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttcttcttt ttctacagc tctgggcaa cgtgctggtt attgtgctgt ctcattcattt
1920*

*tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980*

*cgactctcca ccatgtaccg catgcagctt cttcttctta tcgctctttc tcttgctctt
2040*

*gtgaccaatt ctgaagtgca gcttgtggaa tctggcggcg gccttgtgca gccgggcggc
2100*

tctcttcgcc tttcttgccg tgcttctggc tacgacttta cccattacgg catgaattgg
2160

gtgcgccagg ctccgggcaa gggccttgaa tgggtgggct ggatcaatac
ctacaccggc 2220

gaaccgacct acgctgctga cttaagcgc cgctttacct tttctcttga cacctctaag
2280

tctaccgctt accttcagat gaattctctt cgcgctgaag acaccgctgt gtactactgc
2340

gctaagtacc cgactacta cggcacctct cattggtact ttgacgtgtg gggccagggc
2400

acccttgatga ccgtgtcttc tgcttctacc aagggcccgt ctgtgtttcc gcttgctccg
2460

tcttctaagt ctacctctgg cggcaccgct gctcttggct gccttgatga ggactacttt
2520

ccggaaccgg tgaccgtgtc ttggaattct ggcgctctta cctctggcgt gcataccttt
2580

ccggctgtgc ttcagtcttc tggcctttac tctctttctt ctgtggtgac cgtgccgtct
2640

tcttctcttg gcacccagac ctacatctgc aatgtgaatc ataagccgtc taataccaag
2700

gtggacaaga aggtggaacc gaagtcttgc gacaagaccc atcttcgcaa
gcgccgcgct 2760

ccggtgaagc agacccttaa tttgacctt ctttaagcttg ctggcgacgt ggaatctaata
2820

ccgggcccga tgtaccgcat gcagcttctt cttcttatcg ctctttctct tgctcttgtg
2880

accaattctg acatccagct taccagctt ccgtcttctc tttctgcttc tgtgggacgac
2940

cgcgtagacca tcacctgctc tgcttctcag gacatctcta attaccttaa ttggtaccag
3000

cagaagccgg gcaaggctcc gaagggtgctt atctacttta cctcttctct tcattctggc
3060

gtgccgtctc gcttttctgg ctctggctct ggcaccgact ttacccttac catctcttct
3120

cttcagccgg aagactttgc tacctactac tgccagcagt actctaccgt gccgtggacc
3180

tttgccagg gcaccaaggt ggaaatcaag cgcaccgtgg ctgctccgtc tgtgtttatc
3240

tttccgccgt ctgacgaaca gcttaagtct ggcaccgctt ctgtggtgtg ccttcttaat
3300

aattttacc cgcgcgaagc taagggtcag tggaagggtg acaatgctct tcagtctggc
3360

aattctcagg aatctgtgac cgaacaggac tctaaggact ctacctactc tctttcttct
3420

acccttacc ttttaaggc tgactacgaa aagcataagg tgtacgcttg cgaagtgacc
3480

catcagggcc tttcttctcc ggtgaccaag tctttaatc gcggcgaatg ctgataaagc
3540

ggccgcggta cctctagagt cgacccgggc ggcctcgagg acggggtgaa
ctacgcctga 3600

ggatccgata ttttccctc tgccaaaaat tatggggaca tcatgaagcc ccttgagcat
3660

ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720

tgtctctcac tcggaagcaa ttcgttgatc tgaatttcga ccaccataa taccattac
3780

cctggtagat aagtagcatg gcgggtaat cattaactac aaggaacccc
tagtgatgga 3840

gttggccact ccctctctgc gcgctcgctc gctcactgag gccgggacgac caaaggctgc
3900

ccgacgcccc ggctttgccc gggcggcctc agtgagcgag cgagcgcgca g
3951

<210> 37

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv6.rBG.ITR*

<220>

<221> *repeat_region*

<222> *(1)..(130)*

<223> *5'ITR*

<220>

<221> *promoter*

<222> *(198)..(579)*

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> *(279)..(538)*

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> *(582)..(862)*

<223> *CB promoter*

<220>

<221> *Intron*

<222> *(956)..(1928)*

<223> *chicken beta-actin intron*

<220>

<221> *5'UTR*

<222> *(1940)..(1987)*

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2053)..(2421)

<223> *aVEGFv6 VH*

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv6 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *Rabbit globin poly A*

<220>

<221> *repeat_region*

<222> (3822)..(3951)

<223> *3'ITR*

<400> 37

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgctg ggcgacctt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatgggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggccccct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgccca agtacgcccc ctattgacgt
480

caatgacggt aatggccccg cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcac cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctccccccc ctccccaccc ccaattttgt atttatttat
660

ttttaatta tttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720

ggcggggcgcg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

gagcggcgcg ctccgaaagt ttcctttat ggcgaggcgg cggcggcggc
ggccctataa 840

aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc ctcgccccg
tgccccgctc 900

cgccgccgcc tcgcgccgcc cgccccggct ctgactgacc gcgttactcc cacaggtagag
960

cgggcgggac ggcccttctc ctccgggctg taattagcgc ttggttaat gacggcttgt
1020

ttcttttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080

gagcggctcg ggggggtgctg gcgtgtgtgt gtgcgtgggg agcgccgcgt
gcggctccgc 1140

gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200

gcgcgagggg agcgcgccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260

caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320

cgggctgcaa cccccctgc acccccctcc ccgagttgct gagcacggcc cggcttcggg
1380

tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcggca 1440

ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggcg 1500

*cggcggcccc cggagcgccg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccctctag cgggcgcggg gcgaagcgg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttctctttt ttctacagc tcctgggcaa cgtgctggtt attgtgctgt ctcattcatt
1920*

*tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980*

*cgactctcca ccatgtaccg gatgcagctg ctgctgctga tcgccctgag cctggccctg
2040*

*gtgaccaaca gcgaggtgca gctggtggag agcggaggag gactggtgca
gccaggagga 2100*

*agcctgcggc tgagctgcgc cgccagcggga tacgatttca cccactacgg
aatgaactgg 2160*

*gtgcggcagg ccccaggaaa gggactggag tgggtgggat ggatcaacac
ctacaccgga 2220*

*gagccaacct acgccgccga tttcaagcgg cggttcacct tcagcctgga taccagcaag
2280*

*agcaccgcct acctgcagat gaacagcctg cgggccgagg ataccgccgt
gtactactgc 2340*

*gccaagtacc catactacta cggaaccagc cactggtact tcgatgtgtg
gggacaggga 2400*

*accctgggtga ccgtgagcag cgccagcacc aagggaccaa gcgtgttccc
actggcccca 2460*

*agcagcaaga gcaccagcgg aggaaccgcc gccctgggat gcctggtgaa
ggattacttc 2520*

*ccagagccag tgaccgtgag ctggaacagc ggagccctga ccagcggagt
gcacaccttc 2580*

*ccagccgtgc tgcagagcag cggactgtac agcctgagca gcgtggtgac
cgtgccaagc 2640*

*agcagcctgg gaaccagac ctacatctgc aacgtgaacc acaagccaag
caacaccaag 2700*

*gtggataaga aggtggagcc aaagagctgc gataagaccc acctgcggaa
gcggcgggcc 2760*

*ccagtgaagc agaccctgaa cttcgatctg ctgaagctgg ccggagatgt
ggagagcaac 2820*

ccaggaccaa tgtaccggat gcagctgctg ctgctgatcg cctgagcct ggccttggtg
2880

accaacagcg atatccagct gaccagagc ccaagcagcc tgagcgccag
cgtagggagat 2940

cggtgacca tcacctgag cgccagccag gatatacga actacctgaa
ctggtaccag 3000

cagaagccag gaaaggcccc aaaggtgctg atctacttca ccagcagcct
gcacagcggg 3060

gtgccaagcc gggtcagcg aagcgggagc ggaaccgatt tcacctgac
catcagcagc 3120

ctgcagccag aggatctgc cacctactac tgccagcagt acagcaccgt gccatggacc
3180

ttcggacagg gaaccaaggt ggagatcaag cggaccgtgg ccgccccaaag
cgtgttcac 3240

ttccaccaa gcatgagca gctgaagagc ggaaccgcca gcgtggtgtg
cctgctgaac 3300

*aacttctacc cacgggaggc caaggtgcag tggaaggtgg ataacgccct
gcagagcgga 3360*

*aacagccagg agagcgtgac cgagcaggat agcaaggata gcacctacag
cctgagcagc 3420*

*accctgacct tgagcaaggc cgattacgag aagcacaagg tgtacgcctg
cgaggtgacc 3480*

*caccagggac tgagcagccc agtgaccaag agcttcaacc ggggagagtg
ctgataaagc 3540*

*ggccgcggta cctctagagt cgacccgggc ggcctcgagg acggggtgaa
ctacgcctga 3600*

*ggatccgatc ttttccctc tgccaaaaat tatggggaca tcatgaagcc ccttgagcat
3660*

*ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720*

tgtctctcac tcggaagcaa ttcgttgatc tgaatttcga ccaccataa taccattac
3780

cctggtagat aagtagcatg gcgggtaat cattaactac aaggaacccc
tagtgatgga 3840

gttggccact ccctctctgc gcgctcgctc gtcactgag gccgggagac caaaggcgc
3900

ccgacgcccg ggctttgccg gggcggcctc agtgagcgag cgagcgcgca g
3951

<210> 38

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv7.rBG.ITR*

<220>

<221> *repeat_region*

<222> (1)..(130)

<223> 5'ITR

<220>

<221> *promoter*

<222> (198)..(579)

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> (279)..(538)

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> *5'UTR*

<222> (1940)..(1987)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1988)..(1993)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2053)..(2421)

<223> *aVEGFv7 VH*

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv7 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *rabbit globin poly A*

<220>

<221> *repeat_region*

<222> *(3822)..(3951)*

<223> *3'ITR*

<400> 38

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgtcg ggcgaccttt
60

ggtcgccccg cctcagtgag cgagcgagcg cgcagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatgggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgcca agtacgcccc ctattgacgt
480

caatgacggt aatggcccc cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcat cgctattacc atggtcgagg tgagccccac
600

gttctgctt actctcccca tctcccccc ctccccacc ccaattttgt atttattat
660

ttttaatta ttttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720

ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

*gagcggcgcg ctccgaaagt ttcctttat ggcgaggcgg cggcggcggc
ggcctataa 840*

*aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900*

*cgccgcccgc tcgcgcccgc cgccccggct ctgactgacc gcgttactcc cacagggtgag
960*

*cgggcgggac ggccttctc ctccgggctg taattagcgc ttggttaat gacggcttg
1020*

*ttctttctg tggctgcgtg aaagcctga ggggctccgg gagggccctt tgtgcggggg
1080*

*gagcggctcg gggggtgcgt gcgtgtgtgt gtgcgtgggg agcgcgcgt
gcggctccgc 1140*

*gctgccccgc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200*

*gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260*

*caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320*

*cgggctgcaa cccccctgc accccctcc ccgagttgct gagcacggcc cggcttcggg
1380*

*tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcggca 1440*

*ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggcg 1500*

*cggcggcccc cggagcgccg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccctctag cgggcgcggg gcgaagcggg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttcttcttt ttctacagc tctgggcaa cgtgctggtt attgtgctgt ctcattcattt
1920*

*tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacaccccg
agcaaggacg 1980*

*cgactctcca ccatgtaccg gatgcagctg ctgctgctga tcgccctgag cctggccctg
2040*

*gtgaccaaca gcgaggtgca gctggtggag agcggaggag gactggtgca
gccaggagga 2100*

*agcctgcggc tgagctgcgc cgccagcggg tacgatttca cccactacgg
aatgaactgg 2160*

*gtgcggcagg ccccaggaaa gggactggag tgggtgggat ggatcaacac
ctacaccgga 2220*

*gagccaacct acgccgccga tttcaagcgg cggttcacct tcagcctgga taccagcaag
2280*

*agcaccgcct acctgcagat gaacagcctg cgggccgagg ataccgccgt
gtactactgc 2340*

*gccaagtacc catactacta cggaaccagc cactggtact tcgatgtgtg
gggacaggga 2400*

*accctggtga ccgtgagcag cgccagcacc aagggaccaa gcgtgttccc
actggcccca 2460*

*agcagcaaga gcaccagcgg aggaaccgcc gccctgggat gcctggtgaa
ggattacttc 2520*

*ccagagccag tgaccgtgag ctggaacagc ggagccctga ccagcggagt
gcacaccttc 2580*

*ccagccgtgc tgcagagcag cggactgtac agcctgagca gcgtggtgac
cgtgcccaagc 2640*

*agcagcctgg gaaccagac ctacatctgc aacgtgaacc acaagccaag
caacaccaag 2700*

*gtggataaga aggtggagcc aaagagctgc gataagaccc acctgcggaa
gcggcgggcc 2760*

*ccagtgaagc agaccctgaa cttcgatctg ctgaagctgg ccggagatgt
ggagagcaac 2820*

*ccaggaccaa tgtaccggat gcagctgctg ctgctgatcg cctgagcct ggcctggtg
2880*

*accaacagcg atatccagct gaccagagc ccaagcagcc tgagcgccag
cgtgggagat 2940*

*cgggtgacca tcacctgcag cgccagccag gatatcagca actacctgaa
ctggtaccag 3000*

*cagaagccag gaaaggcccc aaaggtgctg atctacttca ccagcagcct
gcacagcgga 3060*

*gtgccaagcc ggttcagcgg aagcggaagc ggaaccgatt tcaccctgac
catcagcagc 3120*

*ctgcagccag aggatttcgc cacctactac tgccagcagt acagcaccgt gccatggacc
3180*

*ttcggacagg gaaccaaggt ggagatcaag cggaccgtgg ccgccccaaag
cgtgttcac 3240*

*ttcccaccaa gcgatgagca gctgaagagc ggaaccgcca gcgtggtgtg
cctgctgaac 3300*

*aacttctacc cacgggaggc caaggtgcag tggaaggtgg ataacgcctt
gcagagcgga 3360*

*aacagccagg agagcgtgac cgagcaggat agcaaggata gcacctacag
cctgagcagc 3420*

*accctgacc tgagcaaggc cgattacgag aagcacaagg tgtacgcctg
cgaggtgacc 3480*

*caccagggac tgagcagccc agtgaccaag agcttcaacc ggggagagtg
ctgataaagc 3540*

*ggccgcggta cctctagagt cgacccgggc ggcctcgagg acggggtgaa
ctacgcctga 3600*

*ggatccgatac ttttccctc tgccaaaaat tatggggaca tcatgaagcc ccttgagcat
3660*

*ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720*

*tgtctctcac tcggaagcaa ttcgttgatc tgaatttcga ccaccataa taccattac
3780*

*cctggtagat aagtagcatg gcgggtaat cattaactac aaggaacccc
tagtgatgga 3840*

*gttggccact ccctctctgc gcgctcgctc gctcactgag gccgggagac caaaggctgc
3900*

ccgacgcccc ggctttgccc gggcggcctc agtgagcgag cgagcgcgca g
3951

<210> 39

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv8.rBG.ITR*

<220>

<221> *repeat_region*

<222> *(1)..(130)*

<223> *5'ITR*

<220>

<221> *promoter*

<222> *(198)..(579)*

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> *(279)..(538)*

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> *(582)..(862)*

<223> *CB promoter*

<220>

<221> *Intron*

<222> *(956)..(1928)*

<223> *chicken beta-actin intron*

<220>

<221> *5'UTR*

<222> *(1940)..(1987)*

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1988)..(1993)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2053)..(2421)

<223> *aVEGFv8 VH*

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv8 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *rabbit globin poly A*

<220>

<221> *repeat_region*

<222> (3822)..(3951)

<223> *3'ITR*

<400> 39

ctgcgcgctc gctcgtcac tgaggccgcc cgggcaaagc ccgggcgctc ggcgaccttt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgagagagg gagtggccaa
ctccatcact 120

aggggttcct thtagttaat gattaacccg ccatgctact tatctaccag ggtaatgggg
180

atcctctaga actatagcta gtcgacattg attattgact agtattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgctt ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgccca agtacgcccc ctattgacgt
480

caatgacggt aaatggcccg cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcac cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctccccccc ctccccaccc ccaattttgt atttatttat
660

ttttaatta ttttgtgcag cgatgggggc gggggggggg ggggggagcg
cgccaggcgg 720

ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

gagcggcggc ctccgaaagt ttccttttat ggcgagggcg cggcggcggc
ggccctataa 840

aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900

cgccgccgcc tcgcgccgcc cgccccggct ctgactgacc gcgttactcc cacaggtgag
960

cgggcgggac ggcccttctc ctccgggctg taattagcgc ttggttaat gacggcttgt
1020

ttcttttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080

gagcggctcg gggggtgcgt gcgtgtgtgt gtgcgtgggg agcgccgcgt
gcggctccgc 1140

gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200

gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260

caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggc 1320

cgggctgcaa cccccctgc accccctcc ccgagttgct gagcacggcc cggcttcggg
1380

*tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcgcca 1440*

*ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggcg 1500*

*cggcgggccc cggagcgccg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccctctag cgggcgcggg gcgaagcgg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860

cttcttcttt ttctacagc tcttgggcaa cgtgctgggt attgtgctgt ctcatcatt
1920

tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980

cgactctcca ccatgtaccg catgcagctg ctgctgctga tcgctctgtc actggctctg
2040

gtgaccaatt cagaagtgca gctggtggaa tcaggtggtg gtctggtgca
gcccggtggt 2100

tcactgcgcc tgtcatgcgc tgcttcaggt tacgacttta cccattacgg tatgaattgg
2160

gtgcgccagg ctcccggtaa gggctctggaa tgggtgggtt ggatcaatac
ctacaccggt 2220

gaaccacct acgctgctga cttaagcgc cgctttacct tttactgga cacctcaag
2280

tcaaccgctt acctgcagat gaattcactg cgcgctgaag acaccgctgt gtactactgc
2340

gctaagtacc cctactacta cggtaacctca cattggtact ttgacgtgtg gggtcagggt
2400

accctgggtga ccgtgtcatc agcttcaacc aagggtcctt cagtgtttcc cctggctccc
2460

tcatcaaagt caacctcagg tggtaaccgct gctctggggtt gcctgggtgaa ggactacttt
2520

cccgaacccg tgaccgtgtc atggaattca ggtgctctga cctcaggtgt gcataccttt
2580

cccgctgtgc tgcagtcatc aggtctgtac tcaactgtcat cagtgggtgac cgtgccctca
2640

tcatcactgg gtacccagac ctacatctgc aatgtgaatc ataagccctc aaataccaag
2700

*gtggacaaga aggtggaacc caagtcatgc gacaagacc atctgcgcaa
gcgccgcgct 2760*

*cccgtgaagc agaccctgaa tttgacctg ctgaagctgg ctggtgacgt ggaatcaaat
2820*

*cccgggccca tgtaccgcat gcagctgctg ctgctgatcg ctctgtcact ggctctggtg
2880*

*accaattcag acatccagct gaccagtca ccctcatcac tgtcagcttc agtgggtgac
2940*

*gcgctgacca tcacctgctc agcttcacag gacatctcaa attacctgaa ttggtaccag
3000*

*cagaagcccc gtaaggctcc caaggctgctg atctacttta cctcatcact gcattcaggt
3060*

*gtgccctcac gcttttcagg ttcaggttca ggtaccgact ttaccctgac catctcatca
3120*

*ctgcagcccc aagactttgc tacctactac tgccagcagt actcaaccgt gccctggacc
3180*

tttggtcagg gtaccaaggt ggaaatcaag cgcaccgtgg ctgctccctc agtgtttacc
3240

tttccccct cagacgaaca gctgaagtca ggtaccgctt cagtgggtgtg cctgctgaat
3300

aatthttacc cccgcgaagc taaggtgcag tggaaggtgg acaatgctct gcagtcaggt
3360

aattcacagg aatcagtgac cgaacaggac tcaaaggact caacctactc
actgtcatca 3420

accctgacct tgtcaaaggc tgactacgaa aagcataagg tgtacgcttg
cgaagtgacc 3480

catcagggtc tgtcatcacc cgtgaccaag tcatttaatc gcggtgaatg ctgataaagc
3540

ggccgcggta cctctagagt cgacccgggc ggcctcgagg acgggggtgaa
ctacgcctga 3600

ggatccgatac ttttccctc tgccaaaaat tatggggaca tcatgaagcc ccttgagcat
3660

ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720

tgtctctcac tcggaagcaa ttcgttgatac tgaatttcga ccaccataa taccattac
3780

cctggtagat aagtagcatg gcgggtaat cattaactac aaggaacccc
tagtgatgga 3840

gttggccact ccctctctgc gcgctcgtc gctcactgag gccgggacgac caaaggctcg
3900

ccgacgcccg ggctttgccc gggcggcctc agtgagcgag cgagcgcgca g
3951

<210> 40

<211> 3957

<212> DNA

<213> Artificial Sequence

<220>

<223> *ITR.CB7.Cl.alphaVEGFv9.rBG.ITR*

<220>

<221> *repeat_region*

<222> *(1)..(130)*

<223> *5'ITR*

<220>

<221> *promoter*

<222> *(198)..(579)*

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> *(279)..(538)*

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> *5'UTR*

<222> (1946)..(1993)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1994)..(1999)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1999)..(2058)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2059)..(2427)

<223> *aVEGFv9 VH*

<220>

<221> *misc_feature*

<222> (2428)..(2748)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2752)..(2763)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2764)..(2835)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2836)..(2895)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2896)..(3216)

<223> *aVEGFv9 VL*

<220>

<221> *misc_feature*

<222> (3217)..(3537)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3538)..(3543)

<223> *Stop Cassette*

<220>

<221> *polyA_signal*

<222> (3613)..(3739)

<223> *rabbit globin poly A*

<220>

<221> *repeat_region*

<222> (3828)..(3957)

<223> *3'ITR*

<400> 40

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgctc ggcgacctt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgctt ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgcca agtacgcccc ctattgacgt
480

caatgacggt aatggcccc cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcat cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctccccccc ctccccaccc ccaattttgt atttatttat
660

*ttttaatta tttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720*

*ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780*

*gagcggcgcg ctccgaaagt ttcctttat ggcgaggcgg cggcggcggc
ggccctataa 840*

*aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900*

*cgccgccgcc tcgcgccgcc cgccccggct ctgactgacc gcgttactcc cacaggtgag
960*

*cgggcgggac ggccttctc ctccgggctg taattagcgc ttggttaat gacggcttgt
1020*

*ttctttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080*

*gagcggctcg gggggtgctg gcgtgtgtgt gtgctgggg agcggcgcgt
gcggctccgc 1140*

*gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg cttgtgctg tccgcagtgt
1200*

*gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260*

*caaaggctgc gtgcggggtg tgtgctggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320*

*cgggctgcaa cccccctgc acccccctcc ccgagttgct gagcacggcc cggcttcggg
1380*

*tgcggggctc cgtacggggc gtggcgcggg gctcggcgtg ccgggcgggg
ggtggcggca 1440*

*ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggca 1500*

*cggcggcccc cggagcggcc gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccctctag cgggcgcggg gcgaagcgg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttcttcttt ttctacagc tcctgggcaa cgtgctggtt attgtgctgt ctcattcatt
1920*

*tggcaaagaa ttcgctagag ctagcgggca cttgcactg gaacttaca
cacccgagca 1980*

aggacgcgac tctccacat gtatagaatg cagcttcttc ttcttatcgc tcttagcctt
2040

gctcttgta caaacagcga agtgcagctt gtggaaagcg ggggggggct
tgtgcagcct 2100

ggggggagcc ttagacttag ctgtgctgct agcgggatg attttacaca ctatgggatg
2160

aactgggtga gacaggctcc tgggaagggg cttgaatggg tggggtgat
caacacatat 2220

acaggggaac ctacatatgc tgctgatttt aagagaagat ttacatttag ccttgataca
2280

agcaagagca cagcttatct tcagatgaac agccttagag ctgaagatac
agctgtgtat 2340

tatttgcta agtatcctta ttattatggg acaagccact ggtattttga tgtgtggggg
2400

caggggacac ttgtgacagt gagcagcgt agcacaaggg ggcctagcgt gtttcctctt
2460

*gctcctagca gcaagagcac aagcgggggg acagctgctc ttgggtgtct
tgtgaaggat 2520*

*tattttcctg aacctgtgac agtgagctgg aacagcgggg ctcttacaag cggggtgcac
2580*

*acatttcctg ctgtgcttca gagcagcggg ctttatagcc ttagcagcgt ggtgacagtg
2640*

*cctagcagca gccttgggac acagacatat atctgtaacg tgaaccacaa
gcctagcaac 2700*

*acaaaggtgg ataagaaggt ggaacctaag agctgtgata agacacacct
tagaaagaga 2760*

*agagctcctg tgaagcagac acttaacttt gatcttctta agcttgctgg ggatgtggaa
2820*

*agcaaccctg ggccatgta tagaatgcag cttcttcttc ttatcgctct tagccttgct
2880*

cttgtgacaa acagcgatat ccagcttaca cagagcccta gcagccttag cgctagcgtg
2940

ggggatagag tgacaatcac atgtagcgct agccaggata tcagcaacta
tcttaactgg 3000

tatcagcaga agcctgggaa ggctcctaag gtgcttatct atttacaag cagccttcac
3060

agcggggtgc ctagcagatt tagcgggagc gggagcggga cagattttac
acttacaatc 3120

agcagccttc agcctgaaga ttttgctaca tattattgtc agcagtatag cacagtgctt
3180

tggacatttg ggcaggggac aaaggtggaa atcaagagaa cagtggctgc
tcctagcgtg 3240

tttatctttc ctctagcga tgaacagctt aagagcggga cagctagcgt ggtgtgtctt
3300

cttaacaact tttatcctag agaagctaag gtgcagtgga aggtggataa cgctcttcag
3360

*agcgggaaca gccaggaag cgtgacagaa caggatagca aggatagcac
atatagcctt 3420*

*agcagcacac ttacacttag caaggctgat tatgaaaagc acaagggtga
tgcttgtaa 3480*

*gtgacacacc aggggcttag cagccctgtg acaaagagct ttaacagagg
ggaatggtga 3540*

*taaagcggcc gcggtacctc tagagtcgac ccgggcggcc tcgaggacgg
ggtgaactac 3600*

*gcctgaggat ccgatctttt tcctctgcc aaaaattatg gggacatcat gaagcccctt
3660*

*gagcatctga cttctggcta ataaaggaaa tttattttca ttgcaatagt gtgttggaat
3720*

*ttttgtgtc tctcactcgg aagcaattcg ttgatctgaa tttcgaccac ccataatacc
3780*

cattaccctg gtagataagt agcatggcgg gttaatcatt aactacaagg aaccctagt
3840

gatggagttg gccactccct ctctgcgcgc tcgctcgctc actgaggccg ggcgaccaa
3900

ggtcgcccga cgcccgggct ttgcccgggc ggcctcagtg agcgagcgag cgcgcag
3957

<210> 41

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv10.rBG.ITR*

<220>

<221> *repeat_region*

<222> (1)..(130)

<223> 5'ITR

<220>

<221> *promoter*

<222> (198)..(579)

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> (279)..(538)

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> 5'UTR

<222> (1940)..(1987)

<223> c-myc 5'UTR

<220>

<221> misc_feature

<222> (1988)..(1993)

<223> kozak

<220>

<221> misc_feature

<222> (1993)..(2052)

<223> leader

<220>

<221> misc_feature

<222> (2053)..(2421)

<223> aVEGFv10 VH

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv10 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *rabbit globin poly A*

<220>

<221> *repeat_region*

<222> (3822)..(3951)

<223> 3'ITR

<400> 41

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgtcg ggcgaccttt
60

ggtcgccccg cctcagtgag cgagcgagcg cgcagagagg gagtggccaa
ctccatcact 120

aggggttcct thtagttaat gattaacccg ccatgctact tatctaccag ggtaatgggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacgggggtca ttagttcata gccatataat ggagttccgc gttacataac ttacggtaaa
300

tggccccgct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgcca agtacgcccc ctattgacgt
480

caatgacggg aaatggcccc cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcat cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctccccccc ctccccacc ccaattttgt atttatttat
660

ttttaatta ttttgtgcag cgatgggggc gggggggggg ggggggagcg
cgccaggcgg 720

ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

gagcggcggc ctccgaaagt ttcttttat ggcgagggcg cggcggcggc
ggcctataa 840

*aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900*

*cgccgccgcc tcgcgccgcc cgccccggct ctgactgacc gcgttactcc cacaggtagag
960*

*cgggcgggac ggcccttctc ctccgggctg taattagcgc ttggttaat gacggcttg
1020*

*ttcttttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080*

*gagcggctcg gggggtgcgt gcgtgtgtgt gtgcgtgggg agcgcgcgct
gcggctccgc 1140*

*gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200*

*gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260*

*caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320*

*cgggctgcaa cccccctgc acccccctcc ccgagttgct gagcacggcc cggcttcggg
1380*

*tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcgga 1440*

*ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggcg 1500*

*cggcggcccc cggagcgccg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca ccccctctag cgggcgcggg gcgaagcggg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttcttcttt ttctacagc tctgggcaa cgtgctggtt attgtgctgt ctcattcatt
1920*

*tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980*

*cgactctcca ccatgtatcg gatgcaactg ctgctgctga ttgccctgag cctggccctg
2040*

*gttaccaaca gcgaagtca actggttgaa agcggcggcg gcctggttca
acccggcggc 2100*

*agcctgcggc tgagctgtgc cgccagcggc tatgacttta cccattatgg catgaactgg
2160*

*gttcggcaag cccccggcaa gggcctggaa tgggttggt ggattaacac
ctataccggc 2220*

*gaaccacct atgccgccga cttaagcgg cggtttacct ttagcctgga caccagcaag
2280*

*agcaccgcct atctgcaaat gaacagcctg cgggccgaag acaccgccgt ttattattgt
2340*

*gccaagtatc cctattatta tggcaccagc cattggtatt ttgacgtttg gggccaaggc
2400*

*accctggta ccgtagcag cgccagcacc aagggcccca gcgttttcc cctggcccc
2460*

*agcagcaaga gcaccagcgg cggcaccgcc gccctgggct gtctggtaa
ggactatfff 2520*

*cccgaaccg ttaccgtag ctggaacagc ggcgccctga ccagcggcgt tcatacfff
2580*

*cccgccgttc tgcaaagcag cggcctgtat agcctgagca gcgttggtac cgttcccagc
2640*

*agcagcctgg gcacccaac ctatattgt aacgtaacc ataagcccag
caacaccaag 2700*

*gttgacaaga aggttgaacc caagagctgt gacaagacc atctgcggaa
gcggcgggcc 2760*

*cccgtaagc aaaccctgaa cttgacctg ctgaagctgg ccggcgacgt
tgaaagcaac 2820*

*ccggcccca tgtatcggat gcaactgctg ctgctgattg ccctgagcct ggcctggt
2880*

*accaacagcg acattcaact gacccaagc cccagcagcc tgagcgccag
cgttggcgac 2940*

*cggttacca ttacctgtag cgccagccaa gacattagca actatctgaa ctggtatcaa
3000*

*caaaagcccg gcaaggcccc caaggttctg atttatttta ccagcagcct gcatagcggc
3060*

gttcccagcc ggtttagcgg cagcggcagc ggcaccgact ttaccctgac cattagcagc
3120

ctgcaacccg aagactttgc cacctattat tgtcaacaat atagcaccgt tcctggacc
3180

tttgccaag gcaccaaggt tgaattaag cggaccgttg ccgccccag cgtttttatt
3240

ttcccccca gcgacgaaca actgaagagc ggcaccgcca gcgttgttg tctgctgaac
3300

aacttttatc cccgggaagc caaggtcaa tggaaggttg acaacgcctt
gcaaacgaggc 3360

aacagccaag aaagcggtac cgaacaagac agcaaggaca gcacctatag
cctgagcagc 3420

accctgacc tgagcaaggc cgactatgaa aagcataagg ttatgcctg tgaagttacc
3480

catcaaggcc tgagcagccc cgttaccaag agctttaacc ggggcgaatg
ttgataaagc 3540

*ggccgcggta cctctagagt cgacccgggc ggcctcgagg acggggtgaa
ctacgcctga 3600*

*ggatccgata ttttccctc tgccaaaaat tatggggaca tcatgaagcc cttgagcat
3660*

*ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720*

*tgtctctcac tcggaagcaa ttcgttgata tgaatttcga ccaccataa taccattac
3780*

*cctggtagat aagtagcatg gcgggtaat cattaactac aaggaacccc
tagtgatgga 3840*

*gttgccact ccctctctgc gcgctcgtc gtcactgag gccgggagac caaaggctgc
3900*

*ccgacgcccg ggctttgcc gggcggcctc agtgagcgag cgagcgcgca g
3951*

<210> 42

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv11.rBG.ITR*

<220>

<221> *repeat_region*

<222> (1)..(130)

<223> *5'ITR*

<220>

<221> *promoter*

<222> (198)..(579)

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> (279)..(538)

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> *5'UTR*

<222> (1940)..(1987)

<223> *c-myc 5' UTR*

<220>

<221> *misc_feature*

<222> (1988)..(1993)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2053)..(2421)

<223> *aVEGFv11 VH*

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2754)

<223> *furing cleavage site*

<220>

<221> *misc_feature*

<222> (2755)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv11 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *rabbit globin poly A*

<220>

<221> *repeat_region*

<222> (3822)..(3951)

<223> *3' ITR*

<400> 42

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgctc ggcgacctt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgcagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatgggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgccca agtacgcccc ctattgacgt
480

caatgacggt aatggcccg cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcac cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctccccccc ctccccaccc ccaattttgt atttatttat
660

ttttaatta tttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720

ggcggggcgcg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

gagcggcgcg ctccgaaagt ttccttttat ggcgaggcgg cggcggcggc
ggccctataa 840

aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900

cgccgccgcc tcgcgccgcc cgccccggct ctgactgacc gcgttactcc cacaggtgag
960

cgggcgggac ggcccttctc ctccgggctg taattagcgc ttggttaat gacggcttgt
1020

ttcttttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080

gagcggctcg ggggggtgctg gcgtgtgtgt gtgcgtgggg agcgccgcgt
gcggctccgc 1140

gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200

gcgcgagggg agcgcgcccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260

caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320

cgggctgcaa cccccctgc acccccctcc ccgagttgct gagcacggcc cggcttcggg
1380

tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcggca 1440

ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggcg 1500

*cggcggcccc cggagcgccg gcggctgtcg aggcgcggcg agccgcagcc
attgccttt 1560*

*atggaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccctctag cgggcgcggg gcgaagcgg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttctcttt ttctacagc tcctgggcaa cgtgctggtt attgtgctgt ctcattcatt
1920*

*tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980*

*cgactctcca ccatgtatag aatgcagctg ctgctgctga tcgctctgtc actggctctg
2040*

*gtgaccaact cagaggtgca gctggtggag tcaggaggag gactggtgca
gcccggagga 2100*

*tcactgagac tgtcatgcbc tgcttcagga tatgatttca cccattatgg aatgaactgg
2160*

*gtgagacagg ctcccggaaa aggactggag tgggtgggat ggatcaacac
ctataccgga 2220*

*gagcccacct atgctgctga tttcaaaaga agattcacct tctcactgga tacctcaaaa
2280*

*tcaaccgctt atctgcagat gaactcactg agagctgagg ataccgctgt gtattattgc
2340*

*gctaaatata cctattatta tggaacctca cattggtatt tcgatgtgtg gggacagggg
2400*

accctgggtga ccgtgtcatc agcttcaacc aaaggaccct cagtgttccc cctggctccc
2460

tcatcaaaat caacctcagg aggaaccgct gctctgggat gcctggtgaa agattatttc
2520

cccgagcccc tgaccgtgtc atggaactca ggagctctga cctcaggagt gcataccttc
2580

cccgtgtgtc tgcagtcatc aggactgtat tcaactgtcat cagtggtgac cgtgccctca
2640

tcatcactgg gaaccagac ctatatctgc aacgtgaacc ataaaccctc
aaacaccaaa 2700

gtggataaaa aagtgagacc caaatcatgc gataaaacc atctgagaaa
aagaagagct 2760

cccgtgaaac agaccctgaa cttcgatctg ctgaaactgg ctggagatgt
ggagtcaaac 2820

cccggaccca tgtatagaat gcagctgctg ctgctgatcg ctctgtcact ggctctggtg
2880

accaactcag atatccagct gacccagtca ccctcatcac tgtcagcttc agtgggagat
2940

agagtgacca tcacctgctc agcttcacag gatattctcaa actatctgaa ctggtatcag
3000

cagaaacccg gaaaagctcc caaagtgctg atctatttca cctcatcact gcattcagga
3060

gtgccctcaa gattctcagg atcaggatca ggaaccgatt tcaccctgac catctcatca
3120

ctgcagcccg aggatttcgc tacctattat tgccagcagt attcaaccgt gccttgacc
3180

ttcggacagg gaaccaaagt ggagatcaaa agaaccgtgg ctgctccctc
agtgttcac 3240

ttccccct cagatgagca gctgaaatca ggaaccgctt cagtgggtg cctgctgaac
3300

*aacttctatc ccagagaggc taaagtgcag tggaaagtgg ataacgctct
gcagtcagga 3360*

*aactcacagg agtcagtgac cgagcaggat tcaaaagatt caacctattc actgtcatca
3420*

*accctgaccc tgtcaaaagc tgattatgag aaacataaag tgtatgcttg cgaggtgacc
3480*

*catcagggac tgtcatcacc cgtgacccaaa tcattcaaca gaggagagtg
ctgataaagc 3540*

*ggccgcggta cctctagagt cgacccgggc ggcctcgagg acggggtgaa
ctacgcctga 3600*

*ggatccgatc ttttccctc tgccaaaaat tatggggaca tcatgaagcc ccttgagcat
3660*

*ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720*

tgtctctcac tcggaagcaa ttcgttgatc tgaatttcga ccaccataa taccattac
3780

cctggtagat aagtagcatg gcgggtaat cattaactac aaggaacccc
tagtgatgga 3840

gttggccact ccctctctgc gcgctcgctc gctcactgag gccgggagac caaaggctcg
3900

ccgacgcccg ggctttgccc gggcggcctc agtgagcgag cgagcgcgca g
3951

<210> 43

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv12.rBG.ITR*

<220>

<221> *repeat_region*

<222> (1)..(130)

<223> 5'ITR

<220>

<221> *promoter*

<222> (198)..(579)

<223> *CMV IE promoter*

<220>

<221> *enhancer*

<222> (279)..(538)

<223> *C4 enhancer*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chicken beta-actin intron*

<220>

<221> *5'UTR*

<222> (1940)..(1987)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1988)..(1993)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2053)..(2421)

<223> *aVEGFv12 VH*

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furing cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv12 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *rabbit globin poly A*

<220>

<221> *repeat_region*

<222> *(3822)..(3951)*

<223> *3'ITR*

<400> 43

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgtcg ggcgacctt
60

ggtcgccccg cctcagtgag cgagcgagcg cgcagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgct ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgcca agtacgcccc ctattgacgt
480

caatgacggt aatggcccc cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcat cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctccccccc ctccccacc ccaattttgt atttatttat
660

ttttaatta ttttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720

ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780

*gagcggcgcg ctccgaaagt ttcctttat ggcgaggcgg cggcggcggc
ggcctataa 840*

*aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900*

*cgccgcccgc tcgcgcccgc cgccccggct ctgactgacc gcgttactcc cacagggtgag
960*

*cgggcgggac ggccttctc ctccgggctg taattagcgc ttggttaat gacggcttg
1020*

*ttctttctg tggctgcgtg aaagcctga ggggctccgg gagggccctt tgtgcggggg
1080*

*gagcggctcg gggggtgcgt gcgtgtgtgt gtgcgtgggg agcggccgct
gcggctccgc 1140*

*gctgccccgc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200*

*gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260*

*caaaggctgc gtgcggggtg tgtgcgtggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320*

*cgggctgcaa cccccctgc accccctcc ccgagttgct gagcacggcc cggcttcggg
1380*

*tgcggggctc cgtacggggc gtggcgcggg gctcgccgtg ccgggcgggg
ggtggcggca 1440*

*ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggcg 1500*

*cggcggcccc cggagcgccg gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccctctag cgggcgcggg gcgaagcggg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttcttcttt ttctacagc tctgggcaa cgtgctggtt attgtgctgt ctcattcattt
1920*

*tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980*

*cgactctcca ccatgtaccg tatgcagctg ctgctgctga tagccctgag cctggccctg
2040*

*gtgacaaata gcgaggtgca gctggtggag agcggggggg ggctggtgca
gcccgggggg 2100*

agcctgcgtc tgagctgtgc cgccagcggg tacgacttta cacattacgg gatgaattgg
2160

gtgcgtcagg cccccgggaa agggctggag tgggtggggg ggataaatac
atacacaggg 2220

gagcccacat acgccgccga ctttaaactg cgttttacat ttagcctgga cacaagcaaa
2280

agcacagcct acctgcagat gaatagcctg cgtgccgagg acacagccgt
gtactactgt 2340

gccaaatacc cctactacta cgggacaagc cattggtact ttgacgtgtg
ggggcagggg 2400

acactggtga cagtgagcag cgccagcaca aaagggccca gcgtgtttcc
cctggccccc 2460

agcagcaaaa gcacaagcgg ggggacagcc gccctggggg gtctggtgaa
agactacttt 2520

cccgagcccg tgacagtgag ctggaatagc ggggccctga caagcggggg
gcatacattt 2580

*cccgccgtgc tgcagagcag cgggctgtac agcctgagca gcgtggtgac
agtgccccagc 2640*

*agcagcctgg ggacacagac atacatatgt aatgtgaatc ataaaccag
caatacaaaa 2700*

*gtggacaaaa aagtggagcc caaaagctgt gacaaaacac atctgcgtaa
acgtcgtgcc 2760*

*cccgtaaac agacactgaa ttttgacctg ctgaaactgg ccggggacgt
ggagagcaat 2820*

*ccggggccca tgtaccgtat gcagctgctg ctgctgatag ccctgagcct ggcctggtg
2880*

*acaatatagcg acatacagct gacacagagc cccagcagcc tgagcgccag
cgtgggggac 2940*

*cggtgacaaa taacatgtag cgccagccag gacataagca attacctgaa
ttggtaccag 3000*

*cagaaacccg ggaaagcccc caaagtgctg atatacttta caagcagcct
gcatagcggg 3060*

*gtgcccagcc gtttagcgg gagcgggagc gggacagact ttactgac
aataagcagc 3120*

*ctgcagcccg aggactttgc cacatactac tgtcagcagt acagcacagt gccctggaca
3180*

*ttgggcagg ggacaaaagt ggagataaaa cgtacagtgg ccgccccag
cgtgtttata 3240*

*ttcccccca gcgacgagca gctgaaaagc gggacagcca gcgtggtgtg
tctgctgaat 3300*

*aatthttacc cccgtgaggc caaagtgcag tggaaagtgg acaatgcct
gcagagcggg 3360*

*aatagccagg agagcgtgac agagcaggac agcaaagaca gcacatacag
cctgagcagc 3420*

*acactgacac tgagcaaagc cgactacgag aaacataaag tgtacgcctg
tgaggtgaca 3480*

*catcaggggc tgagcagccc cgtgacaaaa agctttaatc gtggggagtg
ttgataaagc 3540*

*ggccgcggta cctctagagt cgacccgggc ggcctcgagg acggggtgaa
ctacgcctga 3600*

*ggatccgata ttttccctc tgccaaaaat tatggggaca tcatgaagcc ccttgagcat
3660*

*ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720*

*tgtctctcac tcggaagcaa ttcgttgatc tgaatttcga ccaccataa taccattac
3780*

*cctggtagat aagtagcatg gcgggtaat cattaactac aaggaacccc
tagtgatgga 3840*

*gttggccact ccctctctgc gcgctcgctc gctcactgag gccgggacgac caaaggctcg
3900*

ccgacgcccc ggctttgccc gggcggcctc agtgagcgag cgagcgcgca g
3951

<210> 44

<211> 3951

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CB7.Cl.alphaVEGFv13.rBG.ITR*

<220>

<221> *repeat_region*

<222> *(1)..(130)*

<223> *5'ITR*

<220>

<221> *misc_feature*

<222> *(131)..(167)*

<223> *part of AAV*

<220>

<221> *promoter*

<222> *(198)..(579)*

<223> *CME IE promoter*

<220>

<221> *misc_feature*

<222> *(204)..(233)*

<223> *promoter start*

<220>

<221> *enhancer*

<222> *(279)..(538)*

<223> *C4 enhancer with 2 mismatches*

<220>

<221> *misc_feature*

<222> *(561)..(584)*

<223> *CMV promoter end*

<220>

<221> *promoter*

<222> (582)..(862)

<223> *CB promoter*

<220>

<221> *misc_feature*

<222> (585)..(615)

<223> *begin promoter*

<220>

<221> *TATA_signal*

<222> (836)..(839)

<220>

<221> *Intron*

<222> (956)..(1928)

<223> *chichen beta-actin intron*

<220>

<221> *misc_feature*

<222> (1814)..(1830)

<223> *end of intron*

<220>

<221> *5'UTR*

<222> (1940)..(1987)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1988)..(1993)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1993)..(2052)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2053)..(2421)

<223> *aVEGFv13 VH*

<220>

<221> *misc_feature*

<222> (2422)..(2742)

<223> *CH1*

<220>

<221> *misc_feature*

<222> (2746)..(2757)

<223> *furin cleavage site*

<220>

<221> *misc_feature*

<222> (2758)..(2829)

<223> *F2A linker*

<220>

<221> *misc_feature*

<222> (2830)..(2889)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2890)..(3210)

<223> *aVEGFv13 VL*

<220>

<221> *misc_feature*

<222> (3211)..(3531)

<223> *CL*

<220>

<221> *misc_feature*

<222> (3532)..(3537)

<223> *stop cassette*

<220>

<221> *polyA_signal*

<222> (3607)..(3733)

<223> *rabbit globin poly A*

<220>

<221> *misc_feature*

<222> (3785)..(3821)

<223> *part of AAV*

<220>

<221> *repeat_region*

<222> (3822)..(3951)

<223> *3'ITR*

<400> 44

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgctc ggcgacctt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctaccag ggtaatggg
180

atcctctaga actatagcta gtcgacattg attattgact agttattaat agtaatcaat
240

tacgggggtca ttagttcata gcccatatat ggagttccgc gttacataac ttacggtaaa
300

tggcccgctt ggctgaccgc ccaacgaccc ccgcccattg acgtcaataa tgacgtatgt
360

tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggagt atttacggta
420

aactgcccac ttggcagtac atcaagtgta tcatatgcca agtacgcccc ctattgacgt
480

caatgacggt aatggcccc cctggcatta tgcccagtac atgaccttat gggactttcc
540

tacttggcag tacatctacg tattagtcat cgctattacc atggtcgagg tgagccccac
600

gttctgcttc actctcccca tctcccccc ctccccacc ccaattttgt atttattat
660

*ttttaatta tttgtgcag cgatgggggc gggggggggg ggggggcgcg
cgccaggcgg 720*

*ggcggggcgg ggcgaggggc ggggcggggc gaggcggaga ggtgcggcgg
cagccaatca 780*

*gagcggcgcg ctccgaaagt ttcctttat ggcgaggcgg cggcggcggc
ggccctataa 840*

*aaagcgaagc gcgcggcggg cgggagtcgc tgcgcgctgc cttcgccccg
tgccccgctc 900*

*cgccgccgcc tcgcgccgcc cgccccggct ctgactgacc gcgttactcc cacaggtgag
960*

*cgggcgggac ggcccttctc ctccgggctg taattagcgc ttggttaat gacggcttgt
1020*

*ttctttctg tggctgcgtg aaagccttga ggggctccgg gagggccctt tgtgcggggg
1080*

*gagcggctcg gggggtgctg gcgtgtgtgt gtgctgggg agcggcgcgt
gcggctccgc 1140*

*gctgcccggc ggctgtgagc gctgcgggcg cggcgcgggg ctttgtgcgc tccgcagtgt
1200*

*gcgcgagggg agcgcggccg ggggcggtgc cccgcggtgc ggggggggct
gcgaggggaa 1260*

*caaaggctgc gtgcggggtg tgtgctggg ggggtgagca gggggtgtgg
gcgcgtcggg 1320*

*cgggctgcaa cccccctgc acccccctcc ccgagttgct gagcacggcc cggcttcggg
1380*

*tgcggggctc cgtacggggc gtggcgcggg gctcggcgtg ccgggcgggg
ggtggcggca 1440*

*ggtgggggtg ccgggcgggg cggggccgcc tcgggccggg gagggctcgg
gggaggggag 1500*

*cggcggcccc cggagcggcc gcggctgtcg aggcgcggcg agccgcagcc
attgcctttt 1560*

*atggtaatcg tgcgagaggg cgcagggact tcctttgtcc caaatctgtg
cggagccgaa 1620*

*atctgggagg cgccgccgca cccctctag cgggcgcggg gcgaagcgg
gcggcgccgg 1680*

*caggaaggaa atgggcgggg agggccttcg tgcgtcgccg cgccgccgtc
cccttctccc 1740*

*tctccagcct cggggctgtc cgcgggggga cggctgcctt cgggggggac
ggggcagggc 1800*

*ggggttcggc ttctggcgtg tgaccggcgg ctctagagcc tctgctaacc atgttcatgc
1860*

*cttcttcttt ttctacagc tcctgggcaa cgtgctggtt attgtgctgt ctcattcatt
1920*

*tggcaaagaa ttcgctagcg ggcactttgc actggaactt acaacacccg
agcaaggacg 1980*

cgactctcca ccatgtaccg gatgcagctg ctgctgctga tcgccctgag cctggccctg
2040

gtgaccaaca gcgaggtgca gctggtggag agcggggggg ggctggtgca
gcccgggggg 2100

agcctgcggc tgagctgcgc cgccagcggg tacgacttca cccactacgg
gatgaactgg 2160

gtgcggcagg cccccgggaa ggggctggag tgggtggggg ggatcaacac
ctacaccggg 2220

gagcccacct acgccgccga cttcaagcgg cggttcacct tcagcctgga caccagcaag
2280

agcaccgcct acctgcagat gaacagcctg cgggccgagg acaccgccgt
gtactactgc 2340

gccaagtacc cctactacta cgggaccagc cactggtact tcgacgtgtg
ggggcagggg 2400

accctggtga ccgtgagcag cgccagcacc aaggggccca gcgtgttccc cctggccccc
2460

*agcagcaaga gcaccagcgg ggggaccgcc gccctgggggt gcctggtgaa
ggactacttc 2520*

*cccgagcccc tgaccgtgag ctggaacagc ggggccctga ccagcgggggt
gcacaccttc 2580*

*cccgccgtgc tgcagagcag cgggctgtac agcctgagca gcgtggtgac
cgtgcccagc 2640*

*agcagcctgg ggaccagac ctacatctgc aacgtgaacc acaagcccag
caacaccaag 2700*

*gtggacaaga aggtggagcc caagagctgc gacaagaccc acctgcggaa
gcggcgggcc 2760*

*cccgtgaagc agaccctgaa cttcgacctg ctgaagctgg ccggggacgt
ggagagcaac 2820*

*cccgggcccc tgtaccggat gcagctgctg ctgctgatcg ccctgagcct ggccctggtg
2880*

*accaacagcg acatccagct gacccagagc cccagcagcc tgagcgccag
cgtgggggac 2940*

*cgggtgacca tcacctgacg cgccagccag gacatcagca actacctgaa
ctggtaccag 3000*

*cagaagcccg ggaaggcccc caaggtgctg atctacttca ccagcagcct
gcacagcggg 3060*

*gtgcccagcc ggttcagcgg gagcgggagc gggaccgact tcaccctgac
catcagcagc 3120*

*ctgcagcccg aggacttcgc cacctactac tgccagcagt acagcacctg gccctggacc
3180*

*ttcgggcagg ggaccaaggt ggagatcaag cggaccgtgg ccgccccag
cgtgttcac 3240*

*ttcccccca gcgacgagca gctgaagagc gggaccgcca gcgtggtgtg
cctgctgaac 3300*

*aacttctacc cccgggaggc caaggtgacg tggaaggtgg acaacgccct
gcagagcggg 3360*

*aacagccagg agagcgtgac cgagcaggac agcaaggaca gcacctacag
cctgagcagc 3420*

*accctgacct tgagcaaggc cgactacgag aagcacaagg tgtacgcctg
cgaggtgacc 3480*

*caccaggggc tgagcagccc cgtgaccaag agcttcaacc ggggggagtg
ctgataaagc 3540*

*ggccgcggta cctctagagt cgacccgggc ggcctcgagg acggggtgaa
ctacgcctga 3600*

*ggatccgatc ttttccctc tgccaaaaat tatggggaca tcatgaagcc ccttgagcat
3660*

*ctgacttctg gctaataaag gaaatttatt ttcattgcaa tagtgtgttg gaatttttg
3720*

*tgtctctcac tcggaagcaa ttcgttgatc tgaatttcga ccaccataa taccattac
3780*

*cctggtagat aagtagcatg gcgggtaat cattaactac aaggaacccc
tagtgatgga 3840*

*gttggccact ccctctctgc gcgctcgctc gctcactgag gccgggacgac caaaggctcg
3900*

*ccgacgcccc ggctttgcc gggcggcctc agtgagcgag cgagcgcgca g
3951*

<210> 45

<211> 3809

<212> DNA

<213> Artificial Sequence

<220>

<223> ITR.CMV.PI.alphaVEGFv7.eMCV.IRES.SV40.ITR

<220>

<221> repeat_region

<222> (1)..(130)

<223> *ITR*

<220>

<221> *promoter*

<222> (191)..(932)

<223> *human CMV I.E. enhancer & promoter*

<220>

<221> *Intron*

<222> (1047)..(1179)

<223> *Promega chimeric intron*

<220>

<221> *5'UTR*

<222> (1248)..(1295)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1299)..(1307)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1305)..(1364)

<223> *leader*

<220>

<221> *misc_feature*

<222> (1365)..(1685)

<223> *aVEGFv7 VL*

<220>

<221> *misc_feature*

<222> (1686)..(2006)

<223> *CL*

<220>

<221> *enhancer*

<222> (2018)..(2608)

<223> *IRES*

<220>

<221> *misc_feature*

<222> (2606)..(2665)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2666)..(3034)

<223> *aVEGFv7 VH*

<220>

<221> *misc_feature*

<222> (3035)..(3355)

<223> *CH1*

<220>

<221> *polyA_signal*

<222> (3384)..(3615)

<223> *SV40 late polyadenylation signal*

<220>

<221> *repeat_region*

<222> (3680)..(3809)

<223> *ITR*

<400> 45

ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc ccgggcgctc ggcgacctt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctacgta gccatgctct
180

aggaagatct tcaatattgg ccattagcca tattattcat tggttatata gcataaatca
240

atattggcta ttggccattg catacgttgt atctatatca taatatgtac atttatattg
300

gctcatgtcc aatatgaccg ccatgttggc attgattatt gactagttat taatagtaat
360

caattacggg gtcattagtt catagcccat atatggagtt ccgcgttaca taacttacgg
420

taaatggccc gcctggctga ccgcccacg acccccgccc attgacgtca ataatgacgt
480

atgttcccat agtaacgcca atagggactt tccattgacg tcaatgggtg gagtatttac
540

ggtaaactgc ccacttgga gtacatcaag tgtatcatat gccaaagtccg ccccctattg
600

acgtcaatga cggtaaattg cccgcctggc attatgccca gtacatgacc ttacgggact
660

ttctacttg gcagtacatc tacgtattag tcatcgctat taccatgggtg atgcggtttt
720

ggcagtacac caatgggagt ggatagcggg ttgactcacg gggatttcca agtctccacc
780

ccattgacgt caatgggagt ttgttttggc accaaaatca acgggacttt caaaatgtc
840

*gtaataaccc cgccccgttg acgcaaattg gcggtaggcg tgtacggtgg
gaggctata 900*

*taagcagagc tcgtttagtg aaccgtcaga tcactagaag ctttattgcg gtagtttatc
960*

*acagttaaat tgctaacgca gtcagtgctt ctgacacaac agtctcgaac ttaagctgca
1020*

*gaagttggtc gtgaggcact gggcaggtaa gtatcaaggt tacaagacag
gtttaaggag 1080*

*accaatagaa actgggcttg tcgagacaga gaagactctt gcgtttctga taggcaccta
1140*

*ttggtcttac tgacatccac tttgcctttc tctccacagg tgtccactcc cagttcaatt
1200*

*acagctctta aggctagagt acttaatacg actcactata ggctagcggg gactttgcac
1260*

*tggaacttac aacacccgag caaggacgcg actctagacc caccatgtac
cggatgcagc 1320*

*tgctgctgct gatcgccctg agcctggccc tggtagacaa cagcgatatac cagctgaccc
1380*

*agagcccaag cagcctgagc gccagcgtgg gagatcgggt gaccatcacc
tgcagcgcca 1440*

*gccaggatat cagcaactac ctgaactggt accagcagaa gccaggaag
gccccaaagg 1500*

*tgctgatcta cttcaccagc agcctgcaca gcggagtgcc aagccggttc
agcgggaagcg 1560*

*gaagcggaac cgatttcacc ctgacatca gcagcctgca gccagaggat ttcgccacct
1620*

*actactgcca gcagtacagc accgtgccat ggaccttcgg acaggggaacc
aaggtggaga 1680*

*tcaagcggac cgtggccgcc ccaagcgtgt tcatttccc accaagcgat gagcagctga
1740*

*agagcggaac cgccagcgtg gtgtgcctgc tgaacaactt ctaccacgg
gaggccaagg 1800*

*tgcagtggaa ggtggataac gccctgcaga gcggaaacag ccaggagagc
gtgaccgagc 1860*

*aggatagcaa ggatagcacc tacagcctga gcagcacctt gaccctgagc
aaggccgatt 1920*

*acgagaagca caaggtgtac gcctgcgagg tgaccaccca gggactgagc
agcccagtga 1980*

*ccaagagctt caaccgggga gagtgctgat aaggccggcc cctctcctc cccccccct
2040*

*aacgttactg gccgaagccg cttggaataa ggccggtgtg cgtttgctta tatgttattt
2100*

*tccaccatat tgccgtcttt tggcaatgtg agggcccgga aacctggccc tgtcttcttg
2160*

acgagcattc ctaggggtct ttcccctctc gccaaaggaa tgcaaggtct gttgaatgtc
2220

gtgaaggaag cagttcctct ggaagcttct tgaagacaaa caacgtctgt agcgaccctt
2280

tgcaaggcagc ggaaccccc acctggcgac aggtgcctct gcggccaaaa
gccacgtgta 2340

taagatacac ctgcaaaggc ggcacaacc cagtgccacg ttgtgagttg
gatagttgtg 2400

gaaagagtca aatggctctc ctcaagcgta ttcaacaagg ggctgaagga
tgcccagaag 2460

gtacccatt gtatgggatc tgatctgggg cctcgggtaca catgctttac atgtgttag
2520

tcgaggtaa aaaaacgtct aggcccccg aaccacgggg acgtggtttt cctttgaaaa
2580

acacgatgat aatatggcca caaccatgta ccgcatgcag ctgctgctgc tgatcgcctt
2640

*gagcctggcc ctggtgacca acagcgaggt gcagctggtg gagagcggag
gaggactggt 2700*

*gcagccagga ggaagcctgc ggctgagctg cgccgccagc ggatacgatt
tcaccacta 2760*

*cggaatgaac tgggtgcggc aggccccagg aaagggactg gagtgggtgg
gatggatcaa 2820*

*cacctacacc ggagagccaa cctacgccgc cgatttcaag cggcggttca ccttcagcct
2880*

*ggataccagc aagagcaccg cctacctgca gatgaacagc ctgcgggccg
aggataccgc 2940*

*cggtgtactac tgcgccaagt acccatacta ctacggaacc agccactggt acttcgatgt
3000*

*gtggggacag ggaaccctgg tgaccgtgag cagcgccagc accaagggac
caagcgtgtt 3060*

*cccactggcc ccaagcagca agagcaccag cggaggaacc gccgccctgg
gatgcctggt 3120*

*gaaggattac ttcccagagc cagtgaccgt gagctggaac agcggagccc
tgaccagcgg 3180*

*agtgcacacc ttcccagccg tgctgcagag cagcggactg tacagcctga
gcagcgtggt 3240*

*gaccgtgcc aagcagcagcc tgggaacca gacctacatc tgcaacgtga
accacaagcc 3300*

*aagcaacacc aaggtggata agaaggtgga gccaaagagc tgcgataaga
cccacctgaa 3360*

*gtgatgaaag cttgcggccg cttcgagcag acatgataag atacattgat
gagtttgac 3420*

*aaaccacaac tagaatgcag tgaaaaaaaaat gctttatttg tgaatttgt gatgctattg
3480*

*ctttatttgt aaccattata agctgcaata aacaagttaa caacaacaat tgcattcatt
3540*

ttatgtttca ggttcagggg gagatgtggg aggttttta aagcaagtaa aacctctaca
3600

aatgtggtaa aatcgataag gatcttccta gagcatggct acgtagataa
gtagcatggc 3660

gggttaatca ttaactacaa ggaacccta gtgatggagt tggccactcc ctctctgcgc
3720

gctcgctcgc tactgaggc cgggcgacca aaggtcgccc gacgcccggg cttgcccgg
3780

gcggcctcag tgagcgagcg agcgcgcag 3809

<210> 46

<211> 3686

<212> DNA

<213> Artificial Sequence

<220>

<223> *ITR.CMV.PI.alphaVEGFv7.fmdIRES.SV40.ITR*

<220>

<221> *repeat_region*

<222> *(1)..(130)*

<223> *ITR*

<220>

<221> *promoter*

<222> *(191)..(932)*

<223> *human CMV I.E. enhancer and promoter*

<220>

<221> *TATA_signal*

<222> *(897)..(901)*

<220>

<221> *Intron*

<222> *(1047)..(1179)*

<223> *Promega chimeric intron*

<220>

<221> 5'UTR

<222> (1248)..(1295)

<223> c-myc 5'UTR

<220>

<221> misc_feature

<222> (1299)..(1307)

<223> kozak

<220>

<221> misc_feature

<222> (1305)..(1313)

<223> leader

<220>

<221> misc_feature

<222> (1314)..(1364)

<223> leader

<220>

<221> *misc_feature*

<222> (1365)..(1685)

<223> *aVEGFv7 VL*

<220>

<221> *misc_feature*

<222> (1686)..(2006)

<223> *CL*

<220>

<221> *misc_feature*

<222> (2021)..(2482)

<223> *FMDV*

<220>

<221> *misc_feature*

<222> (2501)..(2542)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2543)..(2911)

<223> *aVEGFv7 VH*

<220>

<221> *misc_feature*

<222> (2912)..(3232)

<223> *CH1*

<220>

<221> *polyA_signal*

<222> (3261)..(3492)

<223> *Sv40 late polyadenylation signal*

<220>

<221> *repeat_region*

<222> (3557)..(3686)

<223> *ITR*

<400> 46

ctgcgcgctc gctcgtcac tgaggccgcc cgggcaaagc ccgggcgctc ggcgaccttt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgcagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctacgta gccatgctct
180

aggaagatct tcaatattgg ccattagcca tattattcat tggttatata gcataaatca
240

atattggcta ttggccattg catacgttgt atctatatca taatatgtac atttatattg
300

gctcatgtcc aatatgaccg ccatgttggc attgattatt gactagttat taatagtaat
360

caattacggg gtcattagtt catagcccat atatggagtt ccgcgttaca taacttacgg
420

taaatggccc gcctggctga ccgccaacg acccccgccc attgacgtca ataatgacgt
480

atgttcccat agtaacgccca atagggactt tccattgacg tcaatgggtg gagtatttac
540

ggtaaactgc ccacttgcca gtacatcaag tgtatcatat gccaaagtccg ccccctattg
600

acgtcaatga cggtaaattgg cccgcctggc attatgccca gtacatgacc ttacgggact
660

ttcctacttg gcagtacatc tacgtattag tcatcgctat taccatgggtg atgcggtttt
720

ggcagtacac caatgggcgt ggatagcggg ttgactcacg gggatttcca agtctccacc
780

ccattgacgt caatgggagt ttgttttggc accaaaatca acgggacttt ccaaaatgtc
840

gtaataaccc cgccccgttg acgcaaatgg gcggtaggcg tgtacgggtg
gaggctata 900

taagcagagc tcgttagtg aaccgtcaga tactagaag cttattgcg gtagttatc
960

acagttaaat tgctaacgca gtcagtgctt ctgacacaac agtctcgaac ttaagctgca
1020

gaagttggtc gtgaggcact gggcaggtaa gtatcaaggt tacaagacag
gtttaaggag 1080

accaatagaa actgggcttg tcgagacaga gaagactctt gcgtttctga taggcaccta
1140

ttggtcttac tgacatccac tttgccttc tctccacagg tgtccactcc cagttcaatt
1200

acagctctta aggctagagt acttaatacg actcactata ggctagcggg gactttgcac
1260

tggaacttac aacacccgag caaggacgcg actctagacc caccatgtac
cggatgcagc 1320

tgctgctgct gatcgccctg agcctggccc tggtagacaa cagcgatatc cagctgaccc
1380

*agagcccaag cagcctgagc gccagcgtgg gagatcgggt gaccatcacc
tgcagcgcca 1440*

*gccaggatat cagcaactac ctgaactggt accagcagaa gccaggaaag
gccccaaagg 1500*

*tgctgatcta cttcaccagc agcctgcaca gcggagtgcc aagccggttc
agcgggaagcg 1560*

*gaagcggaac cgatttcacc ctgaccatca gcagcctgca gccagaggat ttcgccacct
1620*

*actactgcca gcagtacagc accgtgccat ggaccttcgg acaggggaacc
aaggtggaga 1680*

*tcaagcggac cgtggccgcc ccaagcgtgt tcattttccc accaagcgat gagcagctga
1740*

*agagcggaac cgccagcgtg gtgtgcctgc tgaacaactt ctaccacgg
gaggccaagg 1800*

*tgcagtggaa ggtggataac gccctgcaga gcggaacag ccaggagagc
gtgaccgagc 1860*

*aggatagcaa ggatagcacc tacagcctga gcagcacct gaccctgagc
aaggccgatt 1920*

*acgagaagca caaggtgtac gcctgcgagg tgaccaccca gggactgagc
agcccagtga 1980*

*ccaagagctt caaccgggga gagtgctgat aaggccggcc agcaggttc
cccaactgac 2040*

*acaaaacgtg caacttgaaa ctccgcctgg tctttccagg tctagagggg taacacttg
2100*

*tactgcgttt ggctccacgc tcgatccact ggcgagtgtt agtaacagca ctgttgcttc
2160*

*gtagcggagc atgacggccg tgggaactcc tccttggtaa caaggaccca
cggggccaaa 2220*

*agccacgccc acacgggccc gtcattgtgtg caaccccagc acggcgactt
tactgcgaaa 2280*

cccactttaa agtgacattg aaactggtac ccacacactg gtgacaggct aaggatgcc
2340

ttcaggtacc ccgaggtaac acgggacact cgggatctga gaaggggact
ggggcttcta 2400

taaaagcgct cggtttaaaa agcttctatg cctgaatagg tgaccggagg tcggcacctt
2460

tcctttacaa ttaatgacct taatgtaccg catgcagctg ctgctgctga tcgccctgag
2520

cctggccctg gtgaccaaca gcgaggtgca gctggtggag agcggaggag
gactggtgca 2580

gccaggagga agcctgcggc tgagctgcgc cgccagcgga tacgatttca
cccactacgg 2640

aatgaactgg gtgcggcagg ccccaggaaa gggactggag tgggtgggat
ggatcaacac 2700

ctacaccgga gagccaacct acgccgccga tttcaagcgg cggttcacct tcagcctgga
2760

taccagcaag agcaccgcct acctgcagat gaacagcctg cgggccgagg
ataccgccgt 2820

gtactactgc gccaaagtacc catactacta cggaaccagc cactggtact tcgatgtgtg
2880

gggacagga accctggtga ccgtgagcag cgccagcacc aagggaccaa
gcgtgttccc 2940

actggcccca agcagcaaga gcaccagcgg aggaaccgcc gccctgggat
gcctggtgaa 3000

ggattacttc ccagagccag tgaccgtgag ctggaacagc ggagccctga
ccagcggagt 3060

gcacaccttc ccagccgtgc tgcagagcag cggactgtac agcctgagca
gcgtggtgac 3120

cgtgccaaag agcagcctgg gaaccagac ctacatctgc aacgtgaacc
acaagccaag 3180

*caacaccaag gtggataaga aggtggagcc aaagagctgc gataagaccc
acctgaagtg 3240*

*atgaaagctt gcggccgctt cgagcagaca tgataagata cattgatgag
tttgacaaa 3300*

*ccacaactag aatgcagtga aaaaaatgct ttatttgtga aatttgtgat gctattgctt
3360*

*tatttgaac cattataagc tgcaataaac aagttaaca caacaattgc attcatttta
3420*

*tgtttcaggt tcagggggag atgtgggagg tttttaag caagtaaac ctctacaaat
3480*

*gtggtaaaat cgataaggat cttcctagag catggctacg tagataagta
gcatggcggg 3540*

*ttaatcatta actacaagga acccctagtg atggagttgg ccactccctc tctgcgcgct
3600*

cgctcgctca ctgaggccgg gcgaccaaag gtcgcccgac gcccgggctt
tgcccgggcg 3660

gcctcagtga gcgagcgagc gcgcag 3686

<210> 47

<211> 3619

<212> DNA

<213> *Artificial Sequence*

<220>

<223> *ITR.CMV.PI.alphaVEGFv7.cMycIRES.SV40.ITR*

<220>

<221> *repeat_region*

<222> *(1)..(130)*

<223> *ITR*

<220>

<221> *promoter*

<222> (191)..(932)

<223> *human CMV I.E. enhancer and promoter*

<220>

<221> *TATA_signal*

<222> (897)..(901)

<220>

<221> *Intron*

<222> (1047)..(1179)

<223> *Promega chimeric intron*

<220>

<221> *5'UTR*

<222> (1248)..(1295)

<223> *c-myc 5'UTR*

<220>

<221> *misc_feature*

<222> (1299)..(1307)

<223> *kozak*

<220>

<221> *misc_feature*

<222> (1305)..(1313)

<223> *leader*

<220>

<221> *misc_feature*

<222> (1314)..(1364)

<223> *leader*

<220>

<221> *misc_feature*

<222> (1365)..(1685)

<223> *aVEGFv7 VL*

<220>

<221> *misc_feature*

<222> (1686)..(2006)

<223> *CL*

<220>

<221> *misc_feature*

<222> (2021)..(2415)

<223> *IRES c-myc*

<220>

<221> *misc_feature*

<222> (2226)..(2273)

<223> *mini c-myc IRES*

<220>

<221> *misc_feature*

<222> (2275)..(2275)

<223> *this C to T mutaion increases expression*

<220>

<221> *misc_feature*

<222> (2434)..(2475)

<223> *leader*

<220>

<221> *misc_feature*

<222> (2476)..(2844)

<223> *aVEGFv7 VH*

<220>

<221> *misc_feature*

<222> (2845)..(3165)

<223> *CH1*

<220>

<221> *polyA_signal*

<222> (3194)..(3425)

<223> *SV40 late polyadenylation signal*

<220>

<221> *repeat_region*

<222> (3490)..(3619)

<223> *ITR*

<400> 47

ctgcgcgctc gctcgtcac tgaggccgcc cgggcaaagc ccgggcgctc ggcgaccttt
60

ggtcgcccgg cctcagtgag cgagcgagcg cgagagagg gagtggccaa
ctccatcact 120

aggggttcct tgtagttaat gattaacccg ccatgctact tatctacgta gccatgctct
180

aggaagatct tcaatattgg ccattagcca tattattcat tggttatata gcataaatca
240

atattggcta ttggccattg catacgttgt atctatatca taatatgtac atttatattg
300

gctcatgtcc aatatgaccg ccatgttggc attgattatt gactagttat taatagtaat
360

caattacggg gtcattagtt catagcccat atatggagtt ccgcgttaca taacttacgg
420

taaatggccc gcctggctga ccgccaacg acccccgccc attgacgtca ataatgacgt
480

atgttcccat agtaacgccca atagggactt tccattgacg tcaatgggtg gagtatttac
540

ggtaaactgc ccacttgcca gtacatcaag tgtatcatat gccaaagtccg ccccctattg
600

acgtcaatga cggtaaattgg cccgcctggc attatgccca gtacatgacc ttacgggact
660

ttcctacttg gcagtacatc tacgtattag tcatcgctat taccatgggtg atgcggtttt
720

ggcagtacac caatgggcgt ggatagcggg ttgactcacg gggatttcca agtctccacc
780

ccattgacgt caatgggagt ttgttttggc accaaaatca acgggacttt ccaaaatgtc
840

gtaataaccc cgccccgttg acgcaaatgg gcggtaggcg tgtacgggtg
gaggctata 900

taagcagagc tcgtttagtg aaccgtcaga tcaactagaag ctttattgcg gtagtttatc
960

acagttaa at tgctaacgca gtcagtgctt ctgacacaac agtctcgaac ttaagctgca
1020

gaagttggtc gtgaggcact gggcaggtaa gtatcaaggt tacaagacag
gtttaaggag 1080

accaatagaa actgggcttg tcgagacaga gaagactctt gcgtttctga taggcaccta
1140

ttggtcttac tgacatccac tttgcctttc tctccacagg tgtccactcc cagttcaatt
1200

acagctctta aggctagagt acttaatacg actcactata ggctagcggg gactttgcac
1260

tggaacttac aacacccgag caaggacgcg actctagacc caccatgtac
cggatgcagc 1320

tgctgctgct gatcgccctg agcctggccc tggtgaccaa cagcgatatc cagctgaccc
1380

*agagcccaag cagcctgagc gccagcgtgg gagatcgggt gaccatcacc
tgcagcgcca 1440*

*gccaggatat cagcaactac ctgaactggt accagcagaa gccaggaaag
gccccaaagg 1500*

*tgctgatcta cttcaccagc agcctgcaca gcggagtgcc aagccggttc
agcgggaagcg 1560*

*gaagcggaac cgatttcacc ctgaccatca gcagcctgca gccagaggat ttcgccacct
1620*

*actactgcca gcagtacagc accgtgccat ggaccttcgg acaggggaacc
aaggtggaga 1680*

*tcaagcggac cgtggccgcc ccaagcgtgt tcattttccc accaagcgat gagcagctga
1740*

*agagcggaac cgccagcgtg gtgtgcctgc tgaacaactt ctaccacgg
gaggccaagg 1800*

*tgcagtggaa ggtggataac gccctgcaga gcggaacag ccaggagagc
gtgaccgagc 1860*

*aggatagcaa ggatagcacc tacagcctga gcagaccct gaccctgagc
aaggccgatt 1920*

*acgagaagca caaggtgtac gcctgcgagg tgaccaccca gggactgagc
agcccagtga 1980*

*ccaagagctt caaccgggga gagtgctgat aaggccggcc aattccagcg
agaggcagag 2040*

*ggagcgagcg ggcggccggc tagggtggaa gagccgggcg agcagagctg
cgctgcgggc 2100*

*gtcctgggaa gggagatccg gagcgaatag ggggcttcgc ctctggccca
gccctcccgc 2160*

*tgatcccca gccagcggtc cgcaaccctt gccgcatcca cgaaacttg cccatagcag
2220*

*cgggcgggca cttgcaactg gaactataa cacccgagca aggacgcgac
tctcccagc 2280*

cggggaggct attctgcca ttggggaca cttccccgcc gctgccagga cccgcttctc
2340

tgaaaggctc tccttgacg tgcttagacg ctggatTTTT ttcgggtagt ggaaaaccag
2400

cagcctcccg cgacgatgta ccgcatgcag ctgctgctgc tgatcgccct gagcctggcc
2460

ctggtgacca acagcgagg gcagctggtg gagagcggag gaggactggt
gcagccagga 2520

ggaagcctgc ggctgagctg cgccgccagc ggatacgatt tcaccacta
cggaatgaac 2580

tgggtgcggc aggccccagg aaagggactg gagtgggtgg gatggatcaa
cacctacacc 2640

ggagagccaa cctacgccgc cgatttcaag cggcggttca ccttcagcct ggataccagc
2700

*aagagcaccg cctacctgca gatgaacagc ctgcgggccg aggataccgc
cgtgtactac 2760*

*tcgccaagt acccatacta ctacggaacc agccactggt acttcgatgt gtggggacag
2820*

*ggaaccctgg tgaccgtgag cagcgccagc accaagggac caagcgtgtt
cccactggcc 2880*

*ccaagcagca agagcaccag cggaggaacc gccgccctgg gatgcctggt
gaaggattac 2940*

*ttcccagagc cagtgaccgt gagctggaac agcggagccc tgaccagcgg
agtgcacacc 3000*

*ttcccagccg tgctgcagag cagcggactg tacagcctga gcagcgtggt
gaccgtgcca 3060*

*agcagcagcc tgggaacca gacctacatc tgcaacgtga accacaagcc
aagcaacacc 3120*

*aaggtggata agaaggtgga gccaaagagc tgcgataaga cccacctgaa
gtgatgaaag 3180*

*cttgcggccg cttcgagcag acatgataag atacattgat gagtttggac
aaaccacaac 3240*

*tagaatgcag tgaaaaaat gctttatttg tgaatttgt gatgctattg ctttatttg
3300*

*aaccattata agctgcaata aacaagtaa caacaacaat tgcattcatt ttatgttca
3360*

*ggttcagggg gagatgtggg aggttttta aagcaagtaa aacctctaca
aatgtggtaa 3420*

*aatcgataag gatcttcta gagcatggct acgtagataa gtagcatggc
gggttaatca 3480*

*ttaactacaa ggaacccta gtgatggagt tggccactcc ctctctgcgc gctcgctcg
3540*

*tcactgaggc cgggcgacca aaggtcgccc gacgcccggg cttgcccgg
gcggcctcag 3600*

tgagcgagcg agcgcgcag

3619

<210> 48

<211> 738

<212> PRT

<213> Unknown

<220>

<223> AAV8 capsid

<400> 48

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser

1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Lys Pro

20 25 30

Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro

35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro

50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp

65 70 75 80

Gln Gln Leu Gln Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala

85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly

100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro

115 120 125

Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg

130 135 140

Pro Val Glu Pro Ser Pro Gln Arg Ser Pro Asp Ser Ser Thr Gly Ile

145 150 155 160

Gly Lys Lys Gly Gln Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln

165 170 175

Thr Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro

180 185 190

Pro Ala Ala Pro Ser Gly Val Gly Pro Asn Thr Met Ala Ala Gly Gly

195 200 205

Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser

210 215 220

Ser Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val

225 230 235 240

Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His

245 250 255

Leu Tyr Lys Gln Ile Ser Asn Gly Thr Ser Gly Gly Ala Thr Asn Asp

260 265 270

Asn Thr Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn

275 280 285

Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn

290 295 300

Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Ser Phe Lys Leu Phe Asn

305 310 315 320

Ile Gln Val Lys Glu Val Thr Gln Asn Glu Gly Thr Lys Thr Ile Ala

325 330 335

Asn Asn Leu Thr Ser Thr Ile Gln Val Phe Thr Asp Ser Glu Tyr Gln

340 345 350

Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe

355 360 365

Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn

370 375 380

Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr

385 390 395 400

Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Thr Tyr

405 410 415

Thr Phe Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser

420 425 430

Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu

435 440 445

Ser Arg Thr Gln Thr Thr Gly Gly Thr Ala Asn Thr Gln Thr Leu Gly

450 455 460

Phe Ser Gln Gly Gly Pro Asn Thr Met Ala Asn Gln Ala Lys Asn Trp

465

470

475

480

Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Thr Thr Thr Gly

485

490

495

Gln Asn Asn Asn Ser Asn Phe Ala Trp Thr Ala Gly Thr Lys Tyr His

500

505

510

Leu Asn Gly Arg Asn Ser Leu Ala Asn Pro Gly Ile Ala Met Ala Thr

515

520

525

His Lys Asp Asp Glu Glu Arg Phe Phe Pro Ser Asn Gly Ile Leu Ile

530

535

540

Phe Gly Lys Gln Asn Ala Ala Arg Asp Asn Ala Asp Tyr Ser Asp Val

545

550

555

560

Met Leu Thr Ser Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr

565

570

575

Glu Glu Tyr Gly Ile Val Ala Asp Asn Leu Gln Gln Gln Asn Thr Ala

580

585

590

Pro Gln Ile Gly Thr Val Asn Ser Gln Gly Ala Leu Pro Gly Met Val

595

600

605

Trp Gln Asn Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile

610

615

620

Pro His Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe

625

630

635

640

Gly Leu Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val

645

650

655

Pro Ala Asp Pro Pro Thr Thr Phe Asn Gln Ser Lys Leu Asn Ser Phe

660

665

670

Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu

675

680

685

Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr

690

695

700

Ser Asn Tyr Tyr Lys Ser Thr Ser Val Asp Phe Ala Val Asn Thr Glu

705

710

715

720

Gly Val Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg

725

730

735

Asn Leu

<210> 49

<211> 2217

<212> DNA

<213> Unknown

<220>

<223> *Nucleic acid sequence of AAV8 capsid*

<400> 49

atggctgccg atggttatct tccagattgg ctcgaggaca acctctctga gggcattcgc
60

gagtgggtggg cgctgaaacc tggagccccg aagcccaag ccaaccagca
aaagcaggac 120

gacggccggg gtctggtgct tcctggctac aagtacctcg gacccttcaa cggactcgac
180

aagggggagc ccgtcaacgc ggcggacgca gcggccctcg agcacgacaa
ggcctacgac 240

cagcagctgc aggcgggtga caatccgtac ctgcggtata accacgccga
cgccgagttt 300

caggagcgtc tgcaagaaga tacgtctttt gggggcaacc tcgggcgagc
agtcttccag 360

gccaagaagc gggttctcga acctctcggc ctggttgagg aaggcgctaa
gacggctcct 420

ggaagaaga gaccggtaga gccatcacc cagcgttctc cagactcctc
tacgggcatc 480

*ggcaagaaag gccaacagcc cgccagaaaa agactcaatt ttggtcagac
tggcgactca 540*

*gagtcagttc cagaccctca acctctcgga gaacctccag cagcgcctc tgggtgggga
600*

*cctaatacaa tggctgcagg cggtggcgca ccaatggcag acaataacga
aggcgccgac 660*

*ggagtgggta gttcctcggg aaattggcat tgcgattcca catggctggg
cgacagagtc 720*

*atcaccacca gcacccgaac ctgggccctg cccacctaca acaaccacct ctacaagcaa
780*

*atctccaacg ggacatcggg aggagccacc aacgacaaca cctacttcgg
ctacagcacc 840*

*ccctgggggt attttgactt taacagattc cactgccact ttcaccacg tgactggcag
900*

cgactcatca acaacaactg gggattccgg cccaagagac tcagttcaa gctttcaac
960

atccaggta aggaggtcac gcagaatgaa ggcaccaaga ccatcgccaa
taacctcacc 1020

agcaccatcc aggtgtttac ggactcggag taccagctgc cgtacgttct cggctctgcc
1080

caccagggct gcctgcctcc gttcccggcg gacgtgttca tgattcccca gtacggctac
1140

ctaactca acaacggtag tcaggccgtg ggacgctcct ctttctactg cctggaatac
1200

tttccttcgc agatgctgag aaccggcaac aactccagt ttacttacac ctcgaggac
1260

gtgcctttcc acagcagcta cgcccacagc cagagcttgg accggctgat gaatcctctg
1320

attgaccagt acctgtacta cttgtctcgg actcaaaca caggaggcac
ggcaaatacg 1380

*cagactctgg gcttcagcca aggtgggcct aatacaatgg ccaatcaggc
aaagaactgg 1440*

*ctgccaggac cctgttaccg ccaacaacgc gtctcaacga caaccgggca
aaacaacaat 1500*

*agcaactttg cctggactgc tgggaccaa taccatctga atggaagaaa ttcattggct
1560*

*aatcctggca tcgctatggc aacacacaaa gacgacgagg agcgtttttt tcccagtaac
1620*

*gggatcctga ttttggcaa acaaaatgct gccagagaca atgcggatta
cagcgatgtc 1680*

*atgctcacca gcgaggaaga aatcaaaacc actaacctg tggctacaga
ggaatacgg 1740*

*atcgtggcag ataacttgca gcagcaaac acggctcctc aaattggaac
tgtcaacagc 1800*

*cagggggcct tacccggtat ggtctggcag aaccgggacg tgtacctgca
gggtcccatc 1860*

*tgggccaaga ttctcacac ggacggcaac ttccaccgt ctccgctgat gggcggcttt
1920*

*ggcctgaaac atcctccgcc tcagatcctg atcaagaaca cgctgtacc tgcggatcct
1980*

*ccgaccacct tcaaccagtc aaagctgaac tctttcatca cgcaatacag caccggacag
2040*

*gtcagcgtgg aaattgaatg ggagctgcag aaggaaaaca gcaagcgtg
gaaccccgag 2100*

*atccagtaca ctcctaacta ctacaaatct acaagtgtgg actttgctgt taatacagaa
2160*

*ggcgtgtact ctgaaccccg cccattggc acccgttacc tcaccgtaa tctgtaa
2217*