



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2016/11/21
(87) Date publication PCT/PCT Publication Date: 2017/05/26
(85) Entrée phase nationale/National Entry: 2018/05/17
(86) N° demande PCT/PCT Application No.: CN 2016/106598
(87) N° publication PCT/PCT Publication No.: 2017/084630
(30) Priorités/Priorities: 2015/11/20 (CN PCT/CN2015/095166);
2016/11/08 (CN PCT/CN2016/105042)

(51) Cl.Int./Int.Cl. *C07D 245/04* (2006.01),
A61K 31/395 (2006.01), *A61P 31/04* (2006.01)

(71) Demandeurs/Applicants:
RQX PHARMACEUTICALS, INC., US;
GENENTECH, INC., US

(72) Inventeurs/Inventors:
CHEN, YONGSHENG, CN;
SMITH, PETER ANDREW, US;
ROBERTS, TUCKER CURRAN, US;
HIGUCHI, ROBERT I., US;
PARASELLI, PRASUNA, US;
...

(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : ANTIBIOTIQUES MACROCYCLIQUES A LARGE SPECTRE
(54) Title: MACROCYCLIC BROAD SPECTRUM ANTIBIOTICS

(57) **Abrégé/Abstract:**

Provided herein are antibacterial compounds, wherein the compounds in some embodiments have broad spectrum bioactivity. In various embodiments, the compounds act by inhibition of bacterial type 1 signal peptidase (SpsB), an essential protein in bacteria. Pharmaceutical compositions and methods for treatment using the compounds described herein are also provided.

(72) **Inventeurs(suite)/Inventors(continued)**: KOEHLER, MICHAEL F. T., US; SCHWARZ, JACOB BRADLEY, US;
CRAWFORD, JAMES JOHN, US; LY, CUONG Q., US; HANAN, EMILY J., US; HU, HUIYONG, US; YU, ZHIYONG, CN

(51) **International Patent Classification:**
C07D 245/04 (2006.01) *A61P 31/04* (2006.01)
A61K 31/395 (2006.01)

(21) **International Application Number:**
PCT/CN2016/106598

(22) **International Filing Date:**
21 November 2016 (21.11.2016)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
PCT/CN2015/095166
20 November 2015 (20.11.2015) CN
PCT/CN2016/105042
8 November 2016 (08.11.2016) CN

(71) **Applicants:** **RQX PHARMACEUTICALS, INC.** [US/US]; 11099 N. Torrey Pines Road, Suite 290, La Jolla, CA 92037 (US). **GENENTECH, INC.** [US/US]; 1 DNA Way, South San Francisco, CA 94080 (US).

(72) **Inventors:** **CHEN, Yongsheng**; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **SMITH, Peter Andrew**; 260 King Street Unit 909, San Francisco, CA 94107 (US). **ROBERTS, Tucker Curran**; 536 Maple Street, Apt. 301, San Diego, CA 92103 (US). **HIGUCHI, Robert I.**; 434 Marview Drive, Solana Beach, CA 92075 (US). **PARASELLI, Prasuna**; 6323 Oleander Way, San Diego, CA 92130 (US). **KOEHLER, Michael F. T.**; 516 Georgia Avenue, Palo Alto, CA 94306 (US). **SCHWARZ, Jacob Bradley**; 7211 Balmoral Way, San Ramon, CA 94582 (US). **CRAWFORD, James John**; 1501 Church St #102, San Francisco, CA 94131 (US). **LY, Cuong Q.**; 801 Maple Ave, Burlingame, CA 94010 (US).

HANAN, Emily J.; 1904 Woodside Road, Redwood City, CA 94061 (US). **HU, Huiyong**; 34121 Susa Terrace, Fremont, CA 94555 (US). **YU, Zhiyong**; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN).

(74) **Agent:** **AFD CHINA INTELLECTUAL PROPERTY LAW OFFICE**; Suite B, 1601A, 8 Xue Qing Rd., Haidian, Beijing 100192 (CN).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:
— with international search report (Art. 21(3))

(54) **Title:** MACROCYCLIC BROAD SPECTRUM ANTIBIOTICS

(57) **Abstract:** Provided herein are antibacterial compounds, wherein the compounds in some embodiments have broad spectrum bioactivity. In various embodiments, the compounds act by inhibition of bacterial type 1 signal peptidase (SpsB), an essential protein in bacteria. Pharmaceutical compositions and methods for treatment using the compounds described herein are also provided.

WO 2017/084630 A1

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 358

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 358

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

MACROCYCLIC BROAD SPECTRUM ANTIBIOTICS

CROSS-REFERENCE

[0001] This application claims the benefit of PCT International Application

5 PCT/CN2016/105042, filed November 8, 2016, and PCT International Application PCT/CN2015/095166, filed November 20, 2015, both of which are incorporated by reference in their entirety.


BACKGROUND OF THE INVENTION

[0002] Antibiotic resistance is a serious and growing phenomenon in contemporary medicine
10 and has emerged as a major public health concern of the 21st century. Therefore, novel classes of broad-spectrum antibiotics, especially those that target novel mechanisms of action, are needed to treat multidrug-resistant pathogens.

SUMMARY OF THE INVENTION

15 [0003] Described herein are novel macrocyclic compounds for the treatment of microbial infections, such as for the treatment of bacterial infections. In various embodiments, the present disclosure provides lipopeptide macrocyclic compounds for the treatment of bacterial infections. In various embodiments, the present disclosure provides classes and subclasses of chemical compounds structurally related to arylomycin for the treatment of bacterial infections. In
20 various embodiments, the macrocyclic compounds act by inhibition of bacterial type 1 signal peptidase (SpsB), an essential protein in bacteria.

[0004] In one aspect described herein is a compound of Formula (I):



Formula (I);

25 wherein:

R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-$

- C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-N(R²³)C(O)(C₁-C₆)alkylNR²¹R²², or -(C₁-C₆)alkyl-C(O)N(R²³)(C₁-C₆)alkyl, or optionally substituted heterocycloalkyl;
- R³ is H, or -(C₁-C₆)alkyl;
- 5 R⁴ is H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OH, -(C₃-C₆)cycloalkyl, or -C(O)NH₂; or R³ and R⁴ are combined to form a heterocycloalkyl ring;
- R⁵ is H, or -(C₁-C₆)alkyl; or R⁴ and R⁵ and the carbon atom to which they are attached form a cyclopropyl ring;
- R⁶, R⁷, and R⁸ are each independently H, or -(C₁-C₆)alkyl;
- 10 R⁹ is H, -(C₁-C₆)alkyl, -(C₁-C₆)haloalkyl, or -(C₃-C₆)cycloalkyl;
- R¹⁰ is H, or -(C₁-C₆)alkyl;
- R¹¹ and R¹² are each independently H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)heteroalkyl-CO₂H, -(C₁-C₆)alkyl-S(O)-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-
- 15 N(H)CH=NH, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted -(C₁-C₆)alkyl-N(H)heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring, and R¹² is H;
- R¹⁵, R¹⁶, R¹⁷, and R¹⁸ are each independently H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;
- 20 X is optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₃-C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;
- Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;
- 25 Z is H, halogen, -NH₂, -CN, -CF₃, -CO₂H, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl-, -(C₂-C₁₂)alkynyl, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl-, -N(R²⁴)(C₁-C₁₂)alkyl-, -N(R²⁴)C(O)(C₁-C₁₂)alkyl-, optionally substituted -(C₃-C₇)cycloalkyl-, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;
- 30

each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)heteroalkyl$, $-(C_1-C_6)alkyl-CO_2H$, $-C(O)(C_1-C_6)alkyl$, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)alkyl$; or two R^{31} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R^{23} is independently H or $-(C_1-C_6)alkyl$;

each R^{24} is independently H or $-(C_1-C_6)alkyl$;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl$; or R^{25} and R^{26} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R^{27} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$, or optionally substituted $-(C_1-C_6)heteroalkyl$;


each R^{28} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$, or optionally substituted $-(C_1-C_6)heteroalkyl$;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[0005] In another aspect described herein is a compound of Formula (I'):



Formula (I');

wherein:

R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl-heterocycloalkyl$, $(C_1-C_6)heteroalkyl$ or optionally substituted heterocycloalkyl;
 R^3 is H, or $-(C_1-C_6)alkyl$;

R^4 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OH$, $-(C_3-C_6)cycloalkyl$, or $-C(O)NH_2$; or R^3 and R^4 are combined to form a heterocycloalkyl ring;

R^5 is H, or $-(C_1-C_6)alkyl$; or R^4 and R^5 and the carbon atom to which that are attached form a cyclopropyl ring;

5 R^6 , R^7 , and R^8 are each independently H, fluoro, hydroxyl, amino, optionally substituted alkyl or heteroalkyl or $-(C_1-C_6)alkyl$;

R^9 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;

R^{10} is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;

or R^9 and R^{10} are combined to form a heterocycloalkyl or cycloalkyl ring

10 R^{11} and R^{12} are each independently H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)heteroalkyl-CO_2H$, $-(C_1-C_6)alkyl-S(O)-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-N(H)CH=NH$, $-(C_1-C_6)alkyl-C(NH_2)=NH$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-N(H)SO_2NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(H)-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted $-(C_1-C_6)alkyl-N(H)heterocycloalkyl$, or $-(C_1-C_6)alkyl-heteroaryl$; or R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring, and R^{12} is H;

R^{15} , R^{16} , R^{17} , and R^{18} are each independently H, $-(C_1-C_6)alkyl$, $-(C_3-C_6)cycloalkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$;

20 X is optionally substituted $-(C_1-C_6)alkyl-$, $-(C_2-C_6)alkenyl-$, $-(C_2-C_6)alkynyl$, $-(C_3-C_7)cycloalkyl-$, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-O-(C_1-C_6)alkyl-$, $-N(R^{24})(C_1-C_6)alkyl-$, $-N(R^{24})(C_6-C_{10})aryl-$, or $-SO_2(C_1-C_6)alkyl-$;

25 Y is a bond, optionally substituted $-(C_1-C_6)alkyl-$, $-(C_2-C_6)alkenyl-$, $-(C_2-C_6)alkynyl$, $-(C_1-C_6)alkyl-N(R^{24})(C_1-C_6)alkyl-$, $-O-(C_1-C_6)alkyl-$, $-O(C_6-C_{10})aryl-$, $-N(R^{24})(C_1-C_6)alkyl-$, $-N(R^{24})SO_2(C_1-C_6)alkyl-$, $-N(R^{24})C(O)(C_1-C_6)alkyl-$, $-C(O)(C_1-C_6)alkyl-$, $-S(C_1-C_6)alkyl-$, $-SO_2(C_1-C_6)alkyl-$, $-C(O)NH(C_1-C_6)alkyl-$, $-(C_3-C_7)cycloalkyl-$, optionally substituted $-C(O)N(R^{24})aryl-$, optionally substituted $-N(R^{24})C(O)aryl-$, optionally substituted $-N(R^{24})SO_2aryl-$, optionally substituted aryl, or optionally substituted heteroaryl;

30 Z is H, halogen, $-NH_2$, $-CN$, $-CF_3$, $-CO_2H$, $-(C_1-C_{12})alkyl$, $-(C_2-C_{12})alkenyl$, $-(C_2-C_{12})alkynyl$, $-C(O)NR^{25}R^{26}$, $-O-(C_1-C_{12})alkyl$, $-N(R^{24})(C_1-C_{12})alkyl$, $-N(R^{24})C(O)(C_1-C_{12})alkyl$, optionally substituted $-(C_3-C_7)cycloalkyl$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted aryl, or optionally substituted heteroaryl;

each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)heteroalkyl$, $-(C_1-C_6)alkyl-CO_2H$, $-C(O)(C_1-C_6)alkyl$, $-C(O)O(C_1-C_6)alkyl$, $-C(O)O(C_1-C_6)haloalkyl$, $-C(=NH)(C_1-C_6)alkyl$, -

$C(=NH)N(R^{31})_2$, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)alkyl$; or two R^{31} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

5 each R^{23} is independently H or $-(C_1-C_6)alkyl$;

each R^{24} is independently H or $-(C_1-C_6)alkyl$;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl$; or R^{25} and R^{26} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{27} is independently halogen, $-NR^{23}R^{24}$, $-NC(O)R^{23}$, $-NC(O)NR^{23}R^{24}$, nitro, hydroxyl,

10 optionally substituted $-(C_1-C_6)alkyl$, optionally substituted $-(C_1-C_6)heteroalkyl$, $-(C_1-C_6)alkoxy$, $-C(O)(C_1-C_6)alkyl$, or $-S(O)_2(C_1-C_6)alkyl$;


each R^{28} is independently halogen, $-NR^{23}R^{24}$, $-NC(O)R^{23}$, $-NC(O)NR^{23}R^{24}$, nitro, hydroxyl, optionally substituted $-(C_1-C_6)alkyl$, optionally substituted $-(C_1-C_6)heteroalkyl$, $-(C_1-C_6)alkoxy$, $-C(O)(C_1-C_6)alkyl$, or $-S(O)_2(C_1-C_6)alkyl$;

15 p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[0006] In another embodiment described herein is a compound of Formula (I) or (I') having the structure of Formula (Ia):




20

Formula (Ia).

[0007] In another embodiment described herein is a compound of Formula (I), (I'), or (Ia) wherein R^6 , R^7 , and R^8 are H. In another embodiment described herein is a compound of Formula (I), (I'), or (Ia) wherein R^{15} and R^{16} are H.

25 **[0008]** In another embodiment described herein is a compound of Formula (I) or (I') having the structure of Formula (Ib):




Formula (Ib).

[0009] In one embodiment is a compound of Formula (I), (I'), (Ia), or (Ib) wherein R¹⁷ is -CH₃.

In a further embodiment is a compound of Formula (I), (I'), (Ia), or (Ib) wherein R¹⁸ is H. In

5 further embodiments is a compound of Formula (I), (I'), (Ia), or (Ib) wherein R⁵ is H. In a further embodiment is a compound of Formula (I), (I'), (Ia), or (Ib) wherein R⁴ is H. In a further embodiment is a compound of Formula (I), (I'), (Ia), or (Ib) wherein R⁴ is -(C₁-C₆)alkyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), or (Ib) wherein R⁴ is -(C₃-C₆)cycloalkyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), or (Ib) wherein R⁴ and R⁵ and the carbon atom to which they are attached form a cyclopropyl ring. In a further embodiment is a compound of Formula (I), (I'), (Ia), or (Ib) wherein R⁹ is -(C₁-C₆)alkyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), or (Ib) wherein R⁹ is -CH₃.

[0010] In another embodiment described herein is a compound of Formula (I) or (I') having the structure of Formula (Ic):




Formula (Ic).

[0011] In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R¹¹ is -(C₁-C₆)alkyl-OR²³. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R¹¹ is -CH₂CH₂OH. In a further embodiment is a compound of Formula (I), (I'),

20 (Ia), (Ib), or (Ic) wherein R¹¹ is -(C₁-C₆)alkyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R¹¹ is -(C₁-C₆)alkyl-NR²¹R²². In a further

embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^{11} is $-(C_1-C_6)alkyl-NH_2$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^{11} is $-CH_2NH_2$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^{11} is $-CH_2CH_2CH_2NH_2$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^{11} is $-CH_2CH_2CH_2CH_2NH_2$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^{11} is $-CH_2CH_2NH_2$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^1 and R^2 are each independently H, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^1 and R^2 are each independently $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^1 and R^2 are each $-CH_2CH_2NH_2$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$ and R^2 is H. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^1 is $-CH_2CH_2NH_2$ and R^2 is H. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^1 is H and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), or (Ic) wherein R^1 is H and R^2 is $-CH_2CH_2NH_2$.

[0012] In another embodiment described herein is a compound of Formula (I) or (I') having the structure of Formula (Id):



Formula (Id);

wherein R^{11} is $-CH_2NH_2$, $-CH_2CH_2NH_2$, or $-CH_2CH_2CH_2NH_2$.

[0013] In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is optionally substituted aryl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is optionally substituted phenyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is optionally substituted

heteroaryl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is heteroaryl which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is heteroaryl which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In a further

5 embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is pyridine which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is pyridine which is

10 unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is pyridine which is substituted once with methyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is

15 pyrimidine which is substituted once with methyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is pyrimidine which is substituted twice with methyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein X is optionally substituted $-(C_1-C_6)alkyl$ -. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Y is optionally substituted aryl. In a further


20 embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Y is optionally substituted phenyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Y is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Y is optionally substituted $-(C_1-C_6)alkyl$ -. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Y is $-O-(C_1-C_6)alkyl$ -. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Y is $-N(H)-(C_1-C_6)alkyl$ -. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Y is a bond. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is optionally substituted aryl. In

25


30

a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is optionally substituted phenyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is phenyl substituted once or twice with $-(C_1-C_8)$ alkyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is phenyl substituted once with n-butyl, isobutyl, or tert-butyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is phenyl substituted once with n-butyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is phenyl substituted once with isobutyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is phenyl substituted once with tert-butyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is optionally substituted $-(C_3-C_7)$ cycloalkyl. In a further embodiment is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), or (Id) wherein Z is halogen.

[0014] In a further embodiment is a compound of Formula (I), (Ia), (Ib), (Ic), or (Id) wherein -X-Y-Z is



[0015] In another aspect described herein is a compound of Formula (II):



Formula (II);

wherein:

- 5 R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, or $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, or optionally substituted heterocycloalkyl;
- 10 R^3 is H, or $-(C_1-C_6)alkyl$;
 R^4 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OH$, $-(C_3-C_6)cycloalkyl$, or $-C(O)NH_2$; or R^3 and R^4 are combined to form a heterocycloalkyl ring;
 R^5 is H, or $-(C_1-C_6)alkyl$; or R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring;
- 15 R^6 , R^7 , and R^8 are each independently H, or $-(C_1-C_6)alkyl$;
 R^9 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;
 R^{10} is H, or $-(C_1-C_6)alkyl$;
 R^{11} and R^{12} are each independently H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)heteroalkyl-CO_2H$, $-(C_1-C_6)alkyl-S(O)-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-N(H)CH=NH$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted $-(C_1-C_6)alkyl-N(H)heterocycloalkyl$, or $-(C_1-C_6)alkyl-heteroaryl$; or R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring, and R^{12} is H;
- 20 R^{13} and R^{14} are each independently H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-heterocycloalkyl$, or $-(C_1-C_6)alkyl-$

heteroaryl; or R¹³ and R¹⁹ are combined to form an optionally substituted heterocycloalkyl ring, and R¹⁴ is H;

R¹⁵, R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are each independently H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;

5 X is optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₃-C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;

10 Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

15 Z is H, halogen, -NH₂, -CN, -CF₃, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl-, -(C₂-C₁₂)alkynyl-, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl-, -N(R²⁴)(C₁-C₁₂)alkyl-, -N(R²⁴)C(O)(C₁-C₁₂)alkyl-, optionally substituted -(C₃-C₇)cycloalkyl-, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

20 each R²¹ and R²² is independently H, -(C₁-C₆)alkyl-, -(C₁-C₆)heteroalkyl-, -(C₁-C₆)alkyl-CO₂H, -C(O)(C₁-C₆)alkyl-, -C(O)N(R³¹)₂, or -SO₂N(R³¹)₂; or R²¹ and R²² and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R³¹ is independently H or -(C₁-C₆)alkyl; or two R³¹ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R²³ is independently H or -(C₁-C₆)alkyl;

25 each R²⁴ is independently H or -(C₁-C₆)alkyl;

each R²⁵ and R²⁶ is independently H or optionally substituted -(C₁-C₆)alkyl; or R²⁵ and R²⁶ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R²⁷ is independently halogen, optionally substituted -(C₁-C₆)alkyl, or optionally substituted -(C₁-C₆)heteroalkyl;

30 each R²⁸ is independently halogen, optionally substituted -(C₁-C₆)alkyl, or optionally substituted -(C₁-C₆)heteroalkyl;


n is 0 or 1;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[0016] In another aspect described herein is a compound of Formula (II'):



Formula (II');

5 wherein:

R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted $(C_1-C_6)heteroalkyl$ or optionally substituted heterocycloalkyl;

R^3 is H, or $-(C_1-C_6)alkyl$;

R^4 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OH$, $-(C_3-C_6)cycloalkyl$, or $-C(O)NH_2$; or R^3 and R^4 are combined to form a heterocycloalkyl ring;

R^5 is H, or $-(C_1-C_6)alkyl$; or R^4 and R^5 and the carbon atom to which that are attached form a cyclopropyl ring;

R^6 , R^7 , and R^8 are each independently H, or $-(C_1-C_6)alkyl$;

R^9 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;

R^{10} is H, or $-(C_1-C_6)alkyl$;

R^{11} and R^{12} are each independently H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)heteroalkyl-CO_2H$, $-(C_1-C_6)alkyl-S(O)-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-N(H)CH=NH$, $-(C_1-C_6)alkyl-C(NH_2)=NH$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-N(H)SO_2NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(H)-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted $-(C_1-C_6)alkyl-N(H)heterocycloalkyl$, or $-(C_1-C_6)alkyl-heteroaryl$; or R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring, and R^{12} is H;

R^{13} and R^{14} are each independently H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-heterocycloalkyl$, or $-(C_1-C_6)alkyl-heteroaryl$; or R^{13} and R^{19} are combined to form an optionally substituted heterocycloalkyl ring, and R^{14} is H;

R^{15} , R^{16} , R^{17} , R^{18} , and R^{19} are each independently H, $-(C_1-C_6)alkyl$, $-(C_3-C_6)cycloalkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$;

X is optionally substituted $-(C_1-C_6)alkyl-$, $-(C_2-C_6)alkenyl-$, $-(C_2-C_6)alkynyl$, $-(C_3-C_7)cycloalkyl-$, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-O-(C_1-C_6)alkyl-$, $-N(R^{24})(C_1-C_6)alkyl-$, $-N(R^{24})(C_6-C_{10})aryl-$, or $-SO_2(C_1-C_6)alkyl-$;

Y is a bond, optionally substituted $-(C_1-C_6)alkyl-$, $-(C_2-C_6)alkenyl-$, $-(C_2-C_6)alkynyl$, $-(C_1-C_6)alkyl-N(R^{24})(C_1-C_6)alkyl-$, $-O-(C_1-C_6)alkyl-$, $-O(C_6-C_{10})aryl-$, $-N(R^{24})(C_1-C_6)alkyl-$, $-N(R^{24})SO_2(C_1-C_6)alkyl-$, $-N(R^{24})C(O)(C_1-C_6)alkyl-$, $-C(O)(C_1-C_6)alkyl-$, $-S(C_1-C_6)alkyl-$, $-SO_2(C_1-C_6)alkyl-$, $-C(O)NH(C_1-C_6)alkyl-$, $-(C_3-C_7)cycloalkyl-$, optionally substituted $-C(O)N(R^{24})aryl-$, optionally substituted $-N(R^{24})C(O)aryl-$, optionally substituted $-N(R^{24})SO_2aryl-$, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, $-NH_2$, $-CN$, $-CF_3$, $-(C_1-C_{12})alkyl$, $-(C_2-C_{12})alkenyl$, $-(C_2-C_{12})alkynyl$, $-C(O)NR^{25}R^{26}$, $-O-(C_1-C_{12})alkyl$, $-N(R^{24})(C_1-C_{12})alkyl$, $-N(R^{24})C(O)(C_1-C_{12})alkyl$, optionally substituted $-(C_3-C_7)cycloalkyl$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted aryl, or optionally substituted heteroaryl;

each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)heteroalkyl$, $-(C_1-C_6)alkyl-CO_2H$, $-C(O)(C_1-C_6)alkyl$, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)alkyl$; or two R^{31} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{23} is independently H or $-(C_1-C_6)alkyl$;

each R^{24} is independently H or $-(C_1-C_6)alkyl$;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl$; or R^{25} and R^{26} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{27} is independently halogen, $-NR^{23}R^{24}$, $-NC(O)R^{23}$, $-NC(O)NR^{23}R^{24}$, nitro, hydroxyl, optionally substituted $-(C_1-C_6)alkyl$, optionally substituted $-(C_1-C_6)heteroalkyl$, $-(C_1-C_6)alkoxy$, $-C(O)(C_1-C_6)alkyl$, or $-S(O)_2(C_1-C_6)alkyl$;

each R^{28} is independently halogen, $-NR^{23}R^{24}$, $-NC(O)R^{23}$, $-NC(O)NR^{23}R^{24}$, nitro, hydroxyl, optionally substituted $-(C_1-C_6)alkyl$, optionally substituted $-(C_1-C_6)heteroalkyl$, $-(C_1-C_6)alkoxy$, $-C(O)(C_1-C_6)alkyl$, or $-S(O)_2(C_1-C_6)alkyl$;


n is 0 or 1;

5 p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[0017] In another embodiment described herein is a compound of Formula (II) or (II') having the structure of Formula (IIa):




10

Formula (IIa).

[0018] In another embodiment described herein is a compound of Formula (II), (II'), or (IIa) wherein R^6 , R^7 , and R^8 are H. In another embodiment described herein is a compound of Formula (II), (II'), or (IIa) wherein R^{15} and R^{16} are H.

15 **[0019]** In another embodiment described herein is a compound of Formula (II) or (II') having the structure of Formula (IIb):




Formula (IIb).

[0020] In another embodiment described herein is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^{18} is H. In another embodiment described herein is a compound of Formula (II),

20

(II'), (IIa), or (IIb) wherein R^{19} is H. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^{17} is $-\text{CH}_3$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^5 is H. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^4 is H. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^4 is $-(\text{C}_1\text{-C}_6)\text{alkyl}$. In another embodiment is a compound of (II), (II'), (IIa), or (IIb) wherein R^4 is $-(\text{C}_3\text{-C}_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^9 is $-(\text{C}_1\text{-C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^9 is $-\text{CH}_3$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^1 and R^2 are each independently H, or $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^1 and R^2 are each independently $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IIb) wherein R^1 and R^2 are each $-\text{CH}_2\text{CH}_2\text{NH}_2$.

[0021] In another embodiment described herein is a compound of Formula (II) or (II') having the structure of Formula (IIc):




Formula (IIc);

wherein R^1 and R^2 are each independently H or $-\text{CH}_2\text{CH}_2\text{NH}_2$.

[0022] In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{11} is $-(\text{C}_1\text{-C}_6)\text{alkyl-OR}^{23}$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{OH}$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{11} is $-(\text{C}_1\text{-C}_6)\text{alkyl}$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{11} is $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{11} is $-(\text{C}_1\text{-C}_6)\text{alkyl-NH}_2$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{11} is $-\text{CH}_2\text{NH}_2$. In another embodiment described herein is a

compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{13} is $-(\text{C}_1-\text{C}_6)\text{alkyl}-\text{OR}^{23}$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{13} is $-\text{CH}_2\text{OH}$. In another embodiment described herein is a compound of Formula (II), (II'), (IIa), (IIb), or (IIc) wherein R^{13} is $-\text{CH}_2\text{CH}_2\text{OH}$.


[0023] In another embodiment described herein is a compound of Formula (II) or (II') having the structure of Formula (IId):



Formula (IId).

[0024] In one embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^{17} is $-\text{CH}_3$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^5 is H. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^4 is H. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^4 is $-(\text{C}_1-\text{C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^4 is $-(\text{C}_3-\text{C}_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^9 is $-(\text{C}_1-\text{C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^9 is $-\text{CH}_3$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^1 and R^2 are each independently H, or $-(\text{C}_1-\text{C}_6)\text{alkyl}-\text{NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^1 and R^2 are each independently $-(\text{C}_1-\text{C}_6)\text{alkyl}-\text{NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (II), (II'), (IIa), or (IId) wherein R^1 and R^2 are each $-\text{CH}_2\text{CH}_2\text{NH}_2$.

[0025] In another embodiment described herein is a compound of Formula (II) or (II') having the structure of Formula (IIe):




Formula (IIe);

wherein R^1 and R^2 are each independently H or $-\text{CH}_2\text{CH}_2\text{NH}_2$.

[0026] In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein X is optionally substituted aryl. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein X is optionally substituted phenyl. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein X is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein X is optionally substituted $-(\text{C}_1\text{-C}_6)\text{alkyl-}$. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Y is optionally substituted aryl. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Y is optionally substituted phenyl. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Y is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Y is optionally substituted $-(\text{C}_1\text{-C}_6)\text{alkyl-}$. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Y is $-\text{O}-(\text{C}_1\text{-C}_6)\text{alkyl-}$. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Y is $-\text{N}(\text{H})-(\text{C}_1\text{-C}_6)\text{alkyl-}$. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Y is a bond. In a further embodiment is a compound of Formula (II), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Z is $-(\text{C}_1\text{-C}_6)\text{alkyl}$. In a further embodiment is a compound of Formula (II), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Z is optionally substituted aryl. In a further embodiment is a compound of Formula (II), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Z is optionally substituted phenyl. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Z is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Z is optionally substituted $-(\text{C}_3\text{-C}_7)\text{cycloalkyl}$. In a further

embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein Z is halogen.

[0027] In a further embodiment is a compound of Formula (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) wherein -X-Y-Z is



[0028] In another aspect is a pharmaceutical composition comprising a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), or a pharmaceutically acceptable salt, pharmaceutically acceptable solvate, or pharmaceutically acceptable prodrug thereof, and a pharmaceutically acceptable excipient thereof.

[0029] In another aspect is the use of a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (I'), (IIa), (IIb), (IIc), (IId), or (IIe), or a pharmaceutically acceptable salt, pharmaceutically acceptable solvate, or pharmaceutically acceptable prodrug thereof, for the preparation of a medicament for the treatment of a bacterial infection in a patient.

[0030] In another embodiment is a method for treating a bacterial infection in a mammal comprising administering to the mammal a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), or a pharmaceutically acceptable salt, pharmaceutically acceptable solvate, or pharmaceutically acceptable prodrug thereof, at a frequency and for a duration sufficient to provide a beneficial effect to the mammal. In another embodiment, the bacterial infection is an infection involving *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Pseudomonas acidovorans*, *Pseudomonas alcaligenes*, *Pseudomonas putida*, *Stenotrophomonas maltophilia*, *Burkholderia cepacia*, *Aeromonas hydrophilia*, *Escherichia coli*, *Citrobacter*

freundii, Salmonella typhimurium, Salmonella typhi, Salmonella paratyphi, Salmonella enteritidis, Shigella dysenteriae, Shigella flexneri, Shigella sonnei, Enterobacter cloacae, Enterobacter aerogenes, Klebsiella pneumoniae, Klebsiella oxytoca, Serratia marcescens, Francisella tularensis, Morganella morganii, Proteus mirabilis, Proteus vulgaris, Providencia alcalifaciens, Providencia rettgeri, Providencia stuartii, Acinetobacter baumannii, Acinetobacter calcoaceticus, Acinetobacter haemolyticus, Yersinia enterocolitica, Yersinia pestis, Yersinia pseudotuberculosis, Yersinia intermedia, Bordetella pertussis, Bordetella parapertussis, Bordetella bronchiseptica, Haemophilus influenzae, Haemophilus parainfluenzae, Haemophilus haemolyticus, Haemophilus parahaemolyticus, Haemophilus ducreyi, Pasteurella multocida, Pasteurella haemolytica, Branhamella catarrhalis, Helicobacter pylori, Campylobacter fetus, Campylobacter jejuni, Campylobacter coli, Borrelia burgdorferi, Vibrio cholerae, Vibrio parahaemolyticus, Legionella pneumophila, Listeria monocytogenes, Neisseria gonorrhoeae, Neisseria meningitidis, Kingella, Moraxella, Gardnerella vaginalis, Bacteroides fragilis, Bacteroides distasonis, Bacteroides 3452A homology group, Bacteroides vulgatus, Bacteroides ovalus, Bacteroides thetaiotaomicron, Bacteroides uniformis, Bacteroides eggerthii, Bacteroides splanchnicus, Clostridium difficile, Mycobacterium tuberculosis, Mycobacterium avium, Mycobacterium intracellulare, Mycobacterium leprae, Corynebacterium diphtheriae, Corynebacterium ulcerans, Streptococcus pneumoniae, Streptococcus agalactiae, Streptococcus pyogenes, Enterococcus faecalis, Enterococcus faecium, Staphylococcus aureus, Staphylococcus epidermidis, Staphylococcus saprophyticus, Staphylococcus intermedius, Staphylococcus hyicus subsp. hyicus, Staphylococcus haemolyticus, Staphylococcus hominis, or Staphylococcus saccharolyticus.

[0031] In another embodiment the bacterial infection is an infection involving a Gram-negative bacteria. In another embodiment, administering comprises a topical administration.

[0032] In a further embodiment are methods of treating a mammal in need of such treatment comprising administering to the mammal a second therapeutic agent. In another embodiment, the second therapeutic agent is not an SpsB inhibitor. In another embodiment, the second therapeutic agent is an aminoglycoside antibiotic, fluoroquinolone antibiotic, β -lactam antibiotic, macrolide antibiotic, glycopeptide antibiotic, rifampicin, chloramphenicol, fluoramphenicol, colistin, mupirocin, bacitracin, daptomycin, or linezolid. In another embodiment, the second therapeutic agent is a β -lactam antibiotic. In another embodiment, the β -lactam antibiotic is selected from penicillins, monobactams, cephalosporins, cephamycins, and carbapenems. In another embodiment, the β -lactam antibiotic is selected from Azlocillin, Amoxicillin, Ampicillin, Doripenem, Meropenem, Biapenem, Cefamandole, Imipenem,

Mezlocillin, Cefmetazole, Cefprozil, Piperacillin/tazobactam, Carbenicillin, Cefaclor, Cephalothin, Ertapenem, Cefazolin, Cefepime, Cefonicid, Cefoxitin, Ceftazidime, Oxacillin, Cefdinir, Cefixime, Cefotaxime, Cefotetan, Cefpodoxime, Ceftizoxime, Ceftriaxone, Faropenem, Mecillinam, Methicillin, Moxalactam, Ticarcillin, Tomopenem, Ceftobiprole, 5 Ceftaroline, Flomoxef, Cefiprome, and Cefozopran. A further embodiment comprises administering a β -lactamase inhibitor.

INCORPORATION BY REFERENCE

10 [0033] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE FIGURE

15 [0034] Figure 1 depicts the activity of Compound 135 in a neutropenic thigh infection model.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0035] As used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise.

20 [0036] The term "about" as used herein, when referring to a numerical value or range, allows for a degree of variability in the value or range, for example, within 10%, or within 5% of a stated value or of a stated limit of a range.

[0037] All percent compositions are given as weight-percentages, unless otherwise stated.

25 [0038] All average molecular weights of polymers are weight-average molecular weights, unless otherwise specified.

[0039] As used herein, "individual" (as in the subject of the treatment) means both mammals and non-mammals. Mammals include, for example, humans; non-human primates, e.g. apes and monkeys; and non-primates, e.g. dogs, cats, cattle, horses, sheep, and goats. Non-mammals include, for example, fish and birds.

30 [0040] The term "disease" or "disorder" or "malcondition" are used interchangeably, and are used to refer to diseases or conditions wherein a bacterial SPase plays a role in the biochemical mechanisms involved in the disease or malcondition such that a therapeutically beneficial effect can be achieved by acting on the enzyme. "Acting on" SPase can include binding to SPase and/or inhibiting the bioactivity of an SPase.

[0041] The expression “effective amount”, when used to describe therapy to an individual suffering from a disorder, refers to the amount of a compound described herein that is effective to inhibit or otherwise act on SPase in the individual's tissues wherein SPase involved in the disorder is active, wherein such inhibition or other action occurs to an extent sufficient to produce a beneficial therapeutic effect.

[0042] "Substantially" as the term is used herein means completely or almost completely; for example, a composition that is "substantially free" of a component either has none of the component or contains such a trace amount that any relevant functional property of the composition is unaffected by the presence of the trace amount, or a compound is "substantially pure" if there are only negligible traces of impurities present.

[0043] “Treating” or "treatment" within the meaning herein refers to an alleviation of symptoms associated with a disorder or disease, or inhibition of further progression or worsening of those symptoms, or prevention or prophylaxis of the disease or disorder, or curing the disease or disorder. Similarly, as used herein, an “effective amount” or a “therapeutically effective amount” of a compound refers to an amount of the compound that alleviates, in whole or in part, symptoms associated with the disorder or condition, or halts or slows further progression or worsening of those symptoms, or prevents or provides prophylaxis for the disorder or condition. In particular, a "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount is also one in which any toxic or detrimental effects of compounds described herein are outweighed by the therapeutically beneficial effects.

[0044] By "chemically feasible" is meant a bonding arrangement or a compound where the generally understood rules of organic structure are not violated; for example a structure within a definition of a claim that would contain in certain situations a pentavalent carbon atom that would not exist in nature would be understood to not be within the claim. The structures disclosed herein, in all of their embodiments are intended to include only "chemically feasible" structures, and any recited structures that are not chemically feasible, for example in a structure shown with variable atoms or groups, are not intended to be disclosed or claimed herein.

[0045] When a substituent is specified to be an atom or atoms of specified identity, "or a bond", a configuration is referred to when the substituent is "a bond" that the groups that are immediately adjacent to the specified substituent are directly connected to each other in a chemically feasible bonding configuration.

[0046] All chiral, diastereomeric, racemic forms of a structure are intended, unless a particular stereochemistry or isomeric form is specifically indicated. Compounds described herein can

include enriched or resolved optical isomers at any or all asymmetric atoms as are apparent from the depictions, at any degree of enrichment. Both racemic and diastereomeric mixtures, as well as the individual optical isomers can be isolated or synthesized so as to be substantially free of their enantiomeric or diastereomeric partners, and these are all within the scope of the invention.

5 [0047] The inclusion of an isotopic form of one or more atoms in a molecule that is different from the naturally occurring isotopic distribution of the atom in nature is referred to as an "isotopically labeled form" of the molecule. All isotopic forms of atoms are included as options in the composition of any molecule, unless a specific isotopic form of an atom is indicated. For example, any hydrogen atom or set thereof in a molecule can be any of the isotopic forms of
10 hydrogen, i.e., protium (^1H), deuterium (^2H), or tritium (^3H) in any combination. Similarly, any carbon atom or set thereof in a molecule can be any of the isotopic form of carbons, such as ^{11}C , ^{12}C , ^{13}C , or ^{14}C , or any nitrogen atom or set thereof in a molecule can be any of the isotopic forms of nitrogen, such as ^{13}N , ^{14}N , or ^{15}N . A molecule can include any combination of isotopic forms in the component atoms making up the molecule, the isotopic form of every atom forming
15 the molecule being independently selected. In a multi-molecular sample of a compound, not every individual molecule necessarily has the same isotopic composition. For example, a sample of a compound can include molecules containing various different isotopic compositions, such as in a tritium or ^{14}C radiolabeled sample where only some fraction of the set of molecules making up the macroscopic sample contains a radioactive atom. It is also
20 understood that many elements that are not artificially isotopically enriched themselves are mixtures of naturally occurring isotopic forms, such as ^{14}N and ^{15}N , ^{32}S and ^{34}S , and so forth. A molecule as recited herein is defined as including isotopic forms of all its constituent elements at each position in the molecule. As is well known in the art, isotopically labeled compounds can be prepared by the usual methods of chemical synthesis, except substituting an isotopically
25 labeled precursor molecule. The isotopes, radiolabeled or stable, can be obtained by any method known in the art, such as generation by neutron absorption of a precursor nuclide in a nuclear reactor, by cyclotron reactions, or by isotopic separation such as by mass spectrometry. The isotopic forms are incorporated into precursors as required for use in any particular synthetic route. For example, ^{14}C and ^3H can be prepared using neutrons generated in a nuclear reactor.
30 Following nuclear transformation, ^{14}C and ^3H are incorporated into precursor molecules, followed by further elaboration as needed.

[0048] The term "amino protecting group" or "N-protected" as used herein refers to those groups intended to protect an amino group against undesirable reactions during synthetic procedures and which can later be removed to reveal the amine. Commonly used amino

protecting groups are disclosed in Protective Groups in Organic Synthesis, Greene, T.W.; Wuts, P. G. M., John Wiley & Sons, New York, NY, (3rd Edition, 1999). Amino protecting groups include acyl groups such as formyl, acetyl, propionyl, pivaloyl, t-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, o-nitrophenoxyacetyl, α -chlorobutyryl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, 4-nitrobenzoyl, and the like; sulfonyl groups such as benzenesulfonyl, p-toluenesulfonyl and the like; alkoxy- or aryloxy-carbonyl groups (which form urethanes with the protected amine) such as benzyloxycarbonyl (Cbz), p-chlorobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxycarbonyl, t-butyloxycarbonyl (Boc), diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl (Alloc), 2,2,2-trichloroethoxycarbonyl, 2-trimethylsilylethylloxycarbonyl (Teoc), phenoxycarbonyl, 4-nitrophenoxycarbonyl, fluorenyl-9-methoxycarbonyl (Fmoc), cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl, phenylthiocarbonyl and the like; aralkyl groups such as benzyl, triphenylmethyl, benzyloxymethyl and the like; and silyl groups such as trimethylsilyl and the like. Amine protecting groups also include cyclic amino protecting groups such as phthaloyl and dithiosuccinimidyl, which incorporate the amino nitrogen into a heterocycle. Typically, amino protecting groups include formyl, acetyl, benzoyl, pivaloyl, t-butylacetyl, phenylsulfonyl, Alloc, Teoc, benzyl, Fmoc, Boc and Cbz. It is well within the skill of the ordinary artisan to select and use the appropriate amino protecting group for the synthetic task at hand.

[0049] The term "hydroxyl protecting group" or "O-protected" as used herein refers to those groups intended to protect an OH group against undesirable reactions during synthetic procedures and which can later be removed to reveal the amine. Commonly used hydroxyl protecting groups are disclosed in Protective Groups in Organic Synthesis, Greene, T.W.; Wuts, P. G. M., John Wiley & Sons, New York, NY, (3rd Edition, 1999). Hydroxyl protecting groups include acyl groups such as formyl, acetyl, propionyl, pivaloyl, t-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, o-nitrophenoxyacetyl, α -chlorobutyryl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, 4-nitrobenzoyl, and the like; sulfonyl groups such as benzenesulfonyl, p-toluenesulfonyl and the like; acyloxy groups (which form urethanes with the protected amine) such as benzyloxycarbonyl (Cbz), p-chlorobenzyloxycarbonyl,

p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxycarbonyl, t-butyloxycarbonyl (Boc), diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl (Alloc), 2,2,2-trichloroethoxycarbonyl, 2-trimethylsilylethylloxycarbonyl (Teoc), phenoxycarbonyl, 4-nitrophenoxycarbonyl, fluorenyl-9-methoxycarbonyl (Fmoc), cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl, phenylthiocarbonyl and the like; aralkyl groups such as benzyl, triphenylmethyl, benzyloxymethyl and the like; and silyl groups such as trimethylsilyl and the like. It is well within the skill of the ordinary artisan to select and use the appropriate hydroxyl protecting group for the synthetic task at hand.

[0050] In general, “substituted” refers to an organic group as defined herein in which one or more bonds to a hydrogen atom contained therein are replaced by one or more bonds to a non-hydrogen atom such as, but not limited to, a halogen (i.e., F, Cl, Br, and I); an oxygen atom in groups such as hydroxyl groups, alkoxy groups, aryloxy groups, aralkyloxy groups, oxo(carbonyl) groups, carboxyl groups including carboxylic acids, carboxylates, and carboxylate esters; a sulfur atom in groups such as thiol groups, alkyl and aryl sulfide groups, sulfoxide groups, sulfone groups, sulfonyl groups, and sulfonamide groups; a nitrogen atom in groups such as amines, hydroxylamines, nitriles, nitro groups, N-oxides, hydrazides, azides, and enamines; and other heteroatoms in various other groups. Non-limiting examples of substituents that can be bonded to a substituted carbon (or other) atom include F, Cl, Br, I, OR', OC(O)N(R')₂, CN, NO, NO₂, ONO₂, azido, CF₃, OCF₃, R', O (oxo), S (thiono), C(O), S(O), methylenedioxy, ethylenedioxy, N(R')₂, SR', SOR', SO₂R', SO₂N(R')₂, SO₃R', C(O)R', C(O)C(O)R', C(O)CH₂C(O)R', C(S)R', C(O)OR', OC(O)R', C(O)N(R')₂, OC(O)N(R')₂, C(S)N(R')₂, (CH₂)₀₋₂N(R')C(O)R', (CH₂)₀₋₂N(R')N(R')₂, N(R')N(R')C(O)R', N(R')N(R')C(O)OR', N(R')N(R')CON(R')₂, N(R')SO₂R', N(R')SO₂N(R')₂, N(R')C(O)OR', N(R')C(O)R', N(R')C(S)R', N(R')C(O)N(R')₂, N(R')C(S)N(R')₂, N(COR')COR', N(OR')R', C(=NH)N(R')₂, C(O)N(OR')R', or C(=NOR')R' wherein R' can be hydrogen or a carbon-based moiety, and wherein the carbon-based moiety can itself be further substituted.

[0051] When a substituent is monovalent, such as, for example, F or Cl, it is bonded to the atom it is substituting by a single bond. When a substituent is more than monovalent, such as O, which is divalent, it can be bonded to the atom it is substituting by more than one bond, i.e., a

divalent substituent is bonded by a double bond; for example, a C substituted with O forms a carbonyl group, C=O, which can also be written as "CO", "C(O)", or "C(=O)", wherein the C and the O are double bonded. When a carbon atom is substituted with a double-bonded oxygen (=O) group, the oxygen substituent is termed an "oxo" group. When a divalent substituent such as NR is double-bonded to a carbon atom, the resulting C(=NR) group is termed an "imino" group. When a divalent substituent such as S is double-bonded to a carbon atom, the results C(=S) group is termed a "thiocarbonyl" group.

[0052] Alternatively, a divalent substituent such as O, S, C(O), S(O), or S(O)₂ can be connected by two single bonds to two different carbon atoms. For example, O, a divalent substituent, can be bonded to each of two adjacent carbon atoms to provide an epoxide group, or the O can form a bridging ether group, termed an "oxy" group, between adjacent or non-adjacent carbon atoms, for example bridging the 1,4-carbons of a cyclohexyl group to form a [2.2.1]-oxabicyclo system. Further, any substituent can be bonded to a carbon or other atom by a linker, such as (CH₂)_n or (CR'₂)_n wherein n is 1, 2, 3, or more, and each R' is independently selected.

[0053] C(O) and S(O)₂ groups can be bound to one or two heteroatoms, such as nitrogen, rather than to a carbon atom. For example, when a C(O) group is bound to one carbon and one nitrogen atom, the resulting group is called an "amide" or "carboxamide." When a C(O) group is bound to two nitrogen atoms, the functional group is termed a urea. When a S(O)₂ group is bound to one carbon and one nitrogen atom, the resulting unit is termed a "sulfonamide." When a S(O)₂ group is bound to two nitrogen atoms, the resulting unit is termed a "sulfamate."

[0054] Substituted alkyl, alkenyl, alkynyl, cycloalkyl, and cycloalkenyl groups as well as other substituted groups also include groups in which one or more bonds to a hydrogen atom are replaced by one or more bonds, including double or triple bonds, to a carbon atom, or to a heteroatom such as, but not limited to, oxygen in carbonyl (oxo), carboxyl, ester, amide, imide, urethane, and urea groups; and nitrogen in imines, hydroxyimines, oximes, hydrazones, amidines, guanidines, and nitriles.

[0055] Substituted ring groups such as substituted cycloalkyl, aryl, heterocyclyl and heteroaryl groups also include rings and fused ring systems in which a bond to a hydrogen atom is replaced with a bond to a carbon atom. Therefore, substituted cycloalkyl, aryl, heterocyclyl and heteroaryl groups can also be substituted with alkyl, alkenyl, and alkynyl groups as defined herein.

[0056] By a "ring system" as the term is used herein is meant a moiety comprising one, two, three or more rings, which can be substituted with non-ring groups or with other ring systems, or both, which can be fully saturated, partially unsaturated, fully unsaturated, or aromatic, and

when the ring system includes more than a single ring, the rings can be fused, bridging, or spirocyclic. By "spirocyclic" is meant the class of structures wherein two rings are fused at a single tetrahedral carbon atom, as is well known in the art.

[0057] As to any of the groups described herein, which contain one or more substituents, it is understood, of course, that such groups do not contain any substitution or substitution patterns which are sterically impractical and/or synthetically non-feasible. In addition, the compounds of this disclosed subject matter include all stereochemical isomers arising from the substitution of these compounds.

[0058] Selected substituents within the compounds described herein are present to a recursive degree. In this context, "recursive substituent" means that a substituent may recite another instance of itself or of another substituent that itself recites the first substituent. Because of the recursive nature of such substituents, theoretically, a large number may be present in any given claim. One of ordinary skill in the art of medicinal chemistry and organic chemistry understands that the total number of such substituents is reasonably limited by the desired properties of the compound intended. Such properties include, by way of example and not limitation, physical properties such as molecular weight, solubility or log P, application properties such as activity against the intended target, and practical properties such as ease of synthesis.

[0059] Recursive substituents are an intended aspect of the disclosed subject matter. One of ordinary skill in the art of medicinal and organic chemistry understands the versatility of such substituents. To the degree that recursive substituents are present in a claim of the disclosed subject matter, the total number should be determined as set forth above.

[0060] Alkyl groups include straight chain and branched alkyl groups and cycloalkyl groups having from 1 to about 20 carbon atoms, and typically from 1 to 12 carbons or, in some embodiments, from 1 to 8 carbon atoms. Examples of straight chain alkyl groups include those with from 1 to 8 carbon atoms such as methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl, and n-octyl groups. Examples of branched alkyl groups include, but are not limited to, isopropyl, iso-butyl, sec-butyl, t-butyl, neopentyl, isopentyl, and 2,2-dimethylpropyl groups. As used herein, the term "alkyl" encompasses n-alkyl, isoalkyl, and anteisoalkyl groups as well as other branched chain forms of alkyl. Representative substituted alkyl groups can be substituted one or more times with any of the groups listed above, for example, amino, hydroxy, cyano, carboxy, nitro, thio, alkoxy, and halogen groups.

[0061] The term "alkylene" means a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of one to six carbon atoms

unless otherwise stated, such as methylene, ethylene, propylene, 1-methylpropylene, 2-methylpropylene, butylene, pentylene, and the like.

[0062] The term “carbonyl” means C=O.

[0063] The terms “carboxy” and “hydroxycarbonyl” mean COOH.

5 **[0064]** Cycloalkyl groups are cyclic alkyl groups such as, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl groups. In some embodiments, the cycloalkyl group can have 3 to about 8-12 ring members, whereas in other embodiments the number of ring carbon atoms range from 3 to 4, 5, 6, or 7. Cycloalkyl groups further include polycyclic cycloalkyl groups such as, but not limited to, norbornyl, adamantyl, bornyl,
10 camphenyl, isocamphenyl, and carenyl groups, and fused rings such as, but not limited to, decalanyl, and the like. Cycloalkyl groups also include rings that are substituted with straight or branched chain alkyl groups as defined above. Representative substituted cycloalkyl groups can be mono-substituted or substituted more than once, such as, but not limited to, 2,2-, 2,3-, 2,4-
15 2,5- or 2,6-disubstituted cyclohexyl groups or mono-, di- or tri-substituted norbornyl or cycloheptyl groups, which can be substituted with, for example, amino, hydroxy, cyano, carboxy, nitro, thio, alkoxy, and halogen groups. The term “cycloalkenyl” alone or in combination denotes a cyclic alkenyl group.

[0065] The terms “carbocyclic,” “carbocyclyl,” and “carbocycle” denote a ring structure wherein the atoms of the ring are carbon, such as a cycloalkyl group or an aryl group. In some
20 embodiments, the carbocycle has 3 to 8 ring members, whereas in other embodiments the number of ring carbon atoms is 4, 5, 6, or 7. Unless specifically indicated to the contrary, the carbocyclic ring can be substituted with as many as N-1 substituents wherein N is the size of the carbocyclic ring with, for example, alkyl, alkenyl, alkynyl, amino, aryl, hydroxy, cyano, carboxy, heteroaryl, heterocyclyl, nitro, thio, alkoxy, and halogen groups, or other groups as are
25 listed above. A carbocyclyl ring can be a cycloalkyl ring, a cycloalkenyl ring, or an aryl ring. A carbocyclyl can be monocyclic or polycyclic, and if polycyclic each ring can be independently be a cycloalkyl ring, a cycloalkenyl ring, or an aryl ring.

[0066] (Cycloalkyl)alkyl groups, also denoted cycloalkylalkyl, are alkyl groups as defined above in which a hydrogen or carbon bond of the alkyl group is replaced with a bond to a
30 cycloalkyl group as defined above.

[0067] Alkenyl groups include straight and branched chain and cyclic alkyl groups as defined above, except that at least one double bond exists between two carbon atoms. Thus, alkenyl groups have from 2 to about 20 carbon atoms, and typically from 2 to 12 carbons or, in some embodiments, from 2 to 8 carbon atoms. Examples include, but are not limited to

vinyl, $-\text{CH}=\text{CH}(\text{CH}_3)$, $-\text{CH}=\text{C}(\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{C}(\text{CH}_3)=\text{CH}(\text{CH}_3)$, $-\text{C}(\text{CH}_2\text{CH}_3)=\text{CH}_2$, cyclohexenyl, cyclopentenyl, cyclohexadienyl, butadienyl, pentadienyl, and hexadienyl among others.

[0068] Cycloalkenyl groups include cycloalkyl groups having at least one double bond between 2 carbons. Thus for example, cycloalkenyl groups include but are not limited to cyclohexenyl, cyclopentenyl, and cyclohexadienyl groups. Cycloalkenyl groups can have from 3 to about 8-12 ring members, whereas in other embodiments the number of ring carbon atoms range from 3 to 5, 6, or 7. Cycloalkyl groups further include polycyclic cycloalkyl groups such as, but not limited to, norbornyl, adamantyl, bornyl, camphenyl, isocamphenyl, and carenyl groups, and fused rings such as, but not limited to, decalinyl, and the like, provided they include at least one double bond within a ring. Cycloalkenyl groups also include rings that are substituted with straight or branched chain alkyl groups as defined above.

[0069] (Cycloalkenyl)alkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of the alkyl group is replaced with a bond to a cycloalkenyl group as defined above.

[0070] Alkynyl groups include straight and branched chain alkyl groups, except that at least one triple bond exists between two carbon atoms. Thus, alkynyl groups have from 2 to about 20 carbon atoms, and typically from 2 to 12 carbons or, in some embodiments, from 2 to 8 carbon atoms. Examples include, but are not limited to $-\text{C}\equiv\text{CH}$, $-\text{C}\equiv\text{C}(\text{CH}_3)$, -

$\text{C}\equiv\text{C}(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{C}\equiv\text{CH}$, $-\text{CH}_2\text{C}\equiv\text{C}(\text{CH}_3)$, and $-\text{CH}_2\text{C}\equiv\text{C}(\text{CH}_2\text{CH}_3)$ among others.

[0071] The term "heteroalkyl" by itself or in combination with another term means, unless otherwise stated, a stable straight or branched chain alkyl group consisting of the stated number of carbon atoms and one or two heteroatoms selected from the group consisting of O, N, and S, and wherein the nitrogen and sulfur atoms may be optionally oxidized and the nitrogen heteroatom may be optionally quaternized. The heteroatom(s) may be placed at any position of the heteroalkyl group, including between the rest of the heteroalkyl group and the fragment to which it is attached, as well as attached to the most distal carbon atom in the heteroalkyl group. Examples

include: $-\text{O}-\text{CH}_2-\text{CH}_2-\text{CH}_3$, $-\text{CH}_2-\text{CH}_2\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_3$, $-\text{CH}_2-\text{S}-\text{CH}_2-\text{CH}_3$, $-\text{CH}_2\text{CH}_2-\text{S}(=\text{O})-\text{CH}_3$, and $-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_3$. Up to two heteroatoms may be consecutive, such as, for example, $-\text{CH}_2-\text{NH}-\text{OCH}_3$, or $-\text{CH}_2-\text{CH}_2-\text{S}-\text{S}-\text{CH}_3$.

[0072] A "heterocycloalkyl" ring is a cycloalkyl ring containing at least one heteroatom. A heterocycloalkyl ring can also be termed a "heterocyclyl," described below.

[0073] The term “heteroalkenyl” by itself or in combination with another term means, unless otherwise stated, a stable straight or branched chain monounsaturated or di-unsaturated hydrocarbon group consisting of the stated number of carbon atoms and one or two heteroatoms selected from the group consisting of O, N, and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. Up to two heteroatoms may be placed consecutively. Examples include -CH=CH-O-CH₃, -CH=CH-CH₂-OH, -CH₂-CH=N-OCH₃, -CH=CH-N(CH₃)-CH₃, -CH₂-CH=CH-CH₂-SH, and -CH=CH-O-CH₂CH₂-O-CH₃.

[0074] Aryl groups are cyclic aromatic hydrocarbons that do not contain heteroatoms in the ring. Thus aryl groups include, but are not limited to, phenyl, azulenyl, heptalenyl, biphenyl, indacenyl, fluorenyl, phenanthrenyl, triphenylenyl, pyrenyl, naphthacenyl, chrysenyl, biphenylenyl, anthracenyl, and naphthyl groups. In some embodiments, aryl groups contain about 6 to about 14 carbons in the ring portions of the groups. Aryl groups can be unsubstituted or substituted, as defined above. Representative substituted aryl groups can be mono-substituted or substituted more than once, such as, but not limited to, 2-, 3-, 4-, 5-, or 6-substituted phenyl or 2-8 substituted naphthyl groups, which can be substituted with carbon or non-carbon groups such as those listed above.

[0075] Aralkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to an aryl group as defined above. Representative aralkyl groups include benzyl and phenylethyl groups and fused (cycloalkylaryl)alkyl groups such as 4-ethyl-indanyl. Aralkenyl group are alkenyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to an aryl group as defined above.

[0076] Heterocyclyl groups or the term “heterocyclyl” includes aromatic and non-aromatic ring compounds containing 3 or more ring members, of which, one or more is a heteroatom such as, but not limited to, N, O, and S. Thus a heterocyclyl can be a heterocycloalkyl, or a heteroaryl, or if polycyclic, any combination thereof. In some embodiments, heterocyclyl groups include 3 to about 20 ring members, whereas other such groups have 3 to about 15 ring members. A heterocyclyl group designated as a C₂-heterocyclyl can be a 5-ring with two carbon atoms and three heteroatoms, a 6-ring with two carbon atoms and four heteroatoms and so forth. Likewise a C₄-heterocyclyl can be a 5-ring with one heteroatom, a 6-ring with two heteroatoms, and so forth. The number of carbon atoms plus the number of heteroatoms sums up to equal the total number of ring atoms. A heterocyclyl ring can also include one or more double bonds. A heteroaryl ring is an embodiment of a heterocyclyl group. The phrase “heterocyclyl group” includes fused ring species including those comprising fused aromatic and non-aromatic groups.

For example, a dioxolanyl ring and a benzodioxolanyl ring system (methylenedioxyphenyl ring system) are both heterocyclyl groups within the meaning herein. The phrase also includes polycyclic ring systems containing a heteroatom such as, but not limited to, quinuclidyl.

Heterocyclyl groups can be unsubstituted, or can be substituted as discussed above.

5 Heterocyclyl groups include, but are not limited to, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, pyrrolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, pyridinyl, thiophenyl, benzothiophenyl, benzofuranyl, dihydrobenzofuranyl, indolyl, dihydroindolyl, azaindolyl, indazolyl, benzimidazolyl, azabenzimidazolyl, benzoxazolyl, benzothiazolyl, benzothiadiazolyl, imidazopyridinyl, isoxazolopyridinyl, thianaphthalenyl, purinyl, xanthinyl, adeninyl, guaninyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, quinoxalinyl, and
10 quinazolinyl groups. Representative substituted heterocyclyl groups can be mono-substituted or substituted more than once, such as, but not limited to, piperidinyl or quinolinyl groups, which are 2-, 3-, 4-, 5-, or 6-substituted, or disubstituted with groups such as those listed above.

[0077] Heteroaryl groups are aromatic ring compounds containing 5 or more ring members, of
15 which, one or more is a heteroatom such as, but not limited to, N, O, and S; for instance, heteroaryl rings can have 5 to about 8-12 ring members. A heteroaryl group is a variety of a heterocyclyl group that possesses an aromatic electronic structure. A heteroaryl group designated as a C₂-heteroaryl can be a 5-ring with two carbon atoms and three heteroatoms, a 6-ring with two carbon atoms and four heteroatoms and so forth. Likewise a C₄-heteroaryl can be
20 a 5-ring with one heteroatom, a 6-ring with two heteroatoms, and so forth. The number of carbon atoms plus the number of heteroatoms sums up to equal the total number of ring atoms. Heteroaryl groups include, but are not limited to, groups such as pyrrolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, pyridinyl, thiophenyl, benzothiophenyl, benzofuranyl, indolyl, azaindolyl, indazolyl, benzimidazolyl, azabenzimidazolyl, benzoxazolyl, benzothiazolyl, benzothiadiazolyl, imidazopyridinyl, isoxazolopyridinyl, thianaphthalenyl, purinyl, xanthinyl, adeninyl, guaninyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, quinoxalinyl, and quinazolinyl groups. Heteroaryl groups can be unsubstituted, or can be substituted with groups as is discussed above. Representative substituted heteroaryl groups can be substituted one or more times with groups such as those listed above.

30 [0078] Additional examples of aryl and heteroaryl groups include but are not limited to phenyl, biphenyl, indenyl, naphthyl (1-naphthyl, 2-naphthyl), N-hydroxytetrazolyl, N-hydroxytriazolyl, N-hydroxyimidazolyl, anthracenyl (1-anthracenyl, 2-anthracenyl, 3-anthracenyl), thiophenyl (2-thienyl, 3-thienyl), furyl (2-furyl, 3-furyl), indolyl, oxadiazolyl, isoxazolyl, quinazolinyl, fluorenyl, xanthenyl, isoindanyl, benzhydryl, acridinyl, thiazolyl, pyrrolyl (2-pyrrolyl),

pyrazolyl (3-pyrazolyl), imidazolyl (1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl), triazolyl (1,2,3-triazol-1-yl, 1,2,3-triazol-2-yl, 1,2,3-triazol-4-yl, 1,2,4-triazol-3-yl), oxazolyl (2-oxazolyl, 4-oxazolyl, 5-oxazolyl), thiazolyl (2-thiazolyl, 4-thiazolyl, 5-thiazolyl), pyridyl (2-pyridyl, 3-pyridyl, 4-pyridyl), pyrimidinyl (2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl), pyrazinyl, pyridazinyl (3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl), quinolyl (2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl), isoquinolyl (1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl), benzo[b]furanyl (2-benzo[b]furanyl, 3-benzo[b]furanyl, 4-benzo[b]furanyl, 5-benzo[b]furanyl, 6-benzo[b]furanyl, 7-benzo[b]furanyl), 2,3-dihydro-benzo[b]furanyl (2-(2,3-dihydro-benzo[b]furanyl), 3-(2,3-dihydro-benzo[b]furanyl), 4-(2,3-dihydro-benzo[b]furanyl), 5-(2,3-dihydro-benzo[b]furanyl), 6-(2,3-dihydro-benzo[b]furanyl), 7-(2,3-dihydro-benzo[b]furanyl), benzo[b]thiophenyl (2-benzo[b]thiophenyl, 3-benzo[b]thiophenyl, 4-benzo[b]thiophenyl, 5-benzo[b]thiophenyl, 6-benzo[b]thiophenyl, 7-benzo[b]thiophenyl), 2,3-dihydro-benzo[b]thiophenyl, (2-(2,3-dihydro-benzo[b]thiophenyl), 3-(2,3-dihydro-benzo[b]thiophenyl), 4-(2,3-dihydro-benzo[b]thiophenyl), 5-(2,3-dihydro-benzo[b]thiophenyl), 6-(2,3-dihydro-benzo[b]thiophenyl), 7-(2,3-dihydro-benzo[b]thiophenyl), indolyl (1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl), indazole (1-indazolyl, 3-indazolyl, 4-indazolyl, 5-indazolyl, 6-indazolyl, 7-indazolyl), benzimidazolyl (1-benzimidazolyl, 2-benzimidazolyl, 4-benzimidazolyl, 5-benzimidazolyl, 6-benzimidazolyl, 7-benzimidazolyl, 8-benzimidazolyl), benzoxazolyl (1-benzoxazolyl, 2-benzoxazolyl), benzothiazolyl (1-benzothiazolyl, 2-benzothiazolyl, 4-benzothiazolyl, 5-benzothiazolyl, 6-benzothiazolyl, 7-benzothiazolyl), carbazolyl (1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl), 5H-dibenz[b,f]azepine (5H-dibenz[b,f]azepin-1-yl, 5H-dibenz[b,f]azepine-2-yl, 5H-dibenz[b,f]azepine-3-yl, 5H-dibenz[b,f]azepine-4-yl, 5H-dibenz[b,f]azepine-5-yl), 10,11-dihydro-5H-dibenz[b,f]azepine (10,11-dihydro-5H-dibenz[b,f]azepine-1-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-2-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-3-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-4-yl, 10,11-dihydro-5H-dibenz[b,f]azepine-5-yl), and the like.

[0079] Heterocyclalkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of an alkyl group as defined above is replaced with a bond to a heterocyclalkyl group as defined above. Representative heterocyclalkyl groups include, but are not limited to, furan-2-yl methyl, furan-3-yl methyl, pyridine-3-yl methyl, tetrahydrofuran-2-yl ethyl, and indol-2-yl propyl.

[0080] Heteroarylalkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to a heteroaryl group as defined above.

[0081] The term "alkoxy" refers to an oxygen atom connected to an alkyl group, including a cycloalkyl group, as are defined above. Examples of linear alkoxy groups include but are not limited to methoxy, ethoxy, propoxy, butoxy, pentyloxy, hexyloxy, and the like. Examples of branched alkoxy include but are not limited to isopropoxy, sec-butoxy, tert-butoxy, isopentyloxy, isohexyloxy, and the like. Examples of cyclic alkoxy include but are not limited to cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, and the like. An alkoxy group can include one to about 12-20 carbon atoms bonded to the oxygen atom, and can further include double or triple bonds, and can also include heteroatoms. For example, an allyloxy group is an alkoxy group within the meaning herein. A methoxyethoxy group is also an alkoxy group within the meaning herein, as is a methylenedioxy group in a context where two adjacent atoms of a structures are substituted therewith.

[0082] The term "thioalkoxy" refers to an alkyl group previously defined attached to the parent molecular moiety through a sulfur atom.

[0083] The term "glycosyloxyoxy" refers to a glycoside attached to the parent molecular moiety through an oxygen atom.

[0084] The term "alkoxycarbonyl" represents as ester group; i.e. an alkoxy group, attached to the parent molecular moiety through a carbonyl group such as methoxycarbonyl, ethoxycarbonyl, and the like.

[0085] The terms "halo" or "halogen" or "halide" by themselves or as part of another substituent mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom, preferably, fluorine, chlorine, or bromine.

[0086] A "haloalkyl" group includes mono-halo alkyl groups, poly-halo alkyl groups wherein all halo atoms can be the same or different, and per-halo alkyl groups, wherein all hydrogen atoms are replaced by halogen atoms, such as fluoro. Examples of haloalkyl include trifluoromethyl, 1,1-dichloroethyl, 1,2-dichloroethyl, 1,3-dibromo-3,3-difluoropropyl, perfluorobutyl, and the like.

[0087] A "haloalkoxy" group includes mono-halo alkoxy groups, poly-halo alkoxy groups wherein all halo atoms can be the same or different, and per-halo alkoxy groups, wherein all hydrogen atoms are replaced by halogen atoms, such as fluoro. Examples of haloalkoxy include trifluoromethoxy, 1,1-dichloroethoxy, 1,2-dichloroethoxy, 1,3-dibromo-3,3-difluoropropoxy, perfluorobutoxy, and the like.

[0088] The term “(C_x-C_y)perfluoroalkyl,” wherein $x < y$, means an alkyl group with a minimum of x carbon atoms and a maximum of y carbon atoms, wherein all hydrogen atoms are replaced by fluorine atoms. Preferred is -(C₁-C₆)perfluoroalkyl, more preferred is -(C₁-C₃)perfluoroalkyl, most preferred is -CF₃.

5 **[0089]** The term “(C_x-C_y)perfluoroalkylene,” wherein $x < y$, means an alkyl group with a minimum of x carbon atoms and a maximum of y carbon atoms, wherein all hydrogen atoms are replaced by fluorine atoms. Preferred is -(C₁-C₆)perfluoroalkylene, more preferred is -(C₁-C₃)perfluoroalkylene, most preferred is -CF₂-.

10 **[0090]** The terms "aryloxy" and "arylalkoxy" refer to, respectively, an aryl group bonded to an oxygen atom and an aralkyl group bonded to the oxygen atom at the alkyl moiety. Examples include but are not limited to phenoxy, naphthyloxy, and benzyloxy.

[0091] An “acyl” group as the term is used herein refers to a group containing a carbonyl moiety wherein the group is bonded via the carbonyl carbon atom. The carbonyl carbon atom is also bonded to another carbon atom, which can be part of an alkyl, aryl, aralkyl cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl group or the like. In the special case wherein the carbonyl carbon atom is bonded to a hydrogen, the group is a “formyl” group, an acyl group as the term is defined herein. An acyl group can include 0 to about 12-20 additional carbon atoms bonded to the carbonyl group. An acyl group can include double or triple bonds within the meaning herein. An acryloyl group is an example of an acyl group. An acyl group can also include heteroatoms within the meaning here. A nicotinoyl group (pyridyl-3-carbonyl) group is an example of an acyl group within the meaning herein. Other examples include acetyl, benzoyl, phenylacetyl, pyridylacetyl, cinnamoyl, and acryloyl groups and the like. When the group containing the carbon atom that is bonded to the carbonyl carbon atom contains a halogen, the group is termed a “haloacyl” group. An example is a trifluoroacetyl group.

25

[0092] The term “amine” includes primary, secondary, and tertiary amines having, e.g., the formula N(group)₃ wherein each group can independently be H or non-H, such as alkyl, aryl, and the like. Amines include but are not limited to R-NH₂, for example, alkylamines, arylamines, alkylarylamines; R₂NH wherein each R is independently selected, such as dialkylamines, diarylamines, aralkylamines, heterocyclylamines and the like; and R₃N wherein each R is independently selected, such as trialkylamines, dialkylarylamines, alkyl diarylamines, triarylamines, and the like. The term "amine" also includes ammonium ions as used herein.

30

[0093] An "amino" group is a substituent of the form -NH₂, -NHR, -NR₂, -NR₃⁺, wherein each R is independently selected, and protonated forms of each, except for -NR₃⁺, which cannot be

protonated. Accordingly, any compound substituted with an amino group can be viewed as an amine. An "amino group" within the meaning herein can be a primary, secondary, tertiary or quaternary amino group. An "alkylamino" group includes a monoalkylamino, dialkylamino, and trialkylamino group.

- 5 [0094] An "ammonium" ion includes the unsubstituted ammonium ion NH_4^+ , but unless otherwise specified, it also includes any protonated or quaternarized forms of amines. Thus, trimethylammonium hydrochloride and tetramethylammonium chloride are both ammonium ions, and amines, within the meaning herein.

[0095] The term "amide" (or "amido") includes C- and N-amide groups, i.e., $-\text{C}(\text{O})\text{NR}_2$, and $-\text{NRC}(\text{O})\text{R}$ groups, respectively. Amide groups therefore include but are not limited to primary carboxamide groups ($-\text{C}(\text{O})\text{NH}_2$) and formamide groups ($-\text{NHC}(\text{O})\text{H}$). A "carboxamido" or "aminocarbonyl" group is a group of the formula $\text{C}(\text{O})\text{NR}_2$, wherein R can be H, alkyl, aryl, etc.

[0096] The term "azido" refers to an N_3 group. An "azide" can be an organic azide or can be a salt of the azide (N_3^-) anion. The term "nitro" refers to an NO_2 group bonded to an organic moiety. The term "nitroso" refers to an NO group bonded to an organic moiety. The term nitrate refers to an ONO_2 group bonded to an organic moiety or to a salt of the nitrate (NO_3^-) anion.

[0097] The term "urethane" ("carbamoyl" or "carbamyl") includes N- and O-urethane groups, i.e., $-\text{NRC}(\text{O})\text{OR}$ and $-\text{OC}(\text{O})\text{NR}_2$ groups, respectively.

20 [0098] The term "sulfonamide" (or "sulfonamido") includes S- and N-sulfonamide groups, i.e., $-\text{SO}_2\text{NR}_2$ and $-\text{NRSO}_2\text{R}$ groups, respectively. Sulfonamide groups therefore include but are not limited to sulfamoyl groups ($-\text{SO}_2\text{NH}_2$). An organosulfur structure represented by the formula $-\text{S}(\text{O})(\text{NR})-$ is understood to refer to a sulfoximine, wherein both the oxygen and the nitrogen atoms are bonded to the sulfur atom, which is also bonded to two carbon atoms.

25 [0099] The term "amidine" or "amidino" includes groups of the formula $-\text{C}(\text{NR})\text{NR}_2$. Typically, an amidino group is $-\text{C}(\text{NH})\text{NH}_2$.

[00100] The term "guanidine" or "guanidino" includes groups of the formula $-\text{NRC}(\text{NR})\text{NR}_2$. Typically, a guanidino group is $-\text{NHC}(\text{NH})\text{NH}_2$.

[00101] The term "ring derived from a sugar" refers to a compound that forms a ring by removing the hydrogen atoms from two hydroxyl groups of any sugar.

30 [00102] A "salt" as is well known in the art includes an organic compound such as a carboxylic acid, a sulfonic acid, or an amine, in ionic form, in combination with a counterion. For example, acids in their anionic form can form salts with cations such as metal cations, for example sodium, potassium, and the like; with ammonium salts such as NH_4^+ or the cations of various

amines, including tetraalkyl ammonium salts such as tetramethylammonium, or other cations such as trimethylsulfonium, and the like. A "pharmaceutically acceptable" or

"pharmacologically acceptable" salt is a salt formed from an ion that has been approved for human consumption and is generally non-toxic, such as a chloride salt or a sodium salt. A

5 "zwitterion" is an internal salt such as can be formed in a molecule that has at least two ionizable groups, one forming an anion and the other a cation, which serve to balance each other. For example, amino acids such as glycine can exist in a zwitterionic form. A "zwitterion" is a salt within the meaning herein. The compounds described herein may take the form of salts. The

term "salts" embraces addition salts of free acids or free bases which are compounds described

10 herein. Salts can be "pharmaceutically-acceptable salts." The term

"pharmaceutically-acceptable salt" refers to salts which possess toxicity profiles within a range that affords utility in pharmaceutical applications. Pharmaceutically unacceptable salts may nonetheless possess properties such as high crystallinity, which have utility in the practice of the present disclosure, such as for example utility in process of synthesis, purification or

15 formulation of compounds of the present disclosure.

[00103] Suitable pharmaceutically-acceptable acid addition salts may be prepared from an inorganic acid or from an organic acid. Examples of inorganic acids include hydrochloric, hydrobromic, hydriodic, nitric, carbonic, sulfuric, and phosphoric acids. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic,

20 carboxylic and sulfonic classes of organic acids, examples of which include formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, 4-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, trifluoromethanesulfonic, 2-hydroxyethanesulfonic, p-toluenesulfonic, sulfanilic,

25 cyclohexylaminosulfonic, stearic, alginic, β -hydroxybutyric, salicylic, galactaric and galacturonic acid. Examples of pharmaceutically unacceptable acid addition salts include, for example, perchlorates and tetrafluoroborates.

[00104] Suitable pharmaceutically acceptable base addition salts of compounds of the present disclosure include, for example, metallic salts including alkali metal, alkaline earth metal and

30 transition metal salts such as, for example, calcium, magnesium, potassium, sodium and zinc salts. Pharmaceutically acceptable base addition salts also include organic salts made from basic amines such as, for example, *N,N'*-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. Examples of pharmaceutically unacceptable base addition salts include lithium salts and cyanate salts.

Although pharmaceutically unacceptable salts are not generally useful as medicaments, such salts may be useful, for example as intermediates in the synthesis of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIE) compounds, for example in their purification by recrystallization. All of these salts may be prepared by conventional means from the
5 corresponding compound according to Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIE) by reacting, for example, the appropriate acid or base with the compound according to Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIE). The term "pharmaceutically acceptable salts" refers to nontoxic inorganic or organic acid and/or base addition salts, see, for example, Lit et al., Salt Selection for Basic Drugs (1986), *Int*
10 *J. Pharm.*, **33**, 201-217, incorporated by reference herein.

[00105] A "hydrate" is a compound that exists in a composition with water molecules. The composition can include water in stoichiometric quantities, such as a monohydrate or a dihydrate, or can include water in random amounts. As the term is used herein a "hydrate" refers to a solid form, i.e., a compound in water solution, while it may be hydrated, is not a hydrate as the term is
15 used herein.

[00106] A "solvate" is a similar composition except that a solvent other than water replaces the water. For example, methanol or ethanol can form an "alcoholate", which can again be stoichiometric or non-stoichiometric. As the term is used herein a "solvate" refers to a solid form, i.e., a compound in solution in a solvent, while it may be solvated, is not a solvate as the
20 term is used herein.

[00107] A "prodrug" as is well known in the art is a substance that can be administered to a patient where the substance is converted in vivo by the action of biochemicals within the patient's body, such as enzymes, to the active pharmaceutical ingredient. Examples of prodrugs include esters of carboxylic acid groups, which can be hydrolyzed by endogenous esterases as
25 are found in the bloodstream of humans and other mammals. Further examples of prodrugs include boronate esters which can be hydrolyzed under physiological conditions to afford the corresponding boronic acid. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

[00108] In addition, where features or aspects of the present disclosure are described in terms of Markush groups, those skilled in the art will recognize that the presently described compounds are also thereby described in terms of any individual member or subgroup of members of the Markush group. For example, if X is described as selected from the group consisting of bromine, chlorine, and iodine, claims for X being bromine and claims for X being
30

bromine and chlorine are fully described. Moreover, where features or aspects of the present disclosure are described in terms of Markush groups, those skilled in the art will recognize that the present disclosure is also thereby described in terms of any combination of individual members or subgroups of members of Markush groups. Thus, for example, if X is described as
5 selected from the group consisting of bromine, chlorine, and iodine, and Y is described as selected from the group consisting of methyl, ethyl, and propyl, claims for X being bromine and Y being methyl are fully described.

[00109] If a value of a variable that is necessarily an integer, e.g., the number of carbon atoms in an alkyl group or the number of substituents on a ring, is described as a range, e.g., 0-4, what
10 is meant is that the value can be any integer between 0 and 4 inclusive, i.e., 0, 1, 2, 3, or 4.

[00110] In various embodiments, the compound or set of compounds, such as are used in the inventive methods, can be any one of any of the combinations and/or sub-combinations of the above-listed embodiments.

[00111] In various embodiments, a compound as shown in any of the Examples, or among the
15 exemplary compounds, is provided. Provisos may apply to any of the disclosed categories or embodiments wherein any one or more of the other above disclosed embodiments or species may be excluded from such categories or embodiments.

[00112] The present disclosure further embraces isolated compounds according to Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe). The expression "isolated
20 compound" refers to a preparation of a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), or a mixture of compounds according to Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), wherein the isolated compound has been separated from the reagents used, and/or byproducts formed, in the synthesis of the compound or compounds. "Isolated" does not mean that the preparation is technically pure
25 (homogeneous), but it is sufficiently pure to compound in a form in which it can be used therapeutically. Preferably an "isolated compound" refers to a preparation of a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a mixture of compounds according to Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), which contains the named compound or mixture of compounds according to Formula
30 (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) in an amount of at least 10 percent by weight of the total weight. Preferably the preparation contains the named compound or mixture of compounds in an amount of at least 50 percent by weight of the total weight; more preferably at least 80 percent by weight of the total weight; and most preferably at least 90 percent, at least 95 percent or at least 98 percent by weight of the total weight of the preparation.

[00113] The compounds described herein and intermediates may be isolated from their reaction mixtures and purified by standard techniques such as filtration, liquid-liquid extraction, solid phase extraction, distillation, recrystallization or chromatography, including flash column chromatography, or HPLC.


5 Isomerism and Tautomerism in Compounds Described Herein

Tautomerism

[00114] Within the present disclosure it is to be understood that a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a salt thereof may exhibit the phenomenon of tautomerism whereby two chemical compounds that are capable of facile
 10 interconversion by exchanging a hydrogen atom between two atoms, to either of which it forms a covalent bond. Since the tautomeric compounds exist in mobile equilibrium with each other they may be regarded as different isomeric forms of the same compound. It is to be understood that the formulae drawings within this specification can represent only one of the possible tautomeric forms. However, it is also to be understood that the present disclosure encompasses
 15 any tautomeric form, and is not to be limited merely to any one tautomeric form utilized within the formulae drawings. The formulae drawings within this specification can represent only one of the possible tautomeric forms and it is to be understood that the specification encompasses all possible tautomeric forms of the compounds drawn not just those forms which it has been convenient to show graphically herein. For example, tautomerism may be exhibited by a
 20 pyrazolyl group bonded as indicated by the wavy line. While both substituents would be termed a 4-pyrazolyl group, it is evident that a different nitrogen atom bears the hydrogen atom in each structure.



[00115] Such tautomerism can also occur with substituted pyrazoles such as 3-methyl, 5-methyl, or 3,5-dimethylpyrazoles, and the like. Another example of tautomerism is amido-imido (lactam-lactim when cyclic) tautomerism, such as is seen in heterocyclic compounds bearing a ring oxygen atom adjacent to a ring nitrogen atom. For example, the equilibrium:



is an example of tautomerism. Accordingly, a structure depicted herein as one tautomer is intended to also include the other tautomer.

30 *Optical Isomerism*

[00116] It will be understood that when compounds of the present disclosure contain one or more chiral centers, the compounds may exist in, and may be isolated as pure enantiomeric or diastereomeric forms or as racemic mixtures. The present disclosure therefore includes any possible enantiomers, diastereomers, racemates or mixtures thereof of the compounds described herein.

[00117] The isomers resulting from the presence of a chiral center comprise a pair of non-superimposable isomers that are called “enantiomers.” Single enantiomers of a pure compound are optically active, *i.e.*, they are capable of rotating the plane of plane polarized light. Single enantiomers are designated according to the *Cahn-Ingold-Prelog* system. The priority of substituents is ranked based on atomic weights, a higher atomic weight, as determined by the systematic procedure, having a higher priority ranking. Once the priority ranking of the four groups is determined, the molecule is oriented so that the lowest ranking group is pointed away from the viewer. Then, if the descending rank order of the other groups proceeds clockwise, the molecule is designated (*R*) and if the descending rank of the other groups proceeds counterclockwise, the molecule is designated (*S*). In the example below, the *Cahn-Ingold-Prelog* ranking is $A > B > C > D$. The lowest ranking atom, D is oriented away from the viewer.



[00118] The present disclosure is meant to encompass diastereomers as well as their racemic and resolved, diastereomerically and enantiomerically pure forms and salts thereof. Diastereomeric pairs may be resolved by known separation techniques including normal and reverse phase chromatography, and crystallization.


[00119] “Isolated optical isomer” means a compound which has been substantially purified from the corresponding optical isomer(s) of the same formula. Preferably, the isolated isomer is at least about 80%, more preferably at least 90% pure, even more preferably at least 98% pure, most preferably at least about 99% pure, by weight.

[00120] Isolated optical isomers may be purified from racemic mixtures by well-known chiral separation techniques. According to one such method, a racemic mixture of a compound described herein, or a chiral intermediate thereof, is separated into 99% wt.% pure optical isomers by HPLC using a suitable chiral column, such as a member of the series of DAICEL[®]

CHIRALPAK[®] family of columns (Daicel Chemical Industries, Ltd., Tokyo, Japan). The column is operated according to the manufacturer's instructions.


Rotational Isomerism

[00121] It is understood that due to chemical properties (*i.e.*, resonance lending some double bond character to the C-N bond) of restricted rotation about the amide bond linkage (as illustrated below) it is possible to observe separate rotamer species and even, under some circumstances, to isolate such species (see below). It is further understood that certain structural elements, including steric bulk or substituents on the amide nitrogen, may enhance the stability of a rotamer to the extent that a compound may be isolated as, and exist indefinitely, as a single stable rotamer. The present disclosure therefore includes any possible stable rotamers of formula (I) which are biologically active in the treatment of cancer or other proliferative disease states.



Regioisomerism


[00122] In some embodiments, the compounds described herein have a particular spatial arrangement of substituents on the aromatic rings, which is related to the structure activity relationship demonstrated by the compound class. Often such substitution arrangement is denoted by a numbering system; however, numbering systems are often not consistent between different ring systems. In six-membered aromatic systems, the spatial arrangements are specified by the common nomenclature "para" for 1,4-substitution, "meta" for 1,3-substitution and "ortho" for 1,2-substitution as shown below.



[00123] In various embodiments, the compound or set of compounds, such as are among the inventive compounds or are used in the inventive methods, can be any one of any of the combinations and/or sub-combinations of the above-listed embodiments.

Compounds

[00124] In one aspect described herein are compounds of Formula (I):



Formula (I);

wherein:

R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$,
 5 $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, or $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, or optionally substituted heterocycloalkyl;

R^3 is H, or $-(C_1-C_6)alkyl$;

10 R^4 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OH$, $-(C_3-C_6)cycloalkyl$, or $-C(O)NH_2$; or R^3 and R^4 are combined to form a heterocycloalkyl ring;

R^5 is H, or $-(C_1-C_6)alkyl$; or R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring;

R^6 , R^7 , and R^8 are each independently H, or $-(C_1-C_6)alkyl$;

15 R^9 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;

R^{10} is H, or $-(C_1-C_6)alkyl$;

R^{11} and R^{12} are each independently H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)heteroalkyl-CO_2H$, $-(C_1-C_6)alkyl-S(O)-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-N(H)CH=NH$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted $-(C_1-C_6)alkyl-N(H)heterocycloalkyl$, or $-(C_1-C_6)alkyl-heteroaryl$; or R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring, and R^{12} is H;

R^{15} , R^{16} , R^{17} , and R^{18} are each independently H, $-(C_1-C_6)alkyl$, $-(C_3-C_6)cycloalkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$;

25 X is optionally substituted $-(C_1-C_6)alkyl$ -, $-(C_2-C_6)alkenyl$ -, $-(C_2-C_6)alkynyl$ -, $-(C_3-C_7)cycloalkyl$ -, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally

substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;

Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, -NH₂, -CN, -CF₃, -CO₂H, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl-, -(C₂-C₁₂)alkynyl-, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl-, -N(R²⁴)(C₁-C₁₂)alkyl-, -N(R²⁴)C(O)(C₁-C₁₂)alkyl, optionally substituted -(C₃-C₇)cycloalkyl-, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

each R²¹ and R²² is independently H, -(C₁-C₆)alkyl-, -(C₁-C₆)heteroalkyl-, -(C₁-C₆)alkyl-CO₂H, -C(O)(C₁-C₆)alkyl-, -C(O)N(R³¹)₂, or -SO₂N(R³¹)₂; or R²¹ and R²² and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R³¹ is independently H or -(C₁-C₆)alkyl; or two R³¹ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R²³ is independently H or -(C₁-C₆)alkyl;

each R²⁴ is independently H or -(C₁-C₆)alkyl;

each R²⁵ and R²⁶ is independently H or optionally substituted -(C₁-C₆)alkyl; or R²⁵ and R²⁶ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R²⁷ is independently halogen, optionally substituted -(C₁-C₆)alkyl, or optionally substituted -(C₁-C₆)heteroalkyl;

each R²⁸ is independently halogen, optionally substituted -(C₁-C₆)alkyl, or optionally substituted -(C₁-C₆)heteroalkyl;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00125] In one embodiment is a compound of Formula (I) wherein R⁶, R⁷, and R⁸ are H.

[00126] In another embodiment is a compound of Formula (I) wherein R¹⁵ and R¹⁶ are H.

[00127] In one embodiment is a compound of Formula (I) wherein R¹⁷ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I) wherein R¹⁷ is -CH₃. In another embodiment is a compound of Formula (I) wherein R¹⁷ is -CH₂CH₃. In another embodiment is a compound of Formula (I) wherein R¹⁷ is -(C₃-C₆)cycloalkyl. In another embodiment is a compound of

Formula (I) wherein R^{17} is cyclopropyl. In another embodiment is a compound of Formula (I) wherein R^{17} is $-(C_1-C_6)\text{alkyl}-C(O)OR^{23}$. In another embodiment is a compound of Formula (I) wherein R^{17} is $-\text{CH}_2\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (I) wherein R^{17} is $-(C_1-C_6)\text{alkyl}-NR^{21}R^{22}$. In another embodiment is a compound of Formula (I) wherein R^{17} is $-\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (I) wherein R^{17} is H.

[00128] In another embodiment is a compound of Formula (I) wherein R^{18} is H.

[00129] In another embodiment is a compound of Formula (I) wherein R^3 is H.

[00130] In another embodiment is a compound of Formula (I) wherein R^5 is H.

[00131] In another embodiment is a compound of Formula (I) wherein R^4 is H. In another

embodiment is a compound of Formula (I) wherein R^4 is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (I) wherein R^4 is $-\text{CH}_3$. In another embodiment is a compound of Formula (I) wherein R^4 is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (I) wherein R^4 is $-(C_1-C_6)\text{alkyl}-\text{OH}$. In another embodiment is a compound of Formula (I) wherein R^4 is $-\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (I) wherein R^4 is $-(C_3-C_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (I) wherein R^4 is cyclopropyl. In another embodiment is a compound of Formula (I) wherein R^4 is $-\text{C(O)NH}_2$.

[00132] In another embodiment is a compound of Formula (I) wherein R^3 , R^4 , and R^5 are H.

[00133] In another embodiment is a compound of Formula (I) wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring.

[00134] In another embodiment is a compound of Formula (I) wherein R^{10} is H.

[00135] In another embodiment is a compound of Formula (I) wherein R^{10} is H and R^9 is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (I) wherein R^{10} is H and R^9 is $-\text{CH}_3$. In another embodiment is a compound of Formula (I) wherein R^{10} is H and R^9 is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (I) wherein R^{10} is H and R^9 is $-(C_1-C_6)\text{haloalkyl}$. In another embodiment is a compound of Formula (I) wherein R^{10} is H and R^9 is $-\text{CH}_2\text{F}$. In another embodiment is a compound of Formula (I) wherein R^{10} is H and R^9 is $-\text{CHF}_2$. In another embodiment is a compound of Formula (I) wherein R^{10} is H and R^9 is $-(C_3-C_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (I) wherein R^{10} is H and R^9 is cyclopropyl. In another embodiment is a compound of Formula (I) wherein R^{10} is H and R^9 is H.

[00136] In another embodiment is a compound of Formula (I) wherein R^{12} is H.

[00137] In another embodiment is a compound of Formula (I) wherein R^{12} is H and R^{11} is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (I) wherein R^{12} is H and R^{11} is $-\text{CH}_3$. In another embodiment is a compound of Formula (I) wherein R^{12} is H and R^{11} is $-(C_1-$

C₆)alkyl-OR²³. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -CH₂OH. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -CH₂CH₂OH. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -
 5 -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-NH₂. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -CH₂NH₂. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -CH₂CH₂NH₂. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂CH₂NH₂. In another embodiment is a compound of
 10 Formula (I) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-CN. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -CH₂CN. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -CH₂C(O)NH₂. In
 15 another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -CH₂CH₂C(O)NH₂. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-heteroaryl. In another embodiment is a compound of Formula (I) wherein R¹² is H and R¹¹ is H.

[00138] In another embodiment is a compound of Formula (I) wherein R¹¹ and R¹⁸ are
 20 combined to form an optionally substituted heterocycloalkyl ring and R¹² is H.

[00139] In another embodiment is a compound of Formula (I) wherein p is 1 and R²⁷ is halogen. In another embodiment is a compound of Formula (I) wherein p is 1 and R²⁷ is optionally substituted -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I) wherein q is 0, p is 1 and R²⁷ is halogen. In another embodiment is a compound of Formula (I) wherein q is 0, p is 1 and R²⁷ is optionally substituted -(C₁-C₆)alkyl. In another embodiment is a
 25 compound of Formula (I) wherein q is 1 and R²⁸ is halogen. In another embodiment is a compound of Formula (I) wherein q is 1 and R²⁸ is optionally substituted -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I) wherein p is 0, q is 1 and R²⁸ is halogen. In another embodiment is a compound of Formula (I) wherein p is 0, q is 1 and R²⁸ is optionally
 30 substituted -(C₁-C₆)alkyl.

[00140] In another embodiment is a compound of Formula (I) wherein p is 0, and q is 0.


[00141] In another embodiment is a compound of Formula (I) wherein R¹ and R² are each independently H, or -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (I) wherein R¹ and R² are each H. In another embodiment is a compound of Formula (I)

wherein R^1 and R^2 are each independently $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (I) wherein R^1 is H, and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (I) wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$, and R^2 is H. In another embodiment is a compound of Formula (I) wherein R^1 is H, and R^2 is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (I) wherein R^1 is $-CH_2CH_2NH_2$, and R^2 is H. In another embodiment is a compound of Formula (I) wherein R^1 and R^2 are each $-CH_2CH_2NH_2$. In a further embodiment is a compound of Formula (I) wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$ and R^2 is H. In a further embodiment is a compound of Formula (I) wherein R^1 is $-CH_2CH_2NH_2$ and R^2 is H. In a further embodiment is a compound of Formula (I) wherein R^1 is H and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In a further embodiment is a compound of Formula (I) wherein R^1 is H and R^2 is $-CH_2CH_2NH_2$.


[00142] In another embodiment is a compound of Formula (I) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (I) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (I) wherein X is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (I) wherein X is heteroaryl which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I) wherein X is heteroaryl which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (I) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In a further embodiment is a compound of Formula (I) wherein X is pyridine which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I) wherein X is pyridine which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I) wherein X is pyridine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (I) wherein X is pyrimidine which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I) wherein X is pyrimidine which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I) wherein X is pyrimidine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (I) wherein X is pyridine which is substituted once with methyl. In a further embodiment is a compound of Formula (I) wherein X is pyrimidine which is substituted once with methyl. In a further embodiment is a compound of Formula (I) wherein X is pyrimidine which is substituted twice with methyl. In another embodiment is a compound of Formula (I) wherein X is optionally substituted $-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (I)

wherein Y is optionally substituted aryl. In another embodiment is a compound of Formula (I) wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (I) wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (I) wherein Y is optionally substituted $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (I) wherein Y is $-O-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (I) wherein Y is $-N(H)-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (I) wherein Y is a bond. In another embodiment is a compound of Formula (I) wherein Z is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (I) wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (I) wherein Z is optionally substituted phenyl. In a further embodiment is a compound of Formula (I) wherein Z is phenyl substituted once or twice with $-(C_1-C_8)alkyl$. In a further embodiment is a compound of Formula (I) wherein Z is phenyl substituted once with n-butyl, isobutyl, or tert-butyl. In a further embodiment is a compound of Formula (I) wherein Z is phenyl substituted once with n-butyl. In a further embodiment is a compound of Formula (I) wherein Z is phenyl substituted once with isobutyl. In a further embodiment is a compound of Formula (I) wherein Z is phenyl substituted once with tert-butyl. In another embodiment is a compound of Formula (I) wherein Z is optionally substituted heteroaryl. In another embodiment is a compound of Formula (I) wherein Z is optionally substituted $-(C_3-C_7)cycloalkyl$. In another embodiment is a compound of Formula (I) wherein Z is halogen.

[00143] In another embodiment is a compound of Formula (I) wherein -X-Y-Z is



[00144] In one aspect described herein are compounds of Formula (I'):



Formula (I');

wherein:

- 5 R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl-heterocycloalkyl$, $(C_1-C_6)heteroalkyl$ or optionally substituted heterocycloalkyl;
- 10 R^3 is H, or $-(C_1-C_6)alkyl$;
 R^4 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OH$, $-(C_3-C_6)cycloalkyl$, or $-C(O)NH_2$; or R^3 and R^4 are combined to form a heterocycloalkyl ring;
 R^5 is H, or $-(C_1-C_6)alkyl$; or R^4 and R^5 and the carbon atom to which that are attached form a cyclopropyl ring;
- 15 R^6 , R^7 , and R^8 are each independently H, fluoro, hydroxyl, amino, optionally substituted alkyl or heteroalkyl or $-(C_1-C_6)alkyl$;
 R^9 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;
 R^{10} is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;
 or R^9 and R^{10} are combined to form a heterocycloalkyl or cycloalkyl ring
- 20 R^{11} and R^{12} are each independently H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)heteroalkyl-CO_2H$, $-(C_1-C_6)alkyl-S(O)-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-N(H)CH=NH$, $-(C_1-C_6)alkyl-C(NH_2)=NH$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-N(H)SO_2NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(H)-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted $-(C_1-C_6)alkyl-N(H)heterocycloalkyl$, or $-(C_1-C_6)alkyl-heteroaryl$; or R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring, and R^{12} is H;
- 25

R^{15} , R^{16} , R^{17} , and R^{18} are each independently H, $-(C_1-C_6)alkyl$, $-(C_3-C_6)cycloalkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$;

X is optionally substituted $-(C_1-C_6)alkyl$ -, $-(C_2-C_6)alkenyl$ -, $-(C_2-C_6)alkynyl$ -, $-(C_3-C_7)cycloalkyl$ -, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-O-(C_1-C_6)alkyl$ -, $-N(R^{24})(C_1-C_6)alkyl$ -, $-N(R^{24})(C_6-C_{10})aryl$ -, or $-SO_2(C_1-C_6)alkyl$ -;

Y is a bond, optionally substituted $-(C_1-C_6)alkyl$ -, $-(C_2-C_6)alkenyl$ -, $-(C_2-C_6)alkynyl$ -, $-(C_1-C_6)alkyl-N(R^{24})(C_1-C_6)alkyl$ -, $-O-(C_1-C_6)alkyl$ -, $-O(C_6-C_{10})aryl$ -, $-N(R^{24})(C_1-C_6)alkyl$ -, $-N(R^{24})SO_2(C_1-C_6)alkyl$ -, $-N(R^{24})C(O)(C_1-C_6)alkyl$ -, $-C(O)(C_1-C_6)alkyl$ -, $-S(C_1-C_6)alkyl$ -, $-SO_2(C_1-C_6)alkyl$ -, $-C(O)NH(C_1-C_6)alkyl$ -, $-(C_3-C_7)cycloalkyl$ -, optionally substituted $-C(O)N(R^{24})aryl$ -, optionally substituted $-N(R^{24})C(O)aryl$ -, optionally substituted $-N(R^{24})SO_2aryl$ -, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, $-NH_2$, $-CN$, $-CF_3$, $-CO_2H$, $-(C_1-C_{12})alkyl$ -, $-(C_2-C_{12})alkenyl$ -, $-(C_2-C_{12})alkynyl$ -, $-C(O)NR^{25}R^{26}$, $-O-(C_1-C_{12})alkyl$ -, $-N(R^{24})(C_1-C_{12})alkyl$ -, $-N(R^{24})C(O)(C_1-C_{12})alkyl$ -, optionally substituted $-(C_3-C_7)cycloalkyl$ -, $-(C_1-C_6)alkyl-heterocycloalkyl$ -, optionally substituted aryl, or optionally substituted heteroaryl;

each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl$ -, $-(C_1-C_6)heteroalkyl$ -, $-(C_1-C_6)alkyl-CO_2H$ -, $-C(O)(C_1-C_6)alkyl$ -, $-C(O)O(C_1-C_6)alkyl$ -, $-C(O)O(C_1-C_6)haloalkyl$ -, $-C(=NH)(C_1-C_6)alkyl$ -, $-C(=NH)N(R^{31})_2$ -, $-C(O)N(R^{31})_2$ -, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)alkyl$; or two R^{31} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{23} is independently H or $-(C_1-C_6)alkyl$;

each R^{24} is independently H or $-(C_1-C_6)alkyl$;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl$; or R^{25} and R^{26} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{27} is independently halogen, $-NR^{23}R^{24}$ -, $-NC(O)R^{23}$ -, $-NC(O)NR^{23}R^{24}$ -, nitro, hydroxyl, optionally substituted $-(C_1-C_6)alkyl$ -, optionally substituted $-(C_1-C_6)heteroalkyl$ -, $-(C_1-C_6)alkoxy$ -, $-C(O)(C_1-C_6)alkyl$ -, or $-S(O)_2(C_1-C_6)alkyl$;

each R^{28} is independently halogen, $-NR^{23}R^{24}$ -, $-NC(O)R^{23}$ -, $-NC(O)NR^{23}R^{24}$ -, nitro, hydroxyl, optionally substituted $-(C_1-C_6)alkyl$ -, optionally substituted $-(C_1-C_6)heteroalkyl$ -, $-(C_1-C_6)alkoxy$ -, $-C(O)(C_1-C_6)alkyl$ -, or $-S(O)_2(C_1-C_6)alkyl$;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00145] In one embodiment is a compound of Formula (I') wherein R^6 , R^7 , and R^8 are H.

[00146] In another embodiment is a compound of Formula (I') wherein R^{15} and R^{16} are H.

[00147] In one embodiment is a compound of Formula (I') wherein R^{17} is $-(C_1-C_6)$ alkyl. In

5 another embodiment is a compound of Formula (I') wherein R^{17} is $-CH_3$. In another embodiment is a compound of Formula (I') wherein R^{17} is $-CH_2CH_3$. In another embodiment is a compound of Formula (I') wherein R^{17} is $-(C_3-C_6)$ cycloalkyl. In another embodiment is a compound of Formula (I') wherein R^{17} is cyclopropyl. In another embodiment is a compound of Formula (I') wherein R^{17} is $-(C_1-C_6)$ alkyl- $C(O)OR^{23}$. In another embodiment is a compound
10 of Formula (I') wherein R^{17} is $-CH_2CH_2OH$. In another embodiment is a compound of Formula (I') wherein R^{17} is $-(C_1-C_6)$ alkyl- $NR^{21}R^{22}$. In another embodiment is a compound of Formula (I') wherein R^{17} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (I') wherein R^{17} is H.

[00148] In another embodiment is a compound of Formula (I') wherein R^{18} is H.

15 **[00149]** In another embodiment is a compound of Formula (I') wherein R^3 is H.

[00150] In another embodiment is a compound of Formula (I') wherein R^5 is H.

[00151] In another embodiment is a compound of Formula (I') wherein R^4 is H. In another embodiment is a compound of Formula (I') wherein R^4 is $-(C_1-C_6)$ alkyl. In another embodiment is a compound of Formula (I') wherein R^4 is $-CH_3$. In another embodiment is a compound of
20 Formula (I') wherein R^4 is $-CH_2CH_3$. In another embodiment is a compound of Formula (I') wherein R^4 is $-(C_1-C_6)$ alkyl-OH. In another embodiment is a compound of Formula (I') wherein R^4 is $-CH_2OH$. In another embodiment is a compound of Formula (I') wherein R^4 is $-(C_3-C_6)$ cycloalkyl. In another embodiment is a compound of Formula (I') wherein R^4 is cyclopropyl. In another embodiment is a compound of Formula (I') wherein R^4 is $-C(O)NH_2$.

25 **[00152]** In another embodiment is a compound of Formula (I') wherein R^3 , R^4 , and R^5 are H.

[00153] In another embodiment is a compound of Formula (I') wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring.

[00154] In another embodiment is a compound of Formula (I') wherein R^{10} is H.

[00155] In another embodiment is a compound of Formula (I') wherein R^{10} is H and R^9 is $-(C_1-C_6)$ alkyl. In another embodiment is a compound of Formula (I') wherein R^{10} is H and R^9 is $-CH_3$. In another embodiment is a compound of Formula (I') wherein R^{10} is H and R^9 is $-CH_2CH_3$. In another embodiment is a compound of Formula (I') wherein R^{10} is H and R^9 is $-(C_1-C_6)$ haloalkyl. In another embodiment is a compound of Formula (I') wherein R^{10} is H and R^9 is $-CH_2F$. In another embodiment is a compound of Formula (I') wherein R^{10} is H and R^9 is -
30

CHF₂. In another embodiment is a compound of Formula (I') wherein R¹⁰ is H and R⁹ is -(C₃-C₆)cycloalkyl. In another embodiment is a compound of Formula (I') wherein R¹⁰ is H and R⁹ is cyclopropyl. In another embodiment is a compound of Formula (I') wherein R¹⁰ is H and R⁹ is H.

- 5 [00156] In another embodiment is a compound of Formula (I') wherein R¹² is H.
- [00157] In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₃. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-OR²³. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₂OH. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₂CH₂OH. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-NH₂. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₂NH₂. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₂CH₂NH₂. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-CN. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₂CN. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₂C(O)NH₂. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -CH₂CH₂C(O)NH₂. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-heteroaryl. In another embodiment is a compound of Formula (I') wherein R¹² is H and R¹¹ is H.

[00158] In another embodiment is a compound of Formula (I') wherein R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring and R¹² is H.

- [00159] In another embodiment is a compound of Formula (I') wherein p is 1 and R²⁷ is halogen. In another embodiment is a compound of Formula (I') wherein p is 1 and R²⁷ is optionally substituted -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I') wherein q is 0, p is 1 and R²⁷ is halogen. In another embodiment is a compound of Formula (I') wherein q is 0, p is 1 and R²⁷ is optionally substituted -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I') wherein q is 1 and R²⁸ is halogen. In another embodiment is a

compound of Formula (I') wherein q is 1 and R²⁸ is optionally substituted -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I') wherein p is 0, q is 1 and R²⁸ is halogen. In another embodiment is a compound of Formula (I') wherein p is 0, q is 1 and R²⁸ is optionally substituted -(C₁-C₆)alkyl.

5 [00160] In another embodiment is a compound of Formula (I') wherein p is 0, and q is 0.

[00161] In another embodiment is a compound of Formula (I') wherein R¹ and R² are each independently H, or -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (I') wherein R¹ and R² are each H. In another embodiment is a compound of Formula (I') wherein R¹ and R² are each independently -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (I') wherein R¹ is H, and R² is -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (I') wherein R¹ is -(C₁-C₆)alkyl-NR²¹R²², and R² is H. In another embodiment is a compound of Formula (I') wherein R¹ is H, and R² is -CH₂CH₂NH₂. In another embodiment is a compound of Formula (I') wherein R¹ is -CH₂CH₂NH₂, and R² is H. In another embodiment is a compound of Formula (I') wherein R¹ and R² are each -


15 CH₂CH₂NH₂. In a further embodiment is a compound of Formula (I') wherein R¹ is -(C₁-C₆)alkyl-NR²¹R²² and R² is H. In a further embodiment is a compound of Formula (I') wherein R¹ is -CH₂CH₂NH₂ and R² is H. In a further embodiment is a compound of Formula (I') wherein R¹ is H and R² is -(C₁-C₆)alkyl-NR²¹R²². In a further embodiment is a compound of Formula (I') wherein R¹ is H and R² is -CH₂CH₂NH₂.


20 [00162] In another embodiment is a compound of Formula (I') wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (I') wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (I') wherein X is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (I') wherein X is heteroaryl which is unsubstituted or substituted once or twice with -(C₁-C₆)alkyl.

25 In a further embodiment is a compound of Formula (I') wherein X is heteroaryl which is unsubstituted or substituted once with -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I') wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In a further embodiment is a compound of Formula (I') wherein X is pyridine which is unsubstituted or substituted once or twice with -(C₁-C₆)alkyl. In a further embodiment is a compound of Formula (I') wherein X is pyridine which is unsubstituted or substituted once with -(C₁-C₆)alkyl. In a further embodiment is a compound of Formula (I') wherein X is pyridine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (I') wherein X is pyrimidine which is unsubstituted or substituted once or twice with -(C₁-C₆)alkyl. In a further embodiment is a compound of Formula (I') wherein X is


pyrimidine which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (I') wherein X is pyrimidine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (I') wherein X is pyridine which is substituted once with methyl. In another embodiment is a compound of Formula (I') wherein X is optionally substituted $-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (I') wherein Y is optionally substituted aryl. In another embodiment is a compound of Formula (I') wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (I') wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (I') wherein Y is optionally substituted $-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (I') wherein Y is -O- $(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (I') wherein Y is -N(H)- $(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (I') wherein Y is a bond. In another embodiment is a compound of Formula (I') wherein Z is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (I') wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (I') wherein Z is optionally substituted phenyl. In a further embodiment is a compound of Formula (I') wherein Z is phenyl substituted once or twice with $-(C_1-C_8)alkyl$. In a further embodiment is a compound of Formula (I') wherein Z is phenyl substituted once with n-butyl, isobutyl, or tert-butyl. In a further embodiment is a compound of Formula (I') wherein Z is phenyl substituted once with n-butyl. In a further embodiment is a compound of Formula (I') wherein Z is phenyl substituted once with isobutyl. In a further embodiment is a compound of Formula (I') wherein Z is phenyl substituted once with tert-butyl. In another embodiment is a compound of Formula (I') wherein Z is optionally substituted heteroaryl. In another embodiment is a compound of Formula (I') wherein Z is optionally substituted $-(C_3-C_7)cycloalkyl$. In another embodiment is a compound of Formula (I') wherein Z is halogen.

[00163] In another embodiment is a compound of Formula (I') wherein -X-Y-Z is





[00164] In another embodiment is a compound of Formula (I) having the structure of Formula (Ia):



Formula (Ia):

wherein:

R¹ and R² are each independently H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -CH₂CH(OH)CH₂NH₂, -CH₂CH(heterocycloalkyl)CH₂NH₂, -CH₂C(O)NH₂, -CH₂C(O)N(H)CH₂CN, -(C₁-C₆)alkyl-
10 C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-N(R²³)C(O)(C₁-C₆)alkylNR²¹R²², or -(C₁-C₆)alkyl-C(O)N(R²³)(C₁-C₆)alkyl, or optionally substituted heterocycloalkyl;

R³ is H, or -(C₁-C₆)alkyl;

15 R⁴ is H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OH, -(C₃-C₆)cycloalkyl, or -C(O)NH₂; or R³ and R⁴ are combined to form a heterocycloalkyl ring;

R⁵ is H, or -(C₁-C₆)alkyl; or R⁴ and R⁵ and the carbon atom to which they are attached form a cyclopropyl ring;

R⁶, R⁷, and R⁸ are each independently H, or -(C₁-C₆)alkyl;

R⁹ is H, -(C₁-C₆)alkyl, -(C₁-C₆)haloalkyl, or -(C₃-C₆)cycloalkyl;

20 R¹⁰ is H, or -(C₁-C₆)alkyl;

R¹¹ and R¹² are each independently H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-

C(O)NR²⁵R²⁶, -(C₁-C₆)heteroalkyl-CO₂H, -(C₁-C₆)alkyl-S(O)-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-N(H)CH=NH, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted -(C₁-C₆)alkyl-N(H)heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring, and R¹² is H;

5 R¹⁵, R¹⁶, R¹⁷, and R¹⁸ are each independently H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;

X is optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₃-C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;

Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, -NH₂, -CN, -CF₃, -CO₂H, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl, -(C₂-C₁₂)alkynyl, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl, -N(R²⁴)(C₁-C₁₂)alkyl, -N(R²⁴)C(O)(C₁-C₁₂)alkyl, optionally substituted -(C₃-C₇)cycloalkyl, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

each R²¹ and R²² is independently H, -(C₁-C₆)alkyl, -(C₁-C₆)heteroalkyl, -(C₁-C₆)alkyl-CO₂H, -C(O)(C₁-C₆)alkyl, -C(O)N(R³¹)₂, or -SO₂N(R³¹)₂; or R²¹ and R²² and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R³¹ is independently H or -(C₁-C₆)alkyl; or two R³¹ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R²³ is independently H or -(C₁-C₆)alkyl;

each R²⁴ is independently H or -(C₁-C₆)alkyl;

each R²⁵ and R²⁶ is independently H or optionally substituted -(C₁-C₆)alkyl; or R²⁵ and R²⁶ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R²⁷ is independently halogen, optionally substituted -(C₁-C₆)alkyl, or optionally substituted -(C₁-C₆)heteroalkyl;

each R²⁸ is independently halogen, optionally substituted -(C₁-C₆)alkyl, or optionally substituted -(C₁-C₆)heteroalkyl;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00165] In one embodiment is a compound of Formula (Ia) wherein R^6 , R^7 , and R^8 are H.

[00166] In another embodiment is a compound of Formula (Ia) wherein R^{15} and R^{16} are H.

5 **[00167]** In one embodiment is a compound of Formula (Ia) wherein R^{17} is $-(C_1-C_6)$ alkyl. In another embodiment is a compound of Formula (Ia) wherein R^{17} is $-CH_3$. In another embodiment is a compound of Formula (Ia) wherein R^{17} is $-CH_2CH_3$. In another embodiment is a compound of Formula (Ia) wherein R^{17} is $-(C_3-C_6)$ cycloalkyl. In another embodiment is a compound of Formula (Ia) wherein R^{17} is cyclopropyl. In another embodiment is a compound of Formula (Ia) wherein R^{17} is $-(C_1-C_6)$ alkyl- $C(O)OR^{23}$. In another embodiment is a compound of Formula (Ia) wherein R^{17} is $-CH_2CH_2OH$. In another embodiment is a compound of Formula (Ia) wherein R^{17} is $-(C_1-C_6)$ alkyl- $NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ia) wherein R^{17} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ia) wherein R^{17} is H.

15 **[00168]** In another embodiment is a compound of Formula (Ia) wherein R^{18} is H.

[00169] In another embodiment is a compound of Formula (Ia) wherein R^3 is H.

[00170] In another embodiment is a compound of Formula (Ia) wherein R^5 is H.

[00171] In another embodiment is a compound of Formula (Ia) wherein R^4 is H. In another embodiment is a compound of Formula (Ia) wherein R^4 is $-(C_1-C_6)$ alkyl. In another embodiment is a compound of Formula (Ia) wherein R^4 is $-CH_3$. In another embodiment is a compound of Formula (Ia) wherein R^4 is $-CH_2CH_3$. In another embodiment is a compound of Formula (Ia) wherein R^4 is $-(C_1-C_6)$ alkyl-OH. In another embodiment is a compound of Formula (Ia) wherein R^4 is $-CH_2OH$. In another embodiment is a compound of Formula (Ia) wherein R^4 is $-(C_3-C_6)$ cycloalkyl. In another embodiment is a compound of Formula (Ia) wherein R^4 is cyclopropyl. In another embodiment is a compound of Formula (Ia) wherein R^4 is $-C(O)NH_2$.

[00172] In another embodiment is a compound of Formula (Ia) wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring.

[00173] In another embodiment is a compound of Formula (Ia) wherein R^3 , R^4 , and R^5 are H.

30 **[00174]** In another embodiment is a compound of Formula (Ia) wherein R^{10} is H.

[00175] In another embodiment is a compound of Formula (Ia) wherein R^{10} is H and R^9 is $-(C_1-C_6)$ alkyl. In another embodiment is a compound of Formula (Ia) wherein R^{10} is H and R^9 is $-CH_3$. In another embodiment is a compound of Formula (Ia) wherein R^{10} is H and R^9 is $-CH_2CH_3$. In another embodiment is a compound of Formula (Ia) wherein R^{10} is H and R^9 is -

(C₁-C₆)haloalkyl. In another embodiment is a compound of Formula (Ia) wherein R¹⁰ is H and R⁹ is -CH₂F. In another embodiment is a compound of Formula (Ia) wherein R¹⁰ is H and R⁹ is -CHF₂. In another embodiment is a compound of Formula (Ia) wherein R¹⁰ is H and R⁹ is -(C₃-C₆)cycloalkyl. In another embodiment is a compound of Formula (Ia) wherein R¹⁰ is H and R⁹ is cyclopropyl. In another embodiment is a compound of Formula (Ia) wherein R¹⁰ is H and R⁹ is H.

[00176] In another embodiment is a compound of Formula (Ia) wherein R¹² is H.

[00177] In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₃. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-OR²³. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₂OH. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₂CH₂OH. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-NH₂. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₂NH₂. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₂CH₂NH₂. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-CN. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₂CN. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₂C(O)NH₂. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -CH₂CH₂C(O)NH₂. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-heteroaryl. In another embodiment is a compound of Formula (Ia) wherein R¹² is H and R¹¹ is H.

[00178] In another embodiment is a compound of Formula (Ia) wherein R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring and R¹² is H.

[00179] In another embodiment is a compound of Formula (Ia) wherein p is 1 and R²⁷ is halogen. In another embodiment is a compound of Formula (Ia) wherein p is 1 and R²⁷ is optionally substituted -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (I) wherein q is 0, p is 1 and R²⁷ is halogen. In another embodiment is a compound of Formula (I)

wherein q is 0, p is 1 and R^{27} is optionally substituted $-(C_1-C_6)$ alkyl. In another embodiment is a compound of Formula (Ia) wherein q is 1 and R^{28} is halogen. In another embodiment is a compound of Formula (Ia) wherein q is 1 and R^{28} is optionally substituted $-(C_1-C_6)$ alkyl. In another embodiment is a compound of Formula (Ia) wherein p is 0, q is 1 and R^{28} is halogen. In another embodiment is a compound of Formula (Ia) wherein p is 0, q is 1 and R^{28} is optionally substituted $-(C_1-C_6)$ alkyl.

[00180] In another embodiment is a compound of Formula (Ia) wherein p is 0, and q is 0.


[00181] In another embodiment is a compound of Formula (Ia) wherein R^1 and R^2 are each independently H, or $-(C_1-C_6)$ alkyl- $NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ia) wherein R^1 and R^2 are each H. In another embodiment is a compound of Formula (Ia) wherein R^1 and R^2 are each independently $-(C_1-C_6)$ alkyl- $NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ia) wherein R^1 is H, and R^2 is $-(C_1-C_6)$ alkyl- $NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ia) wherein R^1 is $-(C_1-C_6)$ alkyl- $NR^{21}R^{22}$, and R^2 is H. In another embodiment is a compound of Formula (Ia) wherein R^1 is H, and R^2 is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ia) wherein R^1 is $-CH_2CH_2NH_2$, and R^2 is H.


In another embodiment is a compound of Formula (Ia) wherein R^1 and R^2 are each $-CH_2CH_2NH_2$. In a further embodiment is a compound of Formula (Ia) wherein R^1 is $-(C_1-C_6)$ alkyl- $NR^{21}R^{22}$ and R^2 is H. In a further embodiment is a compound of Formula (Ia) wherein R^1 is $-CH_2CH_2NH_2$ and R^2 is H. In a further embodiment is a compound of Formula (Ia) wherein R^1 is H and R^2 is $-(C_1-C_6)$ alkyl- $NR^{21}R^{22}$. In a further embodiment is a compound of Formula (Ia) wherein R^1 is H and R^2 is $-CH_2CH_2NH_2$.

[00182] In another embodiment is a compound of Formula (Ia) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (Ia) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (Ia) wherein X is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (Ia) wherein X is heteroaryl which is unsubstituted or substituted once or twice with $-(C_1-C_6)$ alkyl. In a further embodiment is a compound of Formula (Ia) wherein X is heteroaryl which is unsubstituted or substituted once with $-(C_1-C_6)$ alkyl. In another embodiment is a compound of Formula (Ia) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In a further embodiment is a compound of Formula (Ia) wherein X is pyridine which is unsubstituted or substituted once or twice with $-(C_1-C_6)$ alkyl. In a further embodiment is a compound of Formula (Ia) wherein X is pyridine which is unsubstituted or substituted once with $-(C_1-C_6)$ alkyl. In a further embodiment is a compound of Formula (Ia) wherein X is pyridine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a


compound of Formula (Ia) wherein X is pyrimidine which is unsubstituted or substituted once or twice with $-(C_1-C_6)\text{alkyl}$. In a further embodiment is a compound of Formula (Ia) wherein X is pyrimidine which is unsubstituted or substituted once with $-(C_1-C_6)\text{alkyl}$. In a further embodiment is a compound of Formula (Ia) wherein X is pyrimidine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (Ia) wherein X is pyridine which is substituted once with methyl. In a further embodiment is a compound of Formula (Ia) wherein X is pyrimidine which is substituted once with methyl. In a further embodiment is a compound of Formula (Ia) wherein X is pyrimidine which is substituted twice with methyl. In another embodiment is a compound of Formula (Ia) wherein X is optionally substituted $-(C_1-C_6)\text{alkyl}$ -. In another embodiment is a compound of Formula (Ia) wherein Y is optionally substituted aryl. In another embodiment is a compound of Formula (Ia) wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (Ia) wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (Ia) wherein Y is optionally substituted $-(C_1-C_6)\text{alkyl}$ -. In another embodiment is a compound of Formula (Ia) wherein Y is $-O-(C_1-C_6)\text{alkyl}$ -. In another embodiment is a compound of Formula (Ia) wherein Y is $-N(H)-(C_1-C_6)\text{alkyl}$ -. In another embodiment is a compound of Formula (Ia) wherein Y is a bond. In another embodiment is a compound of Formula (Ia) wherein Z is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (Ia) wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (Ia) wherein Z is optionally substituted phenyl. In a further embodiment is a compound of Formula (Ia) wherein Z is phenyl substituted once or twice with $-(C_1-C_8)\text{alkyl}$. In a further embodiment is a compound of Formula (Ia) wherein Z is phenyl substituted once with n-butyl, isobutyl, or tert-butyl. In a further embodiment is a compound of Formula (Ia) wherein Z is phenyl substituted once with n-butyl. In a further embodiment is a compound of Formula (Ia) wherein Z is phenyl substituted once with isobutyl. In a further embodiment is a compound of Formula (Ia) wherein Z is phenyl substituted once with tert-butyl. In another embodiment is a compound of Formula (Ia) wherein Z is optionally substituted heteroaryl. In another embodiment is a compound of Formula (Ia) wherein Z is optionally substituted $-(C_3-C_7)\text{cycloalkyl}$. In another embodiment is a compound of Formula (Ia) wherein Z is halogen.

[00183] In another embodiment is a compound of Formula (Ia) wherein -X-Y-Z is





[00184] In another embodiment is a compound of Formula (I) having the structure of Formula (Ib):



Formula (Ib);

wherein:

- R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, or $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, or optionally substituted heterocycloalkyl;
- R^4 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OH$, $-(C_3-C_6)cycloalkyl$, or $-C(O)NH_2$;
- R^5 is H, or $-(C_1-C_6)alkyl$; or R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring;
- R^9 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;
- R^{11} is H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)heteroalkyl-$

CO₂H, -(C₁-C₆)alkyl-S(O)-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-N(H)CH=NH, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted -(C₁-C₆)alkyl-N(H)heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring;

5 R¹⁷ and R¹⁸ are each independently H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;

X is optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₃-C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;

Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, -NH₂, -CN, -CF₃, -CO₂H, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl-, -(C₂-C₁₂)alkynyl-, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl-, -N(R²⁴)(C₁-C₁₂)alkyl-, -N(R²⁴)C(O)(C₁-C₁₂)alkyl-, optionally substituted -(C₃-C₇)cycloalkyl-, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

each R²¹ and R²² is independently H, -(C₁-C₆)alkyl, -(C₁-C₆)heteroalkyl, -(C₁-C₆)alkyl-CO₂H, -C(O)(C₁-C₆)alkyl, -C(O)N(R³¹)₂, or -SO₂N(R³¹)₂; or R²¹ and R²² and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R³¹ is independently H or -(C₁-C₆)alkyl; or two R³¹ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R²³ is independently H or -(C₁-C₆)alkyl;

each R²⁴ is independently H or -(C₁-C₆)alkyl; and

each R²⁵ and R²⁶ is independently H or optionally substituted -(C₁-C₆)alkyl; or R²⁵ and R²⁶ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

30 or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00185] In one embodiment is a compound of Formula (Ib) wherein R¹⁷ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (Ib) wherein R¹⁷ is -CH₃. In another embodiment is a compound of Formula (Ib) wherein R¹⁷ is -CH₂CH₃. In another embodiment is a compound of Formula (Ib) wherein R¹⁷ is -(C₃-C₆)cycloalkyl. In another embodiment is a

compound of Formula (Ib) wherein R^{17} is cyclopropyl. In another embodiment is a compound of Formula (Ib) wherein R^{17} is $-(C_1-C_6)\text{alkyl}-C(O)OR^{23}$. In another embodiment is a compound of Formula (Ib) wherein R^{17} is $-\text{CH}_2\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (Ib) wherein R^{17} is $-(C_1-C_6)\text{alkyl}-NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ib) wherein R^{17} is $-\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (Ib) wherein R^{17} is H.

[00186] In another embodiment is a compound of Formula (Ib) wherein R^{18} is H.

[00187] In another embodiment is a compound of Formula (Ib) wherein R^5 is H.

[00188] In another embodiment is a compound of Formula (Ib) wherein R^4 is H. In another embodiment is a compound of Formula (Ib) wherein R^4 is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (Ib) wherein R^4 is $-\text{CH}_3$. In another embodiment is a compound of Formula (Ib) wherein R^4 is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (Ib) wherein R^4 is $-(C_1-C_6)\text{alkyl}-\text{OH}$. In another embodiment is a compound of Formula (Ib) wherein R^4 is $-\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (Ib) wherein R^4 is $-(C_3-C_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (Ib) wherein R^4 is cyclopropyl. In another embodiment is a compound of Formula (Ib) wherein R^4 is $-\text{C}(O)\text{NH}_2$.

[00189] In another embodiment is a compound of Formula (Ib) wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring.

[00190] In another embodiment is a compound of Formula (Ib) wherein R^9 is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (Ib) wherein R^9 is $-\text{CH}_3$. In another embodiment is a compound of Formula (Ib) wherein R^9 is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (Ib) wherein R^9 is $-(C_1-C_6)\text{haloalkyl}$. In another embodiment is a compound of Formula (Ib) wherein R^9 is $-\text{CH}_2\text{F}$. In another embodiment is a compound of Formula (Ib) wherein R^9 is $-\text{CHF}_2$. In another embodiment is a compound of Formula (Ib) wherein R^9 is $-(C_3-C_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (Ib) wherein R^9 is cyclopropyl. In another embodiment is a compound of Formula (Ib) wherein R^9 is H.

[00191] In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-(C_1-C_6)\text{alkyl}$.

In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-\text{CH}_3$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-(C_1-C_6)\text{alkyl}-OR^{23}$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound

of Formula (Ib) wherein R^{11} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-(C_1-C_6)alkyl-NH_2$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-CH_2NH_2$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-CH_2CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-(C_1-C_6)alkyl-CN$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-CH_2CN$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-CH_2C(O)NH_2$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-CH_2CH_2C(O)NH_2$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is $-(C_1-C_6)alkyl-heteroaryl$. In another embodiment is a compound of Formula (Ib) wherein R^{11} is H. **[00192]** In another embodiment is a compound of Formula (Ib) wherein R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring.


[00193] In another embodiment is a compound of Formula (Ib) wherein R^1 and R^2 are each independently H, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ib) wherein R^1 and R^2 are each H. In another embodiment is a compound of Formula (Ib) wherein R^1 and R^2 are each independently $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ib) wherein R^1 is H, and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ib) wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$, and R^2 is H. In another embodiment is a compound of Formula (Ib) wherein R^1 is H, and R^2 is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ib) wherein R^1 is $-CH_2CH_2NH_2$, and R^2 is H. In another embodiment is a compound of Formula (Ib) wherein R^1 and R^2 are each $-CH_2CH_2NH_2$. In a further embodiment is a compound of Formula (Ib) wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$ and R^2 is H. In a further embodiment is a compound of Formula (Ib) wherein R^1 is $-CH_2CH_2NH_2$ and R^2 is H. In a further embodiment is a compound of Formula (Ib) wherein R^1 is H and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In a further embodiment is a compound of Formula (Ib) wherein R^1 is H and R^2 is $-CH_2CH_2NH_2$.

[00194] In another embodiment is a compound of Formula (Ib) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (Ib) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (Ib) wherein X is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (Ib) wherein X is heteroaryl which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (Ib) wherein X is heteroaryl which is


unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (Ib) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In a further embodiment is a compound of Formula (Ib) wherein X is pyridine which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (Ib) wherein X is pyridine which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (Ib) wherein X is pyridine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (Ib) wherein X is pyrimidine which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (Ib) wherein X is pyrimidine which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (Ib) wherein X is pyrimidine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (Ib) wherein X is pyridine which is substituted once with methyl. In a further embodiment is a compound of Formula (Ib) wherein X is pyrimidine which is substituted once with methyl. In a further embodiment is a compound of Formula (Ib) wherein X is pyrimidine which is substituted twice with methyl. In another embodiment is a compound of Formula (Ib) wherein X is optionally substituted $-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (Ib) wherein Y is optionally substituted aryl. In another embodiment is a compound of Formula (Ib) wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (Ib) wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (Ib) wherein Y is optionally substituted $-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (Ib) wherein Y is $-O-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (Ib) wherein Y is $-N(H)-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (Ib) wherein Y is a bond. In another embodiment is a compound of Formula (Ib) wherein Z is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (Ib) wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (Ib) wherein Z is optionally substituted phenyl. In a further embodiment is a compound of Formula (Ib) wherein Z is phenyl substituted once or twice with $-(C_1-C_8)alkyl$. In a further embodiment is a compound of Formula (Ib) wherein Z is phenyl substituted once with n-butyl, isobutyl, or tert-butyl. In a further embodiment is a compound of Formula (Ib) wherein Z is phenyl substituted once with n-butyl. In a further embodiment is a compound of Formula (Ib) wherein Z is phenyl substituted once with isobutyl. In a further embodiment is a compound of Formula (Ib) wherein Z is phenyl substituted once with tert-butyl. In another embodiment is a compound of Formula (Ib) wherein Z is optionally substituted heteroaryl. In another embodiment is a

compound of Formula (Ib) wherein Z is optionally substituted -(C₃-C₇)cycloalkyl. In another embodiment is a compound of Formula (Ib) wherein Z is halogen.

[00195] In another embodiment is a compound of Formula (Ib) wherein -X-Y-Z is



[00196] In another embodiment is a compound of Formula (I) having the structure of Formula (Ic):



Formula (Ic);

wherein:

R¹ and R² are each independently H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -CH₂CH(OH)CH₂NH₂, -CH₂CH(heterocycloalkyl)CH₂NH₂, -CH₂C(O)NH₂, -CH₂C(O)N(H)CH₂CN, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-N(R²³)C(O)(C₁-C₆)alkylNR²¹R²², or -(C₁-C₆)alkyl-C(O)N(R²³)(C₁-C₆)alkyl, or optionally substituted heterocycloalkyl;

- R^{11} is H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)heteroalkyl-CO_2H$, $-(C_1-C_6)alkyl-S(O)-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-N(H)CH=NH$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted $-(C_1-C_6)alkyl-N(H)heterocycloalkyl$, or $-(C_1-C_6)alkyl-heteroaryl$;
- X is optionally substituted $-(C_1-C_6)alkyl-$, $-(C_2-C_6)alkenyl-$, $-(C_2-C_6)alkynyl$, $-(C_3-C_7)cycloalkyl-$, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-O-(C_1-C_6)alkyl-$, $-N(R^{24})(C_1-C_6)alkyl-$, $-N(R^{24})(C_6-C_{10})aryl-$, or $-SO_2(C_1-C_6)alkyl-$;
- Y is a bond, optionally substituted $-(C_1-C_6)alkyl-$, $-(C_2-C_6)alkenyl-$, $-(C_2-C_6)alkynyl$, $-(C_1-C_6)alkyl-N(R^{24})(C_1-C_6)alkyl-$, $-O-(C_1-C_6)alkyl-$, $-O(C_6-C_{10})aryl-$, $-N(R^{24})(C_1-C_6)alkyl-$, $-N(R^{24})SO_2(C_1-C_6)alkyl-$, $-N(R^{24})C(O)(C_1-C_6)alkyl-$, $-C(O)(C_1-C_6)alkyl-$, $-S(C_1-C_6)alkyl-$, $-SO_2(C_1-C_6)alkyl-$, $-C(O)NH(C_1-C_6)alkyl-$, $-(C_3-C_7)cycloalkyl-$, optionally substituted $-C(O)N(R^{24})aryl-$, optionally substituted $-N(R^{24})C(O)aryl-$, optionally substituted $-N(R^{24})SO_2aryl-$, optionally substituted aryl, or optionally substituted heteroaryl;
- Z is H, halogen, $-NH_2$, $-CN$, $-CF_3$, $-CO_2H$, $-(C_1-C_{12})alkyl$, $-(C_2-C_{12})alkenyl$, $-(C_2-C_{12})alkynyl$, $-C(O)NR^{25}R^{26}$, $-O-(C_1-C_{12})alkyl$, $-N(R^{24})(C_1-C_{12})alkyl$, $-N(R^{24})C(O)(C_1-C_{12})alkyl$, optionally substituted $-(C_3-C_7)cycloalkyl$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted aryl, or optionally substituted heteroaryl;
- each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)heteroalkyl$, $-(C_1-C_6)alkyl-CO_2H$, $-C(O)(C_1-C_6)alkyl$, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;
- each R^{31} is independently H or $-(C_1-C_6)alkyl$; or two R^{31} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;
- each R^{23} is independently H or $-(C_1-C_6)alkyl$;
- each R^{24} is independently H or $-(C_1-C_6)alkyl$; and
- each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl$; or R^{25} and R^{26} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;
- or a pharmaceutically acceptable salt, solvate, or prodrug thereof.
- [00197]** In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_3$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-(C_1-C_6)alkyl-OR^{23}$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_2OH$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_2CH_2OH$. In another embodiment is a

compound of Formula (Ic) wherein R^{11} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-(C_1-C_6)alkyl-NH_2$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_2NH_2$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_2CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-(C_1-C_6)alkyl-CN$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_2CN$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_2C(O)NH_2$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-CH_2CH_2C(O)NH_2$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is $-(C_1-C_6)alkyl-heteroaryl$. In another embodiment is a compound of Formula (Ic) wherein R^{11} is H.

[00198] In another embodiment is a compound of Formula (Ic) wherein R^1 and R^2 are each independently H, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ic) wherein R^1 and R^2 are each H. In another embodiment is a compound of Formula (Ic) wherein R^1 and R^2 are each independently $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ic) wherein R^1 is H, and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (Ic) wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$, and R^2 is H.


In another embodiment is a compound of Formula (Ic) wherein R^1 is H, and R^2 is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (Ic) wherein R^1 is $-CH_2CH_2NH_2$, and R^2 is H. In another embodiment is a compound of Formula (Ic) wherein R^1 and R^2 are each $-CH_2CH_2NH_2$. In a further embodiment is a compound of Formula (Ic) wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$ and R^2 is H. In a further embodiment is a compound of Formula (Ic) wherein R^1 is $-CH_2CH_2NH_2$ and R^2 is H. In a further embodiment is a compound of Formula (Ic) wherein R^1 is H and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In a further embodiment is a compound of Formula (Ic) wherein R^1 is H and R^2 is $-CH_2CH_2NH_2$.

[00199] In another embodiment is a compound of Formula (Ic) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (Ic) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (Ic) wherein X is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (Ic) wherein X is heteroaryl which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (Ic) wherein X is heteroaryl which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In another embodiment is a compound of


Formula (Ic) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In a further embodiment is a compound of Formula (Ic) wherein X is pyridine which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (Ic) wherein X is pyridine which is unsubstituted or substituted once with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (Ic) wherein X is pyridine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (Ic) wherein X is pyrimidine which is unsubstituted or substituted once or twice with $-(C_1-C_6)alkyl$. In a further embodiment is a compound of Formula (Ic) wherein X is pyrimidine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (Ic) wherein X is pyridine which is substituted once with methyl. In a further embodiment is a compound of Formula (Ic) wherein X is pyrimidine which is substituted once with methyl. In a further embodiment is a compound of Formula (Ic) wherein X is pyrimidine which is substituted twice with methyl. In another embodiment is a compound of Formula (Ic) wherein X is optionally substituted $-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (Ic) wherein Y is optionally substituted aryl. In another embodiment is a compound of Formula (Ic) wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (Ic) wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (Ic) wherein Y is optionally substituted $-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (Ic) wherein Y is $-O-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (Ic) wherein Y is $-N(H)-(C_1-C_6)alkyl$ -. In another embodiment is a compound of Formula (Ic) wherein Y is a bond. In another embodiment is a compound of Formula (Ic) wherein Z is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (Ic) wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (Ic) wherein Z is optionally substituted phenyl. In a further embodiment is a compound of Formula (Ic) wherein Z is phenyl substituted once or twice with $-(C_1-C_8)alkyl$. In a further embodiment is a compound of Formula (Ic) wherein Z is phenyl substituted once with n-butyl, isobutyl, or tert-butyl. In a further embodiment is a compound of Formula (Ic) wherein Z is phenyl substituted once with n-butyl. In a further embodiment is a compound of Formula (Ic) wherein Z is phenyl substituted once with isobutyl. In a further embodiment is a compound of Formula (Ic) wherein Z is phenyl substituted once with tert-butyl. In another embodiment is a compound of Formula (Ic) wherein Z is optionally substituted heteroaryl. In another embodiment is a

compound of Formula (Ic) wherein Z is optionally substituted $-(C_3-C_7)$ cycloalkyl. In another embodiment is a compound of Formula (Ic) wherein Z is halogen.

[00200] In another embodiment is a compound of Formula (Ic) wherein -X-Y-Z is



[00201] In another embodiment is a compound of Formula (I) having the structure of Formula (Id):



Formula (Id);

wherein:

R^{11} is $-CH_2NH_2$, $-CH_2CH_2NH_2$, or $-CH_2CH_2CH_2NH_2$;

X is optionally substituted $-(C_1-C_6)$ alkyl-, $-(C_2-C_6)$ alkenyl-, $-(C_2-C_6)$ alkynyl-, $-(C_3-C_7)$ cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally

15

substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;

Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, -NH₂, -CN, -CF₃, -CO₂H, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl-, -(C₂-C₁₂)alkynyl-, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl-, -N(R²⁴)(C₁-C₁₂)alkyl-, -N(R²⁴)C(O)(C₁-C₁₂)alkyl, optionally substituted -(C₃-C₇)cycloalkyl-, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

each R²⁴ is independently H or -(C₁-C₆)alkyl; and

R²⁵ and R²⁶ is independently H or optionally substituted -(C₁-C₆)alkyl; or R²⁵ and R²⁶ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00202] In one embodiment is a compound of Formula (Id) wherein R¹¹ is -CH₂NH₂. In another embodiment is a compound of Formula (Id) wherein R¹¹ is -CH₂CH₂NH₂. In another embodiment is a compound of Formula (Id) wherein R¹¹ is -CH₂CH₂CH₂NH₂.

[00203] In another embodiment is a compound of Formula (Id) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (Id) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (Id) wherein X is optionally substituted heteroaryl. In a further embodiment is a compound of Formula (Id) wherein X is heteroaryl which is unsubstituted or substituted once or twice with -(C₁-C₆)alkyl.


In a further embodiment is a compound of Formula (Id) wherein X is heteroaryl which is unsubstituted or substituted once with -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (Id) wherein X is optionally substituted pyridine or optionally substituted pyrimidine.


In a further embodiment is a compound of Formula (Id) wherein X is pyridine which is unsubstituted or substituted once or twice with -(C₁-C₆)alkyl. In a further embodiment is a

compound of Formula (Id) wherein X is pyridine which is unsubstituted or substituted once with -(C₁-C₆)alkyl. In a further embodiment is a compound of Formula (Id) wherein X is pyridine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (Id) wherein X is pyrimidine which is unsubstituted or substituted once or twice with -(C₁-C₆)alkyl. In a further embodiment is a compound of Formula (Id) wherein X


is pyrimidine which is unsubstituted or substituted once with $-(C_1-C_6)\text{alkyl}$. In a further embodiment is a compound of Formula (Id) wherein X is pyrimidine which is unsubstituted or substituted once or twice with methyl. In a further embodiment is a compound of Formula (Id) wherein X is pyridine which is substituted once with methyl. In a further embodiment is a compound of Formula (Id) wherein X is pyrimidine which is substituted once with methyl. In a further embodiment is a compound of Formula (Id) wherein X is pyrimidine which is substituted twice with methyl. In another embodiment is a compound of Formula (Id) wherein X is optionally substituted $-(C_1-C_6)\text{alkyl}$ -. In another embodiment is a compound of Formula (Id) wherein Y is optionally substituted aryl. In another embodiment is a compound of Formula (Id) wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (Id) wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (Id) wherein Y is optionally substituted $-(C_1-C_6)\text{alkyl}$ -. In another embodiment is a compound of Formula (Id) wherein Y is $-O-(C_1-C_6)\text{alkyl}$ -. In another embodiment is a compound of Formula (Id) wherein Y is $-N(H)-(C_1-C_6)\text{alkyl}$ -. In another embodiment is a compound of Formula (Id) wherein Y is a bond. In another embodiment is a compound of Formula (Id) wherein Z is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (Id) wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (Id) wherein Z is optionally substituted phenyl. In a further embodiment is a compound of Formula (Id) wherein Z is phenyl substituted once or twice with $-(C_1-C_8)\text{alkyl}$. In a further embodiment is a compound of Formula (Id) wherein Z is phenyl substituted once with n-butyl, isobutyl, or tert-butyl. In a further embodiment is a compound of Formula (Id) wherein Z is phenyl substituted once with n-butyl. In a further embodiment is a compound of Formula (Id) wherein Z is phenyl substituted once with isobutyl. In a further embodiment is a compound of Formula (Id) wherein Z is phenyl substituted once with tert-butyl. In another embodiment is a compound of Formula (Id) wherein Z is optionally substituted heteroaryl. In another embodiment is a compound of Formula (Id) wherein Z is optionally substituted $-(C_3-C_7)\text{cycloalkyl}$. In another embodiment is a compound of Formula (Id) wherein Z is halogen.

[00204] In another embodiment is a compound of Formula (Id) wherein $-X-Y-Z$ is





[00205] In another aspect described herein is a compound of Formula (II):



Formula (II);

wherein:

R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, or $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, or optionally substituted heterocycloalkyl;

R^3 is H, or $-(C_1-C_6)alkyl$;

R^4 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OH$, $-(C_3-C_6)cycloalkyl$, or $-C(O)NH_2$; or R^3 and R^4 are combined to form a heterocycloalkyl ring;

R^5 is H, or $-(C_1-C_6)alkyl$; or R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring;

R^6 , R^7 , and R^8 are each independently H, or $-(C_1-C_6)alkyl$;

R^9 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;

R¹⁰ is H, or -(C₁-C₆)alkyl;

R¹¹ and R¹² are each independently H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)heteroalkyl-CO₂H, -(C₁-C₆)alkyl-S(O)-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-

5 N(H)CH=NH, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted -(C₁-C₆)alkyl-N(H)heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring, and R¹² is H;

R¹³ and R¹⁴ are each independently H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-

10 C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹³ and R¹⁹ are combined to form an optionally substituted heterocycloalkyl ring, and R¹⁴ is H;

R¹⁵, R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are each independently H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;

15 X is optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₃-C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;

Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

25 Z is H, halogen, -NH₂, -CN, -CF₃, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl-, -(C₂-C₁₂)alkynyl-, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl-, -N(R²⁴)(C₁-C₁₂)alkyl-, -N(R²⁴)C(O)(C₁-C₁₂)alkyl-, optionally substituted -(C₃-C₇)cycloalkyl-, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

each R²¹ and R²² is independently H, -(C₁-C₆)alkyl, -(C₁-C₆)heteroalkyl, -(C₁-C₆)alkyl-CO₂H, -C(O)(C₁-C₆)alkyl, -C(O)N(R³¹)₂, or -SO₂N(R³¹)₂; or R²¹ and R²² and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R³¹ is independently H or -(C₁-C₆)alkyl; or two R³¹ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R²³ is independently H or -(C₁-C₆)alkyl;

each R^{24} is independently H or $-(C_1-C_6)alkyl$;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl$; or R^{25} and R^{26} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R^{27} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$, or optionally substituted $-(C_1-C_6)heteroalkyl$;

each R^{28} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$, or optionally substituted $-(C_1-C_6)heteroalkyl$;

n is 0 or 1;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00206] In one embodiment is a compound of Formula (II) wherein n is 0. In another embodiment is a compound of Formula (II) wherein n is 1.

[00207] In another embodiment is a compound of Formula (II) wherein R^6 , R^7 , and R^8 are H.

[00208] In another embodiment is a compound of Formula (II) wherein R^{15} and R^{16} are H.

[00209] In one embodiment is a compound of Formula (II) wherein R^{17} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (II) wherein R^{17} is $-CH_3$. In another embodiment is a compound of Formula (II) wherein R^{17} is $-CH_2CH_3$. In another embodiment is a compound of Formula (II) wherein R^{17} is $-(C_3-C_6)cycloalkyl$. In another embodiment is a compound of Formula (II) wherein R^{17} is cyclopropyl. In another embodiment is a compound of Formula (II) wherein R^{17} is $-(C_1-C_6)alkyl-C(O)OR^{23}$. In another embodiment is a compound of Formula (II) wherein R^{17} is $-CH_2CH_2OH$. In another embodiment is a compound of Formula (II) wherein R^{17} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (II) wherein R^{17} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (II) wherein R^{17} is H.

[00210] In another embodiment is a compound of Formula (II) wherein R^{18} is H.

[00211] In another embodiment is a compound of Formula (II) wherein R^{19} is H.

[00212] In another embodiment is a compound of Formula (II) wherein R^3 is H.

[00213] In another embodiment is a compound of Formula (II) wherein R^5 is H.

[00214] In another embodiment is a compound of Formula (II) wherein R^4 is H. In another embodiment is a compound of Formula (II) wherein R^4 is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (II) wherein R^4 is $-CH_3$. In another embodiment is a compound of Formula (II) wherein R^4 is $-CH_2CH_3$. In another embodiment is a compound of Formula (II) wherein R^4 is $-(C_1-C_6)alkyl-OH$. In another embodiment is a compound of Formula (II) wherein

R^4 is $-\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (II) wherein R^4 is $-(\text{C}_3-\text{C}_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (II) wherein R^4 is cyclopropyl. In another embodiment is a compound of Formula (II) wherein R^4 is $-\text{C}(\text{O})\text{NH}_2$.

[00215] In another embodiment is a compound of Formula (II) wherein R^3 , R^4 , and R^5 are H.

5 **[00216]** In another embodiment is a compound of Formula (II) wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring.

[00217] In another embodiment is a compound of Formula (II) wherein R^{10} is H.

[00218] In another embodiment is a compound of Formula (II) wherein R^{10} is H and R^9 is $-(\text{C}_1-\text{C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (II) wherein R^{10} is H and R^9 is -

10 CH_3 . In another embodiment is a compound of Formula (II) wherein R^{10} is H and R^9 is -

CH_2CH_3 . In another embodiment is a compound of Formula (II) wherein R^{10} is H and R^9 is -

$(\text{C}_1-\text{C}_6)\text{haloalkyl}$. In another embodiment is a compound of Formula (II) wherein R^{10} is H and

R^9 is $-\text{CH}_2\text{F}$. In another embodiment is a compound of Formula (II) wherein R^{10} is H and R^9 is -

CHF_2 . In another embodiment is a compound of Formula (II) wherein R^{10} is H and R^9 is $-(\text{C}_3-$

15 $\text{C}_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (II) wherein R^{10} is H and R^9 is cyclopropyl. In another embodiment is a compound of Formula (II) wherein R^{10} is H and R^9 is H.

[00219] In another embodiment is a compound of Formula (II) wherein R^{12} is H.

[00220] In another embodiment is a compound of Formula (II) wherein R^{12} is H and R^{11} is -

20 $(\text{C}_1-\text{C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (II) wherein R^{12} is H and R^{11} is

$-\text{CH}_3$. In another embodiment is a compound of Formula (II) wherein R^{12} is H and R^{11} is $-(\text{C}_1-$

$\text{C}_6)\text{alkyl}-\text{OR}^{23}$. In another embodiment is a compound of Formula (II) wherein R^{11} is $-\text{CH}_2\text{OH}$.

In another embodiment is a compound of Formula (II) wherein R^{12} is H and R^{11} is $-\text{CH}_2\text{CH}_2\text{OH}$.

In another embodiment is a compound of Formula (II) wherein R^{12} is H and R^{11} is $-(\text{C}_1-\text{C}_6)\text{alkyl}$.

25 In another embodiment is a compound of Formula (II) wherein R^{12} is H and R^{11} is $-(\text{C}_1-$

$\text{C}_6)\text{alkyl}-\text{NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (II) wherein R^{12} is H and

R^{11} is $-(\text{C}_1-\text{C}_6)\text{alkyl}-\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{12} is

H and R^{11} is $-\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{12} is H

and R^{11} is $-\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{12} is

30 H and R^{11} is $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein

R^{12} is H and R^{11} is $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula

(II) wherein R^{12} is H and R^{11} is $-(\text{C}_1-\text{C}_6)\text{alkyl}-\text{CN}$. In another embodiment is a compound of

Formula (II) wherein R^{12} is H and R^{11} is $-\text{CH}_2\text{CN}$. In another embodiment is a compound of

Formula (II) wherein R^{12} is H and R^{11} is $-(\text{C}_1-\text{C}_6)\text{alkyl}-\text{C}(\text{O})\text{NR}^{25}\text{R}^{26}$. In another embodiment is

a compound of Formula (II) wherein R^{12} is H and R^{11} is $-\text{CH}_2\text{C}(\text{O})\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{12} is H and R^{11} is $-\text{CH}_2\text{CH}_2\text{C}(\text{O})\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{12} is H and R^{11} is $-(\text{C}_1\text{-C}_6)\text{alkyl-heteroaryl}$. In another embodiment is a compound of Formula (II) wherein R^{12} is H and R^{11} is H.

5 [00221] In another embodiment is a compound of Formula (II) wherein R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring and R^{12} is H.

[00222] In another embodiment is a compound of Formula (II) wherein R^{14} is H.

[00223] In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-(\text{C}_1\text{-C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_3$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-(\text{C}_1\text{-C}_6)\text{alkyl-OR}^{23}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-(\text{C}_1\text{-C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-(\text{C}_1\text{-C}_6)\text{alkyl-NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-(\text{C}_1\text{-C}_6)\text{alkyl-CN}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{CN}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-(\text{C}_1\text{-C}_6)\text{alkyl-C}(\text{O})\text{NR}^{25}\text{R}^{26}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{C}(\text{O})\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{CH}_2\text{C}(\text{O})\text{NH}_2$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is $-(\text{C}_1\text{-C}_6)\text{alkyl-heteroaryl}$. In another embodiment is a compound of Formula (II) wherein R^{14} is H and R^{13} is H.

30 [00224] In another embodiment is a compound of Formula (II) wherein R^{13} and R^{19} are combined to form an optionally substituted heterocycloalkyl ring and R^{14} is H.

[00225] In another embodiment is a compound of Formula (II) wherein p is 1 and R^{27} is halogen. In another embodiment is a compound of Formula (II) wherein p is 1 and R^{27} is optionally substituted $-(\text{C}_1\text{-C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (II) wherein q is 0, p is 1 and R^{27} is halogen. In another embodiment is a compound of Formula (II)

wherein q is 0, p is 1 and R²⁷ is optionally substituted -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (II) wherein q is 1 and R²⁸ is halogen. In another embodiment is a compound of Formula (II) wherein q is 1 and R²⁸ is optionally substituted -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (II) wherein p is 0, q is 1 and R²⁸ is halogen. In another embodiment is a compound of Formula (II) wherein p is 0, q is 1 and R²⁸ is optionally substituted -(C₁-C₆)alkyl.

[00226] In another embodiment is a compound of Formula (II) wherein p is 0, and q is 0.

[00227] In another embodiment is a compound of Formula (II) wherein R¹ and R² are each independently H, or -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (II) wherein R¹ and R² are each H. In another embodiment is a compound of Formula (II) wherein R¹ and R² are each independently -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (II) wherein R¹ is H, and R² is -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (II) wherein R¹ is -(C₁-C₆)alkyl-NR²¹R²², and R² is H.

In another embodiment is a compound of Formula (II) wherein R¹ is H, and R² is -CH₂CH₂NH₂.

In another embodiment is a compound of Formula (II) wherein R¹ is -CH₂CH₂NH₂, and R² is H.

In another embodiment is a compound of Formula (II) wherein R¹ and R² are each -CH₂CH₂NH₂.

[00228] In another embodiment is a compound of Formula (II) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (II) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (II) wherein X is optionally substituted heteroaryl. In another embodiment is a compound of Formula (II) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In another embodiment is a compound of Formula (II) wherein X is optionally substituted -(C₁-C₆)alkyl-.

In another embodiment is a compound of Formula (II) wherein Y is optionally substituted aryl.


In another embodiment is a compound of Formula (II) wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (II) wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (II) wherein Y is optionally substituted -(C₁-C₆)alkyl-.

In another embodiment is a compound of Formula (II) wherein Y is optionally substituted -(C₁-C₆)alkyl-. In another embodiment is a compound of Formula (II) wherein Y is -O-(C₁-C₆)alkyl-.


In another embodiment is a compound of Formula (II) wherein Y is -N(H)-(C₁-C₆)alkyl-.

In another embodiment is a compound of Formula (II) wherein Y is a bond. In another embodiment is a compound of Formula (II) wherein Z is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (II) wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (II) wherein Z is optionally substituted phenyl. In another embodiment is a compound of Formula (II) wherein Z is optionally substituted

[00229] In another embodiment is a compound of Formula (II) wherein -X-Y-Z is



[00230] In another aspect described herein is a compound of Formula (II'):



Formula (II');

wherein:

R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-$

C₆)alkyl-heterocycloalkyl, optionally substituted (C₁-C₆)heteroalkyl or optionally substituted heterocycloalkyl;

R³ is H, or -(C₁-C₆)alkyl;

R⁴ is H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OH, -(C₃-C₆)cycloalkyl, or -C(O)NH₂; or R³ and R⁴ are

5 combined to form a heterocycloalkyl ring;

R⁵ is H, or -(C₁-C₆)alkyl; or R⁴ and R⁵ and the carbon atom to which that are attached form a cyclopropyl ring;

R⁶, R⁷, and R⁸ are each independently H, or -(C₁-C₆)alkyl;

R⁹ is H, -(C₁-C₆)alkyl, -(C₁-C₆)haloalkyl, or -(C₃-C₆)cycloalkyl;

10 R¹⁰ is H, or -(C₁-C₆)alkyl;

R¹¹ and R¹² are each independently H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)heteroalkyl-CO₂H, -(C₁-C₆)alkyl-S(O)-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-N(H)CH=NH, -(C₁-C₆)alkyl-C(NH₂)=NH, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-

15 N(H)SO₂NR²⁵R²⁶, -(C₁-C₆)alkyl-N(H)-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted -(C₁-C₆)alkyl-N(H)heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring, and R¹² is H;

R¹³ and R¹⁴ are each independently H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-

20 C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹³ and R¹⁹ are combined to form an optionally substituted heterocycloalkyl ring, and R¹⁴ is H;

R¹⁵, R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are each independently H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;

25 X is optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₃-C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;

Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, -NH₂, -CN, -CF₃, -(C₁-C₁₂)alkyl, -(C₂-C₁₂)alkenyl, -(C₂-C₁₂)alkynyl, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl, -N(R²⁴)(C₁-C₁₂)alkyl, -N(R²⁴)C(O)(C₁-C₁₂)alkyl, optionally substituted -(C₃-C₇)cycloalkyl, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

5 each R²¹ and R²² is independently H, -(C₁-C₆)alkyl, -(C₁-C₆)heteroalkyl, -(C₁-C₆)alkyl-CO₂H, -C(O)(C₁-C₆)alkyl, -C(O)N(R³¹)₂, or -SO₂N(R³¹)₂; or R²¹ and R²² and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R³¹ is independently H or -(C₁-C₆)alkyl; or two R³¹ and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

10 each R²³ is independently H or -(C₁-C₆)alkyl;

each R²⁴ is independently H or -(C₁-C₆)alkyl;

each R²⁵ and R²⁶ is independently H or optionally substituted -(C₁-C₆)alkyl; or R²⁵ and R²⁶ and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R²⁷ is independently halogen, -NR²³R²⁴, -NC(O)R²³, -NC(O)NR²³R²⁴), nitro, hydroxyl,

15 optionally substituted -(C₁-C₆)alkyl, optionally substituted -(C₁-C₆)heteroalkyl, -(C₁-C₆)alkoxy, -C(O)(C₁-C₆)alkyl, or -S(O)₂(C₁-C₆)alkyl;

each R²⁸ is independently halogen, -NR²³R²⁴, -NC(O)R²³, -NC(O)NR²³R²⁴), nitro, hydroxyl, optionally substituted -(C₁-C₆)alkyl, optionally substituted -(C₁-C₆)heteroalkyl, -(C₁-C₆)alkoxy, -C(O)(C₁-C₆)alkyl, or -S(O)₂(C₁-C₆)alkyl;

20 n is 0 or 1;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00231] In one embodiment is a compound of Formula (II') wherein n is 0. In another
25 embodiment is a compound of Formula (II') wherein n is 1.

[00232] In another embodiment is a compound of Formula (II') wherein R⁶, R⁷, and R⁸ are H.

[00233] In another embodiment is a compound of Formula (II') wherein R¹⁵ and R¹⁶ are H.

[00234] In one embodiment is a compound of Formula (II') wherein R¹⁷ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (II') wherein R¹⁷ is -CH₃. In another
30 embodiment is a compound of Formula (II') wherein R¹⁷ is -CH₂CH₃. In another embodiment is a compound of Formula (II') wherein R¹⁷ is -(C₃-C₆)cycloalkyl. In another embodiment is a compound of Formula (II') wherein R¹⁷ is cyclopropyl. In another embodiment is a compound of Formula (II') wherein R¹⁷ is -(C₁-C₆)alkyl-C(O)OR²³. In another embodiment is a compound of Formula (II') wherein R¹⁷ is -CH₂CH₂OH. In another embodiment is a compound of Formula

(II') wherein R^{17} is $-(C_1-C_6)\text{alkyl}-NR^{21}R^{22}$. In another embodiment is a compound of Formula (II') wherein R^{17} is $-\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (II') wherein R^{17} is H.

[00235] In another embodiment is a compound of Formula (II') wherein R^{18} is H.

5 [00236] In another embodiment is a compound of Formula (II') wherein R^{19} is H.

[00237] In another embodiment is a compound of Formula (II') wherein R^3 is H.

[00238] In another embodiment is a compound of Formula (II') wherein R^5 is H.

[00239] In another embodiment is a compound of Formula (II') wherein R^4 is H. In another embodiment is a compound of Formula (II') wherein R^4 is $-(C_1-C_6)\text{alkyl}$. In another

10 embodiment is a compound of Formula (II') wherein R^4 is $-\text{CH}_3$. In another embodiment is a compound of Formula (II') wherein R^4 is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (II') wherein R^4 is $-(C_1-C_6)\text{alkyl}-\text{OH}$. In another embodiment is a compound of Formula (II') wherein R^4 is $-\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (II') wherein R^4 is $-(C_3-C_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (II') wherein R^4 is cyclopropyl. In another embodiment is a compound of Formula (II') wherein R^4 is $-\text{C}(\text{O})\text{NH}_2$.

[00240] In another embodiment is a compound of Formula (II') wherein R^3 , R^4 , and R^5 are H.

[00241] In another embodiment is a compound of Formula (II') wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring.

20 [00242] In another embodiment is a compound of Formula (II') wherein R^{10} is H.

[00243] In another embodiment is a compound of Formula (II') wherein R^{10} is H and R^9 is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (II') wherein R^{10} is H and R^9 is $-\text{CH}_3$. In another embodiment is a compound of Formula (II') wherein R^{10} is H and R^9 is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (II') wherein R^{10} is H and R^9 is $-(C_1-C_6)\text{haloalkyl}$. In another embodiment is a compound of Formula (II') wherein R^{10} is H and R^9 is $-\text{CH}_2\text{F}$. In another embodiment is a compound of Formula (II') wherein R^{10} is H and R^9 is $-\text{CHF}_2$. In another embodiment is a compound of Formula (II') wherein R^{10} is H and R^9 is $-(C_3-C_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (II') wherein R^{10} is H and R^9 is cyclopropyl. In another embodiment is a compound of Formula (II') wherein R^{10} is H and R^9 is H.

30

[00244] In another embodiment is a compound of Formula (II') wherein R^{12} is H.

[00245] In another embodiment is a compound of Formula (II') wherein R^{12} is H and R^{11} is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (II') wherein R^{12} is H and R^{11} is $-\text{CH}_3$. In another embodiment is a compound of Formula (II') wherein R^{12} is H and R^{11} is

(C₁-C₆)alkyl-OR²³. In another embodiment is a compound of Formula (II') wherein R¹¹ is -CH₂OH. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -CH₂CH₂OH. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-NH₂. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -CH₂NH₂. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -CH₂CH₂NH₂. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-CN. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -CH₂CN. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -CH₂C(O)NH₂. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -CH₂CH₂C(O)NH₂. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-heteroaryl. In another embodiment is a compound of Formula (II') wherein R¹² is H and R¹¹ is H.

[00246] In another embodiment is a compound of Formula (II') wherein R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring and R¹² is H.

[00247] In another embodiment is a compound of Formula (II') wherein R¹⁴ is H.

[00248] In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -CH₃. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -(C₁-C₆)alkyl-OR²³. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -CH₂OH. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -CH₂CH₂OH. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -(C₁-C₆)alkyl-NH₂. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -CH₂NH₂. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -CH₂CH₂NH₂. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (II') wherein R¹⁴ is H and R¹³ is -CH₂CH₂CH₂CH₂NH₂. In another

embodiment is a compound of Formula (II') wherein R^{14} is H and R^{13} is $-(C_1-C_6)\text{alkyl-CN}$. In another embodiment is a compound of Formula (II') wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{CN}$. In another embodiment is a compound of Formula (II') wherein R^{14} is H and R^{13} is $-(C_1-C_6)\text{alkyl-C(O)NR}^{25}\text{R}^{26}$. In another embodiment is a compound of Formula (II') wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{C(O)NH}_2$. In another embodiment is a compound of Formula (II') wherein R^{14} is H and R^{13} is $-\text{CH}_2\text{CH}_2\text{C(O)NH}_2$. In another embodiment is a compound of Formula (II') wherein R^{14} is H and R^{13} is $-(C_1-C_6)\text{alkyl-heteroaryl}$. In another embodiment is a compound of Formula (II') wherein R^{14} is H and R^{13} is H.

[00249] In another embodiment is a compound of Formula (II') wherein R^{13} and R^{19} are combined to form an optionally substituted heterocycloalkyl ring and R^{14} is H.

[00250] In another embodiment is a compound of Formula (II') wherein p is 1 and R^{27} is halogen. In another embodiment is a compound of Formula (II') wherein p is 1 and R^{27} is optionally substituted $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (II') wherein q is 0, p is 1 and R^{27} is halogen. In another embodiment is a compound of Formula (II') wherein q is 0, p is 1 and R^{27} is optionally substituted $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (II') wherein q is 1 and R^{28} is halogen. In another embodiment is a compound of Formula (II') wherein q is 1 and R^{28} is optionally substituted $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (II') wherein p is 0, q is 1 and R^{28} is halogen. In another embodiment is a compound of Formula (II') wherein p is 0, q is 1 and R^{28} is optionally substituted $-(C_1-C_6)\text{alkyl}$.

[00251] In another embodiment is a compound of Formula (II') wherein p is 0, and q is 0.


[00252] In another embodiment is a compound of Formula (II') wherein R^1 and R^2 are each independently H, or $-(C_1-C_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (II') wherein R^1 and R^2 are each H. In another embodiment is a compound of Formula (II') wherein R^1 and R^2 are each independently $-(C_1-C_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (II') wherein R^1 is H, and R^2 is $-(C_1-C_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (II') wherein R^1 is $-(C_1-C_6)\text{alkyl-NR}^{21}\text{R}^{22}$, and R^2 is H. In another embodiment is a compound of Formula (II') wherein R^1 is H, and R^2 is $-\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (II') wherein R^1 is $-\text{CH}_2\text{CH}_2\text{NH}_2$, and R^2 is H. In another embodiment is a compound of Formula (II') wherein R^1 and R^2 are each $-\text{CH}_2\text{CH}_2\text{NH}_2$.

[00253] In another embodiment is a compound of Formula (II') wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (II') wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (II') wherein X is


optionally substituted heteroaryl. In another embodiment is a compound of Formula (II') wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In another embodiment is a compound of Formula (II') wherein X is optionally substituted $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (II') wherein Y is optionally substituted aryl.

- 5 In another embodiment is a compound of Formula (II') wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (II') wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (II') wherein Y is optionally substituted $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (II') wherein Y is $-O-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (II') wherein Y is $-N(H)-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (II') wherein Y is a bond. In another embodiment is a compound of Formula (II') wherein Z is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (II') wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (II') wherein Z is optionally substituted phenyl. In another embodiment is a compound of Formula (II') wherein Z is optionally substituted heteroaryl. In another embodiment is a compound of Formula (II') wherein Z is optionally substituted $-(C_3-C_7)cycloalkyl$. In another embodiment is a compound of Formula (II') wherein Z is halogen.

[00254] In another embodiment is a compound of Formula (II') wherein $-X-Y-Z$ is



[00255] In another embodiment described herein is a compound of Formula (II) having the structure of Formula (IIa):



Formula (IIa);

wherein:

- R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$,
 5 $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, or $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, or optionally substituted heterocycloalkyl;
- R^3 is H, or $-(C_1-C_6)alkyl$;
- 10 R^4 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OH$, $-(C_3-C_6)cycloalkyl$, or $-C(O)NH_2$; or R^3 and R^4 are combined to form a heterocycloalkyl ring;
- R^5 is H, or $-(C_1-C_6)alkyl$; or R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring;
- R^6 , R^7 , and R^8 are each independently H, or $-(C_1-C_6)alkyl$;
- 15 R^9 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;
- R^{10} is H, or $-(C_1-C_6)alkyl$;
- R^{11} and R^{12} are each independently H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)heteroalkyl-CO_2H$, $-(C_1-C_6)alkyl-S(O)-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-N(H)CH=NH$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted $-(C_1-C_6)alkyl-N(H)heterocycloalkyl$, or $-(C_1-C_6)alkyl-heteroaryl$; or R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring, and R^{12} is H;
- 20 R^{13} and R^{14} are each independently H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(H)C(NH)NH_2$, $-(C_1-C_6)alkyl-heterocycloalkyl$, or $-(C_1-C_6)alkyl-heteroaryl$; or R^{13} and R^{19} are combined to form an optionally substituted heterocycloalkyl ring, and R^{14} is H;
- 25

R^{15} , R^{16} , R^{17} , R^{18} , and R^{19} are each independently H, $-(C_1-C_6)alkyl$, $-(C_3-C_6)cycloalkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$;

X is optionally substituted $-(C_1-C_6)alkyl$ -, $-(C_2-C_6)alkenyl$ -, $-(C_2-C_6)alkynyl$ -, $-(C_3-C_7)cycloalkyl$ -, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-O-(C_1-C_6)alkyl$ -, $-N(R^{24})(C_1-C_6)alkyl$ -, $-N(R^{24})(C_6-C_{10})aryl$ -, or $-SO_2(C_1-C_6)alkyl$ -;

Y is a bond, optionally substituted $-(C_1-C_6)alkyl$ -, $-(C_2-C_6)alkenyl$ -, $-(C_2-C_6)alkynyl$ -, $-(C_1-C_6)alkyl-N(R^{24})(C_1-C_6)alkyl$ -, $-O-(C_1-C_6)alkyl$ -, $-O(C_6-C_{10})aryl$ -, $-N(R^{24})(C_1-C_6)alkyl$ -, $-N(R^{24})SO_2(C_1-C_6)alkyl$ -, $-N(R^{24})C(O)(C_1-C_6)alkyl$ -, $-C(O)(C_1-C_6)alkyl$ -, $-S(C_1-C_6)alkyl$ -, $-SO_2(C_1-C_6)alkyl$ -, $-C(O)NH(C_1-C_6)alkyl$ -, $-(C_3-C_7)cycloalkyl$ -, optionally substituted $-C(O)N(R^{24})aryl$ -, optionally substituted $-N(R^{24})C(O)aryl$ -, optionally substituted $-N(R^{24})SO_2aryl$ -, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, $-NH_2$, $-CN$, $-CF_3$, $-(C_1-C_{12})alkyl$ -, $-(C_2-C_{12})alkenyl$ -, $-(C_2-C_{12})alkynyl$ -, $-C(O)NR^{25}R^{26}$, $-O-(C_1-C_{12})alkyl$ -, $-N(R^{24})(C_1-C_{12})alkyl$ -, $-N(R^{24})C(O)(C_1-C_{12})alkyl$ -, optionally substituted $-(C_3-C_7)cycloalkyl$ -, $-(C_1-C_6)alkyl-heterocycloalkyl$ -, optionally substituted aryl, or optionally substituted heteroaryl;

each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl$ -, $-(C_1-C_6)heteroalkyl$ -, $-(C_1-C_6)alkyl-CO_2H$ -, $-C(O)(C_1-C_6)alkyl$ -, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)alkyl$; or two R^{31} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R^{23} is independently H or $-(C_1-C_6)alkyl$;

each R^{24} is independently H or $-(C_1-C_6)alkyl$;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl$; or R^{25} and R^{26} and

the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R^{27} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$ -, or optionally substituted $-(C_1-C_6)heteroalkyl$;

each R^{28} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$ -, or optionally substituted $-(C_1-C_6)heteroalkyl$;

n is 0 or 1;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00256] In one embodiment is a compound of Formula (IIa) wherein n is 0. In another embodiment is a compound of Formula (IIa) wherein n is 1.

[00257] In another embodiment is a compound of Formula (IIa) wherein R^6 , R^7 , and R^8 are H.

[00258] In another embodiment is a compound of Formula (IIa) wherein R^{15} and R^{16} are H.

5 [00259] In one embodiment is a compound of Formula (IIa) wherein R^{17} is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (IIa) wherein R^{17} is $-\text{CH}_3$. In another embodiment is a compound of Formula (IIa) wherein R^{17} is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (IIa) wherein R^{17} is $-(C_3-C_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (IIa) wherein R^{17} is cyclopropyl. In another embodiment is a compound of Formula (IIa) wherein R^{17} is $-(C_1-C_6)\text{alkyl}-\text{C}(\text{O})\text{OR}^{23}$. In another embodiment is a compound of Formula (IIa) wherein R^{17} is $-\text{CH}_2\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (IIa) wherein R^{17} is $-(C_1-C_6)\text{alkyl}-\text{NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (IIa) wherein R^{17} is $-\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (IIa) wherein R^{17} is H.

15 [00260] In another embodiment is a compound of Formula (IIa) wherein R^{18} is H.

[00261] In another embodiment is a compound of Formula (IIa) wherein R^{19} is H.

[00262] In another embodiment is a compound of Formula (IIa) wherein R^3 is H.

[00263] In another embodiment is a compound of Formula (IIa) wherein R^5 is H.

[00264] In another embodiment is a compound of Formula (IIa) wherein R^4 is H. In another embodiment is a compound of Formula (IIa) wherein R^4 is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (IIa) wherein R^4 is $-\text{CH}_3$. In another embodiment is a compound of Formula (IIa) wherein R^4 is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (IIa) wherein R^4 is $-(C_1-C_6)\text{alkyl}-\text{OH}$. In another embodiment is a compound of Formula (IIa) wherein R^4 is $-\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (IIa) wherein R^4 is $-(C_3-C_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (IIa) wherein R^4 is cyclopropyl. In another embodiment is a compound of Formula (IIa) wherein R^4 is $-\text{C}(\text{O})\text{NH}_2$.

[00265] In another embodiment is a compound of Formula (IIa) wherein R^3 , R^4 , and R^5 are H.

30 [00266] In another embodiment is a compound of Formula (IIa) wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring.

[00267] In another embodiment is a compound of Formula (IIa) wherein R^{10} is H.

[00268] In another embodiment is a compound of Formula (IIa) wherein R^{10} is H and R^9 is $-(C_1-C_6)\text{alkyl}$. In another embodiment is a compound of Formula (IIa) wherein R^{10} is H and R^9 is $-\text{CH}_3$. In another embodiment is a compound of Formula (IIa) wherein R^{10} is H and R^9 is -

CH₂CH₃. In another embodiment is a compound of Formula (IIa) wherein R¹⁰ is H and R⁹ is - (C₁-C₆)haloalkyl. In another embodiment is a compound of Formula (IIa) wherein R¹⁰ is H and R⁹ is -CH₂F. In another embodiment is a compound of Formula (IIa) wherein R¹⁰ is H and R⁹ is -CHF₂. In another embodiment is a compound of Formula (IIa) wherein R¹⁰ is H and R⁹ is -(C₃-C₆)cycloalkyl. In another embodiment is a compound of Formula (IIa) wherein R¹⁰ is H and R⁹ is cyclopropyl. In another embodiment is a compound of Formula (IIa) wherein R¹⁰ is H and R⁹ is H.

[00269] In another embodiment is a compound of Formula (IIa) wherein R¹² is H.

[00270] In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is - (C₁-C₆)alkyl. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -CH₃. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is - (C₁-C₆)alkyl-OR²³. In another embodiment is a compound of Formula (IIa) wherein R¹¹ is -CH₂OH. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -CH₂CH₂OH. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-NH₂. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -CH₂NH₂. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -CH₂CH₂NH₂. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -CH₂CH₂CH₂CH₂NH₂. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-CN. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -CH₂CN. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -CH₂C(O)NH₂. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -CH₂CH₂C(O)NH₂. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is -(C₁-C₆)alkyl-heteroaryl. In another embodiment is a compound of Formula (IIa) wherein R¹² is H and R¹¹ is H.

[00271] In another embodiment is a compound of Formula (IIa) wherein R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring and R¹² is H.

[00272] In another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H.

[00273] In another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H and R¹³ is - (C₁-C₆)alkyl. In another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H and R¹³


- is -CH₃. In another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H and R¹³ is -
(C₁-C₆)alkyl-OR²³. In another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H
and R¹³ is -CH₂OH. In another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H
and R¹³ is -CH₂CH₂OH. In another embodiment is a compound of Formula (IIa) wherein R¹⁴ is
5 H and R¹³ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (IIa) wherein R¹⁴
is H and R¹³ is -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula (IIa)
wherein R¹⁴ is H and R¹³ is -(C₁-C₆)alkyl-NH₂. In another embodiment is a compound of
Formula (IIa) wherein R¹⁴ is H and R¹³ is -CH₂NH₂. In another embodiment is a compound of
Formula (IIa) wherein R¹⁴ is H and R¹³ is -CH₂CH₂NH₂. In another embodiment is a compound
10 of Formula (IIa) wherein R¹⁴ is H and R¹³ is -CH₂CH₂CH₂NH₂. In another embodiment is a
compound of Formula (IIa) wherein R¹⁴ is H and R¹³ is -CH₂CH₂CH₂CH₂NH₂. In another
embodiment is a compound of Formula (IIa) wherein R¹⁴ is H and R¹³ is -(C₁-C₆)alkyl-CN. In
another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H and R¹³ is -CH₂CN. In
another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H and R¹³ is -(C₁-C₆)alkyl-
15 C(O)NR²⁵R²⁶. In another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H and R¹³
is -CH₂C(O)NH₂. In another embodiment is a compound of Formula (IIa) wherein R¹⁴ is H and
R¹³ is -CH₂CH₂C(O)NH₂. In another embodiment is a compound of Formula (IIa) wherein R¹⁴
is H and R¹³ is -(C₁-C₆)alkyl-heteroaryl. In another embodiment is a compound of Formula (IIa)
wherein R¹⁴ is H and R¹³ is H.
- 20 **[00274]** In another embodiment is a compound of Formula (IIa) wherein R¹³ and R¹⁹ are
combined to form an optionally substituted heterocycloalkyl ring and R¹⁴ is H.
- [00275]** In another embodiment is a compound of Formula (IIa) wherein p is 1 and R²⁷ is
halogen. In another embodiment is a compound of Formula (IIa) wherein p is 1 and R²⁷ is
optionally substituted -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (IIa)
25 wherein q is 0, p is 1 and R²⁷ is halogen. In another embodiment is a compound of Formula
(IIa) wherein q is 0, p is 1 and R²⁷ is optionally substituted -(C₁-C₆)alkyl. In another
embodiment is a compound of Formula (IIa) wherein q is 1 and R²⁸ is halogen. In another
embodiment is a compound of Formula (IIa) wherein q is 1 and R²⁸ is optionally substituted -
(C₁-C₆)alkyl. In another embodiment is a compound of Formula (IIa) wherein p is 0, q is 1 and
30 R²⁸ is halogen. In another embodiment is a compound of Formula (IIa) wherein p is 0, q is 1
and R²⁸ is optionally substituted -(C₁-C₆)alkyl.
- [00276]** In another embodiment is a compound of Formula (IIa) wherein p is 0, and q is 0.
- [00277]** In another embodiment is a compound of Formula (IIa) wherein R¹ and R² are each
independently H, or -(C₁-C₆)alkyl-NR²¹R²². In another embodiment is a compound of Formula


(IIa) wherein R^1 and R^2 are each H. In another embodiment is a compound of Formula (IIa) wherein R^1 and R^2 are each independently $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (IIa) wherein R^1 is H, and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (IIa) wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$, and R^2 is H.

5 In another embodiment is a compound of Formula (IIa) wherein R^1 is H, and R^2 is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIa) wherein R^1 is $-CH_2CH_2NH_2$, and R^2 is H. In another embodiment is a compound of Formula (IIa) wherein R^1 and R^2 are each $-CH_2CH_2NH_2$.


[00278] In another embodiment is a compound of Formula (IIa) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (IIa) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (IIa) wherein X is optionally substituted heteroaryl. In another embodiment is a compound of Formula (IIa) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In another embodiment is a compound of Formula (IIa) wherein X is optionally substituted $-(C_1-C_6)alkyl-$.
 10 In another embodiment is a compound of Formula (IIa) wherein Y is optionally substituted aryl. In another embodiment is a compound of Formula (IIa) wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (IIa) wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (IIa) wherein Y is optionally substituted $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IIa) wherein Y is $-O-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IIa) wherein Y is $-N(H)-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IIa) wherein Y is a bond. In another embodiment is a compound of Formula (IIa) wherein Z is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIa) wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (IIa) wherein Z is optionally substituted phenyl.
 15 In another embodiment is a compound of Formula (IIa) wherein Z is optionally substituted heteroaryl. In another embodiment is a compound of Formula (IIa) wherein Z is optionally substituted $-(C_3-C_7)cycloalkyl$. In another embodiment is a compound of Formula (IIa) wherein Z is halogen.

[00279] In another embodiment is a compound of Formula (IIa) wherein $-X-Y-Z$ is





[00280] In another embodiment described herein is a compound of Formula (II) having the structure of Formula (IIb):



Formula (IIb);

wherein:

- R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, or $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, or optionally substituted heterocycloalkyl;
- R^4 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OH$, $-(C_3-C_6)cycloalkyl$, or $-C(O)NH_2$;
- R^5 is H, or $-(C_1-C_6)alkyl$; or R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring;
- R^9 is H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)haloalkyl$, or $-(C_3-C_6)cycloalkyl$;
- R^{11} is H, $-NH_2$, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-SR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-CN$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)heteroalkyl-$

CO₂H, -(C₁-C₆)alkyl-S(O)-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-N(H)CH=NH, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted -(C₁-C₆)alkyl-N(H)heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹¹ and R¹⁸ are combined to form an optionally substituted heterocycloalkyl ring;

5 R¹³ is H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹³ and R¹⁹ are combined to form an optionally substituted heterocycloalkyl ring;

R¹⁷, R¹⁸, and R¹⁹ are each independently H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;

X is optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₃-C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;

15 Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -

20 N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, -NH₂, -CN, -CF₃, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl-, -(C₂-C₁₂)alkynyl-, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl-, -N(R²⁴)(C₁-C₁₂)alkyl-, -N(R²⁴)C(O)(C₁-C₁₂)alkyl-, optionally substituted -(C₃-C₇)cycloalkyl-, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

25 each R²¹ and R²² is independently H, -(C₁-C₆)alkyl, -(C₁-C₆)heteroalkyl, -(C₁-C₆)alkyl-CO₂H, -C(O)(C₁-C₆)alkyl, -C(O)N(R³¹)₂, or -SO₂N(R³¹)₂; or R²¹ and R²² and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R³¹ is independently H or -(C₁-C₆)alkyl; or two R³¹ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

30 each R²³ is independently H or -(C₁-C₆)alkyl;

each R²⁴ is independently H or -(C₁-C₆)alkyl; and

each R²⁵ and R²⁶ is independently H or optionally substituted -(C₁-C₆)alkyl; or R²⁵ and R²⁶ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00281] In one embodiment is a compound of Formula (IIb) wherein R^{17} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIb) wherein R^{17} is $-CH_3$. In another embodiment is a compound of Formula (IIb) wherein R^{17} is $-CH_2CH_3$. In another embodiment is a compound of Formula (IIb) wherein R^{17} is $-(C_3-C_6)cycloalkyl$. In another embodiment is a compound of Formula (IIb) wherein R^{17} is cyclopropyl. In another embodiment is a compound of Formula (IIb) wherein R^{17} is $-(C_1-C_6)alkyl-C(O)OR^{23}$. In another embodiment is a compound of Formula (IIb) wherein R^{17} is $-CH_2CH_2OH$. In another embodiment is a compound of Formula (IIb) wherein R^{17} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (IIb) wherein R^{17} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{17} is H.

[00282] In another embodiment is a compound of Formula (IIb) wherein R^{18} is H.

[00283] In another embodiment is a compound of Formula (IIb) wherein R^{19} is H.

[00284] In another embodiment is a compound of Formula (IIb) wherein R^5 is H.

[00285] In another embodiment is a compound of Formula (IIb) wherein R^4 is H. In another

embodiment is a compound of Formula (IIb) wherein R^4 is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIb) wherein R^4 is $-CH_3$. In another embodiment is a compound of Formula (IIb) wherein R^4 is $-CH_2CH_3$. In another embodiment is a compound of Formula (IIb) wherein R^4 is $-(C_1-C_6)alkyl-OH$. In another embodiment is a compound of Formula (IIb) wherein R^4 is $-CH_2OH$. In another embodiment is a compound of Formula (IIb) wherein R^4 is $-(C_3-C_6)cycloalkyl$. In another embodiment is a compound of Formula (IIb) wherein R^4 is cyclopropyl. In another embodiment is a compound of Formula (IIb) wherein R^4 is $-C(O)NH_2$.

[00286] In another embodiment is a compound of Formula (IIb) wherein R^4 and R^5 are H.

[00287] In another embodiment is a compound of Formula (IIb) wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring.

[00288] In another embodiment is a compound of Formula (IIb) wherein R^9 is $-(C_1-C_6)alkyl$.

In another embodiment is a compound of Formula (IIb) wherein R^9 is $-CH_3$. In another embodiment is a compound of Formula (IIb) wherein R^9 is $-CH_2CH_3$. In another embodiment is a compound of Formula (IIb) wherein R^9 is $-(C_1-C_6)haloalkyl$. In another embodiment is a compound of Formula (IIb) wherein R^9 is $-CH_2F$. In another embodiment is a compound of Formula (IIb) wherein R^9 is $-CHF_2$. In another embodiment is a compound of Formula (IIb) wherein R^9 is $-(C_3-C_6)cycloalkyl$. In another embodiment is a compound of Formula (IIb) wherein R^9 is cyclopropyl. In another embodiment is a compound of Formula (IIb) wherein R^9 is H.

[00289] In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_3$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-(C_1-C_6)alkyl-OR^{23}$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_2OH$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_2CH_2OH$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-(C_1-C_6)alkyl-NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_2CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-(C_1-C_6)alkyl-CN$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_2CN$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_2C(O)NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-CH_2CH_2C(O)NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is $-(C_1-C_6)alkyl-heteroaryl$. In another embodiment is a compound of Formula (IIb) wherein R^{11} is H.

[00290] In another embodiment is a compound of Formula (IIb) wherein R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring and R^{12} is H.

[00291] In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-CH_3$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-(C_1-C_6)alkyl-OR^{23}$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-CH_2OH$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-CH_2CH_2OH$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-(C_1-C_6)alkyl-NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-CH_2CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-(C_1-C_6)alkyl-CN$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-CH_2CN$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-(C_1-C_6)alkyl-$

$C(O)NR^{25}R^{26}$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is -
 $CH_2C(O)NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is -
 $CH_2CH_2C(O)NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is $-(C_1-$
 $C_6)alkyl-heteroaryl$. In another embodiment is a compound of Formula (IIb) wherein R^{13} is H.


5 **[00292]** In another embodiment is a compound of Formula (IIb) wherein R^{13} and R^{19} are
combined to form an optionally substituted heterocycloalkyl ring and R^{14} is H.

[00293] In another embodiment is a compound of Formula (IIb) wherein R^1 and R^2 are each
independently H, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula
(IIb) wherein R^1 and R^2 are each H. In another embodiment is a compound of Formula (IIb)


10 wherein R^1 and R^2 are each independently $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a
compound of Formula (IIb) wherein R^1 is H, and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another
embodiment is a compound of Formula (IIb) wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$, and R^2 is H.
In another embodiment is a compound of Formula (IIb) wherein R^1 is H, and R^2 is -
 $CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIb) wherein R^1 is -
15 $CH_2CH_2NH_2$, and R^2 is H. In another embodiment is a compound of Formula (IIb) wherein R^1
and R^2 are each $-CH_2CH_2NH_2$.

[00294] In another embodiment is a compound of Formula (IIb) wherein X is optionally
substituted aryl. In another embodiment is a compound of Formula (IIb) wherein X is
optionally substituted phenyl. In another embodiment is a compound of Formula (IIb) wherein
20 X is optionally substituted heteroaryl. In another embodiment is a compound of Formula (IIb)
wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In another
embodiment is a compound of Formula (IIb) wherein X is optionally substituted $-(C_1-C_6)alkyl-$.
In another embodiment is a compound of Formula (IIb) wherein Y is optionally substituted aryl.
In another embodiment is a compound of Formula (IIb) wherein Y is optionally substituted
25 phenyl. In another embodiment is a compound of Formula (IIb) wherein Y is optionally
substituted heteroaryl. In another embodiment is a compound of Formula (IIb) wherein Y is
optionally substituted $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IIb)
wherein Y is $-O-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IIb) wherein
Y is $-N(H)-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IIb) wherein Y is
30 a bond. In another embodiment is a compound of Formula (IIb) wherein Z is $-(C_1-C_6)alkyl$. In
another embodiment is a compound of Formula (IIb) wherein Z is optionally substituted aryl. In
another embodiment is a compound of Formula (IIb) wherein Z is optionally substituted phenyl.
In another embodiment is a compound of Formula (IIb) wherein Z is optionally substituted
heteroaryl. In another embodiment is a compound of Formula (IIb) wherein Z is optionally

[00295] In another embodiment is a compound of Formula (IIb) wherein -X-Y-Z is



[00296] In another embodiment described herein is a compound of Formula (II) having the structure of Formula (IIc):



wherein:

R¹ and R² are each independently H or -CH₂CH₂NH₂;

15 R¹¹ is H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-SR²³, -(C₁-C₆)alkyl-C(O)OR²³,
-(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)heteroalkyl-
CO₂H, -(C₁-C₆)alkyl-S(O)-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-N(H)CH=NH, -(C₁-C₆)alkyl-

- N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted -(C₁-C₆)alkyl-N(H)heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl;
- R¹³ is H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-
- 5 N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl;
- X is optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₃-C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;
- 10 Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -
- 15 N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;
- Z is H, halogen, -NH₂, -CN, -CF₃, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl-, -(C₂-C₁₂)alkynyl, -C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl-, -N(R²⁴)(C₁-C₁₂)alkyl-, -N(R²⁴)C(O)(C₁-C₁₂)alkyl, optionally substituted -(C₃-C₇)cycloalkyl, -(C₁-C₆)alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;
- 20 each R²¹ and R²² is independently H, -(C₁-C₆)alkyl, -(C₁-C₆)heteroalkyl, -(C₁-C₆)alkyl-CO₂H, -C(O)(C₁-C₆)alkyl, -C(O)N(R³¹)₂, or -SO₂N(R³¹)₂; or R²¹ and R²² and the nitrogen atom to which they are attached form a heterocycloalkyl ring;
- each R³¹ is independently H or -(C₁-C₆)alkyl; or two R³¹ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;
- 25 each R²³ is independently H or -(C₁-C₆)alkyl;
- each R²⁴ is independently H or -(C₁-C₆)alkyl; and
- each R²⁵ and R²⁶ is independently H or optionally substituted -(C₁-C₆)alkyl; or R²⁵ and R²⁶ and the nitrogen atom to which they are attached form a heterocycloalkyl ring;
- or a pharmaceutically acceptable salt, solvate, or prodrug thereof.
- 30 **[00297]** In another embodiment is a compound of Formula (IIc) wherein R¹¹ is -(C₁-C₆)alkyl. In another embodiment is a compound of Formula (IIc) wherein R¹¹ is -CH₃. In another embodiment is a compound of Formula (IIc) wherein R¹¹ is -(C₁-C₆)alkyl-OR²³. In another embodiment is a compound of Formula (IIc) wherein R¹¹ is -CH₂OH. In another embodiment is a compound of Formula (IIc) wherein R¹¹ is -CH₂CH₂OH. In another embodiment is a

compound of Formula (IIc) wherein R^{11} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-(C_1-C_6)alkyl-NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-CH_2NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-CH_2CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-(C_1-C_6)alkyl-CN$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-CH_2CN$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-CH_2C(O)NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-CH_2CH_2C(O)NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is $-(C_1-C_6)alkyl-heteroaryl$. In another embodiment is a compound of Formula (IIc) wherein R^{11} is H.


[00298] In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_3$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-(C_1-C_6)alkyl-OR^{23}$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_2OH$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_2CH_2OH$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-(C_1-C_6)alkyl-NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_2NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_2CH_2CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-(C_1-C_6)alkyl-CN$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_2CN$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_2C(O)NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-CH_2CH_2C(O)NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is $-(C_1-C_6)alkyl-heteroaryl$. In another embodiment is a compound of Formula (IIc) wherein R^{13} is H.


[00299] In another embodiment is a compound of Formula (IIc) wherein R^1 and R^2 are each H. In another embodiment is a compound of Formula (IIc) wherein R^1 is H, and R^2 is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IIc) wherein R^1 is

$\text{CH}_2\text{CH}_2\text{NH}_2$, and R^2 is H. In another embodiment is a compound of Formula (IIc) wherein R^1 and R^2 are each $-\text{CH}_2\text{CH}_2\text{NH}_2$.

[00300] In another embodiment is a compound of Formula (IIc) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (IIc) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (IIc) wherein X is optionally substituted heteroaryl. In another embodiment is a compound of Formula (IIc) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In another embodiment is a compound of Formula (IIc) wherein X is optionally substituted $-(\text{C}_1\text{-C}_6)\text{alkyl-}$. In another embodiment is a compound of Formula (IIc) wherein Y is optionally substituted aryl. In another embodiment is a compound of Formula (IIc) wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (IIc) wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (IIc) wherein Y is optionally substituted $-(\text{C}_1\text{-C}_6)\text{alkyl-}$. In another embodiment is a compound of Formula (IIc) wherein Y is $-\text{O}-(\text{C}_1\text{-C}_6)\text{alkyl-}$. In another embodiment is a compound of Formula (IIc) wherein Y is $-\text{N}(\text{H})-(\text{C}_1\text{-C}_6)\text{alkyl-}$. In another embodiment is a compound of Formula (IIc) wherein Y is a bond. In another embodiment is a compound of Formula (IIc) wherein Z is $-(\text{C}_1\text{-C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (IIc) wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (IIc) wherein Z is optionally substituted phenyl. In another embodiment is a compound of Formula (IIc) wherein Z is optionally substituted heteroaryl. In another embodiment is a compound of Formula (IIc) wherein Z is optionally substituted $-(\text{C}_3\text{-C}_7)\text{cycloalkyl}$. In another embodiment is a compound of Formula (IIc) wherein Z is halogen.

[00301] In another embodiment is a compound of Formula (IIc) wherein $-\text{X-Y-Z}$ is





R¹ and R² are each independently H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -CH₂CH(OH)CH₂NH₂, -CH₂CH(heterocycloalkyl)CH₂NH₂, -CH₂C(O)NH₂, -CH₂C(O)N(H)CH₂CN, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-N(R²³)C(O)(C₁-C₆)alkylNR²¹R²², or -(C₁-C₆)alkyl-C(O)N(R²³)(C₁-C₆)alkyl, or optionally substituted heterocycloalkyl;

R⁵ is H, or -(C₁-C₆)alkyl; or R⁴ and R⁵ and the carbon atom to which they are attached form a cyclopropyl ring;

R¹⁷ is H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;

(C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-C₆)alkyl-;

Y is a bond, optionally substituted $-(C_1-C_6)alkyl-$, $-(C_2-C_6)alkenyl-$, $-(C_2-C_6)alkynyl-$, $-(C_1-C_6)alkyl-N(R^{24})(C_1-C_6)alkyl-$, $-O-(C_1-C_6)alkyl-$, $-O(C_6-C_{10})aryl-$, $-N(R^{24})(C_1-C_6)alkyl-$, $-N(R^{24})SO_2(C_1-C_6)alkyl-$, $-N(R^{24})C(O)(C_1-C_6)alkyl-$, $-C(O)(C_1-C_6)alkyl-$, $-S(C_1-C_6)alkyl-$, $-SO_2(C_1-C_6)alkyl-$, $-C(O)NH(C_1-C_6)alkyl-$, $-(C_3-C_7)cycloalkyl-$, optionally substituted -

5 $C(O)N(R^{24})aryl-$, optionally substituted $-N(R^{24})C(O)aryl-$, optionally substituted $-N(R^{24})SO_2aryl-$, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, $-NH_2$, $-CN$, $-CF_3$, $-(C_1-C_{12})alkyl-$, $-(C_2-C_{12})alkenyl-$, $-(C_2-C_{12})alkynyl-$, $-C(O)NR^{25}R^{26}$, $-O-(C_1-C_{12})alkyl-$, $-N(R^{24})(C_1-C_{12})alkyl-$, $-N(R^{24})C(O)(C_1-C_{12})alkyl-$, optionally substituted $-(C_3-C_7)cycloalkyl-$, $-(C_1-C_6)alkyl-heterocycloalkyl-$, optionally substituted aryl, or

10 optionally substituted heteroaryl;

each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl-$, $-(C_1-C_6)heteroalkyl-$, $-(C_1-C_6)alkyl-CO_2H$, $-C(O)(C_1-C_6)alkyl-$, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)alkyl-$; or two R^{31} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

15 each R^{23} is independently H or $-(C_1-C_6)alkyl-$;

each R^{24} is independently H or $-(C_1-C_6)alkyl-$; and

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl-$; or R^{25} and R^{26} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;

20 or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

[00303] In one embodiment is a compound of Formula (IId) wherein R^{17} is $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IId) wherein R^{17} is $-CH_3$. In another embodiment is a compound of Formula (IId) wherein R^{17} is $-CH_2CH_3$. In another embodiment is a compound of Formula (IId) wherein R^{17} is $-(C_3-C_6)cycloalkyl-$. In another embodiment is a compound of Formula (IId) wherein R^{17} is cyclopropyl. In another embodiment is a compound of Formula (IId) wherein R^{17} is $-(C_1-C_6)alkyl-C(O)OR^{23}$. In another embodiment is a compound of Formula (IId) wherein R^{17} is $-CH_2CH_2OH$. In another embodiment is a compound of Formula (IId) wherein R^{17} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$. In another embodiment is a compound of Formula (IId) wherein R^{17} is $-CH_2CH_2NH_2$. In another embodiment is a compound of Formula (IId) wherein R^{17} is H.

[00304] In another embodiment is a compound of Formula (IId) wherein R^5 is H.

[00305] In another embodiment is a compound of Formula (IId) wherein R^4 is H. In another embodiment is a compound of Formula (IId) wherein R^4 is $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IId) wherein R^4 is $-CH_3$. In another embodiment is a

compound of Formula (IId) wherein R^4 is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (IId) wherein R^4 is $-(\text{C}_1\text{-C}_6)\text{alkyl-OH}$. In another embodiment is a compound of Formula (IId) wherein R^4 is $-\text{CH}_2\text{OH}$. In another embodiment is a compound of Formula (IId) wherein R^4 is $-(\text{C}_3\text{-C}_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (IId) wherein R^4 is cyclopropyl. In another embodiment is a compound of Formula (IId) wherein R^4 is $-\text{C}(\text{O})\text{NH}_2$.

[00306] In another embodiment is a compound of Formula (IId) wherein R^4 and R^5 are H.

[00307] In another embodiment is a compound of Formula (IId) wherein R^4 and R^5 and the carbon atom to which they are attached form a cyclopropyl ring.

[00308] In another embodiment is a compound of Formula (IId) wherein R^9 is $-(\text{C}_1\text{-C}_6)\text{alkyl}$. In another embodiment is a compound of Formula (IId) wherein R^9 is $-\text{CH}_3$. In another embodiment is a compound of Formula (IId) wherein R^9 is $-\text{CH}_2\text{CH}_3$. In another embodiment is a compound of Formula (IId) wherein R^9 is $-(\text{C}_1\text{-C}_6)\text{haloalkyl}$. In another embodiment is a compound of Formula (IId) wherein R^9 is $-\text{CH}_2\text{F}$. In another embodiment is a compound of Formula (IId) wherein R^9 is $-\text{CHF}_2$. In another embodiment is a compound of Formula (IId) wherein R^9 is $-(\text{C}_3\text{-C}_6)\text{cycloalkyl}$. In another embodiment is a compound of Formula (IId) wherein R^9 is cyclopropyl. In another embodiment is a compound of Formula (IId) wherein R^9 is H.

[00309] In another embodiment is a compound of Formula (IId) wherein R^1 and R^2 are each independently H, or $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (IId) wherein R^1 and R^2 are each H. In another embodiment is a compound of Formula (IId) wherein R^1 and R^2 are each independently $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (IId) wherein R^1 is H, and R^2 is $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$. In another embodiment is a compound of Formula (IId) wherein R^1 is $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$, and R^2 is H. In another embodiment is a compound of Formula (IId) wherein R^1 is H, and R^2 is $-\text{CH}_2\text{CH}_2\text{NH}_2$. In another embodiment is a compound of Formula (IId) wherein R^1 is $-\text{CH}_2\text{CH}_2\text{NH}_2$, and R^2 is H. In another embodiment is a compound of Formula (IId) wherein R^1 and R^2 are each $-\text{CH}_2\text{CH}_2\text{NH}_2$.

[00310] In another embodiment is a compound of Formula (IId) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (IId) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (IId) wherein X is optionally substituted heteroaryl. In another embodiment is a compound of Formula (IId) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In another embodiment is a compound of Formula (IId) wherein X is optionally substituted $-(\text{C}_1\text{-C}_6)\text{alkyl}$.

In another embodiment is a compound of Formula (IId) wherein Y is optionally substituted aryl.

In another embodiment is a compound of Formula (IId) wherein Y is optionally substituted

phenyl. In another embodiment is a compound of Formula (IId) wherein Y is optionally

substituted heteroaryl. In another embodiment is a compound of Formula (IId) wherein Y is

5 optionally substituted $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IId)

wherein Y is $-O-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IId) wherein

Y is $-N(H)-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (IId) wherein Y is

a bond. In another embodiment is a compound of Formula (IId) wherein Z is $-(C_1-C_6)alkyl$. In

another embodiment is a compound of Formula (IId) wherein Z is optionally substituted aryl. In

10 another embodiment is a compound of Formula (IId) wherein Z is optionally substituted phenyl.


In another embodiment is a compound of Formula (IId) wherein Z is optionally substituted

heteroaryl. In another embodiment is a compound of Formula (IId) wherein Z is optionally


substituted $-(C_3-C_7)cycloalkyl$. In another embodiment is a compound of Formula (IId) wherein

Z is halogen.

15 **[00311]** In another embodiment is a compound of Formula (IId) wherein $-X-Y-Z$ is



20 **[00312]** In another embodiment described herein are compounds of Formula (II) having the structure of Formula (IIe):



Formula (Ile);

wherein:

R^1 and R^2 are each independently H or $-\text{CH}_2\text{CH}_2\text{NH}_2$;

5 X is optionally substituted $-(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-(\text{C}_2\text{-C}_6)\text{alkenyl-}$, $-(\text{C}_2\text{-C}_6)\text{alkynyl-}$, $-(\text{C}_3\text{-C}_7)\text{cycloalkyl-}$, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-\text{O}-(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{N}(\text{R}^{24})(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{N}(\text{R}^{24})(\text{C}_6\text{-C}_{10})\text{aryl-}$, or $-\text{SO}_2(\text{C}_1\text{-C}_6)\text{alkyl-}$;

10 Y is a bond, optionally substituted $-(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-(\text{C}_2\text{-C}_6)\text{alkenyl-}$, $-(\text{C}_2\text{-C}_6)\text{alkynyl-}$, $-(\text{C}_1\text{-C}_6)\text{alkyl-N}(\text{R}^{24})(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{O}-(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{O}(\text{C}_6\text{-C}_{10})\text{aryl-}$, $-\text{N}(\text{R}^{24})(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{N}(\text{R}^{24})\text{SO}_2(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{N}(\text{R}^{24})\text{C}(\text{O})(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{C}(\text{O})(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{S}(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{SO}_2(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6)\text{alkyl-}$, $-(\text{C}_3\text{-C}_7)\text{cycloalkyl-}$, optionally substituted $-\text{C}(\text{O})\text{N}(\text{R}^{24})\text{aryl-}$, optionally substituted $-\text{N}(\text{R}^{24})\text{C}(\text{O})\text{aryl-}$, optionally substituted $-\text{N}(\text{R}^{24})\text{SO}_2\text{aryl-}$, optionally substituted aryl, or optionally substituted heteroaryl;

15 Z is H, halogen, $-\text{NH}_2$, $-\text{CN}$, $-\text{CF}_3$, $-(\text{C}_1\text{-C}_{12})\text{alkyl-}$, $-(\text{C}_2\text{-C}_{12})\text{alkenyl-}$, $-(\text{C}_2\text{-C}_{12})\text{alkynyl-}$, $-\text{C}(\text{O})\text{NR}^{25}\text{R}^{26}$, $-\text{O}-(\text{C}_1\text{-C}_{12})\text{alkyl-}$, $-\text{N}(\text{R}^{24})(\text{C}_1\text{-C}_{12})\text{alkyl-}$, $-\text{N}(\text{R}^{24})\text{C}(\text{O})(\text{C}_1\text{-C}_{12})\text{alkyl-}$, optionally substituted $-(\text{C}_3\text{-C}_7)\text{cycloalkyl-}$, $-(\text{C}_1\text{-C}_6)\text{alkyl-heterocycloalkyl-}$, optionally substituted aryl, or optionally substituted heteroaryl;

each R^{24} is independently H or $-(\text{C}_1\text{-C}_6)\text{alkyl}$; and

20 R^{25} and R^{26} is independently H or optionally substituted $-(\text{C}_1\text{-C}_6)\text{alkyl}$; or R^{25} and R^{26} and the nitrogen atom to which they are attached form a heterocycloalkyl ring;


or a pharmaceutically acceptable salt, solvate, or prodrug thereof.


[00313] In one embodiment is a compound of Formula (Ile) wherein R^1 and R^2 are each H. In another embodiment is a compound of Formula (Ile) wherein R^1 is H, and R^2 is $-\text{CH}_2\text{CH}_2\text{NH}_2$.

25 In another embodiment is a compound of Formula (Ile) wherein R^1 is $-\text{CH}_2\text{CH}_2\text{NH}_2$, and R^2 is H. In another embodiment is a compound of Formula (Ile) wherein R^1 and R^2 are each $-\text{CH}_2\text{CH}_2\text{NH}_2$.

[00314] In another embodiment is a compound of Formula (Ile) wherein X is optionally substituted aryl. In another embodiment is a compound of Formula (Ile) wherein X is optionally substituted phenyl. In another embodiment is a compound of Formula (Ile) wherein X is optionally substituted heteroaryl. In another embodiment is a compound of Formula (Ile) wherein X is optionally substituted pyridine or optionally substituted pyrimidine. In another embodiment is a compound of Formula (Ile) wherein X is optionally substituted $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (Ile) wherein Y is optionally substituted aryl. In another embodiment is a compound of Formula (Ile) wherein Y is optionally substituted phenyl. In another embodiment is a compound of Formula (Ile) wherein Y is optionally substituted heteroaryl. In another embodiment is a compound of Formula (Ile) wherein Y is optionally substituted $-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (Ile) wherein Y is $-O-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (Ile) wherein Y is $-N(H)-(C_1-C_6)alkyl-$. In another embodiment is a compound of Formula (Ile) wherein Y is a bond. In another embodiment is a compound of Formula (Ile) wherein Z is $-(C_1-C_6)alkyl$. In another embodiment is a compound of Formula (Ile) wherein Z is optionally substituted aryl. In another embodiment is a compound of Formula (Ile) wherein Z is optionally substituted phenyl. In another embodiment is a compound of Formula (Ile) wherein Z is optionally substituted heteroaryl. In another embodiment is a compound of Formula (Ile) wherein Z is optionally substituted $-(C_3-C_7)cycloalkyl$. In another embodiment is a compound of Formula (Ile) wherein Z is halogen.

[00315] In another embodiment is a compound of Formula (Ile) wherein -X-Y-Z is





[00316] In another aspect are hydrates or metabolites comprising any of the aforementioned compounds.

5 **[00317]** In another aspect are pharmaceutical compositions comprising any of the aforementioned compounds together with a pharmaceutically acceptable excipient.

[00318] In another aspect described herein is the use of a compound described herein in the manufacture of a medicament for treatment of a bacterial infection in a patient.

[00319] In another aspect are methods of treating a mammal in need of such treatment
 10 comprising administering to the mammal an antibacterial effective amount of any of the aforementioned compounds at a frequency and for a duration sufficient to provide a beneficial effect to the mammal. In one embodiment, the mammal has a bacteria-related infection that is resistant to treatment with arylomycin A2. In a further embodiment, the causative bacteria species of the bacteria infection is an infection involving *Pseudomonas aeruginosa*,
 15 *Pseudomonas fluorescens*, *Pseudomonas acidovorans*, *Pseudomonas alcaligenes*, *Pseudomonas putida*, *Stenotrophomonas maltophilia*, *Burkholderia cepacia*, *Aeromonas hydrophila*, *Escherichia coli*, *Citrobacter freundii*, *Salmonella typhimurium*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella enteritidis*, *Shigella dysenteriae*, *Shigella flexneri*, *Shigella sonnei*, *Enterobacter cloacae*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Klebsiella oxytoca*,
 20 *Serratia marcescens*, *Francisella tularensis*, *Morganella morganii*, *Proteus mirabilis*, *Proteus vulgaris*, *Providencia alcalifaciens*, *Providencia rettgeri*, *Providencia stuartii*, *Acinetobacter baumannii*, *Acinetobacter calcoaceticus*, *Acinetobacter haemolyticus*, *Yersinia enterocolitica*, *Yersinia pestis*, *Yersinia pseudotuberculosis*, *Yersinia intermedia*, *Bordetella pertussis*, *Bordetella parapertussis*, *Bordetella bronchiseptica*, *Haemophilus influenzae*, *Haemophilus parainfluenzae*, *Haemophilus haemolyticus*, *Haemophilus parahaemolyticus*, *Haemophilus ducreyi*, *Pasteurella multocida*, *Pasteurella haemolytica*, *Branhamella catarrhalis*, *Helicobacter pylori*, *Campylobacter fetus*, *Campylobacter jejuni*, *Campylobacter coli*, *Borrelia burgdorferi*, *Vibrio cholerae*, *Vibrio parahaemolyticus*, *Legionella pneumophila*, *Listeria monocytogenes*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Kingella*, *Moraxella*, *Gardnerella vaginalis*,

Bacteroides fragilis, Bacteroides distasonis, Bacteroides 3452A homology group, Bacteroides vulgatus, Bacteroides ovalus, Bacteroides thetaiotaomicron, Bacteroides uniformis, Bacteroides eggerthii, Bacteroides splanchnicus, Clostridium difficile, Mycobacterium tuberculosis, Mycobacterium avium, Mycobacterium intracellulare, Mycobacterium leprae, Corynebacterium diphtheriae, Corynebacterium ulcerans, Streptococcus pneumoniae, Streptococcus agalactiae, Streptococcus pyogenes, Enterococcus faecalis, Enterococcus faecium, Staphylococcus aureus, Staphylococcus epidermidis, Staphylococcus saprophyticus, Staphylococcus intermedius, Staphylococcus hyicus subsp. hyicus, Staphylococcus haemolyticus, Staphylococcus hominis, or Staphylococcus saccharolyticus. In another embodiment the bacterial infection is an infection involving a Gram-negative bacteria. In a further embodiment, the bacterial infection is an infection involving a Gram-positive bacteria.

[00320] In a further embodiment are methods of treating a mammal in need of such treatment comprising administering to the mammal a second therapeutic agent to any of the aforementioned methods of treatment. In another embodiment, the second therapeutic agent is a not an SpsB inhibitor. In another embodiment, the second therapeutic agent is an aminoglycoside antibiotic, fluoroquinolone antibiotic, β -lactam antibiotic, macrolide antibiotic, glycopeptide antibiotic, rifampicin, chloramphenicol, fluoramphenicol, colistin, mupirocin, bacitracin, daptomycin, or linezolid.

[00321] In some embodiments is a method for treating a bacterial infection in a patient, preferably a human, where the treatment includes administering a therapeutically or pharmacologically effective amount of a combination of 1) a β -lactam antibiotic; and 2) a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), or a pharmaceutically acceptable salt thereof; and 3) a pharmaceutically acceptable carrier. In embodiments where a β -lactam antibiotic is used in combination with a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), the β -lactam antibiotic may be a carbapenem, cephalosporin, cephamycin, monobactam or penicillin. Exemplary carbapenem antibiotics useful in the methods of the invention include ertapenem, imipenem, biapenem, and meropenem. Exemplary cephalosporin antibiotics useful in the methods of the invention include, ceftobiprole, ceftaroline, Cefiprome, Cefozopran, cefepime, Cefotaxime, and ceftriazone. Exemplary penicillin antibiotics useful in the methods of the invention include ampicillin, amoxicillin, piperacillin, oxacillin, cloxacillin, methicillin, and nafcillin. In some embodiments of the invention, the β -lactam may be administered with a β -lactamase inhibitor. In some embodiments of the invention, the carbapenem may be administered with a DHP inhibitor, e.g., cilastatin.

[00322] In various embodiments of the invention where a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) and a β -lactam antibiotic are used in combination, the β -lactam antibiotic and compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) can be administered sequentially or concurrently.

5 Preferably, the β -lactam antibiotic and compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) are administered together. When administered concurrently, the β -lactam antibiotic and compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) may be administered in the same formulation or in separate formulations. When administered sequentially, either the β -lactam or compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) may be administered first.

10 After administration of the first compound, the other compound is administered, for example, within from 1 to 60 minutes, e.g., within 1, 2, 3, 4, 5, 10, 15, 30, or 60 minutes. In one aspect of the invention, when a β -lactamase inhibitor is used, it may be administered separately, or in a formulation with the compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) and/or β -lactam antibiotic. In one aspect of the invention, when a DHP inhibitor is used to improve the stability of a carbapenem, it may be administered separately, or in a formulation with the compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) and/or carbapenem.

[00323] Further described herein are pharmaceutical compositions comprising a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), a pharmaceutically acceptable carrier, and optionally a β -lactam antibiotic. In embodiments where a combination is used, the β -lactam antibiotic and the compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), are present in such amounts that their combination constitutes a therapeutically effective amount. Due to the potentiating effects of the compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), the amount of β -lactam antibiotic present in a combination may be less than that of a β -lactam antibiotic used alone. In certain embodiments, the composition further comprises a β -lactamase antibiotic.

[00324] In further embodiments where the β -lactam antibiotic is a carbapenem, is provided a pharmaceutical composition comprising a carbapenem antibiotic, a DHP inhibitor, a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe), and a pharmaceutically acceptable carrier. In some embodiments where the β -lactam antibiotic is a carbapenem, the carbapenem antibiotic is preferably selected from the group consisting of ertapenem, imipenem, and meropenem.

[00325] In some embodiments is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId) for use in treating a bacterial infection. In some embodiments is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId), in combination with one or more additional therapeutical agents including a β -lactam antibiotic, for use in treating a bacterial infection. In some embodiments is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId), in combination with one or more additional therapeutical agents including a β -lactam antibiotic, for use as a medicament for treating a bacterial infection. In some embodiments is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId), in combination with one or more additional therapeutical agents including a β -lactam antibiotic, for use as a medicament for treating a bacterial infection. In some embodiments is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId), in combination with one or more additional therapeutical agents including a β -lactam antibiotic, for use in the preparation of a medicament for treating a bacterial infection. In some embodiments is a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId), in combination with one or more additional therapeutical agents including a β -lactam antibiotic, for use in the preparation of a medicament for treating a bacterial infection.

[00326] In some embodiments described herein, a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId) can enhance the activity of a β -lactam antibacterial agent by inducing susceptibility to the antibacterial agent in a drug-resistant strain such as MRSA. In some embodiments, a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId) can enhance the activity of a β -lactam antibacterial agent by reducing the dosage of the antibacterial agent need for a therapeutic effect in a drug-sensitive strain. For example, if a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId) reduces the Minimum Inhibitory Concentration (MIC) of an antibacterial agent (where the MIC is the minimum concentration of antibacterial agent which will completely inhibit growth) in a susceptible strain, then such treatment may be advantageous to enable a reduction in the amount of antibacterial agent administered (could reduce side effects of an antibiotic), or to decrease the frequency of administration. In some embodiments, compounds of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId) can enhance the activity of an antibacterial agent such as a carbapenem to prevent the emergence of a resistant sub-population in a heterogeneous bacterial population with a resistant sub-population.

[00327] Potentiators can be used to enhance the activity of antibacterial agents whose clinical efficacy has been limited by the increasing prevalence of resistant strains. In some embodiments described herein, a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IId) is used as a potentiator wherein a compound of Formula (I), (I'),

(Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) can be administered together with a β -lactam antibiotic (either concurrently or sequentially) to allow effective treatment of an infection involving a resistant bacterium, or to reduce the amount of antibacterial agent necessary to treat an infection.

- 5 **[00328]** In one embodiment, is a compound described herein which displays antibiotic activity useful in the treatment of bacterial infections, such as by way of example only, various strains of *S. aureus*, *S. pneumoniae*, *E. faecalis*, *E. faecium*, *B. subtilis* and *E. coli* including species that are resistant to many known antibiotics such as methicillin-resistant *S. aureus* (MRSA), vancomycin-resistant *Enterococcus* sp. (VRE), multidrug-resistant *E. faecium*, macrolide-resistant *S. aureus* and *S. epidermidis*, and linezolid-resistant *S. aureus* and *E. faecium*.

Methicillin-Resistant *Staphylococcus aureus*

- [00329]** *Staphylococcus aureus* (*S. aureus*), a spherical bacterium, is the most common cause of staph infections. *S. aureus* has been known to cause a range of illnesses from minor skin infections, such as pimples, impetigo, boils, cellulitis folliculitis, furuncles, carbuncles, scalded skin syndrome, abscesses, to life-threatening diseases such as pneumonia, meningitis, osteomyelitis endocarditis, toxic shock syndrome, and septicemia. Further, *S. aureus* is one of the most common causes of nosocomial infections, often causing postsurgical wound infections.

- [00330]** Methicillin was introduced in the late 1950s to treat infections caused by penicillin-resistant *S. aureus*. It has been reported previously that *S. aureus* isolates had acquired resistance to methicillin (methicillin-resistant *S. aureus*, MRSA). The methicillin resistance gene (*mecA*) encodes a methicillin-resistant penicillin-binding protein that is not present in susceptible strains. *mecA* is carried on a mobile genetic element, the staphylococcal cassette chromosome *mec* (SCC*mec*), of which four forms have been described that differ in size and genetic composition. The methicillin-resistant penicillin-binding protein allows for resistance to β -lactam antibiotics and obviates their clinical use during MRSA infections.

- [00331]** In one aspect is a method for treating a subject having a resistant bacterium comprising administering to the subject a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof. In one embodiment, the bacterium is a Gram-positive bacteria. In another embodiment, the Gram-positive bacterium is *S. aureus*. In further embodiment, the *S. aureus* is resistant or refractory to a beta-lactam antibiotic. In yet a further embodiment, the beta-lactam antibiotic belongs to the class of penicillins. In a further embodiment, the beta-lactam antibiotic is methicillin. In yet another embodiment, the subject has a methicillin-resistant *S. aureus* bacteria. In one embodiment the

beta-lactam antibiotic is flucloxacillin. In another embodiment is a method for treating a subject having a dicloxacillin-resistant bacteria comprising administering to the subject a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt,

5 stereoisomer, tautomer or prodrug thereof wherein the subject is refractory to dicloxacillin. Also disclosed herein is a method for treating a subject having a methicillin-resistant bacteria comprising administering a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the subject has
10 been determined to have a methicillin-resistant bacteria. In one embodiment the subject is screened for methicillin-resistant bacteria. In another embodiment, the subject screening is performed through a nasal culture. In a further embodiment the methicillin-resistant bacteria is detected by swabbing the nostril(s) of the subject and isolating the bacteria. In another embodiment, Real-time PCR and/or Quantitative PCR is employed to determine whether the
15 subject has a methicillin-resistant bacteria.

[00332] In one embodiment is a method for treating a subject having a first-generation cephalosporin-resistant bacteria comprising administering a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof
20 wherein the subject is refractory to a first-generation cephalosporin. In one embodiment, the bacteria is resistant to a first-generation cephalosporin. In a further embodiment, the bacteria is resistant to cefacetrile. In another embodiment, the bacteria is resistant to cefadroxil. In yet another embodiment, the bacteria is resistant to cefalexin. In one embodiment, the bacteria is resistant to cefaloglycin. In another embodiment, the bacteria is resistant to cefalonium. In
25 another embodiment, the bacteria is resistant to cefaloridine. In yet another embodiment, the bacteria is resistant to cefalotin. In a further embodiment, the bacteria is resistant to cefapirin. In yet a further embodiment, the bacteria is resistant to cefatrizine. In one embodiment, the bacteria is resistant to cefazaflur. In another embodiment, the bacteria is resistant to cefazedone. In yet another embodiment, the bacteria is resistant to cefazolin. In a further embodiment, the bacteria
30 is resistant to cefradine. In yet a further embodiment, the bacteria is resistant to cefroxadine. In one embodiment, the bacteria is resistant to ceftazidime.

[00333] In one embodiment is a method for treating a subject having a second-generation cephalosporin-resistant bacteria comprising administering a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt,

ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the subject is refractory to a second-generation cephalosporin. In another embodiment, the bacteria is resistant to a second-generation cephalosporin. In a further embodiment, the bacteria is resistant to cefaclor. In another embodiment, the bacteria is resistant to cefonicid. In yet another embodiment, the bacteria is resistant to cefprozil. In one embodiment, the bacteria is resistant to cefuroxime. In another embodiment, the bacteria is resistant to cefuzonam. In another embodiment, the bacteria is resistant to cefmetazole. In yet another embodiment, the bacteria is resistant to cefotetan. In a further embodiment, the bacteria is resistant to cefoxitin.

[00334] In one embodiment is a method for treating a subject having a third-generation cephalosporin-resistant bacteria comprising administering a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the subject is refractory to a third-generation cephalosporin. In another embodiment, the bacteria is resistant to a third-generation cephalosporin. In a further embodiment, the bacteria is resistant to cefcapene. In another embodiment, the bacteria is resistant to cefdaloxime. In yet another embodiment, the bacteria is resistant to cefdinir. In one embodiment, the bacteria is resistant to cefditoren. In another embodiment, the bacteria is resistant to cefixime. In another embodiment, the bacteria is resistant to cefmenoxime. In yet another embodiment, the bacteria is resistant to cefodizime. In a further embodiment, the bacteria is resistant to cefotaxime. In yet a further embodiment, the bacteria is resistant to cefpimizole. In one embodiment, the bacteria is resistant to cefpodoxime. In another embodiment, the bacteria is resistant to cefteram. In yet another embodiment, the bacteria is resistant to ceftibuten. In a further embodiment, the bacteria is resistant to ceftiofur. In yet a further embodiment, the bacteria is resistant to ceftiole. In one embodiment, the bacteria is resistant to ceftizoxime. In another embodiment, the bacteria is resistant to ceftriaxone. In yet another embodiment, the bacteria is resistant to cefoperazone. In yet a further embodiment, the bacteria is resistant to ceftazidime.

[00335] In one embodiment is a method for treating a subject having a fourth-generation cephalosporin-resistant bacteria comprising administering a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the subject is refractory to a fourth-generation cephalosporin. In another embodiment, the bacteria is resistant to a fourth-generation cephalosporin. In a further embodiment, the bacteria is resistant to cefclidine. In another embodiment, the bacteria is resistant to cefepime. In yet another embodiment, the bacteria is resistant to ceftuprenam. In one embodiment, the

bacteria is resistant to cefoselis. In another embodiment, the bacteria is resistant to ceftazidime. In another embodiment, the bacteria is resistant to ceftiofur. In yet another embodiment, the bacteria is refractory to cefquinome.

[00336] In one embodiment is a method for treating a subject having a carbapenem-resistant bacteria comprising administering a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the subject is refractory to a carbapenem. In another embodiment, the bacteria is resistant to a carbapenem. In a further embodiment, the bacteria is resistant to imipenem. In another embodiment, the bacteria is resistant to meropenem. In yet another embodiment, the bacteria is resistant to ertapenem. In one embodiment, the bacteria is resistant to faropenem. In another embodiment, the bacteria is resistant to doripenem. In another embodiment, the bacteria is resistant to panipenem. In yet another embodiment, the bacteria is resistant to biapenem,

Vancomycin-Intermediate and Vancomycin-Resistant Staphylococcus aureus

[00337] Vancomycin-intermediate Staphylococcus aureus and vancomycin-resistant staphylococcus aureus are specific types of antimicrobial-resistant Staph bacteria that are refractory to vancomycin treatment. S. aureus isolates for which vancomycin MICs are 4-8 µg/mL are classified as vancomycin-intermediate and isolates for which vancomycin MICs are ≥16 µg/mL are classified as vancomycin-resistant (Clinical and Laboratory Standards Institute/NCCLS. Performance Standards for Antimicrobial Susceptibility Testing. Sixteenth informational supplement. M100-S16. Wayne, PA: CLSI, 2006).

[00338] As used herein, the term “minimum inhibitory concentration” (MIC) refers to the lowest concentration of an antibiotic that is needed to inhibit growth of a bacterial isolate in vitro. A common method for determining the MIC of an antibiotic is to prepare several tubes containing serial dilutions of the antibiotic, that are then inoculated with the bacterial isolate of interest. The MIC of an antibiotic is determined from the tube with the lowest concentration that shows no turbidity (no growth).

[00339] In one aspect is a method of treating a subject having a bacterial infection comprising administering to the subject a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the bacterial infection comprises a vancomycin-intermediate Staphylococcus aureus bacterium. In one embodiment, the vancomycin-intermediate Staphylococcus aureus bacterium has a MIC of between about 4 to about 8 µg/mL. In another embodiment, the vancomycin-intermediate

Staphylococcus aureus bacterium has a MIC of about 4 µg/mL. In yet another embodiment, the vancomycin-intermediate Staphylococcus aureus bacterium has a MIC of about 5 µg/mL. In a further embodiment, the vancomycin-intermediate Staphylococcus aureus bacterium has a MIC of about 6 µg/mL. In yet a further embodiment, the vancomycin-intermediate Staphylococcus aureus bacterium has a MIC of about 7 µg/mL. In one embodiment, the vancomycin-intermediate Staphylococcus aureus bacterium has a MIC of about 8 µg/mL.

[00340] In another aspect is a method of treating a subject having a bacterial infection comprising administering to the subject a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the bacterial infection comprises a vancomycin-resistant Staphylococcus aureus bacterium. In one embodiment, the vancomycin-resistant Staphylococcus aureus bacterium has a MIC of between about 16 µg/mL. In another embodiment, the vancomycin-resistant Staphylococcus aureus bacterium has a MIC of about ≥ 16 µg/mL. In yet another embodiment, the vancomycin-resistant Staphylococcus aureus bacterium has a MIC of about 20 µg/mL. In a further embodiment, the vancomycin-resistant Staphylococcus aureus bacterium has a MIC of about 25 µg/mL.

[00341] In one embodiment, conditions treated by the compounds described herein include, but are not limited to, endocarditis, osteomyelitis, meningitis, skin and skin structure infections, genitourinary tract infections, abscesses, and necrotizing infections. In another embodiment, the compounds disclosed herein are used to treat conditions, such as, but not limited to, diabetic foot infections, decubitus ulcers, burn infections, animal or human bite wound infections, synergistic-necrotizing gangrene, necrotizing fasciitis, intra-abdominal infection associated with breaching of the intestinal barrier, pelvic infection associated with breaching of the intestinal barrier, aspiration pneumonia, and post-operative wound infections. In another embodiment, the conditions listed herein are caused by, contain, or result in the presence of VISA and/or VRSA.

Vancomycin-Resistant Enterococci

[00342] Enterococci are bacteria that are normally present in the human intestines and in the female genital tract and are often found in the environment. These bacteria sometimes cause infections. In some cases, enterococci have become resistant to vancomycin (also known as vancomycin-resistant enterococci or VRE.) Common forms of resistance to vancomycin occur in enterococcal strains that involve the acquisition of a set of genes encoding proteins that direct peptidoglycan precursors to incorporate D-Ala-D-Lac instead of D-Ala-D-Ala. The six different

types of vancomycin resistance shown by enterococcus are: Van-A, Van-B, Van-C, Van-D, Van-E and Van-F. In some cases, Van-A VRE is resistant to both vancomycin and teicoplanin, while in other cases, Van-B VRE is resistant to vancomycin but sensitive to teicoplanin; in further cases Van-C is partly resistant to vancomycin, and sensitive to teicoplanin.

5 **[00343]** In one aspect, is a method of treating a subject having a vancomycin-resistant enterococci comprising administering to the subject a compound of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the enterococci has developed resistance to vancomycin. In one embodiment, the subject has
10 been previously treated with vancomycin for a sustained period of time. In another embodiment, the subject has been hospitalized. In yet another embodiment, the subject has a weakened immune system such as patients in Intensive Care Units or in cancer or transplant wards. In a further embodiment, the subject has undergone surgical procedures such as, for example, abdominal or chest surgery. In yet a further embodiment, the subject has been colonized with
15 VRE. In one embodiment, the subject has a medical device such that an infection has developed. In another embodiment, the medical device is a urinary catheter or central intravenous (IV) catheter.

[00344] In another embodiment, is a method of treating a subject having a vancomycin-resistant enterococci comprising administering to the subject a compound of Formula (I), (I'),
20 (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the enterococcus has Van-A resistance.

[00345] In another embodiment, is a method of treating a subject having a vancomycin-resistant enterococci comprising administering to the subject a compound of Formula (I), (I'),
25 (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the enterococcus has Van-B resistance.

[00346] In another embodiment, is a method of treating a subject having a vancomycin-resistant enterococci comprising administering to the subject a compound of Formula (I), (I'),
30 (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe) or a pharmaceutically acceptable salt, ester, solvate, alkylated quaternary ammonium salt, stereoisomer, tautomer or prodrug thereof wherein the enterococcus has Van-C resistance.

Administration and Pharmaceutical Composition

[00347] Pharmaceutical compositions described herein comprise a therapeutically effective amount of a compound described herein (i.e., a compound of any of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe)) formulated together with one or more pharmaceutically acceptable carriers. As used herein, the term "pharmaceutically acceptable carrier" means a non-toxic, inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Some examples of materials which can serve as pharmaceutically acceptable carriers are sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator. The pharmaceutical compositions described herein can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, or as an oral or nasal spray, or a liquid aerosol or dry powder formulation for inhalation.

[00348] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms optionally contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[00349] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions are optionally formulated according to the known art using suitable dispersing or

wetting agents and suspending agents. The sterile injectable preparation is optionally a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that are optionally employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[00350] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00351] In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This is optionally accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is optionally accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are optionally prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

[00352] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compound described herein (i.e., a compound of any of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe)) with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[00353] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic

acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption
5 accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, acetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form optionally comprise buffering agents.

10 **[00354]** Solid compositions of a similar type are optionally employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[00355] The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings known in the
15 pharmaceutical formulating art. They optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

[00356] Solid compositions of a similar type are optionally employed as fillers in soft and hard-
20 filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[00357] The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling
25 coatings and other coatings known in the pharmaceutical formulating art. In such solid dosage forms the active compound is optionally admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms optionally comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such a magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the
30 dosage forms optionally comprise buffering agents. They optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

[00358] Dosage forms for topical or transdermal administration of a compound described herein include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as are optionally required.

5 Ophthalmic formulations, ear drops, and the like are also contemplated.

[00359] The ointments, pastes, creams and gels may contain, in addition to an active compound described herein, excipients such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

10 **[00360]** Compositions described herein are optionally formulated for delivery as a liquid aerosol or inhalable dry powder. Liquid aerosol formulations are optionally nebulized predominantly into particle sizes that can be delivered to the terminal and respiratory bronchioles where bacteria reside in patients with bronchial infections, such as chronic bronchitis and pneumonia. Pathogenic bacteria are commonly present throughout airways down
15 to bronchi, bronchioli and lung parenchyma, particularly in terminal and respiratory bronchioles. During exacerbation of infection, bacteria can also be present in alveoli. Liquid aerosol and inhalable dry powder formulations are preferably delivered throughout the endobronchial tree to the terminal bronchioles and eventually to the parenchymal tissue.

[00361] Aerosolized formulations described herein are optionally delivered using an aerosol
20 forming device, such as a jet, vibrating porous plate or ultrasonic nebulizer, preferably selected to allow the formation of a aerosol particles having with a mass medium average diameter predominantly between 1 to 5 μ . Further, the formulation preferably has balanced osmolarity ionic strength and chloride concentration, and the smallest aerosolizable volume able to deliver effective dose of the compounds described herein compound described herein (i.e., a compound
25 of any of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe)) to the site of the infection. Additionally, the aerosolized formulation preferably does not impair negatively the functionality of the airways and does not cause undesirable side effects.

[00362] Aerosolization devices suitable for administration of aerosol formulations described herein include, for example, jet, vibrating porous plate, ultrasonic nebulizers and energized dry
30 powder inhalers, that are able to nebulize the formulation into aerosol particle size predominantly in the size range from 1-5 μ . Predominantly in this application means that at least 70% but preferably more than 90% of all generated aerosol particles are within 1-5 μ range. A jet nebulizer works by air pressure to break a liquid solution into aerosol droplets. Vibrating porous plate nebulizers work by using a sonic vacuum produced by a rapidly vibrating porous

plate to extrude a solvent droplet through a porous plate. An ultrasonic nebulizer works by a piezoelectric crystal that shears a liquid into small aerosol droplets. A variety of suitable devices are available, including, for example, AeroNeb™ and AeroDose™ vibrating porous plate nebulizers (AeroGen, Inc., Sunnyvale, California), Sidestream® nebulizers (Medic-Aid Ltd.,
5 West Sussex, England), Pari LC® and Pari LC Star® jet nebulizers (Pari Respiratory Equipment, Inc., Richmond, Virginia), and Aerosonic™ (DeVilbiss Medizinische Produkte (Deutschland) GmbH, Heiden, Germany) and UltraAire® (Omron Healthcare, Inc., Vernon Hills, Illinois) ultrasonic nebulizers.

[00363] In some embodiments, compounds described herein compound described herein (i.e., a
10 compound of any of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe)) are formulated for use as topical powders and sprays that contain, in addition to the compounds described herein, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays optionally contain customary propellants such as chlorofluorohydrocarbons.

15 **[00364]** Transdermal patches have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

20 **[00365]** According to the methods of treatment described herein, bacterial infections are treated or prevented in a patient such as a human or lower mammal by administering to the patient a therapeutically effective amount of a compound described herein, in such amounts and for such time as is necessary to achieve the desired result. By a “therapeutically effective amount” of a compound described herein is meant a sufficient amount of the compound to treat bacterial
25 infections, at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions described herein will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the
30 activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors known in the medical arts.

[00366] The total daily dose of the compounds described herein compound described herein (i.e., a compound of any of Formula (I), (I'), (Ia), (Ib), (Ic), (Id), (II), (II'), (IIa), (IIb), (IIc), (IId), or (IIe)) administered to a human or other mammal in single or in divided doses can be in amounts, for example, from 0.01 to 50 mg/kg body weight or more usually from 0.1 to 25 mg/kg body weight. Single dose compositions may contain such amounts or submultiples thereof to make up the daily dose. In general, treatment regimens described herein comprise administration to a patient in need of such treatment from about 10 mg to about 2000 mg of the compound(s) described herein per day in single or multiple doses.

Examples

[00367] Compounds disclosed herein are made by the methods depicted in the reaction schemes shown below. Procedures are provided herein that, in combination with the knowledge of the synthetic organic chemist of ordinary skill in the art, are in some embodiments used to prepare the full range of compounds as disclosed and claimed herein.

[00368] The starting materials and reagents used in preparing these compounds are either available from commercial suppliers such as Aldrich Chemical Co., (Milwaukee, Wis.), Bachem (Torrance, Calif.), or Sigma (St. Louis, Mo.) or are prepared by methods known to those skilled in the art following procedures set forth in references such as Fieser and Fieser's Reagents for Organic Synthesis, Volumes 1-17 (John Wiley and Sons, 1991); Rodd's Chemistry of Carbon Compounds, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989); Organic Reactions, Volumes 1-40 (John Wiley and Sons, 1991), March's Advanced Organic Chemistry, (John Wiley and Sons, 4th Edition) and Larock's Comprehensive Organic Transformations (VCH Publishers Inc., 1989). These schemes are merely illustrative of some methods by which the compounds disclosed herein are in some embodiments synthesized, and various modifications to these schemes can be made and will be suggested to one skilled in the art having referred to this disclosure. The starting materials and the intermediates, and the final products of the reaction may be isolated and purified if desired using conventional techniques, including but not limited to filtration, distillation, crystallization, chromatography and the like. Such materials may be characterized using conventional means, including physical constants and spectral data. Compounds are typically isolated as formic acid salts by reverse phase HPLC using AcCN/H₂O with formic acid as an additive. In some instances, purifications are conducted without formic acid, and the compounds are isolated as the free base.

[00369] The methods of LCMS analysis are as follows:

[00370] LCMS (Method 5-95 AB, ESI): ESI, 5% AcCN/H₂O, 0.7 min; to 95% AcCN/H₂O, 0.4 min; 1.5 mL/min, Merck RP-18e, 2 x 25 mm.

[00371] LCMS (Method 10-80AB, 2 min, ESI): ESI, 10% AcCN/H₂O (0.04% TFA), 0.9 min to 80% AcCN/H₂O (0.04% TFA), then held for 0.6 min; 1.2 mL/min, Xtimate C18, 3 μm, 2.1 x 30 mm).

[00372] LCMS (Method 10-80AB, 7 min, ESI): ESI, 10% AcCN/H₂O (0.04% TFA), 6 min to 80% AcCN/H₂O (0.04% TFA), then held for 0.9 min; 0.8 mL/min, Xtimate C18, 3 μm, 2.1 x 30 mm).

Some abbreviations used herein are as follows:

DIPEA: diisopropylethylamine

DMAP: 4-dimethylaminopyridine

10 DMF: dimethylformamide

DCM: dichloromethane

TFA: trifluoroacetic acid

EDC: 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide

HATU: *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate

15 HCTU: *O*-(6-Chlorobenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate

HOBt: hydroxybenzotriazole

pyBOP: (Benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate

DMDO: 3,3-Dimethyldioxirane

DMP: Dess-Martin periodinane

20 THF: tetrahydrofuran

MeOH: methanol

EtOAc: ethyl acetate

Trt resin: 2-Chlorotrityl chloride resin

Rink amide resin: Rink amide (aminomethyl)polystyrene

25 Boc: t-butoxycarbonyl

CBz: benzyloxycarbonyl

Fmoc: [(9*H*-fluoren-9-yl)methoxy]carbonyl


Teoc: Trimethylsilylethoxycarbonyl

CDI: 1,1'-Carbonyldiimidazole

30 HFIP: 1,1,1,3,3,3-hexafluoropropan-2-ol

TLC: thin-layer chromatography

Example 1: Synthesis of (S)-methyl 2-amino-3-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propanoate



5 **[00373]** Step 1: To a solution of (S)-methyl 2-((tert-butoxycarbonyl)amino)-3-(4-hydroxyphenyl)propanoate (100 g, 0.323 mol) in acetone (2.0 L) was added K_2CO_3 (37 g, 0.34 mol). After the addition, MeI (32 mL, 0.97 mol) was added dropwise, and the reaction mixture was stirred at room temperature for 72 h and monitored by TLC. The reaction had not yet gone to completion, so NaOH (0.1 eq) was added to the reaction mixture. And after 2 h, the reaction

10 was completed. The solid was filtered and the solvent was removed. The residue was taken up in ethyl acetate and washed with H_2O , extracted with ethyl acetate (300 mL x 3). The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated to give (S)-methyl 2-((tert-butoxycarbonyl)amino)-3-(4-methoxyphenyl)propanoate (100 g, 95.4%).

[00374] Step 2: To a solution of (S)-methyl 2-((tert-butoxycarbonyl)amino)-3-(4-methoxyphenyl)propanoate (80 g, 40 g each x 2, run in two separate batches, 259 mmol overall) in methanol (1.5 L in each of the two flasks) was added sequentially Ag_2SO_4 (85 g, 272 mmol, 1/2-added to each flask) and I_2 (72 g, 283 mmol, 1/2-added to each flask). The reaction mixture was stirred at room temperature for 2 h. The reaction was monitored by LCMS. When all (S)-methyl 2-((tert-butoxycarbonyl)amino)-3-(4-methoxyphenyl)propanoate had been consumed,

20 then a solution of 10% (w/w) sodium thiosulfate was added until the reaction turned pale yellow. The solid was filtered and most of the methanol was evaporated by rotary evaporation. Water and ethyl acetate were added to each batch. The aqueous layer was extracted with ethyl acetate (3 x 200 mL). The combined organic layers were washed with brine, dried over sodium sulfate and concentrated. The crude material was combined for the two batches and they were


25 purified together by flash column chromatography on silica gel (25% then 35% then 40% ethyl


acetate in hexanes) to give (S)-methyl 2-((tert-butoxycarbonyl)amino)-3-(3-iodo-4-methoxyphenyl)propanoate (97 g, 89%).

[00375] Step 3: (S)-Methyl 2-((tert-butoxycarbonyl)amino)-3-(3-iodo-4-methoxyphenyl)propanoate (92 g, 46 g each run in two separate batches, 211 mmol) was dissolved in anhydrous DMSO (1.5 L, ½-added for each batch) under argon and to the solution was added bis(pinacolato) diboron (80.5 g, 317 mmol, ½-added for each batch) and KOAc (103 g, 1.05 mol, ½-added for each batch). This mixture was degassed with argon for twenty minutes, then Pd(dppf)Cl₂ (4.6 g, 6 mmol, ½-added for each batch) was added. The mixture was degassed with argon five times, then kept under argon and heated to 80°C for 3 h. TLC showed that the reaction was complete, and the reaction mixture was cooled to room temperature and filtered. The reaction mixture was dissolved in EA and washed with H₂O. The aqueous layer was extracted ethyl acetate (3 x 200 mL). The combined organic layers were dried over sodium sulfate, filtered and concentrated to give the crude product. The batches were then combined and purified together by flash column chromatography on silica gel (3% ethyl acetate in hexanes, then 20% to 25% ethyl acetate in hexanes to give (S)-methyl 2-((tert-butoxycarbonyl)amino)-3-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propanoate (70 g, 76%).

[00376] Step 4: (S)-Methyl 2-((tert-butoxycarbonyl)amino)-3-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propanoate (22 g, 50.6 mmol) was dissolved in dichloromethane (150 mL) and treated with trifluoroacetic acid (50 mL). The reaction mixture was stirred at room temperature and the reaction was monitored by HPLC. When all of the starting material had been consumed, the solvents were evaporated, DCM was added and Na₂CO₃ was added to neutralize the TFA. The mixture was filtered, and the solution was concentrated. DCM was added to the concentrated oil, and the mixture was cooled at 0°C for 1 hr, whereupon the solid precipitates that formed were filtered. The filtrate was concentrated to give (S)-methyl 2-amino-3-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propanoate. The material was used without further purification.

Example 2: Synthesis of (S)-2-((tert-butoxycarbonyl)amino)-2-(4-hydroxyphenyl)acetic acid





[00377] Step 1: To a stirred mixture of (S)-2-amino-2-(4-hydroxyphenyl)acetic acid (100 g, 0.6 mol, 1 eq) in a mixture of acetone (400 mL) and water (400 mL) was added di-*tert*-butyl dicarbonate (130.5 g, 0.6 mol, 1 eq) and NaHCO_3 (75.4 g, 0.9 mol, 1.5 eq). The mixture was allowed to stir at 25°C overnight. After HPLC showed the reaction was complete, the mixture was acidified with 5% citric acid (pH ~ 3). The mixture was filtered and the filter cake was washed with water, then dried to give (S)-2-((tert-butoxycarbonyl)amino)-2-(4-hydroxyphenyl)acetic acid (140 g, 87.5%). The crude product was used directly without further purification.


[00378] Step 2: To a solution of (S)-2-((tert-butoxycarbonyl)amino)-2-(4-hydroxyphenyl)acetic acid (45 g, 0.17 mol) in dry benzene (500 mL) was added paraformaldehyde (75.6 g, 0.84 mol, 5 eq) and *p*-toluenesulfonic acid (1.6 g, 8.5 mmol, 0.05 eq). A Dean-Stark apparatus with an attached condenser was then fit to the top of the flask and the mixture was heated at approximately 120°C until LC-MS showed the reaction was complete. The reaction was then cooled and the benzene was evaporated. The residue was taken up in ethyl acetate, washed with saturated NaHCO_3 (2 x 150 mL), then dried over sodium sulfate, and filtered. The solvent was removed to give (S)-*tert*-butyl 4-(4-hydroxyphenyl)-5-oxooxazolidine-3-carboxylate (36 g, 76.5%).

[00379] Step 3: (S)-*tert*-Butyl 4-(4-hydroxyphenyl)-5-oxooxazolidine-3-carboxylate (36 g, 0.13 mol, 1 eq) was dissolved in trifluoroacetic acid (75 mL) at 0°C then treated with triethylsilane (80 mL, 4 eq). The mixture was stirred at room temperature overnight. After LC-MS showed the reaction was complete, TFA was then evaporated to afford (S)-2-(4-hydroxyphenyl)-2-(methylamino)acetic acid, which was used without further purification.

[00380] Step 4: The resultant (S)-2-(4-hydroxyphenyl)-2-(methylamino)acetic acid was dissolved in water (85 mL), and to this solution was added solid NaHCO_3 until the pH reached 7. The solution was cooled to 0°C, then Na_2CO_3 was added until pH reached 9. A solution of di-*tert*-butyldicarbonate (28.3 g, 1.0 eq) in THF (75 mL) was added to the mixture. The mixture

was allowed to warm to room temperature then stirred overnight. After HPLC showed the reaction was complete, THF was then evaporated. The aqueous solution was extracted 2x with hexanes and then acidified with citric acid to pH ~3-4. The acidified solution was then extracted with ethyl acetate (200 mL x 3). The combined organic layers were washed with brine, dried over sodium sulfate, filtered and concentrated to give (S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(4-hydroxyphenyl)acetic acid (35 g, 97% via 2 steps).

Example 3: Synthesis of Compound 101-B



- 10 **[00381]** Step 1: To a solution of (S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(4-hydroxyphenyl)acetic acid (35 g, 0.12 mol) in DMF (300 mL) was added triethylamine (18.4 mL, 0.14 mol, 1.1 eq), HOBt (16.2 g, 0.12 mol, 1 eq), Ala-OMe HCl (19.5 g, 0.14 mol, 1.1 eq) and EDC (26.7 g, 0.14 mol, 1.1 eq) and the reaction was stirred overnight. After LC-MS

showed the reaction was complete, water and EtOAc were added. The aqueous layer was extracted with EtOAc (3×150 mL), and the combined organic layers were washed with 5% citric acid (pH – 3), saturated NaHCO₃ (aq), water and brine. The combined organic layers were then dried over sodium sulfate, filtered and concentrated to give (S)-methyl 2-((S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(4-hydroxyphenyl)acetamido)propanoate (30 g, 65.8%) as a white foam. The crude product was taken on to the next step directly without further purification.

[00382] Step 2: To a solution of (S)-methyl 2-((S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(4-hydroxyphenyl)acetamido)propanoate (30 g, 82 mmol) in acetone (400 mL) was added K₂CO₃ (56.6 g, 0.41 mol, 5 eq) and iodomethane (20.8 mL, 0.41 mol, 5 eq) and the reaction was stirred at reflux overnight. After LC-MS showed the reaction was complete, the reaction was then cooled to room temperature and the mixture was filtered. The filtrate was concentrated and the residue was taken up in water and ethyl acetate. The aqueous phase was extracted with EtOAc (3 x 150 mL). The combined organic layers were dried over sodium sulfate, filtered and concentrated to give (S)-methyl 2-((S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(4-methoxyphenyl)acetamido)propanoate (28 g, 90%), as a white foam.


[00383] Step 3: To a solution of (S)-methyl 2-((S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(4-methoxyphenyl)acetamido)propanoate (85 g, 0.22 mol, 1 eq) in methanol (1000 mL) was added sequentially Ag₂SO₄ (72.6 g, 0.23 mol, 1.05 eq) and I₂ (59.6 g, 1.05 eq). After LC-MS showed the reaction was complete, a solution of 10% (w/w) sodium thiosulfate was added until the reaction turned pale yellow. Most of the methanol was evaporated by rotary evaporation and then water and ethyl acetate were added. The aqueous layer was extracted with ethyl acetate (3 x 300 mL). The combined organic layers were washed with brine, dried over sodium sulfate and concentrated to give (S)-methyl 2-((S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(3-iodo-4-methoxyphenyl)acetamido)propanoate (100 g, 88.5%).

[00384] Step 4: To (S)-methyl 2-((S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(3-iodo-4-methoxyphenyl)acetamido)propanoate (25 g, 49.4 mmol, 1 eq) in THF (300 mL) was added 0.2 M LiOH (500 mL, 98.8 mmol, 2 eq). The solution was stirred until TLC showed all starting material had been consumed. 5% citric acid (pH – 3) was added to pH – 3 and then the THF was evaporated by rotary evaporation. The aqueous layer was extracted with EtOAc (3 x 100 mL). The combined organic layers were washed with brine, dried over sodium sulfate, filtered and concentrated to give (S)-2-((S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(3-iodo-4-methoxyphenyl)acetamido)propanoic acid (23 g, 94.6%), which was used directly without further purification.

[00385] Step 5: To a solution of (S)-methyl 2-amino-3-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propanoate (6.5 g, 19.4 mmol, 1 eq) and (S)-2-((S)-2-((tert-butoxycarbonyl)(methyl)amino)-2-(3-iodo-4-methoxyphenyl)acetamido)propanoic acid (10 g, 20.3 mmol, 1.05 eq) in acetonitrile:DMF (2.2:1, 168 mL) was added HOBt (6.5 g, 48.5 mmol, 2.5 eq) and EDC (8.1 g, 42.7 mmol, 2.2 eq). The reaction was stirred at room temperature overnight. After LC-MS showed the reaction was complete, diluted citric acid (pH- 3) was added and the aqueous was extracted with EtOAc (3 x 150 mL). The combined organic layers were then washed with saturated NaHCO₃ solution, brine and dried over sodium sulfate. The mixture was filtered and the filtrate was concentrated to give the crude product (6S,9S,12S)-methyl 6-(3-iodo-4-methoxyphenyl)-12-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzyl)-2,2,5,9-tetramethyl-4,7,10-trioxo-3-oxa-5,8,11-triazatridecan-13-oate, which was used directly without further purification.

[00386] Step 6: (6S,9S,12S)-Methyl 6-(3-iodo-4-methoxyphenyl)-12-(4-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzyl)-2,2,5,9-tetramethyl-4,7,10-trioxo-3-oxa-5,8,11-triazatridecan-13-oate (16 g, 19.4 mmol, 1 eq) and NaHCO₃ (16.3 g, 0.19 mol) were sealed in a flask with a condenser and put under an atmosphere of argon. DMF (600 mL) in a round bottle flask was purged several times via cycling with vacuum and Ar. PdCl₂(dppf) (3.3 g, 4.5 mmol) was then added to the DMF. The DMF solution was then degassed with Ar for 15 minutes. The solution of PdCl₂(dppf) dissolved in DMF was then transferred via syringe to the flask containing the substrate and NaHCO₃. The resulting mixture was submitted to several more cycles of vacuum and Ar then heated to 120°C overnight. After LCMS showed the reaction was completed, DMF was evaporated under vacuum. The crude material was subjected to abbreviated column chromatography (40% EA in PE) to remove most of the Pd species and then purified by prep HPLC to give Compound 101-A (2.1 g, 19.5% over two steps).

[00387] Step 7: To a stirred solution of Compound 101-A (2.1 g, 3.78 mmol) in DCM (25 mL) was added TFA (2 mL). The reaction was monitored via TLC and when starting material was consumed, the solvent was evaporated under vacuum. The residue was then dissolved in EtOAc and the organic layer was washed with saturated NaHCO₃ (10 mL), dried over sodium sulfate and concentrated to give Compound 101-B (1.7 g, 98.8%). MS (ESI) *m/z* 456.2 (M + H)⁺.

Example 4: Synthesis of Compound 101-G

[00388] Step 1: The removal of the methoxy protecting groups is described and is referred to as General Method 1. To a solution of Compound 101-B (5.0 g, 11.0 mmol) in EtSH (116 mL, 1.61 mol), AlBr₃ (165 mL, 165 mmol) was added slowly at 0°C under N₂. The mixture was stirred for 18 h. The volatiles were removed under reduced pressure and the residue was quenched by water (50 mL), which was further washed by DCM (20 mL x 3). The aqueous layer was purified by prep-HPLC (acetonitrile 1–20% / 0.1% TFA in water) to give Compound 101-C (4.5g, 99.2% yield) as a white solid.

[00389] Step 2: To a solution of Compound 101-C (4.7 g, 8.9 mmol) in 1,4-dioxane/H₂O (9:1, 165 mL) was added 1 N NaOH dropwise until pH~11. A solution of Cbz-OSu (6.66 g, 26.7 mmol) dissolved in 1,4-dioxane (50 mL) was then added. After stirring for 1 h, NaOH (1.07 g, 26.7 mmol) was then added to the reaction followed by MeOH (60 mL). This resulting mixture was allowed to stir for 20 mins. To the reaction was then added dilute citric acid (10% v/v, 50 mL), the aqueous layer was extracted with EtOAc (3 x 150 mL) and the combined organic layers were washed with brine (3 x 100 mL), dried over Na₂SO₄ and concentrated to give the crude product. The residue was diluted with DCM (50 mL) and the suspension was filtered to give desired compound (3.2 g). The DCM phase was concentrated and the residue was purified by silica gel column (eluting 10~20% methanol in EtOAc) to give the desired compound (1.0 g). The combined batches gave Compound 101-D (4.2 g, 86.1% yield) as a white solid.


[00390] Step 3: To Compound 101-D (4.3 g, 7.85 mmol) was added a solution of 1.25M HCl in MeOH (128 mL) and the reaction was stirred at 0°C. The volatiles were removed to afford Compound 101E (4.15 g, 94.1% yield) as a white solid, which was used directly in the next step.

[00391] Step 4: The bis-alkylation of phenol groups is described and is referred to as General Method 2. To a solution of Compound 101-E (3.9 g, 6.94 mmol) and K₂CO₃ (14.4 g, 104 mmol) in DMF (50 mL) was added tert-butyl 2-bromoethylcarbamate (15.6 g, 69.5 mmol) at 0°C. The mixture was stirred at room temperature for 48 h. The mixture was filtered and the filtrate was diluted with EtOAc (500 mL). The EtOAc layer was washed with brine (2 x 400 mL), dried over Na₂SO₄, concentrated and purified by chromatography on silica (solvent

gradient: 0-60% EtOAc in petroleum ether) to afford Compound 101-F (4.8 g, 81.5% yield) as a white solid.


[00392] Step 5: The hydrogenation of Cbz protecting groups is described and is referred to as General Method 3. To a solution of Compound 101-F (4.8 g, 5.7 mmol) in MeOH (100 mL), 10% Pd/C (1.26 g, 1.18 mmol) on carbon was added at room temperature. The reaction mixture was stirred for 1 h at the same temperature under hydrogen atmosphere (15 psi). The filtrate was then concentrated to afford Compound 101-G (4.0 g, 99% yield) as a white solid.

Example 5: Synthesis of Compounds 101-I, 101-J, 101-K, and 101-L




[00393] Step 1: The coupling of a Cbz-protected amino acid to an amine is described and is referred to as General Method 4. To a solution of Compound 101-G (3.5 g, 4.9 mmol) and (S)-2-(((benzyloxy)carbonyl)amino)-6-((tert-butoxycarbonyl)amino)hexanoic acid (2.4 g, 6.4 mmol) in DCM (30 mL) at 0 °C, HATU (3.7 g, 9.8 mmol) and DIPEA (1.9 g, 14.7 mmol) was added. The resulting mixture was allowed to gradually warm up to room temperature and stirred for 2 h. The reaction mixture was diluted with DCM (100 mL), which was washed with brine (100 mL x 3). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified by silica column chromatography to afford Compound 101-H (5.3 g, 99% yield) as a white solid.

[00394] Step 2: The hydrogenation step was performed using General Method 3 (Example 4) using Compound 101-H (1.5 g, 1.4 mmol) to afford Compound 101-I (1.2 g, 93% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 0.711$, $[M + H]^+ = 942.6$.




[00395] Compound 101-J was prepared from Compound 101-G and (S)-2-(((benzyloxy)carbonyl)amino)-5-((tert-butoxycarbonyl)amino)pentanoic acid using the conditions in Example 5. LCMS (Method 5-95 AB, ESI): $t_R = 0.841$, $[M + H]^+ = 928.4$.




5


[00396] Compound 101-K was prepared from Compound 101G and (S)-2-(((benzyloxy)carbonyl)amino)-4-((tert-butoxycarbonyl)amino)butanoic acid using the conditions in Example 5. LCMS (Method 5-95 AB, ESI): $t_R = 0.838$, $[M + H]^+ = 914.5$.



10 **[00397]** Compound 101-L was prepared from Compound 101G and (S)-2-(((benzyloxy)carbonyl)amino)-3-((tert-butoxycarbonyl)amino)propanoic acid using the conditions in Example 5. LCMS (Method 5-95 AB, ESI): $t_R = 0.833$, $[M + H]^+ = 900.5$.

Example 6: Synthesis of 3-((tert-butoxycarbonyl)(decyl)amino)propanoic acid







[00398] To a solution of methyl acrylate (2.2 g, 26 mmol) in THF (20 mL) was added a solution of decan-1-amine (6 g, 38 mmol) in THF (20 mL) at 0°C. The reaction mixture was stirred at 30°C for 48 h. The resulting solution was concentrated to obtain methyl 3-(decylamino)propanoate (6.4 g).

[00399] Step 2: To a solution of crude methyl 3-(decylamino)propanoate (6.4 g, 15 mmol) and Et_3N (4 g, 40 mmol) in DCM (30 mL) was added dropwise a solution of Boc_2O (5.7 g, 26 mmol) in DCM (20 mL) at 0°C. The reaction mixture was then allowed to warm to 30°C gradually and stirred for 18 h. After the reaction was completed, H_2O (50 mL) was added and the resulting aqueous layer was further extracted with DCM (50 mL*2). The combined organic layers were concentrated and the residue was purified by silica gel column (PE/EtOAc=50/1~20/1) to give methyl 3-((tert-butoxycarbonyl)(decyl)amino)propanoate (6.5 g, 73%) as a colorless oil.

[00400] Step 3: To a solution of methyl 3-((tert-butoxycarbonyl)(decyl)amino)propanoate (8.2 g, 23.9 mmol, crude) in EtOH (40 mL) was added a solution of LiOH (1.15 g, 48 mmol) in H_2O (20 mL) at 0°C. The reaction mixture was then allowed to warm to 30°C gradually and stirred for 18 h. After the reaction was complete, EtOH was removed under reduced pressure. The remaining aqueous solution was then adjusted to pH=2~3 with 6 N HCl , followed by the extraction with EtOAc (50 mL *3). The combined EtOAc layers were dried over Na_2SO_4 , and concentrated to give 3-((tert-butoxycarbonyl)(decyl)amino)propanoic acid (7 g, 88.6%) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 3.47-3.43 (t, $J=6.8\text{Hz}$, 2H), 3.19-3.15 (t, $J=7.2\text{Hz}$, 2H), 2.61 (brs, 2H), 1.51-1.39 (m, 11H), 1.24-1.22 (m, 14H), 0.88-0.84 (t, $J=6.8\text{Hz}$, 3H).

Example 7: Synthesis of Compound 101





[00401] Step 1: General Method 4 (Example 5) was applied to Compound 101-I (1.0 g, 1.27 mmol) and 3-((tert-butoxycarbonyl)(decyl)amino)propanoic acid (504 mg, 1.53 mmol) to afford
5 Compound 101-M (1.3 g, 82% yield) as a white solid.


[00402] Step 2: The lithium hydroxide hydrolysis of an ester to an acid is described and is referred to as General Method 5. To a solution of Compound 101-M (1.3 g, 1.04 mmol) in THF/H₂O (40 mL, 1:1) was added LiOH monohydrate (87 mg, 2.07 mmol) at 0°C. The mixture was allowed to gradually warm up to room temperature and stirred for 1 h. Most THF was
10 removed under reduced pressure and the resulting mixture was adjusted pH=5 with saturated citric acid, which was further extracted by EtOAc (30 mL x 3). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄ and concentrated to give Compound 101-N (1.1 g, 86% yield) as a white solid.

[00403] Steps 3 and 4: The coupling of an aminonitrile to a carboxylic acid and subsequent
15 Boc-deprotection is described and is referred to as General Method 6. To a solution of Compound 101-N (180 mg, 0.15 mmol), aminoacetonitrile hydrochloride (31 mg, 0.33 mmol) and DIPEA (38 mg, 0.29 mmol) in DCM/DMF (3 mL, 2:1) at 0°C, HATU (56 mg, 0.15 mmol) was added while stirring. The resulting mixture was stirred at room temperature for 1h. Most DCM was removed under reduced pressure and the residue was poured into water (10 mL),
20 which was extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, concentrated to the residue, which was purified by flash chromatography column to afford Compound 101-O (140 mg, 76%) as a white solid.

[00404] Compound 101-O (130 mg, 0.10 mmol) was added to 5% TFA in HFIP (6.5 mL) and the mixture was stirred at room temperature for 2 h. Volatiles were removed under reduced
25 pressure and the resulting crude was re-dissolved with DMF (5mL), which was neutralized with


solid NaHCO₃. The filtrate was then purified by HPLC to afford Compound 101 (54 mg, 60% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 0.710$, $M+H^+ = 877.6$; ¹HNMR (400 MHz, MeOH-*d*₄) δ 8.51 (brs, 2H, HCOOH), 7.28-7.25 (m, 2H), 7.20 (d, $J=8$ Hz, 1H), 7.18 (d, $J=8$ Hz, 1H), 6.90 (brs, 1H), 6.84 (brs, 1H), 6.37 (s, 1H), 4.82-4.79 (m, 3H), 4.28-4.20 (m, 4H), 4.21 (s, 2H), 3.33-3.26 (m, 2H), 3.26-3.16 (m, 5H), 3.16-3.12 (m, 1H), 3.11-2.95 (m, 2H), 2.95-2.91 (m, 2H), 2.90 (s, 3H), 2.73-2.66 (m, 2H), 1.75-1.65 (m, 6H), 1.64-1.51 (m, 1H), 1.50-1.16 (m, 18H), 0.92 (t, $J=6.8$ Hz, 3H).

Example 8: Synthesis of Compound 102




- 10 [00405] Compound 102 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.766$ min, $[M + H]^+ = 834.4$.


Example 9: Synthesis of Compound 103




- 15 [00406] Compound 103 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.747$ min, $[M + H]^+ = 820.4$.

Example 10: Synthesis of Compound 104


[00407] Compound 104 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.725$ min, $[M + H]^+ =$
 5 806.5.

Example 11: Synthesis of Compound 105


[00408] Compound 105 (formic acid salt) was prepared in 50% yield as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.709$ min,
 10 $[M + H]^+ = 792.3$.

Example 12: Synthesis of Compound 106


[00409] Compound 106 (formic acid salt) was prepared in 43% yield as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.681$ min,
 15 $[M + H]^+ = 778.3$.

Example 13: Synthesis of Compound 107


[00410] Compound 107 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.652$ min, $[M + H]^+ =$
 5 764.4.

Example 14: Synthesis of Compound 108


[00411] Compound 108 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 902$ $[M + H]^+$; ^1H NMR (500 MHz, DMSO- d_6) δ 9.21 (d, $J =$
 10 7.7 Hz, 1H), 9.00 (d, $J = 5.1$ Hz, 1H), 8.80 (dd, $J = 23.6, 7.8$ Hz, 1H), 8.39 (d, $J = 7.9$ Hz, 1H),
 8.01 (dd, $J = 15.0, 8.4$ Hz, 2H), 7.78 (d, $J = 8.4$ Hz, 2H), 7.68 (d, $J = 8.3$ Hz, 2H), 7.52 (d, $J =$
 8.4 Hz, 2H), 7.21 (d, $J = 8.4$ Hz, 1H), 7.15 (s, 2H), 7.04 (dd, $J = 21.3, 7.0$ Hz, 1H), 6.72 (s, 2H),
 6.35 (s, 1H), 4.84 (dd, $J = 13.5, 7.8$ Hz, 1H), 4.79 – 4.74 (m, 1H), 4.69 (dd, $J = 15.3, 8.2$ Hz,
 1H), 4.48 (d, $J = 6.8$ Hz, 1H), 4.17 (d, $J = 5.6$ Hz, 2H), 4.11 – 4.02 (m, 4H), 3.19 (d, $J = 16.0$
 15 Hz, 1H), 3.01 (d, $J = 17.3$ Hz, 1H), 2.98 – 2.87 (m, 5H), 2.80 (s, 3H), 2.74 (dd, $J = 18.2, 7.5$ Hz,
 2H), 1.94 – 1.72 (m, 2H), 1.68 – 1.55 (m, 2H), 1.53 – 1.37 (m, 2H), 1.33 (d, $J = 3.7$ Hz, 9H),
 1.20 (d, $J = 6.6$ Hz, 3H).

Example 15: Synthesis of Compound 109

[00412] Compound 109 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 906 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 8.97 (d, $J = 8.0$ Hz, 1H), 8.76 – 8.68 (m, 2H), 8.37 (d, $J = 9.0$ Hz, 1H), 8.16 (s, 1H), 8.03 – 7.97 (m, 2H), 7.82 – 7.76 (m, 2H), 7.33 – 7.27 (m, 2H), 7.25 – 7.19 (m, 1H), 7.16 (s, 1H), 7.07 (d, $J = 8.7$ Hz, 2H), 6.73 (d, $J = 2.1$ Hz, 2H), 6.37 (s, 1H), 4.86 (q, $J = 7.3$ Hz, 1H), 4.80 – 4.67 (m, 2H), 4.18 (d, $J = 5.9$ Hz, 2H), 4.10 (dq, $J = 12.3, 5.9, 5.1$ Hz, 4H), 3.84 (d, $J = 23.8$ Hz, 6H), 3.17 (s, 1H), 3.01 (dt, $J = 10.6, 5.6$ Hz, 4H), 2.80 (s, 3H), 2.77 (d, $J = 1.4$ Hz, 2H), 1.78 (d, $J = 7.8$ Hz, 2H), 1.65 – 1.53 (m, 2H), 1.51 – 1.38 (m, 2H), 1.21 (d, $J = 6.7$ Hz, 3H).

Example 16: Synthesis of Compound 110


[00413] Compound 110 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 872 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 8.95 (d, $J = 8.0$ Hz, 1H), 8.70 (t, $J = 6.5$ Hz, 2H), 8.35 (d, $J = 9.0$ Hz, 1H), 7.95 (d, $J = 8.2$ Hz, 2H), 7.75 – 7.70 (m, 2H), 7.68 – 7.62 (m, 2H), 7.45 – 7.38 (m, 3H), 7.36 – 7.30 (m, 2H), 7.25 – 7.21 (m, 1H), 7.16 (d, $J = 1.3$ Hz, 2H), 7.08 (d, $J = 8.5$ Hz, 1H), 6.72 (s, 2H), 6.37 (s, 1H), 4.84 (d, $J = 7.2$ Hz, 1H), 4.72 (dd, $J = 17.1, 9.2$ Hz, 2H), 4.19 (d, $J = 5.9$ Hz, 2H), 4.17 – 4.06 (m, 4H), 3.18 (s, 1H), 3.05 (dt, $J = 10.9, 5.9$ Hz, 5H), 2.80 (s, 3H), 2.77 (s, 2H), 1.77 (d, $J = 7.7$ Hz, 2H), 1.59 (d, $J = 7.3$ Hz, 2H), 1.48 (s, 2H), 1.21 (d, $J = 6.7$ Hz, 3H).

Example 17: Synthesis of Compound 111

[00414] Compound 111 (formic acid salt) was prepared utilizing the methods in Example 7

(Compound 101). LC-MS: $m/z = 914 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 8.96 (d, $J =$


5 8.0 Hz, 1H), 8.83 (d, $J = 7.8$ Hz, 1H), 8.70 (t, $J = 5.6$ Hz, 1H), 8.37 (d, $J = 9.0$ Hz, 1H), 8.07 – 8.01 (m, 2H), 7.94 – 7.88 (m, 2H), 7.83 (dd, $J = 5.6, 1.9$ Hz, 2H), 7.68 (t, $J = 1.8$ Hz, 1H), 7.22 (d, $J = 8.8$ Hz, 1H), 7.15 (d, $J = 1.3$ Hz, 2H), 7.07 (d, $J = 8.5$ Hz, 1H), 6.73 (d, $J = 1.9$ Hz, 2H), 6.37 (s, 1H), 4.86 (d, $J = 7.3$ Hz, 1H), 4.72 (dd, $J = 16.4, 9.0$ Hz, 2H), 4.18 (d, $J = 5.9$ Hz, 2H), 4.15 – 4.06 (m, 4H), 3.17 (s, 1H), 3.01 (q, $J = 5.3$ Hz, 5H), 2.80 (s, 3H), 2.79 (d, $J = 2.2$ Hz, 10 2H), 1.83 – 1.74 (m, 2H), 1.64 – 1.55 (m, 2H), 1.46 (s, 2H), 1.20 (d, $J = 6.7$ Hz, 3H).

Example 18: Synthesis of Compound 112


[00415] Compound 112 (formic acid salt) was prepared utilizing the methods in Example 7

(Compound 101). LC-MS: $m/z = 914 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 8.97 (d, $J =$


15 8.0 Hz, 1H), 8.82 (d, $J = 7.8$ Hz, 1H), 8.70 (t, $J = 5.5$ Hz, 1H), 8.38 (d, $J = 9.0$ Hz, 1H), 8.10 – 8.05 (m, 2H), 7.98 (d, $J = 8.2$ Hz, 2H), 7.88 (dd, $J = 8.4, 5.5$ Hz, 4H), 7.22 (dd, $J = 8.5, 2.3$ Hz, 1H), 7.15 (d, $J = 1.4$ Hz, 2H), 7.07 (d, $J = 8.5$ Hz, 1H), 6.73 (s, 2H), 6.37 (s, 1H), 4.90 – 4.82 (m, 1H), 4.80 – 4.66 (m, 2H), 4.18 (d, $J = 5.8$ Hz, 2H), 4.12 – 4.05 (m, 4H), 3.17 (s, 1H), 2.99 (dt, $J = 9.7, 5.3$ Hz, 5H), 2.81 (s, 3H), 2.81 – 2.77 (m, 2H), 1.79 (q, $J = 7.5$ Hz, 2H), 1.65 – 1.54 20 (m, 2H), 1.51 – 1.38 (m, 2H), 1.20 (d, $J = 6.7$ Hz, 3H).

Example 19: Synthesis of Compound 113


[00416] Compound 113 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 880 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 8.98 (d, $J = 7.7$ Hz, 1H), 8.92 (d, $J = 7.9$ Hz, 1H), 8.73 (t, $J = 5.8$ Hz, 1H), 8.39 (d, $J = 9.0$ Hz, 1H), 8.21 (t, $J = 1.9$ Hz, 1H), 7.93 – 7.80 (m, 4H), 7.64 – 7.57 (m, 3H), 7.19 (ddd, $J = 15.7, 8.7, 2.5$ Hz, 2H), 7.04 (dd, $J = 12.0, 8.5$ Hz, 2H), 6.72 (t, $J = 2.3$ Hz, 2H), 6.36 (s, 1H), 4.93 – 4.85 (m, 1H), 4.79 – 4.68 (m, 2H), 4.18 (d, $J = 5.7$ Hz, 2H), 4.02 (dd, $J = 11.1, 5.5$ Hz, 4H), 3.19 (d, $J = 18.3$ Hz, 1H), 3.01 – 2.88 (m, 5H), 2.78 (s, 5H), 1.77 (d, $J = 16.9$ Hz, 2H), 1.59 (q, $J = 7.4$ Hz, 2H), 1.46 (s, 2H), 1.20 (d, $J = 6.8$ Hz, 3H).

Example 20: Synthesis of Compound 114


[00417] Compound 114 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 880 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 8.97 (d, $J = 8.0$ Hz, 1H), 8.79 (d, $J = 7.5$ Hz, 1H), 8.71 (t, $J = 5.7$ Hz, 1H), 8.36 (d, $J = 9.0$ Hz, 1H), 7.56 (ddd, $J = 7.6, 6.2, 2.7$ Hz, 1H), 7.50 – 7.42 (m, 7H), 7.23 (dd, $J = 8.6, 2.3$ Hz, 1H), 7.10 – 7.02 (m, 2H), 6.99 (dd, $J = 8.7, 2.4$ Hz, 1H), 6.76 – 6.69 (m, 2H), 6.47 (s, 1H), 4.81 – 4.68 (m, 2H), 4.61 – 4.54 (m, 1H), 4.19 (d, $J = 5.9$ Hz, 2H), 4.17 – 4.08 (m, 4H), 3.19 (s, 1H), 3.12 – 3.00 (m, 5H), 2.78 (d, $J = 8.5$ Hz, 2H), 2.74 (s, 3H), 1.67 – 1.49 (m, 4H), 1.38 – 1.28 (m, 2H), 1.21 (d, $J = 6.7$ Hz, 3H).

Example 21: Synthesis of Compound 115


[00418] Compound 115 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 880 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 8.96 (d, $J = 8.0$ Hz, 1H), 8.81 (d, $J = 7.9$ Hz, 1H), 8.70 (t, $J = 5.6$ Hz, 1H), 8.35 (d, $J = 8.9$ Hz, 1H), 8.04 – 7.98 (m, 2H), 7.64 – 7.59 (m, 1H), 7.59 – 7.54 (m, 2H), 7.49 – 7.42 (m, 3H), 7.26 – 7.20 (m, 1H), 7.16 (d, $J = 1.3$ Hz, 2H), 7.08 (d, $J = 8.6$ Hz, 1H), 6.73 (d, $J = 3.3$ Hz, 2H), 6.39 (s, 1H), 4.90 – 4.83 (m, 1H), 4.80 – 4.67 (m, 2H), 4.19 (d, $J = 6.0$ Hz, 2H), 4.13 (dd, $J = 10.5, 5.2$ Hz, 4H), 3.18 (s, 1H), 3.06 (dq, $J = 9.7, 5.4$ Hz, 5H), 2.82 (s, 3H), 2.80 (s, 2H), 1.78 (d, $J = 7.6$ Hz, 2H), 1.59 (d, $J = 7.1$ Hz, 2H), 1.50 (s, 2H), 1.21 (d, $J = 6.8$ Hz, 3H).

Example 22: Synthesis of Compound 116


[00419] Compound 116 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 896 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 8.99 (d, $J = 7.9$ Hz, 1H), 8.74 (t, $J = 5.6$ Hz, 1H), 8.66 (d, $J = 7.7$ Hz, 1H), 8.36 (d, $J = 8.9$ Hz, 1H), 8.01 – 7.95 (m, 2H), 7.53 – 7.46 (m, 2H), 7.25 – 7.19 (m, 1H), 7.16 – 7.04 (m, 7H), 6.76 – 6.69 (m, 2H), 6.35 (s, 1H), 4.83 (t, $J = 7.2$ Hz, 1H), 4.79 – 4.67 (m, 2H), 4.18 (d, $J = 5.9$ Hz, 2H), 4.09 (dd, $J = 10.9, 5.6$ Hz, 4H), 3.19 (d, $J = 16.6$ Hz, 1H), 2.98 (dt, $J = 11.0, 5.4$ Hz, 5H), 2.79 (s, 3H), 2.78 (s, 2H), 1.75 (q, $J = 7.6$ Hz, 2H), 1.58 (dq, $J = 15.2, 7.4$ Hz, 2H), 1.44 (dq, $J = 15.0, 7.3$ Hz, 2H), 1.20 (d, $J = 6.7$ Hz, 3H).

Example 23: Synthesis of Compound 117




[00420] Compound 117 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 894 [M + H]^+$.

5 Example 24: Synthesis of Compound 118

[00421] Compound 118 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 922 [M + H]^+$.


Example 25: Synthesis of Compound 119

[00422] Compound 119 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 796 [M + H]^+$.




= 7.6, 4.4 Hz, 2H), 8.70 (t, $J = 5.6$ Hz, 1H), 8.42 (d, $J = 8.8$ Hz, 1H), 8.05 (d, $J = 8.5$ Hz, 2H), 7.91 – 7.79 (m, 5H), 7.77 – 7.73 (m, 2H), 7.54 – 7.49 (m, 2H), 7.44 – 7.38 (m, 2H), 7.22 (dt, $J = 8.6, 2.4$ Hz, 1H), 7.16 (d, $J = 8.6$ Hz, 2H), 7.07 (d, $J = 8.5$ Hz, 1H), 6.73 (t, $J = 2.3$ Hz, 2H), 6.33 (s, 1H), 5.02 (q, $J = 7.3$ Hz, 1H), 4.79 – 4.70 (m, 2H), 4.18 (d, $J = 5.9$ Hz, 2H), 4.13 – 4.05 (m, 4H), 3.20 (d, $J = 18.2$ Hz, 1H), 3.06 – 2.92 (m, 9H), 2.77 (s, 3H), 2.07 (dd, $J = 20.0, 12.5$ Hz, 2H), 1.21 (d, $J = 6.7$ Hz, 3H).

Example 29: Synthesis of Compound 123




[00426] Compound 123 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 804$ $[M + H]^+$.


Example 30: Synthesis of Compound 124




[00427] Compound 124 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 822$ $[M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 9.28 (d, $J = 7.5$ Hz, 1H), 8.96 (d, $J = 7.9$ Hz, 1H), 8.70 (t, $J = 5.7$ Hz, 1H), 8.39 (d, $J = 9.0$ Hz, 1H), 7.51 (d, $J = 8.2$ Hz, 2H), 7.34 – 7.29 (m, 2H), 7.22 (dd, $J = 8.7, 2.3$ Hz, 1H), 7.18 – 7.12 (m, 2H), 7.08 (d, $J = 8.5$ Hz, 1H), 6.72 (d, $J = 2.4$ Hz, 2H), 6.34 (s, 1H), 4.83 (s, 1H), 4.73 (td, $J = 15.9, 14.1, 9.1$ Hz, 2H), 4.18 (d, $J = 5.8$ Hz, 2H), 4.16 – 4.06 (m, 4H), 3.19 (d, $J = 16.2$ Hz, 1H), 3.01 (dt, $J = 10.3, 5.4$ Hz, 5H), 2.88 (dd, $J = 10.0, 6.0$ Hz, 2H), 2.77 (s, 3H), 2.67 – 2.59 (m, 2H), 2.05 – 1.88 (m, 2H), 1.61 – 1.53 (m, 2H), 1.35 – 1.26 (m, 2H), 1.20 (d, $J = 6.7$ Hz, 3H), 0.90 (t, $J = 7.3$ Hz, 3H).

Example 31: Synthesis of Compound 125

[00428] Compound 125 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 792 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 9.08 (d, $J = 7.8$ Hz, 1H), 8.98 (d, $J = 7.8$ Hz, 1H), 8.71 (t, $J = 5.6$ Hz, 1H), 8.42 (d, $J = 9.0$ Hz, 1H), 8.08 – 7.95 (m, 4H), 7.64 (ddt, $J = 10.3, 5.2, 3.7$ Hz, 3H), 7.23 (t, $J = 9.0$ Hz, 1H), 7.15 (s, 2H), 7.06 (d, $J = 8.8$ Hz, 1H), 6.73 (d, $J = 4.0$ Hz, 2H), 6.33 (d, $J = 0.9$ Hz, 1H), 5.04 (t, $J = 7.1$ Hz, 1H), 4.80 – 4.70 (m, 2H), 4.18 (d, $J = 5.9$ Hz, 2H), 4.08 (s, 4H), 3.19 (d, $J = 16.4$ Hz, 1H), 2.98 (d, $J = 16.3$ Hz, 5H), 2.78 (s, 3H), 2.68 – 2.64 (m, 2H), 2.17 – 2.01 (m, 2H), 1.21 (d, $J = 6.7$ Hz, 3H).


Example 32: Synthesis of Compound 126

[00429] Compound 126 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 790 [M + H]^+$.

Example 33: Synthesis of Compound 127

[00430] Compound 127 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from Compound 101-J and 4-butylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.545$, $[M + H]^+ = 812.2$.


Example 34: Synthesis of Compound 128



5

[00431] Compound 128 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 0-60AB): $t_R = 0.850$ min/ 2 min, $[M+H]^+ = 770.5$.


Example 35: Synthesis of Compound 129



10


[00432] Compound 129 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from 3-((tert-butoxycarbonyl)(decyl)amino)propanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.557$, $[M + H]^+ = 849.5$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.49 (brs, 2H, HCOOH), 7.29-7.24 (m, 2H), 7.17 (d, $J=8$ Hz, 1H), 7.10 (d, $J=8$ Hz, 1H), 6.88 (brs, 1H), 6.81 (brs, 1H), 6.29 (s, 1H), 4.91-4.76 (m, 3H), 4.27-4.15 (m, 4H), 4.19 (s, 2H), 3.36-3.23 (m, 6H), 3.17-3.01 (m, 6H), 2.86 (s, 3H), 2.76-2.71 (m, 2H), 2.20-2.10 (m, 1H), 2.05-1.95 (m, 1H), 1.71 (brs, 2H), 1.38-1.31 (m, 17H), 0.90 (t, $J=6.4$ Hz, 3H).

15




Example 36: Synthesis of Compound 130**[00433]**


[00434] Compound 130 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.699$, $[M + H]^+ = 835.6$;

5 ^1H NMR (400 MHz, MeOH- d_4) δ 8.45 (brs, 1H), 7.34-7.20 (m, 2H), 7.16 (d, $J=8.5$ Hz, 1H), 7.10 (d, $J=8.5$ Hz, 1H), 6.88 (brs, 1H), 6.80 (brs, 1H), 6.22 (s, 1H), 5.21-5.13 (m, 1H), 4.86-4.79 (m, 2H), 4.30-4.17 (m, 4H), 4.19 (s, 2H), 3.41-3.32 (m, 3H), 3.29-3.22 (m, 5H), 3.22-3.09 (m, 2H), 3.06-2.99 (m, 2H), 2.80 (s, 3H), 1.77-1.65 (m, 2H), 1.46 - 1.23 (m, 19H), 0.90 (t, $J=6.8$ Hz, 3H).

10 **Example 37: Synthesis of Compound 131**

[00435] Compound 131 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from 3-((tert-butoxycarbonyl)(decyl)amino)propanoic acid and Compound 101-G. LCMS (Method 5-95 AB, ESI): $t_R = 0.585$, $[M + H]^+ = 749.3$; ^1H NMR (400
15 MHz, MeOH- d_4) δ 8.47 (brs, 2H, HCOOH), 7.29-7.23 (m, 2H), 7.17 (d, $J=8.0$ Hz, 1H), 7.09 (d, $J=8.0$ Hz, 1H), 6.87 (brs, 1H), 6.80 (brs, 1H), 6.31 (s, 1H), 4.81-4.76 (m, 2H), 4.26-4.18 (m, 4H), 4.21 (s, 2H), 3.34-3.28 (m, 2H), 3.25-3.15 (m, 4H), 3.15-3.10 (m, 2H), 3.07-3.03 (m, 2H), 2.94-2.90 (m, 2H), 2.78 (s, 3H), 1.74-1.70 (m, 2H), 1.39-1.32 (m, 17H), 0.90 (t, $J=6.4$ Hz, 3H).






[00439] To a solution of nonanal (600 mg, 4.22 mmol) in DCM (25 mL) at 0°C was added methyl 4-aminobutanoate (988 mg, 8.44 mmol) and HOAc (1 mL), followed by the addition of NaBH₃CN (398 mg, 2 mmol). The mixture was stirred at 15°C for 12h. After the reaction was complete, H₂O (20 mL) was added and the aqueous layer was extracted by DCM (30 mL*2). The combined organic layers were concentrated to obtain methyl 4-(nonylamino) butanoate. LCMS (Method 5-95 AB, ESI): $t_R = 0.770$, $[M + H]^+ = 244.0$


[00440] The N-Boc formation and LiOH ester hydrolysis was performed in a manner similar to Example 6 to afford 0.46 g of 4-((tert-butoxycarbonyl)(nonyl)amino)butanoic acid. ELSD-LC/MS 352.3 $[M + Na]^+$.

[00441] Compound 133 (formic acid salt) was prepared as a white solid utilizing the methods in Example 37 (Compound 131). LCMS (Method 5-95 AB, ESI): $t_R = 0.717$, $[M + H]^+ = 749.5$; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.48 (brs, 3H, HCOOH), 7.29-7.20 (m, 2H), 7.16 (d, $J=8.4$ Hz, 1H), 7.09 (d, $J=8.4$ Hz, 1H), 6.91(brs, 1H), 6.77 (brs, 1H), 6.30 (s, 1H), 4.82-4.74 (m, 2H), 4.29-4.12 (m, 4H), 4.19 (s, 2H), 3.38-3.33 (m, 1H), 3.24-3.05 (m, 7H), 3.04-2.95 (m, 3H), 2.78 (s, 3H), 2.73-2.57 (m, 3H), 2.07-1.94 (m, 2H), 1.74-1.63 (m, 2H), 1.50-1.23 (m, 15H), 0.90 (t, $J=6.6$ Hz, 3H).


Example 40: Synthesis of Compound 134



[00442] Compound 134 (formic acid salt) was prepared as a white solid utilizing the methods in Example 39 (Compound 133). LCMS (Method 5-95 AB, ESI): $t_R = 0.562$, $M + H^+ = 749.3$.

Example 41: Synthesis of Compound 135

- 5 **[00443]** Step 1: A solution of 1-bromo-4-*n*-butylbenzene (100 g, 0.472 mol), 4-(methoxycarbonyl)phenylboronic acid (82.0 g, 0.456 mol), 2 M Na₂CO₃ (150 g, 1.42 mol) in toluene/EtOH (900 mL/300 mL) was degassed with N₂ three times, then Pd(PPh₃)₄ (27.2 g, 23.6 mmol) was added. The resulting mixture was degassed with N₂ three times and then heated to reflux for 5h. After TLC showed the reaction was complete, toluene and EtOH was removed
- 10 under vacuum. The residue was extracted with EA (3x). The combined organic layers were washed with brine and dried with Na₂SO₄. The solvent was removed to give the crude product. The crude product was purified by column chromatography on silica gel eluted with PE: EA (150:1). The solvent was removed to give methyl 4'-butyl-[1,1'-biphenyl]-4-carboxylate (105 g, 86.0%) as a white solid.
- 15 **[00444]** Step 2: A mixture of methyl 4'-butyl-[1,1'-biphenyl]-4-carboxylate (89.0 g, 0.332 mol), NaOH (26.6 g, 0.664 mol) in THF/H₂O (500 mL/100 mL) was heated to reflux overnight. After TLC showed the reaction was complete, THF was removed. The residue was adjusted pH=3~4 with 2 N HCl solution. The resulting mixture was filtered and the cake was washed with water, and dried to give 4'-butyl-[1,1'-biphenyl]-4-carboxylic acid (60.0 g, 71.1%) as a white solid.
- 20 **[00445]** Compound 135 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from Compound 101-J and 4'-butyl-[1,1'-biphenyl]-4-carboxylic acid. LCMS (Method 5-95 AB, ESI): *t*_R = 0.757, [M + H]⁺ = 888.4; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.47 (brs, 1H, HCOOH), 7.91 (d, *J*=8 Hz, 2H), 7.71 (d, *J*=8.0 Hz, 2H), 7.53 (d, *J*=8 Hz, 2H), 7.36-7.20 (m, 4H), 7.25-7.18 (m, 2H), 6.90 (brs, 1H), 6.81 (brs, 1H), 6.50 (s, 1H),
- 25 5.10-5.05 (m, 1H), 4.85-4.72 (m, 2H), 4.32-4.20 (m, 4H), 4.21 (s, 2H), 3.35-3.10 (m, 6H), 3.08-2.95 (m, 2H), 2.92 (s, 3H), 2.70-2.60 (m, 2H), 2.05-1.75 (m, 4H), 1.70-1.55 (m, 2H), 1.45-1.30 (m, 2H), 1.37 (d, *J*=7.2 Hz, 3H), 0.95 (t, *J*=6.8 Hz, 3H).

Example 42: Synthesis of Compound 136


[00446] Step 1: To a stirred solution of (R)-2-((tert-butoxycarbonyl)amino)propanoic acid (10.0 g, 52.8 mmol), NMM (5.88 g, 58.1 mmol) in THF (50 mL) was added ethyl chloroformate (8.86 g, 81.6 mmol) dropwise at -10°C and the mixture was stirred at the same temperature for 20 mins, followed by the addition of 10N ammonia in THF (50 mL, 500 mmol) slowly. The resulting mixture was stirred at -10°C for another 2 h. The volatiles were removed and the residue was taken up with EtOAc (100 mL), which was washed with 1N KHSO₄ solution and brine (50 mL each). The organic layer was dried over Na₂SO₄ and concentrated to give (R)-tert-butyl (1-amino-1-oxopropan-2-yl)carbamate (6.5 g, 65.3% yield) as a white solid.

[00447] Step 2: To a solution of (R)-tert-butyl (1-amino-1-oxopropan-2-yl)carbamate (3.5 g, 18.6 mmol) and pyridine (4.4 g, 55.8 mmol) in anhydrous THF (50 mL) was added TFAA (5.9 g, 27.9 mmol) dropwise at -10°C and the mixture was stirred at the same temperature for 2 h. The volatiles were removed and the residue was taken up with EtOAc (100 mL), which was washed with 1N KHSO₄ solution and brine (50 mL each). The organic layer was dried over Na₂SO₄ and concentrated to crude product, which was further re-crystallized using 100 mL (20% EtOAc in petroleum ether) to afford (R)-tert-butyl (1-cyanoethyl)carbamate (1.5 g, 47.4% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 4.82 (brs, 1H), 4.61 (brs, 1H), 1.53 (d, J=7.6 Hz, 3H), 1.45 (s, 9H).

[00448] Step 3: (R)-Tert-butyl (1-cyanoethyl)carbamate (100 mg, 0.59 mmol) was added in portions into HCOOH (1.0 mL) at 0°C and the mixture was allowed to warm to 20°C while stirring and stirred at the same temperature for 3 h. The volatiles were removed to afford (R)-2-aminopropanenitrile (65 mg, 95.3% yield) as a white solid, which was used directly in the next step.


[00449] Compound 136 (formic acid salt) was prepared utilizing the methods in Example 41 (Compound 135) except (R)-2-aminopropanenitrile hydrochloride is used in the coupling step. LC-MS: $m/z = 880 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 8.91 (d, $J = 7.9$ Hz, 1H), 8.82 (d, $J = 7.5$ Hz, 1H), 8.74 (d, $J = 7.9$ Hz, 1H), 8.37 (d, $J = 8.9$ Hz, 1H), 8.01 (d, $J = 8.4$ Hz, 2H), 7.80 – 7.75 (m, 2H), 7.69 – 7.63 (m, 2H), 7.33 (d, $J = 8.1$ Hz, 2H), 7.19 (d, $J = 8.6$ Hz, 1H), 7.15 (s, 2H), 7.09 – 7.02 (m, 1H), 6.74 (s, 2H), 6.37 (s, 1H), 4.84 (dd, $J = 16.0, 7.4$ Hz, 2H), 4.72 (d, $J = 12.6$ Hz, 2H), 4.05 (s, 4H), 2.94 (s, 5H), 2.80 (s, 3H), 2.78 (d, $J = 7.9$ Hz, 2H), 2.65 – 2.62 (m, 2H), 1.78 (s, 2H), 1.60 (t, $J = 7.6$ Hz, 4H), 1.44 (d, $J = 7.2$ Hz, 5H), 1.38 – 1.31 (m, 2H), 1.20 (d, $J = 6.7$ Hz, 3H), 0.96 – 0.88 (m, 3H).

10 Example 43: Synthesis of Compound 137



[00450] Compound 137 (formic acid salt) was prepared utilizing the methods in Example 41 (Compound 135). LC-MS: $m/z = 903 [M + H]^+$; ^1H NMR (400 MHz, DMSO- d_6) δ 9.00 (d, $J = 7.8$ Hz, 1H), 8.89 (d, $J = 8.0$ Hz, 1H), 8.74 (t, $J = 5.7$ Hz, 1H), 8.39 (d, $J = 9.0$ Hz, 1H), 8.20 (t, $J = 1.8$ Hz, 1H), 7.90 – 7.81 (m, 2H), 7.72 – 7.66 (m, 2H), 7.57 (t, $J = 7.8$ Hz, 1H), 7.35 (d, $J = 8.2$ Hz, 2H), 7.18 (ddd, $J = 16.6, 8.6, 2.5$ Hz, 2H), 7.03 (t, $J = 8.7$ Hz, 2H), 6.72 (d, $J = 2.6$ Hz, 2H), 6.35 (s, 1H), 4.88 (q, $J = 7.5, 7.0$ Hz, 1H), 4.79 – 4.68 (m, 2H), 4.18 (d, $J = 5.8$ Hz, 2H), 3.99 (dt, $J = 10.6, 5.5$ Hz, 4H), 3.18 (d, $J = 17.8$ Hz, 1H), 3.02 (d, $J = 12.1$ Hz, 1H), 2.89 (q, $J = 5.6, 4.3$ Hz, 4H), 2.78 (s, 3H), 2.76 (s, 2H), 2.68 – 2.62 (m, 2H), 1.77 (d, $J = 13.1$ Hz, 2H), 1.65 – 1.55 (m, 4H), 1.45 (s, 4H), 1.40 – 1.29 (m, 2H), 1.20 (d, $J = 6.7$ Hz, 3H), 0.92 (t, $J = 7.3$ Hz, 3H).

Example 44: Synthesis of Compound 138



5


NCCNC(=O)[C@H](N)C(=O)N[C@@H](Cc1ccc2c(c1)Oc3ccc(cc3)OCCN)C(=O)N[C@@H](C)C(=O)N[C@@H](C)C(=O)N[C@@H](Cc4ccc(cc4)C(=O)NCC5=CC=CC=C5)C(=O)NCC#N

10

NCCNC(=O)[C@H](N)C(=O)N(C)[C@@H](Cc1ccc(OCCN)cc1)c2c(Cl)ccc(cc2C3=CC=C4CCCCC4)C(=O)N[C@@H](CN)C(=O)N[C@@H](C)C(=O)NCC#N

140

15


Example 47: Synthesis of Compound 141

[00454] The synthesis of a biaryl tail via Pd-catalyzed Suzuki coupling of an aryl boronic acid and an aryl halide, followed by base hydrolysis of the ester is described and is referred to as General Method 7.


[00455] Step 1: To a solution of 4-*t*-butylphenylboronic acid (151.6 mg, 0.85 mmol) in 1,4-dioxane (5 mL) and water (1 mL) were added potassium carbonate (181.0 mg, 1.31 mmol), 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (47.9 mg, 0.07 mmol), and methyl 4-bromo-2-methylbenzoate (150.0 mg, 0.65 mmol). The reaction mixture was stirred at 100 °C for 2 h under N₂ and concentrated. The residue was taken up in EtOAc (20 mL) and washed with water (20 mL x 2) and brine (20 mL). The organic layer was dried over MgSO₄ and concentrated. The residue was purified by flash column chromatography (5% EtOAc in petroleum ether, R_f = 0.7) to afford methyl 4-(4-*tert*-butylphenyl)-2-methylbenzoate (120 mg, 64.9% yield) as a colorless oil. LCMS (5-95AB_1.5min): t_R = 0.972 min, [M + H]⁺ = 281.9.

[00456] Step 2: Methyl 4-(4-*tert*-butylphenyl)-2-methylbenzoate (120.0 mg, 0.430 mmol) was hydrolyzed using General Method NaOH to give 4-(4-*tert*-butylphenyl)-2-methylbenzoic acid (100 mg, 0.3726 mmol, 87.7% yield) as a white solid.

[00457] Compound 141 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): t_R = 0.633 min, [M + H]⁺ = 902.4.


Example 48: Synthesis of Compound 142

[00458] Compound 142 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from 4-(4-butylphenyl)-2-methylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.643$ min, $[M + H]^+ = 902.4$.

Example 49: Synthesis of Compound 143


[00459] Compound 143 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from 4-(4-trifluoromethylphenyl)-2-methylbenzoic acid.

LCMS (Method 5-95 AB, ESI): $t_R = 0.588$ min, $[M + H]^+ = 900.3$.


Example 50: Synthesis of Compound 144

[00460] Compound 144 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from 3-methyl-[1,1':4',1''-terphenyl]-4-carboxylic acid. LCMS


(Method 5-95 AB, ESI): $t_R = 0.615$ min, $[M + H]^+ = 908.4$.


Example 51: Synthesis of Compound 145

[00461] Compound 145 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.632$ min, $[M + H]^+ = 874.6$.

Example 52: Synthesis of Compound 146

[00462] Compound 146 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from 4-(4-isobutylphenyl)-2-methylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.653$ min, $[M + H]^+ = 888.5$. ^1H NMR (400 MHz, CD_3OD): δ 7.56-7.50 (m, 4H), 7.48-7.44 (m, 1H), 7.35-7.31 (m, 1H), 7.26-7.17 (m, 4H), 7.10-7.08 (m, 1H), 6.92-6.90 (m, 1H), 6.79 (br s, 1H), 6.39 (s, 1H), 5.19-5.15 (m, 1H), 4.81-4.79 (m, 3H), 4.30-4.15 (m, 5H), 3.36-3.33 (m, 1H), 3.25-3.05 (m, 7H), 2.95 (s, 3H), 2.53 (d, $J = 7.6$ Hz, 2H), 2.49 (s, 3H), 2.35-2.20 (m, 1H), 2.20-2.05 (m, 1H), 1.95-1.80 (m, 1H), 1.36 (d, $J = 7.2$ Hz, 3H), 0.94 (d, $J = 6.4$ Hz, 6H).


Example 53: Synthesis of Compound 147



[00463] An oven-dried three neck flask (500 mL) was charged with (4-chlorophenyl)boronic acid (12 g, 74.7 mmol), ethyl 4-iodobenzoate (14.1 g 51.2 mmol), $\text{Pd}_2(\text{dba})_3$ (4.68 g, 5.12 mmol), PCy_3 (1.43 g, 5.12 mmol) and K_2CO_3 (21.21 g, 153.5 mmol). DMF (100 mL) was added and the reaction mixture was purged with N_2 . The mixture was stirred at 80°C for 12 h. The reaction mixture was cooled to room temperature and filtered to remove K_2CO_3 . The solvent was removed and the brown residue was purified by column (1% to 5% EtOAc in petroleum ether) to give ethyl 4'-chloro-[1,1'-biphenyl]-4-carboxylate (9.52 g, 71.4%).


[00464] To a solution of ethyl 4'-chloro-[1,1'-biphenyl]-4-carboxylate (9.52 g, 36.6 mmol) in a mixture of THF (150 mL) and H_2O (20 mL) was added NaOH (4N, 5.86 g, 146 mmol). After the mixture was stirred at 70°C for 10 h, the organic solvent was removed under reduced pressure, and pH was adjusted to 3 with 4M HCl. The product was collected by filtration, washed with water and dried to give 4'-chloro-[1,1'-biphenyl]-4-carboxylic acid (8.5 g, 100%). ^1H NMR (400 MHz, CDCl_3) δ 8.10 (d, $J = 8.4$ Hz, 2 H), 7.70 (d, $J = 8$ Hz, 2 H), 7.65 (d, $J = 8.8$ Hz, 2 H), 7.47 (d, $J = 8.8$ Hz, 2 H).

[00465] Compound 147 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from 4'-chloro-[1,1'-biphenyl]-4-carboxylic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.710$, $[\text{M} + \text{H}]^+ = 852.4$; ^1H NMR (400 MHz, $\text{MeOH}-d_4$) δ 8.48 (brs, 2H, HCOOH), 7.97 (d, $J=8.4$ Hz, 2H), 7.77 (d, $J=8.4$ Hz, 2H), 7.68 (d, $J=8.4$ Hz, 2H), 7.50 (d, $J=8.4$ Hz, 2H), 7.32 (d, $J=8.0$ Hz, 1H), 7.24 (d, $J=8.0$ Hz, 1H), 7.16 (d, $J=8.0$ Hz, 1H), 7.09 (d, $J=8.0$ Hz, 1H), 6.89 (brs, 1H), 6.82 (brs, 1H), 6.34 (s, 1H), 5.22-5.14 (m, 1H), 4.83-4.79 (m, 2H), 4.32-4.14 (m, 4H), 4.20 (s, 2H), 3.37-3.33 (m, 2H), 3.27-2.95 (m, 7H), 2.89 (s, 3H), 2.37-2.30 (m, 1H), 2.22-2.14 (m, 1H), 1.36 (d, $J=6.4$ Hz, 3H).


Example 54: Synthesis of Compound 148

[00466] Compound 148 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from 4'-chloro-3-methyl-[1,1'-biphenyl]-4-carboxylic acid.


5 LCMS (Method 5-95 AB, ESI): $t_R = 0.560$, $[M + H]^+ = 866.2$.

Example 55: Synthesis of Compound 149

[00467] Compound 149 (free base) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.661$, $[M + H]^+ = 832.5$.


Example 56: Synthesis of Compound 150

[00468] Compound 150 (free base) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from 3-methyl-[1,1'-biphenyl]-4-carboxylic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.649$, $[M + H]^+ = 846.4$.

Example 57: Synthesis of Compound 151


[00469] Compound 151 (free base) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.735$, $[M + H]^+ = 860.5$;

5 ^1H NMR (400 MHz, MeOH- d_4) δ 7.45-7.38 (m, 2H), 7.36-7.16 (m, 7H), 7.06-6.98 (m, 2H), 6.84 (brs, 1H), 6.71 (brs, 1H), 6.50 (s, 1H), 5.20-5.13 (m, 1H), 4.78-4.71 (m, 1H), 4.54-4.46 (m, 1H), 4.23-3.97 (m, 4H), 4.21 (s, 2H), 3.10-2.80 (m, 5H), 3.00 (s, 3H), 2.67 (q, $J=6.8$ Hz, 2H), 2.33 (s, 3H), 2.16-2.06 (m, 1H), 2.06-1.96 (m, 1H), 1.36 (d, $J=6.8$ Hz, 3H), 1.26 (t, $J=6.8$ Hz, 3H).

Example 58: Synthesis of Compound 152


[00470] Compound 152 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from 4'-ethyl-3-methyl-[1,1'-biphenyl]-4-carboxylic acid.

10 LCMS (Method 5-95 AB, ESI): $t_R = 0.698$, $[M + H]^+ = 874.2$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.50 (brs, 1H, HCOOH), 7.50-7.44 (m, 5H), 7.35-7.33 (m, 1H), 7.29 (d, $J=8$ Hz, 2H), 7.22 (d, $J=8$ Hz, 2H), 7.12 (d, $J=8$ Hz, 1H), 6.92 (brs, 1H), 6.74-6.58 (m, 2H), 5.11-5.08 (m, 1H), 4.85-4.81 (m, 2H), 4.31-4.21 (m, 6H), 3.28-3.24 (m, 5H), 3.15-3.01 (m, 3H), 3.04 (s, 3H), 2.71 (q, $J=7.2$ Hz, 2H), 2.46 (s, 3H), 2.08-1.70 (m, 4H), 1.39 (d, $J=7.2$ Hz, 3H), 1.29 (t, $J=7.2$ Hz, 3H).

Example 59: Synthesis of Compound 153


[00471] To a solution of ethyl 2-amino-4-methylpyrimidine-5-carboxylate (4.0 g, 22 mmol) in
 5 CHBr₃ (66 mL) was added isopentyl nitrite (44 mL) and the mixture was stirred at 85°C for 4 h. The volatiles were removed and the residue was taken up by EtOAc (100 mL), which was washed by brine (100 mL x 2). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified by silica gel flash column to give ethyl 2-bromo-4-methylpyrimidine-5-carboxylate (3.0 g, 55.5% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃): δ 8.93 (s, 1H),
 10 4.41 (q, *J* = 7.2 Hz, 2H), 2.82 (s, 3H), 1.41 (t, *J* = 7.0 Hz, 3H).

[00472] Compound 153 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from ethyl 2-bromo-4-methylpyrimidine-5-carboxylate. LCMS (Method 5-95 AB, ESI): *t_R* = 0.736 min, [M+H]⁺ = 904.7.


Example 60: Synthesis of Compound 154

15


[00473] Compound 154 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): *t_R* = 0.661 min, [M+H]⁺ = 905.6.

Example 61: Synthesis of Compound 155

[00474] Compound 155 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.681$ min, $[M+H]^+ = 862.4$.


Example 62: Synthesis of Compound 156

[00475] Compound 156 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.617$ min, $[M+H]^+ = 890.4$. ^1H NMR (400 MHz, CD_3OD): δ 8.77 (s, 1H), 8.53 (s, 1H), 8.32 (d, $J = 8.4$ Hz, 2H), 7.32 (d, $J = 8.0$ Hz, 2H), 7.19 (d, $J = 8.4$ Hz, 2H), 7.07 (d, $J = 8.4$ Hz, 1H), 6.89 (s, 1H), 6.74 (s, 1H), 6.45 (s, 1H), 5.25-5.15 (m, 1H), 4.85-4.75 (m, 2H), 4.30-4.10 (m, 6H), 3.35 (s, 2H), 3.30-3.20 (m, 1H), 3.20-3.05 (m, 5H), 2.96 (s, 3H), 2.80-2.60 (m, 5H), 2.35-2.20 (m, 1H), 2.20-2.0 (m, 1H), 1.75-1.60 (m, 2H), 1.50-1.25 (m, 5H), 0.98 (t, $J = 7.2$ Hz, 3H).

Example 63: Synthesis of Compound 157


[00476] Compound 157 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.607$ min, $[M+H]^+ = 890.5$. ^1H NMR (400 MHz, CD_3OD): δ 8.70 (s, 1H), 8.10 (d, $J = 7.6$ Hz, 2H), 7.30-7.05 (m, 4H), 7.00-6.80 (m, 3H), 6.74 (s, 1H), 6.28 (s, 1H), 5.25-5.10 (m, 1H), 4.80-4.65 (m, 2H), 4.50-4.35 (m, 1H), 4.30-4.05 (m, 4H), 4.05-3.90 (m, 1H), 3.10-2.75 (m, 11H), 2.60-2.40 (m, 5H), 2.15-2.05 (m, 1H), 2.05-1.85 (m, 2H), 1.34 (d, $J = 6.8$ Hz, 3H), 0.97 (d, $J = 6.0$ Hz, 6H).

Example 64: Synthesis of Compound 158




[00477] Compound 158 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.722$ min, $[M+H]^+ = 910.4$.


Example 65: Synthesis of Compound 159




[00478] Compound 159 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.770$ min, $[M+Na]^+ = 940.0$.

Example 66: Synthesis of Compound 160

[00479] Compound 160 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.710$ min, $[M+H]^+ =$
 5 876.8.

Example 67: Synthesis of Compound 161


[00480] Compound 161 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.587$ min, $[M+H]^+ =$
 10 876.6.

Example 68: Synthesis of Compound 162

[00481] Compound 162 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.757$ min, $[M+H]^+ =$
 15 904.6. ^1H NMR (400 MHz, CD_3OD): δ 8.78 (s, 1H), 8.50 (br s, 2H), 8.35 (d, $J = 8.4$ Hz, 2H), 7.35-7.25 (m, 3H), 7.23 - 7.18 (m, 2H), 7.08 (d, $J = 8.4$ Hz, 1H), 6.90 (s, 1H), 6.79 (s, 1H), 6.40 (s, 1H), 5.19 - 5.16 (m, 1H), 4.80-4.70 (m, 2H), 4.21 - 4.18 (m, 6H), 3.15 - 3.13 (m, 7H), 2.95 (s,


3H), 2.72 - 2.68 (m, 6H), 2.32 - 2.14 (m, 2H), 1.70 - 1.66 (m, 2H), 1.38 - 1.35 (m, 7H), 0.92 (t, $J = 6.8$ Hz, 3H).

Example 69: Synthesis of Compound 163




5 [00482] Compound 163 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.711$ min, $[M+H]^+ = 890.5$. ^1H NMR (400 MHz, CD_3OD): δ 8.78 (s, 1H), 8.48 (br s, 2H), 8.36 (d, $J = 8.4$ Hz, 2H), 7.55 (d, $J = 8.4$ Hz, 2H), 7.32 (d, $J = 8.8$ Hz, 1H), 7.25-7.15 (m, 2H), 7.08 (d, $J = 8.8$ Hz, 1H), 6.90 (s, 1H), 6.79 (s, 1H), 6.39 (s, 1H), 5.19 - 5.16 (m, 1H), 4.80-4.70 (m, 2H), 4.25 - 4.10 (m, 10 6H), 3.20 - 3.10 (m, 8H), 2.95 (s, 3H), 2.75 - 2.65 (m, 4H), 2.32 - 2.14 (m, 1H), 1.40 - 1.30 (m, 12H).

Example 70: Synthesis of Compound 164




15 [00483] Compound 164 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.727$ min, $[M+H]^+ = 906.5$. ^1H NMR (400 MHz, CD_3OD): 8.74 (s, 1H), 8.50 (br s, 2H), 8.36 (d, $J = 8.8$ Hz, 2H), 7.32 (d, $J = 8.4$ Hz, 1H), 7.20 (d, $J = 8.4$ Hz, 2H), 7.09 (d, $J = 8.4$ Hz, 1H), 7.02 (d, $J = 8.8$ Hz, 2H), 6.89 (s, 1H), 6.75 (s, 1H), 6.44 (s, 1H), 5.25 - 5.15 (m, 1H), 4.85-4.75 (m, 2H), 4.30 - 4.15 (m, 6H), 4.08 (d, $J = 6.4$ Hz, 2H), 3.25 - 3.10 (m, 8H), 2.96 (s, 3H), 2.67 (s, 3H), 2.35 - 2.25 (m, 20 1H), 2.25-2.10 (m, 1H), 1.85-1.75 (m, 2H), 1.60-1.50 (m, 2H), 1.36 (d, $J = 6.8$ Hz, 3H), 1.02 (t, $J = 7.4$ Hz, 3H).

Example 71: Synthesis of Compound 165


[00484] A mixture of ethyl diacetoacetate (4.0 g, 23.23 mmol) and Cs_2CO_3 (15.2 g, 46.46 mmol) in acetonitrile (50 mL) was stirred at 25°C for 30 min and cooled to 0°C. MeOTf (3.81 g, 23.23 mmol) was added dropwise and the mixture was stirred at 25°C for 1.5 h. The mixture was diluted with ethyl ether and filtered, and the filtrate was concentrated. The residue was taken up in ethyl ether (100 mL), washed with 2N NaOH (10 mL x 2), dried over Na_2SO_4 and concentrated to give crude ethyl (E)-2-acetyl-3-methoxy-but-2-enoate (2.7 g, 62.4% yield) as a yellow oil, which was used directly in the next step.

[00485] A mixture of ethyl (E)-2-acetyl-3-methoxy-but-2-enoate (2.7 g, 14.5 mmol), guanidine hydrochloride (3463.0 mg, 36.25 mmol) and NaOMe (1566.6 mg, 29 mmol) in ethanol (20 mL) was stirred at 80°C for 12 h, and cooled to 0°C and filtered. The filtrate was concentrated under reduced pressure. The residue was taken up in EtOAc (100 mL), washed with brine (50 mL x 3), dried over Na_2SO_4 and concentrated. The residue was purified by silica gel column (10% EtOAc in petroleum ether, $R_f = 0.45$) to give ethyl 2-amino-4,6-dimethyl-pyrimidine-5-carboxylate (820 mg, 29% yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 6.96 (s, 2H), 4.26 (q, $J = 7.2$ Hz, 2H), 2.30 (s, 6H), 1.29 (t, $J = 7.4$ Hz, 3H).


[00486] Compound 165 (formic acid salt) was prepared as a white solid utilizing the methods in Example 59 (Compound 153). LCMS (Method 5-95 AB, ESI): $t_R = 0.707$ min, $[\text{M}+\text{H}]^+ = 890.4$. ^1H NMR (400 MHz, CD_3OD): 8.50 (br s, 2H), 8.30-8.20 (m, 2H), 7.30-7.28 (m, 3H), 7.19 (d, $J = 8.4$ Hz, 2H), 7.08 (d, $J = 8.4$ Hz, 1H), 6.89 (s, 1H), 6.74 (s, 1H), 6.46 (s, 1H), 5.25 - 5.20 (m, 1H), 4.80-4.70 (m, 2H), 4.30 - 4.10 (m, 6H), 3.25 - 3.10 (m, 8H), 3.00 (s, 3H), 2.67 (t, $J = 7.4$ Hz, 2H), 2.53 (s, 6H), 2.35 - 2.20 (m, 1H), 2.20-2.05 (m, 1H), 1.73-1.67 (m, 2H), 1.35 (d, $J = 6.8$ Hz, 3H), 0.98 (t, $J = 7.2$ Hz, 3H).

Example 72: Synthesis of Compound 166


[00487] Compound 166 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.613$ min, $[M+H]^+ =$
 5 895.5.

Example 73: Synthesis of Compound 167


[00488] Compound 167 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.604$ min, $[M+H]^+ =$
 10 889.9.

Example 74: Synthesis of Compound 168


[00489] Compound 168 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.745$ min, $[M+H]^+ =$
 15 889.4.

Example 75: Synthesis of Compound 169


[00490] Compound 169 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.644$ min, $[M+H]^+ = 889.5$. ^1H NMR (400 MHz, CD_3OD): δ 7.88 (d, $J = 7.6$ Hz, 2H), 7.83 (d, $J = 8.0$ Hz, 1H), 7.72 (d, $J = 8.0$ Hz, 1H), 7.35 - 7.25 (m, 3H), 7.25 - 7.15 (m, 2H), 7.10 (d, $J = 8.4$ Hz, 1H), 6.90 (s, 1H), 6.76 (s, 1H), 6.44 (s, 1H), 5.20 - 5.10 (m, 1H), 4.85 - 4.75 (m, 2H), 4.30 - 4.15 (m, 6H), 3.25 - 3.05 (m, 8H), 2.96 (s, 3H), 2.67 (s, 3H), 2.56 (d, $J = 7.2$ Hz, 2H), 2.35 - 2.20 (m, 1H), 2.20 - 2.10 (m, 1H), 2.00 - 1.85 (m, 1H), 1.36 (d, $J = 6.4$ Hz, 3H), 0.95 (d, $J = 6.4$ Hz, 6H).

Example 76: Synthesis of Compound 170


[00491] Compound 170 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.737$ min, $[M+H]^+ = 906.5$. ^1H NMR (400 MHz, CD_3OD): δ 8.82 (s, 1H), 8.50 (br s, 2H, *FA*), 8.43 (d, $J = 8.0$ Hz, 2H), 7.69 (d, $J = 8.0$ Hz, 2H), 7.34-7.32 (m, 1H), 7.24 (d, $J = 8.4$ Hz, 1H), 7.19 (d, $J = 8.8$ Hz, 1H), 7.09 (d, $J = 8.4$ Hz, 1H), 6.92 (br s, 1H), 6.82 (br s, 1H), 6.39 (s, 1H), 5.18 - 5.17 (m, 1H), 4.82 - 4.79 (m, 2H), 4.25-4.15 (m, 6H), 3.50-3.30 (m, 3H), 3.15 - 3.12 (m, 5H), 2.96 (s, 3H), 2.72 (s, 3H), 2.40-2.10 (m, 2H), 1.36 (d, $J = 6.8$ Hz, 3H), 0.32 (s, 9H).


Example 77: Synthesis of Compound 171

[00492] Compound 171 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.697$ min, $[M+H]^+ = 902.7$.

Example 78: Synthesis of Compound 172

[00493] Compound 172 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.769$ min, $[M+H]^+ = 924.5$.

Example 79: Synthesis of Compound 173




[00494] Step 1: A mixture of potassium acetate (649.2 mg, 6.55 mmol), 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (159.7 mg, 0.22 mmol),
 5 bis(pinacolato)diboron (1108.5 mg, 4.37 mmol) and methyl 4-bromo-2-methylbenzoate (500.0 mg, 2.18 mmol) in 1,4-dioxane (20 mL) was stirred at 110 °C for 4 h under N_2 protection, and evaporated to dryness. The residue was taken up in EtOAc (30 mL), washed with water (20 mL x 2) and brine (20 mL), dried over MgSO_4 and concentrated. The residue was purified by flash
 10 column chromatography (10% ethyl acetate in petroleum ether, $R_f = 0.6$) to afford methyl 2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (400 mg, 66.4% yield) as a yellow oil.

[00495] Step 2: To a solution of 5-bromo-2-chloropyridine (200.0 mg, 1.04 mmol) in toluene (10 mL) and water (2 mL) were added cesium carbonate (1693.1 mg, 5.2 mmol), isobutylboronic acid (158.9 mg, 1.56 mmol) and tetrakis(triphenylphosphine)palladium(0)
 15 (120.1 mg, 0.10 mmol). The reaction mixture was stirred at 110 °C for 16 h and filtered. The filtrate was diluted with H_2O (20 mL) and extracted with EtOAc (40 mL x 2). The combined organic layers were washed with water (80 mL x 3) and brine (80 mL), dried over Na_2SO_4 and concentrated. The residue was purified by prep-TLC (7.5% EtOAc in petroleum ether) to obtain 2-chloro-5-isobutylpyridine (110 mg, 62.4% yield) as a yellow oil. LCMS (Method 5-95 AB,
 20 ESI): $t_R = 0.872$ min, $[\text{M}+\text{H}]^+ = 169.8$.

[00496] Step 3: A mixture of 2-chloro-5-isobutylpyridine (50.0 mg, 0.29 mmol), 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (21.6 mg, 0.03 mmol), methyl 2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (122.1 mg, 0.44 mmol) and cesium carbonate (288.1 mg, 0.88 mmol) in toluene (5 mL) and water (1 mL) was stirred at 110 °C
 25 for 16 h under nitrogen. The mixture was diluted with H_2O (10 mL) and extracted with EtOAc (10 mL x 2). The combined organic layers were washed with water (20 mL x 4) and brine (20 mL), dried over Na_2SO_4 and concentrated. The residue was purified by prep-TLC (7.5% EtOAc in petroleum) to obtain methyl 4-(5-isobutylpyridin-2-yl)-2-methylbenzoate (74 mg, 88.6% yield) as a yellow oil. LCMS (Method 5-95 AB, ESI): $t_R = 0.816$ min, $[\text{M}+\text{H}]^+ = 283.9$.

[00497] Compound 173 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.639$ min, $[M+H]^+ = 889.5$.


Example 80: Synthesis of Compound 174



5

[00498] Compound 174 (formic acid salt) was prepared as a white solid utilizing the methods in Example 79 (Compound 173) from 5-bromo-2-chloropyrimidine. LCMS (Method 5-95 AB, ESI): $t_R = 0.553$ min, $[M+H]^+ = 890.9$.


Example 81: Synthesis of Compound 175



10


[00499] Compound 175 (formic acid salt) was prepared as a white solid utilizing the methods in Example 79 (Compound 173) from 5-bromo-2-chloropyrimidine. LCMS (Method 5-95 AB, ESI): $t_R = 0.699$ min, $[M+H]^+ = 890.5$.

Example 82: Synthesis of Compound 176




15

Example 83: Synthesis of Compound 177



Example 84: Synthesis of Compound 178




-169-

(7.5% EtOAc in petroleum ether) to obtain methyl 4-(5-bromopyrazin-2-yl)-2-methylbenzoate (150 mg, 58.1% yield) as a white solid. ^1H NMR (400 MHz, CDCl_3): δ 8.81 (s, 1H), 8.75 (d, $J = 1.2$ Hz, 1H), 8.04 (d, $J = 8.4$ Hz, 1H), 7.88 (s, 1H), 7.85 (d, $J = 8.4$ Hz, 1H), 3.93 (s, 3H), 2.70 (s, 3H).


- 5 **[00503]** To a solution of isobutylboronic acid (99.6 mg, 0.98 mmol) in toluene (3 mL) and water (0.3 mL) were added tetrakis(triphenylphosphine)palladium(0) (56.4 mg, 0.05 mmol), potassium carbonate (202.5 mg, 1.47 mmol) and methyl 4-(5-bromopyrazin-2-yl)-2-methylbenzoate (150.0 mg, 0.49 mmol). The reaction mixture was stirred at 100 °C for 16 h and filtered. The filtrate was diluted with H_2O (20 mL) and extracted with EtOAc (40 mL x 2). The
10 combined organic layers were washed with water (80 mL x 3) and brine (80 mL), dried over Na_2SO_4 and concentrated. The residue was purified by prep-TLC (9.5% EtOAc in petroleum ether, $R_f = 0.4$) to obtain methyl 4-(5-isobutylpyrazin-2-yl)-2-methyl-benzoate (52 mg, 37.4% yield) as a yellow oil. LCMS (Method 5-95 AB, ESI): $t_R = 0.956$ min, $[\text{M}+\text{H}]^+ = 284.9$.

- [00504]** Compound 178 (formic acid salt) was prepared as a white solid utilizing the methods
15 in Example 47 (Compound 141). LCMS (Method 5-95 AB, ESI): $t_R = 0.679$ min, $[\text{M}+\text{H}]^+ = 890.5$.


Example 85: Synthesis of Compound 179




- [00505]** Compound 179 (formic acid salt) was prepared as a white solid utilizing the methods
20 in Example 84 (Compound 178) from 5-bromo-2-chloropyrimidine. LCMS (Method 5-95 AB, ESI): $t_R = 0.657$ min, $[\text{M}+\text{H}]^+ = 890.2$.

Example 86: Synthesis of Compound 180


[00506] Compound 180 (formic acid salt) was prepared as a white solid utilizing the methods in Example 84 (Compound 178) from 3,6-dibromopyridazine. LCMS (Method 5-95 AB, ESI): $t_R = 0.633$ min, $[M+H]^+ = 890.4$.

Example 87: Synthesis of Compound 181


[00507] Compound 181 (formic acid salt) was prepared as a white solid utilizing the methods in Example 84 (Compound 178). LCMS (Method 5-95 AB, ESI): $t_R = 0.503$ min, $[M+H]^+ = 889.4$.

Example 88: Synthesis of Compound 182

[00508] Compound 182 (formic acid salt) was prepared as a white solid utilizing the methods in Example 79 (Compound 173) from 2-chloro-5-isobutylpyrimidine and methyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-naphthoate. LCMS (Method 5-95 AB, ESI): $t_R = 0.596$ min, $[M+H]^+ = 926.3$.

Example 89: Synthesis of Compound 183

[00509] Compound 183 (formic acid salt) was prepared as a white solid utilizing the methods in Example 47 (Compound 141) from 4,4,5,5-tetramethyl-2-(4-(tert-pentyl)phenyl)-1,3,2-dioxaborolane. LCMS (Method 5-95 AB, ESI): $t_R = 0.743$ min, $[M+H]^+ = 905.2$. ^1H NMR (400 MHz, CD_3OD): δ 8.79 (s, 1H), 8.52 (br s, 3H), 8.38 (d, $J = 8.4$ Hz, 2H), 7.50 (d, $J = 8.8$ Hz, 2H), 7.33 - 7.30 (m, 1H), 7.22 (d, $J = 8.8$ Hz, 1H), 7.18 (d, $J = 8.8$ Hz, 1H), 7.08 (d, $J = 8.4$ Hz, 1H), 6.90 (br s, 1H), 6.80 (br s, 1H), 6.39 (s, 1H), 5.19 - 5.17 (m, 1H), 4.82 - 4.75 (m, 1H), 4.20-4.13 (m, 7H), 3.15 - 3.05 (m, 8H), 2.95 (s, 3H), 2.70 (s, 3H), 2.30-2.10 (m, 2H), 1.74 (q, $J = 7.3$ Hz, 2H), 1.36 (s, 6H), 1.30-1.20 (m, 3H), 0.71 (t, $J = 7.6$ Hz, 3H).

Example 90: Synthesis of Compound 184

[00510] The Sonogashiro coupling of an acetylene to an aromatic halide, followed by reduction of the alkyne and hydrolysis of the ester is described and is referred to as General Method 8. A mixture of methyl 4-bromo-2-fluorobenzoate (500.0 mg, 2.15 mmol), oct-1-yne (702.9 mg, 6.44 mmol), bis(triphenylphosphine)palladium(II)dichloride (75.3 mg, 0.11 mmol) and copper(I)


iodide (20.4 mg, 0.11 mmol) in triethylamine (9.83 mL, 70.9 mmol) was stirred at 100 °C for 2 h under nitrogen atmosphere. LCMS (5-95AB/1.5min): $t_R = 1.006$ min, $[M + H]^+$ 262.9 showed 60% of DP. The reaction was quenched with water (15 mL) and extracted with dichloromethane (3×25 mL). The combined organic extracts were washed with brine (2×25 mL), dried over anhydrous Na_2SO_4 and filtered. The filtrate was concentrated and the residue was purified by column chromatography on silica gel (eluting with 5% ethyl acetate in petroleum ether, $R_f = 0.5$) to afford methyl 2-fluoro-4-(oct-1-yn-1-yl)benzoate (550 mg, 97.7% yield) as a brown solid. LCMS (5-95AB_1.5 min): $t_R = 1.006$ min, $[M + H]^+$ 262.9.


[00511] To a solution of methyl 2-fluoro-4-(oct-1-yn-1-yl)benzoate (550.0 mg, 2.1 mmol) in methanol (25 mL) was added 10% Palladium on carbon (111.56 mg, 0.10 mmol). The mixture was stirred at 30 °C under hydrogen (40 psi) for 16h. The reaction was filtered over a pad of Celite and concentrated. The residue was purified by column chromatography on silica gel (eluting with petroleum ether / ethyl acetate from 100:1 to 10:1) to afford methyl 2-fluoro-4-octylbenzoate (500 mg, 89.5% yield) as a yellow solid. LCMS (5-95AB_1.5min): $t_R = 1.033$ min, $[M + H]^+$ 266.

[00512] To a solution of methyl 2-fluoro-4-octylbenzoate (500.0 mg, 1.88 mmol) in methanol (5 mL) was added NaOH (1000.0 mg, 25 mmol) in water (5 mL). The mixture was stirred at 100 °C for 2h, cooled to RT and hydrochloric acid (1.0 M) was added until pH = 3-4. The mixture was extracted with ethyl acetate (3×30 mL). The combined organic extracts were washed with brine (2×30 mL), dried over sodium sulfate and filtered. The filtrate was concentrated to give 2-fluoro-4-octylbenzoic acid (450 mg, 95% yield) as a yellow solid. ^1H NMR (400 MHz, $CDCl_3$): δ 7.94 (t, $J = 8.0$ Hz, 1H), 7.06 (d, $J = 8.4$ Hz, 1H), 6.99 (d, $J = 12.0$ Hz, 1H), 2.66 (t, $J = 7.4$ Hz, 2H), 1.65 - 1.62 (m, 2H), 1.31 - 1.28 (m, 10H), 0.89 (t, $J = 6.8$ Hz, 3H).

[00513] Compound 184 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from 2-fluoro-4-octylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.775$ min, $[M + H]^+ = 872.4$.

Example 91: Synthesis of Compound 185






[00514] To a solution of sodium thiomethoxide (45.3 mg, 0.65 mmol) in N,N-dimethylformamide (3 mL) was added methyl 2-fluoro-4-octylbenzoate (86.0 mg, 0.32 mmol) and the mixture was stirred at 20 °C for 16 h. The mixture was diluted with water (5 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were dried over sodium sulfate, filtered and concentrated. The residue was purified by prep-TLC (10% EtOAc in petroleum ether, $R_f = 0.7$) to give methyl 2-(methylthio)-4-octylbenzoate (30 mg, 31.6% yield) as a yellow oil. ^1H NMR (400 MHz, CDCl_3): δ 7.90 (d, $J = 8.0$ Hz, 1H), 7.03 (s, 1H), 6.95 (d, $J = 8.0$ Hz, 1H), 3.88 (s, 3H), 2.62 (t, $J = 7.6$ Hz, 2H), 2.45 (s, 3H), 1.69 - 1.56 (m, 2H), 1.27 (m, 10H), 0.86 (t, $J = 6.6$ Hz, 3H).

[00515] Methyl 2-(methylthio)-4-octylbenzoate (30.0 mg, 0.10 mmol) was hydrolyzed with LiOH according to General Method 5 (Example 5) to give crude 2-(methylthio)-4-octylbenzoic acid (22 mg, 77% yield) as a yellow solid.

[00516] Compound 185 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from 2-(methylthio)-4-octylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.676$ min, $[\text{M} + \text{H}]^+ = 900.4$.

Example 92: Synthesis of Compound 186




[00517] To a solution of 2-fluoro-4-octylbenzoic acid (120.0 mg, 0.48 mmol) in tetrahydrofuran (3 mL) was added ethylmagnesium bromide (0.55 mL, 1.66 mmol) in Et_2O (3 M) dropwise at 0 °C. The mixture was stirred at 15 °C for 16 h. The reaction was quenched with water (2 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were washed

with saturated brine (15 mL x 3), dried over Na₂SO₄ and concentrated. The residue was purified by prep-TLC (20% ethyl acetate in petroleum ether, R_f = 0.5) to give 2-ethyl-4-octylbenzoic acid (70 mg, 56.1% yield) as a white solid.

[00518] Compound 186 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from 2-ethyl-4-octylbenzoic acid. LCMS (Method 5-95 AB, ESI): t_R = 0.787 min, [M + H]⁺ = 882.5.


Example 93: Synthesis of Compound 187




[00519] 3-Methyl-5-octylpicolinic acid was prepared using the methods from Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): t_R = 0.730 min, [M + H]⁺ = 250.0.

[00520] Compound 187 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184) from 3-methyl-5-octylpicolinic acid. LCMS (Method 5-95 AB, ESI): t_R = 0.658 min, [M + H]⁺ = 869.6.

Example 94: Synthesis of Compound 188




[00521] Compound 188 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): t_R = 0.614 min, [M + H]⁺ = 872.7.

Example 95: Synthesis of Compound 189


[00522] 4-Heptyl-2,6-dimethylbenzoic acid a colorless oil, was prepared using the methods from Example 90 (Compound 184). ¹H NMR (400 MHz, CDCl₃) δ 6.76 (s, 2H), 2.44 (t, J = 7.6 Hz, 2H), 2.27 (s, 6H), 1.55-1.45 (m, 2H), 1.30-1.15 (m, 8H), 0.816 (t, J = 5.8 Hz, 3H).

[00523] Compound 189 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184) from 4-heptyl-2,6-dimethylbenzoic acid. LCMS (Method 5-95 AB, ESI): t_R = 0.776 min, [M + H]⁺ = 869.4.

Example 96: Synthesis of Compound 190

[00524] 2,6-Difluoro-4-heptylbenzoic acid, a yellow solid, was prepared using the methods from Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): t_R = 0.873 min, [M + H]⁺ = 257.9.

[00525] Compound 190 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184) from 2,6-difluoro-4-heptylbenzoic acid. LCMS (Method 5-95 AB, ESI): t_R = 0.636 min, [M + H]⁺ = 876.4.


Example 97: Synthesis of Compound 191

[00526] 4-Heptyl-2-(trifluoromethyl)benzoic acid, a brown solid, was prepared using the methods from Example 90 (Compound 184). ¹H NMR (400 MHz, CD₃OD): δ 7.73 (d, J = 8.0

5 Hz, 1H), 7.61 (s, 1H), 7.53 (d, J = 8.0 Hz, 1H), 3.89 (s, 3H), 2.74 (t, J = 7.8 Hz, 2H), 1.70 - 1.60 (m, 2H), 1.40 - 1.20 (m, 8H), 0.90 (t, J = 6.8 Hz, 3H).

[00527] Compound 191 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184) from 4-heptyl-2-(trifluoromethyl)benzoic acid. LCMS (Method 5-95 AB, ESI): t_R = 0.640 min, [M + H]⁺ = 908.7.


10 **Example 98: Synthesis of Compound 192**




[00528] 4-Octyl-2-(trifluoromethyl)benzoic acid, a yellow solid, was prepared using the methods from Example 90 (Compound 184). ¹H NMR (400 MHz, CD₃OD): δ 7.76 (d, J = 8.0 Hz, 1H), 7.59 (s, 1H), 7.51 (d, J = 7.2 Hz, 1H), 2.74 (t, J = 7.6 Hz, 2H), 1.70 - 1.60 (m, 2H),

15 1.40 - 1.20 (m, 10H), 0.90 (t, J = 7.0 Hz, 3H).


[00529] Compound 192 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): t_R = 0.664 min, [M + H]⁺ = 922.4.

Example 99: Synthesis of Compound 193

[00530] Compound 193 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.637$ min, $[M + H]^+ = 871.3$.


Example 100: Synthesis of Compound 194

[00531] Compound 194 (formic acid salt) was prepared as a white solid (42.7 mg) utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.633$, $[M + H]^+ = 868.5$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.51 (brs, 2H, HCOOH), 7.36-7.27 (m, 2H), 7.24 (d, $J=8.0$ Hz, 1H), 7.17 (d, $J=8.0$ Hz, 1H), 7.13-7.06 (m, 3H), 6.90 (brs, 1H), 6.83 (brs, 1H), 6.42 (s, 1H), 5.04-5.00 (m, 1H), 4.81-4.77 (m, 2H), 4.27-4.15 (m, 4H), 4.20 (s, 2H), 3.36-3.30 (m, 1H), 3.22-3.10 (m, 5H), 3.05-2.93 (m, 2H), 2.99 (s, 3H), 2.61 (t, $J=7.6$ Hz, 2H), 2.45 (s, 3H), 2.03-1.95 (m, 2H), 1.90-1.80 (m, 3H), 1.65-1.58 (m, 2H), 1.40-1.25 (m, 10H), 0.90 (t, $J=6.4$ Hz, 3H).

Example 101: Synthesis of Compound 195


[00532] Compound 195 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.590$, $[M + H]^+ = 826.4$.

Example 102: Synthesis of Compound 196




5 **[00533]** Compound 196 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.704$, $[M + H]^+ = 826.5$.

Example 103: Synthesis of Compound 197



10 **[00534]** Compound 197 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.584$, $[M + H]^+ = 840.5$.


Example 104: Synthesis of Compound 198



15 **[00535]** Compound 198 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.753$, $[M + H]^+ = 854.4$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.49 (brs, 1H), 7.34 (d, $J=8.0$ Hz, 2H), 7.26 (d, $J=8.0$ Hz, 1H), 7.13 (brs, 1H), 7.11 (d, $J=8.0$ Hz, 2H), 6.92 (brs, 1H), 6.83 (brs, 1H), 6.35 (s, 1H), 5.17-5.14 (m,

2H), 4.83-4.81 (m, 1H), 4.30-4.20 (m, 4H), 4.22 (s, 2H), 3.32-3.13 (m, 8H), 2.95 (s, 3H), 2.64 (t, $J=7.2$ Hz, 2H), 2.43 (s, 3H), 2.40-2.25 (m, 1H), 2.15-2.02 (m, 1H), 1.64 (brs, 2H), 1.40-1.20 (m, 11H), 0.92 (t, $J=6.8$ Hz, 3H).



Example 105: Synthesis of Compound 199



5

[00536] Compound 199 was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.596$, $M+H^+ = 868.4$.

Example 106: Synthesis of Compound 200



10


[00537] Steps 1-3: Starting from ethyl 2-bromo-4-methylpyrimidine-5-carboxylate, the standard procedure of Sonogashiro coupling, hydrogenation and ester hydrolysis (General Method Sonogashiro) afforded 4-methyl-2-pentylpyrimidine-5-carboxylic acid (105 mg) as a yellow solid. ^1H NMR (400 MHz, DMSO- d_6) δ 8.97 (s, 1 H), 2.85 (d, $J=7.6$ Hz, 2H), 2.69 (s, 3H), 1.79-1.72 (m, 2H), 1.32-1.28 (m, 4H), 0.89-0.83 (t, $J=6.8$ Hz, 3H).

15

[00538] Compound 200 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.492$, $[M + H]^+ = 828.1$;


^1H NMR (400 MHz, $\text{MeOH-}d_4$) δ 8.70 (s, 1H), 8.50 (brs, 2H, HCOOH), 7.33 (d, $J=8.0$, 1H), 7.26 (d, $J=8.0$, 1H), 7.20 (d, $J=8.4$, 1H), 7.11 (d, $J=8.4$, 1H), 6.92 (brs, 1H), 6.84 (brs, 1H), 6.39 (s, 1H), 5.19-5.15 (m, 1H), 4.80-4.70 (m, 2H), 4.27-4.18 (m, 4H), 4.22 (s, 2H), 3.33-3.12 (m, 9H), 2.96 (s, 3H), 2.95 (t, $J=8.0$ Hz, 2H), 2.65 (s, 3H), 2.29-2.19 (m, 1H), 2.17-2.05 (m, 1H), 1.86-1.78 (m, 2H), 1.40-1.25 (m, 7H), 0.94 (t, $J=6.4$ Hz, 3H).

Example 107: Synthesis of Compound 201




[00539] Compound 201 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184) from 6-hexylnicotinic acid. LCMS (Method 5-95 AB, ESI): t_R = 0.537, $[\text{M} + \text{H}]^+ = 827.3$; ^1H NMR (400 MHz, $\text{MeOH-}d_4$) δ 8.92 (d, $J=2$ Hz, 1H), 8.48 (brs, 2H, HCOOH), 8.21 (d, $J=8.0$ Hz, 1H), 7.46 (d, $J=8.0$ Hz, 1H), 7.32 (d, $J=8.0$ Hz, 1H), 7.25 (d, $J=8.0$ Hz, 1H), 7.17 (d, $J=8.0$ Hz, 1H), 7.09 (d, $J=8.0$ Hz, 1H), 6.88 (brs, 1H), 6.81 (brs, 1H), 6.34 (s, 1H), 5.18-5.14 (m, 2H), 4.8-4.77 (m, 1H), 4.26-4.10 (m, 4H), 4.20 (s, 2H), 3.37-3.33 (m, 1H), 3.22-3.17 (m, 5H), 3.14-3.10 (m, 3H), 2.89-2.85 (m, 4H), 2.33-2.20 (m, 1H), 2.20-2.05 (m, 1H), 1.76-1.60 (m, 3H), 1.40-1.20 (m, 8H), 0.91 (t, $J=6.6$ Hz, 3H).


Example 108: Synthesis of Compound 202




[00540] Compound 202 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.749$ min, $[\text{M} + \text{H}]^+ = 840.5$.

Example 109: Synthesis of Compound 203

[00541] Compound 203 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.641$ min, $[M+H]^+ =$ 872.4.

Example 110: Synthesis of Compound 204

[00542] Compound 204 (formic acid salt) was prepared as a white solid utilizing the methods in Example 90 (Compound 184). LCMS (Method 5-95 AB, ESI): $t_R = 0.636$ min, $[M+H]^+ =$ 866.6.


Example 111: Synthesis of Compound 205

[00543] Compound 205 (formic acid salt) was prepared utilizing the methods in Example 90 (Compound 184) except the alkyne reduction was performed as follows. To a solution of

methyl 2-chloro-4-(hept-1-yn-1-yl)-6-methylbenzoate (250.0 mg, 0.90 mmol) in toluene (10 mL) was added chlorotris(triphenylphosphine)rhodium(I) (124.5 mg, 0.13 mmol). The mixture was stirred under 15 psi hydrogen pressure at 60 °C for 16 h, filtered and evaporated. The residue was purified by chromatography on silica (0-5% EtOAc in petroleum ether) to afford methyl 2-chloro-4-heptyl-6-methylbenzoate (200 mg, 78.9% yield) as a yellow solid. ¹H NMR (400 MHz, CD₃OD): δ 7.10 (s, 1H), 7.03 (s, 1H), 3.90 (s, 3H), 2.58 (t, J = 7.8 Hz, 2H), 2.27 (s, 3H), 1.65 - 1.55 (m, 2H), 1.35 - 1.25 (m, 8H), 0.90 (t, J = 7.0 Hz, 3H).

[00544] Data for Compound 205: LCMS (Method 5-95 AB, ESI): *t_R* = 0.640 min, [M+H]⁺ = 888.5.

10 Example 112: Synthesis of Compound 206




[00545] Step 1: To a mixture of 3-bromo-2-methylbenzoic acid (5.0 g, 23 mmol) in MeOH (80 mL) was added SOCl₂ (11.0 g, 93 mmol) at 20°C. The mixture was stirred for 1.5h at 70°C. The volatiles were removed and the residue was taken up by EtOAc (100 mL), which was washed sequentially with saturated NaHCO₃ and brine (each 100 mL). The EtOAc layer was dried over Na₂SO₄, concentrated and the residue was purified by flash column chromatography to give methyl 3-bromo-2-methylbenzoate (5.3g, 99% yield) as a red solid.

[00546] Step 2: A solution of methyl 3-bromo-2-methylbenzoate (500 mg, 2.2 mmol), *n*-butyl boronic acid (890 mg, 8.7 mmol), Pd(PPh₃)₄ (252 mg, 0.22 mmol) and K₂CO₃ (905 mg, 6.6 mmol) in toluene (20 mL) was stirred at 100°C for 4 h. After filtration, the filtrate was washed with brine (20 mL x 3), dried over Na₂SO₄ and concentrated. The residue was purified by HPLC to give methyl 3-butyl-2-methylbenzoate (120 mg, 27% yield) as colorless oil. LCMS (Method 5-95 AB, ESI): *t_R* = 0.871, [M + H]⁺ = 206.9.

[00547] Step 3: The ester hydrolysis method with NaOH (General Method NaOH) previously described (Example 47) was applied to methyl 3-butyl-2-methylbenzoate (120 mg, 0.58 mmol) to afford 3-butyl-2-methylbenzoic acid (110 mg, 98% yield) as a white solid.


[00548] Compound 206 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from 3-butyl-2-methylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.541$, $[M + H]^+ = 826.3$.

Example 113: Synthesis of Compound 207




[00549] Compound 207 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from 5-butyl-2-methylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.554$, $[M + H]^+ = 826.2$.


Example 114: Synthesis of Compound 208




[00550] Compound 208 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): $t_R = 0.546$, $[M + H]^+ = 812.3$.

Example 115: Synthesis of Compound 209

[00551] Compound 209 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from 2-methoxy-4-octylbenzoic acid. LCMS (Method 5-95
5 AB, ESI): $t_R = 0.782$ min, $[M/2+H]^+ = 442.9$.

Example 116: Synthesis of Compound 210


[00552] Compound 210 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): $t_R = 0.772$ min, $[M/2+H]^+ =$
10 444.8.

Example 117: Synthesis of Compound 211


[00553] Following similar procedures as described for Compound 206 (Example 112), 2-bromo-4-octylbenzoic acid (180 mg, 0.5747 mmol, 94% yield) was obtained as a white solid
15 from 2-bromo-4-iodobenzoic acid. ^1H NMR (400 MHz, CD_3OD): δ 7.74 (d, $J = 8.0$ Hz, 1H), 7.52 (s, 1H), 7.30 - 7.20 (m, 1H), 2.64 (t, $J = 7.8$ Hz, 2H), 1.65 - 1.55 (m, 2H), 1.35 - 1.20 (m, 10H), 0.90 (t, $J = 6.8$ Hz, 3H).

[00554] Compound 211 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from 2-bromo-4-octylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.626$ min, $[M + H]^+ = 932.3$.

Example 118: Synthesis of Compound 212




5



[00555] Step 1: To a solution of methyl 2-bromo-4-octylbenzoate (100.0 mg, 0.31 mmol) in dimethyl sulfoxide (2 mL) were added copper(I) iodide (174.6 mg, 0.92 mmol) and sodium methanesulfinate (93.6 mg, 0.92 mmol). The mixture was stirred at 100°C for 16 h under N_2 , diluted with water (10 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by prep-TLC (20% EtOAc in petroleum ether, $R_f = 0.3$) to afford methyl 2-methylsulfonyl-4-octylbenzoate (80 mg, 80.2% yield) as a white solid. ^1H NMR (400 MHz, CDCl_3): δ 7.95 (s, 1H), 7.65 (d, $J = 7.6$ Hz, 1H), 7.48 (d, $J = 7.2$ Hz, 1H), 3.97 (s, 3H), 3.37 (s, 3H), 2.72 (t, $J = 7.8$ Hz, 2H), 1.70-1.60 (m, 2H), 1.40-1.20 (m, 10H), 0.89 (t, $J = 6.4$ Hz, 3H).

[00556] Step 2: 2-(Methylsulfonyl)-4-octylbenzoic acid was prepared from methyl 2-(methylsulfonyl)-4-octylbenzoate utilizing General Method NaOH.

[00557] Compound 212 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from 2-(methylsulfonyl)-4-octylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.759$ min, $[M + H]^+ = 932.5$.


Example 119: Synthesis of Compound 213

[00558] 4-Methyl-2-octylthiazole-5-carboxylic acid, a white solid, was prepared using the methods in Example 112 (Compound 206) from ethyl 2-bromo-4-methylthiazole-5-carboxylate.

5 ^1H NMR (400 MHz, CDCl_3): δ 3.10 – 2.90 (m, 2H), 2.74 (s, 3H), 1.90 – 1.70 (m, 2H), 1.46 – 1.24 (m, 10H), 0.89 (t, 6.6 Hz, 3H).


[00559] Compound 213 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): $t_R = 0.775$ min, $[\text{M} + \text{H}]^+ = 875.6$.

10 **Example 120: Synthesis of Compound 214**




[00560] 4-Cyclobutyl-2-methylbenzoic acid, a white solid, was prepared using the methods in Example 112 (Compound 206). ^1H NMR (400 MHz, CDCl_3): δ 8.01 (d, $J = 8.0$ Hz, 1H), 7.14 – 7.11 (m, 2H), 3.57 (p, $J = 8.8$ Hz, 1H), 2.65 (s, 3H), 2.41 – 2.37 (m, 2H), 2.20 – 2.17 (m, 2H), 2.16 – 2.07 (m, 1H), 1.95 – 1.84 (m, 1H).


[00561] Compound 214 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from 4-cyclobutyl-2-methylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.682$ min, $[\text{M} + \text{H}]^+ = 810.7$.

Example 121: Synthesis of Compound 215

[00562] Compound 215 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from 4-cyclopentyl-2-methylbenzoic acid. LCMS (Method 5-
5 95 AB, ESI): $t_R = 0.550$ min, $[M + H]^+ = 824.3$.

Example 122: Synthesis of Compound 216


[00563] Compound 216 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from 4-cyclopropyl-2-methylbenzoic acid. LCMS (Method 5-
10 95 AB, ESI): $t_R = 0.479$ min, $[M + H]^+ = 797.2$.

Example 123: Synthesis of Compound 217

[00564] Step 1: To a solution of tetrakis(triphenylphosphine)palladium(0) (151.3 mg, 0.130 mmol) in toluene (10 mL) and water (2 mL) were added 3-methyl-1-butylboronic acid (759.4 mg, 6.55 mmol), methyl 4-bromo-2-methylbenzoate (300 mg, 1.31 mmol) and sodium carbonate (694 mg, 6.55 mmol). The reaction mixture was stirred at 110 °C for 16h. The mixture was filtered and the filtrate was diluted with H₂O (30 mL) and extracted with EtOAc (30 mL x 2). The organic layers were combined and washed with water (60 mL x 2) and brine (30 mL). The organic layers were separated, dried over Na₂SO₄ and concentrated. The residue was purified by preparative TLC (5% EtOAc in petroleum ether) to obtain methyl 4-isopentyl-2-methylbenzoate (250 mg, 1.1348 mmol, 86.6% yield) as a colorless oil.


[00565] Step 2: Methyl 4-isopentyl-2-methylbenzoate (250 mg, 1.13 mmol) was hydrolyzed as previously described (General Method NaOH) to give crude 4-isopentyl-2-methylbenzoic acid (200 mg, 0.970 mmol, 85.4% yield) as a yellow solid.

[00566] Compound 217 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.732$ min, $[M + H]^+ = 826.5$.

Example 124: Synthesis of Compound 218


[00567] 4-Isobutyl-2-methylbenzoic acid a white solid, was prepared using the methods from Example 112 (Compound 206). ¹H NMR (400 MHz, CDCl₃): δ 8.00 (d, J = 8.0 Hz, 1H), 7.14 – 6.94 (m, 2H), 2.65 (s, 3H), 2.50 (d, J = 6.8 Hz, 2H), 1.95-1.80 (m, 1H), 0.92 (d, J = 6.5 Hz, 6H).

[00568] Compound 218 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): t_R = 0.701 min, [M + H]⁺ = 812.4.

Example 125: Synthesis of Compound 219


[00569] 2-Methyl-4-neopentylbenzoic acid, a white solid, was prepared using the methods from Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): t_R = 0.807 min, [M + H]⁺ = 206.8.

[00570] Compound 219 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from 2-methyl-4-neopentylbenzoic acid. LCMS (Method 5-95 AB, ESI): t_R = 0.573 min, [M + H]⁺ = 826.3.

Example 126: Synthesis of Compound 220

[00571] To a degassed mixture of 4-tert-butyl-2-methylbenzoic acid (192.0 mg, 1 mmol),
 5 sodium persulfate (1.19 g, 4.99 mmol) and Selectfluor (1.77 g, 4.99 mmol) in acetonitrile (4 mL) and water (4 mL) in dry-ice acetone bath was added silver nitrate (17.0 mg, 0.10 mmol). The mixture was degassed by three freeze-pump-thaw cycles and heated at 80 °C for 16 h. Water (10 mL) was added and the mixture was extracted with EtOAc (15 mL x 2). The combined organic layers were concentrated and the residue was purified by prep-TLC
 10 (petroleum ether/EtOAc/HOAc 2/1/0.01, $R_f = 0.3$) to give 4-(tert-butyl)-2-(difluoromethyl)benzoic acid (75 mg, 32.9% yield) as a white solid.


[00572] Compound 220 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.593$ min, $[M + H]^+ = 848.3$.

Example 127: Synthesis of Compound 221

[00573] Methyl 2-(difluoromethyl)-4-octylbenzoate, a colorless oil, was prepared following similar procedures as described in Example 126 (Compound 220) and Example 112 (Compound 206). ¹H NMR (400 MHz, CDCl₃) δ 7.94 (d, J = 7.6 Hz, 1H), 7.61 (s, 1H), 7.52 (t, J = 55.8 Hz, 1H), 7.32 (d, J = 8.0 Hz, 1H), 3.90 (s, 3H), 2.68 (t, J = 7.8 Hz, 2H), 1.64 - 1.58 (m, 2H), 1.40 - 1.10 (m, 10H), 0.86 (t, J = 6.6 Hz, 3H). Methyl 2-(difluoromethyl)-4-octylbenzoate was hydrolyzed as as previously described (General Method NaOH) to give 2-(difluoromethyl)-4-octylbenzoic acid.


[00574] Compound 221 (formic acid salt) was prepared as a white solid utilizing the methods in Example 126 (Compound 220) and Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): t_R = 0.780 min, [M + H]⁺ = 904.8.

Example 128 Synthesis of Compound 222




[00575] Compound 222 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from 4-methyl-2-pentylthiazole-5-carboxylic acid. LCMS (Method 5-95 AB, ESI): t_R = 0.643, [M + H]⁺ = 833.5; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.49 (brs, 2H, HCOOH), 7.33 (d, J=8.0 Hz, 1H), 7.27 (d, J=8.0 Hz, 1H), 7.19 (d, J=8.0 Hz, 1H), 7.11 (d, J=8.0 Hz, 1H), 6.91 (brs, 1H), 6.84 (brs, 1H), 6.34 (s, 1H), 5.12-5.09 (m, 1H), 4.82-4.81 (m, 2H), 4.28-4.20 (m, 4H), 4.21 (s, 2H), 3.35-3.25 (m, 1H), 3.22-3.09 (m, 7H), 3.03 (t, J=7.2 Hz, 2H), 2.90 (s, 3H), 2.64 (s, 3H), 2.31-2.25 (m, 1H), 2.15-2.11 (m, 1H), 1.84-1.80 (m, 2H), 1.42-1.37 (m, 7H), 0.98 (t, J=6.8 Hz, 3H).

Example 129: Synthesis of Compound 223



[00576] Compound 223 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): $t_R = 0.760$ min, $[M+H]^+ = 869.6$.


Example 130: Synthesis of Compound 224



5

[00577] Compound 224 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): $t_R = 0.758$ min, $[M+H]^+ = 868.4$.


Example 131: Synthesis of Compound 225



10

[00578] Compound 225 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): $t_R = 0.753$ min, $[M+H]^+ = 869.5$.



Example 132: Synthesis of Compound 226



15

[00579] Compound 226 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206). LCMS (Method 5-95 AB, ESI): $t_R = 0.753$ min, $[M+H]^+ = 908.4$.


Example 133: Synthesis of Compound 227




[00580] A solution of methyl 6-bromonicotinate (500 mg, 2.3 mmol), n-butyl boronic acid (708 mg, 6.9 mmol), Pd₂(dba)₃ (212 mg, 0.23 mmol), RuPhos (108 mg, 0.23 mmol) and K₃PO₄ (1.47 g, 6.9 mmol) was stirred for under N₂ at 100 °C overnight. The volatiles were removed and the resulting residue was purified by silica gel flash column to afford methyl 6-butylnicotinate (330 mg, 73.8% yield) as colorless oil. LCMS (Method 5-95 AB, ESI): $t_R = 0.733$, $[M + H]^+ = 194.0$.

[00581] Compound 227 (formic acid salt) was prepared as a white solid utilizing the methods in Example 112 (Compound 206) from methyl 6-butylnicotinate. LCMS (0-60 AB, 2 min, ESI): $t_R = 0.828$, $[M + H]^+ = 799.5$.

Example 134: Synthesis of Compound 228



[00582] Compound 228 (formic acid salt) was prepared as a white solid utilizing the methods in Example 133 (Compound 227). LCMS (Method 5-95 AB, ESI): $t_R = 0.566$ min, $[M+H]^+ = 856.6$.

Example 135: Synthesis of Compound 229


[00583] Methyl 2-amino-4-pentylbenzoate, a yellow oil, was prepared using the methods from Example 90 (Compound 184) from 2-amino-4-bromobenzoic acid. ^1H NMR (400 MHz, CDCl_3) δ 7.76 (d, $J = 8.8$ Hz, 1H), 6.56 – 6.43 (m, 2H), 5.66 (br s, 2H), 3.86 (s, 3H), 2.51 (t, $J = 7.8$ Hz, 2H), 1.62 – 1.57 (m, 2H), 1.33 – 1.30 (m, 4H), 0.89 (t, $J = 6.6$ Hz, 3H).

[00584] Step 1: To a solution of hydrobromic acid (1.08 mL, 37.08 mmol) in water (4 mL) was added methyl 2-amino-4-pentylbenzoate (400.0 mg, 1.8075 mmol). The solution was cooled to 0°C and sodium nitrite (249.4 mg, 3.62 mmol) in water (0.1 mL) was added dropwise. The mixture was stirred at 0°C until no gas emission, then heated at 80°C for 16 h, and extracted with dichloromethane (40 mL x 2). The combined organic layers were washed with saturated NaHCO_3 (40 mL) and brine (40 mL), dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified by prep-TLC (10% EtOAc in petroleum ether, $R_f = 0.6$) to give methyl 2-bromo-4-pentylbenzoate (90 mg, 17.5% yield) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 7.74 (d, $J = 8.0$ Hz, 1H), 7.49 (s, 1H), 7.16 (d, $J = 8.0$ Hz, 1H), 3.92 (s, 3H), 2.61 (t, $J = 7.8$ Hz, 2H), 1.67 – 1.59 (m, 2H), 1.36 – 1.27 (m, 4H), 0.90 (t, $J = 7.0$ Hz, 3H).

[00585] Steps 2 and 3: Following similar procedures as described in Example 120 (Compound 214), 2-cyclopropyl-4-pentylbenzoic acid (40 mg, 94% yield) was obtained as a white solid. ^1H NMR (400 MHz, CDCl_3) δ 7.90 (d, $J = 8.0$ Hz, 1H), 7.05 (d, $J = 8.0$ Hz, 1H), 6.83 (s, 1H), 2.60 (t, $J = 7.6$ Hz, 2H), 1.63 – 1.57 (m, 3H), 1.35 – 1.29 (m, 4H), 1.06 – 1.00 (m, 2H), 0.90 (t, $J = 6.6$ Hz, 3H), 0.75 – 0.69 (m, 2H).

[00586] Compound 229 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.733$ min, $[M + H]^+ = 852.5$.

Example 136: Synthesis of Compound 230




[00587] Step 1: General Method 4 (Example 5) was applied to Compound 101-J (200 mg, 0.22 mmol) and 4-bromo-2,6-dimethylbenzoic acid (99 mg, 0.44 mmol) to afford Compound 230-A (186 mg, 76% yields) as a white solid.

10 **[00588]** Step 2: General Method Alkyl Boronic Acid was applied to Compound 230-A (186 mg, 0.16 mmol) and n-butyl boronic acid (67 mg, 0.65 mmol) to afford Compound 230-B (160 mg, 88% yield) as a pale-yellow solid.

[00589] Compound 230 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from Compound 230-B. LCMS (Method 5-95 AB, ESI): $t_R =$
 15 0.556, $[M + H]^+ = 840.4$. ^1H NMR (400 MHz, MeOH- d_4) δ 8.48 (brs, 2H, HCOOH), 7.28-7.22 (m, 2H), 7.19 (d, $J=8.0$ Hz, 1H), 7.11 (d, $J=8.0$ Hz, 1H), 6.89 (brs, 2H), 6.86 (brs, 1H), 6.73 (brs, 1H), 6.53 (s, 1H), 5.10-5.08 (m, 1H), 4.89-4.78 (m, 2H), 4.29-4.23 (m, 4H), 4.22-4.20 (m, 2H), 3.35-3.33 (m, 1H), 3.28-3.20 (m, 4H), 3.15-3.08 (m, 2H), 3.02 (s, 3H), 2.98 (t, $J=7.2$ Hz,


2H), 2.53-2.46 (m, 2H), 2.26 (s, 6H), 1.99-1.94 (m, 1H), 1.90-1.78 (m, 3H), 1.57-1.49 (m, 2H), 1.37-1.27 (m, 2H), 1.36 (d, $J=7.2$ Hz, 3H), 0.92 (t, $J=6.8$ Hz, 3H).

Example 137: Synthesis of Compound 231




- 5 [00527] Compound 231 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from (S)-2-(((benzyloxy)carbonyl)amino)-3-(tert-butoxy)propanoic acid and dodecanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.653$, $[M + H]^+ = 807.3$. LCMS (Method 5-95 AB, ESI): $t_R = 0.653$, $[M + H]^+ = 807.3$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.48 (brs, 2H, HCOOH), 7.28-7.23 (m, 2H), 7.15 (d, $J=8.0$ Hz, 1H), 7.09 (d, $J=8.0$ Hz, 1H), 6.87 (brs, 1H), 6.81 (brs, 1H), 6.35 (s, 1H), 5.00-4.75 (m, 3H), 4.25-4.15 (m, 4H), 4.19 (s, 2H), 3.89-3.84 (m, 1H), 3.72-3.67 (m, 1H), 3.40-3.25 (m, 1H), 3.18-3.09 (m, 5H), 2.92 (s, 3H), 2.30-2.26 (m, 2H), 1.65-1.52 (m, 2H), 1.40-1.30 (m, 19H), 0.90 (t, $J=6.4$ Hz, 3H).
- 10

Example 138: Synthesis of Compound 232





- 15 [00590] Compound 232 (formic acid salt) was prepared as a white solid using the methods in Example 137 (Compound 231) from decanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.747$, $[M + H]^+ = 779.4$.

Example 139: Synthesis of Compound 233

[00591] Compound 233 (formic acid salt) was prepared as a white solid (17.3 mg) utilizing the methods in Example 137 (Compound 231) from 4'-chloro-[1,1'-biphenyl]-4-carboxylic acid.

- 5 LCMS (Method 5-95 AB, ESI): $t_R = 0.616$, $[M + H]^+ = 839.2$. ^1H NMR (400 MHz, MeOH- d_4) δ 8.49 (brs, 2H, HCOOH), 7.96 (d, $J = 8.4$ Hz, 2H), 7.72 (d, $J = 8.4$ Hz, 4H), 7.47 (d, $J = 8.4$ Hz, 2H), 7.35-7.20 (m, 2H), 7.17-7.06 (m, 2H), 6.89 (s, 1H), 6.79 (s, 1H), 6.48 (s, 1H), 5.20-5.10 (m, 1H), 4.80-4.76 (m, 2H), 4.32-4.13 (m, 4H), 4.17 (s, 2H), 4.06-3.98 (m, 1H), 3.94-3.86 (m, 1H), 3.37-3.33 (m, 1H), 3.26-3.08 (m, 5H), 3.00 (s, 3H), 1.36 (d, $J = 4.8$ Hz, 3H).

10 **Example 140: Synthesis of Compound 234**




- [00592]** Step 1: A mixture of methyl 2-chloropyrimidine-5-carboxylate (2.0 g, 12.9 mmol) and (4-chlorophenyl)boronic acid (2.0 g, 12.9 mmol), Pd(PPh₃)₂Cl₂ (752 mg, 1.07 mmol) and
- 15 Na₂CO₃ (1.1 g, 10.7 mmol) in DME (50 mL) was stirred at 80 °C for 24 h. Water (250 mL) was added into the reaction, which was extracted with EtOAc (100 mL x 3). The combined organic layers were combined, concentrated and the residue was purified by chromatography on silica to give methyl 2-(4-chlorophenyl)pyrimidine-5-carboxylate (250 mg, 8.9% yield) as a white solid. (Typical Suzuki coupling condition)

[00593] Step 2: To a solution of THF (15 mL) and H₂O (4 mL) was added methyl 2-(4-chlorophenyl)pyrimidine-5-carboxylate (250 mg, 1.0 mmol) and NaOH (103 mg, 2.6 mmol). The mixture was stirred at 80°C overnight. The mixture was concentrated and the residue was adjusted to pH~5 with 2N HCl solution. The resulting precipitate was collected, which was washed with water (20 mL) to give 2-(4-chlorophenyl)pyrimidine-5-carboxylic acid (200 mg, 85% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 0.839$, $[M + H]^+ = 234.7$ (Typical ester hydrolysis procedure, NaOH/THF/H₂O)


[00594] Compound 234 (formic acid salt) was prepared as a white solid utilizing the methods in Example 137 (Compound 231) from 2-(4-chlorophenyl)pyrimidine-5-carboxylic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.698$, $[M + H]^+ = 841.5$.


Example 141: Synthesis of Compound 235



[00595] Compound 235 (formic acid salt) was prepared as a white solid utilizing the methods in Example 137 (Compound 231) except (S)-2-(((benzyloxy)carbonyl)amino)-3-(tert-butoxy)propanoic acid is used in the amino acid coupling step. LCMS (Method 5-95 AB, ESI): $t_R = 0.805$ min, $[M + H]^+ = 856.5$.

Example 142: Synthesis of Compound 236





[00596] Step 1: To a solution of (S)-4-amino-2-(((benzyloxy)carbonyl)amino)-4-oxobutanoic acid (75 mg, 0.28 mmol) in THF (15 mL) at 0°C, DEPBT (252 mg, 0.84 mmol) and NaHCO₃ (71 mg, 0.84 mmol) was added. The mixture and stirred at the same temperature for 10 min, followed by the addition of Compound 101-G (100 mg, 0.14 mmol). The resulting mixture was stirred at room temperature for 4 days. The volatiles were removed and the resulting residue was taken up in EtOAc (50 mL), which was washed with brine (50 mL x 2). The EtOAc layer was dried over Na₂SO₄, concentrated and the residue was purified by flash column chromatography to afford Compound 236-A (95mg, 71% yield) as a white solid.


[00597] Step 2: General Method Hydrogenation (Pd/C, H₂) was applied to Compound 236-A (95 mg, 0.10 mmol) to afford Compound 236-B (84 mg, 99% yield) as a white solid.

[00598] Step 3: The coupling of an amine to an acid chloride is described and is referred to as General Method Acid Chloride. To a solution of Compound 236-B (84 mg, 0.10 mmol) in DCM (5 mL) was added decanoyl chloride (23 mg, 0.12 mmol) and Et₃N (20 mg, 0.20 mmol) at 0°C. The mixture was stirred at room temperature for 1h. The reaction was adjusted pH to 5 with saturated citric acid solution, which was extracted with DCM (10 mL x 3). The combined organic layers were washed with brine (20 mL). The DCM layer was dried over Na₂SO₄, concentrated and the residue was purified by prep-TLC to give Compound 236-C (87 mg, 90% yield) as a white solid.

[00599] Compound 236 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from Compound 236-C. LCMS (Method 5-95 AB, ESI): t_R = 0.896, [M + H]⁺ = 806.8. ¹H NMR (400 MHz, MeOH-d₄) δ 7.28-7.22 (m, 2H), 7.14-7.06 (m, 2H), 6.87 (brs, 1H), 6.81 (brs, 1H), 6.33 (s, 1H), 5.22-5.18 (m, 1H), 4.80-4.70 (m, 2H), 4.21 (brs, 2H), 4.12-4.09 (m, 4H), 3.19-3.05 (m, 2H), 3.00 (brs, 4H), 2.92 (s, 3H), 2.76-2.65 (m, 1H),


2.61-2.50 (m, 1H), 2.26-2.15 (m, 2H), 1.64 (brs, 2H), 1.36-1.33 (m, 15H), 0.92 (t, $J=7.2$ Hz, 3H).

Example 143: Synthesis of Compound 237




- 5 **[00600]** Compound 237 (formic acid salt) was prepared as a white solid utilizing the methods in Example 142 (Compound 236) from Compound 236-B. LCMS (Method 5-95 AB, ESI): $t_R = 0.631$, $[M + H]^+ = 854.3$. ^1H NMR (400 MHz, MeOH- d_4) δ 8.51 (brs, 2H, HCOOH), 7.35-7.28 (m, 2H), 7.25 (d, $J=8.0$ Hz, 1H), 7.16 (d, $J=8.0$ Hz, 1H), 7.12-7.03 (m, 3H), 6.90 (brs, 1H), 6.80 (brs, 1H), 6.38 (s, 1H), 5.38-5.33 (m, 1H), 4.81-4.74 (m, 2H), 4.25-4.12 (m, 4H), 4.19 (s, 2H),
- 10 3.20-3.06 (m, 4H), 2.99 (s, 3H), 2.83-2.76 (m, 2H), 2.73-2.64 (m, 2H), 2.63-2.57 (m, 2H), 2.39 (s, 3H), 1.66-1.54 (m, 2H), 1.40-1.24 (m, 9H), 0.90 (t, $J=5.2$ Hz, 3H).


Example 144: Synthesis of Compound 238




- 15 **[00601]** Compound 238 (formic acid salt) was prepared as a white solid utilizing the methods in Example 142 (Compound 236) from Compound 236-B. LCMS (Method 5-95 AB, ESI): $t_R = 0.587$, $[M + H]^+ = 826.4$.

Example 145: Synthesis of Compound 239

[00602] Compound 239 (free base) was prepared as a white solid utilizing the methods in Example 142 (Compound 236). LCMS (Method 5-95 AB, ESI): $t_R = 0.813$, $[M + H]^+ = 882.7$.

5 Example 146: Synthesis of Compound 240


[00603] Compound 240 (free base) was prepared as a white solid utilizing the methods in Example 142 (Compound 236). LCMS (Method 5-95 AB, ESI): $t_R = 0.782$, $[M + H]^+ = 902.5$.

Example 147: Compound 241

[00604] Compound 241 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from Compound 101-G and (S)-2-(((benzyloxy)carbonyl)amino)propanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.679$, $[M + H]^+ = 791.3$. ^1H NMR (400 MHz, MeOH- d_4) δ 8.47 (brs, 2H, HCOOH), 7.30 (d, $J=8.0$ Hz, 1H), 7.24 (d, $J=8.0$ Hz, 1H), 7.16 (d, $J=8.4$ Hz, 1H), 7.09 (d, $J=8.4$ Hz, 1H), 6.89 (brs, 1H), 6.80 (brs,

1H), 6.35 (s, 1H), 4.79-4.72 (m, 3H), 4.30-4.15 (m, 4H), 4.19 (s, 2H), 3.40-3.34 (m, 1H), 3.24-3.09 (m, 5H), 2.88 (s, 3H), 2.25 (t, $J=7.6$ Hz, 2H), 1.69-1.53 (m, 3H), 1.50-1.22 (m, 21H), 0.90 (t, $J=6.6$ Hz, 3H).


Example 148: Synthesis of Compound 242



5

[00605] Compound 242 (formic acid salt) was prepared as a white solid utilizing the methods in Example 147 (Compound 241). LCMS (Method 5-95 AB, ESI): $t_R = 0.616$, $[M + H]^+ = 783.3$.


Example 149: Synthesis of Compound 243



10

[00606] Compound 243 (formic acid salt) was prepared in 29% yield as a white solid utilizing the methods in Example 7 (Compound 101) from (S)-2-(((benzyloxy)carbonyl)amino)propanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.707$ min, $[M + H]^+ = 839.3$.


Example 150: Synthesis of Compound 244



15


[00607] Compound 244 (formic acid salt) was prepared as a white solid utilizing the methods used in Example 7 (Compound 101) from Compound 101-G and (S)-5-amino-2-(((benzyloxy)carbonyl)amino)-5-oxopentanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.782$, $[M + H]^+ = 848.6$. ^1H NMR (400 MHz, MeOH- d_4) δ 8.53 (brs, 2H, HCOOH), 7.31-7.26 (m, 2H), 7.18 (d, $J=8.0$ Hz, 1H), 7.12 (d, $J=8.0$ Hz, 1H), 6.90 (brs, 1H), 6.84 (brs, 1H), 6.36 (s, 1H), 5.00-4.76 (m, 3H), 4.30-4.20 (m, 4H), 4.21 (s, 2H), 3.40-3.22 (m, 5H), 3.20-3.10 (m, 1H), 2.94 (s, 3H), 2.39-2.36 (m, 2H), 2.29-2.27 (m, 2H), 2.20-2.05 (m, 1H), 1.93-1.82 (m, 1H), 1.70-1.54 (m, 2H), 1.38-1.33 (m, 19H), 0.92 (t, $J=7.2$ Hz, 3H).

Example 151: Synthesis of Compound 245




[00608] Compound 245 (formic acid salt) was prepared as a white solid utilizing the same methods used in Example 150 (Compound 244). LCMS (Method 5-95 AB, ESI): $t_R = 0.732$, $[M + H]^+ = 840.5$.


Example 152: Synthesis of Compound 246




[00609] Compound 246 (formic acid salt) was prepared as a white solid utilizing the methods in Example 150 (Compound 244). LCMS (Method 5-95 AB, ESI): $t_R = 0.637$, $[M + H]^+ = 868.9$.

Example 153: Synthesis of Compound 247



[00610] Compound 247 (formic acid salt) was prepared as a white solid utilizing the methods in Example 150 (Compound 244). LCMS (Method 5-95 AB, ESI): $t_R = 0.835$, $[M + H]^+ =$
 5 896.6.

Example 154: Synthesis of Compound 248


[00611] Compound 248 (formic acid salt) was prepared as a white solid utilizing the methods in Example 150 (Compound 244). LCMS (Method 5-95 AB, ESI): $t_R = 0.786$, $[M + H]^+ =$
 10 916.5.

Example 155: Synthesis of Compound 249

[00612] Compound 249 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) except (S)-2-(((benzyloxy)carbonyl)amino)-3-hydroxypropanoic acid is used in the amino acid coupling step. LCMS (Method 5-95 AB, ESI): $t_R = 0.659$ min, $[M + H]^+ = 861.8$.
 15




[00615] Compound 250 (formic acid salt) was prepared as a white solid utilizing the same methods used in the preparation of Compound 231 (Example 137) except 6-(4-chlorophenyl)nicotinic acid was used in the coupling step. LCMS (Method 5-95 AB, ESI): $t_R = 0.640$, $[M + H]^+ = 840.0$.

Example 157 Synthesis of Compound 251


[00616] Compound 251 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from (R)-2-(((benzyloxy)carbonyl)amino)propanoic acid.

5 LCMS (Method 5-95 AB, ESI): $t_R = 0.851$ min, $[M + H]^+ = 839.5$.

Example 158: Synthesis of Compound 252


[00617] Compound 252 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from 2-(((benzyloxy)carbonyl)amino)acetic acid. LCMS

10 (Method 5-95 AB, ESI): $t_R = 0.723$ min, $[M + H]^+ = 825.2$.

Example 159: Synthesis of Compound 253

[00618] Compound 253 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from (S)-2-(((benzyloxy)carbonyl)amino)-5-(tert-butoxy)-5-

15 oxopentanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.799$, $[M + H]^+ = 849.4$



Example 160: Synthesis of Compound 254 and Compound 255

- 5 **[00619]** Step 1: Compound 254-A was prepared according to General Method 4 (Example 5) from Compound 101-G (104 mg, 0.28 mmol) and Fmoc-L-methionine to afford Compound 254-A (100 mg, 66.9%) as a light yellow solid after silica column chromatography.
- 10 **[00620]** Step 2: Compound 254-A was subject to the standard Fmoc-deprotection conditions. To a solution of Compound 254-A (100 mg, 0.18 mmol) in DCM (4 mL) was added piperidine (1.0 mL) at 0°C and the mixture was stirred at room temperature under N₂ for 2 h. The volatiles were concentrated and the residue was re-dissolved in EtOAc (30 mL), which was washed by brine (2 x 30 mL). The organic layer was dried over MgSO₄, concentrated and purified by prep-TLC (eluting 3% MeOH in DCM) to give Compound 254-B (50 mg, 63.1% yield) as a yellow solid.
- 15 **[00621]** Steps 3 and 4: Compound 254-C was prepared as previously described from decanoyl chloride using triethylamine as a base in DCM followed by LiOH ester hydrolysis.

[00623] Compound 255 (2 mg, white solid) was isolated as a by-product which occurs from oxidation of the sulfide during the preparation of Compound 254. Data for Compound 255: LCMS (Method 5-95 AB, ESI): $t_R = 0.653$, $[M + H]^+ = 839.2$.


CCCCCCCCCCCC(=O)NC(=O)[C@H](O)[C@@H](NC(=O)N[C@@H](C)C(=O)NC(=O)N[C@@H](C)C(=O)NC(=O)C#N)C1=CC=C(C(=C1)C(=CC2=CC=C(C=C2)C(=CC3=CC=C(C=C3)OCCN)OCCN)C4=CC=C(C=C4)C(=CC5=CC=C(C=C5)C(=CC6=CC=C(C=C6)C(=CC7=CC=C(C=C7)C(=CC8=CC=C(C=C8)C(=CC9=CC=C(C=C9)C(=CC10=CC=C(C=C10)C(=CC11=CC=C(C=C11)C(=CC12=CC=C(C=C12)C(=CC13=CC=C(C=C13)C(=CC14=CC=C(C=C14)C(=CC15=CC=C(C=C15)C(=CC16=CC=C(C=C16)C(=CC17=CC=C(C=C17)C(=CC18=CC=C(C=C18)C(=CC19=CC=C(C=C19)C(=CC20=CC=C(C=C20)C(=CC21=CC=C(C=C21)C(=CC22=CC=C(C=C22)C(=CC23=CC=C(C=C23)C(=CC24=CC=C(C=C24)C(=CC25=CC=C(C=C25)C(=CC26=CC=C(C=C26)C(=CC27=CC=C(C=C27)C(=CC28=CC=C(C=C28)C(=CC29=CC=C(C=C29)C(=CC30=CC=C(C=C30)C(=CC31=CC=C(C=C31)C(=CC32=CC=C(C=C32)C(=CC33=CC=C(C=C33)C(=CC34=CC=C(C=C34)C(=CC35=CC=C(C=C35)C(=CC36=CC=C(C=C36)C(=CC37=CC=C(C=C37)C(=CC38=CC=C(C=C38)C(=CC39=CC=C(C=C39)C(=CC40=CC=C(C=C40)C(=CC41=CC=C(C=C41)C(=CC42=CC=C(C=C42)C(=CC43=CC=C(C=C43)C(=CC44=CC=C(C=C44)C(=CC45=CC=C(C=C45)C(=CC46=CC=C(C=C46)C(=CC47=CC=C(C=C47)C(=CC48=CC=C(C=C48)C(=CC49=CC=C(C=C49)C(=CC50=CC=C(C=C50)C(=CC51=CC=C(C=C51)C(=CC52=CC=C(C=C52)C(=CC53=CC=C(C=C53)C(=CC54=CC=C(C=C54)C(=CC55=CC=C(C=C55)C(=CC56=CC=C(C=C56)C(=CC57=CC=C(C=C57)C(=CC58=CC=C(C=C58)C(=CC59=CC=C(C=C59)C(=CC60=CC=C(C=C60)C(=CC61=CC=C(C=C61)C(=CC62=CC=C(C=C62)C(=CC63=CC=C(C=C63)C(=CC64=CC=C(C=C64)C(=CC65=CC=C(C=C65)C(=CC66=CC=C(C=C66)C(=CC67=CC=C(C=C67)C(=CC68=CC=C(C=C68)C(=CC69=CC=C(C=C69)C(=CC70=CC=C(C=C70)C(=CC71=CC=C(C=C71)C(=CC72=CC=C(C=C72)C(=CC73=CC=C(C=C73)C(=CC74=CC=C(C=C74)C(=CC75=CC=C(C=C75)C(=CC76=CC=C(C=C76)C(=CC77=CC=C(C=C77)C(=CC78=CC=C(C=C78)C(=CC79=CC=C(C=C79)C(=CC80=CC=C(C=C80)C(=CC81=CC=C(C=C81)C(=CC82=CC=C(C=C82)C(=CC83=CC=C(C=C83)C(=CC84=CC=C(C=C84)C(=CC85=CC=C(C=C85)C(=CC86=CC=C(C=C86)C(=CC87=CC=C(C=C87)C(=CC88=CC=C(C=C88)C(=CC89=CC=C(C=C89)C(=CC90=CC=C(C=C90)C(=CC91=CC=C(C=C91)C(=CC92=CC=C(C=C92)C(=CC93=CC=C(C=C93)C(=CC94=CC=C(C=C94)C(=CC95=CC=C(C=C95)C(=CC96=CC=C(C=C96)C(=CC97=CC=C(C=C97)C(=CC98=CC=C(C=C98)C(=CC99=CC=C(C=C99)C(=CC100=CC=C(C=C100)C(=CC101=CC=C(C=C101)C(=CC102=CC=C(C=C102)C(=CC103=CC=C(C=C103)C(=CC104=CC=C(C=C104)C(=CC105=CC=C(C=C105)C(=CC106=CC=C(C=C106)C(=CC107=CC=C(C=C107)C(=CC108=CC=C(C=C108)C(=CC109=CC=C(C=C109)C(=CC110=CC=C(C=C110)C(=CC111=CC=C(C=C111)C(=CC112=CC=C(C=C112)C(=CC113=CC=C(C=C113)C(=CC114=CC=C(C=C114)C(=CC115=CC=C(C=C115)C(=CC116=CC=C(C=C116)C(=CC117=CC=C(C=C117)C(=CC118=CC=C(C=C118)C(=CC119=CC=C(C=C119)C(=CC120=CC=C(C=C120)C(=CC121=CC=C(C=C121)C(=CC122=CC=C(C=C122)C(=CC123=CC=C(C=C123)C(=CC124=CC=C(C=C124)C(=CC125=CC=C(C=C125)C(=CC126=CC=C(C=C126)C(=CC127=CC=C(C=C127)C(=CC128=CC=C(C=C128)C(=CC129=CC=C(C=C129)C(=CC130=CC=C(C=C130)C(=CC131=CC=C(C=C131)C(=CC132=CC=C(C=C132)C(=CC133=CC=C(C=C133)C(=CC134=CC=C(C=C134)C(=CC135=CC=C(C=C135)C(=CC136=CC=C(C=C136)C(=CC137=CC=C(C=C137)C(=CC138=CC=C(C=C138)C(=CC139=CC=C(C=C139)C(=CC140=CC=C(C=C140)C(=CC141=CC=C(C=C141)C(=CC142=CC=C(C=C142)C(=CC143=CC=C(C=C143)C(=CC144=CC=C(C=C144)C(=CC145=CC=C(C=C145)C(=CC146=CC=C(C=C146)C(=CC147=CC=C(C=C147)C(=CC148=CC=C(C=C148)C(=CC149=CC=C(C=C149)C(=CC150=CC=C(C=C150)C(=CC151=CC=C(C=C151)C(=CC152=CC=C(C=C152)C(=CC153=CC=C(C=C153)C(=CC154=CC=C(C=C154)C(=CC155=CC=C(C=C155)C(=CC156=CC=C(C=C156)C(=CC157=CC=C(C=C157)C(=CC158=CC=C(C=C158)C(=CC159=CC=C(C=C159)C(=CC160=CC=C(C=C160)C(=CC161=CC=C(C=C161)C(=CC162=CC=C(C=C162)C(=CC163=CC=C(C=C163)C(=CC164=CC=C(C=C164)C(=CC165=CC=C(C=C165)C(=CC166=CC=C(C=C166)C(=CC167=CC=C(C=C167)C(=CC168=CC=C(C=C168)C(=CC169=CC=C(C=C169)C(=CC170=CC=C(C=C170)C(=CC171=CC=C(C=C171)C(=CC172=CC=C(C=C172)C(=CC173=CC=C(C=C173)C(=CC174=CC=C(C=C174)C(=CC175=CC=C(C=C175)C(=CC176=CC=C(C=C176)C(=CC177=CC=C(C=C177)C(=CC178=CC=C(C=C178)C(=CC179=CC=C(C=C179)C(=CC180=CC=C(C=C180)C(=CC181=CC=C(C=C181)C(=CC182=CC=C(C=C182)C(=CC183=CC=C(C=C183)C(=CC184=CC=C(C=C184)C(=CC185=CC=C(C=C185)C(=CC186=CC=C(C=C186)C(=CC187=CC=C(C=C187)C(=CC188=CC=C(C=C188)C(=CC189=CC=C(C=C189)C(=CC190=CC=C(C=C190)C(=CC191=CC=C(C=C191)C(=CC192=CC=C(C=C192)C(=CC193=CC=C(C=C193)C(=CC194=CC=C(C=C194)C(=CC195=CC=C(C=C195)C(=CC196=CC=C(C=C196)C(=CC197=CC=C(C=C197)C(=CC198=CC=C(C=C198)C(=CC199=CC=C(C=C199)C(=CC200=CC=C(C=C200)C(=CC201=CC=C(C=C201)C(=CC202=CC=C(C=C202)C(=CC203=CC=C(C=C203)C(=CC204=CC=C(C=C204)C(=CC205=CC=C(C=C205)C(=CC206=CC=C(C=C206)C(=CC207=CC=C(C=C207)C(=CC208=CC=C(C=C208)C(=CC209=CC=C(C=C209)C(=CC210=CC=C(C=C210)C(=CC211=CC=C(C=C211)C(=CC212=CC=C(C=C212)C(=CC213=CC=C(C=C213)C(=CC214=CC=C(C=C214)C(=CC215=CC=C(C=C215)C(=CC216=CC=C(C=C216)C(=CC217=CC=C(C=C217)C(=CC218=CC=C(C=C218)C(=CC219=CC=C(C=C219)C(=CC220=CC=C(C=C220)C(=CC221=CC=C(C=C221)C(=CC222=CC=C(C=C222)C(=CC223=CC=C(C=C223)C(=CC224=CC=C(C=C224)C(=CC225=CC=C(C=C225)C(=CC226=CC=C(C=C226)C(=CC227=CC=C(C=C227)C(=CC228=CC=C(C=C228)C(=CC229=CC=C(C=C229)C(=CC230=CC=C(C=C230)C(=CC231=CC=C(C=C231)C(=CC232=CC=C(C=C232)C(=CC233=CC=C(C=C233)C(=CC234=CC=C(C=C234)C(=CC235=CC=C(C=C235)C(=CC236=CC=C(C=C236)C(=CC237=CC=C(C=C237)C(=CC238=CC=C(C=C238)C(=CC239=CC=C(C=C239)C(=CC240=CC=C(C=C240)C(=CC241=CC=C(C=C241)C(=CC242=CC=C(C=C242)C(=CC243=CC=C(C=C243)C(=CC244=CC=C(C=C244)C(=CC245=CC=C(C=C245)C(=CC246=CC=C(C=C246)C(=CC247=CC=C(C=C247)C(=CC248=CC=C(C=C248)C(=CC249=CC=C(C=C249)C(=CC250=CC=C(C=C250)C(=CC251=CC=C(C=C251)C(=CC252=CC=C(C=C252)C(=CC253=CC=C(C=C253)C(=CC254=CC=C(C=C254)C(=CC255=CC=C(C=C255)C(=CC256=CC=C(C=C256)C(=CC257=CC=C(C=C257)C(=CC258=CC=C(C=C258)C(=CC259=CC=C(C=C259)C(=CC260=CC=C(C=C260)C(=CC261=CC=C(C=C261)C(=CC262=CC=C(C=C262)C(=CC263=CC=C(C=C263)C(=CC264=CC=C(C=C264)C(=CC265=CC=C(C=C265)

Example 162: Synthesis of Compound 257




-209-


Example 163: Synthesis of Compound 258




-210-

Example 164: Synthesis of Compound 259

[00629] Compound 259 (formic acid salt) was prepared as a white solid utilizing the methods in Example 162 (Compound 257). LCMS (Method 5-95 AB, ESI): $t_R = 0.598$, $[M + H]^+ = 853.5$.


Example 165: Synthesis of Compound 260

[00630] Compound 260 (free base) was prepared as a white solid utilizing the methods in Example 162 (Compound 257). LCMS (Method 5-95 AB, ESI): $t_R = 0.810$, $[M + H]^+ = 855.5$; ^1H NMR (400 MHz, MeOH- d_4) δ 7.31-7.22 (m, 2H), 7.22 (d, $J=7.5$ Hz, 1H), 7.14 (d, $J=7.5$ Hz, 1H), 7.07-7.04 (m, 3H), 6.84 (brs, 1H), 6.65-6.58 (m, 2H), 5.20-5.18 (m, 1H), 4.80-4.70 (m, 2H), 4.21 (s, 2H), 4.16-4.01 (m, 4H), 3.74 (t, $J=5.6$ Hz, 2H), 3.32-3.28 (m, 1H), 3.15-2.91 (m, 5H), 3.02 (s, 3H), 2.61-2.52 (m, 2H), 2.36 (s, 3H), 2.16-2.05 (m, 1H), 2.00-1.90 (m, 1H), 1.57 (brs, 2H), 1.40-1.20 (m, 11H), 0.92 (t, $J=6.4$ Hz, 3H).

Example 166: Synthesis of Compound 261

[00631] Compound 261 (formic acid salt) was prepared as a white solid utilizing the methods in Example 162 (Compound 257). LCMS (Method 5-95 AB, ESI): $t_R = 0.794$, $[M + H]^+ = 889.6$.


Example 167: Synthesis of Compound 262




5

[00632] Compound 262 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.684$ min, $[M + H]^+ = 869.4$.

Example 168: Synthesis of Compound 263



10



[00633] Compound 263-A is an intermediate in the preparation of Compound 262 (Example 167). To a solution of Compound 263-A (360.0 mg, 0.31 mmol) in tetrahydrofuran (8 mL) was added tetrabutylammonium fluoride (2M; 2.0 mL, 4 mmol) in tetrahydrofuran. The solution was stirred at 20 °C for 2 h and evaporated to dryness. The residue was taken up in EtOAc (20 mL), washed with water (20 mL x 2) and brine (20 mL), dried over MgSO₄ and concentrated. The residue was purified by flash column chromatography (5% methanol in DCM, R_f = 0.5) to afford Compound 263-B (320 mg, 98.6% yield) as a white solid. LCMS (Method 5-95 AB, ESI): t_R = 1.122 min, [M + H]⁺ = 1045.4.


[00634] To a solution of Compound 263-B (170.0 mg, 0.16 mmol) in dichloromethane (10 mL) was added Dess-Martin periodinane (138.0 mg, 0.33 mmol). The mixture was stirred at 20 °C for 2 h, quenched with saturate NaHCO₃ solution and extracted with EtOAc (20 mL). The organic layer was washed with water (20 mL x 2) and brine (20 mL), dried over MgSO₄ and concentrated to afford crude Compound 263-C (160 mg, 94.3% yield) as a white solid. LCMS (Method 5-95 AB, ESI): t_R = 0.974 min, [M + H]⁺ = 1043.5.

[00635] To a solution of Compound 263-C (400.0 mg, 0.38 mmol) and tert-butyl 2-aminoacetate hydrochloride (128.5 mg, 0.77 mmol) in methanol (20 mL) was added sodium cyanoborohydride (48.2 mg, 0.77 mmol). The reaction was stirred at 20 °C for 2 h and evaporated to dryness. The residue was taken up in EtOAc (10 mL), washed with water (10 mL x 2) and brine (10 mL), dried over MgSO₄ and concentrated. The residue was purified by flash column chromatography (5% methanol in DCM, R_f = 0.5) to afford Compound 263-D (300 mg,

67.5% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 1.043$ min, $[M + H]^+ = 1159.3$.

[00636] Compound 263 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from Compound 263-D. LCMS (Method 5-95 AB, ESI): $t_R =$
5 0.642 min, $[M + H]^+ = 927.4$.

Example 169: Synthesis of Compound 264




[00637] Step 1: To a solution of (S)-4-(((benzyloxy)carbonyl)amino)-5-methoxy-5-oxopentanoic acid (1.0 g, 3.4 mmol) in THF (3.5 mL) at 0°C was added 1M BH_3/THF (6.7 mL, 6.7 mmol) and the mixture was stirred at the same temperature for 1 h. The reaction was quenched by addition of 1M $NaHSO_4$ (10 mL), followed by the removal of most THF under reduced pressure. The resulting mixture was diluted with water (20 mL), which was extracted by EtOAc (30 mL x 3). The combined organic layers were washed with brine (90 mL), dried over
15 Na_2SO_4 , concentrated and the residue was purified by silica gel flash column to afford (S)-methyl 2-(((benzyloxy)carbonyl)amino)-5-hydroxypentanoate (550 mg, 60% yield) as colorless oil. LCMS (Method 5-95 AB, ESI): $t_R = 0.729$, $M+Na^+ = 303.9$

[00638] Steps 2 and 3: Typical TBS protection and ester hydrolysis (LiOH) procedure was followed to afford (S)-2-(((benzyloxy)carbonyl)amino)-5-((tert-butyl-
20 dimethylsilyl)oxy)pentanoic acid (640 mg) as a colorless oil.

[00639] Compound 264 (formic acid salt) was prepared as a white solid utilizing the methods in Example 162 (Compound 257). LCMS (Method 5-95 AB, ESI): $t_R = 0.658$, $[M + H]^+ = 835.6$; ^1H NMR (400 MHz, $MeOH-d_4$) δ 8.49 (brs, 2H, $HCOOH$), 7.35-7.25 (m, 2H), 7.17 (d, $J=8.0$ Hz, 1H), 7.12 (d, $J=8.0$ Hz, 1H), 6.92 (brs, 1H), 6.80 (brs, 1H), 6.41 (s, 1H), 4.95-4.88 (m, 1H), 4.82-4.70 (m, 2H), 4.40-4.22 (m, 4H), 4.20 (s, 2H), 3.65-3.58 (m, 2H), 3.48-3.35 (m, 1H),
25

3.31-3.22 (m, 4H), 3.20-3.10 (m, 1H), 2.93 (s, 3H), 2.32-2.20 (m, 2H), 2.00-1.90 (m, 1H), 1.80-1.60 (m, 5H), 1.50-1.20 (m, 20H), 0.92 (t, $J=6.8$ Hz, 3H).

Example 170: Synthesis of Compound 265



[00640] Step 1: To a solution of 4-tert-amylphenol (3.0 g, 18.27 mmol) in dichloromethane (20 mL) was added dropwise bromine (0.94 mL, 18.27 mmol) at 0 °C and stirred for 30 min. Then the mixture was allowed to warm to 20 °C and stirred for 16 h. The reaction mixture was diluted with DCM (20 mL), washed with water (50 mL x 3), dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified on silica gel column eluted with ethyl acetate/petroleum ether (1:10) to afford 2-bromo-4-(1,1-dimethylpropyl)phenol (2.6 g, 58.5% yield) as a white solid. ^1H NMR (400 MHz, CDCl_3): δ 7.37 (d, $J = 2.0$ Hz, 1H), 7.16 (dd, $J = 8.4, 2.4$ Hz, 1H), 6.94 (d, $J = 8.4$ Hz, 1H), 5.35 (s, 1H), 1.58 (q, $J = 7.5$ Hz, 2H), 1.23 (s, 6H), 0.66 (t, $J = 7.4$ Hz, 3H).

[00641] Step 2: A mixture of 2-bromo-4-(1,1-dimethylpropyl)phenol (500 mg, 2.06 mmol), palladium(II) acetate (46.2 mg, 0.21 mmol), RuPhos (192.0 mg, 0.41 mmol) and potassium carbonate (710.6 mg, 5.14 mmol) in toluene (10 mL) and water (1 mL) was treated with trimethylboroxine (516.3 mg, 4.11 mmol) and heated at 80 °C for 16 hours under nitrogen. The reaction mixture was diluted with water (10 mL) and extracted with ethyl acetate (20 mL x 3). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by Prep-TLC (10% ethyl acetate in petroleum ether) to afford 4-(1,1-dimethylpropyl)-2-methylphenol (220 mg, 60% yield) as a colorless oil. ^1H NMR (400

MHz, CDCl₃): δ 7.05 - 6.95 (m, 2H), 6.69 (d, J = 8.4 Hz, 1H), 4.51 (s, 1H), 2.24 (s, 3H), 1.61 - 1.56 (m, 2H), 1.23 (s, 7H), 0.66 (t, J = 7.4 Hz, 3H).

[00642] Step 3: A solution of 4-(1,1-dimethylpropyl)-2-methyl-phenol (220.0 mg, 1.23 mmol) and pyridine (292.9 mg, 3.7 mmol) in dichloromethane (5 mL) was treated with trifluoromethanesulfonic anhydride (417.8 mg, 1.48 mmol) at 0 °C. The resulting mixture was stirred at 15°C for 1 hour under nitrogen. The reaction mixture was adjusted to pH = 5 using 2N HCl and extracted with dichloromethane (20 mL x 3). The combined organic layers were dried over sodium sulfate and filtered. The filtrate was concentrated to afford [4-(1,1-dimethylpropyl)-2-methyl-phenyl] trifluoromethanesulfonate (360 mg, 94% yield) as a brown oil, which was used in the next step without purification.


[00643] Step 4: The carbonylation of an aryl triflate is described and is referred to as General Method Carbonylation. This method can also be used with aryl bromides and iodides.

[00644] A solution of [4-(1,1-dimethylpropyl)-2-methyl-phenyl] trifluoromethanesulfonate (240 mg, 0.770 mmol) and triethylamine (234.8 mg, 2.32 mmol) in methanol (10 mL) was treated with 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (56.6 mg, 0.08 mmol). The resulting mixture was stirred at 60 °C for 16 hours under CO (35 psi). The reaction mixture was concentrated and the residue was purified with silica gel column (3% ethyl acetate in petroleum) to afford methyl 4-(1,1-dimethylpropyl)-2-methyl-benzoate (40 mg, 23.5% yield) as a colorless oil.

[00645] Step 5: 4-(1,1-Dimethylpropyl)-2-methyl-benzoate (40.0 mg, 0.18 mmol) was hydrolyzed as as previously described (General Method NaOH) to give crude 4-(1,1-dimethylpropyl)-2-methyl-benzoic acid (30 mg, 80.1% yield) as a white solid.


[00646] Compound 265 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): t_R = 0.602 min, $[M + H]^+ = 826.2$.

Example 171: Synthesis of Compound 266




[00647] Compound 266 (formic acid salt) was prepared as a white solid utilizing the methods in Example 170 (Compound 265) starting from 4-(tert-butyl)-2-methylphenol. LCMS (Method 5-95 AB, ESI): $t_R = 0.562$ min, $[M + H]^+ = 812.5$.

Example 172: Synthesis of Compound 267



5




[00648] Step 1: To a solution of 2-chloro-4-hydroxybenzoic acid (200.0 mg, 1.16 mmol) in methanol (5 mL) was added thionyl chloride (413.6 mg, 3.48 mmol) dropwise at 0 °C. The reaction mixture was stirred at 70 °C for 2 h and concentrated. The residue was diluted with water (15 mL) and extracted with EtOAc (15 mL x 2). The organic layers were combined and washed with water (30 mL x 2) and brine (20 mL). The organic layers were separated, dried over Na₂SO₄ and concentrated to obtain methyl 2-chloro-4-hydroxy-benzoate (200 mg, 92.5% yield) as a yellow solid which was used directly without further purification.

[00649] Step 2: To a solution of methyl 2-chloro-4-hydroxy-benzoate (100.0 mg, 0.54 mmol) in N,N-dimethylformamide (3 mL) was added 1-bromoheptane (2879.6 mg, 16.08 mmol) and potassium carbonate (2222.1 mg, 16.08 mmol). The mixture was stirred at 20 °C for 4 h, diluted with water (20 mL) and extracted with EtOAc (20 mL x 2). The organic layers were combined and washed with water (40 mL x 4) and brine (20 mL). The organic layers were separated, dried over Na₂SO₄ and concentrated. The residue was purified by preparative TLC (5% EtOAc in petroleum ether) to obtain methyl 2-chloro-4-heptoxy-benzoate (152 mg, 99.6% yield) as a yellow oil. For more sterically hindered alkyl halides, elevated temperatures are used.

[00650] Step 3: Methyl 2-chloro-4-heptoxy-benzoate (152.0 mg, 0.53 mmol) was hydrolyzed previously described (General Method NaOH) to give crude 2-chloro-4-heptoxy-benzoic acid (110 mg, 76.1% yield) as a yellow solid.

[00651] Compound 267 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.647$ min, $[M + H]^+ = 890.4$.


Example 173: Synthesis of Compound 268




[00652] Following similar procedures as described in Example 172 (Compound 267), 4-(heptyloxy)-2-methylbenzoic acid (605 mg, 96.1% yield) was obtained as a yellow solid from 4-hydroxy-2-methylbenzoic acid. ^1H NMR (400 MHz, $CDCl_3$) δ 8.09 - 8.04 (m, 1H), 6.79 - 6.73 (m, 2H), 4.01 (t, $J = 6.4$ Hz, 2H), 2.64 (s, 3H), 1.84 - 1.73 (m, 2H), 1.49 - 1.27 (m, 8H), 0.90 (t, $J = 7.0$ Hz, 3H).

[00653] Compound 268 (formic acid salt) was prepared as a white solid utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): $t_R = 0.608$ min, $[M + H]^+ = 870.4$. ^1H NMR (400 MHz, $MeOH-d_4$) δ 8.46 (brs, 2H), 7.38 (d, $J=8.0$ Hz, 1H), 7.32 (dd, $J=8.4$, 1.8 Hz, 1H), 7.25 (d, $J=8.0$ Hz, 1H), 7.18 (d, $J=8.0$ Hz, 1H), 7.10 (d, $J=8.0$ Hz, 1H), 6.90 (d, $J=1.8$ Hz, 1H), 6.85 - 6.77 (m, 3H), 6.35 (s, 1H), 5.14-5.10 (m, 1H), 4.82-4.76 (m, 2H), 4.36-4.18 (m, 4H), 4.20 (s, 2H), 3.99 (t, $J=6.4$ Hz, 2H), 3.35-3.20 (m, 5H), 3.17-3.08 (m, 3H), 2.91 (s, 3H), 2.40 (s, 3H), 2.30-2.24 (m, 1H), 2.20-2.05 (m, 1H), 1.81-1.74 (m, 2H), 1.53-1.31 (m, 11H), 0.92 (t, $J=6.6$ Hz, 3H).

Example 174: Synthesis of Compound 269






[00654] Following similar procedures as described in Example 172, 4-butoxy-2-chlorobenzoic acid (160 mg, 0.70 mmol) was obtained as a white solid from 2-chloro-4-hydroxybenzoic acid.

¹H NMR (400 MHz, CD₃OD): δ 7.90 (d, J = 9.2 Hz, 1H), 7.02 (d, J = 2.0 Hz, 1H), 6.91 (dd, J = 9.0, 2.6 Hz, 1H), 4.04 (t, J = 6.4 Hz, 2H), 1.80 - 1.70 (m, 2H), 1.55 - 1.45 (m, 2H), 0.99 (t, J = 7.6 Hz, 3H).

[00655] Compound 269 (formic acid salt) was prepared as a white solid utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): t_R = 0.574 min, [M + H]⁺ = 848.5.

10 **Example 175: Synthesis of Compound 270**




[00656] Following similar procedures as described in Example 172, 2-chloro-4-

(pentyloxy)benzoic acid (160 mg, 0.6593 mmol, 94% yield) was obtained as a yellow solid. ¹H

¹H NMR (400 MHz, CD₃OD): δ 7.90 (d, J = 8.8 Hz, 1H), 7.02 (d, J = 2.8 Hz, 1H), 6.92 (dd, J = 8.8, 2.4 Hz, 1H), 4.04 (t, J = 6.4 Hz, 2H), 1.85 - 1.75 (m, 2H), 1.50 - 1.35 (m, 4H), 0.96 (t, J = 7.2 Hz, 3H).

[00657] Compound 270 (formic acid salt) was prepared as a white solid utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): t_R = 0.591 min, [M + H]⁺ =

862.6.


Example 176: Synthesis of Compound 271

[00658] Following similar procedures as described in Example 172, 2-chloro-4-


5 (hexyloxy)benzoic acid (670 mg, 2.61 mmol, 88.3% yield) was obtained as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 0.926$ min, $[M + H]^+ = 256.9$.

[00659] Compound 271 (formic acid salt) was prepared as a white solid utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): $t_R = 0.735$ min, $[M + H]^+ = 876.6$.


10 **Example 177: Synthesis of Compound 272**




[00660] Compound 272 (formic acid salt) was prepared as a white solid utilizing the methods in Example 172 (Compound 267) from 2-chloro-4-(octyloxy)benzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.649$ min, $[M + H]^+ = 904.6$.

Example 178: Synthesis of Compound 273


[00661] Compound 273 (formic acid salt) was prepared as a white solid utilizing the methods in Example 172 (Compound 267) from 4-(heptyloxy)-2-(trifluoromethyl)benzoic acid. LCMS
 5 (Method 5-95 AB, ESI): $t_R = 0.634$ min, $[M + H]^+ = 924.4$.

Example 179: Synthesis of Compound 274


[00662] Compound 274 (formic acid salt) was prepared as a white solid utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): $t_R = 0.582$ min, $[M+H]^+ =$
 10 842.3.

Example 180: Synthesis of Compound 275


[00663] Compound 275 (formic acid salt) was prepared utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): $t_R = 0.601$ min, $[M+H]^+ = 868.5$.

Example 181: Synthesis of Compound 276

[00664] Compound 276 (formic acid salt) was prepared utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): $t_R = 0.679$ min, $[M+H]^+ = 828.5$.


5 Example 182: Synthesis of Compound 277

[00665] Compound 277 (formic acid salt) was prepared utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): $t_R = 0.613$ min, $[M+Na]^+ = 878.4$.


Example 183: Synthesis of Compound 278

10


[00666] Compound 278 (formic acid salt) was prepared utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): $t_R = 0.706$ min, $[M+H]^+ = 904.5$.

Example 184: Synthesis of Compound 279


[00667] Compound 279 (formic acid salt) was prepared utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): $t_R = 0.713$ min, $[M+H]^+ = 862.4$.

5 Example 185: Synthesis of Compound 280


[00668] Compound 280 (formic acid salt) was prepared as a white solid utilizing the methods in Example 172 (Compound 267) from 2-methyl-4-(neopentyloxy)benzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.575$ min, $[M/2+H]^+ = 421.9$.

10 Example 186: Synthesis of Compound 281

[00669] Compound 281 (formic acid salt) was prepared as a white solid (38 mg) utilizing the methods in Example 172 (Compound 267) from 4-(heptyloxy)-2-methylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.755$, $[M + H]^+ = 884.5$.

Example 187: Synthesis of Compound 282

[00670] Compound 282 (formic acid salt) was prepared as a white solid (87 mg) utilizing the methods in Example 172 (Compound 267) from 2-chloro-4-(heptyloxy)benzoic acid. LCMS
 5 (Method 5-95 AB, ESI): $t_R = 0.760$, $[M + H]^+ = 904.9$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.45 (brs, 1H, HCOOH), 7.46 (d, $J=8.0$ Hz, 1H), 7.32-7.24 (m, 2H), 7.18 (d, $J=8$ Hz, 1H), 7.11 (d, $J=8$ Hz, 1H), 7.02 (brs, 1H), 6.94 (d, $J=8.0$ Hz, 1H), 6.89 (brs, 1H), 6.81 (brs, 1H), 6.41 (s, 1H), 5.07-5.03 (m, 1H), 4.81-4.75 (m, 2H), 4.29-4.17 (m, 4H), 4.20 (s, 2H), 4.01 (t, $J=4.8$ Hz, 2H), 3.38-3.34 (m, 1H), 3.28-3.24 (m, 4H), 3.15-3.05 (m, 1H), 3.01-2.94 (m, 1H), 2.96 (s, 3H), 2.02-
 10 1.97 (m, 1H), 1.90-1.75 (m, 5H), 1.52-1.30 (m, 11H), 0.92 (t, $J=6.8$ Hz, 3H).


Example 188: Synthesis of Compound 283

[00671] To a stirred solution of methyl 2-chloro-4-hydroxy-benzoate (200.0 mg, 1.07 mmol),
 15 hept-2-yn-1-ol (120.23 mg, 1.07 mmol), and diphenyl-2-pyridylphosphine (338.63 mg, 1.29 mmol) in dry tetrahydrofuran (10 mL) at 0 °C under a nitrogen atmosphere was added dropwise a solution of di-*t*-butyl azodicarboxylate (296.17 mg, 1.29 mmol) in THF (2 mL) over a period of 5 min. The reaction was stirred for 1 h and the solvent was evaporated under reduced pressure. The resulting oil was diluted with water (30 mL) and extracted with EtOAc (25mL x
 20 3). The combined organic layers were washed with 1N HCl (10 mL x 3) and brine (20 mL x 3),

dried over Na_2SO_4 and concentrated. The residue was purified by prep-TLC (10% EtOAc in petroleum ether, $R_f = 0.1$) to give methyl 2-chloro-4-hept-2-ynoate (220 mg, 0.7836 mmol, 73.1% yield) as a colorless oil. HNMR (400 MHz, CDCl_3): 7.86 (d, $J = 8.8$ Hz, 1H), 7.04 (d, $J = 2.8$ Hz, 1H), 6.89-6.87 (m, 1H), 4.69 (s, 2H), 3.88 (s, 3H), 2.20 (t, $J = 6.8$ Hz, 2H), 1.49-1.43 (m, 2H), 1.39-1.35 (m, 2H), 0.87 (t, $J = 7.2$ Hz, 3H).

[00672] Compound 283 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.712$ min, $[\text{M}+\text{H}]^+ = 886.4$.

Example 189: Synthesis of Compound 284



[00673] To a solution of methyl isobutyrate (2000.0 mg, 19.58 mmol) in tetrahydrofuran (50 mL) cooled at -78°C under N_2 protection was added lithium diisopropylamide (11.75 mL, 23.5 mmol) and stirred at 0°C for 3 h. Propargyl bromide (2562.5 mg, 21.54 mmol) was added. After being stirred at 20°C for 2 h, the reaction was quenched with saturated NH_4Cl solution (10 mL) and extracted with EtOAc (20 mL). The organic layer was washed with water (20 mL x 2) and brine (20 mL), dried over MgSO_4 and concentrated. The residue was purified by flash column chromatography (10% EtOAc in petroleum ether) to give methyl 2,2-dimethylpent-4-ynoate (500 mg, 3.5668 mmol, 18.2% yield) as a yellow oil.

[00674] To a solution of methyl 2,2-dimethylpent-4-ynoate (4000.0 mg, 28.54 mmol) in tetrahydrofuran (5 mL) cooled at 0°C was added lithium aluminum hydride (1600.0 mg, 42.8 mmol) in portions, and stirred at 0°C for 3 h. The reaction was quenched sequentially with water (1 mL), 10% NaOH solution (1 mL), and water (1 mL). The mixture was filtered and concentrated to afford 2,2-dimethylpent-4-yn-1-ol (3000 mg, 26.745 mmol, 93.7% yield) as a


yellow oil. ^1H NMR (400 MHz, CDCl_3) δ 3.44 (s, 2H), 2.18 (d, $J = 2.4$ Hz, 2H), 2.01 (t, $J = 2.6$ Hz, 1H), 0.99 (s, 6H).

[00675] A mixture of 2,2-dimethylpent-4-yn-1-ol (500.0 mg, 4.46 mmol), 4-dimethylaminopyridine (54.5 mg, 0.45 mmol), triethylamine (2255.3 mg, 22.29 mmol) and p-toluenesulfonyl chloride (1699.7 mg, 8.92 mmol) in dichloromethane (20 mL) was stirred at 35 °C for 16 h and concentrated to dryness. The residue was taken up in EtOAc (20 mL), washed with water (20 mL x 2) and brine (10 mL), dried over MgSO_4 and concentrated. The residue was purified by flash column chromatography (10% EtOAc in petroleum ether) to afford 2,2-dimethylpent-4-ynyl 4-methylbenzenesulfonate (900 mg, 3.38 mmol, 75.8% yield) as a yellow oil.


[00676] A mixture of methyl 4-hydroxy-2-methylbenzoate (100.0 mg, 0.60 mmol), cesium carbonate (588.23 mg, 1.81 mmol) and 2,2-dimethylpent-4-ynyl 4-methylbenzenesulfonate (320.6 mg, 1.2 mmol) in N,N-dimethylformamide (10 mL) was stirred at 120 °C for 16 h. The reaction was quenched with water (20 mL) and extracted with EtOAc (20 mL). The organic layer was washed with water (20 mL x 2) and brine (20 mL), dried over MgSO_4 and concentrated. The residue was purified by flash column chromatography (10% EtOAc in petroleum ether) to afford methyl 4-(2,2-dimethylpent-4-ynoxy)-2-methylbenzoate (150 mg, 95.7% yield) as a yellow oil. Ester hydrolysis with NaOH (General Method NaOH) afforded 4-((2,2-dimethylpent-4-yn-1-yl)oxy)-2-methylbenzoic acid.

[00677] Compound 284 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.600$ min, $[\text{M}+\text{H}]^+ = 867.5$.

Example 190: Synthesis of Compound 285




[00678] Compound 285 (formic acid salt) was prepared as a white solid utilizing the methods in Example 189 (Compound 284) except allyl bromide was used in the initial alkylation step with methyl isobutyrate. LCMS (Method 5-95 AB, ESI): $t_R = 0.599$ min, $[\text{M}+\text{H}]^+ = 868.5$.

Example 192: Synthesis of Compound 287


[00679] Compound 287 (formic acid salt) was prepared as a white solid utilizing the methods in Example 189 (Compound 284) starting from (1R,4S)-bicyclo[2.2.1]heptan-2-ylmethanol.

5 **[00680]** LCMS (Method 5-95 AB, ESI): RT = 0.723, [M+H]⁺ = 880.5.



Example 193: Synthesis of Compound 288

[00681] To 2,2-dimethylpentanoic acid (500 mg, 3.84 mmol) was added 1M borane (19.2 mL, 19.2 mmol) in tetrahydrofuran at 0°C, and stirred for 16 h under nitrogen. The reaction was quenched by MeOH at 0°C and concentrated to obtain crude 2,2-dimethylpentan-1-ol (540 mg, 4.6472 mmol) as a yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 3.31 (s, 2H), 1.35-1.15 (m, 4H), 0.90 (t, J = 6.8 Hz, 3H), 0.86 (s, 6H).

[00682] Compound 288 (formic acid salt) was prepared as a white solid utilizing the methods in Example 189 (Compound 284). LCMS (Method 5-95 AB, ESI): t_R = 0.749 min, [M+Na]⁺ = 892.8.

Example 194: Synthesis of Compound 289

[00683] Compound 289 (formic acid salt) was prepared as a white solid utilizing the methods in Example 193 (Compound 288) starting from 1-methylcyclohexanecarboxylic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.636$ min, $[M+Na]^+ = 904.5$.


Example 195: Synthesis of Compound 290

[00684] Step 1: 1,3-Dichloro-5-heptoxy-benzene was prepared using the methods in Example 172 (Compound 267).

[00685] Step 2: To a solution of 1,3-dichloro-5-heptoxy-benzene (200.0 mg, 0.77 mmol) in tetrahydrofuran (10 mL) was added lithium diisopropylamide (0.37 mL, 0.92 mmol) and stirred at -78°C for 1 h. CO_2 was bubbled and stirred at -78°C for 2 h. The reaction was quenched with water (40 mL) and extracted with EtOAc (30 mL x 3). The organic layers were washed with water (50 mL x 2) and brine (50 mL), dried over MgSO_4 and concentrated. The residue was purified by prep-TLC (20% EtOAc in petroleum ether, $R_f = 0.5$) to obtain 2,6-dichloro-4-heptoxy-benzoic acid (150 mg, 64.2% yield) as a white solid. ^1H NMR (400 MHz, CDCl_3): δ 6.72 (s, 2H), 3.83 (t, $J = 6.4$ Hz, 2H), 1.69 - 1.64 (m, 2H), 1.35 - 1.19 (m, 8H), 0.82 (t, $J = 6.4$ Hz, 3H).


[00686] Compound 290 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.754$ min, $[M+H]^+ = 924.5$.

Example 196: Synthesis of Compound 291




- 5 **[00687]** Compound 291 (formic acid salt) was prepared utilizing the methods in Example 195 (Compound 290). LCMS (Method 5-95 AB, ESI): $t_R = 0.756$ min, $[M+H]^+ = 909.6$.

Example 197: Synthesis of Compound 292




- 10 **[00688]** Methyl 4-(heptyloxy)-2,6-dimethylbenzoate was prepared by the carbonylation conditions from Example 170 (Compound 265) from 2-bromo-5-(heptyloxy)-1,3-dimethylbenzene.


[00689] Compound 292 (formic acid salt) was prepared utilizing the methods in Example 170 (Compound 265). LCMS (Method 5-95 AB, ESI): $t_R = 0.752$ min, $[M+H]^+ = 884.5$.

Example 198: Synthesis of Compound 293

[00690] Compound 293 (formic acid salt) was prepared utilizing the methods in Example 197 (Compound 292). LCMS (Method 5-95 AB, ESI): $t_R = 0.631$ min, $[M+H]^+ = 908.5$.


5 Example 199: Synthesis of Compound 294

[00691] Compound 294 (formic acid salt) was prepared utilizing the methods in Example 197 (Compound 292). LCMS (Method 5-95 AB, ESI): $t_R = 0.761$ min, $[M+H]^+ = 884.4$.



Example 200: Synthesis of Compound 295

10

[00692] Compound 295 (formic acid salt) was prepared utilizing the methods in Example 197 (Compound 292). LCMS (Method 5-95 AB, ESI): $t_R = 0.635$ min, $[M+H]^+ = 906.4$.

Example 201: Synthesis of Compound 296

[00693] Compound 296 (formic acid salt) was prepared utilizing the methods in Example 197 (Compound 292). LCMS (Method 5-95 AB, ESI): $t_R = 0.738$ min, $[M + H]^+ = 878.4$.

5 Example 202: Synthesis of Compound 297

[00694] Step 1: Borane-THF complex (31.85 mL, 31.85 mmol) was added to 4-bromo-3-chlorobenzoic acid (1.5 g, 6.37 mmol) at 0 °C, and stirred at 0 °C for 10 h. The reaction was quenched by methanol (50 mL) and concentrated under reduced pressure to give (4-bromo-3-chlorophenyl)methanol (1.3 g, 5.8696 mmol, 92.1% yield) as a light yellow oil.


[00695] Step 2: To a solution of (4-bromo-3-chlorophenyl)methanol (1.3 g, 5.87 mmol) in 1,2-dichloroethane (20 mL) at 25 °C was added manganese dioxide (5.1 g, 58.7 mmol), and stirred at 25 °C for 18 h. The mixture was filtered and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column (5-10% EtOAc in petroleum ether) to give 4-bromo-3-chlorobenzaldehyde (600 mg, 46.6% yield) as a light yellow oil.

[00696] Step 3: To a suspension of anhydrous ferric chloride (14.2 mg, 0.09 mmol) and 4-bromo-3-chlorobenzaldehyde (462.4 mg, 2.11 mmol) in nitromethane (10 mL) were successively added tert-butyl(hexyloxy)dimethylsilane (380.0 mg, 1.76 mmol) and triethylsilane (449.1 mg, 3.86 mmol) at 0 °C under nitrogen. After stirring at room temperature for 30 min, the reaction was quenched with phosphate buffer till pH = 7 and extracted with DCM (45 mL x 3). The combined organic layers were washed with brine (15 mL x 3), dried over Na₂SO₄, filtered and concentrated. The residue was purified by prep-TLC (5% EtOAc in petroleum ether) to give 1-bromo-2-chloro-4-((hexyloxy)methyl)benzene (450 mg, 83.9% yield) as a colorless oil.

[00697] Step 4: Following the methoxycarbonylation procedure (General Method Carbonylation), 1-bromo-2-chloro-4-((hexyloxy)methyl)benzene (200.0 mg, 0.65 mmol) was converted to methyl 2-chloro-4-((hexyloxy)methyl)benzoate (125 mg, 67.1% yield) as a colorless oil. ¹H NMR (400 MHz, CDCl₃): δ 7.80 (d, *J* = 8.0 Hz, 1H), 7.42 (s, 1H), 7.25 (d, *J* = 8.0 Hz, 1H), 4.49 (s, 2H), 3.91 (s, 3H), 3.46 (t, *J* = 6.6 Hz, 2H), 1.64 - 1.61 (m, 2H), 1.37 - 1.29 (m, 6H), 0.88 (t, *J* = 6.6 Hz, 3H).

[00698] Compound 297 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): *t*_R = 0.719 min, [M+H]⁺ = 890.9.

Example 203: Synthesis of Compound 298




[00699] To a solution of 4-bromo-2-methylbenzoic acid (3.0 g, 13.95 mmol) in tetrahydrofuran (30 mL) at -78 °C was added 2M nBuLi (13.95 mL, 27.9 mmol) in hexanes and stirred at -78 °C for 3 h. DMF (2.0 g, 27.9 mmol) was added and the mixture was stirred at -78 °C for 1.5 h. The reaction was quenched with 1N HCl (20 mL) and extracted with EtOAc (200 mL x 4). The combined organic layers were dried over Na₂SO₄ and concentrated. The resulting yellow solid was washed with petroleum ether to give crude 4-formyl-2-methylbenzoic acid (900 mg, 5.4825 mmol, 39.3% yield), which was used directly in the next step.

[00700] To a solution of 4-formyl-2-methyl-benzoic acid (0.9 g, 5.48 mmol) in N,N-dimethylformamide (5 mL) at 20°C were added K₂CO₃ (2269.8 mg, 16.45 mmol) and iodomethane (2.84 g, 20.01 mmol) and stirred for 10 h. The mixture was diluted with 1N HCl (15 mL) and extracted with EtOAc (30 mL x 3). The combined organic layers were washed with
5 brine (30 mL x 3), dried over Na₂SO₄ and concentrated. The residue was purified by silica gel column (5% EtOAc in petroleum ether) to give methyl 4-formyl-2-methyl-benzoate (900 mg, 5.0511 mmol, 92.1% yield) as a yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 10.03 (s, 1H), 8.01 (d, J = 8.4 Hz, 1H), 7.74-7.72 (m, 2H), 3.92 (s, 3H), 2.65 (s, 3H).

[00701] Compound 298 (formic acid salt) was prepared utilizing the methods in Example 202 (Compound 297). LCMS (Method 5-95 AB, ESI): t_R = 0.726 min, [M+H]⁺ = 870.5.

Example 204: Synthesis of Compound 299




[00702] Step 1: To a solution of methyl 4-bromo-2-chloro-6-methylbenzoate (300 mg, 1.14 mmol) and bis(pinacolato)diboron (361.4 mg, 1.42 mmol) in dimethyl sulfoxide (5 mL) were added 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (83.3 mg, 0.11 mmol) and potassium acetate (290.5 mg, 2.96 mmol), and stirred at 100 °C for 16 h under N₂. The reaction mixture was diluted with water (5 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. The
20 residue was purified by prep-TLC (9% EtOAc in petroleum ether, R_f = 0.5) to give methyl 2-chloro-6-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (260 mg, 73.5% yield). ¹H NMR (400 MHz, CDCl₃): δ 7.63 (s, 1H), 7.51 (s, 1H), 3.92 (s, 3H), 2.30 (s, 3H), 1.32 (s, 12H).

[00703] Step 2: To a solution of methyl 2-chloro-6-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (260.0 mg, 0.84 mmol) in tetrahydrofuran (3 mL) were added hydrogen peroxide (187.1 mg, 1.93 mmol) and 1M aq. sodium hydroxide (0.84 mL, 0.84 mmol),

and stirred at 28 °C for 16 h. The mixture was diluted with water (5 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated to give crude methyl 2-chloro-4-hydroxy-6-methylbenzoate (150 mg, 89.3% yield) as a yellow solid. LCMS (Method 5-95 AB, ESI): t_R = 0.654 min, $[M+H]^+$ = 200.9.

- 5 **[00704]** Compound 299 (formic acid salt) was prepared as a white solid utilizing the methods in Example 172 (Compound 267). LCMS (Method 5-95 AB, ESI): t_R = 0.756 min, $[M+H]^+$ = 904.5.


Example 205: Synthesis of Compound 300 and 301



[00705] To a solution of ethyl 2-hydroxy-4-methylpyrimidine-5-carboxylate (200.0 mg, 1.1 mmol) in N,N-dimethylformamide (5 mL) were added cesium carbonate (1073.1 mg, 3.29 mmol) and 1-bromoheptane (636.03 mg, 3.29 mmol). The mixture was stirred at 20 °C for 16 h and concentrated to dryness. The residue was taken up in EtOAc (50 mL), washed with water (50 mL x 2) and brine (50 mL), dried over MgSO₄ and concentrated. The residue was purified by prep-TLC to afford ethyl 2-(heptyloxy)-4-methylpyrimidine-5-carboxylate (55 mg, 0.1962 mmol, 17.9% yield; 5% EtOAc in petroleum ether, R_f = 0.3) and ethyl 1-heptyl-4-methyl-2-oxo-1,2-dihydropyrimidine-5-carboxylate (130 mg, 0.4637 mmol, 42.2% yield; 50% EtOAc in petroleum ether, R_f = 0.3), both as a white solid.

- 20 **[00706]** Compound 299 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from ethyl 2-(heptyloxy)-4-methylpyrimidine-5-carboxylate. LCMS (Method 5-95 AB, ESI): t_R = 0.722 min, $[M+H]^+$ = 872.4.


- [00707]** Compound 300 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from ethyl 1-heptyl-4-methyl-2-oxo-1,2-dihydropyrimidine-5-carboxylate. LCMS (Method 5-95 AB, ESI): t_R = 0.646 min, $[M+H]^+$ = 872.6.
- 25


Example 206: Synthesis of Compound 302

[00708] Step 1: A degassed mixture of 4-tert-butylphenol (275.4 mg, 1.83 mmol), methyl 4-bromo-2-methylbenzoate (350.0 mg, 1.53 mmol), (dimethylamino)acetic acid (86.7 mg, 0.84 mmol), cuprous chloride (75.6 mg, 0.76 mmol) and cesium carbonate (995.7 mg, 3.06 mmol) in 1,4-dioxane (10 mL) was heated at 105 °C for 16 h under nitrogen. After cooling down, the reaction was diluted with EtOAc (15 mL) and filtered. The filtrate was concentrated and the residue was purified by prep-TLC (2% EtOAc in petroleum ether, $R_f = 0.5$) to give methyl 4-(4-(tert-butyl)phenoxy)-2-methylbenzoate (240 mg, 52.6% yield) as a colorless oil.

[00709] Step 2: Methyl 4-(4-(tert-butyl)phenoxy)-2-methylbenzoate (240.0 mg, 0.80 mmol) was hydrolyzed as previously described (General Method NaOH) to give 4-(4-(tert-butyl)phenoxy)-2-methylbenzoic acid (210 mg, 91.8% yield) as a white solid. ^1H NMR (400 MHz, CDCl_3): δ 8.06 (d, $J = 8.4$ Hz, 1H), 7.41 (d, $J = 8.8$ Hz, 2H), 7.01 (d, $J = 8.4$ Hz, 2H), 6.89 – 6.78 (m, 2H), 2.64 (s, 3H), 1.36 (s, 9H).

[00710] Compound 302 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.777$ min, $[\text{M} + \text{H}]^+ = 904.6$.


Example 207: Synthesis of Compound 303



[00711] Following similar procedures as described in Example 206, 4-(4-butylphenoxy)-2-methylbenzoic acid (180 mg, 94.4% yield) was obtained as a white solid. ^1H NMR (400 MHz, CDCl_3): δ 8.06 (d, $J = 8.8$ Hz, 1H), 7.20 (d, $J = 8.4$ Hz, 2H), 6.99 (d, $J = 8.4$ Hz, 2H), 6.86 – 6.77 (m, 2H), 2.67 – 2.58 (m, 5H), 1.67-1.59 (m, 2H), 1.42-1.36 (m, 2H), 0.96 (t, $J = 7.2$ Hz, 3H).

[00712] Compound 303 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.756$ min, $[\text{M} + \text{H}]^+ = 904.4$.

10 Example 208: Synthesis of Compound 304



[00713] A mixture of ethyl 2-hydroxy-4-methylpyrimidine-5-carboxylate (50.0 mg, 0.27 mmol), PyAOP (485.5 mg, 1.1 mmol) and Cs_2CO_3 (179 mg, 0.55 mmol) in N,N-dimethylformamide (5 mL) was stirred at 25 °C for 1 h, then 4-isobutylphenol (164.9 mg, 1.1 mmol) and Cs_2CO_3 (179 mg, 0.55 mmol) were added. The reaction was stirred in the microwave at 75 °C for 1 h, diluted with water (20 mL) and extracted with EtOAc (25 mL x 3). The combined organic layers were washed with brine (25 mL x 3), dried over Na_2SO_4 and concentrated. The residue was purified by prep-TLC (10% EtOAc in petroleum ether) to give ethyl 2-(4-isobutylphenoxy)-4-methylpyrimidine-5-carboxylate (110 mg, 63.7% yield).


[00714] Step 2: Ethyl 2-(4-isobutylphenoxy)-4-methylpyrimidine-5-carboxylate (110.0 mg, 0.35 mmol) was hydrolyzed with LiOH (General Method LiOH) to afford 2-(4-isobutylphenoxy)-4-methylpyrimidine-5-carboxylic acid (60 mg, 59.9% yield) as a white solid.

^1H NMR (400 MHz, CDCl_3): δ 9.07 (s, 1H), 7.20 (d, $J = 8.8$ Hz, 2H), 7.11 (d, $J = 8.4$ Hz, 2H), 2.82 (s, 3H), 2.50 (d, $J = 7.2$ Hz, 2H), 1.95-1.80 (m, 1H), 0.94 (d, $J = 6.4$ Hz, 6H).

[00715] Compound 304 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.722$ min, $[\text{M}+\text{H}]^+ =$

5 906.4.


Example 209: Synthesis of Compound 305



[00716] Compound 305 (formic acid salt) was prepared as a white solid utilizing the methods in Example 208 (Compound 304). LCMS (Method 5-95 AB, ESI): $t_R = 0.706$ min, $[\text{M}+\text{H}]^+ =$

10 906.5.


Example 210: Synthesis of Compound 306




[00717] Compound 306 (formic acid salt) was prepared as a white solid utilizing the methods in Example 208 (Compound 304). LCMS (Method 5-95 AB, ESI): $t_R = 0.718$ min, $[\text{M}+\text{H}]^+ =$

15 906.7.

Example 211: Synthesis of Compound 307






[00718] The preparation of alkynyl aromatic compounds is accomplished by Sonogashiro coupling and base hydrolysis. Methyl 2-methyl-4-(pent-1-yn-1-yl)benzoate is an intermediate in the preparation of Compound 197 (Example 103).


- 5 Ester hydrolysis with NaOH according to Example 47 afforded 2-methyl-4-(pent-1-yn-1-yl)benzoic acid.

[00719] Compound 307 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): $t_R = 0.686$, $[M + H]^+ = 822.4$.

Example 212: Synthesis of Compound 308



10




[00720] Step 1: To a solution of methyl 2-methyl-4-(pent-1-yn-1-yl)benzoate (Example 103) (300 mg, 1.4 mmol) in MeOH (50 mL) was added Lindlar Pd catalyst (30 mg, 0.46 mmol) and the mixture was stirred at room temperature under H_2 (15 psi) for 1 h. The filtrate was concentrated to afford (Z)-methyl 2-methyl-4-(pent-1-en-1-yl)benzoate (200 mg, 65% yield) as colorless oil. LCMS (5-95 AB, 2 min, ESI): $t_R = 1.464$, $[M + H]^+ = 219.2$.

[00721] Step 2: Ester hydrolysis procedure with NaOH (Example 47) was applied to (Z)-methyl 2-methyl-4-(pent-1-en-1-yl)benzoate (100 mg, 0.45 mmol) to afford (Z)-2-methyl-4-(pent-1-en-1-yl)benzoic acid (80 mg, 85.4% yield) as a white solid.

[00722] Compound 308 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): $t_R = 0.573$, $[M + H]^+ = 824.4$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.49 (brs, 2H, HCOOH), 7.39 (d, $J=8.0$ Hz, 1H), 7.32 (d, $J=8.0$ Hz,


1H), 7.24 (d, $J=8.0$ Hz, 1H), 7.18 (brs, 1H), 7.16 (d, $J=8.0$ Hz, 2H), 7.09 (d, $J=8.0$ Hz, 1H), 6.90 (brs, 1H), 6.82 (brs, 1H), 6.43 (d, $J=12.0$ Hz, 1H), 6.33 (s, 1H), 5.80-5.70 (m, 1H), 5.12-5.16 (m, 1H), 4.82-4.78 (m, 2H), 4.36-4.16 (m, 4H), 4.19 (s, 2H), 3.37-3.10 (m, 8H), 2.93 (s, 3H), 2.43 (s, 3H), 2.33-2.15 (m, 3H), 2.12-2.00 (m, 1H), 1.52-1.46 (m, 2H), 1.36 (d, $J=6.8$ Hz, 3H), 0.94 (t, $J=7.2$ Hz, 3H).

Example 213: Synthesis of Compound 309




[00723] Compound 309 (formic acid salt) was prepared as a white solid utilizing the methods in Example 211 (Compound 307). LCMS (Method 5-95 AB, ESI): $t_R = 0.567$ min, $[M+H]^+ = 822.7$.

Example 214: Synthesis of Compound 310




[00724] Compound 310 (formic acid salt) was prepared as a white solid utilizing the methods in Example 211 (Compound 307). LCMS (Method 5-95 AB, ESI): $t_R = 0.743$ min, $[M+H]^+ = 862.5$.

Example 215: Synthesis of Compound 311



[00725] Compound 311 (formic acid salt) was prepared as a white solid utilizing the methods in Example 211 (Compound 307). LCMS (Method 5-95 AB, ESI): $t_R = 0.743$ min, $[M+H]^+ = 898.7$.


Example 216: Synthesis of Compound 312



5

[00726] Compound 312 (formic acid salt) was prepared as a white solid utilizing the methods in Example 211 (Compound 307). LCMS (Method 5-95 AB, ESI): $RT = 0.662$ min, $[M+Na]^+ = 887.7$.


Example 217: Synthesis of Compound 313




10

[00727] Compound 313 (formic acid salt) was prepared as a white solid utilizing the methods in Example 133 (Compound 227). LCMS (Method 5-95 AB, ESI): $RT = 0.611$ min, $[M+Na]^+ = 878.4$.

Example 218: Synthesis of Compound 314



15



[00728] Step 1: Compound 101-D (2.0 g, 3.65 mmol) was added to a solution of 1.25N HCl/MeOH (150 mL) and the mixture was stirred at 0°C for 4 h. The volatiles were removed to give the crude as a white solid.

[00729] Step 2: The above crude was dissolved in DCM (5 mL) and the mixture was added Boc_2O (0.93 g, 4.27 mmol) and TEA (1.08 g, 10.7 mmol). The resulting mixture was stirred for at room temperature for 16 h. The volatiles were removed and the residue was purified by silica gel flash column to obtain Compound 314-A1 and 314-A2 as a mixture of regioisomers (1.8 g, 76.4% yield) as a white solid. LCMS (5-95 AB, ESI): $t_R = 0.880$, $[\text{M} + \text{H}]^+ = 684.6$.


To a mixture of Compound 314-A1 and 314-A2 (1.8 g, 2.72 mmol) and *t*-butyl (2-bromoethyl)carbamate (3.0 g, 13.6 mmol) in DMF (5 mL) was added K_2CO_3 (3.8 g, 27.2 mmol) and the reaction mixture was stirred at room temperature for 3 h. The reaction mixture was added with DCM (50 mL), which was washed with 2N HCl, saturated NaHCO_3 and brine (20 mL each). The organic layer was then dried over Na_2SO_4 , concentrated and the residue was purified on silica gel flash column to afford the mixture of regioisomers, which was further purified by SFC (OD, 250mm x 30mm, 5um) to afford Compound 314-B1 (80 mg, 3.6% yield) and Compound 314-B2 (1.6 g, 73.2% yield) as a white solid.

[00730] Compound 314 (free base) was prepared utilizing methods similar to those from

Example 5 and Example 7 from Compound 314-B2. LCMS (5-95 AB, ESI): $t_R = 0.782$, $[\text{M} + \text{H}]^+ = 811.5$; ^1H NMR (400 MHz, $\text{MeOH}-d_4$) δ 7.28-7.15 (m, 2H), 7.10 (d, $J=8.0$ Hz, 1H), 7.09-7.01 (m, 3H), 6.92 (brs, 1H), 6.83 (d, $J=8.0$ Hz, 1H), 6.61 (brs, 1H), 6.55 (s, 1H), 5.12-5.09 (m, 1H), 4.81-4.78 (m, 1H), 4.68-4.66 (m, 1H), 4.19 (s, 2H), 4.18-4.10 (m, 2H), 3.24-3.20 (m, 1H),


3.14-3.06 (m, 1H), 3.02-2.96 (m, 2H), 2.98 (s, 3H), 2.87-2.80 (m, 2H), 2.53-2.48 (m, 1H), 2.33 (s, 3H), 2.10-1.90 (m, 1H), 1.57-1.54 (m, 2H), 1.36-1.29 (m, 12H), 0.91 (t, $J=7.2$ Hz, 3H).

Example 219: Synthesis of Compound 315




- 5 **[00731]** Compound 315 (free base) was prepared utilizing methods similar to those from Example 5 and Example 7 from Compound 314-B1. LCMS (5-95 AB, ESI): $t_R = 0.788$, $[M + H]^+ = 811.4$.

Example 220: Synthesis of Compound 316




- 10 **[00732]** Compound 316 (formic acid salt) was prepared as a white solid utilizing the methods in Example 37 (Compound 131) from Compound 314-B2. LCMS (5-95 AB, ESI): $t_R = 0.794$, $[M + H]^+ = 706.3$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.53 (brs, 1H), 7.28 (d, $J=8.0$ Hz, 1H), 7.15-7.11 (m, 2H), 6.94 (brs, 1H), 6.90 (d, $J=8.0$ Hz, 1H), 6.78 (brs, 1H), 6.36 (s, 1H), 4.83-4.87 (m, 3H), 4.32-4.36 (m, 2H), 4.31 (s, 2H), 3.33-3.26 (m, 2H), 3.16-3.05 (m, 4H), 2.93 (brs, 2H),
15 2.76 (s, 3H), 1.66-1.80 (m, 2H), 1.35-1.42 (m, 19H), 0.92 (t, $J=6.4$ Hz, 3H).

Example 221: Synthesis of Compound 317



[00733] Compound 317 (formic acid salt) was prepared as a white solid utilizing the methods in Example 37 (Compound 131) from Compound 314-B1. LCMS (5-95 AB, ESI): $t_R = 0.640$, $[M + H]^+ = 706.3$

Example 222: Synthesis of Compound 318



[00734] Step 1: To a stirred solution of Compound 318-A (211 mg, 0.2 mmol) in dry MeOH (5mL) at 0°C , was slowly added SOCl_2 (0.2 mL) dropwise. The reaction mixture was stirred at rt overnight. The solvent was removed under reduced pressure and the residue was dried under high vacuum and used directly in the next reaction to afford Compound 318-B. LCMS: MS (ESI) for $\text{C}_{41}\text{H}_{44}\text{ClN}_5\text{O}_8$: m/z 770.1 ($M + H$) $^+$.


[00735] Step 2: To Compound 318-B (0.3 mmol) was dissolved in CH_2Cl_2 (8 mL) and Et_3N (140 μL , 1.0 mmol, 5eq), to this stirred solution was added $(\text{Boc})_2\text{O}$ (106 μL , 0.46 mmol, 2.3

eq). The reaction mixture was stirred at rt overnight. After the reaction was complete, brine solution was added and the mixture was extracted with ethyl acetate. The combined organic layers washed with brine, dried over anhydrous Na₂SO₄, filtered and solvent was removed under reduced pressure. The residue was purified by flash chromatography (DCM - 5% DCM-MeOH) to afford 152 mg (78%, over 2 steps) of Compound 318-C as a white solid. MS (ESI) for (C₅₁H₆₀ClN₅O₁₂): m/z 970.2 (M + H)⁺.

[00736] Step 3: To a stirred solution of Compound 318-C (145 mg, 0.15 mmol) in dry DMF (5 mL) was added K₂CO₃ (42 mg, 0.3 mmol, 2 eq) followed by tert-butyl (2-bromoethyl)carbamate (100 mg, 0.45 mmol). The reaction mixture was stirred at rt overnight. After completion of the reaction, crushed ice was added and the resultant white cloudy mixture was extracted with EtOAc. The combined organic layers were washed with brine and dried over anhydrous Na₂SO₄, filtered and the solvent was removed under vacuum. The residue was purified by flash chromatography (DCM - 5% DCM-MeOH) to afford 118 mg (71%) of the Compound 318-D as a white solid. MS (ESI) for (C₅₈H₇₃ClN₆O₁₄): m/z 1113.4 (M + H)⁺.


[00737] Compound 318 (formic acid salt) was prepared as a white solid utilizing the methods in Example 235 (Compound 331) from Compound 318-D. LCMS (ESI): [M + H]⁺ = 837.4.

Example 223: Synthesis of Compound 319





[00738] (S)-2-Aminopropanenitrile was prepared in a manner similar to (R)-2-aminopropanenitrile (Example 42 except that (S)-2-((tert-butoxycarbonyl)amino)propanoic acid is used as the starting material.

[00739] Compound 319 (formic acid salt) was prepared as a white solid utilizing the methods in Example 235 (Compound 331) from (S)-2-aminopropanenitrile. LCMS (ESI): [M + H]⁺ = 851.4.

Example 224: Synthesis of Compound 320

[00740] Compound 320 (formic acid salt) was prepared as a white solid utilizing the methods in Example 222. LCMS (ESI): $[M + H]^+ = 834.6$.

5 Example 225: Synthesis of Compound 321


[00741] Compound 321-A was prepared utilizing the methods in Example 218.

[00742] A solution of 321-A (410.0 mg, 0.38 mmol) and piperidine (1 mL, 11.74 mmol) in dichloromethane (4 mL) was stirred at 25 °C for 18 h. The reaction was diluted with DCM (50 mL) and acidified by 10% aq. citric acid. The organic layer was washed with water (20 mL x 2) and brine (20 mL), dried over anhydrous sodium sulfate, and concentrated in vacuo to give 321-B (300 mg, 0.3021 mmol, 80.5% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 1.062$ min, $[M+H]^+ = 993.6$.

[00743] A mixture of 321-B (50.0 mg, 0.05 mmol), 2-bromoethanol (188.7 mg, 1.51 mmol) and potassium carbonate (208.7 mg, 1.51 mmol) in N,N-dimethylformamide (1.5 mL) was stirred at 25 °C for 48 h. The mixture was poured into ice-water (10 mL) and extracted with EtOAc (20 mL x 2). The combined organic layers were washed with water (15 mL x 2) and
 5 brine (20 mL), dried over MgSO₄ and concentrated. The residue was purified by prep-TLC (10% methanol in DCM, R_f = 0.3) to afford 321-C (30 mg, 57.5% yield) as a white solid. LCMS (Method 5-95 AB, ESI): t_R = 1.055 min, [M+H]⁺ = 1038.2.


[00744] Compound 321 (formic acid salt) was prepared as a white solid utilizing methods similar to those from Example 7 using Compound 321-C. LCMS (Method 5-95 AB, ESI): t_R =
 10 0.787 min, [M+H]⁺ = 861.7.

Example 226: Synthesis of Compound 322




[00745] Compound 322 (formic acid salt) was prepared as a white solid utilizing the methods in Example 225 (Compound 321). LCMS (Method 5-95 AB, ESI): t_R = 0.804 min, [M+H]⁺ =
 15 859.8.


Example 227: Synthesis of Compound 323




[00746] Compound 323 (formic acid salt) was prepared as a white solid utilizing the methods in Example 225 (Compound 321). LCMS (Method 5-95 AB, ESI): t_R = 0.823 min, [M+H]⁺ =
 20 847.5.

Example 228: Synthesis of Compound 324


[00747] Compound 324 (formic acid salt) was prepared as a white solid utilizing the methods in Example 225 (Compound 321). LCMS (Method 5-95 AB, ESI): $t_R = 0.823$ min, $[M + H]^+ = 834.7$.

Example 229: Synthesis of Compound 325

[00748] Compound 325 (formic acid salt) was prepared as a white solid utilizing the methods in Example 225 (Compound 321). LCMS (Method 5-95 AB, ESI): $t_R = 0.816$ min, $[M + H]^+ = 849.4$.


Example 230: Synthesis of Compound 326

[00749] Compound 326 (formic acid salt) was prepared as a white solid utilizing the methods in Example 225 (Compound 321). LCMS (Method 5-95 AB, ESI): $t_R = 0.692$ min, $[M + H]^+ = 819.5$.


Example 231: Synthesis of Compound 327 and Compound 328

[00750] Compound 327 (formic acid salt) was prepared as a white solid utilizing the methods in Example 225 (Compound 321). LCMS (Method 5-95 AB, ESI): $t_R = 0.822$ min, $[M + H]^+ = 863.4$.

Compound 328 (formic acid salt) was isolated as a white solid as a result of incomplete deprotection of the Boc-intermediates during the preparation of Compound 327. LCMS (Method 5-95 AB, ESI): $t_R = 0.717$ min, $[M+H]^+ = 919.4$.


Example 232: Synthesis of Compound 329


[00751] Compound 329 (formic acid salt) was prepared as a white solid utilizing the methods in Example 225 (Compound 321) except that the first alkylation of the phenol is performed with (2-bromoethoxy)(tert-butyl)dimethylsilane and the second alkylation is performed with tert-butyl (2-bromoethyl)carbamate. LCMS (Method 5-95 AB, ESI): $t_R = 0.825$ min, $[M + H]^+ = 849.9$.

Example 233: Synthesis of Compound 330

[00752] Compound 330 (formic acid salt) was prepared as a white solid utilizing the methods in Example 225 (Compound 321) except that the first alkylation of the phenol is performed with bromoethane and the second alkylation is performed with tert-butyl (2-bromoethyl)carbamate. LCMS (Method 5-95 AB, ESI): $t_R = 0.839$ min/ 1.5 min, $[M + H]^+ = 833.6$.

5 **Example 234: Synthesis of Compound 331-G**





[00753] Step 1: To a solution of Compound 101-E (800 mg, 1.76 mmol) in DMF (15 mL) was added (S)-2-(((benzyloxy)carbonyl)amino)-6-((tert-butoxycarbonyl)amino)hexanoic acid (735 mg, 1.93 mmol), 3-[(E)-ethylazo]-N,N-dimethyl-propan-1-amine hydrochloride (946.77 mg, 5.27 mmol), 1-hydroxybenzotriazole (711.94 mg, 5.27 mmol), and N,N-diisopropylethylamine (681 mg, 5.27 mmol). The mixture was stirred at 30 °C for 16 h. TLC showed the start material was consumed (50% ethyl acetate in petroleum ether, $R_f = 0.5$). The mixture was poured into water (30 mL). The precipitate was filtered, washed with water, re-dissolved in methanol, and concentrated to give Compound 331-A (1200 mg, 1.45 mmol, 83.5% yield) as a yellow solid.

[00754] Step 2: To a solution of Compound 331-A (1200 mg, 1.47 mmol) in methanol (15 mL) was added Pd/C (200.0 mg, 1.47 mmol), and the mixture was stirred at 30 °C under

hydrogen (50 psi) for 16 h. The catalyst was filtered off and the filtrate was concentrated to give Compound 331-B (900 mg, 1.12 mmol, 81.6% yield) as a white solid. LCMS (5-95AB_1.5min_1500): $t_R = 0.782$ min, $[M + H]^+$ 684.4.

[00755] Step 3: The coupling of an acid chloride to an amine is General Method Acid Chloride and is described for this example. A mixture of 4-(4-butylphenyl)benzoic acid (200 mg, 0.79 mmol) in thionyl chloride (5.0 mL) was stirred at 60 °C for 16 h. The solution was concentrated and dissolved in dichloromethane (2 mL). To the solution of Compound 331-B (500 mg, 0.73 mmol) and triethylamine (74 mg, 0.73 mmol) in dichloromethane (15 mL) was added the above solution of 4-(4-butylphenyl)benzoyl chloride in dichloromethane. The reaction mixture was stirred at 25 °C for 3 h. LCMS showed that all of start material was consumed completely. TLC (10% dichloromethane in methanol, $R_f = 0.4$). The reaction was concentrated to dryness and the residue was purified by flash column chromatography (eluted with 5% dichloromethane in methanol). The desired fractions were concentrated to afford Compound 331-C (650 mg, 0.71 mmol, 96.6% yield) as a white solid. LCMS (5-95AB/1.5 min): $t_R = 0.951$ min, $[M + H]^+$ 921.4. Alternatively, this coupling reaction can be performed using 4'-butyl-[1,1'-biphenyl]-4-carboxylic acid using General Method HATU conditions in Example 4.

[00756] Step 4: A mixture of aluminium chloride (2.8 g, 21.19 mmol) and 1-dodecanethiol (4.3 g, 21.19 mmol) in dichloromethane (12 mL) was stirred at 26 °C for 5 min, and then cooled to 0 °C. Then Compound 331-C (650 mg, 0.71 mmol) was added slowly. The solution was stirred at 26 °C for 2 h. LCMS showed that all of start material was consumed completely. The solution was quenched by 1N hydrochloride acid, and filtered. The filter cake was dried to afford crude Compound 331-D as a white solid. LCMS (5-95AB/1.5 min): $t_R = 0.828$ min, $[M + H]^+ = 778.4$.

[00757] Step 5: A solution of Compound 331-D (500 mg, 0.64 mmol) and thionyl chloride (229 mg, 1.93 mmol) in methanol (10 mL) was stirred at 60 °C for 1 h. LCMS showed that all of start material was consumed completely. The solution was concentrated to afford Compound 331-E (500 mg, 0.63 mmol, 98.2% yield) as a yellow solid. LCMS (5-95AB/1.5 min): $t_R = 0.856$ min, $[M + H]^+ = 792.8$.


[00758] Step 6: To the solution of Compound 331-E (500 mg, 0.63 mmol) and sodium bicarbonate (10.6 mg, 0.13 mmol) in 1,4-dioxane (6 mL) and water (2 mL) was added di-tert-butyl dicarbonate (138 mg, 0.63 mmol). LCMS showed that all of start material was consumed completely. TLC (5% dichloromethane in methanol, $R_f = 0.2$). The reaction was concentrated to dryness and the residue was taken up in ethyl acetate (50 mL). It was washed with water (20 mL x 2) and brine (10 mL), dried (sodium sulfate) and concentrated. The crude was purified by flash


column chromatography (eluted with 5% dichloromethane in methanol). The desired fractions were concentrated in vacuo afford Compound 331-F (500 mg, 0.56 mmol, 88.8% yield) as a white solid. LCMS (5-95AB/1.5 min): $t_R = 1.048$ min, $[M + H]^+ = 892.4$.

[00759] Step 7: A mixture of Compound 331-F (500 mg, 0.56 mmol), tert-butyl 2-

5 bromoethylcarbamate (1.25 g, 5.61 mmol) and potassium carbonate (2.32 g, 16.82 mmol) in N,N-dimethylformamide (20 mL) was stirred at 26 °C for 96 h. LCMS showed that all of start material was consumed completely. The reaction was quenched with ice-water (5 mL), and the mixture was taken up in ethyl acetate (20 mL). The organic layer was washed with water (20 mL x 2) and brine (10 mL), dried (sodium sulfate) and concentrated. The crude was purified by
10 flash column chromatography (eluted with ethyl acetate). The desired fractions were concentrated to afford Compound 331-G (450 mg, 0.38 mmol, 68.1% yield) as a colorless oil. LCMS (5-95AB/1.5min): $t_R = 0.995$ min, $[M + H]^+ = 1179.0$.

Example 235: Synthesis of Compound 331





[00760] Step 1: A mixture of Compound 331-G (80 mg, 0.07 mmol) and aqueous lithium hydroxide hydrate (0.41 mL, 0.2 mmol, 0.5 M) in 1, 4-dioxane (2 mL) and water (1 mL) was stirred at 26 °C for 2 h. The solution was quenched with 5% aqueous potassium bisulfate solution to pH = 6, and the mixture was taken up in ethyl acetate (20 mL). The organics were washed with water (20 mL x 2) and brine (10 mL), dried (sodium sulfate) and concentrated to afford Compound 331-H (50 mg, 0.043 mmol, 63.2% yield) as a white solid. LCMS (5-95AB/1.5min): t_R = 1.103 min, $[M + H]^+$ 1164.9.


[00761] Step 2: Starting from Compound 331-H (50.0 mg, 0.04 mmol) and 2-aminoacetonitrile hydrochloride (39.7 mg, 0.43 mmol), General Method HATU (Example 5) was used to prepare Compound 331-I (30 mg, 58.2% yield) as a white solid. LCMS (5-95AB_1.5min): t_R = 1.103 min, $[M + H]^+$ 1204.1.

[00762] Step 3: The deprotection of an N-Boc group using neat formic acid is described and is referred to as General Method FA. A mixture of Compound 331-I (30.0 mg, 0.025 mmol) in formic acid was stirred at 15 °C for 1 hour, then the mixture was lyophilized to a solid.

Preparative HPLC (acetonitrile/water with 0.22% formic acid) afforded Compound 331 (3.7 mg, 16 %) as a white solid. ¹H NMR (400 MHz, CD₃OD): δ 8.49 (s, 3H), 7.94 (d, J = 8.4 Hz, 2H), 7.74 (d, J = 8.0 Hz, 2H), 7.57 (d, J = 7.6 Hz, 2H), 7.32-7.11 (m, 5H), 6.90-6.82 (m, 2H), 6.45 (s,

1H), 5.06-4.95 (m, 1H), 4.80-4.79 (m, 2H), 4.29-4.15 (m, 5H), 3.18-3.10 (m, 5H), 2.99-2.85 (m, 5H), 2.71-2.64 (m, 3H), 2.01-1.92 (m, 2H), 1.82-1.45 (m, 6H), 1.43-1.38 (m, 6H), 0.97 (t, J = 7.2 Hz, 3H). LCMS (Method 5-95 AB, ESI): $t_R = 0.753$ min, $[M + H]^+ = 903.1$.


Example 236: Synthesis of Compound 332



5

[00763] Compound 332 (formic acid salt) was prepared as a white solid utilizing the methods from Example 235 (Compound 331), from Compound 331-H and (S)-2-aminopropanenitrile formic acid salt. LCMS (Method 5-95 AB, ESI): $t_R = 0.770$ min, $[M + H]^+ = 916.7$.


Example 237: Synthesis of Compound 333



10

[00764] Compound 333 (formic acid salt) was prepared as a white solid utilizing the methods from Example 236 (Compound 332) using tetradecanoic acid. LCMS (Method 10-80AB): $t_R = 2.785$ min/ 7 min, $[M + H]^+ = 891.8$.


Example 238: Synthesis of Compound 334



15


[00765] Compound 334 (formic acid salt) was prepared as a white solid utilizing the methods from the preparation of Compound 331 (Example 235). LCMS (Method 5-95 AB, ESI): t_R = 0.769 min, $[M + H]^+ = 876.5$.

Example 239: Synthesis of Compound 335




[00766] Compound 335 (formic acid salt) was prepared as a white solid utilizing the methods in Example 235 (Compound 331). LCMS (Method 5-95 AB, ESI): $t_R = 0.753$ min, $[M + H]^+ = 868.6$.

Example 240: Synthesis of Compound 336




[00767] Compound 336 (formic acid salt) was prepared as a white solid utilizing the methods in Example 235 (Compound 331). LCMS (Method 5-95 AB, ESI): $t_R = 0.783$ min, $[M + H]^+ = 862.9$.

Example 241: Synthesis of Compound 337



[00768] Compound 337 (formic acid salt) was prepared in as a white solid utilizing the methods in Example 235 (Compound 331). LCMS (Method 5-95 AB, ESI): $t_R = 0.771$ min, $[M + H]^+ = 848.7$.


Example 242: Synthesis of Compound 338



5

[00769] Compound 338 (formic acid salt) was prepared as a white solid utilizing the methods in Example 235 (Compound 331). LCMS (Method 5-95 AB, ESI): $t_R = 0.746$ min, $[M + H]^+ = 874.7$.


Example 243: Synthesis of Compound 339



10


[00770] Compound 339 (formic acid salt) was prepared as a white solid utilizing the methods in Example 235 (Compound 331) from 2-methyl-4-octylbenzoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.778$ min, $[M+Na]^+ = 890.5$.

Example 244: Synthesis of Compound 340




15


[00771] Compound 340 (formic acid salt) was prepared as a white solid utilizing the methods from the preparation of Example 236 (Compound 332) LCMS (Method 10-80AB): $t_R = 0.885$ min/ 2 min, $[M + Na]^+ = 896.3$.

Example 245: Synthesis of Compound 341


[00772] Compound 341 (formic acid salt) was prepared as a white solid utilizing the methods in Example 235 (Compound 331). LCMS (Method 5-95 AB, ESI): $t_R = 0.733$ min, $[M + H]^+ =$
 5 860.8.

Example 246: Synthesis of Compound 342


[00773] Compound 342 (formic acid salt) was prepared as a white solid utilizing the methods in Example 236 (Compound 332). LCMS (Method 5-95 AB, ESI): $t_R = 0.813$ min, $[M + H]^+ =$
 10 848.4.

Example 247: Synthesis of Compound 343

[00774] Compound 343 (formic acid salt) was prepared as a white solid utilizing the methods in Example 235 (Compound 331). LCMS (Method 5-95 AB, ESI): $t_R = 0.802$ min, $[M + H]^+ =$
 15 834.5.

Example 248: Synthesis of Compound 344

[00775] Compound 344 (formic acid salt) was prepared as a white solid utilizing the methods in Example 235 (Compound 331). LCMS (Method 5-95 AB, ESI): $t_R = 0.754$ min, $[M + H]^+ = 840.8$.

Example 249: Synthesis of Compound 345

[00776] Step 1: To a mixture of (S)-2-(((benzyloxy)carbonyl)amino)-3-hydroxypropanoic acid (10.0 g, 42 mmol) and imidazole (5.7 g, 84 mmol) in DMF (500 mL), TBSCl (6.3 g, 42 mmol) was added at 0°C. The reaction mixture was warmed to room temperature slowly and stirred for another 16 hr. The volatiles were removed and the residue was taken up in EtOAc (1000 mL) and the EtOAc layer was washed sequentially with 1N HCl, saturated NaHCO₃ and brine (500 mL each). The EtOAc layer was dried over Na₂SO₄, concentrated and the resulting residue was purified by flash column chromatography to afford (S)-2-(((benzyloxy)carbonyl)amino)-3-((tert-butyl)dimethylsilyl)oxy)propanoic acid (7.0 g, 48% yield) as colorless oil.


[00777] Step 2: Typical HATU coupling condition was applied to Compound 101-G (Example 4) (100 mg, 0.14 mmol) and (S)-2-(((benzyloxy)carbonyl)amino)-3-((tert-butyl)dimethylsilyl)oxy)propanoic acid (99 mg, 0.28 mmol) to afford an off-white solid (120 mg, 82% yield).

[00778] Step 3: Typical hydrogenation condition (Pd/C, H₂) was applied to (120 mg, 0.11 mmol) to afford Compound 345-A (100 mg, 96% yield) as a white solid.



[00779] Step 4: To a solution of Compound 345-A (100 mg, 0.11 mmol) and octane-1-sulfonyl chloride compound (93 mg, 0.44 mmol) in DCM (5 mL) was added DIPEA (71 mg, 0.55 mmol) at 0°C. The resulting mixture was stirred at 0°C for 1 h. The volatiles were removed and the resulting residue was taken up in EtOAc (50 mL), which was washed with brine (50 mL x 2). The EtOAc layer was dried over MgSO₄, concentrated and the residue was purified by flash column chromatography to afford Compound 345-B (98 mg, 83% yield) as a white solid.


[00780] Step 5: To a solution of Compound 345-B (98 mg, 0.09 mmol) in DCM (5mL) was added TBAF hydrate (47 mg, 0.18 mmol). The resulting mixture was stirred for 1 h. The volatiles were removed and the residue was purified by flash column chromatography to afford Compound 345-C (80 mg, 92% yield) as a white solid. LCMS (5-95 AB, ESI): t_R = 0.857, [M + H]⁺ = 1000.3.

[00781] Compound 345 was prepared as a white solid utilizing the methods for Compound 101 (Example 7) from Compound 345-C. LCMS (5-95AB_30 min, ESI): t_R = 14.41, [M + H]⁺ = 801.3. ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.51 (brs, 2H, HCOOH), 7.31-7.26 (m, 2H), 7.19 (d, *J*=8.0 Hz, 1H), 7.12 (d, *J*=8.0 Hz, 1H), 6.91 (d, *J*=8.0 Hz, 1H), 6.83 (d, *J*=8.0 Hz, 1H), 6.45 (s, 0.6H), 6.23 (s, 0.4H), 4.80-4.33 (m, 2H), 4.28-4.20 (m, 4H), 4.21 (s, 2H), 3.86-3.80 (m, 1H), 3.78-3.70 (m, 1H), 3.30-3.00 (m, 6H), 2.94 (s, 3H), 2.75 (brs, 2H), 1.86-1.70 (m, 2H), 1.50-1.25 (m, 13H), 0.91 (t, *J*=6.8 Hz, 3H).

Example 250: Synthesis of Compound 346

[00782] Compound 346 (formic acid salt) was prepared as a white solid utilizing the methods in Example 249 (Compound 345) from Compound 101-K (Example 5). LCMS (5-95 AB, ESI): $t_R = 0.765$, $[M + H]^+ = 842.6$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.49 (brs, 2H, HCOOH), 7.27-7.23 (m, 2H), 7.17 (d, $J=8.0$ Hz, 1H), 7.09 (d, $J=8.0$ Hz, 1H), 6.89 (brs, 1H), 6.82 (brs, 1H), 6.34 (s, 1H), 4.80-4.75 (m, 2H), 4.60-4.56 (m, 1H), 4.39-4.13 (m, 4H), 4.19 (s, 2H), 3.23-3.03 (m, 8H), 2.87 (s, 3H), 2.80-2.70 (m, 1H), 2.67-2.65 (m, 1H), 2.13-1.90 (m, 2H), 1.85-1.77 (m, 2H), 1.54-1.23 (m, 17H), 0.90 (t, $J=6.0$ Hz, 3H).

10 Example 251: Synthesis of Compound 347




[00783] Step 1: To a solution of 1-bromo-3-methylbenzene (500 mg, 2.9 mmol) in CHCl₃ (10 mL) was added chlorosulfonic acid (1.2 mL) slowly at 0°C and the reaction was stirred at the same temperature for 4 h. The mixture was poured into crushed ice (50 mL), which was
 5 extracted by CHCl₃ (30 mL x 3). The combined organic layers were washed with brine (60 mL x 2), dried over Na₂SO₄ and concentrated to afford 4-bromo-2-methylbenzene-1-sulfonyl chloride (600 mg, 76% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.91 (d, *J*=8.4 Hz, 1H), 7.58 (d, *J*=1.6 Hz, 1H), 7.55 (dd, *J*=8.4, 1.6 Hz, 1H), 2.75 (s, 3H).

[00784] Step 2: The sulfonamide formation procedure from Example 249 (Compound 345) was applied to Compound 101-K (200 mg, 0.22 mmol) to afford Compound 347-A (160 mg, 63.7% yield) as a white solid.

[00785] Steps 3 and 4: Sonogashira coupling and reduction of Compound 347-A (General Method Sonogashiro) afforded Compound 347-B.

[00786] Compound 347 (formic acid salt) was prepared as a white solid utilizing the methods for Compound 101 (Example 7). LCMS (5-95 AB, ESI): *t*_R = 0.731, [M + H]⁺ = 876.6; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.47 (brs, 2H, HCOOH), 7.85 (d, *J*=8.0 Hz, 1H), 7.29 (brs, 1H), 7.25-7.16 (m, 2H), 7.14-7.04 (m, 2H), 6.83-6.74 (m, 3H), 6.00 (s, 1H), 4.76-4.66 (m, 2H), 4.36-4.30 (m, 1H), 4.29-4.15 (m, 4H), 4.18 (s, 2H), 3.27-2.96 (m, 8H), 2.76-2.72 (m, 2H), 2.67 (s, 3H), 2.61 (s, 3H), 2.13-1.85 (m, 2H), 1.75-1.62 (m, 2H), 1.46-1.24 (m, 9H), 0.90 (t, *J*=6.8 Hz, 3H).

Example 252: Synthesis of Compound 348

[00787] Step 1: To a solution of 2-chloroethanesulfonyl chloride (0.64 mL, 6.1 mmol) in

5 DCM (10 mL) was added pyridine (0.97 g, 12.2 mmol) at -78°C and the resulting mixture was stirred at the same temperature for 20 min. The reaction was warmed to room temperature while stirring and stirred for another 20 min at the same temperature.

[00788] Step 2: To a solution of Compound 348-A (290 mg, 0.31 mmol) and Et_3N (474 mg, 4.7 mmol) in DCM (20 mL) was added the above solution at 0°C . The resulting mixture was

10 warmed to room temperature and stirred for 1 h at the same temperature. DCM (50 mL) was then added and the mixture was washed with saturated citric acid, saturated NaHCO_3 and brine (50 mL each). The organic layer was dried over anhydrous Na_2SO_4 , concentrated and the residue was purified by pre-TLC to afford 348-B (210 mg, 66% yield) as a yellow solid. LCMS (5-95 AB, ESI): RT = 1.056, $\text{M}+\text{Na}^+ = 1041.5$.


15 **[00789]** Step 3: To a solution of 348-B (0.21 g, 0.206 mmol) in MeOH (2mL) was added 1-aminodecane (0.31 g, 2 mmol) at 0°C and the mixture was warmed and stirred at room temperature for 16 h. The volatiles were removed and the residue was purified by flash column

chromatography to afford Compound 348-C (180 mg, zz % yield). LCMS (5-95 AB, ESI): t_R = 1.051, $[M + H]^+ = 1176.6$.

[00790] Step 4: Typical Boc protection condition (Boc₂O, 1.5 equiv, Et₃N) was applied to Compound 348-C (180 mg, 0.15 mmol) to afford 348-D (170 mg, 87% yield) as a white solid.


5 **[00791]** Compound 348 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): $t_R = 0.748$, $[M + H]^+ = 887.0$. ^1H NMR (400 MHz, MeOH- d_4) δ 8.50 (brs, 2H, HCOOH), 7.33-7.11 (m, 3H), 7.12 (d, $J=8.5$ Hz, 1H), 6.92 (d, $J=8.5$ Hz, 1H), 6.84 (d, $J=8.5$ Hz, 1H), 6.40 (s, 0.7H), 5.98 (s, 0.3H), 4.80-4.54 (m, 3H), 4.29-4.20 (m, 4H), 4.21 (s, 2H), 3.75-3.65 (m, 2H), 3.54-3.42 (m, 4H), 3.26-3.03 (m, 7H), 2.90 (s, 3H), 2.73(brs, 1H), 2.08-1.94 (m, 2H), 1.80-1.71 (m, 3H), 1.42-1.32 (m, 18H), 0.91 (t, $J=4.0$ Hz, 3H).

Example 253: Synthesis of Compound 349




[00792] Compound 349 (free base) was prepared as a white solid utilizing the methods in Example 348. LCMS (5-95 AB, ESI): $t_R = 0.583$, $[M + H]^+ = 899.5$; ^1H NMR (400 MHz, MeOH- d_4) δ 7.27-7.17 (m, 2H), 7.10 (d, $J=8.4$ Hz, 1H), 7.02 (d, $J=8.4$ Hz, 1H), 6.85-6.75 (m, 2H), 6.38 (s, 1H), 4.78-4.73 (m, 2H), 4.42 (brs, 1H), 4.20 (s, 2H), 4.12-3.95 (m, 4H), 3.26-3.20 (m, 2H), 3.16-3.09 (m, 2H), 3.09-2.96 (m, 3H), 2.94-2.84 (m, 2H), 2.89 (s, 3H), 2.84-2.67 (m, 3H), 2.65-2.56 (m, 2H), 1.94-1.63 (m, 6H), 1.58-1.45 (m, 3H), 1.43-1.23 (m, 14H), 0.90 (t, $J=6.4$ Hz, 3H).

Example 254: Synthesis of Compound 350





[00793] Compound 350 (formic acid salt) was prepared as a white solid utilizing the methods in Example 252 (Compound 348). LCMS (5-95 AB, ESI): $t_R = 0.572$, $[M + H]^+ = 885.7$.

Example 255: Synthesis of Compound 351



5 [00794] Compound 351 (formic acid salt) was prepared as a white solid utilizing the methods in Example 252 (Compound 348) from Compound 101-G. LCMS (5-95 AB, ESI): $t_R = 0.747$, $[M + H]^+ = 785.3$.

Example 256: Synthesis of Compound 352



10 [00795] To a stirred solution of decanoyl chloride (500 mg, 2.6 mmol) in THF (5 mL) was added (S)-2-aminopentanoic acid (461 mg, 3.9 mmol) and 2N NaOH (5.0 mL) at 0°C and the resulting mixture was stirred at 0°C for 1 h. The pH of the mixture was adjusted to pH=2 using 1N HCl, which was extracted with EtOAc (20 mL x 3). The combined organic layers were


15 washed with brine (50 mL x 2), dried over Na₂SO₄ and concentrated to afford (S)-2-decanamidopentanoic acid (630 mg, 88.5% yield) as a white solid, which was used directly in the next step. LCMS (5-95 AB, ESI): $t_R = 0.904$, $[M + H]^+ = 272.0$.

[00796] Compound 352 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): $t_R = 0.620$, $[M + H]^+ = 891.4$; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.50 (brs, 3H, HCOOH), 7.24-7.16 (m, 2H), 7.14 (d, *J*=8.0 Hz, 1H), 7.08 (d, *J*=8.0 Hz, 1H), 6.87 (brs, 1H), 6.81 (brs, 1H), 6.26 (s, 1H), 5.00-4.97 (m, 1H), 4.85-4.77

20

(m, 3H), 4.23-4.15 (m, 4H), 4.19 (s, 2H), 3.34-3.33 (m, 1H), 3.17-3.03 (m, 7H), 2.82 (s, 3H), 2.28-2.19 (m, 5H), 2.05-1.90 (m, 2H), 1.67-1.62 (m, 5H), 1.35-1.30 (m, 14H), 0.98 (t, $J=7.6$ Hz, 3H), 0.91 (t, $J=7.6$ Hz, 3H).


Example 257: Synthesis of Compound 353



5


[00797] Compound 353 (formic acid salt) was prepared as a white solid utilizing the methods in Example 256 (Compound 352). LCMS (5-95 AB, ESI): $t_R = 0.733$, $[M + H]^+ = 891.6$.

Example 258: Synthesis of Compound 354




10 **[00798]** Compound 354 (formic acid salt) was prepared as a white solid utilizing the methods in Example 256 (Compound 352). LCMS (5-95 AB, ESI): $t_R = 0.730$, $[M + H]^+ = 911.4$.

Example 259: Synthesis of Compound 355




15 **[00799]** Compound 355 (formic acid salt) was prepared as a white solid utilizing the methods in Example 256 (Compound 352). LCMS (5-95 AB, ESI): $t_R = 0.732$, $[M + H]^+ = 911.5$.

Example 260: Synthesis of Compound 356

[00800] Compound 356 (formic acid salt) was prepared as a white solid utilizing the methods in Example 256 (Compound 352) from (2S,3R)-3-(tert-butoxy)-2-decanamidobutanoic acid.


5 LCMS (5-95 AB, ESI): $t_R = 0.720$, $[M + H]^+ = 893.5$.

Example 261: Synthesis of Compound 357-P1 and Compound 357-P2

[00801] Compound 357-P1 and Compound 357-P2 (formic acid salts) were prepared as white solids utilizing the methods in Example 256 (Compound 352) from (D,L)-(erythro)-2-decanamido-3-hydroxybutanoic acid. The compounds are derived from racemic-allo threonine and the stereochemical assignment at these positions is arbitrary. They are distinguished by the elution from reverse phase HPLC with Compound 357-P1 as peak 1 and Compound 357-P2 as peak 2.

[00802] Data for Compound 357-P1: LCMS (5-95 AB, ESI): $t_R = 0.574$, $[M + H]^+ = 893.5$.


15 **[00803]** Data for Compound 357-P2: LCMS (5-95 AB, ESI): $t_R = 0.709$, $[M + H]^+ = 893.7$; ^1H NMR (400 MHz, MeOH- d_4) δ 7.21-7.18 (m, 2H), 7.08 (d, $J=8.0$ Hz, 1H), 7.01 (d, $J=8.0$ Hz, 1H), 6.82 (brs, 1H), 6.77 (brs, 1H), 6.31 (s, 1H), 4.83-4.75 (m, 2H), 4.37-4.35 (m, 1H), 4.32-4.28 (m, 1H), 4.19 (s, 2H), 4.10-3.95 (m, 5H), 3.35-3.15 (m, 1H), 3.16-3.13 (m, 2H), 2.90-2.75 (m, 7H), 2.34-2.32 (m, 3H), 2.13-2.08 (m, 2H), 1.92-1.88 (m, 2H), 1.65-1.62 (m, 3H), 1.33-1.19 (m, 16 H), 0.91 (t, $J=6.6$ Hz, 3H).

Example 263: Synthesis of Compound 359


[00804] Step 1: Decanoic acid and (R)-methyl 2-amino-3-hydroxypropanoate hydrochloride
 5 was subjected to General Method HATU to afford (R)-methyl 2-decanamido-3-hydroxypropanoate.

[00805] Step 2: (R)-Methyl 2-decanamido-3-hydroxypropanoate was subjected to General Method LiOH to afford (R)-2-decanamido-3-hydroxypropanoic acid.


[00806] Compound 359 (formic acid salt) was prepared as a white solid utilizing the methods
 10 in Example 7 (Compound 101) from (R)-2-decanamido-3-hydroxypropanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.576$ min, $[M/2+H]^+ = 440.3$.

Example 264: Synthesis of Compound 360


[00807] Compound 360 (formic acid salt) was prepared as a white solid utilizing the methods
 15 in Example 263 (Compound 359) from (S)-2-decanamido-3-hydroxypropanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.709$ min, $[M + H]^+ = 879.7$.

Example 265: Synthesis of Compound 361


[00808] Compound 361 (formic acid salt) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from 3-(nonylamino)-3-oxopropanoic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.718$ min, $[M + H]^+ = 850.6$.

Example 266: Synthesis of Compound 362

[00809] Compound 362 (formic acid salt) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from 2-(2-decanamidoacetamido)acetic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.719$ min, $[M + H]^+ = 906.5$.


Example 267: Synthesis of Compound 363

[00810] Compound 363 (formic acid salt) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from (S)-2-(2-decanamidopropanamido)acetic acid. LCMS (Method 5-95 AB, ESI): $t_R = 0.718$ min, $[M + H]^+ = 920.6$.

Example 268: Synthesis of Compound 364


[00811] Compound 364 (formic acid salt) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from (S)-3-(2-decanamidopropanamido)propanoic acid.

5 LCMS (Method 5-95 AB, ESI): $t_R = 0.726$ min, $[M + H]^+ = 934.7$.

Example 269: Synthesis of Compound 365


[00812] Compound 365 (formic acid salt) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from (R)-3-(2-decanamidopropanamido)propanoic acid.

10 LCMS (Method 5-95 AB, ESI): $t_R = 0.729$ min, $[M + H]^+ = 934.5$.


Example 270: Synthesis of Compound 366

[00813] Compound 366 (formic acid salt) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from (R)-2-(2-decanamidopropanamido)acetic acid. LCMS


15 (Method 5-95 AB, ESI): $t_R = 0.609$ min, $[M + H]^+ = 920.5$.

Example 271: Synthesis of Compound 367


[00814] Compound 367 (formic acid salt) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from 3-decanamidopropanoic acid. LCMS (Method 5-95 AB,
5 ESI): $t_R = 0.578$ min, $[M + H]^+ = 863.3$.

Example 272: Synthesis of Compound 368

[00815] Compound 368 (formic acid salt) was prepared as a white solid utilizing the methods described previously from 2-(3-decanamidopropanamido)acetic acid. LCMS (Method 5-95 AB,
10 ESI): $t_R = 0.573$ min, $[M + H]^+ = 920.4$.


Example 273: Synthesis of Compound 369

[00816] Compound 369 (formic acid salt) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from 4-(nonylamino)-4-oxobutanoic acid. LCMS (Method 5-
15 95 AB, ESI): $t_R = 0.713$ min, $[M + H]^+ = 863.5$.




Example 274: Synthesis of Compound 370


[00817] 2-(4-Butyl-2-methylbenzamido)acetic acid, a white solid, was prepared from methods previously described. ¹H NMR (400 MHz, CD₃OD): δ 7.35 (d, J = 7.6 Hz, 1H), 7.10 - 7.00 (m, 2H), 4.05 (s, 2H), 2.60 (t, J = 7.8 Hz, 2H), 2.41 (s, 3H), 1.65 - 1.55 (m, 2H), 1.40 - 1.30 (m, 2H), 0.93 (t, J = 7.2 Hz, 3H).

[00818] Compound 370 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): t_R = 0.593 min, [M + H]⁺ = 870.3.

Example 275: Synthesis of Compound 371


[00819] Compound 371 (formic acid salt) was prepared as a white solid utilizing the methods previously described from 2-(2-(4-butyl-2-methylbenzamido)acetamido)acetic acid. LCMS (Method 5-95 AB, ESI): t_R = 0.599 min, [M + H]⁺ = 926.3.



Example 279: Synthesis of Compound 375


[00823] Compound 375 (formic acid salt) was prepared as a white solid utilizing the methods previously described from 3-(2-decanamidoacetamido)propanoic acid. LCMS (Method 5-95


5 AB, ESI): $t_R = 0.588$ min, $[M+Na]^+ = 943.1$.

Example 280: Synthesis of Compound 376

[00824] Compound 376 (formic acid salt) was prepared as a white solid utilizing the methods previously described from 3-(3-decanamidopropanamido)propanoic acid. LCMS (Method 5-95

10 AB, ESI): $t_R = 0.584$ min, $[M + Na]^+ = 956.6$.

Example 281: Synthesis of Compound 377




[00825] Step 1: A mixture of 4-butyl-2-methylbenzoic acid (150.0 mg, 0.78 mmol), diphenylphosphoryl azide (300 mg, 1.09 mmol), benzyl alcohol (252 mg, 2.34 mmol), and triethylamine (220 mg, 2.19 mmol) in toluene (10 mL) was stirred at 100°C for 12 h. The reaction mixture was diluted with water (20 mL) and extracted with EtOAc (30 mL x 3). The combined organic layers were washed with water (20 mL x 2) and brine (20 mL), dried over MgSO₄ and concentrated. The residue was purified by chromatography on silica gel (10% EtOAc in petroleum ether, R_f = 0.5) to obtain benzyl N-(4-butyl-2-methyl-phenyl)carbamate (100 mg, 0.34 mmol, 64.6% yield) as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.87 (s, 1H), 7.40 - 7.35 (m, 5H), 7.21 (d, J = 8.0 Hz, 1H), 6.99 - 6.95 (m, 2H), 5.12 (s, 2H), 2.50 (t, J = 7.6 Hz, 1H), 2.16 (s, 3H), 1.55 - 1.48 (m, 2H), 1.31 - 1.26 (m, 2H), 0.89 (t, J = 7.6 Hz, 1H).

[00826] Step 2: To a solution of benzyl N-(4-butyl-2-methyl-phenyl)carbamate (150 mg, 0.50 mmol) in methanol (10 mL) was added 10% palladium (53.68 mg, 0.05 mmol) on carbon. The reaction mixture was stirred at 15°C for 2 h under H₂ at 15 psi. The mixture was filtered and concentrated to obtain 4-butyl-2-methyl-aniline (80 mg, 0.49 mmol, 97.1% yield) as a white solid.


[00827] Step 3: To a mixture of 4-butyl-2-methylaniline (70 mg, 0.43 mmol) and triethylamine (0.18 mL, 1.29 mmol) in dichloromethane (10 mL) was added methyl 4-chloro-4-oxobutyrates (64.56 mg, 0.43 mmol). The reaction mixture was stirred at 20°C for 12 h. The mixture was concentrated to obtain methyl 4-(4-butyl-2-methyl-anilino)-4-oxo-butanoate (100 mg, 0.36 mmol, 84.1% yield) as a white solid. LCMS (5-95AB_1.5min): t_R = 0.919 min, [M + H]⁺ 278.0.

[00828] Step 4: Methyl 4-(4-butyl-2-methyl-anilino)-4-oxo-butanoate (90 mg, 0.32 mmol) was hydrolyzed as usual to obtain 4-(4-butyl-2-methyl-anilino)-4-oxo-butanoic acid (80 mg, 0.30 mmol, 93.6% yield) as a white solid.


[00829] Compound 377 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): t_R = 0.707 min, [M + H]⁺ = 883.7.

Example 282: Synthesis of Compound 378


[00830] Compound 378 (formic acid salt) was prepared as a white solid utilizing the same methods used in the preparation of Compound 101 (Example 7) except an additional peptide coupling and deprotection step is added to the sequence, starting from (S)-2-(((benzyloxy)carbonyl)amino)-4-((tert-butyl(dimethyl)silyl)oxy)butanoic acid. LCMS (5-95 AB, ESI): $t_R = 0.715$, $[M + H]^+ = 893.7$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.50 (brs, 2H, HCOOH), 7.33-7.24 (m, 2H), 7.19 (d, $J=8.4$ Hz, 1H), 7.11 (d, $J=8.4$ Hz, 1H), 6.90 (brs, 1H), 6.83 (brs, 1H), 6.29 (s, 1H), 5.03-4.76 (m, 3H), 4.40-4.38 (m, 1H), 4.32-4.18 (m, 4H), 4.21 (s, 2H), 3.74-3.62 (m, 2H), 3.32-2.95 (m, 7H), 2.84 (s, 3H), 2.34-2.14 (m, 3H), 2.13-1.82 (m, 3H), 1.66-1.63 (m, 2H), 1.39-1.27 (m, 16H), 0.92 (t, $J=6.4$, 3H).

Example 283: Synthesis of Compound 379


[00831] Compound 379 (formic acid salt) was prepared as a white solid utilizing the methods in Example 282 (Compound 378) from (R)-2-(((benzyloxy)carbonyl)amino)-4-((tert-butyl(dimethyl)silyl)oxy)butanoic acid. LCMS (5-95 AB, ESI): $t_R = 0.712$, $[M + H]^+ = 893.6$.

Example 284: Synthesis of Compound 380


[00832] Compound 380 (formic acid salt) was prepared as a white solid utilizing the same methods used in the preparation of Compound 378 (Example 282). LCMS (5-95 AB, ESI): $t_R =$
 5 0.713, $[M + H]^+ = 883.5$.

Example 285: Synthesis of Compound 381


[00833] Compound 381 (formic acid salt) was prepared as a white solid utilizing the methods in Example 282 (Compound 378) from (R)-2-(((benzyloxy)carbonyl)amino)-4-((tert-butyl-
 10 butyldimethylsilyl)oxy)butanoic acid. LCMS (5-95 AB, ESI): $t_R = 0.694$, $[M + H]^+ = 913.7$.

Example 286: Synthesis of Compound 382


[00834] Compound 382 (formic acid salt) was prepared as a white solid utilizing the methods in Example 282 (Compound 378) from (S)-2-(((benzyloxy)carbonyl)amino)-4-((tert-butyl-
 15 butyldimethylsilyl)oxy)butanoic acid. LCMS (5-95 AB, ESI): $t_R = 0.689$, $[M + H]^+ = 913.9$.

Example 287: Synthesis of Compound 383


[00835] Compound 383 (formic acid salt) was prepared as a white solid utilizing the methods in Example 282 (Compound 378) from Cbz-Ser(O-*t*-Bu)-OH. LCMS (5-95 AB, ESI): $t_R = 0.741$, $[M + H]^+ = 927.8$

Example 288: Synthesis of Compound 384


[00836] Compound 384 (formic acid salt) was prepared as a white solid utilizing the methods in Example 282 (Compound 378). LCMS (5-95 AB, ESI): $t_R = 0.751$, $[M + H]^+ = 907.6$.

Example 289: Synthesis of Compound 385


[00837] Compound 385 (free base) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from (S)-5-amino-2-(((benzyloxy)carbonyl)amino)-5-oxopentanoic acid. LCMS (5-95 AB, ESI): $t_R = 0.573$, $[M + H]^+ = 921.0$.


Example 290: Synthesis of Compound 386

[00838] Compound 386 (formic acid salt) was prepared as a white solid utilizing the methods in Example 263 (Compound 359) from (S)-4-acetamido-2-
 5 (((benzyloxy)carbonyl)amino)butanoic acid. LCMS (5-95 AB, ESI): $t_R = 0.716$, $[M + H]^+ = 934.7$.

Example 291: Synthesis of Compound 387

[00839] Compound 387 (formic acid salt) was prepared as a white solid utilizing the methods in Example 282 (Compound 378). LCMS (Method 5-95 AB, ESI): $t_R = 0.757$ min, $[M+H]^+ = 921.6$.


Example 292: Synthesis of Compound 388



[00840] Starting from Compound 101-K, typical amide coupling (HATU) with (S)-2-(((9H-fluoren-9-yl)methoxy)carbonyl)(methyl)amino)-3-(tert-butoxy)propanoic acid and Fmoc removal (piperidine) procedure was followed to afford Compound 388-A (110 mg) as a white solid.

[00841] Compound 388 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from Compound 388-A. LCMS (5-95 AB, ESI): $t_R = 0.708$, $[M + H]^+ = 893.5$.

Example 293: Synthesis of Compound 389




[00842] To a solution of methyl 4-fluoro-2-methylbenzoate (200.0 mg, 1.19 mmol) in N, N-dimethylformamide (10 mL) were added 1-heptanethiol (0.91 mL, 5.95 mmol) and cesium carbonate (1937.6 mg, 5.95 mmol), and stirred at $60^\circ C$ for 16 h. The reaction was quenched by 5% aq. $KHSO_4$ till pH = 6, diluted with H_2O (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic layers were washed with water (40 mL x 2) and brine (20 mL), dried over Na_2SO_4 and concentrated. The residue was triturated with petroleum ether. The resulting white solid was filtered, washed with petroleum ether and dried to obtain 4-(heptylthio)-2-methylbenzoic acid (198 mg, 62.5% yield). ^1H NMR (400 MHz, CD_3OD): δ 7.84 (d, $J = 8.8$ Hz,

1H), 7.15-7.05 (m, 2H), 3.00 (t, $J = 7.2$ Hz, 2H), 2.55 (s, 3H), 1.75-1.60 (m, 2H), 1.50-1.35 (m, 2H), 1.35-1.20 (m, 6H), 0.90 (t, $J = 7.0$ Hz, 3H).

[00843] Compound 389 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): RT = 0.649 min, $[M+Na]^+ =$
5 908.5.


Example 294: Synthesis of Compound 390



[00844] To a solution of 4-heptylsulfanyl-2-methyl-benzoic acid (50.0 mg, 0.19 mmol) in
10 acetone (3 mL) was added potassium peroxymonosulfate (230.8 mg, 0.38 mmol) in water (1 mL). The mixture was stirred at 0°C for 2 h, quenched by saturated aqueous Na₂SO₃ solution and extracted with EtOAc (20 mL x 2). The combined organic layers were washed with water (40 mL x 2) and brine (20 mL), dried over Na₂SO₄ and concentrated to give 4-heptylsulfonyl-2-methyl-benzoic acid (50 mg, 89.3% yield) as a white solid which was used directly.



15 [00845] Compound 390 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.590$ min, $[M+H]^+ =$ 918.4.

Example 295: Synthesis of Compound 391



[00846] Compound 391 (formic acid salt) was prepared as a white solid utilizing the methods in Example 293 (Compound 389). LCMS (Method 5-95 AB, ESI): $t_R = 0.637$ min, $[M+Na]^+ = 894.4$.



Example 296: Synthesis of Compound 392



[00847] Step 1: To a solution of octanoic acid (5.0 g, 34.7 mmol) in THF (100 mL) was added potassium 3-methoxy-3-oxopropanoate (6.2 g, 38.1 mmol) and the mixture was stirred at room temperature for 2 h, followed by the addition of $MgCl_2$ (3.3 g, 34.7 mmol) and CDI (5.7 g, 36.4 mmol). The resulting mixture was heated to $60^\circ C$ for 3 h. After cooling to room temperature, the filtrate was concentrated and the residue was taken up by EtOAc (150 mL), which was washed with brine (150 mL x 2). The organic layer was dried over Na_2SO_4 , concentrated and the residue was purified by silica gel flash column to give methyl 3-oxodecanoate (4.4 g, 63.4% yield) as yellow oil.

[00848] Step 2: The General Method NaOH (NaOH ester hydrolysis) procedure was applied to methyl 3-oxodecanoate (200 mg, 1.0 mmol) to give 3-oxodecanoic acid (150 mg, 81% yield) as a white solid.



[00849] Compound 392 (formic acid salt) was prepared as a white solid utilizing the same methods used in the preparation of Compound 231 (Example 137) except 3-oxodecanoic acid was used in the coupling step. LCMS (5-95 AB, ESI): $t_R = 0.574$, $[M + H]^+ = 793.2$.

Example 297: Synthesis of Compound 393-P1 and Compound 393-P2

[00850] Step 1: To a mixture of KCN (2.59 g, 39.7 mmol) in EtOAc/MeOH (80 mL, v/v=1:1) was added sequentially with HOAc (2.39 g, 39.7 mmol) and nonanal (5.0 g, 35.2 mmol) at 0°C and the reaction was stirred at room temperature for 16 h. The volatiles were removed and the residue was poured into H₂O (50 mL), which was extracted with EtOAc (40 mL x 3). The combined organic layers were dried over Na₂SO₄, concentrated and the residue was purified by silica gel flash column to give (±)-2-hydroxydecanenitrile (5.8 g, 97.5% yield) as colorless oil.

[00851] Step 2: The acidic hydrolysis of a nitrile to an acid is described and is referred to as General Method Nitrile Hydrolysis. To a solution of (±)-2-hydroxydecanenitrile (2.0 g, 11.8 mmol) in 1,4-dioxane (10 mL) was added with 36% HCl solution (10 mL) and the mixture was heated to 100°C while stirring and stirred at the same temperature for 12 h. The volatiles were removed to afford (±)-2-hydroxydecanoic acid (1.6 g, 60.3% yield) as a white solid, which was used directly for the next step.



[00852] Compound 393-P1 (formic acid salt) and Compound 393-P2 were prepared as white solids utilizing the same methods used in the preparation of Compound 231 (Example 137) except (±)-2-hydroxydecanoic acid was used in the coupling step. The diastereomers were separated on prep-HPLC. Data for Compound 393-P1: LCMS (5-95 AB, ESI): $t_R = 0.727$, $[M + H]^+ = 795.4$. Data for Compound 393-P2: LCMS (5-95 AB, ESI): $t_R = 0.730$, $[M + H]^+ = 795.4$.

Example 298: Synthesis of Compound 394

[00853] Step 1: To a solution of (±)-2-hydroxydecanenitrile (3.0 g, 17.7 mmol) in DCM (20 mL) was added DAST (6.4 g, 39.9 mmol) dropwise at -78°C within 0.5 h. The reaction was warmed up gradually to the room temperature and stirred at the same temperature for 36 h. The volatiles were removed and the residue was taken up by EtOAc (100 mL), which was washed with brine (100 mL x 2). The organic layer was dried with Na₂SO₄, concentrated and the residue was purified by silica gel flash column to afford (±)-2-fluorodecanenitrile (1.5 g, 49.4% yield) as colorless oil.

[00854] Step 2: (±)-2-Fluorodecanenitrile (490 mg, 2.9 mmol) was subject to General Method Nitrile hydrolysis to afford (±)-2-fluorodecanoic acid (450 mg, 83% yields) as a light yellow solid.

[00855] Compound 394-P1 (formic acid salt) and Compound 394-P2 were prepared as white solids utilizing the same methods used in the preparation of Compound 231 (Example 137) except (±)-2-fluorodecanoic acid was used in the coupling step. The diastereomers were separated on prep-HPLC. Data for Compound 394-P1: LCMS (5-95 AB, ESI): *t_R* = 0.760, [M + H]⁺ = 797.6. Data for Compound 394-P2: LCMS (5-95 AB, ESI): *t_R* = 0.773, [M + H]⁺ = 797.4.



Example 299: Synthesis of Compound 395

[00856] Step 1: To a solution of hexanamide (500 mg, 4.4 mmol) in TBME (40 mL) was added phosphorous penta-sulfide (241 mg, 1.1 mmol) at room temperature and the mixture was stirred at the same temperature for 18 h. The volatiles were removed and the residue was purified on silica gel flash column to afford hexanethioamide (420 mg, 73.7% yield) as a pale yellow solid.

[00857] Step 2: To a solution of hexanethioamide (420 mg, 3.2 mmol) in EtOH (6 mL) was added ethyl 3-bromo-2-oxobutanoate (669 mg, 3.2 mmol) dropwise under N₂ at 0°C and the mixture was gradually warmed up and stirred at room temperature for 18 h. The volatiles were removed and the residue was taken up by EtOAc (50 mL), which was washed with brine (50 mL). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified on silica gel flash column to afford ethyl 5-methyl-2-pentylthiazole-4-carboxylate (390 mg, 50.5% yield) as pale green oil.

[00858] Step 3: The NaOH ester hydrolysis procedure (General Method NaOH) was applied to ethyl 5-methyl-2-pentylthiazole-4-carboxylate (390 mg, 1.62 mmol) to afford 5-methyl-2-pentylthiazole-4-carboxylic acid (250 mg, 72.5% yield) as a pale yellow solid.

[00859] Compound 395 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): t_R = 0.568, [M + H]⁺ = 833.3.

Example 300: Synthesis of Compound 396

[00860] Step 1: To a solution of (S)-methyl 2-amino-4-((tert-butoxycarbonyl)amino)butanoate (2.8 g, 12.0 mmol), 2-nitrobenzenesulfonyl chloride (4.0 g, 18.0 mmol) in DCM (40 mL) was added Et₃N (3.7 g, 36.0 mmol) at 0°C and the mixture was stirred at the same temperature for 2h. The reaction mixture with then washed with 1 N HCl (40 mL), saturated NaHCO₃ solution (40 mL) and brine (40 mL). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified by silica gel flash column to give (S)-methyl 4-((tert-butoxycarbonyl)amino)-2-(2-nitrophenylsulfonamido)butanoate (4.5 g, 89.4% yield) as pale-green oil.

[00861] Step 2: To a solution of (S)-methyl 4-((tert-butoxycarbonyl)amino)-2-(2-nitrophenylsulfonamido)butanoate (2.3 g, 5.5 mmol), K₂CO₃ (3.8 g, 27.5 mmol) in DMF (10 mL) was added methyl iodide (3.9 g, 27.5 mmol) and the mixture was stirred for at room temperature for 1.5 h. The reaction was added with EtOAc (100 mL) and the filtration was washed with brine (100 mL x 3). The organic layer was dried over Na₂SO₄, concentrated to give (S)-methyl 4-((tert-butoxycarbonyl)amino)-2-(N-methyl-2-nitrophenylsulfonamido)butanoate (2.3 g, 96.7% yield) as a pale-yellow oil.

[00862] Step 3: To a solution of (S)-methyl 4-((tert-butoxycarbonyl)amino)-2-(N-methyl-2-nitrophenylsulfonamido)butanoate (2.2 g, 5.1 mmol), K₂CO₃ (4.2 g, 30.6 mmol) in DMF (15 mL) was added thiophenol (6.3 g, 57.1 mmol) and the mixture was stirred at room temperature for 3 h. TLC(50% ethyl acetate in petroleum ether, R_f=0.1) showed the start material was


consumed. The reaction was added with EtOAc (100 mL) and the filtration was washed with brine (100 mL x 3). The organic layer was dried over Na₂SO₄, concentrated to give (S)-methyl 4-((tert-butoxycarbonyl)amino)-2-(methylamino)butanoate (830 mg, 66.1% yield) as yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 5.15-5.21 (m, 1H), 3.73 (s, 3H), 3.17-3.32 (m, 3H), 2.35 (s, 3H), 1.84-1.89 (m, 1H), 1.66-1.73 (m, 1H), 1.43 (s, 9H)

[00863] Step 4: To a solution of (S)-methyl 4-((tert-butoxycarbonyl)amino)-2-(methylamino)butanoate (730 mg, 2.96 mmol) and Et₃N (750 mg, 7.41 mmol) in DCM (50 mL) was added benzyl chloroformate (758 mg, 4.45 mmol) dropwise at 0°C and the mixture was warmed up to temperature while stirring and stirred at the same temperature for 2 h. The mixture was washed 1N HCl (30 mL) and brine (30 mL x 2) sequentially. The organic layer was dried over Na₂SO₄, concentrated and the residue was purified by silica gel flash column to give (S)-methyl 2-(((benzyloxy)carbonyl)(methyl)amino)-4-((tert-butoxycarbonyl)amino)butanoate (1.0 g, 88.7% yield) as a pale-yellow oil.


[00864] Step 5: Typical ester hydrolysis (NaOH/MeOH/H₂O) was applied to (S)-methyl 2-(((benzyloxy)carbonyl)(methyl)amino)-4-((tert-butoxycarbonyl)amino)butanoate (1.0 g, 2.63 mmol) to afford (S)-2-(((benzyloxy)carbonyl)(methyl)amino)-4-((tert-butoxycarbonyl)amino)butanoic acid (800 mg, 83% yield) as pale-yellow oil.

[00865] Compound 396 (6.0 mg) as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): t_R = 0.739, [M + H]⁺ = 854.6; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.49 (brs, 2H), 7.35-7.22 (m, 6H), 7.19 (d, *J*=8.4 Hz, 1H), 7.09 (d, *J*=8.4 Hz, 1H), 6.90 (brs, 1H), 6.81 (brs, 1H), 6.24 (s, 1H), 5.57 (s, 2H), 4.77-4.75 (m, 1H), 4.26-4.17 (m, 4H), 4.20 (s, 2H), 3.30-3.20 (m, 1H), 3.18-3.14 (m, 4H), 3.08-3.03 (m, 2H), 3.00 (s, 3H), 2.82 (s, 3H), 2.72-2.64 (m, 3H), 2.40-2.25 (m, 1H), 2.20-2.16 (m, 1H), 1.64 (s, 3H), 1.39-1.25 (m, 10H), 0.90 (t, *J*=6.6 Hz, 3H).


Example 301: Synthesis of Compound 397




[00866] Compound 397 (formic acid salt) was prepared as a white solid utilizing the methods in Example 300. LCMS (5-95 AB, ESI): t_R = 0.688, [M + H]⁺ = 826.5.


Example 302: Synthesis of Compound 398

[00867] Compound 398 (formic acid salt) was prepared as a white solid utilizing the methods in Example 300. LCMS (5-95 AB, ESI): $t_R = 0.753$, $[M + H]^+ = 834.6$.

5 Example 303: Synthesis of Compound 399

[00868] Compound 399 (formic acid salt) was prepared as a white solid utilizing the methods in Example 300. LCMS (5-95 AB, ESI): $t_R = 0.585$, $[M + H]^+ = 893.6$

Example 304: Synthesis of Compound 400





[00869] Compound 400-A is an intermediate in the preparation of Compound 231 (Example 137).

[00870] To a stirred solution of Compound 400-A (150 mg, 0.16 mmol) in DMF (5 mL) was added 1-bromodecane (697 mg, 3.15 mmol) and K_2CO_3 (435 mg, 3.15 mmol) and the mixture was stirred at room temperature for 2 d. The reaction mixture was then added with EtOAc (50 mL), which was washed with brine (50 mL x 2). The organic layer was dried over Na_2SO_4 , concentrated and the residue was purified by prep-TLC to afford Compound 400-B (70 mg, 44.6% yield) as a white solid. LCMS (5-95 AB, ESI): $t_R = 0.851$, $[\text{M} + \text{H}]^+ = 997.5$

[00871] Compound 400 (formic acid salt) was prepared as a white solid utilizing the same methods used in the preparation of Compound 101 (Example 7) starting from Compound 400-B. LCMS (5-95 AB, ESI): $t_R = 0.597$, $[\text{M} + \text{H}]^+ = 765.3$; ^1H NMR (400 MHz, $\text{MeOH}-d_4$) δ 7.31-7.28 (m, 2H), 7.20 (d, $J=8.0$ Hz, 1H), 7.12 (d, $J=8.0$ Hz, 1H), 6.91 (brs, 1H), 6.82 (brs, 1H), 6.39 (s, 1H), 4.80-4.75 (m, 1H), 4.27-4.15 (m, 6H), 4.21 (s, 2H), 4.05-3.95 (m, 1H), 3.85-3.77 (m, 1H), 3.25-3.16 (m, 7H), 2.93-2.80 (m, 1H), 2.89 (s, 3H), 1.73 (brs, 2H), 1.41-1.20 (m, 17H), 0.93 (t, $J=6.0$ Hz, 3H). LCMS (5-95 AB, ESI): $t_R = 0.597$, $[\text{M} + \text{H}]^+ = 765.3$; ^1H NMR (400 MHz, $\text{MeOH}-d_4$) δ 7.31-7.28 (m, 2H), 7.20 (d, $J=8.0$ Hz, 1H), 7.12 (d, $J=8.0$ Hz, 1H), 6.91 (brs, 1H), 6.82 (brs, 1H), 6.39 (s, 1H), 4.80-4.75 (m, 1H), 4.27-4.15 (m, 6H), 4.21 (s, 2H), 4.05-3.95 (m, 1H), 3.85-3.77 (m, 1H), 3.25-3.16 (m, 7H), 2.93-2.80 (m, 1H), 2.89 (s, 3H), 1.73 (brs, 2H), 1.41-1.20 (m, 17H), 0.93 (t, $J=6.0$ Hz, 3H).

Example 305: Synthesis of Compound 401






[00872] Compound 401-A is an intermediate in the synthesis of Compound 401 using the methods previously described.


[00873] To a stirred solution of Compound 401-A (100 mg, 0.11 mmol) in DMF (3 mL) was added tert-butyl (3-bromopropyl)carbamate (34 mg, 0.16 mmol), K₂CO₃ (37 mg, 0.27 mmol).

The mixture was then stirred at room temperature for 72 h. The reaction mixture was added with EtOAc (30 mL), which was washed with brine (30 mL x 2). The organic layer was dried over MgSO₄, concentrated and the residue was purified by prep-TLC to give Compound 401-B (30 mg, 25.7% yield) as a white solid. LCMS (5-95 AB, ESI): t_R = 0.833, [M + H]⁺ = 1087.7

[00874] Compound 401 (formic acid salt) was prepared as a white solid utilizing the same methods used in the preparation of Compound 257 (Example 162). LCMS (5-95 AB, ESI): $t_R = 0.737$, $[M + H]^+ = 927.1$.

Example 306: Synthesis of Compound 402






[00875] To a solution of Compound 101-K (100 mg, 0.20 mmol) in DCM (2 mL) was added CDI (20 mg, 0.20 mmol) at room temperature and the mixture was stirred overnight. To the mixture was then added n-C₈H₁₇NH₂ (42 mg, 0.33 mmol) at the same temperature and the resulting mixture was stirred for another 20 h at room temperature. The reaction mixture was diluted with DCM (50 mL), and then washed sequentially with saturated citric acid, saturated NaHCO₃ and brine (30 mL each). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified on Pre-TLC to obtain Compound 402-A (85 mg, 72.9% yield) as a white solid. LCMS (5-95 AB, ESI): t_R = 1.072, [M + H]⁺ = 1070.0.



[00876] Starting from Compound 402-A, Compound 402 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): t_R = 0.751, [M + H]⁺ = 793.9; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.50 (brs, 2H, HCOOH), 7.38-7.25 (m, 2H), 7.20-7.11 (m, 2H), 6.90 (brs, 1H), 6.83 (brs, 1H), 6.31 (s, 1H), 5.00-4.79 (m, 3H), 4.30-4.20 (m, 4H), 4.21 (s, 2H), 3.30-2.88 (m, 10H), 2.87 (s, 3H), 2.20-2.10 (m, 1H), 2.00-1.85 (m, 1H), 1.55-1.45 (m, 2H), 1.40-1.30 (m, 13H), 0.93 (t, *J*=6.8 Hz, 3H).

Example 307: Synthesis of Compound 403




[00877] Compound 403 (formic acid salt) was prepared as a white solid utilizing the methods in Example 306 (Compound 402). LCMS (5-95 AB, ESI): $t_R = 0.763$, $[M + H]^+ = 835.5$.

Example 308: Synthesis of Compound 404



5



[00878] Step 1: A solution of 4-hexyl-2-methylbenzoic acid (300 mg, 1.4 mmol), diphenylphosphoryl azide (450 mg, 1.6 mmol), benzyl alcohol (442 mg, 4.1 mmol) and Et₃N (386 mg, 3.8 mmol) in toluene (10 mL) was stirred at 100 °C for 16 h. The reaction mixture was diluted with water (30 mL), which was extracted by EtOAc (30 mL x 2). The combined organic layers were washed with brine (50 mL x 2), dried over MgSO₄, concentrated and the residue was purified by silica gel flash column to give benzyl (4-hexyl-2-methylphenyl)carbamate (270 mg, 60.9% yield) as a white solid. LCMS (5-95 AB, ESI): $t_R = 1.099$, $[M + H]^+ = 326.0$.

[00879] Step 2: The standard hydrogenation (Pd/C) procedure (General Method Hydrogenation) was applied to benzyl (4-hexyl-2-methylphenyl)carbamate (270 mg, 0.83 mmol) to afford 4-hexyl-2-methylaniline (125 mg, 79% yield) as a white solid.


[00880] Step 3: To a solution of 4-hexyl-2-methylaniline (30 mg, 0.16 mmol) in THF (1 mL) was added phenyl chloroformate (27 mg, 0.17 mmol) in portions and the mixture was stirred at room temperature for 1 h. The reaction was quenched with water (5 mL), the mixture was extracted by EtOAc (5 mL x 3). The combined organic layers were dried over Na₂SO₄,

concentrated and the residue was purified by silica gel flash column to afford phenyl (4-hexyl-2-methylphenyl)carbamate (40 mg, 81.9% yield) as colorless oil. LCMS (5-95 AB, ESI): t_R = 0.960, $[M + H]^+ = 311.9$

[00881] Step 4: To a solution of Compound 101-K (80 mg, 0.09 mmol) in CHCl₃ (3 mL) was added Et₃N (27 mg, 0.26 mmol) and phenyl (4-hexyl-2-methylphenyl)carbamate (35 mg, 0.11 mmol) and the mixture was heated to 80°C while stirring and stirred at the same temperature for 3 h. The volatiles were removed and the residue was purified by prep-TLC to give Compound 404A-A (90 mg, 90.9% yield) as an off-white solid. LCMS (5-95 AB, ESI): t_R = 0.983, $[M + H]^+ = 1132.5$.

[00882] Starting from Compound 404A-A, Compound 404 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): t_R = 0.627, $[M + H]^+ = 855.5$; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.46 (brs, 2H, HCOOH), 7.35-7.27 (m, 2H), 7.24 (d, $J=8.0$ Hz, 1H), 7.15 (d, $J=8.0$ Hz, 1H), 7.09 (d, $J=8.0$ Hz, 1H), 7.03 (brs, 1H), 6.98 (d, $J=8.0$ Hz, 1H), 6.90 (brs, 1H), 6.81 (brs, 1H), 6.31 (s, 1H), 4.80-4.76 (m, 3H), 4.28-4.17 (m, 4H), 4.19 (s, 2H), 3.27-3.20 (m, 4H), 3.18-3.04 (m, 4H), 2.88 (s, 3H), 2.55 (t, $J=7.2$ Hz, 2H), 2.26-2.15 (m, 1H), 2.24 (s, 3H), 2.03-1.95 (m, 1H), 1.63-1.55 (m, 2H), 1.37-1.26 (m, 9H), 0.89 (t, $J=6.4$ Hz, 3H).

Example 309: Synthesis of Compound 405





[00883] Step 1: To a solution of 4-bromo-2-methylbenzoic acid (10.7 g, 50 mmol) in THF (250 mL) at -78°C was added 2M n-BuLi in hexane (6.0 mL) dropwise and the mixture was stirred at the same temperature for 3 h. DMF (3.85 mL, 50 mmol) was then added to the mixture at -78°C and the resulting mixture was warmed up to room temperature slowly while stirring
 5 and stirred at the same temperature for 1.5 h. The reaction was quenched with 1N HCl (200 mL), which was extracted by EtOAc (200 mL x 3). The combined organic layers were dried over Na₂SO₄ and concentrated to give the crude, which was triturated with petroleum ether to afford 4-formyl-2-methylbenzoic acid (3.0 g, 36.7% yield) as an off-white solid.

[00884] Step 2: To a solution of 4-formyl-2-methylbenzoic acid (762 mg, 4.6 mmol), nonan-5-one (600 mg, 4.2 mmol) in DCM (10 mL) was added boron trifluoride diethyl etherate (1.07 mL, 8.44 mmol) and the mixture was stirred at 45°C for 16 h. DCM (30 mL) was added to the reaction mixture, which was then washed with brine (50 mL x 2). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified by HPLC to afford (E)-2-methyl-4-(pent-1-en-1-yl)benzoic acid (400 mg, 46.4% yield) as an off-white solid. LCMS (10-80 AB 2
 15 min, ESI): t_R = 1.221, [M + H]⁺ = 205.2.

[00885] Compound 405 (free base) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): t_R = 0.700, [M + H]⁺ = 824.5.

Example 310: Synthesis of Compound 406





[00886] To a solution of methyl isobutyrate (1.0 g, 9.8 mmol) in THF (15 mL) was added 2M LDA in THF (5.9 mL, 11.8 mmol) dropwise at -78°C and the mixture was stirred for at the same temperature for 0.5 h. 1-bromodecane (2.6 g, 11.8 mmol) and HMPA (2.1 g, 11.8 mmol) was then added at -78°C and the resulting mixture was slowly warmed up to room
 25 temperature while stirring and stirred for another 3 h at room temperature. The reaction was

quenched with saturated NH_4Cl solution (30 mL), which extracted with EtOAc (30 mL x 3). The combined organic layers were dried over Na_2SO_4 , concentrated and the residue was purified by silica gel flash column to give methyl 2,2-dimethyldodecanoate (1.7 mg, 71.6% yield) as colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 3.64 (s, 3H), 1.50-1.46 (m, 2H), 1.30-1.23 (m, 16H), 1.14 (s, 6H), 0.86 (t, $J = 6.4$ Hz, 3H).

[00887] Step 2: Ester hydrolysis procedure with NaOH (General Method NaOH) was applied to methyl 2,2-dimethyldodecanoate (500 mg, 0.45 mmol) to afford 2,2-dimethyldodecanoic acid (330 mg, 70.1% yield) as a light yellow solid.

[00888] Compound 406 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): $t_R = 0.609$, $[\text{M} + \text{H}]^+ = 848.5$

Example 311: Synthesis of Compound 407



[00889] Step 1: A solution of methyl 4-bromo-2-hydroxybenzoate (600 mg, 2.8 mmol) and octylboronic acid (874 mg, 5.6 mmol), $\text{Pd}(\text{PPh}_3)_4$ (320 mg, 0.28 mmol) and K_2CO_3 (1.15 g, 8.3 mmol) in toluene (16 mL) was stirred at 100°C under N_2 for 16 h. The volatiles were removed and the residue was purified by silica gel flash column to give methyl 2-hydroxy-4-octylbenzoate (706 mg, 96.6% yield) as colorless oil.



[00890] Step 2: To a solution of methyl 2-hydroxy-4-octylbenzoate (300 mg, 1.13 mmol) in DMF (8 mL), 2-bromo acetamide (188 mg, 1.36 mmol) was added Cs_2CO_3 (740 mg, 2.27 mmol) and the mixture was stirred at room temperature for 16 h. The reaction mixture was diluted with

EtOAc (50 mL), which was washed by brine (50 mL x 2). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified by silica gel flash column to obtain methyl 2-(2-amino-2-oxoethoxy)-4-octylbenzoate (172 mg, 47.2% yield) as a white solid. LCMS (5-95 AB, ESI): $t_R = 1.025$, $[M + H]^+ = 322.0$

- 5 **[00891]** Step 3: Typical ester hydrolysis (NaOH/THF) was applied to methyl 2-(2-amino-2-oxoethoxy)-4-octylbenzoate (70 mg, 0.22 mmol) to afford 2-(2-amino-2-oxoethoxy)-4-octylbenzoic acid (68 mg, 99% yield) as a white solid.

- [00892]** The methods in Example 7 (Compound 101) were followed to afford the Compound 407 (21.4 mg) as a white solid. LCMS (5-95 AB, ESI): $t_R = 0.756$, $[M + H]^+ = 927.5$; ¹H NMR (400 MHz, MeOH-*d*₄) δ 7.81 (d, $J=8.0$ Hz, 1H), 7.32 (d, $J=8.0$ Hz, 1H), 7.23 (d, $J=8.0$ Hz, 1H), 7.15 (d, $J=8.0$ Hz, 1H), 7.09 (d, $J=8.0$ Hz, 1H), 6.97 (d, $J=8.0$ Hz, 1H), 6.93 (brs, 1H), 6.88 (brs, 1H), 6.81 (brs, 1H), 6.34 (s, 1H), 4.90-4.75 (m, 5H), 4.23-4.18 (m, 4H), 4.19 (s, 2H), 3.35-3.30 (m, 4H), 3.20-3.13 (m, 4H), 2.90 (s, 3H), 2.71-2.65 (m, 2H), 2.28-2.15 (m, 2H), 1.70-1.55 (m, 2H), 1.40-1.20 (m, 13H), 0.89 (t, $J=6.0$ Hz, 3H).

15 **Example 312: Synthesis of Compound 408**




- [00893]** Step 1: Na (179 mg, 7.8 mmol) was added to EtOH (10 mL) while stirring in portions and the mixture was stirred at room temperature for 2 h, followed by the sequential addition of diethyl oxalate (1.14 g, 7.8 mmol) and octan-2-one (1.0 g, 7.8 mmol). The resulting mixture was stirred at room temperature for 16 h. The reaction mixture was added with EtOAc (50 mL), which was washed with brine (50 mL x 2). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified on silica gel flash column to afford ethyl 2,4-dioxodecanoate as a light yellow oil (900 mg).

[00894] Step 2: A solution of ethyl 2,4-dioxodecanoate (700 mg, 3.1 mmol) and methyl hydrazine (155 mg, 3.4 mmol) in EtOH (10 mL) was stirred at 80°C for 3 h. The volatiles were removed and the residue was taken up in EtOAc (50 mL), which was washed with brine (30 mL x 2). The organic layer was dried over MgSO₄, concentrated and the residue was purified by silica gel flash column to afford ethyl 3-hexyl-1-methyl-1H-pyrazole-5-carboxylate (280 mg, 38.3% yield) as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 6.61 (s, 1H), 4.31 (q, *J*=7.2 Hz, 2H), 4.12 (s, 3H), 2.61-2.54 (m, 2H), 1.66-1.56 (m, 2H), 1.34 (t, *J*=7.2 Hz, 3H), 1.33-1.25 (m, 6H), 0.89 (t, *J*=6.8 Hz, 3H).

[00895] Step 3: Typical ester hydrolysis (NaOH/THF) procedure was applied to ethyl 3-hexyl-1-methyl-1H-pyrazole-5-carboxylate (260, 1.1 mmol) to afford 3-hexyl-1-methyl-1H-pyrazole-5-carboxylic acid (220 mg, 96% yield) as a white solid.

[00896] The methods in Example 7 (Compound 101) were followed to afford Compound 408 (9.0 mg) as a white solid. LCMS (5-95 AB, ESI): *t*_R = 0.686, [M + H]⁺ = 830.5.

Example 313: Synthesis of Compound 409




[00897] Compound 409-A is prepared during the synthesis of Compound 193 (Example 99) utilizing the methods in Example 112 (Compound 206). LCMS (Method 5-95AB): *t*_R = 1.017 min, [M + H]⁺ = 1171.0.

[00898] A mixture of Compound 409-A (30.0 mg, 0.03 mmol), potassium carbonate (10.6 mg, 0.08 mmol) and tert-butyl bromoacetate (15.0 mg, 0.08 mmol) in N,N-dimethylformamide (3 mL) was stirred at 15 °C for 2 h. The reaction was poured into water (3 mL) and extracted with EtOAc (10 mL). The organic extract was washed with water (10 mL x 2) and brine (10 mL),
 5 dried over MgSO₄ and concentrated. The residue was purified by prep-TLC (5% methanol in DCM, R_f = 0.4) to afford Compound 409-B (20 mg, 60.7% yield) as a white solid. LCMS (Method 5-95AB): t_R = 1.035 min, [M + H]⁺ = 1285.2.


[00899] Compound 409 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95AB): t_R = 0.627 min, [M + H]⁺ = 929.0.

10 Example 314: Synthesis of Compound 410



[00900] Compound 410-A is prepared using the previously described methods from 2-(2-(tert-butoxy)-2-oxoethoxy)-4-octylbenzoic acid. During coupling to the carboxylic acid utilizing
 15 General Method 4 (Example 5), bis-addition of the nitrile occurred to form Compound 410-B.

[00901] Compound 410 (formic acid salt) was prepared as a white solid from Compound 410-B utilizing the TFA hydrolysis conditions in Example 7. LCMS (Method 5-95AB): t_R = 0.617 min, [M + H]⁺ = 966.3.


Example 315: Synthesis of Compound 411


[00902] Step 1: A mixture of pyrazole (100 mg, 1.47 mmol), methyl 4-fluoro-2-

5 methylbenzoate (123.51 mg, 0.73 mmol) and potassium carbonate (609.0 mg, 4.41 mmol) in N,N-dimethylformamide (5 mL) was stirred at 60 °C for 16 h. The reaction was poured into water (5 mL) and extracted with EtOAc (20 mL). The organic extract was washed with water (20 mL x 2) and brine (10 mL), dried over MgSO₄ and concentrated. The residue was purified by prep-TLC (33% ethyl acetate in petroleum ether, R_f = 0.3) to afford methyl 2-methyl-4-(1H-pyrazol-1-yl)benzoate (50 mg, 15.7% yield) as a white solid. LCMS (Method 5-95AB): t_R = 0.732 min, [M + H]⁺ = 216.9.

[00903] Step 2: Methyl 2-methyl-4-(1H-pyrazol-1-yl)benzoate was hydrolyzed as previously described (General Method NaOH) to give 2-methyl-4-(1H-pyrazol-1-yl)benzoic acid.

[00904] Compound 315 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 0-60AB): t_R = 0.830 min/ 2 min, [M + H]⁺ = 822.3.

Example 316: Synthesis of Compound 412



[00905] Step 1: A mixture of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.15 g, 6.01 mmol), N,N-diisopropylethylamine (2.48 mL, 15.03 mmol), 1-hydroxybenzotriazole (0.81 g, 6.01 mmol), 4-bromo-3-chlorobenzoic acid (1.18 g, 5.01 mmol) and N,O-dimethylhydroxylamine hydrochloride (0.98 g, 10.02 mmol) in N,N-dimethylformamide (10 mL) was stirred at 20 °C for 16 h. The reaction was quenched with water (20 mL), and extracted with EtOAc (25 mL x 2). The combined organic layers were concentrated and purified by column (30% EtOAc in petroleum ether, $R_f = 0.3$) to give 4-bromo-3-chloro-N-methoxy-N-methyl-benzamide (1.38 g, 98.9% yield) as a yellow oil.

[00906] Step 2: To a solution of heptylmagnesium bromide (2.37 mL, 2.37 mmol) in tetrahydrofuran (3 mL) was added 4-bromo-3-chloro-N-methoxy-N-methyl-benzamide (300.0 mg, 1.08 mmol) under N_2 at 0 °C and stirred at room temperature for 16 h. The reaction mixture was quenched with aq. NH_4Cl (1 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were dried over anhydrous sodium sulfate, and concentrated in vacuo. The residue was purified on silica gel column eluted with ethyl acetate/petroleum ether (1:10) to afford 1-(4-bromo-3-chloro-phenyl)octan-1-one (310 mg, 90.6% yield) as a colorless oil. ^1H NMR (400 MHz, $CDCl_3$): δ 8.00 (d, $J = 1.6$ Hz, 1H), 7.75 - 7.60 (m, 2H), 2.90 (t, $J = 7.6$ Hz, 2H), 1.70 (quint, $J = 7.3$ Hz, 2H), 1.45 - 1.20 (m, 8H), 0.87 (t, $J = 7.0$ Hz, 3H).

[00907] Step 3: A mixture of 1-(4-bromo-3-chloro-phenyl)octan-1-one (150.0 mg, 0.47 mmol) in deoxo-fluor(R) (2.14 mL, 11.81 mmol) was stirred at 80 °C for 16 h. The mixture was quenched with aq. $Na_2S_2O_3$ (3 mL) and aq. $NaHCO_3$ (3 mL) at 0 °C and extracted with petroleum ether (10 mL x 3). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated. The residue was purified by prep-TLC (5% EtOAc in petroleum ether, $R_f = 0.5$) to afford 1-bromo-2-chloro-4-(1,1-difluorooctyl)benzene (54 mg, 33.7% yield) as a colorless oil.



[00908] Step 4: A solution of 1-bromo-2-chloro-4-(1,1-difluorooctyl)benzene (50.0 mg, 0.15 mmol) and triethylamine (44.7 mg, 0.44 mmol) in methanol (10 mL) was treated with 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (10.8 mg, 0.01 mmol). The resulting mixture was stirred at 80 °C for 16 hours under CO (45 psi). The reaction mixture was

concentrated and the residue was purified by silica gel column (10% ethyl acetate in petroleum) to afford methyl 2-chloro-4-(1,1-difluorooctyl)benzoate (30 mg, 63.9% yield) as a colorless oil.

[00909] Step 5: Methyl 2-chloro-4-(1,1-difluorooctyl)benzoate (30.0 mg, 0.09 mmol) was hydrolyzed as previously described (General Method NaOH) to give crude 2-chloro-4-(1,1-difluorooctyl)benzoic acid (25 mg, 87.2% yield) as a white solid.

[00910] Compound 412 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95AB): $t_R = 0.664$ min, $[M+Na]^+ = 946.4$.


Example 317: Synthesis of Compound 413




[00911] 1-(4-Bromo-3-chloro-phenyl)octan-1-one has been described in Example 316. Using the methods in Example 316 (carbonylation followed by NaOH ester hydrolysis with General Method NaOH), 2-chloro-4-octanoylbenzoic acid was synthesized.

[00912] Compound 413 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95AB): $t_R = 0.745$ min, $[M+H]^+ = 902.7$.

Example 318: Synthesis of Compound 414






[00913] Step 1: To a solution of methyl 2-chloro-4-octylbenzoate (350.0 mg, 1.24 mmol) in 1,4-dioxane (2 mL) were added t-butyl acrylate (174.5 mg, 1.36 mmol), bis(tri-
 5 butylphosphine)palladium(0) (9.5 mg, 0.02 mmol), tris(dibenzylideneacetone)dipalladium(0) (34.0 mg, 0.04 mmol) and N,N-dicyclohexylmethylamine (265.9 mg, 1.36 mmol). The mixture was stirred at 25 °C for 16 h under N₂ and filtered. The filtrate was diluted with water (10 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The residue was purified by prep-TLC (10% EtOAc in petroleum
 10 ether, R_f = 0.7) to afford methyl 2-[(E)-3-tert-butoxy-3-oxo-prop-1-enyl]-4-octylbenzoate (290 mg, 62.6% yield) as a yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 8.35 (d, J = 16.0 Hz, 1H), 7.86 (d, J = 8.0 Hz, 1H), 7.39 (s, 1H), 7.22 (d, J = 8.0 Hz, 1H), 6.23 (d, J = 15.9 Hz, 1H), 3.91 (s, 3H), 2.64 (t, J = 7.8 Hz, 2H), 1.65-1.55 (m, 2H), 1.54 (s, 9H), 1.33 - 1.22 (m, 10H), 0.88 (t, J = 6.6 Hz, 3H).

[00914] Step 2: To a solution of methyl 2-[(E)-3-tert-butoxy-3-oxo-prop-1-enyl]-4-octyl-
 15 benzoate (290.0 mg, 0.77 mmol) in 1,2-dichloroethane (2 mL) was added trimethyltin hydroxide (1400.2 mg, 7.74 mmol) and the mixture was stirred at 80 °C for 16 h. LCMS (5-95AB/1.5min): t_R = 1.158 min, [M+Na]⁺ 383.1 showed 36% of DP and 51% of SM. The mixture was diluted with 0.1N KHSO₄ (5 mL) and extracted with EtOAc (10 mL x 3). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified
 20 by prep-TLC (50% EtOAc in petroleum ether, R_f = 0.2) to afford 2-[(E)-3-tert-butoxy-3-oxo-prop-1-enyl]-4-octylbenzoic acid (80 mg, 28.7% yield) as a yellow solid.

[00915] Compound 414 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95AB): t_R = 0.760 min, [M+H]⁺ = 924.5.

Example 319: Synthesis of Compound 415




[00916] Compound 415-A is prepared using methods previously described.


[00917] Compound 415-A (57.6 mg, 0.05 mmol), was dissolved in anhydrous THF (2 mL) and cooled the reaction mixture to 0 °C in ice bath. Isobutyl chloroformate (10 µL, 0.075 mmol) followed by N-methyl morpholine (16 µL, 0.15 mmol) was added under N₂ atm. The reaction mixture was stirred for 30 min. After the starting material was consumed (monitored by TLC) (R)-2-aminopropanenitrile (7 mg, 0.1 mmol) in THF (1.0 mL) was added and the reaction mixture was stirred for 4h. After completion of the reaction (monitored by LCMS) the reaction mixture was quenched with saturated NH₄Cl solution (1.0 mL). The reaction mixture was diluted with brine (2 mL) and extracted with EtOAc and the combined organic layers washed with brine. The organic layer was dried over anhydrous Na₂SO₄, filtered and the solvent was removed under vacuum.

[00918] The residue was subjected for global Boc de-protection (TFA/HFIP) procedure (Example 7) was followed to afford the title compound (5.0 mg) as a white solid. LCMS

(Method 5-95 AB, ESI): $t_R = 0.718$, $[M + H]^+ = 891.6$; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.52 (brs, 4H), 7.27-7.22 (m, 2H), 7.17 (d, *J*=8.0 Hz, 1H), 7.09 (d, *J*=8.0 Hz, 1H), 6.87 (brs, 1H), 6.83(brs, 1H), 6.34 (s, 1H), 4.80-4.76 (m, 3H), 4.27-4.22 (m, 5H), 3.27-3.23 (m, 6H), 3.16-3.12 (m, 2H), 3.02 (t, *J*=7.6 Hz, 2H), 2.94 (t, *J*=7.6 Hz, 2H), 2.88 (s, 2H), 2.74 (brs, 2H), 1.87(brs, 2H) 1.71(m, 6H), 1.54 (d, *J*=8.0 Hz, 3H), 1.38-1.31(m, 18H), 0.91-0.88(m, 3H).

Example 320: Synthesis of Compound 416








[00919] Step 1: To a solution of Compound 416-A (500 mg, 0.5 mmol) in dioxane (20 mL) was added 1M NaOH (10 mL, 10 mmol) and (Boc)₂O (1.2 mL, 5 mmol). The reaction mixture was stirred at room temperature overnight. The dioxane was removed under reduced pressure and the mixture was acidified with 1M HCl. The resultant white pasty material was dried to afford Compound 416-B (507 mg, 96%). MS (ESI) for (C₅₅H₆₆ClN₅O₁₄): *m/z* 1056 (M + H)⁺.

[00920] Steps 2 and 3: 416 (formic acid salt) was prepared utilizing the methods previously described. LCMS (5-95 AB, ESI): *t_R* = 0.658, [M + H]⁺ = 794.0; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.50 (brs, 1H), 7.75 (d, *J* = 8.0 Hz, 2H), 7.60 (d, *J* = 8.0 Hz, 2H), 7.47 (d, *J* = 8.0 Hz, 2H), 7.39 (d, *J* = 8.4 Hz, 2H), 7.19 (d, *J* = 8.0 Hz, 1 H), 6.94 (d, *J* = 8.0 Hz, 1 H), 6.91 (brs, 1H), 6.90 (brs, 1H), 6.82 (d, *J* = 8.0 Hz, 1H), 6.72 (brs, 2H), 4.95-4.91 (m, 1H), 4.79-4.72 (m, 1H), 4.58-4.50 (m, 1H), 4.18 (s, 2H), 3.10-2.80 (m, 4H), 2.94 (s, 3 H), 2.00-1.90 (m, 2H), 1.65-1.45 (m, 4H), 1.37 (d, *J* = 6.8 Hz, 3 H).

Example 321: Synthesis of Compound 417



[00921] Compound 417 (formic acid salt) was prepared utilizing the methods previously described from (S)-2-aminopropanenitrile. LCMS (5-95 AB, ESI): *t_R* = 0.790, [M + H]⁺ = 808.7.


Example 322: Synthesis of Compound 418


[00922] Step 1: (S)-methyl 2-((tert-butoxycarbonyl)amino)-3-((tert-

5 butyldimethylsilyl)oxy)propanoate (1.0 g, 3 mmol) was added to a solution saturated NH_3 in MeOH (50 mL) and the mixture was stirred at the sealed flask for 18 h. The volatiles were removed to give (S)-tert-butyl (1-amino-3-((tert-butyldimethylsilyl)oxy)-1-oxopropan-2-yl)carbamate (950 mg, 99.5% yield) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.35 (brs, 1H), 7.16 (brs, 1H), 6.46 (d, $J=8.4$ Hz, 1H), 4.02-4.06 (m, 1H), 3.76-3.63 (m, 2H), 1.42(s, 10 9H), 0.88(s, 9H), 0.06 (s, 6H).

[00923] Steps 2 and 3: (S)-tert-Butyl (1-amino-3-((tert-butyldimethylsilyl)oxy)-1-oxopropan-2-yl)carbamate was subject to the same dehydration and hydrolysis conditions as in Example 42 (Compound 136).

[00924] Compound 418 (formic acid salt) was prepared utilizing the methods previously 15 described. LCMS (5-95 AB, ESI): $t_R = 0.793$, $[\text{M} + \text{H}]^+ = 824.5$; ^1H NMR (400 MHz, $\text{MeOH}-d_4$) δ 8.51(brs, 1H), 7.87 (d, $J=8.0$ Hz, 2H), 7.69 (d, $J=8.0$ Hz, 2H), 7.59 (d, $J=8.0$ Hz, 2H), 7.45 (d, $J=8.0$ Hz, 2H), 7.12 (d, $J=8.0$ Hz, 1H), 7.05 (d, $J=8.0$ Hz, 1H), 6.99-6.94 (m, 2H), 6.90-6.80 (m, 2H), 6.57 (s, 1H), 5.05-5.01 (m, 1H), 4.85-4.79 (m, 2H), 4.70-4.55 (m, 1H), 3.80-3.70 (m, 2H), 3.09-2.90 (m, 2H), 2.95 (m, 3H), 2.00-1.93 (m, 3H), 1.75-1.58 (m, 5H), 1.36 (d, $J=6.8$ Hz, 3H).

Example 323: Synthesis of Compound 419



[00925] Step 1: A mixture of Compound 419-A (2.6 g, 8.0 mmol) and 1-(bromomethyl)-2-(trifluoromethyl)benzene (2.5 g, 10.4 mmol) in toluene (24 mL) was stirred at 100°C for 14h. After cooling to room temperature, the precipitated solid was collected, washed with pentane, which was further purified by HPLC to afford Compound 419-B (1.2 g, 27% yield) as a yellow solid.


[00926] To a solution of *t*-butyl carbamate (4.0 g, 34 mmol) in THF (14 mL) was added sodium benzenesulfinate (5.6 g, 34 mmol), propionaldehyde (2.18 g, 37.6 mmol) and formic acid (9.5 g, 205 mmol) sequentially and the mixture was stirred at room temperature for 18 h. The resulting precipitate was filtered, which was crystallized under Hex/EtOAc=4:1 to give tert-butyl (1-(phenylsulfonyl)propyl)carbamate (5.0 g, 48.9% yield) as a white solid. LCMS (5-95 AB, ESI): RT = 0.816, $\text{M}+\text{Na}^+ = 321.9$

[00927] To a solution of tert-butyl (1-(phenylsulfonyl)propyl)carbamate (3.0 g, 10.0 mmol) in toluene (100 mL) was added Compound 419-B (565 mg, 1.0 mmol) and acetone cyanohydrin (1.7 g, 20.0 mmol). After the above solution was cooled to -20°C, K_2CO_3 (6.9 g, 50.0 mmol) in H₂O (7.0 mL) was added in one portion and the resulting mixture was then vigorously stirred at the same temperature for 20 min, followed by gradually warming up to room temperature while stirring and further stirring for 48 h. The mixture was quenched by 5% NaHCO_3 solution (60 mL), which was extracted with EtOAc (100 mL x 2). The combined organic layers were washed with brine (100 mL x 3), dried over Na_2SO_4 , concentrated and the residue was purified by silica gel column to afford (S)-tert-butyl (1-cyanopropyl)carbamate (1.4 g, 76% yield) as a light yellow solid. ^1H NMR (CDCl_3 , 400 MHz) δ 4.87 (brs, 1H), 4.50 (brs, 1H), 1.86-1.74 (m, 2H), 1.44 (s, 9H), 1.08 (t, $J=7.6$ Hz, 3H).

[00928] Starting from (S)-tert-butyl (1-cyanopropyl)carbamate, the Boc removal (FA) procedure using methods previously described followed to afford (S)-2-aminobutanenitrile (70 mg, 67% yield) as yellow oil.


[00929] Compound 419 (formic acid salt) was prepared utilizing the methods previously described. LCMS (5-95 AB, ESI): RT = 0.802, $M+H^+ = 835.5$.

Example 324: Synthesis of Compound 420




5 [00930] Compound 420 (formic acid salt) was prepared utilizing the methods in Example 323 (Compound 419). LCMS (5-95 AB, ESI): RT = 0.799, $M+H^+ = 847.5$.

Example 325: Synthesis of Compound 421




10 [00931] Compound 421 (formic acid salt) was prepared utilizing the methods in Example 90 (Compound 184). LC-MS: $m/z = 882 [M + H]^+$.

Example 326: Synthesis of Compound 422






[00932] Compound 422 (formic acid salt) was prepared utilizing the methods in Example 90 (Compound 184). LC-MS: $m/z = 896 [M + H]^+$.

Example 327: Synthesis of Compound 423



[00933] A mixture of 4-bromo-2-fluoro-benzoic acid (250 mg, 1.14 mmol),
 5 bis(triphenylphosphine)palladium(II) dichloride (120.18 mg, 0.17 mmol), and cuprous iodide (32.61 mg, 0.17 mmol) in TEA (8 mL) was degassed with N₂. Ethynylbenzene (349.74 mg, 3.42 mmol) was added, capped in a glass vial, heated in oil bath at 60°C overnight. It was diluted with sat NH₄Cl, washed with iPrOAc (2x). The aqueous layer was acidified with 1N HCl, extracted with iPrOAc (2 x 30 mL), dried over MgSO₄, filtered, concentrated in vacuo,
 10 dried under high vacuum to give 271 mg (99%) of 2-fluoro-4-(2-phenylethynyl)benzoic acid as brown solid. LC-MS: *m/z* = 241 [M + H]⁺.

[00934] Compound 423 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: *m/z* = 860 [M + H]⁺. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.97 (d, *J* = 7.8 Hz, 1H), 8.89 (d, *J* = 7.4 Hz, 1H), 8.71 (t, *J* = 5.6 Hz, 1H), 8.40 (d, *J* = 9.0 Hz, 1H), 7.68 (t, *J* = 7.8 Hz, 1H), 7.62 – 7.57 (m, 2H), 7.53 – 7.45 (m, 5H), 7.23 (dd, *J* = 8.6, 2.2 Hz, 1H), 7.19 – 7.15 (m, 2H), 7.08 (d, *J* = 8.5 Hz, 1H), 6.73 (t, *J* = 2.9 Hz, 2H), 6.32 (s, 1H), 5.03 – 4.96 (m, 1H), 4.79 – 4.69 (m, 2H), 4.19 (d, *J* = 5.8 Hz, 2H), 4.15 – 4.06 (m, 4H), 3.20 (d, *J* = 15.8 Hz, 1H), 3.03 (dd, *J* = 9.6, 4.8 Hz, 5H), 2.93 (t, *J* = 8.0 Hz, 2H), 2.80 (s, 3H), 2.08 (d, *J* = 5.7 Hz, 1H), 2.02 – 1.94 (m, 1H), 1.21 (d, *J* = 6.7 Hz, 3H).



Example 330: Synthesis of Compound 426

[00937] Compound 426 (formic acid salt) was prepared utilizing the methods previously described. LC-MS: $m/z = 886 [M + H]^+$.

5 Example 331: Synthesis of Compound 427


[00938] Step 1: A mixture of 4-(4-chlorophenyl)benzaldehyde (1.0 g, 4.6 mmol) and methyl 2-(triphenyl- λ_5 -phosphanylidene)acetate (2.0 g, 6.0 mmol) in acetonitrile (12 mL) was stirred at room temperature overnight. It was purified with ISCO, 24 g column, eluted with 0-30% iPrOAc/heptane to give 1.18 g (94%) of methyl (E)-3-(4'-chloro-[1,1'-biphenyl]-4-yl)acrylate as white solid. LC-MS: $m/z = 273 [M + H]^+$.

[00939] Step 2: A mixture of methyl (E)-3-[4-(4-chlorophenyl)phenyl]prop-2-enoate (1.18 g, 4.33 mmol) and LiOH (327 mg, 13.0 mmol) in 1,4-dioxane (16 mL) and water (4 mL) was stirred at room temperature overnight. It was concentrated in vacuo, suspended in water, acidified with 1N HCl to pH~5. The white precipitate was collected by filtration, washed with

water, dried to give 1.11 g (99%) of (E)-3-(4'-chloro-[1,1'-biphenyl]-4-yl)acrylic acid as off white solid. LC-MS: $m/z = 259 [M + H]^+$.


[00940] Compound 427 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LC-MS: $m/z = 878 [M + H]^+$.

5 Example 332: Synthesis of Compound 428



[00941] Compound 428 (formic acid salt) was prepared utilizing the methods in Example 331 (Compound 427). LC-MS: $m/z = 824 [M + H]^+$.


Example 333 Synthesis of Compound 429




10

[00942] Compound 429 (formic acid salt) was prepared utilizing the methods in Example 47 (Compound 141) except (E)-styrylboronic acid is used in the Suzuki coupling. LCMS (Method 5-95 AB, ESI): $t_R = 0.705$ min, $[M+H]^+ = 858.9$.

Example 334: Synthesis of Compound 430



15




[00943] Step 1: To a mixture of palladium(II) acetate (29.4 mg, 0.13 mmol), 2-(3,3-dimethylbut-1-en-2-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (328.6 mg, 1.57 mmol), potassium phosphate, tribasic (695.0 mg, 3.27 mmol) and SPhos (107.5 mg, 0.26 mmol) in DMF (3 mL) and water (0.3 mL) was added methyl 4-bromo-2-methylbenzoate (300 mg, 1.31 mmol). The resulting mixture was heated at 70 °C for 16 hours under nitrogen. The reaction mixture was diluted with water (10 mL) and extracted with ethyl acetate (20 mL x 3). The combined organic layers were dried over sodium sulfate and filtered. The filtrate was concentrated. The residue was purified by prep-TLC (10% ethyl acetate in petroleum ether) to afford methyl 4-(3,3-dimethylbut-1-en-2-yl)-2-methylbenzoate (246 mg, 80.9% yield) as a colorless oil.

[00944] Step 2: A mixture of methyl 4-(3,3-dimethylbut-1-en-2-yl)-2-methylbenzoate (246 mg, 1.06 mmol) and 10% palladium (56.3 mg, 0.05 mmol) on carbon in methanol (15 mL) was stirred at 15 °C under H₂ (50 psi) for 16 h. The catalyst was filtered off and the solvent was evaporated to give methyl 4-(3,3-dimethylbutan-2-yl)-2-methylbenzoate (220 mg, 88.7% yield) as a white solid.

[00945] Step 3: Methyl 4-(3,3-dimethylbutan-2-yl)-2-methylbenzoate (220.0 mg, 0.94 mmol) was hydrolyzed according to General Method NaOH to afford 4-(3,3-dimethylbutan-2-yl)-2-methylbenzoic acid (190 mg, 91.9% yield) as a white solid. ¹H NMR (400 MHz, CD₃OD): δ 7.90 (d, J = 8.0 Hz, 1H), 7.06 - 6.93 (m, 2H), 2.57 (s, 3H), 2.51 (q, J = 7.6 Hz, 1H), 1.19 (d, J = 7.2 Hz, 3H), 0.81 (s, 9H).

[00946] Compound 430 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95AB): t_R = 0.721 min, [M + H]⁺ = 840.5.

Example 335: Synthesis of Compound 431


[00947] A mixture of methyl 4-bromo-2-methylbenzoate (50.0 mg, 0.22 mmol), 1,1'-

5 bis(diphenylphosphino)ferrocene palladium dichloride (16.0 mg, 0.02 mmol), potassium carbonate (60.3 mg, 0.44 mmol) and 1-cyclohexen-1-yl boronic acid (41.2 mg, 0.33 mmol) in toluene (5 mL) was stirred at 120 °C for 2 h under N₂ and evaporated to dryness. The residue was taken up in EtOAc (20 mL), washed with water (20 mL x 2) and brine (20 mL), dried over MgSO₄ and concentrated. The residue was purified by flash column chromatography (5%
10 EtOAc in petroleum ether, R_f = 0.7) to afford methyl 4-(cyclohexen-1-yl)-2-methylbenzoate (40 mg, 79.6% yield) as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.88 (d, J = 8.8 Hz, 1H), 7.28 - 7.23 (m, 2H), 6.23-6.21 (m, 1H), 3.88 (s, 3H), 2.61 (s, 3H), 2.41 - 2.39 (m, 2H), 2.23-2.21 (m, 2H), 1.80-1.77 (m, 2H), 1.68-1.65 (m, 2H).


[00948] 4-Cyclohexyl-2-methylbenzoic acid was prepared utilizing General Methods

15 Hydrogenation and NaOH from methyl 4-(cyclohexen-1-yl)-2-methylbenzoate.


[00949] Compound 431 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101) from 4-cyclohexyl-2-methylbenzoic acid. LCMS (Method 5-95AB): t_R = 0.565 min, [M + H]⁺ = 838.3.

Example 336: Synthesis of Compound 432

Example 337: Synthesis of Compound 433




5




[00952] Steps 2 and 3: Methyl 2-methyl-4-(prop-1-en-2-yl)benzoate was hydrogenated and hydrolyzed as described previously to afford 4-isopropyl-2-methylbenzoic acid (340 mg, 96.5% yield) as a yellow solid. ¹H NMR (400 MHz, CDCl₃): δ 8.02 (d, *J* = 8.0 Hz, 1H), 7.20 – 7.10 (m, 2H), 3.00 – 2.85 (m, 1H), 2.66 (s, 3H), 1.28 (d, *J* = 7.2 Hz, 6H).

[00953] Compound 433 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95AB): $t_R = 0.657$ min, $[M + H]^+ = 798.5$.

Example 338: Synthesis of Compound 434

[00954] Compound 434 (formic acid salt) was prepared as a white solid utilizing the same methods in Example 433. LCMS (Method 5-95AB): $t_R = 0.694$ min, $[M + H]^+ = 812.5$.

5 Example 339: Synthesis of Compound 435


[00955] To a solution of methyl 2-methyl-4-(prop-1-en-2-yl)benzoate (1.0 g, 5.26 mmol) in dichloromethane (50 mL) was added diethylzinc (1 M in toluene; 157.7 mL, 157.7 mmol) at 0 °C, followed by a solution of diiodomethane (6.35 mL, 78.85 mmol) in dichloromethane (10 mL). The reaction was allowed to warm to 20 °C and stirred for 16 h, quenched with aq. NH_4Cl and extracted with EtOAc (50 mL x 3). The combined organic layers were washed with brine (50 mL) and dried over Na_2SO_4 and concentrated. The residue was purified by flash column chromatography (1% EtOAc in petroleum ether, $R_f = 0.7$) as a mixture of DP and STM, which was separated by prep-HPLC (acetonitrile 67–77% / 0.1% TFA in water) to give methyl 2-methyl-4-(1-methylcyclopropyl)benzoate (80 mg, 7.5% yield) as a colorless oil.

[00956] Methyl 2-methyl-4-(1-methylcyclopropyl)benzoate (80.0 mg, 0.39 mmol) was hydrolyzed as usual (General Method NaOH) to afford 2-methyl-4-(1-methylcyclopropyl)benzoic acid (70 mg, 93.9% yield) as a white solid. ^1H NMR (400 MHz,

CD₃OD): δ 7.83 (d, J = 8.0 Hz, 1H), 7.15 - 7.10 (m, 2H), 2.56 (s, 3H), 1.42 (s, 3H), 0.95 - 0.85 (m, 2H), 0.85 - 0.75 (m, 2H).

[00957] Compound 435 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95AB): t_R = 0.557 min, $[M + H]^+ = 810.3$.

5 Example 340: Synthesis of Compound 436



[00958] Step1: A mixture of methyl 2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (400.0 mg, 1.45 mmol), sodium periodate (309.8 mg, 1.45 mmol) and ammonium acetate (800.0 mg, 10.38 mmol) in acetone (8 mL) and water (4 mL) was stirred at 20 °C for 16 h, and evaporated to dryness. The residue was taken up in EtOAc (20 mL), washed with water (20 mL x 2) and brine (20 mL), dried over MgSO₄ and concentrated. The residue was purified by flash column chromatography (33% ethyl acetate in petroleum ether) to afford (4-methoxycarbonyl-3-methyl-phenyl)boronic acid (200 mg, 71.2% yield) as a white solid.

[00959] Step 2: A solution cycloheptanone (2000.0 mg, 17.83 mmol) and 4-methylbenzenesulfonylhydrazide (3320 mg, 17.83 mmol) in ethanol (10 mL) was stirred at 60 °C for 16 h. The reaction was diluted with additional ethanol (10 mL) and filtered. The filtrate was concentrated to afford N-(cycloheptylideneamino)-4-methyl-benzenesulfonamide (4000 mg, 80% yield) as a white solid.

[00960] Step 3: A mixture of cesium carbonate (313.8 mg, 0.96 mmol), (4-methoxycarbonyl-3-methyl-phenyl)boronic acid (186.8 mg, 0.96 mmol) and N-(cycloheptylideneamino)-4-

methyl-benzenesulfonamide (180.0 mg, 0.64 mmol) in 1,4-dioxane (5 mL) was stirred at 110 °C for 16 h under N₂ protection, and evaporated to dryness. The residue was taken up in EtOAc (30 mL), washed with water (30 mL x 2) and brine (30 mL), dried over MgSO₄ and concentrated.


The residue was purified by prep-TLC (10% EtOAc in petroleum ether, R_f = 0.6) to afford

methyl 4-cycloheptyl-2-methylbenzoate (80 mg, 50.6% yield) as a colorless oil. LCMS (Method 5-95 AB, ESI): t_R = 0.976 min, [M + H]⁺ = 246.9.

[00961] Step 4: Methyl 4-cycloheptyl-2-methylbenzoate was hydrolyzed as previously described (General Method NaOH) to give 4-cycloheptyl-2-methylbenzoic acid.



[00962] Compound 436 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): t_R = 0.654 min, [M + H]⁺ = 852.4.

Example 341: Synthesis of Compound 437



[00963] Compound 437 (formic acid salt) was prepared utilizing the methods in Example 340 (Compound 436) and Example 195 (Compound 290). LCMS (Method 5-95 AB, ESI): t_R = 0.625 min, [M+H]⁺ = 892.4.

Example 342A: Synthesis of Compound 438



[00964] Step 1: To a mixture of 2-bromo-3-methylthiophene (226.7 mg, 1.28 mmol) and aluminum chloride (375.6 mg, 2.82 mmol) in 1,2-dichloroethane (3 mL) was added octanoyl


chloride (229.1 mg, 1.41 mmol) and stirred at 20 °C for 12 h. LCMS (5-95AB/1.5min): RT = 0.991 min, $[M+H]^+$ 302.9 showed 45% of DP. The mixture was poured onto ice and extracted with EtOAc (30 mL). The organic layer was washed with brine, dried over Na_2SO_4 and concentrated. The residue was purified by prep-TLC (5% EtOAc in petroleum ether, R_f = 0.5) to give 1-(5-bromo-4-methyl-2-thienyl)octan-1-one (160 mg, 41.2% yield) as a yellow oil. ^1H NMR (400 MHz, $CDCl_3$): δ 7.60 (s, 1H), 2.86 (t, J = 7.4 Hz, 2H), 2.22 (s, 3H), 1.67 (quint, J = 7.3 Hz, 2H), 1.43 - 1.22 (m, 8H), 0.90 (t, J = 7.0 Hz, 3H).


[00965] Step 2: A solution of 1-(5-bromo-4-methyl-2-thienyl)octan-1-one (160.0 mg, 0.53 mmol) and triethylamine (0.22 mL, 1.58 mmol) in methanol (10 mL) was treated with 1,1'-bis(diphenylphosphino)ferrocene palladium dichloride (38.6 mg, 0.05 mmol). The resulting mixture was stirred at 80 °C for 10 h under CO (45 psi) and concentrated. The residue was purified by prep-TLC (7% ethyl acetate in petroleum) to afford methyl 3-methyl-5-octanoyl-thiophene-2-carboxylate (120 mg, 80.5% yield) as a colorless oil.

[00966] Step 3: To a solution of methyl 3-methyl-5-octanoyl-thiophene-2-carboxylate (50.0 mg, 0.18 mmol) in 2,2,2-trifluoroacetic acid (1 mL) was added triethylsilane (82.4 mg, 0.71 mmol) dropwise. The mixture was stirred at 20 °C for 16 h and concentrated. The residue was purified by prep-TLC (5% EtOAc in petroleum ether, R_f = 0.6) to afford methyl 3-methyl-5-octyl-thiophene-2-carboxylate (38 mg, 80% yield) as a yellow oil. LCMS (Method 5-95 AB, ESI): RT = 1.180 min, $[M+H]^+$ = 269.1.

[00967] Compound 438 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): t_R = 0.785 min, $[M+H]^+$ = 874.4.

Example 342B: Synthesis of Compound 439 and Compound 440







[00968] Step 1: To a solution of 4-formyl-2-methylbenzoic acid (450.0 mg, 2.74 mmol) in N,N-dimethylformamide (5 mL) at 30 °C were added K₂CO₃ (1.13 g, 8.22 mmol) and iodomethane (1.3 mL, 20.85 mmol) and stirred for 10 h. The reaction was diluted with 1N HCl (15 mL) and extracted with EtOAc (50 mL x 3). The combined organic layers were washed with brine (30 mL x 3), dried over Na₂SO₄ and concentrated. The residue was purified by silica gel column (5% EtOAc in petroleum) to give methyl 4-formyl-2-methylbenzoate (400 mg, 81.9% yield) as a yellow oil.


[00969] Step 2: To a solution of tosyl hydrazide (418.1 mg, 2.24 mmol) in methanol (2 mL) was added methyl 4-formyl-2-methylbenzoate (400.0 mg, 2.24 mmol) dropwise at 60 °C and stirred at 60 °C for 2 h. The solution was concentrated to give crude methyl 2-methyl-4-((2-tosylhydrazono)methyl)benzoate (580.0 mg) as a white solid, which was used directly in the next step.

[00970] Step 3: To a solution of methyl 2-methyl-4-((2-tosylhydrazono)methyl)benzoate (580.0 mg, 1.67 mmol) in dioxane (4 mL) were added styrene (348.8 mg, 3.35 mmol) and potassium carbonate (347.1 mg, 2.51 mmol). The reaction was quenched with water (10 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were washed with water (20 mL) and brine (20 mL), dried over MgSO₄ and concentrated. The residue was purified first by prep-TLC (8% ethyl acetate in petroleum ether), then by reverse phase chromatography (acetonitrile 70-100/0.225% formic acid in water) to afford (±)-anti-methyl 2-methyl-4-(2-phenylcyclopropyl)benzoate (91 mg, 20.4% yield) and (±)-syn-methyl 2-methyl-4-(2-phenylcyclopropyl)benzoate (40 mg, 9% yield), both as a colorless oil.

[00971] Compound 439 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): t_R = 0.715 min, [M+H]⁺ = 872.9.

[00972] Compound 440 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): t_R = 0.708 min, [M+H]⁺ = 872.6.



Example 343B: Synthesis of Compound 442


[00976] Step 1: To a mixture of 2-bromo-5-methoxybenzaldehyde (2000.0 mg, 9.3 mmol) in dichloromethane (10 mL) was added boron tribromide (2M in DCM; 4.65 mL, 9.3 mmol) slowly at 0 °C. The reaction was warmed up to 25 °C and stirred for 3 h. The reaction was quenched with water (10 mL) at 0°C and extracted with EtOAc (50 mL). The organic layer was washed with water (50 mL x 2) and brine (50 mL), dried over MgSO₄ and concentrated. The residue was purified by flash column chromatography (petroleum ether) to afford 2-bromo-5-hydroxybenzaldehyde (1.7 g, 90.9% yield) as a colorless oil.

[00977] Step 2: To a solution of 2-bromo-5-hydroxybenzaldehyde (100.0 mg, 0.50 mmol) in N,N-dimethylformamide (3 mL) were added 1-bromoheptane (267.3 mg, 1.49 mmol) and potassium carbonate (206.3 mg, 1.49 mmol). The mixture was stirred at 20 °C for 16 h and concentrated to dryness. The residue was taken up in EtOAc (50 mL), washed with water (50 mL x 2) and brine (50 mL), dried over MgSO₄ and concentrated. The residue was purified by prep-TLC (10% EtOAc in petroleum ether, R_f = 0.3) to afford 2-bromo-5-(heptyloxy)benzaldehyde (100 mg, 67.2% yield) as a yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 10.32 (s, 1H), 7.52 (d, *J* = 9.2 Hz, 1H), 7.41 (d, *J* = 2.8 Hz, 1H), 7.03 (dd, *J* = 8.8, 3.2 Hz, 1H), 3.99 (t, *J* = 6.6 Hz, 2H), 1.85 - 1.75 (m, 2H), 1.50 - 1.20 (m, 8H), 0.90 (t, *J* = 6.8 Hz, 3H).


[00978] Step 3: A mixture of 2-bromo-5-(heptyloxy)benzaldehyde (100.0 mg, 0.33 mmol) in deoxo-fluor (2.0 mL, 1 mmol) was stirred at 80 °C for 16 h and concentrated to dryness. The residue was taken up in EtOAc (50 mL), washed with water (50 mL x 2) and brine (50 mL), dried over MgSO₄ and concentrated. The residue was purified by prep-TLC (5% EtOAc in petroleum ether, R_f = 0.3) to afford 1-bromo-2-(difluoromethyl)-4-(heptyloxy)benzene (80 mg, 74.5% yield) as a yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 7.44 (d, *J* = 8.8 Hz, 1H), 7.15 (d, *J* = 3.2 Hz, 1H), 7.00-6.70 (m, 2H), 3.94 (t, *J* = 6.6 Hz, 2H), 1.80 - 1.70 (m, 2H), 1.50 - 1.20 (m, 8H), 0.88 (t, *J* = 6.6 Hz, 3H).

[00979] Compound 442 (formic acid salt) was prepared utilizing the methods in Example 170 (Compound 265). LCMS (Method 5-95 AB, ESI): $t_R = 0.781$ min, $[M+H]^+ = 906.4$.

Example 344: Synthesis of Compound 443 and Compound 444




5



[00980] A mixture of methyl 3-methyl-1H-pyrazole-4-carboxylate (250.0 mg, 1.78 mmol), 4-isobutylphenyl boronic acid (412.9 mg, 2.32 mmol), pyridine (282.2 mg, 3.57 mmol) and copper(II) acetate (486.0 mg, 2.68 mmol) in N,N-dimethylformamide (6 mL) was stirred at 25°C for 14 h. The reaction was diluted with EtOAc (120 mL), washed with brine (50 mL x 3), dried over Na₂SO₄ and concentrated. The residue was purified by prep-TLC (20% EtOAc in petroleum ether, $R_f = 0.45$) to give methyl 1-(4-isobutylphenyl)-3-methyl-1H-pyrazole-4-carboxylate (350 mg, 72% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 0.996$ min, $[M+H]^+ = 272.9$.

[00981] Compound 443 and Compound 444 (formic acid salt) was prepared utilizing the methods in Example 5 and Example 7. Prior to TFA deprotection, the isomers were separated on preparative TLC. Data for Compound 443: LCMS (Method 5-95 AB, ESI): $t_R = 0.726$ min, $[M+H]^+ = 878.5$. Data for Compound 444: LCMS (Method 5-95 AB, ESI): $t_R = 0.720$ min, $[M+H]^+ = 878.5$.


20

Example 345: Synthesis of Compound 445

[00982] Step 1: A mixture of methyl 4-fluoro-2-methylbenzoate (300.0 mg, 1.78 mmol), 4-bromo-1H-pyrazole (393.3 mg, 2.68 mmol) and potassium carbonate (739.7 mg, 5.35 mmol) in N,N-dimethylformamide (20 mL) was stirred at 70°C for 12 h. The reaction mixture was diluted with water (100 mL) and extracted with EtOAc (100 mL x 3). The combined organic layers were washed with water (50 mL x 2) and brine (50 mL), dried over MgSO₄ and concentrated. The residue was purified by prep-TLC (20% EtOAc in petroleum ether, R_f = 0.5) to obtain methyl 4-(4-bromopyrazol-1-yl)-2-methyl-benzoate (200 mg, 38% yield) as a white solid. ¹H NMR (400 MHz, DMSO-d₆): δ 8.87 (s, 1H), 7.94 - 7.91 (m, 2H), 7.81 (s, 1H), 7.76 - 7.74 (m, 1H), 3.80 (s, 3H), 2.56 (s, 3H).


[00983] Step 2: A mixture of methyl 4-(4-bromopyrazol-1-yl)-2-methyl-benzoate (200.0 mg, 0.68 mmol), isobutyl boronic acid (345.41 mg, 3.39 mmol), sodium carbonate (359.1 mg, 3.39 mmol) and tetrakis(triphenylphosphine)palladium(0) (783.1 mg, 0.68 mmol) in toluene (5 mL) and water (1 mL) was stirred at 110 °C for 12 h and filtered. The filtrate was concentrated and purified by prep-TLC (30% EtOAc in petroleum, R_f = 0.7) to obtain methyl 4-(4-isobutylpyrazol-1-yl)-2-methyl-benzoate (40 mg, 21.7% yield) as a white solid. LCMS (Method 5-95 AB, ESI): t_R = 1.010 min, [M+H]⁺ = 272.9.

[00984] Compound 445 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): t_R = 0.585 min, [M+H]⁺ = 878.5.

Example 346: Synthesis of Compound 446

[00985] A mixture of methyl 4-(4-bromopyrazol-1-yl)-2-methylbenzoate (150.0 mg, 0.51 mmol), bis(triphenylphosphine)palladium(II) dichloride (17.9 mg, 0.03 mmol), 3-methyl-1-butyne (103.9 mg, 1.52 mmol) and copper(I) iodide (4.9 mg, 0.03 mmol) in triethylamine (4.19 mL, 30.03 mmol) was stirred at 100 °C for 18 h under nitrogen. The reaction was quenched with water (10 mL) and extracted with dichloromethane (10 mL x 3). The combined organic extracts were washed with brine (10 mL x 2), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep-HPLC to afford methyl 2-methyl-4-[4-(3-methylbut-1-ynyl)pyrazol-1-yl]benzoate (35 mg, 24.4% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 1.014$ min, $[M+H]^+ = 283.2$.

[00986] Compound 446 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): $t_R = 0.709$ min, $[M+H]^+ = 888.6$.

Example 347: Synthesis of Compound 447

[00988] Example 348: Synthesis of Compound 448

[00989] Step 1: A solution of 5-bromo-2,3-dihydro-1H-inden-1-one (1.0 g, 4.7 mmol), Pd(PPh₃)₄ (274 mg, 0.24 mmol), K₂CO₃ (1.3 g, 9.5 mmol) and octylboronic acid (1.1 g, 7.1 mmol) in toluene (15 mL) and H₂O (1mL) was stirred under N₂ at 100°C for 16 h. The reaction mixture was added with EtOAc (100 mL), which was washed by brine (100 mL). The organic layer dried over Na₂SO₄, concentrated and the residue was purified by silica gel flash column to give 5-octyl-2,3-dihydro-1H-inden-1-one (1.1 g, 95% yield) as yellow oil.

[00990] Step 2: To a solution of 5-octyl-2,3-dihydro-1H-inden-1-one (1.1 g, 4.5 mmol) in Et₂O (15 mL) was added 4M HCl/MeOH (5 mL), followed by the addition of a solution of 1-nitrobutane (511 mg, 5.0 mmol) in Et₂O (15 mL), and the mixture was stirred at room temperature for 2 h. The resulting precipitate was then collected, washed with Et₂O, and dried to give 2-(hydroxyimino)-5-octyl-2,3-dihydro-1H-inden-1-one (800 mg, 65% yield) as a yellow solid. LCMS (Method 5-95 AB, ESI): t_R = 0.920, [M + H]⁺ = 274.0

[00991] Step 3: A solution of 2-(hydroxyimino)-5-octyl-2,3-dihydro-1H-inden-1-one (800 mg, 2.9 mmol) and TsOH (725 mg, 3.8 mmol) in H₂O (10 mL) was heated to 50°C, followed by the addition of NaOH (468 mg, 11.7 mmol). The resulting mixture was heated further to 80°C for another 15min. After cooling to room temperature, the precipitate was removed from the mixture and the filtrate was acidified with concentrated HCl. The resulting precipitate was collected and purified by silica gel flash column to give 2-(cyanomethyl)-4-octylbenzoic acid (700 mg, 87.5% yield) as a yellow solid.

[00992] Step 4: Typical amide coupling (HATU/DIEA) procedure was applied to Compound 101-K (100 mg, 0.11 mmol) and 2-(cyanomethyl)-4-octylbenzoic acid (39 mg, 0.14 mmol) utilizing the methods previously described to afforded Compound 449-A (120 mg, 94% yield) as a white solid.


[00993] Step 5: To a solution of Compound 449-A (120 mg, 0.10 mmol) in EtOH/H₂O (3.0 mL, v/v=4/1) was added acetaldoxime (24 mg, 0.41 mmol), catalytic amount of Ph₃P (5.4 mg) and Pd(OAc)₂ (0.5 mg) and the mixture was stirred under N₂ at 80°C for 3h. H₂O (15 mL) was added to the reaction and the mixture was extracted with EtOAc (15 mL x 3). The combined organic layers was dried with Na₂SO₄ and concentrated to give Compound 449-B (110 mg, 90.3% yield) as a yellow solid, which was used directly in the next step. LCMS (Method 5-95 AB, ESI): t_R = 0.964, [M + H]⁺ = 1187.6.

[00994] Compound 448 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): t_R = 0.630, [M + H]⁺ = 911.6; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.47 (brs, 2H, HCOOH), 7.50 (d, *J*=8.0 Hz, 1H), 7.35 (d, *J*=8.0 Hz, 1H), 7.28-7.18 (m, 4H), 7.11 (d, *J*=8.0 Hz, 1H), 6.91 (brs, 1H), 6.83 (brs, 1H), 6.37(s, 1H), 5.11-5.08 (m, 1H),

4.81-4.75 (m, 2H), 4.28-4.19 (m, 4H), 4.22 (s, 2H), 3.87-3.77 (m, 2H), 3.37-3.35(m, 1H), 3.27-3.10 (m, 7H), 2.94 (s, 3H), 2.69-2.65 (m, 2H), 2.29-2.24 (m, 1H), 2.18-2.12 (m, 1H), 1.67-1.63 (m, 2H), 1.43-1.23 (m, 13H), 0.89 (t, $J=6.8$ Hz, 3H).

Example 349: Synthesis of Compound 449

5



[00995] Step 1: A solution of 5-bromoisobenzofuran-1(3H)-one (2.0 g, 9.4 mmol), K_2CO_3 (3.9 g, 28.2 mmol), $Pd(PPh_3)_4$ (1.1 g, 0.94 mmol) and octylboronic acid (3.0 g, 18.8 mmol) in toluene (40 mL) was stirred under N_2 at $100^\circ C$ for 24 h. The reaction mixture was added with EtOAc (100 mL), which was washed by brine (100 mL). The organic layer dried over Na_2SO_4 ,

concentrated and the residue was purified by silica gel flash column to give 5-octylisobenzofuran-1(3H)-one (1.26 g, 54% yield) as colorless oil.

[00996] Step 2: To a solution of 5-octylisobenzofuran-1(3H)-one (1.26 g, 5.1 mmol) in MeOH (22 mL) was added KOH (430 mg, 7.7 mmol) at room temperature and the mixture reaction was heated to 70°C while stirring and stirred at the same temperature for 6 h. The volatiles were removed and the residue was re-dissolved with water (20 mL), which was then acidified to pH=4 by 1M KHSO₄ solution. The resulting precipitate was collected, washed with water and dried in vacuo to afford 2-(hydroxymethyl)-4-octylbenzoic acid (852 mg, 63.1% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 0.977$, $M+Na^+ = 287.9$


[00997] Step 3: To a solution of 2-(hydroxymethyl)-4-octylbenzoic acid (752 mg, 3.0 mmol) in DCM (18 mL) was added MnO₂ (2.6 g, 30 mmol) and the mixture was stirred at room temperature for 5 h. The filtrate was then concentrated to obtain 3-hydroxy-5-octylisobenzofuran-1(3H)-one (565 mg, 71.7% yield) as yellow oil, which was used directly in the next step. LCMS (Method 5-95 AB, ESI): $t_R = 0.982$, $[M + H]^+ = 262.9$

[00998] Step 4: To a solution of 3-hydroxy-5-octylisobenzofuran-1(3H)-one (565 mg, 2.15 mmol) and diethyl-cyanomethyl phosphonate (382 mg, 2.15 mmol) in DME (11 mL) was added 60% NaH in oil (172 mg, 4.30 mmol) at 0°C and the mixture was stirred at the same temperature for 1 h. The reaction mixture was quenched with H₂O (30 mL), which was extracted by EtOAc (30 mL x 2). The combined organic layers were dried over Na₂SO₄, concentrated and the residue was purified by HPLC to obtain (E)-2-(2-cyanovinyl)-4-octylbenzoic acid (60 mg, 9.8% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 8.38 (d, $J=16$ Hz, 1H), 8.05 (d, $J=8$ Hz, 1H), 7.34-7.31(m, 2H), 5.77 (d, $J=16$ Hz, 1H), 2.67 (t, $J=8.0$ Hz, 2H), 1.69-1.66 (m, 2H), 1.30-1.26 (m, 10H), 0.87 (t, $J=6.8$ Hz, 3H).


[00999] Typical amide coupling (HATU) procedure was applied to Compound 101-K (150 mg, 0.16 mmol) and (E)-2-(2-cyanovinyl)-4-octylbenzoic acid (56 mg, 0.20 mmol) to afford Compound 449-A (152 mg, 78% yield) as a white solid.

[001000] A solution of Compound 449-A (122 mg, 0.10 mmol), 10% Pd/C (22 mg, 0.02 mmol) in MeOH (12 mL) was stirred under H₂ at room temperature. LCMS was used to monitor the reaction to ensure no over-reduction of nitrile. After LCMS indicated the completion of the reaction, the filtrate was concentrated to give Compound 449-B (116 mg, 94.9% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 1.139$, $[M + H]^+ = 1183.8$.


[001001] Compound 449 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): $t_R = 0.769$, $[M + H]^+ = 907.5$

Example 350: Synthesis of Compound 450


[001002]Compound 450 (free base) was prepared utilizing similar methods in Example 348 (Compound 448). LCMS (Method 5-95 AB, ESI): $t_R = 0.764$, $[M + H]^+ = 923.5$.

5 Example 351: Synthesis of Compound 451

[001003]Compound 451 (free base) was prepared utilizing similar methods in Example 348 (Compound 448). LCMS (Method 5-95 AB, ESI): $t_R = 0.776$, $[M + H]^+ = 925.8$.

Example 352: Synthesis of Compound 452

10





[001004]To a solution of methyl 3-(decylamino)propanoate (380 mg, 1.36 mmol) in DMF (3 mL) was added K_2CO_3 (560 mg, 4.07 mmol) and *t*-butyl 2-bromoethylcarbamate (300 mg, 1.36

mmol) at 0°C and the mixture was stirred at room temperature for 48 h. The reaction was added with H₂O (30 mL), which was extracted with EtOAc (30 mL x 2). The combined organic layers were dried over Na₂SO₄, concentrated and the residue was purified by silica gel flash column to give methyl 3-((2-((tert-butoxycarbonyl)amino)ethyl)(decyl)amino)propanoate (166 mg, 31.6% yield) as colorless oil. LCMS (Method 5-95 AB, ESI): t_R = 0.874, [M + H]⁺ = 387.3.

[001005]Compound 452 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): t_R = 0.700, [M + H]⁺ = 792.5.

Example 353: Synthesis of Compound 453





[001006]To a mixture of Compound 101-G (300 mg, 0.42 mmol), Cbz-Asn-OH (224 mg, 0.84 mmol), HATU (319 mg, 0.84 mmol), MgSO₄ (252 mg, 2.1 mmol) in DMF/DCM (20 mL, v/v=1:3) was added DIEA (217 mg, 1.68 mmol) at 0 °C and the mixture was warmed to room temperature while stirring and stirred for 2 h. The volatiles were removed and the residue was taken up by EtOAc (100 mL), which was washed by brine (50 mL x 2). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified by prep-TLC to afford Compound 453-A (100 mg, 25.2% yield) as a white solid. During the coupling, dehydration of the amide occurs.

[001007]Compound 453 (free base) was prepared utilizing the methods previously described from Compound 453-A. LCMS (Method 5-95 AB, ESI): t_R = 0.686, [M + H]⁺ = 816.2; ¹H NMR (400 MHz, MeOH-d₄) δ 7.27-7.24 (m, 2H), 7.14 (d, J=8.0 Hz, 1H), 7.09 (d, J=8.0 Hz, 1H), 6.87 (brs, 1H), 6.79 (brs, 1H), 6.21(s, 1H), 5.18-5.15 (m, 1H), 4.83-4.77 (m, 2H), 4.21-4.17

(m, 4H), 4.19 (s, 2H), 3.16-3.13 (m, 5H), 2.93-2.90 (m, 2H), 2.81 (s, 3H), 2.28-2.25 (m, 3H), 1.65-1.61 (m, 3H), 1.33-1.29 (m, 18H), 0.90 (t, $J=6.4$ Hz, 3H).

Example 354: Synthesis of Compound 454



[001008] To a solution of (S)-methyl 2-(((benzyloxy)carbonyl)amino)-4-hydroxybutanoate (900 mg, 3.4 mmol) and Et₃N (0.94 mL, 6.8 mmol) in DCM (10 mL) was added MsCl (0.39 mL, 5.1 mmol) at 0°C and the reaction was warmed up to room temperature while stirring and stirred at the same temperature for 3 h. The pH of the reaction mixture was adjusted to pH=5 with saturated citric acid solution, which was extracted with DCM (30 mL x 2). The combined organic layers were dried over Na₂SO₄, concentrated and the residue was purified by HPLC to afford (S)-methyl 2-(((benzyloxy)carbonyl)amino)-4-((methylsulfonyl)oxy)butanoate (500 mg, 43% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 0.780$, $[M + H]^+ = 345.9$.



[001009] A solution of (S)-methyl 2-(((benzyloxy)carbonyl)amino)-4-

((methylsulfonyl)oxy)butanoate (200 mg, 0.58 mmol), K₂CO₃ (160 mg, 1.16 mmol), 1,2,4-triazole (80.0 mg, 1.16 mmol) in DMF (2 mL) was stirred at 60°C for 16 h. Water (10 mL) was added to the reaction, which was then extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (50 mL), dried over MgSO₄, concentrated and the residue was purified by prep-TLC to afford (S)-methyl 2-(((benzyloxy)carbonyl)amino)-4-(1H-1,2,4-triazol-1-yl)butanoate (100 mg, 54.2% yield). LCMS (Method 5-95 AB, ESI): $t_R = 0.728$, $[M + H]^+ = 318.9$.

[001010] Compound 454 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): $t_R = 0.672$, $[M + H]^+ = 872.4$; ¹H NMR (400 MHz,


MeOH-*d*₄) δ 8.51 (s, 1H), 8.49 (brs, 2H, HCOOH), 7.97 (s, 1H), 7.25-7.22 (m, 2H), 7.17 (d, *J*=8.0 Hz, 1H), 7.07 (d, *J*=8.0 Hz, 1H), 6.86 (brs, 1H), 6.78 (brs, 1H), 6.29 (s, 1H), 4.80-4.74 (m, 3H), 4.35 (t, *J*=6.4 Hz, 2H), 4.18 (brs, 6H), 3.16-3.13 (m, 5H), 2.71 (s, 3H), 2.40-2.30 (m, 2H), 2.26 (t, *J*=6.0 Hz, 2H), 2.25-2.15 (m, 2H), 1.68-1.64 (m, 3H), 1.34-1.28 (m, 17H), 0.89 (t, *J*=6.8 Hz, 3H).

Example 355: Synthesis of Compound 455



[001011] To a solution of (S)-methyl 2-(((benzyloxy)carbonyl)amino)-4-hydroxybutanoate (400 mg, 1.16 mmol) in DMF (5 mL) was added NaCN (62 mg, 1.27 mmol) at room temperature and the reaction mixture was warmed to 75 °C while stirring and stirred at the same temperature for 2 h. The volatiles were removed and the residue was taken by EtOAc (40 mL), which was washed by brine (40 mL). The organic layer was dried over Na₂SO₄, concentrated and the residue was purified by Prep-TLC to give (S)-methyl 2-(((benzyloxy)carbonyl)amino)-4-cyanobutanoate (210 mg, 65.6% yield) as colorless oil. LCMS (Method 5-95 AB, ESI): *t*_R = 0.779, *M*+Na⁺ = 298.9.

[001012] Compound 455 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): *t*_R = 0.814, [*M* + H]⁺ = 830.4.

Example 356: Synthesis of Compound 456


[001013] Step 1: To a mixture of Bn_2NH (1.0 g, 5.1 mmol) and ethyl 4-bromobutanoate (1.0 g, 5.1 mmol) in DMF (5 mL) was added K_2CO_3 (3.5 g, 25.6 mmol) at room temperature and the mixture was stirred at 100 °C for 16 h. The reaction mixture was added EtOAc (50 mL), which was washed with brine (50 mL x 3). The organic layer was dried over Na_2SO_4 , concentrated and the residue was purified on silica gel flash column to obtain ethyl 4-(dibenzylamino)butanoate (520 mg, 32.6% yield) as pale oil.

[001014] Step 2: To a solution of ethyl 4-(dibenzylamino)butanoate (4.0 g, 12.8 mmol) in THF (40 mL) was added 2 N LDA in THF (6.4 mL, 12.8 mmol) dropwise at -78°C and the reaction mixture was stirred at the same temperature for 0.5 h, followed by the addition of a solution of tert-butyl 2-bromoacetate (3.0 g, 15.4 mmol) and HMPA (2.3 g, 12.8 mmol) in THF (40 mL). The resulting mixture was stirred for 3 h at -78°C and the reaction was warmed up to room temperature while stirring and stirred for another 3 h. The reaction mixture was added with EtOAc (200 mL), which was washed with saturated NH_4Cl and brine (each 200 mL). The organic layer was dried over Na_2SO_4 , concentrated and the residue was purified on silica gel flash column to obtain 4-tert-butyl 1-ethyl 2-(2-(dibenzylamino)ethyl)succinate (800 mg, 14.6% yield) as pale oil. ^1H NMR (400 MHz, CDCl_3) δ 7.36-7.21 (m, 10H), 4.05 (q, $J=7.0$ Hz, 2H), 3.61 (d, $J=16$ Hz, 2H), 3.47 (d, $J=16$ Hz, 2H), 2.92-2.86 (m, 1H), 2.46-2.39 (m, 3H), 2.17-2.11 (m, 1H), 1.98-1.90 (m, 2H), 1.42 (s, 9H), 1.17 (t, $J=7.2$ Hz, 3H).

[001015] Step 3: Standard acidic hydrolysis of the t-butyl ester with TFA in DCM afforded 5-(dibenzylamino)-3-(ethoxycarbonyl)pentanoic acid.


[001016]Compound 456 (formic acid salt) was prepared utilizing the methods previously described as a mixture of diastereomers. LCMS (Method 5-95 AB, ESI): $t_R = 0.719$, $[M + H]^+ = 826.5$; ^1H NMR (400 MHz, MeOH- d_4) δ 7.39-7.21 (m, 3H), 7.14-7.03 (m, 4H), 6.89 (brs, 1H), 6.83 (brs, 1H), 6.34 (s, 1H), 4.25-4.19 (m, 4H), 4.21 (s, 2H), 3.21-2.97 (m, 9H), 2.84 (s, 3H),
 5 2.74-2.70 (m, 1H), 2.63-2.59 (m, 2H), 2.23 (s, 3H), 2.10-1.98 (m, 2H), 1.65-1.57 (m, 2H), 1.63-1.35 (m, 6H), 1.22-1.17 (m, 1H), 0.96 (t, $J=6.8$ Hz, 3H).

Example 357: Synthesis of Compound 457




[001017]Compound 457 (formic acid salt) was prepared utilizing the methods previously described using tert-butyl (3-bromopropyl)carbamate in the phenol alkylation step. LCMS (Method 5-95AB): $t_R = 0.772$ min/ 1.5 min, $[M+H]^+ = 896.6$.
 10

Example 358: Synthesis of Compound 458



[001018]Compound 458 (formic acid salt) was prepared utilizing the methods in Example 7 (Compound 101) using (2-bromoethoxy)(tert-butyl)dimethylsilane in the phenol alkylation step. LCMS (Method 5-95AB): $t_R = 0.723$ min/ 1.5 min, $[M + H]^+ = 850.8$.
 15


Example 359: Synthesis of Compound 459

[001019]Compound 459-A was prepared utilizing the methods in Example 7.


5 **[001020]**Step 1: Compound 459-A (70 mg, 0.07 mmol) was treated according to General Method Formic Acid to afford Compound 459-B (40 mg, 67.4% yield) as a white solid. LCMS (5-95 AB, ESI): $t_R = 0.678$, $[M + H]^+ = 811.2$.

[001021]Step 2: To a stirred solution of Compound 459-B (40 mg, 0.05 mmol) and HCHO (28 mg, 0.93 mmol) in MeOH (1 mL) was added Et₃N (14 mg, 0.14 mmol) and NaBH₃CN (59 mg, 0.93 mmol) and the mixture was stirred at room temperature for 1 h. The volatiles were removed and the residue was purified by prep-TLC to give Compound 459-C (20 mg, 49.4% yield) as a light yellow solid. LCMS (5-95 AB, ESI): $t_R = 0.806$, $[M + H]^+ = 867.9$.

[001022]Starting from Compound 459-C, Compound 459 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (5-95 AB, ESI): $t_R =$
15 0.639 , $[M + H]^+ = 835.3$.


Example 360: Synthesis of Compound 460

[001023]Compound 460 (formic acid salt) was prepared utilizing methods in Example 359 (Compound 459). LCMS (5-95 AB, ESI): $t_R = 0.728$, $[M + H]^+ = 804.4$.


5 Example 361: Synthesis of Compound 461

[001024]3-((Tert-butoxycarbonyl)((4'-chloro-[1,1'-biphenyl]-4-yl)methyl)amino)propanoic acid was prepared in a manner similar to 3-((tert-butoxycarbonyl)(decyl)amino)propanoic acid (Example 6). LCMS (5-95 AB, ESI): $t_R = 1.015$, $[M+Na]^+ = 412.1$.


[001025]Compound 461 (formic acid salt) was prepared utilizing methods previously described. LCMS (5-95 AB, ESI): $t_R = 0.586$, $[M + H]^+ = 809.3$.

Example 362: Synthesis of Compound 462

[001026]Compound 462 (formic acid salt) was prepared utilizing methods previously described. LCMS (5-95 AB, ESI): $t_R = 0.747$, $[M + H]^+ = 783.8$.


5 Example 363: Synthesis of Compound 463

[001027]Compound 463 (formic acid salt) was prepared utilizing methods previously described from 2-decanamidoacetic acid. LCMS (5-95 AB, ESI): $t_R = 0.601$, $[M + H]^+ = 749.5$.


Example 364: Synthesis of Compound 464

10


[001028]Compound 464 (formic acid salt) was prepared utilizing methods previously described from 3-decanamidopropanoic acid. LCMS (5-95 AB, ESI): $t_R = 0.619$, $[M + H]^+ = 763.3$.

Example 365: Synthesis of Compound 465

[001029]Compound 465 (formic acid salt) was prepared as a white solid utilizing methods previously described from (S)-2-(((benzyloxy)carbonyl)amino)butanoic acid. LCMS (Method 5 5-95AB): $t_R = 0.864$ min/ 1.5 min, $[M + H]^+ = 833.4$.

Example 366: Synthesis of Compound 466


[001030]Compound 466 (formic acid salt) was prepared utilizing methods previously described. LCMS (5-95 AB, ESI): $t_R = 0.835$, $[M + H]^+ = 754.5$

Example 367: Synthesis of Compound 467

[001031]To a stirred solution of Compound 466 (20 mg, 0.03 mmol) in pH=7 buffer containing 1N NaH_2PO_3 and 1N Na_2HPO_3 was added 1N NaOH solution dropwise until pH=8.5, followed by the addition of imido-formamide hydrochloride (43 mg, 0.53 mmol). The resulting mixture was stirred at 0°C for 2 h. MeOH (2 mL) was added to the reaction mixture. After filtration, the filtrate concentrated and the residue was purified by HPLC to afford Compound 467 (1.6 mg,


7.7% yield) as a white solid. LCMS (5-95 AB, ESI): $t_R = 0.838$, $[M + H]^+ = 781.4$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.55 (brs, 1H), 7.86 (s, 1H), 7.36-7.29 (m, 1H), 7.13-6.93 (m, 6H), 6.92-6.86 (m, 1H), 6.85-6.79 (m, 1H), 6.40 (s, 1H), 5.16-5.08 (m, 1H), 4.79-4.74 (m, 2H), 4.19 (s, 2H), 3.52-3.41 (m, 2H), 3.18-3.00 (m, 2H), 2.92 (s, 3H), 2.62-2.50 (m, 2H), 2.38 (s, 3H), 2.29-2.07 (m, 2H), 1.65-1.50 (m, 2H), 1.40-1.21 (m, 9H), 0.90 (t, $J=6.8$ Hz, 3H).

Example 368: Synthesis of Compound 468




[001032] Compound 468 (formic acid salt) was prepared utilizing methods previously described. LCMS (5-95 AB, ESI): $t_R = 0.842$, $[M + H]^+ = 762.4$.

10 Example 369: Synthesis of Compound 469




[001033] Compound 469 (formic acid salt) was prepared utilizing methods previously described. LCMS (5-95 AB, ESI): $t_R = 0.671$, $[M + H]^+ = 750.2$.

Example 370 : Synthesis of Compound 470




15


[001034] Compound 470 (formic acid salt) was prepared as a white solid utilizing methods previously described. LCMS (Method 5-95AB): $t_R = 0.905$ min/ 1.5 min, $[M+H]^+ = 825.0$.

Example 371: Synthesis of Compound 471

[001035]Compound 471 (formic acid salt) was prepared as a white solid utilizing the methods previously described. LCMS (Method 5-95AB): $t_R = 0.905$ min/ 1.5 min, $[M+H]^+ = 810.5$.


5 Example 372: Synthesis of Compound 472

[001036]Compound 472 (formic acid salt) was prepared as a white solid utilizing the methods previously described. LCMS (Method 5-95AB): $t_R = 0.876$ min/ 1.5 min, $[M+H]^+ = 831.6$.


Example 373: Synthesis of Compound 473

10


[001037]Compound 473 (formic acid salt) was prepared as a white solid utilizing the methods previously described. LCMS (Method 5-95AB): $t_R = 0.895$ min/ 1.5 min, $[M+H]^+ = 840.6$.

Example 374: Synthesis of Compound 474




[001038]Compound 474 (formic acid salt) was prepared utilizing methods previously described. LCMS (Method 5-95AB): $t_R = 0.838$ min/ 1.5 min, $[M+H]^+ = 762.5$.

5 Example 375: Synthesis of Compound 475

[001039]Compound 475 (formic acid salt) was prepared utilizing methods previously described. LCMS (Method 5-95AB): $t_R = 0.848$ min/ 1.5 min, $[M+H]^+ = 748.5$.


Example 376: Synthesis of Compound 476

[001040]Compound 476 (formic acid salt) was prepared utilizing methods previously described. LCMS (Method 5-95AB): $t_R = 0.821$ min/ 1.5 min, $[M+H]^+ = 754.5$.



[001043]Compound 479 (formic acid salt) was prepared utilizing methods previously described from (S)-2-((((9H-fluoren-9-yl)methoxy)carbonyl)amino)-3-(pyridin-4-yl)propanoic acid. LC-MS: $m/z = 888 [M + H]^+$.


Example 380: Synthesis of Compound 480



5

[001044]Compound 480 (formic acid salt) was prepared utilizing methods previously described from (2S,4S)-1-((benzyloxy)carbonyl)-4-((tert-butoxycarbonyl)amino)pyrrolidine-2-carboxylic acid. LC-MS: $m/z = 852 [M + H]^+$.


Example 381: Synthesis of Compound



10

[001045]Compound 481 (formic acid salt) was prepared utilizing methods previously described from (2S,4S)-1-((benzyloxy)carbonyl)-4-((tert-butoxycarbonyl)amino)pyrrolidine-2-carboxylic acid. LC-MS: $m/z = 860 [M + H]^+$.


Example 382: Synthesis of Compound 482



15



[001046] Compound 482 (formic acid salt) was prepared utilizing methods previously described. LCMS (ESI): $[M + H]^+ = 880.4$.

Example 383: Synthesis of Compound 483



5 [001047] Compound 483 (formic acid salt) was prepared utilizing methods previously described. LCMS (ESI): $[M + H]^+ = 894.4$.

Example 384: Synthesis of Compound 484



10 [001048] A solution of methyl 2-(((benzyloxy)carbonyl)amino)-2-(dimethoxyphosphoryl)acetate (1.0 g, 3.0 mmol) and DBU (0.35 mL) in DCM (5 mL) was stirred at 0°C for 20 min, followed by the addition of a solution of (S)-tert-butyl 2-formylpyrrolidine-1-carboxylate (500 mg, 2.5 mmol) in DCM (5 mL). The resulting mixture was then warmed up to room temperature while stirring and stirred for 20 h at the same temperature. The volatiles were removed and the residue


15 was purified by silica gel flash column to give (S,E)-tert-butyl 2-(2-(((benzyloxy)carbonyl)amino)-3-methoxy-3-oxoprop-1-en-1-yl)pyrrolidine-1-carboxylate (700 mg, 69% yield) as colorless oil. LCMS (Method 5-95 AB, ESI): $t_R = 0.801$, $M + Na^+ = 427.0$

[001049] A solution of (S,E)-tert-butyl 2-(2-(((benzyloxy)carbonyl)amino)-3-methoxy-3-oxoprop-1-en-1-yl)pyrrolidine-1-carboxylate (500 mg, 1.24 mmol) and 10% Pd/C (132 mg, 0.12 mmol) in MeOH (10 mL) was stirred at room temperature for 2 h under H₂ (15 psi). The filtrate was concentrated and the residue was purified by HPLC (10mM NH₄HCO₃-ACN, 30%-50%) to give (S)-tert-butyl 2-((S)-2-(((benzyloxy)carbonyl)amino)-3-methoxy-3-oxopropyl)pyrrolidine-1-carboxylate (200 mg, 60% yield) as a white solid. Optical purity was confirmed by SFC. LCMS (Method 5-95 AB, ESI): t_R = 0.639, [M + H]⁺ = 272.9

[001050] To a solution of (S)-tert-butyl 2-((S)-2-(((benzyloxy)carbonyl)amino)-3-methoxy-3-oxopropyl)pyrrolidine-1-carboxylate (900 mg, 3.3 mmol) in DCM (10 mL) was added Et₃N (501 mg, 5.0 mmol), followed by the addition of a solution of Cbz-OSu (820 mg, 3.3 mmol) in DCM (25 mL). The resulting mixture was stirred at room temperature for 16 h. The volatiles were removed and the residue was purified by silica-gel flash column to give (S)-tert-butyl 2-((S)-2-(((benzyloxy)carbonyl)amino)-3-methoxy-3-oxopropyl)pyrrolidine-1-carboxylate (900 mg, 67% yield) as colorless oil.

[001051] Compound 484 (formic acid salt) was prepared utilizing methods previously described. LCMS (Method 5-95 AB, ESI): t_R = 0.780, [M + H]⁺ = 860.4; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.50 (brs, 3H, HCOOH), 7.33-7.23 (m, 2H), 7.18 (d, *J*=8.4 Hz, 1H), 7.11 (d, *J*=8.4 Hz, 1H), 6.93 (brs, 1H), 6.79 (brs, 1H), 6.28 (s, 1H), 4.35-4.14 (m, 4H), 4.21 (s, 2H), 3.63-3.48 (m, 1H), 3.26-3.04 (m, 5H), 2.85 (s, 3H), 2.40-1.94 (m, 8H), 1.86-1.56 (m, 4H), 1.47-1.19 (m, 20H), 0.92 (t, *J*=6.4 Hz, 3H).


Example 385: Synthesis of Compound 485



[001052] To a stirred solution of (S)-4-amino-2-(((benzyloxy)carbonyl)amino)butanoic acid (200 mg, 0.8 mmol) in DMF (5 mL) was added tert-butyl (((tert-butoxycarbonyl)amino)(1H-pyrazol-1-yl)methylene)carbamate (492 mg, 1.6 mmol) and DIEA (205 mg, 1.6 mmol) and the mixture was stirred at room temperature for 3 h. The volatiles were removed and the residue was purified by silica gel flash column to afford (S)-2-(((benzyloxy)carbonyl)amino)-4-(2,3-bis(tert-butoxycarbonyl)guanidino)butanoic acid (220 mg, 56.1% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 0.894$, $[M + H]^+ = 495.1$

[001053] Compound 485 (formic acid salt) was prepared utilizing methods previously described. LCMS (Method 5-95 AB, ESI): $t_R = 0.757$, $[M + H]^+ = 882.4$.



Example 386: Synthesis of Compound 486




[001054] Compound 486-A is an intermediate in the preparation of Compound 223 (Example 129).

[001055] A mixture of Compound 486-A (100.0 mg, 0.09 mmol), triethylorthoformate (14.2 mg, 0.10 mmol), and p-toluenesulfonic acid (1.5 mg, 0.01 mmol) in ethyl acetate (5 mL) was stirred at 50°C for 2 h. The mixture was concentrated and purified by prep-TLC (7% methanol in dichloromethane, $R_f = 0.5$) to obtain Compound 486-A (70 mg, 69.4% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 1.139$ min, $[M + H]^+ = 1156.0$.

[001056] Compound 486 (formic acid salt) was prepared as a white solid utilizing the methods in Example 7 (Compound 101). LCMS (Method 5-95 AB, ESI): $t_R = 0.759$ min, $[M + H]^+ = 879.9$.





[001058] To a solution of 5-octyl-3H-isobenzofuran-1-one (200.0 mg, 0.81 mmol) in methanol (8.5 mL) and water (1.5 mL) was added potassium hydroxide (68.33 mg, 1.22 mmol). The mixture was stirred at 80 °C for 3 h and concentrated. The residue was taken up in EtOAc (20 mL), washed with water (20 mL x 2) and brine (20 mL), dried over MgSO₄ and concentrated to give crude 2-(hydroxymethyl)-4-octylbenzoic acid (200 mg, 93.2% yield) as a white solid.

[001059] A mixture of 2-(hydroxymethyl)-4-octylbenzoic acid (189.9 mg, 0.72 mmol), manganese dioxide (187.4 mg, 2.16 mmol) in dichloromethane (5 mL) was stirred at 20 °C for 16 h and filtered. The filtrate was concentrated and purified by flash column chromatography (5% methanol in DCM, R_f = 0.4) to afford 2-formyl-4-octylbenzoic acid (150 mg, 79.6% yield) as a white solid.

[001060] A mixture of 2-formyl-4-octylbenzoic acid (100.0 mg, 0.38 mmol), iodomethane (108.2 mg, 0.76 mmol) and potassium carbonate (158.1 mg, 1.14 mmol) in N,N-dimethylformamide (3 mL) was stirred at 20 °C for 16 h. The reaction was diluted with EtOAc (20 mL), washed with water (20 mL x 2) and brine (20 mL), dried over MgSO₄ and concentrated. The crude was purified by flash column chromatography (10% ethyl acetate in petroleum ether, R_f = 0.5) to afford methyl 2-formyl-4-octylbenzoate (100 mg, 94.9% yield) as a colorless oil. ¹H NMR (400 MHz, CDCl₃): δ 10.65 (s, 1H), 7.91 (d, J = 8.0 Hz, 1H), 7.75 (s, 1H), 7.46-7.43 (m, 1H), 3.97 (s, 3H), 2.70 (t, J = 7.6 Hz, 2H), 1.66-1.60 (m, 2H), 1.30-1.24 (m, 10H), 0.88 (t, J = 6.8 Hz, 3H).

[001061] To a solution of Compound 101-K (100.0 mg, 0.11 mmol), methyl 2-formyl-4-octylbenzoate (60.47 mg, 0.22 mmol) and acetic acid (0.30 mL) in methanol (10 mL) was added sodium cyanoborohydride (8.3 mg, 0.13 mmol). The solution was stirred at 20 °C for 6 h and concentrated. The residue was taken up in EtOAc (20 mL), washed with water (20 mL x 2) and


5 brine (20 mL), dried over MgSO₄ and concentrated. The crude was purified by prep-TLC (66% EtOAc in petroleum ether, R_f = 0.4) to afford Compound 488-A (100 mg, 77.8% yield) as a white solid. LCMS (Method 5-95 AB, ESI): t_R = 0.888 min, [M + H]⁺ = 1174.6.

[001062] A solution of Compound 488-A (100.0 mg, 0.09 mmol) in 1,2-dichloroethane (5 mL) was stirred at 60 °C for 16 h and concentrated to afford crude Compound 488-B (99 mg) as a

10 white solid. LCMS (Method 5-95 AB, ESI): t_R = 1.015 min, [M + H]⁺ = 1164.9.

[001063] Compound 488 (formic acid salt) was prepared utilizing methods previously described. LCMS (Method 5-95 AB, ESI): t_R = 0.633 min, [M + H]⁺ = 866.5.

Example 389: Synthesis of Compound 489





[001064] Step 1: A mixture of methyl 2-chloro-4-octylbenzoate (160.0 mg, 0.57 mmol), tricyclohexylphosphine (47.6 mg, 0.17 mmol), tris(dibenzylideneacetone)dipalladium(0) (51.8 mg, 0.06 mmol), (E)-1-ethoxyethene-2-boronic acid pinacol ester (123.3 mg, 0.62 mmol) and potassium phosphate tribasic (240.2 mg, 1.13 mmol) in 1,4-dioxane (4 mL) was stirred at 100

20 °C for 16 h and concentrated. The residue was taken up in EtOAc (20 mL), washed with water (20 mL x 2) and brine (20 mL), dried over MgSO₄ and concentrated. The crude was purified by flash column chromatography (20% ethyl acetate in petroleum ether, R_f = 0.7) to afford presumably methyl 2-(1-ethoxyvinyl)-4-octylbenzoate (150 mg, 83.3% yield) as a yellow oil, instead of expected regioisomer.

[001065] Step 2: To a solution of methyl 2-(1-ethoxyvinyl)-4-octylbenzoate (150.0 mg, 0.47 mmol) in dichloromethane (5 mL) was added trifluoroacetic acid (0.21 mL, 2.83 mmol), stirred at 20 °C for 2 h and concentrated. The residue was taken up in EtOAc (20 mL), washed with water (10 mL x 2) and brine (10 mL), dried over MgSO₄ and concentrated to give crude methyl 2-acetyl-4-octylbenzoate (130 mg, 95% yield) as a yellow oil.

[001066] Compound 489 (formic acid salt) was prepared utilizing the methods in Example 388 (Compound 488). LCMS (Method 5-95 AB, ESI): $t_R = 0.676$ min, $[M+Na]^+ = 902.5$.

Example 390: Synthesis of Compound 490-P1 and Compound 490-P2




[001067] Step 1: To a solution of (S)-4-amino-2-(((benzyloxy)carbonyl)amino)butanoic acid (2000.0 mg, 7.93 mmol) in N,N-dimethylformamide (50 mL) were added 2-nitrobenzenesulfonylchloride (0.46 mL, 23.78 mmol) and triethylamine (4.42 mL, 31.71 mmol) dropwise. The reaction mixture was stirred at 25 °C for 12 h and filtered. To the filtrate was added water (50 mL) and the resulting precipitate was collected to obtain (S)-2-(((benzyloxy)carbonyl)amino)-4-(2-nitrophenylsulfonamido)butanoic acid (2000 mg, 4.5723 mmol, 57.7% yield) as a white solid. LCMS (Method 5-95 AB, ESI): $RT = 0.790$ min, $[M + H]^+ = 437.0$.

[001068] Step 2: To a solution of (S)-2-(((benzyloxy)carbonyl)amino)-4-(2-nitrophenylsulfonamido)butanoic acid (800.0 mg, 1.83 mmol) in N,N-dimethylformamide (5 mL) were added iodomethane (4.59 mL, 73.43 mmol) and potassium carbonate (758.3 mg, 5.49 mmol). The mixture was stirred at 25°C for 14 h, diluted with H₂O (20 mL) and extracted with EtOAc (35 mL x 3). The combined organic layers were washed with water (30 mL x4) and brine (40 mL), dried over Na₂SO₄ and concentrated. The residue was purified by flash column

chromatography (30% EtOAc in petroleum ether, $R_f = 0.3$) to obtain (S)-methyl 2-(((benzyloxy)carbonyl)amino)-4-(N-methyl-2-nitrophenylsulfonamido)butanoate (610 mg, 71.7% yield) as a yellow oil. LCMS (Method 5-95 AB, ESI): $RT = 0.813$ min, $[M + H]^+ = 466.1$.

[001069] Steps 3 and 4: Standard NaOH hydrolysis of the ester afforded (S)-2-

5 (((benzyloxy)carbonyl)amino)-4-(N-methyl-2-nitrophenylsulfonamido)butanoic acid was followed by standard HATU coupling (Example 5) afforded Compound 390-A.




[001070] Step 5: To a solution of Compound 490-A (220.0 mg, 0.19 mmol) in acetonitrile (3 mL) were added thioglycolic acid (0.09 mL, 1.27 mmol) and DBU (0.29 mL, 1.94 mmol). The reaction mixture was stirred at 25 °C for 4 h, diluted with H₂O (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic layers were washed with saturated aq. NaHCO₃ (40 mL x 2), water (40 mL x 2) and brine (40 mL), dried over Na₂SO₄ and concentrated. The residue was purified by preparative TLC (10% methanol in dichloromethane, $R_f = 0.4$) to obtain Compound 490-B (170 mg, 92.1% yield) as yellow solid. LCMS (Method 5-95 AB, ESI): $RT = 0.720$ min, $[M + H]^+ = 962.5$.

[001071] Step 6: Standard Boc-conditions afforded Compound 490-C.


[001072] Compound 490-P1 and Compound 490-P2 (formic acid salt) was prepared utilizing methods previously described and were isolated as pure diastereomers by reverse phase HPLC.

Data for Compound 490-P1: LCMS (Method 5-95 AB, ESI): $RT = 0.642$ min, $[M + H]^+ =$


868.6. Data for Compound 490-P2: LCMS (Method 5-95 AB, ESI): $RT = 0.658$ min, $[M + H]^+ = 868.6$.

Example 391: Synthesis of Compound 491


[001073]Compound **491** (formic acid salt) was prepared as a white solid utilizing methods previously described using octylbenzoic acid. LCMS (Method 5-95AB): $t_R = 0.756$ min/ 1.5 min, $[M + H]^+ = 882.5$.

Example 392: Synthesis of Compound 492

[001074]Compound 492 (formic acid salt) was prepared utilizing methods previously described. LC-MS: $m/z = 792$ (M+H).


Example 393: Synthesis of Compound 493

[001075]Compound 493 (formic acid salt) was prepared utilizing methods previously described. LC-MS: $m/z = 802$ (M+H).


Example 394: Synthesis of Compound 494

[001076]Compound 494 (formic acid salt) was prepared utilizing methods previously described from 1-aminocyclopropanecarbonitrile hydrochloride. LC-MS: $m/z = 928$ (M+H).


5 **Example 395: Synthesis of Compound 495**



[001077]Compound 495 (formic acid salt) was prepared utilizing methods previously described from 1-aminocyclopropanecarbonitrile hydrochloride. LC-MS: $m/z = 885$ (M+H).

Example 396: Synthesis of Compound 496

[001078]Compound 496 (formic acid salt) was prepared utilizing methods previously described from 2-amino-2-methyl-propanenitrile. LC-MS: $m/z = 930$ (M+H).

Example 397: Synthesis of Compound 497-P1 and Compound 497-P2

- 5 **[001079]** To a stirred solution of 60% NaH (1.0 g, 25 mmol) in THF (30 mL) was added 2-(2-bromoethyl)isoindoline-1,3-dione (5.6 g, 25 mmol) and the mixture was stirred at 0°C for 0.5 h, followed by the addition of diethyl malonate (4.0 g, 25 mmol). The resulting mixture was then stirred at room temperature for additional 16 h. The reaction mixture was diluted with EtOAc (100 mL), which was washed with brine (100 mL x 3). The organic layer was dried over Na₂SO₄,
- 10 concentrated and the residue was purified on silica gel flash column to afford compound diethyl 2-(2-(1,3-dioxoisindolin-2-yl)ethyl)malonate (1.2 g, 15.8% yield) as light yellow oil.

[001080] To a stirred solution of diethyl 2-(2-(1,3-dioxoisindolin-2-yl)ethyl)malonate (1.0 g, 3.0 mmol) in HOAc (9.0 mL) was added 10% HCl solution (3.0 mL) and the mixture was heated to 90°C while stirring and stirred for 2 h. The volatiles were removed and the residue was taken up in EtOAc (50 mL), which was washed with brine (30 mL x 2). The organic layer was dried over MgSO₄, concentrated and the residue was purified by HPLC to afford 4-(1,3-dioxoisindolin-2-yl)-2-(ethoxycarbonyl)butanoic acid (250 mg, 27.3% yield) as yellow oil. LCMS (Method 5-95 AB, ESI): $t_R = 0.941$, $[M + H]^+ = 306.1$.


[001081] Typical amide coupling (HATU, Example 5)) and ester hydrolysis with HOAc/HCl procedure was applied to afford 4-(1,3-dioxoisindolin-2-yl)-2-(undecylcarbamoyl)butanoic acid (150 mg) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 1.327$, $[M + H]^+ = 431.3$

[001082] Typical amide coupling (HATU, Example 5)) procedure was applied to Compound 101-G (150 mg, 0.21 mmol) and 4-(1,3-dioxoisindolin-2-yl)-2-(undecylcarbamoyl)butanoic acid (90 mg, 0.21 mmol) to afford Compound 497-A (220 mg, 93% yield) as a white solid.



[001083] Typical ester hydrolysis (LiOH/THF) procedure was applied to Compound 497-A (150 mg, 0.13 mmol) to yield the crude product. A solution of the product, hydrazine monohydrate (27 mg, 0.53 mmol) in MeOH (2 mL) was stirred at 80 °C for 16 h. The reaction mixture was diluted with DCM (30 mL), which was washed with brine (30 mL x 2). The organic layer was dried over Na₂SO₄, concentrated to give a residue. And the typical Boc addition procedure was applied to the residue to afford Compound 497-B (60 mg) as a white solid. LCMS (Method 5-95 AB, ESI): $t_R = 1.104$, $[M + H]^+ = 1082.7$.

[001084] Compound 497-P1 (4.4 mg) and Compound 497-P2 (1.9 mg) (formic acid salt) was prepared utilizing methods previously described and were isolated as separated diastereomers. Data for Compound 497-P1 and Compound 497-P2: LCMS (Method 5-95 AB, ESI): $t_R = 0.766$, $[M + H]^+ = 821.0$ (both are identical under these HPLC conditions).

Example 398: Synthesis of Compound 498




[001085] Compound 498 (free base) was prepared utilizing methods previously described. LCMS (Method 5-95 AB, ESI): $RT = 0.759$, $[M + H]^+ = 852.5$.

Example 399: Synthesis of Compound 499

[001086] Starting from Compound 101-F (Example 4), typical procedure including Boc removal (HCl/MeOH), amide coupling with 2-((tert-butoxycarbonyl)amino)acetic acid (HATU/DIEA), affords Compound 499-A.


[001087] Compound 499 (formic acid salt) was prepared utilizing methods previously described. LCMS (Method 5-95 AB, ESI): RT = 0.751, $[M + H]^+ = 969.7$; ^1H NMR (400 MHz, MeOH- d_4) δ 8.52 (brs, 3H), 7.34 (d, $J=8.0$ Hz, 2H), 7.28 (d, $J=8.0$ Hz, 1H), 7.21 (d, $J=8.0$ Hz, 1H), 7.12-7.04 (m, 4H), 6.85 (brs, 1H), 6.73 (brs, 1H), 6.39 (s, 1H), 5.17-5.13 (m, 1H), 4.84-4.79 (m, 2H), 4.22 (s, 2H), 4.20-4.05 (m, 6H), 3.58-3.43 (m, 8H), 3.17-3.10 (m, 3H), 2.97 (s, 3H), 2.62 (t, $J=6.8$ Hz, 2H), 2.41 (s, 3H), 2.33-2.24 (m, 1H), 2.19-2.10 (m, 1H), 1.68-1.51 (m, 2H), 1.40-1.25 (m, 11H), 0.92 (t, $J=6.8$ Hz, 3H).

Example 400: Synthesis of Compound 500


[001088] Compound 500-A is made utilizing the methods in Example 5 except (S)-4-((((9H-fluoren-9-yl)methoxy)carbonyl)amino)-2-(((benzyloxy)carbonyl)amino)butanoic acid is used in the coupling followed by Fmoc deprotection with piperidine.

[001089] To a solution of Compound 500-A (120 mg, 0.13 mmol) and Et₃N (53 μ L, 0.38 mmol) in DCM (5 mL) was added trimethylsilyl isocyanate (44 mg, 0.38 mmol) at 0°C. The resulting mixture was warmed up to room temperature while stirring and stirred at the same temperature for 2 h. The volatiles were removed and the residue was purified by prep-TLC to afford Compound 500-B (90 mg, 72% yield) as a white solid. LCMS (Method 5-95 AB, ESI): RT = 0.759, [M + H]⁺ = 991.7.


[001090] Starting from Compound 500-B, typical hydrogenation, amide coupling (HATU/DIEA) and global Boc removal (TFA/HFIP) procedure (Examples 5 and 7) was followed to afford Compound 500 as a white solid. LCMS (Method 5-95 AB, ESI): RT = 0.803, [M + H]⁺ = 911.6; ¹H NMR (400 MHz, MeOH-*d*₄) δ 8.51 (brs, 2H), 7.42-7.31 (m, 2H), 7.26 (d, *J*=8.4 Hz, 1H), 7.19 (d, *J*=8.4 Hz, 1H), 7.14-7.03 (m, 3H), 6.90 (brs, 1H), 6.81 (brs, 1H), 6.46 (s, 1H), 5.02-4.98 (m, 1H), 4.81-4.76 (m, 2H), 4.33-4.22 (m, 4H), 4.21 (s, 2H), 3.43-3.35 (m, 2H), 3.28-3.14 (m, 6H), 2.98 (s, 3H), 2.61 (t, *J*=7.6 Hz, 2H), 2.43 (s, 3H), 2.17-2.03 (m, 1H), 1.97-1.84 (m, 1H), 1.62 (brs, 2H), 1.39 - 1.26 (m, 13H), 0.92 (t, *J*=6.8 Hz, 3H).

Example 401: Synthesis of Compound 501

[001091] Compound 501 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95 AB, ESI): RT = 0.805, $[M + H]^+ = 910.5$


5 Example 402: Synthesis of Compound 502

[001092] Compound 502 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95AB): $t_R = 0.729$, $[M + H]^+ = 846.5$.

Example 403: Synthesis of Compound 503


10

[001093] Compound 503 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95AB): $t_R = 0.756$, $[M + H]^+ = 847.6$

Example 404: Synthesis of Compound 504


[001094] Compound 504 (formic acid salt) was prepared utilizing the methods previously described. LCMS (Method 5-95AB): $t_R = 0.793$, $[M + H]^+ = 875.6$.

Example 405: Synthesis of Compound 505




- 5 [001095] Compound 505 (free base) was prepared utilizing the methods previously described. LCMS (Method 5-95AB): $t_R = 0.719$, $[M + H]^+ = 819.4$.

Example 406: Synthesis of Compound 506



- [001096] Compound 506 (free base) was prepared utilizing the methods previously described .
 10 MS (ESI) for ($C_{53}H_{75}N_{11}O_9$): m/z 1010.7 $[M + H]^+$. HPLC t_R 2.73 min (10% AcCN/ H_2O – 90% AcCN/ H_2O , 3.0 min, 1.0 mL/min Kinetix C18, 4.8 x 50 mm).

Example 407: Synthesis of Compound 507



- [001097] Compound 507 (formic acid salt) was prepared utilizing the methods previously
 15 described. LCMS (Method 5-95AB): $t_R = 0.722$, $[M + H]^+ = 906.7$.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 358

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 358

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

R^{15} , R^{16} , R^{17} , and R^{18} are each independently H, $-(C_1-C_6)alkyl$, $-(C_3-C_6)cycloalkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$;

X is optionally substituted $-(C_1-C_6)alkyl$ -, $-(C_2-C_6)alkenyl$ -, $-(C_2-C_6)alkynyl$ -, $-(C_3-C_7)cycloalkyl$ -, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-O-(C_1-C_6)alkyl$ -, $-N(R^{24})(C_1-C_6)alkyl$ -, $-N(R^{24})(C_6-C_{10})aryl$ -, or $-SO_2(C_1-C_6)alkyl$ -;

Y is a bond, optionally substituted $-(C_1-C_6)alkyl$ -, $-(C_2-C_6)alkenyl$ -, $-(C_2-C_6)alkynyl$ -, $-(C_1-C_6)alkyl-N(R^{24})(C_1-C_6)alkyl$ -, $-O-(C_1-C_6)alkyl$ -, $-O(C_6-C_{10})aryl$ -, $-N(R^{24})(C_1-C_6)alkyl$ -, $-N(R^{24})SO_2(C_1-C_6)alkyl$ -, $-N(R^{24})C(O)(C_1-C_6)alkyl$ -, $-C(O)(C_1-C_6)alkyl$ -, $-S(C_1-C_6)alkyl$ -, $-SO_2(C_1-C_6)alkyl$ -, $-C(O)NH(C_1-C_6)alkyl$ -, $-(C_3-C_7)cycloalkyl$ -, optionally substituted $-C(O)N(R^{24})aryl$ -, optionally substituted $-N(R^{24})C(O)aryl$ -, optionally substituted $-N(R^{24})SO_2aryl$ -, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, $-NH_2$, $-CN$, $-CF_3$, $-CO_2H$, $-(C_1-C_{12})alkyl$ -, $-(C_2-C_{12})alkenyl$ -, $-(C_2-C_{12})alkynyl$ -, $-C(O)NR^{25}R^{26}$, $-O-(C_1-C_{12})alkyl$ -, $-N(R^{24})(C_1-C_{12})alkyl$ -, $-N(R^{24})C(O)(C_1-C_{12})alkyl$ -, optionally substituted $-(C_3-C_7)cycloalkyl$ -, $-(C_1-C_6)alkyl-heterocycloalkyl$ -, optionally substituted aryl, or optionally substituted heteroaryl;

each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl$ -, $-(C_1-C_6)heteroalkyl$ -, $-(C_1-C_6)alkyl-CO_2H$ -, $-C(O)(C_1-C_6)alkyl$ -, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)alkyl$; or two R^{31} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{23} is independently H or $-(C_1-C_6)alkyl$;

each R^{24} is independently H or $-(C_1-C_6)alkyl$;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl$; or R^{25} and R^{26} and

the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{27} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$ -, or optionally substituted $-(C_1-C_6)heteroalkyl$;


each R^{28} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$ -, or optionally substituted $-(C_1-C_6)heteroalkyl$;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

2. A compound of Formula (I'):



Formula (I');

wherein:

R¹ and R² are each independently H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -CH₂CH(OH)CH₂NH₂,
 5 -CH₂CH(heterocycloalkyl)CH₂NH₂, -CH₂C(O)NH₂, -CH₂C(O)N(H)CH₂CN, -(C₁-C₆)alkyl-
 C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-N(R²³)C(O)(C₁-
 C₆)alkylNR²¹R²², -(C₁-C₆)alkyl-C(O)N(R²³)(C₁-C₆)alkyl, -(C₁-C₆)alkyl-C(O)N(R²³)(C₁-
 C₆)alkyl-heterocycloalkyl, (C₁-C₆)heteroalkyl or optionally substituted heterocycloalkyl;

R³ is H, or -(C₁-C₆)alkyl;

10 R⁴ is H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OH, -(C₃-C₆)cycloalkyl, or -C(O)NH₂; or R³ and R⁴ are
 combined to form a heterocycloalkyl ring;

R⁵ is H, or -(C₁-C₆)alkyl; or R⁴ and R⁵ and the carbon atom to which that are attached form a
 cyclopropyl ring;

15 R⁶, R⁷, and R⁸ are each independently H, fluoro, hydroxyl, amino, optionally substituted alkyl or
 heteroalkyl or -(C₁-C₆)alkyl;

R⁹ is H, -(C₁-C₆)alkyl, -(C₁-C₆)haloalkyl, or -(C₃-C₆)cycloalkyl;

R¹⁰ is H, -(C₁-C₆)alkyl, -(C₁-C₆)haloalkyl, or -(C₃-C₆)cycloalkyl;

or R⁹ and R¹⁰ are combined to form a heterocycloalkyl or cycloalkyl ring

10 R¹¹ and R¹² are each independently H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-
 20 SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-
 C(O)NR²⁵R²⁶, -(C₁-C₆)heteroalkyl-CO₂H, -(C₁-C₆)alkyl-S(O)-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-
 N(H)CH=NH, -(C₁-C₆)alkyl-C(NH₂)=NH, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-
 N(H)SO₂NR²⁵R²⁶, -(C₁-C₆)alkyl-N(H)-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-heterocycloalkyl, optionally
 substituted -(C₁-C₆)alkyl-N(H)heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹¹ and R¹⁸ are
 25 combined to form an optionally substituted heterocycloalkyl ring, and R¹² is H;

R¹⁵, R¹⁶, R¹⁷, and R¹⁸ are each independently H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-
 C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;

X is optionally substituted $-(C_1-C_6)alkyl-$, $-(C_2-C_6)alkenyl-$, $-(C_2-C_6)alkynyl-$, $-(C_3-C_7)cycloalkyl-$, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-O-(C_1-C_6)alkyl-$, $-N(R^{24})(C_1-C_6)alkyl-$, $-N(R^{24})(C_6-C_{10})aryl-$, or $-SO_2(C_1-C_6)alkyl-$;

- 5 Y is a bond, optionally substituted $-(C_1-C_6)alkyl-$, $-(C_2-C_6)alkenyl-$, $-(C_2-C_6)alkynyl-$, $-(C_1-C_6)alkyl-N(R^{24})(C_1-C_6)alkyl-$, $-O-(C_1-C_6)alkyl-$, $-O(C_6-C_{10})aryl-$, $-N(R^{24})(C_1-C_6)alkyl-$, $-N(R^{24})SO_2(C_1-C_6)alkyl-$, $-N(R^{24})C(O)(C_1-C_6)alkyl-$, $-C(O)(C_1-C_6)alkyl-$, $-S(C_1-C_6)alkyl-$, $-SO_2(C_1-C_6)alkyl-$, $-C(O)NH(C_1-C_6)alkyl-$, $-(C_3-C_7)cycloalkyl-$, optionally substituted $-C(O)N(R^{24})aryl-$, optionally substituted $-N(R^{24})C(O)aryl-$, optionally substituted $-N(R^{24})SO_2aryl-$, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, $-NH_2$, $-CN$, $-CF_3$, $-CO_2H$, $-(C_1-C_{12})alkyl-$, $-(C_2-C_{12})alkenyl-$, $-(C_2-C_{12})alkynyl-$, $-C(O)NR^{25}R^{26}$, $-O-(C_1-C_{12})alkyl-$, $-N(R^{24})(C_1-C_{12})alkyl-$, $-N(R^{24})C(O)(C_1-C_{12})alkyl-$, optionally substituted $-(C_3-C_7)cycloalkyl-$, $-(C_1-C_6)alkyl-heterocycloalkyl-$, optionally substituted aryl, or optionally substituted heteroaryl;

- 15 each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl-$, $-(C_1-C_6)heteroalkyl-$, $-(C_1-C_6)alkyl-CO_2H$, $-C(O)(C_1-C_6)alkyl-$, $-C(O)O(C_1-C_6)alkyl-$, $-C(O)O(C_1-C_6)haloalkyl-$, $-C(=NH)(C_1-C_6)alkyl-$, $-C(=NH)N(R^{31})_2$, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)alkyl-$; or two R^{31} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{23} is independently H or $-(C_1-C_6)alkyl-$;

each R^{24} is independently H or $-(C_1-C_6)alkyl-$;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl-$; or R^{25} and R^{26} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

- 25 each R^{27} is independently halogen, $-NR^{23}R^{24}$, $-NC(O)R^{23}$, $-NC(O)NR^{23}R^{24}$, nitro, hydroxyl, optionally substituted $-(C_1-C_6)alkyl-$, optionally substituted $-(C_1-C_6)heteroalkyl-$, $-(C_1-C_6)alkoxy-$, $-C(O)(C_1-C_6)alkyl-$, or $-S(O)_2(C_1-C_6)alkyl-$;

each R^{28} is independently halogen, $-NR^{23}R^{24}$, $-NC(O)R^{23}$, $-NC(O)NR^{23}R^{24}$, nitro, hydroxyl, optionally substituted $-(C_1-C_6)alkyl-$, optionally substituted $-(C_1-C_6)heteroalkyl-$, $-(C_1-C_6)alkoxy-$,


- 30 $-C(O)(C_1-C_6)alkyl-$, or $-S(O)_2(C_1-C_6)alkyl-$;

p is 0, 1, or 2; and

q is 0, 1, or 2;


or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

3. The compound of claim 1 or 2 having the structure of Formula (Ia):




Formula (Ia).

4. The compound of any one of claims 1-3, wherein R^6 , R^7 , and R^8 are H.
5. The compound of any one of claims 1-4, wherein R^{15} and R^{16} are H.
6. The compound of any one of claims 1-3 having the structure of Formula (Ib):




Formula (Ib).

7. The compound of any one of claims 1-6, wherein R^{17} is $-\text{CH}_3$.
8. The compound of any one of claims 1-7, wherein R^{18} is H.
9. The compound of any one of claims 1-8, wherein R^5 is H.
10. The compound of any one of claims 1-9, wherein R^4 is H.
11. The compound of any one of claims 1-9, wherein R^4 is $-(\text{C}_1\text{-C}_6)\text{alkyl}$.
12. The compound of any one of claims 1-9, wherein R^4 is $-(\text{C}_3\text{-C}_6)\text{cycloalkyl}$.
13. The compound of any one of claims 1-8, wherein R^4 and R^5 and the carbon atom to which that are attached form a cyclopropyl ring.
14. The compound of any one of claims 1-13, wherein R^9 is $-(\text{C}_1\text{-C}_6)\text{alkyl}$.
15. The compound of claim 14, wherein R^9 is $-\text{CH}_3$.
16. The compound of claim 6 having the structure of Formula (Ic):



Formula (Ic).

17. The compound of any one of claims 1-16, wherein R^{11} is $-(C_1-C_6)alkyl-OR^{23}$.
18. The compound of claim 17, wherein R^{11} is $-CH_2CH_2OH$.
19. The compound of any one of claims 1-16, wherein R^{11} is $-(C_1-C_6)alkyl$.
20. The compound of any one of claims 1-16, wherein R^{11} is $-(C_1-C_6)alkyl-NR^{21}R^{22}$.
21. The compound of any one of claims 1-16, wherein R^{11} is $-(C_1-C_6)alkyl-NH_2$.
22. The compound of claim 21, wherein R^{11} is $-CH_2NH_2$.
23. The compound of claim 21, wherein R^{11} is $-CH_2CH_2NH_2$.
24. The compound of claim 21, wherein R^{11} is $-CH_2CH_2CH_2NH_2$.
25. The compound of claim 21, wherein R^{11} is $-CH_2CH_2CH_2CH_2NH_2$.
26. The compound of any one of claims 1-25, wherein R^1 and R^2 are each independently H, or $-(C_1-C_6)alkyl-NR^{21}R^{22}$.
27. The compound of any one of claims 1-25, wherein R^1 and R^2 are each independently $-(C_1-C_6)alkyl-NR^{21}R^{22}$.
28. The compound of any one of claims 1-25, wherein R^1 and R^2 are each $-CH_2CH_2NH_2$.
29. The compound of any one of claims 1-25, wherein R^1 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$ and R^2 is H.
30. The compound of any one of claims 1-25, wherein R^1 is $-CH_2CH_2NH_2$ and R^2 is H.
31. The compound of any one of claims 1-25, wherein R^1 is H and R^2 is $-(C_1-C_6)alkyl-NR^{21}R^{22}$.
32. The compound of any one of claims 1-25, wherein R^1 is H and R^2 is $-CH_2CH_2NH_2$.
33. The compound of claim 16 having the structure of Formula (Id):




Formula (Id);


wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{NH}_2$ or $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$.

34. The compound of any one of claims 1-33, wherein X is optionally substituted aryl.
- 5 35. The compound of any one of claims 1-33, wherein X is optionally substituted phenyl.
36. The compound of any one of claims 1-33, wherein X is optionally substituted heteroaryl.
37. The compound of claim 36, wherein X is heteroaryl which may be unsubstituted or substituted once or twice with $-(\text{C}_1\text{-C}_6)\text{alkyl}$.
38. The compound of claim 36, wherein X is heteroaryl which may be unsubstituted or substituted once with $-(\text{C}_1\text{-C}_6)\text{alkyl}$.
- 10 39. The compound of claim 36, wherein X is optionally substituted pyridine or optionally substituted pyrimidine.
40. The compound of claim 36, wherein X is pyridine which may be unsubstituted or substituted once or twice with $-(\text{C}_1\text{-C}_6)\text{alkyl}$.
- 15 41. The compound of claim 36, wherein X is pyridine which may be unsubstituted or substituted once with $-(\text{C}_1\text{-C}_6)\text{alkyl}$.
42. The compound of claim 36, wherein X is pyridine which may be unsubstituted or substituted once or twice with methyl.
43. The compound of claim 36, wherein X is pyrimidine which may be unsubstituted or substituted once or twice with $-(\text{C}_1\text{-C}_6)\text{alkyl}$.
- 20 44. The compound of claim 36, wherein X is pyrimidine which may be unsubstituted or substituted once with $-(\text{C}_1\text{-C}_6)\text{alkyl}$.
45. The compound of claim 36, wherein X is pyrimidine which may be unsubstituted or substituted once or twice with methyl.
- 25 46. The compound of claim 36, wherein X is pyridine substituted once with methyl.

47. The compound of any one of claims 1-33, wherein X is optionally substituted $-(C_1-C_6)alkyl-$.
48. The compound of any one of claims 1-47, wherein Y is optionally substituted aryl.
49. The compound of any one of claims 1-47, wherein Y is optionally substituted phenyl.
- 5 50. The compound of any one of claims 1-47, wherein Y is optionally substituted heteroaryl.
51. The compound of any one of claims 1-47, wherein Y is optionally substituted $-(C_1-C_6)alkyl-$.
52. The compound of any one of claims 1-47, wherein Y is $-O-(C_1-C_6)alkyl-$.
53. The compound of any one of claims 1-47, wherein Y is $-N(H)-(C_1-C_6)alkyl-$.
- 10 54. The compound of any one of claims 1-47, wherein Y is a bond.
55. The compound of any one of claims 1-54, wherein Z is $-(C_1-C_6)alkyl$.
56. The compound of any one of claims 1-54, wherein Z is optionally substituted aryl.
57. The compound of any one of claims 1-54, wherein Z is optionally substituted phenyl.
58. The compound of any one of claims 1-54, wherein Z is phenyl substituted once or twice
15 with $-(C_1-C_8)alkyl$.
59. The compound of any one of claims 1-54, wherein Z is phenyl substituted once with n-butyl, isobutyl or tert-butyl.
60. The compound of any one of claims 1-54, wherein Z is phenyl substituted once with n-butyl.
- 20 61. The compound of any one of claims 1-54, wherein Z is phenyl substituted once with isobutyl.
62. The compound of any one of claims 1-54, wherein Z is phenyl substituted once with tert-butyl.
63. The compound of any one of claims 1-54, wherein Z is optionally substituted heteroaryl.
- 25 64. The compound of any one of claims 1-54, wherein Z is optionally substituted $-(C_3-C_7)cycloalkyl$.
65. The compound of any one of claims 1-54, wherein Z is halogen.
66. The compound of any one of claims 1-33, wherein $-X-Y-Z$ is



5 67. A compound of Formula (II):



Formula (II);

wherein:

- R^1 and R^2 are each independently H, $-(\text{C}_1\text{--C}_6)\text{alkyl}$, $-(\text{C}_1\text{--C}_6)\text{alkyl-OR}^{23}$, $-\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{NH}_2$,
 10 $-\text{CH}_2\text{CH}(\text{heterocycloalkyl})\text{CH}_2\text{NH}_2$, $-\text{CH}_2\text{C}(\text{O})\text{NH}_2$, $-\text{CH}_2\text{C}(\text{O})\text{N}(\text{H})\text{CH}_2\text{CN}$, $-(\text{C}_1\text{--C}_6)\text{alkyl-}$
 $\text{C}(\text{O})\text{OR}^{23}$, $-(\text{C}_1\text{--C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$, $-(\text{C}_1\text{--C}_6)\text{alkyl-C}(\text{O})\text{NR}^{25}\text{R}^{26}$, $-(\text{C}_1\text{--C}_6)\text{alkyl-N}(\text{R}^{23})\text{C}(\text{O})(\text{C}_1\text{--}$
 $\text{C}_6)\text{alkylNR}^{21}\text{R}^{22}$, or $-(\text{C}_1\text{--C}_6)\text{alkyl-C}(\text{O})\text{N}(\text{R}^{23})(\text{C}_1\text{--C}_6)\text{alkyl}$, or optionally substituted
 heterocycloalkyl;
 R^3 is H, or $-(\text{C}_1\text{--C}_6)\text{alkyl}$;
 15 R^4 is H, $-(\text{C}_1\text{--C}_6)\text{alkyl}$, $-(\text{C}_1\text{--C}_6)\text{alkyl-OH}$, $-(\text{C}_3\text{--C}_6)\text{cycloalkyl}$, or $-\text{C}(\text{O})\text{NH}_2$; or R^3 and R^4 are
 combined to form a heterocycloalkyl ring;

R^5 is H, or $-(C_1-C_6)$ alkyl; or R^4 and R^5 and the carbon atom to which that are attached form a cyclopropyl ring;

R^6 , R^7 , and R^8 are each independently H, or $-(C_1-C_6)$ alkyl;

R^9 is H, $-(C_1-C_6)$ alkyl, $-(C_1-C_6)$ haloalkyl, or $-(C_3-C_6)$ cycloalkyl;

5 R^{10} is H, or $-(C_1-C_6)$ alkyl;

R^{11} and R^{12} are each independently H, $-NH_2$, $-(C_1-C_6)$ alkyl, $-(C_1-C_6)$ alkyl-OR²³, $-(C_1-C_6)$ alkyl-SR²³, $-(C_1-C_6)$ alkyl-C(O)OR²³, $-(C_1-C_6)$ alkyl-NR²¹R²², $-(C_1-C_6)$ alkyl-CN, $-(C_1-C_6)$ alkyl-C(O)NR²⁵R²⁶, $-(C_1-C_6)$ heteroalkyl-CO₂H, $-(C_1-C_6)$ alkyl-S(O)-(C₁-C₆)alkyl, $-(C_1-C_6)$ alkyl-N(H)CH=NH, $-(C_1-C_6)$ alkyl-N(H)C(NH)NH₂, $-(C_1-C_6)$ alkyl-heterocycloalkyl, optionally substituted $-(C_1-C_6)$ alkyl-N(H)heterocycloalkyl, or $-(C_1-C_6)$ alkyl-heteroaryl; or R^{11} and R^{18} are combined to form an optionally substituted heterocycloalkyl ring, and R^{12} is H;

10 R^{13} and R^{14} are each independently H, $-NH_2$, $-(C_1-C_6)$ alkyl, $-(C_1-C_6)$ alkyl-OR²³, $-(C_1-C_6)$ alkyl-SR²³, $-(C_1-C_6)$ alkyl-C(O)OR²³, $-(C_1-C_6)$ alkyl-NR²¹R²², $-(C_1-C_6)$ alkyl-CN, $-(C_1-C_6)$ alkyl-C(O)NR²⁵R²⁶, $-(C_1-C_6)$ alkyl-N(H)C(NH)NH₂, $-(C_1-C_6)$ alkyl-heterocycloalkyl, or $-(C_1-C_6)$ alkyl-heteroaryl; or R^{13} and R^{19} are combined to form an optionally substituted heterocycloalkyl ring, and R^{14} is H;

R^{15} , R^{16} , R^{17} , R^{18} , and R^{19} are each independently H, $-(C_1-C_6)$ alkyl, $-(C_3-C_6)$ cycloalkyl, $-(C_1-C_6)$ alkyl-OR²³, $-(C_1-C_6)$ alkyl-C(O)OR²³, or $-(C_1-C_6)$ alkyl-NR²¹R²²;

15 X is optionally substituted $-(C_1-C_6)$ alkyl-, $-(C_2-C_6)$ alkenyl-, $-(C_2-C_6)$ alkynyl, $-(C_3-C_7)$ cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-O-(C_1-C_6)$ alkyl-, $-N(R^{24})(C_1-C_6)$ alkyl-, $-N(R^{24})(C_6-C_{10})$ aryl-, or $-SO_2(C_1-C_6)$ alkyl-;

20 Y is a bond, optionally substituted $-(C_1-C_6)$ alkyl-, $-(C_2-C_6)$ alkenyl-, $-(C_2-C_6)$ alkynyl, $-(C_1-C_6)$ alkyl-N(R²⁴)(C₁-C₆)alkyl-, $-O-(C_1-C_6)$ alkyl-, $-O(C_6-C_{10})$ aryl-, $-N(R^{24})(C_1-C_6)$ alkyl-, $-N(R^{24})SO_2(C_1-C_6)$ alkyl-, $-N(R^{24})C(O)(C_1-C_6)$ alkyl-, $-C(O)(C_1-C_6)$ alkyl-, $-S(C_1-C_6)$ alkyl-, $-SO_2(C_1-C_6)$ alkyl-, $-C(O)NH(C_1-C_6)$ alkyl-, $-(C_3-C_7)$ cycloalkyl-, optionally substituted $-C(O)N(R^{24})$ aryl-, optionally substituted $-N(R^{24})C(O)$ aryl-, optionally substituted $-N(R^{24})SO_2$ aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

25 Z is H, halogen, $-NH_2$, $-CN$, $-CF_3$, $-(C_1-C_{12})$ alkyl-, $-(C_2-C_{12})$ alkenyl-, $-(C_2-C_{12})$ alkynyl-, $-C(O)NR^{25}R^{26}$, $-O-(C_1-C_{12})$ alkyl-, $-N(R^{24})(C_1-C_{12})$ alkyl-, $-N(R^{24})C(O)(C_1-C_{12})$ alkyl-, optionally substituted $-(C_3-C_7)$ cycloalkyl-, $-(C_1-C_6)$ alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

each R^{21} and R^{22} is independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)heteroalkyl$, $-(C_1-C_6)alkyl-CO_2H$, $-C(O)(C_1-C_6)alkyl$, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)alkyl$; or two R^{31} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{23} is independently H or $-(C_1-C_6)alkyl$;

each R^{24} is independently H or $-(C_1-C_6)alkyl$;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)alkyl$; or R^{25} and R^{26} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{27} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$, or optionally substituted $-(C_1-C_6)heteroalkyl$;

each R^{28} is independently halogen, optionally substituted $-(C_1-C_6)alkyl$, or optionally substituted $-(C_1-C_6)heteroalkyl$;


n is 0 or 1;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

68. A compound of Formula (II'):



Formula (II');

wherein:

R^1 and R^2 are each independently H, $-(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-OR^{23}$, $-CH_2CH(OH)CH_2NH_2$, $-CH_2CH(heterocycloalkyl)CH_2NH_2$, $-CH_2C(O)NH_2$, $-CH_2C(O)N(H)CH_2CN$, $-(C_1-C_6)alkyl-C(O)OR^{23}$, $-(C_1-C_6)alkyl-NR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)NR^{25}R^{26}$, $-(C_1-C_6)alkyl-N(R^{23})C(O)(C_1-C_6)alkylNR^{21}R^{22}$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl$, $-(C_1-C_6)alkyl-C(O)N(R^{23})(C_1-C_6)alkyl-heterocycloalkyl$, optionally substituted $(C_1-C_6)heteroalkyl$ or optionally substituted heterocycloalkyl;

R³ is H, or -(C₁-C₆)alkyl;

R⁴ is H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OH, -(C₃-C₆)cycloalkyl, or -C(O)NH₂; or R³ and R⁴ are combined to form a heterocycloalkyl ring;

R⁵ is H, or -(C₁-C₆)alkyl; or R⁴ and R⁵ and the carbon atom to which that are attached form a
5 cyclopropyl ring;

R⁶, R⁷, and R⁸ are each independently H, or -(C₁-C₆)alkyl;

R⁹ is H, -(C₁-C₆)alkyl, -(C₁-C₆)haloalkyl, or -(C₃-C₆)cycloalkyl;

R¹⁰ is H, or -(C₁-C₆)alkyl;

R¹¹ and R¹² are each independently H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-
10 SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-
C(O)NR²⁵R²⁶, -(C₁-C₆)heteroalkyl-CO₂H, -(C₁-C₆)alkyl-S(O)-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-
N(H)CH=NH, -(C₁-C₆)alkyl-C(NH₂)=NH, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-
N(H)SO₂NR²⁵R²⁶, -(C₁-C₆)alkyl-N(H)-C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-heterocycloalkyl, optionally
substituted -(C₁-C₆)alkyl-N(H)heterocycloalkyl, or -(C₁-C₆)alkyl-heteroaryl; or R¹¹ and R¹⁸ are
15 combined to form an optionally substituted heterocycloalkyl ring, and R¹² is H;

R¹³ and R¹⁴ are each independently H, -NH₂, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OR²³, -(C₁-C₆)alkyl-
SR²³, -(C₁-C₆)alkyl-C(O)OR²³, -(C₁-C₆)alkyl-NR²¹R²², -(C₁-C₆)alkyl-CN, -(C₁-C₆)alkyl-
C(O)NR²⁵R²⁶, -(C₁-C₆)alkyl-N(H)C(NH)NH₂, -(C₁-C₆)alkyl-heterocycloalkyl, or -(C₁-C₆)alkyl-
heteroaryl; or R¹³ and R¹⁹ are combined to form an optionally substituted heterocycloalkyl ring,
20 and R¹⁴ is H;

R¹⁵, R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are each independently H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₁-
C₆)alkyl-OR²³, -(C₁-C₆)alkyl-C(O)OR²³, or -(C₁-C₆)alkyl-NR²¹R²²;

X is optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₃-
C₇)cycloalkyl-, optionally substituted heterocycloalkyl, optionally substituted aryl, optionally
25 substituted heteroaryl, -O-(C₁-C₆)alkyl-, -N(R²⁴)(C₁-C₆)alkyl-, -N(R²⁴)(C₆-C₁₀)aryl-, or -SO₂(C₁-
C₆)alkyl-;

Y is a bond, optionally substituted -(C₁-C₆)alkyl-, -(C₂-C₆)alkenyl-, -(C₂-C₆)alkynyl, -(C₁-
C₆)alkyl-N(R²⁴)(C₁-C₆)alkyl-, -O-(C₁-C₆)alkyl-, -O(C₆-C₁₀)aryl-, -N(R²⁴)(C₁-C₆)alkyl-, -
N(R²⁴)SO₂(C₁-C₆)alkyl-, -N(R²⁴)C(O)(C₁-C₆)alkyl-, -C(O)(C₁-C₆)alkyl-, -S(C₁-C₆)alkyl-, -
30 SO₂(C₁-C₆)alkyl-, -C(O)NH(C₁-C₆)alkyl-, -(C₃-C₇)cycloalkyl-, optionally substituted -
C(O)N(R²⁴)aryl-, optionally substituted -N(R²⁴)C(O)aryl-, optionally substituted -
N(R²⁴)SO₂aryl-, optionally substituted aryl, or optionally substituted heteroaryl;

Z is H, halogen, -NH₂, -CN, -CF₃, -(C₁-C₁₂)alkyl-, -(C₂-C₁₂)alkenyl, -(C₂-C₁₂)alkynyl, -
C(O)NR²⁵R²⁶, -O-(C₁-C₁₂)alkyl, -N(R²⁴)(C₁-C₁₂)alkyl, -N(R²⁴)C(O)(C₁-C₁₂)alkyl, optionally

substituted $-(C_3-C_7)$ cycloalkyl, $-(C_1-C_6)$ alkyl-heterocycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl;

each R^{21} and R^{22} is independently H, $-(C_1-C_6)$ alkyl, $-(C_1-C_6)$ heteroalkyl, $-(C_1-C_6)$ alkyl- CO_2H , $-C(O)(C_1-C_6)$ alkyl, $-C(O)N(R^{31})_2$, or $-SO_2N(R^{31})_2$; or R^{21} and R^{22} and the nitrogen atom to which

that are attached form a heterocycloalkyl ring;

each R^{31} is independently H or $-(C_1-C_6)$ alkyl; or two R^{31} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{23} is independently H or $-(C_1-C_6)$ alkyl;

each R^{24} is independently H or $-(C_1-C_6)$ alkyl;

each R^{25} and R^{26} is independently H or optionally substituted $-(C_1-C_6)$ alkyl; or R^{25} and R^{26} and the nitrogen atom to which that are attached form a heterocycloalkyl ring;

each R^{27} is independently halogen, $-NR^{23}R^{24}$, $-NC(O)R^{23}$, $-NC(O)NR^{23}R^{24}$, nitro, hydroxyl, optionally substituted $-(C_1-C_6)$ alkyl, optionally substituted $-(C_1-C_6)$ heteroalkyl, $-(C_1-C_6)$ alkoxy, $-C(O)(C_1-C_6)$ alkyl, or $-S(O)_2(C_1-C_6)$ alkyl;

each R^{28} is independently halogen, $-NR^{23}R^{24}$, $-NC(O)R^{23}$, $-NC(O)NR^{23}R^{24}$, nitro, hydroxyl, optionally substituted $-(C_1-C_6)$ alkyl, optionally substituted $-(C_1-C_6)$ heteroalkyl, $-(C_1-C_6)$ alkoxy, $-C(O)(C_1-C_6)$ alkyl, or $-S(O)_2(C_1-C_6)$ alkyl;


n is 0 or 1;

p is 0, 1, or 2; and

q is 0, 1, or 2;

or a pharmaceutically acceptable salt, solvate, or prodrug thereof.

69. The compound of claim 67 or 68 having the structure of Formula (IIa):




Formula (IIa).

70. The compound of any one of claims 67-69, wherein R^6 , R^7 , and R^8 are H.


71. The compound of any one of claims 67-70, wherein R^{15} and R^{16} are H.

72. The compound of any one of claims 67-69 having the structure of Formula (IIb):



Formula (IIb).

73. The compound of any one of claims 67-72, wherein R¹⁸ is H.
74. The compound of any one of claims 67-73, wherein R¹⁹ is H.
- 5 75. The compound of any one of claims 67-74, wherein R¹⁷ is -CH₃.
76. The compound of any one of claims 67-75, wherein R⁵ is H.
77. The compound of any one of claims 67-76, wherein R⁴ is H.
78. The compound of any one of claims 67-76, wherein R⁴ is -(C₁-C₆)alkyl.
79. The compound of any one of claims 67-76, wherein R⁴ is -(C₃-C₆)cycloalkyl.
- 10 80. The compound of any one of claims 67-75, wherein R⁴ and R⁵ and the carbon atom to which that are attached form a cyclopropyl ring.
81. The compound of any one of claims 67-80, wherein R⁹ is -(C₁-C₆)alkyl.
82. The compound of claim 81, wherein R⁹ is -CH₃.
83. The compound of any one of claims 67-82, wherein R¹ and R² are each independently H,
- 15 or -(C₁-C₆)alkyl-NR²¹R²².
84. The compound of any one of claims 67-82, wherein R¹ and R² are each independently -(C₁-C₆)alkyl-NR²¹R²².
85. The compound of any one of claims 67-82, wherein R¹ and R² are each -CH₂CH₂NH₂.
86. The compound of any one of claims 67-69 having the structure of Formula (IIc):



Formula (IIc);

wherein R^1 and R^2 are each independently H or $-\text{CH}_2\text{CH}_2\text{NH}_2$.

87. The compound of any one of claims 67-86, wherein R^{11} is $-(\text{C}_1\text{-C}_6)\text{alkyl-OR}^{23}$.

88. The compound of claim 87, wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{OH}$.

89. The compound of any one of claims 67-86, wherein R^{11} is $-(\text{C}_1\text{-C}_6)\text{alkyl}$.

5 90. The compound of any one of claims 67-86, wherein R^{11} is $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$.

91. The compound of any one of claims 67-86, wherein R^{11} is $-(\text{C}_1\text{-C}_6)\text{alkyl-NH}_2$.

92. The compound of claim 91, wherein R^{11} is $-\text{CH}_2\text{NH}_2$.

93. The compound of claim 91, wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{NH}_2$.

94. The compound of claim 91, wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$.


10 95. The compound of claim 91, wherein R^{11} is $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$.

96. The compound of any one of claims 67-95, wherein R^{13} is $-(\text{C}_1\text{-C}_6)\text{alkyl-OR}^{23}$.

97. The compound of claim 96, wherein R^{13} is $-\text{CH}_2\text{OH}$.

98. The compound of claim 96, wherein R^{13} is $-\text{CH}_2\text{CH}_2\text{OH}$.

99. The compound of any one of claims 67-69 having the structure of Formula (IId):



15

Formula (IId).

100. The compound of claim 99, wherein R^{17} is $-\text{CH}_3$.

101. The compound of claim 99 or claim 100, wherein R^5 is H.

102. The compound of any one of claims 99-101, wherein R^4 is H.

20 103. The compound of any one of claims 99-101, wherein R^4 is $-(\text{C}_1\text{-C}_6)\text{alkyl}$.

104. The compound of any one of claims 99-101, wherein R^4 is $-(\text{C}_3\text{-C}_6)\text{cycloalkyl}$.

105. The compound of claim 99 or claim 100, wherein R^4 and R^5 and the carbon atom to which that are attached form a cyclopropyl ring.

106. The compound of any one of claims 99-105, wherein R^9 is $-(\text{C}_1\text{-C}_6)\text{alkyl}$.


25 107. The compound of claim 106, wherein R^9 is $-\text{CH}_3$.

108. The compound of any one of claims 99-107, wherein R^1 and R^2 are each independently H, or $-(\text{C}_1\text{-C}_6)\text{alkyl-NR}^{21}\text{R}^{22}$.

109. The compound of any one of claims 99-107, wherein R^1 and R^2 are each independently - $(C_1-C_6)alkyl-NR^{21}R^{22}$.

110. The compound of any one of claims 99-107, wherein R^1 and R^2 are each $-CH_2CH_2NH_2$.

111. The compound any one of claims 67-69 having the structure of Formula (IIe):



Formula (IIe);

wherein R^1 and R^2 are each independently H or $-CH_2CH_2NH_2$.

112. The compound of any one of claims 67-111, wherein X is optionally substituted aryl.

113. The compound of any one of claims 67-111, wherein X is optionally substituted phenyl.

10 114. The compound of any one of claims 67-111, wherein X is optionally substituted heteroaryl.

115. The compound of claim 114, wherein X is optionally substituted pyridine or optionally substituted pyrimidine.

15 116. The compound of any one of claims 67-111, wherein X is optionally substituted $-(C_1-C_6)alkyl-$.

117. The compound of any one of claims 67-116, wherein Y is optionally substituted aryl.

118. The compound of any one of claims 67-116, wherein Y is optionally substituted phenyl.

119. The compound of any one of claims 67-116, wherein Y is optionally substituted heteroaryl.

20 120. The compound of any one of claims 67-116, wherein Y is optionally substituted $-(C_1-C_6)alkyl-$.

121. The compound of any one of claims 67-116, wherein Y is $-O-(C_1-C_6)alkyl-$.

122. The compound of any one of claims 67-116, wherein Y is $-N(H)-(C_1-C_6)alkyl-$.

123. The compound of any one of claims 67-116, wherein Y is a bond.

25 124. The compound of any one of claims 67-123, wherein Z is $-(C_1-C_6)alkyl-$.

125. The compound of any one of claims 67-123, wherein Z is optionally substituted aryl.

126. The compound of any one of claims 67-123, wherein Z is optionally substituted phenyl.

130. The compound of any one of claims 67-111, wherein -X-Y-Z is



134. The method of claim 133, wherein the bacterial infection is an infection involving *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Pseudomonas acidovorans*, *Pseudomonas alcaligenes*, *Pseudomonas putida*, *Stenotrophomonas maltophilia*, *Burkholderia cepacia*,

- Aeromonas hydrophilia, Escherichia coli, Citrobacter freundii, Salmonella typhimurium, Salmonella typhi, Salmonella paratyphi, Salmonella enteritidis, Shigella dysenteriae, Shigella flexneri, Shigella sonnei, Enterobacter cloacae, Enterobacter aerogenes, Klebsiella pneumoniae, Klebsiella oxytoca, Serratia marcescens, Francisella tularensis, Morganella morganii, Proteus mirabilis, Proteus vulgaris, Providencia alcalifaciens, Providencia rettgeri, Providencia stuartii, Acinetobacter baumannii, Acinetobacter calcoaceticus, Acinetobacter haemolyticus, Yersinia enterocolitica, Yersinia pestis, Yersinia pseudotuberculosis, Yersinia intermedia, Bordetella pertussis, Bordetella parapertussis, Bordetella bronchiseptica, Haemophilus influenzae, Haemophilus parainfluenzae, Haemophilus haemolyticus, Haemophilus parahaemolyticus, Haemophilus ducreyi, Pasteurella multocida, Pasteurella haemolytica, Branhamella catarrhalis, Helicobacter pylori, Campylobacter fetus, Campylobacter jejuni, Campylobacter coli, Borrelia burgdorferi, Vibrio cholerae, Vibrio parahaemolyticus, Legionella pneumophila, Listeria monocytogenes, Neisseria gonorrhoeae, Neisseria meningitidis, Kingella, Moraxella, Gardnerella vaginalis, Bacteroides fragilis, Bacteroides distasonis, Bacteroides 3452A homology group, Bacteroides vulgatus, Bacteroides ovalus, Bacteroides thetaiotaomicron, Bacteroides uniformis, Bacteroides eggerthii, Bacteroides splanchnicus, Clostridium difficile, Mycobacterium tuberculosis, Mycobacterium avium, Mycobacterium intracellulare, Mycobacterium leprae, Corynebacterium diphtheriae, Corynebacterium ulcerans, Streptococcus pneumoniae, Streptococcus agalactiae, Streptococcus pyogenes, Enterococcus faecalis, Enterococcus faecium, Staphylococcus aureus, Staphylococcus epidermidis, Staphylococcus saprophyticus, Staphylococcus intermedius, Staphylococcus hyicus subsp. hyicus, Staphylococcus haemolyticus, Staphylococcus hominis, or Staphylococcus saccharolyticus.
135. The method of claim 133 or 134, wherein the bacterial infection is an infection involving a Gram-negative bacteria.
136. The method of any one of claims 133-135, wherein administering comprises topical administration.
137. The method of any one of claims 133-136 further comprising administering a second therapeutic agent.
138. The method of claim 137, wherein the second therapeutic agent is not an SpsB inhibitor.
139. The method of claim 138, wherein the second therapeutic agent is an aminoglycoside antibiotic, fluoroquinolone antibiotic, β -lactam antibiotic, macrolide antibiotic, glycopeptide antibiotic, rifampicin, chloramphenicol, fluoramphenicol, colistin, mupirocin, bacitracin, daptomycin, or linezolid.
140. The method of claim 138, wherein the second therapeutic agent is a β -lactam antibiotic.

141. The method of claim 140, wherein the β -lactam antibiotic is selected from penicillins, monobactams, cephalosporins, cephamycins, and carbapenems.

142. The method of claim 141, wherein the β -lactam antibiotic is selected from Azlocillin, Amoxicillin, Ampicillin, Doripenem, Meropenem, Biapenem, Cefamandole, Imipenem,

5 Mezlocillin, Cefmetazole, Cefprozil, Piperacillin/tazobactam, Carbenicillin, Cefaclor, Cephalothin, Ertapenem, Cefazolin, Cefepime, Cefonicid, Cefoxitin, Ceftazidime, Oxacillin, Cefdinir, Cefixime, Cefotaxime, Cefotetan, Cefpodoxime, Ceftizoxime, Ceftriaxone, Faropenem, Mecillinam, Methicillin, Moxalactam, Ticarcillin, Tomopenem, Ceftobiprole, Ceftaroline, Flomoxef, Cefiprome, and Cefozopran.

10 143. The method of claim 141, further comprising administering a β -lactamase inhibitor.

1/1

FIGURE 1

