



US00PP15393P2

(12) **United States Plant Patent**  
**Chou et al.**

(10) **Patent No.:** **US PP15,393 P2**

(45) **Date of Patent:** **Nov. 30, 2004**

(54) **IMPATIENS PLANT NAMED**  
**'BALFUSINRED'**

(50) Latin Name: *Impatiens walleriana*×*auricom*  
Varietal Denomination: **Balfusinred**

(75) Inventors: **Tau San Chou**, Batavia, IL (US);  
**Kristin Barry**, Gainesville, FL (US)

(73) Assignee: **Ball Horticultural Company**, West  
Chicago, IL (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/741,994**

(22) Filed: **Dec. 19, 2003**

(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./317**

(58) **Field of Search** ..... **Plt./317**

*Primary Examiner*—Kent Bell

(74) *Attorney, Agent, or Firm*—Wood, Phillips, Katz, Clark  
& Mortimer

(57) **ABSTRACT**

A new and distinct cultivar of *Impatiens* plant named  
'Balfusinred', characterized by its single, cupped flowers,  
medium green-colored foliage, upright and mounded habit,  
and excellent basal branching.

**1 Drawing Sheet**

**1**

Latin name of genus and species of plant claimed: *Impa-*  
*tians walleriana*×*auricom*.

Variety denomination: 'Balfusinred'.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct *Impa-*  
*tians* plant botanically known as *Impatiens walleriana*×  
*auricom* and hereinafter referred to by the cultivar name  
'Balfusinred'.

The new cultivar was developed by the inventors in a  
controlled breeding program during March 1999 at Elburn,  
Ill. The objective of the breeding program was to develop  
*Impatiens* cultivars with numerous flowers, new and unique  
flower shape and colors, excellent basal branching and  
upright compact to moderate growth habit.

The female (seed) parent of 'Balfusinred' was the propri-  
etary *Impatiens walleriana* selection designated '12660'  
(not patented) characterized by its upright growth habit,  
white-colored flowers, and medium green-colored foliage.  
The male (pollen) parent of 'Balfusinred' was the propri-  
etary *Impatiens auricom* selection designated '10449-1-2'  
(not patented) characterized by its upright growth habit,  
yellow-colored flowers, and dark green-colored foliage.  
'Balfusinred' was discovered and selected as a single flow-  
ering plant within the progeny of the above stated cross in  
February of 2000 at Elburn, Ill. and initially designated PAS  
'1-08'.

Asexual reproduction of the new cultivar by shoot tip or  
stem cuttings taken since February 2000 at Elburn, Ill. and  
West Chicago, Ill. has demonstrated that the new cultivar  
reproduce true to type with all the characteristics, as herein  
described, firmly fixed and retained through successive  
generations of such asexual propagation.

**SUMMARY OF THE INVENTION**

The new cultivar has not been observed under all possible  
environmental conditions to date. Accordingly, it is possible  
that the phenotype may vary somewhat with variations in the  
environment, such as temperature, light intensity, and day  
length without, however, any variance in genotype.

It was repeatedly found that the cultivar of the present  
invention:

**2**

1. Exhibits single, slightly cupped, flowers,
2. Forms medium green-colored foliage,
3. Exhibits an upright and mounded growth habit, and
4. Exhibits a good basal branching character.

Plants of the new cultivar differ from plants of the female  
parent primarily in flower color, flower shape and leaf shape  
and from plants of the male parent primarily in flower color,  
flower shape, leaf size and branching habit. Plants of the new  
cultivar can be compared to 'Balfusheat' disclosed in U.S.  
Plant patent application Ser. No. 10/741,696. 'Balfusheat'  
and 'Balfusinred' were derived from crossing the same two  
cultivars, however, they exhibit very different flower colors.

Plants of the new cultivar can be compared to plants of  
'96-009-4' (U.S. Plant Pat. No. 10,973). In a side-by-side  
comparison conducted in West Chicago, Ill., plants of the  
new cultivar differed from plants of '96-009-4' primarily in  
the following characteristics:

1. Plants of the new cultivar have a more compact growth  
habit than plants of '96-009-4', and
2. Plants of the new cultivar have larger flowers than  
plants of '96-009-4'.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographs show, as nearly true as it  
is reasonably possible to make the same in color illustrations  
of this type, typical flower and foliage characteristics of the  
new cultivar. Colors in the photographs differ from the color  
values cited in the detailed description, which accurately  
describe the colors of 'Balfusinred'. The plants were grown  
for 13 weeks in a greenhouse at West Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and  
flowering habit of 'Balfusinred'.

FIG. 2 illustrates a close-up view of a single flower of  
'Balfusinred'.

**DETAILED DESCRIPTION**

The chart used in the identification of colors described  
herein is The R.H.S. Colour Chart of The Royal Horticul-  
tural Society, London, England, 1995 edition, except where  
general color terms of ordinary significance are used. The

color values were determined on Sep. 3, 2003 between 9:00 and 10:45 a.m. under natural light conditions.

The following measurements and comparisons describe plants produced from shoot tip or stem (with two nodes) cuttings taken from stock plants and grown under greenhouse conditions comparable to those used in commercial practice in West Chicago, Ill. The plants were grown in 10 cm pots for 15 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 72° F. during the day and approximately 65° F. during the night. Greenhouse light levels were maintained at approximately 4,000–6,000 footcandles during the day.

Botanical classification: *Impatiens walleriana* × *auricom*a cultivar 'Balfusinred'.

Parentage:

*Female (seed) parent*.—Proprietary *Impatiens walleriana* selection designated '12660' (not patented).

*Male (pollen) parent*.—Proprietary *Impatiens auricom*a selection designated '10449-1-2' (not patented).

Propagation:

*Type cutting*.—Shoot tip or stem (with two nodes).

*Time to initiate roots*.—Approximately 7–14 days.

*Time to develop a rooted cutting*.—Approximately 21 days.

*Root description*.—Fine, fibrous.

*Rooting habit*.—Freely branching.

Plant description:

*Habit of growth*.—Moderately vigorous with good basal branching. A mature plant, 15 weeks after the planting of a rooted cutting, commonly measures approximately 23.7 cm in height and approximately 38.5 cm in width.

*Plant form*.—Upright and mounded.

*Lateral branches*.—Quantity: Approximately 4. Length: Approximately 17.1 cm. Diameter: Approximately 8 mm. Texture: Glabrous. Color: 145A with faint overlay of 45A. Internode length: Approximately 3.3 cm.

*Foliage*.—Type: Simple. Arrangement of lower leaves: Alternate. Arrangement of upper leaves: Whorls. Shape: Elliptic. Apex: Acuminate/cuspidate. Base: Attenuate. Margin: Crenate with ciliation. Texture of upper and lower surface: Glabrous. Venation pattern: Pinnate, arcuate. Size of mature foliage: Length: Approximately 7.2 cm. Width: Approximately 3.9 cm. Color of mature foliage: Upper surface is 137A with veins and mid-vein of 145A. Lower surface is 138B with veins and mid-vein of 145A. Petiole: Diameter: Approximately 2 mm. Length: Approximately 2.7 cm with 2 mm stipules along edges. Texture: Upper and lower surface: Glabrous. Color: Upper and lower surface: 145A with overlay of 46A.

Flower description:

*Flower type*.—Single, not fragrant with self-cleaning petals.

*Flowering habit*.—Freely flowering with flowers positioned above the foliage and facing outward.

*Natural flowering season*.—Year round in greenhouse environment. Flowering is continuous from spring until fall in the garden.

*Flower longevity on plant*.—Approximately 5 days.

*Quantity of flowers*.—Approximately 8 flowers and 7 buds per stem at any one time.

*Mature flower buds (just before opening)*.—Shape: Ovate. Length: Approximately 1.2 cm. Diameter: Approximately 9 mm. Texture: Glabrous. Color: 38B.

*Flower shape*.—Oval, cupped when first open becoming more flat with age.

*Flower size*.—Length: Approximately 3.1 cm. Width: Approximately 2.5 cm. Depth: Approximately 1.3 cm.

*Petals*.—Number: Five per flower. Texture: Glabrous. Appearance: Iridescent.

*Superior petal*.—Length: Approximately 1.2 cm. Width: Approximately 1.2 cm. Shape: Cucullate. Margin: Entire. Apex: Emarginate. Base: Truncate. Color of upper surface: 55B with overlay of 37A. Color of lower surface: 39C.

*Lateral petals*.—Length: 1.3 cm. Width: 1.1 cm. Shape: Obovate. Margin: Entire. Apex: Obtuse. Base: Attenuate. Color of upper surface: Margin at lower edge is 55A, rest of margin is 37A. There is central spot of 33C and base of N155B. Color of lower surface: 29B.

*Lower petals*.—Length: 1.8 cm. Width: 1.1 cm. Shape: Obovate. Margin: Entire. Apex: Obtuse. Base: Attenuate. Color of upper surface: 55A with center of petal having a small area of N30B. Color of lower surface: Background of 6D with marginal overlay of 55A and central spot of 33B.

*Peduncles*.—Strength: Strong. Angle to stem: Acute. Length: Approximately 2.1 cm. Diameter: Approximately 1 mm. Texture: Glabrous. Color: 144C.

*Calyx*.—Consists of 3 sepals. Lateral sepals: Length: Approximately 9 mm. Width: Approximately 6 mm. Texture: Glabrous. Color of both surfaces: 154D with apex of 144B. Lower sepal: Length: Approximately 1.3 cm. Width: Approximately 1 cm. Texture: Glabrous. Color of both surfaces: 16C with apex of 1447B. Base of lower sepal is modified to form spur.

*Spur*.—Quantity: One per flower. Double tipped. Length: Approximately 1.3 cm. Diameter at flower: Approximately 2 mm. Diameter of tips: Approximately 1 mm. Aspect: Moderately curved downward. Color: 150D.

*Reproductive organs*.—Androecium: Five stamens, anthers are fused together forming one organ that surrounds the pistil. Filament color: 8B. Anther shape: Hooded. Anther length: Approximately 2 mm. Anther color: N78D. Pollen: None observed. Gynoecium: Pistil number: One per flower. Pistil length: 6 mm. Stigma shape: Five pointed star. Stigma length: 0.5 mm. Stigma color: Colorless, translucent. Style length: 1.5. Style color: Colorless, translucent. Ovary length: 4 mm. Ovary diameter: 2 mm. Ovary texture: Glabrous. Ovary color: N144C.

Seed and fruit development: Neither seed nor fruit production has been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Impatiens* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Impatiens* plant named 'Balfusinred' substantially, as herein shown and described, which:

1. Exhibits single, cupped, flowers,
2. Forms medium green-colored foliage,
3. Exhibits a good basal branching character, and
4. Exhibits an upright and mounded growth habit.

\* \* \* \* \*



Figure 1




Figure 2